

Enteric fevers and inoculation in the French Army during the Great War : 1914-1918.

Publication/Creation

[S.l. : s.n., ca.1920?]

Persistent URL

<https://wellcomecollection.org/works/kkwa5ktn>

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

*Enteric fevers and inoculation in the French Army
during the Great War: 1914 - 1918*

<i>Period</i>	<i>Cases</i>	<i>Total</i>	<i>Remarks</i>
<i>Aug. to Oct. 1914</i>	<i>18,867</i>	<i>18,867</i>	<i>No inoculation</i>
<i>Nov. to Dec. 1914</i>	<i>26,220</i>		<i>Antityphoid inoculation adopted and gradually became practically universal.</i>
<i>1st. Qtr. 1915</i>	<i>27,597</i>		
<i>2.nd. Qtr. 1915</i>	<i>10,684</i>		
<i>3. rd Qtr. 1915</i>	<i>17,653</i>	<i>82,154</i>	
<i>4. th. Qtr. 1915</i>	<i>11,114</i>		<i>Triple typhoid-paratyphoid vaccine substituted for simple antityphoid vaccine.</i>
<i>1916</i>	<i>12,482</i>		
<i>1917</i>	<i>1,678</i>		
<i>1918</i>	<i>757</i>	<i>26,031</i>	

Figures taken from "Le Service de Santé pendant la guerre. 1914-1918." Tome IV pp 707-712.