

**Udderine : a wonderful remedy for udder ill, milk fever, garget, chills, fever and inflammations of all kinds, in horses, cattle & sheep ...**

**Publication/Creation**

[S.l : s.n., 189-?]

**Persistent URL**

<https://wellcomecollection.org/works/vrsjebf6>

**License and attribution**

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection  
183 Euston Road  
London NW1 2BE UK  
T +44 (0)20 7611 8722  
E [library@wellcomecollection.org](mailto:library@wellcomecollection.org)  
<https://wellcomecollection.org>

### **DIRECTIONS.**

Keep the bottle upright and well corked, and in all cases of treatment with Udderine, let the animal be kept warm.

#### **SHEEP,**

SCOUR, GARGET & UDDER ILL, Give half a drench every two hours, with a quarter of a pint of water. CHILLS AND INFLAMMATION AFTER LAMBING. A dose every three hours until relieved, then at longer intervals.

#### **HORSES,**

GRIPES AND COLIC, Give one mark on the bottle every half hour if the pain be violent, in mild cases a dose every two hours.

CHILLS, COLDS AND INFLAMMATION OF THE LUNGS & KIDNEYS, A dose of Udderine every half hour until relieved, then every two or three hours.

# **Udderine**

a wonderful remedy for Udder Ill, Milk Fever, Garget, Chills, Fever and Inflammations of all kinds, in

**HORSES, CATTLE & SHEEP.**

#### **For Cows**

with Milk Fever, Hard or Swollen Udders, Garget, Chills and Inflammation of the Lungs and Kidneys.

#### **For Horses**

with Chills, Gripes, Colic and Inflammation.

#### **For Sheep**

with Scour, Garget, Udder Ill, and Inflammation after Lambing.

### **DIRECTIONS.**

This bottle contains eight drenches or doses.

Each mark on it represents one dose.

#### **COWS,**

MILK FEVER, Give a dose undiluted every quarter of an hour until an improvement is seen, up to four doses, then a dose every hour or two until the fever diminishes. As a preventative of Milk fever give two doses in an equal quantity of water about an hour before calving, and again three hours after.

#### **SCOUR,**

CHILLS, UDDER ILL, GARGET AND INFLAMMATION OF THE LUNGS OR KIDNEYS. Give two doses every three hours with half a pint of cold water.