

Protect your pet from worms / Hoechst Animal Health.

Contributors

Hoechst Animal Health.
Hoechst UK Ltd.

Publication/Creation

Milton Keynes : Hoechst Animal Health, 1992.

Persistent URL

<https://wellcomecollection.org/works/xy8p9q7v>

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

Panacur.

Panacur is recommended by vets, and has been used by pet breeders for many years.

It controls the common roundworms of dogs and cats – including *Toxocara*.

It is the only wormer licensed for *Toxocara* control in pregnant bitches. When given as directed it reduces the transfer of *Toxocara* from the bitch to her unborn pups by 98%.

You can give Panacur in food: it comes in odourless, tasteless granules. Or you can give small quantities by mouth using a suspension.

Moreover, there is no need to fast your dog or cat before each treatment.



1 gram sachets

Pre-measured for cats, kittens, small dogs and puppies. Packs of 5.

3 gram sachets

Pre-measured for dogs and puppies. Packs of 5.



10% Suspension

Pre-measured for kittens and pregnant bitches. A 1ml pump dispenser is available for easy dosing. 100ml bottles.



Hoechst 

Hoechst Animal Health, A Division of Hoechst UK Limited,
Walton Manor, Walton, Milton Keynes, Bucks, MK7 7AJ.

Active ingredient*: fenbendazole.

Contra-indications*: nil.

Further information available on request. *See packs for full details.

PANACUR® IS A TRADE MARK



Protect your pet from worms.



Hoechst 

Protecting your dog or cat against worms is as much a part of pet care as a good diet and the right exercise.

Worms are important because they affect the condition of your pet, and can be a health risk for you and your family.

This leaflet will help you be aware of the major worm problems. It will also explain how to control them.

So you can be sure to keep your pet healthy and happy.



Worm problems.

Two kinds of worm can affect dogs and cats: roundworms and tapeworms.

Roundworms

There are several kinds of roundworm – including Hookworms, Whipworms and Lungworms. But the most important, and most common, is called *Toxocara*.

This worm causes most problems for young puppies. Indeed, virtually all puppies are born infected with *Toxocara canis*. They can also become infected from their mother's milk or the environment.

A heavily infected puppy may be pot-bellied, have occasional diarrhoea, or pass worms in vomit or faeces.

The health risk

T. canis eggs can survive for many years on grass or in the soil.

Young children can become infected if they accidentally swallow developed *T. canis* eggs.

And they may develop eye problems; total blindness in one eye has sometimes been attributed to this worm.

Toxocara cati, the cat roundworm, does not infect kittens before birth, but it is not known if it poses a threat to human health.

Tapeworms

Dipylidium caninum is the most common tapeworm (but there are others too). It can be picked up through your pet eating fleas or lice.

Signs of tapeworm infection include a slight loss of condition, and egg segments that resemble rice grains under the animal's tail or in its faeces.

Prevention.

This is, of course, the best 'cure'. And regular treatment with an effective worm control product like Panacur from Hoechst, is the answer.

How to treat your pet.

Puppies

Give Panacur Wormer granules (or Panacur 10% Suspension – 1ml by mouth or in feed for every kg of bodyweight) at 2, 4, 6 and 12 weeks of age. Then treat 3-monthly up to 1 year of age.

Dogs

Treat with Panacur Wormer granules at 3-6 monthly intervals.

Pregnant bitches

Use Panacur 10% Suspension or Wormer granules daily from about day 40 of pregnancy until 2 days after whelping. Consult your

veterinary surgeon for the correct dose.


Kittens

Treat with Panacur 10% Suspension (1ml by mouth or in feed for every kg of bodyweight) at 2-3 weeks of age and fortnightly up to 8 weeks. Then give Panacur Wormer granules at 3, 6 and 9 months, and 6-9 monthly thereafter.

Adult cats

Treat with Panacur Wormer granules at 3-6 monthly intervals. Additional treatments may be necessary if there is a tapeworm problem.

Free Poster Offer!



This attractive colour poster can be yours absolutely free with your first purchase of Panacur Wormer. Simply send the pack top in an envelope (no stamp required) to: **Hoechst Pet Care, FREEPOST, PO Box 216, AYLESBURY, Bucks HP20 1BR.** (Offer limited to one per customer).

Panacur is available from your veterinary surgeon.