A baby boy with a scarred and swollen face with a warning in spanish about the risk of HIV mothers transmitting the AIDS disease to babies; a poster from the America responds to Aids advertising campaign. Lithograph.

Contributors

United States. Department of Health and Human Services.

Publication/Creation

[Washington (D.C.)]: Department of Health and Human Services, [between 1990 and 1999]

Persistent URL

https://wellcomecollection.org/works/qt4bv6r9

License and attribution

You have permission to make copies of this work under a Creative Commons, Attribution, Non-commercial license.

Non-commercial use includes private study, academic research, teaching, and other activities that are not primarily intended for, or directed towards, commercial advantage or private monetary compensation. See the Legal Code for further information.

Image source should be attributed as specified in the full catalogue record. If no source is given the image should be attributed to Wellcome Collection.



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org

El SIDA y los bebés



na mujer infectada puede transmitir el virus del SIDA a su bebé antes de que nazca, o durante el parto. Si una mujer está infectada, su hijo tiene aproximadamente un 50% de probabilidad de nacer con el virus.

Usted debería recibir consejo y hacerse una prueba de sangre antes de quedar embarazada.



Prepared by the U.S. Department of Health and Human Services, through the U.S. Public Health Service and the Centern for Disease Congol Para más información, llame gratis al 1-800-344-SIDA. (1-800-344-7432)