

## **Salt and the kidneys / CASH.**

### **Contributors**

Consensus Action on Salt & Health.

### **Publication/Creation**

[London] : CASH, 2010.

### **Persistent URL**

<https://wellcomecollection.org/works/u56qvsmt>

### **License and attribution**

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection  
183 Euston Road  
London NW1 2BE UK  
T +44 (0)20 7611 8722  
E [library@wellcomecollection.org](mailto:library@wellcomecollection.org)  
<https://wellcomecollection.org>

# Salt and the Kidneys

## 1. Kidney Disease

Your body removes unwanted fluid by filtering your blood through your kidney via osmosis, to draw excess water out of your blood. This requires a balance of sodium and potassium to pull the water across the wall from the bloodstream into a collecting channel in the kidney. A high salt diet will alter this sodium balance, causing the kidneys to have reduced function and remove less water resulting in higher blood pressure. This puts strain on the kidney and can lead to kidney disease (*Blood Pressure Association, 2008*).

A high salt intake has been shown to increase the amount of protein in the urine, which is a major risk factor for the decline of kidney function and there is increasing evidence that a high salt intake may also increase deterioration of kidney disease. The water retention that occurs with a higher salt diet increases blood pressure, which further increases the risk of kidney disease.

Over 3 million people in the UK are at risk of Chronic Kidney Disease. 3% of the NHS budget is spent treating kidney failures (National Kidney Federation 2009) and it is believed that 37,800 adults in the UK are receiving renal replacement therapy. 1,800 kidney replacements are carried out each year, although 6,909 are on the waiting list (Kidney Research UK, 2008). 1 to 4 in a 1,000 people are believed to be effected by chronic kidney disease in the UK (NHS, 2007).

### Who is at risk?

South Asians and people of African origin are 3-5 times more likely to suffer kidney failure (requiring dialysis) compared to white Caucasians. South Asian patients with diabetes are 10 times more likely to go on to have kidney failure (*Kidney Research UK, 2008*). High blood pressure also puts the kidney under excess stress leading to deterioration of function (*Kidney Research UK, 2008*).

### Evidence

In animal studies, increasing salt intake has been shown to increase the amount of protein excreted in the urine and markedly increases the rate of deterioration of renal function in experimental forms of renal disease. Studies where salt intake has been reduced in animals with experimental renal disease show a slowing of the rate of progression of the renal disease.

Studies in humans have now shown that salt intake increases the amount of urinary protein or albumin (*du Calier et al, 2002, Verhave et al, 2004*) and this is a major risk factor for developing kidney disease and cardiovascular disease. Reducing salt intake from 10 to 5g/day in a double blind study was shown to reduce urine protein excretion by 19.4% (Fig 1. *Swift et al, 2005*). A further double blinded study in a larger number of individuals showed that even a more modest reduction in salt intake, i.e. from 9.7g to 6.5g/day reduced 24hr urinary albumin excretion in all three ethnic groups with mildly raised blood pressure (*He et al, 2009*).

Therefore, individuals with kidney disease should restrict their salt intake because in nearly all forms of kidney disease the kidney retains sodium and water in the body, causing further deterioration of renal function.

## Salt Reduction and Urine Protein Excretion




Figure 1. Change in urinary sodium and protein excretion with a modest reduction in salt intake in 40 hypertensive blacks. (Adapted from Swift et al, 2005)



## 2. Kidney (Renal) Stones

Renal stones are relatively common. Over a lifetime, 6% of women and 12% of men will have renal stones at least once. Although common, renal stones are painful and can cause nausea, difficulty passing urine and may progress to kidney disease if there is a blockage (Kidney Research UK).

A high salt diet has also been associated with renal stones. Urinary calcium, which is the main constituent of renal stones, is increased by a high salt diet and this increases the risk of stones forming. A number of studies have successfully shown that a reduction in salt consumption can reduce calcium excretion, and reduce reoccurrence of renal stones (Borghi *et al*, 2003). Hypercalciuria is present in 80% of renal stone patients (Cappuccio *et al*, 2000). Studies have also shown that individuals with raised blood pressure are more likely to develop renal stones (Cappuccio *et al*, 2000, Cappuccio *et al* 1999). A reduction in salt intake may therefore particularly benefit to these people as it not only lowers blood pressure but can also reduce urinary calcium excretion. A diet designed to reduce hypertension (the DASH diet) has been found to be associated with a marked decreases in kidney stone risk (Taylor *et al*, 2009).

### Who is at risk?

People with high blood pressure, persistent urinary infections and Crohns disease are at greater risk of renal stones. Also white British men between the ages of 30-60, and those with a family history of kidney stones, are at greater risk (NHS, 2007).

### Current Salt Intake & Dietary Advice

Almost everyone in the UK (and the rest of the Western world) eats too much salt. The daily recommended amount in the UK is no more than 6 grams a day, the current average salt intake is 8.6g salt a day although many people are eating more than this.

People with or considered at risk of kidney disease or renal failure should ensure that they keep their salt intake below the recommended maximum of 6g. This can be achieved by simple changes, such as consuming less processed foods and checking product labels before purchase.

For more information and advice on how to reduce your salt intake, please visit our website [www.actiononhealth.org.uk](http://www.actiononhealth.org.uk)

### References

- Borghi L, Schianchi T, Meschi T, Guerra A, Allegri F, Maggiore U, Novarini A. Comparison of two diets for the prevention of recurrent stones in idiopathic hypercalciuria. *The New England Journal of Medicine*. 2002;346(2):77-84
- Cappuccio FP, Kalaitzidis R, Dunclift S, Eastwood JB. Unravelling the links between calcium excretion, salt intake, hypertension, kidney stones and bone metabolism. *Journal of Nephrology*. 2000;13:169-177
- du Cailar G, Ribstein J, Mimran A. Dietary sodium and target organ damage in essential hypertension. *Am J Hypertens*. 2002;15:222-9.
- He FJ, Marciniak M, Visagie E, et al: Effect of modest salt reduction on blood pressure, urinary albumin, and pulse wave velocity in white, black, and Asian mild hypertensives. *Hypertension* 54: 482-488, 2009.
- Kidney Research UK. 2008. kidney health information PDF.
- Kidney research UK fact Sheet. 2009
- Kidney Research UK. Renal Stones Fact Sheet. <http://www.kidneyresearchuk.org/assets/asset281.pdf> [accessed 04/09/09]
- National Kidney Federation. 2009. <http://www.kidney.org.uk/campaigns/Transplantation/ukt-press-dec03.html> [04/09/2009]
- NHS, 2007. <http://www.nhs.uk/Conditions/Kidney-disease-chronic/Pages/Introduction.aspx?url=Pages/WhatIsIt.aspx> [accessed 14/04/2009]
- NHS, 2008. <http://www.nhs.uk/Conditions/Kidney-stones/Pages/Introduction.aspx> [accessed 14/09/09]
- Swift P A, Markandu N D, Sagnella G A, He F J, Macgregor G A. Modest Salt Reduction Reduces Blood Pressure and Urine Protein Excretion in Black Hypertensives. A Randomized Control Trial. *Hypertension*. 2005; 46:308-12
- Taylor En, Fung TT, Curhan GC. DASH-style diet associates with reduced risk for kidney stones. *J Am Soc Nephrol*. 2009. 20(10). 2253-9
- Verhave J C, Hillege H L, Burgerhof J G, Janssen W M, Gansevoort R T, Navis G J, de Zeeuw D, de Jong P E. Sodium intake affects urinary albumin excretion especially in overweight subjects. *J Intern Med*. 2004;256:324-30.

For further information contact CASH  
Email [cash@qmul.ac.uk](mailto:cash@qmul.ac.uk)  
Website [www.actiononsalt.org.uk](http://www.actiononsalt.org.uk)