

Illustrated guidelines on treating malaria at home in Kenya. Colour lithograph by Ministry of Health, ca. 2000.

Publication/Creation

Nairobi : Wizara ya Afya, [2000?]

Persistent URL

<https://wellcomecollection.org/works/jrdn5m6m>

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).

**wellcome
collection**

Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

Matumizi ya Madawa za Dukani KUTIBU MALARIA NYUMBANI

Malaria

Ni nini chanzo chake?



Mbu anayeuma watu wakati wa usiku

Ni nani yuko hatarini?



Ni jikinge vipi?



Tumia neti iliyowekwa dawa kujizuia malaria

Malaria ya kawaida

Dalili

- Homa ama mwili moto
- Kutokwa na jasho
- Kutetemeka
- Kuumwa na kichwa
- Kuumwa na mgongo ama viungo
- Kuumwa na tumbo
- Kutapika
- Kuhara

➤ Malaria ya kawaida inaweza kutibiwa salama nyumbani na dawa za AMODIAQUINE

Matibabu

Jinsi ya kutumia madawa ya AMODIAQUINE ni kama ifuatavyo:

UMRI	SIKU YA 1	SIKU YA 2	SIKU YA 3	JUMLA
Chini ya miezi 6	☐	☐	☐	$\frac{3}{4}$
Miezi 6 hadi 11	☐	☐	☐	$1\frac{1}{4}$
Mwaka 1 hadi 3	☐	☐	☐	2
Miaka 4 hadi 5	☐	☐	☐	$2\frac{1}{2}$
Miaka 6 hadi 12	☐	☐	☐	$3\frac{3}{4}$
Miaka 13 hadi 15	☐	☐	☐	$6\frac{1}{4}$
Zaidi ya miaka 15	☐	☐	☐	$7\frac{1}{2}$

Dalili za Malaria Hatari



Kufitika au kuzirai

Viganja vya mikono na ngozi ndani ya kope kuwa vyeupe



Kulegea...kulala-lala


Shida ya kupumua

➤ Kila wakati malaria hatari inaonekana, lazima mtoto apelekwe kwa daktari **MARA MOJA!** Usijaribu kumtibu nyumbani

Mifano ya dawa za aina ya AMODIAQUINE ni: **Amobin • Camoquin • Diaquin • Malaratab • Uniquin**



Wizara ya Afya-GFATM



Shirika la kutadhi vito chidi ya malaria ulimwenguni