

'Barbidex' : 'ionexten' brand oral depot tablets (resin-bonded dexamphetamine plus resin-bonded amylobarbitone) : one morning dose for day-long relief of anxiety & depression, for day-long appetite control / Clinical Products Ltd.

Contributors

Clinical Products, Ltd.

Publication/Creation

Richmond, Surrey : Clinical Products, [1957?]

Persistent URL

<https://wellcomecollection.org/works/e5f4pug2>

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

PHYSICIAN'S SAMPLE

'barbidex'

'IONEXTEN' BRAND

oral depot tablets

*(resin-bonded dexamphetamine plus
resin-bonded amylobarbitone)*

ONE MORNING DOSE

FOR DAY-LONG RELIEF OF
ANXIETY & DEPRESSION

FOR DAY-LONG
APPETITE CONTROL

**an 'ionexten' tablet
works 'round-the-clock'**

ORAL DEPOT THERAPY

barbidex 'IONEXTEN' TABLETS (RESIN-BONDED DEXAMPHETAMINE AND AMYLOBARBITONE)

In **barbidex 'ionexten'** tablets the prolonged effect does NOT depend on enteric coating; it is inherent in the drug resinate. Controlled ionic exchange feeds dexamphetamine and amylobarbitone into the blood-stream evenly and without interruption at an optimum level for 10-12 hours, independent of varying physiological factors.

ADVANTAGES :

1. Even, uninterrupted effect lasting 10-12 hours with ONE MORNING DOSE.
2. Optimal blood levels (never dangerously high — never ineffectively low).
3. No forgetting of conventional t.d.s. doses.
4. Economy of drug, time and money.

INDICATIONS :

barbidex 'ionexten' tablets are indicated in obesity, anxiety and depression, especially in cases where dexamphetamine alone causes over-stimulation.

DOSAGE :

Half to one tablet at breakfast. Tablets are quarter scored for dosage adjustment.

CONTRA-INDICATIONS :

Sensitivity to sympathomimetic drugs or barbiturates, marked hypertension.

FORMULA :

10 mg. dexamphetamin. sulph. B.P.C. (as resinate).
65 mg. amylobarbiton. B.P.C. (as resinate).

PRESENTATION :

Pocket pack of 30 tablets, each sealed in polythene-coated film.

POISON P.I. S1. S4

AN ORIGINAL



'IONEXTEN' BRAND

oral depot tablets are made only by

CLINICAL PRODUCTS LTD
RICHMOND, SURREY, ENGLAND



Rx for

DIRECTIONS

.....

.....

PRINTED IN ENGLAND