

Copy of a printed x-ray diffraction exposure image referenced as "X-ray fibre photograph of oriented Lupein thymonucleate (Astbury)"

Contributors

Wilkins, Maurice Hugh Frederick, 1916-2004

Publication/Creation

August 1951

Persistent URL

<https://wellcomecollection.org/works/xan4qxr8>

License and attribution


You have permission to make copies of this work under a Creative Commons, Attribution, Non-commercial license.

Non-commercial use includes private study, academic research, teaching, and other activities that are not primarily intended for, or directed towards, commercial advantage or private monetary compensation. See the Legal Code for further information.

Image source should be attributed as specified in the full catalogue record. If no source is given the image should be attributed to Wellcome Collection.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>



X-ray "fibre photograph" of oriented clupein thymonucleate. (Direction of stretching — the fibre axis, — vertical. Distance from specimen to photographic film = 4 cm. Cu K α rays).

transition from Fig. 2 to Fig. 3 suggests. It is known that the clupein molecule is an open peptide chain of up to 28 residues, of which as many as 20 are arginine residues (39), so this interpretation fits in perfectly with interaction between

a vis
stirri
to the
Whe
moun
graph
did n
thym
natur
is ne
inary
being
produ
with
globu
molec
react
under
the th
thus
above
gation
simpl
nucle
Oth
their
studie
The