

Table referenced as "H3-Thymidine in seminal vesicle"

Contributors

Pelc, Stephen Richard

Publication/Creation

February 1962

Persistent URL

<https://wellcomecollection.org/works/a4dvsbag>

License and attribution

You have permission to make copies of this work under a Creative Commons, Attribution, Non-commercial license.

Non-commercial use includes private study, academic research, teaching, and other activities that are not primarily intended for, or directed towards, commercial advantage or private monetary compensation. See the Legal Code for further information.

Image source should be attributed as specified in the full catalogue record. If no source is given the image should be attributed to Wellcome Collection.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

Incorporation of H³-thymidine in the seminal vesicle

| <u>Time</u> <u>(hours)</u> | <u>% positive cells</u> | | <u>low/heavy</u> | <u>grains</u> |
|-------------------------------|-------------------------|------------|------------------|---------------|
| | <u>heavy</u> | <u>low</u> | | |
| 5½ | 1.2 | 0.7 | 0.6 | 20 |
| 24 | 1.2 | 1.9 | 1.6 | 20 |
| 31 | 0.5 | 1.4 | 2.8 | 16 |
| 40 | 0.5 | 0.4 | 0.8 | 15 |
| 48 | <u>1.8</u> | <u>5.5</u> | <u>3.1</u> | 15 |
| mean | 1.04 | 1.98 | 1.8 | |

low/heavy: the ratio of weakly to heavily labelled nuclei.

grains: average number of grains per weakly labelled nucleus.