

Copy of a printed graph referenced as "Flow birefringence DNA"

Contributors

Gratzer, W. B. (Walter Bruno), 1932-

Publication/Creation

February 1963

Persistent URL

<https://wellcomecollection.org/works/azaqvnwt>

License and attribution

You have permission to make copies of this work under a Creative Commons, Attribution, Non-commercial license.

Non-commercial use includes private study, academic research, teaching, and other activities that are not primarily intended for, or directed towards, commercial advantage or private monetary compensation. See the Legal Code for further information.

Image source should be attributed as specified in the full catalogue record. If no source is given the image should be attributed to Wellcome Collection.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>




TABLE I

Sample history	Solvent	$[\eta]$
ative (Sample H-II, salmon)	0.2 M NaCl	68
ative	47.6% (v/v) methanol, 10^{-3} M NaCl 10^{-3} M	70

methanol with 61% methanol (all solution 10^{-3} M NaCl, 10^{-3} M Tris, pH 7) to give 79.5% methanol is at least 90% complete within 5 sec. at 25°C. When DNA-95 methanol and water (same ionic composition) are mixed to give a 70% methanol solution, the absorbance reduction is more than 90% complete within 20 sec. at 25°C. procedure (such as gradual mi