

**Graph referenced as "Helix coil transition in polylysine (Applequist and Doty)"**

**Contributors**

Gratzer, W. B. (Walter Bruno), 1932-

**Publication/Creation**

February 1963

**Persistent URL**

<https://wellcomecollection.org/works/w8c597dq>

**License and attribution**

You have permission to make copies of this work under a Creative Commons, Attribution, Non-commercial license.


Non-commercial use includes private study, academic research, teaching, and other activities that are not primarily intended for, or directed towards, commercial advantage or private monetary compensation. See the Legal Code for further information.

Image source should be attributed as specified in the full catalogue record. If no source is given the image should be attributed to Wellcome Collection.



Wellcome Collection  
183 Euston Road  
London NW1 2BE UK  
T +44 (0)20 7611 8722  
E [library@wellcomecollection.org](mailto:library@wellcomecollection.org)  
<https://wellcomecollection.org>

# Helix Fo



on from hel  
olypeptide  
f the decre  
between resid  
the degree  
al and coil  
measured

$\alpha$

$= 1 - f_h$ , an  
from Fig  
the titrat  
here the h  
is portion  
s to titrat  
behave acc  
trolytes (3

$K + \log \frac{1}{1 - \alpha}$

to determi  
10.44, as  
and Eisenb  
40 per c