

**Graph referenced as "Solvent-induced helix coil transitions"**

**Contributors**

Gratzer, W. B. (Walter Bruno), 1932-

**Publication/Creation**

February 1963

**Persistent URL**

<https://wellcomecollection.org/works/bas5jyjs>

**License and attribution**


You have permission to make copies of this work under a Creative Commons, Attribution, Non-commercial license.

Non-commercial use includes private study, academic research, teaching, and other activities that are not primarily intended for, or directed towards, commercial advantage or private monetary compensation. See the Legal Code for further information.

Image source should be attributed as specified in the full catalogue record. If no source is given the image should be attributed to Wellcome Collection.



Wellcome Collection  
183 Euston Road  
London NW1 2BE UK  
T +44 (0)20 7611 8722  
E [library@wellcomecollection.org](mailto:library@wellcomecollection.org)  
<https://wellcomecollection.org>



concentration  
parallel  
r. From t  
that the n  
eity of he  
much the s  
ein and  
lysine-L-g

ry.—E  
s of L-l  
e acid  
weights  
ere prep  
a found  
ical con  
ent of 1  
d 0.15  
int increas  
at pH 4 a  
pH 12 or u  
the NaC  
is consiste  
vious obs  
er compar  
the poly-  
weaker th