

Graph referenced as "Helix-coil transition in keratin fibers in Li Cl [lithium chloride] solution (Mandelkern)"

Contributors

Gratzer, W. B. (Walter Bruno), 1932-

Publication/Creation

February 1963

Persistent URL

<https://wellcomecollection.org/works/zwdeuz6r>

License and attribution

You have permission to make copies of this work under a Creative Commons, Attribution, Non-commercial license.

Non-commercial use includes private study, academic research, teaching, and other activities that are not primarily intended for, or directed towards, commercial advantage or private monetary compensation. See the Legal Code for further information.

Image source should be attributed as specified in the full catalogue record. If no source is given the image should be attributed to Wellcome Collection.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

the fact that con-
er a very narrow
perature at which
te is well defined,
ature interval the
normal. Plots of
first-order transi-
microscopic phases.
igned the role of an
ble which under-
the transformation
ete harmony with
n other extensive
n as the enthalpy
ation of this type.
ne transformation
ature of the fiber
l the LiBr compo-
xed LiBr composi-
ous work.^{14-16,18,21}
er is observed over
of the liquid med-
rature is, however,
entration. Transi-
observed which are
icipated from con-
ished reports.^{14,18}
slow heating rates

