

Microscope image referenced as "DNA replication"

Contributors

Lee, Dr. S.

Publication/Creation

February 1965

Persistent URL

<https://wellcomecollection.org/works/gmzsrn8z>

License and attribution

You have permission to make copies of this work under a Creative Commons, Attribution, Non-commercial license.

Non-commercial use includes private study, academic research, teaching, and other activities that are not primarily intended for, or directed towards, commercial advantage or private monetary compensation. See the Legal Code for further information.


Image source should be attributed as specified in the full catalogue record. If no source is given the image should be attributed to Wellcome Collection.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

thymidine
time will
which one
radioactive
autoradiog
porating
during a
It follow
inter dupl
strands rad
strand of
active. Be
that the
ng, but th
(that is,
exes) rem
of initial
limb of t
the grain
to carry a
e label of
nder of t
rom Cair
ission of
ers of the
Biology. C
y provide

duplex
e cross
the mi-
plication
of the
d have
while only
duplex is
lographs
icates by
s of the
daughter
at the
lication.
be seen
specially
he radio-
nd of the
by kind
and the
Molecu-
otograph
Cairns)



tion in a molecu
of transfer to the lig
therefore, replicat
'hybrid' molecules
the first generation
started to replicate
along their ^{14}N str
(^{14}N - ^{14}N) molecu
generation is over
begin to appear o
is that once a par
complete generati
again. A possible
always starts at o
scheme, light DNA
whole: generation
Fig. 104 shows.
A second interest
follows. The gener
we have seen that
If synthesis begins
some, then the rep
in the gradient, b
DNA will take abo
during the first ge
should therefore c
a whole spectrum
fact that sharply
p. 219) thus sugg