#### A practical textbook of the diseases of women / by Arthur H.N. Lewers.

#### **Contributors**

Lewers, Arthur H. N.

#### **Publication/Creation**

London: H.K. Lewis, 1897.

#### **Persistent URL**

https://wellcomecollection.org/works/n6b5g44a

#### License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.




Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org

# DISEASES OF WOMEN



LEWERS







Digitized by the Internet Archive in 2015

LEWIS'S PRACTICAL SERIES

DISEASES OF WOMEN

### LEWIS'S PRACTICAL SERIES.

In Crown 8vo Volumes, Illustrated.

- ON THE TREATMENT OF PULMONARY CONSUMPTION.
  A PRACTICAL MANUAL. By V. D. HARRIS, M.D. (Lond.),
  F.R.C.P., Physician to the City of London Hospital for Diseases
  of the Chest, and E. C. BEALE, M.A., M.B. (Cantab.), F.R.C.P.,
  Physician to the City of London Hospital for Diseases of the Chest.
  105. 6d.
- THE SURGICAL DISEASES OF CHILDREN. By D'ARCY POWER, M.A., M.B. (Oxon.), F.R.C.S. (Eng.), Surgeon to the Victoria Hospital for Children. 10s. 6d.
- DISEASES OF THE NOSE AND THROAT. By F. DE HAVILLAND HALL, M.D., F.R.C.P. (Lond.), Physician to Out-Patients at the Westminster Hospital. 10s. 6d.
- PUBLIC HEALTH LABORATORY WORK. By HENRY R. KENWOOD, M.B., D.P.H., F.C.S., Assistant Professor of Public Health, University College, etc. Second Edition, 10s. 6d.
- MEDICAL MICROSCOPY. By FRANK J. WETHERED, M.D., F.R.C.P., Medical Registrar to the Middlesex Hospital. 9s.
- MEDICAL ELECTRICITY. By H. LEWIS JONES, M.A., M.D., F.R.C.P., Medical Officer in Charge of the Electrical Department in St. Bartholomew's Hospital. Second Edition, 10s. 6d.
- HYGIENE AND PUBLIC HEALTH. By LOUIS C. PARKES, M.D., D.P.H. (Lond. Univ.), Lecturer on Public Health at St. George's Hospital Medical School. Fifth Edition, 10s. 6d.
- A PRACTICAL TEXTBOOK OF THE DISEASES OF WOMEN. By ARTHUR H. N. LEWERS, M.D. (Lond.), Obstetric Physician to the London Hospital. Fifth Edition, 10s. 6d.
- ANÆSTHETICS, THEIR USES AND ADMINISTRATION.
  By DUDLEY W. BUXTON, M.D., B.S., Administrator of
  Anæsthetics in University College Hospital. Second Edition, 5s.
- MANUAL OF OPHTHALMIC PRACTICE. By C. HIGGENS, F.R.C.S., Ophthalmic Surgeon to Guy's Hospital. 6s.
- ON FEVERS, THEIR HISTORY, ETIOLOGY, DIAGNOSIS, PROGNOSIS, AND TREATMENT. By A. COLLIE, M.D. 8s. 6d.
- HANDBOOK OF DISEASES OF THE EAR. By URBAN PRITCHARD, M.D. (Edin.), F.R.C.S. (Eng.), Professor of Aural Surgery at King's College. Third Edition, 6s.
- A PRACTICAL TREATISE ON DISEASES OF THE KIDNEYS AND URINARY DERANGEMENTS. By C. H. RALFE, M.A., M.D., F.R.C.P., Physician to the London Hospital. 10s. 6d.
- DENTAL SURGERY. By ASHLEY W. BARRETT, M.B., M.R.C.S., L.D.S., Dental Surgeon to the London Hospital. Third Edition, 3s. 6d.
- BODILY DEFORMITIES AND THEIR TREATMENT. By H. A. REEVES, F.R.C.S. (Edin.), Senior Assistant Surgeon to the London Hospital. 8s. 6d.

LONDON: H. K. LEWIS, 136 GOWER STREET, W.C.

I.H. March.

A

## PRACTICAL TEXTBOOK

OF THE

# DISEASES OF WOMEN

BY

### ARTHUR H. N. LEWERS, M.D. LOND.

OBSTETRIC PHYSICIAN TO THE LONDON HOSPITAL, AND JOINT TEACHER OF PRACTICAL OBSTETRICS IN THE LONDON HOSPITAL MEDICAL SCHOOL;
EXAMINER IN MIDWIFERY AND DISEASES OF WOMEN AT THE CONJOINT BOARD OF THE ROYAL COLLEGE OF PHYSICIANS OF LONDON AND OF THE ROYAL COLLEGE OF SURGEONS OF ENGLAND;

UNIVERSITY SCHOLAR AND GOLD MEDALLIST IN OBSTETRIC MEDICINE, LONDON UNIVERSITY

#### FIFTH EDITION

WITH 174 ILLUSTRATIONS, FOUR COLOURED PLATES, AND 71 ILLUSTRATIVE CASES

LONDON

H. K. LEWIS, 136 GOWER STREET, W.C. 1897


12552

30312363

#### LONDON

H. K. LEWIS, 136 GOWER STREET, W.C

M19701



# Sir JOHN WILLIAMS, Bart., M.D. Lond., F.R.C.P. Lond.,

TO WHOSE TEACHING

IT WOULD BE IMPOSSIBLE FOR

THE AUTHOR

TO OVER-ESTIMATE HIS OBLIGATION,
THIS BOOK IS INSCRIBED.

# PREFACE TO THE FIFTH EDITION.

written, and the operative treatment of uterine an experience in his own practice of more than

## PREFACE TO THE FIFTH EDITION.

THE author has taken the opportunity afforded by the demand for a new edition to thoroughly revise the book, and to enlarge it considerably. The section on extra-uterine gestation has been practically rewritten, and the operative treatment of uterine fibroids by hysterectomy, either with intra-peritoneal treatment of the stump, or by the so-called "panhysterectomy" in which there is no stump-the operation which in suitable cases the author himself prefers—has been fully described. A short account is given of the disease known as "deciduoma malignum," an example of which the author was fortunate enough to have in his own practice recently, in a stage allowing it to be treated by vaginal hysterectomy. The author has now had an experience in his own practice of more than sixty radical operations for cancer of the uterus. More than thirty of these have been supra-vaginal amputations of the cervix, and among these there

has been no mortality. Thirty of the operations have been vaginal hysterectomies, and among this series there have been three deaths.

Among those cases treated by the supra-vaginal amputation of the cervix, then nineteen in number, which formed the subject of his paper read before the Royal Medical and Chirurgical Society in 1892, there are still four, Nos. 6, 12, 16, and 17, which have remained well for ten years, nine years, eight years, and seven and a half years respectively. The corresponding specimens were shown when the paper was read.

Many new figures and illustrative cases have been inserted in the present edition, with the publication of which the issue of the work reaches ten thousand copies.

72 HARLEY STREET,
CAVENDISH SQUARE, LONDON, W.

November 1897.

# CONTENTS.

		-			
Synopsis of Illustrat	rive Cases .	e e			PAGE
Introduction	81 10 4				xv
	CHAPTER	T			
On the Mode of Investiga					
	CHAPTER	II.			
Menstruation and its Diso					45
	CHAPTER	III.			
Bleeding					55
	CHAPTER	IV.			
Dysmenorrhœa					76
	CHAPTER	V.			
On Diseases of the Pudeno					90
	CHAPTER				
Rupture of the Perineum					115
				1,55	3
	CHAPTER				120
Diseases of the Vagina.					132
	CHAPTER V				
Prolapse of the Vagina—P	rolapse of the	Uterus			144
	CHAPTER				
Diseases of the Uterus—In	flammation.				158
	I and				

# CHAPTER X. CHAPTER XI. CHAPTER XII. Carcinoma of the Body of the Uterus-Sarcoma of the Uterus . 273 CHAPTER XIII. CHAPTER XIV. Pelvic Inflammation (Pelvic Peritonitis, Pelvic Cellulitis) . . 322 CHAPTER XV. Pelvic Hæmatocele—Extra-Uterine Gestation . . . . 345 CHAPTER XVI. Diseases of the Ovaries . 384 CHAPTER XVII. Flexions and Versions-Inversion of the Uterus . . . CHAPTER XVIII. Disorders of Micturition . . . . . . . . . . . . . . . 472 CHAPTER XIX. 488 APPENDIX A. Note on the Systematic Treatment of Nerve Prostration and Hysteria—The Weir Mitchell Treatment . . . . 507 APPENDIX B. Notes of Seven Cases in which the Cervix was rapidly dilated by means of Hegar's Dilators . . . . . . . . . 510

# SYNOPSIS OF ILLUSTRATIVE CASES.

	motes of the blody of the filters sareona of the literal	
CASI	E	AGE
I.	Case of mucous polypus of cervix in a virgin of eighteen .	30
2.	Case of amenorrhœa for which no satisfactory cause could be assigned	50
3.	Case of hæmorrhage at the fifth month of pregnancy	57
	(The patient had had regular monthly losses up to the time she came under observation, and did not think she was pregnant.)	Pel
4.	Case of hæmorrhage after incomplete abortion	58
5.	Case of hæmorrhage at the fifth month of pregnancy, due to placenta prævia	59
6.	Case of vesicular or hydatidiform mole, which was expelled spontaneously. No trace of fœtus	63
7.	Case of vesicular mole showing no tendency to spontaneous expulsion, and therefore removed by operation. No trace of fœtus	64
8.	Case of vesicular mole in a primipara. Uterus distended to size of pregnant uterus about the eighth month. Dilatation of cervix; removal of mole. No trace of fœtus	
9.	Case of severe hæmorrhage due to laceration of hymen at first coitus	71
10.	Case of diphtheritic vulvitis in a child aged seven	93
	Case of diabetic vulvitis	94
	Case of primary syphilitic sore on the vulva. Normal induration	2000
	Case of lunus of the sulse	107
	Case of carciname of the	110
	Case of ulcorous vacinities	135
	Case of thick-walled abscess in recto-vaginal septum obstruct-	-33
	ing delivery	141

CAS		PAGE
	Case of villous or fungous endometritis	167
18,	Case of corporeal endometritis and subinvolution after a difficult labour	168
19.	Case of corporeal endometritis due to retention in utero of a piece of sponge-tent	170
20.	Case of super-involution of the uterus: justo-minor pelvis .	181
21.	Case of super-involution of the uterus	182
22.	Case of fibroid tumour complicating pregnancy	193
23.	Case of ovarian tumour of the semi-solid variety simulating uterine fibroid	
24.	Case of large fibro-cystic tumour of the uterus treated by hysterectomy	200
25.	Case of intra-uterine fibroid polypus, size of orange. Os only admitted finger. Cervix incised with Paquelin's cautery .	208
26.	Case of large interstitial uterine fibroid terminating fatally from hæmorrhage. No operative treatment had been employed. An exception to the general rule that fibroids	
27.	rarely cause death	
28.	Case of uterine fibroid (Mrs. D.) removed by combined pan-	
	hysterectomy	
	Case of pelvic abscess simulating soft uterine fibroid	227
30.	Case of carcinoma of the cervix. Small "cauliflower" growth.  Supra-vaginal amputation. Known to be free from recurrence two years afterwards.	258
31.	Case of carcinoma of the cervix. Conical ulcer. Supra- vaginal amputation. Recurrence about a year after at some distance from stump, which remained unaffected to the last	250
32.	Case of carcinoma of the cervix complicating pregnancy at	-39
	the fourth month. Large "cauliflower" growth. Supravaginal amputation	262
33.	Case of advanced carcinoma of the cervix. Large mushroom- shaped growth. Treated by scraping, cautery, and chloride	270
	of zinc	-10

SYNOPSIS OF ILLUSTRATIVE CASES.

xiii

## xiv SYNOPSIS OF ILLUSTRATIVE CASES.

	PAGE
51. Case of tubal mole with intra-peritoneal hæmorrhage from	0
the open fimbriated end of the tube. Operation. Recovery	378
52. A successful case of intra-venous injection of salt solution	
after laparotomy in a case of ruptured tubal gestation .	379
53. Case of ovarian tumour producing prolonged metrorrhagia	
after the menopause. Ovariotomy, followed by complete	
cessation of the metrorrhagia	396
54. Case of tubercular peritonitis. Diagnosis uncertain before	
operation. Laparotomy. Recovery	415
55. Case of retroversion of the gravid uterus causing retention	
of urine	474
56. Case of scrofulous kidney with cystitis, and practically incon-	
tinence of urine. Nephrectomy. Recovery	479
57. Case of congenital incontinence of urine	482
58. Case of scrofulous kidney. Sudden death from syncope.	
Pelvis kyphotic	. 484
59. Case of malignant disease affecting a movable kidney .	. 485
60. A case of double uterus with submucous fibroid tumour	492
61. Case of ill-developed uterus, imperforate hymen, and probably	7
	. 494
62. Case of hæmatokolpos due to septum in the vagina .	. 496
-5.	. 497
64. Case of double uterus with double hæmatometra and com-	-
plete absence of the vagina. Operation. Recovery .	. 497
65-71. Cases illustrating some of the conditions suitable for rapid	i
dilatation of the cervix with Hegar's dilators	. 510

# INTRODUCTION.

54. Case of inhercular peritonitis. Diagnosis uncertain before

53. Case of ovarian tumour producing prolonged metrorrhagia

IT may be safely said that no subject presents greater difficulties to the beginner than Gynæcology.

The opportunities of becoming practically acquainted with it are necessarily less than in most other branches of medicine. Take, for example, diseases of the heart: a beginner, if he listens to the heart in as many cases as possible, spends some three or four weeks before he becomes able to appreciate even well-marked murmurs.

Frequent examination of cases is just as necessary, in order to acquire accuracy in physical signs, in gynæcology, as in diseases of the heart; but whereas an almost unlimited number of observers can listen in a case of chronic heart disease, a gynæcological case can only be examined by two, or at most three, on any one occasion.

It is therefore proportionately important that the very most be made of each clinical opportunity.

Before taking a clinical clerkship in the department for diseases of women, the student should refresh his knowledge of the relations of the pelvic organs by attendance in the post-mortem room, and examining for himself the position of parts when the abdomen has been opened, and the intestines are held up out of the way.

Particular attention should be paid to the uterus, noticing its size, relation to the bladder, and testing its mobility by seeing how far it can be drawn up out of the pelvis; to the broad ligaments, and the relations of the structures projecting from them; and to the pouch of Douglas, bounded laterally by the utero-sacral ligaments, and at its lowest part in close relation with the posterior vaginal wall.

The pelvic organs should then be taken out en masse and examined more closely.

First the urethra and bladder should be laid open, the length of the urethra should be noticed, and the position of the orifices of the ureters. Probes should be passed along them. Next the vaginal portion of the cervix should be looked at, and a probe passed into the uterus. Finally the uterus should be slit up, noticing the relation

of the bladder to the anterior aspect of the cervix—the cavity of the cervix marked by the folds of mucous membrane constituting the *arbor vitæ*—and the cavity of the body. Fine bristles should be passed from the uterus along the Fallopian tubes.

The various named structures shown in Fig. 110, on page 297, should be identified.

If even one set of organs is examined in this way, the student will find it in the end a great saving of time, as his clinical progress will be more rapid and intelligent than it otherwise could be.

Another point of importance is by no means to neglect examining cases where the physician pronounces everything normal; and again, to take every opportunity of recognising physical signs when a case is being examined under an anæsthetic. Those commencing the study of gynæcology should more especially concentrate their attention on acquiring proficiency in the Bimanual examination, and in the diagnosis of pregnancy, especially in the earlier months.

I have endeavoured to arrange the divisions of

the subject, as far as possible, in what may perhaps be called their natural order from a clinical point of view, first dealing with the history, then with the physical examination, then with diseases of the external parts, diseases of the vagina, diseases of the uterus, and so on.

Numerous illustrative cases have been inserted at various parts of the book; they are, I think, a considerable help in learning the subject, and, moreover, they relieve the monotony incidental to systematic description.

# A PRACTICAL TEXTBOOK

first dealing with the history

OF THE

# DISEASES OF WOMEN.

### CHAPTER I.

ON THE MODE OF INVESTIGATING A CASE.

Before discussing systematically the more important of the diseases peculiar to women, it is desirable to consider shortly the proper way of investigating a case in this special department of practice.

By the expression "proper way" is meant the mode of procedure adopted by a practical physician in endeavouring to arrive at a diagnosis in any case that comes before him.

The materials available for diagnosis fall naturally under two heads:—

First.—We have all the patient can tell us about herself—the History.

Second.—The physical signs we can observe for ourselves—the Present State.

### THE HISTORY.

The following should be noted:—

Patient's name, address, age, occupation.

Whether married or single.

If married, how long married.

Number of children, if any, with date when the last was born.

Whether she has had any miscarriages; if any, date of each, and the period to which pregnancy had advanced when the miscarriage occurred.

Character of each confinement, easy or instrumental; if flooding, severe abdominal pain, vomiting, or fever (shivering fits, thirst, etc.) occurred after the confinement. A rough guess may be formed as to the nature of the confinement by asking how long she was compelled to remain in bed after it.

We now ask the patient:—
What do you complain of?

Her answer should be taken down as far as possible in her own words, leading questions being avoided, and after each answer merely enquiring "anything else?"

The duration of each symptom should be noted.

Some difficulty at times arises in fixing the date when the present illness began. Here we can often succeed by asking:—

How long is it since you think you were in perfectly good health?

We next enquire how menstruation has been performed since the beginning of the illness.

We ask :-

Are you quite regular now?

If the patient says "Yes," we next say :-

Every four weeks?

Because women often say they are "quite regular" when they mean they menstruate every fortnight or three weeks; in fact, by "regular" they mean they do not go more than four weeks without menstruating.

How many days does it last?

Have you any pain with it?

Enquiry is to be made whether the pain is before the flow begins, or during the flow, or after it.

If, as often happens, the pain is not sharply limited to one or other of these periods, we should find out when it is at its worst.

Also ascertain the seat of the pain, e.g., the back, hypogastrium, or one or other ovarian region.

The following questions should also be asked :-

Is the pain constant, or worse at times?

Does the pain shoot down the legs?

Is it relieved by lying down?

Particularly enquire whether the patient has always had the pain since she first began to menstruate, or whether it has only affected her for a limited time, e.g., since the birth of the last child, or since she married.

Is the discharge a good colour? i.e., red.

Do you lose much? Two women may menstruate for the same number of days each period, but one may lose much more than the other each day.

Are there any clots or shreds in the discharge?

Having learned how menstruation has been performed since the beginning of the illness, we now go on to ascertain how the function was performed previously, to see if there has been any alteration.

We ask :-

When were you first poorly in your lifetime?

Supposing the patient to say she was fifteen at the time, we ask "between fourteen and fifteen?" or "fifteen and sixteen?"

Did you come on quite regular at first, or after seeing it once did it leave you for some months?

Either alternative is equally physiological.

We then ask as before, whether she was quite regular

every four weeks when the function had become established, or how it was. The duration, quantity lost, colour, clots or shreds in it, if attended by pain or not, are noted as in the former case.

Many diseases disturb the regular performance of the menstrual function, and the object of these enquiries is to be able to compare the performance of the function since the patient has been ill, with its performance when the patient was in thoroughly good health.

There is no absolute standard applicable to all women. Individual variations within the limits of health are very common; but the way the function was performed when the patient was quite well is the standard with which to compare the way it has been performed since she has been ill.

## Short enquiries are made as to:-

### I. Micturition.

(a) Have you any trouble with your water?

(b) Are you obliged to pass it too frequently? and if so, how many times have you to get up at night to pass it?

(c) Have you any pain in passing it?

Common instances of micturition being affected by neighbouring local disease are:—

(i) Retroversion of the gravid uterus, causing retention and subsequent dribbling over of the urine.

(ii) Pelvic peritonitis, which is very often a cause of frequent desire to pass water.

## II. The digestive system.

(a) Is the appetite good?

(b) Are the bowels regular every day, or confined?

(c) Have you pain when you pass your motions?

### III. Nutrition.

Have you got thinner lately?

- IV. Any other system, e.g., the nervous system, if the patient's account of her illness suggests the desirability of doing so.
- V. Previous illnesses.

This concludes the taking of the History.

The next step is to ourselves observe all we can as to the patient's condition; the result of our observations constituting what is called:—

### THE PRESENT STATE.

First the general aspect is to be noted.—Thin or well-nourished. Healthy-coloured or pale. Whether looking obviously ill or not. Presence or absence of cedema.

Note also the state of the tongue and pulse; and if there is any indication of fever, take the temperature.

Guided by what we have learned from the history, we may, or may not, at this stage examine the chest.

The further examination is of two kinds:-

- (i) Examination of the abdomen.
- (ii) Vaginal examination.
- I. Examination of the abdomen.—The methods employed are:—Inspection. Palpation. Percussion. Auscultation.

The patient should lie on her back, all clothes fastening round the waist being loosened; it is best to have the surface of the abdomen exposed.

Inspection.—We notice:—The size of the abdomen, whether distended or not. The shape of the abdomen—localised bulging, e.g., in the flanks. The condition of the umbilicus. If the umbilicus is depressed, there is usually no very considerable tumour or accumulation of fluid within the peritoneum. Any apparent enlargement of the abdomen in such a case is simply due to accumulation of

fat in the abdominal walls, with or without distention of the intestines by flatus.

The position of the umbilicus—whether exactly in the middle line or displaced to one side, whether nearer the pubes or the xiphisternal articulation. This can of course be more accurately ascertained by measurement.

The presence of pigmentation, particularly between the umbilicus and the pubes.

The presence or absence of "skin-cracks"—lineæ albicantes—in themselves only evidence of over-distention of the abdominal walls; but the commonest cause of the over-distention is pregnancy.

On the other hand, absence of skin-cracks is no evidence that the patient has not had a child at term—it merely establishes a probability in that direction; for in exceptional cases, even when the patient has had a child at full term, skin-cracks may be absent.

The presence of enlarged veins under the skin of the abdominal wall.

Alterations during respiration.—Supposing the presence, for example, of the pregnant uterus at an advanced period of gestation, say at the seventh month; if we ask the woman to take a deep breath and to let it all out while we are looking at the abdomen in a good light, during inspiration, we see the upper border of the swelling formed by the pregnant uterus descend, and during expiration we see it ascend. The same phenomenon is seen in cases of ovarian tumour.

The appearance of the breasts.—Although not strictly forming any part of the abdominal examination, still practically it is at this stage of the examination that it is convenient to note their condition.

We observe:—their size, whether plump or flabby.

If the breast looks plump, this may be due either to

the presence of fat, or to a real hypertrophy of the gland tissue. In the latter case the breast has a nodular feeling, which is absent when the plumpness is due to fat.

Presence or absence of skin-cracks.—The nipples, well formed or not, e.g., depressed, either flat, or actually tucked in at the middle.

The primary areola, i.e., that immediately round the nipple, its size, shade of pigmentation, whether studded with little prominences (enlarged follicles) or not. Whether hairs grow on it, or round it.

The secondary areola; this is usually only to be seen in the later months of pregnancy, and for a variable time after labour. A secondary areola may occasionally be seen in brunettes altogether apart from pregnancy; when present, it occupies an area usually a finger's breadth in width (though sometimes it is much wider) around the primary areola. Its appearance is usually described by supposing a white surface to be painted brown, and then further supposing that a shower of water-drops washes away the pigment where the drops fall, exposing the underlying white. It really consists of numerous white circles about  $\frac{1}{10}$  to  $\frac{1}{8}$  of an inch in diameter, separated from one another by brown intervening spaces.

The presence of enlarged veins on the breasts, as seen in pregnancy.

The presence or absence of secretion.—This is observed by squeezing the breast in the whole hand pretty firmly in the direction of the nipple. The fluid squeezed out may be clear or milky.

'In doubtful cases the presence of a little secretion in the breasts is of small value for diagnosis. Much more reliance may be placed on the appearance of the breast. For instance, if in a woman who has had no children it looks plump, and there are veins clearly seen coursing over it, if the primary areola is deeply pigmented and studded with enlarged follicles, and if there is also a secondary areola, and the breast has the nodular feeling previously referred to, there is a strong probability that the woman is pregnant.

Palpation.—The whole area of the abdominal wall should be carefully palpated, using both hands.

Light "springy" palpation gives more information than any effort to overcome the resistance of the abdominal muscles by force.

The patient should be told to keep the mouth open; to breathe deeply and rather slowly, and to let the abdominal wall (or rather, as she understands it, "the stomach") be as loose as she can.

Great assistance will be obtained in some cases, where the patient is very nervous, by taking every trouble to reassure her, and to persuade her, that she is not going to be hurt in any way. Distracting her attention by a few questions is another help to attain the end in view, namely, as little resistance as possible from the abdominal muscles.

Light palpation frightens the patient least, and does not give rise to any discomfort; it should always be employed first. We may employ deep palpation later on, when we have learned all we can by light palpation, and when time has been given for the patient to become a little accustomed to the examination.

The kidneys.—It is good practice to try and feel the kidneys in each case. The right one is more easily felt than the left. To succeed in this the patient must not be too fat, the abdomen must be fairly relaxed, and there must be a fair amount of space between the ribs and the iliac crest. In trying to feel the lower end of the right kidney, press the fingers of the right hand

pretty steadily and firmly backwards and a little upwards in the interval between the iliac crest and the ribs. The fingers of the left hand at the same time being placed under the patient should make pressure forwards towards the fingers of the right hand.

It is best to press the fingers of the right hand steadily, and to make tilting movements forwards with the fingers of the left hand.

In a great many cases, far more than any one who has not tried it would imagine, the lower end, or more, of the right kidney can in this way be recognised.

If it is not felt at first, we may succeed by getting the patient to take a deep breath.

The lower end of the left kidney is not so often to be made out.

Other points to be noted at this stage of the examination are:—

- (a) Whether or not the recti have been separated by overdistention of the abdomen.—If the patient be asked to raise herself into a half-sitting posture, the edges of the recti can be felt with an interval of varying width between them, if they have been separated; at the same time, a pouching out of the tissues between the separated recti can be seen in such cases. Sometimes, on the other hand, the tissues between the separated recti are tucked in by atmospheric pressure as she raises herself.
- (b) Whether any part is tender to light or deep palpation.— A little care is necessary to avoid being misled. If the patient says it hurts her at one particular place, try several others; and when the attention has been distracted from the part first pressed, press it again; or press it while taking off her attention by talking to her.
- (c) Whether any abnormal swelling can be felt.—If such is recognised, endeavour to make out if it is hard or soft; if

it is elastic, or even distinctly fluctuating; if its surface is smooth or irregular; its exact situation; central or more to one side than the other. Also

The part from which it seems to spring.—For instance, if it comes from the pelvis, we shall usually \* be unable to get the fingers completely under it in that direction; though able to do so in every other; and if it springs from some part of the abdomen, e.g., from the kidney, we shall be able to separate it from the pelvis. Of course we are now speaking of fairly simple cases; of these, the above holds true. In complicated cases, e.g., large tumours coexisting with ascites, diagnosis may be difficult, or impossible.

- (d) Sometimes a crackling feeling is communicated to the fingers on palpating the abdomen, rather like what is felt when air has got into the subcutaneous connective tissue—surgical emphysema. The sensation referred to is felt in the abdomen in some cases of peritonitis, being due to the friction between the opposed surfaces covered with lymph; † the rubbing together of these being caused by the palpation.
- (e) Hardening of a tumour may be felt while it is being manipulated; the tumour in that case is almost certain to be the pregnant uterus.
- (f) Movements may be felt against the palpating hand, due to movements of the fœtus in utero.
- (g) Supposing the tumour to be the pregnant uterus at term, it is usually very easy to get the hard fœtal head
- \* Sometimes a small ovarian tumour with a long thin pedicle may be very completely separated from the pelvis when palpating in this way.
- † In some cases, however, where this peculiar sensation is experienced there is no peritonitis. I remember it being well marked in a case of ovarian tumour; yet, at the operation soon after, no evidence of peritonitis was found.

between the fingers of the two hands in the hypogastric region.

There is another sign of great value in the diagnosis of pregnancy. If the fingers make sudden light pressure at various parts of the pregnant uterus—for example, at the fifth month—they displace a layer of fluid, and often come on some hard part of the fœtus, which recedes as it is touched. The sensation so produced, when one has become familiar with it, is almost pathognomonic of pregnancy.

(h) When a hard tumour is present in the abdomen and there is also ascites, if we make sudden pressure inwards with the tips of the fingers on the hard tumour, a peculiar sensation is experienced, due to displacement of the fluid lying between the abdominal wall and the surface of the tumour.

The above are some of the common points to be attended to in palpation.

Percussion.—The whole of the abdominal surface should be percussed. Particular attention should be given to the flanks, and to any area beneath which any mass has been felt.

- (a) If dulness be found in one flank, the patient should be turned on the opposite side, and after a few seconds the previously dull area should be percussed again. If the dulness first noticed was due to fluid free in the abdominal cavity, when the patient is turned on the opposite side it will be replaced by resonance.
- (b) Dulness over the central region—umbilical and hypogastric regions—shading off laterally into resonance so that the flanks are resonant, points to some tumour centrally situated, such as an ovarian tumour—the pregnant uterus—a large fibroid tumour of the uterus—a distended bladder, and the like.
  - (c) A tumour in the lumbar region, the tumour being

distinct and either feeling solid or elastic, over which a distinct line of resonance (colon) is made out, is probably renal: such as hydronephrosis, or malignant disease of the kidney.

(d) Dulness is also obtained over the irregular lumps rising out of the pelvis, produced by pelvic peritonitis, or pelvic cellulitis, or a large pelvic hæmatocele.

Auscultation.—The positive information obtained by auscultation is practically limited to two cases.

- 1. Pregnancy, uterine or extra-uterine.
- 2. Large fibroid tumours of the uterus.

### Sounds heard over the pregnant uterus.

a watch heard through a pillow. Two sounds correspond to each beat of the fœtal heart; but often, in the earlier months, only the first sound is audible. Every opportunity should be taken of becoming thoroughly familiar with these sounds. When they are heard, we have conclusive evidence of pregnancy; our not being able to hear them by no means excludes pregnancy; nor should it even lead us to infer the death of the fœtus, unless we fail to hear the sounds after several observations at different times. The sounds become clearly audible in the last fortnight of the fifth month—from the 18th to the 20th week; practised observers may sometimes detect them a month earlier, but the date named is the time they usually become appreciable to ordinary observers.

Their frequency is about 140 or 150 to the minute; they have been said to be faster in the female than in the male, but this is unreliable in particular cases.

2. The uterine souffle.—This sound is synchronous with the mother's pulse; it is a blowing murmur produced in the enlarged arteries of the uterus. It is not peculiar to pregnancy, but is also occasionally to be heard over large fibroid tumours, and still more rarely over some ovarian tumours. It is distinguished from a mere pressure murmur by the fact that its intensity varies in a sort of rhythmical manner, gradually attaining a maximum, and then diminishing so that the sound becomes nearly inaudible.

3. The funic or umbilical souffle.—This is produced by pressure on the umbilical cord, when this is accidentally pressed between the stethoscope and some hard part of the fœtus. It is synchronous with the fœtal heart, not with the mother's.

It is only to be heard in a small proportion of cases; and is therefore of comparatively little importance.

This concludes the examination of the abdomen.

We next ask the patient to turn on her left side, to push the hips well towards the edge of the couch, and draw up the knees on the abdomen—in fact, bending herself nearly double. It is an advantage to place the patient's left arm and hand behind her. This allows her to lie more on her face and chest than would otherwise be the case.

We can then go on with the special local examination.

- II. The vaginal examination.—The first step is the inspection and examination of the external parts.—To do this properly, the fingers of each hand are placed at the sides of the vulva and the labia separated. We should notice:—
- I. The state of the hymen.—Its shape. Is it perfect (i.e., untorn)? or torn? or is it only represented by the fleshy eminences, about the size of split peas, in the situation where the hymen ought to be? These fleshy eminences—the carunculæ myrtiformes—being what remains of the hymen after the damage done to it by the passage of a child's head at or near term, or of any other body of similar size, e.g., a fibroid polypus, through the orifice. It is important to distinguish between the carunculæ myrtiformes and a merely torn hymen.

The carunculæ myrtiformes are separated from one another by intervals in which no trace of hymen remains, intervals of  $\frac{1}{8}$  inch or more wide (Fig. 1, B).

In the case of a simply torn hymen the fragments are only separated from one another by linear tears, no actual interval of any measurable width existing between the fragments (Fig. 1, A).

The hymen may be torn by coitus, vaginal examination, use of a vaginal pipe of a Higginson's syringe, and, it is said, also by merely stretching the legs.




FIG. I.—To illustrate the difference between a merely torn hymen, A, and the carunculæ myrtiformes, B. In A the hymen has been torn in pieces, which are only separated from one another by linear intervals. In B the carunculæ myrtiformes are seen, which represent the hymen after the passage of a large body, the size of the fœtal head at or near term, through the orifice of the vagina. They form projections, which are separated from one another by wide intervals. a, The orifice of the urethra.

Carunculæ myrtiformes when seen are evidence either that the woman has had a child at or near full term, or that a large body (e.g., a fibroid polypus) the size of a child's head has passed the vulva, or that the orifice has been stretched to a degree that would have allowed a body of that size to pass through it. Hence the importance of being able to identify them, and distinguish them from a simply torn hymen. From the latter condition we learn that the woman has not had a child at or near full term.

It should be remembered that an intact hymen does not exclude the possibility of pregnancy, for in some cases the tissue of which the hymen consists is elastic, so that it stretches instead of tearing; and, besides, pregnancy may occur without penetration. I saw a typical case of this some years ago at the London Hospital, and another, precisely similar, recently, sent up to me by Dr. Randall of Forest Gate. The note taken of the former was as follows:—

A young single woman was brought to see me on account of enlargement of the abdomen, and some other symptoms. On examining the abdomen, it was easily ascertained that the patient was six months pregnant, the fœtal heart being clearly heard. On inspecting the vulva, the hymen was found perfect, and the orifice bounded by it small. My impression was that a digital examination would tear it, and this proved to be the case. Examination with one finger in the usual way, and with no unusual force, caused a deep tear of the hymen posteriorly, which bled freely. A little perchloride of iron and glycerine (1-4) was applied to the laceration, and quickly arrested the bleeding.

Here, then, was a case where the patient had become pregnant without having allowed penetration.

2. The colour of the mucous membrane of the vulva should be noted.—It may be redder than normal, angry-looking, and secreting pus; such is the case in vulvitis, the local affection par excellence met with in young children.

The mucous membrane may be pale; this is its normal condition in old women; frequently in these cases there are patches of dark red on the general pale surface.

- 3. The orifice of the urethra.—If of normal size, and free from abnormal growths—warty growths—vascular caruncles, and the like. If pus is seen in the orifice, pressure should be made with the finger in the vagina along the course of the urethra downwards, to see if any more pus can be pressed out of the meatus.
- 4. The presence of any abnormal swelling or growth, e.g., a swelling as large as a walnut or larger in the posterior half

of a labium majus, feeling elastic, is often met with, this being a cyst formed from the occlusion of the orifice of the duct of Bartholin's gland on that side, and a gradual dilatation of the duct in consequence till it attains the size and position mentioned.

5. Swellings or tumours may appear in the vulva and be seen on inspection, that have arisen elsewhere.—Such are prolapse of the vaginal walls, prolapse of the uterus, tumours of the vagina, and uterine polypi protruding out of the vagina

and appearing externally.

6. If nothing is visible beyond the ordinary norma parts, it is well to ask the patient to bear down or strain. Then if there is any prolapse, or tendency to prolapse, of the vaginal walls, the anterior, or the posterior, or both, will be seen to descend, and perhaps project some distance from the vagina. Perhaps even the vagina may be completely inverted, and form a lump as big as a cocoa-nut outside the vulva, the surface of the lump showing the external os uteri at one point, and the body of the uterus being clearly felt in the lump posteriorly; this condition is called procidentia uteri.

Sometimes on asking the patient to strain, if there are certain abnormal conditions of the urethra or bladder, a

little urine will escape.

Having learned all we can by examination of the vulva,

we now pass on to the

Vaginal examination proper.—The fore-finger of the right hand is oiled or lubricated (a convenient preparation for this purpose is glycerine containing I part of corrosive sublimate dissolved in 1000 parts of glycerine), and passed along the perineum forwards over the anterior edge of the perineum, and so into the vagina. A small point, but one worth attending to, is to take care that the fingernail is trimmed short, and does not project over the soft

end of the finger; and it is also of importance to pass the finger into the vagina in the manner described, so that the more sensitive parts in front of the vaginal orifice may not be touched. Any unnecessary pain, apart from any consideration for the patient, renders the examination more difficult, and less likely to give any useful information.

Points for special attention are:—1. On attempting to pass the finger into the vagina, it may be, that so much pain is caused, that the patient extends her legs, at the same time pressing the buttocks close together, so that the examination has to be given up: in such a position nothing whatever can be made out.

This is likely to occur when there is acute, or even sub-acute, inflammation of the mucous membrane of the vulva or vagina—vulvitis or vaginitis: in cases also of urethral caruncle and labial abscess. Apart from such causes, it is likely to be met with in nulliparous women of emotional temperament. In such cases the patient must be completely anæsthetised before a satisfactory examination can be made.

2. The condition of the vagina.—Whether it embraces the finger closely, or is lax and capacious.

Its length: the canal is notably shortened in old age.

Presence or absence of the vaginal rugæ.

Tenderness on pressure in any particular direction, e.g., pressure on the anterior vaginal wall causes pain in acute and subacute cystitis, while pressure in any other direction causes no pain in such cases.

Presence of foreign bodies, e.g., forgotten pessaries, pieces of sponge, etc.

Abnormal growths.—For instance, tumours growing from the vaginal walls; or from the uterus—fibroid polypi—or malignant growths—occupying the vagina.

The temperature of the vagina.

Having noted the state of the vagina, we pass on to examine the uterus.

Attention is first to be directed to the vaginal portion of the cervix—that part of the cervix projecting into the upper part of the vagina.

We attend to:-

- (i) The direction in which the external os uteri is pointing, forwards or backwards. Normally it looks backwards and downwards.
- (ii) Whether the cervix is in the centre of the pelvis, or to one side.
- (iii) The length of the vaginal cervix—normally about one-third of an inch—its shape, long and markedly conical for instance, or short and cylindrical. Sometimes the vaginal portion of the cervix is hypertrophied, and may even extend to the vaginal orifice. This latter condition is a congenital peculiarity.
- (iv) The shape of the external os and its size.—Whether, for instance, it is a little round, or transverse, aperture some three-sixteenths of an inch across, as in nulliparæ, the edges of it free from irregularities; or a wide aperture, half an inch or more across, with irregular edges admitting the tip of the finger, as in women who have had children. These irregularities in the outline of the external os are due to the slight lacerations that occur in every first labour. In old women the os uteri is often on a level with the roof of the vagina, owing to atrophy of the cervix; the cervix does not then project into the vagina at all.
- (v) The surface of the vaginal portion should be felt all over by the examining finger, starting from the external os; thus we may detect the slightly granular feeling of an erosion round the os, its peripheral limit being a slightly raised margin beyond which the surface is felt to be perfectly smooth.

Little shotty lumps may be sometimes felt on the surface of the vaginal portion, or when the os is patulous often some way within the cervical canal.

Such little prominences may be due either to early malignant disease, or to occlusion of the apertures of gland-ducts; these then becoming distended by their retained secretion.

In this last case, if the prominence be punctured with a needle through a speculum, the lump will disappear. Retention cysts of this kind are called ovula Nabothi.

(vi) We also note whether the cervix has been lacerated deeply on one or both sides. When there is a deep laceration, say on the left side, it will be found that the finger passes from the cervical canal (thus laterally laid open) to the vagina on the same side without anything intervening between the two; whereas on the opposite (right) side, the finger starting from the cervical canal has first to traverse the right half of the vaginal portion of the cervix before coming to the vaginal wall.

The bimanual examination.—This can be made equally well whether the patient lie on the left side, or on the back. It is merely a question of practice. The position on the side is far less unpleasant to the patient, and

should therefore be generally adopted.

To perform it, the left hand is laid on the hypogastric region, and is pressed down into the pelvis as low as the state of the abdominal walls will allow. The hand should be so placed that the ulnar border from the first lies more deeply than the radial. Every endeavour should be made, as previously directed, to get the patient to relax the abdominal muscles. The left hand thus pressed down is kept quite passive while the finger of the right hand in the vagina, placed on the external os, tilts the cervix upwards; this of course imparts at the same time an upward impulse to the body and fundus of the uterus, so that the palmar

aspect of the left hand pressed down in the way described receives an impulse, and so becomes informed of the whereabouts of the fundus uteri.

This being known, if the uterus is in its normal position, its body can be grasped between the fingers of the outside hand, and the finger in the vagina, the latter now being carried in front of the vaginal portion of the cervix (Fig. 2).

If the bladder is full, even apart from any abnormal condition, the body of the uterus cannot be felt bimanually.




FIG. 2.—THE BIMANUAL EXAMINATION (Schroeder).

The body of the uterus is seen grasped between the fingers of the left hand pressing down in the hypogastrium, and the fore-finger of the right hand in the vagina. Note that the bladder is empty.

If we are unable to feel the body of the uterus, a catheter should always be passed. After emptying the bladder, we are often able to feel the uterus bimanually in cases where we had previously failed to do so.

Even if the bladder is empty, we cannot, as a rule, feel the body of the uterus bimanually if the uterus be retroflexed, or strongly retroverted. If the abdominal walls are thin and very lax, we may, however, feel the body of the uterus bimanually, even when retroflexed or retroverted.

The mobility of the uterus will have been estimated when the finger in the vagina made the upward tilting movement on the vaginal cervix.

The normal range of mobility is about  $1\frac{1}{2}$  inches in the upward direction; the uterus may be absolutely immovable, or less movable than normal, or freely movable.

At the same time, a fair estimate may be made of the weight of the uterus, whether heavier than normal, or not.

It is of great importance, during the bimanual examination, to be able to form an estimate of the size of the uterus (particularly with a view to the diagnosis of early pregnancy), also to ascertain whether its surface is smooth, or the seat of irregular projections (as in the case of sub-peritoneal fibroid tumours).

Supposing the body of the uterus cannot be felt bimanually, even when the bladder is empty, the internal finger should explore the region behind the cervix. Normally, when the rectum is empty, no lump should be felt in this situation. If we feel a lump there, it may be one of many things, e.g., it may be the body of a retroflexed uterus, it may be a hæmatocele, or a distended Fallopian tube, or an enlarged and prolapsed ovary. The passing of the sound will settle whether the lump is the body of the uterus or not.

The finger having felt in front of the vaginal portion and behind it—explored the regions called the anterior and posterior vaginal *fornices*—now goes on to examine the spaces at the sides of the cervix—the lateral fornices; and both without and with the aid of the bimanual method, note is taken of any masses to be felt in these regions. For instance, in parametritis (pelvic cellulitis) of one side, a lump is felt on that side of the cervix.

For the purposes of description it is usual to divide

the pelvis into quarters, and to speak of a lump being felt in the right, or left, posterior quarter of the pelvis, as the case may be. The boundaries, for instance, of the right posterior quarter of the pelvis are:—the right broad ligament, the right utero-sacral ligament, and externally, the pelvic wall.

On withdrawing the finger from the vagina, it should be noticed whether there is any blood on it, or not. If there was no blood in the vagina to begin with, the occurrence of bleeding after gentle examination is suggestive of malignant disease.

To complete the examination we may use the speculum, and, if specially indicated, the uterine sound.




Fig. 3.—Fergusson's Speculum.

The two forms of speculum in common use are Fergusson's tubular speculum, and Sims's duckbill speculum.

Fergusson's is the one which will be found most generally useful.

Sims's speculum, on the other hand, is essential for operations on the vagina and cervix. It has the advantage that we can touch the cervix while the speculum is in position.

Other specula that are occasionally useful are Barnes's crescent speculum, and Cusco's bivalve speculum.

Directions for passing Fergusson's speculum.—Care should be taken to choose a size suitable to the capacity of the vagina, as already determined by the finger.

The fore-finger of the left hand is passed a short way into

the vagina, and retracts the perineum; at the same time the middle finger of the left hand draws the right labium to the right. The speculum is held in the right hand, and its end, previously oiled (not dipped in oil), is insinuated into the orifice of the vagina. Special care should be taken not to tuck in any of the parts, e.g., the nymphæ, or hairs. It is




FIG. 4.—SIMS'S DUCKBILL SPECULUM.

then pushed on, upwards, but with a strong inclination backwards, till it is completely passed.

Then, looking up the tube, we see if the vaginal portion of the cervix is in view as it should be. If not, the speculum is turned round, or passed up a little further.



Fig. 5.—Retractor used with Sims's Speculum for pressing forwards the anterior vaginal wall, while the speculum presses back the posterior vaginal wall and perineum. An assistant at the same time should draw the patient's right buttock upwards.

If the cervix is still not in view, the speculum should be withdrawn a little way, and again passed up.

The speculum enables us to see:—I. The condition of the vaginal mucous membrane—redder than normal, and if so whether the redness is uniformly distributed, or confined to certain parts of the vagina, e.g., the lower part or the upper part of the vagina, or, as in other cases, to the summits of the ridges of the mucous membrane, the intervals between the ridges being pale—or again, whether the whole mucous membrane is spotted over with red spots, the intervals between the spots being pale, as in "spotty" vaginitis.




FIG. 6.—BARNES'S CRESCENT SPECULUM.

One blade is first introduced, and then the other, so that, when in position, the blades lie as in the figure.

- 2. Presence of ulcerations in the mucous membrane.—These may be syphilitic, or produced, as is often seen, by the use of hard pessaries. Sometimes we cannot assign any cause for them.
- 3. The presence and character of various discharges in the vagina.—Merely, as is natural, a little whitish fluid just sufficient to moisten the surface; or a copious white, or yellow, or greenish-yellow fluid, or blood.

4. The secretion in the external os.—Normally clear, transparent, and viscid, like unboiled white of egg. In catarrh of the cervix, it loses its transparency and becomes opaque,



Fig. 7.—Barnes's Crescent Speculum.

The blades packed for carrying, one within the other.




FIG. 8.—Cusco's BIVALVE SPECULUM.

and either white, or yellow, according to the intensity of the inflammation. Often a red, raw-looking, slightly granular, or even villous area is seen of varying width immediately

around the external os. Such areas constitute the various forms of erosion of the cervix, and in the more marked forms are probably indications of coexisting inflammation of the cervical mucous membrane.

5. The actual size and shape of the external os.

The speculum is also useful for the application of various medicated fluids to the surfaces within reach.

It will often be found that what appears to be a wide erosion, when the speculum is pressed well up, becomes much diminished in width if the speculum is withdrawn a




FIG. 9.—THE UTERINE SOUND (Simpson's).

The side of the handle corresponding to the concavity of the sound is roughened; the other side of the handle should be smooth. There is an angular projection  $2\frac{1}{2}$  inches from the point of the sound, this being what is called "the normal distance" for the sound to pass into the uterus.

little way; the former appearance being due in part to an eversion of the mucous membrane of the cervix.

The examination may with advantage end here in many cases. If the art of making the bimanual examination has been thoroughly mastered, the cases where the use of the sound will add much to our knowledge are comparatively few.

The sound should not be used where there is the least likelihood of pregnancy, or in cases of pelvic inflammation (pelvic peritonitis and cellulitis), or in cases of malignant disease, or during a menstrual period. Nor should it be used when there is vaginitis, nor, in general terms, when there is any obviously offensive or morbid matter in the vagina.

There are three safeguards against passing the sound into a pregnant uterus.




Fig. 10.—To illustrate the mode of passing the sound described in detail in the text (Hart and Barbour).

First.—It is always well to ask when the patient menstruated last.

Second.—The examination of the abdomen will prevent the risk of passing the sound in advanced pregnancy. Third.—The bimanual examination, by enabling us to estimate the size of the uterus, will put us on our guard as to early pregnancy.

There are many patterns of uterine sound. Simpson's is, on the whole, the one to be preferred. It is certainly as good as any, and better than most.

Directions for passing the uterine sound.—The fore-finger of the right hand being on the external os, the sound, warmed and dipped in some antiseptic lubricant, is passed with its concavity looking backwards along the palmar aspect of the finger, keeping the point of the sound especially in close contact with the finger, till it reaches the external os. It is then gently slipped into the os; it will find its way easily as far as the internal os.

Here a slight check is usually met with. If the previous examination has shown the body of the uterus to be in its normal position of slight anteflexion, the direction in which the concavity of the sound looks will have to be changed, so that it may look forwards; then steady gentle pressure will cause it to pass into the body of the uterus.

If, however, the uterus be retroverted, or retroflexed, there will be no necessity for altering the direction in which the concavity of the sound looks. It will only be necessary to carry the handle of the sound somewhat forwards towards the pubes as it passes into the uterus, its concavity looking backwards as at first.


It is very important to thoroughly understand the manœuvre for changing the direction in which the concavity of the sound looks.

The student should take a sound and follow the directions to be given, looking at Fig. 11.

Place the sound flat on the table in the position a, b, c. Now hold the points a and b closely against the table, so that the part of the sound a b cannot be displaced laterally.

Now take hold of the handle c with the other hand.

Raise it off the table, and bring it down again to the position c'. It will be found that the handle c passes through a semicircle to the position c', without causing any displacement of the part ab, which merely rotates on its own axis. This altering of the direction in which the concavity of the sound looks, without displacement of the terminal part ab, is called the "tour de maître."



When we consider that there is often septic matter in the vagina, it is obvious that if we pass the sound in the manner described in the last paragraph, there is considerable danger of carrying some of it into the cavity of the uterus with the sound. If this is done, pelvic inflammation will probably be set up, and may even end fatally. This is one reason for restricting the use of the sound to cases where it is absolutely necessary for either diagnosis or treatment—a relatively small number.

In every case where the sound is about to be used, it

should be made scrupulously clean, and dipped into an efficient antiseptic solution, such as 1-1000 corrosive sublimate lotion.

Rectal examination.—Sometimes it is an advantage to examine with the finger in the rectum. In this way we notice:—

- 1. Whether the rectum itself is diseased, e.g., stricture; or, as in a case I remember, impaction of a large gall-stone in the rectum.
- 2. It is ascertained for certain whether masses felt to the left and behind are fæcal, or due to some pathological condition of the pelvic organs, e.g., distinction between fæces and an inflamed left ovary.
- 3. In young girls and unmarried women, where some examination is called for, examination should be made per rectum first; the condition of the pelvic organs can thus be ascertained without rupturing the hymen.

For example, an unmarried girl, about 18, was brought to me on account of a more or less constant red discharge from the vagina. On examination per rectum, I felt a mucous polypus the size of an almond hanging from the os uteri. As the only effectual treatment would be removal of the polypus, a vaginal examination was then made, confirming the result already obtained. Of course if nothing abnormal had been found on rectal examination, no vaginal examination would have been made.

The catheter.—The frequent necessity of passing the catheter in the course of a physical examination of the pelvic organs has been already alluded to.

Indeed, provided that he has a catheter to ensure the bladder being empty, the experienced gynæcologist can as a rule obtain all the information he requires by the digital and bimanual methods of examination alone. To put it another way, the catheter is a more indispensable instru-

ment than the speculum, and a far more indispensable one than the uterine sound.

Moreover, after many operations, e.g., ovariotomy, operations for ruptured perineum, supra-vaginal amputation of the cervix, etc., the catheter has to be passed two or three times in the twenty-four hours.

It is therefore important to remember that cystitis is very apt to be set up by its use, unless special care is taken.

Precautions to be observed in using the catheter.—1. If there is any discharge about the external parts, the orifice of the urethra must be wiped with a pledget of cotton-wool soaked in 1-1000 perchloride lotion to avoid carrying some of the discharge into the bladder.

#### KROHNE & CO LONDON

#### FIG. 12.

- 2. The kind of catheter used is an important point. It is impossible to feel sure that the ordinary catheter with the eye at the side, and a space beyond the eye, is clean, for it may be taken as certain that dirt will accumulate in the part beyond the eye. It is best to employ a male celluloid catheter having an eye at the side, but the part of the catheter beyond the eye solid (Fig. 12).
- 3. Before use the catheter should be dipped in corrosive sublimate lotion (1-1000), then in a solution of perchloride of mercury in glycerine (1-1000); after use it should be immediately washed by passing a stream of water through, if possible by holding it under a tap. Then it should be dipped once more in sublimate lotion, and again washed with water; it may then be allowed to dry. Celluloid catheters will stand boiling for a period not exceeding five minutes. This is the best way to sterilise them after use.

On dilatation of the cervix.—There are two methods of dilating the cervix:—1. The rapid method by means of Hegar's dilators. 2. The gradual (slow) method by means of tents.

The choice between them depends on the kind of case in which dilatation is to be employed.

The rapid method is especially preferable where dilatation is required at no long time after a confinement or miscarriage; and it is also far the best method when dilatation of the cervix is undertaken during pregnancy, for the purpose of removing the ovum, either on account of its degeneration into a mole, or on account of some other pathological condition.

In the class of cases referred to the cervix is soft, and dilates readily to the extent necessary to allow the finger to pass the internal os uteri. Apart from pregnancy, the cervix is often similarly soft and dilatable in some cases of uterine fibroids.

On the other hand, when the cervix is not soft, rapid dilatation, to the extent needed to allow the finger to pass into the uterus, is almost certain to be attended by more or less laceration of the cervix; in fact, in such cases rapid dilatation means laceration. As a general rule, therefore, when dilatation is required for exploration, if the patient is not pregnant, or if she has never been pregnant, or if some long period has elapsed since the last pregnancy, or in any case where the cervix feels rigid, the rapid method should not be employed. In such cases the best plan is to use one or more specially prepared laminaria tents. For example, say these are passed at five o'clock on one afternoon; then the next afternoon at two o'clock we can remove the tents, and perhaps find sufficient dilatation has been produced to allow the finger to pass; but if this is not quite the case, the cervix has been softened and rendered easily

dilatable by the action of the tents, and we can readily complete the dilatation with Hegar's dilators without any risk of laceration. The laminaria tents I am in the habit




FIG. 13.—HEGAR'S DILATORS.

of using are kept in a 1 per cent. solution of corrosive sublimate in absolute alcohol; they are thus rendered both aseptic and antiseptic. Just before inserting them they are dipped in a 1-1000 solution of corrosive sublimate in glycerine,

Dilatation by means of tents, unless the most careful antiseptic precautions are observed, is not uncommonly followed by more or less severe attacks of pelvic inflam-




FIG. 14.—THE VOLSELLA.

The instrument figured has a catch at the handles like Spencer Wells's forceps.

mation, peri- or parametritis, and several deaths from their use have been recorded. There is especially great risk in inserting a second tent or series of tents when sufficient dilatation has not been produced by those inserted at first,

I. Rapid dilatation of the cervix.—Hegar's dilators are the instruments most commonly employed for rapid dilatation. They are slightly curved, ebonite cylinders, about three inches and a half long, the distal end forming a blunt cone, and the proximal being fitted with a handle. There are twenty-six sizes usually supplied, the transverse diameter varying from one-twelfth of an inch to one inch (see Fig. 13). The method of using them is as follows:—The patient is put in the lithotomy position, and the vagina thoroughly syringed with some efficient antiseptic, such as 1-1000 corrosive sublimate solution. Sims's speculum is introduced, and the anterior lip of the cervix seized with a volsella, preferably one having a catch at the handles, like Spencer Wells's forceps (Fig. 14). The direction of the uterine cavity is then ascertained with an ordinary sound. The dilators should have been previously placed, in numerical order, in a shallow porcelain tray, and covered with 1-1000 perchloride lotion. One of the dilators corresponding to the supposed calibre of the cervical canal is then dipped in sublimate glycerine (1-1000), and passed through the cervix, which is held steady by the volsella in the left hand. If the dilator used only passes with difficulty, it is held in position a minute or two before withdrawing it. It is important to have the next larger size ready to pass at once after withdrawing each dilator. In this way the dilators are passed, one after the other, till the cervix is sufficiently open to admit the finger. This degree of dilatation is usually obtained after No. 19 of the series has passed. If any morbid condition is discovered in the cavity of the body of the uterus, this is treated by suitable means. Polypi, for instance, or pieces of placenta can be removed.\* If a

<sup>\*</sup> I once removed a piece of sponge from the cavity of the body of the uterus. A sponge tent had been used some weeks before by some one else; afterwards the patient had a profuse, purulent, and somewhat

growth of doubtful character is discovered, a small portion may be removed for microscopical examination. Whether anything abnormal has been found in the interior of the uterus or not, it is of great importance to apply pure tincture of iodine to the endometrium, and afterwards wash out the uterus with hot iodine water (3ij. Tr. iodi to the pint of water). For washing out the uterus nothing answers better than the double-channelled tube shown in Fig. 15. No matter how tightly the tube may be grasped by the uterus,




FIG. 15.—TUBE FOR WASHING OUT THE UTERUS.

The india-rubber tube of an ordinary Higginson's syringe fits on A. The horse-shoe shape of the tube on transverse section at any part (say B) is shown. The fluid injected at A passes between the walls of the tube as at C. This fluid, after escaping from the holes at the distal end, returns along the groove D, which runs on the under surface of the tube for its whole length.

the fluid injected readily escapes along the deep groove on the under surface. These tubes are made of celluloid, or glass. Those I use in gynæcological work are made of glass. For washing out the uterus after delivery I also use a glass tube of the same pattern. The tubes made of celluloid cannot be effectually cleaned. The objection to the

offensive discharge. I dilated the cervix with Hegar's dilators, and found a flattened piece of sponge, about \(^3\) of an inch long by \(^1\) of an inch wide, lying loose in the uterine cavity.

glass tube that it is easily broken can be met by having a plain wooden box made to carry the tube when not in use. Subsequently the patient should have antiseptic vaginal




FIG. 16.—UTERINE TENTS.

a. Sponge. b. Laminaria. c. Tupelo.

douches three times a day for a few days; iodine water (3ij. Tr. iodi—Oj. water) is the antiseptic I usually employ for this purpose. If the dilatation is performed soon after a

miscarriage, or soon after labour, no anæsthetic is needed, but in other cases it is necessary.




FIG. 17.—UTERINE FORCEPS.

2. Dilatation of the cervix by tents.—Tents are made of sponge, laminaria (sea-tangle), or tupelo, a kind of wood.

They possess the property of absorbing moisture, and

expanding as they do so, exercise a dilating force on any part into which they have previously been tightly fitted.




FIG. 18.—SPONGE-HOLDER.

If dilatation by tents has been decided on, it is, to begin with, essential that the patient should remain in bed, not only during the dilatation, but for some days afterwards. A

vaginal douche of corrosive sublimate solution 1-1000 is given. The cervix is now exposed with Sims's speculum, and, if necessary, steadied with a tenaculum. The tent, previously anointed with sublimate glycerine 1-1000, is held




FIG. 19.—EXPANSION OF A TUPELO TENT (Mundé).

The larger figure shows the tent after expansion. The constriction indicates where it has been in contact with the internal os uteri.

with a pair of uterine forceps (Fig. 17), or better with a long sponge-holder having a catch at the handle (Fig. 18), and passed into the cervix so that its highest point may be well

beyond the internal os. If there is room for more than one tent, another is inserted by the side of the first, and so on. It is best to insert a plug of iodoform gauze into the vagina to keep the tents in place.

Tents are left in about twelve hours. When tents specially prepared, as mentioned above (p. 33), are used, and every other antiseptic precaution adopted, they may be left in twenty to twenty-four hours without any ill effect. When the gauze is removed, an antiseptic douche should be given before removing the tents. After removal of the tent, or tents, the finger is at once passed into the cavity of the body of the uterus; the recognition and treatment of any morbid condition discovered is conducted as described on p. 35 under rapid dilatation, not omitting the final washing out of the uterus through the intra-uterine tube (Fig. 15) with an antiseptic lotion.

Formerly, when the necessity for strict asepsis was not understood, dilatation often caused great pain. This is practically unknown nowadays, when tents are used as described above. At most the patient may experience a little discomfort.

Another way of inserting a tent is to fit it on a special introducer—the tent is perforated nearly to its upper end to allow of its being carried on the introducer (Fig. 20). This method is rarely adopted now.

Sterilisation of instruments and appliances.—Instruments should, as far as possible, be made of metal or glass, so that they may be either boiled for a quarter of an hour, or subjected to dry heat at 150° C. in a hot-air bath for an hour. Boiling is the more generally convenient plan. The water should contain 1 per cent. of soda. For use in hospital, gas is generally used to heat the steriliser (Fig. 21); but for operations in private a portable steriliser is required, in which the water is boiled by means of a spirit lamp




FIG. 20.—BARNES'S TENT INTRODUCER.

A tent is seen fitted on the end ready for introduction. When the tent has been placed in position, the stylet on which it is mounted is withdrawn through the larger tube, which is held steady till the stylet is quite free from the tent.

with a large flame. The instruments are taken direct out of the steriliser, and placed in a porcelain tray containing cold carbolic lotion. Of course they must not be touched unless the hands have been sterilised as completely as possible. A satisfactory plan for rendering the hands aseptic is the following:—The nails must be kept short, and never cleaned out with a knife or other pointed instrument, which only causes an appearance of cleanliness. The nails, hands,




FIG. 21.—PORTABLE STERILISER.

and forearms are to be thoroughly scrubbed with soap and hot water and a clean nail-brush. An excess of tincture of iodine should be put into the water used to wash the hands. A little is of course decolorised by the soap, but it is easy to have an excess of free iodine. At least five minutes should be spent on scrubbing the hands and nails. The hands are then rinsed in plain water to get rid of all the soap, and then are to be thoroughly scrubbed in perchloride of mercury lotion 1-1000. Another nail-brush, never used for soap, is employed for scrubbing in the perchloride.

The hands should be scrubbed in the perchloride lotion for at least two minutes. After taking the hands out of the perchloride lotion they are ready for work, and must not be dried, or allowed to touch anything that is not aseptic.

Hegar's dilators cannot be boiled, but after they have been thoroughly scrubbed with soap and water they are rinsed in plain water to get rid of the soap, and then placed in a tray containing sublimate lotion 1-1000.

All instruments must be thoroughly scrubbed with soap and water after use before being sterilised.

Silk and fishing gut should be boiled for an hour, and, if it is intended to keep them in stock, they should be placed in absolute alcohol containing perchloride of mercury 1-1000. They must be placed in carbolic lotion 1-20 for some hours before use to get rid of the alcohol.

Catgut cannot be boiled, as water spoils it, but it may be sterilised by dry heat at 150°. I do not use catgut in abdominal operations. For ordinary vaginal work, such as operations for ruptured perineum, etc., I find a practically satisfactory condition of asepsis is obtained by keeping the catgut, after it has been freed from fat with ether, in a solution of sublimate in absolute alcohol 1-1000, and using it as required direct from the alcoholic solution.

Sponges cannot be boiled, but they may be sterilised by dry heat as above described without injury, if they are dried first.

I prefer using artificial sponges made of Gamgee tissue. These can be boiled. New ones can be made each time, as the material is so cheap. For abdominal work, pieces of Gamgee tissue are cut out of suitable size, and the layers of netting are sewn together round the margin to prevent pieces of the wool getting loose. I have used such artificial sponges for some time, and find them very satisfactory.

### CHAPTER II.

## MENSTRUATION AND ITS DISORDERS.

Normal type.—In this country menstruation usually first appears between the ages of fifteen and sixteen. It is not very rare for it to appear as early as ten, or to be delayed as late as twenty. In hot countries it appears a little earlier, and in cold countries a little later, than in temperate climates.

Menstruation generally ceases between forty-five and fifty.

When menstruation has become thoroughly established, it recurs every four weeks, *i.e.*, from the *beginning* of one period to the *beginning* of the next is four weeks; sometimes it occurs every three weeks without there being any abnormality.

Both at the time of the establishment of the function, and for some time before its cessation, some irregularity as to periodicity is commonly observed. For instance, after the first menstruation the girl often "sees nothing" for some months before it reappears, and there may be a similar interval before it recurs a third time; finally she becomes "regular" every four weeks. This is altogether within physiological limits.

Each period lasts from three days to a week. The quantity of blood lost each time is about three ounces. Normally there are no clots passed. If the discharge is only moderate in amount, clotting is prevented by the mixing of the acid vaginal secretion with the blood. If the quantity lost is much greater than normal, or if the

blood is retained in the cavity of the uterus for some time, clots are formed.

The blood discharged during menstruation comes from the body of the uterus, the mucous membrane of which undergoes fatty degeneration, and becomes disintegrated, with rupture of its capillaries, and consequent escape of blood.

After menstruation is over, a new mucous membrane begins to be developed, starting from the internal os, and gradually the whole of the cavity of the body of the uterus is once more lined by a mucous membrane. This attains its greatest thickness just before the next menstrual period; if pregnancy does not occur, it, in its turn, undergoes the degeneration and disintegration already described.

Ovulation, *i.e.*, the maturation of a Graafian follicle and discharge of its ovum into the Fallopian tube, is, as a general rule, associated with menstruation. It is at present uncertain whether the follicle ruptures before, during, or after the menstrual flow.

Exceptionally:—1. Ovulation may occur without menstruation, as when pregnancy occurs in women who have never menstruated, or who become pregnant during suckling.

2. Menstruation may occur without the maturation and rupture of a Graafian follicle, i.e., ovulation, for in some cases where death has occurred during, or soon after, a menstrual period, no Graafian follicle, either about to rupture, or recently ruptured, has been found.

At the time of menstruation there is congestion of the ovaries and uterus, and there is increased secretion from the glands of the cervical canal. This increased secretion is the only part taken by the cervix in menstruation. The breasts may swell, and become painful.

General phenomena observed in connection with menstrua-

tion.—(a) There is increased vascular tension for some days before the period, which falls during, and after it.

(b) The temperature is a little raised (about half a degree) for some days before the flow, similarly falling again during, and after it.

(c) Some feeling of fulness in the pelvis, perhaps slight backache, and some sense of fatigue, are usually met with, even in healthy subjects.

# ABNORMALITIES OF MENSTRUATION.

Menstruation may never appear, or having appeared, may become suppressed. If it has never appeared, we call the condition *Primary Amenorrhæa*; if it has appeared and subsequently become suppressed, we call it *Secondary Amenorrhæa*.

Again, menstruation may be attended with pain, and we have the condition called *Dysmenorrhæa*.

Lastly, either the quantity of blood lost at each period may be excessive, or menstruation may recur at too short intervals; in either case we have the condition termed *Menorrhagia*.

Amenorrhœa.—It has already been mentioned that menstruation begins in most cases between fifteen and sixteen, but that it may be delayed as late as twenty without there being necessarily anything wrong. At the same time it is to be remembered that the longer its first appearance is delayed beyond the usual time, the greater is the probability of there being some constitutional or local disease present.

# APPARENT AMENORRHICA.

Menstruation may seem to be absent where, in reality, the function is being regularly performed every month.

This is due to some physical obstruction preventing the escape of the blood. Such cases are rare. The obstruction is either due to an imperforate hymen, a congenital or acquired occlusion of the vagina, or occlusion of the external os uteri.

Acquired occlusion of the vagina occurs occasionally after severe specific fevers, such as scarlet fever; the vaginal walls slough, and the granulating surfaces left cohere, so as to close the canal. This may also occur after difficult labour. The possibility that the amenorrhoea may be due to one of these causes should always be thought of when a girl, who has arrived at the age of puberty, but who has never menstruated, experiences pain in the hypogastrium recurring every month. After a time gradual dilatation of the cavities above the obstruction occurs, and a tumour is generally noticed rising out of the pelvis. It is also found that the swelling so formed increases in size when the pains in the hypogastrium occur each month, and diminishes again after the pains subside.

The increase in size corresponds to the menstrual period, when more blood is added to that already in the dilated uterus, or dilated vagina, as the case may be; after the period is over, some of the blood is reabsorbed, and hence the diminution noticed in the intervals between the periods.

The *treatment* proper for such cases will be referred to in the chapter on Diseases of the Pudendum.

# AMENORRHŒA PROPER.

1. In most cases true amenorrhæa is to be looked on as an indication, and as a result, of a bad state of the general health.

—It is in this way that amenorrhæa results from anæmia,

and particularly the peculiar form of anæmia called *chlorosis*; from *overwork*, especially *overwork indoors* (as in the case of domestic servants, shop-girls, etc.); that it follows severe illnesses, and various organic diseases, especially *phthisis*.

In all these instances failure of the general health is the cause of the amenorrhœa.

2. Sometimes, on the other hand, deficient ovarian activity, with which is associated amenorrhæa, seems to be the cause of the failure of the general health, and resulting anæmia or chlorosis.—Here the fault lies with the ovaries, which perform their functions sluggishly. The close connection between the functional activity of the ovaries and menstruation, is seen from the fact that girls whose ovaries have been removed before the age of puberty never menstruate at all; and that women from whom both ovaries are removed, as a rule cease to menstruate.

If the changes in the whole body, which occur at the age of puberty, are to be accomplished without injury to the general health, the demand made on the strength by the rapid development in progress must be met by a corresponding activity in the processes of nutrition.

It is generally agreed that the proper action of the ovaries gives an important impetus to nutrition at this juncture; if this impetus is not supplied, owing to sluggishness of the ovaries, nutrition is not sufficiently active, the strength is unequal to the demands made on it, the health fails, and chlorosis supervenes.

Sometimes when there is deficient ovarian activity and also amenorrhoea, there may be a condition of plethora, and this may after a time be replaced by chlorosis.

3. There is an irregular group of cases where we cannot be certain what the cause of the amenorrhæa is.

Thus amenorrhoea sometimes follows:—A change in the mode of living, particularly change of air from the country

to London, and vice versa; or some disappointment in love.

Occasionally it may be impossible to assign any satisfactory reason for amenorrhoea, as in the following case:—

On May 19th I attended a lady in her third confinement, which was in every way perfectly normal. She suckled the baby till August 1st, and then weaned it. She did not "see anything" while nursing, but she was surprised, and somewhat alarmed, to find that menstruation did not reappear after the child was weaned. Her general health was quite good, except for some feeling of blood to the head and flushing. September, October, and November passed by without menstruation having occurred, and she came to see me on December 3rd with a view to ascertain whether she was pregnant. On examination I found that the uterus was not enlarged. On December 17th I heard from her that menstruation had at last reappeared. The only possible explanation of the amenorrhoea was that she was staying in the country most of the time; but, on the other hand, this appears to have been her usual custom during that part of the year. Menstruation in this case was rather late in starting; it began between sixteen and seventeen, but she was not really regular till she was twenty-one.

- 4. Primary amenorrhæa may be due to congenital defects of the generative organs. Such as the following:—
- (a) There may be no uterus, or only a rudimentary uterus, the ovaries being normal.
- (b) The ovaries may be absent or rudimentary, the uterus being normal.
- (c) The infantile condition of uterus and ovaries may persist throughout life.
- •(d) The ovary and other appendages of one side may be absent.
- 5. Local morbid conditions causing, or associated with, amenorrhaa. Such are:—
- (a) Ovarian tumours; sometimes, however, there is menorrhagia in such cases.
  - (b) Pelvic inflammation (Perimetritis, Parametritis), espe-

cially when ending in abscess. The amenorrhoea is partly, no doubt, due to the depression of the general health. Sometimes, however, there is menorrhagia and metrorrhagia in cases of pelvic inflammation.

- (c) Exposure to cold during menstruation is very apt to lead to pelvic peritonitis and amenorrhoea; cold may stop the flow, even if it does not set up pelvic inflammation.
  - (d) Ovaritis.
  - (e) Superinvolution of the uterus.
- 6. Lastly, amenorrhæa is physiological during pregnancy, and while suckling. When a healthy-looking girl complains of amenorrhæa, having been quite regular up to the time of the last menstruation, there is considerable probability that the cause of the amenorrhæa may be a physiological one.

Treatment.—A. In cases of sudden suppression by exposure to cold during, or just before, menstruation, the patient should have a hot hip bath with mustard in it, and be put to bed, lying between the blankets. A glass of gin, or whisky in hot water, will be useful as a diaphoretic.

If there is much pain, a morphia suppository (gr. ½) may be ordered. Sudden suppression of menstruation by exposure to cold is very likely to result in an attack of pelvic peritonitis, and the patient should be kept in bed for some days as a precaution, even if no symptoms of peritonitis arise.

When the time comes round at which the next menstruation should appear, the patient should have a hot mustard and water hip bath for three or four nights running before the day on which the period is due; if this does not bring it on, three or four leeches may be applied round the anus, or to the inside of the thighs.

In between the dates when the periods are due, attention

should be directed to the general health; and in most cases a mixture containing small doses of saline purgatives will be useful, such as the following:—

B. Mag. Sulph., 9j.
Sod. Bicarb., gr. xv.
Sp. Chlor., mv.
Infus. Gent. Co. ad 3j.

twice or three times a day, according to the effect produced on the bowels.

B. In cases of chronic amenorrhæa associated with anæmia, whether the anæmia has preceded and brought about the amenorrhæa, or whether we have reason to think that ovarian inactivity by failing to stimulate nutrition has led to the anæmia, the main thing is to improve the general health.

Out-door exercise, cold baths, regular hours, and a sufficiency of food are of great value; and I have found Valentine's meat juice useful in these cases.

As regards medicine, if the tongue is furred, some such prescription as that given above is suitable to begin with; and as there is nearly always obstinate constipation, it should include a sufficient dose (for example, a drachm) of sulphate of magnesia, to make the bowels act regularly.

If, however, the tongue is clean, we may at once prescribe iron, and it is best to try a mild preparation at first, such as the ammonio-citrate. If this is well borne, we may soon change it for the astringent preparations of iron, which are by far the best for anæmia as soon as the stomach can bear them.

Such are the sulphate of iron, and the liquor ferri perchloridi.

The sulphate is best given in the form of pill, combined with a small quantity of watery extract of aloes.

R. Ferri Sulph. Exsicc., gr. ij.
 Ext. Aloes Aq., gr. 4.
 M. ft. pil. j., thrice daily after meals.

Or, if we wish to give the liquor ferri perchlor., a suitable combination is:—

R. Liq. Ferri Perchlor., mxx.
Tr. Nuc. Vom., mv.
Mag. Sulph., 9j.
Sp. Chlor., mv.
Aq. ad 3j.
Thrice daily after food.

Occasionally, where iron does not effect our purpose, small doses of arsenic are useful, such as one or two drops of the liquor arsenicalis, three times a day, after food. Levico water, which contains arsenic, is also useful in the same way. The prepared black oxide of manganese, in twenty-grain doses, is also highly spoken of. The cases, however, where iron fails are not many, especially after preliminary treatment with saline purgatives.

C. Where amenorrhæa is only a minor symptom in the course of grave organic disease, such as phthisis, it of course requires no special treatment.

D. In cases where there is no constitutional condition discoverable to account for the amenorrhoa, and where no local morbid condition, other than that the uterus is smaller than it should be, can be detected, it has been recommended that we should resort to local treatment. The local treatment referred to is the dilatation of the cervical canal by a series of bougies, as hereafter described under dysmenorrhoa, or the introduction of a stem pessary into the uterus, the stem being formed partly of zinc and partly of copper—the galvanic intra-uterine stem pessary. It is easy of course to make the inner surface of the uterus

bleed by mechanical irritation, such as treatment of this kind necessarily involves. Menstruation is, however, a very complex process, of which external hæmorrhage is only one of the symptoms. Making the endometrium bleed produces one of the symptoms, but it is a long way from producing the process. Most of the cases, therefore, included in this class, if not all, had better be left entirely without local interference.

Sometimes the administration of permanganate of potash—gr. j.-ij. in pill three times a day—seems to bring about the reappearance of the catamenia.

#### CHAPTER III.

#### BLEEDING.

ONE of the symptoms most commonly met with in organic disease of the pelvic organs is bleeding—using the term to mean bleeding other than that occurring in normal menstruation.

Excessive loss of blood at the menstrual periods is known as menorrhagia. Bleeding occurring between the menstrual periods, or not distinctly associated with menstruation, is known as metrorrhagia. Clinically, however, these distinctions cannot always be made; practically what we must take account of is, whether the patient is losing too much blood—either because her periods recur too often, for example, every fortnight, or because, although the periods do not recur too often, she loses an excessive amount each time; further, these conditions may be combined—the patient menstruating too often, and losing too much blood on each occasion.

Finally, bleeding may occur in the intervals between the periods.

CLASSIFICATION OF THE COMMONER CAUSES OF BLEEDING.

## I. Bleeding in connection with pregnancy.

Abortion, or miscarriage—threatened, or inevitable.

Incomplete abortion, or miscarriage—portions of the ovum being retained, and causing bleeding.

Missed abortion, or miscarriage.

(Note.—When the fœtus dies in the earlier, or middle, periods of pregnancy, the ovum is usually expelled at no long interval after; exceptionally the ovum is retained in utero for a considerable time after the death of the fœtus, and during that time the patient is in a state of "missed abortion" or "miscarriage," according to the period to which pregnancy had advanced when the fœtus died—abortion being the term used to denote expulsion of the ovum up to the end of the third month—miscarriage, its expulsion from the end of the third month to the end of the seventh lunar month—the twenty-eighth week.)

Subinvolution of the uterus.

Placenta prævia.

Accidental hæmorrhage, i.e., hæmorrhage from the detachment of a normally situated placenta.

Post-partum hæmorrhage. Primary. Secondary.

Extra-uterine pregnancy.

Malignant disease of the cervix complicating pregnancy.

Rupture of varicose veins of the vulva, or vagina, during pregnancy.

Molar pregnancy.

Placental polypus.

Fibrinous polypus.

## II. Bleeding not distinctly connected with pregnancy or labour.—Common causes of this are:—

Malignant disease of the cervix uteri.

Fibroid tumours and fibroid polypi of the uterus.

Mucous polypi of the uterus.

The age of puberty, and of the menopause.

Ovarian tumours.

Over-lactation.

The stimulus of recent marriage causing congestion of the pelvic organs.

Pelvic inflammation, in some cases.

Slight bleeding, or a discharge sometimes blood-stained, may be due to:—

Foreign bodies in the vagina (ill-fitting pessaries, hair-pins, pieces of sponge, etc.).

Prolapse of uterus (from ulcers of the inverted vagina).

Vascular caruncle of the urethra.

Less common causes of bleeding are:-

Malignant disease of the body of the uterus.

Malignant disease of the vagina, or of the external parts.

Diseases of the liver and heart causing venous congestion.

Endometritis of the body of the uterus.

Inversion of the uterus.

The hæmorrhagic diathesis.

Laceration of the hymen at the first coitus.

### I. BLEEDING IN CONNECTION WITH PREGNANCY.

Clinically one of the most important facts to grasp is, that in spite of menstruation occurring regularly, or rather in spite of the fact that the patient has bleeding recurring about every month, which she takes for menstruation, she may nevertheless be pregnant—perhaps as much as five or six months, as in the following case:—

C. P., age 37, married eighteen years, three children, the last nine years ago, seven miscarriages, the last a year and a half ago at the end of the third month, was admitted to the London Hospital, complaining of bleeding for the last six weeks. Every day of the last six weeks she had lost something, and more the last three weeks than before. She had been regular up to the commencement of the bleeding, which began at one of her periods. On examination, a uniform, elastic swelling was found rising out of the pelvis, and reaching up to the umbilicus. A souffle could be heard to the right of the tumour, but no fœtal heartsounds. The patient did not think she was pregnant. Vaginal examination found the cervix softened, and blood escaping from the os uteri.

The bimanual examination showed that the swelling in the abdomen was connected with the uterus—in fact, felt like the pregnant uterus. The sound was passed six inches, with the intention of emptying the uterus (this was justifiable, because the patient had been losing blood for six weeks, and in considerable quantities for three weeks). Forty-eight hours later, as no progress had been made, and the patient was still losing, the membranes were ruptured with the sound. Six hours after, the patient miscarried. The fœtus was of about five months' development.

This case, not by any means an uncommon one, shows the necessity of keeping the possibility of pregnancy in mind, although the patient believes herself to have been menstruating regularly, and has no suspicion that she is pregnant.

Incomplete abortion.—The next case illustrates bleeding due to an incomplete abortion, parts of the ovum having been left behind.

E. G., age 39, married sixteen years, seven children, came to the London Hospital complaining of losing a large quantity of blood from time to time, ever since she had had a miscarriage five months previously.

She was three months pregnant when the miscarriage occurred. For some weeks before admission she had been losing blood three or four days out of every week.

Examination showed that the uterus was somewhat large; the sound passed three and a half inches. Some blood was escaping from the os uteri. Under ether, the cervix was rapidly dilated by means of Hegar's dilators (a method that has been already described, see p. 35), and the finger passed into the uterus. Several irregular prominences were felt scattered over the mucous membrane of the uterus, varying in size from that of a split pea up to that of half a cherry. They were scraped away as completely as possible, chiefly with the finger nail. Pure carbolic acid was applied to the interior of the uterus. The little masses removed looked like pieces of placenta that had retained their vitality. After their removal, the patient menstruated regularly, and not excessively.

Placenta prævia.—Bleeding due to placenta prævia usually occurs at an advanced period of pregnancy, when

the fact of pregnancy existing is unlikely to be overlooked. Exceptionally, and this is especially likely where the case is one of central placenta prævia, bleeding occurs comparatively early—as in the following case. Here the patient was unaware of the pregnancy.

Mrs. F., age 37, married thirteen years, four children, the last five years old, no miscarriages, came to me on June 23rd, 1886, with the following history. She was regular till the first week in January, when her last proper menstrual period occurred.

At the beginning of February, just when she was expecting her period, she caught a cold, and the period did not come on; she remaining in bed, on and off, for two or three weeks. Early in March she had a hæmorrhage lasting five weeks; after that she was a month free from bleeding. Then it came on again, and from that time till she came to me the bleeding continued more or less. Since February she had also suffered from attacks of pain across the lower part of the abdomen. On examination the uterus was found reaching up to the umbilicus; bleeding was still going on in small quantity. The patient was anæmic; and, having regard to the time bleeding had been in progress, I had no hesitation in advising her to have the cervix dilated. Under an anæsthetic I accordingly dilated the cervix with Hegar's dilators. As soon as I could get my finger in, I felt the placenta over the os; I detached it all round, ruptured the membranes, brought a leg down, and completed delivery in an hour and twenty minutes from the time of commencing to dilate the cervix. The uterus was washed out afterwards with iodine water, and vaginal douches of the same were used twice a day for a week after. The patient made an uninterrupted recovery.

Extra-uterine gestation.—Bleeding in connection with extra-uterine gestation is marked rather by *irregularity* than by profuseness. A common history in cases of tubal gestation (the commonest variety of extra-uterine gestation) before rupture of the gestation sac, is that the patient misses a period, or goes two or three weeks over her time, and then a hæmorrhage comes on, during which a decidual cast of the uterine cavity may be passed.

Such a history as that, especially if the patient had also had attacks of pain in one or other iliac region, would raise a presumption in favour of the case being one of extra-uterine pregnancy; and some approach to certainty in the diagnosis would be reached if the physical examination discovered a tense, elastic, more or less fixed swelling at one side of the uterus, the size of a hen's egg. It should be mentioned that some cases of extra-uterine pregnancy are attended, like ordinary pregnancy, by amenorrhoea throughout. In a case of interstitial gestation that came under my care at the hospital, the patient thought she was five months advanced in an ordinary pregnancy, and had "seen nothing" for five months. Then rupture of the sac occurred with a fatal result.

Malignant disease of the cervix complicating pregnancy.—
It is important to bear in mind the not infrequent association of these conditions. Otherwise during the early months the pregnancy is very likely to be overlooked. When a woman with malignant disease of the cervix says she has been bleeding for more than twelve months (as in a case which I remember where the patient was four months pregnant), one is apt to forget that she may be pregnant. A careful bimanual estimation of the size of the uterus as a routine method, in every case where any vaginal examination is made at all, is the best safeguard against overlooking early pregnancy, in this and other cases.

Molar pregnancy.—There are two varieties of mole:—

- 1. The vesicular or hydatidiform mole.
- 2. The blood-mole, and its later stage—the fleshy mole.
- 1. The vesicular mole.—This is produced by a cystic degeneration of the chorion villi. Among the symptoms it gives rise to is bleeding. The patient has a red discharge, perhaps with white cysts in it, varying in size from that of a grape downwards. The presence of these cysts

is of course pathognomonic, but in most cases of vesicular mole there is only a red discharge without any of the cysts in it. Another point to which attention must be paid is, that the uterus is generally larger than would correspond to the supposed month of pregnancy. I have seen, however, one well-marked exception in this respect.

Apart from the occurrence of bleeding and the presence of characteristic cysts, however, in any particular case where the size of the uterus on examination is greater than corresponds to the supposed period of pregnancy, it is much more likely that the patient is "out in her count," than that she has a vesicular mole.

Cases of hydatidiform mole may be divided into two classes: (1) Where the case first comes under observation at a time when the diagnosis has yet to be made, where we can only have a strong suspicion as to the nature of the condition present; and (2) where the case is first seen when the os uteri is well dilated, and the mole is in process of being discharged. In such cases the diagnosis is of course obvious, and the treatment, if any is needed, merely consists in facilitating the expulsion of the mass. It is in the first group of cases that difficulty arises, and to arrive at a probable diagnosis requires a careful consideration of the history and physical signs present. The history is usually somewhat as follows:-In the first instance, the patient missed one or more periods, and in consequence believed herself to be pregnant; but this interval of amenorrhœa has been followed by irregular hæmorrhagic discharges. If she is observant, she may have noticed a rapid increase in the size of her abdomen. Usually this is all the history. No doubt it may happen that the characteristic cysts may be found in the vaginal discharge, and if this is so (and we see the cysts ourselves) there is an end of the difficulty. But much more frequently

no cysts are observed in the discharge, either by the patient or by trained nurses who have been specially informed as to the importance of finding the cysts, or by ourselves when examining the patient. In other words, the only local symptom is in most cases the presence of a red vaginal discharge. The physical signs may be described as those of pregnancy, except such as depend on the actual presence of the fœtus; that is to say, there are the usual changes in the breasts, and on examining the abdomen we find a uniform, elastic tumour, centrally situated, and rising out of the pelvis, over which, as a rule, a loud uterine souffle may be heard; but nothing like the parts of a fœtus can be felt on palpating the tumour, nor can the fœtal heart be heard even by repeated observations. It is of some value if the size of the tumour is much greater than would correspond to the supposed duration of the pregnancy. Per vaginam, similarly, we find the signs of pregnancy, except those that depend on the presence of the fœtus. So that we may have blueness of the vulva and vagina, softening of the cervix, the recognition usually that the tumour observed in the abdomen is the body of the uterus; but, on the other hand, ballottement cannot be obtained, nor can anything like a part of a fœtus be felt through any of the vaginal fornices. As a rule, the speculum shows a reddish or brown discharge coming from the os uteri.

Given a case with the history and physical signs mentioned, a diagnosis of hydatidiform mole may be made with a considerable degree of confidence, and the right treatment is to empty the uterus. I have found the most satisfactory method of doing this is to dilate the cervix with Hegar's dilators till the os will admit the finger, and then to remove the mole, partly with the fingers and partly with polypus forceps. When the bulk of the mass has been taken away and the uterus has contracted down, it is well to go

over the whole of the endometrium gently with a blunt curette (such as Récamier's), so as to remove any small fragments remaining. The uterus is then washed out with hot iodine water through a double-channelled intra-uterine tube, and the operation is at an end. Antiseptic vaginal douches are given for a few days subsequently.

As regards the three cases notes of which follow, in the first the mole was discharged spontaneously; in the second the uterus showed no tendency to empty itself, although the patient was kept under observation in the hospital for nearly a month before any operative treatment was adopted. In this case the continuance of the hæmorrhage furnished the chief indication for interference. In the third case the cervix was dilated, and the uterus emptied two days after the patient was first seen. Here the indications for immediate treatment were: (1) the great size of the uterus, which was as large as the pregnant uterus between the seventh and eighth months; (2) the fact that the patient had been suffering for about three weeks from obstinate vomiting, so that she had been rapidly losing flesh. This third case is also remarkable in that the patient had had no children (and indeed had only been married seven months), whereas hydatidiform mole most frequently occurs in those who have had children.

(Note.—In a minority of cases, where the cystic change has only affected a relatively small portion of the chorion, a foetus may be present with the hydatidiform mole.)

### ILLUSTRATIVE CASES.

1. E. R., age 27, married ten years, three children, the last one year and nine months old, no miscarriages, was admitted to the London Hospital on December 7th, 1887.

History of the present illness.—On admission she said that she believed herself to be between three and four months pregnant. At

the time when, according to her calculation, she would have been two and a half months pregnant, bleeding came on, and sickness, which she had been suffering from previously, became more troublesome. She went to a doctor, who gave her some medicine, after taking which the bleeding and sickness stopped. At this time also, she felt something move inside her, falling to the lower part, and causing her much discomfort in walking.

December 8th.—On examination the uterus was found reaching to within a finger's breadth of the umbilicus; there was some brownish discharge in the vagina, but no recent blood. The cervix was apparently shortened, but was not very soft.

Menstrual history.—Before the present illness she was regular every four weeks; for the last three or four months she has not menstruated properly, but has had occasional slight bleeding, lasting sometimes an hour and a half; the bleeding in question was accompanied by pain in the abdomen.

On admission she was given Ext. Ergot. Liq. mx. and Pot. Brom. 9j. three times a day.

January 4th, 1888.—She has had slight bleeding and pain from time to time since the last note. The uterus is within an inch of the umbilicus. The last two or three days the loss of blood has increased in quantity.

On the evening of the 4th, between 6 and 7 p.m., pains like labour pains were felt; they increased in force till midnight, when a hydatidiform mole was passed. The mass was the size of the placenta at full term. No trace of a fœtus could be found. The patient made an uninterrupted recovery.

2. S. C. P., age 22, married four years, two children, the last born in January 1887, one miscarriage, three months after marriage, was admitted to the London Hospital on August 8th, 1888.

History of the present illness.—She has not been properly "poorly" for four months, but for the last six weeks she has had a red discharge; she only noticed the swelling in her abdomen three weeks ago. She has had pain in the lumbar region, and both iliac regions, worse on the right side. Three months ago she began to suffer from morning sickness. On July 30th she felt something move in the tumour, and continued to do so for four days; since then no movement has been felt. Patient suckled her second child up to March last (fourteen months).

Menstrual history.—Catamenia appeared at fourteen, and she has been regular every four weeks; no pain at the time.

Present state, August 13th.—Patient is slightly anæmic. Temperature 99°. Pulse 98, small and weak. A centrally situated swelling is felt in the abdomen, rising out of the pelvis, and reaching up to the umbilicus. The tumour is felt to harden, and again to become softer, alternately. A loud uterine souffle is heard over the tumour, but no fœtal heart sounds.

Vaginal examination.—The vaginal mucous membrane is only slightly blue, the cervix is soft, and the os not patulous, the cervix is blue, there is no blood seen about the os to-day, though on examination on August 11th some was seen.

August 23rd.—The patient has had a red discharge continuously since the date of the last note. The loss is more abundant at night.

She has felt no movement in the tumour.

No fœtal heart sounds to be heard.

The tumour (measured with the tape) reaches a height of  $6\frac{3}{4}$  inches above the pubes.

On vaginal examination no blood is found, nor is the os patulous.

September 1st.—Daily red discharge has continued, and last night a good deal of blood was lost.

September 3rd.—Discharge has continued as before. Patient is now decidedly anæmic. The tumour (measured with the tape) now rises 8 inches above the pubes. Nothing but a loud souffle to be heard over it.

On vaginal examination some blood-stained mucus is seen in the os uteri.

On account of the continuous loss of blood, which had now rendered the patient markedly anæmic, it was decided to dilate the cervix.

Accordingly chloroform was given, and I dilated the cervix with Hegar's dilators, beginning with No. 13 and continuing up to No. 28 (the first of a larger series specially made for me). As soon as the finger could be passed in, this was done; some partially decolorised clot came away, and some portion of a vesicular mole as the finger was withdrawn. It was now of course decided to completely empty the uterus. This was done by passing two fingers of the right hand into the uterus, and pressing down the uterus from the outside with the left hand. Ovum forceps and Récamier's curette were also used in getting the mole away. The whole mass in a graduated glass measured 30 ounces. The cavity of the uterus was washed out with

iodine water, and then the endometrium was swabbed with pure tr. iodi; a hypodermic injection of ergotin was also given.

The uterus contained nothing but the hydatidiform mole; no trace of a feetus could be found.

The patient did quite well, and went home on November 1st, 1888.

3. Mrs. F., a young married lady, came under my observation on December 29th, 1889. She had been married in May 1889. The catamenia were regular every month up to the middle of September last; she then "saw nothing" for about three months, and thought she was pregnant. For about three weeks before I saw her she had been suffering from obstinate vomiting every day, and also for about the same time she had had a brownish-red discharge from the vagina. She was aware of the tumour present in the abdomen, and she is very positive that a month ago it was nothing like its present size. On examination there was a tumour in the abdomen reaching several inches above the umbilicus, and about the size of the pregnant uterus between the seventh and eighth month. It was centrally situated, freely fluctuating, and seemed to rise out of the pelvis. Nothing like any hard part of a feetus was felt anywhere in the tumour. On auscultation I could hear nothing over the tumour; but as I had had half an hour's drive in an open trap, and there was a hard frost at the time, I think it quite possible that a souffle might have been heard under more favourable conditions. A catheter was passed, but only about an ounce of urine was drawn off. On vaginal examination the cervix was typically soft; it was blue, and there was some reddishbrown discharge coming from it. The anterior vaginal fornix was bulged down by a convex elastic swelling, apparently continuous with the cervix. Nothing hard like a part of a fœtus was felt anywhere in the swelling. The breasts had the appearance characteristic of pregnancy at three or four months, and contained a little secretion. I had no doubt the patient was pregnant, and that there was some pathological condition of the ovum as indicated by the persistence of a red discharge from the uterus. It seemed to me that it was either a case of hydatidiform mole, and that the whole tumour was the uterus containing it; or, that it might, perhaps, be a case of threatened miscarriage at about the third or fourth month, complicated by the presence of an ovarian tumour. I certainly thought the balance of probability in favour of vesicular mole. Under the circumstances the best course seemed to be to empty the uterus. Accordingly, on

December 31st the patient was put under the influence of the A.C.E. mixture, and I dilated the cervix with Hegar's dilators till I could pass my finger into the uterus. On withdrawing the finger a fragment of a hydatidiform mole, showing the characteristic vesicles, came away with it. The diagnosis being settled, I proceeded to empty the uterus, using my fingers and from time to time the ovum forceps. The earlier part of the operation was the more difficult, because, owing to the great size of the uterus, the ovum forceps had to be passed in deeply. There were two or three sharp attacks of hæmorrhage at this time. It was checked by injecting hot iodine water into the uterus, but it could not, of course, be completely stopped till the uterus was empty and able to contract down. The removal of the mole was therefore hastened as much as possible, the uterus ultimately contracting well, and the hæmorrhage then ceased. I carefully scraped the endometrium with Récamier's curette, and once more washed out the uterus with iodine water. The whole operation lasted about an hour. No trace of a feetus was found; the cavity of the uterus contained nothing but the mole. I have since heard from the patient's medical attendant that she made an uninterrupted recovery.

(Note.—Not very long afterwards this patient became pregnant again, and was delivered at full term of a healthy child.)

I may add, as an example of pregnancy at an advanced age, that I saw a case of hydatidiform mole, with Dr. F. M. Corner, of Poplar, in a patient who was in her fifty-third year.

2. The blood-mole and fleshy mole.—These are produced by blood finding its way between the layers of the membranes of the ovum—between the vera and reflexa, or the reflexa and chorion, or between the chorion and amnion. If the mole is expelled while the blood-clot is comparatively fresh, it is called a "blood-mole." If not discharged till the blood-clot has become tough, and partially decolorised, it is called a "fleshy mole."

A fleshy mole is usually the size and shape of an orange; on bisecting it, it is found to have a central cavity lined by the amnion; this is raised up into irregular projections by the clot underneath. The fœtus may either have disappeared,

or be recognisable as a little body an inch or less long, attached by a short stalk (the umbilical cord) to some part of the amniotic surface. The wall of the mole may be an inch think, and is made up of the layers of the membranes with blood-clot, more or less altered, between them.

It is not always that a mole in utero gives rise to hæmorrhage spread over any considerable period. I remember a case where a patient had not menstruated for seven months, and she thought she must be seven months pregnant. She had had some yellow discharge for two months. On examination the uterus was found to be of the size corresponding to the beginning of the fourth month of pregnancy. A few weeks later she passed a fleshy mole the size of an orange; its expulsion was only preceded by bleeding for a few hours, as in an ordinary abortion. Not infrequently, however, the presence of a mole gives rise to irregular hæmorrhages; and if the mole is decomposing, the discharge is offensive. A mole is rarely retained beyond the time that would have been full term if the pregnancy had been an ordinary one.

Now as to diagnosis: whereas in the vesicular mole the size of the uterus is generally greater than corresponds to the supposed period of pregnancy, in the case of the fleshy mole the size of the uterus is less, as in the case just recorded, where a woman who thought she was seven months pregnant had a uterus only the size of the fourth month. Again, examining such a case at intervals, e.g., every fortnight, for a few weeks would show that the size of the uterus was stationary.

Placental polypus.—Bleeding dating from a particular confinement may be due to some portion of the placenta having been retained, the normal involution of the uterus being thereby interfered with. Layers of fibrin are deposited on the portion of placenta retained, and ultimately

a sort of polypus is formed, which on examination is often to be felt presenting at the os uteri. The polypus is sometimes called also "fibrinous" polypus, but a fibrinous polypus may originate apart from pregnancy. I have known it develop on the slight irregular projection left in utero after the removal of a fibroid polypus with the écraseur.

# II. BLEEDING NOT DISTINCTLY CONNECTED WITH PREGNANCY OR LABOUR.

Malignant disease of the cervix uteri.—The commonest cause of bleeding in this category is malignant disease of the cervix uteri; in these cases careful enquiry will usually elicit that, besides periodical excessive losses, there is also a more or less constant discharge stained with blood between the periodical losses. The patient rarely goes many days without seeing some blood in the vaginal discharge.

Bleeding occurring some years after the menopause is, in a very large proportion of such cases, due to malignant disease.

Fibroid tumours and fibroid polypi.—These are another common cause of excessive bleeding; the interstitial and sub-mucous fibroids and fibroid polypi producing the most bleeding, and the sub-peritoneal the least, often none at all. Fibroid tumours range from the size of a pea up to masses weighing several pounds, as large as, or larger than, the pregnant uterus at term. A common size for fibroid polypi is that of a hen's egg, or an orange; they may, however, be much larger, or much smaller.

Mucous polypi grow from the mucous membrane of the cervix or body of the uterus. They are much smaller than fibroid polypi; a maximum size for a mucous polypus would be that of a large strawberry; an average one is the size, and somewhat the shape, of a small raisin.

Menorrhagia of puberty and of the menopause.—
The time when the menstrual function is being established

(puberty), and the time when the menstrual function is about to cease altogether, are both periods when excessive losses of blood are common, apart from any discoverable organic disease, local or general.

Vascular caruncle of the urethra.—This is another common cause of slight losses of blood at irregular times. Another much rarer disease of the urethra, malignant disease beginning round the orifice of the urethra, is also a cause of slight bleedings. If the patient happen also to have a yellow or white vaginal discharge, her account of it will be that there are streaks of red in the discharge from time to time.

Foreign bodies in the vagina (pessaries, pieces of sponge, hair-pins, etc.) may cause a more or less red (and often a more or less offensive) vaginal discharge. The bleeding comes in these cases from ulcerations of the vaginal mucous membrane, resulting from the pressure or irritation of the foreign body.

Excessive and irregular losses soon after marriage are not uncommon. The patient usually becomes regular spontaneously, as she becomes more accustomed to married life.

Over-lactation is another cause of excessive bleeding.

Hæmorrhage due to general venous congestion from disease of the heart, or liver.—It is only very rarely that uterine hæmorrhage can be satisfactorily traced to such causes.

Ovarian tumours.—More commonly the disturbance of menstruation caused by the development of ovarian tumours is in the direction of amenorrhoa, the intervals between the periods being lengthened, and the loss on each occasion less than formerly. Exceptionally, the growth of an ovarian tumour is attended with menorrhagia, or metrorrhagia.\*

\* A short paper by the author, "On the Occurrence of Metrorrhagia after the Menopause in Cases of Ovarian Tumour," was published in the Lancet of January 2nd, 1897.

Endometritis of the body of the uterus.—This is another cause of excessive loss of blood. The condition, however, is one not well understood, and it is not easy, except in the case of villous endometritis, to be certain of its presence in actual practice. Points to be attended to as presumably indicating endometritis of the body of the uterus are, slight increase of the size of the body ascertained on bimanual examination, tenderness of the body, and great tenderness on passing the sound. There is a yellow, or sanguineous, discharge from the os uteri, but this, of course, is also found in inflammation limited to the cervical mucous membrane, a far commoner affection.

It is not to be thought that endometritis of the body of the uterus is not a real disease; what is meant is, that in actual clinical work it is comparatively rare to be able to isolate it as the cause of the patient's symptoms. A villous condition of the mucous membrane of the body of the uterus has been termed villous endometritis; this variety is especially likely to be a cause of bleeding, and can be recognised clinically after dilating the cervix.

Laceration of the hymen at the first coitus.—This laceration, as a rule, only causes very trifling bleeding: exceptionally, it may cause severe hæmorrhage; of this I have seen a well-marked case:—

Y. S., a German Jewess, age 20, was admitted into the London Hospital on the afternoon of August 20th, 1889, on account of hæmorrhage. She had been married the previous day. As a result of coitus severe bleeding came on, and as it continued the woman was brought to the hospital. I saw her at once, as I happened to be in the hospital at the time. She was deathly pale, and there was a great deal of recent blood on her clothes. Some blood was still coming from the parts, but she was so nervous that I could not examine her without an anæsthetic. She was then put under ether, and a hot vaginal douche having been given, the parts were carefully examined. There was a deep tear of the hymen posteriorly, but bleeding had stopped;

probably the hot douche just given had stopped it. No blood was coming from the os uteri, nor was there any laceration of the vagina. As a precaution the vagina was packed with carbolic gauze, and a T bandage put on. The gauze was taken out next day, and the patient went home on the 23rd, there having been no recurrence of the bleeding.

# DIFFERENTIAL DIAGNOSIS OF THE VARIOUS AFFECTIONS CAUSING BLEEDING.

Here only some remarks will be made on the general principles to be followed in forming an opinion in such cases.

In the first place, always examine the abdomen, in these as in all other gynæcological cases. If any tumour exist there, keep the possibility of pregnancy in mind, manipulate and ascertain if it has the feeling of the pregnant uterus, particularly noticing if the tumour hardens during manipulation. Listen over it at various parts, and determine whether the uterine souffle, or the fœtal heart sounds, can be heard at any part of it, or not. Observe the condition of the external genitals, whether congested, bluish, and very moist as in pregnancy, or not; on vaginal examination, notice whether the cervix is hard or soft, and above all carefully employ the bimanual method of examination to ascertain the size, shape, and consistence of the body of the uterus. In fact, in all cases in the first instance, consider the question of pregnancy, and endeavour to accumulate proof either of its presence, or absence; the latter conclusion will only be justified if you are certain that the uterus, as estimated bimanually, is not the least bit increased in size.

If pregnancy has been excluded, consider the various causes of bleeding not distinctly connected with pregnancy given above. Some of them are diagnosed by inspection, e.g., vascular caruncle, or malignant disease of the external parts; others, such as foreign bodies or polypi in the vagina, are detected at once by the examining finger, as also is malignant disease of the cervix uteri, if at all advanced.

The bimanual exploration of the regions at the sides of the uterus will show whether small ovarian tumours, or swellings due to inflammation of the uterine appendages, are present. The speculum will aid in settling whether the bleeding be due to some condition of the vaginal portion of the cervix—as, for instance, malignant disease of the cervix in an early stage, and affecting the region round the os uteri. If neither inspection of the external parts, nor the vaginal examination, nor the bimanual examination, discover any sufficient cause for the bleeding, it will remain for consideration whether the cause is some condition in the cavity of the uterus itself; this can only be positively ascertained by dilating the cervix, and exploring the cavity of the uterus with the finger. The age of the patient should also be considered, for if no local condition to account for bleeding has been discovered, it may be accounted for by the approach of the menopause.

It need hardly be said that in young, unmarried women, if nothing is found on examination of the abdomen, medicinal treatment should be fully tried before making any local examination.


Treatment.—From a consideration of the various causes of bleeding that have been enumerated, it will be evident that the treatment will depend on the cause. Here it is only intended to mention the management of cases where, so far as can be ascertained, there is no local organic disease, and where consequently no *special* local treatment is required. The menorrhagia of puberty and of the menopause are examples of such cases.


If the bleeding is at all considerable, rest in bed should be advised. Vaginal douches of hot water (as hot as the patient can bear it) should be given three or four times a day. For medicine, some preparation of ergot should be given; we may choose either ergotin (gr. ij.-v., in the form of pill, three times a day), or the liq. ergot. ammon. (Martindale) (mxx. three times a day), or the ext. ergot. liq. B.P. (3ss. three times a day). It is often beneficial to give bromide of potassium or ammonium in scruple doses as well, particularly in the menorrhagia of the menopause.

Other drugs from which benefit may be expected are nux vomica and quinine.

Gallic acid in 20-grain doses every three or four hours; aromatic sulphuric acid (mxx. quartis horis); tincture of hamamelis (mv. three times a day); tincture of hydrastis (mxx. three times a day); infusion of vinca major (two ounces of the leaves are infused in a pint of boiling water, and 3j. is given every three or four hours), form a supplementary list to choose from if the hæmorrhage is not controlled by the drugs previously mentioned.

If the hæmorrhage is alarming, and the cervix sufficiently patent to allow of the proceeding, the cavity of the uterus may be plugged with a long narrow strip of iodoform gauze. To do this properly the os is exposed with Sims's speculum, and its anterior lip seized with a volsella; the gauze is then to be packed in by means of uterine forceps, or Récamier's curette. Should there be a difficulty in doing this owing to narrowness of the cervical canal, a few sizes of Hegar's dilators, say up to No. 7 or 8, may be first passed. We are of course supposing the hæmorrhage to be really serious. The plug is, however, only a temporary measure, and is not to be left in more than twenty-four hours. Thorough exploration of the endometrium should be insisted on subsequently.





HÆMORRHAGES NOT CONNECTED WITH PREGNANCY.

### CHAPTER IV.

### DYSMENORRHŒA.

THE patient complains of pain occurring in association with her menstrual periods. The pain may begin a few days before the period, or may only occur during the flow, or even only occur after the cessation of the flow. Usually the pain is not so distinctly limited as to occur only before, during, or after the flow. Enquiry should be made as to when the pain is at its worst.

It would be incorrect to speak of every pain occurring at the menstrual period as *dysmenorrhæa*. The pain should be contained within "the genital sphere," *i.e.*, within the regions affected by a well-marked disease of the pelvic organs, such as cancer (Champneys). The boundaries of this region are, above, a line level with the iliac crests in front and behind, and below, the level of the knees.

Cases of dysmenorrhœa may be divided into three groups:—

- I. Cases where a careful local examination discovers no abnormality whatever in the uterus or its appendages, and where also there is no gross abnormality in the process of menstruation itself, such as the passage of a cast of the uterus or of pieces of membrane. (N.B.—Flexions of the uterus are not counted as abnormalities.) The dysmenorrhœa in this group of cases is known as spasmodic, and also as neuralgic dysmenorrhœa.
  - II. Cases where some abnormality is detected on physical

examination. This group includes, among others, the cases known as inflammatory, or congestive, dysmenorrhoea.

III. Cases where some gross abnormality exists in the process of menstruation itself, the only known instance of this being the passage of a membranous cast of the uterus, either in one piece, or broken up into several pieces—membranous dysmenorrhæa.

Diagnosis.—In considering into which group we must put any particular case of dysmenorrhoea, it will usually be easy to decide at once whether it belongs to Group III. or not—it is only important not to mistake a case of membranous dysmenorrhoea for an early abortion; the difficulty will be to decide whether it should go into Group I. or Group II.

If the most careful physical examination can detect no abnormality, the case must be classed in Group I. If, on the other hand, something abnormal is discovered, then the case goes into Group II. It will be evident that we cannot ever be certain that a case belongs to Group I.; the most we can affirm is that our examination discovers nothing abnormal. Physical examination, even in the most skilled hands, is far from enabling us to discover every morbid condition during life that a post-mortem examination would detect. "Many cases have now been recorded where abdominal section showed the Fallopian tubes to be dilated with pus to the size of coils of small intestine, although the most careful bimanual had failed to detect their presence" (Hart and Barbour). And if so marked a degree of disease can escape detection by physical examination, we may be sure that morbid conditions of a less extensive character are frequently overlooked.

Filmy and filamentous adhesions, the remains of a previous attack of pelvic peritonitis, are especially to be thought of in connection with this point. For the present we must be content to take it, that cases of dysmenorrhoea exist apart from any physical abnormality, but we ought to regard such cases with suspicion, remembering that physical abnormalities may be present, though we cannot discover them.

The chief abnormalities alluded to, which at once place the case in Group II., are:—

Imperfectly developed uterus.

Interstitial inflammation of the uterus—metritis.

Pelvic inflammation (pelvic peritonitis and pelvic cellulitis), evidenced by feeling the mobility of the uterus impaired, perhaps so much so that the uterus is completely fixed, and by the presence of lumps in various parts of the pelvis; if at the sides, in the posterior quarters of the pelvis, such lumps are often really made up of distended Fallopian tubes, matted by adhesions to the ovaries and adjoining parts. (This subject will be considered in detail under Pelvic Inflammation.)

Fibroid tumours.

Ovarian tumours in some cases.

Retroflexion, or retroversion, with incarceration of the body of the uterus in Douglas's pouch.

## CLINICAL FEATURES OF SPASMODIC DYSMENORPHOEA.

GROUP I.—In this group of cases the dysmenorrheea is usually, but not always, *primary*, *i.e.*, it dates from the establishment of the function of menstruation. The pain is usually paroxysmal, and referred to the hypogastric region.

As a rule, the pain begins a few hours before the flow, and continues for the first day or two of the period. The flow then usually increases in quantity, and the pain diminishes, or ceases. The pain is at its worst while the flow

is scanty. The blood may contain clots or shreds of membrane. As time goes on the patient usually gets worse. Marriage tends to increase her trouble, unless pregnancy ensues; after which the dysmenorrhœa is often permanently cured. Many patients with spasmodic dysmenorrhœa are, however, unfortunately sterile. Retching or vomiting often occurs at the periods in cases of spasmodic dysmenorrhœa.

It is often found that the severity of the symptoms—pain, retching, vomiting, etc.—varies greatly at different menstrual periods. Sometimes one or two periods may be passed through without the occurrence of any symptoms. This fact should be remembered when estimating the effects of any particular treatment.

In many cases pain similar to that complained of at the periods may be produced at any time by passing the sound, and it occurs in typical cases just as the sound passes the internal os uteri. The pain is, however, only produced if the sound used is large enough to *stretch* the internal os uteri.

The pain in spasmodic dysmenorrhœa is certainly due to "colicky" contractions of the uterus. This kind of dysmenorrhœa is, as Dr. Champneys says, "essentially a neurosis, and has motor phenomena (colic), and vaso-motor phenomena (scanty or greatly varying flow)."

## CLINICAL FEATURES OF INFLAMMATORY DYSMENORRHŒA,

GROUP II.—So far as relates to the cases known as congestive or inflammatory dysmenorrhoea the affection is not primary, but dates from some well-recognised antecedent of pelvic inflammation, e.g., labour or abortion. The pain is of a constant, aching character, and precedes

the flow, being relieved when this occurs, and especially is this so when the loss is free. The pain may begin a variable time before the period, days or even weeks, reaching its greatest intensity just before the flow. As to the causation of the pain, in these cases it should be remembered:—

- 1. That uterine contractions take place during the process of menstruation.
- 2. That at the menstrual period there is an afflux of blood to the reproductive organs, uterus and ovaries particularly, and that this afflux of blood causes these organs to be distinctly increased in size at this time.

Coming now to theory, it may be suggested that any condition which will prevent the uterus and ovaries readily accommodating this increased amount of blood sent to them, and any condition which will interfere with the alterations, in position and size, that the uterus undergoes during its contractions, will afford an intelligible explanation of the pain.

To take an example, if there is interstitial inflammation of the uterus (metritis), it may be taken that the excess of blood endeavouring to find its way into the vessels of the uterus at the menstrual period, will be less readily accommodated than if the tissues of the uterus were healthy. The uterine tissue being in a state of disease cannot easily permit the increase of size and change of shape needed to accommodate the excess of blood coming to it at the menstrual period. This view, so far as metritis is concerned, is not very different from the theory, modified from Fritsch, put forward by Hart and Barbour, viz., that the pain is due to the flushing of a diseased tissue with blood. But I think that we must not only regard interstitial changes in the uterus as offering resistance to the afflux of blood, and so accounting for the pain; conditions of equal importance

outside the uterus, and affecting the ovaries and Fallopian tubes, also offer resistance to the afflux of blood, and take their share in producing pain. More particularly, adhesions between the uterus and adjoining parts, or between the ovaries and the Fallopian tubes, require to be mentioned as causes acting in this direction. For the sake of argument take an extreme case, where there are dense adhesions, not only between the peritoneal coat of the uterus, the ovaries, and Fallopian tubes, but also running on to the walls of the pelvic cavity, matting the organs firmly together, and binding them also to the pelvic walls.

Imagine now that the vessels of the uterus and ovaries are suddenly called upon to accommodate a much larger quantity of blood than they have previously held. In order to do so the uterus and ovaries must swell up, so as to be obviously larger than before, and in doing so they will necessarily undergo some alteration in position, but the adhesions mentioned will prevent the changes taking place as they should do. The resistance offered to these changes of size and position, that the uterus and ovaries naturally ought to undergo at the menstrual period, may account for the pain.

Adhesions will also interfere with the contractions of the uterus that occur at the menstrual period, and some of the pain will be due to this interference. Similarly it is probable that interstitial inflammation of the uterus or ovaries by offering resistance to the menstrual afflux of blood will cause pain.

We may then state this theory in general terms as follows:—

Any condition, either in the tissues themselves of the uterus or ovaries, or external to them, which renders these organs less able to undergo the changes of size, shape, and position, necessary to accommodate the increased supply of blood coming

to them at the menstrual periods, will be a cause of dysmenorrhaa, as also will be any condition impeding the contractions of the uterus that occur at the periods.

Inflammation of the tissues of the uterus (metritis), and similar inflammation of the ovaries, will be conditions in the former category; pelvic peritonitis producing adhesions in various parts of the pelvis is the cause referred to as external to the uterus and ovaries. Either metritis, or pelvic peritonitis producing adhesions round the uterus, is likely also to impede the contractions of the uterus, and so in that way cause pain.

Clinically we often meet with cases of dysmenorrhoea dating from a particular confinement, previous to which there had been no pain at the menstrual periods. We usually find evidence of old pelvic inflammation in these cases, thickening round the uterus, diminished mobility of the uterus, and so on. It is evident that such cases are explained by the theory just above referred to.

The presence of a growing fibroid tumour of the uterus, as is well known, is often associated with dysmenorrhoea; this might be explained partly on the view that the presence of a fibroid tumour in the wall of the uterus interferes with the uterine contractions, acting indeed somewhat like a foreign body, and partly on the view that a growing fibroid is the cause of an excess of blood, over and above what would normally come to the uterus at each period, coming to it, and that this excess cannot easily be accommodated, and hence we have pain.

CLINICAL FEATURES OF MEMBRANOUS DYSMENORRHŒA.

GROUP III.—Here the dysmenorrhœa is essentially primary, *i.e.*, in the large majority of cases it begins at the time menstruation is established.

The pain begins just before, or at the beginning of, the flow, and is "colicky" or paroxysmal in character.

The flow is profuse, but in many cases diminishes in 24 to 36 hours, the pain becoming greater; then the membrane is passed, the pain is relieved, and the flow once more becomes free.

Cases in this group differ from cases of spasmodic dysmenorrhœa (Group I.) in that the flow is generally profuse and the pain is not relieved by the flow, but only by the passage of the membrane. The pain in cases of membranous dysmenorrhœa is probably not due to obstruction of the canal by the membrane. For at the beginning of the period there is the pain, but the canal is certainly not plugged then. Possibly plugging of the canal may be the reason why the pain becomes more intense in many cases at the end of 24 or 36 hours; but here again we have another and, I think, a better explanation, viz., the passage of a foreign body (the membrane) over a specially sensitive spot—the internal os uteri.

A point of much practical importance is to distinguish cases of membranous dysmenorrhœa from cases of early abortion.

A case to be a genuine example of membranous dysmenorrhoea must be one in which a membrane is passed regularly at monthly intervals for a considerable time. Where a membrane is only passed occasionally with postponement of menstruation, the case is probably an instance of early abortion; and there is the same probability if a membrane is only passed when the patient is cohabiting with her husband and ceases to be passed when marital relations are interrupted.

Anatomically the sac of an early abortion is ovoid and more vascular, the membrane of membranous dysmenorrhoea is more triangular and less vascular. In thick-




Fig. 22.—A Dysmenorrhoeal Membrane as seen under Water (Sir J. Y. Simpson).

The late Dr. Matthews Duncan wrote to me that he was sure this figure represented an early abortion, and went on to say: "The membrane of dysmenorrhœa never reaches that state of size and development. I have seen many like that on p. 84, no ovuline structures, merely decidua, and the proof is—cured by marital separation."

ness menstrual membranes vary from  $\frac{1}{25}$  to  $\frac{1}{12}$  inch, rarely  $\frac{1}{6}$  inch. Abortive deciduæ vary in thickness from  $\frac{1}{5}$  to  $\frac{2}{5}$  inch (Champneys).

The membrane passed in dysmenorrhœa, as a rule,

corresponds only to the body of the uterus, and its length varies from  $1\frac{3}{4}$  to 2 inches. A specimen decidedly exceeding this measurement is probably an early abortion.

Microscopical investigation may enable us to make a diagnosis, if sections of the substance passed show the presence of chorionic villi.

As to the obstructive theory of dysmenorrhoea, which was formerly put forward as explaining most cases of dysmenorrhoea, it may be said that it is now to a great extent discredited by the best authorities. The theory was, that either on account of congenital stricture at the external or internal os uteri, or on account of flexion, an obstruction existed to the flow of blood out of the uterus, and that consequently excessive and painful contractions were excited.

Narrowing of the external os is met with, but it is not at all common; that stricture ever occurs at the internal os, before the menopause, has not been proved. After the menopause narrowing or actual occlusion of either the external or the internal os, or at both orifices, is far from uncommon.

As regards flexion producing obstruction, we may also say that it has not been proved that it does so, or at least only in the slightest degree; on the other hand, it has been shown in many of the worst cases of dysmenorrhoea, by actually passing the sound at the time that no obstruction existed.

And, further, it should be remembered, that in some cases where there is a narrow external os—"pin-hole os"—and in others where a membranous cast is thrown off, there is no dysmenorrhæa at all.

Treatment.—In slight cases the patient should be advised to avoid over-exertion and any exposure to chill at the menstrual periods. Care should be taken that the bowels are kept acting a little more freely than usual, when

the menstrual period is at hand. Putting the feet in hot water with mustard in it, and hot drinks at bedtime, are also useful. For medicine we may try cannabis indica (from  $\frac{1}{2}$  to I grain of the extract in pill) thrice daily for a few days before, as well as during, the period. It may be advantageously combined with camphor (gr. ij.-iij.) and a small dose, gr.  $\frac{1}{8}$ , of extract of belladonna. Bromide of potassium and aromatic spirits of ammonia are also useful, gr. x. to xx. of the former, with  $\mathfrak{M}$  xx. of the latter, thrice daily at the periods, at the same time doing all we can between the periods to improve the patient's general health by exercise, particularly riding, tonics, and especially careful attention to the state of the bowels.

As Dr. Champneys humorously puts it, "with constipation none of this class improve, even if treated by the newest alkaloids."

Guaiacum, sulphur, antipyrin, ichthyol, and castoreum form a supplementary list to choose from. Of these I have found most benefit from antipyrin and ichthyol. Tabloids, containing 5 grains of antipyrin, are convenient. Two should be taken when the pain is felt to be coming on; and if the pain is not relieved in two hours, another two may be taken. Not more than four tabloids should be taken in any one day.

Ichthyol is conveniently given in coated pills, containing gr.  $1\frac{1}{2}$  in each. One of these may be given twice or thrice daily for a few days before each period, and during the period.

Whether cases are slight, or severe, the line of treatment sketched out is that to be fully tried first, more especially for obvious reasons in unmarried women. After it has been tried and failed, the next thing to think of is dilatation of the cervix by bougies. Starting from the size (No. 9) that will normally pass (Matthews Duncan), or a smaller

one, if that will not pass, the bougies are passed, one after the other, going up two or three sizes on each occasion, the dilatations being continued at intervals of a few days. simpler and better plan is to anæsthetise the patient, and dilate on one occasion as much as seems desirable, say up to No. 18 of Matthews Duncan's bougies, the instruments usually employed.\* If necessary, the cervix is fixed with a tenaculum, or volsella, during the dilatation. This measure must always be adopted before deciding that the ordinary sound will not pass the external or internal os in any particular case. The treatment by dilatation is suitable for the cases in Group I., where there are no physical signs, and for cases of narrow external os. Sometimes the dilatation may be completed at one sitting without an anæsthetic. In the cases associated with the presence of inflammatory conditions of the pelvis, as evidenced by feeling lumps, or thick firm bands, in the neighbourhood of the uterus, and by the mobility of the uterus being less than normal, dilatation of the cervix is usually contra-indicated.

The choice here lies between being content with palliative treatment, and entertaining the question of removing the uterine appendages; it need hardly be said that the consequences—sterility—and the risks of the operation should be fairly laid before the patient; particularly she should be asked if she would not rather put up with the pain, than have an operation—to which some considerable risk often attaches—done to relieve it. This line of treatment is only indicated in really bad cases, when palliative treatment has failed, and where distinct physical signs, such as lumps at the sides of the uterus, are present, and, lastly, where the

<sup>\*</sup> Hegar's dilators are equally suitable for the purpose. No larger size than No. 12 need be used when dilating for dysmenorrhoea. Strict antiseptic principles should be observed in this, as in all other operations on the cervix.

operation is to be done by a specialist skilled in abdominal surgery.

It is possible that a sort of "massage" of the pelvic organs may give relief, where there are adhesions of only moderate strength around the uterus limiting its mobility. Manipulation under an anæsthetic by breaking and stretch-




Fig. 23.—A Dysmenorrhæal Membrane laid open (Coste).

If a portion of such a membrane is placed on a slide, and looked at with a low power under the microscope, it will be found to be perforated with numerous small holes corresponding to the openings of the uterine glands.

ing some of the adhesions might perhaps give relief; but treatment of this kind is undesirable, for it is obviously open to objection; besides, it may be exceedingly dangerous, if there should happen to be a pyo-salpinx present.

As to treatment of the cases in Group III., it is to be remembered, that it is not always that the passage of casts of the uterus, or pieces of membrane, gives rise to any

dysmenorrhoea. When there really is pain in these cases, we know no very satisfactory treatment. The usual round of remedies already mentioned may be used. If these fail, dilatation with bougies, as already described, may be tried, if the other conditions present do not contra-indicate it.

Scraping the endometrium, in addition to dilatation, has also been found useful, at least for a time.

(Those wishing to pursue the subject further are advised to read the *Harveian Lectures on Painful Menstruation*, by Dr. Champneys.\*)

<sup>\*</sup> Published by H. K. Lewis, 136 Gower Street, London.

#### CHAPTER V.

ON DISEASES OF THE PUDENDUM.

#### VULVITIS.

**Definition.**—An inflammation of the labia majora, and parts lying between them, the anterior limit of the inflammation being the Mons Veneris, and the posterior limit the fourchette (*i.e.*, the anterior margin of the perineum).

Suppurative vulvitis, without any vaginitis, is the commonest disease of the genital organs met with in children. The mother brings the child on account of its having "a discharge from the private parts," and, as is well known, this may have raised a suspicion, usually without any foundation in fact, of the child having been criminally assaulted.

There is usually pain in passing water, owing to the orifice of the urethra being involved. On inspection, the surface of the external genitals is found to be redder than normal, and secreting pus. The parts are very tender. Occasionally more or less extensive ulceration of the inflamed surface is seen. The ulcers are shallow, well defined, and vary in size from that of a florin downwards.

In children, it is important to remember that the vagina is not implicated in the inflammation, or at most only the very entrance of it—there is vulvitis without vaginitis. This may be shown by first wiping away all the pus from the surface, and then with the little finger in the rectum, pressing downwards and forwards, and observing that no pus can be squeezed out of the vagina.

Causation in children.-The fact that almost in-

variably the hymen is seen to be perfect, and that there are no lacerations of the perineum, sufficiently disposes of the




Fig. 24.—External Genitals in a Virgin.

I I. Right labium majus; 2. The fourchette; 3 3. Right labium minus, or nympha; 4. Glans clitoridis; 5. Urethral orifice; 6. Vestibule (a triangular space, the sides of the triangle formed by the nymphæ, the apex by the clitoris, and the base by a transverse line, which is bisected by the urethral orifice); 7. Orifice of the vagina; 8 8. The hymen, running in this case all round the vaginal orifice, but still, broader posteriorly; 9. Opening of the duct of Bartholin's gland on the right side—this is represented too far forward, it should be about as far back as the posterior 3 in the figure; 10. Mons Veneris (Thorburn).

notion that the disorder is the result of an assault. If there were lacerations, the possibility of this would require careful consideration.

Vulvitis may be caused by cold, or by dirt, but not infrequently it arises without any assignable cause in weakly children. The late Dr. Matthews Duncan was strongly opposed to accepting the presence of worms as a cause of vulvitis, though it is one commonly given. I have seen one case where it seemed to me that indirectly the vulvitis




Fig. 25.—Vertical Antero-posterior Mesial Section of the External Genitals (Henle).

a. Rectum; b. Perineal body; c. Vagina; d. Urethra; e. Labium minus; f. Prepuce of the clitoris; g. The fossa navicularis, bounded in front by the hymen and behind by the fourchette (the anterior margin of the perineum).

was due to the presence of worms in the rectum. The orifice of the vagina was wider than normal, and less perfectly closed by the hymen. In this case, there certainly were thread-worms in the rectum, and it seemed to me at least probable that the child had passed her finger into the vagina, in her endeavour to allay the uneasy sensations caused by the worms in the rectum, and that the vulvitis was due to this mechanical irritation.

Diphtheria sometimes affects the vulva, as in the following case:—

E. B., age 7, was brought to the London Hospital on January 22nd, 1887. On January 18th she had fallen over a heap of stones, and hurt




Fig. 26.—Diagram to Illustrate the Relations of the External Genitals. A in front view, B in antero-posterior section (Matthews Duncan).

a. Urethral orifice; b. The fossa navicularis; c. The fourchette, the anterior margin of the perineum, hh; dd. The openings of the ducts of Bartholin's glands (sometimes they open rather farther forwards); e e e e. Margin of vaginal orifice, internal to which is seen the hymen, broader behind than in front—it has been torn near m; ff. Posterior extremities of the labia majora, which are not united posteriorly, but merely connected by the perineum; g. The anus. The lettering in both figures is the same. The "inevitable" laceration, i.e., the one that occurs in every first labour, extends into the fossa navicularis, b (see chapter on Rupture of the Perineum).

herself in the private parts. Next morning her mother noticed that she screamed on passing her water, and that the parts were swollen. On January 20th the child complained of sore throat, and she had a cough. A "few spots of blood" were brought up with the cough; on examination (January 22nd) the external genitals were found to be swollen, and the whole surface of the vulva was covered with false membrane of a greyish-yellow colour. On looking at the throat, the fauces were seen to be covered with false membrane also. The child died rather suddenly on January 28th, after seeming to be going on well.

Causation in adults.—Suppurative vulvitis in grown-up people is a part only of an inflammation affecting the vagina as well—there is vaginitis as well as vulvitis. An inflammation of this kind may be caused by gonorrhæa, or by injuries inflicted on the parts during a difficult labour. In acute cases the labia are enormously swollen, so as to be shining and ædematous, forming a sausage-shaped swelling on each side of the vulvar cleft.

Vulvitis without suppuration.—In these cases the inflammation is characterised by increased redness only (erythema), or by the presence of papules, or vesicles, as well, scattered over the inflamed surface.

Vulvitis of this kind is caused by the passing of irritating discharges over the parts; we get it, therefore, in cases of incontinence of urine; or, apart from incontinence, where the quality of the urine causes it to be unusually irritating, as in diabetes.

In this last case there is also a constitutional predisposition to inflammation of the skin, not only of the vulva, but elsewhere.

#### CASE OF DIABETIC VULVITIS.

H. T., age 70, six children, the last twenty-seven years ago, came to the London Hospital complaining of great irritation of the private parts, and "throbbing" there. She "does not know how to sit down" on account of it. The present attack has lasted a month, but she suffered from the same thing four years ago, when it lasted three months.

She is very thirsty, and drinks a great deal of water. Has been losing flesh for the last twelve months. She has scalding on passing water, and has to get up twice at night to pass it.

Patient is very fat; about a hand's breadth above the pubes the abdominal wall is thrown into a transverse fold; where the skin-surfaces are in contact there is a simple inflammation of the skin, its surface being red and moist, but there are no vesicles or papules on the affected surface. The labia majora are hypertrophied, the whole of the external genitals are purplish-red and raw-looking. In the fold between the labia and the thighs on each side, and in the fold of the nates, extending to about three inches behind the anus, there is similar inflammation of the skin.

Urine, sp. gr. 1040, contains no albumen, but a good deal of sugar.

Vulvitis is an early symptom in diabetes, often the earliest to attract attention. The appearance of the vulva is very characteristic (see the case just recorded), and, to any one who has seen it, at once suggests the advisability of testing the urine.

The absence of vulvitis does not, however, prove the urine to be free from sugar. I have seen cases where the appearance of the vulva was normal, yet the urine contained a large amount of sugar.

Again, the discharge from the vagina in cases of cancer, uterine or vaginal, the presence of pediculi pubis, or any other source of itching causing the patient to scratch the parts, will set up a simple non-suppurative vulvitis.

Treatment.—In children it is sufficient to have the inflamed surface bathed night and morning with weak lead lotion; the mother should be instructed to separate the labia, so that the lotion may reach the whole area affected. Sitting in a hot bath for a quarter of an hour at bedtime every night, or two or three times a week, is also useful. If the child has thread-worms, these should be got rid of by rectal injections of salt and water. The general health

should be improved as much as possible by tonics, cod-liver oil, and proper diet. When ulcers are present, an effectual, but rather painful plan, is to rub them over with solid nitrate of silver, or sulphate of copper.

In adults suppurative vulvitis requires similar treatment, but as vaginitis is also present, vaginal injections must be used (see Vaginitis). In the non-suppurative vulvitis of adults, treatment will depend on the cause. Pediculi may be got rid of by smearing the parts well with terebene oil (1-6), or washing the vulva with corrosive sublimate lotion (1-1000). When the cause is incontinence of urine, or the discharge from malignant disease, the treatment of any coexisting vulvitis is a very minor matter, but soothing applications, such as lead lotion, zinc ointment, etc., may be used, if treatment seems desirable. The urine should always be tested for sugar in cases of non-suppurative vulvitis occurring in adults. The youngest patient, in whom diabetic vulvitis has led me to the diagnosis of diabetes, was sixteen years old.

## VASCULAR CARUNCLE OF THE URETHRA.

This is a bright red body, either sessile, or more or less distinctly pedunculated, attached just within the margin of the urethral orifice, but projecting beyond it. Sometimes it does not project, owing to being some distance up the urethra, and it will then not be readily discovered by mere inspection. In size it varies from that of a pin's head to that of a small grape; on an average, it is about the size of a pea. There may be two or three of various sizes present in the same case.

Vascular caruncle is more common in women accustomed to sexual intercourse than in virgins; there is often some urethritis present, as evidenced by a discharge from the urethra.

The symptoms it causes are :-

Frequent desire to pass water, and pain in passing it. (It is noticed in some cases that the pain on passing water only occurs at the menstrual periods—that is, at a time when the pelvic viscera are specially congested.)

Pain on sitting down.

Pain on connection (dyspareunia).

Sometimes a little bleeding from the surface of the caruncle.

Occasionally one accidentally discovers a caruncle that has not caused any symptoms.

Structure.—A vascular caruncle is composed of capillaries, nerves, and a small quantity of connective tissue.

Treatment.—The best treatment is to burn it away with Paquelin's cautery.

Sometimes these growths recur after removal. They are more likely to do so after removing them with the knife or scissors than after cauterisation.

## LABIAL ABSCESS AND CYSTS OF THE PUDENDUM.

These may conveniently be considered together. The glands of Bartholin are two in number, and lie one on each side of the middle line, the situation of each being marked by taking a point on the skin surface midway between the posterior extremity of the labium majus and the adjoining tuber ischii (Matthews Duncan). The duct is rather more than half an inch long, and runs forwards and inwards to open either in the fossa navicularis (the space between the anterior margin of the perineum and the attachment of the hymen), or rather further forwards, in the angle made by the attachment of the hymen with the adjoining part of the vulva. The commonest abscess met with in the pudendum is an

abscess of the duct just mentioned, and similarly the commonest cyst of the vulva is a retention cyst of the same duct.

Labial abscess.—Inflammation spreads from the vulva along the duct. The orifice of the duct may become occluded, or remain open; even if it remains open, it may not allow a sufficiently free escape of the pus, and con-




Fig. 27.—Abscess of the Duct of Bartholin's Gland on the Right Side (Huguier).

The drawing might also stand for a cyst of the duct, which causes a swelling of similar size and position.

sequently the part of the duct behind the orifice dilates. The swelling so formed is commonly as large as a plover's egg, and occupies the posterior half of the labium majus on the affected side (Fig. 27). Such abscesses are often a complication of gonorrhœa, but they may arise from a simple, non-gonorrhœal inflammation. If the orifice of the duct is open, pressure on the swelling will cause pus to ooze out.

Labial cysts arise by occlusion of the orifice of the duct, and accumulation of secretion in the duct, which dilates and forms a swelling in the same position as that caused by the labial abscess (posterior half of the labium majus, Fig. 27).

It may be mentioned that, in addition to Bartholin's




FIG. 28.—Large Cyst of the Duct of Bartholin's Gland dissected out entire. Natural size after some months in spirit. A square opening, A, has been cut in the anterior wall of the cyst. At the operation it appeared much larger, and had a rough resemblance to a piece of distended intestine. The larger end of the cyst lay backwards.

glands, there are smaller glands in the vulva, viz., mucous glands, very numerous in the vestibule, and sebaceous glands, very numerous on both aspects of the nymphæ; it is evident, then, how small retention cysts, either mucous or sebaceous, or small abscesses, may originate in these situations.

Treatment.—In the case of an abscess a free incision should be made on the inner aspect of the swelling, and the cavity dressed with a strip of iodoform gauze. In the case of a cyst, a piece the size of a sixpence should be cut out of the wall of the swelling on its inner aspect. Pure Tr. Iodi should be applied to the lining membrane of the cavity to destroy its secreting power. It may then be dressed as in the preceding case. A better plan, however, is to dissect out the cyst entire. Several pairs of small Wells's artery forceps should be at hand, as there are generally a good many bleeding points to secure.

In connection with this subject it must be remembered that rare forms of hernia are met with in the vulva. The one which is commonest is a hernia running along the unobliterated canal of Nück, that is, a finger-shaped prolongation of the peritoneum running along the round ligament into the labium majus. The practical point being, that before opening any swelling in the vulva, every care should be taken to make sure that it is not a hernia, e.g., by trying if it can be reduced, by percussion, and so on.

## WARTS.

Warts are very commonly met with on the vulva and adjacent skin. They are sometimes found on the lower part of the vaginal mucous membrane as well, and in such cases they may also be found on the vaginal portion of the cervix, especially when warts are situated at the upper part of the vagina, and so in contact with the vaginal portion of the cervix. They may be caused by gonorrhæa, or be only due to dirt. The best treatment is to burn them off with Paquelin's cautery.

## THE PRIMARY SYPHILITIC SORE.

This is not often seen on the vulva. It has been said that the primary sore in this situation not infrequently lacks the hardness characteristic of the initial sore elsewhere. This is a very difficult matter to be sure of. It is almost impossible to decide in a patient, who is certainly suffering from early secondary symptoms, whether a soft sore at some part of the vulva is the primary sore, or a secondary effect merely. It is also rare to get a reliable history in such cases. Observations in the case of prostitutes are complicated by the probability that the patients may have had syphilis at some previous time, and that whatever symptoms are present may be due to a revived activity of the disease. I have, at all events, seen several cases where the initial sore on the vulva, or just within the vagina, was typically hard—as in the following case:—

Elizabeth D., age 24, was admitted to Davis Ward of the London Hospital on May 25th, 1887, with a sore place in the private parts.

She had only been married about six weeks, having been married on Easter Sunday last (April 10th).

About ten days before admission (i.e., thirty-five days after marriage), she noticed a pimple in the private parts. It itched and smarted, and gradually increased in size.

State on admission.—In the situation of the fossa navicularis, there is an irregularly shaped ulcer; from this a somewhat elevated and distinctly indurated sore extends to the fourchette, and for half an inch behind it over the perineum. The surface of the raised patch is superficially ulcerated. The sore lies chiefly to the left of the middle line. The inguinal glands are enlarged on both sides, but are not tender. No rash on the body, or sore throat. She was ordered a warm bath every night, the sore to be bathed with lead lotion, and mercury was given internally.

On June 4th I saw the patient's husband. He had numerous ulcerated patches on the glans penis, and there was also a well-marked cicatrix of an older sore. Glands enlarged both sides, not tender; there was a

rash on the chest—the man said it had been much brighter, but that it was now fading. There were mucous patches on both tonsils. He ascribed his local symptoms to a blow from a cricket-ball. I sent him round to the surgical out-patient department, where I know he was put on a course of mercury.

To return to his wife's case: the sore rapidly improved under treatment, and the patient was made an out-patient, but she was still kept on mercurial treatment. In October 1887 she came up to the hospital, and said she was three months pregnant. Her tonsils were then

greatly enlarged, and superficially ulcerated.

She suffered from sore throat and ulcerated tongue more or less during the year she was under my observation. She went her full time, and the child was born alive. It subsequently had erythema of the buttocks and genitals, which Mr. Moullin, to whom I sent the child on account of phimosis, regarded as certainly syphilitic. The mother had been taking mercury for three months at a time during the whole of her pregnancy, and it will be seen that at the time she became pregnant her husband was also under the influence of mercury.

## MUCOUS TUBERCLES.

These are frequently met with, and are invariably of syphilitic origin. They appear as flat, raised patches, more or less circular or oval, of a greyish-red colour, and with a moist surface. They occur not only on the vulva, but round the anus, as in the male sex. Mucous tubercles are distinguished from warts by the fact that they have no pedicle or any indication of one—no more pedicle, in fact, than a penny laid flat on the surface of the skin would have. Locally, cleanliness and blackwash will cure them; but as they are only an evidence of syphilis in an active stage, constitutional treatment is necessary.

## ULCERS.

It has already been mentioned that extensive superficial ulceration is occasionally noticed on the pudendum in the simple vulvitis of children. In the vulvitis of adults it is

not at all uncommon to see more or less ulceration affecting the opposed surfaces of the inflamed labia.

Apart from vulvitis, we may find two, or three, or more ulcers, on the inner surface of the labia, or in the fossa navicularis, as a result of secondary or tertiary syphilis.

#### THE HYMEN.

The hymen is usually either:—

(1) Crescentic in shape with the horns of the crescent forwards;




Fig. 29.—Diagrammatic Representation of the Various Forms of Hymen.

- 1. Crescentic form (the commonest); 2. Circular form, nearly as common as 1; 3. Cribriform hymen; 4. Imperforate hymen; 5. Hymen biseptus; 3, 4, 5, are all rare. a. The urethral orifice. The shaded part of each figure indicates the hymen. I have seen several examples of 4 and 5, but have never seen 3.
  - (2) Or circular, and of the same width, or nearly so, all the way round the vaginal orifice.

Rarely:-

- (3) It may be cribriform;
- (4) Or imperforate;
- (5) Or double.

Imperforate hymen (4).—In this condition the vaginal orifice is closed, owing to the hymen taking the form of a complete membrane stretched across it.

**Symptoms.**—Attention will probably not be called to the abnormality till the age of puberty.

The girl will then have colicky pains in the hypogastrium every month, but will not "see anything." Ultimately, if the case go on, a tumour will appear in the hypogastrium, and the patient may notice that this increases in size every month when she has the pain referred to, and diminishes in the intervals, some of the fluid being reabsorbed. The tumour in the hypogastrium may be due either:—

- (1) To the uterus, not increased in size, being perched on the top of the distended vagina, the menstrual fluid only occupying the vagina.
- (2) To the uterus itself being dilated by backward pressure, so that the cavity of the uterus and vagina communicate much as they do in the second stage of labour. The retained fluid in this case occupies the vagina and uterus.

If the case go on long enough, the Fallopian tubes may also become dilated.

Diagnosis.—On inspection the hymen is seen to form a complete septum, closing the vaginal orifice, and, if the case is one of some duration, the hymen is bulged outwards, having much the appearance of half a purple plum.

**Treatment.**—It may either be treated by free incision with rapid evacuation, or by aspirating with a fine needle and slow evacuation.

The plan to be adopted depends on whether the uterus is dilated as well as the vagina, or whether the vagina alone is dilated.

In the latter case free incision with Paquelin's cautery into the hymen may be chosen, the blood being allowed to run out without any pressure being made above the pubes. Vaginal injections should *not* be used. The patient should be kept absolutely at rest in bed for ten days or a fortnight,

A pad of some antiseptic wool should be kept applied to the vulva to receive the fluid.

If the uterus is dilated as well as the vagina, opinions differ as to the best line of treatment. In some such cases the Fallopian tubes are dilated with blood; and if the uterus is aspirated from below, the disturbance of its position or the manipulation incidental to the aspiration, may cause rupture of the Fallopian tubes, and fatal peritonitis. Therefore some authorities advise a preliminary abdominal section, in order to ascertain the state of the Fallopian tubes, and if they are dilated to remove them. Others think it best to draw off a small quantity of the fluid with an aspirator needle, at intervals of a day or two, till the fluid has been nearly all removed, and then to make a free incision into the hymen. The object of this gradual withdrawal of the fluid is to enable the uterus to accommodate itself to its diminishing contents. If rapidly emptied, contractions are set up, some of the fluid, with septic matter from the air, may be forced along the Fallopian tubes into the peritoneum, and death from peritonitis result. Several fatal cases of this kind are recorded (see a case referred to in Chapter XIX.).

## FISSURE OF THE VULVA—INFLAMED PORTIONS OF HYMEN.

It is not uncommon to see one or more fragments of the hymen inflamed and painful, giving rise to dyspareunia, perhaps so great as to prevent coitus altogether. There may be noticed an unhealed fissure between the fragments of the hymen, or a fissure in the fossa navicularis, giving rise to the same symptoms. These conditions should be treated at first by sedative lotions, such as dilute lead lotion; should the latter fail to relieve them, then, in the case of inflamed portions of hymen, these should be dissected off

under an anæsthetic; in the case of fissure, forcible dilatation of the vaginal orifice with the thumbs will probably cure it.

#### VARICOSE VEINS.

These are met with forming a swelling in one, or both, of the labia majora. The condition usually originates during pregnancy. They may be ruptured during labour, or even by slight injuries apart from labour. If the blood escapes only into the connective tissue, we have what is called "thrombus or hæmatoma of the labium"; \* during labour the tumour so formed may obstruct the passage of the child, and may, on that account, require to be emptied; otherwise, unless it suppurates, it is best left alone. If, when the veins rupture, there is an external opening, profuse bleeding occurs, which may be fatal unless assistance is obtained promptly.

## ADHESION OF THE LABIA MAJORA.

This is an affection not uncommon in infants. The adhesion does not extend forwards so as to occlude the orifice of the urethra. The treatment is to break down the adhesions with the fingers, and direct the mother to separate the labia for some time to come, to prevent them reuniting.

\* This, however, is not the usual origin of thrombus of the vulva. On this point Spiegelberg says: "Varicose veins of the vulva undoubtedly occasionally coexist, but they cannot be brought into causal connection with the formation of hæmatomata, since they do not occur more often in conjunction with the latter than at other times; further evidence is derived from the fact that a thrombus is relatively more common in primiparæ, who do not often suffer from varicose veins, than in multiparæ" (Textbook of Midwifery, New Sydenham Society's translation, vol. ii., p. 316).

#### HYPERTROPHY.

This may affect the nymphæ, the clitoris, or the labia majora. Hypertrophy of the nymphæ is sometimes congenital, or it may be the result of masturbation. That of the clitoris is usually congenital. Hypertrophy of the labia majora to a moderate extent is common, as a result of chronic inflammation; an extreme degree of hypertrophy, affecting one, or both, of the labia majora, so that they form large tumours, occurs in hot countries, and is called elephantiasis.

In some cases hypertrophy of the external genitals seems to be due to late syphilis; while in others it seems to be due to a disease *sui generis—lupus* of the vulva, or esthiomène.

This is an extremely chronic affection, and is more especially characterised by hypertrophy of the external genitals and ulceration affecting various parts of the hypertrophied tissues.

The most prominent of the masses may be removed with Paquelin's cautery; the ulcers may also be cauterised. This treatment generally causes improvement for a time, but generally after a while the disease makes fresh progress, with renewed hypertrophy and ulceration.

\* C. W., aged 22, has been married six years, but has had no children or miscarriages; she was admitted to Davis Ward of the London Hospital on August 24th, 1888, complaining of having a swelling in the private parts.

History of the present illness.—She had a yellow discharge from the vagina two years ago. Soon afterwards she noticed a small growth in the vulva, which has gradually increased to its present size.

When the yellow discharge began, the private parts swelled up suddenly, and she had pain in passing water.

<sup>\*</sup> Obst. Trans., London, vol. xxxi., p. 326.

No history of buboes. Of late there has been no pain or trouble in micturition.

The catamenia began at fifteen, and have always been irregular, and accompanied with pain.

The family history was of no special interest.

Present state.—On inspection of the vulva there is seen a somewhat spherical pendulous lump, smooth and of a whitish colour on its inner surface (as at C in Fig. 30), irregular and somewhat nodular on its outer aspect, which has a brownish tint.

The measurements of the growth are as follows:-

Extreme antero-posterior diameter . . . =  $3\frac{1}{2}$  inches. Maximum transverse diameter . . . = 2 ,, Greatest circumference . . . = 8 ,, Projection from the vulva . . . =  $2\frac{3}{8}$  ,,

The tumour grows from the left labium minus, but its attachment is a broad one, and also involves the anterior upper part of the right nympha and the prepuce of the clitoris.

Anteriorly and externally there is a secondary projection, the size of half a walnut, from the general surface of the growth.

The labia majora are thickened by a kind of solid cedema, so that each one measures one inch across. They do not pit on pressure.

Behind and externally the labia majora are studded with small warty prominences, and there are similar ones on the perineum, extending as far back as the anus. There is also one the size of a halfpenny on the mons Veneris.

Where the affected surfaces are in contact there is superficial ulceration. The parts are not tender.

The vagina is healthy as far as the finger can reach.

The glands in the groins are not tender, and are not enlarged.

There is no history of syphilis to be obtained, nor are there any signs of it about the patient. She distinctly says the cause of the yellow discharge was that "her husband gave her the bad disorder," and there is every reason to believe that by this she means gonorrhoea, and not syphilis.

Operation.—August 28th the patient was put under ether, and the pedicle of the tumour was clamped with two pairs of Wells's large pressure forceps. The pedicle was then cut through with Paquelin's cautery.

All the small warty prominences already described were also burnt off,

The large pressure forceps were then taken off, and the cut surface of the pedicle thoroughly seared with the cautery. There was some bleeding even when this was done; and as the tissue of the pedicle seemed too rotten to bear a ligature, two pairs of small pressure forceps were left on the bleeding points for twenty-four hours.




Fig. 30.—Lupus of the Vulva (about natural size).

Tumour removed in the case of C. W., given in the text. The inner aspect is shown. The line by which it was attached is seen at AB. C points to the whitish area corresponding to the inner aspect of the nympha.

The patient did quite well, and left the hospital in about three weeks' time. The hypertrophy of the labia majora remained as before.

The patient came up to the hospital regularly to see me; and although the labia majora remained hypertrophied as before, there were no ulcerations seen till the beginning of January 1889. There was then found to be a lump the size of an almond lying beneath the mucous membrane, just inside the vulva, on the right side. It was hard, and the surface was red and ulcerated.

The patient was readmitted; and after trying the application of iodoform to the patch without benefit, she was put under ether, and the lump pinched up and dissected out with Paquelin's cautery.

The wound healed up soundly, and the patient went home. I have seen her frequently since; the labia majora remain in the hypertrophied condition already described, and up to about the end of May, when I last saw her, there had been no recurrence; but I think this may be expected, sooner or later.

Remarks.—The large growth appears to be made up of an enormously hypertrophied labium minus, as suggested by the whitish inner surface, and the pigmented outer surface. Sections of it show that it is chiefly made up of fibrous tissue.

The tendency of the disease in this case is towards hypertrophy combined with ulceration, and there has been recurrence after an apparently complete removal of the hypertrophied and ulcerated tissue. The general health of the patient seems to have been almost, or completely, unaffected by the local disease.

The late Dr. Matthews Duncan saw this specimen, and was of opinion that the case was one of lupus, and that the specimen had no resemblance to any syphilitic growth.

#### CARCINOMA.

Carcinoma of the vulva is much rarer than carcinoma of the cervix; its frequency would rather correspond to that of primary carcinoma of the body of the uterus. It may originate at any part of the vulva. In the following case it began round the urethral orifice:—

E. D., age 49, married, three children, the last fifteen years ago, came to the London Hospital complaining of a lump in the private parts varying in size at different times, also of difficulty and pain in micturition, and some constant pain in the left side; she had had these symptoms for seven weeks.

The catamenia ceased two years ago, but for the last two years she has had a yellow discharge streaked with blood.

Has been getting thinner lately.

On examination, in the situation of the urethral orifice there was an irregularly-shaped ulcerated cavity, admitting the tip of the finger. The walls of the cavity were formed by hard tissue, and induration extended up the anterior vaginal wall a distance of two inches. The surface of the cavity bled easily on touching it. Two glands in both groins were enlarged.

As regards three other cases I can call to mind, the disease began in the fossa navicularis in one, and in the remaining two on a labium majus.

The patients are generally advanced in life; most of the cases I have seen have been in women over sixty. I recently operated on a woman eighty-four years of age with epithelioma of the vulva, sent to me by Dr. Alexander Grant. The operation included removing two enlarged lymphatic glands from the right groin. She did well after the operation.

Treatment.—When the disease is seen sufficiently early—that is to say, when it appears possible to take away the growth completely, with a fair margin of apparently healthy tissue—an operation should be advised. The glands in one groin or in both groins are generally involved, but this does not contra-indicate operation, if it appears feasible to remove the affected glands at the same time. In removing the growth from the vulva I prefer to use Paquelin's cautery. In order to remove the glands from the groin an incision parallel to Poupart's ligament, three inches or more in length, is made over the affected glands, and these are dissected out, taking away a fair amount of the fat surrounding them at the same time. A few vessels generally require tying with fine catgut, and the incision is closed with silkworm-gut sutures.

Gangrene of the vulva is a rare complication of acute specific fevers in children; and in adults, is a rare complication of puerperal fever.

Fibroid tumours and fibroid polypi of the vulva are described, but are exceedingly rare.

#### PRURITUS OF THE VULVA.

By this is meant itching of the external genitals.

Etiology.—Sometimes no cause can be discovered to account for it; more frequently it occurs in association with some one of the following conditions:—

- 1. Pregnancy.
- 2. Cancer.
- 3. Vulvitis and vaginitis, particularly the early and late stages of these affections. Even if there was no vulvitis to begin with, scratching the parts, from which, as a rule, the patient cannot refrain, will soon set it up.
  - 4. Diabetes.
  - 5. Pediculi pubis.
  - 6. Thread-worms.

Sometimes it only occurs at the menstrual periods. Again, at the menopause it is not an uncommon symptom.

The itching is usually worse at night, and it may prevent the patient sleeping.

Treatment.—We try to remove any morbid condition that may be present—for example, vulvitis or vaginitis, pediculi pubis; when no local cause exists to account for the pruritus, or when it is due to some constitutional state, we have only to rely on empirical treatment. This consists partly of internal remedies, and partly of local applications.

The internal remedies that should be tried are:—Bromide of potassium in scruple doses, and belladonna, best given as recommended by Dr. West in the form of pill with camphor (Ext. belladonnæ, gr. ½, camphor., gr. iij., M. ft. pil. j.); if dryness of the throat occurs, the dose of the belladonna is reduced.

Local applications.—Some one of the following may be tried:—

- 1. Chloroform and olive oil (3j. to 3j.).
- 2. A lotion containing acetate of morphia (gr. ij. to 3j.).
- 3. Borax lotion (9j. to 3j.).
- 4. Lead and opium lotion (4 grains of extract of opium to 3j. of dilute lead lotion).
- 5. Glycerine and belladonna (9j. of extract of belladonna to 3j. of glycerine).
  - 6. Carbolic acid lotion (1 to 60).
  - 7. Liq. carbonis detergens (3ij. to 3iij., to Oj. water).
- 8. A somewhat messy application, but often an effectual one, is Fuller's earth made into a paste, and applied to the vulva.

The preparation used should be applied to the whole of the vulva frequently, or a piece of lint soaked in the application may be placed between the labia. In the few cases where I have tried cocaine for pruritus the result has been disappointing.

Other applications that may be tried are :-

NR Sod. Bibor., \( \)ij.
Ol. Menth. Pip., \( \)ji.
Ft. pulv. \( \)j. to Oj. warm water for bathing the parts.

Or-

R Menthol, 3j. Ol. Olivæ ad 3j.

Or-

R Acid. Salicyl., 9j.

Zinc. Oxid.
Pulv. Amyli
Vaseline, 3ij.
M.

Or-

R Calamin, zjss.
Ol. Amygdal., zij.
Aq. Calcis ad zviij.
Ft. lotio.

The pruritus of diabetes is best relieved, however, by placing the patient on a strict diet, and giving her a pill containing codeine (gr.  $\frac{1}{4}$  to gr.  $\frac{1}{2}$ ) twice daily.

#### CHAPTER VI.

## RUPTURE OF THE PERINEUM.

The inevitable laceration.—In every first labour there is a tear of the posterior part of the vaginal orifice, that is to say, there is a tear running into the fossa navicularis. This is the "inevitable" laceration. The tear may, of course, go much further, but in every first labour it must go as far as that.

Partial rupture of the perineum.—By this is commonly meant a rupture not involving the sphincter ani, as contrasted with

Complete rupture of the perineum.—By which is understood a tear involving the sphincter ani.

All the ruptures above mentioned are caused by want of proportion between the size of the fœtal head and the size of the vaginal *orifice* at the moment the head is born.

The rupture to be next considered—central rupture of the perineum—is caused by want of proportion between the size of the fœtal head and the size of the vaginal canal. In a complete central rupture there would be a communication between the canal of the vagina, say at the point a, and the perineum, say at b. A probe passed in at b would be felt in the vagina at a (Fig. 31).

The rupture, however, may not be complete, and the cause acting may produce merely:—

(1) A tear in the vaginal mucous membrane in the neighbourhood of a;

Or (2) a tear of the skin of the perineum in the neighbourhood of b;

Or (3) a tear of the tissues of the perineal body between a and b, there being neither laceration of the vaginal mucous membrane, nor of the perineal skin.

From what has been said, it will be clear that, in point of time, central rupture of the perineum occurs at an earlier stage of labour than ordinary rupture, for the former occurs while the head is still in the vaginal canal, while the latter does not occur till the head is passing the vaginal orifice.




Fig. 31.—Diagrammatic antero-posterior section of the perineal body, to illustrate central rupture of the perineum.

a. Vaginal mucous membrane. b. Perineum. c. Rectal mucous membrane.

Further, the conditions causing both central rupture and ordinary rupture may be present in the same case, and there will first be produced central rupture, and when the head reaches the vaginal orifice, an ordinary rupture also, running back into the central one. Unless the case has been watched throughout, we shall not then be able to know whether, or not, a central rupture preceded the ordinary rupture.

Treatment.—The laceration, if of any extent, should be sewn up immediately labour is over. A pad of Gamgee tissue, wrung out of perchloride lotion (1-1000), should be passed into the vagina to keep back the discharge while the

raw surfaces are being united. It is best that the patient should lie on her back in the lithotomy position with the legs separated; but the ordinary position on the left side will do if there is an assistant to draw the right buttock upwards, and so expose the raw surface. The raw surfaces should be washed with carbolic acid (1-40), all clots par-




FIG. 32.—To illustrate immediate perineorrhaphy (Thorburn).

ticularly being cleared away. I prefer to use an ordinary well-curved surgical needle held in a needle-holder, and to use fishing gut for the sutures. It is an advantage to use a needle-holder having a catch at the handles, like Wells's artery forceps. Another simple method of passing the sutures is to use a nearly straight needle in a handle, as recommended by the late Dr. Thorburn (Fig. 32).

Supposing that we have to deal with an incomplete rupture, i.e., one not involving the sphincter ani. A B C D is the raw surface (Fig. 33).

Pass the needle unthreaded from a point a quarter of an inch from the edge of the skin (as suture 1 is passed) on the




Fig. 33.—Diagrammatic representation of the raw surface in the preceding figure, to make the mode of passing the sutures clearer.'

patient's left side beneath the raw surface, bringing it out at the point 2 on the same side.

Thread it now with silver wire, or fishing gut, and draw back the needle.



FIG. 34.—STRAIGHT NEEDLE FOR PERINEORRHAPHY.

Now pass the needle similarly on the opposite side (the patient's right), bringing it out at 2,—the operation has reached this stage in Fig. 32; then thread the free end of the suture hanging from 2 (on the patient's left), and draw back the needle. There will then be a suture in the position 1, 2, 2, 1, as seen in Fig. 33.

The other sutures are passed in a similar manner. It is important to bring the needle out exactly at the junction of the vaginal mucous membrane and the raw surface in passing the sutures, or, when tying them, some of the vaginal mucous membrane will be tucked into the wound. Two, or at most three, sutures are sufficient in stitching up the perineum immediately after labour, when there has only been an incomplete rupture. The raw surface always seems to indicate a much more extensive tear than has




Fig. 35.—Diagrammatic representation of rupture of the perineum, involving the sphincter ani.

a b c. Triangular aperture in front wall of rectum; b c correspond to the ends of the torn sphincter. A suture (the purse-string suture) has been passed, as explained in the text, embedded all the way in the tissue bounding the triangular interval.

really occurred. The sutures are tied, beginning with the posterior one, care being taken to bring the surfaces into contact without unnecessary tightness. The pad of Gamgee tissue is now withdrawn from the vagina.

The patient's knees are kept tied together for ten days or a fortnight. The sutures are taken out in a week's time.

If the laceration has involved the sphincter ani, in addition to the raw surface in the above figure, there will be a triangular aperture in the front wall of the rectum, the apex of the triangle being upwards, and the ends of the

torn sphincter one at each end of the base line (Fig. 35). In passing the first suture it should be passed embedded all the way in the tissue immediately bounding the triangular interval, so that, when it is tied, the three points of the triangle will be approximated, just as the mouth of a bag with a string running round it is closed by drawing on the string. This is sometimes known as the "purse-string" suture.




Fig. 36.—The suture tightened and twisted, the points a b c approximate so as to close the gap. The rest of the raw surface is then dealt with as in Fig. 33.

The triangular gap in the rectum may also be closed by stitching it from the apex downwards with interrupted catgut sutures. They are passed so that the knot of each suture lies in the rectum.

The remaining raw surface is then treated as already described.

# SECONDARY OPERATIONS FOR RUPTURE OF THE PERINEUM.

When extensive lacerations of the perineum have not been sutured directly after labour, or when, having been sutured, union has not occurred, it is desirable in certain cases to perform a plastic operation to restore the previous

condition of the parts.

Such an operation should not be done until sufficient time has elapsed since the rupture for the parts to have passed into a quiescent condition, say at least six weeks after labour.

The indications for operation are:—1. Incontinence of fæces, or flatus, owing to the rupture having involved the sphincter ani. 2. When it is desirable, on account of prolapse of the vagina, or uterus, for the patient to wear a vaginal pessary, such as the ordinary india-rubber ring.

If there has been an extensive rupture of the perineum, such a pessary will not stop in. The object of the operation in a case of this kind is to narrow the lower end of the vagina, so as to enable the patient to retain a pessary.

Preparation of the patient.—She should remain in bed for some days before the operation, and should take a mixture containing sufficient sulphate of magnesia to ensure thorough evacuation of the bowels. If there is any discharge from the vagina, this should be cured, or at all events checked, by suitable treatment. Two nights before the operation she should take a draught, such as:—


R. Mag. Sulph., gr. 160.Infus. Sennæ ad 3j.M. ft. haust., h.s.s.

On the morning of the operation a copious enema should be given.

The operation.—Three assistants are advisable, one to give the anæsthetic, and two to help the operator. The patient is secured in the lithotomy position by means of Clover's crutch (Fig. 37).

The vagina is now syringed out with carbolic water, I in 40. An assistant stands at each side of the patient.

The next step is to mark out the surface to be made raw. The shape of this depends on the exact nature of the case. If the rupture has not involved the sphincter ani, Fig. 38




. Fig. 37.—CLOVER'S CRUTCH.

The leg pieces fasten round the legs just below the knees. The long strap passes under the patient's neck and under *one* arm, and is adjusted so as to flex the thighs on the abdomen to the required extent. The bar between the leg pieces consists of an outer tube, and an inner rod sliding in it; the rod is drawn out of the tube till the knees are sufficiently separated, and is then fixed by the screw.

shows the shape and position of the surface to be freshened. The points B and C correspond to the posterior extremity of the labium minus on each side, and it is well to fix a pair of Spencer Wells's forceps on B and C, where the skin and mucous membrane meet, to serve as landmarks.

The point A is taken on the middle line of the posterior vaginal wall about an inch, or an inch and a half, up the vagina; a pair of Wells's forceps should be fixed on it also.

An incision, mucous membrane deep, is now made with a scalpel from A to B, and A to C, and also along the line B D C, just where the mucous membrane of the posterior vaginal wall meets the perineal skin. Beginning posteriorly, the mucous membrane is now dissected off the



area marked out. During the whole proceeding the parts are made tense by each assistant pulling towards himself the labium majus and adjoining skin on his side. Sometimes it is convenient to put a pair of Wells's forceps on the point D; traction on this pair of forceps helps to make the parts tense, and therefore facilitates the dissection. The sutures are inserted as in the figure, beginning behind. 1, 2, 3, 4, 5, are passed by means of a nearly straight or slightly curved needle in a handle. For these, either fishing gut, or silver wire, may be used. It is an advantage to stain the fishing gut a reddish colour, as it is then more easily seen. These

sutures should be completely buried beneath the raw surface. While they are being passed the fore-finger of the left hand is in the rectum as a guide. Two or three sutures should be passed as at 6 and 7 in the figure. These




FIG. 39.—NEEDLE-HOLDER.

are inserted by means of an ordinary needle in a needle-holder \* (Fig. 39), and the material used for them should

<sup>\*</sup> Hagedorn's needle-holder and needles are very convenient in this and similar operations.

be catgut, as they will not then require removal. Sutures 6 and 7 should also, if possible, be buried completely under the raw surface; it is important that they should enter at




FIG. 40.—HAGEDORN'S NEEDLE-HOLDER.

the very edge of the cut mucous membrane to avoid any of it being tucked in when they are tied. All blood and clots being carefully sponged away, sutures 6 and 7 are tied first,





FIG. 41.—HAGEDORN'S NEEDLES.

and then 1, 2, 3, 4, and 5 in numerical order. Before they are tied the screw of the crutch should be loosened, and the knees brought together. It is important not to tie the sutures too tight. The ends of the sutures should be left

long, say two inches. The patient's knees are tied together, and she is put to bed. No T-bandage should be used.

When rupture has involved the sphincter ani, the raw surface to be made is shown in Fig. 42.

In this figure A, B, and C are the same points as in Fig. 38. D and E correspond to the extremities of the torn sphincter. F indicates the apex of the triangular interval in the front of the rectum caused by the rupture of the sphincter. Incisions as before are made along the lines A B D and A C E; in addition, an incision is made from



F to D, and F to E. The mucous membrane is then dissected off the whole surface marked out. The chief difference from the former case is in passing the first suture—the "purse-string" suture. It is passed buried completely in the tissues. When it is tightened, the points E F D are brought into contact, just as a bag is closed by drawing on a string running round its neck.

Instead of using the "purse-string" suture, the side F E may be stitched to the side F D with interrupted catgut sutures, the knots being tied so as to lie in the rectum.

After-treatment.—The patient should lie on her side,

not on her back, so that any discharge may escape more readily. The vagina may be gently douched twice daily with carbolic water (1-60), if we have a nurse who can be trusted to do this without any traction on the parts. The catheter is passed two or three times in the twenty-four hours. The bowels should not be confined, as used to be the custom. On the contrary, it is well to give enough of some simple aperient to keep them acting easily from the first. The diet should be very light; I prefer not to allow meat or fish till the end of the first week.




Fig. 43.—Lines of incision in Mr. Tait's operation for ruptured perineum (Lawson Tait).

Mr. Lawson Tait's Operation for Rupture of the Perineum.

The distinguishing features of this operation are :-

- 1. The raw surface is made by splitting the recto-vaginal septum with scissors, and by dissecting laterally as described below; no tissue is removed.
- 2. The sutures are not passed through the skin at all, their points of entry and final exit being in the raw surface made, and about  $\frac{1}{8}$  inch internal to the edge of the skin on each side.

3. The sutures are not completely buried, but appear in the bottom of the wound to the extent shown by the black lines below B and C in Fig. 44.

The patient being in the lithotomy position, the rectovaginal septum is split by scissors entered at the point F at one side, and carried across to the F at the other side. From the point F to C on each side an incision running forwards and a little outwards is also made with the scissors: the length of it is about an inch. Similarly, incisions are made about ½ inch long, running backwards and outwards from F to E on each side. Fig. 43 shows the lines of incision.

The shallow trenches resulting from the original incisions are gradually deepened by the scissors (working in a direction at right angles to the general surface of the perineum), till the points F can be raised with pressure forceps upwards, and approximated in the middle line; a triangular flap is thus turned upwards and inwards on each side. Similarly the points D are drawn downwards (the patient being in the lithotomy position) and inwards, so as also to lie close together in the middle line.

The resulting raw surface is shown in Fig. 44. A B C D is the boundary of it towards the vagina. The sides in the figure are the edges of the skin. Fishing-gut sutures are passed either with a curved needle in a handle, or with an ordinary surgical needle held in a needle-holder. The dotted lines indicate the extent to which the sutures are buried beneath the raw surface; when the sutures are tied, Mr. Tait claims that a sort of flap valve is formed towards the vagina and rectum respectively. The sutures are left in for a fortnight.

The vagina is washed out daily with a weak antiseptic lotion.

I have now tried this operation in a large number of

cases, and have been well satisfied with it. Probably every one who operates at all frequently introduces some slight modifications in this and other operations.

I find it generally convenient to remove most of the vaginal flap dissected up. The piece removed is somewhat triangular. In this way a neater result has seemed to me to be obtained.

Also I think it is decidedly desirable, in cases of rupture involving the sphincter ani, to stitch the gap in the rectum




Fig. 44.—Raw surface and position of sutures in Mr. Tait's operation for ruptured perineum (Lawson Tait).

separately with interrupted catgut sutures, tied so that the knots lie in the rectum.

I find that students and some others have a great deal of difficulty in understanding the descriptions of the "flap-splitting" operation (Tait's operation). I have accordingly had some figures prepared, which I hope may make the matter clearer. I will therefore again describe the "flap-splitting" operation, with the modifications I find advantageous, and ask the reader to follow it with the coloured diagrams (Plates I. and II., Figs. 1, 2, 3, 4, 5, 6).

# The modified "flap-splitting" operation (Tait's operation).

The patient is supposed to be in the lithotomy position as before (Plate I., Fig. 1).

Pushing the point of the scissors in at B a depth of  $\frac{1}{2}$  inch or so, an incision is made in the recto-vaginal septum from B to A.

Similarly an incision is made from A to E at the junction of the skin and mucous membrane; this is actually about an inch or rather more in length. The same is done on the opposite side from B to F. Also incisions are made from C to G, and from D to H. These are actually rather shorter than the anterior incisions A E and B F, though they are shown of the same length in Fig. 1.

Now a pair of Wells's small forceps is placed on the corner A and on the corner B, and these points are gradually raised by dissecting till two somewhat triangular flaps can be turned forwards, as seen in Fig. 2. Similarly a pair of Wells's forceps is placed on the corner C, and one on the corner D (Fig. 1), and these are gradually raised and turned backwards towards the rectum, as shown in Fig. 2.

The dissection is continued till A and B are turned so far forwards that, owing to the laxity of the vaginal flap, the border E A B F is actually somewhat convex. Also, by continuing the dissection, the corners C and D are brought to lie as in Fig. 3. I now cut away the greater part of the anterior flap along the dotted white line E K F (Fig. 3).

The next point is to stitch up the triangular gap C J D. The method of passing the sutures is shown in Fig. 5. In Fig. 6 the gap in the rectum is seen closed, and one of the stitches (catgut) to bring the cut edges of the vaginal mucous membrane together is seen in position. Also in Fig. 6 the deep sutures (of fishing gut) are seen in position.





Plate 1.









Plate 2.





When these are tied, the edge E G is brought into apposition with the edge F H.

The vaginal and rectal sutures need not be removed: the deep fishing-gut sutures are taken out in ten days or a fortnight.

The union is often so complete that the knots are buried in new tissue, and it is necessary to break through about  $\frac{1}{16}$  inch of intervening tissue before they can be reached.

### CHAPTER VII.

# DISEASES OF THE VAGINA.

#### VAGINITIS.

INFLAMMATION of the vagina may be *local* or *constitutional*— that is to say, it may be due to strictly local causes, or it may occur as a complication, a mere subordinate part, of some condition affecting the whole system.

#### Varieties.

Vaginitis may be:-

Specific, i.e., due to gonorrhœa; or

Simple, i.e., not gonorrhoeal.

Simple vaginitis may be due to any of the following local causes:—

- 1. Mechanical injury produced by foreign bodies (hairpins, pieces of sponge, ill-fitting pessaries, etc.); excessive coitus; difficult labour—in this case more or less sloughing, with perhaps perforation into the bladder or rectum, is not uncommon.
- 2. Irritating discharges, e.g., the discharges in cases of uterine cancer, or the lochia after delivery.
  - 3. Vaginal injections used too hot or too strong.

As examples of *constitutional states* in which vaginitis is likely to arise as a subordinate complication may be mentioned:—*diabetes*, *alcoholism*, *specific fevers*, *diphtheria*.

Further, in *old age* vaginitis is liable to occur, usually without other apparent cause, so that we speak of a "senile vaginitis."

Again, during pregnancy and menstruation, vaginitis is

likely to be set up by slight causes, which often cannot be discovered.

#### ACUTE VAGINITIS.

Symptoms.—The patient complains of :—

A sense of heat and throbbing in the vagina.

Pain in passing water, and frequent desire to pass it.

Often also of pain in walking, if there is co-existing vulvitis.

A greenish-yellow discharge.

A sense of general malaise.

Physical signs.—If there is vulvitis as well (as is usually the case), there will be the appearances described under Acute Vulvitis.

Great swelling, redness, and perhaps excoriation of the external parts.

Tenderness on attempting to pass the finger into the vagina.

If a small speculum can be introduced, the vaginal mucous membrane will be seen to be red and raw-looking, and secreting a greenish-yellow discharge, perhaps slightly blood-stained. The vaginal mucous membrane feels hot, and is greatly swollen, but its rugæ are not obliterated. It bleeds easily on examination. The urethra will be found swollen, and pressure along it will probably cause pus to exude from the meatus.

The thermometer may show slight elevation of temperature.

Such are the characteristics of acute vaginitis, whether simple or specific (gonorrhœal), and it is important to notice that we cannot yet distinguish by mere clinical examination between simple and specific vaginitis.

It is, however, possible for an expert in bacteriology to distinguish between them.

On this point, viz., as to the possibility of identifying any given discharge as gonorrhoeal, I have consulted Dr. Klein, who has kindly sent me his opinion, with permission to publish it.

19 EARL'S COURT SQUARE, S.W. June 2nd, 1897.

DEAR DR. LEWERS,-

Within recent years the gonococcus has by new methods become more amenable to differential cultivation, and it is possible for an expert to diagnose and identify the gonococcus more readily than was possible some years ago. The appearances of the growth on the new media are very characteristic, and by these the gonococcus can be distinguished from other non-gonorrheal cocci.

Very sincerely yours, E. KLEIN.

An acute vaginitis runs its course in ten days or a fortnight, and either gets well or lapses into a chronic stage.

# CHRONIC VAGINITIS.

Vaginitis may be chronic from the first, as, for example, in the senile variety; or chronic vaginitis may be the sequel of an acute attack.

Symptoms.—There may be a slight itching or sense of heat in the vagina, and slight smarting on passing water; often, however, the patient complains of little except that she has a yellow discharge.

Signs.—The speculum shows the mucous membrane to be redder than normal, and we see the discharge. In old people the mucous membrane is often seen to have become smooth, the rugæ having been obliterated. There is little or no tenderness.

Variation in the distribution of vaginitis.—Whether the inflammation be acute or chronic, it will be found sometimes that the whole vagina is affected, sometimes only a part of it. For instance, the inflammation may affect only the lower half, or only the upper half of the vagina; or, again, the summits of the ridges may be inflamed, while the depressions between them escape. Sometimes the appearance presented is that of red spots dotted over the surface, separated from one another by pale areas, "spotty vaginitis."

Granular vaginitis is the name given to that variety of vaginitis in which we find the surface of the vagina studded with hemispherical elevations, the size of a pin's head. It may result from any of the other forms of vaginitis, but is rarely seen apart from pregnancy (Thomas). The elevations referred to are said to be hypertrophied mucous glands.

Ulcerous vaginitis.—Sometimes we find ulcers on the vaginal walls in cases where we have no reason for suspecting the presence of syphilis, as in the following instance:—

J. R., single, age 25, came to the London Hospital, August 1886, complaining of pain across the lower abdomen, a sense of soreness in the vagina, and of a whitish-yellow discharge. There was no history of syphilis, nor any sign of it. On examination the hymen was found to be torn. Through the speculum two ulcers were seen at the upper end of the vagina, one on the anterior and one on the posterior wall; these were about ½ inch by ¾ inch in extent, and there were one or two smaller ulcers. She was given lead lotion as an injection, and general tonic treatment. November 3rd, ulcers much as before described; she was given 5 grs. of iodide of potassium thrice daily for the next four months. In April 1887 the following note was made: "There is a semicircular ulcer, depressed 1 inch, with a sharply defined edge on the anterior lip of the cervix, and a similar but rather triangular ulcer on the right lateral fornix, its largest side about an inch long. The base of this ulcer is dry and yellowish." The iodide of potassium was omitted, and she was given an injection of glycerine of subacetate of lead. A month later the patches had healed, the situation they had occupied being puckered and a little depressed.

Complications.—An acute vaginitis, particularly if due to gonorrhoea, is likely to be complicated by urethritis and cystitis; and the inflammation may also extend from the vagina to the cervix, thence to the body of the uterus, and so along the Fallopian tubes to the pelvic peritoneum; thus we may have cervical endometritis, endometritis of the body of the uterus, salpingitis, and pelvic peritonitis, all resulting from an acute vaginitis. There will also, almost certainly, be vulvitis, often also abscess of the ducts of Bartholin's glands. Further, buboes may occur on account of the vulvitis.

In cases where the inflammation has spread to the pelvic peritoneum, the fimbriated ends of the Fallopian tubes will almost certainly be sealed up, so that no trace of the fimbriæ remains, the outer end of each tube having become permanently adherent to the corresponding ovary. When this happens on both sides (as it usually does when it happens at all), an incurable sterility is necessarily produced.

Another complication of gonorrhoeal vaginitis is arthritis. It is a very rare complication. I have seen one case, sent to me by Dr. Downes, of Hornsey, in which the wrist was the joint affected.

Treatment.—In acute cases hot hip baths and rest in bed for the first few days, with vaginal injections of warm water, or barley water, with some laudanum (3j. to Oj.), may be advised. A smart saline purge will also be useful. As the acute stage subsides, astringent lotions may be used, beginning with glycerine of subacetate of lead (3ss. to Oj. water), followed later by alum, tannic acid, or sulphate of zinc (3j. to Oj. water). An injection of sulpho-carbolate of zinc (3ij. to Oj.) is also useful. If there is pain on micturition, this may be relieved by drinking barley water, and taking a mixture containing tincture of hyoscyamus and

bicarbonate of potash. In chronic cases we may proceed with the astringent applications enumerated from the first. In many cases after apparent cure the case relapses when treatment has been left off; again, many cases do not improve in spite of suitable treatment; this is often because the patient does not use the vaginal douche in an effectual manner. Vaginal douches are best given by means of the




Fig. 45.—Hydrostatic Douche Tin for giving Vaginal Injections.

When the patient is to use it herself, it is necessary to have a tap close to the glass vaginal tube.

While using an injection the patient should lie on her back with the hips somewhat elevated; if expense be no object, the ladies' bed-bath may be placed under her; otherwise an ordinary bed-pan may be used. The quantity of fluid injected should be at least a quart.

hydrostatic douche tin, fitted with india-rubber tubing and a glass vaginal pipe (see Fig. 45).

Another very effectual method of treatment, even from the first, is to douche with weak perchloride lotion (1-5000),

followed immediately by a douche of permanganate of zinc (gr.  $2\frac{1}{2}$ -5 to the pint).

It is often an advantage in subacute or chronic cases to make applications to the vaginal mucous membrane, once or twice a week, through a Fergusson's speculum.

I have found the following method useful:-

Pass Fergusson's speculum, and pour in warm corrosive sublimate solution (1-1000). Manipulate the speculum (by partially withdrawing it and passing it completely again) so that the fluid comes in contact with the whole surface. Pour off the sublimate solution, and apply a solution of sulphate of copper (10 per cent.) similarly. The patient uses vaginal injections herself in addition to this treatment.

When there is much soreness or irritation present, she should be directed, after each injection, to inject, by means of a ball syringe with a long nozzle, about 3j. of water in which oxychloride of bismuth is suspended.

#### VAGINISMUS.

This name is given to a condition where painful spasm of the muscles round the vaginal orifice is set up by attempts at sexual intercourse. Coitus may thus be either rendered difficult, or impossible. Sometimes merely touching the vulva is sufficient to cause the spasm. In bad cases the spasm extends to the muscles of the body generally. Vaginismus may be either *primary* or *secondary*.

In primary vaginismus no local cause can be discovered to account for the condition. The disease is, in fact, a neurosis.

In secondary vaginismus some local cause is present.

On examination we find some one of the following conditions:—

Fissures in the neighbourhood of the vaginal orifice, e.g., in the fossa navicularis.

An inflammation of part of the hymen.

A urethral caruncle.

Vulvitis, or vaginitis, or both.

Little ulcers round the vaginal orifice.

All these conditions may occur without causing vaginismus. It is only when the pain caused by touching the diseased part is so severe as to set up spasm of the muscles round the vaginal orifice, that we call the condition vaginismus.

The subjects of it are usually of an emotional temperament.

Treatment.—In secondary vaginismus we endeavour to remove the cause. Vulvitis, vaginitis, and urethral caruncle are to be treated as advised elsewhere. If there is a fissure, the patient should be placed under the influence of an anæsthetic, and the vaginal orifice forcibly dilated with the fingers, or by means of vaginal dilators. If little ulcers are present, they should be touched with the actual cautery; this may cure them, but in some cases they reappear. In primary vaginismus forcible dilatation may be tried, but with only a very moderate expectation of a cure resulting; in some cases even the degree of dilatation incidental to the birth of a child fails to cure. Sometimes primary vaginismus disappears spontaneously.

In all cases of vaginismus there is dyspareunia (pain on coitus); but dyspareunia is not, in all cases, accompanied by vaginismus.

I have seen two moderately severe cases of dyspareunia, accompanied by vaginismus, cured by dilating the vaginal orifice. In these two cases no cause for the dyspareunia, other than the vaginismus, could be discovered.

## TUMOURS OF THE VAGINA.

All tumours originating in the vagina are rare, and we need do no more than mention that the following may be met with:—

Cysts, usually found on the anterior wall of the vagina, and about the size of a walnut. They contain a clear yellowish fluid. Sometimes the fluid is of a brownish colour, suggesting that the origin of the cyst may have been an extravasation of blood.

I once saw a case where a cyst of the anterior vaginal wall, about the size of a duck's egg, was present as a complication of pregnancy at rather more than the eighth month; it projected outside the vulva. As it seemed not unlikely that it might occasion some trouble at the confinement, it was decided to treat it at once by incision and the application of pure carbolic acid to the lining membrane of the cyst. The interesting point is that this trivial operation seemed to bring about labour, as this came on two days afterwards. Both mother and child did quite well. The case shows that even the slightest operations on the vagina during pregnancy should be postponed, at all events, to a late period, when, if labour should come on, the child would be likely to survive.

Fibroid tumours of the vagina.—These are very rare. I happen to have met with one case.\* The tumour was situated (as they generally are) on the anterior wall.

Primary malignant disease.—Either carcinoma or sarcoma may affect the vagina. More frequently, when there is malignant disease of the vagina, it is secondary to malignant disease of the uterus.

Abscess of the vagina. - Abscesses due to a suppurative

<sup>\*</sup> London Obstetrical Society's Transactions, vol. xxix.

pelvic cellulitis, or a suppurating hæmatocele, often open into the vagina (see case of S. J., in Chapter XIV., for example). But the course of events there is plain. Here it is only intended to mention abscesses that have not originated, as far as can be ascertained, from any of the ordinary conditions causing pelvic abscess.

The following case will illustrate what is meant:-

Sarah G., age 29, married twelve years, six children, the last born November 30th, 1888; five miscarriages, the last three years ago; was admitted to the London Hospital on January 30th, 1889, on account of a tumour in the vagina.

At the last confinement but one the child was stillborn, and at the last confinement only lived forty-eight hours. Chloroform was given on each occasion.

About four months ago she felt a lump about the size of an egg in the front passage; she says it came suddenly; she was standing up when she first noticed it; she experienced great pain at the time, and felt faint.

Her last confinement was an extremely difficult one; she was attended by Dr. Waller and Dr. Turtle, who found a lump in the passage obstructing the descent of the child.

Present state—Abdominal examination.—There is a hard body to be felt in the hypogastrium in the middle line about on a level with the brim of the pelvis. This was shown by subsequent examination to be the uterus.

Vaginal examination.—On inspection a convex smooth lump is seen just within the orifice of the vagina. It is found to lie beneath the mucous membrane of the posterior vaginal wall. The lump extends up almost as far as the finger can reach, forming a mass between the rectum and vagina at least 1½ inches thick. The lowest part of the tumour extends to within an inch of the vaginal orifice. Everywhere the vaginal mucous membrane over the mass has its normal colour.

The uterus can with difficulty be reached; it is very high up, and quite separate from the tumour, though pushed up by it. There are numerous cicatricial bands in the upper part of the vagina (no doubt due to the difficult delivery already referred to).

The lower end of the mass has some mobility, but the attached part

is fixed. The tumour appears to spring from the right side of the pelvis between the vagina and rectum, and it reaches almost to the left side.

As regards consistence the lump is hard in parts, and in parts softer.

The case was seen by many observers, among others by Dr. John Williams and Dr. Herman, and the general opinion was that the tumour was a malignant one.

She gained 6 lb. in weight between February 7th and March 21st.

On the latter date a trocar was inserted into the tumour (which was exactly in the same condition as first described), and about an ounce of

yellow pus, free from any smell, came out.

On March 25th.—A trocar was again inserted, and after two unsuccessful attempts, the abscess cavity was found. A director was now passed alongside of the canula, and a bistoury guided along it then made an opening large enough to admit the finger. Two yellow lumps about the size of half a walnut came out of the cavity besides a small amount of pus. The lumps had the consistence of recent blood-clot, but they were yellow. The cavity having been washed out with iodine water, a drainage tube about three to four inches long was inserted, the lower end of the tube being stitched to the edges of the opening.

The wall of the abscess was remarkably thick, so that some difficulty

was met with in actually reaching the cavity with the trocar.

The patient did quite well, and went out on April 27th. I examined her in November 1889; there was thickening in the position occupied by the abscess. She was quite well in herself.

An abscess may also form occasionally between the anterior vaginal wall and the urethra; and it may open into the urethra. I have mentioned a case of this kind in Chapter XVIII.

Treatment.—Vaginal cysts may sometimes be conveniently dissected out entire. Large vessels may be met with, particularly if the cyst is situated high up in the vagina. In dissecting out a cyst the size of a Tangerine orange from beneath the posterior vaginal wall, with some of its attachments high up near the cervix, I was obliged to leave several

pairs of Wells's forceps on bleeding points for forty-eight hours.

The case was sent to me by Dr. Reynolds, of Dulwich, and did quite well.

Vaginal cysts may also be treated by cutting out a piece of the cyst wall, and applying pure carbolic acid, or tincture of iodine, to the interior of the cyst.

Fibroid tumours may be dissected out entire, as in my case referred to above.

When there is reason to think that malignant growths found in the vagina are the primary and only ones present, and their situation is such as to admit of their removal without much risk, this may be done.

#### CHAPTER VIII.

PROLAPSE OF THE VAGINA. PROLAPSE OF THE UTERUS.

Prolapse of the vaginal walls.—This is an exceedingly common condition. Either the anterior vaginal wall alone, or the posterior vaginal wall alone, may prolapse; or both may come down simultaneously.

Prolapse of the anterior vaginal wall is called cystocele, because the attachment of the bladder to the anterior vaginal wall is so intimate, that when the latter prolapses the bladder must descend at the same time.

Prolapse of the posterior vaginal wall may or may not be rectocele, because the attachment between the posterior vaginal wall and the rectum is not so intimate as that between the anterior vaginal wall and the bladder; it is possible for the posterior vaginal wall to prolapse without carrying a pouch of the rectum with it.

When the vaginal walls prolapse, more or less *secondary* prolapse of the uterus generally occurs as a consequence, as will be explained further on.

Prolapse of the uterus. Procidentia.—By this is meant that the uterus occupies a lower position than natural.

As regards degree, when the uterus has partly, or wholly, passed the orifice of the vulva, the case is called one of "procidentia."

When the uterus, though lower than normal, still lies wholly within the vagina, the case is one simply of prolapse. In extreme cases of procidentia the whole uterus lies outside the body, being contained, as it were, in a bag formed by the inverted vagina. In such cases there is a mass, the size perhaps of a cocoanut, outside the vulva. The os uteri is seen somewhere near the middle of the mass, and the body of the uterus can be felt in the hinder half of it; often both ovaries can be made out as well (Figs. 46 and 48).




Fig. 46.—Complete Procidentia Uteri (Schroeder).

Note.—I. The retroflexed position of the uterus; this position is the invariable rule, except when the uterus has been fixed in a position of anteflexion previously by adhesions. Practically the uterus is always retroflexed in cases of procidentia.

- 2. The complete inversion of the vagina.
- 3. The position of the bladder.
- 4. The position of Douglas's pouch.


Etiology of prolapse of the uterus.—Suppose we compare the peritoneal cavity to an oval elastic bag, with the uterus attached to the wall at the lower part of the bag; then, on looking at the diagram (Fig. 47), it will be evident

that any force acting in the direction of the arrow will tend to cause descent of the uterus. Now, the force acting in the direction of the arrow may be

Increased weight of the uterus, due, for instance, to fibroid tumours in its walls; or it may be

Traction on the uterus from below, as when the vaginal walls are prolapsed, and tend to drag the uterus after them; or, again, force in the required direction may be produced by

Increase of the general intra-abdominal pressure, such as occurs in all acts of straining.



e couses will ac

Obviously, any of these causes will act to greater advantage if the tissues round the uterus are in the lax condition natural to them after labour.

Rupture of the perineum only facilitates prolapse; it is not an exciting cause. The late Dr. Matthews Duncan compared descent of the uterus in cases of prolapse to descent of the head in the second stage of labour, after the os is fully dilated. In the latter case an intact perineum does not prevent the birth of the head; time only is needed, if there are good pains, for the perineum to stretch sufficiently to allow the head to pass; and so it is in cases of prolapse of the uterus,—an intact, unruptured

perineum only delays the descent of the uterus; time is required for the perineum to stretch; but given this, it will do so, and the uterus pass out, just as the head does in labour: the motive power in the latter case being the uterine contractions; whereas, in cases of prolapse, it is some one of the three causes given above, as producing force in the direction of the arrow in the diagram.

Etiology of prolapse of the vaginal walls.—Similarly, rupture of the perineum only facilitates prolapse of the vaginal walls, the only essential exciting cause of their prolapse being increased intra-abdominal pressure, as, for example, during muscular efforts. Other predisposing causes of prolapse of the vagina are, in the case of the anterior vaginal wall, an habitually full bladder, and, in the case of the posterior vaginal wall, an habitually overloaded rectum. Most cases of prolapse of the uterus are secondary to prolapse of the vaginal walls.

Symptoms and diagnosis.—The symptoms are often slight and unimportant; there is usually more or less "bearing down" and feeling of weight in the pelvis.

But in extreme cases, where the whole uterus and its appendages lie outside the body in the inverted vagina, the patient often complains of little, except the inconvenience of having a big lump between her legs. As the late Dr. Matthews Duncan pointed out, this fact (which any one can verify for himself, as the cases are common enough) makes it very hard for us to believe that grave symptoms can be caused by comparatively slight displacements of the uterus, for in this, the greatest of all displacements, symptoms are usually so slight.

Some difficulty in emptying the bladder is another symptom often met with.

The patient has not uncommonly found out for herself that she can manage better when she presses the lump up. It is remarkable that in cases of procidentia, contrary to what we should expect, menstruation is not usually excessive.

As regards diagnosis, this is usually at once obvious. When there is a moderate prolapse of the anterior vaginal wall, it might possibly be taken for a cyst or other tumour growing in the anterior vaginal wall; but any doubt can at




Fig. 48.—Complete Procidentia Uteri (Winckel).

The dotted line indicates the position of the body of the uterus. Note the somewhat crescentic ulceration on the vaginal wall posterior to the os uteri. The patient is lying on her left side.

once be cleared up by passing the sound into the bladder, and being able to feel the point at every part of the swelling, with only the wall of the bladder and vagina between the sound and the finger.

In cases of procidentia, the presence of the os uteri makes the nature of the case at once evident. Often there are extensive ulcerations on the inverted vaginal walls from the friction against the clothes (Fig. 48).

On passing the sound into the uterus it usually passes a good deal farther than the normal distance, often four to four and a half inches, or more. The elongation is partly due to hypertrophy of the uterus, but most of it is due to stretching.

The usual order of events in cases of procidentia has been, first, prolapse of the vaginal walls, which causes traction on the cervix at the vaginal reflection. Now the uterus does not descend freely under the influence of this traction; it is for a time held back by the broad ligaments, and utero-sacral ligaments. In time, however, these stretch to an extent sufficient to allow the whole uterus and its appendages to lie outside the vulva. The only limit to the descent of the uterus is the degree to which the walls of the inverted vagina will stretch.

On this factor it depends whether, in cases of procidentia, the whole uterus lies outside, or only part of it; if the vaginal walls are short, so that, when completely inverted, the vagina forms a bag of little depth, and the uterus is large, the sound passing four and a half inches or more, then there will not be room for the whole uterus in the bag; if, on the other hand, the vaginal walls are long, or the uterus small, the bag formed by the inverted vagina will contain the whole uterus.

Treatment.—In slight cases, either of prolapse of the uterus, or of prolapse of the vaginal walls, a ring pessary of suitable size is all that is needed. In bad cases, such as cases of procidentia, a ring is not usually retained. The condition that must be satisfied for a ring pessary to suit the case is, that the vagina be narrower below than it is above; if either on account of rupture of the perineum, or by stretching of an unruptured perineum, it has come about

that the vagina is as capacious below as it is above, it practically forms a cylindrical cavity, like a cylindrical jar open at the bottom, and a ring that will just go easily into the top has nothing to prevent it falling out at the bottom. It will be noticed, when a ring is being inserted, that it is compressed, so as to pass it by artifice through the narrow vaginal orifice; now, so long as the vagina is normally capacious above and narrow below, force has to be exercised to make it come out, force that either compresses the ring or dilates the orifice; if no such force is in opera-




Fig. 49.—Ring pessary, made of watch spring covered with indiarubber. In the figure it is seen compressed between the finger and thumb, as when about to be introduced. It is important to keep it compressed till it has passed the orifice of the vagina, and also to pass it well backwards. A ring pessary may be left in for three months at a time, the patient using douches night and morning.

tion, or if the force in operation is insufficient, the ring is retained, and keeps up the prolapsed parts. Hodge's pessary may be used for prolapse in the same class of cases as the ring, but it generally possesses no advantages over the ring.

In some cases of prolapse of the vaginal walls, especially in old women, the cradle pessary (invented originally by

Graily Hewitt for anteflexion) is more suitable than the ring pessary or Hodge's pessary.

For those cases in which a ring pessary or Hodge's pessary is useless, Zwanke's pessary, sometimes known as the "butterfly" pessary, may be tried. It is well not to use it if a Hodge's pessary or a ring pessary will suit the case, because its use is not free from risk. It must be taken out daily by the patient. If this is not done, ulceration of the vaginal walls is not unlikely to occur as a result of pressure. The ulceration may be so extensive as to produce a vesico-vaginal fistula. Still, with proper precautions,




Fig. 50.—Cup and stem pessary (vulcanite), with bands for attaching to the waist-belt, seen in the next figure.

The patient takes out this pessary at bedtime, and washes it, replacing it in the morning when she gets up.

Zwanke's pessary is certainly capable of relieving many cases where Hodge's pessary and the ring pessary have failed.

In still more severe cases of prolapse, where neither Hodge's pessary, nor a ring pessary, nor Zwanke's pessary is efficacious, the patient has a choice of two things—either she may use a cup and stem pessary (Fig. 50), which takes its support from a waist-belt, or she may have some operation done. The mildest operative procedure is an operation to restore the original shape of the vagina—that is to say, to make it once more narrow below. When this

has been done, a ring pessary or Hodge's pessary will stop in.

No plastic operation will cure cases of procidentia; no matter how complete the success of the operation may appear at the time, unless the patient wears a pessary, the displacement will most probably return as badly as ever. If,




FIG. 51.—The waist-belt, with the bands seen in the preceding figure fastened to it. Two fasten behind, and two in front; only the last two can be seen in the figure.

however, she wears a ring, a permanent condition of comfort is obtained.

I have some patients with procidentia, treated in this way more than ten years ago, that have remained comfortable up to the present time.

The best plastic operation of the kind referred to is that known as *posterior colporrhaphy*. The patient being in the lithotomy position, a triangular piece of mucous membrane

is removed from the posterior vaginal wall. The base of the triangle is, of course, at the junction of the posterior vaginal wall with the skin of the perineum, and the apex is high up the posterior vaginal wall, not far from the cervix. Stitches are then passed so that the raw surface A B D is in apposition with the raw surface A C D. The effect will clearly be a considerable narrowing of the lower half of the vagina.




Fig. 52.—Diagrammatic representation of raw surface to be made on the posterior vaginal wall in the operation of posterior colporrhaphy.

In some cases where there is a very considerable prolapse of the anterior vaginal wall, and little or no prolapse of the posterior wall, the area of the prolapsed wall will usually need to be diminished by a plastic operation before a satisfactory result can be obtained. For in such cases the prolapsed wall will project past any pessary, or through it, and so appear externally.

Alexander's Operation.—This operation, shortening

the round ligaments, has been recommended (1) for prolapse of the uterus, (2) for retroversion of the uterus.

As regards doing this for retroversion, we shall see in the chapter on Versions and Flexions that retroversion is only of importance (a) as a part of prolapse, and (b) when the retroverted uterus is incarcerated in Douglas's pouch.

Both these conditions can generally be remedied by the use of a suitable pessary (usually the watch-spring ring), the uterus being first replaced by the sound. If a vaginal pessary will not stop in, on account of wideness of the lower part of the vagina, this can be remedied by artificially narrowing the lower part of the vagina by a plastic operation.

As regards Alexander's operation for prolapse of the interus. Take, for the sake of argument, an ordinary case of procidentia uteri, where there is a large mass outside the vulva —in such a case the sound passes four or five inches. The fundus of the uterus is no doubt to a greater or less degree below its normal level in the pelvis, for it may be outside the vulva in the inverted vagina. Still, supposing the fundus of the uterus to be pulled up to its normal position by shortening the round ligaments (drawing them up through the inguinal canal, fixing them there, and cutting off the superfluous portions), this will not alter the fact that the uterus is four or five inches long. Therefore, though the fundus may be at its proper level, the cervix will still be low down, and the vaginal walls will still probably protrude outside the vulva; the operation, therefore, is far from being a complete remedy for the condition.

The operation is, besides, one that is by no means free from danger. The conditions, for which the operation is suggested, have no tendency to shorten life, and at most make the patient uncomfortable; moreover, several harmless methods of treatment that will secure the patient's comfort are at our disposal. For these reasons I have never felt

justified in recommending a patient to have the operation performed, and have never done it. I myself know of two cases ending fatally, one in two or three days after the operation.

Hysteropexy.—This is an operation which may sometimes be advisable in cases of procidentia uteri. There are various methods of performing it. The essential principle is to suture the fundus uteri to the abdominal wall. The abdomen is opened in the middle line in the usual way. The fundus uteri is pulled up with a volsella, or with some similar instrument. A silk suture is then passed with a curved needle, held in a needle-holder, at the lower angle of the wound, as follows:-first through the rectus and parietal peritoneum of one side, then transversely through the fundus, rather nearer to the anterior than the posterior aspect of the uterus. Care is to be taken not to pass the needle so deeply as to enter the uterine cavity. After emerging from the uterus, the needle is then passed through the parietal peritoneum and rectus of the opposite side. Sometimes one suture is enough. If not, another may be passed near the first, and in a similar manner. The uterus being held in the desired position, the suture is tied and the ends cut short. The rest of the abdominal wound is closed in the usual way. It is important, in the case of a woman who may possibly become pregnant subsequently, to pass the suture or sutures rather on the anterior aspect of the fundus than on the posterior aspect. If the sutures are passed on the posterior aspect, and the patient should become pregnant, the anterior wall of the uterus may be prevented rising up as it should do in the course of the development natural in pregnancy; yet it nevertheless undergoes great hypertrophy. The result has been in some cases that the hypertrophied anterior wall of the uterus has formed a tumour, blocking up the entrance to the pelvis, and

producing an insuperable obstacle to the passage of the child. In some such cases Cæsarean section or Porro's operation has had to be performed.

When the operation is performed as here described, the ultimate result is to produce a ribbon-shaped adhesion some two inches long between the fundus and the abdominal wall. The suture or sutures have been found encysted in the middle of this adhesion.

A consideration of the condition of the parts in a marked case of procidentia will make it clear that hysteropexy alone will rarely be an absolute cure for the condition. Probably some form of colporrhaphy to narrow the lower part of the vagina will be necessary, in consequence of the relaxed and hypertrophied condition of the vaginal walls usually met with in cases of the kind.

TABLE SHOWING THE ETIOLOGY OF PROLAPSE OF THE UTERUS AND PROLAPSE OF THE VAGINAL WALLS.

## Prolapse of the uterus.

Exciting cause.—Force acting in direction A B in the diagram, page 146, which may be either:—

- 1. Increase of the general intra-abdominal pressure.
- 2. Traction from below, as by the prolapsing vaginal walls.
- 3. Increase of the weight of the uterus.

### Predisposing causes:-

Rupture of the perineum.

Relaxation of the soft parts by recent labour.

Absorption of fat from the soft parts in the pelvis in old age.

Laborious employments, chronic cough, etc.

## Prolapse of the vaginal walls.

Exciting cause.—Increase of the general intra-abdominal pressure.

Predisposing causes :-

Habitually full bladder (the presence of a small ovarian tumour acts similarly; see Chapter XVI. for an instance of this).

Habitually loaded rectum, in addition to the predisposing causes of prolapse of uterus given above.

### CHAPTER IX. ·

### DISEASES OF THE UTERUS.

#### INFLAMMATION.

Cervical endometritis. Corporeal endometritis.— Inflammation may affect the mucous membrane of the cervix only, when it is known as cervical endometritis; or it may affect the mucous membrane of the body of the uterus, and it is then called corporeal endometritis.

#### CERVICAL ENDOMETRITIS.

Etiology.—This disease is much commoner in women who have had children than in nulliparæ, the starting-point in most cases being injuries inflicted on the cervix during labour. In such cases the whole thickness of the cervix is necessarily more or less involved in the inflammation, whereas, in the cervical endometritis of nulliparæ, the inflammation is more strictly limited to the mucous membrane.

Predisposing causes are those which predispose to inflammation of mucous membranes in other situations, such as the strumous, rheumatic, or gouty constitution.

Exciting causes are :-

- 1. Injury, either during labour, or at some other time, by the use of instruments, the sound, tents, etc.
  - 2. Extension from vaginitis, simple or gonorrhœal. While there is no doubt that this extension often occurs,

it is very striking to observe how a purulent vaginitis will sometimes remain strictly limited to the vagina, the secretion in the os uteri remaining typically healthy; and this, although the vaginal portion is lying constantly in a pool of pus.

3. Cold, especially during a menstrual period.

Symptoms.—The only essential one is the presence of a discharge\* which attracts the patient's attention. There may be many other symptoms present in the case due to associated morbid conditions, but the only symptom caused by the cervical endometritis itself is a discharge.

Normally, the secretion of the cervical glands is clear and viscid, like unboiled white of egg; when there is cervical endometritis, this secretion changes its character, and becomes either (a) opaque and white, in cases of slight catarrh; or (b) opaque and yellow, if the inflammation is more severe.

**Diagnosis.**—This is made by passing a speculum, and observing the character of the secretion within the external os. Often some opaque-white viscid discharge will be seen in the vagina, when the secretion just within the external os is healthy, i.e., transparent. It seems to be the case that when the healthy secretion of the cervical glands lies in the vagina it becomes opaque. Our diagnosis of cervical endometritis rests on the kind of secretion actually within the external os. If this be opaque and white, there is catarrh of the cervical mucous membrane; if opaque and yellow, the inflammation is more severe.

If the thickness of the cervix is much involved in the

<sup>\*</sup> The term "leucorrhœa" is often used in describing this symptom. Etymologically, of course, it ought to mean a white discharge, but practically it has come to be used rather loosely, and to mean a white or yellow discharge. It is better, therefore, not to use it, but rather the English adjective describing the actual discharge.

inflammation, the finger notices increased size of the vaginal portion, and, in chronic cases, induration of it. Little prominences, the size of shot, are often to be felt not only immediately around the external os, but also on the vaginal portion at some distance from the os. Through the speculum these are seen to have a pearly-grey colour. They are little retention cysts, and are called *ovula Nabothi*; their origin will be considered further on.

Erosion of the cervix.—In many women the speculum shows a red, slightly depressed area, of greater or less extent, immediately round the external os. This was formerly called "ulceration" of the cervix, and was vigorously treated with various remedies. It must be admitted that such areas are deviations from the normal type, and that, in many cases at least, they may be considered as originating in an inflammation affecting the area occupied by the erosion, and at the same time affecting the whole cervical mucous membrane.

On the other hand, slight erosions are very commonly observed where we have the secretion within the external os quite transparent, and therefore healthy.

The character of this secretion must be taken as a test of their importance in any particular case. If we see a slight erosion, but yet have the secretion within the external os healthy, we may, if we like, regard the erosion as an abnormality, but it certainly is not one that requires any treatment. If, on the other hand, the secretion is morbid, opaque-white, or opaque-yellow (and in such cases the erosion is usually of greater extent than in the cases where the cervical secretion is healthy), we are justified in applying the local treatment about to be described, and such treatment must be directed, not only to the erosion, but to the whole of the cervical canal as far up as, but not beyond, the internal os.

Pathology of erosions.—Normally, the cervical canal is lined by a single layer of cubical epithelium (ciliated on the summits of the ridges, but not in the depressions between them). The epithelium maintains this character till within a line or two of the external os. The epithelium immediately round the external os is squamous and stratified. Further, it is necessary to remember that while there are numerous racemose glands opening into the cervical canal, there are no glands on the vaginal surface of the vaginal portion. When an erosion exists, however, we find that over the area occupied by it there is only a single layer of cubical epithelium, in place of the several layers of squamous epithelium normally found round the external os.

While opinions differ as to the origin of this single layer of epithelium found on erosions (some thinking that it is derived from the glandular epithelium of the cervix spreading down over the erosion, all the epithelium normally there having been shed; others believing it to be the deepest layer of the epithelium of the rete Malpighii, only the epithelium superficial to this having gone), there is no doubt that the first step towards the production of an erosion is the shedding of the squamous epithelium in the neighbourhood of the external os under the influence of an inflammatory process, which at the same time is affecting more or less of the cervical canal. Further, whatever its origin, the single layer of epithelium on erosions takes on a glandular function.

# Varieties of erosion.—These are :-

- 1. The simple.
- 2. The papillary or villous.
- 3. The follicular.
- 1. The simple is that variety where the surface of the erosion is only slightly granular.
  - 2. The papillary is that where the epithelium dips down

pretty deeply, so as to make the surface more or less villous (Fig. 53).




Fig. 53.—Papillary erosion (Schroeder).

3. If the entrance to the depressions between the papillæ becomes obliterated, closed cavities are formed; these may become distended by the secretion of the epithelium (which,




Fig. 54.—Follicular erosion (Schroeder).

as just now remarked, takes on glandular functions, and the retention cysts resulting may burst. There is then a follicular erosion (Fig. 54).

EVERSION OF THE CERVICAL MUCOUS MEMBRANE.

When the vaginal portion of the cervix has been deeply lacerated during labour, the lips of the cervix gape, and expose the cervical mucous membrane. The greatest degree of eversion is reached when there has been laceration on both sides of the cervix reaching up to the vaginal reflection. The exposed mucous membrane is red, and




Fig. 55.—Bilateral laceration of the cervix with eversion of the cervical mucous membrane (Schroeder).

the appearance presented is very similar to that of a rather extensive erosion. Its true nature will be recognised, first, by noticing with the finger that there has been laceration. Suppose, for example, that there has been a deep laceration on the left side (the most usual position, on account of the frequency of the first vertex presentation), then the finger, placed in the situation of the external os and adjoining part of the cervical canal, finds nothing between it and the vaginal wall on that side. Secondly, we may draw together the gaping lips of the cervix with two hooks, using

a Sims's speculum, and find that the apparent erosion has disappeared, if it was all eversion; but if there was erosion as well as eversion, then that the extent of the red surface exposed has greatly diminished.


The everted cervical mucous membrane is often in a state of chronic inflammation, and consequently in such cases there is a white or yellow discharge. There is no satisfactory proof that laceration of the cervix in itself causes any other symptom except this discharge.

Treatment.—First we have to decide that the case is one where treatment is justifiable; further, the patient's desire to be cured of the discharge ought to be pronounced before we undertake to treat what is, after all, except for the annoyance that the discharge may occasion, a very trivial ailment in itself; it is well also to warn her, before commencing a course of treatment, that it is quite possible we may not be able to cure her, though we may pretty certainly diminish the discharge for the time to a great extent; and, again, it must not be forgotten that after apparent cure a relapse often takes place. When active treatment has been decided on, we pass Fergusson's speculum, and then ascertain the exact direction of the cervical canal with the sound. A solution of sulphate of copper \* is then poured into the speculum, so as to cover the vaginal portion; next we pass through the fluid a Playfair's probe, thinly dressed with wool, several times up and down the cervical canal, not passing it farther than the internal os. The wool on the probe takes up some of the solution each time; the first time or two the copper solution thus carried up acts on the secretion in the cervical canal, coagulating it, and so rendering it easy to remove; afterwards any more of the solution

<sup>\*</sup> The strength of the solution recommended is 9ij. sulphate of copper to 3j. water.

introduced acts on the cervical mucous membrane laid bare by the coagulation and removal of the secretion previously in the canal. This is the best mode of applying local remedies. The ordinary method of exposing the cervix as before, then removing the secretion in the cervical canal with a series of Playfair's probes dressed with dry wool, and then applying some remedy, e.g., pure carbolic acid, or tincture of iodine, to the canal, is less satisfactory, because it is impossible by simple mechanical means to remove the viscid secretion completely, and to ensure the remedy reaching the mucous membrane.

Any one can prove this on the dead body, when, if the uterus is cut open, it will be found difficult, by rubbing the cervical canal with wool, to get away the tenacious mucus. Such applications as have been described should be made twice a week for a time, and the patient should use hot vaginal douches night and morning as well. If this treatment does not succeed in three or four weeks, we may try a rather more severe measure, for which the patient must remain in bed a few days. A stick of fused zinc sulphate and alum (equal



Absorbent wool is wrapped tightly round the terminal inch or so; the probe is then ready for use.

parts) is passed into the cervical canal, but no farther, and left there to melt, the upper part of the vagina being packed with tampons soaked in carbonate of soda to neutralise the acid solution as it comes away; in three hours' time (Matthews Duncan) the tampons are removed, and a vaginal douche given; if this treatment does not succeed, the case had better be set down as incurable, and the patient recommended to be content to keep the discharge from being very great by using hot douches containing some astringent, such as alum or tannic acid.

It may be added, that though cervical endometritis is regarded by some as a cause of sterility, it has probably little or no share in producing it. It is a common experience to find patients becoming pregnant in whom we know there is a cervical endometritis. Considering how often labour is the starting-point of the disease, we may take it that, if cervical endometritis caused sterility, it would be rather a rare thing to find a woman who had had more than one child.

### CORPOREAL ENDOMETRITIS.

Etiology.—Inflammation of the mucous membrane of the body of the uterus due to:—

Labour or abortion, especially if some part of the ovum or clots are left behind.

• Extension from inflammation, gonorrhoeal or simple, lower down.

Direct injury, e.g., by the sound, or other instrument, such as intra-uterine stem pessaries, or tents.

Intra-uterine growths.—Submucous fibroids, or mucous polypi.

Cold during menstruation.

Certain fevers.

Old age.

Diagnosis.—It is often impossible to say whether or not there is corporeal endometritis in any particular case. When really present, it causes a discharge, white or yellow, according to the severity of the inflammation, but we only see this as it comes from the external os, and, therefore, we cannot be sure that it does not come from the cavity of the cervix only.

Certain features in the case may, however, incline us to the opinion that there is corporeal endometritis. These are:—

- 1. A history of menorrhagia.
- 2. On bimanual examination finding the body of the uterus somewhat enlarged and tender.
- 3. Finding that the sound passed gently causes much more pain than usual, and that perhaps a blood-stained discharge follows its withdrawal.

All these, taken together, point to inflammation involving the mucous membrane of the body of the uterus.

In some few cases we can make the diagnosis certain by exploration of the cavity of the uterus with the finger, after, if necessary, dilating the cervix.

In the following cases there was no doubt of the presence of corporeal endometritis.

CASE I.—M. W., aged 35. Married sixteen years; twins a year after marriage; two miscarriages, the last eleven and a half years ago. Admitted to the hospital on account of menorrhagia of fifteen months' duration. She had been under my care in the out-patient department, but had obtained no relief. On examination the uterus was found to be enlarged, and the sound passed four inches. The uterus was freely movable. The vaginal portion of the cervix was healthy. The cervix was rapidly dilated under ether with Hegar's dilators. On passing the finger into the body of the uterus, soft, irregular projections, particularly extensive on the right side, were met with. These were thoroughly scraped away with a blunt-edged spoon, and tincture of iodine applied to the interior of the uterus. Portions

of the substance removed, on microscopical examination afterwards, showed a glandular structure; very large irregularly shaped cavities lined with columnar epithelium, in a single layer, were seen in the sections. The temperature after dilatation remained normal. I have seen this patient at intervals since; she is now regular, and does not lose too much.

This case is an example of that variety of endometritis called *villous* or *fungous endometritis*, localised hypertrophy of the uterine mucous membrane giving rise to irregular projections, or fungosities; hence the name. Some authorities refuse to recognise this condition as an endometritis, regarding the condition as one produced by new growth rather than by inflammation.

CASE II.—S. J., a Jewess, admitted to the London Hospital in labour, ineffectual attempts at delivery having been made outside. Had been in labour from half-past two on the afternoon of May 12th, 1886, till six p.m. on the evening of the 13th, when I saw her. The ring of the Bandl could be felt a hand's breadth above the pubes. Foetal heart not heard. Forceps having failed, delivery by perforation and cephalotripsy.

May 31st.—Ever since delivery there has been a lump to be felt reaching up to the umbilicus, at first of course taken to be the uterus, but latterly, as the patient has had more or less fever ever since the confinement, thought to be due, in part at least, to inflammatory exudation. To-day patient was examined on the couch, and it was then found that a large sound could be passed up to the highest point of the tumour in the abdomen, going in 5½ inches, thus showing the tumour to be the body of the uterus. The uterus was then washed out with carbolic lotion; the fluid that returned first was very foul.

June 13th.—To-day a yellow, leathery mass, having an intensely feetid odour, two inches long by one broad, was found hanging from the cervix; it was twisted off with cervix forceps.

June 24th.—The uterus still being much about the size it was on May 31st, and the patient's general condition unsatisfactory, fever still persisting, she was put under an anæsthetic, and the interior of the uterus thoroughly scraped with a Recamier's curette. Numerous yellow flakes, intensely offensive, about 10 of an inch in thickness,

were removed by this proceeding, perhaps altogether enough to fill two or three tablespoons.




FIG. 57.—RECAMIER'S CURETTE. is required for scraping.

FIG. 58. Useful when a blunt instrument Small serrated curette for scraping the interior of the uterus.

From this time she improved rapidly, her highest temperature on June 27th being 100°, and from that time on quite normal.

By July 14th, just three weeks after the scraping, the uterus had involuted to its ordinary size, and the sound passed the normal distance. The uterus was freely movable.

This case illustrates endometritis of the body of the uterus dating from labour, and also the effect that a morbid condition of the endometrium may have in delaying involution of the uterus. It is not certain whether the yellow, leathery, offensive masses referred to were entirely clots, or whether they may not perhaps have been partly sloughs from the interior of the uterus, due to damage received in the very long labour. The membranes had been ruptured, and the os fully dilated, at least eighteen hours.

The next case is an example of undoubted corporeal endometritis, due to a portion of a sponge tent having been left in the body of the uterus for more than three months:—

CASE III.—Mrs. X., the wife of a medical practitioner, had certain symptoms, which induced her husband to pass a sponge tent into the cervix on July 18th. This was left in for twelve hours; it was then withdrawn, and a second sponge tent was inserted; at the end of eighteen hours this "came away in pieces"; a third tent was then inserted; after it had been in place six hours "the pain was so great that it was withdrawn."

A few days after there was "a slight discharge of mucous matter." She used a vaginal douche of Condy's fluid twice daily, and went to the seaside on August 3rd.

About this time the vaginal discharge increased in quantity, and became distinctly purulent and offensive. Menstruation began on August 19th, and lasted a week, the patient losing twice as much as usual. The next began on September 14th, and lasted five days; it was also profuse. All this time the purulent, offensive discharge continued. Soon after this I was asked to see the case. The uterus was a little enlarged, and a little tender, and there was a purulent discharge escaping from the os uteri. I advised her to rest in bed and to have

hot vaginal douches several times a day to see if she would get well spontaneously. This was tried for two or three weeks without any improvement. Accordingly I dilated the cervix with Hegar's dilators till I could pass my finger into the uterus. I then felt something lying loose in the cavity of the body of the uterus. On removing it, I found this to be a piece of sponge, three-quarters of an inch long by about a quarter of an inch wide. The uterus was washed out in the usual way. I have since heard from her husband that she made an uninterrupted recovery, and that the discharge ceased entirely within a fortnight.

There seems to have been very little constitutional disturbance during the time the piece of sponge was in utero, but there was, I know, a slight elevation of temperature on one occasion at least. I consider the patient very fortunate to have survived; no less than three sponge tents had been used consecutively, and a piece of one left in utero three months; a good many patients so treated would have died of septicæmia.

It remains to say a word on the diagnosis of corporeal endometritis from primary malignant disease of the body of the uterus.

In the latter disease the patient is usually old, most cases being between fifty and sixty years of age.

Then the course of the disease will help us; corporeal endometritis runs a chronic course, and deterioration of the general health from accompanying menorrhagia is only very gradual, whereas in malignant disease there is rapid loss of strength and weight. Again, in malignant disease of the body of the uterus, severe pain is an early and marked symptom; whereas in corporeal endometritis, pain, if present, is much less severe. Further, whereas the starting-point of corporeal endometritis is often labour, or abortion, malignant disease of the body of the uterus usually begins long after the menopause, and usually affects

those who have been either actually sterile, or who have had only one or two children.

Lastly, in cases of doubt we can scrape away portions of the masses projecting into the uterine cavity, and examine sections of them with the microscope.

In the corporeal endometritis of old women the surface of the mucous membrane of the body of the uterus is smooth; sometimes there is occlusion of the internal or external os; sometimes, even without occlusion, there is




Fig. 59.—Section of a granulation, from a case of corporeal endometritis (De Sinéty).

I. Stroma. 2. Dilated glands,

a considerable accumulation of purulent secretion, often intensely fœtid, in the cavity of the body of the uterus. This condition is known as *pyometra*.

Pathology.—In acute cases the mucous membrane is said to be swollen, and secreting pus.

In chronic cases, which are those we have most frequently to deal with in practice, there is hypertrophy of the mucous membrane of the body of the uterus, either a localised hypertrophy, as in Case I. above, or a general hypertrophy. Localised hypertrophy would give the variety called "fungous endometritis"; in these cases definite irregular

projections can be detected with the finger after dilatation of the cervix.

The hypertrophy of the mucous membrane is said to affect all the elements of the mucous membrane, the glands, the blood-vessels, and the connective tissue. In Case I. (above) the glands were considerably dilated.

Treatment.-When the general features of the case lead us to determine on local treatment, e.g., persistent menorrhagia for which no other cause can be found, and which resists ordinary treatment by ergot, hot douches, etc., or recurring abortion, for which no reason can be found, the best practice is to dilate the cervix, and examine the surface with the finger to render our knowledge of the case as precise as possible; then, with a blunt-edged curette, to systematically scrape the whole of the interior of the uterine body, afterwards to apply carbolic acid, or tincture of iodine (pure), to the whole of the surface, and wash away any excess with a douche of weak iodine water through the intra-uterine tube. This was the treatment adopted in Case I., and she was permanently cured. Corporeal endometritis is certainly a cause of sterility, or, if pregnancy occurs, a predisposing cause of abortion. Cervical endometritis probably has little or no power to cause either sterility or abortion.

### METRITIS.

By this is meant interstitial inflammation of the uterus, inflammation of the uterine tissue lying between the peritoneal surface on the one hand, and the mucous surface on the other. Although metritis is a real condition, it cannot be diagnosed apart from the inflammation of the mucous surface, or inflammation of the peritoneal surface, of the uterus. When either of these conditions

is present in marked degree, it may be taken that there is more or less metritis present as well. The case is analogous to pericarditis and endocarditis in relation to myocarditis.

### SUBINVOLUTION OF THE UTERUS.

Immediately after delivery the uterus weighs about twenty-eight ounces.

Under normal conditions its weight falls in six weeks' time to about one ounce and a half. The process by which this change is accomplished is known as the *Involution of the Uterus*.

The uterus of a woman who has borne children always, however, remains permanently somewhat larger than the uterus of a nullipara.


DIFFERENCES BETWEEN THE UTERUS OF A VIRGIN AND THE UTERUS OF A WOMAN WHO HAS BORNE CHILDREN.

# The uterus of a virgin (Fig. 60).

Length.—Its whole length, measured externally from the highest point of the fundus to the external os, is about two inches and a half. A slight constriction, visible externally, divides it into two parts, nearly equal in length—the upper and slightly longer portion comprising the fundus and body of the uterus, the lower portion being the cervix.

Cavity.—On coronal section (Fig. 60, C) we see that, in the virgin, the cavity of the body is rather shorter than the cavity of the cervix.

In shape the cavity of the body is triangular, the sides of the triangle being convex inwards. The cavity of the cervix is fusiform, and is marked by a longitudinal ridge, anteriorly and posteriorly, from which secondary ridges



2, 2, angles; 3, cervix; 4, site of the os internum; 5, vaginal portion of the cervix; 6, external os; 7, 7. vagina. B, 1, 1, profile of the anterior surface; 2, vesico-uterine cul-de-sac; 3, 3, profile of the posterior surface; 4, body; 5, neck; 6, isthmus; 7, cavity of the body; 8, cavity of the cervix; 9, os internum; 10, anterior lip of the os externum; 11, posterior lip; 12, 12, vagina. C, 1, cavity of the body; 2, lateral wall; 3, superior wall; 4, 4, cornua; C, I, cavity of the body; 2, lateral wall; 3, superior wall; 4, 4, cornua; Fig. 60.—Virgin Uterus. A, anterior view; B, median section; C, lateral section (Sappey). 5, os internum; 6, cavity of the cervix; 7, arbor vitæ of the cervix; 8, os externum; 9, 9, vagina.

spring, directed obliquely upwards; this arrangement is called the arbor vitæ.

The length of the cavity of the uterus in virgins averages 2'1 inches (Barnes). Owing to stretching of the tissues, however, the sound generally passes about 2½ inches in virgins.

External os.—In virgins this is either round, or transverse; its outline is free from irregularities, and its longest diameter varies from half a line to two lines,  $\frac{1}{24}$  to  $\frac{1}{6}$  of an inch (Lusk).

### The uterus of a woman who has borne children.

Length.—Its whole length externally is about three inches, two inches of which belong to the fundus and body, and one inch to the cervix.

Cavity.—On section the cavity of the body is seen to be distinctly longer than that of the cervix. In shape the cavity of the body is more ovoid, the sides being rather concave inwards, than convex as in the virgin.

The length of the cavity of the uterus in parous women is not less than  $2\frac{1}{2}$  inches.

External os.—This is either round, or transverse; its outline is irregular from the slight lacerations that occur physiologically in labour. The os is often large enough to admit the tip of the finger, and its longest diameter measures about half an inch.

Etiology of subinvolution.—The process of involution is interfered with by any of the following causes:—

- 1. Pelvic inflammation (perimetritis, parametritis).
- 2. The retention of portions of placenta or membranes, or of clots, after labour or abortion. In such cases corporeal endometritis is likely to co-exist. (See page 168 for a very typical case of subinvolution in connection with corporeal endometritis.)
  - 3. The presence of fibroid tumours of the uterus.- I have

seen several examples of subinvolution associated with, and apparently due to, the presence of uterine fibroids.

4. Passive congestion of the uterus due to getting up too soon after labour, and more particularly getting up too soon after abortion or miscarriage—the patient not thinking it necessary to remain in bed so long after the latter as after delivery at full term.

The way in which passive congestion occurs in such




FIG. 61.—Multiparous Uterus, anterior surface (Sappey).

Note.—The peritoneum should only be figured covering the uterus as low as the internal os on the anterior surface.

cases is as follows:—The uterus is still larger than normal, and the tissues around it lax. When the patient gets up the uterus descends—there is a certain degree of prolapse; this of itself, by dragging on the veins in each broad ligament, causes increased difficulty in the venous blood passing off, *i.e.*, produces passive congestion.

The descent of the uterus is associated with some degree

of retroversion, either alone, or combined with retroflexion; and, owing to the size of the uterus, the *incarceration of* the uterus in Douglas's pouch, described in the chapter on Versions and Flexions, is very apt to occur, still further increasing venous congestion, and retarding the process of involution.

The symptoms of subinvolution are:—
More or less bearing-down pain, and backache.




FIG. 62.—Multiparous Uterus, uterine cavity (Sappey).

Menorrhagia, and often also metrorrhagia. A yellow discharge, which may be offensive.

On examination we find the body of the uterus larger than it ought to be, perhaps very much larger (as in the case already referred to on page 168). Very often we find the uterus lower than normal, and retroverted or retroflexed; and it is particularly in cases of this kind that incarceration of the uterus in Douglas's pouch is commonly

met with. As this condition is fully described later on, it is unnecessary to say more about it here. Sometimes a polypoid mass may be found just within the external os.

The sound passes an increased distance,  $3\frac{1}{4}$ ,  $3\frac{1}{2}$ , or 4 inches. In bad cases it may pass five or more inches according to circumstances. It is important to notice that the bimanual examination often shows that the uterus is larger than it should be, that there is subinvolution, when the sound only passes the normal distance, or a very little more. Here the increased bulk of the uterus is due to excessive thickness of the walls.

Before passing the sound, it is of course necessary to make certain that the enlargement of the uterus is not due to pregnancy; as regards the history, the symptoms dating from a particular confinement or miscarriage will usually help us, and as regards physical signs, the feeling of the body of the uterus in subinvolution is different from the feeling it has in pregnancy. In subinvolution, though the uterus is enlarged, it feels firm and hard on bimanual examination; its shape is distinctly flattened from before back, although the thickness of the walls is increased. In early pregnancy the shape of the enlarged body is more globular, its outline is less definite, and it has an elastic feeling.

Other signs likely to be present in pregnancy, such as purple discoloration of the vaginal mucous membrane, and of the vaginal portion of the cervix, and softening of the cervix, will also aid us in diagnosis.

Treatment.—Preventive treatment.—When we look at the etiology of the disease, we see at once that it usually arises from preventable causes. Thus, as regards pelvic inflammation occurring after labour or abortion, we know that this is usually, if not always, of septic origin, and moreover in most cases the septic element is introduced by dirty

hands, finger-nails, or instruments—careful regard to antiseptic principles in midwifery practice is therefore of the first importance. This is more fully discussed under Pelvic Peritonitis.

Careful attention to see that all the placenta and membranes have come away is another element in the preventive treatment of subinvolution. Again, it is well to give ergot for a week after labour as a matter of routine in all cases. The patient should be advised to suckle, in the absence of any strong reason to the contrary. Another point is to see that the patient does not get up too soon after labour. The continued presence of a red discharge after delivery or abortion is to be looked on as an indication that the patient should remain in the recumbent position, although not necessarily in bed. When the time for preventive treatment has gone by, and a case of subinvolution comes before us, if there is no offensive discharge, and if there is nothing felt just within the external os, such as placental polypus, the treatment should consist of replacing the uterus, if necessary, with the sound, and inserting a ring pessary to keep it in good position. The patient should remain in bed for a time. Hot vaginal douches should be given three times a day. For medicine, she should take some preparation of ergot thrice daily. Under this treatment many cases get well. If, however, the case does not improve after this treatment has been tried for three weeks or so, the next thing to do is to dilate the cervix and examine the interior of the uterus for a piece of adherent placenta, or other abnormality of the endometrium, and if we find anything of the kind, to remove it.

If there is an offensive discharge from the uterus, with persistent subinvolution, as in the case described on page 168, the treatment adopted there is the right one: scraping the whole of the endometrium thoroughly

with a spoon, if necessary, after previous dilatation of the cervix.

In all intra-uterine manipulations the vagina should first be douched with an antiseptic lotion, and the hands and instruments carefully rendered aseptic; after we have finished, the cavity of the uterus should be thoroughly washed out through a double-channelled tube with an antiseptic lotion.

When there is evidence of acute, or subacute, pelvic inflammation, this, rather than the subinvolution, must of course be regarded as the important element of the case, and treatment suitable for it must be adopted.

#### SUPERINVOLUTION OF THE UTERUS.

**Definition.**—Excessive involution of the uterus. The disturbance of the involution is here in the opposite direction from subinvolution; the process goes too far, and leaves the uterus atrophied.

We know little of the causes of this condition. The symptoms are amenorrhoea and sterility, and the diagnosis depends on finding by bimanual examination, and on passing the sound, that the uterus is much smaller than it should be.

Nothing can be done in the way of cure; and if the patient's general health is good, the case is best left alone.

Notes of two cases which came under my care at the London Hospital are appended:—

CASE I.—A. S., age 33; married nine years; three children, the last three years previously, and still-born. No miscarriages. All the confinements difficult; forceps used in second confinement.

Has not menstruated since the last confinement. She lost more blood after the last confinement than after the others, "as the after-birth had grown to her side."

For the last five months has suffered from headache. There has

been no epistaxis, or other loss of blood. Every month she feels as if menstruation were coming on, and has some yellow discharge.

The catamenia appeared when she was fifteen, and she was always regular, though the loss was scanty.

Has had sore throat, on and off, the last twelve months.

Bowels confined as a rule.

Pelvic measurements.—Between anterior superior spines =  $8\frac{1}{4}$  inches; between iliac crests =  $9\frac{1}{4}$  inches; external conjugate =  $6\frac{3}{4}$  inches.

On bimanual examination the uterus was found to be very small in all directions; the sound passed rather less than 2 inches.

(N.B.-In women who have had children the sound passes at least  $2\frac{1}{2}$  inches, often  $2\frac{3}{4}$  or 3 inches.) Nothing else abnormal was found.

CASE II.—S. C., age 29; married six years; one child, five years ago; no miscarriages. Confinement normal; no post-partum hæmorrhage. She suckled the child six months. The child died when it was ten months old. Has not menstruated since the confinement. She is well in herself, except that sometimes she feels "heavy in the head." The catamenia appeared at fifteen; she was regular every month, always losing very little; it lasted two or three days, and she had no pain to speak of at her periods.

V.E.—Uterus quite small, as estimated bimanually.

Cervix projects very little into the vagina. Sound passes just two inches. Nothing else abnormal.

Remarks.—In both cases menstruation had always been scanty. In Case I, the patient had had three children, so that in her case scanty menstruation could hardly be taken as evidence of deficient ovarian activity, at least as regards the highest function of the ovary—ovulation.

The pelvis in this case was an example of the "justominor" pelvis—there being a slight degree of general contraction.

#### CHAPTER X.

#### DISEASES OF THE UTERUS.

#### FIBROID TUMOURS.

Etiology.—It is only during the period of menstrual life, between the ages of fifteen and forty-five, that fibroid tumours of the uterus *begin* to grow. Clinically, however, it is rare to meet with them till after the age of five-and-twenty.

Influence as regards conception and labour.—In general terms it may be said that a woman who has fibroid tumours in her uterus is less likely to become pregnant than other women (about one-third of married women with fibroid tumours are sterile). If she should become pregnant, she has a special liability to abort. If, however, she goes to full term, then special dangers may arise:—

- 1. During labour, from mechanical obstruction to the passage of the child, varying with the exact position of the fibroid, the greatest obstruction occurring with a large interstitial fibroid in the cervix. In such a case delivery per vias naturales may be impossible.
- 2. Immediately after labour, from post-partum hæmor rhage.
- 3. Later, from secondary hæmorrhage, or septic fever, if the fibroid should slough owing to injuries received during labour.

Some persistent cases of subinvolution are due to the presence of fibroids, the uterus for weeks remaining about

as large as it was immediately after labour. I have seen several cases of this.

During pregnancy fibroids grow, sharing in the general hypertrophy of the uterus.

Very rarely after labour, under the influence of the process of involution, fibroids may disappear altogether, but well-authenticated cases of this are on record.




FIG. 63.—Typical .Varieties of Fibroids (Thorburn). (Diagrammatic.) I. Subperitoneal; 2. Submucous; 3. Interstitial; 4 in the figure has a distinct stalk, i.e., is a polypus. It is explained in the text that, so long as fibroid tumours are entirely in the cavity of the body of the uterus, and not partly in the cervix, they do not acquire a distinct stalk, though they may have a neck; 4 in the figure, therefore, being a fibroid still entirely in the cavity of the body of the uterus, should not have a distinct pedicle; 5. Submucous fibroid of the cervix,

#### VARIETIES.

As regards position.—All fibroids begin as interstitial fibroids, i.e., at first they lie embedded in the thickness of the uterine wall. As they grow they make their way either towards the peritoneum, or towards the mucous membrane of the uterus, and are then termed subperitoneal or submucous fibroids respectively. A fibroid ceases to be interstitial only when its largest diameter has passed beyond the uterine wall, and when its attachment to the uterus is by a more or less constricted part, or neck. Gradually, in many cases, more and more of the tumour is expelled from the uterine wall till at last the original tumour hangs by a distinct stalk; it has then become a subperitoneal, or submucous, fibroid polypus, as the case may be. The late Dr. Matthews Duncan drew particular attention to the distinction between a submucous fibroid tumour of the uterus, and a submucous fibroid polypus. The former may lie wholly in the cavity of the body of the uterus; it is attached to the mucous membrane by a neck only, not a stalk.

When there is a distinct stalk to a submucous fibroid, when it has become a polypus, the tumour always lies, partly at least, in the cavity of the cervix, and not wholly in the cavity of the body.

Number.—Fibroids are usually multiple; it is exceptional for there to be only one.

Seat.—They most commonly grow from, or in, the walls of the body of the uterus, and are more common in the posterior than in the anterior wall. Less commonly they grow from the supra-vaginal cervix, and still less commonly from the vaginal portion of the cervix.

Structure.—Fibroid tumours are composed of involuntary muscular fibre and fibrous tissue, in varying proportions.

As regards consistency they may be either :-

- I. Hard, or
- 2. Soft.
- I. Hard fibroids.—These are much commoner than soft, and in them there is an excess of the fibrous tissue element, with a comparatively small amount of involuntary muscle. To the naked eye, a hard fibroid has, on section, a greyish-




Fig. 64.—Subperitoneal Fibroid Tumour of the Uterus (London Hospital Museum). The tumour is only attached by a thin pedicle to the fundus uteri (Barnes).

white colour, with a pearly lustre, and owing to the wavy arrangement of the fibres, its appearance has been aptly compared to that of a ball of cotton. As the bundles of fibres may be arranged round several centres, an apparently single fibroid may, on section, have an appearance like several balls of cotton placed in contact with one another. Hard fibroids have a sort of capsule formed of the tissues

of the uterus in which they are embedded, and they are connected with their capsules by a loose connective tissue, which permits a tumour of this variety to be easily shelled out when the capsule has been divided. The vessels supplying a hard fibroid lie under the capsule round the circumference of the tumour, few or none penetrating its substance (see Fig. 66).

2. Soft fibroids—the rarer variety—have no distinct capsule, their tissues being continuous at the circumference with the tissues of the uterus. They have a pale pink colour. Microscopically they are found to have an excess of involuntary muscular fibre, with but little fibrous tissue.

Natural history.—Fibroids may remain for years unaltered and stationary, or they may continue to grow.

There are certain changes which fibroid tumours are liable to undergo:—

- 1. They may calcify.
- 2. They may soften into a puriform mass, which may, in the case of interstitial fibroids, be discharged through an opening formed towards the mucous membrane.
- 3. They may (in the case of submucous polypi) become gangrenous, and be discharged if the patient survive, which is unlikely.
  - 4. Occasionally they undergo malignant degeneration.
- 5. An opening may be formed in the capsule by ulceration, and the tumour be extruded through it *en masse*.

It is to be remembered that the natural tendency of fibroid tumours is not, as a general rule, to kill the patient; they may cause great suffering, but they only rarely kill.

In this respect they stand in marked contrast to ovarian tumours.

• 6. They may become cystic. How this occurs is not yet settled.

Influence of menstruation on fibroids.—Fibroids

are at their largest immediately before the commencement of the flow, and at their smallest at the end of the flow;




Fig. 65.—Uterine Fibroids, one of which has become cystic (Schroeder).

between the periods they gradually enlarge, attaining a maximum just before the next period.

Symptoms.—There may be none, and it is more particularly in the case of subperitoneal fibroids that this occurs. Subperitoneal fibroids are more likely to have attention directed to them on account of their size than on account of symptoms, such as occur with the other varieties of fibroid tumours. These are:—

- 1. Bleeding.—Menorrhagia and perhaps also metrorrhagia.
  - 2. Painful menstruation.
  - 3. A yellow discharge.
  - 4. Pressure symptoms.

**Pressure symptoms.**—In slight cases there is merely a sense of weight in the pelvis, with a feeling of "bearing down." More severe symptoms may arise: (a) if the tumour be of great size, or (b) even with a small tumour, if on account of its original position it becomes incarcerated in the pelvis.

Micturition.—Frequency of micturition is a common symptom, particularly if the tumour is in the anterior wall. Retention of urine is not uncommon in cases of fibroids, whereas in cases of ovarian tumours it is rare.

Defæcation.—Very rarely complete obstruction may occur from pressure of an incarcerated fibroid on the rectum. If a woman is the subject of uterine fibroids, and intestinal obstruction occurs, there is a strong probability against the obstruction being due to the fibroids.

Pressure on veins may cause the veins of the lower extremities to become varicose, and there may also be some of the violet tinting of the vaginal mucous membrane from venous congestion, like that observed during pregnancy.

Pressure on nerves.—Severe neuralgic pains may occur, particularly in cases where the fibroid is incarcerated in the pelvis.

Pressure on the ureters occasionally causes them to become dilated, and ultimately hydronephrosis may be produced.

Finally, in the case of very large tumours, respiration and digestion may be interfered with.

Diagnosis.—Fibroid tumours have to be diagnosed from:—

- 1. Pregnancy.
- 2. Ovarian tumours.
- 3. Pelvic inflammation.
- 4. Pelvic hæmatocele.
- 5. Retroflexion and anteflexion.
- 6. Some cases of carcinoma.
- 1. Pregnancy.

As regards the history.—In pregnancy there is usually complete amenorrhoea, whereas in fibroids there is menorrhagia. Sometimes, however, menstruation may go on during the first three months of pregnancy; and even as late as the fifth or sixth month, there may be hæmorrhages at intervals of a month, which the patient takes for ordinary menstruation. The explanation of hæmorrhages so late as this must be that there is really a threatened abortion each month, and that the bleeding occurs at intervals of a month on account of the monthly congestion of the uterus, which probably goes on even during pregnancy.

Again, there is a history of morning sickness, and if the pregnancy has reached the end of the fourth month, the history of quickening, whereas in fibroids these symptoms are absent.

Physical signs.—Supposing the tumour in the abdomen is up to the umbilicus, if it be the pregnant uterus it feels elastic, and has a uniform outline and surface, and we may feel it alternatively harden and soften; if it be a fibroid of

the common hard variety, it feels hard, not elastic, and as fibroids are usually multiple, the tumour will probably have




FIG. 66.—Uterus containing Fibroid Tumour, from a case which terminated fatally through hæmorrhage. Large venous sinuses are seen in the capsule, one of which ruptured at the point (a) (Matthews Duncan).—Edin. Med. Jour., 1867, p. 634.

an irregular outline and surface; even if there is only one fibroid, it is very rarely situated so centrally as to have the symmetry of the pregnant uterus at the fifth month. Again, on auscultation, we may expect to hear the fœtal heart if the child is alive; we may also hear the uterine souffle, but this is of less value, as it is occasionally heard over fibroid tumours. Still, a well-marked souffle is very suggestive of pregnancy.

Practically, the greatest difficulty arises in cases where, with no history of amenorrhoea, there is a tumour in the abdomen rising out of the pelvis, and reaching perhaps up to the umbilicus, centrally situated, soft and elastic, and evidently connected with the uterus, and yet over which no foetal heart can be heard.

The tumour may be a *soft* fibroid of the uterus, or it may be the pregnant uterus, the fœtus having died, the ovum being for some reason retained—the woman, in fact, being in a condition of missed abortion, or rather, at this period, of "missed miscarriage": or, again, the case might be one of hydatidiform mole; or it might be a case of placenta prævia.

As regards other points to which attention must be paid, there is the woman's own opinion, either that she is, or is not, pregnant, which is worth something, but not much. Patients are often mistaken in this matter, the non-pregnant thinking herself pregnant, and the pregnant having no idea of her condition.

The condition of the breasts, the purple colour of the vaginal mucous membrane, and the vaginal portion of the cervix, the softening of the cervix, are all signs to which due weight must be given; but it must be borne in mind that there may be milk in the breasts of a woman who is not, and never has been, pregnant; that the purple coloration of the vagina is mainly due to pressure, and

may, therefore, be caused by other tumours; and that the cervix may be softened in some exceptional cases of fibroid tumours.

As a matter of experience these cases almost always turn out to be cases of pregnancy—"missed abortion," or "missed miscarriage"; hydatidiform mole; or placenta prævia.

Finally, it must not be forgotten that both conditions may be present together—normal pregnancy and fibroid tumours of the uterus.




Fig. 67.—Multiple Fibroids of the Uterus (Winckel).

a. Fibroid tumour in right broad ligament. b. and e. Subperitoneal tumours. c. Interstitial fibroid. d. Submucous fibroid. f. External os. g. Right ovary. h. Right Fallopian tube.

Note how the uterine cavity is encroached upon, and rendered unsymmetrical by the projection of the tumours.

In such cases there is amenorrhoea, the uterine tumour is probably irregular in outline and surface, and is larger than would correspond to the supposed period of pregnancy; for certain diagnosis we must wait until the foetal heart is heard.

I remember a case very well, where a woman who had been a widow for some years, and was over forty years of age, came with a history of three months' amenorrhoea, and where there was an elastic irregular tumour connected with the uterus much larger than the uterus at the third month of pregnancy. The tumour rapidly enlarged, and finally, when it was several inches above the navel, after about five months' amenorrhoea, we heard the foetal heart. At this time the surface of the tumour was markedly irregular, some three inches of the upper part being hard, and separated by a deep transverse groove from a soft and more uniformly regular, lower portion, which was no doubt the body of the uterus containing the ovum.

Hegar's sign of pregnancy.—Where the size of the uterine tumour does not exceed that of the pregnant uterus at the fourth month, Hegar's sign of pregnancy is of much value in differential diagnosis. This sign depends on the fact that in pregnancy from about the sixth week onwards the segment of the uterus at the level of the internal os, that is the part at the junction of the cervix with the body of the uterus, becomes markedly softened. To obtain the sign the left hand is placed on the abdomen, and is pressed down so as to be on the posterior aspect of the uterus at the level of the internal os. The index finger of the right hand in the vagina is pressed upwards towards the anterior aspect of the uterus at the corresponding level-that of the internal os. When Hegar's sign is well marked, the fingers of the two hands seem almost to meet, as if the cervix were separate from the swelling formed by the body of the uterus. Indeed, it is said that sometimes the cervix has been mistaken for the whole uterus, and the enlarged body of the pregnant uterus for a tumour separate from the uterus, such as an ovarian tumour, in cases of pregnancy where the observer has not been aware of the conditions on which Hegar's sign depends. The sign is available from the sixth to about the sixteenth week. After that time it is generally

impossible to get the upper hand depressed sufficiently low behind the uterus to recognise the sign.

2. Ovarian tumours.—It has already been said that soft fibroids which feel elastic and almost fluctuating (and which are therefore more likely to be mistaken for ovarian tumours) are comparatively rare. Again, hard, solid-feeling tumours of the ovary are rare, and, when they do occur, are likely to be mistaken for the common hard fibroid, especially the subperitoneal fibroid with a long stalk. Nevertheless, as a general rule, great importance should be attached to the consistence of the tumour. Dr. Barnes says, "If you find a smooth solid tumour, beware; it is uterine."

As regards history, while menorrhagia is the rule with fibroids, it is the exception with ovarian tumours; but still we must remember that in cases of ovarian tumour, both small and large, profuse menstruation does occasionally occur; more commonly, however, menstruation in cases of ovarian tumour is not excessive, and either regular, or if its regularity is disturbed, it is in the direction of amenorrhæa, the intervals between the periods being increased, and the flow lasting for a shorter time, or being less in quantity, than formerly.

On vaginal examination, in cases of large ovarian tumour, the finger, as a rule, reaches the vaginal portion of the cervix easily, often more easily than usual, because the whole uterus of the somewhat pushed down by the pressure of the tumour; it is exceptional for the cervix to be drawn up, so that the os is only reached with difficulty, or not at all, though this does occur sometimes. In large fibroid tumours, on the other hand, it often happens that the vaginal portion of the cervix cannot be reached with the finger, sometimes being so drawn up that even when a full-sized Fergusson's speculum has been passed to its full extent, it is yet an inch or two too short to bring the os into view.

In less extreme cases the vaginal portion may be drawn up so that the os is flush with the vaginal roof.

I remember a case of Dr. Graily Hewitt's of this kind, where the sound could only be passed through a speculum, as the situation of the os could not be detected by the finger alone.

The sound, held by the handle in a long pair of cervix forceps, then passed till only the handle was left projecting beyond the os uteri!

The sound.—As a rule in cases of ovarian tumour the sound passes only the normal distance; whereas in cases of fibroids it passes more than two and a half inches, often much more, as in the case just mentioned. In rare cases the uterus is stretched upwards by the growth of an ovarian tumour adherent to it, and then we do have moderate lengthening of the uterine cavity, the sound passing three or four inches. This is, however, quite the exception.

The bimanual examination.—In cases of large abdominal tumours this often gives no additional information.

When there is a large solid, or chiefly solid, ovarian tumour, it is apt to be mistaken for a uterine fibroid. Sometimes in such cases the sensation imparted to the finger by tilting the cervix enables one to recognise that there is no such close connection between the uterus and the tumour as there would be if the tumour were a uterine fibroid. While the finger touches the cervix it is useful in such cases to have the abdominal tumour pulled upwards as much as possible by an assistant. This may show the tumour not to be so closely connected with the uterus as a uterine fibroid would be.

There are other points that help one in a difficult case. For example, a large tumour that has grown rapidly, if it be a fibroid of the uterus, is almost certain to be attended with a good deal of menorrhagia, and again, the sound will

usually pass much more than the normal distance. In the following case the tumour had been previously diagnosed by another gynæcologist as a uterine fibroid, and it had been treated by the Apostoli treatment before it came under my observation.

Sarah V., aged 35, married seventeen years, five children, the last eight and a half years ago, came under my observation at the London Hospital on August 11th, 1889. She had then been in the Hospital about five weeks.

History of the illness.—She was under medical treatment in March and April of this year, "suffering from pain in the abdomen with sickness and swelling of the abdominal walls." I have ascertained that she was then supposed to be suffering from chronic constipation, and that the swelling was believed to be a fæcal accumulation.

The swelling in the abdomen was first noticed about Christmas, 1888. Since April she has been in good health, but the swelling in the abdomen has not lessened; indeed, she thinks it has gradually increased in size since it was first noticed (Christmas, 1888).

For about eleven weeks previous to the time I first saw her she had been losing too much. During the time she had been losing too much she had constant pain in the abdomen. The pain was not confined to one place; sometimes it was in the right iliac region, at other times in the left groin.

Menstrual history.—The catamenia appeared when she was sixteen years and two months old; she was regular every four weeks, the flow sometimes lasting a week, but unaccompanied by pain.

Since the birth of her last child, eight years ago, till last Christmas she was quite regular. From Christmas to the end of March the flow came on every fortnight without pain. Somewhere about April or May she went over the month without seeing anything. She was then feeling well, and free from pain.

There has been no inter-menstrual discharge.

Other symptoms.—Micturition has been slightly painful the last eleven weeks. Has had piles the last three months. Defæcation has been somewhat difficult since the present illness began. She thinks she has been losing flesh during the last four months.

The patient told me she was sure that twelve months ago she was quite regular, and not losing too much.

Her weight on April 9th, 1889, was 10 st. 9 lb.; on July 6th it was 10 st. 1 lb.

Present state, August 11th. - The patient is not anæmic to any marked degree.

Abdominal examination.—A tumour is felt rising out of the pelvis, and reaching (measured with callipers) at the highest point, which is to the right of the middle line, a distance of 10½ inches above the upper margin of the pubic symphysis.

The tumour for the most part feels as if it were solid, but it certainly has in places an elastic feeling, though distinct fluctuation cannot be obtained.

The vaginal examination.—The vaginal portion of the cervix has its normal range of mobility, and on tilting it with the finger it gives the impression of belonging to a uterus the body of which is not much enlarged, and also the impression that it is not closely incorporated with the large tumour felt in the abdomen.

The sound passes three inches. While the sound was in the uterus the tumour in the abdomen was raised as much as possible away from the pelvis by an assistant. Doing this did not appear to alter the position of the sound in the least. I came to the conclusion that the case was one of ovarian tumour, probably of the semi-solid variety.

Operation, August 29th, 1889.—On opening the abdomen a small quantity of fluid escaped. The appearance of the tumour was unlike that of a fibroid; and, on passing the hand in, the uterus could be felt to be of about the normal size and quite distinct from the tumour.

Wells's trocar was thrust into a part of the tumour that seemed comparatively soft, but only a small quantity of dark grumous fluid came away. Accordingly, the incision was extended upwards to about three inches, or rather more, above the umbilicus, and the tumour was lifted out of the abdominal cavity whole. The peritoneum was washed out with weak iodine water, and a Keith's tube having been inserted, the wound was closed in the usual way.

The tube was taken out in thirty-six hours' time.

The patient made a perfectly satisfactory recovery. On September 2nd the wound measured 7½ inches.

Now, the points in the case which inclined me to believe that the tumour was not a uterine fibroid, but an ovarian tumour, were:—

(a) As regards the history.—The short time that had

elapsed since the beginning of the illness. Before Christmas, 1888, the patient had been in her ordinary health, and noticed nothing amiss. Yet in August 1889 we found a tumour in the abdomen about the size of the pregnant uterus at the eighth month.

The great probability would be that a uterine fibroid of that size would have given rise to symptoms for two or three years or more.

(b) As regards physical signs.—The absence of anæmia; a uterine fibroid of the size and consistence (it was distinctly elastic in parts) presented by the tumour would almost certainly have caused profuse hæmorrhage, and made the patient anæmic.

The feeling imparted to the finger on tilting the vaginal portion of the cervix gave the impression that the uterus had no close connection with the tumour and was of the ordinary size. Again, the sound only passed three inches.

It may be objected that the tumour might in spite of this have been a subperitoneal fibroid connected by only a thin pedicle to the uterus; but the consistence of the tumour, apart from other features in the case, would be strongly against that view; subperitoneal fibroids of that kind are typically hard, and have a history extending over years, whereas in this case the tumour was distinctly soft and elastic in parts, and the history only extended over about eight months.

When the tumour is of moderate size, say reaching up to the umbilicus, or small and entirely in the pelvis, and the abdominal parietes are lax, then, in cases of ovarian tumour, we can often isolate the uterus by the bimanual examination from the tumour; whereas in ordinary cases of fibroids we should not be able to do this; the whole tumour would move on giving an impulse to the cervix.

Very rarely a subperitoneal fibroid may be met with having a long thin stalk permitting us to isolate the uterus from it as completely as if it were an ovarian tumour.

Fibro-cystic tumours of the uterus cannot be generally diagnosed from ovarian tumours. They are extremely rare compared to ovarian tumours. The following case is an example of a large fibro-cystic tumour of the uterus mistaken for an ovarian tumour:—

Miss S., aged 45, was brought to see me by Dr. Warren, of Enfield Highway, in December 1893. The abdomen had been noticed to be enlarging for about three years. Till two years ago patient had been regular every four weeks, and always lost very little. For the last two years she had not been regular. Latterly she had menstruated every fortnight, but the quantity lost each time had been very small. On examination there was found a thin-walled fluctuating swelling, reaching well above the umbilicus and rising out of the pelvis. The sound only passed two and a half inches. I diagnosed the case as one of ovarian tumour. At the operation, however, after tapping and emptying the cyst (which was unilocular and contained a thin straw-coloured fluid) I found that both ovaries were normal, and that the tumour was uterine. The operation was undertaken at the patient's own house at Enfield Highway; fortunately I had the serre-nœud and transfixion pins with me, and accordingly performed a supra-vaginal hysterectomy, fixing the pedicle externally in the lower angle of the wound. The pedicle so fixed was, of course, composed of uterine tissue, but I did not see the cavity of the uterus at all, and believe that it was not opened. The parts removed were the cystic tumour and a portion of the fundus of the uterus. The ovaries were so situated that there was no need to interfere with them. The patient made an absolutely uneventful recovery, and there is no sinus at the lower angle of the wound, which is soundly healed throughout. The patient, when I last heard of her, had been regular since the operation. I showed the specimen at a meeting of the Obstetrical Society of London in October 1894.

## 3. Pelvic inflammation.

History.—The symptoms would be found to date from either labour, abortion, or exposure during menstruation, or to have followed an acute vaginitis, especially one of

gonorrhœal origin—in fact, to have been preceded by some of the ordinary antecedents of pelvic inflammation.

The symptoms would be abdominal pain, perhaps vomiting, and more or less high fever.

Physical signs.—Abdominal examination.—In cases of pelvic inflammation, large hard swellings may be formed by gluing together of the intestines and omentum by adhesive lymph. Such swellings have a less definite outline than swellings due to fibroids; they are also fixed, whereas fibroids usually admit of a certain amount of displacement. Again, in pelvic inflammation the tumour is tender, whereas in fibroids it is not tender.

Vaginal examination.—Bimanual examination.—In acute cases of pelvic inflammation the vagina will be found hot, and the uterus more or less fixed. Bimanual examination is often impossible on account of tenderness. In fibroids the vagina is not hot, and the uterus is not fixed—unless it happens that there are adhesions, which are rarely sufficiently extensive to fix the uterus, or unless lateral fibroid outgrowths have encroached on the broad ligaments, so diminishing, or obliterating, the amount of "slack" which allows the uterus to move.

In practice there is usually little difficulty in diagnosing fibroids from pelvic inflammation.

## 4. Pelvic hæmatocele.

History.—Here there is a history of sudden onset—pain in the lower abdomen coming on suddenly, with the appearance of a red vaginal discharge, often when, as regards menstruation, the patient is a week or two over her time, with faintness, nausea, perhaps vomiting, followed, after twenty-four hours, by some fever. In fibroids there is no history of this kind; the symptoms present have usually all come on gradually.

Abdominal examination.—Hæmatocele may, if large, form a swelling reaching a variable distance upwards in

the abdomen according to its size, say three or four inches above the pubes. Here there is tenderness, and the swelling is fixed.

Vaginal examination and bimanual examination.—The uterus, in hæmatocele, can often be isolated from the tumour, and is usually pushed close behind the pubes, and fixed there, partly by the pressure of the swelling, but chiefly by the adhesive peritonitis in the neighbourhood of the hæmatocele. After the first forty-eight hours, in hæmatocele, we usually have a hard swelling bulging down Douglas's pouch, and symmetrically situated as regards the middle line. The swelling cannot be displaced out of the pouch.

Here again the only case of fibroids likely to give rise to similar physical signs is that where we have fibroids of the posterior wall incarcerated in Douglas's pouch, either by having grown so as to be fitted tightly into the pouch, or having been fixed there by pelvic peritonitis.

In practice cases of hæmatocele are more likely to be mistaken for solid ovarian tumours than for fibroids.

5. Anteflexion and retroflexion.—We have to be careful not to mistake anteflexion or retroflexion for fibroids of the anterior or posterior uterine wall respectively.

Anteflexion.—Here the body of the uterus forms a swelling felt on vaginal examination in the anterior fornix; bimanually, the body can be grasped between the internal and external fingers, and its size, shape, and particularly its outline, regular or irregular, may be pretty easily determined. If there were a fibroid in the anterior wall, the thickness of the body of the uterus from before back would be greater than normal; again, the outline of the anterior surface of the body of the uterus would be irregular, according to the extent to which the fibroid projected towards the peritoneal surface.

On passing the sound, and holding it in position, if the case is simply one of anteflexion, the tumour in the anterior fornix will be found to form a much less obvious projection than before; if an assistant holds the sound in place while a bimanual examination is made, it becomes evident, on grasping the uterus between the fingers, that there is only the thickness of the uterine wall between the sound and the anterior aspect of the tumour; and, further, after recognising by gently tilting the sound whereabouts its point lies, we can make out that there is no convex swelling on the anterior surface of the uterus below this level. Careful attention to these points will make the case clear.

Retroflexion.—Here the body of the uterus forms a swelling felt in the posterior fornix. On passing the sound it will be found to enter with the concavity backwards, and its point may be recognised as lying in the centre of the tumour with no more thickness of tissue between it and the examining finger than would be accounted for by the posterior wall of the uterus.

Again, by raising the uterus with the sound, the tumour will disappear from behind the cervix, and by completing the replacement we can feel the body of the uterus bimanually through the anterior fornix; previously to using the sound we should have been able to make our fingers meet immediately in front of the vaginal portion, without anything like the body of the uterus between them.

If the tumour were a fibroid of the posterior wall:-

The sound would pass with the concavity more or less forwards, and by gently tilting the sound we could make out its point through the abdominal wall, while the tumour was still to be felt, as distinctly as ever, in the posterior fornix.

## DIAGNOSIS OF FIBROID POLYPI.

## A. When they have passed the os uteri.

Malignant disease of the cervix.—This sometimes takes the form of a polypoid mass, perhaps the size of a large pear, which seems to spring from the vaginal portion of the cervix by a sort of a pedicle, thus simulating fibroid polypus of the cervix; if the mass is malignant, the pedicle is never so definitely circumscribed and distinct, as in a case of fibroid polypus.

Again, a malignant growth of this kind is soft, friable, easily bleeding when touched, whereas a fibroid polypus, unless sloughing, has a shining, pinkish capsule, and is of firm consistence; many other considerations would make the diagnosis clear, such as duration of the symptoms, etc., but the physical signs referred to are quite sufficient; even a fœtid, sloughing fibroid polypus is not generally friable—the sloughing only affects the surface, the bulk of the tumour still remaining hard.

In certain rare cases, a malignant growth springing either from some part of the cervical canal, or even from the body of the uterus, may project through the external os uteri into the vagina. The nature of the growth is recognised by the characters just given—softness, friability, etc. Such growths may have a pretty distinct pedicle.

Inversion of the uterus.—It must be remembered that a fibroid polypus attached to the fundus may produce an inversion. So that both conditions may be present.

From inversion simply, we diagnose a fibroid polypus:-

1. By passing the sound.—In fibroid polypus we shall be able with care to pass the sound at least the normal distance, and usually farther. In inversion we shall be unable to pass the sound the ordinary distance, no matter on which side of the polypoid mass we try; this is mentioned because

in some cases of fibroid polypi the pedicle may become adherent at some part of its circumference to the os, but it is always free at some part, permitting the passage of the sound.

2. Bimanually, if the case is one of inversion, we may make out absence of the body of the uterus from its usual




Fig. 68.—Fibroid Polypus of the Uterus (Churchill and Leblond).

position. With the sound in the bladder, and the finger in the rectum, care being taken to keep the sound exactly in the middle line, the absence of the uterus from its usual position can be made out.

3. Inspection of the surface of the mass.—The inverted uterus is red and bleeds easily, and it may be possible to see the openings of the Fallopian tubes. There is a specimen in University College Museum where this could

probably have been done. The appearance of a fibroid polypus has been just mentioned above.

4. The mass formed by the inverted uterus is tender, while a fibroid polypus is not.

It is well to remember that inversion of the uterus is very rare, while fibroid polypi are common.

B. When they have not passed the os uteri.

If the os uteri is sufficiently dilated to admit the finger, the sensation imparted to it by coming on a hard, smooth, somewhat movable body within the uterus, combined with the presence of the fundus uteri in its usual position as ascertained bimanually, and also with the fact that the sound passes farther than normal, will make the diagnosis clear.

If the os were not sufficiently open to allow the finger to pass, and a consideration of all the circumstances made it most probable that the enlargement of the uterus was due to the presence of an intra-uterine fibroid tumour, or fibroid polypus, the proper course to pursue would be to dilate the cervix sufficiently to admit the finger, and so settle the question.

Treatment.—This may be considered under two heads:—

- 1. When there is a submucous fibroid tumour, or fibroid polypus (for distinction between them see p. 185).
  - 2. When the tumour is interstitial, or subperitoneal.
- 1. As regards fibroid polypi the right treatment always is to remove them, and the same is true of submucous fibroid tumours, which are attached by a narrow attachment, or neck, to the wall of the uterus, even though they have not yet acquired a distinct stalk.

When the polypus has passed the os uteri, it is a very easy matter, having seized it with the volsella, to pass the loop of an écraseur wire over the tumour, and manipulate it over the largest diameter of the tumour, so that,




FIG. 69.—THE WIRE ÉCRASEUR.

The length of the wire is so adjusted that, when the handle is completely screwed up, the loop has all been drawn back through the aperture at the end of the instrument.

One or two spare wires should always be at hand, as it is not uncommon for the wire to break; also a pair of pincers, for adjusting another wire if the first one breaks. as the wire is tightened, it shall find its way to the pedicle; this should be done gradually, to avoid the risk of bleeding. No traction is to be made with the volsella in tightening the wire. If, however, the polypus is still within the cavity of the uterus, and the os is just large enough to admit the finger, the case is a very different one. In such cases it is recommended to dilate the os further with Barnes's bags, till it is sufficiently open to allow the tumour to pass. I can only say that in a very typical case I gave a most thorough trial to this method, and found it quite useless.

In this case the cervix was stretched over the convexity formed by the lower surface of the polypus, and the os just admitted the finger easily.

A Barnes's bag was introduced several times; but each time, as soon as the bag was distended with water, it simply came out into the vagina. Finally, I incised the thinned cervix posteriorly with Paquelin's cautery, till I judged there was sufficient room for the tumour to pass, and then removed the polypus with the écraseur in the usual way.

In a future case of the kind, I should try to dilate the cervix with a series of specially large Hegar's dilators. The ordinary series stops at No. 26. The larger series which Messrs. Kröhne & Sesemann have made for me runs up to No. 40, which is 1\frac{3}{4} inches in diameter.

In all cases before commencing the operation the vagina should be thoroughly douched out with an efficient antiseptic, e.g., iodine water (3ij. Tinct. Iodi to Oj. water), and if, as in the case just mentioned, the operation has been intra-uterine, the cavity of the uterus should be washed out with the same solution through the intra-uterine tube (Fig. 15, p. 36).

2. When the tumour is interstitial or subperitoneal.—Here the treatment will depend largely on the

symptoms; if bleeding is the chief symptom, we should first try what rest in bed with large doses of ergot will do, preferably using ergotine, gr. ij.-v., in the form of pill, thrice daily. Hamamelis may also be tried (five to ten drops of the tincture three times a day), or hydrastis canadensis (mxx. of the tincture thrice daily), or the infusion of vinca major (see p. 74), at the same time using frequently a vaginal douche of water as hot as the patient can bear it. If such treatment control the bleeding, well and good; but if not, the next step is to dilate the cervix, and make certain that we are not dealing with a case of fibroid polypus, or submucous fibroid tumour, capable of removal with the écraseur. We will take it that, after dilating the cervix, the finger in the uterus finds that the tumour forms a convex projection inwards towards the mucous membrane, but that the bulk of the tumour is in the thickness of the uterine wall—that it is an interstitial fibroid. Formerly the treatment known as enucleation would have been considered proper for such a case; this operation briefly consisted of making an incision over the convex swelling formed by the tumour towards the mucous surface of the uterus, and endeavouring to "get it out somehow" through the opening made. The operation was a very dangerous and unsatisfactory one, and may now be considered obsolete.

In such cases, prior to resorting to radical operation, it is well to try the effect of scraping the endometrium with a curette, the cervix having been previously dilated. After curetting, pure Tr. Iodi may be applied to the whole of the endometrium. Incision of the capsule of the tumour, with or without curetting of the rest of the endometrium, is also often of use in this class of cases.

In the majority of cases treatment of the kind indicated enables the patient either to reach a condition of practical recovery, or at least to hold her ground. Attention to the patient's weight, and examination of the blood by the hæmacytometer and hæmoglobinometer will afford valuable indications as to the efficacy of the treatment. When it is clear that the patient is losing ground, radical treatment must be adopted.

The operative procedures available are:-

- 1. Removal of the uterine appendages with the object of bringing on the menopause. This operation is only suitable for cases where the tumour is of moderate size, not reaching above the umbilicus. In the case of tumours much larger than this it is usually impossible to find the appendages, or, at all events, to remove them. The mortality of this operation is probably from 5 to 10 per cent.
- 2. Enucleation of the tumour, or tumours, from the uterine wall after abdominal section.—The bed of the tumour is then sutured as completely as possible. In this operation the uterine cavity may or may not be opened. The danger of the operation is greater when the cavity is opened. Thus Martin,\* in 24 cases of this operation, the uterine cavity being opened, lost 8; while in 72, where the cavity of the uterus was not opened, only 10 died. In favourable cases this operation leaves the uterus functionally perfect, i.e., capable of child-bearing.
- 3. Supra-vaginal hysterectomy.—This operation is suitable where the tumour is very large, and where the hæmorrhages are so profuse that it is practically certain that the patient must die if left alone, all the palliative treatment previously referred to having failed to control the bleeding. After opening the abdomen, the whole tumour, with most of the uterus, is amputated, the cervix alone being left, and usually clamped in the lower angle of the abdominal wound . (Fig. 70). It is the exception to meet with cases where supra-vaginal hysterectomy is absolutely required; it is a

<sup>\*</sup> Cassell's Yearbook of Treatment, 1891, p. 310.




Fig. 70.—The figure shows the appearance seen at the conclusion of a supra-vaginal hysterectomy. The body of the uterus has been cut away, and the cervix, tightly constricted by the wire of the clamp (a a), is seen fixed in the lower angle of the abdominal wound. Two strong steel pins have been thrust through the cervix, and their points have been guarded by the caps seen on the right of the figure. The pins are above the wire loop (nearer the surface of the stump). The rest of the abdominal wound has been closed by stitches (seen at the upper part of the figure) in the usual way. At the stage of the operation represented, nothing remains to be done except to put on the dressings (Doran).

The figure might also stand for the final stage of Porro's operation, i.e., removal of the fœtus by opening the abdomen and incising the uterus, with subsequent fixation of the cervix in the lower angle of the abdominal wound, and amputation of the body of the uterus.

The pedicle pins are shown separately in the next figure.

more dangerous operation than ovariotomy. In all but a very small proportion of cases ovarian tumours run a fatal course, whereas in all but a very small proportion of cases fibroid tumours left to themselves do not end fatally. They do sometimes, however, as in the following case that came under my observation:—

Miss C., age 39, said to be suffering from a large fibroid. I saw her in consultation on July 22nd, 1890. Her chief trouble has been with the water; twelve months ago she had retention of urine, and a catheter had to be used. The urine runs away from her on walking. She is regular every month; the period lasts 8-9 days; there is a good deal of pain latterly at the periods in the stomach and back.




Fig. 71.—Pedicle Pins used in Supra-Vaginal Hysterectomy.

The periods only lasted five days till a year ago. She has a yellowish-white discharge between the periods. There was a large irregular tumour to be felt in the abdomen, reaching above the umbilicus. As she was very nervous, and the bowels were greatly loaded, it was decided to examine her under chloroform a week later, the bowels to be thoroughly cleared out in the meantime. This was accordingly done, and on July 29th a complete examination under chloroform was made. The tumour felt in the abdomen was uterine; the os uteri was high up close to the pubes, the anterior lip being thin and crescentic, the posterior lip deformed by the presence of a large interstitial cervical fibroid. The sound passed several inches (6 or 8).

Patient was a stout-looking woman, and not the least anæmic. I advised palliative treatment, ergot, hot douching, etc. If she held her ground, well and good; if not, then the question of operative treatment was to be considered.

About four weeks later she suddenly became much worse, a great deal of hæmorrhage occurring. Her doctor asked me then to take her into the London Hospital with a view to operation; but her condition rapidly (within two or three days) became so much worse that he thought she could not bear the journey, and she died a day or two later, hæmorrhage being the cause of death.

The case illustrates the exceptional case of uterine fibroid, viz., where operative treatment gives the patient the only chance of life. It is remarkable on account of the rapidity with which the patient, who was enjoying average good health when I saw her, passed into, first a serious, and then a hopeless condition. I think most probably a large venous sinus must have ruptured, as in the specimen shown in Fig. 66.

There are three principal methods of performing supravaginal hysterectomy:—

- I. The extra-peritoneal method—that just referred to, in which, after opening the abdomen, the body of the uterus, generally with the uterine appendages of both sides, is removed, and the cervix fixed in the lower angle of the wound by means of the clamp (Fig. 70) and transfixion pins (Fig. 71). The abdominal wound above the clamp is closed in the usual way.
- 2. Hysterectomy with intra-peritoneal treatment of the stump—conveniently known as Baer's method. In this operation the abdomen is opened in the usual way. The uterus is then, if possible, brought outside. The first step is to ligature the ovarian vessels on each side. In Fig. 72, at a, a small hole has been made in the left broad ligament and a ligature passed. On the opposite side a further stage is shown, the opening in the broad ligament having been stretched by opening a pair of Wells's forceps in it to the dimensions shown at c. The object is to have room to tie a second ligature as there shown, to prevent reflux bleeding from the uterine side when the tissues between the ligatures have been divided.

At b, Fig. 72, a second hole lower down in the broad ligament is shown ready to be treated as in the other case, so as to secure a further segment of the broad ligament. In Fig. 73 several further stages are shown. On the reader's left of the figure the highest set of ligatures securing the ovarian vessels have been tied, and the tissues divided. The second pair of ligatures are passed ready for tying. On the reader's right of Fig. 73 the second set of ligatures




FIG. 72.—Intra-Peritoneal Hysterectomy.

have been tied, and the part of the broad ligament intervening between them divided. The next step is to make peritoneal flaps in front and behind. This is done by running the knife across the uterus in front and behind at the level of the dotted line  $c\,d$ . The knife is only to cut through peritoneum. The flaps so formed are turned down, one in front and one behind, to about the level of the internal os.

The next step is to tie the uterine artery on each side. This is shown in Fig. 74 diagrammatically. On the reader's left the ligature passed ready for tying; on the right, the

ligature tied. This ligature may be passed either so as to include peritoneum, or it may be passed within the layers




Fig. 73.—Intra Peritoneal Hys erectomy.




Fig. 74.—Intra-Peritoneal Hysterectomy.

of the broad ligament so as not to include peritoneum. When the ligatures on the uterine arteries have been tied, the body of the uterus is cut away—a little above the

level where the ligatures on the uterine arteries have been tied. Fig. 75 roughly represents the condition of things at this stage in antero-posterior section through the middle of the stump. Ca, ca, are the peritoneal flaps. The shaded triangular pieces indicate wedge-shaped pieces of the cervix



Fig. 75.—Intra-Peritoneal Hysterectomy.

in front and behind, which some operators cut out of the stump.

Some operators then stitch uterine tissue to uterine tissue, after cutting out the portion of the stump indicated.




Fig. 76.—Intra-Peritoneal Hysterectomy. Appearance of stump at the end of the operation.

Whether this is done or not, the final step, though not considered essential by Baer, is to stitch the peritoneal flaps together over the stump, so that, at the end of the operation, there is seen a transverse line of sutures across the floor of the pelvis, running from the neighbourhood of the ligatures on the broad ligament on one side to the corresponding point on the other side (Fig. 76).

The abdominal wound is then closed.

3. Panhysterectomy.—In this operation the uterus is removed entirely from above, the cervix as well as the body. The stages of the operation have been well described by




FIG. 77.—Panhysterectomy (from Mr. Christopher Martin's paper).

Mr. Christopher Martin in the Transactions of the Edinburgh Obstetrical Society, vol. xxi., p. 56. The earlier steps—the turning out of the uterus, the securing of the upper parts of the broad ligaments—are much the same as in the operation just described.

An incision is then carried across the anterior surface of the tumour transversely at a safe distance from the bladder. The knife only goes through peritoneum and the tissue immediately under it. This flap is then stripped down, and the bladder separated from the anterior aspect of the cervix. A pair of Wells's large pressure forceps is pushed up the vagina just in front of the cervix. The vagina is then




Fig. 78.—Panhysterectomy (from Mr. Christopher Martin's paper).

opened from above by cutting on these forceps. The finger is then passed through the opening into the vagina, and used to push up the posterior vaginal fornix, which is then opened on the finger. The uterus is still attached on each side by the lower segment of the broad ligament containing the uterine artery. The next step is to tie the uterine arteries. It is necessary to keep close to the uterus to




Fig. 79.—Panhysterectomy (from Mr. Christopher Martin's paper).





Fig. 80.—Panhysterectomy (from Mr. Christopher Martin's paper).

avoid the risk of tying the ureters; but, on the other hand, there must be sufficient tissue beyond the ligature to prevent it slipping. The uterus is now cut away. Any bleeding points on the cut vaginal walls are tied. The ligatures on the uterine arteries are left long, and are drawn down into the vagina. The ligatures on the upper part of the broad ligament are cut short. A fairly thick piece of iodoform gauze is drawn into the vagina from above. About one inch of it is to be left projecting into the peritoneal cavity. The steps of the operation are shown in the accompanying figures from Mr. Martin's paper (Figs. 77-80). Fig. 81 is a drawing from a specimen I removed by this operation. The patient was sent to me by Dr. Blakiston, of Harley Street, W., and made a good recovery.

Panhysterectomy by the combined method.—This appears to offer no advantages over panhysterectomy performed entirely from above. Fig. 82 is a drawing of a uterus with fibroids which I removed by this method in the London Hospital in 1896. The abdomen was first opened, and the body of the uterus with the tumour removed. The abdominal wound was then closed. The patient was then put in the lithotomy position, and the cervix removed per vaginam. Convalescence was tedious, owing to swelling of the left lower limb. The patient is, however, now quite well, and able to follow her usual occupation.

The difficulty of deciding as to whether a radical operation, such as removal of the uterine appendages, enucleation after laparotomy, or hysterectomy, is the proper treatment in any given case, is very considerable; because, on the one hand, if palliative treatment will suffice for the patient's comfort, it is our duty to employ it; while, on the other hand, it would greatly prejudice the prospect of success, should the case be one requiring radical treatment, if we delay this till the patient is extremely anæmic, and her



strength reduced by repeated floodings. Palliative treatment should not be continued if the patient is becoming

paler and paler in spite of it. This may, to a great extent, be taken as indicating whether the case is one requiring operative treatment or not.




Fig. 82.—Uterus with Fibroids, removed by Combined Panhysterectomy. (Case of Mrs. D.)

A. Cut edge of peritoneum. B. Surface left when the uterus was cut off from the cervix. C. Upper limit of cervix. D. Os externum. F. Ligature on uterine appendages, to control meflux bleeding during the operation.

THE ELECTRICAL TREATMENT OF UTERINE FIBROIDS.

The treatment of fibroids by electricity has been reintroduced, with modifications, by Dr. G. Apostoli, of Paris.

Dr. Apostoli's method is as follows:—The patient being in the lithotomy position, a cake of clay, about ten inches by five, and one inch thick, is placed on the abdominal wall. The clay cake is made by working up potter's clay with water to a firm consistence, and enclosing it in muslin. A flat metal plate, about five inches by three, is embedded in the clay. One of the wires from the battery is attached



Fig. 83.—Platinum Sound with insulated sheaths of different lengths, so that the length of the unprotected metal passed into the uterus can be varied according to the length of the uterine cavity. The reverse end of the sound is trocar-pointed for use when puncture is indicated.

to the metal plate. The clay thus forms one of the electrodes, and is, as nearly as possible, inert. It is important that the clay should everywhere be in close apposition to the skin, and it is, therefore, well to fix it by a few turns of a broad bandage. The other electrode is either:—

- 1. A blunt-pointed platinum sound passed into the uterus, or
- 2. A sharp-pointed trocar made of steel, which is made to penetrate the substance of the tumour to a slight depth, viâ the uterus. In the instrument figured, (1) and (2) are combined in the same instrument (Fig. 83).

By means of a galvanometer (Fig. 84) the exact strength of the current employed can be measured; from 100 to 150 milliampères is that usually employed. The current should be gradually increased up to the highest intensity we mean

to use; it is kept stationary there for a few minutes, say five or ten minutes, and is then *gradually* reduced again to o; the electrodes are then removed, and the sitting is over.

The vagina must be syringed with an antiseptic lotion before and after the treatment on each occasion.

The current may be obtained from a Stöhrer's battery.

The intra-uterine electrode may be made either positive or negative at will.

For cases of fibroids where bleeding is the prominent




Fig. 84.—Horizontal Galvanometer, graduated to register 250 milliampères.

symptom the intra-uterine pole should be positive. For cases of fibroids not attended with bleeding the negative pole is recommended.

Dr. Apostoli advises puncture (1) as a matter of necessity, when for various reasons the sound cannot be introduced; (2) as a matter of choice, "when we see that we can advantageously combine punctures with intra-uterine cauterisation, so as to expedite and make sure of the effects that, with the cauterisation only, we should tardily or perhaps

imperfectly realise." The puncturing instrument is attached to the negative pole.

It is claimed that under this treatment fibroid tumours are markedly reduced in size, and as regards clinical results that "ninety-five times out of every hundred they comprise the suppression of all the miseries constituting the fibroidal symptomatology, which may be thus categorically enumerated:—Hæmorrhages, the troubles of menstruation, dysmenorrhæa, amenorrhæa, nervous disturbances, the direct pains in the growth itself, and from mechanical pressure, and the harassing series of reflex actions."

The minimum time necessary for the treatment is one month, and the applications are made twice a week. All the instruments necessary are supplied by Messrs. Coxeter; they tell me that an improvement of importance is the use of the water rheostat (Fig. 85), "which simplifies the whole operation and renders a *shock* impossible."

Present position of this treatment.—It is desirable that the cases in which it is tried should be carefully chosen; more particularly cases should be taken:—

1. Where the diagnosis of fibroid tumour is beyond question.

In many of the so-called successes reported it has not been at all clear that the cases treated were really cases of uterine fibroid at all. In several, for example, the physical signs reported have not been inconsistent with a diagnosis of subinvolution of the uterus. In other words, many of the cases have not been, to say the least of it, typical cases of uterine fibroids.

By a typical case of uterine fibroid is here meant a case where the uterus is enlarged so as to reach up to, or near, the umbilicus; where the sound passes four or five inches or more, and where hæmorrhage is the prominent symptom.

I have tried the treatment, using the blunt electrode only, in a few typical cases of this kind at the London Hospital. The results have not been such as to convince me of the efficacy of the treatment.




Fig. 85.—Water Rheostat to be introduced in the circuit at an indifferent point for regulating the strength of the current (Coxeter).

The glass is filled with distilled water, and the current is first "turned on" with the poles of the instrument unscrewed to the utmost. As the poles are approximated, the galvanometer indicates the increase of current, which is slow at first, but rapidly increases as the poles are brought near together, necessitating as slow a movement of the screw as in using a microscope with a high power. The poles must be again widely diverged before the sound is withdrawn.

2. Where palliative treatment by rest, hot douching, and drugs has failed. The Apostoli treatment is obviously a more severe measure than palliative treatment of this kind, and it is our duty to relieve our patients by the least severe

means at our disposal. Cases then should only be treated by Dr. Apostoli's method, if at all, when palliative treatment has been tried and failed.

I think there can be no question that puncture of the tumour recommended in certain cases by Dr. Apostoli is a very dangerous and unjustifiable proceeding.

The question being what is the value of electricity in typical cases of uterine fibroid, puncture only introduces a complication. If the case is successful, it is open to the explanation that the success was due to the puncture, and not to the electricity.

It must also be remembered that the Apostoli treatment (without puncture) is one involving considerable risk; several deaths have followed its application; and I myself know of two cases where the patients died soon after the treatment.

It may be added that Dr. Apostoli uses the *faradic* current applied either to the interior of the uterus, or merely to the vagina, for various conditions attended with pain, particularly pelvic inflammation, pains in the ovarian regions, and vaginismus.

I have tried it in several cases of chronic pelvic inflammation at the London Hospital; none of the cases so treated were benefited in the slightest degree.

Case of Pelvic Abscess in an Unusual Position, Simulating Soft Fibroid Tumour of the Uterus.

E. M., aged 36, married, having had eight children and three miscarriages (all before the fourth child was born), was admitted into the London Hospital on June 23rd, 1885, complaining of a swelling in the abdomen, and of a red discharge from the vagina. Her last labour (four weeks previously) was a quick one; she lost a good deal of blood after the child was born, but otherwise the labour was quite normal, and she seemed to recover as usual, getting up on the tenth

day, without having any pain, shivering, or feverishness since her confinement. On the tenth day when she got up she noticed the swelling in the abdomen for the first time; it was then much smaller than it was on admission, but had gradually increased to its present size. On the day she got up she felt so weak that she was obliged to return to bed. When she moved, even in bed, she had a dragging pain in the hypogastrium, but she had no pain as long as she lay still. She had had a red discharge from the vagina ever since her confinement.

On admission the patient was thin and anæmic. The abdomen was somewhat distended. A smooth, elastic swelling was felt rising up from the pelvis to the level of the umbilicus; it was not tender. The swelling was symmetrically situated with regard to the middle line. It was dull on percussion; laterally, beyond the limits of the swelling, the abdomen was resonant. Nothing could be heard on auscultation over the tumour. The patient was put under the influence of ether in order that the relations of the swelling might be more thoroughly determined. Through the speculum a sanious discharge was seen to be issuing from the os externum. A catheter having been passed to make certain the bladder was empty, a swelling was felt in front of the cervix depressing the anterior fornix.

Bimanually this swelling was found continuous with that already noted in the hypogastric region. The uterus was movable; every upward impulse given to the cervix moved the tumour with it. The body of the uterus could not be made out distinct from the tumour. It was thought that the latter extended rather farther towards the left than to the right. The sound was not used on this occasion. The temperature was 101'4° on admission, and varied for some six weeks subsequently from 102° at night (on one occasion 103'5°) to 99° in the morning.

On July 13th the sound was passed; it reached a depth of three inches and a half, and passed towards the right. The tumour was thought to be a soft fibroid.

August 13th.—The patient had been treated since the last note with ergot, at first by the mouth, and afterwards hypodermically, without any alteration taking place in the size of the tumour. A catheter being passed into the bladder, it was found per vaginam that it would be impossible to puncture the tumour in that situation without puncturing the bladder. An aspirator needle was therefore put into the swelling in the middle line, about midway between the umbilicus and the pubes, and a pint of extremely offensive pus drawn off, the

tumour entirely disappearing. Five days afterwards the swelling was as large as at first; it was tapped again on August 22nd, and twenty-two ounces of pus were drawn off, similar in character to the last. After each tapping the abdomen was firmly bandaged.

29th.—The swelling has filled up again, and is, if anything, larger than ever. To-day an opening was made under antiseptic precautions into the swelling, the edges of the abscess wall stitched to the skin, and a large drainage-tube five inches long inserted.

From this time to September 15th the patient went on without a bad symptom, the temperature only once reaching 100°. The wound was dressed about every other day, the cavity of the abscess being washed out at each dressing with iodine water, and some iodoform placed in the deep end of the drainage-tube before reinserting it. The discharge, which had at first been very offensive, became quite sweet. The abdomen was firmly bandaged after each dressing, large pads of lint being placed on each side of the wound.

On September 15th, at 9 a.m., the patient had a rigor lasting seven minutes, the temperature rising to 101°. She complained of pain in her limbs and sore-throat. The pulse was very small and frequent (148). There was no pain in the abdomen, and the wound was looking well. There was a slight erythematous rash on the chest and arms. In the absence of any other discoverable cause, it was thought that these symptoms were due to the absorption of iodoform, which had been very freely introduced into the abscess cavity. The cavity was therefore washed out with carbolic lotion, and iodoform and iodine water omitted. The next day the temperature was normal, and all the symptoms had disappeared.

November 9th.—The temperature has remained normal since last note. The drainage-tube (which had been shortened from time to time, and replaced by narrower ones) was left off altogether to-day. Since October 5th the patient has gained eleven pounds in weight. She feels and looks very well. On November 4th the catamenia came on, and ceased to-day, no red discharge having previously taken place from the uterus for more than three months. The uterus is freely movable, the body of it being drawn to the left side. The sound passes two inches and a half. There is still a small sinus, an inch and a half deep, in the abdominal wall, but there is little or no discharge from it. A fine probe, coated with silver nitrate, was passed into it.

December 31st.—The patient has had a fortnight at Eastbourne. The sinus is the same length as before, but only admits a very fine probe.

She says the last menstrual period lasted a fortnight, but otherwise she has remained quite well, and has maintained her weight.

This patient came to me some months afterwards, as she thought herself pregnant. She had been suffering from pains in the lower part of the abdomen, which she had never before had in her pregnancies.

On examination, she was found to be about four months pregnant. The sinus in the abdomen had completely closed. I consider the pains were due to stretching of inflammatory adhesions round the situation where the abscess had been, as the uterus enlarged, owing to the pregnancy.

#### UTERINE POLYPI.

These are :-

- 1. Fibroid polypi.
- 2. **Mucous polypi.**—According to the late Dr. Matthews Duncan there are three varieties:—
- (i) Adenomatous, where the polypus is composed chiefly of the hypertrophied glands of the mucous membrane.
- (ii) *Molluscous*, where the polypus is formed chiefly by hypertrophy of the connective-tissue element of the mucous membrane.
- (iii) Cystic, where retention cysts are formed from the uterine glands, either of the cervix or the body, and where the little mass resulting, composed of one or several retention cysts, acquires a pedicle.
  - 3. Placental polypi.
  - 4. Fibrinous polypi.
  - 5. Malignant growths of polypoid form.

# FIBROID POLYPI.

These have already been sufficiently considered under Fibroid Tumours, of which they are only a variety.

#### MUCOUS POLYPI.

### Position. Size. Number.

Cystic mucous polypi are merely Nabothian ovules that have become pedunculated, their mode of origin and appearance have been described under Cervical Endometritis. They are mentioned here for the sake of completeness.

The *adenomatous* and *molluscous* varieties are what are usually meant when mucous polypi are spoken of. They occur both in the cervix and in the body, but are commoner




Fig. 86.—Microscopic Section of a Mucous Polypus (De Sinéty). g. Dilated glands; e. Epithelium; mf. Muscular fibre; v. Bloodvessel; ct. Connective tissue.

in the cervix, especially just within the os externum. They have a deep red colour during life, and are extremely soft to the touch. Usually more than one is present at a time. They vary in size from that of a small raisin to that of a large strawberry.\*

\* The largest I have met with was fusiform in shape. After several weeks in spirit its measurements were as follows:—Length,  $2\frac{3}{8}$  in.; Breadth, I in.; Thickness,  $\frac{1}{2}$  in. Before removal about half an inch of it projected beyond the vaginal orifice. It was attached to the cervical canal near the external os. There was a second one, the size of a cob-nut, attached near it.

Mucous polypi are most commonly met with during menstrual life—from fifteen to forty-five. Occasionally, however, they are found in women past the menopause. I have seen several cases of this kind.




Fig. 87.—Four Mucous Polypi growing in the Cervix Uteri (Sir J. Y. Simpson).

**Symptoms.**—Small mucous polypi of the cervix hanging from the os externum are often discovered accidentally, having given rise to no trouble.

The symptoms likely to be present are:—

1. Bleeding.

- 2. A white or yellow discharge.
- 3. Sterility.
- 4. Dysmenorrhæa.

And such symptoms, particularly the last two, may be considered due to mucous polypi with most confidence, when these are large and situated in the body of the uterus.

The bleeding in the case of mucous polypi is said to come from the polypus itself, but probably it comes also in part from the general surface of the mucous membrane of the body of the uterus, at all events when the polypus is attached there. The white and yellow discharge is due to associated cervical or corporeal endometritis.

Diagnosis.—Mucous polypi hanging from the os externum are at once diagnosed by the finger, and brought into view with the speculum. Mucous polypi of the body of the uterus can only be recognised by dilating the cervix, and passing the finger into the uterine cavity. In fact, in cases of hæmorrhage coming from the cavity of the body of the uterus, apart from pregnancy, or any suspicion of malignant disease of the cervix or body of the uterus, we may say the general rule is:—(1) try rest, ergot, and hot vaginal injections; and if these fail after a fair trial, (2) dilate the cervix, and examine the interior of the uterus.

Treatment.—Polypi hanging from the cervix may be twisted off with a pair of tumour forceps (Fig. 88), which should have a catch like Spencer Wells's forceps; or their attachment may be cut through with scissors.

Intra-uterine polypi, recognised only after dilating the cervix, should be either twisted off with tumour forceps, or scraped away with a curette—antiseptic douches being given before and after the operation, and the uterus itself also washed out with a similar lotion through the intra-uterine tube (Fig. 15, p. 36).

PLACENTAL POLYPI-FIBRINOUS POLYPI.

These are not the same. A placental polypus no doubt is usually in part composed of fibrin as well as of placental remains, but it has its origin in labour or abortion, some




FIG. 88.—TUMOUR FORCEPS FOR TWISTING OFF POLYPI.

portions of the placenta or membranes remaining adherent to the uterus, and forming a base to which layers of fibrin attach themselves.

Fibrinous polypus may originate apart from labour or

abortion: for instance, I have known it form after removal of a fibroid polypus attached to the body of the uterus, the fibrin being deposited on the stump left after section of the pedicle. In this case a highly offensive, yellowish, polypoid mass of fibrin protruded from the os externum some days after the fibroid had been removed. Often fragments of the placenta left behind are sessile, attached to the uterine mucous membrane without any stalk.

Symptoms—Treatment.—Bleeding is the symptom that usually leads to investigation; and if the polypus is decomposing, there will also be an offensive discharge. On examination, the uterus will be found to be larger than normal, and we may, perhaps, feel the polypus presenting at the external os, though this is far from being always the case.

The treatment consists in dilating the cervix, and scraping away the polypus thoroughly, or twisting it off, with all the precautions mentioned when describing the treatment of mucous polypi.

# MALIGNANT GROWTHS OF POLYPOID FORM.

These were referred to when considering the diagnosis of fibroid polypi. A simple, non-malignant, papillary growth from the vaginal portion of the cervix has been described; but practically a papillary growth of the vaginal portion of the cervix that is soft, and bleeds easily on touching it, is almost invariably malignant. Malignant growths from the body of the uterus sometimes assume a polypoid form, and may occasionally project through the os uteri into the vagina. I have described further on a case in which I extirpated the uterus for a growth of this kind.

### CHAPTER XI.

## CARCINOMA OF THE CERVIX.

#### ETIOLOGY.

Age.—It is to be noted that carcinoma of the cervix is not rare in comparatively young women; we meet with it not very infrequently in patients of twenty-six or twenty-seven, and after thirty it becomes quite common.

Taking Gusserow's statistics (quoted by Hart and Barbour), based on 2270 cases, the exact percentages are as follows:—

Between the ages of 20 and 30 ... 3.5 per cent.

Fertility.—Most cases occur in those who have had large families. Winckel, basing his calculations on 130 cases, gives the average fertility at 5.6 children; and says, moreover, that only 1.7 per cent. of women with carcinoma of the uterus are absolutely sterile.

How far, if at all, the lacerations of the cervix that occur during labour predispose to carcinoma is not yet decided.

Heredity.—The number of cases in which heredity can be traced is so small—only 7 or 8 per cent.—that it cannot be regarded as proved to have much influence.

Depression of the general health.—This is regarded as a predisposing cause; the fact that carcinoma is more common among the poor than among the well-to-do is explained on this ground.

Race.—Carcinoma is much less frequent among the black races than among the white.

#### PATHOLOGY.

It would be out of place to enter into the question as to whether carcinoma originates from connective-tissue cells, or from pre-existing epithelium. The view taken here will be that which obtains most acceptance in this country, viz., that carcinoma always originates from pre-existing epithelium. Accordingly, as regards the cervix, we may have it arising from the squamous epithelium of the vaginal portion, or from the cubical epithelium within the cervical canal, either that on the surface, or in the deeper parts of the cervical glands.

# Clinical types.

The cauliflower excrescence.—This originates from the squamous epithelium of the vaginal portion, and in its early stages forms a flat, or somewhat papillary, growth from one or both lips of the cervix. These become thickened, and the growth tends to spread to the vaginal walls. Ultimately a mass, the size of the fist or larger, is produced, occupying the upper part of the vagina. The surface of it has a considerable resemblance to the surface of a cauliflower, hence the name.\* The position of the external os in such

<sup>\*</sup> The term "cauliflower excrescence" was originally given to malignant growths, where the papillary, or villous, nature of the surface was a prominent feature. Such growths are much rarer than the form to which the name cauliflower excrescence is given in the text, which moreover is actually more like the white part of a cauliflower,

cases is often difficult to find. Sometimes a mass such as that described has an indistinct pedicle.

The mushroom-shaped variety.—This also is a squamous epithelioma, probably, indeed, it is often a stage towards the production of the typical cauliflower excrescence. Still it is a variety often met with clinically. The vaginal portion is enlarged at its lowest part by thickening of the lips of the




Fig. 89.—Cauliflower Excrescence of the Cervix (Simpson). (A more villous growth than those commonly met with.) Compare this specimen, where amputation of the vaginal portion of the cervix only has been performed, with figures at the end of the chapter where supra-vaginal amputation was done.

cervix, and by the spreading over them of the flat raised epitheliomatous growth from the neighbourhood of the external os outwards towards the circumference. The finger passed outwards from the situation of the os, comes at the periphery of the growth to an overhanging margin, separated by a shallow depression from the part of the cervix not yet affected. The whole vaginal portion, therefore, has a rough resemblance to an inverted mushroom

(Fig. 92). Sometimes one lip only of the cervix is affected by a growth of this character.

The conical ulcer.—In the situation of the external os and replacing it, the finger finds an ulcer, more or less conical in shape, its walls irregular, and formed of hard tissue. The diameter of the ulcer often exceeds considerably the




Fig. 90.—Cauliflower Excrescence growing from one lip of the Cervix Uteri (Sir J. Y. Simpson).

diameter of the healthy vaginal portion. This is due to enlargement of the vaginal portion from the malignant infiltration, followed by ulceration spreading from the centre outwards in all directions. Sometimes, indeed, the whole vaginal portion has in this way been destroyed, and the ulcer is bounded by a hard ring of tissue formed by infiltration of the vaginal walls.

The nodular variety.—Little hard nodules form beneath the mucous membrane, and project either towards the



Fig. 91.—Carcinoma of the Cervix (Schroeder); ulcerating form. The disease is in an early stage. Later, the conical, or crater-like, ulcer described in the text would be produced in such a case.




Fig. 92.—Carcinoma of the Cervix (Schroeder); mushroom-shaped variety. The disease affects both lips of the cervix, so as to cause the vaginal portion to have a rough resemblance to an inverted mushroom.

cervical canal, or elevate the squamous epithelium of the vaginal portion in the neighbourhood of the external os. As they increase in size and reach the surface, they




Fig. 93.—Carcinoma of the Cervix (Schroeder); nodular variety. To the right is seen a section of the cervix through the nodule.




Fig. 94.—Microscopic Section of a portion of the Cervix Uteri seen in the last figure. a. Squamous epithelium in several layers; cn. carcinomatous nodule; between these is seen a portion of inflamed mucous membrane covered with a single layer of epithelium (Schroeder).

ulcerate. This variety is less common clinically than the preceding, partly no doubt because it is a phase of the disease at an early stage.

When ulceration has taken place, it merges sooner or later into the excavated form described in the last paragraph as the "conical ulcer."

If the nodules project only towards the cervical canal, we have an example of carcinoma beginning within the canal, and such a case might, on hurried examination, easily escape detection, as the vaginal portion and os would appear normal. Usually, even where it begins within the canal, it is not far above the os externum, and on everting the lips of the cervix with tenacula something of the disease will be seen. Such cases can rarely, if ever, be diagnosed, previous to the time when ulceration has begun.

# DIRECTIONS IN WHICH CARCINOMA OF THE CERVIX TENDS TO SPREAD.

It may spread:-

- 1. Downwards on to the vaginal wall.
- 2. Outwards into the connective tissue round the cervix, into the broad ligaments and utero-sacral ligaments.
- 3. Upwards along the cervical canal into the body of the uterus.

There is usually a greater tendency for the growth to spread to the vagina, and to the peri-cervical connective tissue, than to the body of the uterus. In a few cases the growth does extend upwards to the body of the uterus before it has spread to the vaginal walls, or the peri-cervical connective tissue; these cases are quite the exception, and it is for them especially that extirpation of the whole uterus for malignant disease of the cervix is to be preferred to supra-vaginal amputation of the cervix. In advanced cases

the ureters become dilated from obstruction, partly due to pressure, and partly to the walls of the ureters themselves being infiltrated; ultimately hydronephrosis may be produced.

As the growth that has involved the anterior vaginal wall ulcerates, we may have a vesico-vaginal fistula; more rarely, from similar processes involving the posterior wall—recto-




Fig. 95.—Carcinoma of the Cervix, an advanced case where ulceration has been a prominent feature. The bladder and rectum communicate freely with the vagina (Farre).

vaginal fistula: still more rarely, perforation into the peritoneal cavity may occur.

In advanced cases where ulceration has been a prominent feature, the bladder and rectum may communicate freely with the vagina (Fig. 95).

Symptoms.—The special local symptoms of carcinoma of the cervix are:—

1. Bleeding.

- 2. A discharge, which sooner or later becomes offensive.
- 3. Pain.

Other local symptoms not infrequently present are:—
painful, difficult, or frequent micturition, and pruritus of the
vulva.

Sometimes wasting precedes all other symptoms, the patient noticing that she is getting thinner without apparent cause. In any case, once local symptoms are present, she rapidly loses weight, and becomes conscious of a sense of weakness and general ill-health, till ultimately she presents the appearance known as the cancerous cachexia.

It should be noted, however, that cachexia is only present in cases that are far advanced. The patient may look quite well for a considerable time in cases of cancer of the cervix. Nothing can be more misleading than to conclude that a patient cannot have cancer of the cervix because, as regards the face, she looks well.

Bleeding.—In most cases hæmorrhage, occurring at some time other than a menstrual period, is the earliest symptom to attract attention. Occasionally a persistent white or yellow discharge, increasing in quantity, and ultimately becoming offensive, may be the symptom that first attracts attention, there being no unusual bleeding till the disease has reached a somewhat advanced stage. This is, however, quite the exception. Hæmorrhage after coitus is particularly significant of carcinoma of the cervix. It may be taken as a general rule that hæmorrhage coming on some years after the menopause is almost always due to malignant disease.

The discharge.—In early cases before ulceration has taken place, the discharge is not offensive; at this time it is likely to be yellowish, and more or less streaked with blood. Later, when ulceration has set in, the discharge does become horribly offensive, and at this time it is usually thin,

and like dirty water. It should be especially noted that an offensive discharge is not pathognomonic of carcinoma. A sloughing fibroid polypus, for instance, produces a discharge every bit as offensive as any that occurs in cases of carcinoma; and on the other hand, as mentioned above, in early cases of carcinoma, the discharge is not offensive.

Pain.—This is a late symptom in carcinoma of the cervix; when at length it appears, it is extremely severe, and usually worse at night, so that the patient cannot sleep without morphia.

Fever.—When ulceration has occurred, and the discharge has become offensive, there is fever, due to absorption of the chemical products of putrefaction—septic intoxication, or sapræmia. The result of scraping away the diseased tissue as completely as possible shows this; for whereas before the scraping the temperature chart has shown the presence of moderate fever, after the scraping the temperature falls, and often remains normal till fresh portions of the growth necrose and putrefy.

In advanced cases a condition of one or both lower extremities, indistinguishable from the phlegmasia dolens of puerperal women, is not uncommon.

Physical signs.—On vaginal examination with the finger and speculum, we may meet with any of the conditions already sufficiently described, viz.:—

The cauliflower excrescence.

The mushroom-shaped variety.

The conical ulcer.

The nodular variety.

Even gentle examination with the finger almost always causes some bleeding. The speculum should, as a rule, only be used in early cases; if it is desirable to pass a speculum in advanced cases, Sims's speculum is the best

for the purpose, care being taken to avoid touching the diseased tissue as far as possible.

In cases where the disease has commenced a little distance above the os uteri, it may be necessary to evert the lips of the cervix with tenacula, in order to see any evidence (e.g., ulceration) of the disease.

In such cases, when the finger is pressed firmly against the suspected area, the tissue will usually be found to break down if affected by a malignant growth.

Mobility of the uterus.—In early cases the uterus is freely movable. Later, as the growth spreads to the vagina and connective tissue round the cervix, it becomes less and less movable; finally, in advanced cases, when the infiltration has advanced along the broad ligaments, and utero-sacral ligaments, to the pelvic wall, the uterus becomes completely fixed.

When the disease has reached this stage, on abdominal palpation we can easily feel a hard mass in the hypogastric region occupying the brim of the pelvis.

Diagnosis.—In most cases the diagnosis is easily made, because most cases come under observation at a comparatively advanced stage, when the physical signs are characteristic. In early cases carcinoma of the cervix has to be distinguished from the following:—

· Non-malignant erosions of the cervix, especially those that bleed easily when touched.

Chronic inflammatory induration of the cervix with occluded follicles.

Sloughing fibroid polypi of the cervix. Condylomata of the cervix.

Chancre of the cervix.

Non-malignant papilloma of the cervix. Small interstitial fibroid of the cervix. Corroding ulcer of the cervix. These in practice rarely give rise to any difficulty in diagnosis. The first two are the only conditions in which any real difficulty is likely to arise.

I. Non-malignant erosions usually improve rapidly, even if they cannot be completely cured, by suitable local treatment. The tissue of an erosion does not break down when pressed upon by a blunt instrument such as the sound.

2. Chronic inflammatory induration of the cervix with occluded follicles.—This may occasionally suggest the nodular variety of carcinoma, the little hard lumps formed by the ovula Nabothi simulating nodules of carcinoma. If such a lump is an occluded follicle, puncture gives exit to fluid, and the hard lump disappears. Again, the history is different, extending often over a long period. Here also, in case of doubt, weighing the patient will settle the question very soon.

It is as well to say here, that most cases, in which any doubt is felt by one familiar with the subject, turn out to be non-malignant. This is particularly worth noticing in connection with the two conditions just referred to, which are those in which temporary difficulty in diagnosis most often arises.

Chancre on the cervix or condylomata on the cervix are rare. Suitable local and constitutional treatment, e.g., injections of lead lotion, and mercury internally, would soon make the diagnosis clear. In the case of condylomata there would probably be other evidences of syphilis present at the same time elsewhere, e.g., round the vulva, anus, in the throat, etc.

Sloughing fibroid polypus.—I have known a polypoid malignant growth of the cervix taken for a sloughing fibroid polypus. The differential diagnosis has been described in the chapter on Fibroid Tumours (see page 204).

Non-malignant papilloma.—If there is such a thing, it is extremely rare, and certainly it is right to treat every case

of papilloma of the cervix as if it were malignant. The only exception to this is where there are warts, evidently of a simple nature, also present on the vaginal walls and on the vulva.

Small interstitial fibroid of the cervix.—So long as no ulceration has occurred, a positive diagnosis cannot at once be made. Watching the case for a short time, and paying attention to the weight of the patient, would soon settle the matter. Fibroids in this situation are not at all common.

Corroding ulcer of the cervix.—This is a very rare condition, the exact nature of which is doubtful; it would be right, however, to treat it as if it were a carcinomatous ulcer.

To sum up as regards differential diagnosis.— In practice diagnosis is seldom difficult; in cases where we are in doubt the condition almost always turns out to be non-malignant. When in doubt, careful attention to the patient's weight at short intervals, with suitable local treatment to the suspicious surface, combined with constitutional treatment, if there is a suspicion of syphilis, will soon clear up matters.

Duration of the disease.—The duration of life from the time the patient first seeks advice, may be taken on an average as from a year to a year and a half. Some cases only live a few months, others may live as long as three or four years. Winckel mentions three years and eight months as the longest duration he has met with.

Causes of death.—These are:—

Peritonitis in 25 per cent. of cases (Winckel).

Uræmia "45 " " "

Exhaustion in most of the remaining cases.

Rarely death may be immediately due to:—

Hæmorrhage.

Embolism.

# Treatment.—In early cases.

The supra-vaginal amputation of the cervix.—When we meet with a case early enough to be able to remove the whole of the obviously diseased part, plus a surrounding shell of apparently healthy tissue (the broader the better), the right treatment is by the operation known as the supra-vaginal amputation of the cervix. The question is often a nice one as to whether, in a particular case, the disease is too far advanced or not for this operation.

We may consider it indicated in cases where the disease is limited to the tissues of the cervix, and has not extended either on to the vaginal walls, or into the connective tissue round the cervix. Only those who are in the habit of performing the operation can decide in doubtful cases whether the disease is too far advanced for it to be possible to make the lines of section sufficiently wide of the disease.

Extirpation of the uterus.—In the small minority of cases where the disease has extended up the cervical canal to the body of the uterus, without spreading either to the vaginal walls, or into the peri-cervical connective tissue, the right treatment is extirpation of the whole uterus. For the large majority of cases of malignant disease of the cervix, where the tendency, as already mentioned, is to spread to the vagina, and connective tissue round the cervix, and not to the body of the uterus, it would seem à priori to be illogical to remove the whole uterus. Yet it is believed by some authorities that recurrence is more likely to take place at an early period after supra-vaginal amputation than after extirpation of the whole uterus (see Berry Hart, in Year-book of Treatment, 1889). This question has yet to be settled.

As, however, the mortality of vaginal hysterectomy has diminished, it has become justifiable to perform the operation in cases of cancer of the cervix where the patient

seems well able to stand a severe operation, where all the conditions are favourable, and where the diagnosis is absolutely indisputable.

In reference to this point the following quotation from a leading article in the *Lancet* of December 27th, 1890, is well deserving of attention:—

"There appears to be a movement in Germany in favour of hysterectomy rather than supra-vaginal amputation, but the enormous numbers of cases recorded seem to us somewhat suspicious. We venture to say that such numbers recorded in towns which are mere villages compared to London, would not be equalled by the records of all the London Medical Schools put together."

## DESCRIPTION OF THE SUPRA-VAGINAL AMPUTATION.

The patient being in the lithotomy position, a vaginal douche of some efficient antiseptic (e.g., iodine water) is given. Sims's speculum is then passed, and the cervix drawn down to the vulva.

Drawing down the uterus.—For drawing down the cervix, if the disease affects both lips, it is best to use volsellæ; at least two should be at hand. It is very important that the traction should be as steady as possible when once a good hold has been obtained. A glance at Fig. 96 will show the relation of the cervix to the ureters.

## Anterior incision.

Separation of the bladder from the supra-vaginal cervix.

—The cervix is carried well back towards the perineum, care being taken to keep it in the middle line; a sound is now passed into the bladder, and the limits of the bladder towards the anterior lip of the cervix ascertained. Guided by this information, a transverse incision is made along the line a b (Fig. 97) through the anterior vaginal wall;

the incision has a slight convexity towards the pubes. There should be at least 1 cm. ( $\frac{2}{5}$  inch) of apparently healthy mucous membrane in each direction, between the incision and the diseased area. I prefer to use blunt-pointed scissors for making the incision. It is important




Fig. 96.—Diagram to illustrate relations of the ureters to the cervix when the patient is in the lithotomy position, with special reference to the operations of supra-vaginal amputation of the cervix, and extirpation of the uterus.

In seven specimens examined with reference to the distance ab, between the orifices of the ureters in the bladder, it was found that this varied from  $\frac{9}{16}$  inch to 2 inches, average value of  $ab = 1\frac{1}{14}$  inch.

In three specimens the distance dc was carefully measured, *i.e.*, the distance from the cervix to a horizontal line joining the orifices of the ureters; in two specimens it measured  $\frac{3}{4}$  inch, and in one specimen I inch.


In two specimens the distances ef were carefully determined, *i.e.*, the distance from the side of the cervix to the ureter in a horizontal line passing through the middle of the cervix; in one specimen ef was  $\frac{3}{4}$  and in the other "rather more than  $\frac{3}{4}$  inch."

On referring to the figure, p. 252, illustrating the incisions for the supra-vaginal amputation of the cervix, it will be seen that there is ample room for them without any risk of injuring the ureters.

to feel one's way at frequent intervals with the sound in the bladder during this step of the operation. In some cases, where the disease affects the anterior lip, the separation of the bladder from the cervix is rendered easier by dilating the urethra, and passing the forefinger of the left hand into the bladder. There is an interval of comparatively loose connective tissue between the supra-vaginal cervix and the bladder; and when this has been entered, most of the necessary separation can be effected with the finger. When we judge that we have cleared the cervix sufficiently in front, we proceed to the next step.

#### Posterior incision.

Separation of the peritoneum from the posterior aspect of the cervix.—The cervix is now carried well forwards towards the pubes. A transverse incision along the line cd (Fig. 97) is then made through the posterior vaginal wall; and keeping close to the cervix, we dissect the peritoneum from its



posterior aspect as high as is necessary to correspond to the height to which the anterior aspect has been cleared. At this stage it is very likely that the peritoneal cavity will be opened—button-holed; but this is not of much importance, and fear of doing it should not prevent a sufficiently high separation being made. The cervix has now been cleared, anteriorly and posteriorly, but as yet the lateral attachments remain.

# Preliminary lateral incisions.

Lateral incisions should be made on each side, along the lines ac, bd; but they should at first be only mucous membrane deep; made, in fact, merely as an indication where the actual separation is to be made when the time comes.

Lateral ligatures.—A strong blunt-pointed pedicle needle armed with a strong silk ligature should then be passed from behind forwards, so as to include the whole lateral attachment. It should be made to encircle the whole lateral attachment of the cervix up to the height at which it is proposed to amputate it. This is done on each side.

The ligatures are tied as tightly as possible; and as they are very apt to slip before the second hitch is made, the best plan is to hold the first knot with Wells's forceps while the second hitch is being tied. The cervix can now be freed from its lateral attachments without fear of hæmorrhage, if the ligatures have been properly placed, and securely tied. It is, however, not uncommon for the ligatures to loosen, and it is desirable to have Wells's forceps ready to seize any large vessel that bleeds as the lateral attachment is divided. Large branches of the uterine artery lie in this attachment, and it is to control them that the precautionary ligatures described are used. Instead of using ligatures, Wells's forceps may be used before cutting the lateral attachments; they may be left on for forty-eight hours.

Removal of the cervix.—The cervix has now been made clear all round. To remove it, we incise the anterior wall till the lumen of the cervical canal is seen (the part of the cavity exposed should be at the level of the internal os); we then pass one or two silver wire sutures with an Emmet's needle, from the anterior margin of the stump through the cut edge of the anterior vaginal wall, and either twist them, or, better, fix them with Aveling's coil and shot. We have thus a hold on the stump. We can now complete the removal of the cervix, and stitch the posterior margin of the stump to the cut edge of the posterior vaginal wall in a similar manner. Stitching as described is not essential.

The cervix removed should be slit up, and examined,

to make sure we have removed it sufficiently high above the disease.

Any Wells's forceps on big vessels may be left on for forty-eight hours. The vagina is once more douched out, a strip of iodoform gauze is placed in the vagina, and the patient sent back to bed.

After the gauze is removed, antiseptic douches may be given twice a day. I prefer not to douche till the end of the week in cases where the peritoneum has been opened.

The operation of extirpation of the uterus will be described in the treatment of carcinoma of the body of the uterus.

## Other modes of treatment.

The supra-vaginal amputation of the cervix, just described, is of comparatively recent origin. Before it was introduced, amputation of only the vaginal portion of the cervix with the écraseur was considered the best that could be done.

When we remember that the whole cervix is on an average about an inch long, and the vaginal portion of the cervix only about  $\frac{1}{3}$  inch long, it will be evident that any operation by which the vaginal portion only is cut off, can very rarely even remove all the obviously diseased tissue, let alone leave a good width of healthy tissue between the line of section and the diseased structures.

So that cases suitable for the supra-vaginal amputation, include all in which amputation of the vaginal portion only would be suitable, and many others where the latter operation would be useless.

In certain cases the écraseur may be used merely as a preliminary step with advantage: for example, where a large cauliflower-like mass blocks the vagina, and yet appears to spring from the cervix by a constricted part, or spurious pedicle. Here it is convenient to first get away the bulk

of the growth with the écraseur, and then examine the condition carefully, to see if all the diseased parts can be removed by proceeding to the supra-vaginal amputation. When using the écraseur as described, it is important to make no downward traction on the mass, or a part of the bladder may be caught in the wire loop; similarly a portion of Douglas's pouch may be nipped in, and removed. In one case of mine, where a cauliflower growth had extended on to the posterior vaginal wall, this latter accident occurred; a piece of Douglas's pouch, an inch broad from side to side, was removed with the malignant mass. The opening was stitched with silver wire, bringing the edges of the peritoneum together, and no harm resulted. The supravaginal amputation is a difficult operation, but not a very dangerous one. Hofmeier (quoted in the Yearbook of Treatment, 1884) gives statistics of 105 operations done in Schroeder's Clinique, with the mortality of 12'3 per cent. I have done the operation 31 times, the patient in each case making a good recovery.

My rule has been to perform the operation whenever there seemed a fair chance that the incisions necessary to remove the cervix could be made through apparently healthy tissue, and so it has happened that in several of my cases the disease has been in a considerably advanced stage when the operation was done. I am still inclined to think, however, that the rule I have adopted is the right one, because, of two cases operated on, the disease being apparently at the same degree of advancement in each, in one recurrence may take place in a few months, while in the other there may be an interval of one or two years before the disease returns. For example, in Case 12, in my series, it is now nine years since the operation, and the patient remains quite well. In this case the disease chiefly affected the posterior lip, and was so far advanced

that, judging from other cases, I anticipated recurrence in six months at latest.

As regards recurrence in the other cases :-

No. 3 was known to be in good health, and free from recurrence, two years after operation. I have not seen her, or heard of her, since. Disease in an early stage when operated on.

No. 6 was still in good health, and free from recurrence, six years after the operation. I have not heard from her very recently.

This specimen and sections of it were shown at the meeting of the Royal Medical and Chirurgical Society in December 1892, when my paper was read "On Some Points in the Supra-vaginal Amputation of the Cervix Uteri for Cancer, with Special Reference to the Methods adopted in those Cases where for Two Years and upwards the Disease has not recurred."

No. 9 was in good health two years after operation. Disease moderately advanced at time of operation. Symptoms of recurrence appeared about two and a half years after the operation.

No. 12, as already mentioned, is still free from recurrence, though it is now *nine years* since the operation. Disease considerably advanced at time of operation. The specimen from this case and sections of it were exhibited at the meeting of the Royal Medical and Chirurgical Society on December 13th, 1892. The growth was a squamous-celled epithelioma.

No. 4 was in good health and free from recurrence till nearly twelve months after operation. Recurrence took place in the pelvis some distance from the stump, which remained unaffected to the last. Disease considerably advanced at time of operation.

No. 8 was in good health, and free from recurrence for

nine months after operation. Disease moderately advanced at time of operation.

Case 16 is still quite well and free from recurrence, it being eight years since the operation. This patient came for several years about once a fortnight to see me at the London Hospital, and I still see her there occasionally, and she is examined from time to time. There has never been the slightest sign of recurrence. The specimen and sections of it were shown at the meeting of the Royal Medical and Chirurgical Society in December 1892, when the paper already alluded to was read. The growth is a well-marked example of columnar-celled epithelioma.

Case 17 is, I have reason to believe, quite well and free from symptoms of recurrence, it being more than seven years since the operation. The growth was a columnar-celled epithelioma. The specimen and sections of it were shown on the same occasion as the specimen from Case 16.

Case 18 was well for more than a year, then symptoms of recurrence appeared, and the last time I saw her there was a large, evidently malignant, ulcer at the top of the vagina. A point of special interest in this case is that the microscopical appearances of the growth were not such as to make it clear that the case was one of cancer. The sections only showed a glandular structure. The clinical characters of the growth were, however, decidedly malignant. The recurrence places the malignancy beyond doubt. The case shows that it is not always safe to pronounce upon the nature of a growth upon the microscopical appearances alone; the clinical characters must be taken into consideration as well.

Case 20 is still quite well—four years and a half since the operation.

As regards Cases 22-31, my intention is to record them fully elsewhere, so that I will say no more about them here.

# Cases of the Supra-vaginal Amputation of the Cervix for Carcinoma.

In the following case the patient remained free from any sign of recurrence for a year and two months after supravaginal amputation of the cervix had been performed; then recurrence took place on the anterior part of the stump, and adjacent anterior vaginal wall; this was cauterised very freely with Paquelin's cautery, and two years from the date of the original operation there was no evidence of disease to be seen, and the patient was quite well.

(No. 3 in the series of 31). E. H., age 41, married twice, had two children by her first husband, no children by her second husband, to whom she had been married fifteen years. She had never had any miscarriages. Admitted to the London Hospital on February 6th, 1886, complaining chiefly of a thick yellow discharge, which she had had for four or five years; during the last six months, however, this discharge had been coloured red, at times when she was not menstruating.

On examination, a soft, papillary growth was seen growing from the posterior lip of the vaginal portion of the cervix. The area occupied by the growth was about the size of a sixpence. The growth bled readily on touching it. The sound passed a distance of 23 inches.

February 12th, 1886.—Supra-vaginal amputation of the cervix was performed. The piece removed was 1½ inch long. The patient recovered without any bad symptom, and went home. Subsequently she came up regularly to the hospital, and nothing suspicious was seen till April 25th, 1887. On examination then, I found a red patch on the scar, about ½ inch wide, immediately to the left of the entrance to the uterine cavity; it had not been there previously. This was watched carefully; it gradually increased in size, and on October 8th, 1887, the patient was readmitted. At this time there was a papillary growth over an area the size of a florin; the growth involved the anterior part of the uterine stump and adjacent anterior vaginal wall. The edge of the growth was overhanging. Gentle examination caused bleeding, and brought away fragments of the growth the size of shot.

On October 13th, 1887, a careful examination under ether was made. It was found to be impossible to cut out the recurrent patch, as most of it was on the anterior vaginal wall, and cutting it out completely would have involved cutting out a piece of the bladder. The whole area occupied by the growth, together with a margin of healthy tissue round it, was, therefore, very freely cauterised with Paquelin's cautery.

On January 24th, 1888, she came up to see me; she has been regular since leaving the hospital, and has had no inter-menstrual discharge of any kind. On examination, I found that the ulcer caused by the cautery had completely healed, and there was no trace of the disease either there or elsewhere.




Fig. 98.—Cervix removed by the supra-vaginal amputation in the case of E. L. a points to the healthy mucous membrane on the anterior lip; b, to a mark left by Spencer Wells's forceps on the specimen; c, to the situation of the continuation of the cervical canal at the top of the conical ulcer; d. The posterior lip.

In the next case the disease was at a more advanced stage at the time of the operation.

(No. 4 in the series of 31). E. L., age 33, six children, the last about four years ago, and one miscarriage, which occurred four months ago, when she was six months pregnant, was admitted to the London Hospital on July 19th, 1886, complaining of having been losing blood ever since the miscarriage. The loss has never stopped more than a day or two during this time.

State on admission, July 19th, 1886.—In the situation of the external

os is a cavity admitting the finger as far as the root of the nail (Fig. 98). The walls of the cavity are formed by hard tissue. Examination causes a little bleeding, and the finger on withdrawal has an offensive odour. In passing the speculum the appearance shown in Fig. 98 is seen. The uterus is freely movable. Most of the excavation is in the anterior lip of the cervix, but there is superficial ulceration of the posterior lip also.

Operation on July 20th.—The supra-vaginal amputation of the cervix was performed. At first sight it appeared as if the disease had extended so close to the bladder that separation would be impossible. I



Fig. 99.—Side view of the same specimen. Case of E. L. a. Anterior lip; b points by three lines to thickening in the anterior wall of the cervix corresponding to the extent of the excavation seen in the preceding figure.

therefore dilated the urethra with Hegar's dilators, and held the wall of the bladder between the left fore-finger and thumb, while the cervix was drawn backwards by an assistant; in this way the bladder was dissected off the anterior aspect of the cervix. No permanent incontinence of urine resulted. In clearing the peritoneum from the posterior aspect of the cervix, Douglas's pouch was opened, the opening being a quarter of an inch long.

The operation was done at 10 a.m. Soon after its completion the patient had a rigor, and at 4 p.m. the temperature rose to 103° (the

pulse, however, only being 76), falling to 98° by 8 a.m. on July 21st. In the evening of July 21st the temperature was 100'2°; after that it remained normal.

The patient left the hospital on August 17th; her weight then was 11 st.  $9\frac{1}{2}$  lb. November 18th, weight 12 st. 2 lb. December 2nd, weight 12 st.  $\frac{1}{2}$  lb.

The patient was seen in January and March 1887. The cicatrix was perfectly healthy, and her general health good. She next came on July 16th, 1887, and complained that she was suffering from great pain in the right hip, and down the right leg. This had come on quite recently, and was preceded by numbness in the leg. The cicatrix was perfectly healthy, and the remaining part of the uterus freely movable. In the right posterior quarter of the pelvis there was felt a lump the size of a cherry, hard and fixed. There was an inch or more of healthy tissue between this lump and the uterus.

Readmitted to the hospital, September 11th, 1887.—Has continued to suffer severe pain in the right hip and leg. For some weeks past the right lower extremity has been swollen. On admission it was seen to be considerably larger than the left; great tenderness in the region of Scarpa's triangle. There was, in fact, phlegmasia dolens of the right lower limb. The scar left by the operation was still quite healthy, and the uterus freely movable; the lump in the right posterior quarter of the pelvis was, however, now the size of a walnut, but there was an interval of healthy tissue between it and the uterus.

From this time she gradually went downhill, and was never free from pain except when under the influence of morphia. She often required as many as five injections of one-third of a grain each in the twenty-four hours. On October 18th she weighed only 8 st. 13 lb.

November 7th.—A little hard nodule was noticed in the skin on right side below the breast, freely movable over the subcutaneous structures, but adherent to the skin, which was red over it.

On November 14th she weighed 8 st.  $5\frac{1}{2}$  lb. December 20th, left hemiplegia, and facial paralysis of left side. Sensation apparently normal. Patient has not been noticed to be unconscious. She is not very clear in her understanding. December 29th, 1887, died. No post mortem. There was never any recurrence in the scar left by the operation, nor any lump perceptible on simple abdominal palpation alone, but the lump in the right side of the pelvis became somewhat larger before her death. She had been troubled with occasional vomiting since her readmission.

The next case illustrates the treatment of cancer of the cervix complicated by pregnancy, the malignant growth not having passed beyond the reach of radical treatment. It also shows that Douglas's pouch may be opened freely with impunity, if proper precautions be observed.




FIG. 100.—Case of L. E. M. Cauliflower Excrescence removed in the preliminary operation with the écraseur, December 29th, 1887.

This specimen is an example of the common variety of cauliflower excrescence, which has a convex, slightly irregular surface. The tissue is soft, and can easily be broken down by the finger.

(No. 10 in the series). L. E. M., age 39, married twelve years, has had seven children, and one miscarriage at the sixth month of pregnancy. The miscarriage was nine months ago. A month afterwards menstruation appeared; it occurred again after an interval of three weeks, and again after an interval of two weeks. Since then she has been bleeding

every few days. There have been floodings which last a day or two, and between the floodings she has had a brown watery discharge. She has had no pain to speak of. There has been no trouble with micturition or defectation, and the appetite has been good. Latterly, she has got thinner. She was admitted to the London Hospital on December 21st, 1887. On examination, a globular swelling, reaching a hand's breadth above the pubes, was felt rising out of the pelvis.




FIG. 101.—Cervix removed by the supra-vaginal amputation. Case of L. E. M. a. The diseased part to which the cauliflower growth seen in the preceding figure was attached. The disease affects the posterior lip. b b b. Cut edge of healthy vaginal mucous membrane. This edge is continuous all round, but it cannot be seen posteriorly, owing to the oblique position of the specimen. c. Bristle passed into the cervical canal. d. Upper end of the specimen corresponding to the level of the internal os.

It may be said at once that this was the pregnant uterus, the pregnancy having advanced to about the end of the fourth month, though the patient had no idea she was pregnant.

A large cauliflower-like mass was felt in the vagina, springing from

the posterior lip of the cervix, its attachment being rather to the left of the middle line. Anteriorly the mass was attached to the cervix by a sort of spurious pedicle, but posteriorly it was continuous without any constriction with the posterior lip of the cervix. The growth bled very freely on touching it. The vaginal portion, where not involved by the growth, was of a purple colour.

December 29th, 1887.- I removed the greater part of the mass in the




FIG. 102.—Cervix removed by supra-vaginal amputation. Case of L. E. M. Posterior aspect of the specimen. ppp. Quadrilateral piece of peritoneum removed with the cervix. This piece of peritoneum formed part of Douglas's pouch. c. Bristle passed through the cervical canal.

vagina with the écraseur, with the intention of proceeding at once to the supra-vaginal amputation. The vascularity of the parts was, however, so great, that I judged it would be safer to empty the uterus first, and allow time for the involution of the uterus and vagina to diminish the blood supply. Accordingly I dilated the cervix rapidly with Hegar's dilators, and removed the fœtus, placenta, and membranes at the same sitting; altogether the patient was about an hour

under the anæsthetic. Antiseptic douches were used before, during, and after the operation, the uterus also being washed out. No bad results followed. To promote involution the patient took ergotine, and had hot douches.

On January 16th, 1888 (eighteen days after emptying the uterus), I performed supra-vaginal amputation of the cervix. I opened Douglas's pouch to an extent of about two inches in a transverse direction, and was thus enabled to get well above the disease. A somewhat quadrilateral piece of the peritoneum covering the posterior aspect of the supra-vaginal cervix was removed, adherent to the specimen (Fig. 102).

The interval allowed for involution to proceed before undertaking the operation, must no doubt have diminished the blood supply; nevertheless, at the time of the operation, the parts were still very vascular; this will be evident from the fact that the patient went back to bed with three pairs of Spencer Wells's large pressure forceps, and six pairs of the ordinary size, left on. They were removed forty-eight hours later. It should be mentioned that the opening in Douglas's pouch was closed with three silver sutures. Great care was taken to do everything as antiseptically as possible, but of course the spray was not used. The patient had moderate fever for a few days after the operation, but after the first two days there was nothing in her condition to occasion anxiety. The sutures were removed on the tenth day.

Fig. 100 shows the appearance of the cauliflower mass removed with the écraseur on December 29th, 1887.

Fig. 101 shows the cervix after removal by the supra-vaginal amputation on January 16th, 1888.

Fig. 102 shows the posterior aspect of the same specimen, and the quadrilateral piece of peritoneum removed with it.

# Twenty-one cases of the supra-vaginal amputation of the cervix for malignant disease.

No.	DATE OF OPERATION.	AGE.	Complete or Incomplete.	STAGE OF DISEASE.	RECURRENCE.
1	July 27th, 1885.	29	Complete	Moderately advanced.	Remained well for 6 months, then re- currence in pelvis.

_					
No.	DATE OF OPERATION.	AGE.	Complete or Incomplete.	STAGE OF DISEASE.	RECURRENCE.
2	August 6th, 1885.	43	Incomplete.	Considerably advanced.	The second
*3	February 12th, 1886. (For full notes, see p. 258.)	41	Complete.	Early.	Quite well on Jan- uary 24th, 1888. Not seen or heard of her since.
4	July 20th, 1886. (For full notes, see p. 259.)	33	Complete.	Considerably advanced (deep conical ulcer).	Remained well and free from recurrence nearly 12 months; then recurrence in pelvis at some dis-
5	January 7th,	39	Incomplete.	Moderately advanced.	tance from the stump.
*6	March 17th, 1887.	42	Complete.	Early.	Was known to be well 6 years after the operation; not heard from quite recently.
7	April 5th, 1887.	51	Complete.	Moderately advanced.	Well for about 6 months, then recurrence in stump.
8	September 15th, 1887.	52	Complete.	Moderately advanced.	Remained well and free from recurrence 9 months, then recurrence in stump.
*9	December 6th, 1887.	65	Complete.	A large soft brittle growth protruding through os uteri, and attached a little way up cervical canal, bleeding readily on touching it.	Remained free from symptoms of recur- rence for more than 2 years. Recurrence
10	January 16th, 1888. (For full notes, see p. 262.)	39	months pregnant	31	Came up to the hospital soon after the operation, not examined on that occasion; not seen since. I heard that she died in a fainting fit some months later.
11	April 17th, 1888.	39	Apparently complete; but a previous operation had been done with the écraseur not long before by another practitioner. This made it more difficult to distinguish sound from diseased tissue.	Apparently fairly early.	In about 6 months.

-					
No.	DATE OF OPERATION.	AGE.	Complete or Incomplete.	STAGE OF DISEASE.	RECURRENCE.
*12	May 31st, 1888.	43	Complete.	Moderately advanced; soft cauliflower growth over area size of half-acrown on posterior lip of cervix.	years since opera-
13	September 22nd, 1888.	32	plete. The opera-	terior lip.	months in stump.
14	March 28th, 1889.	37	Complete. Douglas's pouch and utero-vesical pouch of perito- neum opened. Cervix cut off with Paquelin's cautery	Advanced.	Well 3 months after operation; not seen since.
15	May 2nd, 1889.	31	Complete.	Moderately advanced; soft growth chiefly affecting posterior lip.	Not seen since leav- ing hospital.
*16	August 26th, 1889.	40	months pregnant	Moderately large cauliflower chiefly affecting posterior lip.	years since the oper-
*17	February 18th, 1890.	53	Complete.	Moderately advanced cauliflower growth size of half a walnut springing from right of cervia just within os uteri.	Still quite free from symptoms; 7½ years since the operation.
18	May 7th, 1891.	38	Complete.	Moderately advanced. Anterior vaginal wall infiltrated slightly. Portion so affected removed. Naked-eye appearance that of cancer. Microscopic examination showed only a glandular structure. Probably clinically malignant, as vaginal wall infiltrated by same kind of growth.	Well for more than a year; then recurrence in stump.

No.	DATE OF OPERATION.	AGE.	Complete or Incomplete.	Stage of Disease.	RECURRENCE.
19	February 18th,	-	Incomplete.	addigate 4	
20	January 29th, 1893.	37	Complete.	Cauliflower moder- ately advanced.	Well at present; more than 4 years since the operation. Sent me by Dr. Garry Simpson, of Acton.
21	February 8th, 1893.	60	Complete.	Nodular form. Early stage.	Well at present. I am doubtful as to malignancy, though symptoms and signs suggestive of cancer. See Case 18 supra as to this point.

Cases 9, 11, 13, 17, 20, and 21 were private cases; the others were operated on at the London Hospital.

Excluding Case 21 as doubtful, it will be seen that, of the twenty cases in this series, there are five cases (Cases 6, 12, 16, 17, and 20), that it has been possible to follow, which have remained well for periods of 10 years, 9 years, 8 years,  $7\frac{1}{2}$  years, and 4 years respectively after the operation. All the twenty-one cases recovered so far as the operation itself is concerned.

The specimens from 6, 12, 16, and 17, and sections of them, were shown at the meeting of the Royal Medical and Chirurgical Society on December 13th, 1892, when my paper on the subject was read. While, in my opinion, the majority of cases of cancer of the cervix that are curable by operation can be adequately dealt with by the supra-vaginal amputation, I quite recognise that there are exceptional cases which require total extirpation; and twenty-one out of my thirty total extirpations of the uterus for cancer have been for cancer of the cervix. Cancer of the vaginal portion of the cervix, when the disease is in a very early stage, certainly requires only the supra-vaginal amputation; cancer

of the supra-vaginal cervix is less readily diagnosed, and in many cases has spread outwards too much for the case to be suitable for either supra-vaginal amputation or total hysterectomy. When it is met with in a fairly early stage, vaginal hysterectomy is the operation that had better be performed.

Treatment in advanced cases. — Here we can only treat symptoms; that is to say, our treatment is

directed :-

1. To checking the hæmorrhage.

- 2. To diminishing the offensiveness of the discharge; and
- 3. To relieving the pain.

To check the hæmorrhage.

General treatment.—Ergot, either in the form of ergotine pills, or of the liquid extract, may be prescribed; the patient should remain in bed whenever the bleeding is considerable.

Local treatment.—Plugging.—If the bleeding is very profuse, it will be right, as a temporary measure, to plug the vagina. This should be done through a Sims's speculum, not passed in far.

The vagina is then filled with tampons of salicylic wool, dipped in glycerine of tannic acid, each tampon having a string attached.

The curette.—When there is a soft growth, such as a cauliflower excrescence, even when it has advanced too far for the case to be suitable for supra-vaginal amputation and complete removal of the diseased tissue, much temporary improvement may at times be secured by thoroughly scraping away all that the curette can be made to remove. Simon's sharp spoon (Fig. 103) answers well for this purpose. After using the curette, we may obtain somewhat better results by:—

(a) Applying the actual cautery to the surface that has been scraped; or,

(b) By applying caustics to it.

Marion Sims recommended chloride of zinc solution (300 gr. to 3j.) for the purpose.

Tampons of wool squeezed fairly dry out of this solution are packed into the cavity; below them other tampons are packed, soaked in bicarbonate of soda to prevent the caustic injuring the vagina. All the plugs are removed in twenty-four hours. Subsequently the vagina should be douched thrice daily with an antiseptic lotion, till the slough caused by the caustic separates; this occurs in about a week's time.

The following case illustrates both the kind of case suitable for the application of caustics, and the details of the method.

J. P., age 47, married twenty-six years, one child, twenty-three years ago, no miscarriages, was sent up to me by Dr. Corner, on February 4th, 1888, as a case of advanced malignant disease of the cervix, with a request that I would do whatever could be done for her in the way of local treatment.

History of the present illness.—Menstruation has been irregular during the last seven years, severe flooding alternating with periods of amenorrhoea. During the early months of 1887 she had no discharge between her periods. Since June 1887 she has been losing some blood every day, and there have been floodings also during this time.

Some time between June 1887 and Christmas 1887 she began to have pain in the left iliac region, which has persisted. She has noticed herself getting thinner since June 1887. There has never been any offensive discharge.



SIMON'S SHARP POON.

Note on admission.—In the situation of the vaginal portion of the cervix, there is a mushroom-shaped, soft growth, convex downwards. It bleeds freely on touching it. The uterus is much less movable than it should be, and the growth has evidently extended to the vaginal walls. The case was, therefore, too far advanced to be suitable for the supra-vaginal amputation.

February 14th, 1888.—Under ether I scraped away all the soft part of the growth with Simon's sharp spoon; the result of this was to leave a crater-like cavity in the situation previously occupied by the growth. I now applied Paquelin's cautery freely to the whole of the diseased surface, and painted it over with pure carbolic acid.

On February 20th, as the patient had experienced no inconvenience from the treatment, I was encouraged to apply caustics. The cavity left after the scraping was packed with plugs soaked in chloride of zinc solution (300 grains to 3j.), and the vagina filled with plugs squeezed out of strong solution of carbonate of soda. The plugs were removed in about thirty hours (having, indeed, been accidentally left in rather longer than was intended).

February 27th.—A yellowish-white slough was removed from the vagina. It was in one piece, and formed a complete cast of the ulcerated cavity in the situation of the cervix, and of the upper part of the vagina. Its measurements were as follows:—Length 3\frac{3}{4} inches, breadth 2\frac{3}{8} inches, thickness \frac{1}{4} to \frac{3}{8} inch. The weight of the slough was I oz. 165 grains. It had no smell whatever. The diseased area from which it had separated had the appearance of a healthy granulating surface.

The patient had exceedingly little discomfort following the application of the caustic, but from February 20th to February 27th she had moderate fever, the temperature reaching 101.4° at night.

After separation of the slough the temperature became normal.

Weight on February 4th, 1888, 7 st.  $5\frac{1}{2}$  lb.; weight on March 5th, 7 st.  $4\frac{1}{2}$  lb. (allowing for the fever, and the hæmorrhage inseparable from the scraping, this is satisfactory). Before the present illness began she weighed 9 st.

June 4th, 1888.—Has been very comfortable since leaving the hospital. Has had no discharge or bleeding to speak of, and feels her strength and general health greatly improved.

The cavity left by the treatment adopted has contracted very much, so as just to admit the finger easily. The edges of the cavity show a papillary red growth, only projecting about  $\frac{1}{16}$  inch from it.

Weight 7 st. 13 lb.

July 5th.—There has been a little more discharge since she was last seen. Weight 7 st. 9 lb.

By the use of the curette, followed either by cautery, or by the application of caustics, the bleeding is diminished, and the discharge either ceases to be offensive, or becomes much less offensive, till enough time has elapsed for fresh portions of the growth to die, and become putrid.

The fever, from which these patients suffer, also disappears till the discharge again becomes offensive.

To diminish the offensiveness of the discharge.— Injections of iodine water (3ij. to Oj. water) two or three times a day, or of weak carbolic (one to sixty), or merely of lead lotion, will be useful for the purpose. An injection of water containing sanitas (3ss. to the pint) is also useful. As already mentioned, scraping with the curette, or sharp spoon, while it checks the hæmorrhage, is also the most effectual means of lessening the offensiveness of the discharge.

To relieve pain.—For this purpose we use either suppositories of morphia (containing \( \frac{1}{4} \) grain of morphia), or hypodermic injections of morphia. It is well to postpone the use of morphia as long as possible, till in fact the pain is really severe, because the doses have soon to be increased, or repeated at more frequent intervals, to obtain relief.

Soreness of the external parts produced by the irritating discharge may be relieved by applying ung. zinci oleati (B.P.), or some similar soothing preparation.

### CHAPTER XII.

CARCINOMA OF THE BODY OF THE UTERUS.

This is a much rarer disease than carcinoma of the cervix. Less than 2 per cent. of cases of carcinoma of the uterus are cases of primary carcinoma of the body (Schroeder).

Origin.—1. It may begin in the deeper parts of the glands, and so form nodules which are at first in the thickness of the uterine wall; as they grow, they form projections either towards the peritoneum, or towards the mucous membrane.

2. It may begin more superficially, and then produces usually *papillary masses* projecting into the cavity of the uterus; this is the form that the disease most commonly assumes (Fig. 104).

Rarely, it causes a general thickening of the mucous membrane.

Etiology.—Age.—Most cases occur in women between the ages of fifty and sixty.

Fertility.—Among patients suffering from cancer of the body there is a much larger proportion of sterile women than in the case of those suffering from cancer of the cervix.\*

Symptoms.—These are the same as in carcinoma of the cervix.

Hæmorrhage;

A discharge, which sooner or later becomes offensive;

Pain.—But whereas in carcinoma of the cervix pain is

\* See On Cancer of the Uterus, by Sir John Williams, p. 105.

a late feature, in cases of primary carcinoma of the body it usually occurs early, and is extremely severe.

Physical signs.—In comparatively early cases we find the vaginal portion of the cervix normal; on bimanual examination, the body of the uterus is found to be enlarged, and the uterus is freely movable. The speculum




Fig. 104.—Carcinoma of the Body of the Uterus (Sir J. Y. Simpson).

will show a sanious discharge from the external os, and portions of brain-like material are often mixed with it. The discharge at this stage is usually offensive. Gentle passage of the sound shows that the uterus is enlarged, and its withdrawal is followed by a little more bleeding than was taking place before using it. Even at this comparatively early stage, the patient may present the appearance of the cancerous cachexia.

In advanced cases.—The uterus is fixed, partly by peritonitis, but also by extension of the disease along the broad ligaments.

Diagnosis.—Certainty in diagnosis can only be attained by dilating the cervix. The finger is then passed into the cavity of the body of the uterus, and recognises some one of the conditions described above—usually soft, friable papillary masses. Sometimes we may find that the endometrium is hard and irregular, the growth being tough, so that nothing can be removed by a curette for examination Here the diagnosis is to be made by taking all the circumstances into consideration, and by recognising that there is no non-malignant condition known to us clinically that would account for the state of the inner surface of the uterus. I have had two cases of this kind, and on the grounds referred to, having concluded that the disease was malignant, performed vaginal hysterectomy. Subsequent examination with the microscope showed the growth in each case to be a typical carcinoma. If doubt exist as to the growth, a portion can sometimes, but certainly not always, be removed for subsequent examination with the microscope; to be of any use such a piece must be large enough to be hardened, and to allow of sections being cut. In practice, taking the history, the age of the patient, and so on, with the presence of such masses as those mentioned in the body of the uterus, there can be little doubt as to their being malignant.

Among conditions which might be mistaken for malignant disease of the body of the uterus are :—

In old people.—Senile endometritis.

In younger people.—

Fibroids.

Retained products of conception.

Fungous endometritis.

Senile endometritis.—This is a condition in which, clinically, a differential diagnosis is frequently required. In such cases there is a purulent discharge, which may be slightly blood-stained from time to time, and the discharge may be offensive.

But in senile endometritis:-

As to the symptoms.—There is no wasting. Pain is either absent, or slight.

As to the physical signs.—The body of the uterus is slightly, if at all, enlarged, and indeed it may be quite small; after dilating the cervix, the finger finds the surface of the mucous membrane quite smooth.

In younger people there is always a strong probability against the presence of malignant disease of the body of the uterus.

Fibroids.—The diagnosis is sometimes a matter of some difficulty. Generally, however, in the case of fibroids, there is a history of symptoms extending over a long period—at all events over a longer period than in cases of carcinoma. The pain also is often much more severe in those suffering from cancer of the uterine body than in cases of uterine fibroids. Where there is any doubt, it is well not to delay dilating the cervix. In one of my own cases successfully treated by extirpation of the whole uterus, I was for some time inclined to regard the case as one of fibroids. Fortunately I dilated the cervix, and felt the friable papillary masses diagnostic of cancer.

As to retained products of conception.—The mere fact that the patient has been pregnant establishes a probability against malignant disease of the body of the uterus, a probability which is stronger the more children she has had. There will not be the rapid wasting, and probably not the severe pain of a carcinoma of the body. Nor will the discharge be found to contain the uebris of brain-like

appearance and consistence common in carcinoma of the body.

Dilating the cervix and removing the mass thoroughly with the finger or curette (with all the precautions essential in every intra-uterine manipulation) will be followed by a rapid recovery. Whereas, if the intra-uterine substance had been malignant, this treatment would produce only an improvement of a very temporary nature.

Fungous endometritis.—A doubt occurred at one time in the case narrated at page 167, as to whether it might not be malignant disease of the body; the age of the patient, the duration of the symptoms, and the absence of offensive discharge and pain, were the points relied on in excluding it.

Subsequently a microscopical examination of the material removed (and the complete recovery of the patient) showed that it was a non-malignant formation.

Treatment.—As in the case of carcinoma of the cervix, the treatment differs according as the disease is in a comparatively early stage, or in an advanced stage.

In early cases.—Given that the case is in an early stage, that is to say, that we are as certain as a careful examination under an anæsthetic can make us:—

- I. That the uterus is freely movable;
- 2. That no thickenings can be felt in the direction of the broad ligament, or utero-sacral ligaments, such as would probably mean extension of the growth in those directions; and
- 3. That the enlargement of the body of the uterus is not so great as to preclude its removal per vaginam,—

The right treatment is to extirpate the uterus through the vagina.

The mortality of this operation, according to Dr. W. Duncan's statistics, published in 1885, was 28.6 per cent.\*

<sup>\*</sup> Trans. Obst. Soc. London, vol. xxvii.

A greatly diminished rate of mortality for this operation has been reported since then by different operators, e.g., 10'5 per cent. by Bokelmann; but many of the cases (all Bokelmann's) were cases of carcinoma of the cervix. In such cases the difficulties of the operation are very much less, because the body of the uterus is of normal size. I have removed the whole uterus per vaginam thirty-three timesin twenty-three of the cases for cancer of the cervix, and in ten cases for primary malignant disease of the body of the uterus. Among this whole series of thirty-three vaginal hysterectomies there have been three deaths. All the cases in which the operation was performed for primary malignant disease of the body recovered. My last fifteen consecutive vaginal hysterectomies for cancer have all recovered. In spite of this it seems to me that extirpation of the uterus is a severe operation, and a severe trial of the vital powers of the patient. It should be a sine quâ non, when the operation is contemplated, that the patient's constitution, so far as can be ascertained, is thoroughly sound.

## EXTIRPATION OF THE WHOLE UTERUS PER VAGINAM.

The first steps of the operation are the same as in supravaginal amputation of the cervix, viz., the drawing down of the uterus, the dissection of the bladder from the anterior aspect of the cervix, the incision through the posterior vaginal wall, and the separation of the lateral attachments of the cervix as high as the internal os. Throughout careful antiseptic precautions are employed. What I do is to squeeze a sponge filled with hot I to 40 carbolic lotion over the parts from time to time during the operation.

Douglas's pouch is now opened with scissors close to the cervix, and the opening enlarged transversely till two fingers can be passed through into the peritoneal cavity. The

vesico-uterine pouch of the peritoneum is then to be opened, the fingers of the left hand in the peritoneal cavity indicating the proper situation, *i.e.*, close to the cervix in front.

Now the uterus is clear in front and behind, and is attached only by the broad ligaments on each side. possible, the uterus is now to be retroverted, and its fundus brought out through the opening in Douglas's pouch. A blunt pedicle needle, threaded with a long piece of No. 5 Chinese twist, is now thrust through a suitable place in the broad ligament, from behind forwards; the loop is caught anteriorly, and the needle withdrawn. One of the loose ends of the ligature is now carried up with a pair of Wells's forceps behind the broad ligament and over its upper border; this ligature is caught, taken out of the forceps, and brought over the anterior aspect of the broad ligament, passed through the loop already there, and tied to the other free end of the ligature. This is done on each side, and then by cutting through the broad ligaments the uterus is freed. As an additional precaution, pressure forceps may be used to clamp the broad ligaments. Some operators, myself among them, rely on pressure forceps entirely, and use no ligatures. The parts are now thoroughly douched with hot I to 40 carbolic; some close the wound in the peritoneal cavity, some leave it open. I prefer to leave it open, and to pack the vagina with iodoform gauze. The gauze is taken out in about four days' time; any pressure forceps left on are removed forty-eight hours after the operation. I prefer myself to leave them on for four days.

If during the operation it happens, as is very likely to be the case in cancer of the body, that the uterus cannot be retroverted and brought through the opening in Douglas's pouch, the operator must tie or clamp the lateral attachments of the uterus, and cut through them piece by piece, as best he can manage. This was the method I had to adopt in my first case, the body of the uterus being so large.

The following is an account of the first case in which I extirpated the uterus for cancer of the body:—

A. R., age 58, a washerwoman, married in 1859; the husband left her three and a half months after marriage; had one child stillborn. No miscarriages. Admitted to the London Hospital, February 25th, 1886, complaining of having been constantly "unwell" for ten months, and for the last two months of having had very severe pain at the bottom of her stomach, reaching down the thighs to the knees. Also she had had a watery discharge slightly coloured with blood, and a little offensive at times.

Family history. - No history of cancer or phthisis.

History of the present illness.—The symptoms first appeared ten months ago, as above mentioned. Shortly before the commencement of her illness she had lost some money in her business, and fretted a good deal about it.

In May 1885 the first thing she noticed was that she became "unwell" very suddenly; the discharge was of a deep-red colour, and came away in clots. She had no pain, and the discharge continued for six weeks or two months; it then left her for a day or two, but came on again as badly as ever. She lost flesh, and latterly has lost her appetite. The pain and the watery reddish-yellow discharge came on about two months before admission, the pain a little before the discharge. For the last month the discharge has been offensive.

The pain soon became very severe indeed; she felt it most in the hypogastric region, and down the inside of her thighs as far as the knees.

First of all it was of a throbbing character, and latterly like something cutting her severely. The pain was always much worse at night, and kept her awake; she was often "doubled up" with the pain; it was never relieved by lying down, but, if anything, worse.

Menstrual history.—The only feature of interest was that the menopause occurred comparatively early, in 1866, when she would be only thirty-eight. Since then she had "seen nothing" till ten months ago.

Present state, February 26th, 1886.—Abdomen a little prominent below umbilicus; a little, but not very, tender; abdominal examination detected nothing else abnormal.

Vulva.—Some erythematous vulvitis, such as is produced by irritating discharges. On asking the patient to strain, a watery yellow discharge escaped from the vagina.

Vaginal examination.—Vagina short; one very sharp "bridle" runs from the left side of the cervix to adjacent vaginal wall. Vaginal portion of cervix normal. An irregular hard lump the size of a cobnut felt posteriorly, apparently in supra-vaginal cervix. Examination causes a great deal of pain. Chloroform was therefore given, and a thorough examination made. The uterus is freely movable. The body can be felt bimanually lying to the right of the middle line, the position being one of anteversion; body enlarged. Through the




Fig. 105.—Primary Carcinoma of the Body of the Uterus. This is a drawing of the uterus extirpated per vaginam in the case of A. R. The uterus has been laid open along the left side. The growth in the cavity of the body has a papillary surface. The cervix is healthy. a. Vaginal portion of the cervix; b. Cut surface of uterine wall; c. Part of the mucous membrane of the body that has not been involved by the growth; d. The malignant growth; pp. Cut edge of peritoneum.

speculum, before passing the sound, some blood-stained discharge was seen escaping from the external os. Small fragments of soft, brainlike material appeared in the discharge escaping after passing the sound.

March 1st, 1886.—Uterus extirpated per vaginam. The operation lasted one hour and forty-eight minutes. The uterus on removal was found to weigh seven ounces (the weight of the uterus in a woman

of fifty-eight would be normally one ounce or less); on opening it an extensive papillary growth was seen projecting into the cavity of the body (Fig. 105). Sections of the growths in the uterus were examined by Professor Victor Horsley, and found to be carcinomatous. Sections of the nodule left in the supra-vaginal cervix were also examined by him. It proved to be an ordinary fibroid.

Subsequent progress.—Highest temperature on the evening of March 2nd, 102'2°. From and including March 5th, temperature was normal, and patient made an uninterrupted recovery.

Previous to the operation the temperature reached about 100° at night, falling to normal in the morning.

The patient was seen from time to time subsequently. She enjoyed good health, and was free from pain. In October 1886 an examination was made, and no return was found. She was not seen again until the end of January 1887, when, though about her work as usual (mangling), she had been getting thinner again, and for about a month had had bad pains across lower part of the back, and down the left leg, and a profuse watery discharge, at times red; but she has not lost much blood; she has been suffering from boils. She would not be examined on that occasion, and it was not for some week or two after that an examination could be obtained. On vaginal examination then, a lump the size of an orange was felt filling the upper part of the vagina; it was fixed, and the surface towards the vagina was ulcerated, and bled readily.

Patient was readmitted to the hospital, and gradually went from bad to worse, dying on July 7th, 1887—sixteen months and seven days after the operation.

For some weeks before her death a hard mass could be felt in the umbilical region.

Post mortem.—There was found some recent adhesive peritonitis in the pelvis; a hard mass occupied the greater part of the pelvic cavity, particularly on the left side. The mass in the umbilical region was found lying on the lumbar vertebræ, and was the size of an orange; it appeared to be due to secondary deposit in the lumbar glands. Hydronephrosis of the left kidney. No secondary deposits in the liver, lungs, or other organs.

Right kidney fairly normal, but its capsule could not be separated without causing some tearing. There had been no symptoms of uræmia.

Remarks.—This patient, then, had about ten months

of renewed health and comfort, such as enabled her to resume her ordinary work (mangling). The pain had been exceedingly severe before the operation, and the complete relief afforded from it by the operation was a very gratifying feature in the case. It will be seen that symptoms had been present for ten months before the operation, so that the case was not by any means an early one. I certainly think that her life was prolonged by the operation; for during the period of non-recurrence, she was freed from the continuous loss of blood, from the pain that kept her from sleeping—and her appetite returned; and I think it fair to conclude that, if she had not had the operation performed, she would have died much sooner.

I have seventeen cases out of the thirty-three in which I have removed the uterus for cancer in which I believe the patients at the present time to be free from recurrence. Eight of these are more or less recent cases; but there are nine cases in which intervals varying from two to nearly eight years have elapsed without recurrence. Thus:—

No. 24, Mrs. H., is well 2 years since the operation. I saw her recently. (Cancer of body.)

No. 23, Mrs. H., is well 2 years and 2 months since the operation. I heard from her recently. (Cancer of cervix.)

No. 22, Mrs. de B., is well 2 years and 4 months since the operation. I see her regularly at the London Hospital. (Cancer of cervix.)

No. 19, Mrs. S., is well nearly 3 years since the operation. I see her regularly at the London Hospital. (Cancer of cervix.)

No. 15, Mrs. J., is well  $3\frac{1}{2}$  years since the operation. (Cancer of cervix.)

No. 8, Mrs. S., is well 4 years and 2 months since the operation. (Cancer of the body.)

No. 7, Mrs. D., is well  $4\frac{1}{2}$  years since the operation. (Cancer of the body.)

No. 4, Mrs. D., operated on in October 1890, has attended for several years at intervals, and shows no sign of recurrence.

No. 3, Mrs. W., operated on in November 1889, was seen in October 1893, and was then well. I have not seen her recently. (Cancer of the body.)

In advanced cases.—Palliative treatment such as was recommended for carcinoma of the cervix is all that can be done.

#### SARCOMA OF THE UTERUS.

This is a rare disease; it is impossible to distinguish between carcinoma and sarcoma of the uterus, except by the microscope; so that clinically these two diseases may be grouped together as malignant disease of the uterus. Similar treatment is required whether the disease be sarcoma or carcinoma. Sarcoma of the body of the uterus is commoner than sarcoma of the cervix. When it occurs in the body of the uterus, it may be found as circumscribed nodules in the uterine walls, or as a diffuse growth, forming more or less irregular projections on the surface of the mucous membrane.

Etiology.—Fertility.—Twenty-five out of sixty-three in Gusserow's series of cases were sterile.

Age. — In the same series the ages were as follows:—

2 under 20.

3 between 20 and 30.

14 ,, 30 ,, 40.

26 ,, 40 ,, 50.

14 ,, 50 ,, 60.

2 ,, 60 ,, 70.

1 over 70.

The symptoms in advanced cases are the same as in carcinoma.

In early cases the discharge is said to be less offensive, and pain less marked.

Diagnosis.—The question will be, in each particular

case, is the disease malignant? (i.e., carcinoma or sarcoma), or non-malignant? And the non-malignant conditions from which sarcoma has to be diagnosed are the same as those mentioned under carcinoma. The following rare case was one of circumscribed sarcoma of the body of the uterus, with secondary nodules in the vagina, and also in the lungs.

E. L., æt. 50, married twenty-nine years, eight children, the last eleven years ago, no miscarriages, was admitted to the London Hospital, June 27th, 1885. She complained of having had a sudden attack of flooding about a month ago; it lasted an hour, and she lost a pint of blood. Two days after she had another similar attack; since then she was confined to bed till she came to the hospital. She once tried to walk across the room about a week after the second attack of bleeding, but the exertion was immediately followed by a third flooding. After each flooding she fainted and vomited, but the attacks were not attended by any pain. She has been losing flesh the last three months, and for the last three or four months has had a watery discharge from the vagina of a "dirty" colour. A month ago (that is, about the same time the flooding came on) she noticed a lump protruding from the vulva.

Previous history.—Patient began to menstruate at sixteen; she was quite regular till her marriage; the flow lasted a week, and she had no pain at her periods. During the last nine years, patient says she had altogether six attacks of profuse flooding; the one which began a month ago is the sixth. The first attack of bleeding occurred nine years ago, and confined her to bed for five weeks. The second attack was three years after this. Between the attacks the patient says she was quite regular.

Present condition, June 29th, 1885.—Vulva.—On inspection, without separating the parts, a greenish-black mass is seen between the labia majora posteriorly. By separating the labia the mass is seen to be oval, with its long diameter from before back, measuring  $1\frac{5}{8}$  inch; transversely I inch.

The exposed surface of the mass is indistinctly divided by radiating furrows into three lobes. On tracing it up, the mass is found to be attached to the posterior vaginal wall by a broad pedicle, which extends downwards as far as the posterior edge of the vaginal orifice.

The part of the tumour that is within the vagina is covered with vaginal mucous membrane of a pale pink colour, like that elsewhere in the vagina. But this covering ceases abruptly where the mass is exposed to the air, and, as already mentioned, the surface of the exposed portion is of a greenish-black colour.

About half an inch up the vagina on the right side, a second mass the size of a walnut is to be felt beneath the mucous membrane. At the




FIG. 106.—Circumscribed Sarcoma of the Body of the Uterus (Author's case). The uterus is laid open from behind. In the recent state the sarcomatous masses were dark red.

most prominent part of this mass is an area the size of a threepenny piece, of a dark purple colour. Here the vaginal mucous membrane has become attached to the subjacent growth, and has ulcerated. Elsewhere the mucous membrane covering the tumour is normal. The mass has an elastic, semi-fluctuating feeling.

Uterus.—The condition of parts in the vagina interferes with a satisfactory examination of the uterus, but it feels heavier than normal, and freely movable.

The temperature since admission has varied from normal to 100.4°. Patient is thin and sallow; her general appearance suggests malignant disease; a gland can be felt in each groin, but neither of them is enlarged.

On July 2nd, 1885, the growths were removed, and after their removal (which was easily effected) a careful examination was made, the patient being still under the influence of an anæsthetic.

The uterus was freely movable; its surface appeared smooth, and free from any irregularity. The sound passed  $3\frac{1}{2}$  inches without causing any bleeding.

After the operation the patient had an attack of septicæmia, from the effects of which she sank on July 7th.

Post mortem.—Three distinctly circumscribed growths, of a deep red colour and soft consistence, were found in the walls of the uterus. The largest occupied the thickness of the uterine wall at the fundus. It measured vertically 1\frac{3}{4} inch, and horizontally 2\frac{1}{2} inches. The peritoneal surface of the uterus opposite this growth was discoloured, and of a purple tint, but was quite smooth. The lowest growth was partly in the wall of the cervix, partly in that of the body of uterus (see Fig. 106). There was no peritonitis. There were numerous secondary growths about the size of peas, but individually varying considerably in size, in both lungs. None elsewhere. The kidneys were granular.

Microscopical examination of the growths showed that they were mixed round and spindle-celled sarcomata.

### DECIDUOMA MALIGNUM.

This is the name given to a primary malignant disease of the body of the uterus, occurring in connection with labour or abortion. It runs a rapid course, and secondary deposits occur in other parts, particularly in the lungs and vagina. The exact nature of the disease is at present a matter of some uncertainty. Some observers regard it as having an essentially causal relation with pregnancy. Others consider the relation to pregnancy as accidental, and believe the growth to be essentially a sarcoma. Microscopically, to quote from the excellent paper published by Dr. Herbert Spencer on the subject, "it is characterised by large cells

of various shapes, of which some occur singly or in groups, and others in fused masses, with large deeply staining nuclei resembling enormous giant cells, the so-called 'syncytium.' . . . It resembles closely the syncytium or layer of fused cells which forms the outer coating of the normal chorionic villus, and it seems probable that the syncytium of deciduoma malignum develops from the syncytium of the chorionic villus, though this is denied by some authorities." \* syncytium is vacuolated. The growth in deciduoma malignum is also characterised by large extravasations of blood into it. Dr. Eden, who has devoted much attention to the subject, believes that the cases of deciduoma malignum "ought really to be regarded as rapidly growing sarcomata." He had examined, with Dr. Kanthack, sections of secondary growths from a case of sarcoma of the testis. Both these observers considered that in these sections were to be seen

I have met with two cases bearing on the question. One is that described on page 285, which occurred in 1885; it obviously presents some important clinical features in common with cases described as deciduoma malignum. Thus it was a primary malignant growth of the body of the uterus, not carcinomatous, and secondary nodules were present in the vagina and lungs. There had, however, been no pregnancy for eleven years in that case.

"plasmodial masses which show almost all the points laid

down by writers on deciduoma malignum as characteristic

The second case came under my notice in December 1896. It is undoubtedly identical with the cases that have been described as deciduoma malignum. Dr. Herbert Spencer, Dr. Eden, Mr. Targett, and Mr. Bland Sutton, all of whom have examined the uterus, and sections of it, agree that it is so. That being admitted, an important question arises

<sup>\*</sup> Quarterly Medical Journal, July 1896.

as to the evidence of pregnancy having occurred recently in the case. It appears to me, although it is impossible to prove that the patient had not been pregnant, that there is a considerable probability that she never had been pregnant. I read a paper on this case before the Obstetrical Society of London in July 1897, and it will be found in vol. xxxix. for 1897 of the Society's *Transactions*. My own view is that this case is one of primary sarcoma of the body of the uterus, arising independently of pregnancy; and my impression is that pregnancy is not an essential antecedent for the occurrence of the disease known as deciduoma malignum.

The notes of this case are as follows:-

E. W., a married woman, twenty-four years of age, was admitted under my care into the London Hospital on December 19th, 1896, complaining of pain in the back, bearing-down pain, and a constant red discharge since a supposed miscarriage nine weeks previously.

Previous history.—She has had a cough since she was fifteen years of age, and has attended Victoria Park Hospital. She says her left lung is affected. She had small-pox eight years ago, and since then has been a little deaf. She has been married twelve months, and thinks that she has had three miscarriages. The first miscarriage was in April 1896, when she believed herself to be two months pregnant; the second was in August 1896, when she thought herself to be five or six weeks pregnant; and a third was nine weeks before she was admitted to the hospital.

Menstrual history.—The catamenia began when she was ten years old. She then "saw nothing" for ten months, and was very irregular till the time of her marriage, there being intervals of two, three, and four months between the periods. The periods were scanty, and attended with pain in the hypogastric region and back. She sometimes had to keep her bed during her periods. She has had no trouble with her water. The appetite is generally bad, and the bowels generally constipated.

Family history.—Her father is alive. All the family have suffered from "weak chests."

History of the present illness .- The patient says that she has had a

red vaginal discharge ever since the first miscarriage, which occurred in April 1896. The discharge was more profuse when the patient was up and about. She has also had more or less pain since the first miscarriage. Nine weeks before her admission, when the last miscarriage occurred, she supposed herself to be three weeks pregnant. She was attended by a doctor, and "a small piece of flesh came away."

At the time of her admission the case appeared to be an ordinary one of metrorrhagia, due to incomplete abortion. The note made at this time was as follows:—

Nothing abnormal can be felt on abdominal examination. On vaginal examination, there is recent blood about the external parts, and in the vagina. The os is not patulous. Fresh blood is seen in the os. The uterus is freely movable, and evidently somewhat enlarged.

December 21st, 1896.—Two special laminaria tents were inserted on the evening of December 20th, and at 3.30 p.m. on the 21st the dilatation of the cervix was completed with Hegar's dilators. The finger was then passed into the uterus, and came upon a projecting mass about the size of a walnut, part of which was easily detached. The mass was situated on the posterior wall near the fundus, and to the right side. Most of the mass was easily detached with the finger, but there was no line of demarcation between the mass and the underlying wall of the uterus. As successive portions of the mass were scraped away, the finger kept going more and more deeply into the substance of the uterine wall, so that the condition was altogether unlike that found in ordinary cases of incomplete abortion. The material removed was reddish yellow, and somewhat friable. I had on one previous occasion come upon a condition having some resemblance to that found in this case. This was in a case of hydatidiform mole. The bulk of it came away spontaneously, and it was some weeks afterwards, when the interior of the uterus was explored on account of a persisting metrorrhagia, that the condition referred to was met with. Portions of the hydatidiform mole were projecting at the fundus, and were easily removed, but the finger did not come down on to firm healthy uterine wall. As more and more of the separable material was detached, the finger kept getting more and more deeply into the wall of the uterus, till it was very nearly, if not quite, in contact with the peritoneal coat. In the case of hydatidiform mole referred to, although one felt doubtful as to whether the whole of the mole had been removed, the patient ultimately made a completely satisfactory recovery.

In the present case (E. W.) pure tincture of iodine was applied to

the whole of the endometrium. The uterus was washed out, and a strip of iodoform gauze was left in the vagina. This was removed on the following day. The temperature subsequent to the dilatation did not rise above normal, but the red discharge persisted more or less during the patient's stay in the hospital on this first occasion, although she was constantly taking ergot, and having hot vaginal douches. She went out on January 4th, 1897. At that time her weight was 6 st. 7 lb.




Fig. 107.—Uterus from Case of E. W. Primary Sarcoma of the Body of the Uterus (Deciduoma Malignum). The uterus has first been laid open along the anterior wall, and then the posterior wall has been cut vertically almost up to the peritoneum. A. Scattered nodules of sarcomatous growth. B. Fibroid in posterior wall. C. Anterior cut edge of posterior wall. D. Cut anterior wall of uterus.

Readmission.—The patient was readmitted on January 13th, 1897, as the metrorrhagia had persisted, and there had been two floodings during the few days she was at home. On one occasion it was said that half a chamberful of blood with clots came away.

Examination, January 21st, 1897.—The patient was not markedly anæmic. On examining the abdomen, nothing abnormal could be felt. On vaginal examination, there was a little blood about the vaginal

orifice, and the uterus was considerably enlarged, being about the size of the pregnant uterus at seven or eight weeks. It was freely movable and soft. She was treated with hot vaginal douches, and took 3 grs. of ergotine three times a day up till the 29th. During this time there was no improvement. She suffered from a good deal of pain at the



FIG. 108.—Microscopic Section from the Author's Case of Deciduoma Malignum (E. W.), showing the plasmodial masses (syncytium) and the large multinucleated cells.

bottom of the stomach, and backache, and pain down the left leg as far as the knee; and the red discharge persisted. On January 30th the cervix was again dilated, and the following note was made as to the condition of the endometrium:—

Tracing the anterior wall of the uterus from left to right, with the finger in the uterus, it was found to be smooth. The posterior wall was smooth on the left side, but about the middle it became rough, and

on the right side came to an edge, beyond which there was an irregular chasm, on the floor of which several brittle irregular masses could be felt, and portions were easily broken away with the finger. Bimanually, when the finger was on the floor of this chasm, there seemed to be little or nothing beyond peritoneum between the hands. At this time I said



Fig. 109.—Microscopic Section from the Author's Case of Deciduoma Malignum (E. W.), showing the plasmodial masses (syncytium) and the large multinucleated cells.

to those who were present, that but for the age of the patient I should have no hesitation in saying that the case was one of carcinoma of the body of the uterus, and that I felt practically certain that the disease was malignant, and that it had considerable resemblance to cases that have been described under the title "deciduoma malignum."

Operation, February 11th, 1897.—The patient was anæsthetised with ether, and placed in the lithotomy position. Vaginal hysterectomy

was then performed, the principal steps of the operation being as follows:-The anterior fornix was first incised and the bladder separated as high as the internal os. Then the posterior fornix was opened, and the cervix cleared to the same height. The anterior and posterior transverse incisions were then joined by lateral incisions, bleeding vessels being secured with pressure forceps. Douglas's pouch was opened. Then the vesico-uterine pouch of the peritoneum was opened, and the lateral attachments of the uterus were secured with pressure forceps and divided. The left side of the uterus was freed first and delivered into the vagina, and afterwards the right broad ligament was secured and divided. The uterus was then removed. Iodoform gauze was packed into the vagina, and the upper part of the gauze projected into Douglas's pouch. The patient made an uninterrupted recovery. The pressure forceps were removed on the fourth day, the temperature on that occasion reaching 101°, the highest point it reached throughout the whole period subsequent to the operation. After the tenth day the temperature never rose above 99.2°. She was examined on March 15th. On abdominal examination, nothing abnormal was to be felt. On vaginal examination, a granulating surface the size of a sixpence was to be seen at the top of the vagina, looking quite healthy. The patient went home on March 16th, 1897.

It is important to notice that these cases are rare, and that they are at first liable to be regarded as cases of subinvolution, with metrorrhagia.

Floodings, with more or less persistent metrorrhagia, are the chief symptoms. When a woman has a flooding, and passes large clots, she naturally is apt to think she has had a miscarriage.

Deciduoma malignum (or primary sarcoma of the body of the uterus) may occur in quite young women (this patient of mine, for instance, was only twenty-four), and in that respect it stands in sharp contrast to primary carcinoma of the body of the uterus, which is essentially a disease of advanced life.

Early diagnosis, followed by vaginal hysterectomy, is the chief thing to be desired in the management of these cases.

Now that attention has been directed to it, although the disease can hardly be common—for it is unlikely that the characteristic secondary deposits have been generally overlooked—still, it may be found not to be so extremely rare as it at present appears to be.

Those who wish to go fully into the subject are referred to papers by Dr. Herbert Spencer and Mr. Rutherford Morison, and to the discussion on them to be found in the London Obstetrical Society's Transactions, vol. xxxviii., and to a paper by the author in vol. xxxix. of the same Transactions.


### CHAPTER XIII.

DISEASES OF THE FALLOPIAN TUBES.

Congenital peculiarities.—The chief one is the existence of two or more openings surrounded by fimbriæ, two or more accessory fimbriated extremities, instead of one. In the hundred cases I examined, this condition was met with six times; in most of these, but not in all, the accessory opening communicated with the main tube.

Inflammation of the Fallopian tubes.—Salpingitis.

—This is an extremely important affection, as proved both by post-mortem room investigation and by clinical observation. For example, in cases of pelvic peritonitis following labour, it is certain that in most, perhaps in all cases, the sequence of events is-endometritis, salpingitis, and extension of the inflammation from the tube through the open fimbriated extremity, producing pelvic peritonitis. If the patient recovers, we cannot prove the existence of the salpingitis; but if she dies, we may actually see it, as I remember doing in a case of puerperal fever, fatal a short time after delivery. There was evidence, post-mortem, of corporeal endometritis; the fimbriated ends of the tubes were open, pus could be pressed out of them, and patches of lymph were seen distributed over the pelvic peritoneum generally. In this case it is to be noted that the fimbriated ends of the tubes were open. This must have been because the inflammation of the pelvic peritoneum immediately external to the fimbriated extremity of the tube was of too



outer end of the ovary, Lo. Ovarian ligament;

the broad ligament.

neum,

surface, and extends on to the upper fifth of the vagina. ip. The infundibulo-pelvic ligament, i.e., that part of the anterior surface of the uterus as low as the level of the internal os; it covers almost the whole of the posterior upper free border of the broad ligament not occupied by the Fallopian tube. It runs outwards to the brim of the

pelvis.

intense a character to allow of the formation of adhesive lymph in the neighbourhood of the opening of the tube, which would have sealed it up.

In chronic cases the fimbriated opening has generally




FIG. III.—Uterus and its appendages, showing the state of the latter when matted together by the adhesions resulting from pelvic peritonitis. The ovaries are not visible, being concealed by the adhesions. During life, the physical signs would be, diminished mobility of the uterus, and the presence of "thickening" at the sides of the uterus. As the fimbriated extremities of both tubes are sealed up, a patient whose organs are in the condition represented is incurably sterile (Heitzmann).

become obliterated by adhesive pelvic peritonitis gluing it to surrounding parts. Had this occurred in the case mentioned, the inflammation might have remained localised, and the patient have recovered. Etiology and morbid anatomy.—From what has been said, it will be evident that the etiology of salpingitis is to a great extent, if not entirely, that of corporeal endometritis.

Therefore we may expect to meet with salpingitis:-

- 1. Following labour or abortion.
- 2. In cases of gonorrhœal or simple vaginitis, by extension.
- 3. In cases where the endometrium has been directly injured by tents, or other instruments.
- 4. As one of the morbid conditions produced by cold during menstruation.
  - 5. In certain fevers.
- 6. A tubercular salpingitis is also met with. In such cases there is usually tubercular mischief in other organs.


In fact, whenever there is corporeal endometritis, salpingitis is likely to occur by extension of the inflammation along the tubes.

When the inflammation has reached the fimbriated extremity of the tube, pelvic peritonitis is set up in its neighbourhood. Provided the pelvic peritonitis be of the adhesive variety, what happens is, that the fimbriated end of the tube is closed up, being as it were glued to the ovary, and all trace of the fimbriæ is usually lost.

Subsequently the tube may become dilated by accumulation of the inflammatory secretion. Whether in a particular case dilatation occurs, or not, seems to depend on whether there is free escape for the secretion towards the uterus, or not.

In many cases of dilated tubes, we do find that there is a communication open towards the uterus—that the canal is not closed in this direction, but in such cases we find the channel of communication very fine and tortuous, so that there is practical occlusion. When the channel towards the uterus is narrow, we can easily understand that

there is little tendency for the secretion to escape into the uterus; for as the Fallopian tube becomes dilated, it tends



to fall behind the uterus into Douglas's pouch; the escape of the fluid is then hindered, partly by gravity, and partly

by the kinking of the channel. The outer two-thirds of the tube is the part most liable to become dilated; the sausage-like swelling formed is largest at its outer part, and the whole swelling is usually more or less distinctly subdivided by annular constrictions into two or three compartments, which, however, communicate freely with one another.

Dilated tubes are classified as either hydrosalpinx, pyosalpinx, or hæmatosalpinx, according to the nature of their contents.

In hydrosalpinx the fluid is usually clear and yellow; occasionally it is milky. So long as the contents are not distinctly purulent, and are not composed of blood, it is best to classify a specimen as one of hydrosalpinx.

In pyosalpinx the dilated tube contains pus.

In hæmatosalpinx the tube contains blood.

Frequency.—I examined the condition of the pelvic organs with special reference to this question in a series of a hundred bodies taken as consecutively as possible in the post-mortem room of the London Hospital; dilatation of the Fallopian tubes was found in seventeen of these cases.

In 5 cases there was pyosalpinx.

In 8 ,, hydrosalpinx alone.

In 4 ,, hæmatosalpinx.

There was evidence of pelvic peritonitis in all the cases.

Whatever be the cause of salpingitis, it is a cause which tends to affect both tubes. In thirteen out of the seventeen cases both tubes were dilated; and of the remaining four (where there was dilatation only on one side), in two only was the Fallopian tube on the undilated side normal; taking as the essential characters of a normal Fallopian tube the presence of an open fimbriated extremity, and of a channel in communication with the cavity of the uterus.




Fig. 113.—Double Hydrosal, inx, with broad adhesions between the dilated tubes and adjacent parts (Winckel). The uterus and its appendages are seen from behind. On the left side there is a small ovarian cyst as well as a dilated Fallopian tube. Tuba sin. Left tube; Ovarium sin. Left ovary; Ovariencyste. Ovarian cyst; Adhaesionen. Adhesion; hydrops tubae. Dropsy of the tube; Ovarium d. Right ovary.

Relation of hydrosalpinx to pyosalpinx.—These conditions are probably merely stages of the same disease; the fact that in some cases we find that on one side there is a hydrosalpinx and on the other a pyosalpinx favours this view.

Relation of hæmatosalpinx to hydro- and pyosalpinx.— Hæmatosalpinx may be produced by hæmorrhage into a tube, the fimbriated extremity of which has been previously obliterated, or by an accidental hæmorrhage into a preexisting hydro- or pyosalpinx.


In many cases, however, hæmatosalpinx is the result of extra-uterine pregnancy occurring in the Fallopian tube. In other words, in some cases hæmatosalpinx is due to the presence of a blighted ovum, successive hæmorrhages into the tube taking place from time to time, similar to the hæmorrhages that accompany pathological conditions of the ovum when the pregnancy is intra-uterine. In some specimens of hæmatosalpinx, due to extra-uterine pregnancy, the fimbriated extremity of the tube remains patent—a fact which is in keeping with the pathology of the condition just mentioned. Mr. Doran showed a specimen of hæmatosalpinx at the Obstetrical Society of London, which was probably due to a blighted ovum in the tube. In this specimen the fimbriated end of the tube was open. I have myself met with similar cases.

It is well to bear in mind, then, that what appears to be merely a hæmatosalpinx may really be a blighted ovum—in fact, a tubal mole—in the Fallopian tube. This may be proved by finding chorionic villi in sections of the mass examined under the microscope.

Importance to be attached to these conditions.—This is a point which cannot even yet be regarded as fully determined.

As to the facts, in the writer's series, in one case certainly, and probably in another, the dilated tubes were the cause of death.

Both these were cases of pyosalpinx, and the patients died of acute peritonitis. In one of them the place where the wall of the dilated tube had given way, allowing its



contents to escape into the peritoneal cavity, was clearly seen.

In one case, that of a woman, age sixty-two, the tubes contained cheesy masses; this probably should be classified as a case of pyosalpinx that had undergone natural cure. The condition of a patient who is the subject of a pyosalpinx is certainly a serious one. How far hydro- or hæmatosalpinx constitutes a danger to life, we are not yet in a position to say. The diagnosis of pyosalpinx from hydro- or hæmatosalpinx is often practically impossible. The presence of more or less severe fever from time to time would be in favour of pyosalpinx; but there may be sometimes high fever where there is only hydrosalpinx. Again, patients with even double pyosalpinx, the pus being horribly fœtid, may have long periods of quiescence—periods in which they are free from fever, and perhaps from pain.

**Diagnosis.**—Symptoms.—These are not very definite; but a case presenting the following group of symptoms may, with a good deal of probability, be set down as one of dilated Fallopian tubes.


The patient complains of pain across the lower part of the abdomen, and it may, or may not, be worse on one side. She has dysmenorrhœa, and the regularity of the catamenia is disturbed. Menorrhagia, and sometimes metrorrhagia, are commonly met with in these cases. If the dysmenorrhœa dates from a particular confinement or abortion, or if a history of gonorrhœa can be obtained, the probability of the tubes being diseased is strengthened. Again, the history that the patient has had several previous attacks of pelvic peritonitis is strongly suggestive of disease of the Fallopian tubes. Such cases have been termed cases of "recurrent peritonitis." Patients themselves often speak of pelvic peritonitis as "inflammation of the bowels." A history of attacks of "recurrent peritonitis," or of several attacks of "inflammation of the bowels," particularly in a woman who is absolutely sterile, or has been so during the time covered by the history of these attacks, points strongly to chronic salpingitis.

Physical signs.—In some cases where the swelling formed by the dilated tube is very small, particularly if the part of the tube affected is that just external to the uterus, it may be impossible to recognise any swelling. Such a case is, however, exceptional; dilatation usually affects the outer part of the tube, leaving a portion of undilated tube between the uterus and the tumour. Again, in most cases both tubes are dilated, and in all cases there is pelvic peritonitis, varying in extent. It has been mentioned that the tumour formed by a dilated tube tends to fall behind the uterus; still, however, keeping rather to its own side of the middle line.

Given the history such as that sketched out above, if we find the uterus less movable than normal (on account of the accompanying pelvic peritonitis), and if we find a sausage-like tumour lying in Douglas's pouch, a little to one side of the middle line, that feels as if it contained fluid, and is partially fixed, still more if this condition exists on both sides, the probability is that the case is one of dilated tubes.

Diagnosis by puncture is not generally to be recommended, and may be dangerous; nor would it be conclusive, e.g., the fluid might come from a small ovarian cyst. Considering the frequency of dilated tubes among the general population, as evidenced by the frequency they were met with in a consecutive series of bodies, coming from all parts of a general hospital (in my series 17 per cent.), it is, I think, a fair inference that, if it were possible to examine the organs in a series of patients attending the gynæcological department in any of our hospitals, we should find the tubes diseased in a still larger proportion. For instance, where there is lessened mobility of the uterus, with the presence of either a distinct lump, more or less fixed, in one or both posterior

quarters of the pelvis, or even where there is "thickening" such as is usually taken as evidence of a previous pelvic inflammation—peri- or parametritis—such cases would, I



think, if it were possible to examine the organs, yield a high percentage of dilated tubes (see Fig. 111). In cases where these physical signs are present, and where the abdomen has been opened, it is usually, if not invariably, found that the tubes are diseased.

**Treatment.**—If it were possible to be certain of the presence of a pyosalpinx, as distinguished from a hydroor hæmatosalpinx, there would be no hesitation in recommending an operation for its removal by a skilled specialist. Pyosalpinx certainly constitutes a danger to life; but hydrosalpinx or hæmatosalpinx are not always dangerous, though they may sometimes become so.




FIG. 116.—Showing the appendages grasped by large elbowed pressure forceps. The pedicle-needle bearing a loop of silk, perforates the broad ligament below the forceps and near the uterus (Doran).

The ligature is then to be adjusted ready for tying, and the appendages are seized with two pairs of small Wells's forceps just above the large pressure forceps; this is removed and the ligature tied; the appendages are then cut away. The small forceps are merely to retain a hold of the pedicle, to allow of its being inspected just before the closing of the abdominal wound.

When there is very severe dysmenorrhoea, as well as more or less pain between the periods, associated with the physical signs of dilated tubes, removal of the diseased tubes (and ovaries), by one specially skilled in abdominal surgery, is a justifiable operation, provided palliative treatment (counter-irritation, hot douching, etc.) has been fairly tried, and provided also that sufficient time—for example, eighteen months or two years—has been allowed to elapse to give the patient the opportunity of recovering spontaneously. Of course, the exact nature of the operation must be explained to the patient, and, if she is married, to her husband. In such cases the dilated tubes and ovaries may be found so densely matted together and to the surrounding parts, that the operation for removing them would be an exceedingly dangerous one (e.g., involving risk of tearing some of the big veins, or the ureters), or may even be altogether impossible.

The proportion of cases in which the removal of the appendages is found to be impossible diminishes greatly as the experience of the operator increases. There are very few cases indeed in which an experienced operator finds it impossible to complete the operation.

In my series of seventeen cases, removal of the diseased organs would have been absolutely impossible in one case, and in another could only have been done by removing at the same time the body of the uterus; in the remaining fifteen cases removal would have been possible in all, and easy in most.

The best thing to do, when an operation has been decided on, is to commence the operation merely as an exploratory abdominal section, with the full understanding that, if the organs are found so densely adherent that their removal would be a very hazardous matter, the operator is not to attempt removal, but to close the wound. It sometimes happens that after such an operation, *i.e.*, merely an exploratory abdominal section, the patient's pains are relieved to a greater or less extent. Such a case occurred

to me several years ago at the London Hospital. The patient had swellings on each side of the pelvis such as may be caused by dilated tubes; on opening the abdomen, the organs on each side were so firmly adherent in the pelvis that I decided it would be unjustifiable to attempt to remove them, and I closed the wound. The patient made a good recovery from the operation, and, singularly enough, although she was informed that the tumour we had hoped to remove had not been removed, she has on several occasions since assured me that the pain she has had since the operation is not worth speaking of as compared to what she had before it. Possibly, in examining the relation of the organs during the operation, some adhesions may have been broken down that had something to do with her pain.

The following are notes of the first two cases in which I removed the appendages for chronic inflammatory mischief:—

CASE I. unfortunately contracted syphilis in the summer of 1890, and I have not seen her since; up to that time she had been free from symptoms since the operation.

Case II. expressed herself as being much better for about a year after the operation; but within the last few months her various pelvic pains have returned, and she is now nearly, if not quite, as bad as when I first saw her.

The notes of these cases are as follows:-

CASE I.—N. G., age 28, was admitted into the London Hospital under my care on May 12th, 1888.

She had been married ten years, and had one child, eight and a half years ago. She thought she had had a miscarriage four years before admission.

She had never been well since her confinement, which was said to have been a difficult one. She had been attended at the time by a midwife, and was very ill for some time after it.

Ever since the confinement she had suffered from constant pain and aching in the left iliac region, with numbness down the left leg.

The pain had been much worse for the five months preceding her admission.

Since her confinement, also, menstruation had been very irregular, occurring at intervals of five weeks, seven weeks, and once ten weeks. Sometimes it was a mere "show"; at most, it lasted two days. She had very great pain for a week beforehand, somewhat relieved when the flow began. The pain was like labour pains, and was felt throughout the pelvis, but worse in the left iliac region.

Before the confinement she had always been regular every four weeks, losing a good quantity, and having no pain at the periods.

On May 17th, 1888, she was examined under an anæsthetic. The uterus was fairly movable, and a swelling, also fairly movable, was felt to the left side of it. This was the size of a large egg.

She was advised to try palliative treatment, and did so for six months without finding herself any better. She was accordingly readmitted, and I removed the uterine appendages on January 21st, 1889. Some dense adhesions on the left side had to be separated before the appendages on that side could be removed. Both the Fallopian tubes were slightly enlarged, and the fimbriated ends had been closed by adhesions, the outer ends of the tubes being firmly adherent to the corresponding ovary. I have seen this patient at intervals since the operation; she was quite free from the constant pain which had been her chief trouble for about fifteen months after the operation. She then contracted syphilis, and I have lost sight of her. Curiously, she had had a periodical discharge of blood from the uterus, at shorter intervals since the operation than before it.

CASE II.—C. S., age 30, was sent up to me by Dr. Thompson of Woolwich, and was admitted into the London Hospital under my care in May 1888.

She had been married twelve years, and had had three children, the last nine years before admission.

The confinements were said to have been easy; but her symptoms date from ten years ago. Her chief complaint is of constant pain in the left iliac region, extending down the left leg during that time.

She had been in three other hospitals before coming to the London, and had worn many pessaries.

Before the present illness the catamenia occurred every three weeks, and were unattended with pain; since the illness there has been a good deal of irregularity as regards menstruation; on the whole, the irregularity has been rather in the direction of menorrhagia.

She was examined under an anæsthetic. The uterus was found to be retroflexed, and so bound down by adhesions that it could not be replaced with the sound. A swelling about the size of a walnut was felt to the left of the uterus; this was also more or less fixed. Here, as in the last case, the diagnosis was chronic inflammation of the uterine appendages, and the patient was given palliative treatment. This was continued for more than a year; but as she was then no better, she was readmitted into the hospital, and on August 19th, 1889, I removed the uterine appendages.

There was some dilatation of the left tube, and its outer end was closed and matted to the ovary in the usual way. There were also dense adhesions between the left appendages and the pelvic wall.

I have seen this patient at intervals, and she was completely free from the pain for about a year after the operation. She also has had some periodical discharge of blood from the uterus since the operation; there has been, however, no pain with it.

## ABNORMAL PATENCY OF THE INNER END OF THE FALLOPIAN TUBE.

As a rule, the inner part of the Fallopian tube is so narrow that an extremely fine bristle is needed, and some expenditure of time and patience, to pass it from the uterus along the tube. This is the rule in dead bodies. Very rarely a specimen may be met with when the inner end of the tube is much more open than this, and may allow a common surgical probe to pass along it, even post mortem, as in a case narrated by the late Dr. Matthews Duncan, where, during life, there was reason to think the sound had passed along the Fallopian tube. During life the inner part of the tube has not probably an unvarying calibre, but may dilate to some extent, perhaps sufficiently to allow an ordinary uterine sound to pass along the tube. It is in this way that cases are explained where the size of the uterus, as ascertained by bimanual examination, is not increased, and yet the sound can be passed several inches in the direction of one of the tubes. In rare cases this may be

the true explanation; but in most cases where it happens it is far more probable that the sound has perforated the walls of the uterus, and passed along under the peritoneum between the layers of the broad ligament, and is not in the tube at all. Out of the body I have several times tried pressing the point of an ordinary uterine sound against the uterine end of the tube in the direction of the tube. What happens is, that when the pressure has reached a certain point, the sound perforates the mucous and muscular coats of the uterus, and slips into the space between the layers of the peritoneum forming the broad ligament, outside the Fallopian tube altogether. The peritoneum is loose in this situation; and unless the sound is pressed with great force, or passed for an extreme distance, the peritoneum is not perforated, which may account for no serious mischief resulting when this has been done during life.

Notes of four recent cases of dilated Fallopian tubes that have been under my care, and illustrate the clinical aspect of such cases, follow here:—

Hydrosalpinx on Left Side. Severe Pain in Left Iliac Region and Metrorrhagia of Five Months' Duration. Removal of Dilated Tube, followed by Complete Relief.

E. F., age 30, was admitted into the London Hospital under my care on December 18th, 1895. She was sent to me by Dr. Rhodes of Plumstead. She had been married ten years, and had had no children, but two miscarriages at about the second and third month of pregnancy, the last one being nearly two years ago. She complained that for the last five months she had had severe pain, more especially in the left iliac region, but also in the lower part of the abdomen generally, and in the lower part of the back. For the last month she had experienced a sensation of movements in the abdomen. For the last five months there had also been a slightly blood-stained vaginal discharge, practically continuous for the whole of that time. Before

five months ago she had been quite regular, but during the last five months she had not been able to tell, on account of the continuous red discharge, when she was menstruating and when she was not. During the same time she has had pain and nausea after food. She often vomited, and this gave relief. She has a perforation of the soft palate: this occurred ten years ago, at a time when she suffered from severe ulceration in the throat. She gives no history of any rash or sore, and there are no other signs of syphilis.

On examination (December 28th, 1895) the uterus was found to be of the ordinary size, in the normal position, and fairly movable. An elastic, somewhat sausage-shaped swelling was felt in the left posterior quarter of the pelvis; it seemed about the size of a duck's egg, and was fairly movable. The breasts showed no sign of activity. The diagnosis come to was, that the swelling was probably a dilated Fallopian tube, or perhaps a small ovarian cyst. The possibility of extra-uterine foetation was thought of, but excluded on account of there being no sympathetic changes in the breasts, and no enlargement of the uterus.

On January 10th, 1896, the patient was anæsthetised with the A.C.E. mixture, and the abdomen opened in the usual way. The swelling above described was found to be the left Fallopian tube dilated, and containing clear watery fluid. There was nothing to be seen of the fimbriated extremity of the tube, the outer end being firmly adherent to the corresponding ovary. Only one adhesion had to be tied. The left broad ligament was then transfixed and tied in two halves, one separate ligature being also tied round the whole pedicle. The dilated tube, with a portion of the left ovary, was then removed, and the wound was closed completely. Nothing was done to the appendages on the right side, as they seemed healthy. The patient made a satisfactory recovery. She came up to the Hospital on February 20th, and said that she had been completely free from pain since the operation. There had been no return of the metrorrhagia.

Remarks.—An interesting point in the case was the definite and simple nature of the symptoms—pain in the left iliac region, and a persistent metrorrhagia for five months before the operation. It so often happens, in cases of disease of the appendages, that symptoms are extremely vague, and are not rarely complicated by an obviously neurotic temperament in the patient. This was not so in the present case; the patient was a very sensible woman, having two distinct sources of complaint—the pain and the metrorrhagia; and the removal of the apparent cause has been followed by complete relief from the symptoms. I think

315

persistent pain is a more constant symptom in cases of dilated tubes than is the metrorrhagia, though the latter is by no means uncommon. In my experience metrorrhagia is also present in a considerable proportion of cases in which the cause of the dilatation of the Fallopian tube is the presence of an early extra-uterine pregnancy.

## Left Pyosalpinx. Right Hæmatosalpinx and Small Ovarian Cyst.

O. A., a married woman, 39 years of age, was admitted into the London Hospital under my care on October 26th, 1896.

Previous history.—She married at the age of 20, and had a premature confinement about a year afterwards. The child was born alive, but died in half an hour. This is the only pregnancy.

When she was 19, she had a sore on the inner aspect of the left labium. It was burnt with "caustic"; about two months afterwards she had an ulcerated sore throat. She is positive that she never had a rash.

Two years after the confinement she had an attack of gonorrhoea, attended with frequent micturition and scalding.

She had scarlet fever at the age of 23, and has had several attacks of "rheumatism."

She has had three separate attacks of "inflammation of the bowels," at the ages of 29, 34, and 39 respectively.

Menstruation began at the age of 14. She was quite regular, and has never lost much at her periods.

Present illness.—This began in June 1896, with an attack believed to be "rheumatic fever." She then went to St. Thomas's Hospital. When she had been in the hospital three weeks, the abdomen became very tender, especially on the left side. The question of operation was then apparently discussed, but she was not thought at the time to be strong enough to undergo it. The pain referred to the left iliac region has persisted up to the present time. She also complains of a constant bearing down pain and of some swelling of the right leg.

She was quite regular up to Christmas 1895; the periods then began to diminish in quantity, and finally ceased in May. There have been two scanty periods since then—one a month ago, and one two days before her admission to the London Hospital.

State on admission.—She is a well-nourished, strong-looking woman. The pupils are unequal and irregular, the right pupil almost the size

of a pinhole. They react to light and accommodation. Pulse 102, and temperature 99°.

No cardiac murmur was heard; the urine was normal.

October 31st.—As regards the state of the pelvic organs, she was examined under an anæsthetic, and the following note was made:—

On examination of the abdomen, a swelling is felt in the hypogastric region. On vaginal examination, a swelling is felt somewhat to the right of the middle line, which is found to be the body of the uterus moderately enlarged. The sound passes  $3\frac{1}{2}$  inches. To the left of the uterus there is a swelling the size of an orange, feeling soft and elastic. Both the uterus and this swelling have a fair degree of mobility. The outline of the uterus is fairly regular.

Operation, November 19th, 1896.—The patient was anæsthetised with the A.C.E. mixture, and the abdomen was opened by the usual median incision. The tumour to the left of the uterus was found to be a pyosalpinx adherent to the corresponding ovary, and also adherent to the uterus, rectum, and the pelvic wall. The adhesions were separated with the finger, and the dilated tube, with the ovary, was removed, the pedicle being tied in three sections. While pulling up the pyosalpinx the tube burst and pus escaped. It was mopped up as completely as possible with Gamgee pads. On the right side the Fallopian tube was also dilated, its contents being bloody fluid; and the right ovary was also enlarged, so as to form a small cyst containing similar blood-stained fluid. The appendages on the right side were also removed. The patient was put in Trendlenberg's position for a short time, so that the deeper part of the pelvis, from which the adhesions had been separated, could be looked at. She was then put down flat again, and the peritoneum was washed out with saline fluid. An iodoform gauze drain about 30 inches long (it was measured after removal) was put in, and the rest of the wound closed with silk sutures. The gauze drain was removed on the fourth day, and her subsequent progress was uneventful. She went out on January 2nd, 1897.

The following case illustrates very well the impossibility of exact diagnosis as between a dilated tube and a small ovarian cyst. Both tubes were dilated and contained pus, and there was an ovarian cyst also on each side. On the left side the ovarian cyst itself contained pus:—

Mrs. G. was seen in consultation with Dr. Piercy Fox of Clapham,

on January 9th, 1897. She was 30 years of age, and had been married twelve years. She had had four children, the last four and a half years ago. She was inclined to think there had been one miscarriage eighteen months ago, as she had then gone ten weeks without "seeing anything," and then bleeding came on, but no lump was passed. After that she was fairly regular, the periods lasting a week, and not much being lost. She had no special pains at the times except back-ache.

History of present illness .- She had not been well for several months before I saw her. In the summer of 1896 she fell when getting out of a 'bus, and from that time she had experienced pain in the left groin. Soon after this fall she was unable to hold her water for three days, but this trouble soon passed away. About three months before I saw her she caught a chill; she was menstruating at the time. That period only lasted three days instead of a week, and she has practically been in bed ever since, with constant pain in the left iliac region, running down the thigh to the knee. About the time that she caught the chill her temperature was taken, and was found to be 103.5°. She had violent pains in the abdomen, which was very tender, and she vomited after everything she took. The pain was constant for about seven weeks from the beginning of the illness. She had been regular up to the time when she caught the chill. She missed two periods after the chill. About three weeks before Christmas, 1896, she was seen by another consulting physician, after which the pain was also felt in the right iliac region, but for about two weeks before Christmas all her pain had very much diminished, and she got up the day after Christmas. Her pains, however, came on more acutely than before, causing her to go back to bed, and pain in the lower abdomen has been constantly present, especially in the left iliac region, up till the time when I saw her (January 9th, 1897). As already mentioned, she missed two periods after the chill; but soon after the consultation three weeks before Christmas the period came on again, and lasted for a fortnight.

At the time I saw her the temperature was 101°. The lower abdomen was tender, and an indistinct swelling could be felt in the hypogastric and left iliac regions. On vaginal examination the uterus was fixed, and fixed swellings could be felt in the right and left posterior quarters of the pelvis. The swelling in the left was distinctly elastic, and I felt sure that it contained fluid. Taking into consideration the long period during which she had been ill (over three months), the fact that she had been feverish on and off during the whole of that time, and the persistence of the pain for the same period, I came to the conclusion

that the case was either one of suppurating ovarian cyst or pyosalpinx, possibly bilateral, with a good deal of pelvic peritonitis round the tumour or tumours. I advised that an exploratory abdominal section should be performed, and the swellings removed, if the opinion as to their nature which I had formed was correct.

The patient and her mother had been reading the case of Beatty v. Cullingworth, and they were fully alive to the questions that might arise as regards the removal of the ovaries. The patient expressed herself at the time when I first saw her as decidedly averse to the removal of the ovaries. I explained to her that almost certainly the operation contemplated would involve the removal of the left ovary, and it was extremely probable that the right was also extensively diseased, and would probably also have to be taken away. As she had shown herself to be averse to the removal at all events of both ovaries on this occasion, I refused to have anything to do with the case unless I had the express permission of her husband and self in writing to remove either one or both ovaries, according as might seem necessary at the time of the operation. I have never previously had the express consent in writing from any patient for any operation, but under the circumstances it seemed to me desirable to have a written authority on this occasion. The permission obtained ran as follows :-

"January 11th, 1897.

"I hereby give Dr. Lewers permission—in the operation that is requisite for my recovery—to have full discretion to remove one or both of my ovaries, should he, during the operation, consider such to be quite necessary.

"ALICE G.

Operation, January 14th, 1897.—The patient was anæsthetised with the A.C.E. mixture. Dr. Piercy Fox gave the anæsthetic, and I was assisted at the operation by Mr. L. A. Smith, who was at the time resident accoucheur at the London Hospital. On opening the abdomen in the usual way, the swelling to the right of the uterus at once came into notice. The swelling was made up of an ovarian cyst the size of a Tangerine orange, containing a thin brownish-red fluid, and of the right Fallopian tube, which was enlarged so as to be as thick as the thumb. Its fimbriated extremity could not be distinguished either at the operation or subsequently. The right Fallopian tube contained pus. Both the right ovarian cyst and the right Fallopian tube were very extensively adherent in the pelvis. The attachments were ultimately

separated, during which both the cyst and tube burst, and the pedicle was transfixed and tied with silk as usual. The swelling to the lef side, which had been the more conspicuous one before the operation consisted of the left Fallopian tube thickened to the size of the thumb, intimately adherent to the left ovary, which had become cystic and reached the size of an ordinary orange. The Fallopian tube was intimately adherent to the ovarian cyst, and they both contained pus, which escaped during the separation of the adhesions in the pelvis, which, as on the other side, were extremely extensive and firm. A fairly satisfactory pedicle was ultimately established on the left side and tied. The left appendages were then removed. The peritoneum was washed out with hot saline solution, and a long strip of iodoform gauze, a yard or more, in one continuous piece, was packed into the pelvis, one inch of the gauze being left projecting at the lower angle of the wound. The rest of the wound was closed in the usual way, and the patient put back to bed. During the first two days after the operation she appeared to be in a very critical state, the pulse being exceedingly weak, although not excessively rapid; 112 to 120 indicated the range of frequency. Several hypodermic injections of strychnine were given, which improved the pulse considerably. She was fed by nutrient enemata. On the 16th there was a decided improvement in the pulse, and the improvement was maintained on the 17th, when the iodoform gauze was removed. I felt pretty certain on that day that, apart from accidents, the case was going to do well. A small piece of gauze was passed down for some two or three inches, just to keep the cavity from which the gauze plug had been removed properly drained.

There is very little more to say about the case. The highest temperature from the 14th to the 26th was 100°, after which it was normal, and she ultimately made a very satisfactory recovery.

The following case was one of uterine fibroid, complicated by a small suppurating ovarian cyst on the left side. The uterine appendages were removed on both sides, but the fibroid was left:—

A. L., a single woman, employed as a florist, age 37, was admitted into the London Hospital under my care on January 11th, 1897, at the request of Dr. Deane, of Leytonstone.

Previous history.—She has always enjoyed good health until the present trouble began about a month ago.

Menstruation.—She began to menstruate at fourteen, and has been regular every four weeks till the beginning of the present illness. She had very little pain at the periods, and the quantity lost at each period has been moderate up till two years ago, since which time up till November 1896 it has been relatively excessive. Since then there has been a more or less constant sanious discharge.

Present illness.—The present trouble began about November 1896, when she noticed a swelling of both legs, principally the left, and aching of the legs. Both these conditions were relieved by lying down. During the same time she has complained of pain in the lower part of the abdomen—chiefly on the left side—of a dull aching character, somewhat relieved by lying down. This pain has been in the left iliac region, never above the umbilicus. She has also complained since the beginning of the illness of night sweats. She was apparently quite well till about Christmas, 1895, and, except for a feeling of general malaise, she noticed nothing particularly wrong until about November 1896, when the legs began to swell, as already mentioned. The catamenia were quite regular up till November 1896, every four weeks, but since then she has been losing blood on and off.

State on admission, January 11th, 1897.—The patient is sallow-looking, somewhat anæmic and thin, with no indication of anything wrong with the heart and lungs. The urine was normal.

Abdominal examination.—A swelling is felt in the hypogastric and left iliac regions, reaching up about half-way between the umbilicus and pubes, and reaching outwards on the left side within two inches of the anterior superior iliac spine. The surface of the swelling is somewhat irregular. The swelling extends very slightly also to the right of the middle line.

Vaginal examination.—The nymphæ are large and pendulous. On bimanual examination the swelling in the middle line is apparently the body of the uterus considerably enlarged, and a good deal less movable than it should be. There is also a swelling in the left posterior quarter of the pelvis, which suggests the presence of inflammatory mischief affecting the left uterine appendages. This swelling is more or less fixed. The right lateral fornix appears normal. The sound passes with some difficulty 3½ inches.

The diagnosis made was uterine fibroid, with inflammatory mischief

of the left uterine appendages, probably purulent.

The patient was in the hospital for twenty-four days before the operation. During that time she had a feverish attack lasting about

five days, the temperature reaching 101°. She was carefully examined again on February 1st, 1897, when the physical signs were just the same as those already noted.

Operation, February 4th, 1897.—The patient was anæsthetised with the A.C.E. mixture. The abdomen was opened in the usual way. It was then found that there was a large fibroid tumour of the uterus, and also a tumour outside the uterus in the region of the left uterine appendages. On further examination this swelling was found to be composed of an ovarian cyst the size of a large orange, and the left Fallopian tube, which was enlarged so as to be as thick as the thumb. No sign of the fimbriated end of the tube could be seen, the tube simply ending on the cyst. The ovarian cyst was intimately adherent for an inch or more to the sigmoid flexure, and there were also adhesions between the cyst and the pelvic wall. During the separation of these deep adhesions the cyst burst, and its contents, which were purulent, escaped, some of the pus going into the peritoneum. The ovarian cyst and the dilated tube on the left side were ultimately freed and drawn up out of the wound, the pedicle transfixed and tied with two interlocking ligatures, and a single silk suture tied round the whole pedicle. The cyst and dilated tube were then removed. The right uterine appendages were healthy, but they were removed in a similar manner, with a view of influencing the fibroid tumour of the uterus. (The patient had given her consent to allow either the removal of the uterus with the fibroid tumour, or the removal of the uterine appendages, whichever should seem in the course of the operation to be the more desirable.) The peritoneal cavity was then washed out with warm salt solution, and an iodoform gauze drain was packed into Douglas's pouch, one end of it being left out at the lower end of the incision. The rest of the wound was then closed in the usual way. The gauze was removed on the fourth day after the operation, and the patient made an uneventful recovery, leaving the hospital on March 6th, 1897. Her temperature after the operation never exceeded 100.6°. It remained in the neighbourhood of 100° to about the end of the fourth day, after which it was practically normal.

I have seen her several times since she left the hospital. There has been no return of the menstrual periods, and she has been free from pain.

## CHAPTER XIV.

# Pelvic Inflammation (Pelvic Peritonitis, Pelvic Cellulitis).

The expression "pelvic inflammation" includes pelvic peritonitis and pelvic cellulitis; for though in a particular case we are usually able to say it is a case of pelvic peritonitis, or a case of pelvic cellulitis, according to circumstances, they are usually present together, the one, however, predominating over the other, and giving its name to the case.

Pelvic peritonitis is also known as *perimetritis*, and pelvic cellulitis as *parametritis*.

## PELVIC PERITONITIS—PERIMETRITIS.

Etiology.—Cases of pelvic peritonitis can always be placed in one of the following groups:—

- 1. Cases starting from labour and abortion.
- 2. Cases arising in connection with menstruation.
- 3. Cases following gonorrhaa.
- 4. Cases due to traumatic causes, the injury permitting the entry of septic matter, e.g., rough use of the sound, etc.
- 5. There remain some cases of pelvic peritonitis which do not belong to any of the above groups; such are:—

Cases of pelvic peritonitis arising in connection with:

- a. Ovarian tumours.
- b. Fibroid tumours.
- c. Malignant disease and tubercle.

It will be seen that the etiology of pelvic peritonitis, as regards the first four groups of cases, is identical with the etiology of corporeal endometritis; and indeed, in these four groups, with the doubtful exception of the cases arising in connection with menstruation, there is no doubt that the pelvic peritonitis arises by extension of the inflammation from the body of the uterus along the tubes to the peritoneum, the sequence being corporeal endometritis—salpingitis—pelvic peritonitis.

As regards the group of cases arising in connection with menstruation.—In these, pelvic peritonitis may arise either:—

(a) By extension from the endometrium, exposure to cold having set up corporeal endometritis; or

(b) Simultaneously with corporeal endometritis.

It should be remembered that at the menstrual periods the whole of the pelvic viscera are intensely congested physiologically, and it is easy to see how exposure to cold may in such a case cause a general pelvic peritonitis, and at the same time an inflammation of the whole mucous tract of the genital organs, *i.e.*, salpingitis, corporeal endometritis, cervical endometritis, vaginitis.

(c) There is another way in which to account for pelvic peritonitis occurring during menstruation. There may be an escape of a small quantity of blood into the pelvic peritoneum, either regurgitating from the uterus along the Fallopian tubes, or coming from the ovary, when the Graafian follicles rupture, and this will set up pelvic peritonitis.

Of course, when the quantity of blood in such cases is considerable, we have what is known as pelvic hæmatocele, the symptoms and signs of which are well defined. But when the quantity of blood escaping into the peritoneum is small, the characteristic symptoms of hæmatocele, on which

we chiefly depend for diagnosing it, will be absent, but yet the quantity of blood may be quite sufficient to set up pelvic peritonitis. This is probably the way to explain cases arising during menstruation, when there has been no exposure to cold, or other discoverable cause.

Morbid anatomy.—Pelvic peritonitis may be either:—

1. Simple.—Here the peritoneum is injected, and the surface has lost is lustre, but there is no perceptible exudation of lymph upon it.

2. Adhesive.—Here there is a layer of lymph exuded on the surface of the inflamed peritoneum. Such a layer is commonly as thick as a piece of thickish blotting paper.

3. Serous.—This is similar to the last variety, except that the lymph-covered surfaces are separated by serous fluid to a variable extent.

4. Purulent.—In this variety the fluid separating the

inflamed peritoneal surfaces is pus.

In serous perimetritis and purulent perimetritis, the fluid is encysted, its boundaries being formed partly perhaps by some natural boundary, e.g., Douglas's pouch, and elsewhere by adhesive peritonitis between the neighbouring coils of intestine and other adjacent organs, shutting off the fluid from the general peritoneal cavity.

It should be noticed that a pelvic abscess may be due to a perimetritis as well as to a parametritis (pelvic cellulitis). Probably such abscesses due to perimetritis are really either actually examples of pyosalpinx, or are collections of pus (localised by adhesions between adjoining viscera) that have been caused by a small escape of pus from a minute rupture of a pyosalpinx.

Symptoms.—In acute cases there is pain across the lower part of the abdomen, vomiting, and the ordinary symptoms of fever, anorexia, thirst, headache, and so on; there may be rigors; frequent desire to pass water, and

pain in passing it, are common symptoms, and there may be pain on defæcation.

In chronic cases the prominent symptom is pain across the lower part of the abdomen, either right across the abdomen, or more or less localised in one or other iliac region, and backache. Dysmenorrhœa is common in such cases, and there may be sterility.

There is usually disturbance of the regular course of menstruation, either in the direction of amenorrhœa, or of menorrhagia.

Dyspareunia is often present in the chronic form, such as produces adhesions. In cases arising in connection with ovarian tumours, there is sometimes no pain.

Signs.—In acute cases.

General.—There is rapidity of the pulse, and a high temperature. The patient lies on her back with her legs drawn up.

Local.—The abdomen is more or less distended, and is very tender on palpation; owing to the rigidity of the muscles, we may not be able to make much out by abdominal examination. In other cases, as soon as sufficient time has elapsed to allow of adhesions taking place (say forty-eight hours), we may feel a lump, perhaps reaching up to the umbilicus or higher; such a lump is fixed, hard, tender, and may be more or less resonant. It is formed by matting together of the intestines by adhesive peritonitis.

Vaginal examination.—During the first few hours of the attack we only find great tenderness on examination, and that the vagina is hot; perhaps an indistinct fulness in the posterior fornix may also be detected at this stage.

Later, say after forty-eight hours, when the exudation has had time to coagulate, and the adjacent parts have become fixed by adhesive peritonitis, the physical signs in typical cases are either:—

- I. The uterus is fixed, and occupies the centre of the pelvis; all round it we feel hardness, so that, to quote the usual simile, it feels as if plaster of Paris had been poured into the pelvis, and had set.
- 2. In another equally characteristic condition, we find, as before, the uterus fixed, but pushed forward, so as to




Fig. 117.—Successive accumulations of lymph in Douglas's pouch in pelvic peritonitis, shown in section (Thorburn).

lie close to the pubes; as before, there is hardness all round, but behind, the hardness takes the form of a definite lump, which has pushed the uterus forwards.

This lump is formed by accumulation of the exudation while still fluid in Douglas's pouch, owing to the action of gravity (Fig. 117).

In chronic cases.—One of the most constant physical signs is lessened mobility of the uterus; considerable

experience is necessary to say when the uterus is less movable than it ought to be.

It is observed by endeavouring to tilt the uterus as a whole upwards, and noticing whether it moves as much, and as easily, as it does when the parts are healthy.

More or less thickening may be felt at some parts of the vaginal roof, either behind, or in the right or left posterior quarter of the pelvis. It has already been mentioned, in the account of diseases of the Fallopian tubes, that, judging both from post-mortem room experience, and the experience of abdominal operators, such "thickenings" are in many cases composed of a more or less dilated Fallopian tube matted by adhesions to the ovary, uterus, and parts around. This probability is increased if the thickening is felt in one or other posterior quarter of the pelvis, and is still greater if it occurs in both posterior quarters.

## COURSE.

It is said that the inflammatory products may be entirely absorbed, and no doubt this is true of the greater portion of such products in each case; but judging from the frequency with which adhesions are met with in the postmortem room and during operations, between the various parts of the pelvic viscera, it is difficult to believe that the inflammatory products, once the adhesion stage has been reached, are ever so entirely absorbed as to leave no trace whatever in the form of adhesions.

When the fimbriated extremities of the Fallopian tubes are closed by adhesive peritonitis, we have necessarily an absolute sterility; and we have the condition established for dilatation of the tubes to occur, according as the secretion in the tube can escape easily through the uterine end of the tube or not.

In the chapter on dysmenorrhœa it has been suggested how the persistent adhesions after an attack of pelvic peritonitis may cause painful menstruation.

If the inflammation goes on to suppuration, we have a pelvic abscess, the contents of which may occasionally be absorbed; much more frequently, the abscess bursts either externally, or into the vagina, bladder, or bowel.

## PROGNOSIS.

Pelvic peritonitis usually ends in convalescence; but from what has been said, it is clear that the patient will be unusually fortunate if she escape some of the common sequelæ of pelvic peritonitis—chronic pelvic pain, dysmenor-rhæa, dyspareunia, and sterility. Further, relapses are common; particularly from imprudence during menstruation, over-exertion, exposure to cold, and so on.

Certain cases following labour illustrate the tendency of the inflammation to be lighted up again by slight causes. It is not uncommon to see a case where, after labour, nothing of sufficient importance occurred to attract attention till the patient got up, probably about the tenth day, and that then well-marked symptoms and signs of pelvic peritonitis developed. The explanation of such cases is probably that a slight pelvic peritonitis really existed from the first, but so long as the patient remained at rest in bed the symptoms were ill-marked, pain perhaps being absent, or being taken for after-pains. When she got up, the tendency to congestion accompanying the upright position, and the exertion, caused the pelvic peritonitis to become more severe, and to be accompanied by definite symptoms. When the inflammation becomes general-when general peritonitis results from pelvic peritonitis-the prognosis is very grave.

#### DIAGNOSIS.

In acute cases.—The chief conditions from which pelvic peritonitis has to be distinguished are:—

Pelvic cellulitis.

Pelvic hæmatocele.

Extra-uterine fœtation.

In all these cases there is pelvic peritonitis as a minor complication, but there are usually sufficient grounds for enabling us to decide which is the most important condition present.

Pelvic cellulitis.—As regards *etiology*, this more often follows wounds of the cervix, by laceration during labour, or in gynæcological operations.

Symptoms.—Pain, nausea, and vomiting are less marked. The patient lies with only one leg drawn up.

Physical signs.—In pelvic cellulitis there is usually a lump at one side of the uterus pushing it to the opposite side. The finger feels, on the unaffected side, the normal arching of the lateral fornix, while on the affected side it comes on a mass bulging downwards, and so producing a convexity in the lateral fornix, instead of the normal concavity. Thus the vaginal portion of the cervix appears shortened on the affected side. The uterus and tumour formed by the exudation are less absolutely fixed than in typical cases of pelvic peritonitis.

Pelvic hæmatocele.—The history here is the chief point—sudden onset—sudden pain in the pelvis, with a sense of nausea, faintness, and perhaps vomiting. Such conditions are particularly significant if they happen when, a menstrual period having been a week or so overdue, metrorrhagia occurs.

Physical signs.—In intra-peritoneal pelvic hæmatocele, the one we are now considering, the lump in Douglas's pouch

is, as a rule, much larger than that formed in simple pelvic peritonitis, and the uterus is pushed farther forwards.

Extra-uterine fœtation.—Many cases of extra-uterine fœtation have in their earlier progress been mistaken for pelvic peritonitis. The history is such cases usually helps us—the patient has probably gone two or three weeks over her proper time without being poorly, and then comes on bleeding again—a decidual cast of the uterus may be passed. A tumour, formed by the sac of an extra-uterine fœtation, is before rupture usually more to the right, or left, of the middle line, than the lumps formed by pelvic peritonitis.

### TREATMENT.

Preventive treatment.—A consideration of the etiology will suggest many precautions for guarding against pelvic peritonitis.

As regards labour and abortion, the utmost care should be taken to avoid introducing septic matter on the examining finger; the nails should be kept short, the greatest care should be taken to cleanse them thoroughly with a nailbrush before making a vaginal examination; a knife should never be used for this purpose, as one constantly sees done. The point of the knife merely produces a naked-eye appearance of cleanliness, and is utterly useless for producing the absolute cleanliness which is alone efficient. The use of a knife in this way deepens the space under the nail, and increases its capacity of collecting dirty material, while it can only remove the more obvious portion of such material. It may seem unnecessary to dwell upon this apparently small matter, but it is in reality one of great importance. A reliable method for rendering the hands aseptic has been already described (p. 43). Again, over-frequent examinations during labour should be avoided. When possible, an

antiseptic vaginal douche should be given before labour and after delivery.

Acute vaginitis, gonorrhæal or other, should be cured as soon as possible. In using injections for vaginitis care should be taken to see that there is a free return for the fluid used. I think the hydrostatic douche a much safer apparatus than Higginson's syringe. I have seen three cases where a rigor, and a sudden high temperature, with pain across the lower part of the abdomen, followed soon after the use of a Higginson's syringe in vaginitis; it seems at least possible that some small quantity of the injection mixed with some of the secretion in the vagina may have passed into the peritoneum along the Fallopian tubes. There is obviously much greater danger of this accident happening if an intra-uterine injection is being given; a tube with a double channel attached should always be used in such cases, and the hydrostatic douche apparatus, not Higginson's syringe. Some air always gets in when douches are given with the latter. This can be shown by using a glass vaginal pipe fitted on to a Higginson's syringe; during the "diastole" of the bulb, air will be seen in the upper inch of the glass tube.

As regards all intra-uterine operations, passing the sound, dilating the cervix, scraping the endometrium, etc., all possible antiseptic precautions should be taken.

Treatment in acute cases.—The patient must be kept at perfect rest in bed, and the treatment must be chiefly symptomatic. If there is *vomiting*, the patient should have small pieces of ice to suck; bismuth and hydrocyanic acid may be prescribed. Vomiting is often best relieved by giving small quantities of weak brandy and soda-water iced, or small quantities of iced champagne.

Pain.—If the patient can bear to lose a little blood, half a dozen leeches applied to the hypogastrium will often

relieve it; otherwise, small quantities of morphia hypodermically ( $\frac{1}{5}$  grain), and hot fomentations to the abdomen, should be ordered.

Diet.—Beef-tea, milk and soda-water, toast-water, should be given if they can be retained; if not, nutrient enemata should be tried till the vomiting subsides. Brandy is often necessary, and the state of the pulse will be a guide as to the quantity; it may be given either by the mouth, or in the nutrient enema.

The temperature.—Quinine in powders stirred up with a drachm of milk should be given, e.g., gr. iij., three times a day. Sometimes an ice-bag to the head is useful in reducing the temperature.

Surgical treatment.—There are some cases where probably the best thing to do is to open the abdomen, and wash out the peritoneal cavity. It is impossible to say precisely in what cases this would be right. There are some, however, where there should be no hesitation in adopting it, e.g., if a patient is known to be the subject of an ovarian tumour, or if there is considerable probability that she has dilatation of the Fallopian tubes; and if in either case sudden symptoms of acute peritonitis arise, there should be no hesitation in opening the abdomen.

It is rather in cases following labour or abortion that difficulty arises. Probably some cases of puerperal peritonitis that die might have been saved by opening the abdomen; on the other hand, many apparently very severe cases recover. At present we have not sufficient data to enable us to lay down any rule. Each case must be judged on its own merits. It is well, however, to remember that the risk of an exploratory incision, performed by some one experienced in abdominal surgery, is not in itself great; and in the cases of puerperal peritonitis that would be benefited, probably nothing more than opening the peritoneal

cavity, washing it out, and draining it would be necessary. Treatment of this kind has proved very beneficial in some cases of tubercular peritonitis where laparotomy has been performed, the fluid let out, and the wound closed without drainage. In America several cases have been recorded where abdominal hysterectomy has been performed during the puerperium for septic mischief, but the results are not very encouraging.

Treatment in chronic cases .- Here the pain and dysmenorrhœa are the chief symptoms requiring treatment. For the pain, blistering over the seat of the pain produces improvement, usually, however, only of a temporary character. Painting with iodine paint (Tr. iodi, Lin. iodi, āā partes æquales) is useful in a similar way. Hot vaginal douches, the douche being as hot as the patient can bear it, usually about 110° F., are also useful. The use of the glycerine plug every night may be recommended.\* In a large number of cases, however, improvement is only temporary, and the symptoms return. In some of these cases, where the symptoms and physical signs point probably to disease of the Fallopian tubes, it becomes a question whether removal of the uterine appendages (ovaries and Fallopian tubes, which in such cases are usually more or less adherent to one another) may not be the right treatment. This has been discussed in the chapter on "Diseases of the Fallopian Tubes."

#### PELVIC CELLULITIS—PARAMETRITIS.

Etiology.—Injury, especially to the cervix, but also to the vagina and perineum, is the fundamental antecedent to

<sup>\*</sup> A glycerine plug is simply a piece of absorbent cotton-wool the size of a walnut, with a string four or five inches long tied round it. The plug is dipped in glycerine, squeezed rather dry, and passed into the upper part of the vagina.

pelvic cellulitis. Septic matter gains an entrance at the point of injury, and inflammation is set up in the connective tissue; the connective tissue of the pelvis is primarily affected, though by extension the inflammation may spread to connective tissue far away from the pelvis, e.g., behind the kidney.

The injuries referred to are usually produced during labour or abortion; laceration of the cervix being the most important. It is not the laceration in itself that causes the cellulitis, but the laceration, inasmuch as it admits septic matter, which sets up the inflammation.

Wounds of the cervix during operations are a frequent cause of pelvic cellulitis, unless the operations are done with every antiseptic precaution. For instance, in performing the supra-vaginal amputation of the cervix, where the connective tissue at the sides of the cervix is opened up in a most extensive manner, provided the operation is done antiseptically, no pelvic cellulitis follows. In thirty cases of my own where this operation was done no cellulitis followed. In a case of fibroid polypus too big to remove through the os, where I incised the cervix posteriorly with the cautery, a phlegmon of the right broad ligament followed; I think, because the slough caused by the cautery could not be kept aseptic. Parametritis is very similar to whitlow. Clean punctures and cuts on the fingers heal without any cellulitis; but if dirt (i.e., septic matter) gets into them, whitlow will probably be the result.

Morbid anatomy.—There are three varieties of pelvic cellulitis; that is, three conditions in which we may see it in the post-mortem room.

These are:-

- 1. Phlegmon.
- 2. Abscess.
- 3. Gangrene.

1. **Phlegmon.**—When the exudation has not passed into the stage of suppuration, it is called *phlegmon*.

It is exceedingly rare to have an opportunity of seeing parametritis in the stage of phlegmon in the post-mortem room; cases of parametric phlegmon are common enough, but either the patients recover, or if they die, the cellulitis has usually by that time passed from the stage of phlegmon to that of abscess. The following is an account of a case, and a description of a specimen of a parametric phlegmon, which I had an opportunity of examining post-mortem.

## POST-MORTEM APPEARANCES OF A PHLEGMON OF THE BROAD LIGAMENT.

S. H., aged 39, was admitted to the London Hospital a few days after her confinement, suffering from mania and parametritis.

The physical signs of parametritis were well marked, there being a well-defined lump in the situation of the right broad ligament, displacing the uterus to the opposite side. The patient had also a good deal of bronchitis, and I think her death, which occurred on the tenth day after delivery, was chiefly due to the bronchitis.

On post-mortem examination, the layers of peritoneum forming the right broad ligament were found to be separated by exudation between them, so that from before back the broad ligament from peritoneal surface to peritoneal surface measured one inch and a half. The separation of the layers of peritoneum forming the right broad ligament began at the lower border of the Fallopian tube, and extended downwards as far as the broad ligament extends. Externally the separation by exudation extended to the pelvic wall. The Fallopian tube was, as it were, stretched over the convex upper surface of the swelling formed by the exudation between the layers of the broad ligament.

On cutting into the swelling, the cut surface had an appearance like that of a somewhat coarse sponge, there being seen holes of various sizes separated from one another by solid tissue.

The cavities referred to were filled with a sero-sanguinolent fluid, but none of them contained pus.

The largest holes would admit a No. 16 catheter.

The right ovary measured two and a quarter inches long, one inch

and a quarter high, and three-eighths of an inch thick. Its surface was adherent to the adjacent peritoneal surface of the broad ligament by recent lymph. On section the ovary was found to contain an abscess, holding about half a drachm of pus.

On the left side the broad ligament was normal, when held up to the light, and looked at from before back, having its normal translucency.

The left ovary was two and a quarter inches long, five-eighths of an inch high, and three-sixteenths of an inch thick. It did not contain the corpus luteum.

The *uterus* was about six and three-eighths inches long, and measured about four and a half inches between the points of entry of the Fallopian tubes. The placental site was on the posterior wall, and presented a mammillated appearance. Elsewhere the surface of the uterine cavity was nearly smooth.

The exudation takes place into the connective tissue at the side of the cervix; from this situation it passes between the layers of the broad ligament to the side of the pelvis. Normally, the layers of the broad ligament, if held up to the light, are found to be translucent; when phlegmon of the broad ligament occurs, the layers of the peritoneum forming the broad ligament are separated for a considerable distance by the exudation, so that the broad ligament may measure from before back an inch or two inches in thickness.

The phlegmon forms a convex lump, over the upper surface of which are spread in order, from before back, the round ligament, the Fallopian tube, and the ovary. The appearance on section has already been described in the account of the case just narrated. It is to be noticed that some pelvic peritonitis affects the peritoneum overlying the inflamed pelvic connective tissue, as mentioned in that case. Sometimes the connective tissue in the utero-sacral ligaments is implicated (utero-sacral cellulitis); and sometimes there may be a cellulitis of the connective tissue between the bladder and cervix.

- 2. Abscess.—If the inflammation go a stage further than phlegmon, an abscess is formed.
- 3. Lastly, the inflammation may be of so intense a character as to produce sloughing of the tissues affected—gangrenous parametritis. This is very rare.

Directions in which pelvic cellulitis spreads.—
Here we must distinguish between the course taken by a phlegmonous cellulitis and a purulent cellulitis.

A phlegmon starting at the side of the cervix affects the connective tissue between the layers of the peritoneum forming the broad ligament; it may stop here, but if it spreads, it passes either (1) downwards along the round ligament to the groin, or (2) upwards to the connective tissue round the kidney, or (3) into the iliac fossa, or (4) sometimes it may extend upwards in the sub-peritoneal tissue of the abdominal wall.

A parametric abscess may spread in almost any direction; as contrasted with phlegmon, it is important to remember that it may spread over the brim of the pelvis down the thigh, or pass through the sciatic notch to the buttock, or through the obturator foramen to open at the upper and inner part of the thigh. Thus the spreading of a parametric abscess is a mechanical process, while the spreading of the inflammation in the stage of phlegmon is a vital process, not to be explained by mechanical considerations.\*

Parametric abscess may open externally, commonly in the groin, above or below Poupart's ligament, or into the vagina, rectum, or bladder, very rarely into the peritoneum. Sometimes it may open in two directions, e.g., there being one opening into the vagina, and another above Poupart's ligament; this is often due to separate foci of suppuration.

<sup>\*</sup> See Clinical Lectures on Diseases of Women (Matthews Duncan), 3rd edit., p. 236.

Remote parametritis.—When a cellulitis has spread to some distance from the pelvis, it may happen that while the inflammation and the signs of it at a distance are still evident enough, yet all signs of the cellulitis in the pelvis have disappeared. The distant cellulitis is known in such cases as "remote" pelvic cellulitis, or "remote" parametritis (Matthews Duncan).

The term "remote" is applied in the same way in cases of perimetritis.




Fig. 118.—Diagrammatic representation of the lateral position of the exudation in parametritis (Thorburn).

**Symptoms.**—The symptoms are similar to those met with in pelvic peritonitis, with the exception that pain is less marked, and vomiting occurs less commonly. The patient lies with only one leg drawn up.

Physical signs in recent cases.—The exact signs vary according as we are dealing with the disease in the stage of phlegmon, or in the stage of abscess; in either case the markedly unilateral, asymmetrical character of the signs should be observed.

In the stage of phlegmon.—There is a lump on one side of the cervix, pushing the uterus over to the unaffected

side. The lump obliterates more or less completely the concavity of the lateral fornix on the affected side, and produces an apparent shortening of the cervix on that side. The lump can very probably be reached on bimanual examination, and differentiated from the body of the uterus. The uterus and the lump have a certain small amount of mobility, differing from the absolute immobility in typical cases of pelvic peritonitis.

In the stage of abscess.—There are the physical signs of fluid, fluctuation, and dulness—that is, in cases where there is sufficient formation of pus—and it is so situated that these signs can be made out. In certain cases, where at one stage we have recognised a phlegmonous inflammation, say in the iliac region, we notice, as it reaches the suppurative stage, a softening, a "bogginess," over the area of the lump which had previously been hard, rather than actual fluctuation. As regards dulness, it may happen that intestine intervenes between the lump and the abdominal wall, and then there will be a tympanitic note on percussion.

After an abscess has burst a fistula remains open for a long while, weeks or months. A probe may often be passed two, three, or more inches along a fistula of this kind for a long time after all acute symptoms have passed away, and when convalescence is well established; in time it will close up by itself, and is best left alone. I have several times tried passing a probe, coated with nitrate of silver, along such fistulæ to hasten their healing, but have not been satisfied that it had much effect. Sooner or later, however, they have nearly always healed.

Complications.—Acute inflammation may attack the knee-joint on the affected side, or on the opposite side, and the fluid in the joint may be serous or purulent. Phlegmasia dolens is a common complication of parametritis, affecting the leg of the side corresponding to the parametritis.

#### LATE RESULTS OF PARAMETRITIS. .

As absorption proceeds, the uterus may be gradually drawn over towards the affected side in cases where there has been cellulitis in one broad ligament. In the less common cases, where there is cellulitis of the utero-sacral ligaments, contraction in this situation, pulling on the uterus somewhere near the junction of the cervix and body of the uterus in a backward direction, causes a marked anteflexion.

Clinically it is extremely difficult, if not impossible, if a case is seen at this stage for the first time, to be sure, if we find the uterus a little to one side of the middle line, that its position is not due to one broad ligament being congenitally shorter than the other, a condition that is common enough, rather than to previous cellulitis. Laceration of the cervix on the side towards which the uterus deviated would be in favour of a previous cellulitis. regards old cellulitis in the utero-sacral ligaments producing anteflexion, a difficulty of a somewhat similar character arises. For what is the evidence of old cellulitis of the utero-sacral ligaments? Merely feeling bands running in the situation of these ligaments, and concluding that the bands are a little thicker and more defined, and perhaps shorter than usual. I have noticed in many cases, where a retroflexed uterus was found incarcerated in Douglas's pouch and replaced, that after replacement these bands, the utero-sacral ligaments, were particularly easily defined, and this in cases where there was no reason to suppose there had been any previous utero-sacral cellulitis; so that, while admitting that cellulitis in this situation would cause anteflexion, as absorption proceeded, it must also be said that in practice we can rarely be certain that we have before us a case of old cellulitis of these ligaments.

Post-mortem room evidence in the case of old cellulitis is on quite a different footing from the same evidence in cases of old pelvic peritonitis; in the case of the latter the evidence in the shape of adhesions, instead of the uniform glistening character of the peritoneum, is unmistakable. If we find adhesions, there has been peritonitis; but in the case of cellulitis, long after the acute stages have passed, and absorption is as complete as it ever will be, the nature of the case renders it almost impossible to say there has at some time or other been cellulitis. Take, for instance, a case where there has been cellulitis years before on one side of the cervix, the question would be as to whether the connective tissue found in that situation was more abundant and denser than in the normal condition; and any one who will examine the connective tissue normally present there will see the difficulty or impossibility of deciding such a question. In the one hundred specimens I examined, I was unable to say positively in a single case that there had been old pelvic cellulitis. No doubt there had been pelvic cellulitis years ago in some of them; but its traces were not definite and unmistakable, as they are in pelvic peritonitis.

**Treatment.**—The treatment is similar to that in pelvic peritonitis, with such slight modifications as are at once obvious (see p. 330). Pelvic cellulitis terminates in abscess much more frequently than pelvic peritonitis. When a pelvic abscess has formed, whether it be due to perimetritis or parametritis, the best treatment is to open it with antiseptic precautions at whatever point seems most convenient. The operation is one, however, that should only be done by some one with special experience in abdominal and pelvic surgery.

Case of parametritis following labour, illustrating extension of the inflammation along the right round ligament

to the labium majus, and complicated with suppuration in the left knee-joint.

S. A. H., age 35, nine children, the last eight days ago. The last confinement was difficult, and instruments were used; this had never been the case before. Admitted to the London Hospital, December 20th, 1886, complaining of pain in the right iliac region, and of great weakness since the confinement.

Temperature on admission 102°.

On the day after her confinement she was seized with shivering fits, which occurred twice the same evening, and on the three succeeding nights. She has also had pain on defectation, and trouble on micturition; twice the urine has had to be drawn off.

December 23rd.—Rigor. Temperature 101.8°. The left knee-joint is very painful, and contains fluid. A swelling is felt in the hypogastrium reaching 2 inches above the pubes, and extending outwards about equally on each side of the middle line.

The right labium majus is swollen, so as to form a sausage-shaped swelling  $1\frac{7}{8}$  inch across, and  $4\frac{1}{2}$  inches long. The swelling reaches upwards in the direction of the inguinal canal. Left side of the vulva is normal.

Vaginal examination.—Most of the swelling in the hypogastrium is the uterus. Some indistinct thickening is felt to the right of the uterus, much less than was found in that situation on admission. Uterus fairly movable. Sound passes 3 inches. The swelling in the right labium was aspirated, and, as pus came out, a free incision was made into it antiseptically.


December 29th.—Left knee-joint aspirated; pus came out. The case was now transferred to Mr. Tay. Further treatment consisted of keeping the joint at rest on a splint, and aspirating it twice. The patient did well, and the joint recovered completely.

# Case of suppurative parametritis following labour. —Abscesses opening into the vagina, and above Poupart's

ligament.

S. J., age 32, seven children, the last a month ago, admitted to the London Hospital, June 7th, 1887.

Confinements have all been difficult. Last child delivered with forceps.






FIG. 119.—Temperature chart of S. J., case of suppurative parametritis following labour. The case was one much lower till the middle of July, when the abscess in left iliac region was forming; this was opened on July of about the average duration. June 27th. \*-Discharge of pus from the vagina; the temperature then remained 28th.† After August 11th the temperature remained normal (the word "line" is a misprint).

When a month pregnant in her last pregnancy, had rheumatic fever (there is a well-marked mitral systolic murmur), and had to remain in bed seven months. The joints chiefly affected were those of the left leg and arm.

On admission temperature 101° (see chart for subsequent temperature). Complained of forcing pain on passing her water.

Note on June 9th.—Abdomen distended. Umbilicus level with the skin. A distinct hard mass is felt occupying the left iliac region, reaching three fingers' breadths above Poupart's ligament; the mass does not extend beyond the middle line.

Measurements of the pelvis.—Between ant. sup. il. sp. =  $9\frac{1}{2}$  inches. Maximum distance between the iliac crests =  $10\frac{1}{2}$  inches. External conjugate =  $6\frac{3}{4}$  inches.

It was therefore a slightly generally contracted (or justo-minor) pelvis. Patient cannot extend the left thigh. Says she has not been able to do so for eight months.

Left labium majus larger than right.

Uterus nearly fixed. Cervix lacerated on both sides. Hardness extends outwards from the left side of the cervix to the pelvic wall, and is continuous with the lump felt in the left iliac region.

The finger in the vagina feels a somewhat cylindrical swelling, posteriorly parallel to the vagina; this is not satisfactorily accounted for by fæces in the rectum, and seems to be due to a cellulitis of the connective tissue between the vagina and rectum.

July 2nd.—On June 27th a discharge of very offensive pus in considerable quantity came from the vagina during the night. On June 29th the pus became sanious, and has continued so since.

July 4th.—With Sims's speculum the opening from which the pus is coming can be distinctly seen; it is situated on the posterior wall of the vagina, about an inch up. The swelling in the left iliac region is softer than before.

July 28th.—Where the hard mass had been in the left iliac region there is now an abscess on the point of bursting; it was therefore opened, and a drainage tube inserted.

September 5th.—The temperature has been normal since August 9th. Patient is looking fat and well. Uterus is still nearly fixed, and there is still difficulty in extending the left thigh.

#### CHAPTER XV.

PELVIC HÆMATOCELE.—EXTRA-UTERINE GESTATION.

Pelvic hæmatocele.—There are two varieties of pelvic hæmatocele:—

- 1. The intra-peritoneal, and
- 2. The extra-peritoneal.

In the intra-peritoneal variety the blood is effused into the peritoneal cavity; in the extra-peritoneal into the subperitoneal connective tissue.

#### CAUSATION.

Age.—Most cases occur between the ages of twenty-five and thirty-five (Schroeder).

Fertility.—It is more common in women who have had children.

Mode of production.—I. Rupture of an extra-uterine gestation sac is one of the commonest causes of it. If we take, for example, the case of tubal gestation, we know that, as a rule, the sac will rupture before the end of the third month of pregnancy. According to the position of the rent there will result either an intra-peritoneal or an extra-peritoneal pelvic hæmatocele. If the rupture occurs in the part of the expanded tube covered by peritoneum, then the hæmatocele will be intra-peritoneal. If, on the other hand, at the lower part of the tube, the part not covered by peritoneum, the blood will be poured into the connective tissue

between the layers of peritoneum forming the broad ligament, and the resulting hæmatocele will be extra-peritoneal.

2. Reflux of blood along the Fallopian tubes during menstruction. The menstrual group of cases.

Till about ten or fifteen years ago this group was considered to comprise almost all cases of pelvic hæmatocele. There was always great difficulty in explaining why the menstrual blood, instead of following its usual channel, should, in certain cases, regurgitate along the Fallopian tube into the peritoneum. The probability now is that, although such a thing is theoretically possible, it very rarely, if ever, occurs. Indeed, for practical purposes, clinically there need be little hesitation in regarding cases of pelvic hæmatocele as symptomatic of extra-uterine pregnancy occurring in the Fallopian tube.

The general congestion of all the pelvic viscera that occurs normally at each menstrual period might be supposed to act as a powerful predisposing cause to the occurrence of hæmatocele, and so far therefore we should not be surprised if many cases occurred in connection with menstruation.

In cases of this class blood is supposed to regurgitate from the cavity of the uterus along the Fallopian tubes into the peritoneal cavity. Why it should do so, rather than escape as usual into the vagina, we cannot explain in a satisfactory manner. Certain considerations render the occurrence more intelligible. Such are:—

- a. The fact that the inner ends of the Fallopian tubes are in very rare cases, even post mortem, found much larger than usual, admitting the ordinary sound, instead of only a fine bristle.
- b. The polarity of the uterus. When the lower part is in a state of spasm, the body and fundus are in a state of relaxation. Now, it is very likely that such relaxation may

also affect the inner ends of the Fallopian tubes, causing them to be abnormally patulous.

Causes which may produce hæmatocele during menstruation are:—

Violent exercise.

Lifting heavy weights.

Fright. (I saw a case where there were well-marked signs of hæmatocele and a history pointing to it, the onset having occurred during the earthquake at Nice.)

Coitus.

Exposure to cold.

In many cases, however, supposed to belong to the menstrual group, the vaginal discharge of blood is not really due to true menstruation, but it is the metrorrhagia so commonly met with in early cases of extra-uterine pregnancy.

Rupture of a Graafian follicle may be attended by a more abundant escape of blood than usual, and so cause hæmatocele.

- 3. Rupture of a vein, either in the broad ligament or under the peritoneun covering the uterus, may be the source of hæmatocele, either intra- or extra-peritoneal. Such a rupture would be more likely to occur if the vein were varicose, and during the congestion of a menstrual period.
  - 4. Rupture of an ovarian cyst; rupture of the uterus.

Probably almost all, if not quite all, cases of pelvic hæmatocele are due to rupture of an extra-uterine gestation.

#### MORBID ANATOMY.

Intra-peritoneal hæmatocele.—When the blood is effused into the peritoneal cavity, it gravitates into Douglas's pouch and coagulates. In twenty-four hours the coagulum has become enclosed by adhesive peritonitis set up round

it. The boundaries, for instance, of a hæmatocele just big enough to fill Douglas's pouch would be below, of course, the pouch, and above, coils of intestine glued to one another and to the edge of Douglas's pouch by adhesive peritonitis. If the effusion is considerable, the blood not only fills Douglas's pouch, but the rest of the cavity of the pelvis, and may, when coagulated and surrounded by adhesive peritonitis, form a lump that can be felt a variable distance above the pubes: for example, as high as the umbilicus. It is certain that such is the course of events when no peritonitis, old or recent, existed previous to the escape of blood; but no doubt in many cases peritonitis has existed previously and led to adhesions, so that when the blood escapes it finds itself limited by the adhesions. Thus in one set of cases the first thing is the escape of blood into the cavity of the peritoneum and its coagulation there, and then follows adhesive peritonitis in the neighbourhood of the clot; and in another set of cases there has been at some previous time a pelvic peritonitis leading to adhesions which cut off a part of the general peritoneal cavity from the rest; then, if the causes producing hæmatocele come into operation, blood is effused into the small compartment of the peritoneum so shut off.

Extra-peritoneal hæmatocele.—The blood here escapes into the connective tissue at some part of the pelvis, usually into that of the broad ligament, or at the back of the supra-vaginal cervix. The potential space being obviously limited, the quantity of blood so effused cannot be great. Exceptionally after the effusion has occurred into the subperitoneal tissue the tension of the peritoneum over it becomes so great that the peritoneum itself ruptures, and bleeding then occurs into the general peritoneal cavity.\*

<sup>\*</sup> I have seen an instance of this in a case of partial rupture of the uterus, limited to the peritoneal covering, occurring during labour.

Symptoms.—It has been already mentioned that, apart from the history, it is impossible in many cases to distinguish between pelvic peritonitis and pelvic hæmatocele. What, then, is the history?

We find that the patient was suddenly seized with severe pain in the lower part of the abdomen, that she became at the same time faint, perhaps actually unconscious. There may have been a sense of nausea only, or nausea followed by vomiting. If the quantity of blood effused has been considerable, the patient is markedly anæmic. Such a history points strongly to there having been a pelvic hæmatocele, and it becomes even more suggestive if there is a slight metrorrhagia at the time of the seizure, especially if the patient has missed the last menstrual period. From the time of onset to the time the patient is seen there will have been the ordinary symptoms of pelvic peritonitis, pain in the lower part of the abdomen, and very likely trouble in connection with micturition or defæcation. Sometimes bleeding from the vagina persists for some time after the onset.

The history to be sought which would point to the presence of extra-uterine pregnancy is something of this kind: The patient had gone two or three weeks over her time for being poorly, and fancied she might be pregnant; she then became, as she thought, poorly again. She may have passed a decidua from the uterus. There will have been more or less pain in one or other iliac region. So far the history is only that of an extra-uterine fœtation previous to rupture of the sac, and has nothing to do with hæmatocele; but if now the sac ruptures, so as to produce intraperitoneal hæmatocele, there will be the symptoms mentioned above, sudden pain in the pelvis, faintness, nausea or vomiting, collapse, and anæmia, with small rapid pulse—120 or more. During the first twenty-four hours the

temperature will probably be sub-normal, but afterwards there will be more or less fever. If the hæmorrhage is profuse, death follows; if the bleeding ceases before a fatal quantity has been lost, the blood poured into the peritoneum behaves as in the other cases, and becomes




Fig. 120.—Retro-uterine Hæmatocele (Barnes). U. Uterus; R. Rectum; A. Blood-clot. Note how the uterus is pushed forward and elevated.

encysted by surrounding pelvic peritonitis. Sometimes the sac ruptures, not into the peritoneum, but into the sub-peritoneal connective tissue, and the symptoms are then much less definite.

As regards the cases dependent on rupture of a vein.—If the result is a sub-peritoneal hæmatocele, the symptoms are

vague; if, on the other hand, it causes an intra-peritoneal hæmatocele, they are well marked, and the same as those produced in other cases by escape of blood into the peritoneal cavity.

Physical signs.—Intra-peritoneal hæmatocele.—At first, within a few hours of the occurrence, there can only be felt a sense of fulness in Douglas's pouch. In one case seen while the blood was not enclosed by adhesions, I was able distinctly to feel fulness in this position. In another, where a copious intra-peritoneal effusion of blood occurred from rupture of the uterus, abdominal palpation gave one a sense of there being something unusual in the general peritoneal cavity, without one feeling sure what it really was (till the post-mortem examination). In cases due to tubal pregnancy I have often noticed a special degree of tenderness in the corresponding iliac region on abdominal palpation; and I think this point a most suggestive one. After forty-eight hours the coagulated blood is enclosed by adhesive peritonitis, matting together the viscera in the immediate neighbourhood. Now on abdominal examination we shall feel a lump rising out of the pelvis to a variable distance above the pubes. If the hæmatocele be only a small one, we may not detect any lump by mere abdominal The abdomen is somewhat distended and palpation. tender.

On vaginal examination we find the uterus pushed forwards, close behind the pubes, but not displaced laterally. In some cases the uterus is raised, while in others it seems lower than normal. Behind it, through the posterior vaginal wall, we feel a lump more or less hard, and symmetrically situated as regards the middle line. The uterus is fixed. It is usually quite easy on bimanual examination to identify the fundus of the uterus just behind the pubes. It will be seen, on referring to the physical signs of pelvic peritonitis,

that they may be precisely the same; hence the importance of the history, on which alone we can found a diagnosis.

Extra-peritoneal hæmatocele.—The position of the lump formed by the thrombus is similar to that of the exudation in cases of pelvic cellulitis, *i.e.*, at one side of the uterus, displacing it to the other side.

Diagnosis.—As regards the intra-peritoneal variety, it will be seen that diagnosis depends on a distinct history of sudden onset being obtained following on symptoms pointing to extra-uterine pregnancy.

Marked enlargement of the uterus, softening of the cervix, and a patulous condition of the os uteri are strongly in favour of extra-uterine pregnancy.

As regards the extra-peritoneal variety, it is impossible to diagnose the existence of it with any certainty, unless there is a history of extra-uterine fœtation.

Prognosis.—If the hæmorrhage is considerable, and the blood escapes freely into the general cavity of the peritoneum, death may follow in an hour; if the quantity of blood lost is not so great as to cause death from hæmorrhage, the clot, as we have seen, becomes enclosed by adhesions, and the prognosis is the same as in pelvic inflammation (perimetritis, parametritis). Ultimately, after a period of some months, in most cases the clot becomes absorbed; exceptionally, suppuration may occur, and the case become one of pelvic abscess; and as regards prognosis and treatment, what has been said under suppurative paraand perimetritis applies.

Treatment.—If the case is seen soon after the escape of the blood, very careful consideration should be given to the question as to whether the case is really one of ruptured extra-uterine pregnancy. Here the history may help us, the patient having perhaps gone some weeks over her time without menstruating, having suffered from pain

in one or the other iliac region, having herself thought that she was pregnant. If on vaginal examination, within twelve hours after the onset of the symptoms, a tumour can be felt at one or other side of the uterus, and if the tumour is more or less fixed, there is considerable probability that the case is one of rupture of an extra-uterine gestation, because such a tumour cannot be due to hæmatocele, sufficient time not having elapsed for one to form.

In such a case an exploratory abdominal section with strict antiseptic precautions, if it can be done by some one of experience in abdominal surgery, is sometimes, but by no means always, the best practice. Each case must be judged on its own merits.

If, both on account of the history and of the physical signs (absence of a tumour in the pelvis, and merely a feeling of fulness in Douglas's pouch), we conclude that it is not a case of ruptured extra-uterine gestation, or, being a case of extra-uterine pregnancy, if we decide not to operate, the patient should be merely kept at rest in bed, with an ice-bag over the hypogastric region. Stimulants must be given according to the degree of collapse present; and if they cannot be given by the mouth, may be given by enema (3j. of brandy, with 3iij. of coffee, or with 3iij. of beef-tea). Pain must be treated by hypodermic injections of morphia.

Later treatment.—It is best not to interfere with the tumour formed by the hæmatocele unless very clear evidence of suppuration is present; then the case becomes one of pelvic abscess, and the remarks made as to treatment of this condition at p. 339 apply equally here. Opening a hæmatocele that has not suppurated is unnecessary, and indeed adds dangers of its own, one of which is recurrence of bleeding from some part of the interior of the hæmatocele; and another is, that if the coagula inside the cavity of the

hæmatocele are interfered with, there is some risk of breaking down adhesions that separate it from the general cavity of the peritoneum.

The following case is an example of hæmatocele due to the rupture of an extra-uterine gestation sac. I think it would have been better not to have opened the hæmatocele.

P. R., age 27, married eight years, two children, the last four years ago, no miscarriages, was admitted to the London Hospital on April 17th, 1886, complaining of pain in the right iliac and lumbar regions, also in the hypogastrium.

History.—Patient was last "poorly" on March 28th, 1886; the period lasted seven days, and she passed a clot one inch and three-quarters long. Before this she had seen nothing for two months. She had a dull aching pain in the lower part of the abdomen, on and off, like that which she had three years ago. The pain was somewhat relieved on March 28th, when the period began.

It seems that the pain referred to first came on three years ago when she was suckling her second child, which she did for twelve months.

She thinks it was caused by catching cold just before she expected to be unwell.

The period did not come on, and she saw nothing for six or seven weeks; then she became "poorly," and passed a clot exactly like the one she passed recently in March last.

Catamenia first appeared when she was thirteen, occurred regularly every four weeks, lasting three or four days, and were unattended by any pain till three years ago.

Since then she has had a pain for two or three days before each period.

Since marriage, has menstruated every three weeks, and has lost much more than before.

State on admission (April 17th, 1886).

Uterus movable, but less so than normal; an elastic swelling the size of a large egg, somewhat fixed, is felt occupying the right posterior quarter of the pelvis. Sound passed three inches. Cervix torn bilaterally; there is a granular erosion, or eversion, roundethe external os Temperature 100°.

From April 17th to April 24th temperature normal.

April 24th.—Temperature rose to 101°. Patient was nearly doubled

up with severe pain in the lower part of the abdomen. Twelve leeches were applied to the hypogastrium, after which the pain was relieved considerably; no marked collapse.

April 26th.—Patient began to lose blood from the vagina. Still some pain at short intervals in the same place as before.

May 3rd.—Since April 24th the temperature has gone up at night to 100° or 101°, falling to normal in the morning.

Vaginal examination (May 3rd).—The uterus is very low down, and pushed close behind the pubes. A tumour is now felt behind the uterus, bulging down the posterior vaginal wall, and extending laterally across the pelvis.

May 6th.—Temperature at night has been high since the last note. It was thought that suppuration was taking place. An incision an inch across was made into the tumour through the posterior vaginal wall close to the cervix. Some dark blackish fluid and chocolate-coloured clot escaped from the opening.

May 10th.—The finger was passed cautiously into the cavity to see if any loose pieces of clot were present. What was thought to be a large piece of loose clot was hooked out, but on subsequent examination it was seen to be a fœtus (which was shown at a meeting of the Obstetrical Society of London, and is now in the London Hospital Museum).

Great difficulty was experienced in keeping the cavity sweet, in spite of frequent irrigation of carbolic acid lotion (1-40) and the introduction of iodoform into it; and on May 16th, the temperature having been 104° for the preceding two nights, an attempt was made under ether to remove any portions of the decomposing clot that seemed ready for removal. While doing this a very sharp attack of hæmorrhage from the cavity of the hæmatocele occurred; it was only controlled by plugging the cavity with gauze. The gauze was removed two days after.

Subsequently the patient did very well, and left the hospital in the second week in July. A shallow depression in the posterior fornix with rather sharp edges remains to indicate the position of the incision. There is a little thickening round it. The uterus is freely movable.

I have thought it well to insert here an outline of the subject of extra-uterine gestation, as it is so closely related to that of pelvic hæmatocele.

#### EXTRA-UTERINE GESTATION.

It may now be considered certain that, whenever a fertilised ovum develops in any other place than the uterus, its situation is primarily in some part of the Fallopian tube. Thus all cases of extra-uterine gestation are primarily examples of tubal pregnancy.

The development of an ovum in the Fallopian tube most commonly occurs in the outer and wider part of the tube, termed the *ampulla*; less frequently in the narrower inner portion, called the *isthmus*; and very rarely in that part of the tube, called the *interstitial* portion, which passes through the uterine wall.

Etiology.—Antecedent Sterility.—As a matter of clinical experience, we generally find, in cases of tubal pregnancy, either that the patient has been previously absolutely sterile, or that a period of sterility of some years' duration has preceded the occurrence of the extra-uterine gestation. To this general rule, however, there are numerous exceptions. In one of my own cases, for instance, the patient had borne a child at term only nineteen months previously.

Age.—As might perhaps have been anticipated, the period of life during which patients are most liable to become the subjects of extra-uterine pregnancy is also that during which normal pregnancy is most likely to occur. The time in question is from twenty-five to thirty-five. In regard to this point it is interesting to observe that this period of life is also that in which pelvic hæmatocele had been observed most commonly to occur before the connection between extra-uterine pregnancy and hæmatocele had been worked out. Several of my own cases have been about twenty-eight years old. Very few cases occur after forty or before twenty-five.

As regards the causation of extra-uterine pregnancy, Mr.

Lawson Tait suggests that there has been a previous desquamative salpingitis, depriving the mucous membrane of the Fallopian tube of its ciliated epithelium. According to this view the ovum is thereby retarded in the tube instead of passing on into the uterus. Similarly the access of spermatozoa to the tube will be facilitated by the loss of the ciliated epithelium. This explanation would accord well with the fact that the patients in whom extra-uterine pregnancy occurs have generally been sterile, either absolutely or for some considerable time. Pelvic peritonitis, the result of salpingitis, is undoubtedly one of the commonest causes of sterility. In many such cases the fimbriated openings of the tubes are completely sealed up by adhesive peritonitis, and then, of course, the patient is absolutely sterile. But short of this, adhesions round the ovaries and in the neighbourhood of the openings of the tubes must interfere to some extent with the passage of ova into the Fallopian tubes when discharged from the Graafian follicles.

Mr. Bland Sutton believes "that a healthy Fallopian tube is more liable to become gravid than one which has been inflamed." It would seem to follow from this that a healthy normal woman is more liable to an abnormality than an unhealthy abnormal woman, which, though possible, seems unlikely. Nor does Mr. Sutton's dictum go well with the undoubted fact that in many cases there is an antecedent period of sterility. For in that case it would appear to follow that women with normal Fallopian tubes are more liable to long periods of sterility than those with unhealthy tubes, which is certainly not the fact. On the whole it appears to me, therefore, to be premature to dismiss the occurrence of previous salpingitis from the list of causes predisposing to extra-uterine pregnancy.

### Course and Progress.

When a fertilised ovum has lodged in the Fallopian tube, it may either—

I. From an early period degenerate into a mole. Thus there are tubal moles as well as uterine moles.

According to Mr. Bland Sutton, who has devoted much attention to this part of the subject, and done much to elucidate it, tubal moles are due to "blood extravasated from the circulation of the embryo into the sub-chorionic chamber." Or—

2. Tubal abortion may take place. When the ovum is lodged in the outer part of the Fallopian tube, changes generally take place in the neighbourhood of the fimbriated opening of the tube, which as a rule lead to its closure. This, according to Mr. Sutton, generally occurs before the eighth week. While the fimbriated end of the tube continues open, it is easy to imagine how the ovum may be discharged through it into the peritoneum. When this occurs, the patient is said to have had a tubal abortion. The ovum may be discharged entire into the peritoneal cavity, with more or less severe intra-peritoneal hæmorrhage; or several successive intra-peritoneal hæmorrhages may occur, the ovum still remaining in the Fallopian tube. This would be an instance of threatened tubal abortion, analogous to what occurs when abortion is threatening in a case of uterine pregnancy.

In tubal moles there may or may not be an amniotic cavity to be seen on cutting the mole across; and, as in uterine moles, there may or may not be some trace of the fœtus to be seen. When there is no amniotic cavity visible, the real nature of the mass may be ascertained by cutting sections of it and examining them for chorionic villi. (See Plates III. and IV.) In some cases shaggy projections

consisting of chorionic villi can be distinctly recognised with the naked eye on the outer surface of the mole. In cases where rupture of the tube has occurred these shaggy projections may often be seen protruding from the rent (Fig. 124). Or—

3. The ovum may continue to develop in a fairly normal manner for a time. At length, owing to thinning of the Fallopian tube, and continued increase in the size of the ovum, a time comes when the wall of the Fallopian tube

gives way.

According to Mr. Lawson Tait the gravid Fallopian tube always ruptures before the fourteenth week; but it often ruptures at a much earlier date. The rupture may be intra-peritoneal or extra-peritoneal. If the tube ruptures at some part of its circumference covered by the peritoneum, then we have intra-peritoneal rupture, and hæmorrhage occurs into a space the capacity of which may for practical purposes be regarded as unlimited. On the other hand the tube may rupture at that part of its circumference which is not covered by peritoneum, and lies towards the space between the layers of the broad ligament. We then have an extra-peritoneal rupture, and the blood is poured out into a relatively small space the capacity of which is limited, so that the amount lost is inconsiderable compared with what is lost in the former case.

Mr. Sutton says that intra-peritoneal rupture is three times as common as the extra-peritoneal variety.

As regards intra-peritoneal rupture, in early cases the ovum may be expelled with more or less blood into the peritoneum. If the quantity of blood is not great, the patient does not die. The effused blood forms a pelvic hæmatocele, and the whole mass is ultimately absorbed.

In more advanced cases the ovum is not always expelled, at all events completely, and more or less hæmorrhage may

continue for some time. In some cases of my own, where the fœtus has been of eight to ten weeks' development, the fœtus has been expelled into the peritoneum, while the placenta has remained wholly or partially attached within the tube. It is important to observe that rupture of the Fallopian tube may be met with in cases of tubal mole as well as in those where the tube contains a recognisable fœtus. It should also be borne in mind that rupture may occur even when the fimbriated end of the tube is patent.

As regards extra-peritoneal rupture, hæmorrhage occurs into the space between the layers of the broad ligament, and the ovum, gradually, as Mr. Sutton insists, finds its way into the loose tissue lying between those layers. It may now die and cause no more danger. This is probably what generally happens in cases of extra-peritoneal rupture. Exceptionally, however, the ovum retains its vitality, and continues its development between the layers of the broad ligament. All advanced cases of extra-uterine pregnancy have originated in this way—that is, the ovum has survived an extra-peritoneal rupture of the tube, and has continued its growth outside the peritoneum. As the resulting extra-peritoneal gestation develops it pushes the peritoneum before it, and may thus detach the peritoneum from the anterior abdominal wall.

In cases of advanced extra-uterine gestation, which originate as described, it is a matter of great importance whether the placenta is situated below, with the fœtus above; or whether, on the other hand, the placenta is at the highest part of the sac, and the fœtus below it. The latter relation is much the more dangerous for the mother.

In both cases the sac is liable to an intra-peritoneal rupture. If the placenta is situated above, at the highest point of the sac, this accident is attended with very severe, and probably fatal, hæmorrhage. If the placenta is situated below, rupture of the sac allows the escape of the liquor amnii and of the fœtus into the peritoneum. Thus the fœtus may be found, and has been found, free among the intestines.

In the event of an advanced extra-uterine gestation proceeding to term without fatal rupture a kind of spurious labour occurs, after which the fœtus dies, and undergoes mummification, or, by the deposit of calcareous salts, conversion into a lithopædion.

A case is on record where a patient survived, carrying within her an extra-uterine fœtus of this kind for fifty-two years.

In many of these advanced cases, however, when the death of the fœtus has occurred, after a longer or shorter time the extra-uterine sac becomes infected from the intestines, and undergoes suppuration. The resulting abscess ultimately opens spontaneously: it may burst into the bowel, the bladder, the vagina, or even through the uterus. It may also open externally. Fœtal bones are discharged at intervals through the opening. I have myself removed portions of the parietal bones which had become impacted in the rectum in a case of this kind. Sometimes the spontaneous opening of such abscesses is spoken of as secondary extra-peritoneal rupture.

The following is an account of a case of gestation occurring in the interstitial portion of the Fallopian tube:—

A. D., age 34, had had six children, the last about a year and ten months before admission. At the time the accident happened she believed herself to be five months pregnant. She had not menstruated for five months. Nothing had occurred during the pregnancy different from what she had experienced on other occasions. On October 12th, about 3 p.m., while going upstairs, she was suddenly seized with severe pain in the abdomen, and became rapidly blanched and collapsed. When the pain seized her, she felt as if something had given way in the abdomen, and stated that she heard a crack. Vomiting occurred soon after. She was seen by some one from a neighbouring dispensary; but

after a time, the friends becoming alarmed, Dr. Wheeler Brown was called in. This was about II p.m. He at once recognised the gravity of the case, and brought the patient up to the London Hospital. She was admitted about one o'clock in the morning, and shortly afterwards Mr. Blaxland, the resident accoucheur, sent for me. When I saw the




Fig. 121.—The Uterus from the Author's Case of Interstitial Tubal Gestation, described in the text.

The uterus has been laid open along the anterior aspect; a glass rod (E) has been passed from the cavity of the gestation sac into the cavity of the uterus. F.F. point by dotted lines to the thick decidua lining the cavity of the uterus. The specimen is in the London Hospital Museum.

I was told she had been some time before. She was able to answer questions, and in fact supplied most of the history that has been already given. The pulse was small and compressible (120). She looked pale and anxious, and complained of pain over the abdomen. The

abdomen was tender. The patient was very fat (subsequently it was found there were two inches of fat in the abdominal wall), and nothing definite was made out by palpation. She was then examined under chloroform, her own consent and that of her husband having been previously obtained to an operation if it should be thought desirable. Even under chloroform the examination was unsatisfactory. The abdomen, on palpation, had an indistinctly "doughy" feeling. On vaginal examination the uterus was made out to be much smaller than would correspond to a five months' pregnancy, and it had not the globular shape of the pregnant uterus. There was slight oozing of blood from the os uteri. According to the history there had not been more than a mere trace of blood lost from the vagina.

This being the state of the case, it will be seen that the diagnosis presented very great difficulty. First, as regards the existence of pregnancy at all, there was nothing to rely on but the patient's own statement. It is true there, was a little milk in the breasts; but in a woman who had had a child only twenty months previously, and had suckled it, this of course was worth very little. The fœtus could not be felt on palpation of the abdomen; and by vaginal examination it was certainly ascertained that the uterus did not, at the time of examination, contain an ovum of anything like five months' development. Again, considering the question of tubal fœtation, according to the patient's account, she was five months pregnant, and had not menstruated during that time, nor suffered from any pain in the abdomen; whereas, had it been a case of tubal fœtation, we should expect rupture to have taken place at least two months earlier, and also that there would have been a history of irregular hæmorrhage and more or less pain.

I came to the conclusion that the best plan would be to make an exploratory incision, as, even if the diagnosis should turn out to be wrong, the patient's risk would not thereby be much increased. On opening the peritoneum, it was found to contain a large quantity of blood and recent clot, and the hand came upon a fœtus and placenta lying loose among the intestines. The fœtus lay much nearer the liver than the uterus, the placenta being midway between the fœtus and the uterus. The membranes, which were twisted into the form of a band, passed downwards and to the left, and were found to lead to a tear in the left side of the fundus uteri. The rent was still bleeding a little. The fœtus, placenta, and blood-clots were removed, and the tear in the uterus stitched with deep sutures of silver wire and superficial catgut sutures. A glass drainage-tube was inserted at the lower angle of the

abdominal wound, and this was closed in the usual way. The patient was very cold when removed to bed. Hot bottles were applied to the feet, and enemata of brandy given. She recovered consciousness about 9.30 a.m., and complained of pain, but gradually sank, and died about 12 (noon), seven hours after the operation.




FIG. 122.—Part of the Uterus and Appendages of the Left Side, seen from above and behind, from the Author's Case of Interstitial Gestation.

The ruptured gestation is seen above, with the glass rod. The ovary has been cut open, and the corpus luteum is well seen. A. Cavity of ruptured gestation sac. B. Corpus luteum. C. Fallopian tube. D.D.D. Peritoneal coat of uterus. E. Glass rod passed from the gestation sac through the uterine opening of the tube into the cavity of the uterus.

A careful examination of the uterus was made after death. The sutures used for the tear at the operation were removed. On opening

the uterus, it became at once evident that the fœtus had never been in utero at all. The cavity of the body of the uterus was lined by a decidua from one quarter to three-eighths of an inch thick, in places partly detached. A glass rod passed from the cavity of the uterus through an aperture corresponding in position to the inner end of the left Fallopian tube, and appeared at the inner end of the tear in the fundus. Evidently then the ovum had developed in that part of the left Fallopian tube which passes through the wall of the uterus; that is to say, it was a case of interstitial gestation. The tear extended from the point where the left Fallopian tube entered the uterus inwards, and a little backwards across the fundus. The tear was two and a quarter inches long and two inches deep. At the inner end of the cavity formed by the tear, looked at from above, was a round hole, about three-sixteenths of an inch in diameter, which led into the cavity of the uterus. The whole uterus measured six and a quarter inches in length. The width between the Fallopian tubes was about five inches. The fœtus removed was exactly eight inches long.

Diagnosis and treatment.—Before rupture of the tube, it not infrequently happens that there are absolutely no symptoms leading the patient to seek advice, so that no opportunity is given us of making a diagnosis. In the case just described, a typical example of interstitial gestation, there were no symptoms of anything being wrong, and the patient thought herself five months advanced in a normal pregnancy when the fatal rupture occurred.

Still, in some cases there are symptoms; and these, in conjunction with certain physical signs, to be more particularly referred to immediately, will sometimes enable us to make a shrewd guess at the cause.

The symptoms that may be expected are symptoms common in early pregnancy, uterine or extra-uterine: as pains in the breasts and morning sickness; menstruation may be completely absent, as in normal pregnancy (it was so in the case of interstitial gestation referred to above), or there may be more or less irregular hæmorrhages; colicky pain in one or other iliac fossa is another symptom of some

significance. Perhaps a decidual cast of the uterus may be discharged.

If a patient were to come with the symptoms referred to, and if in addition we found the vagina and vaginal portion of the cervix bluish, the cervix enlarged and soft, the os uteri patulous, and the uterus somewhat enlarged, and perhaps less movable than normal; and if, besides, bimanually a swelling were to be detected in the pelvis, especially to one side, and if also there were pulsation to be felt in the region of the swelling; if also there were changes in the breasts, and if the patient is over twenty-five and has never been pregnant, or at all events some considerable time has elapsed since the last pregnancy,—the probability of the presence of an unruptured extra-uterine gestation sac will be sufficiently strong to warrant a careful consideration of operative interference—exploratory abdominal section—and if a tubal foetation be found, it can be dealt with.

I have once had the good fortune to meet with a case of this kind; and having made the diagnosis of tubal pregnancy, I operated on it successfully before rupture of the tube had occurred. The case was recorded in the *Lancet* (1895, vol. i., p. 858), and the notes are as follows:—

Case of Tubal Gestation, diagnosed before Rupture. Operation: Removal of Fætus and Sac. Recovery.

L. W., a married woman, aged 34, was admitted into Davis Ward of the London Hospital on February 24th, 1894, complaining of severe pain in the back and left iliac region.

Family history.—Her father and mother both died of "fits"; two sisters and one brother are living and healthy. There is no history of tubercle.

Personal history.—The patient is thirty-four years old. She now lives in Bow, and has always lived in London. She was married at the age of nineteen, and has had two children. The first child was born ten months after marriage, and died in eight days. The second child was born

fourteen years ago, and is still living. Both confinements were easy. Seven years ago she had one miscarriage when two months pregnant. She does no work beyond her housework.

Catamenia appeared at thirteen; were regular every four weeks and normal in amount and colour up to two months ago, since when she has seen nothing. She menstruated regularly while suckling.

Present illness.—This began about January 1st, 1894; she then felt very sick, retching, but not actually vomiting. About this time she had severe pain starting in the left iliac region, extending to the back, and going across to the right side of the abdomen. She remained in Bromley Sick Asylum from January 19th for three weeks, but obtained no relief. A week later she came to the London Hospital, and was admitted. The pain in the lower abdomen has been felt throughout; it appears constant, not worse at night than in the day, and it does not shoot down the thighs, but is particularly severe when patient goes to stool. On February 12th she first noticed a discharge, slight in quantity at first, of a reddish-brown colour, and not offensive. This has continued up to the present time. It has not increased in quantity, and as regards colour it has remained the same. She thinks she is occasionally bigger in the left iliac region than at other times, but she has not herself noticed any definite swelling. She feels sare her breasts have been getting bigger lately. She is sick at times; the sickness occurs at night and during the day, without any relation to meals. No pain is felt in the vagina, but there is considerable pain in the "back passage."

State on admission.—She is rather pale, but there is no marked anæmia. The pulse is 70, and regular; the tongue is clean and moist; the bowels are not confined; and the temperature is normal. Urine is acid, amber, s.g. 1022; it contains no albumen or sugar. On abdominal examination nothing abnormal can be felt. On vaginal examination the uterus is found to be distinctly enlarged, apparently uniformly; the cervix is soft and blue; the uterus is freely movable; the sound passes  $3\frac{5}{8}$  inches, with the curve forwards. A swelling about the size of an orange, but less regular in outline, occupies the left side of the pelvis: this swelling has a fair range of mobility. The breasts are firm; on both sides fluid can be squeezed out—that on the left side is distinctly milky.

Progress, March 5th.—She is quite definite in her statement that she was regular up to Christmas, 1893, when the last menstrual period occurred. The quantity was slight, the period lasting three days; previously the period usually lasted seven days. She is under the impression that the breasts were larger in January than now. Since Christmas she

has had severe pain occasionally in the left side, which resembles labour pains, lasting two hours. She has had two intervals of a week without pain.

A diagnosis of unruptured tubal pregnancy was made, and operation advised.

Operation, March 8th.—The abdomen was opened in the usual way.




Fig. 123.—Tubal Gestation. Case of L. W. Diagnosed and operated on before rupture.

The lower figure represents the Fallopian tube removed with the ovum. The upper figure to the right shows the swelling A in the lower figure cut across; it is due to a detached portion of the placenta developing in that part of the tube. Above and to the reader's left is a drawing of one of the loose pieces of placental tissue that came out on the sponges.

No blood was seen on opening the peritoneal cavity. The uterus early came into view, of purplish-red colour; it was considerably enlarged, feeling pulpy, and suggesting a pregnant uterus. On passing the hand down to the left side of the pelvis a swelling was felt, about the size of an orange, which burst on attempting to draw it out, and a feetus (about three months) was, so to speak, shot out of the wound, a little clear fluid escaping at the same time. The Fallopian tube was held up, the broad ligament was transfixed and tied with the Staffordshire knot, and a single ligature was passed round the whole pedicle. The tube was then cut away. One or two separate pieces of placenta about the size of a penny came up on sponges as the peritoneal cavity was sponged out (one of these is figured above). The wound was completely closed without drainage. The patient made an uneventful recovery.

As to the diagnosis at the time of rupture of the sac, this will usually be sufficiently clear if the rupture takes place into the peritoneal cavity. The patient will be suddenly seized with severe pain in the abdomen, and will probably soon show the usual symptoms of serious internal hæmorrhage. This will be quite enough to warrant an immediate exploratory operation, if it appears that the condition of the mother is not too unfavourable, and all the circumstances for doing an abdominal section are convenient: In regard to these cases Mr. Tait says: "Absolute accuracy of diagnosis in the abdomen is very far from being possible; only the ignorant assert that it is." It must be remembered, when the question of operation is being considered, that undoubtedly many cases left to nature recover, since typical cases of pelvic hæmatocele (now known to be due to extra-uterine pregnancy) generally recover. Again, when the condition of the patient is actually extremely bad, when she first comes under observation, an immediate operation of a severe character destroys the very slender chances she might have had of spontaneous recovery. The kind of case I mean is when there is profound anæmia, and a rapid, feeble pulse, say 120

to 140, and especially when from the history and signs it appears probable that the intra-peritoneal hæmorrhage has occurred some time, a day or more, before the case is seen. Probably the bleeding has stopped, and the best thing for the patient is not to operate, but to keep her at absolute rest, and to try to improve her condition in every possible way. The cases in which laparotomy is justifiable are especially those where the symptoms and signs make the nature of the case clear, and where the rupture has only occurred very recently, a very few hours before she is seenwhere also, although the hæmorrhage has been serious, still the condition of the patient is yet far from desperate. Even then one cannot avoid the conclusion that many such cases would recover well without operation. There is no doubt a widespread belief that most cases of extrauterine pregnancy require abdominal section. Nothing could be more erroneous; and if such were to be the rule of practice, it would have been better for the patients if the pathology of extra-uterine pregnancy had remained in obscurity. These remarks do not apply to operation before rupture of the tube.

I think most probably both Mrs. S. M. and Mrs. F. F., and perhaps Mrs. C., whose cases are given here, might have recovered without operation, and in similar cases in


future I should not advise operation.

The following case was one in which a tubal mole caused an intra-peritoneal rupture of the tube:—


Tubal Mole. Intra-Peritoneal Rupture, with Profuse Hæmorrhage. Abdominal Section. Recovery.

Mrs. C., 34 years of age, who had been married eleven years, was seen by me in consultation with Dr. R. Ambrose, M.P., on March 19th, 1896, at 3.30 p.m. She had had two children, the last six years ago; both her confinements were quite natural. She had had one miscarriage at

Plate 3.



Section of tubal mole showing chorionic villi cut across transversely or somewhat obliquely; from case of Mrs. C. (High Power).








Plate 4.



Section of tubal mole showing chorionic villi cut across transversely or somewhat obliquely; from case of Mrs. C. (Low Power).



about the sixth or eighth week four years ago. She got up immediately after the miscarriage, but went back to bed two days afterwards, and remained in bed for six weeks. She was then admitted into a hospital, and remained there for six weeks longer on account of what she was told was peritonitis. As regards her present illness, she had menstruated regularly every four weeks till about two months previously. January 25th was the last day of the last period. From that time she "saw nothing" until March 19th, the day on which I was called in to see




Natural size of original.

Fig. 124.—Tubal Mole (Case of Mrs. C.). The ruptured tube (right) and corresponding ovary removed.

A. Dilated Fallopian tube. C. Cut edge, where tube has been opened after removal: the dark portion (D) to the right is the mole. E. Fringes of placental tissue projecting through the rent (F) that occurred spontaneously. G. Narrow end of Fallopian tube, where it was cut off at the operation. O. Right ovary. P. Probe passed through fimbriated end of tube, which was open. H. Fimbriæ.

her. For two weeks previous to this date she had had constant pain in the right iliac region, and several attacks of slight faintness, but she had not been obliged to keep in bed. On March 19th, when she got up, while she was dressing she was seized with severe pain in the right iliac region, accompanied by a severe attack of faintness, and she had to be assisted back into bed. At this time she had a slight loss of blood, this being the first occurrence of any hæmorrhagic discharge since the last menstrual period (January 25th). I saw her for the first time at 3.30 p.m. on March 19th, when the particulars of the case already given were obtained. The patient at that time was extremely pale, the pulse was 108, and there was a marked tendency to syncope on the slightest exertion. It happened that the head of the bed was towards the window, and, in order to make a more satisfactory examination, I had the patient moved so that her head was at the foot of




Natural size of original.

Fig. 125.—Case of Mrs. C. Right Fallopian tube laid open above, showing D, the mole. E. Part of the same, consisting of spongy tissue protruding through the rent that occurred spontaneously. C.C. Cut edges of dilated Fallopian tube. G. Inner end of tube.

the bed. While doing this she almost fainted away, the pulse stopping for a few seconds. On examining the abdomen there was some fulness about the hypogastric region, and an indistinct sensation of a swelling there and in the right iliac region. But the lower abdomen was extremely tender, so that no very complete examination could be made. On vaginal examination, there was a little recent blood in the vagina, the uterus was distinctly less movable than normal, and there was an indistinct swelling to the right of it. The breasts had an active

appearance. It appeared to me that the case was one of ruptured tubal pregnancy, and as the surroundings of the patient were somewhat inconvenient in the event of operation I had the patient removed in an ambulance to the London Hospital. I saw her again at 9 p.m. on the same evening; her condition then was decidedly worse than when I had seen her earlier in the afternoon; the pulse especially was weaker and more frequent. It had been 108 in the afternoon at half-past three, whereas at nine o'clock it was 120 and decidedly smaller. I thought that abdominal section was the right procedure, and the




Natural size of original.

Fig. 126.—Case of Mrs. C. The Uterine Appendages of the Left Side, also removed at the same operation.

A. Left Fallopian tube dilated and containing blood-clot (hæmato-salpinx). H. Traces of fimbriæ belonging to outer end of tube, which was closed. O. Left ovary. G. Inner cut end of tube.

patient and her friends readily agreed that it should be done. On opening the abdomen, there was a large quantity of blood free in the peritoneal cavity; on passing the hand down to the right, the right tube was drawn up to the wound, and was seen to be considerably enlarged. There was a rent in it somewhat anteriorly, from which a spongy substance was protruding; at this point bleeding was still going on. The uterine appendages on the right side were then

removed in the usual way; the uterus was observed to be considerably and uniformly enlarged, and indeed, as one has observed in several similar cases, had all the appearance of the pregnant uterus. On examining the appendages on the left side, it was found that these were by no means normal, for the left Fallopian tube was considerably dilated, being an inch or more in diameter, and accordingly I also removed the appendages on the left side. The abdomen was washed out with warm saline solution repeatedly, till all the clot and blood which had been effused seemed to have been washed away. A Keith's



Natural size of original.

Fig. 127.—Case of Mrs. C. Left Fallopian Tube dilated and containing Blood-clot.

G. Inner end of tube. C.C. Cut edges of dilated tube where it has been laid open. The walls of the dilated tube are thin, and the clot is not adherent to the inner surface of the tube.

tube was inserted, and the rest of the abdominal wound closed in the usual manner. The patient made an uninterrupted recovery; the temperature after the operation was only once above Ioi° F., and that on March 22nd. I saw this patient on May 28th, when she came to report herself at the hospital; she was then quite well, and the abdominal wound was soundly healed.

On examining the appendages removed, there was a large corpus

luteum in the right ovary about three-quarters of an inch in diameter; the right tube had, as already mentioned, ruptured anteriorly, and from the rent a small portion of spongy tissue was projecting. Subsequent examination of a section of this spongy tissue showed that it contained an abundance of chorionic villi. (See Plates III. and IV.) The tube itself contained a hard clot, which, before the tube was opened, felt as if it might have been a fœtus; but no fœtus was found. The fimbriated extremity of the right tube was open. The left Fallopian tube, which was considerably dilated also, was found on section to contain a firm blood-clot, not attached to the inner surface of the tube. The wall of the left tube was extremely thin, very much thinner than the wall of the right Fallopian tube. The left ovary was somewhat large, but contained no corpus luteum. The fimbriated extremity of the left tube was closed.

The following case is an example of tubal gestation with intra-peritoneal rupture at about the third month:—

Mrs. N., a married lady, aged 28, was seen by me in consultation on April 1st, 1893. Her history was as follows:-She had been married twice—the second time a year and eight months ago. She had had one confinement seven or eight years ago, and was said to have had three miscarriages since her second marriage. She came on "poorly" on January 5th, 1893, instead of the 13th, which would have been the proper time. Her husband left her on January 28th, and was away till March 25th. She menstruated again on February 9th, and, except for an interval of two days, she had a red vaginal discharge from that time till I saw her in April. Pain in the right iliac region had been a symptom from about the beginning of March. On April 1st, at 3 a.m., she was seized with severe pain in the stomach and back, and felt faint and sick. I saw her about 5 p.m. the same day; she was in bed, but she had rallied a good deal since the early morning, and was able to tell me most of the history that has been given above. The breasts had an appearance of activity, and contained milk; she was sure they were larger than they had been. An indistinct fulness could be felt in the hypogastrium, but tenderness interfered a good deal both with this and with the subsequent vaginal examination. The vagina and cervix had a distinct blue tinge. The uterus was considerably enlarged, and there was a soft swelling, with an indefinite outline, to the right of the uterus. It seemed to be very probable that the case

was one of extra-uterine pregnancy, although it was possible to explain the various features of the case by supposing it to be one of threatened miscarriage or molar pregnancy; on that supposition the swelling to the right would be an accidental complication such as a small ovarian tumour or dilated Fallopian tube. Soon afterwards the patient was removed to Fitzroy House. I was again sent for to see her on April 17th, about 6 p.m., as she had been once more seized with great pain in the right iliac region and faintness. The red discharge had persisted more or less. I advised that she should be examined under chloroform and that the cervix should be dilated, so that the condition of the interior of the uterus might be ascertained. This was accordingly done on the next day. I found the uterus to be uniformly enlarged, and its cavity empty. The sound passed four inches. The interior of the uterus was soft and pulpy; there was nothing to account for its enlargement except the view that the swelling felt to the right of it was an extra-uterine foetation. At this time the patient was extremely anæmic; the abdomen was moderately distended and tender. I advised abdominal section, and it was arranged to perform it on April .22nd. Between the 18th and the 22nd the distention of the abdomen increased somewhat, but it was never extreme; there was great pain, especially marked in the right iliac region. The pulse was 104, and there was no rise of temperature. Morphia and hot fomentations were used in order to relieve the pain.

The operation was undertaken on April 22nd, at 10.30 a.m. On opening the abdomen the peritoneal cavity was found to contain a large quantity of blood and clot. Some of the clot was more recent than other portions. The fluid blood was dark and of a somewhat chocolate colour. A feetus of about three months' development was found to be lying among the intestines; the umbilical cord was ruptured about two inches from the fœtus, but I am not sure whether this may not have been accidentally done when exploring the condition of things with the hand in the first instances. The right Fallopian tube was found to have been the seat of the pregnancy. It had ruptured towards the outer part and anteriorly; there was a second rupture more posteriorly, out of which a shaggy piece of placenta, the size of half a walnut, projected into the peritoneal cavity. There was no evidence whatever of peritonitis, there being no lymph, and the surface of the peritoneum everywhere was smooth and shining. The ruptured sac was held up, and the broad ligament below it was transfixed and tied. Another ligature was then tied round the whole pedicle, and the sac was cut away. The peritoneum was then washed out very thoroughly with hot iodine water, containing one drachm of the tincture to each quart of water. About twelve jugfuls of this were used. After about five jugfuls had been used, and the lower part of the peritoneum seemed to be quite clean, a large quantity of dark blood came from the region of the liver, having been shut off from the effect of the previous washing by the left hand holding back the intestines. Finally, the water used for washing came back quite clean, and the wound was completely closed in the usual way.

The subsequent progress of the patient was as follows:-Convalescence was delayed much longer than is usual in uncomplicated cases of abdominal section, such as ordinary cases of ovariotomy. The sutures were taken out a week after the operation-i.e., on April 29th-and the wound was found to be completely healed. This was about her worst day; she was very restless and excitable. At 8 p.m. on that day her pulse was 140, and the temperature 102.8° F. She had slight hiccough and retching, bringing up a little dark bitter fluid into her mouth, but there was no actual vomiting. On May 3rd there was swelling of the right submaxillary and parotid glands, which persisted for some days, but afterwards disappeared without suppuration taking place. On the 7th the wound spontaneously reopened at the lower end, and a good deal of thin bloodstained fluid came away. On the next day the discharge became more purulent. The probe passed two inches at the lower angle of the wound, and a small drainage-tube was inserted there. This was gradually shortened from time to time, and was finally discontinued on the 19th. The temperature was highest in the third week after operation, when it reached 104° on three occasions. It was during this week that the wound broke down at the lower end as already mentioned. On June 12th the patient was allowed to sit up for three hours, and on the 15th she went for a drive. On the 18th there was some yellow discharge from the vagina, which continued for some days. On the 22nd she left London to visit her mother; at that time there was still a small sinus at the lower angle of the wound. After an absence of two months she returned to London and came to see me; the sinus had been for some time closed, and when I examined her there was a sound cicatrix at the situation of the wound. She had completely regained her ordinary health.

An interesting point about this case is that about a year afterwards she suffered from somewhat similar symptoms,

and a diagnosis of tubal gestation was again made, and abdominal section decided on. The day before the proposed operation the patient passed a blighted ovum the size of a hen's egg from the uterus, which had been known to be enlarged. The operation was undertaken as arranged, and the swelling to the left of the uterus which had been thought to be a tubal gestation was found to be a small ovarian tumour about the size of a cocoanut. It was removed, and the patient made a good recovery. Some portion of the left ovary (which was degenerated so as to form the cyst) must nevertheless have furnished the ovum, which formed the blighted ovum discharged from the uterus the day before the operation. The case is interesting also as an instance of two abdominal sections on the same patient.

Another case, interesting as one of tubal abortion (threatened), is the following:—

Tubal Mole. Threatened Tubal Abortion, with Intra-Peritoneal Hæmorrhage from the Open Fimbriated End of the Tube. Laparotomy: Removal of the Dilated Tube. Recovery.

S. M., aged 28, was admitted under my care into the London Hospital on March 24th, 1894. She had been married two years and a half, but had had no children or miscarriages.

History of the present illness.—She says her illness started at the beginning of March with pain in the left iliac region, which has steadily increased in severity up to the time of her admission. She thought there was a lump in the side where she felt the pain.

Menstrual history.—She menstruated first at the age of thirteen, and has always been regular every month up till a week before Christmas, 1893, when the last proper period occurred.

On March 10th, 1894, she passed something "like a piece of flesh," and about an hour afterwards lost a good deal of blood. More or less metrorrhagia continued till she was admitted into the hospital.

On examination then the breasts presented an active appearance,

and contained milk. The uterus was distinctly and uniformly enlarged, and was soft. The cervix was blue and soft. There was a swelling the size of a Tangerine orange felt in the left side of the pelvis. The temperature and pulse were normal.

Taking all the points of the case into consideration, it seemed very probable that the case was one of tubal pregnancy, and I advised

abdominal section, to which the patient consented.

Operation, March 27th, 1894. - The patient walked to the operation table. After she was anæsthetised she was placed in Trendlenberg's position. On opening the abdomen there was no blood diffused in the general peritoneal cavity; but when the intestines were held out of the way, a mass, estimated at a teacupful, of recent "currant-jelly" clot could be seen in the left posterior quarter of the pelvis. The uterus was considerably enlarged, and was very soft; it presented the bluishred colour seen either in the ordinary pregnant uterus, or in the uterus when sympathetically enlarged in cases of extra-uterine pregnancy. To the left of it was the swelling referred to above, which proved to be the left Fallopian tube. This was held up, and the broad ligament below it transfixed, and tied with the Staffordshire knot; another ligature was then tied round the whole pedicle, and the tumour removed. The peritoneum was washed out with several jugs of weak iodine water, and a Keith's tube having been inserted in Douglas's pouch, the rest of the wound was closed in the usual way.

As regards the specimen, the tube had not ruptured, and the fimbriated end of it was open. The clot seen in the pelvis had evidently come from the tube through the open fimbriated extremity.

The patient made an uninterrupted recovery, and was discharged on April 17th.

The specimen was referred to a Committee of the Obstetrical Society of London, consisting of Mr. Doran, Mr. Bland Sutton, and myself. The Committee agreed that it was a genuine example of tubal mole.

A Successful Case of Intra-Venous Injection of Salt Solution after Laparotomy in a Case of Ruptured Tubal Gestation.

On April 20th, 1896, I was called in by Dr. Fennell, of Dalston, to see Mrs. F. F., aged 30 years, when the following history was

obtained:-She had been married nine years, and had had four children, the last nineteen months previously. There had been no miscarriages. Instruments were used at the first confinement, but not at the others. She generally got up ten days after her confinements, but after the last confinement she was in bed nearly two months on account of what the medical attendant told her was peritonitis, Menstruation was quite regular up to February 26th, 1896, but she "saw nothing" in March. She took some pills on April 4th with a view to bringing on the period; these gave her great pain. On April 6th a vaginal discharge containing blood began, and she had been losing blood more or less ever since up to the day when I first saw her (April 20th). The patient said that a solid substance came away from the vagina on April 11th when passing urine. She had had some pain in passing urine, and had suffered from constipation since April 6th. She said that she had had pain at the lower part of the abdomen, especially over the left iliac region, for some days before I saw her, and there was an especially sharp attack of pain on the early morning of April 20th. On examination of the abdomen there was some resistance in the left iliac region and some tenderness there. The breasts did not appear active. On vaginal examination there was a little blood to be seen about the external parts. The uterus was anteverted and slightly flexed and distinctly larger than normal, although the sound only passed three inches. A swelling was felt behind and to the left of the uterus, occupying the left posterior quarter of the pelvis. It was distinctly elastic, and could be displaced a little by upward pressure. The case appeared to me in all probability to be one of extra-uterine pregnancy. The question arose as to whether the substance passed by the patient a few days before I saw her might not have been an ovum, and the case one of ordinary miscarriage, with possibly some peri-uterine inflammation; but, on the whole, it seemed to me much more probable that the case was really one of tubal pregnancy, and that the substance in question said to have been passed, if anything more than a clot, was probably a decidual cast of the uterus. The arrangements of the patient's home were not very convenient for performing abdominal section, and accordingly I had her removed to the London Hospital, where she was admitted under my care on April 21st.

Operation.—On April 23rd she was anæsthetised with the A.C.E. mixture; the abdomen was opened in the usual way, and was found to contain blood and masses of clot. The left Fallopian tube was considerably enlarged and had ruptured. The left ovary contained a large

corpus luteum with a well-marked yellow convoluted border. The uterine appendages on the right side were also slightly enlarged, and the uterus was considerably enlarged. The appendages on both sides were removed in the usual way, and the peritoneum well washed out with saline fluid. A Keith's tube was inserted, and the rest of the abdominal incision was closed. When this had been done, it was observed before the dressing was put on that the Keith's tube had become full of dark blood. This was withdrawn once or twice, but the tube very soon became again full of blood. The abdomen was therefore reopened to discover the source of the bleeding; the stumps of the broad ligaments were pulled up and inspected, but no blood was coming from them. On looking at the back of the uterus there were some quite superficial tears in the peritoneal coat near the level of the internal os, from which blood was coming rather freely. Several attempts were made to control this bleeding by passing silk ligatures on a curved needle under the tissues at the bleeding point and tying, and also by applying Wells's forceps; but the tissues were so friable that the more manipulation took place in the neighbourhood of the tears the worse the bleeding became. I therefore decided to perform supravaginal hysterectomy; the uterine arteries were ligatured on each side outside the uterus, and then the body of the uterus was cut away. A large pair of Wells's forceps was left on, but the surface of the stump oozed rather freely, and two silk ligatures were passed right and left from before backwards and tied firmly to check it. Some iodoform gauze was packed over the stump, a small piece being left sticking out at the lower end of the wound. A Keith's tube was inserted, and the remainder of the wound closed in the usual way. The reopening of the wound and the various measures that were required to permanently control the bleeding had occupied a considerable time, and, besides, a good deal of blood had been lost, taking into consideration what was extravasated in the abdomen by the extra-uterine sac before operation, and also reckoning the amount that was lost from the superficial tears on the back of the uterus. How these were produced I am unable to say positively. My impression is, however, that some of the masses of clot removed were partially adherent to the uterus, and that in removing them the peritoneal coat of the uterus must have been slightly torn, and then the tissues were so friable that almost any kind of manipulation sponging, pulling up the uterus, and the like-seemed to increase the laceration and the hæmorrhage. At all events, at the end of the operation, when the patient was put back to bed, her face was quite

white, her hands and feet quite cold, and no pulse could be felt at the wrist; and I have no doubt whatever that had she been left without further treatment she would have died, probably without regaining consciousness at all. I decided, however, to try the effect of injecting salt solution into the veins. Our stock of boiled water had been exhausted in washing out the abdomen during the operation, so that the water used in making the salt solution for intra-venous injection was merely the water out of the kettle and out of the tap. The salt solution, however, was sterilised, because we keep a concentrated solution of it, which is carefully prepared and boiled before each operation. A glass reservoir was filled with salt solution at the temperature of 100° F., and containing a drachm of common salt to each pint. The injection was made into the median cephalic vein of the left arm, the necessary pressure being obtained simply by hanging up the glass reservoir on the wall about three feet above the patient's head. After one pint had been injected the patient's pulse at the wrist could be felt. After four pints had been injected the pulse was as good as it had been before the operation. The patient made a good recovery. I have seen her several times since she left the hospital; the abdominal wound is soundly healed, and she remains quite well.

As regards the treatment to be adopted in advanced cases of extra-uterine gestation, at, or near, or past the full time of gestation, the right course will generally be to operate. As to the time for interference, as the fœtus, even when alive, is almost always imperfect in some way, and little like to survive in any case, the operation should be performed at the time when all the conditions are likely to be the most favourable for the mother. This is some weeks after the death of the fœtus. If the operation is undertaken when the fœtus is living, it will be wise not to interfere with the placenta: this is especially so when the placenta is in the pelvis below the fœtus. The cord is cut short close to the placenta, and this is emptied of blood as much as possible. The sac is then thoroughly washed out, and the abdominal wound completely closed, in the hope that the placenta may be absorbed. If symptoms of

septicæmia arise, a second operation may be undertaken to remove the placenta. After the death of the fœtus the placenta may probably be removed without any great risk at the same time as the fœtus.

Among works that may be advantageously consulted by those wishing to pursue the subject further are:—Surgical Diseases of the Ovaries and Fallopian Tubes, by Mr. Bland Sutton, 2nd edit.; Lectures on Ectopic Pregnancy and Pelvic Hæmatocele, by Mr. Lawson Tait; The Diagnosis and Treatment of Extra-Uterine Pregnancy, by Dr. Strahan; Diseases of the Fallopian Tube, by Dr. Cullingworth; and various papers to be found in the Transactions of the Obstetrical Society of London, more especially in those of the last ten years.

## CHAPTER XVI.

DISEASES OF THE OVARIES.

## OVARIAN TUMOURS.

Tumours of the ovary may be divided into :-

- I. Solid.
- 2. Cystic.
- 1. Solid ovarian tumours.—All solid tumours of the




Fig. 128.—Myoma of the Ovary (Doran).

ovary are comparatively rare:—It is sufficient here to mention that they are occasionally met with.

Non-malignant solid tumours are fibro-myomata (Fig. 128).

Malignant solid tumours are either sarcomata, or carcinomata. (See p. 197 for case of ovarian tumour for the most part solid.)

2. Cystic ovarian tumours,—Origin.—It would be out of place here to discuss in any detail the various theories that have been put forward regarding the origin of these tumours. Small cystic tumours of the ovary may certainly originate from fully developed Graafian follicles



Natural size of original.

Fig. 129.—Small Multilocular Ovarian Tumour.

B.B.B. Irregular projections due to separate cysts. A indicates where the tumour was cut off after the pedicle had been tied. (From a case of the Author's.)

by a process of dropsical distention. Probably that shown in Fig. 129 may have originated in this way. Most probably, however, though not certainly, that is not the way in which the ordinary large cystic tumours are formed. Considering the enormous number of Graafian follicles that may be seen in the parenchyma of the fœtal ovary, and the much smaller number that can be recognised in the ovary at

puberty, it seems certain that by far the greater number of these follicles are destined to undergo a process of atrophy, without ever becoming mature, and rupturing. Certain appearances are seen in sections of an adult ovary under the microscope which are probably due to relics of those follicles that have atrophied without passing through the stages of maturation and rupture. It is highly probable that the ordinary multilocular ovarian cyst originates from some disturbance of the process of atrophy that normally affects the majority of the Graafian follicles seen at birth in the ovary. Certain of the follicles that should atrophy, instead of doing so, take on a process of growth and give rise to ovarian cysts. This theory, then, supposes some failure or defect in the process of involution.

In a paper in the Pathological Society's Transactions for 1886, Mr. F. Eve suggests that cystic disease, or cystic adenoma of the ovary, may originate from certain remains of the germinal epithelium of the ovary, which are not immediately converted into Graafian follicles. In regard to these remains Klein writes: -All appearances are in favour of the view that these masses of epithelium increase in the adult, and that their cells change into ova some time after birth. Eve supports his suggestion by reference to a specimen of small cystic ovary in which he thought the development of cysts from such groups of epithelium was observable. According to this view the epithelial growth stops short of the formation of Graafian follicles, but cysts are produced, from the walls of which tubular prolongations spring; these become dilated, and form secondary cysts. This theory supposes the fault to be something erratic in the process of evolution.

There are reasons for thinking that ovarian tumours, in the interior of which papillary growths develop, have originated from the remains of the Wolffian body found in the hilum of the ovary (see Fig. 130). In small tumours of this kind the parenchyma of the ovary can be recognised at



Fig. 130.—Incipient, Papillomatous Cyst of the Hilum of the Ovary.

The free portion of the ovary projects above, posteriorly (Doran).




Fig. 131.—Small Multilocular Ovarian Cyst, slightly reduced from natural size (Doran). The, Fallopian tube and its fimbriated extremity are seen at the upper part of the figure.

some part of the circumference of the tumour little, if at all, changed.

Dermoid cysts are produced by an abnormal inclusion of the epiblast.

Either the connective tissue elements in the walls of the cysts, or the epithelioid elements on the inner surface, may take on a malignant development. Speaking generally, the more solid tissue there is entering into the composition of an ovarian tumour, the more likely it is to be malignant.

## Varieties.

- 1. The variety due to dropsy of the Graafian follicles.
- 2. Proliferating ovarian cysts.
- A. Those originating in the parenchyma, with no papillary growths on the inner surface of any of the cavities, are known as *proliferating glandular cysts*.
- B. Those originating in the hilum of the ovary, and showing papillary growths in the interior of the cysts, are called *proliferating papillary cysts*.
  - 3. Dermoid cysts.
  - 4. Malignant cysts.

As has been mentioned, it is certain that some small ovarian cysts originate by dropsical distention of fully developed Graafian follicles, while it is highly probable that this is not the mode of origin of the common multilocular ovarian tumours (proliferating glandular cysts).

According to Doran, small tumours due to dropsy of the Graafian follicles are distinguished on inspection from small multilocular ovarian cysts of the proliferating glandular kind by the fact that, in the former variety, "the cysts bulge from the free border of the ovary," while in the latter "the ovary enlarges in a uniform manner." Again, according to the same authority, cysts originating from the remains of the Wolffian body in the hilum (proliferating papillary ovarian cysts) differ from proliferating glandular cysts, in

that, in the former, the parenchyma of the ovary, comparatively unaltered, can be recognised at some part of the circumference of the tumour (this is, however, not always the case); while in the latter it cannot be so recognised,—the enlargement of the ovary in the last case having been uniform.




Fig. 132.—Dermoid Cyst of the Ovary (Doran).

THE FLUID CONTAINED IN OVARIAN CYSTS.

This is usually glairy in consistence, and as regards colour it may be either almost colourless, grey, or yellowish-grey; green or reddish-brown, or dark and tarry. The material may be of a colloid consistence, that will not run through a canula.

According to Doran, secondary cysts in which papillary growths are found contain a clear non-glairy fluid.

## CELLS FOUND IN OVARIAN FLUIDS.

Cylindrical epithelium cells may be found, and these are the most important microscopical constituents, as indicating that the fluid is ovarian. Small corpuscles (Drysdale's corpuscles) about the size of pus cells, with granular contents, but no nucleus, are found in ovarian fluids; they are now believed to be free nuclei of epithelium cells that have undergone fatty degeneration. They have been found in other fluids than those taken from ovarian cysts, and are not pathognomonic, although suggestive.

Microscopic structure.—A section of the cyst-wall of an ovarian tumour shows a single layer of epithelium on the peritoneal surface, flat in the case of large cysts, cubical if the cyst is small. The substance of the cyst-wall is composed of fibrous tissue, either mature or immature, and on the inner surface is a layer, in the case of large cavities flattened, or in the case of smaller cavities of cylindrical, epithelium cells. The papillary growths found in the interior of some ovarian cysts are mainly formed of connective tissue, covered towards the free aspect of the papillæ with cylindrical epithelium cells. Whether we find cylindrical or flattened epithelium, appears to be chiefly a question of pressure.

Dermoid cysts are generally unilocular. The inner surface is lined by a tissue like skin, provided with sebaceous follicles, sometimes with sweat glands and hairs. The cavity of the cyst is filled with a putty-like material, in which are loose hairs that have been shed. The hairs are usually reddish or light, rarely dark.

In the tissue of the cyst-wall bones are often found; teeth also may be present, either attached to bones, or isolated in the connective tissue of the cyst-wall. Teeth are also found loose in the cavity of the cyst. Dermoid cysts may also contain cholesterine crystals, nerve and brain substance, and non-striated muscular fibre. According to Winckel, striped muscular fibre and nails have not been found in them.

Dermoid cysts are usually unilateral, sometimes bilateral. In size they vary from that of a walnut to that of a football. Sometimes dermoid cysts are found co-existing with ordinary cystic degeneration in the same ovary. Recently I had a case of this kind, in which there was a large multilocular




Fig. 133.—Section from a Carcinomatous Ovary (Doran). 2-inch and \frac{1}{8}-inch objectives.

ovarian tumour of the usual kind on the left side, and on the right side the ovary was enlarged to the size of a breakfast cup, partly by cystic change of the same kind as that on the other side, and partly by the presence in it of a unilocular dermoid cyst the size of a Tangerine orange.

Ordinary ovarian tumours may be composed of one large cyst, or a few large cysts, or of a great many cysts. Certainly ovarian tumours are met with clinically where there is only one cyst, as far as can be discovered by the naked eye. This has occurred either from the tumour being formed by the growth of one cyst; or by the tumour

having, at an earlier stage of its existence, been formed of several cysts, the septa between them having at a later period become broken down. Traces of this are sometimes to be seen in apparently unilocular tumours.

CHANGES THAT MAY OCCUR IN AN OVARIAN TUMOUR.

It may inflame;

Its pedicle may become twisted;

Hæmorrhage may occur into its interior; or

It may rupture.

Inflammation may occur, and if it affects the outer surface of the tumour, lead to peritonitic adhesions. Adhesions to the omentum, to the parietal peritoneum, and to intestine are the most common. Deep adhesions in the pelvis are not at all rare. Among the least common are adhesions to the bladder. If inflammation affect the interior of the cyst, it may lead to suppuration. This is particularly likely to occur after tapping (though it often occurs independently of it), especially if air gain entrance into the cyst. It also often occurs owing to infection from the bowel when the cyst is adherent to it.

Twisting of the pedicle.—This may, in rare cases, be sufficient to cause gangrene of the tumour.

Hæmorrhage into the interior of the tumour, either spontaneously, or after twisting of the pedicle, may occur.

Rupture.—Ovarian cysts occasionally burst, either from mere distention, or from mechanical violence. The contents escape into the peritoneum, and in most cases quickly set up fatal peritonitis; rarely, if the fluid contained in the cyst was thin and unirritating, it may be absorbed, and in that case rupture is to be regarded as a process of natural cure.

Symptoms.—The symptoms that commonly attract the patient's attention in cases of ovarian tumour are:—

Increase in the size of the abdomen.

Pain at some part of the abdomen.

Disturbance of menstruation.

Usually the first thing noticed is the increasing size of the abdomen.

Sometimes, however, pain in the abdomen is the first thing noticed, and sometimes some disturbance of menstruation. Sometimes trouble connected with micturition, or rarely even retention of urine, may be the first symptom; and in one of my cases prolapse of the vaginal walls, occurring in an unmarried girl of twenty-three, was the first thing to lead to an investigation.

Menstruation is frequently disordered in cases of ovarian tumour; usually the disturbance is in the direction of amenorrhæa, the quantity lost on each occasion being less than formerly, or the intervals between the periods being increased, or both of these conditions may be present. But it is not so very uncommon to meet with cases where the disturbance of the menstrual function is in the other direction, i.e., where there is menorrhagia. I have seen several instances of this. Sometimes again, in cases of ovarian tumour, menstruation is painful; sometimes, on the other hand, whereas menstruation had been painful from the time when the function became established, from the time the tumour was first observed menstruation has become painless. This was so in one of my cases, and it was also one where there was menorrhagia. Sometimes we meet with cases where menstruation continues to be performed in a perfectly normal manner.

A Note on the Occurrence of Metrorrhagia after the Menopause in Cases of Ovarian Tumour.

During the years of menstrual life—that is to say, from the age of puberty, which may be taken as about fifteen, up to the menopause, which is normally established between the ages of forty-five and fifty years—the influence on menstruation exerted by the development of an ovarian tumour is well known. In about one-third of all such cases menstruation is unaffected; as regards the remainder, in the majority there is a tendency in the direction of amenorrhoea, the periods occurring either less frequently or the quantity lost each time becoming diminished, or both these conditions may be combined. In a minority of cases the development of an ovarian tumour is marked by the occurrence of menorrhagia. When, however, we come to enquire about cases where an ovarian tumour commences to develop after the menopause, as to whether hæmorrhagic discharges from the uterus may be caused by the ovarian tumour and be rightly regarded as symptomatic of it, we find little or no information on the point in the majority of books to which reference would most naturally be made. Thus, for instance, in the works on the Diseases of Women by Barnes, Thomas, Hart and Barbour, and Skene, there appears to be no notice of the point, and the same applies to several other works. In West and Duncan's Lectures on the Diseases of Women there is an analysis of ninety-four cases of ovarian tumour with reference to the way in which menstruation was affected. It is merely stated there that "in seven cases menstruation had ceased before the disease began." Similarly in the System of Gynacology by Allbutt and Playfair recently published there is an analysis of a hundred and eighteen cases as regards this point, and it is mentioned that twenty were cases before puberty or after the menopause, apparently implying that no menstruation or pseudomenstruation was to be expected as the result of the growth of the tumour in cases of this kind. It may fairly be concluded that the occurrence of bleeding from the uterus after the menopause in cases of ovarian tumour is a somewhat rare event. That this symptom may occur is, however, mentioned by Pozzi in the following passage: "After the menopause congestive phenomena have been noted in the uterus leading to the reappearance of a more or less irregular bloody discharge, which makes the patient believe menstruation has returned." At this passage there is a reference to a paper by Terrier in the Revue de Chirurgie, 1884, p. 1; but, on looking up this paper, it appears that he is speaking of the appearance of bleeding subsequent to the operation for removing the tumour, which is quite a different thing. The passage by Terrier is as follows: "To resume, in women already aged, accustomed, so to speak, to the menstrual flow, the periods have a certain tendency to reappear after the removal of both ovaries. But this red discharge only lasts a few months, and is not slow to disappear altogether."\* In Garrigues' Diseases of Women, 1894, p. 565, the point in question is clearly but very briefly mentioned. He says: "On the other hand, even after the menopause, new hæmorrhagic discharges from the uterus may occur."

A special importance attaches to the causes of bleeding after the menopause, since in such a large proportion of cases, as is well known, this symptom is due to the presence of a malignant growth at some part of the genital tract. It is interesting, therefore, to keep in mind that this symptom may in rare cases be due to the fact that an ovarian tumour is in process of development. While the tumour is still small, especially in a very stout patient, the occurrence of bleeding after the menopause may not improbably excite a suspicion of cancer. Without doubt in all such cases the possibility of the presence of cancer should always be thought of first; but when cancer has been excluded by careful and repeated examination—and in some cases this is only possible after dilatation of the cervix—it is satisfactory to bear in mind

<sup>\*</sup> Revue de Chirurgie, 1882, p. 367.

that there are causes for metrorrhagia after the menopause other than cancer, and that the development of an ovarian tumour is, in rare instances, one of them. The following case is an example of it:—

P. P. H., aged 54 years, was admitted into the London Hospital under my care on January 6th, 1896. She was a widow, and had had seven children. She was sent to me by Mr. W. H. Piercy Fox. She had first noticed a swelling in the abdomen about six months previously, but she thought that it might have been present longer without her noticing it, as she was very stout. The swelling in the abdomen had increased very rapidly, she thought, during the last six weeks, so that, in her opinion, the abdomen was now twice the size it was three months before. For the last three months she had suffered very much from dragging pains and general discomfort in the abdomen with dyspnœa. She was only fairly comfortable when lying on her right side. About three years previously the catamenia ceased, and she "saw nothing" for two years. There was then a period of bleeding, some blood coming away daily for six months. Then there was an interval of three months without bleeding. During the last month there had been a continuous red discharge daily. Up to three years previously she had been regular. She had "suffered from her chest and heart" for some years, with a good deal of shortness of breath. On examination the physical signs were those of an ovarian tumour reaching six and a half inches above the umbilicus. The circumference of the abdomen was forty-five inches at the umbilicus. On vaginal examination some brown discharge was seen about the external genitals, and the speculum showed some sanious discharge in the external os, but the cervix appeared normal. The sound passed three inches. Tilting the uterus with the finger gave one the impression that it did not form part of the large tumour in the abdomen.

On January 23rd the patient was anæsthetised with the A.C.E. mixture, and the abdomen opened by the usual incision. On opening the peritoneum a good deal of free fluid was seen; this was soapy and of thick consistence, and at once suggested that the cyst had burst, as was found to be the case. Most of the loculi of the tumour were small, and the size of the tumour could not be materially reduced by puncturing with the trocar. The incision was therefore extended above the umbilicus, and the tumour removed practically entire. An interesting

point was the presence of recent adhesive peritonitis generally over the intestines, and in fact over whatever part of the peritoneal surface came into view. The patient had had no acute symptoms before the operation, no rise of temperature, no acceleration of pulse, vomiting, or more pain than the mere distention of the abdomen would sufficiently account for, and she walked to the operation table. Yet the cyst had burst, and the extravasated contents had set up the general adhesive inflammation of the peritoneum above described. I had a similar case about five years ago, where, in spite of the rupture, there were no symptoms. In the present case the peritoneum was irrigated with warm saline fluid, and a Keith's tube having been inserted, the rest of the wound was closed in the usual way. The tube was removed on the third day, and the patient made an uninterrupted recovery.

Apart from the bursting of the cyst before operation, an interesting point in the case is the return of a sanguineous vaginal discharge after complete cessation of the catamenia for two years. Generally of course after so long a period of amenorrhoea the reappearance of blood suggests malignant disease of the uterus; but so far as could be ascertained, without dilating the cervix, the uterus in this case was perfectly healthy. It seems certain that the metrorrhagia was to be accounted for by the development of the ovarian tumour exciting a sympathetic congestion of the uterine mucous membrane. I have seen the patient several times since leaving the hospital, and up till the present time (September 1897) she has never had any return of the sanguineous discharge, and she remains quite well.

Pressure symptoms.—As the tumour attains a large size, its pressure on the various organs leads to disturbance of their several functions.

Pressure on the bladder causes frequent and painful micturition, and this arises because the bladder cannot become distended in the natural way; rarely retention of urine may occur. Scanzoni, quoted by Dr. West, records a case where this was due to obstruction of the ureters, so that the retention of urine could not be relieved by a catheter. The tumour in this case was a cystic sarcoma, i.e., a sarcomatous tumour, partly solid, partly cystic; the solid part pressed on the ureters, and obstructed them, so that they became

greatly dilated, one being 2 inches, the other 11 inch in diameter.

I have seen retention of urine occur in a case of small ovarian tumour that had developed downwards and outwards between the layers of the broad ligament, so as to become fixed, although there were no adhesions between it and surrounding parts.\*

Pressure on the rectum may cause obstinate constipation, and even prevent the escape of flatus (West). Occasionally even intestinal obstruction may be produced. When this occurs, it may be a mere pressure symptom, or more probably it may be due to compression or twisting of the gut by the dragging of peritoneal adhesions formed during the growth of the tumour.

Pressure on the stomach causes dyspeptic symptoms, and sometimes vomiting. Wasting necessarily follows the interference with the functions of the stomach.

Pressure on the lungs and heart.—Difficulty of breathing is thus produced, and even asphyxia, as in a case recorded by Dr. Barnes. In that case a young woman with a large ovarian cyst died suddenly, with symptoms of lung distress, in the hospital, while awaiting further treatment. At the post-mortem "the diaphragm was driven up so as to confine the heart and lungs within the narrowest space." Dr. Barnes concluded "that under the impetus of some excitement or exertion, the heart and lungs were suddenly taxed beyond their feeble powers of adaptation, and that thus asphyxia was induced."

Pressure on large veins in advanced cases may lead to cedema of the legs; sometimes even thrombosis of the main vein of a limb may occur, as in one of my cases.

In that case cedema of the left lower limb, from the groin downwards, occurred suddenly, with pain in the limb

<sup>\*</sup> The same case is referred to later in this chapter.

and fever, while the patient was in the hospital awaiting operation. At the same time a number of superficial veins on the left side of the abdominal wall became dilated, that had not previously been so, and could be traced as communicating with a plexus in the skin over Scarpa's triangle.

Thrombosis in one of the trunk veins had no doubt. occurred. The patient had a moderately high temperature for about a week. The leg was kept well raised; it remained considerably enlarged for some two or three weeks longer, after which it diminished in size, and in six weeks from the time when œdema first occurred it had returned nearly to its previous condition. Measurement, however, showed that it was still a little larger than the right. I then allowed her to get up; and as in a few days she seemed none the worse for doing so, decided to operate. At the operation I was agreeably surprised to find no complication of any kind in the pelvis. The tumour was a moderately large one, about the size of the pregnant uterus at the eighth month. It contained fluid of a dark brown colour, and in consistence like treacle. The patient made a typically good recovery.

Prognosis.—This depends greatly on the nature of the tumour.

Dermoid cysts may remain quiescent for many years. Probably the variety due to dropsical distention of Graafian follicles has little tendency to grow beyond a moderate size, or to destroy life. The proliferating glandular and proliferating papillary ovarian cysts will on the average run on to a fatal termination within three years from the time they first attract attention. Malignant ovarian tumours run a course to be measured by months.

Non-malignant solid tumours (fibroids) of the ovary grow very slowly, and may not endanger life for many years, or perhaps not at all.

Since, however, the determination of the exact variety of ovarian tumour present cannot, as a rule, be made with certainty till after its removal, and since by far the larger number of ovarian tumours tend to destroy life within a fixed time, the rule is that every ovarian tumour ought to be removed.

Diagnosis.—We consider the question of diagnosis either in:—

- (1) The early stages, before the tumour has grown sufficiently to form an abdominal swelling; or in
- (2) The later stages, when the tumour has caused a more or less considerable enlargement of the abdomen.

## I. IN THE EARLY STAGES.

The conditions from which a small ovarian tumour has to be distinguished are:—

Fæcal accumulation (more likely to give rise to mistake if situated to the right of the uterus).

Sub-peritoneal fibroid of uterus.

Dilated Fallopian tube.

Pelvic peritonitis.

Pelvic cellulitis.

Pelvic hæmatocele.

Retro-peritoneal cyst of congenital origin.

Malignant tumours when small.

Extra-uterine fætation.

It may be said at once that in some cases the diagnosis of small ovarian tumours is easily made.

As a rule they are globular, elastic, freely movable, and can be, as it were, chased about the half of the pelvis to which they belong backwards and forwards with great ease, particularly if the patient is under the influence of an anæsthetic.

In such cases we can easily define the uterus as separate from the tumour, there being a distinct interval between them.

The only other tumours that give physical signs at all like these are a sub-peritoneal fibroid with a long pedicle allowing it to have a great range of mobility, and a dilated Fallopian tube; but sub-peritoneal fibroids are characteristically hard, whereas small ovarian cystic tumours are elastic. Moreover, we know that fibroid tumours of the uterus are, as a rule, multiple; there is usually more than one present. If no irregularity of outline could be made out as regards the uterus, this would be of some value in excluding fibroids. Occasionally a dilated Fallopian tube (hydrosalpinx, hæmatosalpinx, pyosalpinx) may be so little adherent as to have as much mobility as a small ovarian cyst; and in such cases absolute diagnosis between them may be impossible prior to operation. As a rule it is only in the case of small ovarian cysts, partly or completely adherent, so that they are either quite fixed, or at least not freely movable, that real difficulty arises.

As regards pelvic peritonitis, or cellulitis, or pelvic hæmatocele, the history will be of great assistance. If we rely on the physical signs alone, there may be much difficulty. It may be said in general terms that the outline of an ovarian tumour, even with a number of adhesions around it, is more definite than the outline in peritonitis, cellulitis, and hæmatocele. When there is encysted serous effusion, or encysted purulent effusion, as a result of peritonitis, cellulitis, or suppurating hæmatocele, we must rely chiefly on the history; but even here the more diffused character of the physical signs will help us: for instance, in the case of a suppurating pelvic peritonitis, the pus being encysted in Douglas's pouch, there will probably be hardness around the uterus occupying the pelvis pretty generally,

except where the pus is actually present, *i.e.*, in this instance, in Douglas's pouch. It must be remembered that "suppurating pelvic peritonitis" may in many, perhaps in most, cases really mean that there is a pyosalpinx with adhesions. Again, in peritonitis, cellulitis, and hæmatocele there will usually be fever, or there will have been fever at some period of the case; whereas in most cases of ovarian tumour there is no fever: even when an ovarian cyst contains fœtid pus there may be little or no fever. Though there may be intervals in cases of suppurating ovarian cysts without fever, periods during which there is more or less fever are sure to be met with, if the case is sufficiently long under observation. (See case of Mrs. G., p. 316).

Dilated tubes.—Hydrosalpinx. Pyosalpinx. Hæmato-salpinx.—These form swellings in the right and left posterior quarters of the pelvis, perhaps encroaching to a greater or less extent on Douglas's pouch. They are more often bilateral than unilateral. As there is always some peritonitis in their neighbourhood, it is usual for the swellings they give rise to, to be more or less fixed; only exceptionally are they found as freely movable as a small ovarian cyst. When there is an elastic, somewhat sausage-shaped swelling felt in the posterior fornix just reaching the middle line, and then a vertical groove, and then another elastic swelling on the other side of the middle line, the swellings being little if at all movable, there is considerable probability that they are dilated Fallopian tubes.

Extra-uterine fœtation.—Here the history, usually at first some interval of amenorrhœa, and then irregular hæmorrhages, perhaps with passing of a decidual cast of the uterus, the pain in one or other iliac region, and some of the sympathetic symptoms of pregnancy, e.g., vomiting,

help us. As regards physical examination, the presence of a tumour in the pelvis, somewhat elastic, usually more or less fixed, and situated in one or other posterior quarter of the pelvis; the enlargement of the uterus with blue discoloration of the vaginal portion and vaginal walls, and arterial pulsation in the vicinity of the tumour, together perhaps with sympathetic changes in the breasts, enable us to arrive at a very probable diagnosis of extra-uterine pregnancy.

Retroflexion of the gravid uterus.—The elastic tumour formed by the body of the uterus might be mistaken for an ovarian cyst; more usually the resemblance is to an extra-uterine fœtation; and to distinguish between these conditions may be difficult, unless the patient is under the influence of an anæsthetic: there will then be usually no difficulty in pushing up the body of the retroflexed uterus and satisfying ourselves that it is the body of the uterus. Besides, the history of the two conditions is different.

# II. DIAGNOSIS IN ADVANCED CASES WHEN THERE IS AN OBVIOUS ENLARGEMENT OF THE ABDOMEN.

The conditions from which a large ovarian tumour has to be distinguished are :—

- 1. Pregnancy.
- 2. Phantom tumours.
- 3. Distended bladder.
- 4. Ascites.
- 5. Fibroids.
- 6. Hæmatocele.
- 7. Encysted serous perimetritis and purulent perimetritis.
- 8. Hydronephrosis and pyonephrosis.
- 9. Hydatids.

- 10. Advanced extra-uterine pregnancy.
- 11. Tubercular peritonitis.
- 12. Malignant disease of the peritoneum.

Pregnancy has over and over again been mistaken for an ovarian tumour. Every obstetric physician in large practice sees examples of this mistake from time to time.

It will be well to mention shortly the physical signs that would usually be present in the case of a large multilocular ovarian tumour (the commonest variety) about the size of the uterus at full term.

Inspection.—We notice that the abdomen is distended, the umbilicus not depressed—either level with the skin, or actually pouched out (if there is much co-existing ascites); on asking the patient to breathe slowly and deeply, if the abdomen is exposed to a good light we can often see the upper border of the tumour more or less distinctly outlined. We may notice that the enlargement of the abdomen is not symmetrical. Irregular projections due to the presence of secondary cysts may be visible at parts of the tumour, if the abdominal wall is thin.

Large subcutaneous veins will be seen at various parts of the abdominal wall, and skin-cracks similar to those met with in advanced pregnancy will usually be present.

The following measurements should be taken:—1. Maximum girth of the abdomen. 2. From the xiphisternal articulation to the umbilicus. 3. From the umbilicus to the upper border of the pubes. 4. From the umbilicus to each anterior superior iliac spine. 5. From the spinal column to the umbilicus on each side, so as to compare the semi-circumferences.

The greatest circumference of the abdomen in ovarian tumours is usually below the umbilicus—two or three inches below it, for example. In ascites, the greatest circumference is at the umbilicus. Again we measure the distance from the xiphisternal articulation to the umbilicus, and also that from the umbilicus to the pubes. Normally the umbilicus is an inch or so nearer to the top of the pubes than to the xiphisternal articulation; in cases of ovarian tumour this relation is often altered, so that these measurements are equal, or the lower measurement may even be greater than the upper, the relation of the measurements thus being inverted. Further, in ovarian tumours the umbilicus is not usually equidistant from each anterior superior iliac spine, as it is normally, or in cases of ascites.

In some cases of ovarian tumour, these measurements may not be in accordance with what has just been said, so that too much importance must not be attached to them in differential diagnosis.

Palpation.—We recognise the presence of a tumour, and are able to define its limits above and laterally, more or less completely—below, we cannot separate it from the pelvis. The surface of the tumour is often felt to be irregular, owing to the projection of secondary cysts. Fluctuation can be obtained over the area occupied by the tumour, or at parts of it. We may be able to feel the movements of the tumour during respiration, if the whole abdomen is not too tense.

Percussion.—The area of dulness occupies the middle region of the abdomen, shading off at the flanks, and at the epigastrium, into resonance. The whole of the hypogastric region is dull. The dulness described is that of any large centrally situated tumour—an ovarian tumour, the pregnant uterus, a distended bladder, or a large fibroid tumour of the uterus. If we define the exact line at which dulness begins from above downwards during quiet respiration, and then ask the patient to take a deep breath and hold it, the displacement downwards of the tumour on inspiration causes the line at which dulness commences

to be displaced downwards, so that percussing in the same place we get a resonant note in place of the dull note obtained before.

Auscultation.—Nothing but gurgling sounds (produced in the intestines) can be heard, as a rule, at any part of the tumour. Sometimes a crackling sound can be heard, which may indicate a localised peritonitis at the spot, but does not necessarily do so. Occasionally a sound having some resemblance to the uterine souffle may be heard over ovarian tumours; it is, however, never so distinct as in cases of uterine tumours.

On local examination.—There may be some bluish discoloration of the vaginal mucous membrane, but not approaching in degree that met with in advanced pregnancy. The vaginal portion of the cervix is neither softened nor enlarged, and is not of a particularly blue colour. It is commonly quite as easy to reach it as usual, sometimes even more easy to reach it, owing to the uterus being pushed downwards. Exceptionally, owing to adhesions between the tumour and uterus, the latter is so drawn up that it may be impossible to reach the os uteri with the finger at all.

It is not uncommon for the uterus to be a little to the right or left of the middle line.

Often nothing of the tumour can be felt on vaginal examination; sometimes lumps of perhaps the size of a walnut can be felt behind the uterus in Douglas's pouch. Supposing the rectum to have been emptied, these are, in all probability, secondary projections from the surface of the tumour, occupying Douglas's pouch, and they may not improbably be found adherent there at the subsequent operation.

The sound passes the normal distance ( $2\frac{1}{2}$  to 3 inches); if much more, it is usually in cases where the cervix does not project into the vagina, the whole uterus being displaced upwards, and to some degree elongated by stretching.

It is often possible in cases of ovarian tumour to retrovert the uterus with the sound, and with the finger in the rectum to define the body of the uterus as quite distinct from the tumour.

## DIFFERENTIAL DIAGNOSIS.

Pregnancy.—What has been said on this subject at page 190, on the differential diagnosis of fibroids, applies equally here, and the reader is referred to it to avoid repetition. In addition, we may say that in advanced pregnancy (as we are now considering only the case of large abdominal tumours) it is easy to recognise the parts of the fœtus, the head particularly; and if the child is alive, we shall be sure, on repeated examination at all events, to hear the fœtal heart-sounds, and most probably feel the movements of the fœtus. Further, we can on palpating in a leisurely manner over the whole of the tumour recognise that it alternately hardens and softens.

Pregnancy has not very uncommonly been mistaken for ovarian tumour, and occasionally an ovarian tumour is mistaken for pregnancy. I saw a case of ovarian tumour (there happened to be menorrhagia) where the case was supposed to be one of pregnancy with placenta prævia. Attempts, necessarily unsuccessful, had been made to induce labour. Almost always such mistakes might be easily avoided if only all the methods of physical examination at our disposal were employed as a matter of routine in every case where an abdominal tumour is present.

There are two cases of real difficulty in connection with the diagnosis of pregnancy, and these are not common. They are:—

- 1. Pregnancy with hydramnios.
- 2. Pregnancy complicated with an ovarian tumour.

I. Pregnancy with hydramnios.—The fact of pregnancy will be ascertained by the history of amenorrhoea, and other symptoms of pregnancy; and, as regards physical signs, by the condition of the breasts, and the softening of the cervix.

The alternate hardening and softening that occurs in the walls of the pregnant uterus will be recognised as affecting the whole surface of the tumour. Galabin says he has known cases of pregnancy with hydramnios tapped on the supposition that they were cases of ovarian tumour.

2. Pregnancy complicated with an ovarian tumour.—In early pregnancy there is a danger of the ovarian tumour alone being recognised. We must rely for diagnosis partly on the history, and, if the ovarian tumour is not very large, it may be possible to carry out the bimanual examination, and recognise the enlargement of the body of the uterus. The condition of the cervix should also be noticed. In advanced cases there will be the usual symptoms and physical signs of pregnancy to guide us, and we may recognise that alternate hardening and softening only occur in part of the swelling—that, of course, corresponding to the uterus.

Phantom tumours.—By this expression we mean enlargement of the abdomen, due (1) to the presence of flatus, or (2) to a large quantity of fat in the abdominal wall, or (3) to contraction of the muscles, so as to produce prominence of the abdomen. All these conditions are often present together. The history is not of much value, as the patient is often firmly persuaded she is pregnant, or that she has a tumour. Often by getting her to keep her mouth open, and breathe slowly and deeply, while we palpate the abdomen, we can sink the hands deeper and deeper at each expiration till we come down on the vertebræ.

Bimanually we can make out there is no tumour between

the fingers, and that the uterus is not enlarged. Phantom tumours and pregnancy are, I think, the conditions which most often lead to mistakes: such mistakes might always be avoided by careful examination.

In cases where there is any uncertainty, the administration of an anæsthetic will always enable us to come to a definite conclusion.

I have known one case where a wide separation of the recti, allowing a large hernial protrusion of the intestines through the interval, was mistaken for an ovarian tumour.

Distended bladder.—In all cases of abdominal tumour a catheter should be passed, when, if it be the distended bladder, it will of course disappear as soon as all the urine has been drawn off.

Ascites.—The circumference of the abdomen is usually greatest at the level of the umbilicus in ascites. The umbilicus is equidistant from each anterior superior iliac spine, and maintains its normal position as regards the pubes and xiphisternal articulation, being about an inch nearer the former than the latter; no tumour with a definite outline can be either seen or felt. The flanks are somewhat bulged out, and the front of the abdomen somewhat flattened as the patient lies on her back.

In ascites, when the distention is only moderate, a distinct tremor of the surface is often visible, due to oscillations of the fluid during respiration.

The front of the abdomen in ordinary cases is resonant, while the flanks are dull. If, however, the colon on either side is much distended with flatus, the note will be more or less resonant on that side. If there is a short mesentery, the intestines may not be able to float upwards, and the front of the abdomen may then be dull. This may also happen if the intestines are adherent, or the distention extreme. If the patient is turned on one side, the upper-

most flank, previously dull, becomes resonant. If the fluid is enclosed by adhesions, diagnosis will be very difficult.

Fibroids.—The diagnosis between fibroid tumours and ovarian tumours has been fully considered at page 195, to which the reader is referred.

Hæmatocele.—I have seen several cases where pelvic hæmatocele has been mistaken for a solid ovarian tumour. The history of sudden onset of the symptoms, characteristic of hæmatocele, will usually prevent mistake. Then, again, the tumour in hæmatocele is firmly fixed; whereas the solid ovarian tumour, unless it is malignant, is more likely to be movable: even when malignant, it may be movable at an early stage. Malignant ovarian tumours are usually attended by severe pelvic pain, with rapid loss of flesh, and ascites is not long in making its appearance. Whereas in hæmatocele, when the acutest stage is over, and the patient at rest in bed, pain is not usually severe; further, wasting and ascites do not occur.

When doubt arises, careful enquiry should be made for a history of extra-uterine fœtation (see p. 356), as almost all cases of hæmatocele are produced by rupture of a tubal gestation-sac.

Encysted serous or purulent perimetritis.—The history will help us to distinguish these conditions from ovarian tumours; the illness will date from some of the well-known causes of pelvic inflammation, e.g., labour or abortion; there will be the history of pelvic pain, perhaps rigors and vomiting; there will perhaps be some fever present at the time the case comes under observation. As regards physical signs the tumour formed in either of the cases under consideration rarely has the defined outline usual in cases of ovarian tumour, and it is usually much more fixed than ordinary cases of ovarian tumour. Again, on vaginal examination there will usually be clear evidence

of inflammatory exudation in the pelvis round the uterus, and this organ will be more or less fixed. In cases of ovarian tumour nothing abnormal is necessarily felt round the uterus; and though its mobility may be restricted by the pressure of the tumour above it, the uterus is usually not anything like fixed. Occasionally, if we consider the case only on its physical signs, a centrally situated encysted collection of pus, due to perimetritis, may simulate an ovarian tumour; but the history and the presence of well-marked fever will help us to avoid error. In some cases it will be impossible before operation to distinguish between the cases here under consideration and cases of pyosalpinx, or of suppurating ovarian tumour.

Hydronephrosis and pyonephrosis.—There will be a history of a tumour growing from above downwards; there will usually be disturbances pointing to kidney mischief, frequency of micturition, blood, pus, or albumen in the urine; and unless the tumour is large, we may expect to be able to separate it from the pelvis, i.e., on palpation get the hand under it. We shall usually be able to trace the colon resonance across the tumour; and no line of resonance will be found between the tumour and the spinal column on the side from which the tumour grows. cystic renal tumours the outline is regular, uniformly smooth and convex. On the other hand, an ovarian tumour is first noticed below, and has extended upwards; the intestines are behind it; there is usually a line of resonance in each flank between the tumour and the spinal column, unless the tumour is very large; again, the outline of the tumour is often irregular; while, of course, the urine is, so far as the ovarian tumour is concerned, normal; and menstruation is, as a rule, disturbed.

In cases of hydronephrosis, if the case is some time under observation, the size of the tumour may sometimes be found to vary, the tumour diminishing, or indeed at times almost vanishing, owing to intermittent emptying of its contents into the ureter.

Hydatids only rarely occur in the pelvis; usually there will be a history of the swelling having spread from above down. When they do occur in the pelvis, they are almost certain to be mistaken for some more common pelvic tumour, such as a small ovarian cyst. Tapping, which would probably enable us to make the diagnosis, is undesirable for other reasons.

Advanced extra-uterine pregnancy.—Here there will be the history, differing in all probability from that of ordinary pregnancy. The exact symptoms vary a good deal in different cases; but periods of amenorrhœa, broken by irregular hæmorrhages, pain in the lower part of the abdomen, perhaps expulsion of a decidua, are symptoms to be expected at some time. As regards physical signs in advanced cases, the parts of the fœtus can be felt with unusual distinctness through the abdominal wall; and if the fœtus is alive, we shall hear the fœtal heart. In actual practice early cases of extra-uterine pregnancy are most likely to be overlooked, or mistaken for pelvic cellulitis, or peritonitis; while advanced cases have to be diagnosed from ordinary pregnancy. The only ovarian tumour at all resembling advanced extra-uterine pregnancy is a large dermoid cyst with bony projections and masses at various parts of it, producing some resemblance to the projections of fœtal limbs, or the fœtal skull. With care it will usually be easy to make out that the resemblance is only superficial, that the projections met with in the dermoid cyst do not correspond either in number or position with those caused by the presence of a fœtus.

Enlargement of the spleen.—The edge of the spleen looking to the right and upwards, and having a notch in it,

and the history of the tumour having grown from above down, will usually serve to distinguish this from an ovarian tumour.

The tumour is found to have a sharp edge with a notch in it, this edge looking partly upwards, owing to some rotation having occurred as the spleen enlarged. The history will be of a tumour beginning above and spreading downwards. In most cases diagnosis is easy. Sometimes the error is in taking an ovarian tumour for a splenic tumour. I remember a case of this, where, at a special hospital, the tumour had been thought to be an enlarged spleen. At the London Hospital the tumour was thought to be ovarian, and at the operation was found to be so.

Tubercular peritonitis—Malignant disease of the peritoneum.—Both these conditions have not rarely been mistaken for ovarian tumour, and the diagnosis may be very difficult, and indeed is sometimes impossible, prior to exploratory operation. My object here is only to call attention to points that are practically useful in diagnosis, rather than to enumerate differences that—on paper—make the diagnosis appear easy.

As regards tubercular peritonitis.—There is often a considerable quantity of free fluid in the peritoneum: this, however, is by no means very rare in cases of ovarian tumour. Through the fluid it may be possible to feel solid masses; and a point of practical importance is that these masses are much more irregularly diffused than is the case generally with the irregular projections formed by the development of secondary cysts in the case of the ordinary ovarian tumour. On vaginal examination hard fixed swellings, or a general hardness of the vaginal fornices, is what may be expected in cases of tubercular peritonitis. The indefinite character of the hard deposits felt on vaginal examination in cases of tubercular peritonitis is a point of

some importance. In large ovarian tumours often very little positive information is obtained on vaginal examination, except as to the exact length of the uterus.

Similarly, in cases of malignant disease of the peritoneum, the rapid formation of a large quantity of free fluid, with wasting and cachexia, is what may be expected. The masses felt are also often irregularly distributed, with wider intervals between them than would be the case if the masses were integral parts of an ovarian tumour or projections from it. There is not generally fever in cases of malignant disease of the peritoneum; while in tubercular peritonitis, and in cases of ovarian tumour, if the tumour has become inflamed, fever is common.

Also on vaginal examination in the cases under consideration hard fixed masses are felt irregularly distributed, or lining the whole of the vaginal roof, and also having the indefinite outline which has already been referred to as characteristic in cases of tubercular peritonitis.

It will be seen that generally tubercular peritonitis and malignant disease of the peritoneum resemble one another more than they do the general run of ovarian tumours.

As regards exploratory laparotomy—in cases of tuber-cular peritonitis merely opening the abdomen—letting out the fluid and closing the wound are often followed by a great improvement in the condition of the patient. In cases of malignant disease of the peritoneum, on the other hand, exploratory laparotomy is by no means an innocuous proceeding. There is considerable risk that a low form of peritonitis may follow the operation, to which the patient will succumb. It appears as if the mere exposure of the peritoneum infiltrated with the malignant masses to the air was enough to lead to a fatal result. The healthy peritoneum easily disposes of any organisms, or of blood or exudations following an exploratory incision; but the

unhealthy peritoneum infiltrated with new growth often appears to be unequal to this task, so that peritonitis follows, and the patient dies. Whenever it is possible, therefore, it is highly desirable to distinguish tubercular peritonitis from malignant disease of the peritoneum. In the former case laparotomy is often very good treatment, whereas in the latter it is necessarily useless, and often fatal.

In malignant disease of the peritoneum the growth has no doubt often started in the ovaries—the order of events being malignant degeneration of the ovaries, with subsequent diffused infection over the peritoneum generally.

The following are notes of a case of tubercular peritonitis treated by laparotomy. Before the operation the diagnosis was uncertain, and lay between (1) an ovarian tumour, (2) tubercular peritonitis, and (3) malignant disease of the peritoneum:—

H. C., aged 33; married two years; has had two children, the last three months ago. Suckled the child till two weeks ago. Was admitted into the London Hospital under my care on August 6th, 1897. Her general health has been good, and there is no tubercular history.

Present illness.—Abdomen has been swelling for three weeks. Severe sharp pain in abdomen and back. Vomiting almost every day. No appetite. Constipation. Painful micturition. No cedema of legs.

Note on admission.—No apparent enlargement of liver. Heart nil. Tongue swollen, "beefy," tremulous, and sore. Patient looks very ill, and is wasted.

Abdominal examination.—Abdomen is prominent generally. Umbilicus not depressed. Circumference at umbilicus, 37 inches; circumference 2 inches below umbilicus, 36½ inches; circumference 2 inches above umbilicus, 36 inches; umbilicus to pubes, 6 inches; umbilicus to xiphisternal joint, 8¾ inches. On palpation there is evidently a large quantity of fluid in the abdomen. Resonance in both flanks. Tip of xiphoid cartilage can be felt. Area corresponding to central part of abdomen, starting from umbilicus, with a radius of several inches, is dull. No outline of any tumour can be felt. Free fluctuation from side to side. Skin of abdomen slightly cedematous.

Vaginal examination. - Labia majora are superficially inflamed. Os

uteri not very patulous. Cervix rather closer to pubes than normally. A hard mass felt behind and to the left, apparently fixed, or nearly so. Right side nearly free. Sound enters with curve forward 3 inches. Examination causes great pain.

Per rectum.—The mass is evidently outside and in front of rectum.

There is a small amount of fæcal accumulation.

Temperature.—Between 98.5° and 100° for first six days; between 100° and 101.5° for next six days.

Occasional vomiting.

Operation on August 16th.—Laparotomy wound of 5 inches made. Peritoneum found to be adherent to abdominal wall. Peritoneum incised. About two gallons of clear fluid escaped. Both parietal and visceral peritoneum were found to be studded with miliary tubercle. A mass about the size of a large cocoanut was seen rising out of the pelvis, apparently cystic. There were solid cake-like masses separate from the pelvis. The wound was then sewn up, no washing out of the abdominal cavity being done.

The patient stood the operation well, there being no vomiting after fifteen or sixteen hours. The temperature for two and a half weeks after the operation ranged between 98.5° and 100°, and after this was normal. The vomiting became much less frequent, the pulse-rate diminished, and the patient had much less pain, and began to put on flesh. She left the hospital four weeks after the operation much improved. A large mass could still be felt in the lower abdomen, but there was no evidence of any fresh collection of fluid.

Treatment.—At the present time it is generally recognised that an ovarian tumour should be removed as soon as a diagnosis has been made. The natural history of these tumours is known; the large majority will run on to a fatal termination in three years, and they are liable to many accidents, which may place the patient in immediate peril at any moment. Tapping is undesirable (apart from the risk of setting up inflammation in the tumour, which, if nothing worse happens, will probably render its subsequent removal much more difficult by causing the tumour to become more adherent to surrounding parts), because if the tumour be tapped, and happen to contain papillary

growths, some of the cells are very likely to become detached from these, and escaping into the general peritoneal cavity, set up similar growths there. No one should, under ordinary conditions, perform ovariotomy unless circumstances render it probable that he will have the opportunity of performing the operation a large number of times, since the mortality varies with the experience of the operator. A perfectly uncomplicated ovariotomy is an easy enough operation. It is, however, quite impossible to be sure beforehand that a particular case is a simple one. Unexpected complications and accidents are frequently met with, and can only be adequately dealt with by those whose experience in abdominal work has been considerable.

### OVARIOTOMY.

Preliminary considerations.—The room for the operation.—As regards the room, it is to be remembered that there will be two persons constantly in it, subsequent to the operation—the patient and the nurse—and that, according to the rules of hygiene, each of them requires at least 3000 cubic feet of air per hour. The air can only be changed three or four times an hour without draughts. There should be therefore 2000 cubic feet of space at least in the room; this will suffice if the ventilation is good. A room fourteen feet by twelve feet, and twelve feet high, contains roughly about the cubic space mentioned. There should be a fireplace in the room, and a very efficient system of ventilation is by means of a large Tobin's tube. A good light is of course indispensable.

If the size of the room will allow of it, the bed and the operation table should both be in the room at the time of the operation; if the room is too small to allow of this conveniently, the bed may be brought in afterwards, the

patient, well covered with blankets, in the meantime remaining on the operation table. This is the system followed in the room which I have for ovariotomy at the London Hospital—the room being a small one.

Assistants.—Two are needed, one to give the anæsthetic, and one to stand opposite the operator, and assist him with the operation. A third assistant is often useful. It is important that the anæsthetist should have had experience in abdominal cases; for if vomiting or coughing occur during the operation, the risk of the intestines coming out (and therefore, of course, the danger of the operation) is considerably increased. I prefer to have the patient anæsthetised with the A.C.E. mixture. Only one nurse is absolutely necessary; where two can be had as easily as one, it is better to have two. The nurse also should have had previous experience in abdominal cases.

Preparation of the patient.—She should be confined to bed for a few days previous to the operation, and care should be taken that the bowels are acting regularly. Two nights before the operation a purgative is given, such as that recommended on page 121.

Ten grains of the pil. col. e. hyoscyami will do as well. On the morning of the operation a copious enema of soap and water should be given.

The patient should have long warm stockings, flannel drawers, and a flannel vest with long sleeves, also a flannel jacket to wear over her nightdress, the object of this being to avoid any unnecessary chilling during the operation. In cold weather I like the patient to wear also a "Gamgee jacket" round her chest. The night before the operation the patient should have a hot bath, and wash thoroughly with carbolic soap, particularly the surface of the abdomen. If the operation is fixed for ten o'clock in the morning, the patient should have a cup of strong beef-tea at six.

It used to be thought that ovariotomy should not be performed during a menstrual period; this is now known to make little, if any, matter. In two of my cases, for instance, the patient was menstruating at the time of the




FIG. 134.—BLUNT-POINTED SCISSORS BENT ON THE FLAT.

operation. Both patients made good recoveries. In the first of these I was not told till afterwards that menstruation had begun. In the second case it appeared probable that mental emotion had something to do with the occurrence




FIG. 135.—Wells's small Pressure Forceps.

of the flow; the patient had previously had the operation postponed on account of menstruation, when she took scarlet fever, and was away some ten weeks at the Fever Hospital. During that time she "saw nothing"; two days before the operation the menstrual flow began.

The sponges.—Twelve sponges are sufficient, three should

be small, and there should be one large flat sponge: the remaining eight are of the ordinary size.

Some prefer twenty sponges. The greater the total number used, however, the more the chance of the nurse making some mistake in counting them. The nurse must be specially informed of the importance of being absolutely certain as to the number of sponges she has. On no pretence whatever should any other sponge be brought into the operation room, nor should any sponge be torn in two,




FIG. 136.—Wells's large Pressure Forcers. Sometimes it is an advantage to have forceps of the same size and strength made with the terminals seen in the next figure.


nor thrown away during the operation. No matter how good the nurse, I prefer to count the sponges before the operation myself, and have them counted over before me previous to closing the wound. The nurse should not allow any spectator to touch the sponges; it is better for the nurse, after wringing out a sponge, to put it in a small basin containing hot 1-40 carbolic lotion, and for the operator, or his principal assistant, to take the sponge when wanted direct out of this basin.

Artificial sponges.—For some time past I have given up using sponges in abdominal cases. I use Gamgee pads, cut

of various sizes, and stitched round to avoid the cottonwool escaping from between the layers of the netting. These pads are boiled before the operation. All that has been said as regards the number of sponges, etc., applies to the pads used in their place. New pads are of course used for each operation.

List of instruments, etc., needed for ovariotomy.—In an operation like ovariotomy, for which so many things are required, the only way to avoid omissions is to have a list of everything needed, and prepare for the operation in accordance with the list.

The ultimate responsibility for everything being right rests



with the operator; but as, from the very circumstances of the case, many matters of the greatest importance, both before and during the operation, have to be attended to by others, it may be well said that a successful ovariotomy is a triumph of *organisation* quite as much as of *operation*.

The anæsthetic.

Steriliser (see p. 43).

Porcelain trays (4).—They should be sufficiently deep to allow the instruments to lie well covered with carbolic lotion without being too full.

Sponges (12).—See paragraph on sponges, supra.

Waterproof sheet with oval opening.—On one side for about two inches round the opening the sheet is coated

with plaster. Just before the operation the side spread with plaster is held to the fire, and then made to adhere to the surface of the abdomen. The object of having this sheet is merely to keep the patient dry—fluid from the cyst, etc., that would otherwise wet her clothes, running off on the waterproof. Many operators have given up this waterproof sheet.

Footbath or pail to catch the fluid.

Two glass kidney-shaped receivers, useful for catching fluid. Sterilised towels (4 at least).—To be wrung out of hot 1-40 carbolic, and placed over the waterproof sheet—one above and one below. As these get soiled during the operation they are changed for clean ones.

Scalpel.—This should not be too small.

Probe-pointed bistoury (which some operators use for enlarging the incision upwards; scissors do equally well for the purpose, should this be required).

Scissors (blunt-pointed).—One straight pair, one pair bent on the flat.

A pair of large blunt hooks, the hook forming a segment of a circle the size of a penny.

Dissecting forceps.

A pair of longer forceps with teeth.—Useful for pinching up the layers of the abdominal wall to be divided after the skin and fat have been cut through.

Bull-dog artery forceps.

Twelve pairs of Spencer Wells's small pressure forceps.— It is convenient to have six pairs of straight forceps, and six bent on the flat.

Three pairs of small T-shaped Wells's forceps are sometimes useful.

Six pairs of Spencer Wells's large pressure forceps.—Two of the largest straight, two bent on the flat, and two straight pairs of intermediate size.

Pedicle needle for transfixing the pedicle. Some use this made with a sharp point. An instrument with a point that is not actually sharp, and yet not very blunt, is the one

I prefer. The eye should take No. 5 Chinese twist, when wet, easily.

Cyst forceps .- Two pairs.

Wells's trocar, or Thornton's.—Wells's, being larger, is suitable for tumours containing a large quantity of fluid. Thornton's is useful when the quantity of fluid to be evacuated is not great. It is easier with the latter to prevent escape of the fluid by the side of the trocar.

Volsellæ (2).

Two strong retractors.

Twelve pairs of straight needles for passing the deep sutures through the abdominal wall.

It is convenient to have two sets of these straight needles—a shorter set for thin patients, and a longer set for cases where the abdominal wall is very thick that will take a thicker silk.


Twelve pieces of silk about eighteen inches long (Spencer Wells) are cut, and threaded at each end on one of the needles.

Silkworm-gut sutures for the abdominal wall are now used by most operators instead of silk.

Needle-holder and three small curved

needles for passing superficial sutures, if any are needed.

Hagedorn's needle-holder, and needles of various sizes, to be used with it, may be added to the list.






FIG. 139.—SPENCER WELLS'S TROCAR.

A few fine straight needles for suturing intestine, etc.
Aneurism needle.
Bladder sound.
Uterine sound.

Two long peritoneal sponge-holders, similar in pattern to that figured on page 39, but a little longer.

Curved needle in handle.

Carbolised silk (Chinese twist, or braided silk—the latter is very strong).\*

All the silk to be used should be boiled in a solution of carbolic acid (1-40) the day before the operation for a quarter of an hour. It should then be wound on glass reels, and left in 1-20 carbolic lotion till required. It is of course essential that the hands of whoever is going to wind off the silk should be most carefully rendered aseptic.

Two mackintosh aprons are required, one for the operator, the other for his assistant. They should be long enough to reach from the neck to the feet, and must be thoroughly washed after each operation.

A laryngoscopic mirror and lamp should be at hand for throwing light into the pelvis if necessary.

Two ordinary mackintoshes.

Lister's dressings.

Two packets of iodoform gauze (20 per cent.).

A flannel binder to fasten over the dressing. It is well to have lint sewn at the part where the binder comes in contact with the patient's back.

Strips of flannel bandage to pass round each thigh, and pin to the lower edge of the abdominal binder, so as to keep it in place. These strips should be changed after the operation by the nurse, as often as they become dirty.

Six basins, besides those used for washing the hands-two

<sup>\*</sup> The following sizes should be prepared:—Braided silk, No. 10 and No. 12. Chinese twist, No. 0, 2, 3, 4: No. 0 for superficial sutures, for fine ligatures, or for suturing intestine; No. 2 or 3 for the abdominal wall, or for tying adhesions; No. 4 for thin pedicles. No. 10 or 12 braided silk for stouter pedicles.

for the nurse to use in washing the sponges, one smaller for handing the sponges as described above, one placed conveniently for the operator, that he may rinse the hands from time to time during the operation, one containing plain water for rinsing off the soap after washing the hands, and one containing 1-1000 perchloride and a nail-brush for the final disinfection of the hands.

At the London Hospital I have a large tin vessel holding two or three gallons of a 1-40 carbolic acid solution, heated by a large spirit lamp underneath, to ensure a plentiful supply of hot water throughout the operation, without the nurse having to obtain it from outside.

Three new wooden nail-brushes.—Two for washing the hands and nails, and one to be kept free from soap and used with the perchloride solution.

A catheter.

An injection bottle, in case it is requisite to give brandy per rectum.

Razor and sublimate glycerine (1-1000) for shaving the pubes.

Hypodermic syringe for giving morphia, if necessary, after the operation.

Trendlenberg's operating table, or one on a similar principle. This is useful in case it be desirable to get a good view of the deeper parts of the pelvis. The patient's head is towards the window. With Trendlenberg's table there is a hinge joint nearer the foot than the head, which enables the patient's pelvis to be raised to the desired extent till a good view is obtained.

Certain other instruments must be at hand in reserve.

1. In case drainage may be necessary:—

Keith's glass drainage-tubes (different sizes).

India-rubber sheeting for the same.

Glass tube about 12 inches long, rather narrow, for drawing up fluid from the drainage-tube. This is better than sucking up the fluid with a syringe.

2. In case it may be necessary to fix the pedicle in the lower

angle of the wound :-

The clamp, with wires, and pincers, for adjusting the wire.

Transfixion pins with caps.

Solid perchloride of iron for applying to the stump.

3. Morris's kidney retractors.

4. Two quart jugs for use in washing out peritoneum, or a glass irrigation apparatus with terminal glass tube.

5. Transfusion apparatus.

Lotions, etc., necessary :-

Carbolic (1-20).—Four Winchester quarts at least ready mixed.

Tincture of iodine (3x.).

Sublimate lotion (1-500).—Two Winchester quarts at least; or the tabloids for making 1-1000.

Sublimate glycerine (3ij. of 1-1000).

Methylated spirit for the steriliser.

Hot water.—An unlimited supply.

Solution of sodium chloride (3ij. to 3j.).—3viij. should be at hand. This is boiled the day before, and what evaporates made up with boiled water.

A gallon of boiled water should be prepared for the operation. If the peritoneum is to be washed out, 3j. of the sterilised salt solution is added to a quart of the boiled water—this is then the right strength (3j. of sodium chloride to the pint). A convenient plan is to add 3j. of the salt solution to a pint of cold boiled water, and then make it up to two pints, partly with boiling water from the kettle, and partly with more cold boiled water, till it is of the desired temperature.

## THE OPERATION.

Before the patient is brought into the room all the instruments are covered with a towel.

The nurse should see that the patient passes her water immediately before she is brought in. It is well to have a catheter passed.

No spectators should be allowed in till the patient is under the influence of the anæsthetic.

As the patient lies on the table her clothes are drawn up round the waist, so as to be out of the way; the legs are then tightly wrapped in a blanket, arranged so that it can be tucked in below the feet.

As soon as the patient is unconscious the pubic hair is shaved off, so far as it seems likely to be in the way. When she is not nervous, it saves time to have this done the night before by the nurse. An ordinary mackintosh is now placed over the blanket covering the lower extremities.

The operator and his assistant take off their coats, and turn up their shirt-sleeves, well above the elbow. It is convenient to put on a white linen coat, and roll up its sleeves well above the elbow. The mackintosh apron is then put on. They now wash their hands and arms thoroughly, paying special attention to the nails (the water should contain an excess of tincture of iodine, enough to make the water brown in spite of the soap); and after washing rinse off the soap in plain water. Then the hands and arms are to be scrubbed for two minutes in 1-1000 perchloride lotion with a nail-brush not used The mackintosh with the oval opening, which for soap. has been in the meantime held to the fire by the nurse to melt the plaster, is then carefully adapted to the surface of the abdomen, so that the aperture is symmetrically situated with regard to the middle line from the umbilicus

to the pubes. The lower end of the opening in the mackintosh should be at the upper margin of the pubes. Many operators no longer use this mackintosh.

The operator now sponges the skin with 1-1000 perchloride lotion, and also the mackintosh for a few inches round in each direction. The sterilised towels are now wrung out of hot 1-40 carbolic, and placed over the mackintosh, one above and one below.

The incision.—The operator now takes a scapel and makes the incision.

This should be in the middle line, and should begin about one inch or an inch and a half below the umbilicus, and not extend lower than to within two inches of the pubes. As soon as the skin and fat are divided, the assistant should press a sponge firmly into the wound for a few seconds; the operator then fixes Wells's pressure forceps on any bleeding points. Unless the recti should have been separated, the sheath of one or other rectus must be opened, as there is no linea alba below the umbilicus (Treves). The operator pinches up the white aponeurosis in view with the toothed forceps, and makes an opening in it with the knife held flat. The aponeurosis is then divided to the full extent of the wound by means of scissors bent on the flat. The recti are not to be cut; the operator must look for the interval between them, and divide the tendinous structures found there by pinching these up, and cutting towards himself, with the knife held flat, till the yellow sub-peritoneal fat comes into view. Spencer Wells's forceps are then put on any points that may be bleeding, and the wound sponged clean. The sub-peritoneal fat and peritoneum are then pinched up and divided, with the knife held flat as before. As soon as the opening made in the peritoneum is large enough to admit the finger, the fore-finger of the left hand is introduced, and the peritoneum opened to the

extent of the wound by means of scissors bent on the flat, passed along the finger.

The surface of the cyst, which usually has a greyish-blue colour, is now seen. The wound should be large enough to allow the operator to introduce his hand and part of his arm into the peritoneal cavity, so that he may thoroughly explore the relations of the tumour.

If the cyst is small, so that it can be removed without puncturing it, either through the existing wound, or by only enlarging this a little, it is better to remove it entire, as it is impossible to be sure beforehand that the contents may not be feetid pus. As a rule, however, the next step of the operation is puncturing the cyst.

The assistant steadies the cyst by placing his hands one on each side of the abdomen. The operator then takes the trocar and plunges it into the cyst, using no more force than is necessary. As soon as the trocar is in the cyst cavity, the canula must be projected beyond the trocar. As a rule the claws on Wells's trocar do not afford a very satisfactory hold, so that it is better to seize the cyst wall near the point of puncture with a volsella. Great care should be taken to avoid allowing any of the contents of the cyst to escape into the peritoneal cavity. If the tumour is composed chiefly of a single cyst, it will soon have collapsed sufficiently to allow of its being drawn out of the wound; if, on the contrary, the tumour is made up of several large cysts, besides the one punctured, it will not come out. It will then be necessary either to bring these in turn, so long as the tumour cannot be drawn out, up to the wound, and puncture them; or the puncture first made may be enlarged, so that the operator can pass his hand and arm into the interior of the cyst, and break up the secondary cysts.

If the operator's hands are fouled with the contents of the cyst, he must wash them thoroughly before again using them within the cavity of the peritoneum. As the tumour is gradually drawn out of the wound, its surface is carefully watched with a view to the recognition of adhesions.

Adhesions to the omentum are the least serious. If the omentum is very extensively adherent, it must be tied in sections with carbolised silk. If too large a piece of it is trusted to one ligature, there is great danger of secondary hæmorrhage.

Adhesions to the intestine are much more serious. An attempt must be made to separate the cyst from the intestine by pressing with a sponge, or, better, with the finger. If this does not succeed, the cyst-wall must be divided, and a thin lamella of it left adherent to the intestine.




FIG. 140.—PEDICLE NEEDLE FOR TRANSFIXING THE PEDICLE.

Adhesions deep down in the pelvis also require very careful management.

According to their length and firmness, it will be right either to tie them with carbolised silk, or to split the cystwall, and leave a thin portion of it attached.

Ligature of the pedicle.—The assistant holds up the tumour so that the pedicle is well in view. The operator, holding the pedicle between the thumb and fingers of the left hand, selects the point through which to pass the pedicle needle, specially avoiding any vessel. He then transfixes it with the pedicle needle (Fig. 140), threaded with the silk previously selected for the purpose. The loop of the ligature is seized on the far side of the pedicle, and held while the pedicle needle is withdrawn. There are at least three good methods of proceeding from this stage.

I. The loop is drawn through till it is large enough to allow of its being carried over the whole cyst towards the operator; he then places one of the free ends above the loop, and holding the pedicle with his left fore-finger and thumb, draws the two free ends of the ligature taut, so as to constrict the pedicle. They are then tied as tightly as is possible without breaking the ligature. In making the first tie the one end should be hitched twice over the other.

This mode of managing the ligature—drawing back the loop over the tumour—passing one of the free ends above the loop, drawing the ends tight, and tying, is the one




FIG. 141.—THE STAFFORDSHIRE KNOT.

introduced by Mr. Lawson Tait, and he has named the knot "the Staffordshire knot" (Fig. 141).

It is suitable where the tumour is small—for example, not larger than a cocoanut—and where the pedicle is thin.

Whatever method of ligature is adopted, when the knots are being drawn tight, the assistant must lower the tumour, so that the pedicle may not be on the stretch.

2. Another simple way of ligaturing the pedicle is to transfix as before, leaving a loop on the far side, then to carry one of the free ends round the pedicle, pass it through the loop, draw the two free ends tight round the other side of the pedicle, and tie them together (Figs. 142 and 143). This is an extremely convenient method, and I have used it in a great many cases. This knot also

answers well for tying the broad ligaments in vaginal hysterectomy, and I have referred to it when describing that operation.

It is important while tying the ligature, although aiming at tying it tight, to stop well short of using the degree of




Fig. 142 (Doran).

force that will break it: apart from the waste of time in applying another, such an accident may lead to a tearing of the pedicle, and this may be very troublesome. In one of my early cases the ligature broke: I immediately turned to get another, and proceeded to transfix and tie what




Fig. 143 (Doran).

appeared to be the same pedicle as before. After the ligature was applied there was an area outside it, from which pretty free bleeding occurred; and as there had been no adhesions in the case, I was at first puzzled to account for it. It was really the outer part of the original pedicle which had been torn, or, as it were, had the peritoneum "scalped" off it by the first ligature. As soon as

I made out what had happened I transfixed the broad ligament, and tied the bleeding surface with the Staffordshire knot. The patient made a good recovery.

Some recommend tying a preliminary ligature round the outer part of the pedicle, which contains the ovarian vessels, before applying the main ligature. It certainly appears to be an additional safeguard against secondary hæmorrhage; it cannot be considered as absolutely necessary; but it may be advantageously used when the ovarian vessels are very large, and the whole pedicle is thick and broad.

It is well also to tie a ligature round the whole pedicle, taking care to let the silk lie in the groove already made by the other ligatures. All the ligatures are cut short.

3. Another method of tying the pedicle, and perhaps the one most generally used, is, after transfixing as before, to cut the loop on the far side of the pedicle, and tie the two resulting ligatures one round each half of the pedicle. Before tying, care should be taken to see that the ligatures interlock, otherwise the pedicle may be split.

Two pairs of Wells's small forceps are now fixed one on each side of the pedicle, about three-quarters of an inch beyond (on the tumour side of) the ligature, and the pedicle is cut through just beyond them. These forceps are to enable us to bring the pedicle into view, without disturbing the ligatures, just before the wound is closed.

A sponge fixed in a pair of sponge forceps (somewhat resembling Wells's large pressure forceps) is now guided down along the fingers of the left hand into Douglas's pouch. If there is no bleeding, it is best to get on with the introduction of the sutures into the abdominal wall. A large flat sponge is passed into the wound, so as to keep back the intestines, and soak up the small amount of blood escaping as the sutures are being passed. A large blunt hook is now introduced into the upper and lower angle of

the wound: these hooks are held by the assistant; they facilitate the even introduction of the sutures, and the neat apposition of the edges of the wound. No needle-holder is necessary; the operator begins at the lower end of the wound, pinches up the abdominal wall with the fore-finger and thumb of the left hand, and passes the needle from the peritoneum, about a quarter of an inch from the cut edge, through all the structures of the abdominal wall, muscle included, and brings out the needle on the skin about a quarter of an inch from the cut edge. The needle already threaded on the other end of the suture is then passed similarly through the abdominal wall on the other side. When as many sutures have been passed as seem necessary, the free ends on each side are gathered together, and fixed with a pair of Wells's forceps; the loops of the sutures lying across the wound are then pulled long enough to allow of their being caught and held out of the way by the blunt hooks in the upper and lower angles of the wound. The flat sponge is then removed. If much blood or fluid from the cyst have escaped into the peritoneum, it is best to wash out the peritoneal cavity with sterilised normal salt solution (3j. to Oj.) heated to blood heat. The edges of the wound are held together, and the abdomen is kneaded, so that the fluid may wash the peritoneum thoroughly. The time occupied in the process, and the quantity of fluid to be introduced, will depend entirely on the degree to which the peritoneum has been fouled. For example, if fœtid pus from a suppurating cyst has got into the peritoneum, the washing must be done very thoroughly indeed. Some operators, however, do not wash out even in such cases; they rely entirely on sponging, and consider that washing out rather tends to diffuse the irritating material through the peritoneal cavity generally. No doubt washing out may be easily overdone. Many operators get very good

results who never adopt it. The water is then sponged out—just at the last, the patient's shoulders being raised off the table, to allow all the fluid to flow into the pelvis. The last sponge should come up clean.

The operator should not forget to look at the other ovary; if it should be in a state of cystic degeneration, it also must be removed.

The sponges and forceps must now be carefully counted before closing the wound; it is best that the counting be done under the operator's eye, so that he may feel personally certain that the numbers are correct.

The abdominal sutures are then tied, beginning below, the edges of the skin being carefully adjusted, and the sutures not tied too tightly. If necessary, superficial sutures of fine silk are inserted with a small curved needle held in a needle-holder. If the waterproof sheet has been used, it is now removed, the surface of the abdomen cleaned, and the wound dressed. Several pads of iodoform gauze are placed over and around the wound. Some loose carbolic gauze or Gamgee tissue may be used to fill the hollow produced by removal of the tumour, if it has been a large one. The outside dressing is then put on, and the flannel binder pinned tightly over all. The patient is then put back to bed. There should be a hot bottle for her feet ready in bed.

Special difficulties during the operation.—There may at the outset be some difficulty in ascertaining when the peritoneum has been reached; this is likely to occur when there are dense adhesions between the anterior surface of the tumour and the parietal peritoneum. Under such circumstances it is best to enlarge the wound in an upward direction, so as to endeavour to strike the peritoneal cavity above the adherent part of the tumour.

The tumour may be found to have no pedicle, having

grown between the layers of the broad ligament beneath the pelvic peritoneum. Sometimes after tapping it becomes possible to get a sort of broad pedicle, and tie it in the usual way. This happened to me in one case where the tumour had grown outwards between the layers of the left broad ligament to the wall of the pelvis. There were no adhesions; but before tapping the tumour was perfectly immovable. The patient made a good recovery.

Enucleation.—When no pedicle can be established even after tapping, the opening made by the trocar must be enlarged somewhat, and the cyst shelled out of its capsule. Large vessels must be seized with Wells's forceps and tied. Care must be taken not to lacerate the capsule during this proceeding.

As regards the capsule left after enucleation of the cyst, it may happen:—

- 1. That it can be drawn up, its attachment transfixed, and tied like an ordinary pedicle, and the rest of the capsule removed; or,
- 2. If this cannot be done, the capsule may be drawn up and stitched to the lower part of the abdominal wound, any superabundant portion beyond the stitches being cut off. If the capsule has not been torn in shelling out the cyst, the cavity of the capsule will then be completely shut off from the general peritoneal cavity. A glass drainage-tube is inserted into the cavity of the capsule. The peritoneal cavity is treated in the ordinary way, as already described.
- 3. After enucleating a cyst, even a large one, if all the vessels in the capsule have been secured, and there is nothing but capillary bleeding, the walls of the capsule will sometimes fall together, and need not be drawn up and stitched to the abdominal wall as above described. The abdominal wound may simply be closed in many of these cases without drainage.

On opening the abdomen the tumour may prove to be malignant, and so densely adherent in every direction that removal of it would be impossible. In such a case the wound must be closed. Apart from malignancy, an ordinary ovarian tumour may be so universally adherent that complete removal is impossible. This is very rare when the operator has had much experience. In such a case the opening made by the trocar in the cyst is enlarged, the hand introduced into the cyst cavity, and as much as possible of the solid material in the cyst, if any, removed; the edges of the cyst are then stitched to the abdominal wall, so as to shut off the cyst cavity from the general peritoneal cavity, and a glass drainage-tube inserted.



FIG. 144.—KEITH'S GLASS DRAINAGE-TUBE.

Drainage.—In the account given of an ordinary complete ovariotomy, drainage is not mentioned, because in the large majority of cases it is not required. It is a very difficult matter in certain cases to decide whether to insert a drainage-tube in the lower angle of the wound, or not. Most cases certainly do not require it. Moreover, it adds to the difficulty of the after-treatment, and unless very skilfully managed is an additional source of danger in itself. The more manipulation that has been necessary within the peritoneal cavity, the more likely is it that a drainage-tube ought to be inserted. When, therefore, the operation has been very long and difficult, when a very large number of adhesions have had to be dealt with, and particularly when the separation of adhesions has left a large oozing surface, a glass drainage-tube (Fig. 144) should be inserted in the

lower angle of the wound; it should be passed well down into Douglas's pouch. The exposed end of the tube has a projecting collar; this is passed through a small hole in the centre of a piece of india-rubber sheeting, about a foot square. The hole is so small that the end of the tube can only be passed through when the india-rubber is on the stretch; it thus grasps the tube firmly. A sponge is placed over the end of the tube, and the india-rubber wrapped round the sponge, so as to exclude the air. The upper part of the wound is dressed in the usual way. The sponge has to be changed and wrung out of carbolic lotion (1-40) every four hours for the first twenty-four hours, every six hours for the second twenty-four hours, and after that every twelve hours till it is removed. It is better to empty the drainage-tube at each dressing by passing down it a narrow glass tube, then putting a finger on the upper end of this, and so withdraw the fluid a little at a time till all is removed. This is far better than to suck up the fluid with a syringe. Omentum is more likely to be sucked into the holes at the end of the glass tube if a syringe is used. When the colour of the discharge is no longer red, and the quantity of it very little, the tube may be taken out. When I do use a drainage-tube, I leave it in till the end of the third day.

Drainage by iodoform gauze.—Instead of using a drainagetube a strip of iodoform gauze may be passed down into Douglas's pouch. The upper end is left protruding about an inch from the wound. I have used an iodoform gauze drain in several cases. It is taken out at the end of the third or fourth day; a short strip is then inserted, so that the cavity left after removing the first piece of gauze may fill up from the bottom.

The objections to the use of iodoform gauze are, first, that it often causes a good deal of disturbance, especially

rapidity of pulse, during the first two days; and, secondly, that taking out the gauze is rather painful.

After-treatment.—The patient has nothing given her by the mouth for the first thirty-six hours except pieces of ice to suck, or a teaspoonful of iced soda water. Generally small quantities of hot water seem more grateful to the patient. If the patient is feeble, it is as well to begin giving nourishment per rectum from the beginning. The nutrient enema should not be more than four ounces in quantity; it may consist of two ounces of peptonised beef-tea, one ounce of coffee, and one ounce of brandy. Great care is necessary to avoid (1) giving the enemata too frequently, or (2) persisting with them longer than is absolutely necessary. I have known a very troublesome diarrhœa to be set up in this way. A tube should be passed a little way into the rectum to allow flatus to escape before giving the enema. Hypodermic injections of morphia are given if the patient is in much pain: they are not always necessary. After the first thirty-six hours, if there is no sickness, she may begin to take fluid nourishment by the mouth, commencing with a teaspoonful of a mixture of equal parts of milk and lime water, or of milk and soda water.

It is a great mistake to give large quantities during the first three or four days after an abdominal section.

The stitches are taken out on the eighth day, the wound being well sponged with 1-40 carbolic lotion before doing so. A pad of iodoform gauze is laid on and fixed by broad bands of strapping to support the cicatrix. From neglecting to do this the wound burst open in one of my cases. The patient was the subject of chronic bronchitis. The edges of the wound were drawn together with strapping, and the patient did well. The bursting open of the wound was not attended by any special symptoms, although a coil of intestine could be seen at the bottom of the wound.

This immunity was, I think, due to the fact that the general cavity of the peritoneum was shut off by adhesions, and also to the fact that the case was being treated on rigidly antiseptic principles.

At the end of the week the bowels should be opened by an enema. No solid food should be given till the bowels

have acted.

When the case is going to do badly after an operation, unfavourable symptoms usually set in about the third day.

Rise of temperature, rapidity of the pulse, vomiting, hiccough, and abdominal distention may occur.

The more of these symptoms that are present together,

the worse the prognosis.

Another early symptom, if the case is going to do badly, is a change in the appearance of the patient's face; it acquires a pinched look. The patient may be slow in apprehending what is said to her, and there may be some degree of restlessness.

A high temperature must be treated by applying ice, or ice-cold water, to the head—the former by means of an ice-bag, the latter by means of Thornton's ice-cap. An ice-bag does very well. Brandy must be given according to the state of the pulse.

If all the unfavourable symptoms mentioned are present, very little can be done for the patient, and she is almost certain to die before the eighth day.

In my experience as regards prognosis the most important thing to judge by is the pulse, both as regards its frequency and its quality.

Mr. Lawson Tait writes as follows: "Concerning fatal cases, I am altogether of Dr. Keith's opinion, that the very first search to be made for an explanation should be in the details of the operation." \*

<sup>\*</sup> Diseases of the Ovaries, 4th edit.

When the patient is convalescent, an abdominal belt should be ordered for her before she gets out of bed. She should wear it for a year after the operation. Another point of importance is to warn her to be exceedingly careful to rest, and to avoid any exposure at the menstrual period for some time to come. Neglect of this precaution may lead to very alarming symptoms, high temperature and rapid pulse particularly, which, however, usually pass off in a few days.

Mortality of ovariotomy.—Although long runs of cases are recorded from time to time without any fatality, yet it may be taken that if the rule of operating on every case of ovarian tumour as it comes is followed without any rejection of cases, there will be about five deaths in a series of one hundred consecutive cases, even in the best hands.

#### OVARITIS.

Etiology.—The conditions usually given as causing ovaritis are:—

Labour and abortion.

Gonorrhæa.

Suppression of menstruation by cold.

Operations on the uterus.

Recent marriage.

Certain fevers.

Alcoholism (Matthews Duncan).

On looking through this list, it will be seen that most of these causes are causes of pelvic inflammation (pelvic peritonitis and cellulitis).

It is difficult to imagine what may be called an isolated ovaritis, *i.e.*, an inflammation of the ovary without any implications of the structures in the immediate neighbourhood. Probably in most cases ovaritis is secondary to

pelvic peritonitis or cellulitis. In the case of ovaritis secondary to pelvic peritonitis, the inflammation has spread along the Fallopian tube to the peritoneum, producing pelvic peritonitis, which involves the peritoneal aspect of the ovary, producing adhesions between it and surrounding parts, e.g., the Fallopian tube: this is periovaritis. Just as in pelvic peritonitis the tissue immediately under the peritoneum must participate to some extent in the inflammation, so in periovaritis no doubt the superficial layers of the parenchyma are at the same time to some extent involved. In the case of ovaritis secondary to pelvic cellulitis (interstitial ovaritis), the inflammation has spread along the connective tissue in the fold of the broad ligament forming the mesovarium to the parenchyma of the ovary. Under pelvic cellulitis I have referred to a specimen I had the opportunity of examining, where the inflammation in the broad ligament was in the stage of phlegmon, while that in the adjoining ovary had gone on to abscess. A follicular form of ovaritis is described where the contents of the Graafian follicles become turbid, or purulent, the parenchyma of the ovary being more or less involved at the same time. It is sufficient, however, to speak of periovaritis and interstitial ovaritis.

Ovaritis may be acute or chronic. In the acute form there is more or less enlargement of the ovary. In the chronic form the organ may be enlarged; but, on the other hand, it is said that it may be smaller than normal, the change being of a cirrhotic nature, with destruction of follicles. It is well to remember that healthy ovaries vary considerably in size. The ovaries even in the same body are often different in size; and, again, the same ovary varies in size at different times, enlarging during menstruation and pregnancy.

Except where there is a great deal of fat in the abdominal

wall, the ovaries can be felt by those practised in the bimanual examination, even when the ovaries are of normal size. This can certainly be done with the aid of an anæsthetic, and not rarely even without it in those people who submit well to the examination. We must not conclude, therefore, merely because the ovaries are to be felt, that there is necessarily anything abnormal—any inflammation or degeneration of the ovary.

Symptoms and signs.—In acute cases these are the same as for the pelvic peritonitis or cellulitis, to which the ovaritis is in the large majority of cases, if not in all, secondary.

In chronic cases the patient complains of pain at the affected side; dysmenorrhæa, pain on coitus, sometimes pain on micturition or defæcation. There is sometimes menorrhagia. Whether this is present or not depends on whether there is co-existing endometritis of the body of the uterus or not (Matthews Duncan).

As regards the physical signs, the uterus is probably less movable than normal, owing to the persistent adhesions; and there is a swelling about the size of a walnut in the situation of the ovary, this swelling often being more or less fixed.

Such swellings, when they come to be examined after removal by operation or post-mortem, are often found to consist of the Fallopian tube (more or less dilated, its fimbriated extremity not to be recognised, the tube ending on the surface of the ovary), matted to the ovary by adhesions; and there are also adhesions between the swelling, formed by the tube and ovary, and adjacent parts. Such a condition is not uncommonly bilateral. I have known cases where swellings, believed to be enlarged ovaries, as the result of physical examination, were found to be chiefly due to dilatation of the Fallopian tubes at a subsequent operation.

Treatment.—The treatment of acute ovaritis is that of acute pelvic inflammation, and the treatment of chronic ovaritis is very similar to that for chronic pelvic inflammation, e.g., blistering or painting the area to which pain is referred with iodine; hot vaginal douches; the use of a glycerine plug every night; and regulating the bowels. If there is menorrhagia, ergot, or bromide of potassium in scruple doses, may be tried. Though many cases improve under such treatment, the improvement is often only of a temporary character. This is especially so where the patient is of a markedly neurotic temperament. Apart from this, many cases undoubtedly get well if the course of palliative treatment is sufficiently prolonged. When this is not so, the question of removal of the ovaries and tubes may have to be considered. The patient must thoroughly understand what it is proposed to do-the risks of the operation itself, and the possibility that she may not be cured by it; the sterility necessarily produced (though many such cases are per se sterile, owing to occlusion of the tubes). The operation is a more difficult one than ovariotomy, owing to the dense adhesions often met with, and should only be undertaken by a specialist skilled in abdominal surgery.

# DISPLACEMENTS OF THE OVARY.

It must be remembered that the position of the ovary is constantly varying, according to the position of the uterus. This follows from the intimate connection of the two organs, so that there is no one position entitled to be considered the normal position of the ovary. In reality there are several normal positions for it.

# HERNIA.—PROLAPSE.

Hernia.—The ovary may be found in a hernial sac, usually that of an inguinal hernia. Cases of this kind are usually congenital. The treatment consists of protecting the organ from pressure by a suitable shield. If this cannot be done, and the suffering is considerable, it may be removed by operation.

Prolapse.—The common form of this is where the ovary descends into Douglas's pouch. It does this in marked retroversion or retroflexion of the uterus. In such cases both ovaries can usually be felt in the posterior fornix, one on each side of the swelling produced by the uterus. Again, in cases of procidentia uteri, when the whole uterus lies outside the vulva, the ovaries can often be felt at the sides of the uterus.

The ovary may, however, come to lie in Douglas's pouch if its attachment to the broad ligament becomes stretched, without any displacement of the uterus. Such stretching is likely to occur when the ovary enlarges from any cause, and remains freely movable.

Symptoms.—Pain on coitus is present, and perhaps also pain on defæcation, or there may be more or less constant pain aggravated on such occasions. Menstruation may be irregular. This will depend rather on other changes, e.g., commencing cystic degeneration, or inflammation affecting the ovary, than on its mere prolapse.

Diagnosis.—The swelling, suppose it to be the size of a walnut, is felt in the posterior fornix. It may be movable or fixed, according to the presence or absence of adhesions. It must be distinguished from the body of the retroflexed uterus. This is easily done either by the bimanual examination, feeling the body of the uterus in front, or by passing

the sound. The sickening pain caused by pressure on the ovary also serves to identify it.

Treatment.—If the prolapsed ovary is movable, it may be pushed up, and an effort made to keep it up by inserting




Fig. 145.—Diagram of the Structures in, and adjacent to, the Broad Ligament (Doran).

1. Framework of the parenchyma of the ovary, seat of 1a, simple or glandular, multilocular cyst. 2. Tissue of hilum, with 3, papillomatous cyst. 4. Broad ligament cyst, independent of parovarium and Fallopian tube. 5. A similar cyst in broad ligament above the tube, but not connected with it. 6. A similar cyst developed close to 7, ovarian fimbria of tube. 8. The hydatid of Morgagni. 9. Cyst developed from horizontal tube of parovarium; cysts, 4, 5, 6, 8, and 9 are always lined internally with a simple layer of endothelium. 10. The parovarium; the dotted lines represent the inner portion, always more or less obsolete in the adult. II. A small cyst developed from a vertical tube; cysts that have this origin, or that spring from the obsolete portion, have a lining of cubical or ciliated epithelium, and tend to develop papillomatous growths, as do cysts in 2, tissue of the hilum. 12. The duct of Gartner, often persistent in the adult as a fibrous cord. 13. Track of that duct in the uterine wall; unobliterated portions are, according to Coblenz, the origin of papillomatous cysts in the uterus.

an elastic ring pessary of suitable size, or a Hodge's pessary, preferably one with its upper end made to contain glycerine, so as to produce a sort of soft cushion. If palliative measures fail, we may remove the ovary by operation, if the patient is anxious for this to be done, after the nature of the proposed operation has been explained to her.

Malformations.—It is sufficient to mention that the ovaries may in rare cases be congenitally absent. Sometimes one ovary and the other uterine appendages of the same side are wanting. The corresponding kidney may also be absent. A more frequent anomaly is that the ovaries are present, but are imperfectly developed, and remain so throughout life.

Parovarian cysts.—Fig. 145 shows diagrammatically the relations of the parovarium and neighbouring structures in the broad ligament. A parovarian cyst is formed by distention of one of the tubules of the parovarium. Sometimes parovarian cysts contain papillary growths.

The fluid in a parovarian cyst is thin, and of low specific gravity. It contains a little salt and a trace of albumen.

As these cysts sometimes contain papillomatous growths, they should not be tapped, for fear of affecting the peritoneum. They should be removed in the same way as an ovarian tumour. Indeed it is not generally possible to diagnose a parovarian cyst from an ovarian tumour; but as the treatment is similar, this is a matter of no great importance.

# CHAPTER XVII.

# FLEXIONS AND VERSIONS.

The normal position of the uterus.—The accompanying diagram (Fig. 146) shows what may be taken as the normal position of the uterus when the bladder is empty. The position is one of slight anteflexion.

It must not be forgotten, however, that although it is customary to speak of that position of the uterus as more particularly *the* normal position, there are in reality several normal positions—several other positions which the uterus may occupy that are to be regarded as physiological, not pathological.

For example, as the bladder fills the uterus is pushed backwards as a whole, rotating round an imaginary transverse axis. In technical language, it becomes retroverted. In Fig. 147, B, C, D, indicate the positions successively occupied by the uterus as the bladder becomes more and more distended. When the bladder is emptied, the uterus returns to its original position of slight anteflexion, A.

Flexion of the uterus.—By a flexion of the uterus we mean that the long axis of the body of the uterus makes an angle, more or less obtuse according to the degree of flexion, with the long axis of the cervix. The angle is usually situated at the internal os.

In Fig. 146 the long axis of the body of the uterus meets the long axis of the cervix at about an angle of 120°.

Anteflexion.-When the angle formed by the meeting of




Fig. 146.—Showing the normal shape and position of the virgin uterus when the bladder is empty (Schultze).

these two axes is on the anterior aspect of the uterus, we have anteflexion.




Fig. 147.—B, C, D, positions successively occupied by the uterus as the bladder fills (Van de Warker).

Retroflexion.—When the angle is on the posterior aspect of the uterus, we have a retroflexion.

Version of the uterus.—When the whole uterus is rotated round an imaginary transverse axis, without the position of the cervix and body relative to one another being altered, we have what is called a version of the uterus. If the rotation is forwards, an anteversion; if backwards, a retroversion.

Version and flexion may be combined in the same case. For instance, a uterus may be anteflexed and retroverted at the same time, the axis of the cervix making an angle with the axis of the body on the anterior aspect of the uterus, and then the whole organ being rotated backwards on an imaginary transverse axis.

**Diagnosis.**—Normally the external os is found looking downwards and backwards; and on bimanual examination, if the bladder is empty, the body of the uterus can be grasped between the internal finger and the fingers of the external hand.

In other words, on bimanual examination, when the uterus is in the normal position, the body of it can be felt through the anterior fornix; that is to say, the uterus is anteverted or anteflexed. Conversely, if the body of the uterus cannot be felt through the anterior fornix bimanually, the position of the uterus is not one of anteversion.

Two precautions are necessary to avoid error:-

I. The external hand must make pressure well above the situation where the body of the uterus may be expected to be if the position of the uterus be one of anteversion or anteflexion. The fingers of this hand must not, for example, press downwards close behind the pubes, or they may very probably be altogether in front of the body of the uterus, and meet the internal finger without the body of the uterus being grasped, even though it be anteverted or anteflexed. This mistake is a common one. Another somewhat similar mistake is omitting to make the pressure with the external

fingers in the middle line. In this case also the external and internal fingers may be made to meet without grasping the body of the uterus, although it really lies in front.

2. The internal finger should make its pressure upwards *immediately* in front of the vaginal portion of the cervix, not some distance in front, or it will easily meet the external fingers without the body of the uterus being grasped.

It is well for the beginner to satisfy himself, in an undoubted case of anteversion or anteflexion, how all these mistakes can be made. They are certainly all commonly made by students beginning the study of the subject. The diagnosis may be confirmed by passing the sound, and finding that it enters with the concavity forwards. When the body of the uterus lies to the front, it is usually quite unnecessary to pass the sound for this purpose. The information obtained by the bimanual examination in practised hands is quite conclusive.

If an examination is made with the precautions mentioned, and the body of the uterus cannot be felt through the anterior fornix, but the internal and external fingers meet with only the abdominal wall, the vaginal wall, and the walls of the bladder between them, the body of the uterus does not lie to the front. It must, therefere, be either retroverted (Fig. 151), anteflexed and retroverted (Fig. 150), or retroflexed and retroverted (Figs. 152 and 153).

Retroversion.—When this exists alone (Fig. 151), the physical signs are as follows:—The os uteri looks upwards and forwards; we can feel the body of the uterus through the posterior fornix; and if the abdominal walls are very lax, we may be able to grasp the body of the uterus bimanually, even in this position. If there is no co-existing retroflexion, we find no angle at the point where the axis of the cervix meets the axis of the body of the uterus.

Retroflexion and retroversion.—Instead of the os uteri looking downwards and backwards, as it does normally, we find it looking more forwards than usual, but yet




FIG. 148.—Slight anteflexion—the normal position. a. Posterior aspect; b. Anterior aspect.




Fig. 149.—Anteflexion with slight retroversion.




Fig. 150.—Anteflexion with considerable retroversion.




FIG. 151.—Retroversion without any retroflexion.




FIG. 152.—Retroversion and slight retroflexion.




Fig. 153.—Slight retroversion and marked retroflexion.

Retroversion may occur :-

Alone (Fig. 151), or

Associated with anteflexion (Figs. 149 and 150), or

Associated with retroflexion (Figs. 152 and 153).

Retroflexion always has some retroversion combined with it (Figs. 152 and 153).

not so directly upwards and forwards as it does in pure retroversion; we feel the body of the uterus through the posterior fornix, and we notice that there is an angle produced where the axis of the cervix meets that of the body (Figs. 152 and 153).

To make sure that a lump felt through the posterior fornix is the body of the uterus, we rely on the following considerations:—

- 1. The absence of the body of the uterus from the front as ascertained bimanually.
- 2. On the lump felt through the posterior fornix moving with the cervix, and feeling continuous with it.
- 3. On information obtained by passing the sound. If the lump is the body of the uterus, the sound passes with its concavity directed backwards, and we can recognise that it has entered the lump felt through the posterior fornix; moreover, if the lump is the body of the uterus, and there are no adhesions fixing it, we can replace the uterus with the sound (as shown in Fig. 154) into a position of anteversion, or anteflexion. After this has been done it will be found that the lump previously felt through the posterior fornix has disappeared, and that on bimanual examination the body of the uterus can be grasped through the anterior fornix.

Attention to these points is needed to make certain that a lump felt through the posterior fornix is the body of the uterus, and not, for instance:—

A fæcal accumulation, or

A fibroid tumour growing from the posterior wall of the uterus, or

An enlarged and prolapsed ovary, or

A dilated Fallopian tube, or

A swelling due to hæmatocele or inflammatory exuda-




Fig. 154.—Replacement of the Retroverted or Retroflexed Uterus by the Sound (Thorburn). The first figure on the reader's left shows the sound passed into a retroflexed uterus.

The second figure on the left shows the first step of replacement, the handle of the sound carried back towards

The third figure shows the "tour de maître" (see p. 29), the handle of the sound being carried through a wide the perineum without reversing the curve (from  $\alpha$  in the second figure to  $\delta$ ), thus raising the uterus.

semi-circle from a to b, so as to bring the concavity of the sound forwards.

The replacement is completed by carrying the handle of the sound once more back towards the perineum, as in the end figure on the right. Retroversion or retroflexion with incarceration of the body of the uterus in Douglas's pouch.—In many cases of retroversion and retroflexion we find that the uterus is as movable as it normally should be. If, however, it happens that the size of the body of the uterus is such that when forced down into Douglas's pouch the uterus fits tightly there, we have the condition known as retroversion, or retroflexion, with incarceration. The uterus in such cases is fixed. The utero-sacral ligaments, which form the edges of Douglas's pouch, play an active part in retaining the body of the uterus in its incarcerated position. They contain muscular fibres, and can therefore exercise a certain amount of grip on the uterus, keeping it in Douglas's pouch, when it has once been pushed there.

#### ETIOLOGY.

Forward displacements.—The view taken here that forward displacements of the uterus (anteversion and anteflexion) have in themselves no pathological significance, renders it unnecessary to dwell at length on their etiology. It has already been said that the normal position of the uterus in the virgin, when the bladder is empty, is one of anteversion together with anteflexion, so that the axis of the body of the uterus forms an angle with the axis of the cervix on the anterior aspect of the uterus, the angle being an obtuse angle, and its value about 120°. Sometimes the angle is a much smaller one, and it may even be an acute angle.

An exaggerated anteflexion of this kind is met with :-

- I. In cases where the uterus is ill-developed.—Here we shall be able to make out that the uterus as a whole is small, e.g., only one inch and a half long.
- 2. In cases where there has been pelvic peritonitis, and the excessive anteflexion has been produced by the dragging of

adhesions. Here there will be probably a history of the inflammatory attack to guide us, and some evidence of it on physical examination: for instance, diminished mobility of the uterus.

3. In cases where there has been cellulitis in the utero-sacral ligaments.—As the inflammatory exudation contracts, it pulls on the uterus at the junction of the cervix and the body, the force acting in a backward direction; it is obvious how this will tend to produce an exaggerated degree of anteflexion. The mobility of the uterus will probably here also be diminished.

In the first group of cases it is not the anteflexion, but the want of development, that is of chief importance; and in the second and third groups it is not the anteflexion, but the pelvic inflammation, that is the pathological factor to be reckoned with.

Backward displacements.—It has been already said that retroversion to a moderate degree is physiological, in-asmuch as it occurs every time the bladder fills. We may speak of this as "physiological retroversion." Retroflexion is occasionally congenital, but more rarely so than excessive anteflexion.

Retroversion in excess of the physiological retroversion just spoken of, and retroflexion, are usually acquired.

They generally imply, at least when present to any marked degree, a certain amount of descent of the uterus also; their mode of causation is therefore in many cases identical with that of uterine prolapse (see Chapter VIII.). Retroversion and retroflexion may also be caused by the dragging of adhesions, the result of pelvic peritonitis; in such cases the mobility of the uterus will be diminished or lost, and it will be impossible without using an undue amount of force to replace the uterus into a position of anteversion or anteflexion with the sound.

Significance of versions.—Anteversion in itself has no pathological significance; if the mobility of the uterus is diminished, there has probably been pelvic inflammation, which is to be considered the important feature in the case.

Retroversion in excess of physiological retroversion, as we have seen, usually indicates some descent of the uterus, which under ordinary circumstances becomes retroverted as it prolapses. If the mobility of the uterus is lost, or diminished, and the uterus cannot be replaced with the sound, we are justified in considering that there has been pelvic peritonitis, and that the altered position of the uterus has probably been caused by it; however this may be, it is the pelvic peritonitis, and not the retroversion, that is the important feature of such a case.

So long as the uterus is freely movable, and not enlarged to any extent, retroversion of the non-pregnant uterus usually leads to no symptoms; there are some cases, however, where disorders of micturition (frequent desire to pass water, or incontinence of urine) are met with, associated with retroverted uterus; in cases of this kind the uterus is, as a rule, not quite of the normal size, but somewhat enlarged. In such cases it is desirable at all events to try the effect of replacing the uterus, and to insert a ring pessary to keep it from becoming again retroverted.

When the retroverted uterus is incarcerated in Douglas's pouch, some of the following symptoms are usually present:

Bearing-down pain in the lower part of the abdomen and back.

Dysmenorrhœa.

Menorrhagia.

Trouble connected with micturition, or defæcation.

Here it is the *incarceration* of the uterus, not the mere retroversion, that is important.

Whether marked symptoms exist, or not, will depend on

the tightness with which the uterus is gripped in Douglas's pouch.

Significance of flexions.—Flexions have been considered to have a pathological significance chiefly for two reasons:—

I. Because it has been thought that at the angle of flexion there was an obstruction to the circulation, so that the body and fundus of the uterus became congested—




Fig. 155.—Diagram to show the Position of the Blood Vessels supplying the Uterus (after Hyrtl).

O.A. Ovarian artery; U.A. Uterine artery; c d. Lateral arterial channel on each side; ab. One of the transverse branches given off from this lateral channel; xy. Indicates roughly the position of the internal os; us, us, show position of pressure on each broad ligament when the uterus is incarcerated in Douglas's pouch.

flexion of the uterus was thought to cause congestion of the body and fundus.

2. It was supposed that the canal of the uterus was narrowed at the angle of flexion, usually at the internal os uteri, so that an obstruction was produced at that point—flexion was thought to cause obstruction of the uterine channel.

- I. The first proposition—that flexion causes uterine congestion—must be regarded as disproved chiefly for the following reasons \*:—
- 1. Owing to the arrangement of the vessels supplying the uterus, it is impossible that flexion can interfere in any way with the return of the blood from the body and fundus of the uterus along the broad ligaments. This will be clear from the accompanying diagram (Fig. 155).

The ovarian artery reaches the uterus near the fundus on each side; as it nears the uterus it divides into two




FIG. 156.—Transverse Section of Uterus, showing the arrangement of uterine arteries, the arterial circles formed by their primary branches, and the branches of the latter supplying the mucous membrane (Sir John Williams).

The large white dot at each side of the figure corresponds to a transverse section of the lateral arterial channel, cd, in the preceding figure.

branches, one of which descends to join the uterine artery.

Thus there is, as it were, a lateral arterial channel, cd, cd, on each side, from which branches are given off that have a general transverse direction over the uterus with a slight inclination upwards.

These transverse branches join corresponding branches from the opposite side. From these transverse branches

\* The account in the text is based on a paper by Sir John Williams, "On the Circulation in the Uterus," in vol. xxvii, of the Transactions of the Obstetrical Society of London.

secondary smaller branches are given off towards the mucous surface of the uterus in a direction perpendicular to the plane of the paper. The veins have a similar arrangement. It will thus be seen that each transverse section of the uterus has its own vascular supply, and that a constriction in the situation of xy, about the level of the internal os, the usual situation of flexion, cannot affect the circulation in the uterus above or below.

When the uterus is incarcerated in Douglas's pouch, congestion of the body and fundus does occur, but here the obstruction is to the return of the blood along the broad ligament on each side—the edges of the pouch pressing on the broad ligaments in the situation us, us, on each side. It is not the retroflexion causing obstruction at the angle of flexion, but the pressure of the edges of Douglas's pouch on the broad ligament, that causes obstruction.

- 2. If flexion caused congestion, we should expect to have excessive menstruation in cases of flexion. Now, cases are constantly met with in practice where there is marked flexion, retro- or anteflexion, and yet there is no menorrhagia.
- 3. Though we speak of the angle of flexion, it is rather because the term is a convenient one than that there is any actual angle; what is usually spoken of as "an angle" is really rather a simple curve; owing to the thickness of the walls of the uterus, when the body is bent on the cervix, the intervening part of the uterus forms a curve, not an angle.
- 4. In the paper already referred to it was shown that when the fundus of the uterus was stitched to the cervix the vessels could be as well injected artificially as when nothing of the sort had been done.

II. As to the relation of flexion to obstruction of the uterine canal.

What has been said as to the angle of flexion being rather a curve than an angle applies here also. The direction of the canal is altered by the flexion, but that is all. The late Dr. Matthews Duncan, speaking of a specimen of an acutely flexed uterus, remarked that the flow of the menses out of it would not be "nearly so much obstructed as the passage of the water along a bend of the river Thames."

Obstruction produced by flexion has been considered of importance:—

- 1. As a cause of dysmenorrhœa.
- 2. As a cause of sterility.
- I. The supposed obstruction as a cause of dysmenorrhæa.—
  The explanation of the pain on this hypothesis is that it is caused by obstruction; excessive contractions of the uterus are set up to drive the menstrual fluid past the obstruction.

It has been shown above, from the examination of specimens, that in flexed uteri the direction of the channel is altered, but that the channel itself is not appreciably obstructed. Clinical evidence shows also that flexion is about equally common in patients with dysmenorrhæa, and in cases without dysmenorrhæa, as the statistics given in the following passage from Hart and Barbour show:—

"Herman and Vedeler have shown that the connection between anteflexion and dysmenorrhæa has been overestimated. In his very interesting paper on the cause of dysmenorrhæa, Vedeler reports on a large number of cases (observed by himself) of patients with and without dysmenorrhæa. To ascertain the relation of this symptom to anteflexion, we extract from his tables all the cases of nulliparæ with uterus to the front; we take nulliparous cases only, because parity in itself affects anteflexion; and

consider cases with uteri to the front as we are dealing with anteflexion only. We find that 37'3 per cent. (25 out of 67) of patients with dysmenorrhœa had a well-marked anteflexion, and that 33'3 per cent. (46 out of 138) of patients without dysmenorrhœa also had well-marked anteflexion" (Manual of Gynæcology, 3rd edit., p. 335).

Further, in many of the worst cases of dysmenorrhoea the sound can be passed quite easily, showing the absence of any obstruction.

2. As a cause of sterility.—Just as there is practically no obstruction to the exit of menstrual fluid at the angle of flexion, so there is none to the entrance of spermatozoa. Dilatation of the cervix probably only increases the chance of conception to a very slight degree.

Treatment of versions and flexions.—Anteversion and anteflexion require no treatment. Any symptoms present must be referred to some associated morbid condition, e.g., pelvic inflammation, recent or old, and not to the anteversion or anteflexion. When there is retroversion or retroflexion, with incarceration of the uterus in Douglas's pouch, the condition is to be looked upon as a pathological one, and remedied. The bladder should be emptied, and the uterus replaced with the sound; a suitable pessary, either an elastic ring or a Hodge's pessary, should then be inserted to prevent the displacement recurring. So long as the uterus is freely movable, any existing flexion should be regarded as having in itself no pathological importance.

Marked retroflexion, without incarceration of the uterus in Douglas's pouch, is usually to be taken as an indication that the uterus is lower than normal—that it is somewhat prolapsed. In such cases the treatment is that suitable for a case of slight prolapse; it is well to replace the uterus, so that the body of it lies to the front, before inserting a pessary. We shall then be able to know whether the pessary is

efficient or not by subsequent examination. If it is efficient, the uterus will be to the front; if it is not efficient, the displacement will have recurred.

It is well to understand clearly that no vaginal pessary, whether it be a ring pessary, or a Hodge's pessary, or any other, can straighten or keep straight a flexed uterus. The only pessary that can do this (were it of any importance to do it) is an intra-uterine stem pessary, *i.e.*, a pessary that has a straight rigid rod occupying the canal of the cervix and body of the uterus.

If a patient, complaining of pelvic pain, says that it is distinctly relieved by lying down, it is often worth trying if a pessary that tends to keep the uterus at a higher level will relieve her. For this purpose the ordinary ring pessary is the best.

In such cases the relief obtained is due to removal of passive congestion by raising the uterus as a whole, and by lessening the tension on the various uterine supports.

Some general remarks on vaginal pessaries.—For all ordinary cases, where the use of a vaginal pessary is indicated, either the ring pessary made of watch spring, covered with india-rubber, or Hodge's pessary, will be found to meet every requirement.

Several sizes of each should be at hand. Hodge's pessary is made of many different materials: for example, vulcanite, celluloid, or copper wire, covered with india-rubber.

The action of the watch-spring pessary, and the mode of inserting it, have been mentioned on page 150.

Position of Hodge's pessary.—Hodge's pessary should lie with the concavity of the upper curve directed forwards. The pessary should not be pressing firmly against any bony part, but should simply be grasped by the vaginal walls.

In the figure (157) of Hodge's pessary one end is shown as narrower than the other. In many Hodge's pessaries

there is no difference in the width of the upper and lower ends. So long as the concavity of the part lying highest in the vagina is directed forwards, it does not matter whether the width of the pessary is the same all the way down, or whether one end is narrower than the other. The point is to use a shape that fits the patient comfortably and acts efficiently.

Action of Hodge's pessary.—The upper limb of Hodge's pessary pulls the vaginal wall at its insertion posteriorly into the cervix in a backward direction, and so tends to




FIG. 157.—HODGE'S PESSARY. Side view and front view.

throw the weight, the body of the uterus, forwards, the fulcrum lying somewhere between these two parts.

To insert a Hodge's pessary.—One finger of the left hand retracts the perineum slightly; the pessary held in the right hand is passed through the orifice of the vulva, the bar closing the upper end of the pessary parallel to the cleft between the labia. When the pessary is in the vagina, it is turned round through a quarter of a circle, so that the upper bar lies transversely in the vagina. The upper end has a tendency to find its way into the anterior fornix; and care must, therefore, be taken to hook it with the fore-finger behind the cervix into the posterior fornix.

Whenever a pessary has been inserted, the patient should be told that she is wearing an instrument, and shown one of the kind she is wearing; and she should be warned that it should not on any account be left unchanged for a longer period than three months at a time. In general it is best to see her again in the course of a few days, to know if the pessary is comfortable. If then all is right, she need not be seen again for three months. A patient wearing a pessary should always be directed to use a vaginal douche of some antiseptic lotion night and morning.

If a pessary be used that fits too tightly, it will very likely cause ulcerations; and in neglected cases even vesicovaginal fistula may be produced. It is well to see that the diameter of the pessary at every part of it is a relatively broad one, so as to distribute the pressure over a broad surface rather than over a narrow one.

In fact, the nearer any part of a pessary approaches n character to anything like an "edge," the greater the probability of it causing ulceration.

Zwanke's pessary, sometimes known as the butterfly pessary, has two wings attached to metal rods, the ends of which can be fixed together by a screw-cap. When this is the case, the "wings" of the pessary are expanded. When the wings are closed, the ends of the rods are widely divergent. It is introduced in this state into the vagina; then the wings are opened by bringing together the rods and fixing them by the screw. The patient takes out the pessary at night, and replaces it in the morning. It is a very efficacious instrument for many cases of prolapse. The edges of the "wings" should be very thick and rounded. The patient should be warned of the danger of leaving the pessary in for an indefinite time.

I have known a large vesico-vaginal fistula produced by leaving a Zwanke's pessary for a long period unattended to in the vagina. The patient was an old woman, who had worn the pessary many years for prolapse with great comfort.

Then she got rheumatoid arthritis in her fingers, and was unable to turn the screw of the instrument; and so it came about that the pessary was worn many months without being removed. About three weeks before I saw her she began to find her water coming away, making her constantly wet. She then went to a medical man, who was unable to remove the instrument. She then came to the London Hospital. One wing of the pessary was found lying in the bladder, and encrusted with phosphates. On separating the metal rods widely, the pessary was easily removed.

I have also had to remove a Greenhalgh's pessary that had been in the vagina ten years without being taken out.

This form of pessary is shaped like Hodge's, but the lower cross-bar is formed of india-rubber *only*, the rest of the pessary consisting of copper wire covered with india-rubber. In this case one lateral half of the pessary was deeply embedded in the soft tissues, and had to be, as it were, dug out. After removal, a trench was left as broad and deep as to contain the little finger; but it is worth noticing that in this case no vesico-vaginal fistula had been produced.

Inversion of the uterus.—This is a very rare displacement. It may be either acute or chronic. The acute variety almost always occurs as a complication of the third stage of labour; exceptionally it may be produced by the dragging of a submucous fibroid tumour, or fibroid polypus, attached to the fundus of the uterus, or by traction on such a polypus while it is being removed.

The chronic form is a sequel of the acute, when the patient has survived the immediate consequences of the accident.

Pathology.—Relaxation of the fundus and neighbouring part of the uterus is a necessary condition for the production of inversion. Given this relaxation, then either

pressure from above, or traction on the fundus from below, may cause inversion. Instances of traction from below are (a) pulling on the cord in cases of adherent placenta, (b) dragging of a fibroid polypus adherent to the fundus, or its neighbourhood. An example of pressure from above, causing inversion, is where sudden pressure is made on the uterus with the hand in the third stage of labour, the uterus at the moment happening to be relaxed. Hence the importance of the practical rule in regard to expression




Fig. 158.—Instrument for replacing an inverted uterus by constant elastic pressure.

of the placenta, viz., only to make efforts to express it when the uterus is felt to be hard, *i.e.*, contracted.

Degrees of inversion.—It is usual to divide cases of inversion into groups:—

- 1. Where an inversion exists, but the inverted fundus has not passed the os uteri;
- 2. Where the inverted fundus has passed through the os; and
- 3. Where the whole uterus is turned inside out, cervix as well as body.

Symptoms.—In the acute variety there is severe shock,

and there may be hæmorrhage. In the chronic form symptoms may in rare cases be absent;\* usually, however, there is more or less bearing down, and the patient loses an excessive amount of blood (menorrhagia and metrorrhagia). When she is not losing blood, there is a yellow discharge,




Fig. 159.—White's Repositor, with spiral spring to place against the operator's chest (Thorburn).

owing to inflammation of the exposed mucous membrane of the uterus.

Physical signs and diagnosis.—These have been considered under the diagnosis of fibroid polypi (p. 204).

Treatment.—The treatment of acute cases occurring during labour belongs to Midwifery. In chronic cases we may try to effect replacement, either with the hand alone,

<sup>\*</sup> As in a case (quoted by Dr. West) observed by Madame Boivin.

or by pressure exerted by some instrument guided by the hand, or we may try the effect of constant elastic pressure on the inverted uterus.

It seems preferable to try the last-mentioned method first. An instrument on the principle of, and somewhat resembling, a cup and stem pessary is used. The cup is made of vulcanite, and is adapted to the inverted fundus. Any desired degree of upward pressure is then obtained by tightening up the elastic bands that pass to the waist belt. The best instrument for the purpose is that known as Aveling's repositor. This method may be used for two or three weeks, the patient remaining in bed, and the instrument being frequently examined to see it is in proper position; pain is, if necessary, relieved by hypodermic injections of morphia.

Should this treatment fail, we may try manipulation under chloroform, and we may either use the hand alone or use White's repositor, guided by the hand. If the hand alone is used, we should try to replace first the part last inverted. Dr. West refers to a case of thirteen years' standing, where Noeggerath succeeded in replacing the uterus by manipulation with the hand alone.

In very rare cases, where all means at our disposal for replacing the uterus have failed, and where the patient is becoming exhausted by the constant loss of blood, it is justifiable to remove the uterus by operation.

#### CHAPTER XVIII.

#### DISORDERS OF MICTURITION.

The significance of the following symptoms will be shortly considered:—

Frequent desire to pass water.

Pain on passing it.

Difficulty in passing it.

Retention of urine.

Incontinence of urine.

It is important to bear in mind that such symptoms may be due either to:—

- (1) Some disease or abnormality of the urinary tract itself, including alterations in the quantity or character of the urine.
- Or (2) morbid conditions of other parts—the urinary tract itself not being diseased, or at least not primarily so.
- I. Abnormal conditions of the urinary organs, or of the urine, causing disorders of micturition.
  - a. Diseases of the urethra.

Urethritis
Vascular caruncle

Malignant disease
Stricture

Common.

Rare.

Frequent desire to pass water and pain on passing it are the chief symptoms; occasionally there may be retention, which is partly voluntary in some cases from fear of the pain accompanying micturition. b. Diseases of the bladder.

Cystitis and its various causes, gonorrhœa, foreign bodies in the bladder (hair-pins, etc.), vesical calculi.

Malignant disease.—Primary (rare), secondary to malignant disease of uterus (common).

Frequent desire to pass water and pain in passing it are the prominent symptoms.

Vesico-vaginal or vesico-uterine fistula.—The characteristic symptom here is, of course, incontinence of urine.

c. Alterations in the urine.

Quantity.—If an excessive quantity of urine is secreted, as in diabetes, and some cases of hysteria, the bladder being frequently filled, there will be frequent calls to empty it. In these cases there is frequent micturition, but not painful micturition.

Quality.—When the urine contains "gravel," the bladder is irritated mechanically; in such cases micturition is painful as well as frequent.

d. Diseases of the pelvis of the kidney, or of the kidney itself.—Scrofulous disease of the kidney and pelvis of the kidney are perhaps the conditions most likely for a time to be overlooked, and the case regarded as one of cystitis.

Two cases in illustration of this are added at the end of the chapter. Malignant disease of the kidney also requires mention under this heading.

- II. Abnormal conditions outside the urinary tract causing disorders of micturition.
- a. Pelvic inflammation, including pelvic peritonitis, pelvic cellulitis, and ovaritis.

These may lead to pain on passing water, and frequent desire to pass it.

b. Pressure.—When the uterus is not enlarged, neither its anteflexion nor retroflexion disturbs the function of micturition; when the uterus is enlarged, however, the case is different.

If the enlarged uterus occupy a position of anteversion or anteflexion, we may have frequent desire to pass water, and sometimes involuntary passage of a small quantity of urine. Cases of this kind are met with during pregnancy, and also where the uterus is enlarged owing to the presence of fibroid tumours.

If the enlarged uterus occupy a position of retroversion or retroflexion, we may have disturbances of micturition, viz., frequent micturition, incontinence of urine, or retention.

The best example of micturition being interfered with by an enlarged retroverted uterus is retention of urine in cases of retroverted gravid uterus.

When there is retroversion or retroflexion of the gravid uterus, the uterus may rise out of the pelvis at the proper time (the beginning of the fourth month), and so the malposition be spontaneously rectified; sometimes, however, it does not do so, and retention of urine occurs.

This is due partly to the urethra being dragged upwards by the cervix, which is displaced high up behind the pubes, and partly to direct pressure on the urethra. Often there is some dribbling over of urine from the distended bladder, which may mislead the patient into thinking she is passing her water properly. In cases of this kind the degree of distention reached may be very great. For example, in several cases of the kind I have drawn off more than 100 ounces of urine, and in the following case, in every way a very typical one, 126 ounces:—

J. R., age 35, has had nine children, the last two years and two months ago. She was brought to the London Hospital on Wednesday, April 15th, 1891, with the following history:—She had been in her usual health up to 2 a.m. on the morning of the previous Sunday, April 12th, when on getting out of bed to pass her water she found herself unable to do so. She had pain in the lower abdomen for about half an hour before making the attempt to pass water. The pain increased after the ineffectual effort to micturate. She went back to

bed and stayed there. A doctor was called in, who gave her some medicine. She was only able to pass water once, and then very little and after great straining, between 2 a.m. on the 12th, and the afternoon of the 15th, when she was brought to the hospital. On inquiry it appeared that she had suckled the last child for one year and eight months, and that she had been last "unwell" soon after Christmas, 1890, since which time she had not been poorly. Before Christmas she had been for some time regular every three weeks, the period lasting three days.

April 15th, 1891.—Abdominal examination.—There is a tense, uniform, centrally situated swelling reaching about a hand's breadth above the umbilicus, dull on percussion and containing fluid. Nothing heard over it. Patient is evidently in great pain.

Vaginal examination.—Vulva and vaginal mucous membrane blue. Vaginal portion of the cervix high up above the pubes, and out of reach. Posterior vaginal wall bulged down by a soft swelling.

A catheter was easily passed, and 126 ounces of urine were drawn off, after which the tumour in the abdomen was found to have disappeared. The patient was admitted into the hospital. On the following morning, April 16th, the resident accoucheur, Mr. Calthrop, found that the uterus was still retroverted, and 80 ounces of urine were drawn off. At my visit in the afternoon of the same day I examined the patient, and found that the uterus had of itself gone up into its proper position, the os uteri then looking downwards and backwards. After this the patient was able to pass her water naturally. The urine contained no albumen (except a trace on one occasion), but hyaline and granular casts were found several times.

After emptying the bladder in cases of this kind we may either (1) replace the uterus immediately, under chloroform if necessary, or (2) keep the patient at rest in bed for a few days, at the same time taking care that the bladder is regularly emptied every six hours. In most cases, as in the one of which an account has just been given, the uterus will then rise out of the pelvis of itself. In some cases the urine first drawn off is dark, like porter, and contains an abundance of blood, probably from sloughing of the vesical mucous membrane.

Ovarian tumours very rarely cause retention of urine, but

they do occasionally. I have met with some instances of this (see p. 393).

Fibroid tumours of the uterus, on the other hand, often cause retention; and when this has been relieved, it does not necessarily recur, at least for some time.

Retention of urine may occur in advanced cases of malignant disease of the cervix involving the vagina. I have met with a case of this kind where five pints of urine were drawn off. The retention did not recur during the time the case was under observation.

Malignant tumours originating in the pelvis may cause frequent and painful micturition.

Tumours in the vagina, whether originating there or elsewhere, may also interfere with micturition.

- c. Hysteria may cause retention of urine, or lead to frequent micturition, owing to an unusually large quantity of urine being secreted; sometimes, however, in such cases absolutely less urine than normal is secreted.
- d. Labour.—Difficulty in passing water, or complete inability to pass it, may occur after labour, due to the bruising or laceration of the parts in the neighbourhood of the urethra.
- e. Procidentia.—Difficulty in passing water is not infrequently present; often the patient has found that by pressing the parts up she can pass water more easily.

Diagnosis.—The chief point to be considered at first is whether the symptoms are due to morbid conditions of the urinary tract, or of the urine; or to morbid conditions elsewhere,—whether, in fact, the cause is in Group I. or Group II.

Some of the causes in Group I., such as vascular caruncle, stricture of the urethra, and urethritis, are diagnosed by inspection.

Careful examination of the urine is of great importance,

and it should be drawn off with a catheter for the purpose, to avoid contamination with vaginal discharges.

Bleeding caused by passing the catheter probably means malignant disease of the bladder; fragments of the growth in such cases may come away in the eye of the catheter. The diagnosis would be completed by dilating the urethra with Hegar's dilators and passing the finger into the bladder. It is easy also to dilate the urethra with the fingers, first passing the little finger.

The sound \* should be passed into the bladder, and search made for foreign bodies or calculi. The distance the sound can be passed into the bladder, measured from the external orifice of the urethra, should be noted; in health it passes four and a half inches. This measurement is diminished in various diseases, particularly acute and subacute cystitis.

Tenderness on passing the sound also indicates disease of the bladder.

On vaginal examination, if we find there is special tenderness when pressure is made on the anterior vaginal wall, this may indicate cystitis.

When the urine contains pus, and is *acid*, the cause may be acute or sub-acute cystitis, or pyelitis. In chronic cystitis the urine is alkaline.

It must not be forgotten that pus in the urine may be due to a pelvic abscess opening into the bladder, or to a peri-urethral abscess opening into the urethra. I have seen a case of this; the abscess formed a swelling about the size of a walnut in the anterior wall of the vagina below the urethra. Pressure on the swelling caused pus to flow from the urethra. A counter opening was made at the most dependent part through the anterior vaginal wall. The cavity soon filled up, and no incontinence of urine resulted.

<sup>\*</sup> The ordinary uterine sound answers the purpose very well.

When for various reasons we suspect that the pus comes, in part at least, from the urinary tract above the bladder, it becomes a matter of importance to determine if possible whether the pus comes from one side or both sides, the possibility of relief by nephrectomy being kept in mind. To decide this point catheterisation of the ureters has been proposed and carried out by various observers; the practical difficulties of this proceeding are, however, great, and it is at present only successfully practised by a few who have given special attention to the subject. A method which I adopted successfully in one case many years ago is as follows:-The urethra is dilated; then one piece of a Bryant's rectal speculum is passed along the urethra into the bladder, and so placed that, seen from the front, it occupies one lateral half of the urethra and bladder beyond. When in this position, the speculum divides the bladder into two compartments: for example, supposing the speculum occupies the right lateral half of the urethra and the bladder, then the orifice of the left ureter is in view. The surface is gently mopped with cotton-wool, and then the character of the urine escaping from the ureter is observed. As it collects in the hollow of the speculum, some of it may be taken up with a syringe, and tested in the usual way. Having finished the examination of the left ureter, the speculum is now quickly turned round, so as to lie in the left lateral half of the urethra, thus bringing the orifice of the right ureter into view. When this was done in the case about to be narrated, a little fountain of clear urine about a quarter of an inch high escaped from the right ureter, the exit of the urine from it having evidently been prevented by pressure of the speculum, while the left ureter was being observed. Here also, if the obvious characters of the urine do not at once settle the point at issue, some may be collected and tested. The electric light is necessary for

illumination, unless direct sunlight is available. In the following case, observations made in the manner described, showed that suppuration was taking place on one side only, as pus was seen steadily oozing from the left ureter, and clear urine came from the right. Guided by the oozing of the pus from one particular point, a hollow probe was passed two inches along the left ureter, but I was not able to pass the probe along the right ureter. The evidence already obtained would not, however, have been rendered any more conclusive by catheterisation of the ureters. I think, then, that in women, when the urine contains pus,




FIG. 160.—BRYANT'S RECTAL SPECULUM.

and when, owing to other considerations—for instance, wasting, fever, failure to improve under local treatment, such as washing out the bladder with antiseptic lotions—we have reason to believe that the presence of pus is not to be accounted for by cystitis alone, either the procedure above described should be employed, or the ureters should be catheterised by Kelly's method, with a view to give the patient the benefit of surgical treatment if the affection prove to be unilateral. The following is an abstract of the case referred to:—

A. G., age 18, shop assistant, was admitted to the London Hospital on May 15th, 1886, complaining of inability to hold her urine. She first began to experience micturition trouble in November 1885. She noticed then that she had to void urine more frequently than she had to

do before, and had to get up at night to pass it. In January 1886 she began to have some pain during micturition, which became more and more frequent, and at last the bladder was so irritable, that for some time before she came to the hospital she had been quite unable to hold her urine. She had never passed blood with her urine. On one occasion, in April 1886, the patient had a pain across the small of the back and in the stomach. The pain was severe and shooting; it came on suddenly and went away suddenly; it did not cause nausea or vomiting, nor did it pass down towards the legs.

On admission, the patient was anæmic, but fairly well nourished, the temperature normal. There was no tenderness in the abdomen, or anything abnormal to be felt there. Vulva and skin adjacent to it red, and sore-looking. Hymen perfect, but lax. On vaginal examination, pressure on the anterior vaginal wall caused pain, much more than the same pressure on the posterior vaginal wall. The uterus and its appendages were normal. The sound passed into the bladder a distance of three inches, or rather less, the measurement being taken from the external orifice of the urethra. The passage of the sound caused pain. About a drachm of urine was obtained by passing a catheter; reaction amphoteric; \* it contained pus in abundance, and some shreds or flakes; no casts; only pus-cells and bladder epithelium were seen under the microscope. The other organs were healthy.

The case was thought to be one of cystitis only for some time, and the bladder was washed out daily with liquor carbonis detergens (one drachm to a pint), and afterwards with boracic acid lotion, without, however, any improvement taking place. May 31st .- The temperature had hitherto been normal; but from this date onwards the patient suffered from attacks of high fever running an irregular course, the temperature on several occasions being as high as 104°. July 8th.-Examined under ether to-day. No tumour to be felt in the region of either kidney. August 21st.—The patient has lost 13 lb. in weight since June 14th. 31st.—There have been general pains during the attacks of fever, but no pain specially referred to the region of the kidneys. For the last three weeks increased resistance has been felt in the region of the left kidney, palpation causing some pain, and exciting the abdominal muscles to contract. When the patient lay on her face, dulness over the left kidney in the back was found to reach two fingers' breadths lower than over the right kidney. Under ether a

<sup>\*</sup> I.e., turns red litmus blue, and blue litmus red.

lump was easily felt in the situation of the left kidney. The observations on the character of the urine flowing from the right and left ureters respectively were then made, as described above in this chapter. My colleague, Mr. Treves, saw the case with me, and it was agreed that it was probably one of scrofulous degeneration of the left kidney, and that the right kidney was healthy, as evidenced by the urine flowing from its ureter. Although the patient's condition was most unfavourable, yet, as the rapid loss of weight and other grave symptoms made it certain that the case must end fatally in a short time if left alone, we felt it right to suggest operative interference to the patient and her friends, at the same time putting the risk fairly before them. Consent having been readily given, Mr. Treves operated. The patient did very well after the operation, and the temperature was persistently lower than it had been previously, though still febrile. The urine contained (three weeks after the operation) a small quantity of pus, due no doubt to the cystitis which still remained. The amount of pus in the urine was trifling, however, when compared to the large quantity passed before removal of the kidney.

Whatever the ultimate result may be in this particular case, there is no doubt that a diagnosis sufficiently probable to justify an exploratory operation could have been made by the method described months earlier when no tumour was to be felt in the abdomen, and before the patient's strength had been so much reduced as to render her a very unfavourable subject for operative interference at all.

Note on the operation by Mr. Treves.—"The kidney was approached by the ordinary lumbar incision. It was embedded in an extensive mass of tough adhesions. It was subsequently found that this perinephritic inflammation was the main constituent of the 'renal' tumour, since the gland itself was but little enlarged. The adhesions were broken down with difficulty. It was found impossible to separate the kidney from the capsule, or from the supra-renal body. The latter structure, with the entire kidney, was therefore removed. A surgical pedicle could not be established. The ureter, distended with caseous and purulent matter, gave way; the renal vessels were clamped, and subsequently ligatured. The kidney was entirely occupied by a series of tubercular abscesses. The wound was stuffed with sponge covered with iodoform, and the cavity left to granulate up."

January 1888.—The patient now looks fat and well. The wound completely healed some time ago. She still cannot hold her water properly, but she has recently acquired the power of holding it for a

short time, which is a great improvement. It is as certain as anything can be that she must have died long ago had the operation not been performed.

January 1890.—Still quite well; urine normal.

Dr. Howard Kelly, of Baltimore, has successfully practised catheterisation of the ureters, and has devised a special cystoscope and sounds for the purpose. With the aid of the cystoscope the ureteral sound can be passed into the orifice of the ureter under the direct guidance of the eye.

The existence of a vesico-vaginal or vesico-uterine fistula, and its position, can be demonstrated by pouring milk into the bladder while we are looking into the vagina with a Sims's speculum.

Coming now to Group II., we need only say that pelvic peritonitis and cellulitis, enlargements of the uterus, ovarian and other tumours, will necessarily be discovered by a careful physical examination; and as regards hysteria, we shall be guided to a diagnosis by an absence of local physical signs, combined with the general appearance and bearing of the patient.

Sometimes, apart from hysteria and apart from any discoverable disease, local or general, congenital incontinence is met with, persisting even up to seventeen or eighteen, or later, and resisting all treatment, as in the following case:—

J. W., age 17, admitted to the London Hospital on account of inability to hold her water since birth.

December 29th, 1888.—She attended first in the out-patient department for a short time. The uterus was found retroverted, not enlarged, and freely movable. It was placed in a position of anteversion with sound, and a ring pessary inserted. Incontinence continued as badly as before. Ring removed and patient admitted. There was simple erythematous vulvitis, and also soreness and redness of the skin round the vulva, so that there was no doubt as to the reality of the incontinence. (In some cases where incontinence is complained of, there is no such vulvitis or

condition of skin; then probably we may take it that the incontinence is trifling and may be curable.)

January 5th.—Bed always wet; patient passes her water voluntarily (I oz.) every quarter of an hour; during night, three times. Blistering above the pubes. Large doses Tr. Belladon. tried without effect, also cautery to urethral orifice and anterior vaginal wall for about one inch up. This latter caused a slight improvement till the soreness produced by the cautery disappeared, but the improvement was only temporary.

January 17th.—Constant current; one pole, lumbar spine; the other over pubes ten minutes; same treatment till February 3rd: no better.

N.B.—Use of cautery after failure of electricity. Went home to Blackwater absolutely without the least improvement. In other ways she seemed a sensible girl. (Urine, acid; albumen, o; sugar, o; sp. gr., 1010; sound to bladder, 4½ to 5 inches.)

Treatment.—The treatment suitable in each case will depend on the cause: for example, vascular caruncle must be burnt off; vesico-vaginal fistula operated on with a view to closing the fistula; ovarian tumours removed; foreign bodies, or calculi in the bladder, extracted; a scrofulous kidney, if the disease is proved to be unilateral, may be extirpated. In cases of cystitis for which no cause can be discovered, rest in bed, with frequent hot hip baths, is useful. In such cases the bladder should be washed out two or three times a day with water containing 3ij. of liq. carbonis detergens to the pint.\* The best way of doing this is to use a catheter made of india-rubber, with a glass funnel fitted on the other end of the catheter. The bladder is first emptied with another catheter; then the indiarubber catheter is passed, and the solution to be used poured into the glass funnel. The quantity that can be introduced depends on the irritability of the bladder: perhaps only two or three drachms, or less, may be tolerated.

<sup>\* 1-10,000</sup> corrosive sublimate lotion also answers very well, used once a day.

Large quantities of demulcent drinks, such as barley water, are useful. Where pain on passing water is a marked symptom, and is due to disease of the bladder, or urethra, tincture of hyoscyamus, decoction of pareira, bicarbonate of potash, spiritus ætheris nitrosi and camphor water, separately or in various combinations, may be prescribed. Oil of sandalwood in capsules mxv. thrice daily is also useful. It is often advantageous to give besides a mixture containing benzoate of ammonia in twenty-grain doses in combination with Tinct. Collinsonia Canadensis (mxx.-xxx.), and Buchu.

Goodell speaks highly of belladonna "in almost every form of vesical irritation."

His favourite formula is:-

R. Atropin. Sulph., gr. ss. Aq. destillat., 3iv.

Four drops to be taken in a wineglassful of water before each meal. To be increased or diminished according to the constitutional effect.

For nocturnal incontinence in young girls we attend carefully to the state of the bowels. If the girl suffers from worms, suitable treatment is adopted; we advise the mother not to let her drink any fluids for two hours before she goes to bed; and she should be wakened to pass water when the mother is going to bed. Sometimes a blister applied above the pubes is efficacious; sometimes benefit is derived from small doses of tincture of belladonna. Of course we are now speaking of cases where the urine is healthy, and there seems no reason for suspecting organic disease.

The following case is an example of the latent character of scrofulous disease of the kidney. It was not suspected during life.

M. K., age 30, was admitted to the London Hospital complaining of pain on micturition, and of having to pass her water frequently for the

previous six weeks. The urine passed during this time had been milky. She had had to get up at night to pass her water.

Has been troubled with cough and shortness of breath five months.

Had one child born dead; she thinks it was at full term.

On admission the urine was found to contain a small quantity of pus, and was acid.

The case was thought to be one of sub-acute cystitis, and treated accordingly.

She died suddenly of syncope, apparently due to fright, while an operation was being performed in the ward.

On post-mortem examination there was found advanced scrofulous degeneration of the left kidney. The right kidney contained two small abscesses, but was for the most part healthy.

She had a typical kyphotic pelvis, the measurements of which are appended.

Conjugate of the brim, 54 inches.

Transverse, 51 inches.

Conjugate at the outlet with the coccyx pushed back, 31 inches.

Between ischial tuberosities, 3 inches.

Between ischial spines, 315 inches.

Diagonal conjugate (i.e., from the sacral promontory to the lower border of the symphysis pubis), 6 inches.

The following case is an example of malignant disease affecting a movable kidney. The prominent symptoms were occasional hæmaturia; sometimes pain on micturition, apparently only when clots had to be passed; wasting, and the presence of a tumour in the abdomen.

M. B., age 62, married at the age of 27, a widow since she was 40, has had six children. She was admitted to the London Hospital under my care on December 29th, 1887, complaining of a tumour in her right side, and having passed blood with her water.

History of the illness.—Nine or ten months ago she passed "blood instead of water." The next morning after getting up she was attacked with severe pain in the right iliac region and in the stomach. She passed a motion, but this did not relieve the pain. A doctor who was called in declared the lump in her side to be a fæcal accumulation, and ordered her some aperient medicine, which gave her some relief.

Two or three months afterwards she again passed a large quantity of dark-coloured blood, together with a few clots. The passing of it was accompanied by a good deal of pain. Since then she has been losing flesh and feeling weak.

She first felt pain on passing her water seven months ago, and has suffered from it occasionally since, especially six weeks ago, when it was of a severe cutting character. Until a month ago she had to get up at night to pass water, but lately she has not had to pass her water with undue frequency.

December 29th.—In the right lumbar region there is felt a hard mass moving on respiration, and easily separable from the liver. It is very freely movable on manipulation. The mass feels solid, and like a movable kidney, only it is larger than a healthy kidney, and in places the surface of the tumour is somewhat nodular.

It is not tender.

When the finger is laid lightly on the abdomen over the tumour, there is resonance, but less than in the corresponding iliac region. When the finger is pressed down on the surface of the tumour, the note obtained on percussion is dull.

Bladder.—The sound passed  $4\frac{1}{2}$  inches into the bladder, measured from the external orifice of the urethra. No pain caused by passing it.

Urine.—Alkaline, sp. gr. 1008, smoky in colour, contained albumen, and on standing a deposit formed, which was found to contain blood corpuscles and pus corpuscles.

On another occasion the urine was acid, and on standing gave a deposit in which blood corpuscles were seen, though the guaiacum test gave a negative result. I thought the case was one of malignant disease affecting a movable kidney, and transferred the case to the surgical side for operation.

The tumour was removed on January 20th, 1888, and found to be what had been expected.

On cutting the kidney open there was seen a partially decolorised clot, forming a cast of the pelvis of the kidney. The malignant growth was in the form of white circumscribed nodules, the projection of which caused the unevenness of the surface referred to above.

The patient unfortunately died of peritonitis a few days later.

Hæmaturia in the newly-married.—I have seen one remarkable case in which marked hæmaturia occurred soon after marriage. Of course, as a general rule, such a symptom

is merely due to admixture of blood (resulting from lacerations of the hymen) with the urine; but in the case I refer to the urine was drawn off with a catheter by myself, and found to contain a good deal of blood. It is not easy to explain how this hæmaturia originated—I merely record it as a matter of clinical interest.

# CHAPTER XIX.

### MALFORMATIONS.—STERILITY.

Malformations.—These will be merely mentioned, as they are all rare, and most of them very rare.




FIG. 161.

1. Of the vulva.—All malformations of the vulva are instances of arrested development. The following five figures from Schroeder illustrate stages in the development of the vulva, terminating in the normal condition of parts



FIG. 162.

(Fig. 165). Any one of the conditions represented in the first four figures may persist, and constitute a malformation. In figure 161 we have the allantois, the rectum, and Müller's ducts communicating, but shut off from the exterior.

This condition, absence of the vulva, is found in some monstrous fœtuses.

In figure 162 a depression of the skin has established a




FIG. 163.

communication with the exterior; but the rectum, bladder, and vagina open into a single passage or cloaca.

In figure 163 the tissues between the rectum and bladder




FIG. 164.

are seen to be descending, so as to divide the common opening into an anterior part, the uro-genital sinus, and a posterior part, the anus.

In figure 164 the perineum has now been completely





Fig. 165.

formed, but the bladder and vagina still open into a common opening, the uro-genital sinus.

In figure 165 the uro-genital sinus is divided, by descent of the tissues between the bladder and vagina, into an upper part, the urethra, and a lower part, which forms the vestibule, thus completing the normal arrangement of the parts.

2. Of the uterus.—It is easy to understand the various



malformations that occur, if it be borne in mind that the Fallopian tubes, the uterus, and vagina are developed from two tubes—Müller's ducts. These remain separate above, forming the Fallopian tubes; but below they coalesce, the

septum between them being absorbed, to form the uterus and vagina.




Fig. 167.—Uterus duplex (Thomas).




Fig. 168.—Uterus septus (Schroeder, after Kussmaul).

The following malformations are described :-

- 1. Uterus unicornis.—Here one of Müller's ducts has not developed (Fig. 166).
  - 2. Uterus duplex. -Both Müller's ducts have developed,

but they have not coalesced, the result being two almost distinct uteri, merely joined where they lie in contact.

3. Uterus septus.—The external appearance is that of a normal uterus, but on section the uterine cavity is seen more or less distinctly subdivided into two. The vagina may be single or double (Fig. 168).

# A Case of Double Uterus with Submucous Fibroid Tumour.

A married woman, 38 years of age, was sent to me by Mr. Snowman, and admitted into the London Hospital under my care on October 16th, 1895. She had been married fourteen years, but had never been pregnant. The catamenia appeared when she was fifteen, but she was not regular till she was twenty-one, several months intervening at times between the periods. Since she was twenty-one she had been quite regular every four weeks till seven months ago, but nearly always had some pain during menstruation. She complained that for the last seven months she had been losing too much at her periods, and for the last six months there had been a constant red discharge between the periods, but till recently she could still tell when the menstrual periods occurred. For the last six months she had passed clots during menstruation, but had never done so previously. The dysmenorrhœa had been relieved during the last seven months by lying down, but this is said to have had no effect in lessening the excessive flow. She had had some pain and smarting during micturition ever since her marriage, but this had been worse during the last seven months. She had been losing flesh during the last six months. On vaginal examination the uterus was found considerably enlarged. A curious falciform fold of mucous membrane (F, Fig. 169) was felt at the upper part of the vagina running from before backwards and from right to left. The sound passed four and a half inches. At this time it was not recognised that there was a second vaginal portion, as the finger passed to the right of the fold just described. After disinfecting the vagina, a laminaria tent, specially rendered aseptic by prolonged immersion in absolute alcohol containing I per cent. of corrosive sublimate, was inserted into the cervix (A)—that on the right side—on October 16th, at 5 p.m., a strip of · iodoform gauze being packed below the tent to keep it in position. The next day, at 2 p.m., an anæsthetic was given, and the patient

placed in the lithotomy position. The iodoform gauze was removed, and a vaginal douche of perchloride of mercury (1-1000) given. The upper part of the vagina and the cervix into which the tent had been inserted were then thoroughly exposed with Sims's speculum and




Fig. 169.—Uterus septus (Author's case).

A. B. Two separate vaginal portions, each with its external os uteri. E. Spur above which the septum has been absorbed. C. Cavity of uterus. D. Submucous fibroid attached at fundus. F. Falciform fold of mucous membrane described in the notes of the case. V. Vagina.

retractor. It was then seen that there was a second vaginal portion (B) to the left side. It was perfectly well formed, but slightly smaller than the vaginal portion on the right side. The falciform fold of vaginal mucous membrane (F) passed between the two vaginal portions, running, as

already mentioned, from before backwards and slightly from right to left. The tent was now removed from the right cervix. The uterine sound was passed into each cervix and passed the same distance on each side-viz., four and a half inches. The right cervix (A) was now further dilated by means of Hegar's dilators till the finger could be passed through it into the uterus. The submucous fibroid (D), the size of a large walnut, was felt attached at the fundus and to the right side. It was seized with polypus forceps and twisted till it seemed loose, but it could not be brought away till the cervix was further dilated up to The condition of the uterus was then the size of No. 24 Hegar. examined more closely. The uterine sound was passed into the left cervix, and the finger into the cervix on the right side, through which the fibroid tumour had just been removed. The condition of things figured in the illustration was then found-that is, the finger felt the sound in the cavity (C) of the body of the uterus, showing that there was a single cavity for the body of the uterus, into which led the two separate cervical canals. There was a sort of spur (E) felt a little above the level of the internal os. Bimanually the upper part of the uterus was felt to be slightly irregular, there being three distinct prominences, as figured in the illustration, most probably due to the presence of sub-peritoneal fibroids. The patient made the uneventful recovery usual nowadays after operations of this nature.

4. Uterus bicornis.—The external appearance here is not that of a normal uterus, as there is a sulcus at the fundus, dividing it into two horns (Fig. 170).

The cavity may be completely divided into two by a septum, or incompletely, there being two cavities above and a single cervix.

In rare cases the uterus may be rudimentary or absent. In such cases the ovaries are usually absent, but they may be present.

The following case was one of ill-developed uterus, with imperforate hymen, and probably imperforate vagina:—

Kate L., age 20, single, a general servant, came to the London Hospital complaining of "general debility," and of having never "seen anything." On examination it was found that the hymen was

imperforate, but not bulged, as it is when there is a retention of menstrual fluid above it. The orifice of the urethra was rather small, but she had no trouble as to micturition.

She was a patient in whom bimanual examination was very easy and




Fig. 170.—Uterus bicornis (Schroeder).

satisfactory. On examining her per rectum in this way, the only representative of any uterus to be felt was a small solid body about one inch long and perhaps a quarter of an inch thick. The whole pelvis was thoroughly searched bimanually, but nothing else was found.




Fig. 171.—Infantile Uterus (Schroeder).

Infantile uterus.—Here the uterus in the adult retains its infantile characters; that is to say, the cervix is relatively

well-developed as compared with the body, which forms only a quarter or thereabouts of the whole length of the uterus. The whole uterus is small, one and a half to two inches long. In such cases the vagina and ovaries are also usually incompletely developed.

Congenital atrophy of the uterus.—Here the relative proportions of the body and cervix are normal, but the whole uterus is small. Usually there is a very marked anteflexion of the uterus, and the vaginal portion of the cervix points forwards. In such cases, if menstruation occurs at all, it is irregular and painful, and the function is late in beginning. The menopause occurs early.

# Malformations of the vagina.

The vagina may be absent.—The uterus and its appendages may in such cases be either rudimentary or well-developed; in the latter case the non-development of the vagina causes retention of menses (Hæmatokolpos, Hæmatometra). (Consult pages 103 and 104.)

I have seen one case of this due to a complete septum about one inch and a half from the vaginal orifice. The external organs of generation were normal.

The following is an abstract of the case:-

S. K., aged 18, single, complained of pains in the abdomen that had lasted two years, and of pain in the left groin of two months' duration; she had occasionally great difficulty in passing her water for two years.

Menstrual history.—She had never "seen anything," but she has every month a feeling of malaise, and the pains in the abdomen are worse at that time.

She had been in bed for a month before I saw her, and she felt ill in herself.

Appetite poor. Pains at the epigastrium after food. Bowels very loose, acting two or three times a day.

On abdominal examination an elastic swelling is felt occupying the hypogastrium; at its upper part a small knob-like projection can be felt (uterus?). The whole swelling is about the size of the uterus at the end

of the fourth month of pregnancy. External genitals normal. The vagina is completely occluded by a transverse septum about one inch and a half from the orifice. Bimanually it can be recognised that there is a collection of fluid between the hand on the tumour in the hypogastrium and the finger in the vagina. I heard subsequently from her medical attendant that she had been operated on, and the usual treacly fluid evacuated, but that the patient died of septic peritonitis about a month after the operation.

I have also seen a case where there was a somewhat similar septum, apparently at first sight complete, but in which careful examination discovered a fine aperture just admitting an ordinary surgical probe.

The details of the case were as follows: -

Edith M., age 20, married two years, no children or miscarriages, came to the London Hospital complaining of pain in the left iliac region, worse at the periods; she had had this for a long time, but the pain has been worse at the last three periods.

She has had a white vaginal discharge for two years.

Menstrual history.—Catamenia appeared at thirteen, regular every three weeks, scanty and pale, always attended with much pain, which, as mentioned above, has been worse the last three times.

On examination a septum was found across the vagina about one inch and a half from the orifice. The external organs were normal. Per rectum the uterus could be felt to be of the normal size and freely movable.

I took her into the hospital and examined her under ether. A careful examination of the septum showed the presence of a minute aperture in it, just admitting an ordinary probe. The septum was then cut through with scissors, and the opening enlarged by dilating with the fingers. Nothing else abnormal was detected. The patient did well subsequently.

The following case was one of complete absence of the vagina, with double uterus and double hæmatometra. I believe the case to be unique in respect of the double hæmatometra:—

L. T., a single girl, aged 17, was first seen by me on November 26th, 1893, at Reading, at the request of my friend Dr. Arthur Roberts, of

Reading, who was attending her. On that occasion a very careful examination was made, the patient being under the influence of an anæsthetic administered by Dr. Rows, of Reading. The condition of things discovered at that examination is described below.

Family history.—Her father is alive and well. Her mother died during the patient's infancy. She has no sisters or brothers, but she has half-sisters and half-brothers.

Personal history.—She is seventeen, and will be eighteen on the 24th of May next. She has lived all her life at Southampton, and has had no illness of any kind before the present trouble began.

Special history.—She has never "seen anything" at all, but for some time past (the time when the pain began was not given very accurately) she has complained of great pain in the hypogastric region and back, which comes on about once a month, and lasts about fourteen days. Severe pain is practically her only symptom.

At the examination on November 26th, 1893, the following facts were noted:—The patient is a tall, well-developed girl; the breasts are well developed. On examining the abdomen no swelling could be felt in the hypogastric region, nor was there any swelling to be felt there at any time subsequently, though she was examined on many occasions extending over a period of more than two years, during which she was kept under observation before the operation.

Vulva.—When the labia are not separated the appearance of the external genitals is quite normal; the mons Veneris, labia majora, and pubic hair present their normal appearance. On separating the labia majora, the nymphæ, glans clitoridis, vestibule, urethral orifice, fourchette, and fossa navicularis are normal, but where the vaginal orifice should be the condition is as follows:—There is a slightly raised median ridge running from before back from the posterior margin of the urethral orifice to where the posterior margin of the vaginal orifice should be. On each side of the raphe described there is a slight depression of the surface, to the extent of  $\frac{1}{32}$  of an inch or so, and round each of these depressions there is a distinct but rather rudimentary hymen. That is to say, where the vaginal orifice should be there is a central antero-posterior ridge, with a very shallow depression surrounded by a rudimentary hymen on each side of it (see Fig. 172).

On rectal examination a uterus can be felt bimanually; it is in the normal position and fairly movable. As regards its size, the uterus appears to be rather larger than the uterus of a virgin aged seventeen, but not very much larger. Certainly it is not so much enlarged as to

lead one to feel sure there is any accumulation of menstrual fluid inside it. With the sound in the bladder and the finger in the rectum, the partition between the rectum and the bladder is felt to be very thin, and especially so in the neighbourhood of the cervix. A dimple is felt at the end of the cervix, which is taken to be the external os uteri; it points backwards and somewhat to the right. The distance from the anus to the position of the os was measured as carefully as possible with




Fig. 172.—Case of L. T. Double Uterus, with Double Hæmatometra and Complete Absence of the Vagina.

The dark area immediately in front of the anus is the artificial vagina formed by the operation. In it, high up, are seen the two openings, one into each half of the double uterus, made at the operation. In front of the oval dark area (artificial vagina) is seen the condition of the external genitals before (and after) operation.

the finger, and found to be about 2 inches. This is of interest in the light of what occurred at the operation, as the dissection had to be carried about 3½ inches or more from the surface before the dimple in question was reached. As regards the opinion of the case arrived at

on this occasion, I came to the conclusion that there was no vagina, but that, on the other hand, there was no evidence on physical examination of distention of the uterus. As the patient only complained of pain, it did not seem to me at that time that making an artificial passage to the os uteri would necessarily relieve it, as the pain did not seem to depend on distention of the uterus. I therefore advised postponing the question of any operation till there was definite evidence of accumulation in the uterus. The patient was given ichthyol in pills for some time, and it seemed for a while to relieve her. I should say that I took her into




FIG. 173.—Diagrammatic representation of the uterus in case of L. T. before operation. The uterus is seen in coronal section (i.e. a section parallel to the coronal suture).

the London Hospital on January 12th, 1894, and kept her in till March 2nd, 1894. During that time there was no rise of temperature; the only symptoms were the pain described and headache. There were intervals when she was free from any discomfort.

Examination during her stay in the hospital confirmed the results already found when she was examined at Reading.

I saw her at intervals during 1894 and 1895. Sometimes she was free from pain for long periods, while at other times the pain was very severe. She was readmitted into the London Hospital on October 31st, 1895. The attacks of pain had been irregular in their occurrence, and

not every month. For instance, the last onset of pain was six weeks before her readmission, and it continued up to her admission. The pain is often worse at night. The pain is said to be rather to the left side of the hypogastric region, and to extend down the left leg as far as the knee, as well as being felt in the lower part of the back.

November 14th, 1895.—An anæsthetic was given, and she was again thoroughly examined. There was no swelling to be felt in the lower




Fig. 174.—Diagram to show the relation of the uterus to the adjacent parts in mesial sagittal section in the case of L. T.

part of the abdomen even when she was deeply under the influence of the anæsthetic. On bimanual examination per rectum the uterus was felt to be distinctly enlarged, and about the size of the pregnant uterus at the eighth week. There was a swelling to the left of the uterus, which could not, however, be separated from the uterus, and it seemed like an enlargement of the uterus in that direction.

I felt sure now that the uterus was distended, and accordingly decided to operate. Extensive alterations were being made in the ward at the

time, causing a great deal of noise, so that the operation was postponed till January 2nd, 1896. In the meantime the patient was often in great pain, and hypodermic injections of morphine were frequently needed.

As to the nature of the proposed operation, the question as to whether a preliminary abdominal section should be undertaken, as has often been suggested (recently again by Dr. J. C. Webster, in an article on "Atresia of the Vagina," in the American Journal of Obstetrics for October 1895), was carefully considered. It is well known that dilated Fallopian tubes not rarely complicate cases of hæmatometra. As I had carefully watched the case for two years, and seen the patient both when the uterus was little if at all distended, and again when the uterus was clearly enlarged, it seemed to me that most probably in this instance the Fallopian tubes were not dilated. I therefore decided to attempt to make an artificial passage up to the uterus without any preliminary abdominal operation.

Operation, January 2nd, 1896.—The patient was anæsthetised with the A.C.E. mixture, and placed in the lithotomy position. The parts having been thoroughly sponged with corrosive sublimate lotion (I in 1000), a transverse incision about 21 inches long was made in the perineum just behind the fourchette. A sound was held in the bladder, and the left forefinger frequently introduced into the rectum during the operation. The wound was gradually deepened, partly by tearing, but principally with the knife and scissors. The sides of the wound were held apart by strong retractors right and left. The point aimed at was the dimple felt on examining per rectum. The dissection had to be carried much further from the surface than had been anticipated; and its depth was ultimately at least 3\frac{1}{2} inches. The strong traction on each side of the wound had gradually torn the perineum back to the very margin of the anus, and a few fibres of the sphincter ani were torn. The cavity produced by the dissection was of a capacity to allow of its containing the fist. It was at length possible to recognise that the immediate neighbourhood of the uterus had been reached, a part uncovered by bladder. There was no trace of any pre-existing cavity: an interesting point, because in cases of atresia of the vagina the atresia is often confined to the lower part of the vagina—the lower third or half, for instance. When the obstruction has been opened up to that extent, a normal vagina is found above. There was nothing of the kind in the present case, nor was there any trace of the normal smooth covering of the vaginal portion. The part of the uterus corresponding to the vaginal portion, as regards its aspect towards the vagina, merely presented the raw surface made by the dissection. As the dissection proceeded it had gradually come to my knowledge that there was something in the nature of a second dimple felt to the right of the one that had been recognised before the operation.

When the amount of tissue over the principal dimple, that to the left (of the patient), was felt to be very little, a pair of Wells's small pressure forceps was thrust into it, evidently at once entering a cavity; the usual dark treacly fluid met with in cases of retained menses at once began to escape. The opening was enlarged by separating the blades of the forceps till the finger could be passed into the cavity. This was felt to be slightly loculated at one place, and to extend some distance to the left side.

Towards the middle line this cavity presented a vertical partition, towards the left its wall was concave. The dimple to the right side was now localised as well as possible, and a pair of Wells's forceps similarly thrust into it. It entered a larger cavity than that on the left side, and the puncture was followed by a fresh escape of retained fluid. The opening was enlarged with the forceps, and the finger passed into this right cavity. This cavity seemed to have been about the size of a duck's egg. Each cavity was now washed out with hot iodine water, a strip of iodoform gauze was packed loosely into it, an inch or so being left hanging from each os uteri into the vagina. The wound (vagina) was now packed with iodoform gauze. Two stitches were put in to unite the tissues for a quarter of an inch in front of the anus, and the patient was put back to bed.

Subsequent progress.—The patient went on very comfortably from January 2nd, the date of the operation, till January 13th, having little or no pain, and no rise of temperature. The gauze was still in; a little of the lower end of the part used to plug the vagina had been cut away on two occasions.

On January 15th the temperature rose for the first time since the operation, reaching 101°. The gauze was removed from the vagina, and also the two strips from the two compartments of the uterus. A good deal of pus followed the removal of the vaginal plug, and some also the removal of each uterine plug.

On the 16th the temperature was still up (102°). As the patient was extremely tender, an anæsthetic was given in order to wash out the two uterine cavities thoroughly. This was done with iodine water. The finger was passed into each os uteri. Each uterine cavity was

mopped out with pure tincture of iodine, the excess being washed away, and iodoform was introduced with a Recamier's curette freely into each cavity; a good deal of iodoform was also left in the vagina, but no plug of gauze was used on this occasion.

The rest of the patient's progress was uneventful. The cavity contracted rapidly, but was kept from becoming too small by passing No. 26 of Hegar's bougies daily. A pewter ring was modelled to fit the vagina, and the patient left the hospital on February 22nd. Examination before she went out showed that the uterus had become very much smaller than it was found to be any time before the operation. When examined with a Sims's speculum, the two apertures leading to the uterine cavities could be seen about half an inch apart at the top of the vagina. The vagina was about two inches deep from the surface. Some of its surface was still granulating, but over a good deal of the lowest part the epithelium had spread in. The patient had complete control over the bladder and rectum. She was seen again a week afterwards, and a small ring pessary, 15 inch in diameter, was put in instead of the pewter pessary.

There has been a great deal of difficulty in keeping the artificial vagina open. When an ordinary small ring pessary is left, the parts cohere through the opening of the ring. I therefore had a small vulcanite ring made with a vulcanite diaphragm, perforated by a few small holes. This acts efficiently. The patient menstruates regularly and painlessly. I have seen her regularly since the operation, the last time in October 1897.

The vagina may be double.—There is a septum dividing the vagina wholly or partially into two separate parts. When there is a double vagina, and also a double uterus, on one side there may be occlusion of the lower part of the vagina, the other opening as usual; on the occluded side there will be retention of menses. This is called "unilateral hæmatokolpos."

Atresia or occlusion of the genital canal.—This may be congenital or acquired.

In congenital cases it may be due to either imperforate hymen, occlusion of the vagina, or less commonly occlusion at some part of the cervix. Acquired occlusion is due to inflammation, followed by adhesion of the inflamed surfaces. Injuries to the vagina and cervix during labour, and sloughing of the vaginal walls after acute specific fevers, account for most of these cases.

#### STERILITY.

The late Dr. Matthews Duncan showed that there is no reason for regarding a woman as probably sterile till three complete years of married life have passed without the occurrence of pregnancy.

Etiology.—The local causes to which sterility may be ascribed are:—

- I. Pelvic peritonitis which has obliterated the openings of the Fallopian tubes.—When this has occurred on both sides, so that both Fallopian tubes are occluded, it is of course impossible for an ovum to reach the uterus, and the sterility is absolute. This is by far the commonest local cause of sterility.
  - 2. Ovarian tumours (if double).
  - 3. Morbid conditions of the endometrium.

    Corporeal endometritis.

    Fibroid and other polypi.
  - 4. Imperfect development of the uterus or ovaries.

As regards the above three sets of causes we can have no doubt of their influence.

5. All causes of pain on intercourse.—Particularly if it is severe enough to give rise to vaginismus.

We must not forget that the fault may be with the husband,\* and it is a fact that the same woman may be sterile with one husband and fertile with another.

\* M. Pajot, basing his observations on notes of 400 cases, concludes that, of 100 sterile marriages of from two to fourteen years' duration, the husband is at fault in from fifteen to twenty cases. (See Sterility and Impotence, Ultzmann, translated by Arthur Cooper.)

There is a very strong probability against any narrowing of the cervical canal, short of actual occlusion, being the cause of sterility in any particular case. This is shown by Pallen's statistics, quoted by Galabin, where in 337 cases of sterility the cervix was incised, only 13 or 14, however, becoming pregnant afterwards, which, as Dr. Galabin says, is not a greater proportion than might be accounted for by coincidence. Still, a cervical canal dilated, say up to the size of No. 12 in Hegar's series of dilators, must, one would think, allow a larger proportion of semen to enter the uterus, than a canal that only admits the uterine sound. If nothing abnormal can be made out on careful examination, and the patient has been married three years, and is anxious to have her chance of having a family increased, dilatation of the cervix to the extent mentioned may be tried.

Treatment.—If the patient has not been married three years, we need only attend to the general health; till then, there is no reason for her to think she will have no children. If, however, there is pain on coitus, we endeavour to discover the cause, and treat it in a suitable manner. If the pain on coitus lead to secondary vaginismus, we may expect to cure the vaginismus by removing the cause. In primary vaginismus, *i.e.*, where we can discover no local cause, the prospect is unsatisfactory, though we may try the treatment recommended on page 139.

In face of the strong probability against narrowness of the cervical canal at any part being the cause of sterility in any given case, it is undesirable to resort to incision of the internal or external os. If the patient suffers from spasmodic dysmenorrhœa (which is common in cases of sterility), and there is nothing abnormal to be made out by physical examination, we may endeavour to cure the dysmenorrhœa by dilating the cervix as recommended when discussing dysmenorrhœa.

#### APPENDIX A.

Note on the Systematic Treatment of Nerve Prostration and Hysteria.—The Weir Mitchell Treatment.

This plan of treatment was originally introduced by Dr. Weir Mitchell, of Philadelphia.

The cases for which it is suitable are those where careful examination discovers no organic disease of the nervous system, and where some one or more of the symptoms commonly known as hysterical are present. The patient is a "worn and wasted, often bedridden, woman, who has broken down either from some sudden shock, such as grief, or money losses, or excessive mental or bodily strain. At first, perhaps, there may have been only a debility, constantly, however, on the increase, daily more and more yielded to, until at last all power of effort is lost, fostered too often by injudicious sympathy, and the constant nursing of devoted relatives and friends. Coincident with this is the total loss of appetite, the profound anæmia, and the consequent wasting of the tissues, so characteristic of these cases. On the soil so prepared are often developed the graver protean forms of hysterical disease, such as paresis or paralysis, vomiting, disorder of motion, hystero-epilepsy, etc." \*

<sup>\*</sup> The Systematic Treatment of Nerve Prostration and Hysteria, by W. S. Playfair, M.D.

The chief points in the treatment are :-

- I. Removal of the patient from her surroundings to lodgings or a hospital for private patients, where she can be isolated, no one being allowed to see her but her doctor and nurse.
- 2. The patient at first is to be kept absolutely at rest in bed.
- 3. The use of mechanical tonics to produce muscular waste. This is done partly by systematic kneading and rubbing of the muscles by a trained nurse, and partly by making the muscles contract by applying the faradic current.

In this way the patient's power of assimilating food is greatly increased.

4. Systematic over-feeding.

It is essential that the patient should be removed from her surroundings; any attempt to carry out the treatment at the patient's own home is likely to end in failure.

Another point of importance is the selection of a suitable nurse. She must be capable of exerting a moral control over the patient; and yet, on the other hand, she must be kind, and able to make herself an agreeable companion. It may be necessary to change the nurse, if in a few days the case is not going on satisfactorily; and in this way success has been the result where failure appeared imminent.

For three or four days the patient is kept on a milk diet alone. Then massage is commenced, at first for about twenty minutes, afterwards for an hour and a half, night and morning. The faradic current is also used for from ten to twenty minutes, night and morning. By about the tenth day the patient is taking about two quarts of milk, and three full meals in addition, as in the following diet sheet taken from Dr. Playfair's book.

Breakfast, consisting of a plate of porridge and cream, fish or bacon, toast and tea, coffee or cocoa.

Lunch, at one p.m., of fish, cutlets or joints, and a sweet, such as stewed fruit and cream, or a milk pudding.

Dinner, at seven p.m., consisting of soup, fish, joint, and sweets; and, in addition, a cup of raw meat soup at seven a.m. and eleven p.m.

Dr. Playfair finds professional rubbers unsatisfactory. He prefers to choose a strong, intelligent woman, who knows nothing of massage, and train her according to the directions given in Dr. Weir Mitchell's book.

The treatment lasts from six to eight weeks.

#### APPENDIX B.

Notes of Seven Cases in which the Cervix was rapidly dilated by means of Hegar's Dilators.

CASE I.-E. S., aged 39, married sixteen years, seven children. Admitted into the London Hospital complaining of losing a large quantity of blood on and off ever since a miscarriage she had five months previously, when about three months pregnant. On admission it was found that some blood was escaping from the external os, and that the uterus was enlarged. The sound passed three inches and a half. Four days later, under ether, the cervix was rapidly dilated; the finger passed into the uterine cavity came on several irregular prominences. These were removed as completely as possible. They looked like pieces of placenta. Pure carbolic acid was applied to the interior of the uterus. The temperature after the dilatation was normal for four days. On the evening of the fifth day there was a rise to 101'4°; the next day the temperature was again normal, and continued so. No cause for the rise on that one day was discovered, the patient's general condition being perfectly satisfactory otherwise. Three months later she came to see me. Since leaving the hospital she had menstruated regularly, the periods lasting four days only, and recurring at the proper time.

Case II.—I. L., aged 40, married twenty years, six children, the last nine years ago, four miscarriages, the last before the birth of her fifth child. Admitted into the hospital complaining principally that for the previous six months her periods had occurred every fortnight; before that she had been quite regular. No cause for the bleeding was discovered on examination, and accordingly the cervix was dilated rapidly under ether. Nothing, however, was found in the cavity of the uterus to account for the bleeding, and it seemed probable that the menorrhagia was due to the menopause being about to occur earlier

than usual. In this case the temperature after dilatation of the cervix never rose above normal.

CASE III.-M. W., aged 35, married sixteen years, twins a year after marriage, two miscarriages, the last eleven years and a half ago. Admitted to the hospital on account of menorrhagia of fifteen months' duration. She had been under treatment in the out-patient department, but had obtained no relief. On examination the uterus was found to be enlarged, and the sound passed four inches. The uterus was freely movable. The vaginal portion of the cervix was healthy. The cervix was rapidly dilated under ether. On passing the finger into the body of the uterus, soft irregular projections, particularly extensive on the right side, were met with. These were thoroughly scraped away with a spoon, and tincture of iodine applied to the interior of the uterus: portions of the substance removed, on microscopical examination afterwards, showing a glandular structure; sections showed very large irregular cavities lined with columnar epithelium in a single layer. The temperature after dilatation remained normal. I have seen this patient at intervals since; she is now regular, and does not lose too much.

CASE IV.—E. B., aged 45, married twenty-three years ago, three children, the last sixteen years ago, three miscarriages. Admitted chiefly on account of having lost too much at her periods for the last eleven months. During this time the flow once lasted nine weeks, and once five weeks, and often a fortnight. On examination there was found an erosion of the anterior lip of the cervix; for a time it was thought doubtful whether it was not commencing malignant disease. Under local treatment, however, the erosion improved rapidly, but the menorrhagia was still as bad as ever; the cervix was therefore rapidly dilated. A little sessile lump, the size of a split pea, was found in the cavity of the uterus near the point of entrance of the right Fallopian tube; it did not admit of removal. The temperature after dilatation remained normal in this case also. This was probably one of the menorrhagic cases commonly met with, apart from local disease, about the time of the menopause.

CASE V.—H. H., aged 37, married fifteen years, no children, one miscarriage ten years ago. Admitted complaining of having lost too much at her periods for the last six years, sometimes bleeding for three weeks at a time. The cervix was rapidly dilated till the finger could be passed into the uterine cavity. Nothing abnormal was detected. In this case it is worth noticing that the time occupied in

dilating the cervix was only twenty minutes. The patient had had no children, as the external os was small. The temperature on the second day was 100.8° in the evening, but except on that occasion was normal throughout.

CASE VI.-E. M., aged 39, married eight years, had one child when she was fifteen years old, none since. Admitted into the hospital complaining of having had a red vaginal discharge for three months. Three months previously she had one of her ordinary menstrual periods; but instead of this passing off in the usual way, she continued to have a red discharge till she came to the hospital. In the outpatient department no cause was found for the hæmorrhage, and she was therefore taken in, that further investigation might be made. When the anterior lip of the cervix was seized with a volsella preparatory to dilatation, the lip was necessarily somewhat everted, and it was at once evident that there was malignant disease within the cervical canal that had not extended beyond the external os. The cervix was rapidly dilated, and it was found that there was extensive malignant disease in the cervical canal. The cavity of the body of the uterus was healthy. As here we are only concerned with the case so far as it was affected by dilatation of the cervix, I need only say that on the evening of the day when dilatation was performed the temperature was 100'2°; but after that it remained normal.

In the preceding six cases dilatation of the cervix was performed in the non-pregnant condition; in the following case it was done at the sixth month of pregnancy for the purpose of terminating the pregnancy, on account of persistent hæmorrhage that was seriously affecting the patient:—

CASE VII.—Mrs. F., aged 37, married thirteen years, had had four children, the last five years ago, no miscarriages. She was brought to me by Dr. Cockell, of Dalston, complaining of attacks of pain from time to time across the lower part of the abdomen and back, and of irregular hæmorrhages. Three months before I saw her she had an attack of bleeding lasting five weeks; then she went a month free from bleeding; then it came on again, small clots being passed. The bleeding continued till the time I saw her. The uterus was about the size it reaches at the sixth month of pregnancy. The patient was

a good deal pulled down by the bleeding, and it was clearly desirable that the uterus should be emptied as soon as possible. Accordingly the cervix was dilated with Hegar's dilators. As soon as it was sufficiently open to admit my finger, I felt the placenta attached right over the internal os; in fact, it turned out to be a case of placenta prævia. I separated the placenta all round as far as I could reach, and then ruptured the membranes, brought down a leg, and so delivered. The whole period occupied, from the time she began to take the anæsthetic till the uterus was emptied, was one hour and twenty minutes. Strict antiseptic precautions were used, the vagina being washed out with iodine water before beginning, and the uterus washed out afterwards with the same antiseptic. Two or three injections of ergotine were given hypodermically during the operation. Dr. Cockell wrote to me some time after to say that the case had done extremely well, and that the temperature had never risen above normal.

Note.—The date of this case (No. VII.) was June 1886. Since that time, eleven years ago, I have ceased to rely exclusively on the rapid method of dilatation. For further information the reader is referred to page 32.

I have never had any bad result from rapid dilatation in my own practice.

Still, fatal consequences are not unknown, and have probably been due to deep lacerations of the cervix in the neighbourhood of the internal os. The action of a dilator is twofold: when merely pushed forward, it acts as a wedge; but when a boring movement is imparted to it, it acts as a lever as well—the fulcrum being at one side of the cervix, and the weight (or force actually being exerted on the cervix) a little in advance of the fulcrum, and at a point opposite to it. It is important, therefore, not to use a long instrument, because the force acting on the cervix is in direct proportion to the distance of the handle from the fulcrum.

## INDEX.

Abdominal belt, 442
— examination, 5
— pressure, 146
Abortion, as a cause of bleeding,
55, 75
Abscess of the labium, 97
—— pelvic, in perimetritis, 324
——— in parametritis, 334
— in hæmatocele, 352
———— in salpingitis, 305
Absence of ovaries, 50
of uterus, congenital, 50
Adhesion of the labia in infants,
106
Adhesions in the causation of dys-
menorrhœa, 77, 81
in ovariotomy, treatment of,
431
Age, in regard to bleeding, 69, 75
—— carcinoma, 236, 273
———— fibroids, 183
vaginitis, 132
Alcoholism as a cause of vaginitis,
132
Alexander's operation, 154
Amenorrhœa, 47
— apparent, 47
in superinvolution, 181
Amputation of the cervix, 250
Ani, sphincter, rupture of perineum
involving, 115, 119, 126
Anteflexion of uterus, 449, 457
5

```
Anteversion of uterus, 452, 459
Antiseptic precautions, importance
  of, in midwifery practice, 330; in
  operative work generally, 41
Apostoli treatment for fibroids,
  223
Arbor Vitæ, 176
Areola, primary, 7
- secondary, 7
Ascites, diagnosis of, from ovarian
  tumour, 409
Aspiration in cases of pelvic ab-
  scess, 228
--- in suppurative inflammation
  of the knee-joint, 342
Atresia (occlusion) of the vulva, 103,
  489
--- of the vagina, 496, 497
— of the uterus, 497
- of the genital canal, sym-
 ptoms of, 104
 --- --- treatment of, 105,
  497
Auscultation, 12
Bandl, ring of, 168
Barnes's crescent speculum, 24
- tent-introducer, 42
Bartholin, the glands of, diseases
  affecting, 97
```

Baths in amenorrhœa, 51

- in vulvitis, 95

Bimanual examination, 19 Bivalve speculum, 25 Bladder, exploration of, 477  — importance of emptying for the bimanual examination, 20  — inflammation of, 473, 477  — irritability of, 4, 472  — prolapse of, 144  — separation of, from cervix, in the supra-vaginal amputation, 250  Bleeding, 55  — in connection with pregnancy, 55, 75  — not distinctly connected with pregnancy, 56, 69, 75
Bivalve speculum, 25 Bladder, exploration of, 477  — importance of emptying for the bimanual examination, 20  — inflammation of, 473, 477  — irritability of, 4, 472  — prolapse of, 144  — separation of, from cervix, in the supra-vaginal amputation, 250  Bleeding, 55  — in connection with pregnancy, 55, 75  — not distinctly connected with pregnancy, 56, 69, 75
Bladder, exploration of, 477  — importance of emptying for the bimanual examination, 20  — inflammation of, 473, 477  — irritability of, 4, 472  — prolapse of, 144  — separation of, from cervix, in the supra-vaginal amputation, 250  Bleeding, 55  — in connection with pregnancy, 55, 75  — not distinctly connected with pregnancy, 56, 69, 75
<ul> <li>importance of emptying for the bimanual examination, 20</li> <li>inflammation of, 473, 477</li> <li>irritability of, 4, 472</li> <li>prolapse of, 144</li> <li>separation of, from cervix, in the supra-vaginal amputation, 250</li> <li>Bleeding, 55</li> <li>in connection with pregnancy, 55, 75</li> <li>not distinctly connected with pregnancy, 56, 69, 75</li> </ul>
the bimanual examination, 20  — inflammation of, 473, 477  — irritability of, 4, 472  — prolapse of, 144  — separation of, from cervix, in the supra-vaginal amputation, 250  Bleeding, 55  — in connection with pregnancy, 55, 75  — not distinctly connected with pregnancy, 56, 69, 75
<ul> <li>inflammation of, 473, 477</li> <li>irritability of, 4, 472</li> <li>prolapse of, 144</li> <li>separation of, from cervix, in the supra-vaginal amputation, 250</li> <li>Bleeding, 55</li> <li>in connection with pregnancy, 55, 75</li> <li>not distinctly connected with pregnancy, 56, 69, 75</li> </ul>
<ul> <li>irritability of, 4, 472</li> <li>prolapse of, 144</li> <li>separation of, from cervix, in the supra-vaginal amputation, 250</li> <li>Bleeding, 55</li> <li>in connection with pregnancy, 55, 75</li> <li>not distinctly connected with pregnancy, 56, 69, 75</li> </ul>
— prolapse of, 144 — separation of, from cervix, in the supra-vaginal amputation, 250 Bleeding, 55 — in connection with pregnancy, 55, 75 — not distinctly connected with pregnancy, 56, 69, 75
<ul> <li>separation of, from cervix, in the supra-vaginal amputation, 250</li> <li>Bleeding, 55</li> <li>in connection with pregnancy, 55, 75</li> <li>not distinctly connected with pregnancy, 56, 69, 75</li> </ul>
the supra-vaginal amputation, 250 Bleeding, 55 —— in connection with pregnancy, 55, 75 —— not distinctly connected with pregnancy, 56, 69, 75
Bleeding, 55  — in connection with pregnancy, 55, 75  — not distinctly connected with pregnancy, 56, 69, 75
Bleeding, 55  — in connection with pregnancy, 55, 75  — not distinctly connected with pregnancy, 56, 69, 75
<ul> <li>in connection with pregnancy,</li> <li>55, 75</li> <li>not distinctly connected with pregnancy, 56, 69, 75</li> </ul>
55, 75 — not distinctly connected with pregnancy, 56, 69, 75
— not distinctly connected with pregnancy, 56, 69, 75
pregnancy, 56, 69, 75
- differential diagnosis of various
affections causing, 72
treatment of, 73
Blood-mole, 67
Blunt hooks, large, in ovariotomy,
422, 434
Bougies, for dilating the cervix in dysmenorrhœa, 86
Braided silk for ovariotomy, 425
Breasts, condition of, to be noted
during examination of the abdo-
men, 6
Broad ligament, anatomy of, 297,
447
inflammation of connective
tissue in, 333
— phlegmon of, 335
0.1:0 ( 0 0 1
Calcification of fibroid tumours,
187
Cancer (see Carcinoma)
Cannabis Indica in dysmenorrhœa,
86
Carbolic acid in cervical catarrh,
165
in corporeal endometritis, 173

Carbolic acid in pruritus of the vulva, 113 Carcinoma of the bladder, 477 - of the body of the uterus, 273 - of the cervix, 236 - of the external genitals, 110 - occurring in fibroid tumours, 187 - of the kidney, 485 --- of the ovaries, 385, 388, 415 - of the vagina, 140 - causing pruritus, 112 Caruncle, vascular, of urethra, 96 Carunculæ myrtiformes, 14 Case, the mode of investigating a, I Castoreum, in dysmenorrhœa, 86 Catarrh of the cervix, 158 Catheter, precautions in using, 31 Cauliflower excrescence, 237 Caustics in carcinoma, 270 Cautery, in advanced carcinoma, 270 - in recurrent carcinoma, 259 Cellulitis, pelvic, 333 --- etiology, 333 - morbid anatomy, 334 --- producing anteflexion, 458 - in utero-sacral ligaments, 340, 458 --- symptoms and signs, 338 - treatment of, 341 - varieties of, 334 Central rupture of the perineum, Cervical endometritis, 158 Cervix uteri, amputation of, 250 --- anatomy of, 174, 251 - atresia (occlusion) of, in old age, 172 - atrophy of, in old age, 18 - carcinoma of, 236 — catarrh of, 158

Cervix uteri, dilatation of, 32	Decidua, passage of a, in mem-
glands of, 161	branous dysmenorrhœa, 84
— hypertrophy of, 18	Deciduoma malignum, 287
——————————————————————————————————————	Dermoid cysts, 390
— mucous membrane of,	Diabetes as a cause of pruritus, 112
161, 175	— of vaginitis, 132
— papilloma of, 246	of vulvitis, 94
—— position of, 18, 450	Diet after ovariotomy, 440
relation of ureters to,	- in the treatment of nerve-
251	prostration, 508
supra-vaginal amputation	Dilatation of the cervix, 32
of, 250	— for diagnostic purposes, 73
— vaginal portion of, 254	— for dysmenorrhœa, 86
Chloride of zinc in carcinoma, 270	- of the vaginal orifice for va-
Clover's crutch, 122	ginismus, 139
	— rapid method, 35
Cocaine for pruritus, 113	— slow method, 38
Colporrhaphy, 152	Dilators, Hegar's, 33
Corporeal endometritis, 166	Diphtheritic vaginitis, 132
Crescent speculum, Barnes's, 24	
Cribriform hymen, 103	— vulvitis, 93
Cup and stem pessary, 151	Direction of the external os uteri,
Curette, 169	normally, 18
— in carcinoma, 269	— in retroversion, 454
- in fibrinous and placental	Discharge in cervical endometritis,
polypi, 235	159
— in fungous endometritis, 168	fœtid, in carcinoma of the
— in mucous polypi, 233	cervix, 244
- in subinvolution of the uterus,	in carcinoma of the
168, 180	body, 274
Cystic degeneration of fibroids, 187,	- in sloughing fibroid po-
200	lypi, 204
	in corporeal endometri-
Cystitis, 473, 477	tis, 167, 168, 170
Cystocele, 144	Displacements of the uterus, 144
Cysts, ovarian, 385	156, 449
— parovarian, 448	caused by hæmatocele, 351
— pudendal, 97	— ovarian tumours, 406
—— vaginal, 140	
	parametritis, 338
	———— perimetritis, 326
Dangers of tents, 34	— diagnosis, 147, 452
Decidua, passage of a, in extra-	- relative importance of, 459
uterine pregnancy, 59, 356	460

Displacements of the vaginal walls,
144, 156
Distention of the Fallopian tubes,
299
— of the uterus, 104, 172, 497
of the vagina, 104, 496
Douche, the hydrostatic, 137
Douglas's pouch, incarceration of
the uterus in, 457
— opening in supra-vaginal
amputation of the cervix, 252
Drainage in ovariotomy, 438
— in pelvic abscess, 341
Duckbill speculum, 23
Dysmenorrhœa, 76
— in cases of fibroids, 78, 189
— inflammatory or congestive, 79
— spasmodic, 78
theories of the causation of, 79
— treatment of, 85
Dyspareunia, in vascular caruncle,
97
various causes of, 138
— various causes of, 138
— various causes of, 138  Écraseur, use of, in certain cases of
— various causes of, 138  Écraseur, use of, in certain cases of supra-vaginal amputation, 254
— various causes of, 138  Écraseur, use of, in certain cases of supra-vaginal amputation, 254 — for fibroid polypi, 206
<ul> <li>— various causes of, 138</li> <li>Écraseur, use of, in certain cases of supra-vaginal amputation, 254</li> <li>— for fibroid polypi, 206</li> <li>Electricity, use of, in certain painful</li> </ul>
<ul> <li>— various causes of, 138</li> <li>Écraseur, use of, in certain cases of supra-vaginal amputation, 254</li> <li>— for fibroid polypi, 206</li> <li>Electricity, use of, in certain painful conditions, 227</li> </ul>
<ul> <li>— various causes of, 138</li> <li>Écraseur, use of, in certain cases of supra-vaginal amputation, 254</li> <li>— for fibroid polypi, 206</li> <li>Electricity, use of, in certain painful conditions, 227</li> <li>Electrolysis for fibroid tumours, 223</li> </ul>
<ul> <li>— various causes of, 138</li> <li>Écraseur, use of, in certain cases of supra-vaginal amputation, 254</li> <li>— for fibroid polypi, 206</li> <li>Electricity, use of, in certain painful conditions, 227</li> <li>Electrolysis for fibroid tumours, 223</li> <li>Elephantiasis of the vulva, 107</li> </ul>
<ul> <li>— various causes of, 138</li> <li>Écraseur, use of, in certain cases of supra-vaginal amputation, 254</li> <li>— for fibroid polypi, 206</li> <li>Electricity, use of, in certain painful conditions, 227</li> <li>Electrolysis for fibroid tumours, 223</li> <li>Elephantiasis of the vulva, 107</li> <li>Endometritis, cervical, 158</li> </ul>
<ul> <li>— various causes of, 138</li> <li>Écraseur, use of, in certain cases of supra-vaginal amputation, 254</li> <li>— for fibroid polypi, 206</li> <li>Electricity, use of, in certain painful conditions, 227</li> <li>Electrolysis for fibroid tumours, 223</li> <li>Elephantiasis of the vulva, 107</li> <li>Endometritis, cervical, 158</li> <li>— diagnosis and symptoms,</li> </ul>
<ul> <li>— various causes of, 138</li> <li>Écraseur, use of, in certain cases of supra-vaginal amputation, 254</li> <li>— for fibroid polypi, 206</li> <li>Electricity, use of, in certain painful conditions, 227</li> <li>Electrolysis for fibroid tumours, 223</li> <li>Elephantiasis of the vulva, 107</li> <li>Endometritis, cervical, 158</li> <li>— diagnosis and symptoms, 159</li> </ul>
Ecraseur, use of, in certain cases of supra-vaginal amputation, 254 — for fibroid polypi, 206 Electricity, use of, in certain painful conditions, 227 Electrolysis for fibroid tumours, 223 Elephantiasis of the vulva, 107 Endometritis, cervical, 158 — diagnosis and symptoms, 159 — pathology, 161
<ul> <li>— various causes of, 138</li> <li>Écraseur, use of, in certain cases of supra-vaginal amputation, 254</li> <li>— for fibroid polypi, 206</li> <li>Electricity, use of, in certain painful conditions, 227</li> <li>Electrolysis for fibroid tumours, 223</li> <li>Elephantiasis of the vulva, 107</li> <li>Endometritis, cervical, 158</li> <li>— diagnosis and symptoms, 159</li> <li>— pathology, 161</li> <li>— treatment, 164</li> </ul>
<ul> <li>— various causes of, 138</li> <li>Écraseur, use of, in certain cases of supra-vaginal amputation, 254</li> <li>— for fibroid polypi, 206</li> <li>Electricity, use of, in certain painful conditions, 227</li> <li>Electrolysis for fibroid tumours, 223</li> <li>Elephantiasis of the vulva, 107</li> <li>Endometritis, cervical, 158</li> <li>— diagnosis and symptoms, 159</li> <li>— pathology, 161</li> <li>— treatment, 164</li> <li>— corporeal, 166</li> </ul>
<ul> <li>— various causes of, 138</li> <li>Écraseur, use of, in certain cases of supra-vaginal amputation, 254</li> <li>— for fibroid polypi, 206</li> <li>Electricity, use of, in certain painful conditions, 227</li> <li>Electrolysis for fibroid tumours, 223</li> <li>Elephantiasis of the vulva, 107</li> <li>Endometritis, cervical, 158</li> <li>— diagnosis and symptoms, 159</li> <li>— pathology, 161</li> <li>— treatment, 164</li> <li>— corporeal, 166</li> <li>— as a cause of bleeding,</li> </ul>
Ecraseur, use of, in certain cases of supra-vaginal amputation, 254 — for fibroid polypi, 206 Electricity, use of, in certain painful conditions, 227 Electrolysis for fibroid tumours, 223 Elephantiasis of the vulva, 107 Endometritis, cervical, 158 — diagnosis and symptoms, 159 — pathology, 161 — treatment, 164 — corporeal, 166 — as a cause of bleeding, 71, 75
Ecraseur, use of, in certain cases of supra-vaginal amputation, 254 — for fibroid polypi, 206 Electricity, use of, in certain painful conditions, 227 Electrolysis for fibroid tumours, 223 Elephantiasis of the vulva, 107 Endometritis, cervical, 158 — diagnosis and symptoms, 159 — pathology, 161 — treatment, 164 — corporeal, 166 — as a cause of bleeding, 71, 75 — diagnosis of, 166
Ecraseur, use of, in certain cases of supra-vaginal amputation, 254 — for fibroid polypi, 206 Electricity, use of, in certain painful conditions, 227 Electrolysis for fibroid tumours, 223 Elephantiasis of the vulva, 107 Endometritis, cervical, 158 — diagnosis and symptoms, 159 — pathology, 161 — treatment, 164 — corporeal, 166 — as a cause of bleeding, 71, 75

Endometritis, corporeal, senile, 172 ---- traumatic, case of, 170 — treatment, 173 Enucleation of fibroid tumours per vaginam, 209 - after laparotomy, 210 —— spontaneous, 187 --- of ovarian tumours, 437 Epoch as a cause of bleeding, 69, Ergot and ergotine, for fibroid tumours, 209 —— for subinvolution, 180 - use of empirically where no local organic disease, 74 Erosion of the cervix, 160 — treatment, 164 - varieties of, 161 Esthiomene, 107 Eversion of cervical mucous membrane, 163 Examination, the bimanual, 19 Exploration of the bladder, 477 External genitals, anatomy of, 91, 92, 93 --- os uteri, direction of, 18 Extirpation of the uterus for fibroids, 210-222 - for carcinoma of the cervix, 249 ---- for carcinoma of the body, 278 - for deciduoma malignum, 293 Extra-uterine gestation, 356 - as a cause of bleeding, 56, 59 - as a cause of hæmatocele, 345 - diagnosis of ovarian cysts from, 403, 412

Exudation in the pelvis, in perime-
tritis, 324, 326
— in parametritis, 336
Fallonian tubes anatomy con
Fallopian tubes, anatomy, 297
catheterisation of, 312
congenital peculiarities,
296
disease of, in relation to
dysmenorrhœa, 81, 87, 88
in relation to sterility, 505
— distention of, 299
— divisions of, 297
frequency of pathological
conditions of, 301
inflammation of, 296
Fergusson's speculum, 22
Fibrinous polypus, 69, 234
Fibro-cystic tumours of uterus, 188,
200
Fibroid tumours of the uterus, 183
of the ovary, 384
of the vagina, 140
- of the vulva, III
Fimbriated extremity of Fallopian
tube, 297
Fissures of the vulva, 105
Fleshy mole, 67
Flexions of the uterus, 449
Fœtal heart sounds, 12
Follicular erosion of cervix, 162
Foreign bodies in the vagina as a
cause of bleeding, 70
Fossa navicularis, 91, 93, 115
Fourchette, 91, 93
Friction sounds over abdominal
tumours, 406
Fungous endometritis, 167
Funic souffle, 13
Galvanometer, 224

Gangrene of the vulva, III

```
Glands, Bartholin's, diseases of, 97,
Glycerine plug, 333
Gonorrhœa, causing cystitis, 473
- Klein on the diagnosis of, 134
Gonorrhœal endometritis, 158, 166
--- peritonitis, 322
- salpingitis, 299
- urethritis, 136
-- vaginitis, 136
- vulvitis, 94
Graafian follicles, dropsical disten-
  tion of, 385
Granular vaginitis, 135
Hæmatocele, pelvic, 345
---- morbid anatomy of, 347
- physical signs, 351
---- symptoms of, 349
- treatment of, 352
- and ovarian tumour, 401
Hæmatokolpos, 104, 496
Hæmatoma (or thrombus) of the
  labium, 106
Hæmatometra, 104, 497
Hæmatosalpinx, 301
Hæmorrhage, classification of con-
ditions causing, 55, 75
---- prevention of, in supra-vaginal
  amputation of cervix, 253
--- rules of treatment in cases of
  uterine hæmorrhage, 73
- into an ovarian cyst, 392
Hair in dermoid cysts, 390
Hegar's dilators, 33, 87, 477
- sign of pregnacy, 194
Heredity, influence of, in carcinoma,
Hernia into the vulva, 100
--- of the ovary, 446
History, the, in case taking, I
Hodge's pessary, 465, 466
```

Hot douches, in cases of hæmorrhage, 73, 209 ---- in chronic peritonitis, 333 - in subinvolution, 180 Hydatid tumours, diagnosis of, 412 Hydatidiform mole, 60 Hydramnios, 407 Hydronephrosis, 411 Hydrosalpinx, 301, 313 Hydrostatic douche, 137 Hymen, atresia of, 103, 504 --- forms of, 103 - imperforate, 103, 504 - inflamed portions of, in vaginismu, 105 Hypertrophy of the external genitals, 107 - of the mucous membrane of the uterus in endometritis, 168 Hysterectomy, for fibroids, 210 —— for carcinoma, 249, 278 Hysteria, treatment of, by the Weir Mitchell system, 507 Hysterical retention of urine, 476 Hysteropexy, 155

Ice-bag, after ovariotomy, 441

— in hæmatocele, 353
Imperfectly developed uterus, 490

— associated with dysmenorrhæa, 78
Imperforate hymen, 103, 504
Incomplete abortion, 58
Incontinence of fæces and flatus in complete rupture of perineum, 121

— of urine, 472

— as a cause of vulvitis, 94
India-rubber ring pessary, 150
Inevitable laceration of perineum, 115

Infantile uterus, 495 Infundibulo-pelvic ligament, 297 Inguinal glands in carcinoma of the vulva, III Injections, vaginal, 137 —— intra-uterine, 36 Inspection of abdomen, 5 --- of abdomen in cases of ovarian tumour, 404 - of breasts, 6 - of external genitals, 13 Interstitial inflammation of uterus, - as a cause of dysmenorrhœa, 80 Intestinal obstruction in cases of fibroid tumours, 189 Intestines, adhesions of cyst to, in ovariotomy, 431 Intra-abdominal pressure in prolapse, 146, 156 Intra-uterine electrode in the Apostoli treatment of fibroids, 223 Inversion of uterus, 206, 468 Involution of uterus, 174 - importance of securing prior to operations on the cervix, 264 Iodine in cervical catarrh, 165 - in corporeal endometritis, 167, 173 --- paint to abdomen in chronic peritonitis, 333 - water as an antiseptic injection, 36, 272 - for the hands before operations, 43, 428 Iodoform, cases of supposed poisoning by, 229 Irritability of the bladder, 472 Irritating discharges as a cause of vulvitis, 94 Isthmus of Fallopian tube, 297

Keith's drainage-tube for ovariotomy, 438
Kidney, scrofulous disease of, 473,
479, 484
— tumours of the, diagnosis from
ovarian tumours, 411
Kidneys, palpation of, 8
— malignant disease of, 485
Klein on the diagnosis of gonorrhœa, 134

Labium, abscess of, 97 --- cyst of, 97 - thrombus of, 106 Labour in the etiology of inflammation of the pelvic organs :---- of cervical endometritis, 158 ---- of corporeal endometritis, 166 --- of pelvic cellulitis, 334 — of pelvic peritonitis, 322 --- of salpingitis, 299 --- of vaginitis, 132 Laceration of cervix, 163 Lactation, importance of, 180 - protracted, as a cause of menorrhagia, 70 Laminaria tents, 37 Leeches in pelvic peritonitis, 331 Leucorrhœa, 159 Ligaments, the broad ligaments, 297 --- the ovarian ligament, 297 - the round ligaments, 297 Ligature, in ovariotomy, 431 - in the supra-vaginal amputation of the cervix, 253 Lineæ albicantes, 6 Lymph, accumulation of, in Douglas's pouch, 326

Malformations of uterus, vagina, and vulva, 488 Malignant disease, as a cause of bleeding, 69, 75 - of the body of the uterus, 273, 284, 287 - of the cervix, 236 --- of the ovary, 385 - of the peritoneum, 413 - of the vagina, 140 - of the vulva, 110 Massage in hysteria, 507 - local, in chronic pelvic inflammation, 88 Medication, intra-uterine, 164, 167 Membranous dysmenorrhœa, 83 Menopause, artificial production of, - bleeding occurring at, 69 - bleeding occurring years after, 69 Menorrhagia and metrorrhagia, 55 Menstruation, 45 — abnormalities of, 47 - during pregnancy, 57 Metritis, 173 Microscopic appearances of carcinoma of the cervix, 241 — of carcinomatous ovarian tumour, 391 -- of erosion of the cervix, --- of granulations in endometritis, 172 — of mucous polypi, 231 - of ovarian fluids, 390 Micturition, disorders of, 472 - in cystitis, 473 - in fibroid tumour, 189 — in hysteria, 476 - in malignant disease of the kidney, 485

Micturition, in ovarian tumour, 393
— in pelvic inflammation, 324
— in prolapse, 147
— in retroversion of the gravid
uterus, 474
— in scrofulous kidney, 479, 484
— in vaginitis, 133
in vascular caruncle, 97
Missed abortion, 56
Mobility of the uterus, 21
Mode of introducing tents, 40
of investigating a case, I
— of passing the finger into the
vagina, 16
Molar pregnancy, 60
Mortality of hysterectomy, for fib-
roids, 212
— of extirpation of the uterus for
cancer, 249, 278
— of ovariotomy, 442
of supra-vaginal amputation of
the cervix, 255
Mucous membrane of cervix, 174
—— polypi, 231
as a cause of bleeding, 69
Mucous tubercles, 102
Müller, ducts of, 490
Multiparous uterus, 176
Myoma of the ovary, 384
— of the uterus, 183
— of the dierus, 103
Nahathi amla sea
Nabothi, ovula, 160
Needle, straight in handle, for
perineorrhaphy, 118
Needle-holder, 124, 125
Needles, Hagedorn's, 125
— for ovariotomy, 423
Nelaton's cyst forceps, 423
Neuralgic dysmenorrhœa, 76
— treatment of, 87
Nitrate of silver, 96
Nück, canal of, 100
2. 10.1., 04.141 01, 100

Nulliparous uterus, 174 Nymphæ, 91, 107

Obstructive dysmenorrhœa, 85 Old age as a cause of vaginitis, 132 Oöphorectomy for dysmenorrhœa, 87 - for fibroids, 210 - for inflammation of the uterine appendages, 308 Os externum and dysmenorrhœa, - in multiparous uterus, 176 - in virgin uterus, 176 Os internum, in regard to passing the sound, 28 Ovarian artery, 214, 460 --- corpuscles of Drysdale, 390 - fimbria, 297 - ligament, 297 --- tumours, 384 - as a cause of bleeding, 70, 396 —— —— diagnosis, 400 —— —— prognosis, 399 ---- structure, 385 ---- symptoms, 392 ---- treatment, 416 ---- varieties, 388 Ovaries, absence of, 50 - adhesions of, to surrounding structures, 81, 298, 442 --- carcinoma of, 385 --- cysts of, 385 --- dermoid cysts of, 390 - development of, imperfect, 50 - displacements of, 445 - fibroids of, 384 ---- hernia of, 446 - inflammation of, 442 --- position of, 445

Ovaries, prolapse of, 446

— proliferous cysts of, 388

— sarcoma of, 385

Ovariotomy, 417

Ovaritis, 442

Over-lactation as a cause of bleeding, 70

Ovula Nabothi, 160

Ovulation and menstruation, 46

Pain in carcinoma of the cervix, 245 - of the body of the uterus, --- ovarian tumours, 393 Palpation of the abdomen, 8 Papillary erosion, 161 Papilloma of the cervix, 246 Papillomata of the vulva, 100 Parametritis, 333 Parovarian cysts, 448 Pedicle, treatment of, in ovariotomy, 43I Pelvic abscess, 227, 301, 324, 337 --- cellulitis, 333 - inflammation, 322 --- as a cause of displacements, 457 - as a cause of dysmenorrhœa, 77 - as a cause of sterility, - hæmatocele, 345 --- peritonitis, 322 Percussion, 11 Perimetritis, 322 Perineal body, 116 Perineorrhaphy, immediate, 116 - secondary, 120 Perineum, rupture of, 115 - central rupture of, 115 - relation of ruptured perineum to prolapse, 146

Peritoneum, relation of, to uterus and vagina, 175 ---- folds of, forming the broad ligaments, 297 forming the utero-sacral ligaments, 457 - opening in ovariotomy, 429 - opening Douglas's pouch in supra-vaginal amputation of the cervix, 252 - in procidentia uteri, 145 Peritonitis, 322 Pessary, cup and stem, 151 — Hodge's, 466 - india - rubber watch - spring, 150 --- general rules to be observed in using, 465 Phantom tumours, diagnosis, 408 Placenta, portions of retained, 58 — prævia, 58 Placental polypus, 68 Playfair's probe, 165 Polypi of the uterus, 230 Position of the uterus, the normal, 449 - for vaginal examination, 13 Posterior fornix, 21 Posture of patient in para- and perimetritis, 338, 325 Pouch of Douglas, descent of, in procidentia uteri, 145 - incarceration of uterus in, 457 - -- opening in operations on the cervix, 252 --- prolapse of ovary into, Pregnancy complicated with fibroid tumour, 193 --- complicated with ovarian tumour, 407

Pregnancy complicated with carcinoma of the cervix, 60, 262 - diagnosis of, from fibroid tumours, 190 — diagnosis of, from ovarian tumours, 407 --- menstruation during, 57 - as a cause of pruritus, 112 Present state, the, in case-taking, 1, 5 Procidentia uteri, 144 Prolapse of the ovary, 446 --- of the uterus, 144 - of the vaginal walls, 144 Proliferous cysts of the ovary, 388 Pruritus vulvæ, 112 Puberty, the age of, as a cause of menorrhagia, 69 - symptoms of atresia at, 104 Purse-string suture, 119 Pyelitis, 473 Pyometra, 172 Pyonephrosis, 473, 479 Pyosalpinx, 301

## Quarters of the pelvis, 22

Rapid dilatation of the cervix, 35
Recamier's curette, 169
Rectal examination, 30
Recti, separation of, 9
Rectocele, 144
Remote parametritis, 338
— perimetritis, 338
Renal tumours and ovarian, 411
Replacement of uterus with sound, 456
Repositor for inversion of the uterus, 469, 470
Retained menses (hæmatokolpos, hæmatometra), 104, 496

Retention of urine, due to fibroids, 189, 476 —— — due to hysteria, 476 - due to ovarian tumours, 475 - in retroversion of the gravid uterus, 474 Retroflexion, 454 Retro-peritoneal cysts, 400 Retro-uterine hæmatocele, 345 Retroversion, 453 — physiological, 449 - in prolapse, 145 Ring pessary, 150 Rodent (or corroding) ulcer of cervix, 246 Room for ovariotomy, 417 Round ligaments, shortening the, Rupture of the perineum, 115 - influence of in cases of prolapse, 146 Salpingitis, 296 - as a result of vaginitis, 136 Sarcoma of the body of the uterus, 284, 287 - of the cervix uteri, 284 — of the ovary, 385 —— of the vagina, 140, 285 Scissors bent on the flat for ovariotomy, 419 Secretion in the breasts, significance of, 7 Senile vaginitis, 132 Septic matter, introduced by the nails, 330 - introduced by the sound, 29 - - causing cellulitis, 334 - causing peritonitis, 322 330

524 INDEX.

Septicæmia from the use of tents, 34 Serous peritonitis, 324 Sickness after ovariotomy, 441 - in pelvic peritonitis, 331 Simon's sharp spoon, 270 Sims's speculum, 23 Size of the uterus, estimation of the, Skin-cracks, 6 Souffle, funic, 13 - uterine, 12 Sound, the uterine, 26 - in anteflexion, 453 - by no means always necessary, 26 - directions for passing, 28 — in fibroids, 196 - in inversion, 204 - in retroversion and retroflexion, 455, 456 - in subinvolution, 179 — in superinvolution, 181 Sounds heard over the pregnant uterus, 12 Speculum, Barnes's crescent, 24 — Bryant's rectal, 479 - Cusco's bivalve, 25 - Fergusson's, 22 —— Sims's, 23 Spencer Wells's pressure forceps, 419, 420 ---- trocar, 424 Sphincter ani, operation to restore, 121, 130 — rupture of, 115 Splenic tumours, 412 Sponge tent, 37, 170 Sponges for ovariotomy, 419 Spotty vaginitis, 135 Staffordshire knot, 432 Steriliser, 43

Sterility, 505 Sticks of zinc sulphate and alum for cervical catarrh, 165 Straight needle for perineorrhaphy, 118 Subinvolution of uterus, 174 — as a cause of bleeding, 68, 75, 178 —— case of, 168 Sulphate of copper for catarrh of the cervix, 164 Sulphocarbolate of zinc for vaginitis, 136 Superinvolution of the uterus, 181 Suppression of menstruation, 47 Suppuration in dilated Fallopian tubes, 301, 324 - in hæmatocele, 352 - in knee-joint complicating parametritis, 342 --- in ovarian tumours, 392 - in pelvic cellulitis, 337 --- peritonitis, 324 Supra-vaginal amputation of the cervix, 250 Sutures in ovariotomy, 423, 425 Syphilis of the vulva, 101, 102, 107 Syringe, Higginson's, 331

Tait's operation for rupture of the perineum, 127

— Staffordshire knot, 432

Tangle tent, 37

Tapping in dilated tubes, 306

— in ovarian tumours, 416

— in parovarian cysts, 448

— in purulent pelvic inflammation, 228

Teeth in dermoid cysts, 390

Temperature, high, after ovario-

tomy, treatment of, 441

Tenderness on palpating the abdomen, 9 Threadworms as a cause of vulvitis, — as a cause of pruritus, 112 Thrombosis complicating ovarian tumour, 398 Thrombus of the labium, 106 Tour de maître, 29 Transverse section of uterus, 461 Trocar for ovariotomy, 424 Tube for washing out the uterus, 36 Tubo-ovarian cyst, 304 Tumours of the ovary, 384 — of the uterus, 183, 230, 236, 273 - of the vagina, 140 Tupelo tent, 37 Type of menstruation, 45 Ulceration, so-called, of cervix, 160 - carcinomatous, of cervix, 239, 243, 259 --- corroding, of cervix, 246 - syphilitic, of cervix, 246 - of the pudendum, 90, 102 Ulcerous vaginitis, 135 Ulcers of the inverted vagina in procidentia, 148 Umbilical souffle, 13 Umbilicus, condition of, to be noted in examination of the abdomen, 5, 6

Uræmia in carcinoma, 248

cinoma, 243

mours, 397

Ureters, relation of, to cervix uteri,

- dilatation of, in cases of car-

- in cases of ovarian tu-

—— inspecting the orifices of, 478

—— catheterisation of, 478, 479

Urethra, malignant disease of, 110, ---- position of, 91, 92, 93 ---- stricture of, 472 - vascular caruncle of, 96, 472 Urethritis, 90, 133, 472 Urine, incontinence of, 472 ---- retention of, 472, 474, 476 Urogenital sinus, 489 Utero-sacral ligaments, 457 (and see Introduction) Uterine artery, 214, 215, 460 - appendages, removal of, for dysmenorrhœa, 87 \_\_\_\_ for fibroids, 210 --- contractions, in diagnosis, 10, 190, 407 --- forceps, 38 - souffle, 12 --- sound, 26 — tube, 36 Uterus, absent or rudimentary, 50, - amputation of, for fibroids, --- anatomy of, 174, 214, 251, 297, 460 ---- anteflexion of, 449, 456, 457 — anteversion of, 452, 456, 457 - arterial supply of, 214, 215, 460 — bicornis, 494 - carcinoma of body of, 273 --- carcinoma of cervix of, 236 - congenital atrophy of, 496 --- congestion of, not caused by flexion, 461 - corroding ulcer of, 246 --- curetting, cases illustrating, - extirpation of, per vaginam, 278

Uterus, fibroid tumours of, 183
— infantile, 495
inflammation of, 173
—— inversion of, 204, 468
malformations of, 490
mobility of, 21
peritoneum of, 175
— polypi of, 230
position of, the normal, 449
prolapse of, 144
retroflexion of, 451, 454
- retroversion of, 145, 452
sarcoma of, 284
septus, 492
shape of, in multipara, 176
in virgin, 174
— subinvolution of, 174
— superinvolution of, 181
— unicornis, 490
Vagina, absence of, 496
- atresia (occlusion) of, 496
— capacity of, 17
— carcinoma of, 140
- cicatrisation in, after labour,
505
— cysts of, 140
— diseases of, 132
— distention of, in atresia, 104
— fibroids of, 140
— glands of, 135
inflammation of, 132
— plugging in carcinoma, 269
— prolapse of, 144
sarcoma of, 140, 285
septum in, 496
Vaginal examination, 13
portion of cervix, 175, 254
Vaginismus, 138, 505
Vaginitis, 132
Varicose veins of the vulva, 106
El El

Vascular caruncle of urethra, 96 - as a cause of bleeding, 70 --- as a cause of vaginismus, 139 Venous sinuses round fibroid tumours, 191 - thrombosis, in carcinoma, 245, 261 — in ovarian tumours, 398 Versions of uterus, 452 Vesico-vaginal fistula, 473 Vesicular mole, 60 Vestibule, boundaries of, 91 - development of, 489 Villous endometritis, 168 - as a cause of bleeding, 71, 75 Volsella, 34 Vulva, abscess of, 97 --- anatomy of, 91, 92, 93 — atresia of, 103, 489 --- carcinoma of, 110 --- cysts of, 97 - elephantiasis of, 107 - fibrous tumour of, III —— fissure of, 105 --- hæmatoma of, 106 - inflammation of, 90 --- malformations of, 488 - pruritus of, 112 --- varicose veins of, 106 Vulvitis, 90

Warts of the vulva, 100
Washing out the uterus, tube for,
36
Weir Mitchell treatment for hysteria, 507
White's repositor for inversion,

# SELECTED LIST

OF

# NEW AND RECENT WORKS

PUBLISHED BY

# H. K. LEWIS,

# 136 GOWER STREET, LONDON, W.C.

(ESTABLISHED 1844).

\*\* For full list of works in Medicine and Surgery published by H. K. Lewis see complete Catalogue sent post free on application.

SYDNEY RINGER, M.D., F.R.S.

Holme Professor of Clinical Medicine in University College; Physician to University College Hospital,

AND

HARRINGTON SAINSBURY, M.D., F.R.C.P.

Physician to the Royal Free Hospital and the City of London Hospital for Diseases of the Chest, Victoria Park.

A HANDBOOK OF THERAPEUTICS. Thirteenth [Now ready.

H. ALDERSMITH, M.B. LOND., F.R.C.S. Medical Officer, Christ's Hospital, London.

RINGWORM AND ALOPECIA AREATA: Their Pathology, Diagnosis, and Treatment. Fourth Edition, enlarged, with new Illustrations, demy 8vo, 10s. 6d. [Now ready.

A. BROCA, M.D. Chirurgien des Hôpitaux des Paris, &c.

AND

F. LUBET-BARBON, M.D.

MASTOID ABSCESSES AND THEIR TREAT-MENT. Translated and edited by Henry J. Curtis, B.S. and M.D. (Lond.), F.R.C.S. (Eng.), Assistant to the Professor of Pathology, University College, London, &c. With coloured Illustrations, cr. 8vo, 6s.

[Now ready.]

E. W. GOODALL, M.D. LOND.

Medical Superintendent of the Eastern Hospital of the Metropolitan Asylums
Board; Formerly Medical Registrar to Guy's Hospital;

J. W. WASHBOURN, M.D. LOND., F.R.C.P.

Physician to the London Fever Hospital; Assistant Physician to Guy's Hospital, and Lecturer in the Medical School.

A MANUAL OF INFECTIOUS DISEASES.

Illustrated with Plates, Diagrams, and Charts, 8vo, 15s.

[Now ready.

A. C. ABBOTT, M.D.

Professor of Hygiene, and Director of the Laboratory of Hygiene, University

THE PRINCIPLES OF BACTERIOLOGY: A Practical Manual for Students and Physicians. Fourth Edition, with 106 Illustrations, 19 being coloured, post 8vo, 12s. 6d. nett. [ Fust ready.

JAMES ANDERSON, M.D., F.R.C.P.

Late Assistant Physician to the London Hospital, &c. NOTES ON MEDICAL NURSING; from the Lectures given to the Probationers at the London Hospital. Edited by E. F. LAMPORT, Associate of the Sanitary Institute. Third Edition, crown 8vo, 2s. 6d. [Now ready.

#### E. CRESSWELL BABER, M.B. LOND.

Surgeon to the Brighton and Sussex Throat and Ear Hospital.

GUIDE TO THE EXAMINATION OF THE NOSE WITH REMARKS ON THE DIAGNOSIS OF DIS-EASES OF THE NASAL CAVITIES. With Illustrations, small 8vo, 5s. 6d.

#### FANCOURT BARNES, M.D., M.R.C.P.

Physician to the Chelsea Hospital; Obstetric Physician to the Great Northern Hospital, &c.

GERMAN-ENGLISH DICTIONARY OF WORDS AND TERMS USED IN MEDICINE AND ITS COGNATE SCIENCES. Square 12mo, Roxburgh binding, 9s.

H. CHARLTON BASTIAN, M.A., M.D., F.R.S. Examiner in Medicine at the Royal College of Physicians; Physician to University College Hospital, etc.

PARALYSES: CEREBRAL, BULBAR, AND SPI-NAL. A Manual of Diagnosis for Students and Practitioners. With numerous Illustrations, 8vo, 12s. 6d.

II. VARIOUS FORMS OF HYSTERICAL OR FUNC-TIONAL PARALYSIS. Demy 8vo, 7s. 6d.

RUBERT BOYCE, M.B., M.R.C.S. Professor of Pathology in University College, Liverpool.
TEXTBOOK OF MORBID HISTOLOGY FOR STUDENTS AND PRACTITIONERS. With 130 coloured figures, royal 8vo, 31s. 6d.

## CHARLES H. BURNETT, A.M., M.D.

Clinical Professor of Otology in the Woman's Medical College of Pennsylvania; Aural Surgeon to the Presbyterian Hospital, Philadelphia.

SYSTEM OF DISEASES OF THE EAR, NOSE AND THROAT. By 45 Eminent American, British, Canadian and Spanish Authors. Edited by Charles H. Burnett. With Illustrations, in two imperial 8vo vols., 48s. nett.

HARRY CAMPBELL, M.D., B.S. LOND., M.R.C.P.
Physician to the North-West London Hospital.

THE CAUSATION OF DISEASE. An exposition of the ultimate factors which induce it. Demy 8vo, 12s. 6d.

FLUSHING AND MORBID BLUSHING, THEIR PATHOLOGY AND TREATMENT. With plates and wood engravings, royal 8vo, 10s. 6d.

DIFFERENCES IN THE NERVOUS ORGANISA-TION OF MAN AND WOMAN, PHYSIOLOGICAL AND PATHOLOGICAL. Royal 8vo, 15s.

HEADACHE AND OTHER MORBID CEPHALIC SENSATIONS. Royal 8vo, 12s. 6d.

R. E. CARRINGTON, M.D., F.R.C.P.

NOTES ON PATHOLOGY. With an Introductory Chapter by J. F. GOODHART, M.D., F.R.C.P., Physician to Guy's Hospital. Edited by H. E. CROOK, M.D. Lond., F.R.C.S. Eng., and G. MACKESON, L.R.C.P., M.R.C.S. Cr. 8vo, 3s.

ALFRED H. CARTER, M.D. LOND., F.R.C.P.

Senior Physician to the Queen's Hospital; Professor of Therapeutics, Mason College, Birmingham, &c.

ELEMENTS OF PRACTICAL MEDICINE. Seventh Edition, crown 8vo, 10s.

FRANCIS HENRY CHAMPNEYS, M.A., M.B. OXON., F.R.C.P.
Physician-Accoucheur and Lecturer on Obstetric Medicine at St. Bartholomew's Hospital, &c.

LECTURES ON PAINFUL MENSTRUATION.

Royal 8vo, 7s. 6d.

WALTER S. COLMAN, M.D., F.R.C.P.
Assistant Physician to the National Hospital for the Paralysed and Epileptic,

SECTION CUTTING AND STAINING: A Practical Introduction to Histological Methods for Students and Practitioners. Second Edition, with Illustrations, crown 8vo, 3s. 6d.

E. TREACHER COLLINS, F.R.C.S.

Assistant Surgeon to the Royal London Ophthalmic Hospital, Moorfields;
Hunterian Professor, Royal College of Surgeons, England, 1893-94

RESEARCHES INTO THE ANATOMY AND PATHOLOGY OF THE EYE. With 10 Plates and 28

Figures in the text, demy 8vo, 6s.

W. H. CORFIELD, M.A., M.D. OXON., F.R.C.P. LOND. Professor of Hygiene and Public Health in University College, London.

DWELLING HOUSES: their Sanitary Construction and Arrangements. Fourth Edition, with Illustrations, crown 8vo.

[In preparation.

DISEASE AND DEFECTIVE HOUSE SANITATION.
With Illustrations, crown 8vo, 2s.

SIDNEY COUPLAND, M.D., F.R.C.P.
Physician to the Middlesex Hospital, and Lecturer on Practical Medicine in the Medical School, etc.

NOTES ON THE CLINICAL EXAMINATION OF THE BLOOD AND EXCRETA. Third Edition, 12mo, 1s. 6d.

H. RADCLIFFE CROCKER, M.D. LOND., B.S., F.R.C.P.
Physician for Diseases of the Skin in University College Hospital.

DISEASES OF THE SKIN: THEIR DESCRIPTION,
PATHOLOGY, DIAGNOSIS, AND TREATMENT.
Second Edition, with Illustrations, 8vo, 24s.

EDGAR M. CROOKSHANK, M.B.
Professor of Comparative Pathology and Bacteriology, King's College,
London.

A TEXT-BOOK OF BACTERIOLOGY: Including the Etiology and Prevention of Infective Diseases, and a short account of Yeasts and Moulds, Hæmatozoa, and Psorosperms. Fourth Edition, illustrated with 22 coloured plates and 273 illustrations in the text, med. 8vo, 21s. nett. [Now ready.

HISTORY AND PATHOLOGY OF VACCINATION.
2 vols., royal 8vo, coloured plates, 20s. nett.

#### HERBERT DAVIES, M.D., F.R.C.P.

Late Consulting Physician to the London Hospital, etc.

THE MECHANISM OF THE CIRCULATION OF THE BLOOD THROUGH ORGANICALLY DIS-EASED HEARTS. Edited by ARTHUR TEMPLER DAVIES, B.A., M.D. Cantab., M.R.C.P. Crown 8vo, 3s. 6d.

HENRY DAVIS, M.R.C.S. ENG.

Teacher and Administrator of Anæsthetics at St. Mary's Hospital, and Anæsthetist at the Dental Hospital of London.

GUIDE TO THE ADMINISTRATION OF ANÆS-THETICS. Second Edition, fcap. 8vo, 2s. 6d.

ROBERT W. DOYNE, F.R.C.S.

Surgeon to the Oxford Eye Hospital; Ophthalmic Surgeon to St. John's Hospital, Cowley, etc.

NOTES ON THE MORE COMMON DISEASES OF

THE EYE. With test types, crown 8vo, 2s.

Now ready.

## DR. A. DUHRSSEN.

Late first Assistant in the Obstetric Clinic of the Charité Hospital in Berlin.

MANUAL OF GYNÆCOLOGICAL PRACTICE FOR STUDENTS AND PRACTITIONERS. Translated and Edited from the fourth German edition, by JOHN W. TAYLOR, F.R.C.S., Surgeon to the Women's Hospital, Birmingham, &c.; and FREDERICK EDGE, M.D. LOND., F.R.C.S., Surgeon to the Women's Hospital, Wolverhampton. With 120 Illustrations, cr. 8vo, 6s.

MANUAL OF OBSTETRIC PRACTICE FOR STUDENTS AND PRACTITIONERS. Translated and edited from the sixth German edition by JOHN W. TAYLOR and Frederick Edge. With Illustrations, cr. 8vo, 6s. [ Fust published.

> W. ELDER, M.D., F.R.C.P. EDIN. Physician to Leith Hospital.

APHASIA AND THE CEREBRAL SPEECH ME-CHANISM. With Illustrations, demy 8vo, 10s. 6d. Now ready. W. SOLTAU FENWICK, M.D., B.S. LOND., M.R.C.P. Physician to Out-patients at the Evelina Hospital for Sick Children, &c.

THE DYSPEPSIA OF PHTHISIS: Its Varieties and Treatment, including a Description of Certain Forms of Dyspepsia associated with the Tubercular Diathesis. Demy 8vo, 6s.

DISORDERS OF DIGESTION IN INFANCY AND CHILDHOOD. With illustrations, demy 8vo, 10s. 6d.

[Now ready.

# J. MILNER FOTHERGILL, M.D.

INDIGESTION AND BILIOUSNESS. Second Edition, post 8vo, 7s. 6d.

GOUT IN ITS PROTEAN ASPECTS.
Post 8vo, 7s. 6d.

THE TOWN DWELLER: HIS NEEDS AND HIS WANTS. Post 8vo, 3s. 6d.

ERNEST FUCHS.

Professor of Ophthalmology in the University of Vienna.

TEXTBOOK OF OPHTHALMOLOGY. Translated from the Second German Edition by A. Duane, M.D., Assistant Surgeon, Ophthalmic and Aural Institute, New York. With 190 Illustrations, large 8vo, 21s.

PROFESSOR DR. PAUL FÜRBRINGER. Director of the Friedrichshain Hospital, Berlin, &c.

TEXTBOOK OF DISEASES OF THE KIDNEYS AND GENITO-URINARY ORGANS. Translated by W. H. GILBERT, M.D., Physician in Baden-Baden, &c. Vol. I., demy 8vo, 7s. 6d.

SIR DOUGLAS GALTON, K.C.B., HON. D.C.L., LL.D., F.R.S., ormerly Secretary Railway Department Board of Trade; Assistant Inspector-General of Fortifications, &c.

HEALTHY HOSPITALS. OBSERVATIONS ON SOME POINTS CONNECTED WITH HOSPITAL CONSTRUCTION. With Illustrations, 8vo, 10s. 6d.

JOHN HENRY GARRETT, M.D.

Licentiate in Sanitary Science and Diplomate in Public Health, Universities of Durham and Cambridge, &c.

THE ACTION OF WATER ON LEAD: being an inquiry into the cause and mode of the action and its prevention. Crown 8vo, 4s. 6d.

ALFRED W. GERRARD, F.C.S. Examiner to the Pharmaceutical Society.

ELEMENTS OF MATERIA MEDICA AND PHAR-MACY. With the New Official Remedies, 1890. Crown 8vo, 8s. 6d.

JAMES F. GOODHART, M.D. ABERD., F.R.C.P.

Physician to Guy's Hospital, and Consulting Physician to the Evelina

Hospital for Sick Children.

ON COMMON NEUROSES; OR THE NEUROTIC

ELEMENT IN DISEASE AND ITS RATIONAL

TREATMENT. Second Edition, crown 8vo, 3s. 6d.

JOHN GORHAM, M.R.C.S.

TOOTH EXTRACTION: A manual on the proper mode of extracting teeth. Fourth edition, fcap. 8vo, 1s. 6d.

## GEORGE M. GOULD, A.M., M.D.

THE STUDENT'S MEDICAL DICTIONARY: Including all the words and phrases generally used in Medicine, with their proper pronunciation and definitions. Tenth Edition, 8vo, 14s. nett. [Now ready.

A POCKET MEDICAL DICTIONARY, giving the Pronunciation and Definition of about 12,000 of the Principal Words used in Medicine and the Collateral Sciences. 32mo, 4s. nett.

LANDON C. GRAY, M.D.

Professor of Nervous and Mental Diseases in the New York Polyclinic, &c.

A TREATISE ON NERVOUS AND MENTAL DISEASES FOR STUDENTS AND PRACTITIONERS

OF MEDICINE. With 168 Illustrations, 8vo, 21s.

DR. JOSEF GRUBER.

Professor of Otology in the University of Vienna, &c.

A TEXT-BOOK OF THE DISEASES OF THE EAR.

Translated from the German, and Edited by EDWARD
LAW, M.D., C.M. EDIN., M.R.C.S. ENG., Surgeon to the London
Throat Hospital for Diseases of the Throat, Nose and Ear; and
COLEMAN JEWELL, M.B. LOND., M.R.C.S. ENG. Second edition,
with 165 Illustrations, and 70 coloured figures, royal 8vo, 28s.

W. S. HEDLEY, M.D.

Medical Officer in charge of the Electro-Therapeutic Department of the London Hospital.

THE HYDRO-ELECTRIC METHODS IN MEDI-CINE. Second Edition, with Illustrations, demy 8vo,

CURRENT FROM THE MAIN: The Medical Employment of Electric Lighting Currents. With Illustrations, demy 8vo, 2s. 6d.

PRACTICAL MUSCLE - TESTING; AND THE TREATMENT OF MUSCULAR ATROPHIES. With Illustrations, demy 8vo, 3s. 6d. [Now ready.

BERKELEY HILL, M.B. LOND., F.R.C.S. Professor of Clinical Surgery in University College,

ARTHUR COOPER, L.R.C.P., M.R.C.S. Surgeon to the Westminster General Dispensary, &c.

SYPHILIS AND LOCAL CONTAGIOUS DISOR-DERS. Second Edition, entirely re-written, royal 8vo, 18s.

NORMAN KERR, M.D., F.L.S.

President of the Society for the Study of Inebriety; Consulting Physician,
Dalrymple Home for Inebriates, etc.

INEBRIETY: ITS ETIOLOGY, PATHOLOGY, TREATMENT, AND JURISPRUDENCE. Third Edition, 8vo, 21s.

## F. CHARLES LARKIN, F.R.C.S. ENG. Surgeon to the Stanley Hospital,

RANDLE LEIGH, M.B., B.SC. LOND.

Senior Demonstrator of Physiology in University College, Liverpool.

OUTLINES OF PRACTICAL PHYSIOLOGICAL

CHEMISTRY. Second Edition, with Illustrations,

crown 8vo, 2s. 6d. nett.

## LEWIS'S PRACTICAL SERIES.

In Crown 8vo Volumes, with Illustrations.

- THE TREATMENT OF PULMONARY CONSUMPTION.

  By VINCENT D. HARRIS, M.D. Lond., F.R.C.P., Physician to the City of London Hospital for Diseases of the Chest, etc., and E. CLIFFORD BEALE, M.A., M.B., Cantab., F.R.C.P., Physician to the City of London Hospital for Diseases of the Chest, etc. 10s. 6d. [Now ready.
- THE SURGICAL DISEASES OF CHILDREN AND THEIR TREATMENT BY MODERN METHODS. By D'ARCY POWER, M.A., M.B. Oxon., F.R.C.S. Eng., Demonstrator of Operative Surgery at St. Bartholomew's Hospital, &c. 10s. 6d.
- DISEASES OF THE NOSE AND THROAT.

  By F. de HAVILLAND HALL, M.D., F.R.C.P. Lond., Physician in Charge of the Throat Department at the Westminster Hospital, &c. 10s. 6d.
- PUBLIC HEALTH LABORATORY WORK.

  By H. R. KENWOOD, M.B., D.P.H., F.C.S., Assistant Professor of Public Health, University College, &c. Second Edition, 10s. 6d.
- MEDICAL MICROSCOPY.

  By FRANK J. WETHERED, M.D., M.R.C.P., Medical Registrar to the Middlesex Hospital. 98.
- MEDICAL ELECTRICITY.

  By H. LEWIS JONES, M.A., M.D., F.R.C.P., Medical Officer, Electrical Department, St. Bartholomew's Hospital. Second Edition, 10s. 6d.
- HYGIENE AND PUBLIC HEALTH.

  By LOUIS C. PARKES, M.D., D.P.H. Lond. Univ., Lecturer on Public Health at St. George's Hospital, etc. Fifth Edition, 10s. 6d.
- MANUAL OF OPHTHALMIC PRACTICE.

  By C. HIGGENS, F.R.C.S., Lecturer on Ophthalmology at Guy's Hospital Medical School, &c. 6s.
- A PRACTICAL TEXTBOOK OF THE DISEASES OF WOMEN. By ARTHUR H. N. LEWERS, M.D. Lond., M.R.C.P., Obstetric Physician to the London Hospital. Fifth Edition, 10s. 6d.
- ANÆSTHETICS, THEIR USES AND ADMINISTRATION.

  By DUDLEY W. BUXTON, M.D., B.S., M.R.C.P., Administrator of Anæsthetics at University College Hospital, etc. Second Edition, 5s.
- ON FEVERS: THEIR HISTORY, ETIOLOGY, DIAGNOSIS, PROGNOSIS, AND TREATMENT. By A. COLLIE, M.D. 8s. 6d.
- HANDBOOK OF DISEASES OF THE EAR.

  By URBAN PRITCHARD, M.D. (Edin.), F.R.C.S., Professor of Aural
  Surgery at King's College, London, etc. Third Edition, 6s.
- A PRACTICAL TREATISE ON DISEASES OF THE KID-NEYS AND URINARY DERANGEMENTS. By C. H. RALFE, M.A., M.D. Cantab., F.R.C.P., Physician to the London Hospital. 10s. 6d.
- DENTAL SURGERY FOR GENERAL PRACTITIONERS AND STUDENTS OF MEDICINE. By ASHLEY W. BARRETT, M.B. Lond., M.R.C.S., L.D.S., Consulting Dental Surgeon to the London Hospital. Third Edition, 3s. 6d. [Now ready.]
- BODILY DEFORMITIES AND THEIR TREATMENT.

  By H. A. REEVES, F.R.C.S. Ed., Senior Assistant Surgeon and Teacher

  of Practical Surgery at the London Hospital. 8s. 6d.

J. WICKHAM LEGG, F.R.C.P.

Formerly Assistant Physician to Saint Bartholomew's Hospital.

A GUIDE TO THE EXAMINATION OF THE URINE. Seventh Edition, edited and revised by H. Lewis Jones, M.D., Medical Officer in charge of the Electrical Department in St. Bartholomew's Hospital. With Illustrations, fcap. 8vo, 3s. 6d.

LEWIS'S POCKET MEDICAL VOCABULARY. Second Edition, 32mo, limp roan, 3s. 6d.

WILLIAM THOMPSON LUSK, A.M., M.D.

Professor of Obstetrics in the Bellevue Hospital Medical College, &c.

THE SCIENCE AND ART OF MIDWIFERY. Fourth
Edition, rewritten, with numerous Illustrations, 8vo, 18s.

JEFFERY A. MARSTON, M.D., C.B., F.R.C.S., M.R.C.P. LOND.

Surgeon General Medical Staff (Retired).

NOTES ON TYPHOID FEVER: Tropical Life and its Sequelæ. Crown 8vo, 3s. 6d.

WILLIAM MARTINDALE, F.C.S.
W. WYNN WESTCOTT, M.B. LOND.

THE EXTRA PHARMACOPŒIA, with Medical References, and a Therapeutic Index of Diseases and Symptoms. Eighth Edition, limp roan, med. 24mo, 9s.

A. STANFORD MORTON, M.B., F.R.C.S. ENG.
Surgeon to the Moorfields Ophthalmic Hospital, &c.

REFRACTION OF THE EYE: Its Diagnosis, and the
Correction of its Errors. Sixth Edition, small 8vo, 3s. 6d.

C. W. MANSELL MOULLIN, M.A., M.D. OXON., F.R.C.S. ENG. Surgeon and Lecturer on Physiology at the London Hospital, &c.

SPRAINS; THEIR CONSEQUENCES AND TREAT-MENT. Second Edition, crown 8vo, 4s. 6d.

ENLARGEMENT OF THE PROSTATE: its Treatment and Radical Cure. With plates, 8vo, 6s.

WILLIAM MURRAY, M.D., F.R.C.P. LOND.

ROUGH NOTES ON REMEDIES. Second Edition, crown 8vo, 3s. 6d.

ILLUSTRATIONS OF THE INDUCTIVE METHOD IN MEDICINE. Crown 8vo, 3s. 6d.

WILLIAM MURRELL, M.D., F.R.C.P.

Lecturer on Materia Medica and Therapeutics at Westminster Hospital.

MASSOTHERAPEUTICS; OR MASSAGE AS A MODE OF TREATMENT. Fifth Edition, crown 8vo, 4s. 6d.

WHAT TO DO IN CASES OF POISONING. Eighth Edition, royal 32mo, 3s. 6d.

#### G. OLIVER, M.D., F.R.C.P.

PULSE-GAUGING: A Clinical Study of Radial Measurement and Pulse Pressure. Illustrations, fcap. 8vo, 3s. 6d.

ON BEDSIDE URINE TESTING: a Clinical Guide to the Observation of Urine in the course of Work. Fourth Edition, fcap. 8vo, 3s. 6d.

THE HARROGATE WATERS: Data Chemical and Therapeutical, with notes on the Climate of Harrogate. Crown 8vo, with Map of the Wells, 3s. 6d.

DR. A. ONODI.

THE ANATOMY OF THE NASAL CAVITY, AND ITS ACCESSORY SINUSES. An Atlas for Practitioners and Students, translated by St Clair Thomson, M.D. Lond., F.R.C.S. Eng., M.R.C.P. Lond. Plates, small 4to, 6s. nett.

WILLIAM OSLER, M.D., F.R.C.P. LOND.

Professor of Medicine, Johns Hopkins University, &c.

CHOREA AND CHOREIFORM AFFECTIONS.

Large 8vo, 5s.

LOUIS C. PARKES, M.D. LOND., D.P.H.
Lecturer on Public Health at St. George's Hospital, &c.

INFECTIOUS DISEASES, NOTIFICATION AND
PREVENTION. Fcap. 8vo, cloth, 2s. 6d., roan, 4s. 6d.

R. DOUGLAS POWELL, M.D., F.R.C.P.,

Physician Extra-ordinary to H.M. the Queen; Physician to the Middlesex
Hospital; Consulting Physician to the Hospital for Consumption and
Diseases of the Chest at Brompton.

DISEASES OF THE LUNGS AND PLEURÆ IN-

CLUDING CONSUMPTION. Fourth Edition, with coloured plates and wood-engravings, 8vo, 18s.

DR. THEODOR PUSCHMANN. Public Professor in Ordinary at the University of Vienna. HISTORY OF MEDICAL EDUCATION FROM THE MOST REMOTE TO THE MOST RECENT TIMES. Translated by Evan H. Hare, M.A. (Oxon.), F.R.C.S. (Eng.), F.S.A. Demy 8vo, 21s.

SAMUEL RIDEAL, D.SC. (LOND.), F.I.C., F.C.S., F.G.S. Fellow of University College, London.

PRACTICAL ORGANIC CHEMISTRY. The detection and properties of some of the more important organic compounds. 12mo, 2s. 6d.

PRACTICAL CHEMISTRY FOR MEDICAL STU-DENTS, Required at the First Examination of the Conjoint Examining Board in England. Fcap. 8vo, 2s.

J. JAMES RIDGE, M.D., B.S., B.A., B.SC. LOND.

Medical Officer of Health, Enfield.

ALCOHOL AND PUBLIC HEALTH. Second Edition, Crown 8vo, 2s.

FREDERICK T. ROBERTS, M.D., B.SC., F.R.C.P. Professor of the Principles and Practice of Medicine in University College; Physician to University College Hospital, &c.

THE THEORY AND PRACTICE OF MEDICINE. Ninth Edition, with Illustrations, large 8vo, 21s.

THE OFFICINAL MATERIA MEDICA. Second Edit., entirely rewritten in accordance with the latest British Pharmacopæia, with the Additions made in 1890, fcap. 8vo, 7s. 6à.

H. D. ROLLESTON, M.D., F.R.C.P. Fellow of St. John's College, Cambridge; Assistant Physician and Lecturer on Pathology, St. George's Hospital, London;

AND A. A. KANTHACK, M.D., M.R.C.P. PRACTICAL MORBID ANATOMY. Crown 8vo, 6s. [CAMBRIDGE NATURAL SCIENCE MANUALS. ROBSON ROOSE, M.D., F.R.C.P. EDIN.

I.

GOUT, AND ITS RELATIONS TO DISEASES OF THE LIVER AND KIDNEYS. Seventh Edition, crown 8vo, 4s. 6d.

NERVE PROSTRATION AND OTHER FUNC-TIONAL DISORDERS OF DAILY LIFE. Second Edition, demy 8vo, 18s.

WILLIAM ROSE, B.S., M.B. LOND., F.R.C.S.

Professor of Surgery in King's College, London, and Surgeon to King's College Hospital, &c.

ON HARELIP AND CLEFT PALATE. Demy 8vo, with Illustrations, 6s.

BERNARD ROTH, F.R.C.S.
Fellow of the Medical Society of London.

THE TREATMENT OF LATERAL CURVATURE
OF THE SPINE. With Photographic and other Illustrations, demy 8vo, 5s.

DR. C. SCHIMMELBUSCH.

Privat-docent and Assistant Surgeon in Prof. v. Bergmann's University Clinic at Berlin.

THE ASEPTIC TREATMENT OF WOUNDS.

Translated from the Second German Edition by A. T. RAKE,
M.B., B.S. London, F.R.C.S. Eng., Registrar and Pathologist to
the East London Hospital for Children. With Illustrations,
crown 8vo, 5s.

G. E. SHUTTLEWORTH, B.A., M.D.

Late Medical Superintendent, Royal Albert Asylum for Idiots and Imbeciles of the Northern Counties, Lancaster, &c.

MENTALLY - DEFICIENT CHILDREN: THEIR TREATMENT AND TRAINING. With Illustrations, crown 8vo, 4s.

ALEXANDER J. C. SKENE, M.D.

Professor of Gynæcology in the Long Island College Hospital, Brooklyn.

TREATISE ON THE DISEASES OF WOMEN.

Second Edition, with 251 engravings and 9 chromo-lithographs, medium 8vo, 28s.

J. LEWIS SMITH, M.D.

Physician to the New York Foundling Asylum; Clinical Professor of Diseases of Children in Bellevue Hospital Medical College.

A TREATISE ON THE MEDICAL AND SURGICAL DISEASES OF INFANCY AND CHILDHOOD.

Eighth Edition, with Illustrations, large 8vo, 21s. [Now ready.]

E. HUGH SNELL, M.D., B.SC., LOND.

Diplomate in Public Health of the University of Cambridge; London County

Council Medical Officer to the Blackwall Tunnel.

COMPRESSED AIR ILLNESS, OR SO-CALLED CAISSON DISEASE. Demy 8vo, 10s. 6d. [Now ready.

JOHN KENT SPENDER, M.D. LOND.
Physician to the Royal Mineral Water Hospital, Bath.

THE EARLY SYMPTOMS AND THE EARLY TREATMENT OF OSTEO-ARTHRITIS, commonly called Rheumatoid Arthritis. With special reference to the Bath Thermal Waters. Small 8vo, 2s. 6d.

JAMES STOCKEN, L.D.S. ENG.

Pereira Prizeman for Materia Medica; Late Dental Surgeon to the National Dental Hospital.

DENTAL MATERIA MEDICA AND THERAPEU-TICS. Fourth edition, revised by Leslie M. Stocken, L.R.C.P., M.R.C.S., L.D.S.; and J. O. Butcher, L.D.S. Eng., Assistant Dental Surgeon to Guy's Hospital. Fcap. 8vo, 4s.

ADOLF STRUMPELL.

Professor and Director of the Medical Clinique at Erlangen.

A TEXT-BOOK OF MEDICINE FOR STUDENTS
AND PRACTITIONERS. Second edition translated
by Dr. H. F. VICKERY and Dr. P. C. KNAPP, with Editorial Notes
by Dr. F. C. SHATTUCK, Visiting Physician to the Massachusetts
General Hospital, &c. With 119 Illustrations, imp. 8vo, 28s.

C. W. SUCKLING, M.D. LOND., M.R.C.P.

Professor of Materia Medica and Therapeutics at the Queen's College,
Physician to the Queen's Hospital, Birmingham, etc.

ON THE DIAGNOSIS OF DISEASES OF THE BRAIN, SPINAL CORD, AND NERVES. With Illustrations, crown 8vo, 8s. 6d.

ON THE TREATMENT OF DISEASES OF THE NERVOUS SYSTEM. Crown 8vo, 7s. 6d.

J. BLAND SUTTON, F.R.C.S.

Assistant Surgeon to the Middlesex Hospital.

LIGAMENTS: THEIR NATURE AND MORPHOLOGY. Second Edition, wood engravings, post 8vo, 4s. 6d.

[Now ready.

HENRY R. SWANZY, A.M., M.B., F.R.C.S.I.

Examiner in Ophthalmology to the University of Dublin; Surgeon to the

A HANDBOOK OF DISEASES OF THE EYE AND THEIR TREATMENT. Sixth Edition, Illustrated with Wood Engravings, Colour Tests, etc., large post 8vo, 12s. 6d. [Now ready.

ALBERT TAYLOR.

Associate Sanitary Institute; Chief Sanitary Inspector to the Vestry of St. George, Hanover Square, etc.

THE SANITARY INSPECTOR'S HANDBOOK.
Second Edition, with Illustrations, cr. 8vo, 5s. [Now ready.

E. G. WHITTLE, M.D. LOND., F.R.C.S. ENG.

Senior Surgeon to the Royal Alexandra Hospital for Sick Children, Brighton.

CONGESTIVE NEURASTHENIA, OR INSOMNIA

AND NERVE DEPRESSION. Crown 8vo, 3s. 6d.

SIR JOHN WILLIAMS, M.D., F.R.C.P.

Professor of Midwifery in University College, London, &c.

CANCER OF THE UTERUS: BEING THE HARVEIAN LECTURES FOR 1886. Illustrated with Lithographic Plates, royal 8vo, 10s. 6d.

E. T. WILSON, M.B. OXON., F.R.C.P. LOND.

Physician to the Cheltenham General Hospital, &c.

DISINFECTANTS AND ANTISEPTICS: HOW TO

USE THEM. In Packets of one doz. price is., by post

[Fust thoroughly revised.]

BERTRAM C. A. WINDLE, D.SC., M.D., M.A. DUBL.

Professor of Anatomy in Mason College, Birmingham, &c.

A HANDBOOK OF SURFACE ANATOMY AND
LANDMARKS. Second Edition, revised with the collaboration of T. Manners-Smith, M.A., M.R.C.S., Lecturer on
Osteology, Mason College, Birmingham. With illustrations, post
8vo, 3s. 6d.

[Now ready.

EDWARD WOAKES, M.D. LOND.

Senior Aural Surgeon, London Hospital; Lecturer on Diseases of the Ear,
London Hospital Medical College.

ON DEAFNESS, GIDDINESS, AND NOISES IN
THE HEAD. Fourth Edition, Part I., with Illustrations, 8vo, 10s. 6d.

[Now ready.

OSWALD ZIEMSSEN, M.D.

Knight of the Iron Cross, and of the Prussian Order of the Crown.

THE TREATMENT OF CONSTITUTIONAL SYPHILIS. Post 8vo, 3s. 6d.

EWIS'S DIET CHARTS. A Suggestive set of Diet Tables for the use of Physicians, for handing to Patients after Consultation, modified to suit Individual Requirements; for Albuminuria, Anæmia and Debility, Constipation, Diabetes, Diarrhœa, Dyspepsia, Eczema, Fevers, Gall Stones, Gout and Gravel, Heart Disease (chronic), Nervous Diseases, Obesity, Phthisis, Rheumatism (chronic); and Blank Chart for other diseases. 5s. per packet of 100 charts, post free. A special leaflet on the Diet and Treatment of Infants is

sold separately. 12, 1s.; 100, 7s. 6d., post free.

I EWIS'S FOUR-HOUR TEMPERATURE CHART. Meets the requirements of a chart on which the temperature and other observations can be recorded at intervals of of four hours. Each chart will last a week. 20, 1s.; 50, 2s.; 100, 3s. 6d.; 500, 14s.; 1000, 25s.

CHART FOR RECORDING THE EXAMINATION OF URINE. Designed for the use of medical men, analysts and others making examinations of the urine of patients, affording a convenient method of recording the results of the examination. 10, 1s.; 100, 7s. 6d.; 250, 15s.; 500, 25s.; 1000, 40s.

CLINICAL CHARTS FOR TEMPERATURE OBSERVATIONS, ETC. Arranged by W. RIGDEN, M.R.C.S. 12, 18.; 100, 78.; 250, 158.; 500, 28s.; 1000, 50s.

Each Chart is arranged for four weeks, and is ruled at the back for making notes of cases; they are convenient in size, and are suitable both for hospital and private cases.

I EWIS'S CLINICAL CHART, SPECIALLY DESIGNED FOR USE WITH THE VISITING LIST. This Temperature Chart is arranged for four weeks, and measures 6 × 3 inches. 12, 6d.; 25, 1s.; 100, 3s. 6d.; 500, 16s. 6d.; 1000, 30s.

[ EWIS'S NURSING CHART. Printed on both sides. 20, 1s.; 50, 2s.; 100, 3s. 6d.; 500, 14s.; 1000, 25s.

\*.\* Mr. Lewis is in constant communication with the leading publishing firms in America and has transactions with them for the sale of his publications in that country. Advantageous arrangements are made in the interests of Authors for the publishing of their works in the United States.

Mr. Lewis's publications can be procured of any Bookseller in any part of the world.

Complete Catalogue of Publications post free on application.

