

Manual of gynecology / by D. Berry Hart and A.H. Freeland Barbour.

Contributors

Hart, D. Berry 1851-1920.

Barbour, A. H. Freeland 1856-1927.

Publication/Creation

Edinburgh ; London : W. & A.K. Johnston, 1897.

Persistent URL

<https://wellcomecollection.org/works/hbxstuh>

License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>



The Library of the
Wellcome Institute for
the History of Medicine

MEDICAL SOCIETY
OF
LONDON
DEPOSIT

Accession Number


Press Mark

HART, David Berry
and BARBOUR, Alexander
Hugh Freeland.



22101112283

WILLIAM BRYCE,
STUDENTS' BOOKSELLER,
54 and 56, LOTHIAN STREET.





MANUAL OF GYNECOLOGY

MANUAL
OF
GYNECOLOGY

BY

D. BERRY HART, M.D., F.R.C.P.E., F.R.S.E.

LECTURER ON MIDWIFERY AND DISEASES OF WOMEN, SURGEONS' HALL, EDINBURGH;
PHYSICIAN, ROYAL MATERNITY AND SIMPSON MEMORIAL HOSPITAL, EDINBURGH;
ASSISTANT PHYSICIAN FOR DISEASES OF WOMEN, ROYAL INFIRMARY, EDINBURGH;

AND

A. H. FREELAND BARBOUR, M.A., B.Sc., M.D., F.R.C.P.E., F.R.S.E.

LECTURER ON MIDWIFERY AND DISEASES OF WOMEN, SCHOOL OF MEDICINE, EDINBURGH;
ASSISTANT PHYSICIAN, ROYAL MATERNITY AND SIMPSON MEMORIAL HOSPITAL, EDINBURGH;
ASSISTANT PHYSICIAN FOR DISEASES OF WOMEN, ROYAL INFIRMARY, EDINBURGH;
PHYSICIAN FOR DISEASES OF WOMEN TO THE COWGATE DISPENSARY,
AND TO THE WOMEN'S DISPENSARY, EDINBURGH


WITH 13 LITHOGRAPHS AND 381 WOODCUTS

FIFTH EDITION

W. & A. K. JOHNSTON
EDINBURGH AND LONDON

1897

PRINTED BY W. AND A. K. JOHNSTON, EDINBURGH AND LONDON.



M19700

WELLCOME INSTITUTE LIBRARY	
Coll.	weIMOmec
Call No.	WP100
	1897
	H32m
	119

TO

OUR FRIEND AND TEACHER

ALEXANDER RUSSELL SIMPSON

M.D., F.R.S.E.

PROFESSOR OF MIDWIFERY AND DISEASES OF WOMEN AND CHILDREN IN THE
UNIVERSITY OF EDINBURGH



PREFACE TO THE FIFTH EDITION.

THIS edition has been carefully revised and brought up to date, the portions dealing with operative Gynecology being in great part re-written and illustrated by new plates.


To make room for numerous additions to the text, the Index of Gynecological Literature has been withdrawn. References to the most important recent papers on operative work are given in foot-notes.

To Mr Chas. Donald, M.B., we are indebted for much assistance in preparing this edition for the press.

D. BERRY HART.

A. H. FREELAND BARBOUR.

EDINBURGH, October 1897.



PREFACE TO FIRST EDITION.

IN writing this Manual we have tried to keep before our eyes the great principle that the Anatomy, Physiology, and Pathology of the Pelvic Organs form the foundation of good Clinical work. As students we felt the want of a text-book based on this principle and embodying the most recent views from the various literatures instead of giving those of one school. This want we have endeavoured to supply.

Our thanks are due to Professor Simpson for his kind advice in matters of difficulty: and specially to Mr J. A. Melville, for the literary revision of the text and the preparation of the copious Table of Contents and Indexes.

Messrs W. & A. K. Johnston have executed the lithographs with their well-known accuracy and finish: and to Mr James Bayne we are indebted for the care and fidelity with which he has drawn on the wood the majority of the engravings. We have in all cases acknowledged the source of every illustration not specially prepared for this work.

D. BERRY HART.

A. H. FREELAND BARBOUR.

EDINBURGH, July 1882.




TABLE OF CONTENTS.

PART I.

ANATOMY, PHYSIOLOGY, AND METHODS OF EXAMINATION OF THE FEMALE PELVIC ORGANS.

SECTION I. ANATOMY AND PHYSIOLOGY OF THE FEMALE PELVIC ORGANS.

	PAGE
CHAP. I. General Anatomy of External Genitals and Contents of Pelvis	4
External Genitals as observed clinically	5
The Pelvic Floor and Organs resting on it, con- sidered as a whole	10
The Pelvis considered in detail	11
Pelvic Floor dissected from below	11
Pelvic Floor dissected from above	14
The Uterus and its Annexa	16
Fallopian Tubes	24
Ovaries	26
Vagina	30
Bladder	33
Rectum	40
Perineal Body	42
Peritoneum	42
Connective Tissue of Pelvis	46
CHAP. II. The Sectional Anatomy of the Female Pelvis	48
Sagittal Mesial Section	49
Sagittal Lateral Section	50
Transverse or Horizontal Section	51
Coronal Section	51
Axial Coronal Section of Pelvis	54

	PAGE
CHAP. III. The Position of the Uterus and its Annexa, and the Relation of the Superjacent Viscera	55
The Normal Form and Position of the Uterus	57
The Local Divisions of the Pelvic-Floor Peritoneum as viewed through the Pelvic Brim, and the position of the Uterine Annexa	59
The Physiological Changes in the position of the Uterus	61
The Relation of the Small Intestine to the Pelvic Floor and to the Uterus with its Annexa	62
CHAP. IV. Structural Anatomy of the Female Pelvic Floor	64
In Sagittal Mesial Section	65
Pubic Segment	65
Sacral Segment	65
Segments Contrasted	65
In Axial Coronal Section	67
The entire Displaceable Portion	67
The entire Fixed Portion	67
Functions of the Female Pelvic Floor	68
Pelvic Floor Projection	69
CHAP. V. Bloodvessels of the Pelvis	72
Arterial Supply	73
to Uterus, Ovaries, etc.	73
to Perineal Region	74
Venous Supply	74
Lymphatics of the Pelvis	76
Lymphatic Glands	76
Lymphatic Vessels	76
of External Genitals	76
of Vagina and Cervix Uteri	76
of Uterus	76
Relation between Glands and Vessels	78
Nerves of the Pelvis	79
Spinal	79
Sympathetic	79
Development of Pelvic Organs	79
CHAP. VI. Physics of the Abdomen and Pelvis, with special reference to the Semiprone and Genupectoral Postures	85
The effect of Intra-abdominal Pressure on the Female Pelvic Floor	85
Results brought about by change of Posture, especially by the Genupectoral Posture	87
The effect on Uterine position of Digital Pressure in the Vaginal Fornices	87

	PAGE
CHAP. VII. Menstruation and Ovulation	92
Preliminary Considerations	93
General Phenomena of Menstruation	94
Local Phenomena of	94
Ovulation	95
Corpus Luteum	95
Source of Discharge, and Changes in the Uterine Mucous Membrane	96
SECTION II. PHYSICAL EXAMINATION OF THE FEMALE PELVIC ORGANS.	
CHAP. VIII. External Abdominal Examination	102
Inspection of External Genitals	102
Vaginal Examination	102
Bimanual or Abdomino-Vaginal Examination	102
Other Varieties of Abdominal Examination	114
Simple Rectal, Abdomino-Rectal and Abdomino- Recto-Vaginal	114
Simon's Method of passing the Hand into the Rectum	116
CHAP. IX. The Volsella	117
Description of Instrument	117
Methods of Use	117
Mechanism of Displacement it causes	117
Uses in Diagnosis	117
in Treatment	119
Contra-indications	119
Sims' Tenaculum	120
CHAP. X. Vaginal Specula	121
Spatular Speculum—the Sims	121
Tubular Speculum—the Fergusson	121
Bivalve Speculum—the Neugebauer	121
the Cusco	121
Uses and Comparative value of the various Specula	126
CHAP. XI. The Uterine Sound	128
Nature	128
Preliminaries to its use; contra-indications	129
Method of Use	130
Employment for Diagnosis	134
Treatment	135
Dangers attending its use	135
Sound combined with Bimanual	135
Relation of Sound to Bimanual and Rectal Ex- amination	137

	PAGE
CHAP. XII. Tents and other Uterine Dilators	138
Tents	138
Material	138
Purposes for which used	139
Preliminaries to and method of use	140
Dangers in use and contra-indications	143
Hard Rubber Dilators—Tait's, Hegar's	143
CHAP. XIII. The Curette	145
Varieties	146
Cases in which useful	146
Method of Use	146
Cautions and Dangers	147
Relation of Posture to Examination and Treatment	147
CHAP. XIV. Instruments	148
Knives	148
Scissors	148
Needles	149
Sutures	149
Vaginal Syringes and Douches ; Uterine douche	149
Cautery	152
Anæsthetics	153
Action of Chloroform	153
Uses of Chloroform	154
Method of administration	155
Dangers	156
Cocaine	157
CHAP. XV. Antisepsis	158
Activity of various Antiseptics	160
Asepsis	162
Disinfection of operating-room, hands	163

PART II.

DISEASES OF THE FEMALE PELVIC ORGANS.

SECTION III. THE PERITONEUM AND CONNECTIVE TISSUE.

CHAP. XVI. The Etiology of Gynecological Diseases	174
The special Anatomical Relations of the Pelvic Organs	175
Their Development and its occasional Defects	175
The Functions of Menstruation, etc.	176
Micro-organisms	176
Methods for detecting Micro-organisms	177

	PAGE
CHAP. XVII. Pelvic Peritonitis (Perimetritis)	182
Pathological Anatomy and Varieties	182
Etiology	182
Symptoms and Physical Signs	185
Differential Diagnosis	186
Course and Results	186
Prognosis	187
Treatment of Acute	187
Prophylactic	187
General	188
Local	191
Chronic	191
Tubercular Peritonitis	191
Malignant Peritonitis	193
 Pelvic Cellulitis (Parametritis)	193
Pathological Anatomy and Varieties	193
Etiology	194
Symptoms	194
Physical Signs	195
Differential Diagnosis between Pelvic Peritonitis and Cellulitis	196
Course and Results	196
Prognosis	197
Treatment	197
 Effects on Uterus of Pelvic Peritonitis and Cellu- litis	199
Displacements caused by Pelvic Peritonitis	200
Pelvic Cellulitis	201
 Parametritis Chronica Atrophicans	202
Circumscripta	202
Diffusa	203
Reflex Disturbances in	203
 CHAP. XVIII. Pelvic Hæmatocele and Hæmatoma	205
Preliminary Considerations	206
Terminology	206
Nature of Pelvic Hæmatocele	206
Pathological Anatomy	206
Etiology: Sources of Hæmorrhage and Varieties	209
Symptoms	212
Physical Signs	212
Diagnosis and Differential Diagnosis	213
Course and Results	213
Prognosis	213
Treatment	214
At onset of Hæmorrhage	214
After Suppuration has occurred	214

CHAP. XVIII. Pelvic Hæmatocele and Hæmatoma—*continued*.

New Growths in Peritoneum and Connective Tissue (Broad and Round Ligaments)	215
Tumours of Broad Ligament	215
Hydrocele of Round Ligament	215
Tumours of the Round Ligament	216
Echinococci in Pelvic Organs	216
Tumours of Pelvic Connective Tissue	217

SECTION IV. AFFECTIONS OF THE FALLOPIAN TUBES AND OVARIES.

CHAP. XIX. Affections of the Fallopian Tube	220
Abnormalities	224
Sinuosity, Stricture and Occlusion of the Tubes	224
Patent Condition of the Tubes	225
Catarrhal Conditions without Distension	225
leading to Hydrosalpinx	226
Purulent Salpingitis—Pyosalpinx	226
Infectious Granulomata	228
Hæmatosalpinx	229
New Formations: Tubo-ovarian Cysts	230
Affections of the Parovarium	231
CHAP. XX. Diagnosis of Tubal Disease	233
Catarrhal Conditions, Septic and Gonorrhœal	233
Pyosalpinx and Hydrosalpinx	233
Tubercular Disease	235
Treatment of Tubal Disease	236
Tait's Operation	236
CHAP. XXI. Malformations of Ovary	238
Ovaritis	239
Pathological Anatomy	239
Etiology	240
Symptoms and Physical Signs	240
Differential Diagnosis	241
Progress and Results	241
Treatment	241
Periovaritis	242
Displacements of the Ovary—Hernia	242
Etiology	242
Diagnosis and Differential Diagnosis	242
Treatment	242

	PAGE
CHAP. XXI. Malformations of Ovary— <i>continued</i> .	
Prolapsus	242
Pathological Anatomy	243
Etiology	243
Symptoms	243
Physical Signs	243
Treatment	243
Development of Operations for the Removal of the Appendages	244
CHAP. XXII. Pathology of Tumours, of Ovary, Parovarium and Broad Ligament	246
Preliminaries	247
Varieties of Ovarian Cyst	249
Naked-eye Anatomy	249
Microscopic Anatomy	249
Nature of Ovarian Fluid	252
Mode of Origin of Ovarian Cysts	253
Solid Ovarian Tumours	256
Non-Malignant	256
Malignant	257
Parovarian Cysts	259
Other Broad-ligament Cysts	259
CHAP. XXIII. Diagnosis of Ovarian Tumours	262
When Small (Pelvic in position)	262
Lateral to Uterus	263
Posterior to Uterus	264
When Large, Multilocular, and Pediculated (chiefly Abdominal in position)	264
Symptoms	264
Physical Signs	264
Differential Diagnosis	266
When Large and Extra-peritoneal (often papillo- matous)	268
Diagnosis of Adhesions	268
Co-existence of Pregnancy and Ovarian Tumour	268
CHAP. XXIV. Operative Treatment of Ovarian Tumours	269
Ovariectomy	270
Vaginal method	270
Abdominal method	270
Requisites	271
Preliminaries	272
The Incision	272
Evacuation of Cyst	273
Drawing out of the Cyst from Abdomen	273
Securing of the Pedicle	274
Treatment of Adhesions and Bleeding	276
Peritoneal Toilette	277
Closure of Wound	277
Drainage	277
Dressing of Wound	277

	PAGE
CHAP. XXIV. Operative Treatment of Ovarian Tumours— <i>continued</i> .	
After-Treatment: Treatment of Complications	278
Abdominal Method when the Tumour is Papillo-	
matous and Extra-peritoneal	279
Enucleation	279
Relation of Listerism to Ovariectomy	279
Ovariectomy when Pregnancy is present	280
Contra-indications to Ovariectomy	280
Course and Result of Ovarian Tumours when left	
alone	281
Adhesions	281
Torsion of Pedicle	281
SECTION V. AFFECTIONS OF THE UTERUS.	
Periods during which Morbid Conditions of the	
Uterus arise	283
CHAP. XXV. Malformations of the Uterus	285
Relations of Malformations to Development	285
Pathology	286
Etiology and Classification	291
Symptoms	292
Diagnosis	293
Prognosis	296
Treatment	296
CHAP. XXVI. Small Os Externum ; Rigidity, Stenosis, and	
Atresia of Cervix	297
Etiology and Pathology	297
Symptoms	299
Diagnosis	301
Prognosis	301
Treatment	301
Dilatation	301
Division	302
CHAP. XXVII. Atrophy of the Cervix and Uterus	305
Superinvolution of the Uterus	306
Pathology	306
Etiology	306
Symptoms	308
Diagnosis	308
Prognosis	309
Treatment	309
CHAP. XXVIII. Hypertrophy of the Cervix : Amputation	310
Forms of Hypertrophy of the whole Uterus	310
Two forms of Hypertrophy of the Cervix	310

CHAP. XXVIII. Hypertrophy of the Cervix : Amputation—*continued*.

Hypertrophy of the Cervix—Vaginal Portion	PAGE 311
Pathology	311
Etiology	311
Symptoms	312
Diagnosis	312
Treatment—Amputation	312
Circular and Flap Operations	313
Hypertrophy of the Cervix—Supra-vaginal portion	314
Treatment	317

CHAP. XXIX. Laceration of the Cervix and its Consequences 318

Introductory	319
Pathology	319
Etiology	321
Symptoms	322
Diagnosis	323
Treatment	324
Emmet's Operation	324

CHAP. XXX. Chronic Cervical Catarrh 329

Pathology	330
Etiology	335
Symptoms	336
Physical Signs	336
Diagnosis and Differential Diagnosis	337
Prognosis	338
Treatment	338

CHAP. XXXI. Endometritis 343

Pathology	344
Relation of Micro-organisms to Endometritis	348
Etiology	350
Symptoms of Acute Endometritis	351
of Chronic Endometritis	351
Physical Signs of Acute Endometritis	352
of Chronic Endometritis	353
Diagnosis : Differential Diagnosis	353
Prognosis	354
Treatment of Acute Endometritis	354
of Chronic Endometritis	354

CHAP. XXXII. Acute Metritis 360

Pathology	360
Etiology	360
Symptoms	361
Physical Signs	361
Progress and Termination	361
Diagnosis	362
Prognosis	362
Treatment	362

 Chronic Metritis 362

Pathology	363
Etiology	365

CHAP. XXXII. Acute Metritis—*continued*.

	PAGE
Symptoms	366
Physical Signs, Diagnosis	367
Differential Diagnosis	367
Treatment	368

CHAP. XXXIII. Displacements of the Uterus 371

Preliminaries	371
Definitions	373
Etiology	373
Frequency	373
Symptoms	374
Anteflexion	374
Pathology	374
Etiology	375
Symptoms	377
Obstruction and Congestion theories of Dysmenorrhœa	377
Physical Diagnosis	378
Differential Diagnosis	379
Prognosis	379
Treatment	380
Anteversio	380
Pathology	380
Etiology	380
Retroversion	381
Pathology and Etiology	381
Symptoms	382
Diagnosis	383
Treatment	383
Retroflexion = Retroversion + Retroflexion	383
Pathology	383
Etiology	386
Symptoms	387
Diagnosis	388
Differential Diagnosis	390
Prognosis	391
Treatment	392
Treatment of Associated Inflammatory Conditions	392
Replacement and Retention by Pessaries	392
Operative Treatment: Vaginal Fixation	402
Ventral Fixation	403
Alexander Adams' Operation	405

CHAP. XXXIV. Inversion of the Uterus 408

Pathology	408
Etiology and Frequency	412
Symptoms	413
Diagnosis	414
Differential Diagnosis	415
Course and results of Chronic Inversion	417
Prognosis	417

CHAP. XXXIV. Inversion of the Uterus—*continued*.

	Treatment	PAGE 417
	Reposition	418
	with the hand alone or aided	418
	by continuous slight elastic pressure	420
	Hysterectomy	421
CHAP. XXXV.	{ Tumours of the Uterus	422
	Classification	422
	{ Fibroid Tumours	423
	{ Pathology	423
	Situation	423
	Structure	423
	Mode of Growth, Varieties	426
	Changes in the Uterus	429
	Degenerative Changes	432
	Fibroid Tumours of the Cervix	433
	{ Etiology	434
CHAP. XXXVI.	Fibroid Tumours of the Uterus—(<i>continued</i>)	437
	Symptoms	437
	Menorrhagia, Irregular Hæmorrhages	437
	Painful Menstruation	439
	Sensations due to size and weight of Tumour	439
	Pressure Symptoms	439
	Sterility and Abortion	440
	Progress and results	440
	Physical Signs : Differential Diagnosis	441
	Of small Fibroid Tumours	441
	Of large Fibroid Tumours	443
	Prognosis	445
CHAP. XXXVII.	Fibroid Tumours of the Uterus—(<i>continued</i>)	446
	Medical Treatment	446
	Treatment by Electricity	448
	Surgical Treatment	450
	Historical sketch	451
	Operations by Vaginal Method	453
	Removal of polypi	453
	Enucleation	453
	Colpotomy	454
	Vaginal Hysterectomy	454
	Operations by Abdominal Method	454
	Removal of Subserous Tumours	455
	Enucleation	455
	Abdominal Hysterectomy	456
	Treatment of stump—Extra-peritoneal	456
	Intra-peritoneal	460
	Retro-peritoneal	462

CHAP. XXXVII. Fibroid Tumours of the Uterus—*continued*.

	PAGE
Operations by Combined Method	462
Pan-hysterectomy	462
Result of the three methods	464
Removal of Uterine Appendages	465

CHAP. XXXVIII. Fibro-Cystic Tumour of the Uterus 466

Pathology	466
Symptoms	468
Diagnosis : Differential Diagnosis	468
Treatment	469

CHAP. XXXIX. Polypi of the Uterus 470

Varieties	470
Symptoms	476
Diagnosis	478
Differential Diagnosis	479
Prognosis	480
Treatment	480

Tuberculosis Uteri 481

CHAP. XL. Carcinoma Uteri (of Cervix) 483

Pathology	484
Classification	484
Origin	484
Position	487
Progress	488
Extension through Lymphatics	489
„ to neighbouring Organs	491

Etiology 494

General predisposing Causes	494
Local predisposing Causes	496

CHAP. XLI. Carcinoma Uteri (of Cervix)—(*continued*) 499

Symptoms	499
Local Symptoms	499
Hæmorrhage	499
Offensive discharge	500
Pain	500
General Symptoms	501
Diagnosis	501
Differential Diagnosis	503
Prognosis	505
Causes of death	506

	PAGE
CHAP. XLII. Carcinoma Uteri (of Cervix)—(<i>continued</i>) . . .	509
Introductory	509
Treatment of Symptoms	510
Hæmorrhage	510
Offensive discharge	511
Pain	511
General Treatment	511
Treatment of the Disease	512
Principles of Treatment	512
Scraping out of diseased tissue, with and without Caustics	513
Amputation of the Cervix	514
Ecraseur or Galvano-cautery	514
Knife and Scissors	514
Supra-vaginal Amputation	515
Excision of the whole Uterus	516
By Abdominal Incision (Freund's method)	516
Through the Vagina	516
With Clamps	517
,, Ligature	520
Results	521
Metastasis and Recurrence	523
CHAP. XLIII. Carcinoma of the Body of the Uterus	525
Pathology and Etiology	525
Symptoms and Diagnosis	526
Treatment	527
Adenoma Malignum	528
CHAP. XLIV. Sarcoma Uteri	529
Pathology	529
Etiology and Frequency	533
Symptoms	534
Diagnosis	534
Prognosis	536
Treatment	536
Deciduoma Malignum	536
Clinical Features	539
Pathology	543
Origin and Nomenclature	545
SECTION VI. AFFECTIONS OF THE VAGINA.	
CHAP. XLV. Atresia Vaginæ	548
Pathology	548
Atresia Hymenalis	548
Atresia Vaginalis	550
Etiology	551
Symptoms	552
Diagnosis	553
Prognosis	555

CHAP. XLV. Atresia Vaginæ—*continued*.

	PAGE
Treatment	555
Dangers of Operation	555
Operation for Imperforate Hymen	556
for Atresia of the Vagina	557
for Atresia of the Cervix	558
Atresia of one half of a Septate Uterus and Vagina	558

CHAP. XLVI.	Vaginitis	561
	Nature and Relation to Micro-organisms	561
	Varieties	563
	Pathology	564
	Etiology	565
	Symptoms	566
	Diagnosis	566
	Treatment	567
	Vaginismus	568
	Etiology	568
	Symptoms and Diagnosis	569
	Treatment	569
	Sims' Operation	570
	Tumours of the Vagina	571
	Cysts	571
	Fibroid Tumours	573
	Carcinoma	573
	Sarcoma	574
	Tuberculosis	575

SECTION VII. AFFECTIONS OF THE VULVA AND PELVIC FLOOR.

CHAP. XLVII.	The Vulva: Malformations	578
	Development	578
	Hermaphroditism	580
	True	580
	False	580
	Inflammation of the Vulva (Vulvitis).	582
	Pruritus Vulvæ	583
	Eruptions on the Vulva	585
	Ulcus Serpiginosum	585
	Tumours of the Vulva	586
	Cysts of Bartholinian Glands	586
	Elephantiasis	587
	Neuroma	588
	Fibroma	588
	Lipoma	588
	Carcinoma and Sarcoma	588
	Lupus	588
	Kraurosis or Atrophy	589
	Pudendal Hernia	590
	Varix	590
	Hæmatoma	590
	External Hæmorrhage	590

	PAGE
CHAP. XLVIII. Rupture of the Perineum	592
Preliminaries and Nomenclature	592
Pathology and Varieties	593
Etiology	594
Significance	595
Treatment	596
Prophylactic	596
Operative—immediate	596
deferred	596
for restoration of function	
of sphincter ani	597
for rupture of perineum	602
CHAP. XLIX. Displacements of the Pelvic Floor	603
Preliminaries	603
Undue Yielding or Bulge	604
Prolapsus Uteri	605
Definition	605
Preliminaries	605
Etiology	605
Nature	606
Symptoms and Physical Signs	607
Mechanism	607
Summary of Displacement in Prolapsus	609
Diagnosis and Differential Diagnosis	609
Treatment by Pessaries	610
Treatment by Operation	610
Preliminary considerations as to Opera-	
tive Technique	614
Repair of Sacral Segment by Perine-	
orrhaphy, etc.	614
Operations that aim at narrowing	
Vaginal Walls (Colporrhaphy)	616
Shortening Round Ligaments	616
Vaginal Enterocoele	617

SECTION VIII. DISTURBANCES OF MENSTRUAL FUNCTION.

CHAP. L. Amenorrhœa	620
Causes, Local and Constitutional	620
Symptoms	620
Treatment	621
Menorrhagia	622
Causes, Local and Constitutional	622
Treatment	622
Dysmenorrhœa	623
Varieties	624
Treatment	626

SECTION IX. DISTURBANCES OF REPRODUCTIVE FUNCTION.

CHAP. LI.	Sterility	629
	Relative	630
	Absolute	630
	Etiology	631
	General Causes	631
	Local Causes	631
	Treatment	632

SECTION X. AFFECTIONS OF BLADDER AND RECTUM.

CHAP. LII.	The Bladder	636
	Anatomy and Physiology	637
	Methods of Exploring the Urethra	638
	Methods of Exploring the Bladder—	
	by Catheter and Sound	639
	by Finger and Speculum	640
	by Posture and Ureteric Catheterisation	643
	by Electric Endoscope	645
CHAP. LIII.	Affections of Urethra and Bladder	647
	Malformations	647
	Diseases of the Urethra	647
	Displacements	647
	Neoplasms; Urethral Caruncle	648
	Urethritis	649
	Dilatation and Stricture	649
	Diseases of the Bladder	650
	Displacements	650
	Cystocele	650
	Neoplasms	650
	Cystitis	651
	Calculi and other Foreign Bodies	654
	Functional Diseases	655
CHAP. LIV.	Vesico-vaginal Fistula	658
	Pathological Anatomy and Varieties	658
	Etiology	661
	Symptoms	662
	Diagnosis	662
	Prognosis	663
	Treatment	663
	Essentials	664
	Operation—Preparatory	666
	Paring of edges	667
	Adaptation of edges with sutures	668
	After-treatment	674
	Ureteric Fistula	674
	Closure of the Vagina: Kolpoplexis	676

	PAGE
CHAP. LV. The Rectum	678
Physiology	678
Examination	680
Diseases	680
Displacements of	681
Fissure of the Anus	681
Piles	583
Functional Disturbance of Rectum—Constipation .	684
Coccygodynia	685

APPENDIX.

Abdominal Section	687
Preliminaries	688
Antiseptics	688
The Abdominal Incision	690
Exploration of Abdomen and Pelvis and Removal of Tumours .	690
Possible Accidents during Laparotomy	694
Peritoneal Toilette; Closure of Wound	695
Anterior and Posterior Colpotomy: Vaginal Hysterectomy .	695
Electricity in Gynecology: The Apostoli Method of Treatment .	697
History	697
Note on Electrical Terms used	697
Units of Measurement	699
Action of different Currents and Poles	701
Action of the Galvanic Current	701
Action of the Faradic Current	702
Apparatus and Instruments	702
The Current: its strength, duration, and frequency of application .	703
Pathological Conditions in which Electricity is used in Gynecology .	703
Systematic Treatment of Nerve Prostration	704
Seclusion	705
Absolute rest in bed	705
Systematic extra-feeding	705
Use of Massage and Electricity	706
Hysteria	706
Massage	708
Case-Taking	710
Classification of Diseases of Women	714
Sources of Gynecological Literature	719

CLASSIFIED LIST OF ILLUSTRATIONS.

To facilitate study, we have grouped the illustrations under the following heads.

Anatomy—naked eye
Sectional Anatomy.
Anatomy microscopic.
Pathology—naked eye
Pathology—microscopic.

Charts of etiology.
Gynecological examination.
Instruments.
Operations.

ANATOMY—NAKED EYE.

PLATE III. Distribution of ovarian, uterine and vaginal
arteries facing p. 73

Fig.	Page	Fig.	Page
1 External genitals	6	43 Schultze's diagram of position of uterus	57
2 External genitals in section	7	45 Contents of female pelvis viewed through pelvic brim	60
3 Hymen of virgin with vertical slit	9	46 Position of fundus uteri and lie of ovaries	61
4 Hymen of virgin with oval opening	9	47 Diagram of position of uterus, according to distension of bladder	62
5 Crescentic hymen	9	49, 51 Diagrams of pelvic floor projection	69, 71
6 Outlet of bony pelvis	10	52 Venous supply of uterus and vagina	75
7 Dissection of perineal region	12	53 Lymphatics and lymphatic glands of pelvis and lower part of abdomen	77
8 Diagram of perineal muscles	12	54 Lymphatics of uterus	78
9 Muscles of clitoris and bulb	13	61 Diagram of intra-abdominal pressure	86
10 Oblique coronal section through external genitals	14	62 Outline of figure in genupectoral posture	87
11 Dissection of pelvis	15	64, 65 Diagrams of uterus before and after menstruation	97
12 Levator ani and coccygeus	16	124 Diagram of structures in broad ligament	231
13 Virgin uterus, from front and in section	17	164 Nulliparous os uteri	303
14 Multiparous uterus, from front and in section	18	165 Multiparous os uteri	303
15 Diagram of divisions of cervix	19	308 Sacral segment of the pelvic floor	594
16 Coronal section of uterus	20	332 Bladder in systole	637
20 Fallopian tube, ovary and parovarium	25	346 Normal relations of the cervix, the ureters, and the urethra	660
21 Section of a seal's ovary with peritoneal capsule and tube in it	27	376 Direction of rectum and of anus in relation to intra-abdominal pressure	679
23 Vagina on vertical section	29		
24 Anterior vaginal wall and multiparous cervix	30		
25 Diagram of vertical mesial section of pelvis	31		
26 Horizontal section of pelvic floor at pelvic outlet	32		
29 Urethral glands of Skene	34		
31 Course of the ureters	36		
34a The rectum inflated, showing sphincters	39		

SECTIONAL ANATOMY OF PELVIS.

PLATE I. Position of uterus and ovaries	facing p. 48
„ II. Coronal sections of left half of pelvis	„ p. 52

Fig.	Page	Fig.	Page
32 Vertical mesial section, showing Y-shape of bladder—frozen	37	48 Vertical mesial section, during parturition—frozen	66
33 Vertical section of pelvis, with bladder contracted—frozen	38	55 Model of organs at end of second month, p. 56, 57, 58, 59, 60, 80, and 128. Transverse sections of six weeks foetus, showing development of pelvic organs	81, 82, 83, 248
34b Coronal section through anus	39	63 Vertical mesial section of pelvis in genupectoral posture—frozen	90
36 Vertical mesial section showing peritoneum—frozen	43	264 Vertical mesial section of pelvis from a case of carcinoma uteri—spirit-hardened	492
37 Vertical mesial section, with bladder distended—frozen	45	265 Vertical mesial section of pelvis from a case of carcinoma uteri et vaginæ—spirit-hardened	492
38 Lateral sagittal section—spirit-hardened	50		
39 Coronal section of pelvis—frozen	52		
40, 41 Axial coronal sections of pelvis	53		
42 Transverse section of pelvis in line of pyriform muscles	56		
44 Vertical mesial section with bladder distended	58		

ANATOMY—MICROSCOPIC.

PLATE V. Mucous membrane of Fallopian Tube	facing p. 221
--	---------------

Fig.	Page	Fig.	Page
17 and 18 Vertical section through mucous membrane of uterus	21, 22	28 Transverse section of urethra	34
19 Vertical section through mucous membrane of cervix	23	30 Epithelial cells from vesical mucous membrane	36
22 Section of human ovary	28	35 Perpendicular section through end of rectum	41
27 Squamous cells of vagina on papilla of connective tissue	33	66 Mucous membrane of menstruating uterus	98

PATHOLOGY—NAKED EYE.

PLATE VI. Purulent and tubercular salpingitis.	facing p. 228
„ VII. Diagram of mode of origin and growth of multilocular and papillomatous tumours.	„ p. 259
„ VIII. Figs. 1 and 2, Erosion and Laceration of Cervix as seen in the Speculum	„ p. 332
„ XIII. Ulcus serpiginosum vulvæ	„ p. 586

Fig.	Page	Fig.	Page
94 Cervical canal dilated by a poly-pus	141	120 Retro-uterine hæmatocele, with pouch of Douglas not previously obliterated	209
116 Uterus bound down by peritonitic adhesions	186	121 Ante- and retro-uterine blood effusion	210
117 Uterus retroverted and fixed with adhesions	187	122 Vertical mesial section of recent hæmatocele in pouch of Douglas in case of ruptured sac of extra-uterine gestation	211
118 Uterus drawn to one side by peritonitic adhesions	200	123 Sarcoma of pelvic connective tissue	217
119 Vertical mesial section of a hæmatoma felt as a retro-uterine tumour in case of extra-uterine gestation	208	125, 126, 127 Diagrams of pyosalpinx	234, 235
		130 Multilocular ovarian cyst	249

Fig.	Page	Fig.	Page
131 Papillomatous cyst from hilum of ovary	250	248 Fibro-cystic tumour of uterus	467
138 Myoma of ovary	256	249 Fibrous polypus laid open to show its identity in structure with a fibroid tumour	471
142 Simple broad ligament cyst	260	250 Intra-uterine submucous fibroid becoming vaginal	472
149, 150 Rudimentary uterus	286, 287	251 Submucous fibroid which has come to be wholly in the vagina	473
151 Uterus bipartitus	287	252 Group of mucous polypi growing in the cervix uteri	474
152 Uterus didelphys	288	254 Non-malignant papilloma of cervix	475
153 Uterus unicornis	288	255 Pediculated submucous fibroid	477
154 Uterus bicornis	289	259 Carcinomatous nodule	487
155 Uterus septus	289	263 Carcinoma of cervix uteri producing fistula	491
156 Infantile uterus	290	264, 265 Vertical mesial sections of pelvis in carcinoma uteri	492
157 Primary atrophy of uterus	290	268 Cauliflower excrescence growing from cervix uteri	502
158 Fœtation in detached horn of uterus	294	270 Kidney and ureters, with hydro-nephrosis from a case of carcinoma	506
159 Uterus septus, puerperal	295	271 Carcinoma of the cervix leading to occlusion of os uteri	507
160 Normal and pinhole os in speculum	297	272 Mode of the spreading of carcinoma	512
161 Conical vaginal portion	298	278 Uterus extirpated for cancer of body	524
167 Uterus and ovaries from a case of superinvolution	306	279 Carcinoma of the body of the uterus	526
168, 169 Hypertrophied vaginal portion of cervix	311, 312	280 Sarcoma uteri with tumours in vagina	530
175 Hypertrophy of intermediate portion of cervix	315	282 Sarcoma uteri invading Fallopian tubes	532
176 Hypertrophy of supra-vaginal portion of cervix	315	285 Uterus from a case of deciduoma malignum	543
177 Hypertrophy of whole uterus secondary to prolapsus	316	288 Atresia vaginæ, seen from behind	549
178 Single laceration of cervix	320	289 Case of double atresia	550
179 Multiple or stellate laceration of cervix	320	290 Atresia hymenalis	553
197 Antelexion with stenosis at os externum	375	291 Atresia vaginæ—lower third	553
198 Diagram to show antelexion produced by cicatrization of utero-sacral ligaments	376	292 Atresia of cervix at os externum	554
199 Myoma of anterior wall	379	293 Atresia of cervix at os internum	554
203 Uterus retroverted and bound back by peritonitic adhesions	382	295 Atresia in a septate uterus	559
204 Extreme retroflexion of uterus	384	300 to 304 Normal development of external organs of generation	579
205 Congenital retroflexion	385	305, 306 Spurious hermaphroditism	581
219 Inversion of uterus	409	307 Abscess of the Bartholinian gland	587
220 Inversion of uterus+inversion of vagina, caused by a small sub-mucous fibroid	411	309 Central rupture of the perineum	595
221 Inversion of uterus	412	317 Hernial nature of prolapsus uteri	606
230 Section of a large fibroid tumour, fibres round several centres	424	329 Posterior vaginal enterocele	617
231 Section of a fibroid tumour, showing wavy bundles of fibrous tissue	425	330 Dysmenorrhœal membrane seen under water	625
232 Section of a fibroid tumour, showing spaces between bundles of fibrous tissue	425	331 Dysmenorrhœal membrane laid open	626
233 Uterus with elongated cavity due to presence of several fibroids	428	340 Caruncle at urethral orifice	648
234 Interstitial fibroid tumour	429	342 Large stone extracted by vaginal lithotomy	654
235 Submucous fibroid tumour projecting into uterine cavity	430	343 Chief varieties of urinary fistula	659
236 Pediculated submucous fibroid in process of extrusion	431	344, 345 Deep and superficial vesico-vaginal fistulae	659
237 Cervical polypus having apparently two pedicles	434	347 Relation of peritoneum to a fistula	661
239 Uterus containing large fibroid tumour	438	377 Rectocele	681
240 Case of two-and-a-half months' pregnancy associated with two large fibroids	442		

PATHOLOGY—MICROSCOPIC.

PLATE IV. The staphylo-coccus, strepto-coccus and Gono-	
coccus ; and their relation to the tissues facing p.	177
„ V. Figs. 3, 4, 5, Chronic catarrhal, pseudo-	
follicular, and purulent salpingitis	p. 221
„ VIII. Fig. 3, Section of cervix with simple erosion	p. 332
„ IX. Mucous membrane of Uterus in Endo-	
metritis	p. 345

Fig.	Page	Fig.	Page
129 Cellular bodies—the source of ovarian cysts	248	196 Section of uterine tissue in chronic metritis	364
132 Papillæ covered with columnar epithelium from ovarian cyst	251	253 Section of a mucous polypus of the cervix	475
133 Round-celled sarcoma from dermoid cyst	251	257 Cancer of the vaginal portion	486
134 Cells from ovarian fluid	253	258 Cancer of the cervix proper	486
135 Diseased blood-vessels in ovary	254	260 Section of a portion of cervix uteri with carcinomatous nodule	488
136 Epithelial tubes—the source of ovarian cysts	254	261 Section of a flat canceroid of the cervix	489
137 Cyst wall of multilocular ovarian tumours	255	262 Plug of cancer cells in a lymphatic	490
139 Cancer of ovary	257	269 Scraping from carcinoma of the cervix	504
140 Spindle-celled sarcoma of ovary	257	281 Section of mucous membrane in sarcoma	531
141 Alveolar sarcoma of ovary	258	283 Scraping from a fibroid tumour	535
141a Cell groups from ascitic fluid round malignant omental tumour	258	284 Scraping from a spindle-celled sarcoma	535
183 Papillary form of erosion	331	286 Section of deciduoma malignum	544
184 Follicular form of erosion	332	287 Tissues which may give rise to deciduoma malignum	545
185 True ulceration of the cervix	333	296 Granular vaginitis—acute form	564
186 Healing of catarrhal patch	338	297 Granular vaginitis—chronic form	564
191 Hypertrophied glands in endometritis	345	298 Colpitis emphysematosa	565
192, 193 Mucous membrane in endometritis fungosa	346	299 Section of vaginal cyst	572
194 Granulation from endometritis composed of embryonic tissue	347	306* Section of ulcer serpinosum	586

CHARTS OF ETIOLOGY.

Fig.	Page	Fig.	Page
238 Influence of age on development of fibroid tumour	435	267 Influence of age on development of carcinoma	497
266 Influence of sex on development of carcinoma	495		

GYNECOLOGICAL EXAMINATION.

Fig.	Page	Fig.	Page
67, 68 Abdominal regions	103, 104	85 First stage of passing sound	130
69, 70 Right hand in Bimanual	110, 111	86 Second stage of passing sound in retroverted uterus	131
71 Left hand in Bimanual	112	87 Proper contrasted with improper method of turning the sound	132
72 Displacement of pelvic floor and abdominal wall in Bimanual	113	88 Second stage of passing sound with uterus to the front	132
73 Right hand in abdomino-recto-vaginal examination	115	89 Sound arrested in antelexion	133
74 Uterus drawn down by volsella	118	90 Sound combined with Bimanual	136
79 Method of holding Sims' speculum	123		

Fig.	Page	Fig.	Page
95 Introduction of tangle tents . . .	142	225, 226 Diagnosis of partial inversion and intra-uterine polypus . . .	416
143 Area of dulness in ovarian tumour and ascites . . .	265	227 Uterine polypus+Inversion . . .	416
199 Diagnosis between fibroid and ante flexion . . .	379	290-293 Diagnosis of various forms of atresia . . .	553, 554
206 Diagnosis of retroflexion by Bimanual . . .	389	336 Catheterisation of ureters . . .	633
222, 223, 224 Diagnosis of inversion and polypus in vagina . . .	414	337, 338 Kelly's method of examining the bladder . . .	644
		378 Use of anal speculum . . .	682
		379-381 Diagrams for case-taking . . .	711-714

INSTRUMENTS.

Fig.	Page	Fig.	Page
50 Callipers for measuring pelvic floor projection . . .	70	202 Thomas' anteversion pessary . . .	381
75 Sims' tenaculum . . .	119	210 Hodge pessary . . .	396
76, 77 Sims' speculum . . .	122	211 Albert Smith pessary . . .	396
78 Bozeman's speculum . . .	122	212 Side view of Albert Smith pessary . . .	396
80 Speculum for hysterectomy . . .	124	215 Position and support of pessary illustrated . . .	399
81 Fergusson's speculum . . .	125	216 Position and action of pessary in vagina . . .	400
82 Cusco's speculum . . .	126	217 Schultze's pessary . . .	401
83 Sir J. Y. Simpson's sound . . .	128	218 Meadow's compound stem pessary . . .	401
84 A. R. Simpson's sound . . .	129	243 Koeberlé's serre-nœud . . .	456
91 Laminaria tents before and after expansion . . .	139	244 Keith's clamp for pedicle of fibroids . . .	458
92 Tupelo tents before and after expansion . . .	140	245 Cintrat's serre-nœud . . .	458
93 Tupelo tent expanded . . .	141	256 Forceps with catch for mucous polypi . . .	480
96 Tait's dilators . . .	143	273 Simon's sharp spoon . . .	514
97 Hegar's dilator . . .	144	275 Doyen's clamp for vaginal hysterectomy . . .	517
98 Loop curette . . .	145	294 Perforated glass plug used after operation for atresia vaginae . . .	557
99 Simon's scoop . . .	146	318 Greenhalgh's pessary with transverse bars . . .	610
100 Thomas' curette, modified by A. R. Simpson . . .	146	319 Ring pessary with diaphragm . . .	610
101 Recamier-Roux curette . . .	146	320 Simple elastic ring pessary . . .	610
102 Bozeman's scissors . . .	148	321 Ring pessary <i>in situ</i> . . .	611
103 Emmet's needles . . .	149	322 Zwanck's pessary . . .	611
104 Needle-holder . . .	149	334, 335, Simon's urethral specula . . .	642
105 Higginson's syringe . . .	150	339 Kelly's speculum for finding ureter . . .	645
106, 107 Vaginal douche . . .	150	341 Skene-Goodman's self-retaining catheter . . .	653
108 Fritsch's catheter for washing out the interior of the uterus . . .	151	351-353 Knives for operating on fistulae . . .	667
109 Cones for cauterization . . .	152	354 Sponge-holder . . .	667
110 Chloroform drop-cork . . .	155	357-358 Bozeman's fork and method of use . . .	668, 669
111-113 Schimmelbusch's sterilising apparatus . . .	164-166	361 Bozeman's suture adjuster . . .	670
144 Spencer Well's trocar . . .	273	362-363 Coghill's wire-twister and method of use . . .	671
145 Ordinary trocar . . .	273	372, 373 Sims' stationary catheter . . .	674
146 Nélaton's forceps . . .	274	378 Anal speculum . . .	682
147 Spencer Well's clamp . . .	274		
162 Schultze's dilator . . .	302		
163 Marion Sims' dilator . . .	302		
200 Greenhalgh's intra-uterine stem . . .	380		
201 Graily Hewitt's cradle pessary . . .	381		

OPERATIONS.

PLATE X. Operative treatment of backward displacements of the uterus . . .		facing p. 403		
„	XI. Hysterectomy for fibroid tumours of the uterus . . .	p. 457		
„	XII. Vaginal hysterectomy with clamps . . .	p. 518		
Fig.	Page	Fig.	Page	
114	Patient in lithotomy posture with operating tables . . .	166	Bilateral division of cervix with Kuchenmeister's scissors . . .	303
148	Ligature of pedicle by the simple, Tait's and Bantock's knots . . .	170	Circular amputation of the cervix . . .	313
		171-174	Flap amputation of the cervix . . .	314

Fig.		Page	Fig.		Page
180	Emmet's operation denuded surface	325	315	Continuous catgut suture in anterior colporrhaphy	601
181	Emmet's operation — introduction of sutures	326	316	Emmet's operation for ruptured perineum	602
182	Emmet's operation — tying of sutures	326	320	Introduction of ring pessary	610
187	Forceps dressed with cotton wadding for applications to cervical canal	339	323-326	Operation for repair of perineum	612-614
188, 189	Schroeder's excision of cervical mucous membrane	340	327	Various forms of raw surface made on posterior vaginal wall in operation for prolapsus	615
190	Martin's method of excising the mucous membrane of the cervix	341	328	Raw surface as made by Martin	616
195	Sound dressed with wadding for intra-uterine applications	355	333	Emmet's button-hole operation on the urethra	639
207	Reposition of retroflexed uterus by finger in rectum	393	344	Closure of superficial vesico-vaginal fistula	659
208	Reposition of retroverted uterus with the sound	394	345	Closure of deep vesico-vaginal fistula	659
209	Reposition of uterus with volsella, and finger in rectum	395	347	Relations of peritoneum to extensive fistula	661
213	Introduction of pessary	397	348	Simon's method of paring edges of fistula	664
214	Pessary carried on by finger	398	349	Sutures passed to close fistula	665
228	Reposition of inverted uterus with White's repositor	419	350	American and German methods of paring edges of fistula contrasted	665
229	Cup with stem for gradual reduction of inversion	420	355, 356	Splitting and flap operations for fistula	668
241	Mode of passing chain-ligature	452	357, 358	Fork used to prevent sutures from cutting	668, 669
242	Martin's operation for enucleation of fibroid from wall of uterus	455	359	Counter-pressure made with blunt hook	669
246	Extra-peritoneal treatment, with elastic ligature, of pedicle of fibroids	459	360-363	Method of twisting sutures	670-671
247	Different modes of suturing the pedicle after removal of a uterine fibroid	460	364-365	Bozeman's method of fixing sutures with plates and shot	671
274	Schroeder's supravaginal amputation of cervix	515	366	Removal of sutures	671
276, 277	Vaginal extirpation of the uterus	519, 520	367-368	Closure of a four-cornered fistula	672
310, 311	A. R. Simpson's operation for complete rupture of the perineum	597	369	Anterior lip of cervix used to close fistula	672
312, 313	Tait's operation for rupture into anus	598	370	Anterior lip divided to close in vertically a fistula	673
			371	Operation for atresia urethræ in fistula	673
			374-375	Simon's operation for closure of vagina: kolpoplexis	675-676
			378	Division of base of an anal fissure	682



PART I.

ANATOMY, PHYSIOLOGY, AND METHODS OF EXAMINATION OF THE FEMALE PELVIC ORGANS.

Section I. Anatomy and Physiology of the Female Pelvic Organs.

„ II. Physical Examination of the Female Pelvic Organs.



ERRATA.

- Page 23, line 13—"Non-stripped" in Fig. 19 should be "non-striped."
Page 182, line 22—"Parimetritis" should be "Perimetritis."
Page 180, lines 12 and 13—"After inoculation they are placed," etc.,
should come after line 8 "6 % glycerine agar."

ERRATA.

Page 28, line 13. "and striped" to "it should be 'unstriped'."
Page 182, line 22. "Portsmouth" should be "Petersburg."
Page 186, line 12 and 13. "After mentioning that the house" should
be "should give after line 12, 'of the same name'."

SECTION I.

ANATOMY AND PHYSIOLOGY OF THE FEMALE PELVIC ORGANS.

IN order to give a comprehensive idea of the Anatomy and Physiology of the Female Pelvic Organs, it will be advisable to consider them in the following manner.

CHAPTER I. General Anatomy of External Genitals and Contents of Pelvis.

CHAPTER II. The Sectional Anatomy of the Female Pelvis.

CHAPTER III. The position of the Uterus and its Annexa, and the Relation of the Superjacent Viscera.

CHAPTER IV. The Structural Anatomy of the Pelvic Floor ; Pelvic-Floor Projection.

CHAPTER V. The Blood-vessels, Lymphatics, and Nerves of the Pelvis. Development of Pelvic Organs.

CHAPTER VI. Physics of the Abdomen and Pelvis, with special reference to the Semiprone and Genupectoral Postures.

CHAPTER VII. Ovulation and Menstruation.

CHAPTER I.

GENERAL ANATOMY OF EXTERNAL GENITALS AND CONTENTS OF PELVIS.

LITERATURE.

- EXTERNAL GENITALS. *Ballantyne*—The Labia Minora and Hymen : Edin. Med. Jour., Nov. 1888, p. 425. *Budin*—Recherches sur l'hymen et l'orifice vaginal : Paris, 1881. *Carrard*—Beitrag zur Anatomie und Pathologie der kleinen Labien : Ztschrift für Geburtshülfe und Gynäkologie, Bd. X., Hft. 1. *Cullingworth*—A Note on the Anatomy of the Hymen and that of the Posterior Commissure of the Vulva : Jour. of Anat. and Phys., Vol. XXVII., p. 343. *Matthews Duncan*—Papers on the Female Perineum : Churchill, London, 1880. *F. P. Foster*—A Contribution to the Topographical Anatomy of the Uterus and its Surroundings : Am. J. of Obst., Vol. XIII., p. 30. *Garrigues*—The Obstetric Treatment of the Perineum : Am. J. of Obst., Vol. XIII., p. 231. *Hart, D. Berry*—Note on the Naked-eye Anatomy of the Female External Genitals : Ed. Med. Jour., 1882. Atlas of Female Pelvic Anatomy, Edinburgh, 1884. *Henle*—Handbuch der Eingeweidelehre des Menschen : Braunschweig, 1866. *Klein*—The External Genitals of the Male and Female, by E. Klein : Stricker's Manual of Human and Comparative Histology, Vol. II., (Syd. Soc. Tr., 1872). *Klein, G.*—Entstehung des Hymens : Festschrift von der deutschen Gesellschaft für Gynäkologie, 1894. *Klein and Smith*—Atlas of Histology : Smith, Elder, & Co., London, 1880. *Sutton*—On the Nature of the Hymen ; Brit. Gyn. Jour., 1888. *Webster*—The Nerve-endings in the Labia Minora and Clitoris, etc., Labor. Reports, R.C.P.E., 1891.
- MUSCLES OF PELVIC FLOOR. *Cunningham*—The Dissector's Guide ; Abdomen : Mac-lachlan & Stewart, Edinburgh, 1880. *Doran*—A Dissection of the Muscles of the Female Pelvis and Perineum : Lond. Obst. Tr., 1886, p. 274. *Henle*—op. cit. *Luschka*—Die Musculatur am Boden des weiblichen Beckens : Wien, 1861. *Savage*—Female Pelvic Organs, 2nd Edition : London, 1870. *Turner*—An Introduction to Human Anatomy : A. & C. Black.
- UTERUS, AND ANNEXA ; ORGAN OF ROSENMÜLLER AND GARTNER'S CANAL ; VAGINA. *Barnes*—The Diseases of Women : London, 1878. *Breisky*—Die Krankheiten der Vagina : Billroth's Handbuch, Stuttgart, 1879. *Coe*—The Anatomy of the Female Pelvic Organs : Amer. Syst. of Gynecology, Vol. I., p. 95. *Cruveilhier*—Traité d'Anatomie Descriptive : Paris, 1871. *Döderlein*—Das Scheidensekret u. seine Bedeutung für das Puerperalfieber : Leipzig, Besold, 1892. *Engelmann*—The Mucous Membrane of the Uterus : Am. J. of Obst., Vol. VIII., p. 30. *Farre*—The Uterus and its Appendages : Todd's Cyclopædia, Vol. V. *Hart*—Structural Anatomy of Female Pelvic Floor : Mac-lachlan & Stewart, 1880. *Henle*—op. cit. *Hennig*—Der Katarrh der inneren weiblichen Geschlechtstheile : Leipzig, 1862. *Küstner*—Das untere Uterinsegment und die Decidua cervicalis : Jena, 1882. *Rainey*—On the Structure and Use of the Ligamentum Rotundum Uteri : Lond. Phil. Tr. 1880, p. 515. *Ridder*—Ueber die Gartner'schen (Wolff'schen) Kanäle beim menschlichen Weibe : Virch. Archiv. Bd. 96. (This paper gives literature and

references to papers by Gartner, Dohrn, Freund, and others.) *Robinson, A.*—On the Peritoneal Relations of the Mammalian Ovary: *Jour. of Anat. and Phys.*, Vol. XXI., p. 169. *Routh, A.*—On cases of associated Parovarian and Vaginal Cysts: *London Obst. Tr.*, Vol. XXXVI. *Ruge and Veit*—Zur Pathologie der Vaginal Portion: *Stuttgart*, 1878. *Sappey*—*Traité d'Anatomie Descriptive*: Paris, 1873. *Schroeder*—*Handbuch der Krankheiten der weiblichen Geschlechtsorgane*: Leipzig, 1879. *Sutton, J. B.*—*Surgical Disease of the Ovaries and Fallopian Tube*: Cassell & Co., London, 1896.

BLADDER. *H. J. Garrigues*—Remarks on Gastro-Elytrotomy: *Am. Gynecol. Tr.*, Vol. III., p. 212. *Griffiths*—Observations on the Urinary bladder and Urethra: *Jour. of Anat. and Phys.*, Vol. XXV., p. 535. *Polk, W. M.*—Landmarks in the Operation of Laparo-Elytrotomy: *N. Y. Med. Jour.*, May 1882. *Skene*—Diseases of the Bladder and Urethra: New York, 1870. The Anatomy and Pathology of Two Important Glands in the Female Urethra: *Am. J. of Obst.*, Vol. XIII. *Winckel*—Die Krankheiten der weiblichen Harnröhre und Blase: *Billroth's Handbuch*, Stuttgart, 1877.

RECTUM. *Braune*—Topographisch-anatomischer Atlas, Zweite Auflage: Leipzig, Veit and Co., 1872. *Chadwick*—The Function of the Anal Sphincters so-called: *Am. Gyn. Tr.*, Vol. II., p. 43. *Hart*—Physics of Rectum and Bladder: *Edin. Med. Jour.*, 1882. *Otis*—Anatomical Researches on the Human Rectum: Leipzig, 1887. *Pirogoff*—*Anatome Topographica etc. sectionibus per corpus humanum congelatum*: Petropoli, 1859. *Ruedinger*—Topographisch-chirurgische Anatomie des Menschen: IV. Abtheilung. *Symington*—Rectum and Anus: *Jour. of Anat. and Phys.*, Vol. XXIII.

PERINEAL BODY. *Hart, Henle, Savage*, op. cit. *Ranney*—The Female Perineum: New York *Med. Jour.*, Vol. XXXVI., Nos. 1 and 2. *T. G. Thomas*—The Female Perineum; its Anatomy, Physiology, and Pathology: *Am. J. of Obst.*, Vol. XIII., p. 312.

PERITONEUM AND CELLULAR TISSUE. *Bandl*—Die Krankheiten der Tuben, der Ligamente, und des Beckenperitonäums: Stuttgart, 1879. *Barnes*—St George's Hospital Reports, Vol. VII., p. 57. *Freund*—Anatomische Lehrmittel zur Gynäkologie: Beiträge zur Geburtshülfe und Gynäkologie. Herausgegeben von der Gesellschaft für Geburtshülfe in Berlin, IV. Band, 1 Heft. S. 58. *Guérin*—Sur la Structure des Ligaments Larges: *Comptes Rendus*, 1879, 1364. *Le Bec*—Contribution à l'étude des Ligaments Larges (au point de vue de l'anatomie et de la pathologie): *Gaz. Hebdomadaire*, 15 Avril 1881. *Von Preuschen*—Ueber Cystenbildung in der Vagina: *Virchow Archiv*. Bd. 70. *De Sinéty*—Gynécologie: Paris, 1880. *Spiegelberg*—Remarks upon Exudations in the neighbourhood of the Female Genital Canals; German Clinical Lectures (New Syd. Soc. Tr.), p. 169. *Braune, Cunningham, Hart, Pirogoff*, op. cit.

EXTERNAL GENITALS AS OBSERVED CLINICALLY.

UNDER the term external genitals are comprised the structures known as Labia Majora, Fourchette, Labia Minora, Clitoris with its prepuce, Vestibule, and Fossa Navicularis. For clinical convenience the urethral orifice and hymen also are described with these; although the urethral orifice belongs to the urinary system, and the hymen separates anatomically the external genitals (vulva) from the vagina. External
Genitals.

The *Labia Majora* (fig. 1, *a*) are two thick folds of hair-clad skin, extending from the symphysis pubis backwards between the thighs, and meeting each other nearly in the middle line and about 2·7 cm. (1 inch) in front of the anus; their blunted posterior ends can be seen most distinctly in the foetus. Each labium has an outer and inner surface, and consists of a thick fold of skin enclosing a quantity of fat, blood- Labia
Majora.

vessels, and dartos. Superiorly, where they are best developed, they form by their junction—anterior commissure—the structure known as the mons veneris (*vide* fig. 1); while posteriorly they stop short of the fold of skin known as the *Fourchette* or posterior commissure. The fat and connective tissue are almost entirely wanting at the fourchette, which is not a distinct structure, but the posterior junction of the




FIG. 1.

EXTERNAL GENITALS OF VIRGIN, with Diaphragmatic Hymen. The Labia Majora and Minora are drawn apart, and the prepuce drawn back. The cadaver is in the lithotomy posture. (Modified from *Sappey*.)

a Labium majus; *b* Labium minus; *c* Vestibule just above urethral orifice; *d* Glans clitoridis; *e* Praeputium clitoridis; *f* Mons Veneris. (4)

thinned-out labia minora. Both labia majora are, in the adult, covered with crisp hair which is abundant over the mons veneris and outer surface but very much less so on the inner.

The *Labia Minora* (fig. 1, *b*) are two small oblique folds of hairless skin, one on the inner surface of each labium majus. Posteriorly each is lost in the labium majus at about its middle, while anteriorly they converge and each divides into two small branches—an upper and

Labia
Minora.

a lower. The upper branches meet to form the prepuce of the clitoris (fig. 1, *e*), while the lower in a similar way form its suspensory ligament. As a rule the labia minora do not, in the adult, project beyond the labia majora. Sebaceous glands are present on both labia. Microscopically the labia minora have the structure of skin, enclosing fibro-elastic tissue, a venous plexus, nerves, and lymphatics; nerve end organs are found in them and in the glans clitoridis. As above stated, the labia minora may be continued into the fourchette.

The *Clitoris*, covered by its prepuce, lies in the middle line and at the apex of the smooth piece of mucous membrane known as the vestibule. Only that part analogous to the glans penis is seen (fig. 1, *d*). The clitoris proper consists of two crura which arise from the rami of the ischium and pubes and unite superiorly to form the body of the clitoris, which lies beneath the mucous membrane. The glans clitoridis is not




FIG. 2.

VERTICAL MESIAL SECTION OF EXTERNAL GENITALS (*Henle*).

a Anus; *b* Perineal body; *c* Vagina; *d* Urethra; *e* Labium Minus; *f* Prepuce of Clitoris; *g* Fossa Navicularis, with Hymen in front and Fourchette behind. (†)

directly continuous with the body, but joins it through the pars intermedia of the bulb (*vide post*, p. 13).

The *Vestibule* (fig. 1, *c*) is a triangular smooth mucous surface bounded superiorly by the clitoris, laterally by the labia minora, and inferiorly by the upper margin of the vaginal orifice. In the middle line, at its base, the dimple of the urethral orifice can be distinctly felt 1 inch (2–2.5 cm.) in front of the fourchette. Small depressions and mucous glands open on its surface. It is covered with a multiple squamous epithelium.

The *Vaginal Orifice* lies in the middle line between the base of the vestibule and the fossa navicularis. Its orifice is guarded by the *hymen*, a thin fold of mucous membrane enclosing some connective tissue, blood-vessels, and nerves (?). The hymen is usually described as being crescentic in shape, attached to the posterior margin of the vaginal orifice and with a free edge towards the base of the vestibule (figs. 2 and 5);

or diaphragmatic, attached all round the vaginal orifice but with a small hole (figs. 1 and 4) or vertical slit (fig. 3) in it. Sometimes it is not so perforated, constituting a pathological condition. It must be noted, however, that if the hymen be examined clinically without disturbing its relations, it will be found to surround a vertical slit, the varying shapes of which, as seen in figs. 4 and 5, can only be recognised on separation of the edges. (*Farre, Cullingworth.*)

The point as to whether the Hymen belongs developmentally to the external genitals or vagina is disputed. Budin believes that the hymen is simply the thinned-out inferior margins of the anterior and posterior vaginal walls. One specimen we have examined certainly supports his statement that the vaginal columns run on the inner aspect of the hymen. Matthews Duncan has pointed out the interesting fact that in atresia vaginæ the hymen may be present, *i.e.* may be present although the vaginal walls are absent. More recently Pozzi has described cases of mal-development of the sexual organs, and brought out some interesting facts. One case was that of a male hypospadiac with external genitals simulating a female type, *i.e.* with a pseudo-vulva, a distinct hymen, and a fourchette. Pozzi found a ridge passing from the base of the glans penis, encircling the meatus urinarius and becoming continuous with the hymen: this he terms the male vestibular band. In a female with atresia vaginæ he found a similar band passing from the clitoris, surrounding the urethral orifice, and blending with the hymen. He advances the view that the hymen is vulvar in its origin and alleges that in women the "male vestibular band" can be seen on careful examination. In the hypospadiac already described this band was the remnant of the *corpus spongiosum*, so that he believes the hymen to be the analogue of the bulb in man.

Ballantyne and Sutton support the view that the hymen is vulvar in its origin. Ballantyne has also confirmed Pozzi's view (*vide* development of organs, p. 83, for fuller discussion).

Fossa
Navicu-
laris.

Fossa Navicularis.—Normally, the inner aspect of the fourchette is in contact with the outer and lower surface of the hymen. When the fourchette is pulled back by the finger, a boat-shaped cavity is made—the fossa navicularis. Its posterior boundary is, therefore, the inner aspect of the fourchette; its anterior is the external aspect of the hymen. These two are in contact unless artificially separated (fig. 2).

From behind forwards, in the female ano-vulvar region there lie in the middle line the following structures.

- (1.) Anus.
- (2.) Skin over base of Perineal Body.
- (3.) Fourchette.
- (4.) Fossa Navicularis.
- (5.) Vaginal orifice, with Hymen or its remains.
- (6.) Urethral orifice.
- (7.) Vestibule.
- (8.) Clitoris with its prepuce.

Laterally, we have the labia majora and minora.

The following points should be carefully noted. In the nude erect female only the mons veneris is seen, and the labia majora and minora lie in a plane nearly parallel to the horizon. The well-developed labia

majora have their inner surfaces always in contact, and are only slightly separated by the widest divergence of the knees. The labia minora are always in contact, and require to be artificially separated in order to see their inner surfaces. The untouched fourchette forms a U-shaped loop, with the limbs of the loop in contact by their inner edges. The fossa navicularis only exists when artificially opened up, and therefore, to see the external genitals fully, the labia must be separated and the prepuce and fourchette drawn back.

A line running as follows separates mucous membrane from skin. Starting from the base of the inner aspect of the right labium minus, it passes *down* beside the base of the outer aspect of the hymen, *up* along the base of the inner aspect of the left labium minus, *in* beneath the prepuce of the clitoris, and *down* to where it first started from.

The vulvar slit is sagittal, and lies in the middle line between the labia majora and minora.

The virginal vaginal orifice is also a vertical slit, only exists as an Hymen, opening when artificially made, and is anatomically defined by the hymen




FIG. 3.

FIG. 4.

FIG. 5.

HYMEN OF VIRGIN, with Vertical Slit. (†) HYMEN with Oval Opening. (†) CRESCENTIC HYMEN. (†)

which separates the external genitals from the internal genitals. The sharp line between skin and mucous membrane can be distinctly seen on the living subject. The labia minora are skin, thin and fine, and not mucous membrane as often alleged.

The following measurements by Foster are useful for reference :—

	Tip of Coccyx to Anus.	Anus to Fourchette.
Average distance in nulliparæ, . . .	4.5 cm. . .	2.7 cm.
" " multiparæ, . . .	4.7 cm. . .	2.5 cm.
Meatus urinarius, 2—2.5 cm. from fourchette, in nulliparæ; 2—3.1 cm., in women who have borne children.		

In a healthy woman who has experienced complete coitus, the hymen is torn or often only stretched. It admits two fingers without pain.

In a woman who has borne full-time children, the vaginal orifice is always torn, although the fourchette and all behind it may be intact. The carunculæ myrtiformes are probably the remains of the hymen. In addition, the passage of the child's head may cause tears of the posterior vaginal wall, perineal body, or even anterior wall of anus.

THE PELVIC FLOOR AND ORGANS RESTING ON IT CONSIDERED AS A WHOLE.

The outlet of the bony female pelvis is filled in by what is generally described as the "soft parts." This term, however, should not be employed, as it is misleading, especially in obstetrics. It is better named the pelvic floor or pelvic diaphragm.

Pelvic
Floor.

The pelvic floor is a thick fleshy elastic layer, dovetailed all round to the bony pelvic outlet (fig. 6). It may be considered as an irregularly-edged segment of a hollow sphere, with an outer *skin* aspect




FIG. 6.

BONY PELVIC OUTLET, with transverse line showing Rectal and Urethral Triangles
(D. J. Cunningham). (4)

and an inner *peritoneal* one. On the outer skin aspect lie the external genitals already described. On the inner peritoneal surface we have the organ known as the uterus, and its appendages the Fallopian tubes and ovaries. The vagina runs, in the erect female, at an angle of about 60° to the horizon, from the vaginal orifice upwards to the mouth of the womb, as a transverse slit in the pelvic diaphragm. In front of the vagina lies the bladder, while behind it the rectum is placed; these structures, along with muscles, connective tissue, blood-vessels, nerves, and lymphatics, making up the pelvic diaphragm.

Figure 1 shows, accordingly, the pelvic floor seen from its convex, skin aspect; fig. 45 gives it and the organs resting on it as viewed from its concave, peritoneal side; while fig. 32 displays it as seen in sagittal mesial section.

THE PELVIS CONSIDERED IN DETAIL.

PELVIC FLOOR DISSECTED FROM BELOW.

If a female cadaver be placed in the lithotomy posture and a transverse line drawn just in front of the ischial tuberosities, the perineal region will be divided into a posterior rectal triangle and an anterior urethral one (fig. 6). The former contains the anus, the latter the external genitals.

The fascia of the pelvic floor and its relations demand a few words here.

(1.) *The superficial fascia.*

(2.) *The deep layer of the superficial fascia.*

(3.) *The triangular ligament in two layers.*

(1.) *The superficial fascia* lies beneath the skin, and is simply the continuation over the pelvic floor of the general superficial fascia of the body.

(2.) *The deep layer of the superficial fascia* has the following attachments:—Laterally and above, it is joined to the pubic arch; while posteriorly it passes round the transverse perineal muscles to join the base of the anterior layer of the triangular ligament. If air be injected beneath this deep layer, its passage is limited by the attachments given, and a sac is made—the pudendal sac. Into this sac an inguinal hernia may push its way, and in it the round ligaments of the uterus end.

(3.) *The triangular ligament* consists of two layers of fascia, filling in the pubic arch. They are termed anterior and posterior. The following table may be omitted at present, until the whole anatomy is mastered.

<i>Between skin and superficial fascia.</i>	{	Supfl. hæmorrhoidal vessels and nerves.
		Supfl. perineal artery and nerve.
<i>Between deep layer of superficial fascia and anterior layer of triangular ligament.</i>	{	Transversus perinei.
		Bulbo-cavernosus.
		Erector clitoridis.
		Transverse perineal blood-vessels and nerves.
		Venous plexuses.
		Bulbs of vagina.
		Pudendal sacs.
<i>Between the layers of the triangular ligament.</i> (v. also p. 14.)	{	Dorsal artery and vein of clitoris.
		Compressor urethræ.
		Vagina—in part.
		Urethra—in part.
		Pudic vessels and nerves.

By suitable incisions the skin and superficial fascia can be removed around the anus, and the ischio-rectal fossa defined. This is a small pyramidal cavity on each side of the rectum, bounded externally by the obturator internus muscle, internally by the levator ani. Its apex is formed by the junction of these muscles, while its base is partially closed in by the transversus perinei and the edge of the gluteus maximus muscle (fig. 7). If axial-transverse sections of the fossa be made (Pl. II. fig. 2, and fig. 41), we see that it is merely the passage of the subcutaneous fat between the gluteus maximus, levator ani, and obturator internus muscles. The gluteus maximus forms the posterior and inferior boundary.

Ischio-
rectal
Fossa.

On transverse sections from before backwards it can be noted that its boundaries vary. At the level of the ischial tuberosity it is bounded as follows: inside, levator ani; outside, lower half of obturator internus; while the gluteus floors it in incompletely. About an inch posterior to the tuberosity, we find the boundaries change as follows: inside, we have still the levator ani; outside, a small portion of the obturator internus; while the gluteus maximus floors it in completely. At the posterior margin of the fossa, the levator ani is the inner and upper boundary, the gluteus maximus the outer and lower, the fossa here being quite below the level of the obturator internus.




FIG. 7.

DISSECTION OF PERINEAL REGION (*Savage*).

a is just above Transversus Perinei; *b* Base of Perineal body; *c* Bulbo-cavernosus; *d* lies on Levator Ani and in Ischio-rectal Fossa; *e* Erector Clitoridis; *f* Bulb of Vagina; *g* Bartholinian Gland; *h* Vestibule; *J* Glans Clitoridis. ($\frac{1}{2}$)

If the skin and superficial fascia be now removed from the urethral triangle, the following muscles, etc., will be exposed (fig. 7).

Muscles
beneath
superficial
fascia (deep
layer).




FIG. 8.

Perineal muscles.—On each side of the vaginal orifice three muscles lie, viz., the bulbo-cavernosus (fig. 8, *b c*), erector clitoridis or ischio-cavernosus (fig. 8, *e c*), and transversus perinei (fig. 8, *t p*).

The *Bulbo-cavernosi* consist of two muscular slips, one on each side of the vaginal orifice, which spring behind from the perineal body and pass round the vaginal orifice, partially covering the bulb and the vagina (fig. 7, *c*). The anterior end of each slip splits into three portions which end as follows:—

One passes to the under surface of the corpus cavernosum of the clitoris, a second goes to the posterior surface of

the bulb, and a third blends with the mucous membrane between the clitoris and urethral orifice (*Henle, v. fig. 9*).

The *Erector Clitoridis* arises from the inside of the ischial tuberosity and is inserted into the back and sides of the crus clitoridis (fig. 9, *e*).

The *Transversus Perinei* arises from the ramus of the ischium, and passes to the perineal body. It is difficult to define practically in dissection (fig. 7, *a*).

Now that these muscles have been described, we are in a position to localise more important structures.




FIG. 9.

a Symphysis Pubis, showing muscles in connection with Clitoris and Bulb. The Clitoris, *c, c''*, is cut across near its point, and thrown down with the vestibular mucous membrane (*Henle*). *b* Bulb. *c* Clitoris. *d* Urethra. *e* Erector Clitoridis; *f* Bulbo-cavernosus with its three insertions; *d* Branch to Dorsal Vein of Clitoris. (4)

The *Bulbi Vaginæ* (*corpora cavernosa urethræ*) are small masses of Bulbi erectile tissue about the size of a bean, lying one on each side of the Vaginæ vaginal orifice and partly under cover of the bulbo-cavernosus muscle. Each rests on the triangular ligament, and has internally the mucous membrane of the vagina; while, as already said, they are partly covered by the bulbo-cavernosus muscle. Anteriorly each blends with its fellow, and this *pars intermedia* becomes continuous with the clitoris (fig. 7, *f*).

Bartholinian
Glands.

The *Bartholinian Glands* lie one on each side of the vaginal orifice close to the posterior end of the bulb, and in front of the posterior layer of the triangular ligament (figs. 7, *g*, and 10, *e*). Each has a long duct opening on the labia minora close to the outer and anterior aspect of the hymen. Ranney asserts that these glands lie behind the posterior layer of the triangular ligament. They are compound mucous glands.

Perineal
body.

Between the lower third of the posterior wall of the vagina and the anterior wall of the anus, is an angular interspace (fig. 2, *b*) filled up by the structure known as the perineal body. This will be more fully described afterwards. At the present stage of the dissection only




FIG. 10.

OBLIQUE SECTION, parallel to the Anterior Pelvic Wall and through the External Genitals (*Henle*).
a Vagina; *b* Urethra; *c* Corpus Cavernosum Clitoridis, covered by its Erector; *d* Bulbus Vaginae covered by Bulbo-cavernosus Muscle; *e* Bartholinian Gland.

its base is seen, with the following muscles taking origin from or having an insertion into it,—sphincter ani, transversus perinei, bulbo-cavernosus, levator ani (fig. 7).

Between
layers of
triangular
ligament.

Between the layers of the triangular ligament lie the urethra, a portion of the vagina, compressor urethræ, dorsal vein of the clitoris, internal pudic vessels and nerves, the artery to bulb, dorsal nerve of clitoris, and Bartholinian glands (*Cunningham*).

The dissection of the urethral triangle has now been considered until the bladder has been exposed as it lies behind the pubes, from which it is separated by a considerable amount of loose fatty tissue. In order to complete the consideration, we have now to take up the muscles not yet described, viz., levator ani, coccygeus and the obturator internus.

THE PELVIC FLOOR DISSECTED FROM ABOVE.

The pelvic floor must now be looked at from its internal concave or peritoneal aspect. If the peritoneum and connective tissue beneath it, with the nerves and blood-vessels, be removed on one side of the pelvis,

say the right, the two muscles known as the coccygeus and levator ani will be exposed. These spring from the middle of the inner side of the true pelvis, and blending partly directly and partly indirectly with one another, form what may be termed the diaphragmatic muscles of the pelvic floor. If looked at through the pelvic brim, they are seen to form on both sides a concave arrangement analogous to the thoracic diaphragm (fig. 11).




FIG. 11.

DISSECTION OF PELVIS from above (*Savage*).

a Sacrum ; *b* Urethra ; *c* Vagina ; *d* Rectum ; *e* Levator Ani ; *f* Coccygeus ; *g* Obturator internus. ($\frac{1}{2}$)

The *Coccygeus* springs from the spine of the ischium and is inserted into the side of the lower part of the sacrum, and side and front of coccyx. There are two coccygei, one on each side (figs. 11 and 12).

The *Levator Ani* has an extensive origin. It springs in front from the back of the body and horizontal ramus of the pubes, from the pelvic fascia (white line) and the spine of the ischium. From this the muscle sweeps downwards and inwards to become attached in the middle line from before backwards as follows,—to the vagina, the rectum, its fellow of the opposite side, and finally to the tip of the coccyx (fig. 12). The

pubic fibres blend "with the posterior half of the upper border of the sphincter vaginæ" (Doran).

The levator ani can act on the vagina, elevating and compressing it, and is also believed to aid the sphincter ani (*v.* Pl. II. and figs. 40 and 41).

The *Obturator internus* has the following origin: deep surface of obturator membrane except at its lowest part; fibrous arch completing canal for obturator vessels and nerves; and surface of true pelvis bounded above by iliopectineal eminence, posteriorly by great sciatic notch, inferiorly by ischial tuberosity (fig. 41). Its relations are well shown in axial-transverse sections (*v.* Chap. II. and figs. 40 and 41). In fig. 41, its inferior half bounds the ischiorectal fossa; its upper half,




FIG. 12.

LEVATOR ANI and COCCYGEUS seen from without, after removal of part of hip bone and clearing out of Ischiorectal Fossa (*Luschka*).

a Fibres of Levator Ani on Vagina; *b* Anus, with Sphincter. (4)

the bladder and levator ani. It can also be seen that it lies in relation to the broad ligaments, *i.e.*, it bounds them where the peritoneal laminae diverge.

We have now to take up the consideration of the generative organs. It is difficult to describe these without alluding to structures not fully considered until further on. The student may, therefore, not entirely grasp some of the points until the whole anatomy of the organs has been mastered.

THE UTERUS AND ITS ANNEXA.

The Uterus *The Uterus* is a triangular body, with a truncated apex downwards, placed between the bladder and rectum, and with the appearance seen at figs. 13, *A* and 14, *B*. In describing it we take up its external appearance, its nature on section, and its structure and relations.

On external examination we find the parts known as the body (fig. 13, *A*, *c*), and neck (fig. 13, *A*, *a*, *b*). Keeping in mind its description as a triangle, we see the neck occupying the apex and the uterine orifices of the Fallopian tubes at the other two angles. Between the Fallopian tubes lies the fundus uteri. The anterior surface of the uterus is almost flat; the posterior is convex at its upper part, as is well seen in fig. 13, *B*. Where the body passes into the cervix there is a slight depression noticed on the posterior surface. This corresponds to the isthmus.

On making a vertical mesial section, we observe that the uterus is a hollow organ possessing a cavity with the anterior and posterior walls in apposition (fig. 13, *B*). In order to see the cavity it is advisable to

Corpus
Uteri.

Cavity of
Uterus.




FIG 13.

A. VIRGIN UTERUS (front view) (*Sappey*). The Appendages and Vagina are cut away.

a Cervix (vaginal portion); *b* Isthmus; *c* Body; *a b* Cervix proper.

B. The SAME in vertical mesial section.

a is anterior surface, and lies just above where peritoneum passes on to bladder.

C. The SAME with cavity exposed by coronal section.

e Os Externum; *d* Os Internum; *f* Uterine Opening of Fallopian Tube. (3)

look at the uterus in coronal section, *i.e.*, a section which, passing through the cavity, divides the uterus into an anterior and a posterior half, as shown in fig. 13, *C*, fig. 14, *A*. This latter section enables us more fully to understand the division of the uterus into body proper and cervix, and the division of the uterine cavity into cavity of the body proper and cervical cavity.

Cavity of Body.—This is a triangular slit in the uterus with the apex downwards, and with anterior and posterior walls. At each angle there is an opening, viz., at the lower angle we have the *os internum* opening

into the cervical canal (fig. 13, *C, d*), and at the upper angle the uterine openings of the Fallopian tubes (fig. 13, *C, f*). The lining of the cavity is known as its mucous membrane.

Cavity of the Cervical Canal.—This is spindle-shaped or conical (fig. 13, *B, C*), and has two openings, viz., os internum above and os externum below. The former opens into the uterine cavity, the latter into the vagina.

Cervix
Uteri.

The Cervix is divided into two portions, the vaginal and the supra-vaginal. The vaginal portion is within the vagina, and appears as a conical mass of the size and shape seen at fig. 13, *A, a*. The os




FIG. 14.

A. MULTIPAROUS UTERUS in coronal section to show cavity.
B. MULTIPAROUS UTERUS from front (*Sappey*). (3)

externum is in virgins a mere dimple, and feels to the examining finger like the tip of the nose. In women who have borne children it is transverse (fig. 14, *B*), and in most cases has its lips fissured more or less deeply, and the mucous membrane of the cervical canal partially everted. The supra-vaginal portion is continuous with the body through the isthmus.

The length of the whole unimpregnated uterus is, speaking generally, about 3 inches; the length of the cavity of cervix and body about $2\frac{1}{2}$ inches.

Measurements with the sound on the living female are a little in excess of those obtained in sections on cadavera, owing probably to the sound's elongating the uterus somewhat.

	Virgin.	Nulliparæ.	Multiparæ.
Length of uterus	2.35 in.	2.50 in.	2.70 in.
Width	1.50 „	1.55 „	1.70 „
Thickness	0.85 „	0.90 „	1.00 „
Vertical diameter of cavity .	1.80 „		<i>Sappey.</i> 2.44 in.
Transverse „ „ „	0.60 „		1.24 „
Length of entire organ in young women			<i>Richet.</i> 5.6 cm.
Do. body of uterus			3.35 „
Do. cervix			2.3 „
Do. vaginal portion of cervix			55-6 „
Capacity of uterus in nulliparæ = 2.3 c.cm.; in multiparæ 3.5 c.cm.			<i>Hennig.</i> <i>Sappey.</i>

Various authors divide the cervix uteri more minutely as follows. Divisions of Cervix Uteri.
They consider it as made up of—

- a. a vaginal portion ;
- b. an intermediate or middle portion ;
- c. a supravaginal portion. (Fig. 15).

This view is of importance in relation to the seat and extent of the changes in the size of the uterus in prolapsus uteri.




FIG. 15.

DIAGRAM of UTERUS to show divisions of Cervix (*Schroeder*).

a Vaginal portion ; b Intermediate portion ; c Supravaginal portion ; Bl Bladder ; P Peritoneum.
The dotted line shows peritoneum.

The question as to the precise position of the os internum in the Position of
unimpregnated uterus is at present much disputed. Küstner, who Os Inter-
has examined the point carefully, places the os internum at the num.
narrow part where the lumen of the cervical canal becomes continuous
with that of the uterine cavity proper. This part lies at the level of
the isthmus uteri (*v. fig. 16*) and is also the point where the com-
plicated uterine musculature passes into the simpler cervical muscular
arrangement. The folds of the arbor vitæ sometimes cease at this
point, but may pass above it or in multiparæ may end below it.

Küstner also alleges that for $\frac{1}{2}$ cm. ($\frac{1}{5}$ in.) below the os internum as defined by him the cervical substance and mucous membrane are like that of the uterine body, and that this special part of the cervical canal participates in the menstrual and pregnancy changes; and he therefore terms this the "inferior uterine segment," and speaks of a "cervical decidua." The os internum is believed by some to be at the level where the peritoneum passes on to the bladder.

Lower
Uterine
Segment.

While the two great divisions of the uterus are the *body* and *cervix*, it is of importance to keep in mind that in pregnancy we distinguish a special part of the body as the Lower Uterine Segment. It has the following characteristics: that the peritoneum is loosely attached over it, the muscular wall thinner there and the muscular bundles more separable; further, it plays in labour a passive rôle, and comes to be




FIG. 16.

CORONAL SECTION OF UTERUS (Küstner).

a. a. Uterine opening of Fallopian tubes; o. i. Os internum; o. e. Os externum.

marked off from the part above by a thickening in the wall known as the contraction or retraction ring. We may show the relations of all the divisions in the following scheme:—

Body	. (in pregnancy)	{ Upper portion (upper two-thirds).
		{ Lower Uterine Segment (lower one-third).
Cervix	.	{ supravaginal portion.
		{ intermediate „
		{ vaginal „

Küstner, as we have seen, speaks of the inferior uterine segment as cervical in origin.

Structure
of Uterus.

Structure of the Uterus.—If the uterus be viewed in vertical mesial section, it will be seen to be made up of three distinct elements, viz., peritoneum, unstriped muscular fibre, and mucous membrane (fig. 13, B.).

The peritoneum covers, partially, its external surface ; the mucous membrane lines the cavity of the body and cervix ; while the muscular fibre, by far the largest constituent, forms the tissue lying between these.

The Peritoneum of the Uterus clothes its posterior surface (except the vaginal and middle portions of the cervix), but only dips down on the front surface as far as the isthmus, at which level it is reflected

Periton-
eum of
Uterus.




FIG. 17.

Mucous membrane of uterus to show glands and interglandular tissue ; the mucous membrane is thicker here than normal.

on to the bladder (fig. 13, *B, a*). At the sides of the uterus the peritoneum on the anterior and posterior surfaces runs out to the wall of the pelvis, thus forming the important structures known as the broad ligaments.

The *Ligaments* of the uterus are—

- Broad ligaments ;
- Round ligaments ;
- Utero-sacral and Utero-vesical.

Ligaments
of Uterus.

Round
Ligaments.

The broad ligaments are described under the peritoneum. (See p. 44)

The round ligaments are two in number. According to Rainey, each springs by three fasciculi of tendinous fibres—the inner from the tendons of the internal oblique and transversalis, the middle from the superior column of the external abdominal ring near its upper part, and the outer fasciculus from just above Gimbernath's ligament. These unite into a rounded cord which crosses in front of the deep epigastric artery and passes between the layers of the broad ligament backwards, downwards, and inwards to the anterior and superior part of the uterus. Striped and unstriped muscle, blood-vessels, etc., are found in each.




FIG. 18.

MUCOUS MEMBRANE OF UTERUS (2nd). (Gulland.)

Utero-
sacral
Ligaments.

The utero-sacral ligaments are peritoneal folds, two in number, enclosing connective tissue and unstriped muscular fibre, passing from the lower, lateral part of the body of the uterus outwards and backwards towards the second sacral vertebra. They are known as the folds of Douglas, and form part of the upper, lateral boundaries of the pouch of Douglas. They are of the highest importance practically. The peritoneum, as it passes between uterus and bladder, constitutes the utero-vesical ligaments.

Utero-
Vesical
Ligaments.
Muscula-
ture of
Uterus.

The *Musculature of the Unimpregnated Uterus* is of little importance in Gynecology, and needs only a passing notice. Three coats are described:—a thin subperitoneal coat passing into the round ligaments, broad ligaments, utero-sacral and utero-vesical ligaments; a middle coat; and an inner concentric and very abundant layer which surrounds the

Fallopian tubes, os externum, and os internum. The student should not forget that the arrangement of the muscular fibres is of the highest importance in practical obstetrics.

The *Mucous Membrane* of the cavity of the body of the uterus is a thin reddish-gray layer, about 1 mm. ($\frac{1}{25}$ inch) thick in the unimpreg-
nated but fully developed organ. It is set on the inner aspect of the muscular layer of the uterus without the intervention of any submucous layer, is made up of ciliated columnar epithelium on a basis of connective tissue, and has numerous glands—the utricular glands. On section and microscopic examination, the glands, lined by the ciliated epithelium, lying, according to some, on a thin *membrana propria* can be seen

Mucous
Membrane
of Uterus.




FIG. 19.

CERVIX UTERI.—Semidiagrammatic. *gld*, Gland. *ep*, Squamous epithelium of vaginal portion. *art*, Artery. *vn*, Vein. *ct*, Connective tissue. *m*, Non-stripped muscle. (299). (Gulland.)

coursing down obliquely from the free surface and ending at, as well as sometimes dipping for a short distance into the muscular fibre. Fig. 17 shows the glands cut irregularly at various levels; they are more abundant than usual in this specimen (fig. 18). The glands usually bifurcate at their lower ends, and two may have a common mouth. Some of them penetrate the muscular fibre and can thus regenerate the mucous membrane after complete curetting.

The connective tissue in which the glands are embedded consists of delicate, round, and spindle-shaped cells, the former being more abundant near the surface, the latter deeper. Fibrillated bundles of connective tissue lie also between the cells and pass out between the muscular fibre of the uterine wall. According to Leopold, the connective tissue is in the form of a plexus of fine bundles, covered with endothelial plates each with a nucleus. The spaces between these bundles form lymph sinuses.

Mucous
Membrane
of Cervix.

The *mucous membrane* lining the cervix is different in arrangement and structure from that lining the cavity of the uterus. It is thrown into numerous folds, presenting to the naked eye the appearance known as the *arbor vitæ*, which consists of a longitudinal mesial ridge on the anterior and posterior walls, from both sides of which secondary ridges branch off obliquely. It is lined throughout with a single layer of epithelium (fig. 19), which is ciliated on the elevated portion of the ridges, but is columnar in the depressed portions (*de Sinéty*).

The upper boundary of the *arbor vitæ* varies. The boundary lies about midway between *os externum* and *fundus*. Before puberty, the folds pass up into the cavity of the body. In *multiparæ*, they do not pass up so far as in *nulliparæ* (*Küstner*).

The glands are of the racemose type, and consist of elongated, repeatedly branching ducts, which extend deeply into the connective tissue, and are somewhat dilated at their extremities (*Ruge and Veit*). They are lined by columnar epithelium, resting on a *membrana propria*, and open on the ridges and furrows of the mucous membrane.

There is a sharp line of demarcation between this single layer of epithelium (columnar and ciliated) which lines the cervical canal and the epithelial covering of the external surface of the vaginal portion, and this line of demarcation corresponds in the adult to the *os externum*. Beyond the *os externum*, the epithelial covering has all the characters of skin; it consists of vascular papillæ covered with many layers of squamous epithelium. The vascular papillæ are not easily recognised without the help of reagents (*Ruge and Veit*). The epithelial cells are like those found in the skin, and dovetail into one another by denticulate edges.

It is a disputed question whether glands are present on the vaginal aspect of the normal cervix. *De Sinéty* says he has never met with them except in the neighbourhood of the *os externum*, and their occurrence there he attributes to an eversion of the mucous membrane of the canal. *Ruge and Veit* also consider the existence of glands as a pathological condition, which is, however, easily induced.

The normal histology of the cervix uteri has an important bearing on the pathology of the so-called ulcerations and on laceration of the cervix and ectropium.

FALLOPIAN TUBES.

Fallopian
Tubes.

The *Fallopian tubes* are two tubes, one on each side of the uterus, running sinuously from its upper angles out towards the side of the pelvis (figs. 20 and 45). They lie enclosed in the upper free margin of the broad ligaments, and vary in length from 10 to 11 cm. (4 to 4 $\frac{2}{3}$ inches). They are not of equal length, the right being frequently longer than the left.

The Fallopian tube, the uterus lying to the front (*anteverted*), has

been found by His to pass first outwards and then upwards over the ovary, the fimbriated end lying on the posterior aspect of the ovary (Pl. I., fig. 2). Three parts come up for consideration—the isthmus, the ampulla, and the pavilion or fimbriated end.

The *isthmus* is the straight narrow part of the tube (fig. 20, *b*), which at its internal end opens into the uterine cavity, and has a lumen barely admitting a bristle. On transverse section the diameter of the whole thickness is about 2 to 3 mm.

The *ampulla* is the curved and thick part of the tube (fig. 20, *c*), having an average diameter of about 6-8 mm., with a lumen admitting the ordinary uterine sound.

The free *fimbriated end* of the Fallopian tube is expanded and funnel-




FIG. 20.

VIEW from behind of the LATERAL ANGLE of the UTERUS, with part of the Left Broad Ligament, Fallopian Tube, Ovary, and Parovarium (Henle).

a Uterus; *b* Isthmus of Fallopian Tube; *c* Ampulla; *g* has Parovarium to the right, and Fimbriated end of Fallopian Tube and Ovarian Fimbria just below it; *d* Parovarium; *e* Ovary; *f* Ovarian Ligament; *l* Infundibulo-pelvic Ligament (†). The topographical relations are disturbed here.

shaped (infundibulum); and it is provided with primary and secondary fimbriæ surrounding the opening of the tube to which they converge. One special fimbria runs to the ovary (fig. 20, *g*).

On section the Fallopian tube is seen to be made up of three layers from without inwards: viz., peritoneum, longitudinal and circular unstriped muscular fibres (the latter being inner), and mucous membrane lined with ciliated columnar epithelium. Connective tissue and elastic fibres lie between the peritoneal and muscular layers. No glands exist in the mucous membrane which is much folded in a longitudinal direction, especially in the ampulla. Bland Sutton, however, believes the folded

Structure
of Fallop-
ian Tube.

mucous membrane to be glandular, and to afford an albuminous covering to the ovum as it passes down.

It is remarkable that the ciliated epithelium lining the Fallopian tube and pavilion should be continuous with the squamous epithelium of the peritoneum; and that, further, there is direct continuity between the vagina, uterus, Fallopian tubes, and peritoneum,—so that the peritoneal sac in the female is not closed as in the male. The importance of this anatomical fact cannot, from a pathological point of view, be over-rated.

Par-
ovarium.

Parovarium or Organ of Rosenmüller. If the broad ligament be held between the light and the observer's eye, this rudimentary structure will be seen enclosed in its folds in the space between the ovary and ampulla (fig. 20, *d*). It consists of closed tubules lined with ciliated epithelium, which converge towards the ovary, and are united by a longitudinal one.

In the cow and sow the longitudinal tube or Wolffian duct persists, extending in the latter animal from a point a little above the division of the uterus into its cornua down the side wall of the vagina and opening into the vagina at the sides of the urethral orifice. These are named Gartner's canals after their chief investigator, and they correspond to the vas deferens, etc., in the male. Beigel has shown that these canals may be found in the uterus of the human foetus, a statement verified by Kölliker, Dohrn, and others. According to Rieder, they may persist either as a closed muscular epithelium-lined tube or as a muscular bundle without epithelium. The epithelial lining consists of a single or double layer of cylindrical cells (cells = 16μ): this is surrounded by connective tissue and by three coats of unstriped muscular fibre (inner and outer longitudinal and middle circular). It may produce one form of cervical or vaginal cyst as was shown by von Preuschen (*v. chapters on Development, Ovarian Pathology, and Vaginal Cysts*).

OVARIES.

Ovaries.

The ovaries, two in number, lie one on each side of the uterus, projecting markedly through the posterior layer of the broad ligament.

Form, Size, and Relations.—The ovary is a small oval-shaped body about the size of an almond, the weight of which varies from 60 to 135 grains. According to Farre its measurements are as follows:—

	Longitudinal Diameter.	Transverse Diameter.	Perpendicular Diameter.
Greatest	2 in.	$1\frac{1}{2}$ in.	$\frac{1}{2}$ in.
Smallest	1 in.	$\frac{1}{2}$ in.	$\frac{1}{4}$ in.
Average	$1\frac{1}{3}$ in.	$\frac{3}{4}$ in.	$\frac{3}{8}$ in.

The ovary has an anterior and posterior border, and an upper and lower surface. The posterior border is convex and free, the anterior flattened and attached to the broad ligament. It should be noted that this anterior border is called the hilum, and that the blood-vessels and nerves enter there.

The position of the ovary will be discussed afterwards (p. 61), but at

present it is sufficient to consider it as lying behind the broad ligament and on the side wall of the true pelvis, suspended as it were by the infundibulo-pelvic ligament so that its long axis lies more or less parallel to the axis of the brim of the pelvis.

Ovarian Fossa.—The ovary lies in a shallow fossa of the broad ligament, the ovarian fossa. In the rat, seal, and other animals the ovary is surrounded by a peritoneal fold, and thus completely cut off from the peritoneal cavity (fig. 21). It is possible this may happen in the human female, giving rise to one form of tubo-ovarian cyst.

Ligaments of the Ovary.—In addition to the attachment which the broad ligament gives to the ovary, two important ligaments are described—the ovarian ligament and the infundibulo-pelvic ligament. Ligaments of Ovary.

The *Ovarian Ligament* (fig. 20, *f*) is about 3 cm. ($1\frac{1}{5}$ inch) long, and extends from the inner end of the ovary to the corresponding upper Ovarian Ligament.




FIG. 21.

Section of a seal's ovary with peritoneal capsule, and tube in it.

angle of the uterus, just below the uterine origin of the Fallopian tube. It is a longitudinal fold of the peritoneum into which the unstriped muscular fibre of the uterus is prolonged.

The *Infundibulo-Pelvic Ligament* (fig. 20, *l*) is about 2 cm. long, and runs from the outer end of the Fallopian tube to the side wall of the pelvis. It is simply that part of the upper margin of the broad ligament unoccupied by Fallopian tube. Infundibulo-Pelvic Ligament.

The *Ovarian Fimbria* (fig. 20, *g*) prevents the separation of the ovary and infundibulum tubæ. Ovarian Fimbria.

Thus the ovary is kept in position by its attachment to the broad ligament, by the ovarian and by the infundibulo-pelvic ligaments. Its own specific gravity has also a share, *i.e.*, the ovary floats at a certain level.

Structure of the Ovary.—The ovary is covered with epithelium differing from the squamous epithelium of the peritoneum in being made up of Structure of Ovary.

columnar nucleated cells with a dull lustre. It is continuous at the hilum, however, with the peritoneal epithelium, the line of contact being marked by a whitish and elevated line—the white line of Farre. The epithelium covering the ovary is known as the germ-epithelium of Waldeyer. This distinctive term is of importance in connection with the development of the ova, and will be more particularly alluded to afterwards. A tunica albuginea made up of condensed connective tissue lies below the germ-epithelium.

On section and microscopical examination, the ovary is found to consist of connective tissue with the structures known as the Graafian




FIG. 22.

SECTION through the CORTICAL part of the OVARY (Turner).

e Germ Epithelium; *ss* Ovarian Stroma; 1, 1, large-sized Ovarian Follicles; 2, 2, middle-sized; and 3, 3, smaller-sized Graafian Follicles; *o* Ovum within Graafian Follicles; *v*, *v*, Blood-vessels in the Stroma; *g* Cells of Membrana Granulosa.

follicles embedded in it, along with blood-vessels, nerves, lymphatics, and some unstripped muscular fibre. These are enclosed in the epithelial covering already described. The connective tissue is divided into a cortical and medullary layer; the former lying beneath the peritoneum, the latter being at and near the hilum (fig. 21). The medullary layer is very vascular, and has some unstripped muscular fibre round the branches of the ovarian artery (fig. 22).

The Graafian follicles are scattered through the whole substance of the ovary. The following points should be carefully noted.

a. The younger and smaller Graafian follicles lie in the cortical layer. Their diameter is generally about $\frac{1}{100}$ th in., and they exist in immense numbers. According to careful estimates, the ovary of a female infant may contain 40,000 to 70,000 such follicles.

b. The larger follicles are much fewer in number and lie deeper in the ovary. Diameter $\frac{1}{30}$ th to $\frac{1}{100}$ th in.

c. There are also still larger follicles nearer the surface than the latter. These have advanced from the deeper layer (*vide* under Menstruation).




FIG 23.

A SECTION OF WHOLE VAGINA passing through Lateral Fornix; and B SECTION OF UPPER THIRD passing through the Cervix Uteri (*Hart*).

P. D. Pouch of Douglas; ut Uterus; o.e. Os Externum; Vg Vagina; p.f. Posterior Fornix; a.f. Anterior Fornix; V.u.p. Vesico-uterine Peritoneum; Bl. Bladder.

Structure of a Graafian Follicle. This consists of

1. A Tunica fibrosa and Membrana propria;
2. The Membrana granulosa, a layer of nucleated columnar epithelial cells forming the discus proligerus at one part;
3. Fluid—the liquor folliculi.

The *ovum* (diameter $\frac{1}{100}$ to $\frac{1}{130}$ in.) lies in the discus proligerus; it has the following structure:—

1. External envelope—zona pellucida, a homogeneous membrane,
2. Yolk protoplasm,
3. Germinal vesicle ($\frac{1}{700}$ th in. diameter),
4. Germinal spot $\frac{1}{3000}$ th in. diameter).

THE VAGINA.

Vagina—
Position.

The vagina is a mucous slit in the pelvic floor, extending from the hymen to the cervix uteri, and lying between the urethra and bladder in front and the rectum behind. In the upright posture it makes an angle of about 60° with the horizon, *i.e.*, it is nearly parallel to the brim conjugate.

Vaginal
Walls.

The vagina has two walls, an anterior and posterior, which are continuous at their sides. The anterior vaginal wall is triangular in shape,




FIG. 24.

ANTERIOR VAGINAL WALL AND MULTIPAROUS CERVIX, looked at from behind (*Henle*).

a Urethral Orifice; *b* Anterior Vaginal Column; *c* Cervix Uteri. (†)

the base being above. Its lower limit is marked out by the hymen. At its upper end it is reflected down to a small extent on the anterior lip of the cervix uteri, the anterior fornix being thus formed (fig. 23). It is closely incorporated with the urethra, but between it and the posterior aspect of the bladder there is loose connective tissue. Its length is about 5 cm., *i.e.*, 2–2½ inches.

The mucous membrane of the wall is arranged in folds roughly transverse. At its lower end is a vertical mesial single or double thickening of the mucous membrane, about 2 cm. long, known as the anterior vaginal column (fig. 24, *b*). This begins near the urethral orifice, or about $1\frac{1}{2}$ cm. above it. According to Budin, the columns are prolonged on the hymen.

The posterior vaginal wall is triangular in shape, and extends from the vaginal orifice upwards to the cervix uteri, upon which it is reflected, thus forming the posterior fornix vaginae, which is deeper than the anterior one. Its length is about $7\frac{1}{2}$ cm. (3 inches) *i.e.*, about $2\frac{1}{2}$ cm.




FIG. 25.

DIAGRAM OF VERTICAL MESIAL SECTION OF FEMALE PELVIS, showing Sigmoid curve of posterior Vaginal Wall (*Schultze*). (4)

(nearly an inch) longer than the anterior. It is also transversely rugous, and has a posterior column analogous to the anterior, but smaller. The vaginal rugæ can also be seen on the inner aspect of the hymen.

While the direction of the anterior vaginal wall is almost straight, that of the posterior vaginal wall is sigmoid (fig. 25). The curve varies, however, according to the position of the uterus and the fulness or emptiness of the adjacent bladder and rectum.

When the bladder and rectum are empty, we find the direction of the vagina parallel to the pelvic brim. When the bladder is distended, the

vagina is, chiefly at its upper part, driven nearer the sacrum ; while, if the rectum be distended, the vaginal axis may be almost perpendicular.

Structure
of Vagina.

Structure of Vagina.—The vaginal wall, on section and microscopical examination, is found to consist of mucous membrane, made up of epithelium (fig. 27) (the superficial layer being squamous and nucleated, the deeper layer cylindrical and with elongated nuclei), connective tissue, elastic tissue, and some unstriped muscular fibre. The superficial layer of the connective tissue forms papillæ, into which blood-vessels project. The epithelium is therefore ridged. External to this lie two layers




FIG. 26.

HORIZONTAL SECTION OF THE PELVIC FLOOR AT THE PELVIC OUTLET (*Henle*).
Ua Urethra ; *Va* Vagina ; *R* Anus ; *L* Levator Ani.

of unstriped muscular fibre ; the inner longitudinal, the outer circular (*Henle*). Breisky alleges the inner to be circular. Von Preuschen has described glands in the vagina, but they are very few in number. He found the ducts lined with squamous epithelium and the deeper part with ciliated epithelium—the latter being continuous with the cylindrical deep cells of the vagina. Gland-like crypts and lymph follicles also exist (*Löwenstein*). The whole is surrounded by loose connective tissue, containing the outer venous plexus of the vagina.

According to Döderlein, the normal vaginal secretion consists of vaginal epithelium, lymph corpuscles ; and a bacillus is present which gives an acid reaction (lactic acid) to the discharge.

As already said, the vagina is a mere slit in the pelvic floor, although it is often erroneously described as a tube or cavity. On vertical section, as fig. 23 shows, it appears as a mere linear slit ; while on transverse section it is H-shaped, or crescentic (figs. 26 and 41). The H-shape is said to be caused by the coalescence of the Wolffian ducts which, with part of the ducts of Müller, form the vaginal canal. The vagina is eminently dilatable and its walls separable, as will be more




FIG. 27.

SQUAMOUS CELLS OF VAGINA ON PAPILLA OF CONNECTIVE TISSUE.

Note active cells next papilla.

fully considered under the structural anatomy of the pelvic floor ; but this dilatation or separation is the result of posture with manipulation, or of parturition. Under mere changes of posture the vagina retains its slit-like form.

THE URETHRA AND BLADDER.

Position.—The empty female bladder lies behind the pubes and in front of the vagina.

The urethra is a straight slit (some describe it as sigmoid) about $1\frac{3}{8}$ inches long, with thick walls closely incorporated with the anterior vaginal wall behind. It runs parallel to the plane of the pelvic brim. Its lower opening is known as the meatus urinarius, the position of which has been already considered in the section on the External Genitals ; its upper opening is at the neck of the bladder. On section and microscopical examination, its mucous membrane is found covered with squamous epithelium in its lower part ; while higher up it is like that

Urethra—
Position.

Micro-
scopic
Structure.

of the bladder, and is very rich in elastic fibres. There is a double layer of *unstripped* muscular fibre, the longitudinal layer being internal and the circular outside; and, according to Uffelman, a circular (inner) and longitudinal layer of *striped* muscle, which stretches from the neck of the bladder to within $\cdot 6$ in. ($1\frac{1}{2}$ cm.) of the meatus urinarius. Luschka also describes a special sphincter of the vaginal and urethral orifices. It should be further noted that the mucous membrane is folded longitudinally, and contains mucous glands lined with cylindrical epithelium, papillæ, and lacunæ, and also villous tufts near the meatus; and that




FIG. 28.

TRANSVERSE SECTION of URETHRA much enlarged (Skene).

a Urethral Canal; *b b* Glands described by Skene; *c* Vein; *d* Artery.

Skene's
Tubules.

there is a submucous layer between the mucous membrane and unstripped muscle, containing many veins. Skene of New York has described two tubules in the female urethra. They lie on each side (figs. 28 and 29), "near the floor of the female urethra, and extend up from the meatus urinarius for about $\frac{3}{4}$ inch. They lie beneath the mucous membrane, and in the muscular walls of the urethra." We have in section of the female urethra:—

Mucous membrane;

Submucous layer;

Muscular layer, longitudinal and circular, unstripped;

do. do. striped (Uffelman).

External to these, there is the anterior vaginal wall behind, and loose tissue in front.

According to Henle, the closed urethral slit is on section transverse near the bladder, sagittal at the meatus, and star-shaped between these two points.

In the bladder proper we have three openings—the internal orifice of the urethra and the orifices of the two ureters. The latter lie one on each side, about $1\frac{1}{2}$ inches from the internal orifice. These openings give us the landmarks for the division of the bladder into neck, base, and body. All above the lines joining the ureteric openings and the centre of the symphysis is the body; all below is the base, and that portion between the ureteric openings and the internal orifice is the




FIG. 29.

URETHRA LAID OPEN from above, showing glands with probes passed in (*Skene*).

trigone. The trigone is formed by a blending together of the innermost bands of muscular fibre and fibrous connective tissue. Just above the ureters is the *bas fond*.

The wall of the bladder is made up of three layers, viz., a mucous, a muscular, and a peritoneal.

The mucous membrane consists of connective tissue lined by several layers of transitional or multiform epithelium (fig. 30). It is arranged in folds, except over the trigone and openings. The folds or rugæ are due to the laxity of the submucous coat.

The muscular coat of the bladder is of the unstriped variety, and has a complicated arrangement. There are external longitudinal fibres, oblique and transverse fibres within these, and an internal longitudinal layer on which rests the submucous coat. The so-called *detrusor urinæ*

Structure
of Bladder.

consists of two longitudinal bands, about 2 inches wide, on the anterior and posterior surfaces of the bladder. It is disputed whether




FIG. 30.

EPITHELIAL CELLS from the MUCOUS MEMBRANE of the BLADDER. Those in the upper row are the superficial squamous cells; those in the lower row are the peculiar cells of the middle stratum (Turner).




FIG. 31.

RELATION OF URETER ON THE RIGHT SIDE OF A DISSECTED PELVIS (Holl).

V Vagina; C Cervix; B Bladder; Ur, Ur, Ureter.

- | | | |
|------------------------|--------------------------|--------------------------|
| 1 Common iliac artery; | 2 External iliac artery; | 3 Internal iliac artery; |
| 4 Uterine artery; | 5 Pudic artery; | 6 External iliac vein. |

there is a sphincter at the neck of the bladder. Probably there is not; but the puckering of the mucous membrane at the neck is alleged to have a valve-like function.

The peritoneal covering of the bladder will be considered subsequently.

The relations of the ureters are of importance with regard to inflammatory exudations, fistulæ, and excision of the uterus for cancer. Ureters.

To Freund and Joseph, Luschka, Garrigues, Holl, and Polk, we are




FIG. 32.

VERTICAL MESIAL SECTION OF FEMALE PELVIS, showing Y-shape of Bladder (*Fürst*).
a uterus, b bladder, c rectum. ($\frac{1}{2}$)

indebted for anatomical researches as to the course of the ureter in the pelvis. We give Holl's drawing of the right ureter (fig. 31), and follow in the main his description.

Its course may be conveniently described in four portions.

(1.) *From the brim of the pelvis to the origin of the uterine from the internal iliac artery.*
About '6 in. ($1\frac{1}{2}$ cm.) below the division of the common iliac artery into its external and

internal branches, the Ureter passes over the external iliac vessels, and lies in front of the internal iliac artery and then in the space between the internal iliac artery and external iliac vein. So far, the portion described is at or about the level of the pelvic brim.

The Ureter next passes down into the true pelvis, and at the origin of the obturator, vesical, and uterine arteries begins to describe a bow-shaped portion 3·6 inches (9 cm.) long, with the greatest convexity of the bow where the uterine artery crosses it. By this crossing, the bow-shaped portion of the Ureter is divided into an upper and a lower part.

(2.) *From the origin of the uterine artery to where the Ureter is crossed by it.* This is the upper part of what is known as the bow- or spindle-shaped portion.

(3.) *From where the Ureter is crossed by the uterine artery to the bladder*—the lower part of the spindle-shaped portion.




FIG. 33.

VERTICAL MESIAL SECTION OF FEMALE PELVIC FLOOR, showing contracted bladder in a suicide (*Braune*). The peritoneum descends in front of the uterus to *b* and behind it to *d*; *b a* and *d c* are loose extra-peritoneal tissue. (1)

The uterine artery as it crosses the Ureter is separated from it by a venous plexus. In this way, a distance of about ¼ inch (1 cm.) separates Ureter and uterine artery at this point.

At the level of the os uteri externum the uterine artery crosses the Ureter to reach the uterus, and at this point the Ureter is ⅙ inch (1½ cm.) distant from the cervix. The course of this portion is of great importance. It is 1·6 inch (4 cm.) long, lies in relation to the side of the vagina, and then for the last two centimetres, before it pierces the bladder, lies between the anterior vaginal wall and the posterior wall of the bladder. The Ureter does not pass lower, therefore, than about the middle of the anterior vaginal wall.

(4.) *The portion piercing the bladder.* The Ureter runs through the bladder wall obliquely downwards and inwards for from ⅙ to ⅘ inches (1·5 to 2 cm.).

Shape and
Position of
Bladder.

Shape of empty Bladder and changes in its position.—The empty female bladder lies completely behind the pubes, and has its fundus covered by peritoneum. When empty and viewed in mesial section it may present one of two shapes. In the large majority of specimens figured, it forms with the urethra a Y-shape on sagittal mesial section.

The oblique legs of the Y may be about equal in size, or the posterior may be shorter (fig. 32). This form is so common that it has been accepted hitherto by all authors as the normal one. In certain cases, however, but not in so many as the former, the empty bladder cavity forms with the urethra a continuous tube on vertical mesial section (fig. 33). In such cases, it is oval in shape, corrugated, and firm to the touch. This latter shape is the one always found in the lower animals, such as the rabbit and dog, and is the only one seen




FIG. 34 a.

RECTUM INFLATED (*Chadwick*).
a b Sphincter tertius; c Ampulla of Rectum.




FIG. 34 b.

CORONAL SECTION THROUGH ANUS (*Symington*).
r rectum; i s internal sphincter; e s external
sphincter; l a levator ani; v vagina.

in the human foetus. If, therefore, the pelvic floor be viewed on its peritoneal aspect, the fundus of the empty bladder will be found to be very often large and concave, while in some cases it is small and convex. In the former case, the inner surface of the upper segment of the bladder, large in area, is in contact with the surface of the lower segment; in the latter, the anterior and posterior walls, small in area, touch one another.

It is probable that when the bladder has the Y-shape on section, it is relaxed and empty (fig. 32); and when the oval shape (fig. 33), it has been caught in systole. The bladder contracts to expel the urine and then relaxes. Between the acts of urination the bladder is therefore only a flaccid sac. Some additional facts as to the position and disten-

tion of the bladder are best considered further on, under the structural anatomy of the pelvic floor. We may here state, however, that (1) when empty, in the non-parturient female, it is behind the pubes (fig. 32); (2) it is drawn partly above the pubes in the parturient female (fig. 48); (3) it is tilted above the pubes in retroversion of the gravid uterus.

RECTUM.

Rectum.

The *Rectum* is not separated by any division from the sigmoid flexure, but may be defined as extending from the left sacro-iliac synchondrosis to the anus. It curves downwards, backwards, and inwards, to about the third sacral vertebra. This is known as the first part of the rectum; it is completely covered by peritoneum, which forms the mesorectum. The peritoneum is reflected from the rectum on to the upper part of the vaginal wall, about 3 inches above the vaginal orifice. Thereafter, the rectum lies in relation anteriorly to the posterior vaginal wall to which it is loosely attached until about $1\frac{1}{2}$ inches from the anus.

Micro-
scopic
Structure
of Rectum.

The rectum is made up of peritoneal investment; unstriped muscular fibre in two layers, longitudinal and circular, the former being the outer; a submucous coat; and a mucous lining with its muscularis mucosae, columnar epithelium, no villi, but with Lieberkuhnian follicles closely set together. At the upper limit of the anus, the circular fibres are very well marked, and constitute the sphincter ani internus (fig. 35).

Certain oblique folds in the rectum—consisting of mucous, submucous, and circular unstriped muscular coats—are of special interest. One exists $1\frac{1}{2}$ inches from the anus, another is near the sacral promontory, and one is intermediate (*Turner*). The lowest (the valve of Houston or sphincter ani tertius of Hyrtl) has been described by Chadwick of Boston, as being not an entire circular fold, but made up of two semi-circular constrictions, one on the anterior wall, and one on the posterior an inch higher up (fig. 34 a).

Anus.

The *Anus* is that part of the rectum at its external orifice. It is about an inch long, and has its long axis directed backwards and cutting the axis of the vagina at about a right angle. The rectum, therefore, when in contact with the posterior vaginal wall closely follows its direction, but at a little above the anus turns sharply backwards. There is thus left between it and the last $1\frac{1}{2}$ inch of the posterior vaginal wall, an angular interspace to be filled up by the structure known as the perineal body.

During life, the anus is closed by its sphincters in such a way that the lateral walls are in contact (*Symington*). This explains that the apparent gaping of the anus in sagittal mesial sections is approximately right (*v. Plate I.*), and that the appearance given at fig. 51 is wrong.

Fig. 35, from *Ruedinger*, shows the arrangement of voluntary and involuntary muscle in the anus. The division of the external sphincter

into two parts, and the separation of the lower division (5) into compartments by fibres from the longitudinal unstriated layer (9), are noteworthy. Similarly the internal sphincter (7) is divided into compartments by fibres from the muscularis mucosae (13). Near the anal orifice the




FIG. 35.

PERPENDICULAR SECTION through the end of the RECTAL WALL enlarged (*Ruedinger*).

- 1 Mucous Membrane of the Rectum; 2 boundary between Mucous Membrane and skin of buttock; 3 Fat; 4 Levator Ani; 5 Sphincter Ani externus; 9 Fibres of Longitudinal Layer separating external Sphincter into parts; 7 Sphincter Ani internus; 8 Longitudinal Fibres of muscular coat, which radiate outwards at 9; 13 Longitudinal Fibres of Muscularis mucosae which radiate outwards at 12; 11 Circular Fibres of muscular coat; 6, 10, and 14 Slips of muscular fibre passing into tissue beyond.

mucous membrane has certain perpendicular folds in it known as the Columnae Morgagni, with depressions between these—the Sinus Morgagni (fig. 2, a).

PERINEAL BODY.

Perineal
Body.

The posterior vaginal wall is in contact with the anterior rectal wall, for about $1\frac{1}{2}$ inches above the apex of the perineal body, there being only loose tissue between. The anus has its long axis directed backwards, while the vaginal axis runs forwards; we thus get a pyramidal space filled up by the structure known as the Perineal body (*Henle and Savage*).

The Perineal body is made up of muscular insertions and origins (striped and unstriped), and fibrous and elastic tissue. Its base is covered by the skin lying between the anus and vagina; its anterior side is in great part below the level of the posterior vaginal wall: its posterior side lies in front of the anterior rectal wall and anus; while laterally, it is bounded by fat. The voluntary muscles passing into it are the sphincter ani, transversus perinei, bulbo-cavernosus, and levator ani (fig. 7).

This Perineal body measures about $1\frac{1}{2}$ inches (4 cm.) vertically, the same transversely, and $\frac{3}{4}$ in. antero-posteriorly. If a straight line be made to join the tip of the coccyx and the subpubic ligament, it will just clear the apex of this structure.

Its functions are important, but have been both exaggerated and underrated. It gives a fixed point for many muscles, prevents pouching of the anus forwards, and strengthens that part of the pelvic floor which has no posterior bony support.

Its special significance, however, will be considered further on.

At present, the nomenclature in regard to the "Perineal region" is exceedingly vague—the term Perineum being used in this general sense by accoucheurs, especially in regard to the tears caused by parturition. It is better to speak of these as tears of the hymen, fourchette, and perineal body, instead of saying "perineal tears." The surface between the anal and vaginal orifices is, strictly speaking, not the perineum but the "skin over the base of the perineal body" and "the fourchette."

PERITONEUM.

Pelvic
Periton-
eum.

This is the thin serous covering of the concave surface of the pelvic floor and the organs resting on it. A knowledge of its disposition is of the highest importance to the gynecologist. This is best considered as follows.

1. *The Pelvic Peritoneum followed in a Vertical Mesial Section and from before backwards.*—The Peritoneum of the anterior abdominal wall is reflected, at a point a little above the symphysis pubis, on to the fundus of the empty bladder (fig. 36). It passes downwards and backwards over the bladder, from which it crosses on to the anterior surface of the uterus at a point about the level of the os internum. From

this it passes up over the anterior surface of the uterus. Thus there is formed a vesico-uterine pouch, containing no small intestine either when the bladder is in systole or in diastole (fig. 36). When the bladder has the Y-shape in pathological antelexion, the peritoneum passes directly



FIG. 36.

FROZEN SECTION showing Peritoneum (*Fürst*). The dotted line indicates Peritoneum in this and figs. 37-42. *a* Anus; *b* Vagina; *c* Bladder; *d* Uterus; *e* below pouch of Douglas; *f* Symphysis Pubis. (!)

backwards across the fundus of the bladder and on to the anterior surface of the uterus at or below the level of the os internum. There is thus produced a utero-abdominal pouch.

The peritoneum covers the whole of the anterior surface of the uterus above the os internum, passes over the fundus, and down the posterior surface which it covers almost completely. From this it descends still

deeper, on to the posterior aspect of the posterior vaginal wall for about half an inch (fig. 36). The depth of the peritoneal pouch formed behind the uterus is greater on the left side than on the right. The amount of its dip varies. In one section by Pirogoff the peritoneum runs down on the posterior vaginal wall till within about an inch from the vaginal orifice. This extent of posterior peritoneal duplicature is abnormal. This variation in depth is quite evident in sections: in some it ends at the level of the posterior fornix, while in others it is seen passing as deeply as has been already described. This descent of the peritoneum behind the uterus is of the highest importance practically, and forms the well-known pouch of Douglas. This pouch is defined as follows:—Its upper lateral boundaries are the utero-sacral ligaments; its anterior boundary is the uppermost half inch of the posterior vaginal wall and posterior aspect of the supra-vaginal portion of cervix; its posterior boundary is the sacrum and rectum, covered by peritoneum. It is the lowest part of the peritoneal cavity, and from its relation to the posterior vaginal wall can be explored through the posterior vaginal fornix. It is partially filled by intestine when the uterus lies to the front, which becomes displaced when the uterus is retroverted or retroflexed.

Pouch of
Douglas.

Broad
Ligaments.

2. *The Disposition of the Pelvic Peritoneum at the sides of the Uterus: Broad Ligaments.*—At the sides of the uterus, the peritoneum clothing its anterior and posterior surfaces passes outwards and somewhat backwards to the sides of the pelvis in front of the sacro-iliac synchondrosis. In this way we get two laminae of peritoneum nearly in apposition, which become more separated at their junction with the pelvic floor and sides of the pelvis; the space between the laminae is, at its outermost part, in relation to the obturator internus muscle (*v. Chap. II.*). These are the broad ligaments of the uterus. The posterior lamina has the larger area, and has its outer part applied to the pelvic side wall.

Immediately within their upper free margin, the Fallopian tubes are placed. That part of the free margin not occupied by Fallopian tube forms the infundibulo-pelvic ligament of the ovary (fig. 20 and Pl. I.). Projecting through the posterior lamina of the broad ligament is the ovary, covered by its germ-epithelium. The ovarian ligament and parovarium have already been described under the ovary and Fallopian tube.

Between the layers of the broad ligament lie connective tissue, unstriped muscle, blood-vessels, and lymphatics. According to M. Guérin, the broad ligaments enclose a small space shut off from the rest of the cellular tissue of the pelvis, and he denies that as yet there is proof of any special diagnosable inflammatory affection of the broad ligaments. Guérin alleges that, by inflation, it can be demonstrated

that the broad ligaments are thus shut off—a fact denied by other observers.

The position of the broad ligaments varies according to that of the uterus. When the uterus is normal in position, *i.e.*, lying to the front, their posterior surfaces look upwards and somewhat backwards, and they run outwards and backwards as already described. Displacement of the uterus backwards causes their coincident displacement, and in pregnancy they are drawn up and lie almost vertically. Pathologically, they cicatrize after inflammatory attacks and cause unilateral deviations of the uterus.




FIG. 37.

RELATION OF BLADDER AND PERITONEUM WHEN BLADDER DISTENDED (*Pirogoff*).

a Vagina; *b* Uterus; *c* Anus; *d* Bladder; *e* Symphysis.

3. *The Pelvic Peritoneum on the side walls of the Pelvis.*—The pelvic peritoneum clothes the side walls of the pelvis. It dips down least at the sides of the bladder, and most at the utero-sacral ligaments. periton-
eum on
side walls
of Pelvis.

Although the pelvic peritoneum has been described in three sections, it must of course be kept in mind that it is a continuous membrane with no breaks in its continuity.

Some special facts about the peritoneum should now be noted.

1. *As to the Bladder.*—Over the bladder and anterior abdominal wall, the peritoneum is easily separable. According to Spiegelberg, posteriorly it is closely blended with the uterus above the os internum, below this quite loosely attached. When the bladder is distended, the peritoneum is stripped off the lower part of the anterior abdominal wall to an extent varying with the distention Relation to
Bladder
and
Rectum.

(fig. 37). During parturition, the peritoneum is drawn off the bladder (*Hart*).

2. *As to the Rectum*.—Its upper part is completely invested by peritoneum; the second part is only partially covered, *i.e.* the peritoneum gradually leaves the rectum, quitting first the posterior surface, then the sides, and finally passing from the anterior surface on to the posterior vaginal wall.

See also Chapter II. on The Sectional Anatomy of the Female Pelvis, and especially Chapter III., p. 59.

Periton-
eum in
relation to
operations.

Practical Points.—Although the vesico-uterine pouch can be reached by a transverse incision through the anterior fornix, it will not be cut into in operations on the anterior vaginal wall. In the upper third or so of the posterior vaginal wall the peritoneum may be opened into. When the fingers are passed into the posterior fornix vaginae, only about $\frac{1}{3}$ inch of tissue intervenes between them and the peritoneum.

CONNECTIVE TISSUE OF PELVIS.

By this we understand (I.) the Fascia described so elaborately by the human anatomist as the Pelvic Fascia; and (II.) the loose Connective Tissue padding the interstices between the muscles, lying round the cervix uteri, and spreading out beneath the pelvic peritoneum.

Pelvic
Fascia.

I. The *Pelvic Fascia* of the anatomist is carefully described in the ordinary systematic and dissecting-room manuals, to which the student is therefore referred (*v.* also p. 11 and Chap. II.).

Pelvic Con-
nective
Tissue.

II. The *loose connective tissue* found lying subperitoneally, surrounding the cervix uteri and spreading out between the layers of the broad ligament, is of the highest importance pathologically, as in it and in the pelvic peritoneum occur those inflammatory exudations so common in women. Of late years our knowledge of the disposition of this tissue has been rendered much more accurate, and accordingly our discrimination of pelvic inflammatory attacks made much more precise.

Methods of
studying
it.

The distribution and relations of the pelvic connective tissue may be studied in various ways. The most valuable information is obtained by considering sections of frozen or spirit-hardened pelvises. This gives the precise position of the tissue, its amount, and distribution. Another valuable method of investigation is to inject air beneath the peritoneum, between the layers of the broad ligament, and at other points. By this we learn the varying attachments of the pelvic peritoneum to the subjacent tissue, and the lines of cleavage, as it were, of the pelvic connective tissue, along which pus will burrow. Instead of air we may inject plaster of Paris or water; plaster of Paris will be found the most useful.

We therefore consider—

- a. Results obtained by the injection of water, air, plaster of Paris ;
- b. Results obtained by section.

a. Results obtained by injections of water, air, or plaster of Paris.

The best summary of these results is given by Bandl, to whom on this point we are indebted for much valuable information.

König in his researches employed the bodies of women who had died a short time after labour from non-puerperal diseases, and injected air or water. The following briefly are his results :—

Connective
Tissue in-
vestigated
by injec-
tions.

(1.) Water injected between the layers of the broad ligament, high up in front of the ovary, passed first into the tissue lying at the highest part of the side wall of the true pelvis. It then passed into the tissue of the iliac fossa, lifting up the peritoneum, and followed the course of the psoas, passing only slightly into the hollow of the iliac bone. Lastly, it separated the peritoneum from the anterior abdominal wall for some little distance above Poupart's ligament, and from the true pelvis below it.

(2.) On injection beneath the base of the broad ligament to the side and in front of the isthmus, the deep lateral tissue became filled first; then the peritoneum became lifted up from the anterior part of the cervix uteri. The separation passed thence first to the tissue near the bladder, and ultimately the fluid passed along the round ligament to the inguinal ring. There it separated the peritoneum along the line of Poupart's ligament, and passed into the iliac fossa.

(3.) An injection at the posterior part of the base of the broad ligament filled the corresponding tissue round Douglas' pouch, and then passed on as described at (1.).

Schlesinger has followed out these results in more elaborate researches.

b. Results obtained by section.

The Sectional Anatomy of the Pelvis has now become a subject of such importance that it demands consideration in a separate chapter. The student will find at pp. 51, 52, reference made specially to the distribution of the connective tissue.

CHAPTER II.

THE SECTIONAL ANATOMY OF THE FEMALE PELVIS.

LITERATURE.

Barbour, A. H. F.—Spinal Deformity in Relation to Obstetrics: W. & A. K. Johnston, Edinburgh, 1884. *The Anatomy of Labour as exhibited in Frozen Sections*: W. & A. K. Johnston, Edinburgh, 1889. *Braune*—Topographisch-anatomischer Atlas, Zweite Auflage: Veit & Co., Leipzig, 1872. *Dwight*—Frozen Sections of a Child: Wood & Co., New York, 1883. *Hart, D. Berry*—Atlas of Female Pelvic Anatomy: W. & A. K. Johnston, Edinburgh, 1884. Supplement to Atlas of Female Pelvic Anatomy: W. & A. K. Johnston, Edinburgh, 1884. *Legendre*—Anatomie Homolographique: Paris, 1868. *His*—Ueber Präparate zum Situs Viscerum, etc.: Arch. für Anat., 1878. Die Lage der Eierstöcke, Arch. für Anat. 1881. *Lusckha*—Die Anatomie des menschlichen Beckens: Tübingen, 1864. *Pirogoff*—Anatome Topographica sectionibus per corpus humanum congelatum, etc.: Petropoli, 1859. *Ruedinger*—Topographisch-chirurgische Anatomie des Menschen: Stuttgart, 1873. *Simpson and Hart*—The Relation of the Abdominal and Pelvic Organs in the Female: W. & A. K. Johnston, Edinburgh, 1881. *Veit*—Die Anatomie des Beckens: Enke, Stuttgart, 1887. Pirogoff and Braune's Atlases are the great storehouse of Sectional Anatomy. Accounts of the method of freezing are given by Braune, Barbour, Simpson and Hart.

WHILE dissections are valuable in ascertaining the anatomy of any region, it must be remembered that they involve displacement of relations and therefore may lead into error or exaggeration. These may be corrected and additional accuracy obtained by making sections of frozen bodies or parts of them. If a body or a pelvis be covered with mackintosh and embedded in a mixture of salt and finely pounded ice or snow, it will in three or four days become as firm and solid as marble, and may then be sawn in any direction necessary. Tracings of the sawn surface may be made while it is still frozen; and in this way an accurate and trustworthy drawing may be obtained on which valuable measurements can be made.

We have said that the sections may be sawn in any direction, but usually they are made in special and definite lines as follows:—

(1.) *Sagittal Mesial*, i.e. parallel to the sagittal suture so that the body or pelvis is divided into right and left halves;

(2.) *Sagittal Lateral*, i.e. parallel and to one or other side of the sagittal mesial plane;




FIG. 1.




FIG. 2.

W. H. A. Johnston, Edinburgh & London.

POSITION OF UTERUS AND OVARIES.

FIG. 1. Sagittal Mesial Section of Pelvis (*Hart*).

FIG. 2. Fundus Uteri and Ovaries—Seen through the Pelvic Brim (*His*).



(3.) *Transverse or Horizontal*, i.e. at right angles to the long axis of the body, and with surfaces upper and lower;

(4.) *Coronal*, i.e. parallel to the coronal suture, dividing the body or pelvis into anterior and posterior portions, with surfaces anterior and posterior.

In sections of the pelvis alone, the axis of the brim is taken instead of the long axis of the body. We have therefore the following:—

(5.) *Axial coronal*, i.e. a section cut parallel to the axis of the brim and from side to side, with sawn surfaces anterior and posterior;

(6.) *Axial transverse*, i.e. at right angles to the axis of the brim and with surfaces therefore upper and lower.

We now take up the consideration of certain special sections.

1. *Sagittal Mesial Section.*

Plate I., fig. 1, shows a frozen sagittal mesial section of the pelvis with the uterus in position, the bowel and bladder naturally empty and the small intestine removed from the pouches so as to display the Fallopian tube and ovary. This section brings out the following facts: the uterus is not mesial but displaced somewhat to the left; the empty bladder is Y-shaped in sagittal mesial section; the urethra, vagina, and rectum are nearly parallel to one another and to the conjugate of the brim; the anus cuts these axes at right angles. The intestines have been removed from the Pouch of Douglas and vesico-uterine pouch. The nearness of the anterior abdominal wall to the promontory of the sacrum is well shown. The Perineal body is seen in section, and it should be noted that the greater part of it lies below the Hymen. Those gynecologists who exaggerate its functions usually draw it as being entirely behind the lower part of the posterior vaginal wall. Plate I. and fig. 23 shows that it does not do this. The student should note the peritoneal relations.

Plate I. also shows the relations of the Fallopian tube and ovary. When freshly cut, the intestines filled the peritoneal cavity; but after the section had been hardened in spirit, these were carefully lifted out so as to expose the ovary and Fallopian tube. The ovary lies with its long axis vertical, as His has pointed out. The preparation bears out his views completely with regard to the position of the ovaries, for on the other side of the body the ovary had its long axis somewhat transverse; and he has found that when the uterus was laterally displaced, the ovary of the side towards which the uterus was displaced lay vertical while the other ovary was somewhat transverse. In this cadaver the uterus lay to the left side and it is the left ovary which has its long axis vertical. The Fallopian tube does not form a loop enclosing the ovary as His found in his specimens (Plate I., fig. 2).

2. *Sagittal Lateral Section.*

Sagittal
Lateral
Section.

By this section a specially valuable view is obtained. Fig. 38 shows a drawing of a section at the junction of the uterus and broad ligaments; in it, although the pubes is divided mesially, the pelvic contents are cut to one side of the mesial plane. It should be noted that the amount of retropubic tissue is less than in the sagittal mesial one; that at the junc-




FIG. 38.

SAGITTAL MESIAL SECTION OF PELVIS cutting at Junction of Broad Ligament and Uterus.

a Vagina with its walls separated; *b* Bladder; *c* Symphysis; *d* Broad Ligament; *e* Ovary; *f* Fallopian Tube. In this specimen the Uterus was laterally displaced.

tion of the broad ligaments with the uterus there is a large amount of tissue with large blood-vessels; and specially that the finger placed in the lateral fornix vaginæ touches the base of the broad ligament there.

Connective Tissue of Broad Ligaments. This fact is valuable as to diagnosis. On section, the boundaries of the space between the broad ligaments are seen: superiorly the cut section of the Fallopian tube, anteriorly and posteriorly the peritoneum, and

inferiorly the vaginal fornix. The assertion by Guérin and Le Bec as to the insignificance of the tissue here is not borne out.

Sections made nearer the side pelvic wall display specially the lessening tissue between the layers of the broad ligaments and show sections of the ovary.

3. *Transverse or Horizontal Section.*

These give results confirming those above stated. Pirogoff gives several sections in his Atlas, but these are not clearly defined in their connective-tissue relations. Freund has published a very valuable series of preparations in his gynäkologische Klinik. The most valuable sections are those at the level of the supra-vaginal portion of the cervix, which show the tissue lying here all round it.

This is the best place to draw special attention to what Virchow first termed the parametric tissue. By this term he meant the loose fatless tissue ($\cdot 8$ in. thick), with abundant blood-vessels and lymphatics, surrounding "the lower portion of the uterus and the upper portion of the vagina" (*Spiegelberg*). This is the parametric tissue proper. Some extend the meaning of the term parametric tissue so as to include all the connective tissue in the pelvis.

4. *Coronal Section.*

Plate II. fig. 1, shows a coronal section of the pelvis passing through the base of the sacrum and the great trochanter. We note that the sacro-iliac joint runs from above downwards and inwards. The body of the sacrum bulges downwards, and the ischial tuberosity projects inwards so that the side wall of the pelvis is not straight: both of these are abnormalities. The anterior portion of the sacro-sciatic notch is seen. The levator ani is seen arising from the pelvic fascia over the obturator internus, and passing down to be inserted into the perineal body. The muscles of the perineum are also exposed. The body of the retroverted uterus is seen in great part, and lies perpendicular to the horizon; the frozen intestines have been removed so as to expose the fundus; the left Fallopian tube and round ligament have been divided as they pass forwards from the uterus. The left ovary has been partially cut across, and the removal of the intestines has exposed it entirely. Some cellular tissue is also exposed in the broad ligament; and there is some fatty cellular tissue external to this and continuous with the sub-peritoneal fatty tissue which lies external to the ovary and in the region of the sacro-sciatic notch. The uterus in this cadaver lay perpendicular to the horizon, and the ovary has the vertical position already described as a common one. The connective tissue between the bladder and the rectum is well seen as also its continuity with that in the broad ligament. This section explains clearly how a cellulitis when supplicated

may open into the vagina or pass through the sciatic notch to the hip. The levator ani and transversus perinei ending in the perineal body are clearly seen.

This section of the sacral plane does not show the bite or joggle described by Matthews Duncan ; but it is well seen in the next figure.

Plate II. fig. 2, shows a coronal section $\frac{3}{4}$ inch behind the preceding. In the bony pelvis we note, as has been said, that the sacro-iliac



FIG. 3

CORONAL FROZEN SECTION OF PELVIS (*Ruedinger*).



a Fundus uteri ; *b* Bladder ; *d* Labium minus ; *e* Labium majus.

joint shows the bite or joggle. The spine of the ischium has been divided where it gives origin to the levator ani ; the tuberosity is cut through in its posterior part, where it gives origin to the muscles. The levator ani is seen arising from the ischial spine and passing downwards to be inserted into the rectum at the external sphincter. External to it lies the ischio-rectal fossa, which extends upwards as far as the ischial spine ; internal to it, a well-marked layer of the pelvic fascia is displayed. The uterus has been sliced across from the ovarian ligament

CORONAL SECTIONS OF LEFT HALF OF PELVIS—Seen from the front (Barbour).

FIG. 1.

FIG. 2.





to below the utero-sacral ligament ; the intestines seen above it occupy




FIG. 40.

AXIAL CORONAL SECTION OF PELVIS—seen from behind (Hart).




FIG. 41.

AXIAL CORONAL SECTION OF PELVIS—seen from behind (Hart).

the highest part of the pouch of Douglas. The peritoneum of the pouch of Douglas has been cut across in two places,—where it covers the body

of the uterus about the level of the ovarian ligaments, and also 1.3 cm. ($\frac{1}{2}$ in.) above the bottom of the pouch of Douglas.

We observe in this section the boundaries of the ischiorectal fossa, and the continuity of the tissue in the broad ligament with that in front of the sacrum.

At fig. 39 is shown the relations of the pelvic organs in Ruedinger's coronal section of a female cadaver.

5. Axial Coronal Section of Pelvis.

Axial
Coronal
Section.

Fig. 41 is an axial coronal section made $1\frac{1}{2}$ inch behind the pubes and passing through the hip joints. This pelvis was not normal, as there was a cellulitis of the left broad ligament and a displacement of the bladder to the right side. The section is viewed from behind. Owing to a slight distention of the bladder the uterus lay in the axis of the brim and has been divided coronally. The left broad ligament has been divided similarly, so that its side relations to the obturator internus are displayed. The vagina is a crescentic slit, the side limits of the vaginal portion of the cervix being marked *xx*. The levatores ani are seen springing from the pelvic fascia and curving downwards and inwards below the rectum. We see that here the boundaries of the ischiorectal fossa are gluteus maximus, below; levator ani, above and to the inner side; and obturator internus, above and to the outer side. On the right side, the ureter has been cut as it lies in the bladder wall, $\frac{7}{8}$ inch distant from the vagina. On the left side it is about one inch from the vagina. This section exhibits the side relations of the broad ligament, the continuity of the connective tissue between the layers of the broad ligament with that in front of the iliacus muscle, and the accurate packing, as it were, of the pelvic viscera.

Fig. 40 gives a section similar in direction to the preceding, but about one inch farther back so that it grazes the posterior surface of the uterus.

The Pouch of Douglas is cut into at one part. The left broad ligament is shortened by the cellulitis already mentioned. The ischiorectal fossa is seen at its most posterior part, and is very small, being roofed in by the levator ani, its floor being formed by the gluteus maximus. The divided ureters are seen lying in the loose fatty tissue outside the broad ligaments.

CHAPTER III.

THE POSITION OF THE UTERUS AND ITS ANNEXA, AND THE RELATION OF THE SUPERJACENT VISCERA.

LITERATURE.

Bandl—Ueber die normale Lage und das normale Verhalten des Uterus und die pathologisch-anatomischen Ursachen der Erscheinung Antelexio: Arch. f. Gynäk., Bd. XXII., S. 408. *Braune*—Topograph. Anatom. Atlas, Zweite Auflage: Leipzig, Veit & Co., 1872. *Claudius*—On the Position of the Uterus: Med. Times and Gazette, 1865, p. 5. *Credé*—Beiträge zur Bestimmung der normalen Lage der gesunden Gebärmutter: Archiv f. Gynäkologie, Bd. I., S. 84. *Foster*—A Contribution to the Topographical Anatomy of the Uterus and its Surroundings: Am. J. of Obst. XIII., p. 30. *Hart, D. B.*—Atlas of Female Pelvic Anatomy: W. & A. K. Johnston, Edinburgh, 1884. See also Supplement to Atlas. *Hasse*—Beobachtungen über die Lage der Eingeweide im weiblichen Beckeneingange: Archiv. f. Gynäk., Band viii., S. 402. *His*—Ueber Präparate zum Situs Viscerum u.s.w.: Arch. für Anat., 1878, S. 53. Also, Die Lage der Eierstöcke in der weiblichen Leiche: Arch. für Anat., 1881, S. 398. *Pirogoff*—Anatome Topograph. etc.: Petropoli, 1859. *Sappey*—Traité d'Anatomie Descriptive: Paris, 1873. *Schroeder*—Handbuch der Krankheiten der weiblichen Geschlechtsorgane: Leipzig, 1879. *Schultze*—Die Pathologie und Therapie der Lageveränderungen der Gebärmutter: Berlin, 1881. *Symington, J.*—The Topographical Anatomy of the Child: Edinburgh, Livingstone, 1887. *Waldeyer*—An admirable account of the subject will be found in Van de Warker's articles on a study of the Normal Movements of the Unimpregnated Uterus: N. Y. Medical Journal, Vol. XXI., p. 337. Also, On the Normal Position and Movements of the Unimpregnated Uterus: Am. J. of Obst., Vol. XI., p. 314.

THE amount of literature, chiefly French and German, on this subject is much too extensive even to be mentioned here, for the position of the uterus has given rise to much discussion. This is partly due to the inherent difficulty of accurate clinical observations, to the erroneous opinions advanced by many eminent anatomists, and to arbitrary demands as to the normal uterine position made by gynecologists with strong opinions on anteversion.

Thus, in the well-known works of Braune, Luschka, Cruveilhier, and Henle, the uterus is figured from actual sections as normal with the fundus in the hollow of the sacrum, *i.e.*, retroposed. Claudius of Marburg, also an anatomist, is uncompromising on this point. He states, indeed, that the uterus is normal only when, with its broad ligaments, its posterior surface touches the sacrum as closely as the lungs do the

Difference
in opinions
as to posi-
tion of
Uterus.

ribs (fig. 42). Now, almost all gynecologists agree, from clinical observation, that the body of the uterus lies over on the bladder, with the os uteri looking more or less back. This divergence of opinion is extraordinary; and it leads to this interesting practical observation, that what the anatomist considers a uterus normal in position, the gynecologist believes to be abnormal. That is, the retroverted uterus—considered normal in cadavera by the anatomist—is, in certain cases, when found




FIG. 42.

TRANSVERSE SECTION of PELVIS in line of PYRIFORM MUSCLES (*Luschka*). The Peritoneum has been removed on the right side. *a* 3d Sacral Vertebra; *b* Bladder; *c* Ureter; *d* Levator Ani; *e* Rectum; *f* Anterior Layer of Broad Ligament; *g* Uterus; *h* Pyriform Muscle. Note that here the uterus is retroverted, and the pouch of Douglas without intestine.

in the living subject, replaced by the gynecologist so that it lies with its body over the bladder.

There can be no doubt that the uterus lies normally to the front with its anterior surface resting on the bladder. Great refinement is exercised, quite unnecessarily, by many gynecologists in settling what they believe to be the exact angle which the long axis of the uterus should make with the horizon, when a woman is in the erect posture; and this refinement has been greatly stimulated by the mechanical treatment of what is known by many as anteversion of the uterus.

In treating of this vexed question, we shall consider—

1. The normal form and position of the uterus;

2. The local divisions of the pelvic-floor peritoneum as viewed through the pelvic brim, and the position of the uterus and its annexa ;
3. The physiological changes in the position of the uterus ;
4. The relation of the small intestine to the pelvic floor and to the uterus and its annexa.

THE NORMAL FORM AND POSITION OF THE UTERUS.

The question of the *form* of the uterus we consider only in the limited Normal aspect of the angular relation of the long axis of the uterus to the long form of Uterus.



FIG. 43.

DIAGRAM to show Normal Form and Position of VIRGIN UTERUS (Schultze).

axis of the cervix. These are not in the same straight line, but, when the bladder and rectum are empty, lie at an obtuse angle of varying value. This angle is more open in multiparous women (fig. 25), than in nulliparæ (fig. 43).

The question as to whether in the normal uterus the cervix and body are in the same straight line or meet at an angle opening anteriorly, is much disputed and by no means easy to settle. Bimanually, the normal uterus is fairly often found anteverted, but the question arises whether the Bimanual examination has not brought about or at any rate exaggerated the anteversion. Bandl asserts that when the uterus is removed

and examined *post mortem*, ante flexion is rarely found, the normal uterine axis being straight. It should be remembered however that the removal of the uterus from the body involves the cutting of the utero-sacral ligaments and the absence of intra-abdominal pressure, *i.e.*, removes the conditions in the living subject which keep up "physiological ante flexion;" so that a uterus somewhat ante flexed during life may be




FIG. 44.

SECTION OF PELVIS, showing UTERUS driven back by distended Bladder, and Peritoneum disturbed (Kohlrausch). This is not a normal condition of parts by any means.

straightened by removal *post mortem*. The best way to ascertain the existence of ante flexion in the living woman is to use simple vaginal examination. The question really is as to the normal form of the uterus in the living woman with the peritoneal folds intact and intra-abdominal pressure in action. Under these conditions there is a normal degree of

anteflexion which is called "Physiological anteflexion," in contrast with Schultze's "Pathological anteflexion," so commonly caused by utero-sacral cellulitis (*v.* also Chap. on Displacements of the Uterus).

The *position* of the uterus, with empty bladder and rectum, is such that it lies with its anterior surface touching the posterior aspect of the bladder, no intestine usually intervening; the os externum uteri looks downwards and backwards: and the uterus is slightly twisted as a whole on its long axis, so that the uterine end of the right Fallopian tube is nearer the symphysis than that of the left. We have expressly said with bladder and rectum empty. According to Schultze, the long axis of the uterus is nearly parallel to the horizon. This is probably exaggerated, as Schultze's researches were conducted in a way that certainly anteverted the uterus unduly (figs. 25 and 43). Many authors figure the uterus nearly vertical to the horizon, for this purpose distending the bladder until the uterus is elevated to what they consider the proper angle (fig. 44). It is needless to say how absurd this is. Kohlrausch's diagram, so often quoted in support of this allegation, really shows, if it show anything, the position of the uterus when the bladder is well distended. The student should note this point, as Kohlrausch's section is the favourite diagram of those who treat as pathological what is really a normal uterus.

Normal position of Uterus.

It is important to know how results as to the uterine position have been obtained. The chief methods are as follows:—

(1.) *By frozen, spirit-hardened, or chromic-acid sections.*—Results obtained in this way are valuable, if we make allowance for some *post-mortem* change in the uterine position not yet thoroughly understood.

Methods of investigating position of Uterus.

(2.) *By the bimanual examination of the pelvic contents.*—This is probably the best method, although it exaggerates the normal anteversion of the uterus in a way that will be readily understood when the chapter on the Bimanual has been studied.

(3.) *By the use of the sound*, or by a more elaborate means described by Schultze. Space does not permit of a full description of the latter, but a good account of it is given in Foster's paper.

THE LOCAL DIVISIONS OF THE PELVIC-FLOOR PERITONEUM AS VIEWED THROUGH THE PELVIC BRIM, AND THE POSITION OF THE UTERINE ANNEXA.

For valuable papers and sections on this subject, we are indebted to Hasse, Waldeyer, and others (fig. 45). Hasse froze not quite thoroughly a female cadaver in the upright posture, cut through the abdomen transversely, and then lifted out the softened viscera until the pelvic contents were exposed undisturbed. The bladder was moderately distended.

Pelvic
Contents
as seen
through
the Brim.

Fig. 45 shows Hasse's drawing. The fundus of the uterus lying on the bladder is well seen. In front of the broad ligament—of which the infundibulo-pelvic ligament is the only portion visible in fig. 45—we have, on each side, the paravesical pouch of the peritoneum. Behind it, lies the lateral pouch of Douglas; while just behind the uterus and




FIG. 45.

FEMALE PELVIS and CONTENTS viewed through the Pelvic Brim (*Hasse*).

v Bladder; *II* Paravesical Pouch; *u* Uterus; *o* Ovary; *t* Fallopian Tube; *d* Pouch of Douglas; *I* Lateral Pouch of Douglas; *ip* Infundibulo-pelvic Ligament; *lr* Round Ligament; *p u* Position of Ureter; *lo* Ovarian Ligament; *r* Rectum; *c* Colon.

bounded on each side by the utero-sacral ligament is the pouch of Douglas proper. The Fallopian tubes lie in the true pelvis, in the paravesical pouch. Each broad ligament sweeps outwards and backwards to near the sacro-iliac synchondrosis of its own side. The position of the ureter is well indicated.

According to Hasse the long axis of each ovary runs outwards and forwards, forming with the transverse axis of the uterus an angle open to the front. Part of each ovary (the half) projects above the plane of the pelvic brim. Schultze figures the ovaries as having their long axes almost antero-posterior (fig. 46), and His in his cases found the long axes

Direction
of Ovaries.




FIG. 46.

POSITION OF FUNDUS UTERI and lie of OVARIES. Bladder distended (*Schultze*).

nearly vertical. In recent sections, the authors found the ovary lying nearly vertical as His describes (*v. Pl. I.*). The long axis of the ovary on the side to which the uterus is displaced is nearly vertical, while the ovary of that side from which the uterus is displaced is more transverse (*v. page 26, and Pl. I., fig. 2*).

THE PHYSIOLOGICAL CHANGES IN THE POSITION OF THE UTERUS.

The mobility of the uterus is one of its most characteristic features. With every movement of respiration, in singing, in walking, and in all violent movements, the uterine position is changed. Van de Warker has studied, in a valuable paper, the influences bringing about these changes in position; this may be consulted for details of his method of investigation and results obtained.

Of the greatest importance is the effect of the distended bladder on the uterine position. As the bladder fills, the uterus becomes retroposed to an extent shown at figs. 44, 45, and 47. The intestines are forced out of the upper part of Douglas' pouch, and the height of the peritoneal reflection from the anterior abdominal wall is considerably increased. All these points are well illustrated by fig. 37 from Pirogoff. As the urine is evacuated, the uterus passes forward to its normal anteverted

Effect of
Bladder
on position
of Uterus.

condition and the intestines pass back into Douglas' pouch. Rectal distension displaces the uterus forwards and to the right side.

THE RELATION OF THE SMALL INTESTINE TO THE PELVIC FLOOR
AND TO THE UTERUS WITH ITS ANNEXA.

Relation
of small
Intestines
to Uterus.

The small intestine lies resting on the uterus, ovaries, Fallopian tubes, and broad ligaments. There is usually no small intestine in the vesico-




FIG. 47.

POSITION OF UTERUS. *A* with bladder and rectum empty; *B, C, D* according to distention of bladder (*Van de Warker*).

uterine pouch. When the bladder is empty and the unimpregnated uterus to the front, there is small intestine in Douglas' pouch except at its very lowest part. The pouch of Douglas becomes emptied of intestine as the bladder distends, and has no intestine in it when the uterus is retro-

verted. Many authors assert that there is never small intestine in Douglas' pouch. This opinion is undoubtedly wrong, as any one can satisfy himself by studying sections. Often Douglas' pouch contains serum, and this displaces the intestine. Figures 36 and 37 bear out these opinions; fig. 39 should be carefully studied as illustrating the position of the superjacent intestines. The paravesical pouch probably contains intestine when the uterus lies to the front, and certainly contains it when the uterus is pathologically retroverted. Occasionally, the omentum may interpose between the small intestine and the pelvic viscera.

To sum up briefly :—

- a. The uterus and bladder behave practically as one organ *quâ* position (i.e., they move together), when the uterus is to the front.
- b. The exact angle which the uterus makes with the horizon cannot be fixed, and knowledge on this point is not necessary.
- c. The uterus lies normally to the front, but has a range of mobility indicated in fig. 47. The posterior lip of the cervix is .6 to 1.2 in. (1.5 to 3 cm.) above the tip of the coccyx. By digital pressure the uterus can be raised considerably.

Summary
as to position of
Uterus.

CHAPTER IV.

THE STRUCTURAL ANATOMY OF THE FEMALE PELVIC FLOOR: THE PELVIC-FLOOR PROJECTION.

LITERATURE.

- STRUCTURAL ANATOMY. *Hart*—The Structural Anatomy of the Female Pelvic Floor : Edinburgh, 1881. Atlas of Female Pelvic Anatomy : 1884. Supplement to ditto : 1885. W. & A. K. Johnston, Edinburgh and London. *Symington*—A Contribution to the Normal Anatomy of the Female Pelvic Floor : Edin. Med. Jour., March, 1889.
- PELVIC-FLOOR PROJECTION. *Foster*—Topographical Anatomy of Uterus, etc. : Am. J. of Obst., Vol. XIII., p. 30. *Schröder*—Noch ein Wort über die normale Lage und die Lageveränderungen der Gebärmutter : Arch. f. Gynäk., Bd. IX., S. 68. *Schultze*—Zur Kenntniss von der Lage der Eingeweide im weiblichen Becken : Archiv f. Gyn., Bd. IX., S. 262. *Simpson and Hart*—The Relation of the Abdominal and Pelvic Organs in the Female : W. & A. K. Johnston, Edinburgh and London, 1881.

THE STRUCTURAL ANATOMY OF THE FEMALE PELVIC FLOOR.

Structure of Pelvic Floor with regard to function.

HITHERTO we have regarded the pelvic floor in detail as made up of bladder, vaginal walls, rectum, connective tissue, and peritoneum. In this chapter we purpose considering it in its structural aspect. In its formation, the following functions have been provided for. As compared with the floor of the male pelvis, the female pelvic floor differs in having in it the cleft known as the vagina. Then further, women have to undergo parturition in which the child is born through the vagina, which is then greatly distended. At the same time a woman has resting on her pelvic floor the same abdominal viscera as the male, and her pelvic floor is also subjected to the same strain from intra-abdominal pressure. Thus we have to explain how the female pelvic floor has been constructed so as to allow of parturition and the rectal and vesical functions and yet remain strong enough to resist ordinary intra-abdominal pressure. The question is a *structural* or architectural one. We study it in this present chapter just as we should study the structure of a box or chair.

In order to understand this question, we must study the pelvic floor as seen both in *sagittal mesial* and in *axial coronal* section.

a. Sagittal Mesial Section.

In this view (*cf.* Pl. I.) we see the pelvic floor or diaphragm stretching from symphysis pubis to sacrum. The anus is to be imagined closed as in life. The first thing to note is the vagina, which is seen as a cleft running upwards in the pelvic floor from hymen to cervix uteri. Its walls are in close apposition (*vide* figs. *passim*). They are often erroneously represented apart; in order, as it were, to let the student see the vagina. This is wrong, however. It is no more necessary to figure the vaginal walls always apart, than it would be always to sketch a man with his mouth open to render it visible. The first idea one gets on looking at a frozen section is that, owing to the apposition of the vaginal walls, the pelvic-floor is unbroken; and that the vaginal cleft, the introduction of which does weaken the floor somewhat, cuts it not perpendicularly to the horizon but obliquely at an angle of about 60°. Its appearance in Sagittal Mesial Section.

The pelvic floor, as seen in this section, is made up of two segments which are known as the *pubic* and *sacral* segments. It is of importance to define these exactly.

The *Pubic Segment* is made up of loose tissue, viz., bladder, urethra, anterior vaginal wall, and bladder-peritoneum. It is attached in front to the symphysis pubis. This attachment is a loose one; the bladder and urethra, meeting one another at right angles, are separated from the pubes by the pyramidal deposit of loose fat already described as the retropubic fat deposit. Note specially that the retropubic fat deposit as seen in this section—that of a subject in the dorsal or the erect posture—is triangular; and that the peritoneum passes from the anterior abdominal wall on to the fundus of the bladder, just a little above the top of the symphysis. Below the pubic arch, the urethra becomes blended with the perineal muscles there. The Pubic Segment.

The *Sacral Segment* is attached to the coccyx and sacrum; it consists of rectum, perineum, posterior vaginal wall, and strong tendinous and muscular tissue. The inferior portion of this segment, the perineum lies about 1½ inches from the symphysis. The Sacral Segment.

In addition to the retropubic fat deposit, it should be noted that—

- a. The posterior wall of the bladder is *loosely* attached to the anterior vaginal wall;
- b. The urethra and anterior vaginal wall are *closely* blended;
- c. The posterior vaginal wall and anterior rectal wall are *loosely* connected as far down as the apex of the perineal body (fig. 33).

The two segments, as seen in sagittal mesial section, are thus anatomically contrasted:— The Segments contrasted.

The pubic segment is made up of loose tissue, and is loosely attached to the pubic symphysis; the sacral segment is made up of dense tissue and is firmly dovetailed into the sacrum and coccyx.

They are further contrasted *functionally*:—

The pubic segment is drawn up during labour; the sacral segment is driven down.

The proof of this functional contrast is too elaborate to be given here, but will be found in detail in Hart's Atlas. Briefly stated it




FIG. 48.

PELVIC FLOOR differentiated in parturition (*Braune*). The Pubic Segment is drawn up and the Sacral one driven down. Note position of bladder and its peritoneum.

is that during labour the pubic and sacral segments as seen in a sagittal mesial section may be likened to two folding doors. Uterine action pulls up the pubic segment, and drives the child down against the sacral one. This action is analogous to the way one passes out through two

folding doors, when he pulls the one door towards him and pushes the other from him. As the result of this elevation of the pubic segment, the bladder is drawn in part above the pubes and its peritoneum stripped off (fig. 48).

The various components of the pubic segment are definitely displaced in its movements. Thus the retropubic fat is—

Displacement of the Pubic Segment.

1. Behind the pubes in the non-parturient female (fig. 36);
2. Above it in the parturient female (fig. 48);
3. Below it in prolapsus uteri;
4. Below it in the extra pelvic-floor projection of pregnancy;
5. Partially above the symphysis in the genupectoral posture (fig. 63).

The peritoneum is—

1. Reflected on to the top of the empty bladder in the non-parturient female;
2. Stripped off the bladder during parturition;
3. Reflected on to fundus of empty bladder, at a higher level above symphysis, in the genupectoral posture.

Thus the peritoneum over the bladder is movable; the peritoneum over the sacral segment is fixed.

b. Axial Coronal Section.

If now we study axial coronal sections, we shall find these views (based on sagittal mesial) both enlarged and modified. If actual sections such as are shewn in figs. 40 and 41, be examined it will be found that, owing to the presence of loose tissue, a line of cleavage runs within the obturator internus, upper part of the levator ani, and rectum, separating these structures from the vagina. We thus find a complete ring of loose tissue of which part has been seen in sagittal mesial section and part in axial coronal section. This ring of loose tissue runs as follows:—beginning behind the pubes (retropubic fat), it passes on the internal aspect of the obturator internus and upper portion of Levator ani of the left side; between the posterior vaginal and anterior rectal walls; on the inner aspect of the obturator internus and upper portion of the Levator ani of the right side; and then back to the retropubic fat. This ring of loose tissue divides the pelvic floor into two portions:—

Axial Coronal Section.

a. The entire displaceable portion;

b. The entire fixed portion.

a. The entire displaceable portion comprises bladder, urethra, and vaginal walls. It has resting upon it the uterus, broad ligaments, Fallopian tubes, and ovaries; and lies within the ring of loose tissue.

b. The entire fixed portion lies outside of the ring of loose tissue. If

the entire displaceable portion were cut out of the pelvic floor, then on looking through the pelvic brim, we should see, *in front*, the posterior aspect of the pubes, sloping downwards and backwards; *at the sides*, the inner aspects of the obturator internus sloping downwards and inwards; and *behind*, the anterior rectal wall and sacrum sloping downwards and forwards. We should, in fact, be looking down into a funnel whose walls all sloped towards a central point. This funnel forms the entire fixed portion of the pelvic floor.

It will now be understood that the entire fixed portion supports the entire displaceable portion; and that consequently on these two combined (*i.e.*, the whole pelvic floor) the uterus and annexa and the abdominal viscera rest.

Divisions
of Pelvic
Floor.

The terminology given need not confuse if it be remembered that the terms "pubic segment and sacral segment" apply to sagittal mesial sections only, and are applicable to the mechanism of parturition; while "entire displaceable and entire fixed portions" apply to transverse sections, and are to be used for the general physics of the pelvic floor and for prolapsus uteri. The relation between the two views given by sagittal mesial section and by transverse (or by axial coronal) section may be represented as follows:—

Sagittal Mesial Section.

Transverse or Axial Coronal Section.

Pubic Segment.	{ Bladder and urethra, Anterior vaginal wall,	} Entire displaceable portion.
Sacral Segment.	{ Posterior vaginal wall, Tissue attached to sacrum, Bowel in pelvic floor, All outside of inner aspects of levator ani.	
		} Entire fixed portion.

Functions
of Pelvic
Floor.

The chief functions demanded of the female pelvic floor are—

- a. *Support of Intra-abdominal Pressure,*
- b. *Vesical and rectal functions,*
- c. *Parturition.*

a. *Support of Intra-abdominal Pressure.* The abdominal and pelvic viscera rest on the pelvic floor; more correctly, these viscera (along with the entire displaceable portion of the pelvic floor) rest on the entire fixed portion of the pelvic floor, the inward convergence of whose parts enables them to support these. Prolapsus uteri is thus, as we shall afterwards see, not a mere uterine descent, but a downward displacement of the abdominal and pelvic viscera along with the entire displaceable portion of the pelvic floor.

b. Vesical and rectal functions. The loose tissue round the rectum and bladder allows of the contraction and diminution in bulk of these organs which are necessary for the expulsion of their contents.

c. Parturition. This is the great function of the pelvic floor, and is provided for structurally as follows. The child is driven through the vagina (*i.e.* through the entire displaceable portion) by the upward tension of the uterine muscle attached to the top of the vaginal walls and by the dilating pressure of the foetal head. This upward movement of the entire displaceable segment is allowed by the ring of loose tissue of which we have spoken. We are now able to understand the full significance of the statement already made that the pubic segment of the pelvic floor is pulled up partly into the abdominal cavity while the sacral segment is driven downwards and backwards. In addition, the levatores ani will be pressed outwards.

The result of parturition is (1) To dilate the vaginal walls and render them more easily everted, (2) to tear the inferior margin of the sacral segment, *i.e.* the perineum, (3) to elongate and slacken the ring of loose tissue uniting the entire displaceable and the entire fixed portions. In this way, it favours that driving downwards and outwards of the entire displaceable portion which happens in Prolapsus uteri.

PELVIC-FLOOR PROJECTION.

By this is understood the amount of projection of the pelvic floor, in Definition sagittal mesial section, *beyond the straight line joining the tip of the coccyx and the subpubic ligament—i.e., beyond the conjugate of outlet* (fig. 49). of Pelvic Floor Projection.




FIG. 49.

DIAGRAM to show what is meant by PELVIC-FLOOR PROJECTION. *ap*=conjugate of outlet. A perpendicular bisecting *ap* and cutting the arc gives the greatest pelvic-floor projection (*F. P. Foster*).

Definite results have not as yet been obtained, but this is one special reason why attention should be directed to it.

Schroeder measured the conjugate at the outlet with callipers; and then passed a measuring line from the coccyx to the apex of the pubic arch, the tape following the curve of the pelvic floor. The subjoined table gives some of his results.

	Distance from tip of coccyx to lower border of symphysis.	
	By Tape Measure.	By Callipers.
Average of the pregnant woman	cm. 13.35	9.15
„ „ gynecological patients	„ 12.6	8.27
„ „ nulliparæ	„ 13.2	9.75

Schroeder's deduction is that the average projection of the pelvic floor beyond the plane of the pelvic outlet is 4.1 cm. There is no doubt that this is an excessive average.

Mode of
measuring
pelvic
floor pro-
jection.

F. P. Foster of New York has written ably on this subject, and made a large series of observations. Fig. 50 shows the callipers he




FIG. 50.

CALLIPERS for measuring PELVIC-FLOOR PROJECTION (*Foster*).

employed. The ends of the limbs (*a* and *b*) are placed on the tip of the coccyx and lower border of the symphysis pubis, respectively. The horizontal bar between these limbs is graduated in cm., and the limb (*a*) glides along it in a groove. A movable upright (*c*), also graduated, has its upper point placed against the most projecting part of the pelvic floor. If now the whole apparatus be removed and laid flat on a sheet of paper, the conjugate and amount of projection can be read off at once. Greater accuracy is ensured by noting, before removing the

apparatus, the point on the transverse bar at which the upright (*c*) stands as well as the reading which it gives.

Foster's average (2.5 cm.) of the pelvic-floor projection is less than Schroeder's. He placed the patient semiprone, however; a position in which the pelvic-floor projection is slightly diminished. Fig. 51 shows Foster's diagram of pelvic-floor projection. The uterus is more anteverted than in Foster's original drawing.

Measurements made on frozen sections must be used with caution. Schroeder has justified his average by such measurements, but has taken no account of the existence of pregnancy in some of the cases.




FIG. 51.

DIAGRAM OF PELVIC-FLOOR PROJECTION and position of uterus, modified from *Foster*.
The anterior and posterior walls of the anus are not in apposition, as shown in the diagram, and the uterus is too high in the pelvic cavity.

We might tentatively advance the following statements:—

- (1.) The pelvic-floor projection is over-estimated by Schroeder;
- (2.) Foster's average is nearer the mark;
- (3.) The pelvic-floor projection is increased by advanced and even by early pregnancy (Braune's Plates).

Summary
as to pelvic
floor pro-
jection.

The whole inquiry needs further investigation in order to settle also other points, among which we may mention the relation of the vagina to the pelvic outlet and the varying amount of pelvic-floor projection in different postures.

CHAPTER V.


THE BLOOD-VESSELS, LYMPHATICS, AND NERVES OF THE PELVIS: DEVELOPMENT OF PELVIC ORGANS.


LITERATURE.

- BLOOD-VESSELS.** *Henle*—Handbuch der Eingeweidelehre des Menschen : Braunschweig, 1866. *Hyrtl*—Die Corrosions-anatomie und ihre Ergebnisse : W. Braumüller, Wien, 1873. *Klein and Smith's Atlas*. *Luschka*—Die Musculatur am Boden des weiblichen Beckens : Wien, 1861. *Quain*—Anatomy, 8th Ed. : London, 1882. *Sappey*—Traité d'Anatomie Descriptive : Paris, 1873. *Savage*—Female Pelvic Organs, 5th Ed. : London, 1882. *Tait, Lawson*—The Pathology and Treatment of Diseases of the Ovary, p. 8. : Birmingham, 1883. *Sir J. Williams*—On the Circulation in the Uterus, with some of its Anatomical and Pathological Bearings : Obstet. Soc. of London, 1885.
- LYMPHATICS.** *Bourguery and Jacob*—Traité Complet de l'Anatomie de l'Homme : Paris, 1839. *Championnière*—Lymphatiques Utérines et Lymphangite Utérine : Bull. de la Soc. Med. des Hôpitaux de Paris, Vol. VII. *Curnow*—The Lymphatic System and its Diseases : Lancet, 1879. *Klein*—The Anatomy of the Lymphatic System : London, 1873. *Le Bec*—Contributions à l'Etude des Ligaments larges, etc. : Gaz. Heb., 15 Avril, 1881. *G. Leopold*—Die Lymphgefäße des normalen nicht schwangeren Uterus : Archiv für Gynäk., Bd. VI., S. 1. *W. T. Lusk*—Puerperal Fever : International Congress Tr., Philad., 1877. *Poirier*—Du rôle des Lymphatiques dans les inflammations de l'utérus des annexes et du péritoine pelvien. Le Progrès Medical, Janv. 25th, 1890, et seq. *Sappey, Savage*—Op. cit.
- NERVES.** *T. Snow Beck*—The Nerves of the Uterus : Phil. Tr., 1846. *Flower*—Diagrams of the Nerves of the Human Body : J. and A. Churchill, London, 1872. *Frankenhäuser*—Die Nerven Der Gebärmutter und ihre Endigung in den glatten Muskelfasern : Jena, 1867. *Gawronsky, V.*—Ueber Verbreitung und Endigung der Nerven in der weiblichen Genitalien. Arch. für Gynäk. Bd. 47, S. 271. *Griffiths*—Observations on the Urinary Bladder and Urethra in Part II., The Nerves, in Part I. : Jour. of Anat. and Phys., Oct. 1894. *Head*—On Disturbance of Sensation with special reference to the pain of visceral disease : Brain, 1893. *Jastrebow*—On the normal and pathological Anatomy of the ganglion cervicale uteri : Lond. Obstet. Trans., 1881. *R. Lee*—The Anatomy of the Nerves of the Uterus : London, Bailliére, 1841. *Tiedemann*—Tabulæ Nervorum Uteri : Heidelberg, 1822.
- DEVELOPMENT OF THE PELVIC ORGANS.** *Foulis*—The Development of the Ova, and the Structure of the Ovary in Man and the other Mammalia : Tr. R. S., Edin., Vol. XXVII. *Hart, D. Berry*—A Preliminary Note on the Development of the Clitoris, Vagina, and Hymen : Jour. of Anatomy and Physiology, Oct. 1896 ; also Edin. Obst. Trans., 1896. *Klein and Smith, Quain, Turner*—Op. cit.

BLOOD-VESSELS.

Preliminary Remarks:—The blood supply to the pelvic organs and perineum is derived from the ovarian arteries (which are branches of the abdominal aorta), and from the uterine, vaginal, and internal pudic





arteries (which are all branches of the anterior division of the internal iliac).

We shall first consider the arterial supply of the uterus, ovary, Fallopian tubes, vagina, bladder, rectum, and that of the perineal region ; and then the venous distribution.

ARTERIAL SUPPLY.

(1.) *Arterial supply to uterus, ovary, etc.*—The *Ovarian artery* of each side (corresponding to the spermatic of the male) is a branch of the abdominal aorta. Its relations when in the abdomen do not concern us here. In the pelvis it passes between the layers of the broad ligament, running tortuously towards the upper angle of the uterus. Near this it divides into two branches. The upper supplies the fundus uteri ; the lower anastomoses at the side of the uterus with the uterine artery (Plate III. *c, d*). Arterial supply to Uterus and Ovary.

The Ovarian artery gives off—

Branches to the ampulla of the Fallopian tube (Plate III. *a' a'*),

Branches to the isthmus (*b'*).

Numerous branches to the ovary (*c' c' c'*).

Branch to the round ligament (*b*).

The *Uterine Artery* (Plate III. *e*) springs from the anterior division of the internal iliac, and passes downwards and inwards towards the cervix uteri. It then passes upwards between the layers of the broad ligament by the side of the uterus, in an exceedingly tortuous manner well shown in Plate III., to anastomose with the lower branch of the ovarian. The course of the blood-vessels in the uterine wall has been studied and described by Sir J. Williams with special reference to some anatomical and pathological points. The primary branches after entering the uterine tissue have a somewhat superficial course, being separated from the peritoneum by only a thin layer of muscular fibres. From these, secondary branches run towards the mucous surface in a direction perpendicular to that surface ; these anastomose freely and end in capillary loops in the mucous membrane. All internal to the primary branches—the greater part of the muscular wall—belongs, according to Williams, to the mucous membrane, *i.e.*, to the muscularis mucosæ. The *Vaginal arteries* (*g g g*) usually spring immediately from the anterior division of the internal iliac artery, but sometimes arise from the uterine or middle hæmorrhoidal. A special branch of the uterine artery to the cervix joins with its fellow at the isthmus to form the circular artery, and with those of the vagina to form the azygos artery of the vagina (*h h*). The vaginal arteries of one side anastomose freely with those of the other. Plate III., from Hyrtl, illustrates beautifully the free anastomosis of branches of the aorta with the ovarian, uterine, and vaginal arteries. It should be noted that, in operation

for removal of the uterus, ligature of the broad ligament controls all hæmorrhage.

From the same anterior division of the internal iliac proceeds the blood supply to the bladder and rectum.

Arterial
supply of
Perinum.

Arterial supply to the perineal region.—This comes from the internal pudic. The superficial perineal branch supplies the labia; the artery to the bulb supplies the bulbus vaginæ; the terminal branches go to the clitoris.

VENOUS SUPPLY.

Veins of
Pelvis.

The venous supply of the pelvis is very abundant, and exists in the form of numerous plexuses freely communicating with one another. The veins are unprovided with valves; hæmorrhage from a wound is therefore often exceedingly profuse, especially during pregnancy, when the whole pelvic vascular system is hypertrophied.

The following is a summary of the main facts as to the venous supply of the female pelvis.

The *Vesical plexus* lies external to the muscular coat of the bladder. The *Hæmorrhoidal plexus* lies below the mucous membrane of the lower part of the rectum.

The veins of the *labia* correspond in distribution to the arteries, and those from the outermost parts drain into the pudic which opens into the common iliac vein. Large veins from the labia minora open into the pars intermedia of the bulb.

The veins from the *glans* and *corpus clitoridis* pass into the dorsal vein of the clitoris, which communicates with the vesical plexus.

The veins of the *bulb* pass into the vaginal plexus.

The *Vaginal plexuses*—one outside the muscular coat and one in the submucous tissue—are most abundant at the lower part of the vagina, communicate with the hæmorrhoidal and vesical plexuses, and open into the internal iliac vein.

The *Uterine plexus* is very abundant, as is well shown in one of Hyrtl's plates; it ultimately opens into the ovarian veins (fig. 52), which pass on the right side to the inferior vena cava, on the left to the left renal vein. The right ovarian vein has a valve where it pierces the coat of the inferior vena cava (*Brinton*, quoted by *Lawson Tait*). The veins are small, lie in the outer muscular coat, and run longitudinally; in the middle layer of that coat they open into large sinuses (surrounded by circular unstriped muscle) with which the capillary vessels communicate. This is an arrangement like that in the corpus spongiosum of the penis (*Klein*).

The *Ovarian plexus*, otherwise known as the pampiniform plexus, lies between the folds of the broad ligament and communicates with the uterine plexus (fig. 52). Some apply this term to all the veins in the

broad ligament. The ovarian plexus opens into the inferior vena cava. Just at the hilum of the ovary lies the collection of veins known as the bulb of the ovary.

Beneath the peritoneum and between the layers of the broad ligaments are vast venous plexuses. Knowledge on this point is of the highest importance in relation to pelvic hæmatocele.




FIG. 52.
UTERUS and VAGINA with their venous supply, seen from behind—(Luschka). *f* Fallopian tube; *pp* pampiniform; *o* ovarian; *v* vaginal plexuses.

The vesical, hæmorrhoidal, and vaginal plexuses, with the pudic veins open into the internal iliac vein which joins the inferior vena cava.

From the hæmorrhoidal plexus, the superior hæmorrhoidal vein passes into the portal system; and thus we get a communication between the pelvic and portal venous systems.

In the vaginal mucous membrane, clitoris and uterus, we have erectile tissue, *i.e.*, veins in connective tissue with unstriated muscular fibre.

LYMPHATICS.

Lymphatic
Glands.

Under this we take up—

a. The Lymphatic glands ;

b. The Lymphatic Vessels.

a. The Lymphatic Glands.—These are (1) the *inguinal glands*, which lie parallel to and just below Poupart's ligament ; and (2) the *pelvic glands*. These latter consist of the following :—

(*a*) A gland at the isthmus uteri (*Championnière*) ;

(*b*) Hypogastric or iliac glands, which lie subperitoneally in the space between the external and internal iliac vessels ;

(*c*) Sacral, on the lateral aspect of the anterior surface of the sacrum and in the mesorectum ; and

(*d*) A gland or collection of small glands at the obturator foramen—the obturator gland of Guérin.

These all pour into the lumbar glands, which lie in front of the lumbar vertebræ and discharge into the thoracic duct.

Lymphatic
Vessels of
External
Genitals.

b. The Lymphatic Vessels. (1.) *Of External Genitals.*—Numerous vessels form a network on the internal aspect of the labia majora, over the labia minora, and round the vaginal and urethral orifices, vestibule, and clitoris ; all of these open into the inguinal glands. From this arrangement, the enlargement of the inguinal glands in syphilis and vulvar cancer is intelligible. The lymphatics of the *lower fourth of the vagina* also open into these glands.

Of Vagina.

(2.) *Of Vagina (upper three-fourths) and Cervix Uteri.*—These lymphatics open into the hypogastric glands.

So far we have followed Sappey's description. Le Bec, however, asserts that the lymphatics of the vagina pour into a series of trunks at the level of the isthmus uteri, and that those of the cervix join them ; and that the conjoined lymphatics then pass below the base of the broad ligament to the obturator gland, from which vessels communicate with others from the thigh and even from the epigastrium.

The relation between lymphatics and glands is as follows :—

(*a*) Those of the external genitals pass into the inguinal glands ;

(*b*) The lymphatics of the bladder, vagina, and cervix pass to the hypogastric glands (*Sappey*). According to Le Bec, they pass to the obturator gland.

Of Uterus.

(3.) *Of Uterus.*—The lymphatics of the body of the uterus pass through the broad ligaments ; and, along with those from the ovary and Fallopian tube, enter the lumbar glands. If Le Bec be right, the lymphatics from the cervix pass *below* the broad ligament and those from

the uterus along the *upper* part of the same. Some of the uterine lymphatics pass along the round ligament to the groin.

Leopold, who has investigated the lymphatics in the unimpregnated uterus, considers "the mucous membrane of the uterus as a lymphatic surface which contains no special lymphatic vessels, but consists of lymph sinuses covered with endothelium.

"The lymph passes from the lymphatic spaces of the mucous membrane, through the mucous membrane hollows, into the lymph spaces and vessels of the muscular coat, surrounds here all the bundles up to the serous covering, and flows into the larger vessels which enter the broad ligament in the neighbourhood of the blood-vessels" (*loc. cit.*, S. 31).




FIG. 53.

LYMPHATICS and LYMPHATIC GLANDS of PELVIS and lower part of ABDOMEN (*Poirier.*)

These are matters not of mere anatomical detail, but of the very highest pathological and practical importance. The richness of blood and lymphatic supply to the vagina, cervix, and uterus explains the extraordinary rapidity with which septic matter spreads through the body, and the extreme danger which may attend even an insignificant lesion of the internal genital organs, when septic matter is present and is absorbed. We may remark here that septic matter will of course follow the lymphatic routes already laid down, and that bacteria can penetrate the walls of blood-vessels and pass into the general circulation. It should not be forgotten, however, that the bacteria passing along the lymphatic vessels

may penetrate them, pass into the peritoneal cavity, and thence spread through the diaphragm to set up the pleurisy and pericarditis so common in septicæmia (*Lusk*). Thorough comprehension of lymphatic distribution and knowledge of the evil effects of septic matter are of the first importance to the student.




FIG. 54.
LYMPHATICS OF UTERUS (*Poirier*).

Relation
between
Glands
and Lym-
phatics.

The lymphatics of the *Rectum* lie in two layers (mucous and muscular), and open into the glands of the mesorectum or into the sacral glands.

The stomata of the peritoneum of the pelvis communicate with lymph capillaries lying in the subendothelial tissue.

The *Inguinal Glands* (parallel to Poupart's ligament) receive the lymphatics of the vulva, lower $\frac{1}{4}$ th of vagina, and urethra.

The *Hypogastric or Internal Iliac* receive those of the bladder, upper $\frac{3}{4}$ ths of vagina, and neck of uterus.

The *Sacral Glands* receive those from the rectum.

The *Lumbar Glands* receive the lymphatics from the pelvic glands, body of the uterus, Fallopian tubes, and ovaries (*v.* figs. 53 and 54).

NERVES.

These are (a) Spinal; (b) Sympathetic.

(a) *Spinal*. The pelvic muscles are supplied as follows: *Levator and Sphincter ani* by inferior hæmorrhoidal branch of pudic, 4th and 5th sacral, and coccygeal nerves; *Coccygeus*, by 4th and 5th sacral and coccygeal nerves; *Muscles of Perineum and Clitoris*, by the branches of pudic nerve. Pelvic Nerves.

(b) *Sympathetic*. The hypogastric plexus lies between the common iliac arteries; it gives off branches which, reinforced by branches from the lumbar and sacral ganglia and sacral nerves, form the *inferior hypogastric plexuses*—one on each side of the vagina. From these, filaments proceed to the vagina, uterus, Fallopian tube, and ovary.

Frankenhäuser describes a ganglion at the cervix uteri and also a vesical one. Jastrebow found the cervical ganglion to be a plexus with a ganglion enclosed in it.

The pelvic, splanchnic, and the hypogastric nerves are motor and sensory to the bladder. The pudic is motor to the sphincter ani and all the striped muscles of the vagina and clitoris: it is sensory to the skin of the perineum (*Griffiths*).

The terminations of the nerves in the muscular layers of the uterus have been studied by Frankenhäuser, who figures them passing to the nuclei of the unstriped muscle. Those entering the mucous membrane are said to end in ganglia. Numerous end bulbs have been found in the clitoris and vagina.

Gawronsky has found that in the vagina the nerves end in the epithelium, and in the uterus in the glands and epithelium. In the tube there are two concentric plexuses ending in the epithelium and in the nerve cells of the submucosa; while in the ovary they pass to the Graafian follicles and to the cells of the membrana granulosa.

DEVELOPMENT OF THE PELVIC ORGANS.

This is best considered for our practical purposes in the fœtus at the sixth and eighth weeks, and between the third and fourth months.

At the sixth and eighth weeks we have the following structures developed, viz., the ducts of Müller, the Wolffian bodies and duct, and the Ovary.

Fig. 55, from Keibel, shows a drawing of a model of the organs late

in the second month, and shows very well the relation of the organs in sagittal mesial section. The length of the urino-genital sinus should be noted. At its upper end is to be seen the eminence of Müller, the projection where the ducts of Müller end, and below the level of which the Wolffian ducts open into the sinus.

To see further details we must study transverse sections of a six weeks' foetus cut serially by the paraffin method.

Below the level of the liver and above that of the bladder, we see in fig. 56 the Wolffian bodies, one on each side of the vertebral column, with the Wolffian duct in the centre of each.

The origin of the Wolffian duct is disputed, some alleging that it arises from the Mesoblast. We believe in its origin from the Epiblast, as has been shown by several investigators.




FIG. 55.

MODEL OF ORGANS AT END OF SECOND MONTH (*Keibel*). The x indicates the top of the urino-genital sinus, and Müller's eminence where the hymen develops.

Lower down another ridge develops to the outer side of the Wolffian body; it is in connection with this ridge that the duct of Müller and the ovary arise. This ridge has two portions, an outer and inner (fig. 57). The outer has the ducts (Wolffian and Müllerian) in it, and curves in front of the inner to form the broad ligaments. The inner soon assumes the characteristic shape of the ovary. At first the union between the ridges is quite thin (fig. 57).

The section below this now shows us the broad ligament fully formed with the two ducts of Müller near its centre, the Wolffian duct on each side, and the ovary attached to its posterior surface, while at the hilum of the ovary we see the remains of the Wolffian tubules (fig. 58).

The outer surface of the ovary is covered with columnar epithelium, the germ epithelium of Waldeyer. This passes into the substance of the ovary, and gives rise to the ova and membrana granulosa.

At the level of the pelvic cavity, in the pelvic floor in fact, we come on the genital cord of Thiersch, containing three canals, the two outer being the Wolffian ducts, the central one the coalesced Müllerian ducts (fig. 59). In an earlier foetus (fourth to fifth week) the ducts of Müller have not blended, and we thus get four canals on section.




FIG. 56.

T.S. of six weeks' foetus showing Wolffian body on each side.




FIG. 57.

Shows outer ridge where Ovary and Duct of Müller develop.

In the figs. 59 and 59a we see the urino-genital fissure as a crescentic section with the eminence of Müller on the back wall of fig. 59. Here the ducts of Müller end blindly. Section 59a shows the Wolffian ducts opening into the urino-genital sinus.

This account gives only the development of the genital tract as far

down as the hymen, external genitals, and anus, and does not include an account of the origin of the last three structures.




FIG. 58.

T.S. showing Ovary and Broad Ligaments.




FIG. 59.

T.S. showing Genital Cord.

The Fallopian tubes are the non-coalesced portions of the ducts of Müller; the uterus proper is formed by the union of the ducts of Müller

to form one organ, while the remainder of the Müllerian ducts makes up the vagina down to, but not including, the hymen.




FIG. 59a.
T.S. showing Urino-Genital Sinus.

We have now therefore to consider the development of the hymen, but defer that of the external genitals and of the anus till afterwards (*v. Sect. on Vulva*).

DEVELOPMENT OF THE HYMEN.

To study the development of the hymen one must examine serial sagittal mesial and lateral sections of a female foetal pelvis between the third and fourth months. It will then be found that two epithelial bulbs form at the site of the hymen, that they are at first solid and have a structure the same as the epithelium of the fully developed vagina. The cells at the periphery of the bulb are the same as those in the deep layers of the vaginal epithelium. According to some these bulbs arise from the ducts of Müller, but the Wolffian ducts have been traced into them, and they therefore appear to arise from the Wolffian ducts and are thus Epiblastic. Whatever be their origin they perform a very remarkable part in development. Their epithelium passes into that of the Müllerian vagina and, filling it up, renders it at first solid. The central cells of the bulbs and of the now solid vagina are, however, superficial and more liable to break down, and thus a lumen forms again. The urino-genital sinus on the outer surface opposite these bulbs now sends in processes, and in this way the hymeneal opening is formed (*fig. 60*).

The epithelial cells of these bulbs not only pass into the vagina but map out the fornices and pass at first into the lower part of the cervical canal.

The genital tract down to the hymen is, therefore, Müllerian, the hymen itself mainly Wolffian.

This view explains the skin-like structure of the vagina and vaginal




FIG. 60.

Shows the Wolffian bulbs formed and breaking down in the centre. The involution from the urino-genital sinus is also seen. Above, the involution of the Epidermis to form the prepuce of the Clitoris can also be noted.

portion of the cervix, and the different structure of the genital tract above the os externum.

It has been for some time discussed whether the hymen is to be considered vulvar or vaginal. The real fact of the matter seems to be that it is vaginal and in part Wolffian.

Further details in development will be considered under the special organs.

CHAPTER VI.

PHYSICS OF THE ABDOMEN AND PELVIS, WITH SPECIAL REFERENCE TO THE SEMIPRONE AND GENUPECTORAL POSTURES.

LITERATURE.

W. Braune—Die Oberschenkelvene des Menschen in anatomischer und klinischer Beziehung : Leipzig, Veit and Co., 1871. *J. M. Duncan*—On the Retentive Power of the Abdomen : Researches in Obstetrics, p. 409. *Hart*—The Structural Anatomy of the Female Pelvic Floor. *Schatz*—Beiträge zur physiologischen Geburtskunde : Archiv für Gynäk., Bd. IV., S. 191. Einfluss der Lehre vom intraabdominalen Drucke auf die Gynäkologie : Archiv für Gynäk., Bd. V., S. 227. *Simpson and Hart*—The Relations of the Abdominal and Pelvic Organs in the Female. *Van de Warker*—A Study of the Normal Movements of the Unimpregnated Uterus : N. Y. Med. Jour., Vol. XXI., p. 337.

In this chapter it is proposed to give a brief sketch of a subject of the highest importance but still in its infancy. The *resumé* must be restricted, from want of space, to certain practical points of which we consider here the following :—

1. *The effect of intra-abdominal pressure on the female pelvic floor ;*
2. *The results brought about by change of posture, especially by the genupectoral posture ;*
3. *The effect on uterine position of digital pressure in the vaginal fornices.*

THE EFFECT OF INTRA-ABDOMINAL PRESSURE ON THE FEMALE PELVIC FLOOR.

We suppose the body to be in the upright posture. For simplicity, the pelvic floor is considered as being under fluid pressure. Fig. 61 shows the effect of this on the pelvic-floor segments. Fluid pressure acts at right angles to the limiting surface, which in this case is the pelvic peritoneum. Thus, if the perpendiculars be counted, starting from the symphysis, it can readily be seen that the first three will press the pubic segment against the symphysis ; that the fourth and fifth will do this also, but will further have a resultant tending to drive the pubic

Effect of
intra-ab-
dominal
pressure.

past the sacral segment; that the sixth and seventh will, directly, tend to do this last; and that the others will drive it partly past the sacral segment, and partly against it. From want of rigidity in the pubic segment, this driving-down tendency is partly lost. Thus the effect of ordinary intra-abdominal pressure is to press the pubic against the sacral segment. Increased intra-abdominal pressure displaces downwards a definite portion of the pelvic floor, viz., all lying in front of the anterior rectal wall.




FIG. 61.

DIAGRAM to illustrate effect of intra-abdominal pressure on the segments of the pelvic floor (Hart).

a Uterus; b Bladder; c Retropubic fat; d Labium majus; e Symphysis;
f Perineal body; g Rectum.

There is in the pelvic floor a definite line of cleavage at which it yields, which line runs between the anterior rectal and posterior vaginal walls (*see* p. 67). This definite downward displacement causes the lesion known as prolapsus uteri.

From this we see that the female pelvic floor is not equally strong throughout. It would be, were the sacral segment prolonged and attached to the symphysis pubis. But then parturition would be an impossibility. It has been constructed not only *quâ* intra-abdominal pressure, but also *quâ* parturition and the vesical and rectal functions.

THE RESULTS BROUGHT ABOUT BY CHANGE OF POSTURE, ESPECIALLY
BY THE GENUPECTORAL POSTURE.

The abdominal walls, along with the viscera bounded by them, are often spoken of as the abdominal cavity with its contained viscera. We must, however, keep in mind that this cavity is always perfectly full. There is never any vacuum in it. The viscera are always in apposition, with only a little fluid as a film separating them. The abdominal walls are yielding, and any tendency to a vacuum is counteracted by atmospheric pressure on the walls. In no posture, is there ever a vacuum in the abdominal cavity. Even if the trunk were inverted, the small intestines would still touch the uterus as they do in fig. 39. The abdominal walls and viscera enclosed by them behave, therefore, like a plastic viscous fluid—like so much thick gum or treacle.

In the *upright posture*, the viscera bulge above the symphysis pubis, more or less, according to the development of the subject. The bulging




FIG. 62.

OUTLINE OF FEMALE FIGURE IN GENUPECTORAL POSTURE. The dotted line indicates the contour when the vaginal orifice is unopened; the continuous line, the change in contour after air is admitted into the vagina (*Simpson and Hart*).

is slight in a well-formed female, but excessive if the woman is fat. Just below the sternum, the antero-posterior diameter of the abdomen is lessened. The pelvic floor is convex as seen from without, *i.e.*, the pelvic-floor projection is well marked. Atmospheric pressure is acting equally all over the abdominal and pelvic surfaces; but the pelvic-floor, bearing the weight of the viscera probably bulges more than the other boundaries of the abdomen. A fluid contained in a bag suspended from a fixed point is pyriform, with the bulb nearer the earth. This shape is due to the weight of the fluid.

If a woman be made to assume the posture known as the *genupectoral*

(better *genufacial*), the bulge is at the sternum. The following points should be noted in regard to this posture (fig. 62):—

1. The antero-posterior diameter of the abdominal cavity is increased at the sternum ;
2. It is diminished above the pubes and in the iliac fossæ ;
3. The pelvic-floor projection is diminished ;
4. The pubic and sacral segments are still in contact, and the abdominal viscera always in contact with the uterus and one another.

Let us now contrast these postures.

	<i>Upright posture</i>	<i>Genupectoral posture</i> (fig. 62).
Upright and Genupectoral Postures contrasted.	1. Greatest antero-posterior (<i>a-p</i>) diameter of abdomen in hypogastrium.	1. Greatest antero-posterior diameter at sternum.
	2. Least <i>a-p</i> diameter at sternum.	2. Least <i>a-p</i> diameter in hypogastrium.
	3. Pelvic-floor projection at its maximum.	3. Pelvic-floor projection diminished.
	4. Pelvic-floor segments in contact.	4. Pelvic-floor segments in contact.

In the latter posture, on inspection of the genitals, the labia can be seen to be furrowed and the skin over the ischiorectal fossa slightly hollowed. If now the labia majora and minora be separated and the fourchette lifted up, no further change as yet takes place ; but when the hymen is opened up, air passes into the vagina (often with a distinct hiss), and the vaginal walls become separated, enclosing a somewhat large cavity. The bulge at the sternum is now slightly increased, while the diameter in the hypogastrium is diminished (*see fig. 62*). *It is only when the anatomical entrance of the vagina (the hymeneal orifice) is opened up that the vagina distends with air.*

It has been shown by A. R. Simpson and D. Berry Hart, that the segments of the pelvic floor separate from each other when a woman assumes the genupectoral posture and the hymenal orifice is opened. The pubic segment passes down with the viscera ; the sacral segment remains behind, recoiling slightly upwards. *Thus, functionally, the pubic segment is visceral, the sacral one is vertebral.*

They have shown further that there is a definite displacement of the pubic segment constituents, viz. :—

- a.* The empty bladder is partly above the pubes ;
- b.* The peritoneum passes from abdominal wall to bladder, at a point $1\frac{1}{2}$ inches above the symphysis ;
- c.* The retropubic fat is partly above and partly below the top of the symphysis. We may now once more contrast these postures.

<i>Upright posture.</i>	<i>Genupectoral posture</i> (vagina distended with air) (fig. 63).	Result of distention of Vagina with Air.
1. Pubic and sacral segments in apposition and vagina a slit.	1. Pubic and sacral segments separated and vaginal walls bounding a cavity.	
2. Retropubic fat behind pubes.	2. Retropubic fat partly above pubes.	
3. Empty bladder behind pubes.	3. Empty bladder partly above pubes.	
4. Peritoneum passes from anterior abdominal wall to fundus of empty bladder, immediately above symphysis.	4. Peritoneum passes from anterior abdominal wall to fundus of empty bladder, $1\frac{1}{2}$ inches above symphysis.	
5. Urethra and bladder meet at a right angle.	5. Urethra and bladder almost in same line.	

The reason why the pubic segment passes downwards when the vaginal orifice is opened is, that atmospheric pressure now acts on the vaginal aspect of the pubic segment (with its movable attachment to the pubes) and drives it further down. As the result of this posture, changes take place in the length and direction of the vaginal walls and in the position of the uterus.

1. *Vagina*.—(a.) Both walls elongate.

(b.) The anterior follows the direction of the posterior aspect of the symphysis; the posterior, the curve of the sacrum.

2. *Uterus*.—(a.) The normally placed uterus passes nearer the sacrum and nearer the thoracic diaphragm.

(b.) The retroverted uterus, fixed or unfixed, becomes more retroverted.

(c.) The retroverted unfixed uterus does not become replaced so as to lie anteverted.

The results given have been obtained as follows:—

a. By observation on living patients, aided by silhouettes of the outlines of the nude body in the upright and genupectoral postures;

b. By study of frozen sections of the female pelvis, and especially by study of a frozen section of a cadaver placed in the genupectoral posture.

For further details on this subject Simpson and Hart's atlas may be consulted.

An important practical result follows from these observations. *The vagina dilates, or, more properly, the segments of the pelvic floor separate exposing their free margins—the vaginal walls—when a patient assumes the genupectoral posture and the hymeneal orifice is opened so as to admit*

air. If a patient be so placed opposite a good light, and the sacral segment be drawn up, a complete view of the vaginal walls and cervix is obtained. The same results can be got by placing the patient in the posture known as the *semiprone*. On this last fact is based the use of the vaginal speculum known as Sims' or duckbill speculum (*v.* Chap. X).




FIG. 63.

PELVIS FROM FROZEN SECTION OF CADAVER IN GENUPECTORAL POSTURE. *A* anus; *P* perineum; *R* rectum; *V* vagina; *u* urethra; *B* bladder; *f* retropubic fat; *U* retroverted uterus; *pp* peritoneum. Between the small intestine and peritoneum is fatty omentum. (*Simpson and Hart.*)

In the genupectoral posture the rectum can be distended if the anus be dilated so as to admit air. The same holds good in the case of the bladder as Pawlik and Kelly have shown.

This may be put as follows. Owing to the loose attachment of the bladder to the pubes, the vagina, rectum or bladder can be distended with air in the genupectoral posture, the special dilatation depending on whether the anus, vaginal entrance, or urethra is opened up.

THE EFFECT ON UTERINE POSITION OF DIGITAL PRESSURE IN THE
VAGINAL FORNICES.

This is a subject of great practical importance.

If, when a patient is lying on her left side, the index finger of the examiner's right hand is passed into the vagina as far as the posterior fornix, and pressure made there, the following results may be noted:—

Effect of
Digital
Pressure
in the
Fornices.

(1.) The posterior vaginal wall is elongated, the cervix drawn back, and the uterus, if anteverted, becomes more so.

(2.) If the uterus is retroflexed, the flexion is not remedied. Should the fundus be fixed, the retroflexion is increased as the cervix is drawn back while the fundus remains.

Similarly, if pressure be made in the anterior fornix:—

(1.) The uterus becomes elevated and slightly rotated backwards, because the cervix is pulled forwards.

(2.) If the uterus is anteflexed, the flexion is not diminished.

By pressure in these fornices, therefore, we only act on the cervix, unless the uterus is very much retroverted or anteverted. The body of the uterus is acted on only indirectly, through its union with the cervix.

Consequently, no vaginal pessary can undo the flexion of a retroflexed or anteflexed uterus.

CHAPTER VII.

MENSTRUATION AND OVULATION.

LITERATURE.

- Aveling*—On Nidation in the human female: *Obstet. Jour. of Gt. Brit. and Irel.*, 1879.
- Beigel*—Die Krankheiten des weiblichen Geschlechtes: F. Enke, Stuttgart, 1875.
- Bonnet*—Veränderungen des uterus während Brunst u. Trächtigkeit; in *Ellenbergers Vergleich. Physiologie d. Haussäugethiere*. *Dalton*—Report on the Corpus Luteum *Am. Gyn. Tr.*, Vol. II., p. 11. *Physiology*, 6th edition, J. & A. Churchill, 1876.
- Engelmann*—The Mucous Membrane of the Uterus, with especial reference to the Development and Structure of the Decidua: *Am. J. of Obst.*, Vol. VIII., p. 30.
- Frey's Histology*—Barker's Tr., 1874. *Goodman*—The Cyclical Theory of Menstruation: *Am. Jour. of Obstet.*, Oct. 1878. *Heape*—The Menstruation of *Sennopithecus Entellus*: *Phil. Tr. Roy. Soc. Lond.*, Vol. 185 (1894), B. pp. 44-471. *Heil*—Der Fimbrienstrom und die Überwanderung des Eies: *Arch. für Gynäk.*, 1893. *Hofmeier*—Zur Kenntniss der normalen Uterusschleimhaut: *Cent. für Gyn.*, 1893. *Jacobi*—The Question of rest for women during Menstruation: *Smith, Elder & Co.*, London, 1878. *Kinthead*—On the passage of the Ovum from the Ovary into the Fallopian Tubes: *Med. Press*, September 14, 1881. *Kundrat*—Untersuchungen über die Uterusschleimhaut: *Stricker's Jahrbuch*, 1873. (Kundrat and Engelmann were co-workers.) *Leopold*—Studien über die Uterusschleimhaut während Menstruation, Schwangerschaft und Wochenbett: *Archiv für Gynäk.*, Bd. XI., S. 1091. Untersuchungen über Menstruation und Ovulation: *Archiv f. Gyn.*, Bd. XXI., S. 347. *Leopold und Mironoff*—Beiträge zur Lehre von der Menstruation und Ovulation: *Arch. für Gynäk.*, Bd. XLV. *Loewenhardt*—Die Berechnung und die Dauer der Schwangerschaft: *Archiv für Gynäk.*, Bd. III., S. 456. *Martin, C.*—The Nerve Theory of Menstruation: *Brit. Gyn. Jour.*, Vol. IX. *Möricke*—Die Uterusschleimhaut in den verschiedenen Altersperioden und zur Zeit der Menstruation: *Ztschr. für Geburtshülfe und Gynäk.*, VII. Band, 1. Heft, 1881. *Pinner*—Über den Uebertritt des Eies aus dem ovarium in die Tube beim Säugetier: *Arch. für Physiol.*, 1880. *Ploss*—Das Weib in der Natur und Volkerkunde, 1887. *Reinl*—Die Wellenbewegung der Lebensprocesse des Weibes: *Volkmann's Sammlung*, etc., No. 243. *Ritchie, C. G.*—Contributions to Ovarian Physiology and Pathology: *Churchill*, London, 1865. *Simpson, A. Russell*—Emmenologia; Contributions to Obstetrics and Gynecology: *Edinburgh*, A. & C. Black. *Stephenson*—On the Menstrual Wave: *Amer. Jour. of Obstet.*, April 1882. *Sutton, J. B.*—Menstruation in Monkeys: *Brit. Gyn. Jour.*, Vol. II., 1880. *Tait, Lawson*—*Br. Med. Journ.*, June 4, 1881. Notes on the Relations of Ovulation and Menstruation: *Medical Times*, London, 1884. *Underhill*—Note on the Uterine Mucous Membrane of a Woman who died immediately after Menstruation: *Ed. Med. J.*, 1875. *Sir J. Williams*—On the Structure of the Mucous Membrane of the Uterus, and its periodical changes: *London Obst. Jour.*, Vol. II., p. 681. *Wyder*—Das Verhalten der Mucosa Uteri während der Menstruation: *Zeit. f. Geburtshülfe und Gynäk.*, Bd. IX., Hft. 1. For further literature see Heape's paper.

THE subject of Menstruation is not as yet well known, and on many points eminent and trustworthy observers are at variance. The nature of the process is at present *sub lite*. The old theories of its being due to plethora or its being a disease are now exploded, but Heape believes that there is evidence in the monkey (*Semnopithecus Entellus*) that an irritant is present at a certain stage in the congested blood-vessels below the surface epithelium. The modern view, termed the ovulation theory, asserts that the starting-point in menstruation is the bursting of a Graafian follicle. But in cases of abdominal section performed between the menstrual periods, as has been specially observed by Tait and Leopold, Graafian follicles have been found on the point of bursting, showing that ovulation may in certain cases occur remote from menstruation. An objection that may be urged to this is that abdominal-section cases are not normal. It is of interest to note that Heape, in examining the uteri of menstruating monkeys, found in forty-five specimens only two with a recent corpus luteum.

Recent observations by Leopold and Mironoff have tended, however, to show that ovulation in the human female may be considered as often preceding and initiating menstruation.

Jacobi, Stephenson, and Reinl (working on Goodman's cyclical theory) have given good proof that a woman in her full sexual vigour seems to pass through a series of cyclical changes, of each of which the menstrual period is the climax. Jacobi found that, during the few days before the flow, the excretion of urea is increased; the temperature is slightly raised; and that, in regard to the pulse, there is a rhythmic wave beginning at a minimum point 1 to 4 days after the cessation of the flow and gradually rising to a maximum 7 or 8 days before menstruation. So far as our present knowledge goes, the following is a brief *resumé*.

PRELIMINARY CONSIDERATIONS.

Definition.—A cyclical change with constitutional disturbances whose Prelimi-
most marked local phenomena are periodical flow of blood from the uterine naries.
cavity, with shedding of the superficial layers of its mucous membrane, accompanying (according to the hitherto accepted theory) the discharge of an ovum from the ovary, occurring in properly developed women between the ages of 14 and 44, and interrupted by uterogestation and lactation.

Period of its Onset.—Menstruation begins, in this country, usually at the age of 13 to 15 (puberty). It may be delayed till 16, 17, or 20; but this is unusual. Its onset is said to be earlier in warm countries, later in cold ones; earlier in delicately nurtured girls. Observations of medical women in India, mainly those of Dr Pechey-Phipson, have

shown, however, that early onset is not the rule there, puberty occurring at the same period as in temperate climates.

Period of its Cessation.—With the interruptions of pregnancy and lactation, it continues in healthy women until the age of 44 to 50. The period of its final cessation is known as the menopause. As a general rule the menopause is early when menstruation has begun early, and *vice versa*.

GENERAL PHENOMENA OF MENSTRUATION.

General
Pheno-
mena.

Changes at Puberty.—At this period of life, when the girl becomes the woman, we find certain well marked general changes occurring. The bust and mons veneris develop and the whole contour of the body becomes more rounded and attractive; hair appears on the genitals. The romping carriage of the girl becomes subdued, and greater shyness characterises her conduct to the opposite sex.

Phenomena premonitory to each menstrual flow.—There is usually a feeling of weight in the pelvis and increase of sexual inclination. Many women, however, have very little uneasiness during the whole flow; while others are always considerably distressed,—this distress being still outside the boundary of actual disease.

Periodicity and duration of Discharge.—When once established it recurs, in the large majority of cases (about 87 p. c. of the whole), with great regularity: the most common intervals are 28 days (in 71-p. c.) and 30 days (in 14-p. c.); less frequent are 21 days (in 2-p. c.) and 27 days (in 1 + p. c.). We speak therefore of the 21 day *type* and so on. The discharge lasts for a number of days, varying from 2 to 8; if below 2 or above 8 it is abnormal; but of course other points besides mere duration must be taken into account. The duration of the flow and the “type” give the woman’s “habit” in menstruation.

LOCAL PHENOMENA.

Local
Pheno-
mena.

Three periods are usually distinguished from a clinical point of view: 1. Invasion; 2. Persistence; 3. Decline.

1. *Invasion.*—Discharge pale.

2. *Persistence.*—Discharge bright red, non-coagulable from its mixture with mucus. It consists microscopically of epithelium from vaginal, cervical, and uterine cavities; mucous globules; compound granular corpuscles; and red and white blood-corpuscles.

3. *Decline.*—Discharge lessens in amount and becomes lighter in colour.

The total *quantity* varies from 2 to 8 ounces.

Thus far we have related facts fairly well ascertained and not much disputed. We now enter on more debateable ground, in considering—

- I. Ovulation ;
- II. The Corpus luteum ;
- III. Source of discharge, and changes in the uterine mucous membrane.

I. *Ovulation*.—According to the ovulation theory, ovulation forms the starting-point of the process of menstruation. We have already considered the structure and development of the ovary, and now describe

The changes in the Ovary at each Menstrual Period.—A Graafian follicle enlarges and moves nearer the surface. Probably this produces, through a nervous mechanism, a hyperæmia of the whole pelvic contents,—peritoneum, connective tissue, uterus, ovaries, Fallopian tubes, and vagina. It is alleged, as yet on insufficient grounds, that the fimbriated end of the Fallopian tube grasps the ovary, and that the ovum from the ruptured Graafian follicle passes into it and along the tube to the uterine cavity. In the Fallopian tube and uterus the action of the cilia is towards the external os (Hofmeier), and we thus get a current in the peritoneal serum which carries the ovum into the uterine cavity. However it reaches the Fallopian tube and uterus, its further development depends on its fertilisation or non-fertilisation. In the latter case it passes off unnoticed in the menstrual discharge; in the former it develops into the fœtus.

II. *The Corpus luteum*.—After the rupture of the Graafian follicle, we get its cavity filled up by the structure known as the corpus luteum. It varies according as pregnancy does or does not follow its formation. The difference is well given in Dalton's table, which we subjoin.

	CORPUS LUTEUM OF MENSTRUATION.	CORPUS LUTEUM OF PREGNANCY.
End of 3 weeks.	12 by 13 mm. in diameter ; central clot reddish, convoluted wall pale.	
One month.	Smaller ; convoluted wall bright yellow ; clot still reddish.	Larger ; convoluted wall bright yellow ; clot still reddish.
Two months.	Insignificant cicatrix.	12 by 22 millimetres in diameter ; convoluted wall bright yellow ; clot perfectly decolorized.
Four months.	Absent or unnoticeable.	18 by 22 millimetres in diameter ; clot pale and fibrinous ; convoluted wall dull yellow.
Six months.	Absent.	Still as large as at the end of the second month ; clot fibrinous ; convoluted wall paler.
Nine months.	Absent.	10 by 13 millimetres in diameter ; central clot converted into a radiating cicatrix ; external wall tolerably thick and convoluted, but without any bright yellow colour.

The corpus luteum is formed by proliferation of the cells of the membrana granulosa, by the sprouting of new capillaries with migratory cells into the hypertrophied convoluted epithelium. The central portion degenerates into gelatinous tissue, the cortical into fatty tissue (*Klein and Smith*). It thus consists of a vascular framework, with a yellow pigmentary and cellular substance.

Source of Discharge.

III. *Source of Discharge and Changes in the Uterine Mucous Membrane.*—All observers are agreed that the mucous membrane of the uterine cavity is the source of the discharge, *i.e.*, that it comes from the area limited by the uterine ends of the Fallopian tube and the os internum.

Now begins the divergence.

Williams' View.

(1.) Sir J. Williams holds that "uterine contraction drives the blood from the muscular wall into the mucous membrane; the vessels of this membrane, having undergone fatty degeneration, give way, and extravasation of blood results. This extravasation takes place always near the surface, for in that situation the degenerative change has most advanced. The rush of blood into the vessels of the mucous membrane expels the contents of the glands, together with the greater part of their lining epithelium. . . . When hæmorrhage has taken place into the membrane, it undergoes rapid disintegration, and becomes entirely removed." The new mucous membrane "is produced by proliferation of the elements of the muscular wall of the organ: the muscular fibres producing the fusiform cells; the connective tissue, the round cells; and the groups of round cells in the meshes formed by the muscular bundles, the glandular epithelium." These "groups of round cells" may be the terminations of the uterine glands.

In a more recent paper,¹ Williams has modified the statement of his view by affirming that the greater portion of the muscular wall of the uterus represents the muscularis mucosæ. According to this, only the glandular portion of the mucous membrane is shed.

Entire removal of the mucous membrane down to the muscular fibre, and its regeneration from groups of round cells in the muscular coat, are the essentials of Williams' view.

Kundrat and Engelmann's View.

(2.) Kundrat and Engelmann thus describe the changes.

Mucous membrane becomes swollen and pulpy, and measures in thickness 3–6 mm. The thickness is most marked at the fundus and central portions of the anterior and posterior surfaces. The surface is puffy and injected; glands are distinctly seen on section as fine spirals.

Microscopically, this increase in thickness is seen to be due to a proliferation of the round cells of the stroma, an enlargement of all the cell

¹ On the Circulation of the Uterus, etc.: Lond. Obs. Trans., 1885.

elements in the superficial layers, and an increase of the intercellular substance. This superficial layer has grown far above the original gland openings, causing the funnel-shaped depressions or small pits seen on surface view. The glands are increased in thickness and length. The vessels are enlarged and gorged with blood. Fig. 66 shows the mucous membrane of the menstruating uterus magnified 40 times.

The increase of the thickness of the mucous membrane begins as the time of menstruation approaches, is most marked during the period itself, and gradually decreases after the cessation of the catamenial flow.




FIG. 64.

DIAGRAM of UTERUS just before MENSTRUATION. The shaded portion represents the MUCOUS MEMBRANE (J. Williams).




FIG. 65.

DIAGRAM of UTERUS when MENSTRUATION has just ceased, showing the cavity of the body deprived of MUCOUS MEMBRANE (J. Williams).

Fatty degeneration takes place in the cells of the interglandular tissue, blood-vessels, and glandular and surface epithelium.

They hold that "the hæmorrhage is always confined to the surface of the lining membrane, and is due to the fattily degenerated tissue being unable to resist the blood pressure;" and *they therefore maintain, what is most probably the case, that only the superficial layer of the mucous membrane is shed at a menstrual period.*

(3.) Leopold denies the existence of any fatty degeneration of the superficial layers of the mucous membrane. He believes that an extravasation of red and white blood corpuscles from the superficial capillaries takes place especially towards the superficial layer, undermining the uppermost layer of cells; and that, finally, the copious supply of blood

reaching these capillaries from the numerous arteries causes rupture and bleeding. The mucous membrane is regenerated by an upward growth of the glandular epithelium.

Möricke's
View.

Williams, Kundrat, Engelmann, and Leopold examined uteri from *post-mortem* cases. Möricke has curetted the uteri of living women at various stages of menstruation, and microscopically examined what he removed. He asserts "that during menstruation the mucous




FIG. 66.

MUCOUS MEMBRANE OF MENSTRUATING UTERUS (*Kundrat and Engelmann*). (4°)

membrane disappears neither partially nor fully." This shows how widely microscopists vary. Williams says all the mucous membrane down to the uterine muscle is removed; Kundrat, Engelmann, and Leopold say only the superficial layers are removed; and Möricke says none is removed.

We have deemed it best to lay these views before the student. The

subject is difficult to investigate, and one on which the authors are not qualified to give an opinion. They incline, however, to the views of Kundrat, Engelmann, and Leopold.

A dispute still exists as to which ovum is fertilised when pregnancy occurs—the ovum of the last menstruation, or that of the first period missed. Many observers believe in Loewenhardt's theory, viz., that the ovum fertilised is that of the first period missed.

Lately the dominant influence of the ovary in menstruation has been questioned by some, notably by Lawson Tait. The operation known as Battey's operation, where both ovaries are removed, does not always cause a cessation of bleeding. It is a broad fact, however, that if the ovaries be completely removed and no ovarian tissue left, not even in the bite of the ligature, menstruation usually ceases. The influence of the removal of the tubes cannot be satisfactorily ascertained, as in all operations they are only partially cut away.

Leopold's monograph is illustrated by many valuable lithographs, and the same may be said in regard to Dalton's work on the Corpus Luteum.

Menstruation in Monkeys.—It has long been known that monkeys menstruate. Recently more exact details have been given by Bland Sutton and Heape. Sutton examined the uteri of menstruating monkeys (*Macacus Rhesus*) in the London Zoological Gardens, and found blood in the cavity, but no loss of epithelium.

Heape, who obtained his material in Calcutta, investigated the uteri of *Semnopithecus Entellus* (fifty specimens). The mucosa of the resting uterus is a plasmodium of anastomosing nuclei with surface and glands lined with columnar epithelium. During the period of growth he found the stroma increased, and the blood-vessels behind the epithelium engorged. The hypertrophied vessels undergo amyloid, not fatty degeneration, and the number of leucocytes increase in the blood-vessels below the surface, probably from the presence in the blood of some noxious material. The blood extravasated by rupture of these blood-vessels collects in spaces or lacunæ, which then rupture through the surface. In this way the superficial layer of the mucosa is cast off (*mucosa menstrualis*), and then regeneration takes place. Some of the new epithelium seems to be formed from the stroma, the rest from pre-existing epithelium, and leucocytes take no part in the formation of the new lining of the uterus. Many other interesting points are discussed, and the whole monograph is most valuable. He divides the phenomena of menstruation as follows:—

- A.—*Period of Rest.*
 - Stage I. Resting Stage.
- B.—*Period of Growth.*
 - Stage II. Growing Stroma.
 - „ III. Increase of vessels.
- C.—*Period of Degeneration.*
 - Stage IV. Breaking down of vessels.
 - „ V. Formation of lacunæ.
 - „ VI. Rupture of lacunæ.
 - „ VII. Formation of Menstrual Clot.
- D.—*Period of Recuperation.*
 - Stage VIII. The Recuperation Stage.



SECTION II.

PHYSICAL EXAMINATION OF THE FEMALE PELVIC ORGANS.

IN this section we have to take up the physical examination of the female pelvic organs—that is, exploration by the hands and instruments of the gynecologist. This will be considered in the following manner :—

CHAPTER VIII. Abdominal Examination ; Vaginal Examination ; the Bimanual Examination, with its various modifications ; Examination per Rectum.

CHAPTER IX. The Volsella.

CHAPTER X. Vaginal Specula.

CHAPTER XI. The Uterine Sound.

CHAPTER XII. Tents and other Uterine Dilators.

CHAPTER XIII. The Curette.

CHAPTER XIV. Knives ; Scissors ; Needles ; Sutures ; Douches and Syringes ; Anæsthetics.

CHAPTER XV. Antiseptis ; Asepsis.

CHAPTER VIII.

ABDOMINAL EXAMINATION; VAGINAL EXAMINATION THE BIMANUAL EXAMINATION, WITH ITS VARIOUS MODIFICATIONS; EXAMINATION PER RECTUM.

LITERATURE.

Hegar—Die operative Gynäkologie, zweite Auflage: Stuttgart, 1881. *Kelly, H. A.*—A New Method of Examination and Treatment of Diseases of the Rectum and Sigmoid Flexure, *Annals of Surgery*, April 1895. *Mundé*—Minor Gynecology: Wood & Co., New York, 1881. *Cunningham, D. J.*—Delimitation of the Regions of the Abdomen, *Jour. of Anatomy and Physiology*, Vol. XXVII., 257.

IN a female patient whose symptoms point to a pelvic cause, it is necessary to investigate the case by what is commonly known as a vaginal examination. A mere vaginal examination, however, gives very little information. The proper method is first to make an external abdominal examination and then the vaginal examination, the latter being only a stage of the more complete method of investigation known as the bimanual. Special cautions as to cases unsuitable for pelvic exploration are given under the head of vaginal examination. We consider the examination in the following order:—

- I. External abdominal examination;
- II. Inspection of external genitals (only when necessary);
- III. Vaginal examination;
- IV. The bimanual (abdomino-vaginal) examination.

EXTERNAL ABDOMINAL EXAMINATION.

External
Abdominal
Examina-
tion.

The patient should lie on the back, with knees drawn up, and head supported on a pillow. The bowels and bladder should be empty. The abdominal surface should be exposed from the epigastrium downwards; no part of the mons veneris should be uncovered. The most delicate method of accomplishing this is as follows. A sheet or blanket is thrown over the recumbent patient; beneath this she raises up her dress as far as the pit of the stomach; the examiner then places his one hand on

the sheet, a little above the mons veneris, and turns it down over it with his other hand. The abdominal surface is examined in four ways, viz., inspection, palpation, percussion, auscultation.

A. *Inspection*.—The form, colour, equality or inequality of bulge of the abdominal surface should be noted; the presence or absence of the linea nigra, lineæ albicantes (fresh and old), pigmentary deposits, fat streaks, and skin eruptions. The linea nigra has little significance. The lineæ albicantes indicate that the patient's abdominal cavity is or has been distended beyond the normal. They are not specially




FIG. 67.

characteristic of pregnancy. Fresh lineæ albicantes are glistening and pearly; old ones have a dull white or scarred appearance.

B. *Palpation* should be performed with both hands. For this purpose the hands, well warmed, are laid flat on the abdominal surface; and the whole area is manipulated between them. One hand alone is of little use. By this method the abdominal contents may be compressed and moved between the hands. The feeling given normally is that of manipulating a plastic fluid. Tapping with one index finger so as to give a fluctuating impulse to the other hand is of great value. Circumscribed nodules or tumours, fluid collections, thickening of the skin, should be noted and mapped out on the scheme given in the chapter on case-taking.

Abdominal
regions.

For the more exact localisation of the normal and abnormal abdominal contents, anatomists divide the anterior abdominal surface into definite regions by vertical and transverse lines. The lower transverse line is drawn at the level of the most prominent lateral point of the iliac crest when viewed from the front (Cunningham)—this is the inter-tubercular plane; the upper one joins the most prominent parts of the tenth costal cartilages—(subcostal plane). A vertical line joining the cartilage of the eighth rib with the middle of Poupart's ligament on each




FIG. 68.

side, completes the division into nine areas, which are named in order as follows (*vide* Figs. 67 and 68.)

- | | | |
|-------------------------|-----------------|------------------------|
| 1. Right Hypochondriac. | 2. Epigastric. | 3. Left Hypochondriac. |
| 4. „ Lumbar. | 5. Umbilical. | 6. „ Lumbar. |
| 7. „ Iliac. | 8. Hypogastric. | 9. „ Iliac. |

In fixing the lower transverse line we have followed Cunningham, who places this line a little higher than usual.

In these regions the following structures approximately are found:—

- Epigastric Region.*—The greater part or the whole of the left lobe, and part of the right lobe of the liver, with the gall-bladder, part of the stomach, including both orifices, the first and second parts of the duodenum, the duodeno-jejunal flexure, the pancreas, upper or inner end of the spleen, parts of the kidneys, and the suprarenal bodies.
- Right Hypochondriac.*—The greater part of the right lobe of the liver, the hepatic flexure of the colon, and part of the right kidney.
- Left Hypochondriac.*—Part of the stomach, with the greater portion of the spleen and the tail of the pancreas, the splenic flexure of the colon, part of the left kidney, and sometimes a part of the left lobe of the liver.
- Umbilical.*—The greater part of the transverse colon, the third part of the duodenum, some convolutions of the jejunum and ileum, with portions of the mesentery and great omentum, and part of the right, or sometimes of both kidneys.
- Right Lumbar.*—The ascending colon, part of the right kidney, and sometimes part of the ileum.
- Left Lumbar.*—The descending colon, part of the jejunum, and sometimes a small part of the left kidney.
- Hypogastric.*—The convolutions of the ileum, the bladder in children, and when distended in adults also, the uterus when in the gravid state, and behind, the sigmoid loop and upper part of the rectum.
- Right Iliac.*—The cæcum with the vermiform appendix, and the termination of the ileum.
- Left Iliac.*—The sigmoid colon, convolutions of the jejunum and ileum (*Quain*).

The student must remember that the bladder and the uterus and appendages in the non-gravid conditions lie within and below the pelvic brim.

For the relations of the lower regions of the abdomen to the pelvic contents, the student may consult fig. 45, which shows very well the latter as seen through the brim.

In palpating the normal abdomen, the sensation given is one of impulse communicated generally through a plastic fluid. When free fluid is in the abdominal cavity, the impulse is more distinct. When the fluid is encysted, the impulse and tense feeling are localised.

When any large body is felt in the abdominal cavity, the first point to be determined is whether the body is pelvic or abdominal. This is easily

done by attempting to press the hand downwards just above the symphysis pubis. If the tumour is pelvic, and rising up into the abdomen, the hand cannot be so pressed; and conversely.

The next point is to ascertain with which of the organs the tumour is connected; and, for this, perfect familiarity with the topography of the viscera is of the highest importance. The student should ask himself what structures are normally present in the region, and then to which of these the tumour is to be referred; with regard to the iliac regions he should bear in mind the frequency of inflammatory and suppurative deposits in the peritoneum and cellular tissues. Appendicitis must also be kept in mind.

In all tumours, the existence or non-existence of intermittent contractions should be carefully noted. Their presence indicates a uterine tumour—pregnancy or soft fibroid.

The following general points should be kept in mind. The bladder is only in the hypogastric region when distended or displaced upwards; if empty, it is behind the pubes and in the true pelvis; a distended bladder may be as large as a six months' pregnancy. Ovarian tumours are more or less lateral; uterine tumours generally central, although the pregnant uterus has usually a right lateral obliquity. In advanced pregnancy, the parts of the fœtus can be distinctly palpated. Finally, it should be kept in mind that in cases of cystic tumours the catheter may require to be passed into the bladder, for an obvious reason.

CASE.—Mrs A. was sent for consultation as to removal of internal tumour. On examination, a cystic tumour was felt mesially in the abdomen and reaching up to umbilicus. Vaginal and bimanual examinations were exceedingly painful. A catheter passed into the bladder evacuated a large amount of urine. The uterus was now found to be retroverted and gravid $3\frac{1}{2}$ months, and the cystic tumour had disappeared.

Palpation
of Groin.

Palpation of the inguinal region is of great importance and should never be omitted. Glandular and other enlargements in this position may be the following:—

(1.) Glands enlarged from gonorrhœa. There are usually one or two—large, painful, and often suppurating.

(2.) Glands enlarged from syphilis. These are multiple, hard, small, painless, and never suppurate in an uncomplicated case.

(3.) Glands enlarged from vulvar malignant disease, or malignant disease of vagina (lowest $\frac{1}{4}$) or urethra; from sarcoma of pelvis.

(4.) Femoral or inguinal hernia.

(5.) Thrombosis of femoral vein.

Percussion.

C. *Percussion* is to be made in the usual way. To perform this thoroughly, the patient should be percussed (*a*) when on her back; (*b*) when on the left side; (*c*) when on the right side; (*d*) when sitting up. Changes in the percussion note on the patient changing her posture

should be carefully noted, as they are of great value (*vide* under Ovarian Tumours and Ascites).

D. *Auscultation* is performed with the ordinary stethoscope. The ^{Ausculta-}foetal heart, uterine souffle, and friction may be heard by it. The im-^{tion.}portance of auscultation is evident. Foetal heart-sounds indicate pregnancy; the point of greatest intensity of the heart-sounds indicates the lie of the child. Uterine souffle and no heart-sounds (after $4\frac{1}{2}$ months) indicate either pregnancy and child dead, or fibroid tumour. Ovarian cysts have no souffle.

Before finishing abdominal examination, the patient should be made to raise her shoulders by grasping the examiner's hands. When there is no encysted abdominal tumour, the recti can be seen to flatten the abdominal contour; if, however, a solid or cystic tumour be present, the contour is unaltered. An exception should be made in the case of thin-walled cysts not tensely filled, where the recti do flatten the contour.

INSPECTION OF EXTERNAL GENITALS.

This should not be made a routine practice. As a general rule, inspec-^{Inspection}tion of the genitals should only be made when there is local tenderness, ^{of External}where syphilis or gonorrhœa is suspected, or where it is said by the ^{Genitals.}patient that something comes down at the vaginal orifice. Soft chancres, hard chancres (almost never seen in females), mucous patches, condylomata; urethral caruncles; irritable spots causing vaginismus; labial abscess; parturition tears of perineum and labia; prolapsed pelvic organs; external or internal piles, may be found.

VAGINAL EXAMINATION.

Preliminaries.—Vaginal examination should not be made on girls ^{Vaginal}below or little beyond the age of puberty, unless the symptoms are ^{Examina-}urgent, *e.g.* mechanical retention of menstrual fluid from atresia. In such cases the consent of the parents or guardians should be obtained, and they should be present at the time of examination. In the case of unmarried women it should not be performed unless specially necessary. In both classes of patients the value of a rectal examination should be kept in mind. The vaginal examination should be made on married women whose symptoms point to a pelvic cause. Finally, no woman should be examined vaginally when menstruating normally, unless under exceptional circumstances.

Special cases require consideration: viz., that of a mistress who requests a medical man to examine her servant, who is suspected of pregnancy; or of a young woman, who, owing to a malicious report, requests examination as to her condition and a certificate that she is not pregnant.

In the first case, it is better for the medical man not to examine the patient, as he may be liable to an action for assault.

In the second case, the medical man should advise the patient against being examined. This latter case is quite different from that of an unmarried woman who, having run the risk of impregnation, requests examination to settle whether she is pregnant. In this instance the medical man investigates the case in the usual way.

After settling these preliminaries, and having obtained the patient's consent to "examine" (a term which will readily be understood by her as meaning a vaginal examination), the next point is to determine the posture the woman is to occupy while the examination is being made.

Position of
Patient.

In this country it is customary to place the patient on her left side for the vaginal examination, and in the dorsal posture for the bimanual. The patient lies on a convenient couch, with knees well drawn up and clothes loose. The examiner carefully oils or soaps the index and middle finger of his right hand. With his left hand he clears away the clothes from the hips so as to make a passage for the examining fingers, which he passes onwards till he reaches the cleft between the buttocks. He next passes them forwards over the anus, skin over base of perineum and fourchette, until the pulp of the finger rests at the vaginal orifice. In multiparous women, the lax vaginal orifice is easily felt. When in doubt, he passes his fingers cautiously on until he touches the vestibule, which is always smooth. Carrying his fingers back, he will then reach the vaginal orifice at the base of the vestibule.

The tyro must be careful not to pass his finger into the rectum by mistake. He should remember that the vaginal axis passes backwards, the anal axis forwards; that no force is required to pass the finger into the vagina where the hymen has been ruptured, whereas some force is necessary to overcome the resistance of the sphincter ani. The clitoris, lying at the apex of the vestibule, should never be touched on vaginal examination.

The two fingers, being now at the vaginal orifice, should be carried backwards into the vagina until its upper limits are felt. In doing so, the following points should be noted.

What to
note.

1. *State of Vaginal Orifice*: patulous or narrow, presence or absence of painful spots, presence or absence of spasm.

2. *Walls*: shape and length; presence or absence of rugæ; moisture, heat, secretion, tumours attached to them; fistulæ; foreign bodies, such as pessaries, glycerine plug, oakum plug.

3. *Cervix*: direction, size, shape, and consistence. Note whether thickened, expanded, and fixed; drawn to one or other side; mobile and not fixed; or whether split and with cicatrices radiating from it to vaginal roof.

4. *Os*: size, shape, consistence of lips. Thus, it may be a dimple, as in nulliparæ; transverse, as in parous women (figs. 13 and 14); or the cervix may be split on one or both sides, and thus no *os externum* be present but the cervical canal be more or less exposed. Bodies pro-

jecting through it should be noted : these may be polypi, fragments of abortion, cancerous masses, stem pessaries.

5. *Posterior fornix* is concave when felt from below. It has normally a feeling like that of the inside of the angle of the mouth. Note if any lump can be felt through it, projecting downwards in Douglas' pouch, rendering the fornix convex. A body or resistance felt through the posterior fornix may be the following :—

- | | |
|---|--|
| (1.) Fæces or tumours in the rectum ; | Bodies felt
through
posterior
fornix. |
| (2.) Acute or chronic inflammatory deposit in the peritoneum or cellular tissue ; | |
| (3.) Retroverted corpus uteri (non-gravid or gravid) ; | |
| (4.) Blood effusion ; | |
| (5.) Fibroid attached to posterior wall of uterus ; | |
| (6.) Ovary inflamed or cystic ; | |
| (7.) Ascitic fluid ; | |
| (8.) Extra-uterine foetation or hydatid (rare). | |

Anterior fornix.—Note if there is any body felt through it. If so, it is most probably the corpus uteri, normal or enlarged from pregnancy or fibroid. There may be also inflammatory or blood effusions, or a tender ovary, but these are rare here.

7. *Lateral fornices*.—Note cicatrices, prolapsed or cystic ovary, lateriflexed uterus, inflammatory or blood effusion or gestation in broad ligament, dilatation of Fallopian tubes, fibroids placed laterally.

The vaginal examination has now been completed. The student should keep in mind that he really learns very little from a vaginal examination, just as he can learn very little as to the size and relation of any object by touching it with the fingers on a but limited area. Vaginal examination is thus only the preliminary to the bimanual or abdomino-vaginal.

BIMANUAL (ABDOMINO-VAGINAL) EXAMINATION.

This method of examination is the all important one in gynecology, Bimanual. and is the one which the student and practitioner will find most valuable, so that its practice should precede all other methods of internal investigation. As the practitioner's experience increases, he will find that he relies more upon this and becomes less dependent on other means of examination.

Method of performing Bimanual. Posture of Patient.—The patient must now be placed in the dorsal posture. The head and shoulders should be supported and the knees drawn up.

Arrangement of Examiner's hands.—The *internal* hand (the right) is placed as follows : The fingers (index and middle) are in the vagina, the thumb rests in the fold between a labium majus and the thigh or

Position of
Hands in
Bimanual.

upon the symphysis, and the other fingers lie in the cleft of the nates (fig. 73), or flexed on the palm (fig. 70). The whole hand is then rotated backwards so as to bring its long axis as nearly as possible into the axis of the brim, and is then pushed up towards the brim of the pelvis. Thus the pubic segment, uterus with annexa, and posterior vaginal wall are lifted up towards the brim. The middle finger is placed over the os and the index one in the anterior fornix, so that the uterus as it is pushed up becomes more anteverted. The right hand while examining therefore has the appearance at fig. 69. The external hand (the left) is placed on the abdominal wall just above the pubes. It is now steadily depressed until the abdominal wall below it is markedly cupped (figs. 70 and 71) and moulded over the uterus and appendages, which have been elevated by the inner hand. In this way the two




FIG. 69.

RIGHT HAND IN BIMANUAL EXAMINATION.

hands estimate the size and relations of the pelvic contents, just as one would estimate the size of a watch covered with a cloth. The student should note specially that the upper hand should be steadily and not spasmodically depressed; that he should always keep the ulnar edge of the hand (rather than the palm) towards the abdominal surface, so that he may not retrovert the uterus; and that he should palpate all the abdominal areas along the pelvic brim so as not to miss anything. *His first object in the bimanual examination is to determine where the uterus is,* as this greatly simplifies the recognition of abnormal products in the pelvis. He then bimanually explores the fornices, moving the internal fingers appropriately and noting what he feels. At first his diagnosis should be simply physical, *e.g.*, "uterus felt to front and a large firm lump behind it;" or, "uterus felt retroverted and a small moveable tumour on its left side."

It is of importance that the student should know what a "normal bimanual" is. The following is a description of the condition found in a nulliparous married woman on vaginal and bimanual examination. Normal condition on Bimanual.

"Ostium vaginae patulous, and admits two fingers; vaginal walls moist, rugous, with no abnormalities. Vaginal portion of cervix normal in size (fig. 13); os uteri felt like a dimple, looking downwards and backwards. No bodies are felt through the lateral and posterior fornices which are concave on their vaginal aspects, and have the feeling, on pressure, of the angle of one's mouth. In the anterior fornix a body is felt, which on bimanual examination is discovered to be the uterus lying to the front and not enlarged. The body and cervix meet at a very




FIG. 70.

BIMANUAL EXAMINATION. The upper hand is not shown. (Hart).

obtuse angle. Bimanual exploration of the fornices reveals nothing distinctly palpable.¹ The patient complains of no pain during the whole examination."

Cases where the Bimanual is difficult.—The student will soon find that Difficult Bimanual. the bimanual can be performed in certain cases with great facility and accuracy, while in others it is exceedingly unsatisfactory.

The best case for a bimanual is in a patient a fortnight or three weeks after delivery. The reasons for this are evident: The ostium vaginae and vaginal walls have been rendered lax by the child's head; the pubic

¹ One practised in the Bimanual can feel the normal ovaries.

segment has been drawn up and its attachments slackened; the abdominal walls have had their elasticity diminished by the full-time uterus, and the uterus itself is not involuted to its normal size. In such a case, there are evidently all the requisites for a good bimanual.

Difficult bimanual cases are found in stout nulliparous women, and in cases of pelvic inflammation. In such the rectal examination (with or without the use of the volsella) is indicated.

Students at first find the bimanual unsatisfactory. By perseverance,




FIG. 71.

ANTERIOR ABDOMINAL SURFACE with upper hand placed for Bimanual (*ad naturam*).

The hand is really turned more round towards the middle line than appears in the cut, and pressure is made with the tips of all the fingers so that they are almost perpendicular to the abdominal surface.

however, they will obtain by means of it an accuracy in diagnosis which is astonishing. It is not only the best means of investigation, but one from which no possible harm can arise. In no cases is it contra-indicated except those of advanced cancer or of acute inflammation.

We have described the simple abdomino-vaginal examination. It will be readily understood that we may have others, as follows:—

- (1.) Recto-abdominal (finger in rectum and left hand above);

- (2.) Recto-vagino-abdominal (middle finger in rectum, index finger in vagina, and left hand above);
- (3.) Vesico-vagino-abdominal (middle finger in vagina, index in bladder, and left hand above);

Of these the third is never now practised.

Note that in the Bimanual the pubic segment with uterus and its annexa are elevated, the sacral segment shortened, and the abdominal wall depressed (fig. 72).

Anatomy
of Bi-
manual.

Before and after the bimanual or other examination, the examiner should scrupulously cleanse his hands, not only to avoid carrying sepsis, but also to guard against his own accidental inoculation, by no means




FIG. 72.

DISPLACEMENT OF PELVIC-FLOOR SEGMENTS AND ABDOMINAL WALL IN BIMANUAL (Hart).

infrequent, with syphilis. There are no better substances for this than turpentine and ordinary soap. The odour is by no means disagreeable, and if found objectionable can be easily covered by vinegar, which in itself is a good cleanser. The hands should finally be vigorously scrubbed with a nail brush (without soap) in corrosive sublimate, 1 in 2000 or 3000. Carbolic lotion (1-40) and soap give also thoroughly good and convenient disinfection.

In examining cancerous cases, where the odour is exceedingly penetrating and persistent, it is a good plan to dip the fingers in turpentine prior to the examination. (*v.* Chap. XV. Antisepsis.)

Rectal
Examina-
tion.

The results obtained by a vaginal examination are limited by the fact that the reflection of the vaginal walls to form the fornices, prevents the finger being pushed up to a sufficient distance. This defect is compensated for by the downward pressure of the upper hand in the Bimanual; but in cases where the abdominal walls are unyielding or the pubic segment stiff, due pelvic exploration by an abdomino-vaginal examination alone is impossible. In such cases, rectal exploration and the abdomino-rectal or abdomino-recto-vaginal examination are invaluable; they give better information than the more commonly practised abdomino-vaginal.

Methods.

The usual methods are the following:—

- (1.) Simple rectal, abdomino-rectal, abdomino-recto-vaginal;
- (2.) Passage of the *whole* hand into the rectum (Simon's method).

SIMPLE RECTAL; ABDOMINO-RECTAL; ABDOMINO-RECTO-VAGINAL.

Prelimi-
naries.

Preliminaries.—The patient should be told that it is necessary to examine the bowel. If the rectum is loaded the examination should be deferred till next day, and the patient instructed to use a purgative at night and an enema in the morning.

Manner of
Perform-
ance.

The following points should be especially noted. The examiner should thoroughly soap the fingers and nails. A vaginal examination may be made first; and then, the index finger being kept in the vagina, the middle one is passed into the rectum (fig. 73). If the patient is virginal, and it is wished to avoid a vaginal examination, then the index finger alone is passed into the rectum. When the finger or fingers are withdrawn from the rectum the hands should be at once cleansed; there can be nothing more hurtful to a patient's feelings than the passing of the uncleansed fingers from the rectum into the vagina. The patient lies first on the left side and then on the back.

Anatomy
of Rectal
Examina-
tion.

The finger passed into the rectum goes forwards; when passed into the vagina, the direction is backwards. After overcoming the resistance of the strong external sphincter it enters the rectal ampulla (fig. 34), which is often expanded by flatus. Passing the finger onwards and to the left side, a confused mass of tissue is felt in which we may detect the opening betwixt the segments of the sphincter tertius.

What to
note.

As we pass the finger inwards we note piles (internal and external), fissures, polypi, ulcers, stricture (specific and malignant).

We next turn the pulp of the examining finger so that it lies on the anterior rectal wall. Through this can be felt the cervix. Note that the whole cervix is felt, and that it is much larger than the vaginal portion felt on vaginal examination. Be sure not to mistake it for the body of the uterus. If the uterus lies to the front, its forward direction can be noted; if to the back, then the body will be felt on passing the finger

further up. Pushing the finger well upwards and passing it first to the right and then to the left, we feel the ovaries (more distinctly when enlarged) as small oval tender bodies (figs. 73 and 45).

Fig. 89 shows a common condition of the uterus which is frequently mistaken for and treated as a retroversion. We allude to the uterus anteflexed and drawn back by cellulitis of the utero-sacral ligaments. As such patients are usually nulliparous and have therefore somewhat unyielding abdominal walls which cause a difficult bimanual, and as a lump is felt through the posterior fornix, the diagnosis of retroversion

Diagnosis
of Ante-
flexion.




FIG. 73.

ABDOMINO-RECTO-VAGINAL EXAMINATION. Upper hand not shown. Note prolapsed ovary.

is often made. The rectal examination, however, clears up the case; as the finger feels the knee of the flexion and the uterine body going forwards from it.

The *upper hand* is used during the rectal examination just as in the Bimanual, *i.e.*, the examination is abdomino-recto-vaginal or abdomino-rectal. The simple rectal (with the finger in the rectum unaided by the other hand) does not give much information as to the condition of the uterus.

Where, from rigidity of the abdominal walls, it is difficult to press

down or fix the uterus with the external hand, this may be done with the volsella in the vagina. The use of the volsella enables us to draw the uterus better within reach of the finger in the rectum. This examination per rectum aided by the volsella will be considered in the next chapter.

Value of
Rectal
Examina-
tion.

Of all manual examinations of the pelvis, the abdomino-rectal or abdomino-vagino-rectal is the most thorough. In retroversion, prolapsed ovaries, and pathological ante flexion, it is of special value. A patient may object to it and refuse to allow it; and, of course, the practitioner must keep this in mind.

SIMON'S METHOD OF PASSING THE HAND INTO THE RECTUM.

Simon's
Method.

This consists in passing the whole hand through the sphincter ani into the rectum, and even up to the transverse colon. The patient is deeply narcotised; the hand is passed cautiously through, by inserting first two fingers and the others successively until the entire hand is passed; incision of the sphincter ani may be necessary. Sometimes an incurable incontinence of fæces has resulted.

The unanimous opinion of gynecologists is that this severe method of examination is unnecessary. Careful bimanual examination, aided when necessary by anæsthetics, gives equally good results.

Injection
of air.

For specialists it is of use to know that valuable results in minute precise rectal examination can be got by first injecting air into the rectum. The whole rectum up to the sigmoid flexure can be dilated, the sphincters made out and the bony pelvic wall carefully explored. It is necessary to add, however, that this is an adjunct to the rectal method of examination of use only in certain very rare instances.

This method we recommended in the first edition of the Manual, and we have often simply elevated the hips with the patient in the lithotomy posture and exposed the rectal walls with the ordinary Sims' speculum. Recently, Kelly of Baltimore has advocated the use of the genu-pectoral posture, the admission of air and the use of short tubular specula, and special scoops to aid in the examination.

CHAPTER IX.

THE VOLSELLA.

LITERATURE.

Goodell—Some Practical Hints for the Treatment and the Prevention of the Diseases of Women : Medical and Surgical Reporter, January, 1874. *Hegar*—Zur Gynäkologischen Diagnostik : Die combinirte Untersuchung, Volkmann's Sammlung, No. 105. *Simpson, A. R.*—The Use of the Volsella in Gynecology : Contributions to Obstetrics and Gynecology, p. 183. The literature is fully given in A. R. Simpson's paper.

WE have already seen that one of the most striking anatomical features Volsella and properties of the uterus is the considerable range of its mobility in almost every direction. It can be pushed upwards from its normal position $1\frac{1}{2}$ or 2 inches, and is displaceable forwards or laterally in a very marked degree. If laid hold of with the instrument known as a volsella, it can be drawn downwards (by a force not exceeding five or six pounds) until the os externum lies close to the vaginal orifice. This procedure facilitates, in suitable cases, diagnosis and treatment of gynecology so much that it is well worthy of the allotment of a special chapter to its discussion. We consider the following points:—

1. Description of instrument (fig. 74);
2. Method of use;
3. Mechanism of the displacement it causes;
4. Uses;
5. Contra-indications.

1. *Description of Volsella.*—Probably the best pattern of volsella is the one used in Germany, and known as the bullet forceps. It is a straight instrument, is provided with a catch, and has a bite that does not tear. There are, however, very many varieties of volsellæ, and the practitioner in choosing one should note that the bite is not too fine, and should consider a simple catch indispensable. As it is generally the anterior lip of the cervix that is laid hold of, and the volsella lies along the straight anterior vaginal wall, the slight pelvic curve usually given to the blades is unnecessary.

Method
of Use.

2. *Method of Use.* (a). *Without previous passage of Speculum.*—The patient is placed in the ordinary left lateral posture. Two fingers of the right hand are passed into the vagina, and the anterior lip of the cervix touched. The volsella, held in the left hand, is guided along between the index and middle exploring fingers; the anterior lip of the cervix is seized and drawn down. Rectal examination is now made. This plan is also useful in cases where the hymen is rigid, and where dilatation of the cervix is being employed for dysmenorrhœa. (Section VIII.)

(b.) *With the Speculum.*—For this see Chapter X.




FIG. 74.

MECHANISM OF DISPLACEMENT OF PELVIC-FLOOR SEGMENTS when Volsella is used.

Mechanism
of the Dis-
placement
caused.

3. *Mechanism of the displacement it causes.*—The uterus is drawn down so as to lie behind the symphysis pubis. If drawn down fully, as it may be in exceptional cases, it has its long axis in the vagina and the os externum near the vaginal orifice.

The vaginal walls are inverted: *i.e.*, when the os externum is at the vaginal orifice, we have a deep pouch behind and in front of the uterus.

The relations of the bladder and rectum are given in fig. 74.

Use in
Diagnosis.

4. *Uses.* (a) *In diagnosis.*—(1.) The cervix, which may seem “ulcerated,” as it is commonly called, is easily demonstrated by the volsella to be singly or doubly lacerated. For this purpose the anterior and

posterior lips are laid hold of, and when brought together the ulceration is seen to be due to laceration with eversion.

(2.) Abdominal tumours can be shown to be connected with the uterus or not as the case may be. If the patient be placed in the dorsal posture and the tumour be laid hold of by an assistant, then, when the uterus is drawn down, the tumour can be felt to descend, if fixed to it.

(3.) To the examination *per rectum* the volsella is a valuable addition. If one finger be placed in the rectum and the cervix laid hold of with a volsella and drawn down, the mobility of the uterus can be estimated; the whole posterior uterine surface may be palpated for small fibroids. The ovaries are made more accessible; and the uterus, especially if small, can have its length estimated by the rectal finger.

This method of examination of the uterus by rectum and volsella, judiciously conducted, is of the very greatest value.

It is evident that it will also help one as to the diagnosis of displacements of the uterus; but its value in this respect is somewhat lessened



FIG. 75.

SIMS' TENACULUM.

by the displacement its use causes. Thus it makes a retroversion less retroverted; an ante flexion less ante flexed; an anteversion less anteverted.

(b) *In treatment.*—In this the volsella is one of the most useful instruments the gynecologist possesses. Thus it helps greatly in the examination of the aborting uterus; in replacement of the gravid or non-gravid retroverted uterus; in insertion of sponge or tangle tents. In operations such as Emmet's for repair of the cervix, amputation of vaginal portion of cervix, excision of the uterus through the vagina, it is indispensable. Use in Treatment.

Details of its uses in these cases will be given under the special descriptions of the operations; and it will also be shown, in the Chapter on Specula, that by using the volsella the speculum may be dispensed with in certain cases.

5. *Contra-indications.*—It should not be used in acute peritonitic or cellulitic attacks, in distended Fallopian tubes, in hæmatocele or in advanced cancerous disease. No pain should be caused by its use provided only the vaginal aspect of the cervix is laid hold of. Contra indications.

The amount of traction to be made will vary with the necessities of the case. In many instances only a mere steadying action is requisite; in others the cervix has to be drawn half-way down the vagina. In Amount of traction to be used.

special cases the cervix is drawn down to the vaginal orifice or beyond it, as in amputation of the cervix or excision of the uterus.

For simply steadying the cervix, *Sims' tenaculum* is of great service (fig. 75). This is a form of sharp hook with a delicately made stem, diminishing to the point which is set on the stem almost at a right angle; the hook should be only very slightly curved in. In operating on carcinoma cervicis uteri, the volsella is occasionally unsuitable as the tissue is too friable. A hook may be passed into the cervical canal in such cases so as to draw down the uterus sufficiently.

CHAPTER X.

VAGINAL SPECULA.

LITERATURE.

Barnes—Diseases of Women: London, 1878. *Goodell*—Lessons in Gynecology: Philadelphia, 1880. *Hart*—Structural Anatomy: Edin., 1880. *Mundé*—Minor Gynecology: Wood & Co., New York. *Reid, W. L.*—History of the Vaginal Speculum: Glasgow Medical Jour., 1896, p. 161. *Sims, J. Marion*—Clinical Notes on Uterine Surgery: Hardwicke & Co., London, 1866. *Thomas*—Diseases of Women: Philadelphia, 1881.

WE have already seen that the segments of the pelvic-floor are separable when a woman assumes certain postures; that the sacral segment can be hooked up, and that by this means we get a view of the vaginal boundaries of these segments and of the os uteri. This is the natural method of opening up the pelvic floor; or the natural specular method.

Gynecologists had used various instruments for enabling them to look into the vagina: but all these proved unsatisfactory until Marion Sims, noting the natural postural dilatation of the vagina, introduced his famous duckbill speculum.

We take up the consideration of three types of speculum, viz.:—

Varieties.

1. Spatular—the duckbill or Sims speculum;
2. Tubular—The Fergusson speculum;
3. Bivalve—the Neugebauer, Cusco and other modifications.

We note under each its nature, the method of employing it, and the theory of its action and uses.

1. The SIMS OR DUCKBILL SPECULUM is shown at figs. 76 and 77.

Sims
Speculum
Nature.

Its Nature.—Each instrument in reality consists of *two specula*, which are of different size and connected by a handle; usually, however, we speak of these specula as the *blades* of the *speculum*. The real Sims speculum is light, has each blade slightly concave on its anterior aspect, and has the blades at *right angles* to the intermediate handle.

Modifications of Sims' speculum are numerous. Indeed, it seems difficult for gynecologists to resist modifying an instrument, and rare to find them improving it. The most widely known modification is Boze-Bozeman's. man's; it is heavier than Sims', has the blades meeting the handle at an

Modifica-
tions.

acute angle, and the blades more concave on the anterior aspect. (Figs. 77 and 78.)

One curious fact about almost all specula is, that they are too long. Sims' blade is 4 inches long, though the posterior vaginal wall measures only $3\frac{1}{2}$ inches. Thus, as we wish to expose only the anterior vaginal wall and cervix uteri, a 3-inch length of blade is sufficient.

The most valuable improvements in specula have been made to meet the necessities of such operations as vaginal hysterectomy and amputa-



FIG. 76.
SIMS' SPECULUM.

tion of the cervix. Thus a speculum for use in the lithotomy posture in these operations has been developed from Sims' and become indispensable in such work. It is shorter, broader, and may have various blades fitting one handle with a convenient catch, or the blade and handle may be in one piece (fig. 80). These allow the drawing down of the uterus, and dilate the vaginal entrance for freer manipulation in a way the Sims speculum cannot, and are thus an absolute necessity in certain cases.



FIG. 77.
SIMS' SPECULUM.



FIG. 78.
SIMS' SPECULUM modified by BOZEMAN.

Method
of Use.

The method of employing Sims' speculum.—Under this it is important to note :—(a) How to place the patient, (b) How to pass the speculum, and (c) How to hold it when passed.

Position of
Patient.

(a). *How to place the patient.*—The patient must be placed in the Sims or semiprone posture. This is briefly as follows: the patient lies almost on the breast; the lower left arm is over the edge of the couch next the gynecologist; the hips are close to the edge; the knees are well drawn up; and the upper or right knee touches the couch with its inner aspect. The posterior aspect of the sacrum is therefore oblique to the horizon.

As the result of this posture—a modified genupectoral one—the vaginal walls separate when air is admitted; the pubic segment passing down with the viscera, the sacral one remaining behind.

(b). *How to pass the speculum.*—Choose the blade which is of the proper size to pass the vaginal orifice; warm it, and oil it with the fingers on its convex aspect only. The concave surface must be dry to reflect light, and therefore the speculum should never be oiled by dipping it. Hold it by the other blade in the left hand, as shown at fig. 79. Then pass the index and middle fingers of the right hand into the vagina to separate the labia; carry in the speculum between them; push it onwards, following the curve of the posterior vaginal wall, until the beak of the instrument lies in the posterior fornix. Now draw the instrument back as a whole, in a direction at right angles to the posterior vaginal wall; then turn the beak forwards, so as to bring the cervix more

Passage of Speculum.




FIG. 79.

One method of holding the SIMS SPECULUM.

into view. Finally, tilt the blade so that the beak lies on a lower level than the proximal end of the blade; this keeps up the upper labium.

(c). *How to hold the speculum when passed.*—When passed, the speculum may be held as shown in fig. 79, or the hand may rest on the patient's hip while grasping the instrument; the cervix may be drawn down with a volsella. Various attempts have been made to add to the Sims speculum a means of rendering it self-retaining; the majority of these are by no means successful, and therefore we need not describe what is seldom used.

How it is held.

Theory of action and uses of the Sims speculum.—The Sims speculum is based on the effects consequent on the genupectoral posture. When the patient is semiprone and the vaginal orifice opened, the segments of the pelvic floor separate; and then the Sims speculum is a simple means of hooking the sacral segment well back.

Action and Uses of Sims' Speculum.

The Sims speculum is, on the whole, by far the most useful speculum for examination and for operative work, not requiring downward traction on the uterus and great dilatation of the vaginal entrance.

Method of use of the modified Sims speculum with removable handles.—The patient is placed in the lithotomy posture, and the appropriate blade fixed on the handle and passed as already described under (b). It is then pulled downwards, the cervix taken hold of with a volsella, and the operation proceeded with. The lithotomy posture has all the advantages of the Sims posture, and is more convenient for operative work.

If the operator is single-handed in performing, *e.g.*, such an operation as curetting, he may fasten the volsella with a tape to the sheet on which the patient lies. He can then hold the speculum with his left hand and this leaves the right for manipulation.




FIG. 80.

MODIFIED SIMS SPECULUM in one piece (*Doyen*).

Fergusson
Speculum.

2. The FERGUSSON SPECULUM is seen at fig. 81. It is made in three suitable sizes; and may be described as a glass tube, with a proximal trumpet and a distal bevelled end. It is made of glass, silvered on the inside and coated with caoutchouc. The bevelling of the distal end makes a shorter anterior side and a longer posterior one. The maker's name is usually placed at the trumpet end, at the foot of the anterior side, and serves to indicate that side when the speculum is in the vagina.

How used.

Mode of employment of the Fergusson speculum.—The patient lies in the left lateral position with hips raised. Warm the speculum, and oil it on the outside. Take it by the trumpet end with the right hand and

pass it into the vaginal orifice previously opened up by index and middle fingers of the left ; now push it in, short side to the front, until arrested. By looking along it, the practitioner can note if the cervix is in view. It is generally not so, but may be snared by the following manœuvres : carry the trumpet end well back towards the perineum, and then depress the distal end first to the left and then to the right, finally turning it round if these fail. In multiparæ with lax vagina it is easy to pass the Fergusson ; but it is more difficult in nulliparæ.

The Fergusson is a favourite speculum with many. It is useless in vaginal and cervical surgery, but with it applications to the cervix can be made very well and easily. When used for making applications to the endometrium, it is advisable to pull the cervix well down with a volsella after the speculum is passed, and to use a straight sound covered with cotton wool.

3. Of *bivalve specula* there are various forms ; the Neugebauer with its modification—the Crescent Speculum of Barnes ; the Cusco, which is often called the Bivalve Speculum ; and other varieties.




FIG. 81.

FERGUSSON SPECULUM.

The NEUGEBAUER is like a Sims speculum divided transversely at the middle of the handle. It is also made in suitable sizes.

Mode of employment.—Warm and oil *two blades*. Introduce one blade (the broader one) with its convexity touching the posterior vaginal wall. Then introduce the other with its convexity touching the anterior vaginal wall and so that its edges fit within the edges of the posterior vaginal wall blade. The beak of the posterior blade is thus in the posterior fornix ; that of the anterior blade in the anterior fornix. From their contact a leverage is obtained on approximating the handles, by which traction is made on the fornices, and the cervical canal more or less everted.

The Neugebauer and Crescent specula are useful in making cervical and endometric applications, and are better specula than the Fergusson.

The CUSCO OR BIVALVE SPECULUM is shown at fig. 82. It is composed of two blades jointed on to one another at their bases. The blades are opened to the desired distance by pressure on the thumb-piece, and kept open by a screw. It is introduced with its blades right and left,

and then turned so that they lie anterior and posterior, that with the screw being posterior. It is then pushed onwards, and the blades opened and fixed by the screw. Care should be taken not to catch any of the hair in the screw; and, in withdrawing it, not to pinch up the vaginal walls.

The Cusco speculum is self-retaining and useful in cervical and endometric applications.

W. L. Reid of Glasgow has introduced another variety of bivalve speculum which he has found useful. In it the blades are separable and move on parallel bars.

If the patient be placed in the genupectoral or semiprone posture, the posterior vaginal wall hooked back with the fingers and the cervix drawn down with a volsella, a useful view can be obtained without the aid of any speculum.




FIG. 82.

CUSCO'S SPECULUM.

USES AND COMPARATIVE VALUE OF THE VARIOUS SPECULA.

The Sims is undoubtedly the best and most scientific type of speculum we possess. When properly used, and aided by the volsella or tenaculum, it leaves nothing to be desired in certain cases. The operative development of the past few years has led to the abandonment of the Sims posture by most gynecologists, and the almost exclusive employment of the lithotomy one for operative work. The specula now mainly used are modified Sims' specula (v. fig. 80), *i.e.*, the blades are shorter and broader. They thus allow of the cervix being well drawn down, and broaden the vaginal entrance. For operative cases their use is imperative.

The Fergusson is easily passed, involves only slight exposure, and is good in very minor gynecology. It gives only a limited view of the

vaginal walls. The student should note that it brings the flaps of a split cervix together and somewhat conceals the lesion.

The Neugebauer, on the other hand, opens up a cervical split, and may do this so effectually as to give the impression that there is none. The Fergusson and Cusco are *self-retaining*.

CHAPTER XI.

THE UTERINE SOUND.

LITERATURE.

Simpson, A. R.—The Uterine Sound : Ed. Med. Journal, 1882. *Simpson, Sir J. Y.*—Memoir on the Uterine Sound, Selected Obst. Works : A. & C. Black, Edinburgh, 1871.—See *Munde's* Minor Gynecology and *Thomas* as to Huguier & Lair.

Uterine
Sound.

WE shall consider this important gynecological instrument as follows :—

1. Its nature ;
2. Preliminaries to its use, contra-indications ;
3. Method of use, difficult cases ;
4. Employment for diagnosis and treatment ;
5. Dangers attending its use ;
6. Relation to bimanual and rectal examination.

NATURE.

Nature.

The sound of Sir James Simpson is not only the classical instrument, but, taken all in all, is probably the best. We describe it, therefore, as a type of the instrument, and then consider its modifications.



FIG. 83.

SIR J. Y. SIMPSON'S SOUND.

Simpson's sound is a rod of flexible metal 12 inches long, specially graduated, and provided with a suitable handle (fig. 83). It is made of copper, nickel-plated ; this is sufficiently pliable to be moulded, and yet sufficiently stiff to retain any special shape given to it. Instrument-makers often make this sound too unyielding. It should be always pliable enough to be bent with two fingers.

The handle has the shape shown at fig. 83. Note that it is roughened

on the same side as that towards which the point of the instrument lies. Consequently, when the sound is in the uterus, we can tell the direction of the point by noting this roughness on the handle.

The graduation is important. $2\frac{1}{2}$ inches from the point is a rounded knob: this is the length of the fully-developed unimpregnated uterine cavity. Other markings are $3\frac{1}{2}$ inches, $4\frac{1}{2}$ inches, $5\frac{1}{2}$ inches, and so on up to $8\frac{1}{2}$ inches. The notch, $1\frac{1}{2}$ inches from the point, is of little use and weakens the instrument.

The modifications of this instrument are numerous. The changes are chiefly in its flexibility, lightness, and in the use of another material.

A. R. Simpson has modified the instrument by making it shorter, abolishing the $1\frac{1}{2}$ inch notch, and squaring the handle (fig. 84): this gives a very handy and useful instrument.




FIG. 84.

A. R. SIMPSON'S SOUND.

PRELIMINARIES TO ITS USE: CONTRA-INDICATIONS.

No instrument should have the preliminaries to its use more carefully considered. The rash and careless use of the sound may do immense mischief to the patient. Note, then, *when not to use it*. Prelimi-
naries to
Use.

- (1.) The sound is not to be passed during an ordinary menstrual period.
- (2.) It is not to be passed in an acute inflammatory attack of uterus, ovaries, pelvic peritoneum, or connective tissue.
- (3.) It is not to be passed in cases of cancer of the cervix or body of the uterus.
- (4.) It is not to be passed if the patient has missed a menstrual period. This is a safe rule, but admits of limitation, as we shall see afterwards.

Before using it, attend to the following points.

- (1.) Ascertain that the patient has not missed a period.
- (2.) Do the bimanual carefully. If in doubt, use the rectal examination aided by the volsella.
- (3.) Place the patient in the left lateral posture.
- (4.) Give the sound the curve you find the uterus to have.
- (5.) Never pass the sound without preliminary disinfection of the external parts and vagina. This restricts its use in the consulting room.




FIG. 85.

FIRST STAGE OF PASSING THE SOUND.

METHOD OF USE.

Method of
Use.

After the preliminaries mentioned above, take the sound, previously disinfected, in the hand. Pass the index finger of the right hand into the vagina and touch the anterior lip of the cervix, *i.e.*, in front of the os. Guide the sound along the vaginal finger and make the point enter the os uteri (fig. 85). Pass it in for an inch or so, to fix it.

When
Uterus Re-
troverted.

If the uterus be retroverted then carry the handle towards the symphysis, when the point of the instrument will glide into the uterine cavity

until arrested by the fundus (fig. 86). No force is needed. If force seems necessary, the instrument should be withdrawn and a more careful Bimanual performed.

If the uterus lie to the front, the procedure is different. Pass the sound ^{When Uterus to front.} as already described until it has entered the cervix for an inch or so (fig. 85). Note now that the point of the sound looks back, whereas the





FIG. 86.

SECOND STAGE OF PASSING THE SOUND when UTERUS is Retroverted.

fundus lies to the front. Clearly, we must make the point look to the front. This is done by turning the handle so that its roughened surface looks to the front. To do this we do not twist round the handle on its long axis, but make it sweep round the arc of a wide semi-circle as in fig. 87. The point, during this manœuvre, remains fixed or nearly so. Now carry the handle back to the perineum when the point glides into the cavity (fig. 88).

Another way of passing the sound, when the uterus lies *to the front*, is as follows. Place the patient well across the bed. Do Bimanual and



(a.) Proper method of TURNING THE SOUND, contrasted with improper method (b.).

curve sound appropriately. Take the sound in the right hand. Pass two fingers of the left hand, palmar surface forward, into the vagina, and




FIG. 88.

SECOND STAGE OF PASSING THE SOUND when UTERUS is to the Front.

touch the posterior lip of the cervix. Carry the sound, point looking forwards, into the vagina ; make it enter the os, and then carry the handle

towards the perineum, when the point will glide on. This method avoids the sweeping round of the handle, and is useful if the uterus is very much anteverted.




FIG. 89.

SOUND ARRESTED (before Rotation) in a case of Anteversion.

The sound may be passed after the uterus is drawn down with a volsella, or after the Sims speculum has been introduced.

The practitioner will find it, however, better and safer to replace the uterus by bimanual manipulation than with the sound.

Difficult
Cases for
Sound.

Difficult Cases.—These are chiefly found in markedly anteflexed uteri. The sound passes in so far (fig. 89), but when turned has its point looking too directly upwards. In such cases first draw the cervix down with a volsella, now pass the sound, and should it still stop at the flexion make pressure with a finger in the anterior fornix to push up the fundus. Then get an assistant to carry the handle of the sound towards the perineum.

EMPLOYMENT OF THE SOUND FOR DIAGNOSIS AND TREATMENT.

(A) DIAGNOSIS.

Use of
Sound in
Diagnosis.

(1.) *Length of uterine cavity.* This varies in different pathological conditions. Thus the cavity is

(a) Lessened in Superinvolution of uterus,
Atrophic uteri ;

N.B.—The sound easily perforates the thin wall of the super-involuted uterus ; this may do no harm. It may also pass along the Fallopian tube.

(b) Increased in Subinvolution of uterus,
Hypertrophy of uterus,
Cervical hypertrophy,
Endometritis,
Submucous fibroids,
Interstitial fibroids,
Small uterine polypi,
Prolapsus uteri.

(2.) *Direction of uterine axis ;* whether retroverted, anteverted, lateri-verted.

(3.) *Relation of axis of uterine body to that of cervix ;* whether we have anteflexion or retroflexion.

(4.) *Stenosis and atresia at os internum and os externum ; tenderness of fundus,* as in endometritis.

(5.) *Mobility of uterus.* This should be ascertained in the following way. Pass the sound as already described. Make the patient turn on her back, and then place two fingers in the vagina, palmar surface upwards and touching the posterior lip of the cervix. The sound lies on the palm of the hand, is steadied with the thumb, and can be used to move the uterus gently about as desired.

(6.) *Rough condition of endometrium ;* often associated with bleeding when sound is passed.

(7.) *Differential diagnosis between uterine polypi projecting into vagina, and inverted uterus, etc.*—When we have a polypus to deal with, the

sound passes in through the cervix for more than the usual distance because the uterine cavity is enlarged. In inversion, it passes for only a short distance into the cervix and is then stopped by its reflexion. Sometimes, however, the neck of the polypus is adherent all round to the cervical canal, thus simulating inversion: and in some very rare cases the mucous membrane of the uterus becomes separated and expelled from the uterine cavity, simulating inversion of the whole uterus owing to the separation stopping at the os internum. It is evident that in these last two cases the bimanual clears up the diagnosis, the upper hand feeling the body of the uterus in its normal position in both of them. The sound is only confirmatory of the bimanual.

(B) TREATMENT.

- (1.) *Rectification of abnormal angular relation between the uterine body and cervix (anteflexion, retroflexion); dilatation of uterine canal as a whole, or of stricture at os internum.* Use of Sound in Treatment.
- (2.) *Replacing of retroverted unfixed uterus.*
- (3.) *Application of acids to endometrium on the sound covered with cotton wool.*

DANGERS ATTENDING ITS USE.

The great danger to the patient from the passage of the uterine sound are abortion, and abrasion of the uterus membrane with absorption of septic matter and resulting pelvic cellulitis or peritonitis. Dangers of Sound.

The former untoward result must be very carefully guarded against. One valuable caution is never to omit the question as to the menstruation, and to ask if it was the usual amount, as some women have a slight discharge of blood at the first period after they conceive. The best safeguard is the careful performance of the bimanual. This soon teaches the practitioner to know whether he has an unimpregnated uterus between his hands, or one at the second or third month of gestation. Special care should be taken when the uterus is retroverted: it may be also gravid; and the pregnancy may, by causing pressure, have induced the patient to consult a medical man. As the bimanual is often difficult, an unwary use of the sound may make the diagnosis disagreeably evident.

The means to avoid setting up any inflammatory disturbance are—to perform the bimanual carefully, to curve and oil the sound properly, and to pass it gently and with antiseptic precautions.

SOUND COMBINED WITH BIMANUAL.

The importance of this method of examination has been pointed out by A. R. Simpson. For its performance the short sound with the square handle (fig. 84) is necessary. It is of such a length that, Use of Sound in Bimanual.

when the middle finger is at the knob, the flat surface of the handle rests on the ball of the little finger, against which it is steadied by the flexed little and ring fingers.

The sound is introduced into the uterus in the ordinary way. The fingers are passed into the vagina as for a vaginal examination, and the sound grasped as in fig. 84. Or the sound may be steadied with the middle finger while the index is used to feel the uterus through the




FIG. 90.

SOUND combined with BIMANUAL Examination (*A. R. Simpson*).

anterior fornix (fig. 90). The external hand is placed as in the bimanual.

This method is specially useful (*a*) when the uterus is flaccid; the sound stiffens it, and enables the external hand to define it: (*b*) when, from the presence of small fibroids or pelvic deposits, there is doubt as to what is the fundus uteri; the sound felt by the external hand in the uterus, indicates the fundus.

RELATION OF SOUND TO BIMANUAL AND RECTAL EXAMINATION.

Before Sir James Simpson introduced the use of the sound, gynecological examination was confined to the exploration of the vagina and cervix.

Relation of
Sound to
Bimanual
and Rectal
Examina-
tion.

Simpson gave an immense impulse to gynecology, by placing in the hands of gynecologists an instrument which explored the uterine cavity above the cervix, and enabling them to obtain a perfection of diagnosis before undreamed of; thus gynecological examination came to consist of a vaginal examination, and then a passage of the sound, due attention being given to the non-existence of pregnancy. He recommended, further, the elevation of the uterus with the sound, and its definition with the upper hand.

The next step in gynecology was the use of the two hands—the bimanual and rectal examinations—which in the last twenty years has developed immensely. Consequently the use of the sound has become more limited. The teaching in this chapter has been based on a recognition of this fact, and the use of the sound is recommended only in the rare cases where after the bimanual, rectal, and volsellar examinations have been carefully employed, the necessary information was not obtained. Even then the sound should not be passed in the consulting room where cleansing of the patient is impracticable, but only where suitable precautions can be used. The skilled gynecologist will restrict the use of the sound, with few exceptions, to the operating table. Thus in curetting, it is advantageous to pass the sound after all antiseptic precautions have been used, and the uterus has been drawn down so that the direction of the cavity may be ascertained exactly. The drawing down of the uterus has changed its axis, and the preliminary passage of the sound gives the operator accuracy in passing the dilator and curette.

Much therefore of this chapter is mainly of historical interest, as the use of the sound has been markedly limited by advances in the use of the bimanual, and by a clearer knowledge of the cause of pelvic inflammatory attacks.

CHAPTER XII.

TENTS AND OTHER UTERINE DILATORS.

LITERATURE.

Hegar und Kaltenbach.—Die Operative Gynäkologie: Stuttgart, 1881. *Landau*—Ueber Erweiterungsmittel der Gebärmutter: Volkmann's Sammlung, No. 187. *Lewers*—On Rapid Dilatation of the Cervix Uteri: Lancet, 1887, II., p. 507. *Mundé*—Minor Gynecology: New York, 1881. *Simpson, J. Y.*—Selected Obst. Works, Vol. I.: Edinr. 1871. *Sims, J. M.*—Uterine Surgery: London, 1867. See also Index of Recent Gynecological Literature in Appendix.

Uterine
Dilators.

HITHERTO we have considered only the means which have placed the vagina and cervix within range of digital examination. In this section we take up the methods by which we get digital examination of the uterine cavity—methods of the highest practical value, which, like the sound, we owe to the genius of Sir James Simpson.

We therefore consider the following methods of dilating the cervical canal:—

- I. *Slow dilatation with sponge tents, tangle tents, tupelo tents;*
- II. *Rapid dilatation with graduated hard-rubber dilators—Tait's, Hanks', and Hegar's;*
- III. *Dilatation by incision and screw dilators (v. Chap. XXVI.).*

DILATATION BY SPONGE, TANGLE, AND TUPELO TENTS.

Sponge-
tent—
material.

1. *Material*.—The sponge tent is a cone of good, unbroken, thoroughly dried sponge, impregnated with some antiseptic, and then firmly compressed into small transverse bulk, its original length being preserved. When thus prepared and placed under conditions where it can absorb moisture, it swells up; and in thus expanding dilates any dilatable structure which may grasp it.

Good sponge tents of various sizes may be had from all chemists. In order to prevent the antiseptic from volatilizing, the sponge tents are covered with grease. They are provided with a tape at the base to aid their extraction from the cervix after use.

Tents are also made from the ordinary sea-tangle (*laminaria digitata*) (fig. 91), and from tupelo wood (*nyssa aquatilis*). It is certainly the case that the tupelo expands more rapidly than either tangle or sponge. Fig. 92 shows its power in this respect. Tangle tents may be had hollow; this facilitates the imbibition of moisture but weakens their expanding power.

2. *Purposes for which used.*

(1.) To restrain hæmorrhage in cases of abortion, and at the same time dilate the cervix for further interference. Uses of
Tents.




FIG. 91.

Shows on the left a straight and a curved tangle tent, and on the right these tents after expansion. Note how one has been gripped by the os internum (*Mundé*).

(2.) To dilate the cervix and uterine cavity, and enable the practitioner to ascertain and remove the cause of pathological uterine hæmorrhage, whether due to endometritis, sarcomata, polypi, or incomplete abortion.

(3.) To correct pathological flexions of the uterus, or to dilate a stenosed cervix. Their use for this is not only unnecessary but dangerous.

Scope of
Tangle and
Tupelo
Tents.

Tangle tents have the same scope as sponge tents. They do not, however, expand so well and thoroughly. Their special advantages are due to their smaller size, and the fact that several may be passed at the same time into the cervix. They may be used, therefore, in cases of narrow cervix and in flexions. Tupelo tents are very good; they are easily passed and, from their rapid expansion, preferable to sponge tents.




FIG. 92.

DIAGRAM to show relations between size of Tupelo Tent, before and after expansion. The dotted outside line indicates the size of the tent after expansion (*Mundé*).

Preliminaries and Mode of Use.

3. *Preliminaries to and Method of use.*—Tents should not be passed during an ordinary menstrual period, although they often require to be used when pathological bleeding is going on. They should always be passed at the patient's own house; and she should be kept strictly in bed during their use, and for some time after. Before their use, the vagina should be thoroughly washed out with warm carbolic lotion (1-40), or with corrosive sublimate (1-2000). Schultze, in passing tangle tents for flexions, first ascertains the uterine curve with the sound; if blood follows its use, he postpones the introduction of the tent for forty-eight hours, in the meantime applying pure carbolic acid to the endometrium.

Before using the sponge tent, it is advisable to remove most of the grease covering it.

Sponge tents may be used in various ways.

(1.) The patient is placed in the genufacial, or better, in the semi-prone posture. Sims' speculum is passed, the anterior lip of the cervix ^{How} passed.



FIG. 93.

EXPANDED TUPELO TENT with construction at os internum (*Mundé*).

laid hold of with a volsella and drawn down. The sponge or tangle tent, held in forceps, can then be passed into the cervix (fig. 95).

(2.) The tent is fixed on the spike of an appropriate instrument, and is then passed like the uterine sound; *i.e.*, with the patient placed in the left lateral position, the index and middle fingers carried into the




FIG. 94.

SPONGE TENT POLYPUS OF SIR JAMES SIMPSON. (†)

Drawing of the uterus which contained a polypus—obtained from a patient of Sir James Simpson's, who died from the hæmorrhage it caused. It was this preparation which suggested to him the sponge tent.

vagina and placed on the anterior lip of the cervix. The tent, fixed on the spike, is passed along these fingers and its point made to enter the cervix. The handle is then rotated and carried to the perineum.

(3.) The patient is placed on her left side and athwart the bed. Pass

the volsella, draw the anterior lip of the cervix down. The volsella is not always needed. Place the tent between the index and middle fingers of the left hand with the thumb at the base, carry these fingers into the vagina with their dorsum on the posterior vaginal wall, make the point of the tent enter the cervix and push it on with the thumb.

Another way is to use the volsella as above described, but to fasten it to the bed. Then pass Sims' speculum, holding it with the left hand, so that the tent held in the right hand can be passed into the cervix without difficulty.

Occasionally, difficulty is experienced in passing a tent owing to marked anteversion of the uterus. If the cervix be drawn down with




FIG. 95.

SIMS' DIAGRAM ILLUSTRATING PASSAGE OF TANGLE TENT. Patient is semiprone, Sims' speculum passed, and cervix steadied with tenaculum. The tent is passed with forceps.

a volsella, the difficulty may be overcome; or it may be necessary to partially retrovert the uterus bimanually prior to passing the tent.

Moulding
of Tangle
and Tupelo
Tents.

Tangle and Tupelo Tents.—The same instructions as for sponge tents hold good. Tangle tents, however, when used to correct flexions must first be moulded as follows:—Ascertain the curve of the uterus by Bimanual and sound, select a suitable tent and dip it for a few seconds in boiling water, then mould it to uterine curve and pass it as already explained.

Tents require to be left in the cervix for a period varying from twelve to fifteen hours, and the vagina should be frequently douched with carbolic lotion during this time. At the end of this period the tent should be removed. During the removal no great force should be used. Sometimes the removal is difficult owing to constriction by the os internum or to irregularities in the mucous membrane.

The cervix is generally then sufficiently dilated to admit of digital examination of the endometrium.

4. *Dangers of sponge and tangle tents and contra-indications.*—The practitioner must keep prominently before him that the use of a tent may prove by no means a harmless measure. Cases of death from septi-
cæmia after the careful and proper use of *one* tent have occurred. The patient runs a risk proportionate to the number used; and, therefore, it is not advisable to use more than two consecutively unless under special circumstances. They are not to be used if acute or sub-acute pelvic inflammation, pyosalpinx, ovaritis (acute or chronic), carcinoma cervicis, or pelvic hæmatocele be present.

Dangers
and
Contra-
indica-
tions.

The reason why sponge tents may prove dangerous is only too apparent. The uterine mucous membrane is a lymphatic surface absorbing most rapidly. We cannot insert sponge tents with Listerian precautions; and, in addition, we have the expanding pressure of the tent forcing septic matter into the mucous membrane.




FIG. 96.

TAIT'S DILATORS.

To sum up briefly, tents are highly useful in necessary cases—no means at the disposal of the gynecologist gives him in proper cases such valuable help; but he should not forget the risks occasionally arising from their use—risks which should make him cautious but not timid.

RAPID DILATATION BY GRADUATED HARD-RUBBER DILATORS—
TAIT'S, HEGAR'S.

The statement already made as to the dangers attending the use of slowly expanding tents would lead one to expect that attempts at rapid dilatation have been made. For this purpose, graduated vulcanite dilators have been employed by Tait, and Hegar.

Hard
Rubber
Dilators—
Tait's,
Hegar's.

Tait's dilators consist of graduated vulcanite cones (fig. 96) which can be screwed on to a suitable handle. The proximal end of the handle is perforated for elastic bands which, passing in front and behind, are attached to a suitable belt round the patient's waist. Thus the elasticity of the bands causes the cone gradually to pass up into the cervix, dilating it as it goes. By this apparatus, Tait claims to avoid septic infection and to dilate rapidly. The obvious objection is the

amount of watching it entails and the absence of the pelvic curve on the handle.

Hegar's dilators consist of a series of slightly curved stems $4\frac{3}{4}$ in. to $5\frac{1}{2}$ in. (12–14 cm.) in length, with a short flat handle 2 in. long, numbered from 1 to 30 and with diameters ranging from about $\frac{1}{12}$ in. to $1\frac{1}{2}$ in. (2–30 mm.).

There is little doubt that, to prevent sepsis, vulcanite dilators are the best. For dilating the cervical canal quickly in order to explore the uterine cavity with the finger, for the removal of polypi, or for curetting, they are specially indicated and are to be used as follows. In a case, for instance, where the cervical canal is to be dilated in order to gain access for the removal of a polypus, the patient is chloroformed, placed in the




FIG. 97.

HEGAR'S DILATOR. The lower figure represents the dilator (No. 15) complete, reduced to one-third scale; the two upper figures show cross sections of the smallest (No. 1) and the largest (No. 30) sizes.

lithotomy posture and the vaginal douche employed. Hegar's dilators, which are lying in a solution of corrosive sublimate 1 in 2000, are then passed, until sufficient dilatation is obtained. The polypus is then removed, and the uterine cavity carefully douched.

We recommend therefore the use of the tupelo tents in cases of threatened abortion where the practitioner has not sufficient assistance to enable him to use the vulcanite dilators. Where, however, this assistance can be procured, especially for exploration, curetting, and endometric applications, Hegar's dilators are the safest and best.

CHAPTER XIII.

THE CURETTE.

LITERATURE.

Ballantyne, J. W.—Curettag of the Uterus: Edin. Medical and Edin. Obst. Trans., 1896. (This paper gives the literature fully.) *Hart, D. Berry*—Uterine Curetting: Manchester Medical Chr., 1895. *Mundé*—The Dull Wire Curette in Gynecological Practice: Ed. Med. Jour., XXIII., p. 819. *Recamier*—Memoire sur les Productions Fibreuses et les Fongosités Intra-utérines: Union méd., IV., p. 66, 1850. *Simon*—Die Auslöfflung breitbäsiger weicher sarkomatöser und carcinomatöser Geschwülste aus Körperhöhlen: Beiträge zur Geburtshülfe von der Gesellschaft in Berlin, 1872. *Sims, J. Marion*—Clinical Notes on Uterine Surgery: London, 1866. *Thomas*—Diseases of Women: London, 1882. *Werth, R.*—Untersuchungen über die Regeneration der Schleimhaut nach Ausschabung der Uteruskörperhöhle: Arch. für Gynak., Bd. 49, 1895.

THE curette is an instrument, provided with a cutting or a dull edge, Curette. which can be introduced into the uterine cavity (previously dilated by tents if necessary) for the purpose of scraping off or removing abnormal endometric granulations, sarcoma of the mucous membrane, carcinoma of the cervix, or the remains of an incomplete abortion. This instrument has had a somewhat chequered career. Originally introduced by Recamier, whose instrument was stiff and sharp, it did good work in some cases; but fell into disrepute, undoubtedly deserved, after the




FIG. 98.
LOOP CURETTE. (4)

record of certain instances where its use had caused perforation of the uterus. Marion Sims and Simon recommend a modified instrument which, owing to its stiff unyielding nature, did not at first find much favour with the profession. Thomas then introduced his flexible dull wire curette, but this has now been found too feeble and a return has been made to stronger instruments.

There are four varieties of curette—(1.) Loop with sharp edge (fig. 98); (2) Simon's Spoon (fig. 99); (3.) Thomas' Dull Wire (fig. 100); (4.) Roux's (fig. 101). Of these we recommend Roux's.

Uses of
Curette.

Cases in which the Curette is useful.—The curette may be used to remove a piece of intrauterine tissue for aid in diagnosis. It is most frequently employed to remove abnormal tissue, in abortion, sarcomatous or carcinomatous diffuse growth, and endometric conditions.




FIG. 99.

SIMON'S SPOON. (3)

Simon's
Spoon.

Method of Use.—We take curetting for incomplete abortion as a type of procedure. The instruments necessary are Sims' speculum, volsella, Hegar's dilators, sound or probes armed with cotton wool, and Fritsch's uterine double catheter (fig. 108). The instruments are sterilised and placed in carbolic lotion (1-40). The patient is placed semiprone or, if



FIG. 100.

THOMAS' DULL WIRE CURETTE, with knob added by A. R. Simpson. (4)

Thomas'
Curette.

chloroform is given, in the lithotomy posture: the speculum is passed and the cervix steadied with a volsella. Hegar's dilators are now introduced until the cervical canal is patulous enough to admit the curette easily. The curette is then employed by being passed systematically over the anterior and posterior surfaces from above downwards,



FIG. 101.

RECAMIER-ROUX CURETTE.

Roux's
Curette.

and the mucous membrane down to the musculature removed. No force is required, and the finger can make out by the feeling of the curette when the resistant muscle is reached.

In cases of incomplete abortion, it is of importance to dilate sufficiently to enable the finger to be passed in. If not, detached and somewhat large pieces of the deciduæ may be retained.

The cavity of the uterus is then washed out with a mercurial lotion, and pure carbolic acid applied.

We have in the chapter on the Etiology of Pelvic Disease described a class of case under the term of chronic-infected cases, where we have multiple minor non-suppurative lesions, often depending on previous abortion and where endometritis is a prominent condition. In such, curetting is very valuable. This point will, however, be discussed again afterwards.

In endometritic conditions, curetting is invaluable, and should supersede all minor intra-uterine medication. The practitioner will find curetting in suitable cases one of the most satisfactory operations in gynecology.

Cautions and dangers.—The same precautions should be used as given under sponge tents. The dangers are *nil* when the operation is properly performed. Cautions
and
Dangers.

RELATION OF POSTURE TO EXAMINATION AND TREATMENT.

We have already mentioned several postures as being the proper ones for certain manipulations; and we here sum up briefly what it is of use to know in regard to these.

The *lateral posture*, where the patient lies on her side in the ordinary way, is convenient for vaginal examination; for the use of Fergusson's, Neugebauer's, or Cusco's speculum, and the passage of the sound and catheter.

The *dorsal posture* is imperative for abdominal examination and the bimanual.

The *semiprone* is the best posture for passage of Sims' speculum or for vesico-vaginal fistula operation.

The *lithotomy posture* is specially valuable for operations on the perineum, vaginal walls, cervix, and uterus.

The *genupectoral posture* is used in replacement of the retroverted gravid uterus and for examination of the bladder.

CHAPTER XIV.

KNIVES; SCISSORS; NEEDLES; SUTURES; DOUCHES AND SYRINGES; CAUTERY; ANÆSTHETICS.

KNIVES.

Knives. For vaginal and cervical surgery, long-handled knives with the blade straight or at an angle to the shaft are required (*v.* under operation for vesico-vaginal fistula), but the main use of the knife to the gynecologist is in abdominal section. The handle should be made of metal so as to allow of sterilisation by boiling in soda solution.

SCISSORS.

Scissors. These are of the greatest use to the gynecologist and have superseded the knife in all perineal operations. Curved scissors are necessary for fistula cases (fig. 102), Bozeman's being specially good. They are right




FIG. 102.

SIMPLE CURVED SCISSORS.

and left, but no woodcut gives a proper idea of their curves. For cervical operations, stout and sharp scissors are necessary as the vaginal portion of the cervix is tough. In perineal operations a pair of short angled scissors is best. For Hysterectomy, probe pointed scissors are very useful, but sharp pointed ones, with locking blades, are also employed.

NEEDLES.

We need only note that for cervical and fistula operations strong Needles. short needles either curved or perfectly straight are needed. The cervical tissue is so dense that markedly curved needles snap when slight. They are passed with a needle-holder, of which fig. 104 shows a Needle simple form. Curved or tubular needles set on handles are also Holders. useful.




FIG. 103.

FORMS OF NEEDLE (*Emmet*).

Hagedorn's needles are flattened laterally and full-curved. A special needle-holder is necessary for them. For perineal operations strong full-curved needles are necessary.




FIG. 104.

NEEDLE-HOLDER.

SUTURES.

These may be silver wire, carbolized silk, catgut, silk-worm gut, or horse-hair. For fistulæ, deep stitches, and cervical lacerations, *silver wire* or *silk* may be used. Silk-worm gut has not in our hands given good results. *Catgut* is valuable in the rectal stitches of complete rupture of the sphincter ani; and is now largely used instead of silk for operations on the cervix, vagina, and perineum, as it obviates the necessity of removing the stitches afterwards. Chromic acid gut is excellent, and juniper gut is also a thoroughly reliable suture. Practically we rely on silk for the ovariectomy pedicle, ligature of the broad ligaments, and for the abdominal wound; for other purposes catgut is best (*v.* also Chap. XV.).

VAGINAL SYRINGES AND DOUCHES: UTERINE DOUCHE.

For the purpose of applying antiseptic and astringent lotions to the vagina and split cervix, for hot-water injections, and for merely cleansing purposes, the vaginal syringe and douche are employed.

Vaginal
Syringe.

Vaginal Syringes.—Fig. 105 shows the well-known Higginson syringe. Valuable as this is, it is difficult for ordinary patients to manage single-handed. For them we should therefore recommend the

Vaginal
Douche.

Vaginal Douche.—A convenient form of this is shown at fig. 106. It can be hung up after being filled, and a gentle flow is thus obtained by




FIG. 105.

HIGGINSON'S SYRINGE.

gravitation. The overflow from the vagina is received into any suitable receptacle on which the patient sits.

For patients in bed its use is equally easy. The nurse or attendant should be instructed to make the patient lie on her back, her hips being well raised with a pillow. The pillow itself should be covered with a




FIG. 106.

VAGINAL DOUCHE.




FIG. 107.

SYPHON DOUCHE.

waterproof or folded blanket. An ordinary basin is then slipped below the hips to receive the overflow.

Instead of the douche, a single tube working by syphon action may be employed. This consists of a "sinker," a long piece of gutta percha tubing with a bent piece of glass tubing inserted so as to render it rigid where

it passes over the edge of the vessel containing the fluid, and a terminal vaginal tube. The "sinker" should be large and hollow, so that when inverted it may serve as a cup by which the tube may be filled with water; once filled, the tube is temporarily compressed while the sinker is being dropped into the jug or pail full of water ready for use. The great advantage of the douche is its simplicity.

The material for injection is various. Hot water at a temperature of 110°-120° F. is invaluable in inflammatory conditions. Medicated
Injections.

Hot carbolic lotion (equal parts of hot water and 1-20 lotion so as to give a temperature of 110°-120° F.) is admirable for cleansing purposes in abortion cases.

In so-called leucorrhœal conditions, the discharge is not vaginal as a rule, except in old women, but comes from the uterus. In such the vaginal douche merely cleanses the vagina without diminishing the discharge. Boracic acid is an excellent mild antiseptic in the proportion of ʒi to the pint. The crystals should be used, not the fine powder. Alum, sulphate of copper, or sulphate of zinc (ʒss. to Ojj.) may be



FIG. 108.

FRITSCH'S CATHETER FOR WASHING OUT THE INTERIOR OF THE UTERUS.

substituted for boracic acid, if a local astringent action is thought necessary.

It is a good plan to make the patient first douche with hot water in the dorsal posture, with a slipper bed pan or bed bath below the hips, and end with the special lotion. After it is finished the dorsal posture should be maintained for ten minutes, and the last of the injection expelled by sitting up.

The *Uterine Douche* is to be employed only after the cervical canal and uterine cavity have been so far dilated as to admit the index finger. Uterine
Douche. An ordinary vaginal douche or Higginson syringe may be employed. In giving a uterine douche in the removal of abortion or fibroid polypus, the external genitals should be cleansed with soap and carbolic lotion, the hair shaved if necessary, and the vagina thoroughly douched. Care must be taken to give the uterine douche gently and slowly, allowing free exit of the fluid, and carefully excluding air from the apparatus. The size of the uterine tube should never be such as to fill the cervical canal, and the douche should be held not much above the level of the patient's hips. The best uterine tube is Fritsch's (fig. 108), or some of its modifications, as the double canula entirely obviates any retention of fluid. Passage of the fluid through a patent Fallopian

tube into the peritoneal cavity is one of the risks, but can usually be avoided by giving the injection gently.

The uterine douche is used once only, immediately after the operation, unless septic symptoms arise. In the after treatment, the vaginal douche is sufficient.

CAUTERY.

Cautery—
Paquelin's.

The ordinary cautery may be employed in the treatment of the pedicle in ovariectomy. Details are postponed till that subject is considered.




FIG. 109.

VARIOUS FORMS OF PAQUELIN'S CONES. *A* rectangular; *B* curved; *C* straight.

In the well-known Paquelin's cautery, the vapour of benzoline is pumped through a slender, hollow cone of platinum, which has been previously heated in a gas flame or spirit lamp. It speedily becomes red or white hot by the combustion of the vapour, and can then be used.

Note as to its use: (1) To be careful with the benzoline as it is exceedingly inflammable; (2) To heat the platinum cone first (in outermost zone of the flame) before pumping in the benzoline. If the vapour is pumped in before the platinum is hot enough to ignite it, the cone is cooled by its cold stream.

The cautery should be used at a dull heat. When white hot it causes bleeding, because it thoroughly burns the tissues and thus leaves no char to act as a hæmostatic.

When used to cauterize the cervix, care is necessary that the hot metal rod does not touch the vaginal walls. Various plans have been tried to prevent this accident. Thus the rod may be covered except at its terminal two inches with a wooden case which must not touch the metal. Fig. 109 shows some of the various rods of Paquelin's cautery.

ANÆSTHETICS.

LITERATURE. *Brunton, T. L.*—Remarks on one of the causes of Death during the Extraction of Teeth under Chloroform: *Br. Med. J.*, II., 1875, p. 395. *Chiene*—Chloroform: *London Practitioner*, January 1877. *Hart, D. B.*—On Death from Insufficient administration of Chloroform: *Ed. Med. J.*, 1879. *Lancet*—The *Lancet* and the Hyderabad Commissions on Chloroform: *Lancet Offices*, London, 1896. *Lister*—Chloroform: *Holmes' System of Surgery*, Vol. V. *Report of Br. Med. Ass. Committee*: *Br. Med. J.*, Vol. I., 1879. *Murray, R. Milne*—The Cessation of Respiration under Chloroform and its Restoration by a new method: *Edin. Med. J.*, 1885.

THE chief anæsthetics are chloroform and ether. Other agents or mix-
tures have been tried—viz., ethidene; mixtures of alcohol, ether, and
chloroform: nitrous oxide; bichloride of methylene: the results have
not been satisfactory with these. In the British Medical Association's
Report on the action of anæsthetics, ethidene is strongly recommended.
Chloroform and ether, however, still remain our most trustworthy agents.

Action of Chloroform.—Chloroform when administered to a patient
has a perfectly definite effect on the nervous system. Sensation is first
abolished, and then reflex action. This is all the effect wished for in
any case. If, however, the chloroform be pushed further, the respiratory
centre becomes paralysed so that breathing ceases; and finally the heart
stops from paralysis of its ganglia. In almost all cases this is the
sequence in the susceptibility to chloroform of the parts of the nervous
system regulating sensation, reflex action, respiration, and the circulation.
Rarely have we the heart affected before the respiratory centre. When
first administered, it causes a transient rise in the blood pressure; and
then a gradual irregular fall. The more recent investigators on this

Anæsthe-
tics.

Action of
Chloro-
form.

point (see the British Medical Report) found that in dogs, chloroform reduced the blood pressure more rapidly and to a greater extent than ethidene, and that ether did not cause any appreciable depression. As the blood pressure is the resultant of the force and frequency of the heart's action and the state of dilatation of the small blood-vessels, it is evident that chloroform when administered to dogs slowed the heart and weakened the vasomotor centre more than ethidene or ether. It should be kept in mind, however, that dogs are very susceptible to the action of chloroform and easily killed by it.

Death
not always
due to
overdose.

It is wrong to suppose that in every death under chloroform the fatal result is caused by an overdose, or by the action of the drug on a fatty heart. This is a very common view, but an exceedingly erroneous one.

To prevent the patient's feeling, though one of the most gratifying results in anæsthesia, is not by any means *the* great object in operative cases. One of the most essential aims of its administration is to prevent the reflex transmission of powerful nervous impulses from the part operated on to the heart, or their direct transmission to the respiratory or vasomotor centres. If chloroform be administered to a limited extent so that sensation alone is abolished, and any large nervous trunk like the Fifth, or large nervous area like the splanchnic, be irritated, then we may have reflex inhibition of the heart or paralysis of the vasomotor and respiratory centres; in man, death may result. There are reliable clinical reports that this reflex inhibition of the heart has caused its stoppage in man. It is sometimes urged against this that no amount of stimulation of the lower end of the cut vagus in a rabbit can permanently stop its heart; in man, however, the conditions are not the same as in the rabbit. Goltz, quoted by Lauder Brunton, gives some most interesting facts in this connection. A frog was decapitated, its heart exposed, and the animal hung with its legs downwards. On tapping the intestines pretty hard, the heart stopped through reflex inhibition of the vagus but soon resumed again. It contracted vigorously but had no blood in it to propel. The irritation of the splanchnics had not only inhibited the heart but so lowered the tone of the vasomotor centre that the veins of the abdominal cavity were widely dilated; and thus the blood, when the animal was vertical, did not reach the opening of the inferior vena cava into the right auricle. When the frog was laid on its back, however, the blood flowed at once to the heart.

This then gives us the proper view of the administration of chloroform in all cases where cutting operations or operations involving large nervous trunks are being performed: *the chloroform must be pushed until sensation and reflex action are abolished, and this state is to be kept up during the operation.*

Uses.

Uses.—Chloroform is used in all cutting operations except very slight

ones: where the straining of the patient prevents the manipulation necessary for accurate diagnosis and treatment; in phantom tumours; and also, when necessary, in cases where vaginal examination of virgins is indicated.

Method of administration.—The patient should have no food for three or four hours prior to the operation. Just before the administration of chloroform is begun, half an ounce of brandy may be given. Method of Administration.

The patient lies on the back with all fastenings unloosed, and




FIG. 110.

CHLOROFORM DROP CORK.

should not sit up. A towel or napkin folded square is taken and chloroform poured on it. Fig. 110 shows a convenient and economical drop-cork which can be fitted into any bottle. A very simple plan, however, is to place a safety pin between the cork and the edge of the bottle mouth. The amount does not matter. We judge of the amount of chloroform required not by the quantity poured on the cloth but by the effect on the patient. If reflex action be not abolished, even though a quart has been used, the patient has not had enough; while if respiration be affected after a few whiffs, she has had too much.

The face of the patient should look to the side, and the chin should be kept well away from the sternum. The administrator keeps the chin forward with his right hand. This has the additional advantage of allowing him to feel the puff of the breath on the palm.

The cloth is to be held not too closely over the face and the patient directed to take long breaths.

The administrator has to keep two points before him. He is to watch the breathing most narrowly, and to ascertain when reflex action is abolished.

He can watch the breathing well by feeling the puff of the breath constantly on his hand. The abolition of reflex action is generally tested by touching the conjunctiva; when the patient is not fully under, the orbicularis contracts. This is not a perfect test, but the best we have.

When reflex action is abolished, no more chloroform is to be given; should it show signs of returning, fresh chloroform is put on the cloth.

DANGERS.

These are the following:—

(1.) *Asphyxia*;

(2.) *Reflex inhibition of heart or respiratory or vasomotor centres.*

Dangers.

(1.) *Asphyxia*.—This may arise early from fainting, muscular relaxation allowing the tongue to fall back on the pharynx; or from closure of the glottis, owing to paralysis of its intrinsic muscles. The marked extension of the head already insisted on prevents the second from happening. If it arise, the tongue is to be pulled well forward with a pair of forceps. Foulis recommends that the tongue be pressed forward by a spatula or spoon applied at its root.

When asphyxia arises from paralysis of the respiratory centre owing to an overdose of chloroform, the treatment is immediate stoppage of the administration of the chloroform *and artificial respiration by Sylvester's or Howard's method for hours if necessary*. The head should be kept hanging over the edge of the table, so as to send blood to the respiratory centre; or the patient may be inverted (Nelatonized). Nitrite of amyl is useful in counteracting cerebral anæmia, and hypodermics of ether and strychnia are also valuable.

Reflex Inhibition.

(2.) *Reflex inhibition of the heart or respiratory or vasomotor centres*.—This can only happen when there has not been given sufficient chloroform to abolish reflex action. It is by no means an uncommon thing, therefore, for the patient to die because sufficient chloroform has not been administered; sensation alone had been abolished when the operation began. The usual account is that "the patient gave a start when the first incision was made, and died." In some cases this has happened

after only a teaspoonful had been poured on the cloth. Yet this is often called "a death from chloroform."

Contra-indications.—Every patient on whom an operation is to be performed may have chloroform; if the operation is indicated, so is chloroform. If the patient has a weak heart, then chloroform is imperative for any major operation; it must be given till reflex action is abolished, as reflex inhibition of the heart is specially dangerous here. Contra-indications.

Occasionally, chloroform causes severe vomiting after the operation. For this reason Keith always uses ether. Vomiting during the operation is dangerous only when any solid matter regurgitates back into the trachea; tracheotomy may then be necessary. Vomiting.

Sickness after the operation is treated by the application of a mustard leaf to the pit of the stomach and abstinence from food. Washing out the stomach is advocated by some in bad cases.

COCAINE, introduced by Koller as a local anæsthetic, is coming to be much used in gynecology, especially in the removal of urethral caruncles, Emmet's operation, ligature of piles, and plastic operations on the perineum. A solution of the hydrochlorate (4–20 p. c.) is the one usually employed. It may be used hypodermically even in an abdominal section, but is only useful when one operates in a case *in extremis*. We can then inject in the middle line 40 minims of a 3–5 p. c. solution, and also freeze the skin with the spray from a cylinder of ethyl chloride.

Despite all criticisms and attempted improvements, chloroform administered as directed here, *i.e.*, by the method long ago advocated by Sir James Simpson, Syme, and Lister, holds its ground as the best anæsthetic yet discovered.

CHAPTER XV.

ANTISEPSIS: ASEPSIS.

LITERATURE.

- Barbour*—Puerperal Septicæmia : Edin. Med. Journ., Nov. 1885 ; Discussion, Feb. 1886.
- Bockhart*—Beitrag zur Ätiologie und Pathologie des Harnröhrentrippers : Vierteljahrschrift f. Derm. 1883, Heft 1. *Brieger*—Untersuchungen über Ptomaine : III., Berlin, 1886. *Brunner*—Ueber Catgutinfektion, Beiträge zur Klin. Chir., Bd. VI., S. 98, p. 190. *Brunton, T. L.*—Pharmacology, Therapeutics, etc. ; London, 1885.
- Buist, J. B.*—Vaccinia and Variola : London, Churchill, 1887. In investigating the above conditions, Buist, of Edinburgh, found *Staphylococcus albus* and *aureus* present in vaccine lymph. These may have been derived from the skin. *Bumm*—Beitrag zur Kenntniss der Gonorrhoe der weiblichen Genitalien : Arch. f. Gynäk., Bd. XXIII. S. 327. *Cheyne, W. W.*—Antiseptic Surgery : London, 1882. Manual of the Antiseptic Treatment of Wounds : London, 1885. *Cheyne, W. W.*—Suppurative and Septic Diseases : Pentland, Edin. and Lond., 1889. *Davaine*—Recherches relative à l'action des substances antiseptiques sur le virus de la septicémie : Com. Rend. Soc. de Biolog., Tom. I., 6th Series, p. 25. *De Ruyter*—Zur Iodoformfrage : Langenbeck's Archiv, 1887, S. 984. *Döderlein*—Das Scheidensekret und seine Bedeutung für das Puerperalfieber : Leipzig, Besold, 1893. *Fehleisen*—Die Ätiologie des Erysipel : Berlin, 1883. *Geppert*—Zur Lehre von den Antiseptics : Berl., klin. Woch., Bd. 28, 1889, pp. 789 and 919. *Klein*—Micro-organisms and Diseases : Lond. 1884. *Koch*—Article "Ueber Desinfection" in Mittheilungen aus dem kaiserlichen Gesundheitsamte : Berlin, 1881. *La Place*—Säure Sublimatlösung als desinficirendes Mittel und ihre Verwendung in Verbandstoffen : Deutsch. Med. Woch., Leipz., 1887, XIII., 866. *Lister, Sir Joseph*—On Corrosive Sublimate as a Surgical Dressing : Lancet, 1884, p. 223. Also an address on the Antiseptic management of wounds : Brit. Med. Jour., 1893. *Lomer*—Our present Knowledge of the Relations between Micro-organisms and Puerperal Fever : Am. Journ. of Obstet., Vol. XVII., p. 673. *Lusk*—Nature, Origin, and Prevention of Puerperal Fever : Intern. Congr. Tr., 1876. Also article on "Puerperal Fever" in Pepper's Practical Medicine, 1885. *Neisser*—Die Mikrokokken der Gonorrhoe : Deutsch. Med. Woch., 1882, No. 20, S. 279. *Petit*—Sur les Substances anti-fermentesciles : Comptes Rendus de l'Academie des Sciences, 1872. *Rosenbach*—Micro-organismen bei den Wund-infections Krankheiten des Menschen : Wiesbaden, 1884. *Russell, J. A.*—Article "Disinfection:" Quain's Dict. of Medicine, 1883. *Schimmelbusch*—The Aseptic Treatment of Wounds : Lond., Lewis, 1884. *Stadfeldt*—Sind als Desinficiens in der Geburtshülfe Sublimatlösungen der Karbolsäure vorzuziehen : Centralbl. f. Gyn. 1884, No. 7. *Tarnier*—Contribution to Discussion on Spiegelberg's paper on "Antisepsis in Geburtshülfe:" Transactions of London Intern. Congress, Vol. IV., 1881, p. 390. *Woodhead, G. S.*—Notes on the use of Mercuric Salts in Solution as Antiseptic Surgical Lotions : Proc. of Roy. Soc. Edinr., 1888, v. also Reports of Laboratory, R.C.P.E., 1888. *Woodhead and Hare*—Pathological Mycology : Edinr., 1885. *Ziegler*—Pathologische Anatomie, Bd. II. S. 805. The U.S.A. Catalogue gives full references under Antiseptics and Disinfection.

ANTISEPSIS.

By an antiseptic, we understand, strictly speaking, an agent capable of destroying or inhibiting the growth, or neutralising the products of micro-organisms causing sepsis, such as the micrococci known as streptococcus pyogenes, and staphylococcus pyogenes aureus. Antiseptics of course can also destroy any form of micro-organism, and the term is thus used in this more extended sense.

Lister's standpoint was that septic infection of wounds was due to the introduction of "germs" from the wound-surroundings, *viz.*, from the air, the instruments used in operating, and the hands of the operator. The spray of carbolic lotion (1-40) was used to render the air pure; and for instruments and hands, carbolic lotion (1-40 and 1-20) was employed. Lister's main ideas have in all respects been confirmed, except that air infection was exaggerated. The present standpoint is that wound infection is due to micro-organisms introduced to an inappreciable extent from the air, but mainly by unclean touch from hands and instruments, and that these require a more thorough cleansing than was at first believed.

Formerly the antiseptic properties of any substance was considered sufficient if it kept a wound free from fœtor and caused no blackening of the green silk protective placed over the wound. Owing, however, to increased knowledge as to the nature of micro-organisms arrived at by improved methods of isolation and cultivation on gelatine or peptonised jellies, more exact information has been gained as to the trustworthiness of our many antiseptic agents.

The most elaborate and exact researches have been made by Koch, and his results have been found to agree with subsequent clinical trial.

Koch's method was as follows: he dipped sterilised threads in cultivations of bacilli not containing spores, and others in those containing spores; the former were then immersed in a solution of carbolic acid (1 p. c.) for two minutes, and thereafter placed on some of the materials used for cultivation, and he found they did not grow; the latter (*i.e.*, those with spore-bearing bacilli) were however unaffected after being steeped even for two days in a 2 p. c. solution of carbolic acid. Immersion in even a 5 p. c. aqueous solution of carbolic acid did not render the spores incapable of development. 5 p. c. solutions in alcohol and in oil were ineffective on the spores even after 70 to 110 days' immersion; similar solutions destroyed the bacilli after six days immersion.

The most powerful germicide was found to be corrosive sublimate, which in weak solutions (1 in 20,000) killed spore-bearing bacilli almost immediately, and inhibited their growth when of a strength of only 1 in 30,000. An evident difference exists between micro-organisms in

relation to their resistance to antiseptics: bacilli without spores, and micrococci, are readily killed by a 1-20 aqueous solution of carbolic acid, while spores resist immersion in 1-20 carbolic lotion even for days.

Carbolic oil and alcoholic solutions of carbolic acid have proved inefficient as antiseptics, and should therefore be discarded in practice.

These researches give a guide in determining what antiseptics we should use, but require, as we shall see, to be accepted with some modification.

Activity of
various
Antiseptics.

The following is taken from a table given by Koch of the activity of various antiseptics. The double underlining means that after that number of days the spores of the bacillus anthracis were taken out of the fluid and found to be no longer capable of development. When the numeral is not so underlined it means that after immersion for the special number of days the spores were still capable of growth.

FLUID.	PERIOD (in days) OF THE IMMERSION OF THE SPORES IN THE FLUID.	REMARKS.
Absolute alcohol	1 3 5 . . . 110	
Æther	1 5 8* 30	*Incomplete growth.
Oil of Turpentine	1* <u>5</u> <u>10</u>	*Isolated but well-marked development.
Chlorine water	<u>1</u> <u>5</u>	
Bromine (2 % in water)	<u>1</u> <u>5</u>	
Iodine water	<u>1</u>	
Iron chloride	2* <u>6</u>	*Delayed but well developed.
Sublimate (1 % in water)	<u>1</u> <u>2</u>	
Thymol (5 % in alcohol)	1 6 10 15	
Salicylic acid (5 % in alcohol)	1 6 10 15	

In regard to thymol and salicylic acid it should be noted that alcoholic solutions were used, which, like oily solutions of antiseptics, are less effective than aqueous ones: *e.g.*, an alcoholic is less active than an aqueous solution of iodine. Koch's results have, however, been criticised in the following way. It has been shown by Geppert that if the thread with bacilli or spores were dipped in corrosive sublimate and then in ammonium sulphide so as to precipitate and render inert the

corrosive sublimate on the thread, the organisms grew when transferred to gelatine. This tends to show that the micro-organisms were not killed by the antiseptic, but that their growth was inhibited by the corrosive sublimate present in the thread at its time of transfer. It also shows, however, that this dipping in certain antiseptics is reliable through the slight amount of antiseptic retained—an amount not prejudicial at all to the tissues.

We must now consider our chief antiseptics from the clinical standpoint.

Carbolic acid is in many respects one of our most trustworthy anti-septics. A watery solution of 1 in 20 is thoroughly effective except in the case of spore-bearing bacilli, and can be relied on in operative work. From its not acting on metals and having no injurious action on sponges, it is useful for cleaning these as well as for skin cleansing. A solution of 1 in 20 if prolonged in its use has, however, a disagreeable action on the skin and the odour is pronounced. It can be used with soaps. Carbolic Acid.

Corrosive sublimate was recommended in 1874 by Davaine, used by Tarnier in obstetrics prior to 1880, and was very many years ago the favourite antiseptic of the late A. B. Stirling, assistant-curator in the Edinburgh Anatomical Museum, so well known for his freezing-microtome and microscopic work. Since Koch found it the only germicide for the spores of bacillus anthracis, it has come into great prominence. Corrosive Sublimate.

Solutions of 1 in 2000, 1 in 4000, 1 in 8000 are very effective; it is undoubtedly a valuable addition to antiseptics, as it is rapid in action, very soluble, odourless, and non-irritating to the hands. Its corrosive action on instruments and injury to sponges are the drawbacks to its use.

Some important facts as to the action of corrosive sublimate on soaps and blood albumin must be kept in mind. With ordinary soaps, albumin, or blood, we get insoluble and inert compounds formed. Thus if 5 c.c. blood be added to 50 c.c. corrosive sublimate (1-1000), nearly all the mercury is thrown down as albuminate of mercury. This precipitation of the mercury is prevented however by the addition of tartaric acid or common salt, so that $\frac{3}{4}$ p. c. to 1 p. c. salt solution should be used in making 1 to 1000 corrosive sublimate (Woodhead).

Messrs Duncan, Flockhart & Co. have made a special bottle (containing five ounces) with a cupped glass stopper of one drachm capacity. The solution of corrosive sublimate is of such a strength that one cup added to four tumblers of water (one quart) gives a solution 1 in 2000. This strong solution contains $5\frac{3}{4}$ grains of corrosive sublimate and 3 grains common salt to a drachm of water.

It may be ordered thus :

R. Lotionis Hydrargyri Perchloridi. $\bar{3}$ v.
 (5 $\frac{3}{4}$ grs. Hydrargyri Perchloridum, and 3 grs.
 Sodii Chloridum to 1 drachm of water).

To be put in a special bottle with cupped stopper.

Sig. Poison : for external use.

Biniiodide
of mercury.

Biniiodide of mercury is also very effective, and is believed to be better than corrosive sublimate, as it is doubly effective, and does not form insoluble compounds nor corrode metals much. These antiseptics can also be had as compressed pellets made up with tartaric acid in the case of the corrosive pellets. These are useful for the practitioner, and prevent mistakes on the part of nurses. Tartaric acid should not be added to the strong solutions of corrosive as it converts the latter into calomel in about a fortnight (Dott).

Iodoform, although it has little action on micro-organisms, seems to neutralise their products, and as a powder or gauze is most valuable in dressing wounds.

ASEPSIS.

Asepsis.

By this we mean a condition where no septic causes are present, this result being due to sterilisation by boiling water or moist steam. This condition can be brought about only in instruments and dressings, for evident reasons, and it is a decided advance in operative work from its simplicity and efficiency. The methods we discuss presently, but we may say that for operative work, one has to rely on a combination of antiseptics and sterilisation—antiseptics for the operator's hands and patient's skin, and sterilisation for instruments, wound-dressings, and the operator's and assistant's operating tunic, and for towels, etc., used.

The only difference between antisepsis and asepsis is in the cause. Antisepsis is asepsis aimed at by the use of antiseptics. Asepsis is brought about by boiling or steaming, and really attained in this way. It is doubtful if we can attain asepsis by the use of antiseptics.

We have therefore to discuss the question of how we can render free from contaminating material all that is within the operating field. This comprises the consideration of (1) the operating room ; (2) the purification of the operator's hands, and those of his assistants and nurses ; (3) the purification of the skin of the part to be operated on ; (4) the purification and cleansing of towels, tampons, operating tunics, and dressings ; (5) the preparation of sponges and ligatures ; (6) the sterilisation of instruments. It is of importance to keep in mind that our antiseptic and aseptic treatment need only be directed against certain micro-organisms, viz., those found normally in skins, and certain pathogenic organisms occasionally present. As a type of the former we may

mention a micrococcus, the staphylococcus epidermidis albus described by Welch as normally present in skin. In the vagina a bacillus has been discovered by Döderlein, which he believes to have an acid secretion, and to act as a protective against micrococcal infection.

Pathogenic organisms may be present also on and in the deeper layers of the skin—usually, as in the vagina, some form of streptococcus or staphylococcus.

All these forms are readily killed by ordinary antiseptic precautions, and it is not necessary to use such stringent antiseptics as, for instance, is required to destroy spore-bearing bacilli like the bacillus tuberculosis.

Various micrococci may be found in the lower segment of the genital tract, but the vagina, uterine cavity, and Fallopian tubes, when healthy, are practically aseptic. Organisms may of course be introduced by careless examinations with fingers or instruments not previously cleansed.

(1.) *The operating room.*—In a hospital the operating room should be so contrived that the floor is of an impermeable material, that the wall and roof are covered with a washable material, and that the corners are rounded so as not to lodge dust. The floor is therefore covered with impermeable cement, the walls with enamel varnish or some washable substance. An abundant supply of hot water is arranged for, with suitable hand-basins, with taps worked by the feet, and also apparatus for providing boiled water. There should be suitable tables for instruments, either with glass or metal tops, and on large castors so that they can be moved by the operator's foot. *The arrangements should not be too elaborate, and we must remember that in private practice great simplicity of appliance is requisite. There the careful scrubbing of an ordinary wooden table, and its protection with sterilised sheets: the surrounding of the operating field with clean towels wrung out of hot carbolic lotion, or boiled in soda solution has often to suffice, and gives excellent results. The field of mischief is practically bounded by the operator's hands.

(2.) *The disinfection of the operator's hands and those of his assistants and nurses.*—Thorough cleansing of the hands is one of the most important parts of antiseptic treatment. It is now recognised that the mere dipping of the hands in an antiseptic lotion is insufficient, and that more thorough precautions are necessary. There can be little doubt, however, that extravagant statements are made on this point to the extent that practically no means of cleansing the hands can rid them of all micro-organisms. Various micrococci may be found normally in the skin and sweat glands, usually innocent organisms; and the operator may of course in touching putrid wounds infect his skin with pathogenic micrococci. It is of great importance, therefore, to make the cleansing of the hands a very thorough procedure. They should be first thoroughly

The operating room.

Disinfection of hands.

scrubbed with soap and hot water by means of a sterilised nail-brush that is preserved in 1-2000 sublimate lotion. Nails should be short, and special care must be taken with the subungual spaces. To free the surface from greasy matter, turpentine or lysol is very useful, and finally a thorough friction with sublimate lotion (1-2000) for five minutes completes the process. Other methods may be employed, viz. (1) use of hot water and soap, followed by (2) immersion in alcohol, and (3) scrubbing with carbolic lotion (1-50). Immersion of the hands in saturated permanganate of potash after the use of soap and water, and the discharge of the deep staining with saturated oxalic acid solution,




FIG. 111.

DIAGRAM OF STERILISER TO CONTAIN SCHIMMELBUSCH'S BOX. The arrows show the direction of the steam. (*Sänger.*)

followed by sterilised lime-water, sterilised water, and finally immersion in sublimate lotion (1-500) for five minutes, has been also recommended as very thorough (Robb).

When the operator and his assistants have thus cleansed the hands and arms prior to operation, they must not accidentally in forgetfulness soil them by touching unclean articles.

The same care must be taken with the hands of the assistants and nurses, and it is advisable that all engaged within the operating field should be provided with tunics or blouses sterilised in a way to be described afterwards.

(3.) Great care must be taken to disinfect *the part to be operated on*. Disinfection of patient.

The abdominal surface in laparotomies should be shaved and cleansed with turpentine and soap and water and carbolic lotion (1-40) the night before the operation. Special care should be taken with the navel. The operator should himself cleanse the patient's skin before operation with soap and carbolic lotion (1-40), finally drying it with a sterilised towel.

For the perineal region one uses the same precautions, viz., a sitz-bath the previous evening, shaving the genitals, washing them with turpentine, soap and water, finally scrubbing with carbolic lotion (1-30) before operation. The vagina should be well brushed with a handled sterilised brush




FIG. 112.

SCHIMMELBUSCH'S BOX. It can be filled with tampons, operating tunics, and placed in Steriliser. (Sänger.)

and carbolic lotion. Special directions for cervical and other operations will be given under the separate operations.

(4.) *The purification and cleansing of tampons, towels, operating tunics, and dressings.*—For this purpose it is now held rightly that sterilisation with steam is best. Dry heat at a temperature of 120°-140° C. has been used, but it is more difficult to carry out, and, owing to its want of penetration, not so effective. Purification of dressings.

For sterilisation by steam we require a special apparatus, and of these many varieties are made. The operator may use a simple one such as that of Schimmelbusch, but more elaborate and fixed ones are employed for large hospitals (fig. 111).

In these sterilisers the various articles to be enumerated presently are placed, and subjected to the action of nascent steam for half-an-hour. The air is thus driven out, and the contents of the steriliser thoroughly

impregnated with vapour, and all micro-organisms completely destroyed. The steam should enter from above, and the dressing be warmed before it is passed in. The dressings do not require to be dried after the sterilisation is over.

The articles that can be so sterilised are gauze, gauze tampons, operating tunics, sheets, and towels.

Gauze dressings and tampons are suitably prepared and placed in dressing boxes. These dressing boxes are provided with lids, and have perforations that can be open or shut, so that when filled with gauze the holes can be uncovered and the steam penetrates. When thoroughly sterilised the box is closed, and can then be kept until required for use (*v.* fig. 112). These boxes are of great use, and provide a ready stock of dressings for private and country work.




FIG. 113.

SCHIMMELBUSCH'S STERILISER FOR INSTRUMENTS. In this they can be boiled in a 1 per cent. soda solution.

If necessary, iodoform can be sprinkled on the gauze moistened with sterilised water, and pressed into the dressing with a cleansed rod or glass presser. Gauze tampons can be made of gauze sewn round pads of salicylic wool. They are useful instead of sponges for perineal operations, and by some operators are used instead of sponges in abdominal section.

It will be readily understood that operating tunics, towels, and sheets can be sterilised in the same manner.

(5.) *The preparation of sponges and ligatures.*—Sponges after operation should be thoroughly washed with abundant water, then soaked in very hot soda solution (1 per cent.) for half-an-hour, but they are not to be boiled, as this destroys them. They are then rinsed in boiled water, soaked in carbolic lotion (1–20) for twenty-four hours, and either kept permanently in this, or dried and soaked in carbolic lotion (1–20) prior to operation.

Sponges
and liga-
tures.

New sponges require the sand removed from them. They should be beaten well, soaked in hydrochloric acid and water, sufficiently bitter to be unpleasant to the taste, and then treated as above. The greatest care requires to be exercised in regard to sponges. Good sponges are expensive, but with care will last for some time.

Silk ligatures should be boiled in soda solution (1 per cent.) along with the instruments, and are thus rendered aseptic completely. *Catgut* is the submucous coat of the sheep's intestine. Chromic acid and juniper gut are the chief forms employed, and are sold ready for use. Chromic acid gut is dry, while juniper gut is usually preserved in alcohol, after being soaked in sublimate and alcohol (sublimate 10,




FIG. 114.

Arrangement for operation on patient in Lithotomy posture (Sänger). *a*, Limbs in leg-holder (*h*) and covered with sterilised stockings; *b*, sterilised gauze pad; *c* is air-pad with gauze napkin, *d*, and flap, *e*; *f* is bath for overflow; *i*, chloroform mask; *k*, movable glass shelf; *l*, table for instrument clerk; *m* is a lower table within reach of the operator; *n*, box for sterilised tampons; *o* has sterilised wool; *p*, dish for sublimate lotion, and *q* forceps.

absolute alcohol 800, distilled water 200), thrice renewed, for forty-eight hours. Glycerine added to the alcohol (20 per cent.) makes it suppler (Schimmelbusch). Prior to use, catgut should be soaked in carbolic lotion. When used in needles these should be threaded prior to soaking, as it swells. The perfect disinfection of catgut is a difficult matter, and anthrax has arisen from the use of imperfectly sterilised specimens. A method recently recommended is to expose the catgut to dry heat at a temperature of 80 degrees C., and then to boil it in cumol, a hydrocarbon with a high boiling point (145° C.).

It is then transferred to sterilised tubes. Wire and silk-worm gut can be disinfected by soaking in carbolic lotion.

Instru-
ments.

(6.) *The Sterilisation of Instruments.*—Sterilisation of instruments can be most effectively carried out by boiling them in a simple apparatus filled with soda solution.

The steriliser for instruments (fig. 113) is a box made of tin or copper, with a tightly fitting lid and four legs, removable. Heat is supplied by a spirit lamp. The instruments are placed on a perforated tray with a handle at each end, and this is put into the steriliser with soda solution. The lid should be firmly closed, the lamp lighted, and the instruments




FIG. 115.

PAD (Price, Kelly), modified by Säger. Kelly's pad is seen on the Trendelenburg table.

in this way boiled for thirty minutes. They are then thoroughly aseptic, and are ready for operation when cooled either by adding boiled water or 1-20 carbolic lotion to the steriliser. During the operation they can remain in the steriliser or be placed in shallow trays in 1-40 carbolic lotion.

The operator's aim is, therefore, to have hands thoroughly cleansed by antisepsis, to operate on a thoroughly pure surface, to have such a disinfection by sterilisation of operating tunics, of the necessary towels and sheets surrounding the patient, and of the instruments, sponges, ligatures, etc., that everything touching the wound or touched by the operator, is free from septic causes, *i.e.*, carries no micro-organismal contamination. His purpose is finally completed by the application of dry

sterilised dressing impregnated, if necessary, with iodoform, so that no septic organism can contaminate the wound secretions.

This requires great vigilance and care on the part of the operator, careful drilling of assistants, and a thorough knowledge of bacteriological results, all tempered with common sense.

Fig. 114 shows a patient arranged for such an operation as Hysterec-tomy or one of the perineal operations. A convenient pad for operations is seen at fig. 115 (*Price, Kelly*).

1871

1. The first of the year was a very cold day, with a heavy frost, and the wind from the north.

2. The second day was a very warm day, with a heavy rain, and the wind from the south.

3. The third day was a very cold day, with a heavy frost, and the wind from the north.

4. The fourth day was a very warm day, with a heavy rain, and the wind from the south.

5. The fifth day was a very cold day, with a heavy frost, and the wind from the north.

6. The sixth day was a very warm day, with a heavy rain, and the wind from the south.

7. The seventh day was a very cold day, with a heavy frost, and the wind from the north.

8. The eighth day was a very warm day, with a heavy rain, and the wind from the south.

9. The ninth day was a very cold day, with a heavy frost, and the wind from the north.

10. The tenth day was a very warm day, with a heavy rain, and the wind from the south.

PART II.

DISEASES OF THE FEMALE PELVIC ORGANS.

After considering the etiology and classification of the diseases of the female pelvic organs, we shall adopt a classification based on anatomy, grouping the diseases under the structure affected. We shall devote one section to each group of affections as follows :—

- Section III. The Peritoneum and Connective Tissue ;
- „ IV. The Fallopian Tubes and Ovaries ;
- „ V. The Uterus ;
- „ VI. The Vagina ;
- „ VII. The Vulva and the Pelvic Floor.

Further, we shall consider under special sections disturbances of the following functions :—

- Section VIII. The Menstrual function ;
- „ IX. The Reproductive function.

Finally, we shall devote one section to affections of the other pelvic organs :—

- Section X. The Bladder and the Rectum.

In an Appendix there will be considered the Classification of Gynecological Diseases ; Abdominal Section and Colpotomy ; Electricity in Gynecology ; the Systematic Treatment of Nerve Prostration ; Case-taking ; and Gynecological Literature.

PART II.

DIRECTOR OF THE NATIONAL BUREAU OF STANDARDS

A few months after the establishment of the National Bureau of Standards, the following functions were assigned to the Director:

1. To establish and maintain the national standards of length, mass, and time.

2. To establish and maintain the national standards of temperature.

3. To establish and maintain the national standards of electrical units.

4. To establish and maintain the national standards of mechanical units.

5. To establish and maintain the national standards of chemical units.

6. To establish and maintain the national standards of physical constants.

Section VII. The National Bureau of Standards.

IX. The National Bureau of Standards.

The following functions were assigned to the Director of the National Bureau of Standards:

Section VIII. The National Bureau of Standards.

In an effort to bring about a more complete and accurate knowledge of the physical constants, the following functions were assigned to the Director of the National Bureau of Standards:

1. To establish and maintain the national standards of length, mass, and time.

SECTION III.

AFFECTIONS OF PERITONEUM AND CONNECTIVE TISSUE.

- CHAPTER XVI. Etiology and Classification of Gynecological Diseases.
,, XVII. Pelvic Peritonitis and Pelvic Cellulitis (Parametritis).
,, XVIII. Pelvic Hæmatocele and Hæmatoma: New Growths
in the Pelvic Peritoneum and Connective Tissue.

CHAPTER XVI.

THE ETIOLOGY AND CLASSIFICATION OF GYNECOLOGICAL DISEASES.

LITERATURE.

Bennet, J. Henry—On Inflammation of the Uterus : London, Churchill, 1845. *Bernutz and Goupil*—Clinical Memoirs of the Diseases of Women, Vol. II. : New Sydenham Society, Meadow's translation, 1866. *Cullen*—Beschleunigtes Verfahren zur Färbung frischer Gewebe mittelst Formalins : Cent. für allgemeine Path. und path. Anat., 1895. *Hart, D. Berry*—The Pathological Classification of Diseases of Women : Edinburgh Hospital Reports, Pentland, 1894. *Headley, D. Balls*—The Evolution of the Diseases of Women : Smith, Elder & Co., London, 1894. *Hewitt*—The Diseases of Women : London, Longmans, Green & Co. *Pick*—Eine Methode der Schnellanfertigung gefärbter Dauerpräparate für die "Stückchendiagnose"—Cent. für Gyn., No. 40, 1896.

GYNECOLOGY was at first a branch of general medicine, and the existence of special pelvic disease was inferred usually from reflex pains in the left inframammary, sacral or iliac regions, and from disturbance of the menstrual functions, or excess of leucorrhea. Little progress was made until the discovery of the use of instruments of precision (sound, speculum, etc.); but even then any operative treatment was minor, so that real pathological advances were wanting until the abundant abdominal, vaginal, and uterine surgery of the past twenty years gave fresh material for investigation and research, and actual demonstration of hitherto obscure pelvic conditions. Gynecology has also shared in the valuable work of the bacteriologist, and in the advances of antiseptic surgery, so that we are now bound to reconsider some essential points, viz., the etiology of uterine diseases, and the question of how we are to classify them.

THE ETIOLOGY OF GYNECOLOGICAL DISEASES.

It is very remarkable that attempts to give a complete etiological account of Diseases of Women were made early in the history of Gynecology. Dr J. Henry Bennet, who published his Treatise on Inflammation of the Uterus in 1845, a work characterised by great

ability, attempted then and in subsequent years to found a system of uterine pathology, and asserted that in cervical inflammation and ulceration we had the conditions which formed the most common pelvic lesions in women, and that many other diseases could be distinctly traced as secondary to them. Other observers merely substituted another diseased organ for the cervix, and made some of its diseased conditions the principal ones to which many others were secondary. Thus Tilt assigned the leading place to inflammatory conditions of the ovary: Hewitt asserted that given a flexion or a version, you had conditions important in themselves and originating almost all gynecological diseases; while not a few held that uterine disorders were really due to constitutional conditions. All these attempts were more akin to the efforts of deductive philosophy, where the philosopher started with some apparently impregnable definition and explained the world and mankind in a series of elaborate deductions from it, than to scientific pathology.

These attempts, most praiseworthy at the time, and in the state of knowledge then, probably the only possible, have failed, and one must now recognise that the etiology of gynecological disease is the same as that of the diseases of other organs, only modified by structure and function.

In the Etiology of Diseases of Women, the following may be considered important factors, not sharply separated, but combining with one another:—

Factors
in the
Etiology
of Disease.

- (1.) The special anatomical relations of the pelvic organs;
- (2.) Their development and its occasional defects;
- (3.) The functions of menstruation, the sexual relations, pregnancy, labour, the puerperal state;
- (4.) Micro-organismal causes.

(1.) *The special anatomical relations of the pelvic organs.*—The most important of these are the abundant vascular supply of the pelvic tissues, its specially rich lymphatic arrangements, and above all the fact that the peritoneal cavity is not a closed sac as in the male, but that there is actual continuity of peritoneal and tubal epithelium, with that of the genital tract below. The preponderance of pathological conditions behind the broad ligament and in the pouch of Douglas, is undoubtedly to be correlated with this, and also with the fact that the ostium abdominale of the tube opens in that neighbourhood. We have thus from this continuity, and also the lymphatic distribution, access to the peritoneum for pathogenic organisms.

(2.) *Their development and its occasional defects.*—We need only mention here the development of uterus and vagina from the ducts of Müller, the early rôle of the Wolffian ducts and bodies, and the peculiar development of the ovary, and the great fact that disturbances may

arise in these, and give rise to malformations and new growths (*v.* Chap. on Ovarian Pathology).

(3.) *The functions of menstruation, the sexual relations, pregnancy, labour, and the puerperal state.*—The functions of menstruation may be disturbed by congenital malformation, and when once inflammatory conditions have been set up, we get interference with the turgescence and erection of the organs during the period, and pain more or less severe as a result.

With the sexual relations is mainly to be associated gonorrhœal infection. So far as is known, syphilis is not, apart from its own lesions, an important factor in Gynecology, although in the production of abortion it is all-powerful.

The micro-organismal nature of gonorrhœa and the anatomical relations of the parts already alluded to, give most favourable opportunities for its spread, and render its cure when present most difficult.

The actual lacerations in normal and abnormal labour, septic conditions arising from incomplete expulsion of membrane during the third stage, and the mismanagement of abortion cases by the neglect of the patient or medical attendant, form by far the most fruitful source of Gynecological trouble. It is not too much to say that careful management of labour and abortion would reduce gynecological cases by a very large fraction.

In extrauterine gestation we have a factor, recently more thoroughly recognised, in the production of blood-effusions in the tubes, connective tissues, and peritoneum.

Micro-organisms as causes.


(4.) *Micro-organismal Causes.*—This is one of the most important factors in the Etiology of Gynecology, and it is in the region of bacteriology that the most useful and permanent advances have been made. The result of bacteriological investigation has not only been of the greatest value in itself, but has completely overthrown much absurd pathology and noxious meddlesome treatment.

The following micro-organisms are recognised as causing specific diseases :—*Bacillus tuberculosis* of Koch ; the *Gonococcus* of Neisser ; *Streptococci* ; and one rare fungus—the *Actinomyces bovis* may also be noted. We do not here consider parasitic pediculi, parasites causing skin diseases, or *tæniæ* causing pelvic hydatids.

Bacillus of tubercle.

Bacillus tuberculosis.—The discovery of this micro-organism by Koch in 1882 has given great precision to our knowledge of genital tuberculosis. This bacillus is in the form of a rod with rounded ends 1·5 to 3·5 μ . in length. It probably forms spores. The special microscopic anatomy will be considered under the organs, but we may note here that by the discovery of the bacillus we now know accurately that tuberculosis occurs rarely in the vulva, more frequently in the vagina and bladder, quite frequently in the body of the uterus (although rare





MICRO-ORGANISMS AND THEIR RELATION TO THE TISSUES, v. p. 177. (DOLÉRIE AND PUCHEVIN)

I. *Staphylococcus pyogenes aureus*; II. *Staphylococcus pyogenes aureus* in the liver; III. *Streptococci* of septicemia; IV. *Streptococci* in epithelium and leucocytes; V. *Gonococci* in pus; VI. *Gonococci* enlarged and diagrammatic; VII. *Gonococci* in epithelium; VIII. *Gonococci* in epithelium; IX. *Gonococci* in epithelium.

in the cervix), most frequently of all in the tubes. In the ovary it is fairly common, and quite common in the peritoneum.

Genital tuberculosis arises by the spread from neighbouring affected tissues (viz., peritoneum, intestine) through the blood, by autoinfection, from tubercular stools, and probably by coitus.

The Gonococcus.—The discovery of the gonococcus as the cause of gonorrhœa by Neisser in 1879 has proved a most important advance. Gonococcus.

Neisser's gonococcus exists in the form of diplococci—double micrococci 1.5μ . in diameter, and so placed that their concave surfaces are close together. It grows on human blood serum between 30° and 40° C., and pure cultivations inoculated in the healthy human urethra give rise to gonorrhœa (Bumm). It is the undoubted cause of gonorrhœa.

Bumm has alleged that the gonococcus grows only on columnar epithelium, but according to Wertheim, it can also penetrate deeper, and burrow in connective tissue.

Streptococci have been found in tubal conditions, in cellulitis, and in peritonitic exudations, and thus play an important part in septic inflammatory conditions. They also form toxines. Streptococcus.

Actinomyces bovis.—This rare fungus has been found in a very few cases to be the cause of suppurative conditions in the ovary. Its structure is that of granulation tissue containing masses which stain deeply with aniline dyes, and are composed of club-shaped bodies. Actinomyces.

Plate IV., from Doléris and Pichevin, shows the most important micro-organisms, and their appearance in the tissues. I. *Staphylococcus pyogenes aureus* (Crookshank). II. Metastatic colony of *staphylococcus pyogenes aureus* in the liver (Crookshank). III. *Streptococci* of septicæmia and erysipelas (Doléris). IV. *Streptococci* in epithelium and in leucocytes (Cornil and Babes). V. *Streptococci* in the muscular coat of the uterus and in the vessels: *b*, muscle; *v*, vein; *l*, lymphatic (Widal). VI. *Streptococci* in the puerperal mucosa: *b*, chorion; *y*, muscle (Widal). VII. *Gonococci* in pus (Cornil and Babes). VIII. *Gonococci* (diagrammatic). IX. *Gonococci* in the epithelium of the conjunctiva (Cornil and Babes).

METHODS FOR DETECTING THESE MICRO-ORGANISMS.

It is of great importance that all fluids and specimens obtained clinically should be examined by the gynecologist both microscopically and bacteriologically. We give here, therefore, a summary of the methods used at the laboratory of the Royal College of Physicians, Edinburgh, for which we are indebted to Mr J. Hume Patterson, Laboratory Assistant there. Bacteriological Methods.
Examination for Staphylococcus and Streptococci.

Staphylococcus pyogenes albus, aureus, and Streptococcus pyogenes.—Receive the pus or fluid to be examined into a sterilised flask or test-

tube. With a platinum needle, which has been heated in the flame of a Bunsen burner or a spirit-lamp, lift a little of the fluid and smear it over the surface of a slightly alkaline agar tube, and, without recharging the needle, inoculate other two agar tubes in succession.

Label the tubes 1, 2, and 3, and place in an incubator at a temperature of 35° C. over night.

In the morning growths should appear in each of the tubes, but they may be so numerous in No. 1 as to make it quite impossible to isolate and examine them separately. This should be easily done with the second or third dilution.

The staphylococcus pyogenes albus appears in small round white colonies with moist surface, and raised on the surface of the agar agar.

The staphylococcus pyogenes aureus appears as small round cream-coloured colonies with moist surface, and raised on the surface of the agar. The colonies after a day or so become golden in colour.

The streptococcus pyogenes appears as small round greyish transparent looking colonies, and is easily distinguished from the staphylococcus aureus and albus.

With a sterilised platinum needle inoculate gelatine (10%) tubes, and keep at the temperature of the room; growths should appear in a few days. The staphylococcus albus and aureus liquefy the gelatine in three or four days after commencement of growth. The streptococcus *does not* liquefy gelatine.

Also make cover-glass preparations from the agar tubes, by rubbing a small quantity of the growth on a cover-glass with a drop of distilled water. To fix the film, allow to dry, and pass three times through the flame of a Bunsen burner with the back of cover glass to the flame.

Stain as follows. Put one drop of saturated watery solution of methyl green and one drop of saturated watery solution of dahlia into a watch-glass, and dilute with distilled water. Place a drop of the stain on a slide, and put the cover-glass preparation face down on the stain, and examine.

A permanent preparation may be made after examining, by washing the coverslip in water, allowing to dry, and mounting in xylol balsam.

Cover-glass preparations may also be stained in a saturated watery solution of methyl blue or gentian violet for a few minutes, washed in water, allowed to dry, and mounted in xylol balsam.

Examina-
tion for
Gonococci.

Gonococcus.—Make cover-glass preparations from the discharge, or, if a urine, centrifuge first, and prepare coverslips from the deposit.

Stain by Gram's method (alcoholic solution of gentian violet 1 part, aniline oil water 8 parts; shake well up, and filter into a watch-glass; stain films for three or four minutes: rinse very slightly in absolute alcohol, place in Gram's solution for five or ten minutes, and

decolorise in absolute alcohol). Then counter-stain with weak solution of saffranine in water for three or four minutes, wash in water, allow to dry, and mount in xylol balsam.

The gonococcus and the pus corpuscles are decolorised with Gram's solution, and are stained red with the saffranine, while the other organisms are stained violet.

The gonococci are seen in groups either between or in the pus cells. The gonococci are sometimes so few in number that it is difficult to detect them in this way, and in that case cultures should be made in the following manner. With a platinum loop-needle, which has been sterilised, rub some blood, taken under sterile precautions, over the surface of an agar tube. The blood can be obtained either by sterilising one's own finger with soap and water, corrosive sublimate (1-2000), and methylated spirit, and then pricking it with a sterilised needle or scalpel; or the ear of a rabbit may be used.

Inoculate the tubes with the pus or deposit, and place in the incubator at 35° C.

The growth appears as small round greyish transparent-looking colonies, similiar in appearance to the streptococcus pyogenes.

Coverslips are made and stained by Gram's method and saffranine.

Tubercle Bacillus.—Make cover-glass films by taking a small quantity of the sputum or centrifuged urine with a needle and placing on a cover-glass; then place a clean coverslip on the top and squeeze the two together; wipe round the edges with a piece filter-paper, which should be burned afterwards; separate the cover-slips by sliding the top one off. Pass three times through a Bunsen flame with back of cover-glass to the flame, in order to fix the film.

Examina-
tion for
Tubercle
Bacillus.

If sputum is thin, centrifuge first and use the deposit. Centrifuge also in the case of a urine.

Stain in the following manner: Filter some Ziehl-Neelsen fuchsine (carbolic acid crystals, 5 parts; aqua dest., 100 parts; absolute alcohol, 10 parts; fuchsine, 2 parts) into a watch-glass, and place coverslips face down on the stain, and warm gently for a few minutes. Wash in water.

Decolorise in 25% sulphuric acid. Wash in water.

Counter-stain in methyl blue (saturated watery solution of methyl blue with an equal part of aqua dest.). Wash in water, allow to dry, and mount in xylol balsam.

The tubercle bacilli are stained red.

Sections of tissues are treated in a similar way, but are stained for an hour in a warm solution of the stain, or twelve hours in a cold solution.

After being stained with the methyl blue, they are washed in water,

absolute alcohol, cleared up with clove oil, xylol, and mounted in xylol balsam.

Pus.—The bacilli are so few in number, that it is useless to stain.

The best method is to inject some of the pus into the peritoneal cavity of a guinea-pig, and kill it in ten days after, when the *post-mortem* will show whether tubercle is present.

Cultures of the tubercle bacillus may be made on blood serum or 6% glycerine agar agar.

Examina-
tion of
Fresh
Tissues.

Rapid Method of cutting Fresh Tissues.—Place piece tissue $1 \times .5 \times .2$ cm. for two hours in 10% watery solution of formaline. Freeze on ice microtome and cut.

After inoculation they are placed in an incubator from four to six weeks.

Place the sections for three minutes in 50% formaline solution, then three minutes in 50% alcohol, and one minute in absolute alcohol; then wash sections and stain.

The tissue may also be at once frozen and cut, and the sections treated as above.

The fresh tissue may indeed be frozen at once, the sections hardened in formaline (2%) for a few minutes, washed in water, stained in alum carmine, and then passed through water and alcohol (80%, two to three minutes), absolute alcohol (ten seconds) carbol-xylol, and balsam.

THE CLASSIFICATION OF GYNECOLOGICAL DISEASES.

All writers of gynecological text-books adopt an *anatomical* classification of the diseases considered. They either begin with peritonitis and cellulitis, taking up the diseases of the other genital organs in a regular downward anatomical sequence; or, they begin with vulvar disease, and reverse the order.

The first plan is the better one, and either indeed will be our only feasible one for many years yet. The former has lost some of its value, since the importance of tubal disease, in relation to peritonitis especially, has become known,—since, in fact, the importance of primary peritonitis has been curtailed. But all the same we cannot at present adopt any other method without confusion, and without serious gaps in the arrangement.

The anatomical arrangement has further this supreme convenience, that it groups together the diseases of the organs compactly, and is thus the only plan of arrangement suitable for the student.

Another system of classification is the *pathological*, but we defer consideration of this to the Appendix.

CHAPTER XVII.

PELVIC PERITONITIS AND PELVIC CELLULITIS (PARAMETRITIS).

LITERATURE.

- Bandl*—Die Krankheiten der Tuben, etc.: Billroth's Handbuch, Stuttgart. *Barnes*—Diseases of Women: London, 1878. *Cullingworth*—The Value of Abdominal Section in Certain Cases of Pelvic Peritonitis: Tr. of Lond. Obst. Soc., 1892. *Doherty*—On Chronic Inflammation of the Uterine Appendages after Childbirth: Dub. J. of Med. Sc., 1843. *Duncan, J. Matthews*—A Practical Treatise on Perimetritis and Parametritis: Edinburgh, A. & C. Black, 1869. On Albuminuria with Parametritis: Lond. Roy. Med. and Chir. Tr., Vol. XLVII. *Emmet, J. D.*—The Principles and Practice of Gynæcology: London, Churchill, 1879. *Flügge*—Die Miknooyanismen, etc.: Leipzig, F. C. W. Vogel, 1896. *Freund, W. A.*—Anatomische Lehrmittel, etc.: Gesellschaft f. Geburtshülfe, Berlin, Band iv., Hft. 1. Das Bindegewebe im weiblichen Becken und seine pathologische Veränderungen, etc.: Gynäkologische Klinik, Strassburg, 1885. *Freund, H. W.*—Ueber die feineren Veränderungen der Nervenapparate im Parametrium bei einfacher und parametritischer Atrophie: Cent. f. Gyn., IX. S. 644. *Heitzmann*—Die Entzündung des Beckenbauchfells beim Weibe: Wien, 1883. *Lusk*—Puerperal Fever: Internat. Cong. Trans., 1877. *MacDonald, Angus*—Three Cases of Parametritis, with observations on its Diagnosis and Treatment: Ed. Med. J., 1880, p. 1060. *Marmorek*.—Le Streptocoque et le serum antistreptococcique: Annales de l'Institut Pasteur, tome neuvième, 1895. *Noeggerath*—Latent Gonorrhœa especially with regard to its influence on Fertility in Women: Am. Gyn. Tr., Vol. I. *Olshausen*—On Puerperal Parametritis and Perimetritis: New Syd. Soc. Transl., 1876. *Priestley, W. O.*—Pelvic Cellulitis and Pelvic Peritonitis: Reynolds' System of Medicine, Vol. V. *Schroeder*—Krankheiten der weiblich, Geschlechtsorgane: Leipzig, 1879. *Schultze, B. S.*—Ueber die pathologische Anteflexion der Gebärmutter und die Parametritis posterior: Hirschwald, Berlin, 1875. *De Sinéty*—Gynécologie: Paris, 1879. *Simpson, Sir J. Y.*—On Pelvic Cellulitis and Pelvic Peritonitis, Clinical Lectures on the Diseases of Women (edited by A. R. Simpson): Edinburgh, A. & C. Black, 1872. *Simpson, A. R.*—Quarterly Report of the Royal Maternity and Simpson Memorial Hospital; Ed. Med. Journal, Vol. XXVI., p. 1059. *Spiegelberg*—Remarks upon Exudations in the Neighbourhood of the Female Genital Organs; Second Series of German Clinical Lectures, Translated by New Sydenham Society, 1877. *Tait, Lawson*—On the Treatment of Pelvic Suppuration by Abdominal Section and Drainage: Tr. of Lond. Roy. Med. and Chir. Soc., 1880, p. 307. *Tilt*—Handbook of Uterine Therapeutics: London, 1863. *Williams, John*—Serous Perimetritis; Lond. Obstet. Trans., 1885. *Winckel*—Die Pathologie der weiblichen Sexual-organe: Leipzig, 1881. This last gives photolithographs of great value. *Woodhead, G. S.*—A Lecture on the Diagnosis and Anti-toxic Serum Treatment of Diphtheria: Lancet, Dec. 15, 1894. *Ziegler*—A Text-book of Pathological Anatomy and Pathogenesis, Macalister's translation: London, Macmillan & Co., 1896.

Pelvic
Periton-
itis.

In treating of the subjects of pelvic peritonitis and pelvic cellulitis it will be convenient to take up some preliminary matter and then to consider separately each condition under the following heads:—

Nature,	Diagnosis and differential
Pathological anatomy and	diagnosis,
varieties,	Course and results,
Etiology,	Prognosis,
Symptoms,	Treatment.
Physical Signs,	

PELVIC PERITONITIS.

The student must specially note that pelvic peritonitis is a secondary condition. The old idea of it as a primary one, mainly, and as resulting from an initial congestion due to mere vascular disturbance, must be given up, and with it much of the antiphlogistic treatment that has sunk so deeply into our therapeutics. In the peritoneal area too, we must keep in mind that the pelvic peritoneum is a part where the inflammatory disturbance occurs with less virulence than in other sites, such as the peritoneum over the intestinal coils.

It is to be noted too that a peritonitis beginning in the pelvis may become general. We do not, however, take up this question at our present stage.

SYNONYMS.—Parimetritis: Pelveo-peritonitis.

NATURE.—An acute or chronic inflammatory condition affecting chiefly the pelvic peritoneum.

PATHOLOGICAL ANATOMY AND VARIETIES.

Patholo-
gical
Anatomy.

In the early stages, the peritoneum is injected and the epithelial cells dull in lustre. Soon, in marked cases, fibrinous or serous fluid is poured out: the former stiffens the peritoneum and often causes extensive adhesions between uterus and rectum, Fallopian tubes and ovary; the latter either remains free in the cavity, or becomes encysted by the false membranes already alluded to, often making Douglas' pouch to bulge down. In bad cases pus is formed. We may therefore speak

Varieties.

of simple pelvic peritonitis, adhesive pelvic peritonitis, and serous or purulent pelvic peritonitis. These, however, are mere varieties. Tubercular and malignant peritonitis will be considered by themselves.

ETIOLOGY.

Etiology.

The causes of pelvic peritonitis are numerous. They are chiefly the following.

1. The existence of pelvic cellulitis, pelvic hæmatocele, ovaritis, ovarian tumour, fibroid tumour, tubercle, or carcinoma.

2. Childbirth and abortion.
3. Gonorrhœa.
4. Latent gonorrhœa in the male.
5. A chill, especially during menstruation.
6. Venereal excess.
7. Instrumental examination by the sound, stem pessaries, sponge or tangle tents.
8. Tubal disease.

1. *The existence of pelvic cellulitis, pelvic hæmatocele, ovaritis, ovarian tumour, fibroid tumour, tubercle, or carcinoma.*

We have already noted that marked pelvic cellulitis is always associated with some pelvic peritonitis. The pelvic peritoneum and cellular tissue are adjacent and intimately connected with one another in their vascular, nervous, and especially in their lymphatic supply; we have already seen how the stomata of the peritoneum communicate with subendothelial lymphatics. In the same way we can understand a pelvic peritonitis arising secondarily from ovaritis. A hæmatocele is always followed by inflammatory changes in the peritoneum.

Ovarian tumours often set up pelvic peritonitis after being tapped as well as from their mere mechanical pressure or from torsion of their pedicle—a fact of the highest importance as regards the operation of ovariectomy. Occasionally we get general peritonitis from suppuration of a small ovarian tumour and its perforation with escape of pus into the peritoneal cavity. Small fibroids, tubercle, and cancer do the same, and thus give rise to considerable difficulty in diagnosis.

2. *Childbirth and abortion.* When an inflammatory lesion follows these, it is generally cellulosic and, as we shall afterwards see, probably septic. Pelvic peritonitis often enough follows, and is then probably likewise septic. The organisms causing septic attacks are now considered to be bacilli or cocci, and pass either by the lymphatics or blood-vessels. In mischief arising from the intestine, the bacterium coli commune is the organism.

3. *Gonorrhœa* is one great cause of peritonitis. It results from actual spread of the gonorrhœal virus. The purulent infection probably passes along the Fallopian tubes and out at the fimbriated end, setting up a severe peritonitis; or it passes by the tissues themselves, spreading along the lymphatics. In puerperal women, gonorrhœa is by no means innocent, as the following case by A. R. Simpson shows:—

“J. C., primipara, prostitute, æt. 18, was admitted to the hospital and delivered of a male child. On the afternoon following, severe peritonitis set in which proved fatal in ten days. On *post-mortem* the abdomen contained $\frac{3}{4}$ viii. of yellow pus. Surface of intestines covered with recent

fibrinous lymph becoming purulent. Mucous membrane of bladder much congested and in certain areas rough and granular. . . . On squeezing the Fallopian tubes a large quantity of pus was expelled, and the tubes appeared to be much distended with it. Mucous membrane much congested." (Report by D. J. Hamilton.)

4. *Latent gonorrhœa in the male.*—By this term Noeggerath of New York, who first directed attention to the subject, means a gonorrhœa in the male apparently cured, which sometime after—even two years—infects a healthy genital tract, causing discharge and pelvic peritonitic disturbance. The authors have seen cases bearing out Noeggerath's views, and this theory is now generally admitted.

5. *Chill, especially during menstruation.*—It is an old view that the pelvic congestion of menstruation may under undue exposure to cold pass into peritonitis; but proof is lacking.

6. *Venereal excess* in prostitutes may, for evident reasons, have peritonitis as its sequel, although exact proof of this is difficult.

7. *Instrumental manipulation.*—This is alluded to under the various instruments and needs mere mention here.

8. *Tubal disease.*—This is now recognised as an important cause of pelvic peritonitis, and has been above alluded to under *Gonorrhœa*. The facts that the genital tract communicates with the peritoneal cavity through the Fallopian tubes, and that gonorrhœa and septic diseases are due to micro-organisms, explain, in many instances, the causation of peritonitis. Tubal disease and peritonitis are mutually related, inasmuch as occlusion of the tube may be set up after the peritonitis and thus tubal distention follow. Gonorrhœal pus sets up, according to Bumm, limited peritonitis, the explanation given being that the gonococcus, its specific organism, does not flourish on squamous as it does on cylindrical epithelium. Wertheim has however shown that Bumm's view is untenable, and that the gonococcus can spread by tissue and not only superficially. The micrococci found in septic pus set up violent peritonitis when introduced into the peritoneal cavity.

Bernutz's
Analysis.

We append Bernutz's analysis of the causes of pelvic peritonitis in ninety-nine cases:—

43	occurred in puerperæ.
28	„ after gonorrhœa.
20	„ during menstruation.
8 traumatic	{ 3 due to venereal excess.
	{ 2 „ syphilitic diseases of cervix.
	{ 2 „ introduction of the uterine sound.
	{ 1 „ use of vaginal douche.

SYMPTOMS AND PHYSICAL SIGNS.

A. Acute Peritonitis.

Symptoms. Increased, full, and bounding pulse; increased temperature; rigor; shooting pains very severe. Symptoms.

Physical Signs. On palpation of lower part of abdomen the patient complains of pain; and the abdominal muscles, apart from the patient's volition resist pressure. She lies usually on the back, and with both legs drawn up. Physical Signs.

On vaginal examination the vagina feels hot and tender, and pulsating vessels may be felt in the fornices.

After exudation is present, we may feel one or other of the following conditions:—

1. A flat hard non-bulging condition of the fornices round the cervix, which is not displaced to one or other side but is immobile. The usual simile, and a very good one, is that it feels as if plaster of Paris had been poured into the pelvis.

2. An indistinct fulness high up in the pelvis. This may be from serous exudation.

3. A bulging tumour behind the uterus displacing it to the front; or a tense fluid laterally, apparently in the site of the broad ligament (fig. 38). The former is due to encysted serous effusion in the pouch of Douglas, the latter to encysted serous fluid behind the broad ligaments displacing it forwards. As a general rule these effusions are high in the pelvis and symmetrical. Sometimes the bulging retro-uterine tumour feels nodulated after a time; this is probably from extension of the inflammatory condition into the subjacent connective tissue.

Note that the bimanual is often impossible owing to the rigid condition of the fornices and abdominal muscles. The bimanual estimation of effusion is often misleading, owing to the fact that we feel the rigid peritoneal membrane through the fornices, and from the rigidity of the abdominal wall draw the conclusion that there is effusion between. Careful examination under chloroform is of great value in such cases.

B. Chronic Peritonitis.

Symptoms. These are chiefly backache, sideache, leucorrhœa, increased menstruation and sterility. Pain is the most marked symptom, and is felt most on vaginal examination or coitus. Symptoms.

Physical Signs. On vaginal examination, obscure thickening is felt in the fornices. The uterus, if displaced, is often markedly anteverted from cicatrisation of the peritoneum in the pouch of Douglas. Very frequently it is retroverted and bound down by adhesions, which may, however, allow of a certain range of mobility. Physical Signs.

The chronic form may occur as a sequel to the acute ; most frequently it develops as the result of previous tubal and uterine diseases.

DIFFERENTIAL DIAGNOSIS.

This will be considered under Cellulitis.

COURSE AND RESULTS.

Course and Results. Very often the inflammatory condition clears up. The adhesive form leaves its mark in the shape of pathological anteversions and retroversions bound down (*v. figs. 116, 117*). The Fallopian tubes may have their ovum-conducting power so interfered with that an incurable sterility results. When they are not injured to this extent, conception may occur ;




FIG. 116.

PERITONEAL BANDS binding down the Uterus, Tubes, and Ovaries—result of chronic pelvic peritonitis (*Heitzmann*).

and the adhesions may ultimately yield to the stretching brought to bear on them by the developing uterus. They may, however, resist this and cause abortion.

Occasionally, pelvic peritonitis becomes general and may then be rapidly fatal. We get distension of the abdomen, vomiting of a very severe type, with high pulse, 140 to 160, and often little rise of temperature, this is the form seen in cases of septic infection, the result of abdominal section.

Serous exudations may become absorbed or perforate into the bowel or vagina.

PROGNOSIS.

Each case must be judged on its own merits. We give, therefore, Prognosis, only general hints.

As to life.—Pelvic peritonitis is not usually fatal. If it becomes general and is septic or gonorrhœal in its origin, then the prognosis is very grave. A high and persistently rapid pulse, with a temperature not in the same ratio, also makes prognosis grave. We cannot too strongly impress on the practitioner the importance of pulse rate as affecting prognosis. A pulse below 100 indicates a favourable prognosis; above 120, or when “running,” the prognosis is of the most serious description (*v.* under ovariectomy).

As to sterility.—This is difficult to give, and often time alone settles the point. The mechanical closure by pressure of the Fallopian tube—a condition not diagnosable—and ovaritis rendering ovulation impossible,




FIG. 117.

UTERUS retroverted and bound back by peritonitic adhesions (*Winckel*). *a a* adhesions; *b* bladder; *v* vagina; *u* uterus; *r* rectum. (1)

are conditions often produced and both incurable. Prognosis as to conception should always be cautious, and never absolute when the peritonitis has been extensive.

TREATMENT.

- A. Acute pelvic peritonitis.*—*a. Prophylactic.*
b. General.
c. Local.

Treat-
ment.

a. Prophylactic.—The importance of prophylaxis cannot be overrated. In all operations the strictest antiseptic precautions must be employed. It is attention to this that gives the operator freedom from septic mischief, and it is in this that the limiting of disease finds its best results. The dangers of the consulting-room use of the sound have

Prophy-
lactic
Treat-
ment.

been already alluded to. Care should also be taken by patients during menstruation, and it is also evident that gonorrhœa needs judicious treatment.

General
Treat-
ment.

b. General.—Our views as to the treatment of pelvic peritonitis have undergone considerable modification in view of our knowledge as to its etiology. Formerly, as may be well seen on consulting the earlier editions of the present work the treatment was mainly medical, although the influence of sepsis was recognised, and was moulded on a markedly antiphlogistic type. We have now, however, to consider the matter more broadly, and to take up each case on its own merits, keeping in mind the possibility of surgical interference.

In a case where no septic source is known, or where it cannot be combated as above indicated, the practitioner must have recourse to general treatment.

Diet.

(1.) *Diet.*—In the early stages of inflammation this should be chiefly milk, iced or mixed with aerated lime water or potash water.

Stimu-
lants.

When the patient's strength is reduced and the pulse flagging, nutritious stimulating food must be frequently given. Milk should be still continued; but beef tea or strong soups every two or three hours must be added. Stimulants are requisite at this stage, viz., brandy, champagne, gin, or whisky. Care must be taken to give these in their stimulating doses, *e.g.*, for brandy, a table-spoonful every two or three hours.

Regulation
of the
Bowels.

The regulation of the bowels is not requisite in the early stages; but in the later periods must be looked after. Gentle aperients such as compound liquorice powder, colocynth and hyoscyamus pills, castor oil, etc., can be used; and occasional enemata are of service. Enemata should not, however, be used exclusively, as that might lead to the formation of troublesome scybala.

Tonics.

When suppuration is tedious, it should be seen that no bed sores form; and iron and quinine should be administered.

R. Ferri et Quininæ Citratis gr. lxxx.
Aquæ ʒij.
Sig. Teaspoonful thrice daily in water.

or

R. Ferri et Ammoniaë Citratis gr. lxxx.
Aquæ ʒij.
Sig. Teaspoonful thrice daily in water.

The bitterness is best masked by dilution with water and not with orange or other syrups which derange the stomach.

Treatment
against
Sepsis.

(2.) *To combat any septic condition.*—The first point to determine here is the fact of any source of septic absorption. When this is ascertained to be in any part of the uterine or vaginal tract, prompt disinfection,

with antiseptics, such as the Perchloride of Mercury lotion (1 in 3000 to 4000) is imperative. The Tamponnade with Iodoform gauze is also of great value. Of course this treatment may be, and indeed often is necessary before any distinct symptoms or signs of peritonitis exist, and where the diagnosis is merely that of septic absorption from the genital tract.

In cases where the condition is pronounced, the pulse high, and the temperature not up in the same ratio, or where the peritonitis is threatening to become general, the practitioner has to face the question of having abdominal section performed. He must consider the possibility of there being a ruptured tubal condition, a torsion, or a suppurative condition of a small ovarian cyst. There is no doubt that in such cases the patient's only chance for life lies in the prompt performance of laparotomy, followed by drainage.

We know of no specific medicine for sepsis (*v.*, however, p. 198). A favourite one is the muriate of iron of the Edinburgh Pharmacopœia.

R. Tincturae Ferri Muriatis (Ed. Phar.) ʒij.

Sig. Thirty drops thrice daily in a glass of water. Water should be drunk freely after the dose is given, and the mouth thoroughly rinsed with bicarbonate of soda and water.

Quinine may be used for the same purpose.

R. Quininæ Sulphatis	gr. xxxvi.
Acidi Sulphurici diluti	ʒij.
Aquam ad	ʒvj.

Sig. Tablespoonful thrice daily in water.

(3.) *To alleviate pain.*—Nothing is so good for this as the hypodermic Treatment of Pain.
injection of morphia.

R. Morphinae Bimeconatis	gr. viij.
Spiritus Vini Rectificati	miiij.
Aquæ	̄i.

Sig. For Hypodermic injection. Fifteen minims contain $\frac{1}{4}$ grain of morphia.

The bimeconate is a good preparation and causes less sickness than other forms; as one drachm of this preparation contains one grain of morphia, and as the hypodermic syringe holds only 30 minims, it is impossible to give an overdose to an adult.

When doses larger than half-a-grain are needed, the hypodermic solution of the acetate of morphia (B. P.) may be employed. Twelve minims contain 1 grain, and therefore 3 minims is the first dose for an adult.

It is a good plan for the practitioner to keep the ordinary 8 gr. to 3i.

solution, and to prescribe the stronger solution only for any patient requiring it; in this way he avoids carrying two solutions of different strength by which mistakes might arise.¹ The *stronger* solution is prescribed as follows :—

R. Injectionis Morphinae Hypodermicae (B. P.) ℥ij.

Sig. For Hypodermic injection. Ten minims contain 1 grain
Acetate of Morphia. Dose, 1 to 5 minims.

Chlorodyne (25 min.); extractum opii liquidum (10 min.) or laudanum (tinctura opii, 25 min.) may be used. More useful than these are morphia suppositories.

R. Morphinae Hydrochloratis gr. $\frac{1}{3}$

Fiat suppositorium. Mitte tales vj.

Sig. As directed.

It is a good plan to quiet the pain rapidly with the hypodermic injection; and to keep up the good effect by suppository, in $\frac{1}{3}$ grain doses every six hours, beginning 6 to 8 hours afterwards. See that the patient or attendant understands that the suppositories are to be passed into the empty bowel.

Opiates should not be used indiscriminately, as they are apt to mask symptoms. The patient cannot be allowed to suffer, however, and therefore the medical man's hands are forced in this respect.

Treatment
of High
Pulse and
Tempera-
ture.

(4.) *To bring down pulse and temperature.*—In early stages, tincture of aconite or veratrum viride was at one time considered of great value, but is little employed now.

R. Tincturae Aconiti ℥ij.

Sig. Six drops are to be put in a wine glass containing six teaspoonfuls of water. Give a teaspoonful every quarter of an hour.

If the temperature keep high, quinine in 5–10 grain doses may be given. The salicylate of quinine is a good preparation and is given just as quinine is. When the stomach is irritable the quinine, in 20 grain doses, suspended in an ounce of mucilage, may be given per rectum.

Antipyrin (10–15 grains) and antifebrin (5–10 grains) are useful. The former also aids in headache, but the latter tends to produce cyanosis, and though very effective requires to be watched for undue depression. Alcohol may be given with it (*v. Leech, Med. Chron., Vol. VIII., p. 297*).

After the fever has subsided and suppuration threatens, the strength must be kept up by tonics (such as quinine and iron) and by nutritious food with a judicious amount of stimulant, claret for example.

¹ Morphia is also made up in compressed Hypodermic Tabloids, containing various doses. They are readily dissolved in a few drops of water, are both reliable and portable, and should supersede fluid preparations. There can be no accidental overdose with them, a matter of importance.

c. *Local Treatment*.—Ice is not generally used as a local application in this country, and has its disadvantages. Local Treatment.

Of greater use are large hot linseed poultices. They should be made very hot, a layer of flannel intervening between them and the skin, and should be covered with a layer or two of cotton. Such a poultice will be effective for 2 or 3 hours. Blisters and turpentine stupes are good, but soon render the skin so sore that after-treatment by poultices is difficult.

The hot vaginal douche (as directed at page 150) with carbolic acid added in septic cases, should on no account be omitted.

Encysted serous collections should, as a general rule, be left to be absorbed. When troublesome from pressure, they may be tapped by Matthieu's aspirator. A clear serous fluid, often coagulable, is then drawn off, so like urine that the almost involuntary first thought is that the operator has tapped the bladder by mistake.

The treatment of suppuration will be best considered under pelvic cellulitis.

The modern standpoint as to pelvic peritonitis may be summed up as follows. Ascertain the source of the mischief and deal with it by local disinfection or by surgical means when possible. Use palliative treatment carefully, and be especially cautious in regard to antiphlogistics, and opiates.

B. *Treatment of chronic pelvic peritonitis*.—When adhesions are extensive, and tubal disease present, the question of abdominal section has to be considered. Cullingworth's papers give valuable facts in regard to this point. When the uterus is retroverted, it may ultimately be replaced by bimanual manipulation. Massage is good in such cases, but its employment will be considered afterwards when we speak of the systematic treatment by rest and food (*v.* Appendix). Treatment when Chronic.

Of late, since our knowledge of the nature of tubercle has been rendered more exact by Koch's discovery of the tubercle bacillus, *tubercular* peritonitis has been found to be by no means rare; and the bacillus tuberculosis has now been discovered, sparingly and in giant cells, by several observers. We may also have *malignant* peritonitis, due usually to rupture of papillomatous ovarian cysts. In both the tubercular and the malignant form we get ascitic fluid, but characteristic cells in the latter only.

TUBERCULAR PERITONITIS.

LITERATURE. *Hegar*—Die Entstehung, Diagnose und chirurgische Behandlung der Genitaltuberculose des Weibes: Stuttgart, Enke, 1886. *Poten*—Ein Fall geheilter Bauchfelltuberculose: Cent. für Gyn., 1887, S. 33. *Schwarz*—Ueber die palliative Incision bei Peritonitis tuberculosa: Wien. Med. Wochens., No. 13, 1887. *Tait*, *Lawson*—Diseases of the Ovaries, fourth edit., p. 334: Birmingham, 1883. *Wells*,

Sir T. S.—Ovarian and Uterine Tumours, p. 100: Churchill, London, 1882.
Williams, J. Whitridge—Tuberculosis of the Female Generative Organ: Johns Hopkins Reports, III., p. 85. For history and further literature see Williams, Schwarz, or Cassel's Year Book for 1888.

Preliminary Remarks.—The serious results of tubercular disease of the lungs, meninges, and mucous tracts, render the comparatively good prognosis in tubercular peritonitis as remarkable as it is at present inexplicable. In Wells' historical case in 1862, abdominal section was performed for ascites due to peritoneal tuberculosis, miliary tubercles were found studding the bowel surface; and yet, as the result of the section and evacuation of fluid only, complete recovery took place, the patient being well nineteen years afterwards (1881). Since then, equally good results have been obtained by others.

Pathological Anatomy.—The condition is secondary, and may exist as miliary tubercle, acute or in a caseous ulcerating condition; or as chronic fibro-tuberculosis.

Symptoms.—It is remarkable that this condition may be latent, may be only discovered by accident, and that the temperature may be sub-normal or high.

The patient's general health may be good, with no rise of temperature if the peritoneum alone is affected. It must be kept in mind, however, that the lungs may be simultaneously implicated.

Physical Signs.—We may have fluid in the abdomen so encysted as to simulate ovarian cyst, or there may be free fluid but with impaired mobility and with irregular lumps, due to matting of bowels and omentum.

The *Differential Diagnosis*, which is chiefly from typhoid fever, ovarian cyst, and malignant peritonitis, is difficult and may be cleared up only by exploratory incision.

Treatment.—Abdominal Section, with complete evacuation of fluid and careful peritoneal toilette so as to dry out as thoroughly as possible, is all that is requisite; the use of antiseptic irrigations or the applications of iodoform to the peritoneum before the wound is closed has been found unnecessary, and the same may be said of drainage. In 17 cases collected by Schwarz, the general age was seventeen to thirty-three; youngest, four; oldest, fifty-seven. Immunity was found in these to range from two to ten years, but one case of complete cure has been recorded by Wells. A phthisical condition of lung, if not too far advanced is not a contra-indication.

When no fluid, or only a matted condition, is present, or where the masses are solid, linimentum hydrargyri may be freely applied to the skin of the abdomen and the patient put under favourable general conditions, viz., dry hill air, and abundant use of fattening food.

MALIGNANT PERITONITIS.

By this we mean a condition where the peritoneum is more or less invaded by papillomatous growths, secondary usually to rupture of papillomatous cysts of the ovary (*v.* Pathology of Ovarian Tumours, Chap. XXII.).

Symptoms.—The patient is not at first cachectic, and the only thing attracting attention is the distention of the abdomen from fluid. The condition is not necessarily fatal, and we have seen one case where the patient lived for three or four years. It may, however, soon cause death when pleuritic or pericardial effusions come on.

The *Physical Signs* are abdominal distention, irregularly encysted fluid, irregular masses felt in the abdominal cavity on palpation, with occasionally secondary nodules in the pelvic or iliac glands, and characteristic cell groups in the fluid drawn off. These render diagnosis fairly easy (fig. 141A).

The *Treatment* is palliative by tapping.

PELVIC CELLULITIS (PARAMETRITIS).

SYNONYM.—Parametritis, a term sometimes limited to inflammation of the cellular tissue round the cervix and upper part of vagina—Virchow's ^{Pelvic Cellulitis.} parametric tissue. At the close of this chapter, we shall have to notice specially a variety of this described by W. A. Freund as *Parametritis chronica atrophicans circumscripta et diffusa*.

NATURE.—An acute or chronic inflammatory affection, usually septic, affecting the cellular tissue of the pelvis.

PATHOLOGICAL ANATOMY AND VARIETIES.

It is the rare exception to examine a multiparous female pelvis without finding some trace of a previous cellulitis or peritonitis. Thus split ^{Pathological Anatomy and Varieties.} the cervix, so common in women who have borne children, is almost always associated with some cellulitis at the base of the broad ligaments. The uterus is rarely central, but is often drawn to the one side by the cicatrization of some previous lateral cellutic inflammation of the broad ligament; the traction may even be so great that it lies at right angles to its proper axis. We have seen that the utero-sacral ligaments are peritoneal folds containing connective tissue and unstriped muscular fibre. Inflammatory attacks in one or both of these folds (combined pelvic peritonitis and pelvic cellulitis) are very common. Schultze calls this "parametritis posterior," but utero-sacral cellulitis is a more accurate term. The cicatrization of these ligaments after such inflammation, causing traction just above the isthmus, brings about the most common cause of dysmenorrhœa and sterility—pathological ante flexion

of the uterus (*v.* Antelexion of the Uterus). It is evident that in this way, too, we get the uterus antelexed and drawn to one side, or antelexed and drawn back (fig. 89).

Sometimes pelvic abscesses are found in localities to be afterwards alluded to. Often the uterus and ovaries are in an atrophic condition owing to compression of the vessels and nerves by the cellutic attack; this quite agrees with the clinical fact that many women with bad pathological antelexion do not suffer much at their periods, because the withered condition of the organs produces scanty menstruation. According to some, we can have no cellulitis in the broad ligaments and no formation of pus—abscess of the broad ligaments. Clinical, anatomical, and pathological evidence is in favour of the occurrence of both. At the same time, it is almost impossible clinically to distinguish abscess of the broad ligament from an encysted serous pelvic peritonitis behind it pushing it forwards, or in some cases from a distended tube.

ETIOLOGY.

Etiology. In *parous women* the great cause of pelvic cellulitis is probably septic matter (*i.e.*, either cocci or bacilli, or their products) absorbed by the lymphatics from the torn perineum, vagina, or cervix. This passes along the abundant lymphatics and blood vessels in the cellular tissue beneath and in the broad ligaments, causing inflammation of the glands and proliferation of the connective tissue in which these are embedded. Thus we find *childbirth*, *premature labour*, and *abortion*, often followed by cellutic attacks for obvious reasons. In parturition we have the cervix, for instance, torn vertically at one side; and septic matter deposited there often speedily spreads along the lymphatic stream (*v.* page 77).

In *nulliparæ*, cellulitis may arise from the same causes as are given under pelvic peritonitis, *e.g.*, exposure to cold during menstruation.

Pelvic peritonitis, in a minor degree, is always associated with cellulitis as already mentioned. So far as we have considered the etiology of pelvic inflammatory affections, we have associated them with some virus, most frequently septic. We do not believe that mere traumatic injury, apart from septicity and tension, can cause an inflammatory attack.

SYMPTOMS.

Symptoms. The patient has a rigor or chill. Pain is felt over the lower part of the abdomen, but it is not so intense as in peritonitis. The pulse and temperature are raised, the pulse being lower and the temperature higher, than in pelvic peritonitis. After exudation has taken place, the patient may have one thigh alone drawn up.

PHYSICAL SIGNS.

There is pain on palpation of the abdomen ; and after exudation has taken place, we feel a fulness at one side of the uterus or in the iliac fossa. Physical Signs.

Bimanual examination, always difficult, reveals at first nothing but increased heat and tenderness. After exudation has occurred, it is found in the following positions :—

- (1.) As a bulging at the side of the uterus, depressing the lateral fornix and pushing the uterus usually to the other side ;
- (2.) In the upper portion of the broad ligament, and therefore not bulging downwards ;
- (3.) In the iliac fossa ;
- (4.) Very rarely, behind the uterus ;
- (5.) Still more rarely, between uterus and bladder.

We have seen pus pointing in the inguinal region on one side, and with no dipping down into the pelvis or immediate connection with the side of the uterus. The intermediate deposit has cleared away, or it may be that the parametritis was remote from the uterus. When pus is present in large amount, the fluctuation can be felt bimanually. When it forms in the centre of a large inflammatory exudation, an obscure boggy feeling may or may not be made out. Aspiration prior to evacuation may help here very much. But it is better to incise where, from the fact that there is an evening rise of temperature with occasional rigors and sweatings, pus is diagnosed.

The course of these exudations, inflammatory and purulent, is explained in two ways. Explanation of course of Exudations.

(a.) By the course of the lymphatics, which run, as we have seen, from the uterus outwards beneath and between the layers of the broad ligament to the glands in the lumbar region.

(b.) By the lines of cleavage in the cellular tissue of the pelvis. The student should refer back to the description of cellular tissue of the pelvis given in Chap. II., and especially to König's researches (page 47). Based on these, and on clinical work, König holds that—

- (1.) An exudation in the broad ligament, near the tube and ovary, passes first along the psoas and iliacus and then sinks into the true pelvis ;
- (2.) Exudations which begin primarily in the deeper cellular tissue on the antero-lateral aspect of the cervix, pass first on to the cellular tissue of the true pelvis at the side of the uterus and bladder, then with the round ligament to Poupart's ligament beneath the inguinal canal, and then they pass outwards and backwards into the iliac fossa ;
- (3.) Abscesses, developing from the posterior aspect of the broad ligaments, fill first the postero-lateral part of the pelvis and then pass as in (1.).

DIFFERENCES AND DIFFERENTIAL DIAGNOSIS BETWEEN ACUTE PELVIC
PERITONITIS AND CELLULITIS.

Differences and Differ- ential Diagnosis.	<i>Differences.</i>	
	<i>Pelvic Peritonitis.</i>	<i>Pelvic Cellulitis.</i>
	(1.) Inflammatory affection of pelvic peritoneum chiefly.	(1.) Inflammatory affection of pelvic cellular tissue chiefly.
	(2.) Usually general, round the uterus.	(2.) Usually lateral.

Differential Diagnosis.

<i>Pelvic Peritonitis.</i>	<i>Pelvic Cellulitis.</i>
(1.) Pain very severe.	(1.) Pain not so severe.
(2.) Patient's legs drawn up on both sides.	(2.) Usually only one leg drawn up.
(3.) Firm flat effusion not bulging into fornices, and situated round the uterus ; or a mesial bulging of serous effusion behind uterus. Cervix (vaginal portion) is of normal length.	(3.) Firm effusion, bulging usually into fornix of one side. Thus cervix (vaginal portion) apparently shortened on one side.
(4.) Does not spread along round ligament or into iliac fossa, but may affect all peritoneum.	(4.) Exudation or pus spreads in definite directions, and is usually localised.
(5.) Uterus displaced to front, or unaltered in position.	(5.) Uterus usually displaced to one side.
(6.) Vomiting more frequent.	(6.) Vomiting less frequent.

It is often very difficult to differentiate these ; and therefore in some cases the diagnosis must be pelvic inflammation—probably cellullitic or probably peritonitic or tubal, as the case may be.

COURSE AND RESULTS.

Course and Results.	Very often the attack passes off and leaves no trace. The septic poison is too small in amount to do harm ; or it sets up some inflammatory exudation, which mechanically arrests progress, and then becomes absorbed. The vitality or health of the tissues and the strength of the poison have also their share in determining its progress. Exudation may take place and may be absorbed almost completely, may suppurate slowly, and only to a limited extent, and may form a large abscess. This abscess may open into the bowel or bladder, or pass below Poupart's ligament, or upwards beneath the kidney. Rarely does it appear in the perineum, or pass through the sciatic notch to the buttock. In one case where the
---------------------	---

last occurred, the patient complained of a very deep-seated pain just over the notch.

It is valuable to note how rarely the abscess perforates into the peritoneal cavity. The peritoneal surfaces of the abdominal contents are in contact; and as the inflammatory attack spreads, it sets up a peritonitis which glues the adjacent surfaces together. When pus does enter the peritoneal cavity, it sets up a rapidly fatal peritonitis.

Matthews Duncan has pointed out that albuminuria is often present in pelvic cellulitis but not in pelvic peritonitis; it was present in 6 out of 16 cases (37·5 p. c.) of cellulitis, but absent in 32 cases of peritonitis.

PROGNOSIS.

This depends on the extent of the inflammatory attack, and its effect on the patient's health. Its septic origin usually causes anxiety; but it does not spread so rapidly as peritonitis. Resolution of inflammatory deposits is slow. Pathological ante flexion gives rise to troublesome dysmenorrhœa and sterility. Prognosis should always be guarded as to complete recovery.

TREATMENT.

The general and the local treatment are exactly the same as in pelvic peritonitis. The occurrence of suppuration is indicated by rigors, and should be hastened by the hot douche and poultices. We may have only part or parts of the exudation suppurating, so that in a cellulitic swelling we may have inflammatory exudation containing separate abscess cavities. Within the past few years, as the result of the work of Péan and his pupils, especially Ségond, Hysterectomy has been practised in cases of pelvic suppuration around the cervix. The results have been good, and there is no doubt the use of this method of interference will increase. We do not discuss it at present, as Hysterectomy is now proposed in many cases where Abdominal Section has hitherto been the recognised procedure, and this renders it advisable to discuss the whole question in a separate chapter (*v.* Abdominal Section in Appendix).

When pus is present in large quantity, the treatment varies according to the part at which it points.

(1.) If it point above or below Poupart's ligament, in the buttock, or behind the kidney, it is to be opened with antiseptic precautions, and a drainage tube inserted. Results by this method are admirable. In certain cases abdominal section may be performed, as in abscess occupying the broad ligament, and the opening in the ligament stitched to the abdominal incision and drained.

Treatment
of Pelvic
Abscess.

(2.) If it bulge behind the cervix, it may be opened as follows:—pass Sims' speculum, and open into the cavity with Paquelin's cautery at a dull heat; make the opening big enough to admit two good-sized drainage tubes. Daily irrigate the cavity with weak carbolic lotion (1-100) or boracic lotion (1-30). If the discharge is profuse, it may be received into pads of sublimated wood-wool wadding placed over the vulva; oakum or marine lint may be used among the poor.

According to the position of the bulging the operator may make an incision with the knife in the anterior fornix (rare) or in the lateral fornix. In the former instance he speedily reaches the loose tissue in front of the cervix, and gets into the abscess cavity, care being taken not to injure the bladder. In the case of the lateral fornix a preliminary silk ligature should be passed with Olshausen's needle, or the tissue may be clamped with pressure forceps after a superficial incision has been made close to the cervix, before the deeper tissues are cut.

The drainage tubes should be double, and with a small piece at the end at right angles which prevents their slipping out. They should not be perforated, as this prevents the washing out. Straight tubes can be fastened with a stitch to the edge of the incision.

The practitioner will very often find the remains of cellulitis as an indistinct thickening in the fornices. For these, blisters in the iliac regions, the glycerine plug, and hot douche, are useful (*v.* under Chronic Ovaritis).

TREATMENT OF SEPTIC CONDITIONS BY ANTITOXINES.

In a patient who has septic organisms in the tissues of the genital tract we get a formation of toxins, and by the absorption of these she becomes poisoned, and ultimately may succumb. We attribute to the toxins the general functional disturbances, the rise of temperature, and above all the increased rate of the pulse. No one can see a case of septicaemia after an abdominal section or abortion without feeling that the patient is under the influence of a cardiac poison. In a case likely to be fatal the pulse rises to 120, goes on to 140, begins to flag and become irregular; and when this irregularity comes on, the end is not far off. The student now understands that in inflammatory affections of the tube and ovaries, of the peritoneum and cellular tissue, cocci and bacilli play an essential part; that in fact, they are the cause of the infection in the septic group of these conditions. The question of treatment of such causes naturally arises, and hitherto that has been most successful in the way of prevention (*v.* Chapter XV. on Antisepsis, etc.). In giving the treatment of pelvic peritonitis at page 189, the statement is made, "We know of no specific medicine for sepsis." This statement requires modification, however, in view of recent work.

Micro-organisms kill not so much by their actual presence in the blood and tissues, but by poisons they secrete, viz., toxines.

The remarkable researches of Löffler, Behring, Roux, S. Martin, and others, have shown that diphtheria is due to the presence of a bacillus, and that it secretes a poisonous albumose or toxine. This toxine can be obtained from pure cultures of the bacillus, and when injected hypodermically into the horse, there can be obtained from the serum of its blood after a time, an antitoxine which neutralizes the toxine. We give only the barest outline of this as it does not belong to our subject. The special application comes now.

We have seen that streptococci are often the organisms of sepsis, and it is of the greatest importance to ascertain if a toxine is produced by them, and if an antitoxine or anti-streptococcic serum can be obtained as in the case of diphtheria. This has been accomplished by Marmorek, who obtained toxines from cultivations of streptococci, and also, by injecting this toxine into the donkey, an anti-streptococcic serum from its blood.

We have thus reached a most important stage in the treatment of septic conditions, and may put the matter thus. In a patient with septicaemia the condition is probably due to a local development of streptococci which are secreting toxines, and poisoning the patient. We say "probably due to a local development of *Streptococci*" as other micro-organisms, notably staphylococci and the bacterium coli commune may be the cause of the blood poisoning, and against them the anti-streptococcic serum is inert. The treatment and cure of such a case must therefore not only be by local disinfection (*v.* Chapter XV.) and general treatment, but we may neutralize the toxine by the hypodermic injection of the anti-streptococcic serum. This may be used in hypodermic doses of 10 ccs. (about 3-4 drachms) in 24 hours; and already gratifying results have been obtained. There is marked reduction of pulse and temperature, and unless the local infection is too far advanced or actual disinfection impossible, the results are likely to prove most satisfactory.

The far-reaching nature of such researches needs no remark.

EFFECTS OF PELVIC PERITONITIS AND CELLULITIS ON THE UTERUS.

It is unfortunate that uterine displacements have of late years bulked so largely in gynecology—we mean by this that many regard a uterine displacement in itself as a condition sufficient to account for symptoms of bearing down pain, leucorrhœa, or even for sterility and dysmenorrhœa. It is a well-ascertained fact that uterine displacements are in many cases the result of antecedent peritonitis or cellulitis, are mere physical signs of these affections, and therefore secondary lesions of far less importance than the pelvic inflammation which was the primary one.

Effects of
Peritonitis
and Cellu-
litis on the
Uterus.

These displacements might be grouped under the two heads:—

- A. those caused by Pelvic Peritonitis;
- B. those caused by Pelvic Cellulitis.

A. Displacements caused by Pelvic Peritonitis.

Displace- From the lymph effused and the resultant bands formed in pelvic
ments from peritonitis, the uterus becomes bound to the adjacent peritoneum on the
Peritonitis. rectum (retroversion and retroposition); or more rarely, to that on the
bladder (anteversion); sometimes it is twisted on its long axis or matted
to the coil of intestine surrounding it. Figs. 116, 117, 118, illustrate
these conditions.




FIG. 118.

PERITONITIC ADHESIONS DRAWING THE UTERUS TO ONE SIDE (*Heitzmann*).

The *Diagnosis* of such adhesions is made by digital pressure through the rectum in the case of retroversion, and through the anterior fornix in anteversion. In the former case, the immobility of the uterus is felt; and when pushed up so as to be manipulated by the abdominal hand, replacement is found to be impossible; or if partially successful, the displacement returns almost immediately. Sometimes the retroverted uterus when not adherent is replaced with difficulty, owing to the cohesion of the peritoneum on the posterior uterine surface with the peritoneum behind it, and this point has to be borne in mind. The sound should certainly not be employed in cases with adhesions, as, by its leverage, vascular adhesions may be torn, and hæmorrhage produce hæmatocele with subsequent pelvic peritonitis.

B. *Displacements caused by Pelvic Cellulitis.*

These are two in number: viz. (a.) Lateriversion; and (b.) Pathological Ante-<sup>Lateri-
version.</sup>flexion due to Utero-sacral Cellulitis.

(a.) *Lateriversion* is the result of cellulitis in one broad ligament, subsequent absorption of the inflammatory effusion, and cicatrisation of the ligament. The *Diagnosis* of this condition is easy. There is often a split of the cervix at the side corresponding to the displacement as well as scarring in the fornix with coincident displacement of the cervix. Bimanually, the uterus is felt drawn to the one side, fixed, and sometimes the body is lateriflexed as it were on the cervix. Bimanual displacement of the uterus to the non-affected side causes pain. The pathology of this displacement in many cases is that cellulitis, probably septic, has spread after parturition from the split cervix along the lymphatics at the base of and in the broad ligament; effusion of lymph, perhaps of pus, has followed; finally there result the incomplete resolution and cicatrisation already mentioned.

(b.) *Pathological Ante-<sup>Patho-
logical
Ante-
flexion.</sup>flexion due to Utero-sacral Cellulitis* is one of the most important, most intractable, and most misunderstood of lesions. Its nature may be thus described. A cellulitis, in or in the neighbourhood of the utero-sacral ligaments, has gone on to cicatrisation,—producing fixation of the uterus and, along with the action of intra-abdominal pressure, ante-^{flexion.}flexion (*v.* Chap. XXXIII. Displacements of the Uterus). This cellulitis is often the result of abortion, more rarely of full-time parturition; it is frequently found in nulliparæ, and may in some cases be due to the zymotic diseases of childhood.

This condition is *diagnosed* as follows: on vaginal examination, the cervix is found high up, because drawn back, and pointing usually downwards and forwards; through the anterior fornix the body of the uterus is felt. Bimanually, the uterus is recognised as ante-^{flexion.}flexed (see under Ante-^{flexion.}flexion). Through the posterior fornix we feel thickening and fixation of the tissue in the neighbourhood of the utero-sacral ligaments, or we may sometimes feel the thickened ligaments themselves running in a direction forwards and inwards. The rectal examination gives valuable information, as the thickening is more distinctly felt, the ante-^{flexion.}flexion is more accurately mapped out and ovaritis or other inflammatory thickening discovered.

The amount of fixation should be estimated by bimanual movement of the uterus, as this helps in prognosis. Often the cellulitis affects one side of the parametric tissue and gives a displacement of the uterus towards the posterior extremity of an oblique diameter of the pelvis.

We shall have again to consider the symptoms and treatment of these conditions in the chapter on Displacements of the Uterus. From what

has been said, however, it will be evident that their treatment should be simply that of chronic peritonitis and cellulitis.

PARAMETRITIS CHRONICA ATROPHICANS.

Para-
metritis
Chronica
Atrophica.
cans.

We have already described some of the results of acute pelvic peritonitis and cellulitis in causing pathological retroversions and ante-flexions. W. A. Freund of Strassburg has drawn attention to a condition of the pelvic connective tissue, similar in some of its results but differing from what we have described in not having an acute stage. He terms it *Parametritis Chronica Atrophicans Circumscripta et Diffusa*. His researches are very valuable and explain results usually ascribed to mere displacements of the uterus or the pathological condition of the cervix; they also give a basis for treatment or at least show the futility of much of the mechanical treatment by pessaries.

a. Parametritis Chronica Atrophicans Circumscripta.

Nature.—A circumscribed chronic inflammatory process affecting chiefly the fascial and aponeurotic thickenings of the fatless connective tissue, and causing changes analogous to those in cirrhosis of the liver, kidney, and spleen.

Etiology.—The primary cause may lie in bladder, rectum, or uterus. When in the bladder, there has been some ulcerative process from which irritation has passed causing paracystitis chronica atrophicans (inflammation of the connective tissue near the bladder). From the side of the bladder, thickenings in the connective tissue pass outward and forward and by their ultimate atrophy bring about uterine displacement in the opposite direction: thus, left paracystitis will cause retro-dextro-flexion of the uterus, while right paracystitis will bring about retro-sinistro-flexion.

In the rectum, the starting-point may be dysenteric or simple follicular ulceration at the level usually of the anterior fold of mucous membrane forming part of the sphincter tertius. The cellulitic irritation runs in the utero-sacral ligaments and causes pathological ante-flexion. This effect of rectal disease has not been sufficiently recognised in this country and is worthy of clinical and pathological investigation.

Freund records two interesting post-mortems of chlorotic women, 19 and 23 years of age respectively: the heart, large arteries, and kidneys were hypoplastic (*i.e.* insufficiently developed); the ovaries were small and cystic; chronic pelvic peritonitis was present in Douglas' pouch; and finally, there was follicular ulceration above the sphincter tertius, and chronic paraproctitis (chronic inflammation of the connective tissue near the rectum) with shortening of utero-sacral ligaments.

In the uterus, split cervix is one great cause; we have, radiating from the split, chronic thickening running along the base of the broad ligament behind the cervix and down to the fornix. By the atrophy

and cicatrisation of these chronic inflammatory thickenings, there result ultimately displacements of the uterus, compression of the veins, and therefore catarrh of the cervix with reflex pains due to alterations in the sympathetic filaments distributed in the connective tissue.

In *diagnosis*, careful examination (vaginal, rectal, and bimanual) reveals the thickening due to the chronic parametritis, and the consequent displacement; the initial lesion in bladder, rectum, or uterus, may be made out.

b. Parametritis Chronica Atrophicans Diffusa.

We have here a condition whose pathology is not so evident as that of the circumscribed form. It is said to begin in the base of the broad ligaments and to pass out to the pelvic walls. Ultimately, the whole pelvic tissue becomes dense, the veins partly narrowed and partly dilated, the arteries contracted and the ureters distorted. Hyperæmia of the urethra, the neck of the bladder, and rectum, is present, causing catarrh; while the uterus, at first enlarged and catarrhal, finally atrophies; the Fallopian tubes and ovaries also become atrophied; the vagina is shortened and the external genitals withered.

On microscopic examination, perineuritis of the sympathetic plexuses in the connective tissue has been found (H. W. Freund). The *etiology* is obscure. It may be due to sexual excess or frequent child-bearing and excessive suckling in women with hypoplasia of the genital organs and blood vessels.

Diagnosis is based on careful bimanual examination and determination of the changes above described, by attention to the history and carefully noting the conditions of menstruation (at first profuse and painful, and then scanty), as well as the catarrhal processes going on in the bladder, cervix uteri, and rectum.

Reflex disturbances arise from both varieties of Parametritis Atrophicans, due to the changes (from inflammation and pressure) in the sympathetic filaments. We may speak of these as Sympathetic, Spinal, and Cerebral Hysteria. Reflex disturbances in Parametritis Atrophicans.

In the Sympathetic form, we have neuralgia of the stomach and intestines, aching kidneys, vesical pains, palpitation of the heart, and disturbances of the respiration.

In the Spinal group, there are painful spots over the spinous processes of the cervical, dorsal, and lumbar vertebræ; the pains may radiate laterally and we may get pains in the extremities. Hysterical paralysis may ultimately develop.

In the Cerebral group, there is neuralgia of the fifth nerve, hemi-crania, and fixed boring pains.

The *Prognosis* is fairly good in the circumscribed form but not hopeful in the diffuse.

Treatment.—In the circumscribed form, the cause (in bladder, rectum, or cervix) must, when possible, be treated. The vaginal hot douche and bimanual massage to set up absorption and perhaps stretch nerve filaments (as in Nussbaum's nerve-stretching for sciatica) have done good. The influence of stitching cervical lacerations (Emmet's operation) may be beneficial.

The uselessness of treatment by pessaries of the uterine displacements caused by inflammation is evident.

In the diffuse form and when nervous symptoms arise, we must rely on nervous remedies, chiefly bromide of potassium. For the neuralgia, the constant current and systematic massage may be tried; and, for the paralysis, the interrupted form. The patients ultimately get well, but it is a question of years.

CHAPTER XVIII.

PELVIC HÆMATOCELE AND HÆMATOMA: NEW GROWTHS IN THE PELVIC PERITONEUM AND CONNECTIVE TISSUE.

LITERATURE.

HÆMATOCELE AND HÆMATOMA.—*Aitken, Lauchlan*—Case of Pelvic Hæmatocele: Ed. Med. J., 1862, p. 104. *Bandl*—Die Krankheiten der tuben, etc., Billroth's Handbuch: Stuttgart, 1879. *Barbour*—The Diagnosis of Advanced Extra-uterine Gestation after the Death of the Fœtus: Edin. Med. Jour., 1882. *Barnes*—Diseases of Women: London, 1878, p. 590. *Bernutz and Goupel*—Clinical Memoirs of the Diseases of Women, Vol. II.: New Syd. Soc., Meadows' Tr., 1866. *Bernutz*—Hématocèle utérine: Archiv de Tocologie, 1884, p. 978. *Bourdon*—Tumeurs fluctuantes du petit bassin: Rev. Méd., 1841. *Credé*—Monatsschrift f. Geburtskunde, Bd. IX., S. 1. *Duncan, Matthews*—Uterine Hæmatocele: Ed. Med. J., 1862, p. 418: Diseases of Women, Churchill, Lond., 1883. *Düvelius*—Beitrag zur Lehre von der operativen Behandlung des Hæmatoma periuterinum extraperitoneale: Archiv für Gynäk., Bd. XXIII., S. 107. *Freund*—Die Gynäkologische Klinik: Strassburg, 1885. *Fritsch*—Die retrouterine Hæmatocele: Volkmann's Sammlung, No. 56. *Gusserow*—Ueber Hæmatocele periuterina: Arch. für Gynäk., Bd. XXIX., Hft. 3. *Hart and Carter*—The Frozen Sectional Anatomy of Advanced Extra-uterine Gestation: Ed. Med. Jour., Oct. 1887. *Hunter, Wm.*—Intraperitoneal Blood Transfusion and the Fate of Absorbed Blood: Jour. of Anat. and Phys., 1887. *Imlach*—Pelvic Hæmatocele treated by Abdominal Section: Brit. Med. Jour., 1885, I., 933; and 1886, I., 339. *Kuhn*—Ueber Blutergüsse in die breiten Mutterbänder und in das den Uterus umgebenden Gewebe: Zurich, 1874. *M'Clintock*—Diseases of Women: 1853. *Macdonald, Angus*—Uterine Hæmatocele, Ed. Med. J., Jan. 1883. *Martin*—Das extraperitoneale periuterine Hæmatom: Stuttgart, Enke, 1881. *Nélaton*—Gaz. des Hôpitaux, 1851 and 1852. *Olshausen*—Ueber Hæmatocele und Hæmatometra: Arch. für Gynäk., Bd. I., S. 24. *Pelletan*—Clinique Chirurgicale: Paris, 1810. *Priestley, W. O.*—Pelvic Hæmatocele: System of Gynecology: London, 1896. *Schroeder*—Handbuch der Krankheiten, etc.: Leipzig, 1878, S. 453. Ueber die Bildung der Hæmatocele retrouterina und anteuterina: Arch. f. Gyn., Bd. V., S. 348. *Simpson, Sir J. Y.*—Periuterine or Pelvic Hæmatocele, Collected Works, Vol. III., p. 121: A. & C. Black, Edinburgh. *Sutton, J. Bland*—Surgical Diseases of Tubes and Ovaries: Cassell & Co., 1896. *Tait, Lawson*—Lectures on Ectopic Pregnancy and Pelvic Hæmatocele: Birmingham, 1888. *Tilt*—Pathology and Treatment of Sanguineous Tumours: Lond. 1853. *Voisin*—De l'hématocèle Rétro-utérine: Thèse, Paris, 1858. Traité de l'hématocèle: Paris, 1860. *Webster, J. C.*—Ectopic Pregnancy: Pentland, 1895. *Zweifel*—Zur Behandlung der Blutergüsse hinter der Gebärmutter: Archiv f. Gyn., Bd. XXII., S. 185. Zur Behandlung der Blutergüsse hinter der Gebärmutter: Arch. f. Gynäk., Bd. XXIII., S. 414. For Tubal Abortion, see Bland Sutton and J. C. Webster.

PELVIC HÆMATOCELE AND HÆMATOMA.

Prelimi-
naries.

Preliminary Considerations.—The abundant venous supply of the pelvic organs, the congestion induced by menstruation, the hæmorrhage accompanying the monthly rupture of the Graafian follicle, and especially the rupture of an early extra-uterine gestation, render women peculiarly liable to hæmorrhages into the pelvic cavity. Yet it is astonishing that it is only since 1850 that this subject has really attracted gynecologists' attention. It was in that year that Nélaton gave the subject due prominence; although Recamier (1831), Bourdon, Velpeau, and Bernutz had all recorded cases—under such titles as “Bloodgush from an aneurism of the ovary,” “Blood cysts of the pelvic cavity.” Nélaton had diagnosed his case as an abscess, and opened it with a bistoury; the blood and blood clots escaping from the incision showed its real nature unmistakably. Since that time pelvic hæmatocele has taken its place in Gynecology as a serious and important *symptom*.

Terminology.—The hæmorrhage is either intra-peritoneal or extra-peritoneal, but both forms may be present. The terminology is at present unsettled. “Hæmatocele” means “hæmorrhage into the peritoneal cavity,” but we may use the phrase “pelvic hæmatocele” as including both varieties, and add “intra-peritoneal” or “extra-peritoneal” where the diagnosis can be made. “Hæmatoma” is sometimes used instead of “extra-peritoneal hæmatocele.” “Retro-uterine” hæmatocele is employed when the bulging is distinctly behind the uterus.

NATURE.—*An effusion of blood into the pelvic peritoneum or connective tissue.*

Pelvic hæmatocele is thus not a disease. It is only a symptom of some previously existing pathological condition of the pelvic organs, just as hæmoptysis is not a disease but usually a symptom of some lung condition.

PATHOLOGICAL ANATOMY.

Patho-
logical
Anatomy.

Our knowledge on this point is extremely defective, although of late some light has been thrown on it by information gained from abdominal section, and more especially by the admirable work of William Hunter. From experiments on the lower animals by intra-peritoneal transfusion of blood, he has arrived at the following conclusions:—

“The results of the foregoing experiments may be regarded as definitely proving, that in the case of the peritoneal cavity at least the fate of extravasated blood is not so entirely a merely local one as has hitherto been generally supposed. On the contrary, a very considerable, sometimes even a large, proportion of the red corpuscles may escape a local fate altogether, becoming absorbed mainly through the lymphatics of the diaphragm into the circulation, where they continue, for a certain time at least, to perform their functions as before.

"The rapidity with which this absorption takes place is always both relatively and absolutely greatest during the earlier hours after the effusion, especially in the case of entire blood, the absorption extending, however, over a period of twenty-four hours or even longer according to the amount of the effusion.

"The maximum increase is attained to on the third or fourth day after the injection, the time depending partly on the quantity of blood transfused, partly on its fluidity. . . .

"The actual absorption of corpuscles which takes place during the earlier hours after the transfusion can, however, never be accurately determined, even by enumeration of the corpuscles in the circulating blood, still less by estimation of the hæmatoglobin. For owing to the serous effusion which almost always occurs into the abdomen as the immediate result of the injection, the number of corpuscles in the circulating blood as determined by enumeration, is always apparently much increased; and it is not till this effused serum, along with the injected serum, has become reabsorbed, and the injected serum has become removed from the circulating blood, that the actual amount of absorption of corpuscles which has taken place becomes apparent.

.
 "A slight inflammatory reaction always occurs for a few hours after the injection, resulting in an effusion of serum containing leucocytes, more or less marked according to the amount of irritation. This effusion is, however, of short duration, ceasing generally in the course of the first few hours, after which the effused serum along with that of the injected blood becomes reabsorbed back into the circulation.

"The irritation produced by the presence of clots is probably of more consequence, as it certainly is longer lasting. The resulting inflammation, however, is generally localised. In no instance at least in these experiments was it such as in any way to endanger life.

.
 "It is in the neighbourhood of the female generative organs, and in connection with pathological conditions of these organs, that such extravasations most frequently occur. A few considerations only need be presented here.

"If the extravasation take place extraperitoneally, *e.g.*, between the layers of the broad ligament, as is probably the case in the great majority of instances, it is clear that most of the conditions will be present, especially as regards the more or less definite boundaries of the extravasated blood, to ensure the early coagulation of the blood, and that, too, *en masse*. As any absorption of corpuscles which may then occur can only take place through the ordinary lymphatic channels of the pelvis, through which the absorption of corpuscles as such is but slight, by far the greater proportion of the corpuscles will thus be doomed to a local fate.

"If, on the other hand, the effusion of blood occur not only extraperitoneally, but also in part into the peritoneal cavity itself, as is probably not unfrequently the case, the ultimate fate of the blood may be different. Its coagulation may then be more or less delayed, and its absorption greatly facilitated by the special action of the diaphragm in promoting absorption.

"The distribution of the blood in such cases will naturally be, in the first instance at least, in the neighbourhood of the pelvic organs, although the peristaltic action of the intestines will tend to distribute it more or less amidst the coils of intestine. However clear may be the part played by the diaphragm in absorption in the case of animals, in whom the quantity of blood injected, relative to the size of the abdomen, is so great, the case is otherwise in the human subject, where the quantity of blood, relative to the size of the abdomen, may be very small, and the blood itself is generally situated at that part of the abdomen most distant from the diaphragm. It became of interest, therefore, to determine what part the diaphragm played in the absorption of small quantities of fluid.

"In two of my experiments on rabbits, in which death took place within a period of 24-36 hours after the injection, the inflammation was observed to be most intense over the under surface of the diaphragm and upper surface of the liver, these surfaces being covered with a thickish layer of fibrinous lymph, with, at parts, larger nodules of fibrin

and leucocytes. It seemed as if the septic poison introduced had acted most virulently at the seat of its absorption. It has already been seen that it was in this neighbourhood that fluid blood was always found most abundant, if examination were made shortly after its injection." ¹

It is of the highest pathological importance to note that in a very large proportion of the cases diseased ovaries have been found ; changes




FIG. 119.

HÆMATOMA FELT AS A RETRO-UTERINE TUMOUR IN CASE OF EXTRA-UTERINE GESTATION IN RIGHT BROAD LIGAMENT (*Hart and Carter*).

in the Fallopian tubes (dilatation and filling with blood or pus) being less common.

The effused blood undergoes changes in course of time ; so that blood crystals, granular corpuscles, and oil drops are found as traces of the

¹ Loc. cit., pp. 461-465.

previous blood effusion. In most cases of recovery, it becomes entirely absorbed. As the result of abdominal section for ruptured Fallopian-tube gestation, it has been noted that the effused blood becomes increased in specific gravity and stains sponges deeply.

In the extra-peritoneal effusions, the fate of the extravasated blood is to a great extent local. The blood-clot is formed into connective tissue, and large areas of blood crystals are found.

The practical deduction from all this is that in intra-peritoneal Prognosis. effusions the majority of cases can be tided over until the effused blood is absorbed. Ruptured Fallopian-tube gestations require abdominal section in most instances. In extra-peritoneal effusions the immediate




FIG. 120.

RETRO-UTERINE HÆMATOCELE. Pouch of Douglas not previously obliterated (*Schroeder*).

prognosis is much less grave, and ultimate recovery usually takes place.

ETIOLOGY: SOURCES OF HÆMORRHAGE AND VARIETIES.

The table quoted below shows that pelvic hæmatocele is most common Etiology. in women between the ages of 25 and 35—that is, women in their period of full menstrual and sexual vigour. Out of 43 cases, the ages, according to *Schroeder*, were as follows:—

In	3 cases, or 7.0 p. c., the ages were . . .	22-25
„ 14	„ 32.5 „ „ . . .	25-30
„ 13	„ 30.2 „ „ . . .	30-35
„ 9	„ 20.9 „ „ . . .	35-40
„ 3	„ 7.0 „ „ . . .	40-43
„ 1	„ 2.2 „ „ . . .	53

It is more common in parous women ; there is considerable difference of opinion as to its frequency, Olshausen placing it as high as 4 p. c. of all female diseases, while Schroeder estimates it only at .7 p. c.

The following are the chief causes of hæmorrhage and its anatomical sources.

1. *Predisposing causes.* Profuse menstruation ; violent exercise during menstruation, such as dancing ; violent coitus during menstruation ; varicose conditions of the subperitoneal veins ; purpura ; scorbutus ; hæmophilia.

2. *Anatomical sources.* (a.) *Pelvic Peritoneum.*—There may be rupture of veins of the pampiniform plexus, or of the veins below the uterine




FIG. 121.

COPIOUS BLOOD-EFFUSION ANTE- AND RETRO-UTERINE.

peritoneum. In the former case, we may get the blood pouring directly into the peritoneal cavity ; or first passing between the layers of the broad ligament, and either remaining enclosed there or rupturing into the peritoneal cavity. The hæmorrhage, according to Virchow, may arise from vessels developed in the false membranes of pelvic peritonitis. Credé of Leipzig quotes a case where he tapped a tumour and first got serum, then blood-stained serum, and finally blood. In two days, a fresh tapping gave first putrid blood and then fresh blood in abundance.

(b.) *Connective tissue.*—Rupture of veins occurs here also.

(c.) *Uterus.*—We may have regurgitation in menorrhagia from the uterus along the dilated Fallopian tubes. Rupture of interstitial extra-uterine pregnancy is another cause of hæmorrhage.

(d.) *Fallopian tube*.—Blood may come from its hyperæmic mucous membrane and pass into the peritoneal cavity. Intra-peritoneal hæmatocele is often the result of the rupture of an early Fallopian-tube gestation intra-peritoneally, or we may have the condition known as tubal abortion causing it. By this we mean that an early tubal




FIG. 122.

RECENT HÆMATOCELE IN POUCH OF DOUGLAS from rupture of a gestation-sac lying in it; the uterus, the cavity of which is not cut into, is closely incorporated with anterior wall of sac (*Barbour*).

pregnancy has been converted into a "mole" by hæmorrhage, and expelled or partially expelled with intraperitoneal hæmorrhage into the peritoneal cavity. When it develops between the layers of the broad ligament, hæmatoma is the result.

(e.) *Ovary*.—Here it results from rupture of congested vessels, or of the Graafian follicles.

Of all these causes, rupture of veins below the peritoneum, and rupture of Fallopian-tube pregnancies are the most common. The student will now clearly see the *symptomatic nature* of hæmatocele.

SYMPTOMS.

Symptoms. The chief symptoms are menorrhagia, sudden onset, sudden bloodlessness, pain. The pulse may become feeble from anæmia, and the temperature is not above normal. Menorrhagia is not always present, and the bloodlessness may not be very well marked; sometimes patients have a sudden faint feeling. In cases of copious effusion from rupture of an extra-uterine pregnancy, the symptoms are often like those of irritant poisoning: viz., sudden onset, prostration, vomiting. The marked anæmia, however, points to some internal hæmorrhage; inquiry should then be made as to menstruation, and this always followed by bimanual examination. In Fallopian-tube gestation the decidua may be discharged from the uterus before actual rupture.

In retro-uterine hæmatocele, we find frequent painful micturition and difficulty in evacuation of the bowels. There is sometimes retention of urine.

PHYSICAL SIGNS.

**Physical
Signs.**

These differ according as the effusion is intra- or extra-peritoneal.

Intra-peritoneal Hæmatocele.—When blood is poured out near the pouch of Douglas, and below adhesions, we may get the following. On abdominal percussion, dulness may be present. On vaginal examination, a resistant bulging tumour is felt, varying in size from that of a billiard ball to that of a child's head, and sometimes filling up a large part of the pelvic cavity; the os uteri is pressed close behind the symphysis, looks downward, and is often almost inaccessible (figs. 120 and 122). A good plan to get at it is to turn the index finger palmar surface to the symphysis, and push it well up. On bimanual examination, the fundus uteri is felt unusually distinct, beneath the abdominal walls and above the pubes, and generally to one or other side. This settles the point that the tumour is retro-uterine and not the uterus. Considerable bleeding into the peritoneal cavity gives dulness on percussion but no bulging of the fornices.

Extra-peritoneal Hæmatocele: Pelvic Hæmatoma.—When the blood-effusion is poured out between the layers of one of the broad ligaments, we get displacement of the uterus to the opposite side, arched dulness on abdominal percussion to one or other side of the hypogastric region with bulging more or less marked in the lateral or posterior fornices (fig. 119). When the effusion is peri-uterine, we get the abdominal

dulness more extensive and the bulging in the fornices all round the uterus. Pelvic peritonitis is often a result of the intra-peritoneal form of blood effusion.

All that has been given here is only how to recognise intra-pelvic hæmorrhage, which is merely a symptom or sign of some lesion. The diagnosis of the lesion causing the hæmorrhage is, except in the case of extra-uterine pregnancy, as yet beyond our clinical knowledge.

DIAGNOSIS AND DIFFERENTIAL DIAGNOSIS.

Pelvic hæmatocele requires to be diagnosed from—

<i>Pelvic peritonitis followed by enclosed serous effusion in pouch of Douglas,</i>	Diagnosis and Differential Diagnosis.
<i>Pelvic cellulitis,</i>	
<i>Fibroid on posterior wall of uterus,</i>	
<i>Ovarian cyst behind uterus,</i>	
<i>Extra-uterine pregnancy,</i>	
<i>Retention of blood in horn of a malformed uterus,</i>	
<i>Retroversion of non-gravid or gravid uterus.</i>	

Of these we consider at present only pelvic peritonitis and pelvic cellulitis. The others will be treated of each under its respective head.

In these two purely inflammatory affections we have the inflammatory symptoms from the first; without a history of sudden onset, of menorrhagia, or of the symptoms of internal hæmorrhage. Further, the difference in etiology of hæmatocele and peritonitis will help us. The history is the most important aid in diagnosis.

When we have the rare form of hæmatoma due to blood effusion between anterior rectal wall and posterior vaginal wall there is great straining, and marked swelling felt *per rectum*.

COURSE AND RESULTS.

In many cases ($\frac{4}{5}$ according to Voisin) the blood effused becomes entirely absorbed, in a time varying from 2 to 10 months.

The tumour, with partially clotted or purulent contents, may burst into the rectum, vagina, or peritoneal cavity; in the last case, fatal peritonitis may follow.

When the blood effusion is very large, death may be rapid.

PROGNOSIS.

As to life.—This is, as a rule, settled soon. The most fatal cases are extra-uterine pregnancies, and those in which there are no peritonitic adhesions to limit the blood effusion. After peritonitis is set up, the prognosis is much as in pelvic peritonitis.

TREATMENT.

(1.) *At onset of hæmorrhage.*(2.) *When suppuration occurs.*

Treatment. (1.) *At onset of hæmorrhage.*—The treatment here is expectant. The patient is to be put at complete rest, with ice bags to the abdomen. Ergotine should be injected into the buttock. The ice-bag is to be kept on for several days, as this will limit the subsequent peritonitis. If the patient is collapsed, then stimulants and hypodermic injections of sulphuric ether or whisky must be freely used; a large mustard poultice over the abdomen is often serviceable, both as a blood derivative and in allaying vomiting.

In most cases, the source of the bleeding is unknown; the present state of knowledge does not enable us to lay down any rule as to the opening of the abdominal cavity and the attempt to ascertain and secure the bleeding source. In Fallopian-tube pregnancies which have burst, however, the abdomen has been opened and the tube ligatured on either side of the rupture; Lawson Tait has operated successfully on forty-two cases of rupture of Fallopian-tube gestation, but always at some period after the rupture. Sinclair, Herman, and Berry Hart in this country, Johnstone in America, and many others, have also operated successfully, and the justifiability of this treatment is now fully admitted.

Martin has performed laparotomy in four cases successfully. He opens the abdomen, incises the sac, clears out clots, ties vessels, and drains. When possible, the opening of the blood sac should be stitched to the abdominal wound. Imlach of Liverpool has also recorded cases where he opened the abdomen and tied the Fallopian tubes along which blood had regurgitated. Accordingly, we may now look forward to an extension of more active interference by abdominal section. Zweifel has in several cases incised the tumour per vaginam, turned out the clots and drained the cavity. In Hæmatoma, when absorption is very slow, Gusserow has had good results by incising through the vagina, washing out, and draining. While absorption is going on, the treatment is the same as in pelvic peritonitis.

(2.) *After suppuration has occurred.*—The tumour is to be opened and drained, as recommended at p. 197 for suppurating pelvic cellulitis.

Recently, Lawson Tait has recommended that some pelvic abscesses be opened by abdominal section, as we often get very tedious cases when they perforate into the bowel. The following was the treatment in one of six cases in which he performed it. "I determined to open it from above. . . . I found a large cavity containing about two pints of fœtid pus with decomposing blood-clots. This I carefully cleansed out, and after having united the edges of the opening into the

cyst carefully to the abdominal wound, I fixed in one of Kœberlé's drainage tubes five inches long. . . . The patient went home cured on the thirtieth day." Tait's cases were chiefly suppurating hæmatoceles (*Tr. of Lond. Roy. Med. and Chir. Soc., vol. lxi.*).

NEW GROWTHS IN THE PELVIC PERITONEUM AND CONNECTIVE TISSUE (BROAD AND ROUND LIGAMENTS).

LITERATURE.

PERITONEUM AND CONNECTIVE TISSUE: BROAD LIGAMENT. *Cobbold, W. A.*—Parasites: Churchill, 1879. *Freund, W. A.*—Das Bindegewebe im weiblichen Becken und seine pathologischen Veränderungen mit besonderer Berücksichtigung der parametritis chronica atrophicans und der Echinococcuskrankheit: Gynäkologische Klinik, Strassburg, 1885. *Freund and Chadwick*—Four cases of Echinococcus in the Female Pelvis: *Am. J. of Obstet.*, 1874. *Hart, D. Berry*—Sarcoma of pelvic connective tissue in *Atlas of Female Pelvic Anatomy*: Edinburgh, 1884. *Neisser*—Die Echinococcuskrankheit: Berlin, 1877. *Schatz*—Die Echinokokken der Genitalien und des kleinen Beckens beim Weibe: *Cent. f. Gyn.*, 44, 1886. *Simmons*—Malignant disease of female Sexual Organs: *Edin. Med. Jour.*, Dec. 1885. Cobbold and Neisser give the literature on Echinococci fully and references to cases by Hewitt, Corrigan, Turner, Wynne, and others.

ROUND LIGAMENT.—*Goodell*—Lessons in Gynecology: Philadelphia, 1880. *Sänger*—Weitere Beiträge zur Lehre von den primären desmoiden Geschwülsten der Gebärmutterbänder, besonders der Ligamenta rotunda: *Archiv f. Gyn.*, XXI. 279 and XXIV. 1. *Schroeder*—Krankheiten der weiblichen Geschlechtsorgane: Leipzig, 1878, S. 417. *Thomas*—Diseases of Women: Philadelphia, 1880, p. 136. *Wile*—Hydrocele in the Female: *Am. J. of Obst.*, July 1881, which see for further literature. *Winckel*—Lehrbuch der Frauenkrankheiten: Stuttgart, 1886.

TUMOURS OF THE BROAD LIGAMENT.

Hæmatoma and inflammatory conditions of the broad ligament have been already considered. We need only further mention that we may have cysts, fibroids (rare), phleboliths, cancer, and tuberculosis; the last two are only parts of the general peritoneal affection. Ovarian cysts may develop into the Broad Ligament, and cysts may develop in the Broad Ligament independently of the Parovarium. Cysts of the Broad Ligament will be considered along with Ovarian Tumours.

HYDROCELE OF THE ROUND LIGAMENT.

Nature and Pathological Anatomy.—This is a rare malady, and may exist as encysted fluid about the round ligament (extra-peritoneal), or in the canal of Nuck—a process of peritoneum extending from the internal inguinal ring in the labium majus. It may be closed at the internal ring, thus forming a cyst; or it may communicate with the peritoneal cavity.

The fluid is serous in its nature; it may be olive green in colour.

Physical Signs.—(a) *Of encysted hydrocele of the cord.* An oval trans-

lucent swelling exists in the inguinal canal. It cannot be returned into the abdominal cavity, has usually existed for some time, is not tender on pressure, and gives rise to no symptoms. It must be differentiated from an ovary in the inguinal canal, and from incarcerated hernia.

(b) *Of hydrocele in the labium majus.* The labium majus is distended with a fluctuating tumour, dull on percussion and of translucent appearance; usually, the contents cannot be returned into the abdominal cavity. Aspiration gives a clear fluid. It is to be diagnosed from hernia in the usual way.

Treatment.—Aspiration and drainage; or aspiration and injection of a few drops of tincture of iodine. Goodell recommends that when the labial form communicates with the abdominal cavity, the internal ring should first be firmly compressed and the injected fluid then sucked out.

TUMOURS OF THE ROUND LIGAMENT.

Tumours
of Round
Ligament.

Fibrous, myomatous, sarcomatous tumours, and their combinations, have been described in the round ligament by Säger. They may develop in any part of its course: intra-peritoneally; within the inguinal canal; or extra-peritoneally—in the abdominal wall, the pelvic cellular tissue or the labia majora. Such tumours are rare, those of the third group (extra-peritoneal) being the most frequent. They may be removed unless dipping down into the pelvis.

ECHINOCOCCI IN THE PELVIC ORGANS.

Echino-
cocci in
Pelvic
Organs.

Echinococci or Hydatids are the sexually immature forms of the *Tænia echinococcus*, a small tapeworm found in the intestines of the dog. When present in the human body, they form elastic tumours and may occur in the female pelvic organs.

Freund, in 25 years, met with 19 cases—of which 7 were in the pelvic connective tissue: while Schatz met with 6 out of 7000 gynecological and obstetric cases (1 in 1166). Schatz has also collected 66 cases of *Echinococcus* disease in the female pelvic organs and found the frequency as follows: 14 in uterus, 14 at pelvic brim, 10 in Douglas' pouch, 7 in ovary, 7 in broad ligament, 7 in pelvic connective tissue, 5 between rectum and vagina, 2 between bladder and vagina.

They may remain many years without symptoms or may perforate into bowel or bladder. When large, they cause pressure symptoms on bladder and rectum. The *physical signs* are those of a tense elastic tumour without pain; at first, usually situated near the rectum; and ultimately, when increased in size, displacing the pelvic organs as an ovarian tumour would when developing between the layers of the broad ligament, *i.e.*, first forwards and then upwards. The *diagnosis* is

often difficult and tapping may be requisite. When they project sufficiently into the abdomen, the treatment is laparotomy with shelling out of the tumour; or incision of the sac, with careful cleansing and stitching the edges to the abdominal incision (*v.* Abdominal Section in Appendix). When pelvic, the sac is opened and drained (*v.* pp. 197, 198). Hydatids are rare in this country, but common in Iceland and Australia (Cobbold).

TUMOURS OF THE PELVIC CONNECTIVE TISSUE.

We may have fibromyomata, sarcomata, or dermoid cysts as primary conditions in the pelvic connective tissue.

Tumours
of Pelvic
Connective
Tissue.




FIG. 123.

SARCOMATOUS TUMOUR OF THE PELVIC CONNECTIVE TISSUE (*Hart*).

A Tumour, B Uterus, Bl Bladder, Ov Ovary, c c Inguinal, and c' c' Sacral Lymphatic Glands.

Fig. 123 shows the pelvis from an interesting case of primary sarcoma which began in the connective tissue at the left side of the uterus and spread through the lymphatic glands. This case presented the following points of interest.

A. B., æt. twenty-seven, was an undersized, wretchedly thin girl, who had felt unable for her usual occupation of a domestic servant; but the medical man whom she had consulted had been unable at first to find anything tangible to account for her condition. Afterwards, however, the inguinal glands of the left groin (those parallel to Poupart's ligament) began to be enlarged, and the left leg was painful and somewhat swollen. In

CHAPTER XIX.

PATHOLOGY OF FALLOPIAN TUBE AND PAROVARIUM.

LITERATURE.

Ballantyne and Williams—The Pathology and Histology of the Fallopian Tubes : Brit. Med. Jour., 1891. *Bandl*—Die Krankheiten der Tuben, etc., Billroth's Handbuch : Stuttgart, 1879. *Barnes*—Diseases of Women : London, 1878, p. 376. *Bumm*—Über die Bedeutung der gonorrhöischen Infection für die Entstehung schwerer Genitalaffectionen der Frau. Verh. der deutsch. Gesell. für Gynäk., Leipzig, 1892. *Cullingworth*—Clinical Illustrations of the Diseases of the Fallopian Tubes : 2nd edit. with many Drawings : Rivington, Percival & Co., London, 1896. *Doran*—Clinical and Pathological Observations on Tumours of the Ovary, Fallopian Tube, and Broad Ligament : London, 1884, v. also Trans. Path. Soc., Lond., 1887 ; and Trans. of Lond. Obst. Soc., 1887 : also his article in Playfair and Allbutt's System of Gynecology. *Fowler, J. K.*—Hydro- and Pyo-Salpinx : Med. Times and Gazette, 1884. *Freund*—Ueber die Indikationen zur operativen Behandlung der erkrankten Tuben : Volk. Samml., No. 323. *Griffiths*—Tubo-ovarian Cysts : Trans. Lond. Obst. Soc., 1887. *Hennig*—Krankheiten der Eileiter und die Tubarschwangerschaft : Stuttgart, 1876. *Klob*—Pathologische Anatomie der weiblichen Sexualorgane : Wien, 1874. *Lewers*—On the Frequency of Pathological Conditions of the Fallopian Tubes : Trans. of Lond. Obst. Soc., 1887. *Martin, A.*—Die Krankheiten der Eileiter : Leipzig, 1895. The various sections are taken by *Martin, Wendeler, Kossmann, Orthmann, Sänger* and *Barth*. This is one of the most important works that has recently appeared on this subject. *Noeggerath*—Die latente Gonorrhoe im weiblichen Geschlecht, VIII., p. 123 : Bonn, Cohen & Sohn, 1872. *Pinner*—Über den Übertritt des Eies aus dem Ovarium in die Tube beim Säugethier. Arch. für. Phys., 1880. *Savinoff*—Ein Fall von Salpingitis chronica productiva vegetans. Arch. für Gynäk., 34, 1889. *Schroeder*—Handbuch der Krankheiten, etc. : Leipzig, 1878, S. 329. *Simpson, Sir J. Y.*—Diseases of Women (Edit. A. R. Simpson), p. 539. *Stewart, Sir T. G. S. and R. Muir.*—Notes of a Case of Actinomycosis of Ovaries and Liver : Edin. Hospital Reports, Vol. I., 1893. *Stratz*—Gynäkologische anatomie : Berlin, H. Kornfeld, 1892. *Sutton, J. Bland*—Surgical Diseases of the Ovaries and Fallopian Tubes : Cassell & Co., 1896. *Tait*—Menstrual Fluid retained in the Left Fallopian Tube simulating Ovarian Tumour : Br. Med. J., 1878, p. 677. The Pathology and Treatment of Diseases of the Ovaries : Birmingham, 1883. Note on Chronic Inflammatory Disease of the Uterine Appendages : Ed. Med. Jour., Sept. 1885. *Wells, Sir T. S.*—Diseases of Ovarian and Uterine Tumours : London, 1882. *Wertheim*—Die ascendierende Gonorrhoea beim Weibe : Arch. für Gynäk., 1892. *Williams*—Ovarian Tumours : Reynolds' System of Medicine, Vol. V. *Winckel*—Die Pathologie der weiblichen Sexualorgane : Leipzig, 1881. *Wylie, W. G.*—Operations for Salpingitis : N. Y. Medical Record, Aug. 29th, 1885. *Zemann*—Ueber die Aktinomykose des Bauchfells und der Baueingeweide beim Menschen : Medicin. Jahrbücher der K. K. Gesellschaft der Aertzte in Wien, 1883, S. 477. For other literature see *Bandl, Doran, and Martin*, whose works and that of *Hennig* we mainly follow.



Fig. 1.




Fig. 2.



Fig. 3.





Fig. 5.



Fig. 4.



MUCOUS MEMBRANE OF FALLOPIAN TUBE,
NORMAL AND DISEASED (Martin).

- FIG. 1. Foetal tube, showing sinuities.
 FIG. 2. Normal mucous membrane.
 FIG. 3. Mucous membrane in chronic catarrhal salpingitis.
 FIG. 4. " " pseudo-follicular "
 FIG. 5. " " purulent "
 m. Muscular wall. n. Mucous membrane.

FALLOPIAN TUBE.

Preliminary Considerations.—Our present knowledge of tubal disease is practically of less than twenty years' growth. While attention had been drawn to the connection between inflammatory and uterine disease by several authors (E. Martin, Buhl, 1859, Hennig), and while Cruveilhier's famous atlas (1843) pictured conditions to which both the obstetrician and gynecologist obstinately shut their eyes, the importance of tubal conditions did not practically influence Gynecology until the abdominal sections, initiated mainly by the bold theories of Battey and Tait, found in the living woman conditions of the tubes which had never been dreamt of. Before this epoch the gynecologist occupied himself in trying to determine whether the inflammatory swellings he discovered clinically were peritonitic or cellulitic, but there was no active operative treatment, whichever he determined them to be.

Noeggerath did most excellent service by his speculation as to latent gonorrhœa in the male being the cause of much pelvic disease in women, and in this he has been followed by various observers, A. MacDonald, Sânger, W. J. Sinclair, and many others.

While the discovery of tubal disease has come late into gynecology it must be kept in mind that gynecology itself is not much older than half a century, and it is really only now that such conditions could be fairly and fully understood. It must not, indeed, be forgotten that without the knowledge we now possess of septic infection, of the micro-organismal nature of gonorrhœa and of tubercle, Noeggerath's theory would have remained a speculation, and probably a much derided one. Thanks, however, to many factors, to the clinical work of the operative gynecologist, to improved microscopical technique, and, above all, to Bacteriology, we are now in a position to investigate and treat a series of diseases too long unknown. The general causation of acute and chronic inflammatory tubal disease has now been shown to be due, in the main, to micro-organisms, of which the chief are gonococci (for gonorrhœa); streptococci (for septic cases): the contents of the tube may be sterile. In 206 cases Wertheim found the gonococcus in 56 cases; streptococci in 11; staphylococci in 6; pneumococcus lanceolatus in 1; and in 122 instances, the results of bacteriological examination, were negative.

Anatomy and Physiology of the Tube.—This has been already fully considered, and we here discuss only a few points.

The tubal lining is a much folded one (*v.* Plate V., fig. 2), and these folds are exactly analogous to those in the uterine mucous membrane proper. The question therefore arises as to whether we may consider them as glands. The opinion that they are so, is held by

Preliminary
Considerations.

Anatomy
and Physi-
ology of
Tubes.

Bland Sutton, who also alleges that they furnish an albuminous covering to the ovum during its descent. At present, however, it is best to consider it as a much folded mucous membrane, and to defer the question as to its real nature until sufficient facts have been collected.

The tube probably does not take any part in menstruation, *i.e.*, it furnishes neither blood nor epithelium during that period. The real and effective source of the menstrual discharge is the triangular uterine area, bounded above by the internal openings of the tubes, and below by the *os internum*. The student may, however, ask how the occasional existence of tubes distended with blood can be explained if it is not due to retained menstrual fluid. The answer to this is furnished, however, by the occurrence of other pathological conditions in the tubes, *viz.*, early tubal gestation and atresia vaginæ.

Function
of Tubes.

The main *functions* of the tubes are :—

(1) To act as a duct between ovary and uterus; and (2) to act as a canal for the passage of the spermatozoa after these have been placed in the genital tract by coitus.

(1) As we have already seen, an ovum is periodically shed at or near a menstrual period, and is carried along the tube to the uterus. How this is accomplished is much disputed. The cilia of the tube and uterus work towards the *os externum*, and it is evident that we have thus a mechanism for the transport of the ovum which cannot be gainsaid. The experimental observations of Pinner, who injected salt solution with corpuscular elements into the peritoneal cavity of rabbits and found this fluid in the lower genital tract, are of importance, although it may be urged that this might be a mere oozing into the tract due to increased intra-abdominal pressure. The ova of *Ascaris lumbricoides* were injected by Lode, and although these are only half the size of the rabbits' ova they were detected in the tube some hours afterwards. Ciliary action, therefore, seems the main factor in carrying on the ovum, and we must assign a very minor part, if any at all, to the tubal peristalsis itself, or to the alleged actual grasping of the ovary by the fimbriated end of the tube at the time of the escape of the ovum.

(2) The passage of the spermatozoa along the tube is undoubtedly effected in the main by their independent movements, and it is stated that they can progress 1 to 6 mm. in one minute.

The question has been much disputed as to where the single spermatozoon necessary for fertilisation meets the ovum, whether in the tube or in the uterine cavity. It is impossible to determine the question accurately in the human female. A much more important point is the site of grafting of the impregnated ovum. As a matter of fact this normally occurs within a certain definite area, which may be

termed the "area of safe implantation." The three points determining this area are the inner openings of the Fallopian tubes and the os internum. This also determines the area from which the epithelium is shed at a menstrual period, and establishes a causal relation between menstruation and the area of safe grafting of the ovum, not yet quite thoroughly understood.

When the ovum grafts in the tube, *i.e.*, outside the internal opening, we then get the abnormal early gestation known as tubal gestation.

The *route* of infective action is of great importance. This may be by the epithelial lining of the tube but more often it is by the lymphatics of the tubal and uterine tissue. From the abundance of the lymphatic supply and its free anastomosis it can be readily understood that we do not get a purely tubal infective condition, but a combination in which mainly tube and ovary take part—the so-called Salpingo-oophoritis, *i.e.*, an inflammatory condition affecting and matting together tube and ovary.

It is through the tubes that the pelvic peritoneum is also in many cases infected, *i.e.*, pelvic peritonitis is usually secondary to tubal disease.

CLASSIFICATION OF TUBAL DISEASES.

The classification of tubal diseases is admittedly difficult, but the most scientific yet brought forward is that of Martin in his valuable work; this we follow with only some slight modification. The diseases are grouped as follows:—

- (1) Abnormalities.
- (2) Sinuosity, stricture, and occlusion.
- (3) Patent tube.
- (4) Catarrhal conditions, without distention, comprising—
 - (a) Acute endosalpingitis;
 - (b) Chronic diffuse interstitial salpingitis, including salpingitis isthmica nodosa, and salpingitis chronica productiva vegetans.
- (5) Catarrhal conditions associated with occlusion and leading to distention; hydrosalpinx.
- (6) Purulent salpingitis; pyosalpinx.
- (7) Infectious granulomata—syphilitic, tubercular, actinomycotic.
- (8) Hæmatosalpinx.
- (9) New formations and tubo-ovarian cysts.

We shall first consider methods of examination, then the pathological conditions, in the present chapter, and take up the *Diagnosis, Prognosis, and Treatment* in Chapter XX.

Bimanual Examination of Tubes.

Bimanual
Examina-
tion of
Tubes.

Can the normal Fallopian tubes be palpated in the Bimanual? The student will probably have already noted that, in considering the Bimanual (Chap. VIII.), we did not name the Fallopian tubes as structures whose form and limits he was expected to define. In a very favourable case, the conjoined manipulation may recognise them at their uterine origin, more especially if the rectal examination be made and the uterus be well drawn down with the volsella. Practically, the Fallopian tubes (unless dilated) are not palpable on ordinary examination. It must not be forgotten that many cases have now been recorded, where abdominal section showed the Fallopian tubes to be dilated with pus to the size of coils of small intestine, although a careful Bimanual had failed to detect their presence.

When the tube is not much distended its inner relation to the uterus can be well mapped out both on vaginal and rectal examination.

Catheterisation of the tubes.—In certain undoubted cases the uterine sound has been passed along the Fallopian tube, while in others the supposed sounding of the tube has been really the perforation of the uterine wall. It is impracticable to sound the normal Fallopian tubes to any effect; and the procedure is by no means devoid of danger.

(1) ABNORMALITIES.

Abnormal-
ities.

These are of little practical interest. The chief are, an accessory fimbriated end; defective development; displacement; want of apposition of fimbriæ and accessory tubes to ovary (Lawson Tait). The hydatid of Morgagni (fig. 124, 8) is attached to the tube, and arises from the duct of Müller. It is lined by ciliated epithelium, has unstriped muscle in its walls, and in its stalk always.

(2) SINUOSITY, STRICTURE, AND OCCLUSION OF THE TUBES.

Sinuosity,
Stricture,
and Occlu-
sion of
Tubes.

The tube may be unduly sinuous, a persistence of a foetal condition (Freund). (Plate V., fig. 1).

The tube may have a congenital stricture; or may become closed at the uterine or the fimbriated end, or in the middle. When stricture occurs at the uterine end, it is said to be caused by implantation of the placenta there or by endometritis with adhesion. In the middle, small tumours or adhesions may cause strictures—in the latter case usually partial. At the fimbriated end, the occlusion is due to a catarrh of the tubes which has spread to the peritoneum and set up adhesive peritonitis.

These strictures are of importance in relation to sterility and fluid

accumulations (pus, serum, blood) which they favour ; but in themselves cannot be diagnosed during life.

(3) PATENT CONDITION OF THE TUBES.

By this is meant undue dilatibility. It is of great importance in relation to uterine injections. Even in careful injection of the uterine cavity, post partum or otherwise, fatal results have followed from the fluid's passing along the tube into the peritoneal cavity. "Forcible uterine injections on the cadaver, with the cervix entirely filled up by the syringe, almost always sent fluid along the tubes into the peritoneal cavity. Less forcible injections under like conditions sent the fluid along a less distance (2-3 mm.), and often sent it into the veins ; while gentle injections with a tube not filling the cervical canal sent fluid neither into the tubes nor veins." Bandl, from whom the above is taken, records a case where death resulted from injection of an aborting uterus with perchloride of iron, although the injection pipe was less in diameter than the cervix. Death may be immediate from shock, or some days after from peritonitis. In uterine injections, no more than 1-4 drops should be used.

Winckel has recorded a unique case where a round worm (*Ascaris Lumbricoides*) was found calcified on the posterior surface of the uterus and left broad ligament. It had probably passed from the anus into the vagina and ultimately through the Fallopian tube into the peritoneal cavity.

(4) CATARRHAL CONDITIONS WITHOUT DISTENTION.

This is the most common inflammatory condition of the tubes, and forms about 30 per cent. of tubal cases.

Simple catarrh of the lining of the tube (acute endosalpingitis) is rarely primary, and is, as a rule, secondary to uterine or peritoneal conditions.

The most common cause is certainly micro-organismal—gonorrhoeal and septic ; while as minor causes we may put down irritating injections of iodine, unskilful manipulation of the organs during examination, and the careless use of the unpurified uterine sound or of intra-uterine pessaries.

The infectious fevers (scarlet fever, measles, cholera, etc.) have also some place in producing tubal inflammatory conditions.

The tubes are congested, more tortuous, and their secretion contains white and red blood corpuscles and mucus. According to Hennig, Hyalin, Mucin, and Colloid can be detected in this order of frequency.

The foldings of the mucous membrane are swollen and reddened (Pl. V., fig. 3), the chief changes being in the connective tissue ;

the alterations in the epithelium are less marked, but it may become flattened in character as the change tends to become chronic.

In the *chronic catarrhal conditions* we get small-celled infiltration of the foldings, thickening of the ends of these, loss of their epithelium and a blending of the folds (Pl. V., fig. 3). In this way the recesses of the mucous membrane get snared off, retain their secretion, and a form known as salpingitis pseudo-follicularis may thus arise (Pl. V., fig. 4). When the muscular walls of the tube become involved we have an interstitial salpingitis produced where the connective tissue and muscular fibre are hypertrophied; this has been described as salpingitis chronica productiva vegetans by Sawinoff. The tubes are thickened, and radiating pain is a marked symptom.

A rare form of tubal disease (1 per cent. of bodies examined according to Chiari of Prag) has been termed salpingitis isthmica nodosa. It affects the uterine end of the tube, and manifests itself as small projections, like tiny myomata, consisting, however, of cysts lined with cylindrical epithelium, and having hypertrophied muscular walls. They are the result of catarrh of the lining.

(5) CATARRHAL CONDITIONS ASSOCIATED WITH OCCLUSION AND LEADING TO HYDROSALPINX.

Catarrhal Conditions with Occlusion. *Hydrosalpinx*.—By this we understand a tube dilated with serum, the walls being thinned and translucent. The size of the tube may be considerable, sometimes larger than a child's head. It is the outer end which is distended, and there are usually omental and intestinal adhesions. The muscular walls are thin, the epithelium cuboidal, and there may be calcification of the walls. This is termed *hydrosalpinx simplex*; when the uterine end is patent and allows escape of fluid we have *hydrops tubæ profluens*; there occurs further a hæmorrhagic form, and also, though rarely, an emphysematous variety.

Instead of the term hydrosalpinx, that of sactosalpinx serosa has been proposed by Martin of Berlin (σακτο = tensely filled sac).

The etiology of hydrosalpinx is probably a catarrhal micro-organismal endosalpingitis to begin with, occlusion of tube, followed by marked secretion of fluid and secondary thinning of walls.

(6) PURULENT SALPINGITIS—PYOSALPINX.

Purulent Salpingitis. Purulent salpingitis. Pyosalpinx. Sactosalpinx purulenta (Martin). The catarrhal form may become purulent and chronic, and when occlusion of the tube at the fimbriated and uterine ends takes place, we get the secreted pus retained and a retort-shaped tumour formed, the important and well-known pyosalpinx.

Naked-eye appearances.—The tube is distended mainly at the outer end, and its shape is thus aptly compared to a retort. The size of the tube may vary from that of the thumb to a thickness equal to the fore arm. On longitudinal section of a pyosalpinx, intercommunicating loculi can be noted and the walls found to be thickened, although they are sometimes thinned. Adhesions may form to adjacent viscera, and the tubes may blend together at their ends (fig. 125).

Causation.—On clinical data the causation of purulent salpingitis will be found to be mainly gonorrhœal, and septic after labour; and the same holds good probably for pyosalpinx, although the bacteriological evidence may be less precise.

The appended tables give the etiology of purulent salpingitis on clinical data, and on bacteriological examination:—

AUTHOR.	Gonorrhœa.	Septic Salp.		Tubercular.	Syphilis.	With Uterine Catarrh.	?	TOTAL.
		Septic.	Non-Septic.					
Martin	198	283	—	17	10	1258	—	1766
L. Tait	19	16	11	—	—	—	17	63
Boldt	25	30	1	2	3	24	9	94
V. Rosthorn	37	45	—	—	—	—	73	155
	279	374	12	19	13	1282	99	2078

Accurate bacteriological investigation gives the following result (Wertheim):—

AUTHOR.	Gonococci.		Streptococci.	Staphylococci.	Pneumococci.	Bact. Coli.	Undetermined Bact.	Sterile.	TOTAL.
	Pure.	Mixed.							
Wertheim	32	—	6	1	1	—	4	72	116
Frommel	—	—	—	—	1	—	—	—	1
Charnier	5	4	—	—	—	—	—	6	15
Prochownick	1	—	21		—	—	—	5	27
Schauta	23	—	15	—	—	1	—	69	108
Martin	15	6	7	—	5	2	11	63	109
	76	10	50	—	7	3	15	215	376

The negative results of examination should be noted in the second table. This, of course, does not exclude a micro-organismal cause in these.

The examination of the pus in the way already described (p. 178), and the cultivation of the organisms in various media has given valuable results.

Microscopical Anatomy of Purulent Salpingitis and Pyosalpinx.

In acute septic salpingitis, usually of puerperal origin, we have the mucous lining of the tube thickened, mainly from dilatation of the lymph spaces, which are distended with leucocytes and occasional red blood corpuscles. The secretion in the tube contains streptococci and staphylococci. As the disease spreads the muscular coat becomes infiltrated with small cells. In the gonorrhœal form the gonococci are found in the tissue of the tube (Wertheim), and not only on the columnar epithelium as Bumm asserted. In the chronic form the spaces formed by the folds of the mucous membrane become gradually obliterated by round cell infiltration, the lumen becomes lined with granulation tissue and the wall may thicken.

In pyosalpinx the purulent secretion alters, containing altered leucocytes and red blood corpuscles, and so far as micro-organisms are concerned the result of examination may be negative.

(7) THE INFECTIOUS GRANULOMATA.

Infectious
Granulo-
mata.

By these we understand syphilitic, actinomycotic, and tubercular conditions. The first two are of little practical importance; the third requires more detailed description.

Syphilitic conditions of the tubes are not diagnosable clinically. There is no specific organism for syphilis known, but gummata have been found in the adult and in the syphilitic foetus (Ballantyne and Williams).

Actinomycosis is still more rare, but Grainger Stewart and Muir have recorded its presence in the pelvic organs (ovary and tube).

Tuberculosis.—The student will note here the greater accuracy which Koch's great discovery of the *bacillus tuberculosis* has given to our knowledge on this head.

Pathological anatomy of tubercular tubal disease.—Tubercular disease of the tube is the most common form of tubercular disease of the genital tract.

It may be primary, or secondary to peritoneal tuberculosis.

The sources of infection are not quite clear, but many observers consider that it may take place through the blood, by the spread of tubercular infection from the bowel where it has been started by the ingestion of tubercular milk or meat, and by coitus.

The last source has been strongly supported by many, and a good deal can be said in its favour. The theory of this form of infection

Fimbriated end of tube

Uterine end of tube

Torn adhesions---

Fig. 1.

Uterine end of tube---

Fimbriated end of tube

Hæmorrhagic subserous cyst

Fig. 2.

Fimbriated end of tube

Uterine end of tube

Cut edge of broad ligament

Ovary

Fig. 3.

Uterine end of tube---

Miliary tubercles

Fimbriated end of tube


Ovary

Fig. 4.

PURULENT AND TUBERCULAR SALPINGITIS (CULLINGWORTH)

FIGS. 1-3. Purulent Salpingitis.

FIG. 4. Tubercular disease of tube.



is
at
in
v
lo
m
fo
tr

m
re
in
th
T
at
p
(1
fi

se
p
A
if
an

oe
Fa
m
co
th
th
th
co
Hy

is, that in men with genito-urinary phthisis, early or advanced, bacilli are placed in the female genital tract during coitus and may thus infect the woman locally. From the anatomically dense nature of the vaginal and cervical tissue we would not expect the bacilli to affect a lodgement there; menstrual processes would tend to hinder endometric affection. The tube appears to form the most favourable nidus for the growth of the bacillus, and this may explain the greater frequency of tubal tuberculosis.

The tubes themselves may vary considerably in appearance, from a mere irregularly thickened and beaded condition to considerable distention with pus and caseating masses. The peritoneum is often involved, and distinct masses thus formed. *Microscopically* we may find in the acute forms caseating contents in the tubes, with bacilli. The mucous membrane has round celled infiltration with giant cells and bacilli. In the chronic forms there is greater tendency to supuration. Whitridge Williams recognises three forms in the tubes: (1) Miliary tuberculosis; (2) Chronic diffuse tuberculosis; (3) Chronic fibroid tuberculosis.

(8) HÆMATOSALPINX (SACTOSALPINX HÆMORRHAGICA).

It may be mentioned under this head that hyperæmia and congestion of the tube may occur in cases of measles, scarlet fever, small-pox and cholera. Hæmato-salpinx.

By hæmatosalpinx we understand distention of the tube with blood. According to some the tubes furnish blood at a menstrual period, and if this very doubtful opinion be correct, we can understand that with any obstruction to its exit we may get the tube distended with blood.

More certain causes are developmental malformations leading to occlusion in uterus or vagina and damming back of blood.

Probably the most frequent cause is early disintegration of a Fallopian tube gestation leading to the condition known as tubal mole or tubal abortion. In the former case there is retention of the compressed pregnancy in the tube; in the latter it is expelled into the peritoneal cavity or may be discovered in the fimbriated end of the tube.

The size of the tube varies, the blood is often chocolate coloured, and the walls of the tube usually thickened.

Diagnosis is difficult except in cases of malformation, or where the condition of early Fallopian tube gestation is evident.

The treatment is abdominal section and removal of the tube after ligature (*v. Chap. XX.*).

(9) NEW FORMATIONS AND TUBO-OVARIAN CYSTS :

OVARIAN HYDROCELE.

New
forma-
tions.

Much information has been gained as to these within the past few years, but they are of more interest to the specialist than to the student and practitioner. We therefore briefly tabulate the conditions, referring those working up such specimens, to Säger and Barth's work in A. Martin's Handbook. The conditions recognised are :—

A. *Mucous membrane of tube.*

1. Mucous and decidual polypi.
2. Papillomata.
3. Carcinomata.
4. Deciduoma malignum.
5. Sarcomata.

B. *Muscular wall of tube.*

1. Myomata.
2. Fibromata.
3. Sarcomata.

C. *Tumours of the sub-peritoneal tissue of the tube.* Lipomata.D. *Tumours of the peritoneum covering the tube.*

Papillomata and cystic structures.

E. *Tumours of the fimbriated ends.*

Of all these the most interesting are the papillomatous conditions first described by Alban Doran, and the so-called deciduoma malignum. The latter will be discussed under Carcinoma Uteri.

Of the former there are two forms—simple and cystic. Bland Sutton considers these cases as adenomatous, in accordance with his view that the mucous membrane of the tube is glandular. Hydro-peritoneum is usually present, the fluid being apparently formed in the tube and escaping through the patent ostium abdominale.

Tubo-ovarian cysts result from adhesions between the fimbriated end of the Fallopian tube and the ovary, with degeneration of the corpora lutea of the Graafian follicles thus enclosed. The contents may be poured into the uterus along the tube.

Ovarian hydrocele.—Under this term Bland Sutton has described an interesting case often considered as a tubo-ovarian cyst or hydro-salpinx.

Where such are carefully examined, the tube, tortuous and dilated, is found to open into the cyst. It is probably a complete pouch of peritoneum surrounding the ovary as in the seal (fig. 21) and rat (Robinson), instead of the shallow ovarian fossa in which the human ovary normally lies (*v. p.* 27). The ovary may be seen in the floor of the cyst in small specimens, but in large ones becomes thinned out on the cyst wall.

Plates V. and VI. show the various pathological conditions described.

Plate V.—I. Uterus, tube, and ovary of new-born infant, to show sinuosity of tube (Martin). II. Transverse section of mucous membrane of ampulla of normal tube. III. Chronic catarrhal salpingitis; *m*, muscular coat; *n*, mucous membrane (Martin). IV. Salpingitis pseudo-follicularis: *m*, muscular coat; *n*, mucous membrane; *l*, lumen of tube (Martin).

Plate VI. shows tubercular and purulent tubes (Cullingworth).

PAROVARIIUM.

The diagram shown at fig. 124, taken from Doran's interesting and valuable work shows that the Parovarium, which is the remains of the Wolffian bodies, consists of a horizontal tube and eight or ten well-developed vertical tubes, with five or six in addition represented only




FIG. 124.

DIAGRAM OF THE STRUCTURES IN AND ADJACENT TO THE BROAD LIGAMENT (Doran).

1. Framework of the parenchyma of the ovary, seat of a simple or glandular multilocular cyst.
2. Tissue of hilum, with 3, papillomatous cyst.
4. Broad ligament cyst, independent of parovarium and Fallopian tube.
5. A similar cyst in broad ligament above the tube, but not connected with it.
6. A similar cyst developed close to 7—ovarian fimbria of tube.
8. The hydatid of Morgagni.
9. Cyst developed from horizontal tube of parovarium.
- Cysts 4, 5, 6, 8, and 9 are always lined internally with a simple layer of endothelium.
10. The parovarium; the dotted lines represent the inner portion, always more or less obsolete in the adult.
11. A small cyst developed from a vertical tube; cysts that have this origin, or that spring from the obsolete portion, have a lining of cubical or ciliated epithelium, and tend to develop papillomatous growths, as do cysts in 2—tissue of the hilum.
12. The canal of Gartner, often persistent in the adult as a fibrous cord.
13. Track of that duct in the uterine wall; unobliterated portions are, according to Coblenz, the origin of papillomatous cysts in the uterus.

by fibrous threads. The horizontal tube may be traced (12, fig. 124) to the side of the uterus forming the canal of Gartner already alluded to (p. 26). It is important to observe that the vertical tubes become lost in the hilum of the ovary; the significance of this will

be referred to under Ovarian Tumours. The tubes are lined with cubical or broken-down epithelium, and may give rise to tumours known as parovarian (9, 11, fig. 124).

This form of tumour is usually produced by *the distention* of one or more, usually one, of the tubules; its mode of production may, however, be like that of papillomatous ovarian tumours in which true tumour-growth takes place. The diagnosis and treatment of parovarian tumours will be best considered along with those of ovarian (*v.* Chaps. XXIII. and XXIV.).

CHAPTER XX.

DIAGNOSIS AND TREATMENT OF TUBAL DISEASE.

For Literature see Chapter XIX. and p. 244.

Under this head we take up the symptoms and treatment of :—

- (a) Catarrhal conditions, septic and gonorrhœal.
- (b) Pyosalpinx and hydrosalpinx.
- (c) Tubercular disease.

We limit ourselves to these, as they are the conditions open to clinical recognition. Most of the others afford scope only for post-operative diagnosis.

(a) CATARRHAL CONDITIONS, SEPTIC AND GONORRHŒAL.

Symptoms.—The practitioner will first recognise that the case is a septic or gonorrhœal one, and the point to be considered now is, can the onset of tubal mischief, *e.g.*, its spread from the uterus, be ascertained. Acute spread to the tubes will be signalised by lateral pain and increase of temperature, but it is difficult clinically to separate this from ovarian or cellular tissue inflammatory affection.

In cases favourable to the bimanual, one is able to feel the tubal thickening.

(b) PYOSALPINX : HYDROSALPINX.

It is in pyosalpinx that our diagnosis is most accurate, although the condition may sometimes be missed, especially when the tubes are large and blended.

The history will be that of a previous gonorrhœal attack, or a septic attack following labour. In virginal cases the cause is tubercular usually.

Pain of a dull constant character is often complained of, and great dysmenorrhœa is usually experienced. What strikes one most in such cases is the constant and genuine suffering of the patient.

On bimanual examination the uterus should first be mapped out and the tubes at their uterine origin carefully explored, as they are




FIG. 125.

DOUBLE PYOSALPINX blended and with adhesion to rectum, uterus, and vermiform appendix (*Martin*).

often undilated there, and can be felt passing into the distended portions of the tubes.




FIG. 126.

RIGHT-SIDED PYOSALPINX : left-sided catarrhal condition (*Martin*).

There is usually no rise of temperature in chronic cases. The distention may be considerable, and when the pelvis is filled by

the distended and coalesced tubes, exact diagnosis is only possible on abdominal section.

Figs. 125, 127 indicate the nature of the adhesions that may be present.

In *hydrosalpinx* the diagnosis is much more difficult, as the symptomatology is discordant in recorded cases. Thus the effect on menstruation varies, as some have menorrhagia, some have pain, and in others the menstruation is scanty.

Bimanually one can feel the cystic swelling, and when the uterus is pressed against it a sense of recoil may be felt (Landau).




FIG. 127.

RIGHT AND LEFT PYOSALPINX, left ovary adherent. The right tube is adherent behind and to ovary (Martin).

(c) TUBERCULAR DISEASES.

In early tubal tubercular disease there is usually greater frequency of menstruation and increase in its amount. On examination, the thickened and nodular condition of the tubes may be present.

In more advanced cases where the tubes are distended with caseating matter, the negative conditions as to sepsis and gonorrhœa and the positive tubercular tendency are helpful.

The cases are often chronic, and one is apt to forget that the symptomatology of increased menstruation and some pain may be due to early tubercular tubal disease.

TREATMENT OF TUBAL DISEASES.

Prophylaxis.—The facts already given as to the causes of septicity, and the relation of septic disease in the endometrium to tubal disease,

have given a material basis for antisepsis and asepsis, and dealt a death-blow to all intrauterine manipulations which cannot be carried out with strictest antiseptic precautions. The use of stem pessaries, apart from its irrational nature, is to be absolutely forbidden for this reason alone, and while vaginal pessaries are in a few cases helpful, the fact as to the occasional danger of their foulness must not be overlooked.

In early tubal mischief such as may follow septic conditions of the endometrium (cases due to child-birth, abortion, failure of antiseptic precautions) one relies on intrauterine disinfection and thorough rest to the patient, with rest to the local organs involved. For this purpose light diet is to be given; the bowels are to be regulated by gentle laxatives and enemata, and when necessary small doses of morphia by suppository ($\frac{1}{6}$ - $\frac{1}{4}$ gr.). All undue examination is to be avoided. Should freedom from pain and subsidence of the pulse and temperature follow, we may hope that the mischief has passed off.

When chronic thickenings persist they are to be treated with the ichthyol tampon (*v.* under Ovaritis) and blisters. If the return to a healthy condition is slow, and pain is complained of, then a course of baths and vaginal douching at Woodhall, Kreuznach, Schwalbach, or Schlangenbad is often of the greatest service.

All this applies to cases where the condition is a non-suppurative one.

In inflammatory thickenings of the appendages, not yielding to the above treatment, and especially where there is pyosalpinx, hydrosalpinx, or tubercular disease, the question of abdominal section has to be considered. It has been amply shown that pyosalpinx when neglected may burst and cause death. Thus J. K. Fowler found in the *post-mortem* record for three years of the Middlesex Hospital fifteen cases of pyosalpinx; in eight of these it had been the cause of death.

Tait's
Operation.

It is now the established form of treatment to open the abdomen and remove the uterine appendages—an innovation in gynecology due mainly to the boldness of Lawson Tait. He makes a small abdominal incision, frees adhesions by manipulation with the fingers, and taps any cysts with a long curved trocar guided by the fingers. When adhesions are thus broken down, he brings up the tubes to or through the abdominal incision, ligatures with the Staffordshire knot, cuts away the parts above the ligature, drops the pedicle, and drains with a glass tube. Where he cannot remove the tube, he stitches the opening in it carefully to the abdominal incision. Some operators, especially in Germany, make a larger incision, apply ligatures to adhesions, and do not hesitate to turn out the small intestines (suitably covered with warm towels) to facilitate this.

Fuller details on these matters will be given afterwards in the chapter on Ovariectomy, and in the Appendix, where abdominal section in general is considered.

We must note here, however, that tubal suppurative conditions may be treated by vaginal hysterectomy, the tubes and ovaries being removed with the uterus (*v.* Appendix on Abdominal Section).

CHAPTER XXI.

MALFORMATIONS OF OVARY: OVARITIS: PERIOVARITIS: HERNIA, PROLAPSUS: OPERATIONS FOR REMOVAL OF APPENDAGES.

LITERATURE.

Barnes—Diseases of Women, p. 297; Lond., 1878. On Hernia of the Ovary, and Observations on the Physiological Relations of the Ovary: Am. J. of Obst., XVI., p. 1, 1883. *Engelmann*—The dry Treatment in Gynecology: Amer. Jour. of Obst., June and July 1887. *Englisch*—Oesterr. Med. Jahrbuch, 1871, p. 335; or Sydenham Year Book, 1871-72, p. 293. *Freund*—Die Lage und Entwicklung der Beckenorgane: Breslau, 1863. *Griffith*—Tubercle of the Ovaries: Tr. Path. Soc., London. Vol. XL. Gives literature fully. *Hermann*—Prolapse of the Ovaries: Med. Times and Gazette, 22nd October 1881. *His*—Die Lage der Eierstöcke in der weiblichen Leiche: Archiv für Anatomie und Physiologie, Anat. Abtheilung, 1881. *Klob*—Pathologische Anatomie der weiblichen Sexualorgane: Wien, 1864. *Langton*—Hernia of the Ovary: St. Barth. Reports, Vol. XVIII., p. 198. *Lebedinsky*—Ovarien bei Scharlach: Centralb. f. Gyn. I. *Mundé*—Prolapse of the Ovaries: Am. Gyn. Tr., 1879, p. 164. *Olshausen*—Die Krankheiten der Ovarien: Billroth's Handbuch, Stuttgart, 1879. *Schroeder*—Die Krankheiten der weiblichen Geschlechtsorgane: Leipzig, 1878, S. 341. *Schultze*—Zur Kenntniss von der Lage der Eingeweide im weiblichen Becken: Arch. f. Gynäk. Bd. ix. S. 262. *Slavjansky*—Die Entzündung der Eierstöcke: Arch. f. Gyn. Bd. iii. S. 183. *Stratz, C. H.*—Gynäcologische Anatomie. Circulationsstörungen und Entzündungen der Ovarien und Tuben: Kornfeld, 1892. *Sutton, J. B.*—Surgical Diseases of the Ovaries, etc.: Cassell & Co., London, 1896. *Tait, Lawson*—The Pathology and Treatment of Diseases of the Ovary: Birmingham, 1883.

WE first take up some preliminary considerations.

Examina-
tion of
Ovaries.

Palpation of Normal Ovaries.—After the student has had practice in the Bimanual, he will probably meet with some favourable case where he is able to feel the normal-sized ovary. This is best done as Schultze recommends. To map out the right ovary, use the index and middle fingers of the right hand internally and the left hand externally; for the left ovary, the left hand is used internally and the right externally. The patient should lie on her back, with the knees drawn up and the legs rotated outwards. This rotation of the knees renders the psoas muscles tense, thus making their inner edges (which Schultze gives as a guide to the position of the ovaries) more easily palpable. Normally, they lie at about the level of the pelvic brim, half way between the Fallopian-tube angle of the uterus and the psoas (*v. pp.* 26, 61).

Another method of palpating the ovaries is to draw down the uterus with the volsella, and make the examination with the finger per rectum.

MALFORMATIONS OF OVARY.

Absence of one or both ovaries or rather their very rudimentary development, is generally only part of maldevelopment of the uterus. Malformations of Ovary. Occasionally a third ovary is present—a fact worth keeping in mind in relation to Battey's operation (p. 245).

OVARITIS.

SYNONYM—Oöphoritis.

Ovaritis.

NATURE—An acute or chronic inflammation of the ovary.

Simple *Hypercæmia* of the Ovary may also occur.

PATHOLOGICAL ANATOMY.

Acute ovaritis.—Of this we recognise two forms corresponding to the two subdivisions of ovarian tissue—the follicular or parenchymatous, Pathological Anatomy. and the interstitial. Stratz, however, states that these forms are always combined.

In the *follicular form*, the ovary is not much enlarged; but we find on microscopical examination the peripheral follicles increased in size, their contents turbid or purulent, the cells of the membrana granulosa and the ovum in a state of cloudy swelling. The zona pellucida becomes thickened and folded. Usually the surrounding tissue participates, though to a less marked degree, in the inflammatory changes; and in marked cases the germ-epithelium becomes cloudy and broken down, with fibrinous deposits on its surface.

Lebedinsky has examined the changes in the ovary in scarlet fever. To the naked eye, there was no difference; but on microscopic examination, the Graafian follicles were found altered with cloudy swelling or destruction of the epithelium. The younger follicles were most markedly affected, but the stroma was unaltered. In this way the follicles become destroyed and cicatrized, and the ovarian function thus greatly impaired.

In the *interstitial form*, the ovary is increased in size and its connective-tissue elements are proliferated. Pus may form, and often there are small apoplexies. Slavjansky speaks of the following varieties of the interstitial form: serous, suppurative, hæmorrhagic, and necrotic.

Chronic ovaritis.—As the result of this, we get destruction of the follicles and a cirrhotic condition of the organ, as was found in a case of Tait's examined by Doran. To the naked eye, the ovaries appeared

markedly fissured on the surface. Occasionally the ovary remains distinctly larger. Whether or not we get a super-involution of the uterus as the result of severe and double ovaritis, is not as yet settled. The ovaries may be small and cystic, and according to Tait this form gives rise to severe menorrhagia.

ETIOLOGY.

Etiology. The causes of ovaritis are the following:—

1. Chill at menstrual period ;
2. Gonorrhœa, latent gonorrhœa in the male ;
3. Instrumental exploration of the uterus ;
4. Childbirth and abortion ;
5. Acute febrile disease ;
6. Pelvic peritonitis.

Gonorrhœa.—The ovaries may be inflamed sympathetically, just as the testicles are in gonorrhœa of the male.

Instrumental exploration.—Sometimes after the passage of the uterine sound, especially in difficult cases, the ovary becomes tender.

Childbirth and abortion.—This is a common cause of ovaritis. Thus, in 27 cases at Halle, Olshausen found the ovaries affected in 13. Usually both ovaries are implicated.

Acute febrile diseases.—Cholera, the exanthemata, septicæmia, and phosphorus and arsenic poisoning have ovaritis as one of their results.

Pelvic peritonitis.—It will readily be understood that ovaritis often occurs as part of general pelvic peritonitis.

The follicular form usually occurs in febrile diseases and pelvic peritonitis ; the interstitial form is generally puerperal.

SYMPTOMS AND PHYSICAL SIGNS.

Symptoms and Signs. *Acute ovaritis.*—A case of simple acute ovaritis is not common. The patient usually complains of pain at the side radiating to the back, and of pain on pressure in the iliac regions.

When the Bimanual is made, the ovary or ovaries are unusually accessible, and are felt as mobile, tender, and somewhat enlarged bodies, often about the size of a walnut ; and pressure causes great pain of a sickening character. Owing to adhesions, the mobility may be wanting.

Chronic ovaritis.—The symptoms and physical signs are as in the acute form, but much less marked and with a chronic history. Menorrhagia is often present. Sympathetic pain is sometimes felt below the left mamma. In some cases a form of epilepsy is brought on (menstrual epilepsy), menstruation being in abeyance.

DIFFERENTIAL DIAGNOSIS.

When the ovary is not fixed, there is nothing else with which it can be confounded.

Differential
Diagnosis.

PROGRESS AND RESULTS.

We may have resolution of the affection, adhesion, suppuration and abscess. Sterility is a frequent result of double ovaritis; hysteria is often present.

Progress
and
Results.

TREATMENT.

Acute ovaritis.—A fly blister should be applied over the appropriate iliac region, and the hot vaginal douche frequently used. Bromide of potassium may be given as follows.

Treatment
when
Acute.

R. Potassii Bromidi gr. xxx to ʒi.

Fiat pulv: tales xii.

Sig. One powder at night.

Chronic ovaritis.—The hot douche and occasional blisters are best. The glycerine plug is of value.

Treatment
when
Chronic.

A glycerine plug is made as follows: Take a square piece of absorbent cotton wool about the size of the palm of the hand; pour on its centre about ʒss. glycerine; turn the corners over and squeeze the whole so as to saturate it; lastly, tie a piece of thread about 8 inches long round it. Pass Sims' or Fergusson's speculum, and place the plug in the fornix below the ovary. It should be left in for eighteen to twenty-four hours, and then withdrawn.

Glycerine
Plug.

This plug reduces congestion, owing to the affinity of glycerine for water; has an antiseptic action; and, as we shall afterwards see, forms an admirable pessary. It sets up a watery discharge, so that the patient should be told to wear a diaper.

The tampon may be soaked in ichthyol and glycerine (1 in 20), and this combined with the use of the hot douche is a good routine treatment of chronic cases. The patient can carry it out herself, using the douche at night and the tampon in the morning. If the discharge it sets up be objected to, a dry tampon of non-absorbent cotton wool dipped in bismuth or any mild antiseptic powder may be substituted for the glycerine tampon. It may be passed with the aid of a speculum, and should be smeared at its upper part with vaseline. It does not become hard like the glycerine plug, and the elasticity of the non-absorbent wool is of benefit.

The following mixture is of use.

R. Potassii Bromidi ʒij.

Potassii Iodidi ʒj.

Inf. Gentian. Co. ʒvi.

Sig. Tablespoonful thrice daily.

In menorrhagia uncontrollable by ordinary means, the appendages may be removed (p. 236).

PERIOVARITIS.

Peri-
ovaritis.

By this we understand an inflammatory affection of the tissues surrounding the ovary, which fixes the organ. It is a convenient clinical term for local peritonitic inflammations at the site of one of the ovaries. It is higher up than the usual cellulitic deposit. The treatment is the same as in chronic ovaritis.

DISPLACEMENTS OF THE OVARY—HERNIA.

Hernia of
the Ovary.

The term *Hernia* is limited to those cases where the ovaries are present in the inguinal canals, in the obturator foramen (rare), or as part of an abdominal hernia. Percival Pott's case, where this first condition existed and where he excised both of the displaced organs, is the classical instance of this displacement. The usual form is where they are present in the *inguinal canal*.

ETIOLOGY.

Etiology.

Ovaries in the inguinal canal are usually congenital, having descended along the unobliterated process of peritoneum. In 17 cases out of 23 cases, Englisch found it to be congenital; and in one-third of these, the hernia was double.

DIAGNOSIS AND DIFFERENTIAL DIAGNOSIS.

Diagnosis.

An oval tumour of the size of the ovary, tender on pressure, is found in the inguinal canal. Its connection with the uterus may be demonstrated by drawing the latter down with a volsella.

It requires to be diagnosed from an ordinary hernia, and from hydrocele of the round ligament.

TREATMENT.

Treat-
ment.

A protecting shield may be worn; and where very troublesome, the ovaries may be cut down upon and removed. Reduction is usually impossible, owing to adhesions.

PROLAPSUS.

Prolapse
of Ovary.

We have already considered the support of the ovary. Its attachments to the broad ligament, to its own special ovarian ligament, and to the ovarian fimbria of the Fallopian tube, assist, but its chief support is the infundibulo-pelvic ligament of the Fallopian tube; in addition, its own specific gravity has an influence in determining its level. Its

position is constantly changing. As the bladder fills, it is displaced backwards, and its lower end rises; during pregnancy, it is drawn upwards out of its pelvic position and somewhat enlarged. The ovary is thus an organ liable to displacement, of which the most important is the *downward* one—known as *prolapse of the ovary*.

PATHOLOGICAL ANATOMY.

The ovary lies lower than usual, in the lateral or in the true pouch of Douglas; the uterus may be in its normal position, but oftener it is retroverted. The ovary is usually enlarged, and often fixed by peritonitic adhesions. Patho-
logical
Anatomy.

Mundé considers the varieties of prolapsus as—

- (1.) Retro-lateral, in the lateral pouch of Douglas;
- (2.) Retro-uterine, in the true pouch of Douglas;
- (3.) Ante-uterine, in the utero-vesical pouch, very rare;
- (4.) In the infundibulum of an inverted uterus.

ETIOLOGY.

The conditions present in the puerperium favour displacement of the ovary for two reasons; the normal ascent of the uterus during pregnancy may stretch the ovarian and infundibulo-pelvic ligaments, and the ovary may not return to its normal size after parturition. Simple congestion of the organ may cause it to descend; and it is alleged that sudden jolts may also drive it below its normal site. It is not quite certain whether the congestion is cause or result. Probably it is the cause; but it is also aggravated by the displacement. Etiology.

SYMPTOMS.

These are radiating pains, pain on defæcation and coitus, a dragging sensation, reflex nervous symptoms with general irritability. Symptoms.

PHYSICAL SIGNS.

Bimanually, we feel in the true or in the lateral pouch of Douglas a small body or bodies, exquisitely tender and lying distinct from the uterus. By the rectal examination, the ovary is felt with unusual distinctness. Great care must be taken to be gentle in examination. Cystic small ovaries are often adherent, the adhesion being probably caused by rupture of the cysts, which may be done by even gentle manipulation and cause aggravation of symptoms and fresh adhesions. Physical
Signs.

TREATMENT.

Blisters over the iliac region, hot vaginal douche, and bromide of potassium in fifteen-grain doses thrice daily. The bowels are to be Treat-
ment.

opened by means of saline purgatives, such as the Friedrichshall water or Carlsbad salts. The following mixture is good:—

R. Magnesiae Sulphatis	ʒvj.
Quininae Sulphatis	gr. xxiv.
Acidi Sulph. dil	ʒiij.
Tincturae Capsici	ʒj.
Aquam ad	ʒvj.

Sig. Tablespoonful thrice daily.

This relieves the congestion by unloading the bowels.

A course of treatment at Woodhall, or at Kreuznach or some other German Spa is often of service.

Often the prolapsed and non-fixed organ becomes, after a week of this treatment, distinctly higher in position. The glycerine plug or dry tampon is then of the utmost value.

In the chronic stage, when the uterus is retroverted and not fixed, the ring or the Albert Smith pessary is good (*v.* Retroversion of Uterus).

The cases where the tender ovaries are fixed low down by adhesions are exceedingly difficult to treat. When the uterus is retroverted and fixed and the ovaries below it, we get one of the most troublesome cases possible. Palliative treatment by blisters and the hot douche is best; if the case is not amenable to this treatment and the patient's general health is suffering, the propriety of Battey's operation should be considered.

DEVELOPMENT OF OPERATIONS FOR REMOVAL OF THE APPENDAGES.

LITERATURE.

BATTEY'S OPERATION.—The literature on this operation is too extensive to be given in detail in a student's manual. The best summaries of cases are by Engelmann, Hegar, and Simpson. *Battey*—Battey's Operation: Transactions of International Medical Congress, Lond., 1881. See *Am. J. of Obst.*, October 1881, for discussion. See also *Battey's Operation*: American System of Gynecology edited by Mann, Vol. II., p. 837. *Beatson, G. T.*—On the Treatment of inoperable cases of Carcinoma of the Mamma: *Tr. of Med. Chir. Soc.*, Edinburgh, 1895-96. *Byford*—Removal of the Uterine Appendages, etc., by Vaginal Section: *Am. Journ. of Obstet.*, 1888, pp. 337 and 872. *Engelmann*—The Difficulties and Dangers of Battey's Operation: *Am. Med. Asso. Trans.*, 1878 (date of reprint). Battey's Operation, 3 fatal cases: *Am. J. of Obst.*, July 1878. *Hegar*—Die Castration der Frauen: *Volkmann's Sammlung*, Nos. 136-138. *Simpson, A. R.*—History of a Case of Double Oöphorectomy or Battey's Operation: *Br. Med. J.*, May 24th, 1879. *Sims, J. Marion*—Remarks on Battey's Operation: *Br. Med. Journal*, 1877.

TAIT'S OPERATION.—*Bertram*—Laparotomie bei Tumoren der Tuba Fallopii: *Berliner Klinische Wochenschrift*, Jany. 22, 1883. *Savage*—Diseases of the Fallopian Tubes: *Birm. Med. Rev.*, Jan. 1883. *Tait, Lawson*—The Diagnosis and Treatment of Chronic Inflammation of the Ovary: *Br. Med. Jour.*, July 29, 1882. An Account of 208 consecutive cases of Abdominal Section performed between Nov. 1, 1881, and Dec. 31, 1882: *Br. Med. Jour.*, Feb. 17, 1883. Recent Advances in Abdominal Surgery:

Int. Med. Cong. Tr., Lond., Vol. II., p. 228. The Modern Treatment of Uterine Myoma: Brit. Med. Journal, August 15, 1885. Removal of Uterine Appendages for the arrest of Uterine Hæmorrhage: Am. Journal of Med. Science, 1882. Thomas, T. G.—A Contribution to the Subject of the Removal of the Uterine Appendages (Tait's Operation) for Prolonged Menstrual Troubles with Recurrent Pelvic Inflammations: N. Y. Med. Jour., Jan. 13, 1883. (This gives merely the literature concerned in the development of the operations.)

It will ever be to the renown of Dr Battey of Georgia, U.S.A., that he first recognised that something required to be done towards the treatment of the unsatisfactory cases supposed to be due to chronic inflammatory diseases of the peritoneum and pelvic connective tissue. Sufferers from these conditions were allowed to drag on without help and sufficient investigation until Battey proposed to cure such by removing the ovaries so as to bring on an artificial menopause. His first operation was performed on August 17th, 1872, for severe dysmenorrhœa. On July 27th of the same year, Hegar of Freiburg had removed both ovaries, but did not publish an account of his case. Gradually Battey's example was followed by other operators, and Tait urged that the tubes should also be removed to check menstruation.

Plentiful operation has now demonstrated that inflammatory conditions of the ovaries and tubes are prevalent to an extent not even dreamt of, and also that neither of the operations absolutely checked menstruation in all cases.

Operation in such cases is now considered necessary in order to remove diseased structures causing risk to bodily health, and not merely to check menstruation (*v.* Chapters XIX. and XX.). At the same time, certain indications for Battey's operation have remained, viz.:—removal of ovaries presumably healthy in order to check the hæmorrhage of certain bleeding fibroids; their removal to cure so-called ovarian dysmenorrhœa; also for certain forms of epilepsy, and hystero-epilepsy. In the first two classes of cases it gives good results, but not in the third. Quite recently Beatson of Glasgow has urged Battey's operation in inoperable mammary cancer, but sufficient experience has not been gained as to its efficacy in such cases.

Nomenclature.—By Battey's operation we mean then, removal of ovaries presumably healthy, in order to bring on premature menopause. Tait's operation is removal of ovaries and as much of the tubes as can be grasped in the ligature, for the same object. Battey's operation has also been termed "normal ovariectomy" and "castration," but "oöphorectomy" is the best term of all those suggested.

It is worth noting that Blundell of London proposed this operation in 1823, but no actual case was so treated.

For the details of the operation, the student is referred to p. 236, and to Abdominal Section in the Appendix.

CHAPTER XXII.

THE PATHOLOGY OF TUMOURS OF THE OVARY, PAROVARIIUM, AND BROAD LIGAMENT.

LITERATURE.

- Amann, V. A.*—Kurzegefasstes Lehrbuch der microscopisch gynäkologischen Diagnostik. Bergmann, Wiesbaden, 1897. *Bantock*—On the Pathology of certain (so-called) Unilocular Ovarian Tumours: Lond. Obst. Jour., Vol. I., p. 124. *Coblenz*—Zur Genese und Entwicklung von Kystomen im Bereich der inneren weiblichen Sexualorgane: Virchow's Archiv, Bd. 84; *ibid.*, Bd. 82. See also Ztschrift für Geburtshülfe und Gynäk., Bd. VII.; and Arch. für Gynäk., Bd. XVIII. *Coe*—Fibromata and Cystofibromata of the Ovary: Am. J. of Obst., XV., 561. *Cullingworth*—Fibroma of both Ovaries: Lond. Ob. Tr., XX., p. 276. *De Sinéty*—(v. Malassez). *Donat*—Ein Fall von sogenanntem Pseudomyxoma Peritonei: Archiv für Gynäk., Bd. XXVI. *Doran*—Clinical and Pathological Observations on Tumours of the Ovary, Fallopian Tube and Broad Ligament: London, 1884. (Also v. Harris.) *Drysdale*—On the Ovarian Cell found in Ovarian Fluid: Trans. Americ. Med. Ass. (1873, date of reprint). *Duplay*—Des Kystes du ligament large: Arch. Générales de Médecine, Oct. 1882. *Eichwald*—Colloidartung der Eierstöcke: Wurzb. Med. Z., B.V. 1864, p. 270. *Fischel*—Ueber Parovarialsystem und parovarielle Kystome: Arch. für Gynäk., Bd. 15, S. 198. *Flaischlen*—Ein Fall von Psammomasarcom des Ovarium Zur Lehre der Entwicklung der Papillärkystomen: Zeit. für Geb. und Gyn., Bd. 6, 231. *Foulis*—Cancer of the Ovary: Ed. Med. Jour., 1875, p. 838. The Diagnosis of Malignant Ovarian Tumours, and Malignant Peritonitis: Brit. Med. Jour., 1878, pp. 91 and 658. *Fox, Wilson*—On the Origin, Structure, and Mode of Development of the Cystic Tumours of the Ovary: Lond. Roy. Med. and Chir. Tr., Vol. XLVII., p. 227. *Gabbett*—Colloid Degeneration of the non-cystic Ovary, &c.: Journal of Anat. and Physiology, Vol. XVI. *Garrigues*—Diagnosis of Ovarian Cysts by means of the examination of the Contents: Am. J. of Obst., XV., p. 1. *Griffiths*—Article on Ovary in Playfair and Allbutt's System of Gynecology: Macmillan & Co., 1896. *Gusserow*—Ueber Cysten des breiten Mutterbandes: Archiv f. Gynäk., Bd. IX., S. 478. *Harris and Doran*—The Ovary in Incipient Cystic Disease: Jour. of Anat. and Physiol., Vol. XV., Pt. IV., July 1881. *Howell, S. Y.*—Pathology of Ovarian Tumours: Amer. Syst. of Gynec. and Obst., Vol. II., p. 950. *Jones, Mary A. Dixon*—Another hitherto Undescribed Disease of the Ovaries: New York Med., Vol. LI., p. 511. *Killian*—Zur Anatomie der Parovarialcysten: Arch. für Gynäk., XXVI., S. 460. *Landerer*—Ueber Metastasenbildung bei carcinomatösen ovarialcysten; Zts. für Geb. and Gyn. XXXI., 123. *Leopold*—Die soliden Eierstocksgeschwülste: Arch. f. Gyn., Bd. 6. *Malassez et De Sinéty*—Sur la Structure-l'Origine et le Developement des Kystes de l'Ovaire: Archiv de Physiologie Normale et Pathologique, Vol. V., 1878, p. 343. *Nagel*—Beitrag zur Anatomie gesunder und kranker Ovarien: Arch. für Gyn., 31, 237a. *Næggerath*—The Diseases of Blood-vessels of the Ovary in Relation to the Genesis of Ovarian Cysts: Am. Jour. of Obst., Vol.

XIII., 1880. *Olshausen*—Die Krankheiten der Ovarien: Billroth's Handbuch: Stuttgart. *Patenko*—Ueber die Entwicklung der Corpora Fibrosa in Ovarien: Virchow's Archiv, Bd. 84, 1881. *Rindfleisch*—Pathological Histology, New Sydenham Society Translation, 1873, p. 171. *Schroeder*—Die Krankheiten der weiblichen Geschlechtsorgane: Leipzig, 1879. *Slavjansky*—Zur normalen und pathologischen Histologie des Graaf'schen Bläschens des Menschen: Virchow's Archiv, Bd. 51, 1870. *Sutton, J. Bland*—Surgical Diseases of the Ovaries, etc.: Cassell, London, 1896. *Tait*—Diseases of the Ovaries: Cornish, Birmingham, 1883. *V. Rosthorn*—Zur Kenntniss des Endothelioma Ovarii: Arch. für Gyn., Bd. 41, p. 328. *V. Swiecicki*—Zur Casuistik des Pseudomyxoma Peritonei (Werth): Cent. für Gynäk., No. 44, 1885. *Waldeyer*—Die Eierstockscystome: Archiv f. Gynäk., Bd. 1, S. 252. *Wells, Sir T. S.*—Ovarian and Uterine Tumours: Churchill, London, 1882. *Werth*—Ueber Pseudomyxoma Peritonei: Arch. für Gynäk., Bd. 24. *Whitridge Williams*—Papillomatous Tumours of the Ovary: Johns Hopkins Reports, Vol. III. *Williams*—Ovarian Tumours: Reynolds' System of Medicine, Vol. V. *Olshausen, Schroeder, Sir J. Williams, and Whitridge Williams* give the literature well. *Coe's* and *Cullingworth's* articles give the literature for solid tumours.

THE somewhat complex subject of Ovarian Tumours will be best considered under the following heads:—

Origin of
Ovarian
Cysts.

1. *Preliminaries*;
2. *Varieties of ovarian cysts, their naked-eye and microscopic anatomy*;
3. *The nature of ovarian fluid*;
4. *The mode of origin of ovarian cysts*;
5. *Solid ovarian tumours, malignant tumours and the nature of the ascitic fluid associated with them.*

(1.) PRELIMINARIES.

We must first consider some points in relation to the development of the foetus, and the anatomy and physiology of the ovary and adjacent structures. These we take up under the following divisions:—

- (1.) Development of the genito-urinary organs;
- (2.) Anatomy of the ovary;
- (3.) Physiology of the ovary.

(1.) *Development of the genito-urinary organs.* In the human foetus there are two structures from which the future urinary and sexual organs are to be developed: these are the ducts of Müller and the Wolffian bodies (figs. 55-60). In the female, the ducts of Müller form the Fallopian tubes, uterus and vagina; the Wolffian bodies do not persist, but traces are found normally in the broad ligament forming the parovarium, while we may have further traces in the positions shown in fig. 124, as well as in the hilum of the ovary.

Develop-
ment of
genito-
urinary
organs.

(2.) *Anatomy of the ovary.* In regard to the anatomy of the ovary, we must note two great divisions of it: viz., the Hilum and Parenchyma—the former containing traces of the Wolffian bodies and the latter the characteristic structures known as the Graffian follicles with

Anatomy
of Ovary.

their ova (fig. 124). Another division is into oöphoron, paroöphoron, and epoöphoron (parovarium) (fig. 128). In regard to the development of the follicles, we have already seen that the passage of the germ epithelium into the substance of the ovary (the so-called Pflüger's tubes) gives rise to the ova and membrana granulosa.

A section of a developed ovary shows, further, *cellular structures* (fig. 129), which (according to Waldeyer) are some of Pflüger's ducts that




FIG. 128.

FROM SEVEN WEEKS' FŒTUS (♀).

have not developed as they should have done into Graafian follicles, and which may give origin to ovarian cysts.

It must also be remembered that we have in the ovary a great variety of tissue, viz., fibrous and spindle-celled connective tissue, and unstriped muscle.



FIG. 129.

CELLULAR BODIES alleged by Waldeyer to be enclosed germ epithelium which has not developed into normal Graafian follicles. He believes these to be one source of ovarian tumours (*Noeggerath*).

Physiology
of Ovary.

(3.) *Physiology of the ovary.*—When we consider that every month between puberty and the menopause a Graafian follicle distends and then ruptures, we are led to expect what really does sometimes occur, viz., that the follicle may not rupture but merely distend to form a pathological cyst. When pregnancy occurs, the ruptured follicle has its large corpus luteum filling it; and in this also we may have pathological

development. Of the 30,000 to 75,000 Graafian follicles contained in each ovary, only an insignificant number develop and rupture at each menstrual period. Many of the rest atrophy, forming the corpora fibrosa which are seen on section as fibrous points and contain no vessels; it is alleged that these corpora fibrosa may originate also from ripe follicles or from follicles where there has been hæmorrhage.

(2.) VARIETIES OF OVARIAN CYSTS; THEIR NAKED-EYE AND MICROSCOPIC ANATOMY.

The solid forms are considered at p. 256.




FIG. 130.

A SMALL MULTILOCULAR OVARIAN CYST, slightly reduced from natural size (Museum of the Royal College of Surgeons of England, Pathological Series, No. 275) (*Doran*).

Cystic ovarian tumours may be classified as follows:

- (1.) Hydrops folliculorum;
- (2.) Cystoma ovarii—
 - a.* Cystoma ovarii proliferum glandulare (arising in the parenchyma of the ovary),
 - b.* Cystoma ovarii proliferum papillare,
 - c.* Combination of *a* and *b*;
- (3.) Dermoid cysts;
- (4.) Cystoma malignum.

Varieties
of Ovarian
Cysts.

The two great divisions of the cystic tumour are (*a*) the ordinary multilocular, containing many cysts, and lined with columnar epi-

thelium, and (b) the papillomatous form, where the interior of the cyst is lined with tag-like projections. The former is an adenoma; the latter has its nature very much disputed, and is often malignant.

Naked-eye
Anatomy.

Naked-eye Anatomy.—An ordinary multilocular ovarian tumour is best described as made up of two parts—the cyst and its pedicle. The cyst is always multiple (fig. 130); and the pedicle is usually made up of ovarian ligament, Fallopian tube and broad ligament. In the case of the papillomatous form of ovarian tumour (fig. 131, and Pl. VII., fig. 5), we may still recognise the ovary, as such, continuous with the tumour; but in the ordinary multilocular form this cannot be done.

F. T.




FIG. 131.

A LARGE PAPILOMATOUS CYST springing from the Hilum of the Ovary, the greater part of which organ is not involved in the morbid growth. The cyst has forced its way between the layers of the broad ligament as far as the Fallopian tube; this condition has been made more clear by removal of a part of the ligament over the tube, and another part over the cyst; the corresponding portion of the wall of the cyst has also been taken away to expose the cavity (*Doran*).

In the multilocular form, on section, many cavities are found with glairy or semisolid contents. In papillomatous cysts we have the papillomatous condition seen at fig. 132, where the papillomata are fine tag-like projections, and the fluid usually watery. In the multilocular cysts we may have papillary masses sprouting and coalescing. Occasionally, though very rarely, the multilocular tumour is not formed of coalesced tumours, but is grape-like—*Rokitansky's tumour*. *Tait* figures a specimen in his work on diseases of the ovary; *Winckel* and *Olshausen* record similar cases.

Micro-
scopical
Anatomy.

Microscopical Anatomy.—Externally the cystic tumour is covered with cubical or flat cells, not with peritoneum. Beneath this we have

fibrous tissue in lamellæ, while most internally there is the cyst wall with an endothelial or columnar cell-lining. In the papillomatous tumours the projections are covered with cylindrical epithelium, often ciliated, with a core of connective tissue and blood-vessels (fig. 132).

In some cases of ruptured ovarian cyst it has been pointed out by Werth that, in addition to the presence of the gelatinous cyst-contents among the abdominal viscera, we may get a special condition of the peritoneum set up to which he gives the name *Pseudomyxoma Peritonei*.




FIG. 132.

SECTION THROUGH CYST WALL, showing papillæ covered with columnar epithelium, and sub-epithelial layer of connective tissue (*Rindfleisch*). (290)

In one case microscopic examination of the altered peritoneum showed small-celled infiltration, and extension of blood-vessels as a network through the gelatinous layer so that the latter came to lie in spaces.

Donat has also recorded a case operated on by Säger, analogous to those recorded by Werth, where recovery took place. He urges with




FIG. 133.

ROUND-CELLED SARCOMA FROM A DERMOID CYST, showing the transition from the connective tissue of the firmer portion of the tumour to the collection of round cells, with a trace of fibrillation of the intercellular substance in the softer portion of the tumour (*Doran*).

good reason that the so-called "*Pseudomyxoma Peritonei*" is simply peritonitis set up by the irritation of the effused cyst contents (*Fremdkörper Peritonitis*).

Dermoid cysts are said to be due to abnormal inclusion of the epi-Dermoid blast—i.e., are developmental in their origin. They have an outer Cysts.

fibrous coat and an inner one composed of true skin. They may contain hair, teeth, bone, striped muscle, nervous matter, cholesterine, and sebaceous matter. Doran draws attention to the fact that dermoid cysts may contain malignant new growths, notably sarcomata (fig. 133). When teeth are present, their crowns have been found to slope slightly towards the median plain of the body; in this way the side of the body from which the tumour has arisen can be made out (Holländer v. Olshausen).

The nature of these cysts has been very much disputed. Bland Sutton explains their origin very ingeniously by supposing that the Graafian follicles are really mucous glands, and that as mucous membrane and skin are practically identical, we may have skin-like products in a mucous membrane. This would explain the presence of hair, teeth, etc. The old opinion as to their being inclusions of the epiblast was met with the difficulty as to how the epiblast could get so included. It is an interesting speculation to determine how far the fact that the Wolffian body and duct are probably derived from the epiblast—i.e., from a source which may give rise to skin and its products—will help to support the old view.

The *Cystoma malignum* is a cystic tumour which has undergone malignant degeneration. It is noteworthy that malignant disease often develops after the removal of an apparently simple tumour, notably after papillomatous tumours.

(3.) THE NATURE OF OVARIAN FLUID.

Ovarian
Fluid.

Ovarian fluid varies much in consistence and colour. It is usually viscid, and may be so thick as to be almost gelatinous. Its colour is yellowish or greenish; and the specific gravity, when of the more fluid consistence, varies from 1010 to 1020. Chemically, the fluid is complex. The chemical composition has been investigated by Eichwald, whose paper may be consulted.

Ovarian fluid does not give a flocculent precipitate as ascitic fluid does.

The presence (in ascitic) or absence (in ovarian) of such a precipitate can be most easily determined by suspending, as Foulis has suggested, a soft cotton thread in a bottle containing the doubtful fluid; the thread can then be examined microscopically for the deposit which forms in its interstices.

The corpuscular elements of ovarian fluids are various. There may be oil globules, cholesterine crystals, blood fresh or altered, with large granular cells.

Corpuscle
of Bennett
and Drysdale.

Hughes Bennett of Edinburgh and Drysdale of Philadelphia have described a corpuscle, seen at fig. 134, as characteristic of ovarian fluids. According to Drysdale it "is generally round, delicate, trans-

parent, and contains a number of granules but no nucleus;" its size varies from $\frac{1}{5000}$ of an inch to $\frac{1}{2000}$ of an inch in diameter. Acetic acid added to pus makes the cells larger, and brings nuclei into view; while it only increases the transparency of the ovarian cell, and




FIG. 134.

SOME CELLULAR ELEMENTS OF OVARIAN FLUID. At the upper right hand corner we have red blood corpuscles. Below these lie the granular ovarian cells, and below them free granular matter. At the upper left hand corner is shown an epithelial cell; below it, a pus cell after addition of acetic acid; and below this, pus cells before addition of acetic acid. (*Drysdale*).

makes its granules more evident. Garrigues has also investigated the microscopical nature of ovarian fluids in an able research. He believes Drysdale's cell and Bennett's corpuscle to be the nuclei of epithelial cells fattily degenerated, and that there are no pathognomonic ovarian cells.

(4.) MODE OF ORIGIN OF OVARIAN CYSTS.

Ovarian tumours may arise from the following sources:—

- | | |
|--|---|
| <ul style="list-style-type: none"> (1.) Distention and coalescence of Graafian follicles ; (2.) Degeneration of undeveloped Graafian follicles (ordinary multi-locular tumours) ; (3.) Development of remnants of the Wolffian bodies in the hilum of the ovary (papillomatous tumours) ; (4.) Malignant development of the connective tissue of the ovary. (5.) Certain epithelial tubes running into the ovary. This source is now believed to be an important one, and to be due to an abnormal development of the germ epithelium ; | Mode of
Origin of
Ovarian
Cysts. |
|--|---|

There are other alleged sources for which the evidence is not as yet sufficient: viz.,

- (6.) Degeneration of blood-vessels ;
- (7.) Colloid degeneration of ovarian stroma.

(1.) *Distention and coalescence of Graafian follicles.*—There can be no doubt that small cysts may so originate. The proof of this is positive, as Rokitansky found ova in cysts about the size of a bean. Wilson
Fox's view.

Wilson Fox has attempted to show, in his well-known paper, that all the varieties of cystic tumours may be formed in this way.



FIG. 135.

CELLULAR BODIES which Næggerath believes to be diseased blood-vessels and not germ epithelium as Waldeyer asserts (*Næggerath*).




FIG. 136.

SECTION OF OVARY showing an epithelial tube (at the shaded part of the section). Lower down are seen spaces of varying size, and lined with a single layer of epithelium; these cysts are developed from the epithelial tubes. The connective tissue basis is shown only at the shaded part of section (*De Sinéty*) (25).

(2.) *Degeneration of undeveloped Graafian follicles (ordinary multilocular tumours).*—This is probably an important source for the ordinary multilocular tumours. The normal atrophic changes in the youngest or primordial follicles have been traced by Slavjansky and Patenko,

whose researches are too detailed for quotation here. Changes in the normal retrogression of these—viz., active ingrowth of the ovarian stroma, and breaking down of the relics of the membrana propria of the follicle—are probably important in bringing about the cystic changes.

(3.) *Development of remnants of the Wolffian bodies in the hilum of the ovary (papillomatous tumours).*—As already mentioned when speaking of the origin of the parovarium (v. p. 231), remains of the Wolffian bodies persist at the hilum of the ovary. Coblenz believes that when ovarian tumours show a *papillomatous* development, they are usually extraperitoneal, and have arisen from this portion of the ovary. It is




FIG. 137.

CYST WALL OF MULTILOCULAR OVARIAN CYST; ovarian adenoma. (Schmaus in Amann's Mikroskopisch Gynäkologischen Diagnostik).

a, inner lining of a cyst; b, outer wall of cyst; c, small cysts.

probable that this source has been exaggerated, and as a matter of fact papillomatous cysts are not so often extraperitoneal as has been alleged.

(4.) *Malignant development of connective tissue of ovary.*—In malignant disease of the ovary, ascitic fluid is often formed in which are characteristic cells first described by Foulis of Edinburgh. They will be considered under the ascitic fluid associated with malignant tumours (fig. 140).

(5.) (6.) (7.) *Certain epithelial tubes running into the ovary; degeneration of blood-vessels; colloid degeneration of ovarian stroma.*—Nøggerath of New York first stated his view, that diseased blood-vessels might form a source of ovarian cysts, but this source is not generally believed in now. De Sinéty and Malassez first described certain epithelial tubes (fig. 136) from which ovarian tumours develop; these are not true Pflüger's ducts, and but differ from them in being hollow and having no ovum. They consider them as Malassez.

Pflüger's ducts which have taken on a low type of development. *Colloid degeneration of the ovarian stroma* has been said by Rindfleisch to produce an ovarian tumour.

Variance
of Opinion.

The student will, therefore, see that the cellular structures found on section of ovaries, although considered by all as a source of origin for ovarian cysts, have their nature disputed. Næggerath believes them to be diseased blood-vessels; Waldeyer, Spiegelberg, Schroeder and others think them to be Pflüger's ducts, while Doran considers them to be undeveloped Graafian follicles; De Sinéty and Malassez hold that they are Pflüger's ducts degraded in development. The most probable sources for cystic tumours are undeveloped Graafian follicles and the germ epithelium.

It is a matter for regret that there has been so much speculation as to the origin of ovarian cysts, and so little demonstration. What is most needed is the examination by serial microscopical sections of small diseased ovaries.

(5.) SOLID OVARIAN TUMOURS; MALIGNANT TUMOURS AND THE NATURE OF THE ASCITIC FLUID ASSOCIATED WITH THEM.

Solid and
Malignant
Tumours.

Non-malignant (solid) tumours are rare. Myoma of the ovary (fig. 138) has been described by Doran; and Cullingworth has reported an interesting case of fibroma of both ovaries (Desmoid tumours).




FIG. 138.

MYOMA OF THE OVARY (Doran).

The ovary may be enlarged by the presence of small fibrous nodules arising probably from undeveloped Graafian follicles; they have been termed *gyromata*.

We may also have the form of tumour known as endothelioma. It probably arises from the lymphatic endothelium. These solid tumours are usually small in size.

A tubercular condition of the ovary is found as part of general tuberculosis. According to Griffith the ovary is affected with miliary




FIG. 139.

CANCER OF THE OVARY (2-inch and $\frac{1}{2}$ -inch objectives) (Doran).

tubercle, either superficially or deep. The latter may caseate and go on to abscess.

Malignant disease of the ovary is a comparatively frequent occurrence. It often complicates cystic degeneration, specially in the papillary form of ovarian cyst. It arises also independently, and may occur either as primary Carcinoma or Sarcoma. Fig. 139 shows the character of the growth in a case of scirrhus of the ovary in a girl aged fifteen, described by Thornton and Doran.




FIG. 140.

SPINDLE-CELLED SARCOMA OF THE OVARY, showing the superficial and the more central part of the tumour (Doran).

Sarcoma may occur both in the spindle-celled and alveolar forms. The spindle celled (fig. 140) forms a transition from the simple fibromatous tumour to the alveolar sarcoma (fig. 141).

An important feature is the rapid development of ascites, without the Foulis' Re-existence of cardiac, hepatic, or renal disease to explain it. Of great searches.

importance are the cells in the ascitic fluid associated with malignant ovarian disease. Foulis has investigated this subject, and has brought out results of very great value. Through his kindness we have been able to reproduce in fig. 141A cells he has drawn attention to; and he has kindly furnished us with the following description.




FIG. 141.

ALVEOLAR SARCOMA OF THE OVARY (*Doran*).

“Cell groups found in the deposit from ascitic fluid surrounding a large flat or pancake-shaped tumour of the omentum. The tumour was thought to be ovarian. In the fluid in the pleural sacs exactly similar cells and cell groups were




FIG. 141A.

CELL-GROUPS from ascitic fluid surrounding malignant omental tumour (*Foulis*).

seen, and the pleural surface of the diaphragm was studded over with cancerous nodules.

The cell groups and cells were drawn by the aid of the camera lucida under a power of 350 diameters, with No. 3 ocular.”



Fig. 2

$\frac{2}{3}$ nat. Gr. lin.




DIAGRAM OF MODE OF ORIGIN AND GROWTH OF MULTILOCULAR AND
PAPILLOMATOUS OVARIAN TUMOURS (COBLENZ).

It is probable that these liberated cells found in ascitic fluid graft themselves on the peritoneum, and pass through the diaphragm into the pleura and pericardium.

PAROVARIAN CYSTS.

These tumours are developed from the parovarium, have a separable ^{Parovarian Cysts.} peritoneal covering, are thin-walled, and contain a watery fluid which is little more than a mere solution of salt. They may contain papillomatous growths, however, owing to their Wolffian origin—an argument for their being always removed by abdominal section. Small parovarian tumours are common, but they may also be of very large size. They are seldom lined by ciliated epithelium, but usually by cubical or squamous cells, the flattening being, according to Spiegelberg, due to pressure of contents.

It must be remembered of course that all cysts of the broad ligament are not parovarian in their origin. Parovarian cysts are in the site of the parovarium, with the ampullary portion of the tube and the ovarian fimbria stretched and the ovary intact.

OTHER BROAD LIGAMENT CYSTS (PAROVARIAL CYSTS).

By these we mean cysts developed in the broad ligament, but not from ^{Parovarial Cysts.} the ovary or parovarium. They are, however, identical in origin with Parovarian cysts, as they arise from Wolffian relics; further, they may be papillomatous.

The direction of development of these tumours is of great practical interest as they may spread within the folds of the ligament towards the side of the pelvis, towards the uterus, or down in the direction of Douglas' pouch. This renders their removal troublesome as they have then to be enucleated, owing to the absence of a pedicle (*v.* Plate VII.).

These cysts may rupture and cause infective papillomatous growths of peritoneum and ovary.

Plate VII. from Coblenz will be helpful to the student in enabling him to understand the genesis of ovarian tumours, and will also show him the value of a knowledge of development in clearing up the origin of disease.

Fig. 1 shows diagrammatically the development of the urinary and generative organs in the human foetus—female organs (chiefly developed from the ducts of Müller while the Wolffian bodies are rudimentary) shown to the right of the line *m*, and male organs (chiefly developed from the Wolffian bodies while the ducts of Müller are rudimentary) to the left. The rudimentary organs are coloured blue in the figure. On both sides, we have *nn* supra renal capsule, *n* kidney, *u* ureter, *v* bladder,

ua urethra; to the *right* (female organs) are *O* ovary, *po* parovarium, *wb* part of Wolffian body not forming parovarium, *gc* Wolffian duct persisting in Gärtner's canal, *ot* fimbriated end of tube, *ft* Fallopian tube, *ut* uterus, *vg* vagina, *ur* urachus; to the *left* (male organs) are *T* testis, *ep* epididymis, *vd* vas deferens, *md* duct of Müller rudimentary down to *vp* vesicula prostatica.

Fig. 2 shows the fully-developed generative organs in the female: on the left, the organs found in the normally developed female are given; while, on the right, the coloured portion shows the rudimentary structures from which there may be pathological development. On the left, the broad ligament is supposed to have been removed; on the right, the organs are shown in coronal section ($\frac{1}{3}$ nat. size); *ota* ostium tubæ




FIG. 142.

A SIMPLE BROAD LIGAMENT CYST (*Doran*).

Or. Ovary split open; *F.T.* Fallopian tube; *B.L.C.* Broad ligament cyst.

abdominale, *hm* hydatis Morgagni, *fo* ovarian fimbria, *O* ovary, *lo* ovarian ligament, *po* parovarium, *lr* round ligament, *vg* vagina, *wv* upper wall of vestibule, *cc* corpus cavernosum clitoridos, *u* ureter, *l* labium minus, *lm* labium majus; *wb* Wolffian body in its special separate parts as follows:—

Segment I. parovarium, II., III., IV. normally obliterated parts of Wolffian body and duct. From II. we may get cysts of broad ligament developing as well as papillomatous ovarian ones. From the duct (III. and IV.), we may get cysts of cervix uteri and vagina.

Fig. 3 shows a section (in line *ss* fig. 2) of broad ligament, Fallopian tube, and ovary. The blue line *pp* is the peritoneum, *u* being

posterior layer of broad ligament; the red one, the germ epithelium of ovary; *t* tube, *ov* ovary, *lr* round ligament.

Fig. 4 shows development of ordinary multilocular tumour; *C* cystic and *ov* solid parts of tumour; *aa* line of section when tumour is removed; other letters as before.

Fig. 5 shows a tumour which is multilocular and papillomatous, the latter feature caused by Wolffian remains at hilum of ovary.

Fig. 6 shows papillomatous tumour of the parovarium developing in broad ligament, the ovary being intact.

Fig. 7 shows papillomatous cyst extending within the layers of broad ligament developed from remains of Wolffian body and pushing up posterior layer of broad ligament (*cf.* fig. 3*u*).

The student will see by comparing figs. 3, 4, 5, 6, and 7, how glandular and papillomatous cysts alter the relations of structures in the broad ligament. He will also understand the formation of the pedicle (*v.* figs. 4, 5, and 6), as well as the necessity for enucleation in such a case as fig. 7.

CHAPTER XXIII.

DIAGNOSIS OF OVARIAN TUMOURS.

LITERATURE.

Atlee, W. L.—The General and Differential Diagnosis of Ovarian Tumours: Lippincott and Co., Philadelphia, 1873. On Sarcoma of the Ovaries: Am. Gyn. Tr., Vol. II. *Barnes*—Diseases of Women, p. 400: London, 1878. *Brown, J. Baker*—Surgical Diseases of Women: London, 1866. *Duncan, J. Matthews*—Diseases of Women: London, Churchill, 1886. *Hicks, Braxton*—Further Remarks on the Use of Intermittent Contractions of the Pregnant Uterus as a means of Diagnosis: Lond. Inter. Congress Tr., 1882. Also Tr. of Lond. Obst. Soc., Vol. XIII., and Proc. of Royal Soc. of Lond., 1878. *Jastrebow*—On a point in the Diagnosis of Ovarian Tumours: Lond. Inter. Congress Tr., 1882. *Olshausen*—Die Krankheiten der Ovarien: Billroth's Handbuch, Stuttgart, 1877. *Ritchie, C. G.*—Contributions to Ovarian Physiology and Pathology: Churchill, London, 1855. *Schroeder*—Handbuch der Krankheiten der weiblichen Geschlechtsorgane: Leipzig, 1879. *Tait, Lawson*—Diseases of the Ovary: Cornish, Birmingham, 1883. *Wells, Sir T. Spencer*—Ovarian and Uterine Tumours: London, 1882. *Williams, John*—Ovarian Tumours: Reynolds' System of Medicine, Vol. V. Lectures on the Diagnosis and Surgical Treatment of Abdominal Tumours: Br. Med. Jour., 1878.

FOR convenience we take up the diagnosis and differential diagnosis of ovarian tumours under three heads:—

- A. When small (pelvic in position);
- B. When large, multilocular, and pediculated (chiefly abdominal in position);
- C. When large and extraperitoneal (often papillomatous).

A. WHEN SMALL (PELVIC IN POSITION).

These may be either (a) Lateral to uterus, or (b) Posterior to uterus.

(a) Pelvic Ovarian Tumours lateral to Uterus.

Diagnosis
of Ovarian
Tumours
when
pelvic and
lateral to
Uterus.

1. *Symptoms.*—These are chiefly those of pressure and bearing down, and have no diagnostic value. There is no menorrhagia.
2. *Physical signs.*—Palpation and percussion give evidence of the presence of a tumour only when it projects much above the brim. Auscultation gives negative results. On vaginal examination, the cervix

is found displaced to the side opposite to that where the tumour is. Through the fornix a tense, rounded, fluctuating mass is felt projecting downwards. Bimanually the uterus is felt not enlarged, but is displaced to the one side, and is distinct from the tumour, which can be mapped out between the hands. Usually the uterus and tumour are not very movable, owing to the limited space of the pelvic cavity. When the tumour is tapped, ovarian fluid is got.

3. *Differential diagnosis*.—When *lateral* to the uterus, they require to be differentiated from the following:—

- (1.) Pelvic cellulitis;
- (2.) Pelvic peritonitis (encysted serous effusions);
- (3.) Parovarian cysts;
- (4.) Hydrosalpinx, Pyosalpinx;
- (5.) Fallopian-tube gestation;
- (6.) Fibroid and fibro-cystic tumours of uterus;
- (7.) Blood effusion;
- (8.) Solid ovarian tumours.

(1.) *Pelvic cellulitis*.—With this we have inflammatory history and probable cause (as abortion or labour) to guide us. When the cellulitis has gone on to suppuration, there will be rigors and other indications of suppuration. Cellulitic deposits, unless when in the inner part of the broad ligament, are always fixed; are firm when not purulent, and even when purulent do not give very distinct fluctuation.

(2.) *Pelvic peritonitis*.—This will not cause the fornix to bulge downwards, and the history will help us. Tapping gives serum, and not ovarian fluid. When an ovarian tumour is fixed by peritonitic adhesions, it will be almost impossible to diagnose it from encysted pelvic peritonitic effusion except by examination of the fluid.

(3.) *Parovarian cysts* are not so rounded, and have very distinct fluctuation; their secretion is usually simple salt and water.

(4.) *Hydrosalpinx and pyosalpinx* are high in pelvis, tortuous, elongated from side to side.

(5.) *Extra-uterine gestation*.—The symptoms and signs of pregnancy with a tumour beside the uterus corresponding to the period of amenorrhœa (sometimes masked, however, by irregular hæmorrhages from the uterus) point to extra-uterine gestation.

(6.) *Fibroid and fibro-cystic tumours of uterus* (v. Section V.).

(7.) *Blood effusion* in the broad ligaments is more difficult to diagnose during life, but sudden onset with history of fainting and pallor are found (v. Chap. XVIII.).

(8.) *Solid ovarian tumours* are rare. When malignant, there are often nodules in the fornices and ascitic fluid which shows the cells shown at fig. 141A.

(b) *Pelvic Ovarian Tumours posterior to Uterus.*

Diagnosis
of Pelvic
Ovarian
Tumours
when
small and
posterior
to Uterus.

1. *Symptoms.*—The most noticed ones are associated with urination; there may be either retention or constant desire to micturate. There is no menorrhagia.

2. *Physical signs.*—Palpation, auscultation, and percussion give the same result as when the tumour is lateral. On bimanual examination, the uterus is felt markedly displaced to the front, but is not enlarged; and bulging downwards behind the cervix, the round globular cystic ovary can be grasped. Tapping gives ovarian fluid.

Differential diagnosis.—When *posterior* to the uterus, they require to be differentiated from the following conditions:—

- (1.) Encysted serous peritonitic effusion;
- (2.) Retro-uterine hæmatocele;
- (3.) Fibroid and fibro-cystic tumours of the uterus;
- (4.) Retroverted gravid uterus and extra-uterine foetation;
- (5.) Parovarian cysts.

(1.) *Peritonitic effusion* has an inflammatory history; it is not so rounded nor so well defined above. The fluid is serous.

(2.) *Retro-uterine hæmatocele* has, after the blood has coagulated, a hard feeling, and is more expanded transversely. There is a history of sudden onset, menorrhagia, and subsequent inflammatory symptoms.

(3.) *Fibroid and fibro-cystic tumour of the uterus* (v. Section V.).

(4.) *Retroverted gravid uterus and extra-uterine gestation.*—In both of these there will be the signs and symptoms of pregnancy; the amenorrhœa in the latter case may be masked by hæmorrhages from the uterus.

(5.) *Parovarian cysts.*—The character of the fluid is our only certain guide.

It should be specially noted that these pelvic ovarian tumours are apt to cause *pelvic inflammation*, and thus render the exact diagnosis, unless aided by tapping, very difficult.

B. DIAGNOSIS OF OVARIAN TUMOURS WHEN LARGE, MULTILOCULAR,
AND PEDICULATED (CHIEFLY ABDOMINAL IN POSITION).

Diagnosis
when
large.

1. *Symptoms.*—These are chiefly due to its bulk. The patient's notice is attracted to the fact that she is getting rapidly stout.

2. *Physical signs.*—When the patient lies on her back and the abdominal surface is exposed, the following points can be noted:—

On *inspection* the abdomen is seen to be greatly distended. The distention may be uniform, but is often more or less markedly lateral. The distance from the anterior superior spinous process to the umbilicus

is greater on one side than the other. The superficial abdominal veins may be dilated, and lineæ albicantes are sometimes present.

On *palpation*, the distention is felt to be due to an encysted collection of fluid. A mass is felt in the abdominal cavity which is like a sac filled with fluid. Fluctuation is got by placing one hand at a special part and tapping at an opposite point with the fingers of the other hand. However long the tumour be manipulated, *there is never felt any muscular contraction of the cyst wall.*

The feeling of fluctuation is, however, deceptive, and may be felt, for instance, in soft fibroid tumours of the uterus and solid colloid growths.

On *percussion*, when the patient lies dorsal, a dull note is obtained




FIG. 143.

The shaded portion shows the dull area: left figure, ovarian tumour; right figure, ascites (*Barnes*).

over the tumour (fig. 143); but at the flank where the tumour does not bulge, it is clear and tympanitic, since the intestines are there. When the patient turns on her side, with this flank uppermost, the dullness and tympanitic note do not change in position. This sign shows we have to deal with an *encysted* collection of fluid.

Auscultation gives entirely negative results. No sound is heard unless that of friction over a localised peritonitis.

On *vaginal examination*, the uterus is felt displaced to one or other side, or very much to the front. It is rarely retroverted, and—unless impregnated—is not enlarged. The tumour does not usually bulge down into the fornices, but may be made out bimanually.

In order to ascertain how the pedicle lies, we may make the examination *per rectum*, but as a rule it is quite unnecessary to ascertain this fact prior to operation. The tumour is drawn upwards in the abdominal cavity by an assistant. We now lay hold of the cervix with a volsella, pass the index finger of the right hand into the rectum, make traction on the cervix till the fundus is brought within reach of the rectal finger. We recognise a tense band passing from one angle of the fundus, and the enlarged ovarian artery may be felt pulsating in it. The possibility of both ovaries being cystic (which would produce a pedicle on each side), should not be forgotten, though this is comparatively rare. The examination with the volsella is made easier by placing the patient in the *genupectoral posture*; the weight of the tumour makes it gravitate into the abdomen, and renders the pedicle tense; it is also easier to make the rectal examination in this position.

Differen-
tial Diag-
nosis.

3. *Differential Diagnosis.*

They must be diagnosed from the following conditions:—

- (1.) Pregnancy and Hydramnios,
- (2.) Fibromyoma uteri,
- (3.) Ascitic fluid,
- (4.) Fibrocystic tumours of the uterus,
- (5.) Parovarian tumours,
- (6.) Encysted dropsy,
- (7.) Thickened omentum enclosing intestines by adhesions,
- (8.) Omental tumours,
- (9.) Renal tumours,
- (10.) Hydatid of liver,
- (11.) Pseudocyesis,
- (12.) Distended bladder.

In examining a case of abdominal tumour, the practitioner first makes his examination systematically—in every case what is called the routine examination, noting what he observes. By this means he may get facts enough to warrant his drawing a positive conclusion as to its nature. This, however, is not always the case, and he has then to use diagnosis *by exclusion*: it must be one of a certain fixed number of things, and the possibilities are excluded one by one till a definite diagnosis is reached. When examination is unsatisfactory, it should be repeated under chloroform.

We have stated above that ovarian tumours require to be diagnosed from twelve conditions. On each of these we make some brief remarks.

(1.) *Pregnancy*.—At the period of pregnancy when the uterus is so enlarged as to be above the pelvic brim, certain conditions are present. These are suppression of menstruation for a given period, and size of the uterus corresponding to this; mammary signs; lineæ albicantes, and pigmentation. On palpation, we feel a tumour without distinct fluctuation and *having intermittent contractions*; the foetus can be palpated out. The foetal heart (after the fourth month) and the uterine souffle are heard. The vagina is dark in colour, the mucous secretion increased, and the cervix soft.

We need hardly say that palpation, the foetal heart sounds, bruit

and vaginal changes mark out the pregnancy unmistakably. These points may seem too simple to require mention, but cases have been recorded where the pregnant uterus has been tapped for an ovarian cyst.

Hydramnios may simulate an ovarian cyst. The amenorrhœa will help, and especially the occurrence of intermittent contractions as Braxton Hicks has specially pointed out. In one of his recorded cases, the tumour was the size of a seven months' uterus with distinct fluctuation, and there was amenorrhœa for five months. Palpation gave the uterine hardening. Previous to this it had been tapped as a cystic ovarian tumour.

(2.) *Fibromyoma uteri* (v. Section V.).

(3.) *Ascitic fluid*.—When the patient lies on the back, percussion gives a tympanitic note at the umbilicus and a dull one at the flanks (fig. 143); when on the left side, the note is dull on that side and clear over the right; when on the right, it is dull on that side and tympanitic on the left; when she sits up, the upper limit of the dulness is curved with the convexity downwards.

The reason of this is evident. The intestines float on the fluid at its highest point, and give the tympanitic note accordingly (fig. 143).

In tubercular and malignant peritonitis the change of note as the patient alters her position is less perfect, owing to adhesions limiting the movement of the peritoneal fluid. This is a help in diagnosing such conditions.

(4.) *Fibrocystic tumours of the uterus* are difficult to diagnose. The following points should be noted. Fluctuation is only partial and the consistence is variable; the rate of growth is slower; and the fluid drawn off coagulates spontaneously (*Atlee*). It is often difficult to distinguish these from ovarian tumours, and the best operators have sometimes failed to do so (v. Section V.).

(5.) *Parovarian tumours* have very well-marked fluctuation, have their characteristic fluid, and when once tapped do not usually refill as they are often retention cysts.

(6.), (7.), and (8.) In many cases we can make out that the tumour does not pass down into the pelvis and is not connected with the uterus. Sometimes the case is obscure, and abdominal incision alone clears matters up.

(9.) *Renal tumours* grow downwards and inwards, have all their edges rounded, and do not as a rule project posteriorly. When right-sided, the colon lies between them and the liver. Their fluid contains urea.

(10.) The *hydatid* may be connected with the liver and contains hooklets.

(11.) In *Pseudocystis*, the percussion note is tympanitic and the swelling disappears under chloroform.

(12.) The *distended bladder* is of course emptied by the catheter.

C. WHEN LARGE AND EXTRAPERITONEAL (OFTEN PAPILLOMATOUS).

In this class the tumour is not pediculated, and in its extraperitoneal burrowing growth pushes aside uterus, bladder, or large intestines, so that extreme displacement of these may take place (*v. fig. 7, Pl. VII.*). It is therefore of importance in the diagnosis of large abdominal cysts to ascertain the position of the uterus, and also the percussion note so as to make out if large intestine is displaced. When these tumours develop laterally, the displacement of the uterus is an aid to diagnosis; when posterior to the uterus, however, their diagnosis is less easy, as they may only slightly displace the uterus. They usually then bulge well down into the pelvis, lying below the peritoneal level. Their existence should therefore be suspected—

- (1.) If uterus or bladder is displaced markedly;
- (2.) When over a cyst of size sufficient to displace the small intestine, we get a tympanitic note. This indicates displacement of large intestine, which can only be done by an extraperitoneal cyst.

DIAGNOSIS OF ADHESIONS.

Diagnosis
of Adhe-
sions.

When pelvic, the fixation of the tumour they cause can be felt. Adhesions are often the result of tapping; they may also arise from mere pressure. Careful inquiry should always be made as to the history of inflammatory attacks. On palpating the tumour, one can often feel friction. On making the patient take a deep breath, it should be noted whether the abdominal walls move over the surface of the tumour. Much less importance is attached nowadays to the existence of abdominal adhesions. When pelvic, especially if to the bladder or deep in the pouch of Douglas, they are more serious.

CO-EXISTENCE OF PREGNANCY AND OVARIAN TUMOUR.

Co-exist-
ence of
Preg-
nancy.

It should be kept in mind that pregnancy may co-exist with an ovarian tumour, giving its own special symptoms and physical signs in addition.

CHAPTER XXIV.

OPERATIVE TREATMENT OF OVARIAN TUMOURS.

LITERATURE.

- Atlee*—The General and Differential Diagnosis of Ovarian Tumours: Philadelphia, 1873.
- Bantock*—First Series of Twenty-five Cases of Completed Ovariectomy: Brit. Med. Jour., 1879, p. 766. Specimen illustrating the changes in the Pedicle of an Ovarian Cyst when treated by Ligature: Tr. Lond. Obst. Soc., XIV., 1872. Early Operative Treatment of Ovarian Cysts: Lond., 1882.
- Cheyne, Watson*—Antiseptic Surgery: Smith, Elder, & Co., London, 1882.
- Chiara*—Accidental Puncture of Pregnant Uterus: Annali di Ostetricia, etc., 1885, 412 and 453.
- Freund, H. W.*—Der Gewöhnliche und ungewöhnliche Wanderungsmechanismus wachsender Eierstocks geschwülste: Volk. Samml., 361-362.
- Goodell, Wm.*—Article "Ovariectomy:" in Vol. II., American System of Gynecology and Obstetrics (edited by Mann). See also Amer. Jour. of Obst., Jan. 1888.
- Hegar und Kaltenbach*—Die operative Gynäkologie: Stuttgart, 1881, S. 201.
- Lee, C. C.*—Accidental Puncture of the Pregnant Uterus: Amer. Jour. Obstet., 1885, 190.
- Lister*—Address on the Treatment of Wounds: Lancet, Nov. 19 and 26, 1881.
- Mahomed*—Lancet, 1879.
- Malcolm, J. D.*—The Physiology of Death from Traumatic Fever: London, Churchill, 1893.
- Martin, C.*—The After-treatment of Cases of Abdominal Section: Birmingham, Cornish Brothers, New Street, 1894.
- Miner*—Ovariectomy by Enucleation: Philad. Intern. Cong. Tr., 1877.
- Mundé, P. F.*—The Value of Electrolysis in the Treatment of Ovarian Tumours: Am. Gyn. Tr., Vol. II., p. 348.
- Semeleder*—Twenty Cases of Ovarian Cysts treated by Electrolysis: Am. Jour. of Obst., XV., 515.
- Simpson, A. R.*—Three Cases Illustrative of Herman Freund's Law of Pedicle Torsion in Ovarian Tumours: Edin. Obst. Tr., 1893-94.
- Sims, J. M.*—Thomas Keith and Ovariectomy: Am. Jour. of Obst., April 1880.
- Smith, J. Greig*—Abdominal Surgery: 5th Edition, London, 1896. See also his article in Playfair and Allbutt's System of Gynecology.
- Tait, Lawson*—On Axial Rotation of Ovarian Tumours leading to their Strangulation and Gangrene: London Obst. Tr., Vol. XXII., p. 86.
- The Antiseptic Theory tested in Ovariectomy: Lond. Roy. Med. and Chir. Tr., Vol. LXIII., p. 161.
- Recent Advances in Abdominal Surgery: London Intern. Cong. Tr., 1882.
- Diseases of the Ovary: Birmingham, 1883.
- Thornton, J. K.*—The Silk Ligature as a means of securing the Ovarian Pedicle: Brit. Med. Jour., 1878, p. 125. Also, "Rotation of Ovarian Tumours; its Etiology, Pathology, Diagnosis, and Treatment:" Inter. Jour. of Med. Sc., Vol. XCVI., p. 357.
- Wells, Sir T. S.*—Ovarian and Uterine Tumours: Lond., 1882.
- Recent Advances in the Surgical Treatment of Intraperitoneal Wounds: London Intern. Cong. Tr., 1882.
- Additional Cases of Ovariectomy performed during Pregnancy: London Obst. Tr., Vol. XIX., p. 184.

REMOVAL of the ovarian tumour, or Ovariectomy, is the treatment now practised. Other methods have, however, been employed; a brief resumé of these will be useful to the student.

Treatment
of Ovarian
Tumours.

Exploded
Methods.

These methods have been tapping, tapping and injection of the cyst with iodine, electrolysis, drainage into the peritoneal cavity or through the vagina.

Tapping is not a method of treatment followed by cure, and should be used only when it is absolutely necessary to obtain fluid for diagnosis. It may cure parovarian cysts, but it is best to remove them by abdominal section. Ovarian cysts are not retention cysts but have a proliferating lining membrane, for which reason tapping does not cure them. An additional reason against tapping is that it is a procedure by no means free from danger, even to life. By oozing of the fluid through the puncture, adhesions are set up: in some cases, septic peritonitis has proved fatal. Tapping, further, is only palliative and must be followed by ovariectomy.

Method of Tapping.—See that the bladder is empty. With the patient lying on her back make an incision through skin and fat for about an inch, and midway between umbilicus and pubes. Then plunge in the trocar seen at fig. 145. To the side-tube a long piece of tubing is attached, which dips under water. While the fluid is flowing, the patient lies on her side. No bandage is necessary. Care should be taken to prevent regurgitation of air, and a suitable dressing should be applied to the wound (*vide* under Ovariectomy).

Tapping and injection of the cyst with iodine is a procedure not now practised, owing to the risks and uncertainty attending it.

Electrolysis was at one time advocated as a means of cure. Its pretensions to this are unfounded, and few now practise it. Its use has been carefully considered by Mundé of New York, and Semeleder, city of Mexico, in the articles cited, which may be consulted for details and information.

Drainage into the peritoneal cavity, or through the vagina.—The former is dangerous, and the latter is practised only where the cyst is immovably fixed by adhesions.

One fact must be finally noted. Cases of cure of ovarian cysts by tapping, drainage, or electrolysis, are sometimes recorded. These cysts have probably not been ovarian but cysts of the broad ligament—parovarian. Mere tapping often cures the latter. Electrolysis does the same. Electricity has nothing to do with it, the puncture of the needle is enough.

OVARIOECTOMY.

This used to be performed either by vaginal or abdominal incision. The former is now very rarely employed.

VAGINAL METHOD.

Vaginal
Ovari-
ectomy.

This was practised when the tumour was pelvic and small. Thomas of New York, Goodell of Philadelphia, Gilmore, Hamilton, and others have recorded cases. The following was the plan of procedure.

Chloroform or etherise the patient. Place her semiprone or in the lithotomy posture. Pass the Sims speculum. Incise the posterior vaginal wall behind the cervix, in the middle line. Tap the tumour with an aspirator, and then draw it through the incision with the finger or curved forceps. Ligature the pedicle with thin carbolised silk threaded on a handled needle, and divide it on the side next the tumour. Pass a T-shaped drainage tube into the wound which may be stitched round it or left open. Should the temperature rise or the discharge become foetid, irrigate daily with weak carbolic lotion (1-100).

ABDOMINAL METHOD.

Abdominal
Ovari-
ectomy.

The question used to be discussed as to the best time to operate in a case of ovarian tumour—whether, if small, one should wait until it is large. The opinion now held is that one should operate whenever the tumour is diagnosed without reference to its size.

Let us suppose, then, that the ovariologist has a patient—who is otherwise healthy—with an ovarian tumour free from adhesions, and that her period has occurred ten days before. How is the operation performed?

If the patient has not been in any way confined to bed, it is probably better to delay the operation till another period has passed, in order to accustom her to an invalid's life. A pulse and temperature chart should also be taken for a few days prior to the operation. She is kept on light diet, and has no solid food for six hours previous to the administration of chloroform. On the evening prior to the operation, castor oil should be given and an enema used in the morning.

The following are the requisites for operation :

Requisites
for Opera-
tion.

Chloroform and ether ;
Hypodermic syringe ;
Carbolic lotion ;
Porcelain trays for instruments ;
Sponges (a definite number), some small and fixed on sponge-
holders ;
Knives ;
Probe-pointed curved bistoury ;
Scissors, straight and curved ;
Spatulæ ;
Dissecting and dressing forceps ;
Péan's or Well's artery forceps—a definite number (12) of pairs ;
Tenacula, blunt hooks ;
Needles on fixed handles ;
Aneurism needle ;
Fine catgut for bleeding vessels ;
Carbolised silk (Nos. 3 and 4) ;
Two pairs ovariectomy forceps (Nélaton's or Keith's) ;
Well's trocar ;
Clamp (in reserve) ;
Cautery, actual or Paquelin's ;
Cautery-clamp ;
Curved needles ;
Needle holder with small needles on horse hair sutures ;
Drainage tubes (glass or ordinary) ;
Reflecting mirror ;
Iodoform, iodoform gauze, salicylic wool, flannel bandages.

The assistants necessary are three in number, viz., one for chloroform, one for instruments, one to help the operator. A trained nurse who can pass the catheter and administer purgative or nutritive enemata, is necessary. The patient is placed on an ordinary table, of convenient height and length, and lies on her back. The table is placed so that

the patient's feet are towards the window. The legs and chest are to be warmly covered, and hot-water bottles should be laid at her sides and feet. The room should be comfortably warm. The best position for the operator is to stand on the patient's right side, with his back to her feet and to the window. The question of the use of antiseptics in ovariectomy will be discussed afterwards. The instruments are placed near the operator in shallow porcelain trays, and in 1-40 carbolic solution.

Sponges. The sponges should be soft, fine, and thoroughly clean. Twelve are sufficient. Some are small and on sponge holders; one is large and flat. They should be thoroughly wrung out of warm 1-60 solution. *The sponge assistant should know how many sponges he has, and should be sure that he has recovered them all before the abdominal wound is closed. Sponges should never on any account be torn up during an operation.*

Preliminaries. *Preliminaries.*—The patient, who has had a very light breakfast some hours previously, should be chloroformed or etherised; the skin washed and shaved. She has warm underclothing on. The boundaries of the exposed abdomen are covered with thin mackintosh, and over this are laid towels boiled in soda solution and wrung out dry.

The following are the steps of an ordinary operation:—

1. The abdominal incision;
2. Evacuation of the cyst contents;
3. Drawing out of the cyst from the abdomen;
4. Securing of the pedicle;
5. Treatment of adhesions, and bleeding from them;
6. The peritoneal toilette;
7. Closure of the abdominal wound;
8. Drainage—when necessary;
9. Dressing of the wound;
10. After-treatment—complications.

Incision. 1. *The abdominal incision.*—This is usually four inches long, is made in the middle line, and has its lower limit about an inch above the symphysis. It passes through—

skin,
fat,
linea alba,
extraperitoneal fat,
peritoneum.

Sometimes the linea alba is missed, and the rectus muscle cut into. This is no disadvantage, and the cicatrix is probably stronger. The extraperitoneal fat is a good landmark. All bleeding points are carefully attended to before the peritoneum is opened. They may be seized with Péan's forceps which are left on for a time, or they may be

ligatured with catgut. When the extraperitoneal fat is reached, it is picked up with two Péan's forceps so as to get a short transverse fold; this is cut, and the manœuvre repeated until the peritoneal cavity is opened. The cyst is then exposed.¹

2. *Evacuation of the cyst contents.*—This may be accomplished in various ways. Well's trocar (fig. 144), with its point projected, is plunged Methods of Evacuation.




FIG. 144.

WELLS' TROCAR (A). *a*, sharp point, protected by tube *b*, which is projected by pushing out thumb-piece *d*; *c*, tooth catch to grasp cyst wall; gutta-percha tubing is fitted on to *e*.

in, and the fluid passes along the thick tube to a suitable pail below the table. As soon as the trocar enters the cyst, the shield is pushed out to guard the point. The trocar has teeth for catching up the cyst wall. Keith used a large aspirator, so as to empty speedily. Schroeder used no trocar, but simply cut in with his knife and squeezed the fluid out. The kneed trocar may be used (fig. 145), but a simple large trocar without toothed catch is best. When the fluid is very thick it may not flow, and have to be squeezed or scooped out. Secondary cysts, if large, are also perforated.

While the fluid is being evacuated an assistant keeps up steady pressure on the abdominal walls, in the region of the flanks, in order to prevent fluid from passing in or the intestines from passing out.

3. *Drawing out of the cyst from the abdomen.*—This is accomplished Cyst drawn out. by seizing the collapsed walls of the tumour with Nélaton's (fig. 146) or Keith's forceps, and steadily pulling it out. The assistant still keeps




FIG. 145.

TROCAR FOR TAPPING. Tubing is fitted to side-piece.

up pressure. By this means the operator now has the pedicle at the abdominal incision, and the cyst outside. The assistant by means of Cyst separated. sponges keeps back the intestines should they attempt to protrude.

¹ Sometimes the cyst develops between the layers of the broad ligament, lifts up the anterior lamina, and strips the peritoneum off the anterior abdominal wall. When the operator has cut through the abdominal muscles he is puzzled by finding no peritoneum. Puncture and dragging out the collapsed cyst will, however, clear up matters.

Securing of Pedicle. *Securing of the pedicle.*—This is one of the most important steps of the operation. There are three methods which may be used, viz.—

The clamp,
The cautery,
The ligature.

Of these, the clamp is now seldom used. Keith and others advocate the cautery; but the ligature and dropping back of the pedicle is the favourite and probably the best method. The clamp may be necessary if the pedicle is thick.




FIG. 146.

NÉLATON'S FORCEPS.

By Clamp. *The clamp* was introduced by Jonathan Hutchinson, but, as already said, has yielded to the ligature. The varieties of clamp are numerous. Fig. 147 shows Wells'; it consists of two short arms jointed together and provided with a screw and removable handles. It is used as follows:—

The clamp is held by its handles and made to grasp the pedicle between the cyst and the uterus; the bars of the clamp proper are then approximated, and the screw tightly screwed up. The pedicle is examined to see that it is grasped and equally compressed;




FIG. 147.

WELLS' CLAMP ($\frac{1}{2}$), with removable handles. The serrated part with the screw is the clamp proper.

if one part is thin, Spencer Wells recommends that the pedicle be first secured with a ligature. The pedicle is treated extra-peritoneally with the clamp, which rests on the skin. The great advantage of the clamp is its security against hæmorrhage. Its evident disadvantages are the following:—It does not suit all cases, as it cannot be used when the pedicle is too large or too short; it may cause ventral hernia; it exercises undue traction on the uterus; but, above all, it may cause a slough deeper down than the skin, and the discharges, passing into the peritoneal cavity, may do great mischief. Thus the mortality was high (25 p.c.) in cases where the clamp was used.

The cautery was introduced, as a means of treating the pedicle, by Baker Brown of London. By Cautery.

In order to use the cautery, we need a special cautery clamp and either cautery irons or Paquelin's cautery. Keith used ordinary cautery irons heated in a little charcoal brazier. The *cautery-clamp* has two hinged bars provided with handles; each bar has one surface, which is made of ivory—a non-conductor—and is placed next the skin; the other surface is made of metal; one of the bars has on its metal surface a metal upright running the whole length of the bar. The pedicle is seized with the clamp (ivory side next to the skin), and the screw turned to fix it. Then the cyst is cut off, so as to leave about an inch of the pedicle on the metal side. The cautery iron at white heat, which is hatchet-shaped, is then passed firmly over the surface, in the angle between the horizontal bar and the upright, until the pedicle is seared flush with the clamp. It acts by heating the clamp to a dull heat and cornifying the part grasped. The pedicle is now caught at the under surface of the clamp with two pairs of forceps,




FIG. 148.

LIGATURE OF PEDICLE by (1) interlocking double ligature; (2) Tait's knot; (3) Bantock's knot. These must be completed by artery knot.

and the clamp removed. If all is right, the pedicle is dropped into the abdomen after the peritoneal toilette is finished.

The ligature should be thin carbolised Chinese silk No. 3 or 4. It is used in the following way:— By Ligature.

A double silk ligature is threaded on a *blunt* needle. The pedicle is transfixated with this, and the ligature cut. Thus we have two ligatures through the pedicle; one is passed round the one half of the pedicle, the other round the other half. They interlace first so as to make a figure of eight (fig. 148, 1). Each is tied firmly in a reef knot. The pedicle is then seized with Péan's forceps, one on each side above the ligature; the cyst is clipped off about half an inch on the cyst side of the ligature; while the pedicle is still held up by the forceps it can be carefully examined to see if any bleeding occurs. It should be noted whether the ligature splits the pedicle vertically so as to cause bleeding; if so, the ends of the thread can be made to surround the whole pedicle below this. If there is no bleeding, the ligature is cut short and the pedicle dropped into the pelvis.

When the pedicle is thick and fleshy it may require to be tied in

three portions as follows:—Pass a double thread so that its shorter half will embrace only *one-third* of the pedicle; withdraw the needle, but keep it still running on the thread, and use it to carry the longer half of the thread through a second point so as to embrace the *middle third* of the pedicle; one portion of the longer half thus forms a loop round the middle third, while the other portion embraces the *other third* of the pedicle (fig. 241). Tait's or Bantock's knot may also be used (*v. fig. 148*).

After the pedicle has been secured by one of these methods, the other ovary should be examined, and, if cystic, removed also.

Changes in Pedicle. The distal portion of the pedicle does not slough. According to Thornton, we may have the five following results:—

- (1.) Adhesion of the peritoneal surfaces on opposite sides of the ligature, and absorption of ligature.
- (2.) Lymph effused over ligature and end of stump, formation of new vessels.
- (3.) Adhesion of pedicle raw surface to some neighbouring peritoneal surface, and passage of blood-vessels between.
- (4.) Hæmorrhage from pampiniform plexus at outer edge.
- (5.) No change or sloughing if patient dies soon.

Treatment of Adhesions and Bleeding.

5. *Treatment of adhesions and bleeding.*—Adhesions in certain cases may give a great deal of trouble. They may be at any point of the periphery of the tumour. When close to important viscera (especially the bladder, intestine, or liver), they are serious. Their treatment is best considered as follows:—(*a.*) when short, (*b.*) when long.

(*a.*) When easily separable, they may be detached by sponging. If the cyst is connected with the anterior abdominal wall, it is sometimes cut into. The operator then separates the cyst from the wall by passing his finger in between them where the adhesion ceases; or he may evert the abdominal wall, and strip the cyst off it with dissecting forceps. Spencer Wells recommends in bad cases to evacuate the cyst and then, by seizing the posterior wall of the cyst with a hand passed into the interior, to evert it and afterwards separate the adhesions. Pressure with sponges or ligatures will arrest any bleeding, or the cautery may be applied. If the bleeding is intractable, a good plan is to pinch up the abdominal walls at the bleeding part and pass a long straight needle through this fold, so as to keep the bleeding peritoneal surfaces in apposition.

Adhesions in the region of the sacro-iliac synchondrosis are dangerous owing to the risk of tearing into the large veins or ureter. The possibility of an adhesion to the tip of the vermiform appendix must be kept in mind.

(*b.*) When the adhesions are *long*, they may be ligatured at two points close to the cyst, and divided between these.

When adhesions to the bladder are present great care must be taken, as, in separating them, the bladder may be torn into. If this happens,

the tear should be stitched with fine silk or catgut, and a catheter kept in for some days. (*Vide* under Vesico-vaginal Fistula). When adhesions are inseparable, the adherent portion of the cyst may be ligatured all round with silk, and then cut beyond the ligatures; or it may be simply cut all round the adherent portion, and the edges then cauterised.

For reflecting light into the pelvis or other deep parts, an ophthalmoscopic mirror is invaluable.

6. *The peritoneal toilette*.—This term is a convenient one used by German operators to indicate the *cleansing of the peritoneum*. It must be laid down as a cardinal principle in abdominal section that no serum or blood is to be left in the abdomen. The peritoneum should be thoroughly dry, and no oozing points are to be left. The importance of the toilette cannot be too strongly insisted on. Thomas Keith, whose success in ovariectomy was unrivalled, took the greatest care in this matter, and attributed his success to it. Sims indeed says, "But I think now that it matters very little what we do with the pedicle, whether we use the clamp, the cautery or the ligature, provided we take every care against the exudation of bloody serum into the peritoneal cavity after the closure of the abdominal wound." Peritoneal Toilette.

7. *Closure of the abdominal wound*.—This is done as described under Closure of Wound. Abdominal Section in the Appendix.

8. *Drainage*.—As to drainage, the rule is that none is needed in simple cases. This rule may seem to the student to clash with the invaluable principle that every wound from which there will be discharge ought to be drained. In ovariectomy, however, the peritoneum is an absorbent sac, and the discharge, after a simple operation, is absorbed *before it has time to putrefy* (*Lister*). In complicated cases, as where there have been many adhesions, this drainage by absorption is insufficient; it becomes also dangerous from the amount of serum thrown out, and the risk of its putrefying. *External drainage* is, in such cases, imperative. A perforated glass drainage-tube is passed in at the lower angle of the wound, and down into the pelvis. To keep the patient dry, there is laid over the abdomen a piece of thin rubber sheeting with a slit in it through which the tube passes. Over the end of the tube a sponge or some other absorbent is placed and removed when soaked (Keith). Several pints of serum may thus come away in bad cases. It should not be kept in longer than necessary, and each day should be turned round in the wound. If discharge is sucked up, omentum may pass in at the holes and form little pea-like swellings inside. When the tube has to be removed these detain it, and their narrow necks require to be ligatured and snipped through. The sucking up of discharge may be obviated by placing a strip of iodoform gauze inside the tube, and renewing it occasionally. Drainage.

9. *Dressing of the wound*.—Where there is no drainage, it is sufficient Dressing.

to dust with iodoform and lay on a pad of iodoform gauze or other antiseptic material. The gauze is fastened down with collodion, and above all is placed cotton wool and a broad flannel bandage. Where a drainage tube is used we dust the wound as before, lay over it a piece of protective silk, and then pack round the tube some antiseptic absorbent wool. If the pulse and temperature do not rise and there is no uneasiness, the dressing is left untouched—in simple cases—for eight or nine days. If there is drainage, the dressing should be changed occasionally according to the amount of discharge.

After-treatment
and Complications.

10. *After treatment: treatment of complications.*—Morphia may be given hypodermically, but only when necessary (*vide* p. 189). Little food is allowed for the first thirty-six hours; hot water should be given occasionally, as it helps flatus. At the end of this time, milk and beef-tea are added. An enema may be administered on the third or fourth day. When flatus is troublesome, a tube may be passed into the rectum. Sickness is often great, and should be treated with mustard poultices over the epigastrium and enemata of beef-tea and brandy. If it persists to the third or fourth day, two or three grains of calomel may be given. Tait recommends thirty or forty grains of Epsom salts each hour until the bowels move.

The great guide as to the patient's safety is the state of the pulse. A slow pulse, absence of abdominal distention, and the passage of flatus downwards are the three cardinal symptoms of a safe recovery.

Complications may be—Secondary hæmorrhage;

High temperature;

Septicæmia.

Secondary hæmorrhage, if from the pedicle or adhesions, must be treated by the reopening of the wound and application of ligatures.

For high temperatures the ice cap is good. The Americans recommend the more wholesale method of reduction of temperature by Kibbee's ice cot. Krohne and Seseman of London supply very convenient ice-caps made of block tin pipe. Quinine in fifteen grain doses should be tried. It is probable that some high temperatures, recorded by ovariologists, have been due to the absorption by the peritoneum of carbolic acid used in Listerism.

In cases of septicæmia with peritonitis where drainage has been employed, the peritoneal cavity should be washed with very weak carbolic lotion whenever there seems to be any tension or accumulation of putrid fluid; the abdominal incision may require to be reopened for this purpose. The condition should be further treated by iron and stimulants as needed. (*Vide* Treatment of Pelvic Peritonitis.)

Paralysis of the bowel, with great distention and death, has also been noted (Malcolm); as also death from heart clot (Tait). Tetanus has also occurred.

The patient should, after convalescence, wear an abdominal belt to prevent hernia at the abdominal scar.

ABDOMINAL METHOD WHEN THE TUMOUR IS PAPILLOMATOUS AND EXTRAPERITONEAL.

In such cases (*v. fig. 7, Plate VII.*), a different procedure has to be adopted, viz., Enucleation. The tumour is tapped, drawn on as much as possible, and its peritoneal covering incised, so as to include an elliptical portion. The finger is then used to separate the tumour from its capsule, steady traction facilitating this. Bleeding is arrested with forceps or ligature. Goodell, who has given by far the most graphic description of this method, advises that the uterus and bladder be carefully defined, and the separation begun at the uterine side of the tumour where the large blood-vessels enter. The difficulty in the operation is the separation in the pelvis, since the large veins there (as well as the ureter) are apt to be torn. Injury to the ureter is especially dangerous: it is often not recognised, and, unless a fistula form, is fatal. When enucleation is finished, a large oozing extraperitoneal surface is left. Its edges should be stitched to the abdominal incision, so as to close it off from the peritoneal cavity, and a glass drainage-tube passed in. Some, however, close this opening and drain *per vaginam*.

Cases like these are the really difficult and dangerous ones. The chance of return or peritoneal infection is very great.

The idea of this method of enucleation is due to Miner of Buffalo, although the pathology of this form was not clearly understood then: indeed, Miner's original paper, inasmuch as it seemed to apply to the ordinary ovarian cyst when adherent, was not very intelligible.

THE RELATION OF LISTERISM TO OVARIOTOMY.

The Listerian method of treating wounds is based on the now generally accepted theory that the germ-laden air, or other germ-contaminated substances, coming in contact with a wound leads to putrefactive changes which may end in septicæmia. Lister found carbolic acid destructive to the activity of these germs; and, consequently, Listerism requires that the air in contact with the wound, and all else that touches it, must be purified either with the spray or lotion. Listerism is in no sense a treatment of wounds, but is a *treatment of wound surroundings*. The application of carbolic lotion to a wound is a necessary evil, as carbolic acid is an irritant, and may be absorbed. In the cases treated by the surgeon, Listerism is of the greatest value; and, with drainage, has worked the most mighty revolution in surgery. In peritoneal operations, however, its good is marred by the fact that the peritoneum absorbs the carbolic

Listerism
in Ovari-
otomy.

lotion, and thus its surface is irritated, and often toxic effects ensue. Keith, Tait, and Bantock have therefore abandoned Listerism in abdominal surgery; but Wells and Thornton still carry it strictly out. Listerism has been modified, but only in this, that less importance is now attached to air-contamination of raw surfaces during an operation. Unclean "touch" is the real danger.

Practically most ovariologists at present trust to asepsis and modified antiseptics (*v.* Chap. XV.), and to drainage when necessary.

OVARIOTOMY WHEN PREGNANCY IS PRESENT.

Pregnancy
and Ovari-
otomy.

Although pregnancy co-exists with a large ovarian tumour, ovariectomy should be performed. Bland Sutton gives interesting tables of unilateral ovariectomy during the earlier and later months of pregnancy, of double ovariectomy, and of ovariectomy in the puerperium. These show that before the fourth month the mortality is low, and the risk of abortion small, being greater for parovarian than ovarian tumours. After the fourth month the risk is as in ordinary ovariectomy, but the chance of abortion increases progressively with the advance of pregnancy. Puncture of the gravid uterus during the progress of the operation must be guarded against. This may happen if the pregnancy has not been diagnosed and the pregnant uterus mistaken for a secondary cyst; or it may be as in Lee's case that, owing to a change of the position of the patient from the dorsal to the lateral posture, the ovarian cyst recedes from the abdominal incision and the uterus lies below it without the changes being noted. When this accident occurs, the treatment depends on the depth of the wound. Should the uterine cavity not be opened, then bleeding is arrested by pressure, the wound stitched with continuous silk suture. If the amniotic cavity is opened into, the same treatment may be adopted (*v.* Chiara's case); or the incision may be suitably enlarged, and the foetus, placenta, and membranes extracted. The treatment after this may be removal of the uterus by Porro's operation, simple suture of the walls with silver wire, or the Cæsarean section with the modification introduced by Sänger. The question of the treatment of a labour complicated with an ovarian tumour concerns the obstetrician rather than the gynecologist.

CONTRA-INDICATIONS TO OVARIOTOMY.

Contra-
indica-
tions.

These are universal adhesions and malignant disease. Ordinary ascites, kidney disease, or heart disease, is not a contra-indication unless far advanced. Prognosis should be careful in these cases. In some fatal cases it has been found on post-mortem that the kidneys were small and granular from interstitial inflammation. This may be present while there is no albumen in the urine. There is usually a pulse of high tension and cardiac hypertrophy (*v.* Mahomed's articles).

COURSE AND RESULTS OF OVARIAN TUMOURS WHEN LEFT ALONE.

In some rare cases the operator is unable to remove the cyst after he has begun his operation. He may then stitch the cyst edges to the abdominal walls carefully closing it off from the peritoneum. The best results by this method are got in dermoid and parovarian cysts: they are not good in ordinary ovarian cystomata.

Natural
History of
Ovarian
Cysts.

Adhesions may be set up as the result of chronic peritonitis arising from pressure or tapping. Occasionally the cyst bursts, and in the case of the ordinary ovarian tumour we may get rapid death or the condition termed pseudomyxoma peritonei by Werth (v. p. 251). When parovarian tumours burst, the fluid is usually non-irritating, and is absorbed by the peritoneum, the patient thus becoming cured. Matthews Duncan and others have recorded cases of burst ovarian tumour rapidly becoming fatal. Waxy disease of the liver, kidneys, etc., may result in those cases where the tumour suppurates and discharges into the bowel or through the skin.

Torsion of the pedicle to a slight extent is often noticed in ovarian tumours. When the torsion is so great as to cut off the blood supply from the cyst, we get gangrene of the tumour, and in some cases very serious symptoms, viz., peritonitis, vomiting, and severe abdominal pains. Wiltshire of London was the first to operate for this condition, and recently Lawson Tait has operated successfully in three cases. His paper should be consulted for details. It is interesting to note that the tumours so rotated are usually right-sided, and not necessarily ovarian. The usual explanation of the rotation is that it is caused gradually by the faecal contents passing down the rectum. Tait's book and Thornton's paper may be consulted for fuller details.

According to H. W. Freund, the pedicle twists to the right in right-sided tumours; to the left in left-sided ones.

If peritonitis occur before the tumour is removed, ovariectomy should be at once performed. Keith was the first to do this successfully.

The course and results of ovarian tumours when left alone can fortunately not now be studied. The picture of ovarian disease running its course unchecked, so eloquently described by West, is happily now almost unknown.

"We have symptoms of the same kind as we see towards the close of every lingering disease, betokening the gradual failure, first of one power, then of another; the flickering of the taper, which, as all can see, must soon go out. The appetite becomes more and more capricious, and at last no ingenuity of culinary skill can tempt it, while digestion fails even more rapidly, and the wasting body tells but too plainly how the little food nourishes still less and less. The pulse grows feebler, and the strength diminishes every day, and one by one each customary

exertion is abandoned. At first the efforts made for the sake of the change which the sick so crave for are given up ; then those for cleanliness ; and lastly, those for comfort—till at length one position is maintained all day long in spite of the cracking of the tender skin, it sufficing for the patient that respiration can go on quietly, and she can suffer undisturbed. Weariness drives away sleep, or sleep brings no refreshing. The mind alone, amid the general decay, remains undisturbed ; but it is not cheered by those illusory hopes which gild, though with a false brightness, the decline of the consumptive ; for step by step death is felt to be advancing ; the patient watches his approach as keenly as we, often with acuter perception of his nearness. We come to the sick chamber day by day to be idle spectators of a sad ceremony, and leave it humbled by the consciousness of the narrow limits which circumscribe the resources of our art." (Quoted by Spencer Wells.)

The question of the mortality after ovariectomy is a complex one, owing to differences in cases and also because the use of the clamp in early operations unduly raised the death rate. Of late years the mortality has fallen considerably, chiefly owing to the use of the intraperitoneal treatment of the pedicle (ligature or cautery) and greater care to avoid sepsis.

SECTION V.

AFFECTIONS OF THE UTERUS.

THERE are three periods during which morbid conditions of the uterus arise.

1. *The period of evolution or development*—from the ovum up to puberty. During this stage they appear as anomalies in development—before birth or during childhood. They produce no marked symptoms, but a recognition of their existence is important as regards the future history of the patient.

2. *The period of physiological activity*—from puberty to the menopause. During this stage there occur in the uterus the morbid processes of acute and chronic inflammation, and of new-formation or tumour-growth; on account of its mobility, the uterus is also liable to various forms of displacement. These pathological processes give rise to symptoms of themselves, and also from their effect on the normal functions of the uterus—menstruation, conception, and pregnancy. During parturition the cervix uteri is frequently lacerated, and this may be the starting-point of important pathological conditions.

3. *The period of senile involution or retrogressive development*—from the menopause to death. The term involution is generally used in the restricted sense of the process which occurs after childbirth, but it is the only one which conveniently expresses the retrogressive changes after physiological activity has ceased. During this stage, the most important pathological process is that of malignant new-formation.

Accordingly the following subjects have to be considered in this Section :—

CHAPTER	XXV. Malformations of the Uterus.
„	XXVI. Atresia and Stenosis of the Cervix Uteri.
„	XXVII. Atrophy of the Cervix and Uterus : Superinvolution.
„	XXVIII. Hypertrophy of the Cervix ; Amputation.
„	XXIX. Laceration of the Cervix and its Consequences.
„	XXX. Chronic Cervical Catarrh.

- CHAPTER XXXI. Endometritis.
- „ XXXII. Metritis, Acute and Chronic ; Subinvolution.
- „ XXXIII. Displacements of the Uterus : Ante-
flexion ; Retroversion ; Retroflexion.
- „ XXXIV. Inversion of the Uterus.
- „ XXXV. Tumours of the Uterus. Fibroid Tumour : Patho-
logy and Etiology.
- „ XXXVI. Fibroid Tumour of the Uterus : Symptoms and
Diagnosis.
- „ XXXVII. Fibroid Tumour of the Uterus : Treatment.
- „ XXXVIII. Fibrocystic Tumour of the Uterus.
- „ XXXIX. Uterine Polypi : Tuberculosis.
- „ XL. Carcinoma Uteri (of Cervix) : Pathology and
Etiology.
- „ XLI. Carcinoma Uteri (of Cervix) : Symptoms and
Diagnosis.
- „ XLII. Carcinoma Uteri (of Cervix) : Treatment.
- „ XLIII. Carcinoma Uteri (of Body) : Adenoma Malignum.
- „ XLIV. Sarcoma Uteri : Deciduoma Malignum.

CHAPTER XXV.

MALFORMATIONS OF THE UTERUS.

LITERATURE.

Ballantyne—Malformation of the Genital Organs in Women: Allbutt and Playfair's System of Gynecology: London, 1896. *Dirner*—Ein Fall von Uterus didelphys, etc.: Archiv f. Gyn., XXII., S. 463. *Dos Santos, Las Casas*—Missbildungen des Uterus: Zeit. f. Geb. u. Gyn., XIV., S. 140. *Kussmaul*—Von dem Mangel, der Verkümmerung und Verdoppelung der Gebärmutter, etc.: Würzburg, 1859. *Macdonald, Angus*—Case of Pregnancy in the Left Horn of a Bifurcated Uterus, etc.: Ed. Med. Jour., April 1885. *Mayerhofer*—Die Entwicklungsfehler der Gebärmutter: Billroth's Handbuch für Frauenkrankheiten, Stuttgart, 1878. *Pfannenstiel*—Ueber Schwangerschaft bei Uterus didelphys: Festsch. der deut. Gesellschaft f. Gyn.: Wien, 1894. *Schäffer, O.*—Bildungsanomalien weiblicher Geschlechtsorgane aus dem fötalen Lebensalter u.s.w.: Archiv f. Gyn., S. 199. *Secheyron*—Du Cloisonnement Pelvien Antéro-Postérieur: Annal. de Gyn., XXI., 441 and XXIII., 247 et seq. *Simpson, A. R.*—Case of Double Uterus: Ed. Med. Jour., 1864, p. 957. *Turner*—Malformations of the Organs of Generation: Edin. Med. Jour., June 1865 and May 1866. *Westermayer, H.*—Uterus bicornis et Vagina duplex, etc.: Diss. Inaug. Erlangen, 1894. The standard work is that of Kussmaul. The recent literature is given by Ballantyne, whose monograph has a valuable bibliography with some 400 references.

WHAT is usually described as "malformation" is really a nonformation of one part, involving a relative disproportion. Of this we have an illustration in the uterus. The one-horned uterus is not a "malformation," if by this term we mean that the part which is present is maldeveloped; the condition is a result of the nonformation of the other horn and intervening fundus. It is misleading also to speak of a "double uterus;" the structure thus described is really one uterus, in which the halves have not united into a whole. The word as used, therefore, means an incomplete result, not a defective process. Maldevelopment is a contradiction in terms, there can only be arrested development.

Relation of
Malforma-
tions to
Develop-
ment.

Malformations must be studied in connection with the normal development of the organ. In this way, they become at once intelligible. There are two processes in the progression of an organ to its mature form—*development* and *growth*. There are therefore two causes which together operate in producing malformations—arrested develop-

ment and arrested growth. The period of development of the uterus, by which we mean formation of parts, extends up to the twentieth week; the period of growth is much longer, and extends to the twentieth year.

The student should not pass over this section of the subject as of little importance. To the practical man, malformations seem of little value because he has no power of modifying the result. To the scientific man they are, however, of the greatest interest as furnishing him with permanent impressions of the transitional states of development; they are development caught in the act and fixed permanently for after-investigation. In this chapter we recommend the student to read Etiology before Pathology.

PATHOLOGY.

Uterus
absent or
rudi-
mentary.

Complete absence of the uterus is an extremely rare occurrence, and cannot be demonstrated except on post-mortem examination. It has been described only in cases of foetal monstrosities. A *rudimentary condition* sometimes occurs; in this the uterus is represented by a band of muscular fibre and connective tissue on the posterior wall of the bladder (fig. 149), and the peritoneum forms a single pouch between the bladder and the rectum (fig. 150).




FIG. 149.

RUDIMENTARY UTERUS (*Veit*). Sa Sacrum; U Solid Rudiment of Uterus; h Rudimentary Horn; B Bladder; O Ovary; T Fallopian Tube; r Round Ligament.

In the *uterus bipartitus* (fig. 151), rudimentary horns are present and are solid or hollow. The cervix is represented by a fibrous band which connects the horns with one another and with a rudimentary vagina. The ovaries are sometimes well developed, so that ovulation takes place. The breasts and external genitals may be fully formed.

The *uterus unicornis* (fig. 153) may exist with or without a rudimentary second horn. The vaginal portion of the cervix is small; the palmæ plicatæ within the cervical canal are most marked towards the non-developed side. The body of the uterus is of disproportionate

length and curves towards one side. The fundus, by which we understand the fully-developed horn, is small and tapering; it has only one Fallopian tube and ovary connected with it. On the convex side of the somewhat curved body is the representative of the other horn which is either solid or hollow; it is connected with the developed one by fibrous tissue which may or may not form a previous canal. Connected




FIG. 150.

The same in its relation to the Pelvic Organs. *U* Rudiment of Uterus on the posterior wall of Bladder. The Peritoneum forms one pouch between Bladder and Rectum. (*Schroeder.*)

with this rudimentary horn are the Fallopian tube and ovary of the same side, which are sometimes perfectly developed. In examining preparations of this and other uterine malformations, it is sometimes difficult to determine what is rudimentary horn and what is Fallopian tube. Here development furnishes us with a guide. The insertion of




FIG. 151.

UTERUS BIPARTITUS (*Rokitansky*). *V* Vagina; *U* Uterus; *h* Rudimentary Horn; *o* Ovary; *T* Tube; *r* Round Ligament; *b* Broad Ligament.

the round ligament indicates the point up to which the ducts of Müller are to be formed first into uterine horn and then into corpus uteri. Accordingly, on examining such preparations we determine the point of attachment of the round ligament; all below this is uterine horn, all above it is Fallopian tube. Associated with this malformation we sometimes find absence or rudimentary condition of the kidney of the

Round ligament indicates junction of Uterine Horn and Tube.

same side, since the development of the renal is closely connected with that of the generative system.

Uterus Didelphys. In the *uterus didelphys* the two halves of the uterus remain separate throughout their course; the vagina may be absent, single, or double. It is a rare condition in the living adult female; Pfannenstiel, who has




FIG. 152.
UTERUS DIDELPHYS (Coats).

studied it in relation to pregnancy, has collected eighteen cases, and other cases have been reported by Simon and Löhlein.¹

Fig. 152 shows a uterus described by Paterson and Coats from a patient who died a fortnight after the delivery of a seven months' child.




FIG. 153.

UTERUS UNICORNIS (Schroeder). *R* Right Side; *L* Left Side. The left horn (*h*) is well developed and communicates with the Uterine Cavity. The right horn is in the form of an elongated band; its point of connection with the Fallopian tube is indicated by the insertion of the round ligament which is hypertrophied. Other letters as in preceding diagrams.

There are apparently two uteri, which are separate, but open into a common vagina; they are of nearly equal size—the right which contained the fœtus measuring 5 in. and the left $4\frac{3}{4}$ in. in length, and being respectively $2\frac{1}{2}$ and $1\frac{3}{4}$ in. in breadth.

¹ Centralb. f. Gyn., 1894. S. 997 and 1313.

A case in which there was an accessory or *third* uterus has also been described.¹ There were no uterine appendages connected with it, and it probably originated as a diverticulum on one duct of Müller, the two other uteri arising each from its own duct.

By *uterus bicornis* we understand that the separation into two horns is distinctly visible *externally*. Of this there are various degrees, from a ^{Uterus} Bicornis.




FIG. 154.

UTERUS BICORNIS UNICOLLIS (Schroeder). *r* Round Ligament.

mere depression at the middle of the fundus to a well-marked bifurcation, which rarely extends lower than the os internum; the further down the




FIG. 155.

UTERUS SEPTUS IN VERTICAL TRANSVERSE SECTION (Kussmaul). *U* Uterus placed on septum which divides Cavity into two lateral portions; *T* Fallopian Tubes; *V* Vagina divided into lateral cavities by prolongation of septum downwards.

separation extends, the more obtuse is the angle between the divergent horns. There is occasionally a fold of peritoneum, containing muscular

¹ Holländer, Berlin. klin. Wochens, 1894, p. 452. See also case by Depage, Archiv de Toc., 1894, p. 550.

fibre and blood-vessels, running from the bladder to the rectum in the hollow between the horns. In addition to this external division, the separation is usually carried further down by an internal septum which may extend to the os externum.

Uterus
Septus.

In the *uterus septus* (fig. 155) there is no external indication of the



FIG. 156.

INFANTILE UTERUS (Schroeder).

internal division. The uterus is divided by a septum beginning at the fundus uteri and extending downwards for various distances, sometimes as far as the os externum. It is otherwise normal.

Various other conditions have been described, such as a septate




FIG. 157.

PRIMARY ATROPHY OF THE UTERUS (Virchow).

vagina with two cervices, and apparently a single uterine cavity, or a double cervix with a single uterine cavity above and vagina below.

The presence of *septa* at various points in the vagina or uterus, or of *sacs with an accumulation of blood or pus* alongside of a patulous tract, or

¹ Merttens: *Centralb. f. Gyn.*, 1894, p. 1001.

of transverse septa causing *complete occlusion* usually point to defective development of one or other duct at a point in its course.

The *infantile uterus* (fig. 156) is characterised by shortness of body and disproportionate length of cervix; in fact the relative lengths of body and cervix remain the same as at birth, from which the name "*infantile*" is derived. The cervix ($1\frac{1}{2}$ inches long) is two or even three times the length of the body ($\frac{1}{2}$ in. to $\frac{3}{4}$ in.). The whole uterus is smaller than normal. The walls (specially those of the body) are thin and the cavity is small. Infantile Uterus.

The term *congenital atrophy* is applied to cases in which the proportions of body and cervix are of the normal *virgin* type, while the organ as a whole is atrophied (fig. 157). An excess of connective tissue is present in the walls, which makes their consistence firmer. This malformation occurs in scrofulous and chlorotic patients and with cretinism, and is often associated with hysteria and epilepsy. Congenital Atrophy of Uterus.

ETIOLOGY AND CLASSIFICATION.

Malformations differ according to the period at which development and growth are arrested, and the extent to which they are interfered with. There are five periods in development and growth (*Fürst*), which can be easily remembered when we bear in mind the division of the period of intra-uterine life into ten *lunar months*. In the first period, which extends over the *first* and *second* lunar months (from fertilisation to the eighth week), the septum between the adjacent ducts of Müller is as yet unbroken. By the end of the second period, which corresponds to the *third* month (*i.e.* eighth to twelfth week), the septum has entirely disappeared; but the upper portions of the ducts remain distinctly separate, forming the horns of the uterus and the Fallopian tubes. During the third period, *fourth* and *fifth* months, the angle between the uterine horns disappears so that the base of the uterus becomes flat. In the fourth period, *last five months*, the flattened end of the uterus, between the Fallopian tubes, becomes arched through the development of the fundus. The fifth period extends from *birth to puberty*. During this period no important change takes place till, at puberty, the uterus passes from the infantile to the virgin form. It does not, however, cease to grow till the twentieth year. Five periods in development of Uterus.

We are not yet in a position to refer each malformation in detail to its proper period; but the more perfectly we are able to do this the more satisfactory will our classification be. At present we separate the first four periods from the fifth, and speak of the period of foetal life in contradistinction to the period of childhood. This forms the basis of our classification. Classification of Malformations.

1. MALFORMATIONS ARISING DURING FŒTAL LIFE. Of these there are

the following:—*complete absence* or *rudimentary condition* of the uterus; the *uterus bipartitus*, produced by a development of only the upper parts of the ducts of Müller into rudimentary horns of the uterus and Fallopian tubes; the *uterus unicornis*, due to the development of only one duct; the *uterus didelphys*, due to the development of the ducts separately; without coalescence; the *uterus bicornis*, in which the ducts coalesce below, and the horns remain un-united by a fundus above; the *uterus septus*, in which the coalescence of the ducts and development of the fundus takes place so that the uterus appears normal externally while internally the septum has persisted. These last three are sometimes spoken of as varieties of the double uterus or *uterus duplex*. The association of an antero-posterior reduplication of the peritoneum with some cases of uterus bicornis is of interest from an etiological point of view, pointing back to some mechanical cause which kept the ducts of Müller from blending.¹ It is interesting that a rudimentary condition of the uterus has been observed in more than one member of the same family.

2. MALFORMATIONS ARISING DURING CHILDHOOD. Of these there are the following:—the *uterus infantilis*, in which the uterus does not undergo the development which should take place at puberty, but remains of the same type as it was at birth; *congenital atrophy* of the uterus, in which it assumes the virgin type, but the organ as a whole is atrophied.

SYMPTOMS.

The symptoms of malformation consist in an *impairment of function*, and hence do not appear until puberty.

In the external appearance of the patient there is not necessarily anything to attract attention. The figure, features, temperament, and voice are of the feminine type, even though the uterus is not developed. The mammæ may be fully formed. The external genitals may be found well formed, as their development is independent of the internal organs. It is rare, on the other hand, to find a normal vagina present when the uterus is rudimentary.²

Sometimes local symptoms absent. Complete absence and rudimentary condition of the uterus may give rise to no local symptoms, except the non-appearance of menstruation. If the ovaries are developed, ovulation with associated monthly disturbance is present and the accumulation of menstrual blood in a rudimentary horn may call for operative measures to form a channel for its escape. Even on entering married life the condition need not necessarily attract attention; if the vagina be not well developed, the urethra becomes dilated so as to take its place.

Cause of local symptoms. In the uterus unicornis, menstruation, conception and pregnancy may

¹ See a case reported by Buchanan: Glas. Med. Journ., July 1892.

² As in cases by Kahn-Bensinger, *Centralb. f. Gyn.*, 1887, S. 377; Grechen, *ib.* S. 493; Mundé, *ib.* S. 670; Steinschneider, *ib.* 1888, S. 49; Zweifel, *ib.* S. 474.

go on undisturbed in the developed horn. It is the *imperfectly developed horn which gives rise to symptoms*—the result of the retention of menstrual blood and of the products of conception. If the mucous membrane of this horn discharge blood periodically and there be no communication with the uterus to allow of escape, the blood collects and produces a distended sac—a very rare occurrence. It is of great interest to note that we may have a fertilised ovum growing in the isolated horn; we have not space here to discuss how this interesting condition is produced (fig. 158). Pregnancy may also occur in the one half of a uterus didelphys.¹

Uterus bicornis and uterus septus produce no symptoms, unless one half of the partitioned uterus does not open into the cervical canal—in which case hæmatometra occurs at puberty (*v.* Chap. XLV.). The statement that the patient menstruates regularly throws the practitioner off his guard. He should remember that the menstrual blood *may flow undisturbed from one half of the uterus while it is accumulating in the other*. In both of these forms we have two possible seats for a growing ovum (fig. 159); and this accounts for super-foetation, and those curious cases in which an ovum has been expelled in the course of a pregnancy which went on to full-time.² When the uterus is double, abortion and premature labour are more frequent; the septum also causes difficulty in delivery, and involution progresses more slowly. It has been noted that a decidua forms in the empty half of the uterus, as it does in extra-uterine gestation, and may be expelled in the puerperium.

The anomaly of menstruation during pregnancy has also been thus explained; Henderson found a double uterus in a patient who menstruated regularly during two of her pregnancies—the flow coming probably from the empty cavity.³

The uterus infantilis and the congenitally atrophic uterus are characterised by the absence or scantiness of the menstrual flow and the constitutional nervous disturbance which is usually associated with them.

DIAGNOSIS.

Complete absence of the uterus cannot be diagnosed with certainty in the living subject. A rudimentary condition may be present, and yet not be detected on the most careful examination. To examine cases in which this condition is suspected, we first pass a sound into the bladder and then with one or two fingers of the right hand in the rectum palpate the tissues which lie between the sound and the fingers. It is evident that in such a condition as is represented in fig. 150 the rudiment of the

Diagnosis
of absence
of Uterus.

¹ In twelve out of the eighteen cases collected by Pfannenstiel, to which Simon adds a thirteenth (*Centralb. f. Gyn.*, 1894, S. 1313).

² As in Gray's case (*Glas. Med. Journ.*, XXXI., p. 182) where an abortion took place in the sixth week of a normal pregnancy, and Ross's (*Edin. Med. Journ.*, 1885, p. 131) where there was a twin abortion in the sixth month and a full-time labour three months later.

³ *Glas. Med. Journ.*, XIX., p. 276.

The diagnosis of the one-horned uterus is not easy. The points to rely on are the following: the fundus turns to one side of the pelvis, is tapering, and has only one ovary connected with it. The rudimentary horn and the other ovary lie removed from it. Diagnosis of Uterus Unicornis.

The uterus didelphys is rare. A groove on the external surface of the uterus separating it into lateral halves, so that sounds can be passed into the separate cavities without coming in contact, indicates this condition. Of Uterus Didelphys.

The uterus bicornis is a comparatively frequent condition, and if Of Uterus Bicornis.




FIG. 159.

UTERUS SEPTUS (posterior view) FROM A WOMAN WHO DIED IN THE PUERPERIUM (*Cruveilhier*). The Uterine Cavity is divided by a septum which extends to the os externum. The left half (1) is strongly developed and contained the foetus. The right half (2) was empty.

well marked is easily recognised. Unusual breadth of the fundus, with a slight depression in the centre, points to a minor degree of this deformity.

The uterus septus is easily diagnosed if the septum extend as far as the os externum, so as to be within reach of the examining finger. If the septum does not extend so far, the condition may not be detected as there is no change in the external form to direct attention to the internal malformation. The sound may pass with equal ease into either cavity, or always into the same, and thus furnish no indication. In a case that Diagnosis of Uterus Septus.

came under our own observation the patient was examined frequently during life, bimanually and with the sound, and the uterus pronounced normal. At the post-mortem the external appearance of the uterus was normal; it was only on cutting into it that it was observed that the cavity was divided into two portions by a septum which extended to the os internum.

Of Infantile and Congenitally Atrophic Uterus.

The uterus infantilis and the congenitally atrophic uterus are recognised by their smallness. This is most distinctly made out with the finger in the rectum, the uterus being at the same time drawn down and fixed with the volsella. The well-developed vaginal portion and the unusual length of the cervix, as felt per rectum, enable us to diagnose the infantile from the congenitally atrophic uterus.

Differential Diagnosis.

With regard to *differential diagnosis*, gestation in a detached horn becomes a condition of great importance to the gynecologist when it simulates a fibroid tumour. The occurrence of irregular hæmorrhages from the empty uterine cavity, the absence of the foetal heart and uterine souffle when the foetus is dead, and the difficulty that there may be in palpating foetal parts, mask the existence of pregnancy. In the cases recorded by Angus Macdonald and Werth, the nature of the case was clear only on abdominal section; Macdonald draws attention to such cases as explaining the phenomena of "missed labour," the occurrence of which might sometimes give a clue.

PROGNOSIS.

Prognosis of Malformations.

In prognosis we must keep in view the possibility of ovulation with menstrual molimina, the secretion of menstrual blood and its accumulation in a closed cavity, the probability of conception and of gestation in an isolated horn. The most difficult cases are those in which the practitioner has to decide whether marriage is justifiable or not.

TREATMENT.

Treatment.

Malformations of the uterus lie beyond the range of treatment, except when they give rise to retention of menstrual blood or of the products of conception. The treatment of the former condition will be considered under Atresia of the Vagina (see Section VI.), and reference will be made to the latter in the chapter on Abdominal Section. Extirpation of the ovaries has been performed, and even of the uterus or its detached horn for dysmenorrhœa in cases of rudimentary uterus. Cases of congenital atrophy, associated with chlorosis, are amenable to treatment by feeding-up and iron.

CHAPTER XXVI.

SMALL OS EXTERNUM; RIGIDITY, STENOSIS, AND ATRESIA OF CERVIX.

LITERATURE.

Barnes—Diseases of Women: London, 1878, p. 245. *Burton*—So-called Obstructive Dysmenorrhœa: Brit. Med. Jour., 1884, II., 607. *Chrobak*—Die Untersuchung der weiblichen Genitalien: Billroth's Handbuch, Bd. I., S. 106. *Duke*—On the Rapid Dilatation of the Cervix Uteri (with discussion at Brit. Med. Ass.): Brit. Med. Jour., 1888, II., p. 873. *Duncan, Matthews*—On Sterility: Brit. Med. Jour., 1883, I., 702. *Goodell*—Rapid Dilatation of the Uterine Canal: Amer. Jour. Obstet., 1884, p. 1179. *Greenhalgh*—Intra-uterine Pessary: British Med. Jour., June 1878. *v. Grünewaldt*—Ueber die Sterilität geschlechtskranker Frauen: Archiv f. Gyn., VIII., 415. *Mackintosh*—Practice of Physic: London, 1836, p. 481. *Marckwald*—Ueber die kegelmantelförmige Excision der Vaginalportion und ihre Anwendung: Archiv f. Gyn., Bd. VIII., S. 48. *Müller*—Die Sterilität der Ehe: Billroth's Handbuch, Bd. I., S. 385. *Schultze*—Ueber Indication und Methode der Dilatation des Uterus: Wiener med. Blätter, 1879, Nos. 42, 43, 44, 45. *Simpson, Sir J. Y.*—Diseases of Women: Edinburgh, 1872, p. 245. *Sims, Marion*—On the Surgical Treatment of Stenosis of the Cervix Uteri, and Discussion: Am. Gyn. Trans., 1878, p. 54. *Vedeler*—Ueber Dysmenorrhœe: Archiv für Gyn., XXI., 211.

ETIOLOGY AND PATHOLOGY.

THE various conditions treated of in this chapter have been described mainly from clinical observation and in relation to the symptoms of ^{Etiology} and ^{Pathology}.




FIG. 160.

A NORMAL AND A PIN-HOLE OS, as seen in the SPECULUM (*Schroeder*).

dysmenorrhœa and sterility. Owing to the absence of exact data, there has been room for great difference of opinion as to the pathology and frequency of these conditions.

Small Os Externum.—In a certain number of cases, 6-9 p. c. (Vedeler), the os externum is smaller than the normal size; it may be so narrow as to admit only a fine probe (pin-hole os). The contrast between this and the normal os is shown in fig. 160. The cervix is conical in form (fig. 161) and of unusually firm consistence; sometimes it is hypertrophied, the vaginal portion measuring as much as two inches. The cervical mucous membrane is frequently in a condition of catarrhal inflammation; according to Von Grünewaldt, the conical shape of the cervix is often the result of the accumulation of mucus.

Rigidity of Cervix.—The changes in the cervix resulting from an increase of its connective tissue have been fully described by Scanzoni. A peculiarly rigid condition of the cervical tissue, apart altogether from any contraction of the canal, is observed on passing bougies in cases of dysmenorrhœa (Matthews Duncan). A similar condition has been noted as specially frequent in cases of sterility (Olshausen, Martin and Chrobak).




FIG. 161.

CONICAL VAGINAL PORTION (Barnes).

Stenosis (contraction) of the cervical canal is congenital or acquired. As a *congenital* condition affecting the cervical canal throughout its whole extent, it is a comparatively rare occurrence. It is always associated with smallness of cervix and body, pointing to generally defective development of the uterus (which is further indicated by the scantiness of menstruation). The commonest cause of the *acquired* form is cicatrization—after labour, after amputation of the cervix, or after the repeated application of strong caustics; the last is perhaps the most frequent. Inflammation of the mucous membrane, resulting in adhesions, also produces it.

Atresia of Cervix (ἀ-τρήσις, non-perforation), or occlusion of the canal, is rare as a *congenital* condition, and is due to the presence of a cap of tissue covering the os uteri. The canal is seldom, if ever, imperforate throughout its course. An incomplete transverse septum has been described in a few cases.¹

¹ Budin—Progrès Médical, 1887.

It is more frequently *acquired*, and results from the following causes:—sloughing and cicatrization after labour; cicatrization after the application of caustics, and after amputation of the cervix; adhesion of granulations in cervical catarrh (after menopause), and round the base of tumours.

The practical point for the practitioner to remember is that atresia may follow the repeated application of caustics and amputation of the cervix. It occurs also as part of the physiological changes which take place after the menopause. Twenty-eight per cent. of women above fifty years of age have atresia of the cervix (Hennig).

SYMPTOMS AND DIAGNOSIS.

The symptoms found most frequently associated with these conditions Symptoms.
are—

Dysmenorrhœa,
Sterility.

We say “associated,” because the relation of the symptoms to the pathological condition is as yet not known. There is no subject in Gynecology round which more discussion has raged, and concerning which there are at present more abrupt differences of opinion.

Dysmenorrhœa.—Mackintosh, from a doubtful analogy between the menstruating uterus and the bladder, introduced dilatation with bougies as a treatment of dysmenorrhœa. The theory was that a stricture prevented the discharge of blood in the former case, just as it prevents a discharge of urine in the latter; and that the pain was due to uterine efforts to overcome obstruction. Sir James Simpson showed that stenosis could not be the only factor, since obstructive dysmenorrhœa might be equally present with a patulous cervix; it depended also on the amount of the menstrual discharge and the danger of its clotting while in the uterus, and may be absent where, though the os is small, the flow is scanty. Marion Sims took up the position that painful menstruation was almost wholly due to mechanical causes, and was the great exponent of what is known as “the mechanical theory.” On the other hand, Matthews Duncan, maintained that he had never seen a pin-hole os in cases of dysmenorrhœa; and attributed the pain to irregular contractions of the uterus which had nothing to do with expulsion of its contents. Vedeler’s investigations have shown that a small os externum is as common in patients without as in those with dysmenorrhœa. Emmet, at a discussion on Sims’ Operation before the American Gynecological Society, characterised the mechanical theory of dysmenorrhœa as a myth; in his Gynecology, he says that, unless the flow is scanty, painful menstruation is accompanied by clots, but that their formation does not depend upon obstruction.

Hitherto, conclusions have been drawn almost entirely from the condition of the uterus and cervix between the menstrual periods; and it will be evident from the foregoing how wide is the difference of opinion on the subject. It seems to us that valid conclusions can only be drawn from the condition of the cervix *during menstruation*, and that the diversity of opinion will remain until we have accurate knowledge on this point.

We have called the condition "Small Os Externum" instead of "Stenosis" advisedly; as the latter word implies that there is resistance to the outflow of blood, while the as yet scanty evidence rather seems to show that the canal becomes more patulous during menstruation than at any other time.

Relation of
Stenosis to
Sterility.

Sterility.—When we come to treat of sterility, we shall find that it is frequently associated with dysmenorrhœa. According to the statistics given by Matthews Duncan, as well as those by Marion Sims and Emmet, about one-half of cases of sterility suffer from severe dysmenorrhœa; and two-thirds of Vedeler's cases of dysmenorrhœa in married women were sterile. A narrow os externum, according to the mechanical theory, hinders the upward passage of the spermatozoa just as it retards the downward flow of the menstrual blood. This explanation is evidently open to the criticism that the spermatozoa are microscopic; and that, as Fritsch puts it, a drop of water will fall as easily through a ring of 2 cm. diameter as through a hoop of 100. It is, however, quite possible that a narrow os externum while not absolutely preventing conception may retard it: Müller, in enforcing the very important distinction between absolute and relative sterility, thinks that a contracted os may render conception more difficult, especially where the spermatozoa are scanty in the spermatic fluid. Thus, a counter-illustration to Fritsch's would be that where the drops are few there is more chance of catching them in a bowl than in a thimble. Although there is a general reaction against stenosis *per se* as a cause of sterility, cervical catarrh, when associated with it, is considered by many to play an important rôle through stagnation of the mucous secretion. It has not, however, been proved that a plug of mucus can be an effectual bar to the progress of spermatozoa, and catarrh is a very frequent condition in parous women.

A rigid condition of the cervix has, as already said, been frequently noted as present in cases of sterility. Matthews Duncan suggested that it operates through checking spontaneous dilatation of the cervix during coition.

In studying the complex question of sterility (*v.* Section IX.), the at first too obvious mechanical causes sink into insignificance as soon as we come in sight of the less obtrusive and more subtle physiological and vital considerations; and, after a careful survey of the literature, we

come to the conclusion that any discussion of sterility in which mechanical considerations have a prominent place must be inadequate and will always be bootless.

DIAGNOSIS.

A history of dysmenorrhœa and sterility will lead us to suspect that one of these conditions of the cervix may be present. On vaginal examination, the finger recognises the conical shape and firm consistence of the cervix. In cases of small os externum, the first impression is that it is altogether absent; but more careful examination detects a slight depression. The speculum shows the appearance represented in figs. 160 and 161. The sound is passed with difficulty: but we must remember that difficulty in passing the sound is quite unreliable as a test of the canal's being relatively narrower at a given point; a sharp flexion, a projecting tumour or even a fold of mucous membrane may arrest the sound. Burton by passing the sound in six cases of dysmenorrhœa during the height of the pain made the interesting observation that the canal was more patent then than at any other period.

Diagnosis
of Stenosis
of the
Cervix.

PROGNOSIS.

This must always be guarded, as the etiological relationship between the conditions of the cervix described and these symptoms is still *sub lite*, and the results of our empirical treatment correspondingly uncertain.

Prognosis.

TREATMENT.

The methods of treatment are—

- A. Dilatation.
- B. Division.

Dilatation for stenosis is carried out by passing graduated bougies, by sponge or laminaria tents, by forcible dilatation with instruments. Division is effected by the metrotome or by scissors. We here consider only dilatation for stenosis; its use for intra-uterine medication will be dealt with under the treatment of Endometritis.

A. Dilatation.

Sponge and laminaria tents were formerly used, but are now abandoned because of the dangers of septicæmia.

Treatment
of Stenosis
by Dilata-
tion.

Dilatation by means of *graduated bougies* was brought into prominent notice by Mackintosh, who employed straight metallic bougies of different degrees of thickness. He passed first a small one not thicker than a probe, and then larger ones till the os was rendered quite patu-

lous. A No. 9 bougie is the largest size which will pass through a virgin cervix. We have, therefore, to begin with one of smaller calibre, say 6 or 7, and go up to a No. 11 or 12, as the cervix must be over-stretched to effect a cure. Hegar's dilators (see p. 144) are also used in the same way.¹

Various dilators with *expanding blades* have been devised. Fig. 162 shows the form used by Schultze. He dilates the cervical canal beforehand with laminaria; he then washes it out with a 2 per cent. solution




FIG. 162.

SCHULTZE'S DILATOR.

of carbolic acid, as he attributes many of the serious consequences of forcible dilatation and incision to the absorption of the secretions. The dilator is now introduced, and the blades (which open antero-posteriorly) are forcibly separated. Ellinger has made a dilator so constructed that the blades remain parallel to one another while being separated; Goodell has had very good results from forcible dilation with this instrument both with regard to Dysmenorrhœa and Sterility. The dilator em-




FIG. 163.

MARION SIMS' DILATOR (Sims).

ployed by Marion Sims is seen at fig. 163. Other forms have been introduced by Reid, Duke, and More Madden.

B. Division.

Treatment
of Stenosis
by Divi-
sion.

Sir James
Simpson's
Metro-
tome.

Division of the cervix with the knife was introduced by Sir James Y. Simpson. The instrument which he devised for this purpose was the *metrotome*—a bistoury caché, with a single blade sharp on the outer edge which is unsheathed on compressing the handle. A screw on the handle regulates the extent to which the blade is to be protruded.

¹ Anvard has devised tubular dilators, so constructed that the next size larger can be slipped over the previous one *in situ*: *Archiv de Toc.*, 1894, p. 814.

The effect of division is that the narrow circular os becomes an orifice with gaping lips, and thus resembles that of a multipara (*cf.* figs. 164 and 165).

The objection to the metrotome is that we do not know how deep the incision is being made, and it has been abandoned for the scissors.




FIG. 164.

NULLIPAROUS OS UTERI (*Sir J. Y. Simpson*).




FIG. 165.

PAROUS OS UTERI (*Sir J. Y. Simpson*).

Kuchenmeister has devised scissors with a hook in the external blade to keep it from slipping off the cervix, but a pair of ordinary scissors does perfectly.

The operation is performed as follows. The patient is placed semi-prone. The Sims speculum is passed, and held by an assistant. The vagina should be thoroughly syringed beforehand with an antiseptic. The anterior lip of the cervix is laid hold of with the volsella; the

Operation
for Bilateral
Division of
Cervix.




FIG. 166.

SHOWING THE BILATERAL DIVISION OF THE CERVIX, with Kuchenmeister's Scissors (*Barnes*).

scissors are introduced, the straight blade being passed within the cervical canal; the point or hook of the external blade is carried to about one-third up the vaginal portion of the cervix (see fig. 166) and the section made. In many cases, all that is necessary is to divide the ring round the os externum; when this is divided the cervical canal is sometimes found to be dilated above it. Should hæmorrhage occur, some perchloride of iron is swabbed on the cut surface, and a vaginal tampon of lint soaked in an antiseptic is applied.

As will be evident from what has been said under Symptoms, the scope of this operation is very limited unless we have recourse to it as a stage in treating cervical catarrh in a nullipara.

Treatment after Bilateral Division of Cervix. *More important than the incision is the after-treatment.* The patient must be seen on the following day, and every second day for a fortnight, and the finger passed in on each occasion to prevent union of the cut surfaces and dilate the cervical canal.

Excision of a portion of the cervix is also done with a view to convert the stenosed into a gaping os like that of a multipara. The operation is similar to that done in amputating the cervix (*v. fig. 172*).

Treatment of Atresia. *Atresia of the cervix* is chiefly of importance in regard to the accumulation of menstrual blood or mucus above the obstruction. It is this which produces the Symptoms and calls for Treatment. It will be better to defer the consideration of these till we treat of Atresia Vaginæ (Section VI.).

CHAPTER XXVII.

ATROPHY OF THE CERVIX AND UTERUS: SUPERINVOLUTION.

WE meet with an atrophic condition of the cervix and uterus under five different conditions:—

1. As a congenital condition ;
2. Associated with certain constitutional affections, as phthisis, scrofula, chlorosis ;
3. In acute general affections as scarlet, and typhus fever, and local conditions as parametritis atrophicans ;
4. In the puerperal uterus, as the result of superinvolution ;
5. After the menopause.

Conditions
under
which
Atrophy
of Uterus
occurs.

According to Gottschalk,¹ atrophy occurs physiologically (superinvolution) or pathologically ; and the latter is primary (the cause being uterine, *e.g.*, endometritis, presence of tumours, parametritis posterior), or secondary to affections of the tubes and ovaries or more general conditions.

Should the student find on vaginal examination that the cervix is small and projecting only slightly into the vagina, and on bimanual examination that the body of the uterus is found with difficulty and is smaller than it should be, he must next ascertain which of the above-mentioned causes has produced the atrophy.

The history will enable him to form his diagnosis. With the *congenital condition* there is a history of amenorrhœa or scanty menstruation since puberty, of sterility if the patient has entered married life, and of hysteria and other disturbances of the nervous system which usually accompany imperfect development of the uterus. The *constitutional condition*, and especially the state of the blood and of the lungs, in other cases enables him to account for the condition of the uterus. Probably the small uterus found in chlorotic patients is a congenital condition, and not secondary to the constitutional state. If the atrophic condition be the result of *superinvolution*, there is a history of childbirth or abortion with

¹ Beitrag zur Lehre von der Atrophia Uteri : Samml. klin. Vorträge N.F., No. 49, 1892.

non-appearance of menstruation after it. With regard to the *menopause*, the age of the patient is the chief guide; we must remember the possibility of an early menopause, as early as at the age of thirty-five.

The only atrophic condition which we shall consider here is that occurring in the puerperal uterus as the result of superinvolution.

SUPERINVOLUTION OF THE UTERUS.

LITERATURE.

Engström—Zur Kenntniss der puerperalen Hyperinvolution der Gebärmutter: Festsch. d. 50 jähr. Jub. d. Gesell. f. Geb. u. Gyn. zu Berlin. *Frommel*—Ueber puerperale Atrophie des Uterus: Zeits. f. Geburts. und Gynäk., Bd. vii., H. 2, S. 305. *Jaquet*—Ueber Atrophia Uteri: Berl. Beiträge zur Geburts. und Gynäk., Bd. ii., S. 3. *Johnson, T. J.*—Superinvolution of the Uterus: Am. Gyn. Trans., 1883, p. 1064. *Klob*—Patholog. Anatom. der weib. Sexualorgane: Wien, 1864, S. 205. *Ries*—Ueber die Atrophie des Uterus nach puerperaler Erkrankung: Zeits. f. Geb. u. Gyn., Bd. xxvii., S. 38. *Simpson, A. R.*—Superinvolution of the Uterus: Edin. Med. Jour., May 1883. *Simpson, Sir J. Y.*—Morbid Deficiency and Excess in the Uterus after delivery: Selected Obstetrical and Gynecological Works, 1871, p. 595. On Superinvolution of the Uterus and Amenorrhœa: Diseases of Women, Edin., 1872, p. 597. *Thorn*—Zur Laktationsatrophie des uterus: Zeitsch. f. Geb. u. Gyn., Bd. xvi., and Centralb. f. Gyn. 1894, S. 716.

PATHOLOGY.

The uterus is small. Its external length may be reduced from the normal 3 to $1\frac{3}{4}$ inches. The walls are thin and flaccid, sometimes of a dense and fibrous consistence. The vaginal portion projects only slightly into the vagina, and may be almost flush with the vaginal roof. The os may be relatively patulous, or contracted so as only to admit a probe. The uterine cavity is reduced to $2\frac{1}{4}$, 2, or even $1\frac{1}{2}$ inches in length. The ovaries are atrophied, and sometimes show an increase of fibrous tissue in their structure. The accompanying specimen (fig. 167), described by Sir James Simpson, illustrates these points.

Ries found the mucous membrane atrophied or absent, and a small-celled infiltration of the muscular tissue. In one case the muscular fibre was in great part replaced by connective tissue and thrombi. His preparations were taken from cases of death from puerperal sepsis in which the changes may have been more pronounced than usual.

Engström distinguishes an excentric atrophy in which the walls are thin and flabby, but the cavity not diminished, from a concentric in which the whole organ is shrunk.

ETIOLOGY.

As to the frequency of this condition, A. R. Simpson found it present in 22 out of 1300 cases, that is in about 1·7 per cent.; Frommel estimates its frequency at 1 per cent. The reason why, in certain cases,




FIG. 167.

PREPARATION OF UTERUS AND OVARIES IN A CASE OF SUPERINVOLUTION, *ad naturam*. Weight of parts represented—one ounce, four drachms, twenty-five grains. Uterine cavity measures $1\frac{1}{2}$ in. Thickness of posterior uterine wall (laid open in figure) $\frac{1}{4}$ in. Tissue of uterus, dense and fibrous. Ovaries atrophied, with increase of fibrous tissue and no appearance of Graafian vesicles. (Sir J. Y. Simpson.)

the process of involution during the puerperium goes on till the uterine cavity is reduced to less than $2\frac{1}{2}$ inches in length is not known. A condition of transitory superinvolution is according to Thorn not infrequent. During lactation the uterus becomes atrophied due to a reflex tropho-neurosis, and returns to its normal size afterwards. Engström does not attach much importance to the direct stimulus from nursing, but accounts for atrophy whether transient or permanent by general anæmia which he found present in almost all his cases. In some instances there is a history of great *loss of blood* at the confinement (A. R. Simpson). In other instances *pelvic peritonitis* has occurred during the puerperium: this can produce, we know, atrophy of the ovary through binding it down with adhesions; and atrophy of the ovaries may lead to atrophy of the uterus. As already said *protracted lactation* is also a factor, and any conditions which lead to *Anæmia*. It is also associated with the *tubercular diathesis* (Klob).

The term superinvolution has also been applied to atrophy of the uterus following hypertrophy from causes other than pregnancy, *e.g.*, sub-mucous fibroids, and that following operations on the cervix, but it is best to limit it to cases of atrophy after parturition.

SYMPTOMS.

Continued amenorrhœa is the symptom which leads the patient to seek advice. After she has ceased nursing, she expects the flow to return. It does not do so, however, even after months have passed. Pain in the back, weakness, and hysterical symptoms are sometimes present.

DIAGNOSIS.

The small cervix at once suggests what the condition is. We sometimes have difficulty in making out the uterus bimanually; here the examination per rectum, combined with the volsella, is useful. The best idea of the size of the uterus is gained by pressing the ball of the finger in the rectum against the isthmus of the uterus, and then moving the uterus upwards and downwards upon the finger which can thus estimate accurately its size; having done this, we make more traction on the uterus to bring it as far down as possible, and examine the ovaries.

The sound must be used with care, as it easily perforates the thin walls of the uterus. It does not pass into the uterus as far as the $2\frac{1}{2}$ in. knob.

Differential diagnosis must be made from—

- Congenital malformation;
- Congenital atrophy;
- Senile atrophy.

PROGNOSIS.

This should always be guarded. The curability of the case depends, as Fordyce Barker has pointed out, on the condition of the ovaries—a point, however, exceedingly difficult to determine. When the patient has the menstrual molimina and the menstruation though scanty, still persists, we may hope for improvement even though the uterus is small.

TREATMENT.

From the unsatisfactoriness of treatment, such cases may, as a rule be left alone. Iron and constitutional remedies may be tried, and permanganate of potash as in other cases of amenorrhœa.

Local remedies as douching, galvanism, and massage have been recommended. The use of intra-uterine stem pessaries, introduced by Sir J. Y. Simpson, has been abandoned from the risks attendant on it.

CHAPTER XXVIII.

HYPERTROPHY OF THE CERVIX: AMPUTATION.

LITERATURE.

Byrne—Amputation and Excision of the Cervix Uteri : Trans. Americ. Gyn. Soc., Boston, II., pp. 57 and 110. *Dührssen*—Beitrag zur Anatomie, Physiologie und Pathologie der Portio vaginalis uteri : Archiv f. Gyn., XLI., S. 260. *Galabin*—Lond. Obst. Journ., Sept. 1878. *Goodell*—Clinical Notes on the Elongations of the Cervix Uteri : Am. Gyn. Trans., 1880, p. 268. *Hegar und Kaltenbach*—Operative Gynäkologie : Stuttgart, 1881, S. 445. *Huguier*—Mémoires sur les allongements hypertrophiques du col de l'utérus : Paris, 1860. *Leblond*—Operative Gynécologie : Paris, 1878. *Marckwald*—Ueber die kegelmantelförmige Excision der Vaginalportion, etc. : Archiv f. Gyn., Bd. viii., S. 48. *Müller*—Die Amputatio Colli Uteri : Zeitschrift für Geburt. und Gyn., Bd. ix., S. 178. *Perdrizet*—Des amputations du col de l'utérus : Archiv de Toc., Aug. et Sep. 1894. *Schroeder*—Charité-Annalen, 1878. Zur Technik d. plast. Op. am Cervix Uteri : Zeitschrift für Geburt. u. Gyn., Bd. iii., S. 419; Bd. vi. Hft. 2, S. 218. *Simon*—Monatsch. f. Geburtskunde, xiii., S. 418. *Sims, Marion*—Uterine Surgery, 1866. *Stratz*—Ueber einseitige Hypertrophie des untern Cervicalabschnitts : Zeits. für Geb. und Gyn., Bd. xii., S. 229.

HYPERTROPHY of the whole uterus occurs in two forms :—

1. Hypertrophy of the muscular tissue—in pregnancy ;
2. Hypertrophy of the connective tissue—in subinvolution and chronic metritis, both of which will be considered under Chronic Metritis (Chap. XXXII.).

Hypertrophy of the cervix alone calls for special notice here.

HYPERTROPHY OF THE CERVIX.

Under this head we consider two conditions :—

- A. Hypertrophy limited to the vaginal portion, which is a distinct *primary* lesion ;
- B. Hypertrophy of the supra-vaginal portion, which is usually associated with hypertrophy of the body of the uterus ; this occurs in prolapsus uteri and is probably *secondary* to that condition.

A. HYPERTROPHY OF THE VAGINAL PORTION.

Pathology.—The characteristic of this condition of the cervix is a great increase in length affecting it equally all round¹ (fig. 169). The mucous membrane and the subjacent tissue are not thickened, so that the diameter of the cervix is not much increased. As the result of the increase in length, the conical apex of the cervix comes to lie immediately behind the hymen, and may protrude through the vaginal orifice (fig. 168). The os externum is often small.

Etiology.—This condition is a true hypertrophic growth; it is not very common and the cause of it is unknown. As it occurs in the




FIG. 168.

HYPERTROPHIED VAGINAL PORTION *c* PROTRUDING THROUGH THE VULVA. The Sound has passed very far into the small os *o* (Schroeder).

virgin, it is probably congenital.² Sometimes it does not attract attention till the patient enters married life, when it produces as a rule sterility because the form of the cervix interferes with conception.

The cervix is frequently *thickened* as the result of chronic inflammation consequent on its laceration in childbirth; this is not a true hypertrophic growth, and will be considered under Laceration of the Cervix (Chap. XXIX.).

¹ Only one case of unilateral hypertrophy in a nullipara could be found by Stratz in the literature—a case recorded by Huguier. Partial hypertrophies are less rare in multiparæ and will be referred to under Laceration of the Cervix.

² As in a case recorded by Thomson: *Cent. f. Gyn.*, 1895, S. 415.

Symptoms.—The symptoms are due to the presence of the hypertrophied cervix in the vagina. There is bearing-down as in prolapsus uteri, irritation of the mucous membrane of the vagina and consequent leucorrhœa, discomfort on walking about and on rising suddenly. If the cervix protrude beyond the vulva, ulceration of its mucous membrane and excoriation are produced.

Diagnosis.—This presents no difficulty. The fornices are found in their normal position on vaginal examination (see fig. 169), the fundus uteri at its normal height in the pelvis on bimanual examination. These two clinical facts indicate that the low position of the apex of the cervix is not due to a descent of the fundus but to a hypertrophy of the cervix, and that the hypertrophy of the cervix is limited to the portion which projects into the vagina (*cf.* fig. 169 with fig. 175 and fig. 176).




FIG. 169.

HYPERTROPHY OF VAGINAL PORTION OF CERVIX. Neither fornix is obliterated (*Schroeder*).
Section of Pelvis seen in fig. 168.

The sound may pass five inches or more into the cervical canal; as the patient is usually a nullipara and the abdominal walls therefore firm, it facilitates the Bimanual to do it with the sound in the uterus. The combined recto-vaginal examination shows that the uterus, above the vagina is of normal length.

Treatment.—This consists in amputation of the cervix which is the only course open to us, because the hypertrophy will not diminish but rather increase. Amputation is performed by three methods:—

1. Scissors or knife,
2. Ecraseur,
3. Galvano-caustic wire.

In amputation *with the knife*, we may employ either the circular

method (Sims' and Hegar's, fig. 170), or the flap amputation by wedge-shaped excision of the anterior and posterior lips separately (Simon and Marckwald, figs. 171, 172).

The operation.—The instruments required are the following :—

Antiseptic douche,	Dissecting forceps,
Speculum and spatulæ,	Strong full-curved needles and
Volsellæ,	needle-holder,
Knives and scissors,	Silk and catgut sutures.

Amputa-
tion of the
Cervix for
Simple
Hyper-
trophy.

The patient is placed in the lithotomy posture. The external genitals and vagina are thoroughly cleansed. In this as in all operations on the cervix, continued irrigation with an antiseptic douche is an advantage. The speculum and spatulæ are necessary to expose the cervix thoroughly and give room for operative manipulation. It also makes operating easier to have the cervix well drawn down with volsellæ; but




FIG 170.

Circular amputation (*Hegar u. Kaltenbach*).

in doing this we must be guided by the mobility of the uterus. In all such operations a careful bimanual should be made just before operating, to determine the condition of the parts round the uterus which can be much more satisfactorily ascertained under the anæsthetic.

The cervix is amputated by the circular or flap method, the latter being the one usually employed. In the former the vaginal mucous membrane is stitched to the cervical all round (fig. 170). In the latter the cervix is split into an anterior and posterior lip, a wedge-shaped portion is cut out of each (figs. 171 and 172); cervical mucous membrane is stitched to vaginal in the centre, and vaginal to vaginal at the sides (figs. 173 and 174). Silk may be used for the central sutures; but the advantage of using catgut throughout is that the patient is saved the inconvenience of having the stitches removed. If silk is used, they are taken out in a week.

Amputation with Ecraseur or Galvano-caustic wire.

Amputation with the *Ecraseur* or with the *Galvano-caustic wire* is not such a neat method of operating as with the knife, and there is liability




FIG. 171.

SPLITTING THE CERVIX into an anterior and a posterior lip (*Schroeder*).




FIG. 172.

EXCISION OF WEDGE from each lip.

to closure of the cervical canal through cicatrisation. The method of using these will be described under amputation of the cervix for carcinoma (see Chap. XLII.).




FIG. 173

POSITION OF SUTURES before they are tied.




FIG. 174.

APPEARANCE OF STUMP when Sutures are tied.

B. HYPERTROPHY OF THE SUPRA-VAGINAL PORTION.

Diagnosis of Hypertrophy limited to supra-vaginal portion of Cervix.

The existence of hypertrophy limited to the supra-vaginal portion of the cervix and not affecting the body of the uterus cannot be determined by *clinical* examination alone. The obvious reason is that we have no means of ascertaining in a case of hypertrophy where the precise upper limit of the cervix lies. The position of the os internum cannot be

learned from the sound, and the distance to which the utero-vesical pouch of peritoneum descends can only be ascertained on post-mortem




FIG. 175.

HYPERTROPHY OF INTERMEDIATE PORTION OF CERVIX. The anterior fornix is obliterated (*Schroeder*).

examination. We cannot affirm, therefore, that the hypertrophy is limited to the supra-vaginal portion of the cervix and that it does not affect the body of the uterus as well.




FIG. 176.

HYPERTROPHY OF SUPRA-VAGINAL PORTION OF CERVIX. Both fornices are obliterated (*Schroeder*).

In the present state of our knowledge it is impossible to say whether this hypertrophy is primary or secondary. We believe that in the great proportion of cases it is secondary to prolapsus uteri. It has also been

described as an exceptional occurrence in the early months of pregnancy.¹

By French and by many German gynecologists, however, hypertrophy of the supra-vaginal portion of the cervix is considered a distinct primary lesion. Huguier first drew attention to the increase in the length of the uterine canal in cases described as prolapsus uteri; he affirmed that the fundus uteri always remained in its normal position, and that the os externum came to lie outside the vulva *because* the




FIG. 177.

PROLAPSUS UTERI WITH CERVICAL ELONGATION (*Barnes*); *p, p*, peritoneum.

cervix had increased in length; this hypertrophied condition of the cervix was occasioned by a prolapse of the vaginal walls which made traction on the cervix, and thereby stimulated it to increased growth.

By these gynecologists, three forms of cervical hypertrophy are described according to the portion of the cervix which is hypertrophied. The division of the cervix into three portions—a vaginal, an intermediate, and a supra-vaginal portion—has been already described (see page 18).

Three
forms of
Cervical
Hyper-
trophy.

¹ By Martin—Berliner Gesellschaft f. Geb. u. Gyn., 1880.

The vaginal portion is limited superiorly by the insertion of the anterior fornix; the intermediate by that of the posterior fornix; the supra-vaginal by the os internum. Hypertrophy of the vaginal portion is characterised by the *persistence of both* fornices in their normal position; it has been already described (see fig. 169). In hypertrophy of the intermediate portion the posterior fornix remains, while the *anterior is obliterated* (see fig. 175). In hypertrophy of the supra-vaginal portion *both anterior and posterior fornices are obliterated* (see fig. 176).

Treatment.—While hypertrophy limited to the vaginal portion of the cervix is very rare, that affecting the whole cervix and usually associated with prolapsus uteri is a common condition, and it was for it that the various modes of amputating the cervix were introduced.

Huguier, who first exactly described supra-vaginal hypertrophy, introduced the *conoid amputation*. One incision is made from the posterior fornix obliquely upwards and forwards as far as the cervical canal; a second is made from the anterior fornix upwards and backwards to meet the latter; by this means a wedge-shaped or conical piece of the supra-vaginal portion of the cervix is removed.

The flap operation already described, however, gives the best stump. In amputating for supra-vaginal hypertrophy, the *relations of bladder and peritoneum* of the pouch of Douglas require to be considered (fig. 177). The bladder invariably descends for a varying distance in relation to the front of the hypertrophied cervix. The peritoneum of the pouch of Douglas, inasmuch as it lines the upper part of the posterior vaginal wall, will, when that wall is everted, dip down alongside of the hypertrophied cervix. If the posterior fornix is not obliterated, the peritoneum will not descend alongside of the protruding cervix.

After drawing the cervix down, a sound is passed into the bladder to define its position. The line of reflection of the posterior vaginal wall indicates how much is vaginal portion, and by cutting within that we avoid the pouch of Douglas. If it is cut into, it should be closed with a buried catgut suture.¹

¹ Dührssen advises the use of a buried suture in all cases to prevent retraction of the elastic-tissue layer of the cervix and to lessen the risk of sepsis; but we see no advantage in this, unless the pouch of Douglas be cut into.

CHAPTER XXIX.

LACERATION OF THE CERVIX AND ITS CONSEQUENCES.

LITERATURE.

- Baer*—Analysis of Twenty-seven Operations for the Restoration of the Lacerated Cervix Uteri, with special reference to the effect on sterility and labour; *Amer. Jour. Obstet.*, 1883, pp. 295, 852. *Coe, H. C.*—The present status of Emmet's operation for laceration of the Cervix Uteri: *New York Journ. of Gyn. and Obst.*, Tome iii., p. 453. *Czempin*—Risse des Cervix Uteri ihre folgen und operative Behandlung: *Zeit. für Geb. und Gyn.*, Bd. XII., S. 287. *Emmet, T. A.*—Surgery of the Cervix: *Am. Jour. of Obst.*, Feb. 1869; *ibid.*, Nov. 1874. *American Practitioner*: Indianapolis, Jan. 1877. *Principles and Practice of Gynæcology*, p. 440: Philadelphia, 1884. *Emmet, J. D.*—Emmet's operation for laceration of the Cervix Uteri, its scope and limitations: *Am. Journ. of Obst.*, April, 1895. *Eustache*—La Trachelorrhaphie ou opération d'Emmet: *Archiv de Tocolog.*, 1884, p. 684. *Goodell*—Notes of one hundred and thirteen cases of operation for laceration of the cervix: *Am. Journ. of Obst.*, Jan. 1882. *Houzel*—Note sur l'Opération d'Emmet ou Trachelorrhaphie: *Annal. de Gyn.*, XXX., 241 and 351. *Howitz*—Gynäkologische og. Obstetriciske Meddelelser, Bd. I., Heft 3. *Kaltenbach*—Ueber tiefe Scheiden- u. Cervicalrisse bei der Geburt: *Zeitschrift f. Geb. u. Gyn.*, Bd. II., S. 274. *Klein*—Prager med. Wochenschrift, 1878, No. 24. *Mundé*—*Am. Jour. of Obst.*, Jan. 1879; Death after Trachelorrhaphy, *New York Med. Jour.*, 1883, p. 332. *Murphy*—Observations on the effects of Trachelorrhaphy on Fertility and Parturition: *Amer. Journ. of Obstet.*, Jan. 1883. *Nieberding*—Ueber Ectropium und Risse: Würzburg, 1879. *Noeggerath*—Cervixrisse und Erkrankungen der Gebärmutter: *Berl. klin. Wochenschr.*, 1887, No. 41; and (for Discussion) *Archiv f. Gyn.*, Bd. XXXI., S. 469. *Olshausen*—Zur Pathologie der Cervicalrisse: *Centralbl. f. Gyn.*, 1877, Nr. 13. *Pallen*—*St. Louis Med. and Surg. Jour.*, May 1868; *Richmond and Louisville Med. Jour.*, 1874; *British Med. Jour.*, May 1881. *Park*—Trachelorrhaphy: *Edin. Med. Jour.*, XXXIII., II., p. 1130. *Playfair*—Trachelorrhaphy: *Lond. Obstet. Trans.*, 1883, and *Medical Times and Gazette*, 1885, I. 340. *Roser*—Das Ectropium am Muttermund: *Arch. f. Heilk.* II. Jahrg., 2 H., 1861. *Ruge u. Veit*—Zur Pathologie der Vaginalportion; *Stuttgart*, 1878. *Sänger*—Ueber die klinische Bedeutung der Cervixrisse: *Centralbl. f. Gyn.*, Bd. XII., S. 441. *Simpson, Sir J. Y.*—*Edin. Jour. of Med. Science*, 1851, p. 152. *Spiegelberg*—Ueber Cervicalrisse, ihre Folgen u. ihre operative Beseitigung: *Breslauer ärzt. Zeitschrift*, 1879, No. 1. *Van de Warker*—Etiology of Lacerations; *Am. Journ. of Obst.*, Jan. 1882. Thirty-one operations for Repair of Laceration of the Cervix Uteri: *ibid.*, July 1883. *Vulliet*—Douze cas d'opération de Trachelorrhaphie: *Archiv de Tocolog.*, 1884, p. 1000. *Wells, B. H.*—Possible Dangers of Trachelorrhaphy: *Amer. Jour. Obstet.*, 1884, p. 561. Also, the Etiological Relation of Cervical Laceration to Uterine Disease: *Amer. Jour. Obs.*, 1888, p. 257.

THE student will not have gone far in the clinical study of Gynecology without being surprised at the large number of patients who refer the commencement of their illness to a confinement or miscarriage. They come complaining of various ailments—a weak back, pain in the side, white discharge, losing too much at the monthly time, or general unfitness for work. On physical examination, he finds a variety of conditions—a fissured and thickened velvety cervix, thickenings in the lateral fornices or behind the uterus often displacing it by traction, and the uterus itself enlarged. We do not mean that all of these are present in one case, but that one or more of them may be; nor is any one symptom invariably connected with one lesion. He asks himself why labour is so often the starting-point of female complaints; and one important reason, though by no means the only one, is that the tear of the cervix in labour literally opens the door to a variety of lesions. Cervical catarrh is favoured, if not started (as Emmet says), by the split condition of the cervix; the raw surface has admitted septic matter which leads to chronic inflammation of the parametrium with all the changes in the train of parametritis; and sub-involution is kept up (if not directly by the tear, as Emmet holds) indirectly by the consequent parametritis which Freund has shown to affect the venous and lymphatic circulation in the uterus. It is impossible to consider laceration of the cervix separate from the results which in the great majority of cases follow, and hence this chapter deals with “Laceration of the Cervix and its Consequences.” Many of these latter being distinct lesions in themselves, will be treated of separately in the following chapters and only referred to here in their relation to laceration as an antecedent.

For the recognition of laceration of the cervix as a distinct and important lesion we are indebted to the genius of Emmet of New York, who was the first to insist on its clinical significance and elaborate an operation for its treatment.

J. H. Bennet of London had previously described the changes produced in the cervix by its laceration in labour, unfortunately attributing them to a process of ulceration. Roser of Marburg had described the pathology of the condition; but its importance as a factor in uterine disease was brought into notice by Emmet's first paper which was published in 1869, seven years after he had introduced his operation. Emmet's views as to the importance of lacerations of the cervix have given rise to a great deal of discussion; and their significance is still a *questio vexata* in Gynecology.

PATHOLOGY.

The commonest *seat* of the laceration is to the front and left side¹ of the cervix, probably because the long diameter of the child's head is most frequent.

¹ According to Emmet and Spiegelberg; Klein and Czempin found right-sided laceration more frequent.

commonly in the right oblique diameter of the pelvis, and the thicker end of the wedge is to the front. The next in frequency is a double laceration—to the front and left, and to the back and right sides. Less




FIG. 178.

SINGLE LACERATION. The flaps are held apart with a double tenaculum (*Emmet*).

frequently is the laceration at either end of the left oblique diameter. We have found lacerations to the front and right side in cases where the head presented right occipito-anterior. The *form* of the laceration is various—single (see fig. 178), double (see Plate VIII., fig. 2), or multiple




FIG. 179.

MULTIPLE OR STELLATE LACERATION (*Emmet*).

(see fig. 179). The *extent* of the laceration varies, from a mere indentation of the ring of the os externum to a gaping fissure separating the lips of the cervix up to the vaginal fornices. Occasionally it extends

into the roof of the vagina,¹ and is marked by a cicatricial band drawing the cervix to one side; we have noticed this in forceps cases, specially when the forceps had been applied before the os was dilated.

Among the pathological conditions which are the *consequences of laceration* are the following. One result is that the mucous membrane of the cervical canal is exposed, and the occurrence of *cervical catarrh* favoured (*v. Cervical Catarrh*). The submucous tissue is also thickened and the whole cervix thus hypertrophied.² With these inflammatory changes there is *eversion* of the lips of the cervix, although this is sometimes counteracted by the formation of cicatricial tissue in the cleft. Results.

Another consequence is *cellulitis*; frequently we find, on the same side as the laceration, a localised cellulitis in the shape of a distinct deposit, or a tense condition of the utero-sacral or broad ligament, accompanied with tenderness on pressure through the fornix. This tenderness, as well as the constant pain complained of in the side, is probably due to changes in the sympathetic plexus in the connective tissue already referred to under Parametritis. *Subinvolution* of the uterus is also frequently present; there is a formation of cicatricial tissue, which compresses the veins and lymphatics and leads to passive congestion and hypertrophy. The compression of the vessels seems sometimes to have an opposite result, leading to atrophy through stoppage of nutrition.

ETIOLOGY.

A laceration of the cervix will be found, according to Emmet's statistics, in 32·8 per cent. of parous women; according to Wells, who takes the average of all the various authorities, in 32 per cent. Though it is obvious that lacerations may be produced and heal again so that all trace of them escapes notice, we cannot affirm that the cervix is lacerated with every first full-time labour; but when present, a laceration of the cervix (if we exclude the possibility of the cervix having been divided artificially) *is the most reliable diagnostic of a former parturition*. It must, however, be remembered that a divided condition of the cervix with ectropium of the cervical mucous membrane has been described as a *congenital* condition by Fischel and Küstner; in such cases, the everted mucous membrane is not much altered and retains the *arbor vitæ*. Frequency of laceration.

We should have expected that lacerations would be more readily produced in a rapid labour, in which the os had not time to dilate;

¹ Czempin, in an extremely interesting paper on cases of laceration of the cervix observed in Martin's Clinique at Berlin, draws especial attention to these tears extending into the fornix which he describes as "*Cervix-Laquearissime*." They are not infrequent (having been present in sixty-eight out of his two hundred and eighty-seven cases), usually unilateral, and more frequent with single than with double tears of the cervix itself. Their symptoms are more marked, due to the changes in the parametrium.

² Partial hypertrophies of such a size as almost to form a tumour sometimes, but very rarely, occur. Stratz describes three cases, in one of which the tumour weighed 2 lbs. (*Zeitsch. für Geb. und Gyn.*, Bd. xii., S. 229).

Emmet and Pallen, however, have found that they are more commonly the result of tedious labours. Spiegelberg blames early rupture of the membranes done to hasten labour; while Klein finds them most frequent where there is a short interval between rupture of the membranes and delivery of the child, as also where the child is heavy.

Barker and Mundé both draw attention to the fact that they are less common among the wealthy than among the poor. This is probably explained by the better care and longer rest in the puerperium which the former enjoy.

Produced
during
pregnancy.

Even *during pregnancy*, according to Nieberding, fissuring of the cervix with ectropium is produced. He examined the cases admitted to the lying-in hospital at Wurzburg at three periods—during pregnancy, as shortly as possible after delivery, and on dismissal. Only in 26 per cent. of the primiparæ examined (thirty-eight cases) was the appearance of the cervix normal during pregnancy; in all the others more or less ectropium was present. In 50 per cent. there were in addition small fissures, which made the os stellate or irregular in form.

SYMPTOMS.

Symptoms
of laceration.

It is very important to know what symptoms are referable to a lacerated cervix. Those who revel in operative treatment ascribe every pathological condition in the uterus to lacerations, while others altogether deny that they have any pathological significance.

We advance the following considerations in regard to the symptoms.

1. Lacerations of the cervix *in themselves produce no symptoms*. Hæmorrhage may arise at the time of production, but is not a symptom of the persistence of the laceration.

2. Other pathological conditions arise secondarily as the result of the laceration, of which the most important are cervical catarrh and cellulitis; cicatricial tissue in the cleft produces reflex nervous symptoms.

We sometimes find a well-marked laceration by chance, as it were, the patient having had no symptoms referable to a pelvic cause.

Frequently she complains of *leucorrhœa* and symptoms common to pelvic or uterine inflammation. *Menstruation* is often irregular, increased in 50 per cent. according to Emmet's statistics; this is in many cases due to subinvolution. *Sterility*, when present, is probably due to the accompanying catarrh; and the *tendency to abortion* to the secondary changes in the uterus or parametrium. *Neuralgia* is sometimes present, which may show itself locally in excessive tenderness to touch at the seat of laceration and has been compared to the sensitiveness present in toothache. In other cases it has taken the form of neuralgic pain in the pelvis generally, often in the groin and extending down the leg, or sympathetic neuralgia elsewhere. Emmet and others record cases in

which persistent neuralgia disappeared on excision of the cicatricial plug in a lacerated cervix. Other *reflex disturbances* (such as cataleptic convulsions, persistent salivation, profuse sweating, hysterical anuria) have disappeared after Emmet's operation. *General weakness* and inability to work are present here as in other chronic conditions.

The relation of laceration to malignant disease, of which it seems sometimes to be the starting-point, will be considered under Cancer of the Uterus.

DIAGNOSIS.

This presents, in many cases, no difficulty.

The *finger* feels the indentation or fissuring of the vaginal portion. Sometimes the cervical canal is patulous, and admits the distal phalanx of the finger easily. Difficulty in diagnosis arises when there is much eversion of the mucous membrane of the cervical canal with thickening of the cervical tissue; the fissure is thus obliterated, because the circle of the os is not formed of the os externum but of a higher unfissured portion of the canal. This thickening and the velvety feeling of the everted mucous membrane lead us to suspect the condition. Occasional difficulty in recognition.

The *speculum* shows the cleft in the cervix with, in the great majority of cases, round it appearances which will be more fully described under Cervical Catarrh. We see a bright red irregular patch on one side of or surrounding the os; from its granular appearance, its vascularity, and the fact that it bleeds easily, it resembles an ulcerated surface. For this reason it is often described as "*ulceration*" of the cervix, but it is no more an ulceration than is the inflamed mucous membrane of the conjunctiva. By ulceration we understand a destruction and loss of tissue. The epithelium and subepithelial tissue may be destroyed as an immediate result of injury during labour; but the raw-looking surface, appearing secondary to and also independent of lacerations (*see* Catarrh in Nulliparæ), is not an ulcerated surface and should therefore not be treated as such.

As already mentioned (p. 126), Sims' speculum must be used; the other forms only mask the laceration.

For the appearance presented by the various forms of laceration when seen in the speculum, the student should compare fig. 178 and fig. 179. The difference between the colour of the everted cervical mucous membrane and that of the vagina is represented in Plate VIII., figs. 1 and 2.

The microscopic changes which produce the appearance simulating ulceration will be described under Cervical Catarrh.

The *tenacula* are a valuable adjunct in examination with the speculum. If we place one in the anterior and one in the posterior lip, and roll these in on one another, the raw-looking surface will in many cases disappear. This easily demonstrated fact had not been recognised till

Emmet drew attention to it, and based on it the operation which will be always associated with his name. By thus rolling the lips inwards, we restore the laceration and see the extent of it so as to judge of the possibility of approximating the lips with sutures.

TREATMENT.

From what has been said in the introductory paragraph, and also under "Pathology," it is evident that the treatment of laceration of the cervix means much more than the closure of the split. Emmet in his operative procedure not only closes the laceration but excises the cicatricial tissue; he also makes his patients undergo a long preparatory treatment directed to the cervical catarrh. The cases calling for his operation are much fewer than might at first sight be supposed,¹ because no laceration however well marked calls for treatment unless it is producing symptoms; and there are other operations (Schroeder's and Martin's) for removing the consequences of laceration which are as efficient as Emmet's. Where there is much induration of the cervix, amputation is preferable to Emmet's operation.²

Immediate
operation
for lacera-
tion.

The stitching up of a laceration *immediately after parturition* was first performed by Pallen of New York. Having failed to check by the tampon post partum hæmorrhage from a lacerated cervix, he passed Sims' speculum and sewed up the laceration with silver-wire sutures; this checked the hæmorrhage. We have never had occasion to perform the "immediate" operation; injections of very hot water have always sufficed to check hæmorrhage. Considering the liability to septic inflammation in the puerperal condition, we would be very chary about operating unless the hæmorrhage were considerable and not diminished by hot injections.

Emmet's
Operation.

The paring of the edges of an *old laceration* and uniting of them with sutures is known as "Emmet's operation," which is a simpler and more suggestive name than "Trachelorrhaphy."

Prelimi-
naries.

Preliminaries to Emmet's Operation.—The patient should use hot-water injections for some weeks previous to the operation, and apply a blister if there be any indication of cellulitis. Emmet lays great stress on this preparatory treatment, and says that we should not operate so long as there is any tenderness on pressure in the fornices. He further recommends, in cases where the cervix is thickened and the mucous follicles

¹ Principles and Practice of Gynecology: 1884, p. 483. The conservatism as to this operation which exists in this country is justified by what Emmet said in a letter given in the interesting tabulated record of opinions of the leading operators which Zinke collected as to when and when not the operation is to be performed; the italics are ours. "The Operation has long since passed out of my hands, and so fully endorsed that I have no fear for its future. The great point is to check the abuse, which is fearful. Every one feels competent to perform it; it is done without the proper preparatory treatment, and with no special purpose. I believe in *nine cases out of ten*, where it is done, or attempted, the execution of the operation is defective and without any benefit to the patient."

² See J. D. Emmet's paper as to the limitations of the operation.

enlarged, scarification of the cervix and painting with iodine or tannin and glycerine.

The Operation. The following instruments¹ are required :—

Vaginal douche,	Dissecting forceps,
Sims' speculum,	Short needles (fig. 103) straight
Volsellæ,	and curved,
Tenacula,	Needle holder,
Bistoury and scissors,	Medium silver wire, and catgut.

Emmet's
Operation
for lacer-
ated
Cervix.

The patient is placed under chloroform in the lithotomy posture (in the semiprone posture by Emmet, but this does not give the operator so much room); the sacral segment is drawn back with the speculum by an




FIG. 180.

EXTENT OF DENUDED SURFACE AND COURSE OF SUTURES ACCORDING TO EMMET (*Emmet*).
The sutures are passed in order 1 2 3 4; the course of suture 4 alone is indicated by letters *a b c d*.

assistant, and the cervix is laid hold of with the volsella and drawn down. The uterus may be curetted at this stage. Draw the edges of the laceration together with the tenacula to see how much tissue must be pared from the edges of the cleft to allow it to be sewed up, and then proceed to operate. Wash out the vagina with carbolised water. When possible, continual irrigation is kept up during the operation. Now pare the edges of the laceration with the scissors or knife; scissors are preferable, because they cut with greater ease and rapidity.

¹ It is of great advantage, as Martin has pointed out, to curette the uterus before operating on the cervix; this can be done at the one operation, in which case we need the curette and sounds dressed with cotton-wool dipped in iodine or carbolic acid in addition to the instruments mentioned.

With long-bladed scissors we can remove the tissue from one edge of the laceration with a steady clean cut right into the angle; Emmet lays great stress on the removal of the cicatricial tissue in the angle but uses the bistoury to do this. When the laceration is bilateral this must be done on both sides. Fig. 180 shows the extent of surface denuded by Emmet in a case of bilateral laceration. Great care must be taken to leave a broad strip (broader than represented in fig. 180) undenuded in the middle line to form the walls of the cervical canal. Now introduce the sutures; these if of wire are about eight inches long so that both ends protrude from the vagina, and are well adapted to the eye of the needle so as not to obstruct its passage. Emmet recommends




FIG. 181.

MODE OF PASSING SUTURES *a b* denuded surface as in fig. 180. The sutures are passed in order as numbered.




FIG. 182.

APPEARANCE OF CERVIX WITH SUTURES TWISTED UP. They are left long so as to extend to vaginal orifice and are removed in order as numbered.

the round needle as it makes a smaller hole and is therefore followed by less hæmorrhage; when the tissues are dense, the lance-shaped point perforates more easily. Catgut¹ has the great advantage over silver wire, that the stitches do not require to be removed afterwards; strong sutures are necessary, as some force is needed to tie them tight. If wire or silk has been used the sutures are removed after ten days.

No special regimen is required afterwards, the diet need not be restricted. Secondary hæmorrhage has sometimes followed the operation: it is best checked by passing a suture through the cervix higher up and

¹ Meinert recommends passing the catgut right through the cervix and fixing the ends with shot on plates: Eine sichere Catgutnaht für die Emmet'sche Operation: Archiv f. Gyn., XXXIII., S. 310.

tying it tightly on the side from which the hæmorrhage comes so as to constrict the vessels in the cervix.

The effect of the operation on sterility has given rise to a great deal of discussion. Wells gives in his paper an interesting table of statistics as to subsequent conceptions, and affirms that the operation increases fertility; the proportion (one-fourth) of cases fertile after Emmet's operation is, however, the same as Emmet gives for cases of laceration generally, *i.e.* whether operated on or not.

The cicatrix does not cause difficulty in subsequent parturition. The cervical catarrh may persist after the operation. Sometimes metritis, cellulitis, or peritonitis has unfortunately followed it.

Other methods of closing the cervical laceration by *flap operations* have been proposed. Sanger¹ and Kleinwachter² dissect up a flap of cervical mucous membrane from the faces and apex of the cleft, and turn it in towards the cervical canal. The object of this is to preserve the cervical mucosa, and the result is a wider cervical canal. The operation is analogous to Simpson's one for the repair of the torn perineum, in which the vaginal mucosa is dissected up in a similar way. Duhrssen makes an incision along the middle of each face of the cleft extending into the angle, dissects a flap outwards and inwards, and then everting these flaps brings together the raw surfaces. This operation is similar to what is done in fistulæ, when the margins of the fistula are split, and the flaps turned towards the respective free surfaces. The object of repair of the cervix is not, however, comparable to that of repair of the perineum, or the closure of a fistula. Flap-splitting is called for in these operations to save tissue—to strengthen the perineum or to close in the fistula; while in the case of the cervix we have to do with excess of diseased tissue, either hypertrophied mucosa or new cicatricial formation, which requires to be removed. The cleft is in itself of no significance.

Other operations to meet the consequences of laceration.—Emmet's operation is directed not only against the split but also its consequences, the cicatrization and the cervical catarrh. Simply to close an old split would be as meaningless as shutting the stable door in the proverb. For the treatment of the catarrh, we have also Schroeder's excision of the mucous membrane of the cervix and Martin's amputation and excision, both of which will be described in the next chapter.

For extensive tear into the fornix which has resulted in cicatrization in the parametrium with lateral displacement of the uterus, Martin has introduced as a special operation³ the separation of this cicatricial tissue from the cervix. Under chloroform, in the lithotomy posture, the cervix

¹ Cent. f. Gyn. 1888. No. 49, and Volk. Samm. klin. Vortrage N.F. No 6, July 1890.

² Zeits. f. Geburts u. Gyn., XVII., No. 2.

³ Czempin (*loc. cit.*) gives three cases in which marked symptoms disappeared after this operation, and also a tabular report of nine more recent cases in Martin's clinique with similar good results.

is drawn over with forceps from the affected side and a semilunar incision made in the cicatrix in the fornix, following the contour of the cervix. This may be sufficient; or it may be necessary in addition to cut out a portion of the cicatrised tissue. The antero-posterior incision is then stitched so as to bring front and back together and thus make the line of junction traverse.

CHAPTER XXX.

CHRONIC CERVICAL CATARRH.

LITERATURE.

Baldy—An American Text-book of Gynecology : London, 1895. *Bennett, G. H.*—Practical Treatise on Inflammation, Ulceration, and Induration of the Neck of the Uterus : London, 1845. *Candia*—L'amputazione della porzione dell' utero come cura radicale delle erosioni e della metrite parenchimale cronica : *Annali di Ostet.*, 1888, 463. *Duncan, Matthews*—Diseases of Women : London, 1886. *Fischel*—Ein Beitrag zur Histologie der Erosionen der Portio vaginalis uteri : *Archiv f. Gyn.*, Bd. XV., S. 76. *Fritsch*—Erosionen, Ectropium und Cervicalcatarrh : *Billroth u. Luecke's Handbuch der Frauenkrankheiten*, Stuttgart, 1885. *Hennig*—Der Katarrh der inneren weiblichen Geschlechtstheile : Leipzig, 1862. *Hildebrandt*—Ueber den Katarrh der weiblichen Geschlechtsorgane : *Volkmann's Sammlung klin. Vorträge*, Leipzig, 1872, No. 32. *Hofmeier*—Folgestände des chronischen Cervixkatarrhs und ihre Behandlung : *Zeitsch. f. Geb. u. Gyn.*, Bd. IV., S. 331. *Klotz*—Gynäkologische Studien über die pathologischen Veränderungen der Portio vaginalis Uteri : Wien, 1879. *Küstner*—Beiträge zur Lehre von der Endometritis, Erosion und Ectropium : Jena, 1883. *Mundé*—The treatment and curability of chronic uterine catarrh : *Amer. Journ. of Obstet.*, 1883, p. 857. *Ruge and Veit*—Zur Pathologie der Vaginalportion : *Zeitschrift f. Geb. u. Gyn.*, Bd. II., S. 415. *Schroeder*—Krankheiten der weiblichen Geschlechtsorgane : Leipzig, 1879, S. 122. *Stevenson*—Note on the use of Electrolysis in Gynecological Practice, and Discussion : *Trans. Lond. Obs. Soc.*, June and July 1888. *Vulliet*—De la dilatation permanente de l'utérus : *Arch. de Toc.*, 1886, p. 933.

WITH cervical catarrh we begin the study of inflammation of the uterus, to which this and the two following chapters are devoted, and with regard to which some general observations must be made by way of introduction. Our knowledge of this subject has been slowly extending with improved methods of investigation. After the speculum had concentrated attention unduly on the naked-eye appearances of the cervix, the introduction of the sound and the curette along with the bimanual examination, brought within range the condition of the uterus itself, and its cavity. And to-day abdominal section has still further extended the field of observation to the uterine appendages. With a wider field there have come corresponding changes in gynecological opinion as to the significance of the different lesions.

In studying inflammation of the uterus, it is convenient to separate inflammation of the cervix from inflammation of the body, and analyse the latter into inflammation of the mucous membrane, endometritis ;

and of the body generally, metritis. While pathological differences between these are sufficiently well marked to warrant this sub-division, clinically it is not always easily made out. Inflammation is not limited in its spread to one tissue, and there is a combination of symptoms which is more or less common to all the forms of uterine inflammation. This has led French writers on gynecology to group the subject matter of these three chapters under the general heading of metritis. Although this simplifies the handling of the subject, and saves repetition, it savours of a return to the symptomatic standpoint from which gynecology has been slowly rising.

In considering uterine inflammation, we must bear in mind that the uterus differs functionally from other organs lined by mucous membrane, such as the bladder, stomach, or lungs. These are in constant use, while the reproductive function is only occasional. We must be careful, therefore, in transferring to the uterine mucosa ideas of inflammation derived from other mucous membranes. An endometritis is not comparable to a gastritis or a cystitis. Many conditions grouped under the heading of "endometritis" are more allied to new-formation than inflammation.

On the other hand, while the reproductive function is occasional, the uterine mucosa is subject to a periodic hypertrophy and degeneration in connection with menstruation. This complicates greatly the study of the morbid changes in the mucosa.

These considerations bear on inflammation of the body of the uterus rather than on that of the cervix. The cervical canal is anatomically more like a mucous membrane than the endometrium. Further, while it takes no part in the changes of pregnancy and menstruation, it is continually secreting a characteristic mucus for the lubrication of the vagina. There is thus a cervical catarrh analogous to a bronchial or gastric catarrh, to the consideration of which the following chapter is given.

Acute catarrh of the cervix is known to us only as part of a general catarrh affecting both body and cervix, and will be described under Acute Endometritis. *Chronic catarrh* occurs localised in the cervical mucous membrane; it is a very common condition and one of the most troublesome which the practitioner has to treat.

DEFINITION.—A chronic inflammatory process affecting the mucous membrane lining the cervical canal.

SYNONYMS.—Cervical endometritis, Endo-cervicitis.

PATHOLOGY.

The mucous membrane of the cervical canal is inflamed. When the os externum has been lacerated, the lips gape and the mucous membrane

is thus everted; on bringing the margins of the laceration together, this eversion will disappear. Further, there are granular patches with irregular outline which extend beyond the limits of the os externum; these have a red appearance resembling the cervical mucous membrane, and are therefore sharply defined from the paler mucous membrane which covers the vaginal portion of the cervix.

This last condition was till late years generally held to be an "ulceration" and is still described, even in recent English works, under that name. The term should, however, be discarded as based on an erroneous pathology and suggesting most pernicious treatment. The cause of the error is easily explained: a raw-looking granular surface was seen with the speculum; the raw appearance was ascribed to the loss of the epithelium, and this supposition was supported by the microscopic examination of specimens taken from the dead body, in which the

Pathology
of so-called
Ulceration
of the
Cervix.




FIG. 183.

SECTION OF CATARRHAL PATCH, OR PAPILLARY FORM OF EROSION (from a micro-photograph).

epithelium had been macerated and removed; the granular points were supposed to be the subjacent papillae which had become hypertrophied.

Both of these suppositions have been shown to be erroneous by the careful investigations of Ruge and Veit, who examined specimens of the so-called ulcerations cut fresh from the living subject; they demonstrated

Ruge and
Veit's
investiga-
tions.

(1) that the apparently raw surface is covered with epithelium, (2) that the granular points are new formations and have no connection with the papillae of the mucous membrane.

The microscopic appearance of the mucous membrane described by them is as follows. The surface is covered with a single layer of epithelium; the cells are smaller than those which line the normal cervical canal, and being narrow and long have a palisade-like arrangement; the thin layer of cells allows the subjacent vascular tissue to shine through,

hence the *redness* of colour. The surface is further thrown into numerous folds producing glandular recesses and processes; these processes cause the *granular* appearance of the surface. The condition is well seen in Plate VIII., and constitutes the *simple* erosion: fig. 1 shows such an erosion as seen in the speculum: fig. 3 shows a microscopic section of the same, stained with carmine; the left half of the section corresponds to the deep red portion of fig. 1, the right half to the paler portion outside of this. If the recesses be long and narrow, the surface is split up into distinct papillæ; this constitutes the *papillary* erosion (see fig. 183). If the ducts of the glandular recesses become obliterated, the section will distend the gland below and produce retention-cysts; these will increase in size, and may come to the surface and burst. Thus there is formed the *follicular* erosion (see fig. 184).

The raw-looking surface is therefore a *newly-formed glandular secreting surface*, resembling in structure the cervical mucous membrane. This




FIG. 184.


FOLLICULAR FORM OF EROSION (Schroeder).

addition to the extent of secreting surface increases the leucorrhœal discharge which is the leading symptom.

In studying cervical catarrh we must not, however, limit our view to surface changes of epithelium, or else we shall simply fall into the error of those whose idea of it was limited by what they saw in the speculum. The whole substance of the cervix is affected by the inflammatory process, and it is possible that the epithelial changes are a mere accident, or at most a surface sign of a deeper lesion. According to Abel and Landau it is not of much importance whether the epithelium is squamous or cylindrical, and is present or absent; and the redness is due not so much to the thinness of the epithelium as to the congestion below. The changes seen in the speculum are significant as the signs of an inflammatory condition of the cervix, with which we also have frequently associated like changes in the uterus, in one word, chronic metritis. Pozzi, approaching the subject from a clinical standpoint



1



2



3



makes chronic metritis *the* lesion, of which the speculum changes are only one of many signs.

For this reason it is better not to apply any special name to these red patches, or to call them simply *catarrhal patches*.

The origin of this new epithelial structure is disputed. Ruge and Veit hold that the single layer of small cylindrical cells is produced by proliferation of the cells of the deepest layer of the rete Malpighi, while those of the superficial layer are shelled off (fig. 184). On the other hand, those red patches are generally continuous with the mucous membrane of the cervical canal, and resemble it in their microscopic structure; it is therefore much more probable that they are occasioned by proliferation of the *epithelium which lines the cervical glands*, leading to an extension of the glandular surface beyond the os externum.

The question as to the origin of the cylindrical epithelium found in erosions is rendered more difficult by the fact that the boundary line between the squamous epithelium outside of and the cylindrical within




FIG. 185.

TRUE ULCERATION OF THE CERVIX. At the sides of diagram is seen the normal epithelium, which is prolonged in processes, *e p* between the connective tissue papillæ; *e* is superficial layer of squamous epithelium reduced to a thin layer at *e'*; *c t*, tissue of mucosa infiltrated with small cells; *b v*, blood-vessels surrounded by small-celled infiltration (*Fischel*).

the cervical canal varies at different periods of development, and in different individuals.

The vagina is, as we have seen, originally lined with cylindrical epithelium, that is to say, the genital tract being formed of the ducts of Müller has the same epithelial lining throughout. Only at a subsequent period does the vagina come to have squamous epithelial lining, which may be derived from the Wolffian bulbs (p. 84). Whatever its origin, it extends upwards as a rule to the tip of the cervix where the os externum usually is found. Rarely does it fall short of this, in which case the vaginal aspect of cervix has the same rosy appearance as the lining of the canal. To this condition *Fischel* has applied the term congenital ectropium, since the appearance described resembles that called, later in life, ectropium, but the term is, we think, misleading. It is really a defective extension of the squamous epithelium upwards. More frequently a converse condition is present, the squamous epithelium extending within the cervical canal. This has been described in the child by Lott, and Landau and Abel; and Klotz makes one type of cervix in which in the adult squamous epithelium extends within the cervical canal.

True
Ulcerations.

Sometimes a true ulcerated process—destruction of epithelium with inflammation of connective tissue—does occur; such a condition is represented in fig. 185.

Ovula
Nabothii.

Along with those changes in the mucous membrane, chronic inflammatory changes occur in the other tissues of the cervix. There is increased formation of connective tissue, which produces antero-posterior thickening and sometimes elongation. The secretion in the obstructed glands becomes inspissated, and hence the retention cysts are felt as firm pea-like bodies—ovula Nabothii—in the substance of the cervix or projecting from it; or their contents may suppurate and form small abscesses. As there are no racemose glands on the vaginal portion beyond the limits of the os externum (see Histology of Normal Cervix), these ovula Nabothii must be produced from the glands of the mucous membrane of the cervical canal or from the newly-formed glandular tissue. Fritsch draws attention to the fact that the glands of the cervix are enormously hypertrophied during pregnancy, so that the cervix becomes almost a glandular organ; the persistence of this condition after the puerperium, may explain the increased glandular formation which is described above as the chief pathological element in cervical catarrh.

Cysts in
the Cervix.

Sometimes we find a single large cyst in the cervix, due to obstruction of the mucous glands. When it is in the substance of the wall, the soft bulging into the cervical canal and the accompanying menorrhagia may lead one to suspect commencing sarcomatous infiltration. Puncturing with a trocar removes a clear or straw-coloured fluid, rich in mucous corpuscles.

The microscopic pathology of the cervix has only of recent years been carefully investigated, and there are many points on which definite information has not as yet been obtained. The following is a brief summary of the pathological changes described, which are best understood by comparison with the microscopic structure of the normal vaginal portion.

Normal
and Pathological
conditions
of Cervix
contrasted.

NORMAL CONDITION. The vaginal portion is covered on its vaginal surface with many layers of squamous epithelium, resting on papillæ of connective tissue; there are no mucous follicles. The cervical canal is lined with a *single layer* of *cubical* epithelium (ciliated only on the ridges), folded so as to form shallow recesses which do not branch; there are *racemose mucous glands*, which have branching ducts. The substance of the cervix is made up of *connective* tissue.

PATHOLOGICAL CHANGES. These, according to the extent and duration of the process, affect the three elements—epithelium, glands, connective tissue.

The *epithelium* of the cervical canal may be simply exposed (ectropium after laceration), or it may be inflamed. When inflamed,

the folding of the mucous membrane is greatly increased so that the surface has a papillary or granular appearance. Further, this inflamed mucous surface may be found extending beyond its normal limit (the os externum) in the form of red patches (catarrhal patches) which are smooth or granular. These patches, though the most striking feature of the disease as seen in the speculum, are chiefly of significance as an index of a deeper lesion, just as a furred tongue is of a gastritis.

The *glands* hypertrophy and new glands form as the result of the proliferation of epithelium described above. The openings of the glands are at first restricted to the area covered with a single layer of cubical epithelium, but their branching ends extend below the limiting surface of stratified squamous epithelium. Their ducts become obstructed, and retention cysts form not only on the red patches but also underneath the adjacent, apparently normal, vaginal mucous membrane. They may remain as little nodules in the mucous membrane, or may come to the surface and burst; in the latter case, the cubical epithelium and papillæ on the inner wall of the cystic gland are exposed and, being now on a free surface, proliferate. When the glands are the special seat of the pathological changes, the whole substance of the cervix is converted into a cystic mass.

The *connective tissue* always increases in amount, specially when the process is chronic. This increase constitutes the "areolar hyperplasia" of Thomas.

ETIOLOGY.

The most important cause is, undoubtedly, the injury of the cervix produced in *parturition*; hence cervical catarrh is common in parous women. How this injury produces the inflammatory condition is a disputed point. Emmet refers it to the persistence of the split in the cervix and holds that the exposure of the mucous membrane to friction against the vaginal walls leads to irritation and inflammation; but we frequently see cases of well-marked lacerations without consequent cervical catarrh. It is admitted by all that the existence of lacerations greatly favours the development of catarrh.

Other less important causes are the *spread of inflammation* from the vagina *upwards* (vaginitis, which may be simple or gonorrhœal), and from the endometrium *downwards*. The latter is favoured by the fact that the discharges from the endometrium necessarily flow over the cervix and irritate it.

Cervical catarrh is the most frequent complication of retroflexion of the uterus. The flexion favours gaping of the lacerated cervix, and produces passive congestion of the cervical tissues.

Frequency
of Catarrh
in Multi-
paræ.

SYMPTOMS.

These are—Leucorrhœa ;

Pain in back and loins, increased on exercise ;

Irregular menstruation ;

Sterility.

Leucorrhœa is the prominent symptom. Under normal conditions the secretion from the mucous membrane of the uterus and cervix is not sufficient to attract attention ; when it is excessive, it is termed leucorrhœa (λευκός white, ῥέω to flow) or in popular language “whites.” A transparent leucorrhœa from the cervix and uterus occurs before and after the menstrual flow ; this is a hyper-secretion due to temporary congestion.

Characters
of Cervical
Leucor-
rhœa.

The secretion from the glands of the cervical canal is clear and viscid, resembling unboiled white of egg. It becomes of an opaque white when mucous corpuscles are abundant, yellowish when pus corpuscles are present. Frequently, it is tinged with blood from the blood-vessels of the newly-formed vascular tissue.

Pain in the back and loins is present, as in all uterine disease. It is aggravated on active exercise, such as walking and riding, or whatever causes friction of the cervix against the vaginal walls.

Menstruation is irregular, and often increased in quantity ; this is probably due to extension of inflammation upwards to the endometrium. We must take care not to mistake leucorrhœa tinged with blood for the regular menstrual flow.

Sterility is often present. In nulliparæ with a small os externum, the plug of mucus in the cervical canal is alleged to be a bar to conception. In multiparæ, we have seen conception take place even though there was a deep laceration and well-marked catarrh ; the presence of catarrh, however, though not an obstacle to conception, diminishes its probability.

PHYSICAL SIGNS.

Condition
of Cervix
in Chronic
Catarrh.

On *vaginal examination*, the condition of the cervix is found to vary according as the patient is nulliparous or multiparous and the disease of long or short duration. In a nullipara, the cervix feels puffy and large, the margins of the os soft and velvety (when there is eversion with extension of catarrhal area beyond the os externum) ; or the os and cervix are apparently normal but movement causes pain (when the catarrhal area does not extend beyond the os externum). In a multipara, the existence of a laceration must first be determined and the extent of it noted ; the margins of the os are soft and velvety, and pea-like nodules (Nabothian follicles) are felt on and sometimes round them ; polypoidal projections may be present and, more rarely, the cervix is converted

into a mass of cysts ; the os is usually gaping so that the finger can be passed into the cervical canal, where the mucous membrane has an irregular surface and is often thrown into longitudinal ridges.

The *speculum* is now employed ; its use must always be preceded by a careful examination with the finger to ascertain, when laceration is present, the undisturbed relations of the lips of the cervix. Neither finger nor speculum alone is sufficient, we must employ both, and learn to associate what is felt by the finger (*e.g.*, lacerations, velvety mucous membrane, pea-like follicles) with what is seen with the speculum. The superiority of the Sims speculum for examination is very marked, as it exposes the lips of the cervix without disturbing the relations.

Appear-
ance in
Speculum
of Cervix
in Catarrh.

In a nullipara, we see the os apparently normal but with a tenacious plug of mucus projecting through it ; or there may be red catarrhal patches such as are represented in Plate VIII., fig. 1, which shows very well the contrast between the appearance of these patches and the surrounding mucous membrane ; no chromo-lithograph, however, perfectly displays the natural colours.

In a multipara, a laceration is sometimes evident. Oftener it escapes recognition ; the os appears to be wide and unfissured, while on both lips there is a red velvety surface (Plate VIII., fig. 2) ; if, now, tenacula be fixed in the gaping lips and those rolled in on one another, the red surfaces will disappear and a bilateral laceration become evident. Sometimes, white cicatricial tissue indicates the situation of the laceration. Though the lips are thus approximated, a red surface is often visible because the catarrhal area has spread beyond the os externum. The obstructed Nabothian follicles appear as bluish-red projections from the mucous membrane ; occasionally they form small polypi.

DIAGNOSIS AND DIFFERENTIAL DIAGNOSIS.

The diagnosis between cervical and vaginal catarrh is made clear by using the speculum, for we see in the former case the leucorrhœa coming from the cervix and having the characters above described. Should the discharge not be profuse enough to be seen with the speculum, we may employ the method recommended by Schultze for diagnosing between uterine and vaginal catarrh. The vagina is douched out in the evening, and a tampon soaked in a solution of tannin is placed against the os externum ; in the morning the tampon is removed through the speculum, and we note the quantity and character of the discharge which has accumulated upon it.

Diagnosis
of Cervical
from
Vaginal
Leucor-
rhœa,

The diagnosis between cervical catarrh and endometritis is difficult, and in many cases cannot be made ; when cervical catarrh is present, we cannot be positive that there is not some endometritis as well. Increase in the length of the uterine cavity (especially with tenderness or irregularities of the mucous membrane), ascertained by the sound,

from
Endo-
metritis.

indicates endometritis. When the cervix is much thickened and indurated, we may suspect the commencement of malignant disease; this will be considered under Carcinoma of the Cervix.

PROGNOSIS.

In this we must consider the constitutional health of the patient, the duration of the symptoms, and the extent to which the tissues are affected. According to Thomas, the prognosis is less favourable when there is considerable secretion of mucus with little apparent "granular degeneration." The practitioner will often find that cases of cervical catarrh have already passed through several hands, and he should therefore be on his guard in offering hopes of speedy cure.

TREATMENT.

Constitutional treatment important.

In the first place, special attention must be given to the patient's *general health*; if we trust to local treatment alone, we shall often be



FIG. 186.

HEALING OF A CATARRHAL PATCH TREATED BY ASTRINGENT OR ANTISEPTIC INJECTIONS (*Hofmeier*).
a to *b*, newly-formed squamous epithelium; from *c* to *c'*, is seen alteration of the epithelium at the mouths of the glands; *d, d'*, glands with ducts obliterated; *e*, gland duct which has persisted.

disappointed. We should recommend change of air and light nourishing food. A certain amount of exercise is valuable; but too much of it, specially of riding, is injurious. Tonics (such as arsenic, quinine, and iron) are useful. Disturbances of the digestive system, which are frequent in chronic cases, must be treated as each case indicates. Complete rest from sexual activity is advisable; this can often be secured by recommending that the patient go away from home for a time.

Cervical catarrh is in some cases only a local manifestation of a constitutional state such as tuberculosis or anæmia.

The *local treatment* varies according as the patient is nulliparous or

multiparous. In both cases we must be prepared to carry out a system of treatment which lasts for weeks.

1. In *nulliparæ* we begin with a course of vaginal injections of hot water. These are used freely, from ten minutes to a quarter of an hour, every night. To the simple water, astringents or antiseptics are added : sulphate of zinc (ʒj to the pint) ; sulphate of alumina or sulphate of copper (ʒij to the pint), or corrosive sublimate (1 to 4000). Local treatment in Nulliparæ.

The action of these on the catarrhal patches has been specially investigated by Hofmeier and by Küstner. The former found that such a patch, treated by daily vaginal injections of pyroligneous acid, became gradually encroached on by the surrounding squamous epithelium's creeping in tongue-like processes over the cylindrical epithelium. The more superficial glands became filled up with the squamous epithelial cells ; the deeper ones had their ducts narrowed or even plugged, while the gland-cavity persisted below (fig. 186). Küstner found that similar changes could be produced by antiseptic douches.

If the os be narrow, it is good to notch it bilaterally with the scissors. This acts beneficially by allowing the mucus to escape freely. Mundé recommends the trimming of the lips of the cervix so as to produce a funnel-shaped os.




FIG. 187.

FORCEPS DRESSED WITH COTTON WADDING.

When we find that the secretion continues copious in spite of the frequent injections, we must make a local application to the mucous membrane. Of applications the best are iodine (the tincture or the strong liniment) and carbolic acid, the former in milder and the latter in more severe cases. The liquor hydrargyri pernitratidis is recommended by Heywood Smith, and chromic acid is much praised by De Sinéty. In making these applications we proceed as follows. The mucus, which would prevent the action of the medicament on the mucous membrane, is first thoroughly removed by the forceps dressed with cotton wool as represented at fig. 187. A second pair of forceps, covered merely with a film of cotton wadding, is now dipped in the medicament and applied to the surface. Should the canal be narrow, a sound dressed as for endometric applications (see fig. 195) is preferable. Care is taken that there be no free drop of the solution on the cotton wool, which might fall on the vaginal mucous membrane ; after the application is made, a pledget of cotton wadding with glycerine is placed below the cervix.

Rarely in *nulliparæ* is the pathological process so extensive as to require operative means for removing cervical tissue.

Local
treatment
in Multi-
paræ.

2. *In multiparæ.* Here the cervical catarrh is usually associated with other conditions—retroflexion, subinvolution, and, especially, marked laceration of the cervix. The first treatment indicated is to diminish the passive congestion of the cervix by hot-water injections with astringents or antiseptics, and the use of the glycerine plug. The latter is prepared as already described (p. 241). If the uterus be retroflexed, it should be replaced and kept in position by a pessary. Even where it is not retroverted, a pessary is often useful in lifting the uterus upwards in the pelvis and diminishing passive congestion. In cases where there is a distinct laceration of the cervix, and specially where the catarrhal patches can be made to disappear by rolling the lips inward on each other, Emmet's operation is indicated.

Depletion
by Scarifi-
cation or
Leeches.

Local depletion by scarification is done best through the Fergusson speculum, and with a lancet-shaped bistoury; a number of small punctures are made from a quarter to half-an-inch in depth.




FIG. 188.




FIG. 189.

SCHROEDER'S EXCISION OF THE CERVICAL MUCOUS MEMBRANE in cervical catarrh. Fig. 188 LINE OF INCISION IN MUCOUS MEMBRANE. Fig. 189 MUCOUS MEMBRANE EXCISED and flap *bc* turned in on *ab* (Schroeder).

Scarification is also useful for another object. When there are hard knobby retention cysts producing irritation by the pressure of their contents, the puncturing of these diminishes the chronic inflammation. Paquelin's cautery is also used to puncture the cervix; but this use of it belongs rather to the treatment of the hypertrophy of the cervix in Chronic Metritis.

In very chronic cases, the only remedy is the destruction of the diseased glandular tissue—just as in tonsilitis we partially excise the tonsils. This has been done by the application of strong caustics or the cautery. The zinc-alum sticks introduced by Sköldberg of Stockholm are recommended highly by Matthews Duncan. They are made by fusing together equal parts of sulphate of zinc and sulphate of alumina, and running into moulds. The stick is pushed into the cervix, and a

plug of wadding laid in the vagina to keep it in place and receive the discharge. The student must discriminate this use of a powerful caustic once for all from the repeated touching of the surface with a milder caustic, just as one would touch a slow ulcer—a treatment which cannot be too strongly condemned.

Electricity has been used both in France and this country with the same object, viz., the cauterisation of the cervical glands. An electrode with a rounded end (or a uterine-sound one, if it has to be passed up the canal) is connected with the negative pole of the battery, while the positive pole is placed on the surface of the skin. Several cases¹ have been treated successfully by this method, but it remains to be seen whether it possesses advantages over other forms of cauterisation to compensate for the difficulties in its use.

Thomas recommends the steel curette for the removal of the diseased




FIG. 190.

MARTIN'S METHOD OF EXCISING THE MUCOUS MEMBRANE OF THE CERVIX (*Martin*).

The continuous black line shows line of excision, which is higher up in the fornix than in fig. 188. The dotted line is the course of the suture introduced after the piece of the lip is excised.

glands; it is applied "so forcibly as to remove the arbor vitæ and mucous glands from the os internum to the os externum. Sometimes a second operation in two or three weeks after the first has been necessary, and sometimes even a third."

Schroeder used the knife, and operated as follows. The cervix is laid hold of with two volsellæ, one on each lip, and drawn downwards. It is divided laterally as far as the fornix with the scissors, so as to form an anterior and a posterior lip which are separate as far as the vaginal roof (fig. 188). A transverse incision (seen in section, at *a*, in fig. 188) is made across the base of the anterior lip, dividing the whole thickness of the cervical mucous membrane. He next pierces the point of the lip at *c*, pushing the knife in the direction *bb* till it reaches the cross incision *a*; he carries the blade outwards first to one side and then to the other, so that all outside of the line *a b c* is cut away. The flap

Schroeder's
Operation
for
Cervical
Catarrh.

¹ Lovell Drage and Gibbons—Brit. Med. Journ., 1888, I., p. 1274. Touret—Nouv. Arch. d'Obstét. et de Gyn., April, 1887.

of cervix is now turned in, and stitched as in fig. 189. The advantage claimed for this method of operating is that the degenerated cervical mucous membrane is replaced by vaginal mucous membrane which shows no tendency to degenerate. Schroeder operated thus more than three hundred and fifty times (two deaths), and with very good results as to the cure of the catarrh.

Martin of Berlin in excising the diseased mucous membrane sometimes removes more of the substance of the cervix, as fig. 190 shows, thus combining amputation with excision. He splits the cervix into two lips, cuts through the cervical mucous membrane in the posterior lip above the diseased portion, then removes as much of the lip as is necessary, and stitches it. The anterior is treated in the same way; and then the sides are sutured—the sutures often requiring to be passed deeply to control bleeding. In introducing these last the volsella can be taken out and the cervix held down by the sutures in the two lips.

CHAPTER XXXI.

ENDOMETRITIS.

LITERATURE.

Atthill—On Endometritis: Dublin Jour. of Med. Sc., Jan. 1873. *Baldy*—An American text-book of Gynecology: London, 1895. *Barbour*—Article on Inflammation of the Uterus in Allbutt and Playfair's System of Gynecology: London, 1896. *Barnes*—Diseases of Women: London, 1878, p. 530. *Braun, Carl*—Therapie der Metritis und Endometritis, etc.: Wiener med. Wochenschrift, 1873, Nos. 39-43. *Cornil*—Leçons sur l'anatomie pathologique des Métrites, etc.: Paris, 1889. *De Sinéty*—Gynécologie, p. 327: Paris, 1879. *Dolérís*—De l'Endométrite et de son Traitement (with Discussion): Archiv de Toc., 1887, pp. 97, 145, 193, 314, 460. *Fritsch*—Dilatation of the Uterus and Intra-uterine Therapeutics: Amer. Journ. of Obstet., 1883, p. 113. *Guérin*—De la Métrite aigue: Annales de Gynécolog., Juillet, 1874; and Arch. de Tocologie, Juillet, 1877. *Heinricius*—Ueber die chronische hyperplasirende Endometritis: Arch. f. Gyn., Bd. XXVIII., S. 163. *Heitzmann*—Pathologie und Therapie der nicht puerperalen Endometriden: Centralb. f. die gesammte Therapie, 1886, Nos. 1-5. *Hennig*—Der Katarrh der inneren weiblichen Geschlechtstheile: Leipzig, 1862. *Klob*—Pathologische Anatomie der weiblichen Sexualorgane: Wien, 1864, S. 211. *Küstner*—Beiträge zur Lehre von der Endometritis: Jena, 1883. *Landau and Abel*—Beiträge zur pathologischen Anatomie des Endometrium: Archiv f. Gyn., XXXIV., S., 165. *Olshausen*—Ueber chronische hyperplasirende Endometritis des Corpus Uteri: Archiv f. Gynäkologie, Band. VIII., Heft 1. *Playfair*—Intra-uterine Medication: Brit. Med. Jour., Dec. 1869; Ibid., March 1880; Lancet, January and February, 1873. *Pozzi*—Traité de Gynécologie: Paris, 1897. *Récamier*—Des granulations dans la cavité de l'utérus: Annal. de Therap., Nov. 1846. See also Union Méd. de Paris, Juin 1850. *Routh*—On "Fundal Endometritis:" Lond. Obst. Trans., Vol. XII. Cases of Menorrhagia treated by injections, etc.: Lond. Obst. Trans., Vol. II. *Ruge*—Zur Aetiologie und Anatomie der Endometritis: Zeits. f. Geb. u. Gyn., Bd. V., S. 317. *Schroeder*—Krankheiten der weiblichen Geschlechtsorgane: Leipzig, 1886, S. 108. *Simpson, Sir J. Y.*—Diseases of Women: Edinburgh, 1873, p. 736. *Smith, Tyler*—Pathology and Treatment of Leucorrhœa: London, 1855. *Thomas*—Diseases of Women: London, 1880, p. 268.

THE endometrium differs functionally (as it does anatomically) from the cervical mucosa. While the latter is a secreting organ, the former has to produce a decidua in utero-gestation. As already said, the mucosa of the body of the uterus is not comparable to that of the stomach or lungs, and many of the morbid processes grouped under endometritis are more allied to new-formation than inflammation.

This is specially true of those which are simple in origin, in contradistinction to those which are septic or specific. There is a growing tendency to regard those forms of endometritis, for example the glandular and fungous forms, as having the characters of a new formation rather than an inflammation.

There are, however, truly inflammatory conditions of the uterine mucosa, and the importance of the endometrium as the starting-point of inflammatory conditions in and around the uterus has of recent years been emphasised. As the result of this, salpingitis and even peritonitis are regarded as secondary lesions, the important condition being that of the endometrium. This change of view in operative Gynecology is interesting as seen in three ways. Intra-uterine treatment is being emphasised,¹ pelvic inflammation is regarded as an indication rather than a contra-indication for curetting,² and the more conservative treatment of the uterine appendages is being advocated.³

We now consider inflammation of the mucous membrane of the uterus, although, from what we have said of extension of inflammation beyond the endometrium, we rarely find endometritis as a lesion by itself.

DEFINITION.—Inflammation of the mucous membrane of the uterus.

SYNONYMS.—Uterine catarrh, internal metritis.

PATHOLOGY.

In *acute endometritis* both body and cervix are involved, and usually the underlying muscular coat also. The mucous membrane is swollen and soft, and covered with red-stained mucus or creamy pus. Extravasations of blood are present as red streaks or patches. These changes are not so marked in the cervical mucous membrane as in that of the body; the vaginal portion has the same appearance as during pregnancy, being soft and swollen and showing red catarrhal patches round the os.

The ciliated epithelium is destroyed, and sometimes casts of the epithelium of the glands are found in the discharge (*Schroeder*). The secretion is at first mucous, then purulent.

In *chronic endometritis* the mucous membrane is hypertrophied and marked with patches of old extravasation.

Micro-
scopic
changes in
Chronic
Endo-
metritis.

The microscopic appearances have only of recent years been worked at, and there is considerable difference of opinion both as to the changes produced and the significance of them. To understand these, we must keep in mind the two essential elements of the mucous membrane—the glands and the inter-glandular tissue; and also the view of Leopold, who considers the inter-glandular tissue as made up chiefly of lymphatics.

Accordingly, as the changes affect principally one or other of the two

¹ As in Mackenrodt's paper, *Beitrag zur intra-uterinen Therapie*: Sam. Klin. IV. N. F. No. 45.

² Mackenrodt—*Loc. cit*; also Piqué—*Gaz. des Hôpitaux*. 1891, No. 19.

³ Polk—*New York Jour. of Gyn. and Obstet.*, Feb. 1892.





FIG. 1.




FIG. 2.




FIG. 3.



FIG. 4.



FIG. 5.




FIG. 6.

W. & A. R. Johnston Edinburgh & London.

MUCOUS MEMBRANE OF UTERUS IN ENDOMETRITIS

(FIGS. 1—5, Ruge; FIG. 6, Heinrichs).

- FIG. 1. Normal Mucous Membrane, FIG. 2. Interstitial Endometritis,
FIG. 3. Glandular hypertrophic E., FIG. 4. Glandular hyperplastic E. (all magnified ten times).
FIG. 5. Endometritis after abortion showing group of decidua cells *d c*
FIG. 6. From *E. fungosa* showing nature of changes in interglandular tissue (v. p. 318).

elements of the mucous membrane, Ruge finds a *glandular*, an *inter-* Ruge's three varieties. *stitial*, and a *mixed form*—the last being a combination of the first two.

In the *glandular*, a marked growth and increase of the glandular epithelium occurs. The gland-ducts hypertrophy (Pl. IX., fig. 3), and through multiplication of the epithelium may have bulgings of it into their lumina, making them saw-like instead of tubular in section, or the wall may be thrown into folds (*cf.* appearance of normal glands Pl. IX., fig. 1 with Pl. IX., fig. 3). In addition to hypertrophy there may be hyperplasia (Pl. IX., fig. 4), the glands increasing in number either through lateral branching or the ingrowth of new ones from the surface. In the *interstitial* (Pl. IX., fig. 2), the stroma is affected—in recent cases its cells, in more chronic the intercellular substance. In the recent cases there are abundance of small cells (like nuclei only, from the small quantity of their protoplasm), which, if recovery does not take place, pass into spindle-cells arranged in interlacing bands; sometimes they swell up and take on a decidual character—the nuclei




FIG. 191.

GLANDULAR TYPE OF ENDOMETRITIS (*De Sinéty*). 1. Stroma; 2. Dilated glands. (49)

becoming larger and containing nucleoli. In the chronic cases, the intercellular substance is thickened by exudation and its fibres increased and thickened.

In endometritis after abortion, small islands of decidual cells (which have not undergone retrograde changes as rapidly as the rest of the decidua) are sometimes found with small-celled proliferation going on round them (Pl. IX., fig. 5).

A special form of endometritis was carefully described by Olshausen Olshausen's Endometritis Fungosa. under the name of *E. fungosa*. In it the mucous membrane is hypertrophied to three or four times its normal thickness. It is elevated through its whole extent in a soft cushion-like swelling, or in more localised spongy masses; the hypertrophy does not extend beyond the os internum to the cervix, and thus resembles in its situation a decidual membrane. The portions removed by the curette are unusually thick; one side presents a smooth rose-coloured surface which resembles the

appearance of the mucous membrane of the intestine, and the other has a deep-red raw surface. "The microscopic examination of these scrapings," Olshausen says, "shows that there is great hypertrophy of the mucous membrane with increase of all its elements—moderate dilatation of the lumina of the glands, enlargement of the blood-vessels, and marked cellular infiltration of the connective tissue." The characteristics of this type are, that the glands do not become enlarged so as to produce cystic dilatations, and that the blood-vessels are distended; the latter fact explains the hæmorrhage which is the chief symptom. The small-celled infiltration is shown in the sections given in figs. 192, 193.

In some cases of endometritis fungosa, Zeller found that portions of the exfoliated mucous membrane consisted of squamous epithelium arranged in several layers—a sort of psoriasis uterina. This shows that columnar epithelium may change into squamous, a fact of great



FIG. 192.

SECTION OF ENDOMETRITIS FUNGOSA—
Low power; showing changes,
chiefly interstitial (from a micro-
photograph).




FIG. 193.

SAME—High power, to show the small-celled
infiltration of the connective tissue.

interest with regard to the changes in catarrhal patches described in the preceding chapter.

Heinricius'
view of
Endome-
tritis
Fungosa.

Heinricius has described the scrapings taken from a large number of cases of fungous endometritis. A thin section with sparing infiltration gives under a high power¹ the appearance seen in Pl. IX., fig. 6. The stroma between the glands (the epithelium of which is seen in the corners of the section) consists of a basis of stellate corpuscles with anastomosing processes, upon and between which lie two varieties of cells—large, oval, faintly stained ones, and others, small, round, and deeply stained, the former being the nuclei of an endothelium, the latter lymph corpuscles. He thus agrees with Leopold that the interstitial tissue

¹ Zeiss, Ocular 3, Water immersion K.

Beiträge zur pathologischen Anatomie des Endometrium: Archiv f. Gyn., XXXIV., S. 165.

consists largely of lymph sinuses. When inflammation occurs, the lymph corpuscles and those of the endothelium proliferate and produce an appearance resembling a "small-celled infiltration," for the basal network is obscured by them. He thus comes round to practically the same condition as Olshausen has described, but assigns a different position to the small cells.

Another form of endometritis is described by De Sinéty. "In other

Landau and Abel deny the existence of a hyperplastic glandular form of Endometritis and would recognise only the *E. fungosa*, making the cases of hypertrophied glands a localised *Adenoma simplex*. Their argument is that the changes in any inflammation are primarily in the interglandular tissue, the changes in the epithelium of the glands being so to speak accidental and the result of the hyperæmia. The "cork-screw-like hypertrophy" is a normal condition. Where the glands actually grow, it is an *Adenoma*. Further, as to the interglandular changes, the decidual cells described by Ruge are not peculiar to the uterus, but simply the large epithelial cells (fibro-blasts) which are an intermediate stage in the formation of connective tissue from inflammatory products in any situation.




FIG. 194.

CROSS SECTION OF GRANULATION COMPOSED OF EMBRYONIC ELEMENTS, FROM A CASE OF ENDOMETRITIS. 1. Embryonic tissue; 2. Part undergoing fatty degeneration (*De Sinéty*) (A^o).

cases," he says, "the vegetations are specially constituted of embryonic tissue with few blood-vessels. There are only traces of the glands and some remains of more or less degenerated epithelium. We have to do with a truly inflammatory tissue comparable to that which forms upon an exposed wound. At certain points there are islands of degenerated elements which are not coloured by reagents and are analogous to those observed in foci producing pus. The degeneration of embryonic elements explains to us the abundance of the muco-purulent discharge observed during life" (fig. 194). Slavjansky also has described a villous or papillary form of endometritis, in which the mucous membrane has lost its epithelial covering and has its inner layer composed of embryonic connective tissue.

Villous or
Papillary
form of
Endome-
tritis.

Cornil, who has also described the changes in the mucous membrane in endometritis, maintains that it is a true inflammatory

process from the excess of mucus, the multiplication of the epithelium, and the migration of leucocytes. In the gland-cells, karyokinesis is often seen.

Ultimate
changes in
Endome-
tritis.

When chronic endometritis has persisted for a long time, the mucous membrane becomes atrophied; the ciliated and afterwards the cylindrical epithelium is lost, and small polymorphous cells resembling squamous epithelium take their place; finally, the mucous membrane disappears altogether and the uterine cavity comes to be lined with a layer of connective tissue. The glands fall out so that the mucous membrane becomes mesh-like, or they are constricted to form retention cysts.

Senile
Endo-
metritis.

Senile atresia of the cervical canal is the result of a localised chronic endometritis. This is one of the physiological changes which occur after the menopause. In some cases, however, it becomes pathological; accumulation of mucus, more rarely of blood, takes place above the obstruction.

Relation of Micro-organisms to Endometritis.

Looking at endometritis from a bacteriological standpoint, Winckel¹ has classified the various forms in two groups, according as they are due to micro-organisms or not. In the latter group he places (1) simple catarrh due to disturbance of circulation, as in chlorosis, uterine displacement, faults in dress, mode of life, etc.; (2) hæmorrhagic endometritis, as in acute and infectious diseases; (3) decidual endometritis, after abortion; and (4) exfoliative endometritis. In the former group he places (5) gonorrhœal endometritis; (6) tubercular endometritis; (7) puerperal septic endometritis, usually due to the streptococcus longus, more rarely to a staphylococcus or to the bacterium coli commune; (8) saprophytic endometritis due to combinations of cocci and bacilli, of which the senile purulent endometritis is probably one form; (9) the so-called diphtheritic endometritis which is due to streptococci; (10) syphilitic endometritis—the cervical mucous membrane exposed by laceration being a favourable nidus, but infection of the decidua the more important cause; (11) endometritis due to amœbæ—protoplasmic bodies with nuclei and vacuoles being present in the dilated uterine glands, and causing proliferation of epithelium.

This classification raises the question as to the part played by micro-organisms, which can only be answered by observations made directly on the endometrium. Bumm,² from the examination of the secretions of forty-five cases of endometritis, concludes that the micro-organisms found in some cases of endometritis are accidental, and that neither the hyperplastic nor catarrhal varieties are dependent on them for their development. On the other hand, endometritis may follow an acute septic or gonorrhœal infection.

¹ Centralb. f. Gyn. 1895, S. 695.

² Ibid., S. 698.

Gottschalk and Immerwahr¹ have examined the secretions from sixty cases, and found that in twenty-one, that is in 34·5 per cent., no micro-organisms were present; of the remaining thirty-nine, twenty-eight showed non-pathogenic micro-organisms (mostly diplococci), while eleven showed staphylococci. The results of these observers thus agree with those of Pfannenstiel and Döderlein.

We must also remember that even the presence of micro-organisms does not prove their activity. It is not enough to find them; the question is whether they can grow, that is to say, under what conditions they are hurtful or harmless. In a very interesting paper on the flora of the female genital tract in health and disease,² Menge has maintained that the secretions of the cervical canal destroy bacteria, and that the vagina aids the cervix in its endeavour, so to speak, to keep the upper portion of the genital tract free from germs. He concludes that in the normal endometrium of body and cervix and their secretions, and even in chronic hypertrophied conditions of the mucosa, with or without small-celled infiltration, germs are not present, with the exception of the gonococcus, which even healthy tissues are unable to resist. Staphylococci and streptococci if present during an acute endometritis, disappear in a chronic stage. It is possible, however, that these latter may have a share in the etiology of endometritis through persistent anatomical changes in the endometrium, excited by them during the acute stage. He associates the presence of bacteria with these four conditions:—gonorrhœal endometritis (gonococcus); tubercular endometritis (tubercle bacillus); acute endometritis (streptococci, staphylococci, sapræmic bacteria, bacterium coli commune); septic collections in the uterus, as retained placental products, or after enucleation of polypi.

While evidence is thus accumulating that micro-organisms do not play so important a rôle in the etiology of endometritis as Doléris and others have held, it does not follow that we must minimise antiseptics in its treatment. On the contrary, the greatest antiseptic precautions must always be used lest we introduce micro-organisms which were not there before. Unless the external genitals and vagina be thoroughly cleansed, and aseptic instruments be used, a purulent endometritis may be induced on a simple one. For the same reason, forcible dilatation should not lightly be had recourse to, and all forms of treatment should be avoided which imply repetition in their use. Another practical deduction is that removal of the mucosa by the curette is not called for in all cases of endometritis; and to draw a picture of a germ-laden mucosa being thus removed in order that a healthy one may grow under our antiseptic precautions, is to push the germ theory too far.

¹ Archiv f. Gyn. 1896, Bd. L., Heft 3.

² Centralb. f. Gyn. 1895, S. 795.

ETIOLOGY.

Acute endometritis is a rare condition, and never occurs before puberty. It comes on most frequently in connection with menstruation, when the physiological congestion readily passes into inflammation. It is occasioned by exposure to cold or sexual excess at the periods, and by the extension of gonorrhœal inflammation from the mucous membrane of the vagina. It also occurs in the exanthemata,¹ typhus, scarlet fever, and measles; it has further been observed in cholera (*Slavjansky*), and in certain cases of phosphorus poisoning. In puerperal inflammation, endometritis is of course present.

Chronic endometritis is occasionally the result of acute; most frequently, however, it arises independently. Sometimes it is merely the indication of the constitutional state; in scrofulous and chlorotic cases, the normal leucorrhœa (which precedes and follows menstruation) is increased in quantity and prolonged during the intermenstrual period. This is due to hypersecretion rather than to inflammation. Increased leucorrhœa, with diminished menstrual flow, is quite characteristic of phthisis.

Causes of
Chronic
Endome-
tritis.

Chronic endometritis arises independently from the following causes:—

- Parturition, specially when the uterus has not been completely emptied;
- Exposure to cold during menstruation;
- Polypi or other tumours in the uterine cavity;
- Some uterine displacements, *e.g.*, prolapse.
- Direct injury through incautious use of sound or tent;
- Extension of gonorrhœal or simple inflammation from vagina and cervix.

It has also been found after non-physiological amenorrhœa.

Of these the most important is *parturition*. Endometritis is frequent after abortion; usually this is due to the patient's rising too soon, or to the incomplete emptying of the uterus. Küstner has traced the transition of a portion of decidua, retained after abortion, into a tissue having the structure of a mucous polypus. As to the frequency of this occurrence, he says that, of 112 cases of endometritis, nine were cases of deciduoma. After full-time labour, the seat of the placenta seems to be in many cases the starting-point of the inflammatory process.

Uterine displacements do not necessarily produce endometritis. We sometimes find a retroversion or retroflexion which has produced no symptoms. As a rule, chronic inflammation of the endometrium, as well as of the muscular coat, follows passive congestion.

¹ Massin found acute inflammatory changes in twelve cases of typhus, pneumonia and dysentery: *Archiv f. Gyn.*, Bd. XL., Hft. 1, S. 146.

Brennecke¹ and, more recently, Heinricius² have drawn attention to the occurrence of endometritis following non-physiological amenorrhœa. After irregular menstruation (at longer or shorter intervals), or complete amenorrhœa, profuse bleeding takes place from the uterus. It is most common in patients towards the menopause, but has also occurred in anæmic or poorly nourished girls. They ascribe it to lowered activity of the ovaries so that the hyperæmia at the menstrual period leads only to hyperplasia of the uterine mucous membrane, not to hæmorrhage; hence the mucous membrane becomes hyperplastic, and when hæmorrhage does return it is profuse.

SYMPTOMS.

A. *Of Acute Endometritis.*

These are fever more or less severe, according to the inflammation, pain in the back and lower part of the abdomen with the sensation of weight in the pelvis, and in severe cases vesical and rectal tenesmus. The characteristic symptom is the discharge, which is at first clear and watery, but after a few days becomes creamy and purulent. The menstrual flow is sometimes suppressed, rarely is it increased.

B. *Of Chronic Endometritis.*

The symptoms usually given are the following:—

- Menorrhagia;
- Leucorrhœa;
- Dysmenorrhœa;
- Weakness in the back;
- Pain in pelvis and loins;
- Digestive derangements;
- Sterility;
- Abortion.

Menorrhagia is the characteristic symptom, and may become serious from the anæmia which it produces. It shows itself first in increased duration of the menstrual flow, which becomes gradually prolonged over the intermenstrual period till the loss of blood becomes continuous. *Dysmenorrhœa* is frequently present, but it is more probably due to complications (*e.g.*, cellulitis or chronic metritis) than to the condition of the mucous membrane. Membranous dysmenorrhœa (accompanied with exfoliation of the mucous membrane at the menstrual period) might be considered here, as its pathology is most nearly allied to endometritis; from its peculiar symptoms, however, it is better to consider it in the chapter on Dysmenorrhœa (Section VIII.).

¹ Archiv f. Gyn., XX., S. 455.

² Op. cit.

*Leucorrhœa*¹ is a frequent symptom. The secretion from the body of the uterus is of a watery character, less dense and gelatinous than that from the cervix; usually, however, there is cervical catarrh as well. The uterine secretion has an alkaline reaction, while vaginal leucorrhœa is acid. Sometimes it is tinged with blood, producing an appearance which Bennet compared to the rust-coloured sputum in pneumonia. The blood-stained leucorrhœa must not be confounded with the menstrual flow. In some cases the discharge is purulent, accumulates in the uterine cavity, and is only discharged at intervals.

"*Weakness in the back*" is the common complaint made by the patient. It may amount to actual pain, but more generally it shows itself as feebleness or weariness which incapacitates the patient for her daily work.

Derangements of the digestive and nervous systems invariably follow when the disease has become chronic. There is impaired digestion with loss of appetite, and, as the result, general debility. Whether these are due to the drain on the system produced by the leucorrhœa or to the close connection between the nervous centres for the sexual organs and those for the digestive apparatus, we do not know. Derangements of the nervous system show themselves in frontal headache and depression of spirits amounting sometimes to melancholia.

Anæmia, with its characteristic train of symptoms, is the leading symptom in the hæmorrhagic type (*Olshausen*).

Sterility is frequently present, and has been in certain cases the only symptom complained of. The secretion may destroy spermatozoa, may mechanically prevent them from passing upwards, or the villi of the fertilised ovum may be prevented from finding an attachment in the diseased mucous membrane. Again, the ovum is attached for a time, but from the imperfect formation of the uterine portion of the placenta, *abortion* takes place; repeated abortion is characteristic in chronic endometritis. A vicious circle is thus produced: as mentioned under etiology, endometritis frequently follows abortion; abortion, in its turn, frequently follows endometritis.

PHYSICAL SIGNS.

A. *Of Acute Endometritis.*

There is tenderness on pressure over the lower part of the abdomen due to peritonitis which generally accompanies the acute form. On vaginal examination the cervix is found to be swollen and puffy, the os is dilated and feels velvety from eversion of the mucous membrane, the bimanual is unsatisfactory from sensitiveness to pressure. The speculum shows the vaginal portion to be congested, with catarrhal

¹ We mention this as a symptom usually given, although proof is wanted that the secretion from the uterine mucosa is increased in Endometritis—it may be entirely cervical.

patches round the os and the follicles enlarged and sometimes containing pus. The leucorrhœal discharge already described is seen coming from the os uteri. The sound should not be used, as its introduction causes pain and sometimes hæmorrhage.

B. *Of Chronic Endometritis.*

Tenderness on pressure is not necessarily present, though we frequently find it as the result of complications—peritonitis, cellulitis, ovaritis.

On vaginal examination the vaginal portion of the cervix is normal, or has the characters described under cervical catarrh. The bimanual shows the uterus to be *enlarged*; it is soft and flabby so that its form cannot easily be made out, or of a firm consistence from chronic metritis.

The sound passes *beyond* the $2\frac{1}{2}$ -inch knob to a varying extent, and on withdrawal is frequently tinged with blood. Its introduction may be difficult from irregularities in the mucous membrane, and is sometimes painful from associated peritonitis. It is most useful in demonstrating *irregularities of the mucous membrane*, and their recognition is of great importance: to detect these, the sound is held lightly between the finger and thumb and moved slowly backwards and forwards over the mucous membrane; a grating or catching sensation is felt when they are present. We must note, however, as Olshausen points out, that the spongy irregularities may escape detection by the sound.

In the speculum we see, issuing from the os, the leucorrhœal discharge with the characteristics given above; usually it is mixed with that from the cervix. The appearances described under cervical catarrh are also frequently present.

DIAGNOSIS : DIFFERENTIAL DIAGNOSIS.

The curette is invaluable in diagnosis, especially when its use is followed by *microscopical examination* of the scrapings—the importance of which here cannot be overrated.

Value of microscope in diagnosis.

This throws light on the etiological question, whether the endometritis be due to incomplete emptying of the uterus after *parturition*. In such a case we find among the scrapings large decidual cells or fragments of the villi of the chorion in a state of fatty degeneration.

It enables us to differentiate endometritis from commencing malignant disease—*carcinoma* and *sarcoma*. In carcinoma we see under the microscope abundance of epithelial cells of irregular form and with many nuclei (*v. fig. 269*). In sarcoma we see under the microscope the typical round or spindle-shaped cells. The hæmorrhagic type of endometritis may readily be mistaken for sarcoma uteri, because “it

spreads in a diffuse manner, pre-eminently causes hæmorrhage, produces pain not at all or only late" (*Olshausen*). The microscope, however, settles the diagnosis. Care must be taken not to mistake the small-celled infiltration of the tissue (fig. 193) for round-celled sarcoma. The cells of the latter are characterised by their larger size and oval nuclei (*v.* figs. 281 and 284).

PROGNOSIS.

Endometritis is not a fatal disease in itself, though, when long protracted, it seriously affects the constitution and produces permanent ill-health. In cases of excessive hæmorrhage, the condition becomes grave.

The treatment is often protracted, and the patient should always be warned of this. The occurrence of conception will produce the most favourable conditions; and, if due care be taken to prevent abortion in the early months, and in the management of the puerperium, we may hope for a cure.

When endometritis is associated with a strumous, tubercular, or syphilitic diathesis, it may baffle all our efforts.

TREATMENT.

A. Of Acute Endometritis.

Treatment
of Acute
Endome-
tritis.

Rest in bed, warm fomentations over the abdomen, and the free use of opium if there is much pain, form all the treatment required. Should the bowels not be moved freely before the attack, castor oil with an enema should be given since the loaded rectum presses injuriously on the inflamed uterus. Should the bowels not be loaded, the patient is not to be troubled with purgatives but rather kept under the influence of opium. If there is menorrhagia, ergot is required; when the discharge is free, it is to be given hypodermically. Warm water injections should not be used until the acute stage is passed, the pain and other signs of inflammation have subsided, and the leucorrhœa is abundant.

B. Of Chronic Endometritis.

Of Chronic
Endome-
tritis.

Prophylactic treatment is of great importance. A patient who is subject to endometritis should guard against exposure during the menstrual period. When conception takes place, the practitioner should remember the liability to abortion, the importance of seeing that the uterus be thoroughly emptied after parturition, and that the patient take proper care during the puerperium; during the latter period, ergot is beneficial.

We begin with hot-water injections, and the administration of ergot; this is given as the liquid extract (twenty drops in water three times a day, increased to thirty at the menstrual period) or ergotin—four grains in pill, daily.

If the uterine cavity be enlarged so that the sound moves freely within it, if there be roughness of the endometrium, or if there has been a recent miscarriage or confinement, we employ the *curette followed by the application of carbolic acid*. In the last class of cases the cause of the endometritis has been the incomplete separation of the decidua; if treated while still recent, such cases furnish the most satisfactory instances of an immediate and complete cure. Curetting
of Uterus.

The fixing of the uterus by adhesions or cicatrization does not contraindicate the operation, though these render it more difficult through preventing the uterus from being drawn down by the volsella; when they are present, undue traction must not be made. The time selected for operation is a week after a menstrual period; when the discharge is continuous, the period is indicated by increase in amount.




FIG. 195.

SOUND DRESSED WITH WADDING FOR THE APPLICATION OF CARBOLIC ACID.

Curetting of the Uterus with Application of Carbolic Acid. The following instruments are necessary:—

- Sims' or Fritsch's speculum,
- Three or four sounds dressed with cotton wool,
- Volsella,
- Curette,
- Crystals of carbolic acid liquefied,
- Cotton wadding and glycerine,
- Vaginal and intra-uterine douche.

Though chloroform is not necessary, its use makes the operative manipulation much easier, and allows it to be done more thoroughly.

The sounds should be covered with a thin layer of cotton wool, extending almost to the knob (fig. 195). The sound is dressed as follows:—A film of cotton wadding is laid on the palm of the left hand, the last two and a half inches of the sound are moistened and pressed firmly on the cotton wadding, the left hand is closed over it, the sound is turned twice

or thrice round within the shut hand till the cotton wadding becomes tightly rolled on. The dressing must bite the sound firmly so that it may not come off within the uterine cavity, and must not be too thick to be easily carried in. To remove the cotton wadding afterwards, the dressing is unrolled under water. The crystals of carbolic acid are kept in stoppered bottles, and at the ordinary temperature a portion remains liquid. Tincture of iodine, strong nitric acid, or chromic acid may be substituted for it. Chloride of zinc has been used but it causes coagulation in the vessels and necrosis of the mucosa which is shed along with a thin layer of the muscular coat.¹ Hence carbolic acid is to be preferred to it.

Operation
of Curet-
ting.

The patient is placed semiprone; Sims' speculum is passed and held by an assistant who with the left hand draws back the upper labium—if there be no assistant some form of self-retaining speculum is used; the vagina is washed out with an antiseptic, and the cervix fixed with a volsella. The curette shown at fig. 101 is the best. The uterine wall is scraped from side to side and from above downwards, special care being taken to remove the tissue from the Fallopian-tube angles and the sides of the cavity. The detached fragments are brought down with a raking motion or washed out with an intra-uterine douche through a Fritsch's catheter. A sound, dressed with dry cotton wadding, is passed to clear away the blood and mucus; the same process is immediately repeated with a second, and with a third if necessary. A reserve sound, previously dipped in the carbolic acid so as to be ready for use, is carried in immediately after the last of these has been withdrawn; if there is much bleeding or the uterine cavity is large, a second application should be made; our aim is to apply the carbolic acid to the whole of the raw surface, without its being diluted with blood or mucus. The volsella being withdrawn, a pledget of cotton wadding soaked in glycerine is placed in the upper part of the vagina so as to embrace the cervix; this prevents the carbolic acid from running down into the vagina.

The patient keeps her bed for two days after the operation, the pledget having been removed on the second day. Special care should be taken at the next menstrual period.

Curetting
for Tubal
Inflamma-
tion.

Curetting, and especially curetting followed by packing the uterus with gauze so as to ensure drainage, has recently been advocated on account of its secondary influence on the condition of the Fallopian tubes. Whether it does this by promoting drainage from the tube, or by an osmosis as Polk suggests, or simply by cutting off a source of infection from the endometrium, we cannot say; but undoubtedly certain tubal conditions are benefitted by it. Though we have not yet data for laying down rules, it is evident that discrimination between cases is

¹ Dumontpallier—Gaz. des Hôpitaux, 1890, No. 57, p. 521.

necessary. For example, while curetting would benefit a catarrhal salpingitis, the operative manipulation which it implies might be disastrous in a pyosalpinx. Curetting is now also advocated in conditions where some years ago it was tabooed. Chronic peritonitis and cellulitis, which were before considered as contra-indications, are now in certain cases regarded as a reason for removing the source of infection from the uterus which keeps up the inflammatory condition.

The regeneration of the uterine mucosa after curetting has been recently studied by Werth¹ in uteri removed by extirpation which had been previously curetted. The uterine mucosa is apparently reproduced much more rapidly than one would expect (within five days), and not by the process (formation of granulation tissue) which occurs after, say, the scraping of an ulcer on the skin, but by a process apparently peculiar to the uterine mucosa.

Regenera-
tion of
Mucous
Membrane
after
Curetting.

He examined six uteri that were extirpated after the operation of curetting at periods varying from three to sixteen days. In two of them extirpation had not been resolved on when curetting was done, but became necessary later; in the other four the curetting was done as a preliminary when the patient was being examined some days previous to the extirpation. The microscopic examination showed that the action of the curette varies in different parts of the same uterus, the mucosa being untouched in places (usually at the fundus and in the angles), while in others it is removed down to the muscular wall. It is the mucosa over the anterior wall rather than the posterior, and low down rather than high up in the uterus, that is most thoroughly removed. The uterus five days after curetting shows the regeneration changes most evidently. The interior of the uterus is entirely clothed with the new mucosa, with glands opening freely on the surface, and an unbroken covering of surface-epithelium. This new mucosa is characterised by its richness in fibrillar connective tissue as against cellular elements. The vessels play the chief part in its construction. They extend outwards, carrying with them a mantle of fibrillar connective tissue which follows the division of the vessels, and forms a network of new tissue. The glandular element arises from the fundi of the uterine glands which, embedded in the muscular coat, have escaped the curette. From these grow upwards new glands amongst the fibrillar tissue, usually straight and vertical. The cells are at first shorter towards the surface, and longer towards the bottom of the glands. The connective tissue often grows more rapidly, so that the glands come to open in pits on the surface. Its more rapid growth causes division of the glands, so that they open with more than one orifice, and also the irregularity of the surface of the mucosa. The surface epithelium is formed essentially from the glands, although the cells also multiply on the surface by division. New surface epithelium is found three days after curetting. In the later stages of the mucous membrane degeneration the excess of fibrillar connective tissue is removed by a hyaline degeneration which is apparent even five days after curetting, in the sub-epithelial layer. By the tenth day only a few bundles remain here, their place having been taken by large cells, mostly fusiform, with numerous protoplasmic processes. Where the curette has injured the muscular wall the loss of tissue is evident even seventeen days after, being marked by a layer of necrosed muscle and connective tissue covered by fibrin. Into this extend wide vessels surrounded by round cells, recalling the usual appearance of granulation tissue.

Applications without a previous curetting may be made in cases where there is no history of recent parturition or where the symptoms (menorrhagia) are slight. In all other cases the preliminary use of

Endo-
metric
Appli-
cations.

¹ Archiv f. Gynäk., Bd. XLIX., Hft. 3, S. 369. Baldy (op. cit., p. 231) also figures a uterus three months after curettage; and sections of the mucosa thirteen days after it.

the curette is a distinct advantage, as it removes the fungosities and thus allows the caustic to act more efficiently. Iodised phenol,¹ introduced by Battey, is a very useful and safe application.

Solid
Applica-
tions.

Sir James Simpson applied the nitrate of silver in powder on a porte caustique. The simplest way is to carry an ordinary quill with a nitrate of silver point into the cavity of the uterus; it may be passed in and withdrawn again, or held there till the point melts off; Credé of Leipsic claimed very good results from this mode of treatment. Barnes has devised an ointment positor for introducing ointments or fluids; he applies the iodide of mercury ointment in this way, and also tincture of iodine on a sponge. Dry iodoform has also been recommended by Kugelmann,² the powder being blown in through a curved metal catheter. Iodoform gauze has also been found useful by Polk³ in treating endometritis, especially the hæmorrhagic form; the cervix is dilated and the uterus washed out and then packed, the gauze being removed in twenty-four hours and if necessary re-introduced.

Electricity has been used in endometritis as in other chronic inflammations; this will be considered when the whole subject of Electricity in Gynecology is dealt with in the Appendix.

The importance of constitutional treatment must not be forgotten. The bowels should be moved regularly by saline aperients; the aloes and iron pill is also useful. The preparations of quinine, iron, and strychnine, are valuable in improving the tone of the nervous and digestive systems.

Cold baths and sea-bathing aid greatly in strengthening the constitution. The water of certain mineral springs, such as Ems and Kreuznach, seems to have a special action on the uterine as on other mucous membranes. The regular diet and exercise required at these baths have also, no doubt, their beneficial effect.

The diathesis—strumous, tubercular, or syphilitic—should not be forgotten. In them the treatment must from the first be constitutional.

Intra-
uterine
Injections.

Intra-uterine injections. Applications to the interior of the uterus are also made in the form of a fluid injected with a syringe. The nozzle of the latter is shaped like a sound, so that it may be passed into the uterine cavity; the barrel is of glass, and is graduated (like a hypodermic syringe) so that the quantity injected (not more than a few minims) is exactly known. The solutions used are carbolic or chromic acid, tincture of iodine or perchloride of iron, nitrate of silver, and sulphate of iron or copper. The cervix must be well dilated, to allow the fluid to escape readily past the nozzle of the syringe. To facilitate this reflux, syringes have been devised with a double canula. Injection of fluid *into the non-*

¹ Robert Bell in a paper read at the British Gynecological Society recommends it strongly—the proportions being 320 grs. of iodine dissolved in 8 ounces of liquefied carbolic acid: Brit. Gyn. Trans., 1888, p. 189.

² Centralb. f. Gyn., 1885, Bd. IX., S. 648.

³ Amer. Jour. Obs., 1888, p. 1052.

puerperal uterus is not unattended with risk¹ (*v. p.* 225), and the fact that we have the equally effective and perfectly safe method of intra-uterine medication described above renders it unnecessary. As a means of treating endometritis it is condemned by the general opinion of gynecologists in this country and America; in France and Germany, however, it is extensively practised.²

A method of *dilating the uterine canal for therapeutic purposes* was brought before the French Academy of Medicine by Vulliet.³ It consists in packing the uterus with tampons, varying in size from a pea to an almond, saturated in an ethereal solution of iodoform; the tampons are removed after forty-eight hours and a fresh series inserted, and the operation is repeated eight or ten times until the cavity has become so dilated that it can be explored through its whole length with a speculum, and applications made more thoroughly than after any other method of dilatation.

Taylor of Birmingham has devised an "artificial amnion" (a finger-stall of pure rubber, carried in on a hollow sound and distended with air) for dilating the cervix previous to making applications to the interior of the uterus, and its use as a preliminary to intra-uterine medication has been advocated by Park.⁴

¹ A fatal case has been recorded in the *Lancet*, April 16, 1887.

² For further details of this method the student may consult the following references: *Klemm*—"Die Gefahren der Uterininjection," Leipzig, 1863; *Cohnstein*—"Beiträge zur Therapie der chronischen Metritis," Berlin, 1868; *Leblond*—"Manuel de Gynécologie," p. 220, Paris, 1878; and *Hegar und Kallenbach*—"Operative Gynäkologie," S. 104, Stuttgart, 1881.

³ *Archiv de Toc.*, Oct. 1886.

⁴ *Edin. Med. Journ.*, Sept. 1887.

CHAPTER XXXII.

METRITIS, ACUTE AND CHRONIC: SUBINVOLUTION.

LITERATURE.

Barbour—Article "Inflammation" in Allbutt and Playfair's System of Gynecology: London, 1896. *Barnes*—Diseases of Women, p. 507: London, 1878. *Bennet, J. H.*—Practical Treatise on Inflammation of the Uterus: London, 1853. *Byford*—Medical and Surgical Treatment of Women: Philadelphia, 1881. *Courty*—Diseases of Women (Dr Agnes MacLaren's Translation: London, 1882). *De Sinéty*—Manuel de Gynécologie, pp. 315 and 351: Paris, 1879. *Fritsch*—Die chronische Metritis: Billroth u. Luecke's Handbuch f. Frauenkrankheiten, Stuttgart, 1885. *Gallard*—Traitement de la Métrite Chronique: Bull. gén. de thérapeut., etc., 1879, T. XCVII. 4—12, liv. *Guérin*—Ann. de Gyn., 1878, Tom. II. p. 9. *Klob*—Pathologische Anatomie der weiblichen Sexualorgane, S. 124: Leipzig, 1878. *Pozzi*—Traité de Gynécologie: Paris, 1897, p. 144. *Scanzoni*—Die chronische Metritis: Wien, 1863. *Schroeder*—Krankheiten der weiblichen Geschlechtsorgane, S. 84: Leipzig, 1879. *Simpson, Sir J. Y.*—Diseases of Women, p. 585: Edin., 1872.

DEFINITION.—Inflammation in the muscular coat of the uterus leading, when chronic, to increased formation of connective tissue.

ACUTE METRITIS.

PATHOLOGY.

The uterus is enlarged and may be of the size of a goose's egg; it is thickened, specially antero-posteriorly, and of a doughy consistence. The peritoneal surface is usually covered with lymph.

On section the muscular wall is thickened, but soft and pulpy; the cut surface is of a bright red colour, shows the veins to be engorged, and yields on compression a yellowish-red exudation. The mucous membrane is thickened and vascular, but the cavity of the uterus is not altered in size. Microscopically, the muscular bundles are infiltrated with pus corpuscles.

ETIOLOGY.

Acute metritis is produced by extension of inflammatory action from the mucous or serous lining of the uterus to the intervening muscular tissue. It occurs most commonly as part of the general inflammation

produced by absorption of septic matter during the puerperium. It also arises from exposure to cold at a menstrual period—the active congestion passing readily into acute inflammation,—from gonorrhœal infection and immoderate sexual activity.

Frequently, it is the result of surgical interference:—careless use of sound, intra-uterine injections, pessaries and sponge-tents; scraping the uterus, the removal of submucous fibroids, operations on the cervix.

SYMPTOMS.

There is fever and general constitutional disturbance varying with the intensity of the inflammation. The onset may be marked by rigors. There is a sensation of fulness, weight, and burning heat in the pelvis; pain in the hypogastric and sacral regions, aggravated on movement of the body or the emptying of the bladder and rectum; nausea and vomiting, diarrhœa and tenesmus of rectum and bladder.

Menstruation is suppressed in those cases where the metritis is occasioned by exposure to cold at the menstrual period. In other cases, it is diminished in amount; exceptionally there is menorrhagia.

PHYSICAL SIGNS.

There is tenderness on pressure in the hypogastric region. On vaginal examination, the vaginal walls are hot and dry, the cervix is swollen and movement of it causes pain. The bimanual examination cannot be made on account of the pain and the resistance of the abdominal walls; if the patient be put under chloroform, the uterus will be felt to be enlarged but freely movable unless fixed by old adhesions (fig. 203). The sound should not be used, as it causes hæmorrhage from the vascular mucous membrane.

PROGRESS AND TERMINATION.

The acute symptoms do not last usually more than a week. The fever and pain diminish; there is less heat in the pelvis and vagina, and leucorrhœal discharge becomes free. As complications, there may be catarrh of the bladder, rectum, or vagina.

The acute *usually* passes into the chronic stage to be immediately described; though *sometimes*, under proper treatment and care, there is resolution with absorption of the exudation; rarely does it terminate in abscess formation. Circumscribed abscesses in the uterine walls—recorded by Scanzoni, Reinmann, Bird, Ashford, Schroeder, Macdonald, and others—are sometimes produced and burst into the uterus itself; or adhesions may form and perforation take place into the bladder, vagina, rectum, and intestines, or even through the abdominal walls.

DIAGNOSIS.

The diagnosis that there is acute metritis and *nothing more*, is a refinement to which few would lay claim. But if the symptoms and physical signs are as described above, if the uterus be freely movable and no deposit is felt in the fornices, we may conclude that acute metritis is the prominent lesion. The possibility of abscess-formation should be kept in view. The practitioner may also, though very rarely, see cases where there is acute metritis and endometritis, and nothing else. It is wrong to say that acute metritis is rare. It is often a complication of pelvic peritonitis and cellulitis, with the physical signs masked by these latter diseases.

PROGNOSIS.

The *immediate result* will depend on the extent to which the peritoneum is involved. Even when the attack is not severe, the liability to pass into a chronic intractable condition makes us guarded in giving an opinion as to *complete recovery*.

TREATMENT.

Intra-
uterine
Injections.

If the metritis is supposed to be due to a septic cause, the first measure indicated is the *removal of that cause*. Thus if it come on during the puerperium, if the lochia are foetid and we suspect that a portion of the placenta has been retained, the uterine cavity should be washed out with an injection of 1 to 40 carbolic or 1 to 4000 corrosive sublimate solution. Great care must be taken not to introduce air with the injected fluid.

In all cases of metritis, the patient must be kept *at rest*. This is done by keeping her recumbent. The bowels are evacuated by an enema—not by purgatives—followed by a morphia suppository. Pain is relieved by warm fomentations, to which turpentine may be added, applied over the lower part of the abdomen; but if it be severe, the patient should be kept under the influence of opium as already described in the treatment of pelvic peritonitis. If the temperature be above 102°, quinine should be given—10 grains every two or three hours—till it falls. The sulpho-carbolate of soda (15 grains) is useful in some cases.

CHRONIC METRITIS.

SYNONYMS.—Chronic parenchymatous inflammation (Scanzoni), sub-involution (Sir J. Y. Simpson), diffuse proliferation of connective tissue (Klob), infarct (Kiwisch), areolar hyperplasia (Thomas).

There has been great divergence of opinion among gynecologists as to the term which should be applied to the changes occurring in chronic

metritis. Virchow describes the process as a hyperplasia of fibromuscular tissue, and places chronic metritis alongside of fibroid tumours of the uterus. Klob classes it among the new formations, and characterises it as “die diffuse Bindegewebswucherung”—“diffuse proliferation of connective tissue.” Thomas calls it “areolar hyperplasia,” and Noeggerath has suggested the term “diffuse *interstitial* metritis.”

From a *pathological point of view* the term “metritis” is incorrect, because there has never been demonstrated a chronic inflammation of the muscular fibre of the uterus. The morbid process described as chronic metritis consists in an *increase of connective tissue* out of proportion to that of the muscular fibre, which remains normal or is but slightly increased in quantity. We are not yet in a position to propose a term resting on a sure pathological basis; to do this would require a complete knowledge of the pathological changes, which has not yet been attained. We prefer to retain the term “chronic metritis.”

From a *clinical point of view*, this term is very convenient, including a variety of cases of different origin but presenting the same clinical features on examination.

It may be objected that to apply the term “chronic inflammation” to the process is misleading, as it implies a previous acute stage which is rarely present; the process would be more correctly described as an increased connective-tissue formation dependent on long-continued hyperæmia. But the term chronic inflammation is applied to the process producing similar changes in other organs, as cirrhosis of the liver; chronic metritis produces, in fact, cirrhosis of the uterus.

We have brought “subinvolution of the uterus” under this head, though in other English text-books it is treated as a separate lesion. Subinvolution of Uterus. The term subinvolution is *etiological* and simply expresses one mode, the most important one, in which the condition to be described is produced. *Apart from the history*, it is not possible to diagnose between a subinvolted uterus and one enlarged by chronic metritis alone. Further, the condition of subinvolution is maintained by the process of chronic metritis, that is, by the formation of connective tissue which takes the place of the muscular fibre. Finally, the treatment is the same in both cases.

PATHOLOGY.

The condition of the uterus depends on the duration of the disease. At an early stage (as in cirrhosis of the liver) the organ is enlarged, hyperæmic, and soft; at a later period it is indurated, anæmic, and hard. The peritoneal surface is of normal colour, or shows here and there patches of extravasated blood. The enlargement is uniform, so that the shape of the uterus is not altered.

On section, the tissue is soft and hyperæmic in the early stage; firm,

cartilaginous, and of a whitish colour (from the compression of the capillaries by the cicatricial tissue) in a later stage. The uterine walls are increased in thickness. The uterine cavity is increased in size.

De Sinéty. "In the first period," says De Sinéty,¹ "the dominant lesion is the presence in great number of embryonic elements throughout the whole thickness of the muscular wall. These elements are met with specially round the blood-vessels or form islands of variable dimensions which are more or less apart." The second period is characterised by two changes: (1) Marked dilatation of the lymphatic spaces, and (2) a localised hyperplasia of the connective tissue round the blood-vessels (fig. 196). The sclerosis, for such it may be called, differs from a




FIG. 196.

SECTION OF THE UTERINE TISSUE IN A CASE OF CHRONIC METRITIS ⁴⁹. *ct* connective tissue round the blood-vessels *bv*; *ls* dilated lymphatic spaces; *mf, l*, muscular fibre cut longitudinally; *mf, t*, muscular fibre cut transversely (De Sinéty).

similar change in the kidney or liver in the fact that the formation of connective tissue is localised round the blood-vessels. In the case described by De Sinéty, he says that it was difficult to say whether the muscular tissue was normal or diminished in quantity.

Fritsch. Fritsch² has examined uteri, extirpated for cancer, which showed the naked-eye characters of chronic metritis. He notes the following pathological changes. (1) The disposition of the muscular fibre and connective tissue is less regular than in the normal uterus, and the latter is increased in quantity. (2) The blood-vessels are more numerous and

¹ Gynécologie, p. 354.

² Luecke u. Billroth's Handbuch f. Frauenkrankheiten, Stuttgart, 1885, S. 917.

more tortuous ; the lumen of the vessel is often diminished ; the tunica media is thickened ; the contour of the vessel is masked through a connective tissue degeneration of its wall. (3) The lymphatic spaces appear gaping instead of as narrow clefts. (4) The peritoneum is thickened. Cornil¹ also describes a new formation of connective tissue between the muscular fibres. This tissue does not contract cicatricially, but produces a permanent increase in volume.

Snow Beck² also describes the presence of "an increased amount of round and oval globules, with amorphous tissue in the uterine walls." The increase in the size of the uterus is due to the presence of the soft tissue rather than to an increase in the muscular fibre.

ETIOLOGY.

The causes of chronic metritis may be arranged under two heads :—

- A. Causes which operate through interference with the normal involution of the puerperal uterus ;
- B. Causes which operate through the production of repeated or protracted congestion of the uterus.

A. Causes which operate through interference with the normal involution of the uterus.

- (1.) Retention of portions of placenta, membranes, or blood-clot in the uterus ;
- (2.) Lacerations of the cervix uteri ;
- (3.) Pelvic inflammations, occurring after labour ;
- (4.) Rising too soon after delivery ;
- (5.) Non-lactation ;
- (6.) Repeated miscarriages.

In the process of involution there are two factors, the fatty degeneration of the muscular fibre and the removal of the products of this degeneration. The condition of permanent enlargement or subinvolution is not due to the non-degeneration of muscular fibre, but to the substitution of connective tissue for the products of this degeneration. This seems to be the reason why the process of chronic metritis is met with more frequently in those who have borne children. John Williams³ made the interesting observation that involution was distinctly retarded by removal of the ovaries.

Any source of irritation in or beside the uterus leads to chronic metritis ; in this way we explain the effect of the *retention of portions of placenta or membranes*. An extensive *laceration of the cervix*, Emmet says, favours subinvolution for a similar reason. Continued *cellulitis or peritonitis* acts in the same way, or through interference with the circula-

¹ Leçons sur l'anatomie pathologique des Métrites : Paris, 1889.

² Lond. Obst. Trans., vol. xiii., p. 239.

³ Lancet, July 26, 1884.

tion. If the patient *rise too soon*, the increased weight of the non-involuted uterus leads to passive congestion and formation of connective tissue. Passive congestion will, on the other hand, be diminished by whatever produces uterine contractions; the physiological stimulus of suckling, excited reflexly through the mammæ, favours involution; in *non-lactation* this stimulus is absent. *Abortions* are an important cause; because patients do not take so much care of themselves as after a full-time labour, and the stimulus of lactation is absent. After abortion, conception readily takes place before the uterus has returned to its normal size, and this favours a recurrence of abortion.

B. *Causes which operate through production of repeated or protracted congestion.*

- (1.) Displacements of the uterus;
- (2.) Pressure of tumours in or near the uterus;
- (3.) Causes producing increased flow of blood to the uterus, *e.g.* endometritis or too free use of caustics.

SYMPTOMS.

In the great proportion of cases, the patient *dates her suffering from a confinement*; frequently there is a history of repeated abortions. The patient finds, on rising after the puerperium, that she does not regain her former strength. There is weakness in the back amounting in more severe cases to pain, a sensation of weight and bearing-down in the pelvis and of want of power in the limbs.

Menstruation is irregular and often increased in frequency and quantity, though this is more characteristic of endometritis. There is leucorrhœa from accompanying endometritis or cervical catarrh.

Effect on
Reproduction.

The *reproductive function* is variously affected. Before the structure of the uterus has become permanently altered, pregnancy followed by early *abortion* may repeatedly happen. The cause of the abortion is probably the alteration which is taking place in the structure of the mucous membrane, rendering it unfitted for the development of the placenta; after an abortion, the conditions are peculiarly favourable for a second conception even before the uterus has had time to undergo involution; an excessive development of connective tissue gradually renders the uterus incapable of involution, and thus the condition of subinvolution is perpetuated. Should the pregnancy go on to full time, the presence of an undue proportion of connective tissue in the uterine wall leads in the third stage of labour to atony of the uterus and retention of the placenta; see an interesting case of this reported by Kaschkaroff,¹ who gives the result of his microscopic investigation.

¹ Centralblatt für Gynäkologie, No. 5, 1879.

After the condition has existed for some time there is *sterility*. This is due not so much to the changes in the uterus itself, though the leucorrhœa may prevent fertilisation, as to the ovaritis or pelvic peritonitis which is usually superadded; ovulation may be prevented by change in the structure of the ovary or by its being bound down by adhesions; the Fallopian tubes may be obstructed by cicatricial contractions.

The general constitutional derangements are very important, and it is on account of these that the patients usually seek advice. Chronic metritis is the most important of all the diseases of women; the suffering of the patient in cases of displacement of the uterus is due not so much to the displacement as to the chronic inflammation secondary to it.

PHYSICAL SIGNS, DIAGNOSIS.

The uterus is *equally* enlarged; there is no alteration in its form. The character of the enlargement is best understood by contrasting it with that due to pregnancy. In the second or third month of pregnancy, there is antero-posterior enlargement of the uterus; the vaginal finger comes on the anterior wall springing out from the cervix; the abdominal hand feels the rounding out of the fundus, combined with a softness which prevents us from distinctly defining its outline. In chronic metritis the vaginal finger does not feel any bulging of the anterior wall, and the abdominal hand recognises the fundus to be uniformly thickened; the outline of the latter may be felt with unusual distinctness through the greater firmness of the uterine tissue.

The enlarged uterus may be in its normal position, and freely movable or fixed by adhesions; it is often retroflexed.

The sound passes more than the $2\frac{1}{2}$ inches; it passes *readily*, and is felt to be freely movable in the uterine cavity.

DIFFERENTIAL DIAGNOSIS.

The conditions which are most liable to be confounded with chronic metritis are *early pregnancy* and *small fibroid tumours*.

Diagnosis
of early
Pregnancy.

In a case of early pregnancy, the "having passed a period" will put us on our guard; some patients, however, menstruate after conception. Discolouration of the vagina points to pregnancy, but is often not marked. The softening of the cervix is a more reliable sign, unless pregnancy has occurred in a uterus which has undergone changes of chronic metritis. Our only sure guide is the bimanual examination, which shows us the change in the form and consistence described above. When the abdominal muscles are resistant, the finger can recognise per rectum the bulging and softness of the posterior uterine wall. The interesting question suggests itself in this connection, how soon it is possible to

recognise the changes in the uterus peculiar to pregnancy. *How soon can we diagnose pregnancy?* Before auscultation was known the first reliable signs were foetal movements; the date at which the mother first recognised these varied indefinitely. Auscultation gave us an earlier and more reliable indication in the sounds of the foetal heart; these cannot be heard before the fourth month. The bimanual examination enables us to detect pregnancy from the eighth to the tenth week. We have under very favourable circumstances diagnosed it at the fifth week, and the subsequent history has confirmed our diagnosis.

For the differential diagnosis of chronic metritis from small fibroid tumours, we refer the student to the "Diagnosis of Small Fibroid Tumours" (Chap. XXXVI.).

TREATMENT.

Our first object is to diminish the passive congestion of the pelvic organs. The patient should be instructed to lie down for a few hours every day. Sedentary occupations or those that require the patient to stand for a long time in one position should be avoided. While enjoining a certain amount of rest, we must remember that rest becomes injurious when it interferes with nutrition. A certain amount of exercise, especially in the open air, should be as emphatically prescribed as a certain amount of rest.

Passive congestion is also diminished by giving local support to the uterus by a Hodge pessary; where the vagina is roomy, a soft ring pessary sometimes answers better.

The pelvic circulation is stimulated by vaginal injections; hot water will generally be found to be the most valuable; cold water is a more effectual stimulus, but few patients can stand it. The vaginal injection should be employed just before going to bed; the douche is preferable to Higginson's syringe (*v.* page 150). The injection should be continued from ten minutes to a quarter of an hour. It is a decided advantage to have the douche given with the patient in the *dorsal posture*, as Gallard recommends. Occasional warm baths are useful in some cases; when the patient is in the bath, the vaginal douche can be used at the same time with greater freedom and effect. A cold hip-bath every morning is the best stimulus to the circulation. *Medicinal baths* have a peculiarly beneficial effect in chronic metritis. Amongst those the first place has always been held by Kreuznach, the waters of which are specially rich in bromides and iodides. The baths at Kissingen are rich in carbonates, and are of a lower temperature than those of Wiesbaden and Baden-Baden which contain a smaller proportion of salts.

Mineral
Waters in
Chronic
Metritis.

Further, the *drinking* of medicinal waters is also beneficial. The mineral springs at Ems and Vichy have, from their action upon the mucous membrane, always had a great reputation for the treatment of

chronic uterine inflammation. Where there is much catarrh, they are specially serviceable. In scrofulous and chlorotic individuals, the advantage of waters which are rich in salts of iron is evident. Comparatively few of our patients, however, will be able to enjoy the luxury of a course of treatment at one of these watering-places; but much benefit will be derived from change of air to the sea-side, or to the regular regime and cheerful surroundings of a hydropathic.

Attention to the action of the bowels is all important. Accumulations in the rectum and sigmoid flexure of the colon favour passive congestion, and interfere with the appetite and digestion. The mineral waters—Friedrichshall, Carlsbad and Hunyadi Janos—are the best aperients.

The Carlsbad salts are specially useful in bilious patients; a teaspoonful should be dissolved in a tumblerful of water and drunk in repeated sips during the morning. Friedrichshall and Hunyadi Janos waters act best mixed with an equal amount of hot water; their dose varies from a wineglassful to a tumblerful. A good substitute for these waters is the tonic and aperient prescription given on page 244.

Ergot (twenty drops of the liquid extract thrice daily, increased to thirty at the menstrual period) and the *Hydrastis Canadensis* (same dose of its liquid extract) are very useful, especially when there is menorrhagia.

The iodide and bromide of potassium may also be given internally, as recommended at page 241.

Great care, and in some cases complete rest, should be enjoined at the menstrual period. As exacerbations usually occur at these times, a great deal is done towards a cure by prophylactic measures in regard to this.

Of local treatment the most important is *counter-irritation* by occasional blistering or repeated application of iodine or of croton oil to the iliac regions. French gynecologists recommend the application of the blistering fluid to the cervix. Many gynecologists apply iodine to the cervix and roof of the vagina; Scanzoni recommended a solution of 4 grs. of iodide of potassium in 30 min. of glycerine. The simple tincture of iodine, or a solution of equal parts of iodine and glycerine, may also be applied in this way. *Local depletion* by scarification or leeches, as described under Endometritis, is less frequently employed than formerly.

In speaking of Emmet's operation, we mentioned that it was sometimes followed by diminution in the size of the uterus. Carl Braun¹ has shown that after *amputation of the cervix* for hypertrophy, the uterus sometimes undergoes changes which resemble those which occur physiologically in the puerperal uterus. Martin of Berlin strongly recommends the amputation of the posterior lip.

¹ Zeitschr. d. Ges. d. Wiener Aerzte, 1864, S. 43.

Electricity has also been recommended by Apostoli for chronic metritis; it is more properly a treatment of endometritis, as it is to its cauterising action on the mucous membrane that beneficial results are due. Weir Mitchell's method of treatment by feeding and massage has given good results where the constitutional weakness has been the chief source of trouble. Both of these will be considered in the Appendix.

CHAPTER XXXIII.

DISPLACEMENTS OF THE UTERUS: ANTEFLEXION; ANTEVERSION; RETROVERSION; RETROFLEXION.

LITERATURE.

Bandl—Ueber die normale Lage u.s.w. : Archiv für Gyn., XXII., 408. *Bantock*—On the Use and Abuse of Pessaries : London, 1884. *Barnes*—Diseases of Women, p. 679 : London, 1878. *Campbell, H. F.*—Pneumatic Self-replacement of the Gravid and non-Gravid Uterus : American Gynecological Transactions, Vol. I., 1876. *Croom, J. Halliday*—The Management of Anterior and Posterior Displacements of the Uterus : Brit. Med. Journ. 1888, I., p. 286. *Duncan, Matthews*—Diseases of Women, p. 403 : London, 1886. *Emmet*—Principles and Practice of Gynæcology, pp. 278 and 312 : Philadelphia, 1884. A Study of the Causes and Treatment of Uterine Displacement : Amer. Journ. Obstet., 1887, p. 1040. *Fritsch*—Die Lageveränderungen der Gebärmutter : Billroth und Luecke's Handbuch, Stuttgart, 1885. *Hart*—The Structural Anatomy of the Female Pelvic Floor : Edinburgh, 1881. *Herman*—On the Relation of Antelexion of the Uterus to Dysmenorrhœa : Lond. Obst. Tr., Vol. XXIII., p. 209. Pathological Importance of Flexions : Lancet, 1884, II., pp. 672, 729, 771. *Hewitt Graily*—The Mechanical System of Uterine Pathology : London, 1878. Importance of Flexions and Displacements : Lancet, 1884, I., pp. 1020, 1063, 1110 ; and Lancet, 1885, I., pp. 243, 284. The Early History and Etiology of Flexions of the Uterus : Brit. Med. Journ. 1886, II., p. 913. *Mundé*—The Curability of Uterine Displacements : Amer. Jour. of Obst., Oct. 1881. *Ruge*—Congenitale Retroflexio : Zeitschrift für Geburtshilfe und Gynäkologie, 1878, Band II., S. 24. *Schroeder*—Krankheiten der weiblichen Geschlechtsorgane, S. 140 : Leipzig, 1879. *Schultze, B. S.*—Ueber Versionen u. Flexionen u.s.w. : Archiv f. Gyn., Bd. IV., S. 373. Zur Frage von der patholog. Antelexion der Gebärmutter : ibid., Bd. IX., S. 453. The Pathology and Treatment of Displacements of the Uterus—English Translation by J. J. Macan : London, 1888. *Simpson, A. R.*—Displacements of the Uterus : Allbutt and Playfair's System of Gynecology, London, 1896. *Simpson, Sir J. Y.*—Diseases of Women, pp. 253, 245, and 764, Edin., 1872. *Thomas*—Diseases of Women, pp. 363, 408 : London, 1880. *Van de Warker*—The relation of symptoms to Versions and Flexions of the Uterus : Amer. Gyn. Trans., 1879, p. 334. *Vedeler*—Ueber Dysmenorrhoe : Archiv für Gyn., XXI., 211. *Wylie*—Prevention and Treatment of Antelexion and Anteversion : Amer. Jour. Obstet., 1884, p. 1261 ; and Edin. Med. Jour., XXX., 1148. The recent literature on the Operative Treatment of Displacements will be given under the various operations.

As the uterus is a movable organ within the pelvis, it is subject to Preliminary changes of *position* ; as it is composed of muscular tissue, it ^{varies.} is liable to alterations of its normal *curvature*. Both of these changes

are described in English text-books as "displacements," although, strictly speaking, this term should be applied only to the former.

The normal form, position, and relations of the uterus have been already described (see Chap. II.).

The uterus is constantly exposed to forces producing a temporary displacement. In front there is the *bladder*, the dilatation of which displaces the uterus backwards and somewhat upwards (fig. 37). Behind there is the *rectum*, which normally should have little influence on the position of the uterus; but, owing to inattention to its regular evacuation, it is frequently over-distended and thus acts as a displacing cause operating from above and behind. Above there is the *abdominal pressure*, which is constantly acting on the uterus especially during inspiration. One has only to watch the movements of the anterior vaginal wall during respiration to see that this factor is always operating. Its action is of course increased by whatever increases the intra-abdominal pressure, that is, by any straining efforts which bring the abdominal muscles into play.¹ Below there is the *pelvic floor*, which has a constant action in supporting the uterus against the abdominal pressure.

One of the most important contributions on the normal position of the uterus and displacements produced pathologically is from Ziegenspeck.² He examined the condition of the pelvis *post-mortem* in 56 cases, in 35 of which he had previously noted the condition during life according to Schultze's method. After describing the most important *post-mortem* changes, he mentions that he found the uterus anteflexed *post-mortem* in 24 out of the 56. His conclusions as to normal attachment of the uterus is thus summed up. The pelvic floor almost altogether supports and holds the anteflexed normally fixed uterus; the elastic traction of the vessels of the pelvic organs and of the peritoneum keep it in this anteflexed position. The uterus in this position is to a certain extent incorporated with the pelvic peritoneum, its attachment to the neighbouring organs being only of secondary importance. As to the pathological processes, he concludes that changes in the walls are only the result, never the cause of displacement. The fixation of the uterus was always more marked in cases of retroflexion than in those of pathological anteflexion. Peritonitic changes have little influence on the position of the uterus, while parametric ones are very important, being present in all cases of anterior and of posterior displacement: in anterior, affecting the utero-sacral ligaments; in posterior, the cellular tissue round the uterine vessels and beside the bladder and anterior fornix of the vagina.

Physio-
logical and
Patho-
logical
Displace-
ments.

We must distinguish between *physiological* and *pathological* displacement. The former is transient, and passes away when the cause has ceased to operate; the latter is persistent, and produces permanent alterations in form, position, and structure. It is difficult to draw the line between those two. The pathological condition is frequently due to simple overstepping of the limits of the physiological. Thus the carrying of the uterus backwards into a retroverted position by the

¹ Tight-lacing will intensify this action of the abdominal muscles. Braxton Hicks believes that a concave disposition of the abdominal muscles, found in spare women, prevents the bladder from expanding upwards and forwards and makes it either unduly antevert the uterus (if it be already pathologically anteverted) or retrovert it:—*Lancet*, 1886, I., p. 537.

² Ueber normale und pathologische Anheftungen der Gebärmutter und ihre Beziehungen zu deren wichtigsten Lageveränderungen: *Archiv f. Gyn.*, Bd. XXXI., S. 1.

distention of the bladder is physiological, while its remaining permanently in that position is pathological.

It is evident that the uterus can be displaced in at least three ways: *first*, the different parts of it may alter their position relative to one another; *second*, it may rotate round the transverse axis; *third*, the organ may be displaced as a whole. Any great rotation round the vertical axis is prevented by the attachments of the uterus.

1. Alteration in the relative position of body and cervix constitutes *flexion* of the uterus, in which there is a change in the curvature of the long axis, *i.e.*, in the direction of the uterine canal. Definitions.

2. Rotation of the organ round an imaginary transverse axis constitutes *version* of the uterus.

3. Displacement of the organ as a whole, although frequently observed, has not been described in English works by a precise term. We might use the term *position* with the suitable prefix. Thus when the uterus lies "back as a whole" in the pelvis, it might be described as "a retroposition" or as "retroposed" (*Germ.*, *retroponirt*).

The uterus, in its normal condition, is anteflexed, anteverted, anteposed—placed as far forward as the bladder will allow.

Various deviations from the normal condition may occur.

1. The normal curvature may be exaggerated—anteflexion.

2. The uterus may be straightened, the normal angle becoming less pronounced and thus throwing the cervix more backwards—anteversion.

3. The uterus may be directed backwards—retroversion.

4. It may not only be turned backwards but the normal angle may be reversed, the fundus being bent backwards instead of forwards—retroversion + retroflexion.

5. The uterus may be displaced as a whole, usually by cicatricial contraction. This last condition is the most difficult to treat.

The *etiology* of flexions and versions is a subject of great importance. Etiology.
In a certain number of cases they are congenital, a fact to be borne in mind specially with regard to retroversion. In many cases they result from inflammatory conditions,¹ pelvic peritonitis, and especially cellulitis (*v. p.* 200). A knowledge of etiology guides us in prognosis and treatment. It indicates what cases we may hope to cure, and what cases we should leave alone, and how we can prevent the occurrence of displacements, as, for example, of retroversion in the puerperal condition.

Of the *frequency* of forward displacements we have no data, as there is Frequency
no agreement as to what is to be considered a pathological degree of

¹ Ziegenspeck's researches confirm this from pathological anatomy, and Emmet (*loc. cit.*) has also from a clinical standpoint emphasised the importance of pelvic inflammation as causing versions of the uterus, and would limit the use of pessaries (invaluable in suitable cases) accordingly.

anteflexion or anteversion. As to backward displacements, Fränkel found them in 18 p. c. of gynecological cases.¹

Symptoms. The *symptoms* of displacements have given rise to much discussion, some maintaining that they produce no symptoms at all. We sometimes, on examining a patient, find a retroflexion which has not made its presence felt by any symptoms. This is however the exception; as a rule, backward displacements are followed by a train of symptoms. *This apparent contradiction is to be explained by the fact that flexions and versions, in themselves, give rise to no symptoms primarily.* The symptoms arise *secondarily*: they are due (1) to interference with the functions of menstruation, conception, and pregnancy; (2) to chronic metritis and endometritis, which is produced by the displacement; (3) to pelvic cellulitis and peritonitis; and (4) to accompanying inflammation of the uterine appendages. Bantock, in his interesting monograph on the Use and Abuse of Pessaries, gives very fully the various views held as to the significance of displacements as well as the results of his own experience.

ANTEFLEXION.

PATHOLOGY.

Anteflexion, as has before been stated, is merely an exaggeration of the normal condition. As to its frequency, there is great difference of opinion. The reason of this diversity is that a degree of flexion which would be called pathological by one observer would still be called physiological by another. The question of symptoms does not help us in deciding this; because, on the one hand, we sometimes find an extreme degree of flexion although the patient does not complain of any special symptoms; on the other hand, symptoms often described as characteristic are due to a different cause. It is in fact worthy of consideration whether we should not limit the term anteflexion, as descriptive of a special lesion, to cases of pathological anteflexion resulting from inflammatory conditions of the cellular tissue. Anteflexion is more frequent in nulliparæ, while retroflexion is more common in multiparæ.

The usual *seat of the flexion* is at the upper portion of the cervix, or at its junction with the body. Flexion of the body itself is rare. Sometimes the cervix is bent sharply forwards, so that it lies in the axis of the vagina and forms a distinct right angle with the body which is approximately in its normal position (see fig. 197). In other cases, the uterus is sharply curved on itself (see fig. 198). This last condition is sometimes mistaken for retroversion, because the finger feels through the posterior fornix the supra-vaginal portion curving backwards and the position of the fundus is not ascertained till the bimanual examina-

¹ In 936 of 5180 cases in public and private practice from 1882-85. He found retroflexion commoner than retroversion, as 645 to 291. Ueber die Erfolge der mechanischen Behandlung, u.s.w.: Archiv f. Gyn., Bd. XXIX., S. 316.

tion is made. In such cases the examination with one finger in the rectum is useful, as we can thus get above the point of flexion and feel that the fundus turns forwards.

The vaginal portion is frequently small and the os reduced to a pin hole (congenital cases); sometimes it is high up and difficult to reach, being drawn upwards and backwards by cicatricial bands. As regards the microscopic changes in the tissue, we are still in want of information. Virchow found no fatty degeneration of muscular fibre at the angle of flexion; the tissue was anæmic at this point but congested elsewhere. According to Rokitansky, the connective tissue




FIG. 197.

ANTEFLEXION WITH STENOSIS AT OS EXTERNUM. *V* vagina, *B* bladder, *p* peritoneum of pouch of Douglas (*Winckel*).

framework of the uterus is thinnest at the os internum; hence the liability to flexion at this point.

ETIOLOGY.

Etiologically we distinguish two kinds of ante-flexion, the congenital and the acquired.

In cases in which the ante-flexion is *congenital*, the whole uterus is imperfectly developed, the cervix is small and the pin-hole os looks downwards and forwards. Congenital
Ante-
flexion.

Acquired
Ante-
flexion.

As regards *acquired* antelexion, it is undoubtedly often the result of inflammatory changes behind the uterus. In many cases of antelexion, we observe that the cervix is higher than its normal position and far back in the pelvis; and that the attempt to bring it to its normal position produces pain. The cause of this condition was first brought into notice by Schultze,¹ who ascribes it to a cellulitis in the utero-sacral ligaments; this produces cicatricial contraction so that the cervix is drawn upwards and backwards, and the fundus thrown more forwards. Bandl thinks the first step in the process is a cervical catarrh; and that the inflammation spreads from the mucous membrane to the tissue of the cervix itself, making it more rigid, and thence to the cellular tissue round the cervix. Schroeder, however, held that the retraction of the




FIG. 198.

DIAGRAM TO SHOW ANTELEXION PRODUCED BY CICATRISATION OF UTERO-SACRAL LIGAMENTS. The arrow indicates the direction of the intra-abdominal pressure. (Schultze.)

cervix is produced by adhesions resulting from peritonitis. We draw attention specially to this cause of antelexion, because it can be distinctly made out by careful examination. When it has been made out it is a contra-indication to hasty operative interference, and the prognosis as to cure is unfavourable.

Hewitt's
Views.

Graily Hewitt referred this, as all other flexions, to softness of the uterine tissue and thinness of wall, producing undue flexibility.

It is alleged that a fibroma, or other tumour increasing the weight of the fundus, will favour antelexion if the fundus be directed forwards. In the commencing enlargement of pregnancy, the fundus droops more forwards or is at least more distinctly felt through the anterior fornix.

¹ Loc. cit., S. 414.

Unequal growth of the uterine walls has been given as the cause of congenital flexions, and unequal involution of the walls as the cause of flexions acquired during the puerperium. This is merely an explanation of how it is produced; the cause of this unequal growth requires, in turn, an explanation.

SYMPTOMS.

The most important symptoms of pathological antelexion are—

Dysmenorrhœa,
Sterility.

It will be noted that these are the symptoms of pelvic and uterine inflammation and are not pathognomonic.

In many cases we find a well-marked antelexion giving rise to no symptoms which patients complain of, as they are not accustomed to speak of sterility as a symptom.

Dysmenorrhœa. By this we understand that menstruation is accompanied with pain. The form of dysmenorrhœa present in antelexion has been called "uterine," in contradistinction to "ovarian" (see Dysmenorrhœa, Section VIII.). By "uterine dysmenorrhœa," is meant that the pain is not marked until the menstrual flow has appeared and that it continues as long as the discharge continues. The pain is felt in the small of the back and sometimes in the pelvis generally, but is not localised in one ovarian region.

Two different explanations of this pain have been given. For convenience we describe these as the obstruction and the congestion theories.

1. The *obstruction or mechanical theory.* According to this, the antelexion of the uterus produces a narrowing of the uterine canal at the point of flexion.¹ Hence, when the menstrual decidua and blood are shed, they find an obstacle to their free exit. There is consequent retention and coagulation, and the coagula stimulate the uterus to muscular contractions to effect their expulsion. The mechanical resistance to the outflow of blood and the uterine contractions excited to overcome this, are the cause of the pain. The condition is like that in stricture of the male urethra. The blood, like the urine, collects but cannot be passed without pain; there is dilatation with sometimes secondary hypertrophy of the uterus in the former case, as of the bladder in the latter. It may fairly be objected to this mechanical explanation that the discharge is not always clotted, that in some cases it is very small in quantity, that it is doubtful whether the blood coagulates in the uterus, and that in many cases the pains complained of have not the distinctive character of labour pains.

¹ It is doubtful whether this occurs. Graily Hewitt (Brit. Med. Journ. 1888, I., 461) figures a specimen where the lumen of the tube is flattened out laterally at the angle of flexion.

What has been already said with regard to Dysmenorrhœa ascribed to Stenosis of the Os externum (*v. p.* 299) holds good also here.

Congestion
Theory.

2. The *congestion theory* is clearly stated and advocated by Fritsch.¹ According to this gynecologist, the dysmenorrhœa is not due directly to the bend on the canal. The pain arises from the resistance which the muscular tissue of the uterus offers to the hyperæmia. In normal cases, this tissue yields to the distending vessels; but when the uterus is small or bent on itself, there is an obstruction offered to the flow of blood. The mucous membrane cannot swell up as it does normally. Thus there is undue vascular tension and compression of the nerve endings in the uterus. This last causes the pain.

Whether this explanation harmonises better with the facts it is difficult to say; but we should suggest a modification of Fritsch's view. The flushing of any diseased tissue with blood causes an aggravation of pain, which is increased if the tissue be of a dense structure. The intense pain in periostitis as the affected limb becomes warm in bed, is thus accounted for. Now the tissues of the uterus are frequently in a state of chronic inflammation, and there is sometimes increase of connective tissue making it of less yielding structure; this occurs in retroflexion complicated with subinvolution. The monthly flushing of the pelvis with blood would, under these circumstances, be accompanied with pain. We must also remember that cellulitis and peritonitis are often present with ante flexion; and increase of pelvic congestion will, of course, produce increase of pain.

Sterility.

Sterility is frequently associated with ante flexion; the patient is not so likely to refer to it, as the dysmenorrhœa is the more pressing symptom and that for which she seeks advice. This symptom has been referred to the obstruction in the uterine canal; as the menstrual blood is prevented from passing downwards, so the spermatozoa are prevented from passing upwards (*v. also p.* 300). But it is evident that this mechanical explanation is insufficient, because no mere contraction could prevent the passage of microscopic spermatozoa; and without doubt sterility is frequently the result of the binding down of the ovaries or the Fallopian tubes by concomitant inflammation. On the other hand there is the clinical fact that by passing the sound or dividing the cervix we place the patient under more favourable conditions for conception.

Dyspareunia—pain on sexual intercourse—is occasionally an important symptom, though naturally the patient does not refer to it. In such cases we generally find that there is inflammatory action behind the cervix.

PHYSICAL DIAGNOSIS.

On vaginal examination the cervix is short, conical and with a pin-hole os in congenital ante flexion, high up and drawn backwards in

¹ Loc. cit., S. 35.

acquired, in which also painful bands are felt in the posterior fornix (most distinctly on recto-vaginal examination). On bimanual examination, there is increased forward flexion of the uterus; and the whole uterus is displaced backwards in the pelvis in acquired ante flexion.

The sound is not necessary unless the bimanual is difficult; and should not be used when inflammation is present.

When the uterus is drawn backwards, the bimanual is difficult; hence from the direction of the cervix and the thickening behind it, such cases are often erroneously diagnosed as retroversion of the uterus.

DIFFERENTIAL DIAGNOSIS.

The only conditions which, after this careful examination, might yet be mistaken for an ante flexion are—

Myoma in the anterior uterine wall,

Cellulitis between the cervix and the bladder—a very rare condition.

Differ-
ential
Diagnosis
of Ante-
flexion.




FIG. 199.

SOUND PASSED TO SHOW THAT A MYOMA OF THE ANTERIOR WALL IS NOT AN ANTEFLEXION (*Leblond*).

For a small myoma, the sound is required to tell the direction of the uterine cavity (fig. 199). The bimanual, with the sound in the uterus (v. fig. 90) is very useful in such cases.

Diagnosis
from
Myoma,

The diagnosis from *cellulitis* is less easy, because through the tenderness it is difficult to ascertain whether the body felt in the anterior fornix is the fundus uteri or a cellulitic deposit. A careful bimanual examination will, if it be a cellulitic deposit, show that the fundus uteri is lying in some other position.

From
Cellulitis.

PROGNOSIS.

The prognosis should always be guarded in respect of the disappear-

ance of symptoms. The unfavourable cases are those in nulliparæ, due to utero-sacral cellulitis.

TREATMENT.

Pelvic inflammation, if present, must first be treated. Where the uterus is displaced by cicatricial bands, the stretching of these by massage has been suggested and is worthy of trial.

Treatment
by Sound.

The treatment of congenital ante flexion is symptomatic, that is to say, directed to the relief of dysmenorrhœa and sterility without our being able to say whether they correct any pathological condition or not.

When dysmenorrhœa is prominent, dilatation of the cervix gives the best results ; while for sterility, division is better. Dilatation is effected




FIG. 200.

GREENHALGH'S INTRA-UTERINE STEM.

by expanding dilators, or graduated bougies (see p. 301). When the uterine cavity is enlarged and menstruation profuse, curetting after dilatation does good.

For sterility, division of the cervix is the usual operation (see p. 302). It does good by relieving the retention of cervical mucus, as well as by allowing free ingress for the spermatozoa. Should the cervix be large, the cervical amputation is preferable to simple division.

Treatment
by Stems.

Intra-uterine stem pessaries have also been recommended. Fig. 200 shows a gutta-percha stem which is carried in on a uterine sound.

ANTEVERSION.

PATHOLOGY AND ETIOLOGY.

The *pathological change* consists in a straightening of the uterine axis, so that the normal angle of forward curvature is diminished and the cervix passes more directly backwards. The uterus is usually enlarged and its texture is firmer. In this condition it is movable or fixed. If the former, its position varies with the distention of the bladder ; if the latter, the fixed uterus will press more or less on the bladder as it distends and thus produce one of the symptoms of anteversion.

As Anteversion is the form and position taken up by the uterus when it is enlarged through chronic metritis, it does not require special consideration. Its causes, symptoms, and diagnosis will be found in Chapter XXXII. So also its treatment has been discussed there.

Various pessaries have been devised to support the enlarged uterus




FIG. 201.

GRAILY HEWITT'S CRADLE PESSARY. *a* is in posterior fornix; *b* at vaginal orifice; *c* in anterior fornix (*Barnes*).



FIG. 202.

THOMAS' ANTEVERSION PESSARY. A Hodge pessary with a bar projecting into the anterior fornix, which is believed to hold up the fundus.

through the anterior fornix, of which two are shown in figs. 201 and 202; as it is impossible to do this, their principle is unscientific. Any practical relief which they give is got equally by a ring pessary, the use of which has been referred to under Chronic Metritis.

RETROVERSION.

PATHOLOGY AND ETIOLOGY.

Physiological retroversion occurs whenever the bladder is fully distended (*v.* fig. 37). This is distinguished from the pathological condition by the fact that it is transient, and ceases when the bladder is emptied.

Pathological retroversion is found under the following conditions.

1. It occurs congenitally—which we assume when we find, on examining a virgin or nullipara, the uterus retroverted and either no symptoms or a history of symptoms going back to puberty. This is by no means a rare condition in virgins, as Küstner found it in 21 per cent. of private and 13 per cent. of hospital cases of backward displacement; and Graily Hewitt in 23 per cent. of cases (60 out of 259) noted in his private practice during thirteen years.

2. During the first days of the puerperium the uterus lies retroverted, or at least retroposed. The weight of the uterus and the laxity of its

Physio-
logical
Retro-
version.

Patho-
logical
Retro-
version.

attachments make it occupy this position when the patient is recumbent.

3. It is produced by the mechanism of prolapsus uteri (*v.* Section VII.). The axis of the uterus changes its direction as the organ descends.

4. It is also of importance as a stage in the production of retroflexion—the most frequent and important displacement which calls for treatment. The uterus becomes retroverted, and then acquires a backward flexion.

5. Chronic peritonitis producing obliteration of the pouch of Douglas, and cicatricial bands which drag the uterus backward, maintain, if they




FIG. 203.

UTERUS RETROVERTED AND BOUND BACK BY PERITONITIC ADHESIONS (*Winckel*). *aa* adhesions; *b* bladder; *v* vagina; *u* uterus; *r* rectum ($\frac{1}{2}$).

do not produce, retroversion—as is beautifully shown in the accompanying preparation from *Winckel's Atlas* (fig. 203).

The chief *causes* of retroversion are:—

1. A sudden straining effort, or a violent blow (a very difficult cause to establish);¹
2. Non-return of the uterus to its normal form and position during the puerperium;
3. Inflammatory action behind the uterus, producing adhesions in the pouch of Douglas; or cicatrization of the anterior vaginal wall.²

SYMPTOMS.

The symptoms of retroversion are the same as those found in retroflexion, to be presently described. When it arises during the puerperium,

¹ Graily Hewitt says that in 58 cases of backward displacement in virgins, nearly one-half (28 cases) traced their symptoms back to a severe fall, accident or strain; but this does not establish any of these as the cause.

² This acts by drawing the cervix forwards. Murdoch Cameron mentions a case where after division of a bridge on the anterior vaginal wall, the retroverted uterus became normal.—*Glas. Med. Journ.*, 1887, p. 420.

a late flooding—two to three weeks after labour—is sometimes a prominent symptom; or there is a daily loss of blood in small quantities whenever the patient rises and goes about (*Fritsch*).

DIAGNOSIS.

On vaginal examination, the cervix is low down in the pelvis and the os looks downwards and forwards. The finger feels the supra-vaginal portion of the cervix through the posterior fornix and may be able to reach the fundus, but the posterior surface is straight—there is no angle.

On bimanual examination, the hands can meet in the anterior fornix with nothing but the vaginal and abdominal walls between them. It is difficult to make out the body of the uterus. We may try to do this in two ways. *First*, with one finger in front of the cervix and the other behind it, lift the uterus upwards towards the abdominal walls; the hand placed on the abdomen will feel the anterior surface of the body of the uterus moving under it. *Second*, tilt the cervix well forwards with the index finger in the vagina, and thus increase the retroversion; the middle finger will feel the body of the uterus through the posterior fornix.

Bimanual
in Retro-
version.

The rectal examination is of great service here. The sound will pass as in fig. 86.

The differential diagnosis is the same as in retroflexion. The only point requiring special notice here is that we may have a retroversion with an ante flexion high up. Cases of ante flexion due to cicatrization of the utero-sacral ligaments are often, from the forward direction of the cervix, diagnosed as a retroversion (*v. p. 374*).

TREATMENT.

This consists in (1) removing existing inflammation; (2) replacement of the uterus when not fixed by adhesions; (3) retention of it in its normal position by pessaries: these will all be considered under retroflexion. Congenital cases should be left alone.

When adhesions are present, it is better not to interfere; or we may be content with supporting the retroverted uterus with a pessary.

RETROFLEXION.

For convenience this condition is usually called "Retroflexion," to distinguish it from "Retroversion" already described; strictly speaking, the condition is **RETROVERSION + RETROFLEXION**.

PATHOLOGY.

The pathological changes in the position and structure of the organs

in the pelvis consequent on retroversion + retroflexion, can be learned only from sections made with the organs *in situ*.

The following facts are based more on clinical examination than on pathological study. The changes in the various structures will be considered separately and shortly in a typical case of retroflexion in a multipara.

The *cervix* is directed downwards and forwards, or directly downwards (*v. fig. 206*). We observe clinically that it is much more easily reached. This is due partly to the alteration in its direction and position (being nearer the symphysis pubis it is more within reach), partly to the sinking down of the uterus as a whole in the pelvis. The os is patulous, because retroflexion usually implies previous parturition. If deeply fissured, it may form a gaping cleft which readily admits the tip of the finger. There is often ectropium and cervical catarrh. Some-




FIG. 204.

EXTREME RETROFLEXION OF UTERUS (*Barnes*).

times there is marked hypertrophy of the posterior lip, so that it is mistaken for the projection of the whole vaginal portion.

The *uterus* is flexed on itself, so that the fundus lies in the pouch of Douglas, the depth to which the fundus descends and the acuteness of the angle of flexion varying in different cases (*v. figs. 204 and 206*). If the condition of the uterine walls offers no resistance to flexion, the intra-abdominal pressure will tend to drive the fundus downwards till equilibrium is maintained—that is, till the fundus rests in the bottom of the pouch of Douglas.

Condition
of the
Uterus in
Retro-
flexion.

The size of the uterus is increased, and its cavity measures more than two and a half inches. Since the flexion generally occurs while the uterus is still enlarged through subinvolution, it is difficult to say whether this hypertrophy arises as the direct result of the displacement or through its interfering with the process of involution. What-

ever the cause of this hypertrophy is, its effect is to interfere with the natural cure of the displacement. The thickness of the uterine walls at the angle of flexion varies in different cases. Sometimes neither wall is atrophied at the point of flexion (fig. 204). Barnes says that according to his clinical experience this is the usual condition. On the other hand, Fritsch states that he has found marked thinning of the *posterior* wall at the angle of flexion. It is interesting to note that in a case of *congenital retroflexion* (see fig. 205) described by Ruge it is the *anterior* wall which is atrophied at the angle. The mucous membrane of the uterus is generally in a condition of chronic catarrh.

The microscopic changes consist in a dilated condition of the blood-vessels, with increase of connective tissue—the appearances produced




FIG. 205.

CONGENITAL RETROFLEXION (Ruge). Note the thinning of the anterior wall of the uterus.

by long-continued passive congestion. At the point of flexion, however, an opposite condition has been described; the blood-vessels were compressed and the tissues atrophied.

The *ovaries* follow as a rule the displaced fundus, the thin infundibulo-pelvic ligament stretching more readily than the ovarian. The position of the ovaries will, however, depend on the effects of peritonitic adhesions, which may fix them in any position. Sometimes we feel them below the fundus in the pouch of Douglas. They are frequently enlarged and tender on pressure.

The *bladder* is not necessarily altered in position, but has no longer the uterus resting upon it. The utero-vesical pouch is obliterated in

Ovaries in
Retro-
flexion.

Bladder
in Retro-
flexion.

cases of well-marked retroflexion. The ureters are often compressed or bent, which leads to dilatation; frequently they are found dilated to the thickness of the finger. Fritsch observed in one case the left ureter obliterated by a mass of cicatricial tissue, and the corresponding kidney changed into a sac full of white atheromatous debris.

The *rectum* may have the retroflexed fundus pressing against its anterior wall.

The *peritoneum* is altered in its normal relations as follows. The broad ligaments have their surfaces reversed, that is to say, the anterior, which was formerly inferior, is now superior; from their attachments, they offer no obstacle to retroflexion. The utero-vesical pouch necessarily disappears. The pouch of Douglas will, on the other hand, be distended by the fundus uteri; this implies *stretching of the utero-sacral ligaments* associated with the alteration in position of the cervix.

The *pelvic nerves* are occasionally affected, as shown by weakness in the lower limbs. This loss of power must be produced reflexly; from the anatomical relations, the retroflexed fundus cannot compress the motor nerves of the sacral plexus as is sometimes affirmed.

ETIOLOGY.

Retroflexion is, according to Fränkel's recent statistics, more common than retroversion.¹ As a *congenital* condition, it is not nearly so frequent as antelexion. It is more common in multiparæ than in nulliparæ, because the etiology is specially related to the *puerperal condition*. In this condition the uterus is enlarged and heavy and its walls are soft. The ligaments are lax, and the tissues of the pelvic floor have been recently stretched and have not recovered their tone. Through the distention of the bladder, the uterus is often thrown into a retroverted position.

We sometimes find on examining a patient shortly after her confinement that the uterus is lying back in the pelvis even though the bladder be not distended; we may thus suppose that the *intra-abdominal pressure* (which, when the uterus is in its normal position, is directed upon its posterior surface) comes now to act on the anterior surface and drives the fundus backwards and downwards. If the uterine tissue is soft enough to allow the fundus to be fixed on the cervix, such a flexion will gradually take place when the patient makes straining efforts. Apart from this, the *dorsal posture* and the common practice of *tight bandaging* after confinement will favour backward displacement of the fundus. If the patient *rise too soon* while the uterus is still large and heavy and the uterine supports correspondingly lax and weak, the tendency to displacement is increased.

The cause of retroflexion in nulliparæ is obscure.

¹ In 1882-85 he treated 936 retrodeviations of uterus, of which 645 were retroflexions and 291 retroversions.

SYMPTOMS.

The following are the more important local symptoms, to be found in cases of backward displacement.

Local
Symptoms
of Back-
ward Dis-
placement.

Weakness in the back,
Symptoms of chronic pelvic peritonitis,
Painful defæcation ;

Leucorrhœa,
Dysmenorrhœa,
Menorrhagia ;

Sterility,
Abortion.

In long-standing cases, there may follow the train of general constitutional symptoms consequent on chronic uterine disease.

The symptoms are arranged in three groups :—the first, including those which are more or less continuous ; the second, those which are within the menstrual period, variable or periodic ; the third, those connected with the function of reproduction.

The connection between the symptoms present in cases of retroflexion and the displacement itself has given rise to much discussion and difference of opinion ; and here we must emphasise what was said on page 374 that the symptoms are not due to the lesion immediately but to other pathological changes consequent on or associated with it. Herman¹ would refer the symptoms in displacements entirely “to weakness and over-stretching of the muscular and ligamentous tissues which support the uterus,” but we cannot thus ignore chronic metritis and endometritis and the disturbances of menstruation and reproduction. On the other hand, in judging of the symptoms of retroflexion we must keep before us Vedeler's² statistics, who found in 40 p. c. of cases of retroflexion no symptoms, and concludes that every degree of retroflexion may exist either with or without symptoms.

Weakness in the back is the most common complaint. It may amount to actual pain, which is aggravated on muscular exertion and generally at the menstrual periods. The symptoms of *chronic pelvic peritonitis* are usually present ; the feeling of weight and discomfort in the pelvis is sometimes due to the stretching of old adhesions. The importance of pelvic inflammation, fixing the uterus in its abnormal position and preventing its replacement, we shall consider under treatment. *Painful defæcation* with tenesmus is explained by the relation of the loaded rectum to the retroflexed uterus ; irritation from pressure of the fundus

¹ The Pathological Relationship of Uterine Displacements : Brit. Med. Jour., 1888, I., p. 1213.

² Retroflexio Uteri : Archiv f. Gyn., Bd. XXVIII., S. 228.

against the wall of the rectum may produce straining efforts, but this is very rare.

The *leucorrhœa* is due to chronic inflammation of the mucous membrane. As the result of the displacement, there is passive congestion of all the tissues of the uterus; this leads in the first instance to a simple hypersecretion of mucus, which gradually passes into chronic inflammation. The mucous secretion is more marked immediately after the increased congestion of the menstrual period; but, gradually, it spreads itself over the intermenstrual period. *Dysmenorrhœa* is not so frequent a symptom here as in ante flexion; the explanation is, on the mechanical theory, that retroflexion usually occurs in multiparæ where the cervical canal is patulous. *Menorrhagia* forms one of the more prominent symptoms; it is due partly to the chronic inflammation of the mucous membrane, partly to obstruction to the return of the blood from the uterus.

Affection
of Repro-
ductive
System.

The *reproductive function* is variously and seriously affected. This is brought under our notice when retroflexion occurs in one who has already been pregnant, and presents an obstacle to conception or at least to the growth of a fertilised ovum in the uterus. Sometimes a patient tells us that she had a child several years ago; that she has suffered from pain in the back, leucorrhœa, and irregular menstruation since that time and has never conceived again. With this history, we may find retroflexion of the uterus although often it is the tubes that are at fault.

Abortion
in Retro-
flexion.

The *sterility* may, of course, be due to a variety of causes—the altered position of the cervix, the increased mucous secretion, obstruction of the Fallopian tubes, malposition of the ovaries. We cannot, therefore be sure of curing the sterility by replacing the uterus, although we frequently find that the patient *does* conceive shortly after this treatment. After conception has taken place, there is the further risk of *abortion*; with a history of repeated abortion, we sometimes find retroflexion. Conception probably often takes place in a retroflexed uterus, which afterwards may right itself so that pregnancy goes on to the full time. Abortion is due to the inability of the uterus thus to right itself, or to the pathological condition of the mucous membrane which prevents the ovum from becoming securely attached. When abortion does not occur and the pregnant uterus does not straighten itself so as to grow upwards into the abdomen, it enlarges without the undoing of the flexion; in this case it will expand more and more into the hollow of the sacrum and become wedged below the promontory. This constitutes Retroflexion of the Gravid Uterus.

DIAGNOSIS.

On *vaginal* examination the cervix is felt low down in the pelvis, the cause of which has been explained under Pathology. The os looks

directly downwards. A firm round body is felt in the posterior fornix, continuous with the cervix uteri but separated from it by a groove more or less distinctly marked according to the amount of flexion. Place the forefinger on the cervix, and the middle finger on this body; on moving the former, the latter moves with it.

But a fibroid tumour of the posterior wall would produce similar conditions; therefore make the *bimanual* examination. First place the vaginal fingers in the anterior fornix and make pressure with the external hand until the fingers of both hands meet; there is nothing between them except the abdominal and vaginal walls, the fundus is therefore not to the front. Now put the vaginal fingers into the groove behind




FIG. 206.

DIAGNOSIS OF RETROFLEXION BY BIMANUAL EXAMINATION.

the cervix, or, better still, lay hold of the cervix with the index finger in front of it and the middle finger in the groove behind (see fig. 206), and lift up the uterus as high in the pelvis as possible; make pressure with the external hand until the cervix lies fairly between the hands; the upper surface of the uterus is felt to curve backwards. In a favourable case (with lax abdominal walls) we can do the bimanual examination on a still deeper plane, and get both hands to meet behind or at least fairly embrace the retroflexed fundus. Having ascertained that the fundus uteri is retroflexed, we ask ourselves whether it be fixed or movable—*whether it can be replaced or not.*

Rectal
examina-
tion in
Retro-
flexion.

The *rectal* examination has this advantage, that the finger passes upwards over the free surface of the fundus without displacing it. It is indispensable in cases where the rigidity of the abdominal walls prevents our getting the uterus between the hands in the bimanual. The drawing down of the uterus with the volsella is an additional help in such cases, as it enables the finger in the rectum to reach the fundus.

Uterine
Sound in
Retro-
flexion.

The *sound* confirms the diagnosis in doubtful cases, and tells us further whether the retroflexed uterus is enlarged. Before using the sound, we must palpate the uterus carefully to ascertain that it is not becoming enlarged with a growing ovum, and inquire as to the patient's menstruation. We curve the sound to correspond with the degree of flexion ascertained on bimanual examination. If introduced with the concavity directed backwards, it passes into the uterine cavity without our having to make the rotation (*v. fig. 86*); through the posterior fornix, we feel the end of it in the retroflexed fundus; it usually passes in beyond the two and a half inches. We can also learn from the sound whether the uterus can be replaced or not; but it is better to get the information from the bimanual examination. The sound is of most use in differential diagnosis.

Differ-
ential
Diagnosis
of Retro-
flexion,

Differential diagnosis. The following are the conditions arranged in the order of frequency, which might be mistaken for retroflexion:—

Fæces in the rectum;

Pelvic deposit in the pouch of Douglas { Peritonitis,
Hæmatocele,
Carcinoma;

Cellulitis behind the cervix;

Myoma of the posterior wall;

Prolapsed ovary or small ovarian tumour or dilated tube.

from load-
ed Rectum,

Fæcal matter in the rectum gives rise to difficulty only on superficial examination. We should always decline to give an opinion as to the condition of the pelvic organs when the rectum is loaded. If this be attended to, no mistake in diagnosis will be made under this head.

from
Pelvic
Deposit,

Pelvic deposit in the pouch of Douglas gives rise to more difficulty, because it may closely simulate the condition found in retroflexion—"a body felt through the posterior fornix and moving along with the cervix." Such a deposit will be proved not to be the fundus uteri by our finding the latter in another position. If inflammation is present, it is difficult to make the examination necessary to ascertain this; we may not be justified in using the sound just where it would give us the desired information: such cases present great difficulty in diagnosis,

and the true condition can only be ascertained on repeated examination or after the inflammation has subsided.

Cellulitis behind the cervix is rarely present in such a form as to give rise to a mistake in diagnosis, unless the inflammation renders the necessary examination difficult. from
Cellulitis,

A *myoma* projecting posteriorly from the lower segment of the uterus resembles, in form and firmness, the retroflexed fundus. On bimanual examination, however, we find that we have between the hands a larger body than the uterus alone. The fundus may also be felt to the front, and distinct from the tumour. To ascertain its position, it is best to make the bimanual examination with the sound in the cavity of the uterus. Fig. 199 shows the information given by the sound, if we suppose that the structure to the left of the figure is the rectum. A fibroid tumour accompanied by inflammation presents great difficulty. from
Myoma,

If the *ovary* be *prolapsed*, enlarged through inflammation, and adherent to the posterior aspect of the uterus, it simulates (on vaginal examination) the retroflexed fundus. So also does a small *ovarian tumour* or *dilated tube* lying in the pouch of Douglas, though these are softer and more elastic than the uterus. The bimanual examination, supplemented if necessary by the use of the sound and the drawing down of the uterus with the volsella, enables us to ascertain the exact position of the fundus and its relation to the tumour. from
Prolapsed
Ovary or
Tube.

PROGNOSIS.

The prognosis depends upon the mobility of the uterus, and the possibility of replacing it. It is always less favourable where inflammation is present; though we have seen considerable exudations become after a time absorbed, and the uterus again movable so that it could be replaced. As regards the probability of future conception, our statements should be guarded; though the probabilities are increased if we can replace the uterus.

Whether a permanent cure of the displacement (so that the uterus will keep its normal position after the instrument is removed) is often effected, we have not much definite information. *A priori*, we should not expect that the stretched utero-sacral ligaments would readily become shortened again unless a pregnancy supervene. The curability of the retroflexion depends, according to Mundé, on the *recency of the displacement*; "recent displacements of any variety are the only cases which offer a fair chance of complete recovery by any of the mechanical means at our disposal." The length of time during which a pessary must be worn so as to effect a cure of recent puerperal retroflexion is, according to Mundé, six months to a year. Possibility
of cure of
Retro-
flexion.

TREATMENT.

Here we have to consider :—

- (1.) The treatment of associated inflammatory conditions ;
- (2.) The reposition of the displaced uterus and its retention by means of pessaries ;
- (3.) The operative treatment of these displacements.

1. *The treatment of associated inflammatory conditions.*—Often there is some inflammatory condition of the appendages which may or may not be the result of the displacement, but which is usually the starting-point of the perimetritis, often also present. Further, there is the chronic metritis which is the important factor in the clinical picture of a retroflexion with characteristic symptoms. The displacement *per se* is usually symptomless ; the rôle it plays is to produce a vulnerability of the tissues to inflammatory conditions.

The treatment of these various inflammatory conditions has been already discussed (*v.* chapters on peritonitis, salpingitis, ovaritis, and chronic metritis). It is referred to again here to emphasise the fact that displacements are often of minor importance. Further, so long as active inflammation is present, it is a contra-indication to the reposition and retention by pessaries which has now to be considered.

2. *The reposition of the displaced uterus and its retention by means of pessaries.*—This is only applicable to cases of movable retroflexion, without marked tenderness beside the uterus. Congenital cases should not be thus treated, and it is an open question whether, if retroflexion be found in an unmarried patient local treatment by pessaries is desirable. Further, if the uterus is fixed by adhesions, reposition is impossible unless these are broken down.

Schultze's method of doing this by manipulation under chloroform has not found much favour in this country. He aims not so much at forcible reposition, as the loosening of adhesions through careful bimanual stretching.

Method.—Bladder and rectum are empty ; dorsal posture, thighs flexed and abducted. Irrigate the rectum with warm water. With the index and middle fingers in rectum and the external hand grasping the fundus, lift the uterus carefully up. Slight adhesions yield to pressure of fingers ; broader ones are stretched by the ends of the fingers, although repeated attempts may be necessary. A pessary introduced after reposition.

He also attempts to replace adherent prolapsed ovaries in the same way.

To replace the retroflexed uterus, there are three methods :—

- (1.) By bimanual manipulation ;
- (2.) With the sound ;
- (3.) By genupectoral posture, combined with traction by volsellæ.

(1.) *Bimanual manipulation* is the safest method, but from its discomfort to the patient it is not much used. It may be done with the fingers in the vagina as in fig. 206, or better still with one in the vagina

and the other in the rectum as in fig. 207. Pressure is not made with the upper hand until the uterus has been pushed round to the front.

(2.) The method of *replacement with the sound* will be evident from fig. 208. After passing the sound, as in fig. 208, it is not rotated on its long axis, but swept round in a curve to the position of 2: the rough surface of the handle which looks backward in 1, looks forward in 2. It is then moved directly backwards to 3, the uterus being thus replaced, the sound is withdrawn and a pessary introduced.

Various forms of uterine repositors have been devised by Sims and others. They resemble a sound with the intra-uterine portion jointed to the stem, on which it is moved by a suitable mechanism. They are of no practical value.




FIG. 207.

REPOSITION OF THE RETROFLEXED UTERUS BY THE FINGER IN THE RECTUM.

(3.) The importance of the genupectoral posture in replacing the retroflexed uterus has been brought forward by H. F. Campbell. On placing the patient in this posture, the abdominal contents gravitate downwards and forwards; this displacement withdraws the internal pressure from the pelvic floor, so as to subject it to the atmospheric pressure from without. If the vaginal orifice be now opened, the vaginal cavity becomes distended with air; if the walls are lax, the cavity may be so large that the finger reaches the cervix with difficulty. The position of the uterus changes;¹ but the retroflexed uterus

The Retro-
flexed
Uterus
in Genu-
pectoral
Posture.

¹ For full account of changes produced by the genupectoral posture, the student should consult the Atlas of the "Relations of the Abdominal and Pelvic Organs in the Female:" *Simpson and Hart*, 1881.

does not become replaced, as Campbell supposed. It moves as a whole near the sacrum; and, if already retroverted, it becomes still more so. To effect replacement, we must either push the fundus forwards or draw the cervix backwards. It is best to combine these actions; having laid hold of the cervix with the volsella per vaginam, we draw it downwards while with the index finger of the right hand, per rectum, we press the fundus towards the bladder (see fig. 209). This method of reposition is only used in cases of retroflexion of the gravid uterus.




FIG. 208.

REPLACEMENT OF THE UTERUS WITH THE SOUND. 1, 2, 3, the successive positions of the SOUND and of the UTERUS.

Having replaced the uterus by one of those methods, we have to retain it in its normal position.

The retention of the uterus in its normal position is effected by *vaginal pessaries*. Of these the best forms are the Hodge or, its modification, the Albert Smith.

Material of Pessaries. The *material* of which they are made is vulcanite, which is light and smooth and not affected by vaginal discharges. To bend the vulcanite,

the pessary should be placed in hot, almost boiling, water. It is thus made pliable and can be moulded to the desired form, but becomes firm again on placing it in cold water; this is also effected by oiling the pessary and heating it in a spirit lamp. Pessaries are also made of gutta-percha, which has the advantage of being easily moulded; these cannot, however, be worn for a long time, as the gutta-percha is absorbent and, retaining the secretions, sets up irritation. The patient can wear one for a few weeks till we see that it fits comfortably and is effective, and then we can substitute one of a similar form made of




FIG. 209.

REPLACEMENT OF THE UTERUS WITH THE VOLSELLA AND THE FINGER IN THE RECTUM; the patient is in the genupectoral position.

vulcanite. Celluloid pessaries are now sometimes used instead of vulcanite ones.

The *form* of the Hodge is an elongated horse-shoe, with a straight transverse bar joining the free ends. Seen from the front (fig. 210), it has a curved upper end which is adapted to the posterior fornix; the lower end consists of a straight bar which serves to keep the sides apart, and lies under cover of the symphysis pubis; the external angles of this

The Hodge
Pessary.

The Albert
Smith
Pessary.

end are rounded to prevent their cutting the vagina; the sides run almost parallel. Seen from the side (fig. 212), it is a mould of the vaginal slit; there is an upper sacral curve, which is long and well-marked; there is a lower pubic one, which is not necessarily present or is only slightly marked. The pessary lies so that the concavity of the sacral curve looks forward, that is to say, the upper end of the pessary (like the posterior fornix vaginae) curves forwards. The Albert Smith (fig. 211) contracts in its lower half to a more or less beak-shaped end; seen from the side, it has the pubic curve more marked (fig. 212). Scientifically it is the more correct form, because the posterior wall of the vagina is narrower below than it is above. The lower end should not be too much contracted, otherwise it is apt to interfere with married life; also when the vaginal orifice is wide, it favours the expulsion of the instrument. A second modification of the Hodge is




FIG. 210.

HODGE PESSARY.



FIG. 211.

ALBERT SMITH PESSARY.



FIG. 212.

SIDE VIEW OF ALBERT SMITH PESSARY. The Hodge is similar, but has the lower curve less marked.

recommended by Thomas, in which the upper bar is thicker, the sacral curve more pronounced, and the whole instrument longer.

Choice of
Hodge
Pessary.

The *choice of an instrument* suitable to the case must be made. The pessary should be narrower and shorter than the posterior vaginal wall, so that it produces no tension when it is in position. The upper bar should be of such a size that it can be passed in easily; the lower should be narrower than the upper, but not too narrow for the reasons given above. The proof of a good fitting instrument is that the patient does not feel its presence, nor should it interfere with married life.

Mode of
Introduc-
tion of
Hodge
Pessary.

The *mode of introduction* of the pessary demands special attention. It is important that this apparently simple manœuvre be effected

without causing pain to the patient. From the fact that the vulvar orifice is antero-posterior while the cavity of the vagina is transverse, the instrument must be introduced with its plane surface horizontal (the patient is supposed to be on the side) and afterwards rotated so that this comes to be vertical. From the position of the cervix, the instrument is very liable to run into the anterior fornix. When in position the upper end must curve forwards. Having oiled the instrument, grasp it with the lower end (the square end in the case of the Hodge, the narrower end in the case of the Albert Smith) between the finger




FIG. 213.

INTRODUCTION OF PESSARY, FIRST STAGE.

and thumb of the right hand. Separate the labia with the first and second fingers of the left hand; when the vaginal orifice is narrow, hook back the fourchette with one finger or get the posterior corner of the end which is being introduced within the vaginal orifice; and press back the perineum with it so that the anterior corner is not pushed against the clitoris or vestibule. Now push the pessary backwards in the axis of the vagina till it is half within the cavity (see fig. 213), and rotate it so that the concavity of the sacral curve looks forwards.

Pass the index finger behind the instrument into the vagina, and place the tip of it against the upper bar; carry the pessary onwards, keeping the upper bar well against the posterior vaginal wall to prevent its slipping up in front of the cervix (fig. 214).

How the
Hodge
Pessary
lies when
in situ.

The *position* and *action* of the pessary when *in situ* are as follows. It lies exactly adapted to the vaginal walls (fig. 216); the upper end being in the posterior fornix behind the cervix, the lower just within the vaginal orifice. It is kept in position through its resting on the oblique anterior face of the sacral segment of the pelvic floor, against which it is compressed by the posterior face of the pubic segment.




FIG. 214.

SECOND STAGE: PESSARY CARRIED ON BY FINGER.

The student will readily understand and remember the position of the pessary in the following way. Hold the hand inclined as in fig. 215, with the palm slightly inflexed. It resembles the posterior vaginal wall in the following points:—(1) It is broader above than below; (2) it curves forwards above; (3) from its obliquity, it allows the pessary to sit on it. Now place the pessary on it. It will only lie adapted to the hand when the broad end is above and the upper curve is directed forwards.

Action of
the Hodge
Pessary.

The Hodge pessary does not act as a lever; that is to say, the intra-abdominal pressure does not act specially on the lower bar and depress it, causing the superior one to rise. The intra-abdominal pressure acts

nearly equally on both bars, of which fact the student may satisfy himself clinically. Its action is that the *upper bar gives a point d'appui to the posterior fornix*. The posterior vaginal wall runs round the upper bar as on a pulley, and, as it is inserted into the cervix, the latter is thereby drawn upwards and the fundus thrown forwards (fig. 216). The pessary, therefore, has the same action as the utero-sacral ligaments, if we suppose that these keep the cervix backwards. This is only the action in the case of a retroverted uterus which has been replaced. A vaginal pessary, however, gives relief even though we may not be able to replace the uterus. In this case we may suppose that it acts by supporting the uterus as a whole, thus diminishing tension on the ligaments and passive congestion.¹




FIG. 215.

HAND HOLDING ALBERT SMITH PESSARY.

The after watching of the case is important. The patient should be instructed to return in two days to see that the instrument is in place, and to return at once if it causes pain. After this she should report herself occasionally, say at intervals of a month, when examination is made to ascertain that the uterus keeps its place. If she uses hot-water injections occasionally, it is not necessary to remove the instrument to clean it more frequently than this. After the pessary has been worn for some months, it may be removed to see if the uterus remains in position without it. Sometimes we find that the uterus falls back again into its abnormal position as soon as the instrument is withdrawn; in such a

[¹ See Granville Bantock on *The Use and Abuse of Pessaries*, London, 1884; Hart on *The Structural Anatomy of the Female Pelvic Floor*.

case, it must be introduced again and may have to be worn for years. Should conception occur, the pessary may be worn till the fourth month, after which the uterus rises above the brim and there is no longer reason to fear displacement.

Schultze's
Pessary.

In Germany, Schultze's pessary (fig. 217) is the one in general use. It has the form of a figure of eight, the upper ring embracing the cervix. It is interesting to note that it also goes on the principle that the pessary acts on the cervix, not the body of the uterus.




FIG. 216.

POSITION AND ACTION OF PESSARY.

In some cases the uterine tissue is flaccid at the angle of flexion, and the body falls to the back or front as if it were jointed to the cervix. Here the Hodge, which acts on the body through the cervix, does no good; the intra-uterine stem, along with a Hodge which has transverse bars, is suitable for some of these cases. Wynn Williams, Meadows (fig. 218), and Routh have devised various forms of pessary on this

principle. The general opinion of gynecologists in this country is against intra-uterine stem pessaries.

From what has been said on the action of the Hodge pessary, it is evident that in the treatment of Retroversion + Retroflexion *the version alone is affected by the pessary.* Whether the flexion is remedied will depend on the state of the uterine walls and the effect of intra-abdominal pressure upon them. Hodge's Pessary good only in Retroversion.

For illustrative examples showing the value of pessaries in suitable cases, the student may consult Bantock's monograph, or Macans' translation of Schultze.

3. *The Operative Treatment of Retroversion and Retroflexion.*—There is probably no subject in operative gynecology on which so much has been written during the last few years as the above, unless it be the extirpation of the uterus for fibroids or malignant disease. Operations for Backward Displacements.




FIG. 217.
SCHULTZE'S PESSARY.




FIG. 218.
MEADOWS' COMPOUND STEM PESSARY.

While the latter operation is of the first importance, as it implies the saving of life, the value of the former is more difficult to estimate as it concerns the relief of suffering into which the personal equation largely enters. The causal connection between the displacement and the symptoms is in many cases open to question (see p. 374). That benefit follows an operation does not prove that the displacement was the cause, as it might be, accounted for in another way, *e.g.*, by the psychical effect of the operation. It is just in operations done to relieve reflex pains that there is most room for this effect.

Such prominence has, however, been given to these operations that we must describe them, and leave their validity to the test of time. Three modes of fixing the replaced uterus have been tried, which are best described as vaginal fixation, ventral fixation, and shortening of the round ligaments.

Vaginal
Fixation.

1. *Vaginal fixation*¹ includes all operations which aim at fixing the cervix or body of the uterus to the vagina. This has been tried in various ways. By excising a portion of the anterior wall of the cervix von Rabenau² hoped, from the consequent contraction, to effect a permanent ante flexion. Conversely by amputation of the posterior lip of the cervix and stitching the stump to the posterior vaginal wall, Richelot³ sought to produce anteversion. To excite adhesions in the utero-vesical pouch of the peritoneum, and thus tack the uterus forward to the bladder, Schücking⁴ passed a suture through the anterior wall of the uterus, the utero-vesical peritoneum, and the anterior fornix. The ligature was introduced with a curved needle from the uterine cavity which required to be previously dilated. The risk of injuring the bladder or intestine in this operation led Zweifel⁵ to modify it by opening into the utero-vesical pouch, and Sänger⁶ to stitch the fundus directly to the anterior fornix.

Further, the anterior surface of the uterus has been stitched to the anterior vaginal wall with and without opening into the utero-vesical pouch. This mode of procedure has been elaborated by Mackenrodt and Dührssen, and requires fuller consideration, as it is to their operations or a modification of them, that the term *vaginal fixation* is commonly applied.

Macken-
rodt's
Operation.

Mackenrodt's method as first described by him, at the Berlin Obstetrical Society in May 1892, was as follows.⁷ After drawing down the cervix and defining the bladder he made a transverse incision in the anterior fornix, from the centre of which a longitudinal one was carried to near the urethral orifice. The triangular flaps were dissected off, the bladder separated from the uterus, and sutures passed stitching the uterus to the vaginal flaps. He has since modified the technique on the ground that failure was due to the persistence of the utero-vesical pouch. First he tried to obliterate it by passing deep cat-gut sutures; and subsequently by opening into the pouch so as to stitch the anterior surface of the uterus to the vaginal incision. Latterly, on account of the bad effects on pregnancy and labour from the firmness of the fixation thus produced, he has stitched the uterus not to the vagina, but to the peritoneum over the bladder.⁸

Dührssen's
Operation.

Dührssen⁹ makes a transverse incision in the anterior fornix, separates the bladder from the cervix, cuts into the utero-vesical pouch, and exposes as large an area as possible of the anterior surface of the uterus, passes a temporary suture (or if one is not sufficient others higher and higher up, as more and more of the anterior surface of the uterus comes into view) through the uterus (Plate X., fig. 1, *a,a,a*) with which to draw it down.

¹ The term vagino-fixation though shorter is incorrect. Called also vaginal-hysteropexy, colpo-hysteropexy.

² Ueber eine neue operative Behandlung der Retroflexio Uteri: Cent. f. Gyn., 1886, S. 429. Six cases are recorded, but the ultimate result is not given. Schmitt proposed a similar operation. Cent. f. Gyn., 1888, S. 685.

³ De l'hystéropexie vaginale: Union Medicale. Dec. 1889.

⁴ Eine neue Methode der radicale Heilung der Retroflexio Uteri: Cent. f. Gyn., 1888, S. 181 and 682. Also *ibid.*, 1890, S. 123, and 1891, S. 249. He has operated on 217 cases, 88 of which have been followed, with failure only in 4. The suture is not removed for six weeks, and a pessary is worn for 12. 23 of his cases were subsequently delivered at term.


⁵ Ueber die Vaginalfixatio Uteri: Cent. f. Gyn., 1890, S. 689.

⁶ Cent. f. Gyn., 1888, S. 17 and 34.

⁷ Die Therapie der Retroflexio Uteri: Cent. f. Gyn., 1892, S. 479.

⁸ Ueber die Unzulässigkeit der Vagino-fixation und ihren notwendigen Ersatz durch die Vesico-fixation. Zeitschr. f. Geb. u. Gyn., 1895, Bd. XXXIII., S. 514.

⁹ Ueber Vagino-fixatio Uteri: Zeitschr. f. Geb. u. Gyn., 1892, Bd. XXIV., S. 368, and Cent. f. Gyn., 1893, S. 681.






Fig. 1.



Fig. 2.




Fig. 3.




Fig. 4.




Fig. 5.




Fig. 6.

OPERATIVE TREATMENT OF BACKWARD DISPLACEMENTS.

- FIG. 1. Vaginal fixation ; FIG. 2. Result of same (Dührssen).
 FIG. 3. Indirect ventral fixation ; FIG. 4. Position of uterine sutures (Olshausen).
 FIG. 5. Direct ventral fixation (Leopold).
 FIG. 6. Dissection for Alexander-Adams operation.

Steady traction on this, aided by volsellæ grasping the cervix, is of the first importance. Three sutures are now passed through the raw surface of the anterior margin of the vaginal incision, and then through the uterus near the fundus (Pl. X., fig. 1, *b, b, b*). These are tied and buried. The traction sutures are now withdrawn, and the vaginal wound closed. The result is shown at Pl. X., fig. 2.

The effect of vaginal fixation on subsequent pregnancy and labour must be reckoned with in estimating the value of the operation. If the adhesions yield, the retroversion will return in the puerperium; while if they do not yield, they become a source of danger during pregnancy and labour. They are apparently firmer when the utero-vesical pouch has been opened into, as in Dührssen's operation, than when this is not done. Hence abortion¹ is frequent. Further, the fixation of the anterior wall causes expansion of the uterus to take place at the expense of the posterior, with consequent upward displacement of the cervix. Two cases of obstructed labour (with one death) have been recorded by Strassmann,² one by Graefe,³ in which Cæsarean section was required, another by Wertheim,⁴ and two more recently, by Rühl.⁵ Negri⁶ finds that seventy-seven cases of vaginal fixation with subsequent pregnancy terminated as follows: forty-nine in normal labour, nine required operative interference, and nineteen in abortion.

Effect on
Pregnancy
and
Labour.

2. Under *ventral fixation*⁷ we include various operations which have the object of causing union by adhesion between the uterus and the anterior abdominal wall.

Ventral
Fixation.

To avoid opening into the peritoneal cavity, the uterus has been pushed up from below against the abdominal wall, and transfixed there by a suture passed through the latter. The risk of transfixing intestine at the same time is obvious, and this blind method of procedure has not found many advocates.

Methods.

The usual method, however, is to open the abdomen and stitch the uterus, either indirectly by means of its broad ligaments (Pl. X., figs. 3 and 4), or directly by sutures passed through its substance (Pl. X., fig. 5) to the abdominal incision.

In the direct method the uterus may be stitched to the abdominal wall by catgut sutures, uniting the superficial tissue of the margins of the uterus to the peritoneum and subjacent tissue of the abdominal wall; the abdominal wound being subsequently closed in the usual manner. Or the sutures used to close the wound may be passed likewise through the anterior surface of the uterus. To cut off infection from the skin, the sutures which pass through the uterine wall may stop short of the skin, and thus be buried when the abdominal incision

¹ Dührssen reports six abortions in twenty-four pregnancies—(Strassmann: Zeits. f. Geb. u. Gyn., Bd. XXXIII., S. 510, and Archiv f. Gyn., 1896); Weberstedt also records six abortions out of twenty-four pregnancies from Guesserow's Clinique—(Beobachtungen über die Beziehungen gynäkologischer Erkrankungen zur Schwangerschaft: Dissert. inaug. Berlin, 1895.)

² Loc. cit.

³ Cent. f. Gyn., 1895, S. 470.

⁴ Monats. f. Geb. u. Gyn., 1895, Bd. II., S. 473.

⁵ Cent. f. Gyn., 1896, S. 147.

⁶ Annali di Ostetricia e Gynecologia: Agosto, 1896.

⁷ Called also ventro-fixation, ventri-fixation, gastro-hysteropexy.

is closed. Further, instead of passing straight through the anterior surface of the uterus, they may pass out and in upon its peritoneal aspect, so as to excite adhesions over the whole area of the uterus, and keep it in apposition with the abdominal wall. To promote adhesion, the peritoneal surface of the uterus may be scarified.

We consider now these methods of *ventral fixation* more in detail.

The various modes of suturing the uterus to the wall of the abdomen *without opening the latter*, are worthy only of passing notice. Kelly¹ recommended lifting the uterus upwards by the hand in the vagina until it could be felt through the abdominal wall just above the pubes, and transfixing the wall and fundus uteri with a ligature which was left in for a fortnight. So also Shober² has pushed up the uterus with a canula in the uterine cavity through which a needle was passed, and the uterus stitched to the abdominal wall. An abdominal incision through the superficial fascia was made as a preliminary in the bottom of which the uterine ligature was tied so as to bury it. Further, Foerster³ incised the pouch of Douglas, brought the uterus out, scarified and tranfixed the fundus by two silk threads which were subsequently carried by a canula and needle through the abdominal cavity and abdominal wall.

These methods of tacking the uterus to the abdominal wall in the dark hardly merit the name of operations, but the stitching of the uterus indirectly by means of its broad ligaments, or directly to the abdominal wall, deserves fuller consideration. It was a natural enough experiment when the appendages were removed in cases of retroversion, to stitch the pedicles into the abdominal incision⁴ as was first done by Koeberlé (1869), and Sims (1875).

Ventral Fixation.

The stitching of the broad ligament to the abdominal wall, apart from amputation of the appendages, as a treatment for retroflexion, was first devised by Olshausen,⁵ and constitutes the *indirect* method.

Indirect Method.

He united with silk the round and broad ligaments at the upper angle of the uterus (Pl. X., fig. 4) to the abdominal wall.

It will be noted that in this operation the three sutures (l m n) which unite the angle of the uterus to the parietal peritoneum, are distinct from those which close the abdominal incision, and are of course not removed, but left to be absorbed (Pl. X., fig. 3). Care is required in introducing them not to pierce the Fallopian tube (v. Pl. X., fig. 4) or hypogastric artery in the abdominal wall. Instead of suturing the broad ligament by its anterior aspect, Kelly⁶ has stitched the ovarian ligament to the abdominal wall, which he thinks produces more complete anteversion of the uterus.

Direct Method.

Fixation of the uterus *directly* to the abdominal wall is done by Leopold in the following way.⁷

He passes three of the sutures which close the abdominal incision, also through the anterior surface of the uterus (Pl. X., fig. 5). These dip into its superficial tissue for

¹ Amer. Journ. Obstet., 1889, p. 1066.

² Amer. Journ. Obstet., 1895, p. 843.

³ The Surgical Treatment of Retrodisplacement of the Uterus, with special reference to Vaginal and Vento-fixation. Amer. Journ. Obstet., 1895, Vol. I., p. 747.

⁴ In addition to stitching the pedicle in the abdominal incision, Klotz passed a glass tube behind the uterus into the pouch of Douglas, the presence of which he thinks excites adhesions and favours fixation. See Cent. f. Gyn., 1888, S. 11, and 1891, S. 97, where he records thirty-eight cases treated in this manner.

⁵ Ueber ventrale Operationem bei Lageanomalien: Cent. f. Gyn., 1886, S. 667 and 698.

⁶ Amer. Jour. Obst., 1889, p. 1066. Peterson records seventeen cases done by Kelly's method. He used both silk and catgut, and carried the suture deeply into the muscles and fascia of the abdominal wall. "Suspension of the Retro-displaced Uterus by the Utero-ovarian ligaments"—with interesting discussion. Amer. Jour. Obstet., June 1895, Vol. I., p. 832.

⁷ Cent. f. Gyn., 1888, S. 161, and 1890, S. 185. Also Sammlung klinischer Vorträge. No. 333.

about one-third of an inch, and they run opposite to the insertion of the Fallopian tube, of the round ligament, and the third still lower down. The area of the uterus within the sutures is scraped with the back of the bistoury, but not enough to make it bleed. The sutures are removed in a fortnight. A Hodge pessary is worn for a month. Czerny¹ modifies this procedure in that the sutures which fix the uterus do not penetrate as far as the skin surface of the abdomen, and are of catgut. Terrier² makes the uterine sutures pass out and in on the peritoneal surface of the uterus so as to promote peritonitic adhesion to the abdominal wall.

As to the effect of ventral fixation on pregnancy and labour, Mil-
 änder³ has collected seventy-four cases. Of the sixty-three who have
 been confined, fifty-four went to full time, while six aborted, and three
 had premature labour. Eleven of the labours required assistance, two
 being delivered by Cæsarean section. Küstner⁴ gives the result in one
 hundred and twenty-two cases: seventy-four went to full time, while opera-
 tive interference was called for in many cases. It is evident, therefore,
 that though the risks in subsequent pregnancy and labour are not
 so great in ventral as in vaginal fixation, they are still considerable
 enough to demand attention in judging as to the advisability of
 operation.

Effect on
Pregnancy
and
Labour.

3. *The shortening of the round ligaments*, an operation first proposed
 by Alquié,⁵ but commonly called the Alexander-Adams', after the two
 British operators who reintroduced and elaborated it, is usually done by
 cutting down on the ligaments after they emerge from the external
 abdominal ring. Through its not having the risk of abdominal section,
 and thus being a minor rather than a major operation, it has become
 increasingly popular of recent years, especially in America and France.

Alexander-
Adams'
Operation.

Shortening of the round ligaments has also been done by stitching
 them within the abdominal cavity; this implies, of course, abdominal
 section and increases the risk of the operation.

The technique of Alexander-Adams'⁶ operation is, shortly, as follows.

The pubic hair having been shaved and the skin cleansed, the pubic spine is felt for
 on both sides, and an incision (Pl. X., fig. 6) from 1½ to 2 in. long, according to the
 fatness of the patient, made from it outwards parallel with Poupart's ligament. The super-
 ficial fascia is divided until the pillars of the ring are recognised. In cutting through the
 fibres between these some delicate fat protrudes. On hooking this up with an aneurism
 needle (Pl. X., fig. 6) the ligament is seen as a red cord. The inguinal nerve should
 be cut and fascia cleared from it anteriorly. If the ligament be not recognisable at the
 external ring, the canal can be split up to the internal. The ligament when found on
 the one side is caught in forceps, and an antiseptic tampon placed on the wound; the
 ligament on the other side is found in the same way.

The uterus is now replaced by an assistant, with the sound, and the ligaments drawn out

¹ Beitr. zur klin. Chirurgie, 1888, Bd. IV., Hft. 1, S. 179.

² Dumoret—Laparo-hystéropexie, etc. Thèse de Paris, 1889.

³ Ventrifixation des Uterus; Schwangerschaft und Querlage des Kindes: Zeitschr. f. Geb. u. Gyn. Bd. XXXIII., S. 464.

⁴ Volkmann's Sam. klin. Vorträge, No. 171, Dec. 1896.

⁵ Alquié presented to the Académie de Médecine in 1840, a memoir, "Sur une nouvelle méthode pour traiter les divers déplacements de la matrice," which had not been published until it appeared in a recent monograph by Moreau, "Du raccourcissement des ligaments ronds appliqué à la guérison des déplacements de la matrice." Bruxelles, 1894.

⁶ F. Imlach—On shortening the round ligament of the uterus: Edin. Med. Jour., April 1885. W. Alexander—The operation of correcting some uterine displacements by shortening the round ligaments: Brit. Gyn. Jour., 1885-86.

from 2 to 4 inches. Each is then sewn to one or both of the pillars by catgut suture, and the piece of free ligament cut away. The wounds are closed with two or three sutures.

Shortening of the round ligament within the abdomen, which implies abdominal section, has been recommended by Gill Wyllie¹ and Palmer Dudley.² The former scrapes the inner side of each round ligament about its middle, folds it, so as to bring the raw surfaces into apposition, and stitches with three sutures. The latter denudes the ligaments and stitches them in a loop on to the anterior surface of the uterus.

In America the Alexander-Adams' operation has been warmly advocated by Mundé,³ Cleveland,⁴ and Polk.⁵ In England its value was recently discussed before the British Gynecological Society, in a paper on "Ventre-suspension of the Uterus," by Mayo Robson.⁶ In the discussion, while Alexander refers to the large number of successful cases in which he has performed it, J. W. Taylor (Birmingham), William Duncan, and Mayo Robson give the preference to ventral fixation.

As to the frequency with which vaginal and ventral fixation has been done we note that in Berlin Gusserow⁷ has done only thirteen vaginal fixations, limiting the operation to cases that cannot be treated by pessaries; while Olshausen⁸ prefers vaginal fixation, though he has done twenty-three ventral fixations in six years. In Schauta's Clinique at Vienna thirty-seven vaginal fixations have been done—Dührssen's method alone giving permanent results.⁹ Säger,¹⁰ with some two hundred and thirty retroversions annually in his Clinique at Leipzig, has performed only thirty-eight vaginal fixations in four years. In Werth's Clinique at Keil,¹¹ fifty cases of vaginal fixation have been done—twenty-nine by Mackenrodt's and twenty-one by Dührssen's method; while in the Freiburg¹² Clinique, from 1887 to 1895, only twenty-two ventral fixations were done, mostly after the removal of the diseased appendages. In the Berne¹³ Clinique, Müller has done forty-three vaginal fixations, preferring it to ventral fixation. Kelly¹⁴ has done one hundred and seventy-one ventral fixations. Laphorn Smith, in the Montreal Hospital, has done in five years twenty-one Alexander operations, and twenty-eight ventral fixations, and prefers the latter.

As regards the relative merits of these three operative methods of dealing with the retroverted uterus, vaginal fixation is becoming discredited through the accidents in labour which have followed it, and will become restricted to patients in whom pregnancy is not likely to occur, that is, who are near the menopause. Ventral fixation will be limited to those cases in which the abdomen has to be opened for other reasons, it will be performed as an adjunct to removal of the uterine appendages, although it is questionable whether it is called for here, as the retroverted uterus usually shrinks and ceases to cause pain after

¹ Surgical Treatment of Retroversion of the Uterus with adhesions, with a new method of shortening the round ligaments: Amer. Jour. Obstet., 1889, p. 478.

² A New Method of Surgical Treatment for certain forms of Retrodisplacement of the Uterus with adhesions: Amer. Jour. Obstet., 1890, p. 336.

³ Ten Years' Experience with Alexander's Operation: Amer. Jour. Obstet., 1894, Vol. ii., p. 297, where he reports on sixty-five cases.

⁴ Alexander's Operation: Amer. Jour. Obstet., 1895, Vol. ii., p. 124. He records eighty-four cases of Alexander's Operation which he prefers to ventral fixation which he had done successfully before.

⁵ Discussion on Cleveland's Paper.

⁶ Brit. Gyn. Journ., May 1896.

⁷ Berlin Klin. Woch., S. 750.

⁸ Zeitschr. f. Geb. u. Gyn., Bd. xxxii., S. 1.

⁹ Wertheim. Cent. f. Gyn., 1895, S. 470.

¹⁰ Monat. f. Geb. u. Gyn., Bd. iii., S. 2.

¹¹ Groosdeff—Vratch, 1896, Nos. 21, 22, 23.

¹² Breutleht—Dissertation on Ventrofixatio Uteri.

¹³ Vaginal Fixation: Vratch, 1896.

¹⁴ Suspensio Uteri, Annals of Gyn. and Ped.: Boston, Vol. viii., p. 732.

the appendages have been removed (Pozzi). Where, however, abdominal section has been done to set free a retroverted uterus fixed by adhesions, ventral fixation has a recognised place. While from its comparative safety the Alexander-Adams' operation will continue to have a large number of supporters, it is obvious that it is only applicable to those mobile retroversions, without disease of the uterine appendages, which cannot be treated by pessaries.

CHAPTER XXXIV.

INVERSION OF UTERUS.

LITERATURE.

- Atthill*—Inversion of uterus due to fibroid tumour : Dublin Medical Journal, Feb. 1879. *Aveling*—On Inversion of the Uterus : Brit. Med. Journ. 1886, I., p. 475. *Barnes*—Diseases of Women : London, 1878, p. 721. Med. Chir. Trans., 1869. *Crosse*—An Essay, literary and practical, on Inversio Uteri : Trans. Provincial Med. and Sur. Assoc., London, 1845. *Duncan, Matthews*—On the Production of inverted Uterus : Edin. Med. Jour., May 1867. *Emmet*—Principles and Practice of Gynæcology : Churchill, London, 1884, p. 406. *Fritsch*—Die Lageveränderungen der Gebärmutter : Billroth's Handbuch für Frauenkrankheiten : Stuttgart, 1881. *M'Clintock*—Diseases of Women : Dublin, 1863, p. 76. *Macdonald*—Two cases of chronic inversion of the uterus : Edin. Obst. Trans., Vol. vi., p. 170. *Malins*—Chronic Inversion of the Uterus : Allbutt and Playfair's System of Gynecology, London, 1896. *Spiegelberg*—Zu den Inversionen der Gebärmutter : Archiv f. Gyn., B. iv., S. 350, and B. v., S. 118. *Thomas*—Diseases of Women, p. 453 : Philadelphia, 1880. *Werth*—Ueber partielle Inversion des Uterus durch Geschwülste : Archiv f. Gyn., Bd. xxii., S. 65. The essay by Crosse gives the fullest anatomical description of inversion.

PATHOLOGY.

In inversion, the uterus is turned inside out, so as to form a polypoidal projection into the vagina ; its peritoneal surface is converted into a cup-shaped hollow ; its mucous membrane becomes *everted* so as to lie exposed on all sides in the cervix and vagina.

The mechanism by which this condition is brought about is the following.

1. A portion of the muscular wall of the uterus *having lost its tone*, becomes depressed towards the uterine cavity. In the puerperal condition this is usually that portion of the wall to which the placenta has been attached, and the condition has been described by Rokitansky as "paralysis of the placental seat ;" this partial inversion will be frequently found on abdominal palpation in cases of post-partum hæmorrhage (*Fritsch*). In cases of tumour-growth, fatty degeneration (*Scanzoni*) or malignant infiltration (*A. R. Simpson*) weakens the wall of the uterus round the base of the polypoidal growth, and thus produces an analogous condition.

2. *Muscular contractions* of the non-depressed portion of the uterus, combined with *intra-abdominal pressure*, carry the depressed portion further into the uterine cavity, until the fundus reaches the os internum (fig. 226). In the puerperal condition, muscular contractions occur spontaneously, or are produced by the presence of the placenta; in the case of a polypoidal tumour, they are due to the presence of the foreign body. *Traction from below*, such as the pulling away of the placenta or the tension of the pedicle of a polypus which is being extruded, also produces inversion.

3. The fundus of the uterus, by continuation of the same process, dilates the cervical canal and is "born" into the vagina (fig. 223).

In some cases inversion seems to take place from below upwards with a mechanism similar to that of prolapsus uteri, the lower part of the body of the uterus becoming inverted into the cervical canal (*Taylor*).




FIG. 219.

INVERSION OF UTERUS (half-size, *Barnes* from *Crosse's* essay). The fundus lies in the vagina; the cervix is not inverted; the lips are seen as a flattened out swelling below the angle of inversion. The ovaries (seen from behind) are not in the peritoneal cup.

Matthews Duncan, whose paper was a valuable contribution towards establishing the correct theory of inversion, distinguishes between active and passive inversion. The active is that described above; the passive is produced by inertia of the whole uterus, in which the organ is driven down entirely by intra-abdominal pressure or by traction from below—and not by uterine contractions. Varieties
of Inver-
sion.

It is evident that the process may become arrested at any of these stages and persist as a permanent condition. When it has persisted for a few weeks, it constitutes "chronic inversion;" this is found in the following forms. (1.) Inversion of one horn only, is a rare occurrence. Slight inversion of the uterine wall, at the base of a polypoidal has been more frequently observed. (2.) Partial inversion, when the fundus has descended as far as the os internum, is also found as a

chronic condition. (3.) Complete inversion is the condition most frequently met with.

Anatomy
of Inver-
sion.

An exact knowledge of the relation of parts in *complete inversion* is necessary for diagnosis and treatment. This can only be gained by studying the inverted uterus as seen in section (fig. 219). We must study the position of—

The body of the uterus,
The cervix uteri,
The Fallopian tubes and ovaries,
The peritoneum,
The bladder.

The body of the uterus. The inversion extends, in simple uncomplicated cases, as far as the os internum but *no further*. The uterus lies partly in the vagina, partly in the cervical canal. Its neck is embraced by the os externum, which may lie loosely on it (favouring hæmorrhage) or constrict it firmly (favouring gangrene). After involution takes place, it becomes small, rounded, and of firm consistence, closely resembling a pediculated fibroid tumour; and it has been amputated by mistake for such. It has a rounded form, is of a softer consistence and deeper red colour than a pediculated fibroid, and has a smooth and slippery surface which bleeds freely when handled. The softness may be so marked that the uterus moulds itself to the vaginal cavity and, becoming flattened against the posterior vaginal wall, takes on a mushroom-like form (*Freund*).

The mucous membrane of the uterus may undergo all the changes of any tumour with a constricted base and exposed surface. It is usually congested and bleeds easily; it may become ulcerated and even gangrenous, or may be hypertrophied with polypoidal formations; it may lose its single layer of cubicle epithelium and develop a stratified squamous epithelium. The occurrence of these changes has an important bearing on the necessity of replacing the organ.

The cervix uteri. This is rarely¹ displaced in simple uncomplicated inversion; it forms a broad ring embracing the neck of the tumour. Sometimes the inversion is complicated with prolapsus, or, more properly, the vagina also becomes inverted and the inverted uterus caps the inverted vagina (fig. 220). When this occurs, the cervix uteri is also more or less inverted; a part remains just above the os externum, as a depressed ring which also disappears on making traction on the uterus (*Fritsch*).

The *Fallopian tubes and ovaries*, with some coils of small intestine, may (at first) lie within the inverted cup, which is lined with *peritoneum*; afterwards they retract out of it. In long-standing cases, the

¹ Crosse figures one preparation in which the cervix as well as the body of the uterus was inverted although there was no prolapsus.

rim of the peritoneal cup is contracted by the muscular fibre of the cervix so as scarcely to admit a finger (fig. 221). In a case of six months' standing, in which A. R. Simpson performed Thomas' operation before having recourse to amputation, the contracted ring just admitted the finger; an ovary was caught within it.

Adhesions rarely form between the *peritoneal surfaces*; this is an




FIG. 220.

INVERSION OF UTERUS+INVERSION OF VAGINA, occasioned by a small submucous fibroid (M'Clintock). *Sm F*, submucous fibroid; *U* uterus, *V* vagina, *B* bladder.

interesting fact and is of importance in regard to replacement. We might have expected detachment of the peritoneal lining or tearing of it by the sudden dislocation; the previous stretching of it during pregnancy is perhaps the reason why this has not been noticed. Fritsch says that the lifting up of the fornices by the tumour in the vagina, diminishes the strain on the peritoneum.

The *bladder*, from its relation to the cervix (*v. Chap. IV.*), is not altered in position unless there is prolapsus. When the latter occurs, there is cystocele (*v. fig. 220*). We may therefore contrast the two types of inversion as follows.

Inversion of uterus—cervix and bladder normal in position.

Inversion of uterus + prolapsus (*i.e.*, inversion of vagina)—cervix inverted and cystocele.

ETIOLOGY AND FREQUENCY.

Inversion arises under two different conditions:—

1. In the puerperium—puerperal inversion;
2. Secondary to intra-uterine tumours growing from the fundus.




FIG. 221.

INVERSION OF UTERUS (*Crosse*). The inverted uterus (*U*) lying in the vagina (*V*) is cut open to show the peritoneal sac which does not contain the ovaries (*O*); bristles are passed into uterine orifices of tubes. *b* Broad and *r* round ligaments; *T* tube.

Inversion has also occurred independent of the puerperal condition and of tumour growth; this is quite exceptional.

Etiology of
Puerperal
Inversion.

1. *Puerperal inversion*. This is by far the most frequent form; out of 400 cases, 350 occurred in the puerperal uterus (*Crosse*).

Its former frequency was due to improper management of the third stage of labour. When the uterus was flabby and not contracting and the placenta not coming away, the removal of the latter by traction on

the cord drew down the part of the wall to which it was attached and thus inverted the uterus. This accident was favoured by the situation of the placenta over the fundus (*Hennig*). Since the removal of the placenta by compression (which is best done by the Credé method—with the thumbs of both hands well down behind the fundus so that the uterus may be firmly compressed antero-posteriorly) has been adopted, this accident has become rarer.

A dilated condition of the uterus (distention by blood clots) or a flaccid condition of the walls favours inversion.

2. *Inversion secondary to uterine tumours* is much rarer. Of 400 cases, only forty (ten per cent.) arose in this way (*Crosse*). It has been observed with pediculated fibromata (fig. 220),¹ and will be referred to again when we treat of them (*v.* Chap. XXXV.). *Brewis*² has recorded a case of its occurring spontaneously in a uterus from which a polypus had been previously discharged. It is frequent in sarcoma (*v.* Chap. XLIV.), but very rare in carcinoma uteri. *Tait*³ found it with villous epithelioma, and *Barnes* describes a specimen in which both conditions were present, but does not say which was the primary lesion.

Etiology of
Inversion
due to
Tumours.

SYMPTOMS.

The symptoms produced by inversion at the time of its occurrence, concern the obstetrician rather than the gynecologist. There is the feeling of something giving way in the pelvis, accompanied with pain, hæmorrhage, and sometimes collapse. With complete inversion, there is retention of urine; it often occurs, or at least becomes so marked as to attract the patient's notice, when she has made a straining effort. The cases where the patient says that it first came down several days after labour, are to be explained by supposing that partial inversion occurred after labour but only the final stage attracted attention.

If the uterus be not replaced at the time, the case becomes one of chronic inversion. The symptoms of chronic inversion are—

Hæmorrhage,
Pain in the pelvis of a bearing-down character,
Anæmia and weakness.

Hæmorrhage is the most dangerous symptom. The menstruation is always profuse, as may be easily understood from the fact that the mucous membrane is extended in its area and lies exposed in the cervical canal and vagina. There is also inter-menstrual hæmorrhage, which comes on unprovoked or on straining.

¹ Lee records two cases of its occurrence with fibroid tumours.—*Amer. Journ. Obstet.*, 1888, p. 616

² *Edin. Med. Journ.*, July, 1887.

³ *Brit. Med. Journ.*, 1887, I., p. 66.

The *bearing-down pain* in the pelvis resembles that felt in prolapsus uteri. It varies indefinitely in intensity; sometimes it is very acute, rarely is it so slight that the patient becomes reconciled to her discomfort and is able for work.

The *anæmia* and *weakness* may be so marked as to cause suspicion of malignant disease.

DIAGNOSIS.

Diagnosis
of recent
Inversion.

The diagnosis of *recent inversion* is easy. If the placenta has not yet been born, the hands laid on the fundus to expel it by the Credé method find that the rounded fundus is replaced by a cup-shaped hollow. The cervix is sometimes lifted up by the inverted uterus, so as to be "high above the pubes, even near the umbilicus" (*Crosse*). On passing the hand into the vagina to remove the placenta, care is required to




FIG. 222.

UTERINE POLYPUS (*Thomas*). The uterus in its normal position. Sound passes into uterine cavity.




FIG. 223.

INVERSION OF UTERUS (*Thomas*). A cup-shaped depression is in the place of the uterus. Sound arrested at angle of flexion.




FIG. 224.

UTERINE POLYPUS. Adhesions round pedicle obliterate cervical canal.

recognise what is placenta and what is inverted uterus, and not to increase the inversion in detaching the placenta. If the placenta is already expelled, the hand on the abdomen recognises the same condition; while a large soft body, varying in size according to the extent of the inversion, fills the vagina.

Diagnosis
of Chronic
Inversion.

Chronic Inversion. Before the sound and the bimanual came to the gynecologist's aid in diagnosis, it was impossible to diagnose this condition with certainty. Mistakes were committed by the most eminent surgeons, just because they had not the means of examination which we now possess. Even nowadays mistakes occur through the hasty

making of a diagnosis before all the means of examination have been employed. We therefore describe fully the routine examination.

1. Pass the fingers into the vagina; a rounded and firm or flattened and soft tumour, which bleeds easily, is felt in the vaginal cavity. Sweep the fingers round it, and recognise that it is free on all sides except at its upper extremity. Round this extremity is felt the cervix, the lips and fornices being recognised; or the cervix is thinned out to a ring and the fornices obliterated. If the cervical canal be obliterated by adhesions, the finger will not pass farther up; if it be patulous, it will pass for one-and-a-half to two inches and find that the cervical mucous membrane is reflected equally all round on to the neck of the tumour.

2. With one finger in front of the tumour and the other behind it, lift it up towards the abdominal wall which is depressed with the external hand till the fingers in the vagina are in contact with it. The external hand feels, in the place of the fundus uteri, a truncated body with a depression in the centre (see fig. 223).

3. Now pass one finger into the rectum, which comes first on the body in the vagina: on dragging the latter downwards, the finger in the rectum reaches the upper border of the body and can feel that it ends abruptly, and can pass into the cup-shaped end. Now depress the abdominal walls till they reach the finger in the rectum, or pass a sound into the bladder and direct the point of it backwards till it can be touched by the rectal finger.

4. The sound may be used to probe round the neck of the body where there is not space for the finger to pass upwards. It is most useful, however, in differential diagnosis.

DIFFERENTIAL DIAGNOSIS. Inversion must be differentiated from the following conditions:—

1. Polypus in the vagina, simple or with adherent pedicle;
2. Intra-uterine polypus;
3. Uterine polypus with partial inversion;
4. Prolapsus uteri;
5. Inversion and prolapsus.

Differ-
ential
Diagnosis
of Inver-
sion from
Polypus.

1. In a uterine polypus which lies in the vagina, the fundus will be found to lie somewhere else than in the vagina; it may be retroverted and thus escape recognition in the bimanual; the rectal examination will then discover it. Having found what we suppose to be the fundus, pass the sound along the side of the pedicle; if it is in the uterus, the sound passes more than $2\frac{1}{2}$ inches; if it passes $2\frac{1}{2}$ inches or less, suspect that partial inversion complicates the polypus.

When there are adhesions round the pedicle obliterating the cervical canal, a careful bimanual will reveal the fundus in its normal position

and justify us in breaking down the adhesions with the sound so as to effect a passage into the uterine canal (fig. 224).

2. In a uterine polypus which is still *intra-uterine* the differential diagnosis is more difficult. A case has been recorded in which inversion of one horn of the uterus was diagnosed and amputated as a polypus. A careful examination *per rectum* under chloroform might detect the cup-shaped depression found in partial inversion; the uterine cavity is always enlarged when a polypus is present (fig. 225 and fig. 226).

3. Having satisfied ourselves that there is a polypus, the possibility of there being *partial inversion* of the uterus at its attachment must be kept in view (fig. 227). A careful rectal examination might reveal a depression on the peritoneal aspect of the uterus. The greater sensitive-




FIG. 225.

INTRA-UTERINE POLYPUS
(Thomas).




FIG. 226.

PARTIAL INVERSION (Thomas).




FIG. 227.

UTERINE POLYPUS + PARTIAL
INVERSION (Thomas).

ness of the uterine mucous membrane also helps us; thus if we apply the *écraseur* without chloroform—which is not necessary—to remove the polypus, and the patient has great pain on our tightening up the wire, we may suspect that the loop has embraced the wall of the uterus.¹

4. *Uncomplicated prolapsus uteri* would only on a very superficial examination be mistaken for inversion. The obliteration of the fornices, the presence of the os externum at the end of the protruded tumour, and that of the uterus within it—as demonstrated by the sound and examination *per rectum*—show that it is a case of prolapsus. If, however, the prolapsus be due to a fibroid tumour of the cervix and the os

¹ Faucon noted this in one case even though the patient was under an anæsthetic; the inversion was partial and only of one horn, and could not be recognised before the operation—*Sur une forme particulière d'Inversion polypeuse de l'utérus*, etc.—*Archiv de Toc.*, 1887, p. 1042.

A. Reposition.

The obstacle to reposition is the resistance of the tissue of the lower segment of the uterus; the principle of treatment is to overcome this by steady pressure.

Suppose that we have a case of inversion, how are we to proceed? The patient is kept perfectly at rest for a few days; injections of very warm water are employed twice or thrice daily; nutritious diet is given, and iron is usually required for anæmia. Ergot is required if there is menorrhagia; should it not be the menstrual period, the best thing to check hæmorrhage is injection of very hot water.

Having thus prepared the patient we proceed to reposition. Are we to employ the more rapid manual method or the slower one with an instrument? If the patient does not object to an operation under chloroform, and if we can have assistants to take turns with us in keeping up manual pressure, the former method should certainly be tried first.

Reposition
with the
hand.

(a.) *Reposition with the hand alone or aided by instruments.* For a few days previously, the largest size Barnes bag which the patient can bear is placed in the vagina and distended; this makes space for the operator's hand, and may itself effect the reposition.¹ The patient, under chloroform, is placed in the lithotomy position; pass the right hand into the vagina, and grasp the uterus with the fingers as far into the angle of reflexion as possible. Now press the uterus steadily upwards against the left hand on the abdomen. The fingers may be separated as far as possible so as to open out the cervix.²

Noegger-
ath.

White.

Atthill.

Sometimes the process of re-inversion is started by dimpling inwards one horn of the uterus, and then forcing the depressed horn onwards as a wedge to open up the ring of the cervix.³ As the hand cannot keep up steady pressure for any length of time, a cup is set on a curved iron rod with a spiral spring⁴ to make the pressure equal. A curved wooden rod, with a large cup at one end and a small one at the other, has also been used to keep up pressure.⁵ The end of the instrument is pressed against the operator's chest, and the cup is steadied with the hand in the vagina. It is evident that these instruments require a roomier vagina than when the hand alone is used; and if the cup slips unexpectedly it may rupture the fornix. Counter-pressure is made over the abdomen with the hand, or if the abdominal walls are thin and there is a distinct cup on the peritoneal aspect, with a cone of wood,⁶ which is used to dis-

Thomas.

¹ Kroner has collected six cases of inversion (longest of eleven years standing) replaced by this means; the pressure was applied for periods varying from one to eleven days.—*Archiv f. Gyn.*, B. xiv., S. 270.

² Emmet—*Op. cit.*, p. 418. It is very doubtful whether the constricting cervix has anything to do with preventing reposition, though upward and outward pressure round the neck favours it.

³ Noeggerath.—*Am. Med. Times*, 1862, Vol. iv., pp. 230, 235.

⁴ White—*Intern. Med. Cong. Trans.*, Philadelphia, 1876. Byrne—*New York Med. Journ.*, Oct., 1878.

⁵ Atthill—*Loc. cit.* Braxton Hicks—*Brit. Med. Journ.*, Aug. 1872.

⁶ Thomas—*Op. cit.*, p. 468.

tend the ring of the cervix; the traction can be taken off the vaginal Schroeder walls by fixing the cervix with volsellæ.¹ Counter-pressure may be made per rectum in the following way:—Pass index and middle fingers Courty. of right hand into rectum, draw down the uterus with the left hand until these fingers get fairly above the cervix so as to press on the margins of the peritoneal depression; grasp uterus now with left hand, turning it so that the fundus is towards the symphysis and the cervix towards the sacrum; finally, make pressure with the index and thumb in the angle of reflexion against the two fingers in the rectum.² This manual pressure is, with the help of assistants, to be kept up from half-an-hour to two hours according to the condition of the patient. If not successful in this time, the patient is kept in bed and under the influ-




FIG. 228.

WHITE'S REPOSITOR, WITH ELASTIC SPRING PLACED AGAINST THE OPERATOR'S CHEST. While the right steadies cup and uterus, counter-pressure is made with the left hand or better by an assistant (Thomas).

ence of opium while a Barnes bag is placed in the vagina to maintain the uterus as far as it has been replaced. When the uterus has been so far reinverted that the fundus is above the level of the os externum, the Emmet lips of the latter may be drawn together with wire sutures.³

Abdominal section, so as to allow the operator to get at the constricting rim of the cup Abdominal from its peritoneal side and dilate it with expanding forceps, has been proposed by Section Thomas. It was successful in the first case; a second proved fatal from peritonitis. It for Inver- has been tried unsuccessfully by A. R. Simpson, while Malins,⁴ and Mundé⁵ succeeded sion. so far in dilating the ring, but failed in pulling up the uterus by the ingenious method of passing a thread through the fundus; Schmalfuss⁶ has recently recorded a successful

¹ Schroeder—*Op. cit.*, S. 203. Atthill—*Loc. cit.*

³ Emmet—*Op. cit.*, p. 430.

⁵ *Amer. Journ. Obstet.*, 1888, p. 1279.

² Courty—*Maladies de l'utérus*, 1866.

⁴ *Lancet*, 1885, ii., 401.

⁶ *Centralb. f. Gyn.*, 1886, p. 745.

case. Brown¹ succeeded in dilating the ring by getting at it per vaginam through an incision in the inverted fundus; a dilator was introduced and the rim expanded: the incision in the uterus was stitched before the inverted fundus was pushed up.

Reposition
by Elastic
Pressure.

(b) *Reposition by continuous slight elastic pressure.* If manual reposition has failed, we try the more gradual method; in some cases we employ it from the first. Gradual pressure may be produced by an india-rubber bag placed in the vagina and distended with water from a douche-can so that hydrostatic pressure is brought to bear.² Thiry³ has devised an ingenious bag consisting of a double-walled india-rubber capsule, which is slipped over the uterus; when distended with air, it compresses and pushes up the inverted fundus. Pressure by an inflated bag is not so efficient as that produced by a wooden cup set on a stem⁴ with a vaginal (or, better still, a vaginal and perineal)⁵ curve so that the pressure is made in the axis of the brim. Pressure may also be made by the four elastic bands which pass, two in front and two behind, to a broad




FIG. 229.

CUP WITH STEM AND ELASTIC BANDS which are fixed to an abdominal belt, for gradual reduction of inversion (Thomas).

abdominal bandage; by the tightening of the front or the back bands, the direction of pressure is altered.

In this method there are two points which require careful attention. (1.) The elastic pressure must always act in the line of the axis of the inverted uterus, and likewise of the axis of the pelvic brim; the cup is apt to slip off the uterus, and the handle of the instrument to alter its direction. *Pressure in a wrong direction is injurious, and may produce sloughing.* To prevent these accidents we pad, with wadding soaked in carbolised oil, all round the neck of the inverted uterus and round the cup of the repositor when *in situ*; we watch the position of the instru-

¹ *New York Med. Journ.*, Nov. 24, 1883.

² Runge—*Lancet*, 1887, i., p. 1293. Jaggard records an interesting case of inversion of twenty-one months' standing reduced after thirty-three days' use of the colpeurynter—*Amer. Journ. Obstet.*, 1887, p. 130.

³ *Archiv de Tocolog.*, 1885, p. 925.

⁴ Lawson Tait—*Obstet. Journ.*, Vol. iv., p. 555.

⁵ Aveling—*Loc. cit.*, records ten cases of successful reposition with his cup and stem which has a sigmoid curve.

ment, and remove and re-apply it every day so as to see how it is pressing and whether there is sloughing.

(2.) There must be effective counter-pressure, so as to take the strain off the vaginal walls. This is effected by means of a broad flannel bandage, firmly secured round the loins, under which cotton wool is padded in such a way as to press exactly upon the fundus.

The elastic pressure is kept up from one to three weeks. Cases of reposition at this period, or even after it, are recorded.¹

In cases of inversion due to tumour growth, the tumour—if simple—must be removed in the first instance; we then wait to see if the uterus will replace itself, and if it does not we proceed to replace it. If the tumour be malignant, the propriety of amputating the uterus with the tumour must be considered.

B. Hysterectomy.

Where all means of reposition have been tried and failed, or where the uterus is ulcerated and gangrenous, hysterectomy is called for. Also in cases of inversion associated with malignant disease, *e.g.*, sarcoma. Formerly amputation of the uterus at the level of the cervix was the usual treatment for such cases, but now that vaginal hysterectomy has become a recognised operation without a serious mortality, it is preferable. The operation will be described under hysterectomy for cancer of the uterus.

¹ As by Neugebauer, after three weeks—*Centralb. f. Gyn.* 1887, p. 63.

CHAPTER XXXV.

TUMOURS OF THE UTERUS. FIBROID TUMOURS: PATHOLOGY AND ETIOLOGY.

LITERATURE.

Barnes—Diseases of Women, p. 746: London, 1878. *Doran*—On Myoma and Fibromyoma of the Uterus and allied Tumours of the Ovary: Trans. of Lond. Obs. Soc., 1888, p. 410. On the absorption of Fibroid Tumours of the Uterus, etc.: Trans. of Lond. Obs. Soc., 1893, p. 250. *Duncan, Matthews*—Hæmorrhage from Fibrous Tumours of the Uterus: Edin. Med. Jour., Jan. and Feb. 1867. *Gusserow*—Die Neubildungen des Uterus: Deutsche Chirurgie, Stuttgart, 1885. *Haultain*—Benign Growths of the Uterus: Allbutt and Playfair's System of Gynecology, London, 1896. *Johnston*—A Review of some collected Cases of Fibromata of the Cervix Uteri: Am. Journ. of Obstet., Nov. and Dec. 1885. *Kleinwächter*—Zur Entwicklung der Myome des Uterus: Zeits. f. Geb. u. Gyn., Bd. ix., S. 68. Zur Biologie der Fibromyome des Uterus: Zeits. f. Geb. u. Gyn., 1892-93, S. 164. *Klob*—Pathologische Anatomie der weiblichen Sexualorgane, S. 149: Wien, 1864. *Lee*—Tumours of the Uterus: London, 1847. *M'Clintock*—Diseases of Women: Dublin, 1863. *Reamy*—Case of Fibroid Polypus of the Uterus, with remarks on some points in Etiology (with Discussion): Amer. Jour. Gyn., 1886, pp. 813 and 859. *Routh*—Fibrous Tumours of the Womb: London, 1864. *Schorler*—Ueber Fibromyome des Uterus: Zeits. f. Geb. u. Gyn., Bd. xi., S. 139. *Schroeder*—Krankheiten der weiblichen Geschlechtsorgane, S. 218: Leipzig, 1886. *Simpson, Sir J. Y.*—Diseases of Women, p. 659: Edin., 1872. *Thomas*—Diseases of Women, p. 519: Philadelphia, 1880. *Winckel*—Ueber Myome des Uterus, etc.: Volkmann's Sammlung klin. Vorträge, No. 98, 1876. *Wyder*—Beiträge zur norm. und patholog. Histolog. der Uterusschleimhaut: Archiv f. Gyn., Bd. xiii., S. 35. Die Mucosa uteri bei Myomen: Arch. f. Gyn., Bd. xxix., S. 1. See also references in the text.

UNDER Tumours of the Uterus we have to study—

Fibroids,	Papilloma,
Polypi,	Carcinoma,
Tuberculosis,	Sarcoma,
Adenoma,	Deciduoma malignum.

Intro-
ductory.

The term "polypus" is so convenient clinically that we retain it, but we must remember that it involves cross-classification, including one variety of fibroid tumour—the fibrous polypus—while the mucous polypus is a pediculated adenoma. As adenoma is of importance in relation to commencing carcinoma, we shall refer to it also under that

subject. The so-called "Deciduoma malignum" will be treated of under Sarcoma. In the chapters that follow we shall consider Fibroids and Cancer at some length, the others briefly.

Fibroid tumour is considered first, as in frequency it comes before cancer, although in seriousness the latter is by far the more important. It presents a remarkable contrast with cancer in every respect: it shows itself early in life, while cancer is late; it occurs among the well-to-do, while cancer makes its ravages among the poor and badly fed; it is the tumour of the sterile, while cancer is that of the fertile; it rarely affects life, while the fate of the cancer-patient is almost sealed.

Synonyms.—Myoma or Fibro-myoma Uteri; Fibrous Tumour; Tumeur fibreuse; Hystérôme.

As this tumour is composed of both the connective tissue and muscular elements of the wall of the uterus, it is at once a fibroma and a myoma; the most correct term is therefore *fibro-myoma*. In the majority of cases, however, the fibrous tissue preponderates, so that the tumour resembles a fibroma; the English term fibroid (a term derived from the root of fibroma and *εἶδος*, = like a fibrous tumour) is therefore not inappropriate, and is also more convenient.

PATHOLOGY.

Under this head we shall describe their

Situation;

Structure—naked eye and microscopic;

Mode of growth, varieties;

Changes in uterus;

Degenerative changes.

SITUATION.

They occur much more frequently in the body of the uterus than in the cervix; of seventy-four cases of fibroid tumours recorded by Lee, only four were in the cervix. In the body of the uterus the most common seat is *the posterior wall*; they occur less frequently in the anterior wall, and very rarely at the sides of the uterus. The soft, truly muscular form is most commonly situated at the fundus.

STRUCTURE.

They are composed of the same elements as the muscular wall of the uterus, viz., of non-striped muscular fibre and fibrous tissue. These are both present in every case, as the name for these tumours (fibro-myoma) implies. The proportion of these constituents, however, varies; in some rare cases the muscular tissue preponderates, producing a true *myoma* which is not circumscribed and grows rapidly; more usually there is excess of fibrous tissue producing a *fibro-myoma*, which is dis-

Nomen-
clature.

Naked-eye
Structure
of a Fibroid
Tumour.

tinctly marked off from the wall of the uterus and grows slowly. The naked-eye characters of the myoma are those of a pale, flesh-coloured tumour having a soft consistence, passing gradually into the surrounding uterine wall, and usually single. The fibro-myoma, by far the most frequent form,¹ is of firm consistence which makes it feel like a foreign body in the softer muscular wall; it is of a pale colour, resembling fibrous tissue; it cuts like cartilage, the cut surface having a glistening satin-like appearance and being often uneven through the firmer fibrous tissue forcing out the softer parts between; the bundles of fibrous tissue have a concentric arrangement round one or more centres (fig. 230).




FIG. 230.

SECTION OF A LARGE FIBROID TUMOUR, with the Fibres arranged round several centres
(Sir J. Y. Simpson).

Capsule of
a Fibroid.

The tumour is surrounded by loose fibrous tissue, which with the immediately adjoining muscular layer constitutes the so-called *capsule*; it has a broad connection at one point with the muscular tissues of the wall, or lies free in its capsule. Sometimes the submucous and sub-peritoneal forms develop so near the surface that they have not a layer of the muscular wall to form a capsule and may hence be called "free" in contrast with the encapsulated (Haultain).² The loose fibrous tissue which separates the tumour from the adjoining wall is

¹ Doran (*loc. cit.*) thinks the frequency of fibro-myoma as compared with myoma is over-estimated. Young fibroids are "pure myomata with or without connective tissue."

² *Op. cit.* This monograph contains beautiful illustrations of the various seats and structure of fibroid tumours.

however present in these cases. The looseness of the tissue round the tumour is important in relation to its removal by the process described as *enucleation*. Few blood-vessels penetrate into the substance of the tumour, although the tissue immediately round it is very vascular and often contains enlarged veins which resemble the venous sinuses of the pregnant uterus (fig. 239); nutrition is apparently effected by transudation from the capsule. In some rare cases, however, these tumours possess a cavernous structure consisting of dilated blood-vessels.




FIG. 231.

SECTION OF FIBROID TUMOUR, showing wavy bundles of fibrous tissue † (Gusserow).

Virchow has described this form as “*Myoma teleangiectodes seu cavernosum* ;” cases are recorded by Leopold and Schroeder.

On microscopic examination, the myomatous form has the appearance of muscular fibre of the uterus—the muscle-cells being, according to Doran,¹ larger than those of the uterus in which it grows. The fibromatous form (common fibroid tumour) has the appearance shown



FIG. 232.

SECTION OF FIBROID TUMOUR, showing spaces between bundles of fibrous tissue † (Gusserow).

at fig. 231, in which the wavy bundles of fibrous tissue are well seen. Sometimes the bundles of fibrous tissue are separated by spaces (fig. 232), which Klebs considers to be lymphatic spaces. Nerves have been traced into the substance of these tumours by Lorey; but, as an interesting case recorded by Freund shows, they are not sensitive:—a submucous fibroid was extruded beyond the vulva; the lower third,

¹ *Loc. cit.* He figures a section of a myoma from a pregnant uterus which shows this well, the muscle-cells being still larger than the hypertrophied ones of the uterus.

which protruded beyond its capsule of mucous membrane, was not sensitive to the prick of a needle; the upper two-thirds, from their being still covered by *mucous membrane*, were very sensitive. The mucous membrane covering them is ciliated,¹ like that of the uterus generally; though when it has been exposed for some time (*e.g.* when a fibrous polypus comes to be in the vagina) it becomes squamous.²

Epithelial structures have been found in fibroid tumours by Rieke,³ Orloff,⁴ Huser,⁵ and Schottländer.⁶ Some have seen in this an argument for Cohnheim's hypothesis as to the embryonic origin of these tumours, that this epithelium represents a diverticulum of the duct of Müller, or is a relic of the Wolffian duct. It may, however, also have arisen from the uterine mucosa at a late period, and have been cut off from it by the growth of the tumour. While it thus throws no light on the origin of fibroids, it is of interest with regard to carcinomatous degeneration in them.

MODE OF GROWTH—VARIETIES.

Rate of
Growth.

Fibroid tumours grow slowly; the more they consist of fibrous tissue, the slower the growth. During pregnancy, they increase more rapidly in size; in the puerperium, they may become smaller again and even cease to be recognisable. It is difficult to determine the rapidity of growth. It is unsatisfactory to estimate it from the appearance of symptoms and compare the time elapsed with the present size of the tumour; the only reliable data are got from the examination of the tumour from time to time. Schorler has reported on eighteen cases observed by Schroeder and comes to this conclusion: a tumour will not grow to be for the first time recognisable in less than three months' time, and in a year may not be much larger; in five years it may grow to the size of a man's fist, and in thirteen to the size of the head. On the other hand, Kleinwächter, from the study of forty cases under his own care, affirms that this represents their growth as more gradual than the facts warrant. While for long periods they may remain stationary, at others they increase perceptibly from month to month. He also thinks it exceptional for them to cease growing after the menopause.

After the menopause, their growth is, as a rule, arrested; the menopause is generally late in cases of Fibroids.

Mode of
Growth.

All fibroid tumours are, in the beginning, interstitial or intra-mural. As they increase in size they expand in the substance of the wall or towards one of the free surfaces (peritoneal or mucous), thus becoming subperitoneal or submucous. Hence three varieties are recognised—

¹ Gervis—*Brit. Med. Journ.*, 1886, ii., p. 871.

² Reamy—*Loc. cit.*, p. 817.

³ Beitrag zur Aetiolog. der Uterusgeschwülste—*Arch. f. patholog. Anatom.*, Bd. cxlii., S. 133.

⁴ Zur Genese der Uterus-myoma—*Zeits. f. Heilkunde*, Berlin-Prag., Bd. xvi., S. 311.

⁵ Ueber das Vorkommen Drüsenschläuchen in einem Fibromyoma Uteri: *Münch. med. Wochenschr.* No. 10.

⁶ Ueber drüsige Elemente in Fibromyomen des Uterus: *Zeits. f. Geb. u. Gyn.*, Bd. xxvii.

interstitial, subperitoneal, and submucous. It is evident that these terms are relative, as it is difficult to say when an interstitial fibroid becomes submucous. Gusserow limits the term "submucous" to *pediculated* submucous, and "subperitoneal" to *pediculated* subperitoneal fibroids. A submucous tumour, however, often gives rise to the clinical signs diagnostic of the submucous variety long before it becomes pediculated. Each variety requires short description. For the sake of convenience we describe first the fibroid tumours found in the body of the uterus; the comparatively rare fibroid tumours of the cervix are best noticed separately (p. 433).

A. The Subperitoneal grow outwards into the peritoneal cavity. The thickness of the pedicle varies (fig. 233); its length determines the mobility of the tumour. When the tumour attains a certain size, one of two things happens. (1.) It may grow up into the abdomen and expanding there draw the uterus forcibly upwards, producing by this traction elongation of the cavity (fig. 233) with thinning of the walls. An interesting case is recorded by Tinns,¹ in which the cavity of the body of the uterus was elongated to 6 inches; the cervical canal, extending only 1 inch inwards from the os externum, ended blindly at a point 2 inches distant from the beginning of the cavity of the body; the intervening portion was obliterated so as to form a solid muscular cord. Virchow says that the body may even be torn from the cervix by forcible traction. (2.) The tumour, growing from the first within the pelvis, may through pressure produce the symptoms of incarceration; or, having a long pedicle, may fall down from the abdomen into the pelvis and produce similar symptoms. The point of origin of the tumour and the length of the pedicle determine whether these symptoms can be relieved by pushing the tumour out of the pelvis. Twisting of the pedicle occurs less frequently in fibroid than in ovarian tumours; when it occurs, it leads to œdema or gangrene. Schroeder² mentions a case where, on operating, he found the tumour distended with blood from partial twisting of the pedicle. Timmers records a case, diagnosed as an ovarian tumour with twisted pedicle, treated successfully by laparotomy.³ Adhesions form with other organs, as occurs with all abdominal tumours; these may become new sources of nutrition. Sometimes they lead to detachment of the tumour from the uterus: the tumour is anchored, as it were, to the abdominal walls; and, when the uterus from pregnancy or other causes becomes displaced, the pedicle gives way. Turner⁴ reports a case in which a small calcareous fibroid was found free in the pouch of Douglas; a second was attached to the posterior wall of the bladder and to the pelvis; a third was bound down to

Varieties
of Fibroid
Tumours.

Subperi-
toneal
Fibroids.

Growth
into Abdo-
men.

Incarcera-
tion in
Pelvis.

¹ *Lond. Obst. Trans.*, Vol. ii., p. 34.

² *Op cit.*, S. 230.

³ *Centralt. f. Gyn.*, 1892, S. 242. See also a case by Cappie—*Obstet. Journ.*, ii., p. 302. Twisting of the uterus four times round its axis, is recorded by Pick: *Cent. f. Gyn.*, 1892, S. 445.

⁴ *Edin. Med. Journ.*, 1861, p. 698.

the bladder and the pelvic wall by adhesions, but still retained its connection with the uterus by a thin pedicle. Adhesions to the intestines have produced symptoms of intestinal obstruction.¹ Hernial protrusion of the abdominal walls has been described by Düll:² he reports two cases of this very rare occurrence; in one case, the skin covering the hernial sac became gangrenous, so that the tumour lay exposed.

B. The Interstitial remain in the substance of the uterine wall, and do not become pediculated. The appearance of such a tumour is well

Interstitial
Fibroids.




FIG. 233.

UTERUS WITH ELONGATED CAVITY DUE TO THE PRESENCE OF SEVERAL FIBROID TUMOURS
(Sir J. Y. Simpson).

seen at fig. 234. Usually there are many such tumours present (fig. 233); Schultze counted as many as fifty in one uterus, and Thomas describes the uterus of a negress containing thirty-five.

C. The Submucous are the most important clinically. They lie immediately underneath the uterine mucous membrane, and project into

Sub-
mucous
Fibroids.

¹ Eade—*Lancet*, Dec. 21, 1872.

² Cited by Schroeder, *op. cit.*, S. 233. Lawson Tait mentions the same condition—*Brit. Med. Journ.*, 1888, i., p. 861.

the cavity of the uterus (fig. 235). They are attached over a broad base, or by a pedicle; when they hang free, they are known as fibrous polypi—the most frequent form of uterine polypus (*v.* Chap. XXXIX.). When a fibrous tumour projects into the uterine cavity, it acts as a foreign body and produces uterine contractions. These lead, in some instances, to *pedunculation* of the tumour and even to its extrusion from the uterine cavity; in such a case, it hangs as a polypus in the vagina. In other rare cases, the capsule ruptures and the liberated tumour is expelled in shreds—*spontaneous enucleation*.

The muscular wall hypertrophies, more especially when the tumour is submucous or interstitial. A small fibroid lying in the lower segment ^{Changes in Uterus.}




FIG. 234.
INTERSTITIAL FIBROID TUMOUR (Sir J. Y. Simpson).

of the uterus has caused the whole organ to hypertrophy to the size of a child's head.¹ In submucous fibroids, the *mucous membrane* is also hypertrophied.

Wyder² has studied the hypertrophied mucosa with special reference to the causation of the bleeding, and describes a glandular and interstitial endometritis.

From a comparison of the mucous membrane in subperitoneal as compared with interstitial tumours he comes to the conclusion that the thicker the muscular capsule is the less likely is the tumour to affect the

¹ Tillaux—*Gaz. des Hôp.*, 1867, No. 144.

² *Op. cit.*—He examined twenty uteri extirpated in Gusserow's Clinique for fibroid tumour.

circulation in the mucous membrane. The uterine glands in this case are hypertrophied, but the interglandular tissue little or not at all affected; while the nearer the tumour comes to the uterine cavity, the more does the interglandular connective tissue become affected, and this sometimes at the expense of the glands which atrophy. The bearing of this on bleeding is that it is the cicatrization of the interglandular connective tissue, causing compression of the veins, which leads to congestion and bleeding. Semb¹ finds a hypertrophy of the mucosa which he does not consider inflammatory in nature as Wyder does. This is often followed by atrophy from the pressure of the tumour, or inflammatory and other complications. Bleeding is not due to changes in the mucosa but to general enlargement of the uterus.




FIG. 235.

SUBMUCOUS FIBROID TUMOUR PROJECTING INTO UTERINE CAVITY (Sir J. Y. Simpson).

Changes in the position of the uterus have been already referred to; when subperitoneal fibroids rise up into the abdomen, it is sometimes drawn forcibly upwards by them and may be twisted on itself.² At other times the weight of a subperitoneal or interstitial tumour leads to prolapsus uteri. Inversion of the uterus is also occasioned by sub-

¹ From the examination of twenty-three fibroid uteri extirpated in Leopold's Clinique at Dresden. —Ueber das Verhalten der Uterusschleimhaut bei Myomen: *Archiv f. Gyn.*, 1893, Vol. xliii., S. 200. Marchesi also describes changes which he considers hyperplastic rather than inflammatory—Le alterazioni dell'endometrio in casi di fibromi dell'utero: *Annali di Ostet. et Gyn.*, Dec. 1894.

² As in the case reported by Küster—*Beiträge zur Geb. u. Gyn.*, 1872, i., S. 7; the uterus was twisted two and a half times, so that the broad ligaments formed a spiral. Skutsch records another case operated on by Schultze in which the uterus was twisted half round—*Centralb. f. Gyn.*, 1887, S. 52.

mucous fibroids when these are situated near the fundus and when their pedicle does not admit of their extrusion as polypi.¹

Pressure effects on the ureters and bladder, the rectum, and the pelvic vessels and nerves will be considered in Chap. XXXVI. *Affections of remote organs* occur in large tumours. Thus Bantock² considers fatty




FIG. 236.

PEDICULATED SUBMUCOUS FIBROID IN PROCESS OF EXTRUSION (Sir J. Y. Simpson).

degeneration of the liver a consequence of fibroid tumours, while hypertrophy and sometimes dilatation of the heart is present, as in the case

¹ Kötschau records a case of partial inversion, with what he calls "eversion of the uterine mucous membrane," i.e. its being pushed downwards without the tumour's becoming pediculated—*Centralb. f. Gyn.*, 1887, S. 757.

² *Brit. Gyn. Journ.*, 1887, Vol. ii., p. 84.

of large abdominal tumours generally.¹ *Changes in the uterine appendages*, ovaritis² and salpingitis also occur. Hydro- and hæmatosalpinx³ have been found.

DEGENERATIVE CHANGES.

These are the following:—Softening, Induration, Calcification, Suppuration, and (very rarely) Malignant Degeneration.

Softening. The *softening* may be due to œdema, to fatty degeneration, or to myxomatous degeneration. The occurrence of œdema is unquestioned, and many cases of sudden increase in the size of fibroid tumours may be thus explained. Fatty degeneration is a rare occurrence.⁴ Myxomatous degeneration, resulting in the formation of spaces containing mucus between the layers of the tumour, sometimes occurs. These changes pass into the fibro-cystic tumour (*v. Chap. XXXVIII.*).

Induration. *Induration*, with atrophy or shrinking of the tumour, occurs in some cases after the menopause; the muscular tissue fattily degenerates and disappears, the fibrous tissue contracts.⁵ An infarction has also been found.⁶

Calcification. When *calcification* occurs, lime salts (chiefly phosphates) are deposited in the fibrous tissue and produce the so-called womb-stones.⁷ This deposit usually commences in the centre of the tumour and extends outwards, more rarely in the external layers so as to form a shell round the tumour. Sometimes it is so extensive that the tumour can be cut with the saw, and the cut surface polished; more usually it is incomplete, and forms a coral-like skeleton. Calcification of portions of the tumour is often accompanied by suppuration in others, probably from interference with nutrition.

Suppuration. *Suppuration* occurs frequently in submucous fibroids, as the result of injury from operative interference or from constriction of the pedicle during the process of expulsion. It has also been observed as a rare occurrence in subperitoneal fibroids, accompanying calcification or from torsion of the pedicle. In such a case, the pus either finds its way through the abdominal walls or fatal peritonitis follows.

Malignant Degeneration. Whether *malignant* degeneration specially affects fibroid tumours, is a

¹ Fenwick—On intra-abdominal tumours as a cause of cardiac degeneration: *Brit. Gyn. Journ.*, 1887, Vol. ii., p. 72. Hennig has found circulatory disturbances even with small fibroids: *Cent. f. Gyn.*, 1894, S. 724.

² Bulius has examined the ovaries from fifty cases of fibroids and found increase of the stroma and changes in the vessels with cystic degeneration of the follicles—*Verhalten des Eierstockes bei Fibro-myoma Uteri*: IV. Gynäkolog. Congress zu Bonn, 1891. So also Popow has examined the ovaries from forty cases in Lebedeff's Clinique at St. Petersburg. He finds interstitial oöphoritis frequent, with occasionally cystic degeneration and atresia of the follicles: *Zur Frage über die Veränderungen der Ovarien bei Fibromyoma Uteri*: *Cent. f. Gyn.*, 1890, No. 49. Adeno-epithelioma has also been described (Edebohls, *Am. Journ. Obstet.*, 1891, p. 619).

³ Röse—*Deut. Zeits. f. Chirurg.*, Bd. xxv., Hft. 5, S. 456. Jacobs—*Cent. f. Gyn.*, 1891, S. 117.

⁴ See recent case by Martin—*Cent. f. Gyn.*, 1893, S. 212.

⁵ Sir J. Y. Simpson—*Obst. Mem.*, p. 115.

⁶ By v. Ott. The patient had felt pain over it, ascribed to a local peritonitis—*Centralb. f. Gyn.*, xii., S. 274.

⁷ See a recent case by Thorn—*Zur Casuistik der Uterussteine*: *Zeits. f. Geb. u. Gyn.*, Bd. xxviii.

disputed point. A few cases of carcinomatous¹ changes in the endometrium, and even in the heart of the tumour, have been described, as also of sarcomatous degeneration² in the substance of the tumour. These are so rare, in proportion to the frequency of fibroids, that their occurrence may be a coincidence.

As to the frequency of these various changes, Martin³ gives us the following interesting statistics of his own cases. Of 205 fibroids he found slight retrogressive changes in 70, fatty degeneration in 3, suppuration in 10, œdematous swelling in 11, cystic degeneration in 8, blood-cavities in 3, sarcomatous degeneration in 6, but never carcinoma.

Fibroid Tumours of the Cervix.

The occurrence of fibroid tumours in the cervix is rare; but, when they are present, they often give rise to difficulty in diagnosis on account of the distortion which they produce. They spring from either wall, and grow outwards towards the peritoneal cavity and into the cellular tissue, or downwards into the vagina. In the former case they dissect off the peritoneum, becoming "intra-ligamentous;" and from their situation, embedded in the cellular tissue, and the symptoms produced by their incarceration, they become a source of great danger.⁴ When submucous, they produce elongation of one lip and may form a polypoidal tumour in the vagina (fig. 237); the accompanying distortion of the os externum leads to difficulty in diagnosis. Cases in which a large tumour bulges through the ostium vaginæ have been mistaken for inversion and prolapsus. Sometimes prolapsus is due to the weight of the tumour and disappears after its removal.⁵ The interstitial form is easily mistaken for inversion, when the os is converted into a transverse cleft which escapes observation and the unaffected lip is thinned out to a mere band.

Johnston reports on ninety-six cases of fibroid tumour of the cervix, dealing especially with their effect on pregnancy and labour. He finds that abortion is more frequent with fibroid tumours in the body, premature labour with those in the cervix; he affirms that during pregnancy or labour one-third of the mothers and more than one-half of the children die so that, where the tumour cannot be removed, celibacy is to be recommended.

¹ See Ehrendorfer's interesting paper on this subject—Die primäre carcinomatöse Degeneration der Fibromyome des Uterus: *Cent. f. Gyn.*, 1892, S. 513. Semb (*Archiv f. Gyn.*, Bd. xliii., S. 221) gives a case and figures, also a peculiar papillary hypertrophy of the epithelium. Also Geuer—"Ueber das gleichzeitige Vorkommen von Carcinom und Myom am Uterus:" *Cent. f. Gyn.*, 1894, S. 341.

² See "Sarcoma Uteri," Chap. XLIV. Also Ricker—Beitrag zur Etiolog. der Uterusgeschwülste: *Archiv f. patholog. Anatom.*, Bd. cxlii., S. 193.

³ Ueber Myome: *Archiv f. Gyn.*, Bd. xxxii., S. 470.

⁴ Pozzi—"De la valeur de l'hystérectomie, etc.": Thèse d'agrég., Paris, 1875.

⁵ Barnes—*Obst. Trans.*, iii., p. 211.

ETIOLOGY.

Gusserow, to whose exhaustive article—*Die Neubildungen des Uterus*—in Billroth's *Handbuch* we are greatly indebted in this Chapter, says in regard to etiology, "Ueber die Ursachen der Uterusmyome wissen wir so wenig, wie über die Ursachen der meisten pathologischen Neubildungen, nämlich *Nichts*" (of the causes of fibroid tumours we know as little as of the causes of most pathological new-formations, that is *nothing*). Virchow and Winckel have both made elaborate attempts to assign a cause to the development of fibroid tumours. The number and variety of causes adduced by these observers only show how far we are from the knowledge of the real cause; with such a variety of causes, the difficulty would not be to explain why they are present in some but




FIG. 237.

CERVICAL FIBROUS POLYPUS springing by a pedicle from the region of the os internum, and pushing itself under the whole mucous membrane of the cervical canal; so that its insertion is partly continuous with the tissue of the uterus, partly truly submucous. Between these a cavity has formed through tearing of the mucous membrane, so that the tumour has apparently two pedicles (*Schroeder*).

why they are not present in every case. The development of the true *myoma* has been studied by Kleinwächter. He examined uteri with very small myomata and found that there was a small isthmus of muscular fibre uniting the myomatous mass, lying in its connective tissue capsule, with the muscular tissue around. This isthmus sometimes bifurcates and resembles in form an obliterated blood-vessel (capillary). He also saw some capillaries surrounded with round cells and forms transitional to muscular fibres. Hence he concludes that the true myoma is due to a degeneration of a blood-vessel with its branches. From finding micrococci in them, Galippe and Landouzy¹ have suggested that they are due to the irritation of a parasite. More recently,

¹ *Brit. Med. Journ.*, 1887, i., p. 799.

TABLE AND DIAGRAM SHOWING FREQUENCY OF FIBROID TUMOURS ACCORDING TO AGE OF PATIENT.




FIG. 238.

Gottschalk¹ has also described an endarteritis and proliferation of the connective tissue around the vessels. Fibroids are caused by some irritant leading to active congestion. This irritant may be chemical or parasitic—it is not a coccus; nor is the change an inflammation, but a new formation.

Olshausen² has found pain (sensitiveness to pressure and dysmenorrhœa) and menorrhagia complained of before any tumour could be detected by palpation, and thinks this points to congestion of the uterus as being an early clinical symptom in some cases of myoma.

Fibroids are without doubt the most frequent new-formation in the uterus. Klob says that they are present in 50 p. c. of women who die over fifty years of age; and Bayle, in 20 p. c. of those who die over thirty-five years; both of these estimates are probably beyond the mark.

Develop-
ment of
Fibroids
according
to age.

Their appearing is in some way related to the development of the sexual apparatus. Thus, there are no well-authenticated cases of their arising before puberty³ or after the menopause. The majority of patients are between the ages of thirty and forty when they first seek medical advice, as it is evident from the accompanying table based on statistics collected by Gusserow (fig. 238). Schroeder says that of 196 patients, who during three years of his private practice consulted him for fibroid tumours, 104 were between forty and fifty, and 62 between thirty and forty.

Sexual activity seems to predispose to their development, as a much larger number of patients with fibroid tumours belong to the married than the unmarried class. Of 1876 cases from various authorities collected by Reamy,⁴ we find that 1422 or 75 p. c. of persons with fibroid tumours seeking advice were married. But when we take into account the larger proportion of the married in the population generally, and the still larger proportion in those who seek advice, we find that we cannot affirm that fibroids are more common in the married than the unmarried.⁵ As the presence of a fibroid tumour interferes with conception, we often find sterility present.

¹ Ueber die Histogenese und Etiologie der Uterusmyome: *Archiv f. Gynäk.*, 1895, Bd. xliii., S. 534.

² Notizen ueber das klinische Anfangsstadium der Myome: *Archiv f. Gynäk.*, xxviii. S. 494.

³ Tillaux reports a case of a fibroid tumour of the cervix in a girl of nineteen which had caused symptoms for six years.—*Annales de Gyn.*, xxvi., p. 241.

⁴ *Loc. cit.*, p. 818.

⁵ Schumacher found that in the Basle Clinique the frequency of the married to the unmarried among the cases of fibro-myoma were as two to one; while the relative proportion of married to unmarried who sought advice was as five to one, and in the general population between thirty and fifty years of age as 3·4 to 1: *Beiträge zur Etiologie der Uterus-Fibromyome*, Basle, 1889.

CHAPTER XXXVI.

FIBROID TUMOURS OF THE UTERUS: SYMPTOMS; DIAGNOSIS; PROGNOSIS.

LITERATURE.

See Literature of Chaps. XXXV. and XXXVII.

LIKE other pathological conditions of the uterus, fibroid tumours sometimes produce no symptoms and their presence is discovered accidentally or on *post-mortem* examination. This absence of symptoms is more likely to occur should the tumour be small, or should there be no sexual activity as in unmarried women. In the latter case, although symptoms appear only when the patient enters married life, the tumour may have been already a long time present. Subperitoneal tumours, even when large, may only produce discomfort from undue abdominal distention.

The symptoms usually present may be tabulated as follows:—

1. Menorrhagia, irregular hæmorrhages ;
2. Painful menstruation ;
3. Pelvic sensations due to size and weight of tumour, peritonitic pain ;
4. Symptoms of pressure on bladder and rectum,
blood-vessels and nerves,
ureters ;
5. Sterility and abortion.

1. *Hæmorrhage* is the most characteristic symptom in submucous fibroids, and appears first as a *gradual increase* of the normal menstrual flow ; it never begins with a sudden flooding as in carcinoma uteri. In *menorrhagia*, the hæmorrhage comes from the hypertrophied mucous membrane of the uterine cavity generally ; it does not come from the mucous membrane covering the surface of the tumour which is frequently thinned and atrophied, nor from the substance of the tumour

Hæmor-
rhage in
Fibroids.

itself which as we have seen is sparingly vascular. When, however, the submucous fibroid projects as a polypus, passive congestion and hæmorrhage from the mucous membrane covering it may be occasioned by the constriction of its pedicle. *Irregular hæmorrhages* arise from ulceration of the mucous membrane covering the tumour, or rupture of the dilated veins in its capsule. Fig. 239 shows a case¹ in which,




FIG. 239.

UTERUS CONTAINING FIBROID TUMOUR, from a case which terminated fatally through hæmorrhage. Note the large venous sinuses in the capsule, one of which ruptured at the point *a* (*Matthews Duncan*).

through the rupture of a uterine sinus in the lower part of the tumour, a sudden and fatal hæmorrhage occurred. In subperitoneal fibroids menstruation is not increased, and in certain rare cases is diminished.

¹ Reported by Matthews Duncan—*Edin. Med. Jour.*, 1867, p. 634. He also refers to a case of Cruveilhier's in which death was occasioned in the same way.

2. *Pain* accompanies *menstruation*. In the submucous variety there is often characteristic uterine *dysmenorrhœa*, in which the pain resembles labour pains. The congestion causes the polypus to swell and this produces uterine contractions (*v.* Uterine Polypi). In interstitial and even in subserous fibroids, there is often pain at the menstrual period which cannot be thus explained. In subserous fibroids with a pedicle containing large vessels, as well as in interstitial, Gusserow ascribes the pain to the distention of the tumour with blood. This pain is of a stretching or dragging nature, and is quite different from the pain of uterine contractions.

Pain in
Fibroids.

3. *Increased weight* of the uterus occasions *sensations* of discomfort, which are described as "fulness or weight in the pelvis," "a sensation of dragging," "bearing-down pain." When the tumour is so large that it fills the pelvis and becomes wedged in it, intense pain is produced; this is either always present, or recurs only at the menstrual periods when the tumour is distended by blood. As in carcinoma uteri, peritonitic pains—indicated by local tenderness and reflex contraction of abdominal muscles—may arise at any time from secondary chronic peritonitis. Neuralgic pain is sometimes present locally (see below), but may be also through the whole body.

Weight
Symptoms.

4. Frequency of micturition, due to *pressure on the bladder*, is the most common pressure symptom. *Pressure on the urethra* produces difficulty of micturition and even retention; with some patients, this recurs regularly at the menstrual period. Even very small fibroids, when they are situated in the anterior uterine wall, may press on the neck of the bladder and produce symptoms of cystitis. *Pressure on the rectum* by fibroids in the posterior wall occasions constipation or, more rarely, mucous diarrhœa. Incarcerated fibroids have produced complete obstruction, and led to a fatal result¹ or furnished an indication for colotomy. Intestinal obstruction has also resulted from adhesions between the tumour and the small intestine.² *Pressure on the veins* produces hæmorrhoids and varicose veins in the legs. Interesting cases of neuralgia due to pressure on *pelvic nerves* have been recorded. In these cases the neuralgia entirely disappeared as soon as the tumour was lifted up and supported by a pessary.³ Compression of the *ureters*, with consequent dilatation and hydronephrosis, occurs less frequently in fibroid tumours than in carcinoma. The reason for this is evident; in carcinoma the compression is due to infiltration of the tissue round the ureter, which from the anatomical relation of the ureters to the cervix easily occurs; fibroid tumours in their growth simply press against the ureters, and may push them aside. Several cases of single and double hydronephrosis and of death from uræmia⁴ have been re-

Pressure
Symptoms.

¹ Holdhouse—*Lond. Path. Soc. Trans.*, iii., 371.

² Eade—*Lancet*, Dec. 21, 1872.

³ Kidd—*Dub. Quart. Journ.*, 1872. Jude Hûe—*Annales de Gyn.*, iv., p. 239.

⁴ Gusserow quotes cases from Jude Hûe, Murphy, Hanot—*Neubildungen*, etc., S. 52.

corded. Bright's disease has developed secondarily.¹ In fibroid tumours where pressure symptoms are present, we should always examine the urine.

5. *Sterility* is frequent. Of 149 cases of married women collected by Schroeder, 33 per cent. were sterile and the average number of children to each mother was about three. When conception occurs, fibroid tumour may lead to abortion or complicate labour.

Sterility
in Fibroids.

PROGRESS AND RESULTS.

A *relative cure* usually takes place at the menopause, when the tumour ceases to grow.² In the case of subserous tumours, this may happen even before that time.

Spontaneous disappearance.

Spontaneous disappearance of the tumour has been observed in certain cases, although nothing definite is known as to the means by which it is effected. After a careful sifting of the literature of this subject, Doran³ has collected thirty-seven cases in which this occurred. Of these, thirteen were associated with the puerperium which seems to affect the tumour by a process comparable to involution. In sixteen cases the disappearance occurred apart from the puerperium; and as the patients were under forty-five, the effect of the menopause may be excluded: in six of them there was a history of inflammatory complications, leaving ten cases of apparently spontaneous disappearance. The remaining eight cases occurred after forty-five,⁴ and may be accounted for by the effect of the menopause.

Spontaneous Expulsion.

Complete cure also results from *spontaneous expulsion*. This occurs in three ways:—

- (1.) By pediculation and extrusion of the tumour as a polypus (*v.* under Uterine Polypi);
- (2.) By enucleation, in which the tumour is shelled *en masse* out of its bed;
- (3.) By the breaking-down of its substance and consequent expulsion in fragments.

Spontaneous Enucleation.

Enucleation occurs in submucous and also in interstitial tumours. The mucous membrane of the capsule ulcerates, and the tumour is thus exposed; partly through suppuration, partly through uterine contractions, it becomes detached all along the line of its capsule and, being thus liberated, is expelled. This process is comparatively safe for the patient, though there is always the risk of hæmorrhage from the large veins in the capsule (fig. 239). In spontaneous enucleation, suppuration does not occur in the tumour itself but only in its capsule.

¹ Hubert—*Bul. de la Soc. Anatom.*, 1873, p. 870.

² Exceptionally such tumours continue to grow after the menopause. See paper by Müller and discussion: *Archiv f. Gyn.*, 1891, Bd. xl., S. 340. So also Kleinwächter, *v. ante* p. 426.

³ *Op. cit.*

⁴ In two of these, the age is not given; and in one it is doubtful.

The *breaking-down* of the substance of the tumour is a much more dangerous process for the patient. As it is a slow one, there is a risk of absorption of septic matter. The commencement of this change is indicated by increase in the size of the tumour, which becomes tense and painful to the touch. There is a purulent fœtid discharge from the vagina, and sometimes hæmorrhage. The constitutional symptoms of loss of appetite and hectic fever afterwards develop, and most of such cases end fatally.

Expulsion of the tumour generally takes place per vaginam. As in other tumours we have inflammatory adhesions forming with neighbouring organs, followed by suppuration and perforation by the tumour. Thus calcified fibroids have perforated into the bladder, and have been mistaken for vesical calculi.¹ A fibroid has perforated into the rectum, and has been discharged per anum. In some cases adhesions with the abdominal wall have formed, and the tumour has been thus discharged.

Considering the frequency of fibroid tumours, it is rare that death follows immediately from their presence. A fatal result, however, may follow from (1) suppuration in the tumour producing death from septicæmia, or a septic peritonitis; (2) uræmia, due to compression of the ureters; (3) direct hæmorrhage; (4) acute simple peritonitis.

PHYSICAL SIGNS; DIFFERENTIAL DIAGNOSIS.

The physical signs of fibroid tumours are usually so well marked that diagnosis is easy. In certain cases, however, diagnosis is very difficult; and when inflammation is superadded, certainty is impossible. Physical diagnosis is best considered under two heads: *a.* of small fibroid tumours, up to the size of a walnut or egg; *b.* of larger ones, which rise up as distinct tumours into the abdomen.

a. OF SMALL FIBROID TUMOURS.

1. Pediculated *submucous* fibroids should be easily recognised. When they are small and not projecting through the os, we have to dilate the cervix to ascertain their presence and attachment; when larger and projecting into the vagina, they may readily be mistaken for inversion of the uterus. On sweeping the finger round the base, we recognise the commencement of the cervical canal unless the polypus be adherent at its neck leading to obliteration of the canal (*v.* fig. 224). Further, the bimanual or rectal examination shows the fundus uteri to be in its normal position.

2. Small *interstitial* fibroids when situated *low down* and causing bulging of one lip of the cervix, give rise to difficulty; owing to the great enlargement of one lip, the os is displaced to the other side and

¹ M'Clintock—*Dub. Quart. Jour.*, Feb. 1868.

its form altered to that of a mere slit which easily escapes observation. Such cases have been occasionally mistaken, even by the most experienced for inversion. This mistake is prevented by examination per rectum. Further, the sides and base of the tumour must be carefully scrutinised to discover the os; when this is found, the sound will show the position of the uterine cavity.

3. *Interstitial* fibroids placed *high up* in the uterus, or small *subserous* ones with a *broad base* of attachment, often escape detection. To ascer-




FIG. 240.

CASE OF TWO-AND-A-HALF MONTHS' PREGNANCY ASSOCIATED WITH TWO LARGE FIBROID TUMOURS—one in the anterior, the other in the posterior wall. The uterus and tumours were removed by Laparotomy (*Barnes*).

tain their presence we proceed as follows. Pass the sound; this defines the course of the uterine canal and position of the fundus. Now make the bimanual examination with the sound, as represented in fig. 90; the finger in the anterior fornix detects the thickening of the anterior wall, produced by a small fibroid. Now steady the sound with the left hand, and pass the forefinger of the right hand into the rectum so as to feel the sound lying in the uterus. Should there be a fibroid in the

posterior wall, the finger recognises an unusual thickness of tissue between it and the sound. Carry the sound, firmly grasped by the left hand, towards the symphysis, so as to bring the fundus better within reach of the rectal finger; and, by moving it from side to side, ascertain whether the tumour is intimately connected with the uterus so that it moves along with it. From their being largely composed of fibrous tissue, these tumours are firmer than the uterine wall; the *localised hardness*, therefore, helps us in recognising them.

Small fibroid tumours, when submucous or interstitial, require to be diagnosed from chronic metritis,
 early pregnancy,
 ante- and retro-flexion.

Differ-
 ential
 Diagnosis
 of Small
 Fibroid
 Tumours.

When subperitoneal and pediculated from—

enlarged Fallopian tube or ovary,
 tumour or inflammatory collection in the broad ligament.

In *chronic metritis* the uterus is not globular but flat, and the enlargement is equable; the uterine canal is patulous; the os is everted, and shows catarrhal patches. We must remember that chronic metritis is occasionally present along with a fibroid tumour.

In *early pregnancy*, the uterus is soft and elastic: the cervix is generally softened, while in fibroids it remains hard. Pregnancy, however, may occur in a uterus which is already the seat of a fibroid tumour (fig. 240); and in such a case the diagnosis becomes certain only after the uterus is considerably enlarged. The possibility of pregnancy must specially be kept in mind here, as we involuntarily think of using the sound to aid in detecting fibroids.

Anteflexion is closely simulated by a fibroid in the anterior wall; a body is felt in the anterior fornix, continuous with it, but separated by a groove. Similarly, a fibroid in the posterior wall has all the characters of the *retroflexed fundus*. Examination by the sound (*v.* fig. 199), and especially by the sound plus the bimanual, clears up the case.

Enlarged *Fallopian tube*¹ or *ovary* may closely resemble a pediculated subserous fibroid; they are not so firm and sharply defined, nor do they move so rigidly with the uterus. In the former also there are the history and symptoms of tubal disease. *Inflammatory collections* in the broad ligament are recognised by their history, the fixation of the uterus, and the changes they undergo; but *solid tumours* there cannot be diagnosed from pediculated fibroids except by exploratory incision.

b. OF LARGE TUMOURS.

When the tumour extends into the abdomen, we proceed with the systematic examination as described at page 102.

¹ Horrocks discusses this point in differential diagnosis in the *Brit. Med. Journ.*, 1886, i., pp. 441, 586, 821.

Diagnosis
of Large
Fibroid
Tumours.

Palpation.—The tumour has a well-defined outline, and a firm solid consistence. It is intimately connected with the uterus; this is best ascertained by laying hold of the cervix with the volsella, when the cervix will be found to move along with the abdominal tumour. Subserous fibroids have a certain range of free movement depending on the length of the pedicle. In soft fibroids, there may be intermittent contractions. *Percussion.*—The note is absolutely dull, unless intestines come between the tumour and the abdominal wall. *Auscultation.*—The uterine souffle is heard most distinctly at the sides, sometimes all over the tumour. As the uterine souffle simply means enlarged uterine arteries, there is no souffle when these are not enlarged; hence it is absent in subserous fibroids with a small pedicle. *Vaginal examination.*—Should the tumour be large and lift the uterus into the abdomen, the cervix will be high up; or it may be displaced in various ways, according to the position of the tumour; it has a firm consistence. *Bimanual.*—With pediculated subserous fibroids, the uterus is felt distinct from the tumour; with interstitial and submucous, we simply feel a large mass continuous with the cervix. *The Sound.*—This should not be used till all possibility of pregnancy has been excluded. In doubtful cases, we wait three or four months till the positive signs indicative of pregnancy should have had time to develop. From the use of the sound we learn (1) the length, (2) the direction of the uterine cavity. The length of the cavity is always increased in submucous, and generally in interstitial, but not in subserous tumours; it may measure six or eight inches. The direction of the canal is often tortuous in submucous tumours; hence the passage of the sound is difficult, sometimes impossible. We feel that the sound goes so far and then catches on a hard projection. In such cases, a soft (No. 8) bougie is very useful, as its flexibility allows it to pass the obstruction.

Differ-
ential
Diagnosis.

Large fibroid tumours require to be diagnosed from—

Advanced pregnancy,
Ovarian tumours,
Extra-uterine gestation,
Hæmatocele and inflammatory deposits.

In *advanced pregnancy* the uterus is of softer consistence, and shows ballottement—the indication of a solid within a fluid; further, we can feel the parts of the fœtus. It becomes occasionally harder under the hand, especially if we make the patient change her position; this *variation in consistence* is a most valuable diagnostic, as it is rarely present in fibroid tumours. We hear the uterine souffle and, unless the child be dead, we hear in addition the *fœtal heart*; the possibility of the child's being dead should always be kept in mind. On vaginal examination, there is discoloration of the vaginal walls with free secretion; the cervix

is softened. There is usually amenorrhœa corresponding in duration to the size of the uterus.

The diagnosis is not so easy as it appears on paper; witness a case¹ in which abdominal section was about to be done in a case of four months' pregnancy, which was not recognised, on the most careful examination, until the patient was under the anæsthetic. Such a case shows the necessity, in doubtful cases, of anæsthesia even for examination.

Ovarian tumours are soft and elastic; small ones may be firm. There is no uterine souffle. They only give rise to difficulty in diagnosis when they have become adherent to the uterus, and move along with it. It is sometimes impossible to diagnose between them and cystic fibroid tumours (*v. Fibro-cystic Tumours*).

Extra-uterine gestation presents great difficulty in diagnosis, especially when the gestation is in an undeveloped horn of the uterus. This condition may so closely simulate a fibroid that it may not be diagnosed till Abdominal Section has been made (*v. p. 296*). But we delay its consideration till the chapter on that subject.

In *hæmatocele and inflammatory deposits* we have the history of the attack to guide us. It may be impossible to form a diagnosis on first examination; but after watching the case for a few weeks and noting any change in the deposit in addition to ascertaining its precise situation, we can form a diagnosis. Pelvic peritonitis frequently occurs round a subperitoneal fibroid, or any fibroid producing pressure; and in such a case it is impossible to diagnose between the tumour and the effusion round it. Many cases reported of gradual absorption of a fibroid tumour under treatment were probably cases of mistaken inflammatory exudation.

PROGNOSIS.

In forming our prognosis we must take into account (1) the site of the tumour in the uterus, most favourable when subserous; (2) its position in the pelvis, whether low down and likely to become wedged within it; (3) the symptoms already present, of which *hæmorrhage* is the most important; (4) rapidity of growth, which by itself rarely forms a reason for interference. Though (as already said) they are rarely dangerous to life, they may cause the patient many years of suffering from which she only finds relief at the menopause.

¹ *Brit. Med. Journ.*, 1886, ii., p. 474.

CHAPTER XXXVII.

FIBROID TUMOURS OF THE UTERUS: TREATMENT.

LITERATURE.

Atlee—The treatment of the fibroid tumours of the uterus: Internat. Med. Cong. Trans., Sept. 1876. *Bantock*—British Medical Journal, 1881, p. 426. *Bigelow*—Gastrotomy for Myofibroma of the Uterus: Amer. Jour. Obstet., Nov. and Dec. 1883, Jan. to March 1884. *Duncan, Matthews*—Diseases of Women: London, 1883, p. 289. *Fritsch*—Sechzig Fälle von Laparomyomotomie mit epikritischen Bemerkungen über die Methoden dieser Operation: Volk. Samml., No. 339. *Greenhalgh*—On the use of the actual cautery in the enucleation of fibroid tumours of the uterus: London Med. Chirurg. Trans., Vol. lix. *Hegar u. Kaltenbach*—Die operative Gynäkologie: Stuttgart, 1881, S. 416. *Hofmeier*—Die Myomotomie: Stuttgart, F. Enke, 1884. *Keith*—Surgical treatment of tumours of the abdomen: Edinburgh, 1885. On Supra-vaginal Hysterectomy, with remarks on the principle of the Extra-peritoneal method of treating the pedicle: Brit. Med. Journ., Dec. 8, 1883. Thirteen cases of Hysterectomy, with remarks on Carbolic Acid Spray in Abdominal Surgery: Brit. Med. Journ., Jany. 31, 1885. *Leblond*—Traité élémentaire de Chirurgie gynécologique: Paris, 1878. *Martin*—Pathologie und Therapie der Frauenkrankheiten: Wien und Leipzig, Urban und Schwartzberg, 1887. *Palmer*—Laparotomy and Laparo-Hysterotomy, their indications for fibroid tumours of the uterus: Americ. Gyn. Trans., 1880, p. 361. *Pozzi*—Traité de Gynécologie: Paris, 1897, p. 287. *Schroeder*—Handbuch der Krankheiten der weiblichen Geschlechtsorgane: Leipzig, F. C. W. Vogel, 1886. *Simpson, A. R.*—The treatment of fibroid tumours of the uterus: Contributions to Obstetrics and Gynecology, Edinburgh, 1880. *Sims, Marion*—On intra-uterine fibroids: New York Medical Journal, April 1874. *Thornton*—Cases of Hysterectomy, with remarks on the value of the Carbolic Acid Spray in the operation: Brit. Med. Journ., May 23, 1885. Article—"Hysterectomy" in Allbutt and Playfair's System of Gynecology: London, 1896. *Wells, Spencer*—On Ovarian and Uterine Tumours, etc.: London, 1882. See also references in the text.

THIS is best considered under the heads of medical treatment, including that by electricity, and surgical treatment.

MEDICAL TREATMENT.

Under this head we include the administration of such medicines as ergot and hydrastis canadensis, and the use of electricity.

There is no medicine which acts immediately upon fibroid tumours so as to cause disintegration and absorption. We have, however, a very important remedy in *ergot of rye*; the beneficial effects of which have been

brought forward by Hildebrandt,¹ and by A. R. Simpson, whose paper on the treatment of fibroids may be consulted for illustrative cases.² It acts beneficially in two ways—by checking their nutrition through diminishing the amount of blood circulating to them, and by favouring their pedunculation and expulsion; these are both due to its action on the unstriped muscular fibre of the walls of the uterus and coats of the blood-vessels.³ Success in its use depends, according to Simpson, on securing that the preparation of ergot used be active, that it be properly administered, and that the case be a suitable one. The formula for the preparation which he recommends is—

R Ergotini ℥ii.
Aquaë ℥vi.
Chloral-hydratis ℥ss. M.

Three grains of ergotin are contained in twelve minims of the fluid, which is a good medium dose. Chloral is added to make the solution keep; but even with this it becomes after some weeks unfit for use, and should therefore be made up repeatedly and in small quantities.

Adminis-
tration of
Ergotin
Subcu-
taneously.

It is administered with the ordinary hypodermic needle. Care must be taken that the syringe contains no air; this is best secured by holding it with the needle upwards and squirting out some of the liquid. The injection is made in the gluteal region, which is readily done when the patient is lying on her side; and on the right and left sides alternately, so as to diminish the frequency of punctures in the same region. Insert the needle vertically and plunge it rapidly *deep into the muscle*, the point entering to the depth of from an inch to an inch and a half; now empty the syringe, and quickly withdraw the needle. After use, remember to cleanse the needle with water and to replace the wire in it. The patient soon becomes accustomed to the prick of the needle, and, if it be entered deeply into the muscle, there is little fear of local suppuration; after three years' experience we have seen this in but one case, and this was probably due to a bad preparation of the solution. For the first few weeks the injections may be made twice a week, afterwards only once a week. The treatment is continued for several months until its effect is seen in diminution of the size of the tumour or, at least, of the hæmorrhage from it. The suitable cases are those in which the tumour is intra-mural or submucous; "it must be surrounded by layers of muscular fibre, sufficiently developed to be capable of being excited to contraction."

When the patient cannot be seen frequently by a physician, a friend or a nurse should be instructed how to apply the needle. Ergotin can also be administered in the form of pill, suppository (4 grs. in each) or liquid extract (30 drops thrice daily). When given by the mouth, however, it does not act so quickly or surely as when given hypodermically.

Hydrastis Canadensis,⁴ fifteen minims to one drachm of the tincture or up to four drachms of the liquid extract, is now being used instead of

¹ Berl. klin. Wochenschrift, 1872, No. 25.

² Dobronrawow gives two cases in which size of tumour distinctly diminished—*Centralb. f. Gyn.*, 1886, S. 16.

³ Ringer—*Brit. Med. Journ.*, Jan. 19, 1884.

⁴ Rutherford gives five cases treated by it—it controlled hæmorrhage, but had no effect on size of tumour—*Brit. Med. Journ.*, 1888, ii., p. 123.

ergot; it does not disturb the digestive system by causing constipation as ergot sometimes does.

Bromide
of Potas-
sium in
Fibroids.

Bromide of potassium was recommended by Sir J. Y. Simpson, who believed that it had a marked influence in checking the growth and even in reducing the size of fibroid tumours. Being a nervine sedative, it is useful in cases where the only symptoms are discomfort from the presence of the tumour or neuralgic pain. As a prolonged use of the bromide is generally necessary, small doses (ten grains, three times a day) should be administered.

When the patient can afford it, benefit is undoubtedly derived from a course of treatment of *mineral waters* (such as those of Kreuznach) as recommended for chronic metritis.

In the case of growing tumours, keeping the patient on a low *non-stimulating diet* is beneficial; the full diet and free use of stimulants, to which a patient inclines to make up for the loss of blood, rather favour the growth of the tumour.¹

The symptoms due to the weight of the tumour may be relieved by *artificial support*. Thus patients with a small fibroid often derive great benefit from wearing a Hodge pessary; the discomfort of a large abdominal tumour is materially lessened by wearing a broad flannel bandage.

When the tumour nearly fills the pelvis and is beginning to press injuriously upon the bladder and rectum, we should, when possible, *push it up out of the pelvis* into the abdomen; this is done before the occurrence of pelvic peritonitis, which may hopelessly bind it within the pelvis. The most favourable case for this manipulation is a subserous fibroid with a distinct pedicle.

TREATMENT OF FIBROIDS BY ELECTRICITY.

More than twenty years ago, Tripiér of Paris treated uterine fibromata with Faradisation, and as far back as 1867 Althaus wrote in the *British Medical Journal* on the electrolytic treatment of tumours;² while in America in 1870, Cutter³ began to use galvanism for the treatment of fibroid tumours. It is, however, to Apostoli that the credit is due of elaborating the electrical treatment of fibroids and bringing it prominently forward before the profession.⁴

¹ See J. Knowsley Thornton on the Treatment of Uterine Fibro-myoma—*Lancet*, 1886, ii., p. 811.

² See letter by Althaus in the *British Medical Journal*, 1887, i., p. 1364.

³ *Amer. Journ. Obs.*, 1888, p. 384.

⁴ In his paper read at the Dublin meeting of the British Medical Association in 1887, "On the Treatment of Fibroid Tumours of the Uterus by Electricity with Observations and Complete Statistics of all the Cases so treated from July 1882 to July 1887."—*Brit. Med. Jour.*, 1887, ii., 699. In the *Lancet*, April 6, 1889, he records 278 cases of fibromata or hypertrophy of the uterus treated by his method, and says as to results, "I can affirm that when there has been no negligence, and my advice has been fully acted upon, ninety-five times out of a hundred permanent benefit has been acknowledged." His *Travaux d'Electrothérapie Gynécologique*: Paris, 1894, contains translations of the chief contributions to the literature of this subject in different countries. While these all agree as to the benefit derived from checking hæmorrhage and especially improving nervous tone, its effect as regards diminution of the tumour is only occasional, and its disappearance exceptional. As to its possible harmful effects on reproduction, he has published eighty cases in which pregnancy followed—Suites éloignées du traitement électrique conservateur en gynécologie; grossesses consécutives: *Arch. d'électr. méd. Bordeaux*, pp. 233, 282.

The technique will be more fully described in the chapter on Electricity in the Appendix. Here we need only say that the internal electrode consists of a platinum rod the thickness of a uterine sound, sheathed in a vulcanite tube except over the portion within the uterus. The external electrode consists of a pad of clay laid on the abdomen, having a copper or leaden plate connected with the battery wire. The internal electrode is usually negative unless hæmorrhage is the chief symptom, in which case it is made positive on account of the hæmostatic action of that pole. The current strength used varies from 70 to 100 milliamperes for the first application, increased afterwards to 200 or even 250 milliamperes.

Electricity has now come to have a recognised place in the treatment of fibroids although there is still great divergence of opinion as to its value. It is a noteworthy fact that Keith, who had as great success as any operator in the removal of fibroid tumours by abdominal section, said—

“Apostoli’s treatment puts a woman with a fibrous tumour who suffers much into the position of a woman with a fibrous tumour who does not suffer or may be even unaware of its presence. It does not bring about the disappearance of the tumour, or it does so very rarely, but size is lessened more or less—one-half, one-third, two-thirds. . . . What I now plead for is, that for a time all bloody operations for the treatment of uterine fibroids should cease, and that Dr Apostoli’s treatment as practised by him should have a fair trial.”¹

The diminution in size of the tumour of which Keith speaks has not, however, been found to be so constant a phenomenon that it can be counted on. Dr Milne Murray, who has given much attention to electricity in gynecological therapeutics and the improvement of the operative technique, writes as follows²:—

“It is not alleged that tumours are necessarily dispersed or materially diminished in bulk by electrical treatment, however long or energetically carried out; that both these events happen from time to time is no doubt true, but the symptomatic cure which is claimed as the aim and result of this treatment does not depend on the disappearance or even on a considerable diminution of the tumour. To those who have had even a moderate experience of this method, it is known that a tumour which was a menace to life may cease to give any inconvenience without undergoing any appreciable diminution in size.”

So also Steavenson, who had charge of the Electrical Department of St. Bartholomew’s Hospital, says³—

“In my paper referred to (St. Bartholomew’s Hospital Reports), I have said that ‘compared with other methods it is probably the best short of actual operation.’ I have admitted that the results are not so brilliant as we could have wished, or as we were led to hope they would be. All the palliative modes of treatment of uterine fibroids are

¹ *Brit. Med. Jour.*, June 8, 1889.

² “The Electrical Treatment of Diseases of Women” in Allbutt and Playfair’s *System of Gynecology*: London, 1896, p. 326—a monograph specially valuable for its clear description of the necessary appliances, practical directions as to their use, and well-balanced opinion as to the results, claiming neither less nor more for this treatment than facts warrant.

³ *Lancet*, April 6, 1889.

eminently unsatisfactory, and the profession would have hailed with delight any mode of treatment that would have promised a cure. This certainly electricity does not accomplish, at any rate with tumours of any size; but there is no doubt that in the majority of cases the symptoms are relieved, and one of the most troublesome that yields to electrical treatment is that of hæmorrhage. Improvement will also take place under the administration of ergot and by the imbibition of the iodo-bromine waters of Kreuznach and Woodhall Spa. . . . It certainly is a question whether in their case (*i.e.* hospital patients) the advantage obtained by the electrical treatment is sufficiently great over other modes of treatment as to call for the expenditure of the time and trouble necessary for carrying it out."

Results of
Electrical
Treatment
of Fibroids.

The value of this method was discussed at the British Medical Association in 1889,¹ and also at the New York Academy of Medicine,² and the Berlin Obstetrical Society³ in the same year, in all of which discussions there were as many opponents as advocates of the method. While exceptionally, diminution and even disappearance⁴ of the tumour has been noted, the application of electricity to fibroids is in great measure a *treatment of symptoms*, and hence it finds its place under the medical rather than the surgical treatment of these tumours.

For recent opinions and results of cases the following papers may be consulted.

P. F. Mundé—My recent experience with electricity in Gynecology: Amer. Jour. Obstet., June 1890. *W. Fraser Wright*—Gynecological Cases treated by electricity in Professor Simpson's Clinique: Edin. Obs. Trans., 1889-90, p. 58. *Engelmann* records nine cases with, in some instances, diminution of the tumour, but thinks he has got better results from ergotine than baths (Wien Klin. Woch., 1890, No. 27). *Massin* gives the result of the treatment of the twenty-three cases in Slavjansky's Clinique. (See French translation of his paper in Apostoli's *Travaux d'Electrothérapie*, p. 356.) *Murray, R. Milne*—(twenty-four cases of fibroids) On the treatment of pelvic disease by electricity, with table of forty-five cases grouped according to symptoms: Edin. Obs. Trans., 1889-90, p. 120. *Mandl and Winter*—Report on ninety-four cases treated by electricity in Chrobak's Clinique, of which seventeen were fibroids. Zur gynecologischen Therapie: Wien Klin. Wochen, 1891, S. 955, 983; and 1892, S. 50, 70. *Kellogg*—(sixty cases of fibroid tumours) Summary of my personal experience with electrolysis in treatment of fibroid tumours: Jour. Amer. Med. Assoc., 1892. *Parsons, J. T.*—Twenty cases of fibroma and other morbid conditions of the uterus treated by Apostoli's method: Lancet, 1892, Vol. i., p. 467. *Haultain, F. W. N.*—(twenty-two cases of fibroids) The electrical treatment of uterine fibroids and subinvolution: Edin. Obs. Trans., 1893-94, p. 210. *Corson, E. R.*—(nine cases of fibroids) The Apostoli treatment of uterine fibroids: Amer. Jour. Obst., 1895, Vol. ii., p. 341.

SURGICAL TREATMENT.

Surgical
Treatment
of Uterine
Fibroids.

While the surgical treatment of fibroid tumours⁵ includes the removal of pediculated submucous tumours, as polypi, and the enucleation

¹ Brit. Med. Jour., Oct. 19, 1889. ² Med. News, Jan. 25, 1890. ³ Cent. f. Gyn., 1889, p. 275.

⁴ As by Milne Murray, in which a fibroid as large as a six months' pregnancy disappeared after forty applications (Edin. Obs. Trans., 1893-94, p. 118); and by Parsons who records a similar case (Brit. Gyn. Jour., 1892-93, p. 338). So also Kjoergaarch records disappearance of a tumour the size of a hen's egg: Apostoli's *Travaux d'Electrothérapie*, p. 585.

⁵ *Curetting of the Uterus* is sometimes of service in checking hæmorrhage from fibroid tumours, and might therefore almost have a place under surgical treatment. Its good results are, however, due to its effect on the endometritis which accompanies these tumours, and it can scarcely be considered a treatment of the tumour itself. From the distortion of the uterine cavity curetting is often a difficult operation and not without risk through sepsis from injury of the tumour.

of small interstitial ones, the chief interest gathers round the treatment of those larger interstitial tumours, the removal of which implies the amputation of a portion of the uterus. The development of the operative technique for this is one of the most interesting subjects in operative gynecology. The surgical problems which the treatment of these large tumours present, and the various solutions that have been suggested would require a separate treatise. The multiplicity of operative methods shows that we have not reached a final solution, and makes it impossible to describe the surgical treatment of fibroid as succinctly as that of ovarian tumours. A brief historical sketch and a description of the nature of the problems will put the student in a better position to understand the various operations of which we shall select only the most important.

The operative treatment of fibroids has followed step by step that of ovarian tumours, which here led the way. The abdomen was first opened for a fibroid, by mistake for an ovarian, by Lizars in 1825; but at that time the extirpation of these tumours was not thought of. Success however in the removal of ovarian tumours encouraged the operator to proceed with fibroids, and in 1837 Granville successfully removed a pediculated subserous tumour. The first deliberate hysterectomy for an interstitial fibroid was done by Kimbal in 1855, but it was Koeberlé and Péan who, by their operative methods and success, secured for the operation a recognised place. How to deal with the pedicle is, as we shall see, the problem, and Koeberlé's *serre-nœud* was the first successful solution. Historical Sketch.

The difficulty in dealing with large fibroid tumours is due to the peculiarity of the blood supply, the presence of the peritoneal cavity, and of the uterine cavity communicating with the vagina. Difficulties in operating:

From Plate III. it is evident that the blood supply comes from two sources, the ovarian and the uterine arteries, and that while the former runs in the upper part of the broad ligament, the latter hugs the side of the uterus. The whole blood supply can thus be cut off by a single loop thrown round the isthmus uteri, gathering in the broad ligaments (Pl. XI., fig. 1a). Should the tumour be large or opening out the the ligaments, the latter must be cut through to make a track for the loop that it may constrict the uterus efficiently. This means that the broad ligaments must be tied separately so as to control the ovarian artery (Pl. XI., fig. 1b); the loop of the *serre-nœud* constricts the uterine. arising from Blood-supply;

The judging as to whether this separate ligature of the ligaments is necessary and the best mode of doing it, is of importance in operating. For ligaturing the broad ligaments, and also the pedicle (by Zweifel's method), the chain ligature is useful (see fig. 241). A continuous suture is carried through in loops by a needle on fixed handle: after the loop is carried through, the thread is caught in forceps always on the side of the needle-point farthest from the free end of the thread; the loops are subsequently divided, the

right ends to tie (1 with 1', 2 with 2') can be told by pulling on the thread, and before tying care is taken that the ligature interlocks with the next one (1' with 2).

Further, the bulk of the uterine stump makes the control of the uterine arteries difficult. As the tissue yields, the control slackens. Hence the significance, as improvements in technique, of the *serre-nœud* which can be tightened up, of the elastic ligature, and of the tying of the pedicle of the uterine stump in various portions, or of the arteries by themselves. Hence also the advantage of the extra-peritoneal treatment; in which the pedicle is kept under observation.

from
Peritoneal
cavity ;

The presence of the peritoneal cavity gives rise to peculiar difficulties. Although the peritoneum is not dreaded now as formerly, it is still very much in evidence as is seen from the continued battle between extra- and intra-peritoneal methods, and the more recent appearance of the third "retro- or sub-peritoneal." To drop the pedicle into the abdomen and close it, as in ovariectomy, is the surgical ideal; but the results have made many operators regard it as unattainable.¹ In addition to the risk of concealed hæmorrhage, there is the danger of infection of the




FIG. 241.

MODE OF PASSING CHAIN LIGATURE. The first loop has been divided and interlocked; the second is in the forceps' bite, but not yet divided; the third is running in the needle.

peritoneum by the stump. If the latter is to be kept outside the abdominal wound that bleeding may be seen and controlled, the peritoneal cavity must be carefully shut off. A collar of parietal peritoneum must be carefully adapted round the neck of the pedicle. Further, a trough or gutter must be kept open above this, so as to keep infective matter from burrowing past the peritoneal collar. It is evident also that the drier the stump is, the better: astringent powders are the best dressing. The term "extra-peritoneal" is restricted to the stitching of the stump in the abdominal wound. Another mode of keeping it outside the peritoneum is to leave it lying in the pelvic floor, but to suture the pelvic-floor peritoneum over it, so as to keep it retro-peritoneal, or sub-peritoneal. The stump has also been turned into the vagina, either through the anterior or posterior fornix; this is like the retro-peritoneal method, but means freer drainage.

It is possible, however, that too much has been made of the position

¹ In spite of the remarkable successes of Fritsch, Brennecke, Leopold and von Ott by this method, the extra-peritoneal method has more advocates still.

of the stump relative to the peritoneal cavity, and that a more important point is its vitality; so that if it be ligatured in such a way as not to impair this seriously, it may be safely left in the peritoneal cavity.

Another source of danger is the presence of the cervical canal in the heart of the stump communicating with the vagina, and furnishing a channel for septic infection. The cleansing and draining of this canal has given rise to much thought, but probably the natural plug of mucus is the best safeguard against sepsis. from Sepsis through Cervix.

The problem of the stump has been so great, that of recent years the removal of the cervix with the body of the uterus has found many advocates. Vaginal hysterectomy of course implies that there is no stump, and to this fact its success is in part due. Whether abdominal hysterectomy may not in the end dispose of the stump problem by abolishing it, *i.e.*, by in every case cutting out the whole uterus, only time will show.

The various operations aim at the removal of the tumour alone, or the tumour with part of the uterus, or of the whole uterus. Further, this may be done by the vaginal method, by the abdominal method, or by the combined abdomino-vaginal method. Hence there is a multiplicity of operations, and some confusion of terms, which puzzles the student. Operations performed.

In considering the operative treatment of fibroid tumours, it will be convenient to group them under the three above-mentioned methods.

A. OPERATIONS BY THE VAGINAL METHOD.

1. The removal of a fibroid tumour which has become *pediculated* is a comparatively simple operation, and will be considered under the treatment of fibrous polypi. Removal of Polypi.

2. The removal of a non-pediculated submucous fibroid by *enucleation* is an operation which has a very limited range. The danger of sepsis, and the introduction of other methods, has restricted its use to the few cases for which it is the only resource, such as the cervical fibroid seen in fig. 237. Enucleation.

After the cervix has been dilated, when this is necessary, the mucous membrane over the tumour is incised with a bistoury or thermocautery, which lessens the risk of sepsis. The tumour is then shelled out of its bed—different scoop-like curettes have been devised for this by A. R. Simpson, Thomas and Pozzi. After as much of the tumour as possible has been removed, the cavity is douched with an antiseptic, and packed with iodoform gauze. The risk of sepsis is great, from the sloughing tissue and the abundant lymphatic supply of the uterus. The mortality, according to Kleinwächter, is 15 p. c.;¹ and it is an open question

¹ Twenty-two out of 147 cases collected: *Wien. med. Presse.*, 1887, No. 42. On the other hand, Leopold records 28 operations without a death: *Archiv f. Gyn.*, 1890, Bd. xxxviii., Hft. 1. And Ascher 10 with one death: *Zeits. f. Geb. u. Gyn.*, 1890, Bd. xx., Hft. 2, S. 307.

whether enucleation should not be followed by hysterectomy. The tumour in fig. 237 is a case that might be treated by enucleation.

Colpotomy
for Fib-
roids.

3. *Colpotomy*, in which the tumour is got at by an incision in the fornix, is of very limited application to fibroids. It will be described in the Appendix.

Vaginal
Hysterec-
tomy.

4. *Vaginal Hysterectomy*, or removal of the whole uterus per vaginam has, during the last ten years, received a recognised place in the treatment of bleeding fibroids. While the first operation was done by Kottmann¹ in 1881, to Péan belongs the honour of having established the operation by skilful technique and phenomenal success.² At first sight it appears that only very small tumours can be thus treated, but by "morcellement" fibroid uteri up to the umbilicus may be removed.³

For tumours of considerable size, however, the abdominal or combined will always compete successfully with the vaginal method; and for small tumours, a more conservative surgery has prevailed in this country.⁴ Extirpation of the uterus for small fibroids, can never rank with its extirpation for malignant disease. Cases in which this treatment is justifiable, occasionally present themselves, but they are the exception. We shall therefore defer the consideration of vaginal hysterectomy to the operative treatment of cancer, where it has an unquestioned place.

B. OPERATIONS BY THE ABDOMINAL METHOD.

Operations
by Abdo-
minal
Method.

Here the tumour alone may be removed, or a portion of the uterus with it. When the *whole* uterus (including the cervix) is removed, the vagina is necessarily cut into, which makes it practically an operation by the combined abdomino-vaginal method, under which head we shall consider it.

Terms
employed.

If a pediculated subserous fibroid is removed or a tumour enucleated from the uterine wall, it is a "myomotomy." When the uterus is amputated (usually through the isthmus) with the tumour, it is a "hysterectomy."⁵ The term is incorrectly used, inasmuch as the whole uterus is not cut out; but it came into use before the removal of the entire uterus was much practised, and so became synonymous with "supra-vaginal amputation." The operation in which the cervix also is removed is now called "complete abdominal hysterectomy" or "pan-hysterectomy," to distinguish it from "hysterectomy," which is usually partial.

¹ *Corresp. Schweiz. Ärzte.*, Jan. 1882, No. 2, p. 42.

² In his most recent report, from 1890 to 1895, he has done 248 vaginal hysterectomies with four deaths, a mortality of 1.6 p. c.

³ Ségond reports on 66 cases of fibroid uteri reaching the umbilicus thus extirpated, with only seven deaths. *Neuv. Cong. Fran. de Chir. : Sem. Méd.* 1895, p. 475.

⁴ The first case recorded was done by Heywood Smith, to relieve pain which interfered with patient's work: *Brit. Gyn. Jour.*, 1893-94, p. 440.

⁵ This term used alone means the abdominal operation; when done by the vaginal method it is a "vaginal hysterectomy."

1. When the tumour alone is removed, the methods vary, according as it is subserous and pediculated, or growing between the layers of the broad ligament or into the cellular tissue, or lying within the substance of the wall. Removal of pediculated subserous Tumours.

In the case of subserous pediculated tumours the pedicle can be treated intraperitoneally as in ovariectomy, *i.e.*, transfixed and ligatured in two portions, though it is desirable, in addition, to bring together with catgut the edges of the peritoneum over the end of the stump; or the extra-peritoneal method, to be presently described, may be adopted.

The second class of tumours demands a more serious operation, implying their enucleation from the peritoneum or cellular tissue. The cavity thus produced may be either sewed up with catgut and the abdominal incision closed;¹ or its margins may be stitched to the open abdominal wound, the hollow being packed with iodoform gauze.² Enucleation by Abdominal Method.

The third condition, when the fibroid is in the substance of the wall,




FIG. 242.

MARTIN'S OPERATION FOR ENUCLEATION OF FIBROID FROM WALL OF UTERUS (*Martin*).
a. Shows uterus with temporary elastic ligature round it; the shaded portion of capsule being the extent of incision in it. *b.* Shows how the hollow in uterine wall is closed by sutures.

gives occasion for two quite distinct methods of operation—enucleation from the wall, and hysterectomy.

2. *Enucleation* of the tumour from the uterine wall with sewing up of the hollow thus produced is an operation introduced by Martin of Berlin. The cases in which it can be done are limited; but, where it is possible, it has the double advantage of being a less serious operation than hysterectomy and not mutilating the uterus.

After the uterus has been exposed by abdominal section and drawn forward into the incision, a temporary elastic ligature is thrown round the broad ligaments; this is not necessary in all cases, as with a mesial incision the bleeding may be but slight. A longitudinal incision is made over the tumour which is shelled out of its capsule; the margins of the cavity are then trimmed with scissors, considerable portions of the muscular wall and all the connective tissue portion of the capsule being sometimes

¹ As in recent cases by Baumgärtner and Veit—*Centralb. f. Gyn.*, Bd. xi., S. 771.

² As in Rokitansky's case—*Ibid.*, S. 839.

excised; and the wound is closed by continuous deep and superficial juniper catgut sutures. The uterine cavity may be opened into during the operation, but if it be disinfected or packed with iodoform gauze¹ (extending down into the vagina for ease of removal) which acts as a drain, it does not affect the prognosis. (v. fig. 242.)

3. The removal of a portion of the uterus with the tumour constitutes hysterectomy, in which there is left a stump of cervix, and more or less of body of uterus (with its cavity cut across), according to the height of the tumour in the uterine wall. Strictly speaking, this is only a supra-vaginal amputation, but the term hysterectomy has come into use and is convenient, if we remember that the whole uterus is not cut out.

ABDOMINAL HYSTERECTOMY FOR FIBROIDS.

Here there are two forms of operative technique according as the pedicle is treated by the extra-peritoneal or intra-peritoneal methods. Though the latter is theoretically superior, the former has, till recently, been the favourite one with British operators, having given splendid




FIG. 243.

KœBERLÉ'S SERRE-NŒUD, with pins for transfixing stump.






results in the hands of Keith, Bantock, Tait, and Thornton. We shall describe therefore in detail the operation with extra-peritoneal treatment of the pedicle.

Extra-peritoneal Method.—The instruments required are those for ovariectomy, except that instead of ovariectomy forceps and trocar, we require some six pairs of long pressure forceps (Spencer Wells) and two Kœberlé's serre-nœuds with iron wires looped at one end (fig. 243). Also two or three long-handled curved needles threaded with silk, for tying the broad ligaments; elastic tubing for temporary constriction of the tumour; a long knife for amputating the tumour; two pins with screw-caps for transfixing the pedicle; and drainage tubes. And for dressing the stump, a powder of the per-sulphate or per-chloride of iron, iodoform gauze with antiseptic wool, binder and safety pins are required.

The patient is prepared as for ovariectomy. Some advise that the bladder be not emptied as it is more easily defined when distended. A

¹ As Fränkel did in his two cases—Ueber die Enucleation submucöser oder intraparietaler Myome von der Bauchhöhle aus (Martinsche Operation), etc.: *Archiv f. Gyn.*, Bd. xxxiii., S. 449.





HYSTERECTOMY FOR FIBROID TUMOURS OF THE UTERUS.

- FIG. 1. Control of Blood-supply: broad ligament embraced by serre-nœud in (a), tied separately in (b).
 FIG. 2. Application of serre-nœud: ligaments clamped next uterus and tied off (ovary cut away at b, left at a).
 FIG. 3. Extra-peritoneal, FIG. 4. Intra-peritoneal, FIG. 5. Retro-peritoneal treatment of stump.

long abdominal incision (sometimes from the ensiform cartilage to near the pubes) is necessary—with care to arrest bleeding, and in going through the peritoneum not to injure the bladder or tumour. Bleeding from an incision in the latter is difficult to control. The hand is now passed round the tumour to discover adhesions, but especially the relation of the broad ligaments and appendages. If the relations admit of the *serre-nœud* wire embracing ligaments and tumour (as in Plate XI., fig. 1*a*), it is thrown round the mass, the ovaries also being included.¹ Where this cannot be done without tension or tearing of the ligaments, the latter are tied as in Plate XI., fig. 1*b*; in which case the wire of the *serre-nœud* is carried through the same puncture as the ligature next the uterus. The *serre-nœud* is now tightened up. The elastic ligature is required for temporary constriction to arrest bleeding during the removal of masses which prevent the pedicle from being defined; and may also be used as a permanent substitute for the *serre-nœud* (fig. 246), in which case it is knotted, and the knot tied up with silk, or clamped in forceps for additional security. Great care is required that no intestine or omentum be caught in, and especially that the bladder be not included. Should it be necessary to divide the ligaments, before applying or tightening up the wire, the broad ligament must be temporarily clamped on its uterine side (Pl. XI., fig. 2). In ligaturing the broad ligaments, the ovaries may be left or removed with the tumour (compare the two broad ligaments in Pl. XI., fig. 2). To allow the wire to hug the uterus a transverse incision may be made through the peritoneum anteriorly, joining the broad ligament incisions, and the peritoneum and bladder stripped off with the finger. This procedure prevents the inclusion of the ureters. The pedicle is now transfixed with two pins immediately above the wire, and the tumour amputated (Pl. XI., fig. 3). During the amputation of the tumour, sponges are packed round the pedicle at the abdominal incision to catch any fluid that escapes. Should there be oozing from the pedicle, the *serre-nœud* is tightened further. In the cut surface of the stump the mucous canal is generally small, with a plug of mucus which is left undisturbed. Should, however, the cavity be large or septic, it should be cleansed with 1 to 1000 corrosive sublimate solution. The abdominal incision is now closed, care being taken to adjust the peritoneum carefully round the neck of the pedicle (Pl. XI., fig. 3). Some recommend the stitching of the peritoneal collar round the stump. But this is not necessary; though it may be desirable to carry the abdominal wound suture immediately above and below the pedicle, also through the peritoneum of the latter, so as to insure co-adaptation. For the same reason where drainage is required the tube should be introduced between the sutures higher up

Abdominal
Hysterectomy for
Fibroids.

¹ Thornton recommends that one ovary be left as producing more rapid recovery, less suffering at the change of life, and preventing the depression consequent on removal of both ovaries.

in the wound, not beside the pedicle. A strip of iodoform gauze is tucked in round the stump to secure drainage; the gutter round the pedicle thus formed is of great importance as allowing the discharges connected with the sloughing of the stump to escape freely. The surface of the stump is rubbed with per-sulphate or perchloride of iron. It must be applied carefully so that it may not touch the margins of the abdominal wound, and thoroughly so as to cause tanning of the stump. The drier the latter is made and kept, the less is the risk of sepsis. Pads are placed under the transfixion needles to keep them off the skin, antiseptic wadding put round and over the stump, and a bandage applied.

After-treatment.

In the after-treatment, if the pedicle is thick, it must be watched carefully for oozing, and the *serre-nœud* tightened up. In some cases, however, the dressing need not be disturbed for two or three days. Then it must be changed and the stump dusted with iodoform and bismuth, or borax. The abdominal wound sutures are removed at the end of a week.¹ During the sloughing of the pedicle (which may take three or four weeks or even longer) it may be necessary to tighten the wire occasionally, and clip away bits of dead tissue. The protracted healing of the wound, and danger of sepsis are the great drawbacks to the extra-peritoneal method.



FIG. 244.
KEITH'S CLAMP
for securing the
Pedicle extra-
peritoneally.

Other
Methods
of Extra-
Peritoneal
Treat-
ment.

Instead of the *serre-nœud*, Keith² used the clamp shown at fig. 244, drawing the parts together with a silk ligature before applying it.

Notwithstanding his brilliant success, he latterly preferred the more conservative electrical treatment to the use of the knife—a fact worth remembering when we read of the enormous number of operations now done in which a low mortality, instead of the well-being of the patient, seems to be the aim of the operator.




FIG. 245.
CINTRAT'S SERRE-NŒUD (Hegar).

The *serre-nœud* of Cintrat has been used with great success by Péan. The pedicle is transfixed with a big needle, and a double iron wire

¹ Thornton advises leaving them for a few days longer than in ovariectomy, as the wound is liable to reopen.

² Of sixty-four cases he had a mortality of 15·7 p. c. for hospital (thirty-eight cases), and only 3·8 p. c. (twenty-six cases) for private practice: *Brit. Med. Jour.*, 1887, p. 1257.

drawn through, which is tightened up and twisted by the ingenious instrument seen at fig. 245.

The elastic ligature, introduced by Kleeberg, proved a distinct advance in the treatment of the extra-peritoneal pedicle as its progressive contraction controls bleeding. Its mode of use was elaborated by Hegar, who transfixed the pedicle with a needle specially constructed to carry in the ligature. Another merit of Hegar's was the emphasis of the peri-pedicular trough (*v.* fig. 246). Pozzi recommends the




FIG. 246.

TREATMENT OF FIBROID TUMOURS BY ELASTIC LIGATURE (*Hegar und Kaltenbach*).

a, Abdominal incision with the stump in its lower angle; only the peritoneum is brought together with the lower sutures, while the upper sutures take in the whole abdominal wall. *b*, Same in section, to show the trough floored by the peritoneum round the stump and the position of the elastic ligatures.

elastic ligature without transfixion of the pedicle, and we have referred to it as an alternative to the *serre-nœud* in the hands of British operators.

Before considering the intra-peritoneal method we must note certain attempts at a compromise between them, in which the stump is kept in the *closed* abdominal incision, or fixed to it. They were introduced by Wölfler¹ and von Häcker,² two pupils of Billroth's. And other modifications have been proposed by Sänger, Fritsch,³ and Kelly.⁴ To control

¹ *Zur Technique der supravaginalen Amputatio Uteri*: Wien. Med. Woch., 1885, No. 25, S. 793.

² *Ibid.*, No. 48, S. 1466.

³ *Op. cit.*, in which he records nineteen cases without any deaths.

⁴ *Amer. Jour. Obstet.*, 1889, p. 375.

bleeding Fritsch ties the uterine arteries, and then closes in the muscular tissue of the stump, as in fig. 247.⁵ The stump itself is stitched against the abdominal incision, the sutures closing, the latter being also passed through the stump. In v. Häcker's method the pedicle is stitched into the incision, so that the technique resembles rather the purely extra-peritoneal operation.

Intra-
Peritoneal
Method.

In the *intra-peritoneal method* the pedicle is tied and dropped as in ovariectomy (Pl. XI., fig. 4). This method was first advocated by Schroeder as the surgical ideal. He drew attention to the risks




FIG. 247.

DIFFERENT MODES OF SUTURING THE PEDICLE SO AS TO CONTROL BLEEDING; the peritoneum is also stitched over the end of the stump in these cases.

1. By transfixion to one side of cervical canal and tying in two parts—as in Leopold's method.
2. By transfixion and tying in three or more parts by the chain ligature (cf. fig. 241) as in Zweifel's method—the chain being continued into the broad ligaments.
3. By ligature of uterine arteries alongside of cervix as in Baer's method.
4. By "Stage suture" as in Schroeder's method: the raw faces, produced by a wedge-shaped incision, are united by a continuous catgut suture beginning deeply and rising in "stages."
5. By ligature of uterine arteries in face of stump and apposition of raw surface (*e, d, c*) by suture (*f*) as in Fritsch's method. *aa*, cervix; *b*, its canal; *e*, peritoneum.

of infection through the cervical canal, and recommended careful apposition by the stage-suture of the muscular tissue of the cervix as well as of the peritoneum over it.

While Tait, Bantock and Thornton have advocated the extra-peri-

toneal method, the intra-peritoneal has been slowly gaining ground in this country. Thus we have four successful cases recorded by Milton,¹ ten by Greig Smith,² three by Heywood Smith,³ twelve with one death by Sinclair,⁴ thirteen with two deaths by Mary Scharlieb,⁵ seven with one death by Donald,⁶ eight with one death by Harrison Cripps.⁷

The two essentials in the operation are the shutting off of the stump from the peritoneal cavity by early adhesion of the peritoneum covering it, and the ligaturing of the vessels in such a way as to control bleeding without affecting the vitality of the stump. Of the two, the latter is the more important, and hence the various modes of controlling hæmorrhage require some notice (see fig. 247). This has been done by the "stage-suture"⁸ (Schroeder) in which the surfaces are brought together by a continuous catgut suture drawing together first the deeper parts, and then the more superficial at successive stages; hence the term. Schroeder also passed a silk suture deeply through the whole stump. Or the stump may be transfixed and tied in two portions, the transfixion being made to one side of the cervical canal. Leopold⁹ who deals thus with the stump, lays stress on the enucleation of all fibroid masses until a pedicle of an inch in diameter is obtained. Or the stump may be ligatured in several portions as Zweifel recommends.¹⁰ He carries a single ligature in loops with a sharp needle through the stump just as it is carried through the broad ligament with a blunt needle. Or the uterine arteries may be ligatured individually as they lie beside the cervix, as Baer¹¹ and others have done. The latter method though the most difficult, is surgically the best. But all of them have given equally good results in different hands. The end in view is to prevent sloughing of tissue.

The careful adaptation of the peritoneum over the stump was emphasised by Schroeder. Zweifel takes a peritoneal flap from the anterior wall of the uterus long enough to cover the whole stump (Pl. XI., fig. 4): while others after transfixing and tying the pedicle stitch the anterior and posterior walls of the broad ligaments together right across the pelvis (Pl. XI., fig. 5). The seclusion of the peritoneal cavity is certainly of consequence, but may be brought about by coaptation without ligature where the stump retracts within the ligaments (already tied at the side, as in Baer's operation). Fear of infection through the cervical canal has led many to cauterise it with the actual cautery or

Control of
Bleeding.

Exclusion
of stump
from Peri-
toneal
Cavity.

¹ *Lancet*, Nov. 29, 1890, Sept. 26, 1891.

² *Brit. Gyn. Jour.*, 1892-93, p. 23.

³ *Brit. Med. Jour.*, 1895, p. 69.

⁴ *London Obstet. Trans.*, 1896, p. 41.

⁵ Brennecke has recently done nineteen cases without a death: *Zeits. f. Geb. u. Gyn.*, Bd. xxviii. Hft. 2.

⁶ *Cent. f. Gyn.*, 1894, S. 617, where he records twenty cases without a death.

⁷ *Cent. f. Gyn.*, 1894, S. 321. He has had remarkable success with this method. Of ninety-two hysterectomies + two myomotomies, only three have died, giving a mortality of 3.2 p. c. Of the last forty-three cases only one died, and that from obstruction independent of the operation.

⁸ Supra-vaginal Hysterectomy without ligature of the cervix, etc.: *Amer. Jour. Obstet.*, 1892, ii., p. 489.

⁹ *Brit. Gyn. Jour.*, 1892-93, p. 54.

¹⁰ *Brit. Gyn. Jour.*, 1893-94, p. 49.

¹¹ *Brit. Med. Jour.*, Oct. 24, 1896.

strong carbolic. But Döderlein's researches showing that the cervical mucus is germ-free, have discounted this dread.

A third mode of treating the stump has been called the *sub-peritoneal* or *retro-peritoneal*, but it hardly merits a separate name (Pl. XI., fig. 5), and will be referred to under operations by the combined method.

The term merely implies that certain operators¹ emphasise the fact that the peritoneal cavity should be shut off from the stump—a fact which all advocates of the intra-peritoneal method, from Schroeder downwards have insisted on. In all the methods described, the stump, though spoken of as intra-peritoneal, is retro-peritoneal, inasmuch as the peritoneum is brought over the end of it.

C. OPERATIONS BY THE COMBINED ABDOMINO-VAGINAL METHOD.

Under this we include the removal of the stump *per vaginam*, after the supra-vaginal amputation has been done by the abdominal method, and also complete hysterectomy.

The former is sometimes called the "mixed method,"² as being a combination of the abdominal and vaginal. After the supra-vaginal amputation has been completed, the cervical stump is removed *per vaginam*, as the bladder is more easily separated, and the base of the broad ligaments ligatured or clamped from the vagina. This double operation leaves the parts in the same condition as after complete hysterectomy by the abdominal method. Péan having removed the uterus and tumour and constricted the pedicle by Cintrat's *serre-nœud*, as in the extra-peritoneal operation, returns stump and wire into the abdomen and closes the incision. He then removes the cervix (constricted by the wire) *per vaginam*, clamping the broad ligaments.³

Complete hysterectomy for fibroids by the abdominal method has only recently⁴ been brought into prominence, and chiefly by Martin of Berlin. In this operation, Thornton proceeds as follows.⁵

Thornton's
Method of
Pan-hys-
terectomy.

The broad ligaments being ligatured they are temporarily clamped on the uterine side and divided between the ligature and clamp. The peritoneum in front of and behind the uterus having been incised and peeled back, the finger is pushed down alongside of the uterus to feel

¹ Maunsell drew attention to the importance of dissecting back a peritoneal flap all round the stump, treating it by what he called "the deperitonised extra-peritoneal method": *New Zealand Medical Journ.*, Sept. 1887. See also *Brit. Gyn. Journ.*, 1891-92, p. 100. James Gough, after transfixing and tying the pedicle, ran a continuous catgut suture across the pelvis, bringing together the anterior and posterior layers of the broad ligament over the stump—"a new method—the intra-abdominal but sub-peritoneal method of disposing of the pedicle!" *Am. Jour. Obstet.*, 1890, p. 372. So also Heywood Smith describes some of the above-mentioned operations as "sub-peritoneal hysterectomy": *Brit. Gyn. Jour.*, 1892-93, p. 23.

² Thus Smyly records eleven cases done by the "mixed method": *Brit. Med. Jour.*, Jan. 12, 1895.

³ Of his first series of cases he lost only one out of fifty, and of his second (1890-94) seven out of 120, or 5·8 per cent.: *Anal. de Gyn. et Obst.*, May 1893, and *Sur les Fibromes Uériques. Gaz. des. Hôp.*, 1895, p. 287.

⁴ The first operation in America was in 1888 by Mary A. Dickson-Jones (see *Am. Jour. Obstet.*, 1896, p. 405), and in England by F. B. Jessett (*Brit. Gyn. Jour.*, 1892-93, p. 71).

⁵ Article on Hysterectomy: *System of Gynecology*, London, 1896, p. 626.

the pulsating uterine artery, which is ligatured by transfixion close to the cervix so as to exclude the ureter. The anterior fornix is pushed up with a sound, which is cut down on behind the bladder, and the incision carried quickly round the vaginal fornices with scissors curved on the flat, and uterus and tumour removed. Bleeding points are quickly seized in forceps and tied with fine silk by ligature or transfixion, the peritoneal edges of the broad ligament being drawn together so as to reduce the vaginal opening. The vagina is cleansed and plugged with iodoform gauze, and the abdominal incision closed, a Keith's drainage tube being stitched in it so as to remove quickly fluid accumulating in the pouch of Douglas. This tube is taken out in twenty-four to forty-eight hours as the vaginal plug now drains satisfactorily; the latter remains for five or six days. An antiseptic pad occasionally changed receives discharge from it; while the urine is removed by a catheter for some days. When the vaginal plug is withdrawn, antiseptic douches are begun and the plug is not renewed.

Doyen¹ has recommended the rapid cutting away of the uterus and tumour without previous ligature. After opening the abdomen, he cuts down on a pair of forceps pushed up in the posterior fornix. The cervix, being thus freed behind, is pulled up through the rent by volsellæ, the mucous membrane of the anterior fornix divided with the scissors, and the bladder separated from below upwards with the finger till the utero-vesical peritoneum is divided. One broad ligament is now seized between the finger and the thumb, so as to control the vessels, and cut away from the uterus; the uterine artery and any other vessels caught in forceps and tied. The other ligament is treated in the same way. The peculiarities of this method are the separation of the bladder from below upwards, which is made possible by bending the cervix on itself upwards through the incision in the posterior fornix; and the division of the ligaments before tying, which admit of more rapid operating.

Martin² uses catgut instead of silk, and sutures the margin of the peritoneal wound to the fornix, the left hand being passed into the vagina to direct the needle. He also cuts first into the posterior fornix, and after ligaturing the base of the broad ligaments, separates the uterus from the bladder. The wound in the pelvic floor is closed by a continuous catgut suture, bringing the peritoneum together. He has a lower mortality since he closed the peritoneum than when it was left open. Although he has had better results since he closed the peritoneal cavity, this does not seem necessary, as is seen from the results of Thornton, Christopher Martin, and Le Bec.

¹ *D'Hystérectomie Abdominale et Vaginale*: Paris, 1892, p. 72. Cases by this method have recently been reported by Edge and J. W. Taylor. *Brit. Gyn. Jour.*, 1897, pp. 305, 498, 526.

² Die totale Extirpation des myomatösen Uterus, etc. *Berlin Klin. Woch.*, 1894, S. 625. *Zeits. f. Geb. u. Gyn.*, Bd. liii., S. 218, 486.

Christopher Martin¹ has had ten successful cases of pan-hysterectomy for fibroids, and other operators have had like success.²

Le Bec's
Opera-
tion.

In Le Bec's operation³ the vaginal fornix is included in the ligature of the base of the broad ligament. After the upper portion of the ligaments has been tied in the usual way outside the ovary (see right broad ligament of Pl. XI., fig. 2; also stump in fig. 5) an incision is made in the anterior and posterior fornices by cutting down on a pair of forceps pushed up through the vagina and partially opened so as to make the fornix tense. The forceps are pushed through the one wound, and ligature seized, and then carried with them up through the other wound; a loop is thus formed, including the fornix, and tied from above. All bleeding being thus controlled the tumour is cut away, the stump of the cervix divided mesially, and the halves clipped off. The ligatures tying the base of the ligaments are then drawn down through the vagina, so that the two pedicles lie within it, as in Pl. XI., fig. 5, while the peritoneal cavity is cut off above by catgut sutures.⁴ Hence the stump is infra-peritoneal or retro-peritoneal.

*Results of these Various Methods for the Surgical Treatment of
Fibroid Tumours.*

Choice of
Operations.

In the last edition of this Manual we gave a *précis* of the results of various operators to bring out the mortality of the operation and the merits of various methods.⁵ Individual cases vary to such a degree in the gravity of the operation, according to the situation and size of the tumour, the condition of the patient, and difficulties which may arise in the course of the operation and complications after it, that such statistics are of very little value. Nor can it be said that one method is better than another without a specific case before us. The extra-peritoneal is, generally speaking, more rapid than the combined, and therefore in patients reduced by hæmorrhage might be preferable. Against a more rapid operation, however, must be set the disadvantage of a more protracted convalescence. So that the balance hangs level between the two. Nor can the argument of the surgical ideal be unduly pressed as

¹ *Edin. Med. Jour.*, March 1896; *Brit. Med. Jour.*, 24th Oct. 1896.

² So Delagénière records twenty cases with one death: *Bull. et Mem. de la Soc. de Chirurg.*, 1895, t. xxi., p. 487. *Hystérectomie abdominale totale par ligatures—Annal. de Gyn.*, 1895, xliv., p. 356.

³ *Brit. Med. Journ.*, Oct. 1896, where he records forty cases. The advantage of this method is the ease with which the broad ligaments are tied at their base, while its drawback is the inclusion of more tissue in the ligature.

⁴ Maunsell has suggested the extension of deperitonised extra-peritoneal method to complete extirpation. And the final result (see his diagrams *Brit. Gyn. Jour.*, 1892-93, p. 79) is practically what Le Bec has since described.

⁵ Price gives a table of statistics of Abdominal Hysterectomy by leading operators. *Am. Jour. Obst.*, 1892, Vol. ii., p. 751, compiled from various sources. Cushing has collected by circular the results of hysterectomy in America, according to the different modes of treating the pedicle, and finds a mortality for supra-vaginal hysterectomy of 13 p. c. for extra-peritoneal, 38.7 p. c. for intra-peritoneal, and 8.5 p. c. for infra-peritoneal methods; and for complete abdominal hysterectomy of 14.3 p. c. for myoma, 29.1 p. c. for cancer, and 7.5 p. c. for salpingitis. Under the intra-peritoneal he includes ligature by Schroeder's or Zweifel's methods and the intra-peritoneal separate ligature of the uterine arteries and covering of the stump with peritoneum. (The evolution in America of abdominal hysterectomy: *Annal. of Gyn. and Ped.*, No. 9.)

the ultimate appeal must always be to facts, viz., the recovery and ultimate wellbeing of the patient.¹

A *fibroid of the cervix* may push its way into the cellular tissue and displace the peritoneum. Such an extra-peritoneal tumour may also be removed by laparotomy.²

Sänger³ reports on two cases of abdominal section for fibroid tumour of the cervix: in one, the pedicle was treated by the elastic ligature and dropped back; in the other, the uterus was amputated and the stump stitched by Zweifel's method (*v. p.* 434) and dropped back. Kelly⁴ also cut down on two fibroid tumours of the cervix and removed them with the *écraseur*; no pedicle was tied, but the peritoneal cavity was drained and washed out for some days afterwards. Byford⁵ has removed a subserous fibroid of the cervix *per vaginam*.

D. REMOVAL OF OVARIES OR OF UTERINE APPENDAGES.

The removal of these, as we have seen (*v. p.* 245), usually stops menstruation and induces the menopause. Hence in the case of fibroid tumours this operation does good in two ways—by checking bleeding and stopping the growth of the tumour.⁶ The mortality is also low (under 3 p. c.), so that this operation, were it always practicable, would have a wide field in the treatment of myoma. Unfortunately, it is frequently impossible to get at both ovaries in cases of large myoma; while one is to the front and easily accessible, the other is to the back and sometimes low down towards the pouch of Douglas. The technique is the same as that described in Chapter XXI., with the exception that a long abdominal incision is often necessary to allow the operator to pass the whole hand into the abdomen so as to get at the appendages. As to mortality, Lawson Tait⁷ gives it as 4.4 p. c. for his first 272 cases. While it undoubtedly stops bleeding, its effect on the tumour has been much canvassed. Tait says that in seventeen out of fifty which he followed for some time after the operation, the tumour had entirely disappeared. Wiedow⁸ also reports carefully on fifty-six cases followed for some time, and mentions having found in thirty-nine cessation of bleeding and diminution in the size of the tumour.

¹ As the President said in the discussion before the London Obstetrical Society on "Relative Methods," in which the *serre-nœud* was described as a barbarous procedure; "Each one of us would prefer to have his life saved by a 'barbarous procedure' to being killed by the most advanced surgical triumph."

² As in Thelen's case: *Centralb. f. Gyn.*, 1885, No. 3.

³ *Centralb. f. Gyn.*, Bd. xiii., S. 207.

⁴ *Amer. Jour. Obstet.*, 1886, p. 45.

⁵ *Amer. Jour. Obstet.*, 1888, 1205.

⁶ Thornton says, "There can be no doubt that the operation of removal of the appendages, in suitable cases, is less dangerous to life than that of hysterectomy, and in my own hands its after-results have been excellent. I know of two cases only in which the tumours have not entirely disappeared; and one of those, for reasons too long to enter upon here, is not a test case: the other would, I believe, have recovered, if she had given herself time, but she got into the hands of the electricians;" *op. cit.*, p. 629.

⁷ *Diseases of Women*, p. 194.

⁸ W. Wiedow.—Zur Kastration bei Uterusfibrom (*Cent. f. Gyn.*, 1882, No. 6, S. 81). Die Kastration bei Uterusfibrom (Congrès internat. de Copenhague, 1882): *Archiv f. Gyn.*, 1885, Bd. xxv., Hft. 2, S. 299.

CHAPTER XXXVIII.

FIBRO-CYSTIC TUMOUR OF THE UTERUS.

LITERATURE.

Atlee—Ovarian Tumours: Philadelphia, 1873. *Beates*—Cystic leiomyoma of Uterus: Am. Journ. of Obstet., 1884, p. 753. *Breus*—Ueber wahre epithelführende Cystenbildung in Uterusmyomen: Wien, 1893. *De Sinéty*—Manuel de Gynécologie, Paris, 1879, p. 413. *Diesterweg*—Ein Fall von Cysto-fibroma verum: Zts. f. Geb. und Gyn. ix., S. 191. *Grasskopff*—Zur Kenntniss der Cystomyome des Uterus: Munich, 1884. *Gusserow*—Neubildungen, etc.: Stuttgart, 1885, S. 117. *Haultain*—Benign tumours of the Uterus: Allbutt and Playfair's System of Gynecology, London, 1896. *Heer*—Ueber Fibrocysten des Uterus: Zurich, 1874. *Leopold and Fehling*—Ein Beitrag zur Lehre von den kystischen Myomen des Uterus (Myosarcoma lymphangiectodes uteri): Archiv für Gyn., Bd. vii., S. 531. *Peaslee*—Ovarian Tumours: London, 1873. *Rein*—Beitrag zur Lehre von den lymphangiectatischen Fibromyomen des Uterus in pathologisch-anatomischer und klinischer Beziehung: Archiv f. Gyn., ix., S. 414. *Schroeder*—Die Krankheiten der weiblichen Geschlechtsorgane, S. 213: Leipzig, 1878. *Sir Spencer Wells*—Ovarian and Uterine Tumours: London, 1883. *Spiegelberg*—Die Diagnose der cystischen Myome des Uterus und ihre intraperitoneale Ausschälung, eine neue Operationsmethode derselben: Archiv f. Gyn., vi., S. 341. *Thomas*—Diseases of Women, p. 551: London, 1882.

SYNONYM—Cysto-fibroma.

Attention has only of recent years been directed to this, the rarest form of uterine tumour. Its pathology is now being worked out, and at present we group under this head tumours which may afterwards be shown to be anatomically separable. On the other hand, some do not describe fibro-cystic tumours as a special form, but refer to their production under the heading of "degeneration in fibroids."

PATHOLOGY.

The majority of fibro-cystic tumours are simply fibroid tumours which have become softened. The spaces between the bundles of fibrous tissue open out and contain serum; the trabeculæ between adjoining spaces give way, which allows these to run together to form larger cavities. Fig. 248 shows this in a *subserous fibroid*, the form which most frequently undergoes this change.

The term "cystic," is, it is evident, misleading as applied to this

form of tumour. The cavities are not "cysts," that is, they do not possess a special wall.

Kœberlé was the first to suggest that some forms of fibro-cystic tumour might be due to *dilated lymphatics*. Leopold and Fehling¹ have carefully described a case in which the cavities were lined with endothelium. The fluid from these cavities was of a clear yellow colour, and *coagulated* as soon as it was exposed to the air; fibrin was




FIG. 248.

LARGE THREE-LOBED FIBROID SPRINGING FROM THE FUNDUS BY A SOMEWHAT THIN PEDICLE, of which CF is cystic, while SsF and the dark shaded mass behind the uterus are subserous. This along with two smaller fibroids growing from the posterior surface of the uterus was removed by laparotomy (Schroeder).

present in it. To this form the name of *Fibromyoma lymphangiectodes* has been given. Müller¹ has also described recently a preparation in which he found the epithelial lining present in the smaller cysts. Atlee says this coagulation of the fluid — formation of colourless

¹ Beitrag zur Kenntniss der cystoiden Uterustumoren: *Archiv f. Gyn.*, Bd. xxx., S. 249.

blood-clot—is diagnostic of the fluid from *all* fibro-cystic tumours, and may be relied on to distinguish them from ovarian. Spiegelberg records a case in which this spontaneous coagulation of the fluid was observed, but the most careful microscopic examination could detect no epithelial lining of the cavities. A transition case has been described by Rein, in which the cavities were not themselves lined with endothelium but *communicated* directly with the lymphatic spaces.

Mucoid
Degenera-
tion.

Mucoid degeneration of a fibroid tumour has been described by Virchow as Myxomyoma. In this case the interstitial tissue contained fluid rich in mucin and with numerous nucleated round cells.

Sarcomatous degeneration of a fibroid¹ apparently also produces a cystic condition of a fibroid tumour although this is not a true fibro-cystic tumour.

Cysts with an epithelial lining have been described by Babesin and Diesterweg. The latter removed on two occasions (with two years' interval) a submucous polypus with cysts; the cavities were lined with ciliated epithelium and contained thin brownish blood. Baer on cutting through a similar polypus with the *écraseur* was afraid that he had cut through the peritoneal cup of an inverted uterus, as the appearance of the section of the cyst resembled it.

SYMPTOMS.

These are the same as those of fibroid tumours, with the exception of a more rapid growth. As they are usually subserous, menorrhagia is not often present.

DIAGNOSIS; DIFFERENTIAL DIAGNOSIS.

Their diagnosis is often difficult, as the difference in consistence between the more solid and the fluid parts may escape detection. The connection of the tumour with the uterus is the most important point to establish. Examination under chloroform, with the drawing down of the uterus with the volsella and exploration per rectum is of service.

Differential Diagnosis.—Their diagnosis from ovarian tumours is the most important and, at the same time, the most difficult. As in the majority of cases they are merely altered fibroid tumours, their differentiation from a simple fibroid is merely a matter of degree of softness. In a case described by Beates as one of Cystic Leiomyoma of the uterus, the patient had been tapped twice²; and, as the fluid gave the ovarian cell described by Drysdale (*v. p.* 253), the case was set down

¹ As in Fenger's case (*Amer. Jour. Obstet.*, 1888, p. 1200), and probably also Erich's (*ibid.*, 1886, p. 517). Aslanian records a case of fibro-sarcoma with a dilated condition of the veins—*Archiv de Toc.*, Feb. 1895.

² In Hosack Fraser's case, the patient was tapped twenty-one times—*Brit. Med. Jour.*, 1896, p. 837.

as undoubtedly one of ovarian tumour. The differential diagnosis from ovarian tumour is often not made till the abdomen is opened.¹

TREATMENT.

The treatment consists in removal through the abdominal walls, according to the methods described for fibroid tumours² (*v. p. 456*).

¹ As in a recent case by Lewers, *London Obstet. Trans.*, 1895, p. 270. Also by Mayo Robson, *Brit. Gyn. Jour.*, 1891-92, p. 137.

² Sometimes enucleation is required as in Alban Doran's case—*Brit. Med. Jour.*, 1893, p. 1006. The same patient has been operated on more than once for this condition—*Lancet*, 1888, i., p. 973.

CHAPTER XXXIX.

UTERINE POLYPI: TUBERCULOSIS.

LITERATURE.

Barnes—Diseases of Women, p. 195 : London, 1878. *De Sinéty*—Manuel pratique de Gynécologie, p. 419 : Paris, 1879. *Gusserow*—Die Neubildungen des Uterus, Billroth's Handbuch, S. 179 : Stuttgart, 1885. *Hegar und Kaltenbach*—Die operative Gynäkologie, S. 473 : Stuttgart, 1881. *Hicks, Braxton*—Three cases of very large polypi of the uterus, etc.; Obstet. Journ. of Great Brit., Jan., 1879. *Küstner*—Notiz zur Metamorphose des Uterusepithels : Centralblatt f. Gyn., 1884, p. 321. *Matthews Duncan*—Edin. Med. Journ., July 1871; and Obstet. Journ., 1873, p. 497. *Simpson, Sir J. Y.*—Diseases of Women, p. 704 : Edin., 1872. *Thomas*—Diseases of Women : London, 1880, p. 558. *Underhill*—On the Structure of three cervical Polypi, and the Structure of a true mucous Polypus of the Cervix : Edin. Obst. Soc. Trans., Vol. iv., pp. 231 and 241.

By the term "Polypus" is understood a pediculated tumour attached to the mucous membrane of the uterus. It includes the following tumours, which are anatomically distinct :—

1. Submucous fibroids, which have become pediculated and are in process of extrusion ;
2. Mucous polypi and adenoma ;
3. Pediculated cystic follicles ;
4. Placental polypi ;
5. Papilloma of the cervix.

For clinical reasons, it is convenient to use the term polypus in its general sense as implying an external form alone ; the symptoms produced by these tumours resemble one another, and their exact nature is sometimes not made out till they are removed. Pathologically, the term should be limited to mucous polypi. It is confusing to speak of a fibroid tumour which has a broad base of attachment as a submucous fibroid, and of one which has a pedicle as a fibrous polypus. The polypoidal projections formed by pediculated ovula Nabothii are only pediculated retention cysts. Placental polypi, unless malignant, are not true new-formations.

1. *Pediculated submucous fibroid tumours* form the so-called "fibrous Polypus." They spring from the muscular wall of the uterus, usually from the *body* which, as we have seen, is more commonly the seat of fibroid tumours than the cervix. They are of *firm* consistence, of a size varying from a goose's egg upwards, and are of a rounded or pyriform shape (fig. 249), sometimes elongated and constricted through the pressure of the uterine walls (fig. 236); the surface is smooth or marked with furrows corresponding to the fasciculi of fibrous tissue.




FIG. 249.

FIBROUS POLYPUS LAID OPEN TO SHOW ITS IDENTITY IN STRUCTURE WITH A FIBROID TUMOUR
(Sir J. Y. Simpson).

Sometimes they are of such a size¹ that, although lying in the vagina, they fill the pelvis and press on the bladder and rectum; the uterus is then raised above the pelvic brim (just as it is elevated when the vagina is distended with fluid), and is felt as a smaller body riding on the top of the tumour. Adhesions may form between the surface of the fibroid and the vagina, producing the impression that the tumour springs from the vaginal mucous membrane.²

¹ Koeberlé removed one weighing over 1½ lbs. (*Centralb. J. Gyn.*, 1889, S. 263).

² Braxton Hicks—*Loc. cit.*

The pedicle consists of a narrowing of the calibre of the tumour towards its base of attachment, or of a distinct stalk which may be long enough to allow the fibroid to lie at the vulva. As fibroid tumours are sparingly vascular, the pedicle does not, as a rule, contain large vessels. When a pediculated submucous fibroid lies in the cavity of the uterus, it sets up uterine contractions which lead to its expulsion; there is a




FIG. 250.

INTRA-UTERINE SUBMUCOUS FIBROID WHICH IS BECOMING VAGINAL (*Sir J. Y. Simpson*).

stage at which it lies partly within the uterus (fig. 250), partly in the vagina (the portion constricted by the cervix has been mistaken for a pedicle, and only the lower lobe of the hour-glass tumour removed); finally, the whole tumour lies in the vagina, but still maintains its connection with the uterus through its pedicle (fig. 251). The congestion of the fibroid excites uterine contractions, specially at the menstrual

period, and thus favours its expulsion. At those times, we may have the cervical canal only temporarily dilated and the polypus projecting through it; after the period, the contractions pass off and the polypus is retracted into the uterine cavity. This condition is fully described by French writers under the name of "*polypes à apparitions intermittentes*." Its practical importance is that we should examine sometimes at the menstrual period, when a polypus (not recognisable at other times) may be felt through a dilated cervix.

They have the microscopic structure described at p. 424 (*v. fig. 249*).

2. *Mucous polypi* are developed from the *mucous membrane* of the ^{Mucous} uterus, most frequently from that of *the cervix*. They are of *soft pulpy* ^{Polypi.}



FIG. 251.

SUBMUCOUS FIBROID WHICH HAS COME TO LIE WHOLLY IN THE VAGINA (*Sir J. Y. Simpson*).

consistence, of about the size of an almond—rarely larger—and have a flattened form; usually, there are more than one present (*fig. 252*). They are extremely vascular and have the microscopic structure of the mucous membrane from which they are developed.

The typical cervical polypus has the structure seen at *fig. 253*. From the fact that the gland-ducts appear as channels on the surface, it was described by Oldham as the "*channeled polypus*." Sometimes the polypus shows also the stratified epithelium of the vaginal aspect of the cervix, as in a specimen described by Underhill; he supposes that in this case it sprang from the margin of the *os externum*; he describes also a polypus which sprang from the *vaginal aspect* and

showed only the stratified epithelium. Küstner has shown that stratified epithelium may be found on mucous polypi which have grown high up in the cervical canal; this is another example of how the single-layered uterine epithelium may become changed into stratified epithelium (*cf.* Zeller's observations, p. 346). These polypi sometimes form the starting-point of malignant disease; Underhill traced the commencement of sarcomatous degeneration in one case.

De Sinéty divides them into two groups according as they spring (1) from the cervix, (2) from the body of the uterus. Each has the




FIG. 252.

GROUP OF MUCOUS POLYPI GROWING IN THE CERVIX UTERI (*Sir J. Y. Simpson*).

characteristic epithelium (*v.* p. 22) lining the ducts and cysts; the former have the columnar non-ciliated epithelium of the cervix, the latter the ciliated cylindrical epithelium of the body.

A localised hypertrophy of the glands of the uterus has been described by Schroeder as adenoma polyposum; the changes resemble those of glandular endometritis (*v.* p. 345).

Williams, in his monograph,¹ describes four cases of adenoma of the cervix, two being simple villous growths and two being malignant.

¹ Cancer of the Uterus: London 1888, pp. 40-44.

3. *Pediculated Nabothian follicles* have been already described under cervical catarrh (p. 334).

4. *Placental or fibrinous polypi*. These are produced as the result of Placental Polypi.




FIG. 253.

SECTION OF A MUCOUS POLYPUS OF THE CERVIX ⁴⁰. *g* dilated glands, *e* epithelium, *mf* muscular fibre, *v* blood-vessel, *ct* connective tissue (*De Sinéty*).

incomplete detachment of the placenta; in some cases we can trace placental villi in their structure. On the surface of this irregularity of the mucous membrane, blood coagulates; and thus the fragment of




FIG. 254.

NON-MALIGNANT PAPILLOMA OR FIBROMA PAPILLARE OF CERVIX (*Ackermann*).

placenta grows larger through being coated with fibrin. This increase in size may go on until the polypus is the size of an egg. This form of polypus is not a new-formation and only finds a place here on account of its polypoidal form. After an abortion,¹ they may form

¹ Küstner—Beiträge zur Lehre von der Endometritis: Jena, 1883.

in the same way, a piece of decidua left in the uterus maintaining its structure, vitality and nutritive connection with the tissues below.

Attention has recently been directed to "destructive polypi" of the placental site, especially in connection with deciduoma malignum and its relation to hydatid mole. Fraenkel¹ has studied the changes in hydatid moles, and finds colonies of cells, lying at some distance from the villi, apparently due to a proliferation of the syncytium and the layer of cells beneath. These form the stepping-stone to the malignant tumours arising at the placental site (see Deciduoma Malignum—Chap. XLIV.).

5. *Papilloma of the cervix.* Simple papilloma of the cervix is a very rare form of tumour; the great proportion of papillary tumours found here are malignant (carcinomatous or sarcomatous). Fig. 254 shows such a tumour, described by Ackermann,² which sprang from the anterior lip of the cervix. It consisted of a branching stem of connective tissue, with papillæ covered mostly with squamous but in some places with a single layer of cylindrical epithelium. There was no recurrence after removal. The term "cauliflower" excrescence, introduced by Clarke, describes very well the appearance of these tumours. Virchow has shown that in many of these papillomata we find proliferation of the epithelium, and that they form the first stage of epithelial cancer of the cervix (*v. p.* 488); we must therefore regard the cauliflower excrescence as, in the great proportion of cases, a malignant tumour.

SYMPTOMS.

These are Hæmorrhage,
Leucorrhœa,
Dysmenorrhœal pains,
Sterility,
Irritation and discomfort.

Hæmorrhage.

The hæmorrhage shows itself first as an increase of the ordinary menstrual flow; afterwards, it comes at irregular intervals. In the case of a submucous fibroid, it comes from the uterine mucous membrane which is hypertrophied. In the mucous polypus, it comes from the tumour itself which is vascular and bleeds easily; when the polypus protrudes through the cervix, there may be hæmorrhage³ (*v.* the preparation represented at fig. 94). In other cases the drain of blood, though not directly fatal, may produce profound anæmia; hence the importance of ascertaining and removing the cause of the hæmorrhage. The cachectic appearance of the patient, thus induced, may be such as to lead us to

¹ Die Histologie der Blasenmolen und ihre Beziehungen zu den malignen von den Chorionzotten (Decidua) ausgehenden Uterustumoren: *Archiv f. Gyn.*, xlix. (1895), S. 480.

² *Virchow's Archiv*: Bd. xliii., S. 88.

³ Barnes records the case of a woman of twenty-six years of age in which a polypus the size of a walnut produced a fatal hæmorrhage.

form a strong prepossession in favour of the existence of malignant disease before we proceed to physical examination.

The *leucorrhœa* is due to the endometritis which is always present. ^{Leucorrhœa.} The polypoidal retention-cysts are the result of a chronic catarrh of the cervix or uterus. It is disputed whether mucous polypi are the cause or the result of the inflammatory changes; De Sinéty inclines to the latter view. When the polypus comes to lie in the vagina, it produces an irritating vaginal leucorrhœa.




FIG. 255.

PEDICULATED SUBMUCOUS FIBROID, springing from the fundus, which has not dilated the cervical canal (*Sir J. Y. Simpson*).

The *dysmenorrhœal pains* are due to the muscular efforts of the uterus ^{Dysmenorrhœal Pains.} to expel the polypus, and are most marked when the polypus has descended to the os internum or lies in the cervical canal.

In rare cases the presence of the foreign body in the uterus has produced the sympathetic phenomena of pregnancy—pigmentation of the breasts and abdomen, and morning sickness.

Sterility. *Sterility* is occasioned by the mechanical obstruction of the polypus, or more frequently by the associated endometritis.

A pediculated fibroid may form a serious complication to labour, in preventing the progress of the child's head; such a polypus has been laid hold of with the forceps under the impression that it was the presenting head. They may also give rise to hæmorrhage in the puerperium.¹

DIAGNOSIS.

1. *When the polypus has dilated the os externum*, it will be recognised by the finger per vaginam. If it be larger than a walnut and of firm consistence, and if the uterine cavity be increased in length, it is a pediculated fibroid tumour. If it be small and of a pulpy consistence, it is a true mucous polypus; mucous polypi do not, as a rule, produce hypertrophy of the uterus.

Having learned that there is a pediculated body in the vagina or cervical canal, carry the finger upwards to ascertain its point of attachment; if this be high up in the uterine cavity, the tumour is a pediculated fibroid; if it springs from the cervical mucous membrane, it is probably a mucous polypus.

On bimanual examination, the uterus is found to be enlarged in the case of pediculated fibroids; it is not enlarged with mucous polypi, unless from associated chronic metritis.

The speculum shows that the surface of the true mucous polypus has a bright cherry-red colour, which contrasts with the darker red of the cervical mucous membrane embracing it. The appearance of the fibroid tumour depends on the condition of the investing mucous membrane, which is often ulcerated or sloughing; when the capsule has given way, the fibrous substance of the tumour is seen to be of a paler colour.

2. *When the uterus is enlarged but the os externum not dilated*, the diagnosis is more difficult (fig. 255). If the uterus be markedly enlarged and of firm consistence and (the possibility of pregnancy being excluded), the sound pass for 4 or 5 inches, there is probably a sub-mucous fibroid tumour. It is difficult to determine whether it is pediculated or not. The sound must be used with care as its use is not unattended with risk; laceration of the mucous membrane, with the introduction of septic matter, has resulted from too free and repeated exploration in this way. Dilatation of the cervix and exploration with the finger are sometimes necessary to ascertain whether the fibroid be pediculated, and to what part of the uterus it is attached.

3. *When the uterus is not much enlarged*, the diagnosis is very difficult. The possibility of a fibroid tumour is excluded. A small

¹ See paper by Halliday Croom on Fibrous Polypi complicating the puerperium: *Edin. Med Journ.*, xxii., i., p. 289.

mucous polypus, however, may exist in the uterine cavity and escape detection with the sound. In such a case, it is recognised only on dilating the cervix and exploring the uterine cavity with the finger.

The *curette* is a valuable aid to diagnosis when the actual exploration of the uterine cavity with the finger is not desirable. By its use we diagnose and treat the case at the same time. Thus irregularity of the uterine surface (which is easily detected by the *curette*), and the character of the scrapings removed, may show that we have to do with pediculated retention-cysts or placental polypi.

DIFFERENTIAL DIAGNOSIS.

The characters which distinguish a pediculated fibroid from a mucous polypus are its larger size, firmer consistence, and its springing from the body of the uterus. The uterine cavity is increased in size. We probably find, also, other fibroid tumours, interstitial or subserous.

A pediculated fibroid hanging down into the vagina, may readily be mistaken for the inverted fundus uteri; this is most likely to happen when there is much hæmorrhage from the former, and when concomitant pelvic inflammation makes examination difficult. A true diagnosis here is all-important, as removal of the fibroid may save the patient's life; while amputation of the uterus, under the supposition that it was a fibroid, may lead to disastrous consequences.

Given a tumour the size of a pear hanging down through the cervical canal into the vagina, we wish to make sure that it is not the inverted body. First, sweep the finger carefully round the neck and note whether the mucous membrane of the cervical canal is reflected on to the neck of the tumour; sometimes inflammatory adhesions round the neck produce a condition simulating inversion. Now make the bimanual; if the body in the vagina be a fibroid, the uterus will be in its normal place. The abdomino-vaginal examination is often difficult on account of the body in the vagina; therefore pass the finger into the rectum, through the anterior wall of which we can distinctly feel whether the cervix has a truncated end above (inversion), or passes up into the body of the uterus (fibroid); the abdomino-rectal makes this more evident. When examination is difficult and the diagnosis doubtful, we should not hesitate to give chloroform and make a thorough examination; it is well to be prepared to operate at the same time, if necessary.

Finally, use the sound, which is an important test. Sweep the finger carefully round the neck of the tumour and feel for a depression corresponding to the os, into which endeavour to introduce the sound. If it passes for two and a half inches or more and is then arrested, it is probably in the uterine cavity; make sure of this by pressure with the hand on the abdominal wall, or per rectum.

When the tumour in the vagina fills the pelvis or rides above the brim, so that the finger cannot reach the pedicle or feel whether the os is present, the diagnosis is very difficult. We rely on careful abdominal palpation to ascertain whether the uterus can be felt resting on the top of the tumour.

We must not forget that we may have both conditions present, *i.e.*, pediculated fibroid + a certain amount of inversion.

PROGNOSIS.

The prognosis as to *danger to life* will depend on the hæmorrhage. Wherever a polypus is present, we should advise its removal.

As to *the operation*, the removal of mucous polypi and smaller fibroids is safe and easy. The fear of hæmorrhage from the pedicle of a fibroid tumour, which led to the treatment by ligature, has been found by experience to have been exaggerated. Where there is a rigid cervix to be dilated before we can remove the tumour, where the tumour is large so that it must be removed in portions, where there is a thick pedicle and consequently a larger raw surface, the operation will be a more serious one and the prognosis given more guardedly.

Should there be pregnancy, the polypus may be removed without interrupting its course. If it be of such a size as to interfere with labour, it should be removed as soon as discovered.

TREATMENT.

Whenever it is necessary to dilate the cervix for diagnosis, we should have instruments ready to remove the tumour at the same time. The dilatation is effected by laminaria tents, or by Hegar's graduated dilators. A good method is to place a laminaria tent in the cervix to start the dilatation; after six or eight hours chloroform the patient, fix the cervix with volsellæ, and introduce the graduated dilators in succession till the cervical canal is wide enough to admit the index finger; remove the polypus by the means to be described; and wash out the uterine cavity with an antiseptic solution.

Small *polypoidal projections* are removed with the curette, as described under Endometritis, followed by the application of pure carbolic acid.



FIG. 256.

FORCEPS WITH
CATCH FOR
REMOVING
MUCOUS POLYPI.

Mucous polypi are twisted off with the forceps, shown at fig. 256. It is advantageous to use forceps with a catch, as this keeps a steady hold of the tumour and leaves the operator's fingers free to twist the forceps round. In removing *fibroids*, we first ascertain the seat of insertion and size of the pedicle. When the tumour is small, we

can learn this by the fingers; when so large that we cannot get the fingers past the tumour to the pedicle, we probe round its base with the sound or, laying hold of the tumour with forceps, endeavour to rotate it and thus test the thickness of the pedicle.

The pedicle will yield to torsion with the forceps. This is the simplest method and should always be tried in the first instance. If this fail, divide the pedicle with curved scissors. Make traction with the forceps to render the pedicle tense; too forcible traction might produce inversion. Guarding the uterine wall with the fingers, carry in the curved scissors. In cutting, make the scissors hug the surface of the tumour and thus keep clear of the uterine wall. Strangulation by ligature, formerly widely practised, is now entirely abandoned; the sloughing stump was a fruitful source of septicæmia.

When the pedicle is of considerable thickness, it may be divided with the *écraseur* or with the galvano-caustic wire. The wire *écraseur* is preferable to the chain *écraseur*, as it is more easily applied. For the nature and method of use of the *écraseur*, the student is referred to Treatment of Carcinoma of the Cervix.

When the size of the tumour makes the pedicle inaccessible, it must be diminished. This is best effected by Hegar's method: traction is made on the tumour, which is at the same time incised in a spiral manner with scissors; the tumour is thus (as it were) unwound, till finally the pedicle is reached and divided.

Chloroform is not necessary for the removal of smaller polypi. The section of the pedicle is painless; if pain be present on tightening the *écraseur* round the neck of a polypus, the operator should examine carefully again to make sure that the wire is not constricting the inverted fundus. Where the polypus is large and the operation tedious, it is better to have the patient anæsthetised as the operator has then more freedom.

TUBERCULOSIS UTERI.

LITERATURE. *Hegar*—Die Entstehung, Diagnose, und chirurgische Behandlung der Genitaltuberculose des Weibes: Stuttgart, 1886. *Cornil*—Sur la Tuberculose des Organes génitaux de la femme: Verneuil, 1888. *Pozzi*—Traité de Gynécologie: Paris, 1897, p. 904. *J. D. Williams*—Tuberculous Disease of the Portio Vaginalis: Brit. Med. Jour., 1895, Vol. i., p. 969. The literature is given fully in Williams' paper.

WHILE tuberculosis of the genital tract, though a rare condition,¹ has given rise to considerable discussion as to its production by direct infection, and tuberculosis of the tubes is of special interest with regard to tubercular peritonitis, tuberculosis of the uterus is not of great importance. It is almost always secondary, that is to say, appearing in

¹ Williams found it present in 3 % of 100 *post-mortems* on the female pelvic organs, and cites its frequency in the operating theatre as being, according to Martin 3 %, and Whitridge Williams 8 % in the case of appendages removed for inflammatory disease.

the course of development of tubercle in some other organ, and especially in the lungs. Pozzi refers to three forms, but admits that the distinction between them is theoretical—an acute miliary, an interstitial of very chronic course, and an ulcerating form which is the most frequent.

Williams, in describing two cases of ulcerating tubercular disease of the cervix, draws attention to the curious fact that this portion of the genital tract is sometimes alone secondarily affected.

The diagnosis of tubercular endometritis can, of course, only be made from the discovery of the bacillus in the discharge, which is often a matter of great difficulty (Cornil). The treatment is that of the general condition.

CHAPTER XL.

CARCINOMA UTERI (OF CERVIX): PATHOLOGY AND ETIOLOGY.

LITERATURE.

Barbour—Cases of Carcinoma of the Female Pelvic Organs: Edin. Med. Jour., July 1880. *Barnes*—Diseases of Women, p. 821: London, 1878. *Gusserow*—Die Neubildungen des Uterus, S. 199: Stuttgart, 1885. Ueber Carcinoma Uteri, Volkmann's Samml. klin. Vor., N. 18. *Pozzi*—Traité de Gynécologie: Paris, 1897. *Ruge and Veit*—Zur Pathologie der Vaginal-portion: Stuttgart, 1878. Der Krebs der Gebärmutter: Stuttgart, 1881. *Russell*—The Operative Significance of Metastasis and Post-operative Recurrences in Carcinoma of the Uterus: Am. Jour. Obstet., Dec. 1896. *Schroeder*—Die Krankheiten der weiblichen Geschlechtsorgane, S. 264: Leipzig, 1878. *Seelig*—Path. Anat. Untersuch. über die Ausbreitung des Gebärmutterkrebs: innaug. dissert. Strasburg, 1894. *Simpson, Sir J. Y.*—Diseases of Women: Edinburgh, 1872, p. 140. *Sinclair, W. J.*—Malignant Diseases of the Uterus: Clifford and Allbutt's System of Gynecology, London, 1896. *Tanner*—On Cancer of Female Sexual Organs: London, 1863. *Veit*—Sur Anat. des Carcinoma Uteri: Zeits. f. Geb. u. Gyn., Bd. xxxii. *Virchow*—Ueber Cancroide und Papillargeschwülste, 1850. *Williams*—Cancer of the Uterus: London, 1888. *Winter*—Über die Recidive des Uteruskrebses insbesondere über Imprecidive. Also, Über die Ursache der Krebsrecidive: Zeits. f. Geb. u. Gyn., 1892, S. 141.

THUS far we have considered only the simple or benign tumours in the uterus. We pass now to the malignant; and these present themselves under the two chief types, viz., epithelial and connective-tissue, as carcinoma and sarcoma. Two rarer forms of malignant disease have recently attracted attention—the adenoma malignum and deciduoma malignum. The former will be considered with Carcinoma of the Body of the uterus (Chap. XLIII.), as it is the body which is usually affected by it. The position of the latter is not yet determined, but provisionally we place it alongside of Sarcoma Uteri.

The cervix, as we have seen, differs anatomically from the body of the uterus; it also differs pathologically, *i.e.*, is distinctly marked off from the body of the uterus as regards some of the morbid processes to which it is liable. We have seen that while the body of the uterus is the common seat of fibroid tumours, the cervix is rarely so; in cancer the opposite condition obtains, for the body is rarely, while the cervix

is very often, attacked by it. When cancer of the uterus is spoken of, it is in fact almost always cancer of the cervix that is meant; and it is the latter that we have chiefly to consider here, for only about 2 p. c. of the cases of cancer are in the body, the remaining 98 p. c. being in the cervix.

PATHOLOGY.

On no subject in pathology has more been written and a greater variety of opinion expressed than on carcinoma. We have endeavoured to arrange, in the table on page 485, the facts most important for the student to know.

CLASSIFICATION.

There are three varieties of carcinoma usually given in the English text-books. These are medullary (encephaloid) and scirrhus cancer, and epithelioma. Now the distinction between the first two is merely a question of degree; in the former the cellular element, in the latter the fibrous stroma is in excess. When we say that medullary cancer is frequent but scirrhus rare, we only mean that carcinoma runs a rapid course when it occurs in the uterus. The distinction between these two and epithelioma is more marked, and is therefore given in the table.

From the above it is evident that we are not yet in a position to make a scientific classification. The division according to clinical features into *true carcinoma* and *epithelioma* or *cancroid* (*καρκίνος* and *εἶδος*, *like cancer*) is convenient: as it implies that in some cases progress is more rapid than in others; and that the disease in the one case produces metastatic deposits, in the other remains local.

In the following pages we shall use the term carcinoma as meaning the glandular type of tumour, called by some adeno-carcinoma, in contra-distinction to epithelioma.

ORIGIN.

Virchow's
View.

View of
Thiersch
and
Waldeyer.

As regards the origin, there are two distinct views. That the disease arises from *connective-tissue cells alone*, is the view maintained by Virchow and his followers; while Thiersch and Waldeyer hold that in *all cases* it originates in *epithelial cells*. In the cervix, as possible sources, there are two varieties of epithelium; the squamous on the vaginal aspect, the cubical lining the canal. In the *flat* *cancroid* of the cervical canal, it arises from the cubical epithelium which lines the latter; in the papillary form, it originates in the cells of the rete Malpighi on its outer aspect (*Klebs*). It will be seen that Waldeyer holds the view that, in all cases, it arises from the latter only.

CLASSIFICATION ACCORDING TO CLINICAL FEATURES.	FORMS.	ORIGIN.	POSITION.	PROGRESS.
<p>CARCINOMA</p> <p>progresses rapidly ; produces metastasis, affects connective tissue rapidly.</p>		<p>{ from the cervical epithelium of constricted cervical glands (<i>Krebs</i>) ;</p> <p>{ from plugs of the deepest layers of squamous epithelium on the vaginal aspect of cervix (<i>Waldeyer</i>) ;</p> <p>{ from connective-tissue cells of cervix (<i>Virchow</i>).</p>	<p>in substance of cervix.</p>	<p>produces thickening, then ulceration ;</p>
<p>EPITHELIOMA OR CANCROID</p> <p>progresses slowly ; does not produce metastasis ; spreads by extension.</p>	<p>{ flat (flache cancröid)</p> <p>{ papillary</p>	<p>{ from the cubical epithelium of cervical canal (<i>Krebs</i>) ;</p> <p>{ from plugs of the deepest layers of squamous epithelium on vaginal aspect of cervix (<i>Waldeyer</i>) ;</p> <p>{ from the deepest layers of squamous epithelium on vaginal aspect of cervix (<i>Krebs and Waldeyer</i>).</p> <p>{ from connective-tissue cells (<i>Ruge and Veit</i>).</p>	<p>superficial within cervical canal.</p> <p>superficial outside of cervix.</p>	<p>excavates cervix ;</p> <p>spreads downwards into vagina (cauliflower excrescence).</p>
				<p>When ulceration and breaking down have been produced, these forms are no longer distinguishable.</p>

Ruge and
Veit's
Investiga-
tions.

More recent investigations into the origin of carcinoma are by Ruge and Veit. According to them carcinoma arises, in the majority of cases, from a transformation of the connective-tissue cells; even the




FIG. 257.

CANCER OF THE VAGINAL PORTION (*J. Williams*).

a, Normal squamous epithelium on the vaginal aspect of the cervix; *b*, processes of cancerous cells which have developed from it.

papillary form which produces the so-called cauliflower excrescence, although it apparently springs from the epithelium, is developed from




FIG. 258.

CANCER OF THE CERVIX PROPER (*J. Williams*).

a, Normal columnar epithelium lining a gland within the cervical canal; *b*, cancerous cells derived immediately from it.

the connective-tissue cells. The connective-tissue stroma becomes vascular and almost like granulation tissue. The young cells, which

are apparently produced from the connective-tissue corpuscles, take on an epithelial character. These observers never saw plugs of epithelium extending downwards into the connective-tissue.

Williams, on the other hand, in figuring a specimen like one by Ruge and Veit, says that the hypertrophied connective-tissue papillæ pushing their way through the proliferating cancerous epithelium (the superficial layers of which are shed in places) produce only an appearance of their being the starting-point of the disease.

According to the place in the cervix in which it begins, we distinguish Cancer of the Vaginal Portion from Cancer of the Cervix proper—an important distinction which we owe to Ruge and Veit. It is difficult to draw an imaginary line which would divide the cervix into these two parts; but if we hold to an exclusively epithelial origin for cancer, we can define the former as cancer beginning in the *squamous* epithelium on the vaginal aspect, the latter as cancer beginning in the *columnar* epithelium lining the canal. Figs. 257 and 258 illustrate the origin of cancer-cells from these two sources. Cancer of the vaginal portion is the rarer of the two forms.¹

POSITION.

There are apparently three places in the cervix where carcinoma may first show itself. (1.) It may begin as hard *nodules in the substance of the cervix* underneath the mucous membrane; these increase in size, Three positions in Cervix.




FIG. 259.

CARCINOMATOUS NODULE GROWING IN ONE LIP OF THE CERVIX AND PUSHING THE MUCOUS MEMBRANE OUTWARDS. The figure to the right is a section of the cervix made through the line *x* (Schroeder).

come to the surface of the mucous membrane (fig. 259), and produce ulceration. (2.) More rarely does it commence in the *interior of the cervical canal* and spread along its mucous membrane so as to excavate the canal. (3.) It may appear *on the vaginal aspect of the cervix* as an ulcerating surface (fig. 261) or as an irregular papillary tumour, which, extending downwards into the vagina, attains considerable size.²

¹ Seven undoubted cases of it, and fifteen of cancer of the cervix proper, are described in Williams' monograph.

² So Veit distinguishes cancer of the vaginal portion, cervical cancer, and cancer nodules in the cervix. In addition to papillary, nodular, and excavating forms, Pozzi makes a fourth—the liminary or vaginal, in which it commences in the posterior fornix.

Form of
Slow
Ulceration
not malign-
ant.

It is important to remember that there is a form of slow ulceration on the surface of the vaginal portion which is not malignant. Sir John Williams¹ described this as "corroding ulcer of the os uteri:" it begins at the os and extends symmetrically downwards into the vagina, without hard or thickened edges, extending by simple ulceration or the formation of reddish raised tubercles which ulcerate; in one case, there was calcification of the internal iliac arteries; of three cases observed, the duration was in one for two years and in two for ten years. According to Matthews Duncan, this is a form of lupus, which we shall have to notice specially as an affection of the vulva.

There is also a form of adenoma which, though it is not malignant (*v. p.* 476), tends to become so. Fürst² has recorded a very interesting case of this in which the amputated cervix showed only the appearance of a cysto-adenoma, while eighteen months afterwards the patient died of true cancer of the cervix.

PROGRESS.

While in the initial stage we may recognise these three forms, after ulceration has occurred they pass into one another and are no longer distinguishable.

As regards the further progress, there are *three modes of spreading* of the disease; first, downwards into the vagina; second, upwards into




FIG. 260.

MICROSCOPIC SECTION OF A PORTION OF THE CERVIX UTERI SEEN IN FIG. 259. *e* squamous epithelium in several layers; *c n* carcinomatous nodule; between these is seen a portion of inflamed mucous membrane covered with a single layer of epithelium (*Schroeder*).

the body of the uterus; and, third, into the connective tissue of the pelvis. This distinction is of importance from the operative stand-

¹ *Brit. Med. Jour.*, April 5, 1884.

² Ueber suspectes und malignes Cervix-Adenom: *Zeits. f. Geb. u. Gyn.*, xiv., S. 352.

point, because while the first group gives scope for amputation of the cervix, and the second for extirpation of the uterus, the third shows the difficulties which beset operative treatment and its unsatisfactoriness in the majority of cases. The local dissemination of cancer through the connective tissue has recently demanded attention from its bearing on operative treatment by extirpation of the uterus, which has become a recognised operation. This affection of surrounding tissues has a bearing both on judging as to the suitability of cases for operation, and on the recurrence (or rather the appearance) of the disease after its apparent removal.

The *lymphatics* play the chief rôle in this dissemination, and it is to Poirier's researches as to their course, and the work of Seelig, Winter, Veit, and others, as to their implication in carcinoma, that we are




FIG. 261.

SECTION OF A FLAT CANCROID (EPITHELIOMA) OF THE CERVIX. *e* squamous epithelium, *cc* carcinomatous cells; between these is seen some granulation tissue (Schroeder).

chiefly indebted. From fig. 54 (p. 78) we see that the lymphatics from the body of the uterus pass through the upper part of the broad ligament to the lumbar glands, receiving on their way those from the ovary and Fallopian tube; while those from the cervix and upper part of vagina pass through the lower part of the broad ligament to the iliac or ilio-pelvic glands. A third and less important chain runs down the round ligament to the inguinal glands. The upper and lower lumbar glands receiving the uterine lymphatics are shown in fig. 53, in which the uterus is anteverted so that the group of vessels running to the lumbar glands, which are higher up on fig. 54, appear in fig. 53 as if lower down. These lumbar glands surround the aorta and vena cava, extending as far down as the bifurcation of the common iliac artery

where they become continuous with the iliac or ilio-pelvic glands. These last, which receive the cervical lymphatics, extend from the bifurcation of the common iliac artery downwards—in front of the internal iliac. Cancerous infiltration of the lumbar glands is difficult to recognise, though it may show itself through interference with the




FIG. 262.

PLUG OF CANCER CELLS IN A LYMPHATIC VESSEL (*Seelig*); endothelium adherent to thrombus at *é*, detached at *e*.

circulation. Cancerous ilio-pelvic glands, on the other hand, can be palpated per vaginam or per rectum. The inguinal glands, the most accessible, are very rarely affected in uterine cancer.¹

In an early stage of cancer, nodules may be found separate from the original focus, sometimes in a chain. Seelig has shown that these nodules are plugs of carcinomatous tissue which have grown within the lymphatic vessel from carcinomatous epithelium carried by the lymph stream. The endothelium of the lymphatic vessel is not affected (*v. fig. 262*). In cervical cancer, extension takes place along the lymphatics, and especially in those of the outer muscular layers; while in

¹ Russell figures an interesting specimen with nodules in the round ligament, evidently in the chain of lymphatics which follow it. Enlargement of the inguinal glands is frequent in affections of the vulva, but in such cases, infection is by another route.

cancer of the body, the lymphatics of the mucosa and then those of the inner muscular layers are involved. This slower and more limited extension explains the better results, as to recurrence, in hysterectomy for cancer of the body as compared with that for the cervix.

In cancer of the cervix, Abel and Landau¹ have found changes in the *mucous membrane of the body* also—not only those of chronic inflammation, but also of carcinomatous degeneration; they further found microscopic changes exactly similar to sarcoma, but which might be the first stage of carcinoma of the body.

Eckart,² on the other hand, found only hyperplasia of the glands with papillary proliferation into their lumen, *i.e.*, endometritis glandularis.

Saurenhaus, from the examination of a still larger amount of material,³ has shown that the changes, though extensive, are of a benign character, whether we characterise them as a hyperplastic endometritis or a simple adenoma.

EXTENSION TO NEIGHBOURING ORGANS.

In its further progress, the carcinomatous growth invades the surrounding organs. Pushing its way forwards in the cellular tissue




FIG. 263.

CARCINOMA beginning in the CERVIX UTERI, and ending in the production of recto-vesico-vaginal fistula (Farre).

between the *bladder* and the uterus, it involves the mucous membrane of the former; it first produces vesical catarrh, then sloughing of the

¹ Ueber das Verhalten der Schleimhaut des Uteruskörpers bei Carcinom der Portio vaginalis: *Archiv f. Gyn.*, xxxii., S. 271, and xxxv., S. 214.

² From the examination of ten uteri extirpated by Kaltenbach for cancer of the cervix: *Centralb. f. Gyn.*, 1888, S. 426.

³ Fifty uteri extirpated for cancer: *Centralb. f. Gyn.*, 1888, S. 755.

walls, and finally vesico-vaginal fistula. The bladder is affected in a




FIG. 264.

VERTICAL MESIAL SECTION OF PELVIS, FROM CASE OF CARCINOMA UTERI. *a*, Perineal body; *b*, Symphysis pubis; *c*, Rectum; *d*, Body of Uterus; *e*, Small fibroid; *f*, Urethro-vaginal septum; *g*, Bladder. A small tube passes between bladder and excavated cervix through a fistula (*Barbour*).

considerable proportion of cases; of 311 cases of carcinoma this occurred in 41 per cent., fistula resulting in 18 per cent. (*Gusserow*). From the




FIG. 265.

VERTICAL MESIAL SECTION OF PELVIS, FROM CASE OF CARCINOMA VAGINÆ ET UTERI. *f* points to vagina eroded by disease; *e* is a malignant growth attached to uterus. Other letters as in fig. 264 (*Barbour*).

position of the ureters, they are frequently involved. The carcinoma-

tous growth may press upon the ureters near their point of entrance into the bladder; or it infiltrates their walls, and the consequent thickening produces constriction at the part affected. Dilation of the ureter above thus results, which produces hydronephrosis and finally atrophy of the kidney (see fig. 270). The frequency of this condition will be apparent from the fact that Blau found it present in 57 out of 93 post-mortem examinations. Artaud describes two degrees of kidney affection¹: with moderate pressure the kidney is slightly enlarged and shows hypertrophy of the glomeruli and dilatation of the convoluted tubules with small-celled infiltration round both of these and the arteries; (2) with greater pressure, dilatation of the ureters and atrophy of the kidney. More rarely does the carcinomatous infiltration extend backwards into the *rectum* and produce recto-vaginal fistula; of 282 cases the rectum was affected in 18 per cent., fistula resulting in 8.5 per cent. (*Gusserow*).² When both bladder and rectum have been opened into, a common cloaca is produced as in fig. 263.

Perforation into the *peritoneal cavity* is rare. The peritoneum is not simply pushed forward, but is taken up into the carcinomatous growth. As this process goes on, adhesions are constantly being formed between the layers of the peritoneum in front of the growth so that it does not project free into the cavity beyond. These adhesions further prevent the peritoneal cavity from being opened into when the carcinomatous mass breaks down.

The accompanying sections (figs. 264, 265), made from post-mortem preparations, will serve to illustrate some of the points noted above.

Points to be noted in fig. 264.

1. Seat of disease in the *cervix*;
2. Complete destruction of the cervix and lower segment of the uterus;
3. Production of an irregular cavity from the extension of the disease in *three* directions through the cellular tissue—

Description of two
Pelves with
Cancer of
Cervix.

- (a) Behind the uterus,
- (b) Between the uterus and the bladder,
- (c) Between the vagina and the bladder;

4. The pouch of Douglas entirely obliterated and partially replaced by the carcinomatous excavation, the vesico-uterine pouch shortened by adhesions, perforation into the peritoneal cavity at one point;
5. Bladder small and contracted, carcinomatous fistula;
6. Rectum intact.

¹ Lancereaux considers ascending nephritis a constant phenomenon in cancer, being present from the earliest stages of the disease. *Anat. des Mal. des Org. Génito-urin.*, 1884, p. 417.

² Fere and Carron (Statistics of Complications of Carcinoma Uteri in 51 post-mortems at the Salpêtrière 1881-83) found extension to the bladder with fistula in 18, to the rectum in 7, and to the peritoneum in 9 cases.

Points to be noted in fig. 265.

1. Vagina (as well as cervix) affected, the nymphæ had a cartilaginous consistence, inguinal glands enlarged—although not shown in figure ;
2. Extension of the disease along the mucous membrane of the uterus, excavating it though not destroying the walls to the same extent as in fig. 264 ;
3. Partial obliteration of the pouch of Douglas ;
4. Bladder dilated through pressure on the urethra, its walls apparently not involved ;
5. Rectum intact.

ETIOLOGY.

The female sex is more liable to carcinoma than the male. According to Sir J. Y. Simpson's statistics, the proportion is $2\frac{1}{2}$ to 1. These statistics are drawn from the Annual Reports of the Registrar-General for England during the years 1847-1861. During that time there were 87,348 fatal cases of carcinoma, of which 61,715 were among women and 25,633 among men. For the year 1860, the deaths from carcinoma among men were .97 per cent. of the total male mortality, among women 2.2 per cent. The cause of this greater relative frequency is connected with the development of the sexual organs in the female. Up to puberty, the mortality (from carcinoma) of the sexes is the same ; afterwards, the relative proportion of female to male deaths gradually rises till it attains its maximum about the age of 50, after which it falls away again (fig. 266).

The diagram on page 495 is based on the statistics of 91,058 deaths in Great Britain. It brings out three facts : the total number of deaths in each sex increases with age to a certain point ; the increase among women is relatively the greater ; it reaches its maximum at an earlier age with the female sex.

It is a remarkable fact that while the mortality from phthisis and tubercular disease has diminished, that from cancer has increased by more than four times what it was half a century ago.¹

The most frequent seat is the uterus, where fully one-third of the total cases occur ; the next in frequency is the mamma.

Although the immediate etiology of carcinoma is unknown, there are certain causes general and local which favour its development.

1. The *general predisposing causes* are the following :—

Heredity ;

Age ;

Depreciation of the vital powers.

¹ See Roger Williams on "The continued increase of Cancer, with remarks as to its Causation." *Brit. Med. Jour.*, 1896, ii., p. 318. In 1840, 1 in 129 of the total mortality was due to cancer ; while in 1894 it had risen to 1 in 23. He attributes this to increase of industrialism acting injuriously on the health of the town population, and also to the increased consumption of meat.

TABLE AND DIAGRAM OF COMPARATIVE FATALITY OF CARCINOMA IN MALE AND FEMALE, ACCORDING TO AGE.




FIG. 266.

In the DIAGRAM, the upper line indicates mortality in the female, the lower that in the male.

Race and
Heredity.

The influence of *race* is brought out in Chisholm's statistics, which show that carcinoma is more than twice as frequent among the white population as among the black. As regards *heredity in families*, much less stress is now laid upon this than formerly.

According to Gusserow's statistics, in 1028 cases heredity was proven in only 79, that is in about 7·6 per cent. Schroeder, placing the statistics of Sibley and of Barker together, shows that heredity has been proven in only 8·2 per cent.; Picot places it at 13 per cent. These figures show that we cannot lay much stress on heredity as a predisposing cause. On the other hand, we must remember that these statistics are drawn principally from hospital reports, from a class of people who know little about the former history of their families.

Age.

Age has undoubtedly a considerable influence upon the frequency of this disease. This is evident from the table given on page 495. Gusserow collected statistics of 2210 cases reported by various authorities. The mortality per cent. for various ages is represented by the curve in the diagram on page 495. From the table it is evident that carcinoma is very rare before puberty. The proportion of cases below 20 years (2 in 2270) is so small that it need not be taken into account. The first glance at the diagram would lead one to believe that the increasing frequency of the disease is due to the development of the functional activity of the sexual organs, but a more careful consideration shows that the increase continues and reaches its maximum after the latter has ceased. This table should be compared with that for Fibroid Tumours on page 435.

Deprecia-
tion of
Vital
Powers.

Whatever tends to *depreciate the vital powers* favours the occurrence of this disease. We meet with it more frequently among the poorer classes, where there is insufficiency of food with privation and hardship. Schroeder contrasts, in this respect, the development of carcinoma with that of myoma. In his polyclinic among the poorer classes, the proportion of carcinoma to myoma was as 100 to 61; in his private practice among the wealthier, it was as 100 to 332.

2. The *local predisposing causes* are the following:—

Erosion of the cervix and protracted catarrh;
Repeated parturition.

Influence
of Split
Cervix.

The relation of *erosion and laceration of the cervix* to the development of carcinoma has been recently pointed out by Ruge and Veit, and also by Breisky. We draw attention to this point specially, because the most important differential diagnosis is that between long-standing inflammation and commencing malignant disease; and the possibility that the former may pass into the latter should always be kept in view.¹

¹ Williams, however, in his cases never found the disease starting in a tear, and thinks that there is no evidence that laceration plays any part in the etiology of cancer.

Influence
of repeated
Parturi-
tion.

Repeated parturition has an important influence. Carcinoma is much more frequent in multiparæ. Gusserow finds an average of 5·1 children to every case of carcinoma, which is a high average productivity. Whether this is due to the greater functional activity of the uterus or to the production of fissures with their resulting chronic inflammatory changes, is a more difficult question.

CHAPTER XLI.

CARCINOMA UTERI (OF CERVIX): SYMPTOMS AND DIAGNOSIS.

LITERATURE.

See Literature of Chapters XL. and XLII.

SYMPTOMS.

The *local* symptoms of carcinoma uteri are three—

Hæmorrhage,
Offensive discharge,
Pain.

There are in addition a considerable number of *general* symptoms, which arise secondarily.

As a rule, however, no symptoms are present in the first stage, that is until ulceration sets in. In exceptional cases, when infiltration of the connective tissue or of the walls of the uterus has taken place at an early period, pain may be an early symptom; there is no pain so long as the disease is limited to the cervix. This entire absence of symptoms until the disease has already made considerable progress, is the reason of the great difficulty in ascertaining the period of its probable commencement. From the same cause, the patient does not seek relief till the possibility of eradicating the disease is much diminished.

LOCAL SYMPTOMS.

Hæmorrhage is usually the first symptom noticed by the patient. She observes that menstruation is more profuse than formerly. This, when the disease occurs late in life, she attributes to approach of the menopause. In other cases, profuse hæmorrhage occurs irregularly between and independent of the menstrual periods. Sometimes the hæmorrhage is noticed only after exertion (as straining at stool) or after coitus. Sometimes the patient states that "the menstrual flow never entirely ceases;" which means that the vaginal discharge is always tinged with blood. The explanation of hæmorrhage in these earlier

stages is to be found in the vascularity of the stroma of the new formation.¹ It is rich in delicate vessels which readily rupture. In the later stages, hæmorrhage is not a prominent symptom unless a large vessel be accidentally eaten into. Death from hæmorrhage is rare.

Discharge. The *discharge* characteristic of carcinoma is not present until ulceration has occurred. In the papillary form of epithelioma² (cauliflower excrescence) there is a free discharge before the growth has begun to break down; this is of a watery character, has no odour, and is due simply to the transudation of serum. As soon, however, as ulceration occurs in any of the forms, there is a discharge containing the molecular debris of the breaking down tissue which gives it a characteristic and peculiarly offensive odour. In the rapidly growing forms (medullary) of carcinoma, there is an almost equally rapid molecular death of the newly formed tissue due to fatty degeneration of the epithelial cells. In epithelioma this discharge is less marked, because there is less necrosis of tissue; but in true carcinoma, especially in an advanced stage, it is quite characteristic. In fact, a diagnosis may be sometimes made merely from the odour which hangs about the person. At first the discharge is yellowish-white in colour, but afterwards from the decomposition of the fatty cells it becomes of a reddish-brown; if there is hæmorrhage, it will be tinged with blood.

Pain. *Pain* is not such a constant symptom as is usually supposed. Some cases run their whole course without the patient's complaining specially of pain. It is not present so long as the disease is limited to the cervix; hence it is of no use as a diagnostic of carcinoma of the cervix in its early stage, unless the cellular tissue has been at the same time involved. But as soon as the new growth has extended upwards to the body of the uterus or to the cellular tissue of the pelvis, pain is produced through pressure on or actual lesion of the terminations of the nerves. The character of the pain varies. It is "a dull gnawing pain localised in the pelvis or back," or "a sharp pain shooting through to the back or down the thighs to the knees"; this last is caused by simple pressure on the crural and sciatic nerves, or, in the later stages, from affection of the cellular tissue of the nerve sheaths. Occasionally it is felt in the mammæ or other seats of uterine sympathetic pain. The intensity of the pain varies also in different cases; it is marked where there is more formation of new tissue and less ulceration, that is when there is more pressure on the nerve endings. Thus, if there has been much deposit between the uterus and the bladder accompanied with an increase of

¹ Pozzi attributes the initial hæmorrhages to congestion of the uterus, apart from rupture of the vessels of the new growth; and compares it to hæmoptysis in phthisis.

² Though, as we have said, we have not at present a truly *pathological* classification of the different forms of carcinoma, it is convenient, *clinically*, to use the terms Epithelioma and true Carcinoma. By them we do not imply anything as to the origin of the disease. By epithelioma we understand those forms which begin more superficially, spread more slowly, and do not tend to involve the connective-tissue.

pain, we find that the pain diminishes when the mass breaks down and a vesico-vaginal fistula is formed. We may distinguish between pain due to the development of carcinoma, and that produced by the chronic peritonitis which accompanies it when the peritoneum becomes affected; the latter produces great sensitiveness of the abdominal walls to pressure, and a board-like rigidity from reflex spasm of the muscles.

GENERAL SYMPTOMS.

In addition to these local symptoms which are immediately due to the carcinomatous infiltration and degeneration, there are more general symptoms which arise secondarily.

First we mention *loss of flesh* and *general debility*. The patient may continue healthy and well-looking in the early stages; sometimes, one is surprised to find that the disease is already well advanced in a patient who to outward appearance is in perfect health. But, sooner or later, the drain on the system produces great emaciation. The patient also has a careworn expression, partly from this loss of flesh and partly from the constant pain; from this expression alone, known as the "cancerous facies," the diagnosis may sometimes be made.

The wasting (*marasmus*) is occasioned not only by the drain of the new growth, but also by *disturbances* of the *digestive system* which arise in the course of the disease. Loss of appetite may amount to disinclination for food, and digestion is interfered with. This is produced at first sympathetically, as in other uterine disorders; but latterly it is due to gastric catarrh, constipation, the condition of the blood (anæmia and uræmia), and the unhealthiness of the atmosphere resulting from the offensive discharges.

There is, further, *painful micturition and defæcation* according to the extent to which the bladder and rectum are involved. The latter is always present, as the rectum, whenever it is distended, presses upon the carcinomatous growth. When fistulæ are produced, the urine and fæces pass per vaginam.

Pruritus vulvæ frequently results from the acrid and irritating discharge, and from the dribbling of the urine from a fistula. The *skin* acquires in the later stages a *dingy straw tint*, which when very marked is suggestive of jaundice. That disease may actually be present when there is secondary carcinoma of the liver, but this is rare. The colour is due to the anæmia, or (according to Barnes) to the absorption of decomposed fæcal matter (*copræmia*).

DIAGNOSIS.

As the patient does not seek advice till cancer has begun to ulcerate, the physical signs have by that time become well marked and the diagnosis is usually easy.

Vaginal
Examina-
tion.

On making a *vaginal examination*, the finger feels the enlarged, thickened, irregular, everted lips of the cervix spreading like a mushroom in the vagina (described by Malgaigne as "*champignons cancéreux*"). Sometimes a distinct tumour is present, the form of which is sufficiently indicated by the term *cauliflower excrescence* (see fig. 268). In other cases the finger feels an irregular ulcerated surface in the position of the cervix, soft and friable with hard and unyielding margins. The examining finger is stained with blood, and the odour of the discharge cannot fail to be recognised. If there is any doubt as to diagnosis, a fragment should be removed and examined microscopically. The appearance of a fibrous stroma with alveoli which contain irregular cells




FIG. 268.

CAULIFLOWER EXCRESCENCE GROWING FROM THE CERVIX UTERI (Sir J. Y. Simpson).

of an epithelial type with one or more large nuclei, will confirm the diagnosis of carcinoma.

The value of the microscope in the diagnosis of cancer has given rise to considerable difference of opinion. While Williams¹ and Griffith² emphasise its usefulness, Thornton³ and Herman lay more stress on clinical observation, the former showing the risks of "partial operation," as he characterises the snipping off of portions for microscopic examination. Clinical observation, corrected and supplemented by microscopic examination, is best. While microscopic examination is not in many cases of value, in a certain proportion of cases it is conclusive; and its utility comes in just at that early period where clinical investigation feels its weakness. The microscopical examination of scrapings removed by the curette is of much less value than of pieces excised, for although the histological

¹ *Op. cit.*

² Early Diagnosis of Cancer of the Uterus: *Brit. Med. Jour.*, 1896, Vol. i., p. 264.

³ The Early Diagnosis of Malignant Disease, etc.: *Brit. Med. Jour.*, 1896, Vol. i., p. 261. See also Jessett on the Early Diagnosis of Malignant Disease: *Brit. Gyn. Jour.*, 1896-97, p. 327.

characters of epithelioma are so distinctive that it may be recognised from the fragments, it is difficult to be sure of adeno-carcinoma without proof of the invasion of deeper structures which can only be got from a large piece of tissue.

The *speculum* need not be used for the recognition of carcinoma, Speculum. except in its early stage or to ascertain more exactly the seat and extent of the growth. If the disease be far advanced and the diagnosis certain, the introduction of it causes unnecessary pain and hæmorrhage. Sinclair¹ recommends the use of the *sharp spoon* in diagnosis, as Spoon. chronically inflamed tissues do not break down under it, but this belongs rather to its differential diagnosis from chronic inflammation.

The *rectal examination* is valuable, and in these cases should always be carefully carried out. It gives us important information in two distinct classes of cases. First, in early carcinoma or in cases where there is a suspicion of commencing carcinoma, the cellular tissue of the pelvis should be carefully examined to ascertain whether any localised deposit or enlarged glands can be felt; this can be done most easily by the rectal examination. If it is desirable to introduce two fingers into the rectum or if the examination causes much pain, the patient should be narcotised. Second, in cases of advanced carcinoma where the vaginal examination is difficult on account of the hæmorrhage and pain which it occasions, a more thorough examination can be made per rectum. The finger can reach higher up than per vaginam, and thus we can ascertain the extent of the carcinomatous deposit and the size and mobility of the uterus. The condition of the rectal mucous membrane itself is observed at the same time, to ascertain whether it is already involved in the disease. In some cases the rectal examination is the only one possible, as in the case of carcinoma vaginæ represented at fig. 265 where the deposit round the ostium vaginæ made the introduction of the finger impossible. Rectal Examination.

DIFFERENTIAL DIAGNOSIS.

The following are the most important lesions from which carcinoma is to be differentiated:—

- Hypertrophy of the cervix, with induration and occluded follicles;
- Papillary erosion or ectropium, with cicatricial tissue;
- Syphilitic ulceration, condylomata on the cervix;
- Small fibroid in the cervix, sloughing polypi;
- Retained portions of placenta or membranes;
- Diphtheritic inflammation of the mucous membrane;
- Sarcoma of the cervix.

As regards the first two of these, it is evident that carcinoma resembles them only at an early stage. But it is precisely at this stage

¹ *Loc. cit.*

Import-
ance of
Pelvic
Examina-
tion in
Cancer.

that a correct diagnosis is all important for treatment. We should also remember (as Ruge and Veit have pointed out) that these conditions may be at once the result of chronic inflammation and the starting-point of malignant disease. The statement of the patient that the symptoms have existed for a long time, should not throw us off our guard. In *all* cases in which a patient *over forty years of age* seeks advice *with symptoms referable to the pelvis*, a careful examination should be made. We may thus accidentally discover carcinoma in an early stage, while still within the possibility of radical treatment. If the carcinomatous infiltration be general, it cannot be distinguished, except by microscopical examination, from chronic induration. When localised, the diseased part is distinctly marked off from the adjoining tissue, shows a difference in its level, and is of a slightly yellow colour with granular yellowish-white inequalities.¹ Where there is only suspicion of carcinoma, there is no harm in excising a portion of the suspected part and submitting it to *microscopic investi-*




FIG. 269.

SCRAPING FROM CARCINOMA OF THE CERVIX, STAINED WITH LOGWOOD, $\times 250$; drawn by S. Delépine.

gation. A careful examination per rectum of the pelvic cellular tissue should always be made as mentioned above.

A superficial ulcerating epithelioma might be mistaken for a simple erosion, but has *thickened infiltrated edges*. The latter may, however, pass into the former.

Condylomata on the cervix simulate epithelioma, but they disappear under appropriate treatment. Syphilitic ulceration produces sometimes deep excavation, even a rectal fistula. This at the first glance might be taken for carcinoma, but more careful examination and inquiry into the history of the case will remove all doubt.

Small myomata are more sharply defined than a carcinomatous nodule of the same size, because the surrounding tissue is not infiltrated.

¹ Stratz—Zur Diagnose des beginnenden Carcinoms an der Portio: *Zeits. f. Geb. u. Gyn.*, Bd. xiii., S. 89.

When a small submucous fibroid or a cervical polypus has ulcerated, it presents appearances similar to an ulcerating carcinomatous nodule. The former however is firmer, and fragments cannot be broken off by the finger-nail, while the latter is friable and breaks down easily.

The possibility that carcinoma may be first noticed during the puer- ^{Carcinoma}perium should be remembered. In such a case it is more likely to be ^{in the} in the body of the uterus, as cancer of the cervix affects conception ^{Puer-}perium. Even in the body, it is rare in the puerperium; and cases described as such are usually Deciduoma malignum (*v.* Chap. XLIV.).

Diphtheritic inflammation of the mucous membrane may easily be confounded with ulcerating carcinoma (*Schroeder*). The irregular swelling of the mucous membrane and the offensive discharge tinged with blood, which are present in diphtheritic inflammation, may be suggestive of carcinoma at the time; but this superficial resemblance soon disappears.

Sarcoma of the cervix is a very rare condition. Sarcomatous tumours are softer and grow more rapidly than carcinomatous. A positive diagnosis can only be made after microscopical examination of scrapings taken from the tumour (fig. 269).

PROGNOSIS.

The prognosis in carcinoma is always very grave. The possibility of spontaneous cure is a disputed point. There is one apparently well-authenticated case recorded by Habit.¹ Another is mentioned by Barnes,² in which there is some doubt as to the correctness of diagnosis. The prognosis as to the probable duration of life will depend on the extent to which the disease has already advanced and the possibility of checking its progress or even extirpating it altogether by operative interference. With regard to the results of operative interference, see under Treatment.

As regards the duration of disease if not interfered with, there is a slight difference of opinion. This may be explained by the variable period in the course of the disease at which the symptoms appear. Sir J. Y. Simpson gives the probable duration of life after the detection of the disease as from 2 to 2½ years; Gusserow and Schroeder give it as from 1 to 1½; while, according to Fordyce Barker, it is as long as 3 years and 8 months. The statistics of H. Arnott, drawn from 57 carefully observed cases, give the duration, after the first symptom (usually a flooding), of true cancer as 53·8 weeks; of epithelioma, 82·7 weeks. We may say therefore to the patient's friends that the disease will run a course of from one to two years. It is better not to tell the patient herself what her trouble is, though its serious nature should not be disguised.

¹ *Sydenham Society's Year Book*, 1864, p. 401.

² Barnes, *Diseases of Women*: London, 1878.

CAUSES OF DEATH.

The causes of death, arranged in order of importance, are the following:—

Exhaustion,
Uræmia,
Peritonitis,
Septicæmia,
Hæmorrhage,
Venous thrombosis.

Exhaustion.

Exhaustion, under which we include *marasmus*, is the result partly of the drain on the system and partly of the inability to take food.




FIG. 270.

KIDNEYS AND URETERS, from a case of Cancer of Uterus with Uræmic Convulsions.

Uræmia.

The importance of *uræmia* as a frequent cause of death has only recently been recognised. According to Seyfert,¹ in the majority of cases death results from it. It is due to compression of the ureters, as already described under Pathology. It may be acute, accompanied by coma and convulsions. Fig. 270 shows the kidneys and ureters from a patient who had two convulsions before death, and we have recorded another case in which convulsions were present.² More

¹ Säxinger, *Prager med. Vierteljahrsschrift*, Bd. i., S. 103.

² Edinburgh Hospital Reports, 1895, p. 675.

generally it is chronic, and shows itself in the dulness of the patient, occasional headache, and decreasing sensibility to pain — which diminishes suffering as the disease approaches its termination.

Peritonitis is sometimes the cause of death, but not so frequently *Peritonitis*, as one would suppose; the disease is prevented from extending to the peritoneum generally by the adhesions which are formed. When peritonitis occurs, it is localised and chronic; in some cases, however, a general peritonitis is set up which proves fatal. *Perforation* may take *Perforation*.




FIG. 271.

CARCINOMA OF THE CERVIX LEADING TO OCCLUSION OF OS UTERI, dilatation of uterus and perforation (A. R. Simpson). Uterus and vagina laid open; a quill is passed through the perforation.

place from the sudden giving way of adhesions; the escape of the carcinomatous debris into the peritoneal cavity produces death from shock or septic peritonitis. The preparation shown at fig. 271 was taken from a patient in whom the cause of death was *rupture of the uterus*. The case is reported and the preparation described by A. R. Simpson (op. cit., p. 276). There was carcinoma of the cervix which had contracted the lumen of the canal; the cavity of the uterus was expanded the walls being thinned out; at the fundus "was a small

perforation about the size of a pea, with thin edges," through which fluid had escaped and set up peritonitis which rapidly proved fatal.

Septicæmia.

Septicæmia suggests itself as a likely cause of death. We are familiar with it as produced in the puerperal condition: it is explained by the fact that, at that time, there is abundant means for absorption in the numerous lymphatics, and large veins which have been recently lacerated; hence, whenever septic matter is present, there is great risk of septicæmia. Similar conditions exist in carcinoma, during the progress of which the blood-vessels are eroded and their extremities bathed in putrid matter. Barnes has drawn special attention to this as a source of blood-poisoning; according to Eppinger's¹ observations its occurrence is rare, and this he ascribes to the diminution of the absorptive power of the eroded vessels.

Hæmorrhage.

Hæmorrhage is in very rare instances immediately fatal. As already pointed out, though it is important as an early symptom, it occurs less frequently and is less abundant as the disease advances. If a large vessel be suddenly opened into, a fatal hæmorrhage may follow.

Thrombosis.

Venous thrombosis, due to mechanical compression of the veins, sometimes occurs; and a clot may be detached producing embolism in the lungs. Fatty degeneration of the heart is, sometimes, also present.

Patients with cancer have also died of tetanus,² which has been ascribed to the action of micro-organisms from secretion retained through plugging of the vagina.

¹ *Prager med. Wochenschrift*, 1876, S. 210.

² See case by Hofmeier: *Centralb. f. Gyn.*, Bd. xi., S. 171.

CHAPTER XLII.

CARCINOMA UTERI (OF CERVIX): TREATMENT.

LITERATURE.

Barnes—Diseases of Women, p. 856: London, 1879. *Currier, And. F.*—Removal of the Uterus for Cancer: N. Y. Med. Jour., vol. xxxiv., p. 494. *Duncan, W. A.*—On Extirpation of the entire Uterus: London Obstet. Trans., 1885, p. 8. *Freund*—Samm. klin. Vorträge, Nr. 133; and Centrabl. f. Gyn., N. 12, 1878. *Gusserow*—Die Neubildungen, etc.: Stuttgart, 1885, S. 199. *Hegar und Kaltenbach*—Die Operative Gynäkologie, S. 391. *Hofmeier*—Zur Statistik des Gebärmutterkrebses und seiner operativen Behandlung: Zeitsch. f. Geb. und Gyn., Bd. x., S. 269. *Lewers*—On the supra-vaginal amputation of the cervix uteri for malignant disease, etc.: Lancet, 1888, i., p. 464. *Pawlik*—Zur Frage der Behandlung der Uteruscarcinom.: Wiener Klinik, 1882, Hft. xii., S. 425. Ueber die endgültige Heilung des Carcin. Cervicis Uteri durch Operation—Zeits. f. Geb. u. Gyn., Bd. xiii., S. 360. *Sänger*—Zur vaginalen Totalexstirpation des carcinomatösen Uterus: Archiv f. Gyn., Bd. xxi., Hft. 1. *Schroeder*—Charité Annalen: V. Jahrgang, S. 343. Zeitschrift für Geburtshilfe und Gynäkologie: B. iii., S. 419; B. vi., Heft 2, S. 218. *Simpson, A. R.*—Carcinoma Uteri; Contributions to Obstetrics and Gynecology, p. 261: Edinburgh, A. & C. Black. Case of Extirpation of the Cancerous Uterus through the Vagina: Edin. Obst. Trans., Vol. vii., p. 136. *Simpson, Sir J. Y.*—Diseases of Women, p. 170: Edinburgh, 1872. *Sims, Marion*—The Treatment of Epithelioma of the cervix uteri: American Journ. of Obst., July 1879. *Sinclair, W. J.*—Malignant Diseases of the Uterus: Clifford and Allbutt's System of Gynecology: London, 1896. *Thomas*—Diseases of Women, p. 591; 1882. *Thornton, J. Knowsley*—Article Hysterectomy in Clifford and Allbutt's System of Gynecology: London, 1896. See also references under Operations and Results in the text.

THE treatment of carcinoma ought to be regarded in two aspects: first, as treatment of *the symptoms*; second, as treatment of *the disease*. Again, the treatment of the disease may be either *palliative* or *radical*.

We need not discuss here the vexed question whether carcinoma is a constitutional or a local disease. It cannot be too strongly impressed on the practitioner that, as far as our present experience goes, in attacking the disease itself he must rely upon *surgical* and not on *medical* treatment. Our aim ought to be the removal of the disease and not merely the alleviation of the symptoms. To remove it completely we must recognise it early. Up to the present time successful treatment has been a rare occurrence, because we have failed to recognise carcinoma in its commencing stages. The possibility of treating it successfully in the

future will depend on the possibility of our recognising it in its commencement. Not less important than early recognition is complete removal and that without delay. In the uterus, more readily than in the mamma, does the carcinoma get beyond the reach of the operator. In carcinoma mammæ, we can excise not only the breast but also the axillary glands if these should be already implicated. But, in carcinoma uteri, as soon as the pelvic glands are involved the case is hopeless as regards a radical cure.

We shall consider, first, the treatment of the symptoms; because, in the majority of cases, when the patient comes under our notice, the disease itself has already got beyond our reach.

TREATMENT OF SYMPTOMS.

These are hæmorrhage, offensive discharge, pain.

HÆMORRHAGE.

In the treatment of hæmorrhage, there are two points to be considered: *first*, the instructions to be given to the patient; and, *second*, the means which we can ourselves employ.

Use of
Ergot.

(1.) The patient is instructed to take the liquid extract of ergot in large doses whenever there is much hæmorrhage either during the menstrual period or independent of it. If she is subject to floodings, a friend might be taught how to give the ergotin solution hypodermically. Ice applied to the vagina and injections of cold water check hæmorrhage; a small piece of sponge or tampon of wadding, soaked in perchloride of iron, might be passed into the vagina if cold is not sufficient. The patient is recommended to avoid sexual intercourse as this favours active congestion, and in some cases is the cause of hæmorrhage.

(2.) The means at our own command are the following:—

Simple pressure, effected by complete and thorough plugging of the vagina;

The use of styptics, caustics, or the actual cautery;

The removal of diseased tissue by the curette or other means.

The
Vaginal
Tampon.

The plugging of the vagina should be done whenever we are called in on account of profuse hæmorrhage. The packing is carefully done with pledgets of lint or cotton wadding (with string attached) soaked in carbolic oil; the speculum is introduced carefully and not carried high up.

Of styptics, the best are the perchloride and the pernitrate of iron. Sir J. Y. Simpson recommended a saturated solution of the perchloride in glycerine. A pledget soaked in either of these is introduced, and placed so as to be in contact with the bleeding surface; and the rest of the vagina is packed, as above described, with the pledgets steeped in carbolic oil. The perchloride should be used with great caution in

cases of advanced ulceration, as we have seen it corrode into the tissue so as to reach the peritoneum and produce peritonitis. The use of caustics, cautery, and curette, will be considered under Operative Treatment.

OFFENSIVE DISCHARGE.

This is best treated by astringent and antiseptic injections. These should be used frequently, as it is important to keep down the unpleasant odour and make the patient's surroundings as comfortable as possible. If the discharge be plentiful and not very offensive, as in the cauliflower excrescence, the indication is more for the use of astringents like sulphate of alumina and iron (4 grains to the oz.). Tannin or sulphate of zinc can also be used, and it is well to change the astringent occasionally. If there is much necrosis of tissue with offensive discharge, corrosive sublimate (1 to 2000) is required. Solution of bromine (1 of the B.P., solution to 3 of water) is a good disinfectant, but its odour is disagreeable. Lysol is more pleasant. Condyl's fluid is largely used, but it is only deodorant not disinfectant.

The skin round the external genitals should in all cases be protected from the acrid discharges, as the irritation is a source of discomfort. A lotion of equal parts of olive oil and glycerine or of olive oil and lime water, applied after each vaginal injection, serves this purpose well.

PAIN.

This can be effectually relieved only by some preparation of opium; ^{Use of} it is well to delay the habitual use of this remedy as long as possible, ^{Opium.} as it interferes with digestion and nutrition. It may be given as a morphia suppository ($\frac{1}{4}$ of a grain in each) per rectum, or as the liquor morphinæ hydrochloratis by the mouth. We obtain its action most surely and quickly and with the least disturbance of the digestive system by giving it hypodermically. It is desirable to change the narcotic, as even opium gradually loses its effect; the hydrate of chloral, in 20 grain doses, may be used as a substitute. Various local anodynes have been suggested, but are of little use.

Attention to the *general condition* of the patient is very important. ^{General} The three main points are to give a sufficient quantity of nutritious and ^{Treat-} easily digestible food, to keep the bowels regular, and to have the ^{ment.} atmosphere healthy and the surroundings cheerful. Food should be given in small quantities and frequently; milk, eggs and beef-tea should be substituted for more solid food as soon as digestion fails. In the later stages, the bowels should be evacuated by enemata rather than by purgative medicines. The room should be well ventilated by

day and night, and the vaginal injections repeated frequently. Gusserow recommends that during the night a piece of waterproof sheeting be tied round the patient's waist to keep down the disagreeable odour.

TREATMENT OF THE DISEASE.

As before stated our aim here is extirpation. If complete removal be possible, carcinoma will be no longer the incurable disease which haunts the mind of the patient and baffles the skill of the practitioner. The principles of treatment can be best understood by considering the progress of the disease as consisting of three stages: (1) when the disease is present as a germ infiltrating healthy tissue; (2) when the germ has developed into a tissue having the typical carcinomatous structure; (3) when this newly-formed tissue breaks down. The accompanying diagram (fig. 272) illustrates this progress. The three stages are represented by three zones.

Diagram
of spread
of Cancer.




FIG. 272.

DIAGRAM TO ILLUSTRATE THE SPREADING OF CARCINOMA. 1, Healthy tissue infiltrated with germs of Carcinoma—area of lymphatic infection; 2, Carcinomatous tissue fully developed; 3, Carcinomatous tissue breaking down.

The extent of zone 1 is not well defined, for we have no means, unless with the microscope, of ascertaining how far the surrounding tissue is infiltrated. The existence of this area has been emphasised recently by the work of Seelig and others (see p. 489) who have shown that beyond the area where the disease can be recognised by touch and sight, there is lymphatic affection (see fig. 262). The area of zone 2 is more definite; the line *a b c* is well marked, for the carcinomatous tissue when fully formed has characteristics by which it can be recognised from the surrounding healthy tissue by its physical signs. Zone 3 represents the third stage, in which the immediate danger to the patient lies. It is not the *formation* of the carcinomatous tissue which is dangerous, but its *ulceration* with accompanying hæmorrhage and exhausting discharge.

From these facts we deduce the following principles of treatment. *First*, to effect radical cure we must remove zone 1, as well as zones 2 and 3; *i.e.*, we must remove not only the tissue which is evidently

carcinomatous, but also all the surrounding tissue which may contain germs of the disease. Sometimes by a chance the operator has done this through keeping well clear of the evidently diseased part, and thus we can explain the few recorded cases of cure. *Second*, we may anticipate the natural process of breaking down, with its accompanying exhausting results and risks of a fatal hæmorrhage, by destroying the newly-formed carcinomatous tissue as far as it is recognisable. We shall thus save the patient from the effects of the disease until zone 1 has passed into the condition of zone 2 and is beginning to break down. Thus we explain the temporary benefit (for a period measurable by months) derived from the partial excision of the new growth. *Third*, the application of caustics alone may effect the destruction of area 2; but we are not so sure that we are removing the *whole* up to line *a b c*, as we are when using the knife or other cutting instrument. The latter means is preferable because we can make certain that we have reached this line in all cases where it is attainable by operation. *Fourth*, the use of the knife and the application of caustic to the raw surface will, where the disease has spread far, be more effectual than the use of the knife alone; the caustic will now without doubt operate on the area of zone 1 and destroy so far the germs of the disease:—

There are three methods of operative treatment:—

1. Scraping out of diseased tissues, with or without application of caustics,
2. Amputation of the cervix,
3. Excision of the uterus.

SCRAPING OUT OF DISEASED TISSUE, WITH OR WITHOUT CAUSTICS.

We have recourse to this means of treatment (1) in cases in which the disease is not of a form suitable for amputation—when it does not form a pediculated mass but is spreading along the mucous membrane of the vagina, (2) in cases which are too far advanced for amputation of the cervix. This method has the advantage that the carcinomatous tissue is soft and friable compared with the surrounding connective tissue and can be therefore easily scraped away; on the other hand, there is the danger of exciting metastasis.

After scraping away obviously diseased tissue with a spoon (fig. 273) or burning it out with the cautery, the cavity is plugged with iodoform gauze.

In addition to scraping, strong caustic may be applied, to cause the carcinomatous tissue to slough out. Nitric acid or an alcoholic solution of bromine¹ (1-5) have been used, but chloride of zinc² (as a solution

¹ Recommended by Routh (*Brit. Med. Jour.*, Feb. 1880) and Wynn Williams (*Lond. Obstet. Trans.*, Vol. xii., p. 249).

² Recommended by Van de Warker (*Amer. Jour. Obstet.*, March 1884), and by Fränkel who had a case after its use without recurrence during seven years (*Beitr. zur Geburts. Berlin*, Bd. ii., S. 23).

of 1 in 2 or in paste) is the best. Below the tampon of zinc chloride the vagina must be packed with pledgets soaked in an alkali (bicarbonate of soda) to neutralise the superfluous acid that runs down. The packing is left in for six days; after which antiseptic injections are given until the slough comes away several days later.¹

AMPUTATION OF THE CERVIX.

This operation is called for by two sets of circumstances: (a) when the disease is as yet limited to the cervix and there is a distinct line of demarcation above, so that in operating we can cut through healthy tissues; (b) when it has spread so far that, although we cannot operate upon healthy tissue, we are yet justified in removing as far as possible the projecting mass.

The means of amputation are the following:—

Écraseur, or galvano-cautery;
Knife and scissors.

I. ÉCRASEUR OR GALVANO-CAUTERY.

Both of these possess the advantages that they are easy of application and cause less hæmorrhage than the knife, although with the latter we



FIG. 273.

SIMON'S SHARP SPOON.

Écraseur
and
Galvano-
Cautery
compared.

can follow more certainly the line of demarcation. The écraseur has the advantage that it is easily portable, requires no preparation, and is always ready when wanted. In using these, the cervix is laid hold of with volsellæ and drawn down to the vulvar orifice. The chain or wire (if a galvano-cautery² is used) is placed round the cervix as far above the limits of the disease as possible; and tightened slowly so as to crush or burn through the tissues rather than cut them. After amputation, if there is much hæmorrhage, a styptic may be applied to the stump; and the vagina is firmly packed. The packing is not removed for some days as there is danger of secondary hæmorrhage.

II. KNIFE AND SCISSORS. The advantage claimed for this method of operating is that it allows the operator to follow the line of demarcation between the diseased and the healthy tissues; if in the course of

¹ The whole uterus may slough out as in case recorded by Jessett (*Brit. Gyn. Jour.*, 1895, p. 32) which was still without recurrence after four months.

² Pawlik gives the after-history of 136 cases operated on by C. Braun in the Vienna Clinique; the mortality from the operation was $7\frac{1}{2}$ p. c. 26 of the cases were without recurrence two years after operation—the longest period being 19½ years.

the amputation he finds the carcinomatous new-formation extending higher up than he anticipated, he can remove as much more of the suspected part as may be necessary.¹

The cervix may be amputated at the level of the fornix, as described at p. 313. In the *supra-vaginal* amputation,² the cervix is drawn down with a volsella, the knife carried round the anterior fornix, and the bladder separated from the cervix almost up to the utero-vesical pouch of peritoneum. The cervix is now carried forward and the posterior fornix incised in a similar way, the ends of this incision being made continuous with that made anteriorly. Should the peritoneum of the pouch of Douglas be cut into, it is of no consequence. The clearing of the cervix from the cellular tissue above the lateral fornices may be more difficult, from the firmness of the connective-tissue and the presence of the branches of the uterine artery. Hæmorrhage is best




FIG. 274.

LINE OF INCISION AND POSITION OF SUTURES IN THE SUPRA-VAGINAL AMPUTATION OF THE CERVIX (Schroeder).

prevented by transfixing the tissues with an aneurism needle, and ligaturing before cutting through between the ligature and cervix; the cervix being thus freed, the anterior lip is removed and stitched as in fig. 274, the forceps-grasp of the posterior lip steadying the cervix for this manipulation. The sutures of the anterior lip being tied are now used to hold down the cervix, while the posterior lip is being amputated and stitched. Sometimes bleeding from the cervix cannot be arrested by sutures, in which case forceps must be left on.

¹ In amputating the cervix, Stiles' method may be used to tell whether we have cut through above the limits of the disease. A 5 p. c. solution of nitric acid is washed over the cut surface of the piece removed. This makes the carcinomatous tissue stand out opaque-white in contrast to normal fibrous tissue which becomes translucent.

² After Schroeder's method. Hofmeier (*Zeits. f. Geb. u. Gyn.*, Bd. x., S. 269) reporting on 105 cases done in Schroeder's Clinique gives a mortality of 12.3 p. c. Of forty-seven cases, fifteen were without recurrence two years after operation, and ten had not been heard of; after three years twelve were well, and after four years, five. Lewers (*Lancet*, 1888, Vol. i., p. 464) reports ten cases with no deaths, and three without recurrence for one year.

EXCISION OF THE WHOLE UTERUS.

To Freund of Strassburg is due the credit of having first thought out and carried into execution a method by which the whole uterus can be removed. This method has increased the possibility of a radical cure of malignant disease of the uterus, though the number of cases suitable for extirpation is more limited than we should have supposed. The uterus alone can be removed by it, not the glands or connective-tissue in the pelvis to which the disease in the majority of cases soon spreads. But when the disease has originated in the body of the uterus, or beginning at the cervix has extended upwards into the uterus rather than into the vagina or the connective-tissue, the *extirpation of the uterus* holds out the prospect of a radical cure. This may be done

- A. By abdominal incision,
- B. Through the vagina.¹

Freund's
Method.

A. BY ABDOMINAL INCISION (Freund's method). As the high mortality from this method (72 per cent.) has made most operators abandon it for the vaginal method, we shall merely indicate in what the operation consists.

The abdominal cavity having been opened, the uterus is laid hold of and each broad ligament ligatured in three parts, the lowest ligature passing through the lateral fornix of the vagina. The uterus is now cut away from the broad ligaments; and the knife carried through the peritoneum of the utero-vesical pouch and pouch of Douglas into the anterior and posterior fornices so that the whole organ is thus excised. The ends of the ligatures in the broad ligaments are brought through the hole in the roof of the vagina, in which a drainage tube is also placed.

The results of this method of extirpation are according to Gusserow 148 cases with a mortality of 71.6 per cent., according to Duncan 137 cases with a mortality of 72 per cent.²

B. EXTIRPATION THROUGH THE VAGINA. This is now the recognised operation for removal of the uterus in cancer. Till recently the mortality was so great that it was doubtful whether removal of the uterus would ever take its place alongside of removal of the breast in cancer. By the vaginal operation, however, the mortality has been greatly reduced, and the results as to non-recurrence are as satisfactory as for mammary cancer, if not more so. The technique varies according as the ligature or clamp is used. We describe the latter operation first, as for its simplicity of technique, rapidity, and slight loss of blood it is preferable. Its drawback is the sloughing of tissue in the bite of the clamp, so that, as in the extra-peritoneal treatment of the

¹ A third method, which is a combination of these, has been so seldom used that it requires no notice here.

² Several cases of total extirpation of the pregnant cancerous uterus are on record. Sir Spencer Wells in 1881 and Zweifel in 1888 removed one at sixth month, patients recovering in both cases. Schroeder operated at full-time in two cases, and Bischoff in one. All three died.

stump in fibroids, we have here to balance a more rapid operation against a more tardy convalescence.

(a) *Vaginal hysterectomy with clamp*.—The use of the clamp to arrest hæmorrhage in vaginal hysterectomy was, we believe, first suggested by Spencer Wells.¹ But it was the French operators Péan and Richelot,² who first carried it out successfully and elaborated the technique for its use. In Germany it has been strongly advocated by Abel,³ and by Leopold and Theodor Landau.⁴

We shall describe the operation after Doyen's⁵ method, which has given good results in our hands.⁶

His technique differs from Péan's in that he omits the lateral splitting of the cervix, with the preliminary clamping of the base of the broad ligaments which this necessitated to control hæmorrhage; he has also emphasised the mesial splitting of the anterior uterine wall, as greatly favouring the descent of the fundus. The instruments




FIG. 275.

DOYEN'S CLAMP FOR VAGINAL HYSTERECTOMY.

c, Inner face of blade. a, Blades open; b, closed; d, compressed.

required are bistoury and scissors, vaginal spatulæ—of the form shown in fig. 80 (p. 124), several strong volsellæ-like bullet-forceps, clamps of

¹ Ovarian Tumours: London, 1882, p. 526.

² Richelot had done up to August 1895, fifty-eight cases of hysterectomy for uterine cancer, with six deaths. *Derniers résultats de l'hystérectomie vaginale: Annal. de Gyn.*, Dec. 1895.

³ Seventy-nine cases reported on with four deaths.

⁴ (See their beautifully illustrated monograph "Die vaginal Radicaloperation, Technik und Geschichte." Hirschwald: Berlin, 1896.) They have performed 438 vaginal hysterectomies—191 for inflammatory conditions, with a mortality of 2.6 p. c., and 247 for tumours (cancer, sarcoma, fibroid) with a mortality of about 6 p. c. (Macnaughton Jones on Gynecology in Berlin: *Brit. Gyn. Jour.*, Feb. 1897.) Inasmuch as they emphasise the careful and complete removal of tissue, and the clamps are applied at the close of the operation, they call it the "enucleation method" rather than the "clamp method." This term is however equally applicable to Doyen's method. It is worth noting, however, that the clamps are not used in this operation as "preventive," to facilitate the cutting away of tissue without hæmorrhage, but as definite or consecutive, that is, as a permanent means of controlling it. They also draw attention to the fact that the clamps act as a drain, and do not interfere with the closure of the peritoneal cavity which takes place by natural methods (without any sutures) above them.

⁵ La Castration, totale par le vagin (*Extr. des Archiv. Prov. Chirurgie*, Dec. 1892), where he reports on 112 operations—twenty-eight for fibroid, twenty-three for cancer, and sixty-one for other conditions) with six deaths.

⁶ Barbour—Vaginal Hysterectomy by Doyen's Method: *Scot. Med. and Surg. Journ.*, July 1897.

Doyen's
Method of
Vaginal
Hysterec-
tomy.

the pattern in fig. 275. The blades of the clamps are concave and elastic. From their concavity they come in contact first at the points, and only on firm compression of the handle, throughout. (Compare *b* and *d*.) This device ensures compression of the tissue towards the end of the blade which we shall see corresponds to the base of the broad ligament and uterine artery. Further, the blades are grooved on their inner aspect to near the point (fig. 275, *c*), producing a space into which the tissue bulges, which lessens the risk of the instrument slipping. A broad and narrow-bladed clamp are required for each broad ligament; and it is better to have a few in reserve.

For a few days prior to the operation, the vagina should be frequently douched and packed with some antiseptic gauze. Doyen recommends its distension for forty-eight hours, with an air-bag.


At the operation, the patient is placed in the lithotomy posture, the external genitals shaved and thoroughly cleansed (as also the vagina) with soap and water and turpentine, and then douched with corrosive.

(1) The cervix is seized in two volsellæ, placed laterally and as high up as the fornix will allow; and a circular incision made with the scissors, no deeper than the vaginal mucous membrane, round the cervix (Pl. XII., fig. 1).

(2) The pouch of Douglas is opened into by alternately clipping and feeling with the index finger to tell when the peritoneal cavity is reached. When it is opened into, the incision is enlarged by guiding in the closed scissors and then opening them so as to tear the tissues apart. The finger now explores the posterior surface of the uterus, noting the presence of adhesions or sub-peritoneal fibroids.

(3) The utero-vesical pouch is reached in the same way. Great care is needed in the separation of the bladder which is effected by the finger rather than the scissors (Pl. XII., fig. 5). As soon as the cervix is partly exposed, it is divided in the middle line anteriorly (Pl. XII., fig. 2); and the uterus drawn down by laying hold of it by the margins of the longitudinal section (Pl. XII., fig. 2). It is pulled down "hand over hand" as it were, a new grip being taken higher up with a fresh pair of forceps before the forceps with the lower grip are taken off (see fig. 3 in Pl. XII., in which the split anterior wall of the uterus is shown with the points of the succeeding forceps-bites).


Should the uterus be bulky with fibroid masses, a V-shaped incision may be made instead of a single mesial one, and the anterior wall cut out in small portions—1, 2, 3, 4, 5, 6 in Pl. XII., fig. 4. It is essential before cutting out the piece in the bite of the forceps, to take a hold with another pair higher up—before cutting out 1, the forceps are placed on 2 and 3, and so on. The division of the anterior wall lets the uterus collapse and slip out. Fibroid masses in the wall may have to



F. & A. K. Schuster, Edinburgh & London.

VAGINAL HYSTERECTOMY WITH CLAMPS (DOYEN).

- FIG. 1. Incision in fornices. FIG. 2. Uterus drawn down and anterior wall divided.
 FIG. 3. Splitting of anterior wall; FIG. 4 shows the V-shaped incision. FIG. 5. Separation of bladder. FIG. 6. Application of clamp.



be enucleated and sub-mucous or sub-peritoneal tumours torn away to reduce the size of the uterus.

(4) As soon as the fundus appears, it is grasped in forceps and flexed forwards so as to bring the upper margin of the broad ligament down. The left broad ligament is now taken between the index finger and thumb of the left hand (the index being passed (from above) over the upper margin of the ligament down its posterior aspect to its base, until it touches the thumb which has been slipped along its anterior aspect). A large clamp is guided underneath the finger and thumb, care being taken to feel that it grasps the whole ligament and includes nothing besides. It is then closed tight (Plate XII., fig. 6). A second narrow pair of clamps is placed for safety between the first pair and the uterus which is now cut away on that side. The right ligament is treated in the same way and the uterus cut away. The clamps are




FIG. 276.

VAGINAL EXTIRPATION OF THE UTERUS (*Martin*).

The cervix has been drawn downwards with forceps, the pouch of Douglas opened transversely, and row of sutures passed through vaginal fornix and peritoneum.

now brought down into the axis of the vagina, the broad ligaments being thus twisted from their original direction in the axis of the brim to lie in the axis of the outlet.

(5) A gentle douche of weak boracic may now be given, and a tampon of gauze placed between the forceps and also round the handles to keep them from pressing injuriously on the vaginal wall. A turn of the gauze is passed round the two pairs of forceps on either side so as to keep them together.

After-treatment.—The urine is drawn off for forty-eight hours to keep the tampon clean. There is sometimes considerable pain complained of till the clamps are removed, and morphia suppositories may be

required. The large clamps are taken off in forty-eight hours, and the small ones some twelve hours later. The gauze is removed a day later.

Vaginal
Hysterec-
tomy with
Ligature.

(b) *Vaginal hysterectomy with ligature.*—Different operators have introduced various modifications, but these are only in detail. We describe the operation as performed by Martin of Berlin.

1. Place patient in lithotomy posture, empty bladder and thoroughly disinfect genital tract. Let assistants hold anterior and posterior vaginal specula and lateral retractors in position, draw down cervix with volsella and direct it forwards towards pubes. Make a transverse incision through the junction of vaginal mucous membrane with posterior surface of cervix. The pouch of Douglas is thus opened. Then sew the peritoneum and vaginal mucous membrane together by three or




FIG. 277.

VAGINAL EXTIRPATION OF UTERUS (Martin).

Application of first ligature in lateral fornix to control vessels in base of broad ligament.

four sutures parallel to line of incision and slightly behind it (fig. 276).

2. Pass left index finger into pouch of Douglas and press left broad ligament down against vaginal roof. With a large curved needle pass a suture through anterior part of left lateral fornix, through broad ligament above uterine artery, and out again through the vaginal roof close to the outermost suture of the posterior row (fig. 277). Pass a ligature in the same manner, also, on the right side. Then with scissors cut through the bases of the broad ligaments as high as ligatures reach, keeping close to uterus.

3. Draw cervix backwards, and at the line of junction of vaginal mucous membrane with its anterior surface, make a transverse incision down to muscular substance of uterus. Carefully separate bladder

from uterus and open into utero-vesical pouch of peritoneum. Bring vaginal mucous membrane and peritoneum into close apposition by a transverse row of sutures applied as was done posteriorly.

4. With volsella pull down fundus through pouch of Douglas as far as possible. The broad ligaments, and generally the tubes and ovaries, are thus brought into vagina.

5. Now ligature broad ligaments above the level of the uterine arteries. This is done by two or three sutures passed exactly as in the case of that first applied in the lateral fornix, only at successively higher levels in the vaginal roof. The uterus is then cut away, the tubes and ovaries being also removed when possible.

Thus the sutures are all tied on the vaginal surface, and they approximate the serous surfaces of broad ligaments and pelvic floor to one another as well as to the vaginal mucous membrane.

If a wide opening remains, it can be made smaller by a suture on each side. A rubber drainage-tube may be used or not. Dust wound with iodoform and place an iodoform-gauze tampon in vagina.

Czerny brings down fundus through opened-up *utero-vesical* pouch. Olshausen, Leopold and others do *not* draw fundus down, but gradually cut away uterus from broad ligaments after suturing the latter in successive stages from below upwards.

Fritsch operates in the same way, but ligatures the uterine arteries and cuts through base of broad ligaments before opening into peritoneum.

Extirpation of the uterus has also been done by the sacral method, an operation which, though it allows a more thorough clearing out of diseased tissue and glands, will never compete with the vaginal operation. It was introduced by Hochenegg who has collected ninety-eight cases with eighteen deaths—a mortality of 18 p. c.; he himself having done thirty with three deaths, a mortality of 10 p. c.¹

RESULTS OF HYSTERECTOMY FOR CANCER.

In judging of these we must keep in view both the mortality from the operation, and the ultimate result as to non-recurrence of the disease.

It is only recently that this operation has found favour in this country, and we have not yet had recorded a sufficient number of cases to form a general conclusion.

Jessett² records seventy cases of vaginal hysterectomies for cancer, and sarcoma with six deaths. Forty-three were operated on more than two years previously, and of these twelve were known to be free from recurrence, while eleven were without it when last heard of.

¹ Quoted by Buechler: *Zeits. f. Geb. u. Gyn.*, 1894, Bd. xxx., Hft. 2, S. 394.

² The early Diagnosis of Malignant Disease of the Body of the Uterus, etc.: *Brit. Gyn. Jour.*, 1896-97, p. 327.

Purcell,¹ who did the first vaginal hysterectomy for cancer in England (the patient being alive still, twelve years after operation), has had sixty-three operations with twelve deaths, eighteen of these being without recurrence after two years. Rutherford Morison² records nine successful cases.

Kelly's results are given by Russell;³ forty-seven cases, forty vaginal, four combined, and three abdominal, with a mortality of 10 p. c. in the operation.

Mortality
of Vaginal
Hysterec-
tomy.

On the Continent, however, the operation has been extensively performed, and the mortality has been materially reduced by improved methods. Fehling⁴ finds the mortality for 778 extirpations done by thirteen of the leading German operators to be 9.1 p. c., while in the hands of individual men⁵ it has fallen as low as 3 p. c.

While in this country, cases suitable for operation only exceptionally present themselves, patients are apparently seen abroad at a much earlier period. Thus Krugenberg finds 3.7 p. c. of cases of malignant uterine disease in 24,887 patients, and of these 31.6 p. c. were treated by radical operation; while Kaltenbach was able to operate on 40 p. c. (134 out of 452) of the patients who, from 1887-94, sought advice at the Halle-clinique for cancer. So also in Gusserow's⁶ clinique, 15.4 p. c. of the 577 cases were operable. As to the permanent benefit, the operation fully justifies itself. Though Pozzi says "in my opinion hysterectomy is only a palliative treatment of longer or shorter duration—the average being a year," the results of other operators have been more encouraging. Thus of ninety-two cases operated on by Kaltenbach, which were followed for more than three years, twenty-three were without recurrence; others had died during that period from other causes, without recurrence, while nine had gone for five years without it. Krugenberg⁷ finds after three years 37.5 p. c. without recurrence; after four, 29.5 p. c. This was for cervical cancer, and the results for corporeal are even better, as after five years 66.7 p. c. were without recurrence. Leopold⁸ found that 55.8 p. c. of his 164 cases were without recurrence after four years.

Compari-
son with
Amputa-
tion.

Hysterectomy and Amputation of the Cervix.—The relative advantages of the major operation of extirpation of the uterus, bears to the minor one of amputation of the cervix is *sub lite*. For a time the mortality for extirpation was so high that it could not compete with the minor operation. But now that it has been reduced the question is raised whether extirpation is not to be preferred in all cases. In this country, supra-vaginal amputation is still preferred in suitable cases.

¹ Discussion on Jessett's paper.

² Notes of nine consecutive cases of extirpation of the uterus for cancer:—eight vaginal operations by forceps and ligature, and one abdomino-vaginal.

³ John Hopkins Bull., 1895.

⁴ Quoted by Buecheler: *Zeits. f. Geb. u. Gyn.*, 1894, Bd. xxx., S. 394.

⁵ *e.g.*, Kaltenbach with 138 cases.

⁶ Oskar Müller—Zur Lehre von Carcinoma uteri: *Charité Annal.*, xviii., S. 529.

⁷ Die Resultate der oper. Behand. des Carcin. und Sarcom. Gebärd.: *Zeits. f. Geb. u. Gyn.*, 1892, S. 108.

⁸ Die Dauererfolge der vaginalen Totalexstirpation, etc.: *Geb. u. Gynäk.*, Bd. ii., aus der königl. Frauenklinik, Dresden—Leipzig, 1895.

Thus Thornton¹ favours it. And Jessett² recommends it where the disease is limited; as in seventeen cases there had been no recurrence after operation. Hofmeier also could point to immunity given for at least four years to one-third of the cases of supra-vaginal amputation, which is as good a result as has been got from hysterectomy.

Metastasis and Recurrence after Operation—Considerable attention Metastasis
and Re-
currence. is being directed to this subject at present from its important bearing on operative treatment. The unsatisfactoriness of operations for cancer of the uterus, as for cancer elsewhere, lies in the liability to the reappearance of the disease; and many question the justifiability of subjecting a patient to such a grave operation as extirpation of the uterus unless more satisfactory results as to non-recurrence, for at least a reasonable time, can be obtained. Hence the importance of inquiry as to the frequency of the reappearance of cancer after operation.

The type and seat of the morbid process have both a bearing on this. Thus recurrence is less frequent after epithelioma than after carcinoma. Further, there is more immunity after operation in superficial cancer of the vaginal portion and cancer of the body of the uterus (whatever form the cancer takes) than in ordinary cancer of the cervix. In the former case it may be due in part to the type (epithelioma being more frequent there), but it is also due to a less rapid invasion of the lymphatics, which is the chief cause in the latter case (see p. 491).

Local recurrence may be accounted for in three ways: (1) by incomplete removal—cancer cells being left in the lymphatics of apparently healthy tissue; (2) by unclean operation—cancer cells being inoculated on the raw surfaces during the operation; and (3) by reappearance of the disease from the same cause which originally produced it; further, there may be metastasis in remote organs. The rapidity of the return suggests that it is due to one of the first two causes. The second of these has been emphasised by Winter and Mackenrodt; and has this to be said for it, that in operating for cancer of the cervix the operator is often working in a hole with broken-down carcinomatous débris. Hence the operation is not clean, like the excision of a non-ulcerating breast. Further, the peritoneum, which is peculiarly susceptible to implantation with malignant cells, is also exposed. To diminish the risk, Winter recommends removal of as much of the broken-down tissue as possible before extirpating, and the cleansing of the field with an alcoholic solution of corrosive sublimate.

Incomplete removal of affected lymphatics certainly plays an important part, and the study of the exact seat of the recurrence has a

¹ *Brit. Med. Jour.*, Vol. i., p. 263.

² *Brit. Gyn. Jour.*, 1893-94, p. 389. See also his paper "On twenty-five cases of supra-vaginal amputation" (*Brit. Gyn. Jour.*, 1892-93, p. 353).

peculiar interest in connection with its relation to lymphatic distribution.¹ The suggestion of operations² with more complete removal of the parametric tissue, shows the importance of this.

¹ See the papers by Seelig, Roger Williams and Russell. Stiles in an elaborate research on cancer of the breast, in which he has studied especially its relation to the lymphatics as Seelig has done in the case of the uterus, believes that here also recurrence is due to cancer cells left in the lymphatics. He notes that the endothelium is not affected by these cancer plugs. Further, he believes in inoculation, as he says that a knife once used to cut cancerous tissue, should not be used again at the same operation. "Contribution to the Surgical Anatomy of the Breast:" *Edin. Med. Jour.*, June 1892.

² As by Kelly, who passes a bougie into the ureter and removes the uterus by the abdominal method, ligaturing the uterine artery an inch from the uterus before it gives off the vaginal branch. (Clark—A more radical method of performing hysterectomy, etc.: *Johns Hopkins' Hospital Bull.*, Vol. vi.) He mentions that out of twenty vaginal hysterectomies, only four showed on microscopic examination that the cut surface was healthy.

CHAPTER XLIII.

CARCINOMA UTERI (OF BODY): ADENOMA MALIGNUM.

LITERATURE.

Breisky and Eppinger—Prager med. Wochenschrift, S. 78, 1877. *Gusserow*—Neubildungen des Uterus, S. 254: Stuttgart, 1855. *Hofmeier*—Zur Anatomie und Therapie des Carcinoma Corporis Uteri: Zeitsch. f. Geb. u. Gyn., Bd. xxxii. *Schroeder*—Die Krankheiten der weiblichen Geschlechtsorgane, S. 295. *Simpson, Sir J. Y.*—Selected Obstetrical and Gynecological Memoirs, edited by Watt Black, p. 769. *Veit*—Zeitschrift. für Geburts. und Gyn., Bd. i., S. 467. Zur Kenntniss des Carcinoma Corporis Uteri: Centralb. f. Gyn., Bd. x., S. 173. See also literature of Chaps. XL. and XLII.

PATHOLOGY AND ETIOLOGY.

CARCINOMA affects the body of the uterus much more rarely than the cervix; in only 13 out of 686 cases of uterine cancer, that is in rather



FIG. 278.

UTERUS EXTIRPATED FOR CANCER; no recurrence five years after operation (*Hofmeier*).

less than 2 per cent., was the disease situated in the body of the uterus (*Schroeder*).

As in the cervix, the disease originates either in the substance of the walls of the uterus or in the mucous membrane. In the former case, it begins as localised nodules which grow rapidly and produce bulging of the mucous membrane or of the peritoneal coat, but do not tend to ulcerate. When in the mucous membrane, it causes a uniform swelling (fig. 278) or, more usually, projects in polypoidal masses (fig. 279). Fig. 278 from Hofmeier, shows a uterus extirpated for cancer; the disease had not recurred within five years after the operation.

By Eppinger and Ruge the disease has been directly traced to the epithelium of the uterine glands; these first hypertrophy, and then their proliferating epithelium passes into carcinomatous epithelial cells. The new-formation ulcerates, so that the wall of the uterus becomes converted into an excavated surface with a hard base. Adhesions rapidly




FIG. 279.

CARCINOMA OF THE BODY OF THE UTERUS. The uterine cavity is increased in size, but the cervix is undilated (Sir J. Y. Simpson).

form with neighbouring organs, while secondary deposits may develop in the peritoneal cavity.

As to *Etiology*, what has been said of carcinoma of the cervix applies here with two additional facts: (1) It occurs rather later in life than cancer of the cervix; and (2) is more frequent in nulliparæ.¹

SYMPTOMS AND DIAGNOSIS.

Pain. Again, as in carcinoma of the cervix, the symptoms are pain, hæmorrhage, and foetid discharge. 1. *Pain*, in contrast with carcinoma of the

¹ Taking Veit's two series of cases together, we have out of 80 cases, 31 between 50 and 60, and 21 above 60 years of age (cf. table in fig. 267); and of 72 cases, 38 were childless.

cervix, is always an early symptom. Sir J. Y. Simpson drew attention to periodic attacks of severe pain as characteristic of cancer of the body. This is not always present and is probably due to uterine contractions set up by accumulation of secretion (*Veit*). 2. *Hæmorrhage* is also ^{Hæmorrhage.} present at an early stage; it takes the form of profuse menorrhagia, because the mucous membrane from which the menstrual flow takes place is diseased. 3. *The discharge* is usually profuse and becomes after ^{Discharge.} a time foetid. Sometimes it is watery and not offensive; rarely is it altogether absent.

On vaginal examination, the cervix is found to be either normal (fig. 279) or dilated. The uterus is enlarged, and may be freely movable or may be fixed by adhesions. The sound shows the cavity to be enlarged and may reveal irregularity of the mucous membrane; its introduction is followed by hæmorrhage. The condition of the mucous membrane is more precisely ascertained by examination *with the finger after dilatation of the cervix* with a tent. In the majority of cases, certainty of diagnosis is possible only through *microscopic examination* of fragments removed by the curette. Should these show merely hypertrophied glands, we must remember that this is sometimes a transition stage to malignant disease. Typical carcinomatous cells are seen at fig. 269.

The *Differential Diagnosis* must be made from—

Portions of retained placenta,
Sloughing submucous-fibroid,
Hæmorrhagic endometritis,
Deciduoma malignum.

During the period of sexual activity, differential diagnosis is often difficult, and can be made only by exploring the uterus with the finger, and microscopic examination of the tissue removed. After the menopause, the recurrence of hæmorrhage is an important diagnostic. The microscope is, when available, the most reliable guide.

TREATMENT,

As to the treatment of the symptoms, this is the same as in Carcinoma of the Cervix (*v.* Chap. XLII.). As to the treatment of the disease, the scraping away of the polypoidal masses with the curette or sharp spoon gives temporary relief from the hæmorrhage and discharge. The only hope of cure lies in extirpation of the uterus (*v.* p. 516).¹ The results of hysterectomy are better, as regards non-recurrence, for cancer of the body than for that of the cervix. See p. 491.

¹ Of 7 cases (1 by Schroeder) done by Veit, 1 died after operation; and of 4 cases followed, 1 had recurrence in first year, 2 in second, and 1 not after three years.

ADENOMA MALIGNUM.

LITERATURE—*Williams, Sir J.*—Cancer of the Uterus: London, 1888. *Länderer*—Eine Adeno-carcinom des Corpus Uteri: Zeits. f. Geb. u. Gyn., Bd. xxv., 1892. *Hofmeier*—Zur Anatomie und Therapie des Carcinoma Corporis Uteri: Zeits. f. Geb. u. Gyn., Bd. xxxii., 1895. *Sinclair, W. J.*—Article on Malignant Disease of the Uterus: Clifford and Allbutt's System of Gynecology, London, 1896. *Ruge and Veit*—Zeits. f. Geb. u. Gyn., 1881, Bd. vi., S. 302. *Beyea*—Malignant Adenoma of the Corpus Uteri: Amer. Jour. Obstet., Vol. xxxiii., 1896, p. 196.

It is doubtful whether malignant adenoma should be considered as an affection distinct from carcinoma of the uterus.¹ It usually affects the body of the uterus; being, unlike carcinoma, rare in the cervix.² Its clinical history closely resembles the latter, and the pathological appearances are not clearly marked off—adenoma passing into carcinoma. There is an a-typical hypertrophy, the lumina of the glands becoming enlarged, irregular, and often communicating with each other. The columnar epithelium loses its cilia and becomes stratified, and the gland may be converted into an epithelial plug. It occurs late in life, begins insidiously, and runs a chronic course. While hæmorrhage is present, there is not the foetid discharge so characteristic of cancer. Cases have been described by Matthews Duncan³ and more recently by Länderer, Beyea, and Sinclair.

¹ At the discussion on this subject before the Berlin Gynecological Society opinions were divided; while Hofmeier advocated the distinction, Leopold maintained that the term adenoma can mean only a benign glandular formation: as soon as the glandular development is a-typical, it ceases to be an adenoma. *Cent. f. Gyn.*, 1891, S. 438-440.

² Gebhard could collect only 6 cases in which the cervix was affected: *Zeits. f. Geb. u. Gyn.*, Bd. xxxiii., Hft. 3.

³ Quoted by Williams, *op. cit.*, p. 100.

CHAPTER XLIV.

SARCOMA UTERI: DECIDUOMA MALIGNUM.

By sarcoma we understand a *connective-tissue* tumour of an embryonic type. As we trace back carcinoma to the epithelium and true myoma to the muscular fibre, so we trace back sarcoma to the connective tissue. Nature of Sarcoma.

For the recognition of sarcomata as of connective tissue origin and the limitation of the term to malignant tumours of this type, we are indebted to Virchow. Formerly they were known in English literature as "recurrent fibroids;" the existence of this form of tumour in the uterus was recognised and fully described by Hutchinson (1857).

SARCOMA UTERI.

Literature—Chrobach—Beitrag zur Kenntniss des Uterussarkoms: Archiv f. Gyn., Bd. iv., S. 549. Clay, J.—On diffuse Sarcoma of the Uterus: Lancet, Jan. 1887. Galabin—Lond. Obst. Trans., Vol. xx. Gusserow—Die Neubildungen des Uterus, S. 158: Stuttgart, 1885. Jacobasch—Vier Fälle von Uterussarcom: Zeitschrift f. Geburts. u. Gyn., Bd. vii., Hft. 1. Kahlden, v.—Das Sarkom des Uterus: Zeigler's Beiträge zur Patholog. u. Anatom., etc.: Bd. xiv. Kunert—Ueber Sarcoma Uteri: Arch. f. Gyn., Bd. vi., S. 29. Pick—Ueber Sarcom des Uterus und der Vagina, etc.: Archiv f. Gyn. (1894), xlv., S. 191. Rogivue—Du Sarcôme de l'utérus: Inaug. Dissert., Zürich, 1876. Schroeder—Die Krankheiten der weiblichen Geschlechtsorgane, S. 320; Leipsic, 1886. Simpson, A. R.—Contributions to Obstetrics and Gynecology, p. 240: Edinburgh, 1880. Spiegelberg—Sarcoma Colli Uteri hydropicum papillare: Archiv f. Gyn., Bd. xiv., S. 178. Ein weiterer Fall: Ibid., Bd. xv., S. 437. Thomas—Sarcoma of the Uterus, Lond. Obst. Journ., Vol. ii., 1875, p. 437. Virchow—Die krankhaften Geschwülste: Bd. ii., S. 350. Williams, Whitridge J.—Contributions to the Histology and Histogenesis of Sarcoma of the Uterus: Amer. Jour. Obstet., 1894, Vol. xxix., p. 721. Winkler—Ein weiterer Fall von Sarcoma papillare hydropicum Cervicis et Vaginæ: Arch. f. Gyn., Bd. xxi., S. 309. For a full *résumé* of the literature see Williams' and Pick's papers.

PATHOLOGY.

Unlike carcinoma, sarcoma rarely occurs in the cervix; in the larger proportion of cases it is in the *body* of the uterus.

It occurs in two forms:—

1. Diffuse sarcoma of the mucous membrane;
2. Circumscribed fibrous sarcoma.

Diffuse
Sarcoma.

The *diffuse sarcoma of the mucous membrane* arises from the sub-epithelial connective tissue. It appears as a general swelling of the mucous membrane which becomes soft and crumbly, or as irregular foldings or knobby projections into the uterine cavity; sometimes these projections have a polypoidal and apparently circumscribed character (fig. 280) so that this form passes insensibly into the fibrous. The




FIG. 280.

SARCOMA UTERI WITH TUMOURS IN THE VAGINA—from a specimen in the Pathological Institute at Strassburg (*Gusserow*).

masses have a greyish-white brain-like appearance, and soft pulpy consistence. The mucous membrane may be broken down but is not deeply excavated as in carcinoma. On microscopic examination the mucous membrane is seen to be infiltrated with masses of closely set round cells, more rarely spindle-cells. Epithelial-cell proliferation

may complicate this form of sarcoma and bring it into close relation with carcinoma. Klebs has proposed to call such forms carcino-sarcomata; but v. Kahlden finds no evidence of the occurrence of a mixed form of tumour.

The *circumscribed fibro-sarcoma* arises in the muscular coat; like the fibroid it may be submucous, interstitial, or sub-peritoneal, and is found usually in the body, rarely in the cervix. The tumours are of a firm consistence, and feel like knots in the muscular wall of the uterus or project as polypi into its cavity; they thus resemble small fibroids, but do not usually have a capsule. Microscopically they consist of a localised sarcomatous, generally round-celled, infiltration (fig. 281).

A case of melanotic sarcoma of the uterus has been described by Williams.¹ In some cases sarcoma is apparently a *degeneration of a fibroid tumour*, as in a specimen described by A. R. Simpson.² Similar cases have been reported by Ballantyne,³ Chrobak, and Müller.⁴ Sar-




FIG. 281.

MICROSCOPIC SECTION OF THE MUCOUS MEMBRANE OF THE UTERUS IN A CASE OF SARCOMA (Schroeder).
s Sarcomatous tissue; c small-celled infiltration; g uterine glands.

comatous cells have also been traced to the intima and adventitia of the vessels by Kleinschmidt⁵ in a case of sarcoma of the cervix; and Williams⁶ has traced them to the transformation of the muscular fibre of a myoma.

Secondary nodules may form in the vagina (fig. 280) and peritoneal cavity. Sometimes the peritoneum is affected by continuous spreading of the new growth outwards towards the peritoneal covering; here it causes adhesions, through which the sarcomatous infiltration may extend to other organs (Gusserow). A. R. Simpson records an unique case in which the infiltration spread along the mucous membrane of the Fallopian tubes⁷ (fig. 282), so that from their fimbriated ends there

¹ *Op cit.*, p. 755.

² *Edin. Med. Jour.*, Nov. 1884.

³ Zur operativen Behandlung der Uterusmyome: *Archiv f. Gyn.*, Bd. vi., S. 125.

⁴ Ueber primäres Sarkom der Cervix Uteri: *Archiv f. Gyn.*, Vol. xxxix., p. 1, 1891.

⁵ *Op. cit.*, in which microscopic drawings are given showing the transition.

⁶ Colman describes a case in which the growth extended under the mucosa of the tube and displaced it: *Amer. Jour. Obstet.*, 1893, Vol. xxviii., p. 811.

⁷ *Loc cit.*, p. 243.

projected "round masses, having the appearance of the thrombus projecting from a small vein into a larger trunk." The uterus was of the size of a four-months' pregnancy.

Co-existence of
Inversion.

A. R. Simpson draws attention to the frequency of *inversion of the uterus* as the result of sarcoma. We referred to it as a rare complication of pediculated submucous fibroid tumours. In sarcoma, it appears to occur more frequently—in four out of forty-eight cases. He attributes this to the paralysis of the muscular wall of the uterus through sarcomatous infiltration, and to the peculiar dilatability of the cervix observed in some cases.

Sarcoma of
the Cervix.

Sarcoma of the cervix is rare; in Winkler's paper, eight cases are referred to besides his own. Two of these were spindle-celled, the




FIG. 282.

SARCOMA UTERI INVADING THE FALLOPIAN TUBES and projecting from their fimbriated ends
(A. R. Simpson).

rest round-celled sarcoma. Whitridge Williams has collected only eighteen cases in addition to those of the special "grape-like form" mentioned below. In this respect sarcoma resembles fibroid tumour which we have seen to be relatively rare in the cervix.

A special form has been described as sarcoma papillare hydropicum cervicis. It grows as a papillary tumour which fills the vagina and may project outside the vulva. The cells are embedded in an abundant intercellular substance which stains faintly, is granular, and traversed by delicate threads. It has been erroneously described as a myxosarcoma; in Spiegelberg's cases, it was shown that this intercellular substance was not mucin but coagulated lymph. Other cases have

been described by Pfannenstiel¹ and Pick.² From the form of the tumour, it has also been called "grape-like"—sarcoma botryoides; the dropsical condition being due to interference with the circulation in the necks of the papillæ, which thus become œdematous as they lie free in the vagina. It is of interest as containing sometimes glandular structures, striped muscle, and even hyaline cartilage. It is not clear whether these arise from changes in the sarcoma cells, or whether this tumour may not be developed from embryonic elements.

Metastatic deposits, though rare, are found more frequently in fibro-sarcoma than in diffuse spreading sarcoma. They have been found in the lymphatic glands, lungs, liver, and vertebræ. Perforation of the uterine wall and death from hæmorrhage³ and peritonitis⁴ have also been noted.

ETIOLOGY AND FREQUENCY.

Of the reason why a source of irritation should lead the connective tissue to produce a sarcomatous new-formation, we know as little as why the same cause produces a carcinomatous new-formation from the epithelium.

As to its frequency, a sufficient number of cases has not yet been collected to form any generalisation. It is, however, so rare that every carefully observed case which has been authenticated by microscopic examination should be placed on record.

Age has the same predisposing influence as in fibroma and carcinoma. Of seventy-six cases which we have collected from the literature, we find that

4	were	under	20,
5	„	between	20 and 30,
17	„	„	30 „ 40,
31	„	„	40 „ 50,
19	„	„	50 „ 60,
4	„	„	60 „ 70,
1	was	above	70.

The number of sterile patients among those affected with sarcoma (twenty-five out of sixty-three) is noteworthy; in this respect it contrasts with carcinoma⁵ (*Gusserow*).^a

¹ Das träubige Sarkom des Cervix Uteri: *Virchow's Archives*, cxxvii. (1892), S. 305, where he refers to twelve cases.

² *Op. cit.* Pick's case was in a child two years of age. A case in a child of three has been recorded by J. C. Smith: *Amer. Jour. Obstet.*, 1893, Vol. xxvii., p. 577.

³ *Jacobasch. Op. cit.*

⁴ By Weber and Dressler—quoted by Williams.

⁵ In seventy-four cases of sarcoma, twenty-five were sterile, and sixteen had less than three children.

SYMPTOMS.

The following symptoms characterise the early stage, in which the patient seeks advice :—

1. Hæmorrhage,
2. Absence of pain,
3. Watery non-offensive discharge,
4. Cachexia.

Hæmorrhage.

Hæmorrhage appears first as increase of the menstrual flow, or as irregular hæmorrhages after the menopause. As the new-formation *does not ulcerate* rapidly like carcinoma, the increased menstruation is due to hyperæmia of the mucous membrane (*Clay*).

Pain.

The *absence of pain* in the early stage is remarked on by Clay and A. R. Simpson; in this respect it differs from intra-uterine cancer. According to Gusserow, on the other hand, pain is frequently present and that of an intense and rending character. This apparent discrepancy of opinion may be explained by the varying progress of the infiltration. In the spreading of carcinoma, we noted that pain was most severe when the disease was extending upwards and compressing the nerve endings in the uterus and connective tissue.

Discharge.

The free rice-watery discharge has a slight odour but is not nearly so offensive as in carcinoma; this is due to the fact that there is not the same rapid ulceration and necrosis of tissue. When the disease has progressed further, the discharge becomes equally foetid. The presence in the discharge of *greyish-white shreds*, like particles of brain matter, is diagnostic; under the microscope these are seen to consist of small portions of sarcomatous tissue.

Cachexia.

Cachexia is of importance as it helps us to distinguish developing sarcoma from a non-malignant polypus; the drain from the latter may make the patient gradually anæmic, but there are not the loss of flesh, the loss of appetite, and the rapid failure of strength, which point to malignant disease.

DIAGNOSIS.

If the tumour project through the os, the diagnosis is not difficult. The age of the patient with the symptoms given above and the existence of a *soft friable* pediculated tumour which springs *from the body* of the uterus, will point to the diagnosis; a portion, detached with the nail, shows the characteristic microscopical structure. When nothing projects through the cervical canal, we dilate it by the rapid method described at p. 480. The finger recognises a soft friable condition of the mucous membrane, or a distinct polypoidal tumour, or a localised thickening in the walls.

The uterus is in some cases distinctly enlarged and may reach half-way to the umbilicus or lie retroverted; in the early stages it is movable, but it soon becomes fixed.

The sound shows the cavity to be enlarged; its use causes hæmorrhage.

The differential diagnosis is here often very difficult, as these conditions are also present in—

Differential
Diagnosis.

Chronic endometritis (hæmorrhagic type),
Small fibroid tumours (interstitial or polypoidal),
Carcinoma,
Deciduoma malignum.

Curetting the surface, with microscopic examination of the scrapings, will help us in the first case.




FIG. 283.

SCRAPINGS FROM A FIBROID TUMOUR to show the size and form of the muscular fibre, their rod-shaped nuclei—stained, $\frac{2}{3}\frac{1}{2}$; drawn by S. Delépine.



FIG. 284.

SCRAPINGS FROM A SPINDLE-CELLED SARCOMA to show the larger size of the spindle cells and their oval nuclei—stained, $\frac{2}{3}\frac{1}{2}$; drawn by S. Delépine.

The removal of the polypoidal mass, with the finger nail or nail-curette, will enable us to examine its nature; the possibility of both conditions being present, polypoidal fibroid + commencing sarcomatous degeneration, must be remembered. With an interstitial thickening, we can only watch the progress of the case.

In carcinoma of the fundus, there is generally excavation of the uterine wall and the base of the ragged surface is harder than in sarcoma. The examination of scrapings is not always decisive, as the cells found in sarcoma sometimes closely resemble epithelial cells.

For deciduoma malignum, see p. 542.

PROGNOSIS.

The prognosis is grave. Compared with carcinoma, its development is not so rapid nor are the symptoms of pain and offensive discharge so aggravated in the early stage. In two of the cases recorded by A. R. Simpson the patient survived for four years after the diagnosis of sarcoma was made out, and Gusserow mentions a case where the course was prolonged for ten years.

The temporary relief procured by removal is longer of duration than in carcinoma. When it returns, the development of the new tumour is more rapid than that of the first growth. Cases of non-recurrence, after hysterectomy, are rare.

As to the communication of the prognosis to the patient and friends, see under Carcinoma.

TREATMENT.

The tumour or uterus should be removed as soon as we suspect malignancy.

The cervix should be well dilated so as to allow the finger to pass freely into the uterus. Gradual dilatation is preferable; injury of healthy mucous membrane in dilating or curetting should be avoided as sarcomatous cells have become engrafted on a fresh wound surface.

When circumscribed and polypoidal, the tumour may be removed with the scissors; when diffuse the uterus may be curetted. Curetting, however, gives only such temporary relief that it is now chiefly used as an aid to diagnosis. Extirpation of the uterus per vaginam, as described at p. 516, is the only treatment here, as in cancer.

DECIDUOMA MALIGNUM (SARCOMA DECIDUO-CELLULARE).

Literature—*Apfelstedt and Aschoff* (31), (32)—Ueber bösartige Tumoren der Chorionzotten: *Archiv f. Gyn.*, Bd. I., S. 511. *Bacon* (23)—A Case of Deciduoma Malignum: *Amer. Jour. Obstet.*, 1895, p. 679. *Cazin* (53)—Des déciduomes malins: *La Gynécologie*, 1896, p. 15. *Chiari* (1), (2), (3)—Ueber drei Fälle von primärem Carcinom im Fundus und Corpus des Uterus: *Wien Med. Jahrb.*, vii., 1877. *Chrobak* (47)—Demonstration eines per vaginam extirpirten Uterus: *Cent. f. Gyn.*, 1896, S. 1281. *Cock, J.* (55)—A Case of Deciduoma Malignum, etc.: *Brit. Med. Jour.*, 1896, p. 1819. *Eden*—Deciduoma Malignum, a Criticism: *Lond. Obstet. Trans.*, 1896, p. 149. *Fraenkel* (36)—Das von dem Epithel der Chorionzotten ausgehende Carcinom des Uterus (nach Blasenmole): *Archiv f. Gyn.*, Bd. xlviii., S. 80. *v. Franqué* (44)—Ueber einige bösartige Geschwülste, etc.: *Zeits. f. Geb. u. Gyn.*, 1896, xxxiv., Hft. 2. *Freund* (43)—Ueber bösartige Tumoren der Chorionzotten: *Zeits. f. Geb. u. Gyn.*, 1896, xxxiv., Hft. 2. *Gottschalk* (14)—Das Sarcom der Chorionzotten: *Archiv f. Gyn.*, Bd. xlvi., S. 1, and Ueber das Sarcoma chorion-deciduo-cellulare (Deciduoma Malignum): *Berlin klin. Woch.*, 1893, S. 87 and 116, and *Archiv f. Gyn.*, li., S. 56. *Götze* (42)—Ueber die Tumoren der Eihäute: *Inaug. Diss. Halle*, 1895, and *Cent. f. Gyn.*, 1896, S. 699. *Guttenplan* (15) Ein Fall von hämorrhagischen Sarcom des Uterus und der Vagina, mit Metastasen in den

Lungen: Inaug. Diss. Strassburg, 1883. *Hartmann et Toupet* (52)—Des Conséquences tardives de la rétention partielle ou totale de Placenta, etc.: *Annales de Gyn.*, 1895 (Tome xliii.), p. 285. *Kaltenbach* (16)—Erfahrungen über Uterussarcom, Beil. z. Cent. f. Gyn., 1890, S. 131. *Klien* (34)—Ein Fall von Deciduo-Sarcoma-uteri giganto-cellulare: *Archiv f. Gyn.*, xlvii., 1894, S. 243. *Koettnitz* (51)—Ueber chorio-deciduale Tumoren malignen Characters: *Deut. med. Woch.*, 1893, S. 497. *Kuppenheim* (35)—Ein Fall von Sarcoma deciduo-cellulare: *Cent. f. Gyn.*, 1895, S. 916. *Lebensbaum* (19)—Krebs der Vagina als Impfmetastase bei Corpus carcinom: *Cent. f. Gyn.*, 1893, S. 112. *Lindfors* (49)—Fall von Deciduoma malignum nach Blasenmole: *Cent. f. Gyn.*, 1897, S. 6. *Löhlein* (17)—Sarcom deciduo-cellulare nach vorausgegangenem "Myxoma Chorii:" *Cent. f. Gyn.*, 1893, S. 297, *Ibid.*, 1894, S. 484. *Lönnberg und Mannheimer* (40) (35)—Zur Kasuistik der bösartigen "serotinalen" Uterusgeschwülste: *Cent. f. Gyn.*, 1896, S. 474. *Maier, R.*—Ueber Geschwülstbildungen mit dem Bau des Decidualgewebe: *Virchow's Archives*, Bd. lxxvii., S. 65. *Marchand* (25), (26)—*Monats. f. Geb. u. Gyn.*, 1895, Bd. i., S. 513. *Menge* (20), (21)—Ueber Deciduosisarcoma Uteri: *Zeits. f. Geb. u. Gyn.*, xxx., Hft. 2, S. 323. *Meyer* (5)—Ein Fall von zerstörender Wucherung, etc.: *Archiv f. Gyn.*, Bd. xxxiii., S. 58. *Morison, J. R.* (29)—A Case of Deciduoma Malignum: *Lond. Obst. Trans.*, 1896, p. 130. *Müller* (13)—Ueber das Deciduoma malignum: *Verhandl. d. Deutsch. Gesell. f. Gyn.*, iv., Congress, S. 341. *Neumann* (45), (46)—Ein Fall von Syncytioma malignum: *Cent. f. Gyn.*, 1896, S. 1225 and 1233, and Beitrag zur Lehre von "malignen Deciduom:" *Monats. f. Geb. u. Gyn.*, Bd. iii., Hft. 5, and *Cent. f. Gyn.*, 1897, S. 26. *Nové-Josserand and Lacroix* (18)—Sur le déciduome malin: *Annal. de Gyn.*, et d'Obst., tome xli., Fev. et Mar., 1894. *Pestalozza* (10), (12)—Contributo allo Studio dei Sarcome dell' Utero: *Il Morgani*, Sept. 1891. (11) Di un raro esito remoto della mola vesicolare: *Il Morgani*, Oct. 1891. (37), (38), (39)—Sarcoma deciduo-cellulare: *Annal. di Ostet. et Gyn.*, Nov. 1895, and *Cent. f. Gyn.*, 1896, S. 175. *Pfeifer* (4)—Ueber eine eigenthüm. Geschwulstform des Uterusfundus (Deciduoma malignum): *Praeger med. Woch.*, 1890, No. 26. *Resinelli* (48)—Deciduoma Malignum: *Gaz. des Hôp. de Méd. et de Chir.*, 1896, No. 29. *Runge* (33)—Ein neuer Fall von bösartigem Tumor der Chorionzotten: *Archiv f. Gyn.*, Bd. li., S. 185. *Sänger* (6)—Zwei aussergewöhnliche Fälle von Abortus: *Cent. f. Gyn.*, 1889, S. 132, and Ueber Sarcoma Uteri deciduo-cellulare, etc.: *Archiv f. Gyn.*, xlv., 1893, S. 89. Ueber Deciduoma: *Verhandlungen der Deutschen Gesell. f. Gyn.*, Bd. iv., S. 333. *Schauta* (22)—Ein Fall von Sarkom deciduo-cellulare: *Cent. f. Gyn.*, 1895, S. 248. (54) Demonstration eines Blasenmole: *Cent. f. Gyn.*, 1897, S. 53. *Schmorl* (7), (8), (9)—Ueber malignes Deciduom: *Centralb. f. Gyn.*, 1893, S. 169. (50) *Cent. f. Gyn.*, 1897, S. 90. *Spencer* (28)—A Case of Deciduoma Malignum: *Lond. Obst. Trans.*, 1896, p. 135. *Quarterly Jour. of Med.*, July 1896. *Superno* (27)—*Annali di Ost. e Gin.*, 1895, p. 775. *Tannen* (30)—Ein Fall von Sarcoma uteri deciduo-cellulare: *Archiv f. Gyn.*, Bd. xlix., S. 94. *Whitridge Williams* (24)—*Johns Hopkins' Hospital Reports*, Vol. iv., No. 8, and *Amer. Jour. Obstet.*, 1895, p. 319.

The numbers after the names refer to the cases given in the table on pp. 538-541.

In 1889 Sänger drew attention to a malignant affection of the uterus developing in the puerperium, to which he gave the name of *deciduoma malignum* from its resemblance to decidua in structure. Subsequently he has called it *sarcoma deciduo-cellulare* because clinically it resembles sarcoma, while its pathological structure reproduces only the cells of the decidua, not the other elements. Though the clinical features of this affection are so well marked as to give rise to little discussion, there has been much controversy as to its pathological nature, etiology and nomenclature. As yet we have not sufficient data to come to a conclusion with regard to these. In discussing this subject,

therefore, we must depart from our usual method of study, and consider first the clinical features, and then the pathology and etiology, which will include origin and nomenclature.

No.	Reporter.	Age.	Number of Pregnancies.	Nature and Termination of last Pregnancy.	Bleeding began.
1 ¹	Chiari 1	24	One	Normal ; full time	0
2	" 2	23	Three	" " "	Four weeks
3	" 3	42	(?)	At sixth month	Six days
4	Pfeifer	36	Five	Mole at second month	Nine months
5	Meyer	55	Four	Mole at fifth month	Immediately
6	Sänger	23	One	Abortion at second month	0
7	Schmorl 1	—	—	—	—
8	" 2	25	Three	Normal ; full time	—
9	" 3	—	—	—	Three months
10	Pestalozza 1	25	Three	Abortion "at" "fourth month	—
11	" 2	45	Ten	Mole at fourth month	—
12	" 3	33	Four	Normal	—
13	Müller	30	Seven	Mole at fifth month	Shortly
14	Gottschalk	42	Four	Abortion at two-and-a-half months	0
15	Guttenplan	28	Eight	Mole	—
16	Kaltenbach	—	—	Large mole	—
17	Löhlein	47	Seven	Mole at seventh month	Twenty-one months
18	Nové-Josserand and Lacroix	24	Three	Mole at fourth month	Three to four weeks
19	Lebensbaum	27	Six	Normal	Five weeks
20	Menge 1	36	Ten	Mole at sixth month	Five months
21	" 2	18	—	Mole at second month	Six weeks
22	Schauta	29	Five	Mole at sixth month	One month
23	Bacon	48	Nine	Mole at ninth month	Five weeks
24	Whitridge Williams	35	—	Normal ; full time	(Nodule in right labium in a fortnight)
25	Marchand 1	17	—	Supposed tubal gestation	0
26	" 2	31	Nine	Normal ; full time	Third week
27	Superno	32	Six	Mole at third month	Shortly
28	Spencer	27	Two	Normal ; full time	Twenty-eight days
29	J. R. Morison	35	Nine	" " "	Six weeks
30	Tannen	23	Three	Mole at third month	Eight months or possibly after a later six weeks' abortion
31	Apfelstedt and Aschoff	33	Three	Ovum expelled at fourth month	Two months

¹ The reference will be found in the "literature" under the corresponding number.

CLINICAL FEATURES.

The table given below is based on all the reported cases.¹ In preparing it, we have selected the facts of chief clinical interest.

<i>Diagnosis—Operation.</i> ²	<i>Metastasis.</i>	<i>Duration.</i> ³
"Endometritis; metritis; later pleurisy and lobular pneumonia"	Lungs and alae vesper-tilionis	Death in six months.
"Bleeding from retained placenta"	Lungs.	Death in seven months (eight days after clearing out of uterus).
—	Lungs; right ovary; vagina; pelvic glands	Death in six months.
"Malignant disease of uterus; tubercular of lungs, with endometritis"	Lungs; anterior vaginal wall	Death in fourteen months.
"Destructive growth of retained products"	Only abdomen examined	Death in nine months.
"Chronic pyaemia"	Lungs; diaphragm; rib; iliac fossa	Death in seven months.
Death from typhus; and tumour discovered accidentally	—	Death from typhus.
"Malignant placental polypus"	Lungs; and broad ligament	Death in two - and - a - half months.
—	—	Death in six months.
"Sarcoma-uteri haemorrhagica seu infectiosa"	Lungs	Death in six months.
Pulmonary symptoms before death	Lungs	Death after five months.
"Sarcoma uteri haemorrhagica seu infectiosa"	Lungs and vagina	Death.
"Malignant decidual tumour" from curetting	Vagina, left gluteal region	Death in four months.
"Malignant sarcoma;" hysterectomy six months after abortion	Lungs, spleen, right kidney	Death in thirteen months (seven after operation).
—	Lungs, vagina	Death in three months.
"Sarcoma"	Metastasis	Death in eighteen months.
"Sarcoma uteri" from curetting: hysterectomy after twenty-seven months	Lungs	Death in three-and-a-quarter years (one year after operation).
"Deciduoma malignum:" hysterectomy three months after abortion	Uterine wall	Living a year later.
Adeno-carcinoma: hysterectomy seven months after	Vaginal (excised)	Death in six days.
"Deciduo-sarcoma:" hysterectomy after seven months	Lungs, pericardium, spleen, stomach, jejunum, vagina, vestibule, pelvic connective tissue, femur	Death in thirteen months (six after operation).
"Deciduo-sarcoma"	—	Living one year later.
Decidual sarcoma from curetting: hysterectomy after six weeks	Vaginal (excised)	Well after one - and - a - half years.
—	Right broad ligament and lungs	Death in six months.
"Sloughing haematoma of labium with sepsis"	Lungs, spleen, liver, kidneys, right ovary.	Death in three months.
Bleeding tumour of vaginal wall, and rapidly growing abdominal tumour	Vagina	Death three weeks after vaginal tumour noted.
Uterus cleared out repeatedly: hysterectomy after five months	—	Death in eleven months (six after operation).
Hysterectomy	None	Well one year after
Puerperal sepsis	Cervix, lungs	Death ten-and-a-half weeks after.
"Squamous celled epithelioma" made from scrapings: hysterectomy after fifteen weeks	Lungs (?)	Death in seven months.
At first as "abortion," then "deciduoma malignum," from curettings: hysterectomy after four months	—	No recurrence seven months later.
"Fibrous polypus" removed after two months bleeding; "sarcoma-deciduo-cellulare" from curetting two months later: hysterectomy, eight months	Lungs, liver, pancreas mesentery, intestine, right femur	Death with lung symptoms in nine months (one after operation).

¹ We have not included cases by Paviot, Jeannel, and others, which appear doubtful.
Hysterectomy, and date post partum.

³ Post partum.

TABLE OF CLINICAL FEATURES OF DECIDUOMA MALIGNUM—*contd.*

No.	Reporter.	Age.	Number of Pregnancies.	Nature and Termination of last Pregnancy.	Bleeding began.
32	Same	42	Three	Mole at sixth month (?)	(Labial swelling on eighth day)
33	Runge	44	Multipara	Three years previously	Irregularly for three years
34	Klien	27	Four	Mole at third month	Shortly
35	Kuppenheim	33	Five	Spontaneously at seventh month	Three weeks
36	Fraenkel	25	One	Mole at third month	(Inflammatory symptoms after seven months)
37	Pestalozza 1	32	Seven	—	Shortly
38	„ 2	44	Thirteen	After mole	Shortly
39	„ 3	22	Two	After mole at second month	—
40	Lönnberg and Mannheim	38	Eight	Normal, but previous one mole	Four weeks after
41	„	42	Three	Mole at fourth month	Seven weeks
42	Götze	21	—	Abortion	—
43	Freund	40	Three	Normal	Shortly
44	v. Franqué	32	Six	Normal	Four weeks
45	Neumann 1	35	Three	Normal	Six weeks
46	„ 2	29	Six	Mole at seventh month	—
47	Chrobak	33	Three (?)	Early abortion (?)	Shortly
48	Resinelli	—	Four	Abortion after influenza	(Tumours in vestibule in three months)
49	Lindfors	44	Three	Mole at third month	Shortly
50	Schmorl	—	—	Abortion	Shortly
51	Koettnitz	25	Three	Normal	Shortly
52	Hartmann et Tonpet	25	Two (?)	Abortion (?)	Shortly
53	Cazin	30	Three	Mole at four months	Shortly
54	Schauta	—	Four	Mole	(Vaginal tumour noted before mole expelled)
55	J. Cock	30	Four	Normal	In a fortnight

TABLE OF CLINICAL FEATURES OF DECIDUOMA MALIGNUM—*contd.*

<i>Diagnosis—Operation.</i>	<i>Metastasis.</i>	<i>Duration.</i>
Left labium painful in pregnancy; remained swollen and hard in puerperium; swelling extending along side vagina excised after three months and mole evacuated. Similar tissue found in uterus	Lungs, spleen	Death four months after abortion (one month after operation).
"Myoma;" for which supra-vaginal amputation done, when chorion-carcinoma found	—	Well three months after operation.
"Deciduoma malignum" from curetting	Lungs	Nine months.
"Sarcoma-deciduo-cellulare" from curetting: hysterectomy, two months	—	No recurrence ten months later.
Laparotomy; chorionic masses removed but uterus left, as tumours also in parametrium	Bladder, spleen (other organs not examined)	Death after eleven months.
Hysterectomy within a week after diagnosis from curetting	Vagina, lungs	Death in six months. Well after one year.
Diagnosis from curetting and hysterectomy three months later	—	Well after one year.
"Malignant serotinal tumour" from symptoms and numerous vaginal tumours; hysterectomy twenty-three months; vaginal tumours excised twelve days later	Lungs, liver, spleen, kidneys, abdominal lymphatics	Death six months after.
"Sarcoma-deciduo-cellulare"	—	After six months still well.
"Non-malignant placental polyp;" seven weeks later ulcerating tumour of posterior vaginal wall removed — "sarcoma-deciduo-cellulare;" one-and-a-half months later uterus and two vaginal tumours extirpated	Vagina	Tumour irremovable by laparotomy; and death.
Diagnosis from curetting; hysterectomy, three-and-a-half months	Vagina	Fifteen months after operation well.
"Deciduoma malignum" from curetting; and hysterectomy, three-and-a-half months	None	Well six months after.
Tumours in uterus and vagina; extirpation	Lungs	Death three months after operation.
"Deciduoma malignum" after uterus extirpated, for foul discharge and fever	Vagina	Well eighteen months after operation.
"Sarcoma uteri" with metastasis	—	Well after operation.
"Deciduoma malignum" from clinical history; septic symptoms; hysterectomy, three months	Vagina, lungs	Death shortly.
—	—	Death five days after extirpation.
Placental polypus removed after four weeks; three weeks later two varicose vaginal tumours found and removed	Lungs, kidney, and other organs	Death in seventeen weeks.
"Haemorrhagic endometritis after abortion," then "deciduoma" suspected	Vagina, pelvic cellular tissue and lungs	Death in two-and-a-half months.
"Sarcoma" Hysterectomy after nine months	Only uterus examined	Death in four months from sepsis.
—	Ovaries (removed at operation)	Well two-and-a-quarter years after operation.
"Deciduoma malignum" from vaginal tumour and uterine contents	—	—
"Deciduoma malignum"	Lungs, ovary	Death in two months.

Clinical
Features
of Decidu-
oma Malignum.

Summarising the facts brought out in the foregoing columns we note that the *age* of the patients ranges from seventeen to fifty-five, the *mean being thirty-three*. This disease is not therefore like cancer, an affection of advanced life; and for reasons to be mentioned immediately it necessarily arises within the child-bearing period.

Commence-
ment.

It will be noted that the affection has always developed *shortly after pregnancy*—in this respect it differs from sarcoma or carcinoma, as we ordinarily understand these; but it is an open question whether we may not be dealing with these affections modified by recent pregnancy, rather than with an entirely new pathological condition.

While it may arise after a normal full-time pregnancy, there seems to be a special tendency to it after a *hydatid mole*—twenty-four out of the fifty-five cases giving a history of this. In several cases it has followed on abortion, so that in its commencement it has often been mistaken for the retained products of an abortion.

Symptoms.

The *first symptom* has in almost every case been *hæmorrhage*, which began as early as six days, and as late as twenty-one months after delivery, the usual period being from four to six weeks. Exceptionally, the development of a tumour in the labium, vestibule, or vagina was the first thing to attract notice. Though it is not referred to in the table, the hæmorrhage was often followed by foul smelling discharge and fever, which increased the resemblance of the case to one of “retained products of conception.”

Diagnosis.

While in the earlier cases, mistaken diagnosis was frequent, in the later ones the correct *diagnosis* was usually made—most frequently from *microscopic examination* of the tissue removed by the curette. It is impossible to give in a table the description of the physical signs found in each case, but generally the uterus was found larger on bimanual examination than it should be, and sometimes rapidly increasing in size. Its contents, while friable and vascular were of the nature of a soft tumour growing in the walls, sometimes distinctly polypoidal, and not the loose debris and blood-clot found in cases of retained products of abortion.

Metas-
tasis.

The *metastases* are the most striking feature. In the large proportion of cases they are general—usually in the lungs, pointing to diffusion through the blood-vessels. Only in a few cases are they local, the extension taking place through the lymphatics. It is noteworthy that none of the cases that recovered showed pulmonary symptoms.

Duration.

The *duration* of the disease is *short*, death occurring about six months after the delivery; the most rapid course was two months, and the most protracted three and a quarter years, but the mean duration of life was under six months. When we consider that there is no evidence of the pathological condition having existed during the pregnancy, this is a very short time for a malignant condition to develop and to run

its course. This disease, whatever its nature, is apparently more malignant than cancer or sarcoma, as we ordinarily understand these.

The only successful *treatment* has been *hysterectomy before metastases* Treatment. have developed. Curetting may be necessary for diagnosis, but cannot be considered as a form of treatment, the effect being at most only temporarily beneficial, and sometimes prejudicial by exciting new infection. Of twenty-three cases in which hysterectomy was done, nine died at varying periods from five days to thirteen months after the operation. Of the fourteen other cases, three were reported as well shortly after the operation, one at three months, two at six, and the others at seven, ten, twelve, fifteen, eighteen and twenty-seven months respectively.

PATHOLOGY.

The naked-eye changes in the uterus are well defined, and will be evident from fig. 285.¹ Spencer describes the uterus as enlarged ($4\frac{1}{2}$




FIG. 285.

UTERUS FROM A CASE OF DECIDUOMA MALIGNUM (Spencer)

inches long and $2\frac{1}{2}$ inches broad); and when laid open, showing a gangrenous cavity nearly perforating the fundus, due to the breaking down of the tumour. There were secondary growths in the cervix and lungs. A microscopic section (fig. 286) shows that the tumour consists of "isolated multinucleated masses of large cells (the so-called 'syncytium'), and of a loose tissue between the syncytial masses containing individual cells similar to those of the syncytium; in this tissue hæmorrhages, and in places, aggregations of leucocytes, are seen.

Microscopic Structure of Deciduoma Malignum.

¹ Reproduced here from the London Obstetrical Transactions, by the kind permission of Dr Spencer and the Council of the London Obstetrical Society. This preparation has the special interest of being from the first recorded case of deciduoma malignum in England.

The syncytium is often extensively vacuolated. The growth is penetrating into the muscular wall.

"Under a high power (one-sixth) the syncytium, which in places has the appearance of a huge multinucleated cell, is evidently made up of a number of large cells aggregated or fused together. . . . The loose tissue between the syncytial masses is a delicate reticulum of connective tissue, in the meshes of which are large cells similar to those making up the syncytium."

The new formation sometimes appears as soft villous growths simulating chorionic tissue, or as nodules like beans, or as an ulcerating surface. Bacon and Cazin describe three layers in the uterine wall: blood-clot and debris on the free surface, underneath this necrosed tissue, and then the neoplasm extending as plugs into the uterine tissue. In the "zone of penetration" are seen large cells with deeply-staining nuclei, alone, in rows, and in masses. Sometimes there is a




FIG. 286.

SECTION OF TUMOUR FROM UTERUS IN FIG. 285 (*Spencer*).

row of nuclei in a band of protoplasm, recalling the syncytium of the placenta. The tumour cells break up the muscular bundles and cause their degeneration, penetrate the walls of the vessels, and proliferate in their channel, thus favouring metastases having a similar structure—the cells proliferating so as to form villous masses suggesting metastasis of chorionic villi. Two kinds of cell have been described, polyhedral cells with large nuclei resembling decidual cells, and multinucleated masses of protoplasm resembling giant cells or the syncytium of the placenta. The proportion of these varies both in the original tumour and the metastases. Thus, while Sanger found the decidual-like cells predominating, so that he describes the disease as a sarcoma with cells of decidual type, Gottschalk found both elements in the tumour but epi-

thelial cells alone in the metastasis, while Freund described his case as consisting of syncytium alone.

ORIGIN AND NOMENCLATURE.

A glance at fig. 287 will recall to the student the histological structures found at the placenta site. In the diagram we have given also the various names proposed for this tumour according to the view taken as to its genesis. They are self-evident, but suggest the following criticism.

The clinical resemblance of this tumour to sarcoma is a strong argument for Sanger's position that it is developed from the maternal connective tissue, that is to say, the decidual cells. Against this we have Ruge's view that the decidual cell cannot become malignant. He regards the decidual cell and the sarcoma cell as descending along different lines, the physiological and the pathological, from the connective-tissue cell, and holds that the one cannot pass into the other. If it be a sar-




FIG. 287.

HISTOLOGICAL STRUCTURES found at placental sites which have suggested the NOMENCLATURE of Deciduoma Malignum.

coma, we must go a step farther back and refer its origin to connective-tissue cells which have not become decidual.

The source of the syncytium, or nucleated layer of protoplasm surrounding the villi is disputed, some maintaining that it is maternal as in the dog, cat, and mouse; others that it is foetal. Whether it be foetal or maternal it is *epithelial*, and if this tumour arise from this source it is a *carcinoma*. Hence the term "syncytial carcinoma."

Beneath the syncytium is another layer of epithelial cells which is either a specialised layer of the latter, or the proper chorionic epiblast, according to the view we take of the syncytium. Some have maintained that the round-celled element of this tumour arises from this layer (which is called Langan's layer to distinguish it from the syncytial layer), while the giant-cells arise from the syncytium. Marchand takes this view, and believing that the syncytium is maternal, comes to the conclusion that the tumour must have a compound origin, foetal and maternal. It is difficult to see how this can be, unless we regard the one element as active, and the other as parasitic on it. He

recognises, however, the difficulty of the question by suggesting the provisional name of "serotinal tumour," which implies no theory as to its nature except that it develops in the decidua serotina.

The core of the villus consists of foetal connective tissue. Gottschalk alone refers the origin of part of the tumour to this, making it a foetal sarcoma. He believes that the epithelium of the villus is also involved, thus making the affection a combined sarcoma and carcinoma of foetal tissues only.

It is thus still an open question whether this tumour is a sarcoma or a carcinoma. The weight of argument is in favour of the former;¹ and we have therefore considered it under sarcoma uteri. Whether it merits a special name is also under discussion. Its clinical features are sufficiently well marked to distinguish it from sarcoma uteri as ordinarily understood, and the term deciduoma malignum, though vague, is convenient until we have more definite knowledge as to its structure and origin.

To sum up, we may say that under the name "deciduoma malignum" attention has been directed to a malignant condition developing after child-birth or abortion, with symptoms resembling those of retained placenta followed by sepsis. Its course is rapid, lasting only a few months, pulmonary metastases being frequent. Its pathology and nomenclature are still *sub lite*, and it is possible that different pathological conditions have been described under this head: viz., a malignant affection of the epithelium of the chorionic villi—carcinoma of the chorion, and a larger group of malignant tumours of connective-tissue origin. These may arise from transformed decidual cells, as Sanger holds; or, if we accept Ruge's position that when once connective-tissue cells have become transformed into decidual ones they cannot undergo further transformation into malignant ones, they must be derived from the connective tissue of the mucosa directly.

¹ See an interesting discussion on this subject before the London Obstetrical Society, where it was held that there was nothing in the histological characters of the three cases reported to justify the supposition that the tumour was of decidual origin. It was considered to be ordinary sarcoma uteri: *Lond. Obstet. Trans.*, 1896, p. 171.

SECTION VI.

AFFECTIONS OF THE VAGINA.

THESE we shall consider in the following order:—

CHAPTER XLV. Atresia Vaginæ.

„ XLVI. Vaginitis: Vaginismus: Tumours.

CHAPTER XLV.

ATRESIA VAGINÆ.

LITERATURE.

Breisky—Die Krankheiten der Vagina: Stuttgart, 1879. *Delaunay*—Étude sur le cloisonnement transversal du Vagin, etc.: Paris, 1877. *Dohrn*—Angeborene Atresia vaginalis: Archiv für Gynäk., Bd. x., S. 3. *Duncan, J. Matthews*—Case of so-called Imperforate Hymen: Lond. Obst. Tr., Vol. xxiv. *Emmet*—Principles and Practice of Gynecology, p. 188: Philadelphia, 1884. Congenital Absence and Accidental Atresia of the Vagina, etc.: Trans. Am. Gyn. Soc., ii., p. 437. *Fasola*—Contributo allo studio dell'origine dell'imene, a proposito di un caso di ematocolpo per mancanza delle parte inferiore delle vagina e dell'imene: Annal di Ostet., 1885, p. 146. *Fuld*—Salpingotomie wegen Hämatosalpinx bei Gynatresie: Archiv f. Gyn., Bd. xxxiv., S. 191. *Hart*—Transverse septal Atresia of the lower third of the Vagina: Scot. Med. and Surg. Journ., April 1897. *Leopold*—Blutansammlung im verschlossenen Uterovaginal-kanale und die Salpingotomie: Archiv f. Gyn., Bd. xxxiv., S. 371. *Puech, A.*—Des Atrésies complexes des voies génitales de la Femme: Ann. de Gynécologie: Paris, 1875. *Sänger*—Drei Fälle von Salpingo-oöphorectomia duplex bei Hæmatometra gynatresia: Cent. f. Gyn., 1896, S. 49. *Simpson, Sir J. Y.*—Diseases of Women, p. 256: Edin., 1872. *Simpson, A. R.*—Contributions to Obstetrics and Gynecology, p. 195: Edin., 1880. *Veit*—Erkrankungen der Vagina: Handbuch der Gynäkologie: Wiesbaden, 1897. *Webster, J. C.*—Some observations regarding the Diagnosis and Treatment of Atresia Vaginæ: Amer. Jour. Obstet., 1895, Vol. ii., p. 544. The literature is given fully in Veit's recent monograph.

Definition. ATRESIA (ἀ-τρήσις, non-perforation) has been already defined as occlusion of the genital tract where the obstruction is complete and leads to accumulation of menstrual blood or mucous secretion. This occurs at three places—the *hymen*, the *vagina*, and the *cervix uteri*. Atresia of the *cervix* has been already described (v. Chap. XXVI.). Accumulation of blood in one-half of a *septate uterus* or *vagina* will be considered by itself at the end of this Chapter.

PATHOLOGY.

1. ATRESIA HYMENALIS.—The structure of the normal hymen has been already described (page 7). In atresia hymenalis it forms a continuous membrane, is thicker and of an almost cartilaginous toughness; this explains the rarity of spontaneous cure by rupture of the membrane. This condition is produced by the occurrence of inflammatory

adhesion of the folds after their formation, that is after the nineteenth week of foetal life. When the vagina is distended with menstrual blood, the hymen bulges forwards.¹ As the menstrual blood accumulates, the vagina distends so as to form a tense membranous-walled sac nearly filling the pelvis, and with a smaller firmer body (the undilated uterus) rising from its upper surface (*v. fig. 290*). If the tension be not relieved, the cervix next becomes dilated and may rupture. Finally the uterus itself becomes opened out, though this does not occur till late.

During this period, accumulations of blood may take place in the Fallopian tubes in the form of diverticula, usually situated towards the




FIG. 288.

ATRESIA VAGINÆ, SEEN FROM BEHIND. Thickness of obstruction (through which a probe is passed) 3-4 mm.; of vaginal wall below atresia 2-3 mm., above it (at *x*) 6 mm. Dilatation of the body of the uterus is small compared with the common cavity formed by cervix and upper portion of vagina. Left Fallopian tube markedly dilated, with no distinct flexion on it, and changed at its free end into a thin-walled blood sac which had burst. Right tube undilated. (*Breisky.*)

fimbriated end (figs. 288 and 289). These may be produced by a reflux of the blood from the distended uterus into the tubes, or by hæmorrhage from the mucous membrane of the tubes themselves. Schroeder held that the latter was the only explanation, but recent enquiry as to menstruation show that bleeding from the mucous membrane of the tubes is unlikely. The uterine end of the Fallopian

¹ Blood extravasation occurred into the labia in Davy's case. *Lancet*, 1886, ii., p. 1171.

tube is in some cases closed. Blood may escape gradually from the fimbriated end of the tube, and set up a localised peritonitis matting down the tube and uterus; a hæmatocele is sometimes thus produced. Veit accounts for the occlusion of the fimbriated end by a previously existing catarrh which also causes the atresia of the genital tract below. He refers to the frequency of inflammation in early childhood and thinks that the atresia arises then and not during foetal life.

2. ATRESIA VAGINALIS. The *thickness of the obstruction* varies in different cases, according to the extent of the original obliteration and




FIG. 289.

CASE OF DOUBLE ATRESIA. The *lower* affects the hymen and was *acquired*; above this was a cavity one inch long which contained purulent debris: the *upper* obstruction was one inch thick and was *congenital*; above it, is the dilated uterus and cervix. The Fallopian tubes contain blood-sacs with small rents in their walls (*Breisky*, case reported by *Steiner*).

the thinning produced by the pressure from above. The *dilatation of the vagina* above the obstruction is remarkable; it may form a tumour filling the pelvis, pressing on the bladder and rectum, and raising the uterus above the brim; the walls become *hypertrophied* as is well seen in the preparation represented in fig. 288, taken from a patient who died on the same day as the operation for atresia was performed.

The *seat of the obstruction* is most frequently in the *lower third* of the Seat of Obstruction. vagina. This condition may be mistaken for imperforate hymen; as the wall of the sac, bulging through the hymeneal orifice, becomes adherent to the hymen which appears as a mere fringe on the bulging membrane. There is not, however, the same distention of the vulvar orifice and perineum as in atresia hymenalis. *Atresia of the whole vagina* is usually associated with imperfect development of the uterus (*Breisky*).

Atresia may exist at more than one point in the vagina. The specimen represented in fig. 289 illustrates this. It has this further interest that the lower atresia—at the vaginal orifice—was *acquired*, the result of a fall on a block of wood when the patient was two years old; the upper atresia was *congenital*. The accumulation of menstrual blood in the upper sac called for operative interference when the patient was seventeen years of age. The lower sac contained purulent matter. On the fifteenth day after the operation, death occurred from septic peritonitis.

The character of the *retained menstrual blood* is peculiar. It is of a Character of retained Blood. brownish chocolate-red colour, of a thick treacle-like consistence, and contains no coagula. Microscopically, it shows shrivelled red blood-corpuscles, flat epithelial cells, mucous corpuscles, extravasated blood-pigment, and granular debris. The mucus prevents coagulation; part of the fluid portion is probably reabsorbed, since the quantity removed is less than the sum of what we should expect from the successive periods passed.¹

ETIOLOGY.

1. Atresia may be *congenital*, due to non-development of a part of the Congenital Atresia. canal or its subsequent closure during foetal life.

Atresia hymenalis implies that the hymeneal folds were developed (at the nineteenth week) but afterwards became blended into a continuous membrane.

Atresia of the vagina behind the hymen is, according to Dohrn, due to the fact that (at the eighteenth week of foetal life) the walls of the genital canal become closely approximated behind the site of the hymen, so that closure of the vagina is especially favoured in that part; or, according to Hart, to persistence of a septum between the Wolffian bulbs and the involution from below.

Atresia of the middle or upper third implies the development of the ducts and their coalescence into a vagina, with a subsequent occlusion due perhaps to inflammation (*Breisky*).

¹ Oliver gives Bedson's chemical analysis of the retained blood: "It gave the spectrum of reduced hæmatin, and contained '6 p.c. of urea; 100 cc. contained total solids 7.65 grms., organic compounds 6.93 grms., mineral compounds .72 grms. In the solids were found salts, for example, chlorides, sulphates and phosphates, and such bases as iron, calcium, magnesium and sodium: *Brit. Med. Journ.*, 1888, ii., p. 1160.

Complete absence of the vagina or its representation by a fibrous cord is due to the non-development of the ducts of Müller; *absence of the lower third* is occasioned by the non-extension of the ducts downwards so as to open into the cloaca.

Acquired
Atresia.

2. Atresia is also *acquired*; that is, it arises during life. The most important causes which produce this condition are the following:—

Sloughing and subsequent cicatrisation after labour¹;

Sloughing from impaired vitality in typhus, scarlet-fever, small-pox, and cholera;

Cicatrisation after injuries received in childhood;

Superficial inflammation of the mucous membrane, leading to adhesion of opposed surfaces.²

The commonest form of congenital atresia is due to imperforate hymen; of acquired, is due to cicatrisation of the upper part of the vagina and cervix after labour.

SYMPTOMS.

As congenital atresia is productive of bad results only in so far as it impedes the menstrual flow, symptoms do not arise till puberty. Should menstruation not take place at puberty, the condition may not attract attention till the patient enters married life.³ Cases are, however, on record in which the accumulation of mucus has called for operative interference even in childhood.

Symptoms
arise at
Puberty.

At puberty the patient experiences menstrual molimina without the appearance of a discharge. As the vaginal sac distends, pain is felt in the pelvis at first only at the periods and then more continuously. With this there is also constitutional disturbance. The periods of suffering become more protracted, the intervals of relief shorter. When the dilated vagina presses on the bladder and rectum, it causes difficulty in micturition and defæcation. The abdomen swells and this, with the amenorrhœa, causes suspicion of pregnancy which is sometimes the occasion for seeking advice. If the case is left to itself it terminates fatally through rupture of the uterus or cervix (usually the latter) or of a blood sac in the Fallopian tube, or through a simple or septic peritonitis independently of rupture. In some cases, the obstructing membrane has given way by rupturing (in acquired atresia) or sloughing (in the congenital form). But even this is not a favourable termination, as

¹ As in the cases recorded by Holdsworth (*Lancet*, 1883, i., p. 949) and Cross (*Amer. Journ. Obstet.*, 1883, p. 809, and 1886, p. 802).

² As in the case recorded by More Madden (*Dublin Med. Journ.*, lxxv., p. 158), in which it developed in a multipara after a miscarriage. Heywood Smith found a complete vaginal septum formed by adhesions developing during pregnancy (*Lond. Obstet Trans.*, xxiii., p. 117).

³ Zinnstag records a curious case in which an apparently imperforate hymen was not observed until labour set in; there must have been a perforation (to account for conception) at one time, but it had closed subsequently: *Centralb. f. Gyn.*, xii., S. 219. Doléris reports a similar case: *Archiv de Toc.*, 1886, p. 135.

the risks consequent on operative interference are still more likely to ensue when the hymen ruptures of itself.

DIAGNOSIS.

The importance of physical diagnosis will be evident from the following case. "A. B., æt. 16, unmarried, has for twelve months suffered from pain in the pelvis and back, with occasional acute exacerbations accompanied by nausea and vomiting. She has been treated for inflammation; and mercurial ointment had been applied to a swelling which had appeared in the left groin, on the supposition that it was an enlarged gland." Examination per rectum showed a condition similar to that seen at fig. 291; the swelling in the left groin was the elevated uterus.

The practitioner will often ask himself whether a vaginal examination




FIG. 290.

ATRESIA HYMENALIS (Schroeder).




FIG. 291.

ATRESIA VAGINÆ—lower third (Schroeder).

is necessary. On the patient's returning several times and there being nothing in the constitutional state (phthisis, chlorosis) to explain the amenorrhœa, tell the friends that there is no apparent cause for the non-appearance of menstruation except on the supposition of a mechanical obstruction to its outflow. If there be pain in the pelvis and marked constitutional disturbance, the reasons for demanding an immediate examination will be evident. The conditions found in the various forms of atresia will be easily understood by studying figs. 290 to 293. The external genitals are first examined; a wide urethral orifice may be mistaken at first glance for the vagina, as in *atresia hymenalis* the urethral orifice is more patulous than it is normally (Oldham); the hymen is seen bulging forwards at the ostium vaginæ. The finger is passed into the rectum and feels that the anterior wall is

made to bulge by a tense elastic sac. On bimanual (recto-abdominal) examination, this sac is felt to be equally distended and to fill the pelvis; it may extend into the abdomen as far as the umbilicus. The feeling of the sac is quite characteristic and is like that of a tense india-rubber ball; on its upper surface, the uterus is felt as a small firmer tumour.

In *atresia vaginæ* the condition is the same, except that the hymen does not bulge and that the sac does not extend so low down.

Diagnosis
from Preg-
nancy.

Atresia of the cervix (figs. 292, 293) might be mistaken for early pregnancy; as the amenorrhœa and the distended uterus are present in both cases. But the condition of the cervix, the form of the uterus, and specially the characteristic tense feeling of the tumour, enable us

From Sar-
coma.

to distinguish it from a pregnant uterus. Malignant tumours (sarco-




FIG. 292.

ATRESIA OF CERVIX AT OS EXTERNUM
(Schroeder).




FIG. 293.

ATRESIA OF THE CERVIX AT OS INTERNUM
(Schroeder).

mata) have a similar elastic consistence, but with them we should not have amenorrhœa.

It is not in all cases easy to say whether the atresia be congenital or acquired. The existence of other malformations would favour the former view, of cicatrices beside the obstruction the latter. There will also be a greater thickness of tissue felt between the urethra and rectum in the acquired form, corresponding to the obliterated vaginal canal.

Estimation
of Extent
of Atresia.

In *atresia vaginæ* it is important to estimate the distance to which atresia extends, so that we may know how much tissue we must cut through to reach the sac or the cervix uteri. This is best done by

passing the index finger into the rectum till the tip is on the place where the bulging of the sac begins or where the projection of the cervix is felt; the thumb is at the same time passed into the ostium vaginæ till it reaches the obstructing membrane; the thickness of the latter can thus be estimated.

PROGNOSIS.

If menstrual blood be accumulating, the prognosis is always grave. In atresia of the hymen the prospect of cure by operative treatment is more hopeful than in congenital atresia of the vagina. In acquired atresia of the vagina, if the obstruction be removable, the prognosis is favourable. The unfavourable cases are those in which the vagina is partially or not at all developed; the prognosis as to curability by operation depends on the thickness of the tissue between the urethra and the rectum, which determines the possibility of opening up a vagina.

When menstrual blood has accumulated, while explaining to the patient's friends the necessity of immediate operative treatment, we should inform them also of the dangers attendant on the operation—the immediate danger of rupture of a blood sac in the Fallopian tube, the more remote one of simple or septic peritonitis.

The seriousness of the complication of hæmatosalpinx is seen in Fuld's statistics:¹ of sixty-five which he has collected, more than two-thirds (forty-eight) died; while seventeen were saved by operation.

TREATMENT.

The treatment consists in the formation of a channel to allow the menstrual blood to escape; in the case of imperforate hymen this is easily done by incising the membrane, but in atresia vaginæ we have to construct a new vaginal canal. Two dangers associated with this operation must be kept in view. *First*, too rapid collapse of the sac may lead to rupture of the Fallopian tubes or of vascular adhesions round the uterus. To prevent such accidents, the operator should allow the contents of the sac to escape slowly, and should on no account apply pressure from above to hasten the process. *Second*, the operation is frequently followed by septicæmia. To prevent this, antiseptics should be used. Listerism cannot be carried out here; but by washing out the sac carefully with carbolised water, preventing the entrance of air, and allowing free drainage when fluid collects, we greatly diminish this risk.

Dangers of
Operation.

The danger of rupture of a hæmatosalpinx has only recently been

¹ *Op. cit.* These cases were collected from all sources, and before the operation for hæmatosalpinx was a recognised one.

recognised, and raises the question whether abdominal section should not first be performed in all cases where a dilated tube is recognised.¹ Fuld has collected sixty-six cases of atresia, in which the tubes were dilated and ruptured, forty-eight of whom died. Of these deaths, nine were apart from operation, but thirty-nine after the operation described above of emptying the sac from below.

Another danger, which follows some time after the operation, is the contraction of the new canal which, unless specially guarded against, may lead to its obliteration. Emmet expresses this well when he says "the surface of the canal is essentially a cicatricial one, and will consequently contract to a greater or less extent." To diminish the liability to contraction, he recommends that the tissues be torn with the finger nail or broken up with the scissors rather than divided with the knife; the raw surface is made to heal upon a glass plug.² Crede³ prevented cicatrization by taking a flap from the labium majus and turning it into the vagina so that it could be stitched to the cervix and to the raw surface produced by dividing the old cicatricial tissue in the vagina.

We shall describe shortly the operations for (1) imperforate hymen, (2) atresia of the vagina, (3) atresia of the cervix.

Operation
for Imper-
forate
Hymen.

1. *Imperforate Hymen.* This operation, though apparently simple, should never be performed in the consulting room but always at the patient's house or in hospital. The time chosen should be between two menstrual periods which are indicated by menstrual molimina. The hymen is punctured with a small trocar which has been rendered thoroughly clean and aseptic beforehand. The fluid is allowed to escape slowly. After it has ceased to flow, the opening in the hymen is enlarged with a knife. This incision is made in the form of a cross, or the membrane is pinched up with forceps and an elliptical portion cut out. A. R. Simpson recommends that the opening in the hymen be made with the cautery, which prevents septic absorption by the wound. We can dispense with the trocar if we take care to make at first only a small opening, which can afterwards be enlarged. A stream of warm antiseptic water is now made to flow gently into the cavity; the opening should be large enough to permit the fluid to flow outwards at the same time, so that the sac may be washed out without being subjected to any pressure. A plug of lint soaked in antiseptic oil is placed in the hymeneal orifice, and a larger pad over the vulva. The patient keeps her bed for ten days after the operation. If there be a rise of tempera-

¹ Kehrer has done this once successfully, and Leopold five times—Fuld and Leopold *op. cit.* The diagnosis of hæmatosalpinx may be made, according to Fuld, either by feeling the dilated tube or finding that the amount of fluid evacuated from the vagina does not correspond to the size of the mass felt before on palpation—the latter suggesting rupture into the abdomen. Laparotomy is called for in both cases. Säger has also published three cases—*op. cit.*

² In the *Americ. Journ. Obst.* (1887, p. 1189) he refers to his attending in her second confinement a patient on whom he had operated ten years previously to make an artificial vagina when she was fifteen years old.

³ *Archiv f. Gyn.*, Bd. xxii., S. 229.

ture or other indication of septic inflammation, the vagina should be again washed out.

2. *Atresia of the Vagina.* The patient is placed in the lithotomy position, and the labia are retracted by the fingers of the assistants who hold the thighs. The sound is passed into the previously emptied bladder; it is then held by an assistant in such a way that the urethra and bladder are drawn well upwards towards the pubes. The index finger (with, if necessary, the second) of the left hand is introduced into the rectum; and the thickness of tissue between the finger and the sound, as well as the position of the distended sac above, carefully ascertained: the finger is kept in the rectum during the operation, both to hook that structure backwards so as to prevent its being cut into and to guide in tearing up the septum. Should the operator wish to have both his hands free to use instruments, an assistant can pass the finger into the rectum. The operator now makes with a knife a transverse incision over the hymen, or through the skin between the anus and the urethra. When the sac is reached, it is punctured and washed out with the same precautions as

Operation
for Atresia
Vaginæ.




FIG. 294.

PERFORATED GLASS PLUG TO BE USED AFTER OPERATION FOR ATRESIA VAGINÆ. The left hand figure shows the external end of the tube with the tapes attached.

in the operation for imperforate hymen; it is then carefully and gently packed with strips of lint soaked in antiseptic oil. These are taken out on the following day, but a tightly fitting plug is left in the newly formed portion of the vagina to prevent its contraction; after three or four days, a perforated glass plug (fig. 294) is passed in to keep the new canal dilated. The plugs are made of various thicknesses, and have a rim at the external end to prevent their being pushed in too far. The plug must not be so long as to press on the roof of the vagina, and should be of such a thickness that, while it can be easily slipped out and in by the wearer, it stretches the new canal; it is kept in position by tapes which are fastened to the rim and, before and behind, to an abdominal band. A pessary can be employed subsequently; some instrument may have to be worn constantly for a year or more and where there is continued tendency to contraction, for a short period daily during many years.

In a case operated on by Page, there was an accumulation of fluid in

the vagina, and a second in the uterus itself which did not discharge till the cervix was incised.¹

This operation has been performed even when there has been no accumulation of menstrual blood. The indications for operating are thus given by Thomas: "It should be resorted to (a) if menstrual blood be imprisoned; (b) if a uterus can be distinctly discovered and the patient be suffering from absence of menstruation; (c) if the necessity for sexual intercourse be imperative." Cases have been recorded in which the formation of a vaginal canal has led to the establishment of menstruation when it was formerly absent, to the development of the uterus and ovaries where these were rudimentary (?), or to an improvement in the general health of the patient although there was no indication of further development in the rudimentary uterus and ovaries.

More difficulty is experienced in operating where there is no accumulation of menstrual blood and the vagina is entirely absent or represented by a fibrous cord. In such a case, there is not the same necessity for surgical interference unless it be to satisfy the claims of married life. If the uterus and ovaries be well developed and the patient be anxious to have her condition remedied, the operation is justifiable. Here we have not the distended sac as a guide to the point on which we are to cut down. The cervix, of which the position should be ascertained by a combined recto-abdominal examination, should be fixed as far as possible by an assistant's making firm pressure from above upon the uterus; there is no danger in such pressure if there be no accumulation of menstrual blood. The mode of procedure is the same as that just described.

Operation
for Atresia
Cervicis.

3. *Atresia of the cervix.* Usually the obstruction is so slight that the forcible passage of the sound overcomes it. Should the obstruction resist all efforts to pass the sound we require to use the knife to open the canal. If the uterus be much distended with menstrual blood, it is safer to empty it first with the aspirator-needle passed through one of the fornices; the emptying should be effected slowly and, if the distension be considerable, at more than one sitting; rapid emptying is apt to set up uterine contractions which may produce rupture of a dilated Fallopian tube.

Atresia of one-half of a Septate Uterus and Vagina.

This form of atresia has certain characteristics which distinguish it from the other forms described above.

The chief peculiarity is that it presents the phenomena of free menstruation + those of retained menstruation.

The pathological condition is apparent from fig. 295. Spontaneous rupture of the septum with escape of the retained fluid (in this case

¹ *Lancet*, 1884, i., p. 706.

through the patulous uterus or vagina) occurs more frequently in this than in other forms of atresia; rupture of the Fallopian tube, with its fatal consequences, is also a more frequent occurrence (*Puech*). The spontaneous rupture of the septum does not usually occur at its lowest point; hence there is liability to accumulation of purulent matter in the pouch below the point of perforation, which is a source of septicæmia.

The symptoms are the same as in the other forms of atresia, but they are masked by the presence of a menstrual flow. This visible menstruation is often irregular, and profuse leucorrhœa (from the patulous cavity) is frequently present. Lackie¹ has recorded a curious case in which after operation on an atresic septate uterus a discharge of blood came periodically from the formerly atresic half, but alternating with the




FIG. 295.

SEPTATE UTERUS; the right half is pervious, the left half has been distended with retained menstrual blood (*Schroeder*).

regular menstrual flow which had always taken place from the other half of the uterus.

Physical examination shows a fluctuating tumour lying beside the uterus and alongside of the patulous vaginal canal. Sometimes it winds in a spiral manner round the latter.

The diagnosis is not difficult if the blind sac extend to the ostium vaginæ and be felt running alongside of the vaginal canal or winding round it. If, however, it be limited to the side of the uterus or only extend partially on to the vagina, it may easily be mistaken for other para-uterine tumours—most frequently for hæmatocele. To clear up the diagnosis and also as a step towards treatment, we puncture the sac with the aspiratory-needle.² The character of the discharged blood will indicate the diagnosis.

¹ *Scottish Med. and Surg. Journ.*, 1897, p. 165.

² Kiderlen mentions a case from Martin's Clinique in which about two-and-a-half pints of fluid were evacuated from the dilated right half of the vagina and uterus: *Zeits. f. Geb. u. Gyn.*, B. xv., S. 1.

The treatment consists in slowly but thoroughly evacuating the sac, washing out and establishing a permanent opening from it.

A *septate vagina* is sometimes found with a septate uterus (*v. fig. 155*), both halves being pervious so that there are no symptoms.¹ In rare cases, the one vagina is imperforate. Kleinwächter² records an interesting case of a bulging tumour of the anterior vaginal wall resembling in position a cystocele; it ruptured and pus escaped. On laying open the fistulous tract, its walls had the naked eye and microscopic characters of vaginal mucous membrane in a state of inflammation. Traces of a septate condition may persist as bands.

¹ Cullingworth has recorded two cases of a transverse septum in the lower part of the vagina: *Lancet*, 1889, i., p. 726.

² *Zeits. f. Geb. u. Gyn.*, B. xi., S. 254.

CHAPTER XLVI.

VAGINITIS: VAGINISMUS: TUMOURS.

LITERATURE OF VAGINITIS.

Döderlein—Ueber das Verhalten pathogener Keime zur Scheide: Deuts. med. Woch., 1893, No. 10, p. 157. Also, Das Scheidensecret und seine Bedeutung für das Puerperalfieber: Leipzig, Besold, 1892. *Hennig*—Der Katarrh der weiblichen Geschlechtsorgane. *Hildebrandt*—Monat. f. Geb., Bd. xxxii., S. 128. *Krönig*—Ueber das bakterienfeindliche Verhalten des Scheidensecretes Schwangerer: Deuts. med. Woch., 1894, No. 43, p. 819. *Macdonald, Angus*—Edin. Med. Jour., June, 1873. *Menge*—Ueber ein bakterienfeindliches Verhalten der Scheidensecrete Nichtschwangerer: Deuts. med. Woch., 1894, No. 46-48. *Menge and Krönig*—Bakteriologie des weiblichen Genitalkanals, Vol. i., 1897. *Miller, A. G.*—Four and a Half Years' Experience in the Lock Hospital, Edinburgh: Edin. Med. Journ., 1883. *Næggerath*—Latent Gonorrhoea in the Female Sex: Am. Gyn. Trans., Vol. i., p. 268. *Ruge*—Ueber die Anatomie der Scheidenentzündung: Zeitschrift f. Gyn. u. Geb., Bd. iv., S. 133. *Schroeder*—Die Krankheiten der weiblichen Geschlechtsorgane, S. 460: Leipzig, 1879. *Veit*—Erkrankungen der Vagina: Handbuch der Gynäkologie: Wiesbaden, 1897. *Walther*—Bakteriologische Untersuchungen des weiblichen Genitalsecretes in Graviditate und im Puerperium: Archiv f. Gyn., Bd. xlviii., p. 201. *Winckel*—Colpohyperplasia cystica, etc.: Arch. f. Gyn., Bd. ii., S. 406. *Winter*—Die Mikroorganismen im Genitalkanal der gesunden Frau: Zeits. f. Geb. u. Gyn., Bd. xiv., p. 443.

VAGINITIS.

SYNONYMS.—Colpitis (Gr. κόλπος, *a fold*): Elythrititis (Gr. ἑλυτρον *a sheath*).

NATURE AND RELATION TO MICRO-ORGANISMS.

Vaginitis is an inflammation of the mucous membrane of the vagina. The structure of this mucous membrane has been already described (v. p. 32). From its consisting of connective-tissue papillæ covered with several layers of squamous epithelium, it resembles the structure of the skin rather than that of a mucous membrane, few, if any, mucous glands being present. Consequently, the inflammatory changes are more allied to those of the skin than to those of a mucous membrane. Further, this peculiarity in the anatomical structure is of the first importance with regard to infection. The vagina is even more proof

Relation of
Structure
to Inflamm-
atory
Changes,

and to
Micro-
Organisms.

than the skin against micro-organisms, since its epidermal protection¹ is not weakened by hair follicles, sweat, or sebaceous glands.

The anatomical structure of the vagina and its secretions have been recently specially studied with regard to bacteriology. The most important micro-organisms are shown in Plate IV. (see p. 177). The staphylococcus pyogenes aureus is shown at figs. 1 and 2; the streptococcus and its relation to the epithelium, leucocytes, and the tissues in figs. 3 to 5; and the gonococcus at figs. 7 to 9. Fig. 9 is given because of its bearing upon the infection of the infant in its passage through a gonorrhœal vagina, which produces the most destructive form of "ophthalmia neonatorum."

Vaginal
Secretions.
Döder-
lein's
View.

The normal vaginal secretion has been already referred to on p. 32. Its source and characteristics are still a matter of discussion. Döderlein would distinguish between physiological and pathological secretions: the former being markedly acid and containing a "vaginal bacillus," which causes the acidity by producing lactic acid; the latter feebly acid, neutral, or alkaline, and showing different micro-organisms, saprophytic and pathogenic. Some 50 per cent. of pregnant women have this pathological secretion in which germs flourish, and auto-infection is possible.

Could this distinction be drawn, the study of infection would be much simplified. The discharge becomes alkaline at the menstrual period, in the puerperium, and in many cases of leucorrhœa; and thus conditions arise in the vagina favourable for the growth of micro-organisms, and infection of the genital tract. Unfortunately the results of Krönig, Menge, and Walthard do not support Döderlein's view. The question is much more complicated, and several factors are at work, the analysis of which will require much laborious investigation.

Krönig's
Results.

As the result of investigations on pregnant and puerperal women, which are beyond the scope of a text-book on Gynecology, Krönig has come to the conclusion that the distinction between physiological and pathological secretion does not hold, that all secretions alike contain no pathogenic germs. All secretions are equally germicidal, although there is a difference in the vitality of the germ, the staphylococcus taking twice as long to kill as the streptococcus. A vagina becomes aseptic in two to three days. The cause of this bactericidal power is not evident. It is not chemical alone, because it is as potent whether the secretion be faintly or strongly acid. Nor is it due to a special bacillus, although some micro-organisms may be antagonistic to others. If the leucocytes play a part it must be through some property independent of their con-

¹ The efficiency of this appears from Schultze's investigation as to the gonococcus. Of 104 patients with gonorrhœa, only nine showed the gonococcus in the vagina as against eighty-one in the cervix, and seventy-eight in the urethra. In explaining this, the destructive action of the secretions in these localities must be taken into account, but the like preponderance in the urethra and cervix as against the vagina suggests that the epithelial covering in which the cervix and urethra alike differ from the vagina, is an important factor: *Cent. f. Gyn.*, 1896, S. 744.

tractile power, as a temperature which destroys this does not affect their germicidal action. Nor can it be explained by the want of oxygen in the vagina, as the staphylo- and streptococci are anærobic, that is, growing independent of oxygen, and yet they are killed. Nor is it mechanical, because particles of carbon and mercury are removed more slowly. Although he has thus shown that no one known factor is of itself sufficient to account for the germicidal action, it does not follow that all of these may not jointly operate.

His most important practical observation is that corrosive sublimate injections destroy the natural germicidal action, perhaps through precipitating albumen, while plain water only lessens it. Hence prophylactic injections of corrosive sublimate do harm in cases of normal secretion.

For gynecological work, the investigations of Menge on *non-pregnant* patients are of peculiar interest. To determine the efficiency and rapidity of the germicidal action, he introduced pyogenic organisms into the vagina of eighty cases, and found that the vagina "cleansed itself" from these in periods varying from two-and-a-half hours to three days.¹ Several factors are concerned in this germicidal action, which, ranged according to their potency, are—various forms of bacteria, their products, acidity of secretion, a possible serum-action, leucocytes, and absence of oxygen. A weakening of their activity occurs under the following conditions—in infants, at the menstrual period, in cases of increased secretion from cervix or body of uterus or vagina itself, where the vulva is patulous or uterus prolapsed, and at the menopause.

The effect of a change in the pabulum restoring to micro-organisms a virulence which they had lost, has been for long recognised, and an interesting experiment of Walthard's must be mentioned here for its bearing on gynecological as well as obstetrical work. He inoculated the streptococcus into a rabbit's ear, and found that no serious result followed unless the ear was ligatured so as to lessen the resistance of the tissues. Under these circumstances a vaginal streptococcus became as virulent as that found in puerperal fever. Hence an innocuous streptococcus may become fed up on bruised tissues during the puerperium so as to recover a lost virulency; and the same must hold good for gynecological operations where there has been bruising of tissue, as in the enucleation of fibroids.

VARIETIES.

Until the bacteriology of the vagina is better known we shall not be in a position to classify the forms of vaginitis according to etiology. The distinction, however, between *simple* and *specific* (gonorrhœal) is of

¹ He distinguishes between the vaginal orifice and the vaginal vault. The above is the time taken for the vaginal vault to become germ-free after the artificial inoculation of the secretion, the vaginal orifice naturally takes longer.

great clinical importance owing to the intractability and far-reaching effects of the latter. The distinction between acute and chronic is merely one of degree. Special varieties are the *emphysematous*, the *diphtheritic*, and the *senile vaginitis*, which forms one of the physiological changes of the menopause.

PATHOLOGY.

Gonor-
rheal
Vaginitis.

This has been already touched on in what has been said on micro-organisms. Of these the gonococcus is the most important for the




FIG. 296.

GRANULAR VAGINITIS—acute form (Schroeder).

physician to recognise, as its detection aids diagnosis. It was first described by Neisser, and has the appearance shown at fig. 8 in




FIG. 297.

GRANULAR VAGINITIS—chronic form (Schroeder).

Plate IV. When found it is diagnostic,¹ although its absence does not of course show that the secretion is not gonorrhœal.

Simple
Vaginitis.

As to the *anatomical changes*, we find most frequently in vaginitis slight elevations of the mucous membrane, which produce a granular surface. These granulations consist of groups of papillæ infiltrated with small cells; these swell up and push before them the stratified squamous epithelium, the superficial layers of which are shed (fig. 296). When

¹ Bumm—Beitrag zur Kenntniss der Gonorrhoe der weiblichen Genitalien: *Archiv f. Gyn.*, Bd. xxiii., S. 327.

the condition has existed some time, the surface becomes more level through the thinning of the epithelial covering (fig. 297).

Associated with vaginitis in pregnancy, there is sometimes an *emphysematous condition* of the vaginal mucous membrane. Winckel has described cysts containing gas and fluid; according to Ruge, the air is present in spaces among the cellular tissue (fig. 298), while Zweifel thinks they arise from vaginal glands the ducts of which have been closed by inflammation. Oliver¹ has noted a similar condition after the menopause.

Diphtheritic vaginitis occurs either as localised patches or as an affection of the whole vagina. In the latter case the mucous membrane may be so swollen that the finger scarcely reaches the cervix, which also is found to be thickened and covered with the diphtheritic membrane.

The cicatricial contraction of the vagina observed after the menopause is due to a *senile vaginitis*. The epithelium is shed in patches,




FIG. 298.

COLPITIS EMPHYSEMATOSA (Schroeder).

and the raw surfaces thus produced adhere together (*Hildebrandt*). This process is similar to that which produces occlusion of the cervical canal after the menopause.

ETIOLOGY.

The following are the most important causes:—

- Gonorrhœal infection;
- Irritating discharges from the uterus;
- Injurious vaginal injections, badly fitting pessaries, or other causes which injure the vaginal mucous membrane;
- Exanthemata.

Gonorrhœal infection produces the most intractable form of vaginitis, *Gonorrhœa*, which may extend over months or years. The poison may spread

¹ Oliver, J.—A peculiar crackling (emphysematous) sensation in the vaginal canal: *Brit. Med. Jour.*, 1895, p. 805.

along the mucous membrane of the uterus and Fallopian tubes causing endometritis (p. 350), pyosalpinx (p. 227), and pelvic peritonitis (p. 182).

Endometritis.

Irritating discharges from the uterus, as in endometritis, produce a secondary vaginitis which can only be treated by curing the uterine affection. In carcinoma and vesico-vaginal fistulæ, vaginitis arises secondarily.

Mechanical Irritants.

Among the causes which irritate or injure the vaginal mucous membrane, we mention injections of too hot or too cold water and of substances to produce abortion, badly-fitting pessaries, tampons or pieces of sponge which have been allowed to lie some days in the vagina. Vaginitis may also develop on a patient's entering married life, simply from awkwardness in sexual intercourse; on being consulted about such cases, we must remember that *a simple vaginitis may produce most of the symptoms of one due to gonorrhœa*.

Exanthemata as a cause.

Diphtheritic inflammation occurs usually in the puerperal condition and that through bad hygiene. It has been observed in *typhus*, *small-pox*, and *cholera*, and also in some cases of gonorrhœa. Localised diphtheritic patches are seen in fistulæ, in carcinoma, and round badly-fitting pessaries.

SYMPTOMS.

These are the following:—

- A burning heat in the vagina;
- Pain in the floor of the pelvis;
- Frequent desire for micturition, with a scalding sensation while water is passing;
- Free muco-purulent leucorrhœa.

These symptoms are present both in simple vaginitis and that due to gonorrhœal discharge. In the latter case, the urinary symptoms are more pronounced; there is a distinct period from which all the symptoms commenced, their duration is longer, and they resist treatment; they are often complicated with those of enlarged inguinal glands, endometritis, cystitis, or pelvic peritonitis.

DIAGNOSIS.

On vaginal examination, the finger recognises the discharge which escapes on separating the labia, and, in many cases, the rough condition of the mucous membrane.

The speculum shows that the mucous membrane is inflamed and covered with muco-purulent discharge; the redness is usually in the form of patches but may be diffuse.

The appearance of the cervix must be noted to ascertain that the leucorrhœal discharge does not come from it; the differentiation of discharge from the uterus and that from the vagina, is made as described on page 337.

The *differential diagnosis* between *simple* and *gonorrhœal* vaginitis is often very difficult. The history of a distinct source of infection is the only certain guide, and the ascertaining of this is a very delicate question. Apart from this, the following conditions point to a gonorrhœal origin: sudden development of vaginitis with urinary symptoms, in a patient who has had previously no marked leucorrhœal discharge; absence of any other cause to explain these; protracted duration of symptoms and resistance to treatment. However convinced the practitioner may be in his own mind that the vaginitis is of a specific nature, the social unhappiness caused by his expressing a decided opinion should deter him from giving it in cases where a cause is not admitted.

Diagnosis
of Specific
Vaginitis.

Pelvic abscesses discharging through the roof of the vagina have been mistaken for vaginitis. Such a mistake will not arise when the bimanual and other methods of examination are employed. We must not be satisfied with finding vaginitis; the whole routine examination of the pelvic organs must be made after the pressing symptoms have been relieved.

TREATMENT.

In *acute cases*, rest in bed is necessary. Hot water injections are given three or four times daily: the douche is much more convenient than the syringe; it leaves the hands free, requires less exposure of the patient, and keeps up a steady stream (*v. p.* 150). The stream should run for a quarter of an hour. A piece of gutta-percha tubing, weighted at one end and with a clip at the other, makes a handy douche; the weighted end is placed in a ewer of water above the level of the bed, the tube is coiled up in the water so as to be filled, the clamp is put on at the other end and the tube withdrawn; the syphon-action is started by the column of water in the tube and continues till the ewer is empty. The bowels are freely moved, and then a morphia suppository is given. Complete rest from sexual activity is absolutely necessary.

In *chronic cases* or after the acute stage has passed off, astringents are added to the injections. The vaginal walls having been first thoroughly dried, a solution of nitrate of silver ($\frac{3}{4}$ to $\frac{5}{8}$ of water) is applied and a tampon of antiseptic cotton soaked in glycerine and bismuth introduced to keep the walls apart. Chloride of zinc (2 grs. to $\frac{3}{4}$) is recommended by Fritsch.

Medicated
Pessaries.

Applications to the vagina are usually made by means of medicated pessaries. The following are those most frequently used¹:—

Atropine - - -	Sedative - - -	1-20 grain.
Belladonna - - -	do. - - -	2 do. Alo. Ext.
Morphia - - -	do. - - -	$\frac{1}{2}$ do.
Bismuth Oxide - -	Cicatrising & emollient	15 do.
Borax - - -	do. do.	15 do.
Zinc Oxide - - -	do. do.	15 do.
Tannin - - -	Astringent - - -	10 do.
Alum - - -	do. - - -	15 do.
Acetate of Lead and Opium - - -	do. - - -	5 do. 2 grs. Opium.
Gallic Acid - - -	do. - - -	10 do.
Persulphate of Iron	Hæmostatic - - -	5 do.
Sulphate of Zinc (dried) - - -	Caustic - - -	10 do.
Iodide of Lead - -	Alterative & resolvent	5 do.
Mercurial - - -	do. do.	30 do. (<i>Ung. Hydrarg.</i>)
Carbolic Acid - -	Deodorant - - -	5 do.

Tampons.

Lawton's absorbent cotton² is the best material for vaginal tampons which are to be soaked in glycerine or other medicaments.

VAGINISMUS.

LITERATURE. *Duncan, Matthews*—Diseases of Women, p. 142: Lond., 1883. *Henrichsen*—Strictur des Scheidengewölbes, bewirkt durch Krampf des Musculus Levator Ani: Archiv f. Gyn., Bd. xxiii., S. 59. *Hildebrandt*—Ueber Krampf des Levator Ani beim Coitus: Archiv f. Gyn., Bd. iii., S. 221. *Scanzoni*—Lehrbuch der Krankheiten der weiblichen Geschlechtsorgane, S. 704: Wien, 1875. *Simpson, Sir J. Y.*—Edin. Med. Journ., Dec. 1861. And Diseases of Women, p. 284: Edin., 1872. *Sims*—Cases of Vaginismus: Americ. Med. Times, 1862, Nos. 22 to 25. *Thomas*—Diseases of Women, p. 203: Lond., 1882. *Tilt*—The Lancet, Aug. 1874.

Nature.

By vaginismus, we understand a painful reflex contraction of the muscular fibres surrounding the vaginal orifice—just as laryngismus is applied to the same condition in the larynx. Marion Sims first drew attention to this condition.

ETIOLOGY.

It is found in some patients of a nervous and sensitive temperament without there being any local source of irritation, but this is exceptional.

Usually one of the following conditions is present:—

- An irritable spot in the fossa navicularis;
- An inflamed hymen which has not been ruptured, or irritable carunculæ myrtiformes;
- Fissures in the fourchette or round the vaginal orifice;
- Small ulcers within the hymen;
- Fissure of the anus;
- Urethral caruncle.

¹ As made up and supplied by Messrs Duncan, Flockhart & Co.

² Sold in packets (2 oz— $\frac{1}{2}$ lb.).

SYMPTOMS AND DIAGNOSIS.

Dyspareunia and sterility are the leading symptoms.

By *dyspareunia* (a term introduced by Barnes), we understand painful or difficult sexual intercourse; hence the conditions which produce vaginismus arise on the patient's entering married life. The suffering may be so great that medical advice is at once sought; often a sense of delicacy prevents this till the condition has existed some time. Dyspareunia.

In some cases there is a careworn and anxious expression of countenance, in others a hysterical manner. As the ordinary vaginal examination is painful—the patient involuntarily drawing away as soon as the painful spot is touched—it is best to make inspection of the genitals first. Here we may see any of the conditions mentioned under *Etiology*. Sometimes no local cause is evident; but on carrying the finger into the vagina the reflex contraction of the muscle is felt.

Hildebrandt has shown that this muscular contraction is sometimes noticed in the upper part of the vagina, and is then due to spasm of the levator ani. Henrichsen found well-marked contraction of the levator ani in one case; he refers it to the anterior portion of the muscle which springs from the pubes and passes to the vagina near the vulva.

The possibility that the dyspareunia may be due to some local pathological condition at the roof of the vagina (prolapsed ovary or cellulitis) and not at the ostium, should be kept in mind.

The *prognosis* as to cure is good. From the distressing nature of the symptoms, and the relief obtained by the means to be described, they prove very satisfactory cases for treatment.

TREATMENT.

First remove any cause of local irritation, as urethral caruncle or irritable carunculæ myrtiformes; in some cases it is necessary to clip away carefully the whole hymen. Divide the base of irritable fissures of the anus with the knife, or touch them with the actual cautery. Iodoform in powder or made into an ointment, is the best local application to allay irritation or favour healing. Its penetrating and disagreeable odour makes many patients object to it. This is diminished by keeping Tonquin beans in the powder, and by adding oil of eucalyptus or citronelle (10 m. to ʒi) to the ointment or pessary.

R Iodoform	gr. x.
Olei eucalypti	M. i.
Fiat pessarium	Mitte tales xii.
Sig. As directed.	

Cocaine, 5-20 p. c. solution or ointment, is also useful.

After the cause has been removed, the ostium vaginæ must be dilated.

This is best effected by making the patient wear a vaginal dilator night and morning, for an hour at a time; it may be made of wood or glass, and should have a bulbous end about $1\frac{1}{2}$ in. long. The conical form is not good. The pain caused by the introduction passes off after a time. Dilators of gradually increasing size should be used.

If the dilator cannot be worn, we must have recourse to Sims' operation. In some cases, when the vaginismus is evidently due to the narrowness of the ostium, and specially when a reflex contraction of the muscle is noted, this operation is done without previous use of the dilators.

Sims' operation.

Sims' operation for vaginismus. We have already seen (p. 12) that the bulbo-cavernosi muscles embrace the ostium vaginæ and form a kind of sphincter for it; their position is seen in fig. 7. To divide the superficial fibres of this muscle is the aim of the operation.

The patient being under chloroform, two fingers of the left hand are passed into the vagina so as to stretch the ostium. With an ordinary scalpel, an incision is made on each side of the fourchette; the incision is about 2 inches long, and extends from $\frac{1}{2}$ an inch above the ostium to the raphe of the perineum. The ostium is now thoroughly and firmly plugged with lint which is kept in place with a T-bandage; thorough plugging is essential as there is often smart hæmorrhage from the incisions. Next day the lint is removed and a glass dilator introduced, which must be worn for one or two hours night and morning during a period of several weeks.

Forcible Dilatation.

Instead of dividing the sphincter with the knife, it may be forcibly stretched with the fingers till the muscular fibre is ruptured. This is done by passing the thumbs (*Tilt*) or several fingers (*Hegar*) of each hand into the ostium, and then forcibly separating them till we feel the muscular fibre yield under the traction. The advantage of this method is that it is bloodless, and there is no granulating wound left to heal.

Pozzi's Operation.

Pozzi¹ makes a lateral incision as in Sims' operation, and dissects the mucosa back from it so as to produce a lozenge-shaped raw surface, the long axis of the lozenge corresponding to the incision. He then brings together the margins in such a way that the line of suture lies in the other axis of the lozenge, *i.e.*, at right angles to the direction of the incision, or parallel to the hymeneal orifice which is thus increased in circumference.

With these local measures, we should always combine constitutional treatment. Exercise, fresh air and change of scene are beneficial. It is self-evident that complete rest to the sexual system must be strictly enjoined during any course of local treatment; this should be maintained for some time afterwards, which may be secured by recommending a few weeks' residence from home. Tonics (such as quinine, iron, and arsenic) are given as the case requires.

¹ *Annal. de Gyn.*, 1894, tom. xli., p. 591.

TUMOURS OF THE VAGINA.

Under tumours of the vagina we briefly describe the following:—

Cysts,
Fibroid tumours,
Carcinoma,
Sarcoma,
Tuberculosis.

Syphilitic ulceration does not call for special description. Lipoma has also been described.¹

CYSTS OF THE VAGINA.

LITERATURE. *Breisky*—Die Krankheiten der Vagina, S. 130: Stuttgart, 1879. *De Sinéty*—Manuel pratique de Gynécologie, p. 164: Paris, 1879. *Fischel*—Casuistischer Beitrag zur Lehre von den Scheidencysten: Archiv f. Gyn., xxxiii., S. 121. *Gräfe*—Zehn Fälle von Vaginalcysten: Zts. f. Geb. u. Gyn., Bd. viii., S. 460. *Johnston*—A Contribution to the Study of Cysts of the Vagina: Americ. Jour. of Obstet., 1887, pp. 1121, 1241. *Lebedeff*—Beitrag zur Lehre über Vaginalcysten: Zts. f. Geb. u. Gyn., Bd. viii., S. 324. *Mundé*—Case of Cyst of the Vagina: Americ. Jour. of Obstet., Vol. x., p. 673. *Routh*—On Cases of Associated Parovarian and Vaginal Cysts formed from a distended Gartner's Duct: London Obstet. Trans., 1894, p. 152. *Rutherford, H. T.*—Cysts of the Vagina, their Etiology, Pathology, and Treatment: London Obstet. Trans., 1891, p. 354. *Veit*—Ueber einen Fall von sehr grossen Scheidencysten: Zts. f. Geb. u. Gyn., Bd. viii., S. 471. Also, Erkrankungen der Vagina: Handbuch der Gynäkologie, Wiesbaden, 1897, S. 233. *Von Preuschen*—Ueber Cystenbildung in der Vagina: Virchow's Archiv, Bd. lxx., S. 3. Rutherford's and Johnston's papers discuss fully the literature of the subject.

These, though the commonest of vaginal tumours (Winckel), have not frequently been recorded, probably because they may easily escape detection. Johnston has collected 168, to which Rutherford has subsequently added 52.

Pathology. They are situated most frequently in the anterior vaginal wall, and usually in the lower third but within the ostium. They are generally single, rarely have two or more been found together. They are lined with a single layer of cylindrical epithelium which contrasts with the many layers of squamous epithelium of the vaginal mucous membrane from which they lie separate (fig. 299). We have seen them of the size of a hen's egg. Their contents vary from a clear thin fluid to a gelatinous chocolate-coloured inspissated mucus. Fischel and others have also found cysts lined with an endothelium, and he has demonstrated their connection with the lymphatics; these cysts, which must be regarded as dilated lymphatics, are much rarer than those lined with cylindrical or pavement epithelium. Chéron² found a calculus in a cyst of the anterior wall, which communicated with the urethra; he refers to observations by Priestley, Simon and others, of vaginal cysts associated with urethrocele, and would account for this condition by the coalescence of a cyst with the urethra.

¹ Conrad—Cent. f. Gyn., xii., S. 214.

² Archives de Toc., 1887, p. 539.

Vaginal
Cysts—
Etiology.

Etiology. As there are hardly any mucous glands present in the vaginal mucous membrane, the mode of origin of these cysts is disputed. In some cases they can be traced to crypt-like depressions of the mucous membrane which become shut off (*Von Preuschen*). It has been suggested by Veit that they are due to persistence of the canals of Gartner, rudimentary structures which run alongside of the uterus and vagina (cf. Pl. VII., and p. 260). Routh's case in which the vaginal cyst communicated with the cyst in the layers of the broad ligament is of special interest in this connection; and cases supporting this view have been recorded by Milton¹ and Senn.² Freund has ascribed them to the rudiment of a Müller's duct, but the existence of this with a well developed vagina and normal uterus is improbable. A case of suppurating hydatid of the vagina has been recorded by Porak.³ Thorn⁴ accounts for some cysts by traumatic blood and lymph extravasations.




FIG. 299.

SECTION OF VAGINAL CYST (*Schroeder*). The cyst wall which is lined with a single layer of epithelium is separated by some tissue from the mucous membrane which is covered with many layers of squamous epithelium not detailed in the section.

Symptoms. These are often nil; and such cysts readily escape observation, so that they may be more frequent than is supposed. When of large size, they produce bearing down pain with leucorrhœa, and in some cases dyspareunia.

Diagnosis. Small cysts readily escape detection. When large, their smooth elastic surface and fluctuation make them easily recognised. They must not be confounded with cysts due to obstructed Bartholinian glands, which are situated on the labia minora or at the ostium. Careful examination will easily distinguish them from a pouching of the bladder or rectum.

Treatment. This consists in laying the cyst open and destroying its lining wall, which is best done by the cauter. Or we may cut out a

¹ *Lancet*, 1893, p. 324, II., p. 924.

³ *Archiv. de Tocolog.*, 1884, p. 163.

² *Amer. Jour. Obstet.*, 1895, Vol. i., p. 564.

⁴ *Centralb. f. Gyn.*, 1889, S. 658.

portion of the cyst wall, and stitch the margins of the rest to the adjoining vaginal mucous membrane so that the cyst is taken up into the vagina; this does away with the granulating surface and subsequent cicatrisation which accompany cauterisation. If the patient is past the menopause and the cyst gives no trouble, there is no occasion to interfere.

FIBROID TUMOURS OF THE VAGINA.

LITERATURE. *Breisky*—Die Krankheiten der Vagina: Stuttgart, 1886, S. 162. *A. R. Simpson*—Fibroma Vaginæ, Contributions to Obstetrics and Gynecology, p. 201: Edinburgh, 1880. *Veit*—Erkrankungen der Vagina: Handbuch der Gynäkologie, 1897, S. 349.

Pathology. Fibroid tumours rarely originate in the vagina; *Breisky* has collected only 58 cases out of the literature. Like fibroid tumours of the uterus, they consist chiefly of fibrous tissue with some unstriped muscular fibre; they are usually situated in the anterior wall, in 17 out of 27 cases (*A. R. Simpson*); they are pediculated (forming so-called fibrous polypi) or sessile.

Symptoms. These are produced only when the tumour is large. In the case described by *A. R. Simpson*, in which the tumour was the size of two fists, it interfered with micturition and the escape of the uterine discharges. *Da Costa* records a case in which the tumour projected from the vulva.¹

Diagnosis. This is easy, except in the case of large tumours when the pedicle is difficult to reach. The relation of the bladder should always be carefully ascertained by passage of the sound.

Treatment consists in division of the capsule and enucleation of the tumour when it is sessile, or ligature and division of the pedicle when it is pediculated.

CARCINOMA OF THE VAGINA.

LITERATURE. *Breisky*—Die Krankheiten der Vagina, Billroth's Handbuch: Stuttgart, 1879, S. 151. *Bruckner*—Der primäre Scheidenkrebs und seine Behandlung: Zeitschrift für Geburtshilfe und Gynäk., B. vi., Hft. 1, S. 110. *Fenger*—Total Extirpation of the Vagina for Carcinoma: Amer. Jour. Obst., 1893, p. 218. *Goodell*—Boston Gyn. Jour., Vol. vi., p. 383. *Küstner*—Ueber den primären Scheidenkrebs: Archiv f. Gynäk., Bd. ix., S. 279. *Mackenrodt*—Primäres Carcinom der hinteren Scheidenwand: Cent. f. Gyn., 1892, S. 529. *Oliver, J.*—Cancer of the Vagina at the age of twenty-six: Liverpool Med. Ch. Jour., 1891, p. 272. *Parry*—Primary Cancer of Vagina: Amer. Jour. of Obstet., Vol. v., p. 163: and Philad. Med. Jour., Feb. 1873. *Simpson, A. R.*—Contributions to Obstetrics and Gynecology, p. 205: Edinburgh, 1880.

Pathology. Primary carcinoma occurs very rarely in the vagina—in 14 out of 8287 cases (*Beigel*); in the paper cited above, *Fenger* has collected but 57 cases out of the whole literature. This is the more

¹ The tumour measured, antero-posteriorly, 6½, and transversely, 4 inches: *Amer. Jour. Obstet.*, 1895, Vol. xi., p. 940.

surprising when we remember how very frequently it affects the cervix. It occurs in two forms, either as a localised broad-based papillary swelling seated most frequently in the posterior wall; or as a diffuse infiltration which often constricts the canal in a ring-like manner. The inguinal glands are generally enlarged by carcinomatous infiltration.¹

Symptoms and Diagnosis. As in carcinoma of the cervix, there is hæmorrhage and fœtid discharge: the pain is slight in the early stage. The diagnosis that there is *primary* carcinoma of the vagina is often doubtful, because it is difficult to ascertain the condition of the cervix and uterus; in the specimen represented at fig. 281 it was supposed to be primary until the post-mortem showed that it was secondary to carcinoma of the cervix. The examination per rectum is useful in these cases.

Treatment. This consists in the removal of as much as possible of the diseased tissue with the cautery, spoon, or knife. Operative methods for total extirpation of the vagina have recently been described by Olshausen² and Dührssen;³ and a case has been recorded by Fenger.⁴

SARCOMA VAGINÆ.

LITERATURE. *Breisky*—Die Krankheiten der Vagina: Billroth's Handbuch, S. 150. *Munn*—Sarcoma of the Vagina: Amer. Jour. of Obst., Vol. viii., p. 541. *Oliver, J.*—Two Cases of Primary Sarcoma of the Posterior Wall of the Vagina: Med. Press and Circ., London, 1892, p. 475. *Pick*—Ueber Sarkome des Uterus und der Vagina im Kindesalter und das primäre Scheidensarkom der Erwachsenen: Archiv f. Gyn., Bd. xlv., S. 191. *Simpson, A. R.*—Contributions to Obstetrics and Gynecology, p. 204: Edin., 1880. *Smith*—Amer. Jour. of Obst., Vol. iii., p. 671. *Spiegelberg*—Zu den Sarkomen des Uterus und der Scheide: Arch. f. Gyn., Bd. iv., S. 344. *Veit*—Erkrankungen der Vagina: Handbuch der Gynäkologie, Wiesbaden, 1897, S. 354. The literature is given in Veit's article.

Sarcoma of the vagina is still rarer than sarcoma uteri. It may arise very early in life, being sometimes apparently a congenital condition.⁵ Veit distinguishes sarcoma of the vagina in children from that in adults, the former resembling a teratoma, the latter sarcoma as generally described. He gives references, in the literature, to seventeen cases of the former,⁶ and thirty of the latter. As in the uterus, it is either diffuse or in circumscribed nodules (*v.* fig. 280). The symptoms are the same as in sarcoma uteri; and the treatment consists in removal (more easily effected in the circumscribed form), which, in a case reported by Spiegelberg, resulted in a permanent cure.

Schuckhardt⁷ has recorded three cases of operation for its removal in

¹ See two cases reported by J. Oliver: *Lancet*, 1894, ii., p. 257.

² Ueber Extirpation der Vagina: *Cent. f. Gyn.*, 1895, S. 1.

³ Ueber Extirpation der Vagina: *Cent. f. Gyn.*, 1895, p. 234.

⁴ *Loc. cit.*

⁵ As in a case of Graenicher's where a tumour was first noticed shortly after birth, removed at 15 months, and recurred at 4th year. *Centralb. f. Gyn.*, xiii., S. 591.

⁶ He does not include cases by Power and Butlin: *Brit. Med. Jour.*, 1895, xi., p. 973.

⁷ Ueber Sarkom der Scheide: *Archiv f. Gyn.*, xxxii., S. 400.

children under eight years of age, with the result that one was still without return after two years ; a second died from recurrence, while the third was operated on again for recurrence.

TUBERCULOSIS VAGINÆ.

LITERATURE. *Klob*—Patholog. Anat. d. weibl. Sexualorgane, S. 432: Wien, 1864. *Deschamps*—Etudes sur quelques ulcérations rares et non vénériennes de la vulve et du vagin : Archiv de Tocolog., 1885, p. 19. *Hegar*—Die Entstehung, Diagnose, und chirurgische Behandlung der Genitaltuberculose des Weibes : Stuttgart, 1886.

It is only of importance as part of a general affection, to be treated constitutionally. Hegar divides it into primary and secondary: the former is specially liable to arise after labour when the tissues are soft through direct infection from instruments, examining fingers or coitus; the latter takes place through the blood, or from the outside, *e.g.*, by germs from the stools. Barbier¹ says that the bacilli may be either in the seminal fluid itself or in the discharge from a tubercular epididymitis. Zweigbaum,² in reporting a case of primary tuberculosis of the cervix and vagina with secondary of the lung and intestines, has collected twenty-nine cases of tuberculosis of the vagina and cervix.

¹ *Gaz. Med.*, 1888, No. 39.

² *Brit. Med. Journ.*, 1889, i., p. 93.

177

THE HISTORY OF THE
CITY OF BOSTON
FROM THE FIRST SETTLEMENT
TO THE PRESENT TIME
IN TWO VOLUMES
BY NATHANIEL BENTLEY
OF THE BARR

THE FIRST VOLUME
CONTAINING THE HISTORY
FROM THE FIRST SETTLEMENT
TO THE YEAR 1780
LONDON: PRINTED BY J. JOHNSON, ST. PAULS CHURCH-YARD, 1780.

SECTION VII.

AFFECTIONS OF THE VULVA AND PELVIC FLOOR.

CHAPTER XLVII. The Vulva : Malformations ; Inflammation ; Tumours.

„ XLVIII. Rupture of the Perineum and its Operative Treatment.

„ XLIX. Displacements of the Pelvic Floor ; Prolapsus Uteri ;
Enterocoele.

CHAPTER XLVII.

THE VULVA : MALFORMATIONS ; INFLAMMATION ; TUMOURS.

LITERATURE.

- MALFORMATIONS.** *Hildebrandt*—Die Krankheiten der äusseren weiblichen Genitalien : Stuttgart, 1877, S. 2. *Meyer*—Virchow's Archiv, xi., p. 420. *Schroeder*—Die Krankheiten der weiblichen Geschlechtsorgane, S. 497 : Leipzig, 1879. *Simpson, Sir J. Y.*—Hermaphroditism : Collected Works, Vol. ii., p. 407. *Tait, Lawson*—Am. Gyn. Trans., Vol. i., p. 318.
- INFLAMMATION.** *Barbour*—On a Case of Ulcus Serpiginosum of the Vulva : Edin. Obstet. Trans., Vol. xxi., 1895-96, p. 150. *Hildebrandt*—Op. cit., S. 17 and 64. *Simpson, Sir J. Y.*—Diseases of Women, p. 286. *Thomas*—Diseases of Women, p. 122 : London, 1880.
- TUMOURS.** *Breisky*—Ueber Kraurosis vulvæ, eine wenig beachtete Form von Hautatrophie am Pudendum muliebre : Zeitsch. für Heilkunde, vi., 69. Also Centralb. f. Gynäk., 1885, 359. *Deschamps*—Epithélioma primitif de la vulve : Esthiomène : Archiv de Tocologie, 1885, pp. 120, 221. *Duncan, J. Matthews*—On the Hypertrophy of Lupus of the Female Generative organs : Lond. Obst. Tr., 1885, p. 230. See also Ed. Med. Jour., July 1884, and Clinical Lectures, 1886. *Duncan, J. M. and Thin*—On the Inflammation of Lupus of the Pudendum : London Obst. Tr., 1885, p. 310. *Hildebrandt*—Op. cit. Chap. VII. *Huguier*—Mémoire sur l'Esthiomène : Memoires de l'academie de Médecine, t. xiv., p. 508. *Küstner*—Zur Pathologie und Therapie des Vulvacarcinoms : Zeitsch. f. Geb. u. Gyn., 1882, 70. *Lomer*—Zur Casuistik des Carcinoms der Vulva : Ztschrift. f. Geb. u. Gyn., 1882, 167. *MacDonald, Angus*—Lupus of the Vulvo-anal region, with cases : Ed. Obst. Tr., ix., 49. *Peckham*—A Contribution to the Study of Ulcer Lesions of the Vulva : Am. Journ. Obst., 1887, p. 785. *Simpson, A. R.*—Article "Pruritus" in Quain's Dictionary of Medicine, p. 1767. *Smyly*—Diseases of the External Genital Organs : Playfair and Allbutt's Gynecology : London, 1896. *Tait, Lawson*—Climacteric Diabetes in Women : Practitioner, June 1886. *Taylor, J. E.*—Lupus or Esthiomène of the Vulvo-anal region : Am. Gyn. Tr., vi., 199. *Webster, J. C.*—The nerve-endings in the Labia minora and clitoris, with special reference to the Pathology of Pruritus Vulvæ : Soc. Rep. R.C.P.E., Vol. iii., p. 251. *Zweifel*—Die Krankheiten der äusseren weiblichen Genitalien und die Dammrisse : Handbuch der Frauen-krankheiten, Billroth and Luecke, Bd. iii., Stuttgart, 1886.

MALFORMATIONS.

Develop-
ment.

THESE are easily understood when we remember the normal development of the external organs of generation. 1. At the sixth week of foetal life, the *genital eminence* appears externally ; at this period the rectum, allantois and ducts of Müller communicate with one another

but not with the exterior (fig. 300). 2. At the seventh or eighth week a depression of the skin (known as the *genital cleft*) occurs; this extends inwards till it meets the conjoined allantois and rectum, and thus the cloaca is formed (fig. 301). 3. The tissue between the rectum and the




FIG. 300.

R rectum continuous with All allantois (bladder) and M duct of Müller (vagina). x Depression of skin below genital prominence which grows inwards and forms vulva (Schroeder).




FIG. 301.

The depression has extended inwards and becoming continuous with the rectum and allantois, formed the cloaca cl (Schroeder).

allantois grows downwards, and divides the cloaca into an anterior part (the *urino-genital sinus*), where the ducts of Müller end blindly in the eminence of Müller (v. fig. 55) and a posterior part (the anus): thus the *perineum* is formed (figs. 302 and 303). 4. The urino-genital sinus




FIG. 302.

The cloaca is becoming divided into urino-genital sinus Su and anus u by the downward growth of the perineal septum. The ducts of Müller have united into the vagina V (Schroeder).




FIG. 303.

The perineum is completely formed (Schroeder).

contracts in its upper portion to form the urethra; while the lower part gives rise to the lower third of the vagina, the urethra, and the vestibule (fig. 304). The ducts of Müller coalesce to form the upper two-thirds of the vagina (v. p. 83).




FIG. 304.

The upper part of the urino-genital sinus has contracted into the urethra; the lower portion persists as the vestibule Su (Schroeder).

The parts round the vulva develop, therefore, as follows: the *clitoris* from the genital eminence, the *labia minora* from the margins of the genital cleft, the *vestibule* from the urino-genital sinus.

Malforma-
tions.

The following malformations have been described. 1. *Complete atresia of the vulva* through the non-formation of the depression of the skin (fig. 300); the allantois and rectum either communicate as in fig. 300 or have become separated. This condition has only been found in foetal monstrosities. 2. *Persistence of a cloaca* so that the rectum, vagina and urethra have a common orifice (fig. 301); such cases are sometimes spoken of as atresia of the anus but are really due to non-formation of the recto-vaginal septum. 3. *Persistence of the urino-genital sinus* into which the bladder opens directly as the urethra has not formed (fig. 303); in such cases the vulvar orifice is contracted and opens into a long narrow vestibule which, at its farther end, communicates with the bladder and vagina. This condition is sometimes described as hypospadias.

HERMAPHRODITISM.

For a detailed description of this condition with illustrative cases, the student should consult Sir J. Y. Simpson's exhaustive article on Hermaphroditism (Collected Works, Vol. ii., p. 407).

Of hermaphroditism ('Ερμῆς and 'Αφροδίτη) there are two varieties, true and spurious.

True
Hermaphrodit-
ism.

By *true Hermaphroditism*, we understand that from the Wolffian bodies both ovary and testicles have developed so that both forms of gland co-exist in the same individual. This is an extremely rare occurrence; when it has occurred, there is a tendency towards the better development of one form of organ (determining the sex) while the other is rudimentary. According to Hildebrandt (*loc. cit.*, S. 6), only two authentic cases of bilateral hermaphroditism (ovary and testicle present on each side) have been recorded; of unilateral hermaphroditism (ovary and testicle present on one side), the other side having only one form of gland, a case has been recorded by Bannon; lateral hermaphroditism (ovary on one side and testicle on the other) has been more frequently met with, and cases, confirmed by microscopic examination, have been recorded by Berthold, Barkow, and Meyer.

False
Hermaphrodit-
ism.

By *false or pseudo-hermaphroditism*, is understood a malformation of the external organs so that they simulate those of the opposite sex. This occurs in two forms. 1. The external organs in the female may simulate those of the male. This is due to a hypertrophy of the clitoris and its prepuce, with approximation of the labia majora (simulating a scrotum) and contraction or occlusion of the ostium vaginae; in very rare cases, the clitoris is perforated by the urethral canal. This condition is seen at fig. 305, which represents the pelvis and external organs of an infant christened as a boy; a post-mortem dissection showed that the sex was female.¹

¹ Ramsbotham—*Medical Gazette*, xiii., p. 184.

2. The external organs in the male may simulate those of the female; the non-closure of the lower surface of the urethra and perineum, which constitutes hypospadias, produces an appearance resembling the external organs in the female. Numerous cases are on record in which the sex of males has been mistaken, even by medical experts, and the persons have entered married life as belonging to the female sex. The penis may be small and imperforate, the urethra opening at its base; the perineal fissure, lined by mucous membrane, may closely resemble the vagina; and the halves of the scrotum may appear like labia. This condition is seen at fig. 306: the case is reported by Otto;¹ the person lived in a state of wedlock with three husbands before the true sex was ascertained by medical examination.




FIG. 305.

SPURIOUS HERMAPHRODITISM (Sir J. Y. Simpson).

Pelvis of a female infant in which the external organs simulated those of a male. *c* Uterus and appendages, *b* hypertrophied clitoris with a sulcus at its extremity *a*, which ended blindly, and did not communicate with the urethra.




FIG. 306.

Case of hypospadias in the male, making the external organs simulate those of the female. *aa* Lobes of scrotum; *b* imperforate penis, $1\frac{1}{2}$ inches long; *e* perineal fissures $1\frac{1}{2}$ inches deep, lined with mucous membrane, at bottom of which the urethral orifice *d* is seen; *c* the split urethra, with openings *f* of glands beside it—supposed to be orifices of prostatic ducts, of Cowper's glands, and of seminal canals.

Cases of epispadias, in which the urethra (through defect of the upper portion of the penis) is exposed along with a portion of the bladder, would only on hasty examination be mistaken for the external female organs. The exposed vesical mucous membrane with its skin margins

Epispadias
mistaken
for Her-
maphrodit-
ism.

¹ Sir J. Y. Simpson—*Op. cit.*, p. 427.

resembles the vagina with the labia, but it is situated above the pubis; further, below the penis we find the normal scrotum and testicles.

Diagnosis. In examining a case, proceed as follows. 1. Palpate the supposed labia carefully to ascertain whether testicles are present in them; the possibility of hernia of the ovaries into the labia and of non-descent of the testicle into the scrotum, must be kept in view. 2. Examine per rectum for traces of uterus or ovaries. 3. After puberty watch for the menstrual molimina or hæmorrhage in the female, and for development of sexual powers in the male. 4. Note secondary sexual characters: development of breasts, appearance of face, tone of voice, and inclination towards one or other sex.

Hermaphroditism, like malformations in general, lies beyond treatment.

INFLAMMATORY CONDITIONS OF THE VULVA.

Under this heading we shall consider—

Acute, chronic, and follicular Vulvitis;
Erysipelas, gangrene, progressive gangrene;
Pruritus vulvæ and Ulcus serpinginosum.

Abscess of
Bartho-
linian
gland.

Pathology. In the acute stage, the mucous membrane round the ostium vaginæ and urethra is red, swollen and painful. Sometimes the mucous glands are obstructed, and a form of acne develops; the Bartholinian glands may inflame and suppurate, producing an abscess about the size of a pigeon's egg; the sebaceous glands at the roots of the hair on the labia majora are sometimes specially affected, producing the "Folliculite vulvaire" of Huguier, an excessively rare affection. In the chronic stage, there is abundant secretion of creamy purulent matter; when due to gonorrhœa, papillomata form round the vaginal orifice. Erysipelas or gangrene usually occurs after labour, or in infants after fevers (J. M. Duncan). Progressive gangrene with destruction of parts may occur; and in old or young women we may get recurring boils, for which Duncan recommends rubbing with mercurial ointment.

Etiology. It is often secondary to vaginitis, and accompanies urinary fistula and carcinoma. Want of cleanliness and protracted exercise, specially in hot weather, produce it and that most readily in patients with much adipose tissue. It is sometimes occasioned by awkward coitus and by masturbation. In children, it is not uncommon; it is important to remember this, as the inflamed appearance of the vulva and the profuse discharge make the parents suspect that the child has been violated and has contracted specific disease. It is caused by irritation of urine, want of cleanliness, and the strumous diathesis; sometimes it takes an epidemic form in the children of a family or

district. These last are probably due to spreading of gonorrhœa¹ through want of cleanliness.

The Symptoms and Physical Signs will be apparent from what has been said under Pathology.

Treatment. Strict attention to cleanliness must be enjoined ; frequent bathing with warm water and the application of hot linseed poultices will ease pain. In children, the pain in micturition is relieved by its being done while in a warm bath. Sedative lotions such as acetate of lead and opium may be required :—

R	Tinct. opii.	℥ss.	
	Plumbi acetat.	℥i.	
	Aquam ad	℥vi.	M.

In chronic cases, frequent washing with 2 per cent. sol. of carbolic or with astringent lotion is necessary. In abscess of the glands, the pus is evacuated through the gland ducts on pressure, or by free incision. Occasionally a gonorrhœa of the duct of the Bartholinian gland persists so that the duct requires to be laid open.

PRURITUS VULVÆ.

Definition. An irritable condition of the external genitals producing excessive itchiness.

Pathology. The irritable region is at the upper convergent angle of the labia majora at the mons veneris ; it may extend from that over the vestibule and the vaginal orifice, and sometimes over the mons veneris on to the abdomen. The pathological changes in the skin which produce this irritability are not known, because the cases are not seen in an early stage. By the time that the irritation has become so unbearable that advice is sought, the skin is inflamed and excoriated by continued scratching which masks its original condition.

Etiology. Any irritating discharges from the vagina as in carcinoma, and even simple leucorrhœa as from senile vaginitis, may produce it. It occurs in diabetes—due to irritation from the sugar in the urine (*Friedreich*)—and in affections of the kidney and bladder, just as similar conditions produce irritation of the penis in man. In children, it accompanies vulvitis and has been traced to the passing of the *Oxyuris Vermicularis* from the anus to the vulva. It is also caused by whatever produces congestion of the labia—hence its occurrence at the menstrual period and in early pregnancy ; by irritable skin affections as herpes, eczema, and the parasitic eczema marginatum ; and by pediculi.

Symptoms. The irritation is not continuous but recurs periodically. In some cases, it appears only after taking a long walk or after getting warm in bed ; sometimes it is most marked before the menstrual period.

¹ Pott—*Archiv f. Gyn.*, xxxii., S. 493.

The irritability is slight at first but becomes aggravated by scratching. To obtain this temporary relief, the patient gradually avoids company and this, along with the constant irritation, has led in some cases to nervous depression and melancholia; sometimes the practice of masturbation is learned at the same time, and the consequent nervous symptoms gravely complicate the case.

J. C. Webster states that a fibrosis may be present in the nerve end-organs and nerves in the clitoris and labia minora: the nerves are compressed and destroyed, and sub-acute inflammatory changes are found in the connective-tissue of the affected skin.

Diagnosis. As the most hopeful cases for treatment are those in which a distinct removable cause is found, a thorough examination is necessary: (1) Carefully inspect the external genitals for irritating skin eruptions, and examine scrapings of the affected parts microscopically for parasites; (2) expose the vagina and cervix thoroughly with the speculum to ascertain whether there is irritating leucorrhœa, the plugging of the vagina with cotton wadding to check discharge from the vagina or cervix will help us to exclude this; (3) test the urine for albumen and sugar; (4) examine per rectum for any source of irritation there.

Treatment
of Pru-
ritus.

Treatment. We must first remove the cause. When parasites are present, the mercurial or sulphur ointment is required; with vaginal or cervical catarrh, a tampon of wadding and glycerine (with acetate of lead $\bar{3}$ ii to $\bar{3}$ i) in the vagina will check the irritating discharge. Attention to diet (which should consist largely of vegetables) and to the regular action of the bowels is necessary; when the gouty diathesis (with which pruritus is often associated in old patients) is present, lithia water is useful. It is a safe rule to forbid all stimulants. Frequent vaginal injections or sponging with warm water, followed by the application of boracic ointment or bismuth, will relieve mild cases; in more severe, the patient should have, several times a day, a warm sitz-bath combined with the douche; after this, iodoform is dusted over the vestibule, or, if the patient is recumbent, lint soaked in acetate of lead and opium lotion is laid between the separated labia. In some cases, chloroform and almond oil have given relief (*Scanzoni*).

R Chloroformi $\bar{3}$ ii.
Olei amygdalae $\bar{3}$ ii. M.
Sig. Apply externally as directed.

Preparations of mercury give benefit in other cases.

R Hydrargyri perchloridi $\bar{3}$ ss.
Aquæ $\bar{3}$ vi. M.
Sig. Apply externally as directed.

Schroeder has seen very good results from the application of carbolic acid of varying strength—1 to 40 up to 1 to 10. Solid menthol is also used. Where milder measures have failed, solid nitrate of silver well rubbed into the irritated parts and followed by cold water dressing has given relief. In parasitic cases a lotion of equal parts of sulphurous acid and glycerine may be used. To procure rest at night, morphia and chloral may be necessary; Hildebrandt has found tinct. cannabis Indicæ (m. 10-20) even more effective than these. A 4 per cent. solution of cocaine may be tried. Application of galvanic current has been used with success.¹

When medical treatment fails, as it very often does, a cure may be effected by clipping away the affected skin, and bringing together the raw parts with the continuous catgut suture.

ERUPTIONS ON THE VULVA.

The skin round the vulvar orifice may be affected with any of the eruptions found on other parts of the body. Of these the most important are erysipelas, eczema, prurigo, herpes, acne. These eruptions have the same character as when they occur in other situations, and their treatment is the same. Condylomata may be found on the skin, and mucous patches over mucous surfaces. Eczema is frequently caused by diabetes, according to Lécorché.² Hebra's plates of Skin Diseases illustrate these conditions very well; see also a paper in the *Annales de Dermatologie et Syphilographie* for April 1882, by Gougenheim and Soyer.

ULCUS SERPIGINOSUM.

Pl. XIII. shows the vulva of a patient³ affected with this condition, which is "a form of ulcer commencing in the genital region, and in the course of months and years extending in a perfectly concentric manner over a great part of the neighbouring regions of the extremity; it is completely different from the spontaneously healing *ulcus molle*, although at the commencement it is not easily distinguished from it."⁴

On microscopic section, peculiarly shaped cells are found round the vessels, which are "plasma cells" (Unna)—epithelial cells of connective-tissue origin. At the advancing edge are found short bacilli arranged in chains, which are probably the cause of the ulceration.

Unna regards it as a venereal disease, though less infectious than the "*ulcus molle*." It is a rare condition, only a few cases having been recorded.

¹ Blackwood, *Polyclinic*, 1885, No. 9; and v. Campe, *Central. f. Gyn.*, Bd. xi., S. 521.

² Du diabète dans ses rapports avec la vie utérine, etc.: *Annales de Gyn.*, Oct. 1885.

³ For history of this case see A. H. F. Barbour and Norman Walker—"On *Ulcus Serpiginosum Vulvæ*": *Scot. Med. and Surg. Journ.*, July 1897. The case was considered by Macdonald as lupus, but has since been shown by Norman Walker to be one of "*ulcus serpiginosum*."

⁴ Unna's "*Histopathology*," translated by Norman Walker.

Fig. 306* is a section, under a high power, of the tissue removed from the case seen in Pl. XIII. It shows the large nucleated round cells. The methyl-blue stain brought out, here and there, small bacilli though less abundant than in the cases described by Unna.




FIG. 306.*

MICROSCOPIC SECTION OF TISSUE from the case of *Ulcus Serpiginosum* in Pl. XIII.

This case has the special interest that it is one of those formerly reported by Macdonald as "lupus."

TUMOURS OF THE VULVA.

Under these we shall notice briefly—

Cysts of the Bartholinian glands,
Elephantiasis,
Neuroma,
Fibroma,
Lipoma,
Carcinoma,
Sarcoma,
Lupus,
Kraurosis.

This is also the most convenient place to refer to

Pudendal hernia,
Varix, hæmatoma and hæmorrhage.

Cysts and
Abscess of
Bartho-
linian
gland.

Cysts of the Bartholinian glands. The Bartholinian or vulvo-vaginal glands, which are the analogue of Cowper's glands in the male, are situated at each side of the ostium vaginæ (see fig. 7); their ducts (about 2 cm. long and wide enough to admit a fine probe) run upwards to about the middle of the ostium vaginæ, where their mouths may be seen in front of the hymen.

A cyst may form by dilatation of the *ducts* or of the *glands* themselves. When due to distension of the duct, it has at first an elongated oval form; when the gland itself is affected, there may be multiple cysts or a lobulated swelling. They generally occur on the left side.¹ The contents are thick mucus, which is clear or of a brownish tinge. Suppuration may occur and abscess form (*v. fig. 307*).

The symptoms are due to the discomfort of the swelling, which is most felt on walking. The diagnosis is easy, from the position of the swelling and its fluctuating character; when it has developed during the puerperium, we must differentiate it from hæmatoma (which after a time becomes firm from coagulation) and inflammation after injury.

The treatment consists in complete evacuation of the cyst and destruction of its walls. It is not sufficient to open it and allow the fluid to escape; we must cut out a portion of the wall and then plug




FIG. 307.

ABSCESS OF THE BARTHOLINIAN GLAND (*Huguier*).

the cyst with antiseptic lint. By far the best instrument is the thermo-cautery: we first puncture the cyst with it; when the fluid has escaped, we pick up the outer cyst wall with forceps and lay it fairly open with the cautery; we then cauterise the inner wall also. A piece of antiseptic lint is laid over the wound.

Cysts also occur in the labia minora²; they are very rare and their pathology is not known.

Elephantiasis. This is a common condition in tropical countries, but is comparatively rare in Europe and America although a minor degree of it is occasionally met with. Elephantiasis.

The pathological changes consist in a dilatation of the lymphatic spaces and ducts, with secondary formation of connective tissue and

¹ Bonnet—*Gaz. des Hôpitaux*, 1888, No. 69.

² Smith removed two such cysts: *Brit. Med. Journ.*, 1888, i., 250.

thickening of the layers of the cutis vera; sometimes the papillæ are specially enlarged, producing swellings which resemble condylomata in form. The labia majora are most frequently affected, next in frequency the clitoris; more rarely are the labia minora hypertrophied (*Mayer*).

It develops, according to Mayer, most frequently at ages of from 20 to 30 years—that is in the period of sexual activity. It has been traced to direct injury, but the most fruitful cause of minor degrees of hypertrophy is syphilis.

The symptoms are due to the weight and discomfort of the tumour which may reach to the knees. For drawings of the various forms, Esmarck and Kulenkampff's monograph *Die Elephantiaschenformen* (Hamburg, 1885) may be consulted. The treatment of the larger growths is removal with the thermo-cautery.

Neuroma. *Neuroma*, an exquisitely sensitive red papule which resembles a urethral caruncle, has been described by Sir J. Y. Simpson (see fig. 340); its occurrence, except at the urethral orifice, is extremely rare.

Fibroma. *Fibroma*. This springs from the labia majora, resembles in structure fibroid tumours of the uterus, and, like them, is embedded in cellular tissue or hangs down by a pedicle. Taylor has reported a case of fibroid of the vestibule.¹

Lipoma. *Lipoma* may arise from the fatty tissue of the mons veneris or labia majora. Emmet² describes a case in which the tumour hung down to the patient's knees and was supported in a bag round the waist; Stiegele³ removed one which weighed 10 lbs.

Carcinoma. *Carcinoma* of the vulva is rare in comparison with its frequency in the uterus. In 16,637 cases of tumours of the female sexual organs, Gwilt found that 7479 were cancerous; and of these, 72 (or 1 per cent.) were vulvar. The most frequent form is the cancroïd (*West*). It begins, usually on the inner surface of the labia majora, as small round nodules which elevate the skin; they may remain for a long time unnoticed, as their growth is at first slow and painless. After ulceration they spread more rapidly, and extend forwards and backwards but rarely into the vagina. The inguinal glands are early involved.

Complete removal before the glands are affected, is the only treatment. Küstner has advocated removal of the inguinal glands of the affected side if these are larger than those on the healthy side.

Sarcoma of the vulva is very rare. Geith and Terrillon⁴ have recorded cases of melanotic sarcoma. Haeckel has collected 10 cases of melanotic tumours,⁵ mostly sarcomatous.

Lupus Vulvæ. *Lupus vulvæ* is a condition drawn attention to by Huguier, West, Taylor, Matthews Duncan, Macdonald, and Peckham. Duncan has considered it very fully, and an able histological examination of his

¹ *Americ. Journ. Obstet.*, 1888, p. 434.

² *Op. cit.*, p. 601.

³ *Zeits. f. Chir. u. Geb.*, Bd. ix., S. 243.

⁴ *Ann. de Gyn.*, Bd. xxvi., p. 1.

⁵ *Archiv f. Gyn.*, xxxii., p. 400.

specimens was made by George Thin. It may be defined as a slow chronic hypertrophic condition of the pudenda, prone to ulcerate and erode, causing little pain, lasting long, and not infecting neighbouring glands or causing ill-health.

As to its pathology, it is a hypertrophic condition with tendency to ulcerate and cause stricture of urethra, vagina, or rectum. Pus is secreted by the ulcerated surface, and occasionally considerable destruction of parts is caused. The hypertrophy may be small (*lupus minimus*), large (*lupus hypertrophicus*), or forming irregular masses extending to the hip. Other terms have been used, viz., *lupus prominens*, *lupus serpiginosus*; it was termed by Huguier, "*Herpes l'Esthiomène*."

On microscopic examination, Thin found growth of fibrous tissue (ordinary white fibrous tissue) and absence of any neoplastic structure; exudation cells were also present. Blood-vessels were unusually numerous. The appearances thus differ from *lupus vulgaris*, cancer, or syphilis; they are somewhat analogous to elephantiasis, but differ from that condition in the non-implication of the lymphatics and the presence of inflammatory action.

The symptoms may be slight and not attract the patient's attention unless hæmorrhage or inflammation occurs. The physical signs are those of hypertrophy, ulceration, erosion, lasting for years, not implicating glands, and not markedly affecting the patient's health. Large hypertrophies usually affect the clitoris and labia majora; small ones, the urethral orifice and hymen (*Duncan*). The vagina and uterus may become affected.

The condition is rare, but good drawings are given by Duncan. It must be diagnosed from epithelioma and syphilis. Epithelioma is harder, implicates glands soon, and has shallow ulcerations. It may be mistaken for "*ulcus serpiginosum*" (see p. 585). In syphilis, the history is the great test. Jonathan Hutchinson alleges, however, that this lupus is always due to tertiary syphilis. There is good reason to believe that pudendal lupus is not *lupus vulgaris*, cancer, syphilis, nor elephantiasis, but is an affection *sui generis* whose etiology is unknown. The term "*lupus*" is thus a clinical one.

The prognosis is fairly good. Many can be relieved and some cured. In treatment, hypertrophied or ulcerated portions are removed or cauterised with Paquelin's cautery, and the patient put on arsenic and iron.

Kraurosis vulvæ or Atrophy of the Genitals. In old women, the pudenda shrink; the labia minora become very small; the vestibule atrophies and shrinks, making the urethral orifice patulous and causing painful ulceration (*v. fig. 340*). Urethral caruncle is often present. This can be cauterised, and the patient instructed to apply freely vaseline with 6 per cent. cocaine.

Microscopically, Breisky found the sebaceous glands of the labia few, a cicatricial condition of the papillæ and thinness of the rete Malpighii. The sweat glands were also diminished in number.

Pudendal
Hernia.

Pudendal hernia. This corresponds to scrotal hernia in the male. The round ligaments are the analogues of the spermatic cord, and after emerging from the inguinal canal pass into the substance of the labia majora which correspond to the scrotum; if the process of peritoneum surrounding the round ligaments—known as the canal of Nuck—does not become obliterated at birth, it forms a track for the hernia. In some very rare cases a hernia appears at the lower end of the labium majus, apparently directly from the pelvis.

Though it be very rare, the possibility of a hernia must be kept in mind on examining a tumour of the labia; the crackling feeling, the impulse communicated on coughing, and disappearance on taxis, indicate hernia. The serious consequences of cutting into such a hernia by mistake for an abscess, are self-evident.

Varix.

Varix. The plexus of veins which forms the erectile tissue of the bulbi vaginæ has been already referred to (v. p. 13 and fig. 7). A varicose condition of the veins sometimes occurs in pregnancy and with pelvic tumours. In a case described by Holden,¹ they formed, when the patient was erect, a tumour of the size of a child's head. When these vessels rupture and the blood is effused into the cellular tissue, a hæmatoma is formed.

Hæma-
toma.

Hæmatoma. This condition is also called "Thrombus" and "Hæmatocèle" of the vulva; the former term should be limited to a coagulum within a vein, and the latter to blood effusion into the peritoneal cavity. It arises most frequently during labour, from injury produced by the child's head; the effusion may appear rapidly, as a tumour from the size of a walnut to an orange or larger, or may take place gradually. It has also been known to occur independent of labour or pregnancy, as the result of a blow or violent muscular effort.

The treatment consists in the application of ice to the vulva, and regular evacuation of the bladder and rectum without the patient's being allowed to strain. With this treatment, the mass may be absorbed. Should inflammation occur, poultices are applied and pus is evacuated with the knife; if this occurs in the puerperal condition, special care is required to keep the wound aseptic by repeated washing with carbolic solution and dressing with carbolised lint.

External
Hæmor-
rhage.

External hæmorrhage from ruptured veins sometimes occurs. The rupture may be caused by muscular straining, or by a blow or wound of the vulva. The dilated state of the veins makes such an injury serious during pregnancy, and several cases of a fatal result from a blow or kick have been the subject of a criminal prosecution (*Sir J. Y. Simpson*).

¹ "Immense Vulvar and Vaginal Varix:" *N. Y. Med. Record*, July 1868.

The vascular tissues are forcibly driven against the pubic arch and cut on it. In a case recorded by Hyde,¹ hæmorrhage from a vein ruptured by a fall proved fatal in forty minutes. Those who suffer from varicose veins should lie down for some hours during each day ; should a vein rupture, the patient must lie down at once and apply pressure to the bleeding point.

¹ *Lond. Obst. Trans.*, Vol. x

CHAPTER XLVIII.

RUPTURE OF THE PERINEUM AND ITS OPERATIVE TREATMENT.

LITERATURE.

Bantock, G.—On the treatment of Rupture of the Female Perineum, Immediate and Remote: London, 1878. *Collis*—Dub. J. Med. Sc., May 1861. *Duncan, John*—Concerning the Closure of Abnormal Anus: *Lancet*, 1873, Vol. ii., p. 9. *Duncan, Matthews*—Papers on the Female Perineum: Churchill, London, 1879. *Emmet*—Principles and Practice of Gynecology: Third Edition: Philadelphia, 1884. *Fritsch*—Ueber Perineoplastik: *Centr. für Gynäk.*, 1887, No. 30. *Goodell*—Lessons in Gynecology: Philadelphia, 1880. *Hart, D. B.*—The Structural Anatomy of the Female Pelvic Floor: Edin., 1880. *Heiberg*—Om Perinäoraphi, saerligt med. Hensyn til Lawson Tait's Methode: *Gynäkalog. og. obst. Meddel.*, Bd. vi., Hft. 3, 1877. *Hildebrandt*—Die Krankheiten der äusseren weiblichen Genitalien: Stuttgart, 1877. *Malthe*—Norsk Magazin for Laejevid. enskab, II. Reihe, 24 Bd. (describes Voss' Method). *Sänger*—Ueber Perineorrhaphie durch Spaltung des Septum rectovaginale und Lappenbildung: *Volkmann's Sammlung*, No. 301. *Schroeder*—Die Krankheiten der weiblichen Geschlechtsorgane, S. 542: Leipzig, 1887. *Simpson, Sir J. Y.* Diseases of Women, p. 644. *Tait, Lawson*—A New Method of Operation for Repair of the Female Perineum: *Tr. Lond. Obst. Soc.*, 1880. Diseases of Women and abdominal Surgery, Vol. i., pp. 68-71. *Thomas*—Diseases of Women, p. 165: Philadelphia, 1880. See *Duncan* and *Hildebrandt* for literature.

Prelimi-
naries.

Preliminaries and Nomenclature.—The question as to the significance of rupture of the perineum is still debated, some authors believing it to be of no importance unless involving the anus and leading to incontinence of fæces, others holding that it is an important lesion even when not so extensive as to involve the bowel. The relation of rupture of perineum to prolapsus uteri is discussed in the next chapter: at present we consider rupture apart from this. The views advanced in Chapters II. and IV. must be kept in mind. The student should glance over these and look at the sections in Plates I. and II., and fig. 41.

Complete rupture into the anus is serious as it entails incontinence of fæces, as well as rectocele and some sinking of the pelvic floor from the partial loss of the bracing-up action of the levatores ani (*v. p.* 42).

Another point to be kept in mind is the anatomy of the triangular ligament. This is a piece of sheet fascia filling up the pubic arch and perforated by the vagina and urethra. It strengthens the vaginal walls by its grip and, according to Emmet, prevents their eversion. He

believes that the bearing down complained of by some women and associated with a lax condition of the vaginal walls or the existence of rectocele is due to undue distention of this fascia and separation of its lateral attachments; and he bases on this a special operation to be described shortly.

It will be most convenient to retain the nomenclature already used in the Section on Anatomy. The pelvic floor is made up of pubic and sacral segments, as already defined; in labour, each of these behaves characteristically—the pubic segment is drawn up, the sacral one driven down (Chap. IV. and fig. 48).

In this chapter we are specially concerned with the sacral segment. During parturition it is driven downwards and backwards by the advancing fœtus, and is more or less torn at its inferior angle. The term perineum is often vaguely applied; in this Chapter, however, the perineum is defined as *the inferior angle of the sacral segment* (v. p. 65). Fig. 308 shows the perineum. At its lower end, this part of the pelvic floor is made up of the following:—

1. Posterior vaginal wall in front of upper part of perineal body,
2. Hymen,
3. Fossa Navicularis,
4. Fourchette,
5. Perineal body and skin over its base.

These are mesial structures; laterally, we have the labia majora and minora.

The perineal body lies in greater part below the level of the vaginal entrance, and has as its functions—

- (1.) The union of the following muscles—levator ani, bulbo-cavernosus, transversus perinei, sphincter ani;
- (2.) The directing backwards of the anus;
- (3.) The strengthening of a part much stretched during parturition.

PATHOLOGY AND VARIETIES.

It should be kept in mind that both the vaginal orifice and the vulvar orifice are antero-posterior.

When the fœtal head is passing through the vaginal orifice, it distends it all round; while, when passing through the vulvar orifice, it distends the lower half of this only, *i.e.*, it does not stretch so much those parts of the vulva lying above the level of the meatus urinarius.

As the result of normal and abnormal childbirth, we get certain tears of the inferior end of the perineum. In all primiparæ there is laceration of at least the hymeneal orifice, usually mesial and posterior—the “inevitable laceration” of Matthews Duncan. There may be also laceration of the following structures: (a) the vaginal

Pathology
and
Varieties.

orifice, radiating; (*b*) vestibule; (*c*) fourchette; (*d*) labia minora; (*e*) perineal body to a varying depth, the most extensive involving the sphincter ani. Further, there is sometimes central rupture of the perineum. In this lesion, the skin over the base of the perineal body alone may be involved or only the vagina may be torn. Rarely do we find a lesion of vaginal wall, connective tissue, and skin, with an unruptured band of tissue between it and the fourchette—central tear (fig. 309), or perforation through the inferior angle of the thinned-out sacral segment.

ETIOLOGY.

Etiology. The following causes produce rupture in parturition:—

- (1) Passage of a large head or of an occipito-posterior rotated into sacrum; passage of the shoulders;
- (2) Narrowness of pubic arch;
- (3) Straightness of sacrum, as in flat or rickety pelvis;




FIG. 308.

THE SACRAL OR SUPPORTING SEGMENT OF THE PELVIC FLOOR (*Hart*). *e* Symphysis pubis; *f* perineum or inferior angle of sacral segment; *g* anus.

- (4) Syphilitic ulceration;
- (5) Rigidity of parts in elderly primiparæ;
- (6) Careless use of forceps;
- (7) Too early passage of hand into vagina to bring down arms in turning.

Comment on these would lead us too much into Obstetrics.

SIGNIFICANCE OF RUPTURE OF PERINEUM.

Rupture of the perineum involving the sphincter ani and leading to complete or partial incontinence of fæces is an important lesion and imperatively demands operation.




FIG. 309.

CENTRAL RUPTURE OF THE PERINEUM, the child was born not through the Vulva but through the Ruptured Opening (*Sir J. Y. Simpson*).

Rupture of the perineum alone and not involving the sphincter ani may give rise to no symptoms unless associated with other conditions causing prolapsus uteri. According to Emmet, the real accident in some cases of ruptured perineum is tear of the triangular ligament where

it is perforated by the vagina, but probably tear of muscle there is of greater importance.

TREATMENT.

Treatment. We take this up under the following heads :—

a. Prophylactic ;

b. Operative, immediate and deferred.

Prophy-
lactic.

a. *Prophylactic.* This properly belongs to midwifery. The obstetrician is too apt to think of the perineum as something that delays the exit of the foetal head, and to forget the gynecological aspect—that it is part of the supporting segment of the pelvic floor. Extensive tear of this during labour means not only a larger raw surface for septic absorption, but is also one factor predisposing to prolapsus uteri. The question, therefore, of guarding the head during its passage over the perineum is of importance but belongs to obstetrics. We may note however that the foetal head, in passing through the outlet, drives the sacral segment back and glides forward in a direction parallel to the driven-back posterior vaginal wall. The normal curve of the sacrum favours this latter motion.

The perineum may tear (1) from over-distension of the orifice, or (2) from the too forcible driving of the foetal head against it, *i.e.*, at right angles to the perineum ; (3) from descent of the sinciput owing to fixation of the occiput and thus substitution of the larger diameters of the head for the sub-occipito bregmatic.

Operative. b. *Operative treatment*, (1) immediate and (2) deferred. No practitioner should leave a labour case until he is satisfied, by actual inspection or digital examination, as to the amount of perineal tear. When the sphincter ani is involved, the operation is on no account to be deferred, but must be performed at the conclusion of the third stage. The practitioner should never run the risk of his patient's having incontinence of fæces.

(1.) *Immediate operation.* This belongs to obstetrics.

Deferred
Operation.

(2.) *Deferred operation.* This may be to operate for a rupture through the sphincter or to repair the perineal body. At present we consider only the former.

Preliminary remarks. In complete tear through the anus, the external sphincter, internal sphincter, and levator ani are torn. Fig. 310 shows this clearly, and also explains what has to be done. What is wanted is not skin union, but some operative measure by which the torn muscular ends can be vivified and united.

Diagnosis of long-standing rupture of perineum into anus. The patient complains of inability to control the passage of flatus or of fæcal matter when a call to stool happens ; she is especially troubled when diarrhœa is present. Sometimes there is a certain amount of control, when some of the fibres of the upper margin of the internal sphincter are

intact. Patients in the lower classes occasionally treat this unpleasant condition as of little moment; to a woman of any refinement, the condition is a most distressing one.

On inspection, the practitioner notes that the skin surface between the vaginal and anal apertures is gone, so that these apertures are blended. The finger passed into the rectum feels no muscular constriction, and notes that the anterior and posterior rectal walls are in contact. The perineal body appears to be gone, and a V-shaped projection of cicatrised mucous membrane (apex above) is all that remains of it.

Operation for restoration of function of sphincter ani. The patient's bowels are first freely cleared out by castor oil and enemata so as to ensure that no scybala remain. The nurse should be allowed two or




FIG. 310.




FIG. 311.

LINES OF INCISION IN OPERATION FOR REPAIR OF RUPTURE OF PERINEUM THROUGH SPHINCTER ANI. For letters see p. 598.

PASSING OF SUTURES IN SAME OPERATION. For letters see p. 598. The deep sutures are to be passed nearer the skin edge.

three days for this, and the enemata should be given by means of a long tube, and with the hips elevated.

The instruments requisite are the following:—

Requisites.

- Angled scissors,
- Two pairs of artery forceps,
- Péan's forceps,
- Catgut ligatures,
- Silkworm gut or silver wire,
- Operating douche,
- Fully curved needles, large and small,
- Needle holder.

Methods. The patient is chloroformed and placed opposite a good light in the lithotomy posture. The knees are held by assistants as follows. Each stands facing the light, and places a knee of the patient under the arm-pit next to it; with the hand of the same arm, he controls the patient's foot. With his other hand, the assistant exercises tension on the nates as the operator wishes.

The stages of the operation are—(1) Forming flaps with scissors; (2) Applying the stitches.

A. R.
Simpson's
Operation.

The flaps, as made by A. R. Simpson, are shown in fig. 310. The point of the lower blade of the angled scissors is entered at *b*, pushed up to *a*, and then a clip made so as to expose tissue in line *b a*. The point is next entered at *l* on the left side, and pushed between the




FIG. 312.

DIAGRAM SHOWING RUPTURE INTO ANUS AND LINE OF TAIT'S OPERATION (*f e g d c*). Anterior vaginal wall (*b*); anus (*a*); *g* is on posterior vaginal wall.




FIG. 313.

SHAPE OF RAW SURFACE AFTER FLAPS HAVE BEEN DISSECTED UP AND DOWN; *e* and *f* show relation of stitches to skin edge in Tait's method.

vaginal and rectal mucous surfaces, *i.e.*, along the loose connective tissue between these until the point emerges at *l* on the right side. A clip is then made so as to expose tissue in the line *l S l*. Lastly, the point of the scissors is entered at *b* (right side), and *a b* clipped as already given on the left side. In this way an H-shaped figure is cut out (fig. 312). These clipped-out lines map out four flaps which are now to be raised so as to expose for union the muscular tissue lying beneath. The flaps are best raised as follows:—Lay hold of flap *S l a*

(left) at angle 1, with Péan's forceps, and raise it by clipping: do the same with flap S 1 *a* on right side. While the flap is being raised, the index or middle finger of the left hand is kept on its vaginal aspect so as to regulate its thickness. The rectal flaps S 2 *b* are then treated in the same way, the angle 2 of each being seized with the forceps. In this way a quadrilateral surface is now laid bare, with the muscular ends of the external and internal sphincters as well as the interlacings of the various muscles of the perineal body: fig. 314 will make this clear. The incisions in Tait's operation are similar (fig. 312).

The sutures can now be passed as follows:—The needle armed with silkworm gut may be passed through the skin margin, carried across either




FIG. 314.

CORONAL SECTION THROUGH ANUS (*Symington*).

r rectum; *i s* internal sphincter; *e s* external sphincter; *l a* levator ani; *v* vagina.

completely below the tissue or only above the surface at the apex of the wound, and made to emerge on the other side beyond the skin margin (fig. 311). Tait has improved on this by not including the skin margin in the suture (fig. 313). The lowest stitches should pick up and unite the edges of the external and internal sphincter (*v.* fig. 314). If the sutures when tied do not include the skin they cause no pain to the patient. The vaginal flaps need not be sutured as in fig. 311 but may be left alone. The mucous membrane of the rectal flaps may be sutured with catgut, but it is unnecessary. Bleeding,

which can be checked by a stream of very hot water (110°–120° F.) or by Péan's forceps, should have ceased before we tie these sutures. They should be left in for a week to a fortnight. Their removal is a little troublesome, as they are apt to become buried. The best way to remove them is to have the patient in the lithotomy posture, to lay hold of both ends of the suture and pull it to the one side, picking up the loop with a rake.

The advantages of this method of operating are very great. It can be done very rapidly, ensures muscular union, does not allow skin or mucous membrane to interfere with the union of muscle, and is a great improvement on the old methods. In these the union often seemed sound, but the patient had no additional control from want of muscular union.

This method is not, strictly speaking, that of one operator, but has been evolved as follows:—In 1872 John Duncan closed an artificial anus following gangrenous femoral hernia by dissecting up the mucous membrane round the orifice for more than half an inch, invaginating this dissected portion and bringing the raw surfaces together with interrupted catgut sutures: the margins of the skin were then pared and brought together by wire.

Collis, of Dublin, in 1861, in a case of vesico-vaginal fistula split the edges of the fistula instead of paring them. A. Russell Simpson applied the separation of the mucous membrane introduced by Duncan, to tear the perineum involving the anus, splitting the septum between anus and vagina and sewing similar mucous membranes to each other as well as bringing the deep raw surfaces into union. This procedure really forms vaginal and rectal flaps. Lawson Tait improved on this by the use of angled scissors, and also introduced the method of passing the sutures inside of the skin instead of through it as formerly done.

The use of scissors to form flaps is also applicable in perineum operations where the anus is not torn. According to Sænger, Stein—a Danish surgeon, and Voss—a Norwegian, have employed somewhat similar methods in complete rupture.

The continuous spiral catgut suture, now much used in Germany in such cases, has many advantages, and is preferred by most operators to the method with fishing-gut described above. It is very quickly passed, brings the surfaces well into apposition and does not require to be removed. The catgut used must be specially prepared with oil of juniper and corrosive sublimate so as to be aseptic and last eight or nine days.

Method of using continuous suture. The rectal edges are first secured by interrupted catgut stitches. The rest of the wound is then united by a single spiral or continuous suture on a well-curved needle as

follows. The vaginal flap is held well up with Péan's forceps, and the suture passed near its apex and tied. It is then carried from side to side down and back, then up towards the apex of the vaginal flap at a higher level of course, and finally down and back again. This last tier takes in the skin edge. When the last loop is being made the free end of the catgut is kept long and tension made on it and the needle until taut; the free end and needle end are then securely knotted. While this spiral suture is being applied an assistant keeps the catgut tense




FIG. 315.

CONTINUOUS CATGUT SUTURE IN ANTERIOR COLPORRHAPHY (*Pozzi*).

1. In two stages: the first has been completed and brings together the mucosa at the angles of the wound, the raw surface only in its centre; the second stage will unite the mucosa throughout.
2. In three stages. Here two have been used to contract the raw surface, while the third will close in the mucosa throughout.

by always grasping the suture near the point where it has last emerged from the skin. Fig. 315 shows the continuous suture in an anterior colporrhaphy (*v. p.* 616); its mode of application is the same, if we suppose the raw surface in it to represent the perineum.

After-treatment. The patient's food must be liquid and not too abundant. The bowels are to be confined for three days and then moved by

a small dose of castor oil every second day. Prior to the motion, the nurse must inject a large amount of oil and see that scybala if present are broken down. Unless the nurse is skilled, the operator or his assistant must attend to this. The silkworm gut stitches are removed, as already said, on the eighth to the fourteenth day.

Operation for Rupture of the Perineum, the Sphincter ani not being involved. This is described in the chapter on Prolapsus uteri.

Emmet has devised an operation with the view of restoring the grip of the fascia, forming the triangular ligament, upon the vaginal wall. A double triangular raw surface is made on the posterior vaginal walls. This is shown at fig. 316, in which one of the triangles *a, b, c*, is seized




FIG. 316.

EMMET'S OPERATION FOR RUPTURED PERINEUM (*Dudley*).¹

with tenacula. The sutures are now passed along the upper margin in loops so as to fold the edge *a b* on itself at its central point *d*, which is hooked up in a fourth tenaculum. The third figure shows this done on both sides and these sutures tied. Finally, additional sutures are passed through the edge *b c* so as to unite it with the corresponding part of the other triangular raw surface.

¹ *Pepper's System of Medicine*, Vol. iv., pp. 164, 165.—London: Sampson Low, Marston, Searle, and Rivington, 1886.

CHAPTER XLIX.

DISPLACEMENTS OF PELVIC FLOOR: PROLAPSUS UTERI: VAGINAL ENTEROCELE.

LITERATURE.

Alexander—The Treatment of Backward Displacements of the Uterus and of Prolapsus Uteri by the New Method of Shortening the Round Ligaments: London, Churchill, 1884. *Duncan, Matthews*—Papers on the Female Perineum: Churchill, London, 1879. *Emmet*—Principles and Practice of Gynecology, p. 367: Lond., 1879. *Freund, W. A.*—Ueber die Figur des normalen Lumen vaginae und über Dammplastik mit Demonstrationen: Arch. f. Gyn., Bd. vi., p. 317. *Fritsch*—Die Lageveränderungen der Gebärmutter, Billroth's Handbuch: Stuttgart, 1879. *Goodell*—Lessons in Gynecology, Lesson vii.: Philadelphia, 1879. *Hart*—The Structural Anatomy of the Female Pelvic Floor: Edinburgh, 1880. *Hegar und Kaltenbach*—Operative Gynäkologie, 1881. *Huguier*—Mémoire sur les Allongements Hypertrophiques du Col de l'Utérus: Paris, 1860. *Legendre*—De la chute de l'Utérus. *Martin*—Ueber den Scheiden und Gebärmuttervorfal: Volk. Samml., Nos. 183, 184. *Schatz*—Ueber die Zerreissungen des muskulösen Beckenbodens bei der Geburt: Archiv f. Gyn., xxii., S. 298. *Schroeder*—Die Krankheiten der weiblichen Geschlechtsorgane: Leipzig, 1879. *Schultze*—Die Pathologie und Therapie der Lageveränderungen der Gebärmutter: Berlin, 1881. *Schütz*—Medianschnitt durch das Becken einer Frau mit Scheiden- und Uterus-Vorfall: Arch. f. Gyn., Bd. xiii., S. 262. *Sims, Marion*—Uterine Surgery: London, 1865. *Skene*—Injuries to the Pelvic Floor from Parturition and Other Causes: N. Y. Med. Jour., Vol. xli., 1885. *Spiegelberg*—Ein anderer Medianschnitt durch ein Becken mit Scheiden-gebärmutter-Vorfall: Archiv f. Gyn., Bd. xiii., S. 271. *Thomas*—Diseases of Women, p. 168: Philadelphia, 1880. *Veit*—Klinische Untersuchungen über den Vorfall der Scheide und der Gebärmutter: Zeits. f. Geb. u. Gyn., Bd. i. *Winckel*—Die Pathologie der weiblichen Sexualorgane: Leipzig, 1880.

Preliminary Considerations. The subject of this chapter can only be understood in the light of an accurate knowledge of the normal structural anatomy of the pelvic floor, and a consideration of the changes it undergoes during parturition, and in the displacements to be considered. Our information on the last point leaves, however, much to be desired. The student should read over Chap. IV.

We note here that the pelvic floor is to be considered as made up of the two portions termed the "entire displaceable" and "entire fixed."

Fig. 308 shows a sagittal mesial section of the pelvis with the "entire

displaceable portion" removed and the entire fixed portion left: Pl. II., fig. 2, shows the two portions in axial coronal section.

These two portions are separated by loose connective tissue. During parturition the child is driven through the vagina, *i.e.*, through the pelvic floor, which becomes canalised or opened up through this process. If we regard this process only in sagittal mesial section as shown in Braune's plate, we see that the pubic segment is drawn up and the sacral one driven down and back, and the vagina in addition greatly distended. If considered in axial coronal section we should see the "entire displaceable portion" in part drawn up, the fœtus driven through it and thus the levatores ani and glutei muscles in the "entire fixed portion" driven out and back and the former perhaps torn (*Schatz*) or at any rate elongated, and their slope diminished. The slit in the triangular ligament through which the vagina passes is also dilated, and may be unduly so. The upward traction exercised on the "entire displaceable portion" necessarily elongates or slackens the loose connective tissue joining the two portions and is one factor in bringing about prolapsus uteri. As the result therefore of the structure of the pelvic floor, of lesions caused by parturition, and intra-abdominal pressure, we may get certain conditions, viz.,

- I. Undue yielding or bulge of the pelvic floor;
- II. Prolapse of the "entire displaceable portion" with the uterus and abdominal viscera, in part, past the "entire fixed portion"—so-called prolapsus uteri;
- III. Vaginal enterocele,—anterior and posterior.

1. *Undue yielding or bulge of the whole pelvic floor.* This is a condition to which attention has been drawn by Herman and Skene. Our knowledge of this lesion is however very defective and calls for investigation. In Chap. IV. attention has been called to the normal pelvic-floor projection. In undue bulging of the pelvic floor this is increased. Herman measures with a tape the length of the arc described by the curved skin aspect of the pelvic floor between tip of coccyx and lower margin of symphysis pubis. This average, about four inches, may be increased by straining, in virgin cases, to four and a half inches; but in cases of undue bulge, to about six or more.

Causation. This lesion is due to parturition; we are not yet in a position to give precise details, owing to the complete want of sectional and dissectional work on the pelves of women with such a prolapsed condition. *Schatz* and *Skene* have described certain conditions of laceration of the levator ani muscles, atrophy and permanent paralysis, but all has been based on clinical investigation uncorrected by anatomical examination. The subject however is important, the researches so far suggestive, and further accurate work called for.

The *Symptoms* of undue yielding are bearing-down pain with dragging in loins and hips.

The *Treatment* is the use of an abdominal belt with a perineal band.

PROLAPSUS UTERI.

DEFINITION.

A downward displacement of entire displaceable portion of pelvic floor, uterus and appendages, past entire fixed portion ; with coincident descent of small intestine.

PRELIMINARIES.

The subject of Prolapsus Uteri is a complex one, and has been in part made so by erroneous terminology.

Thus the well-known term Prolapsus Uteri has biassed many observers as to the nature of this lesion, inasmuch as they have considered some change in the uterus as initiating the prolapse. This is a natural error, and is perpetuated in most of our text-books by the writers of these considering prolapsus uteri under affections of the uterus. Prolapsus uteri is, however, considered here under Displacements of the Pelvic Floor, as it is really a hernial displacement of part of the pelvic floor in which the entire displaceable segment of the pelvic floor, uterus, and appendages are driven down by intra-abdominal pressure. There is no doubt that some change takes place in the length of the uterus as the result of the downward displacement. This change is, however, a secondary one, as will presently be explained, and does not initiate the lesion.

The student must therefore use the term prolapsus uteri not in its literal sense, but as equivalent to "sacro-pubic hernia."

Prolapsus uteri is sometimes applied to hypertrophy of the vaginal portion of the cervix. This is wrong, as this hypertrophy is a growth phenomenon.

ETIOLOGY.

The factors producing prolapsus uteri are three in number:—(1) *Deficient support by entire fixed portion* ; (2) *Deficient tone of entire displaceable segment of pelvic floor, and slackening of loose tissue round it* ; (3) *Intra-abdominal pressure*.

Deficient support by entire fixed portion. By this is meant that through parturition the sacral segment has become straightened out or deficient at its lower margin—the perineum—and that the slope of the levatores ani has been lessened or that they have been torn (Schatz). It is wrong to imagine that tear of the perineum is everything in prolapsus uteri ; the perineum may be considerably torn and yet, if the

sacral segment is still sufficiently curved and the intra-abdominal pressure not too great, there will be no prolapsus. Tear of the perineum diminishes the sacral support, and deficient sacral and levator-ani support makes the task of intra-abdominal pressure easier.

The bearing of the *second* and *third* factors is sufficiently evident. Of all the three, increased intra-abdominal pressure is the most important and is sufficient to cause prolapsus in virgins. The first and second are adjuvant.

NATURE.

Prolapsus
Uteri a
Hernia.

The uterus has nothing to do with prolapsus. It is a classical term, but a misleading one. Prolapsus uteri is really *a hernia*; and is analo-




FIG. 317.

TO SHOW THE HERNIAL NATURE OF PROLAPSUS UTERI; *a* peritoneum; *b* bladder; *c* uterus; *d* anterior vaginal wall; *e* anterior rectal wall; *f* perineum; *g* posterior vaginal wall. The dark portions are the coverings of the Hernia (after Schütz).

gous in every point to what we term a surgical hernia (such as inguinal hernia).

Thus it has (1) a sac, the peritoneum; (2) a definite road to travel along, whose boundaries are—*a*. in front, the symphysis pubis, *b*. behind, the portion of the sacral segment of the pelvic floor from anterior wall of rectum back to sacrum, *c*. side walls, viz., obturator internus and levator ani muscles; (3) definite coverings, viz., *a*. pubic segment of

pelvic floor, *b.* the uterus, *c.* posterior vaginal wall. Like all herniæ, its sac contains small intestine (fig. 317).

Huguier alleged wrongly we believe, that, by a hypertrophic elongation of the supra-vaginal portion of the cervix, the bladder and posterior vaginal wall were displaced downwards; and that many cases of alleged prolapsus uteri are really due to this. Such cases differed from prolapsus uteri in the fact that the fundus uteri and fundus of bladder are in position. Many gynecologists hold this view of Huguier, most of them modifying it somewhat. Schroeder's Handbook, Goodell's Gynecology, and Hart's Structural Anatomy may be consulted on this moot point. Huguier's Views.

SYMPTOMS AND PHYSICAL SIGNS.

The discomfort caused by the protrusion and the excoriation of the parts is the prominent symptom. The patient complains of "something coming down in front." Further, there is difficulty in micturition.

The physical signs are distinct. If the prolapsus be *incomplete*, a portion of the anterior vaginal wall has passed out at the vaginal orifice, the os uteri is equally displaced downwards, and the posterior fornix is apparently deeper from the descent of the cervix. The uterus, in addition to being low down, is usually enlarged; it lies with its axis coinciding with that part of the pelvic curve in which it is. If the prolapsus be *complete*, we find the whole anterior vaginal wall outside the vulva, the cervix extruded, and the posterior vaginal wall everted (fig. 177). The student must specially note that this description is based on *clinical* observation.

From the *study of frozen sections*, we further learn that the posterior vaginal and anterior rectal walls are separated by peritoneum driven in between them, and that the uterus with other parts has become hypertrophied through long-standing congestion, and the cervix elongated.

MECHANISM OF PROLAPSUS.

The displaced organs can be replaced—posterior vaginal wall first, then uterus, and lastly pubic segment; on the patient's straining, the mechanism of the displacement is repeated, is seen to be perfectly definite and to occur as follows.

We have first the appearance of the anterior vaginal wall, from below upwards, at the orifice. *Pari passu* with its descent, the uterus and posterior vaginal wall have come down; the cervix tracing out the pelvic curve, while the uterus becomes more and more inclined backwards, until at the vaginal orifice it lies in the vaginal axis; the posterior vaginal wall forms a pouch, the depth of half its own length, behind it. Finally, the uterus is driven outside; the cervix sweeps upwards and Mechanism on Clinical Observa-
tion.

forwards, and the posterior vaginal wall is now completely everted—its lowest part appearing last.

Appear-
ance of
Prolapsus
on Section.

On vertical section, we now find these conditions:—(1) Almost complete extrusion of the anterior or pubic part of the floor, the upper and anterior part of the bladder still behind the symphysis; (2) Complete extrusion of the uterus, which sometimes lies with the fundus below the level of the anus; (3) Rectum in position and only posterior vaginal wall down; the latter has peeled from the rectum downwards as far as the lowest inch-and-a-half (of close connection) which is elongated (fig. 317).

Explana-
tion of
Mechan-
ism.

The *explanation of this mechanism* is as follows. The displacement in prolapsus uteri is caused by intra-abdominal pressure, pushing down that part of the pelvic floor which lies in front of the anterior rectal wall, and inside the obturator internus and upper portion of the levator ani muscles. This part consists of entire displaceable portion of pelvic floor, with uterus and appendages. If we now look at a section of the pelvis such as is seen in Pl. I. (vertical mesial section) we find the posterior angle of the pubic segment is attached to the cervix uteri, and the cervix uteri to the top of the posterior vaginal wall. Thus, if intra-abdominal pressure is excessive, this part when driven down must have the following sequence of protrusion at the vaginal orifice: (a) Anterior vaginal wall from below up; (b) Cervix uteri; (c) Posterior vaginal wall from above downwards.

Our knowledge of the side relations in prolapsus is not yet complete, but from the structure of the normal pelvis, we believe that separation takes place inside the obturator internus and upper portion of the levator ani muscles (*v.* Chap. IV.).

The uterus, while it is being forced down, has the *direction of its long axis* continually altering. This is often expressed by saying that the uterus becomes more and more retroverted, as it is forced down. The real fact is, that, as the pubic segment is forced down, it is stretched—chiefly on its perineal aspect. In this way tension is made on the cervix uteri, with the effect of throwing the fundus back and making it rest on the retrojacent structures. As these have (roughly speaking) the pelvic curve, we get the uterus in this way constantly altering the lie of its axis.

The *enlargement* is not purely cervical, but affects the whole uterus, the pubic segment, and the posterior vaginal wall. This enlargement is a *consequence* of prolapsus uteri, and not a factor in its production. If we view a prolapsed uterus (with the os at the ostium vaginæ) through the pelvic brim, it can be seen that it lies, as it were, at the bottom of a valley—the sides of the valley being the broad ligaments, the bed of the valley the uterus. The parts of the uterus do not lie on the same horizontal plane, the cervix lies low. It is thus probable that the

venous supply of the uterus, having a mechanical disadvantage to its return, may have a tendency to stasis. This may lead to areolar hyperplasia at first, and, so far as our present knowledge goes, partly accounts for the increased size of the uterus in prolapsus. There is further probably a tensile elongation of the cervix produced which increases the uterine length.

SUMMARY OF DISPLACEMENT IN PROLAPSUS.

I. On clinical observation while a complete prolapsus is being reproduced, we note—

- (a) The anterior vaginal wall from below upwards passing down and out at the vaginal orifice ;
- (b) The cervix uteri appearing at the vaginal orifice ;
- (c) The posterior vaginal wall, from above down, coming last.

II. If a frozen section of a cadaver with prolapsus uteri be examined (fig. 317), we note that the pubic segment, uterus and posterior vaginal wall are displaced down and out. Fig. 317 is based on Schütz's drawing of such a frozen section. Axial coronal sections have not as yet been published, but the ureters are displaced down along with the bladder, and through being pressed on by the pubic arch may give rise to uræmia, as in a case recorded by A. E. Barker of University College, London.

III. The combined study of I. and II. shows that

The bladder and uterus are displaced down, the vagina everted or turned inside out, the small intestine coincidentally lowered in the pelvis, the displaced parts congested and hypertrophied, and the cervix uteri elongated secondarily.

DIAGNOSIS AND DIFFERENTIAL DIAGNOSIS.

The diagnosis is made by noticing the relation of the parts extruded, and by passing the sound if necessary into the bladder and uterus.

The differential diagnosis must be made from the following conditions :—

- (1.) *Hypertrophy of the vaginal portion of the cervix ;*
- (2.) *Hypertrophy of the supra-vaginal portion of cervix.*

For both of these conditions the student is referred back to page 317 (see figs. 169, 175, 176).

- (3.) *Cystocele*. Uterus is in position, and displacement is found to be due to bulging back of posterior wall of bladder.
- (4.) *Rectocele*. The finger, passed through the anus, can be pushed into the pouched rectum.
- (5.) *Inversion and polypus* (v. p. 416).

TREATMENT.

- A. Treatment by pessaries,
B. Treatment by operation.

A. *Treatment by pessaries*. In slight cases, where the anterior vaginal wall protrudes only a little, we may use an Albert Smith or Hodge




FIG. 318.

GREENHALGH'S PESSARY, with transverse bars.




FIG. 319.

RING PESSARY, with diaphragm.

pessary, with or without transverse bars at the lower part. If this fails, a ring pessary with spring inside should be tried; this instrument is




FIG. 320.

SIMPLE ELASTIC RING PESSARY, compressed between the fingers for introduction (*De Sinéty*).

useful here, inasmuch as it is shorter vertically than the Albert Smith and therefore does not project over the lower end of the shortened posterior vaginal wall. The instrument may be made of vulcanite, block tin, or india-rubber. The india-rubber forms are best, and may be provided with a perforated diaphragm, but this tends to retain discharge.

The pessary is taken in the right hand, and compressed between the finger and thumb as in fig. 320 while it is being passed through the vaginal orifice; the labia are separated with the fingers of the left hand.

If the ring instrument fail, then others may be tried. Fig. 322 shows

Zwanck's pessary, a bad form. A thin india-rubber bag distended with




FIG. 321.

RING PESSARY *in situ* (Hart).

air and provided with a stop-cock is good. In very bad cases and in old




FIG. 322.

ZWANCK'S PESSARY FOR PROLAPSUS.

women where an operation is out of the question, the patient or her

friends should be instructed how to pack the vagina with marine lint; the packing, if thorough, may remain *in situ* for a week. Some recommend pessaries which are attached externally to an abdominal belt. When there is much congestion and excoriation, rest in bed with the use of alum injections (3i to 0i) and application of boracic or zinc ointments to the raw surfaces, are indicated.

If the patient has good abdominal development, an abdominal belt will be of use; when applied, it should be fairly tight at the lower edge and slack at the upper one.

B. Treatment by operation. We must first consider the *status quo* in




FIG. 323.

LINES OF INCISION IN OPERATION FOR REPAIR OF RUPTURED PERINEUM. For letters see text.




FIG. 324.

SUTURES PASSED IN SAME OPERATION.

an advanced prolapsus. There are the following primary and secondary lesions:—

Condition
Parts in
prolapsus
peri.

Primary

Secondary

- (1) Perineal body usually torn and perineal union of levatores ani, transversi perinei, and bulbo-cavernosi, torn to a greater or less extent;
- (2) Increase of intra-abdominal pressure;
- (3) Congestion with areolar hyperplasia of uterus, pubic segment, and posterior vaginal wall; laxity of everted vagina;
- (4) Separation of anterior rectal and posterior vaginal walls and of vagina and bladder from their lateral relations, with peritoneum clothing the separated surfaces.

These secondary lesions, especially the last, are serious and incurable. In order to restore the pelvic floor to its pristine state we should require (1) to repair the perineal body and narrow the vagina; (2) to restrain increased abdominal pressure; these are possible: (3) to do away with congestion and areolar hyperplasia is probably beyond our powers, while (4) to bring about adhesion of the anterior rectal and posterior vaginal walls and to restore the lateral supports is impossible. *Prolapsus uteri is therefore a condition with serious and irremediable secondary results.*

OPERATIVE TREATMENT OF PROLAPSUS UTERI.

For operative purposes we consider prolapsus uteri as a downward




FIG. 325.

EXTERNAL GENITALS IN A MULTIPARA, WITH TEAR OF PERINEUM SHOWING LINE OF OPERATION (a c b) FOR LAWSON TAIT'S OPERATION.

and outward displacement of the entire displaceable portion of the pelvic floor past the entire fixed portion, with eversion of the vaginal walls.

The various operations may be classified as follows:—

1. Those that aim at giving a support to the prolapsed portions by repairing the lower edges of the sacral segment (Perineorrhaphy) and uniting the lower edges of the labia majora (Episioperineorrhaphy);
2. Those that aim at causing a narrowing of the vaginal walls or bringing about their partial union so that they are less easily everted (Colporrhaphy);
3. Those that combine 1 and 2;
4. The special operation which draws up the entire displaceable

portion by shortening the round ligaments of the uterus (Alexander-Adams' Operation).

Preliminary Considerations as to Operative Technique. It should be noted here that the method of rawing the surfaces has recently undergone a change. Formerly it was done with knife and forceps and the tissue removed: now scissors are almost invariably employed so as to raise flaps, thus exposing a raw surface for union without loss of tissue.

Operations
for Pro-
lapsed.

1. *Those that aim at giving a support to the prolapsed portions by repairing the lower edges of the sacral segment and uniting the lower portions of the labia majora.*

(1) *Perineorrhaphy.* This operation aims at restoring the perineal body, *i.e.*, it freshens and unites the torn surfaces. Perineorrhaphy alone




FIG. 326.

RESULT OF SAME OPERATION.

is only of use as an operation in slight cases, inasmuch as the part restored lies mainly beyond the vaginal walls and therefore in no way hinders their eversion, although it may make the vulvar opening through which they pass somewhat narrower. We describe this operation briefly as it is always combined with union of the lower portions of the labia majora (Episioperineorrhaphy) or some operation causing cicatrisation of the posterior vaginal wall (Colpoperineorrhaphy).

In the operation we chloroform patient, use douche and have knees held as described at page 598; make incision *bc* and *aba* as in fig. 323; dissect up flaps and pass stitches as in figs. 324, 326. (The incision *bc* may be omitted—*cf.* fig. 325.) After-treatment and removal of stitches as at page 600. As already said, this operation in itself is not of

permanent benefit unless it gets union of torn muscles, but it allows a pessary to be retained.

(2) *Episioperineorrhaphy*. In this operation the lower portions of the labia majora, as well as the cicatrised surfaces of perineal body, are vivified and the opposing raw surfaces united with silkworm gut sutures.

Lawson Tait operates with angled scissors as follows. He first notches the cicatrised surface mesially at the anterior portion of the perineum, the scissors being held parallel to the long axis of the patient's body. One blade is entered at right angles to this and pushed




FIG. 327.

TO SHOW VARIOUS FORMS OF RAW SURFACE MADE ON POSTERIOR VAGINAL WALL AND PERINEUM IN OPERATION FOR PROLAPSUS UTERI: 1111, Hegar's; 2222, Bischoff's; 3333, SIMON'S; 444ab Winckel's (*Winckel*.)

up in one labium majus to the base of the labium minus or beyond. The same is done on the opposite side. Thus a U-shaped incision is made (fig. 325). Silkworm or catgut stitches are passed to unite each side as follows. The suture is carried on a handled needle, through the raw surface, entering within the skin-margin and passing through the submucous tissue of the vaginal flap—on the one side. The needle is then withdrawn, and passed at a corresponding point on the other side, entering inside the skin and passing through the submucous tissue, when the thread passed on the left side is threaded into it and drawn through. This everts the raw surface on the one side and brings it into

apposition with the correspondingly everted raw surface on the other side. Three or four sutures are thus passed and tied. The operation is quickly done, but union may not be thorough or broad enough.

It is better, therefore, after having made the flap *a c b* (fig. 312) as already described, to close in the raw surface with the continuous catgut suture (fig. 315). The last tier of catgut need not include the skin edges; these may be united with a fresh catgut or silk suture.

(3) *Colpoperineorrhaphy*. This is a combination of perineal repair with a narrowing of the vagina, to be described below. Figs. 327, 328 show the various forms of raw surface that have been made. It will be noticed that they extend over the perineal region as well as upwards on the vaginal wall. They have been largely replaced by the flap operation.

2. *Those that aim at causing a narrowing of the vaginal walls or bringing about their partial union (Colporrhaphy).*




FIG. 328.

RAW SURFACE AS MADE BY MARTIN. 1 2 3 4, raw surfaces on posterior vaginal wall; II, raw surface, round introitus, upon perineum. The surfaces 1-4 are sutured separately, A to A and B to B. The surface I, II is applied to the corresponding one of opposite side, so that the English and Greek letters are in apposition respectively.

For anterior Colporrhaphy or Elytrorrhaphy the patient is placed in the lithotomy posture. The modified Sims' speculum is passed and the anterior vaginal wall is made tense with volsellæ—two lateral and one mesial, drawing down the cervix. A mesial incision is made and the mucosa dissected back and cut away until an oval raw surface is produced which is closed with continuous catgut suture (fig. 315 on p. 601). A posterior colporrhaphy may be performed in a similar way, but is usually combined with perineal repair as described above.

3. *The special operation which aims at drawing up the entire displaceable segment and uterus by shortening the round ligaments (Aran, Freund, Rivington, Alexander-Adams' Operation).*

This operation, first performed in this country by Rivington of

London and brought into prominence by Alexander of Liverpool and Adams of Glasgow, aims at shortening the round ligaments and fixing them in the inguinal canal so as to draw up and fix somewhat the displaced parts. It has been described under the operative treatment of retroversion (*v. p.* 405, and Pl. X., fig. 6).

The results from Alexander-Adams' operation are less satisfactory in prolapse than in retroversion.

We recommend in treatment—

- (1) Use of a ring in slight cases ;
- (2) Episiotomy or Colporrhaphy anterior and posterior, and amputations of the cervix in cases calling for operation.

In bad cases the uterus may be excised and a plastic perineal and vaginal operation performed afterwards.

The use of massage in prolapsus uteri will be described in the Appendix.

VAGINAL ENTEROCELE.

Of this there are two forms, *anterior* and *posterior*. Excessive intra-abdominal pressure usually displaces all of the pelvic floor that lies in




FIG. 329.

POSTERIOR VAGINAL ENTEROCELE (*Breisky*).

front of the anterior rectal wall. Occasionally, but very rarely, intestine is forced down between the posterior aspect of the bladder and upper part of anterior vaginal wall, or between the anterior rectal and posterior vaginal walls (fig. 329). We thus get a mass bulging into the vagina, but affecting only one wall; the uterus and cervix remain in position. This distinguishes it from prolapsus uteri and cervical elonga-

tion; by rectal examination, the posterior form of *enterocele* can be easily distinguished from *rectocele*.

The *causation* is not well known. In the posterior form, a deep dip of the peritoneum behind the posterior vaginal wall may have existed; but of this there is no evidence.

Treatment. If an Albert Smith or a pessary (as shown in figs. 318, 319) fail, an operation may be tried. In the posterior vaginal *enterocele*, for example, the protrusion should be replaced; a raw surface is then made on the posterior lip of the cervix and on a portion of the posterior vaginal wall about its middle; these surfaces are then stitched.

Prolapsus uteri and both forms of vaginal *enterocele* are therefore essentially the same in nature, viz., hernial. Intra-abdominal pressure usually displaces all in front of the anterior rectal wall; but may also force intestine in front of the anterior vaginal wall or behind the posterior one.

SECTION VIII.

DISTURBANCES OF THE MENSTRUAL FUNCTION.

CHAPTER L. Amenorrhœa : Menorrhagia : Dysmenorrhœa.

SECTION IX.

DISTURBANCE OF THE REPRODUCTIVE FUNCTION.

CHAPTER LI. Sterility.

CHAPTER L.

AMENORRHŒA; MENORRHAGIA; DYSMENORRHŒA.

THE three subjects to which this section is devoted are not diseases, but are symptoms of a large number of the more or less well-ascertained pathological conditions already considered. Theoretically, therefore, they should not come up for special consideration; practically, however, it is of use to the practitioner to summarise the conditions causing these symptoms, and to give some special hints as to their treatment.

AMENORRHŒA.

This means cessation of menstruation during the period between puberty and the menopause. It is normal to have Amenorrhœa during pregnancy and lactation. Amenorrhœa may be caused by the following *Local* conditions:—

Causes.

Congenital	{ Absence or incomplete development of uterus and annexa, atresia of the genital canal (with or without accumulation of the menstrual blood), state of cretinism;
Acquired	{ Superinvolution, simple atrophy of uterus, cystic ovarian disease, extensive inflammatory conditions of uterus and ovaries, morphinism.

Constitutional conditions—such as phthisis, chlorosis, prematurity of menopause—also cause amenorrhœa.

The local conditions have already been fully described under the various heads; we give here only a few hints as to the investigation of the causes of this symptom. When the patient complains of *never having menstruated* and there is no constitutional cause for the amenorrhœa, the question of examination should always be entertained; abdominal palpation and rectal examination are employed to ascertain that there is no retention from atresia. To ascertain the condition of the uterus,

a vaginal examination may be necessary. *Sudden cessation* of menstruation in a woman neither phthisical nor chlorotic is usually due to pregnancy; early sickness, mammary and other signs should be looked for. The sure sign of pregnancy in the early months is the characteristic increase in the size of the uterus, agreeing with the number of periods passed.

In cases where amenorrhœa is due to chlorosis, Bland's pills are indicated. These contain sulphate of iron and carbonate of potash made up as undernoted; as the result of the combination, the carbonate of iron is formed.

R Ferri sulphatis
Potassii carbonatis āā gr. iiss.
Mucilaginis tragacanthæ q.s.
Fiat pilula: mitte tales 96.
Sig. Three, thrice daily.

Nine pills must be taken per diem continuously for six to eight weeks, by which time a complete cure usually results.

Before the pills are given, the state of the tongue and bowels should be looked to. If the tongue is foul and the bowels constipated, we may give the following:—

R Magnesii sulphatis ȝi.
Quininæ sulphatis gr. xxiv.
Acidi sulphurici dil. ȝiij.
Aquam ad ȝvi.
Sig. Tablespoonful twice or thrice daily.

This is taken for a week. The Carlsbad salts or Friedrichshall water may be substituted. This hint as to the preliminary purgation is a good one, and is given by Milner Fothergill; if not attended to, the result will be disappointing as the iron will not be so readily absorbed by the intestinal mucous membranes.

Ringer recommends permanganate of potash. The following is a good formula:

R Potassii permanganatis.
Kaolin āā gr. ij.
Vasellini q.s.
Fiat pilula: mitte tales xxiv.
Sig. One thrice daily.

These pills should not be made with any excipient containing glycerine or with an oxidisable substance as their union would cause combustion.

Oxide of manganese (manganesii oxidum præparatum) in two grain doses thrice daily is also excellent.

MENORRHAGIA.

Menorrhagia is the term applied to excessive hæmorrhage at the menstrual periods; when the hæmorrhage is intermenstrual, it is termed metrorrhagia.

The causes of menorrhagia are the following:—

Causes.	Constitutional Hæmorrhagic diathesis, scorbutic conditions, alcoholism ;
	Local { Ovaritis, small cystic ovaries, endometritis, metritis, subinvolution, retroversion of uterus, inversion of uterus, submucous and interstitial fibroids, polypi, carcinoma uteri, sarcoma uteri, incomplete abortion.

It should not be forgotten that we may have menorrhagia in cardiac disease, and also in hepatic congestion (*Matthews Duncan, Warner*).

Women who are drunkards very often suffer from menorrhagia owing to the liver congestion. This may give the practitioner a hint as to the patient's habits, especially as women who drink always conceal the failing, and often most successfully. When called to such, there is usually found great epigastric pain on pressure, tremulous tongue, and depression of spirits, for which their excuses are quite inadequate.

Treatment. The treatment of menorrhagia is the treatment of the condition producing it. In cardiac disease we give digitalis; and in hepatic disease we may try chloride of ammonium, euonymin or iridin.

R	Ammonii chloridi	℥iij.
	Aquæ	℥vj.
	<i>Sig.</i> Tablespoonful thrice daily.	
R	Euonymin	
	vel	
	Iridin	gr. ii.
	Pil. aloes et ferri	q.s.
	Fiat pilula : mitte tales xij.	
	<i>Sig.</i> One at night.	

In cases where there is menorrhagia due to a simple congested condition or to a flabby state of the uterine muscle, we may give the following at the menstrual periods:—

R	Ergotini	gr. iv.
	Argenti oxidi	gr. $\frac{1}{4}$
	Micæ panis	q.s.
	Fiat pilula : mitte tales xij.	
	<i>Sig.</i> One thrice daily as directed.	

Note that it is well not to write "at the menstrual period" on the prescription, but to put "as directed." When the practitioner is consulted as to menorrhagia in unmarried women or young girls, he should first try the ergotin and oxide of silver pill. If this fail and the case be urgent, he should request a local examination. If this be declined, the responsibility rests with the patient.

R Extracti ergotæ liquidi ʒij.
Sig. Thirty drops as directed
or

R Ergotini gr. iv.
Fiat suppositorium : mitte tales xij.
Sig. As directed.

Inform the patient that two suppositories are to be passed into the rectum each morning after the bowels move.

In some cases the hypodermic injection is required (*v. p.* 189).

DYSMENORRHŒA.

LITERATURE. *Champneys, F. H.*—On Painful Menstruation: London, H. K. Lewis, 1891. *Croom, J. Halliday*—Disorders of Menstruation: Clifford Allbutt and Playfair's System of Gynecology, London, 1896. *Duncan, Matthews*—Clinical Lectures: London, 1886, p. 141. *Goodell*—Lessons in Gynecology: Philadelphia, 1879. *Gusserow*—Menstruation and Dysmenorrhœa: Germ. Clin. Lect., New Syd. Soc. Tr., 1877. *Herman, G. E.*—On the Relation between Backward Displacements of the Uterus and Painful Menstruation: Lond. Obst. Trans., 1882. *Solowieff*—Decidua menstrualis: Archiv f. Gyn., Bd. ii., S. 66. *Schroeder*—Die Krankheiten der weiblichen Geschlechtsorgane: Leipzig, 1887. *Simpson, Sir J. Y.*—Diseases of Women, p. 225: Edin., 1872. *Williams, John*—Pathology and Treatment of Membranous Dysmenorrhœa: Lond. Obst. Tr., 1877.

Dysmenorrhœa may be defined as the occurrence of pain before, during, or after the menstrual period.

The pain of dysmenorrhœa varies greatly in intensity. It may be so severe as to render the sufferer a miserable invalid, it may interfere with her work more or less, or it may cause only marked uneasiness. It is always advisable in cases of dysmenorrhœa to ascertain how much the pain interferes with the patient's occupation or whether it confines her to bed. Note also when the pain occurs—prior to, during, or after the blood-flow; in the purely spasmodic form, it is during the flow.

In order to treat dysmenorrhœa intelligently, we must endeavour to ascertain its cause and try to make out how this condition brings about the pain. We know nothing at all as to the real cause of dysmenorrhœa. We know that in many instances it is associated with certain pathological conditions, but how these actually cause the pain is as yet disputed.

Some facts as to menstruation help us in understanding dysmenorrhœa. The uterus is an erectile organ (*p.* 74), and as the decidua menstrualis is five or six times thicker than the uterine mucous membrane, it is evident that metritis or pathological antelexion when present will

hinder the erection and expansion of the uterus, and cause intense pain analogous to the chordee of the penis in gonorrhœa.

In normal menstruation, a fluid made up of blood and epithelial débris escapes from the uterus. Probably, it does not drain away by mere capillary action, but is expelled by uterine contractions. There is no absolute proof of this, but it is a fair deduction from anatomical facts. If a patient be examined while menstruating, we may feel an arching or slight tension of the fornices indicative probably of uterine action.

Dysmenorrhœa is usually divided into certain forms. It is to be regretted that this has been done, because there have not been collected pathological facts sufficient to warrant a classification. The forms usually given are the following :—

Forms
usually
given.

1. Dysmenorrhœa associated with certain diatheses, such as the gouty and rheumatic ;
2. Spasmodic dysmenorrhœa ;
3. Membranous dysmenorrhœa ;
4. Dysmenorrhœa associated with inflammatory conditions of the uterus, ovary, peritoneum or cellular tissue ;
5. Ovarian dysmenorrhœa.

The last term is applied to certain cases which were supposed to be specially connected with the ovaries, and which could not be classified under the preceding heads. The term is a most unfortunate one. It assumes a cause for dysmenorrhœa which is not, as yet, demonstrated ; and, instead of pathological facts or a confession of our ignorance of them, gives us what we have too much of already—erroneous terminology.

Practical
Varieties.

So far as our present knowledge goes we can speak of four varieties :—

1. Spasmodic dysmenorrhœa ;
2. Congestive dysmenorrhœa ;
3. Membranous dysmenorrhœa ;
4. Dysmenorrhœa associated with mal-development of the sexual organs, pyosalpinx, fibromyoma uteri, rheumatic diathesis, and some other unknown causes.

The Erec-
tion and
Expansion
of the
Uterus
hindered.

1 and 2. *Spasmodic and Congestive dysmenorrhœa.* Of these the most frequent cause is pathological ante flexion, *i.e.*, ante flexion of the uterus produced by inflammation in the utero-sacral ligaments with cicatrization. The pathology, diagnosis and treatment of this affection is given at pp. 374-380. We only remark here that it is a very serious lesion owing to its inflammatory etiology. From the flexion produced, we get spasmodic uterine contraction accompanied with very great pain and expulsion of clots. Two theories of dysmenorrhœa have been already

explained (p. 377). Those who hold the purely mechanical theory seem to forget that fluid blood passes easily through a capillary. Does any one believe that the lumen at the flexion is less than that of a capillary?

Spasmodic contraction of the os internum and constriction of the cervical canal are also advanced as causes.

3. *Membranous dysmenorrhœa*. In this condition, the superficial layer of the mucous membrane is cast off as a triangular sac or in shreds of a more or less firm consistence (figs. 330, 331). This may result from the occurrence of hæmorrhage in the deeper layers of the mucous membrane; and then we can understand that, according to the depth, we




FIG. 330.

SKETCH OF A DYSMENORRHŒAL MEMBRANE AS SEEN UNDER WATER (*Sir J. Y Simpson*).

have present no part of the glands or only their cæcal extremities (*Solowieff and Gusserow*). Microscopically, there is excess of round cells and fibrillated tissue in the membrane.

Sir J. Williams, who has written ably on this subject, believes that, owing to an excess of fibrous tissue in the walls of the uterus, the mucous membrane is expelled in coherent shreds. This excess of fibrous tissue is due to defective evolution, sub-involution, or metritis. The membrane is, further, never a plastic exudation. *It is of the greatest importance to remember that it is not a product of conception, and should not be mistaken for an early abortion.*

4. *Dysmenorrhœa from other causes, as defective development of uterus, pyosalpinx, etc.* Many of these conditions are now being elucidated by abdominal section undertaken for Battey's and for Tait's operation.

TREATMENT.

Cautions
as to
Treatment.

At the outset we are met with a difficulty. As we are usually consulted for *Dysmenorrhœa* in unmarried women, the question of the propriety of a pelvic examination comes up. As Duncan has said—"No rules that I can give you will make up for want of good sense and good feeling on your own part, but I shall give you some hints. The first is that you should, as a rule, not resort to this treatment (by bougies) in an unmarried young woman without the concurrence of three parties—




FIG. 331.

A DYSMENORRHEAL MEMBRANE LAID OPEN (*Coste*).

firstly, your own approval; secondly, that of the mother or guardian of the patient; and, thirdly, that of the patient herself. All of these should be quite aware of the circumstances, and of what it is proposed to do."

Nothing can be more reprehensible than the vaginal examination of unmarried women for trifling ailments. When the *Dysmenorrhœa* is slight, make no examination but order some such mixture as the following.

R Spiritus chloroformi,
 Spiritus ammoniæ aromatici, āā ̄ss.
 Liquoris ammoniæ acetatis ̄i
Sig. Teaspoonful in a wine-glassful of hot water occasionally.

Viburnum prunifolium in capsules containing thirty drops can also be given (thrice daily).

Order a hot hip bath, or the feet to be put in mustard and water. On no account whatsoever allow alcohol in any form to be given. If the mother has been giving whisky and water or gin and water, at once point out the risk the patient is running. Do not give morphia, or other opiate, unless driven to it; always give it yourself and hypodermically, never by the mouth or rectum, and give no prescription for it.

When the Dysmenorrhœa is urgent, then an examination should be advised; the index finger well oiled can usually pass in without much pain.

If pathological ante flexion is found, note the amount of inflammatory disturbance, the degree of flexion, and the implication or non-implication of the tubes and ovaries. Begin by ordering blisters to the iliac regions, bromide of potassium, the glycerine plug, and the hot vaginal douche. See that the bowels are regulated, and soft motions secured by the use of liquorice powder (*Pulv. glycyrrhizæ co.*) and occasional enemata, and that no tight lacing is allowed. Chlorotic patients should be put on Bland's pills and digitalis; and change of air, when requisite, ordered. Note the effect of this for some periods; and then, if unrelieved, pass the sound or graduated bougies or use uterine dilator. This course benefits the Dysmenorrhœa, and it is safer than the use of stem pessaries; the dilatation by bougies seems to act like the stretching of the sphincter ani in fissure of the anus and often gives brilliant results.

Patients with neurasthenia often suffer severely at the menstrual periods. Local treatment is contra-indicated, as the dysmenorrhœa often passes off while the general condition is improving.

If the Dysmenorrhœa is membranous, treatment is of little service. The following prescriptions may be tried.

R Liquoris arsenicalis ℥ij.

Sig. Three drops in water thrice daily after food.

R Liqueur arsenii et hydrargyri iodidi (Donovan's solution) ʒij.

Sig. Five drops in water thrice daily after food.

The action may be analogous to that of arsenic in psoriasis.

Treat any endocervicitis or stenosis of cervix present. The prognosis is unfavourable as to cure. The patients are not necessarily sterile.

In the fourth group of cases, Battey's operation has not given the results anticipated. We have not, as yet, however, facts warranting any dogmatic utterance. Where the ovaries are developed but not the uterus, with serious menstrual molimina resulting in consequence,

Batley's operation is undoubtedly indicated. In cases of pyosalpinx, removal of tubes and ovaries by abdominal section gives good results (*v. p.* 236).

Where any diathesis (rheumatic or gouty) is supposed to influence the Dysmenorrhœa, guaiac, colchicum and such specific drugs may be given.

CHAPTER LI.

STERILITY.

LITERATURE.

Ashby—Influence of Minor Forms of Ovarian and Tubal Disease in the Causation of Sterility : Am. Jour. Obst., 1894, Vol. xxx., p. 161. *Bumm*—Zur Bebandlung der Sterilität bei der Frau : Centralb. f. Gyn., 1893, S. 981. *Duncan, J. Matthews*—Fecundity, Fertility, Sterility and allied topics : Edinburgh, A. & C. Black, 1866. On Sterility in Women : J. & A. Churchill, 1884. *v. Grünewaldt*—Ueber die Sterilität geschlechtskranker Frauen : Archiv f. Gyn., Bd. viii., S. 414. *Herman*—On the Relation between Backward Displacement of the Uterus and Sterility and Abortion : Lond. Obst. Trans., 1891, Vol. xxxiii., p. 499. *Kehrer*—Zur Sterilitätslehre : Beiträge zur klinischen und experimentellen Geburtskunde und Gynäkologie, Bd. ii., S. 76. *Müller*—Die Sterilität der Ehe : Billroth u. Luecke's Handbuch der Frauenkrankheiten : Stuttgart, 1885, S. 297. *Palmer*—Sterility : Amer. Jour. Obst., 1894, i., p. 832. *Seeligmann*—Ueber Sterilitas Matrimonii : Berlin Klin. Woch., 1891, No. 41. *Sims, Marion*—Uterine Surgery : London, 1865. *Simpson, Sir J. Y.*—Obstetrics : Edin., A. & C. Black, 1871, p. 830. *Whitehead*—On the causes and treatment of abortion and sterility : London, 1847.

THE reproductive function is the the most complex and subtle of all the functions of life. If we know little about the simpler function of menstruation so that there is room for great difference of opinion with regard to it, we know still less of the function of reproduction. Of its physiology, we know only that it requires the presence of ova and spermatozoa ; of the constitutional influences affecting the vitality of these two and the conditions favourable for their conjugation, even of the place where this occurs, nothing is known. Nor have we yet data for studying the general laws of fertility for the human female. Much has been done by Darwin and others to elucidate these for plants ; little is known of them for animals, and almost nothing for the human species.

Of the disturbances of the reproductive function, sterility belongs to Gynecology ; abortion, retroflexion of the gravid uterus and extra-uterine gestation belong more properly to Obstetrics.

No simple and yet complete definition of sterility can be given. The word has a quite different meaning as we use it relatively or absolutely. As the opposite of fertility, it includes cases in which a child is not born till many years after marriage or the number of children is comparatively few ; further, inasmuch as the reproductive function covers gestation as

Distinction
between
Absolute
and
Relative
Sterility.

well as the birth of a viable child, sterility includes all cases of intra-uterine disease and death of the embryo or foetus, resulting in abortion, premature labour, or the birth of a non-viable child. None of these cases are absolutely sterile, the sterility is *relative*. The term also necessarily covers all cases in which under circumstances favourable to conception, this either has not occurred at all or the product has not gone the length of even an early abortion. Here the sterility is *absolute*. This raises the question as to when sterility is relative, and when absolute. What is the standard of fertility by which we decide that a woman is relatively sterile and measure the degree of that sterility? When can we say that a patient is absolutely sterile?

Relative
Sterility.

Relative Sterility. At first sight, we should be inclined to regard the period of child-bearing as co-extensive with the period of menstruation. But it is not so. The period of fertility is not co-terminous with the period of menstrual activity: it begins later and ends earlier, its total duration being about fifteen years, during which time births take place about every eighteen or twenty months. Its commencement is determined by the year of marriage, in this country on an average the twenty-fifth year, the first child being born in most cases twenty months after marriage. It ceases usually about thirty-eight, some years before the menopause. Thus, as Whitehead puts it, there is a period of quiescence in the function of reproduction both at the commencement and at the termination of menstruation (*Matthews Duncan*).

Taking the foregoing considerations as giving us a standard of fertility, we learn that relative sterility may show itself in such various ways as these,—not having the first child within twenty months after marriage, having children at intervals of longer than twenty months, ceasing to have children within fifteen years after marriage. In applying these considerations to an individual case, however, we must of course take into account the age of the patient. There seems also to be great variation in the productive power of different individuals. One patient has many children without injury to health, while in another the birth of one child exhausts the reproductive function. Sir James Simpson found that among British peers unproductive marriages are relatively more common (1 in 6, instead of 1 in 10). As the result of relative sterility we find that the number of children to a marriage in Britain is 5.2 or one-half of what it would be if all the conditions favourable to reproduction were fulfilled.

Absolute
Sterility.

Absolute Sterility. The interval between marriage and the birth of the first child averages twenty months, and any protraction of this interval means a degree of sterility; but we cannot speak of absolute sterility until several years of married life have passed without even an abortion. Matthews Duncan found in his statistics of the births in Edinburgh and Glasgow for the year 1855, an average interval of

seventeen months to the first child—two-thirds being born before the end of the second year, and only one-twenty-fourth after the fourth year. Hence, he concludes that there is no ground for the assumption of persistent sterility until the fourth year of married life has been entered upon.

Of the number of absolutely sterile marriages in Britain we have no data. The statistics of Sir J. Y. Simpson, based on the reports of the population of Grangemouth and Bathgate which give the number of sterile marriages as 1 in 10, include abortions and all other cases in which a child would not be registered, so that they cannot be relied upon for data regarding absolute sterility. Seeligmann gives the proportion as 23 in 200, and would ascribe one-half of these to gonorrhœa in the male.

The *Etiology* of sterility is too wide a subject to be exhaustively discussed here. We can only indicate what the causes are, and point out the necessity of taking a broad view of this question. Etiology of Sterility.

Amongst *general influences*, we note first of all the effect of *temperature* and *climate*, and of marriage between *near relatives*. Under *want of sexual agreement* have been placed many cases which have not been explained otherwise (such as the classical one of Napoleon and Josephine). *Age* has an undoubted influence; the period of nubility is from the age of twenty to twenty-five, and marriages before or after this period are less fertile. The influence of *disturbed nutrition* is seen in the association of sterility with obesity; it seems that the taking-on of fat is at the expense of the reproductive function, perhaps through interference with ovulation. *Chlorotic* patients are also sometimes sterile. The association of *Dysmenorrhœa* with sterility has been already referred to (pp. 300 and 378) and is a matter of everyday observation. Matthews Duncan found spasmodic dysmenorrhœa in 47·9 (159 out of 332) of his cases of sterility; while Marion Sims found it in 51·6 p. c. (129 out of 250) of his. Further, these conditions disappear together under treatment, and spasmodic dysmenorrhœa is a rare condition in fertile women.

As to *local causes*, we note that sterility is found associated with the following conditions already described:—vaginismus, p. 569; hypertrophied cervix, p. 311; conical cervix with pin-hole os, p. 297; cervical catarrh, p. 336; antelexion, p. 378; retroflexion (more rarely¹), p. 388; endometritis, p. 352; ovaritis, p. 241; pelvic peritonitis, p. 187. The last three are probably the most important. Taking the function of reproduction instead of the various organs as the standpoint from which to regard sterility, we find that this function may be divided into three processes—insemination, impregnation of the ovum or conception, and gestation. A certain number of cases of sterility are due to defect in

¹ Herman, *op. cit.*

insemination (e.g. all cases of dyspareunia); but the most important group of cases coming under this head are those of absence or deficient vitality of the spermatozoa. As we are dealing here only with sterility in the female, this last cause of sterility is beyond our subject; but it is important to remember that Gross's investigations into male sterility show that it is probably the cause in every sixth case which comes before us. As to the relative importance of *conception* and *gestation*, the investigations of v. Grünewaldt show that interference with the latter is a much more important factor in sterility than is generally supposed. Investigating 500 cases of sterility from the standpoint of the influence that the condition of the uterine tissue has on gestation, he comes to the following conclusion:—Conception forms only one link in the chain of processes involved in the fertility of marriage, and is of slight importance compared with the great number of vital processes implied in gestation; the point of greatest importance in the fertility of woman is her capability of carrying a fertilised ovum, which depends to a great extent on the integrity of the uterine tissue.

Kleinwächter¹ met with one-child sterility in 8·32 p. c. of his cases. The age at which the women married seemed to have nothing to do with it. He finds that the causes are the same as in the case of absolute sterility (apart from congenital malformations), viz. :—

Inflammation after puerperium	17·77 p. c.
„ not „	„	.	.	.	12·22 „
Endometritis	17·77 „
Uterine displacements	12·22 „
„ neoplasms	8·88 „
Constitutional conditions	7·77 „
Male impotence	7·77 „
Uterine atrophy	5·55 „
Ovarian neoplasms	3·33 „
Unknown causes	6·66 „

Treatment. In the *treatment* of sterility, we must take a broad view of the etiology and not allow local conditions to influence us unduly. Attention to the general health, and patient waiting until at least three years of married life have passed is all that is required in the large proportion of cases. Entire cessation of intercourse for several months should be recommended, and can be secured by change of air to some watering-place at home or abroad, according to the patient's means. Where coitus is impossible or painful (as in cases of atresia and vaginismus) operative interference is called for immediately, and such cases offer the most satisfactory results in treatment (see p. 570). In estimating the importance of operations on the cervix (p. 301), we must keep in view the rarity of this indication for treatment and the uncertainty that an operation by dilatation or division will be beneficial. Whether the sterility be due to the rigid condition of the cervix or the smallness of the os externum,

¹ *Centralb. f. Gyn.*, xii., 287.

such cases form only 4 p. c. (*Müller*) or 8 p. c. (*Kehrer*) of the total number of women who seek advice for sterility. In other words, taking *Müller's* statistics the chances are 24 to 1 that the cause of sterility must be sought elsewhere than in the cervix.

For cases of stenosis, of which sterility is the chief feature, *Pozzi*,¹ in addition to splitting the cervix, excises a prismatic piece of tissue from the raw surfaces on each lip, and stitches cervical mucous membrane to vaginal over each of the four surfaces. The cicatrisation which follows simple division (p. 303) is thus prevented. Massage and electricity (the negative pole being intra-uterine) have recently been advocated by *Palmer*, *Bumm*, and *Seeligmann*.

¹ Nouvelle opération applicable à la sténose congénitale du col de l'utérus (*Bull. et Mém. de la Soc. de Chir.*, 1893, t. xix., p. 93 et *Ann. de Gyn.*, 1893, t. xli., p. 407)

THE JOURNAL OF

JOHN R. KELLY, M.D.

OF THE
UNIVERSITY OF CHICAGO
AND
OF THE
HARVARD MEDICAL SCHOOL

CHICAGO, ILL.

S E C T I O N X.

AFFECTIONS OF BLADDER AND RECTUM.

- CHAPTER LII. The Bladder: Anatomy, Physiology, and Methods of Examination.
- „ LIII. Affections of the Urethra and Bladder.
- „ LIV. Vesico-Vaginal Fistula.
- „ LV. The Rectum: Coccygodynia.

A P P E N D I X.

- Abdominal Section: Anterior and Posterior Colpotomy:
Vaginal Hysterectomy.
- Electricity in Gynecology.
- Systematic Treatment of Nerve Prostration.
- Hysteria.
- Massage.
- Case-Taking: Classification of Diseases of Women.
- Sources of Gynecological Literature.

CHAPTER LII.

THE BLADDER: ANATOMY, PHYSIOLOGY AND METHODS OF EXAMINATION.

LITERATURE.

Barlow—Beiträge zur Aetiologie, Prophylaxe und Therapie der Cystitis: Arch. für Dermatologie und Syphilis, xxv., 1893. *Burckhardt*—Endoskopie und endoskopische Therapie: Tübingen, Laupp'schen Buchhandlung. *Chiene*—Bladder Drainage: Ed. Med. Jour., 1880. *Croom, J. H.*—On Retention of Urine in the Female: Ed. Med. Jour., April and May, 1878. *Fenwick, E. H.*—The Electric Illumination of the Bladder and Urethra: London, Churchill, 1888. *Foulis*—An Antiseptic Catheter for washing out the Bladder: Brit. Med. Jour., Jan. 30, 1886. *Fritsch*—Die Krankheiten der weiblichen Blase: Handbuch der Gynäk. bearbeitet von J. Veit., Bd. ii., Wiesbaden, Bergmann, 1897. *Gulland, G. L.*—On the Treatment of Bacteriuria by the internal administration of Drugs: Edin. Hosp. Rep., Vol. iv., 1896, p. 294. *Hart*—Physics of Rectum and Bladder: Ed. Obst. Trans., 1882. *Kelly, H. A.*—The direct Examination of the Female Bladder with Elevated Pelvis, etc.: Amer. Jour. Obstet., Vol. xxix., p. 1. My work on the Diseases of the Urinary Tract in Women: Amer. Jour. Obstet., 1896, Vol. xxxiii., p. 399. (This gives résumé of literature.) *Nitze*—Lehrbuch der Kystoskopie: Wiesbaden, 1889. *Noeggerath*—The Vesico-vaginal and Vesico-rectal Touch: Am. J. of Obstet., viii., 135. *Ogston*—Ed. Med. Jour., 1878. *Pawlik*—Ueber die Harnleitersondirung beim Weibe: Archiv f. klinische Chirurgie, Bd. xxxvi., Hft. 2. *Pawlik v. Kelly*: Amer. Jour. Obstet., 1896, Vol. xxxiv., p. 253. *Power*—Physiology of Micturition: The Practitioner, 1875. *Rubeska*—A Criticism on Professor Howard A. Kelly and his Discoveries in the Domain of Urinary Diseases: Amer. Jour. Obstet., 1896, Vol. xxxiii. *Sänger*—Ueber Tastung der Harnleiter beim Weibe: Archiv f. Gyn., Bd. xxviii., S. 54. *Skene*—Diseases of the Bladder and Urethra in Women: W. Wood & Co., New York, 1878. *Syme, James*—Observations in Clinical Surgery: Edin., Edmonston & Douglas, 1862. Second Edition. *Viertel*—Physikalische Untersuchungsmethoden der Blase: Veit's Handbuch, Bd. ii., Wiesbaden, Bergmann, 1897. *Wallace, D.*—The Electric Cystoscope: Edin. Med. Jour., 1890, pp. 712, 860. Also Cystoscopy, notes from an experience of upwards of Fifty Cases: *Ibid.*, 1891, p. 324. *Winckel*—Die Krankheiten der weiblichen Harnröhre und Blase: Billroth's Handbuch, Stuttgart, 1886.

DISEASES of the bladder are of the greatest importance as they are not only very painful, but, for a reason to be given shortly, very intractable. In a Manual of the present scope, a full consideration of vesical disease is impossible; we therefore give a mere sketch, and refer the practitioner for details to Veit's handbook, or to Skene's or Winckel's monographs.

ANATOMY AND PHYSIOLOGY.

For the anatomy, the student is referred to pp. 33 to 40. We should here only point out that the female bladder, owing to its greater breadth transversely at the base (*v.* fig. 346), is relatively more capacious than that of the male. Physiology
of Urina-
tion.

Urination.—The mechanism of the storage and expulsion of urine from the bladder is full of interest, both from a theoretical and a practical point of view. The urine trickles along the ureters, a result partly due to blood pressure and partly to the peristaltic action of the ureters themselves. It thus reaches the bladder, at this stage an empty flaccid sac with its upper half fitting into the lower calyx-like portion. Gradually




FIG. 332.

BLADDER IN SYSTOLE (*Braune*).

the bladder distends, until at last the activity of the motor centre (whose constant action keeps the urethral muscles contracted) is reflexly inhibited, and the urine is expelled by the muscular contraction of the bladder and intra-abdominal pressure. The bladder is now contracted and, on section, has the shape seen at fig. 332—its shape in systole. The bladder then relaxes, *i.e.*, becomes flaccid—its diastole, and once more the urine trickles into it (fig. 25).

The bladder therefore has, like the heart, its systole and diastole. A knowledge of this is important practically. It explains the intractability of inflammatory conditions of the bladder, since the bladder when inflamed does not get—what every inflamed organ requires—rest.

Composi-
tion of
Urine.

The average amounts of the several urinary constituents passed in twenty-four hours, as given by Parkes, are the following:—

Water	1500·000 Grms.
Total solids	72·000
Urea	33·180
Uric acid	·555
Hippuric acid	·400
Kreatinin	·910
Pigment, etc.	10·000
Sulphuric acid	2·012
Phosphoric acid	3·164
Chlorine	7·000
Ammonia	·700
Potassium	2·500
Sodium	11·090
Calcium	·260
Magnesium	·207

Urine also contains various epithelial scales, a little mucus, nitrogen and carbonic-acid gases.

The reaction is acid, and the specific gravity is 1020.

METHODS OF EXPLORING THE URETHRA.

The urethra is explored by sound, finger, and speculum in the same way as the bladder. We need not therefore go into detail in these, but refer the student to methods of exploring the bladder.

We may remark, however, that the exploration by finger, sound, or speculum is not very satisfactory in the case of the urethra, as polypi become flattened against the urethral wall by finger or speculum and are thus overlooked. In such cases the button-hole operation of Emmet is useful and is performed as follows.

The patient is put in the lithotomy posture and a sound of calibre sufficient to stretch the urethra, passed. The object of the operation is to incise the urethra vertically and mesially but not to touch the meatus urinarius or neck of the bladder. The urethra is $1\frac{3}{8}$ inches long, and therefore an incision of the vaginal tissues over the urethra $\frac{3}{4}$ of an inch in length will avoid the urethral orifice and neck of bladder. The vaginal tissue is caught up with a tenaculum and divided down to its canal. The scissors are now used to extend this up towards the neck of the bladder and down towards the urethral orifice. The incision in the vaginal mucous membrane should be one-third longer than that into the urethral canal, and the extra length should be at the bladder end.

No incontinence of urine is produced if the neck of the bladder be uninjured.

Through this incision polypi can be detected and removed, prolapse of the urethral mucous membrane can be excised, and medicaments applied.

Should the incision be made merely for temporary purposes it can be closed by silver stitches including the mucous membrane of the urethra. When the operator wishes to make a urethro-vaginal fistula for purposes of treatment, he unites the edges of the mucous membrane of the vagina to the corresponding edge of the urethral mucous membrane




FIG. 333.

EMMET'S BUTTON-HOLE OPERATION ON THE URETHRA: the patient is supposed to be on her side and Sims' Speculum passed (*Emmet*).

by means of catgut or silk (Button-hole operation—fig. 333). This fistula can be closed when necessary in the ordinary way.

For dilatation by Simon's specula, see page 641.

METHODS OF EXPLORING THE BLADDER.

A. *By Catheter and Sound.*

Preliminaries. No instrumental investigation of the urethra and bladder is to be lightly undertaken. There is great risk that even with careful antiseptic precautions any instrument if used frequently may carry in pyogenic organisms from the urethra, and set up a cystitis which may ultimately pass to the pelvis of the kidney, or to the kidney itself and the connective tissue adjacent. In the urethra, organisms are said to be invariably present, "often pathogenic and especially pyogenic" (Gulland).

The catheter is passed for the purpose of drawing off the urine, while the sound is usually employed for diagnostic purposes—ascertaining the state of the mucous membrane, the presence of stone or other pathological conditions.

Method of passing the catheter. The instrument usually employed for this purpose is a male gum-elastic catheter, No. 8 or 10, but a silver instrument is better, the great advantage of metal being that it can be boiled prior to use. Battey recommends a long rubber catheter as a very useful instrument. The gum-elastic catheter must first be thoroughly washed with carbolic lotion (1-20), or corrosive sublimate (1-2000), and then its end dipped in glycerine and corrosive sublimate (1-2000). Cleanliness in using the catheter is of the very highest importance, as cystitis, renal disease, and even pyæmia may be caused in cases where the urine has been rendered putrid by the catheter. It is of great importance therefore for the nurse to cleanse the external parts carefully first, and to boil the catheter employed.

The patient lies on the left side square across the couch, with the hips at the edge and the knees drawn up. The pulp of the index finger of the left hand is passed over the base of the perineal body and onwards until it touches the vestibule. It should then be carried a little backwards until we feel the meatus at the base of the smooth vestibule and in the middle line. The catheter is passed with the right hand; the index of the left hand feels, through the anterior vaginal wall, that it passes into the urethra. After the last drop of urine has been expelled, the catheter is withdrawn and the finger held over its proximal end so as to retain the fluid remaining in the catheter until it can be poured into a receptacle.

The catheter may also be passed with the patient lying on the back; the index of the right hand is carried under the drawn-up right thigh to feel the meatus, and the catheter is passed between the thighs with the left. It is best for the nurse to expose the parts.

Battey's catheter, or a silver catheter with long rubber tube, is very convenient, as from its length it reaches to the floor and can be withdrawn without any precaution as to spilling. Further, it is easily cleaned; to do this it is coiled up in a bowl of 1-20 carbolic lotion, and then when one end is brought over the edge it empties by syphon action. The indications for the catheter are the various causes of retention of urine (*v. p.* 656); at present we only remark that it should never be passed unless necessary, and that the greatest care should be taken not to introduce septic matter. Foulis has recommended a special apparatus for washing out the bladder which may be used for drawing off the urine also.

B. *Digital and Specular Exploration of the Bladder.*

Owing to the large amount of muscular and elastic tissue in the urethra, it can be stretched to an extent that permits of digital and specular examination of the urethral and vesical lining membrane.

It is not generally known that as early as 1861 Syme of Edinburgh wrote as follows, in regard to the removal of an encrusted hairpin from the female bladder,—“I dilated the urethra by introducing a series of bougies gradually increased in size until the point of my finger was admitted into the neck of the bladder, where feeling the tense resisting fibres situated there, I made a very slight incision. . . . The finger then readily entered the bladder. . . . The patient suffered no inconvenience, and in the course of a few days, having completely regained the power of retention, was dismissed.”

Digital examination. With the patient lying in the lithotomy position and under chloroform, the tip of the little finger is placed against the meatus and by a rotary motion passed through it in the direction of the urethral axis. The meatus is the most resistant portion of the urethra; therefore, to aid in its dilatation, some recommend to notch it with radiating nicks. This is unnecessary (*A. R. Simpson*). By steady pressure, the little finger is first pushed in and then the index one substituted. Hegar's dilators for the cervix are of great use here also. For exploratory purposes this is sufficient; to complete the examination, however, the bimanual should be performed as shown at fig. 72. This is aided by the middle finger in the vagina, and is therefore termed the vesico-vaginal bimanual. Instead of chloroform, cocaine may be injected locally.

The presence of stone or of tumours, the state of the mucous membrane of the bladder, the nature of obscure bodies in front of the uterus can all be thoroughly ascertained; vesico-vaginal fistulæ can be examined if the vagina has been obliterated; intestino-vesical fistulæ can be detected; calculi, impacted in the vesical portion of the ureters, can be removed; fissures of the neck of the bladder can be stretched; Winckel adds to these that we can open a hæmatometra through the bladder, when its evacuation between the bladder and rectum is impossible—a very rare indication. The Fallopian tubes can be felt with the finger in the bladder (*Noeggerath*); and, in one special instance, Croom proved by this method that the sound had perforated the walls of the thin superinvolved uterus and had not passed along the Fallopian tube.

Simon's methods of specular dilatation of urethra. Simon of Heidelberg drew special attention to the dilatation of the urethra by his specula as a means of treatment. The object is to dilate the urethra sufficiently to allow of the passage of calculi, crushed or uncrushed. By it we also destroy temporarily the sphincteric action of the urethra and thus cause incontinence of urine; this allows to the inflamed mucous membrane, now undisturbed by the frequent muscular contractions which before were necessary to expel the urine, the rest it needs. The difficulty of Simon's method is the risk of causing, by over-stretching, permanent incontinence of urine—a condition as yet incurable.

Simon's specula are shown at fig. 334, and the various sizes at fig. 335. The specula are provided with bulbous plugs, to be used while

they are being introduced and afterwards withdrawn. Simon estimated the limit of safe dilatability for the female urethra at various ages as follows:—*Adults*, 6.6-25 cm. ($2\frac{6}{16}$ - $2\frac{7}{16}$ in.) in circumference, or 1.9-2 cm. ($\frac{3}{4}$ in.) in diameter; *young women* (of 15-20 years), 5.6-6.3 cm. in circumference, or 1.8-2 cm. ($\frac{3}{4}$ in.) in diameter; *girls* (of 11-15 years),




FIG. 334.

SIMON'S URETHRAL SPECULA (*Winckel*).

4.7-5.6 cm. ($1\frac{7}{8}$ - $2\frac{1}{8}$ in.) in circumference or 1.5-1.8 cm. ($\frac{9}{16}$ - $\frac{11}{16}$ in.) in diameter.

Practically, we find that the index finger can be passed with safety; and that any dilatation beyond an inch diameter is dangerous in regard to permanent incontinence.




FIG. 335.

THE VARIOUS SIZES OF SIMON'S SPECULA (*Winckel*).

Persistent incontinence has attended the extraction of stones with a diameter of $1\frac{3}{8}$ in., but Dunlap¹ has recorded a case where a stone $2\frac{1}{8}$ in. in diameter was safely extracted uncrushed through the urethra without consequent incontinence of urine.

The dilators of Simon are graduated, and are passed slowly until the desired limit is reached.

¹ *Am. J. of Obst.*, Vol. xiv., p. 855.

It is doubtful if they can, without risk, be used as Simon recommends.

Specular examination by Skene's Specula. These may be described as Skene's small test tubes which fit into a truncated or fenestrated case of vulcanite. The glass tube projects beyond the outer truncated case; and a small mirror can be carried through the inner tube so as to reflect light.

Skene's directions are to pass the tube (with mirror inside) along the urethra, and to use sun-light or gas-light from a movable bracket. When a large Skene's speculum is used, the urethra should be first dilated with the index finger. When viewed through the speculum, the mucous membrane of the bladder is somewhat pale.

The hard rubber speculum can be used in making applications.

c. Postural Examination of the Bladder: Ureteric catheterisation.

Attempts to examine the interior of the bladder by specula such as Skene's are not very satisfactory. The catheterisation of the ureters




FIG. 336.

FINGER PASSED THROUGH URETHRA INTO BLADDER TO GUIDE HOLLOW PROBE INTO LEFT URETER.
a Internal Sphincter of Urethra, *b* Orifice of right Ureter, *c* Inter-ureteric Ligament (*Winckel*).

after the finger had been passed through the urethra (fig. 336) proved difficult, but mainly by the work of Pawlik and Kelly intravesical ex-

amination has been rendered much easier, and an interesting and comparatively simple method evolved. Controversy has arisen between Pawlik and Kelly as to priority in this method (see the references given in the literature). We follow Kelly's method in our descrip-




FIG. 337.

POSITION OF PATIENT FOR EXPLORATION OF BLADDER AND URETER (*after Kelly*).

tion, without, however, expressing any opinion as to the merits of the dispute.

The principle of the method is, that *if the hips be well elevated or the patient placed in the genupectoral posture, the urethra dilated and a specu-*



FIG. 338.

LEFT URETERAL ORIFICE EXPOSED AND SEARCHER ENGAGED. ELECTRIC LAMP HELD BY ASSISTANT (*after Kelly*).

lum passed in, air will pass into the bladder and its walls will become separated and can be illuminated so that the vesical mucous membrane and especially the ureteric openings can be inspected.

Method practised by Kelly. The patient is anæsthetised. The urethra is dilated by Hegar's dilators up to a diameter of 12 mm. A speculum with an obturator is then introduced and the patient's hips raised by means of cushions 12-16 inches above the level of the table (fig. 337).

The speculum is then withdrawn, air passes in, and by means of a small electric lamp and mirror (fig. 338), light is thrown into the interior of the bladder. Any urine in the bladder will require to be sucked out with tube and suction bulb.

The walls become separated by a distance varying between one and two inches. If the speculum (fig. 339) be sloped 30° to one or other side (as shown by the direction of the limbs of the V marked on its side) the ureteric orifices can be seen, and a catheter passed into them. Slender catheters have indeed been passed up to the pelvis of the kidney, and the ureters have also been defined prior to hysterectomy.

This method promises well but must be carried out with strict anti-sepsis, and only when necessary and other simpler means have failed.




FIG. 339.

SPECULUM MARKED FOR FINDING URETER (Kelly).

The ureteric catheter can also be passed in the genupectoral posture and without any preliminary dilatation of the urethra. The inter-ureteric ligament can be mapped out through the anterior vaginal wall, and thus affords a guide to the introduction of the catheter or bougie into the ureter.

Electric Endoscope.

This handy and convenient instrument has now been used with great success in the diagnosis of vesical conditions. It would take up too much space to describe its construction and use fully; these can be found in the special works on this subject. We may, however, state that the instrument has been brought to its present value chiefly by the labours of Nitze and Leiter, and that the introduction of the small incandescent lamp as the illuminating agent has probably been the greatest improvement.

By this means we can ascertain the position of the ureter in operating

on vesico-vaginal fistula and prior to excision of the cancerous uterus ; and in proposed excision of a kidney we can ascertain the state of the other kidney by examination of the urine from it. For a full description see the papers by Fenwick and Wallace. Viertel's article in Veit's *Handbuch der Gynäkologie*, also gives a good account.

CHAPTER LIII.

AFFECTIONS OF THE URETHRA AND BLADDER.

For LITERATURE, *see* CHAPTER LI.

MALFORMATIONS OF THE URETHRA AND BLADDER.

THESE comparatively rare malformations are easily understood on consideration of the development of the organ.

The bladder is the part of the allantois above the lower openings of the Wolffian ducts (figs. 300 to 304), and the upper part of the urethra, its narrow lower portion. The lower portion may also be urino-genital sinus, or may be formed by fresh invagination of the urino-genital sinus after the latter has been blocked by the Wolffian bulbs (*v. p.* 83), while the lower is formed by an invagination from the genito-urinary sinus. The developmental defects are therefore the following :—

- (1) Total absence of urethra ;
- (2) Defect of external portion of urethra—hypospadias ;
- (3) Defect of internal portion of urethra ;
- (4) Atresia of the urethra (in malformed fœtuses) ;
- (5) Extroversion of the bladder from deficient closure of the embryonic abdominal plates.

We would here only note the rarity of these conditions, and refer the practitioner to Skene or Winckel for details.

DISEASES OF THE URETHRA.

Of these the most important are Displacements, Neoplasms, Urethritis, Dilatation, and Stricture.

DISPLACEMENTS.

These will be easily understood by reference to those of the bladder.

Urethrocele is a pouching of the urethra and vaginal wall allowing the

lodgment of stale urine. It is treated by excising a portion of the urethral wall and uniting the edges by stitches.

Prolapse of the mucous membrane of the urethra through the urethral orifice may be remedied by the button-hole operation. The incision is made down to the submucous tissue, and the mucous membrane pulled through this until the excess at the urethral orifice disappears. The excess at the button-hole is then cut off and the wound stitched.

NEOPLASMS OF THE URETHRA; URETHRAL CARUNCLE.

Urethral
Caruncle.

The urethra is liable to be invaded by papillomata, polypi, sarcomata, cysts, carcinomata, and vascular growths (angiomata).




FIG. 340.

CARUNCLE AT URETHRAL ORIFICE (a) AND, IN ADDITION, NEUROMATA IN SURROUNDING MUCOUS MEMBRANE—see page 588 (*Sir J. Y. Simpson*).

Of these last, the most common is the well-known Urethral Caruncle.

Pathology. This is a vascular excrescence varying in size from a pin head to a strawberry; it consists of dilated capillaries in connective tissue, the whole being covered with squamous epithelium. *Physical Signs.* A cherry-red tumour, exquisitely tender and vascular, is seen at

the urethral orifice (fig. 340). *Symptoms.* These are pain on micturition or even retention of urine, and pain on coitus. *Treatment.* Place the patient under chloroform in the lithotomy posture, and destroy the growth by Paquelin's cautery at a dull heat. If bleeding occurs do not treat it lightly; plug the vagina, bringing the half of the last strips of lint over the urethral orifice and fixing with a perineal band.

As regards the other neoplasms, papillomata are painless, sarcomata very rare, their nature being determined microscopically; while carcinomata appear as hard peri-urethral tubercles which break down (*Skene*). In regard to treatment they may be removed by the curette, or by small loop-snares when high up. Emmet's button-hole operation is probably the best method. Polypi in the urethra may cause great difficulty in micturition and should be suspected in intractable cases, and examination made by incision of urethra. We may also have specific inflammatory changes in Skene's "tubules" (*v. p.* 34) simulating urethral caruncle. These may be gonorrhœal, simple catarrhal, or tubercular. The last is usually found with tubercular disease elsewhere.

The tubules may require to be slit up and cauterised.

URETHRITIS.

Acute urethritis is usually part of a gonorrhœa. When pus is secreted, the urethra can be felt swollen and tender; the pus can be squeezed out of the urethral orifice by pressure from above downwards; on passage of the sound, pain is felt in the urethra although no cystitis be found.

Treatment. Give diluent drinks so as to increase the flow of urine. Copaiba may be given in the form of the well-known Nesbitt's specific:—

R Liquoris Copaibæ Co. (Nesbitt) ℥ij.

Sig. Teaspoonful thrice daily.

Iodoform bougies may be passed in, and counter-irritation applied in the shape of the tincture of iodine over the anterior vaginal wall.

Urethritis is very intractable. Emmet advises his button-hole operation to relieve tension and allow of accurate application of local remedies.

DILATATION, AND STRICTURE OF THE URETHRA.

The urethra may be unusually *dilated*, a condition rarely met with; in some cases the dilatation has been caused by coitus, as in malformations of the vagina (*v. p.* 292). The dilation may be local or general. When it is general, the cautery may be used to burn a vertical furrow, the rest of the urethra being guarded by a speculum.

Stricture of the urethra is a rare condition and readily yields to dilatation by bougies or to incision.

DISEASES OF THE BLADDER.

Of the diseases of the bladder we shall here consider Displacements, Neoplasms, Cystitis, and Stone in the Bladder. Vesico-vaginal fistula will be considered in a separate chapter (Chap. LIV.).

DISPLACEMENTS OF THE BLADDER; CYSTOCELE.

The female bladder when empty lies behind the pubes and usually to one or other side. It is never exactly central.

The
mobility
of the
Bladder.

From its loose attachment to the pubis, it is pre-eminently displaceable. (1) It is drawn up during labour; and (2) is displaced upwards by retroversion of the gravid uterus, pelvic ovarian or fibroid tumours, and pelvic hæmatocele. (3) It may be adherent to the anterior surface of an abdominal ovarian or fibroid tumour, and may thus be cut into on abdominal section. (4) It is displaced downwards in prolapsus uteri and in cystocele. (5) In utero-sacral cellulitis, the bladder is drawn back and fixed; its systole is thus interfered with, which explains some cases of so-called hysterical retention of urine. From this mobility it follows that the height of its fundus above the symphysis gives no indication of the amount of urine in the bladder.

By *cystocele* we understand a pouching of the posterior wall of the bladder downwards and backwards; the uterus and summit of the bladder are in normal position.

Senile
form of
Prolapsus.

Many a case, regarded as cystocele, is really part of a prolapsus uteri; on the other hand, the so-called "senile prolapsus uteri" is really a cystocele; at the menopause the cicatrization of the vaginal walls chiefly affects the posterior one, and thus the bladder tends to bulge outwards at the vaginal orifice.

The diagnosis is easily made by the bimanual and use of the sound. The treatment consists in the use of a ring pessary with diaphragm (fig. 319). Should this fail, the vagina may be packed with oakum; or a raw surface (as shown at fig. 315) may be made and stitches applied.

* NEOPLASMS OF THE BLADDER.

Pathological anatomy. We may have mucous, fibroid or fibro-myxomatous polypi. There may also be sarcomatous or carcinomatous disease of the bladder wall, as well as tubercle. In tubercular disease the ulcerated surface has been removed by Schatz in a supra-pubic operation. The carcinomatous condition is not infrequent, and is termed by some "villous cancer." It is most common at the trigone, and is held by some authorities not to be malignant. The bladder may be secondarily affected in carcinoma uteri (*v. p.* 492).

Symptoms. These are disturbances of micturition, with bloody and phosphatic urine.

Physical signs. The passage of the index finger into the bladder will show the position, shape, and other characters of the growth.

Treatment. This will vary according to the position, nature, and pediculation or non-pediculation of the growth. Thus it may be twisted off by narrow polypus forceps, snared by a loop of fine catgut, or removed by incision into the posterior wall of the bladder and use of the galvano-cautery or curette.

CYSTITIS.

Nature. An acute or chronic inflammatory affection of the mucous membrane of the bladder.

Pathological anatomy. In the acute catarrhal form, we have congestion of the vessels and loss of epithelium; in the chronic catarrhal form, the congestion is duller and there is marked rugosity of the lining of the bladder. The submucous and even the muscular tissues also become affected. The mucous membrane may be ulcerated and the muscular tissue exposed.

Various micro-organisms may be found, usually the tubercle bacillus, bacillus coli communis, pyogenic organisms; details as to these and the methods of investigation to be employed will be found in Gulland's or in Barlow's paper.

The inflammatory process may extend deeper, to the muscular tissue (interstitial cystitis), to the peritoneum (pericystitis), or to the connective tissue near (paracystitis). Occasionally, though rarely, we may have diphtheritic inflammation.

In advanced cases, the patient may be septicæmic, and there is often hydro-nephrosis. In some cases of prolonged retention the mucous membrane may slough off and be passed per urethram, but may be regenerated.

Results of
Cystitis.

Etiology. The causes are as follows:—Gonorrhœa; latent gonorrhœa; exposure to cold; injury from coitus; prolonged parturition; introduction of septic matter by catheter or bougie; prolonged retention of urine; stone.

Symptoms. In *acute cystitis* the patient has very frequent and painful micturition. In *chronic cystitis* also, there is frequent micturition but accompanied with less intense pain; there are, further, shooting pains with secondary phenomena—septic, vascular, and nervous.

Physical signs. (a) *Acute cystitis.* The urine has a lowered specific gravity and acid reaction; the colour is little altered, and mucus is present in excess. On vaginal examination, pain is not felt when pressure is made on the posterior vaginal wall, but is felt severely *when the anterior wall is touched.*

Char-
acters of
Urine in
Cystitis.

(b) *Chronic cystitis.* The urine has a low specific gravity, is usually alkaline, and is often offensive; it contains pus, epithelium, phosphates and bacteria; albumen, derived from the pus, is present. The vaginal examination gives the same results as in acute cystitis. If the finger be passed through the urethra (*v. p.* 641), the roughened condition of the lining membrane is felt; crystals of phosphate and marked rugosities can also be detected.

Genito-urinary phthisis is often diagnosed as chronic cystitis. In the former condition we get at first the symptoms of chronic cystitis, viz., purulent urine, pain, and intractability to treatment. Local examination of the bladder may give no definite result, and if the kidney is not palpated its enlargement and purulent condition may not be noticed until the disease is far advanced.

Prognosis. In both acute and chronic cystitis, the prognosis is not good; the treatment is difficult, and in bad chronic cases the patient's strength sometimes becomes exhausted, and septicæmia may cause death.

Treatment
of Acute
Cystitis.

Treatment. (a) *Acute cystitis.* Put patient on milk diet, and give Friedrichshall or Carlsbad water freely. Diluent drinks may be taken *ad libitum*.

The following prescription is useful.

R	Potassii Bicarbonatis	℥ iss.
	Tincturæ Hyoscyami	℥ i.
	Infusum Buchu	
	vel Pareiræ	
	vel Uvæ Ursi ad	℥ vj.
	<i>Sig.</i> Tablespoonful thrice daily.	

In gonorrhœal cystitis, the following may be substituted:—

R	Liquoris Copaibæ Co. (Nesbitt)	℥ ij.
	<i>Sig.</i> Teaspoonful thrice daily.	

If the pain is very acute give morphia suppositories ($\frac{1}{4}$ grain) at night, omitting the mixture with the hyoscyamus if necessary.

For (b.) *Chronic cystitis*, we recommend the following treatment *seriatim*.

Treatment
of Chronic
Cystitis.

1. Put on milk diet with abundant fluids, and purge freely. Give

R	Acidi Nitrici diluti	℥ iij.
	Tincturæ Hyoscyami	℥ i.
	Infusum Buchu ad	℥ vj.
	<i>Sig.</i> Tablespoonful thrice daily.	

The hyoseyamus eases the pain; and the nitric acid corrects the alkaline phosphatic urine, for which also benzoate of ammonia is admirable.

R	Ammonii Benzoatis	℥iii.
	Aquæ	℥vj.
	<i>Sig.</i> Tablespoonful thrice daily.	

The benzoate of ammonia is converted into hippuric acid and corrects alkalinity. Lithia water, tincture of belladonna, and Nesbitt's specific are also useful.

Salol in 5 to 10 gr. doses thrice daily is also excellent. Boracic acid in 5 to 10 gr. doses needs caution in its use owing to its tendency to set up gastric catarrh.

2. If this fail, then wash out bladder as often as possible by means of double catheter, such as Skene's, using corrosive sublimate (1-5000 or 8000), weak boracic lotion, or carbolic lotion. A simple bladder douche can be made with a glass funnel and india-rubber tubing. By raising it and lowering it, fluid can be made to pass in and out. We strongly



FIG. 341.

THE SKENE-GOODMAN SELF-RETAINING CATHETER; AN INDIA-RUBBER BAG CAN BE WORN WITH IT (*Skene*).

recommend weak corrosive sublimate as a bladder douche. Paint anterior-vaginal wall with tincture of iodine.

3. A long (winged) india-rubber catheter may be kept in the bladder so as to drain off the urine constantly and give the bladder rest. The patient need not remain in bed if the Skene-Goodman catheter (fig. 341) is used.

4. In obstinate cases, the formation of an artificial vesico-vaginal fistula gives excellent results. To do this, chloroform the patient; place her in the lithotomy posture and apply Sims' speculum. Open into the bladder through the antero-vaginal wall, in the middle line, with the scissors, as follows: pass a sound into the bladder, project the point into the vagina, and then by means of a pair of straight scissors open the bladder in the middle line. The mucous membrane of the bladder is then stitched to the adjacent vaginal mucous membrane by means of chromic acid catgut.

The urine trickles through the artificial fistula; in this way, the bladder gets complete rest and can be thoroughly washed out.

After some months the fistula is easily closed, as in the operation for vesico-vaginal fistula. Severe cases of cystitis will tax more than any other disease, the practitioner's patience and knowledge. It is well to keep in mind the reason of this intractability, viz., the inability of the bladder to remain at rest, and the micro-organismal origin of the worst cases.

As can be seen from what has gone before, the principles of treatment are the following:—(1) to correct abnormalities in the urine; (2) to allay the irritability of the bladder; (3) to lessen the congestion of the bladder by purgatives and counter-irritants, and to render the urine bland and lessen the work of the kidney by milk diet; (4) to allay the irritable condition of the bladder and counteract putrefaction or gonorrhœal inflammation by injection; (5) to give it complete rest by a permanent catheter or, in extreme cases, by an artificial fistula.

CALCULI AND OTHER FOREIGN BODIES IN THE BLADDER.

The female bladder is liable to receive foreign bodies from three sources.

A. Calculi from the kidneys—uric acid, oxalates, phosphates or cystine.




FIG. 342.

LARGE STONE WHICH FORMED ROUND A HAIR-PIN AS NUCLEUS, EXTRACTED BY VAGINAL LITHOTOMY (*Angus Macdonald*).

B. Substances from neighbouring organs—pus from pelvic abscess, concretions from the intestines, bones from an extra-uterine foetation, pessaries from the vagina, echinococci and other parasites such as those associated with chyluria.

C. Foreign bodies introduced wilfully into the bladder by patients of a depraved taste; these may form nuclei for stones (fig. 342).

Of these, *calculi* are the most important. Stone is less common in the female than in the male, as small calculi can pass along the dilatable female urethra; occasionally, therefore, the gynecologist has to remove from the urethra small stones impacted there—usually at the meatus urinarius. The introduction of foreign bodies, which act as nuclei, is more common in the female.

Symptoms. These are severe pain in micturition, especially at the close; alterations in character of urine; blood in urine.

Physical Signs. The stone, when at all large, can be easily detected bimanually; when any doubt exists, the use of the sound or the passage of the finger into the bladder renders the diagnosis easy. Diagnosis
of Calculi.

Treatment. Measure the stone: if it be less than an inch, it may be extracted through the urethra dilated first by the finger or Simon's specula; if greater than an inch, then dilate the urethra and crush; if very large or hard or if it have a nucleus, extract by vaginal incision. This incision may be stitched up after the operation, or kept open when the bladder has been much irritated; it can afterwards be stitched as in vesico-vaginal fistula. Supra-pubic lithotomy is sometimes required.

For other foreign bodies, the urethra can be dilated and the substance grasped by polypus forceps or manipulated out. When large, they may be extracted as in the case of large stones.

FUNCTIONAL DISEASES OF BLADDER.

By these we understand derangements of the bladder in regard to urination. Either these are due to causes as yet unascertained, or the same derangement (*e.g.*, retention) is associated with many lesions. Functional
affections
of the
Bladder.

The chief functional diseases are—

Irritability,
Incontinence,
Retention.

In regard to all of them, we may remark that in no case should the diagnosis of a functional disease of the bladder be made until the practitioner is satisfied that there is no organic lesion.

Irritability. In this, frequent micturition associated with a disagreeable feeling is present. It may be due to excessive acidity of the urine, but is often a nervous affection. When it is due to excessive acidity give lithia or potash.

R Lithii Carbonatis gr. v.
Fiat pulv. mitte tales vj.
Sig. One thrice daily.

Incontinence, or inability to retain urine long enough, is most common in little girls; occasionally we meet with it in adults, as the result of prolonged labour, as a permanent condition from infancy, or in oxaluria cases.

In the incontinence of girls, note whether there be any irritability of the genitals (vulvitis) or ascarides. Goltz found that, where section of the spine in the dog above the lumbar enlargement had produced retention of urine, he could make it urinate by sponging the anus with cold water; a reflex impulse passed from the rectum, lessening the activity of the inhibitory centre and allowing bladder contraction. In a child, ascarides in the rectum will act in the same way when it is asleep.

Treatment. Treat the irritating cause—as vulvitis or ascarides. If no irritating cause be detected, then give belladonna.

R Tincturæ Belladonnæ ʒij.

Sig. Three drops thrice daily.

In strumous cases, give syrup of the iodide of iron or cod liver oil.

R Syrupi Ferri Iodidi ʒij.

Sig. Thirty drops thrice daily.

The faradic current is also indicated.

Retention of Urine. Palpation shows a fluctuating mesial tumour rising into the abdomen; the position of the fundus of the bladder gives no indication of the amount of urine, as it may be tilted up by retroversion of the gravid uterus. Remember that a bladder may be distended so as to be as large as a six or eight months' pregnancy, and that constant dribbling-away of the urine may be a symptom of retention. Examine the pelvis for an organic lesion.

Retention may be due to one of three great classes of causes:—

Hysterical,
Reflex,
Mechanical.

1. *Hysterical.* By this we mean that from perversity or a prurient desire to have the catheter passed, a patient feigns inability to pass urine.

The treatment is to give a hot hip bath followed by a cold one; if the catheter is needed, get it passed by a nurse of unsympathetic tendencies.

2. *Reflex* causes are the following:—

- (1) Gonorrhœa;
- (2) Urethritis;
- (3) Irritable caruncle;
- (4) Carcinoma, urethral and vaginal;
- (5) Perineal and especially vestibular tears after labour, tears of cervix;
- (6) Ligature of internal piles.

The treatment is hot appliances in (1), (2), (3), and (5) and (6); and the catheter in (4). Remove the source of irritation when possible.

3. *Mechanical.* These are pressure of fibroids, retroversion of the gravid uterus; ovarian or parovarian tumours (pelvic and retro-uterine).

Where the tumour is impacted in the pelvis, a silver male (No. 10) catheter will pass best. The urethra is compressed, the bladder bulging over the symphysis; accordingly, a rigid instrument whose handle can be carried to the perineum is good.

Final Cautions as to Bladder Cases. 1. Remember that pus in the urine does not necessarily mean that the *bladder* is affected. Keep in mind the possibility of disease in the ureter, in the pelvis of the kidney, and in the kidney itself. Do not forget the possibility of pus forming in the neighbourhood of the kidney and opening into the urinary tract. In a bladder case, always palpate and percuss the abdomen.

2. Make out as much as possible by examination of the urine, by bimanual examination of the bladder, and by the use of the sterilised sound before proceeding to more active methods.

3. Dilatation of the urethra and other instrumental means are excellent but require caution, as they often set up fresh irritation.

4. In cystitis, give drug treatment, and treatment by milk diet a thorough and careful trial.

5. In all operative treatment, take the most strenuous precautions as to antisepsis.

6. In intractable cystitis, the formation of an artificial fistula is a form of treatment more controllable in its ultimate results than extreme urethral dilatation.

CHAPTER LIV.

VESICO-VAGINAL FISTULA.

LITERATURE.

- Baker Brown*—Surg. Diseases of Women, 3rd Ed., p. 133; and *Lancet*, March 1864.
Bandl—*Wiener med. Woch.*, 1875, Nos. 49 to 52; and 1877, Nos. 30 to 32. *Boldt*—Implantation of the ureter into the bladder for the cure of uretero-vaginal fistula: *Am. Journ. of Obstet.*, 1896, xxxiii., p. 846. *Bozeman*—Remarks on Vesico-Vaginal Fistula, 1856; *Americ. Journ. of Med. Science*, July 1870; *Obst. Journ. of Great Britain*, June to Aug. 1878. *Byford*—*Medical and Surgical Diseases of Women*: Philadelphia, 1882. *Champneys*—Description of a new operation for Vesico-Uterine Fistula: *Obst. Trans.*, 1888, p. 348. *Emmet*—On Vesico-Vaginal Fistula: New York, 1868. *Ferguson, A. H.*—Vesico- and Recto-Vaginal Fistulæ: *Brit. Med. Journ.*, 1894, Vol. i., p. 406; and *Amer. Journ. Obst.*, 1895, Vol. xxxi., p. 476. *Hegar und Kaltenbach*—*Die Operative Gynäkologie*, S. 582: Stuttgart, 1881. *Phillips*—*Plastic gynecological Operations: Clifford Allbutt and Playfair's System of Gynecology*, London, 1896. *Simpson, Sir J. Y.*—*Diseases of Women*, p. 30: Edin., 1872. *Sims, Marion*—On the Treatment of Vesico-Vaginal Fistula: *Americ. Jour. of Med. Science*, Jan. 1852. *Silver Sutures in Surgery*: New York, 1858. *Simon*—Ueber die Heilung der Blasenscheidenfisteln; Giessen, 1845; Rostock, 1862; and *Wiener med. Wochenschrift*, 1876, Nos. 27-32. *Webster, J. C.*—Operative Treatment of Vesico-Vaginal Fistula: *Edin. Hosp. Reports*, 1895, p. 689. *Winckel*—Die Krankheiten der weiblichen Harnröhre u. Blase: Stuttgart, 1877, S. 95. *Winternitz*—Zur Behand. der Blasen-fisteln: *Cent. f. Gyn.*, 1895, S. 377.

PATHOLOGICAL ANATOMY AND VARIETIES.

THE septum between the urinary and genital tracts may be broken through at various points. According to their situation, we have the following varieties of urinary fistulæ:—

Urethro-vaginal,
Vesico-vaginal,
Vesico-uterine,
Uretero-vaginal,
Uretero-uterine.

The situation of these is sufficiently indicated by their names, and will be easily understood by reference to fig. 343.

A urethro-vaginal fistula rarely occurs alone, but is sometimes present along with a vesico-vaginal one. It lies in the middle line and is, naturally, of smaller size.

By far the most frequent are the vesico-vaginal fistulæ. They may occur at any point of the vesico-vaginal septum, which measures in

Pathology
of Vesico-
vaginal
Fistula.




FIG. 343.

TO REPRESENT THE CHIEF VARIETIES OF URINARY FISTULA—URETHRO-VAGINAL, VESICO-VAGINAL, AND VESICO-UTERINE. Those with the ureters are not seen. The seat of a recto-vaginal fistula is indicated (*De Sinéty*).

height (from the internal orifice of the urethra to the vaginal fornix) about 5 cm. and in breadth 4 cm. (*Kaltenbach*). Their size varies from




FIG. 344.

SUPERFICIAL VESICO-VAGINAL FISTULA, the Cervix is intact (*Hegar and Kaltenbach*).




FIG. 345.

DEEP VESICO-VAGINAL FISTULA, the anterior lip of the Cervix is destroyed (*H. and K.*).

a pin-point or slit-like hole to a large oval (fig. 348) or four-cornered (fig. 367) aperture. When recent they are of larger size, but after

some months become contracted through the formation of cicatricial tissue. The *margins* of the *fistula* are at first irregular, swollen, and ulcerated; but after a time they become thin and firm, through cicatrisation: these changes have an important bearing on treatment. Jobert divided fistulæ in the anterior fornix into *superficial* and *deep*; in the former (fig. 344) the anterior lip of the cervix was not implicated, in the latter it was more or less destroyed (fig. 345). In cases of fistulæ which allow a free flow of urine, the *bladder* becomes permanently contracted and its walls thickened; in large fistulæ, the mucous membrane protrudes through the opening and is easily recognised from its deep red colour. The normal relation of the openings of the *ureters* to that of the *urethra* and to the cervix uteri (fig. 346) renders them liable to be involved in an extensive fistula, or even in a small one lying to one side of the middle line. Sometimes we can recognise their openings on




FIG. 346.

THE NORMAL RELATION OF THE CERVIX, THE URETERS, AND THE URETHRA (*H. and K.*) From cervix to orifice of ureter measures 3 cm., from orifice of ureter to that of urethra measures 4 cm., from orifice of one ureter to that of the other measures 2.5 to 3 cm. The ureters run through the bladder wall in an oblique direction downwards and inwards, for from 1.5 to 2 cm.

the exposed vesical mucous membrane by means of the urine trickling from the orifices; should the urine be bloodstained, it can be distinguished from blood by its acid reaction to test paper. The *urethra*, through disuse, becomes contracted; sometimes complete atresia is present and seriously complicates treatment, and a portion of the canal may even be completely destroyed by pressure (*v. fig. 371*). The *vagina* is often contracted by cicatricial tissue originating from injuries received during labour. The margins of the fistula are often drawn apart, and sometimes fixed down to the bone, by these cicatrices; this interferes with their closure. Contraction of the vagina below the fistula sometimes makes it impossible to ascertain the condition of the upper part and whether the uterus communicates with the fistulous tract. The rela-

tions of the *peritoneum* to fistula are shown in fig. 347, from which it is evident that only in the repair of very extensive fistulæ would its relations require to be considered. The difficult labour which leads to the production of the fistula is liable to be followed by puerperal peritonitis or cellulitis; these may disturb the normal relation of the peritoneum.

Vesico-uterine fistulæ are rare. From their position they can be recognised only after dilatation of the cervical canal (*v.* fig. 343), and it is evident that they must be very small.

Uretero-vaginal fistulæ are situated in the fornix vaginæ. They are of small size, admitting only the point of the sound, and have either sharp edges or open at the point of a small papilla.

ETIOLOGY.

Malignant disease is the most common cause of fistula (*v.* p. 492); but we place this form aside, as it is beyond treatment and merely indicates a stage in the progress of the malignant growth.




FIG. 347.

RELATIONS OF PERITONEUM, indicated by dotted line, to a fistula which has destroyed the whole of the anterior wall of the cervix and the infra-vaginal part of the posterior wall (*H.* and *K.*).

The most important cases of fistulæ which we have to consider here, arise through *injury received during labour*. This injury may act *directly*, producing laceration of the septum; more frequently it acts *indirectly* producing necrosis secondary to pressure or inflammation. The causes which predispose to fistula are a narrow pelvis and pendulous abdomen, a firm or large head (hydrocephalic), and face presentations. The immediate cause is the compression of the soft parts between the child's head and the bony wall of the pelvis; if this pressure continues for a long enough time, it destroys the vitality of the soft parts which afterwards separate as a slough.

Mode of
production
of Fistula
in Labour.

Fistulæ produced by instruments are situated in the lower part of the vagina, and are accompanied with extensive cicatrices and adhesions; those due to pressure of the fœtal head are in the upper part. In craniotomy, the soft parts have been sometimes lacerated by the in-

struments, or by splinters of foetal bone. Forceps are often cited as a cause of the injury. It is not however *the use* of the forceps after a prolonged labour which is to blame, but the *not using* of them at an early period—before the parts have been destroyed by pressure.

Fistulae have followed diphtheritic inflammation in the puerperium, but this is rare. Inflammation and ulceration round badly fitting pessaries have also produced them. Ureteric fistulae may result from injury to the ureter in the operation of vaginal hysterectomy (*v. p.* 517).

SYMPTOMS.

The leading symptom is the *involuntary flow of urine* from the vaginal orifice. *This will not appear until the slough separates*, that is till about the third or fourth day; its separation may be delayed for three or four weeks, when the necrosis is secondary to puerperal vaginitis (*Byford*). When a direct laceration has been produced, the urine will flow *at once* per vaginam; but even here it may escape notice till the second or third day, as it is masked by the lochial discharge.

The power of retaining varies, in certain cases, with the position of the patient; with a fistula situated high up, the erect posture allows the lower portion of the bladder to be used though the flow is continuous in the recumbent posture. With a urethro-vaginal fistula, there may be perfect continence from a sphincter-like action of the muscular fibre in the wall of the urethra; the patient observes, however, that the urine does not pass by the urethral orifice.

Secondary symptoms are due to a constant wetting of all the surrounding parts with the urine. The urinous *odour* is quite characteristic in urinary fistula; there is *excoriation* round the vulva, the inside of the thigh is red and irritated. *Menstruation* is generally in abeyance, returning after the fistula has been cured. There is usually *sterility*; although cases of conception, often followed by abortion or premature labour, have been recorded. The disagreeable surroundings interfere with the appetite and digestion; there is *constipation*, which Freund has ascribed to increased secretion by the kidneys but which is more probably due to reflex contraction of the muscular fibre of the rectum (*Winckel*). The general health thus becomes seriously impaired so that the patient is willing to submit to any operation which promises relief.

DIAGNOSIS.

The irritated appearance of the external genitals with the characteristic odour at once indicates that there is a fistula, but the diagnosis of its position is often very difficult.

Urethro-vaginal and *vesico-vaginal*. When large, these may be felt by the examining finger; on our passing the sound into the bladder the

finger touches it through the fistula. The speculum shows their position and extent, and reveals smaller ones which escape detection with the finger; by stretching the folds of the mucous membrane with tenacula, we may detect a fistula concealed by them.

To recognise *small vesico-vaginal* fistulæ and to differentiate them from the *vesico-uterine* and *ureteric*, proceed as follows:—pass Sims' speculum, carefully wipe away all mucus from the anterior vaginal wall, clear out the cervical canal with a dressed sound and plug it with a pledget of dry cotton wadding; now pass a catheter, and through it distend the bladder slowly with a coloured fluid such as milk or permanganate of potash; as the bladder distends, watch carefully the anterior vaginal wall for any oozing of the fluid. If there is no oozing, the fistula is not vesico-vaginal. If on withdrawing the plug from the cervix it be found stained with fluid, the fistula is *vesico-uterine*. If neither of these forms be present, the urine must come from a *ureteric* fistula; the rarity of this form should lead us to suspect that the fluid may have been temporarily kept from escaping from the bladder by a valvular action of the mucous membrane, and the examination should be repeated after a time. In a case of uretero-uterine fistula, Bérard collected the urine which escaped per vaginam in one vessel and that in the bladder was drawn off per urethram by a catheter into another; the quantities in a given time were found to be equal. His conclusion was that he had obtained the secretions from each kidney separately, so that the fistula was ureteric.

PROGNOSIS.

A *natural cure* will depend on the recentness of the fistula and its size. Small fistulæ, if kept clean, heal of themselves during the puerperium. Large ones require *operative treatment*; cure by this means depends partly on the size of the fistula, but more on the condition of its margins—whether they contain much cicatricial tissue, and whether they are bound down.

TREATMENT.

There are two essentials for successful operative treatment: (1) complete exposure of the fistula, so that (2) the edges may be thoroughly pared and carefully adapted with sutures. The great difficulty lies in the inaccessibility of the field of operation, to which the failure of the older operative measures is chiefly to be attributed.

Marion Sims (1849) first rendered successful treatment really possible by the complete exposure of the fistula with his *speculum*, and by the *careful adaptation* of its margins with *silver-wire sutures*. Since the introduction of *catgut*, we believe that it will displace silver wire in this operation as it does not need to be removed subsequently. To Simon

of Heidelberg is due the credit of having elaborated the operation, and of having extended its sphere so that almost no form of fistula has in his hands proved incapable of treatment. We may shortly contrast the methods adopted by these two operators as follows: Sims pared the edges of the fistula in a sloping manner (fig. 350) carefully avoiding the mucous membrane of the bladder, then adapted the margins of the fistula




FIG. 348.

METHOD OF PARING THE EDGES OF A FISTULA (*Simon*).

with silver wire, and drained the urine continuously per urethram through a catheter: Simon pared away the edges vertically not specially avoiding the mucous membrane of the bladder, united the edges with silk sutures, and encouraged the patient to pass water unaided from the

first—drawing it off with the catheter only when necessary. Bozeman, a pupil of Sims, has drawn attention to the advantages of the genu-




FIG. 349.

SUTURES PASSED IN A CASE OF URINARY FISTULA (*Simon*).



FIG. 350.

SIMS' AND SIMON'S METHODS OF PARING THE EDGES OF FISTULÆ CONTRASTED: Sims' is shown on the right, Simon's on the left. The mucous membrane of the bladder is above, that of the vagina is below. The edges may be pared first according to Sims' method, and if a raw surface is not thus obtained the tissue can be removed up to the fine line (*Kaltenbach*).

pectoral posture in operating and to the importance of preparatory

treatment by dividing and stretching cicatricial contractions; he fixes the sutures with lateral plates and buttons (fig. 364).

When a fistula has been discovered during the puerperium, our first aim is to *aid the natural effort at cure*. A catheter (fig. 372) is placed in the urethra to carry off the urine by the natural passage; the vagina is syringed out frequently with warm water; the edges of the fistula may be kept together, in some cases, by tampons suitably placed in the vagina.

If the fistula does not close by the natural process, we have recourse to operation.

Operation for Vesico-vaginal Fistula.

There is difference of opinion as to the *time for operating*. According to Hegar and Kaltenbach, the best time is six to eight *weeks* after the confinement; "the lochial discharge has ceased, the necrosis of the tissues is defined, the margins of the fistula are vascular and juicy and are at the same time of sufficient firmness to hold the sutures"; the cicatricial tissue which forms round the margins makes the operation more difficult afterwards. Marion Sims delays the operation for a few months.

Under the operation, we shall describe—

1. Preparatory treatment;
2. The operation, which consists of (1) rawing the margins of the fistula by (a) paring, (b) splitting, or (c) making a flap; and (2) the adaptation with sutures;
3. After-treatment.

1. *Preparatory treatment* is only necessary when there are cicatricial bands drawing the margins of the fistula apart or contracting the field of operation. These must be divided and made to heal over a glass plug, or the vagina must be kept distended with air-bags. Frequent vaginal injections are necessary in all cases, to bring the edges into as good condition as is possible.

2. For *the operation* itself the following instruments are required—

Sims' or Fritsch's speculum,
Spatulæ,
Three or four tenacula,
Blunt-hook,
Vaginal douche for permanent irrigation,
Hot water to check hæmorrhage,
Dissecting and artery forceps,
Small bistouries straight or set at an angle—on long handles,
Bozeman's scissors,
Several small sponges and sponge-holders,
Short curved needles and needle-holder,
Curved needles on fixed handles,
Silver wire and wire twister, catgut, or silk-worm gut.

Good light is essential and as *complete exposure* of the field of operation as is possible; this last will determine the *position* of the patient, according as Sims' or the lithotomy posture allows us to get more readily at the fistula. The drawing down of the cervix with volsellæ or sutures (fig. 348), or the protrusion of the edges of the fistula by a catheter in the bladder, is of use in some cases; where the mucous membrane of the bladder (by prolapsing through the fistula) comes in the way, it can be kept back by the sound in the bladder or a sponge probang pushed through the fistula.



FIG. 351.



FIG. 352.



FIG. 353.

KNIVES FOR PARING A FISTULA. Fig. 351, straight knife; fig. 352, bent knife which is shown laterally at fig. 353 (*Sir J. Y. Simpson*).

Chloroform is always an advantage, as it gives the operator more freedom in exposing the parts and prevents the patient from moving; the actual pain of the operation does not demand it.

Three *assistants* are needed—one to give chloroform, a second to hold the speculum, a third for the sponges; six are better, as two are required with the patient in the lithotomy posture and there is one to



FIG. 354.

SPONGE-HOLDER.

take charge of the instruments. The knives employed are shown at figs. 351–353. The sponges should be very small and fitted on holders of which a convenient form is shown at fig. 354. Fixed *needles* are required when the tissue is dense.

(a.) *The paring of the edges of the fistula.* To produce union, it is essential to have a *continuous raw* surface all round the margin. To procure this, we hook up with a tenaculum the portion of vaginal mucous membrane to be removed and transfix it with the knife (v. fig. 348).

The knife should not pass through the mucous membrane of the bladder, unless there be so much cicatricial tissue that a large piece requires to be cut out; the reason for avoiding the vesical mucous membrane is to prevent after-hæmorrhage into the bladder. In small




FIG. 355.

a. Incision for splitting the margins of a Fistula; b. Free edges turned towards the bladder and vagina respectively; b m. Bladder mucosa; v m. Vaginal mucosa.

FIG. 356.

a. Incision for taking flaps from the vaginal aspect of a fistula; b. Flaps turned in towards the bladder, and united at their margins by deep sutures, the raw surface towards the vagina being drawn together by more superficial ones.

fistulæ, we can remove the tissue in a ring and thus ensure a continuous raw surface.

(b.) Another method of making a raw surface is to *split the edges* of the fistula (fig. 355), so that the vesical mucous membrane is separated



FIG. 357.

BOZEMAN'S FORK, used in drawing through the wires to prevent their cutting the Vaginal Mucous Membrane (Sir J. Y. Simpson).

from that of the vagina. The advantage is that no tissue is lost, but the stitching is less accurate.

(c.) Still more tissue can be gained to bridge across the gap by the *flap operation*.¹ Here an oblique incision is made as for rawing the edges, only it stops short of the bladder-margin of the fistula (fig. 356). The flaps formed are turned in towards the bladder, which thus comes to have a strip of vaginal mucosa as part of its lining along the wound, which is united by a buried suture. The raw surface towards the vagina is then drawn together by a continuous catgut suture.

Hæmorrhage is best checked by hot douche; large bleeding points may require twisting or even ligature.

Passage of
Sutures.

(b.) *The adaptation of the edges with sutures* must be carefully done. To prevent the sutures from cutting the vaginal mucous membrane as they are drawn through, the fork or pulley (figs. 357, 358) can be used.

¹ Ferguson—loc. cit.

The sutures must be pretty close together, and should either not pierce the vesical mucous membrane or should take in only its margin. When the tissues are dense, counter-pressure against the point of the needle may be made with a blunt hook, as in fig. 359. Counter-pressure in dense tissue.

After all the sutures are passed, they are tied or twisted; to bring the wires together we can use Bozeman's suture-adjuster (fig. 361); the




FIG. 358.

METHOD OF USING FORK (*Emmet*).

wire twister (devised by Coghill) is very convenient for twisting the wires close, especially when the fistula is deeply placed and not very accessible (fig. 362). Bozeman uses a plate to fix the sutures. The use of catgut does away with all these appliances. With a triangular fistula the closed wound will be Y-shaped, while a quadrilateral fistula will give an I-shaped wound (figs. 367, 368). Bozeman's method.




FIG. 359.

MODE OF APPLYING COUNTER-PRESSURE TO THE POINT OF THE NEEDLE BY MEANS OF A BLUNT HOOK (*Emmet*).

In the case of fistulæ situated close to the cervix, we make use of the anterior lip to close the fistula; the result is a crescentic wound (fig. 369). Fistulæ close to cervix.

369). Sometimes we have to excise a portion of the cervix to get a




FIG. 360.

SIMS' METHOD OF FIXING AND TWISTING THE SUTURES (*after Sims*)



FIG. 361.

BOZEMAN'S SUTURE-ADJUSTER (*Sir J. Y. Simpson*).

sufficient raw surface (fig. 370). When much of the anterior lip is

destroyed, it may be necessary to use the posterior lip to close the fistula (see fig. 345, and compare it with fig. 344); in this case the uterus will



FIG. 362.




FIG. 363.

COGHILL'S WIRE TWISTER (fig. 362); TWISTING THE WIRE (fig. 263).




FIG. 364




FIG. 365.

WIRES DRAWN THROUGH BOZEMAN'S PLATE, fig. 364; fixed with shot as in fig. 365 (Sir J. Y. Simpson).




FIG. 366.

SPECULUM PASSED FOR REMOVAL OF SUTURES; the patient is on her side (Sir J. Y. Simpson).

communicate with the bladder and the menstrual blood be discharged per urethram.

For *vesico-uterine* fistulæ, the best method is to dissect the bladder

Vesico-
uterine
Fistula.

off the cervix until the fistula is fairly exposed.¹ It is then closed with




FIG. 367.




FIG. 368.

FOUR-CORNERED FISTULA, fig. 367, closed by Sutures in fig. 368 (*Hegar and Kalttenbach*).




FIG. 369.

SUTURES PASSED THROUGH ANTERIOR LIP OF CERVIX SO AS TO CLOSE IN TRANSVERSELY A FISTULA OF THE ANTERIOR FORNIX (*H. and K.*).

sutures as is also the corresponding opening of the fistulous tract in the cervix, and finally the wound in the fornix.

¹ Champneys and Mackenrodt—*loc. cit.*

When there is a *urethral* as well as a vesical fistula, the *former must be closed first*: when there is *atresia* of the urethra, the free margins of the urethral wall above and below are pared and united by sutures so




FIG. 370.

ANTERIOR LIP DIVIDED TO CLOSE IN VERTICALLY A FISTULA CLOSE TO IT: *a b c* shows extent of surface, round the oval fistulous opening, to be made raw; the mucous membrane may have to be incised outside the sutures, along the line *A B*, to relieve tension (*Emmet*).

as to bridge over the atresic portion (fig. 371); the vesical fistula is obliterated by a second operation.




FIG. 371.

VESICAL FISTULA + Atresia of a portion of the urethra *w* just below the symphysis *s*. The latter is first bridged over at 1 and then the vesical fistula closed in at 2 (*Winckel*).

Where the margin of a fistula has become cicatrised to the pubic bone, an incision is made in the labium of the corresponding side,¹ and the adherent tissues separated from the bones; and the fistula then treated

¹ As recommended by Schauta—Über die Oper. fixirter Blasenscheidenfisteln: *Monats. f. Geb. u. Gyn. Jour.*, 1895.

in the usual way. Where there is great defect of tissue, the uterus has been called into requisition to fill up the gap.¹

After-treatment.

3. *After-treatment.* A stationary catheter is placed in the bladder—of the ordinary winged form or those shown at figs. 372, 373. The urine is collected in a long narrow vessel (as a soap-dish) passed between the patient's thighs; two catheters are required, so that they may be changed every day as the salts of the urine readily occlude the tube; the one not in use should be kept thoroughly clean.

After-dangers of Operation.

The after-dangers of the operation are hæmorrhage into the bladder and vesical catarrh. The former is a troublesome complication, as the blood-clots collect in the bladder; when there is marked hæmorrhage distending the bladder, the fistula must be opened up again. Sometimes the ureter has been caught in a stitch and compressed; intense pain, shooting from the kidney downwards along the course of the ureter,



FIG. 372.



FIG. 373.

SIMS' STATIONARY CATHETER: fig. 372, first model; fig. 373, newest model. That in fig. 372 is made of block tin so that it can be bent to any curve; when *in situ*, it must be bent so that the external end has its groove uppermost: that in fig. 373 is of rubber and has tubing attached to it.

with vomiting and other symptoms of uræmia followed but passed off on relaxing the sutures.

Removal of Sutures.

The sutures are removed on the tenth day. The method of removing sutures is shown at fig. 366.

For cases of fistulæ incurable by operation, a rubber urinal may be fitted on to an ordinary ring pessary²; or better still on to a spoon-shaped metallic collector.³

URETERIC FISTULA.

In cases of ureteric fistulæ, the end of the ureter next the bladder is tied, while that towards the kidney is implanted into the bladder. One end of a gum elastic catheter is passed into the ureter to define it and facilitate the stitching of it into the bladder. An elliptical incision is made in the bladder near the usual insertion of the ureter; or, if the ureter be short, at the nearest point. The catheter is now passed through this wound into the bladder and out through the urethra;

¹ As by W. A. Freund (*Sam. klin. Vorträge*, N. F., No. 118, 1895), and v. Rosthorn (*Cent. f. Gyn.*, 1896, S., 107).

² Jay—*Amer. Journ. Obstet.*, 1887, p. 50.

³ Bozeman—*ibid.*, 1892, i., 544.

and the ureter carried on it into the bladder wound into which it is fastened—the sutures which close the wound passing also through the wall of the ureter so as to fix it in the wound.¹

Witzel² describes a case in which he did abdominal section so as to




FIG. 374.

SIMON'S OPERATION FOR KOLPOKLEISIS. The patient is in the lithotomy posture; the sound has been passed through the urethra and fistula, and is seen in the upper portion of the vagina; the perineum is drawn back with the speculum and the labia majora with spatulae. A band-like piece of tissue has been removed from both the vaginal walls above the ostium; the raw surface is left unshaded in the figure. The vaginal mucous membrane is held tense by four pairs of forceps outside the raw surface, the shaded area within the latter is the upper third of the vagina. An end of the last suture has been passed through one raw surface, the second end is being carried through the other raw surface (*H. and K.*).

get down on the ureter extra-peritoneally. Then instead of dissecting it out, so as to bring it nearer to the bladder, he dissected off the

¹ This is Boldt's procedure (*op. cit.*). He got access to the bladder by abdominal section. This has also been done by the vaginal route. Boldt's paper describes various methods and gives the literature.

² Extra-peritoneale Uretero-cystostomie mit Schrägkanalbildung. *Cent. f. Gyn.*, 1896, S. 289.

the bladder so as to drag it over towards the ureter, and thus allow the latter to be introduced obliquely through its wall.

Closure of the Vagina: Kolpokleisis.

Where direct closure of the fistula is impossible, the only means for relieving the patient's discomfort is closure of the vagina below the fistulous opening. The portion of the vagina above this becomes, as it were, an extension of the bladder; the menstrual blood is discharged with the urine.

Vidal de Cassis, who originated this operation, performed it as follows. The inner surfaces of the labia majora were pared and brought together by sutures: the vulva was thus closed in an *antero-posterior* direction. After this operation, there always remained just below the urethral




FIG. 375.

SAME OPERATION AS SEEN IN SECTION TO SHOW RELATION OF RAW SURFACES (shaded dark), position of sutures and common receptacle above for urine and menstrual blood. The bladder and urethra are in upper part of figure (*H.* and *K.*).

orifice a small cleft through which the urine trickled. Unless complete continence is obtained, such an operation is useless.

Simon's
Kolpo-
kleisis.

Kolpokleisis is the name given to the operation introduced by Simon. It consists in obliteration of the vagina *transversely* by making a raw surface on its walls above the level of the ostium vaginæ. It is evident that this operation is justifiable only where closure of a fistula is impossible, either through the binding down of its margins to the bone with cicatricial tissue or through the complete destruction of the urethra. As the closure of the vagina interferes with married life, the nature of the operation should be explained to the patient beforehand, and full permission obtained.

The operation is performed as follows. By pinching up the mucous membrane, ascertain where it is most lax, so that the vaginal walls can be easily approximated; the point of closure should be as high up as possible. Mark out with the knife the ring of tissue to be excised. Lay hold of its lower margin and dissect it from below upwards; with

the finger in the rectum and the sound in the urethra, we can judge of the thickness of tissue to be removed (compare fig. 374 with fig. 375). On each ligature of wire or carbolised silk, two small curved needles are threaded so that *both* ends of the thread may be passed *from above downwards*. The needle must be entered into the vaginal mucous membrane above, carried through the substance of the vaginal wall (without appearing in the wound), and brought out through the vaginal mucous membrane below; it is difficult to prevent these sutures from catching up either bladder or rectum, but this should, if possible, be avoided. Care is required in the introduction of the first mesial suture as it is the guide for the others.

The results of this method are satisfactory as regards the production of complete continence. There is no liability to stagnation of urine or formation of concretions (*Hegar and Kaltenbach*). Hæmatometra will not occur unless there has been atresia of the cervix uteri. If menstruation has been in abeyance, it will probably return after the operation; in a case operated on by A. R. Simpson, the patient had not menstruated for a year, but a few weeks after the operation the menstrual blood appeared in the urine.

CHAPTER LV.

THE RECTUM; COCCYODYNIA.

LITERATURE.

Allingham—Diseases of the Rectum : Churchill, 1871. *Chadwick*—On the Functions of the Anal Sphincters : Am. Gyn. Trans., 1877. *Cripps*—Cancer of the Rectum : Churchill, 1880. *Hart*—Physics of the Rectum and Bladder : Edin. Obst. Trans., 1882. *Ruedinger*—Topographisch-chirurgische Anatomie des Menschen, Vierte Abtheilung : Stuttgart, 1873. *Storer*—the Rectum in its relation to Uterine Disease : Am. Jour. of Obst., Vol. i., p. 66. *Syme*—Diseases of the Rectum : Edin., 1859. *Van Buren*—Diseases of the Rectum : H. K. Lewis, 1881.

Not only is the gynecologist frequently consulted about rectal mischief, but as a matter of fact female patients sometimes refer rectal disease to the uterus or vagina ; therefore, in investigating gynecological cases, one has occasionally to satisfy one's self that the rectum is not the seat of the affection.

Vaginismus may be caused by fissure of the anus, as we have already seen, and pruritus vulvæ by ascarides from the rectum passing into the vagina.

PHYSIOLOGY OF THE RECTUM.

Relation
of Rectal,
Vaginal,
and Ureth-
ral Axes.

The anatomy of the rectum has been already considered (p. 36). The relations of the axes of rectum, anus, vagina and urethra, to one another and to intra-abdominal pressure are of importance. As we have already seen, the vagina and urethra are parallel to one another and to the plane of the brim.

Strictly speaking the surface whose outer boundary is the brim of the bony pelvis is not a plane surface, inasmuch as the various points in the outline of the brim are not on the same level. The vagina is thus, properly speaking, parallel to the internal conjugate of the brim.

The rectum runs, in part of its course, close behind the vagina for $1\frac{1}{2}$ inches and parallel to it ; the anal canal turns directly backwards so as to cut the vaginal axis at right angles. Intra-abdominal pressure acts at right angles to the vaginal walls, as can be noted from the fact that in defæcation the Hodge pessary is not driven out of the vagina.

Consideration of fig. 376 will show that the direction of intra-abdominal pressure on the pelvic floor coincides with the long axis of the anus, so that intra-abdominal pressure will act with its full driving force on any body in the anal canal.

The mechanism of defæcation is probably the following. According to Hilton, in his now classical book on "Rest and Pain," the lower part of the rectum is sensitive, but the upper two-thirds are but slightly so; the rest of the large intestine and the small intestine are non-sensitive. Hilton limits the sensitive portion to the lowest two inches of the rectum—to the part below the so-called sphincter tertius. When there is accumulation of fæcal matter in this portion, pain and uneasiness

Mechanism
of Defæca-
tion.




FIG. 376.

TO SHOW DIRECTION OF RECTUM AND OF ANUS IN RELATION TO INTRA-ABDOMINAL PRESSURE.
a uterus, *b* bladder, *d* vaginal orifice, *f* perineum.

produce the desire to expel these contents. There result the following reflex movements:—

- (1) Relaxation of the sphincter ani;
- (2) Peristaltic contraction of the circular unstriated muscle;
- (3) Shortening of the longitudinal muscle with eversion of the mucous membrane—since the longitudinal fibres have a fixed point below, their contraction will probably pull the rectum more into the line of the anal axis;
- (4) Contraction of the segments of the sphincter tertius.

In this way the lowest portion of the rectum becomes roofed in above by the sphincter *tertius* and open below. Intra-abdominal pressure drives this portion downwards; and the rectal contents, elongated by peristalsis and depressed by intra-abdominal pressure and eversion of the mucous membrane, are finally brought into the relaxed anal canal from which intra-abdominal pressure readily expels them. Ruedinger's diagram (fig. 35) shows well how the levator ani will reinvert the everted mucous membrane.

Inattention to the proper evacuation of the bowels leads to non-sensitiveness of the mucous membrane and is thus one factor in constipation.

EXAMINATION OF THE RECTUM.

This may be done in four ways:—

- (a) By finger (*v. p.* 114),
- (b) By speculum,
- (c) By eversion of the anterior rectal wall through digital pressure in the vagina (*Storer*),
- (d) By posture.

Specular
Examina-
tion of
Rectum.

By Speculum. The anal speculum has usually an oval fenestra; it is passed into the anus in the direction of its long axis, and rotated so that each portion of the anal lining comes opposite the aperture (fig. 378).

Storer's
Method.

Storer's method is as follows. Place the patient on her side; pass two fingers (or one) half way into the vagina, with the pulps of the fingers on the posterior vaginal wall. Then press these downwards and backwards, and thus evert the rectal mucous membrane through the dilatable sphincter ani which is at the same time pressed open with the fingers of the other hand. This method is most easily employed in multiparæ.

The rectum can also be thoroughly examined if the patient's hips be raised and a short cylindrical speculum introduced so as to admit air (*Kelly*).

DISEASES OF THE RECTUM.

Women are especially liable to rectal disease from the distention of parts accompanying parturition, as well as from their habitual neglect of the regular evacuation of the bowels. As rectal diseases often simulate those of the vagina, a sketch of the more important of them is necessary in a Manual of Gynecology. We shall therefore consider the following affections:—

- Displacements of the rectum,
- Fissure of the anus,
- Piles,
- Recto-vaginal fistula,
- Functional disturbance of rectum—constipation.

Displacements of the Rectum.

These are—Rectocele ;

Prolapsus recti (a) of mucous membrane,

(b) of whole thickness of bowel.

For *prolapsus recti*, which is properly surgical, see Van Buren or Prolapsus Recti. Allingham.

Rectocele is a protrusion of the lower part of the anterior wall of the rectum covered by the posterior vaginal wall, into the lumen of the vagina or even through the vaginal orifice. *Etiology.* There are two factors—tear of perineal body and pressure of scybala in rectum. *Diagnosis.* The posterior vaginal wall is seen protruding into the vagina or out at the vaginal orifice. The diagnosis is made by noting the




FIG. 377.

RECTOCELE (Schroeder).

relations of the protruded vaginal wall and by passing the finger through the anus into the pouch (fig. 377). *Treatment.* The patient should wear in the vagina a Hodge or Albert Smith pessary with cross bars; explain the necessity of a regular daily evacuation of the bowels.

Fissure of the Anus.

This is a crack, or ulceration, of the anal skin or of the mucous membrane covering the internal sphincter. In the edges of the crack there is usually a nerve filament, and below the crack lies the powerful sphincter ani. Fissure of Anus.

This apparently insignificant lesion gives rise in most cases to an unbearable and even incredible amount of pain, lasting for hours after

the bowels have moved. Hilton's explanation of this is so good that we give it entire.

Hilton's
explanation
of
pain in
Fissure.

"The reason for this anal ulcer being so very painful is the number of nerves associated with it; and the cause of the continued painful contraction which accompanies it lies in the enduring strength of the sphincter muscle. Thus it happens that exposure of those nervous sensory filaments upon the ulcer causes excito-motory or involuntary and spasmodic contraction of the sphincter, through the medium of the spinal marrow. The sphincter muscle contracts towards its own




FIG. 378.

ANUS *a* WITH ANAL SPECULUM *in situ*: it is turned so as to expose in the fenestra a fissure *b* beneath which a tenotomy knife has been passed (*Hilton*).

centre, and, as long as the muscle is in a state of contraction, it brings the sensitive edges of the ulcer into forced contact; this excites more muscular contraction, and thus by time and exercise, the muscle becomes hypertrophied, massive, and increased in dimensions."

Symptoms. *Symptoms.* The patient complains not so much of pain while the bowels are being moved as of an unbearable pain coming on after the evacuation and continuing for some hours. The pain is described as

unendurable, causing the patient to dread and postpone natural motions. There are often iliac pains and vaginismus; this last symptom is not infrequent.

Physical signs. By speculum or eversion, the crack is seen.

Treatment. Chloroform the patient, pass a tenotomy knife beneath Treatment the base of the ulcer (fig. 378) and cut upwards. This divides the muscular fibre so that the irritated edges can no longer be brought together. The fissure gets rest and heals readily; a cure is thus effected.

Another and very good plan is to chloroform the patient, and introducing the thumbs (with the dorsal surfaces in contact) to stretch the anus by forcibly separating them; this ruptures the muscular fibre and acts just as the knife does, and is especially good when the fissures are multiple.

The bowels are not to be moved for a day or two; the patient has then some pain when the motion is passing, but none after it.

Piles.

Hilton has pointed out that at the anus the line of demarcation between skin and mucous membrane is marked out distinctly by "the white line," as he terms it. This line is of great practical importance, as we shall see.

Piles are small tumours at the anus, on either side of this white line. They consist of dilated veins embedded in connective tissue and covered by skin or mucous membrane. We speak of external piles, *i.e.*, those outside of the white line and covered by skin, and internal piles, *i.e.*, those inside of the white line and covered by mucous membrane. Occasionally we have, as a special form of external pile, a dilated vein outside of the white line and usually containing a clot (venous pile).

Symptoms. Venous piles cause great pain; while external piles, Symptoms unless inflamed, occasion little inconvenience; from internal piles, there is bleeding when the bowels are moved.

Physical signs. The venous pile is a purplish tumour outside of the Signs. white line; external piles are like tags of skin, or are more or less distended; internal piles are cherry red and easily bleed.

Treatment. 1. When venous piles contain a clot, incise and turn out Treatment. clot.

2. For internal piles, employ the following *palliative* treatment. Give sulphur confection when necessary.

R Confectionis Sulphuris ʒij
Sig. Dessertspoonful at night.

Order gall and opium ointment to be applied.

R Unguenti Gallæ c̄ Opio ʒij
Sig. As directed.

For any abrasions, order boracic ointment or bismuth and cocaine suppositories.

The *radical* operative treatment belongs more to the surgeon.

Recto-vaginal Fistula.

The situation of such a fistula is shown in fig. 343. It may be due to carcinomatous or syphilitic ulceration, or to injury received during parturition.¹ The last only can be operated on. It is usually due to a tear, during labour involving the anus, and where the lower part of the laceration has united. The best treatment is to cut through the united portion and operate on it as if it were rupture of the perineum involving the anus.

Functional disturbance of Rectum—Constipation.

Women are usually exceedingly careless in the matter of regulation of the bowels; very often, evacuation is practised once a week or even at longer intervals. The main reason for this is want of exercise.

When consulted for constipation, the medical man should first inquire into the patient's habits as to exercise, diet, and sleep. When necessary, walking, golfing, tennis, or cycling should be advised unless there are any contraindications. The fact of the patient being a woman is no contraindication to any of these, but undue exercise should be avoided at the menstrual period. The aim of the patient should be to lead a natural life. The value of a daily evacuation at a fixed hour should be pointed out; this educates the bowels to demand it regularly. All quack pills should be tabooed as dangerous. The diet should be regulated; bran-bread, porridge and milk, stewed fruit, figs, etc., taken as part of food. The following pill is good.

R Extracti Nucis Vomicae
 Extracti Belladonnæ āā gr. $\frac{1}{4}$
 Pilulæ Colocynthis et Hyoscyami „ iij.
 Fiat pilula: mitte tales vj.

Sig. One occasionally.

The nux vomica and belladonna strengthen the peristalsis of the bowel: the colocynth and hyoscyamus pill is purgative; aloes and iron pill may be substituted for it.

The American drug cascara is very useful. We may give a pill of one to three grains thrice daily until the bowels move; twenty drops of the liquid extract may be taken instead.

R Extracti Cascarae Sagradae gr. iii.
 Pulv. Glycyrrh Co. q.s.
 Fiat pilula: mitte tales xij.

Sig. One thrice daily.

¹ It has been described as a congenital condition—*Torek—Cent. f. Gyn.* 1895 S. 328.

R Extracti Cascaræ Sagradæ Liquidi ℥ij.
Sig. Twenty drops thrice daily.

This drug is tonic to the bowel: its use should be stopped when once the bowels begin to act. It should not be given until the diet is regulated. The pill is more convenient, as the liquid extract is bitter.

The purgative mineral waters are very useful. The best are the Friedrichshall, Hunyadi Janos, Apenta, and Aesculap. The patient should take in the morning a wine-glassful or half-tumblerful with an equal amount of hot water; the taste may be masked by the juice of a lemon with sugar. Rubinat is a strong and active aperient in doses of one ounce. The Carlsbad salts are good and may be used as already directed (p. 369). Very often an enema of cold water is helpful. The medical man should deprecate the habitual use of purgatives, and insist on natural and daily evacuation as the result of judicious diet, work, and exercise.

The aloes and iron pill is good in sluggishness of the lower bowel. Rhubarb is bad as a habitual purgative, owing to its tendency to constipate after purging; the well-known "Gregory's Mixture" should not be used as a habitual purgative, but is good in diarrhœa inasmuch as it first purges and then binds. Fluid magnesia, castor oil, and some of the milder salines (*e.g.*, the easily-taken Seidlitz powder) may be employed. Blue pill should be avoided; euonymin or iridin are better hepatic stimulants (*v.* p. 622).

The injection of pure glycerine (℥j-℥j) into the rectum ensures an evacuation of the lower bowel in a few minutes; and is therefore convenient in certain cases. Suppositories made up in large part of glycerine can also be employed. A small syringe is required for the injection of the fluid glycerine. Abdominal massage is often beneficial.

COCCYGDYNIA.

LITERATURE. *Hildebrandt*—Die Krankheiten der äusseren weiblichen Genitalien, S. 127: Stuttgart, 1877. *Nott*—N. O. Medical Journal, May 1844. *Simpson, Sir J. Y.*—Diseases of Women, p. 202: Edinburgh, 1872. *Thomas*—Diseases of Women, p. 151: London, 1880.

By this we understand a painful condition in the region of the coccyx induced by sitting, walking, and the various muscular contractions associated with defæcation and coitus. When we consider the anatomy of the coccyx, its muscular attachments (to the levator ani, coccygeus, external sphincter ani, and gluteal muscles), as well as the strain put on it when driven back during parturition, we are not astonished that in some cases there should be inflammatory changes around and in it causing pain in its movement.

Symptoms. The chief symptom is pain on sitting, walking, and defæcation.

Physical signs. By digital pressure on the coccyx and examination per rectum, the seat and nature of the pains are made out.

Treatment. (1) Massage and manipulation of the coccyx should be tried first. (2) Pass a tenotomy knife beneath the skin on the posterior aspect of the coccyx, and free its lateral and apical muscular attachments; or (3) amputate the coccyx. To do the latter, make a vertical mesial incision over the posterior aspect of the coccyx; seize its tip and pull it well back; then free its muscular attachments with the knife, keeping close to the bone; finally separate it at the sacro-coccygeal joint. (2) and (3) are rarely necessary.

APPENDIX.

ABDOMINAL SECTION: ANTERIOR AND POSTERIOR COLPOTOMY: VAGINAL HYSTERECTOMY.

LITERATURE.

Barbour—Operative Treatment of Advanced Extra-uterine Gestation : Edin. Obst. Trans., 1896-97. *Donald, Arch.*—A Case of Vaginal Cœliotomy for Tubal Pregnancy : Brit. Med. Journ., 1896, i., p. 79. *Dührssen*—Ueber eine neue Methode der Laparotomia (Köliotomie vaginale): Berl. Med. Woch., 1894, No. 4: v. also Zts. für Geb. und Gynäk., Bd. xxviii., S. 401. *Hart*—The Minute Anatomy of the Placenta in Extra-uterine Gestation : Ed. Med. Journ., Oct. 1889. *Hart and Carter*—The Sectional Anatomy of Advanced Extra-uterine Gestation : Edin. Med. Journ., Oct. 1887. And Laboratory Reports, R.C.P.E. : Edin., 1889. *Jessop*—Case of Extra-uterine Gestation ; removal of living foetus, etc. : Lond. Obstet. Trans., 1876, p. 261. *Keith*—Surgical Treatment of Tumours of the Abdomen : Edin., 1885. *Kocher*—Text-book of Operative Surgery, p. 126 (Stiles' Trans.) : London, A. & C. Black, 1895. *La Torre*—Quel est le meilleur mode de fermeture de l'abdomen : Paris, Levé, 1897. *Langenbuch*—International Medical Congress, London, 1881, Vol. ii., p. 278. *Lister*—On Corrosive Sublimate as a Surgical Dressing : Lancet, 1884, p. 723. *Macdonald*—Record of Cases treated in Ward XXVIII., Royal Infirmary, Edinburgh, Nov. 1883 to April 1884 : Edin. Obstet. Trans., Vol. ix., p. 134 ; same from May to November 1884 : *ibid.*, x., p. 178. *Martin, A.*—Die Colpotomia anterior : Monats. für Geb. und Gynäk., Bd. ii., Hft. 2, 1895. See also Brit. Med. Journ., 1896, i., p. 10. *Maygrier*—Terminaison et Traitement de la Grossesse extra-uterine : Paris, 1886. *Morris*—Surgical Diseases of the Kidney : London, Cassell & Co., 1885. *Robson, Mayo*—Diseases of the Gall-bladder and Bile-ducts : Clifford Allbutt's System of Medicine, London, 1896. *Tait, Lawson*—Diseases of the Ovaries : Birmingham, Cornish Brothers, 1883. Ectopic Pregnancy and Pelvic Hæmatocele : Birmingham, 1888. The Pathology and Treatment of Extra-uterine Gestation : Brit. Med. Journ., 1884, p. 317. *Taylor, J. W.*—On the Opening of the Abdomen from the Vagina, etc. : Brit. Med. Journ., 1896, i., p. 78. *Thornton, J. Knowsley*—Cases of Hysterectomy, etc. : Brit. Med. Journ., May 23, 1885. *Treves, F.*—Intestinal Obstruction : London, Cassell & Co., 1884. *Wells, Sir Spencer*—The Diagnosis of Surgical Treatment of Abdominal Tumours : London, Churchill, 1885. See also Literature of Operative Treatment of Ovarian Tumours, Chap. XXIV. ; of Fibroid Tumours, Chap. XXXVII. ; and of Carcinoma Uteri, Chap. XLII.

SINCE the battle won by Atlee, Keith, and Wells in regard to ovariectomy, great changes have been made as to the routes by which the abdominal cavity can be reached. Abdominal section is the operation indicated in the great majority of instances, but the other routes are also valuable, and in some cases better than it.

We have therefore now to sum up briefly the indications for and methods of performing abdominal section, anterior and posterior colpotomy, and vaginal hysterectomy.

ABDOMINAL SECTION (CÆLIOTOMY).

In the preceding pages, operations necessitating abdominal section, viz., those for abdominal and pelvic tumours, have been described; but it is here intended to gather up consecutively and briefly the main points necessary for the successful performance of Abdominal Section so as to give the operator or his assistant a bird's-eye view of the whole subject, and enable him to meet unexpected emergencies such as often arise even after the utmost care has been taken to avoid mistakes in diagnosis.

Preliminaries. The operation is best performed in the special wards of an hospital or in a private hospital in the case of well-to-do patients. The houses of the poor are quite unfitted for operations; and it is much better for wealthy patients to be under the discipline of a good private hospital and away from the well-meaning but hurtful interference of relatives. It also relieves the operator of the anxieties attendant on their misinterpretation of symptoms.

Prior to any operation the patient's systems should be examined, especially lungs, heart, and kidneys. Ether is better not employed when there is a tendency to bronchitis; and the amount of urine should be noted, the usual tests for albumen and sugar employed, and microscopical examination made of its deposit. The urine is sometimes scanty in cases of large tumours, and therefore some diuretic such as acetate or citrate of potash should be given.

The pulse and temperature should also be taken twice daily for a few days prior to operation, and the patient should, if necessary, be kept in bed for that time, so as to accustom her to it.

The importance of having a specially trained nurse cannot be over-rated. She is required to take the pulse and temperature, and to keep a register of these: to draw off the urine when necessary, and to be capable of giving ordinary and nutritive enemata. She must therefore have good hands, be firm and yet gentle, one who carries out instructions to the letter, and who is thoroughly imbued with the spirit of cleanliness.

ANTISEPTICS.

The operation is to be carried out in the spirit of Listerism. The operator strives to have pure surroundings and everything that touches the part operated on aseptic, either by antiseptics or sterilisation. He must therefore consider means of purifying the air, instru-

ments, sponges, skin of patient adjacent to part operated on, and discharges from wounds.

Purification of the air. This is to be got by ventilation, previous purification of the room by sulphur or chlorine fumigation, and preliminary spraying of carbolic lotion into the air of the apartment. The operator, however, attaches the greatest importance to the absolute asepticity of everything that touches the wound—fingers, knives, and (above all) sponges.

Instruments are readily purified by being boiled in soda solution (1 per cent.). During the operation they should lie in shallow porcelain trays of 1-40 carbolic lotion.

Sponges. This is the part of the operative equipment which requires most careful attention. The utmost cleanliness and purification of sponges is a *sine quâ non* to success. Care must be taken that they do not become friable, and the operator should give them his personal attention.

As an exemplar of what is required, we give Lawson Tait's precautions in regard to them.

"New sponges are first put into a large quantity of water with sufficient muriatic acid to make the water taste disagreeably acid. They remain in this mixture until all cleaning effervescence has ceased and all the chalk is removed. For this purpose it may be necessary to renew the acid several times. The Sponges are afterwards carefully and thoroughly washed to make them as clean as possible and free from every rough particle. After being used at an operation they are first washed free from blood, and then put in a deep jar and covered with soda and water (1 lb. of soda to twelve sponges). They are left in this about twenty-four hours (or longer if the sponges are very dirty), and then they are washed perfectly free from every trace of soda. This takes several hours' hard work, using hot water, squeezing the sponges in and out of the water, and changing the water constantly. Leaving them to soak occasionally for a few hours in very hot water greatly assists in the cleansing. When quite clean they are put into a jar of fresh water containing about one per cent. of carbolic acid, and after being in this for twenty-four hours they are squeezed dry and tied up in a white cotton bag, in which they are left hanging from the kitchen ceiling (being the driest place in the house) till they are wanted."

Prior to an operation they should be carefully washed in very hot water and soaked over night in carbolic lotion (1-20).

They are wrung out of 1-40 for the operation and placed near the operator in a suitably warmed dish.

Sterilised tampons or gauze may be used instead of sponges (*v. p.* 166). Gauze and tampons should only be used on forceps: *this prevents their being left in the abdomen.*

The skin near the part to be operated on should be washed the night before the operation with turpentine, soap, and water. The umbilicus is to be carefully cleansed. A cloth soaked in carbolic lotion (1-40) and covered with pink mackintosh should be left on during the night. When the patient is under chloroform, the skin is again washed with corrosive sublimate (1-2000) and the pubes shaved.

The operator's hands and arms are to be cleansed with turpentine, soap, and water, the nails brushed, and all finally washed with corrosive sublimate (1-2000). One absolute rule is that only the operator or the special assistant should touch the wound, sponges, and instruments.

THE ABDOMINAL INCISION.

This is either *mesial* or *lateral*. The *mesial* incision is the usual one and may vary in length.

For an exploratory incision, 2 inches is sufficient, and this is also, as a rule, enough for the removal of the uterine appendages in the pelvis. Its lower end is 1 inch above the symphysis pubis, but must be higher when removing the uterine appendages in an abdominal fibroid.

For ovariectomy, an incision of 3 to 4 inches in length is usually required.

For large solid tumours, the incision may be very long.

If the first incision into the abdominal cavity is found too short, it can easily be enlarged up and down with straight probe-pointed scissors guided on the finger passed in.

The operator cuts down through the skin and abdominal fat to the aponeurosis. Beneath the aponeurosis is the extra-peritoneal fat and then the peritoneum. A good plan is to lay hold of the structures beneath the aponeurosis with two pairs of Péan's forceps, each one catching a little to the side of the mesial line. In this way a fold is pinched up, running across the middle line at right angles to it: this can be cut without danger to subjacent structures, and the same manœuvre repeated on deeper structures.

The *lateral incision* of Langenbuch is to be recommended in renal tumours. It is made at the outer margin of the rectus abdominis with its centre at the level of the umbilicus, and is advantageous inasmuch as the operator reaches the outer layer of the meso-colon, thus avoiding the blood-vessels running in the inner layer.

Kocher objects to this incision, that it cuts the intercostal nerve branches and leads to a paralysis which may predispose to ventral hernia. He favours instead oblique or transverse incisions when the usual mesial one does not suffice.

EXPLORATION OF ABDOMEN OR PELVIS AND REMOVAL OF TUMOURS.

When the abdominal cavity is opened, the operator either explores in doubtful cases, or removes the tumour he has already diagnosed.

While exploring, the deep anæsthetisation of the patient removes all straining of the abdominal muscles. The operator may find that he has to deal with a malignant case, or with a tumour not removable.

He must then close the incision. One good rule in doubtful cases is not to meddle unless there is a fair chance of finishing the case. It is always unwise for the operator, and highly dangerous to the patient, to nibble, as it were, at a case. There is little or no risk in mere exploratory incision, and there is much more risk in an incomplete operation than in a severe completed one.

The removable tumours or conditions admitting treatment are—

- | | | |
|------------|---|--|
| Abdominal. | { | (1) Ovarian, parovarian, and broad-ligament tumours, |
| | | (2) Fibroid, |
| | | (3) Fibro-cystic, |
| | | (4) Splenic, |
| | | (5) Omental, |
| | | (6) Renal, |
| | | (7) Hydatid, |
| | | (8) Mesenteric, |
| | | (9) Pancreatic, |
| | | (10) Distended gall bladder, |
| Pelvic. | { | (11) Uterine appendages in cases of fibroids, |
| | | (12) Uterine appendages diseased (pyosalpinx, cirrhotic or prolapsed and painful ovaries), |
| | | (13) Pelvic abscess, |
| | | (14) Extra-uterine gestation. |

(1) *Ovarian, parovarian, etc.* The removal of these by Abdominal Section has already been fully described under Ovariectomy, in Chap. XXIV. The operation for a *pediculated* tumour may thus be briefly summarised. The operator taps the tumour, withdraws it from the abdomen, and ties the pedicle with the Staffordshire or the ordinary knot. In certain cases, the clamp and cautery can be employed. The tumour is now cut away: the pedicle whether ligatured or cauterised is dropped back (complete intra-peritoneal treatment) and the abdominal incision closed.

When the tumour (usually papillomatous) has developed between the layers of the broad ligament or beneath the peritoneum and is *not pediculated*, its removal is a much more difficult matter. The best plan is to tap first, then to incise the peritoneum and enucleate the tumour. The part first enucleated with the finger is laid hold of with forceps, drawn well up, and then the operator separates further with his finger, seizing bleeding points with Péan's forceps and tying with catgut. Care must be taken at the side walls of the pelvis not to damage the ureter, as well as at the region of the sacro-iliac joints where the large iliac veins with their many branches lie. The part from which the tumour has been enucleated should be drained if necessary.

(2) (3) *Fibroid and Fibro-cystic.* For full details of Hysterectomy for

Removable
Abdominal
Tumours.

Fibroids, see pp. 454-464. The tumour is turned out of the abdomen through a large incision, clamped, and then cut off. The pedicle is usually treated extra-peritoneally. Many operators prefer pan-hysterectomy (p. 462).

(4) *Splenic*. Cystic splenic tumours have been removed successfully. Severe shock may develop while the pedicle is being tied, and there is great risk afterwards of the ligature slipping. In leucocythæmic cases the spleen should not be removed.

(6) *Renal*. After incising the abdominal walls by Langenbuch's incision,¹ the outer layer of the meso-colon is opened, the tumour enucleated until the hilum is reached, when a silk ligature is applied as in the ovariectomy pedicle. The pedicle is now cut on the tumour side of the ligature and the tumour removed.

The pedicle is now dropped back into the extra-peritoneal space, the peritoneum laid over it and pressed down with the sponge. Thornton recommends that the ureter end be secured in the abdominal incision.

(7) (8) *Hydatids or Mesenteric tumours* are opened, the contents evacuated, and the incision into them stitched to the abdominal wound.

(10) *Distended gall bladder*. The gall bladder when distended owing to obstruction by gall stones, has been opened, the calculi removed (recommended by Jean Louis Petit, Handfeld Jones, and carried into execution by Marion Sims, and especially Lawson Tait). Tait, in one of his cases, made an incision 4 inches in length, in the middle line, with the umbilicus in the centre of the incision. The gall bladder was aspirated after the abdomen was opened, and then cut into at that point; the gall stones were extracted, the opening in the gall bladder stitched to the abdominal wound, and the rest of the wound closed in the usual way. Bile oozed from the wound for some days, but the patient made an excellent recovery.²

(11) *Uterine appendages in case of Fibroids*. When a fibroid is not too large and is growing rapidly or causing exhausting hæmorrhages, the appendages should be removed. A two-inch incision is made through the abdominal wall and the ovary and Fallopian tube on either side brought up to it. The ovary and part of the Fallopian tube are looped up, tied with the ordinary or the Staffordshire knot, and the parts outside the ligature cut off. A larger incision is often required, and in certain cases the ovaries may not be accessible. The operator must then remove the uterus.

(12) *Uterine appendages diseased (pyosalpinx, cirrhotic or prolapsed and painful ovaries)*. The uterine appendages when diseased and causing serious indisposition may be removed. This is not by any means to be

¹ On this subject the student may read Morris' *Surgical Diseases of the Kidney* (London 1885), and also Czerny's paper "Ueber Nierenextirpation," with discussion in the *International Congress Transactions*: London, 1880, Vol. ii., p. 242.

² See also Mayo Robson's article.

done lightly, its exact results as to sterility have to be explained, and the operator should never force it on the patient.

In *Pyosalpinx* the operator first brings up the tube, freeing adhesions with his fingers, ligatures as large a loop as possible and cuts away above. Great care is to be taken to prevent any pus entering the abdomen. This is best done by pressing sponges below the freed tube. Any hæmorrhage is arrested by pressure, ligature, hot water, or by the actual cautery. Most operators prefer to separate adhesions before tapping, and it is better not to tap unless necessary. Should the tube rupture during this, the extravasated contents must be most carefully sponged out and the pelvis thoroughly flushed with hot water.

In difficult cases the abdominal incision should be enlarged and the hand passed in if necessary. The intestines can be kept out of the way by a sponge and long forceps. It may be necessary to turn the intestines out temporarily into hot towels, and the value of the Trendelenburg position in such cases must be kept in mind.

(13) *Pelvic abscess* may be treated by abdominal section when it rises up so as to be near the abdominal walls. After the usual incision through the walls, the operator taps the swelling, then draws up the collapsed walls of the cavity, enlarges the opening, and stitches it with silk to the abdominal wall, the rest of the abdominal incision being closed as usual. A glass drainage tube is passed into the abscess cavity, but the peritoneal cavity is accurately closed. The perforations in the tube should not be above the level of the peritoneum.

(14) *Extra-uterine gestation* may be met with in very many forms:—

Forms of
Extra-
Uterine
Gestation.

- | | | |
|-----------|---|--|
| Early. | { | (a) Entire, small, and still in Fallopian tube; |
| | | (b) Ruptured into the peritoneal cavity, which contains much blood and a small foetus; |
| | | (c) Ruptured through the part of the Fallopian tube bounded by the broad ligament, and developing there; |
| Advanced. | { | (d) Both foetus and placenta near full time but lying in extra-peritoneal tissue; |
| | | (e) Foetus in peritoneal cavity with placenta in extraperitoneal tissue; |
| | | (f) Foetus and placenta in extraperitoneal tissue but suppuration going on and termination as in pelvic abscess; |
| | | (g) In a detached horn. |

(a) *Entire, small, and still in Fallopian tube.* Here the operator tries to remove the entire sac by ligature with silk and cutting away above it.

(b) *Ruptured into the peritoneal cavity which contains much blood and a small foetus.* Such cases may be saved by Abdominal Section. Tait has

recorded no fewer than forty-three cases where he has operated for this with only one death.

In a recent case of abdominal section we found the pelvis filled with tarry-like blood, a small foetus in the abdomen, and a rupture in the Fallopian tube about the size of the tip of the index finger. The foetus was removed, a loop of the tube with the rupture on it secured with the Staffordshire knot, the pelvis sponged and then washed out with hot water (120° F.), to check oozing. It was noted at the time that the omentum became blanched; the water was passed in only for a few seconds and then sponged out. Uninterrupted recovery took place.

(c) *Ruptured through the part of the Fallopian tube bounded by the broad ligament, and developing there.* This gives a complex case not good for abdominal section. The operator's aim should be to open the sac and remove the foetus without disturbing the placenta. The sac may be tamponed with iodoform gauze for several days until the placenta is thrombosed and can be safely removed. The operator may be able to open the sac extra-peritoneally. In all extra-uterine gestation, indeed, it is absolutely imperative to avoid removing the placenta at first, as there is no arrangement of muscular fibre to check hæmorrhage as in normal labour. The cut edge of the sac is to be stitched to the abdominal wound which is left large and a drainage tube inserted.

In a case observed by us the placenta had grown after the death of the foetus; the foetus was very much compressed and any attempt to remove it by abdominal section would have caused fatal hæmorrhage by separating the placenta.

(d) In this form a lateral incision may be employed and access gained without opening the peritoneal cavity. The foetus can be removed and the placenta left.

(e) As in (d) except that the peritoneal cavity is opened by a mesial incision.

(f) Is to be treated as in pelvic abscess.

(g) *Gestation in a detached horn.* This is a very rare condition and is of interest chiefly because of its close resemblance to a fibroid (v. p. 296). It is removed and clamped just like a fibroid.

POSSIBLE ACCIDENTS DURING LAPAROTOMY.

The accidents which may happen during Laparotomy are usually, though not always, due to the non-observance of the rules now laid down by successful operators, and should not occur when these are followed. They may be thus summed up.

- (1) Leaving sponges or instruments in the abdomen,
- (2) Wound of small intestine,
- (3) Injury to tip of vermiform appendix,
- (4) Injury to ureter,
- (5) Injury of iliac veins,
- (6) Tears into bladder or rectum.

Sponges or instruments will not be left in the abdomen if they are carefully counted, and the former never torn up during an operation. A fatal result may follow if such foreign bodies are left, although cases have been recorded where they have been removed on the following day, or even been discharged many days after, the patient recovering; in the last cases they have set up abscesses escaping by the bladder or wound.

Wound of the small intestine should be stitched as follows. First stitch mucous membrane to mucous membrane with catgut and then peritoneum to peritoneum by Lembert's suture. The material to be used for the peritoneum is the finest Chinese twist, passed with a curved needle.¹

PERITONEAL TOILETTE; CLOSURE OF WOUND.

The *peritoneal toilette* must be performed most carefully. All bleeding points are to be arrested and all fluids are to be sponged out thoroughly. The pelvis or abdominal cavity if necessary may be washed out with warm water. The peritoneum should be made thoroughly dry before the wound is closed. Careful peritoneal toilette with scrupulous asepsis is the key to success.

The abdominal wound may be *closed* with silk or catgut. Silk is very good and the stitches may be passed as in an ordinary wound. They should not be far apart (half-an-inch or so between each), and should include the whole thickness of the abdominal walls. The skin if necessary may be more accurately approximated by superficial horsehair stitches.

The abdominal incision may be closed entirely with catgut as follows: Use Hartmann's catgut, No. 4, and needle on handle. Pass the unthreaded needle through both sides of wound, beginning at its lower end, and including only the muscle and peritoneum; thread the catgut, pull back and tie. Pass the needle again and thread one of the tied ends, pull this back and tie. Do this, in all, three times at lower end of incision and cut ends short. Do same at upper end, but leave ends long. Now withdraw sponge and carry out same method in middle third of wound. Finally, tie long ends of upper and middle thirds of incision. Unite skin edges with continuous catgut suture.

Some operators unite the peritoneal edges with catgut and then use silk for muscle and skin.

ANTERIOR AND POSTERIOR COLPOTOMY: VAGINAL HYSTERECTOMY.

Anterior Colpotomy. While vaginofixation has not as yet commended itself as either a satisfactory or a scientific operation, it has certainly

¹ See Treves' "Intestinal Obstruction."

demonstrated that the abdominal cavity can be reached by an incision from the vagina through the loose tissue separating the bladder and cervix uteri, and opening the vesical pouch of peritoneum. We are indebted to Dührssen of Berlin for this operation—"anterior colpotomy" or "vaginal cœliotomy." Martin of Berlin has among others also strongly recommended its employment in suitable cases.

Martin's method of operating is briefly as follows. The patient is placed in the lithotomy posture and the parts cleansed as usual. A modified Sims' speculum with a short broad blade is then introduced into the vagina, and the anterior lip of the cervix uteri seized with a volsella and drawn down. A convenient form of instrument is Orthmanu's forceps, which combines a uterine sound and volsella, and thus defines the uterus during the operation. The anterior vaginal wall is also laid hold of in the middle line with an additional pair of forceps at a point about 3 inches from the external os.

A vertical mesial incision is now made until the loose tissue between bladder and vaginal wall and cervix is reached. This is separated laterally, and the bladder is then pushed up with the finger until the vesico-uterine fold of peritoneum is reached; this is then incised, and the peritoneal cavity thus opened.

The body of the uterus can now be brought into the vagina, and with it the appendages, so that the necessary operation can be performed.

The indications are (1) inflammatory conditions of tubes and ovaries, small cystic ovaries; (2) pyosalpinx and hydrosalpinx, tubal gestation; (3) small fibroids, which can be enucleated by an incision through the anterior wall, or morcellement can be practised. The tumour should not be larger than two fists.

After the necessary operation has been performed, the uterus should be replaced through the opening, and the incision closed with catgut as follows. Pass an interrupted catgut suture (No. 4, Hartmann) through the upper vaginal edge, the loose tissue at base of bladder, peritoneum, anterior uterine wall near fundus, and then back through corresponding edge on opposite side: tie. Similar transverse sutures are passed at a lower level. The continuous sutures may be used in a similar way, either alone or combined with the interrupted. Prior to the operation curetting may be done if necessary.

The results of this operation are excellent, the mortality is small, and the disturbance to the patient less than in abdominal section. In 390 cases Martin has had four deaths.

Posterior Colpotomy. Here the operator opens into the pouch of Douglas, and removes ovaries, etc., as in anterior colpotomy. The student will understand from what has gone before how this is done.

This operation has long been practised (Battey, Goodell, Byford), but has not gained much acceptance.

In *Vaginal Hysterectomy* by forci-pressure we have a means of access due to Péan, and greatly developed by Richelot, Doyen, Segond, the brothers Landau, Jacob, and many others. The procedure has already been sufficiently described (v. p. 516). By this method we can remove the uterus affected by cancer or fibroids, and often actually diseased in inflammatory conditions of the appendages, and at the same time gain access to the peritoneal cavity for the treatment of appendage mischief. The operation is thus more thorough, if more severe.

To sum up, then, we have in abdominal section the most universally applicable method of operation for intraperitoneal conditions, and one by which all forms of intraperitoneal disease can be treated. In large tumours it is practically imperative. Anterior colpotomy has evident advantages in the case of small tumours, ovarian and uterine, and in inflammatory appendage disease; while vaginal hysterectomy has its own rôle in cases of cellulitic fixation of uterus, with or without suppuration, intractable chronic endometritic conditions, and those where we have combined uterine and appendage mischief.

ELECTRICITY IN GYNECOLOGY: THE APOSTOLI METHOD OF TREATMENT.

HISTORY.—Apostoli tells us that he studied the surgical employment of electricity at the *Clinique* of Dr A. Tripiér, whose memoir to the Academy of Science in Paris, on Faradisation in the Treatment of Hypertrophies of the Uterus,¹ opened up the way. Apostoli saw the weak points of Tripiér's practice: among others, that the currents employed were too feeble, their intensity not regulated and measured, the point of application wrongly chosen, and the different effects of the Faradic and the Galvanic (or Voltaic) currents, as well as of the positive and the negative poles not distinguished. He began to work out his own ideas in 1882; and in 1883, he described his electric treatment of Perimetritis, reading a paper on that subject at the Congress of Copenhagen in 1884.² In this same year (1884) he laid a memoir on the subject of treatment of Fibroid Tumours of the Uterus by Electricity before the Academy of Medicine of Paris; the subject, as already mentioned (p. 427), of his paper read at the Dublin meeting of the British Medical Association in 1887. It was also in 1887 that he published a book on the *Electric Treatment of Chronic Metritis and Endometritis*.³

History
of the
Apostoli
Method.

Apostoli's subsequent papers and the more recent literature have been already given in describing the electrical treatment of Fibroid Tumours.⁴ The best English article on Electricity in relation to the Diseases of Women is R. Milne Murray's monograph in Clifford Allbutt and Playfair's *System of Gynecology*, which contains a bibliography.

NOTE ON ELECTRICAL TERMS USED.—In order to make clear the description of Apostoli's method which follows, it will be well first to

¹ Hyperplasies conjonctives des organes contractiles de l'emploi de la faradisation dans le traitement des engorgements et déviations de l'utérus et de l'hypertrophie prostatique: *Comptes Rendus de l'Académie des Sciences*, Août 1859. Leçons de clinique sur les maladies des femmes: Paris, Octave Doin, 1883.

² Sur un nouveau traitement des périmétrites: *Comptes Rendus du Congrès de Copenhague*, Section d'Obstétrique et de Gynécologie, p. 141.

³ Sur un nouveau traitement de la métrite chronique, et en particulier de l'Endométrite, par le Galvano-caustique chimique intra-utérine: Paris, Octave Doin, 1887.

⁴ See p. 449.

explain some of the terms used, so that students may read straight on without the interruption of consulting books on electricity which may not be at hand at the time.

Kinds of
Electric
Current.

In the first place, there are two distinct kinds of electric current spoken of, the *Galvanic* (perhaps more accurately the "Voltaic") and the *Faradic*. The former is the electricity that flows in continuous current through the wires from the zinc and copper plates in a voltaic or galvanic cell or battery when their ends are connected. As sulphuric or other oxidising acid is added to the water in the cell, this kind of current is *chemical* in its origin. When the current flows, the zinc plate is used up, its consumption furnishing the energy to drive the current through the cell and connecting wire: the cell, in fact, has been aptly compared to a sort of chemical furnace in which the fuel is zinc. The faradic current, on the other hand, is an *induction* one, *i.e.*, is a current induced in a closed circuit when a magnet is moved near it or when it is moved across the magnetic field, or when an electric current whose strength is changing is near it. The source of this current is, accordingly, not chemical but electro-magnetic.

Electro-
motive
Force and
Strength
of Current.

That which tends to produce a current, *i.e.*, to move electricity from one place to another, is called *Electro-motive force*; the *Strength of a Current* is the quantity of electricity which flows past any point of the circuit in one second, and is directly proportional to the electro-motive force, and inversely proportional to the resistance which the current has to overcome in its flow. This truth with regard to the strength of an electric current flowing in a circuit is, from the name of its discoverer, known as *Ohm's Law*, which may be formally stated here—"The strength of the current varies directly as the electro-motive force, and inversely as the resistance of the circuit." The terms "strong," "great," and "intense," applied to currents all mean the same thing.

Measure-
ment of
Strength
of Electric
Currents.

To measure the strength of electric currents there is used an instrument called the *Galvanometer*, in which a magnetised needle is deflected by a current passing above and below it through a coil of silk-covered insulated copper wire—the amount of deflection depends upon the strength of the current (though not proportional to it) and a properly graduated dial enables us to ascertain perfectly the strength of the current. The sensitiveness of the instrument is greatly increased by the use of the *astatic* needle, a compound one in which the directive power of the earth is neutralised by the joining of two magnetised needles of equal power connected one above the other by a central pin so that the north pole of the one lies over the south pole of the other, and the south pole over the north pole of the other. The sensitiveness is also increased within certain limits by increasing the number of turns of the coil of silk-covered wire. A galvanometer must be able to measure the quantity of electricity passed, and should be of a degree

of sensitiveness corresponding to the strength of the current to be measured—very sensitive for very small currents, less sensitive for strong currents.

UNITS OF MEASUREMENT.—Every kind of measurement requires a unit: as in measuring length we might take the inch, foot, yard, or mile; and in measuring mass or weight we use the grain, ounce, pound, hundred-weight, or ton. Accordingly, for measuring electricity, we have in the first place a series of what are called *absolute* electric units derived from the fundamental Centimetre-Gramme-Second system (C.G.S.) in which—

The *Centimetre* (.3937 in.) is the unit of length,
The *Gramme* (15.432 grns.) is the unit of mass, and
The *Second* is the unit of time.

There are three *derived* units which it is necessary to bear in mind in order to understand the electric units which follow. These are—

The *Dyne* or unit of force, that force which acting for one second on a mass of one gramme gives to it a velocity of one centimetre per second;

The *Erg* or unit of work, the work done in overcoming unit force through unit distance, *i.e.*, in moving a mass through a distance of one centimetre against the force of a dyne; and

Unit Strength of Magnetic Pole.—The unit magnetic pole is of such a strength that when placed at a distance of 1 cm. in air from a similar pole of equal strength it repels it with a force of one dyne.

We are now in a position to understand the definition of the units referred to in the explanation of Apostoli's method. As that method deals with Current Electricity in which the positive and negative poles are in properties the same as magnetic ones, these units are called Electro-magnetic.

Electro-magnetic Absolute Units.—(1) *Unit Strength of Current* is that of a current such that if one centimetre length of its circuit be bent into an arc of one centimetre radius it will exert a force of one dyne on a unit magnet pole placed at the centre of the circle of which the arc is a part, so as to be always a centimetre away from the current.

(2) *Unit Quantity of Electricity*, that quantity of electricity which is conveyed by current of unit strength in one second.

(3) *Unit of Difference of Potential or of Electro-motive Force* exists between two points when it requires the expenditure of one unit of work (Erg) to bring a unit of + electricity from one point to the other against the electric force.

(4) *Unit of Resistance* is possessed by a conductor when unit difference of potential between its ends causes a current of one unit of quantity per second to flow through it.

The first two of these absolute units were found to be inconveniently small and the last two inconveniently large, accordingly a committee of the British Association devised a system of "practical" units in which they substitute for the fundamental units *centimetre* and *gramme*, the Earth's quadrant (1,000,000,000 centimetres) and $\frac{1}{100,000,000,000}$ of a *gramme*.

Electro-magnetic Practical Units.—(1) The *Volt*¹ is the practical unit of *Electro-motive force* and is 100,000,000 absolute units.

(2) The *Ohm*¹ is the practical unit of *Resistance*, and is 1,000,000,000 absolute units.

(3) The *Ampère*,¹ the practical unit of *Strength of Current*, is that furnished by a Volt through an Ohm and is $\frac{1}{10}$ of the absolute unit. In medical electricity, however, the strength of the current is measured in milliamperes.

(4) The *Coulomb*¹ is the practical unit of *Quantity* of current electricity, and is $\frac{1}{10}$ of the absolute unit.

With the aid of these units, we can now state Ohm's law in more definite language, using "*ampères*" to measure "strength of current," "*volts*" for "electro-motive force," and "*ohms*" for "resistance of circuit." Thus the two forms would run as follows:—

(General Form.) The *strength* of the current varies directly as the *electro motive force* and inversely as the *resistance* of the circuit;

(Definite Form.) The number of *ampères* of current is equal to the number of *volts* of electro-motive force, divided by the number of *ohms* of resistance in the circuit, or more briefly;

The number of *ampères* is equal to the number of *volts* divided by the number of *ohms*.

More than one method has been tried of fixing a standard for these units. Thus, the British Association (B.A.) in 1863 constructed coils of German silver to give the resistance of an ohm, but there was some doubt whether the B.A. unit exactly represented the practical unit of resistance as defined above. Accordingly, it was decided at the International Congress of Electricians in Paris in 1881 that the ohm could be most accurately measured by the resistance offered to the electric current by a column of pure mercury with a cross-section of one millimetre; and, in 1884, it was decided at the Paris Congress that the length of the column should be 106 centimetres. This gives almost exactly² the theoretical ohm, and is a little larger than the B.A. unit.³

In concluding this note on the electric terms used, we may mention that

¹ These four terms commemorate the names of four famous electricians: *Alessandro Volta*, who shares with Galvani the discovery of current electricity; *G. S. Ohm*, whose law regulating the strength of current electricity has been given above; *André Ampère*, the founder of the science of electro-dynamics; and *Charles A. de Coulomb*, the inventor of the torsion balance and demonstrator of the law that electrical attraction and repulsion vary inversely as the square of the distance.

² Lord Rayleigh calculated that the length of column to give the exact ohm should be 106.21 cm.

³ The B.A. ohm is .9887 of the new legal ohm, and the B.A. volt is .9887 of the legal volt.

the ends of the wires leading from the battery are called *Electrodes*; that *Electrolysis* (*i.e.* Electric Analysis) is, strictly speaking, the process of decomposing a liquid by means of an electric current, but is also applied to the disintegrating process said to be set up in tumours or other tissues when a current has been passed through them; and that Apostoli describes his method as *mono-polar* when only one pole is active, *i.e.*, is applied to uterus, vagina, or tissue to be acted upon, and as *bi-polar* when both poles are so applied.

Apostoli in describing his application of the faradic current uses the old phraseology (employed before the discovery of Ohm's law) when he speaks of "currents of quantity" and "currents of tension" or "intensity currents;" meaning by the former a current flowing through a circuit in which there is a very small resistance inside the battery,¹ or in the wire, and by the latter a current which has to overcome greater resistance, and which requires, therefore, a high electro-motive force.² These terms are scientifically misleading as the great resistance tends to counteract the high electro-motive power, and the principal phenomena of electro-magnetism are due not to the mere presence of electricity, however great its tension, but to its state of current or flow. The terms are, however, convenient; and, what is more to the purpose here, Apostoli's whole method is founded upon his declared discovery that the physiological effects of currents in the two conditions are very different.

ACTION OF DIFFERENT CURRENTS AND POLES.

Action of the Galvanic or Continuous Current.

The mode of action of a continuous current is different at the two poles. It is thus lucidly described by Milne Murray. "This will be best understood if we study, in the first place, the effect of the passage of the current through a piece of dead tissue—say a piece of beef. A small block of fresh beef is placed on a dish, and into it two steel sewing needles are inserted at a distance of an inch from each other. One of these is connected to the positive and the other to the negative pole of a battery, and a current of, say, 50 m.a. is transmitted. The following things will be observed. 1st. In a few seconds a frothy effervescence will appear round the negative needle, while the tissue will shrink and condense round the positive needle. 2nd. If, at the end of a few minutes, the negative needle be gently pulled, it will come away without difficulty, leaving an aperture a good deal wider than its own thickness;

¹ The *internal resistance* is diminished by having larger plates or bringing them closer together; the former is usually done by connecting the zincs of several cells, producing practically one large zinc, and the same for the coppers.

² *Brit. Med. Jour.*, 1888, I., p. 64. "No Apparatus for Faradisation," he writes, is "complete without two independent bobbins; which according to the length and thickness of the wires gives currents differing in qualities and characters. The bobbin with short thick wire gives current of quantity, because the wire is less resistant and lets pass a greater volume of electricity. The bobbin with longer and finer wire is called the bobbin of tension; the current along it is called the current of tension."

this aperture opens into a sinus which is filled with a soft frothy scum. 3rd. If the positive needle be similarly pulled, it will not come away without considerable traction, and will leave a small orifice with a dense firm outline. 4th. On examination, the negative needle will be found quite bright, while the positive needle will be dulled and slightly corroded. 5th. If the piece of meat be now carefully cut open, so as to expose the channels formed by the needles, it will be found that the tract of the negative needle is surrounded by a softened loose area of disorganised tissue, while the tract of the positive is surrounded by a condensed area much smaller than that round the negative needle; it is, moreover, paler in colour, and cuts with a somewhat gritty sensation. 6th. If the surfaces so exposed are tested with litmus paper, it will be found that on the negative side an alkaline and on the positive side an acid reaction is given."

In addition to the action at the poles, Apostoli and others have affirmed that there is an interpolar action through the entire uterine substance. Of this there is no evidence. Thus Milne Murray says, "Any so-called experimental proof which has been advanced in favour of the existence of interpolar decomposition can be readily explained on other grounds; and we may take it that there is no proof of any electrolytic action occurring anywhere except round the metallic electrodes."

Action of the Faradic or Induced Current.

This current has, according to Apostoli, "contractile power," but its effects differ as the "current of quantity" or the "current of tension" is used. The former, the direct excitant of *muscular contractility*, is employed to overcome uterine muscular inertia, and produce a temporary vascular activity; it thereby excites circulation where there is congestion and stagnation with consequent arrest of the nutrition of the uterus. The "*current of tension*" acts more on the *sensibility* than on the muscular contractility; it has therefore been used in all cases where *pain* is the leading symptom. This treatment Apostoli strongly recommends for perimetritis, ovarian pain, and intense sensibility about the lower part of the vagina.

THE APPARATUS AND INSTRUMENTS.

Apparatus
and Instru-
ments.

For a detailed description of these, the student should consult Milne Murray's monograph or Apostoli's papers. Here we can only mention them. They are (1) a battery capable of yielding an adequate constant current from 10 to 300 milliamperes; a current regulator;¹ a galvanometer; and electrodes. The internal electrode is passed into the vagina or uterus. The *intra-uterine electrode* is of the shape of the

¹ Milne Murray recommends the carbon rheostat.

uterine sound, the last 2 or 3 inches being made of platinum, and has a celluloid or gum-elastic sheath which moves up and down so as to have more or less of the platinum point exposed. When a sound cannot be passed into the uterus, the tumour may be punctured with a needle such as is used in the treatment of aneurisms. The *cutaneous* electrode used by Apostoli was made of wet potter's earth, spread on a layer about 8 inches square, and $\frac{1}{2}$ inch thick on muslin. Equally serviceable is a fold of flannel of the same size, soaked in a warm salt solution, on which a lead plate (connected with the wire) is laid. Milne Murray recommends one made of a sheet of brass wire cloth spread with a gelatine composition.

The induced or faradic current is not used as much as the continuous one. For it a faradic battery is required.

THE CURRENT: ITS STRENGTH, AND FREQUENCY OF APPLICATION.

The patient being recumbent, a vaginal douche is given, and the intra-uterine electrode passed like the sound. The abdomen is sponged with salt water, and the cutaneous electrode applied, care being taken that the contact is uniform throughout. The strength of current used is determined by what the patient can bear, or the amount passing as shown by the galvanometer. The duration of application is five to ten minutes.

Applications can be made twice a week, except during the menstrual period. When the period is prolonged, they can be resumed after the first three or four days of profuse bleeding are passed. The course of treatment varies. In fibroids the average number of application required is from 15 to 20 (Apostoli). Though the range attainable is up to 300 milliampères, it is not necessary to go beyond from 100 to 150.

Electro-puncture is called for when a sound cannot be passed. The needle is sheathed to within $\frac{1}{2}$ inch of the point, and plunged for $\frac{3}{4}$ inch into the most prominent part of the tumour. The sheathed part comes, for $\frac{1}{4}$ inch, into relation with the vaginal mucosa, which is thus protected from cauterisation by the current.

PATHOLOGICAL CONDITIONS IN WHICH ELECTRICITY IS USED IN GYNECOLOGY.

So far as our present knowledge goes, the suitable cases for Apostoli's method are—

1. *Bleeding Fibroids*.—In these the internal pole is positive, and a current strength of 50 to 150 m.a. may be used.

2. *Impacted or large Fibroids causing pressure symptoms*.—Puncture here with negative needle.

Conditions
for which
Electricity
is used in
Gyne-
cology.

3. *Dysmenorrhœa of pathological ante flexion: membranous dysmenorrhœa.*—Internal electrode negative, and current strength about 50 m.a.

4. *Cellulitis.*—Internal electrode covered with cotton wool and placed vaginally.

5. *Endometritis and Subinvolution.*

The results of electrical treatment have been such that it has now a recognised place in gynecology. They have been referred to already when discussing the different pathological conditions mentioned (see especially Fibroid Tumours, p. 449). Perhaps the most striking results are seen in the absorption of inflammatory deposits. In addition to the local changes, the improvement in the well-being of the patient is noteworthy.

THE SYSTEMATIC TREATMENT OF NERVE PROSTRATION.

LITERATURE.

Bramwell, Byrom—The Diseases of the Spinal Cord: Edin., 1882. *Gaskell*—Preliminary Notice of Investigation on the Action of the Vasomotor Nerves of Striated Muscle: Proc. Roy. Soc., Lond., 1876-77, p. 430. *Goodell*—Lessons in Gynecology, Lesson xxx.: Philadelphia, 1880. *Mitchell, Weir*—Fat and Blood, and how to make them: Lond., 1878. *Playfair, W. S.*—The Systematic Treatment of Nerve Prostration and Hysteria: Lond., 1883.

The gynecologist will not have long practised his specialty before he finds that he has occasionally to deal with a class of patients who are quite *sui generis*. The condition of such puzzles him at first extremely, inasmuch as he can find no tangible disease, but yet is bound to confess that the general condition of health is highly unsatisfactory. Very often these patients have gone the round of all medical and surgical specialists, and have come at last to the gynecologist in the hope that his art may do something to remedy their lamentable state.

The class of patients has the following characteristics:—They are thin, often emaciated, unable for any exertion, suffer from neuralgia, have little or no appetite, and are nursed by some devoted sister or mother or husband. As we have said, there is no local condition to account for their state; but often there is a history of overwork, as in the case of governesses and teachers, or of an improper training. By this latter we mean that a sensitive child of high nervous organisation has been over-cultivated, her mental energies too constantly on the rack, and has ultimately collapsed under the strain. For this class of patients Weir Mitchell of Philadelphia introduced a plan of treatment in his well-known book, the results of this method being in suitable cases highly satisfactory.

The main factors in Weir Mitchell's plan are—

- I. Seclusion of the patient, and absolute exclusion of all but the medical attendant and nurse ;
- II. Absolute Rest in Bed ;
- III. A Systematic extra-feeding of the patient ;
- IV. Use of Massage and Electricity.

I. Seclusion of the patient, and absolute exclusion of all but the medical attendant and nurse.

This is imperative, and the treatment should not be gone on with unless this condition is agreed to absolutely. Very often the friends have devoted themselves to every whim and fancy of the patient so assiduously as to impair their own health without improving that of their tyrannous charge.

The nurse should be thoroughly trained and refined, and should implicitly obey all the medical attendant's orders.

II. Absolute rest in bed.

This means muscular and mental rest, and reduces the force and frequency of the heart's action. The nutrition taken is above the amount worked off, and benefit in this way results. This absolute rest is after a while modified, and the patient allowed to sit up for a little until she may at length go about as usual, with the exception of taking a two-hours' sleep during the day.

III. A systematic extra-feeding of the patient.

This is one of the essential features of the method. Weir Mitchell begins with milk diet, about three ounces every two hours, until two quarts are given during the day. At the end of the first week raw beef soup¹ is given, and gradually the diet is increased until the dietary for one day, in one of Mitchell's cases, was as follows. Coffee at 7 ; at 8 iron and malt. Breakfast—a chop, bread and butter, of milk a tumbler and a half ; at 11, soup ; at 2, iron and malt. Dinner (closing with milk, one or two tumblers) consisted of anything she liked, and with it she took about six ounces of Burgundy or Dry Champagne. At 4, soup. At 7, malt, iron, bread and butter, and usually some fruit, and commonly two glasses of milk. At 9, soup ; and at 10, her aloes pill. At noon, massage occupied an hour. At 4.30 p.m., electricity was used for an hour.

In addition to this diet, iron in the form of Bland's pills (p. 621) and

¹ Chop 1 lb. of raw beef, and place in a bottle with 1 pint of water with 5 mm. strong hydrochloric acid. Place in ice all night, and in the morning set in a pan of water at 110° Fahr. for 2 hours. Strain thoroughly, and give filtrate in portions daily.

maltine may be added to aid the digestion of starchy food. The maltine should be given in cold milk or after dinner. The evident question now arises, How does the patient digest all this? The digestion of this immense mass of food is rendered possible by the last feature of the treatment.

IV. *The use of Massage and Electricity.*

This is most important, and consists in the systematic rubbing of the patient and the application of faradic electricity.

The massage is begun a few days after the milk diet, and consists in the systematic kneading of the skin and muscle of the whole body first for half-an-hour, and afterwards for an hour daily. A special massage nurse is necessary for this, and it should be kept up for six or seven weeks. Cocoa-nut oil should be used to render the manipulations easy, and it will also help in fattening the patient.

Electricity is employed for half-an-hour daily in order to cause muscular action, increase the blood supply to the muscle, and act as a tonic and bracing agent. Mitchell has found that after the electricity the temperature usually rises about $\frac{4}{5}$ ths of a degree. The current should not be painful, and Ziemssen's diagrams of the points of stimulation should be followed as a guide.

For further details, the literature given should be consulted by the practitioner wishing to carry it out.

The results in some cases are wonderful, and as yet no harm has been shown to arise to the kidneys from the over-feeding. The bowels must of course be regulated, and a daily motion secured. Before beginning this treatment in any case, it should be thoroughly ascertained that there is no organic disease, and no obscure and rare form of disease such as Addison's disease, myxœdema, etc. A consultation with a specialist should always be had in cases of doubt.

The patient for whom it is suitable is one where there has been under-feeding or improper food, undue mental strain, and consequent loss of flesh and nervous energy.

HYSTERIA.

LITERATURE. *Bourneville et Regnard*—Iconographie photographique de la Salpêtrière : Paris, 1877. *Bourneville et d'Olier*—Recherches sur l'Épilepsie, l'Hystérie et l'Idiotie : Progrès Médical, 1881. *Charcot*—Diseases of the Nervous System: Sydenham Society's Series, London, 1877. *Fritsch*—Krankheiten der Frauen: Braunschweig, 1881. *Jolly*—Article "Hysteria" in Ziemssen's Cyclopædia of Medicine. *Mills*—Hystero-epilepsy : American Journal of the Medical Sciences, Oct. 1881. *Richer*—Études cliniques sur l'Hystéro-Epilepsie : Paris, 1881.

The frequency of hysteria as a complication of pelvic disease requires that we notice it briefly. We can only indicate the leading points and

refer the student to the literature given above. The connection which exists between hysterio-epilepsy and the ovary also calls for short reference.

As to the pathological changes present in hysteria, little definite is known, except what Freund has described in *Parametritis chronica atrophicans* (v. p. 262). In regard to etiology, we note first the influence of heredity; defective moral education *by a hysterical mother, and the power of imitation in developing hysteria, confirm this influence. A reduced state of the system is also a very important cause, and the one to which treatment must be specially directed. As to the exciting causes usually given (such as dysmenorrhœa, uterine displacements, ovaritis), these are so common that we cannot regard them as a cause of hysteria. The only ascertained facts are that removal of the ovaries has in some cases cured hysteria, and that pressure in an ovarian region does sometimes inhibit a hysterio-epileptic attack.

The symptoms of hysteria are protean. *Sensation* is affected as follows. There may be increased sensitiveness to touch (hyperæsthesia) and to pain (hyperalgesia). Hyperæsthesia of the joints is important as simulating arthritis, from which it is diagnosed by the fact that the pain is around (not in) the joint, and that it is not aggravated on forcing the articular surfaces together. Neuralgia along the spine with tender points simulates disease of the vertebral column. The typical headache (known as the "clavus hystericus" from the localised and intense character of the pain), neuralgia of the muscles generally, localised pain in the breast, in one ovarian region, in the bladder and urethra, and the perversions of the special senses need only be mentioned here. When sensitiveness is impaired, it is usually that to pain; while that to heat and touch remains; one half of the body may be affected, or isolated portions of skin—as the back of the hands and feet. Loss of the muscular sense prevents the patient, if the eyes be closed, from knowing what movements she has made. Anæsthesia of any of the mucous membranes may occur. The special senses are often also impaired.

The *motor* disturbances resulting in convulsions belong rather to hysterio-epilepsy, and are fully described, with characteristic photographs, in Bourneville and Regnard's monograph. The paralysis due to hysteria is very important in regard to its diagnosis from that due to a cerebral or spinal lesion. It varies in distribution, and may affect one limb only, or the arm and leg of one side, or the arm on one side and the leg on the other. In the face, the levator palpebræ superioris is frequently affected; paralysis of the muscles supplied by the facial and hypoglossal nerves is rare. This last fact is of value in diagnosing between hysteria and hemiplegia; further, gradual onset, presence of anæsthesia and its varying distribution, normal reaction to the electric current, the progress of the case with variations in the degree and extent

of the paralysis, warrant us in diagnosing hysteria. The diagnosis of hysterical paraplegia, from multiple sclerosis is more difficult. Paralysis may also affect the laryngeal muscles, producing aphonia, and the muscular wall of the œsophagus, stomach, and intestines.

Of the disturbances of the circulatory system, the most important is palpitation with increased force of the apex beat; in some cases, the heart's action fails and there is syncope. Vaso-motor disturbances are seen in the pale skin which does not bleed when pricked, and in the flushings and profuse sweatings which are often present. Salivation and polyuria often occur after a hysterical attack.

In forming a diagnosis, we must be careful to exclude the possibility of organic, cerebral, or spinal disease. A case reported by Bruce¹ is of interest in this connection; here the patient had symptoms of hysteria, there was no optic neuritis or other indication of cerebral mischief, and yet the post-mortem showed a large tumour in the temporo-sphenoidal lobe.

In treatment, the following points are of importance. Care must be taken in the mental and moral training of the children, where there is a tendency to hysteria.² If the system is below par, Weir Mitchell's method should be tried, and iron given when there is anæmia; cold baths are always beneficial. In grave cases, Battey's or Tait's operation may be suggested but never urged, as the results are not brilliant.

MASSAGE.

LITERATURE. *Profanter*—(1) Die Massage in der Gynäkologie; (2) Die manuelle Behandlung des Prolapsus Uteri: Wien, 1888. *Reibmayr*—Die Massage: Leipzig, 1889. *Resch*—Ueber die Anwendung der Massage bei Krankheiten der weiblichen Sexualorgane: Cent. für Gynäk., No. 32, 1887.

One of the most common cases in Gynecology is that where, as the result of a previous attack of pelvic inflammation, the uterus and ovaries are bound down and fixed by more or less dense adhesions—usually peritonitic. For these cases many forms of treatment, ranging from the hot douche up to abdominal section, are recommended, and will be found described in various parts of this Manual. At present we wish briefly to refer to a method of treatment recently come into vogue—Massage.

By this we mean here Bimanual Massage of the adherent tissues or organs so as to slacken these, promote vascular and lymphatic absorption, and in this way bring about a more healthy condition of the local circulation and relief to the nerve pressure supposed to be exerted by the cicatricial tissues.

The originator of this form of treatment is a Swedish layman,

¹ Brain, Part XXII. : 1883.

² Clouston: Puberty and Adolescence medico-psychologically considered: Edin., 1880.

Brandt, and his work has been taken up by several German gynecologists, among whom are Schultze, Profanter, Schauta, and others.

Before going on more particularly to the question of indications, methods, and results, we may say that we believe there are great difficulties in the way of its general acceptance. The chief one is that it involves undue manipulation of the genital organs. This is a most serious objection, and one which will in all probability be fatal to the method. Then again the manipulation will be dangerous if the diagnosis be wrong—*e.g.*, if a pyosalpinx be chosen for it. There is thus every prospect of its being supplanted in the few cases requiring it by abdominal section.

Prolapsus uteri is one of the cases specially recommended for it. Here, however, it is difficult to understand how it does good, although trustworthy observers have recorded cases of cure.

Indications. Retroversion of uterus bound down by adhesions; adherent ovaries; parametritis posterior causing pathological ante flexion; prolapsus uteri.

Methods. In chronic inflammatory cases the patient occupies the dorsal posture, with knees well drawn up and dress freely loosened. The gynecologist carefully ascertains bimanually the condition of the organs, and then, keeping the two fingers passed into the vagina fixed, he grasps or maps out by the outer hand the adhesions to be stretched, and by movement of the outer hand only, stretches these or exercises a rubbing movement on them. Rectal manipulation may be employed instead of vaginal. This bimanual massage should not be practised for more than a few minutes at each sitting, and the number of sittings must be left to the judgment of the gynecologist.

Schultze has extended this method by advocating and practising, not mere stretching, but actual separation of the adhesions. For this purpose the patient is chloroformed, the condition accurately mapped out, and the adhesions then separated by bimanual manipulation. Schultze's results have been good, but it is evident that the risks in less experienced hands are very great.

In prolapsus uteri the method is more complicated and troublesome. Briefly it is as follows (*Profanter*).

(1) *Position of patient.* The patient has her dress thoroughly loosened and lies on a short couch (4 ft. \times 2 ft. 8 in.) with her chest supported by cushions. In this way she is compelled to slacken the abdominal muscles as much as possible. An assistant passes his fingers into the vagina, replaces and ante flexes the uterus. The masseur then with both hands grasps the uterus and draws it up as far as possible.

The patient now raises the hips from the couch, thus supporting her body on elbows and feet, while the gynecologist forcibly separates her

closed knees and then forcibly approximates them, the patient resisting each time. These manœuvres are repeated thrice.

The object of this so-called pelvic gymnastic is to bring into action the pelvic muscles (levator ani, obturator internus, perineal muscles) and thus strengthen the musculature and fascia of the pelvic floor.

The patient need not be confined to bed during the intervals of the treatment.

CASE-TAKING.

LITERATURE. *Emmet*—Gynecology, p. 57: London, 1880. *Simpson, A. R.*—Contributions to Obstetrics and Gynecology, Method of Case-Taking in Gynecology, p. 317.

It is of importance to give some hints as to case-taking or the investigation of cases of diseases of the female sexual organs.

In hospitals, some form of case-taking card is usually employed; and we describe the method of case-taking adopted by Professor Simpson in the Buchanan Ward (for the Diseases of Women) in the Edinburgh Royal Infirmary.

CASE-TAKING CARD.

ANAMNESIS.

1. NAME; AGE; OCCUPATION; RESIDENCE; MARRIED, SINGLE, OR WIDOW; DATE OF ADMISSION.

2. COMPLAINT AND DURATION OF ILLNESS.

3. GENERAL HISTORY OF—(a) Present attack; (b) Previous Health; (c) Diathesis; (d) Social Condition and Habits; (e) Family Health.

4. SEXUAL HISTORY.

(1) *Menstruation*—

A. Normal—(a) Date of Commencement; (b) Type; (c) Duration; (d) Quantity; (e) Date of Disappearance.

B. Morbid—(a) Amenorrhœa; (b) Menorrhagia; (c) Dysmenorrhœa.

(2) *Intermenstrual Discharge*—(a) Character; (b) Quantity.

(3) *Pareunia*.

(4) *Pregnancies*—(a) Number; (b) Dates of First and Last; (c) Abortions; (d) Character of Labours; (e) Puerperia; (f) Lactations.

5. LOCAL FUNCTIONAL DISTURBANCES—(a) Bladder; (b) Rectum; (c) Pelvic Nerves and Muscles.

6. GENERAL FUNCTIONAL DERANGEMENTS—(a) Nervous System; (b) Respiratory System; (c) Circulatory System; (d) Digestive System; (e) Emunctories.

PHYSICAL EXAMINATION.

1. GENERAL APPEARANCE AND CONFIGURATION.

2. MAMMÆ.

3. ABDOMEN—(a) Inspection; (b) Palpation; (c) Percussion; (d) Auscultation; (e) Mensuration.

4. EXTERNAL PUDENDA.

5. PER VAGINAM—(a) Orifice; (b) Walls and cavity; (c) Roof; (d) Os and Cervix Uteri.

6. BIMANUAL EXAMINATION (Abdomino-vaginal, Recto-vaginal, Abdomino-rectal, Abdomino-recto-vaginal, Abdomino-vesico-vaginal)—

(1) *Uterus*—(a) Size; (b) Shape; (c) Consistence; (d) Sensitiveness; (e) Position; (f) Mobility; (g) Relations.

(2) *Fallopian Tubes*.

(3) *Ovaries*—(a) Size; (b) Situation; (c) Sensitiveness.

(4) *Peritoneum and Cellular Tissue*.

(5) *Bladder*. (6) *Rectum*. (7) *Pelvic Bones*.

7. USE OF—(a) Speculum; (b) *Volsella*; (c) Sound; (d) Curette; (e) Aspiratory Needle; (f) Tent.

8. PHYSICAL CHANGES IN—(a) Nervous, (b) Respiratory, (c) Circulatory, (d) Digestive, (e) Emunctory Organs; (f) Skin; (g) Bones.

DIAGNOSIS.

PROGNOSIS.

TREATMENT.

PROGRESS AND TERMINATION.

We have drawn up a schedule¹ based on this card which will be found very convenient, either in private or in dispensary practice, for recording gynecological cases.

ANAMNESIS.

In taking the history of a case the first thing is to ascertain the patient's complaints. These vary much, but in minor cases are usually those of pain, increased menstruation, painful menstruation, leucorrhœa. The pain is usually sacral, iliac or left infra-mammary. A very important




FIG. 379.

OUTLINE DIAGRAM OF ABDOMEN FOR RECORDING POSITION OF TUMOURS RELATIVE TO THE BODY LANDMARKS.

point is to ascertain the starting-point of the patient's illness, and in many cases it can be traced to labour, abortion, or some specific gonorrhœal attack. Not infrequently it has followed marriage. When menorrhagia is complained of, the amount of discharge, and whether clots or shreds are passed, should be noted; and if pain at the period is present, the time of its occurrence, whether before, during, immediately

¹ Supplied by Messrs W. & A. K. Johnston, Edinburgh, in separate sheets, or in book-form.

after the period, or between the periods should be ascertained. Bleeding after coitus is suspicious of malignant disease. The history of a case should be carefully taken, the patient cross-examined on obscure points, but her statement should always have full weight given to it, and should not be discounted. As a matter of fact the patient is a witness really, and can only have her evidence discarded if grossly improbable. In a case of suspected pregnancy for instance, the patient's statement, if made, that she has menstruated regularly is of the greatest importance, and must not be brushed aside without careful investigation. It may be that she has had a regular discharge, and this therefore should be specially inquired into. Queries under the head of pareunia should not be put unless special indications are present, or the patient volunteers a statement. It is important to note the diathesis—strumous, neurotic, tubercular, gouty. The remaining facts to be ascertained under "Anamnesis" need no further comment.

PHYSICAL EXAMINATION.

Abdominal Examination. The method has been already described (v. p. 104). Girth measurements should be made if necessary, and the routine inspection, palpation, etc., carefully done. This should then be checked by the following rapid methods.

The student has to consider in a case of abdominal distention the following points.

- (1.) Is there any abnormal abdominal content?
- (2.) Is it solid?
- (3.) Is it fluid, and if so (a) encysted and intraperitoneal, or (b) encysted and extraperitoneal.
- (4.) Is the fluid when present, free, or partially encysted.

In this way we do not consider special growths, but only some general conditions common to several.

(1.) *Is there any abnormal abdominal content?* This is best settled by percussion. A tympanitic note is evidence of the absence of any abnormal content of much size. This remark does not apply to small growths, and the examination must be supplemented by bimanual palpation with one hand behind and one in front, and also by the abdomino-vaginal bimanual.

(2.) *Is it solid?* We judge of this by touch. We say the tumour is solid, because it feels firm. Firmness must not be confounded with tension in a cystic tumour. The best type of solid tumour is the ordinary fibromyoma uteri (not the soft fibroid).

(3.) *Is it fluid, and, if so, is it (a) encysted and intraperitoneal, or (b) encysted and extraperitoneal?* An encysted intraperitoneal tumour has tension, fluctuation (perfect in parovarian tumours, less so in ovarian),

symmetrical dulness circular or oval in outline, not altering when the patient's posture is changed, and the *uterus lies usually below, and is not involved in the tumour*. This is best ascertained by recto-vaginal examination, and is a point of great value.

In the encysted extraperitoneal growth we have dulness, not altering when the patient changes her position, tension, *but the uterus is often lifted up: occasionally the large bowel (ascending or descending colon) is displaced towards the middle line*.

(4.) *Is the fluid when present free or partially encysted?* Here we find on percussion, when patient is dorsal, a concave upper boundary to dulness (concavity pointing down), dulness at flank and a fluctuating impulse. In the lateral posture the dull note is on the lower side only.




FIG. 380.

OUTLINE DIAGRAM OF PELVIS FOR FILLING IN POSITION OF UTERUS OR TUMOURS (*A. R. Simpson*).

When a large amount of free fluid is present, the extensive dulness may mislead.

Irregular dulness and less perfect change of note on change of posture mean irregular adhesion, and are suspicious of tubercular or malignant disease. The student should keep in mind that a soft fibroid may simulate fluctuation even when touched directly, as is demonstrated at an abdominal section.

The student now proceeds to the *inspection of the genitals*, to the *vaginal examination* and the complete *bimanual* (*v. Chap. VIII.*), and from the facts he ascertains has to make up his mind as to diagnosis, prognosis, and treatment.

Many cases permit of accurate diagnosis, some only of an opinion as to their probable nature.

One special point to be kept in mind is that the commonest case in minor gynecology is one with multiple lesions, viz., laceration of the cervix, more or less pronounced, often a retroflexion with endometritis, and some varying amount of cellulitic peritonitic or appendage thickening. The history is that this has followed abortion, childbirth, or marriage.

The mistake usually made is that of picking out one lesion and giving it undue prominence. The best reading of such cases is that of minor septic infection through the endometrium or cervix with the resultant secondary lesions already given, and treatment is not so much mere




FIG. 381.

OUTLINE DIAGRAM PELVIS AS SEEN THROUGH THE BRIM, TO FILL IN POSITION OF TUMOURS
RELATIVE TO UTERUS (*Schultze*).

rectification of any displacement, but the treatment of the endometritis by curetting and subsequent attention to the inflammatory conditions by the use of the tampon, blisters, hot douche. In obstinate cases, operative removal of affected tubes and ovaries must be considered.

CLASSIFICATION OF DISEASES OF WOMEN.

As already explained, we have classified the diseases considered on an anatomical basis (p. 180).

The following table shows that they may be classified on a pathological basis, and is given here to show how far our knowledge has gone, and that many gaps in it have yet to be filled up.

We may group the diseases of gynecology under the heads of—

1. *Congestive: Vascular Ruptures.*
2. *Hypertrophy.*
3. *Atrophy.*
4. *Traumatism.*
5. *Simple inflammation.*
6. *Micro-organismal:*
 - A. Due to known microbes still in tissues—
 1. Tubercular,
 2. Gonorrhœal,
 3. Actinomycotic,
 4. Septic.
 - B. Due to microbes and their products, these causes having been apparently completely eliminated.
7. *Herniæ.*
8. *New Growths.*
9. *Developmental errors.*
10. *Neuroses.*

Almost all gynecological diseases can be grouped under these heads as in the following table:—

PATHOLOGICAL CLASSIFICATION OF DISEASES OF WOMEN.

I. CONGESTIVE: VASCULAR RUPTURE—

- (1.) Alleged simple congestion of genital tract (?).
- (2.) Pelvic hæmatocele.
- (3.) Pelvic hæmatoma.
- (4.) Ovarian apoplexy.

II. HYPERTROPHY—

- (1.) Hypertrophy of vaginal portion of cervix.
- (2.) „ middle portion.
- (3.) „ supravaginal portion.
- (4.) „ associated with lacerated cervix.
- (5.) Simple hypertrophy of organs.

III. ATROPHY—

- (1.) Atrophic Pelvic Peritonitis.
- (2.) Parametritis chronica atrophicans, circumscripta et diffusa.
- (3.) Superinvolution of uterus.
- (4.) Kraurosis vulvæ.
- (5.) Certain forms of pruritus vulvæ.
- (6.) Senile changes in organs.

IV. TRAUMATISM—

- (1.) Cervical lacerations.
- (2.) Vaginal lacerations.
- (3.) Perineal, vestibular, and vulvar tears.

V. SIMPLE INFLAMMATIONS—

Alleged simple non-specific inflammatory conditions of all genital organs (?).

VI. MICRO-ORGANISMAL AND PARASITIC—

A. Acute and Chronic Infective Diseases—

1. Tubercular disease of peritoneum, Fallopian tube, ovary, uterus, vagina, vulva.
2. Gonorrhœal inflammation of peritoneum, connective tissue, Fallopian tube, ovary, uterus, vagina, vulva, and its ducts. Pyosalpinx.
3. Actinomycosis of tube, ovary, and connective tissue.
4. Septic diseases : Acute peritonitis, cellulitis, oöphoritis, salpingitis, endometritis, metritis. Septic sources in cervix, vagina, vulva. Pyosalpinx. Hydrosalpinx.

B. Chronic Infected Diseases—

1. *A group of cases where we have multiple lesions ;* usually some enlargement and displacement of the uterus, with various degrees of endometritis and thickenings of an inflammatory nature, either peritonitic, cellulitic, salpingitic, or oöphoritic ; cervical splits are often present.
2. Pathological versions and flexions.
3. *A group with minor single inflammatory conditions* of ovary, tube, peritoneum, connective tissue ; often associated with endometritis.

C. Parasitic—

1. Echinococci (*tænia echinococci*) in connective tissue, peritoneal cavity, uterus, ovary.
2. Parasitic skin diseases and pedicular irritation.

VII. HERNIÆ :

- (1.) Herniæ of uterus and appendages, singly or combined, into inguinal canal.
- (2.) Prolapsus uteri.
- (3.) Vaginal enterocele (anterior and posterior). Vulvar enterocele.

VIII. NEW GROWTHS :

A. *Non-Malignant*—

1. *Overgrowths of normal tissue*, with or without fluid or solid products.

(a) *Unstriped muscle*.—Fibro-myoma of round ligament, in broad ligament, of ovary, of uterus; fibrocystic of uterus; fibro-myomatous polypi—*vide* also (d).

(b) *Fibrous connective tissue and fat*.—Fibrous growths of round ligament, of ovary, of vagina. Lipomata of broad ligament, of vulva.

(c) *Glandular tissue*—

(1) Endometritis.

(2) Cystoma ovarii.

(3) Dermoids of ovary.

(4) Cervical and uterine polypi (mucous).

(d) *Vascular elements*.—Fibroids of uterus—some forms *vide* (a). Endometritis fungosa.

B. *Malignant*—

1. *Sarcoma* of pelvic connective tissue; of round ligament; of tube; of ovary, uterus, vagina, and vulva.
2. *Carcinoma* of ovary, peritoneum, tube, uterus, vagina, and vulva; combinations.

C. *Growths due to Evolutionary Relics*—

- (a) *To morbid development of Graafian follicles*.—Cystic and dermoid ovary.
- (b) *To remains of Wolffian body and duct, or to morbid development of germ epithelium*.—Parovarian and papillomatous tumours;¹ some broad ligament cysts;¹ some vaginal and cervical cysts.
- (c) *To undue development of ovarian fossa*.—Ovarian hydrocele.

IX. DISPLACEMENTS OF UTERUS :

- (1) Inversion of uterus—*vide* (VI.) Group B, and (VII.).

Often malignant.

X. DEVELOPMENTAL ERRORS :

- (1) Malformations in peritoneal folds : of ovary, tube, uterus, vagina, vulva. Hermaphroditism.
- (2) Round ligament hydrocele.
- (3) Allantoic cysts.
- (4) Hæmatosalpinx, hæmatometra, hæmatocolpos. Some forms of tubal distension. Hydrosalpinx.

XI. NEUROSES :

Hysteria ; hystero-epilepsy ; neurasthenia (we may have also neurotic symptoms, viz. reflex pains, some forms of dysmenorrhœa ; menorrhagia).

In the *B* section, under the micro-organismal group, or what we have termed *chronic infected cases*, we have really the majority of the chronic diseases coming continually under the notice of the practitioner and specialist. It is on this group that so much theory and energy have been spent, in the attempt to get at the nature of the associated lesions, and to cure the discharges, aches, discomfort, and disturbance of function they give rise to. The *A* and *B* groups are really one, but are sharply contrasted, as in the former we have the exact scientific pathology ; in the latter, dreary speculative theories, where one does not dig long without coming, in Carlyle's phrase, to water. Group *B* is the dust-heap of Gynecology, where valuables lie hid and obscured.

The view of such cases that we wish to urge is this. Many are the result of a previous septic attack, minor or major. The source of infection has been the cervical laceration or the endometrium bared by abortion or labour, and the invasion of the micrococci along mucous membrane or through lymph spaces has been met by the ever-watchful tissue elements ; the invaders have been captured ; and in the cellulitic thickenings, or in those of the tubes and ovary found lasting for months and years after the primary attack, one has the tumuli, as it were, that mark the old battlefield. Gradually, as these thickenings shrink, we get uterine displacements, nerve pressure, cervical ectropion, and hypertrophy, the stock cases of gynæcology (*v. p.* 714).

SOURCES OF GYNECOLOGICAL LITERATURE.

At the beginning of each subject we have already given a summary of the literature to which we were indebted. The literature given, therefore, represents what we considered important, and what we had in most cases personally studied.

Gynecological Literature is so extensive that a full résumé of it would have occupied several times the space we have allotted to

the whole subject. We wish however to point out here the sources, so that any practitioner who wishes to ascertain the best books and monographs on any special subject may know how and where to begin his search.

The sources of Gynecological Literature are threefold :—

- I. Catalogues, Dictionaries ;
- II. The larger Text-books of Gynecology ;
- III. Articles and Abstracts in the various Gynecological quarterlies, monthlies, and weeklies, with Retrospects and Jahrbücher.

I. CATALOGUES, DICTIONARIES.

- (1.) *Index-Catalogue of the Library of the Surgeon-General's Office, U.S.A.* Washington Government Printing Office. In this splendid work, the authors and works are arranged alphabetically ; its value cannot be over-rated.
- (2.) *Nouveau Dictionnaire de Médecine et de Chirurgie pratique* : Paris, J. B. Bailliére et Fils.
- (3.) *Dictionnaire Encyclopédique des Sciences Médicales* : Asselin et Cie., Paris.
- (4.) *Real-Encyclopädie des gesammten Heilkunde* : Wien.
Wood's Cyclopædia : Edinburgh, Young J. Pentland, 1889.
Annual of the Universal Medical Sciences (edited by Sajous) : Philadelphia.
Buck's Reference Handbook of the Medical Sciences : New York.

II. LARGER MODERN TEXT-BOOKS OF GYNECOLOGY.

ENGLISH.

- Barnes*—Diseases of Women : London, J. & A. Churchill, 1878.
Baldy—An American Text-book of Gynecology : Philadelphia, Saunders, 1894.
Berry Hart—Selected Papers in Gynecology, etc. : Edinburgh, W. & A. K. Johnston, 1893.
Byford—Medical and Surgical Treatment of Women : Philadelphia, 1888.
Duncan, Matthews—Diseases of Women : London, Churchill, 1889.
Emmet—Principles and Practice of Gynecology : Philadelphia, Lea's Son & Co., 1880.
Garrigues—Diseases of Women : Philadelphia, W. B. Saunders, 1894.
Goodell—Lessons in Gynecology : Philadelphia, Brinton.
Hewitt—The Diseases of Women : London, Longmans, Green & Co., 1868.
Macnaughton-Jones—Diseases of Women : London, Bailliére, Tindall & Cox, 1897.
More Madden—Clinical Gynecology : London, Bailliére, Tindall & Cox, 1893.
Mundé—Minor Surgical Gynecology : New York, Wood & Co., 1885.
Simpson, A. R.—Obstetrics and Gynecology : Edinburgh, A. & C. Black, 1880.
Simpson, Sir J. Y.—Diseases of Women : (edited by A. R. Simpson) : A. & C. Black, 1871.
Sims, J. Marion—Uterine Surgery : London, Hardwicke, 1866.
Skene, A. J. C.—The Diseases of Women, Treatise on : London, Lewis, 1892.
Sutton, J. Bland—Surgical Diseases of the Ovaries, etc. : London, Cassell & Co., 1896.
Tait, Lawson—Diseases of Women : Birmingham, Cornish Bros., 1886.
 The Pathology and Treatment of Diseases of the Ovary : Birmingham, 1883.
 Diseases of Women and Abdominal Surgery, Vol. I. : Philadelphia, Lea Brothers & Co., 1889.
Thomas—Treatise on Diseases of Women : Philadelphia, Lea, 1891.
Thorburn—Diseases of Women : Griffin & Co., London, 1885.
Wells, Sir T. S.—Ovarian and Uterine Tumours : London, 1882.
West (Duncan's Edition)—Diseases of Women : London, Churchill, 1864.

A System of Gynecology—Edited by T. Clifford Allbutt and W. S. Playfair: Macmillan & Co., Ltd., London and New York, 1896.

- The Development of Modern Gynecology—*M. Handfield Jones.*
- The Anatomy of the Female Pelvic Organs—*D. Berry Hart.*
- Malformations of the Genital Organs in Women—*J. William Ballantyne.*
- The Etiology of the Diseases of the Female Genital Organs—*W. Balls-Headley.*
- Diagnosis in Gynecology—*Robert Boxall.*
- Inflammation of the Uterus—*A. H. Freeland Barbour.*
- The Nervous System in Relation to Gynecology—*W. S. Playfair.*
- Sterility—*Henry Gervis.*
- Gynæcological Therapeutics—*Amand Routh.*
- The Electrical Treatment of Diseases of Women—*Robert Milne Murray.*
- Disorders of Menstruation—*John Halliday Croom.*
- Diseases of the External Genital Organs—*William J. Smyly.*
- Displacements of the Uterus—*Alexander Russell Simpson.*
- Morbid Conditions of the Female Genital Organs Resulting from Parturition—*George Ernest Herman.*
- Extra-Uterine Gestation—*John Bland Sutton.*
- Pelvic Inflammation—*Charles James Cullingworth.*
- Pelvic Hæmatocele—*William Overend Priestley.*
- Benign Growths of the Uterus—*F. W. N. Haultain.*
- Hysterectomy—*J. Knowsley Thornton.*
- Malignant Diseases of the Uterus—*W. J. Sinclair.*
- Plastic Gynæcological Operations—*John Phillips.*
- Diseases of the Fallopian Tubes—*Alban Doran.*
- Diseases of the Ovary—*W. S. A. Griffith.*
- Ovariectomy—*J. Greig Smith.*
- Chronic Inversion of the Uterus—*Edward Malins.*
- Diseases of the Female Bladder and Urethra—*Henry Morris.*

The American System of Gynecology and Obstetrics—Edited by Matthew D. Mann, Edinburgh: Young J. Pentland, 1887.

- Historical Sketch of American Gynecology—*Edward W. Jenks.*
- The Development of the Female Genitals—*Henry I. Garrigues.*
- The Anatomy of the Female Pelvic Organs—*Henry C. Coe.*
- Malformations of the Female Genitals—*Henry I. Garrigues.*
- Gynecological Diagnosis—*Egbert H. Grandin.*
- General Consideration of Gynecological Surgery—*E. C. Dudley.*
- General Therapeutics—*Alexander J. C. Skene.*
- Electricity in Gynecology—*Alphonso D. Rockwell.*
- Menstruation and its Disorders—*W. Gill Wylie.*
- Sterility—*A. Reeves Jackson.*
- Diseases of the Vulva—*Matthew D. Mann.*
- The Inflammatory Affections of the Uterus—*C. D. Palmer.*
- Subinvolution of the Vagina and Uterus—*Thaddeus A. Reamy.*
- Periuterine Inflammation—*Richard B. Maury.*
- Pelvic Hæmatocele and Hæmatomata—*Ely Van de Warker.*
- The Diseases of the Vagina—*Charles Carroll Lee.*
- The Hystero-Neuroses—*George J. Engelmann.*
- Extra-Uterine Gestation—*T. Gaillard Thomas.*
- Tumours of the Breast—*Samuel W. Gross.*
- Diseases of the Breast other than Tumours—*Roswell Park.*
- Fistulae—*Edward W. Jenks.*
- Diseases of the Bladder and Urethra—*William H. Baker.*
- Non-Malignant Tumours of the Uterus—*R. Stansbury Sutton.*
- The Malignant Diseases of the Uterus—*W. T. Lusk.*
- Lacerations of the Cervix Uteri—*Bache M'Evers Emmet.*
- Chronic Inversion of the Uterus—*Samuel C. Busey.*
- Injuries and Lacerations of the Perineum and Pelvic Floor—*Howard A. Kelly.*
- The Treatment of Ovarian and of Extra-Ovarian Tumours—*William Goodell.*
- Diseases of the Ovaries—*Robert Battey and Henry C. Coe.*
- Diseases of the Fallopian Tubes—*Henry C. Coe.*
- The Pathology of Ovarian Tumours—*Stephen Y. Howell.*
- The Clinical History and Diagnosis of Pelvic Tumours other than Uterine and Tubal—*Matthew D. Mann.*
- Displacement of the Uterus—*George T. Harrison.*

GERMAN.

- Fehling*—Lehrbuch der Frauenkrankheiten : Stuttgart, Enke, 1893.
Fritsch—Krankheiten der Frauen : Braunschweig, Berlin, Wreden, 1896.
Hegar und Kaltenbach—Die operative Gynäkologie : Stuttgart, Enke, 1897.
Hofmeier—Grundriss der Gynäkologischen Operationen : Leipzig, Deuticke, 1892.
Schauta—Lehrbuch der gesamten Gynäkologie : Leipzig, Deuticke, 1896.
Schroeder—Handbuch der Krankheiten der weiblichen Geschlechtsorgane : Leipzig, Vogel, 1893.
Veit—Handbuch der Gynäkologie : Wiesbaden, Bergmann, 1896.
Winckel—Lehrbuch der Frauenkrankheiten : Hirzel, Leipzig, 1890.
Winter—Lehrbuch der gynäkologischen Diagnostik : Leipzig, Hirzel, 1896.
Handbuch der Frauenkrankheiten redigirt von Billroth u. Luecke : Stuttgart, Enke, 1879.
- I. Band. Die Untersuchung der weiblichen Genitalien und allgemeine gynäkologische Therapie—*Chrobak*.
 Die Sterilität der Ehe. Entwicklungsfehler des Uterus—*Müller*.
 Die Lageveränderungen und Entzündungen des Uterus—*Fritsch*.
 - II. Band. Die Neubildungen des Uterus—*Gusserow*.
 Die Krankheiten der Ovarien—*Olshausen*.
 Die Krankheiten der Tuben, der Ligamente, des Becken-peritonäum und des Beckenbindegewebes, einschliesslich der Extrauterinschwangerschaft—*Bandl*.
 - III. Band. Die Krankheiten der weiblichen Brustdrüsen—*Billroth*.
 Die Krankheiten der äusseren Genitalien und die Dammrisse—*Zweifel*.
 Die Krankheiten der weiblichen Harnröhre und Blase—*Winckel*.
 Die Krankheiten der Vagina—*Breisky*.

FRENCH.

- Auvard*—Traité pratique de gynécologie : Paris, O. Doin, 1894.
Bernutz and Goupil—Clinical Memoirs on the Diseases of Women : Sydenham Society Tr., 1867.
Bonnet et Petit—Traité pratique de gynécologie : Paris, Baillière et fils, 1894.
Courty—Traité pratique des Maladies de l'utérus, 2nd Edition : Paris, Asselin : also Dr Agnes Maclaren's Translation, London.
De Sinéty—Manuel Pratique de Gynécologie : Paris, Doin, 1879.
Dolérès et Pichevin—Introduction à la pratique gynécologique : Paris, Reuff et Cie., 1896.
Leblond—Traité élémentaire de Chirurgie gynécologique : Paris, Lauwereyns, 1878.
Pozzi—Traité de Gynécologie : Paris, Masson et Cie., 1897.
Tripier—Leçons cliniques sur les Maladies des Femmes : Paris, Doin, 1883.

III. JOURNALS : RETROSPECTS : INDEXES : JAHRBÜCHER.

- American Journal of Obstetrics : New York, Wm. Wood & Co.
 British Medical Journal : London.
 Cassell's Year Book of Treatment.
 Dublin Journal of Medical Science : Dublin, Fannin & Co.
 Edinburgh Medical Journal : Edinburgh, Young J. Pentland.
 Glasgow Medical Journal : Glasgow, MacDougal.
 International Journal of Medical Sciences : Lea's Son & Co., Philadelphia ; Cassell & Co., London.
 Lancet : London.
 London Medical Record : Smith, Elder & Co.
 Medical Press and Circular : London.
 New York Medical Journal and Obstetrical Review : New York, Appleton & Co. ; and London, Cassell & Co.
 Reference Handbook of Medical Sciences : Wood & Co., New York.
 Scottish Medical and Surgical Journal : Edinburgh, Clay.
 Archiv für Gynäkologie : Berlin, Hirschwald.
 Berliner klinische Wochenschrift.
 Monatsschrift für geburtshülfe und Gynäkologie : Berlin, Imberg und Lefson.
 Centralblatt für Gynäkologie : Leipzig, Breitkopf und Härtel.

- Zeitschrift für Geburtshülfe und Gynäkologie : Stuttgart, Enke.
 Annales de Gynécologie : Paris.
 La Gynécologie : Paris, O. Doin.
 Revue de Gynécologie : Paris, Masson.
 Annali di Ostetricia, Ginecologia e Pediatria : Milano, Pietro Agnelli.
 Braithewaite's Retrospect : London, Simpkin, Marshall & Co.
 Index Medicus : a monthly classified Record of the current Medical Literature of
 the World : G. S. Davis, Boston and Detroit, U.S.A.
 Annual of the Universal Medical Sciences (Edited by Sajous) : Davis, Philadelphia.
 Schmidts's Jahrbücher : Leipzig.
 Supplement to Ziemssen's Cyclopædia : London, Sampson, Low, Marston, Searle,
 & Rivington.
 Revue des Sciences Médicales : Paris, E. Masson.
 Neale's Digest : London, Ledger, Smith & Co., 1882.
 American Gynecological Transactions (Index at end) : Boston, Houghton & Co.
 London Obstetrical Transactions : Longmans, Green & Co.
 Edinburgh Obstetrical Transactions : Oliver & Boyd.

In looking up literature on any special subject, first consult the literature given at the beginning of each chapter and then the index of Recent Gynecological Literature in the Appendix. The list of literature given in Billroth and Luecke's Handbuch, the Index Medicus, Neale's Digest and the U. S. A. Index Catalogue may also be consulted with advantage. The various Retrospects and Jahrbücher mentioned above give abstracts of the papers, and the French and German Cyclopædias give special exhaustive articles on each subject.

INDEXES



INDEX OF AUTHORS.

- ABBÉ, 671.
 Abel, 332, 333, 347, 491, 517.
 Ackermann, 476.
 Adams, 405, 617.
 Alexander, 405, 406, 617.
 Allbutt, 449.
 Allingham, 681.
 Althaus, 448.
 Alquié, 405.
 Ampère, 700.
 Apostoli, 370, 448, 449, 450, 697,
 702, 703.
 Aran, 616.
 Arnott, 505.
 Ascher, 453.
 Artand, 493.
 Ashford, 361.
 Aslanian, 468.
 Atlee, 467.
 Atthill, 418.
 Aveling, 420.

 BABES, 177.
 Babesin, 468.
 Bacon, 544.
 Baer, 461, 468.
 Baldy, 357.
 Ballantyne, 8, 228, 531.
 Bandl, 47, 57, 225, 376.
 Bannon, 580.
 Bantock, Granville, 280, 374,
 401, 431, 456, 460.
 Barbier, 575.
 Barbour, A. H. F., 585.
 Barker, A. E., 609.
 Barker, F., 309, 322, 496, 505.
 Barkow, 580.
 Barnes, 124, 125, 358, 413, 417,
 433, 476, 501, 505, 507, 569.
 Barth, 230.
 Battey, 99, 239, 244, 245, 358,
 640.
 Baumgärtner, 455.
 Bayle, 436.
 Beates, 468.
 Beatson, 245.
 Beck, Snow, 365.
 Bedson, 551.
 Beigel, 26, 573.
 Bell, 358.
 Bennett, Hughes, 252, 253.
 Bennett, J. Henry, 174, 319.
 Bérard, 663.
 Berthold, 580.
 Beyea, 528.
 Billroth, 364, 434, 459.
 Bischoff, 516.
 Bird, 361.
 Blau, 493.
 Blundell, 245.

 Boldt, 227, 675.
 Bourdon, 206.
 Bozeman, 121, 148, 665, 669.
 Braun, Carl, 369, 514.
 Braune, 55, 71.
 Breisky, 32, 496, 551, 573, 590.
 Brennecke, 351, 452, 461.
 Breutlecht, 406.
 Brewis, 413.
 Brinton, 74.
 Brown, 420.
 Brown, Baker, 275.
 Brunton, Lauder, 154.
 Budin, 8, 31.
 Buechler, 521.
 Bulius, 432.
 Bumm, 177, 184, 228, 348, 633.
 Bunsen, 178, 179.
 Burton, 301.
 Byford, 465, 662.
 Byrne, 418.

 CAMERON, 382.
 Campbell, H. F., 393.
 Cappie, 427.
 Carron, 493.
 Cazin, 544.
 Chadwick, 40.
 Championnière, 76.
 Champneys, 672.
 Charnier, 227.
 Chéron, 571.
 Chiara, 280.
 Chiari, 226.
 Chisholm, 496.
 Chrobak, 298, 531.
 Cintrat, 458, 462.
 Clarke, 476.
 Claudius, 55.
 Clay, 534.
 Cleveland, 406.
 Clouston, 708.
 Coats, 288.
 Cobbold, 217.
 Coblenz, 255, 260.
 Coghill, 669.
 Cohnstein, 359.
 Collis, 600.
 Colman, 531.
 Conrad, 533.
 Cornil, 177, 347, 365, 482.
 Coulomb, 700.
 Courty, 417, 419.
 Credé, 358, 413, 556.
 Cripps, Harrison, 461.
 Crookshank, 177.
 Croom, 478, 641.
 Cross, 552.
 Crosse, 409, 412, 414.
 Cruveilhier, 55, 438.

 Cullingworth, S., 191, 256, 560.
 Cunningham, 14, 104.
 Cusco, 121, 125, 126, 127, 147.
 Cushing, 464.
 Cutter, 448.
 Czempin, 321, 327.
 Czerny, 405, 521.

 DA COSTA, 573.
 Dalton, 95, 99.
 Davaine, 161.
 Davy, 549.
 De Cassis, Vidal, 676.
 Delagénère, 464.
 De Sinéty, 24, 255, 256, 339, 347,
 364, 474, 477.
 Diesterweg, 468.
 Dobronrawow, 447.
 Döderlein, 32, 163, 349, 462,
 562.
 Dohrn, 26, 551.
 Doléris, 177, 349, 552.
 Donat, 251.
 Donald, 461.
 Doran, 16, 230, 231, 239, 252,
 256, 258, 425, 440, 469.
 Dott, 162.
 Doyen, 463, 517, 518, 697.
 Drysdale, 252, 253, 468.
 Duke, 302.
 Dührssen, 327, 402, 406, 574.
 Düll, 427.
 Dumontpallier, 356.
 Dumoret, 405.
 Duncan, John, 500.
 Duncan, J. W., 582.
 Duncan, Matthews, S., 52, 281,
 299, 300, 340, 409, 417, 438,
 488, 528, 588, 589, 592, 622,
 626, 630, 631.
 Duncan, W., 406, 516.
 Dunlap, 642.

 EADE, 439.
 Eckart, 491.
 Edebohls, 432.
 Edge, 463.
 Eichwald, 252.
 Ehrendorfer, 433.
 Ellinger, 302.
 Emmet, 119, 157, 299, 300, 319,
 322, 324, 326, 327, 335, 365,
 373, 417, 418, 556, 588, 592,
 595, 602, 638.
 Engelmann, 97, 99.
 Englisch, 242.
 Engström, 306, 308.
 Eppinger, 508, 526.
 Erich, 468.
 Esmarck, 588.

- FARRE, 8, 26, 28.
 Faucon, 416.
 Fehling, 467, 522.
 Fenger, 468, 573, 574.
 Fenwick, 431, 646.
 Fere, 493.
 Fergusson, 121, 124, 126, 127, 147.
 Fischel, 321, 333, 571.
 Foerster, 404.
 Foster, 9, 59, 70, 71.
 Fothergill, Milner, 621.
 Foulis, 156, 252, 255, 258.
 Fox, Wilson, 254.
 Fowler, J. K., 236.
 Fränkel, 374, 386, 456, 476, 513.
 Frankenhauser, 79.
 Freund, H. W., 281.
 Freund, W. A., 281, 319, 410, 425, 516, 545, 572, 662, 674.
 Fritsch, 46, 151, 196, 300, 334, 356, 364, 378, 385, 408, 409, 452, 459, 521, 567.
 Frommel, 227.
 Fuld, 555.
 Fürst, 488.

 GALLIPPE, 434.
 Garrigues, 37, 253.
 Gawronsky, 79.
 Gebhard, 528.
 Geith, 588.
 Geppert, 160.
 Gilmore, 270.
 Goltz, 154, 656.
 Goodell, 270, 279, 302, 607.
 Goodman, 93.
 Gottschalk, 305, 349, 437, 544, 546.
 Gough, James, 462.
 Graenicher, 574.
 Gram, 178, 179.
 Graefe, 403.
 Granville, 451.
 Gray, 293.
 Griffith, 502.
 Griffiths, 79, 257.
 Gross, 632.
 Guérin, 44, 51, 76.
 Gulland, 640.
 Gusserow, 406, 427, 434, 436, 439, 492, 496, 498, 505, 512, 516, 522, 531, 534, 536, 625.
 Gwilt, 588.

 HABIT, 505.
 Hæckel, 588.
 Hagedorn, 149.
 Hamilton, 270.
 Hanot, 439.
 Hart, 46, 66, 88, 89, 393, 551, 607.
 Hasse, 59, 60, 61.
 Haultain, 424.
 Heape, 93, 99.
 Hebra, 585.
 Hegar, 139, 143, 245, 459, 481, 570, 575.
 Heinricius, 346, 351.
 Henderson, 293.
 Henle, 13, 32, 35, 42, 55.
 Hennig, 19, 225, 299, 413, 432.
 Henrichsen, 569.
 Herman, 387, 502, 604.
 Hewitt, Grailly, 175, 376, 381, 382.
 Hicks, Braxton, 372, 417, 418, 471.
 Hildebrandt, 447, 565, 567, 569, 580, 585.
 Hilton, 679, 682.

 Hirschwald, 517.
 His, 25, 49, 61.
 Hoehenegg, 521.
 Hodge, 394, 396.
 Hofmeier, 95, 339, 508, 515, 523.
 Holden, 590.
 Holdhouse, 439.
 Holl, 37.
 Holländer, 252.
 Horrocks, 443.
 Howard, 156.
 Houston, 40.
 Hubert, 440.
 Hüe, 439.
 Huguier, 582, 588, 589, 607.
 Hunter, William, 178.
 Huser, 426.
 Hutchinson, J., 274, 529, 589.
 Hyde, 591.
 Hyrtl, 40, 73, 74.

 IMMERWAHR, 349.
 Imlach, 405.

 JACOBI, 93.
 Jacobash, 533.
 Jaggard, 420.
 Jastrow, 79.
 Jessett, 502, 514, 521, 523.
 Jobert, 660.
 Johnston, 433, 571.
 Johnstone, 214.
 Jones-Dixon, 462.
 Joseph, 37.

 KALTENBACH, 359, 491, 522, 659.
 Kaschkaroff, 366.
 Kehrner, 556, 633.
 Keibel, 79.
 Keith, Thos., 157, 273, 275, 276, 280, 449, 456, 463.
 Kjoergaard, 450.
 Kelly, 90, 116, 169, 404, 459, 465, 522, 644.
 Kidd, 439.
 Kiderlen, 559.
 Kimbal, 451.
 Kiwisch, 362.
 Klebs, 425, 484, 531.
 Kleeberg, 459.
 Klein, 74, 95, 322.
 Klein and Smith, 95.
 Kleinschmidt, 531.
 Kleinwachter, 327, 426, 434, 440, 453, 560, 632.
 Klemm, 359.
 Klob, 308, 362, 436.
 Klotz, 333, 404.
 Koch, 159, 160, 171, 191, 228.
 Koeberle, 451, 456, 467, 471.
 Kohlrausch, 59.
 Koller, 157.
 Kölliker, 206.
 König, 47.
 Kottmann, 454.
 Kötschau, 431.
 Krohne, 278.
 Kroner, 418.
 Krönig, 562.
 Krugenberg, 522.
 Kuchenmeister, 303.
 Kugelmann, 358.
 Kulenkampff, 588.
 Kundrat, 97, 99.
 Küster, 430.
 Küstner, 19, 20, 24, 322, 339, 350, 381, 405, 474, 488.

 LACKIE, 559.

 Lancereaux, 493.
 Landau, 235, 332, 347, 491, 517, 697.
 Landerer, 528.
 Landowzy, 434.
 Langenbuch, 690.
 Lawson, 568.
 Le Bec, 51, 76, 463, 464.
 Lebedinsky, 239.
 Lécorché, 585.
 Lee, 280.
 Leech, 190.
 Leiter, 645.
 Leopold, 23, 77, 93, 98, 344, 347, 404, 425, 452, 461, 467, 517, 522, 528, 556.
 Lewers, 468, 515.
 Lister, 143, 157, 159, 279.
 Lizers, 451.
 Löwenhardt, 99.
 Lorey, 425.
 Lott, 333.
 Löwenstein, 32.
 Luscha, 34, 37, 55.
 Lusk, 78.

 MACAN, 401.
 M'Clintock, 441.
 Macdonald, Angus, 296, 361, 417, 588.
 Mackenrodt, 402, 406, 523.
 Mackintosh, 299, 301.
 Madden, More, 302, 552.
 Mahomet, 280.
 Malassez, 255, 256.
 Malcolm, 278.
 Malgaigne, 502.
 Malins, 419.
 Marchand, 230.
 Marchesi, 430.
 Marckwald, 313.
 Martin, 226, 227, 230, 298, 324, 327, 341, 432, 455, 462, 481, 520, 696.
 Massin, 350.
 Maunsell, 462, 464.
 Mayer, 588.
 Meadows, 400.
 Meinert, 326.
 Meyer, 580.
 Milander, 405.
 Milton, 461, 572.
 Miner, 279.
 Mironoff, 93.
 Mitchell, Weir, 370.
 Möricke, 99.
 Morison, Rutherford, 592.
 Müller, 300, 406, 467, 531, 633.
 Mundé, 243, 270, 322, 339, 391, 406, 419.
 Murphy, 439.
 Murray, Milne, 449, 450, 697, 702.

 NEGRI, 403.
 Neisser, 177, 564.
 Nélaton, 373.
 Neugebauer, 121, 125, 127, 147, 421.
 Nieberding, 322.
 Nitze, 645.
 Nöggerath, 184, 255, 256, 363, 417, 641.

 OHM, 700.
 Oldham, 473.
 Oliver, 565.
 Olshausen, 240, 250, 298, 345, 352, 404, 436, 521, 574.
 Orloff, 426.
 Otto, 581.

- PAGE, 557.
 Pallen, 322, 324.
 Palmer, Dudley, 406, 633.
 Park, 359.
 Parkes, 638.
 Parsons, 450.
 Patenko, 254.
 Paterson, 177, 288.
 Pawlik, 90, 514, 643.
 Péan, 273, 275, 451, 454, 458, 462, 517.
 Pechey-Phipson, 93.
 Peckham, 588.
 Peterson, 404.
 Petit, 692.
 Pfannenstiel, 349, 533.
 Pflüger, 256.
 Pichevin, 177.
 Pick, 427, 533.
 Picot, 496.
 Pirogoff, 44, 51, 61.
 Playfair, 449.
 Polk, 37, 356, 358, 406.
 Poirier, 489.
 Popow, 432.
 Porak, 572.
 Pott, Percival, 242.
 Pozzi, 8, 408, 433, 453, 459, 482, 487, 500, 522, 570, 633.
 Price, 169, 464.
 Priestly, 571.
 Prochownick, 227.
 Profanter, 709.
 Puech, 559.
 Purcell, 522.
- QUAIN, 105.
- RAINEY, 22.
 Ranney, 14.
 Reamy, 426, 436.
 Recamier, 145.
 Reid, 126, 302.
 Rein, 468.
 Reinl, 93.
 Reinmann, 361.
 Richelot, 402, 517, 697.
 Richet, 19.
 Rieke, 426.
 Rieder, 26.
 Ries, 306.
 Ringer, 447.
 Rivington, 616.
 Robb, 165.
 Robinson, 230.
 Robson, Mayo, 406, 469.
 Rokitsky, 250, 253, 375, 453.
 Roser, 319.
 Röse, 432.
 Rosthorn, 227.
 Ross, 293.
 Routh, 400, 513, 572.
 Roux, 146.
 Ruedinger, 40, 54.
 Rüge, 33, 385, 486, 496, 504, 526, 545, 565.
 Rühl, 403.
 Runge, 420, 447.
 Russell, 490.
 Rutherford, 447, 571.
- SÄNGER, 230, 251, 280, 402, 406, 459, 465, 537, 544, 600.
 Sappey, 19, 73.
 Saurenhans, 491.
 Savage, 42.
 Sawinoff, 226.
 Säxinger, 506.
 Scanzoni, 298, 361, 362, 408.
 Schauta, 227, 406, 440, 673.
 Scharlieb, 461.
- Schatz, 604.
 Schimmelbusch, 165, 167.
 Schlesinger, 47.
 Schmalfluss, 419.
 Schorler, 426.
 Schottländer, 426.
 Schroeder, 70, 71, 256, 324, 341, 344, 361, 376, 419, 425, 427, 436, 440, 460, 464, 474, 496, 505, 515, 524, 549, 585, 607.
 Schuckhardt, 574.
 Schücking, 402.
 Schultze, 59, 140, 238, 302, 327, 376, 392, 400, 401, 428, 562.
 Schütz, 609.
 Schwartz, 192.
 Semb, 430.
 Ségond, 454, 697.
 Seelig, 489, 490, 512, 524.
 Seeligmann, 633.
 Semeleder, 270.
 Senn, 572.
 Seseman, 278.
 Seyfert, 506.
 Shober, 404.
 Sibley, 496.
 Simon, 114, 116, 145, 146, 571, 641, 664, 676.
 Simpson, A. R., 88, 89, 129, 135, 183, 306, 327, 393, 408, 411, 419, 447, 453, 507, 531, 532, 534, 536, 556, 573, 598, 600, 641, 677.
 Simpson, Sir J. Y., 128, 137, 157, 299, 302, 306, 308, 358, 362, 432, 448, 494, 505, 510, 527, 580, 588, 590, 630, 631.
 Sims, 90, 120, 121, 145, 299, 300, 404, 568, 631, 663.
 Sinclair, 461, 523.
 Skene, 34, 604, 626, 653.
 Sköldberg, 340.
 Skutsch, 430.
 Slavjansky, 239, 254, 347.
 Smith, Albert, 394, 396.
 Smith, Heywood, 389, 461, 552.
 Smith, Laphorn, 406.
 Smith, J. C., 533.
 Smith, Tyler, 417.
 Smyly, 462.
 Solowieff, 625.
 Spencer, 543.
 Spiegelberg, 45, 51, 256, 259, 468, 532, 574.
 Steavenson, 449.
 Stein, 600.
 Stephenson, 93.
 Stewart, Grainger, 228.
 Stiegele, 588.
 Stiles, 515, 524.
 Stirling, A. B., 161.
 Storer, 680.
 Strassmann, 403.
 Stratz, 239, 502.
 Sutton, 8, 25, 99, 230, 252.
 Sylvester, 156.
 Syme, 157, 641.
 Symington, 40.
- TAIT, LAWSON, 74, 93, 99, 139, 143, 227, 236, 239, 240, 250, 276, 280, 413, 420, 428, 456, 460, 465, 599, 615, 689, 692.
 Tate, 417.
 Taylor, 406, 409, 463, 588.
 Terrillon, 588.
 Thelen, 465.
 Thiersch, 81, 484.
- Thin, 589.
 Thiry, 420.
 Thomas, 145, 146, 270, 338, 362, 417, 419, 453, 558.
 Thorn, 432.
 Thornton, 258, 276, 280, 281, 448, 456, 457.
 Tillaux, 429, 436.
 Tilt, 175, 570.
 Timmers, 427.
 Tinns, 427.
 Tripiet, 448, 697.
 Turner, 40, 427.
- UFFELMAN, 34.
 Underhill, 473.
 Unna, 585.
- VAN BUREN, 681.
 Van de Warker, 61, 513.
 Vedeler, 299, 300, 387.
 Veit, 24, 331, 333, 455, 486, 489, 496, 504, 526, 550, 572, 574, 636.
 Velpeau, 206.
 Virchow, 51, 363, 375, 434, 468, 476, 484, 529.
 Volta, 700.
 Von Grunewaldt, 298, 632.
 Von Kahlen, 531.
 Von Hacker, 459.
 Von Ott, 432, 452.
 Von Preuschen, 26, 32, 572.
 Von Rabenau, 402.
 Voss, 600.
 Vulliet, 359.
- WALDEYER, 28, 59, 81, 248, 256, 484.
 Walker, Norman, 585.
 Wallace, 646.
 Walthard, 562, 563.
 Warner, 622.
 Weberstedt, 403.
 Webster, J. C., 584.
 Wells, Spencer, 192, 273, 274, 276, 280, 282, 327, 456, 516, 517.
 Werth, 251, 281, 296, 357, 406.
 Wertheim, 177, 184, 227, 228, 402, 406.
 West, 281, 588.
 White, 417, 418.
 Whitehead, 630.
 Widai, 177.
 Wiedow, 465.
 Williams, J., 73, 96, 97, 99, 228, 365, 474, 481, 488, 531, 625.
 Williams, Roger, 494.
 Williams, Whitridge, 229, 481, 482, 487, 496, 502, 532.
 Williams, Wynn, 400, 513.
 Wiltshire, 281.
 Winckel, 225, 250, 382, 434, 565, 571, 636, 641, 662.
 Winkler, 532.
 Witzel, 675.
 Wölfler, 459.
 Winter, 489, 523.
 Woodhead, 161.
 Wyder, 429.
 Wyllie, Gyl, 406.
- ZELLER, 346.
 Ziegenspeck, 372, 373.
 Ziemssen, 706.
 Zinke, 324.
 Zinnstag, 552.
 Zwanck, 611.
 Zweifel, 402, 451, 461, 465, 516, 565.
 Zweigbaum, 575.



INDEX OF SUBJECTS.

Note.—As the student will find in the *Table of Contents* the sub-divisions of each special subject, we give only one reference—in bold figures—to the places where each subject is specially treated of. References in other parts of the book are given in detail.

Thus, for the topics considered under "Carcinoma Uteri," see that subject in the *Table of Contents*; for references to Carcinoma, not contained in the chapters on Carcinoma, see Index.

For instruments *figured* in text, see *Instruments in Classified List of Illustrations*.

- ABDOMEN**—distension of, in ovarian tumour, 264.
 physics of, **85**.
Abdominal contents, relation to skin, 104, 105.
 examination, **102**.
 pressure, action on uterus, 372.
 regions, 104.
 Section, **688**.
 possible accidents, 694.
 for pelvic abscess, 214.
 for carcinoma, 516.
 for fibro-cystic tumours, 469.
 for fibroids, 454.
 in hæmatocele, 214.
 for ovarian tumours, 270.
 for removal of ovaries, 270.
Abortion and endometritis, 345, 351, 354.
 in lacerated cervix, 322.
 causing pelvic cellulitis, 194.
 ovaritis, 240.
 peritonitis, 183.
 and fibroids, 423, 437, 440.
 and metritis, 366.
 in retroflexion, 388.
 cause of subinvolution, 366.
Abscess—pelvic, 194, 214, **693**.
 opened by cauterization, 198.
 tapping of, 197.
 of uterine wall, 361.
Actinomyces bovis, 177.
Adenoma, 347, 470, 474, 483, 488.
 malignum, **528**.
Adhesions—in anteversion, 367.
 in fibroids, 427.
 in retroflexion, 387, 392.
 in retroversion, 382.
 diagnosis in ovarian tumours, 268.
 produced by ovarian tumours, 280.
 treatment in ovarian tumours, 276.
Age—influence on carcinoma, 494.
 on sarcoma, 533.
 on fibroid tumours, 436.
 on deciduoma malignum, 542.
 in relation to hæmatocele, 209.
 to menstruation, 93.
Alexander-Adams' operation, 405, 614, 616.
Amenorrhœa, **620**.
 due to atresia, 553, 620.
 in chlorosis, 620.
 and endometritis, 351.
 phthisis, 620.
 physiological, 620.
 in superinvolution, 308.
Amœbæ causing endometritis, 348.
Ampère, 700.
Ampulla of Fallopian tube, 25.
Amputation—of cervix, 317, 369, 514.
 hypertrophied in prolapsus, 316.
 of inverted uterus, 421.
Anatomical factors in pelvic disease, 175.
Anæmia in hæmorrhagic endometritis, 352.
 carcinoma, 501.
 cervical catarrh, 338.
 inversion, 413.
Anæsthetics, **153**.
Anteflexion of uterus, 373, **374**.
 and small fibroid tumour, 443.
 pathological, 115, 201.
 mistaken for retroversion, 374, 383.
 passage of sound in, 134.
Anteversion of uterus, 373, **380**.
 due to chronic metritis, 381.
Antiseptics in gynecology, 159, **688**.
Antiseptic douche, 151, 339.
 injections in carcinoma, 511.
Anti-streptococcic serum, 199.
Anti-toxines in treating septic conditions, 198.
Anus, 40.
 fissure of, 681.
 producing vaginismus, 568.
Apostoli's method of electrical therapeutics, **697**.
Areolar hyperplasia—of cervix, 335.
 of uterus, 363.
Arsenic in membranous dysmenorrhœa, 627.
Ascites and ovarian tumours, 266, 267.
Ascitic fluid in malignant ovarian disease, 252, 255, 257.
Asepsis, **162**.

- Asphyxia from chloroform, 156.
 Atresia of cervix uteri, **297**.
 senile, 348.
 of hymen, 548.
 in septate uterus, **558**.
 of vagina, **548**.
 operation for, 558.
 Atrophy of cervix and uterus, **305**.
 congenital of uterus, 291, 292, 296.
 senile, 308.
 Auscultation, 107.
 in fibroids, 444.
 in ovarian tumour, 265.
 Axial Coronal Section, 54, 67.
 BACTERIA, in cellulitis, 194.
 in vaginitis, 562.
 Bacterium coli commune, 349.
 Baden-Baden waters in chronic metritis, 368.
 Bantock's knot, 276.
 Barnes' ointment-positor, 310.
 Bartholinian glands, 14.
 abscess in, 586.
 cysts of, 586.
 inflammation of, 582.
 Battey's operation, 99, 245.
 in dysmenorrhoea, 627.
 in fibroids, 465.
 Benzoate of ammonia in cystitis, 653.
 Bimanual Examination, **109**.
 in antelexion, 379.
 in early pregnancy, 367.
 in fibroids, 441, 442, 444.
 in inversion, 415.
 position of hands in, 110.
 to replace retroflexed uterus, 392.
 in retroflexion, 389.
 in retroversion, 383.
 with sound, 135, 312.
 for stone in bladder, 655.
 in uterine displacements, 374.
 Biniodide of Mercury as an antiseptic, 162.
 Bipolar method of applying electricity, 701.
 Bladder, Anatomy of, **33, 636**.
 affection of, in carcinoma, 491, 492, 493.
 in amputation of cervix, 317.
 calculi in, 654.
 changes in, after fistula, 660.
 changes in position of, 38.
 diastole of, 39, 637.
 dilated in carcinoma, 494.
 displacements of, **650**.
 in retroflexed gravid uterus, 650.
 distended, and ovarian tumours, 266.
 functional diseases of, **655**.
 injuries to in ovariectomy, 276.
 malformations of, **650**.
 methods of exploring, **639**.
 neoplasms of, **650**.
 openings, 35.
 perforation by calcified fibroids, 441.
 peritoneal relations, 317.
 physiology of, 637.
 position, 33.
 changes in, 38.
 in inversion of uterus, 412.
 pressure on, by fibroid tumour, 439.
 in retroflexion, 372, 385, 386.
 shape when empty, 38.
 in foetus, 39.
 structure, 35.
 systole and diastole of, 637.
 effect of its distension on the uterus, 59, 61, 372, 381.
 Bland's pills in chlorosis, 621.
 Blistering of the cervix uteri, 369.
 Blood effusion into pelvis, 206.
 in broad ligaments, 207.
 and ovarian tumour, 263.
 Blood-vessels of pelvis, **72**.
 Bougies graduated to dilate cervix, 301.
 Bougies in antelexion, 380.
 in fibroids, 444.
 Bozeman's scissors, 148.
 speculum, 121.
 Brandt's pelvic gymnastic, 708.
 Bright's disease in fibroid tumour, 440.
 British Association (B.A.) electro-magnetic units, 700.
 Broad ligament, 44.
 affections of, 215.
 anatomy, 21, 50.
 blood effusion into, 207.
 connective tissue of, 46, 50.
 cysts of, **259**.
 new growths in, **215**.
 position, 50.
 tumours of, 215.
 and fibroids of uterus, 443.
 Bromide of Potassium in chronic metritis, 369.
 in fibroid tumours, 448.
 in ovaritis, 241.
 Bromine solution in carcinoma, 511.
 Bulbo-cavernosi, 12.
 Bulbi vaginae, 13, 83.
 CACHEXIA in carcinoma, 501.
 in sarcoma, 534.
 Calcification of fibroid tumours, 427, 432.
 Calculi in bladder, **694**.
 Cancer (see Carcinoma).
 Carbolic acid—activity of, 161.
 in cervical catarrh, 399.
 in endometritis, 355.
 in metritis, 362.
 in pruritus vulvæ, 585.
 Carcinoma—of bladder, 650.
 of cervix, **483, 499, 509**.
 cure by amputation, 514.
 commencement in cervix, 338.
 development, 487.
 duration of life in, 505.
 and endometritis, 353.
 extension to neighbouring organs, **491**.
 of Fallopian tubes, 230.
 affecting fibroid tumours, 433.
 contrasted with fibroid, 423, 502.
 in male and female, compared, 494.
 of ovary, 257.
 combined with sarcoma, 531.
 diagnosis from sarcoma, 535.
 microscope in diagnosis of, 502.
 spontaneous cure of, 505.
 spread of, 488, 512.
 of body of uterus, **525**.
 of syncytium, 545.
 of vagina, **573**.
 secondary, 503.
 of vulva, 588.
 Carlsbad salts, 369, 621, 685.
 Case-taking, method of, **710**.
 A. R. Simpson's card for, 710.
 Cataleptic convulsions and lacerated cervix, 323.
 Catarrh of cervix, **329**.
 and carcinoma, 496.
 laceration, 321, 335.
 retroflexion, 335.
 diagnosis from vaginal, 336.
 of uterus, 340.
 Catarrhal patches, 333, 352.
 Catgut suture—continuous, 600.
 Catheter—methods of passing, 640.
 Skene-Goodman, 653.
 stationary, for fistula, 674.
 Cauliflower excrescence, 476, 502.
 Caustics producing atresia, 299.
 metritis, 366.
 in carcinoma, 513.
 in cervical catarrh, 341.
 Cautery in dividing cervix, 453.
 in incising capsule of fibroid, 453.

- Cautery in ovariectomy, 275.
 Paquelin's, 153.
 in opening vaginal cysts, 572.
- Cellular tissue of pelvis, hæmorrhage into, 210
 (v. Connective tissue).
- Cellulitis, pelvic, 193.
 and antelexion, 200, 373, 376, 379.
 producing lateriversion, 201.
 and lacerated cervix, 321.
 ovarian tumours, 263.
 subinvolution, 365.
 relation to peritonitis, 182, 196.
 and retroflexion, 390.
 in utero-sacral ligaments, 201,
 376, 379.
- Cervical catarrh in lacerated cervix, 321.
 chronic, 329.
 with retroflexion, 384.
 endometritis, 329.
 mucus, character of, 337.
 polypus, 473.
- Cervix—amputation of, 313, 317.
 flap operation, 313.
 in carcinoma, 514.
 compared with extirpa-
 tion of uterus, 522.
 in prolapsus, 316.
 producing involution,
 369.
- anatomy, 18.
 areolar hyperplasia, 362.
 atrophy of, 305.
 atresia of, 397, 554.
 operation for, 558.
 cancer of (see carcinoma).
 carcinoma of, 483, 499, 509.
 catarrh of, 329.
 chronic thickening of, 298, 311.
 conical, 298, 301, 311.
 closure for vesico-uterine fistula, 671.
 connective tissue, increase of, 355.
 cysts of, 334.
 degeneration, granular and cystic, 335,
 338.
 descent in prolapsus, 507.
 dilatation of, 138, 301, 453, 473.
 division of, 302, 380.
 antero-posterior, 380.
 bilateral, 302, 380.
 examination of, 108.
 excision of, 304, 341.
 fibroid tumour of, 423, 433, 465.
 glands of, 24, 335.
 hypertrophy of, 298, 310, 340.
 three forms of, 316.
 in cancer, 503.
 in prolapsus, 609.
 induration of, in carcinoma, 503.
 inflammation, 335.
 in infantile uterus, 291, 296.
 position in inversion, 410, 414.
 laceration of, 318.
 producing cellulitis, 194.
 cervical catarrh,
 335.
 parametritis, 202.
- lymphatics, 76.
 medication in catarrh, 339.
 mucous membrane of, 24, 331, 335.
 normal structure contrasted with patho-
 logical, 334.
 obliteration of canal, 441.
 papilloma of, 470.
 position in antelexion, 376.
 retroversion+retroflexion,
 384.
 rigidity of, 298.
 rupture in atresia, 549.
 sarcoma of, 527.
 stenosis of, 298.
 stitching, 324.
 in superinvolution, 305.
- Cervix—supra-vaginal amputation of, 515.
 tubercular disease of, 482.
 "ulceration" of, 323, 331.
 ulcerations, true, 334.
 vaginal portion of, 18.
- Champignons cancéreux, 502.
- Channelled polypus, 473.
- Childbirth producing cellulitis, 194.
 peritonitis, 183.
- Chill producing cellulitis, 194.
 ovaritis, 240.
 peritonitis, 184.
- Chloral in carcinoma, 511.
- Chloride of zinc in carcinoma, 514.
 vaginitis, 567.
- Chloroform—action of, 153.
 administration of, 155.
 dangers from, 156.
 uses of, 154.
- Chlorosis, amenorrhœa in, 620.
 leucorrhœa in, 350.
 small uterus in, 291, 305.
- Cholera and endometritis, 350.
- Chronic acid—in cervical catarrh, 339.
 in endometritis, 356.
- Chronic cervical catarrh, 329.
 metritis and fibroid tumour, 443.
- Cintra's serre-nœud, 458.
- Clamp in ovariectomy, 274.
 disadvantages of, 274.
 in laparotomy for fibroids, 458.
- Classification of diseases of women, 714.
- Clitoris—anatomy, 7.
 development of, 579.
- Cloaca, 580.
 persistence of, 580.
- Cocaine, 157.
 in vaginismus, 569.
- Coccygeus, 15.
- Coccygodynia, 685.
- Cœliotomy, 688.
- Colpitis, 561.
 emphysematosa, 565.
- Colpoperineorrhaphy, 614.
- Colporrhaphy, 601, 616.
- Colpotomy anterior and posterior, 695.
- Conception in an undeveloped horn, 293, 296.
- Constant electric current, 698, 701.
- Connective tissue of pelvis, 46, 47, 50, 51.
 as seen in sections, 47, 50, 51.
 methods of studying, 46.
 new growths in, 215.
 relation to cellulitis, 193.
 spread of carcinoma in, 488,
 491, 493.
 tumours of, 217.
 of uterus, increase of in
 metritis, 363.
- Connective-tissue origin of cancer, 484.
 sarcoma, 529.
- Constipation, 684.
 in fistula, 662.
- Copremia, 501.
- Coronal Section, 51.
- Corpus luteum of menstruation and pregnancy,
 95.
- Corrosive sublimate as an antiseptic, 161.
 in metritis, 362.
- Coulomb, 700.
- Cradle-pessary of Graily Hewitt, 381.
- Créde's method of expressing placenta, 413.
- Cretinism and amenorrhœa, 620.
 atrophy of uterus, 291.
- Curette, 145, 355.
 cases in which useful, 146.
 method of use, 146.
 Recamier-Roux, 146.
 Simon's, 145.
 Thomas', 146.
 in carcinoma, 527.
 in cervical catarrh, 341.
 in deciduoma malignum, 543.

- Curette, in endometritis, 353.
in diagnosis of small polypi, 479.
in sarcoma uteri, 535.
- Curetting—regeneration of mucosa after, 357.
- Cystic fibroid tumours, **466**.
leio-myoma, 468.
- Cystitis, **651**.
- Cystocele, 650.
in prolapsus, 610.
- Cysto-fibroma, 466.
- Cystoma malignum, 252.
- Cysts in the cervix, 334.
of vagina, **571**.
of vulva, 586.
- DECIDUA and Endometritis, 350.
decidua malignum, 546.
malignum, 476, **536**.
- Defæcation—mechanism of, 679.
difficult in hæmatocele, 212.
painful in carcinoma, 591.
in retroflexion, 387.
want of control, in ruptured perineum, 595.
- Dermoid cysts, 251.
of genito-urinary organs, 247, 259.
- Development—*intra-uterine* periods of, 291.
of myoma, 434.
of pelvic organs, **79**.
of the genito-urinary organs, 578.
of uterus, period of, 282.
relation to malformations, 283, 291.
- Developmental factors in pelvic disease, 175.
- Diabetes, pruritis in, 583.
- Diathesis—strumous and tubercular, 290, 354, 358.
syphilitic, 354.
- Diet in carcinoma, 511.
in peritonitis, 188.
Weir Mitchell's system of, 370, 705.
- Diffuse proliferation of connective tissue, 362.
- Digestive derangements in anteversion, 367.
carcinoma, 501.
endometritis, 351.
- Digital pressure in vaginal pressure, effects on uterine position, 91.
exploration of bladder, 640.
- Dilatation of cervix, **138**.
by rapid method, 480.
of uterine canal, 359.
- Dilators—Tait's and Hegar's, **143**.
various forms of, 301.
- Diphtheritic inflammation and carcinoma, 503, 505.
of vagina, 566.
- Discharge—fœtid, in carcinoma, 500.
of body of uterus, 527.
watery, in sarcoma uteri, 534.
- Discus proligerus, 29.
- Disinfection of hands and instruments, 163.
- Displacements of pelvic floor, 66, **603**.
of uterus, **371**.
due to cellulitis, 45, 201.
fibro-cystic tumours, 468.
hæmatocele, 212.
ovarian tumours, 264.
peritonitis, 200.
relation to chronic metritis, 366.
endometritis, 350.
diagnosed with sound, 134.
etiology, 373.
physiological and pathological, 372.
- Displacements of uterus produced by volsella, 118.
- Douche, forms of, 149.
in chronic metritis, 368.
- Douglas, pouch of, 44.
- Doyen's clamp for hysterectomy, 517.
speculum, 124.
- Drainage of abscess, 198.
in extirpation of uterus, 521.
for ovarian tumour, 270.
in ovariectomy, 277.
- Dressing of wound in ovariectomy, 277.
- Drop-cork for chloroform, 155.
- Dysmenorrhœa, **623**.
congestive, 624.
mechanical and congestion theories of, 377.
relation to Battey's operation, 245.
in antelexion, 377.
cellulitis, 199.
stenosis of cervix, 299.
endometritis, 351.
fibroid tumours, 436, 439.
uterine polypi, 476.
retroflexion, 387.
membranous, 623.
obstructive, 299, 377.
spasmodic, 624.
with sterility, 631.
- Dyspareunia in antelexion, 378.
in peritonitis, 185.
causes, 569.
definition, 569.
in prolapsed ovary, 245.
treatment, 569.
in vaginismus, 569.
- ECHINOCOCCI of pelvic organs, 216.
- Ecraseur—in amputating cervix, 512, 514.
mode of application, 514.
for removal of polypi, 481.
and galvano-cautery, relative advantages, 514.
- Ectropium of cervix, 321, 333.
- Elastic ligature in laparotomy for fibroids, 459.
- Electric endoscope, 645.
apparatus and instruments, **702**.
- Electricity in cervical catarrh, 341.
endometritis, 358.
fibroids, 448.
gynecology, **697, 703**.
metritis, 370.
nerve prostration, 705.
sterility, 633.
- Electrodes, 701, 702.
- Electrolysis, 701.
in ovarian tumour, 270.
- Electro-magnetic units, 700.
- Elythrytis, 561.
- Elythrorrhaphy, 573, 575.
- Emmet's operation for lacerated cervix, 324.
- Ems, waters of, in endometritis, 358.
in chronic metritis, 368.
- Endocervicitis, 330.
- Endometric applications, 356.
- Endometritis, **343**.
relation to abortion, 345, 351, 354.
in antelexion, 354.
cervical, **329**.
and cervical catarrh, 337.
leading to chronic metritis, 366.
classification, 348.
diagnosis by sound, 134.
fungosa of Olshausen, 345.
hæmorrhagic type and carcinoma of the body, 527.
and sarcoma, 535.
tubercular, 482.
varieties of, 344.
villous or papillary form, 347.

- Endoscope, electric, 645.
 Enterocoele, vaginal, 617.
 Enucleation of fibroid tumours, 425.
 spontaneous, 425, 429, 440.
 of ovarian tumours, 279.
 Episio-perineorrhaphy, 613, 615.
 Epispadias, 581.
 Epithelial cells found in carcinoma, 504.
 fibroids, 426.
 origin of cancer, 484.
 Epithelioma of the cervix, 484.
 and inversion, 413.
 Erector clitoridis, 14.
 Ergot and carcinoma, 510.
 chronic metritis, 369.
 endometritis, 354, 355.
 fibroids, 446.
 methods of administration in menorrhagia, 623.
 Ergotine—hypodermic injection of, 447.
 in treatment of fibroids, 447.
 Erosion—follicular, 332.
 and carcinoma, 496.
 papillary, 332.
 and carcinoma, 503.
 simple, 332.
 Esthiomène of vulva, 585.
 Ether, 153.
 Ethidene, 153.
 Eversion of lips of cervix, 321, 323.
 Examination of cases, 701.
 methods of, 102.
 posture in, 147.
 routine described in inversion, 415.
 Exanthemata and endometritis, 350.
 vaginitis in, 565.
 atresia vaginae after, 552.
 External genitals—anatomy, 5.
 development, 83, 578.
 relation in erect posture, 8.
 examination of, 107.
 lymphatics of, 76.
 Extirpation of cancer, 372.
 uterus—for fibroids, 456.
 through abdominal walls, 516.
 through vagina, 516.
 compared with amputation of cervix, 522.
 for sarcoma, 536.
 Extra-peritoneal blood effusion, 209.
 signs of, 212.
 ovarian tumour, treatment, 279.
 treatment of pedicle, 456.
 Extra-uterine gestation, 693.
 and fibroid tumour, 445.
 causing hæmatocele, 209.
 and ovarian tumours, 264.
 Exudations in the pelvis, 185.
 course of, 195.
 FACIES, cancerous, 501.
 Fallopian Tubes—anatomy, 25.
 abnormalities, 224.
 blood-sacs in, 549.
 catheterisation of, 224.
 development of, 82.
 distension with pus or blood, 226, 233, 277.
 divisions, 25.
 enlarged and fibroid tumours, 443.
 functions, 222.
 hydrops of, 226, 233, 235.
 infectious granulomata of, 228.
 inflammation of, 225, 233.
 new formations of, 238.
 palpation of, 224.
 patulous condition of, 225.
 position, 24, 49.
 in inversion, 410.
 removal by abdominal incision, 229, 237.
 per vaginam, 236.
 affected with sarcoma, 531.
 stricture of, 224.
 structure, 25.
 tubercular disease of, 228, 235.
 tumours of, 230.
 Faradic current, 698, 702, 703.
 Fatty degeneration around polypoidal fibroid, 408.
 in fibroid tumour, 432.
 Fertility, standard of, 630.
 Fevers causing ovaritis, 240.
 Fibro-cystic tumours of the uterus, 466, 691.
 and ovarian tumours, 267.
 Fibroid tumour of uterus, 422, 435, 446, 691.
 producing ante-flexion, 376.
 and affections of appendages, 432.
 and ante-flexion, 379.
 Battey's operation for, 245.
 compared with carcinoma, 502.
 epithelial structures in, 426.
 and chronic metritis, 367.
 and gestation in detached horn, 296.
 + inversion, 413.
 pediculated submucous, 470.
 rapidity of growth, 426.
 relation to age, 436.
 removal through abdomen, 451.
 per vaginam, 453.
 sloughing and carcinoma, 503, 505.
 and retroflexion, 389.
 sarcomatous degeneration of, 531.
 and sarcoma, 535.
 and Tait's operation, 236, 245, 465.
 tumour, spontaneous expulsion of, 473.
 tumours of vagina, 573.
 cervix, 423, 433.
 ovary, 256.
 Fibromata of vulva, 588.
 Fibro-myoma uteri, 423.
 lymphangiectodes, 467.
 Fibrous tumour of uterus (see Fibroid).
 Fimbriated end of Fallopian tube, 25.
 Flexions of uterus, 373.
 Flooding, producing superinvolution, 308
 (see Hæmorrhage).
 Foetal heart—in diagnosing pregnancy, 368, 444.
 Foetal life—malformations arising in, 291.
 Foetus in detached horn of uterus, 293.
 Forceps used for extracting mucous polypi, 480.
 and fistula, 662.
 and ruptured perineum, 594.
 Fornices—anatomy of, 31, condition of, in peritonitis and cellulitis, 195.
 effect on uterus of digital pressure in, 91.
 examination of, 109.
 lateral, 49.
 operation for tear into, 327.
 position of, in hypertrophied cervix, 280, 317.
 Fossa navicularis—anatomy, 8.
 irritability producing vaginismus, 568.
 Fourchette—anatomy, 6.
 fissures in, producing vaginismus, 568.
 Freund's extirpation of uterus, 516.
 Friedrichshall water, 369, 621, 685.
 Fritsch's method of treating pedicle of fibroid, 459.
 GALL BLADDER, laparotomy for distended, 692.
 Galvanic or "Galvano-caustic" current, 698, 701, 703.
 Galvanism, 309.

- Galvano-Cautic Wire in removal of, polypi, 481.
in amputating cervix, 312.
compared with écraseur, 514.
mode of application, 514.
- Galvanometer, 698, 703.
- Galvano-puncture, 702.
- Gangrene of fibroid tumour, 427.
- Gartner's canals, 26, 260.
and vaginal cysts, 572.
- Generative organs, development of, 259.
- Genito-urinary organs, 247.
- Genupectoral posture, 87.
in replacing retroverted uterus, 392, 393.
- Germicidal action of vaginal secretion, 563.
- Germinal vesicle, 29.
- Gestation—Abdominal, 693, 694.
in detached horn, 292, 296, 693, 694.
in Fallopian tube, 693.
in a septate uterus, 308.
- Glycerine plug, making of, 241.
- Gonococcus, 177, 564.
- Gonorrhœa—causing cervical catarrh, 335.
cystitis, 651.
ovaritis, 240.
peritonitis, 183.
salpingitis, 227.
vaginitis, 565.
and endometritis, 348, 350.
latent in the male, 184.
and metritis, 361.
micrococci in, 177, 349, 564.
- Gonorrhœal vaginitis, diagnosis from simple, 567.
- Gouty diathesis and dysmenorrhœa, 628.
- Graafian follicles—degeneration of, and ovarian cysts, 253.
development, 248.
distention of, and ovarian cysts, 253.
number, 28.
position, 28.
rupture of, 95.
structure, 29.
- Greenhalgh's intra-uterine stem, 380.
- Gynecological diseases, Etiology of, 174.
- Gynecology, Progress in, 174.
- HÆMATOCELE—pelvic, 206.
and hæmatoma, 206.
ruptured extra-uterine gestation, 210.
fibroid tumour, 445.
- Hæmatokolpos (see Atresia Vaginæ).
- Hæmatoma, 206, 212; of vulva, 590.
- Hæmatosalpinx, 229.
and fibroid tumours, 432.
- Hæmorrhage in carcinoma uteri, 499, 506, 508.
in carcinoma of body of uterus, 527.
causing death in carcinoma, 206, 508.
in carcinoma, treatment of, 510.
in endometritis, 348, 354, 357.
in deciduoma malignum, 542.
in fibroid tumours, 437, 445.
fatal from fibroid tumours, 431.
into bladder after operation for fistula, 674.
internal, in operation for atresia, 555.
intra-peritoneal and extra-peritoneal, 209.
in inversion of uterus, 413.
into the peritoneum of pelvis, 206.
in polypi, 476.
post-partum, 324.
in retroversion, 383.
- Hæmorrhage in sarcoma uteri, 534.
secondary after ovariectomy, 278.
in uterine polypi, 476.
- Hæmorrhoidal plexus, 74.
- Heart, action of chloroform on, 157.
- Hegar's amputation of cervix, 313.
dilators, 144, 480.
method for treatment of pedicle, 459.
- Heredity, influence on carcinoma, 494.
- Hermaphroditism, 580.
false, 580.
true, 580.
- Hernia and Hydrocele, 216.
of fibroid tumour, 428.
of ovary, 242.
- Hernial nature of prolapsus, 606.
- Hewitt's (Graily) cradle-pessary, 381.
- Higginson's syringe, 150.
- Hodge pessary, 394, 395, 398, 400.
mode of action, 398.
- Houston, valve of, 40.
- Hugnier's conoid amputation of cervix, 317.
- Hunyadi Janos water, 369, 685.
- Hydatid of liver and ovarian tumour, 267.
tumour of pelvis, 216, 692.
mole in deciduoma malignum, 476, 542.
- Hydramnos and ovarian tumour, 267.
- Hydrargyri pernitratil liquor in cervical catarrh, 339.
- Hydrastis Canadensis, 369.
- Hydrate of chloral and carcinoma, 511.
- Hydrocele of round ligament, 215.
- Hydronephrosis in cystitis, 651.
in carcinoma, 493.
- Hydrosalpinx, 226.
and ovarian tumours, 263.
- Hydrops folliculorum, 249.
tubæ, 226.
- Hymen—atresia of, 548.
development of, 83, 551.
forms of, 7, 8, 9, 10.
imperforate, operation for, 556.
persistent or inflamed, producing vaginismus, 568.
small ulcers in, producing vaginismus, 568.
- Hypodermic injection of ergotin, 447.
morphia, 189.
- Hyperplasia areolar of uterus, 363.
of cervix, 335.
- Hypertrophy—primary, of cervix, 310.
secondary, of cervix, 310, 314.
of mucous membrane of uterus, 429.
of muscular wall of uterus, 429.
- Hypospadias, 581.
- Hysterectomy for cancer, 521.
for cancer compared with amputation, 522.
for fibroids, 454, 456.
for inversion, 421.
in deciduoma malignum, 543.
vaginal with clamps, 517.
vaginal with ligatures, 520.
- Hysteria, 706.
due to ovaritis, 241.
in parametritis, 203.
small uterus in, 291.
and Battey's operation, 245.
in superinvolution, 308.
- Hystero-epilepsy, 718.
and Battey's operation, 245.
- ICE-BAGS in hæmatocele, 214.
- Ice-cap, 278.
- Ice in peritonitis, 191.
- Ichthyol, 241.
- Incontinence of urine, 656.
- Induced current of electricity, 698, 702.
- Induration of fibroid, 432.
- Infaret of uterus, 362, 432.
- Inflammation of cervix, 335.

- Inflammation of uterus, 330, 344.
 in muscular coat of uterus, 360.
 of pelvic peritoneum and cellular tissue, 182.
 puerperal, 361.
- Inflammatory deposits and fibroid tumour, 445.
- Infundibulo-pelvic ligament, 27.
- Inguinal glands, 76, 78.
 affection in carcinoma, 494.
- Injections—*intra-uterine* in carcinoma, 511.
 in endometritis, 354, 355, 358.
 and metritis, 361.
 into uterus and tubes, 225.
 dangers of, 225, 566.
 vaginal, 339.
- Interstitial fibroid, 427, 428.
 metritis, 363.
- Intestinal obstruction due to fibroid tumours, 439.
- Intestines—in pouch of Douglas, 61.
 relation to uterus and pelvic floor, 62.
- Intra-abdominal pressure upon pelvic floor, 68, 85.
 in inversion, 409.
 in prolapsus, 606, 608.
 and retroflexion, 386.
 action on rectum, 678.
- Intra-peritoneal blood effusion, 206.
 signs of, 212.
 treatment of pedicle, 460.
- Intra-uterine galvanic stem, 309.
 injections, 358.
 medication, 356.
 stem-pessary, 380, 396.
- Inversion of uterus, 408.
 caused by fibroid tumour, 430.
 diagnosed from fibroid tumour, 443.
 mucous polypus, 479.
 mechanism of, 408.
 and prolapsus, 417.
 in sarcoma, 532.
- Inverted uterus—amputation of, 421.
 replacement of—Atthill's method, 418.
 Barnes' method, 418, 419.
 Courty's method, 419.
 Emmet's method, 419.
 Lawson Tait's method, 420.
 Noeggerath's method, 418.
 Thomas' method, 418.
 White's method, 418.
- Involution and operations on cervix, 361.
 senile, 283.
 puerperal, 365.
 artificially produced by amputation of cervix, 369.
- Iodide of potassium in ovaritis, 241.
 in chronic metritis, 369.
- Iodine in cervical catarrh, 339.
 in endometritis, 356.
- Iodised phenol in endometritis, 358.
- Iodoform—in endometritis, 358.
 in vaginismus, 569.
 in pruritus vulvæ, 584.
- Ischio-rectal fossa, 11, 54.
- Isthmus—of uterus, 17.
 of Fallopian tube, 25.
- KEITH, clamp for fibroids, 458.
- Kelly's method of exploring bladder, 644.
- Kidney disease in carcinoma, 493.
- Kissingen waters in chronic metritis, 368.
- Knives, 148.
- Koeberlé's *serre-nœud*, 456.
- Kolpopleisis, 676.
- Kraurosis vulvæ, 689.
- Kreuznach waters in endometritis, 358.
 chronic metritis, 368.
 treatment of fibroids, 448.
- Kuchenmeister's scissors for dividing the cervix, 303.
- LABIA MAJORA, anatomy, 5.
 hydrocele of, 216.
- Minora, anatomy, 6.
 development, 579.
- Labour—atony of uterus in third stage of, due to chronic metritis, 366.
 "missed," 296.
 rapidity of, and lacerated cervix, 321.
 complicated by polypus, 478.
 protracted, producing fistula, 661.
 and vaginal fixation, 403.
 and ventral fixation, 405.
- Laceration of cervix, 318.
 and carcinoma, 496.
 cellulitis, 194.
 chronic metritis, 365.
 subinvolution 365.
 of perineum, 574.
- Lactation—effect on uterus if protracted, 308.
 and subinvolution, 366.
- Laminaria tents, 139, 142.
- Langenbuch's incision, 690.
- Laparotomy (see Abdominal Section).
- Lateri-version of uterus, 201.
- Leeches—application of, to cervix, 340.
- Lembert's suture, 695.
- Leucorrhœa—as a symptom, 711.
 in cervical catarrh, 332, 336.
 endometritis, 350, 351, 353.
 hypertrophied cervix, 312.
 lacerated cervix, 322.
 retroflexion, 387, 388.
 uterine polypi, 476.
 vaginosis, 566.
 vaginal cysts, 572.
 producing pruritus, 583.
- Levator ani, 15, 52, 54.
- Ligature in ovariectomy, 275.
- Lineæ albicantes, 103.
- Lipomata of vulva, 588.
- Listerism—in gynecology, 159, 688.
 in ovariectomy, 219.
- Literature, Gynecological, Sources of, 719.
- Lithia water in pruritus, 584.
 salts of, in irritable bladder, 655.
- Lithotomy posture, 167.
- Liver, fatty degeneration of, in fibroids, 431.
- Locomotion impaired in uterine inflammation, 366.
- Lupus vulvæ, 488, 588.
 and *ulcus serpiginosum*, 585.
- Lymph, coagulable—from fibroid tumour, 467.
- Lymphatics—relation to cellulitis, 194, 195.
 between glands and vessels 76, 77.
 of external genitals, 76.
 in fibroid tumours, 425.
 dilatation of, 467.
 of rectum, 78.
 and spread of cancer, 489.
 relation to septicaemia, 78.
 of uterus, 76.
 dilatation of metritis, 364.
 of vagina, 76.
- Lymphatic glands of pelvis, 76.
 in carcinoma, 489.
 vessels of pelvis, 76.
- MALFORMATION of uterus, 285.
 rectal examination in, 296.
- Malignant peritonitis, 193, 257.
 tumours of the uterus, 483.
- Marckwald's amputation of cervix, 313.
- Marriages, average productivity of, 630.

- Martin's operation for cervical catarrh, 324, 342.
enucleation of fibroids, 435
vaginal extirpation of cervix, 520.
- Massage, 191, **708**.
general in uterine disease, 370, 706, **708**.
in pathological antelexion, 380.
- Medullary cancer, 484.
- Membrana granulosa, 29.
- Menopause and Battey's operation, 245.
changes in cervix after, 299.
in uterus after, 305, 348.
in vagina after, 563.
influence on fibroid tumours, 426, 445
premature, 620.
- Menorrhagia, **622**.
in carcinoma, 499.
lacerated cervix, 322
metritis, 361.
endometritis, 351.
fibro-cystic tumour, 468.
fibroid tumour, 430, 437.
hæmatocele, 212.
inversion, 413.
ovaritis, 240.
peritonitis, 185.
uterine polypi, 476.
retroflexion, 387.
treatment by Battey's operation, 244.
- Menstruation, **92**.
and Battey's operation, 245.
irregular, in cervical catarrh, 336.
in laceration of cervix, 322.
diminished in chronic cellulitis, 203.
in endometritis, 350.
absence in fistula, 662, 677.
during gestation, 293.
relation to hæmatocele, 207.
in metritis, 361, 366.
periodicity and duration, 94.
descent of polypus during, 473.
in an undeveloped horn, 292.
in bicornuous and septate uterus, 292.
in monkeys, 99.
in atresic septate uterus, 559.
and Tait's operation, 236, 245.
in relation to pelvic disease, 176.
- Menstrual blood—composition, 94.
origin, 96.
quantity, 94.
character of retained, 551.
- Mesenteric tumours, 692.
- Metastatic deposits, 533.
- Metastasis in cancer, 523.
in deciduoma malignum, 542.
- Metritis, 340, 344, **360**.
acute, **360**.
chronic, 340, **360**, 374.
- Metrotome, 702.
Sir James Simpson's, 303.
- Micrococci in wounds, 159, 163, 194.
gonorrhœa, 178, 564.
- Micro-organismal causes of pelvic disease, 176.
- Micro-organisms in gynecology, 159.
in endometritis, 348.
in vaginitis, 562.
- Microscope—in carcinoma uteri, 502, 504, 505, 527.
endometritis, 353.
sarcoma uteri, 535.
- Micturition, 637.
difficult, in fibroid tumour, 439.
hæmatocele, 212.
prolapsus, 607
frequent, 655.
in cystitis, 651.
fibroid tumour, 439
- Micturition, frequent in vaginitis, 566.
painful in carcinoma, 501.
- Mobility of uterus, ascertained by sound, 134.
- Monopolar method of applying electric current 701.
- Morcellement in fibroid tumours, 454.
- Morgagni, columnæ and sinus, 41.
hydatis, 266.
- Morphia—methods of administration, 189.
in carcinoma, 511.
metritis, 362
caution as to use in dysmenorrhœa, 627.
- Mucoid degeneration in fibroid tumour, 468.
- Mucous membrane of bladder, 36.
cervix, 24.
rectum, 40.
urethra, 34.
uterus, 23.
vagina, 31, 561.
polypi, 470, 473.
- Müller, ducts of, 79, 81, 259, 291, 578, 579.
- Musculature of bladder, 35.
rectum, 40.
urethra, 34.
uterus, 22.
- Myoma of ovary, 256.
uteri, **423**.
retroflexion, 390.
and antelexion, 379.
- Myxomatous degeneration of fibroids, 432.
- Myxomyoma, 468.
- NABOTHII ovula, 334.
- Nabothian follicles, 334.
pediculated, 470, 475.
- Needles, **149**, 326, 459.
- Nelaton's forceps, 241.
- Nerve-prostration, systematic treatment of, 704.
- Nerves—of pelvis, **79**.
- Nervous derangements in anteversion, 367.
endometritis, 352.
reflex symptoms, and lacerated cervix, 323.
retroflexion, 386.
system, action of chloroform on, 154.
- Neugebauer's operation for prolapsus, 577.
- Neuralgia from pressure by fibroid tumours, 439.
in lacerated cervix, 322.
- Neuromata of vulva, 586.
- Nitrate of silver in endometritis, 358.
vaginitis, 567.
- Nitric acid in endometritis, 356.
carcinoma, 513.
- OBTURATOR gland, 76.
internus, 16.
- Oedema of fibroids, 432.
- Ohm, 700.
- Ohm's Law, 700.
- Omental tumours, 691.
and ovarian, 266.
cells found in fluid from, 258.
- Oöphorectomy, 245.
- Oöphoritis, **239**, 296.
- Operating room, arrangements for, 163.
- Opiates in dysmenorrhœa, 627.
- Opium in carcinoma, 511.
- Os externum—contracted, **297**, 375.
examination of, 108.
position, 18, 23, 24.
- Os interium and utero-vesical fold of peritoneum, 19, 23, 317.
- Ovarian artery—course, 73.
corpuscles, 252.
fimbria, 25, 27.
fluid, 252.
fossa, 27.
ligament, 25.
plexus of veins, 74.
- Ovarian tumours—cystic, **246**, **262**, **269**.

- Ovarian Tumours—and fibroids, 444.
 amenorrhœa, 620.
 fibrocystic, 468.
 genesis of, 259.
 malignant, 255, **256**.
 position of, 106.
 causing peritonitis, 183.
 solid, **256**, 263.
- Ovaries—Anatomy, **26**, 147.
 arterial supply, 72.
 cellular structures in, 256.
 changes in at each menstrual period, 95.
 colloid degeneration of stroma, 253.
 degeneration of blood-vessels, 253.
 development, 81, 260.
 displacements, **242**.
 epithelial tubes, 253.
 examination of, 115.
 hyperæmia, 239.
 inflammation, 239.
 and amenorrhœa, 620.
 ligaments, 27.
 malformations, 239.
 malignant development of connective tissue, 255.
 disease of, 257.
 measurements, 26.
 menstrual period, changes at, 95.
 menstruation, effect on, 99.
 palpation of, 238.
 physiology of, 248.
 position, 26, 49, 61.
 in inversion, 410.
 in retroflexion, 385.
 prolapse of, **242**.
 prolapsed and fixed by adhesions, 244.
 retroflexion, 390.
 removal by abdominal section, 270, 465, 691.
 per vaginam, 270.
 structure, 27.
 in superinvolution, 308.
 tumours of, **246**.
 in rudimentary uterus, 286, 292.
 in uterus unicornis, 286.
- Ovariectomy, **270**.
 mortality, 282.
 normal, 245.
- Ovaritis, **239**.
 and Battey's operation, 245.
 and fibroid tumours, 432.
 and Tait's operation, 237.
- Ovula Nabothii, 334.
- Ovulation, 95.
 with rudimentary uterus, 286, 292.
- Ovum—formation of membrana granulosa, 29.
 structure, 29.
 passage into uterus, 95.
- PAIN in fissure of anus, 682.
 endometritis, 351, 352.
 carcinoma uteri, 500, 511.
 carcinoma of body of uterus, 526.
 cellulitis, 184.
 cervical catarrh, 336.
 endometritis, 351.
 fibroids, 436, 439.
 inversion, 413.
 retention of menses in atresia, 553.
 metritis, 362.
 ovaritis, 240.
 acute and chronic peritonitis, 185.
 prolapsed ovary, 243.
 retroflexion, 387.
 sarcoma uteri, 534.
 superinvolution, 308.
 vaginitis, 506.
- Palpation, method of, 103.
 of fibroids, 444.
- Pampiniform plexus, 74.
- Pan-hysterectomy or complete hysterectomy for fibroids, 462.
- Papilloma of the cervix, 470, 476.
- Paracystitis, 651.
- Parametric tissue, 51.
- Parametritis, **193**.
 chronica atrophicans, **202**.
 posterior, of Schultze, 193, 376.
- Paravesical pouch of peritoneum, 60, 63.
- Parenchymatous inflammation, 362.
 ováritis, 239.
- Parovarial cysts, 259, 263.
- Parovarium, 26, 231.
 development, 260.
 distension, 232.
- Parovarian fluids, nature of, 259.
 tumours, 231, 232, **259**.
 and ovarian, 267.
- Parturition, effect on pelvic floor, 68.
 and cervical catarrh, 335.
 repeated and carcinoma, 496, 526.
 endometritis, 350, 353.
- Pathological classification of diseases of women, 715.
- Pawlik's method of examining bladder, 644.
- Péan's method of laparotomy for fibroids, 458.
 needle for pedicle, 458.
- Pedicle—changes in, after ligature, 275.
 of fibroids, 429.
 of polypi, 472.
 treatment in fibroid tumours, 456, 459, 460.
 extra- and intra-peritoneal methods compared, 461.
 twisting, 427.
 of ovarian tumour, 250.
 examination, 266.
 torsion, 281.
 treatment in ovariectomy, 274.
- Pediculated cystic follicle, 334, 470, 475.
- Pedication and extrusion of fibroids, 440.
- Pedunculation of fibroid tumours, 429, 447.
- Pelvic abscess, 194, 209, 693.
 bursting of, 196.
 and vaginitis, 567.
- cellulitis, **193**.
 and ovarian tumours, 263.
 connective tissue, **46**, 47, 50, 51.
 deposit in pouch of Douglas, and retroflexion, 390.
 examination, importance of in carcinoma, 504.
- floor—anatomy, general, **10**.
 structural, **64**.
 displaceable and fixed portions, 67, 603.
 displacements, 67, **603**.
 divisions, 68.
 fascia, 11.
 functions of, 64, 68.
 preventing downward displacement, 372.
 affected by change of posture, 87.
 genupectoral posture, 88.
 intra abdominal pressure on, 68, **85**.
 measurements on external surface, 10.
 muscles, 12.
 opening up of, in genupectoral posture, 88.
 in parturition, 66.
 pubic segment, 65.
 relation to parturition, 65, 68.
 sacral segment, 65.
- glands, 76.
 inflammation producing displacement, **200**, 373, 382.
 and subinvolution, 365.
 organs, development, **79**.

- Pelvic ovarian tumours, producing pelvic inflammations, 183.
 peritonitis, 181.
 producing anteversion, 377.
 with anteversion, 200, 382.
 chronic, 185.
 and ovarian tumours, 203.
 producing retroversion, 200.
 producing small uterus, 306.
- Pelvic-floor projection, 69.
 callipers for measuring, 70.
 measurements, 70.
 effect of pregnancy on, 71.
 segments, 65.
 contrast between, 65.
 as affected by genupectoral posture, 88.
 in parturition, 66.
- Pelvis—blood-vessels of, 72.
 axial coronal sections of, 54, 67.
 contents of, 60.
 coronal sections of, 51.
 genupectoral posture, section in, 89.
 sectional anatomy, 48.
 horizontal sections, 51.
 sagittal lateral section, 49.
 lymphatics, 76.
 nervous supply, 79.
 physics of, 85.
 sagittal mesial section, 49.
 venous plexuses, 74.
- Perchloride of iron to check hæmorrhage, 510.
 caution as to use, 510.
 of mercury as antiseptic, 162.
- Percussion, method of, 106.
 of fibroids, 444.
 of ovarian tumour and ascites, 265.
- Perforation into peritoneal cavity in carcinoma, 493, 507.
 bladder and rectum by fibroids, 441.
- Perimetritis, 182.
- Perineal body, 14, 42, 49.
 structure, 42.
 measurements, 42.
 pad for operations, 169.
 muscles, 12.
- Perineum—arterial supply, 74.
 development, 579.
 rupture, 592.
 central rupture, 594.
 operative treatment, 596, 612.
 support in parturition, 596.
- Perineorrhaphy, 614.
- Periovaritis, 242.
- Peritoneal toilette, 277, 695.
- Peritoneum—atomy of, 42.
 of bladder, 42, 46, 67, 317.
 in parturition, 67.
 of broad ligaments, 44.
 in carcinoma, 493.
 amputation of cervix, 317.
 prolapsus, 606.
 fistula, 661.
 inversion, 411.
 hæmorrhage into, 206.
 inflammation of, 182.
 effusion into, with abdominal tumours, 257.
 tumour, 264.
 method of opening into, 688.
 in relation to operations, 46, 452.
 in parturition, 46.
 on sides of pelvis, 45.
 local divisions on pelvic floor, 59.
 and rectum, 45.
 in retroflexion, 386.
 of uterus, 20, 43.
- Peritoneum on posterior vaginal wall, 44.
 vesico-uterine pouch of, 43.
- Peritonitis—pelvic, 182.
 fatal, and carcinoma, 507.
 relation to cellulitis, 182, 196.
 malignant, 193.
 cells, found in fluid from, 258.
 and ovarian tumours, 263.
 superinvolution, 308.
 subinvolution, 365.
 retroflexion, 386.
 after uterine injections, 225.
 producing retroversion, 382.
 tubercular, 191.
- Pernitrate of iron to check hæmorrhage, 510.
- Pessary—intra-uterine, 380.
 in retroflexion, 400.
 Meadow's, 400.
 Routh's, 400.
 Schultze's, 400.
 Williams', 400.
 vaginal—action of, 398.
 Albert Smith, 394.
 in chronic metritis, 368.
 effect on position of uterus, 91.
 in fibroid tumour, 448.
 Hewitt's cradle, 381.
 Hodge, 394, 400.
 choice of best form, 396.
 mode of introduction, 396.
 position and action, 398, 401.
 Meadow's, 400.
 medicated, composition of, 568.
 in pregnancy, 400.
 in retroflexion, 394.
 for prolapsed ovaries, 244.
 ring, 610.
 Thomas', 381.
 Zwanck's, 611.
- Pfäuger's ducts, a source of ovarian tumours, 248, 256.
- Phthisis—amenorrhœa in, 620.
 uterus in, 308.
- Physics of abdomen and pelvis, 85.
- Physiological activity, period of, 283.
- Piles, 683.
- Placenta—portions retained, and carcinoma, 503, 527.
 metritis, 365.
 relation to inversion, 409, 413, 414.
- Placental polypi, 470, 475.
- Polypes à apparitions intermittentes, 473.
- Polyp of uterus, 466.
 with endometritis, 350.
 fibroids, 470.
 diagnosis from inversion, 415.
 producing inversion, 409.
 +inversion, diagnosis of, 415, 479.
 mucous, 470, 473.
 and fibrous contrasted, 478.
 placental, 470, 475.
- Porte-caustique, 358.
- Positive and negative poles, 701.
- Posterior fornix, effect of pressure on, 399.
- Posture—effect on abdomen and pelvic floor, 87.
 relation to examination and treatment, 147.
 of patient in cellulitis, and peritonitis, 196.
- Pouch of Douglas, 44.
 intestines in, 63.
- Pozzi's operation for sterility, 633.
- Pregnancy—and fibroid tumours, 427, 433.
 and deciduoma malignum, 542.
 advanced and fibroid tumour, 444.
 early, diagnosis, 367.
 from fibroid tumour, 443.
 characters of uterus, 367.
 in detached horn, 296.

- Pregnancy—diagnosis from chronic metritis, 367.
 ovarian tumour, 266.
 and uterus dilated from atresia, 552.
 fissuring of cervix in, 322.
 and large fibroid tumours, 444.
 +ovarian tumour, 268, 280.
 hypertrophy of cervical glands in, 336.
 sympathetic phenomena of, in
 uterine polypus, 447.
 and vaginal fixation, 403.
 and ventral fixation, 405.
- Presentation of head and lacerated cervix, 319.
- Prolapse of ovary, 242.
 rectum, 681.
- Prolapsus uteri, 68, 415, 416, **605**.
 due to fibroid, 433.
 and hypertrophied cervix, 372.
 +inversion, 417.
 mechanism, 608.
 retroversion in, 382.
 senile, 650.
- Pruritus vulvæ, 583.
 in carcinoma, 501.
- Pseudocyesis and ovarian tumour, 267.
- Pseudo-myxoma peritonei, 251, 281.
- Puberty—changes at, 94.
 symptoms of atresia at, 562.
- Pubic segment—of pelvic floor, 65, 67.
 displacements, 65.
 in bimanual examination, 113.
 in semiprone posture, 123.
- Pudendal hernia, 590.
- Puerperal inflammation, 350.
 inversion, frequency of, 412.
 involution and metritis, 365.
- Puerperium and deciduoma malignum, 542.
 endometritis, 350.
 inversion, 408, **412**.
 metritis, 365.
 peritonitis, 183.
 retroflexion, 386.
 retroversion, 381.
 subinvolution, 365.
- Pulse—in hæmatocele, 212.
 peritonitis, 185, 190.
- Pyosalpinx, 226, **233**.
 and ovarian tumours, 263.
 and laparotomy, 693.
- RECAMIER-ROUX Curette, 146.
- Rectal examination, 114, **680**.
 in atresia, 553.
 carcinoma, 503.
 fibroids, 441.
 hypertrophied cervix, 312.
 inversion, 415.
 pathological antelexion, 379.
 retroflexion, 390.
 superinvolution, 308.
 uterine malformations, 293.
 Simon's method, 116.
- Recto-abdominal examination, 114.
- Rectocele, 681.
 and prolapsus, 610.
- Recto-vaginal fistula, 684.
- Recto-vagino-abdominal examination, 116.
- Rectum—**anatomy, 40**.
 diseases of, **680**.
 examination, 114, 680.
 fibroid tumour's pressure on, 439.
 inflammation round, 202.
 lymphatics of, 78.
 microscopic structure, 40.
 physiology, **678**.
 position, 40.
 in retroflexion, 386, 387, 390.
 structure, 40.
 ulceration in carcinoma, 493.
- Rectum—displacing uterus, 372.
 connection with the vagina, 40, 65.
- Reflex action and chloroform, 153.
 disturbances from parametritis atrophicans, 203.
- Reinversion of inverted uterus, 417.
- Renal calculi, 654.
 tumours and ovarian, 267.
 laparotomy for, 692.
- Reposition of inverted uterus, 417.
- Respiratory centre in chloroform, 156.
- Retention of uterine, 656.
 producing cystitis, 651.
 due to fibroid tumour, 439.
- Retroflexion, 340, 371, 373, **383**.
 and cervical catarrh, 335, 340.
 congenital, 385, 386.
 and small fibroid tumour, 443.
 relation to retroversion, 382, 383.
- Retro-pubic fat deposit, 65.
- Retro-uterine hæmatocele, 212.
 and ovarian tumour, 264.
- Retroversion, 373, **381, 383**.
 congenital, 381.
 passage of sound in, 130.
 caused by peritonitis, 200, 382.
 action of pessary on, 398, 401.
 physiological and pathological, 372, 381.
 in prolapsus, 608.
 with antelexion, 383.
 +Retroflexion, 373, **383**.
- Retroverted gravid uterus and ovarian tumour, 264.
- Rheumatic diathesis and dysmenorrhœa, 628.
- Ring pessary, 610.
 with diaphragm, 610.
 mode of introduction, 610.
 position in vagina, 610.
- Rosemüller, organ of, 26.
- Round ligament—**anatomy of, 22**.
 hydrocele, **215**.
 and junction of uterine horn and tube, 287.
 new growths in, 216.
 shortening for displacements of uterus, 405, 406.
 prolapsus, 613, 616.
 tumours of, **216**.
 a guide to parts of uterus, 287.
- Rupture of the uterus in carcinoma, 507.
- SACRAL SEGMENT of pelvic floor, 65.
 rupture of inferior angle, 594.
 the supporting one, 68, 596, 605.
 supporting pessary, 398.
- Sagittal mesial sections, 49, 65.
 lateral section, 49.
- Salpingitis, **226**.
- Sarcoma of ovary, 257.
 connective tissue, 217.
 decidua cellulare, 536.
 uterus, 483, **529**.
 and carcinoma, 503, 505.
 endometritis, 353.
 inversion, 413.
 vagina, 574.
 vulva, 588.
- Sarcomatous degeneration in fibroids, 433.
 polypus, 474.
- Scarification of cervix, 340.
- Schroeder's operation for cervical catarrh, 324, 341.
- Schultze's dilator, 302.
- Scirrhus cancer, 484.
- Scissors, **148**.
 Bozeman's, 148.

- Scissors, for division of cervix, 303.
in amputation of cervix, 312.
Kuchenmeister's, 303.
- Serofula, leucorrhœa in, 350.
and congenital atrophy of uterus, 291.
- Sectional anatomy, 48, 64, 67.
- Semiprone posture, 122.
- Septic matter causing cellulitis, 194.
metritis, 361.
- Septicæmia—after operation for atresia, 555.
in carcinoma, 506, 508.
from suppuration of fibroid tumours, 441.
in inversion of uterus, 417.
relation of lymphatics to, 78.
and metritis, 361.
after ovariectomy, 278.
after tents, 143.
- Septic peritonitis, 187, 188.
salpingitis, 226.
- Serre-nœud of Cintrat, 458.
Kœberlé's, 456.
in laparotomy for fibroids, 458.
- Serotal tumour, 545.
- Serous peritonitis, 182, 185.
- Serum, collection in Fallopian tubes, 226.
- Sexual activity and fibroids, 433.
excess and endometritis, 350.
- Sickness after chloroform, 157.
- Silk ligatures, preparation of, 167.
- Silk-worm gut, 149.
- Silver-wire sutures, 149.
- Simon and Marckwald's amputation of cervix, 313.
- Simon's sharp spoon and carcinoma, 513, 527.
urethral specula, 641.
- Simpson's (A. R.) nail-curette, 453.
repair of perineum, 598.
sound, 129.
(Sir J. Y.) division of cervix, 302, 380.
galvanic intra-uterine stem, 309.
metrotome, 302.
sound, 128.
- Sims' (Marion) division of cervix, 380.
amputation of cervix, 312.
curette, 145.
dilator, 302.
speculum, 121, 126, 323, 337.
tenaculum, 120.
operation for vaginismus, 570.
- Skene-Goodman catheter, 653.
- Skene's urethral specula, 643.
- Smith (Albert) pessary, 396.
mode of action, 397, 398.
- Souffle—uterine, in fibroid tumour, 444.
absence of, in ovarian tumour, 444.
- Sound—uterine, 128.
combined with Bimanual, 135.
in antelexion, 380.
hypertrophied cervix, 312.
endometritis, 337, 350, 353, 355.
metritis, 361, 367.
small fibroid tumours, 442.
fistula, 663.
pediculated submucous fibroid, 478.
subserous fibroid, 444.
inversion, 415.
polypus and inversion, 479.
polypoidal fibroid, 442.
chronic metritis, 367.
retroflexion, 390, 393.
superinvolution, 308.
method of dressing, 355.
in replacing retroflexion, 393.
in treating antelexion, 380.
- Specula, vaginal—comparative advantages of various forms, 126.
Barnes' crescent, 125.
Battey's, 122.
Bozeman's, 121.
- Specula, vaginal—Cusco's, 125.
Fergusson's, 125.
Neugebauer's, 125.
Sims', 121.
in cancer, 503.
in endometritis, 353.
in lacerated cervix, 323.
- Speculum—anal, 680.
urethral and vesical, 640, 641.
- Spencer Well's clamp, 274.
trocar, 273.
- Sphincter ani—externus, 40.
internus, 41.
tertius, 40.
operation to restore, 597.
- Splenic tumours, 691, 692.
- Sponge-tents, 138.
- Sponges—in operations, 166.
precautions as to, 272, 689.
- Spray in ovariectomy, 279.
- Staffordshire knot, 236, 275.
- Staphylococci, 177, 199, 348, 562.
- Stem—galvanic intra-uterine, 309.
in antelexion, 380.
- Sterility, 629.
in antelexion, 377, 378.
cellulitis, 199.
cervical catarrh, 336.
hypertrophied cervix, 311.
stenosis and rigidity of cervix, 300.
lacerated cervix, 322.
dyspareunia, 569.
endometritis, 351, 352.
stricture of Fallopian tubes, 224.
fibroid tumour, 440.
fistula, 662.
chronic metritis, 366.
peritonitis, 185.
double ovaritis, 241.
uterine polypi, 478.
retroflexion, 388.
sarcoma, 533.
vaginismus, 569.
- Streptococci, 177, 199, 348, 562.
- Structural anatomy of the pelvic floor, 64.
- Strumous diathesis and endometritis, 354.
- Subinvolution, 362.
and anteversion, 357.
lacerated cervix, 321.
cervical catarrh, 340.
with retroflexion, 384.
- Submucous fibroid, 427, 428, 470.
- Subperitoneal fibroid, 427.
- Subserous pediculated fibroids, 455.
- Subserous fibroids and fibro-cystic tumours, 466.
- Superfoetation, 293.
- Superinvolution, 305.
and amenorrhœa, 620.
ascribed to ovaritis, 240.
- Supravaginal portion of cervix, hypertrophy of, 311, 314.
amputation of the cervix, 515.
- Suppuration after cellulitis, 196.
hæmatocele, 214.
of fibroids, 432.
- Sutures, 149, 326.
removal of, 313, 326.
- Sympathetic pain in ovaritis, 240.
- Syncytial carcinoma, 545.
- Syncytium and deciduoma malignum, 543.
- Syphilitic salpingitis, 227.
ulceration and carcinoma, 503.
ulceration of perineum, 594.
- Syringe, Higginson's, 150.
- TAIT'S graduated dilators, 143.
knot, 276.
operation for removal of uterine appendages, 236, 245, 692.
in fibroids, 465.
dysmenorrhœa, 628.

- Tait's operation for repair of perineum, 599.
ruptured perineum, 615.
- Tangle and tupelo tents, **139**, 142.
- Tapping in cellulitis, 197.
hydrocele, 216.
ovarian tumours, 268.
peritonitis, 193.
- Temperature, reduction of, 190.
after ovariectomy, 278.
- Tenaculum—Sims', 120.
in diagnosing lacerated cervix, 323.
- Tents, **138**.
dangers of, 143, 301, 350.
in stenosis of cervix, 301.
- Tetanus in carcinoma, 508.
- Thomas' anteversion pessary, 381.
- Tightlacing and uterine displacement, 372.
- Torsion of pedicle of ovarian tumour, 281.
for removal of polypi, 481.
- Trachelorrhaphy, 324.
- Transverse section, 51.
- Transversus perinei, 13.
- Treatment—operative, in gynecology, 714.
sound in, 128.
specula in, 126.
 volsella in, 119.
- Triangular ligament, 11.
- Trigone of bladder, 35.
- Trocars for tapping ovarian cysts, 273.
- Tubal disease producing peritonitis, 183.
treatment of, 235.
- Tubercle bacillus, 179, 228, 349, 482.
- Tubo-ovarian cysts, 230.
- Tubercular diathesis in superinvolution, 308.
peritonitis, **191**.
- Tuberculosis of uterus, **481**.
vagina, **575**.
- Type of menstruation, 94.
- ULCERATION of Cervix—so-called, 323, 331.
true, 334.
Os uteri, 488.
- Ulcus serpiginosum vulvae, 585.
- Upright posture, 8, 87, 88.
- Urachus, 260.
- Uræmia in carcinoma, 506.
fibroid tumour, 439.
- Ureter—anatomy of, **37**.
catheterisation, 643.
relation to cervix and urethra, 660.
dilatation in carcinoma, 493.
due to cicatrization, 386.
compression by fibroid tumours, 439.
ureter implantation into bladder, 674.
opening of, 35.
position of, 54.
- Uretero-uterine fistula, 658, 661, 662.
vaginal fistula, 658, 661, 662.
- Urethra—anatomy of, 33.
atresia in fistula, 660, 673.
caruncle, 569, 648.
development, 579.
dilatation, 639, 649.
direction, 33.
displacements, **647**.
exploration, 638.
glands, 34.
malformations, **647**.
neoplasms, **648**.
pressure on, by fibroid tumour, 439.
relation to vaginal wall, 65.
stricture, **649**.
- Urethral orifice, 7.
- Urethritis, 649.
- Urethrocele, 647.
- Urethro-vaginal fistula, 658, 661.
+atresia of urethra, 673.
- Urination, physiology of, 637.
- Urine—normal composition, 638.
character in cystitis, 651.
examination in fibroid, 440.
- Urine—incontinence and retention, 656.
in fistula, 662.
- Uro-genital sinus, 571.
persistence of, 580.
- Utero-sacral ligaments—anatomy of, 22.
cicatricial contraction in, 193, 376.
function, compared with pessary, 399.
stretching of, in retroflexion, 386.
- Uterine appendages, removal of, 236, 692.
artery—course, 72.
in removal of fibroids, 451.
catarrh, 344.
cavity—variations in length of, 134.
contractions—importance in diagnosis, 266, 444.
mucous membrane in long-standing endometritis, 347, 348.
changes in, at menstrual period 96.
plexus of veins, 74.
- Uterus—abscess in wall, 361.
absent, or rudimentary, 286, 292, 293.
amputation of inverted, 421.
adenoma malignum of, 528.
anatomy, **17**.
anteversion, 373, **374**.
anteversion, 373, **380**.
atrophy, **305**.
arterial supply, **73**.
atrophy and amenorrhœa, 620.
bicornis, 289, 292, 295.
bipartitus, 286, 292.
body of, in inversion, 410.
carcinoma, **483**, **499**, **509**, 523.
changes after birth, 291.
congenital atrophy, 291, 292, 296.
and superinvolution, 308.
corroding ulcer of, 488.
curetting, 145, 355.
deciduoma malignum of, 536.
degenerative changes, 432.
development, 82, 283.
didelphys, 288, 292, 295.
digital pressure in fornices, 91.
dilatation in atresia, 549.
for therapeutic purposes, 359.
and perforation in carcinoma, 507.
displacements, **371**.
in large fibrous polypi, 471.
ovarian and uterine tumours, 468.
peritonitis and cellulitis, 200.
operative treatment of, 401.
divisions, 21.
duplex, 292.
enlargement of, in endometritis, 363.
metritis, 367.
extirpation of, through abdominal walls, 516.
through the vagina, 516.
in malformation, 296.
fibroid tumour, **422**, **435**, **446**, 691.
changes with fibroid tumour, 429.
fibro-cystic tumours, **466**, 691.
changes in, at angle of flexion, 374, 384.
glands, 23.
hypertrophy, 310.
in fibroid tumour, 429.
in prolapsus, 607.
impairment of function, in maldevelopment, 292.
infantile, 291, 292, 296.

Uterus—*inflammation* (endometritis, metritis), 330, **343**, **360**.
 and amenorrhoea, 620.
inversion, **408**.
 with fibroid, 430.
 spontaneous reinversion, 417.
lateri-version, 201.
ligaments, 21.
lower segment, 20.
lymphatics, 76.
malformations, **285**.
 and *superinvolution*, 308.
measurements, 19.
mobility, 61, 117.
changes at a menstrual period, 96.
periods at which morbid conditions arise in, 288.
mucous membrane, 23.
musculature, 22.
nervous supply, 79.
one-horned, inversion of, 409.
operations for retaining in position, **401**.
peritoneum of, 21.
polypi, **470**.
position, means of ascertaining, 59.
 changes with fibroid, 430.
 with distended bladder, 59, 61, 372.
 in genupectoral posture, 89.
 normal, **38**, 317.
 opinions as to, 55.
 physiological changes in, **61**, 372.
 relation to viscera, 63.
pressure on bladder, 380.
prolapse, 67, 415, 416, **605**.
replacement of, 380, 381, 383, 392.
retention in position, 392, 394, 401.
retroflexion, 340, 372, 373, **383**.
retroversion, 373, **380**, 383.
sarcoma, **529**.
septus, 290, 292, 295.
 atresia of, 558.
shape, 17, 57.
form in section, 16.
stitching to abdominal wall for displacements, 403, 404.
structure, 20.
subinvolution, 340, 362, 363, 365.
superinvolution, **305**.
support of, 68.
tuberculosis of, **481**.
tumours of, **422**.
 and *metritis*, 366.
unicornis, 286, 292, 295.
 and *fibroid tumours*, 296.
vaginal fixation, 402.
ventral fixation of, 403.
veins of, 74.
virgin type, 291.

VAGINA—Anatomy, 30.

arterial supply, 72.
atresia, **512**.
cancer of, *primary*, **571**.
 secondary, 494.
cicatrization after fistula, 660.
 labour, 552.
 menopause, 565.
artificial closure for fistula, 676.
congenital bands in, 560.
cysts, **571**.
development, 83, 579.
dilatation in atresia, **550**.
direction in upright posture, 30.
fibroid tumours, **571**.
form in section, 32.
in genupectoral posture, 88.
inflammation of, 561.
inversion + inversion of uterus, 412.

Vagina—lymphatics, 76.
 menopause, change after, 565.
 plugging to check hæmorrhage, 510.
 position, 30.
 sarcoma, **574**.
 structure of, **32**.
 tuberculosis, **575**.
 venous plexuses, 74.
 walls, 30.
Vaginal cleft, 65.
 bacillus, 562.
 columns, 31.
 enterocele, **617**.
 examination, **107**, 444.
 and *ovarian tumour*, 265.
 retroflexion, 389.
 glands, 32.
 glass plugs, 557.
 hysterectomy, 454, 517, 697.
 injection, 339.
 method of removal of ovaries, 270, 696.
 of ovariectomy, 270.
 tampons and carcinoma, 510.
 vaginitis, 568.
 orifice—atomy, 7.
 dilatation of, 568.
 division of sphincter, 570.
 fissures around, producing vaginitis, 568.
 portion of cervix, 20.
 amputation of, 515.
 hypertrophy of, 311, 314.
 secretions—physiological and pathological, 562.
 walls—attachments of, 65.
 inversion in prolapsus, 607.
 length, 31.
 and *direction in genupectoral posture*, 89.

Vaginismus, 568.

Vaginitis, 561.

diphtheritic, 565.
gonorrhoeal, 565.
senile, 565.

Varicose veins, due to fibroid tumours, 439.
 hæmorrhage from, in vulva, 590.

Varix of vulva, 590.

Veins, pelvic—compressed by fibroid tumours, 439.

Venereal excess, causing peritonitis, 183.

Venous sinuses round fibroid tumours, 425, 438.

Venous thrombosis in carcinoma, 506.

Ventral fixation of uterus, 403.

Versions of uterus, 373.

Vesical catarrh, 674.

Vesical plexus, 74.

Vesico-uterine fistula, **658**, 661, 662, 666.

Vesico-vaginal fistula, **658**.
 artificial, for cystitis, 653

Vesico-vaginal septum, 659.

Vesico-vagino-abdominal examination, 113.

Vestibule—atomy, 7.

development of, 579.

Vichy waters in chronic metritis, 368.

Volvella—description of, **117**.

examination with, 118.

with speculum, 123.

in superinvolution, 308.

for introducing tents, 141.

in replacing uterus, 394.

Volt, 700.

Voltaic current, 698.

Vomiting in hæmatocele, 212.

peritonitis and cellulitis, 196.

Vulva—atomy of, 7.

atresia of, 580.

carcinoma, 586.

cysts, 586.

direction of, 8.

elephantiasis, 586.

eruptions, 585.

Vulva—fibromata, 588.
 hæmatoma, 590.
 hæmorrhage from varicose veins, 590.
 hernia, 590.
 inflammation, **582**.
 kraurosis, 589.
 lipomata, 588.
 lupus, 588.
 malformations, **578**.
 neuromata, 588.
 pruritus, 588.
 sarcoma, 588.
 ulcus serpiginosum of, 585.
 varix, 590.

Vulvitis, **582**.
 of children, 582.

WEIR MITCHELL'S treatment of nerve prostration, and chronic uterine disease, 370, **704**.
 White's repository for inversion, 418, 419.
 Wiesbaden waters in chronic metritis, 368.
 Wire sutures in laceration of cervix, 283, 326, fistula, 663.
 Wolffian bodies, 79, 80, 83, 231, 255, 259, 260.
 bulbs, 83.

ZONA pellucida, 29.
 Zwanck's pessary for prolapsus, 611.





