On the immediate treatment of stricture of the urethra: by the employment of the stricture dilator / [Barnard Wight Holt].

Contributors

Holt, Barnard 1816-1890.

Publication/Creation

London: John Churchill, 1861.

Persistent URL

https://wellcomecollection.org/works/bcmjhbgh

License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org







S. B. Buckentt & M. W. Work With authors Complements

ON THE


IMMEDIATE TREATMENT

OF

STRICTURE OF THE URETHRA.







evaux, del et lith.

M & N .Hamhart, Impt

REPRESENTS THE PENIS LAID OPEN SHEWING THE DIVISION OF THE TWO STRICTURES ON THE FLOOR OF THE URETHRA.

IMMEDIATE TREATMENT

OF

STRICTURE OF THE URETHRA,

BY THE EMPLOYMENT OF THE

"STRICTURE DILATOR."

BY

BARNARD HOLT, F.R.C.S.,

SENIOR SURGEON TO THE WESTMINSTER HOSPITAL; LECTURER ON SURGERY IN THE WESTMINSTER HOSPITAL SCHOOL OF MEDICINE, ETC.

LONDON:

JOHN CHURCHILL, NEW BURLINGTON STREET.

MDCCCLXI.

X 1.02 - 1.05 -

11300931

M18945

WELLCOME INSTITUTE	
Coll.	welMOmec
Call	
No.	WX 600
	1881
	H 75%
-	

PREFACE.

The greater portion of the following remarks has already appeared in the pages of the 'Medical Times and Gazette,' but in preparing them for re-publication I have appended some observations on the relative merits of the treatment I propose, as contrasted with the various operations by incision; and have also added some cases to illustrate more fully some of the points on which I have insisted.

I have lately been able to examine the effects of forcible dilatation upon a stricture after the patient's death from another cause,—a point of much interest; and the description, with a drawing of the preparation, which was exhibited at the Pathological Society of London, will be found at the end of the book.

The fatality that attends the uninterrupted course of a serious stricture, is but too certain. All practical Surgeons are acquainted with the complications and suffering resulting from a disease, which, if taken in time, is always amenable to treatment. If, therefore, any means can be employed, by which all serious results can be obviated, and the treatment at the same time be made so simple as to be available by the majority of Surgeons, it will, perhaps, be admitted that a step has been gained in the right direction. Hitherto I have not ventured to publish my experience of the plan I adopt, simply for the reason that it is injudicious to enunciate any new method of treatment which has not been subjected to numerous and repeated trials; but having now operated upon more than 100 cases with unvaried success, not only in private but also in Hospital practice, where every opportunity has been afforded to the profession of witnessing the treatment and its results, I now

feel justified in submitting that experience to the judgment of my professional brethren, and shall endeavour to frame such clear rules for their guidance, that, in any intelligent hands the operation may be as successful as it has been in my own. Up to the present time, this method of treatment, although adopted by some Surgeons, has been mainly confined to myself, simply, I believe, from the fact that its general utility never having been published, it has not, as yet, been appreciated. Let me indulge a hope that, should I be fortunate enough to express myself so clearly as to be generally understood, the performance of the operation may become extended.

In conclusion, I would only deprecate the prejudices which exist in the minds of Surgeons, against any novelty in the treatment of a common disorder, and request an unbiased perusal of the following pages.

14 SAVILE Row, December, 1861.

STRICTURE OF THE URETHRA,

&c. &c.

THE benefits which the illustrious brothers William and John Hunter have conferred upon mankind can hardly be exaggerated. By introducing into pathology the finely inductive reasoning which had previously characterised their physiological speculations, they elicited a system of enlightened principles which has thoroughly humanized every branch of the healing art. Nature, these great teachers clearly showed, was the only safe instructress in pathology and in therapeutics, careful and patient study of her processes in health and in disease formed the only reliable basis for successful practice. Out of the school of these eminent medical philosophers issued many distinguished disciples, who, by their writings, lectures, and example, have

spread a knowledge of the Hunterian system throughout these kingdoms, and even awakened a surmise of the nature of these important truths in France and Germany. The preeminent abilities of Baillie, Abernethy, and Denman gradually moulded the practice of medicine, of surgery, and of midwifery in subjection to the Hunterian principles. A careful interpretation of Nature was the rigidly enforced rule. All rash, meddlesome, and coarse methods of treatment were denounced, the vis medicatrix was perpetually appealed to, arte non vi was constantly ejaculated. The late excellent surgeon, Mr Lynn, an eminent pupil of Hunter, used, in his eccentric way, often to say, "Nater, gentlemen, Nater cures the disease."

Without doubt, this habitual subjection of the mind to the careful contemplation of natural processes was at the bottom of the marked improvement which, of late years, has taken place in every department of our art. It is a weakness of the human mind, however, to oscillate from one extreme to the other. In medicine, from the extreme of heroic doses, and a licentious use of the lancet, the practical pendulum vibrated to homeopathic globules and a supinely-expectant treatment. In mid-

wifery, from excessive instrumentation, the accoucheur has occasionally fallen into a helpless acquiescence in avoidable evils; and in surgery even an affected aversion to the knife has too often seduced the operator into a fatal procrastination.

The "native hue of resolution," which is nowhere more necessary than in the practice of surgery, has not, perhaps, in any class of affections been so injuriously "sicklied o'er," as in certain obstinate forms of stricture of the urethra. Mr. Abernethy's denunciations of all violence in catheterism as being inconsistent with the main purpose of the remedyviz., the absorption of the material of the stricture—has had a greater effect than he probably contemplated. A timid and dilatory treatment has been the result, and in too many instances the supineness of the surgeon has permitted the case to drift to a fatal termination. It is a sad truth that "Nater" will not cure a chronic stricture.

My attention was early attracted to the prevailing defects in the treatment of strictures. I could not but observe the tediousness of the treatment by ordinary dilatation, occupying many months before an average instrument could be introduced into the bladder,



and that even when the dilatation was accomplished, the constriction generally returned, so that perpetual surgical care was required.

Being deeply impressed with the unsatisfactory nature of the prevailing methods of curing these distressing maladies, about seven years ago I adopted a more energetic mode of treatment, and invited the notice of the profession to a new "Stricture Dilator." Its use was at first limited to simple dilatation, which was readily effected by graduated tubes passed between the blades without the withdrawal of the original instrument. Experience, however, soon showed me that, as a general rule, when dilatation was carried much beyond the degree produced by ordinary bougies, "stricture fever" was induced. I therefore determined, though with some apprehension as to the

consequences, to split the stricture by passing the largest-sized tube at once, and thus immediately to enlarge the contracted part of the canal, so that it might receive a catheter equal to the normal size of the urethra.

Fearing the effects of the urine being permitted to come in contact with the laceration thus occasioned, I kept a gum elastic catheter in the bladder, but as this measure gave rise to considerable irritation, I determined to content myself with simply splitting the stricture, drawing off the urine, and not again using the catheter till two days after the operation. After that interval, an instrument of the same diameter as that used at the time of the operation was again employed, and its use was continued-first, on alternate days, and, subsequently, at longer intervals. Experience has shown, indeed, that instances occur in which it is necessary to use a catheter one size smaller than that first passed after the operation, but such cases are exceptions.

The instrument by which this simple process is accomplished consists, as is shown in the drawing, of two grooved blades fixed in a divided handle and containing between them a wire welded to their points, and on this wire a tube (which, when introduced between the blades

corresponds to the natural calibre of the urethra) is quickly passed, and thus ruptures or splits the obstruction. The simplicity of this apparatus is obvious to all, and the history of above a hundred cases proves that its use is unattended by any of those serious complications-viz., hæmorrhages, false passages, infiltration of urine, perinæal abscess, fistulæ, swelled testis, &c., &c .- which too often accompany the other operative processes devised for the relief of this malady. The forcible distension caused by the dilator affects the morbid obstruction only; the healthy portion of the canal is not disturbed, the slightly vascular character of the strictural deposition at the same time gives rise to but very inconsiderable bleeding, and perfectly obviates infiltration into the surrounding tissues.

The method of performing this operation may be described in a very few words. The permeability of the canal having been once satisfactorily ascertained, the size of the meatus of the urethra is to be gauged by passing into it a sound that will conveniently fit, and the number of the sound so used is to be the number of the tube to be passed: it is important to ascertain this, so that the urethra may not be stretched beyond its natural limits, for while

the urethra of one person will admit of No. 14, another will not admit more than No. 9.

The dilator having been previously well oiled, is to be introduced with the handle somewhat over the patient's left hip, and by keeping the convex portion gently pressing against the under part of the urethra, the point will glide along the upper portion until it is fairly beyond the triangular ligament, when, by bringing the handle to a right angle with the body, and gradually depressing it-but not so much as in the passage of an ordinary catheter-it will usually slip into the bladder; in fact, the same proceeding is to be adopted as in introducing a lithotrite for the purpose of crushing a calculus. Having reached the bladder, the dilator should be gently rotated, to prove that it is fairly within that viscus, and being thus assured, the Surgeon is next to place the point of the tube he had previously selected, upon the wire between the blades, and thrust it quickly onwards to the end. The stricture being now fairly split, the dilator should be rotated to still further separate the sides of the rent, and then be withdrawn; a catheter corresponding to the number of the tube being substituted, for the purpose of removing the urine. The catheter is then to be taken out,

and the patient sent to bed, with directions to take, every four hours, for the first day and night, a mixture containing in each dose two grains of quinine and ten minims of the tincture of opium. The facility with which this proceeding can be effected will of course depend upon the kind and number of the strictures, and the existence or otherwise of false passages, or fistulæ in perinæo. The urine having been withdrawn, the patient does not require to pass water for some hours, and when compelled to do so, the stream is usually larger, and the urine passes with greater facility than before. On the second day from the operation, the same catheter should be gently introduced; but, if the patient complains of much scalding, it will be better to take one size less. This should be repeated every other day for a week, when the larger one may be substituted, and the patient be taught to pass his own instrument. Of course the time occupied in the after-treatment must vary with the nature of the case, and the more obstinate forms necessitate the employment of the catheter for some time, the intervals being gradually increased until it is not required to be used more than once in three, four, or six months, and in most instances, not more frequently than once a year. The bowels should be relieved by a dose of castor-oil taken early on the morning of the operation, and the patient should be directed not to pass water for two or three hours previously, in order—first, to facilitate the introduction of the dilator; and, secondly, to permit its free movement in the bladder.

A perusal of the following cases, which are extracted as salient examples from a long series, and which were, for the most part, witnessed through their whole career by the students of the Westminster Hospital, and by those Surgeons who favour that institution by their attendance, will, I think, corroborate the points which I wish to establish, and, I hope, justify me in upholding as proved the following conclusions:—

- 1. That the operation is of the most simple kind, and that anyone who can pass a bougie through a difficult stricture is competent to perform it.
- 2. That it is not attended with hæmorrhage, infiltration of urine, abscess, or any serious local mischief.
- 3. That in the majority of instances the relief is immediate.
 - 4. That the occurrence of rigors, or any

other constitutional disturbance, is very rare, and the patient is seldom confined to bed longer than from twelve to twenty-four hours.

- 5. That the urethra is immediately made permeable by a catheter of full size, which may be ever afterwards passed at discretion.
- 6. That this method is available in every kind of stricture where a canula of any size can reach the bladder.
- 7. That when the after-treatment is judicious and attentive, the full capacity of the passage is always maintained.
- 8. That in all cases of neglected aftertreatment, the stricture yields again to this method more promptly than to any other.
- 9. That, it being impossible that any but the diseased tissue can be divided, the splitting of the stricture has a decided superiority over any cutting operation.
- 10. And, to sum up the great advantages in one proposition, that the process is facile, speedy, prompt in its effects, and free from every danger, immediate or remote.

The course of general treatment will naturally vary, according to the kind of obstruction, the number of strictures, and the occasional complications of contracted bladder, enlarged prostate, fistulæ in perinæo, false

passage, &c. In simple stricture, however narrow, the relief will be immediate, but in the more complex forms of these maladies, the size of the stream is not increased so directly as might have been anticipated from the immediate enlargement of the canal. Notwithstanding, however, that the size of the stream may for a short time remain somewhat restricted, the patient is able to empty his bladder much more quickly and effectually than before, and has less frequent micturition. The limitation of the jet evidently depends upon the thickening of the surrounding textures, and swelling of the mucous lining; these morbid states speedily subside, and in a short space of time the patient can void his water in a normal manner.

CASE I.

Stricture of Twenty-five Years' Duration—
Severe Constitutional Disturbance—Recurrence of "Stricture Fever" after each attempt to introduce a Catheter—Retention of
Urine—Hæmorrhage—False Passage—
Operation—Recovery.

Thomas W., aged 50, a labourer, of dissolute habits, was admitted into the Westminster

Hospital under my care, on November 5, 1857. He had been the subject of stricture for twenty-five years, during which period the stream has been gradually contracting, and the urine is now passed guttatim. Five years since he was an in-patient at St. Thomas's Hospital, where, after considerable difficulty, a No. 1 catheter was introduced and retained, being replaced by others, in succession, until No. 6 could be passed, when he left. The contraction speedily recurred, and he was admitted into the Westminster Hospital as above. He is now greatly emaciated, with brown and dry tongue; pulse 110; countenance pallid; appetite defective; abdomen tympanitic; nights restless; and there is a constant escape of fæces during the straining to evacuate the bladder, an effort which he is compelled to repeat every hour and a half. The house-surgeon having twice failed in passing a No. 1 catheter, each attempt being followed by syncope and subsequent rigors, I now saw the patient, and directed the nightly use of the warm bath, and prescribed salines with opium through the day, and castor-oil every morning. At the expiration of a week his general condition was sufficiently improved to justify the attempt to introduce a No. 1

catheter, when a stricture was detected five inches from the meatus, through which it was impossible to penetrate. Although the examination was conducted with the greatest gentleness, it was followed by a severe attack of "stricture fever," which was only relieved by the administration of opium. The urethra being irritable, and the patient much exhausted, another week was permitted to elapse, during which time quinine and stimulants were administered, but the second trial was likewise futile, and followed by precisely the same results as the first. After the lapse of a third week, and while the patient was fully under the influence of opium, a third attempt was made, but with no better success. I therefore determined to place the patient thoroughly under the influence of chloroform, when, after very considerable difficulty, a No. 1 catheter was introduced, and a pint of highly-offensive and turbid urine withdrawn. In the evening he had retention, which, after great difficulty, was relieved by a No. 1 gum catheter; there was considerable hæmorrhage, the patient losing from ten to twelve ounces of blood. He passed a restless night, and being unable to make water, the catheter was again had recourse to. (The house-surgeon having unAs it was found to be impracticable to re-introduce it, the man was again anæsthesiated, and another attempt made, but although the catheter passed to its whole length no urine followed, and it was evident a false passage had been made. Opium and the warm bath were had recourse to, and in a few hours a small quantity of bloody urine escaped and continued to dribble away during the day and night.

When I saw the patient on the following morning, his tongue was dry and brown; pulse feeble and rapid; skin hot, dry, and blanched; and he complained of great tenderness upon pressure over the lower part of the abdomen; he was immediately ordered calomel and opium every four hours, and a turpentine fomentation over the seat of pain. For the next few days he continued in a very precarious state, but the symptoms gradually yielded, and in ten days from the retention the urine was free from blood; his health was, however, so much shattered by the accompaniments of the retention, that a month elapsed before any further instrumental attempt was made. Former experience having shown the impossibility of introducing any

instrument excepting while under chloroform, he was once more anæsthesiated, and the dilator with difficulty passed into the bladder; the large-sized tube was then employed, the stricture fairly split, and the dilator removed. A No. 12 catheter was next inserted, and the urine withdrawn; the bleeding was very trifling. Ordered a mixture of quinine and opium, two grains of the former and ten drops of the latter for a dose. In the evening he felt chilly and uncomfortable, but had no shivering; the urine flowed with moderate ease, and was triflingly tinged with blood.

On the following day he was able to walk about the ward; the urine came away in a small but clear stream; the medicine was omitted, and his ordinary diet resumed.

On the second day after the operation, a No. 12 catheter entered the bladder without the least difficulty; the urethra was tender, but he did not experience so much pain as on former occasions; the after-treatment was continued at increasing intervals, and he shortly left the hospital capable of passing his own instrument.

The foregoing is a case of very considerable interest, as embracing many of the most important points in connection with stricture.

The duration of the disease, the manner in which micturition was accomplished, the deplorable state of the patient's health, the supervention of "stricture fever" after every attempt at catheterism, the after establishment of false passage, and the profuse hæmorrhage, are all evidences of the serious nature of the case, which consequently presented a severe test to the adopted treatment. The history satisfactorily shows the immunity from fatal consequences, rapidity of recovery, and complete restoration of the urethra to its natural size, which are the great characteristics of the operation.

CASE II.

Stricture of Fourteen Years' Standing—Of a Dense Cartilaginous Character—Situated about the Triangular Ligament—Urine, which was Ammoniacal and Purulent, passed Guttatim—Operation—Recovery.

John W., aged 45, a labourer, was admitted in December, 1857, suffering from stricture of the urethra, of fourteen years' duration. Seven years since, the urine having been for some weeks previously passed *guttatim*, he had retention, and applied at St George's Hospital, where after considerable difficulty a

catheter was introduced. He continued an in-patient until No. 4 could be passed, and then left, much improved both in health and manner of micturition. The stricture, however, gradually returned, and after several attacks of retention, and a generally increasing difficulty in relieving himself, he became an in-patient of the Westminster Hospital. At the time of his admission, the urine was passed every hour, with great straining, and frequently in drops; there was considerable hardness in the perinæum, and his health was much damaged, as evinced by emaciation, pallor, loss of appetite, sleepless nights, and general feverishness; the urine was highly ammoniacal, and loaded with mucus and pus. A No. 1 silver catheter was attempted to be passed without success, there was complete obstruction at the triangular ligament, and although the point was firmly grasped, it could not be made to penetrate through the stricture. The patient was kept in bed and the bowels regulated, and on the third day from his admission, another attempt was made in the most gentle manner, but with no better result. Considerable irritative fever followed the two trials, and it was only after the expiration of two months, during which time six endeavours

were made, that a No. 1 catheter was passed. The stricture was of the cartilaginous variety, about an inch in length, and gave that peculiar grating to the catheter so specially characteristic of cartilaginous obstruction. A large quantity of highly-offensive urine was withdrawn, and the catheter was retained. On the following day, a larger size was substituted, and on March 4th, the patient having been placed under the influence of chloroform, the dilator was with considerable difficulty introduced, and the No. 12 tube immediately passed. A No. 12 catheter was then easily slipped in, and the urine withdrawn. On the following day there was no febrile disturbance, and the patient declared he made water better than he had done for many years. On the succeeding day, or second after the operation, the urine was passed in a smaller stream, and with some scalding, consequently a No. 11 catheter was used, the urine being much less fetid than before. The stream continued small for a fortnight, but the urine was expelled in a much shorter time, and the frequency of micturition materially diminished; he now only required to relieve himself three times during the night, and the same during the day. No. 11 having been passed on alternate days, a No. 12 was introduced, and used every third and every fourth day in the manner already described. Six weeks after the operation he left the hospital, making water in a perfectly natural manner, having been taught to pass a No. 12 with ease.

The foregoing, one of a class of strictures that offers the greatest impediment to the introduction of a catheter, is an excellent example of a formidable cartilaginous obstruction situated at a part of the urethra which, corresponding to its curve, gives greater trouble in the introduction of an instrument than any other. The patient had been the subject of a stricture for many years, his urine had for months previous to his admission been passed guttatim, his clothes were constantly saturated, and his health was materially damaged by the hourly necessity for relief-an evil which depended upon the stricture, as proved by the large quantity of urine that was withdrawn when a catheter was introduced. All these difficulties were immediately overcome, and a large-sized catheter could be ever afterwards introduced with perfect facility. The offensive ammoniacal condition of the urine gradually abated, that fluid assumed a normal character, and the bladder being no longer

irritated by its retention, the intervals between the times of micturition were prolonged. In a word, the patient was restored to complete comfort without having been detained in bed more than a day, and without experiencing a single unfavourable symptom; indeed, he did not suffer in the same ratio as during the attempt to simply introduce a catheter. To those who are uninitiated in the treatment of stricture, it might appear that, by keeping a catheter in the bladder, and daily increasing the sizes, the same results would have been obtained, but experience proves this plan of treatment to be utterly futile, and that as soon as the catheter is removed, so soon does the stricture return.

CASE III.

Stricture at the Bulb of Twenty-eight Years' Standing—Frequent Attacks of Retention of Urine—Infiltration—Fistulæ in Perinæo and Scrotum—Enlargement of the Prostate Gland.

H. R., a captain in the Navy, aged 58, consulted me after having for many years been suffering from the effects of stricture. He had had repeated attacks of retention of

urine, and upon three occasions matter formed behind and in the scrotum, which eventually gave rise to fistulæ in those situations. At the time of his coming under my care, the greater portion of the urine escaped through these apertures, so that he was always compelled to make water in a sitting posture, or with the aid of an earthenware slipper held between the legs. His clothes were constantly saturated with these dribblings, and his health was materially damaged by constant straining, and the frequency with which he was compelled to make efforts to relieve the bladder. He had been under the care of many surgeons, both in London and elsewhere, but without obtaining amelioration of his sufferings. Having tried various sized and kinds of instruments, I was fortunate enough, at the expiration of three weeks, to get a 0 0 silver catheter through the obstruction. This instrument was retained, and replaced by a larger size, until No. 3 could be passed, when the dilator was introduced, and the No. 12 tube immediately passed. The same after-treatment was adopted as in the preceding cases, and in three months from his first visit he was enabled to pass No. 12 without difficulty. The stream was good, the intervals much longer, and his health better

than it had been for the previous seven years.

Another very similar case may be addedviz.: A gentleman, a solicitor by profession, was affected with a stricture of many years' standing. He had for ten years experienced great difficulty in emptying his bladder, and the urine was loaded with mucus. Upon the first examination, the catheter was arrested at three inches from the meatus; after some trouble it passed on to a second obstruction, near the membranous portion of the urethra, through which it was impossible at the time to penetrate. Upon subsequent trials, however, I got through the second obstruction, and came upon a third, nearer the bladder, which was eventually overcome. The dilator was at once introduced, and the three strictures split. A No. 12 catheter was immediately passed in, and the urine withdrawn. Considerable pain was for a time experienced in this instance in expelling the urine, which was only effected with great straining. The cause, however, of this suffering was speedily apparent, for the patient passed three small calculi, which had formed behind the several obstructions, and were washed away when the urethra became enlarged.

CASE IV.

Stricture in Front of the Triangular Ligament of Eighteen Years' Duration—Several Attacks of Retention—Infiltration and Abscess at the Root of the Penis—Operation—Subsequent Sloughing—Recovery after the Use of the Dilator.

J. R., a man of sallow aspect, was admitted into the Westminster Hospital under my care for stricture. He has been the subject of stricture for eighteen years, during which period he has had several attacks of retention of urine. The urine for some months has been passed guttatim, and occasionally in the smallest stream; and a week prior to his admission, during a violent effort to relieve himself, he felt a sharp burning pain at the root of the penis, with slight relief to his urgency, but no urine escaped per urethram. In the evening he had a distinct rigor, was feverish, and passed a restless night; and on the following morning noticed that the penis was swollen, of a dusky hue, and painful to the touch; his urine continued to escape in drops. The penis was fomented with warm water; but all his symptoms increasing in severity, at

the expiration of a week he was admitted into the hospital. I detected infiltration of urine, and an abscess at the root of the penis, which was immediately opened; and, to prevent further infiltration, and afford a free outlet for the urine, an incision was made into the urethra in the perinæum, posterior to the stricture, by thrusting in a sharp - pointed bistoury in front of the anus, with the back to the rectum, and cutting upwards and forwards during the straining of the patient. urethra being opened, a gum elastic catheter was pushed through the wound into the bladder, and the urine thus permitted to escape. The catheter was firmly secured, and a poultice applied to the wound. In the course of a few days the inflammation and swelling of the penis had subsided sufficiently to allow a No. 1 to be tried through the penis, but it was found impracticable to reach the bladder. Upon a second attempt, the catheter went farther, but would not penetrate the entire canal; the third attempt was more successful, and after considerable difficulty, the bladder was reached, and the urine withdrawn. The catheter was allowed to remain until the following morning, when it was replaced by the dilator, and the stricture at once split by

passing a No. 12 tube. My usual plan of treatment was adopted, and the urine withdrawn. On the following day, the patient expressed himself as having passed a tranquil night, without rigors, or even chilliness; the urine flowed in a moderate stream, and with but slight scalding.

On the third day from the operation, a No. 12 catheter was passed without difficulty, and this act was repeated on alternate days for a fortnight; the intervals were then gradually extended, and in six weeks from the operation, the perinæal opening having healed, he left the hospital, passing his own instrument, and making water in a full stream.

The above-recited case afforded another excellent opportunity of testing the efficiency of the treatment, and although undertaken at a time when the patient was hardly recovered from the effects of infiltration, it was attended with signal success; he never had a bad symptom, and in six weeks left the hospital, so far cured that the stricture was entirely under his own control.

CASE V.

Stricture of Thirty Years' Standing—Fistulæ in Perinæo—Incontinence of Urine—Temporary Impermeability of Urethra—Subsequent Treatment by Small and Large Dilators—Recovery.

J. R., aged 62, a well-formed and ordinarily a robust man, but now greatly emaciated, was admitted in 1857 under my care. He states he has been the subject of stricture thirty years, and attributes the origin of the disease to frequently-contracted gonorrheas. Fifteen years since he was the subject of retention of urine, which was at that time relieved by the introduction of a very small catheter, but being unable to continue in the hospital where he obtained relief, the contraction speedily became worse, and an abscess subsequently formed in the perinæum, which, upon being opened, gave exit to pus and urine, the latter continuing to escape more freely through the perineal opening than through the urethra. He now became an inmate of a provincial hospital, where every endeavour was made to get an instrument into the bladder without success, and he eventually left unrelieved, the urine continuing to dribble away, partly from the urethra and partly through the perineal opening, so that he was compelled to wear an apparatus specially made for the purpose of hindering the urine from saturating his clothes. No treatment having hitherto been of service, he came to London, and placed himself under the care of the late Mr Charles Guthrie, who eventually requested me to admit him under my care. Upon examination, I found his statement to be true, and that the urine continually escaped. A No. 1 catheter was employed, but it was impossible to pass it beyond the membranous portion of the urethra, which appeared to be so encroached upon as to be almost obliterated. Various attempts were made from time to time without success, and he was, consequently, placed under the influence of chloroform, and a No 1 catheter was forcibly passed into the bladder. This was permitted to remain, and on the following day was replaced by a gum elastic one. The patient did not suffer materially from the forced catheterism, but was relieved by the withdrawal of a large quantity of urine, and in three days after passing the No. 1 silver catheter, I was enabled to introduce the dilator, and split the stricture. The patient was then placed in bed, and the mixture of quinine and opium ordered. In the evening he was attacked with rigors, and experienced con-

siderable scalding in passing his water, which he continued to do in a narrow stream, and in small quantities. On the day following he was better, but the urine did not run freely, and on the second day after the operation, the No. 12 catheter was, with some little difficulty, passed into the bladder. A small quantity of urine escaped, and upon this occasion the catheter was allowed to remain five minutes, and was then withdrawn. The same line of treatment was adopted for many alternate days, but with a No. 11 instead of a No. 12 catheter, the stricture being irritable, and not admitting the larger size. A No. 12 was, however, eventually passed, and being taught to use the catheter for himself, the man returned to the country, making water in a full stream, and declaring himself perfectly cured. This patient was under treatment nine weeks.

Three cases of a precisely similar character have been under treatment, where incontinence of urine, the result of over-distension of the bladder, was present, the patient suffering from the absorption of the deleterious properties of the retained urine, the effects of which were recognised by drowsiness, pallor, nervousness, and tingling or itching of the skin. In each case the patient has been subjected to the same treatment, and with similar success.

CASE VI.

Traumatic Stricture—Severe Hæmorrhage— Elastic and Recurrent after Dilatation— Operation—Recovery.

J. H., aged 48, a master-builder, consulted me in June, 1860, in consequence of continued difficulty in passing his water. He stated that five years since, while walking along a roof, he suddenly fell across a beam. His perinæum was seriously bruised, and he experienced great pain in the urethra, from which a considerable quantity of blood escaped. He was taken home, and a Surgeon passed with some difficulty a gum elastic catheter, which was retained. The hæmorrhage gradually subsided, and the catheter was removed. He, however, soon became aware that his urine was passed with greater difficulty than formerly, and that he was compelled to strain a good deal to get rid of the contents of the bladder. He again consulted a Surgeon, who passed a No. 2 catheter into the bladder, and gradually increased the sizes of his instrument until a No. 12 was attained to. The catheterism being discontinued, the stricture soon returned, and he again consulted the same gentleman, who recommenced the treatment, and soon

arrived at No. 12. The convalescence was, however, of short duration, for upon the treatment being discontinued, the contraction speedily recurred. Upon examination, I found no difficulty in passing No. 2, but the stricture would not admit a No. 4, and No. 3 was very tight. I therefore immediately pushed in the dilator, split the stricture, and afterwards passed a No. 12 with facility. This treatment was perfectly successful. There was never the least hesitation in introducing a No. 12 afterwards, and he continued to eject his water in a copious stream without straining and at the ordinary periods.**

CASE VII.

Stricture of Eight Years' Duration—Difficult Micturition—Occasional Retention of Urine —No Complications.

W. B., aged 50, a clerk in a Government-office, was, in 1858, sent to me by Mr Jones, of

^{*} I may refer to two cases of severe traumatic stricture, treated by the same method by Mr Heath, and published in the 'Lancet,' August 31, 1861, as confirming the applicability of this treatment in such forms of the disease.

Kennington, in consequence of stricture of the urethra, attended with a continual difficulty in micturition, and complete retention of urine after any excess. At the time of my seeing him he was suffering from retention, which was immediately relieved by the introduction of a No. 2 catheter. Although there was no special difficulty, upon a subsequent occasion, in introducing a No. 3, yet as it was very tight, and in order to relieve him from his often-recurring retentions, I advised that the stricture should be split. The patient consented. The dilator was introduced, and the stricture immediately split. He was not subjected to the influence of chloroform. The pain was trifling, and after the operation he walked home. Such after - treatment was adopted as has been already described, and he occasionally presents himself to his surgeon, when a large-sized catheter can be passed with ease.

CASE VIII.

In June, 1859, I was consulted by a gentleman who had come from Paris for the purpose of having his stricture split. He had for many years suffered from this complaint, which had latterly become so aggravated that

his urine would only pass in the smallest stream, and occasionally by drops. Upon his first consulting me various catheters were tried without success. Nothing would penetrate the stricture, the point of the instrument becoming entangled in what he described as an old false passage. He was desired to remain perfectly quiet; to take castor-oil every other morning, have warm baths, and renew his visit in a week. Upon the second trial, and after very considerable difficulty, I was enabled to pass No. 1 into the bladder. A considerable quantity of urine was withdrawn, with great relief. He was recommended to avoid exercise, and to come again in three days, when another attempt was made with the No. 1 that had passed before, but without success. A week was, therefore, permitted to elapse before any further trial, when, after continued perseverance, I succeeded with a No. 1, which was allowed to remain for an hour, and was then replaced by No. 2. In the evening he had retention, but it was speedily relieved by the No. 1 catheter. He continued to attend every third day until No. 2 could be passed with some degree of facility, when the dilator was introduced and the stricture split, and No. 12 catheter immediately introduced: the urine

flowed through; and he walked home, having directions to remain quiet, and take the quinine and opium mixture.

On the second day from the operation a No. 12 was freely slipped into the bladder; he had had neither hæmorrhage nor shivering, and the utmost he complained of was a slight scalding. In ten days from the operation, this gentleman passed his own instrument, and in a fortnight returned to Paris, making water with ease, and using No. 12 without the least difficulty.

I saw this patient at the beginning of the present year. The No. 12 can now be passed with the greatest facility. He has never had either retention or difficulty since the operation, and may be considered well.

The cases which have been already narrated are selected as representing the most severe forms of stricture, and as being attended by complications which would materially militate against any ordinary operation, and must have necessitated months of surgical attendance if the dilating plan had been selected. In every one of the recited cases the treatment was successful, and with but a little instruction the patients were enabled to maintain the after-control of the bladder, by occasionally passing a bougie. Although this treatment is

specially advocated in the extreme forms of the disease, it is equally applicable in the milder or less complicated, more especially in cases where strictures are irritable and unyielding, and where catheterism is attended with difficulty and severe suffering.

In such instances the dilator gives immediate relief, and experience has proved that by splitting the stricture, tension is immediately overcome, and the passage of the catheter is made so facile, as to be deprived of any but the most trifling pain. The following cases will best illustrate this:—

J. R., aged 38, a barrister, consulted me in January, 1859, complaining of stricture of the urethra, from which he had suffered for the last eight years. He stated that he had had three attacks of gonorrhea, the discharge accompanying each attack lingering for a considerable period, and necessitating the use of stimulating injections for its cure. During 1858 his symptoms became much aggravated, he was obliged to strain greatly to get rid of the contents of his bladder, the stream of urine which had become spiral, and occasionally forked, was very small, and his sufferings were further aggravated by the protrusion

of piles during his efforts for relief. He had suffered from retention of urine, which upon one occasion followed a debauch, and on another resulted from inability to relieve himself during a railway journey.

Upon examination, I detected a stricture situated about the membranous portion of the urethra, through which a No. 2 catheter could, without material difficulty, be passed. The stricture was of short extent, apparently cartilaginous, and it grasped the catheter tightly. Having explained to the patient the rationale of the operation, he at once consented to its performance. The dilator was, therefore, immediately introduced, the stricture split, and a No. 12 catheter substituted for the purpose of withdrawing the contents of the bladder. There was the slightest possible bleeding, and he stated that the operation was not more severe than the passing of a catheter. rected him to remain at home during the afternoon, and to take the quinine and opium mixture.

This gentleman continued his attendance on alternate days for a fortnight; he never had the slightest difficulty in passing his water; the pain was very trifling, and a No. 12 catheter slipped into the bladder with the greatest ease. Being now taught to pass his own instrument, he was discharged cured.

J. J., aged 61, a gentleman, residing in Wales, consulted me in December, 1858, for stricture of many years' duration. He stated that during the previous year, the stream of urine had become materially decreased in size, that he was obliged to pass water every two hours, that the urine was highly ammoniacal and loaded with mucus, that the irritable state of his bladder compelled him to make water the instant he had the desire to do so; and that his clothes were constantly wetted, from his inability to eject the urine in a continuous stream. His rest was broken, and his health materially damaged by the absorption of the obnoxious properties of the retained urine. Upon examination, I detected a stricture at the triangular ligament, which would only admit a No. 1 catheter. The obstruction was dense and unyielding, and about half an inch in length. As the patient was somewhat plethoric, I prescribed a purgative, and requested his attendance on the following day, at which visit the dilator was with slight difficulty introduced, and the stricture split. A No. 12 catheter was now substituted for the dilator,

and the urine withdrawn. In this case, as in the former one, the cure was uninterrupted by a single unfavourable symptom; the patient was merely confined to the house one afternoon, and the same plan of after-treatment having been carried out, he, at the end of a fortnight, returned to Wales, capable of passing his own catheter.

I have had several letters from this gentleman since, in all of which he declares himself free from the slightest difficulty in passing his water or introducing his catheter.

The foregoing cases have been briefly described, and no allusion or comparison has been made to the various kinds of treatment already published. Their perusal must have satisfied the most sceptical that, in every instance where any kind of instrument can be passed into the bladder, the urethra may, by the mechanical effect of the dilator, be immediately enlarged to its natural size; and that while in slight cases this enlargement is effected without suffering, so as to render the inhalation of chloroform unnecessary, yet in the more severe forms that agent facilitates the operation, and secures a perfect immunity from pain. It is quite true that if the after-

treatment is not attended to, the stricture will sooner or later recur, but as the patient by passing his own instrument has the control of the bladder, it can only return as a consequence of culpable neglect. The immunity from accidents having been already proved, the surgeon need have no fear of those serious results which so frequently accompany any cutting operation, and it is no less extraordinary than true that whereas rigors ordinarily supervene after rapid dilatation, they form the exception where the stricture has been fairly split.

It may be asked, what advantages accrue from the performance of this operation, when it is necessary that the after-treatment should be continued for a variable period of time, and why the ordinary plan of dilatation should be abandoned? My reply is—1st, that by its performance a large-sized catheter can be immediately passed, and the patient be speedily taught to use his own instrument; 2nd, that the patient avoids all the suffering incidental to gradual dilatation, and the frequent disappointment of not being enabled to increase the sizes of the instrument; 3rdly, that it secures all the advantages that can accrue from the performance of any operation, and with much

less danger to the patient than by any cutting operation; and, lastly, that the period of recovery is shorter, it does not necessitate confinement to bed, and the patient is saved the expense of being constantly under the hands of his surgeon.

These observations apply to those cases where dilatation can be continued in the ordinary manner; but all practical surgeons know that there are a large number of cases where dilatation is ineffectual in advancing beyond a certain gauge; and here some operation must be performed, if the patient is not to continue a surgical annuity.

I am fully aware that in advocating a plan of treatment by which the stricture is split or torn, prejudices have to be overcome, from the fears which naturally arise as to the extent of the rupture, and the consequences, not only to the urethra, but also to the immediately investing structures. What may be the precise limit of this rupture in the living body I have not as yet had an opportunity of ascertaining, the treatment having been attended hitherto with unvarying success; and when already more than 100 cases have been operated upon by myself, in addition to those where the operation has been undertaken by other sur-

geons, in all of which the treatment has been most efficacious, it is not too much to hope that the danger must be very limited, when compared with any of the cutting operations. What are the advantages of incising the urethra in those obstinate forms of stricture which necessitate some operation? Is it a more simple operation, less dangerous, and possessing any peculiar advantages in reference to a recurrence of the disease? I unhesitatingly state that there are no such advantages, but that, on the contrary, the operation is more difficult, and requires considerable experience for its performance. The operator, in only a very limited number of cases, can insure the limitation of his incision to the strictured part, and for its performance it is necessary that some instrument should reach the bladder. Do the records of these operations give that degree of success which has been attained by the operation I advocate; and, setting aside the complications of hæmorrhage, infiltration of urine, abscess, &c., which so very frequently ensue, does it offer any better chance of permanent cure than the division of stricture by rupture? Is not the aftertreatment more tedious, the patient being confined to bed with a retained catheter until the greater portion of the wound has healed; and

when this has united, does it not necessitate the rigid enforcement of the after-passage of the catheter, to ensure a prevention of the contraction?

The only difference between an incised wound and a rent, is that the former, if the edges are left approximated, heals much more quickly than the latter, and so more speedily induces a recurrence of the contraction. It is by stretching the uniting medium of either incision or rent that the cure is to be accomplished, just in the same manner as the orthopædic surgeon extends the soft union of a divided tendon.

Mr Syme, with that ingenuity and surgical skill which have so long placed him in the highest rank of our profession, devoted his attention to the treatment of this most important disease, and devised a plan which, at the time of its publication, offered, in the more obstinate forms of stricture, the best chances of success, more especially when the contraction was situated in front of the anus. But even in the hands of this scientific surgeon, the operation has not been without its attendant evil consequences, and in those of surgeons of less experience, has frequently terminated fatally.

Knowing the dangers incidental to perineal section, various surgeons, both at home and

abroad, have endeavoured to substitute an internal for an external incision, and various instruments have been invented for the purpose of incising the stricture from within; but with all there is the serious difficulty of not knowing with accuracy the exact structure which is being divided, and as the cutting portion of the instrument is usually set to divide a stricture, of the extent of which the surgeon is ignorant, it has been frequently found to cut either beyond the thickness of the stricture, or to have divided a portion of the urethra, either in front of or behind the obstruction; hence, the frequent accompaniment of serious hæmorrhage, infiltration of urine, and abscess.

To accomplish either operation, it is necessary that some instrument should pass through the stricture; and when that can be effected, I maintain that no cutting operation is either necessary or justifiable, and simply from the fact, that where any instrument can be passed, there the dilator can enter, and by the introduction of the tube (which, with the dilator, should only represent the natural diameter of the urethra), we can split or tear that which is the seat of obstruction; and as far as I have yet had an opportunity of judging, the obstruction only. There is also another material advantage

which attaches to splitting in preference to cutting-viz., that the surgeon is not obliged to keep any instrument in the bladder, but the patient passes his urine in a natural manner ever afterwards, the catheter being introduced at first on alternate days, and afterwards at longer intervals, until the canal is so far widened that the occasional passage of the bougie will maintain its calibre. While, however, I am unquestionably of opinion, from the large number of cases that I have operated upon, that the operation by rupture is the most feasible, the easiest to perform, and the least dangerous, I do not place it before the profession as a means of cure, unless the after-passage of the bougie is faithfully carried out. We know not at present of any means by which a stricture can be cured without such aftertreatment, any more than the orthopædic surgeon knows of any means of curing deformities without the application of some extending apparatus after a tendon has been divided. Our extending, or rather distending, means consists in the introduction of the catheter, or sound, and I care not whether the operation be by incision or splitting, the disease will recur if the contracted part of the canal is not kept dilated for a given time, such period varying

with the nature and number of the strictures, the irritability of the patient, and the complications of false passage, fistulæ in perinæo, &c., &c., which may exist.

The advocates for internal incision affirm, that inasmuch as the knife is applied solely to that portion of the urethra which constitutes the obstruction, they divide the stricture only; I must, however, express my doubt as to the accuracy of this statement, and record my belief that it is quite impossible to determine, at six or seven inches from the meatus, which side of the urethra is thickened; the sound, or cutting instrument, is firmly grasped, and unless the seat of thickening, or deposit, is always the same, it will be quite impossible, from the sensation conveyed to the surgeon, to know which part to incise.

The numerous preparations in the different museums prove that, although the under part of the urethra may be the most frequently diseased, yet that the exceptions are so numerous as practically to render that fact of no value, and that where the obstruction is of the obstinate character met with in the cases I have detailed, the thickening may be taken to be tolerably circumferential.

Suppose it were otherwise, and that the

upper segment of the urethra were not involved, what should prevent the onward passage of a catheter when it has entered the contraction, since the upper part, if normal, would certainly yield, and permit of its passing to the bladder? The truth is, we have no security that, in the employment of a cutting instrument, our incisions are confined to the stricture alone, or even that the weakest part of the urethra may not be divided, and thus give rise to infiltration of urine and abscess, an occurrence by no means unfrequent where such means are employed. Hitherto, any treatment beyond ordinary dilatation has been considered applicable only to cases where the most severe complications are met with, and where the difficulties have been such as to compel the patient to place himself under the care of a surgeon who has great experience in the treatment of such cases; but the ordinary circumstances of cases, as they are usually presented to us, offer no obstacle to their immediate subjection to the practice I advocate. As a general rule, a patient seeks professional aid when a No. 3 or No. 4 can be introduced; and here, no matter what may be the exact pathological change, the stricture may be immediately split, the catheter being afterwards passed upon three

or four occasions by the surgeon, when the further treatment of the case may usually be transferred to the patient, who is now capable of passing his own instrument. The three following cases will best illustrate this:—

J. W., aged 45, consulted me in January, 1858, in consequence of difficulty in micturition, the result of stricture at about six inches from the meatus. His urethra would admit a No. 3, but it was exceedingly tight; and as he was desirous of returning to the country, I advised that the stricture should be immediately split. The dilator was at once passed, and the stricture split, so that a No. 12 catheter could be introduced into the bladder. The urine having been withdrawn, the catheter was removed, and he was directed to return home, and remain quiet the same afternoon. He again presented himself on the second day after the operation, stating that he had no constitutional disturbance, and that the stream of urine was improved. A No. 12 catheter was again introduced, and without any drawback; this was repeated on three occasions, when, as he had been previously in the habit of passing a No. 3, he had no difficulty in introducing a No. 12, and he returned to the country. I have had frequent opportunities of seeing this gentleman since, but he has never required to consult me again respecting his stricture.

Mr H., a gentleman of colour, consulted me for stricture, situated at the membranous portion of the urethra. He had suffered from difficult micturition for about eighteen years; there were, however, no complications, and his stricture would admit a No. 3, which for some years he had been in the habit of passing. As he was a nervous person, he was placed under the influence of chloroform, when the dilator was introduced, and the stricture split with the No. 12 tube. The 12 catheter was then introduced, and the urine removed. There was scarcely any bleeding; he never had a bad symptom, or took medicine, and in a fortnight returned to Australia, passing his No. 12 catheter, which, with others, in case of fresh cause of difficulty, he procured from Messrs Whicker and Blaize

Mr B., aged 65, residing at Hampstead, consulted me for stricture of thirty years' duration. He had been occasionally the subject of retention of urine, which was usually relieved by warm baths and opium. When I first saw him, in 1858, I could only introduce No. 1, and as he was in great dread of any operation, I

consented to dilate the stricture in the ordinary manner. By the gradual process, I eventually arrived at No. 8, when, from circumstances, he was compelled to discontinue his attendance; and upon his again consulting me, in the latter part of 1859, the stricture had so far returned that I was compelled to begin de novo. This time I persuaded him to have the stricture split, and, without the aid of chloroform the dilator was passed, and the stricture split to No. 13, his urethra being sufficiently capacious to take that size. Although an old man for his age, he never had a bad symptom, and ceased to continue under my care after my teaching him to pass his large-sized catheter, which he accomplished in about three weeks.

Is there, then, any special class of strictures where the operation by rupture is inadmissible, either from the situation, kind, and number of strictures, or from other complications? I believe not; but that there are a large number where any cutting operation must be attended with considerable danger, is now well known. Let us, for example, take a class of cases frequently met with, where fistulæ in perinæo, or elsewhere, exist, combined with great hyperthrophy of all the tissues

in the immediate contiguity of the stricture, and in which considerable time must be expended, and much skill employed before any instrument can be passed into the bladder. Here, to cut into a small groove requires great coolness, determination, and experience; and even when the operator is successful in reaching the groove, and so far dividing the obstruction as to admit a large catheter, the patient is confined to bed for many weeks before union will take place sufficiently to allow of the withdrawal of the instrument. And, again, what is the advantage of dividing a mass of hypertrophied tissue which does not encroach upon the canal, but necessarily intervenes between the knife and the urethra? If the diameter of the urethra can be enlarged to the same extent without such a proceeding, and a free outlet afforded for the urine, the hypertrophy will diminish, and the parts will eventually be restored to their original integrity.

But let me refer to another class where three or more strictures of varying density exist at different parts of the canal. Is the urethra to be laid open from end to end, or can the three obstructions be divided by internal incision at one and the same operation? I have seen these operations performed by the best

surgeons in London, where no possible objection could be urged against their anatomical knowledge, their surgical experience, or their operative dexterity under the most trying circumstances; and I regret to say that they have not been without a fatal issue. No doubt it looks more surgical to place a patient in the position of lithotomy, and undertake a difficult piece of dissection in his perinæum, than to simply introduce an instrument, and pass a large tube, as it were, blindly between its blades, but the results of such cases warrant me in doubting the wisdom of such a proceeding, when the same result can be attained in another manner without difficulty or danger.

Hitherto I have only considered the applicability of the dilator, so far as rupturing or splitting the stricture is concerned, but it is equally efficacious where dilatation is desired, and possesses an advantage over every kind of bougie or sound in its power of dilating a stricture to any required extent without being withdrawn. All practical surgeons know that, having, for instance, had some difficulty in passing a No. 1, it is frequently impossible to introduce a No. 2 at the same visit, whereas, having once introduced the dilator, its diame-

ter can be increased to any extent the surgeon may desire; for this purpose it is not necessary that consecutive tubes should be passed, as by introducing at once a No. 8 or No. 9, and very gently pressing the tube onwards between the blades, they become separated to a considerable distance from the point of the tube, as is excellently shown in the drawing (page 4). By this means (which, however, occupies some little time) the stricture being acted upon from within, a greater amount of dilatation can be effected at one visit, and with much less pain to the patient than by passing consecutive bougies, which irritate the urethra, and owing to the abrupt increase of size, sometimes will not enter the stricture at all. If, in adopting the gradually dilating plan, the tube is very slowly passed, the pain will be trifling, and it will entirely cease, in the majority of instances, in a minute or two after the progress of the tube is discontinued. I cannot conceive any instrument more fitting, where gradual dilatation is desired; the blades can be expanded with the greatest nicety, in entire obedience to the desire of the surgeon, and the feelings of the patient, and as the dilatation is effected from within outwards, its force is expended in the most advantageous

But, although dilatation may be direction. employed where time is no object, and where the stricture is of a yielding character, it is utterly incapable of effecting what may be accomplished by passing the large tube at once, and splitting the stricture; the increased diameter being maintained by the after-passage of bougies. The rapidity with which the case will terminate, so far as only to require the passage of a bougie twice or thrice in the course of the year, will depend upon the number of strictures, and the irritability of the patient, more than it will upon their density. In some cases, although great force is required to split the obstruction, the urethra is to a certain extent insensible, and bears the after-passage of the bougie without the least manifestation of pain, whilst in others, more especially where there have been three or more impediments, and the urethra is very irritable, it is necessary to recede a size or two, owing to the spasms being so great during the introduction of the larger instrument, as to cause suffering.

I have lately had a marked example of this in an artilleryman, sent to me by my friend, Dr Gallway, Surgeon-major of the Royal Artillery. The patient had been the subject of stricture about seven years; he had been through the Crimean war, and an inmate of the military hospital, more or less, ever since his return. Very many and long-continued attempts were made to get a catheter into his bladder without success, and Dr Gallway requested me to meet him in consultation upon the case. I did so, but was equally unsuccessful in passing a catheter. I therefore requested Dr Gallway to allow him to be admitted into the Westminster Hospital, to which he consented, and after several trials, I at last succeeded in passing No. 1 through four obstructions into the bladder. As the patient was excessively nervous and irritable, I delayed splitting the stricture until No. 2 was attained, when the stricture was split with the No. 10 tube, and a 10 catheter immediately passed: he did not suffer from the operation in the slightest degree, and his urine was passed in an improved stream. On the second day from the operation, I attempted to pass the No. 10, but the spasm was so persistent that I preferred having recourse to No. 8 rather than give him pain; and I have been up to the present, now three weeks after the operation, content with that size, which passes with great ease, but still excites spasm. I have no question that in a little time No. 10 may be again used. The urethra must necessarily remain sore under such circumstances, and thus excites an amount of spasm which, by grasping the catheter, adds to the patient's suffering.


In conclusion, I may add, that in advocating the treatment of stricture by rupture I claim simply that credit which attaches to the publication of a series of interesting cases (examples of many others) which have been subjected to this novel treatment. That the principle upon which the instrument is constructed is as old as the hills, and that the power of the wedge has been known as long as the simplest rules of mechanics have been taught, I freely admit, but I have yet to learn that that principle has been heretofore applied to the treatment of stricture of the urethra in the manner detailed above, and with such highly satisfactory results.

Since the foregoing pages were written, I have had an opportunity of ascertaining what is the exact condition of the urethra when the stricture has been split after death.

A man was admitted under me into the Westminster Hospital in Oct., 1861. He had been the subject of stricture for eighteen years, and for the last twelve years the bladder had been relieved with increasing difficulty until he could only pass his water in drops. I endeavoured to introduce a No. 1 catheter, but found the impediment too contracted to admit even that size, and he was sent to bed, with directions to take castor-oil and remain quiet for a few days. In the meantime he was attacked with fever, and he died on Sunday, Nov. 10th. The friends objecting to a general post-mortem examination, I ultimately succeeded in persuading them to permit the urethra to be examined, which I was exceedingly anxious to do, for the purpose of ascertaining what was the precise effect produced by the passage of the dilator and its large tube. With some difficulty I introduced the dilator through two strictures, one within four inches of the meatus, and the second at the bulb, both apparently about half an inch long. The sensation was

like that I have frequently experienced during life, and the dilator having fairly reached the bladder, I passed the No. 12 tube, split the strictures, and before withdrawing it, rotated the dilator in the manner already described. My friend, Mr Heath, now removed the penis, with a portion of the bladder and rectum, and having laid open the urethra, the two strictures, as is shown in the drawing, were found to be split at the under portion of the canal, the rent being directly in the median line, and limited to the extent of the obstruction. The contiguous structures were uninjured, and the divided mucous membrane gaped to an extent sufficient to permit the passage of a large bougie.





Mr. Churchill is the Publisher of the following Periodicals, offering to Authors a wide extent of Literary Announcement, and a Medium of Advertisement, addressed to all Classes of the Profession. Communications, Books for Review, addressed to the respective Editors, are received and duly forwarded by Mr. Churchill.

THE BRITISH AND FOREIGN MEDICO-CHIRURGICAL REVIEW, AND QUARTERLY JOURNAL OF PRACTICAL MEDICINE AND SURGERY. Price Six Shillings. Nos. I. to LV.

THE QUARTERLY JOURNAL OF MICROSCOPICAL SCIENCE.

Edited by Edwin Lankester, M.D., F.R.S., F.L.S., and George Busk, F.R.C.S.E., F.R.S., F.L.S. Price 4s. Nos. I. to IV. New Series. * * A few Nos. of the Old Series are out of print; the others may be obtained.

THE JOURNAL OF MENTAL SCIENCE.

Published by authority of the Association of Medical Officers of Asylums and Hospitals

Edited by John Charles Bucknill, M.D. Published Quarterly, price Half-a-Crown. New Series. Nos. I. to III.

OF MEDICINE: ARCHIVES

A Record of Practical Observations and Anatomical and Chemical Researches, connected with the Investigation and Treatment of Disease. Edited by LIONEL S. BEALE, M.B., F.R.S. Published Quarterly, price 2s. 6d. from No. IX.; Nos. I. to VIII., 3s. 6d.

THE ROYAL LONDON OPHTHALMIC HOSPITAL REPORTS, AND JOURNAL OF OPHTHALMIC MEDICINE AND SURGERY.

Published Quarterly, price 2s. Nos. I. to XV.

THE MEDICAL TIMES AND GAZETTE.

Published Weekly, price Sevenpence, or Stamped, Eightpence. Annual Subscription, £1.10s., or Stamped, £1.14s. 8d., and regularly forwarded to all parts of the Kingdom.

The MEDICAL TIMES AND GAZETTE is favoured with an amount of Literary and Scientific support which enables it to reflect fully the progress of Medical Science, and insure for it a character, an influence, and a circulation possessed at the present time by no Medical Periodical.

HALF-YEARLY ABSTRACT OF THE MEDICAL SCIENCES.

Being a Digest of the Contents of the principal British and Continental Medical Works; together with a Critical Report of the Progress of Medicine and the Collateral Sciences. Edited by W. H. RANKING, M.D., Cantab., and C. B. RADCLIFFE, M.D., Lond. Post 8vo. cloth, 6s. 6d. Vols. I. to XXXIII.

PHARMACEUTICAL JOURNAL.

New Series. Published Monthly, price One Shilling. * * Vols. I. to XX., bound in cloth, price 12s. 6d. each.

BRITISH JOURNAL OF DENTAL SCIENCE.

Published Monthly, price One Shilling. Nos. I. to LXVI.

THE DUBLIN MEDICAL PRESS.

Published Weekly, Stamped, price Sixpence, free to any part of the Empire.

THE MEDICAL DIRECTORY FOR THE UNITED KINGDOM

Published Annually. 8vo. cloth, 10s. 6d.

A CLASSIFIED INDEX

TO

MR. CHURCHILL'S CATALOGUE.

ANATOMY.	DISEASES of the URINARY	MATERIA MEDICA and
PAGE		
Anatomical Remembrancer 3	and GENERATIVE ORGANS,	
Beale on Liver 5	and SYPHILIS.	PAGE
Flower on Nerves 12		Bateman's Magnacopia 5
Hassall's Micros. Anatomy 14	Acton on Urinary Organs 7	Beasley's Formulary 5
Holden's Human Osteology 15	Do. on Reproductive Organs 3	Do. Receipt Book 5
Do. on Dissections 15	Coote on Syphilis 10	Do. Book of Prescriptions 5
Jones' and Sieveking's Patho-	Coulson on Bladder 10	Pereira's Selecta e Præscriptis 22
logical Anatomy 17	Do. on Lithotomy 10	Pharmacopæia Londinensis 22
Maclise's Surgical Anatomy 19	Gant on Bladder	Prescriber's Pharmacopæia 22
Paget's Catalogue 21	Judd on Syphilis 17	Royle's Materia Medica 24
Sibson's Medical Anatomy 25	Milton on Gonorrhea 20	Steggall's First Lines for Che-
Waters' Anatomy of Lung 29	Parker on Cymbilia	mists 26
Wheeler's Anatomy for Artists 30	Parker on Syphilis 21	Stowe's Toxicological Chart 26
Wilson's Anatomy 30	Wilson on Syphilis 31	Taylor on Poisons 27
********		Wittstein's Pharmacy 31
CHEMISTRY.	*********	********
Abel & Bloxam's Handbook 3	DISTACES OF THOSE OF	MEDICINE.
Bowman's Practical Chemistry 7	DISEASES OF WOMEN	Adams on Rheumatic Gout 3
Do. Medical do. 7	AND CHILDREN.	Addison on Supra-Renal Capsules 3
Fownes' Manual of Chemistry 12		Addison on Cells 3
Do. Actonian Prize 12	Ballard on Infants and Mothers 4	Anderson on Fever 3
Do. Qualitative Analysis 12	Barker on Children 4	Barclay on Medical Diagnosis 4
Fresenius' Chemical Analysis 12	Bennet on Uterus 6	Barlow's Practice of Medicine 4
Galloway's First Step	Do. on Uterine Pathology 6	Basham on Dropsy 4
Do. Analysis 12	Bird on Children 7	Beale on Urine 5
Griffiths' Four Seasons	Blake on the Skin in Children 7	Bird's Urinary Deposits 6
Horsley's Chem. Philosophy 16	Eyre's Practical Remarks 11	Brinton on Stomach 7
Jones.—Mulder on Wine 17	Hood on Scarlet Fever 16	Do. on Ulcer of do 7
Plattner on Blowpipe 22	Kiwisch (ed. by Clay) on Ovaries 9	Budd on the Liver 8
Speer's Pathol. Chemistry 26	Lee's Ovarian & Uterine Diseases 18	Do. on Stomach 8
**********	Lee on Diseases of Uterus 18	Camplin on Diabetes 8
CLIMATE,	Do. on Speculum 18	Chambers on Digestion 8
	Smith on Leucorrhœa 25	Davey's Ganglionic 11
Barker on Worthing 4	Tilt on Diseases of Women 27	Dobell's Germs and Vestiges of
Bennet on Mentone 6	Do. on Change of Life 27	Disease 11
Dalrymple on the Climate of	Underwood on Children 28	Evre on Stomach 11
Egypt 10	West on Women 30	Fuller on Rheumatism 12
Francis on Change of Climate 12		Gairdner on Gout 12
Hall on Torquay 14	********	Gibb on Throat 13
Haviland on Climate 14		Granville on Sudden Death 13
Lee on Climate 18	HYGIENE.	Gully's Simple Treatment 13
McClelland on Bengal 19 McNicoll on Southport 19		Habershon on Stomach 13
Menticoll on Southport 19	Armstrong on Naval Hygiene 4	Do. on Mercury 13
Martin (J. R.) on Tropical 20	Beale's Laws of Health 5	Hall on Apnœa 13
Moore's Diseases of India 20	Do. Health and Disease 5	Hall's Observations 13
Scoresby-Jackson's Climatology 24	Bennet on Nutrition 6	Harrison on Lead in Water 14
Taylor on Pau 27	Blundell's Medicina Mechanica 7	Hassall on Urine 14
*********	Carter on Training 8	Headland on Medicines 14
DEFORMITIES, &c.	Chavasse's Advice to a Mother 9	Hooper's Medical Dictionary 16
Riggs on Doformatition	Do. Advice to a Wife 9	Hooper's Physician's Vade-
Righon on Doformittee	Granville on Vichy 13	Mecum 13
Do. Articulate Sounds 6	Hartwig on Sea Bathing 14 Do. Physical Education 14	Hughes on Blood Disease 16
Brodhuret on Coine		Inman's New Theory 16
Do on Clubbeach	Hufeland's Art 16 Lee's Watering Places of England 18	James on Laryngoscope 17
Godfroy on China	Do Phonish Wetering Places 10	Jones' Animal Chemistry 17
Hare on Spine 13	Do. Rhenish Watering Places 18 Parkin on Disease	Lugol on Scrofula 19
Hugeman on Him Toint	Pickford on Hygiene 22	Marcet on Chronic Alcoholism . 19
Inman on Myalgia 16	Robertson on Diet 22	Parkes on Urine
Tamplin on Spine 26	Routh on Infant Feeding	Peacock on Influenza 21
***************************************	Rumsey's State Medicine 24	Pym on Yellow Fever 23
DENTISTRY.	Wells' Seamen's Medicine Chest 29	Richardson's Asclepiad 23
	Wife's Domain 30	Roberts on Palsy 24
Clark's Odontalgist 9	Wilson on Healthy Skin 31	Robertson on Gout 24
Gray on the Teeth	Do. on Mineral Waters 31	Savory's Compendium 24
Tomes' Dental Surgery 28	Do. on Turkish Bath 31	Semple on Cough 25 Shaw's Remembrancer 25
0	on running parit 31	Shaw's Remembrancer 25

CLASSIFIED INDEX.

	AND THE PARTY OF THE PARTY OF THE PARTY.	
MEDICINE—continued.	OPHTHALMOLOGY.	SCIENCE.
Smee on Debility 25	Cooper on Injuries of Eye 10	Baxter on Organic Polarity
Steggall's Medical Manual 26	Do. on Near Sight 10	Bentley's Manual of Botany
Steggall's Gregory's Conspectus 26	Dalrymple on Eye 10	Bird's Natural Philosophy
Do. Celsus 26	Dixon on the Eye 11	Burnett's Philosophy of Spirits
Thomas' Practice of Physic 27	Hogg on Ophthalmoscope 15	Craig on Electric Tension 1
Thudichum on Urine 28	Holthouse on Strabismus 15	Hardwich's Photography 1
Todd's Clinical Lectures 28	Do. on Impaired Vision 15	Hinds' Harmonies 1
Wegg's Observations 29	Hulke on the Ophthalmoscope 16	Jones on Vision 1
Wells on Gout 29	Jacob on Eye-ball 16	Do. on Body, Sense, and Mind 1
What to Observe 19	Jago on Ocular Spectres 16	Mayne's Lexicon 2
Whitehead on Transmission 30	Jones' Ophthalmic Medicine 17	Pratt's Genealogy of Creation 2
Williams' Principles 30	Do. Defects of Sight 17	Price's Photographic Manipula-
Wright on Headaches 31	Do. Eye and Ear 17	tion 2
	Nunneley on the Organs of Vision 21	Rainey on Shells 2
******	Walton on the Eye 29	Reymond's Animal Electricity 2
	***********	Taylor's Medical Jurisprudence 2
MICROSCOPE.	DITURIOTORY	Vestiges of Creation 2
Beale on Microscope in Medicine 5	PHYSIOLOGY.	Sequel to ditto 2
Do. How to Work 5	Beale on Tissues 5	Unger's Botanical Letters 2
Carpenter on Microscope 8	Carpenter's Human 8	
Schacht on do 24	Do. Comparative 8	*********
401 11 11 21	Do. Manual 8	
A111111111	Heale on Vital Causes 15	SURGERY.
	Hilton on the Cranium 15	
MISCELLANEOUS.	Richardson on Coagulation 23	Adams on Reparation of Tendons
	Virchow's (ed. by Chance) Cel- lular Pathology 9	Do. Subcutaneous Surgery
Acton on Prostitution 3		Anderson on the Skin
Bascome on Epidemics 4	********	Ashton on Rectum
Bryce on Sebastopol 8	PSYCHOLOGY.	
Cooley's Cyclopædia 9	Arlidge on the State of Lunacy 4	Bellingham on Aneurism
Forbes' Nature and Art in Disease 12	Austin on Paralysis 4	Bigg on Artificial Limbs
Guy's Hospital Reports 13	Bucknill and Tuke's Psycholo-	Brodhurst on Anchylosis
Haycock's Veterinary 14	gical Medicine 8	Bryant on Diseases of Joints
Lane's Hydropathy 18	Burgess on Madness 7	Chapman on Ulcers
Lee on Homœop, and Hydrop, 18	Burnett on Insanity9	Do. Varicose Veins
Marcet on Food 19	Conolly on Asylums 9	Cooper (Sir A.) on Testis
Massy on Recruits 20	Davey on Nature of Insanity 11	Do. (S.) Surg. Dictionary
Part's Case Book 21	Dunn's Physiological Psycho-	Curling on Rectum
*********	logy 11	Do. on Testis
	Hood on Criminal Lunatics 16	Druitt's Surgery
NERVOUS DISEASES AND	Millingen on Treatment of In-	Fergusson's Surgery
INDIGESTION.	sane 20	Heath's Minor Surgery and
INDIGESTION.	Monro on Private Asylums 20	Bandaging
Birch on Constipation 6	Noble on Mind 20	Higginhottom on Nitrate of Silver
Carter on Hysteria 8	Williams (J.) on Insanity 30	Hodgson on Prostate
Child on Indigestion 9	Williams (J. H.) Unsoundness of	James on Hernia
Downing on Neuralgia 11	Mind 30	Jordan's Clinical Surgery
Hunt on Heartburn 16		Lawrence on Ruptures
Leared on Imperfect Digestion 18	PULMONARY and CHEST	Liston's Surgery
Lobb on Nervous Affections 19		Macleod's Surgery of the Crimea
Radcliffe on Epilepsy 23	DISEASES, &c.	Maclise on Fractures
Reynolds on the Brain 23	Addison on Healthy and Dis-	Maunder's Operative Surgery.
Do. on Epilepsy 23	eased Structure 3	Nunneley on Erysipelas
Rowe on Nervous Diseases 24	Alison on Pulmonary Consump-	Pemberton on Melanosis
Sieveking on Epilepsy 25	tion 3	Pirrie's Surgery
Turnbull on Stomach 28	Billing on Lungs and Heart 6	Price on Scrofula
********	Blakiston on the Chest 7	Smith on Stricture
***************************************	Bright on the Chest 7	Do. on Hæmorrhoids
OBSTETRICS.	Cotton on Consumption 10	Snow on Chloroform Steggall's Surgical Manual
	Do. on Stethoscope 10	
Barnes on Placenta Prævia 4	Davies on Lungs and Heart 11	Teale on Amputation
Davis on Parturition	Dobell on the Chest 11	Do. on Prostate
Lee's Clinical Midwifery 18	Fenwick on Consumption 11	Toynbee on Ear
Pretty's Aids during Labour 22	Laennec on Auscultation 18	Wade on Stricture
Priestley on Gravid Uterus 22	Markham on Heart 20	Watson on the Larynx
Ramsbotham's Obstetrics 23	Richardson on Consumption 23	Webb's Surgeon's Ready Rules
Do. Midwifery 23	Salter on Asthma 24 Skoda on Auscultation 20	Williamson on Gunshot Injuries 3
Cinclair & Labratanta Midwifform Or	SKOON ON AUSCHIINTION 20	
Sinclair & Johnston's Midwifery 25	Thompson on Consumption 97	Wilson on the Skin
Smellie's Obstetric Plates 25	Thompson on Consumption 27	Wilson on the Skin
Smellie's Obstetric Plates 25 Smith's Manual of Obstetrics 25	Thompson on Consumption 27 Timms on Consumption 28	Do. Portraits of Skin Diseases
Smellie's Obstetric Plates 25	Thompson on Consumption 27 Timms on Consumption 28 Turnbull on Consumption 28	Wilson on the Skin

MR. F. A. ABEL, F.C.S., & MR. C. L. BLOXAM.

HANDBOOK OF CHEMISTRY: THEORETICAL, PRACTICAL, AND TECHNICAL. Second Edition. 8vo. cloth, 15s.

MR. ACTON, M.R.C.S.

A PRACTICAL TREATISE ON DISEASES OF THE URINARY AND GENERATIVE ORGANS IN BOTH SEXES. Third Edition. 8vo. cloth, £1. 1s. With Plates, £1. 11s. 6d. The Plates alone, limp cloth, 10s. 6d.

THE FUNCTIONS AND DISORDERS OF THE REPRODUC-TIVE ORGANS IN YOUTH, IN ADULT AGE, AND IN ADVANCED LIFE. Considered in their Physiological, Social, and Psychological Relations. Second Edition. 8vo. cloth, 7s.

PROSTITUTION: Considered in its Moral, Social, and Sanitary Bearings, with a View to its Amelioration and Regulation. 8vo. cloth, 10s. 6d.

DR. ADAMS, A.M. TREATISE RHEUMATIC GOUT; OR, 0NRHEUMATIC ARTHRITIS. 8vo. cloth, with a Quarto Atlas of Plates, 21s.

MR. WILLIAM ADAMS, F.R.C.S.

ON THE REPARATIVE PROCESS IN HUMAN TENDONS AFTER SUBCUTANEOUS DIVISION FOR THE CURE OF DEFORMITIES. With Plates. 8vo. cloth, 6s.

PRINCIPLES AND PRACTICE SKETCH 0FTHE SUBCUTANEOUS SURGERY. 8vo. cloth, 2s. 6d.

DR. ADDISON.

ON THE CONSTITUTIONAL AND LOCAL EFFECTS DISEASE OF THE SUPRA-RENAL CAPSULES. 4to. cloth. Coloured Plates, 21s.

DR. WILLIAM ADDISON, F.R.S.

CELL THERAPEUTICS. 8vo. cloth, 4s.

ON HEALTHY AND DISEASED STRUCTURE, AND THE TRUE PRINCIPLES OF TREATMENT FOR THE CURE OF DISEASE, ESPECIALLY CONSUMPTION AND SCROFULA, founded on MICROSCOPICAL ANALYSIS. 8vo. cloth, 12s.

DR. SOMERVILLE SCOTT ALISON, M.D. EDIN., F.R.C.P. THE PHYSICAL EXAMINATION OF THE CHEST IN PUL-MONARY CONSUMPTION, AND ITS INTERCURRENT DISEASES. With Engravings. 8vo. cloth, 12s.

THE ANATOMICAL REMEMBRANCER; OR, COMPLETE POCKET ANATOMIST. Fifth Edition, carefully Revised. 32mo. cloth, 3s. 6d.

DR. ANDREW ANDERSON, M.D.

TEN LECTURES INTRODUCTORY TO THE STUDY OF FEVER. Post 8vo. cloth, 5s.

DR. MOCALL ANDERSON, M.D. PARASITIC AFFECTIONS OF THE SKIN. With Engravings. 8vo. cloth, 5s.

→-Ð{-

DR. ARLIDGE.

ON THE STATE OF LUNACY AND THE LEGAL PROVISION FOR THE INSANE; with Observations on the Construction and Organisation of Asylums. 8vo. cloth, 7s.

DR. ALEXANDER ARMSTRONG, R.N.

OBSERVATIONS ON NAVAL HYGIENE AND SCURVY.

More particularly as the latter appeared during a Polar Voyage. 8vo. cloth, 5s.

MR. T. J. ASHTON.

ON THE DISEASES, INJURIES, AND MALFORMATIONS OF THE RECTUM AND ANUS. Third Edition. 8vo. cloth, 8s.

MR. THOS. J. AUSTIN, M.R.C.S. ENG.

A PRACTICAL ACCOUNT OF GENERAL PARALYSIS: Its Mental and Physical Symptoms, Statistics, Causes, Seat, and Treatment. 8vo. cloth, 6s.

MR. THOMAS BALLARD, M.R.C.S.

A NEW AND RATIONAL EXPLANATION OF THE DIS-EASES PECULIAR TO INFANTS AND MOTHERS; with obvious Suggestions for their Prevention and Cure. Post 8vo. cloth, 4s. 6d.

DR. BARCLAY.

A MANUAL OF MEDICAL DIAGNOSIS. Second Edition.

DR. T. HERBERT BARKER.

ON THE HYGIENIC MANAGEMENT OF INFANTS AND CHILDREN. 8vo. cloth, 5s.

DR. W. G. BARKER.

ON THE CLIMATE OF WORTHING: its Remedial Influence in Disease, especially of the Lungs. Crown 8vo. cloth, 3s.

DR. BARLOW.

A MANUAL OF THE PRACTICE OF MEDICINE. Second Edition. Fcap. 8vo. cloth, 12s. 6d.

DR. BARNES.

THE PHYSIOLOGY AND TREATMENT OF PLACENTA PRÆVIA; being the Lettsomian Lectures on Midwifery for 1857. Post 8vo. cloth, 6s.

MR. BARWELL, F.R.C.S.

A TREATISE ON DISEASES OF THE JOINTS. With Engravings. 8vo. cloth, 12s.

DR. BASCOME.

A HISTORY OF EPIDEMIC PESTILENCES, FROM THE EARLIEST AGES. 8vo. cloth, 8s.

ON DROPSY, CONNECTED WITH DISEASE OF THE KIDNEYS (MORBUS BRIGHTII), and on some other Diseases of those Organs, associated with Albuminous and Purulent Urine. Illustrated by numerous Drawings from the Microscope. 8vo. cloth, 9s.

MR. H. F. BAXTER, M.R.C.S.L.

ON ORGANIC POLARITY; showing a Connexion to exist between Organic Forces and Ordinary Polar Forces. Crown 8vo. cloth, 5s.

st between

MR. BATEMAN.

MAGNACOPIA: A Practical Library of Profitable Knowledge, communicating the general Minutiæ of Chemical and Pharmaceutic Routine, together with the generality of Secret Forms of Preparations; including Concentrated Solutions of Camphor and Copaiba in Water, Mineral Succedaneum, Marmoratum, Silicia, Terro-Metallicum, Pharmaceutic Condensions, Prismatic Crystallization, Crystallized Aromatic Salt of Vinegar, Spa Waters; newly-invented Writing Fluids; Etching on Steel or Iron; with an extensive Variety of et cætera. Third Edition. 18mo. 6s.

MR. LIONEL J. BEALE, M.R.C.S.

THE LAWS OF HEALTH IN THEIR RELATIONS TO MIND AND BODY. A Series of Letters from an Old Practitioner to a Patient. Post 8vocloth, 7s. 6d.

CONNECTION WITH AND DISEASE, IN HEALTH GENERAL PRINCIPLES OF HYGIENE. Fcap. 8vo., 2s. 6d.

DR. BEALE, F.R.S.

ON URINE, URINARY DEPOSITS, AND CALCULI: their Microscopical and Chemical Examination; the Anatomy of the Kidney, and General Remarks on the Treatment of certain Urinary Diseases. Numerous Engravings. Post 8vo. cloth, 8s. 6d.

HOW TO WORK WITH THE MICROSCOPE. Illustrated Edition.

Crown 8vo. cloth, 5s. 6d.

THE MICROSCOPE, IN ITS APPLICATION TO PRACTICAL MEDICINE. With a Coloured Plate, and 270 Woodcuts. Second Edition. 8vocloth, 14s.

THE LIVER. Illustrated with 66 Pho-ON THE ANATOMY tographs of the Author's Drawings. 8vo. cloth, 6s. 6d.

ILLUSTRATIONS OF THE SALTS OF URINE, URINARY DEPOSITS, and CALCULI. 37 Plates, containing upwards of 170 Figures copied from Nature, with descriptive Letterpress. 8vo. cloth, 9s. 6d.

VI. THE HUMAN BODY. ON THE SIMPLE TISSUES OF Plates, 8vo. cloth, 7s. 6d.

MR. BEASLEY.

THE BOOK OF PRESCRIPTIONS; containing 3000 Prescriptions. Collected from the Practice of the most eminent Physicians and Surgeons, English and Foreign. Second Edition. 18mo. cloth, 6s.

THE DRUGGIST'S GENERAL RECEIPT-BOOK: comprising a copious Veterinary Formulary and Table of Veterinary Materia Medica; Patent and Proprietary Medicines, Druggists' Nostrums, &c.; Perfumery, Skin Cosmetics, Hair Cosmetics, and Teeth Cosmetics; Beverages, Dietetic Articles, and Condiments; Trade Chemicals, Miscellaneous Preparations and Compounds used in the Arts, &c.; with useful Memoranda and Tables. Fifth Edition. 18mo. cloth, 6s.

THE FORMULARY AND SYNOPSIS POCKET BRITISH AND FOREIGN PHARMACOPŒIAS; comprising standard and approved Formulæ for the Preparations and Compounds employed in Medical Practice. Seventh Edition, corrected and enlarged. 18mo. cloth, 6s.

DR. HENRY BENNET.

NUTRITION IN HEALTH AND DISEASE. Post 8vo. cloth, 5s.

A PRACTICAL TREATISE ON INFLAMMATION AND OTHER DISEASES OF THE UTERUS. Fourth Edition, revised, with Additions. 8vo. cloth, 16s.

A REVIEW OF THE PRESENT STATE OF UTERINE PATHOLOGY. 8vo. cloth, 4s.

MENTONE AND THE RIVIERA AS A WINTER CLIMATE.
Post 8vo. cloth, 3s. 6d.

DR. O'B. BELLINGHAM.

ON ANEURISM, AND ITS TREATMENT BY COMPRESSION.

PROFESSOR BENTLEY, F.L.S.

A MANUAL OF BOTANY. With nearly 1,200 Engravings on Wood. Fcap. 8vo. cloth, 12s. 6d.

MR. HENRY HEATHER BIGG.

THE MECHANICAL APPLIANCES NECESSARY FOR THE TREATMENT OF DEFORMITIES. Post 8vo. cloth, 4s.

ARTIFICIAL LIMBS; THEIR CONSTRUCTION AND APPLICATION. With Engravings on Wood. 8vo. cloth, 3s.

DR. BILLING, F.R.S.

ON DISEASES OF THE LUNGS AND HEART. 8vo. cloth, 6s.

DR. S. B. BIRCH, M.D.

CONSTIPATED BOWELS: the Various Causes and the Rational Means of Cure. Post 8vo. cloth, 2s. 6d.

DR. GOLDING BIRD, F.R.S.

URINARY DEPOSITS; THEIR DIAGNOSIS, PATHOLOGY, AND THERAPEUTICAL INDICATIONS. With Engravings on Wood. Fifth Edition. Post 8vo. cloth, 10s. 6d. 11.

ELEMENTS OF NATURAL PHILOSOPHY; being an Experimental Introduction to the Study of the Physical Sciences. Illustrated with numerous Engravings on Wood. Fifth Edition. By Golding Bird, M.D., F.R.S., and Charles Brooke, M.B. Cantab., F.R.S. Fcap. 8vo. cloth, 12s. 6d.

MR. BISHOP, F.R.S.

ON DEFORMITIES OF THE HUMAN BODY, their Pathology and Treatment. With Engravings on Wood. 8vo. cloth, 10s.

ON ARTICULATE SOUNDS, AND ON THE CAUSES AND CURE OF IMPEDIMENTS OF SPEECH. 8vo. cloth, 4s.

LETTSOMIAN LECTURES ON THE PHYSICAL CONSTI-TUTION, DISEASES AND FRACTURES OF BONES. Post 8vo., 2s. 6d.

10+一次数分号

MR. P. HINCKES BIRD, F.R.C.S.

PRACTICAL TREATISE ON THE DISEASES OF CHILDREN AND INFANTS AT THE BREAST. Translated from the French of M. BOUCHUT, with Notes and Additions. 8vo. cloth. 20s.

MR. ROBERT HOWARTH BLAKE, M.R.C.S.L. A PRACTICAL TREATISE ON DISEASES OF THE SKIN IN CHILDREN. From the French of CAILLAULT. With Notes. Post 8vo. cloth, 8s. 6d.

DR. BLAKISTON, F.R.S.

PRACTICAL OBSERVATIONS ON CERTAIN DISEASES OF THE CHEST; and on the Principles of Auscultation. 8vo. cloth, 12s.

DR. JOHN W. F. BLUNDELL.

MEDICINA MECHANICA; or, the Theory and Practice of Active and Passive Exercises and Manipulations in the Cure of Chronic Disease. Post 8vo. cloth, 6s.

MR. JOHN E. BOWMAN.

PRACTICAL CHEMISTRY, including Analysis. With numerous Illustrations on Wood. Fourth Edition. Foolscap 8vo. cloth, 6s. 6d.

II.

MEDICAL CHEMISTRY; with Illustrations on Wood. Third Edition. Fcap. 8vo. cloth, 6s. 6d.

DR. JAMES BRIGHT.

ON DISEASES OF THE HEART, LUNGS, & AIR PASSAGES; with a Review of the several Climates recommended in these Affections. Third Edition. Post 8vo. cloth, 9s.

DR. BRINTON.

THE DISEASES OF THE STOMACH, with an Introduction on its Anatomy and Physiology; being Lectures delivered at St. Thomas's Hospital. Post 8vo. cloth, 10s. 6d.

AND THE SYMPTOMS, PATHOLOGY, TREATMENT ULCER OF THE STOMACH. Post 8vo. cloth, 5s.

MR. BERNARD E. BRODHURST, F.R.C.S.

ON LATERAL CURVATURE OF THE SPINE: its Pathology and Treatment. Post 8vo. cloth, with Plates, 3s.

ON THE NATURE AND TREATMENT OF CLUBFOOT AND ANALOGOUS DISTORTIONS involving the TIBIO-TARSAL ARTICULATION. With Engravings on Wood. 8vo. cloth, 4s. 6d.

PRACTICAL OBSERVATIONS ON THE DISEASES OF THE JOINTS INVOLVING ANCHYLOSIS, and on the TREATMENT for the RESTORATION of MOTION. Third Edition, much enlarged, 8vo. cloth, 4s. 6d.

MR. THOMAS BRYANT, F.R.C.S.

THE DISEASES AND INJURIES OF THE CLINICAL AND PATHOLOGICAL OBSERVATIONS. Post 8vo. cloth, 7s. 6d.

DR. BRYCE.

ENGLAND AND FRANCE BEFORE SEBASTOPOL, looked at from a Medical Point of View. 8vo. cloth, 6s.

DR. BUDD, F.R.S.

ON DISEASES OF THE LIVER.

Illustrated with Coloured Plates and Engravings on Wood. Third Edition. 8vo. cloth, 16s.

ON THE ORGANIC DISEASES AND FUNCTIONAL DIS-ORDERS OF THE STOMACH. 8vo. cloth, 9s.

DR. JOHN CHARLES BUCKNILL, & DR. DANIEL H. TUKE.

A MANUAL OF PSYCHOLOGICAL MEDICINE: containing the History, Nosology, Description, Statistics, Diagnosis, Pathology, and Treatment of Insanity. Second Edition. 8vo. cloth, 15s.

DR. BURGESS.

THE MEDICAL AND LEGAL RELATIONS OF MADNESS; showing a Cellular Theory of Mind, and of Nerve Force, and also of Vegetative Vital Force. 8vo. cloth, 7s.

DR. BURNETT.

THE PHILOSOPHY OF SPIRITS IN RELATION TO MATTER. 8vo. cloth, 9s.

INSANITY TESTED BY SCIENCE. 8vo. cloth, 5s.

DR. JOHN M. CAMPLIN, F.L.S.

ON DIABETES, AND ITS SUCCESSFUL TREATMENT. Second Edition. Fcap. 8vo. cloth, 3s. 6d.

MR. ROBERT B. CARTER, M.R.C.S.

ON THE INFLUENCE OF EDUCATION AND TRAINING IN PREVENTING DISEASES OF THE NERVOUS SYSTEM. Fcap. 8vo., 6s.

THE PATHOLOGY AND TREATMENT OF HYSTERIA. Post 8vo. cloth, 4s. 6d.

DR. CARPENTER, F.R.S.

PRINCIPLES OF HUMAN PHYSIOLOGY. With numerous Illustrations on Steel and Wood. Fifth Edition. 8vo. cloth, 26s.

PRINCIPLES OF COMPARATIVE PHYSIOLOGY. Illustrated with 300 Engravings on Wood. Fourth Edition. 8vo. cloth, 24s.

A MANUAL OF PHYSIOLOGY. With numerous Illustrations on Steel and Wood. Third Edition. Fcap. 8vo. cloth, 12s. 6d.

THE MICROSCOPE AND ITS REVELATIONS. With numerous Engravings on Wood. Second Edition. Fcap. 8vo. cloth, 12s. 6d.

DR. CHAMBERS.

DIGESTION AND ITS DERANGEMENTS. Post 8vo. cloth, 10s. 6d.

10-一分别人

DR. CHANCE, M.B.

VIRCHOW'S CELLULAR PATHOLOGY, AS BASED UPON PHYSIOLOGICAL AND PATHOLOGICAL HISTOLOGY. With 144 Engravings on Wood. 8vo. cloth, 16s.

MR. H. T. CHAPMAN, F.R.C.S.

THE TREATMENT OF OBSTINATE ULCERS AND CUTA-NEOUS ERUPTIONS OF THE LEG WITHOUT CONFINEMENT. Third Edition. Post 8vo. cloth, 3s. 6d.

VARICOSE VEINS: their Nature, Consequences, and Treatment, Palliative and Curative. Post 8vo. cloth, 3s. 6d.

MR. PYE HENRY CHAVASSE, F.R.C.S.

ADVICE TO A MOTHER ON THE MANAGEMENT OF HER OFFSPRING. Sixth Edition. Foolscap 8vo., 2s. 6d.

ADVICE TO A WIFE ON THE MANAGEMENT OF HER OWN HEALTH. With an Introductory Chapter, especially addressed to a Young Wife. Fourth Edition. Fcap. 8vo., 2s. 6d.

DR. G. C. CHILD.

ON INDIGESTION, AND CERTAIN BILIOUS DISORDERS OFTEN CONJOINED WITH IT. Second Edition. 8vo. cloth, 6s.

MR. J. PATERSON CLARK, M.A.

THE ODONTALGIST; OR, HOW TO PRESERVE THE TEETH, CURE TOOTHACHE, AND REGULATE DENTITION FROM INFANCY TO AGE. With plates. Post 8vo. cloth, 5s.

MR. JOHN CLAY, M.R.C.S.

KIWISCH ON DISEASES OF THE OVARIES: Translated, by permission, from the last German Edition of his Clinical Lectures on the Special Pathology and Treatment of the Diseases of Women. With Notes, and an Appendix on the Operation of Ovariotomy. Royal 12mo. cloth, 16s.

DR. CONOLLY.

THE CONSTRUCTION AND GOVERNMENT OF LUNATIC ASYLUMS AND HOSPITALS FOR THE INSANE. With Plans. Post 8vo. cloth, 6s.

MR. COOLEY.

COMPREHENSIVE SUPPLEMENT TO THE PHARMACOPŒIAS.

THE CYCLOPÆDIA OF PRACTICAL RECEIPTS, AND COLLATERAL INFORMATION IN THE ARTS, PROFESSIONS, MANUFACTURES, AND TRADES, INCLUDING MEDICINE, PHARMACY, AND DOMESTIC ECONOMY; designed as a Compendious Book of Reference for the Manufacturer, Tradesman, Amateur, and Heads of Families. Third and greatly enlarged Edition, 8vo. cloth, 26s.

ON THE STRUCTURE AND DISEASES OF THE TESTIS.
With 24 Plates. Second Edition. Royal 4to., 20s.

1 11 2 100, 200.

MR. W. WHITE COOPER.

ON WOUNDS AND INJURIES OF THE EYE. Illustrated by 17 Coloured Figures and 41 Woodcuts. 8vo. cloth, 12s.

ON NEAR SIGHT, AGED SIGHT, IMPAIRED VISION, AND THE MEANS OF ASSISTING SIGHT. With 31 Illustrations on Wood. Second Edition. Fcap. 8vo. cloth, 7s. 6d.

MR. COOPER.

A DICTIONARY OF PRACTICAL SURGERY AND ENCYCLO-PÆDIA OF SURGICAL SCIENCE. New Edition, brought down to the present time. By SAMUEL A. LANE, F.R.C.S., assisted by various eminent Surgeons. Vol. I., 8vo. cloth, £1. 5s.

MR. HOLMES COOTE, F.R.C.S.

A REPORT ON SOME IMPORTANT POINTS IN THE TREATMENT OF SYPHILIS. 8vo. cloth, 5s.

DR. COTTON.

ON CONSUMPTION: Its Nature, Symptoms, and Treatment. To which Essay was awarded the Fothergillian Gold Medal of the Medical Society of London. Second Edition. 8vo. cloth, 8s.

PHTHISIS AND THE STETHOSCOPE; OR, THE PHYSICAL SIGNS OF CONSUMPTION. Second Edition. Foolscap 8vo. cloth, 3s.

MR. COULSON.

I.

ON DISEASES OF THE BLADDER AND PROSTATE GLAND. The Fifth Edition, revised and enlarged. 8vo. cloth, 10s. 6d.

ON LITHOTRITY AND LITHOTOMY; with Engravings on Wood.

ON THE INFLUENCE OF VARIATIONS OF ELECTRIC TENSION AS THE REMOTE CAUSE OF EPIDEMIC AND OTHER DISEASES. 8vo. cloth, 10s.

MR. CURLING, F.R.S.

OBSERVATIONS ON DISEASES OF THE RECTUM. Second

A PRACTICAL TREATISE ON DISEASES OF THE TESTIS, SPERMATIC CORD, AND SCROTUM. Second Edition, with Additions. 8vo. cloth, 14s.

METEOROLOGICAL AND MEDICAL OBSERVATIONS ON THE CLIMATE OF EGYPT, with Practical Hints for Invalid Travellers. Post 8vo. cloth, 4s.

MR. JOHN DALRYMPLE, F.R.S., F.R.C.S.

PATHOLOGY OF THE HUMAN EYE. Complete in Nine Fasciculi: imperial 4to., 20s. each; half-bound morocco, gilt tops, 9l. 15s.

10-->(例

10+0

MR. CHURCHILL'S PUBLICATIONS.

DR. DAVEY.

THE GANGLIONIC NERVOUS SYSTEM: its Structure, Functions, and Diseases. 8vo. cloth, 9s.

ON THE NATURE AND PROXIMATE CAUSE OF IN-SANITY. Post 8vo. cloth, 3s.

DR. HERBERT DAVIES.

ON THE PHYSICAL DIAGNOSIS OF DISEASES OF THE LUNGS AND HEART. Second Edition. Post 8vo. cloth, 8s.

DR. HALL DAVIS.

ILLUSTRATIONS OF DIFFICULT PARTURITION. Post 8vo.

MR. DIXON.

A GUIDE TO THE PRACTICAL STUDY OF DISEASES OF THE EYE. Second Edition. Post 8vo. cloth, 9s.

DR. DOBELL.

DEMONSTRATIONS OF DISEASES IN THE CHEST, AND THEIR PHYSICAL DIAGNOSIS. With Coloured Plates. 8vo. cloth, 12s. 6d.

LECTURES ON THE GERMS AND VESTIGES OF DISEASE, and on the Prevention of the Invasion and Fatality of Disease by Periodical Examinations. 8vo. cloth, 6s. 6d.

DR. TOOGOOD DOWNING.

NEURALGIA: its various Forms, Pathology, and Treatment. THE JACKSONIAN PRIZE ESSAY FOR 1850. 8vo. cloth, 10s. 6d.

DR. DRUITT, F.R.C.S.

THE SURGEON'S VADE-MECUM; with numerous Engravings on Wood. Eighth Edition. Foolscap 8vo. cloth, 12s. 6d.

MR. DUNN, F.R.C.S.

AN ESSAY ON PHYSIOLOGICAL PSYCHOLOGY. 8vo. cloth, 4s.

SIR JAMES EYRE, M.D.

THE STOMACH AND ITS DIFFICULTIES. Fifth Edition.

PRACTICAL REMARKS ON SOME EXHAUSTING DIS-EASES. Second Edition. Post 8vo. cloth, 4s. 6d.

DR. FENWICK.

ON SCROFULA AND CONSUMPTION. Clergyman's Sore Throat, Catarrh, Croup, Bronchitis, Asthma. Fcap. 8vo., 2s. 6d.

MR. FERGUSSON, F.R.S.

A SYSTEM OF PRACTICAL SURGERY; with numerous Illustrations on Wood. Fourth Edition. Fcap. 8vo. cloth, 12s. 6d.

MR. FLOWER, F.R.C.S.

DIAGRAMS OF THE NERVES OF THE HUMAN BODY, exhibiting their Origin, Divisions, and Connexions, with their Distribution to the various Regions of the Cutaneous Surface, and to all the Muscles. Folio, containing Six Plates, 14s.

NATURE AND ART IN THE CURE OF DISEASE. Second Edition. Post 8vo. cloth, 6s.

MR. FOWNES, PH.D., F.R.S.

A MANUAL OF CHEMISTRY; with numerous Illustrations on Wood. Eighth Edition. Fcap. 8vo. cloth, 12s. 6d.

Edited by H. Bence Jones, M.D., F.R.S., and A. W. Hofmann, Ph.D., F.R.S.

CHEMISTRY, AS EXEMPLIFYING THE WISDOM AND BENEFICENCE OF GOD. Second Edition. Fcap. 8vo. cloth, 4s. 6d.

INTRODUCTION TO QUALITATIVE ANALYSIS. Post 8vo. cloth, 2s.

DR. D. J. T. FRANCIS.

CHANGE OF CLIMATE; considered as a Remedy in Dyspeptic, Pulmonary, and other Chronic Affections; with an Account of the most Eligible Places of Residence for Invalids in Spain, Portugal, Algeria, &c., at different Seasons of the Year; and an Appendix on the Mineral Springs of the Pyrenees, Vichy, and Aix les Bains. Post 8vo. cloth, 8s. 6d.

C. REMIGIUS FRESENIUS.

ELEMENTARY INSTRUCTION IN CHEMICAL ANALYSIS, AS PRACTISED IN THE LABORATORY OF GIESSEN. Edited by LLOYD BULLOCK, F.C.S.

QUALITATIVE. Fifth Edition. 8vo. cloth, 9s. QUANTITATIVE. Third Edition. 8vo. cloth, 16s.

DR. FULLER.

ON RHEUMATISM, RHEUMATIC GOUT, AND SCIATICA: their Pathology, Symptoms, and Treatment. Third Edition. 8vo. cloth, 12s. 6d.

DR. GAIRDNER.

ON GOUT; its History, its Causes, and its Cure. Fourth Edition. Post 8vo. cloth, 8s. 6d.

MR. GALLOWAY.

THE FIRST STEP IN CHEMISTRY. Third Edition. Fcap. 8vo. cloth, 5s.

A MANUAL OF QUALITATIVE ANALYSIS. Third Edition. Post 8vo. cloth, 5s.

MR. F. J. GANT.

THE IRRITABLE BLADDER: its Causes and Curative Treatment. Post 8vo. cloth, 4s 6d.

10+一分数分割

DR. GIBB, M.R.C.P.

THE THROAT, EPIGLOTTIS, AND DISEASES OF WINDPIPE. Post 8vo. cloth, 5s.

MRS. GODFREY.

ON THE NATURE, PREVENTION, TREATMENT, AND CURE OF SPINAL CURVATURES and DEFORMITIES of the CHEST and LIMBS, without ARTIFICIAL SUPPORTS or any MECHANICAL APPLIANCES. Third Edition, Revised and Enlarged. 8vo. cloth, 5s.

DR. GRANVILLE, F.R.S. I.

MINERAL SPRINGS OF VICHY: their Efficacy in the AME BY Treatment of Gout, Indigestion, Gravel, &c. 8vo. cloth, 5s,

ON SUDDEN DEATH. Post 8vo., 2s. 6d.

MR. GRAY, M.R.C.S.

PRESERVATION OF THE TEETH indispensable to Comfort and Appearance, Health, and Longevity. 18mo. cloth, 3s.

MR. GRIFFITHS.

THE FOUR SEASONS—Spring, Summer, 0 ECHEMISTRY Autumn, Winter. Illustrated with Engravings on Wood. Second Edition. Foolscap 8vo. cloth, 7s. 6d.

DR. GULLY.

THE SIMPLE TREATMENT OF DISEASE; deduced from the Methods of Expectancy and Revulsion. 18mo. cloth, 4s.

DR. GUY.

HOOPER'S PHYSICIAN'S VADE-MECUM; OR, MANUAL OF THE PRINCIPLES AND PRACTICE OF PHYSIC. New Edition, considerably enlarged, and rewritten. Foolscap 8vo. cloth, 12s. 6d.

GUY'S HOSPITAL REPORTS. Third Series. Vols. I. to VII., 8vo., 7s. 6d. each.

DR. HABERSHON, F.R.C.P.

OBSERVATIONS ON DISEASES OF THE ALIMENTARY CANAL, ŒSOPHAGUS, STOMACH, CÆCUM, and INTESTINES. 8vo. cloth, 10s. 6d.

ON THE INJURIOUS EFFECTS OF MERCURY IN THE TREATMENT OF DISEASE. Post 8vo. cloth, 3s. 6d.

DR. MARSHALL HALL, F.R.S.

PRONE AND POSTURAL RESPIRATION IN DROWNING AND OTHER FORMS OF APNŒA OR SUSPENDED RESPIRATION. Post 8vo. cloth. 5s.

11. PRACTICAL OBSERVATIONS AND SUGGESTIONS IN MEDI-CINE. Second Zeries. Post 8vo. cloth, 8s. 6d.

DR. C. RADCLYFFE HALL.

TORQUAY IN ITS MEDICAL ASPECT AS A RESORT FOR PULMONARY INVALIDS. Post 8vo. cloth, 5s.

MR. HARDWICH.

A MANUAL OF PHOTOGRAPHIC CHEMISTRY. Sixth Edition. Foolscap 8vo. cloth, 7s. 6d.

MR. HARE, F.R.C.S.

PRACTICAL OBSERVATIONS ON THE PREVENTION, CAUSES, AND TREATMENT OF CURVATURES OF THE SPINE; with Engravings. Third Edition. 8vo. cloth, 6s.

DR. JAMES B. HARRISON.

ON THE CONTAMINATION OF WATER BY THE POISON OF LEAD, and its Effects on the Human Body. Foolscap 8vo. cloth, 3s. 6d.

DR. HARTWIG.

ON SEA BATHING AND SEA AIR. Second Edition. Fcap. 8vo., 2s. 6d.

ON THE PHYSICAL EDUCATION OF CHILDREN. Fcap.

DR. A. H. HASSALL.

MICROSCOPIC ANATOMY OF THE HUMAN BODY,

IN HEALTH AND DISEASE. Illustrated with Several Hundred Drawings in Colour. Two vols. 8vo. cloth, £1. 10s.

II.

THE URINE, IN HEALTH AND DISEASE; or, a Simple Explanation of the Physical Properties, Composition, and Uses of the Urine, of the Functions of the Kidneys, and of the Treatment of Urinary Disorders. With Twenty-four Engravings. Post 8vo. cloth, 5s.

MR. ALFRED HAVILAND, M.R.C.S.

CLIMATE, WEATHER, AND DISEASE; being a Sketch of the Opinions of the most celebrated Ancient and Modern Writers with regard to the Influence of Climate and Weather in producing Disease. With Four coloured Engravings. 8vo. cloth, 7s.

MR. WILLIAM HAYCOCK, M.R.C.V.S.

A TREATISE ON THE PRINCIPLES AND PRACTICE OF VETERINARY MEDICINE AND SURGERY. 8vo. boards, 6s. 6d.

DR. HEADLAND.

ON THE ACTION OF MEDICINES IN THE SYSTEM.

Being the Prize Essay to which the Medical Society of London awarded the Fothergillian Gold Medal for 1852. Third Edition. 8vo. cloth, 12s. 6d.

10->8>*

+⊕{-

DR. HEALE.

A TREATISE ON VITAL CAUSES. 8vo. cloth, 9s.

MR. CHRISTOPHER HEATH, F.R.C.S.

A MANUAL OF MINOR SURGERY AND BANDAGING, FOR THE USE OF HOUSE-SURGEONS, DRESSERS, AND JUNIOR PRACTITIONERS. With Illustrations. Feap. 8vo. cloth, 5s.

MR. HIGGINBOTTOM, F.R.S., F.R.C.S.E.

I.

AN ESSAY ON THE USE OF THE NITRATE OF SILVER IN THE CURE OF INFLAMMATION, WOUNDS, AND ULCERS. Second Edition. Price 5s.

II

ADDITIONAL OBSERVATIONS ON THE NITRATE OF SIL-VER; with full Directions for its Use as a Therapeutic Agent. 8vo., 2s. 6d.

MR. JOHN HILTON, F.R.S.

ON THE DEVELOPMENT AND DESIGN OF CERTAIN PORTIONS OF THE CRANIUM. Illustrated with Plates in Lithography. 8vo. cloth, 6s.

DR. HINDS.

THE HARMONIES OF PHYSICAL SCIENCE IN RELATION TO THE HIGHER SENTIMENTS; with Observations on Medical Studies, and on the Moral and Scientific Relations of Medical Life. Post 8vo., cloth, 4s.

DR. DECIMUS HODGSON.

THE PROSTATE GLAND, AND ITS ENLARGEMENT IN OLD AGE. With 12 Plates. Royal 8vo., cloth, 6s.

MR. JABEZ HOGG.

THE OPHTHALMOSCOPE: an Essay on its value in the Exploration of Internal Eye Diseases. Second Edition. Cloth, 3s. 6d.

MR. LUTHER HOLDEN, FR.C.S.

I.

HUMAN OSTEOLOGY: with Plates, showing the Attachments of the Muscles. Third Edition. 8vo. cloth, 16s.

II.

A MANUAL OF THE DISSECTION OF THE HUMAN BODY. With Engravings on Wood. Second Edition. 8vo. cloth, 16s.

MR. C. HOLTHOUSE.

I.

ON SQUINTING, PARALYTIC AFFECTIONS OF THE EYE, and CERTAIN FORMS OF IMPAIRED VISION. Fcap. 8vo. cloth, 4s. 6d.

LECTURES ON STRABISMUS, delivered at the Westminster Hospital.

FARCO + - 01-

キャーなおびではかっま

DR. W. CHARLES HOOD.

SUGGESTIONS FOR THE FUTURE PROVISION OF CRIMI-NAL LUNATICS. 8vo. cloth, 5s. 6d.

MR. P. HOOD.

THE SUCCESSFUL TREATMENT OF SCARLET FEVER; also, OBSERVATIONS ON THE PATHOLOGY AND TREATMENT OF CROWING INSPIRATIONS OF INFANTS. Post 8vo. cloth, 5s.

DR. HOOPER.

THE MEDICAL DICTIONARY; containing an Explanation of the Terms used in Medicine and the Collateral Sciences. Eighth Edition. Edited by Klein Grant, M.D. 8vo. cloth, 30s.

MR. JOHN HORSLEY.

A CATECHISM OF CHEMICAL PHILOSOPHY; being a Familiar Exposition of the Principles of Chemistry and Physics. With Engravings on Wood. Designed for the Use of Schools and Private Teachers. Post 8vo. cloth, 6s. 6d.

DR. HUFELAND.

THE ART OF PROLONGING LIFE. Second Edition. Edited by Erasmus Wilson, F.R.S. Foolscap 8vo., 2s. 6d.

BLOOD DISEASE. Crown 8vo. cloth, 4s. 6d.

MR. W. CURTIS HUGMAN, F.R.C.S.

ON HIP-JOINT DISEASE; with reference especially to Treatment by Mechanical Means for the Relief of Contraction and Deformity of the Affected Limb. 8vo. cloth, 3s. 6d.

MR. HULKE, F.R.C.S.

A PRACTICAL TREATISE ON THE USE OF THE OPHTHALMOSCOPE. Being the Jacksonian Prize Essay for 1859. Royal 8vo. cloth, 8s.

DR. HENRY HUNT.

ON HEARTBURN AND INDIGESTION. 8vo. cloth, 5s.

DR. INMAN, M.R.C.P.

ON MYALGIA: ITS NATURE, CAUSES, AND TREATMENT; being a Treatise on Painful and other Affections of the Muscular System. Second Edition. 8vo. cloth, 9s.

FOUNDATION FOR A NEW THEORY AND PRACTICE OF MEDICINE. Second Edition. Crown 8vo. cloth, 10s.

DR. ARTHUR JACOB, F.R.C.S.

A TREATISE ON THE INFLAMMATIONS OF THE EYE-BALL. Foolscap 8vo. cloth, 5s.

DR. JAMES JAGO, A.B., CANTAB.; M.B., OXON.

OCULAR SPECTRES AND STRUCTURES AS MUTUAL EXPONENTS. Illustrated with Engravings on Wood. 8vo. cloth, 5s.

600co +05-

MR. J. H. JAMES, F.R.C.S.

PRACTICAL OBSERVATIONS ON THE OPERATIONS FOR STRANGULATED HERNIA. 8vo. cloth, 5s.

DR. PROSSER JAMES, M.D.

SORE-THROAT: ITS NATURE, VARIETIES, AND TREAT-MENT; including the Use of the LARYNGOSCOPE as an Aid to Diagnosis. Post 8vo. cloth, 4s. 6d.

DR. BENCE JONES, F.R.S.

I.

MULDER ON WINE. Foolscap 8vo. cloth, 6s.

II.

ON ANIMAL CHEMISTRY, in its relation to STOMACH and RENAL DISEASES. 8vo. cloth, 6s.

DR. HANDFIELD JONES, F.R.S., & DR. EDWARD H. SIEVEKING.

A MANUAL OF PATHOLOGICAL ANATOMY. Illustrated with numerous Engravings on Wood. Foolscap 8vo. cloth, 12s. 6d.

MR. WHARTON JONES, F.R.S.

I.

A MANUAL OF THE PRINCIPLES AND PRACTICE OF OPHTHALMIC MEDICINE AND SURGERY; illustrated with Engravings, plain and coloured. Second Edition. Foolscap 8vo. cloth, 12s. 6d.

II.

THE WISDOM AND BENEFICENCE OF THE ALMIGHTY, AS DISPLAYED IN THE SENSE OF VISION; being the Actonian Prize Essay for 1851. With Illustrations on Steel and Wood. Foolscap 8vo. cloth, 4s. 6d.

III.

DEFECTS OF SIGHT: their Nature, Causes, Prevention, and General Management. Fcap. 8vo. 2s. 6d.

IV.

A CATECHISM OF THE MEDICINE AND SURGERY OF THE EYE AND EAR. For the Clinical Use of Hospital Students. Fcap. 8vo. 2s. 6d.

v.

A CATECHISM OF THE PHYSIOLOGY AND PHILOSOPHY OF BODY, SENSE, AND MIND. For Use in Schools and Colleges. Fcap. 8vo., 2s. 6d.

MR. FURNEAUX JORDAN, M.R.C.S.

AN INTRODUCTION TO CLINICAL SURGERY; WITH A Method of Investigating and Reporting Surgical Cases. Fcap. 8vo. cloth, 5s.

MR. JUDD.

A PRACTICAL TREATISE ON URETHRITIS AND SYPHI-LIS: including Observations on the Power of the Menstruous Fluid, and of the Discharge from Leucorrhea and Sores to produce Urethritis: with a variety of Examples, Experiments, Remedies, and Cures. 8vo. cloth, £1.5s.

1.

10+

DR. LAENNEC.

A MANUAL OF AUSCULTATION AND PERCUSSION. Translated and Edited by J. B. SHARPE, M.R.C.S. 3s.

DR. LANE, M.A.

HYDROPATHY; OR, HYGIENIC MEDICINE. An Explanatory Essay. Second Edition. Post 8vo. cloth, 5s.

MR. LAWRENCE, F.R.S.

A TREATISE ON RUPTURES. The Fifth Edition, considerably enlarged. 8vo. cloth, 16s.

DR. LEARED, M.R.C.P.

IMPERFECT DIGESTION: ITS CAUSES AND TREATMENT. Second Edition. Foolscap 8vo. cloth, 3s. 6d.

DR. EDWIN LEE.

I.

THE EFFECT OF CLIMATE ON TUBERCULOUS DISEASE, with Notices of the chief Foreign Places of Winter Resort. Small 8vo. cloth, 4s. 6d.

II.

THE WATERING PLACES OF ENGLAND, CONSIDERED with Reference to their Medical Topography. Fourth Edition. Foolscap 8vo. cloth, 7s. 6d.

III.

THE BATHS OF RHENISH GERMANY. Post 8vo. cloth, 3s.

IV.

HOMEOPATHY AND HYDROPATHY IMPARTIALLY AP-PRECIATED. With Notes illustrative of the Influence of the Mind over the Body. Fourth Edition. Post 8vo. cloth, 3s. 6d.

DR. ROBERT LEE, F.R.S.

I.

A TREATISE ON THE SPECULUM; with Three Hundred Cases. 8vo. cloth, 4s. 6d.

CLINICAL REPORTS OF OVARIAN AND UTERINE DIS-EASES, with Commentaries. Foolscap 8vo. cloth, 6s. 6d.

CLINICAL MIDWIFERY: comprising the Histories of 545 Cases of Difficult, Preternatural, and Complicated Labour, with Commentaries. Second Edition. Foolscap 8vo. cloth, 5s.

PRACTICAL OBSERVATIONS ON DISEASES OF THE UTERUS. With coloured Plates. Two Parts. Imperial 4to., 7s. 6d. each Part.

10->积)当

101.1

--DE

MR. LISTON, F.R.S.

PRACTICAL SURGERY. Fourth Edition. 8vo. cloth, 22s.

MR. H. W. LOBB, L.S.A., M.R.C.S.E.

ON SOME OF THE MORE OBSCURE FORMS OF NERVOUS AFFECTIONS, THEIR PATHOLOGY AND TREATMENT. With an Introduction on the Physiology of Digestion and Assimilation, and the Generation and Distribution of Nerve Force. Based upon Original Microscopical Observations. With Engravings. 8vo. cloth, 10s. 6d.

LONDON MEDICAL SOCIETY OF OBSERVATION.

WHAT TO OBSERVE AT THE BED-SIDE, AND AFTER DEATH. Published by Authority. Second Edition. Foolscap 8vo. cloth, 4s. 6d.

M. LUGOL.

ON SCROFULOUS DISEASES. Translated from the French, with Additions by W. H. RANKING, M.D., Physician to the Suffolk General Hospital. 8vo. cloth, 10s. 6d.

MR. M'CLELLAND, F.L.S., F.G.S.

SKETCH OF THE MEDICAL TOPOGRAPHY, OR CLIMATE AND SOILS, OF BENGAL AND THE N. W. PROVINCES. Post 8vo. cloth, 4s. 6d.

DR. GEORGE H. B. MACLEOD, F.R.C.S. (EDIN.)

NOTES ON THE SURGERY OF THE CRIMEAN WAR; with REMARKS on GUN-SHOT WOUNDS. 8vo. cloth, 10s. 6d.

MR. JOSEPH MACLISE, F.R.C.S.

I.

SURGICAL ANATOMY. A Series of Dissections, illustrating the Principal Regions of the Human Body.

The Second Edition, complete in XIII. Fasciculi, imperial folio, 5s. each. Bound in

cloth, £3, 12s.; or in morocco, £4. 4s.

ON DISLOCATIONS AND FRACTURES. This Work is Uniform with the Author's "Surgical Anatomy;" each Fasciculus contains Four beautifully executed Lithographic Drawings. Complete in Nine Fasciculi, imperial folio, 5s. each. Bound in cloth, £2. 10s.; or in morocco, £2. 17s.

DR. MONICOLL, M.R.C.P.

A HAND-BOOK FOR SOUTHPORT, MEDICAL & GENERAL; with Copious Notices of the Natural History of the District. Second Edition. Post 8vo. cloth, 3s. 6d.

DR. MARCET, F.R.S.

ON THE COMPOSITION OF FOOD, AND HOW IT IS ADULTERATED; with Practical Directions for its Analysis. 8vo. cloth, 6s. 6d.

ON CHRONIC ALCOHOLIC INTOXICATION, OR ALCO-HOLIC STIMULANTS IN CONNEXION WITH THE NERVOUS SYSTEM. Foolscap 8vo. cloth, 3s. 6d.

- K884-08-

DR. MARKHAM.

I.

DISEASES OF THE HEART: THEIR PATHOLOGY, DIAGNOSIS, AND TREATMENT. Second Edition. Post 8vo. cloth, 6s.

II.

SKODA ON AUSCULTATION AND PERCUSSION. Post 8vo.

SIR J. RANALD MARTIN, K.C.B., F.R.S.

INFLUENCE OF TROPICAL CLIMATES IN PRODUCING THE ACUTE ENDEMIC DISEASES OF EUROPEANS; including Practical Observations on their Chronic Sequelæ under the Influences of the Climate of Europe. Second Edition, much enlarged. 8vo. cloth, 20s.

DR. MASSY.

ON THE EXAMINATION OF RECRUITS; intended for the Use of Young Medical Officers on Entering the Army. 8vo. cloth, 5s.

MR. C. F. MAUNDER, F.R.C.S.

OPERATIVE SURGERY. With 158 Engravings. Post 8vo. 6s.

DR. MAYNE.

AN EXPOSITORY LEXICON OF THE TERMS, ANCIENT AND MODERN, IN MEDICAL AND GENERAL SCIENCE, including a complete MEDICAL AND MEDICO-LEGAL VOCABULARY, and presenting the correct Pronunciation, Derivation, Definition, and Explanation of the Names, Analogues, Synonymes, and Phrases (in English, Latin, Greek, French, and German,) employed in Science and connected with Medicine. Complete in 10 Parts, price 5s. each. The entire work, cloth, £2. 10s.

DR. MILLINGEN.

ON THE TREATMENT AND MANAGEMENT OF THE IN-SANE; with Considerations on Public and Private Lunatic Asylums. 18mo. cloth, 4s. 6d.

MR. JOHN L. MILTON, M.R.C.S.

PRACTICAL OBSERVATIONS ON A NEW WAY OF TREATING GONORRHEA. With some Remarks on the Cure of Inveterate Cases. 8vo. cloth, 5s.

DR. MONRO.

REFORM IN PRIVATE LUNATIC ASYLUMS. 8vo. cloth, 4s.

A MANUAL OF THE DISEASES OF INDIA. Fcap. 8vo. cloth, 5s.

DR. NOBLE.

THE HUMAN MIND IN ITS RELATIONS WITH THE BRAIN AND NERVOUS SYSTEM. Post 8vo. cloth, 4s. 6d.

10-->10

→**○**{

MR. NUNNELEY, F.R.C.S.E.

I.

ON THE OLGANS OF VISION: THEIR ANATOMY AND PHY-SIOLOGY. With Plates, 8vo. cloth, 15s:

IT.

A TREATISE ON THE NATURE, CAUSES, AND TREATMENT OF ERYSIPELAS. 8vo. cloth, 10s. 6d.

MR. PAGET.

A DESCRIPTIVE CATALOGUE OF THE ANATOMICAL MUSEUM OF ST. BARTHOLOMEW'S HOSPITAL. Vol. I. Morbid Anatomy. 8vo. cloth, 5s.

DITTO. Vol. II. Natural and Congenitally Malformed Structures, and Lists of the Models, Casts, Drawings, and Diagrams. 5s.

MR. LANGSTON PARKER.

THE MODERN TREATMENT OF SYPHILITIC DISEASES, both Primary and Secondary; comprising the Treatment of Constitutional and Confirmed Syphilis, by a safe and successful Method. Fourth Edition, 8vo. cloth, 10s.

DR. PARKES, F.R.C.P.

THE URINE: ITS COMPOSITION IN HEALTH AND DISEASE, AND UNDER THE ACTION OF REMEDIES. 8vo. cloth, 12s.

DR. PARKIN.

THE CAUSATION AND PREVENTION OF DISEASE.

8vo. cloth, 5s.

MR. JAMES PART, F.R.C.S.

THE MEDICAL AND SURGICAL POCKET CASE BOOK, for the Registration of important Cases in Private Practice, and to assist the Student of Hospital Practice. Second Edition. 3s. 6d.

DR. THOMAS B. PEACOCK, M.D.

ON THE INFLUENZA, OR EPIDEMIC CATARRHAL FEVER OF 1847-8. 8vo. cloth, 5s. 6d.

MR. OLIVER PEMBERTON, M.R.C.S.

OBSERVATIONS ON THE HISTORY, PATHOLOGY, AND TREATMENT OF CANCEROUS DISEASES. Part I. — Melanosis. With coloured Plates. Royal 8vo. cloth, 4s. 6d.

KBC-01-

DR. PEREIRA, F.R.S.

SELECTA E PRÆSCRIPTIS: with a Key, containing the Prescriptions in an Unabbreviated Form, and a Literal Translation. Thirteenth Edition. 24mo. cloth, 5s.

DR. PICKFORD.

HYGIENE; or, Health as Depending upon the Conditions of the Atmosphere, Food and Drinks, Motion and Rest, Sleep and Wakefulness, Secretions, Excretions, and Retentions, Mental Emotions, Clothing, Bathing, &c. Vol. I. 8vo. cloth, 9s.

MR. PIRRIE, F.R.S.E.

THE PRINCIPLES AND PRACTICE OF SURGERY. With numerous Engravings on Wood. Second Edition. 8vo. cloth, 24s.

PHARMACOPŒIA COLLEGII REGALIS MEDICORUM LON-DINENSIS. 8vo. cloth, 9s.; or 24mo. 5s.

IMPRIMATUR.

Hic liber, cui titulus, Pharmacopæia Collegii Regalis Medicorum Londinensis. Datum ex Ædibus Collegii in comitiis censoriis, Novembris Mensis 14^{to} 1850.

JOHANNES AYRTON PARIS. Præses.

PROFESSORS PLATTNER & MUSPRATT.

THE USE OF THE BLOWPIPE IN THE EXAMINATION OF MINERALS, ORES, AND OTHER METALLIC COMBINATIONS. Illustrated by numerous Engravings on Wood. Third Edition. 8vo. cloth, 10s. 6d.

DR. HENRY PRATT, M.D., M.R.C.P.

- THE GENEALOGY OF CREATION, newly Translated from the Unpointed Hebrew Text of the Book of Genesis, showing the General Scientific Accuracy of the Cosmogony of Moses and the Philosophy of Creation. 8vo. cloth, 14s.
- THE PRESCRIBER'S PHARMACOPŒIA; containing all the Medicines in the London Pharmacopæia, arranged in Classes according to their Action, with their Composition and Doses. By a Practising Physician. Fourth Edition. 32mo. cloth, 2s. 6d.; roan tuck (for the pocket), 3s. 6d.

DR. JOHN ROWLISON PRETTY.

AIDS DURING LABOUR, including the Administration of Chloroform, the Management of Placenta and Post-partum Hæmorrhage. Fcap. 8vo. cloth, 4s. 6d.

MR. LAKE PRICE.

PHOTOGRAPHIC MANIPULATION: Treating of the Practice of the Art, and its various appliances to Nature. With Fifty Engravings on Wood. Post 8vo. cloth, 6s. 6d.

DR. PRIESTLEY.

LECTURES ON THE DEVELOPMENT OF THE GRAVID UTERUS. 8vo. cloth, 5s. 6d.

10--01

10- -OE)

MR. P. C. PRICE, F.R.C.S.E.

SCROFULOUS DISEASES OF THE EXTERNAL LYMPHATIC GLANDS: their Nature, Variety, and Treatment; with Remarks on the Management of Scrofulous Ulcerations, Scars, and Cicatrices. Post 8vo. cloth, 3s. 6d.

SIR WM. PYM, K.C.H.

OBSERVATIONS UPON YELLOW FEVER, with a Review of "A Report upon the Diseases of the African Coast, by Sir WM. BURNETT and Dr. Bryson," proving its highly Contagious Powers. Post 8vo. 6s.

DR. RADCLIFFE, F.R.C.P. LOND.

ON EPILEPTIC AND OTHER CONVULSIVE AFFECTIONS OF THE NERVOUS SYSTEM. Third Edition. Post 8vo. cloth, 7s. 6d.

MR. RAINEY.

ON THE MODE OF FORMATION OF SHELLS OF ANIMALS, OF BONE, AND OF SEVERAL OTHER STRUCTURES, by a Process of Molecular Coalescence, Demonstrable in certain Artificially-formed Products. Fcap. 8vo. cloth, 4s. 6d.

DR. F. H. RAMSBOTHAM.

THE PRINCIPLES AND PRACTICE OF OBSTETRIC MEDI-CINE AND SURGERY. Illustrated with One Hundred and Twenty Plates on Steel and Wood; forming one thick handsome volume. Fourth Edition. 8vo. cloth, 22s.

DR. RAMSBOTHAM.

PRACTICAL OBSERVATIONS ON MIDWIFERY, with a Selection of Cases. Second Edition. 8vo. cloth, 12s.

DR. DU BOIS REYMOND.

ANIMAL ELECTRICITY; Edited by H. Bence Jones, M.D., F.R.S. With Fifty Engravings on Wood. Foolscap 8vo. cloth, 6s.

DR. REYNOLDS, M.D., LOND.

EPILEPSY: ITS SYMPTOMS, TREATMENT, AND RELATION TO OTHER CHRONIC CONVULSIVE DISEASES. 8vo. cloth, 10s.

II.

THE DIAGNOSIS OF DISEASES OF THE BRAIN, SPINAL CORD, AND THEIR APPENDAGES. 8vo. cloth, 8s.

DR. B. W. RICHARDSON.

ON THE CAUSE OF THE COAGULATION OF THE BLOOD. Being the ASTLEY COOPER PRIZE ESSAY for 1856. With a Practical Appendix. 8vo. cloth, 16s.

THE HYGIENIC TREATMENT OF PULMONARY CONSUMP-TION. 8vo. cloth, 5s. 6d.

THE ASCLEPIAD. Vol. I., Clinical Essays. 8vo. cloth, 6s. 6d.

る・心思公園か・四

MR. WILLIAM ROBERTS.

AN ESSAY ON WASTING PALSY; being a Systematic Treatise on the Disease hitherto described as ATROPHIE MUSCULAIRE PROGRESSIVE. With Four Plates. 8vo. cloth, 7s. 6d.

DR. W. H. ROBERTSON.

THE NATURE AND TREATMENT OF GOUT.

8vo. cloth, 10s. 6d.

A TREATISE ON DIET AND REGIMEN. Fourth Edition. 2 vols. post 8vo. cloth, 12s.

DR. ROUTH.

INFANT FEEDING, AND ITS INFLUENCES ON LIFE; Or, the Causes and Prevention of Infant Mortality. Fcap. 8vo. cloth, 5s.

DR. ROWE.

NERVOUS DISEASES, LIVER AND STOMACH COM-PLAINTS, LOW SPIRITS, INDIGESTION, GOUT, ASTHMA, AND DIS-ORDERS PRODUCED BY TROPICAL CLIMATES. With Cases. Sixteenth Edition. Fcap. 8vo. 2s. 6d.

DR. ROYLE, F.R.S.

A MANUAL OF MATERIA MEDICA AND THERAPEUTICS. With numerous Engrivings on Wood. Third Edition. Fcap. 8vo. cloth, 12s. 6d.

MR. RUMSEY, F.R.C.S.

ESSAYS ON STATE MEDICINE. 8vo. cloth, 10s. 6d.

DR. SALTER, F.R.S.

ON ASTHMA: its Pathology, Causes, Consequences, and Treatment. 8vo. cloth, 10s.

MR. SAVORY.

A COMPENDIUM OF DOMESTIC MEDICINE, AND COMPANION TO THE MEDICINE CHEST; comprising Plain Directions for the Employment of Medicines, with their Properties and Doses, and Brief Descriptions of the Symptoms and Treatment of Diseases, and of the Disorders incidental to Infants and Children, with a Selection of the most efficacious Prescriptions. Intended as a Source of Easy Reference for Clergymen, and for Families residing at a Distance from Professional Assistance. Fifth Edition. 12mo. cloth, 5s.

DR. SCHACHT.

THE MICROSCOPE, AND ITS APPLICATION TO VEGET BLE ANATOMY AND PHYSIOLOGY. Edited by Frederick Currey, M.A. Fcap. 8vo. cloth, 6s.

DR. SCORESBY-JACKSON, M.D., F.R.S.E.

MEDICAL CLIMATOLOGY; or, a Topographical and Meteorological Description of the Localities resorted to in Winter and Summer by Invalids of various classes both at Home and Abroad. With an Isothermal Chart. Post 8vo. cloth, 12s.

loth, 12s.

DR. SEMPLE.

ON COUGH: its Causes, Varieties, and Treatment. With some practical Remarks on the Use of the Stethoscope as an aid to Diagnosis. Post 8vo. cloth, 4s. 6d.

MR. SHAW, M.R.C.S.

THE MEDICAL REMEMBRANCER; OR, BOOK OF EMER-GENCIES: in which are concisely pointed out the Immediate Remedies to be adopted in the First Moments of Danger from Poisoning, Apoplexy, Burns, and other Accidents; with the Tests for the Principal Poisons, and other useful Information. Fourth Edition. Edited, with Additions, by Jonathan Hutchinson, M.R.C.S. 32mo. cloth, 2s. 6d.

DR. SIBSON, F.R.S.

MEDICAL ANATOMY. With coloured Plates. Imperial folio. Fasciculi I. to VI. 5s. each.

DR. E. H. SIEVEKING.

ON EPILEPSY AND EPILEPTIFORM SEIZURES: their Causes, Pathology, and Treatment. Second Edition. Post 8vo. cloth, 10s. 6d.

MR. SINCLAIR AND DR. JOHNSTON.

PRACTICAL MIDWIFERY: Comprising an Account of 13,748 Deliveries, which occurred in the Dublin Lying-in Hospital, during a period of Seven Years. 8vo. cloth, 15s.

MR. ALFRED SMEE, F.R.S.

GENERAL DEBILITY AND DEFECTIVE NUTRITION; their Causes, Consequences, and Treatment. Fcap. 8vo. cloth, 3s. 6d.

DR. SMELLIE.

OBSTETRIC PLATES: being a Selection from the more Important and Practical Illustrations contained in the Original Work. With Anatomical and Practical Directions. 8vo. cloth, 5s.

MR. HENRY SMITH, F.R.C.S.

ON STRICTURE OF THE URETHRA. 8vo. cloth, 7s. 6d.

HÆMORRHOIDS AND PROLAPSUS OF THE RECTUM:

Their Treatment by the Application of Nitric Acid. Second Edition. Fcap. 8vo. cloth, 3s.

DR. W. TYLER SMITH.

A MANUAL OF OBSTETRICS, THEORETICAL AND PRAC-TICAL. Illustrated with 186 Engravings. Fcap. 8vo. cloth, 12s. 6d.

THE PATHOLOGY AND TREATMENT OF LEUCORRHŒA. With Engravings on Wood. 8vo. cloth, 7s.

はする多くなると

~~~~

DR. SNOW.

ON CHLOROFORM AND OTHER ANÆSTHETICS: THEIR ACTION AND ADMINISTRATION. Edited, with a Memoir of the Author, by Benjamin W. Richardson, M.D. 8vo. cloth, 10s. 6d.

#### DR. STANHOPE TEMPLEMAN SPEER.

PATHOLOGICAL CHEMISTRY, IN ITS APPLICATION TO THE PRACTICE OF MEDICINE. Translated from the French of MM. BECQUEREL and RODIER. 8vo. cloth, reduced to 8s.

#### DR. STEGGALL.

STUDENTS' BOOKS FOR EXAMINATION.

I.

A MEDICAL MANUAL FOR APOTHECARIES' HALL AND OTHER MEDICAL BOARDS. Twelfth Edition. 12mo. cloth, 10s.

II.

A MANUAL FOR THE COLLEGE OF SURGEONS; intended for the Use of Candidates for Examination and Practitioners. Second Edition. 12mo. cloth, 10s.

III.

GREGORY'S CONSPECTUS MEDICINÆ THEORETICÆ. The First Part, containing the Original Text, with an Ordo Verborum, and Literal Translation. 12mo. cloth, 10s.

IV.

THE FIRST FOUR BOOKS OF CELSUS; containing the Text, Ordo Verborum, and Translation. Second Edition. 12mo. cloth, 8s.

v.

FIRST LINES FOR CHEMISTS AND DRUGGISTS PREPARING FOR EX-AMINATION AT THE PHARMACEUTICAL SOCIETY. Second Edition. 18mo, cloth, 3s. 6d.

#### MR. STOWE, M.R.C.S.

A TOXICOLOGICAL CHART, exhibiting at one view the Symptoms, Treatment, and Mode of Detecting the various Poisons, Mineral, Vegetable, and Animal. To which are added, concise Directions for the Treatment of Suspended Animation. Eleventh Edition. On Sheet, 2s.; mounted on Roller, 5s.

#### DR. SWAYNE.

OBSTETRIC APHORISMS FOR THE USE OF STUDENTS COMMENCING MIDWIFERY PRACTICE. With Engravings on Wood. Second Edition. Fcap. 8vo. cloth, 3s. 6d.

MR. TAMPLIN, F.R.C.S.E.

LATERAL CURVATURE OF THE SPINE: its Causes, Nature, and Treatment. 8vo. cloth, 4s.

10-->(1)

DR. ALEXANDER TAYLOR, F.R.S.E.

THE CLIMATE OF PAU; with a Description of the Watering Places of the Pyrenees, and of the Virtues of their respective Mineral Sources in Disease. Third Edition. Post 8vo. cloth, 7s.

DR. ALFRED S. TAYLOR, F.R.S.

I.

A MANUAL OF MEDICAL JURISPRUDENCE. Seventh Edition. Fcap. 8vo. cloth, 12s. 6d.

ON POISONS, in relation to MEDICAL JURISPRUDENCE AND MEDICINE. Second Edition. Fcap. 8vo. cloth, 12s. 6d.

MR. TEALE.

ON AMPUTATION BY A LONG AND A SHORT RECTANGULAR FLAP. With Engravings on Wood. 8vo. cloth, 5s.

DR. THEOPHILUS THOMPSON, F.R.S.

CLINICAL LECTURES ON PULMONARY CONSUMPTION.
With Plates. 8vo. cloth, 7s. 6d.

II.

LETTSOMIAN LECTURES ON PULMONARY CONSUMPTION; with Remarks on Microscopical Indications, and on Cocoa-nut Oil. Post 8vo., 2s. 6d.

.....

DR. THOMAS.

THE MODERN PRACTICE OF PHYSIC; exhibiting the Symptoms, Causes, Morbid Appearances, and Treatment of the Diseases of all Climates. Eleventh Edition. Revised by Algernon Frampton, M.D. 2 vols. 8vo. cloth, 28s.

MR, HENRY THOMPSON, F.R.C.S.

I.

STRICTURE OF THE URETHRA; its Pathology and Treatment. The Jacksonian Prize Essay for 1852. With Plates. Second Edition. 8vo. cloth, 10s.

II.

THE DISEASES OF THE PROSTATE; their Pathology and Treatment. Comprising a Dissertation "On the Healthy and Morbid Anatomy of the Prostate Gland;" being the Jacksonian Prize Essay for 1860. With Plates. Second Edition. 8vo. cloth, 10s.

DR. TILT.

I.

ON DISEASES OF WOMEN AND OVARIAN INFLAM-MATION IN RELATION TO MORBID MENSTRUATION, STERILITY, PELVIC TUMOURS, AND AFFECTIONS OF THE WOMB. Second Edition. 8vo. cloth, 9s.

THE CHANGE OF LIFE IN HEALTH AND DISEASE: a Practical Treatise on the Nervous and other Affections incidental to Women at the Decline of Life. Second Edition. 8vo. cloth, 6s.

E000-105

10-->10-

DR. THUDICHUM.

TREATISE ON THE PATHOLOGY OF URINE. Including a complete Guide to its Analysis. With Plates, 8vo. cloth, 14s.

DR. GODWIN TIMMS.

CONSUMPTION: its True Nature and Successful Treatment. Crown 8vo. cloth, 10s.

DR. ROBERT B. TODD, F.R.S.

## CLINICAL LECTURES ON THE PRACTICE OF MEDICINE:-

VOL. I .- PARALYSIS, DISEASES OF THE BRAIN, and other AFFEC-TIONS of the NERVOUS SYSTEM. Second Edition. Foolscap 8vo. cloth, 6s.

VOL. II .- CERTAIN DISEASES OF THE URINARY ORGANS, AND ON DROPSIES. Fcap. 8vo. cloth, 6s.

Vol. III .- ON CERTAIN ACUTE DISEASES. Foolscap 8vo. cloth, 6s.

New Edition, in one Volume, Edited by Dr. Beale, 8vo. cloth, 18s.

MR. TOMES, F.R.S.

A MANUAL OF DENTAL SURGERY. With 208 Engravings on Wood. Fcap. 8vo. cloth, 12s. 6d.

MR. JOSEPH TOYNBEE, F.R.S., F.R.C.S.

THE DISEASES OF THE EAR: THEIR NATURE, DIAG-NOSIS, AND TREATMENT. Illustrated with numerous Engravings on Wood. 8vo. cloth, 15s.

DR. TURNBULL.

AN INQUIRY INTO THE CURABILITY OF CONSUMPTION, ITS PREVENTION, AND THE PROGRESS OF IMPROVEMENT IN THE TREATMENT. Third Edition. 8vo. cloth, 6s.

11.

A PRACTICAL TREATISE ON DISORDERS OF THE STOMACH with FERMENTATION; and on the Causes and Treatment of Indigestion, &c. 8vo. cloth, 6s.

VESTIGES OF THE NATURAL HISTORY OF CREATION. Eleventh Edition. Illustrated with 106 Engravings on Wood. 8vo. cloth, 7s. 6d.

BY THE SAME AUTHOR.

## EXPLANATIONS: A SEQUEL TO "VESTIGES."

Second Edition. Post 8vo. cloth, 5s.

DR. UNDERWOOD.

TREATISE ON THE DISEASES OF CHILDREN. Tenth Edition. with Additions and Corrections by HENRY DAVIES, M.D. 8vo. cloth, 15s.

-∋€-

DR. UNGER.

BOTANICAL LETTERS. Woodcuts. Post 8vo., 2s. 6d.

Translated by Dr. B. PAUL. Numerous

MR. WADE, F.R.C.S.

STRICTURE OF THE URETHRA, ITS COMPLICATIONS AND EFFECTS; a Practical Treatise on the Affections. Fourth Edition. 8vo. cloth, 7s. 6d.

DR. WALLER.

ELEMENTS OF PRACTICAL MIDWIFERY; or, Companion to the Lying-in Room. Fourth Edition, with Plates. Fcap. cloth, 4s. 6d.

MR. HAYNES WALTON, F.R.C.S.

SURGICAL DISEASES OF THE EYE. With Engravings on Wood. Second Edition. 8vo. cloth, 14s.

DR. WATERS, M.R.C.P.

THE ANATOMY OF THE HUMAN LUNG. The Prize Essay to which the Fothergillian Gold Medal was awarded by the Medical Society of London. Post 8vo. cloth, 6s. 6d.

DR. EBEN. WATSON, A.M.

ON THE TOPICAL MEDICATION OF THE LARYNX IN CERTAIN DISEASES OF THE RESPIRATORY AND VOCAL ORGANS. 8vo. cloth, 5s.

THE SURGEON'S READY RULES FOR OPERATIONS IN SURGERY. Royal 8vo. cloth, 10s. 6d.

DR. WEBER.

A CLINICAL HAND-BOOK OF AUSCULTATION AND PERCUSSION. Translated by John Cockle, M.D. 5s.

DR. WEGG.

OBSERVATIONS RELATING TO THE SCIENCE AND ART OF MEDICINE. 8vo. cloth, 8s.

MR. T. SPENCER WELLS, F.R.C.S.

PRACTICAL OBSERVATIONS ON GOUT AND ITS COMPLICATIONS, and on the Treatment of Joints Stiffened by Gouty Deposits. Foolscap 8vo. cloth, 5s.

SCALE OF MEDICINES WITH WHICH MERCHANT VES-SELS ARE TO BE FURNISHED, by command of the Privy Council for Trade; With Observations on the Means of Preserving the Health of Seamen, &c. &c. Seventh Thousand. Fcap. 8vo. cloth, 3s. 6d.

DR. WEST.

LECTURES ON THE DISEASES OF WOMEN. Second Edition.

8vo. eloth, 16s.

MR. WHEELER.

HAND-BOOK OF ANATOMY FOR STUDENTS OF THE FINE ARTS. With Engravings on Wood. Fcap. 8vo., 2s. 6d.

DR. WHITEHEAD, F.R.C.S.

ON THE TRANSMISSION FROM PARENT TO OFFSPRING OF SOME FORMS OF DISEASE, AND OF MORBID TAINTS AND TENDENCIES. Second Edition. 8vo. cloth, 10s. 6d.

DR. WILLIAMS, F.R.S.

PRINCIPLES OF MEDICINE: An Elementary View of the Causes, Nature, Treatment, Diagnosis, and Prognosis, of Disease. With brief Remarks on Hygienics, or the Preservation of Health. The Third Edition. 8vo. cloth, 15s.

THE WIFE'S DOMAIN: the Young Couple—the Mother—the Nurse—the Nurse Post 8vo. cloth, 3s. 6d.

DR. JOSEPH WILLIAMS.

INSANITY: its Causes, Prevention, and Cure; including Apoplexy, Epilepsy, and Congestion of the Brain. Second Edition. Post 8vo. cloth, 10s. 6d.

DR. J. HUME WILLIAMS.

UNSOUNDNESS OF MIND, IN ITS MEDICAL AND LEGAL CONSIDERATIONS. 8vo. cloth, 7s. 6d.

DR. WILLIAMSON, STAFF-SURGEON.

NOTES ON THE WOUNDED FROM THE MUTINY IN INDIA: with a Description of the Preparations of Gunshot Injuries contained in the Museum at Fort Pitt. With Lithographic Plates. 8vo. cloth, 12s.

MR. ERASMUS WILSON, F.R.S.

THE ANATOMIST'S VADE-MECUM: A SYSTEM OF HUMAN

ANATOMY. With numerous Illustrations on Wood. Eighth Edition. Foolscap 8vo. cloth, 12s. 6d.

10-->41>

ANA

MR. ERASMUS WILSON, F.R.S. (continued).

11.

DISEASES OF THE SKIN: A Practical and Theoretical Treatise on the DIAGNOSIS, PATHOLOGY, and TREATMENT OF CUTANEOUS DISEASES. Fourth Edition. 8vo. cloth, 16s.

THE SAME WORK; illustrated with finely executed Engravings on Steel, accurately coloured. 8vo. cloth, 34s.

III.

HEALTHY SKIN: A Treatise on the Management of the Skin and Hair in relation to Health. Sixth Edition. Foolscap 8vo. 2s. 6d.

IV.

PORTRAITS OF DISEASES OF THE SKIN. Folio. Fasciculi I. to XII., completing the Work. 20s. each.

V.

ON SYPHILIS, CONSTITUTIONAL AND HEREDITARY;
AND ON SYPHILITIC ERUPTIONS. With Four Coloured Plates. 8vo. cloth,
16s.

VI.

A THREE WEEKS' SCAMPER THROUGH THE SPAS OF GERMANY AND BELGIUM, with an Appendix on the Nature and Uses of Mineral Waters. Post 8vo. cloth, 6s. 6d.

VII.

THE EASTERN OR TURKISH BATH: its History, Revival in Britain, and Application to the Purposes of Health. Foolscap 8vo., 2s.

#### DR. G. C. WITTSTEIN.

PRACTICAL PHARMACEUTICAL CHEMISTRY: An Explanation of Chemical and Pharmaceutical Processes, with the Methods of Testing the Purity of the Preparations, deduced from Original Experiments. Translated from the Second German Edition, by Stephen Darby. 18mo. cloth, 6s.

#### DR. HENRY G. WRIGHT.

HEADACHES; their Causes and their Cure. Third Edition. Fcap. 8vo. 2s. 6d.

#### MR. YEARSLEY.

I.

DEAFNESS PRACTICALLY ILLUSTRATED; being an Exposition of Original Views as to the Causes and Treatment of Diseases of the Ear. Fifth Edition. Foolscap 8vo., 2s. 6d.

II.

ON THE ENLARGED TONSIL AND ELONGATED UVULA, and other Morbid Conditions of the Throat. Seventh Edition. 8vo. cloth, 5s.

E CARCONOLI

古十七名文をひて見

Fcap. 8vo. cloth, 12s. 6d. each.

"We here give Mr. Churchill public thanks for the positive benefit conferred on the Medical Profession, by the series of beautiful and cheap Manuals which bear his imprint."—

British and Foreign Medical Review.

## AGGREGATE SALE, 121,000 COPIES.

- The ANATOMIST'S VADE-MECUM. A System of Human Anatomy. With numerous Illustrations on Wood. Eighth Edition. By Erasmus Wilson, F.R.C.S., F.R.S.
- BOTANY. By Robert Bentley, F.L.S., Professor of Botany, King's College.
- CHEMISTRY. With numerous Illustrations on Wood. Eighth Edition. By George Fownes, F.R.S. Edited by H. Bence Jones, M.D., F.R.C.P., and A. W. Hofmann, F.R.S.
- DENTAL SURGERY. With 208 Illustrations on Wood. By John Tomes, F.R.S.
- MATERIA MEDICA. With numerous Illustrations on Wood. Third Edition. By J. Forbes Royle, M.D., F.R.S., and Frederick W. Headland, M.D., F.L.S.
- MEDICAL JURISPRUDENCE. Seventh Edition. By ALFRED SWAINE TAYLOR, M.D., F.R.S.
- PRACTICE OF MEDICINE. Second Edition. By G. HILARO BARLOW, M.D., M.A.
- The MICROSCOPE and its REVELATIONS. With numerous Illustrations on Wood. Second Edition. By W. B. CARPENTER, M.D., F.R.S.
- NATURAL PHILOSOPHY. With numerous Illustrations on Wood. Fifth Edition. By Golding Bird, M.D., M.A., F.R.S., and Charles Brooke, M.B., M.A., F.R.S.
- OBSTETRICS. With 186 Engravings on Wood. By W. Tyler Smith, M.D., L.R.C.P.
- OPHTHALMIC MEDICINE and SURGERY. With coloured Engravings on Steel, and numerous Illustrations on Wood. Second Edition. By T. Wharton Jones, F.R.C.S., F.R.S.
- PATHOLOGICAL ANATOMY. With numerous Illustrations on Wood. By C. Handfield Jones, M.B., F.R.C.P., and E. H. Sieveking, M.D., F.R.C.P.
- PHYSIOLOGY. With numerous Illustrations on Steel and Wood. Third Edition. By WILLIAM B. CARPENTER, M.D., F.R.S.
- POISONS. Second Edition. By Alfred Swaine Taylor, M.D., F.R.S.
- PRACTICAL SURGERY. With numerous Illustrations on Wood. Fourth Edition. By WILLIAM FERGUSSON, F.R.C.S.

Printed by W. Blanchard & Sons, 62, Millbank Street, Westminster.



