A guide to the practical examination of urine: for the use of physicians and students.

Contributors

Tyson, James, 1841-1919. University of Toronto

Publication/Creation

Philadelphia: Blakiston, 1893.

Persistent URL

https://wellcomecollection.org/works/w2yspbdv

License and attribution


This material has been provided by This material has been provided by the Gerstein Science Information Centre at the University of Toronto, through the Medical Heritage Library. The original may be consulted at the Gerstein Science Information Centre, University of Toronto. where the originals may be consulted.

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.




Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org





















BY THE SAME AUTHOR.

BRIGHT'S DISEASE AND DIABETES. A Treatise on Bright's 8vo. Price \$3.50. Therapeutics. With Colored Plates and many Wood Engravings. Disease and Diabetes. With Especial Reference to Pathology and

"Dr. Tyson has presented his subjects so clearly and concisely, and has selected his material with such sound judgment, that his work cannot fail to be useful to practitioners and students."—The American Journal of the Medical Sciences.

"The symptoms are clearly defined, and the treatment is exceedingly well described, so that every one reading the book must be profited."—Cincinnati

Lancet and Clinic.

THE CELL DOCTRINE: Its History and Present State, together and other Illustrations. with a copious Bibliography of the Subject. With a Colored Plate

Second Revised Edition. Cloth, price \$2.00.

The present edition shows an increase in size of about fifty pages, and almost every page furnishes evidence of careful revision"—American Journal of the Medical "The first edition of Dr. Tyson's work appeared in 1870, and at once received the favorable reception from the profession to which it was entitled by its merits,


résumé of the origin and advance of the doctrine of cell evolution. The work is calculated to be of decided benefit to all who desire a general knowledge of the history, progress, and peculiar phases of the cell doctrine."—American Prac-"The author in this work presents to medical students a concise and instructive

A MANUAL OF PHYSICAL DIAGNOSIS. For the use of Students and Physicians.

Second Edition. Illustrated. Cloth, price \$1.50.

"The present volume is by a clinical teacher of much experience and a practi-tioner of well-known ability, so that his work has a value outside of a class of undergraduates." - American Journal of the Medical Sciences.





Et-

A GUIDE

PRACTICAL EXAMINATION

OF

URINE.

FOR THE USE OF PHYSICIANS AND STUDENTS.

BY

JAMES TYSON, M. D.,

Professor of Clinical Medicine in 1119 University of Pennsylvania, and Physician to the Hospital of the University; Fellow of the College of Physicians of Philadelphia, etc., etc., etc.


EIGHTH EDITION.


REVISED AND CORRECTED, WITH A COLORED PLATE AND WOOD ENGRAVINGS.

PHILADELPHIA:
P. BLAKISTON, SON & CO.,
NO. 1012 WALNUT STREET.

J. A. CARVETH & CO. PUBLISHERS, TORONTO, - ONT.

1893.







LIST OF ILLUSTRATIONS.

COLORED PLATE.

WOOD ENGRAVINGS.	
	PAGE
Measuring-glasses for measuring urine,	18
Apparatus for determining specific gravity,	26
Small urinometer,	28
Dr. Squibb's new urinometer and floating cylinder,	29
Testing for albumin by nitric acid,	40
Esbach's Albuminometer,	55
Burette stand with different forms of burette,	80
Dr. Johnson's picro-saccharimeter,	06
Laurent's polarizing saccharimeter,	IOI
ter,	102
12. Illustrations of the use of Fleischl's saccharimeter,	103
Crystals of nitrate of urea (from Beale),	155
Burette stand with two forms of burette,	160
Marshall's apparatus for estimation of urea by the hypobro-	
mite process,	164
	99I
Prismatic crystals of sodium urate, spherules of ammonium	
Jo	
	161
Spiculated spherules of ammonium urate along with triple	
(ammonio-magnesian) phosphate and octohedral crystals of	
the oxalate of lime (Ranke),	193
10. More usual forms of uric acid crystals (after Harley),	197

LIST OF ILLUSTRATIONS.

TABLE OF CONTENTS.

	100				ar	
					Q	
		0,			in	
		rin			ent	
		5		65	,A	
		Jo		ii.	-	
		-		5	he	
		tio		e	n t	
12		65		=	-=	
INTRODUCTION.		Reagents and Apparatus Required for Examination of Urine,	Selecting a Specimen of Urine,	of	To Determine the Amount of Solid Matters in the Twenty-four	hours' Urine,
2		xa		LS	att	
		E		cte	M	
ă		for		ara	P	
Ö		P		Ä	10	
2		ire		~	CO	
E		nb	ne,	Ca	of	
Z		Re	E	H	III	
		55	f	he	non	
		atn	0	0	H	
		ar	en	no	-	é,
	ne	bb	Į.	-	the	i.
	T	Y	ec.	ica	9	D
	1	pu	S	ys	ii.	28.
	0	3	es	Ph	T.B.	oni
	OIL	nts	Sil	7	ete	A
	eti	ge	cti	er	A	
	ecr	ea	ele	en	0	
	S	K	S	0	F	

	36	54	58	58	59	62	29	69	70	70	107	107	100	109	110	121	121	130	144	151	152	154	170
										4													
				-																			
										-												1	1
																						4.	
																						1	
												100							-				
										4										-0.			
											e,												
				-					0		Urine		1		es,								
-				Paraglobulin						*					Aceton and Acetone Producing Substances								200
-				pnq					-		.5		-		sta								
LANI		in		010							P				qn			15					
4		n n		rag							En III				S		ŝ	ter					
ч		P		Pa			45				5				Bu		ter	at					
	1	A		1)nc				es				Ci		lat	N					
	efo	of		0			o d			-	nc			5	du		Normal Coloring Matters,	Abnormal Coloring Matters			1		
	er	-	e,	=			pe				sta		se,	SO	20		80	.E					
	中	tio	·E	de			ro		Urine,		qu		lo	ac	-		·E	0		-			
	ts	na	D	H			-		H		S		NA.	H	n		olo	0		Sin			
	es	stin	1	5			0		-		ne		La	or	et	rs,	O	lai		TO			
	=	E	ds	lin			se		.=		H		-	K	Ac	tte	lal	E	s's	T	46		
	nu	ve	ie.	pn			ŭ		nd		ch		r	M	P	Ma	E	no	eid	7	ter		-
	0	ati.	ro	012		-5	pn		Ou		ac		gg	4	an	10	So	Ab	A	an	Int		pid
	ii.	ii.	4	3-u	0,	ne	la.		10	se	S	e,	su	0	00	ing	Total.	7774	-	.8	Z		Ac
	Albumin and Tests therefor, .	Quantitative Estimation of Albumin,	Other Proteids in Urine,	Serum-globulin, Globulin or	Mucin, .	Peptone,	Hemialbumose or Propeptone	Fibrin,	Sugars found in I	Glucose,	Other Saccharine Substances found in	Inosite, .	Fruit-sugar or Lævulose,	Sugar of Milk or Lactose,	eto	Coloring Matters,			Biliary Acids,	Leucin and Tyrosin,	Fatty Matters,	Urea,	Uric Acid,
	=	2	=	Ser	Mu	Pel	He	E	ing	15	3	Inc	Fin	Sug	1c	3			Bill	3	Fa	5	5
	-4	-	-	-	7.00	-	-	-	42			-		42	-				-	-	-	Total Co	

Urates, PAG Chlorides, 17 Chlorides, 17 Chlorides, 17 Phosphates, 17 Nuphates, 17 Uninary Deposits, 17 Unorganized Sediments, 19 Uric Acid, 19 Uric Acid, 19 Uric Acid, 19 Uric Acid, 20 Oxalate of Lime, 20 Earthy Phosphates, 20 Calcium Carbonate, 20 Calcium Sulphate, 21 Leucin and Tyrosin, 20 Calcium Sulphate, 21 Leucin and Tyrosin, 21 Cystin, 22 Cystin, 23 Mucus and Pus, 21 Epithelium, 21 Epithelium, 21 Epithelium, 21 Epithelium, 21 Epithelium, 22 Blood-corpuscles, 22 Fungi, 22 Entozoa, 23 Entozoa, 23	PART II. posits, d Sediments, Acid Compounds, late of Lime, thy Phosphates, ium Carbonate, ium Sulphate, cin and Tyrosin, in, Sediments, sus and Pus, thelium, d-corpuscles, re-crasts, matozoids, Elements of Morbid Growths, zooa, Preservation of Organized Urinary Sediments for Sub- quent Examination, Diagnosis of Renal Diseases, Diagnosis of Renal Diseases, Diagnosis of Calculi, APPENDIX. Reducing the Metric System into the English and versa, and for Converting Degrees Centigrade to		to	0	ad	10	Ë	0		99	100	Ā	79	E.	/er	M	0	H	5	nd	20	50,	197		vic	-		
PART II. posits, d Sediments, Acid, Acid, Acid, Acid Compounds, late of Lime, thy Phosphates, itum Carbonate, itum Sulphate, itum s	PART II. posits, A Sediments, A Sedid Compounds, late of Lime, thy Phosphates, ium Carbonate, ium Sulphate, cin and Tyrosin, in, Sediments, sus and Pus, thelium, d-corpuscles, re-casts, matozoids, Elements of Morbid Growths, sooa, Preservation of Organized Urinary Sediments for Sub- quent Examination, Diagnosis of Renal Diseases, Diagnosis of Renal Diseases, PART III. lculi, Reducing the Metric System into the English and					0								*						C								
II. Urinary Sediments for Subes, Luii. Luii. Luii.	II. Urinary Sediments for Subes, Ulili. Ulili. Ulili.		bd	10	sh	100	B	-	E	0	E.	n	ter	ys		H.	Me	-	E	70	cin	du	Re		fo	Ses	Idi	H
PART II. PART II. Compounds, Lime, Sphates, Irphonate, Iphote, Iphote, Iryrosin, Iryrosin, Ityrosin, It	ments, PART II. PART II. Compounds, Lime, sphates, sphates, liphate, li Tyrosin, Pus, Pus, Pus, sids, auscles, uscles, uscles, uscles, sids, PART III. Composition of Calculi, APPENDIX.	257													on	ati	Bi.	an	馬	an	35	di	10	ec.	f	0	00	K
PART II. PART II. Compounds, Lime, ssphates, photate, photate, pus, suscles, suscles, suscles, ration of Morbid Growths, vation of Organized Urinary Sediments for Examination, Examination, Stamination, PART III. Composition of Calculi,	ments, Compounds, Lime, Sephates, promate, liphate, li Tyrosin, nts, Pus, sids, nts of Morbid Growths, vation of Organized Urinary Sediments for Sub- Examination, sis of Renal Diseases, PART III. Composition of Calculi,											70	×	9	Z	E	AP											
ments, PART II. PART II. Compounds, Lime, ssphates, photate, iphotate, iphotate,	ments, PART II. PART II. Compounds, Lime, Sphates, phonate, liphate, liphate, li Tyrosin, It Tyrosin, sids, Pus, uscles, uscles, uscles, vation of Organized Urinary Sediments for Sub- Examination, Stamination, Samination, PART III.	25											=	lcı	C	9	ň	ILIC	SO	Im	8	he	0	E.	en	et	P	-
s,	s, sylvania significant signif	25																					H	lici	0	ary	Tin.	C
s,	s,												F	-	F	A	P											
PART II. s, sounds, ce, ce, ie, ie, ie, ie, iosin, osin, of Morbid Growths, of Morbid Growths, of Organized Urinary Sediments for Sub- ination,	PART II. s. pounds, tes, tes, tes, ie, osin, of Morbid Growths, of Organized Urinary Sediments for Sub- ination,	24											-	ses	sea	D	2	en	R	of	osis	gn	ia	H	itia	rei	iffe	D
PART II. PART II. S. Sounds, Substitute of the state	PART II. s, sounds, tes, tes, tes, te, te, te, te, te, te, te, te, te, te	24															D,	Eio.	na	B	Exe	H	ier	equ	nn.			
PART II. PART II. s. sounds, c. c. c. c. ites, ate, e. ites, ate, sosin, osin, f Morbid Growths,	PART II. PART II. s. sounds, tes, ate, tes, tes, osin, of Morbid Growths,		F	Su	0	s f	nt	BE	di	Se	7	ma	Or I	2	ize	an	gr(0	0	ion	vat	ser	re	e I	Th			
PART II. PART II. s. sounds, cetes, ates, ates, ates, osin, of Morbid Growths,	PART II. PART II. s, counds, tes, tes, tes, osin, osin, Morbid Growths,	24																					00	oz	En			
PART II. s, sounds, ctes, ate, tes, ate, sosin, s,	PART II. s, c, c, tes, ate, tes, tes, sin,	24											bs	M	iro		bio	101	2	of	nts	me	le	to He	Th			
PART II. s, c, c, tes, ate, tes, tes, sin,	PART IL s, c, c, tes, ate, tes, tes, sosin,	23																						gi.	Fu			
PART II. s, c, c, tes, ate, tes, tes, ssin,	PART IL s, counds, cetes, cetes, cetes, cetes, cosin, cos	23																		3.	bid	020	ate	m	Spe			
PART II. s, counds, ctes, tes, tes, tes, tes, sounds,	PART II. s counds, cs, cs, cs, cs, cs, cs, cs, cs, cs, c	22											1									Sts	Car	be-	Tu			
PART IL s younds, cet, cet, cet, cet, cet, cet, cet, cet	PART II. s, counds, ce, tes, tes, tes, tes, tes, tes, tes, te	22																		les	usc	die	00	Od.	Blo			
PART IL s younds, c c c tes, tes, tes, osin,	PART II. s, ounds, ounds, etc, tes, tes, tes, tes, tes, tes, tes, tes	22																				H	eli	B	Ep			
PART II. s, counds, ce, ce, ce, ce, ce, ce, ce, ce, ce, ce	PART II. s, counds, ces, ces, ces, ces, osin,	21																		us,	P	nd	20	Cu	Mu			
PART II. eposits, ed Sediments, c Acid, c Acid Compounds, alate of Lime, thy Phosphates, cium Carbonate, cium Sulphate, cium Sulphate, cium Sulphate, acin and Tyrosin, tin,	ates,	21																			nts	me	di	· Se	ed	n.	GS	0
PART II. eposits, ed Sediments, c Acid, c Acid Compounds, alate of Lime, thy Phosphates, cium Carbonate, cium Sulphate, cium Sulphate, cium Sulphate, cium Sulphate,	ates,	21														*								Stin	S			
PART II. eposits, ed Sediments, c Acid, c Acid Compounds, alate of Lime, thy Phosphates, cium Carbonate, cium Sulphate,	ates,	21												-				-	Si.	yro	T	nd	n a	ici	Le			
PART II. eposits, ed Sediments, c Acid, c Acid Compounds, alate of Lime, alate of Lime, cium Carbonate,	ates,	21.																	20	at	ldi	Su	B	Ciu	Cal	-		
PART II. eposits, ed Sediments, c Acid, c Acid Compounds, alate of Lime, thy Phosphates,	ates,	21																-	te	ons	urb	0	B	ciu	3			
PART II. eposits, ed Sediments, c Acid, c Acid Compounds, alate of Lime,	ates,	20																	es	hat	dsc	Pho	y I	=	Ea			
PART II. eposits, ed Sediments, c Acid,	ates,	20																	*	me	5	of	e	ala	ox Ox			
s,	ates,	20															S	nd	no	dun	S	E.	A C	0	I.			
s,	ates,	19																	*			id,	1ci	0 1	5			
s,	ates,	19												•						nts	me	di	S	ed	niz	833	100	C
S,	ates,	18																				ts,	OSi	ep	D	ary	En	C
	ates,												H	_	RI	2	H											
	ates,	18																*							tes	ha	al p	S
	ates,	17																						S	ate	dds	nos	1
	ates,	17					4																		Cs.	. Ä	DIG	10
		PAG 17														*										es.	rat	0

PRACTICAL

EXAMINATION OF THE URINE.

INTRODUCTION.

SECRETION OF URINE.

As the result of this, a filtration of the watery constituents the convoluted tubules into which has descended the thin, aqueous filtrate from the Malpighian bodies. Here are It is true that in the capillaries of the Malpighian tuft the blood-pressure is relatively greater on account of the resistof the blood, with some dissolved salts, takes place into the when it reaches the second capillary network embracing the essential elements of a complete osmometer-an animal THE theory which explains the secretion of urine most consistently with the facts, is one which, while it makes the process partly physical, requires also something of the nature of elaboration in the office of the kidney. Nothing can be more attractive at first thought than the theory of Ludwig, according to which the process is a purely physical one-partly a filtration and partly a diffusion, or osmosis. ance to the exit of the blood through the efferent vessel. Malpighian capsule. Thus the blood is greatly thickened

membrane formed by the thin wall of the capillary and the delicate basement membrane of the tubule, having a dense fluid, the blood, on one side, and a thin saline solution on the other. An interchange now takes place, as the result of which a current sets in, of liquid from the tubules to the blood, and of the products of regressive metamorphosis—urea and salts—to the tubules, concentrating the fluid in the latter, making it, in a word, urine; while the albuminous constituents of the blood are retained in it because of their well-known indisposition to osmosis.

The objection formerly made to the physical nature of the act of secretion of urine, on the ground that we cannot by this method account for the formation of an acid fluid from an alkaline one, no longer holds, since Dr. Ralfe, of London, has shown this to be quite possible. Into one limb of a small, U-shaped tube, fitted with a membranous diaphragm at the bend, he introduced an alkaline solution of sodium bicarbonate, and into the other limb a solution of neutral sodium phosphate. He then passed a weak electric current through the solutions. In a short time the fluid in the limb connected with the positive pole became acid from the formation of acid sodium phosphate, while the fluid in the limb connected with the negative pole increased in alkalinity. The changes are represented by the following formula:—

Sodium Neutral sodium Sodium Acid sodium bicarbonate. phosphate, carbonate, phosphate, NaHCO₃ + NaH2PO₄ = Na₂CO₃ + NaH₂PO₄.*

One important fact, however, remains unaccounted for by this theory, clear and simple as it is. This is, that if

^{*} Medical News and Library, October, 1871, from London Lancet, July 4, 1871.

the tubules are stripped of their epithelium, as they often are in disease, urea and other products of regressive metamorphosis are no longer so thoroughly removed, but accumulate in the blood, producing the phenomena of the condition known as *uræmia*. We must therefore admit some elaborating action on the part of the epithelium, as originally suggested by Bowman. Doubtless, however, a part of the act is physical—a process of transudation or filtration, and of diffusion or osmosis.

The experimental researches of Heidenhain* have settled the question in favor of an active elaborating office on the part of the epithelium of the kidney. Heidenhain injected into the blood of animals indigo-carmine, a substance which is promptly separated by the kidneys. He removed these organs at suitable intervals after the operation and examined them minutely. In no instance did he find any of the indigo-carmine in the Malpighian capsules, but the cells lining the convoluted tubules and the looped tubes of Henle were filled with it, as was also the lumen of the tubes, if the animal was killed sufficiently long after the injection. Similar experiments with urate of sodium showed that it is secreted at the same place and in the same manner.

REAGENTS AND APPARATUS REQUIRED FOR QUALITATIVE AND APPROXIMATE ANALYSIS.

It is not a matter of very great importance in what form of bottles reagents are kept. They should hold enough—four ounces is a convenient quantity—and be provided with

^{*} Max Schultze's Archiv, vol. x, 1874, p. 1, and Pflüger's Archiv, vol. ix, 1874, p. 1.

[†] All reagents and apparatus suitable for urinalysis may be obtained of Bullock & Crenshaw, 528 Arch Street, Philadelphia.

16

ground-glass stoppers for the acids, but the alkalies are better kept in bottles with rubber stoppers. Those required are as follows:—

- I. Pure colorless nitric acid (HNO₃).
- 2. Nitroso-nitric acid, the brown, fuming nitrous acid of commerce, —nitric acid containing nitrogen tetroxide (HNO₃ + N₂O₄ or NO₂).
- 3. Pure hydrochloric acid (HCl).
- 4. Pure colorless sulphuric acid (H2SO4).
- 5. Pure acetic acid (C2H4O2).
- Liquor potassæ, U. S. P. The sp. gr. is 1065, and it contains .058 of potassium hydroxide (HKO).
- 7. Solution of caustic potash, or caustic soda, I part to 2 of distilled water, sp. gr. 1330 +, to be spoken of in the text as the "stronger solution of potash." It is the ætzkalilauge (or ætznatronlauge, if soda) of the German Pharmacopœia, and contains from .30 to .31 of the hydrate of potassium (or of sodium).
- Solution of sodium carbonate, I part water and 2 parts of the crystallized salt.

9. Solution of barium chloride, 4 parts crystallized barium chloride,

- 16 of distilled water, and I of hydrochloric acid. to. Liquor ammonia, U. S. P.
- 11. The magnesian fluid, containing of magnesium sulphate and pure ammonium chloride, each 1 part, distilled water 8 parts, and pure liquor ammoniæ 1 part.
- 12. Solution of copper sulphate, say I gram to 30 c.c., or 15 grs. to 13.
- Pavy's or Fehling's copper solutions, made as directed under volumetric analysis for sugar.
- 14. Solution of silver nitrate, I part to 8 of distilled water.
- Solution of neutral lead acetate (sugar of lead), 1 part to 4 of distilled water.
- 16. Solution of basic lead acetate, I part to 4 of distilled water
- 17. Distilled water, a litre or a quart.
- 18. Alcohol, 95 per cent., half a litre or a pint.

Other solutions as required.

Apparatus.

A note and drawing-book.

bases, so that they may stand on a shelf or table, are convenient and desirable; see Fig. 5. Some tubes should be graduated in I dozen test tubes, assorted sizes, some narrow. Some test-tubes with divisions of a centimetre and fractions thereof. They may be

sediment, or of albumin after its precipitation by heat. Test-tube rack and drainer.

used as fluid measures, and to determine the proportion of a

4 conical glasses. (Observe that there is not a convexity at the bottom, and the edge should be ground so that they may be covered with ground-glass covers and thus made air tight.)

2 or 3 smooth wineglasses, with broad bottoms, of the kind sometimes known as "collamore" wineglasses.

Red and blue litmus paper; Swedish filtering paper.

Urinometer and urinometer glass.

4 ground-glass covers, assorted sizes.

Spirit-lamp, or bunsen burner.

3 porcelain capsules.

6 beaker glasses, small and medium sizes. 6 watch-glasses.

I long narrow funnel tube 12 inches long and 1 inch wide, for filtering 3 glass funnels, assorted sizes.

Glass stirring-rods and plain glass pipettes. through animal charcoal.

city of 2000 cubic centimetres or more.

I large receiving glass to measure twenty-four hours' urine, with capa-

I graduated measuring glass holding 500 c.c.

I wash-bottle with distilled water.

I retort stand; water-bath.

I or 2 sheet-iron tripods with wire gauze to cover.

I 100 minim pipette; I volume pipette for 5 c.c., another for 10 c.c.

Platinum spoon; tongs.

Swabs for cleaning test-tubes, etc.

A microscope with two object glasses, a 1 or 1 inch, and a 1 inch or 10 inch; stage micrometer; camera lucida for drawing; glass slides,

thin covers, shallow cells; a double nose-piece for quickly changing objectives is very convenient.

For volumetric analysis are required in addition :-

A full set of volume pipettes, 5, 10, 15, 20, 30, 50 c.c.

I dropping pipette holding I c.c., graduated in Toths, and fractions

2 burettes of 50 c.c. capacity; burette stand. A half-litre flask.

Volumetric solutions as directed under Volumetric Analysis.

they may be, he should be provided with a balance which will turn with a milligram, or with to of a grain if the English system is used. If the solutions are made by the physician himself, as

SELECTING A SPECIMEN OF URINE.

In obtaining a specimen of urine for chemical examina-The same is true of sugar, although a trace apparent in the morning urine, a meal, and sometimes when there is no in urine, it is often increased after a a small quantity of albumin is present modification of this course. Thus, when stances, however, constantly demand a to take a part of the total. Circumthe only accurate method is, therefore, vary during the twenty-four hours, and and other properties are well known to urine, as the specific gravity, reaction, part of the whole twenty-four hours' tion, it should, as far as possible, be a little will be detectable after a meal.



urine is to ask for two specimens, one passed on retiring very good plan in the case of saccharine or about an hour after dinner, and the other passed on rising. For *microscopical* examination, the importance of having a fresh specimen which has not undergone decomposition or change of reaction is paramount. In Fig. 1 are represented forms of glass vessels used for measuring large quaratities of urine.

In warm weather urine decomposes rapidly, becoming alkaline, and it is especially important where microscopic examination is to be made that such decomposition should be obviated, as certain casts and crystalline sediments of uric acid are often dissolved in alkaline urine. Where urine cannot be examined in the fresh state, and especially when it has to be sent any distance, decomposition may be averted by adding five grains of salicylate of sodium to each four ounces of urine.

GENERAL PHYSICAL AND CHEMICAL CHARACTERS OF URINE.

Normal urine may be described as a transparent, aqueous fluid, of a pale lemon-yellow hue, acid reaction, specific gravity of about 1020 when passed in the average quantity of 1500 cubic centimetres (50 ounces) in the twenty-four hours, and possessing an odor which can only be indicated as "characteristic" or "urinous." The odor is sometimes spoken of as "aromatic."

Each one of these characters is, however, liable to some variations within the limits of health, as well as in disease, and with these variations we should be thoroughly familiar before interpreting a given specimen.

I. As to Transparency.—This, although quite a constant, can scarcely be considered an essential character of normal urine, while, on the other hand, it by no means

follows that because a given specimen of urine is transparent it is therefore normal.

Causes of Diminished Transparency.—Diminished transparency may be due to one of three causes.

r. Even urine which is apparently perfectly transparent when passed, commonly exhibits, a few minutes after standing, floating somewhere between the top and bottom, a faint cloud, which is said to be mucus derived from the genito-urinary tract. In the urine of females this cloud is apt to be more distinctly visible, in consequence of a larger amount of epithelium from the vagina and adjacent mucous surfaces in this sex. There is nothing abnormal in the presence of such an amount of mucus as is covered by the above description. The effect of alkalies, heat, and strong acids is to leave the appearance unchanged, but acetic acid may produce a slight increase of the opacity by coagulating the mucin. Mucus can be filtered out, leaving the urine clear.

2. Normal, slightly acid urine may be turbid at the moment when passed, by reason of the presence of the earthy phosphates of calcium and magnesium. These, shortly after passing, begin to subside, and, within half an hour, present an appearance not unlike that of mucus,—that of a flocculent mass, floating somewhere between the top and bottom of the vessel. But still later, generally within an hour, they have approached the bottom, and become a sediment, cloudy and bulky, leaving a transparent supernatant fluid.

To test the nature of such sediment add a few drops of any acid, as nitric, which will cause the prompt disappearance of any earthy phosphate, while the application of heat will increase the deposit, such increase being also rapidly dissipated on adding an acid.

The more or less constant presence of the earthy phos-

phates above mentioned cannot be considered abnormal. Requiring an acid urine to keep them in solution, a diminution of the degree of acidity may result in their precipitation, which is further increased by an alkaline reaction. Such diminished acidity and substitution of alkalinity always takes place from two to four hours after a meal, and the deposit is therefore commonly observed at such time.

3. Urine is sometimes rendered turbid by the presence of the so-called *mixed urates* of sodium, potassium, calcium, and magnesium. The most frequent cause of this precipitation in normal urine is a reduction in the temperature of the urine after being passed. Although highly soluble in water at the temperature of the body, the urates drop promptly from a cold urine, of a temperature such as would prevail in a room without fire in winter.

As in the case of earthy phosphates, such opacity soon diminishes by subsidence of the disseminated urates, which becomes a white or pink *deposit*, less bulky than that of phosphates; urates are also apt to be precipitated on the side of the vessel.

To test the nature of this deposit apply heat, which quickly causes the dissipation of urates, while a precipitate of phosphates is increased by it.

1. **Pathologically**, urine may be opaque or semi-opaque from abnormal degrees of the above conditions, or from the presence of *pus*, which also subsides with a rapidity inversely as the quantity of mucus. If the latter is absent, or present in small quantity, the subsidence is rapid; if, on the other hand, it is large, subsidence is slow, often requiring several hours. The turbidity of such urine is *increased* by the application of heat and acids, in conse-

quence of the precipitation of albumin, which is always a constituent of *liquor puris*.

2. The presence of fat in a state of minute subdivision, as in the so-called *chylous* urine, produces a degree of turbidity ranging from mere cloudiness to absolute milkiness. In such urine the fatty matter is disposed to rise and form a whitish, creamy layer on top of a less turbid fluid. Such urine, not uncommon in tropical countries, is also sometimes met in temperate climates.

II. As to Consistence.—In health, urine is never anything else but aqueous, that is, it drops and flows readily, like water.

Pathologically, it often becomes viscid, glutinous, and separable with difficulty into drops, or not at all. Such state may be due to the presence of an excess of pure mucus, or of a mixture of mucus and pus, and very frequently it is caused by the action upon pus of an alkalinity due to the presence of ammonium carbonate. This will be again alluded to.

In the chylous urine above referred to, the presence of the molecular fat also increases the consistence of the urine.

III. As to Color.—While normal urine may be characterized in general terms as pale-yellow, lemon-yellow, or amber-hued, there may be considerable variation in health. Due to the presence, in solution, of the normal coloring matters, the color is deeper or paler according to the proportion of water dissolving them. After copious libations of beer or water, the quantity of urine discharged being large, it is very pale. On the other hand, circumstances which diminish the proportion of water within the limits of health deepen the color. The complemental relation

of the skin and kidneys is well known. Under the influence of warmth, therefore, when the skin is acting freely, the quantity of urine is smaller, and it is darker, provided, of course, the quantity of water ingested is the same. In winter, the skin being less active, the quantity of urine is larger, and its color less deep. In persons from whom the respiratory exhalation is greater, the urine is likewise less abundant, darker, and vice versa.

Pathologically, the color of urine may be altered, first, by increase or diminution of the normal coloring matters, or, second, by the addition of abnormal ones.

The former is generally due to a change in the proportion of the coloring matters to the watery constituent. Thus we have almost an absence of color in the copious urines of diabetes, hysteria, and convulsions, while the urine of fevers and febrile states is high colored, chiefly because the quantity of water is diminished, but in the latter instance also because of the addition of an abnormal coloring matter known as urverythrin.

2. (a) The addition of abnormal coloring matters is seen in the instances just mentioned—fevers, in urines containing blood or blood-coloring matters, and bile pigment; and in the blue and brown urines of which instances have been reported.

(b) The urine is also colored after the ingestion of certain vegetable matters eliminated by the kidneys, as santonin, which imparts a yellow color.

IV. The reaction of normal mixed urine, that is, the urine of the entire twenty-four hours, is always acid. And, usually, specimens of urine passed at any time of day exhibit this reaction, though there is a difference in its degree,

while three or four hours after a meal the urine may become neutral or even alkaline.

The cause of this change in the reaction is still disputed. Roberts believes that it is due to an admixture with the blood of the elements of food, which are largely alkaline, and that the resulting increased alkalinity affects the reaction of the urine secreted. Bence Jones contended that it is the demand made on the blood for the elements of the acid gastric juice, which thus affects the reaction of the urine secreted during digestion. While neither explanation is altogether satisfactory, the former seems more likely to be correct. In health, urine passed on rising after the night's sleep is always acid.

The cause of the acid reaction of the urine is usually said to be acid sodic phosphate, though it is probably also slightly contributed by other acid constituents, as uric and hippuric acids, and, under certain circumstances, also by lactic and acetic acids.

There is often observed in urine which has been standing for a short time, especially at a moderate temperature, an increased degree of acidity, which sometimes results in a decomposition of urates, and a precipitation, first of acid urates, and later of uric acid crystals. This has been ascribed by Scherer to an acid fermentation, in which, the mucus acting as a ferment, lactic and acetic acids are formed by the decomposition of the coloring matters of the urine. This has not been satisfactorily proven, while the increased acidity is by no means constant.

It is certain, however, that acid urine which has stood for some time, and more rapidly in hot weather, acquires an ammoniacal odor, and becomes alkaline in its reaction; attending this change of reaction is a turbidity with a precipitation of a white amorphous and crystalline sediment, and often also with the formation of an iridescent pellicle on the surface. The cause of these changes has been well determined, and has already been alluded to. Through the action of mucus and other organic matters, acting in their decomposition as a ferment, the urea is converted into ammonium carbonate by the addition of two equivalents of water.

$$CH_4N_2O + 2H_2O = (NH_4)_2CO_3$$

which gives the odor of ammonia and the alkaline reac-


The turbidity and deposits are due to the precipitation of the crystalline triple phosphate of ammonium and magnesium, the amorphous phosphate of lime, urate of ammonium, and to living vegetable organisms known as bacteria.

The alkalinity thus resulting from the presence of volatile alkali (ammonia) is easily distinguished from that due to fixed alkali (potash or soda). On drying the test-paper, which has been rendered blue by volatile alkali, the blue color disappears, and the paper resumes its original red or violet tint; if due to fixed alkali the blue remains after desiccation.

V. The **specific gravity** of normal urine may be put down at 1020 for an average amount of 1500 c.c. (5002.) in the twenty-four hours. But as this quantity is by no means fixed, while that of solid matter remains about the same, the specific gravity must vary accordingly. When,

^{*}Normal urine may have an amphoteric reaction, i.e., changing blue litmus paper to red and red litmus blue, due to the simultaneous presence in the urine of acid and neutral phosphates.

from the action of cold or other cause, the skin is not acting, and after copious use of water and diuretics, the specific gravity may descend to 1010, and even lower, within the limits of health. But, when perspiration is copious, or a drain of water from the economy takes place through some other channel, the urine becomes



concentrated, and may be 1030 or higher in specific gravity.

Pathologically, the specific gravity of urine is increased or diminished, but to be quite reliable, observation should be made on the entire quantity passed in the twenty-four hours. The specific gravity is increased in diabetes mellitus,

where it sometimes reaches 1050. A specific gravity of suspicion of diabetes, and calls for sugar tests. It not safe to infer from a low specific gravity alone the absence more than 1028, if it attend a copious urine, should excite infrequently happens that urines with a specific gravity of 1010, and even lower, contain glucose, and it is not

amount of urea present; lower specific gravity of mixed The specific gravity is also increased in the first stage of the presence of albumin and sugar being eliminated, variain the first stage of acute Bright's disease, from the presence of blood, the higher specific gravity of the latter and the diminished secretion raising that of the mixed fluid. The specific gravity is lowered in hysterical and spasmodic forms of Bright's disease, except the stage of acute notic induration of the kidney, which often attends heart tions in the specific gravity point to variations in the acute fevers, in consequence of the increased amount of solid matters excreted and diminished water separation; and hydruria, though here it attends a proportionate increase of water, and is not of much practical significance. In all nephritis referred to, and in the condition known as cyadisease, there is a tendency to lowering of specific gravity owing to the diminished proportion of urea. Particularly is such reduction of specific gravity significant when it attends diminished secretion of urine. In a general way, urine generally means less urea.

than the picnometer (E, Fig. 2) and balance, it is still sufficiently so when carefully constructed. Every urinometer should first be tested with distilled water at 60° F. (15.54° meter is almost invariably used, and though less accurate To determine specific gravity the so-called urino-

C.), into which it should sink to the mark 0, or 1000. In its graduation the lines indicating the degree should gradually approach each other as the bulb is reached, because allowance must be made for the weight of the stem above water. The English-made urinometers, about five inches long (Fig. 2, C), are generally accurate, but the short German instruments (three inches) are very convenient for small quantities of urine. In the little urinometer of Heller (Fig. 3), in which the "sink"


ently altogether more carefully made, and I have generally "sink" contains mercury instead of shot. It is apparshorter than the original of Heller, in which the ordinary to sink, not to 1000, but to 1001+, proving its inaccuracy. specific gravity of a given specimen of urine 1004, that is, 1014, and so on. Especial care should be taken in testing seven of the ordinary scale. Thus, 1001 = 1007, 1002 = found it accurate. scale is retained on an ivory stem within the tube, and the Another urinometer is imported from Germany even slightly And on testing the former with distilled water it was found by a long-tried English instrument was found to be 1019. 1028 by the ordinary scale, of which the specific gravity Baumé is retained, where one degree corresponds with possession an instrument of this kind which recorded the large one by the ordinary scale. The writer has in his these instruments, as a slight variation in them indicates a consists of leaden shot, the graduation of

Dr. E. R. Squibb, of Brooklyn, N. Y., has recently suggested a small urinometer standardized for 25°C., or 77° F., a temperature much more usual than 60° F., and at

SPECIFIC GRAVITY.

Х.

three points, 1000, or 1030, and 1060, with the variations marked, whence corrections are easily made. The glass jar supplied with it is also fluted, and in this way the instrument is kept from clinging to the side of the glass. This object is further secured by making the air-chamber a double cone, base to base, as seen in a, Fig. 4, instead of a cylinder as in c. In this way, also, there is obtained



bb's Urinometer and Jar, shown also in section at b.

a single point of contact between the urinometer and the jar. Finally, the instrument is provided with a thermometer to secure greater accuracy, but as 4, if water be put at rooo, is about the maximum error which can occur at any temperature at which urine is likely to be tested, it is not really necessary. This is certainly the most accurate urinometer I have seen; and, by reason of the fluting of

the glass, a smaller amount of urine is required than in the ordinary cylindrical jar.

The cylindrical glass vessel usually supplied with the urinometer, or a sufficiently large test-tube, should be about three-fourths filled, the urinometer introduced, and when at rest the specific gravity read off. The cylinder or test-tube should not be too small in relation to the urinometer, lest, in consequence of the capillary attraction between the latter and the walls of the cylinder, the urinometer should not sink as low as it ought. For the same reason the urinometer should not be allowed to impinge against one side of the glass. All these difficulties are provided against in Squibb's urinometer. The scale should be read from above, not below the fluid.

number of volumes used. Hence, it is desirable to use low specific gravity. secured: while any error in reading is multiplied by the so nicely graduated that absolute accuracy in reading is be, if the data are, the urinometers in use are not usually the principle of this method is correct, and the results must of the urine will be $1000 + (4 \times 5) = 1020$. Although the specific gravity of the mixed fluid is 1004, then that to use the urinometer, that is, to make five volumes, and sary to add four times as much water as urine to enable us this method as rarely as possible, especially with urines of calculated that of the urine. volume added. From the sp. gr. of this mixture may be water to fill the cylinder to the required height, noting the fill the cylinder, it may be diluted with enough distilled If the quantity of urine be too small sufficiently to Thus: suppose it is neces-

VI. Quantity.—The average amount of urine in the twenty-four hours is usually put down at 1500 c.c., or

deranged conditions, it is the absence of this relation of color and specific gravity to quantity which gives ence that there is also much variation within the limits of increased quantity of urine. It is with regard to quantity that the complemental relation so well known to exist water separation by the latter, and vice versa. In about 50 fluidounces. I am inclined to think this is a little more than the average; perhaps it would be nearly correct to say between 40 and 50 ounces, or 1200 to 1500 c.c. But enough has already been said to allow the inferhealth. All that has been said of color and specific gravity in this respect is true of the quantity of urine, though in an inverse ratio. That is in health, diminished intensity of color and diminished specific gravity correspond with between the skin and kidneys most palpably shows itself, the increased activity of the former causing diminished significance to either.

pertrophy, in common with all conditions which cause increased blood-pressure, including the ingestion of large amounts of water, the peripheral action of cold, etc., there is an increase of water, and a corresponding reduction in Pathologically, the quantity of urine is increased in sive conditions; in the first with increased specific gravity, and in the remainder with diminished. In cardiac hydiabetes mellitus and insipidus, and hysterical and convul-

In all forms of Bright's disease, except the cirrhotic and lardaceous kidneys, there is a tendency to diminished secretion of urine. Toward the fatal termination, however, it is diminished even in these affections. Any marked diminution of urine in these diseases, particularly if of low specific gravity and color.

specific gravity, which means diminished urea, becomes a grave symptom.

In acute fevers and inflammatory affections, the quantity of urine is very constantly diminished until convalescence sets in, when there is generally observed a marked increase, which, in common with the profuse perspiration often observed at the same time, was long ago characterized by the word "critical."

VII. Odor.—Of the odor, little more can be said than that, in health, it is "peculiar" or "characteristic." It is by some spoken of as "aromatic." There is, however, appreciable difference in its intensity, as most have observed in their own cases. Concentrated urines always exhibit what is described in common language as a "strong odor." This is, undoubtedly, due to urea, though the characteristic odor of urine is not ascribed to urea, but rather to the minute quantities of phenylic, taurylic, and damoluric acids found in it.

Urine which has been standing exposed in warm weather acquires an odor which is at once putrescent and ammoniacal, the former from decomposition of mucus and other organic matters, the latter from the ammonium carbonate derived from the urea. The former is predominant when a large amount of organic matter is present, and is often observed in destructive disease of the kidney or its pelvis, and especially of the bladder.

The odor of urine is very promptly influenced by that of substances separated by the kidney from the blood, illustrated by the well-known odor of violets in the urine of persons taking turpentine. The odor of cubebs, copaiba, and sandal-wood oil is promptly communicated to the urine

of persons taking them. So, too, the use of certain vegetable foods promptly influences the odor of the urine. Among these asparagus is conspicuous.

Pathologically, except the increased intensity of the characteristic odor of concentrated urines, the putridity alluded to, and sweetish and fruity smell which often attends the presence of sugar in the urine, there seem to be no modifications of the "characteristic" odor of urine except in those extremely rare instances where sulphuretted hydrogen has been found in it.

TO DETERMINE THE AMOUNT OF SOLIDS IN THE TWENTY-FOUR HOURS' URINE.

Knowing the quantity of urine passed in the twenty-four hours, and its specific gravity, an approximation to the quantity of solid matters, and thence that of water, may be readily obtained by multiplying the last two figures of the specific gravity by the coefficient of Trapp—which is 2—or that of Hæser, 2.33. This will give approximately the number of grams of solid matters in the 1000 c.c. (33.8 f. 0z).

Thus, suppose the twenty-four hours' urine to be 1200 c.c. the specific gravity to be 1022, then, using Hæser's coefficient,

 $22 \times 2.33 = 51.26$ grams in 1000 c.c.

But the total quantity of urine in twenty-four hours is 1200 c. c., therefore it will contain more than 1000 c. c. contain. Hence,

1000: 1200:: 51.26: $x = \frac{51.26 \times 1200}{1000} = 61.51$ grms. (948.09 grs.)

Now, estimating the twenty-four hours' urine at 1500 c.c.,

the normal amount of solid matters is about 70 grams (1080.1 grs.), showing that, in this instance, rather less than the normal quantity was separated. In this manner, valuable information, bearing upon diagnosis and prognosis, may be obtained in a few seconds. The most striking variations are observed in diabetes and Bright's disease. In the former the solids are increased by the addition of sugar, in the latter they are diminished by loss of urea.

While this method of arriving at the solids is not sufficiently accurate for scientific uses, it answers for ordinary clinical purposes.

PART I.

THE DIFFERENT CONSTITUENTS OF URINE IN HEALTH AND DISEASE.

In the examination of a specimen of urine, the following are the steps which will be found most convenient in actual practice. Observe-

I. The quantity passed in twenty-four hours.

II. Color and transparency.

III. Odor.

IV. Reaction.

V. Specific gravity.

VI. Presence or absence of sediment, its quantity, and characters.

examination. The remaining or supernatant fluid, fillered if In all cases, whether the sediment be appreciable or not, a necessary, should then be further examined for certain organic vessel for twelve hours, in order to collect it for microscopical portion of the fluid should be set aside in a conical glass and inorganic constituents.

Organic Constituents.

VII. Presence or absence of albumin and other proteid substances.

VIII. Presence or absence of the different varieties of

IX. Other saccharine substances.

X. Presence or absence of acetone and diacetic acid.

XI. Coloring matters. { Normal. Abnormal.

XIII. Leucin and tyrosin. XII. The biliary acids,

XIV. Fatty matters.

XVI. Uric acid. XV. Urea.

XVII. Urates.

Inorganic Constituents.

XVIII. Chlorides.

XIX. Phosphates. $\begin{cases} a. \text{ Earthy phosphates.} \\ b. \text{ Alkaline} \end{cases}$ XX. Sulphates.

Examination of Sediment Microscopically and Chemically. I. Unorganized deposits, including crystals and amor-

phous deposits.

such as tube-casts, epithelium, pus, blood-corpuscles, etc. II. Organized deposits, including anatomical elements,

matter, extraneous substances, etc. III. Other morphological elements, as fungi, granular

tion than is involved in the consideration of the "general physical and chemical characters." Nos. I, II, III, IV, V, VI require no further explana-

Organic Constituents.

VII. ALBUMIN AND OTHER PROTEIDS.

Albumin.

It and serum-globulin are precipitated from their solutions The albumin, usually found in urine, is serum-albumin. at a temperature of 73° to 75° C. (163.4° to 167° F.). This is true of no other of the proteids which occur in urine. Other agencies, however, also throw down serumalbumin, and become equally delicate tests for it, although less reliable, because of their precipitating other substances.

In all instances, where the urine used for testing is not perfectly clear, it should be filtered before applying the tests. This may be done in a few minutes by means of filtering-paper and a funnel.

(a) The Test by Heat.

A test-tube is filled to 1/4 to 1/5 its depth with perfectly clear urine, and heat applied until boiling occurs. If a turbidity result, the slightest degree of which becomes visible in an otherwise clear urine held in a good light, it is due either to albumin or earthy phosphates.* If the latter, it promptly disappears on the addition of a few drops of nitric or acetic acid; if albumin, it is permanent. If the urine has not been filtered, and is opaque from the presence of amorphous urates, the first effect of the application of heat is to clear up the fluid, and, as the temperature is increased, albumin, if present, is precipitated.

I am satisfied, after long experience, that the more reliable method is to boil the urine first, and add the acid afterward, lest there be produced by the addition of the acid, acid, acid-albumin. Previous addition of acid may also precipitate mucin from normal urine. For it is to be remembered that, if a drop or two of nitric acid be added to

^{*}A very good way is to half fill the test-tube with clear urine, and apply the heat only to the upper part, when a resulting diminished transparency can be very easily recognized on comparing the two portions of the tube.

alkali-albumin. alkali the precipitate redissolves, being converted into neutralizing with liquor potassæ. With the least excess of its source, the acid-albumin may be readily precipitated on drop or two only should be added at first. Whatever be produce out of serum-albumin a soluble acid-albumin not the natural causes of its acidity. Acetic acid, also, may precipitable by heat, if too much acid is used. Hence, a same thing may happen when urine is very highly acid from to be present may fail to precipitate its albumin. The acid-albumin or syntonin, which is not coagulated by heat. boiled in a test-tube in which a drop of nitric acid happens In like manner and for the same reason, albuminous urine acid, it may happen on boiling the urine that no precipitate a specimen of albuminous urine so as to render it distinctly This is because the serum-albumin has been converted into whatever will appear, although much albumin is present.

It sometimes happens, also, when the precipitate obtained by boiling is albumin, that the addition of two or three drops of nitric acid will be followed by a partial disappearance of the turbidity, but, if a few more drops be added, the full amount is again thrown down. We should, therefore, continue the addition of the acid until 15 or 20 drops have been used. On the other hand, should the quantity of albumin be very small, too much acid will dissolve it.

It is also to be remembered that two or three drops of nitric acid may be added to a specimen of cold or boiled albuminous urine, and although a little cloud of albumin may follow the entrance of each drop into the urine, it will be quickly redissolved, and on the application of heat no precipitate may take place, or a small quantity only will fall. If now more acid be added, after a little time more

albumin will come down until all is precipitated. If nitric acid is used in this way it should be slowly added in considerable excess, yet short of an amount sufficient to dissolve the albumin.

It also occasionally happens, when acetic acid is added to urine naturally acid, and at the same time albuminous, that the albumin is at first only partially precipitated on the application of heat, there being a mere opalescence when the quantity of albumin may equal one-half the bulk of urine tested. After waiting a little while, however, the full amount of albumin is thrown down.

Furthermore, serum-albumin is converted by the continued action of an alkali into alkali-albumin, which is also not coagulated by heat. This may occur in highly alkaline urines. But the alkali-albumin is promptly converted into serum-albumin by the addition of a drop or two of dilute acid. Heat, under favorable conditions, will show one part of albumin to 100,000 of urine.

Mehu* calls attention to the fact that the urine charged with oxalate of lime becomes slightly turbid when heated, even after all the lime possible has been filtered out, and that the turbidity thus produced is not removed by the addition of a few drops of concentrated acetic acid. I have never encountered this source of error.


(b) The Nitric Acid Test.

This is best applied by the contact- or Heller's method. Upon a convenient quantity of pure, colorless nitric acid in a small test-tube (one of those with a foot, seen in Fig. 5, is most suitable), allow to trickle from a pipette down the side of the inclined glass an equal amount of clear

^{*} L'Urine, Normale et Pathologique, Paris, 1880, p. 326.

urine, which will thus overlie the acid. If albumin is present, there appears at the point of contact, between the urine and nitric acid, a sharp white band or some of varying thickness, according to the quantity of albumin present.

The urine may be put into the glass first, if preferred,



Testing for Albumin by Nitric Acid.

and the acid may then be allowed to pass down the side and under the urine. The result is the same, but the former is somewhat more easily practiced.

When nitric acid is allowed to underlie normal urine, there appears between the urine and the acid a brown ring

tated at the same place is similarly tinted. If there is much indican present in the urine, a rose-red or violet tint may be communicated to the albumin; if much bloodcoloring matter, a brownish-red, and if undecomposed action of the acid on the coloring matters. In consequence of this fact, when the urine is highly charged with coloring matters, as it often is in fever cases, the albumin precipiwhich grows in intensity on standing, and is due to the biliary coloring matters, a green hue.

tones or proteid bodies, while strong nitric acid does not precipitate mucin. Practically I do not think mucin is Nitric acid, in any strength, does not precipitate pep-

Nitric acid is said also to show one part of albumin in 100,000 of urine, but in my hands it is decidedly less delicate often a source of error. (See page 42). than heat properly used.

easily controlled; the process is further facilitated if the upper end of which interferes with the success of the test. This difficulty is readily between the middle finger and the thumb, whereby the flow may be the urine to flow from the pipette with sufficient slowness-that is, it Precautions,-I. Much difficulty is often experienced in causing will either not flow at all, or the finger, in the effort to cause it to flow, is raised so much as to permit a sudden fall of the urine into the acid, overcome by rotating the pipette, under the end of the index-finger,

is seen to diffuse itself into the urine above. Further, this layer, if caused by urates, is easily dissipated on the application of heat, although 2. A somewhat similar white zone is formed by the action of nitric acid on the mixed urates if present in excess, by which the more insoluble acid urates are thrown down. This zone might be mistaken for that of albumin, but the acid urates begin to appear not so much at the border between the urine and acid as higher up; nor does the zone on its upper surface remain so sharply defined, but while under examination some care is necessary in this application lest in ebullition the ring be the pipette is slightly roughened.

commingled with the entire mass of fluid and thus lost to view, although not actually dissolved. After some hours have elapsed these amorphous acid urates are completely decomposed by a further action of the nitric acid, and uric acid is then deposited as a characteristic crystalline sediment. Further difficulty arises where, as occasionally happens in very severe cases of fever, a small quantity of albumin coexists with an excess of acid urates. In these cases the urine is of high specific gravity, and the line of albumin, lying immediately on the acid, may be obscured by the broader band and cloud of urates. The difficulty from this source is diminished if the urine is diluted with two or three parts of water, while if the method laid down on page 52 is carefully carried out, a mistake is scarcely possible.

It should be added that Thudichum considers that the "cloud of acid urates here referred to is not urates, but hydrate of uric acid."

This "diffused haze" brought into view toward the upper part of the column of urine is regarded by Dr. Roberts † as mucin. Whether it be acid urates or hydrate of uric acid or mucin, its behavior is very different from that of albumin, which appears just above the line of junction of the two fluids.

3. This method of performing the nitric acid test operates equally well with serum-albumin, acid-albumin, and alkali-albumin, and therefore obviates the possibility of the source of error referred to on page 37, originally pointed out by Bence Jones—first, that, if albuminous urine be acidified by a small quantity of acid, as a drop or two, no precipitation of albumin takes place; also, that arising from the addition of too large a quantity, as an equal bulk, of acid, when the mixture may in like manner remain perfectly clear. Roberts says he has known the latter fallacy to cause the concealment of albumin in urine for months, in a case of Bright's disease.

4. Occasionally, also, it happens that a urine is so highly concentrated—so highly charged with urea—that the simple addition of nitric acid causes a precipitation of crystals of nitrate of urea at the junction

^{*}Thudichum, J. L. W., "Pathology of the Urine," 2d edition, London, 1877, p. 377.

[†] Roberts on "Tests for Albumin in Urine," Medical Chronicle, Oct., 884, p. 1.

of the two fluids. But these are readily distinguished from albumin by their solution on the application of heat, and by their appearance under the microscope, which exhibits them made up of six-sided rhombic tablets. Such urine is always of high specific gravity, while albuminous urine, except in cases of acute Bright's disease, is apt to be of low specific gravity.

lowing method recommended by Hofmann and Ultzmann must be potassæ, warm the mixture, and filter. If the filtrate is still not quite clear, add one or two drops of the magnesian fluid (11, p. 16), warm cence. Under ordinary circumstances, this does not interfere with the test; but if the quantity of carbonate of ammonium be very large, as is the case with some old urines, and that of albumin small, the effervescence is so great as to make the nitric acid test impossible; while the amount of acetic acid required to secure an acidity sufficient to permit the use of the heat test may be so great as to completely hold in solution the small quantity of albumin. Such difficulty is further increased by the fact that these alkaline urines are always more or less cloudy, from the presence of amorphous phosphates and of bacteria, and cannot be cleared up by ordinary filtration. Under these circumstances the folagain, and filter. The fluid is then always clear and transparent, and albumin, if present, may be revealed by Heller's nitric acid test, or by 5. If carbonic acid be abundantly present in urine, either free or combined with ammonium, as after the alkaline fermentation, or with sodium or potassium, during the administration of alkaline carbonates pursued. Add to the urine about a fourth part of its volume of liquor or salts of the vegetable acids, an added acid liberates it with effervesthe cautious addition of acetic acid.

 Occasionally, after the administration of turpentine or balsam copaibæ, resinous matters are found in the urine. These are precipitated by nitric acid in the shape of a yellowish-white cloud, which is, however, redissolved on the addition of alcohol.

Other Tests for Albumin.

It has long been known that other agents besides heat and nitric acid coagulate albumin. Some of these have lately claimed a large amount of attention, and been found

to be extremely delicate tests; some more delicate even than the heat test applied in the most careful manner, and very much more delicate than the nitric acid test. An objection, however, which holds against the most delicate of the reagents is, that they precipitate other substances besides albumin, and although these substances are generally distinguishable from albumin by the aid of certain precautions or additional steps, it dare not be said that the reliability of the tests is not weakened thereby.

As the result of a careful study of these tests, based upon experiment and clinical application, I have come to the conclusion that, for the present at least, it is safer to rely upon them in conjunction with the heat and acid tests, with a view to confirming or extending results attained by the latter. Such use is, however, of the greatest importance; and I shall, therefore, now consider the most delicate of them, including the method of their application, and the precautions to be observed in rendering them reliable. Those I deem most worthy of consideration are in the order in which I have found them most delicate: Picric acid, so-dium tungstate with citric acid, potassio-mercuric iodide,*

^{*}The first three named of these agents are so nearly equal in delicacy that one arranges them in any order at some risk, and I must confess to not invariably having found them delicate in the order named. The experiments of Dr. George Oliver, of London, show that any one of these used in solution by the contact method, will precipitate I part of albumin in 20,000 of urine. Dr. Henry B. Millard's experiments go to show that Tanret's test will show very easily I part of albumin in 20,000 parts of urine; that his own test of phenic acid and potash will show the same even more easily; that the nitric-magnesian test will exhibit I in 150,000, and that heat and nitric acid will each show I part in 100,000, heat being more sensitive than nitric acid. (Millard, "Bright's Disease," 3d ed., 1892, p. 85.)

45

ferrocyanide of potassium, and Dr. Roberts's nitric-magnesian solution. Picric Acid.-This has its ablest and most enthusiastic exponent in stance occurring in either normal or abnormal urine, except albumin, which gives a precipitate with picric acid insoluble by the subsequent Dr. George Johnson, of London, who says, "there is no known subapplication of heat." *

acid. A saturated solution may be made by dissolving 6 or 7 grains of the powder in an ounce of boiling distilled or rain water. A portion An ounce of water at 60° F. retains in solution 5.3 grains of the dry of the acid will crystallize out on cooling, leaving a transparent yellow supernatant liquid. Such a solution has a specific gravity of 1005.

the turbid column increases it. Then, if the tube be placed in a stand position, pour gently an inch of the picric acid solution on the surface of the urine, where, in consequence of its low specific gravity, it mixes only with the upper layer of the urine. As far as the yellow color of the picric acid solution extends, the coagulated albumin renders the liquid turbid, contrasting with the transparent urine below. For the action of the test, there must be an actual mixture, and not a mere surthe turbidity is very slight, the application of heat to the upper part of the coagulated albumin will gradually subside, and, in the course of an hour or so, forms a delicate, horizontal film at the junction of the colpour a four-inch column of urine; then, holding the tube in a slanting face contact. When, in consequence of the scantiness of the albumin, Dr. Johnson's mode of applying the test for the detection of a very minute trace of albumin is as follows: Into a test tube six inches long ored and unstained strata of urine.

No previous acidulation of the urine is usually required, as the picric acid accomplishes this, if needed. If, however, a specimen be highly alkaline and ammoniacal, Dr. Johnson says the safest method is by acetic or citric acid; then filter and add picric acid to the filtrate. The precipitated mucin will remain on the filter.

Precautions .- I. The urine to be tested should be perfectly clear,

^{*} Johnson, "Albumin and Sugar Testing," London, 1884, p. 11.

and if not clear when obtained should be rendered so by filtration, or the processes described on p. 42.

2. Urates, peptones, vegetable alkaloids, as quinine, morphia, etc., are all precipitated by picric acid solution at the point of contact but are promptly redissolved by a degree of heat much lower than that of the boiling-point. Quinine promptly appears in the urine after the administration of ten grains of this drug.

Dr. Oliver correctly claims that the addition of citric acid in the proportion of two drachms to an ounce of the picric solution makes a reagent which gives a more distinct reaction than the plain picric acid solution; but Dr. Johnson says that this is because the citric acid precipitates also the *mucin* which exists in all urines.

Dr. Johnson also insists that mucin is not precipitated from urine by picric acid, and that albumin is the only substance found in the urine which gives with picric acid an opalescence or precipitate insoluble in heat. He is not sustained in this view by others, notably by Roberts,*

Oliver,† and Millard.‡

Sodium Tungstate with Citric Acid.—This solution is made by mixing equal parts of a saturated solution of sodium tungstate (1 to 4) and a saturated solution of citric acid (10 to 6). It has a specific gravity of 1214, and is best used by the overlaying method. It is a test of extreme delicacy, quite as delicate in my hands as picric acid, and has the advantage over picric acid of not precipitating quinine from its solution, but, like the picric acid, precipitates acid urates, peptones, and mucin, which are also promptly dissipated by heat.

The Potassio-Mercuric Iodide.—This test was suggested by M. Charles Tanret, of Paris, and is regarded by Dr. Oliver as the most sensitive test known, discovering, like the picric acid and sodium tungstate, I part of albumin in 20,000 of urine, while Millard says it will recognize I part in 200,000 of urine. In my own experiments, however, I have several times failed with the mercuric iodide when I succeeded both with picric acid and sodium tungstate. I am inclined to believe

^{*} Loc. cit., p. 3.

⁺On "Bedside Urine Testing," 3d Ed., London, 1885, p. 111, note.

[‡] Op. citat., p. 63.

that the age of the preparation and the mode of compounding it have something to do with the results.

The solution is prepared by M. Tanret as follows: Bichloride of mercury, 1.35 grams; iodide of potassium, 3.32 grams; acetic acid, 20 cubic centimetres; distilled water, enough to make 100 cubic centimetres. The bichloride of mercury and iodide of potassium should be separately dissolved in portions of the water, which are then united. The resulting reagent is the double iodide of mercury and potassium, the chloride of potassium being without effect. It is also a heavy fluid, having a specific gravity of 1040 +, and is used by the contact method. The urine requires no previous acidulation.

It coagulates the same substances as picric acid, which are likewise dissipated by heat, or alcohol. On cooling they reappear. With regard to mucin, however, Dr. Oliver says it is not dissipated by heat if a large excess of the reagent be employed, the mercuric salt apparently preventing solution.

Tanret also suggests a quantitative method, which is, however, not

accurate.*

Dr. W. G. Eggleston, of Chicago, suggests, with a view to avoiding the confusion of the mucin reaction, that the potassio-mercuric solution be made without acetic acid, and that the urine be acidulated previous to the testing. If a precipitate results, and we have reason to think that it is mucin, half the acidulated urine may be placed in another tube and the mercuric iodide solution used. If this causes a further precipitate, as shown by comparison, we may be reasonably sure that albumin

is present.

Dr. Eggleston also thinks that, when the solution is made up with acetic acid or any acid, some chemical action must take place sconer or later, as iodide of potassium and bichloride of mercury are comparatively unstable compounds in the presence of an acid.

Ferrocyanide of Potassium.—This test, used in saturated solution, while less delicate than any of the three previously considered, has the advantage over them of not precipitating mucin, peptones, or

^{*} Recherche de l'albumine dans l'urine; Bulletin Général de Thérapeutique, tom. 92, 1887, p. 308.

the alkaloids, but it may throw down acid urates.* It is fully as delicate as nitric acid, but less so than heat. It requires for its operation that the urine shall be acid.

Dr. Roberts's Acidulated Brine Solution.—This solution, which consists of a pint of a saturated solution of common salt to which is added an ounce of hydrochloric acid, and the whole filtered, is about equal in delicacy to nitric acid, but much less so than heat.

It has a high specific gravity, and is used by the contact method. It has the great advantage over nitric acid in that it is less caustic and corrosive, and therefore much pleasanter to work with, but recent experiments have shown it to be fallacious, and it is now, I believe, regarded by Roberts himself as unsatisfactory.

is among the most satisfactory tests he has used, detecting less than I tact method. It cannot be used with urine clarified by liquor potassæ, magnesian fluid has a specific gravity of 1240, and is used by the conof phenic and acetic acids and Tanret's mercuric iodide. The nitric dent. It has received the highest praise from Dr. Henry B. Millard, who a watery, clear solution, which does not fume, nor stain, nor burn the albumin with nitric acid. The test fluid consists of I volume of strong part in 150,000 of water, but he regards as more sensitive his own test says that, as regards delicacy, accuracy, and facility of employment, it the urine, and may be carried in a corked bottle with less risk of accifingers, acts less strongly than the pure acid on the coloring matter of reaction in regard to albumin, mucin, and peptone is similar. It forms nitric acid and 5 of a saturated solution of sulphate of magnesium. urine with magnesium sulphate, filtering and then testing the liquid for suggested by C. Gerhardt in 1856, which consisted in saturating the another modification of the nitric acid test, really a modification of a test This, he says, is more prompt and sensitive than pure nitric acid, and its Roberts's Nitric Magnesian Test .- Dr. Roberts has suggested †

^{*} Dr. George Johnson claims that it, in common with all tests which require the combination or addition of acetic or citric acid, does give with mucin of normal urine an opalescence or haziness not distinguishable from that caused by a minute trace of albumin.

[†] Loc. cit., p. 1.

on account of decomposition induced. My own experience with it is

limited.

Millard's Phenic Acetic Acid Test.—This consists of glacial acid, phenic, (95 per cent.) 5 ij; ac. acet. pur. 3 vij. Mix, and add liquor potasse, 3 ij, 3 vj; then filter. Millard claims the test is equally delicate with Tanret's and the picric acid test, and, like them, precipitates mucin, peptone, and the alkaloids, and requires the same precautions. Millard also claims that it will detect 1 part albumin in 200,000 more clearly than Tanret's double iodide of mercury and potassium test, as it precipitates fewer of the alkaloids.

Trichlor-acetic Acid.—This reagent is regarded by many as a delicate test, and esteemed accordingly. A saturated solution (\$\tilde{5}\$ ss to \$\tilde{5}\$ jo water) of the crystals is made, and it is used by the contact method. It is to be remembered that acetic acid precipitates mucin, and it is not unlikely that the delicate white line sometimes obtained by this reagent when other tests give negative results may be mucin.

Albumin Test-Papers.—All of these reagents, except the acid brine solution, may be used in the shape of the test-papers suggested by Dr. Oliver, which are more especially useful for bedside testing, although Dr. Oliver claims for them some advantages over the solutions even when used in the laboratory.*

Most recently, † Dr. Oliver has rejected all of the albumin papers except the "mercuric" and "ferrocyanic," and recommends these to be used as follows:—

To Use the Papers.—A mercuric or ferrocyanic, and a citric acid paper are dropped into the test-tube, and water added to 60 minims. After gentle agitation for half

^{*} The different forms of test-papers suggested by Dr. Oliver, originally made for him by Wilson & Son, of Harrowgate, London, are now furnished by Parke, Davis & Co., of Detroit, Mich., as are also, in one case with the test papers, the suitably graduated test-tubes.

^{† &}quot;Bedside Urine Testing." London, 1885.

a minute or so, the test-papers are removed, and the transparent solution is ready for testing.

The pipette containing the suspected urine is held in a vertical position over the tube, and the urine is delivered in drops. If four drops of urine added to the mercuric solution, and six to the ferrocyanic, * do not produce a trace of milkiness when the contents of the tube are viewed against a dark background, it may safely be inferred that if albumin is present it is in so small a quantity that nitric acid, applied after the contact method for one minute, will not discover it. If a slight milkiness is apparent, it will represent a trace of albumin detectable by nitric acid.

If there is no response, the dropping should be continued, and, if instead of four drops of the mercuric and six of the ferrocyanic solution there be required 10 of the former and 15 of the latter to produce an appreciable opacity, it will indicate a quantity which can readily be shown by heat and acidulation. If 20 of the former and 30 of the latter are required, it indicates a trace of albumin which can be shown only by careful acidulation and subsequent boiling.

In the case of the mercuric test, if a reaction occurs, the solution should be boiled so as to prove the presence or absence of one of the diffusible proteids, peptone, or hemialbumose. If the opacity is unaffected by heat, or is intensified by it, it is caused by albumin, but if it is diminished or entirely removed, presumptive evidence is afforded of the presence of a peptoid body, either alone or in conjunction with albumin. The mercuric test, supple-

^{*} The ferrocyanic solution should be allowed a minute in which to develop the reaction from a trace of albumin.

21

mented by heat, may be said, therefore, to provide a fuller knowledge of the proteids which may appear in the urine than the ferrocyanic, which precipitates albumin only.

Remarks on Testing for Small Quantities of Albumin. The Author's Method.

To determine the presence of albumin in urine when it is abundantly present is usually a very simple matter. The application of heat will throw down albumin even from an alkaline solution, if the latter is highly charged with it, while the addition of a few drops of acid removes all possibility of error. But it is well known that small quantities of albumin, the significance of which in diagnosis and prognosis is sometimes greater than that of large amounts, often escape detection; while large quantities are sometimes obscured in consequence of peculiarities of combination between albumin and acids, and albumin and alkalies, resulting in the formation of the so-called acid- and alkalialbumins. It is with a view to pointing out the way to avoid such errors that the following paragraphs are written.

Under all ordinary circumstances by far the most striking test for small quantities of albumin is that form of the nitric acid test described as the contact method (p. 39), and in the majority of cases, this test, carefully carried out, even in the hands of the inexperienced, will exhibit the presence of albumin when it would have been overlooked in the ordinary mode of application of heat and nitric acid. But it is not so delicate a test as the latter applied in the manner to be described.*

^{*} See a paper by the author, "Notes on Albuminuria," in the Proceedings of the Medical Society of the State of Pennsylvania for 1881,

and albumin are both amorphous and cannot therefore be unusual proportion of urea. The second and third have acid upon the acid urates, and a resulting complete separaa sediment is present, should be boiled again; and if the which, after cooling and standing from six to twelve hours, thus distinguished. All, however, except albumin, disapeasily recognized under the microscope, but the acid urates well-known forms of crystallization by which they can be which it was combined; the third is found only when the tion of the uric acid from the sodium, potassium, etc., with arise from a partial decomposition of the neutral urates by carefully filtered, this deposit is either, 1st, acid urates; 2d, sediment is not dissolved after such ebullition, it is albufore, urine which has been tried by heat and nitric acid, in pear on the reapplication of heat. In all instances, thereurine happens to be highly concentrated and contains an the nitric acid added; the second by a further action of the uric acid; 3d, nitrate of urea; 4th, albumin. The first aside the urine thus treated, say for twelve hours, or until resulting fluid will be apparently clear, but upon setting clear urine and adding a few drops of nitric acid, the heat and acid test will have observed that after boiling the the next morning, there will sometimes be found a small Many who have tested urine for albumin by the ordinary Supposing the urine before testing to have been

Author's Method.—My own method, therefore, of examining a specimen of urine for albumin is invariably as follows:—

I. Unless *perfectly* clear, it is first filtered, and if not rendered clear by filtration, it is clarified by strong alkalies, or the magnesian fluid, according to the directions on

presence of albumin, the tube must be set away, carefully here-although it is well to put the tube aside, in order that the albumin may subside and be approximately estimated. If, however, there is the least doubt about the protected from dust, for six to twelve hours, in order that any appreciable sediment may subside, and be subsebut not more, lest a small amount of albumin present be remains it is caused by albumin, and the test may end redissolved by the excess of acid. If any degee of turbidity the least diminution of transparency. A drop or two of nitric acid is then added, and if a turbidity which has ensued upon the action of the heat disappears, it is caused by phosphates of lime and magnesium, and not albumin. The addition of the nitric acid should be cautiously conbeing carefully watched in a good light for detection of page 43. A portion of the filtered fluid is then boiled, tinued until a decided excess is added-15 to 20 drops-

quently again tried with heat.

II. A test-tube is now filled to the depth of half an inch with Roberts's nitric magnesian solution or colorless nitric acid. About as much urine is then allowed to fall gently upon it in the manner described on page 39, and the point of junction of the two fluids-carefully examined for the white line. This is best observed by holding the tube against a dark ground, produced by a book, pamphlet, or coat sleeve, so that the light may fall obliquely upon the line of junction of the two fluids, while at the same time it is seen against the dark ground.

When this double test is carefully applied as above described, it is scarcely possible to err as to the presence of albumin. Where it is abundantly present, it is, of course, unnecessary to use either the modified heat and acid test

or the contact method, although the latter is always useful in that it affords one means of approximately estimating the amount of albumin.*

Quantitative Estimation of Albumin.

Gravimetric Method.—It is a matter of extreme importance in the course of Bright's disease that we should be able to compare the quantity of albumin contained in

*The following observation by Dr. C. E. Brown-Sequard, in the first number of his Archives of Scientific and Practical Medicine (1873), illustrates some of the difficulties occasionally encountered in this ordinarily very simple process of testing for albumin: "If we first test, by heat, urine containing albumin (after having ascertained that it is naturally acid), we may not find the least precipitate; and if we add nitric acid to it after it has boiled and become somewhat cold, we may yet not find precipitation of albumin. But if we boil a second time that now acidified urine, the solidification of albumin quickly takes place, and a precipitate soon appears."

He further says: "In three cases in which the microscope showed tubular casts in the urine, the albumin contained by this fluid was so modified by the heat that if the urine (which was naturally acid) was boiled first, the addition of nitric acid in small or in large quantity, at a low temperature or at the degree of boiling, produced no solidification of that proteid substance. But when I added either a small or a large quantity of nitric acid to the fresh unboiled urine and then boiled it, the ordinary coagulation took place, and after some time of rest, the ordinary precipitate appeared. It is evident, therefore, that there is sometimes in the urine a kind of albumin which loses its coagulability by boiling."

The lesson from these facts is that it would seem necessary to apply the heat and acid tests both ways, that is, the acid should first be added to the urine and the mixture then boiled, as well as that the urine should be first boiled and the acid then added. I believe, however, if the method above described is carefully carried out, albumin cannot be overlooked.

the urine from day to day. The only accurate method is by precipitation by acetic acid and boiling, separation by filtration, drying, and weighing by delicately accurate balances, the weight of the filter having been previously determined. This, however, involves too much time for the busy practitioner, and we must fall back on one Fig. 6.

Approximative methods.

Approximate Estimation with Boiling.—
The easiest of these is to boil a given quantity of urine in a test-tube, add a few drops of nitric acid, and set aside for at least twelve hours—shaking the urine once or twice in this period in order to secure a uniform subdivision and precipitation of the particles of albumin. The proportion of bulk occupied—one-fourth, one-eighth, a trace, etc.—is used to indicate the quantity of albumin. Greater accuracy is obtained by previously filtering the urine of urates, epithelium, or extraneous matter, which might unduly increase the bulk of deposit on standing. Nothing more

Estimation with Esbach's AlbuminomeEstimation with Esbach's Albuminomeglass tube like that shown in Figure 6. To use
it, fill with urine to U, and to R with the test
solution, consisting of picric acid, 10 grams; Esbach's
citric acid, 20 grams, and water sufficient to Albuminomake a litre. Close the tube by a rubber stopmake a litre contents cautiously, allow the mixture to
per, and mix the contents cautiously, allow the mixture to
remain undisturbed for 24 hours, and then read off the
quantity of precipitated albumin. Each of the main

lines to which the precipitate reaches indicates one gram of albumin in one litre of urine.

Approximate Estimation with Nitric Acid.—The following method of approximate quantitative estimation by means of Heller's nitric acid method is given by Hofmann and Ultzmann. According to them, if the white zone of albumin has the depth of from 2 to 3 mm. (1 to \frac{1}{2} to \frac{1}{2} inch), is delicate and faintly white in color, has no granular appearance, and appears clearly defined only when placed against a dark background, the quantity is less than \frac{1}{2} per cent., usually \frac{1}{10} per cent. If the zone is 4 to 6 mm. (\frac{1}{2} to \frac{1}{2} inch) in depth, granular, white, opaque, and perceptible without a dark background, the quantity is considerable, \frac{1}{4} to \frac{1}{2} per cent. If, however, the zone of albumin appears granular and flocculent, and sinks in more or less lumpy masses to the bottom, and when by stirring the albumin by means of a glass rod the mixture assumes the consistence and appearance of sour cream, then the quantity is very large, 1 to 2 per cent.

cautiously added, 10 minims at a time, until the opacity of the mixture corresponds exactly with that of the standard solution. A comparison tion, as determined by placing the ruled card behind each, water is to be the card. If the opacity is still greater than that of the standard solusay to 60 minims in all, the tube again shaken, and the test made with such as to obscure the lines completely, water is to be freely added, are poured 20 minims of urine, and a potassio-mercuric iodide testa flattened test-tube, graduated to 200 minims, in 10 minim divisions, lines of various degrees of thickness are printed. If the opacity is resulting turbidity is tested by placing behind the tube a card on which paper is dropped into the tube. It is then well shaken, and the sealed tubes of the required size. The test is applied as follows: Into therefore represents To per cent. albumin. It is placed in hermetically standard solution made with alumina precipitated by ammonia, which I per cent, of albumin. This degree of turbidity is imitated by a by coagulating the albumin from a standard solution containing To of and comparing the resulting diminished transparency with that produced the albumin from a given quantity of urine by a mercuric test-paper, that recently suggested by Dr. Oliver. It consists in precipitating all Oliver's Approximate Method .- A better quantitative method is between the known value of the standard solution and the number of times the urine has been diluted, furnishes the proportions of albumin. Thus the value of the standard opacity being 10 per cent,, five dilutions would be .5 per cent., six, .6 per cent., and so on.

When the lines on the card can be read at once, without any dilution, the quantity of albumin is below 10 per cent.

The Proportion of Albumin Found in Urine. -

which urine can contain, since blood-serum only contains large one A half per cent, is much more common, and many albuminuric urines contain much less than a half per cation given that this is what is meant. In point of fact 3 or 4 per cent, is probably the maximum amount of albumin about 5 per cent., and 2 per cent. albuminuria is a very There is much carelessness of expression among physicians in speaking of the quantity of albumin found in a given specimen of urine. Thus we often read that a specimen contains 25 per cent., or even 50 per cent., of albumin. The proportion in bulk is of course intended, but no indi-

little more than three-fourths of an ounce. Of such a loss Senator well says that half a pound of beef will more than which is a very large amount of albumin, and 40 ounces of urine. Then $455.7 \times .02 \times 40 = 364.5$ grains, or It is not unusual, also, to over-estimate the amount of albumin passed in the 24 hours, and thence to exaggerate the drain upon the system. Suppose, for example, the percentage of albumin by weight is .5 of 1 per cent., and the quantity of urine is 50 ounces in the 24 hours, then, 50 = 113.9 grains, the amount of daily discharge, or about one-quarter of an ounce. Supposing there be 2 per cent., there being 455.7 grains in an ounce, 455.7 × .005 × make up the loss of a week. cent. of the proteid.

Other Proteids Found in Urine.

It has long been recognized that modifications of albumin occur in urine, either alone or in association with it, but since the introduction of the delicate tests, more attention has been paid the subject. Among the most constant of these is:—

Serum-globulin, Globulin, or Paraglobulin.—Globulin is almost always associated with serum-albumin, from which it may be separated by Pohl's method as follows: Render the urine slightly alkaline by ammonium hydrate, and after several hours filter to separate the phosphates. Then add saturated solution of ammonium sulphate in the proportion of one volume to one volume of the filtrate. If a precipitate forms it is globulin.

Globulin is also separated by diluting the urine, after filtration, until the specific gravity is 1003 or 1002. Occasionally a cloudiness appears immediately, due to the separation of some of the globulin. But it is completely separated by passing a stream of carbonic acid through the dilute fluid, for from 2 to 4 hours. In from 24 to 48 hours the globulin falls to the bottom as a milk-white flocculent, substance. The supernatant fluid contains the albumin. Should the urine be neutral or alkaline, it must be rendered slightly acid by a few drops of dilute acetic acid.

This test is based upon the fact, that globulin is held in solution by the sodium chloride, and other neutral salts always present in the urine. When urines are largely diluted with moderately pure water, the percentage of neutral salts is so reduced that the globulin falls out of solution.

Dr. Roberts * suggests the following simple modification of the test: Fill a urine-glass or test-tube with water, and let fall into it a succession of drops of albuminous urine. In many cases, each drop as it falls is followed by a milky train, and when a sufficient number of drops has been added, the water assumes throughout an opalescent appearance, as if a few drops of milk had been added to it. The addition of acetic acid causes the opalescence to disappear; for globulin is soluble in concentrated acetic acid, and also in a one per cent. solution of hydrochloric acid. From its solution in common salt it is completely separated on

Occurrence.—Not only does globulin accompany scrum-albumin in most instances, but, it is said, sometimes exceeds it decidedly, although the proportion in the blood is much less, being to serum-albumin as 1 to 1.5. It may even occur, although very rarely, without serum-albumin. According to Senator, it is most abundant in the urine of lardaceous disease of the kidney. It also occurs in acute nephritis, in the hyperæmia of cantharides poisoning, and in albuminuria associated with deranged digestion.

Mucin.—This proteid, abundant in urine which has passed over irritated urinary passages, is said to be present, to some extent, in all urines. It is not precipitated from solutions by boiling, but is precipitated by alcohol, by dilute mineral acids, and by all vegetable acids except, according to Dr. Johnson, picric acid. But Dr. Oliver does not even except picric acid, and fears that in all of

^{*} Discussion on Albuminuria before the Glasgow Pathological and Clinical Society, p. 17. Reprinted from the Glasgow Medical Journal,

To Test for Mucin.—The tests usually employed for mucin are citric and acetic acids used by the contact method, the acid being introduced first. It is also precipitated by dilute nitric acid. Just above the point of contact a cloud-like coagulum gradually makes its appearance, contrasting with the opaque white coagulum of albumin. When albumin and mucin are both present, the latter appears at the upper part of the column of urine, while the albumin is confined to the point of contact between the two fluids. It differs from the somewhat similar white deposit of acid urates in that it is not dissipated by heat. Or the following method may be followed:—

To one volume of urine add three of strong alcohol and allow to stand for several hours, when mucin and all albuminous bodies will have precipitated. Filter and wash the precipitate with alcohol, then treat it with warm water. The filtrate, which should contain the mucin, is then strongly acidified with acetic acid, and if turbidity results, it is due to mucin. Or the contact method may be used with acetic or citric acids.

The Mucin Reaction of Normal Urine.—Dr. Oliver claims that even when a normal urine is underlaid by citric acid solution, there appears in the course of several minutes, along the plane of contact of the two fluids, a delicate whitish zone which becomes gradually more pronounced, and which is mucin. This reaction, concentrated by the contact method, becomes, when the precipitate is diffused throughout the

^{*&}quot;On Bedside Urine Testing." Third edition, London, 1885, p. 11, note.

61

urine, either totally inappreciable, or appreciable only in the slightest degree. It may be studied by adding a citric acid test-paper to 60 minims of transparent urine. If mucin is present in larger quantity than usual,

a slight milkiness appears.

If, however, to the normal urine, acidified by citric acid, a mercuric test-paper be added, a delicate haze may be detected on holding the urine to the light, and against a dark background. This haziness, according to Dr. Oliver, disappears on applying heat to near the boiling point, and reappears on cooling, to revanish on reheating. If, on the other, hand, a solution of the potassio-mercuric iodide is used by the contact method, the opacity thus produced by mucin does not disappear when heat is applied.* If these observations are correct, heat removes all sources of error in testing for albumin with the potassio-mercuric iodide, when used in the shape of test-paper. This, therefore, Dr. Oliver regards as possessing a distinct clinical advantage over the solution.

heated. The opacity returns as the temperature of the solution falls, and in the cold it greatly exceeds the original amount. Heat will again disperse it as before. If a trace of albumin be added to the mucincharged urine, the test-papers will produce an opacity which heat will clear up only to a certain degree, that which remains over being due This highly muciparous solution is added to albumin-free urine, I to I In any case the urine will then become more highly muciparous than is likely to be met with in practice. Filtration may be omitted, being slow, if observation is checked by comparing the unheated fluid with that experimented with. A citric acid and a mercuric test-paper, added to 60 minims, produces an opacity exactly like that produced by a small quantity of albumin, but it differs from it in completely vanishing when * The student is strongly recommended to study the reactions of should be mixed in equal parts, and one drop of acetic acid, or a citric acid test-paper, added to a 4-inch column, which should be thoroughly boiled, when the milkiness produced by a trace of albumin will appear. mucin by impregnating normal urine with saliva as directed by Dr. Oliver. The clear saliva and a solution of salt, say 20 grains to the ounce, or I to 2 according as the observer may wish to charge it with mucin. to albumin.

Peptone.—The frequent occurrence of peptone in urine has rendered it one of the most important of the possible constitutents demanding attention. Much has been lately added to our knowledge of peptonuria, but it is possible that further changes will be made in such knowledge before it is complete.

cal significance of peptonuria." tuberculosis and other diseases named below under "clinirative affections, phosphorus poisoning and typhus fever, are introduced in various morbid states, especially suppube secreted by the kidneys from the blood, into which they duced in the urine. They may doubtless, however, also example the pyogenic organisms, that they may be probacterial organisms, and it is through these agencies, as for mose. These substances may likewise be the product of product of neutralization, and propeptone or hemialbuwhich the most important are syntonin (acid albumin) the intermediate bodies collectively known as albumoses, of of the process of albuminous digestion there are certain Between albumin at the beginning and peptone at the end continued operation of a temperature of 268° to 290° F. decomposing action of bacteria, as well as the longaction of acids and alkalies, and it is said, also, by the It may also be produced from albumin by the continued which is the final result of gastric and pancreatic digestion. It is well known that peptone is a proteid substance

Peptone differs from albumin and propeptone in that it is *nôt* precipitated by heat or nitric acid, by acetic acid combined with chloride of sodium, or with cyanide of potassium. Like albumin and propeptone, it is precipitated by tannin, corrosive sublimate, phosphortungstic

acid, Millon's reagent, and by sodium tungstate, potassiomercuric iodide and picric acid, the last three being among the recently suggested delicate tests for albumin; but when precipitated by these three reagents it is redissolved by heating. It is further characterized by the purple reaction its solutions strike with sodic hydrate and a little salt of copper. The sensitiveness of this so-called "biuret" reaction is increased by slight coloration of the fluid, and especially a yellow. So delicate is it that a solution as weak as 1 part in 1000 responds promptly. This reaction is shared by propeptone, while albumin produces only a blue coloration—under no circumstances a red or violet. It has been suggested that peptone bears the same relation to albumin as grape sugar to starch—that is, that it is a

into the circulation, in the very act of absorption, and is although the reverse has been alleged by Plosz, Maly, and Adamkiewicz. During health, too, the various organs of These organs also appear to retain any small excess which may have been introduced into the blood from over-ingestion of albuminous food. It would seem, therefore, that it is converted into albumin at the moment it is absorbed incapable of taking the place of albumin in the blood, the body contain traces of it, and it is a constant contraces of it. Injected into the blood, it disappears from it very quickly, a certain proportion reappearing in the urine, Peptone is not present in healthy blood or normal urine, and even during digestion the portal vein contains but while some is taken up by various organs of the body. stituent of pus, as was first shown by Hofmeister. hydrate of albumin.

Tests for Peptone.—The best tests for peptone are those which require some preliminary treatment of urine

suspected to contain it. For these we are chiefly indebted to Hofmeister.* The simplest is the

Phosphor-Tungstate Test.—(1) The urine is decolorized and freed from mucin and phosphates by treating say half a litre or about a pint with solution of neutral acetate of lead until the thick flocculent precipitate ceases to form. It is then filtered.

(2) To a portion of the filtrate add acetic acid and a few drops of a solution of ferrocyanide of potassium. Should there be a cloudiness or precipitate it is due to albumin, and the addition should be continued as long as a precipitate occurs. The albumin must be removed by filtration.

(3) Add to a portion of the filtrate above to wintation. bulk of concentrated acetic acid, and then phosphortungstic acid acidulated with acetic acid, † If the fluid remains clear after standing some time, it does not contain peptone. Should, however, after the lapse of 10 minutes, a cloudiness appear, peptone is present.

The Biuret Test.—Treat the urine with acetate of lead, filter, and add to the entire filtrate concentrated solution of tannic acid, so long as a precipitate takes place. Throw the latter, after 24 hours, on a filter, and wash it with water in which tannic acid and magnesium sulphate are dissolved. The precipitate is now thoroughly rubbed up in a capsule with saturated baryta water, and after the addition of a fragment of solid baryta, retained at a boiling temperature for a few minutes. If care

^{*}Neubauer and Vogel, 8th ed., 1881, p. 138, and Zeitschrift für Physiol. Chemic, 5, 73.

[†] The solution of phosphortungstic acid is made by adding to a boiling watery solution of tungstate of sodium, enough phosphoric acid to produce an acid reaction. After cooling, the fluid is to be made strongly acid with acetic or hydrochloric acid, and, after standing one day filtered.

is not taken to mix intimately the precipitate with the baryta, resinous masses form during the heating, which interfere with the proper action of the baryta. After a few minutes the mixture is filtered, baryta water again added, and thoroughly shaken with it until the fluid filters off from the dark-colored precipitate, either colorless or slightly yellow.

In the filtrate thus obtained the "liuret" reaction is now sought by adding first liquor sode or potasse until a decided alkalinity is produced, and then, drop by drop, a very weak solution of sulphate of copper. If a reddish color appears, the addition of the copper solution is to be continued until the reddish-violet has reached its greatest intensity. If no peptone is present the fluid becomes simply green or bluish-green. The simultaneous baryta precipitate does not interfere with the test, as it rapidly subsides while the supernatant fluid retains the color.

Or the lead filtrate may be treated with balf a volume or its own bulk of concentrated hydrochloric acid, and then with an acid solution of phosphortungstic acid, until there is no longer a precipitate. The latter is then promptly filtered off, because, if allowed to stand, there appears on the surface a second reddish precipitate, which interferes with the subsequent demonstration of the peptone. The precipitate is washed on the filter with a solution of sulphuric acid, three to five per cent. strong, until the filtrate passes through colorless, then turned into a capsule, intimately mixed with baryta in substance, the mixture treated with a little water, and, after being warmed for a short time, filtered. If too strongly heated, a dark-hued filtrate is obtained, which is to be avoided. To

this filtrate the "biuret" test is again applied.

The second method is both shorter and more delicate, detecting, according to Neubauer, peptone in solutions containing but .1 gram to the litre, while the tannic acid method requires .15 to .2 gram. It is the one preferred by v. Jaksch, to whom we are indebted for so much of our knowledge of peptonuria and its clinical significance.

Raife's Test.—A rough test for peptone, which answers very well when a considerable quantity is present, may be made by placing a drachm, or 3.5 c.c., of Fehling's solution in the bottom of a test-tube and gently overlaying it with an equal bulk of urine. At the point of contact a zone of

phosphates forms, while above this, if peptones are present, a delicate rose-colored halo will float. Should the peptones be mixed with albumin, the halo will be purple.

Randolph's Test with Acid Mercuric Nitrate and Polassium Iodide.—This test, suggested by the late Dr. N. Archer Randolph, of Philadelphia, is based upon the fact, that if Millon's * reagent be added to an aqueous solution of iodide of potassium, a red precipitate of mercuric iodide results, but if peptone or bile acids are present, the precipitate is yellow. To 5 c.c. of urine, which must be cold and but faintly acid, add two drops of a saturated solution of iodide of potassium and then 3 or 4 drops of Millon's reagent, when, if peptones or bile acids are present, a yellow precipitate falls. Then the question as to whether it be bile acids or peptones must be settled by the tests for the former.

The Clinical Significance of Peptonuria.—The instances in which peptonuria occurs are numerous. The best determined fact with regard to it is that discovered by Maixner, that it is always present when pus corpuscles are disintegrating somewhere in the body, and in most of the diseased states in which it has been found, such a condition of affairs has been probable. It has been especially studied by Maixner, v. Jaksch, Fenomenow, and Pacancowski. Among the diseases in which it has been found may be named typhoid fever, variola, scarlatina, miliary tuberculosis, erysipelas, acute arthritis, pulmonary tuberculosis, gangrene of the lungs, croupous pneumonia, purulent

^{*} To make Millon's reagent, dissolve one part by weight of quicksilver, by the aid of warmth, in one part by weight of concentrated nitric acid, and dilute with an equal volume of water.

pleurisy, embolism, carcinoma of the gastro-intestinal tract and of the liver, catarrhal jaundice, parametritis, cerebral apoplexy, parotitis, abscess. As above mentioned, it may be the product of the pyogenic organisms, as the pyogenes aureus, or of the pneumonococcus and the streptococcus pyogenes.

Exception to the above explanation may have to be made in some cases of carcinoma of the gastro-intestinal tract, and cancer of the liver and uterus. In cancer of the stomach and small intestine, Maixner early ascribed the peptonuria to absorption by the ulcerated surfaces, of the peptone of digestion, but in cancer of the cesophagus, rectum, and uterus, we must have recourse to disintegration of new-formed tissue as the sole source. The almost invariable association of peptonuria with cancer of the liver, Pacancowski thinks, compels the conclusion that the liver in health has something to do with the conversion of peptone into albumin, an office that in cancer is interfered

Although Senator, Petri, and Poehl assert that albuminuria and peptonuria coexist, Maixner, and more recently Pacancowski, fail to confirm this. The latter failed to find it in four cases of chronic nephritis and one of acute. Their possible coexistence is now, however, generally admitted.

It has also been held that peptone may originate from the conversion of albumin by reason of a sort of fermentative action of the cellular elements of urine, as well as by the agency of bacteria.

Hemialbumose, or Propeptone.—This substance is an intermediate product in the conversion of albumin into peptone during gastric and pancreatic digestion. It is,

therefore, abundantly present along with albumin in the gastric and intestinal contents, and, unlike peptone, is also found in the blood during digestion. It was first found by Bence Jones in the urine in a case of osteomalacia, and later by Kühne in another case of the same disease.

Hemialbumose, like albumin, is insoluble in alcohol; sparingly soluble in cold water, but very easily in hot water and water containing only traces of acids, alkalies, or salts. It is not, therefore, precipitated by heat from its watery solution, as is albumin, but if the solution be made strongly acid and concentrated salt solution be added thereto, hemialbumose is precipitated. If the cloudy fluid be now heated, it becomes transparent, but again turbid on cooling. A further large addition of salt maintains the precipitate in spite of heating. Hemialbumose is precipitated by pure nitric acid, but redissolved with the production of an intense yellow color on being heated, and reprecipitates on cooling. An excess of nitric acid, redissolves the precipitate even in the cold, with the production of the same orange-red color.

In these respects hemialbumose differs strikingly from both peptone and albumin. It is like peptone in that it exhibits the biuret reaction with an alkali and salt of copper. Like albumin, it is precipitated by adding first acetic acid and then ferrocyanide of potassium; also by phosphor-molybdanic, phosphor-wolframic, tannic, and picric acids, the precipitate, except that with picric acid, being undissolved by warmth.

To test the presence of Hemialbumose, the albumin must first be removed. This is accomplished by acidifying the urine with a few drops of acetic acid, adding about one-sixth its volume of concentrated salt solution,

69

boiling, and filtering off the precipitate. Albumin and globulin remain upon the filter. The filtrate is then allowed to cool, and if a turbidity arises or is produced by the further addition of salt solution, which disappears by heating, and reappears with cold, propeptone is present. If desired, the precipitate can be filtered off the filtrate, redissolved in a little water, and reprecipitated by acetic acid and ferrocyanide of potassium.

Hemialbumose is removed from fluids by adding acetate of iron and boiling, or by boiling with hydrated lead oxide.

Fibrin.—Fibrin is found in the wine when there are hemorrhages from the genito-urinary passages, and in intense inflammation of these passages and of the kidneys; also in a condition of fibrinuria which occurs in the Isle of France; finally, in chylous urine fibrin is present.

Recognition.—It is recognized by its spontaneous coagulation, the product of which is, however, not to be confounded with mucus or the glairy substance formed by the action of ammonium carbonate on pus; also by its fibrillar structure as shown by the microscope.

Coagula may be filtered out from urine by means of muslin, and washed with water to free them from urinary constituents. If insoluble in dilute alkalies and in 5 to 10 per cent. solution of sodium chloride, they are fibrin.

The presence of Pepsin and Trypsin in urine has been lately asserted.

Proteids Contrasted.

Albumin, hemialbumose, and peptone are soluble in water; globulin is not. All of these proteids are soluble in alkalies and acids, forming with them soluble binary combinations, behaving to the alkalies as acids and to acids as bases. They are soluble also in basic salts (phos-

phates and carbonates of the alkalies), since they take away from these salts the bases. The combination of albumin with a base is called an albuminate, that of albumin with an acid, acid-albumin (syntonin). No special term is applied to the combinations of globulin with bases or with acids.

If the acid be withdrawn from an acid-albumin in solution, by neutralization with an alkali, the insoluble proteid is liberated, and falls as a precipitate. The same thing happens when a dissolved albuminate is neutralized by an acid.

Hemialbumose, although soluble in water, is even more easily soluble than globulin in alkalies and acids, as well as in neutral salts. Albumin and peptone differ from each other also in that albumin is easily converted into precipitable albumin, while peptone is not.

Fibrin, finally, is soluble neither in water, like albumin, hemialbumose, and peptone, nor in dilute cold acids or alkalies, like precipitable albumin, nor in salt solutions, like globulin.

Mucin, like precipitable albumin, is not soluble in water, but is in alkalies and strong mineral acids; on the other hand, it is not soluble in dilute mineral acids, nor in concentrated organic acids.

Hæmoglobin and methæmoglobin are recognizable by their color.

VIII. SUGARS FOUND IN URINE.

Glucose, C.H 12O6.

While the assertion of Brücke and Bence Jones that glucose is present to a slight degree even in normal urine has been quite generally accepted, and has apparently been confirmed by the more recent researches of Dr. Pavy,* of London, it has been contradicted by Seegen, and very careful investigations by my colleague, Professor Wormley, confirm the results of Seegen. † Dr. George Johnson, ‡

^{*} Pavy, " Points Connected with Diabetes," London, 1879.

[†] Seegen, "Der Diabetes Mellitus," 2 Aufl., s. 224.

[‡] Johnson, G., Brit. Med. Jour., Jan. 8, 1887.

and his son, G. Stillingfleet Johnson, have also lately joined the group who deny the presence of glucose in normal urine. It does not follow, however, that instances may not occur in which small quantities of sugar, barely, if at all, recognizable by the ordinary tests, are present in urine, and that these have no clinical significance.

opinion as to the value of the different tests is due to the unequal experience of different observers with a particular test. Mistakes are less likely to be made by a beginner with a freshly made Fehling's solution than with Trommer's test, but if the latter is done as directed below a doubt constantly arise where the evidence of one is insuffisome notice is demanded; and it is suggested that for practical purposes the student should select some one or more of these and accustom himself to their use, and to the modifications in results to which all are more or less subject. I am confident that much of the difference of tolerably constant result obtains. It is necessary, however, to be familiar with more than one test, because cases of cient. (See especially remarks on qualitative testing, only those will be given which have borne the trial of experience, or, being new, are so highly commended, that Of the large number of tests for the detection of sugar, p. 104.)

Specific Gravity and Quantity.—The specific gravity alone, when 1030 or more, affords a presumption of the presence of sugar, and if at the same time the urine is very pale, and far exceeds 1500 c. c. (50 fluidounces) in twenty-four hours, the probabilities are much increased. These facts at least call for the use of other tests to determine the question.

The Copper Tests.

The copper tests depend upon the power which grapesugar possesses of reducing the oxide of copper to a lower state of oxidation. The oldest of these tests is Trommer's, in which the oxide of copper is set free at the time of its application by liquor potassæ or sodæ, in excess.

Trommer's Test.—r. To 4 or 5 c. c. of the suspected urine, add one quarter its bulk of liquor potasse or sodæ. Then add drop by drop of a 10 per cent. solution of cupric oxide. On adding the first drop, a blue precipitate of hydrated cupric protoxide occurs, which, if sugar is present, is redissolved on agitating the mixture. This is presumptive evidence of the presence of sugar, but is not relied upon. The addition of the copper solution is continued until a slight excess has been added and the precipitate is no longer dissolved. The mixture is then boiled for a few seconds, and if sugar is present, a copious precipitate of yellow cuprous hydroxide takes place. This subsequently loses its water and becomes the red cuprous oxide which falls to the bottom or sides of the test-tube, to which it often closely adheres.

If more than 5 to 10 drops of the cupric oxide are required to produce the turbidity, dilute a portion of the urine with 4 or 9 volumes of water and reapply the test as before.

The precise reaction is not known.

2. A second similarly prepared mixture of these ingredients may be made and set aside without the addition of heat, for from six to twenty-four hours. If sugar is present a similar precipitate of suboxide of copper will take place. If the reaction is at all doubtful it is important that this

check-test should be made, since, as Neubauer points out, most of the other organic substances which reduce the salts of copper do so only when heated or after long boiling.

Precautions.—These apply to all the copper tests and should be carefully studied. I. Albumin must always be removed, as it interferes with the reduction of the cupric

2. Too much of the solution of cupric sulphate or too strong a solution should not be used, because the prolonged boiling of any urine with a large excess of copper will produce a yellow or greenish-yellow color, which may not appear until the mixture cools off.

3. While the fluid must be made to boil for perhaps half a minute, the precipitate should take place without prolonged boiling, as a reduction by other organic substances is induced by prolonged boiling.

4. The flocculent precipitate of earthy phosphates should not be mistaken for the suboxide of copper; it is either transparent or of a pale greenish hue. On the other hand, a mere change of color is not sufficient. Strictly normal urine almost always has a decolorising effect. There must be an actual yellow or red precipitate. If it be desired to eliminate altogether any error due to the precipitation of the earthy phosphates, it may be done by adding the potash solution and filtering before adding the copper.

5. As already stated, cupric protoxide is sometimes reduced by other organic matters found in urine, especially uric acid, by creatinin, hippuric acid, the urates, hypoxanthin, mucus, indican, urochloralic acid found in the urine when chloral is administered internally, turpenoglycuronic acid when turpentine is taken, etc. Creatinin has also the property of redissolving cuprous oxide. Practically, uric

process, described on page 87. tion test, or, if absolute accuracy is required, to Brücke's organic substance or by the sugar itself, and demands that ever, does indicate a partial reduction either by some other mistaken for a precipitate. A yellowish precipitate, howand the substitution of a yellowish tinge is also not to be the urine should be subjected to the bismuth or fermentathe suboxide, while the disappearance of the blue color specific gravity, to the fact that a precipitate of the phosgreen precipitate results. Attention should be paid to the larly. When these partial reductions occur, a yellowishpepsin, peptones, urinary coloring matters, etc., act simiorganic substances generally, including creatin, creatinin, compounds have this latter effect; and not only albumin but monium chloride, ammonium urate, and other ammoniacal phates always takes place which must not be mistaken for certain other substances. Dr. Beale* has shown that amin urine and fail to reduce the oxide in the presence of On the other hand, a small amount of sugar may be present rine urines being for the most part light hued and copious. suspected when the urine is dark colored or scanty, sacchaacid is almost the sole source of error, and this may be

When proper precautions are observed, reliable results may be expected with Trommer's test with saccharine urines containing ½ of 1 per cent.

Salkowski's Modification of Trommer's Test. Solution 1.—Dissolve 100 grams of sodic hydrate, purified in sticks, in 300 c.c. distilled water. If there is any sediment carefully decant the solution.

Solution 2.—Cupric sulphate, chemically pure, 10 grams in 100 c.c. distilled water. To 9 c.c. urine add 3 c.c. of solution 1 in an

^{* &}quot;Kidney Diseases and Urinary Deposits," p. 246.

eight-inch test-tube, then the solution 2, drop by drop, shaking the mixture after each until the copper ceases to dissolve. Slowly heat to a little below the boiling point (be especially careful as to this), and if glucose be present the usual greenish, or yellow, or reddish hydroxide of copper will be seen, first in the upper stratum, then throughout the liquid. Where only a minute quantity of glucose is present the test-tube should be set aside for five to ten minutes, to allow the subsidence of the flocculent phosphates.

Two minutes' maintenance near (but avoiding the actual) boiling point will suffice in all cases.

It is said that this test in experienced, careful hands, will demonstrate 0.05 per cent. of glucose.

Other Copper Test Solutions-Fehling's and Pavy's Fluid.

It has been stated that when an alkali is added to a solution of sulphate of copper an abundant precipitate of hydrated cupric protoxide is thrown down. This is not dissolved by any excess of alkali added, but if some organic matter is present, or is added, an excess of alkali dissolves the protoxide. It is for this reason that, if sugar happens to be present in a suspected fluid to which these have been added, the precipitated protoxide is dissolved and a clear blue fluid results.

These facts enable us to construct a fluid which will hold the protoxide of copper in solution; but, in selecting an organic substance, one must be chosen which will not reduce the oxide of copper as does sugar, else it will make our test inoperative. Such substances are tartaric acid, mannife, and giverine, all of which are employed.

Fehling's Solution.—Dissolve 34.652 grams pure crystallized sulphate of copper in 200 grams distilled water; 173 grams chemically pure crystallized neutral sodic tartrate are then dissolved in 480 grams solution of caustic

soda of specific gravity 1.14 and into this basic solution the copper solution is poured, a little at a time. The clear mixed fluid is diluted to 1 litre, or 1000 c.c.

ro c.c. of this solution will be reduced by .o5 gram, or 50 milligrams, of diabetic sugar. If the Fehling's solution is to be kept some time, it is absolutely essential that it should be placed in smaller bottles holding 40-80 grams, sealed, and kept in the cellar.

Still greater security may be obtained by dissolving the cupric sulphate in 500 c.c. water, diluting the tartrate salt and potash to 500 c.c., and keeping the two solutions separately in rubber-stopped bottles. Equal volumes of the two solutions are united when needed for use.

With a view to avoiding the well-known defect of Fehling's solution—its tendency to spoil on keeping—Schmiedeburg suggested the substitution of 15 grams pure mannite for the sodic tartrate. The mannite should first be dissolved in 100 c.c. of water and 500 grams solution caustic soda, sp. gr. 1145, and the solution completed as above. My experience with this solution fully confirms Schmiedeburg's statement, and I strongly commend it.

Fehling's Solution with Glycerine.—With the same end in view, glycerine is used, as follows: Dissolve 34.652 grams of pure cupric sulphate in 200 c.c. of pure water, add 175 c.c. of pure glycerine, and mix thoroughly. 2. Dissolve 130 grams KHO in 500 c.c. water. Then add slowly solution 1 to solution 2, and dilute the whole to 1000 c.c. The solution is used as is Fehling's.

Pavy's Solution consists of-

The solution is made in the same manner as Fehling's, and roo minims correspond to ½ grain of grape sugar, the formula for grape-sugar being here taken C₆H₁₄O₇, while by Fehling it is taken C₆H₁₂O₆.*

To Use Fehling's and Pavy's Solutions for Qualitative Testing.—The same precautions laid down for the use of Trommer's test are to be observed, since the Fehling's and Pavy's solutions are simple modifications of Trommer's test and subject to the same sources of error.

In using either of the above solutions for qualitative testing, a small quantity, say r c.c., should be placed in a testube, and diluted with about four times its bulk of water. The mixture should then be boiled for a few seconds. If the solution remains clear on this boiling, add immediately the suspected urine drop by drop. If sugar is present in any quantity, the first few drops will usually cause the yellow precipitate; but the dropping may be continued if no precipitate occurs, reapplying the heat occasionally, until an equal volume of the urine has been added. If no precipitate occurs, sugar is absent, clinically speaking.

If a precipitate occurs on boiling the *test fluid alone*, a new supply must be obtained, or a little more soda or potash may be added, the fluid filtered, when it is again fit for use. The precipitate referred to is a cuprous oxide, the result of a reduction of the protoxide, which sometimes occurs when Fehling's or Pavy's solution is kept for some time. It is said to be due to racemic acid, into which tartaric acid is convertible on exposure. Under the influence

^{*} This should be remembered, as, in consequence of it, the same urine in the hands of different observers would yield slightly different results, according as one or the other solution is used.

of heat this acid oxidizes at the expense of the protoxide of copper, and the suboxide is precipitated; hence the necessity of boiling a solution before adding the suspected fluid. I have sometimes noted that boiling a spoiled undiluted Fehling solution does not reduce the copper, while from the diluted solution, when boiled, the suboxide is thrown down; so that boiling a diluted Fehling's solution becomes a more delicate test of its quality than boiling the undiluted solution. All possibility of such source of error may be avoided by keeping the solution of copper separate from that of the potassic tartrate in the caustic soda solution, and mixing them at the moment they are required for use.

The delicacy of Fehling's test is said to be o.oor per cent. glucose in water. With glucose in urine it is less delicate, but still in my hands it is the most delicate and reliable test available.

It will be noted that in the use of Fehling's and Pavy's copper solutions an excess of the test fluid is always used, while in the method described as Trommer's, the effect of adding too much may be to produce a yellow sediment and coloration on boiling with any urine, especially on cooling. It occasionally happens, also, when Fehling's solution is used, that no reaction occurs until considerable urine has been added and the mixture cools down after the boiling, when a yellowness or mikiness makes its appearance. Dr. Roberts * believes this reaction due to sugar. But this is at least doubtful, for I have known it to occur in urine which, when treated by Brücke's process, presently to be described, was found to be without a trace of sugar.


^{* &}quot;Urinary and Renal Diseases," Amer. ed., 1879, p. 190.

It may be due to sugar, but it is as likely to be due to uric acid or to some other of the reducing substances contained in urine. In such a case the only way to settle the question is to resort to Brücke's process described on p. 87. But this may be said, that the quantity of sugar which could occasion such a reaction is of doubtful clinical significance. Filtering through animal charcoal is a less troublesome expedient than the lead process. This gives a perfectly clear fluid to work with, but in such filtration a small quantity of sugar is retained by the charcoal and must be washed out with distilled water. Again, it is almost impossible to obtain charcoal sufficiently free from impurities, even when specially prepared for the purpose.

precipitate watched, and then another 1 added, the heat metre of Fehling's solution. If in doing this, I c.c. of urine has been added, it will have contained just half of 1 per cent. of sugar. If more than I c.c., it will have contained less than one-half per cent., but more than onequarter per cent. If exactly 2 c.c. are used, it will have If, on the other 10 cubic centimetre of the suspected urine is added from a suitably graduated pipette. Heat is then reapplied, the again reapplied, until it is found, after proper subsidence, that all the blue color is removed from the cubic centi-Quantitative Analysis by Fehling's or Pavy's method of analysis by Fehling's or Pavy's solutions, and one which may be used in the consulting room as easily as the laboratory, is the following: One cubic centimetre of Fehling's solution is diluted in a large test-tube with four cubic centimetres of distilled water, and boiled as described for qualitative testing. Its purity being thus ascertained, Solutions. Volumetric Process.-The contained exactly one-quarter per cent.

hand, but half a cubic centimetre is used, it will have contained 1 per cent., one quarter of a cubic centimetre, 2 per cent., and so on.

Fig. 7.



Varieties of Burette Stand.

If the proportion of sugar is large, as indicated by the specific gravity or qualitative test, the urine should be diluted with nine parts of water, and the result multiplied by 10.

The following table, prepared by Prof. Wormley, will greatly aid in the estimation of glucose by this method:—

Percentage of glucose in urine as indicated by the quantity of urine required to exactly decolorize I c.c. of Fehling's solution.

OF DILUTED URINE, I IN 10.	Frine. Pe	0.4 [50+4], 12.5			0.7,			1.0, 5.0		r.4, 3.5		 		 I			4.0, 1.25	The same of	 	7.0, 0.7	 9.0, 0.55	
OF UNDILUTED URINE.	C. C. Urine. Per cent.	0.1 [5+1], 5.0	0.12, 4.2	0.14, 3-5	0.16, 3.1	0.18, 2.7	0.2, 2.5		0.3, 1.66		0.4 1.25	 	0.6, 0.83	 0.8, 0.62	0.0, 0.55	.0						

If it is desired to determine the quantity in English measures, Pavy's solution may be used instead of Fehling's, and roo minims measured off into the test-tube, diluted with four times its bulk of water, and boiled as before. Then the urine, diluted if necessary, is allowed to fall into the liquid, drop by drop, the heat being constantly renewed, until all the blue color has disappeared; and when this has

happened the quantity of urine used will have contained just half a grain of sugar.

until all the blue color disappears. The titration must be c.c. will contain 10 grams or 10 per cent. Pavy's solution 10 times, the 5 c.c. will contain .5 gram sugar, and 100 of urine will have contained exactly this quantity, that is, used to decolorize 10 c.c. of Fehling's solution, the 5 c.c. ance is ascertained and noted. If 5 c.c. of the urine are repeated, if necessary, until the exact point of disappearthe urine is cautiously added, the solution being kept hot, to .05 gram of sugar. If now the urine has been diluted .05 gram, since 10 c.c. of Fehling's solution corresponds may be used similarly. This is allowed to subside, and if any blue color remains burette. Immediately a yellow or red precipitate will fall. urine is allowed to fall into the hot solution from the heated with a gas flame or spirit lamp, when half a c.c. of preliminary trial. The capsule containing the diluted according to the proportion of sugar estimated by a burette to 50 c.c. with urine diluted from 5 or 10 times, sule and diluted with 40 c.c. of distilled water. Fill the metres of Fehling's solution are placed in a porcelain cap-Fehling's solution should be placed on a wire gauze and burette and stand shown in Figure 7. Ten cubic centi-Greater precision may be obtained by the use of the

Cupric Test Pellets.—These were first suggested by Dr. Pavy at a meeting of the Clinical Society of London, in January, 1880, and were first made in this country by McKelway, of Philadelphia, at the suggestion of Dr. Joseph Neff.* They contain the elements of Fehrus

^{*} New York Medical Record, March 23, 1880.

AMMONIATED CUPRIC TEST.

ling's solution in solid form, and are very neatly made in the shape of a compressed pill. As made by Mr. McKelway, each pellet when dissolved in distilled water represents 5 milligrams of diabetic sugar, and they may be used for approximate quantitative as well as qualitative

To use them, dissolve one in a small quantity of water in a test-tube and boil the solution. The purity of the pellet is thus tested, for if it has spoiled, the cupric oxide is reduced spontaneously. If the solution remains clear, the urine is added drop by drop, the temperature being kept up, and if sugar is present the usual precipitate occurs. The quantity of urine which just removes the blue color of the solution in which one pellet is dissolved contains 5 milligrams of sugar.

These pellets are very convenient to carry, but in my experience they are liable to become unfit for use sooner than a Fehling's solution, and unless very carefully made they keep but a short time.

Pavy's Ammoniated Cupric Test.—Dr. Pavy has suggested, in lieu of the usual form of Fehling's solution, an ammoniated cupric test solution which has the advantage of obviating the precipitate of sub-oxide of copper, which is held in solution by the ammonia, so that a simple decolorization takes place as the cupric oxide is reduced; the completed reduction being indicated by the total disappearance of the blue color.

The proportions are as follows:--

METRIC SYSTEM.

rams.	33	**	.c.	
4.158 grams.	20.400 "	. 20.400 "	300 c.c.	
Cupric sulphate,	Potassic sodic tartrate,	Potash caustic,	Strong ammonia (sp. gr. o.880),	Water to I litre.

ENGLISH SYSTEM.

36 1/2 grains.	***	***	6 fluid ounces.
36	178	. 178	9
	*		1
			0
			88
			0
			H
	trate		(SD.
Cupric sulphate, .	Potassic sodic tartrate,	Potash caustic,	Strong ammonia (sp. gr. 0.880),
Cupric	Potassi	Potash	Strong

Water to one pint.

Dissolve the potassic sodic tartrate and potash together in a portion of the water, and the cupric sulphate with the aid of heat in another portion; pour the solution of cupric sulphate into the mixture of potassic sodic tartrate and potash; when cold add the ammonia, and finally, with water, bring the volume of liquid to the bulk specified.

Ten c.c. of the liquid, corresponding to .05 gram or 50 milligrams of sugar, is a convenient quantity to work with, and does not call for the employment of any appliance to obviate the inconvenience arising from the evolved ammonia. In performing the analysis twice the volume of water is added to the 10 c.c. of test employed.

It should be stated that Hehner, Yeshida, and Sutton assert that by this method the ratio of reduction is seriously influenced by the amount of fixed alkali present and by the strength of the ammonia.

Purdy's Modification.—Dr. Charles W. Purdy substitutes in his formula 50 c.c. of glycerin for the sodic tartrate, dissolving the capric sulphate in a part of the water and adding the glycerin, while the caustic potash is dissolved in another portion. The two solutions are then mixed, the ammonia added to make one litre, the whole filtered, and the volume increased. Ten c.c. of this solution correspond to .05 gram or 50 milligrams of sugar. If the English system is used the sodic tartrate is substituted by half an ounce of glycerin and the ammonia is reduced to five and a half ounces. One ounce of this solution is reduced by one-fourth of a grain of grape sugar.

Cupric Test Paper.—Dr. Oliver has also constructed a cupric testpaper with tartrate of cuprammonium, this salt being selected as the only one which is permanent on exposure to the air, and the only one that is stable when boiled with an alkaline carbonate as well as with a caustic alkali.

The test-paper is a compound one, consisting of one charged with the reagent and the other with sodium carbonate, the two being united by a thin layer of rubber.

To Use.—1. Drop a test-paper into 60 minims of soft or distilled water. 2. Boil for a few seconds until the water assumes a greenish tint. 3. Remove the papers. 4. Reboil the solution, and then add one drop of the suspected urine. 5. If glucose be present, reduction will take place without further application of heat, though, if preferred, the boiling may be continued and the reaction hastened. In any case, if

the solution remains transparent for a quarter of a minute, heat should be applied to the boiling point for one minute. If then no opacity whatever appears, it may be safely inferred that glucose is not present in pathological amount.

The Fermentation Test.

An excellent test for the presence of sugar is the fermentation test. The most convenient method of its application is as follows: a small quantity of ordinary baker's or brewer's yeast (about a fluidrachm, or 3 to 4 c.c.) is added to about 4 ounces of urine in a 6-ounce vial, which is lightly corked and subjected to a temperature of 15-25° C. (59-77° F.). If sugar is present, evidences of fermentation will present themselves, generally within twelve hours, by the formation of carbonic acid gas, which passes off, leaving the fluid lighter and reduced in specific gravity in proportion to the quantity of sugar present.

Dr. Roberts early announced that urine containing less than o.5 per cent. or $2\frac{1}{2}$ grains to the ounce, yields no sign to the fermentation test. Claims to much greater delicacy are made, but it is evidently impossible to discover a much smaller proportion, because water will absorb an equal bulk of carbon dioxide, so that all the gas generated in a solution containing $\frac{4}{10}$ per cent. must disappear.

Dr. Roberts has made use of the fact of the lowering of the specific gravity in devising a quantitative method. He has shown by careful experiments that every "degree" in specific gravity lost in fermentation corresponds to one grain of sugar per fluidounce. Thus, if before fermentation the specific gravity of a given specimen is 1050, and after fermentation it is 1020, it will have contained 30

only to make the comparison. and each day, when the physician makes his visit, he has and a comparison made. While some time is required specific gravity of the decanted fermented urine is then grains to the fluidounce. The method recommended by Dr. Roberts is as follows: Four ounces of the saccharine preparation can be made by the patient himself or friends, to complete the fermentation, yet, as Dr. Roberts says, the taken; at the same time, that of the unfermented urine, 4-ounce vial, filled with the same urine, but without any or other warm place. Beside it is placed a tightly-corked escape of the carbonic acid, and set aside on a mantelpiece urine are put in a 12-ounce bottle, and a piece of combe complete, and the scum cleared off or subsided. The yeast. In eighteen to twenty-four hours fermentation will bottle is then covered with a nicked cork to permit the pressed yeast, as large as a small walnut, is added. The

The percentage may be roughly arrived at by multiplying the number of degrees lost by .23.

Appliances have been suggested within the past year by Max Einhorn and S. P. Kramer, with the object of securing greater accuracy in the quantitative estimation by fermentation; but, as the results are at best but approximate and the apparatus even less convenient than that suggested by Roberts, I omit them. They may be purchased, with directions for their use, from Eimer & Amend, New York City.

Böttger's Bismuth Test.

To a given quantity of urine add an equal volume of a concentrated solution of sodium carbonate (crystals 1 part, distilled water 3 parts), then a little subnitrate of bismuth. Shake and boil for a couple of minutes. The sugar possesses the power of reducing the salts of bismuth, and

if present, the black metallic bismuth will shortly be deposited on the side of the test-tube. If the quantity of sugar is small, the bismuth will assume a grayish hue; hence, when this is the case, a very small amount of bismuth should be used in making the test.

This is a brilliant test, and except albumin or other substance containing sulphur, nothing but sugar is supposed to reduce bismuth salts. I do not, however, think as highly of it as I formerly did, as the dirty gray hue said to be characteristic of very small quantities of glucose sometimes presents itself when no sugar is present. Especial care must be taken to remove the albumin before applying

the bismuth test, or we may use—

Brücke's Modification of the Bismuth Test.*—
To meet the difficulty referred to in the above paragraph, due to sulphur occasionally present in the urine causing a black precipitate of sulphide of bismuth, Brücke recommends the use of Frohn's † reagent—iodide of bismuth and potassium—to remove the disturbing elements, as follows:—pour into a test-tube a certain quantity of water, say 10 c.c., and fill another tube to the same level with urine. To the first add a drop of the Frohn's reagent, which will cause a decided cloudiness. Then add, drop by drop, hydrochloric acid until the precipitate is redissolved. In this way we ascertain approximately how much HCI

^{*} Proceedings of American Pharmaceutical Association, 1887, p. 287.
Also Hofmann and Ultzmann, "Analysis of Urine," American translation. New York, 1879, p. 93.

^{† 1.5} gram freshly precipitated subnitrate of bismuth is mixed with 20 grams water and heated to boiling; then 7 grams iodide of potassium and 20 drops of hydrochloric acid are added. The reagent is orange red.

must be added to the urine. Acidify the urine with such quantity; treat it with the reagent until precipitation is complete and filter. The filtrate, which should not now become cloudy on adding HCl or the reagent, is boiled for a few minutes with an excess of a concentrated solution of caustic soda or potash, as in Böttger's test; if a gray or black color results, or such a precipitate is formed, the presence of sugar is proven beyond a doubt.

The delicacy of this bismuth test is said to be 0.4 per cent. glucose in water.

Nylander's Modification of the Bismuth Test.—The solution for this test, also known as Almen's solution, is made as follows: Basic bismuth nitrate, 2 grams; sodic potassic tartrate, 4 grams; sodic hydrate (purified sticks), 8 grams; distilled water, 100 c.c.

Rub the solids to a coarse powder, add a little water to form a paste, then, while stirring briskly, gradually add the balance of the water. Let stand for five minutes to complete the solution. Filtration gives a perfectly clear solution. Add I c.c. of this to IO c.c. of urine, boil for two minutes in an eight-inch test-tube. If a light gray to black precipitate form, it may be, usually is, due to the presence of glucose. If albumin is present, it must first be removed.

This test, it is claimed, will detect .025 per cent. of glucose, while smaller quantities give a grayish tint to the flocculent precipitate. Urines of high specific gravity also reduce the bismuth oxide.

The Picric Acid and Potash Test.

This test, originally suggested in 1865 by C. D. Braun, * and revived by Dr. George Johnson† in 1882, is based upon the fact that grape-sugar, when boiled with picric

^{*} Ueber die Einwandlung der Pikrinsture in Pikraminsture, und über die Nachweisung der Trauben-Zucker.—Zeitschrift für Chimie, 1865.

[†] Lancet, November 18, 1882

acid and potash, reduces the yellow picric acid to the deep red picramic acid, the depth of color depending upon the amount of sugar present.

For qualitative testing, to a fluidrachm of the suspected urine, add 40 minims of a saturated solution of picric acid, and half a drachm of liquor potassæ.* If albumin is present, a turbidity will develop on the addition of picric acid, and thus the presence of this substance be recognized; but it does not interfere with the test. Boil the mixture, and if sugar is present, a dark mahogany-red color will be produced. If normal urine be treated in the same way, a somewhat darker hue is also developed, but not nearly so marked as when sugar is present.


Or the test may be applied, especially at the bedside, in the following manner:—

Into a test-tube, graduated up to three drachms, is put about one-third of a grain of picric acid—as much as can be carried on the point of a penknife; then add half a drachm of water; the acid is dissolved in the water by the heat of a lamp. Now, add half a drachm of urine, and the presence of albumin is ascertained; next add a grain lump of caustic potash, and boil the liquid for a few seconds, and the dark coloration appears.

A delicacy of o.or per cent. of glucose in water is claimed for picric acid.

For quantitative testing Dr. Johnson directs a standard solution made by boiling together a fluidrachm of a solution of grape-sugar of the

^{*} Dr. Charles F. Adams has altered the strength of the solutions so that the same result is obtained by taking 5 c.c. of each. Thus, of urine 5 c.c.; liq. potassee, sp. gr. 1.036, 5 c.c.; sol. picric acid (gr. 3.5 to f_{∞}^{2}) 5 c.c.; water 5 c.c.



of the strength of the British Pharmacopœia.** nium; 3 iv of glacial acetic acid, sp. gr. 1065; sp. gr. 1.44; 3 iv of solution of acetate of ammodrachm of liquor potasse, and forty minims of a to the ounce, four times diluted, retains its color distilled water up to 3 iv. The ingredients are all thoroughly 3 j of solution of perchloride of iron, ever, not permanent, and Dr. Johnson imitates it obtained is that which results from the decomto it by adding distilled water. The color thus is not that of the four-drachm mark, it is raised tube in cold water, and if the level of the liquid red. It is cooled by cautiously immersing the which its pale-yellow becomes a beautiful claretliquid is kept boiling for sixty seconds, during which should be marked at four drachms. The mixture is conveniently made in a large test-tube, creased to four drachms with distilled water. The used for comparison. unchanged, at least when kept in the dark, and is adding 3j of liquor ammoniæ, and diluting with with a solution of ferric acetate made by mixing times diluted. The color of this solution is, howpicric acid by a grain of sugar to the ounce, four position of 40 minims of a saturated solution of saturated solution of picric acid, the whole instrength of one grain to the fluidounce, half a This solution, corresponding to a grain of sugar

In testing for sugar, treat a fluidrachm of urine as indicated for qualitative testing, and dilute the mixture with distilled water up to four fluidrachms. The test-tube, which should

Liquor ferri perchlor. f3j; liq. ammon. acetat. f3 ss; ac. acet glacial. f3 ss; liq. ammon. f3j; aqua destil. ad f3 iv.

^{*} Dr. Adams has also substituted the ingredients by those of the United States Pharmacopecia, much more convenient to the American worker, as follows:—

it stood, to the 20-mark, the quantity of sugar would be two grains; if charimeter devised by Dr. Johnson and illustrated by Figure 8. The tube, the boiled mixture is to be introduced. Should the two agree precisely in color, the suspected urine will have contained one grain of sugar to the ounce. If, on the other hand, the mixture requires to to raise the level of the fluid, say from the 10-division mark, at which seconds. Cool the liquid by carefully immersing in cold water, and if below the four-drachm mark it is to be again raised to it by the addition of distilled water. A comparison is then made between the resulting fluid and the standard solution by means of the picro-sacstoppered tube on the left contains the standard solution. Into the graduated tube on the right, having the same diameter as the standard this should be done with distilled water, and if enough must be added be a large one, may be conveniently marked at four fluidrachms. Raise the liquid to the boiling point and keep it boiling for sixty be diluted in order to make it correspond with the standard solution,

to 40, four grains; to 45, four and a half grains.*

In making an analysis, the picric acid must be added in proportion to the amount of sugar. If the proportion of sugar be as high as six grains per ounce, a drachm of the picric acid solution will be required. If it is higher than this, the urine should be diluted with distilled water in a definite proportion before commencing the analysis, and the dilution borne in mind in making the calculation. When the urine has been diluted ten times, the figures on the saccharimeter indicate the number of grains per ounce. Thus, when the ten-times diluted urine, after boiling with picric acid and potash, is further diluted from 10 divisions

^{*} A more exact comparison of the saccharine liquid with the standard can be made by pouring into a flat-bottomed colorless tube, about six inches long and an inch in diameter, as much of the standard as will form a column of liquid about an inch in height, and an exactly equal column of the saccharine liquid in a precisely similar tube. Looking down on the surface of a white porcelain siab or white paper through both tubes at once, a slight difference in tint is easily recognized; and if the liquid to be analyzed be found darker than the standard, it is returned to the saccharimeter and diluted until the two fluids are found to be identical, when the final reading is made.

to 35, to obtain the standard color, the amount of sugar is 35 grains to the ounce.* We may reduce the amount of sugar per ounce to the proportion per cent. by a simple proportion, in which the first term is 455.7 grs. the weight of an ounce of water at 0° Cent.,† 100 the second term, and the quantity of sugar per ounce the third. Thus, if there he 20 grs. of sugar to the ounce, then as 455.7: 100:: 20: 4.3

It has been said that when normal urine is treated with picric acid, potash, and heat, a slight coloration also takes place, which Dr. Johnson estimates as about equal to the change which would be produced by a solution of glucose containing 0.5 to 0.7 grains to the fluidounce. Dr. Johnson, in common with others, now ascribes this reaction to creatinin.

Dr. Johnson, after much experimental research made in association with his son, G. Stillingfleet Johnson, claims that this method is "as accurate as any other," and that for the estimation of sugar in the urine it is even more accurate than either Fehling's or Pavy's process, because the picric acid is not acted on by uric acid or urates, which reduce the oxide of copper. He also claims that the method by the picro-saccharimeter is more speedy than any other, the materials and apparatus inexpensive, and not liable to undergo rapid change.

^{*}Distilled or clear rain-water should be used, because the turbidity resulting from the action of potash on the salts of lime interferes with the exact estimation of the depth of color.

[†]This is the weight of the ounce of distilled water according to the United States Pharmacopœia. The English fluidounce weighs 437.5 grains.

The Indigo-carmine Test.

When a solution of indigo-carmine alkalized by sodium carbonate is boiled and kept heated, the blue color remains, but when a drop of a solution of glucose or saccharine urine is added to the hot solution, a beautiful play of colors occurs, terminating in pale yellow. Mulder first suggested this reaction as a test for sugar. Dr. Oliver has revived it for both qualitative and quantitative analysis. The indigo-carmine solution is not available, because when the carmine and sodium carbonate are present in solution the fluid undergoes a gradual change, the indigo-blue color gradually passing into a pale green. Nor is it convenient to keep the solutions separate. Dr. Oliver has, however, constructed a stable test-paper, charged with a definite quantity of the two reagents, indigo-carmine and sodium carbonate; while increased sensitiveness is produced by the use of an additional paper charged with sodium carbonate.

In Testing.—I. Unless very recently prepared, one of the carmine test-papers should first be gently heated, say on a knife blade or spatula, by being held for a few seconds just above a flame.* It should then be dropped into a half-inch test-tube, and water should be poured in to the amount of 60 minims.

2. Heat is applied, the tube being gently shaken, and boiling kept up for a second or two. The solution will then be quite blue, and, if the water added was soft or distilled, it will be perfectly transparent. Any turbidity observed will arise from the use of hard water, in which case a sodium-carbonate paper should be dropped into the solution. The test-paper may now be removed or allowed to remain.

3. Not more than one drop of the suspected urine is let fall into the tube from the pipette held in an upright position.

4. The contents of the tube are again boiled for a few seconds; then the tube should be raised an inch or two above the flame, and held without shaking while the solution is kept quite hot, but without ebullition, for exactly one minute. If glucose is present "in abnormal amount,"

^{*} This is necessary, because the test-paper, on being kept for some time, loses its sensitiveness for detecting small quantities of sugar, say I to 2 grs. per ounce.

purple and violet, and finally, of the original blue. When the last color has developed, agitation will cause the return of succession purple, red, reddish-yellow, and, finally, straw-yellow." violet; then, according to the quantity of sugar, there will appear in says Dr Oliver, "the soft rich blue will be seen, first of all, to darken into

the color of the solution at the end of one minute will be unchanged. to 60 seconds. If the urine contains less than 1/2 a grain to the ounce, When small—say from 2 to 3 grs. to the ounce—it may require from 30 inversely with the amount of glucose present. When the latter is large -over 20 grs, to the ounce-it will amount only to a few seconds. The time required for the reaction to commence, after boiling, varies

Precautions.- r. Care should be taken during the heating not to

shake the tube, or to keep up active ebullition.

of white paper propped up an inch or two beyond the tube. contents viewed by the reflected light of some bright object-as a sheet escape detection. The tube should be kept below the eye level, and its between the eye and the sky, for then the early color changes may 2. While keeping the contents of the tube hot, it should not be held

The test is as available by artificial as by daylight.

No amount of agitation will restore the blue removed by these agents. tube containing a trace of either of these substances or of Fehling's olution or the alkaline picric solution, lest a deceptive reaction occurs. blue color of carmine. Hence, care should be taken not to use a test-3. Any caustic alkali, as liquor potassæ or sodæ, will discharge the

carmine and Fehling's solution, found that whenever one drop was subthe indigo, Fehling's solution also gave negative results. mitted to the indigo test and the presence of sugar shown, confirmation On the other hand, whenever one drop of urine gave no reaction with was invariably provided by Fehling's solution used in the ordinary way. Dr. Oliver, comparing the results of experiments with the indigo-

substances apt to appear in urine in disease, albumin, peptone, * pus, of the urine, uric, oxalic, lactic, etc., reduce Fehling's solution. Of ordinary constituents of urine affect the carmine test, but all the free acids In further experimental testings, Dr. Oliver found that none of the

decolorizes the solution just as does iodized starch. * In a letter from Parke, Davis & Co., I am informed that peptone

mucus, blood, bile, leucin, tyrosin do not react with either test; nor does one drop of ammoniacal or decomposing albuminous urine, or weak solution of ammonium sulphide; ** but dextrin and milk-sugar as well as glucose reduce both. Inosit reacts with the carmine, and turns Fehling's solution green—a green precipitate falling, leaving the supernatant fluid blue, which, however, becomes green on reheating.

Of medicinal agents likely to find their way into the urine, iron sulphate, gallic and tannic acids alone react with carmine, as do they, also, with Fehling's solution.

Comparative experiments by Oliver with picric acid and potash showed that whenever the indigo-carmine test-paper afforded a reaction, a correspondent reaction was obtained with the picric solution.

Dr. Oliver has also applied his test to quantitative analysis, but it is rather cumbersome and not likely to be availed of. It is therefore

The Phenyl-hydrazin Test.-The true value of this test, suggested says, † as modified by Ultzmann and Bond, "it is the best single test nally directed as follows: To 50 c.c. (13.5 f 3) of the suspected urine add 2 grms. (30.8 grains) of phenyl-hydrazin hydrochlorate, and 1.5 grm. (23 grains) sodium acetate or 1 grm. (15.4 grains) of the latter, if the urine is not decidedly acid. Add also 20 c.c. (5,4 f3) of water scope. But this is not sufficient. Their nature must be confirmed by determining their melting point, which is 204 to 205° C. For this purpose the dry crystals are obtained by filtering them off the urine, washing with a small quantity of water, dissolving in a small amount of by Emil Fischer, is as yet undetermined, although highly estimated by known," detecting .025 per cent. of sugar in urine. I regard it, however, too troublesome a test to be ever largely availed of. The test was origifor one hour. If sugar is present, needle-shaped crystals of phenylglucosazon will have appeared, which must be recognized by the microsome. Most recently, in a noteworthy paper, Dr. Frank Dudley Blane the urine and reagents is then placed in a water bath and gently warmed unless the urine is nearly colorless. The capsule or beaker containing omitted from this edition.

^{*} Even a weak solution of ammonium sulphide will reduce Fehling's solution.

[†] N. Y. Med. Jour., Jan. 7, 1893.

dilute alcohol, and recrystallizing by evaporating at a low temperature.

After repeating this two or three times the crystals are collected in a
desiccator over sulphuric acid.

Then draw out a piece of thin glass tubing in a Bunsen or spirit lamp flame, so that the sealed extremity is 2 or 3 cm. (.8 or 1.2 in.) from where the tube is of original diameter. The tube is broken by a file near where the contraction begins, and a small quantity of the dry body is introduced into the sealed extremity. The piece of tubing is now attached to a thermometer by a small rubber band. The capillary end of the tube containing the substance is placed adjacent to the bulb of the thermometer, and the tube with the bulb is placed into concentrated sulphuric acid in a beaker which is gradually heated. As the mercury ascends to 204° C., the substance will begin to show evidence of fusion, providing the increase in temperature be gradual and the heat be equally diffused by stirring the acid with a glass rod.

Ultzmann's Modification.—Dr. A. K. Bond, of Baltimore, publishes * the following simple modification of this test, suggested by Ultzmann, of Vienna:—

and, after a proper interval, three to twelve hours, examine the sediment clear or not makes no difference. Then cover the tube and let it stand, an equal height. Upon this pour the urine-clear or cloudy-until the about 10 of an inch, and add crystals of sodium acetate, ground fine, to the constant presence of reddish globules in the field. My friend Dr. salt is always in excess when the test is made in this way is shown by ing power of 200 diameters is sufficient for this study. That the phenyl bright yellow needles of phenyl-glucosazon, which branch out or are with the microscope. boils, and continue boiling for about half a minute-whether it becomes acetate are dissolved; heat gently over a low flame until the mixture acetate, 15 parts urine. Shake the tube until the crystals of sodium following proportions in weight: I part phenyl salt, 2 parts sodium tube is half full. This gives in a test-tube five inches long about the like delicate sprays or sheaves, or radiating from a centre. A magnifyoined by others as they are formed, until the field is dotted with groups Pour into a test-tube five inches long the phenyl salt to the depth of If sugar is present, there will be seen first fine,

^{*} Medical News, Aug. 6, 1887

Purdy informs me that phenyl-hydrazin, either the salt or solution, is apt, when brought in contact with the fingers, to produce a painful

The Alphanaphthol and Thymol Test.—This recent test for sugar as directed by Molisch is made by adding to 1 c.c. of the fluid to be tested 2 drops of a 15 to 20 per cent. solution of alphanaphthol, and after mixing, an excess of concentrated sulphuric acid. Upon shaking, if sugar is present, a deep violet color is developed, and on dilution with water a violet blue precipitate occurs, soluble in alcohol and ether with a yellow color, and in potassium hydrate with a deep yellow. If the alphanaphthol is replaced with thymol a deep red color is produced, and on dilution with water a carmine-red flocculent precipitate, soluble with a pale yellow color, in alcohol, ether, and potassium hydrate, but with a bright yellow in ammonium hydrate. These tests, according to Molisch, are the most delicate known, detecting sugar in solutions containing coor fer cent.

It is claimed that most sugars, to which inosit is an exception, respond to these tests, as do most glucosides except indican. Urea, creatinin, xanthin, uric acid, allantoin, hippuric and succinic acids, phenol and pyrocatechin all give negative results. Normal urine, however, responds when diluted 300 times, from which Molisch concludes it contains sugar. He gives the following method of distinguishing normal from diabetic urine. 1. Dilute a specimen of normal urine and one of the urine to be tested for sugar with 100 volumes of water, and compare the colors resulting from the application of the tests. 2. Dilute two similar specimens with 300 or 400 volumes of water. The diabetic urine will still respond to the test while normal urine fails.

Seegen has carefully examined these tests and declares them less sensitive than Trommer's, while various animal substances and secretions and chemically pure albuminous bodies, as peptone, serum albumin, egg-albumin, and casein all reacted. Hence he says they are of no value as tests for sugar alone, and Molisch's conclusion that glucose is a constituent of normal urine is not justified.

In reply to Seegen, Molisch reasserts his original claim, and says that in dilute solutions it is necessary to employ a small quantity of solid alphanaphthol in place of naphthol solution. With reference to albu-

minous bodies, Molisch says that while these may give results resembling somewhat those obtained with sugar solutions, still, the precipitates obtained upon dilution with water are, excepting in the case of peptone, all of a different color (dirty yellow or yellowish brown) from those produced with sugar solutions. Besides, they are all soluble in hydrochloric acid with a carmine-red or reddish-violet color, while the precipitate obtained with sugar solutions is insoluble in hydrochloric acid.

Molisch also states that in place of sulphuric acid, hydrochloric acid may be employed in these tests, the mixture being subsequently boiled for a minute.

Fibrine, vitellin, serum albumin, egg-albumin, and peptone do not give the reaction when hydrochloric acid is used. Normal urine gives the reaction when boiled with alphanaphthol and hydrochloric acid even if diluted ten times. Molisch still asserts that normal urine must contain sugar, or otherwise some body as yet unknown.

Moore's Test.

Moore's test depends upon the fact that grape-sugar, with which diabetic sugar is identical, becomes oxidized when boiled in contact with caustic alkali, taking the oxygen from the atmosphere.

To a small quantity of urine in a test-tube, add half as much liquor potasse or liquor sodae, and boil. If sugar is present a yellowish-brown color soon makes its appearance, which becomes more intense as the boiling is continued, and which will be the deeper the larger the proportion of sugar, becoming finally almost black if the quantity is very large. The coloration is due to the formation, first, of glucic, and finally of melassic acid, both of which remain in solution. The flaky precipitate which is observed after the addition of the alkali, and is increased on the application of heat, consists of the earthy phosphates, which, if very abundant, may be filtered off before the heat is applied.

If now, to the colored fluid, a few drops of nitric acid be added, the brown discoloration disappears, and the odor of caramel or of burnt molasses is developed.

66

Precautions.—I. Solutions of soda and potash are liable to become impregnated with lead, either from being kept in flint glass bottles, or from the glazed earthenware vessels in which, during preparation, they are evaporated. Such contamination always causes the production of a brown and black color when boiled with organic matter containing sulphur, due to the formation of sulphuret of lead. This error may be avoided by first ascertaining the purity of the alkaline solutions, and afterward keeping them in green glass bottles.

2. If the urine exhibits already a high color, which is, however, very rare with diabetic urines, the coloring matter may be precipitated by solution of acetate (sugar) of lead, or by filtering through animal charcoal. The former throws down a small quantity of sugar, and the

3. The coloring matters of bile in urine, either when pure or decomposed (that is, when they respond neither to Gmelin's nor Heller's test), produce a brown color with liquor potassa or soda without the application of heat.

latter retains a little.

4. Bödeker found in the urine of an adult a substance which he calls alkapton, which, when strong solutions of alkali are added, produces a brown discoloration from above downward. This, according to him, also reduces the salts of copper, but does not affect the bismuth salts.

The delicacy of the test is put down at 0.3 per cent. glucose.

Polarimetry.


The most convenient and the quickest of the methods for the quantitative determination of glucose, when the quantity exceeds 1 per cent., and when all the appliances are at hand, is by polarimetry. Smaller amounts can of course be detected—it is said or of 1 per cent. with the best instruments by those who are expert. The costliness of the apparatus, however, will probably always be in the way of its general use.

The process is based upon the fact that glucose rotates

After 270° it again returns until 360°, or the starting point, is reached. ally diminished until 270° is reached, when it is again totally absent, again passes unobstructed. Continuing the rotation, it is again gradubegins to return in increased quantity until 180° is reached, when it reaches 90°, when it is totally cut off. After 90° is passed, the light rotated on its axis, the light gradually diminishes, until the rotation through the analyzer without obstruction. If, however, the analyzer be such that the ray vibrating in one plane is absorbed while the other prism or analyzer in the eye-piece OH is so placed in relation to the passes through. The latter is therefore polarized. Now if the second ing at right angles to each other, but the construction of the prism is polarizer that their oblique ends are parallel, the polarized ray will pass A ray of light entering the polarizer is divided into two parts vibrat-

it is partly obstructed. By turning this prism on its axis the light may diabetic sugar. sugar. If the tube is 10 cm. long, we have simply to multiply the specific rotary power of glucose, and the length of the tube, together column determine the degree of deviation. Hence, knowing the with the angle of deviation, it is easy to calculate the percentage of again be transmitted. The quantity of sugar and the length of the degrees of rotation by 100 and divide by 53.1, the specific rotation of light is rotated by the sugar so that when it reaches the second prism unobstructed a column of glucose is placed, the plane of vibration of Now if between the prisms so arranged that light passes through

is passing through is so great that some special means is provided for determining the precise moment at which the maximum quantity of light one or the other is turned the prism is turned. But the difficulty of either to the graduated arc of the circle or to the vernier, so that when In most polarizing saccharimeters the analyzing prism is attached



recognizing it. In Laurent's so-called "shadow" instrument, a quartz plate is so interposed that when the prisms are placed in certain relation, half of the field of view is dark and the other light; in other positions both halves are equally light, while in intermediate relationships intermediate shades appear, as in Figure 10.

In using Laurent's instrument the two prisms are adjusted by means of the screw F, which moves the analyzer until the field is uniformly illuminated on both sides. The vernier should then read o. The suspected wrine, freed from albumin, if this should be present, is treated with a solution of basic acetate of lead in the proportion of x to xo of urine, and filtered. The tube of known length is then






filled with the mixture, and placed in position between the two prisms. Immediately light passes through the analyzer in OH, and this must be again turned by means of the screw F, until the light is again stopped and the field is uniformly illuminated. The angle is now read off on the dial-plate C and the percentage calculated—if the tube is 10 cm. long, by multiplying the degrees of rotation by 100 and dividing by 53.1, the specific rotation of diabetic sugar. To this result should be added $\frac{1}{10}$ to allow for the 100 c.c. of lead solution* used.

^{*}The urine may also be clarified by filtering through animal charcoal, in which event, of course, no allowance of this kind need be made.

103


Until recently a sodium flame has been the source of light, but an ordinary gas-flame now suffices. A new vernier is interposed between the analyzer in OH and the

FIG. 11.



dial-plate C, Fig. 9, and we have simply to multiply the reading increased by $\frac{1}{10}$ (allowance for the lead solution) by the figures .2051, a factor arrived at by experiment as

F16, 12,



that which multiplied by the reading gives the percentage when a tube 200 millimetres long is used. Thus,

suppose the reading to be 24.9, then 24.9 + 2.49 = 27.39, and 27.39 \times .2051 = 5.61 per cent.

In the instrument of von Fleischel, in use by the writer, and with which a condensed kerosene flame is used, when adjusted and the vernier reads o, two spectra are placed one above the other, with the colors in each exactly continuous, the dark band proceeding through the centre of both, as shown in Fig. 11.

When the column of sugar is interposed the relationship of the spectra is at once altered, as shown in Fig. 12.

The proper relationship is again restored by moving the vernier to which the analyzing prism is attached, and the *percentage* of glucose is at once read off.

For qualitative testing the polarizing saccharimeter is less convenient than Fehling's solution, indeed will not detect as small quantities. Its greatest convenience appears in quantitative determinations where quantities of over one per cent. are to be measured. This instrument is less costly than Laurent's.

Remarks on the Qualitative Testing of Urine suspected to contain Sugar.

There has been more criticism of the methods employed in testing urine for sugar and the results obtained from them than is justified. It arises partly from the fact that the chemist and the clinical physician view the subject from different standpoints. The former makes his experiments with test-fluids upon pure aqueous solutions of sugar, the latter upon a fluid containing numerous other organic constituents, more than one of which is capable of influencing certain test-solutions, though it may be in a less degree than sugar. Let us suppose the chemist to have made solu-

do it. On the other hand, no one can deny that it is sugar, because a very small amount of sugar will do the same thing. Further, it is easy to show that a quantity of sugar so small that it cannot be detected in urine can be readily detected in water, though it is also true that minute traces of sugar can be extracted from urine by certain processes, one of which will be detailed below. But these nor is it necessary that they should be. For such amounts of sugar have no more clinical significance than has the normal proportion of urea in a specimen. It is only tangible amounts of sugar which are significant, and these are practiced with due precaution. With all tests, there are dilute, when it is boiled with it. Yet no one can claim that it is sugar which produces the decoloration when there may be several organic substances in the urine which can processes are not often available for the practical physician, ordinarily recognizable by any one of the tests named, certain very evident reactions and certain doubtful ones, and to interpret these last experience must often be relied knows that it is sugar which thus reacts, because he has But it is very different with the clinician who tests urine suspected to contain sugar. There is no urine which will not partially decolorize a Fehling's solution sufficiently while pure water will have no such effect. The chemist nothing but the test fluid, sugar, and water in combination. tions of grape-sugar of different strengths which he is testing with Fehling's solution diluted in the manner indicated on p. 77. He tests weaker and weaker solutions, and finally reaches one by which the test fluid is just decolorized

Exact Qualitative Method of Testing for Glucose—Brücke's Lead Process.

While any one of the methods just detailed suffices, if carefully used, to determine qualitatively, with sufficient accuracy for clinical purposes, the presence of sugar in a given specimen of suspected urine, it sometimes happens, especially in watching the course of a case of diabetes mellitus under treatment, that the physician desires to know with absolute certainty whether there is even a trace of sugar present. This can be done by the following method, which is a modification of the one originally proposed by Brücke, and used by Pavy to demonstrate the presence of sugar in normal urine:—

Take 50 c.c. of urine and add 60 c.c. of a ro per cent. solution of neutral acetate of lead. The precipitate which takes place includes sulphates, phosphates, carbonates, coloring matters, and part of the uric acid and creatinine, while the sugar remains in solution. Separate the precipitate on a filter, and treat the filtrate with an excess of ammonium hydroxide. A further precipitate occurs, which contains the sugar in combination with lead as a plumbic saccharate (PbO)₃(C₆H₁₂O₆)₂.

This precipitate is then collected and washed, great care being taken to remove all the ammonia, which is best effected by repeating several times the subsidence and decantation before throwing the precipitate on the filter; after which water is passed through until red litmus paper is no longer turned blue by the filtrate.*

^{*}Dr. Pavy cautions against washing with hot water. Although it expedites the process, there is danger of breaking up the combination between the plumbic oxide and the sugar, thereby losing some of the latter.

The precipitate, suspended in about 100 c.c. of water, is decomposed by passing through it a stream of sulphuretted hydrogen as long as a precipitate of black plumbic oxide is produced. Filtration is then performed and the excess of sulphuretted hydrogen expelled by heat. The filtrate, a colorless fluid, is then evaporated over a waterbath to a volume equal to that of the original urine operated upon, that is, 50 c.c.

Dr. Pavy recommends the testing of the fluid thus obtained by the Fehling's solution, but Prof. Wormley has discovered that it still contains uric acid which is capable of producing a neat reaction with the cupric test. This uric acid he removes by simply allowing the fluid to stand twenty-four hours or longer, at the end of which time a considerable sediment of uric acid has fallen to the bottom of the vessel; and it is this acid which produces the reaction in urine in the hands of Brücke, Bence Jones, and Pavy, and ascribed by them to sugar. When the uric acid is wholly removed from normal urine, no reaction occurs; but if the slightest trace of sugar is present, it occurs, and the sugar is detected.

About 50 per cent. of the glucose is lost in the course of the lead process.

IX. OTHER SACCHARINE SUBSTANCES.

Inosite.

Inosite or muscle sugar is sometimes found in the albuminuria of nephritis as well as in diabetes mellitus, and in the latter disease it has been found occasionally to substitute glucose, especially during convalescence; also in phthisis, the syphilitic cachexia, and in typhus fever. Gallois examined the urine of 102 patients for inosite and found it in

seven only; five times in 30 cases of diabetes along with sugar in variable quantity, and twice in 25 cases of albuminuria.

to cool. If, at the end of 24 hours, groups of inosite crystals appear, they are separated and washed with a little cool alcohol. tate occurs, the hot solution is filtered through a hot funnel and allowed solution is simply poured off. But if a flocculent non-adhesive precipiheavy precipitate adhering to the bottom of the glass, the hot alcoholic boiling with three to four times its volume of alcohol. If there arises a filtered out, the fluid concentrated as far as possible and treated while rated, after standing some time, a small quantity of uric acid. This is posed with sulphuretted hydrogen. Out of the filtrate there is sepacollected after twelve hours, washed, suspended in water, and decom precipitate which contains the inosite combined with the lead oxide is one-fourth its bulk over a water-bath before precipitation. The lead and the warmed filtrate treated with basic acetate of lead so long as a precipitate occurs. It is better, however, to concentrate the urine to neutral acetate of lead until precipitation ceases. It is then filtered, albumin that may be present is removed, the urine is treated with Recognition.-Inosite is obtained from urine as follows: After any

In this case it is advisable, in order that there may be no loss of inosite, to dissolve the precipitate obtained by the addition of hot alcohol in as small a quantity as possible of hot water, and precipitate a second time with three or four times its volume of hot water. If no crystals separate, ether is added to the clear, cold alcoholic filtrate, until on shaking thoroughly a milky cloudness appears, and then the fluid is permitted to stand in the cold 24 hours. If not too small amount of ether is added (an excess does no harm), all the inosite present is precipitated in pearl-like, shiny plates.

Inosite differs from cane-sugar in not undergoing vinous fermentation when treated with yeast, although its solutions readily take on the lactic fermentation when brought in contact with putrefying cheese. It does not reduce cupric tartrate in solution with potassic hydrate, but changes it to an olive-green, and after a while a flocculent precipitate falls, and the supernatant fluid becomes blue, but on again heating the solution the olive-green color is redeveloped. This reaction is sometimes observed in treating urine with Fehling's solution, as was originally pointed out by Dr. Ralfe, and it has been suggested by Dr. Oliver that it may be due

to inosite. Should this be true, inosite would appear to be less rare than has been heretofore supposed, but it is doubtful whether a reaction so indistinctive should be allowed much stress as compared with the exact chemical process above described.

Fruit-sugar, or Lævulose.

Fruit-sugar sometimes occurs in urine, but only in association with diabetic sugar. It is characterized by being non-crystallizable, and turning the plane of polarized light to the left instead of to the right. The specific rotary power diminishes as the temperature rises, while that of grape sugar is independent. It is -108.3 at 0° C.; -99.44 at 14°: -97.1 at 17.5°; -52.5 at 87.2°, according to Tuchschmied. It reduces the salts of copper as does glucose, but in a less degree.

Recognition.—Since levulose is always associated with glucose when it occurs in urine, and both reduce the salts of copper, it is only by their opposite action on polarized light that they can be distinguished. If, therefore, a sugar-holding urine deflects polarized light strongly to the left, or if, after allowing for the fact that normal urine turns it to the left to, and that this property is increased by the ingestion of large doses of benzol, phenol, bromo- and nitro-benzol, chloral, and camphor, polarized light is not deflected at all, we may conclude that leevulose is present. It is to be remembered that these substances also reduce copper salts. Other substances turning polarized light to the left must likewise be excluded.

Sugar of Milk, or Lactose. CuH2011 + H20.

Lactose is sometimes found in small quantity in the urine of nursing women and females of animals. Dr. Raffe refers to a case under his observation in the London Hospital, of a young married woman, aged 29, who was suckling an infant, and suffering with debility and frequent micturition, whose urine contained as much as 3 per cent. of sugar. This occurred in three successive confinements, there being no sugar during pregnancy.

Lactose is characterized by crystallizing in white, or colorless, foursided prisms with acuminated ends bounded by four triangles, by its turning polarized light to the right with a rotating power of $+59.3^{\circ}$,

varying also with the degree of concentration; while that of grape-sugar is + 53.1°; by the fact that it reduces the salts of copper as does grape-sugar, and that it does not undergo the alcoholic fermentation with yeast. The fungi which convert milk-sugar into alcohol are cleft fungi. On the other hand, the lactic acid and butyric acid fermentation are readily entered upon.

Recognition.—A very strong reducing power and an otherwise inexplicable deflection to the right suggest milk-sugar. Especially is this suspicion justified if the urine is that of a nursing woman. It can be recognized with certainty only by isolating it with the urine, for the details of which the student is referred to the works on physiological chemistry by Gorup-Bezanez or Hoppe-Seyler, or the elaborate treatises on urine analysis by Neubauer and Vogel or Salkowski.

X. ACETON AND ACETONE-PRODUCING SUBSTANCES; DIACETIC ACID.

Often closely, though not necessarily associated with glycosuria, are those conditions which respond to the tests for acetone.

Aceton.

A peculiar fruity odor of the urine was occasionally noted by the earlier students of diabetes, who were ignorant, however, of its cause, and it was not until 1857 that Petters* recognized in certain urines, by its peculiar chemical behavior, the substance known as aceton, to which this odor is probably due. Kaulich† extended and completed the work of Petters. Numerous other studies followed, the resultant of which was the fact above men-

^{*}Petters, "Untersuchungen über die Hönigharnruhr." Prager Vierteljahrschrift, 1857, XIV, Band 3, S. 81.

[†] Kaulich, "Ueber Acetonbildung in thiereschen Organismus." Prager Vierteijahrschrift, 1860, XVII, Band 3, S. 58.

tioned, that while in diabetes mellitus aceton often occurs in the urine, it is found also in the urine in other diseases and in small amount even in health. Further, not all urines containing aceton exhibit the fruity odor. Contrary to what was at first thought, aceton does not arise from the glycogen in the blood or urine, but from albumin disintegration.* Such, of course, could only be its origin where there is no glycemia or glycosuria. In confirmation of this is the fact that physiological acetonuria is enormously increased by a meat diet.

Tests for Aceron.—The accurate study of aceton is best secured by working with the distillate from the urine. By Legal's test, however, even moderate quantities may be recognized in the urine itself, to which, indeed, the test is better adapted than to the distillate, because parakresol, which also passes over with the latter, strikes a similar reaction. For this reason it is also employed as a preliminary test, even when the distillate is afterward employed.

Legal's Test.—A fresh, rather strong solution of sodium nitro-prusside is made by dissolving a few fragments in a little water in a test-tube. To three or four centimetres of the suspected urine add enough liquor sodæ or liquor potassæ to secure a distinct alkaline reaction. To the mixture then add a few drops of the nitro-prusside solution, when promptly the whole assumes a red color whether aceton is present or not, said to be produced by creatinine even more rapidly than by aceton. In any event the red color disappears, but if aceton is present, the addition of a few drops of concentrated acetic

^{*}V. Engel, "Mengenverhältnisse des Acetons unter Physiol. und Pathol." Bedingungen, Zeitschrift für klin. Med., Band xx, S. 521, 1802.

acid causes a purple or violet red. If there is no aceton, this final change does not occur. The purple color also fades in a little while, even if caused by aceton.

On standing a dark blue precipitate may fall, and may be hastened if, after the addition of acetic acid, the solution is warmed.

Legal's test may also be applied to the distillate, but for reasons named is less satisfactory.

To Test the Distillate.—For the study of aceton in the distillate, treat a half litre or a litre of urine with phosphoric or hydrochloric acid in the proportion of 3 cubic centimetres of the acid to each 100 c.c. of urine. Thus treated, the urine is partially evaporated in a distilling apparatus, and the process completed in a retort. The addition of the acid is made to prevent the evolution of gases. Von Jaksch prefers phosphoric acid.

From ten to thirty centimetres of the distillate are thus obtained, and to it may be applied the following tests:—

Lieben's Iodoform Test.—To a portion of the distillate add a small quantity of liquor potassæ, then a few drops of a solution of iodine and iodide of potassium.* If aceton is present a yellowish precipitate of iodoform is produced at once. Should alcohol happen to be present in the distillate, the reaction takes place also, but more slowly, but with aceton it is immediate.

Gunning's Modification of Lieben's Iodoform Test.—The latter source of error may be altogether avoided by the modification suggested by Gunning, who uses ammonium hydrate and tincture of iodine. With these, alcohol causes

^{*}Friedlander's formula for this solution is, iodine 1.0; potassium iodide 2; distilled water 50.

no precipitate, while aceton does produce one of iodoform in addition to iodine, which falls as a black sediment, even though there be no aceton. The precipitated iodine is soon redissolved if the urine contains much aceton; if it contain but little, the iodoform crystals may be seen in twenty-four or forty-eight hours, resting in a thin layer upon the black precipitate of iodine.

This test may also be used with the urine itself, but with satisfaction only if the quantity of aceton is large.

Reynold's Test.—This test depends on the power of aceton to dissolve freshly precipitated mercuric oxide. It is performed as follows: From a solution of mercuric nitrate in a test-tube yellow mercuric oxide is precipitated by an alcoholic solution of potassium hydroxide. The solution supposed to contain aceton is added and the mixture shaken. The liquid is filtered, taking care that the filtrate passes through clear. To the latter, ammonium sulphide is now carefully added. If aceton is present some mercuric oxide is dissolved, which passes over in the filtrate and shows itself as a dark ring of mercuric sulphide at the point of contact of the two liquids.

The Indigo Reaction of Bayer and Drewsen.—
Heat a few crystals of nitro-benzaldehyde until dissolved. Allow the solution to cool, when the aldehyde separates as a white cloud. Then add the suspected fluid or preferably its distillate, and make the mixture distinctly alkaline with dilute caustic soda. If aceton is present, there appears first a yellow, then a green color, followed by an indigoblue, in the course of ten minutes. If only traces of aceton are present, the yellow fluid is shaken with a few drops of chloroform, when a distinct blue coloration of the chloroform takes place.

By this method it is said aceton can be easily detected in a dilution of r part to 2500, if the distillate is used. Even with undistilled urine, by the aid of chloroform, it can be detected if present in the proportion of r to rooo.

Pyroracemic acid, aldehyde, and acetophenon are the only other substances producing the indigo reaction, and these have not as yet been found in urine.

Diacetic Acid.

In 1865 Gerhardt ** discovered that the urine in certain cases of diabetes mellitus struck a red reaction with a solution of ferric chloride. This reaction was ascribed to diacetic ether or ethyl diacetate, a substance which splits up readily into aceton, alcohol, and carbonic acid. This appeared reasonable, because both alcohol and aceton had been found in diabetic urine. Deichmuller † and Tollens, † however, eventually showed this substance to be diacetic acid, also obtainable from acidulated urine by the action of ether. Other substances, not usually constituents of urine, which strike the same reaction are said to be formic § and acetic acids and the cinchona salts; also carbolic and salicylic acids, decomposition products of antipyrin, kairin, and thallin. Aceton itself does not respond to the chloride

^{*}Gerhardt, "Diabetes Mellitus und Aceton." Wiener medizinische Presse, 1868, Band vi, No. 28.

[†] Deichmuller, "Ueber diabetische Acetonurie," Liebig's Annalen der Chemie, Band 209, S. 22.

[‡]Tollens, "Ueber Eisenchlorid roth farbende Harne." Liebig's Annalen, Band 209, S. 30.

[¿]Le Noble, who also found formic acid in three out of seven cases of diabetes. American Journal of the Medical Sciences, January, 1887.

of iron test, although diacetic acid responds to the different tests for aceton.

has been boiled shows little or no change, while the chloride of iron reaction with the ethereal extract pales after twenty-four to forty-eight hours, and the urine as well as the distillate is found to contain large quantities of watery solution of chloride of iron. If a precipitate of phosphates is produced, remove it by filtration, and to If a Bordeaux red color develops, heat a portion of the urine to boiling, and acidulate a second portion with sulphuric acid and extract with ether. If the urine which is directed by von Jaksch as follows: To urine, as fresh as the filtrate again add more of the chloride of iron solution. It is plain, from the above, that while the striking of a red reaction on the addition of chloride of iron to urine it is not conclusive. To secure greater certainty, this test possible, add cautiously a few drops of a moderately strong is presumptive evidence of the presence of diacetic acid, The Chloride of Iron Test for Diacetic Acid.aceton, diacetic acid is present.

tion is very vivid, and the conditions favorable to the presence of diacetic acid are present, its presence may be inferred from the primary iron reaction alone. It is further important, however, that the urine should be fresh or decomposition be prevented, because diacetic The precautions described are necessary because of the possible presence of the above-named substances, which exhibit the same reaction. At the same time if the reacacid is so quickly converted into aceton.

fade away, the substance, according to Minkowsky, is If the color reaction with the ethereal extract does not

 β -oxybutyric acid, of which diacetic acid is said to be a further oxidation.

Quantitative Estimation of Aceton.—More than one method of quantitative estimation for aceton has been practiced, but the most satisfactory is that of Messinger, applied to urine by Huppert and further developed by Engel and Devoto.

solution of sulphuric acid (1 to 8) and submitted to original volume is carried over. A portion of the and the flask connected by a long glass tube with the up to 100 c.cm. by the addition of distilled water. Two The distillate is then treated with I c.c. of a diluted presence of aceton, the result is rejected and the process The distillation is continued until nine-tenths of the All the attachments must, of course, be closely made. in front of that a bullet apparatus filled with urine. cooler, in front of which a distilling flask is placed, and c.c. of a fifty per cent. solution of acetic acid are added 100 c.c. of urine are placed in a boiling flask and increased determined. A preliminary test is made by Legal's method, redistillation. recommenced after the addition of more distilled water. residue is tried with Lieben's test, and if it shows the when, according to the result, 20 to 50 c.cm., or at most lected, accurately measured, specific gravity and reaction Process .- The urine for the twenty-four hours is col-

The second distillate is poured into a flask of one-litre capacity, provided with a ground-glass stopper, but filled during distillation with a doubled perforated cork, and having also a bullet apparatus full of water in front of it. Immediately after the distillation the flask is closed

SIGNIFICANCE OF ACETON AND DIACETIC ACID. 117

with its glass stopper and titrated with the one-tenth normal iodide solution and the one-tenth hyposulphite solution. One c.c. of the iodo solution corresponds to 0.967 milligram aceton.

The addition of the 2 c.cm. of fifty per cent. acetic acid solution is made to prevent the passing over of phenol, and the later addition of the 1 c.cm. of the 1 to 8 sulphuric acid solution to prevent the transit of ammonia, both of which are disposed to take up iodine themselves.

Clinical Significance of Aceton and Diacetic Acid.—We are indebted to von Jaksch* for most of our knowledge of the difference in the clinical significance of these two substances, which were formerly thought to be of identical import. Nor do the very recent observations of von Engel referred to, while thorough and exhaustive, much more than confirm and extend von Jaksch's results.

It has already been said that traces of accton may be found in normal urine, constituting physiological acetonuria. Further, aceton is found in cases whose course is usually favorable, and its presence may have little or no significance. Diacetic acid, on the other hand, never occurs in normal urine, and diaceturia is a most dangerous complication.

Acetonuria seems to be a result of continued high temperature, or at least accompanies diseases attended with such temperature, and a lowering of temperature is followed by a fall in the quantity of aceton, while a re-development of high temperature is followed by an increase in aceton. Such acetonuria is known as febrile.

Acetonuria often accompanies diabetes, constituting dia-

*" Acetonurie und Diaceturie," Berlin, 1885.

betic acetonuria, but is not necessarily associated with it or with glycosuria, and while the development of acetonuria in diabetes is sometimes accompanied by very unpleasant symptoms, as headache, loss of appetite, and deranged digestion, all of short duration, it is otherwise of little significance except as a possible precursor of the diaceturia which often succeeds it in this disease.

Among other diseases with which aceton has been found associated are certain forms of carcinoma, which have not yet led to inanition; cerebral psychoses accompanied by mental excitement and derangements of digestion.

Very rarely aceton itself is responsible for a set of symptoms in which restlessness, excitement, and delirium are the most conspicuous, constituting a sort of autointoxication, the symptoms of which pass away entirely with the disappearance of the aceton.

On the other hand, what is commonly known as diabetic coma, according to von Jaksch, is the result, not of aceton in the blood, but of diacetic acid, although it is true that it is often preceded by a long-continued acetonuria. It is commonly observed in cases of far-advanced diabetes, constituting diabetic diaceturia. There is added to the usual feeling of weakness and depression, drowsiness, which may deepen into coma or pass away, the diaceturia continuing. Sometimes these symptoms are preceded by excruciating muscular pain. As a rule, there is no relation between the amount of sugar and diacetic acid eliminated, although a sudden diminution of glycosuria is sometimes followed by the appearance of a large amount of diacetic acid, by coma and death.

There is also a febrile diaceturia which accompanies febrile diseases, among which are the acute exanthemata, peri-

carditis, pleuritis, perityphlitis, typhoid fever, miliary tuberculosis, tubercular phthisis and pneumonia. It is to be remembered, however, that symptoms of diabetic coma may occur without either acetonuria or diaceturia. Von Jaksch proposes to do away with the term "diabetic coma" and substitute "coma diaceticum" for all of those cases of coma, from whatever remote cause, accompanied by diaceturia.

Finally, there appears also to be an *idiopathic diacetemia* or auto-intoxication by diacetic acid, unattended by other grave disease, such as pneumonia or diabetes, and manifested by vomiting, dyspnœa, and jactitation, which soon terminates in coma and death. This condition, very grave, but rare in adults, is said by von Jaksch to be much more frequent in children and correspondingly less serious. In such cases the child feels weak, has a thickly-coated tongue, often slight conjunctival catarrh, there is sometimes vomiting, usually constipation, and very little or no fever. In two or three days all of these symptoms, together with the diaceturia, disappear. In other cases nervous symptoms are more marked. Von Jaksch believes that all of these, as well as a certain number of other convulsive attacks in children, are the result of auto-intoxication with diacetic acid.

It is evident that our knowledge of these substances, aceton and diacetic acid, as well as that of the symptoms developed by their presence, is not yet complete; but the statements just made may be considered as representing as definitely as possible our present information.

β-ΟΧΥΒUTYRIC ACID.—The above statements are not seriously affected by the studies of Hugounenq,* Lepine,

^{*} American Journal of the Medical Sciences, October, 1887.

cetic acid, aceton. The recommendation of Hugounenq the changes may be brought about outside of the body is, and says an exclusive meat diet is to be avoided as favormanent benefit. Lepine also advises an alkaline treatment, glucose, alcohol, aldehyde, aldol, \(\beta\)-oxybutyric acid, diadation of it. Just by what steps it is formed within the diabetic patients, is the homologous superior of lactic acid, ing the development of coma. tralize the acid in the blood has been followed without perto use hypodermic injections of alkaline solutions to neuhowever, very simple; the series then running as follows: body we are unable to discover. The method by which and is formed from diseased muscle, just as lactic acid is vations. Beta-oxybutyric acid is found in the blood of and which are apparently confirmed by more recent obserstance and not diacetic acid is the cause of diabetic coma, from healthy muscle. Diacetic acid is only a further oxi-Stadelmann, and Minkowski, to the effect that this sub-

These observers say, too, the acid found in the urine in many cases of diabetes is not diacetic acid, but oxybutyric acid.

As to the relation between the two substances, aceton and diacetic acid, the conclusions of von Jaksch, published in 1885,* are essentially true to-day, viz., that (1) aceton is an oxidation product of albuminous substances; (2) it is found largely increased in the organism under certain pathological states and then may cause toxic symptoms. If the quantity of aceton produced is enormously large, it unites with certain acids produced in retrograde albuminous metamorphoses, perhaps with formic acid only, produc-

^{* &}quot; Ueber Acetonurie und Diaceturie," Berlin, 1885.

ing diacetic acid, perhaps also in part with another series of similar acids, which may even be such volatile substances as are contained in evaporated urine and out of which by oxidation aceton may arise.

XI. COLORING MATTERS.

The pathological significance of all the coloring matters has not, as yet, been determined. Many of them are, however, of such importance that their consideration commands interest next to that of the proteids and sugars.

I. Normal Coloring Matters.

Notwithstanding the very considerable study given to this subject of late years, there is still much confusion as to the normal coloring matters of urine. The two as to the existence of which there is most unanimity are the chromogen* of Urobillin, and Urine-Indican, the chromogen of indigo, the latter being the uroxanthin of Heller.

Thudichum † makes a single coloring matter, which he calls urockrome. To this matter, according to Thudichum, the urine owes the whole or greater part of its yellow color, while numerous other coloring matters, including the urrhodin of Heller, Scherer's urohæmatin, and the urohæmatin of Harley, he considers mixtures of the products of decomposition of this yellow pigment.

The urohæmatin of Scherer and that of Harley are

^{*} Chromogens are substances which develop a color on the addition to the fluid of some oxidizing agent.

[†] Thudichum, "A Treatise on the Pathology of the Urine," 2d edition, London, 1877.

probably identical, Scherer * admitting that urohæmatin contains iron, and approving of the use of the term by Harley for his coloring matter. The urophain of Heller is probably the same thing. It will at any rate here be so considered. Finally, all of these are probably modifications, due to different methods of treating urine, of the substance known as urobilin, now generally acknowledged to be the most important coloring matter of the urine. Upon the presence of indican (Heller's uroxanthin) in most normal urines, all are agreed, although Thudichum prefers not to consider it a coloring matter, but a chromogen or color generator. For the present I shall retain it among the normal coloring matters, treating, therefore, chiefly of two, viz.:—

 urobilin (Jaffé), hydrobilirubin of Maly, and its modifications; urohæmatin of Harley and Scherer; urophain of Heller.

2. Urine-indican or the uroxanthin of Heller; to which will be added a short account of urochrome as described by Thudichum.

1. Urobilin-Hydrobilirubin-Urophain-Urohamatin.

Urobilin, first extracted by Jaffé, and further studied by Maly, is believed to be the bilirubin or normal coloring matter of the bile—altered after passing into the small intestine, by absorbing water and hydrogen—in a word, reduced bilirubin. Hence Maly names it hydrobilirubin. The reaction may be expressed as follows:—

$$2(C_{16}H_{18}N_2O_3)\,+\,H_2O\,+\,H_6=C_{22}H_4N_4O_7.$$

^{*} Harley, "The Urine and its Derangements." Philadelphia, 1872, from London edition, 1871.

As such it is reabsorbed and excreted by the kidneys. As bilirubin is itself the hæmatin of the blood, reduced by the high-colored fever urines by processes described in the and chloroform. It gives no play of colors with nitric trated they exhibit peculiar spectroscopic properties and a beautiful green fluorescence by reflected light. The spectrum is a dark absorption band between Fraunhofer's lines b and F. Both properties become more distinct by the addition of solution of ammonia and a drop of chloride chloric acid, and the absorption band recedes toward F action of the bile acids, the direct descent of urobilin from the blood is established. Obtained most readily from larger works on urinalysis, it is a brown, resinous mass, easily soluble in water, but more readily in alcohol, ether, acid. Its concentrated solutions are brown; more diluted they are yellow, and still more, rose-red. When concenof calcium, while the fluorescence ceases on adding hydroand becomes more indistinct.

Jaffé has inferred from the absence of these peculiar reactions of urobilin in fresh urine that urobilin is not at first present, but is preceded by a chromogen, which is converted into urobilin on exposure, by absorbing oxygen.

Test for Urobilin or Hydrobilirubin.—Add ammonia until distinctly alkaline, filter, and to the filtrate add a little chloride of zinc solution. The appearance of a green fluorescence and the characteristic absorption band indicates the presence of a considerable amount of bilirubin.

It is especially abundant in the high-colored urine of fever cases, heart and liver diseases, and after sweating.

Heller's test for urophain is as follows: About 2 c.c. (32.4 minims) of colorless sulphuric acid are poured into a small beaker-glass, or, better, a "collamore" wineglass

(p. 17), and upon it in a fine stream, from a height of about four inches, twice as much urine is allowed to fall. The urine mingles intimately with the sulphuric acid, and in normal urine, of which the specific gravity is 1020 and the quantity 1500 c.c. in the twenty-four hours, produces a deep garnet-red coloration.

If the coloring matter is increased, the coloration is no longer garnet-red, but is black and opaque; whereas, if the coloring matter is diminished, the mixture appears pale garnet-red and transparent.

Precautions.—Unfortunately, other conditions than that of increased amount of coloring matter produce the increased intensity of the urophain reaction. Thus diabetic urine produces the same dark opacity through carbonization of the sugar by the sulphuric acid. In like manner, urine containing blood, biliary coloring matters, and uroerythrin (an abnormal coloring matter), gives the same reaction with sulphuric acid. Before relying, therefore, upon this reaction, the above substances must be carefully excluded.

Dr. Harley's test for urohæmatin is as follows: Dilute the twenty-four hours' urine with water till it measures 60 ounces (1800 c.c.), or if the quantity exceeds 60 ounces, concentrate it to this amount; then to about 2 drachms (7.4 c.c.) of it, in a test-tube, add half a drachm (1.8 c.c.) of pure nitric acid, and allow the mixture to stand for some minutes. If the quantity of urohæmatin is normal, the mixture will alter but slightly in tint; whereas, if there be an excess, it will become pink, red, crimson, or purple, according to the amount present. Heating the mixture hastens the change in color, but it is better to do this experiment in the cold, and, if necessary, allow plenty of time for the change to take place.

The acid is added to liberate the coloring matter, which

125

may be so thoroughly concealed that a pale urine often contains a large amount of urohematin.

Harley gives a second method, also easy of application, for determining an excess of urohæmatin in cases of destructive diseases of the blood. Boil 4 ounces (120 c.c.) of urine, and add nitric acid to set the coloring matter free. When cool, put the urine in a six-ounce bottle along with an ounce of ether. Cork the bottle, thoroughly shake it, and place aside for twenty-four hours. At the end of that time the ether will be found to be like a red, tremulous jelly. Such a case, however, he admits to be a bad one.

There is every reason to believe that urohamatin represents the disintegration of red blood corpuscles, and that it fluctuates, therefore, with the rate of their destruction

Urochrome of Thudichum.—Thudichum terms the substance, to which he considers the whole or greater part of the yellow color of the urine is due, urochrome. It is an alkaloid, but not of pronounced basic properties. It has been isolated, but not finally analyzed. Its principal characteristic is that on chemolysis with acids it is split up into several bodies of smaller atomic weight, one of which—uromelanin—seems to be derived from the coloring ingredient of the blood. Urochrome does not show any specific absorption band before the spectroscope when strongly acidified, but by chemolysis probably gives rise to two or three substances having distinct spectrum phenomena which greatly aid in their diagnosis. It is not the chromogen of urobilin.

Thudichum gives (op. cit.) several methods of isolating urochrome, the briefest of which consists in precipitating fresh urine with neutral and basic lead acetate, decompos-

ing the precipitate with sulphuric acid, and precipitating the urochrome and some xanthin-like body from the filtrate by phosphomolybdic acid.

Urine-indican—Uroxanthin of Heller—Indigogen of Thudichum.

of a separate chromogen which he calls urrhodingen. which is said to respond to Trommer's test, but not to the ducts was said to be indigo-glucin, a saccharine substance as the result of putrefaction. Formerly one of these prothis is denied by Thudichum. This is said to occur also said by Kletzinsky to be identical with indigo red; but of Heller), and a red coloring matter (urrhodin of Heller) with acids under warmth into indigo-blue (the uroglaucin tend to show a difference. identical with plant-indican, but more recent investigations Thudichum, urrhodin is the result of chemolysis by acids fermentation test. This is now denied. According to It has a bitter taste, and is easily converted by treatment Uroxanthin or urine-indican was formerly thought to be brown syrup, easily soluble in water, alcohol, and ether. less substance, separable from urine in the shape of a clear Indican, C₂₂H₆₂N₂O₃₄, or uroxanthin, is itself a color-

Heller's test is performed as follows: 4 c.c. or [5] of pure hydrochloric acid are poured into a smooth wine-or a small beaker-glass, and into the same while stirring 10 to 20 drops of urine are dropped. Under the normal conditions indican is present in urine in so small quantity that the acid to which the urine is added is colored pale yellow-ation is violet or blue. The more abundant the indican the more rapidly does the violet or blue coloration take place,

after such an interval indicates but a small quantity of makes the test more delicate, and small amounts of indican are thus recognized. If it is desired to test urine containing the biliary coloring matters for indican, the former or 4 c.c. hydrochloric acid. The blue color does not always make its appearance immediately. It is well then to wait 10 or 15 minutes, but the reaction which appears The addition of 2 or 3 drops of pure nitric acid must be precipitated by solution of lead acetate and filtered and often 1 to 2 drops of urine are sufficient to color f3j

posed by hydrochloric acid and calcic hypochlorite, should avoiding a great excess of the latter, when, if indican is Jaffe's method is more striking in its results, and is 3.24 f3) of urine in a large test-tube add an equal amount shaking, a perfectly fresh saturated solution of calcic hypochlorite (chloride of lime) drop by drop, until the greatest intensity of the blue color is reached. This is then shaken with chloroform, which readily dissolves the freshly formed indigo, and separates from the aqueous solution as a blue fluid, the color being more or less deep according to the amount of indican present. In pale urines, often very rich in indican, this method will serve to determine Dark urines, whose other coloring matters are also decompresent, a good indigo extract can be obtained in this way. even approximately quantitative. To 10 or 15 c.c. (2.7 or of fuming hydrochloric acid, and then, with constant its amount with sufficient accuracy for clinical purposes. first be decolorized by a solution of basic acetate of lead,

Weber's test is commended by v. Jaksch. To 30 c.c. of urine add an equal quantity of hydrochloric acid, r to 3 drops of dilute nitric acid, and heat to boiling. The mix-

ture becomes dark; after cooling, add ether and shake. If indican is present a blue froth appears on the surface, while the ether itself assumes a rose to violet color.

Precaution.—Albumin must always be separated before performing these tests, as it develops a blue color with hydrochloric acid after standing a long time. If the resulting color is red instead of blue, iodine is present, and thus the absorption of iodides ascertained.

Clinical Significance of Indican in the Urine.— Normal urine, according to Jaffé, contains 4.5 to 19.5 milligrams in 1500 c.c., or about 6.6 in 1000 c.c. It is increased by a meat diet, in obstructive diseases of the bowel, in pyelitis, diseases of the spinal cord and its membranes, and especially derangements of the entire central and peripheral nervous system, in urina spastica, after coitus, and in hot weather, probably from concentration of the urine. It is also especially abundant in the urine secreted during the reaction from cholera (Wyss).

It has been found by Neftel in cases of cancer of the liver; and its presence in large quantities in persons affected with malignant tumors he considered pathognomonic of cancer of the liver; by Hoppe-Seyler, in a case of melanotic cancer of the orbit. Jaffé finds indican increased in all diseases attended by intestinal obstruction, cancer of the stomach, lymphoma and lympho-sarcoma in the abdomen, purulent peritonitis, certain forms of diarrhoea, and in various diseases where the latter is a symptom. Rosenstein found indican increased eleven to twelve times in Addison's disease.

I found it markedly increased in two cases of cirrhosis of the liver confirmed by post-mortem examination, and in one of evident malignant disease of some abdominal organ, probably the liver; but the diagnosis was not certain and there was no autopsy. M. Robin has recently announced that he considers the presence of indican a valuable diagnostic sign in typhoid fever.*

thus produced during digestion is absorbed and converted in the blood into urine-indican. Now it is supposed that in ordinary normal intestinal digestion very little indol is ciate it pathognomonically with any disease. But recent physiological observations afford a rational explanation for its increase, which is strikingly confirmed by the clinical is introduced into the blood. It was found by Kühne (Virchow's Archiv, vol. xxxix) that during the artificial fermentation of albumin in the presence of minced pancreas, indol was produced. Jaffé suggested that the indol produced; but wherever digestion is interfered with or delayed, as is evidently likely to be the case in almost absorbed, oxidized, and excreted as indican, thus accounting for its presence in increased amount under the circum-From these facts it is evident that it is difficult to assoobservations above noted. Indican is increased when a substance known as indol (CaH,N), first discovered by Baeyer, all of the conditions above instanced, more is produced, stances.

Uroglaucine.—Apery † has recently announced that he has found *uroglaucine* in every one of twelve cases of scarlet fever, deposited in small, blue masses so distinctive that they can scarcely be confounded with any other substance. Both uroglaucine and urrhodine are sometimes found in the sediment of cystitis and Bright's disease.

^{*} Philadelphia Medical Times, October 22, 1881, p. 63. † Apery, "Les Nouveaux Remèdes," November 1, 1885.

Dr. Harley believes that all the various colored urinepigments are but different grades of oxidation of urohæmatin,* and thus accounts for the various cases of blue, green,
brown, and black urines which have been at different times
reported, a most important fact with regard to which is that
they never exhibit these colors at the moment the urine is
passed, but acquire them after exposure to the air or the
action of chemical reagents. He considers these changes
which occur in urohæmatin out of the body due to prior
changes in the body caused by disease.

He admits, however, in common with others, that some portion of the coloring matter of the urine comes from the food, chiefly vegetable food.†

My friend, Dr. S. Weir Mitchell, has called attention to a peculiar greenish or yellowish-green coloration exhibited by the urine of those who are upon a diet of skimmed milk alone. This coloration is probably such as would be expected when the coloring matter derived from the hæmoglobin of the red blood corpuscles is uninfluenced by coloring matters contained in food, but it is a subject which requires to be investigated.

II. Abnormal Coloring Matters.

Under abnormal coloring matters are included those which never enter into the composition of normal urine, whether found elsewhere in the body or not.

131

by the action of heat. Methæmoglobin is an intermediate gives the same absorption band, in the yellow of the specwhile hæmoglobin gives one band in the yellow and one Hæmatin is a deoxygenated hæmoglobin, into which and a coagulated albuminous substance the latter is converted condition, approaching, however, nearer to hæmatin, and trum between Fraunhofer's lines C and D, but nearer to D, They include (a) the coloring matters of blood, hamoglobin or oxyhæmoglobin, methæmoglobin, and hæmatin. in the green between D and E.

urine be treated by sulphuret of ammonium, it becomes reduced hæmoglobin by loss of its oxygen. Of this, the In fresh urine containing blood-coloring matters the prevailing one is hæmoglobin; but if a specimen of such spectrum gives a single broad band between the lines D and E. Shaking with oxygen or atmospheric air again restores the reduced hæmoglobin to oxyhæmoglobin.

- (b) The uroerythrin of Heller.
- (c) Vegetable coloring matters.
 - (d) Biliary coloring matters.

(a) The Coloring Matters of the Blood-Hamoglobin, Methamoglobin, and Hamatin.

These substances can enter the urine either by direct puscles themselves, after they have entered the urine in transudation, or arise from the dissolution of blood-cordifferent ways. They may be present in the urine in very small quantities without being accompanied by albumin, as was first shown by Dr. F. A. Mahomed.*

^{* &}quot;Transactions of the Royal Medico-Chirurgical Society of London," vol. lvii, 1874, p. 196.

The color of the urine is different according as it contains more hæmoglobin or methæmoglobin, the former being brighter, the latter darker, brownish-red. Hemor-rhages from the larger vessels produce more hæmoglobin; capillary hemorrhages, on the other hand, more methæmoglobin. Heller proposes to account for the difference by the fact that in the hemorrhages which take place from the capillaries in renal disease, the blood is much more slowly and more intimately commingled with the urine, and therefore longer retained with it at the normal temperature of the body. Temperature, the presence of carbonic acid, and the absence of oxygen, may favor the change of hæmoglobin to methæmoglobin.

Detection of Blood-Coloring Matters.

slobin Unaccompanied by Albumin.—Dr. Mahomed (op. cit.) directs as follows: One end of a small slip of white blotting-paper is dipped in the urine and dried over the flame of a spirit-lamp; by this means the dilute solution of the crystalloid is concentrated by evaporation; two drops of the tincture of guaiacum are then dropped on the paper, and after a minute or so, allowed for the spirit to evaporate, a single drop of ozonic ether is let fall in the centre of the guaiacum stain. A blue color appears if hæmoglobin is present. Some time, perhaps a quarter of an hour, will elapse before the reaction becomes visible, especially if it be slight; when it appears it is not permanent; it will begin to fade in a few hours, and will have disappeared in a day or two.

The advantage of this test lies in the fact that the physi-

133

cian can carry a few slips of blotting-paper in his pocketbook, dip one in the urine during his visit, allow it to dry, and make the test at home.

Dr. Stevenson's modification of Dr. Mahomed's test, acknowledged by the latter to be far more brilliant, is as follows: To a drop or two of urine in a small test-tube add one drop of the tincture of guaiacum and a few drops of ozonized ether; agitate and allow the ether to collect at the top, forming an upper layer of fluid. If hæmoglobin be present the ether carries up with it the blue color that is produced, leaving the urine colorless below. In this method the blotting-paper, which is somehow the source of fallacy, is not required.

Precautions.—Saliva, nasal mucus, and a salt of iodine (as happens when the patient is taking iodide of potassium) all strike a blue color with tincture of guaiacum, some without and some after the addition of ozonic ether.

Clinical Applications.—By this test, according to Dr. Mahomed, infinitesimal traces of hæmoglobin can be detected in urine which to the naked eye, the microscope, the spectroscope, and even to the nitric acid test for albumin, affords no indication whatever of abnormality. Indeed, the presence of albumin in any quantity interfers with the test, and it is in the *prealbuminuric* stages of scarlatina, or just after it has disappeared, and where there is a high state of vascular tension, that it is serviceable. It will respond in chronic albuminuria also, where minute traces of blood are present. Where the response precedes the appearance of albuminaria, it fades when the albumin becomes copious, and reappears again as it diminishes or after it disappears.

The most useful application of the test, according to Mahomed, is in the prealbuminuric stage of scarlatina, where it will give us information of a state of affairs in the kidney previous to actual inflammation of the organ, when a brisk purge or copious sweat may avert more serious mischief. In cases of albuminuria produced by intense fever and due to venous congestion, as in enteric fever, pneumonia, and sometimes in the febrile stage of scarlatina, when the fever is intense and the albuminuria only slight, no reaction showing the transudation of the hæmoglobin can be obtained.

such urine, although dark red in bulk, is yellowish and and giving the spectrum of hæmaturia. Again, the color of acid, the fluid becoming tinted reddish to reddish-brown, coagulum by boiling with alcohol containing sulphuric blood, and deposits a less copious sediment. blood corpuscles. It is of lower specific gravity than such more transparent in thin layers than urine containing surface. The color may be removed from the washed small, brownish, coherent coagulum which floats upon the cipitate, as does coagulated serum-albumin, but forms a which quickly coalesce and form a large, white, bulky premethæmoglobin. The former is precipitated, not in flakes up into a coagulated albuminous substance, hæmatin and a solution of hæmoglobin is heated in a test-tube it breaks derived from the decomposition of the hæmoglobin. When disks and the presence of a smaller quantity of albumin, from hæmaturia is determined by the absence of blood 2. The presence of hæmoglobinuria as distinguished

It must not be concluded that, because blood corpuscles are absent from a given specimen of urine containing hæmoglobin, they have never been present; for they are

there be a transudation of serum at the same time with the In such event we must depend upon the smaller amount of globinuria, and the smaller sediment. The urine conwhile the urine of hæmoglobin is acid in reaction. Should hæmoglobin it would, of course, be impossible to distinguish sometimes rapidly dissolved, especially in alkaline urine. albumin just alluded to as characteristic of simple hæmotaining the dissolved corpuscles is more apt to be alkaline, the two.

dichroic. If the urine is already alkaline, and no precipitate of earthy phosphate appears on the addition of liquor potassæ and heat, a precipitate can be artificially produced by the addition of one or two drops of the magnesium fluid, which, with the application of heat, carries down the 3. Heller's test for hæmatin is as follows: Precipicaustic potash and gentle heat, over a flame. The earthy ing matters, and appear, therefore, not white, as in normal urine, but blood-red. When the quantity of coloring matter in urine is very small, the earthy phosphates appear tate from urine in a test-tube the earthy phosphates by phosphates carry with them, as they sink, the blood-color-

salt is carried on the point of a knife to the dried hæmatin a thin glass cover, a hair interposed, and a drop or two of To Prepare Hæmin Crystals.-If the precipitated earthy phosphates are filtered out and placed on an objectglass, and carefully warmed until the phosphates are completely dry, Teichmann's hæmin crystals can be produced therefrom. For this purpose a minute granule of common and earthy phosphate, and thoroughly mixed with it. Any excess of salt is then removed, the mixture is covered with glacial acetic acid allowed to pass under. The slide is then coloring matters; whence it is possible

carefully warmed until bubbles begin to make their appearance. After cooling, hæmin crystals can be seen by aid of the microscope. These, though often very small and incompletely crystallized, are easily recognizable by an amplification of 300 diameters. They are chemically hydrochlorate of hæmatin.

Precautions.—Care must, however, be taken to apply only a gentle heat in precipitating the earthy phosphates with caustic potash solution, and to filter quickly, lest the hæmatin may be decomposed.

It sometimes happens, also, that vesicles develop under the thin glass cover, after the addition of acetic acid, even before heat has been applied. These are carbonic acid. They should be allowed to pass away, and the slide then warmed until the formation of vesicles, that is, to the boiling-point of acetic acid.

Occurrence.—Hæmoglobinuria, that is, the direct passage into the urine from the blood of the coloring matters unaccompanied by the corpuscular element, occurs in certain general diseases, as scurvy, purpura, scarlatina, profound malarial poisoning, etc. Hæmaturic or bloody urine results from the above and from a variety of other causes which require no special mention.

Melanin is sometimes found in the urine of persons having melanotic cancer or sarcoma. It is deposited from urine in the shape of granular particles. These are soluble in liquor potassæ, and their solution is decolorized by passing chlorine through it. Melanin differs from carbon in being soluble in potash, while carbon is not.

(b) Uroerythrin.

Heller ascribes the well-known dark, reddish-yellow, or "high" color of all fever urines to the presence of a substance which he calls uroerythrin, as well as to an increase

of the normal coloring matters. Except that it contains solution of neutral acetate of lead, the precipitate presents a similar "rosy red" or "flesh color," which he attributes being found especially in diseases where there is evident It so far at least corresponds with the urohæmatin of Harley that it is a measure of the destruction of the blood corpuscles, though it will be remembered that the urohæmatin iron, little else that is certain is known with regard to uroerythrin. To it he ascribes the reddish color which so often characterizes the deposits of urates known as "lateritious;" of Harley is looked upon as a normal constituent of urine which may be abnormally increased, while uroerythrin, although a modified hæmatin, is still not considered idenif the supernatant urine in such cases be treated with to the same substance. It is doubtless a modified hæmatin, blood dyscrasia, as in low fevers, septic conditions, etc.

tical by its discoverer.

Neubauer includes uroerythrin among the normal coloring matters, while Hofmann and Ultzmann, following Heller, treat it as abnormal.

Detection.—Uroerythrin is known to be present by its pink coloration of the "lateritious" sediment, or by its precipitation by solution of neutral acetate of lead. Too much lead solution must not be added lest the precipitate be too abundant, and the coloring matter rendered less distinct by its being disseminated over a large amount of deposit. If the urine contains hæmatin or the coloring matter of blood, it must be first removed.

Precautions.—I. The froth of a urine highly charged with uroerythrin may appear yellow, as that of urine containing biliary coloring matter, but the precipitate of the latter by acetate of lead is also yellow and not pink, as is uroerythrin.

Clinical Significance.—Uroerythrin is found in the urine in all febrile affections, however slight; also, it is said, in pyæmia, diseases of the liver, and lead colic. All urine, according to Heller, which contains uroerythrin must be abnormal.

(c) Vegetable Coloring Matters.

The coloring matter of plants, especially chrysophanic acid, found in rhubarb and senna leaves, contributes to alkaline urine a reddish-yellow to a deep-red color. It can be recognized by the fact that the red alkaline urine on adding an acid becomes yellow, and on the addition of an excess of ammonia again takes on the red color.

Precautions.—Such precipitation by heat and potash solution might possibly be taken for blood-coloring matters. But the absence of albumin from the urine, the production of the red color by addition of an excess of ammonia, and its paling on the further addition of an excess of acid, serve to distinguish the vegetable-coloring matter from blood-coloring matter and urocrythrin.

Numerous other vegetable matters color the urine, among which santonin is conspicuous for the bright yellow color it produces in acid urine, while the staining of linen by it closely resembles that of biliary coloring matter. Dr. W. G. Smith (Dublin Quarterly Journal of Medical Science,

November, 1870) has investigated the subject, and found that the addition of alkali causes the development of fine cherry-red or crimson color, according to the amount of santonin present; but it will be observed that this reaction is that of the vegetable coloring matters generally, as above described.

Madder, indigo, gamboge, rhubarb, logwood, carrots, whortleberries, etc., give to urine more or less of their peculiar colors

Salicylic acid, when administered in sufficient doses, gives a smoky hue to the urine, and the urine strikes a blue color when a few drops of a solution of ferric chloride are added.

Carbolic acid introduced into the system in sufficient quantity causes a dark and even black discoloration of

(d) Biliary Coloring Matters-The Detection of Bile in the Urine.

The biliary coloring matters are chiefly bilirubin (C₁₆H₁₈N₂O₃), biliverdin (C₁₆H₂₈N₂O₅), and bilifuscin (C₁₆H₂₈N₂O₅), the last two being derivatives by oxidation of the former. The last is found as such in herbivorous bile, and bilifuscin can be obtained from human gall-stones. None of these give any spectrum unless acted upon by reagents. We have seen that urobilin, the normal coloring matter of urine, is bilirubin altered by taking up, while in the small intestine, water and hydrogen, as the result of which it acquires the spectrum described on page 123. From the intestine it is absorbed, and excreted by the kidness as the normal coloring matter of urine.

When bile is abundantly present in urine, the yellow color of the fluid, and especially of the froth or foam produced by shaking, is sufficient to excite suspicion. Further, if a piece of filtering-paper or a piece of linen be moistened with such urine, it retains a permanent yellow color on drying.

The only positive proof of the presence of the coloring matters of bile in the urine is found in Gmelin's or Heller's test for the unaltered coloring matters.

First. A quantity of urine is placed in a test-tube, and a small quantity of fuming nitric acid (nitrous acid of commerce) is allowed to pass carefully down the sides of the test-tube to underlie the urine, as described in Heller's test for albumin. If biliary coloring matters are present, at the point of union between the urine and the acid will very soon be seen a set of colors, which, if typical, should be green, blue, violet, red, and yellow, or yellowish-green again, in the order named from above downward. Often, however, one or more colors are wanting. The green is most constant and the first green indispensable to prove the presence of bile; but violet, shading into red and yellow, is also very constantly seen. The other colors may be produced by other coloring matters, especially indican.

Second. Equally satisfactory is the test if a few drops of the urine are placed upon a porcelain plate, and as much of the fuming acid is placed adjacent and allowed gradually to approach the urine. The same play of colors occurs.

Fleischl's Test. *- A modification of Gmelin's

^{*} Boston Med. and Surg. Journal, Jan. 13, 1876, from Centralblatt für die medicinischen Wissenschaften, 1875, No. 34-

test, by which it is made more delicate. Instead of having impure nitric acid added in such a way that it will form a separate layer at the bottom, the urine should be thoroughly mixed with pure nitric acid, or, still better, with a concentrated solution of the nitrate of sodium, and then concentrated sulphuric acid should be carefully added so as to form a separate layer at the bottom. The play of colors forms at the junction of the urine and the sulphuric acid, the green appearing first above the acid, but rising gradually and giving place to the blue, violet-blue, and yellow. The advantage of this modification is that the pigment is not oxidized so rapidly, and, therefore, the color is not changed so quickly, remaining often half an hour or

Heller's Test for Bile Pigment.—Pour into a testtube about 6 c.c. (1.6 f3) of pure hydrochloric acid, and
add to it, drop by drop, just sufficient urine to distinctly
color it. The two are mixed and "underlaid" as before
with pure nitric acid, and at the point of contact between
the mixture and the colorless nitric acid a handsome play
of colors appears. If the "underlaid" nitric acid is now
stirred with a glass rod, the set of colors which were superimposed upon one another now appear alongside of each
other in the entire mixture, and should be studied by
transmitted light. Heller further says, if the hydrochloric
acid on addition of the biliary urine is colored readishyellow, the coloring matter is bilirubin; on the other hand,
if it is colored green, it is biliverdin.

If the amount of coloring matter is very small, a large quantity of urine should be shaken with chloroform; the chloroform allowed to separate at the bottom of the vessel in large drops. The yellow-colored chloroform is then

removed by means of a pipette, washed with distilled water, and poured into a beaker-glass containing hydrochloric acid. The yellow drops of chloroform sink to the bottom. If now, while diligently shaking the glass, nitric acid is added, the changes of color can be distinctly observed in the chloroform. In consequence of the slower action of the acid upon the coloring matters dissolved in the urine and the consequent slower transition of colors, this method is peculiarly adapted for demonstration.

Precautions.—I. With neither test should too dark-hued a urine be employed. Very dark urines should first be diluted with water.

2. Should albumin be present, the opaque zone at the point of contact between the urine and acid imbibes the coloring matters and exhibits a green coloration; so that the test is in no way interfered with.

3. Urine rich in indican may, however, deceive, forming at the point of contact a blue layer of indigo, which, along with the yellow urine, in reflected light, may appear green. In these doubtful cases the chloroform modification of the test should be used, or the urine may be precipitated with solution of acetate of lead, and the filtrate examined for indican.

 The earthy phosphates, precipitated from biliary urine by liquor potassee and heat, exhibit a brown coloration.

Oltzmann's Test.—Add to 10 c.c. of urine 3 or 4 c.c. of pure caustic potash solution (1 part of KOH to 3 of H₂O), then shake and add an excess of pure hydrochloric acid. The mixture assumes a beautiful emerald-green color.

Marechalt's Test.—Upon a specimen of urine in a test-tube allow a few drops of tincture of iodine to fall carefully. If biliary pigments are present a green color appears at the point of contact between the two fluids, and remains for some time, even twenty-four hours. In this

test the possibility of confounding indican is said to be

matters which respond to neither Gmelin's nor Heller's Test for Decomposed Biliary Coloring Matters. test, Hofmann and Ultzmann recommend the following:--Should the urine contain only altered biliary coloring

A piece of white linen or filtering-paper is immersed in the suspected urine, and allowed to dry, when it will appear colored brown. A further confirmation that the decomposed coloring matters are present will be found in a low specific gravity and a dark urophain reaction.* If, moreover, the urine be treated with liquor potassæ and heat, to precipitate the earthy phosphates, it becomes darker than before and the phosphates are precipitated

bile-pigments are contained in the precipitate, while the milk of lime, freeing from excess of lime by a current of carbonic acid gas, allowing the whole to stand a few hours, filtering, and washing the precipitate with water. The indican, hæmoglobin, and methæmoglobin are in the Bile-pigments have a property of adhering to precipitates may be in precipitates. Hence Dr. J. F. Tarchanoff (Centralblatt für die medicinischen Wissenschaften, 1875, No. 6) from the urinary pigments, precipitating the urine with much more tenaciously than other pigments, and therefore sometimes cannot be detected in fluid urine when they recommends, in order to separate with certainty the biliary

^{*} It should be remembered that this dark urophain reaction is also produced by sugar and blood-coloring matters. These causes should, therefore, be eliminated.

144

filtrate. The precipitate is then dissolved in acetic acid and tested by Gmelin's test.

In addition to urobilin, the normal coloring matter of urine, directly derived by oxidation from bilirubin, the normal coloring matter of fresh human bile, the latter substance and other derivatives from it are found in the urine when from any cause bile is reabsorbed. No attempt has been here made to separate these from each other, and the tests given are those for bilirubin and its combined derivatives, whatever they may be. Indeed, I do not know that any of these except urobilin has been recognized in urine as a distinct coloring matter, unless it be the one last discovered by Stokvis,*—and apparently not yet named,—as a secondary product in most cases of the oxidation of biliary coloring matter, whereby Gmelin's reaction is produced. This same substance MacMunn† believes he has found in the urine, by the aid of the spectroscope, in certain cases of rheumatic fever, pregnancy, thoracic ancurism, cirrhosis of the liver, and cancer of the pylorus. It is indicated by an absorption band, which occurs on the red side of the band of urobilin.

In testing for it, the liquid is to be precipitated with lead acetate, excess of lead removed by oxalic acid, and the filtrate concentrated and boiled with alkalies and a reducing agent. If no reduction takes place, and if the other tests for biliary coloring matters have given a negative result, their absence may be inferred.

XII. THE BILIARY ACIDS.

From a perusal of almost all of the text-books on physiology, and even of numerous manuals on the examination of urine, the student is led to suppose that the detection of bile acids, if present in urine, by means of what is called

^{*} N. Rep. Pharm., xxi, 123; Watt's Dictionary, 2d Supp., 1875;

^{†&}quot;Spectroscope in Medicine," London, 1880, p. 168.

bath. The residue thus obtained is treated with an excess an excess of ether (12 to 24 times its bulk), by which the bile-acids, if present, are precipitated. These are then is many times too strong) is then added to a drachm or tube is kept in a vessel of cold water, to prevent too great gation of this subject in its practical bearings will be found in a clinical lecture by the writer, in the Philadelphia Medical Times for July 5, 1873, "On a Case of Jaundice, with Remarks on the Availability of Pettenkofer's Test," to which the reader is referred. In these experiments the simplest method of obtaining the biliary acids was found to be as follows: Six or eight ounces (180-240 c.c.) of the of absolute alcohol, filtered, and the filtrate treated with The solution is then decolorized by passing through animal charcoal and the resulting colorless fluid tried by Pettensolution of cane-sugar (simple syrup of the Pharmacopæia phuric acid is then added drop by drop, while the testdetection by the direct application of the elements of Pettenkofer's test in urine, or any other animal fluid, is practically impossible, even if the bile acids are present in considerable amount. Nor have any of the modifications of Pettenkofer's test, recently announced as clinically available, proved such in my hands, even where the elements of bile ox-bile has been used. The results of a complete investisuspected urine are evaporated to dryness over a waterremoved by filtration and redissolved in distilled water. kofer's test as follows: A single drop of a 20 per cent. a rise in temperature, which should not exceed 50°-70° C. Pettenkofer's test, is one of the easiest possible. This is, however, far from being the case, and the fact is that such have been added to the urine, except where inspissated two (3.7-7.4 c.c. in a test-tube or porcelain capsule.

(122°-158° F.). As the quantity added approaches a bulk equal to that of the fluid to be tested, a beautiful cherry-red or purple-violet color should make its appearance. So soon as a yellow color appears, then the sulphuric acid is acting on the sugar, and the cherry-red can no longer be looked for. This carbonizing of the sugar is obviated by keeping the temperature down to the degree mentioned.

other than the color test, for the presence of the biliary genous jaundice), and absent in cases of jaundice from obstruction and consequent reabsorption of bile (hepatoalways present in the urine in cases of jaundice from acids, although undoubtedly occurring, is very rare, and simple, and there is really rarely any necessity for any able to the active practitioner, but there is none more and phosphorus poisoning. On the other hand, traces of destructive diseases of the liver, as acute yellow atrophy value in diagnosis. The only circumstances under which dissolution of the blood (hæmatogenous jaundice) else mined. It is not true, as was once supposed, that they are the circumstances under which they occur are illy deterband at E. investigated. It gives a band outside D, and a broad The bile acids yield a spectrum, which MacMunn has Dragendorf having found .7 to .8 gram in 100 litres.* the bile acids are said to be present in normal urine, they are undoubtedly present in the urine are rapidly would the determination of their presence be of real Even this method involves more time than is often avail-

^{*}The presence of bile-acids in normal urine is also asserted by Dr. Oliver, this assertion being based on experience with his new peptone test considered in the ensuing section.

Acids.—This is founded on the physiological fact that when the products of gastric digestion, peptone and parapeptone, which leave the stomach in an acid solution, meet with the bile, they are thrown down in a tenacious layer over the entire mucous membrane of the duodenum. So, too, albuminous urine or urine charged with peptone is precipitated by a solution of bile salts or of their derivative, cholate of sodium. Hence acidified albuminous urine becomes a test for bile salts, but an acidulated antiseptic solution of peptone is a readier and more delicate reagent. Such a solution is made by Dr. Oliver as follows:—

r. xxx.	r. iv.	L xxx.			
50	00	=	2		
-					
ore					
To					
-					
pur					
500					
ar					
27					
S					
ne					
0					
Pulverized peptone (Savary and Moore), gr. xxx.	Salicylic acid, gr. iv.	Acetic acid,	Distilled water,		

Perfect transparency is secured by repeated filtration.

Application.—The urine should be perfectly clear, rendered so by filtration if necessary, boiled and filtered if bloody, rendered normally acid if alkaline, and finally reduced to a specific gravity of 1008. Twenty minims should then be run into 60 minims of the test solution, and if the proportion of bile salts is normal or subnormal there is no immediate reaction, but in a little while there is a mere tinge of milkiness. If, however, the bile salts are present in excess, a distinct milkiness promptly appears, becoming more intense in a minute or two, the degree of opacity being directly proportionate to the amount of bile derivatives.

On agitation, the opalescence diminishes and perhaps finally vanishes, but is restored on adding more of the test

obviates another source of error, in that concentrated normal urine, may actually contain more than the normal sequent thereon. The reduction in specific gravity also amount of bile salts. specific gravity, though affording a reaction similar to urines often simulate an excess, while urines of low urates as avoids their precipitation and also any error con-1008. By the dilution is also secured such a solution of the proper proportions, as in Dr. Oliver's formula, and of disalts themselves. Hence the importance of securing the the reaction, as also does an excess of proteids, or of the an insufficiency, as well as an excess of acid, interferes with degree of warmth as is sufficient to dissolve urates. Further, citric acid test-paper, and by diminishing but not disappearcompletely on adding a drop or two of acetic acid or a precipitates induced by an acidified reagent, in dissolving solution. The precipitate differs from all other urinary luting the urine to be operated upon to a specific gravity of ing when boiled, but the opacity is not affected by such a

The test may be used in the contact method, by running the solution over the urine reduced in specific gravity to roos. If the bile salts are present in normal amount or less, there is again no immediate response, but in the course of a minute, a delicate, thread-like line makes its appearance, which may increase slightly. If the bile salts are abnormally increased an immediate reaction takes place.

This test, according to Dr. Oliver, is so delicate that there can readily be detected one part of bile salts in at least 18,000 to 20,000 parts of a solution of chloride of sodium. So far, he has been unable to find any other constituent of urine which reacts similarly, and, although it

is true that a concentrated solution of chloride of sodium in the presence of an acid will precipitate a proteid, experiment shows that when the peptone solution is run upon a solution of salt of any specific gravity below 1050, no precipitation takes place. Hence there can be no error from this source in urine.

Mucin may be eliminated as a source of error, because this substance in acid solution is not precipitated by adding more acid, and when it is thrown down in urine of acid reaction, it is highly probable that the acid is not the reagent producing it, but merely supplies the requisite degree of acidity to enable the precipitant already present to operate, and in that event the mucin would only indicate the presence of bile salts.

Quantitative Estimation.—This is based upon a permanent standard of opacity provided by mixing together, in equal proportions, the test solution and normal urine reduced to the specific gravity of 1008.

To 60 minims of the test solution add the suspected urine reduced to a specific gravity of roo8, usually 10 to 20 minims at a time, allowing a minute to elapse after each addition, until the opacity induced is exactly equal to or slightly exceeds that of the standard, the tubes being held to the light, shaded by a dark background, such as a coat-

If fifty or sixty minims bring up the opacity to that of the standard, the proportion of bile salts does not exceed the normal amount. Any smaller quantity required indicates an excess, while the smaller the amount needed, the larger the proportion of bile salts present.

Dr. Oliver has constructed a table showing the percentage of increase indicated by a varying number of drops:—

150

45	40	35	30	25	20	15	IO	Si	4	cs	13	I	Minims.
or	or	10	10	or	or	or	or	Of	or	10	or	or	
											4		UT.
	1	11	11	11	-	11		11	11	11	11	1.	Perc
50	66	83	100	240	300	400	600	1200	1500	2000	3000	6000	Percentage of Increase on the Normal Standard.
													ase on ard.

An increase beyond 700 per cent, over the normal is rarely met, although Dr. Oliver mentions an instance of non-jaundiced urine which showed an increase of over 1500 per cent., which afforded at once a beautiful reaction with Pettenkofer's test.

Peptone Test-Paper.—Dr. Oliver has also constructed a peptone test-paper which he considers permanent and reliable, and best used as follows:—

The peptone paper with half a citric paper is dropped into 60 minims water in a test-tube or wineglass. After the lapse of a minute the solution is slightly agitated, and on being set aside for another minute is ready for use.

The solution thus prepared is taken up by the pipette and carefully run over the transparent urine. If bile salts are present in larger amount than the normal average, an immediate reaction is observed as a pearly-white thread or band. In urine in which there is no such excess, a delicate zone may appear, but only in the course of one or two minutes.

XIII. LEUCIN (C₆H₁₃NO₂) AND TYROSIN (C₉H₁₁NO₃).

Leucin and tyrosin, products of a retrograde metamorphosis of nitrogenous substances, are found physiologically only in certain fetid secretions, as those of the axilla and between the toes, but can be produced by chemical means from some glands, as the liver, pancreas, and spleen, where they also occur in certain pathological states. They are found in urine, chiefly in rapidly destructive diseases of the liver, as acute yellow atrophy or phosphorus poisoning, but occasionally also in typhus and smallpox. They always accompany a large amount of biliary coloring matter and the presence of albumin. When at all abundant, as they generally are in acute yellow atrophy, they are deposited from urine and are found in the sediment, the former in the shape of centrically marked spheres, arranged in warty masses, or druses, the latter in needles. (Fig. 27.)

Schultzen has shown * that in animals poisoned by phosphorus 't urea disappears from the urine and is replaced by leucin and tyrosin, which in the healthy organism are converted into urea." A similar substitution takes place in cases of acute atrophy of the liver, the retained urea accounting for the convulsive attacks which usually precede death in these cases.

Detection.—If the crystals—to be more fully described in treating of sediments—do not present themselves in the spontaneous deposit of such cases, the evaporation of a small quantity of the urine will generally promptly display them.

^{*} Boston Medical and Surgical Journal, July 23, 1884, from Zeitschriff für Biologie, viii, 124, and Berliner klin. Wochenschrift, 1872,

If they are not sufficiently abundant to be thus demonstrated, the method of Frerichs must be pursued to separate them. A large amount of urine is precipitated with basic acetate of lead, filtered, the excess of lead removed from the filtrate by sulphuretted hydrogen, and the clear fluid evaporated to a small volume over a water-bath. In twenty-four hours tyrosin needles will be found to have crystallized out, but leucin spheres will not appear until later, because of the great solubility of the leucin.*

XIV. FATTY MATTERS.

That a trace of fat exists dissolved in normal urine has been shown by Schunk; while the list of reported cases in which fat is present in abnormal quantity is gradually increasing. In such cases are, of course, not included those in which fatty epithelium, fatty casts, and free oil drops are present, as the result of chronic Bright's disease; nor those in which fatty epithelium from the bladder or vagina occurs.

The oil may be present in urine in a state of minute subdivision into small drops and molecules, as in the so-called chylous urine, or in the form of clear fluid oil. In the former instance the admixture probably results from the leakage of a lymph vessel into some part of the urinary tract. In the latter its source has been, in one instance†

^{*}Leucin and tyrosin are more fully treated by the writer in the American Journal of the Medical Sciences for January, 1872. The above is believed to be sufficient for practical purposes.

[†]Dr. E. W. Cushing, in Boston Medical and Surgical Journal, vol. 104, 1880, p. 242.

administration of cod-liver oil, and in the third during the through the urine. I have recently had under my care a municating with the left ureter; and such possible source should always be remembered. It is not impossible also that it may come directly from the kidney in cases of cystic cheesy degeneration of that organ, instances of which I have seen where there has been considerable free oil with compound granule cells and cholesterin plates among the cheesy matter. Dr. Roberts * refers to three cases in which pure yellow oil was present in the urine, in two during the use of an emulsion. Dr. Henderson † reported three cases of heart disease in which free oil globules were suspended young woman in whose urine, always free from albumin, free oil drops and aggregates of oil drops could be found at almost any time, and at times tube-casts containing oil drops, the most scrupulous care being taken to avoid at least, traced to an abscess in the left lumbar region com-

Fat has been found in the urine from cases of calculous disease of the pancreas, and in one case referred to by Dr. George W. Johnston,‡ fat made its appearance in the urine one month before it was detected in the alvine dejecta, and in such quantity as to float, when cool, in greasy flakes on the surface.

The presence of **cholesterin** in the urine is also a possible but very rare occurrence, and may be conceived to occur in such a case of cheesy cystic kidney above alluded

^{*&}quot; Urinary and Renal Diseases," Am. ed., 1879, p. 125.

⁺ British Medical Journal, May 22, 1858.

[†]Inaugural thesis for the degree of Doctor of Medicine in the University of Pennsylvania, 1882. Published in the American Jour. of the Medical Sciences, October, 1883, of which see p. 427.

to. The only well-authenticated case I have ever seen reported is that given by Dr. Roberts.* Dr. Beale has shown that cholesterin may be obtained by treating large quantities of urine from cases of chronic Bright's disease, but this is a different matter from its being contained in urine in the free state.

XV. UREA (CN2H,O).

Urea is the chief organic constituent of urine and the index of nitrogenous excretion. Its quantity fluctuates with changes in the quantity and composition of ingesta, and with the rapidity of tissue metamorphosis in health and disease. A range of at least from 20 to 40 grams (308.6 to 617.2 grains) in 24 hours must be admitted in adults.


Detection and Estimation.—The odor of urine highly charged with urea may be said to be characteristic, but certain evidence of its presence can only be obtained by treating the solution suspected to contain it with nitric or oxalic acid. Though crystallizing itself in glistening needles, it is too soluble to permit of easy detection by its own form. If it be desired to detect its presence in a suspected fluid, a drop or two is placed upon a glass slide, a drop of nitric acid added, the slide carefully warmed over a spirit-lamp, and placed aside to crystallize. If urea is present, the microscope will reveal, singly or in strata, six-sided and quadrilateral plates of nitrate of urea (Fig. 13). The crystals have acute angles measuring about 82°, and are so characteristic as to be

155

easily recognizable; they often overlap each other like the shingles of a roof.

Solution of oxalic acid produces similar but less regular crystals of oxalate of urea.

In ordinary normal urine, this crystallization does not take place unless the urine is concentrated by evaporation. But in some urines highly charged with urea, it is simply necessary to add nitric acid to produce the crystals, and thus is arrived at a rough quantitative estimation of urea.



Crystals of Nitrate of Urea, (After Beale.)

As urea is by far the most abundant solid constituent of the urine, it follows that the specific gravity may become a means of approximately estimating its amount, especially when there is no sugar present, if the quantity of albumin is small and that of the chlorides is normal. A specimen of urine neither albuminous nor saccharine, containing a normal proportion of chlorides, and having a specific

gravity of 1020-4 to a quantity of 1500 c.c. (50 0z.) in twenty-four hours, may be taken as a standard normal specimen containing 2 per cent. to $2\frac{1}{2}$ per cent. of urea. These conditions being observed, a higher specific gravity would indicate an increased proportion, and a lower, a diminished proportion. Under these circumstances, a specific gravity of 1014 indicates about 1 per cent. of urea and of 1028 to 1030 about 3 per cent.

and other febrile diseases, accompanied by an increase in separated to the amount of 10 to 16 grams (154 to 247 and by far the largest proportion of urines in which a maining constituents, uric acid, creatinin, phosphates, sulcent. This is on the supposition, of course, that the repercentage of chlorides replaced by urea be added, 31 per the elimination of urea, then must a specific gravity of these are totally absent, as they often are in pneumonia grains), or 3 to 1 per cent. in the twenty-four hours. If albumin. Next to urea, supposing albumin and sugar knowledge of the amount of urea is important contain which is the fact. phates, etc., have little influence on the specific gravityrozo indicate more than 21 per cent, of urea, or, if the absent, the chlorides most affect the specific gravity, being But the chlorides fluctuate markedly in some diseases,

If albumin is present in small quantity, not exceeding $\frac{1}{16}$ per cent., it has little effect, and it can be thrown out of the question. If, however, the albumin be more abundant, I to 2 per cent., it must first be removed by coagulation and filtration, and the approximate estimation be made from the specific gravity of the filtrate after cooling. Care must of course be taken to wash the coagulum by further addition of water until the quantity of fluid originally

lower than in health, showing-what volumetric analysis has determined more precisely-that in chronic albumioperated with is restored. After such removal of albumin, if not before, the specific gravity will generally be found nuria, at least, the quantity of urea is generally diminished.

generally less, though with increased specific gravity, while the large total quantity of urine in the twenty-four hours may show an increase in the total urea for the day. There the presence of sugar, and the only way to arrive at a Where sugar is present, the percentage of urea is also is no way of allowing here for the specific gravity due to knowledge of the amount of urea is by volumetric analysis.

Volumetric Analysis for Urea.

analysis. Several methods of volumetric analysis for urea have been suggested, of which that of Liebig, with the nitrate of mercury solution, was for a long time solely employed, but of late has been largely superseded by the hypobromite and hypochlorite processes. Liebig's method is based upon the fact that urea produces a precipitate with of urea is required, we must have recourse to volumetric Under any circumstances, when an accurate estimation mercuric nitrate.

The following test solutions are required:-

- 1. The baryta solution, consisting of one volume of cold saturated solution of barium nitrate with two volumes of cold saturated solution of caustic baryta (barium hydrate).
 - 2. A saturated solution of sodium carbonate.
- 3. A standard solution of mercuric nitrate of such strength that I c.c. is precisely equivalent to .010 gram, or 10 milligrams, of urea.

To Prepare the Standard Solution of Mercuric Nitrate.—
I. Dissolve 71.48 grams of pure mercury or 77.2 grams of the mercuric oxide in nitric acid by the aid of heat. The acid fluid is concentrated by evaporating over a water-bath to a syrupy consistence, and then diluted with distilled water to a volume somewhat less than a litre. If on dilution a white precipitate of basic nitrate of mercury fall, allow it to settle, and decant the clear liquid. Then add to the residue a few drops of nitric acid to dissolve the precipitate. Add the solution thus obtained to the former decanted liquid and dilute to exactly one litre. The solution requires to be graduated by.

 The Standard Solution of Urea.—Two grams of pure urea should now be dissolved in 100 c.c. of distilled water, of which 10 c.c. will then contain 0.2 gram or 200 milligrams.

a dense precipitate. When the precipitation seems about complete, urea, are now placed in a beaker-glass. A burette is then filled and the yellow oxide of mercury. with the sodic carbonate, which it does by forming sodic nitrate the urea in combination and left some mercuric nitrate to react that the mercuric nitrate has been added in excess-consumed all with the Na2CO3, until finally a yellow color appears. This proves tion of the mercuric nitrate is continued, also the process of testing precipitated, no change of color takes place. The cautious addion a piece of glass on a dark ground. If the urea is not completely solution of sodium carbonate, of which several are previously ready a drop of the fluid containing it is allowed to fall on a drop of the is then allowed to drop into the beaker, where it will quickly form of the liquid corresponds with the arrow on the burette), which lower edge of the meniscus which forms the upper surface to o with the solution of mercuric nitrate (taking care that the Ten c.c. of the standard solution, containing 200 milligrams of

If now the mercuric solution is correct, it will require exactly 20 c.c. of it to precipitate the whole of the urea in the 10 c.c. of the standard solution of urea, and enough more to react with the sodium carbonate. If the yellow coloration does not occur under these circumstances, there has been some impurity in the mercury compounds employed, or some error in making up the solution.


Process.—Take 40 c.c. urine and 20 c.c. of the baryta solution, and throw them into a beaker-glass. By this means the phosphates, sulphates, and carbonates are precipitated. They are removed by filtration through a *dry* filter, and if the filtrate happens not to be quite clear, it may be passed through a second time.*

off the number of cubic centimetres employed.† The from the mixed baryta fluid and urine, containing of course 10 c.c. of pure urine, are measured off into a small beaker-glass. Into this the mercuric nitrate solution is allowed to fall from the burette, first, a number of cubic centimetres approaching the last two figures of the specific gravity (that is, if the specific gravity is 1017, drop say 15 c.c.) before testing with the sodium solution. If no cautiously, adding a fractional part of a cubic centimetre at a time, and testing with the Na2CO3 until the yellow coloration is obtained. When that point is reached, read number of cubic centimetres of mercury solution thus used, minus 2, multiplied by .oro gram, gives the amount While this is taking place, the burette is filled to o with the mercuric nitrate solution, and 15 c.c. of the filtrate yellow coloration appears on such testing, then proceed

^{*} If the filtrate is not alkaline, the precipitation of the phosphates and sulphates may not have been complete. This may be determined by adding a drop or two of the baryta mixture to the filtrate, when, if a precipitate appears, a fresh quantity of urine must be taken, and a larger proportion of the baryta solution added.

be the transport of yellow at which we cease the titration must of course the tasme as that at which in originally standardizing the nitrate of mercury solution the titration was stopped. It is evident that ceasing the titration, now at a slight tinge and again at a marked yellow coloration, must give rise to an error, which practice will soon teach the student to avoid.

of urea in fractions of a gram contained in 10 c.c. of the urine, when the latter is of average composition—that is, when it contains no abnormal constituent, and the amount of chlorides is nearly normal.



Burette Stand with Two Forms of Burette.

Corrections Explained.—The two cubic centimetres are first subtracted because it takes about this quantity of the reagent to convert the chloride of sodium into the nitrate, and until this combination is complete, the combination

with the urea does not begin. Hence this amount must first be subtracted,

If, however, the chlorides are not of average amount, but diminished or increased, and we wish to be accurate, we must first estimate the amount of chlorides calculated as NaCl in 10 c.c. of the urine, by the process to be explained under chlorides, and from a fresh quantity of urine remove the whole of the chlorides by a standard solution of silver nitrate. For this purpose a solution of nitrate of silver is required, of such strength that 1 c.c. will precipitate 10 milligrams sodium chloride. 29.059 grams of fused nitrate of silver, dissolved in distilled water and diluted to a litre, will be such a fluid.

In 10 c.c. of the original urine we determine with the nitrate of silver solution the chloride of sodium by the method for the determination of the chlorides, p. 178. Suppose there are required for this 17.5 c.c. of the silver solution, this indicates 175 milligrams sodium chloride.

Take now 30 c.c. (containing 20 c.c. of urine) of the filtrate from the mixture of baryta fluid and urine, add a drop of nitric acid, and then 17.5 × 2 c.c. = 35 c.c. of the nitrate of silver solution. This will precipitate all the chlorides, which should be removed by filtration, and the filtrate may be now estimated for urea. It is important always to bear in mind the exact amount of urine operated with after adding the nitrate of silver solution to a mixture of baryta solution and urine, of which only two-thirds are urine. Thus, if 35 c.c. of the silver solution are added to 30 c.c. of the filtered mixture of urine and baryta fluid, of the resulting 65 c.c. only 20 would be urine minus the chlorine, or out of 32.5 c.c. to would be urine minus the chlorine.

If the case be one of inflammation, as pneumonia, where there is a total or almost total absence of chlorides, they may be thrown out of the question altogether.

Further Corrections.—If the number of cubic centimetres of mercury solution added to 15 c.c. of the mixture of urine and baryta fluid exceeds 30—that is, if the amount of urea in the unmixed urine exceeds 3 per cent.—we must, for the number of

c.c. of the mercurial solution required above 30, add half the number of cubic centimetres of water to the urine mixture and make a second titration. Thus, suppose 36 c.c. are required on the first titration, the excess is 6 c.c., therefore 3 c.c. of water must be added to the mixture before making the second titration.

If the unmixed urine contains less than 3 per cent, of urea, then for every 4 c.c. of the test solution used below 30 there should be deducted I c.c. from the entire number of cubic centimetres of the mercurial solution used.

These corrections are rendered necessary by the fact that in standardizing the reagent it was mixed with just half its volume of the urea solution, conditions which we have (in regard to dilution) when 10 c.c. of urine containing 3 per cent. of urea are mixed with 5 c.c. baryta mixture and 30 c.c. of the mercury solution. Hence, any excess of reagent employed above 30 c.c. for 15 c.c. of the urine mixture should be diluted with half its volume of water to reduce such excess to the same degree of dilution as was present in standardizing the reagent. So, on the other hand, if less than 30 c.c. of the regent are required, that employed will be under greater dilution than was present in standardizing the reagents; hence, the reduction mentioned above.

Estimation of Urea by the Hypobromite Process.—The principle on which this process is based—that urea, when brought in contact with hypochlorite of calcium, is decomposed into nitrogen, carbonic anhydride, and water—was suggested many years ago by Davy,* and at the same time by LeConte.† In 1874 Messrs. Russell and West‡ again directed attention to the subject, substituting an alkaline solution of hypobromite of sodium and caustic soda, which yields similar products, the carbonic

^{*} Philosoph. Mag., 1854, p. 345.

⁺ Comptes Rendus, xlvii, 237.

^{\$} Journal of the Chemical Society (London), August, 1874.

anhydride being absorbed by the caustic alkali. The following is the reaction:—

$$CON_2H_4 + 3(NaBrO) = 3(NaBr) + CO_2 + 2H_2O + N_2;$$

the volume of nitrogen disengaged being the measure of the urea.

Many forms of apparatus have been suggested by different experimenters, all based upon the assumption that one gram of urea contains 372 c.c. nitrogen, measured at 0° C. and 760 mm. barometric pressure; or that each c.c. of nitrogen evolved, measured under the conditions stated, represents 0.00282 gram urea.*

The simple apparatus shown in Fig. 15, devised by Dr. John Marshall, † is that in use in the chemical laboratory of the University of Pennsylvania. It consists of a graduated measuring tube, a, a graduated pipette, c, a funnel tube, d, and a saucer-like apparatus, b, which serves as a stand and also as a receptacle for the hypobromite solution which in the operation overflows from the measuring tube. The measuring tube, including the bulbous part, holds about 77 c.c., and can, by means of a perforated cork, be fastened to the saucer-like vessel. The apparatus can easily be taken apart and cleansed of the grease-like material which, in the performance of the hypobromite method, usually accumulates in the measuring tube.


The alkaline hypobromite solution is made by dissolving roo grams of caustic soda in 250 c.c. of water, and adding 25 c.c. of bromine to the solution thus produced.

$$_{2}$$
NaHO + $_{1}$ = $_{2}$ NaBr + $_{2}$ O + $_{3}$ NaBrO.

^{*} See note to p. 165.

⁺ Zeit. f. Physiolog. Chemie, Bd. xi, p. 179.

Process.—The thumb is placed over the opening a, and the hypobromite solution is poured in through b. The latter opening, b, is then closed by a rubber stopper, and any air bubbles in the tube are allowed to pass out at the opening a. The graduated tube is now reversed and the closed oval end of the tube is fastened in the opening in the saucer-like vessel. A measured quantity of urine (one cubic centimetre generally suffices) is allowed to pass



Marshall's Apparatus.

from the pipette C into the hypobromite solution through the opening a. Decomposition of the urea immediately occurs, and the evolved nitrogen collects in the upper part of the graduated tube. The carbon dioxide evolved in the decomposition of the urea is absorbed by the excess of caustic soda in the hypobromite solution. About 20 minutes are required for the complete decomposition of

the atmospheric pressure is equalized by attaching the funnel tube (d) to the opening (a) and pouring into it hypobromite solution or water until the surface of the liquid in both tubes stands at the same level. The number of cubic centimetres of nitrogen, the temperature and the barometric pressure are read off, and the the urea. When the decomposition is completed and all the gas bubbles have collected with the gas in the upper part of the tube, percentage of urea calculated by the following formula :-

$$p = \frac{100 \text{ v. (b-b')}}{760 \cdot 354.33 \text{ a. (1 + 0.00366 t.)}}$$

In which

p represents the weight or urea in grams in 100 c.c. of the urine.

the volume of urine employed.

the observed barometric pressure in millimetres.

the tension of aqueous vapor for the temperature t.

the observed temperature (Centigrade).


0.00282 * gram, representing the weight of urea corresponding to a single c.c. of nitrogen gas. Thus, suppose 9.6 c.c. of gas have For clinical purposes the thermometer and barometer may be ignored, and the reading in centimetres need only be multiplied by been read off. Then the 1 c.c. of urine (or as much as was used gram, whence can be calculated either the percentage or 24 hours' in the operation) must have contained 0.00282 \times 9.6 = 0.027 the volume of nitrogen, in cubic centimetres, obtained. quantity of urea.

+ Zeit. f. Physiolog. Chemie, Bd. i, p. 350.

evolved in the practical application of the method, then 1 c.c. of the * The amount of nitrogen gas which, theoretically, should be evolved from I gram of urea is 372.7 c.c., but Hüfner † has shown that practitherefore, I c.c. of the nitrogen at 0° C. and 760 mm. pressure corresponds to 0.00282 gram of urea. Hence the latter figures are used in the calculation. If the full theoretical quantity of nitrogen were nitrogen, at oo C. and 760 mm. pressure, would be equal to 0.00268 cally by the hypobromite method only 354.33 c.c. are evolved, and, grm. of urea.

Various other forms of apparatus for this purpose have been devised for this process, notably by Hüfner in Germany, Gerrard in England, Dr. William H. Greene,* of Philadelphia, Dr. Charles A. Doremus, of New York, E. R. Squibb, of Brooklyn, and others.

Doremus's apparatus is much used, and consists of a bulb



Ureometer of Doremus

and graduated tube, and a small, curved, nipple-capped pipette to hold one c.c. of urine. The tube is graduated so that each of the small divisions equals .oor gram of

^{*} First described in the Philada. Med. Times for January 12, 1884 p. 278.

urea. It is filled with hypobromite solution of the usual strength to the mark on the long arm of the apparatus, and water added to fill the remainder of the arm and the lower part of the bulb. The pipette is then filled with urine up to the c.c. mark, and the point carefully introduced into the bend as far as it will go, holding the measuring tube perpendicularly. The nipple is then slowly and thoroughly compressed so as to expel all the urine. After the evolution of gas is complete the number of divisions is read off and the result multiplied by 100 to obtain the percentage.

With Squibb's apparatus is used the solution of chlorinated soda, U. S. P., which has the advantage of being always obtainable from the drug stores, while the hypobromite solution has to be made fresh for use. Full directions for the estimations are furnished with the apparatus.

In experiments made by Messrs. West and Russell, Mr. Richard Apjohn, Dr. Dupré, Dr. M. Simpson, Mr. C. O'Keefe, Prof. Theodore G. Wormley, and others, with solutions containing known quantities of urea, astonishingly accurate results were obtained, quite sufficiently so for clinical purposes.

M. Depaine (Journ. de Pharm. d'Auv., 1877) recommends that 4.5 per cent. be deducted from the total amount of urea found, to eliminate the error caused by the simultaneous decomposition of uric acid and creatinin.

It is generally acknowledged that not quite all the nitrogen of the urea in a given solution is liberated by the hypobromite process, but authorities are not at all agreed as to the quantity retained. M. Leconte (Chem. Gaz., 1858) obtained .92; Foster announced (Journ. Chem. Soc., March, 1879) that .92 was obtained; Russell and West obtained .94; others secure a larger proportion, as much as .95; others a still

16° F. with cane sugar and 14.5° C or 26.2° F. with grape sugar. subject, has made an important observation, which may serve to exnitrogen to be read off. This, causing an expansion of the gas, would permit a larger amount of hypobromite a large amount of heat is evolved, as much as 8.9° C. or above alluded to. He finds that on mixing solutions of sugar and into this, that neither the glucose nor cane-sugar solution should be mixed with the hypobromite and urea solution, no additional gas is by the hypobromite, but with a solution of cane sugar only .94. Again, plain some of the discrepancies in the results obtained by the observers fessor Wormley, in some recent experiments in connection with this mixed with the hypobromite solution of urine. My colleague, Propractically almost impossible to obtain pure glucose, it resolves itself cane sugar solutions an appreciable amount is evolved. But as it is evolved in the short time alloted to an ordinary estimation, while with Jay's experiments (Ibid.) go to show that if a solution of glucose is the theoretical quantity of nitrogen is evolved from urea solutions mique, February, 1880) then announced that with a solution of glucose, the hypobromite solution evolves a gas. Fauconnier (Bul. Soc. Chi-Chim., 1880) announced that a solution of glucose alone mixed with Esbach (Journ. Chem. Soc., December, 1879), and M. Jay (Bul. Soc. of nitrogen otherwise retained being now liberated. But a little later mixed with the urine and hypobromite solution, the small amount different proportion. Mehu (Journ. Chem. Soc., November, 1879), addeficiency is overcome if a solution of either cane or grape sugar is mitting that about .08 of nitrogen is retained, first suggested that this

Fowler's Hypochlorite Process for Urea.*—This method is based upon the fact that there is a difference in the specific gravity of urine before and after the decomposition of its urea by the hypochlorites; and that such difference bears a definite relation to the quantity of urea

^{*}Fowler, Prize Essay to the Alumni Association of the College of Physicians and Surgeons, New York. Published in the New York Medical Journal, July, 1887.

present. Dr. Fowler found that every degree of density lost corresponds to .77 of 1 per cent., or about 3½ grains per fluidounce. The hypochlorite solution employed is Squibb's solution of chlorinated soda, or Labarraque's solution, of which seven parts will destroy the urea in one part of urine, unless the amount is very large, in which event the urine should be diluted by an equal bulk of water, and the result multiplied by 2.

Process.—1st. Add to 1 volume of the urine 7 volumes of the hypochlorite solution. Effervescence due to the liberation of nitrogen will immediately take place. Shake the jar containing the mixture occasionally, and stand it aside for two hours, when the urea will have been decomposed. Now take the specific gravity of the quiescent fluid.

2d. Ascertain the specific gravity of the mixed urine and hypochlorite solution before decomposition. To do this, multiply the specific gravity of the pure hypochlorite solution by 7, add this to the specific gravity of the pure urine, and divide by 8. The result is the specific gravity of the mixed fluid. From this subtract the specific gravity of the quiescent mixture after decomposition of the urea, multiply the difference by .77, and the result is the percentage of urea.

As changes of temperature affect the specific gravity and volume of liquids, the hypochlorite solution and urine should be mixed and the jar set aside along with a bottle of the urine and the hypochlorite solution in the same place, subject to the same temperature. When decomposition is complete, the specific gravities can be taken and the calculations and the calculations.

Example.—Suppose the specific gravity of the urine is 1010, and that of the hypochlorite solution 1045, that of the mixed fluid will be

$$\frac{1045 \times 7 + 1010}{8} = 1040.$$

Now suppose the specific gravity of the decomposed fluid is 1038, then (1040 — 1038) \times .77 = 1.54, the percentage of urea.

This very simple and easy process has been found quite accurate and is not interfered with by sugar or albumin.

XVI. URIC ACID (C,H,N,O,).

When uric acid is spoken of as a constituent of normal urine, it is never to its free state that allusion is made, but to its combinations, chiefly with potassium, sodium, and ammonium, but also with calcium and magnesium, usually known as mixed urates. Uric acid itself is so extremely insoluble (one part requiring 14,000 of cold and 1800 of hot water to dissolve it) that it is immediately precipitated on being freed of its bases. In quantity it is found ranging .4 to .8 gram (6.17 to 12.34 grs.) in the twenty-four hours, in health varying part passu with urea, of which it is a stage short in oxidation.

Detection by the Microscope.—Its presence as such is recognized by the microscopic peculiarities of its crystals, which in their typical form may be said to be "lozenge-shaped," or, as described by the Germans, "whet-stone-shaped." They are, moreover, always colored yellowish-red, being with their salts the only urinary deposits thus stained, so that when a sediment is seen of which the elements are thus colored, it may, without hesitation, be put down as composed of uric acid or its combinations. More will be said of these crystals in treating of sediments, where their discussion more properly belongs.

The Murexid Test.—The murexid test for uric acid and its combinations is one of extreme beauty. A small portion of sediment, or the residue after evaporation, is placed on a porcelain plate or piece of platinum, a drop or two of nitric acid added to dissolve it, and the solution carefully evaporated over a spirit-lamp flame. When dry, a drop or two of liquor ammoniæ is added, when there promptly appears a beautiful purple color, which will

gradually diffuse itself as the ammonia spreads. The murexid reaction is believed to depend upon alloxan, alloxantin, and ammonia, which arise under the action of the hot nitric acid. This reaction is also said to occur with tyrosin, hypoxanthin, and xanthoglobulin, and Schiff accordingly recommends the—

Carbonate of Silver Test for Uric Acid.—This is very delicate, and is most conveniently applied as recommended by Harley. Dissolve a little uric acid in a solution of sodium or potassium carbonate, place a drop or two of the solution on paper, and add a solution of nitrate of silver. A distinct gray stain promptly occurring indicates the presence of uric acid. Neither of the tests, however, discriminates between uric acid and urates. The micro-

scope most easily does this.

Quantitative Estimation of Uric Acid.—It has been usual to estimate uric acid by a gravimetric method, as

follows:—

To 200 c.c. add 20 c.c. of hydrochloric or nitric acid, and set aside in a cool place, as a cellar, for twentyfour hours. At the end of that time the uric acid crystals, highly colored, will be found adhering to the sides and at the bottom of the beaker. Collect the uric acid on a weighed filter, wash thoroughly with distilled water. Dry the filter and uric acid at a temperature of 100° C. (212° F.), weigh, and the weight of the two, minus the weight of the filter, will be the weight of the uric acid in 200 c.c., except the small portion retained in the acid and washings. Neubauer advises to add to the result 0.0038 gram uric acid for every 100 c.c. of these fluids.

Haycraft's Method.—Several volumetric processes have recently been suggested. All are lengthy and cum-

bersome as compared with the processes for urea. That of Haycraft, however, published in 1885, is probably least so, and quite accurate results have been obtained by its author with solutions of known strength.

It is based on the fact that uric acid combines with silver as silver urate, which is practically insoluble in water, ammonia, or acetic acid, but easily soluble in nitric acid. The chief difficulty lies in the gelatinous nature of the silver urate, which is difficult to wash. It is, however, overcome by collecting the precipitate on an asbestos filter attached to a filter pump. The filter is made by half filling a small funnel with broken glass, upon which is poured to the depth of one-fourth of an inch asbestos fibres made into a uniform pulp, by shaking the fibres with water in a flask. The same filter may be used repeatedly.

Solutions Required.—r. Centinormal ammonium sulphocyanate, made by dissolving about 8 grams of crystals in a litre of water, and standardizing by the decinormal silver solution (2). Dilute with 9 volumes of water. One cubic centimetre is equivalent to 0.00168 of uric acid.

2. Decinormal silver solution, made by dissolving 10.766 grams of silver in pure nitric acid and diluting up to 1 litre; or 16.966 grams of pure silver nitrate in a litre of distilled water.

 A saturated solution of ammonium ferric sulphate (iron alum).

4. Dilute nitric acid (20 to 30 per cent.). Dilute the commercial acid, boil and preserve from light in a blackened flask.

5. Ammonium hydroxide.

6. Ammoniacal silver solution. Dissolve 5 grams of

silver nitrate in 100 cubic centimetres of water, and add ammonium hydroxide until the solution becomes clear.

easily; finally, a drop or two of sulphocyanate produces The color is best seen by holding the flask so as to catch 50 c.c. are required, the solution must be made stronger in a flask, diluted somewhat with water, and 5 c.c. of the ferric indicator added, together with about 10 c.c. of nitric acid. If the iron solution should cause a yellow color, the nitric acid will remove it. The sulphocyanate is then added from a burette; at first a white precipitate is produced, rendering the fluid milky, and as each drop proaches the precipitate becomes flocculent and settles a faint brown color which no longer disappears on shaking. If the solutions are correctly balanced, exactly 50 c.c. of the reflected light of a white wall. If more or less than Process for Standardizing the Ammonium Sulphocyanate falls in, it produces a reddish-brown cloud which quickly disappears on shaking. As the point of saturation ap-Solution. -- 50 c.c. of decinormal silver solution are placed sulphocyanate solution are required to produce this effect. or diluted, as the case may be.

Process.—Measure off 25 centimetres of urine, and place it in a small beaker with about one gram of bicarbonate of sodium. Add two to three cubic centimetres of the ammonium hydroxide, which will produce a precipitate of ammonio-magnesian phosphate. On adding, one or two cubic centimetres of the ammoniacal silver solution, the uric acid falls as a white gelatinous precipitate of urate of silver.

This is collected on the asbestos filter and carefully washed, the filter pump being attached, until the washing

gives no trace of silver, with a drop of salt solution.* The washing is completed by the aid of a few cubic centimetres of the nitric acid. The silver in the solution is estimated by Volhard's method, as follows:—

Add a few drops of the saturated solution of ferric alum, which is the indicator, and drop in from a burette the centinormal solution of ammonium sulphocyanate. A white precipitate is formed, with a reddish coloration, which latter at first disappears. As soon as the red color is permanent the process is at an end.

The uric acid is then calculated by multiplying the number of cubic centimetres of sulphocyanate used by 0.00168.

If the urine contains albumin, this should previously be removed. If uric acid or urates be present in such quantities as to cause turbidity, the secretion should be warmed and diluted, in order to test the accuracy of the process.

XVII. URATES.

It has already been said that in health, practically all the uric acid of the urine is held in combination with potassium, ammonium, sodium, calcium, and magnesium, of which, according to Bence Jones, those with potassium and ammonium are most abundant. These are very soluble compounds at the temperature of the body, but are pre-

^{*} Haycraft recommends a Sprengel's filter pump, but an ordinary Clarke's filter pump answers very well. The filter pump should only be used when the precipitate is thrown on the filter, and with very little force at first. Before putting on the diluted nitric acid, the pump should be disconnected, as the acid should not run through too fast, or it may not completely dissolve all the silver precipitate.

175

cipitated in amorphous granules when the temperature of the urine is lowered, as in winter weather.

Their physiological and pathological significance depends altogether upon the uric acid they contain, but there are some points of reaction with which the student should be quite familiar. These grow out of the fact that uric acid is a bibasic acid, forming neutral and acid salts, and that acid salts are much less soluble than the neutral, requiring 124 parts of boiling and 1120 parts of cold water for their solution. They form, therefore, the bulk of urate deposits, while urates, which remain in solution after such reduction of temperature as constantly takes place in an apartment, must be, if not neutral, at least less acid than those which form the sediment. And a solution remaining for some time clear under such circumstances must contain urates of sodium, etc., with a large proportion of the alkaline base.

The practical application of this fact is seen in this, that when an acid is added to such solution of neutral urate, by seizing upon a portion of the base, it leaves an acid urate of sodium, which, in consequence of its relative insolubility, is promptly precipitated in a finely granular form, producing a decided opacity. Now, this is precisely what often happens in the nitric acid test for albumin. The urine is highly charged with neutral urates which are held in solution. Nitric acid is added, and down goes a precipitate, not crystalline, but amorphous, which is composed of acid urate of sodium. And if Heller's method is followed, an opaque zone is formed at the point of contact between the acid and urine, which may be mistaken for albumin, but which, besides presenting certain visual characters of its own, which have been described, p. 41, is

opacity gradually passes away, and is substituted by a very of the acid the remainder of the base is entirely withdrawn, readily soluble by heat. If urine presenting this reaction has already been stated that this precipitate by nitric acid leaving the free acid, which is deposited in crystals. It small crystalline sediment of uric acid. By longer action with acid be allowed to stand for some time, the milky hydrated uric acid. is considered by Thudichum to be not acid urates, but

only to be mentioned in this connection. acid, having little practical significance as such, require xanthin, hippuric acid, oxalic acid, lactic acid, and phenylic The remaining organic constituents of urine, creatinin,

with lime will be further considered in treating of sediments. Mucus and the crystalline combination of oxalic acid

striking connecting links between the urine of carnivora, urine of man, .5 to 1 gram (7.7 to 15.4 grs.) is separated. sided prism, with two or four beveled surfaces at its ends, the next morning crystals of hippuric acid will usually be acid. If 10 grains benzoic acid be taken in the evening, It is increased in diabetes, where also it almost replaces uric vegetable diet, it is not wholly absent with animal food. state. But while hippuric acid is increased in man by a have both uric acid and hippuric, that is, an intermediate of the first, while in man, who consumes a mixed diet, we omnivora, and herbivora, replacing in the last the uric acid but from this there are deviations. In the twenty-four hours' found in the urine. The typical form of these is a four-Hippuric Acid is interesting in forming one of the most 177

Inorganic Constituents.

XVIII. THE CHLORIDES.

The chlorides found in the urine are chiefly those of sodium, with a small proportion of chloride of potassium and ammonium.

In health the chlorides of the urine are almost an exact measure of the same substances taken in with the food, and amount to 10–16 grams (154.3 to 246.8 grs.) in the twentyfour hours.

Detection and Approximate Estimation.—If a drop of urine be slowly evaporated on a glass slide, characteristic octohedral crystals and rhombic plates of a combination of urea and chlorine make their appearance, and may be examined by the microscope. But more available for detection and approximate estimation is—

The Nitrate of Silver Test.—Nitrate of silver in solution throws down both the phosphates and chlorides from urine. But if a few drops of nitric acid be first added, the phosphates will be held in solution, and only the chlorides will fall as opaque white chloride of silver.

From normal urine containing $\frac{1}{2}$ to 1 per cent. of chlorides, they are precipitated by a single drop of a solution of nitrate of silver, 1 part to 8, in cheesy lumps, which do not further divide themselves, or make the urine more milky by moving the glass about. If, however, the chlorides are diminished to $\frac{1}{10}$ per cent,, or less, the addition of a single drop of the silver solution no longer produces the white cheesy lumps, but a simple cloudiness, and the entire fluid appears equally milky. If, finally, there should be no precipitate whatever, then the chlorides are totally absent

interfere with the test, but if abundant it must be removed. The presence of albumin in moderate amount does not

and indeed in all acute diseases, their disappearance from they appear abundantly in the saliva. In this affection, monia, where they are often totally absent from the urine, which the chlorides seem to be eliminated. In acute pneuexcept, it is said, intermittent fever. Especially is this the in all febrile conditions, whether of local or general origin, Hence a daily trial of the urine for chlorides becomes before physical or any other signs point to improvement. in the disease may often be detected through their return reappearance an improvement. In pneumonia a decline case where there are any exudations, solid or fluid, in the urine indicates an increment in the disease, and their Clinical Significance.—The chlorides are diminished

Volumetric Process for the Chlorides.

here given. There are requiredtion of chlorides, Mohr's nitrate of silver method is r. A cold saturated solution of neutral chromate of Of several volumetric processes employed for the estima-

potassium.

diluting to a litre. grams pure fused nitrate of silver in distilled water and milligrams NaCl. This is made by dissolving 29.075 2. A solution of nitrate of silver, such that I c.c. = IO

strong heat until the carbon is completely oxidized, and Expose the remainder first to a gentle and afterward to a crucible, dissolve in it 1 or 2 grams potassium nitrate, free from chlorides, and evaporate the whole slowly to dryness. Process .- Put 10 c.c. of the urine into a platinum the residue a white, molten, saline mass. The entire white the excess of lime filtered off. To the mixture 2 or 3 while stirring the mixture, until a distinct red color remains. The color continues canary-yellow until all the chlorides are decomposed. As each drop falls into the urine, it must be carefully watched for the least tinge of red surrounding the precipitate of chloride of silver; the very next drop after the complete decomposition of the chlorides gives a permanent red color, due to the presence of silver chromate. The number of cubic centimetres estimated as NaCl, in 10 c.c. urine, whence the total is glass, and the platinum capsule washed off into it with the of calcium carbonate is introduced to make it neutral, and and the silver solution allowed to flow in from the burette consumed X .oto gram will give the amount of chlorides, mass is then dissolved in a little water, placed in a beakerwash-bottle. Dilute nitric acid is then carefully dropped into the alkaline fluid until it is faintly acid, a small pinch drops of the potassium chromate solution are now added, calculated.

XIX. PHOSPHATES.

The phosphates of the urine are composed partly of earthy and partly of alkaline phosphates. The former are cipitable from it by alkalies. The alkaline phosphates are insoluble in water, but soluble in acids; they are held in solution in acid urine by free carbonic acid, and are presoluble in water, and are not precipitated from solution by alkalies. (a) The earthy phosphates are phosphates of calcium and magnesium, and are contained in urine in but small quantities-1 to 1.5 gram (15.43 to 23.14 grains) in

Detection and Approximate Estimation.—The presence of the earthy phosphates is shown by adding any alkali, as caustic ammonia or potash.

Their quantity may be approximately estimated in the following simple way, directed by Hofmann and Ultzmann. A test-tube, 16 centimetres (6.2992 inches) long and 2 centimetres (.787 inch) wide, is filled one-third with clear or filtered urine, to which a few drops of caustic ammonia or caustic potash solution are added, and warmed gently over a spirit-lamp until the earthy phosphates begin to separate in flakes. It is then placed aside for ten or fifteen minutes for them to subside. If the layer of sediment is one centimetre (.3937 inch) high, the earthy phosphates are present in normal amount; if they occupy 2 to 3 centimetres (.787 to 1.181 inch), they are increased; if, on the other hand, only a few flakes are visible, the earthy phosphates are diminished.

Further, in normal urine the earthy phosphates are precipitated white, but if the urine contains abnormal coloring matter, they fall variously colored. If the urine contains blood-coloring matter, the earthy phosphates appear blood-red or dichroic; if there be present vegetable coloring matters, as rhubarb, senna, etc., they are colored rosy-red to blood-red, and by the biliary coloring matters yellowish-brown, and by uroerythrin, gray.

The earthy phosphates are deposited from alkaline urine, and a most important precaution must here be observed not to take such a *deposit* for an excess of phosphates. The phosphates may really be *diminished*, and yet, in con-

sequence of the reaction of the urine, a copious deposit may be present. The possible precipitation of earthy phosphates by heat alone, as a source of error in testing for albumin, has already been alluded to. This frequently occurs, and is best explained on the supposition of Dr. Brett that the earthy phosphates are held in solution in urine by carbonic acid, which, being dissipated by heat, allows the phosphates to fall. It should be further stated, however, that Dr. Owen Rees believes the phosphates are held in solution by ammonium chloride, which would also be dissipated by heat.

Clinical Significance.—The earthy phosphates are increased in the urine by diseases of the bones, especially if extensive, as in osteomalacia and rickets, in chronic rheumatoid arthritis, in diseases of the nerve centres, and after great mental strain; but especially are the earthy phosphates increased by the food and drink, some contending that all variations in the earthy phosphates are due to this cause. In renal diseases, on the other hand, the phosphates are said to be diminished. Earthy phosphates are often found deposited in conditions of dyspepsia and overwork, but this may generally be traced to changes in the reaction of the urine.

(b) The alkaline phosphates, soluble in water and not precipitated by ammonia or alkalies, form the chief bulk of the phosphates, averaging, according to Breed, 4 grams (61.72 grains) in the twenty-four hours, though Neubauer, by volumetric analysis, has seldom found more than two grams (30.86 grains) in this period. Four grams correspond to two grams phosphoric acid. They are almost wholly made up of acid sodium phosphate, with possible

traces of potassium phosphate. The acid sodium phosphate was believed by Liebig to be the cause of the acid reaction of the urine.

transmitting light distinctly, the phosphates are diminon the other hand, the fluid is but slightly cloudy, if it is denser, more cream-like, there is an increase. If, phosphates may be considered present in normal amount; fluid present a milk-like, cloudy appearance, the alkaline down in the shape of a snow-white deposit. If the entire a beaker-glass about one-third as much of the magnesian disregarded, and to a suitable quantity of urine placed in vary much in disease. Practically, therefore, they are fluid (p. 16) is added. All of the phosphates are thrown in comparatively small quantity, and, secondly, do not this is not necessary, since they are in the first place present filtering them out. For approximate estimation, however, is easily done by precipitating them with ammonia and it is necessary, first, to remove the earthy phosphates, which phates .- Accurately to estimate the alkaline phosphates, Approximate Estimation of Alkaline Phos-

Nitrate of Silver Test.—A solution of nitrate of silver added to urine throws down a yellow precipitate of phosphate of silver, and chloride of silver. Both are soluble in *amunonia*, the silver phosphate also in nitric acid, but not the chloride. If, therefore, a few drops of ammonia be added, they will promptly disappear. If now nitric acid just sufficient to neutralize the ammonia be added, the precipitate will again appear; but the moment the nitric acid is present in excess, the silver phosphate is redissolved, but the chloride remains in suspension. If now enough

183

ammonia be added again to neutralize the nitric acid, the phosphate of silver will again fall; but if an excess be added, the entire precipitate, including the chlorides, will be redissolved.

Clinical Significance of Alkaline Phosphates.—The alkaline phosphates in the urine are influenced chiefly by the food, whence they are mainly derived; phosphorus is also oxidized in the economy, and a small part of the phosphates is doubtless derived from the disintegration of nervous and muscular tissues. Any increased activity of vital processes, as in inflammations and in fevers, would, therefore, favor their increase.

Phosphatic Diabetes.—Increased attention has been recently called to the elimination of phosphates by the researches of Prof. Teissier, of Lyons, and Dr. Charles Henry Ralfe, of London. The former first described a condition which he called phosphatic diabetes, in which there is a continuous and successive discharge of phosphates in the urine attended with symptoms not unlike those of saccharine diabetes. Of the cases of this condition Teissier makes four groups. 1st. It may be observed in cer-It may precede or accompany certain affections of the 3d. It may coexist with glycosuria or alternate with it. 4th. It may run a distinct course of itself. Of 13 cases described by Ralfe, all except 2 occurred in male adults. The symptoms common to all, although varying in degree, were loss of flesh, aching rheumatic pains, chiefly of the lower back and pelvic regions, a dry, harsh skin, a tendency to boils, appetite generally ravenous, but in some tain functional derangements of the nervous system. cases a morbid refusal of food.

Volumetric Process for Phosphoric Acid.

This process is based upon the facts that-

r. When a solution of phosphate acidulated with acetic acid is treated with a solution of nitrate or acetate of uranium, a precipitate falls which is composed of uranium phosphate.

2. When a *soluble* salt of uranium is added to a solution of potassium ferrocyanide, a reddish-brown precipitate or color is developed.

The solutions required are-

1. A standard solution of sodium phosphate, made by dissolving 10.085 grains of well-crystallized sodium phosphate (Na₂HPO₄ + 12H₂O) in distilled water, and diluted to a litre; 50 c.c. then contain .1 gram P_2O_5 .

2. Saturated solution of potassium ferrocyanide.

3. Sodium acetate solution, made by dissolving roo grams sodium acetate in roo c.c. pure acetic acid, and diluting with distilled water to rooo c.c.

4. Solution of uranium acetate,* such that I c.c. will correspond to .005 gram or 5 milligrams phosphoric acid.

To Prepare the Uranium Acetate Solution.—Dissolve 20.3 grams of yellow uranic oxide in strong acetic acid previously diluted with distilled water to nearly a litre. To determine the strength of this solution, place 50 c.c. of the standard solution of

^{*}Uranium nitrate may be substituted, thirty-five grains being directly dissolved in water, but uranium acetate has heretofore been preferred because the ammonium salts in the urine interfere with the nitrate. Commercial uranium nitrate and acetate are apt to be contaminated with uranic oxides. Uranium nitrate has the advantage of keeping longer than uranium acetate.

sodium phosphate in a beaker with 5 c.c. of the solution of to a piece of the filtering paper impregnated with it. If the to 212° F.). The uranium solution is then allowed to run from a burette into the warm mixture until precipitation ceases. Then a drop of the mixture is carried by a glass rod into contact with a drop of the ferrocyanide of potassium solution on a white plate, or ponding to .1 gram of P2O5, whence is calculated the amount of sodium acetate, and heat in a water-bath to 90° to 100° C. (194° reddish-brown of the uranium ferrocyanide does not appear, continue the cautious addition of the uranium solution until the color responds to the test. The quantity used is then read off, being that which is sufficient to decompose sodium phosphate corresdistilled water to be added to make I c.c. correspond to .005 gram of phosphoric acid.

Process .- Take 50 c.c. of urine, add 5 c.c. of the Fill the burette with the uranium solution, and drop it into the mixture while warm, testing with the ferrocyanide solution. The number of cubic centimetres used multiplied by .005 will give the phosphoric acid in sodium acetate solution, and warm in a water-bath as the 50 c.c. of urine, whence calculate the quantity for the twenty-four hours. above.

XX. SULPHATES.

The sulphates found in the urine are those of sodium and potassium, the former preponderating. The quantity in twenty-four hours is three to four grams (46.29 to 61.72 grains), corresponding to two grams (30.86) sulphuric Detection and Approximate Estimation.-This down a white precipitate of barium sulphate. A little is simple with any of the barium compounds, which throw acid, as hydrochloric, should previously be added, in

order to hold in solution the barium phosphate, which is otherwise thrown down, or the acid may be previously added to a solution of barium chloride.

If to a small quantity of urine in a beaker-glass one-third as much of the acidulated solution of barium chloride (1 part to 8 plus ½ a part hydrochloric acid) is added, and there occurs an opaque milky cloudiness, the proportion of sulphates is normal; if the opacity is intense, and the whole mixture has the appearance and consistence of cream, the sulphates are increased; if, on the other hand, there is only a slight cloudiness, so that light is still transmitted, the sulphates are diminished.

Clinical Significance.—The sulphates are derived partly from the food and partly from the tissues, are increased by the introduction of sulphur compounds, sulphuric acid and its soluble combinations, by animal food, and by any causes producing increased rapidity of tissue change—as active exercise, the inhalation of oxygen, by febrile movements, and fevers. The greatest increase has been observed in meningitis, cerebritis, rheumatism, and affections of the muscular system. They are diminished in an exclusively vegetable diet.

The Volumetric Process for Sulphuric Acid.

This depends upon the principle that a solution of chloride of barium will throw down a precipitate from a given quantity of urine, so long as any sulphuric acid is present; and further, that in thus treating a specimen of urine acidulated with HCl, a neutral point is reached at which the filtrate will show a slight opacity as well with the sulphuric acid as with the barium chloride solution. In

If now either barium chloride or potassium sulphate is such a fluid we are to suppose potassium chloride, barium chloride, and potassium sulphate balancing each other. added, it itself is decomposed, and barium sulphate precipitated.

The solutions required are-

r. Solution of barium chloride so concentrated that r c.c. will precipitate exactly 12.25 milligrams H.SO,, or 10 milligrams SO3-prepared by dissolving 30.5 grams dry crystallized chloride of barium, and diluted to a litre.

2. Solution of potassium sulphate, such that I c.c. = 12.25 milligrams H2SO4, or 10 milligrams SO3-prepared by dissolving 21.775 grams chemically pure powdered potassium sulphate, dried at 100° C. (212° F.), and diluting to a litre.

Process.-Place 100 c.c. urine, acidulated with 20 to When boiling, allow 5-8 c.c. of the barium solution to flow in from a burette. Remove the heat and allow the tion to flow in, reapply the heat, and filter 10 to 12 drops of the urine into a small test-tube, add some of the barium the barium solution has been added or not. If, however, tion removed for testing, the latter is returned to the the quantity somewhat by the intensity of the reaction in 30 drops hydrochloric acid, and heat it in a water-bath. precipitate to subside. If the fluid becomes rapidly clear, allow another cubic centimetre or two of the barium solusolution, and observe whether there is a precipitate or not. If not, add to another portion a few drops of the potassium sulphate solution, by which we learn whether an excess of the barium solution still produces a precipitate in the porbeaker and more solution allowed to flow in, determining the test-tube, and the process repeated until no precipitation

188

takes place with the barium, and until a *slight* cloudiness takes place when adding the potassium sulphate to a portion of the filtered mixture. If the latter is an intense reaction, say at 12 c.c., then we know that the correct point is somewhere between 11 and 12, and the process is repeated as far as 11 c.c., when it is continued very cautiously, adding only fractions—10 of a centimetre—

until the right point is reached, whence the calculation is

made as before.

PART II.

URINARY DEPOSITS.

PRELIMINARY REMARKS—SECONDARY DEPOSITS.

Ir has already been said that strictly normal freshly passed urine, of acid reaction, contains no sediment except the faint flocculi of mucus which gradually subside toward the bottom, and entangle a few mucus corpuscles and an occasional epithelial cell. Should the urine, however, be alkaline, as is frequently the case three or four hours after a meal, it may be more or less cloudy at the moment it is passed, and quickly deposits a flocculent precipitate of earthy phosphates, which may occupy considerable bulk. They will be found by microscopic examination to be made up of amorphous granules, and will quickly disappear on the addition of a few drops of any

But even urine which is strictly normal will, in the course of time, form deposits as the result of changed reaction. These deposits differ with the stages of such reaction, and should be perfectly understood by the student before he is ready to interpret any sediment arising from other causes.

1. After normal urine, completely without sediment, has stood for a time, especially at a moderate temperature, there is often observed a precipitate of amorphous granular

matter, readily soluble by heat, which is made up of acid urates of potassium, sodium, and ammonium, with which urates of lime and magnesium are occasionally commingled. (See lower portion of Fig. 17.) A little later they are replaced by rhombic crystals of uric acid, stained yellowish or yellowish-red. These are often associated with octahedral crystals of the oxalate of lime.

The explanation given by Scherer of the occurrence of these deposits is that of the so-called acid fermentation, in which, through the agency of the mucus of the bladder, acting as a ferment, lactic and acetic acids are formed out of the coloring matters. These take away a part of the base from the neutral or alkaline urates, and produce first the more insoluble acid urates named above, which are deposited; later they combine with the remainder of the base also, and leave the crystalline uric acid sediment.

As though favoring this so-called acid fermentation, there are also often found at this stage in urine spores of torula cerevisiae—the yeast fungus—small, oval, transparent, structureless cells, to be again referred to. Sufficient proof that such fermentation takes place is, however, wanting.


A much more satisfactory explanation of the occurrence of these deposits has been offered by Voit and Hofmann,* who attribute the decomposition of the basic urates to the acid phosphate of sodium, the excess of phosphoric acid playing the part of the acetic and lactic acid in the fermentation theory, and decomposing the alkaline urates in the same way and with the same results. They prove their position by an artificial production of the same results by

^{*}Neubauer and Vogel, "Analyse des Harns," vi Aufl., 1872, p. 113. from Zeitschrift für Analyt. Chemie, Bd. 7, p. 397.

191 adding a solution of acid phosphate of sodium to a solution of basic urates. The extent to which the reaction goes will depend upon the quantity of acid phosphate of sodium present and the length of time during which the reaction

has been permitted to proceed. It is possible also for the

latter to begin at the moment of secretion, and to continue



Prismatic Crystals of Sodium Urate, Spherules of Ammonium Urate, and Amorphous Urates, with Octohedral Crystals of Oxalate of Lime. (Ranke.)

in the bladder, causing deposits of acid urates and uric acid to appear as "gravel" or "sand" immediately after the According to these authors, a more rapid action of the acid sodium phosphate produces an amorphous precipitate, and a slower separates the crystalline uric acid. The more rapid urine is passed. Such a condition would be pathological.

reaction may be induced by a more abundant separation of the acid sodium phosphate or a greater concentration of the urine.

In the course of these changes, also, the acidity of the urine is diminished, and it may become neutral and even alkaline before the phenomena of the next stage to be described—the alkaline fermentation—set in.


2. After a still longer but variable period, which is shorter in warm weather and longer in cold, we have the so-called alkaline fermentation, which is a real fermentation. This, in which decomposing mucus is also thought by some to be the ferment, is ascribed by Tieghen* to the action of a little torula, structureless and without a cell-wall, which multiplies by budding, not at the surface, but within the urine or at the bottom of the vessel, where it with the deposited salts form a white sediment. In this fermentation we have the urea converted into carbonate of ammonium, as already explained, by the addition of two equivalents of water. As the result of this conversion, the urine is rendered highly alkaline, and a further change in the character of the sediment takes place. At the very beginning of the

^{*}Neubauer and Vogel, "Analyse des Harns," vi Auflage, 1872, pp. 110 and 130.

[†]An explanation of the delay which sometimes occurs in the appearance of these phenomena is based on the recognition of the multiplication of these spores as the cause of the fermentation. If infusoria are simultaneously developed, the urea is more slowly converted, and if the surface of the urine happens to be covered with other plant vegetation (mildew), as is sometimes the case, the urine may remain acid for months in consequence of the interference with the access of oxygen, on the presence of which the spore is dependent for its growth and multiplication.

193

alkaline, the uric acid crystals begin to dissolve and to prismatic crystals of urate of sodium and dark spheres of reaction, when the urine may still be neutral or even weakly change their form so as to become more or less unrecognizable, while on their fragments may often be seen to adhere urate of ammonium (Fig. 18). As the reaction becomes alkaline, the uric acid altogether disappears, and the field



Spiculated Spherules of Ammonium Urate along with Triple (Ammonio-magnesian) Phosphate and Octobedral Crystals of the Oxalate of Lime, (Ranke.)

becomes crowded with granules of amorphous phosphate crystals) and their modifications, of the triple phosphate of ammonium and magnesium, and opaque black balls of urate of ammonium often beset with spiculæ (Fig. 18); the spores referred to are also often present, while millions of bacteria vibrate slowly along, or form granular aggregaof lime, beautiful triangular prisms ("coffin-lid" shaped

tions about a fragment of organic matter, and an occasional infusorium darts across the field of view with magnified celerity. Commonly, however, the intermediate stage is lost sight of, and the stage just described is the only one seen in the alkaline fermentation. Such urine has an ammoniacal and put rescent odor, is cloudy from the suspended phosphate of lime and bacteria, and exhibits to the naked eye an abundant white deposit.

Either of the above set of changes may take place within the body, that is, in the pelvis of the kidney or in the bladder, and as such form pathological conditions which are constantly met with in practice, the first in the condition of uric acid gravel or calculus, with its incident suffering, and the second in the phenomena of irritation and inflammation, more particularly of the bladder, due to obstruction by stone, stricture, or malignant disease. It also seems to be a matter of modern observation that the germs of the fungi above alluded to, which appear to have a very close relation to the phenomena described, either as cause or effect, may be introduced from without by the use of imperfectly cleansed catheters, sounds, or similar instruments.

With this preliminary knowledge of the rationale of the causation of a large proportion of urinary deposits, we are ready to take up their detailed consideration, previous to which, however, allusion must be made to—

EXTRANEOUS SUBSTANCES FOUND IN URINE.

These are very various, and include, indeed, all substances which are liable to get into vessels containing urine. The

most common among these are fibres of cotton and linen, hair of blankets, worsted, wool, human hair, cats' hair, splinters of wood, oil globules, starch corpuscles, tealeaves, bread-crumbs, etc. With the microscopical appearances of all these the student should familiarize himself before he begins the examination of urinary sediments.

Scratches and marks in the glass slides may also confuse, if not mislead, the beginner, and, if they become filled with coloring matters are more likely to do so. Such markings often found in glass slides, which were so long and so often described by observers as pigment flakes. They are little depressions or scratches in the glass which have become filled with oxide of iron used in the polishing of the glass. Their true character was first pointed out by the late Dr. J. G. Richardson, of this city.

CLASSIFICATION OF URINARY DEPOSITS.

Efforts have been made to classify sediments on different bases, that is, on the ground of their external naked-eye characters as to bulk, color, weight, etc.; again with regard to their nature and origin, whether organized or unorganized, crystalline or amorphous; and finally as to the reaction of the urine in which they are found.

The simplest division is into unorganized and organized. A further division of these groups into crystalline and amorphous seems to separate groups which are naturally associated, and is therefore omitted.

PRACTICAL EXAMINATION OF THE URINE.

196

Unorganized.

I. Uric acid (crystalline).

a. Acid sodium urate (amorphous, occasion ally crystalline).

II. Uric acid compounds.

b. Acid potassium urate (amorphous). c. Acid calcium urate (amorphous).

d. Acid ammonium urate (crystalline).

III. Oxalate of lime (crystalline).

 Calcium phosphate (amorphous and crystalline). a. Ammonio-magnesian phosphate (crystalline).

IV. Earthy phosphates.

VII. Leucin and tyrosin (crystalline). VI. Calcium sulphate (crystalline). V. Calcium carbonate (crystalline).

VIII. Cystin (crystalline).

Organized.

I. Mucus and pus. II. Epithelium.

V. Spermatozoids.

VII. Elements of morbid growths. VI. Fungi and infusoria.

VIII. Entozoa.


III. Blood. IV. Tube-Casts.

Unorganized Sediments.

I. URIC ACID.

be characterized by the terms "sand," "gravel," "redin their aggregation frequently form masses so large as to are often large enough to be seen by the naked eye, and adhering to the side of the glass. The individual crystals of small bulk, sinking to the bottom, but sometimes also Occurrence.—Uric acid presents itself as a sediment pepper grains." This latter term is based upon the red or yellowish-red coloration which uric acid crystals in urine exhibit.

They are found perfect only in acid urine, often at the end of the so-called acid fermentation, in urine concentrated from any cause, and where there is a pathological




More Usual Forms of Uric Acid Crystals. (After Harley.)

increase in the production of uric acid due to imperfect oxidation or assimilation.

Recognition.—The typical shapes of a uric acid crystal may be said to be a rhombic and a six-sided plate.

But it is comparatively seldom that the typical forms are observed, the latter shape being somewhat rare, and the angles of the former being generally so rounded off that

the crystals assume an ovoid or "whetstone" shape, of very different sizes, some being mere points with powers of 200 to 300 diameters, while others are large enough to be seen by the naked eye. Further shapes are those of sections of a barrel, envelope, spear, fan, of a comb with teeth on two sides, quadrilateral prisms with terminal planes, dumb-




More Unusual Forms of Uric Acid Crystals. (After Harley.)

bells, and even other forms. What are commonly called "dumb-bells" of uric acid may be rather compared to a tuft of hay constricted at its middle. These varied forms practice soon teaches one to recognize, even though they may deviate much from the typical shape. Uric acid crystals, as mentioned, are almost invariably colored, and can generally thus be distinguished from other deposits. Dr.

199

Beale* states that two or three instances have come under his notice in which they were not colored, and I have met such crystals. Uric acid crystals are met singly, but very commonly they are aggregated, forming beautiful rosettes and other shapes of aggregation of such size as



Other Unusual Forms of Uric Acid, not unlike Crystals of the Triple Phosphate of Ammonium and Magnesium. \times 150-

to be easily visible to the naked eye,—as the "red pepper grains" already alluded to,—and give pain in their transit through the ureter.

* "Kidney Diseases and Urinary Deposits," Philadelphia, 1869, p. 371.

Fig. 19 exhibits the more usual varieties of uric acid, and Figs. 20 and 21 some of the rarer forms.

Tests for Uric Acid.—Whenever a crystalline deposit is of doubtful character and suspected to be uric acid, if the latter it will respond as follows:—

r. Insoluble in cold or hot water, it will readily dissolve in the alkalies, soda, potash, or ammonia. If then the alkaline solution be treated with an excess of acetic acid, in a few hours typical whetstone-shaped forms will crystallize out.

2. Or the sediment may be placed on a glass slide, and treated with the murexid rest, as described on page 170.

The dumb-bell crystals of uric acid, occasionally met with, may be distinguished from the dumb-bell crystals of the oxalate of lime by the characteristic shape already referred to, by their larger size, their darker color, and their solubility in alkalies.

II. URIC ACID COMPOUNDS.

(a) Sodium urate, mainly amorphous, is sometimes crystalline. It always forms a part, and, according to Bence Jones, a predominant part, in the pulverulent, heavy, variously tinted, and generally bulky deposit of the mixed urates known as "brickdust" or "lateritious" sediment. The degree of coloration of this sediment depends upon that of the coloration of the urine whence it falls. From pale urine of low specific gravity, roro to ror4, an almost white sediment separates, falling very slowly, and producing, therefore, an opaque cloudy appearance in suspension, but readily disappearing on the application of heat; from urine of an amber color, and specific

gravity of about 1018, the urates deposited are fawn-colored; and from high-colored urine of higher specific gravity, we have the true red "brickdust" sediment. The sediment is found in acid urine, or urine in which the acid fermentation has only commenced, and has not been operating so long as completely to remove the base and cause the crystalline uric acid to be deposited. It is found also in urine concentrated from any cause, or where it has cooled down considerably below 37° C.




Spherules and Spiculated Spherules of Urate of Ammonium (Sodium?); Amorphous Granular Urates.

(98½° F.), or where there is defective oxidation or assimilation, as in fevers.

Recognition.—By far most frequently do we find sodium urate in fine amorphous granules, by their shape in no wise distinguishable from other fine granular matters, requiring, therefore, the chemical tests for their discrimination. The adhesion of these fine granules to partially coagulated shreds of mucus sometimes gives rise to an appearance resembling finely granular casts (Fig. 23), which is readily detected by the experienced, but which may mislead the beginner. The careful application of

heat, or the addition of a drop of acetic acid, will promptly dissipate the illusion. These granules of sodium urate also assume a larger size, and become little spherules, sometimes provided with spicules, which are considered by some (G. Bird, Beale) to be spicules of uric acid. (See Fig. 22, from Beale, Kidney Diseases.) Other



Prismatic Crystals of Acid Sodium Urate, Spherules of Ammonium Urate, and Amorphous Urates with Octohedral Crystals of Oxalate of Lime. (Ranke.)

spherules are provided with projecting and curved processes, and are believed by Hassall (second edition, page 75) and Thudichum (second edition, page 81) to be composed of sodium urate throughout. That the *spines* were also urate of sodium, Thudichum considered evidenced by their solubility in water. A modified form of the latter are

203

probably the irregularly star-shaped crystals in Dr. Beale's Fig. 110, from the urine of a patient suffering with peritonitis. But all of these forms of spherules with straight and incurved processes (thorn-apple shapes) are put down by the German observers (Neubauer and Vogel, Hofmann and Ultzmann) as crystalline forms of urate of ammonium, in which I am inclined to concur, at least with regard to those which are found at the stage of reaction intermediate between the acid and alkaline fermentations, or, perhaps, rather at the beginning of the latter, when ammonia makes its appearance, and is accompanied by the ammoniomagnesian phosphate. But any spherules which occur early in the acid reaction, or before it is possible for any ammonia to be present, are probably sodium wrate.

The sodium urate is also rarely found in dumb-bells, which are also striated and broad at the extremities like those of uric acid, but less disposed than the latter to break up at the extremities into individual acicles (Atlas of Hofmann and Ultzmann, Taf. IX). One-half of one of these dumb-bells viewed from above would be fan-

Under the same circumstances, at the end of the acid and at the beginning of the alkaline fermentation, do we also have the true prismatic *crystals* of acid sodium urate arranged in star-like masses (Fig. 23).

(b) Acid potassium urate is also amorphous, very soluble, and occurs under the same circumstances as sodium urate, as a constituent of the mixed urates.

(c) Acid calcium urate occurs very seldom, and in small quantity, as a white amorphous powder, along with the mixed urates. It is with difficulty soluble in water, and

known to have lime for its base, by leaving a residue of

calcium carbonate after incineration.

(d) Acid Ammonium Urate.—This is found, along with amorphous earthy phosphates and crystals of the triple phosphates of ammonium and magnesium, in urine in which the alkaline fermentation has commenced. It is the only urate found in alkaline urine.

Recognition of Ammonium Urate.—It is crystalline, and presents itself in the shape of smooth and characteristic "thorn-apple" spherules (Fig. 22 and 23), which serve easily to distinguish it. The spherules are soluble in hot water, and dissolve in hydrochloric or other acid with the evolution of uric acid crystals. Liquor potassæ, added to them, evolves the odor of ammonia, and they give the murexid reaction with nitric acid and ammonia.

Tests for Acid Urates.—Though the acid urates are much less soluble than the neutral urates, in requiring 124 parts of boiling water and 1150 of cold, they readily dissolve on the application of heat to the slide or test-tube containing them. They are dissolved also by the alkalies, liquor potasse, or sodæ. Treated with nitric, hydrochloric, or acetic acid (the diluted are better on account of their slower action), they dissolve, with the subsequent crystallization of uric acid. They also respond to the murexid test.

III. OXALATE OF LIME.

Occurrence.—The oxalate of lime crystals are most frequently met in acid urine, often therefore alongside of crystals of uric acid, but they may also be met in alkaline urine, along with crystals of the triple phosphate. They are

particularly abundant in the urine after a meal of rhubarb plant, after the use of tomatoes and other vegetables containing oxalic acid. There are no means by which the presence of oxalate of lime may be foretold before a microscopic examination of the urine is made. The first edition of this book contained the following: "It never forms a deposit appreciable to the naked eye, and most commonly the crystals do not descend to the bottom of the glass, but are caught as it were by the flocculi of mucus



Oxalate of Lime Crystals. (After Harley.)

which float toward the bottom, rather than occupy it."

Later and repeated observations have convinced me that in many instances the whole of this cloud-like mass, so much resembling mucus, is made up of oxalate of lime.

Recognition,—Two forms of calcium oxalate crystals are met, the *octohedra* and the *dumb-bell* crystals. The appearance of the former is somewhat different according as they are seen in the longer diameter or in the shorter. They may be said to be made up of two four-sided pyramids, placed base to base, and when viewed in the

longer diameter may readily be detected as such by the microscope. When seen in the opposite direction, their characteristic appearance is that of a square, crossed obliquely by two bright lines, and if the crystals be very small, it will appear as a square with a bright spot in the centre—a characteristic appearance by which one may soon learn to detect them, even when they are very small. They are often seen in aggregations of three, four, or more, closely adherent, and forming, as it were, microscopic calculi.

The dumb-bells, very much more rarely met with, are highly characteristic; and although we have spoken of dumb-bells of uric acid and of ammonium urate, neither of the latter presents the typical dumb-bell appearance like those of the oxalate of lime. In addition to these are found allied forms, circular and oval shapes, with darker or brighter centres, and some with partial concavities at the sides, as though passing over into dumb-bells. Dumb-bells are also met aggregated, forming microscopic calculi, which go far to explain the incipient formation of calculi.

Chemical Characters.—The form of crystals of oxalate of lime is so characteristic that there is seldom occasion to make use of the chemical tests to determine them. The only crystals which at all resemble them are certain forms of the triple phosphate. These are small crystals, modifications of the typical triangular prism, with its beveled ends, in which the body of the prism is exceedingly short, as if it were almost left out, so that the two inclined triangular ends closely approach each other, and form a crystal like that of the octohedron of oxalate of lime. Their nature may, however, be suspected by the shape of the larger crystals around them, for they never occur alone. Moreover,

the dumb-bells of uric acid or urates by the solubility of This in fact explains, perhaps, the shape of the It has been shown by Mr. Rainey and others that the presence of organic matter, as mucus, interferes they are promptly dissolved by acetic acid, while the oxalate of lime is totally insoluble in this acid. The in the vegetable acids, including acetic, but are soluble in action of acetic acid, yield their crystalline matter, leaving a framework, which maintains the original shape of the with the crystallization in the regular manner. The dumbbells of oxalate of lime can readily be distinguished from octohedra are highly insoluble in water, in alkalies, and the mineral acids. The dumb-bells, after the prolonged the latter in alkalies. crystal. crystal.

possesses a power of solution over the oxalate of lime, often holding it in solution, and he gives a method by which the The acid phosphate of sodium, according to Neubauer, latter may be obtained by its agency.*

c.c. (13.3 to 20 f3) of the urine to be tested are treated After twenty-four hours, the precipitate then occurring, which nearly always contains uric acid, is placed on a filter, washed with water, and a few drops of hydrochloric acid poured upon it. The latter dissolves out the oxalate ing manner as directed by Neubauer: † 400 c.c. to 600 with solution of chloride of calcium, supersaturated with of lime present and leaves the uric acid on the filter. Calcic oxalate in solution may be detected in the followammonia, and the precipitate dissolved in acetic acid.

^{*} Cutler's American Translation of the seventh edition of Neubauer and Vogel, p. 167.

^{*} Ibid., p. 168.

The filtrate is then diluted in a test-tube with 15 c.c. (2.83 f3) of water, and overlaid most carefully, by means of a pipette, with very dilute ammonia in sufficient quantity. At rest, the two fluids gradually mingle, and after twenty-four hours the oxalate of lime present will have collected at the bottom, and octohedra of great beauty may be studied with the microscope.

Neubauer says he has many times, in this manner, obtained considerable quantities of oxalate of lime where there was previously no deposit whatever. He has, however, in other instances, with normal urine obtained negative results, so that he is unable to decide whether the oxalate of lime should be considered a normal or abnormal constituent of urine.


undoubtedly the case that deposits of oxalate often make lin, Wöhler, and Frerichs, allege that oxalate of lime is of ammonium. Others, as Owen Rees, Aldrige of Dubsays the oxalate of ammonium is converted into carbonate tion of urine from the oxalate of ammonium, but Neubauer that the oxalate of lime is formed during the decomposiand even in tube-casts. Schunck has attempted to show infrequently found in the uriniferous tubules of the kidney, considering; for both octohedra and dumb-bells are not by the kidneys, and meeting immediately the lime salts, is no doubt that oxalic acid is, at times at least, secreted within the organism and one without. their appearance in urine some time after it has been derived from a decomposition of uric acid and urates. for which it has a strong affinity, forms the crystals we are passed. Two sources must, therefore, be admitted, one Their experiments would seem to show this, and it is Sources of Oxalate of Lime in the Urine.-There Clinical Significance.—There is no disease with which the oxalate of lime is particularly associated, nor can deposits of it be considered indicative of derangement. Abundant deposits of oxalate of lime are found in the urine of persons who are typically healthy. On the other hand, it is apt to occur where there is mal-assimilation, and hence dyspeptics are often found having oxalates in their urine, as a result rather than a cause of the affection from which they suffer.

When there are symptoms of renal calculus descending from the pelvis of the kidney, and oxalates are found in the urine, especially if they form the aggregations referred to, the latter may afford explanation of the nature of the stone. Unfortunately, too often there is no sediment whatever attending the descent of a calculus, and we must, therefore, determine its nature without such aid, or remain in ignorance. A careful examination should, however, always be made of the urine in nephritic colic, as valuable information is at times at least furnished by it, especially in the uric acid lithiasis, where uric acid sediment is often found.

IV. EARTHY PHOSPHATES.

Occurrence.—These deposits are found only in feebly acid or alkaline urine, and are the more abundant the more advanced is the stage of alkaline fermentation. They appear to the naked eye as bulky, opaque white deposits, unless they are accompanied by blood, which then more or less tinges them. The urine itself is apt to be turbid from the presence of amorphous phosphate of lime in suspension, to have an ammoniacal and sometimes a fetid

odor, though not necessarily. They are especially abundant in the urine of all irritative affections of the bladder, and often attend diseases of the spinal cord, because of paralysis of the bladder and consequent retention of urine. The earthy phosphates are the triple phosphate, or ammonio-magnesian phosphate, and the phosphate of line.



Forms of Crystals of the Ammonio-magnesian Phosphate. (After Harley.)

(a) The ammonio-magnesian phosphate (MgNH₄ PO₄6H₂O), or triple phosphate, is a crystalline deposit, of which the typical form is a triangular prism (Fig. 25) with beveled ends, very characteristic and easily recognized.

In addition to this, there is an infinite variety of modifications, with one or more corners removed, the body of the crystals variously shortened, etc. Among these forms are the small crystals already referred to as being possibly

mistaken for the oxalate of lime. There are also sometimes found beautiful, star-shaped, feathery (Fig. 26) crystals of triple phosphate, which gradually undergo conversion into the prisms, and between these two there are many intermediate forms.

(b) Phosphate of Lime (amorphous Ca₃(PO₄)₂, crys-



Stellate and Feathery Crystals of Triple Phosphate, (After Harley.)

talline CaHPO₄).—Phosphate of lime is most frequently found amorphous under the same circumstances under which the triple phosphate occurs. It is, however, frequently deposited from normal urine, by which it is held in solution during the acid reaction by the acid phosphate of sodium, or carbonic acid, or by both. At any rate, let the acid reaction be wanting, as it is three or four hours after a meal,

and a copious deposit of calcium phosphate often takes place, and is increased by boiling. In other instances, a urine may be acid in its reaction, and the boiling, apparently by driving off the carbonic acid, will cause the phosphates to go down. These deposits have more than once



Crystalline and Amorphous Phosphate of Lime.

been spoken of as possible sources of error in testing for albumin, but they promptly disappear on the addition of acids. The color of the phosphate of lime alone is not snow-white, as is that of the triple phosphate, but rather yellowish.

Not unfrequently we meet in urinary deposits crystalline

213

It is frequently associated with octohedra of the oxalate of to take on the alkaline fermentation. The occurrence of crystalline phosphate of lime seems peculiar to certain individuals, and Hofmann and Ultzmann have met persons perfectly healthy who, in the summer months, phosphate of lime (Fig. 27), which occurs sometimes alone and sometimes along with the triple phosphate. It is also met in urine of a weak acid reaction, but strongly disposed have almost daily deposits of crystalline phosphate of lime.

from other surfaces, as the gall-bladder. Dr. Beale figures Recognition.-The isolated crystals of phosphate of which form there are, however, variations. But their characteristic feature is in their arrangement, which is that of a circular rosette, in which the apices of the numerous crystals forming it all point to the centre. Phosphate of lime is also found in the shape of spherules or even dumb-The latter are said by Dr. Beale ("Kidney Diseases and Urinary Deposits," p. 357) to be deposited in decomposing mucus, not only from the urinary tract, but lime may be said to be wedge-shaped or even conical, from

occurring, as it does, in urine of different reaction; but if acid, while the octohedra are untouched by it. Uric acid it were necessary to discriminate them, the former are Chemical Characters.-All of the earthy phosphates are dissolved by acids, but are precipitated by alkalies and heat, whereas the uric acid salts are dissolved by both these agencies. The small triple phosphate crystals, which resemble those of oxalate of lime, dissolve quickly in acetic itself could scarcely ever be confounded with phosphates, Moreover, the such dumb-bells in his plate XXI, Figs. 116 and 118. dissolved by alkalies, the latter not.

PRACTICAL EXAMINATION OF THE URINE.

murexid test will not respond to phosphates, but will to uric acid.

V. CALCIUM CARBONATE

is a very rare deposit in human urine, but is found abundantly in horses' urine. When present, it occurs in small spheres, and is detected by its effervescence with acetic acid.

VI. CALCIUM SULPHATE

is also a very rare urinary sediment, occurring when found in urine of decided acid reaction and rather high specific gravity. The crystals occur in needle-like prisms like those in Fig. 28.



Crystals of Calcium Sulphate. (From Van Nüys.)


VII. LEUCIN AND TYROSIN.

Occurrence.—These crystalline deposits are only found in urine which is loaded with biliary coloring matters,

since they attend only grave destructive diseases of the liver, especially acute yellow atrophy and phosphorus

Recognition.—If suspected in urine presenting the above characters, it may be slightly evaporated, when the crystals will usually be deposited if leucin and tyrosin are

Leucin presents itself in the shape of more or less yellow-



Leucin Spheres and Tyrosin Needles.

tinged, highly refracting spheres, which may at first sight be taken for oil-drops. A little study will show them refracting light not quite so strongly, i. e., not possessing quite so wide a dark border; and by suitable illumination many of them will be found marked with radiating and concentric striae. The spherules further exhibit a peculiar disposition to aggregate, appearing partially to merge where two edges come together.

Chemical Characters.—Leucin spheres, unlike oilglobules, are insoluble in ether, and further are soluble in
caustic alkalies, but not in cold mineral acids. Spherules
of sodium urate resemble somewhat leucin spheres. The
former are, however, soluble on being warmed, and may
also be recognized sometimes by the development of spines
on their surface.

Tyrosin is found in the shape of very fine needles arranged in tufts or "sheaf"-like collections, often crossing each other and intersecting at their constricted central portions (Fig. 29).


Chemical Characters.—Tyrosin may be recognized by Hofmann's test. A suspected deposit is boiled in an excess of water. To the boiling fluid a few drops of a solution of mercuric nitrate are added, and there arises a red precipitate, while the supernatant fluid is colored red to purple-red.

VIII. CYSTIN (C3H,NSO2).

Occurrence and Recognition.—Cystin is a rare urinary sediment. Crystalline, forming a whitish or dirty yellowish-gray deposit, which on microscopic examination is found to be made up of regular six-sided tablets of different sizes, often so arranged that one of smaller size is superimposed on one of larger, and this upon a still larger, and so on; but it also occurs in irregular masses (Fig. 30). It is usually met in a pale urine, both acid and alkaline, developing in decomposition the odor of sulphuretted hydrogen, as well as that of ammonia, the former doubtless derived from the sulphur contained in the cystin. It occurs as a separate urinary deposit as well as accompanying cystin calculus, which seems sometimes to be hereditary.

217

Chemical Characters.—It is soluble in ammonia, and, upon spontaneous evaporation of the ammoniacal solution, the six-sided crystals reappear, showing that it is simply dissolved in the ammonia, and not in chemical combination with it. Now, if the six-sided crystals of uric acid, which so closely resemble it, and which often accompany it, are dissolved in ammonia, and the solution allowed to evaporate, there would be formed ammonium urate, and, on evaporation of the solution, this ammonium urate would



remain as an amorphous residue. Cystin is also insoluble in boiling water, in strong acetic and very weak hydrochloric acids; but it is readily soluble in oxalic and strong mineral acids. It is soluble in potash and insoluble in solution of carbonate of ammonium, and therefore may be precipitated from an acid urine by the alkaline fermentation; under these circumstances it would be accompanied by amorphous phosphate of lime and crystalline phosphate of ammonium and magnesium, with neither of which is it likely to be confounded. In a mixed deposit containing

six-sided crystals, the lime and triple phosphate may be dissolved out with acetic acid, while the plates of cystin will remain. They may then be treated with ammonia, as above, to distinguish them from uric acid.

Cystin contains 26 per cent. of sulphur.

Organized Sediments.

I. MUCUS AND PUS.

Mucus must be present in considerable amount to be recognized by its own properties, since it is so transparent and similar to urine in its refractive index. It is visible partly through the accidental morphological constituents which it more or less constantly holds in suspension. These are the so-called mucus-corpuscles and epithelium from all parts of the genito-urinary tract, as well as crystals of the oxalate of lime, granules of sodium urate, and even crystals of uric acid. In strictly normal urine the first two would alone be present, and in very minute quantity. Mucus, when present in normal amount, appears as a delicate cloud, often barely visible, floating toward the bottom rather than at the bottom of the vessel.

By the action of acetic acid, the mucin, an element of mucus which is comparable to albumin, though not coagulable by heat, is precipitated in the shape of delicate fibrillated bands, which are sometimes tortuous, and again appear as delicate threads, known as mucin threads. If a little iodine and iodide of potassium be added to the acetic acid, they are made even more distinct. Tartaric acid and very dilute solutions of the mineral acids have the same effect, while an excess of the same will redissolve the precipitate; so, too, the mineral acids will dissolve the coagulum of

219

ally very much narrower than the latter, and the addition by the action of the acids developed in the acid fermentation. Under these circumstances they are particularly apt to be studded with granular urates, which may cause them to be mistaken for granular tube-casts; but they are generof a little warmth, hydrochloric acid, or alkali will quickly acetic acid, while an excess of the latter will not dissolve it. These coagula may sometimes be found in urine to which no acids have been added, being probably produced dissolve the granules (see Fig. 23).

or less opaque; but even here the opacity is largely due to urine rather than sink to the bottom, giving the entire fluid, therefore, a glairy character. Mucus, however, selpus, as the causes producing them are but differences of without albumin, so long may pus be said to be absent, as As the result of irritation of any part of the genitourinary tract, mucus is increased in quantity, when it assumes a thicker, more ropy character, and becomes more the increased proportion of cellular elements. Under these circumstances, the opaque clouds of mucus are often enormously increased, and with them the adherent epithemucus is apt to pervade more or less the entire mass of the dom becomes very abundant without being attended by degree. So long, however, as urine containing mucus is lial cells from the seat of irritation. When thus in excess,

inch) in diameter, containing one or more nuclei. In a The Mucus- and Pus-corpuscles.-The mucus-corbuscle, as it appears in urine, is a small, granular, spherical or nearly spherical cell, rather larger than a blood-corpuscle, that is, .008 to .010 millimetre (3000 to 2500 of an healthy condition of mucous membrane, a mucus-corpusmucus itself contains no albumin, while pus does.

term leucocyte or white cell is conveniently applied. cle is, as it were, only accidentally arrested in its developsiologically distinct, they are anatomically the same, the above characters, with one or multiple nuclei, is found the blood, and to the whole class of cells to which the ment. The same resemblance which exists between these into the normal tissue of the part, while the mucus-corpusis a cell too rapidly produced to be allowed to develop physiological difference being in this, that a pus-corpuscle be called a pus-corpuscle. Thus, while the two are phypuscle, while the same cell on a suppurating surface would upon a non-suppurating surface it is called a mucus-corsame way. The fact being that when a cell exhibiting the hibiting the same characters, and may be defined in the (that is, not the subject of fatty degeneration), is a cell exthe difference ceases. . For the pus-corpuscle, when young nucleus may be said to be more characteristic. than it is in the pus-corpuscle, of which the multiple young cell, without morbid impression, simply arrested in minates and where the epithelial cell begins? As such a its size, for who shall say where the mucus corpuscle terit has attained the characters of such cell in its developyoung epithelial cell which has reached the surface before cle, however it originates, is nothing more nor less than a bodies exists between them and the white corpuscles of its normal development, a single nucleus is more common ment. As such, therefore, we must not too closely restrict But here

The Action of Reagents.—The mono-nucleated mucus-corpuscle, which may be considered an older mucus-corpuscle, or young epithelial cell thrown off at a later period, usually exhibits its single nucleus distinctly, without the addition of a reagent; but the majority of leucocytes

have not their nuclei visible until acted upon by certain reagents, of which two acting similarly most interest us. These are water and dilute acetic acid.

1. Action of Water.—When water is added to the pus- or mucus-corpuscle, its first effect is to cause the latter to swell up, sometimes to twice the original size, next to become smooth, the granules gradually disappearing, while the nuclei come out with great distinctness. Finally, after some time the body of the cell becomes almost, and then quite invisible, while the nuclei remain some time longer. The circumstances under which the corpuscle exists in urine are not quite identical, because in it we have a solution of



Mucus- and Pus-corpuscles before and after the addition of Acetic Acid.

organic and inorganic matters considerably denser than water, having a sp. gr. 1015 to 1025, and while the action is somewhat similar, it is very much slower; and if the specific gravity of the urine should be very high, exceeding that of the fluid in the cell, there might be no effect, or a contrary one, i.e., a shrinkage of the cell from an exosmosis of its contents.

2. Acetic Acid.—The action of dilute (20 per cent.) acetic acid is identical with that of water, except that it is very much more rapid, and the stage of distinct nuclei is reached much sooner.

3. The caustic alkalies have a rapidly solvent effect

upon these corpuscles, destroying their morphological identity, and converting them into a gelatinous adherent mass.

Characters of Urine containing Pus.—Urine con-

are already displayed in consequence of the action of water. already described as pus-cells, in many of which the nuclei and the microscope reveals hundreds of the granular cells amorphous phosphate of lime; but the latter is dissipated degree does urine containing pus resemble that containing by acids, while acids also precipitate the albumin from pus, posits albumin under the same circumstances. To a less latter on the application of heat, while purulent urine dewhich both are distinguished by the disappearance of the often resembles that due to the pale granular urates, from pus which it contains. The opacity, as well as the deposit, there is no mucus present. Such urine, when shaken up, rapidly to the bottom, so long as the reaction is acid and taining pus, deposits an opaque white sediment, which sinks becomes more or less opaque, according to the amount of Characters of Urine containing Pus .- Urine, con-

Donné's test for pus is based upon the reaction referred to between the alkalies and pus. It consists in the addition of liquor potassæ to the suspected deposit, after the supernatant urine is poured off. If the deposit is pus, it is promptly converted into a viscid, gelatinous substance resembling mucus, which adheres to the bottom of the test-tube, often permitting its inversion without falling out, and which, when it is forced to flow, does so in a continuous mass, as the albumin runs out of a broken egg. If a portion of this glairy mass be examined under the microscope, the pus-corpuscles will be found to have been destroyed, or, rather, converted into the substance itself. If the action has not been very long, or the proportion of alkali to the pus is small, the nuclei of the corpuscles may still be found

as black dots in the mass, or a certain proportion of the corpuscles may preserve their integrity.

natant fluid will be found to contain albumin, which is The glairy product referred to will be found dotted with should be taken, therefore, to determine the reaction of the urine before a mucoid deposit is decided upon, and if it is prove to be crystals of triple phosphate, while the superscopic examination may show the total absence of pus-corpuscles. These have been dissolved by the alkali. Care alkaline, another of acid reaction should be obtained. glistening points, which, on microscopic examination, has set in. Through the agency of the carbonate of and the urine contains a deposit so firmly adhering to the bottom of the bottle that it is impossible to remove it with a pipette. It must be remembered that this is not mucus, although it so closely resembles it, and although micro-Changes in Urine containing Pus .- On this same reaction is based a most important change which urine containing pus undergoes after the alkaline fermentation ammonium generated, precisely the same change is wrought, wanting in deposits of pure mucus.

Frequently, in diseases of the bladder, these changes take place within the organ, forming a gelatinous mass, which plugs up the urethra and makes it almost impossible to evacuate the bladder, thus greatly increasing the suffering of the patient. In such cases the only remedy is to wash out the bladder with weak acid solutions, and having cleansed it, keep it so by their daily use. Even when acid at the time of being passed, these urines become rapidly at the

alkaline afterward.

Sources of Pus in the Urine.—Pus in the urine may come from any part of the genito-urinary tract. When

descending from the *pelvis* of the kidney, as it often does, where there is impacted calculus, it is less apt to be mingled with mucus, the urine retains its normal reaction, and the pus is, therefore, readily miscible with the urine, and as promptly deposited from it. When coming from the *bladder*, if the urine is not already alkaline, it is apt to become so very quickly, and we have then the phenomena described as incident to the alkaline fermentation, taking place soon after the urine is passed, if not in the bladder itself.

In diseases of the *prostate* are apt to be found long plugs of mucus, which, appearing to the naked eye like fine threads, upon microscopic examination are found made up of aggregated pus-corpuscles, in which are sometimes found the larger round, or nearly round, nucleated cells peculiar to this seat. Similar plugs are found in the pus from gonor-rhoea, and it is said also that in this affection the mucus-corpuscles are distinguished from those derived from the bladder by their larger size, their "glass-like clearness," and diminished granulation. If there be no gonorrhoea, these plugs or threads point almost pathognomonically to inflammation or irritation of the prostate.

In females, pus is apt to obtain in the urine from leucorrhora or other purulent discharge from the vagina. This should not be forgotten.

II. EPITHELIUM.

Epithelium from all parts of the genito-urinary tract is found in the urine, but it is not very often that we are enabled to locate its site beyond the bladder and vagina, partly because of the comparatively slight differences in the epithelium from different locations, and partly because macera-

tion in the urine renders such feeble distinctive points even less marked.

Three varieties of epithelium may, however, be distinguished in urine with tolerable ease: 1st, round cells; 2d, cylindrical or conical and spindle cells; and 3d, squamous

(a) Round epithelial cells (Fig. 32, and a, Fig. 33) arise from the uriniferous tubules, particularly in their convoluted portion, from the deeper layers of the mucous mem-



Round Epithelial Cells from the Convoluted Tubules of the Kidney, found in Unine from a case of Acute Nephritis. \times_{420} . The cells A, E, and H are slightly more granular than in health, and C contains a few oil-drops.

brane of the pelvis of the kidney, from the bladder, and from the male urethra. Some of these cells, originally somewhat flattened by pressure, swell up in the urine and become nearly round (Fig. 32). They are distinguished from pus and mucus-corpuscles by their larger size and their single nucleus, which is distinct without the use of reagents, while the multiple nucleus of the pus-cell requires the use of acetic acid to exhibit it. There is no way of distinguishing the source of these cells more precisely than as stated above, except that if the urine be albuminous,

and there is evidence of renal disease, it may be right to infer them to come from the tubules of the kidney, or from the pelvis if there are symptoms of impacted calculus; otherwise from the urethra, the prostate, Cowper's or Littre's glands, but cells from the latter are rare. If the plugs already referred to, made up of pus-cells with a few larger, nearly round, and distinctly mono-nucleated cells united by mucus, are present, we may infer the round cells to be from the epithelium of the prostate. The round cells from the bladder are considerably larger than those from other sources—twice the diameter of a pus-cell.

(b) Columnar or conical and spindle cells (b, Fig. 33) are derived, the *first* from the superficial layers of the pelves of the kidneys, from the ureters and the urethra; the *latter* from the ureters and urethra.

(c) The flat epithelial cells (c^2 , Fig. 33) arise from the bladder or the vagina. These are flat, but often thicker at the middle, contain a single nucleus, are irregularly polygonal in outline, and often folded on themselves either completely or partially. The epithelial cells of the bladder (c^1) are not generally as large as those of the vagina (c^2), nor so flat; they are less apt to occur in layers or flakes, although also found thus. Frequently it is not safe to attempt to distinguish between the two.

In acid urine these cells remain a considerable length of time, but in alkaline urine they are gradually destroyed, becoming at first swollen and more transparent.

III. BLOOD-CORPUSCLES.

These get into the urine from the tubules and pelves of the kidneys, the bladder, the prostate, and from the uterus and vagina in their various physiological and pathological hemorrhages. They may be so abundant as to be easily distinguished in mass by the naked eye, or they may require the microscope for their detection. Urine containing blood in large amount is impressed with the red color of the latter, but containing the moderate amount most fre-



a, Round Epithelium from Bladder.
 b, Columnar Epithelium from Ureter and Urethra.

cl, Columnar and Squamous Epithelium from Deeper Layers of Epithelium

ca, Squamous Epithelium from Superficial Layers of Epithelium of Vagina. of Bladder.

scribed as "smoke-hued," and which is so characteristic as to enable one who is at all experienced to decide at on its reaction. If the urine is acid, it assumes a peculiar blackish-brown color which has long been dequently encountered in urine, it obtains a color depending

once as to the presence of blood. If, on the other hand, the urine is alkaline in reaction, it assumes the bright-red color of blood. Urine containing blood in any quantity appreciable to the naked eye is albuminous.

If blood-corpuscles are present in numbers sufficient to produce an appreciable deposit, they form a brownish-red pulverulent mass at the bottom of the vial, if they come from the kidneys or ureters. They are more apt to be found in coagula if they come from the bladder or urethra, though this latter is not necessarily the case. On the other hand, moulds of clotted blood are sometimes discharged from the ureters with all the agonies of nephritic colic.

distinctive, while the corpuscle itself shrinks and becomes concentrated, the concavity becomes more marked and appears. If, on the other hand, the urine is highly dissolved out. Ultimately the corpuscle altogether dispuscles will swell up, become biconvex instead of biconcave, of these features. If the urine is a dilute one, the coralthough the effects of long-continued maceration tend yellowish color, will always serve to distinguish them, their evident biconcavity when seen on edge, and their longer occurs, while the coloring matter is more or less finally spherical, and the reversal of light and shadow no to interfere in different degrees with the distinctness of all the slide or removed from it. This, in connection with ness or shadow as the object-glass is approximated to focusing, the centre and periphery alternating in brightcles are recognized under the microscope by the optical properties due to their biconcave centres. This is the reversal of light and shadow which they undergo in Recognition of Blood-corpuscles.-Blood-corpusTUBE-CASTS.

229

smaller, and soon acquires the crenated or horse-chestnut shape (Fig. 34).

In an acid urine the blood-corpuscles maintain themselves



Blood-disks.

for a long time, but in an ammoniacal urine they are soon dissolved, being soluble in alkalies. The hæmatocrystalline and hæmatin are then dissolved in the urine, and may be tested for as already directed.

IV. TUBE-CASTS.

Tube-casts, "cylinders," as they are sometimes called, are moulds of the uriniferous tubules, produced by admis-



Epithelial Casts and Compound Granule-cells.

sion into the latter, by capillary rupture or otherwise, of a coagulable constituent of the blood, which there solidi-

fies, and in this act entangles whatever it may have surrounded in its liquid state; subsequently it contracts and slips out of the tubule into the pelvis of the kidney, whence it is carried to the bladder and voided with the urine. Casts rarely exceed a millimetre ($\frac{1}{25}$ in.) in length.


It should be added, however, that at least two other views as to the mode of formation of casts are entertained, according to one of which the cast is a result of the disintegration and fusion of cells which may be the epithelial lining of the tubules, red corpuscles, or leucocytes; and according to another, of a secretion from these same cells as originally suggested by Rovido. That casts are sometimes formed according to the first, at least, of these two methods is not unlikely, while there is reason also to believe that the so-called "cylindroids" or mucus-casts originate in the second way.

That casts are sometimes found in urine free from albumin is beyond doubt. That persons in whom such casts occur are entirely healthy cannot be as unqualifiedly asserted.

The mechanism of the production of the different varieties of casts, on the supposition of an albuminoid exudation from the blood, is very simple. Thus, suppose a tubule to be filled with detached and loosely attached epithelium at the time the fibrin is poured into it. These elements are entangled, and, as the cast contracts, are carried out in the shape of an "epithelial" cast (Fig. 35). If the tubule should happen to have contained blood, the cast entangling it is called a "blood-cast;" if white corpuscles or leucocytes, a pus-cast (Fig. 36). Casts containing even a few blood-corpuscles are also called blood-casts. The basis substance of blood-casts is most



probably the fibrin of the blood. If the epithelium be firmly attached to the basement membrane of the tube, and remain behind when the cast passes out, or if the tube be entirely bereft of epithelium, then is the cast a "hyaline" (Fig. 37) or structureless cast. In the former instance the cast is of smaller diameter, and in the latter of larger, the diameter in the latter being that of the former plus twice the thickness of an epithelial cell. Fig. 38, a, from Rind-




Hyaline and Granular Casts, Illustrating the Formation of the Former at a. (Rindfleisch.)

fleisch, explains this sufficiently. From causes like these, as well as a subsequent contraction of the cast itself, the diameter of casts may vary considerably, ranging commonly from .or to .o5 mm. ($\frac{1}{2\sqrt{10}}$) to $\frac{1}{5\sqrt{10}}$) in.). A cast is seldom completely hyaline, generally containing a few granules and one or two glistening oil-drops, but it is still called hyaline. Completely hyaline casts do, however, occur. A variety

233


lining of the tubule or of blood-corpuscles, it is called a molten wax, is spoken of as a " waxy cast" (Fig. 39). Some hyaline casts are so delicate as to be overlooked unless the light from the mirror illuminating the field of view be modified by shading with the hand or by manipulation of the mirror itself. If a cast contains granular matter, which is generally the granular debris of the degenerated epithelial of hyaline cast, more solid in appearance, and resembling



Waxy Casts. X 150.

Finally, if a cast is loaded with oil-drops, either free or contained in epithelial cells, it is called an "oil-cast or 'granular" cast, and highly granular (Fig. 38, c), modterial of granular casts is derived from broken-down blooderately granular, slightly or delicately granular, according to the amount of granular matter present. When the macorpuscles, the casts appear yellow or yellowish-red. fatty cast" (Fig. 40).

Casts of smaller diameter are sometimes found within those of larger, the material of the latter having been poured out around that of the former after it has undergone some contraction. This occurs usually with waxy or hyaline casts. In consequence of the mode of formation above referred to, hyaline and waxy casts vary considerably in diameter, some being as narrow as .o25 millimetre (1400) of an inch) and even narrower, while others are as much as



Oil-casts and Fatty Epithelium.

.05 millimetre $(\frac{1}{300})$ of an inch) wide. There is no doubt that some of these are formed in the straight or collecting tubes near their openings on the papillæ. To these a limited number of epithelial cells is sometimes attached.

In addition to the epithelial casts above described, there are found in urine under the same circumstances moulds of the uriniferous tubules made up of simple aggregations of the epithelial cells themselves—simple exfoliations of the

235

proliferation, form a compact cellular mass. In addition to these are sometimes found epithelial casts in which the cells cellular contents of the tubule, which, having increased by are seated on the outside or around the fibrinous mould.

Mucus-casts. - Casts are occasionally found which are 41). Unless covered by accidental elements, as granular urates or phosphates of lime, they are smooth, hyaline, or apparently pure mucus-moulds of the uriniferous tubules (Fig.





Mucus-casts. (After Whittaker.)

from the kidney. Yet there is no albumin, or merely as sometimes attends them. For they are particularly apt to gently fibrillated moulds, especially characterized by their great length, which is often enormous, in the course of as the division proceeds, showing positively that they come much as could be accounted for by the presence of pus which which they divide and subdivide, diminishing in diameter occur where there is irritation of the bladder, which is ap-

parently extended through the ureters to the kidney. Under these circumstances they are frequently met. Dr. Beale says ("Kidney Diseases," etc., p. 342) they are not infrequently passed in cases where the urine has a very high specific gravity, 1030 or higher, containing an excess of urea and urates.

These casts are not identical with the bands of mucin already alluded to (p. 218), which are found in urine of highly acid reaction. The mucin bands are probably precipitated by the acids, are often beset with granular urates, and might on this account be mistaken for casts. At the same time the mucus cast is probably nothing but pure mucus or mucin.

Casts of the Seminal Tubules are sometimes found in the urine, but their origin may be inferred from the presence of spermatozoids in them.

ual disappearance the signs of the presence of albumin disand yet by the most searching examination no casts could in which the urine contained large amounts of albumin, sought. I have, however, had cases under my observation are declared absent simply because they are not carefully the other hand, the presence of albumin means casts in the albumin and casts having been present, during their gradcasts, yet I have met them in a few instances only, where, majority of instances, and many times I am certain they appeared before the last casts had been washed out. On it is not impossible for non-albuminous urine to contain may be found by careful microscopic examination. While furnish no deposit appreciable to the naked eye, and yet urine for casts. They are often so sparsely present as to The greatest caution should be exercised in examining To Prepare Urine for Examination for Casts .- be found. Not a single slide, however, should satisfy the examiner, but two or three should be carefully studied throughout their entire field. Nor is a plain slide sufficient. Urine should be examined in shallow cells, and as those of thin glass are generally too deep, the most suitable are made with gum-dammar or other suitable cement, by means of a turn-table and brush, since in this way they may be obtained sufficiently shallow to allow them to be penetrated by an ordinary one-fifth or one-fourth objective. After being made they should be put away for a month or more to thoroughly dry and harden, else they are washed off with

then be carried to the bottom of the glass with the index finger firmly pressed upon the distal end. When it has the lower strata containing the sediment turned into the reached the bottom, the finger should be raised and immethough casts generally fall to the bottom in a short time, I be allowed to stand for this time in a conical glass, and then examined. If the urine has already been standing some time, the supernatant fluid may be removed, and only conical glass, and allowed further to subside. A pipette, made of a plain glass tube drawn nearly to a point should have known twelve hours to elapse before one could be discovered, and therefore, whenever it is possible, urine should Most casts from their lightness subside slowly, and the more so because the urine is albuminous. As soon as received, therefore, the bottle of urine should be shaken up, poured into a conical glass, and carefully covered.* Althe first cleaning of the slide.

^{*} It is desirable that the upper surface of the conical glass be ground, that it may receive one of the ground glass covers referred to on p. 17. The urine is thus thoroughly protected from the action of the air, which favors decomposition and renders the examination unsatisfactory.

diately returned. In this manner only the lowest drops are obtained, which are most likely to contain the casts. A drop of this fluid is allowed to fall into one of the shallow cells, covered with the thin glass cover, and carefully examined with a one-fourth or one-fifth object-glass and the No. 1 eye-piece. If these precautions are taken, and two or three slides examined, casts will either be found, or they are absent. Only the beginner need be cautioned against linen and cotton fibre, hair, or portions of deal-wood. More likely are the mucin flakes and cast-like granular aggregations of inorganic and organic matter to mislead.

V. SPERMATOZOIDS


frequently occur in the sediment of urine of healthy men. When abundant they form a slight flocculent cloud in the urine, but there is generally nothing in the appearance of urine whence their presence may be suspected. They require a power of 400 diameters (one-fifth object-glass with the No. 2 eye-piece) to show them well, when they may be recognized by the oval head or body and the delicate tail-like prolongation emanating from it. They do not exhibit the vibratile movement after entering the urine.

Their recognition is most important in connection with medico-legal cases—cases of suspected rape. Their presence in vaginal mucus soon after coition, and in stains upon linen, is easy of demonstration. In the former case a drop of mucus from within the vagina is placed upon a slide, a drop of water added if necessary, covered with a thin cover, and examined with the microscope. In the latter a simple piece of the stained linen may be soaked in water or artificial serum in a watch-glass for half an hour

FUNGI IN URINE.

or an hour, and the sediment examined. Beale figures

FIG. 42.



Human Spermatozolds. 1. Magnified 350 diameters. 2. 800 diameters, a, viewed from the side ; b, from the front.

(Fig. 74) some filaments of a vegetable nature resembling spermatozoids.

VI. FUNGI.

Most of the living organisms found in decomposing urine, formerly looked upon as of animal origin, are now acknowledged to be vegetable in their nature, and are generally fungi.

The most frequent among these are bacteria, penicillium glaucum, and the yeast fungus. Sarcinæ are occasionally


r. Bacteria.—In the refined study which has of late years been given to the subject of fungi, a classification has been made of the minute objects which were formerly called

stricted at the joints like the moniliform threads sometimes substance. The last often form swarms but never zooglea of jelly-like masses apparently held together by a gelatinous either isolated or in the so-called zooglea form, consisting formed by the globular bacteria. The first two occur extend entirely across the field of view. It is not conand often forms a long string called leptothrix, which may form, vibrio. Bacillus increases by transverse division, across the field; c, desmobacteria, or filamentous bacteria, including, 1st, a straight form, the bacillus, and, 2d, a curved lines in breadth, sometimes at rest and sometimes vibrating moderate powers, the diameter of a red blood-disk, but mere which appear as minute lines approaching in length, with urine; b, microbacteria, the staff-shaped or rod-bacteria, ating rapidly in all putrid fluids including decomposing ing of little trembling points uniform in size and proliferclassification, a, the sphero-bacteria or micrococci, consistschyzomycetes or cleft fungi. They include, of Cohn's

the aid of the magnesian fluid (p. 16), gentle heat, and the pores of the filter. They may be entirely removed by tration, the bacteria being so small that they pass through it a cloudiness which can be only partially removed by fil-The effect of the presence of bacteria in urine is to give

consists, in the sporule-stage, of transparent oval cells, in the ordinary yeast fungus (saccharomyces cerevisia). The found in saccharine urine, and are probably identical with larger spherical cells, granular and nucleated. They are their longer diameter about the size of a blood-disk, and of 2. The yeast or sugar fungus (saccharomyces urinæ) former small oval cells are often arranged in rows of two, three, or more, as seen in the figure (Fig. 43). According to Hassall, this is a fungus peculiar to saccharine urine, but the small oval cells of the sporule-stage at least cannot be distinguished from the similar stage of—

3. Penicillium glaucum, which occurs in acid urine with or without albumin or sugar. The sporule-stage furnishes cells very similar to those of the yeast fungus, but later, penicillium, by the union of its cells, forms thalli or branches, which are characteristic. So, too, in the stage



of aerial fructification, the penicillium multiplies by simple

linear division of cells.

4. The sarcina urinæ is a fungus rarely met with. I have found it twice in fifteen years, once in acid urine, a second time in urine of unknown reaction. Composed of cubes, it is capable of further separation into smaller cubes. It is similar to, but smaller than, the sarcina ventriculi of Goodsir.

The germs of these fungi doubtless enter the urine, in the vast majority of instances, after it has passed from the bladder, one or the other form being developed according

the presence of the bacteria, but not to the other forms. the urine may possess. Decomposition seems essential to as its germs preponderate, or according to the properties

VII. THE ELEMENTS OF MORBID GROWTHS.

cells of the bladder during inflammation or irritation. abnormal. Indeed, every shape of cell may arise from the even the bladder, and must not, therefore, be considered remembered, may be derived from the ureter, urethra, and diversity of the cell-forms. Spindle-cells, it must be multinuclear character, the large size of the nuclei, and suspected to be of morbid origin by their large size, their broken off, and passed with the urine. The former may be be found, and perhaps fragments of the growth may be These are seldom met in the urine. Possibly cells may

epithelial covering. Fragments suitable for examination illary vessels which make up the villus, with or without the are sometimes withdrawn with the catheter. are generally from the villous kind, and may show the cap-Fragments of cancerous growths which get into the urine

VIII. ENTOZOA.

examining one of the slides containing ova, sent to this privilege, through the kindness of Dr. S. W. Gross, of hæmaturia of the Cape of Good Hope, and I had the ciliated embryos of Bilharsia hamatobia have been found by Dr. John Harley in three patients with the endemic passed in two or three instances recorded. The eggs and Echinococcus cysts, as well as their hooklets, have been Entozoa are seldom found in the urine in this climate. country. The parasite itself is found in the vesical, mesenteric, and portal veins, causing hemorrhages into the intestines, bladder, ureters, and pelves of the kidney. The ova and parasite are figured by Beale, op. citat., p. 402.

The filaria sanguinis hominis, the parasite which has recently been shown to have such an intimate association with chyluria, is sometimes found in urine.

Distoma hamatobium has been found in the bladder, ureters, and pelves of the kidney, especially in Egypt.

The Preservation of Organized Urinary Sediments for Subsequent Examination.

Crystalline urinary sediments are so easily obtained that there is no advantage in attempting their preservation, which is always difficult. Organized sediment, on the other hand, may be preserved tolerably unaltered in several fluids. One of the simplest and best of these is a mixture of glycerin and distilled water in such proportion as to secure the average specific gravity of urine, about 1020, to which is added carbolic acid in the proportion of 1 part to 100. A weak solution of salicylic acid is also efficient, and Dr. W. W. Keen recommends a solution of chloral, ten grains to the ounce. Dr. Beale's naphtha and creasote solution is efficient, but much more troublesome to prepare.

Dr. E. S. Wood, of Boston, recommends very highly a filtered solution of acetate of potassium, specific gravity between roso and roso, to which carbolic acid has been added in the proportion of 4 to 5 c.c. of the deliquesced crystal to one litre of the acetate solution. Instead of carbolic acid, salicylic acid may be added to saturation. When urine is to be transported in hot weather, or kept

several days for any reason, a pinch of salicylic acid added to a four-ounce vial is generally sufficient to prevent decomposition, and in no way impairs the reactions or alters the sediments.

To use any of these fluids, allow the sediment to subside in a conical glass, decant the supernatant fluid, replace the latter with the preservative, stir up the sediment, allow again to subside, again decant, and replace with a fresh portion of the preservative. Repeat this until the urine is thoroughly replaced by the preservative. Then place the sediment and preservative in a well-stopped ordinary vial, in which they may be kept for years without marked change; or the sediment thus permeated by the preservative may be mounted in shallow cells over which a thin glass cover is cemented.

DIFFERENTIAL DIAGNOSIS OF RENAL DISEASES.

While it is quite impossible to determine with absolute certainty, by a mere examination of the urine, all of the different affections to which the kidneys are liable, there is, nevertheless, an association, more or less close, of signs with well-determined conditions. With such association, it is important that we should be familiar, while we should as well recognize the fact that they are subject to variations and exceptions. If these facts are remembered, it is not likely that any one can be led far astray by observing the following:—

I. Acute Parenchymatous Nephritis (Acute Diffuse Nephritis: Scarlatinal Nephritis; Acute Tubal Nephritis).—The urine is scanty, dark, "smoke-hued," so long as it remains acid, but becomes red if alkalized. It is highly albuminous.

tubules, more or less round and nucleated, twice as wide as the blood-disks, cloudy, and more granular than in health, the granules often obscuring the nucleus. Crystals of uric acid are often present. The chlorides are first diminished, also the earthy phosphates. Hæmatin, microscopic examination, is found made up of large epithelial casts, blood-casts, casts formed of leucocytes, hyaline casts, dark-red granular casts; also, numerous red blooddisks, some leucocytes, and free cells from the uriniferous ence of blood. It contains a variable but generally large amount of reddish-brown, pulverulent sediment, which, on Its specific gravity is not constant, but apt to be high-1025 or above-not from an increase in urea, but from the pres-

and, if a child, has had scarlet fever or diphtheria, or, if an adult, has been exposed to rain, or has otherwise The patient is dropsical, much swollen about the face, indican, and uric acid are increased.

scarlatinal nephritis, or acute Bright's disease, and the The disease is acute nephritis, acute diffuse nephritis, become wet while over-heated.

granular casts, hyaline casts, and casts containing fragments of epithelium; also renal epithelium and compound granurine is pale, and of low specific gravity, 1010-1015; its quantity, though variable, generally diminished. Albumin is diminished as compared with (I), but it is still abundant -one quarter to one-half the bulk. The urine often deposits an appreciable white sediment, which by microscopic examination is found made up of black, highly ule-cells (Fig. 35). Probably also there are casts contain-II. Chronic Parenchymatous Nephritis (Tubal Nephritis; Chronic Diffuse Nephritis; Large White Kidney).-The chances for recovery are many.

ing a moderate quantity of oil, and perhaps also partially fatty cells. Waxy casts are also sometimes found in this form of disease. The urea is diminished, the chlorides normal, pigment diminished.

There is also often, but not always, cedema, more or less general, which may, however, subside, but the patient has a pale, almost characteristic waxy look; and toward the unfavorable termination there is much dropsy. The symptoms have existed more than six weeks.

The disease is probably the large white kidney, a chronic continuation of (1), known also as chronic tubal nephritis, and chronic diffuse nephritis, and recovery, though possible, is much less likely to occur than with the acute form

sible, is much less likely to occur than with the acute form.

At other times the sediment contains a large number of oil-casts filled with free oil, and oil contained in epithelial cells. There are numerous free fatty cells and free oil-globules.

Vephritis.—The disease has existed for more than a year, the urine is increased as compared with (II), and may be even increased as compared with health, and the specific gravity varies accordingly. The albumin is diminished, but may be considerable, or it may be very small in amount. The urine deposits a more scanty sediment, made up of broad casts, some dark granular and others waxy, together with a few narrow pale casts. Compound granule cells occur, but are less numerous, and there may be some fatty epithelial cells, but the amount of oil, though distinctive, is not very large. The urea is much diminished. There is often some dropsy, less frequently than in (I) and (II), but more than in (IV), and it may be entirely absent.

Here the large white kidney has probably commenced to

contract, and the resulting organ is also called the fatty and contracting kidney, to distinguish it from the next form, the chronically contracted kidney. One must be cautious about drawing too sharp a line between these (II) and (III). The prognosis as to recovery is unfavorable, but the disease may last many years without inconveniencing the patient.

IV. Interstitial Nephritis (Chronically Contracted Kilney; Granular Kidney; Red Granular Kidney; Cirrhotic Kidney.)—The urine is increased in amount, correspondingly pale, but, while micturition may be a little more frequent, it may not attract attention. The patient may have to rise once in the night. The specific gravity is generally diminished (1010-15), while the quantity of albumin is trifling, never exceeds one-quarter, and often is shown by a mere line of opacity in Heller's test. The urine deposits often no visible sediment, and at all times a trifling one. In this are found delicate hyaline and finely granular casts, often of small diameter. Some of these contain one or two glistening oil-drops, but very minute. Here are found the casts, which are at times almost invisible. The urea is

generally slightly diminished.

There is no dropsy. Hypertrophy of the left ventricle is constant, and there may be nausea and vomiting, especially in the morning, but there are often no obvious symptoms whatever connected with the disease. If any, the patient may complain of a weak, tired feeling, and this symptom should suggest an examination of the urine always. The disease may exist for years without the knowledge of the patient, who may or may not be subject to gout. Nausea is a common symptom in the advanced stages, and drowsiness a serious one. (The urine of gouty patients should be frequently examined.)

The disease is interstitial nephritis, and its product the chronically contracted kidney. If exposure to cold and fatigue be avoided, the patient's life may be scarcely shortened, and yet he is constantly liable to attacks of uræmia, which may suddenly terminate his life, and in advanced stages of the disease the cardiac symptoms often produce great discomfort.

V. Lardaceous or Amyloid Degeneration of the Kidney.—
The urine is increased in quantity, clear, of corresponding specific gravity (1007–1015), of a pale golden color, the color of a dilute urine only, contains at first little, but later considerable albumin, about one-fourth to one-half; urea is diminished. There is very little or no sediment visible. Casts are often wanting, and when present include the broad dark granular as well as the hyaline and waxy casts; occasionally fatty casts are found; the waxy are solid-looking, and sometimes give the characteristic red reaction of the amyloid substance when treated with a watery solution of iodine and iodide of potassium. Here hyaline and waxy casts of large diameter are found, and sometimes within these smaller casts.

While the highly refracting waxy casts are not confined to albuminoid kidney, they always indicate chronic and deep-seated processes.

At first there is no dropsy, but later it is sometimes persistent. Generally, however, except toward the termination of the case, it is amenable to treatment by rest and diuretics. The patient has an enlarged liver or spleen, sometimes obstinate diarrhoa; he has had syphilis, or extensive disease of the bones, or has phthisis.

The disease is lardaceous degeneration of the kidney, and is incurable, though the patient may live many years.

The characters of the urine in amyloid disease are often

the same as those of chronic nephritis, and the same is true

of the symptoms.

VI. Acute Active Hyperamia (Parenchymatous Degeneration of the Kidney; "Cloudy Swelling").—Most frequently the sole symptom is albuminuria, the most careful examination failing to discover casts; when casts are present, they are of the hyaline variety. There is, as a rule, no dropsy. The quantity of albumin (To to the bulk) is generally less than in tubal or parenchymatous inflammation or lardaceous degeneration. Such may sometimes be the albuminuria of pregnancy, or such grave diseases as diphtheria and acute

febrile disorders.

After death, the renal epithelia are often more or less enlarged, their contents cloudy. This condition differs from parenchymatous nephritis in the smaller extent and diminished intensity of the morbid process. It is probably due to the pernicious influence of some poison on the minute structure of the kidney, which may extend to all

the tissues. Recovery is frequent.

The disease was called by Niemeyer parenchymatous

degeneration.

VII. Cyanotic Induration.—Cyanotic induration is a peculiar indurated form of kidney due to a simple hyperplasia of its interstitial tissue, the result of long-continued passive congestion. It is most frequently found accompanying valvular disease of the heart, and is characterized by a bluish color.

In addition to the other symptoms of heart disease there is dropsy and often serous effusion in the great cavities. The urine is scanty, of high specific gravity, often 1030 and above, there is usually a moderate amount of albumin,

and a few small hyaline or faintly granular casts; but casts are often absent.

The prognosis is not good, but under favorable circumstances the patient may improve and become more comfortable.

The above is given as a general guide, and I would again refer to the fact that there are deviations from the conditions laid down. There are still many points quite disputed in the pathology of the kidney. Thus, the older German pathologists contended that there is a constant relation of succession between the acute parenchymatous nephritis, the chronic parenchymatous nephritis (large white kidney), and the contracting stage of the latter, making no distinction between the cirrhotic kidney and the fatty and contracting kidney. And it is held by some to-day that all inflammations of the kidney are diffuse; that is, there is no inflammation in which the epithelial tissue or the connective tissue alone is primarily involved, but that both always share the process from the beginning.

One more fact must be mentioned in this connection, and this is that although the presence of fatty casts and fatty epithelium is an unfavorable symptom, yet it does not follow that such cases are necessarily fatal. I have, on more than one occasion, found oil-casts in the urine of patients, and yet have also found them to disappear altogether. The circumstances under which this has most frequently occurred have been, 1st, where there have been heart disease and kidney disease combined, and there has been some exacerbation of one or both, when the albumin has increased, and oil-casts have made their appearance, which later totally disappeared; 2d, where pregnancy has supervened on existing Bright's disease, and oil-casts have been present, which again disappeared after a successful labor.

PART III.

URINARY CALCULI.

THE qualitative analysis of gravel or calculus is much simpler than is generally supposed. There are but three varieties of calculus at all common, and therefore likely to demand analysis. These are (1) uric acid and its compounds, (2) exalate of lime, and (3) the mixed phosphates. Calculi of xanthin and cystin occur, though very rarely.

i. Uric acid calculi are the most common. They are either red or some shade of red, and usually smooth, but may be tuberculated. They leave a mere trace of residue may be increased to the contract of the co

after ignition.

2. Oxalate of lime calculi are frequently met with.

They are generally of a dark-brown or dark-gray color, and

from their frequently tuberculated surface have been called mulberry calculi. They may, however, be smooth when small—hemp-seed calculi. Considerable residue remains after ignition. The calculus is soluble in mineral acids without effervescence.

3. Calculi of the mixed phosphates, or fusible calculi, are composed of the phosphate of lime and of the triple phosphate of ammonium and magnesium. Phosphates make up the external layers of many calculi of various com-

position, and may form entire calculi, but seldom constitute the nucleus alone, of a calculus. Mixed phosphatic calculi are white, exceedingly brittle, fuse in the blowpipe flame and are soluble in acids, but insoluble in alkalies.

Other rarer forms of calculi are made up of carbonate of lime, of xanthin, cystin, and of urostealith.

Few calculi of large size are of the same composition throughout, and when of any considerable size, these constituents are usually arranged in concentric layers about a nucleus. Oxalate of lime is the most frequent nucleus; uric acid may also serve as a nucleus, but phosphates, almost never. Small masses of organic matter, as bloodclots, frequently form nuclei, and may often be recognized by the odor of ammonia on ignition. Foreign bodies, as pieces of pencil or even glass, introduced into the bladder from without, may become nuclei.

To Determine the Composition of Calculi Qualitatively.

Previous to chemical treatment a calculus should be powdered, and in view of the fact that different layers are often of different composition, wherever the stone is of any size, it should be sawed or coarsely broken and a portion of each layer should be examined separately.

Expose a portion of the powdered calculus, on a piece of platinum foil or a platinum spoon, to a dull red heat for a considerable time. Note whether there is a residue.

A calculus which burns with very little or no fixed residue is composed either of uric acid or ammonium urate, of cystin, of xanthin, or of urostealith. If it does not burn completely it may contain uric acid and uric acid salts, phosphate of lime and phosphate of magnesium, the ammonio-magnesian phosphate, or oxalate of lime.

A. There is a fixed residue. To a portion of the original powder apply the murexid test (p. 170).

A purple color results: uric acid is present.
 Observe whether a portion of the calculus melts on being heated.

a. It melts, and communicates-

 A strong yellow color to the flame of a spiritlamp, or Bunsen burner: sodium wrate.

2. A violet color to the flame: potassium urate.

b. It does not melt. Dissolve the residue after ignition in a little dilute HCl, add ammonia until alkaline, and then ammonium carbonate

solution.

I. A white precipitate falls: calcium urate.

2. No precipitate. Add some hydric sodic phosphate solution; a white crystalline precipitate falls: magnesium urate.

II. No purple color results.

Observe whether a portion of the calculus melts on being exposed to the blowpipe flame.

a. It melts (fusible calculus). Treat the residue with acetic acid; it dissolves. Add to the solution ammonia in excess; a white crystalline precipitate falls: ammonio-magnesium phosphate. In case the melted residue is insoluble in acetic acid, treat with HCl; it dissolves. Add to the solution ammonia; a white precipitate indicates calcium phosphate. Ammonio-magnesian phosphate and basic calcium phosphate ordinarily occur mixed in the same concretion.

b. It does not melt. (r) Moisten the residue with water, and test its reaction with litmus-paper; it

is not alkaline. Treat with HCl; it dissolves without effervescence. Add to the solution ammonia in excess; a white precipitate occurs; calcium phosphate. Calculi of pure calcium phosphate are rare, but may occur.

(2) Treat the powdered calculus with acetic acid; it does not dissolve. Treat the residue, after heating, with acetic acid; it dissolves with effervescence: calcium axalate.* The powdered original calculus dissolves with effervescence when treated with acetic acid; calcium carbonate.

B. There is no fixed residue. Apply the murexid test (p. 170).

I. A purple color is developed.

a. Mix a portion of the powdered calculus with a little lime, and moisten with a little water; ammonia is evolved, and a red litmus-paper suspended over the mass is turned blue: ammonium urate.

b. No ammonia: uric acid.

II. No purple color.

a. But the nitric acid solution turns yellow as it is evaporated, and leaves a residue insoluble in potassium carbonate: xanthin.

b. The nitric acid solution turns dark brown, and leaves a residue soluble in ammonia: cystin.

c. The calculus, soft when fresh, dark-brown and brittle when dry, becomes softer again when

^{*}The calcium oxalate is converted by the heating into calcium carbonate, which dissolves with effervescence. If the heat be much higher than a dull red, the carbonate of lime is converted into quick-lime, which does not effervesce on adding an acid.

warmed. Soluble in ether, the amorphous mass, after evaporation of ether, becomes violet on being heated. It dissolves in nitric acid, with slight evolution of gas and without change of color: urostealith.

A coarser mode of analysis, but still sufficiently accurate for practical purposes in most instances, is the following:—

Powder a portion of the calculus and ignite upon a piece of platinum foil.

A. There is a fixed residue. To a portion of the powdered original calculus apply the murexid test (p. 170):

I. A purple color results. The stone is composed of uric acid or uric acid and its compounds.

II. No purple color results.

Observe whether a portion of the original calculus melts under the flame of the blowpipe.

a. It melts (fusible calculus). The stone is the ammonio-magnesian phosphate, containing, also, probably, some phosphate of lime.

b. It does not melt. (1) Treat some of the original calculus in powder with acetic acid. It does not dissolve. Treat the residue, after heating, with acetic acid. It dissolves with effervescence. The stone is exalate of lime.

(2) Moisten the residue after incineration, with water, and test with litmus paper; it is not alkaline. Treat this residue with HCl; it dissolves without effervescence. Add the solution ammonia in excess; a white precipitate occurs; calcium phosphate.

56 PRACTICAL EXAMINATION OF THE URINE.

(3) The powder of the original calculus dissolves with effervescence. The stone is calcium carbonate.

B. There is no fixed residue. Apply the murexid test (p. 170).

I. A purple color results. The stone is uric acid or uric acid and its compounds.

II. No purple color. (See II, p. 255.)

APPENDIX.

MODE OF RECORDING AN EXAMINATION.

To systematize and facilitate the work of urine examinations, forms of record have been devised by those working in the subject. For ordinary use in hospital and private practice, a modification of the form suggested by Heller recommends itself for its economy and convenience.

The form may be printed, but as recommended by Heller, an ordinary half-sheet of letter-paper is folded in four, and marked in the manner indicated below:—

PHYSICAL PROPERTIES.
Quantity taken in twenty-four hours.
Color and reaction.
Sp. gr., quantity and character of sediment.

Uph. (Urophain). Cl. (Chlorides). Ux. (Uroxanthin). Eph. (Earthy phosphates). U. (Urica). Sulphates. Sulphates.

NORMAL CONSTITUENTS.

ABNORMAL CONSTITUENTS IN SOLUTION.

SEDIMENT.

CONCLUSION.

may be used, and for slight increase or slight diminution, "sl. +" or "sl. -." increase or great diminution, "gr. +" and "gr. -" diminished, and the letter "n" for normal. For great shown, the sign "+" for increased, the sign "-" for Abbreviations for the important constituents are used as

the following results. The word "indican," "ind.," is I have added another space for the opinion or diagnosis. Let us suppose an examination to have been made, with

preferred for "uroxanthin," and substituted.

PHYSICAL PROPERTIES.

Sp. gr., 1005. See Quantity in twenty-four hours, y-tour hours, 500 c.c. rellow. Reaction, acid. Sediment, moderate, flocculent.

NORMAL CONSTITUENTS.

ABNORMAL CONSTITUENTS IN SOLUTION. Albumin, 2 per cent.

SEDIMENT.

Numerous oil-casts, free fatty cells and free oil-globules.

Diagnosis-Chronic Parenchymatous nephritis. Fatty kidney.

Grains to Milligrams.

I = 64.8 (--.000425)

2 = 120.6

3 = 194.4

4 = 259.2

5 = 324.0

6 = 388.8

7 = 453.6

8 = 518.4

9 = 583.2 259 For Reducing the Metric or French System into the English, and vice Minims to Cubic Centin

1 = .0616

2 = .1232

3 = .1848

4 = .2464

5 = .3696

7 = .4312

8 = .4928

9 = .5544 3.7 7.4 11.1 14.8 18.5 22.2 25.9 29.6 33.3 versa, as far as required in Urinalysis. H H H H H H H H H TABLES TABLES. Cubic Centimetres to Fluidrachms.

1 = .27 (+.oco5)

2 = .54

3 = .81

4 = 1.08

5 = 1.35

6 = 1.62

7 = 1.89

8 = 2.16

9 = 2.43 15-43 (+ .0022) 30.86 46.29 61.72 77.15 92.58 108.01 16.2 (= .0293) 32.4 48.6 64.8 81.0 Cubic Centimetres to Minims.

1 = 16.2 (= .0293).

2 = 32.4
3 = 48.6
4 = 64.8
5 = 81.0
6 = 97.2
7 = 113.4
8 = 129.6
9 = 145.8 Grams to Grains.

1 = 15.43 (-2.2)
2 = 30.86
3 = 46.29
4 = 61.72
5 = 77.15
6 = 92.58
7 = 108.01
8 = 123.44
9 = 138.87

PRACTICAL EXAMINATION OF THE URINE.

9 = 228.6	8 = 193.2	7 = 177.8	6 = 152.4	5 = 127.0	4 = 101.6	3 = 76.2	2 = 50.8	I = 25.4 (+.00005)	Inches to Millimetres.	9 = 18.9	8 = 16.8	7 = 14-7	6 = 12.6	5 = 10.5	4 = 8.4	3 = 6.3	2 = 4.2	I = 2.I (+.013188)	Litres to Pints.	9 = 304.2	8 = 270.4	7 = 236.6	6 = 202.8	5 = 169.0	4 = 135.2	3 = 101.4	2 = 67.6	1 = 33.8 (+.011)	Litres to Fluidounces.
935433	8 = .31496	7 = .27559	6 = .23622	5 = .19685	4 = .15748	3 = .11811	2 = .07874	и = .03937	Millimetres to Inches.	9 = 4.257	8 = 3.784	7 = 3.311	6 = 2.838	5 = 2.365	4 = 1.892	3 = 1.419	2 = .946	1 = .473 (+.00022)	Pints to Litres.	9 = 270	8 = 240	7 = 210	6 = 180	5 = 150	4 = 120	3 = 90	2 = 60	I = 30 (4238)	Fluidounces to Cubic Centimetres

Metres to Feet.

I = 3.28 z = 6.56 3 = 9.84 4 = 13.12 5 = 16.40 6 = 19.68 7 = 22.96 8 = 20.52 9 = 29.52TABLES.

Feet to Metres.

I = .3048 (+.0000005) z = 6.56 3 = .9144 4 = 1.2192 5 = 1.5240 6 = 1.8288 7 = 2.36 8 = 2.4348 9 = 29.52

To Convert Degrees of Fahrenheit's Thermometer to Centigrade, and vice versâ

Centigrade to Fahrenheit.

1 = 1.8
2 = 3.6
3 = 5.4
4 = 7.2
5 = 9.0
6 = 10.8
7 = 12.6
8 = 14.4
9 = 16.2

Centigrade.


Fahrenheit to Centigrade.

5 = 5.75 (+.000555)
7 = 1.10
7 = 1.65
7 = 3.330
7 = 3.330
7 = 4.440
9 = 16.2
9 = 4.995

To use this table, convert the given to give this to number of degrees Centigrade into the given num degrees Fahrenheit, and add 32°. degrees Centig

To use this table, subtract 32° from the given number of degrees Fahrenheit, and convert the remainder into degrees Centigrade. (From Dr. Craig's Decimal System.)





Cholesterin, 153 Chylous urine, 22 Chlorides, 177 Casts of the uriniferous tubules, 229 Calcium carbonate, deposits of, 214 Calculi, urinary, 251 Blood-corpuscles in urine, 226 Blood, coloring matters of, in urine, 131 Bismuth test for glucose, 86 264 Biliary coloring matters, 139 Biliary acids, Pettenkofer's test for, 145 Biliary acids, 144 Bacteria, 239 sulphate, 214 nitrate of silver test for, 177 Mohr's nitrate of silver volumetric process for, 178 clinical significance of, 178 detection and approximate estimation of, 177 quantitative estimation by Oliver's method, 149 Oliver's peptone test-papers for testing for, 150 Oliver's new peptone test for, 147 to determine composition of, 252 recognition of, 228 Ultzmann's test for, 142 Heller's test for, 141 Marechalt's test, 142 Gmelin's nitrous acid test for, 140 Fleischl's modification of Gmelin's test, 140 decomposed, test for, 143 INDEX.

Creatinin, 176

Creatin, 176

normal, 121 vegetable, 138 Coloring matters, 121

INDEX.

-pellets, 82

Cupric test-papers, 84

Cystin, chemical characters of, 217 deposits, 216

Deposits, urinary, organized, 189 Diacetic acid, 114

chloride of iron test for, 115 clinical significance of, 117

Dumb-bells of oxalate of lime, 206

Entozoa, 242

Extraneous substances found in urine, 194 Epithelium, 224

Fehling's solution, 75 Fat in urine, 22, 152

Fibrin, 69

Fruit-sugar, 109 Fungi, 239

Globulin, 58

by Roberts' fermentation test, 85 alphanaphthol test for, 97 Glucose, 70

Nylander's modification of the bismuth test for, 88 Brucke's modification of the bismuth test for, 87 Bottger's bismuth test for, 86 copper tests for, 72

with glycerine, for testing and estimating, 76 Fehling's solution for testing and estimating, 75

indigo-carmine test for, 93 fermentation test for, 85 lead process for, 106

Pavy's solution for testing and estimating, 76 phenyl-hydrazin test, 95

265

INDEX.

Glucose, picric acid test for, 88

quantitative estimation by picric acid, 89 Purdy's modification of the ammoniated cupric test, 84

to detect the presence of, by specific gravity and quantity, 71 by polarimetry, 99

Salkowski's modification of Trommer's test, 74 Trommer's test for, 72

Gonorrhœa, pus from, 224 volumetric process for estimating, 79

Hæmatin, 132

Heller's test for, 135

test for, by precipitation of albumin, etc., 136

Hæmaturia, 134

Hæmin crystals, to prepare, 135

Hæmoglobin, 131

clinical application of Mahomed's test for, 133

Mahomed's test for, 132

Hæmoglobinuria, 134 Stevenson's modification of Mahomed's test for, 133

Hemialbumose, 67 Haycraft's method of estimating uric acid, 171

test for, 68

Hydrobilirubin, 122 Hippuric acid, 176

test for, 123

Indican or indigogen, 126

Heller's test for, 126 clinical significance of, in urine, 128

Jaffe's test for, 127

Inosite, 107 Inorganic constituents of urine, 177

Introduction, 13

Kidney, acute inflammation of, 244 hyperæmia of, 249

267 recognition of, as a urinary sediment, 213 phosphate of, as a urinary sediment, 211 Lardaceous degeneration of the kidney, 248 chemical characters, 213 Lead process for testing for glucose, 106 Ludwig, theory of secretion of urine, 13 INDEX. chronically contracted, 247 chemical characters of, 216 as a urinary deposit, 215 fatty contracting, 246 large white, 245 detection of, 215 lardaceous, 248 Lime, carbonate of, 214 Kidney, amyloid, 248 sulphate, 214 Lævulose, 109 Leucin, 151 Lactose, 109

Mahomed's method of testing for hæmoglobin, 132

Morbid growths, elements of, in urine, 242

Methæmoglobin, 131

Melanin, 136

Milk sugar, 109

in normal urine, 59

Mucin, 59

to test for, 60

Nephritis, acute parenchymatous, 244 chronic parenchymatous, 245

Murexid test for uric acid, 170

Mucus-corpuscles, 219

Mucus, 176, 218 Mucus-casts, 235

```
Nephritis, interstitial, 247
Nitric acid test, for albumin, 39
Octohedra of oxalate of lime, 205
Odor of urine, 32
Oil in urine, 22, 152
Organic constituents of urine, 35
Oxalate of lime, 204
chemical characters of, 206
clinical significance of, 209
```

Paraglobulin, 58 Penicillium glaucum, 241

deposits of, 204

recognition of, 205 sources of, in the urine, 208

Penicillium glaucum, 241
Peptone, 62
biuret, test for, 64
phosphor-tungstate test for, 64
Ralfe's test for, 65
Randolph's test for, 66
Peptonuria, clinical significance of, 66
Phosphate of lime, deposits of, 211

Phosphates, 179 their recognition, 213

alkaline, approximate estimation of, 182
clinical significance of, 183
nitrate of silver test for, 182
ammonio-magnesian, deposits of, 210
earthy, 209
as urinary sediments, 209
chemical characters of, 213
clinical significance of, 181

detection and approximate estimation, 180 of lime as a urinary sediment, 211

recognition of, 213

Phosphatic diabetes, 183

Phosphoric acid, volumetric process for, 184
Picric acid test for albumin, 45
Pigmented markings on glass slides, 195
Polarizing saccharimeters, 99
Propertone, 67
Proteids found in urine, 58
Contrasted, 69
Pus, changes in urine containing, 222
character of urine containing, 222
-corpuscles, action of reagents, 220
Pus, Donne's test for, 222
sources of, in urine, 223

Quantity of urine, 30

Reaction of urine, 23
Reagents required for urine examination, 15
Recording an examination, 257
Remarks on testing for sugar, 104
Renal diseases, differential diagnosis of, 244

Sarcina, 241
Sediments, organized, 218
unorganized, 196
Selecting a specimen of urine, 18
Seminal tubules, casts of, 236
Serum-globulin, 58
Solids in urine, 33
Specific gravity of urine, 25

to determine, 27 for very small quantities of fluid, 30 Spermatozoids, 238 Squibb's urinometer, 29 Sugar fungus, 240 of milk, 109

18

variations in, 26

569

Urates, 174 Tables, 259 Sulphuric acid, volumetric process for, 186 Sulphates, clinical significance of, 186 Tyrosin, 151, 216 Tube-casts, 229 Sugars found in urine, 70. (See Glucose.) chemical characters of, 216 detection of, 216 as a urinary deposit, 216 detection and approximate estimation, 185 INDEX.

Urea, detection and estimation, 154 volumetric analysis for, by Fowler's process, 168 deposits of, 200 acid, test for, 204 their test and recognition, 201, 204

Uric acid, 170 by Liebig's method, 157 by the hypobromite process, 162

carbonate of silver test for, 171

compounds of, 200 deposits of, 196 their recognition as sediments, 201

detection by microscope, 170 tests for, 200 recognition of, 197

quantitative estimation of, 171 quantitative estimation by Haycraft's method, 171

murexid test for, 170

Urinary calculi, 251 deposits, 189 rationale of production of certain forms, 189 classification of, 195 organized, 218

unorganized, 196

271

```
general physical and chemical characters of, 19
                                                                                                                                                                                             its transparency, and deviations therefrom, 19
                                                                               consistence of, and deviations therefrom, 22
                                                                                                                                                                                                                                                                                                                                                                                             selecting specimen of, for examination, 18
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     clinical significance of, 138
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  to prepare for examination for casts, 236
                          color of, and deviations therefrom, 22
                                                                                                              extraneous substances found in, 194
                                                                                                                                                                                                                                                                                                                                                                                                                                                         to determine specific gravity of, 27
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       to determine solid matters of, 33
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        detection of, 137
                                                                                                                                                                                                                                                                                                                quantity of, and variations, 30
                                                                                                                                                                     inorganic constituents of, 177
Urine, acid fermentation of, 24, 190
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             Urohamatin, Harley's test for, 124
                                                                                                                                                                                                                                                         order of examination of, 35
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             Uroxanthin, Heller's test for, 126
                                                                                                                                                                                                                                                                                   organic constituents of, 36
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 Uroglaucine, or indigo blue, 129
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                Heller's test for, 123
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  Urochrome of Thudichum, 125
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                Jaffe's test for, 127
                                                           coloring matters of, 121
                                                                                                                                                                                                                                                                                                                                                                                                                                       specific gravity of, 25
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   Uroerythrin in urine, 136
                                                                                                                                                                                                                                                                                                                                                 reaction of, 23
                                                                                                                                                                                                                                                                                                                                                                              secretion of, 13
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          test for, 123
                                                                                                                                                                                                                                odor of, 32
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              Urophain, 122
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  Urobilin, 122
```

Urrhodin, or indigo red, 126

