The Schott methods of the treatment of chronic diseases of the heart : with an account of the Nauheim baths, and of the therapeutic exercises / by W. Bezly Thorne.

Contributors

Thorne, W. Bezly -1917. Harvey Cushing/John Hay Whitney Medical Library

Publication/Creation

London : J. & A. Churchill, 1895.

Persistent URL

https://wellcomecollection.org/works/efwknd36

License and attribution

This material has been provided by This material has been provided by the Harvey Cushing/John Hay Whitney Medical Library at Yale University, through the Medical Heritage Library. The original may be consulted at the Harvey Cushing/John Hay Whitney Medical Library at Yale University. where the originals may be consulted.

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org



THE SCHOTT METHODS OF THE TREATMENT OF CHRONIC DISEASES OF THE HEART

W. BEZLY THORNE, M.D.

and the state of the second state of the secon







THE

SCHOTT METHODS

OF THE

TREATMENT

OF

CHRONIC DISEASES OF THE HEART

WITH

AN ACCOUNT OF THE NAUHEIM BATHS, AND OF THE THERAPEUTIC EXERCISES

ILLUSTRATED

BY

W. BEZLY THORNE, M.D., M.R.C.P.



LONDON J. & A. CHURCHILL 11, NEW BURLINGTON STREET 1895

TALE MEDICA DEC 1958 LIBRARY RM822 N3 895T

PREFACE.

IN March, 1891, I was enabled, by the courtesy of the Editors of the Lancet, to lay before the medical profession a paper by Dr. Theodor Schott, in which were explained the principles and practice of the treatment of chronic diseases of the heart by means of mineral baths and exercises, which had been elaborated by him and his deceased brother. As judged by results, it attracted no notice and the system remained an unknown art in this country. In the early part of the year 1894 I was favoured with a similar opportunity of bringing forward a brief account of my own experience of the Schott system. Since that time I have received so many requests for further and more detailed information, that I am encouraged to meet an increasing demand by the publication of the following pages. They do not pretend to offer a complete or exhaustive exposition either of the science and art of the physical treatment of heart affections, or of the range of its application. That their scope is mainly limited by my own knowledge and observation, is my apology for defects which are only too manifest to myself.

W. BEZLY THORNE.

53, UPPER BROOK STREET, LONDON, W. March, 1895.



CONTENTS.

снар. І.	NAUHEIM AND ITS WATERS	page 7
II.	BATHS .	13
III.	THERAPEUTIC MOVEMENTS	23
IV.	SPECIAL CONDITIONS TO WHICH THE SCHOTT SYSTEM IS APPLICABLE	33
v.	THE EXERCISES, DETAILED DESCRIPTION AND ILLUSTRA-	
	TIONS	41
VI.	ILLUSTRATIVE CASES WITH DIAGRAMS	66



THE SCHOTT METHODS

OF THE TREATMENT OF

CHRONIC DISEASES OF THE HEART.

CHAPTER I.

BAD-NAUHEIM AND ITS WATERS.

SITUATED at the north-eastern extremity of the Taunus range, Nauheim—or, to give it its full name, Bad-Nauheim, the birthplace and headquarters of the treatment about to be considered—lies mainly on the gentle slope, which, looking south south-east, forms the foot of the Johannesberg.

The underground streams which have been brought into requisition for therapeutic drinking and bathing, have been tapped in the lowest part of the township namely, at some little distance on either side of the stream which divides the park into two unequal portions; and it may be said at once that they rise from so great a depth as to preclude the possibility of subterranean communication with that small river.

The following analyses have been compiled from the observations recorded by Beneke, Prof. Will, of Giessen, Doctors August and Theodor Schott, and Dr. Uloth :—

S	
22	
H	
[T]	
-	
-	
4	
K	
WATERS	
-	
-	
NAUHEIM	
-	
-	
-	
H	
-	
5	
-	
-4	
-	
A	
F7]	
E	
E	
HE	
THE	
THE	
THE	
OF THE	
OF	

Quantitative analyses of the Nauheim Waters, by Prof. WILL, Giessen, including the results of analyses by Drs. AUGUST and THEODOR SCHOTT, made especially to ascertain the several proportions of carbonic acid gas. The amounts of solids are given in grammes as contained in 1000 grammes of water.

CONSTITUENTS.	SPRING No. 12. Friedrich-Wilhelms- Quelle.	SPRING No. 7. Grosser Sprudel.	SPRING No. 11. Gas-Quelle.	KURBRUNNEN.	CARLS- BRUNNEN (Dr. Uloth).
Chloride of Sodium	29-2940	21-8245	17-1388	15.4215	9.8600
Lithium	0-0536	0.0492	0-0323	0-0267	Trace
Potassium (Cæsium, Rubidium).	1.1194	0.4974	0.7174	0.5270	0.0726
, Ammonium	0.0712	0.0550	0.0433	0-0371	0.0113
" " Calcium	3.3249	1.7000	1-2598	1.0349	1.0575
" " Magnesium	0-5255	0.4402	0.3682	0.7387	0.2040
Bromide of "	0.0083	0.0060	0.0046	0.0063	0.0014
Iodide " " " · · · ·	Trace	1	i	1	1
Sulphate of Calcium	0.0352	0.0347	0.0190	0.0238	0-2277
", ", Strontium (with Baryta) .	0.0499	0.0390	0.0403	0.0324	0.0087
Bicarbonate of Calcium .	2.6012	2.3541	2.1473	1.1461	0.9515
" " Iron .	0.0484	0.0383	0.0313	0-0262	0.0147
", ", Manganese .	0·0069	0.0065	0.0050	0.0080	Trace
	0.0089	0.0104	9200-0	0200-0	Trace
Silicic Acid	0.0213	0.0325	0.0190	0.0186	0-0087
Arseniate of Iron	0.0002	0.00036	0.0005	0.00016	Trace
Phosphate ,, ,,	2000-0	0.00046	Uncertain	0.00034	0.0002
Organic Substances	Trace	Trace	Trace	Trace	Trace
Amount of Solid Constituents .	35.3573	26.3539	21.1663	18-6935	12.4183
Specific Gravity	1.02757	1.02088	1.01685	1.01475	1.0089
Free Carbonic Acid Gas .	•1.0074	+3-1756	1.4136	1.9622	1.4214
Temperature (Fahrenheit), Celsius	(95.54) 35.3	(88.83) 31.6	(81.68) 27.6	(70.52) 21.4	(59) 15
Outflow in 24 hours (Normal	1725	782			
in Cubic Meters Valves half closed	782	529			
Depth of Well in Meters	180	159.5	28	16	9

+ = 1340.46 C.cm.

= 578'93 C.cm.

.

8

THE SCHOTT METHODS OF THE TREATMENT OF

OF THE LUDWIGS	
ANALYSIS	SPRING.
UANTITATIVE	

G

By Prof. WILL.

Results given in grammes in 1000 grammes of water.

um	Bicarbonate of Calcium	0.369	600.0	1	0.113	0.172	0.028	I	0-341	1	0.001	1	. 0.012	Trace	240.1	ure of	1-254	1.0010
um	ate of Calcium Iron									2 2 ···						under pressu		of 66.92° F.
	ate of Calci " Iron " Mang " Maga " Sodium of Calcium " Sodium " Calcium " Calcium " Calcium " Lithium of Sodium of Sodium " Acid Gas travity at t	mn		ganese	nesium	im .	the second	. m							······································	contained 1		emperature

QUANTITATIVE ANALYSIS OF THE SCHWALHEIM SPRING.

By Prof. von Liebig.

Results given in grains in one pound of water = Grammes in 1000 grammes of water.

Chloride of Sodium	11.9465 gr. = 1.7060	09
Sulphate " " · ·	0.6215 , = 0.0900	8
Chloride of Magnesium .	1.0826 , = 0.1546	46
Carbonate of "	0.4185 , = 0.0600	8
", " Caleium .	43140 , = 0.6160	60
" " Iron .	0.0878 , = 0.0125	25
Silica .	0.1489 , = 0.0212	12
Total amount of Solid Constituents	18.6188 , = 2.6603	03
Amount of Carbonic Acid Gas held		
in solution at the pressure of		
one atmosphere	22.7258 , = 3.2465	65
Specific Gravity	1.0022	
Temperature .	51.8° F. 11.0° C.	

CHRONIC DISEASES OF THE HEART.

9

The springs which are used for bathing purposes are No. 12, No. 7, and No. 11; those employed for drinking purposes, the Kurbrunnen and the Carlsbrunnen, the Ludwigsbrunnen and the Schwalheimerbrunnen—mainly the former two. It will be observed that the bathing waters are endowed by nature with temperatures which suit them admirably to the purpose. As a matter of fact, it is only in exceptional cases that the waters have to be either artificially heated or cooled by ice.

A course of baths generally commences with the waters of the great Sprudel (thermal bath), freed from more or less of their natural gas, but, in any case, to such an extent as to induce a deposit of peroxide of iron and calcium carbonate, which, floating in the water, produces an opaque yellow coloration. To these, after a time, in increasing portions, are added one, two, three or even more litres of Mutterlauge-the uncrystallisable mother-liquor or waste product of the neighbouring works which provide large quantities of salt for the table. It is rich in chloride of calcium, iodine and bromine. The smallest quantity, carried to the tongue with the tip of the finger, produces an intense burning suggestive of vesication. Next in order comes the Sprudel bath drawn from No. 7 or No. 12, according to the temperature desired, containing a residue of natural gas sufficient to retain the whole of the iron in solution, and to coat the body with unbroken relays of globules which, on the bather emerging from the water, are found to have produced, insensibly, a wellmarked rubefacience and an agreeable glow of warmth. Then, finally, come the flowing Sprudel baths, probably the most powerful therapeutic baths known, in which the waters of either No. 7 or No. 12 forcibly enter and, through overflow pipes, leave the receptable during the whole period of immersion. These, with their constantly rising and simmering globules, emerging from moving water of crystalline clearness, convey the impression of a bath of champagne, and induce a sense of exhilaration not unlike that which is associated with that favourite beverage. No patient, in any case, is allowed to take more than two, or three, or at the outside four, successive baths in as many days, a day of interval always being imposed. Where much infiltration or osteoid deposit has taken place, carefully regulated massage is made to succeed each bath.

Speaking generally, the effects of the baths are :-to lower the frequency and increase the force of the pulse, and to induce a sense of refreshment and invigoration which is shortly followed by a no means disagreeable inclination to avail oneself of the hour's rest, in the recumbent position, which is enjoined as the invariable sequel. One of the more remote effects is to cause pain and even swelling of joints, and sometimes of nerve-sheaths, which have been previously affected by the gouty, rheumatic, or so-called rheumatoid processes. Such a condition generally endures for a few days only, but not only may it last longer, but it may be re-induced by each of the succeeding increments of balneological strength above mentioned. On the more permanent influences exercised in the circulatory and respiratory systems I shall enlarge in detail later on.

The range of morbid conditions which may be relieved by the internal administration and outward use of the Nauheim waters is very wide. They may be divided into those articular and numerous other changes which are dependent on the prolonged presence in the blood-stream of uric acid in excess, chronic affections of the heart and blood-vessels, with

12 THE SCHOTT METHODS OF TREATMENT.

one notable exception; congestion of the abdominal and pelvic viscera; and the earlier stages of chronic affections, of the congestive or sub-inflammatory order, of the spinal nerve structures.

I propose, however, in this notice, to confine my observations to the systematic use of the saline baths and of regulated movements of the body in chronic affections of the heart, according to methods elaborated after years of careful study by Dr. Theodor Schott and his deceased brother, Dr. August Schott.

CHAPTER II.

BATHS.

It has already been stated that the immediate effect of immersion in the Nauheim baths is to reduce the frequency and increase the force of the pulse. For example, at a time when my own pulse averaged 74 beats per minute in the recumbent position, and 84 in the sitting, the heart and vessels being sound, I found it, on four separate occasions, to have fallen, within two minutes of immersion in a Sprudel bath, to from 60 to 64. In ten minutes it had risen to from 66 to 68, and there remained during the period of immersion which in no case exceeded fifteen minutes. The exertion of dressing raised it to from 76 to 78; but, after the prescribed recumbent position had been assumed, it returned to from 62 to 66, with increased volume, and so remained during the period of repose. It will, therefore, be observed that the influence of the bath was not limited to the period of immersion.

These observations were made in August, 1893, and were verified a year later in the course of a series of five-and-twenty baths. Dr. John Broadbent was present during the twelfth, and having traced

the area of cardiac dulness before and after immersion, certified to a recession, averaging one third of an inch, in the general outline traced from the sternal to the mammary region.

By way of contrast, the following case may be quoted. A patient, aged forty-six, whose health had been declining for several years, was found to have a pulse of 80 in the recumbent, and of 88 in the sitting, position. While he stood it varied from 100 to 104; and if he walked ten paces it rose to from 120 to 130. The apex was found to beat an inch outside the nipple line. Within two minutes of immersion in his first thermal bath (spring No. 7, divested of the greater part of its carbonic acid gas, temp. 90°.5 F.) the pulse had fallen to 70, and, judged by the finger, appeared to have doubled in volume; at the end of four minutes it was 68; in six minutes 66; in eight minutes 68; and while standing, after dressing, it was 90. Before he left the bath, after an immersion of ten minutes, the apex beat was found to have receded half an inch in the direction of the mesial line; and nails and fingers, which had been snow-white up to the junction of the second with the first phalanx, had assumed a healthy flesh tint.

The immediate effect of the first few baths is to produce a sense of oppression at the præcordia, under the influence of which the patient breathes slowly and deeply for two or three minutes. Respiration then becomes easy and continues slower by from two to four breaths a minute.

The effect on the peripheral vessels is to increase their carrying power. A glowing sense of warmth is experienced in the extremities and in the surface of the body generally. The veins are stimulated to a similar activity. In fact, the general vascular capacity, systemic and pulmonary, is increased, and, without loss of blood, the relief of a general bleeding is afforded to an overloaded and labouring heart.

Such being the results of a carefully graduated and regulated series of immersions in these saline waters, it can scarcely be matter for surprise that in three or four days, especially in cases in which the flow of urine has been scanty, there ensues a free diuresis which may continue for days or weeks; that metabolic change becomes accelerated and improved ; that deeply-seated organs, more especially the liver and pelvic viscera, are relieved of congestion and partake in the general impulse to functional health; and that the heart, relieved of its burden, and contracting fully and without hurry on its contents, derives from an improved coronary circulation materials for the repair of its weakened or damaged tissues. It is suggested by Dr. Schott that these effects are produced partly by the cutaneous excitation induced by the mineral and gaseous constituents of the waters, and partly by a more prolonged stimulation of the nerves of sensation excited by imbibition into the superficial layer of the corium. According to this hypothesis, each sensitive nerve branch distributed over the surface that has been immersed, transmits to its parent centre an influence which is centrifugally reflected to the vasomotor system and to the ganglia which control the action of the heart. That the nerve centres are brought under powerful influence is attested by the remarkable trophic changes which may be observed to follow a course of these baths, unaided by the internal use of mineral waters or pharmaceutical remedies, in cases of anæmia, wasting, neurasthenia, and, above all, in cases of osteoarthritis.

15

The rehabilitation of the trophic, and probably of other central nerve tissues, is so lasting that progressive improvement may be observed for three or four months after the completion of the course. It need hardly be pointed out that such a process of general health restoration is a factor of scarcely secondary importance in cases in which the condition of the heart claims the attention of the physician as the main indication for treatment.

It is necessary here to state that it is not claimed that these waters are unique in their therapeutic influences; on the contrary, from the earliest days, Dr. Schott and his brother have insisted that similar, if not indeed identical, effects may be derived from baths artificially prepared so as to resemble the Nauheim waters in their principal mineral ingredients. Thus, in severe cases, it is recommended that the treatment should commence with a 1 per cent. solution of chloride of sodium, and that the strength should be gradually raised to 2 or 3 per For increasing the cutaneous excitation, cent. chloride of calcium is the salt to be relied on. The initial strength of the bath with regard to that ingredient should be 0.2 per cent., approximatively that of No. 7 spring, and, by increasing additions, it may be raised to 0.3 per cent., that is to about the strength of No. 12 spring, and eventually to 0.5 per cent. Such varying degrees of concentration may be obtained by the proportional use of the crystallised Nauheim bath salt, of the mother-lye, and of Mediterranean sea residue which also contains chloride of sodium and of calcium in the proportions necessary for the preparation of initial baths, together with traces of bromides and iodides.

For the production of carbonic acid effervescence

the action of hydrochloric acid on bicarbonate of soda is relied on. As a state of chemical purity is not required by the circumstances of the case, the articles of commerce are sufficient for the purpose. Two ways of employing the reagents are suggested, the one calculated to induce slow and gradual, the other rapid and almost immediate, effervescence. In the case of the first, the various salts, including the requisite proportion of bicarbonate of soda, having been dissolved, a bottle containing the acid is laid at the bottom of the bath, and the stopper having been withdrawn it is moved about from time to time. The bath will be ready for use in two or three hours. For the more rapid production of effervescence, the stopper of the bottle containing the acid is loosened but retained in position, the bottle having then been inverted and lowered until its mouth is just below the surface of the water, the stopper is withdrawn, and the bottle is moved about so as to diffuse a layer of acid as uniformly as possible over the surface of the bath. By this means the bath will be prepared in about five minutes. It will be useful to employ baths of three degrees of effervescence:

Mild		$\dots \frac{1}{2}$ lb. NaHCO ₃ to $\frac{3}{4}$ lb. HCl (25 per cent.)
Medium		1 lb. NaHCO ₃ to $1\frac{1}{2}$ lb. HCl.
Strong (Sprudel streng	gth)	2 lbs. NaHCO ₃ to 3 lbs. HCl.

Except in the case of porcelain baths it is desirable to ensure a slight excess of alkali in order to prevent corrosion.

17

As evidence of the fact that the virtues which are claimed for the Nauheim waters are not peculiar to them, it may be interesting to record the result of some observations recently made on the effects of baths consisting of the waters of Llangammarch Wells in Breconshire, the mineral constituents of which, according to an analysis made by Dr. Dupré in 1883, are as follows :—

		Parts per 1000.
Chloride of Sodium	189.56	2.708
Chloride of Calcium	84.56	1.208
Chloride of Magnesium	24.31	0.350
Chloride of Barium	6.26	0.090
Carbonate of Calcium	2.80	0.040
Silica	1.40	0.002

308.89 grains per gallon.

Iron Protoxide a trace. Potassium Strontium Lithium Bromine

see ee Bruine per Builor

Mr. S. A. Vasey, F.I.C., F.C.S., kindly permitted the first experiment to be made on himself, and the observations were made by Mr. Hugh Bennett, M.R.C.S., L.S.A., of Builth, and myself. The pulse, before immersion, was 100. From the second to the eighth minute the same oppression of the breathing which ensues on immersion in a Nauheim bath was experienced. At the eighth minute the pulse was found to be 78; after an immersion of ten minutes it was 76; after dressing 86, and an hour later 82. The change which was effected in the area of cardiac dulness is shown in the accompanying diagram (I). The temperature of the bath was 92° F. **NOTE**.—Areas of cardiac dulness and apex beats indicated by red lines and crosses, respectively, refer to observations made after either baths or exercises.

DIAGRAM I.



A A, area of cardiac dulness before immersion in bath of Llangammarch water. Temp. 92° F.

 A^1 A^1 , the same after ten minutes' immersion.

 $A \times$ and $A^{\dagger} \times$, positions of apex beat at corresponding stages.

R.N. and L.N., right and left nipples.

S S, mid-sternal line.





DIAGRAM II.



A A and A¹ A¹, before and after eight minutes' immersion in a Llangammarch-Nauheim bath.

 $A \times$ and $A \times$, positions of the apex beat at corresponding stages. R.N. and L.N., right and left nipples. S S, mid-sternal line.

RN

CHRONIC DISEASES OF THE HEART.

The next observation was made on myself by Mr. Bennett. Before undressing the pulse was 96.

0	n imr	nersion	it fe	ell at o	once to		80
A	t the	end of	the	1st m	ninute it	was	68
	,,	,,	.,	2nd	•,	,,	69
	,,	,,	,,	3rd	••	,,	72
	,,	,,	••	4th	.,	,,	73
	,,	,,	,,	5th	,,	,,	74
	,, ,	,,	.,	6th	,,	,,	70
	,,	,,	,,	7th	,,	,,	68
	,.	.,		Sth	,,	,,	72
	,, .	,,	.,	9th	,,	,,	76
	,,	,,	•,	10th	,,	,,	77

The recession of the area of cardiac dulness at the sternal end of the arc was $\frac{1}{2}$ inch, and at the apex $\frac{3}{8}$ inch. The temperature of the bath was 89° F.

A third observation was made on the following day by myself on a woman aged forty-seven, the subject of aortic stenosis, albuminuria, partial ascites, and œdema of the lower extremities. In this instance the strength of the bath, in chloride of sodium, was increased to that of the Nauheim spring No. 7. Before an immersion of ten minutes the pulse was 92, and after it 88; when the patient was partly dressed it was 84. The temperature of the bath was 92° F. The accompanying diagram (II.) shows the diminution which was effected in the area of cardiac dulness.

In each of the above three cases an increase in the volume of the pulse similar to that effected by the Nauheim baths was found to take place, but the after-glow which followed immersion in the bath, which had been strengthened by the addition of

19

chloride of sodium, appeared to be greater than in the other two instances. These observations tend to prove the correctness of Dr. Schott's disinterested contention that the Nauheim baths enjoy no monopoly of heart-therapy.

The following sphygmographic tracings and notes of pulse pressure, as recorded by the sphygmomanometer, are borrowed from a paper of Dr. Theodor Schott's, which I was the means of laying before the medical profession in Britain in 1891,* and afford evidences, which might be indefinitely multiplied, of the invigorating influence which the baths exercise on the heart and the circulation.

Tracings taken from a patient aged thirty-one, suffering from cardiac weakness.



FIG. 1.

Before bathing: Frequency of the pulse, 94; pressure of the pulse, 120 millimetres of mercury.





After the use of a Nauheim effervescent mineral bath of 87° F., duration fifteen minutes: Frequency of the pulse, 72; pressure of the pulse, 140 millimetres of mercury.

* Lancet, May 23rd and 30th, 1891.

Tracings taken from a patient aged forty-six, affected with stenosis ostii arteriosi sinistri. Exercises were superadded to baths on the ninth day of treatment.

man hant

FIG. 3.

Before the beginning of the treatment the pulse could not be counted (more than 150); pressure of the pulse, 82 millimetres of mercury.



FIG. 4.

After the first bath, containing 1 per cent. of salt, temperature 89.5° F., duration ten minutes.

mmmmmm

FIG. 5.

Eighth day of treatment by baths: Frequency of the pulse, 144; pressure of the pulse, 95 millimetres of mercury.



Ninth day of treatment, after half-an-hour's exercises with resistance: Pressure of the pulse, 110 millimetres of mercury.



FIG. 7.

Fourteenth day of treatment : Frequency of the pulse, 108; pressure of the pulse, 115 millimetres of mercury.

FIG. 8.

After three weeks' treatment : Frequency of the pulse, 108 ; pressure of the pulse, 125 millimetres of mercury.

CHAPTER III.

THERAPEUTIC MOVEMENTS.

THE treatment of cardiac affections, as practised by Dr. Schott, is not, however, limited by the therapeutic influences of the baths. As the result of a series of elaborate and prolonged experiments carried out by him and his deceased brother, Dr. August Schott, a system of exercises has been devised which yields results, if anything, still more remarkable. Their effect is illustrated by the following records, which were made by myself in conjunction with Dr. Hermann, of Charkoff, in August, 1893. The patient was a stout, well-built, fresh-looking man, forty years He brought letters from Professor von of age. Jörgensen of Tübingen and from his brother, who is a medical man, both of which described him as having been addicted to alcoholic excesses and being the subject of cor adiposum. Before the exercises the heart sounds were barely audible through a binaural stethoscope, and the apex beat was inappreciable. There was some ædema of the lower extremities.



Fig. 9. Before exercises.



FIG. 10. After twenty minutes' exercises.



FIG. 11. After thirty-five minutes' exercises.



FIG. 12. After forty-five minutes' exercises.

The following diagram (Fig. 13) gives, on a reduced



FIG. 13.

A, left nipple; B, right nipple; C, fossa jugularis; D, upper level of hepatic dulness; E, ensiform cartilage; T D, total diminution of area of cardiac dulness; X, position of apex beat. scale, the tracings of the area of cardiac dulness which were taken before, and at the conclusion of, the same exercises.

These exercises have been denominated by their inventors "Widerstandsgymnastik," or gymnastic with They may perhaps be more conveniently resistance. termed resisted movements or exercises. They consist of movements calculated to bring into successive and regulated action almost every collective system of voluntary muscles which is comprised in the human frame. Each succeeding movement is resisted by an attendant to such an extent as to oppose without arresting it. They consist of slowly-conducted flexion and extension, adduction and abduction, and rotation, in orderly succession, of the arms, the trunk, and the lower extremities. Each movement must be slowly and evenly made with a definite and uniform effort on the part of the patient. The office of the attendant is not limited to resisting the movements with equally uniform, but slightly inferior, force, but it is his duty to impose a short interval between each movement, to enjoin slow and regular breathing, and, more especially, by observing the rate of breathing and its force as indicated by the alæ nasi, to ensure that no undue strain is placed on the heart and lungs. He is also charged to guard the patient from perspiration and the slightest approach to palpitation of the heart. Either of these indications must be the signal for an interval of repose, during which the part being exercised is either left to hang at rest, or is supported by the hand of the attendant, who, under no circumstances, is allowed to grasp or in any manner constrict any portion of the patient's body. As the course proceeds, the energy of the movements, and consequently the force of the resistance, are gradually increased. Before and after the séance, and, if necessary, at some

intermediate period the physician maps out the dimensions of the heart by percussion, to satisfy himself that the effects of the exercises are satisfactory. The results, in fact, are such as would scarcely be believed by any but an eye-witness. It is by no means uncommon, in cases of dilatation, to see, within one hour, the oblique long diameter of the heart's area of dulness diminish by from three-quarters of an inch to an inch and a quarter, and, perhaps more surprising still, to observe a diminution of as many as two inches, in vertical measurement, of a liver which at first extended to the umbilical level; and to hear the patient, at the conclusion of what cannot be described as an ordeal. volunteer the statement that a load has been removed from the præcordia, that he breathes easier and more deeply, and experiences a sense of general relief.

It is not suggested that the whole of such a gain is permanent, for in the time that intervenes before the next day's exercises or bath, as the case may be, the dilated and congested organs tend to their former dimensions, but, be it well observed, they do not wholly relapse. On the contrary, each contraction ensures a proportional permanent gain, until, at the end of a few weeks, the attenuated and dilated heart and the congested liver have recovered either their normal dimensions, or, at any rate, such contraction and compensatory power in the one case and resolution in the other, as constitute them practically sound.

The resources of Dr. Schott and his brother did not come to an end with the conception of this system of physical treatment. With a view to enabling physicians and patients to maintain treatment of the same kind without resorting to Nauheim, they devised a method in the practice of which the patient is instructed, himself, to supply the resistance for which he would otherwise be dependent on a second person; and, by the aid of these self-resisted movements he is able to carry on and, from time to time if need be, resume a therapeutic process of unquestionable value.

In the course of his memorable lectures on surgical pathology, Sir James Paget quoted the profound observation of Treviranus that "each single part of the body, in respect of its nutrition, stands to the whole body in the relation of an excreted substance." In view of what may be achieved by means of the Schott system of therapeutic exercises, we may go further and say that each part of the body, through its motor nerves, is capable of exercising a health-sustaining, and in some cases a healthrestoring, influence on the heart and circulatory system. It has been shown that, even in health, the heart may present, under the alternating influences of exercise and repose, very appreciable variations in size; and Stokes long ago insisted that exertion, under suitable conditions, may promote the health of a damaged heart. On the other hand, there are not wanting examples of hearts that have been morbidly dilated, and to that extent damaged, either temporarily or permanently, by strains disproportionate to their strength. Drs. August and Theodor Schott enjoy the distinction, and are entitled to the credit, of having brought the physiological relations of exercise, function, and repair, into obedience to a therapeutic system which yields results in the treatment of diseases of the heart hitherto unknown and unlooked for. Such service brings honour to their profession and deserves the gratitude of mankind.

What has been said of the influence of the baths applies equally to the therapeutic exercises, except that retardation of the pulse is not so rapidly effected, even though its force and volume be manifestly increased; and that, in the nature of things, it would

27
not be desirable to prolong immersion beyond fifteen or, even in cases of exceptional tolerance, twenty minutes; while, on the other hand, there are few patients who may not be kept under exercise for half an hour, and some can undergo an hour's treatment without fatigue. There are, therefore, in matters of secondary detail, differences between the baths and the exercises, and it rests with the physician to decide whether one should be brought into requisition to the exclusion of the other, or both be employed at suitable times and intervals.

As with the baths so with the exercises, therefore, the following immediate results may be looked for in the majority of patients afflicted with a damaged or weakened heart: retardation of the pulse and increase of its force; contraction of the heart, generally first on the right side (it is rare for the left ventricle not to share in the contraction); slower and deeper breathing with a sense of lightness and relief in the chest; a better colour of the lips, and improved facial aspect; and, where that organ is congested, a marked diminution in the dimensions of the liver. Systematic administration of the exercises is generally followed in a few days by marked, and often long maintained, diuresis. That the diminution in the area of dulness is not due to increased pulmonary inflation and consequent overlapping of the heart, is shown by the fact that the position of the diaphragm remains unchanged, or, if it moves, it does so in an upward direction. The accentuation and migration towards the mesial line of an apex beat, which is appreciable to both sight and touch, afford evidence which, to most minds, is convincing.

In the course of the first few movements a bruit, due to stenosis, may be observed to become accentuated; before the series has been completed murmurs, resulting from valvular insufficiency other than that caused by actual lesion, may be diminished, then modified to duplication, and finally obliterated; heart sounds which were barely to be heard may become audible; and an apex beat that could not, under any circumstances, be detected is readily appreciable to the touch.

The increase of the general vascular capacity is not less striking in the case of the exercises than the Within a few minutes the size of radial baths. artery, as gauged by the touch, may seem to have doubled, and, before the series of movements has been completed, cheeks and fingers that were cold and either white or bluish-red, glow with warmth and healthy colour. The motor nerves, called into orderly, regulated, and, above all, not exhausting activity, seem to exercise centripetal and reflex influences similar to those which are brought into action by the baths through the nerves of sensation. The increased capacity of the vessels, down to the smallest capillaries, enables the heart so to contract as to empty its cavities at each stroke; while, at the same time, the ganglia, which control its action, seem to enforce a tonic contraction, which, renewed and maintained from day to day, leads to the establishment of a better habit of both function and repair.

It may be well, now, to gather together in brief summary the effects of these simple but wonderworking movements, and I am favoured with permission to do so by quoting the following lines from the admirable and lucid article of Sir Philip C. Smyly, which recently appeared in the *Dublin Journal* of *Medical Science* (September, 1894):—

"Take for example, the four following phenomena:---

1. The colour before the movements is a purple-blue in the cheeks and hands and feet.

2. The forehead, neck, and ears, etc., are a waxy white.

3. The pulse is rapid and blood-pressure low in the arteries.

4. The area of dulness over the heart is large.

1. The colour after twenty minutes or so becomes red, and the blue gradually disappears from the hands and feet.

2. The forehead, neck, and ears, etc., become pink.

3. The pulse slows and becomes full, the bloodpressure rises.

4. The area of dulness diminishes at times as much as an inch or more in diameter.

These results are due to :---

- 1. Increased arterial circulation, due to "the diminution of peripheral resistance." *
- 2. Diminished venous congestion, due to larger quantity of red blood in the arteries.
- 3. Diminished work for the heart, due to the free circulation of the blood in the arteries.

.... there will ever be a feeling against this treatment until it is clearly seen and believed to be true :—

- 1. That the movements relieve the back pressure on the heart.
- 2. That the diminution in the size of the heart is due to the absence of excess of blood in its cavity.
- 3. That this is attained by there being more room in the arteries.

* "Mechano Therapy," A. Symons. Eccles. Pract. Aug., 1894, p. 114.

- 4. That the heart muscle gains strength by having room to contract.
- 5. That the contraction being more complete, it takes a longer time, thus making the pulse slower, and, at the same time, fuller.
- 6. Being able to send on more blood it is ready to receive more, and thus removes venous congestion.
- 7. The strength gained by the heart is due to the freedom to contract fully."

Dr. Schott affirms that benefit may be expected to accrue in all cases of chronic heart disease whether of valvular or parietal incidence, except where the myocardium has reached an advanced state of degeneration, or the vessels are the seat of advanced arterio-sclerosis. I have myself been witness of improvement amounting to practical or actual cure in cases presenting the physical signs usually regarded as indicative of the following affections: stenosis of either the aortic or the mitral orifices; stenosis of both; insufficiency of either or both, with attendant dilatation; dilatation consequent on myocarditis, on habitual hæmorrhage and on constitutional anæmia; fatty heart (interstitial); weakened heart; congenital mitral insufficiency; patent foramen ovale; and angina pectoris of apparently both neurotic and organic causation. It is reasonable to assume that measures, calculated to diminish peripheral resistance, and to promote the nutrition and repair of the cardiovascular tissues, must be applicable to, at least, the early stages of aneurism of the heart and great vessels.

The diagnostic and prognostic value of the exercises must not be overlooked. Familiarity with the effects which they may be expected to produce

healthy and on weakened walls enables the on physician to detect early stages of dilatation, the existence of which it might otherwise be difficult or impossible to recognise. More pronounced dilatation may be readily differentiated from parietal hypertrophy, superincumbent fat, and pericardial effusion or infiltration. The measure of contraction induced by a few exercises readily discloses whether an abnormal area of dulness is to be attributed to dilatation or to a substantial mass of unvielding tissue. As regards prognosis, valuable information may be derived from the rate at which præcordial dulness is reduced, and, after a few days, by ascertaining before the bath or exercises, as the case may be, the amount of more than temporary contraction which has been secured. Lastly, an unsuspected valvular lesion may be betrayed by the development of a bruit while the movements are in progress.

A question of no secondary importance is: May such recovery of heart-power and efficiency, together with the improvement in the general health which is contemporaneously effected by the systems under consideration, be so enduring as to justify, in the greater number of instances, the return of the patient to the cares and labours of an active life? From my own observation, and from the testimony of other observers, I am able to reply that such is the case.

CHAPTER IV.

SPECIAL CONDITIONS TO WHICH THE SCHOTT SYSTEM IS APPLICABLE.

THE value of therapeutic measures which are capable of promptly relieving an over-burdened and labouring heart, without recourse to either general bleeding or drugs the use of which may sooner or later be attended with toxic, or, to say the least, undesired effects, is so obvious that it stands in need of no elucidation. Once it is recognised that the whole circulation, arterial, capillary, and venous, systemic and pulmonary, may thereby be stimulated to healthy activity and normal function, and that such healthrestoring effects may, under careful and judicious direction, be maintained, progressively increased, and eventually confirmed without risk of injury or drawback to the patient, it must be apparent that they may be applied with advantage to the relief of troubles which are not commonly considered to be exclusively cardiac in their origin and incidence. The principal typical affections of the heart to which they are applicable have already been enumerated, and, with reference to them, it need only be pointed out that, speaking generally, the baths, as compared with the exercises, offer special advantages for the treatment of those cases in which a rapid, feeble, and, perhaps, also irregular or intermittent pulse is the

3

expression of cardio-vascular degeneration, whether there be co-existing valvular lesions or no. But it should be pointed out that in some extreme cases in which either the cardiac condition itself, or some dependent or independently existing complication forbids the removal of the patient from his bed, the exercises, modified, if need be, to suit the special circumstances of the case, may prove to be of value in preparing the patient to undergo treatment by baths, more especially if they should not be easy of access. The exercises, on the other hand, may be at once brought into use, with or without the auxiliary influences of the baths, to meet either failure of compensation, or, cases in which degeneration has not led to a marked failure of physical power and general health. In severe cases, and whenever it is desired to produce a speedy effect, the combination of the two methods may be expected to yield results which could not be secured by means of either employed alone.

It now remains to enumerate some of the conditions, not primarily or ostensibly cardiac, to which the methods under consideration may be applied with advantage.

First in order, and perhaps most obviously, come those which are associated with the presence of uric acid in the blood in excess. Such are those in which the sequelæ of acute rheumatism, such as peri-, myo-, and endocardial lesions co-exist with injury to joints and tendons. What has been said of the resolvent, metabolic, and trophic effects of the baths points to them as remedial and restorative agents of the first order of efficacy, especially when their direct effect in contracting the heart and nourishing its tissues is borne in mind. Where myocardial changes have left a legacy of dilatation and feeble

CHRONIC DISEASES OF THE HEART.

action, and the structures, entering into the composition of the joints, have either escaped injury or enjoyed the advantages of effectual resolution, the exercises alone may meet the requirements of the case. The same indications apply to the subjects of acute and chronic gout. Cases of dilatation, weakened heart, or special lesion occurring in the course of osteo- or rheumatoid arthritis stand, however, on different ground. When the central and trophic influences which, according to Dr. W. M. Ord and others, play a prominent part in this affection, as well as the measure of neurasthenia and the profound cachexia which frequently arise in conjunction with the arthritic changes characteristic of the disease, are taken into consideration, it will be apparent that the treatment by baths is calculated to relieve at one and the same time such cardiac troubles as may have arisen, and the other local and general conditions to which allusion has been made.

The weakened heart which influenza so often leaves in its train, probably as the result of myocarditis, is readily amenable to the exercises alone. In some such cases murmurs may be detected, mostly basicsystolic. They are probably due to orificial irregularity or dilatation, for they are generally subdued by the first few movements. Sometimes they recur in the course of the earlier intervals; but I have not met with any, not previously existent, which have survived a full course of the movements (Cases D and E, Ch. VI.).

The effects of both baths and exercises, whether singly or conjointly employed, are very remarkable in anæmia associated with more or less dilatation, whether chlorotic, malarial (Case A, Ch. VI.), or arising from loss of blood or chronic intestinal catarrh. In the first two classes of cases it is

common to see the colour, digestion, spirits, energy, and general health of the patient undergo a notable improvement within three or four days, under the influence of the exercises alone-that is, without either baths or a single dose of arsenic or iron. A course of four or five weeks, combined with due precaution as to diet, exercise, and general hygiene, is usually sufficient to ensure a return to health. In the two other classes of cases, it need not be said that the physical treatment should be combined with measures calculated to arrest the waste which the system has suffered. The effects of both methods, especially when combined, have been no less satisfactory in such cases of œdema, anasarca, serous effusion, and albuminuria associated with deficient heart power as, under my observation, have submitted to the treatment. Case J. (Ch. VI.) is one in point.

The Schott methods bring great relief to such cases of asthma, associated with however little cardiac dilatation, as have come under my treatment. I may mention three typical cases: a, is a lady of middle age who, on taking a drive, or in any way coming near a horse, experienced the following train of symptoms :---intense injection of the ocular and palpebral conjunctivæ, nasal defluxion, hoarseness, and the breathing characteristic of spasmodic asthma. After a week of baths she was able to take a long drive with relative impunity, and, as the course proceeded, the improvement continued until finally the symptoms were scarcely appreciable. A slightly dilated heart had resumed its normal dimensions within the first week, and the pulse had become uniformly stronger and fuller.* β , is a lady, thirty-six years of age, who has been liable, with increasing frequency and severity, to accesses of eczema,

* The benefit has, in this case, been maintained for five months.

intestinal catarrh with abilious stools, bronchitis with profuse muco-purulent expectoration, and asthma with nocturnal exacerbations of great severity. The effects of driving were similar to those experienced by a. When she came under treatment she had not been able to lie down for a fortnight, and could secure only a few snatches of sleep with the aid of the fumes of a well-known anti-asthmatic powder. The lungs were emphysematous, and the apex beat was two inches outside the nipple line. During the day, any slight exertion induced cardiac dyspnœa and præcordial distress. Within three days of commencing a course of exercises, all the symptoms had so improved that she could lie down at night and obtain unbroken sleep for two or three hours at a time. At the end of a fortnight the anti-asthmatic inhalations were discontinued. On the conclusion of a course of five weeks (inclusive of the menstrual interval) she enjoyed good nights, could go up and down stairs without the breathing being affected, take long walks, and drive, with scarcely appreciable inconvenience, through the streets of London in hot, dry, and dusty weather. The apex beat had receded two inches and was in the nipple line.* γ , differs from the preceding in having passed the climacteric period by about two years, and in the emphysema being more pronounced. For seven years she had only obtained sleep by being pillowed up and inhaling the fumes of nitre-papers, and outdoor exercise had been limited to slow rambling walks in the garden of her country residence. The exercises alone were em-The nocturnal asthmatic exacerbation, ployed. from day to day, occurred a little later and lasted a shorter time. In the course of the second week, the nitre-papers were abandoned, and good nights were

* The benefit has been maintained for seven months.

enjoyed in the recumbent position. On the conclusion of a five weeks' course, the apex beat had receded from an inch without to half an inch within the nipple line. When last heard of, eight months later, the patient was taking a country walk of about a mile every day, and leading a fairly active life.

I have mentioned these cases with some detail because they seem to open up a prospect of relief to a class of sufferers whose troubles have hitherto, to a great extent, defied treatment. The results recorded are, however, not surprising when considered in the light of what has been shown to take place in the relief of a burdened heart, and the improvement of the capillary circulation. It can scarcely be doubted that the congested and varicose veinlets which encumber the alveoli in such cases, share in the general change for the better, and that the circulation through the pulmonary circuit is quickened, and the aëration of the blood proportionately facilitated, by the increasing systemic arterial and capillary capacity and activity. Be that as it may, it needs no argument to show that an asthmatic patient is in better case when strong and well-contracting heart-muscles propel the blood-stream through channels which offer a reduced, and perhaps no more than a normal resistance. In this connection I may state that convalescence from acute bronchitis and from pneumonia may be favourably influenced by recourse to the movements, more especially as regards the drying up of moist exudations.

A considerable proportion of the subjects of cardiac dilatation are also affected with frequently recurring distension of the stomach, or with chronic dilatation of that organ. These conditions generally subside *pari passu* with the improvement in the state of the heart without special treatment.

The changes in the general circulation, and more especially in the peripheral vessels and the capillaries, which lead to habitual coldness of the extremities with a deep bluish-red colour of the hands which gives place to a white hue on pressure, but returns the moment the pressure is relaxed, and not infrequently similarly affect the colour of the cheeks and of the tip of the nose, yield equally well to both baths and exercises where they have not become hopelessly confirmed. Two out of three cases which I have treated by the latter method have been apparently cured. In one of them an habitual headache, which, with occasional variations of intensity and a few complete intervals, had existed for about seven years, was effectually relieved. The patient is a lady twentytwo years of age. The one in whom the treatment produced only partial and temporary relief, and no permanent benefit, is over thirty years of age and, also, the subject of habitual headache with occasional accesses of acute hemicrania. A man seventy-four years of age who, for four months, had been troubled with habitual headache associated with the evacuation of uric acid crystals and accesses of lumbago, was relieved of the headache in four days. The exercises were persevered with for a month, and the apex beat which had been found an inch outside, receded to a point half an inch within, the nipple line.

Of six single women, under the age of twenty-five, who habitually suffered acutely during the first hours of the menstrual period, five reported that they were unconscious of discomfort during the initial stage of the first menstrual period which occurred after the exercises had been commenced. In the three cases of which I have been able to obtain subsequent information, the relief has been permanent. I have observed the same result to follow the baths.

THE SCHOTT METHODS OF TREATMENT.

40

This brief, notice of what may be called the secondary or indirect results of a treatment which is more especially directed to the heart, would be incomplete if I were not to allude to its psychological influence. No one can have observed the subjects of cardiac inefficiency, especially those who are affected by either simple dilatation, or by that condition associated with valvular lesion and failure of compensation, without being struck with the nervetension and mental suffering which they endure. Intolerance of sound, irritability, difficulty of mental concentration, lessened power of work, depression amounting, in some cases, to despondency, and night alarm, are of common occurrence. With the rehabilitation of the heart and vessels which these methods of physical treatment are so successful in inducing, all such nerve-suffering vanishes like a dream, and the spirits rise to a plane of hope and energy which is surprising alike to the patient and the physician (Cases C and D, Ch. VI.).

CHAPTER V.

THE EXERCISES.

"Movements without design weaken the heart; movements with design, on the contrary, strengthen the heart."—THEO. SCHOTT.

[For the illustrations contained in this chapter I am indebted to the joint labours of Mr. Macdonald Gregory, one of my assistants, and of Mr. Prendergast Parker, the artist.—W. B. T.]

In approaching the subject of the movements which have been shown to exercise therapeutic influences over the heart and blood-vessels, which place the drugs hitherto relied on completely in the shade and relegate them to the position of occasional auxiliaries, it cannot be too clearly stated that we have not to do with "gymnastics" in the sense in which that word is usually employed in the English language. They do, doubtless, in the end promote the development of the muscles generally, but that is not their primary object. It should be distinctly understood that they are designed to produce regulated movement with *little exertion and no fatique*.

The person who administers them, who may be called the "operator," should strictly observe and enforce the following rules :—

1. Each movement is to be performed slowly and evenly, that is, at an uniform rate.

2. No movement is to be repeated twice in succession in the same limb or group of muscles.

3. Each single or combined movement is to be followed by an interval of rest.

4. The movements are not to be allowed to accelerate the patient's breathing, and the operator must watch the face for the slightest indications of : (a) dilatation of the alæ nasi; (b) drawing of the corners of the mouth; (c) duskiness or pallor of the cheeks and lips; (d) yawning; (e) sweating; and (f) palpitation.

5. The appearance of either of the above signs of distress should be the signal for immediately interrupting the movement in process of execution, and for either supporting the limb which is being moved, or allowing it to subside into a state of rest.

6. The patient must be directed to breathe regularly and uninterruptedly, and should he find any difficulty in doing so, or for any reason show a tendency to hold his breath, he must be instructed to continue counting, in a whisper, during the progress of each movement.

7. No limb or portion of the body of the patient is to be so constricted as to compress the vessels and check the flow of blood.

The following are the movements :---

No. 1.—The arms are to be extended in front of the body on a level with the shoulder joints, the palms of the hands meeting in front of the chest (Fig. 16). The operator places his hands on the outer surface of the patient's wrists in such a manner that the ulnar side of the patient's wrist rests in the fork between his own thumb and forefinger. He places one foot in front of the other so that he may lean forward, without overbalancing himself, while the patient's arms are carried outwards until they are in line with each other, and with the transverse diameter of the chest. The operator then places his hands, with a similar disposition of the thumb and forefinger, on the palmar surfaces of the patient's wrist, and



FIG .16.

FIG. 17.

offers resistance while the arms and hands are being brought back to the position from which they started. (Fig. 17).

No. 2.—The arm and hand of one side at a time are extended in the depending position, with the palm of the hand directed forwards, and the operator, standing at the patient's side, places his open hand on the palmar surface of the patient's wrist, the

thumb only being on the dorsal surface (Fig. 18). The patient then flexes the forearm, without movement of the upper arm, until the fingers come into contact with the shoulder. The operator then places



FIG. 18.



FIG. 19.

the palmar surface of his own hand on the dorsal surface of the wrist, and maintains it there while the flexed arm is being extended to the position from which the movement commenced (Fig. 19).

No. 3.—The arms are extended vertically in the depending position, with the palms of the hands turned forwards. After they have been raised outwards until the thumbs meet over the head, they are brought back to the original position. The operator faces the patient, and resists the upward movement



FIG. 20.



FIG. 21.

on the radial side of the wrist (Fig. 20), and the downward movement on the ulnar side (Fig. 21).

No. 4.—The hands, with fingers flexed from the end of the first phalanx in such a manner that the second phalanges of the respective fingers of the two hands are in apposition with their fellows of the opposite side, are pressed together in front of the lower part of the abdomen. The thumbs are extended, and lie within

the three sides of a rectangle formed by the flexed forefingers, and touch each other at their tips (Fig. 22). The arms are then raised until the hands are on a level with the vertex of the head. Resistance is offered by placing the hands on the radial surface of the wrists. The movement is then reversed.



Before the return movement is performed the operator changes the position of his hands so as to receive the wrists in the fork between his thumb and forefinger, the palmar surface of his fingers being applied to the palmar surface of the patient's wrists (Fig. 23).

No. 5.—The extended arms are placed in the depending position, with the palms of the hands

resting against the thighs. They are then raised in parallel planes until vertically extended. The movement is then reversed. The operator faces the patient, and in order that he may maintain an uniform and effectual resistance, the relation of his hands to the patient's wrists must pass through the



FIG. 24.

following changes: In the first position the fork between his thumb and forefinger must be applied to the radial part of the wrist (Fig. 24). As the arms rise to an angle of 45° to the body, his fingers glide round the wrist until they are lightly folded round the radial surface of the wrists. Before the reverse movement commences he receives the

ulnar aspect of the wrist in the fork between his thumb and forefinger (Fig. 25). While the arms are descending his thumbs move outwards, and, at the same time, the fingers glide round the dorsal surface of the wrist in a direction opposite to that which his thumb is taking, in such a manner, and at

64 FIG. 25.

FIG. 26.

such a rate, that, when the patient's arms are on a level with the shoulders, the ulnar aspect of the wrist rests on a reversed fork formed by the radial aspect of operator's forefingers, and the thumb pushed out to a right angle with the somewhat flexed fingers (Fig. 26). As the hands descend towards the thigh the tips of the operator's fingers gradually glide round to the ulnar aspect of the wrist, so as to resist the downward and backward movement of the arms. This is the operator's *pons asinorum*, but it should be mastered.

No. 6.—The trunk is flexed forward, without the knees being bent, and then brought back to the erect



position. The operator stands at the patient's side with one hand over the upper third of the sternum, and the other supporting the mid-lumbar region. (Fig. 27). The reverse movement is resisted by placing one hand over the junction of the cervical and dorsal portions of the spine (Fig. 28).

4

49

No. 7.—The trunk is rotated, without movement of the feet, as far as it can be carried to one side, say to the right, then to the left, and lastly brought back to face forwards as at starting. The movements are resisted by one hand being placed in front of, and a little above, the advancing axilla, while the other is



FIG. 29.

placed over the receding shoulder (Fig. 29). The operator must, to a limited extent, move round the patient when the second stage of the rotation is being performed, and will be able to do so most evenly and securely by carrying one foot round behind the other, somewhat as is done in performing the skating "outside edge backwards," before shifting the position of the other.

No. 8.—The trunk is flexed laterally, first to one side, secondly completely over to the other, and thirdly brought back to the erect position. The



FIG. 30.

operator stands in front of the patient. When the movement is to the right, his left hand is pressed against the right side of the chest in the axilla, while the right firmly supports the opposite hip, and vice versâ (Fig. 30).

No. 9.—This movement is identical with No. 1, with the exception that while it is being executed the fists are kept firmly clenched.

No. 10. —The arms are flexed in succession as in movement No. 2, with this difference, that the



FIG. 31.

palmar surface is turned outwards and the fist is firmly clenched (Fig. 31).

No. 11.—The arm is extended in the depending position, the palm of the hand lying against the thigh, and then makes a complete revolution from the shoulder joint, forwards and upwards, until it is vertically raised alongside of the ear. Before it descends backwards, the palm of the hand should be turned outwards (Fig. 32). The operator stands at the patient's side with his fingers folded round the

FIG. 32.

radial side of the wrist. His other hand must be ready to receive the wrist when it reaches the vertical position, and to maintain the resistance until the arm has descended to the position from which it started. This movement is performed by one arm at a time.

No. 12.—The arms are extended vertically in the depending position, the palms of the hands resting against the thighs. They are then moved upwards and backwards in parallel planes as far as it is possible



FIG. 33.

to do so without bending the trunk forwards. The upward movement is resisted with the fork of the hand on the ulnar aspect of the wrist, the downward, by folding the fingers round the radial surface (Fig. 33).

CHRONIC DISEASES OF THE HEART.

No. 13.—The patient stands with one hand resting on a chair or table, while the thigh of the opposite side is flexed on the trunk to the extreme limit, and then extended until the feet are side by side. The



FIG. 34.

leg should hang downwards from the knee-joint. The upward movement is resisted by a hand placed immediately above the knee (Fig. 34). The return

55

may be resisted by a hand placed below the lower part of the thigh or under the sole of the foot. No. 14. —The patient, supporting himself with



FIG. 35.

one hand, as in the last movement, bends the whole extended lower extremities in succession, first forwards to the extreme limit of movement, then backwards to the same degree, and finally brings the one foot alongside of the other. The forward movements



FIG. 36.

are resisted in front of and above the ankle (Figs. 35, 36), the backward movements behind.

No. 15.—The patient, supported in front by a chair or table, stands on either foot in succession,



FIG. 37.

while the leg of the other side is flexed on the thigh. The upward movement is resisted by pressure on the heel (Fig. 37), the return movement above the instep.

No. 16.—The patient, resting one hand on a chair and standing on the foot of the same side, raises the extended lower extremities in succession, outwards from the hip joint, and then reverses the movement.



FIG. 38.

The operator resists by means of one hand placed above the ankle (Fig. 38).

No. 17.—The arms, extended horizontally outwards, are rotated from the shoulder-joint to the extreme limits, forwards and backwards. The move-

ments may be resisted by the operator grasping the ulnar edge of the metacarpal portion of the hand



FIG. 39.

(Fig. 39), or by closing his thumb and forefinger in a ring round the wrist.

No. 18.—The hands, in succession, are first extended, then flexed on the forearm to the extreme limits, and lastly brought into line with the arm.



FIG. 40.

The operator's one hand supports the wrist, while the other resists the movements at the metacarpophalangeal junction, first on the dorsal, secondly on the palmar, and thirdly again on the dorsal surface (Fig. 40).

CHRONIC DISEASES OF THE HEART.

No. 19.—The feet, in succession, are flexed and extended to the extreme limits, and then brought back to their natural position. The movements are resisted in the dorsal and plantar surfaces, at about



the level of the metatarso-phalangeal joints (Figs. 41, 42).

Such being the mechanical details of these wonderworking movements, it is not surprising that, on making acquaintance with them, the patient asks, with scarcely veiled scepticism, why a wash in the "waters of Jordan" should not be equally effectual. No question could be more apposite, no allusion more appropriate. Many days have not passed before results too manifest to be mistaken offer an unequivocal reply. At the same time, no greater mistake could be made than to assume that the mastery of mechanical details is a sufficient equipment for either the physician or the operator, for in no two cases is their expert and judicious application likely to be precisely similar.

In the first place it should be understood that there

61

is no magic in the exact sequence which has been adopted in the foregoing description. Many patients are, at the beginning, unable to perform the full series without experiencing what is always to be avoided-namely, fatigue or distress as exhibited by one or more of the symptoms which have been enumerated. Others cannot with advantage submit at once to movements of some special parts, such as the trunk or lower extremities. Some who are confined to bed cannot, in the nature of things, execute a portion of the exercises. In all such respects it devolves on the medical adviser to instruct the operator. The time to be occupied by the several movements, the duration of the interval of rest, and the measure of resistance to be offered, are points on which his judgment should be expressed. For that reason he should always be present when the first exercises are administered, and, in many cases, it will be advisable for him to conduct a few movements himself, and then, having gauged the patient's powers and capacity, to administer them to the assistant in order that his estimate of the required rate of movement and degree of resistance may be placed beyond the possibility of misapprehension. It needs not to be said that the medical attendant should, in all cases, submit the patient to an exhaustive preliminary examination. More especially is this the case where there exists any impediment to the rapid filling of the expanding arteries and capillaries. Should, for example, the pulmonary circuit be obstructed, as it is in cases of emphysema and asthma, and with rigidity or stenosis of the aortic orifice, syncope may be easily induced. In presence of such conditions, the resistance should be limited to feather weight, the movements slowly executed, and the intervals prolonged to allow the heart and vessels time for the adjustment of their mutual relations to the changes which are being rapidly effected in the flow and distribution of the blood. It may even be desirable to enforce the recumbent position lest the pressure in the cerebral vessels be unduly lowered by the imperative requirements of the increasing vascular capacity. If the right side of the heart be overloaded, a down-grade should not, at first, be given to the brachial veins by raising the arms above the level of the shoulders. Briefly, the system under consideration brings such powerful influences to bear on the whole circulation, that, in application, it requires to be adapted to the exact condition of each individual. As with other potent remedies, the "rule of thumb" may easily convert a therapeutic agent into an instrument of mischief. No less care should be taken in the selection, instruction, and supervision of operators. They should be intelligent, light of hand, endowed with powers of observation, and trained to use them. The choice of women is not limited. Trained and even "registered" nurses who possess a knowledge of elementary anatomy and physiology, and whose faculties have been cultivated by hospital service, abound in our country; but suitable men are not easily found. Under no circumstances would it be justifiable to administer a course of either baths or exercises otherwise than under medical sanction and direction. With regard to treatment, although the physical method relegates pharmaceutical remedies to the rank of auxiliaries, their influence is, in some instances, of material value in correcting a special defect of health or in raising the general tone of the system. The patient's daily life often needs regulation, more especially in regard to exercise, fresh air, and the avoidance of undue fatigue, excitement, anxiety, mental distress, and all
64 THE SCHOTT METHODS OF THE TREATMENT OF

other depressing conditions. Diet, however, is a matter of scarcely secondary importance. The condition into which most patients have fallen, and the acceleration of tissue change which the bath and exercises alike induce, demand a liberal supply of muscle-forming nourishment, comprising, generally, animal food, though not of necessity butchers' meat, three times daily. If there be a class of subjects with regard to whom the adjustment of the dietary claims exceptional care, it is that numerous one in which a tendency to the excessive deposit of adipose tissue is the accompaniment of anæmia or of some other dyscrasia. With them, the substitution of animal food for a considerable proportion of the fats and carbohydrates in common use, is a measure of great importance. A judiciously devised "thinning," but not "lowering," diet lightens the corporeal burden, gives free play to the muscles, and strengthens the heart to an extent which can hardly be accounted for by the mere removal of superincumbent fat. It should be added that those who have had the widest experience of the Schott methods attach no importance whatever to special limitation of the quantity of fluids ingested, and that graduated mountain climbing, as recommended by Oertel, should only be resorted to towards the end of the treatment or after it has been brought to a satisfactory conclusion. It then forms the rational complement of the treat-Physical exercise, practised by means of ment. mechanical appliances, forms no part of the system, and introduces principles which are not only foreign to its conception but essentially opposed to it.

It now remains to be said that exercises with "selfimposed resistance" are often found to be of value as an after-treatment, more especially as they are within the competence of everyone who has become acquainted with the movements, and involve no risk of injury by over-exertion. "Selbst-hemmungs-gymnastik" or self-restraining gymnastics, were devised to enable patients to be, if one may so express it, their own operators. The restraint or resistance is effected by that hardening of the muscles of the limbs, or groups of muscles, which execute the movements, of which the condition of the forearm produced by firmly clenching the fist is an example. After a little practice the patient can induce that condition at will, and maintain it throughout the several movements, especially those of the arms and legs.

It would be unbecoming to close these observations without offering a tribute of admiration to the industry and genius which August and Theodor Schott have displayed in devising and elaborating means at once so simple and so effectual for the relief of a large measure of disablement and disease, and, at the same time, acknowledging the generous spirit in which they have, consistently with the most honourable professional ethics, made every effort to bestow the fruit of their labours on the medical profession at large for the benefit of suffering mankind.

.5

CHAPTER VI.

ILLUSTRATIVE CASES.

A., a lady, aged twenty-seven, had resided for four years in one of the semi-tropical States of America. Had suffered frequent accesses of tertian fever, which throughout the summer of 1893 had continued in unbroken series. Presented intense anæmia, dyspnæa on exertion, and sallow complexion. Suffered continuous headache and chronic intestinal catarrh, to which she had been liable for years. At the termination of the course the anæmia was completely relieved. The headache and dyspnæa were relieved by the end of the first week. The intestinal catarrh was pharmaceutically treated and relieved, but showed a tendency to return on slight provocation. Four months after the completion of the course the improvement was found to have been maintained. NOTE.—Areas of cardiac dulness and apex beats indicated by red lines and crosses, respectively, refer to observations made after either baths or exercises.

DIAGRAM A.



- A A and A¹ A¹, areas of cardiac dulness before and after first exercises (1 to 13).
- B B and B¹ B¹, areas of cardiac dulness before and after last series of exercises at end of third week.
- C C, area of cardiac dulness on termination of four weeks' course.
- R.N. and L.N., right and left nipples.

S S, mid-sternal line.

RN





DIAGRAM B.



A A, area of cardiac dulness before first exercises. $A^1 A^1$, the same after twenty minutes' exercises. B B, the same after completion of the course, thirty-one days later. C C, the same eighty days after completion of the course. $\times A$ and \times positions of apex beat at corresponding stages. R.N. and L.N., right and left nipples. S S, mid-sternal line.

RI

67

B., aged sixty-five, had rarely smoked less than twenty cigars a day during a thirty years' residence in India, and was found to be unduly stout, with pale and drawn face, and light bluish lips. He could not walk a hundred yards without stopping to recover his breath, nor ascend a flight of stairs without resting, supported on the banister, for the same purpose. He was dieted to reduce his weight and correct gastric fermentation, and treated by exercises. Before the completion of the course his aspect and expression had changed, and his face and lips became ruddy; he walked daily to and from his club, a distance of five miles in all, and there played billiards for two or three hours, and could run up stairs without becoming In weight he lost a pound a week for six breathless. Nine months after the conclusion of the weeks. treatment, he was fishing and shooting in Norway, and now smokes, on an average, six small cigars a day.

C., aged fifty, had presented symptoms of cardiac failure for at least twenty years, and had been liable for four years to accesses of partial syncope, associated with gastric distension and intense vertigo. The treatment by exercises commenced a few days after recovery from the last such attack. On the conclusion of the course, the patient summarised the change in her condition as follows: "Before the treatment my sensations frequently forced upon me the apprehension of impending death, my digestion was bad, and every exertion of mind and body seemed to be too tiring to be endured. Now I walk for at least an hour twice daily. I can eat and drink anything in reason, and I am a stranger to fatigue and depression of spirits." Four months after the completion of the treatment the improvement was more than maintained. The patient had crossed a mountain pass at an altitude of more than 7,000 feet, in a snowstorm, without the breathing being affected, or experiencing any inconvenience.

DIAGRAM C.



- A A and A¹ A¹, areas of dulness before and after exercises. (Pulse reduced from 84 to 76.)
- B B, area of dulness after twelfth exercises (the fourth, after an interval of fourteen days, necessitated by the menstrual period).
- C C, the same after twenty-fifth and last exercises.
- C×, situation of apex beat on that occasion (not having been appreciable when previous observations were recorded).
- R.N. and L.N., right and left nipples.
- S S, mid-sternal line.

R.N





DIAGRAM D.



- A A and A¹ A¹, areas of cardiac dulness before and after first exercises. (Pulse reduced from 108 to 104.)
- B B, area of cardiac dulness twenty hours after completion of the course. (Pulse 84, after going up and down two flights of stairs.)

 $\times \Lambda^{i}$, A \times , and B \times , positions of apex beat at corresponding stages. R.N. and L.N. right and left nipples.

S S, mid-sternal line.

R.I

D., aged eighteen, 5 ft. 10 in. in height, had a first access of influenza in 1890, and a second in 1892. The latter was followed by loss of ocular accommodation, as well as cardiac weakness and vertigo which became so severe, about three weeks after the termination of the febrile state, that he was obliged to take to his bed, and there lay with a basin at his side, because an incautious movement of the head, or even the automatic fixing of the eyes on a crack in the ceiling, except while wearing convex lenses, brought on an attack of retching. From that time forward he was debarred from participation in all games and sports, as any exertion beyond a leisurely walk brought on palpitation, præcordial pain, and dyspnœa. On the 28th of February, 1894, he commenced a course of exercises which extended to the 24th of March, inclusive, after which he returned to the country. A month later I received the following report :- In active pursuits he is now on a level with other young men of his age. His tutor reports that in power of application, and in memory, he is twice the man he was; but the most remarkable change is in his spirits, for, whereas the word "beastly" used to be freely scattered through his letters, everything in life is now said to be "awfully jolly."

E.—The following case is published by permission of Dr. James Harper, with whom I saw the patient six weeks after the termination of the acute stage of influenza. She was twenty years of age, and had not gained in strength or power of movement from the time of leaving her bed. She was found to be very anæmic, somewhat wasted, and could only move from one room to another adjoining, at the cost of dyspnœa and præcordial pain. A well-marked systolic bruit was audible at the base. I administered exercises very slowly, with gentle resistance, and long intervals, for the space of fifteen minutes in all. The areas of dulness were traced by Dr. Harper, who also recorded the following observations :—

Dec. 12, 1893	Before exercises,	P. 96, murmur distinctly audible.
and the second	After "	P. 84, " scarcely "
" 14, "	Before "	" fairly marked.
	After 20 m. "	" scarcely audible.
,, 18, ,,	Before "	P. 84-120 " audible.
	After 20 m. "	P. 84, ,, not audible.
,, 28, ,,		P. 84, ,, not audible. P. 92, no bruit.
Jan. 1, 1894	. Before "	D 199
	After 24 m. "	P. 84, "
" 4, "	,, ,, ,, ,,	P. 84, "
" 9, "	" 12-hour's "	P. 96, "
" 12, "	Before exercises,	P. 96, "

After the completion of the course the patient travelled by rail from Victoria station to South Kensington, whence she walked to my house. She presented the appearance, and enjoyed the sensations of perfect health. The pulse was 88, and I could discover no bruit.

DIAGRAM E.



- A A and A¹ A¹, areas of cardiac dulness before and after fifteen minutes' exercises, with gentle resistance.
- B B, the same at the conclusion of a course extending over thirtyone days.
- R.N. and L.N., right and left nipples.

S S, mid-sternal line.





DIAGRAM F.



A A, area of dulness before exercises.

A¹, right margin of area of dulness after twenty-five minutes' exercises, with slight resistance.

 A^2 , the same, after five more movements, with strong resistance.

 \times , \times^1 , \times^2 , positions of apex beat at corresponding stages.

 \times ³, apex beat as observed by the writer after twenty-five days' treatment by baths and exercises.

R.N. and L.N., right and left nipples.

S S, mid-sternal line.

F., aged fifty-three, was examined in the presence of Dr. Heinemann, of New York, and myself by Dr. Pawinski, of Warsaw, who kindly sanctions the present use of his observations and tracings. A wellmarked systolic bruit at the base, followed by a diastolic souffle in the same situation, were taken to indicate aortic stenosis with insufficiency. The patient suffered from dyspnœa, œdema of the lower extremities, and enlargement of the liver. When I examined him twenty-five days later, the systolic bruit was of diminished intensity, the diastolic souffle was no longer audible, there was no dyspnœa, and the œdema had passed away.

G., aged sixty-seven, had been known for four years to present symptoms of aortic stenosis, but had good compensation and led an active life. He came under observation again on March 8, 1894, some weeks after suffering from symptoms which suggested His face was drawn and an attack of influenza. anxious, and he complained of dyspnœa on exertion, and of great loss of mental and physical energy. On auscultation the basic-systolic bruit was found to have become louder, and to it was superadded a wellmarked apex-systolic murmur. The first exercises reduced the pulse from 60 to 50, and increased its force and volume. After seven days (B) he left London much improved, both murmurs being audible but reduced in intensity. Resumed the treatment after an interval of twenty-six days (C), enjoying at the time good general health and complete freedom from dyspnœa. The first exercises of this series reduced the pulse from 72 to 44. In ten more days the treatment was brought to a conclusion by the necessity of leaving London again. By that time the apex bruit had been superseded by a sound which, but for a slight lack of definition, was healthy. Eighty-five days later the area dulness was found to be as indicated by D. The basic bruit was reduced to its old intensity; the apex sound remained as when last observed. The pulse was 52. The general condition left nothing to be desired. The greater part of the members of this patient's family have, in health, a pulse of about 50.

DIAGRAM G.



A A and $A^1 A^1$, areas of dulness before and after first exercises.

- B B and B¹ B¹, areas of dulness on seventh day before and after exercises.
- C C and C¹ C¹, areas of dulness before and after exercises after an interval of twenty-six days.
- D D, area of dulness eighty-five days after conclusion of treatment.
- R.N. and L.N., right and left nipples.

S S, mid-sternal line.

- ×A, ×B, etc., positions of apex beat at stages corresponding to letters A, B, C, and D.
- $\times A^{i}$, $\times B^{i}$, etc., positions of apex beat at stages corresponding to letters A^{i} , B^{i} , and C^{i} .





DIAGRAM H.



December 7th. A A, before exercises.

- B B. after twenty minutes.
- 9th. C C, before exercises. ,,
- 12th. D D, before exercises, after being out three and ,, a-half hours and returning in a gale.
 - D D, after half-an-hour's exercises.
- 14th. E E, before exercises, and after six and a-half hours' ,, absence from home on professional duty.
 - 25th. F F, before exercises.

January 3rd. G G, three hours after exercises.

R.N. and L.N., right and left nipples.

S S, mid-sternal line.

,,

H., aged fifty-nine, who had for many years been the subject of heart troubles associated with a loud apex-systolic bruit, and exophthalmos, was found on December 3rd, 1893, to be suffering from acute pulmonary apoplexy. The pulse was bigeminous, two beats corresponding to each complete respiratory act. He was at once instructed to practise the arm exercises with self-imposed restraint, and ordered digitalis and strychnia. On the 7th, he was moving about the drawing rcom, and commenced a course of exercises resisted by a trained operator. On the 12th, he went to his office and was from home for three and a-half hours. He has since been three times under similar treatment for slight œdema of the lower extremities, but continues to lead an active professional life.

I., aged sixteen, is reported to have had carditis in the course of scarlet fever, at the age of five, followed by hæmato-albuminuria and ædema. Whooping cough, at the age of six, was followed by chorea of moderate intensity, which lasted for a year. A second attack, lasting four months, occurred at the age of seven. When first seen he was under treatment by Mr. Barwell for spinal curvature. He had a loud systolic-apex bruit, with a well-marked and diffused thrill, and epigastric pulsation perceptible to sight as well as touch. His parents had been advised to remove him from school, and not to allow him to leave the house otherwise than in an invalid chair or pony-chaise. After the exercises on the twenty-first day of treatment he trotted about a hundred and twenty yards and then walked fifty. Before doing so the pulse was 66, and the respirations were 20; afterwards they were respectively 86 and 19. He experienced no fatigue, and showed no signs of distress. At the conclusion of a course extending over twentyeight days he trotted two hundred yards. Before doing so the pulse was 66, and the respirations were 16; afterwards they were 80 and 18 respectively. He was, at that time, taking walks of one and two hours' duration without fatigue, and, generally, leading an active life, though debarred from running more than a few paces, and from joining in out-door games. The areas of dulness before, and at the conclusion of, the treatment were verified by Mr. Barwell, who also noted a much diminished apex impulse and complete absence of thrill and epigastric pulsation. The bruit had diminished, but was still well marked.

DIAGRAM I.



A A and $A^{1} A^{1}$, areas of cardiac dulness before and after first exercises. A \times and $A^{1} \times$, apex beats before and after.

BB, area of cardiac dulness on completion of course twenty-eight days later.

B×, apex beat.

R.N and L.N., right and left nipples.

S S, mid-sternal line.



J.—I am enabled, by the courtesy of Sir Philip C. Smyly, to quote *in extenso* the following case from his article already alluded to :—*

Miss X., æt. seventeen, had been ailing for some time.

Oct. 20th, 1893.—She had an attack of faintness and swelling of the feet and ankles.

Feb. 24th, 1894.—She came under my care. She was very low and weak; anæmic; hands and feet œdematous and very blue; general anasarca; ascites well marked, and fluid in both pleuræ. The area of the heart dulness was well defined to the right side of the sternum, but could not be outlined below or to the left side on account of the dulness from the pleural effusion. The skin was white and waxy on the forehead, ears, and neck. The cheeks were a dark purple-blue. After a very careful examination the diagnosis arrived at was-Dilated heart with patent foramen ovale (possibly); obstructed arterial circulation, with venous congestion of all the organs. No albumen in the urine. Began the resisted movements for twenty minutes every morning about 11 o'clock, and massage by an experienced masseuse every evening. The colour improved every day. area of dulness diminished, and the pulse became fuller and less frequent. The blue colour improved to a dark red.

March 8th.—In the daily report it was noted :— "She did some additional exercises with more strength. Her pulse was considerably stronger. Her nose bled slightly. She passed a cheerful day."

11th.—" Marked improvement in the pulse. Her colour keeps good all day; very little blue at any time."

* Dublin Journal of Medical Science, September, 1894.

75

17th.—Remarked the healthier appearance of the forehead, ears, and neck.

(From the 20th of March until the 11th of April the movements were omitted—unavoidably—though the massage was continued.)

21st.—" Not a good night—restless and dreaming. Appetite very poor."

April 5th.—"Her colour was very dark with much blue in the morning; got right in the afternoon. Urine very scanty."

6th.—"Swelling of the abdomen greatly increased. Only eleven ounces of urine in twenty-four hours."

11th.—Pain in the right side. Movements begun again, but very slightly, owing to great distress in breathing. Urine, eleven ounces.

12th.--Pain worse. Much swelling; could not lie down in bed. Urine, thirteen ounces. Daily exercises and steady improvement.

20th.—" Better. Urine, thirty-one ounces in twenty-four hours."

22nd.—"The heart rhythm was normal for the first time."

25th.—" The menses showed for the first time since October, and continued slightly for five or six days."

26th.—The patient was moved from one house to another. On being lifted into the carriage she became breathless and very blue. Towards evening she breathed better, but could not lie down. She got little or no sleep, and had to be supported sitting up all night.

28th.—The whole of the right pleura was full. Distress of the breathing rapidly increasing. Assisted by Dr. Cruise, I tapped the chest and drew off sixtytwo ounces of fluid, clear and yellow. She coughed up a quantity of thin mucus during the afternoon. Temperature 100°. At 10 o'clock, p.m., the temperature was 99°. She could lie down in bed with only two pillows.

May 4th.—Consultation with Dr. Cruise. She was very much improved. Movements were resumed.

11th.—" Was very cheerful all day. Colour a little high, but no blue."

21st.—Left Dublin, 7 p.m., for Holyhead. Next day to London, and on Wednesday had a consultation with Dr. Bezly Thorne.

26th.—Arrived at Nauheim better than when she left Dublin.

28th.—Had her first bath.

June 1st.—Pulse before the bath, 116; after, 110.

8th.—Dr. Schott showed the patient's mother "that the water had gone down a hand's breadth over her stomach."

20th.—Began the gymnastics. From standing during the examination, and the marking out the area of dulness, her pulse was 114; after the exercises it fell to 88, and then rose to 96, and then to 104.

21st.—I saw the patient at Nauheim with Dr. Th. Schott. I could not find any sign of fluid either in the chest or in the peritoneum. No œdema; and the heart's action normal.

July 6th.-Dr. Schott roports the heart very well.

29th.—Dr. Bezly Thorne saw the patient in London, and reports—"Wonderfully improved. Cannot detect any wrong sound in her heart."

30th.—Dr. Cruise and I very carefully examined the patient together, and found the heart's action normal, and no swelling anywhere. She returned home to all appearance quite well.

At an early part of this case it was mentioned that it appeared possible, from the extreme cyanosis, that

78 THE SCHOTT METHODS OF THE TREATMENT OF

some patency of the foramen existed. It is, however, quite possible that this may have been remedied by the contraction of the dilated heart, and consequent valvular closure of the foramen.

"93, MERRION SQUARE, "August 9th, 1894.

"DEAR SIR PHILIP,

"Having read your paper on the treatment of enlarged heart by movements of the system of the brothers Schott, I beg to add a short note, which, if you wish, you can publish.

"As you know I saw your patient in the most critical portion of her illness, and learned, for the first time, what can be done by these movements, and in confirmation of what you succeeded in accomplishing in a young girl, I now beg to report my own experience of the treatment in a very aged patient.

"Within the last month I met Dr. O'Donoghoe, of Baldoyle, in relation to the case of a very aged gentleman, who was suffering extreme distress of breathing and loss of sleep from a weak dilated heart.

"In addition to the administration of iron and digitalis, and stimulation by a small blister, I used the resisted movements of the arms, and I showed them to Dr. O'D., who fully appreciated their object and value. He and some of the patient's family have still continued them, and the patient is totally changed for the better—sleeps well, has recovered his appetite, and physically shows increased impulse and diminished area of cardiac dulness.

> "I remain, my dear Sir Philip, "Yours most faithfully, "F. R. CRUISE."

On the 19th of September Sir Philip reported :— "X. is wonderful — riding, driving, and boating." He draws attention to the following points, which this case brings into prominence :

1. The importance of the movements without the baths, followed by such an improvement that the patient was able to undertake the journey to Nauheim.

2. The value of the Nauheim baths in removing the ascites and pleural effusion.

3. The interesting observation that the symptoms returned when the movements were discontinued for some weeks, though massage was continued regularly.

INDEX.

Anæmia, 35 Analyses of waters, 8, 9 Aneurism, 31 Animal food, 64 Aortic stenosis, 62 Artificially prepared baths, 16, 17 Asthma, 36, 37

Bath waters, 8, 9, 16, 17, 18 Bronchitis, 38

Carbonic acid, 16, 17 Cold extremities, 39

Diagnostic value of exercises, 31 Diet, 64 Diuresis, 15, 28 Dysmenorrhœa, 39

Effervescence, production of, 16, 17 Exercises, description of, 41 diagnostic value of, 31 in obstruction of pulmonary circuit, 62 influence of, 28 resistance in administering, 62 with self-imposed resistance, 64

Fluids, limitation of, 64

Gastric dilatation, 38

Hands, pale or blue, 30, 39
Headache, 39
Heart, contraction of, 18, 19, 24
Hypertrophy, differential diagnosis of, 32

Inflation of lungs, 28 Influenza, cardiac sequelæ of, 35, 69, 70 Kidneys, effects of baths on, 15 effects of exercises on, 28

Liver, influence of baths on, 15 influence of exercises op, 28 Llangammarch water, 18

Mechanical appliances, 64 Mountain climbing, 64 Murmurs, changes in, 28, 70

Œdema, 36, 77

Pneumonia, 38
Prognostic value of exercises, 32
Psychological influence of Schott methods, 40
Pulse, influence of baths on, 13, 14, 19, 20, 21
Pulse, influence of exercises on 23, 24, 30

Recumbent position, 63 Resistance, measure of, 62 Respiration, effects of baths on, 14 Rheumatism, sequelæ of, 34 Rheumatoid arthritis, 35

Smoker's heart, 67 Sprudel bath, 10, 11 spring, 8 Syncope, precautions against, 62

Temperatures of natural baths, 8,9

Trophic influence of baths, 15

Urine, increased flow of, 15, 28

Vascular capacity, increase of, 29 system, effects on, 14, 23, 24, 29, 30, 38

Widerstandsgymnastik, 25

Pardon & Sons, Printers, Wine Office Court, Fleet Street, London, E.C.

London, 11, New Burlington Street, November, 1894.

A SELECTION

FROM

J. & A. CHURCHILL'S CATALOGUE,

COMPRISING

MOST OF THE RECENT WORKS PUBLISHED BY THEM.

N.B.-J. & A. Churchill's larger Catalogue, which contains over 600 works, with a Complete Index to their Subjects, will be sent on application.

Human Anatomy :

A Treatise by various Authors. Edited by HENRY MORRIS, M.A., M.B. Lond., F.R.C.S., Surgeon to, and Lecturer on Surgery at, the Middlesex Hospital. Roy. 8vo, with 791 Illustrations, nearly all original, and many of them in several colours, 40s. (In one vol. or in three parts.)

Heath's Practical Anatomy:

A Manual of Dissections. Eighth Edition. Edited by WILLIAM ANDERSON, F.R.C.S., Surgeon and Lecturer on Anatomy at St. Thomas's Hospital, Examiner in Anatomy for R.C.P. and S. Crown 8vo, with 329 Engravings, 155.

- Wilson's Anatomist's Vade-Mecum. Eleventh Edition. By HENRY E. CLARK, M.R.C.S. Eng., F.F.P.S. Glasg., Examiner in Anatomy, F.P.S., and Professor of Surgery in St. Mungo's College, Glasgow. Crown Svo, with 492 Engravings and 26 Coloured Plates, 18s.
- An Atlas of Human Anatomy. By RICKMAN J. GODLEE, M.S., F.R.C.S., Surgeon and late Demon-strator of Anatomy, University College Hospital. With 48 Imp. 4to Plates (112 figures), and a volume of Explanatory Text. 8vo, £4 14s. 6d.

Human Osteology.

By LUTHER HOLDEN, Consulting Sur-geon to St. Bartholomew's Hospital. Seventh Edition, edited by CHARLES STEWART, Conservator of the Museum R.C.S., and ROBERT W. REID, M.D., F.R.C.S., Professor of Anatomy in the University of Aberdeen. 8vo, with 59 Lithographic Plates and 75 Engravings. 16s.

Also.

Landmarks, Medical and Surgical. Fourth Edition. Svo, 3s. 6d.

- The Student's Guide to Surgical Anatomy. By EDWARD BELLAMY, F.R.C.S. and Member of the Board of Examiners. Third Edition. Fcap. 8vo, with SI Engravings. 7s. 6d.
- Diagrams of the Nerves of the Human Body, exhibiting their Origin. Divisions, and Connections, with their Distribution to the Various Regions of the Cutaneous Surface, and to all the Muscles. By Sir W. H. FLOWER, K.C.B., F.R.S., F.R.C.S. Third Edition, with 6 Plates. Royal 4to, 12s.
- Pathological Anatomy of Diseases. Arranged according to the nomenclature of the R.C.P. Lond. (Student's Guide Series). By NORMAN MOORE, M.D., F.R.C.P., Assistant Physician and Lecturer on Pathological Anatomy to St. Bartholomew's Hospital. Fcap. 8vo, with 111 Engravings, Ss. 6d.
- A Manual of Clinical and Practical Pathology. tical Pathology. By W. E. WYNTER, M.D., M.R.C.P., F.R.C.S., Medical Registrar to Middlesex Hospital, and F. J. WETHERED, M.D., M.R.C.P., Assistant Physician to Victoria Park Hospital. With 4 Coloured Plates and 67 Engravings. 8vo, 12s. 6d.

Lectures on Pathology:

Delivered at the London Hospital. By the late HENRY GAWEN SUTTON, M.B., F.R.C.P., Physician to, and Lecturer on Pathology at, the London Hospital. Edited by MAURICE E. PAUL, M.D., and Revised by SAMUEL WILKS, M.D., LL.D., F.R.S. 8vo, 15s.

General Pathology: An Introduction to. By JOHN BLAND-SUTTON, F.R.C.S., Sir E. Wilson Lecturer on Pathology, R.C.S.; Assistant Surgeon to, and Lecturer on Anatomy at, Middlesex Hospital. 8vo, with 149 Engravings, 14s.

No. 3.

LONDON: 11, NEW BURLINGTON STREET.
- Atlas of Pathological Anatomy. By Dr. LANCEREAUX. Translated by W. S. GREENFIELD, M.D., Professor of Pathology in the University of Edinburgh. Imp. 8vo, with 70 Coloured Plates, £5 5s.
- Index Pathologicus, for the Registrations of the Lesions recorded in Pathological Records or Case-books of Hospitals and Asylums. By JAMES C. HOWDEN, M.D., Superintendent of the Royal Lunatic Asylum, Montrose. Fcap. folio, 6s.
- Atlas of the Central Nervous System. From the larger work of Hirschfeld and Léveillé. Edited by HOWARD H. TOOTH, M.D., F.R.C.P., Assistant Physician to the National Hospital for the Paralysed and Epileptic. With 37 Plates carefully coloured by Hand. Large Imp. 8vo, 40s.

The Human Brain:

Histological and Coarse Methods of Research. A Manual for Students and Asylum Medical Officers. By W. BEVAN LEWIS, L.R.C.P. Lond., Medical Superintendent, West Riding Lunatic Asylum. Svo, with Wood Engravings and Photographs, 8s.

- Elements of Human Physiology. (Student's Guide Series.) By ERNEST H. STARLING, M.D., M.R.C.P., Joint Lecturer on Physiology at Guy's Hospital. Fcap. 8vo, with 94 Engravings, 6s. 6d.
- Manual of Physiology:
 - For the use of Junior Students of Medicine. By GERALD F. YEO, M.D., F.R.S., Emeritus Professor of Physiology in King's College, London. Third Edition. Crown 8vo, with 254 Engravings (many figures), and Plate of Spectra, 14s.
- Principles of Human Physiology. By W. B. CARPENTER, C.B., M.D., F.R.S. Ninth Edition. By HENRY POWER, M.B., F.R.C.S. 8vo, with 3 Steel Plates and 377 Wood Engravings, 31s. 6d.
- Practical Lessons in Elementary Biology, for Junior Students. By PEYTON T. B. BEALE, F.R.C.S., Lecturer on Elementary Biology and Demonstrator in Physiology in King's College, London. Crown 8vo, 3s. 6d.
- Medical Jurisprudence: Its Principles and Practice. By ALFRED S. TAYLOR, M.D., F.R.C.P., F.R.S. Fourth Edition, by THOMAS STEVENSON, M.D., F.R.C.P., Lecturer on Medical Jurisprudence at Guy's Hospital. 2 vols. 8vo, with 189 Engravings, 31s. 6d.

By the same Authors.

A Manual of Medical Jurisprudence. Twelfth Edition. Crown 8vo, with 55 Engravings, 14s.

- The Student's Guide to Medical Jurisprudence. By JOHN ABER-CROMBIE, M.D., F.R.C.P., Physician to Charing Cross Hospital. Fcap. 8vo, 7s. 6d.
- Sanitary Examinations

Of Water, Air, and Food. A Vade-Mecum for the Medical Officer of Health. By CORNELIUS B. FOX, M.D., F.R.C.P. Second Edition. Crown 8vo, with 110 Engravings, 12s. 6d.

Microscopical Examination of Drinking Water and of Air. By J. D. MACDONALD, M.D., F.R.S., Ex-Professor of Naval Hygiene in the Army Medical School. Second Edition. 8vo, with 25 Plates, 7s. 6d.

Hygiene and Public Health.

- A Treatise by various Authors. Edited by THOMAS STEVENSON, M.D., F.R.C.P., Lecturer on Chemistry and Medical Jurisprudence at Guy's Hospital; Official Analyst to the Home Office; and SHIRLEY F. MURPHY, Medical Officer of Health of the County of London. In 3 vols., royal 8vo, fully Illustrated. Vol. I., 28s.; Vol. II., 32s.; Vol. III., 20s.
- A Manual of Practical Hygiene. By the late E. A. PARKES, M.D., F.R.S. Eighth Edition, by J. LANE NOTTER, A.M., M.D., F.R.S., Professor of Military Hygiene in the Army Medical School. 8vo, with 10 Plates and 103 Engravings, 18s.
- A Handbook of Hygiene and Sanitary Science. By GEO. WILSON, M.A., M.D., LL.D., F.R.S.E., D.P.H. Camb., Medical Officer of Health for Mid-Warwickshire. Seventh Edition. Crown 8vo, with Engravings, 12s. 6d.
- The Prevention of Epidemics and the Construction and Management of Isolation Hospitals. By ROGER MCNEILL, M.D. Edin., D.P.H. Camb., Medical Officer of Health for the County of Argyll. With numerous Illustrations. 8vo.
- Hospitals and Asylums of the World; their Origin, History, Construction, Administration, Management, and Legislation. By HENRY C. BURDETT. In 4 vols. Super Royal 8vo and Portfolio. Complete, 168s. Vols. I. and II.—Asylums: their History, Administration, and Construction, with Plans and Bibliography, 90s. Vols. III. and IV. —Hospitals, Convalescent Institutions, Nurses' Homes, Medical School Buildings, &c., with Portfolio of Plans, 120s.

Mental Diseases:

Clinical Lectures. By T. S. CLOUSTON, M.D., F.R.C.P. Edin., Lecturer on Mental Diseases in the University of Edinburgh. Third Edition. Crown 8vo, with 13 Plates, 14s.

2

Illustrations of the Influence of the Mind upon the Body in Health and Disease: Designed to elucidate the Action of the Imagination. By D. HACK TUKE, M.D., F.R.C.P., LL.D. Second Edition. 2 vols. crown Svo, 15s.

- Prichard and Symonds in Especial Relation to Mental Science. With Chapters on Moral Insanity. 8vo, with 2 Portraits, 5s. Also.
- Reform in the Treatment of the Insane. Early History of the Retreat, York; its Objects and Influence. With a Report of the Celebrations of its Centenary. 8vo, 4s.
- A Dictionary of Psychological Medicine, giving the Definition, Etymology, and Synonyms of the Terms used in Medical Psychology; with the Symptoms, Treatment, and Pathology of Insanity; and THE LAW OF LUNACY IN GREAT BRITAIN AND IRELAND. Edited by D. HACK TUKE, M.D., LL.D., assisted by nearly 130 Contributors, British, Continental and American. 2 vols., 1,500 pages, royal 8vo, Illustrated. 42s.
- Lunacy Law for Medical Men. By CHARLES MERCIER, M.B., Lecturer on Neurology and Insanity to the Westminster Hospital Medical School, and to the Medical School for Women. Crown 8vo, 5s.
- The Journal of Mental Science. Published Quarterly, by Authority of the Medico-Psychological Association. 8vo, 3s. 6d.
- Mental Affections of Childhood and Youth (Lettsomian Lectures for 1887, &c.). By J. LANGDON DOWN, M.D., F.R.C.P., Consulting Physician to the London Hospital. 8vo, 6s.
- Manual of Midwifery : Including all that is likely to be required by Students and Practitioners. By ALFRED L. GALABIN, M.A., M.D., F.R.C.P., Obstetric Physician to, and Lecturer on, Midwifery, &c., at Guy's Hospital. Third Edition. Crown 8vo, with 261 Engravings, 155.
- The Student's Guide to the Practice of Midwifery. By D. LLOYD ROBERTS, M.D., F.R.C.P., Lecturer on Clinical Midwifery and Diseases of Women at the Owens College; Obstetric Physician to the Manchester Royal Infirmary. Fourth Edition. Fcap. 8vo, with Coloured Plates and Engravings.
- [Preparing. Manual of the Diseases peculiar to Women. By JAMES OLIVER, M.D., F.R.S. Edin., M.R.C.P. Lond., Physician to the Hospital for Women, London. Fcap. 8vo, 3s. 6d.

Obstetric Aphorisms :

For the Use of Students commencing Midwifery Practice. By JOSEPH G. SWAYNE, M.D. Tenth Edition. Fcap. 8vo, with 20 Engravings, 3s. 6d.

- Female Pelvic Organs (The Surgery, Surgical Pathology, and Surgical Anatomy of): in a Series of Plates taken from Nature. With Commentaries, Notes, and Cases. By HENRY SAVAGE, M.D., Consulting Physician to the Samaritan Hospital for Women and Children. Fifth Edition. 4to, Uncoloured, 15s.
- Lectures on Obstetric Operations: Including the Treatment of Hæmorrhage, and forming a Guide to the Management of Difficult Labour. By ROBERT BARNES, M.D., F.R.C.P., Consulting Obstetric Physician to St. George's Hospital. Fourth Edition. 8vo, with 121 Engravings, 128. 6d.

By the same Author.

- A Clinical History of Medical and Surgical Diseases of Women. Second Edition. 8vo, with 181 Engravings, 28s.
- Clinical Lectures on Diseases of Women : Delivered in St. Bartholomew's Hospital, by J. MATTHEWS DUNCAN, M.D., LL.D., F.R.C.P., F.R.Ss. L. & E., late Obstetric Physician to St. Bartholomew's Hospital. Fourth Edition. Svo, 16s.
- Gynæcological Operations: (Handbook of). By ALBAN H. G. DORAN, F.R.C.S., Surgeon to the Samaritan Hospital. 8vo, with 167 Engravings, 155.
- The Student's Guide to the Diseases of Women. By ALFRED L. GALABIN, M.A., M.D., F.R.C.P., Obstetric Physician to Guy's Hospital. Fifth Edition. Fcap. 8vo, with 142 Engravings, 8s. 6d.
- A Practical Treatise on the Diseases of Women. By T. GAIL-LARD THOMAS, M.D. Sixth Edition, by PAUL F. MUNDÉ, M.D., Professor of Gynæcology at the New York Polyclinic and at Dartmouth College. Roy. 8vo, with 347 Engravings, 25s.
- Notes on Diseases of Women: Specially designed to assist the Student in preparing for Examination. By JAMES J. REYNOLDS, L.R.C.P., M.R.C.S. Fourth Edition, Fcap. 8vo., 3s. 6d.

Abdominal Surgery.

By J. GREIG SMITH, M.A., F.R.S.E., Surgeon to the Bristol Royal Infirmary, and Lecturer on Surgery in the Bristol Medical School. Fourth Edition. 8vo, with 82 Engravings, 21s.

By the same Author.

- The Physiology of Death from Traumatic Fever; A Study in Abdominal Surgery. By JOHN D. MAL-COLM, M.B., C.M., F.K.C.S.E., Sur-geon to the Samaritan Free Hospital. Svo, 3s. 6d.
- Notes on Gynæcological Nursing. By JOHN BENJAMIN HELLIER, M.D., M.R.C.S., Lecturer on the Diseases of Women and Children in the Yorkshire College, and Surgeon to the Hospital for Women, &c., Leeds. Crown 8vo, IS. 6d.
- A Manual for Hospital Nurses and others engaged in Attending on the Sick, with a Glossary. By EDWARD J. DOMVILLE, Surgeon to the Exeter Lyingin Charity. Seventh Edition. Crown 8vo, 2s. 6d.
- A Manual of Nursing, Medical and Surgical. By CHARLES J. CUL-LINGWORTH, M.D., F.R.C.P., Obstetric Physician to St. Thomas's Hospital. Third Edition. Fcap. 8vo, with Engravings, 2s. 6d.

By the same Author.

A Short Manual for Monthly Nurses. Third Edition. Fcap. 8vo, 1s. 6d.

Diseases of Children.

For Practitioners and Students. By W. H. DAY, M.D., Physician to the Samaritan Hospital. Second Edition. Crown Svo, 128. 6d.

- The Diseases of Children (Student's Guide Series). By JAS. F. GOOD-HART, M.D., F.R.C.P., Physician to Guy's Hospital. Fifth Edition. Fcap. 8vo, 10s. 6d.
- A Practical Treatise on Disease in Children. By EUSTACE SMITH, M.D., F.R.C.P., Physician to the King of the Belgians, and to the East London Hospital for Children, &c. Second Edition, 8vo, 22s.

By the same Author.

Clinical Studies of Disease in Children. Second Edition. Post 8vo, 7s. 6d.

Also.

- The Wasting Diseases of Infants and Children. Fifth Edition. Post 8vo, 8s. 6d.
- Practical Manual of the A Diseases of Children. With a For-By EDWARD ELLIS, mulary. M.D. Fifth Edition. Crown 8vo, 10s.

Materia Medica:

A Manual for the use of Students. By ISAMBARD OWEN, M.D., F.R.C.P., Lecturer on Materia Medica, &c., to St. Second Edition. George's Hospital. Crown 8vo, 6s. 6d.

Materia Medica,

Pharmacy, Pharmacology, and Thera-peutics. By W. HALE WHITE, M.D., F.R.C.P., Physician to, and Lecturer on Materia Medica and Therapeutics at, Guy's Hospital; Examiner in Materia Medica on the Conjoint Board of the Royal Colleges of Physicians and Surgeons. Fcap. Svo, 7s. 6d. Materia Medica

And Therapeutics. By CHARLES D. F. PHILLIPS, M.D., F.R.S. Edin.

Vegetable Kingdom-Organic Compounds-Animal Kingdom. 8vo, 25s. Inorganic Substances. Second Edition. Svo, 215.

Recent Materia Medica.

Notes on their Origin and Therapeutics. By F. HARWOOD LESCHER, F.C.S., Pereira Medallist. Fourth Edition. 8vo, 2s. 6d

Galenic Pharmacy:

A Practical Handbook to the Processes of the British Pharmacopœia. By R. A. CRIPPS, M.P.S. 8vo, with 76 Engravings, 8s. 6d.

The Galenical Preparations of the British Pharmacopœia: A Handbook for Medical and Pharmaceutical Students. By CHARLES O. HAWTHORNE, M.B., C.M., Lecturer on Materia Medica and Therapeutics, Queen Margaret College, University of Glasgow. Svo.

Practical Pharmacy.

By BARNARD S. PROCTOR, formerly Lecturer on Pharmacy at the College of Medicine, Newcastle-on-Tyne. Third Edition. Svo, with 44 Wood Engravings and 32 Lithograph Fac-Simile Prescriptions, 14s.

Selecta è Prescriptis :

Containing Terms, Phrases, Contractions and Abbreviations used in Prescriptions, with Explanatory Notes, &c. Also, a Series of Abbreviated Prescriptions with Translations and Key. By J. PEREIRA, M.D., F.R.S. Eighteenth Edition, by JOSEPH INCE, F.C.S., F.L.S. 24mo, 5s.

A Companion to the British Pharmacopœia. By PETER SQUIRE, Revised by his Sons, P. W. and A. H. SQUIRE. Sixteenth Edition. 8vo, 12s. 6d.

By the same Authors.

The Pharmacopœias of the London Hospitals, arranged in Groups for Easy Reference and Comparison. Sixth Edition. 18mo. 6s.

The National Dispensatory :

Containing the Natural History, Chemistry, Pharmacy, Actions and Uses of Medicines, &c. By ALFRED STILLÉ, M.D., LL.D., JOHN M. MAISCH, Phar. D., CHAS. CASPARI, JUN., Ph.G., and HENRY C. C. MAISCH, Ph.G., Ph.D. Fifth Edition, with 320 Engravings. Imp. 8vo, 36s.

Pocket Formulary

And Synopsis of the British and Foreign Pharmacopœias. By HENRY BEASLEY. Eleventh Edition. 18mo, 6s. 6d.

By the same Author.

Druggist's General Receipt-Book. Ninth Edition. 18mo, 6s. 6d. Also.

Book of Prescriptions:

Containing upwards of 3,000 Prescriptions from the Practice of the most eminent Physicians and Surgeons, English and Foreign. Seventh Edition. 18mo, 6s. 6d.

The Prescriber's Pharmacopœia:

The Medicines arranged in Classes according to their Action, with their Composition and Doses. By NESTOR J. C. TIRARD, M.D., F.R.C.P., Professor of Materia Medica and Therapeutics in King's College, London. Sixth Edition. 32mo, bound in leather, 3s.

Year-Book of Pharmacy:

- Containing the Transactions of the British Pharmaceutical Conference. Annually. 8vo, 10s.
- Royle's Manual of Materia Medica and Therapeutics. Sixth Edition, including additions and alterations in the B.P. 1885. By JOHN HARLEY, M.D., Physician to St. Thomas's Hospital. Crown Svo, with 139 Engravings, 15s.
- The Student's Guide to Structural, Morphological, and Physiological Botany. By ROBERT BENTLEY, late Emeritus Professor of Botany in King's College and to the Pharmaceutical Society. With 660 Engravings. Fcap. 8vo, 7s. 6d.

By the same Author.

The Student's Guide to Systematic Botany, including the Classification of Plants and Descriptive Botany. Fcap. 8vo, with 350 Engravings, 3s. 6d.

Medicinal Plants:

Being descriptions, with original figures, of the Principal Plants employed in Medicine, and an account of their Properties and Uses. By Prof. BENTLEY and Dr. H. TRIMEN, F.R.S. In 4 vols., large Svo, with 306 Coloured Plates, bound in Half Morocco, Gilt Edges, £11 115. Climate and Fevers of India, with a series of Cases (Croonian Lectures, 1882). By Sir JOSEPH FAYRER, K.C.S.I., M.D. 8vo, with 17 Temperature Charts, 12s.

By the same Author.

- The Natural History and Epidemiology of Cholera: Being the Annual Oration of the Medical Society of London, 1888. 8vo, 3s. 6d.
- A Manual of Family Medicine and Hygiene for India. Published under the Authority of the Government of India. By Sir WILLIAM J. MOORE, K.C.I.E., M.D., late Surgeon-General with the Government of Bombay. Sixth Edition. Post 8vo, with 71 Engravings, 12s.

By the same Author.

A Manual of the Diseases of India: With a Compendium of Diseases generally. Second Edition. Post 8vc, 10s.

Also,

- The Constitutional Requirements for Tropical Climates, &c. Crown 8vo, 4s.
- The Prevention of Disease in Tropical and Sub-Tropical Campaigns. (Parkes Memorial Prize for 1886.) By ANDREW DUNCAN, M.D., B.S. Lond., F.R.C.S., Surgeon-Major, Bengal Army. 8vo, 12s. 6d.

Practical Therapeutics:

A Manual. By EDWARD J. WARING, C.I.E., M.D., F.R.C.P., and DUDLEY W. BUXTON, M.D., B.S. Lond. Fourth Edition. Crown 8vo, 14s.

By the same Author.

Bazaar Medicines of India,

- And Common Medical Plants : With Full Index of Diseases, indicating their Treatment by these and other Agents procurable throughout India, &c. Fourth Edition Fcap. 8vo, 5s.
- A Commentary on the Diseases of India. By NORMAN CHEVERS, C.I.E., M.D., F.R.C.S., Deputy Surgeon-General H.M. Indian Army. 8vo, 245.
- Hooper's Physicians' Vade-Mecum. A Manual of the Principles and Practice of Physic. Tenth Edition. By W. A. GUY, F.R.C.P., F.R.S., and J. HARLEY, M.D., F.R.C.P. With 118 Engravings. Fcap. 8vo, 12s. 6d.

- The Principles and Practice of Medicine. (Text-book.) By the late C. HILTON FAGGE, M.D., and P. H. PYE-SMITH, M.D., F.R.S., F.R.C.P., Physician to, and Lecturer on Medicine in, Guy's Hospital. Third Edition. 2 vols. 8vo, cloth, 40s.; Half Leather, 46s.
- Manual of the Practice of Medicine. By FREDERICK TAYLOR, M.D., F.R.C.P., Physician to, and Lecturer on Medicine at, Guy's Hospital. Third Edition. Cr. 8vo, with Engravings, 15s.
- The Practice of Medicine (Student's Guide Series). By M. CHARTERIS, M.D., Professor of Therapeutics and Materia Medica in the University of Glasgow. Seventh Edition. Fcap. 8vo, with Engravings on Copper and Wood, 10s.
- A Dictionary of Practical Medicine. By various writers. Edited by JAS. KINGSTON FOWLER, M.A., M.D., F.R.C.P., Physician to Middlesex Hospital and the Hospital for Consumption. 8vo, cloth, 21s. ; half calf, 25s.
- How to Examine the Chest: A Practical Guide for the use of Students. By SAMUEL WEST, M.D., F.R.C.P., Assistant Physician to St. Bartholomew's Hospital. Second Edition. With Engravings. Fcap. 8vo, 5s.
- The Bronchi and Pulmonary Blood-vessels: their Anatomy and Nomenclature. By WILLIAM EWART, M.D., F.R.C.P., Physician to St. George's Hospital. 4to, with 20 Illustrations, 21s.
- An Atlas of the Pathological Anatomy of the Lungs. By the late WILSON FOX, M.D., F.R.S., F.R.C.P., Physician to H.M. the Queen. With 45 Plates (mostly Coloured) and Engravings. 4to, half-bound in Calf, 70s.

By the same Author.

- A Treatise on Diseases of the Lungs and Pleura. Edited by SIDNEY COUPLAND, M.D., F.R.C.P., Physician to Middlesex Hospital. Roy. 8vo, with Engravings; also Portrait and Memoir of the Author, 36s.
- The Student's Guide to Diseases of the Chest. By VINCENT D. HARRIS, M.D. Lond., F.R.C.P., Physician to the City of London Hospital for Diseases of the Chest, Victoria Park. Fcap. 8vo, with 55 Illustrations (some Coloured), 7s. 6d.

Uric Acid

as a Factor in the Causation of Disease. Bv ALEXANDER HAIG, M.D., F.R.C.P., Physician to the Metropolitan Hospital and the Royal Hospital for Children and Women. Second Edition. With Illustrations, 8vo, 10s. 6d.

- Medical Diagnosis (Student's Guide Series). By SAMUEL FENWICK, M.D., F.R.C.P., Physician to the London Hospital. Seventh Edition. Fcap. 8vo, with 117 Engravings, 7s. By the same Author.
- Outlines of Medical Treatment. Fourth Edition. Crown Svo, with 35 Engravings, 10s.

Also.

Clinical Lectures on Some Obscure Diseases of the Abdomen. Delivered at the London Hospital. 8vo, with Engravings, 7s. 6d.

Also.

- The Saliva as a Test for Functional Diseases of the Liver. Crown 8vo, 2s.
- The Microscope in Medicine. By LIONEL S. BEALE, M.B., F.R.S., Physician to King's College Hospital. Fourth Edition. 8vo, with 86 Plates, 21s. By the same Author.
- The Liver.

With 24 Plates (85 Figures). 8vo, 5s. Also.

On Slight Ailments:

And on Treating Disease. Third Edition. 8vo, 5s.

- The Physiology of the Carbohydrates; their Application as Food and Relation to Diabetes. By F. W. PAVY, M.D., LL.D., F.R.S., F.R.C.P., Consulting Physician to Guy's Hospital. Royal 8vo, with Plates and Engravings, 105. 6d.
- Medical Lectures and Essays. By Sir G. JOHNSON, M.D., F.R.C.P., F.R.S., Consulting Physician to King's College Hospital. 8vo, with 46 Engravings, 25s.

By the same Author.

An Essay on Asphyxia (Apnœa). ^{8vo, 3s.}

Bronchial Affections:

Pneumonia and Fibroid Pneumonia (their Pathological Histology). An Original Investigation. By A. G. AULD, M.D., Assistant Physician to the Glasgow Royal Infirmary. 8vo, with Illustrations, 7s. 6d. Bronchial Asthma:

Its Pathology and Treatment. By J. B. BERKART, M.D., late Physician to the City of London Hospital for Diseases of the Chest. Second Edition, with 7 Plates (35 Figures). 8vo, 10s. 6d.

Treatment of Some of the Forms of Valvular Disease of the Heart. By A. E. SANSOM, M.D., F.R.C.P., Physician to the London Hospital. Second Edition. Fcap. 8vo, with 26 Engravings, 4s. 6d.

Antiseptic Dry-Air Treatment of Consumption. By JOHN J. HART-NETT, M.D., M.Ch., Q.U.I. With Engravings. Crown 8vo, 3s. 6d.

Guy's Hospital Reports. By the Medical and Surgical Staff. Vol. XXXV. Third Series. Svo. 10s. 6d.

- St. Thomas's Hospital Reports. By the Medical and Surgical Staff. Vol. XXI. New Series. 8vo, 8s. 6d.
- Westminster Hospital Reports. By the Medical and Surgical Staff. Vol. VIII. 8vo, 6s.

The Climate of Rome

and the Roman Malaria. By Professor TOMMASI-CRUDELI. Translated by CHARLES CRAMOND DICK. Crown 8vo, 5s.

Vaccinia and Variola:

A Study of their Life History. By JOHN B. BUIST, M.D., F.R.S.E., Teacher of Vaccination for the Local Government Board. Crown 8vo, with 24 Coloured Plates, 7s. 6d.

Medical Ophthalmoscopy :

A Manual and Atlas. By W. R. GOWERS, M.D., F.R.C.P., F.R.S., Physician to the National Hospital for the Paralysed and Epileptic. Third Edition. Edited with the assistance of MARCUS GUNN, M.B., F.R.C.S., Surgeon to the Royal London Ophthalmic Hospital. With Coloured Plates and Woodcuts. 8vo, 16s.

By the same Author.

A Manual of Diseases of the Nervous System.

Vol. I. Diseases of the Nerves and Spinal Cord. Second Edition. Roy. 8vo, with 179 Engravings, 15s. Vol. II. Diseases of the Brain and

Vol. II. Diseases of the Brain and Cranial Nerves : General and Functional Diseases of the Nervous System. Second Edition. Roy. 8vo, with 182 Engravings, 205.

Also.

Diagnosis of Diseases of the Brain. Second Edition. 8vo, with Engravings, 7s. 6d.

Aiso.

Syphilis and the Nervous System. Being a Revised Reprint of the Lettsomian Lectures for 1890. Delivered before the Medical Society of London. 8vo, 4s.

The Nervous System,

Diseases of. By J. A. ORMEROD, M.D., F.R.C.P., Physician to the National Hospital for the Paralysed and Epileptic. With 66 Illustrations. Fcap. 8vo, 8s. 6d. Handbook of the Diseases of the Nervous System. By JAMES Ross, M.D., F.R.C.P., Professor of Medicine in the Victoria University, and Physician to the Royal Infirmary, Manchester. Roy. 8vo, with 184 Engravings, 18s.

Also.

Aphasia:

Being a Contribution to the Subject of the Dissolution of Speech from Cerebral Disease. 8vo, with Engravings. 4s. 6d.

Diseases of the Nervous System. Lectures delivered at Guy's Hospital. By SAMUEL WILKS, M.D., F.R.S. Second Edition. 8vo, 18s.

Stammering:

Its Causes, Treatment, and Cure. By A. G. BERNARD, M.R.C.S., L.R.C.P. Crown 8vo, 2s.

- Secondary Degenerations of the Spinal Cord (Gulstonian Lectures, 1889). By HOWARD H. TOOTH, M.D., F.R.C.P., Assistant Physician to the National Hospital for the Paralysed and Epileptic. With Plates and Engravings. 8vo, 3s. 6d.
- Diseases of the Nervous System. Clinical Lectures. By THOMAS BUZZARD, M.D., F.R.C.P., Physician to the National Hospital for the Paralysed and Epileptic. With Engravings, 8vo. 15s.

By the same Author.

Some Forms of Paralysis from Peripheral Neuritis: of Gouty, Alcoholic, Diphtheritic, and other origin. Crown 8vo, 5s.

Also.

- On the Simulation of Hysteria by Organic Disease of the Nervous System. Crown 8vo, 4s. 6d.
- Gout in its Clinical Aspects. By J. MORTIMER GRANVILLE, M.D. Crown 8vo, 6s.

Diseases of the Liver:

With and without Jaundice By GEORGE HARLEY, M.D., F.R.C.P., F.R.S. 8vo, with 2 Plates and 36 Engravings, 21s.

Rheumatic Diseases,

(Differentiation in). By HUGH LANE, Surgeon to the Royal Mineral Water Hospital, Bath, and Hon. Medical Officer to the Royal United Hospital, Bath. Second Edition, much Enlarged, with 8 Plates. Crown 8vo, 3s. 6d.

Diseases of the Abdomen,

Comprising those of the Stomach and other parts of the Alimentary Canal, Œsophagus, Cæcum, Intestines, and Peritoneum. By S. O. HABERSHON, M.D., F.R.C.P. Fourth Edition. 8vo, with 5 Plates, 21s. On the Relief of Excessive and Dangerous Tympanites by Puncture of the Abdomen. By JOHN W. OGLE, M.A., M.D., F.R.C.P., Consulting Physician to St. George's Hospital. 8vo, 5s. 6d.

Headaches :

Their Nature, Causes, and Treatment. By W. H. DAY, M.D., Physician to the Samaritan Hospital. Fourth Edition. Crown 8vo, with Engravings, 7s. 6d.

- Health Resorts at Home and Abroad. By M. CHARTERIS, M.D., Professor of Therapeutics and Materia Medica in Glasgow University. Second Edition. Crown 8vo, with Map, 5s. 6d.
- The Mineral Waters of France And its Wintering Stations (Medical Guide to). With a Special Map. By A. VINTRAS, M.D., Physician to the French Embassy, and to the French Hospital, London. Second Edition. Crown 8vo, 8s.

Canary Islands

Health Resorts, in their Climatological and Medical Aspects. By J. CLEASBY TAYLOR, M.D., M.R.C.S., Las Palmas. 8vo, with Maps, 3s. 6d.

Homburg-Spa.

An Introduction to its Waters and their use. By Dr. ARNOLD SCHETELIG. Crown 8vo, with Synoptical Table, 2s. 6d.

IllustratedAmbulance Lectures: To which is added a NURSING LECTURE. By JOHN M. H. MARTIN, M. D., F. R. C. S., Honorary Surgeon to the Blackburn Infirmary. Third Edition. Crown Svo, with 60 Engravings, 2s.

Surgery: its Theory and Practice (Student's Guide). By WILLIAM J. WALSHAM, F.R.C.S., Senior Assistant Surgeon to, and Lecturer on Anatomy at, St. Bartholomew's Hospital. Fourth Edition. Fcap. 8vo, with 335 Engravings, 125.

Surgical Emergencies:

Together with the Emergencies attendant on Parturition and the Treatment of Poisoning. ByW, PAUL SWAIN, F.R.C.S., Surgeon to the South Devon and East Cornwall Hospital. Fourth Edition. Crown 8vo, with 120 Engravings, 5s.

Operations on the Brain (A Guide to). By ALEC FRASER, Professor of Anatomy, Royal College of Surgeons in Ireland. Illustrated by 42 life-size Plates in Autotype, and 2 Woodcuts in the text. Folio, 63s. A Course of Operative Surgery. By CHRISTOPHER HEATH, Surgeon to University College Hospital. Second Edition. With 20 coloured Plates (180 figures) from Nature, by M. LÉVEILLÉ, and several Woodcuts. Large 8vo, 30s.

By the same Author.

The Student's Guide to Surgical Diagnosis. Second Edition. Fcap. 8vo, 6s. 6d.

Also.

Manual of Minor Surgery and Bandaging. For the use of House-Surgeons, Dressers, and Junior Practitioners. Tenth Edition. Fcap. 8vo, with 158 Engravings, 6s.

Also.

Injuries and Diseases of the Jaws. Fourth Edition. By HENRY PERCY DEAN, M.S., F.R.C.S., Assistant Surgeon to the London Hospital. 8vo, with 187 Wood Engravings, 145.

Also.

Lectures on Certain Diseases of the Jaws. Delivered at the R.C.S., Eng., 1887. 8vo, with 64 Engravings, 2s. 6d.

Also.

Clinical Lectures on Surgical Subjects. Delivered in University College Hospital. Fcap. 8vo, with 23 Engravings, 6s.

Surgery.

By C. W. MANSELL MOULLIN, M.A., M.D., Oxon., F.R.C.S., Surgeon and Lecturer on Physiology to the London Hospital. Large 8vo, with 497 Engravings, 34s.

The Practice of Surgery :

A Manual. By THOMAS BRYANT, Consulting Surgeon to Guy's Hospital. Fourth Edition. 2 vols. crown 8vo, with 750 Engravings (many being coloured), and including 6 chromo plates, 32s.

By the same Author.

On Tension: Inflammation of Bone, and Head Injuries. Hunterian Lectures, 1888. 8vo, 6s.

The Surgeon's Vade-Mecum : A Manual of Modern Surgery. By R. DRUITT, F.R.C.S. Twelfth Edition. By STANLEY BOYD, M.B., F.R.C.S. Assistant Surgeon and Pathologist to Charing Cross Hospital. Crown Svo, with 373 Engravings, 16s.

Diseases of Bones and Joints. By CHARLES MACNAMARA, F.R.C.S., Surgeon to, and Lecturer on Surgery at, the Westminster Hospital. 8vo, with Plates and Engravings, 12s.

8

The Operations of Surgery :

Intended for Use on the Dead and Living Subject alike. By W. H. A. JACOBSON, M.A., M.B., M.Ch. Oxon., F.R.C.S., Assistant Surgeon to, and Lecturer on Anatomy at, Guy's Hospital. Second Edition. 8vo, with 235 Illustrations, 30s.

On Anchylosis.

By BERNARD E. BRODHURST, F.R.C.S., Surgeon to the Royal Orthopædic Hospital. Fourth Edition. 8vo, with Engravings, 5s.

By the same Author.

- Curvatures and Disease of the Spine. Fourth Edition. 8vo, with Engravings, 7s. 6d. Also.
- Talipes Equino-Varus, or Clubfoot. 8vo, with Engravings, 3s. 6d.
- Surgical Pathology and Morbid Anatomy (Student's Guide Series). By ANTHONY A. BOWLBY, F.R.C.S., Assistant Surgeon to St. Bartholomew's Hospital. Second Edition. Fcap. 8vo, with 158 Engravings, 9s.

By the same Author.

- Injuries and Diseases of Nerves and their Surgical Treatment. 8vo, with 20 Plates. 14s.
- Illustrations of Clinical Surgery. By JONATHAN HUTCHINSON, F.R.S., Senior Surgeon to the London Hospital. In fasciculi. 6s. 6d. each. Fasc. I. to X. bound, with Appendix and Index, £3 IOS. Fasc. XI. to XXIII. bound, with Index, £4 IOS.

By the same Author.

Archives of Surgery; Quarterly. Nos. i. to xxi. 8vo, 2s. 6d. each. Vols. I. to V., each containing 4 Nos., cloth, 10s. 6d. each.

Clubfoot :

Its Causes, Pathology, and Treatment. By WM. ADAMS, F.R.C.S., Consulting Surgeon to the Great Northern and other Hospitals. Second Edition. Svo, with 106 Engravings and 6 Lithographic Plates, 155.

By the same Author.

Lateral and other Forms of Curvature of the Spine: Their Pathology and Treatment. Second Edition. 8vo, with 5 Lithographic Plates and 72 Wood Engravings, 10s. 6d.

Also.

Contraction of the Fingers:

(Dupuytren's and Congenital Contractions), their Treatment by Subcutaneous Divisions of the Fascia, and Immediate Extension. Also on Hammer Toe; its Curability by Subcutaneous Division. And on The Obliteration of Depressed Cicatrices by a Subcutaneous Operation. 8vo, with 8 Plates and 31 Engravings, 6s. 6d. Treatment of Internal Derangements of the Knee-Joint, by Operation. By HERBERT W. ALLING-HAM, F.R.C.S., Surgeon to the Great Northern Central Hospital, &c. 8vo, with Engravings, 5s.

Short Manual of Orthopædy.

By HEATHER BIGG, F.R.C.S. Ed. Part I. Deformities and Deficiencies of the Head and Neck. 8vo. 2s. 6d.

Face and Foot Deformities.

By FREDERICK CHURCHILL, C.M. Svo, with Plates and Illustrations, 10s. 6d.

The Human Foot:

Its Form and Structure, Functions and Clothing. By THOMAS S. ELLIS, Consulting Surgeon to the Gloucester Infirmary. With 7 Plates and Engravings (50 Figures). 8vo, 7s. 6d.

Royal London Ophthalmic Hospital Reports. By the Medical and Surgical Staff. Vol. XIII., Part 4. 8vo, 5s.

Ophthalmological Society

of the United Kingdom. Transactions, Vol. XIII. 8vo, 12s. 6d.

The Diseases of the Eye

(Student's Guide Series). By EDWARD NETTLESHIP, F.R.C.S., Ophthalmic Surgeon to St. Thomas's Hospital. Fifth Edition. Fcap. 8vo, with 164 Engravings and a Coloured Plate illustrating Colour-Blindness, 7s. 6d.

Diseases and Refraction of the Eye. By N. C. MACNAMARA, F.R.C.S., Surgeon to Westminster Hospital, and GUSTAVUS HARTRIDGE, F.R.C.S., Surgeon to the Royal Westminster Ophthalmic Hospital. Fifth Edition. Crown Svo, with Plate, 156 Engravings, also Testtypes, 10s. 6d.

On Diseases and Injuries of the Eye: A Course of Systematic and Clinical Lectures to Students and Medical Practitioners. By J. R. WOLFE, M.D., F.R.C.S.E., Lecturer on Ophthalmic Medicine and Surgery in Anderson's College, Glasgow. With 10 Coloured Plates and 157 Wood Engravings. 8vo, £1 1s.

Normal and Pathological Histology of the Human Eye and Eyelids. By C. FRED. POLLOCK, M.D., F.R.C.S. and F.R.S.E., Surgeon for Diseases of the Eye to Anderson's College Dispensary, Glasgow. Crown 8vo, with 100 Plates (230 drawings), 15s.

Refraction of the Eye:

A Manual for Students. By GUSTAVUS HARTRIDGE, F.R.C.S., Surgeon to the Royal Westminster Ophthalmic Hospital. Seventh Edition. Crown 8vo, with 98 Illustrations, also Test-types, &c., 6s. By the same Author.

The Ophthalmoscope. A Manual for Students. Second Edition. Crown 8vo, with 67 Illustrations and 4 Plates. 4s. 6d.

Atlas of Ophthalmoscopy.

Composed of 12 Chromo - lithographic Plates (59 Figures drawn from nature) and Explanatory Text. By RICHARD LIEBREICH, M.R.C.S. Translated by H. ROSBOROUGH SWANZY, M.B. Third Edition, 4to, 4os.

Glaucoma:

Its Pathology and Treatment. By PRIESTLEY SMITH, Ophthalmic Surgeon to, and Clinical Lecturer on Ophthalmology at, the Queen's Hospital, Birmingham. 8vo, with 64 Engravings and 12 Zinco-photographs, 7s. 6d.

Eyestrain

(commonly called Asthenopia). By ERNEST CLARKE, M.D., B.S. Lond., Surgeon to the Central London Ophthalmic Hospital, Surgeon and Ophthalmic Surgeon to the Miller Hospital. 8vo, with 22 Illustrations, 5s.

Diseases of the Eye:

A Handbook of Ophthalmic Practice for Students and Practitioners. By G. E. DE SCHWEINITZ, M.D., Professor of Diseases of the Eye in the Philadelphia Polyclinic. With 216 Illustrations, and 2 Chromo-Lithographic Plates. 8vo, 18s.

- Diseases and Injuries of the Ear. By Sir WILLIAM B. DALBY, F.R.C.S., M.B., Consulting Aural Surgeon to St. George's Hospital. Fourth Edition. Crown 8vo, with 8 Coloured Plates and 38 Wood Engravings. 10s. 6d. By the same Author.
- Short Contributions to Aural Surgery, between 1875 and 1889. Second Edition. 8vo, with Engravings, 3s. 6d.

Diseases of the Ear,

Including the Anatomy and Physiology of the Organ, together with the Treatment of the Affections of the Nose and Pharynx which conduce to Aural Disease (a Treatise). By T. MARK HOVELL, F.R.C.S.E., M.R.C.S., Aural Surgeon to the London Hospital, and Lecturer on Diseases of the Throat in the College, &c. 8vo, with 122 Engravings, 18s. Sore Throat :

Its Nature, Varieties, and Treatment. By PROSSER JAMES, M.D., Physician to the Hospital for Diseases of the Throat. Fifth Edition. Post 8vo, with Coloured Plates and Engravings, 6s. 6d.

Hintson Ophthalmic Out-Patient Practice. By CHARLES HIGGENS, Ophthalmic Surgeon to Guy's Hospital. Third Edition. Fcap. 8vo, 3s.

A System of Dental Surgery. By Sir JOHN TOMES, F.R.S., and C. S. TOMES, M.A., F.R.S. Third Edition. Crown 8vo, with 292 Engravings, 15s.

- Dental Anatomy, Human and Comparative: A Manual. By CHARLES S. TOMES, M.A., F.R.S. Fourth Edition. Crown Svo, with 235 Engravings, 12s. 6d.
- A Manual of Nitrous Oxide Anæsthesia, for the use of Students and General Practitioners. By J. FREDERICK W. SILK, M.D. Lond., M.R.C.S., Anæsthetist to the Royal Free Hospital, Dental School of Guy's Hospital, and National Epileptic Hospital. 8vo, with 26 Engravings, 5s.
- A Practical Treatise on Mechanical Dentistry. By JOSEPH RICH-ARDSON, M.D., D.D.S. Sixth Edition revised and Edited by GEORGE W. WARREN, D.D.S. Roy. 8vo, with 600 Engravings, 21s.

Notes on Dental Practice.

By HENRY C. QUINBY, L.D.S.I., President-Elect of the British Dental Association. Second Edition. 8vo, with 92 Illustrations, 8s.

Elements of Dental Materia Medica and Therapeutics, with Pharmacopœia. By JAMES STOCKEN, L.D.S.R.C.S., Pereira Prizeman for Materia Medica, and THOMAS GADDES, L.D.S. Eng. and Edin. Third Edition. Fcap. 8vo, 7s. 6d.

Papers on Dermatology.

By E. D. MAPOTHER, M.D., Ex-Pres. R.C.S.I. 8vo, 3s. 6d.

Atlas of Skin Diseases.

By TILBURY FOX, M.D., F.R.C.P. With 72 Coloured Plates. Royal 4to, half morocco, £6 6s.

Diseases of the Skin:

LONDON: 11, NEW BURLINGTON STREET.

A Practical Treatise for the Use of Students and Practitioners. By J. N. HYDE, A.M., M.D., Professor of Skin and Venereal Diseases, Rush Medical College, Chicago. Second Edition. 8vo, with 2 Coloured Plates and 96 Engravings, 20s.

Leprosy in British Guiana.

By JOHN D. HILLIS, F.R.C.S., M.R.I.A., Medical Superintendent of the Leper Asylum, British Guiana. Imp. 8vo, with 22 Lithographic Coloured Plates and Wood Engravings, £1 11s. 6d.

10

Diseases of the Skin

(Introduction to the Study of). By P. H. PYE-SMITH, M. D., F. R. S., F.R.C.P., Physician to, and Lecturer or Medicine in, Guy's Hospital. Crown 8vo, with 26 Engravings. 7s. 6d.

Sarcoma and Carcinoma: Their Pathology, Diagnosis, and Treatment. By HENRY T. BUTLIN, F.R.C.S.,

Assistant Surgeon to St. Bartholomew's Hospital. 8vo, with 4 Plates, 8s.

By the same Author.

Malignant Disease of the Larynx (Sarcoma and Carcinoma). 8vo, with 5 Engravings, 5s.

Also.

Operative Surgery of Malignant Disease. 8vo, 14s.

On Cancer:

Its Allies, and other Tumours; their Medical and Surgical Treatment. By F. A. PURCELL, M.D., M.C., Surgeon to the Cancer Hospital, Brompton. Svo, with 21 Engravings, 10s. 6d.

Cancers and the Cancer Process: a Treatise, Practical and Theoretic. By HERBERT L. SNOW, M.D., Surgeon to the Cancer Hospital, Brompton. 8vo, with 15 Lithographic Plates. 15s.

By the same Author.

The Re-appearance(Recurrence) of Cancer after apparent Extirpation. 8vo, 5s. 6d.

Also,

- The Palliative Treatment of Incurable Cancer. Crown Svo, 28, 6d.
- Cancerous Affections of the Skin. (Epithelioma and Rodent Ulcer.) By GEORGE THIN, M.D. Post Svo, with 8 Engravings, 5s.

By the same Author.

- Pathology and Treatment of Ringworm. 8vo, with 21 Engravings, 5s.
- Diagnosis and Treatment of Syphilis. By TOM ROBINSON, M.D., Physician to St. John's Hospital for Diseases of the Skin. Crown 8vo, 3s. 6d.

By the same Author.

Eczema: its Etiology, Pathology, and Treatment. Crown Svo, 3s. 6d.

Also.

Illustrations of Diseases of the Skin and Syphilis, with Remarks. Fasc. I. with 3 Plates. Imp. 4to, 5s. [BY SIR HENRY THOMPSON, F.R.C.S.]

- Diseases of the Urinary Organs. Clinical Lectures. By Sir HENRY THOMPSON, F.R.C.S., Emeritus Professor of Clinical Surgery and Consulting Surgeon to University College Hospital. Eighth Edition. 8vo, with 121 Engravings, 10s. 6d.
- Diseases of the Prostate: Their Pathology and Treatment. Sixth Edition. 8vo, with 39 Engravings, 6s.
- Surgery of the Urinary Organs. Some Important Points connected therewith. Lectures delivered in the R.C.S. 8vo, with 44 Engravings. Student's Edition, 2s. 6d.
- Practical Lithotomy and Lithotrity; or, An Inquiryinto the Best Modes of Removing Stone from the Bladder. Third Edition. 8vo, with 87 Engravings, 105.
- The Preventive Treatment of Calculous Disease, and the Use of Solvent Remedies. Third Edition. Crown 8vo, 2s. 6d.
- Tumours of the Bladder: Their Nature, Symptoms, and Surgical
 - Treatment. 8vo, with numerous Illustrations, 5s.
- Stricture of the Urethra, and Urinary Fistulæ: their Pathology and Treatment. Fourth Edition. 8vo, with 74 Engravings, 6s.
- The Suprapubic Operation of Opening the Bladder for the Stone and for Tumours. 8vo, with 14 Engravings, 3s. 6d.
- Introduction to the Catalogue; being Notes of a Thousand Cases of Calculi of the Bladder removed by the Author and now in the Hunterian Museum of the Royal College of Surgeons. Svo, 2s. 6d.
- Electric Illumination of the Bladder and Urethra, as a Means of Diagnosis of Obscure Vesico-Urethral Diseases. By E. HURRY FENWICK, F.R.C.S., Surgeon to London Hospital and St. Peter's Hospital for Stone. Second Edition. 8vo, with 54 Engravings, 6s. 6d. By the same Author.
- The Cardinal Symptoms of Urinary Diseases: their Diagnostic Significance and Treatment. 8vo, with 36 Illustrations. 8s. 6d.
- Atlas of Electric Cystoscopy. By Dr. EMIL BURCKHARDT, late of the Surgical Clinique of the University of Bâle, and E. HURRY FENWICK, F.R.C.S., Surgeon to the London Hospital and St. Peter's Hospital for Stone. Royal 8vo, with 34 Coloured Plates, embracing 83 Figures. 215.

- Lectures on the Surgical Disorders of the Urinary Organs. By REGINALD HARRISON, F.R.C.S., Surgeon to St. Peter's Hospital. Fourth Edition. 8vo, with 156 Engravings, 16s.
- Clinical Chemistry of Urine (Outlines of the). By C. A. Mac-MUNN, M.A., M.D. 8vo, with 64 Engravings and Plate of Spectra, 9s.
- Urinary and Renal Derangements and Calculous Disorders. By LIONEL S. BEALE, F.R.C.P., F.R.S., Physician to King's College Hospital. 8vo, 5s.
- Male Organs of Generation (Diseases of). By W. H. A. JACOBSON, M.Ch. Oxon., F.R.C.S., Assistant Surgeon to Guy's Hospital. 8vo, with 88 Engravings. 225.
- The Surgical Diseases of the Genito - Urinary Organs, including Syphilis. By E. L. KEVES, M.D., Professor in Bellevue Hospital Medical College, New York (a revision of VAN BUREN and KEVES' Text-book). Roy. 8vo, with 114 Engravings, 218.
- Diseases of the Rectum and Anus. By ALFRED COOPER, F.R.C.S., Senior Surgeon to the St. Mark's Hospital for Fistula; and F. SWINFORD EDWARDS, F.R.C.S., Senior Assistant Surgeon to St. Mark's Hospital. Second Edition, with Illustrations. 8vo, 125.
- Diseases of the Rectum and Anus. By HARRISON CRIPPS, F.R.C.S., Assistant Surgeon to St. Bartholomew's Hospital, &c. Second Edition. 8vo, with 13 Lithographic Plates and numerous Wood Engravings, 12s. 6d.

By the same Author.

Cancer of the Rectum.

- Especially considered with regard to its Surgical Treatment. Jacksonian Prize Essay. 8vo, with 13 Plates and several Wood Engravings, 6s.
- The Diagnosis and Treatment of Diseases of the Rectum. By WILLIAM ALLINGHAM, F.R.C.S., Surgeon to St. Mark's Hospital for Fistula. Fifth Edition. By HERBERT WM. ALLINGHAM, F.R.C.S., Surgeon to the Great Northern Central Hospital, Demonstrator of Anatomy at St. George's Hospital. Svo, with 53 Engravings. 10s. 6d.

The Surgery of the Rectum.

By HENRY SMITH, Emeritus Professor of Surgery in King's College, Consulting Surgeon to the Hospital. Fifth Edition. 8vo, 6s.

- A Medical Vocabulary: An Explanation of all Terms and Phrases used in the various Departments of Medical Science and Practice, their Derivation, Meaning, Application, and Pronunciation. By R. G. MAYNE, M.D., LL.D. Sixth Edition by W. W. WAGSTAFFE, B.A., F.R.C.S. Crown 8vo, 10s. 6d.
- A Short Dictionary of Medical Terms. Being an Abridgment of Mayne's Vocabulary. 64mo, 2s. 6d.
- Dunglison's Dictionary of Medical Science: Containing a full Explanation of its various Subjects and Terms, with their Pronunciation, Accentuation, and Derivation. Twenty-first Edition. By RICHARD J. DUNGLISON, A.M., M.D. Royal 8vo, 30s.
- Terminologia Medica Polyglotta: a Concise International Dic tionary of Medical Terms (French, Latin, English, German, Italian, Spanish, and Russian). By THEODORE MAXWELL, M.D., B.Sc., F.R.C.S. Edin. Royal Svo, 16s.
- A German-English Dictionary of Medical Terms. By FREDERICK TREVES, F.R.C.S., Surgeon to the London Hospital; and HUGO LANG, B.A. Crown 8vo, half-Persian calf, 12s.

Chemistry,

Inorganic and Organic. With Experiments. By CHARLES L. BLOXAM. Seventh Edition, by JOHN MILLAR THOMSON, Professor of Chemistry in King's College, London, and ARTHUR G. BLOXAM, Head of the Chemistry Department, The Goldsmiths' Institute, New Cross. 8vo, with 282 Illustrations, 18s.

By the same Author.

Laboratory Teaching;

Or, Progressive Exercises in Practical Chemistry. Sixth Edition. By ARTHUR G. BLOXAM. Crown 8vo, with 80 Engravings, 6s. 6d.

Practical Chemistry

And Qualitative Analysis. By FRANK CLOWES, D.Sc. Lond., Professor of Chemistry in the University College, Nottingham. Fifth Edition. Post 8vo, with 57 Engravings and Frontispiece, 7s. 6d.

Quantitative Analysis.

By FRANK CLOWES, D.Sc. Lond., Professor of Chemistry in the University College, Nottingham, and J. BERNARD COLEMAN, Assoc. R. C. Sci. Dublin; Senior Demonstrator of Chemistry in the University College, Nottingham. Second Edition. Post 8vo, with 94 Engravings, 8s. 6d.

By the same Authors.

Elementary Qualitative Analysis. With Engravings. Post 8vo, 2s. 6d.

Watts' Manual of Chemistry,

Theoretical and Practical. By WILLIAM A. TILDEN, D.Sc., F.R.S., Professor of Chemistry in the Mason College, Birmingham.

- PHYSICAL AND INORGANIC CHE-MISTRY. Second Edition. Crown 8vo, with Coloured Plate of Spectra, and 122 Wood Engravings, 8s. 6d.
- CHEMISTRY OF CARBON COMPOUNDS; or, ORGANIC CHEMISTRY. Second Edition. Crown 8vo, with Engravings, 10s.

Qualitative Analysis.

By R. FRESENIUS. Translated by CHARLES E. GROVES, F.R.S. Tenth Edition. 8vo, with Coloured Plate of Spectra and 46 Engravings, 15s.

By the same Author.

Quantitative Analysis.

Seventh Edition.

- Vol. I., Translated by A. VACHER. 8vo, with 106 Engravings, 15s.
- Vol. II., Parts 1 to 3, Translated by C. E. GROVES, F.R.S. 8vo, with Engravings, 2s. 6d. each.

Practical Chemistry,

Including Analysis. By John E. BOW-MAN and CHARLES L. BLOXAM. Fcap. 8vo. Eighth Edition, with 90 Engravings, 5s. 6d.

Inorganic Chemistry.

By EDWARD FRANKLAND, Ph.D., D.C.L., LL.D., F.R.S., Professor of Chemistry in the Normal School of Science, and FRANCIS R. JAPP, M.A., Ph.D. F.I.C., F.R.S., Professor of Chemistry in the University of Aberdeen. Svo, with numerous Illustrations on Stone and Wood, 24s.

Inorganic Chemistry

(A System of). By WILLIAM RAMSEY, Ph.D., F.R.S., Professor of Chemistry in University College, London. 8vo, with Engravings, 15s.

By the same Author.

Elementary Systematic Chemistry for the Use of Schools and Colleges. With Engravings. Crown 8vo, 4s. 6d.; Interleaved, 5s. 6d.

Organic Chemistry:

(Outlines of). By H. FORSTER MORLEY, M.A., D.Sc., Joint Editor of Watts' "Dictionary of Chemistry." Crown 8vo, 7s. 6d. Valentin's Qualitative Chemical Analysis. Eighth Edition. By W. R. HODGKINSON, Ph.D., F.R.S.E., Professor of Chemistry and Physics in the Royal Military Academy, and Artillery College, Woolwich. 8vo, with Engravings and Map of Spectra, 8s. 6d.

Analytical Chemistry.

Notes for Students in Medicine. By ALBERT J. BERNAYS, Ph.D., F.C.S., F.I.C., late Professor of Chemistry, &c., at St. Thomas's Hospital Medical School. Third Edition. Crown Svo, 4s. 6d.

Volumetric Analysis:

(A Systematic Handbook of); or theQuantitative Estimation of Chemical Substances by Measure, applied to Liquids, Solids, and Gases. By FRANCIS SUTTON, F.C.S., F.I.C., Public Analyst for the County of Norfolk. Sixth Edition. 8vo, with 102 Engravings, 17s. 6d.

Commercial Organic Analysis: A Treatise on the Properties, Modes of Assaying, Proximate Analytical Examination, &c., of the various Organic Chemicals and Products employed in the Arts, Manufactures, Medicine, &c. By ALFRED H. ALLEN, F.I.C., F.C.S., Public Analyst for the West Riding of Yorkshire, the

- Northern Division of Derbyshire, &c. Vol. I.—Alcohols, Neutral Alcoholic Derivatives, Sugars, Starch and its Isomers, Vegetable Acids, &c. With Illustrations. Third Edition. Svo. [Preparing.
 - Vol. II.—Fixed Oils and Fats, Hydrocarbons, Phenols, &c. With Illustrations. Third Edition. 8vo.

[Preparing.

Vol. III.—Part I. Aromatic Acids, Tannins, Dyes, and Colouring Matters. Second Edition, 8vo, 148.

> Part II. Amines and Ammonium Bases, Hydrazines, Bases from Tar, Vegetable Alkaloids. Second Edition. 8vo, 18s.

Cooley's Cyclopædia

of Practical Receipts, and Collateral Information in the Arts, Manufactures, Professions, and Trades : Including Medicine, Pharmacy, Hygiene and Domestic Economy. Seventh Edition, by W. NORTH, M.A. Camb., F.C.S. 2 Vols., Roy. 8vo, with 371 Engravings, 42s.

Chemical Technology:

A Manual. By RUDOLF VON WAGNER. Translated and Edited by WILLIAM CROOKES, F.R.S., from the Thirteenth Enlarged German Edition as remodelled by Dr. FERDINAND FISCHER. 8vo, with 596 Engravings, 32s.

Fuel and its Applications.

By E. J. MILLS, D.Sc., F.R.S., and F. J. ROWAN, C.E. Being Vol. I. of Chemical Technology, or Chemistry in its Application to Arts and Manufactures. Edited by CHARLES E. GROVES, F.R.S., and WILLIAM THORP, B.Sc. Royal 8vo, with 606 Engravings, 30s.

Potable Waters:

Their Organic Analysis. By J. A. BLAIR, M.B., C.M., D.Sc., L.R.C.P. Second Edition. Crown 8vo, 3s. 6d.

Technological Handbooks.

- EDITED BY JOHN GARDNER, F.I.C., F.C.S., and JAMES CAMERON, F.I.C.
 - BREWING, DISTILLING, AND WINE MANUFACTURE. Crown Svo, with Engravings, 6s. 6d.
 - BLEACHING, DYEING, AND CALICO PRINTING. With Formulæ. Crown 8vo, with Engravings, 5s. OILS, RESINS, AND VARNISHES.
 - OILS, RESINS, AND VARNISHES. Crown 8vo, with Engravings. 7s. 6d. SOAPS AND CANDLES. Crown 8vo, with 54 Engravings, 7s.
- The Microscope and its Revelations. By the late WILLIAM B. CAR-PENTER, C.B., M.D., LL.D., F.R.S. Seventh Edition, by the Rev. W. H. DALLINGER, LL.D., F.R.S. With 21 Plates and 800 Wood Engravings. 8vo, 26s. Half Calf, 30s.
- The Quarterly Journal of Microscopical Science. Edited by E. RAY LANKESTER, M.A., LL.D., F.R.S.; with the co-operation of ADAM SEDGWICK, M.A., F.R.S., and W. F. R. WELDON, M.A., F.R.S. Each Number, 105.
- Methods and Formulæ

Used in the Preparation of Animal and Vegetable Tissues for Microscopical Examination, including the Staining of Bacteria. By PETER WYATT SQUIRE, F.L.S. Crown 8vo, 3s. 6d.

The Microtomist's Vade-Mecum: A Handbook of the Methods of Microscopic Anatomy. By ARTHUR BOLLES LEE, Assistant in the Russian Laboratory of Zoology at Villefranche-sur-mer (Nice). Third Edition. 8vo, 14s.

Photo-Micrography

(Guide to the Science of). By EDWARD C. BOUSFIELD, L.R.C.P. Lond. Svo, with 34 Engravings and Frontispiece, 6s.

- An Introduction to Physical Measurements, with Appendices on Absolute Electrical Measurements, &c. By Dr. F. KOHLRAUSCH, Professor at the University of Strassburg. Third Edition, translated from the Seventh German Edition, by THOMAS HUT-CHINSON WALLER, B.A., B.Sc., and HENRY RICHARDSON PROCTER, F.I.C., F.C.S. 8vo, with 91 Illustrations, 12s. 6d.
- The Principles and Practice of Veterinary Medicine. By WILLIAM WILLIAMS, F.R.C.V.S., F.R.S.E., Principal, and Professor of Veterinary Medicine and Surgery at the New Veterinary College, Edinburgh. Seventh Edition. 8vo, with several Coloured Plates and Woodcuts, 30s.

By the same Author.

- The Principles and Practice of Veterinary Surgery. Eighth Edition. 8vo, with 9 Plates and 147 Woodcuts, 305.
- The Veterinarian's Pocket Remembrancer: being Concise Directions for the Treatment of Urgent or Rare Cases, embracing Semeiology, Diagnosis, Prognosis, Surgery, Therapeutics, Toxicology, Detection of Poisons by their Appropriate Tests, Hygiene, &c. By GEORGE ARMATAGE, M.R.C.V.S. Second Edition. Post 8vo, 3s.

Chauveau's Comparative Anatomy of the Domesticated Animals. Revised and Enlarged, with the Co-operation of S. ARLOING, Director of the Lyons Veterinary School, and Edited by GEORGE FLEMING, C.B., LL.D., F.R.C.V.S., late Principal Veterinary Surgeon of the British Army. Second English Edition. 8vo, with 585 Engravings, 31s. 6d.

INDEX TO J. & A. CHURCHILL'S LIST.

Abercrombie's Medical Jurisprudence, 2 Adams (W.) on Clubfoot, 9 on Contractions of the Fingers, &c., 9 Adams (W.) on Clubfoot, 9 on Contractions of the Fingers, &c., 9 Allen's Commercial Organic Analysis, 13 Allingham (H.) on Derangements of Knee-joint, 9 Allingham (W.) on Diseases of the Rectum, 12 Armatage's Veterinary Pocket Remembrancer, 14 Auld's Bronchial Affections, 6 Barnes (R.) on Obstetric Operations, 3 — on Diseases of Women, 3 Beale (L. S.) on Liver, 6 — Microscope in Medicine, 6 — Slight Ailments, 6 — Urinary and Renal Derangements, 12 Beale (P. T. B.) on Elementary Biology, 2 Beasley's Book of Prescriptions, 5 — Druggists' General Receipt Book, 5 — Pocket Formulary, 5 Bellamy's Surgical Anatomy, 1 Bentley and Trimen's Medicinal Plants, 5 Bentley's Structural Botany, 5 Berkart's Bronchial Asthma, 6 Bernard on Stammering, 7 Bernay's Notes on Analytical Chemistry, 1 Bigg's Short Manual of Orthopædy, 9 Blair's Potable Waters, 14 Bloxam's Chemistry, 12 — Laboratory Teaching, 12 Bowsfield's Photo-Micrography, 14 Bowman and Bloxam's Practical Chemistry, 13 Brodhurst's Anchylosis, 9 — Curvatures, &c., of the Spine, 9 — Tainare Formical Chemistry, 13 Brodhurst's Anchylosis, 9 — Curvatures, &c., of the Spine, 9 Elementary Analysis, 12 Clowes' Practical Chemistry, 12 Cooley's Cyclopædia of Practical Receipts, 13 Cooper and Edwards' Diseases of the Rectum, 12 Cripps' (H.) Cancer of the Rectum, 12 Diseases of the Rectum and Anus, 12 Cripps' (R. A.) Galenic Pharmacy, 4 Cullingworth's Manual of Nursing, 4 Short Manual for Monthly Nurses, 4 Dalby's Diseases and Injuries of the Ear, 10 Short Contributions, 10 Day on Diseases of Children, 4 Day on Diseases of Children, 4 — on Headaches. 8 Domville's Manual for Nurses, 4 Doran's Gynaecological Operations, 3 Down's Mental Affections of Childhood, 3 Druitt's Surgeon's Vade-Mecum, 8 Duncan (A.), on Prevention of Disease in Tropics, 5 Duncan (J. M.), on Diseases of Women, 3 Dunglison's Dictionary of Medical Science, 12 Ellis's (E.) Diseases of Children, 4 Ellis's (T. S.) Human Foot, 9 Ewart's Bronchi and Pulmonary Blood Vessels, 6 Fagge's Principles and Practice of Medicine, 6 Fayrer's Climate and Fevers of India, 5 Day on Diseases of Children, 4

Minor Surgery and Bandaging, 8 Operative Surgery, 8 Practical Anatomy, 1 Surgical Diagnosis, 8 Hellier's Notes on Gynæcological Nursing, 4 Higgens' Ophthalmic Out-patient Practice, 10 Hillis' Leprosy in British Guiana, 10 Hirschfeld's Atlas of Central Nervous System 2 Holden's Human Osteology, 1 Mapother's Papers on Dermatology, 10 Martin's Ambulance Lectures, 8 Maxwell's Terminologia Medica Polyglotta, 12 Mayne's Medical Vocabulary, 12 Mercier's Lunacy Law, 3 Microscopical Journal, 14 Mills and Rowan's Fuel and its Applications, 14 Moore's (N.) Pathological Anatomy of Diseases, 1 Moore's (Sir W. J.) Family Medicine for India, 5 Manual of the Diseases of India, 5 [Continued on the next page.

INDEX TO J. & A. CHURCHILL'S LIST-continued.

Moore's (Sir W. J.) Tropical Climates, 5 Morley's Organic Chemistry, 13 Morris's Human Anatomy, 1 Moullin's (Mansell) Surgery, 8 Nettleship's Diseases of the Eye, 9 Ogle on Puncturing the Abdomen, 8 Oliver's Diseases of Women, 3 Ophthalmic (Royal London) Hospital Reports, 9 Ormerod's Diseases of the Nervous System, 7 Owen's Materia Medica, 4 Parkes' Practical Hygiene, 2 Pavy's Carbohydrates, 6 Pereira's Selecta è Prescripts, 4 Phillips' Materia Medica and Therapeutics, 4 Pollock's Histology of the Eye and Eyelids, 9 Proctor's Practical Pharmacy, 4 Purcell on Cancer, 11 Ross's Aphasia, 7 — Diseases of the Nervous System, 7 Royle and Harley's Materia Medica, 5 St. Thomas's Hospital Reports, 7 Sansom's Valvular Disease of the Heart, 6 Bavage's Female Pelvic Organs, 3 Schetelig's Homburg-Spa, 8 Schweinitz's (G. E. de) Diseases of Eye, 10 Short Dictionary of Medical Terms, 12 Silk's Manual of Nitrous Oxide, 10 Smith's (E.) Clinical Studies, 4

Stevenson and Murphy's Hygiene, 2

Stillé and Maisch's National Dispensatory, 5 Stocken's Dental Materia Medica and Therapeutics, 10 Sutton's (H. G.), Lectures on Pathology, 1 Sutton's (J. B.), General Pathology, 1 Sutton's (J. B.), General Pathology, 1 Sutton's (F.) Volumetric Analysis, 13 Swaine's Surgical Emergencies, 8 Swayne's Obstetric Aphorisms, 3 Taylor's (A. S.) Medical Jurisprudence, 2 Taylor's (J. C.), Canary Islands, 8 Thin's Cancerous Affections of the Skin, 11 — Pathology and Treatment of Ringworm, 11 Thompson's (Sir H.) Calculous Disease, 11 — Diseases of the Prostate, 11 — Diseases of the VrinaryOrgans, 11 Introduction to Catalogue, 11 Lithotomy and Lithotrity, 11 Suprapubic Operation, 11 Suprapubic Operation, 11 Surgery of the Urinary Organs, 11 Tumours of the Bladder, 11 Tirard's Prescriber's Pharmacopœia, 5 Tomes' (C. S.) Dental Anatomy, 10 Tomes' (J. and C. S.) Dental Surgery, 10 Tommasi-Crudeli's Climate of Rome, 7 Prichard and Symonds and Mental Science, 3 Reform in the Treatment of the Insane, 3 Valentin and Hodgkinson's Qualitative Analysis, 13 Vintras on the Mineral Waters, &c., of France, 8

N.B.-J. & A. Churchill's larger Catalogue of about 600 works on Anatomy, Physiology, Hygiene, Midwifery, Materia Medica, Medicine, Surgery, Chemistry, Botany, &c. &c., with a complete Index to their Subjects, for easy reference, will be forwarded post free on application.

AMERICA.- J. & A. Churchill being in constant communication with various publishing houses in America are able to conduct negotiations favourable to English Authors.



Date Due			
Call Suger			
	and the		
See Se			
Sale Car			
			a start and a start
		a la contra	
and the second			Contract 24
et a strange			Carlo Carlo
a state			
Contraction of the			P.S. STAN
No.	The States		
an said			
		143 24	
and the second			
11/2 2013		and the second second	The states
Demco 293-5			



