

A practical treatise on the diseases of children : By J. Forsyth Meigs and William Pepper.

Contributors

Meigs, J. Forsyth 1818-1882.

Pepper, William, 1843-1898.

Harvey Cushing/John Hay Whitney Medical Library

Publication/Creation

Philadelphia : P. Blakiston, 1886.

Persistent URL

<https://wellcomecollection.org/works/jt64azcz>

License and attribution


This material has been provided by This material has been provided by the Harvey Cushing/John Hay Whitney Medical Library at Yale University, through the Medical Heritage Library. The original may be consulted at the Harvey Cushing/John Hay Whitney Medical Library at Yale University. where the originals may be consulted.


This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>





A
PRACTICAL TREATISE
ON THE
DISEASES OF CHILDREN.

DISSEMINATION OF CHILDREN

PHYSIOLOGICAL TREATMENT

A

PRACTICAL TREATISE
ON THE
DISEASES OF CHILDREN.

BY

J. FORSYTH MEIGS, M.D.,

LATE ONE OF THE PHYSICIANS TO THE PENNSYLVANIA HOSPITAL; CONSULTING PHYSICIAN TO THE
CHILDREN'S HOSPITAL; FELLOW OF THE COLLEGE OF PHYSICIANS OF PHILADELPHIA;
MEMBER OF THE AMERICAN PHILOSOPHICAL SOCIETY, OF THE ACADEMY
OF NATURAL SCIENCES OF PHILADELPHIA, ETC., ETC.

AND

WILLIAM PEPPER, M.D., LL.D.,

PROVOST, AND PROFESSOR OF THE THEORY AND PRACTICE OF MEDICINE AND OF CLINICAL
MEDICINE IN THE UNIVERSITY OF PENNSYLVANIA; PHYSICIAN TO THE UNIVERSITY,
TO THE PHILADELPHIA, AND TO THE CHILDREN'S HOSPITALS; FELLOW OF THE
COLLEGE OF PHYSICIANS OF PHILADELPHIA, ETC., ETC.

SEVENTH EDITION,

REVISED AND ENLARGED.

PHILADELPHIA:
P. BLAKISTON, SON & CO.,
1012 WALNUT STREET.

1886.

Entered according to Act of Congress, in the year 1882,

By P. BLAKISTON, SON & CO.,

In the Office of the Librarian of Congress, at Washington, D. C.

RJ 45
886M

TO THE LATE

GEORGE B. WOOD, M.D., LL.D.,

*President of the College of Physicians of Philadelphia; Emeritus Professor of the Theory and Practice
of Medicine in the University of Pennsylvania; late one of the Physicians to the
Pennsylvania Hospital, &c., &c.,*

This Work is Dedicated,

AS

A TRIBUTE OF RESPECT FOR HIS HIGH PROFESSIONAL ATTAINMENTS

AND

EMINENT PRIVATE VIRTUES,

AND AS


A MARK OF GRATITUDE FOR HIS VALUABLE INSTRUCTIONS,

BY

THE AUTHORS,

J. FORSYTH MEIGS,

WILLIAM PEPPER.



PREFACE TO THE SEVENTH EDITION.

IN preparing the seventh edition of this work for the press, the entire text has been subjected to a thorough revision. All the statistics have been brought up to date, and the data of recent years have been used in calculating new tables, as in the case of the elaborate table on the relative mortality of croup and diphtheria. A short article on Röteln has been added; and the Section on Skin Diseases has been rearranged and in large part rewritten. The General Diseases have been classified more in accordance with our knowledge of their pathology. With the object of maintaining the position of this work as a safe and practical guide in the treatment of the diseases of children, the remarks upon the management of each affection have been revised with especial care, so as to embody the recent results of other observers as well as of our own experience. The great importance of the subject of Food, and the large share of attention it has of late received, have led us to rewrite the article on Thrush and to add a new article on Food, in which the subject of condensed milk is carefully considered. Altogether it is hoped that the work will be found to merit a continuance of the favorable recognition hitherto extended to it by the profession.

PREFACE TO THE FOURTH EDITION.

It has been some years since the third edition of *Meigs on the Diseases of Children* has been exhausted; and the frequent inquiries which have been made for the work, as well as the increasing interest taken by the profession in the study of the diseases of childhood, have led to the belief that the publication of a new edition would be received with the same kind favor which has been already extended to the three former ones.

The changes and additions which were necessitated by the great advance made during the last decade in our knowledge of a number of the diseases of children, as well as by the unavoidable omission of any consideration of several important subjects in the previous editions of this work, were, however, of so extensive a character that it has been found necessary to associate a collaborator in the preparation of the present edition.

Among the principal of these changes may be mentioned the great enlargement of several articles, and especially of those on thrush, convulsions, chorea, tracheotomy in croup, and parasitic skin diseases. Other articles have been entirely rearranged, or even rewritten, as those upon the diseases of the stomach and intestines, and upon eczematous affections. In addition to such changes, however, there have been no less than seventeen full articles added, embracing the following important subjects: Diseases of the Heart, and Cyanosis; Diseases of the Cæcum and Appendix Vermiformis, and Intussusception; Chronic Hydrocephalus, Tetanus, Atrophic Infantile Paralysis, Facial Paralysis, and Progressive Paralysis with Apparent Hypertrophy of the Muscles; Rheumatism, Diphtheria, Mumps, Rickets, Tuberculosis, and Infantile Syphilis; Typhoid Fever; and Sclerema. These various additions and changes have involved the introduction of more than two hundred pages of new matter. Several extensive tables, exhibiting the mortality in this city of some of the most common and fatal diseases, in connection with the variations of temperature, have been prepared with great care from the records of the office of the Board of Health, which were opened to examination through the courtesy of Mr. Chambers, the Chief Registration Clerk of that office. A copious index has also been supplied, which it is trusted will facilitate reference, and render the work more practically serviceable.

Apart from these changes, however, no alteration has been made in the general plan of the work. As in the composition of the previous editions,

the best and most recent foreign and domestic authorities on the diseases of children have been frequently and carefully consulted, and their views fully quoted whenever they appeared of practical importance. For the most part, however, the opinions expressed in the following pages are those to which the authors have been led by their personal observation, and which they, therefore, believe to have been approved by the most searching of all tests, that of practical application.

It has also been their constant aim, while supplying a sufficient amount of information upon questions of etiology, pathology, and morbid anatomy, to insure a practical character to the work. With this view, an unusual amount of space has been devoted to the discussion of the treatment of the different diseases, and in every instance the conclusions derived by the authors from their own experience have been fully and, it is hoped, clearly stated.

In so doing, it has been necessary to consider somewhat at length the extremely important questions of the employment of venesection, antimony, calomel, and stimulants; and a full expression of opinion upon each of these points will be found in its appropriate place.

In conclusion, the authors would venture to express the hope that their efforts may have been successful in furnishing a work which will aid in rendering the study of the diseases of children more attractive and clear, their recognition more easy, and which may serve as a practical guide in the difficult task of treating these disorders.

•

TABLE OF CONTENTS.

PREFACE TO THE SEVENTH EDITION,	PAGE vii
PREFACE TO THE FOURTH EDITION,	ix
INTRODUCTORY ESSAY,	17

CLASS I.

DISEASES OF THE RESPIRATORY ORGANS.

CHAPTER I.

DISEASES OF THE UPPER AIR-PASSAGES.

SECTION I.

DISEASES OF THE NASAL PASSAGES.

ARTICLE I. Coryza,	52
------------------------------	----

SECTION II.

DISEASES OF THE LARYNX.

GENERAL REMARKS,	61
ARTICLE I. Simple laryngitis without spasm,	62
“ II. Spasmodic simple laryngitis,	69
“ III. Pseudo-membranous laryngitis,	85

CHAPTER II.

DISEASES OF THE LUNGS AND PLEURA.

GENERAL REMARKS,	133
ARTICLE I. Atelectasis pulmonum,	134
“ II. Pneumonia,	157
“ III. Bronchitis,	195
“ IV. Emphysema,	218
“ V. Pleurisy,	231
“ VI. Pneumothorax,	252
“ VII. Hooping-cough,	259

CLASS II.

DISEASES OF THE CIRCULATORY ORGANS.

	PAGE
ARTICLE I. Cyanosis,	280
“ II. Diseases of the Heart,	284

CLASS III.

DISEASES OF THE DIGESTIVE ORGANS.

CHAPTER I.

DISEASES OF THE MOUTH AND THROAT.

ARTICLE I. Food,	300
“ II. Simple or erythematous stomatitis,	329
“ III. Aphthæ,	330
“ IV. Ulcerative or ulcero-membranous stomatitis,	332
“ V. Gangrene of the mouth,	336
“ VI. Thrush,	347
“ VII. Affections of the tonsils,	364
“ VIII. Simple or erythematous pharyngitis,	368
“ IX. Retropharyngeal abscess,	373

CHAPTER II.

DISEASES OF THE STOMACH AND INTESTINES.

GENERAL REMARKS,	376
----------------------------	-----

SECTION I.

FUNCTIONAL DISEASES OR MILD CATARRH OF THE STOMACH AND INTESTINES.

ARTICLE I. Indigestion,	376
“ II. Simple diarrhœa,	387

SECTION II.

DISEASES OF THE STOMACH AND INTESTINES, ATTENDED WITH APPRECIABLE ANATOMICAL LESIONS.

ARTICLE I. Gastritis,	397
“ II. Entero-colitis, or inflammatory diarrhœa,	404

	PAGE
ARTICLE III. Cholera infantum,	441
“ IV. Dysentery,	462
“ V. Diseases of the cœcum and appendix cœci,	467
“ VI. Intussusception,	481

CLASS IV.

DISEASES OF THE NERVOUS SYSTEM.

GENERAL REMARKS,	498
ARTICLE I. Tubercular meningitis,	498
“ II. Simple meningitis,	529
“ III. Cerebral congestion,	537
“ IV. Cerebral hemorrhage,	540
“ V. Chronic hydrocephalus,	548
“ VI. General convulsions or eclampsia,	559
“ VII. Laryngismus stridulus,	577
“ VIII. Contraction with rigidity,	593
“ IX. Tetanus nascentium,	602
“ X. Chorea,	610
“ XI. Atrophic infantile paralysis,	634
“ XII. Facial paralysis,	650
“ XIII. Progressive muscular sclerosis, or pseudo-hypertrophic muscular paralysis,	652
“ XIV. Night terrors,	660

CLASS V.

GENERAL DISEASES RESULTING FROM DERANGEMENTS OF THE NORMAL PROCESSES OF NUTRITION.

INTRODUCTORY REMARKS,	665
ARTICLE I. Rheumatism,	666
“ II. Scrofula,	673
“ III. Tuberculosis,	679
“ IV. Rickets,	694
“ V. Congenital syphilis,	706

CLASS VI.

GENERAL DISEASES RESULTING FROM SPECIAL MORBID AGENTS
OPERATING FROM WITHOUT.

	PAGE
ARTICLE I. Typhoid fever,	715
“ II. Variola and Varioloid,	729
“ III. Vaccinia,	751
“ IV. Varicella,	766
“ V. Scarlatina,	769
“ VI. Rubeola,	831
“ VII. Rötheln,	855
“ VIII. Malarial fever,	857
“ IX. Mumps,	861
“ X. Erysipelas,	866
“ XI. Diphtheria,	873
“ XII. Epidemic cerebro-spinal meningitis,	909

CLASS VII.

DISEASES OF THE SKIN.

INTRODUCTORY REMARKS,	918
---------------------------------	-----

CHAPTER I.

ERYTHEMATOUS AFFECTIONS.

ARTICLE I. Erythema,	920
“ II. Roseola,	925
“ III. Urticaria,	928

CHAPTER II.

VESICULAR OR CATARRHAL INFLAMMATION OF THE SKIN.

ARTICLE I. Eczema,	933
“ II. Herpes,	949
“ III. Miliaria,	955

CHAPTER III.

BULLOUS INFLAMMATION OF THE SKIN.

ARTICLE I. Pemphigus,	956
“ II. Rupia,	959

CHAPTER IV.

PUSTULAR OR SUPPURATIVE INFLAMMATION OF THE SKIN.

	PAGE.
ARTICLE I. Ecthyma,	962
“ II. Impetigo contagiosa,	965

CHAPTER V.

PAPULAR OR PLASTIC INFLAMMATION OF THE SKIN.

ARTICLE I. Lichen,	966
“ II. Prurigo,	969

CHAPTER VI.

SQUAMOUS INFLAMMATION OF THE SKIN.

ARTICLE I. Psoriasis,	971
“ II. Pityriasis rubra,	974

CHAPTER VII.

HYPERTROPHIC DISEASES OF THE SKIN.

ARTICLE I. Ichthyosis,	975
“ II. Sclerema,	977

CHAPTER VIII.

PARASITIC DISEASES OF THE SKIN.

GENERAL REMARKS,	980
ARTICLE I. Favus,	983
“ II. Tinea,	990
“ III. Alopecia areata,	996
“ IV. Scabies,	997

CLASS VIII.

WORMS IN THE ALIMENTARY CANAL.

GENERAL REMARKS,	1003
ARTICLE I. Ascaris lumbricoides,	1007
“ II. Ascaris vermicularis,	1019

CHAPTER IV

THE HISTORY OF THE DISCOVERY OF THE GREAT

THE HISTORY OF THE DISCOVERY OF THE GREAT

CHAPTER V

THE HISTORY OF THE DISCOVERY OF THE GREAT

THE HISTORY OF THE DISCOVERY OF THE GREAT

CHAPTER VI

THE HISTORY OF THE DISCOVERY OF THE GREAT

THE HISTORY OF THE DISCOVERY OF THE GREAT

CHAPTER VII

THE HISTORY OF THE DISCOVERY OF THE GREAT

THE HISTORY OF THE DISCOVERY OF THE GREAT

CHAPTER VIII

THE HISTORY OF THE DISCOVERY OF THE GREAT

THE HISTORY OF THE DISCOVERY OF THE GREAT

CHAPTER IX

THE HISTORY OF THE DISCOVERY OF THE GREAT

THE HISTORY OF THE DISCOVERY OF THE GREAT

CHAPTER X

THE HISTORY OF THE DISCOVERY OF THE GREAT

THE HISTORY OF THE DISCOVERY OF THE GREAT

A
PRACTICAL TREATISE
ON THE
DISEASES OF CHILDREN.

INTRODUCTORY ESSAY.

ON THE CLINICAL EXAMINATION OF CHILDREN.

THE clinical examination of children, and particularly of young infants, cannot be successfully practiced upon the same method as that habitually made use of in the case of adults. The truth of this statement will be readily assented to by all who have had much experience in the treatment of the diseases of the two ages, by those who will reflect for a moment on the great differences in the expressions of the various organs in early and adult life, and by those who are acquainted with the opinions of distinguished writers upon children's diseases. It is proper and useful, therefore, to preface a practical work on the diseases of children, with a sketch or plan of the best method to be pursued in forming a diagnosis of these diseases, and with remarks upon the physiological characters which distinguish the organization of early life from that of maturity.

The difficulties that beset the path of the practitioner in his clinical examination of children are so great that he who has not been prepared by preliminary study to surmount these obstacles, will find it a most uncertain and dubious task to unravel the history and nature of any case that may be set before him. The helpless silence of the infant,—the wilful silence, or the loose and inconsistent answers of the older child, which lead astray the mind rather than guide it to true results,—the agitation and fright produced by the examination, rendering it impossible at times to ascertain the real state of the different functions of the economy,—and lastly, the difficulty of obtaining accurate and reliable accounts of the history of the case from the attendants, all combine to make the duty of the physician most perplexing, and, unless he be gifted with a large share of patient and philosophic calmness, most irksome and trying to the temper.

So great, indeed, are the difficulties encountered by some practitioners who enter upon this branch of the practice of medicine without proper preliminary preparation, that they never overcome them; but, to use the words of Dr. West, "grow satisfied with their ignorance, and will then, with the greatest gravity, assure you that the attempt to understand these affections is useless." That it is possible, in great measure, to overcome these obstacles, and to arrive at a correct diagnosis in nearly all cases, is quite as true as that these obstacles really exist. But, in order to do this, the physician must first be aware that difficulties exist, and must have formed in his mind some plan or method by which to surmount or elude them.

Before proceeding to show what is the best method of examining or exploring disease in children, we must state that our remarks apply chiefly to infants and very young subjects; for, after the age of eight or ten years, the physical and intellectual development have progressed to such a point as to render the method of diagnosis nearly the same as that employed in adults.

The chief causes which render the diagnosis of disease in young children difficult, are the absence of the faculty of speech, and the violent agitation generally caused by the examination, which prevents a proper appreciation of the state of certain organs and functions.

It is easy to understand how much our means of diagnosis are restricted by the absence of the faculty of speech. How many symptoms there are in the case of adults with which we become acquainted only through the patient's own account of his sensations; and, consequently of how many must we be deprived in children by the absence of this account. It might, at first view, seem impossible to detect the nature of the sickness without the assistance of this means, so greatly do we depend upon it in our examination of adults. Nevertheless, we shall find ourselves enabled, by an attentive consideration of other resources in the child, by a close study of its physiognomical expression, its decubitus, the nature of its cry, and by the most rigidly careful physical examination, to form our conclusions with almost as great a degree of precision as in older patients.

The other causes of difficulty,—the violent disturbance, both physical and moral, of the child, its fright, agitation, and cries,—constitute, when they are present in a high degree, much greater embarrassments than the want of speech. To overcome these, the physician must use all his skill, tact, and patience; for, unless they can be avoided by art, or overcome by soothing and gentle persuasion, he can learn but little that will be of essential service to him in making up his opinion. He can neither read the countenance of the little patient, nor judge by its attitudes or decubitus of the state of the various organs, whether internal or external; he will be unable to ascertain the rate, force, or regularity of the circulatory or respiratory functions; he cannot, to any useful purpose, examine the abdomen, to learn whether it be tender on pressure, or whether its contained organs be in their natural condition as to size and position; and lastly, he will find that the physical exploration of the lungs and heart, by auscultation and percussion, yield him at best only imperfect results.

To avoid the difficulties just detailed, it is always useful, if not absolutely necessary, to conduct certain portions of the examination whilst the child is calm and quiet, and certain others whilst it is disturbed and agitated. This distinction of the examination into two periods, or stages, is one of the utmost importance in a practical point of view, and should never be forgotten by the physician during his clinical observation of the various symptoms the patient may present.

By the period of calm is meant a condition of total quiescence, in which the child is undisturbed either by internal or external causes of irritation. This condition is best found in the state of sleep. If this cannot be obtained, the one most nearly approaching to it is that which exists during the act of nursing, or which follows that act. Suckling is usually followed, even in the sick child, by a condition of drowsiness or by a gentle and languid slumber, during which it will allow a careful examination upon many points without agitation. If possible, therefore, the physician should always see the child when asleep, and if the mother or nurse propose, on the occasion of his visit, to hurry upstairs to prepare the child, or to bring it down into the parlor or lower room, he should ask, as a favor, that he may see it asleep.

If, in spite of having just been nursed, the child be awake and fretting, and when, also, it is more advanced in age, we should endeavor, by the attraction of toys, by gentle and soothing words and manners, by fondling or by having it carried about the room, to get it quiet.

Before proceeding to a consideration of the particular means by which we are to judge of the state of health or sickness of young subjects, it is proper to call attention to the great importance of a careful examination of the attendants, in regard to the history of the case, previous to and between the medical visits. In the instance of children, their inability to describe their own symptoms compels us to depend entirely upon the mother or nurse for all detail of the case previous to our first visit, and for all accounts of what may have happened between two subsequent ones. It is, therefore, extremely important that this part of the examination should be conducted with every care and caution. Very much that is useful may be learned from it, if it be well managed. A great deal of skill and art are required in putting the questions, and in sifting the evidence thus collected. We should always bear in mind the character of the persons questioned. Much depends upon their education, and much more on their natural powers of observation, and manner of relating what they may have seen. The degree of credence to be attached to their answers must rest upon their probable intelligence. Nurses and mothers will often give accounts of their charges which must be received with large allowance, and even in some few instances with disbelief. We would, however, in this place, most earnestly caution the young practitioner of medicine to be very careful not to misbelieve, or even mistrust, without well-poised reasons, the account of a sick child given by a mother; for though a foolish, weak woman will often give a false or exaggerated statement of the symptoms of her child an observant and intelligent, and sometimes a foolish and weak one, when guided by maternal instinct, will detect variations from the healthful con-

dition of the child, which may entirely escape the search of the most acute and rigorous medical observer. A mother may perceive a change in the expression of the face, in the manner of the muscular movements, in the temper or conduct of her child, which shall fail to attract the attention of the practitioner; or it may be that the symptom which has caused the parent to take alarm occurs only during the absence of the physician. The medical attendant ought, for these reasons, to listen patiently and kindly to whatever the mother or nurse may have to say, and if unable to detect immediately what they assert they have seen, let him not determine at once that there has been a mistake, that their anxiety has deceived them; but let him examine the patient yet again, and more carefully, or let him pay another visit to learn whether the symptom or symptoms continue, or have occurred again. Our own rule, in a doubtful case, is to listen with religious attention to the mother, and unless she be far beneath the average of human intelligence, our opinion as to the fact of some deviation from the ordinary health of the child is considerably influenced by what she tells us.

The inquiry in regard to the history of the case, previous to the first visit of the physician, should bear particularly upon the causes of the sickness, its precise moment and mode of attack, and its course and symptoms up to the present time. The most important points to be considered in connection with these objects, are the health of the parents, including their ordinary health, or their habitual diseases, the causes and periods of their death, if they are not living, and the state of health of the child at the moment of birth and since. The hygienic conditions in which the patient has been placed ought always to be investigated; the place of habitation; the kind of house, and whether a large well-ventilated room, or a small, narrow, and close one; the clothing; the food; and lastly, whether the infant has been suckled, or brought up on artificial diet. The state of the health just anterior to the attack ought always to be examined into. Has it been good and strong, or feeble and delicate? If delicate, what diseases? If the approach of any of the eruptive fevers be suspected from the character of the symptoms, the question as to whether the child has previously had measles or scarlet fever, or has been vaccinated or had variola, should always be asked.

It is next necessary to fix as accurately as possible, the precise period of the onset of the sickness. If the question, "When was the child taken sick?" be asked, as it usually is, the answer will be, "Oh, several days ago," or, "I don't recollect exactly,—I think yesterday, or the day before," or some such loose answer. The best way to learn the exact period in a recent case, is to go back, day by day, or else to inquire as to some particular day. We may ask, was the child quite well day before yesterday; was it well last Sunday? Did it play and amuse itself? Was it as gay and good-tempered as usual yesterday, or the day before, or the day before that? Did it sleep well night before last, or the night before that? A sick child scarcely ever sleeps well at night, and very often we may learn by close inquiry into this particular, the exact time at which the attack began. In this way, by forcing the attendant to tax her memory, and to

go minutely over the events of the several days previous, we shall nearly always succeed in fixing very precisely the moment of onset.

Having determined these points, we should proceed to inquire in regard to the course of the disease prior to the first visit. This is to be done only by patient and repeated questioning. The questions must be so framed as to elicit free and unbiassed answers. They should be general, and not leading. Lastly, we are to inquire into the treatment of the case up to the present time.

It is best that all these interrogatories should be made previous to seeing the child, in some other room than the nursery, in order to avoid the risk of alarming the child by the presence, during an unnecessary length of time, of a stranger. If, however, the child be well acquainted with the physician, it matters not where the inquiries are made.

Having now obtained from the attendants all the information they can give in regard to the history and nature of the case, the physician must proceed to the personal examination of the patient, in order to determine, by his own observation, the exact nature of the sickness, and the treatment it may require.

The most important points to be attended to during the clinical examination, are the countenance or facies, noting its expression, color, the presence or absence of furrows and wrinkles from pain, from emaciation, or from disordered muscular action, the appearances presented by the nasal orifices, and especially by the *alæ nasi*, and the characters exhibited by the mouth; the sleep; the cry; the state of plumpness or emaciation; the condition of the skin as to color, temperature, moisture or dryness, the presence of swellings of any kind, such as those produced by dropsy or by affections of the joints, and the existence of eruptions; the pulse; impulse of the heart; the respiration; the signs furnished by the state of the mouth and throat, and by the disposition towards and power of sucking, or by the manner in which drinks are taken; and lastly, the state of the abdomen.

THE COUNTENANCE.—The countenance of a young and healthy infant, who is sleeping or perfectly quiet, wears no expression except that of comfort and content. It is composed and still; no movement disturbs its innocent tranquillity, unless, perhaps, some gentle smile light it up from time to time, when we might well believe the happy superstition of the fond mother, who will tell us that angels are whispering to it. In sickness, even when slight, the countenance soon loses this expressionless character. In all acute disorders the alteration is very great, such indeed as to strike the most careless and inexperienced observer. The features become contracted, furrows and wrinkles appear about the forehead, the nostrils are dilated, or pinched and thin, and the mouth becomes drawn and rigid. The extent of the change is generally in proportion to the severity of the attack. The part of the face most altered will depend very much upon the particular system of organs implicated in the disease.

Some authors have endeavored to show that different diseases give to the physiognomy certain peculiar and characteristic *expressions*. This is true only to a certain extent. Thus, the facies is very different in abdom-

inal from that observed in thoracic or cerebral diseases; but though it is *generally* easy for a practiced physician to distinguished by the facies alone between a cerebral and thoracic disorder, it is quite impossible for him to distinguish between any two cerebral, thoracic or abdominal affections. The particular changes impressed upon the face by different diseases cannot, however, be discussed in this place, but must be considered in the separate articles upon each disease. Here it can only be stated in general terms, that in diseases of the brain, the upper part of the face, the forehead, and the eyes are chiefly affected; that in diseases of the thoracic organs, the middle portion of the face, and especially the nostrils; whilst in those of the digestive organs, the lower part of the face, the mouth and lips, are the parts which undergo the greatest changes in their expression.

Pain may almost always be detected by the expression of the face. It gives to the countenance various shades of expression, according to its degree of severity, and its permanency or recurrence at intervals. Pain in the head is said, by Dr. M. Hall, to produce a contracted brow, pain in the belly to occasion an elevation of the upper lip, whilst pain in the chest is chiefly denoted by sharpness of the nostrils. We doubt, however, whether pain in any particular organ imparts an expression to one part of the face rather than to another, for indeed pain in any part of the body, whether the head, chest, abdomen, or limbs, gives rise to a contraction of all the features. Not one part of the face alone, but the forehead, mouth, nose, and the whole face, become changed in expression and contracted, when there is severe pain in any part of the body, so that we deem it impossible from the expression alone, to determine where the painful sensation may be seated. The countenance merely tells us there is pain, but not where it is located. The painful expression will be permanent or occasional, as the pain itself is constant or only paroxysmal.

The *color* of the face becomes often an important means of diagnosis. In all the fevers, phlegmasiæ, and diseases of general excitement, the face will be more or less suffused and red, unless the attack be so severe as to occasion a violent shock to the nervous system, in which event the countenance instead of being suffused, is paler than natural. In such cases the face becomes of a dead white, all traces of red disappear, and the skin at the same time has often a slightly shining or varnished appearance. We have not unfrequently observed this symptom in pneumonia and bronchitis, and also in the later stages of true croup. It is a very striking phenomenon, and one which portends great danger.

In chronic cases of all kinds in which the hematosic and nutritive functions are enfeebled, the face assumes a pallid and waxen hue, which is very characteristic. In the various digestive ailments it becomes muddy or sallow, and in affections of the liver more or less yellow. Lastly, in certain diseases and malformations of the heart or lungs, it becomes bluish or livid, constituting one of the most important signs of what is called *morbus cœruleus*, blue disease, or cyanosis.

In reading the countenance of a sick child, the practitioner should always notice the play of the nostrils, since this reveals, to a certain extent, the state of the lungs. In pneumonia, bronchitis, and pleurisy, the

movements of the *alæ nasi* become rapid and energetic, expressing, by the degree of their violence and extent, the amount of embarrassment under which the respiratory function is laboring.

The nostrils and nasal passages should also always be examined to ascertain the presence of mucous or purulent secretions, or of pseudo-membranous deposits, since these fluids or their inspissated products interfere more or less with the free passage of air through those canals.

OF THE SLEEP.—Much useful information as to the state of health of infants and children may be obtained from a careful consideration of the various phenomena connected with their sleep. Of this we are fully convinced from somewhat long and patient observation. We cannot ascertain, indeed, the nature of the disease under which the child may be laboring, but we can detect, with very great certainty, the existence of a deviation from health. We know of few more certain means of fixing the period at which any attack of illness may have begun, than by inquiring at what time the child began to have restless or broken sleep, or insomnia.

A perfectly healthy infant, within the month, who is suckled at an abundant and wholesome breast, will usually sleep twenty out of the twenty-four hours, waking to nurse every two or three hours during daylight, and twice or three times during the night. After the age of two or three months, the child is much more wakeful during the day, though it will still take a nap of two or three hours in the morning, and a shorter one in the afternoon, while it will sleep from early evening until the following morning, waking but once or twice to suck. Indeed, many perfectly healthy infants, of between three and six or seven months of age, sleep without waking from nine or ten o'clock in the evening until six the next morning. After the latter age the sleep is seldom so unbroken; the child begins to undergo the first considerable trial to its health, dentition, and it is rendered thereby more or less ailing and irritable, and consequently restless and troublesome at night.

Children who have passed through the epoch of dentition, and who are perfectly well, usually go to sleep soon after being put to bed, and never wake until the following morning. Not only so, but they sleep soundly and quietly, without being disturbed by slight sounds, and without tossing or turning much in their sleep.

In healthful sleep the whole appearance of the child, its expression of countenance, its attitude, and its breathing, all declare a most perfect and beautiful ease and tranquillity. Nothing can be more suggestive of the comfort and well-being that naturally attend upon health, than the perfect composure and graceful postures exhibited by a hearty child during profound sleep.

It needs, however, but a slight disturbance of the health of a child to break in upon this ordinarily calm and peaceful sleep, and to render it restless, fitful, interrupted by startings, cries, or dreams, and insufficient. The most trifling irritations, as the pressure of a tooth against the gum, a faulty state of the digestion, the presence in the intestinal canal of imperfectly digested food, or the slightest fever from any cause, are sufficient to produce this effect, and hence it is that the character of the

sleep will often become to a watchful practitioner the first sign of disorder held out by nature.

The degree of disturbance of this function will vary with the nature and severity of the disturbing cause. When slight, the child will continue to sleep throughout the ordinary period, but the sleep will be somewhat uneasy. The countenance will be disturbed. There will be contractions of the brow, and momentary workings of the features, which express the perception of some unhealthful sensation. Often the child will toss and turn, and change its position more frequently than natural. Sometimes it will cry out, and appear distressed by some dream or painful sensation. When the cause of disturbance is more serious, the sleep is more broken, the child wakes often, and lies awake for a longer or shorter time, and it becomes very difficult to lull it to sleep again. Or it has painful dreams or night-terrors, causing it to scream and struggle in sleep, and then to wake in the most violent affright. In severe instances it becomes almost sleepless. We have very often known teething children not to sleep more than half as much as in health, and to wear out, by the long continuance of this sleeplessness, the patience and even the health of their attendants. In some instances they will no longer sleep in the bed or crib, and the nurse is obliged to get up and walk with them, or soothe them by the movement of the rocking-chair or cradle. In other cases, the derangement of the health is shown by grinding of the teeth, and by the most violent tossing and tumbling about the bed. We have frequently seen a child lying with its head where its feet should be, or across the bed, and with all the coverings thrown off, in spite of the most careful arrangement of the bedclothes.

These various disturbances are therefore signs of some alteration in the health of the child. They do not lead to an appreciation of the precise nature of that alteration, but they are invaluable as affording indications of the existence of some morbid condition of the economy. Very often, as above stated, they are the first symptoms of the approach of some more or less serious sickness, and as such will often enable us to determine, with much precision, the moment of onset of the attack.

THE CRY.—Crying is one of the modes of expression of the child. Indeed, this, with the expression of the face, are, according to M. Billard, the only means of expression with which nature has endowed the young infant. This is, however, scarcely correct, since we may also class among its means of expression the various spontaneous muscular movements indicative of uneasiness, or of pain, or pleasure; the manner in which it drinks or sucks, whether eagerly, and with appetite, or languidly, or carelessly, or not at all; the enjoyment it receives from pleasant sounds; and the evident delight it takes in regarding the light. Nevertheless, the cry and the expression of the countenance are the two means on which the physician must chiefly rely for early information of the occurrence of sickness in the young infant. These are the trusty sentinels of nature. By them she first gives notice of the approach of danger, and then measures the amount of mischief that may have been done.

The cry which a child utters during sleep, or even when awake, and

when nothing has been done to excite or disturb it, is always indicative of some uneasiness. If the cry be caused by pain, or by any considerable disturbance, it will be accompanied by certain contractions of the features and movements of the body and limbs, which will still more strikingly show that the pain, or other exciting cause, is of a serious nature. Violent and obstinate crying is almost always caused by severe pain,—such as the pain of earache. Indeed, obstinate and long-continued crying, lasting for hours, is rarely met with except from *earache, hunger, or thirst*. The cry of earache is often incessant and unappeasable, the pain being generally constant and not paroxysmal, as are most other pains. It is to be silenced only by the application of remedies to the ear, or by the internal administration of opiates. We have known an infant, three months old, to scream with earache for two days and nights, with only short lulls of a few hours when brought under the influence of large doses of laudanum. As soon as the ear began to discharge, the cry ceased. We are constantly called to see infants and young children who have been crying most violently for hours, and who are thought to have colic, or to have hurt themselves, but who are, in fact, tortured with that most violent of all pains, earache. We have met with few instances in which such severe and *constant* crying has depended on other causes; for, though children scream violently and obstinately from hunger and thirst, they may always be quieted by the supply of either want, whilst in earache the infant generally refuses the breast, or takes it only for a few instants, and then lets go to resume his almost automatic scream.

To show the difficulty of sometimes determining the cause of crying, we may mention that one of us once attended a nursing baby through a severe attack of bronchitis. Just as the child was recovering from the attack it began to cry without any apparent cause. The cry was so constant, violent, and severe, that, feeling certain from the symptoms that it could not be from any dangerous cause, we concluded, by the method of exclusion, though, to be sure, there was neither tenderness of the ear to touch, redness or swelling of the meatus, nor discharge, that it must be an earache. Hot applications and opiates applied to the ears did no good, and the constant scream set the mother half wild. At length the grandmother came in and said she thought the child wanted the breast. Sure enough, there was the trouble; the child lay at the breast almost continuously for twenty-four hours, and earache, crying and all, vanished.

In not a few instances we have thus known infants to cry very often in the day and night, and sometimes very obstinately, too, from hunger. In such cases the child is thought to have colic, and as it is not unfrequently costive, it is dosed with cathartics, carminatives, and opiates; or it is being brought up partially or wholly upon artificial diet, and, as a consequence, has some disorder of the bowels, which is thought to require other kinds of medicaments for its relief. When the stools are natural in appearance, or merely costive, and when the child does not labor under flatulence, it is easy, by careful questioning of the mother, to discern whether she has milk enough, and by examination of the size and weight of the child, to judge whether growth and nutrition go on in their proper ratio; and if it be

found that the mother is a poor nurse, and that the development of the child is slow and imperfect, we should at once direct an additional supply of nourishment, and the suspension of all mere drugs. We have often been surprised and delighted to find how soon, under the new treatment, the child becomes placid and comfortable, how well and how long it sleeps, and at what a rapid rate it develops its form and size. So, when the circumstances above referred to coincide with a somewhat disordered state of the bowels, we should first choose for the child the diet most appropriate to its age and state of health, and then, if after inquiry it appears that the whole quantity taken in the twenty-four hours is below the proper standard, the amount allowed must be augmented.

Thirst, as we have said above, is not a rare cause of crying in very young children. The cry, in this case, is often misunderstood by both physician and attendants, and is ascribed to pain in some unknown and undiscoverable part, to colic, to the teeth, or to ill-temper,—when, in truth, the child is suffering from the pangs of thirst. This is especially apt to occur in children who have had, or still have diarrhœa, with or without vomiting of their ordinary liquid food. The cry is not the acute, shrill cry of pain, occurring in sharp, short paroxysms, as in colic, or in the pleuritic stitch; nor is it the steady scream of continuous earache. It is rather a constant wail of low tone, accompanied with marked restlessness. It lasts for hours, or even for a day or two, until its cause is discovered, or until the thirst has been removed by the retention of appropriate food.

The crying occasioned by pain in the head, by the pain which accompanies pneumonia or pleurisy, or that which is attendant upon abdominal inflammations, is scarcely ever constant, though it may be violent while it lasts. Pain in the head usually causes a sudden or sharp cry or shriek, which is over as soon almost as heard, and which has been called the *hydrencephalic* cry. The pain of pneumonia, which, it should be remarked, is not unfrequently absent, or so slight as not to be noticed, commonly occasions crying only during coughing, and for a short time after, and is accompanied by distortion or grimacing of the features. In pleurisy, again, the cry is also heard generally at the moment of coughing, but it is produced also by the act of moving the child, and by pressure on the affected side. It is commonly much louder, shriller, and indicative of greater suffering than in pneumonia, and in some cases that we have seen, has been very frequent and difficult to appease.

The cry of intestinal pain may almost always be recognized by the fact that it takes place just before or after a stool, that it is accompanied by wriggling and twisting movements of the trunk, and especially of the pelvis, or, in very young infants, by its coincidence with more or less flatulence, which is revealed by a tympanitic condition of the abdomen, and by frequent regurgitations of gas.

Children not unfrequently cry much and very obstinately from mere fretfulness and general distress or malaise. This kind of crying may be recognized by its peculiar tone, which is short, sharp, and irritable. It is a fret rather than a scream; it is occasioned by the least disturbance offered to the child, by the attempt to move it, to dress it, to attend to any

of its wants, even to look at or notice it; it is moreover possible, generally, to still such a cry by soothing treatment, or by the endeavor to amuse the little thing with toys.

Lastly, a child will sometimes attempt to cry, but is unable to utter any or only a very faint sound. This depends commonly upon some laryngeal impediment, but may be also the result of pure exhaustion; there is not sufficient strength to sound a cry.

The cry of the young child has been divided by M. Billard into the cry proper and the return; and inasmuch as these two portions of the cry are differently affected in different diseased conditions, it is important that we should be aware of their existence, and of the effects produced upon their manifestations by disease.

The cry proper is produced during the act of expiration, while the return occurs during inspiration. The cry proper is sonorous and prolonged, the return is much shorter and sharper. The return is feeble in young infants, and becomes stronger as they advance in age. In different states of health, the mode of crying will vary to a considerable extent. The cry may exist alone, or in combination with the return; or again the return only may be heard, whilst the cry is from some cause suppressed. The difference between the two portions of the cry may always be distinctly perceived in a child who is crying violently from any recent cause, whether ill-temper, fright, or pain, unless one or other has been suppressed by some morbid condition which interferes with the perfect performance of the vocal function. After a time, however, when the infant has become fatigued with its efforts, the cry proper ceases in part, and we have only the return, which is heard from time to time between the sobs. According to M. Valleix, it is the return which becomes enfeebled and disappears first, whenever one portion only of the cry is heard. Towards the fatal termination of all diseases, the return ceases more or less completely, and the cry assumes a peculiar moaning or murmuring character, which must be familiar to all who have been much in the sick-rooms of children.

With a remark upon the condition of the lachrymal secretion in disease, we shall conclude this division of the subject.

The infant does not begin to secrete tears until towards the third or fourth month, and of course this function can furnish no sign previous to that time. After that period, however, the suppression of this secretion becomes, according to M. Trousseau, a valuable aid to prognosis, as this suppression occurs generally in all dangerous acute diseases. The occurrence of this symptom in any acute case should be looked upon, therefore, as one of dangerous augury, while the continuance of the secretion, or its reappearance after it has been suppressed, is, on the contrary, a highly favorable omen.

GENERAL APPEARANCE OF THE CHILD; DEVELOPMENT; EMBON-POINT; STATE OF THE SKIN, ETC.—While occupied in hearing the account of the sickness given by the mother or attendants, and even while asking questions in regard to the present state of the patient, the physician may learn a great deal that is useful by an attentive observation of the general appearance of the child as it lies before him. He should

study its size and development, its state of embonpoint or emaciation, its decubitus and gestures, the color, temperature, and dryness or humidity of the skin, and the presence of eruptions or swellings of any kind. Having remarked these various matters during the early part of the examination, he should proceed to inspect carefully the whole external surface by touch and sight, in order to acquire precise and accurate information upon these points.

A child who has been healthy from its birth ought to have attained a certain average size and development at a certain age. If it be much below the average size, if at three months it look like a new-born child, or at a year old like one of six months, it is very clear that something has acted to determine such slow and insufficient growth, and it becomes the business of the practitioner to discover what the impeding cause has been. Not only ought a child to have a certain size and stature, but it should also be possessed of a certain degree of embonpoint. A perfectly healthy young child, one under four years of age, usually presents a much greater fulness and rotundity of the trunk and limbs than does the adult. Its tissues are firm and solid, its surface of a cool and pleasant temperature, its color of a clear and exquisite white, finely tempered with a faint rosy tint in a warm atmosphere, or slightly marbled with light bluish spots in a colder air. Few marks more certainly indicate a healthful temper of the constitution than the clear and exquisitely tinted pink color of the palmar and plantar surfaces of the hands and feet of a young child. Nothing, indeed, can be more beautiful or perfect in shape or contour than the figure of a fine, hearty young child; nothing more pleasing to the eye than its delicate but vivid coloring; and nothing more expressive of the fulness of health and vitality than its whole appearance.

When, therefore, instead of these marks of a pure and active state of the health, we meet with stunted growth, emaciation, soft and flaccid tissues, sallow and dingy tint of the cutaneous surface, pallid or bluish feet and hands, weak and listless movements,—how easy the conclusion that some jarring agent is at work to hinder and obstruct the machinery of life.

In acute diseases emaciation takes place rapidly, but the tissues still retain some degree of elasticity and firmness. In chronic diseases the emaciation is of course slower, but it is more complete, so that, in some instances, the frame seems to consist merely of the bones wrapped round with a dark and unhealthy skin. The tissues beneath the skin, the cellular, adipose, and muscular, are in great part absorbed, and the skin falls into wrinkles and irregularities on the least movement of the child. In some cases of disease, and particularly in those of the abdomen, the derm loses almost entirely its elasticity, so that when pinched into a fold by the fingers, it retains for some time the form that has been given to it.

The *decubitus* and gestures of the child ought to be noticed. Healthy children are, when awake, almost always in motion. Those who have attained the habit of walking are tempted to active exercise by their various plays and amusements. Infants, though they sleep much more than older children, are also, when awake, constantly moving their limbs; they are

seldom still. When asleep they rest quietly and comfortably, generally upon the side, though often upon the back. How different when the child is laboring under disease of any kind. The disposition to movement is gone; the older child insists upon lying on the lap, or in the cradle or bed, and the infant is to be soothed of its crying and fretfulness only by rocking and fondling in the arms. Instead of the free and spontaneous movements of health, we now see only the sudden, impatient, and causeless tossing on the bed or lap, or the constant changing of position, with fretting or complaining, which constitute the *agitation* of sickness; or else the slow, languid, and hesitating movements of weakness or prostration; or, lastly the stillness and immobility of stupor or of coma.

There is nothing peculiar about the decubitus of pneumonia or bronchitis except when there is severe dyspnœa, in which case the child, if old enough to select its own position, lies high upon the pillows; while those who are younger evidently prefer to rest on the lap of the nurse, with the trunk and head supported in her arms, and express by crying and agitation their discomfort and uneasiness when placed in the recumbent position on the lap, or in the cradle or crib. We have seen several young children affected with severe bronchitis or pneumonia, who have preferred to any other position that of being held in the nurse's arms, with the front of the chest placed against her chest, and the head hanging over her shoulder. When the dyspnœa is so severe as to produce, by slow degrees, a partial asphyxia and consequent dullness of perceptivity, the child becomes soporous or comatose, and lies usually upon the back, as in diseases attended with prostration of strength.

In pleurisy and peritonitis the decubitus is usually dorsal, and the child dislikes to be moved or nursed, often crying violently when touched or disturbed.

In intestinal inflammations the young patient is usually excessively restless at first, and very fretful, unless the attack be grave and threatening, when it often lies still for a time from the prostration of strength which attends violent attacks, but becomes restless, turns and twists in the bed, cries out, and agitates the lower extremities at each evacuation of the bowels.

In the early period of cerebral inflammation there is generally excessive restlessness, and great irritability of all the senses and temper, but as the case goes on, and passes into the stage of coma, the child becomes still and quiet, assuming very often the decubitus called by the French "*en chien de fusil*"; that is to say, on the side, with the inferior extremities strongly flexed, and the arms drawn close to, or crossed over the thorax. This position is especially characteristic of the latter stages of tubercular meningitis.

Extreme restlessness, constant tossing upon the bed, or incessant changing from the arms to the bed, or from bed to bed, is a very bad sign. We have observed it in several different affections; especially in obstinate pneumonia, in long-continued intestinal disorders, and in the secondary inflammations of measles and scarlet fever.

Among the gestures most deserving of attention are the sudden starts,

attended with cries, which indicate the occurrence of some painful sensation, as that of colic, of stitch in the side in pneumonia and pleurisy, and sometimes of shooting pain in the head. The frequent carrying of the hand to the head, or to the ear, ought not to pass unnoticed, as this is often indicative of headache or earache. So also of the constant application of the hand to the mouth, or the introduction of the fingers into that cavity, which often occurs when the child is suffering from the odontalgic pain of dentition. Nor should the physician ever neglect to observe any peculiar and especially any automatic movements of the limbs, and particularly of the fingers or toes. Nature often heralds the approach of a convulsive seizure by certain peculiar muscular movements. The thumbs are drawn into the palms of the hand, and the fingers clasped over them; or the toes are strongly bent towards the sole of the foot, or rigidly extended; sometimes the fingers are for an instant convulsively extended upon the hand and drawn widely apart from each other; or lastly, the muscular movements, instead of being easy, steady, and natural, are badly co-ordinated; they are irregular, uncertain, and tremulous. This last character, tremulousness and uncertainty, we have often noticed.

The occurrence of paralysis will often be unperceived for some length of time by an inattentive observer. It is to be discovered by the failure of the child to move one limb, whilst the others are more or less agitated, or by taking hold of the limb, and comparing the total want of resistance in it, with a certain stiffness and opposition to movement almost invariably present in the healthful condition.

The state of the cutaneous surface is always important, and ought to be carefully and systematically examined. The points most requiring to be noted are its temperature, dryness or moisture, color, and the presence of eruptions or swellings. By the temperature, and dryness or moisture, taken in connection with the rate of circulation, we must judge as to the existence of fever. The inferences to be drawn from the condition of the surface in these respects are the same in children as in adults, and they need therefore no particular consideration in this place.

The color of the skin, on the contrary, owing to its great susceptibility to change in certain affections, becomes, in the diseases of early life, of very considerable importance in diagnosis, and deserves therefore some special remarks.

The physician should be aware, in the first place, that the color of a newborn infant is some shade of red, varying from a deep brick-red tint, to one of a much lighter hue. The red appearance fades away usually in about four or five days, and leaves the surface of a yellowish-white, or in some instances of a decidedly yellow color. The yellow color is sometimes so marked as to impose very readily upon an inexperienced person the idea that it must depend on an affection of the liver, or, in other words, that it constitutes a true jaundice. In a very large majority of cases, however, the conjunctiva retains its natural white tint, the digestive functions go on with perfect regularity, there is no fever, and indeed no marks of decided disorder of the health, so that the icterode hue cannot depend, under these circumstances, on any serious lesion of the liver or its append-

ages, and it is manifestly wrong to regard the case as one of disease, or as requiring any treatment.

Besides the yellow color just described, the cutaneous surface in children, and particularly in the new-born infant, may exhibit different shades of a bluish color, which need some attention. When the whole skin assumes a decidedly bluish tint, the case is one of cyanosis or morbid cœruleus, depending on some malformation or disease of the heart or lungs. In severe cases of this kind, the blue color deepens into a purple or even blackish hue. If this appearance last more than a very few days, there can be little doubt that it depends on some malformation or disease of the heart.

The blue or livid tint recurs sometimes, also, in sudden attacks of collapse of the lungs. In such cases it seldom lasts long, though it may be very marked during the paroxysm.

It is quite common to observe in new-born and very young infants a bluish tint of the hands and feet, and of the parts around the mouth, whilst the rest of the body is pale. These appearances depend usually on some obstruction to the pulmonary circulation, as that caused by atelectasis pulmonum, bronchitis, or pneumonia, and they increase, diminish, or disappear, according to the course of the causative malady. In older children, the blue color of the skin is rarely of any considerable intensity, unless the condition has existed from birth, or soon after; but it is not at all uncommon to meet with faint, but quite perceptible shades of that color, depending on the asphyxiated state which occurs in croup, capillary bronchitis, pneumonia, and sometimes in laryngismus stridulus. It is hardly necessary to add, that a very slight blueness of the fingers and toes is sometimes observed in the cold stage of intermittents.

Occasionally we meet with an excessive harshness, aridity and scurviness, or with a wrinkled appearance of the skin, especially upon the abdomen and thorax. This symptom, when strongly marked, is usually attended with enlargement of the superficial veins of that part, and is then very striking even to a careless observer. It accompanies very generally the abdominal tuberculosis of children, and should not pass unobserved. Though generally indicative of tubercular peritonitis, or of tuberculosis of the mesenteric glands, it is not always so, since in a case that occurred to one of us, and in which it was perfectly well-marked, a post-mortem examination showed it to have been caused by a chronic peritonitis, resulting from inflammation and suppuration of the mesenteric glands, entirely independent of tubercular disease. The peritonitis had given rise to extensive adhesions among the intestines, and the pus had found its way by a tortuous sinus between the intestines into the vagina, through which it was discharged externally.

There is one other alteration in the color of the skin which is deserving of notice in a practical point of view. It is an excessive pallor, occurring sometimes in diseases which obstruct the respiratory function. We have been most struck with it in the capillary bronchitis, or suffocative catarrh, of young children, and in membranous croup. The whole surface assumes a dead white hue, which seems to depend on a total want of blood in the cutaneous capillaries. The nose is white, the ears become white and di-

aphanous, and the only relief the eye meets with in gazing upon what seems an almost alabaster countenance, is the still pink or bluish lips, the dark eyebrows and eyes, and perhaps a somewhat leaden tint of the circumference of the mouth and of the forehead. In strongly marked cases the whole surface, even of the fingers and toes, exhibits this white or blanched appearance. When this condition has lasted for several hours, or a day or two, the hands and feet sometimes assume a bluish look, which may last until death occurs, or until the attack approaches a favorable termination. This condition of the surface, when occurring in cases attended with obstruction of the respiratory function, has always appeared to us an indication of imminent danger to the patient; and, indeed, when it lasts more than one or two days, it has very generally proved the harbingers of death.

The clinical examination of the cutaneous surface cannot be considered complete until it has been made with reference to the presence of eruptions, of swellings from œdema, of inflammation, tumors, and lastly, of diseases of the joints. The inquiry in regard to the presence of eruptions is a very important one, from the fact that children are particularly liable to attacks of the exanthematous and other eruptive affections. Many attacks of sickness, beginning with violent fever and other serious symptoms, which would otherwise remain entirely obscure or unexplained, until a much later period from the onset, may be accounted for at an early period by a minute examination of the skin. So, in the latter stages of long and debilitating maladies, in the disorders which follow scarlatina, and in cardiac and hepatic diseases, a proper inspection of the surface will reveal œdematous effusions that might, if this search were neglected or carelessly prosecuted, remain undiscovered. The same remarks will apply to inflammations of the auricular cavities, to the swelling of the joints produced by rheumatism, and to some obscure suppurative inflammations in the limbs of children. A most instructive example of the necessity of this close examination, occurred some years ago in the practice of one of us. A healthy male infant, five weeks of age, was seized suddenly with most violent fever, the reaction being not unlike in character that of acute rheumatic fever. The only visible disturbance of the health, to explain this violent attack, was a certain amount of digestive derangement, and for this the patient was treated. After three days of most severe illness, with strong tendency to convulsion, and with some stiffening of the lower jaw, we were asked to look at the right thigh. It was largely swelled, especially in its lower half; it was hard to the touch, and the skin over the outside of the limb, just above the knee, had assumed an inflammatory redness. It was clear that the child had been attacked with an acute inflammation of the deep tissues of the thigh, and that this was now approaching the surface and becoming visible. Careful inquiry now brought to light the fact that the baby, all through the sickness, had cried severely, as though in sharp pain, whenever it was moved, and especially when its napkins were changed. The distress observed when the napkins were being changed, had been ascribed to some smarting from the urine. Had the surface of the child been more carefully examined at an early period,

the swelling of the thigh, and the pain on motion, might, no doubt, have been detected then, and the intense febrile reaction, with the nervous symptoms, which were thought too great for simple functional disorder of the digestive functions, would at once have been explained.

It is clear, therefore, that in infants and in children under six or even eight years of age, the physician must depend, in great measure for information as to the nature of the case, on his own unassisted explorations; and, knowing this, he should leave nothing neglected that may aid him to gauge with accuracy the state of health of the individual before him. He should cultivate a *habit* of minute, systematic, and patient investigation, since by accustoming himself to such a method in his daily walks, he will assuredly attain, in the end, a tact and sagacity that will not often be at fault.

THE PULSE.—The pulse of the child, in order to be judged of to any real advantage, must be examined during the state of quiet, and, if possible, it should be felt whilst the child is either asleep or dozing. During the waking state a young infant is in such constant motion, that it is very difficult to perceive the pulsations of the radial artery, and impossible to judge of their force or volume, in consequence of the rising and falling of the flexor tendons of the forearm, and because, also, of the natural softness and delicacy of the pulse at that age. In older children, the moral disturbance occasioned by the visit of the physician in most instances, and the irritability and nervousness accompanying the sickness, will either cause the patient to resist the attempt to touch the arm, or else produce so great an effect upon the rate and force of the circulation, as to render very uncertain and unsatisfactory any conclusions to be drawn from the examination. If possible, therefore, the circulation should be examined during sleep. If this be impracticable, the child ought, when still nursing, to be put to the breast, or, when weaned, it ought to be quieted by soothing treatment, by toys, or by the promise of a toy.

It is essential that we should know what is the average of the healthy pulsations of the child, in order to obtain a standard of comparison by which to judge of any departure from that average in disease. Observers have varied not a little in the results at which they have arrived by their examinations upon this point. By selecting those, however, which appear to have been made with the greatest care, and under the most favorable circumstances, we shall, doubtless, obtain an average entirely worthy of confidence. It will be necessary, also, to obtain averages for different periods of childhood, since the rate of the circulation varies to a very great extent at different ages. We shall, therefore, give the rate of the circulation for new-born children (one to ten days old), for the period from four months to six years, for that from six to nine years, and for those from nine to twelve, and from twelve to fifteen years of age.

The average rate of the circulation in very young infants, is from one hundred and one to one hundred and two in the minute, the former being the result obtained by M. Billard in children from one to ten days old, as nearly as it can be gained from his statements, and the latter the one obtained by M. Roger, in infants from one to seven days old (*De la Tem-*

perature chez les Enfants, Paris, 1844). The physician ought, however, to be aware of the fact that, though the above is the average rate of the circulation at the age mentioned, the pulse may range very much above or below that average, without necessarily indicating a morbid state of the health. Thus, though the average frequency in forty children, from one to ten days old, observed by M. Billard, was one hundred and one, it was less than eighty in eighteen, whilst in fourteen it was between one hundred and one hundred and twenty-five, and in six between one hundred and thirty and one hundred and eighty. All these children, he assures us, presented every mark of good health.

The average frequency of the pulse during the first year may be stated at about one hundred and fifteen; at least such is the result obtained by us from an examination of seven observations by M. Roger of children from four to nine months old. This result, it will be observed, shows that the pulse is not so frequent during the first few days after birth, as it becomes at a somewhat later period, which, moreover, agrees with a previous statement to the same effect made by M. Valleix. This latter author is of the opinion that at seven months of age the pulse is much more frequent than some days after birth, and that it afterwards falls gradually as the child advances in years.

We are not acquainted with any observations upon the rate of the circulation during the second year of life, except those of M. Trousseau, who according to M. Bouchut (*Manuel Prat. des Mal. des Nouv.-Nes*, p. 133, Paris, 1845), gives as the average between one year and twenty-one months, one hundred and eighteen.

M. Becquerel (*Traité Théorique et Prat. des Mal. des Enfants*, Paris, 1842), gives us the result of his observations upon thirty children, between two and six years of age, during sleep and in the waking state. During sleep the average was seventy-six; in the waking state it was ninety-two.

Between six and nine years of age, the same observer found the average during sleep to be from seventy-three to seventy-four, whilst in the waking state it was ninety. Between nine and twelve years, the average was, during sleep, seventy-two, in the waking state, eighty. Between twelve and fifteen years the rate was seventy whilst the children were asleep, and seventy-two when awake. Roger gives seventy-seven as the average between six and fourteen years.

One very striking fact attracts our attention in the above statements: the much greater difference between the rate of the circulation during sleep and during the waking state, in very young children, than in those who are somewhat older. Thus, whilst there is a difference of seventeen pulsations in the minute, in the rate of the circulation during sleep and in those who are awake, between the ages of two and six years, the difference under the two conditions mentioned, amounts to only two pulsations in the minute in children that have reached the age of between twelve and fifteen years.

The circulation is somewhat more rapid in girls than boys. This difference should be borne in mind, but as it amounts to only about five beats

in the minute, it is insufficient to be of any very decided value in diagnosis or prognosis.

After these specifications as to the rate of the circulation in children, we shall pass on to some general remarks upon the method of the examination of the pulse, and upon some other of its important characters.

M. Bouchut (*loc. cit.*, p. 129), remarks that in infants at the breast "the palpation of the pulse is almost impossible. It may be counted, but its force, feebleness, size and hardness, can scarcely be appreciated; the intermittent character is the only phenomenon upon which no doubt need rest; it is, moreover, the only one of any value." These opinions of M. Bouchut, though true in some degree, are much too strongly stated, for we are quite sure that it is very easy to detect great differences in the force, size, and tension of the pulse of the same child in health and in disease, and of different children laboring under different diseased conditions. These differences can be detected by careful observations from a very early age, and after two months may be readily recognized, when the variation from the state of health is at all considerable.

The intermittence of the pulse above alluded to, should rather be expressed by the word *irregularity*, since the pulse is not properly *intermittent*, but merely *irregular* in its rhythm. This is quite a common feature in the pulse of children, and, be it noted, is much more frequently met with during sleep than in the waking state. M. Becquerel met with irregularity of the pulse in twenty-four of one hundred and fifty children examined during the waking state, and in fifty-five of one hundred and fifty during sleep. It is clear, therefore, that mere irregularity of the circulation, independently of other symptoms, is not a sign of disease, since it was present in one-sixth of those awake, and in a little more than a third of those asleep. It should be observed, too, that the greatest irregularity exists when the pulse is lowest (in sleep). The chief practical bearing of this fact is that we should be careful not to lay too much stress upon the slowness and irregularity of the pulse, as signs of tubercular disease of the cerebral meninges, unless they are observed during the waking state, and in connection with other symptoms; particularly with vomiting, constipation, and severe headache.

Another very important characteristic of the circulation of the child, is its extreme irritability, which causes its rate to vary to an extraordinary degree, even in perfect health. This is the more marked in proportion as the child is younger. The slightest disturbance, whether moral or physical, will cause the pulse to rise in a young child from one hundred or one hundred and fifteen, to one hundred and twenty, one hundred and thirty, or even one hundred and fifty. From this circumstance may be drawn the inference also, that the pulse should always be examined, as before stated, during sleep, or during profound quiet.

There is still another reason which makes it necessary to touch the pulse during sleep or profound quiet. This is, that when the child is agitated, it becomes literally impossible, in consequence of the contractions of the flexor tendons of the forearm, and of the movements of pronation and supination, to judge with accuracy the various qualities of the arterial action.

EXAMINATION OF THE HEART.—The examination of the heart by auscultation and percussion ought, and, to be of essential aid in diagnosis, must be performed while the child is still and quiet. It is best made during sleep, especially in infants; when this is impossible, it can be performed with great advantage during the state of quiet that follows nursing, or during that which may often be procured by soothing management, or by taking advantage of the fondness that infants show for a strong light, the view of which will generally suffice to occupy and keep them still.

The sounds of the heart present the same general characters in the child as in the adult. They are, of course, more feeble and more rapid; conditions which make it difficult, in the young infant, to perceive and appreciate any minute change from the healthy sounds. After the age of one or two years, however, when the circulation has become slower and more steady, the signs yielded by the physical examination of the heart become much more valuable and positive; so much so, indeed, as to yield results almost as important as in the adult. The first sound is almost always duller than the second. They succeed each other commonly with perfect regularity, and have the same interval between each in the same child. The cardiac sounds are readily heard by placing the ear over the præcordial region. The extent of surface over which they may be heard will depend on several conditions: particularly the state of quiet or agitation of the child, the presence or absence of fever, the state of the lung as to its consistence (constituting it a better or worse conducting medium of sounds), and the condition of the heart itself as to health or disease.

In a healthy child, who is undisturbed by any cause of irritation, and particularly in one sleeping, the sounds are distinctly audible over the whole præcordial region and under the left clavicle. In many subjects they can be heard over the whole front of the thorax, but become, of course, feebler in proportion as we recede from the præcordial region. Usually they are heard quite as distinctly under the right clavicle as over the nipple of that side, in consequence, no doubt, of their transmission in an upward direction by the aorta. They are never heard over the posterior walls of the chest in children in perfect health, and whose circulation is entirely undisturbed. In those who are awake and agitated, and in those who have been making severe muscular exertions, the cardiac sounds are very loudly audible over the whole front of the thorax, and even through to the back of the chest.

When the lungs are indurated by inflammation, as in pneumonia, they transmit with great distinctness, from having become better conducting media, the cardiac sounds to the back. This circumstance sometimes becomes a valuable aid in the diagnosis of pneumonia. We have been enabled to satisfy ourselves of the existence of pneumonia in the lower lobe of the right lung, in a doubtful case, from the fact that the sounds of the heart were much more clear and distinct over the right inferior, than over the left inferior dorsal region.

The præcordial region is decidedly less sonorous on percussion than the parts of the thorax directly over the lungs. This diminution of sound is distinct enough to be evident to any ordinary ear, but it rarely amounts

to absolute flatness. The region exhibiting this dulness of sound is the same in position as in the older person. It occupies the space corresponding to the cartilages of the fifth, sixth, and seventh ribs, and is situated, therefore, between the left nipple and the left edge of the sternum. Its measurements, as given by MM. Rilliet and Barthez, are one and a half to three inches in a transverse, by one and a half to two and a half in a vertical direction. The region of dulness is described by those observers as being represented by a circle or ellipse, the transverse diameter of which extends from the nipple to the sternum, or more rarely, towards the xiphoid cartilage. In children over six years old, the nipple sometimes lies above the middle line of this space.

THERMOMETRIC OBSERVATIONS IN CHILDREN.—As an indication of the intensity and character of the disease in febrile attacks, we have seen that the frequency of the pulse is little to be depended on. Dr. Forster (*Jour. f. Kind.*, July and August, 1862, in *New Syd. Soc. Year-Book*, 1862, p. 413), who has made an extensive series of observations upon this subject, asserts that variations in the temperature of the body offer far more certain indications. The instrument used was a Reaumur's thermometer, eight and a half inches long, in which slight variations are easily appreciable. The bulb was placed in the axilla.

The results given are those of observations upon healthy children, during the first few days of life.

A constant lowering of the temperature of the body takes place after birth, which reaches its maximum, 28.97° R., on an average within the first two hours after birth.

Hours after birth.	Average Temp. (R.)	Minimum Temp. (R.)
$\frac{1}{4}$ —2,	28.97	28.2
2—6,	29.12	28.1
6—10,	29.49	28.7
10—15,	29.53	29.0
15—20,	29.31	29.8
20—25,	30.04	29.7
25—30,	29.9	29.7
30—36,	30.07	29.7
36—42,	30.04	29.4
42—48,	29.86	29.3

A subsequent elevation always occurs. The average time at which the highest temperature was observed, was from thirty to thirty-six hours after birth, at which time the average was 30.07° R.; maximum 30.4° R.; minimum 29.7° R.

The elevation was noticed equally when the infant had and had not taken food.

During the first nine days of life, the temperature was observed as follows:

Days.	Maximum (R.)	Minimum (R.)	Average (R.)	No. of Observations.
1 — $1\frac{1}{2}$,	30.4	29.7	30.01	22
$1\frac{1}{2}$ —2,	30.5	29.3	29.93	16
2 — $2\frac{1}{2}$,	30.4	29.3	29.87	28
$2\frac{1}{2}$ —3,	30.3	29.2	29.74	16
3 — $3\frac{1}{2}$,	30.3	29.3	29.76	27

Days.	Maximum (R.)	Minimum (R.)	Average (R.)	No. of Observations.
3½—4,	30.2	29.0	29.68	17
4 —4½,	30.4	29.2	29.68	25
4½—5,	30.3	29.2	29.72	18
5 —5½,	30.4	29.2	29.82	23
5½—6,	30.5	29.3	29.81	16
6 —6½,	30.6	29.4	29.83	23
6½—7,	30.3	29.1	29.75	17
7 —7½,	30.4	29.3	29.82	22
7½—8,	30.4	29.0	29.72	11
8 —8½,	30.0	29.4	29.70	8
8½—9,	29.9	29.6	29.75	2

We thus see that from the thirtieth to the thirty-sixth hour after birth the highest temperature is observed. Then a fall takes place, which reaches its maximum at four days after birth (average maximum 29.68° R.). Again, between the fifth and eighth days, a new elevation of temperature occurs; but this new elevation is less in degree than that previously noted. The average maximum was 29.83° R. Some differences were found in the results, according as the children were large and heavy, or the reverse. Large and well-developed children had a slightly higher temperature than those less robust.

Thus the average temperature in the early part of the day was, in children weighing eight pounds and upwards, 29.84° R.; but, in children weighing less than this, the average was 29.65° R. The evening observations, again, gave an average for the heavy children of 29.94° R.; for the others, of 29.77° R. Respecting the temperature at different times of the day, observations showed that, from the second to the ninth day, there was an average elevation of temperature, from morning to evening, amounting to $.11^{\circ}$ R.; the average morning temperature being 29.75° R.; the average evening temperature, 29.86° R.

This interesting subject has been further examined in regard to older children, by Mr. Finlayson (*Proc. of Manchester Med. Soc.*, in *Brit. Med. Jour.*, Jan. 16th, 1869, p. 59).

His results are based on two hundred and eighty-one observations on eighteen different children, of ages varying from twenty months to ten and a half years, and are as follows:

1. The daily range of temperature is greater in the healthy child than that recorded in healthy adults—amounting to 2° F.
2. There is invariably a fall of temperature in the evening, amounting to 1, 2, or 3 degrees.
3. This fall may take place before sleep begins.
4. The greatest fall is usually between 7 and 9 P.M. (at least under the conditions of life in hospital).
5. The minimum temperature is usually observed at or before 2 A.M.
6. Between 2 and 4 A.M. the temperature usually begins to rise, such rise being independent of food being taken.
7. The fluctuations between breakfast and tea-time are usually trifling in amount.
8. There seems to be no very definite relationship between the frequency

of the pulse and respirations, and the amount of temperature; the former being subject to many disturbing influences.

RESPIRATION; ITS RATE AND GENERAL CHARACTERS.—The respiration, like the pulse, to be examined with any advantage to the explorer, must be investigated whilst the child is still and quiet. In the young infant it should be done during sleep as it is only then that we can find the breathing uninfluenced by disturbing causes other than those connected with deranged health. In the older child, the play of whose functions is more steady and regular, and less readily jarred by trivial causes, this part of the clinical exploration may be made during the waking state; but, still, it must be done whilst the patient is quiet and tranquil, else the results obtained will necessarily be less certain and reliable than under the opposite state of things.

When examined during sleep, much may be learned by a careful study of the breathing. In health the child breathes entirely through the nostrils. The mouth is closely shut, and enough air passes through the nasal passages to give the child all the air it needs without any visible effort. So noiseless is the breathing that, in quite healthy infants, no sound can be heard unless the ear is applied close to the face of the child. The inspiration may just be heard, by close attention, as a soft, light souffle; but no sound is heard in the expiration. The inspiration is continuous and gradual, the expiration short and rapid, and after this comes quite a long pause, lasting perhaps two seconds. In perfectly healthy children of six months, the expiration is about 24, and the pulse 108 to 112; at eighteen months, the expiration is about 20, and the pulse 112. This noiseless, easy breathing, with the long pause between the two acts, is an almost infallible sign that there is no disease of the lungs.

The respiration ought always to be counted by the watch, if possible, especially by the young practitioner. This is the only mode in which a perfectly accurate idea of the frequency of the respiration is to be obtained. It sometimes happens that a greatly increased rate of the breathing will pass unnoticed by the physician, from the fact that it continues to be regular and without effort. We have known children to breathe eighty times in the minute, without presenting any appearance of labor or effort in the act; without cough, and without the least wheezing or sound to be heard at a short distance from the patient. Under these circumstances, the great rapidity of the respiration might very well pass unnoticed, especially by inexperienced practitioners, and, be it remarked, this would be particularly apt to happen were the attention of the physician addressed to some other part of the economy than the thorax, as the seat of the sickness. For instance, in latent pneumonia, when this stimulates meningitis, or when it is conjoined with gastro-intestinal symptoms, the failure to note a greatly increased rate of the breathing might very well occur. In many cases of secondary pneumonia, it might also take place. In children who have been long sick with diseases that debilitate and impoverish the health, a sudden aggravation of the symptoms dependent on collapse of the lung, might be misunderstood and falsely explained, for the want of this precaution. It is therefore a good and useful rule, for the young practitioner

always to count the respiration, when he has to do with a case presenting the least obscurity of diagnosis, since this simple habit may guide him to the real seat of disease, which else he might mistake.

The rate of the respiration in children is very different at different ages, a circumstance that should always be recollected in the examination of their diseases. The average frequency of the breathing in new-born children and during the first week of life, is thirty-nine, according to M. Roger. It may rise, however, upon very slight disturbances, to fifty, sixty, or even eighty, while it is not at all unusual to find it at twenty-five or thirty in perfectly healthy infants during sleep. Between the ages of two months and two years the average is about thirty-five. Between two and six years, the average is eighteen during sleep and twenty-three during the waking state; from six to twelve years, the average during sleep is eighteen, and in the waking state twenty-three; from twelve to fifteen years, it is eighteen in the former, and in the latter twenty. It will be observed, therefore, that after the age of two years, the rate of the respiration is nearly the same throughout the remainder of the period of childhood; it changes so little, indeed, that the same average will answer for all practical purposes throughout that period.

The other characters of the respiration require some attention on the part of the practitioner. In the first place, the diaphragm plays a more important part in the process in the child than in the adult. In the young infant, indeed, the function is carried on almost wholly by the action of that muscle, so that the respiration is correctly described by the technical term of abdominal. The walls of the chest are almost motionless. On this account the rate and characters of the breathing can be best studied in young children, by examining the abdomen, the movements of which being strong and marked are much more easily seized by the eye than are those of the thorax.

During perfect quiescence, and especially during sleep, the breathing of a young child is soft, regular, though less so than in the adult, and perfectly noiseless; it is necessary to place the ear close to the face or chest of the child, and to listen attentively, in order to hear it. In the young child, and especially the young infant, the breathing is, in the waking state, very different from that of the adult. It is short, irregular, uneven, and marked by occasional pauses, followed by a hurry and precipitation of the movements. These peculiarities in the respiration of the infant appear to depend on the weakness and imperfect action of the muscular apparatus at that early age, which causes the various movements of the body to be hesitating and uncertain, and without that steadiness and evenness which are characteristic of matured strength. After the age of two years, these irregular and tumultuous movements cease, and the breathing becomes more regular and even, like that of adults.

In the inflammatory affections of the lungs,—pneumonia, bronchitis, and pleurisy,—the respiration is almost invariably accelerated. In extensive pneumonia, and in capillary bronchitis, it becomes very rapid, rising to eighty or one hundred in the minute. We counted it in one case at one hundred and twenty-eight. In pleurisy and simple ordinary bronchi-

tis, it seldom becomes so frequent, not exceeding, usually, forty or fifty. In severe pneumonia, the rhythm of the movement sometimes becomes inverted; the pause occurs at the termination of the inspiration instead of the expiration. The patient makes first a violent and labored expiration, bringing into a kind of convulsive action all the expiratory muscles of respiration; instantly after the expiration follows a rapid and full inspiration; then occurs a momentary pause, and again the respiratory act begins with the labored expiratory effort. This kind of respiration is a very unfavorable symptom, as it is indicative of a most dangerous oppression. It is particularly apt to occur in infants and very young children. It has been called *expiratory* respiration.

The respiration, though almost invariably accelerated in pulmonary inflammation, sometimes retains its normal rate, or even falls below that rate. This occurs, we believe, only under one condition of things; when the forces of the constitution have been sapped by previous disease, or exhausted by the long continuance of the thoracic inflammation. It is therefore met with in cases of secondary inflammation, and in those of the chronic form.

The respiration is very much increased in frequency as a general rule in atelectasis pulmonum, or collapse of the lungs. When, therefore, a young child who has been exposed to the cause of this disease (feebleness at birth, exhausting disease, or debilitating hygienic conditions), is suddenly seized with hurried respiration, slight cough, paleness or blueness, with coldness of the cutaneous surface, and in whom there are but few and unimportant physical signs of pulmonary disease, there is very good reason for supposing that some portion or portions of the lungs have become collapsed, or, in other words, have ceased to admit air.

The respiration often lends some assistance in the diagnosis of cerebral affections. In acute meningitis, accompanied by violent febrile reaction, it is more frequent than natural, but often irregular. When the early stage passes into the stage of coma, the breathing becomes slow and irregular. In tubercular meningitis it is seldom increased in frequency except for a day or two before death, whilst in the middle period of the disorder, it is either continued at its normal rate, or becomes slower. During that period, also, it is almost always extremely irregular, and is interrupted by long and mournful sighs, which, to the ear of the experienced physician, who hears in them the almost certain prognostic of approaching death, have an inexpressibly touching sound, increased tenfold by the consciousness of his utter inability to control the fatal tendency of the malady.

There is a peculiarity of the respiration which occurs in collapse of the lung and also in cases of membranous croup, which ought not to be passed by unnoticed. It is, that during the inspiratory effort, the ribs move inwards and backwards towards the mesial line of the trunk, instead of outwards as in normal respiration; and at the same time there may be recession of the lower part of the sternum, so that a more or less deep sulcus is produced around the base of the thorax. This peculiarity is readily explained, as shown by Rees and Jenner, by reference to the nor-

mal relation which exists between the current of inspired air, the expansion of the lungs, the descent of the diaphragm, and the firmness and resistance of the thoracic walls. If this relation be disturbed in any way, the phenomena we are now considering may be produced. Thus if the diaphragm contract suddenly and violently, the lungs cannot expand with sufficient rapidity, and in order to prevent the occurrence of a vacuum, the thoracic walls must yield to the external atmospheric pressure at their least resisting part, which is under normal conditions, at the base of the chest. The same result must occur, also, when the diaphragm contracts with only normal force, but when the calibre of the larynx is much narrowed, or again, when a considerable portion of lung-tissue is collapsed. In the article on rickets, an affection in which the firmness of the chest-walls is much diminished, a full account will be found of the masterly manner in which Jenner has applied the above principles to the explanation of the deformities of the thorax so characteristic of that disease.

AUSCULTATION AND PERCUSSION OF THE LUNGS.—This portion of the examination of the sick child ought to be performed, if possible, whilst the patient is still and quiet. Unfortunately, however, it happens in a large majority of cases that the disturbance of position necessary to effect the exploration, and the presence of the physician, together with the irritability of nerves and temper occasioned by sickness, almost always cause more or less resistance on the part of the child, and produce violent screaming and struggling. In young infants we have to contend only against the instinctive resistance to any physical disturbance naturally attendant upon sickness and suffering. In older children, who have learned to distinguish between familiar and strange faces, and in whom the will has begun to act, there is added to the instinctive resistance of the infant an opposition of the most strenuous and annoying kind, founded upon the natural fear of a stranger, and upon a mental determination not to be interfered with or incommoded by the movements and changes of position necessary for a careful examination.

For these reasons, the physical exploration of the chest in young subjects is often to be accomplished only with great difficulty; and in the midst of the most violent screaming, struggling and contention. It is clearly important to avoid these obstacles if possible. This can only be done by the employment, on the part of the attendants and physician, of the most soothing, gentle, and patient management; and in this way, let it be remarked, it can be done in a large majority of cases. The possession by the physician of a quiet and yet decided manner, the power to interest and attract the child by entering with active sympathy into its little amusements and pursuits, the skill to engage its attention by the exhibition of some book or toy, or the mere influence he may exert to calm its terror or excited irritability, by a soothing voice and gentle persuasion, will, in many instances, overcome any resistance offered to the examination by children over two years of age. Nevertheless, in very young children, and in not a few that are older, no gentle means whatever will overcome opposition. Here the exploration must be made in the midst of struggles and cries, and though the results obtained will be less clear and

positive that when the child is reasonable and obedient, a great deal of most valuable information can be acquired by a quick and dexterous practitioner. The percussion can be made in the short intervals between the cries, or even during their continuance, and by placing the ear close to the finger by which it is performed, the sounds elicited can be very well heard and judged. The auscultation is more uncertain; but, by watching intently the long and deep inspirations which immediately precede the violent cries, the presence or absence of râles, and their characters, the degree of freedom with which the air enters the lung, and the existence or non-existence of bronchial respiration, can, after some experience, be ascertained and commented upon, so as to give considerable certainty to the diagnosis.

The particular position in which to place the child, during the examination, is of some importance. After the age of three or four years the position may be the same as that selected for the adult, if only the patient be reasonable and tractable. When, on the contrary, the child resists, it should be taken on the lap of the mother or nurse, or else held in the arms, with the head inclined over one shoulder, while its back is presented to the practitioner. Infants within the year may sometimes be examined whilst engaged in the act of sucking; but this is inconvenient, both from the constrained position, and from the circumstance that the inspirations are short and imperfect during the act. The French authors recommend that the very young infant should be laid, with its face downwards, across the hand of the practitioner, who is then to approach the back of the chest to his ear. We have found either one of the three following positions most convenient, as the case may be: the infant laid across the lap of the mother, with its face downwards, and the head hanging a little over one knee; held in the arms, with the front of its body placed against the mother's chest, and the head lying over her shoulder; or, lastly, a favorite position of ours, placed in a sitting posture upon the lap, supported by one hand in front, and by the other holding the occipital portion of the head.

Auscultation should always be performed before percussion, because the latter generally alarms or annoys the child, and occasions crying, which of course would interfere more or less with the auscultation, were this performed after percussion. The auscultation should be made with the ear rather than the stethoscope, for the reason that the instrument terrifies the child, and also because it cannot, when the child resists and struggles, be kept in contact with the chest. Moreover, the instrument is unnecessary, except for the examination of the upper portion of the thorax in front, and it had better, therefore, be dispensed with.

Percussion is best made in children by using a finger of the left hand as the pleximeter, and by striking with one finger of the right. One finger is quite sufficient to elicit all necessary sound in young subjects. The strokes should be light and distinct, consisting sometimes of short and quick, and sometimes of slow and measured taps. By the latter slow strokes the exact characters of the sound are often better developed than by the former.

To perform auscultation and percussion with success, the surface ought to be quiet uncovered. The habit of examining the thorax through one or several thicknesses of cloth, which some persons fall into, is a most careless one, and cannot but lead to uncertain and erroneous results.

As a general rule, it is sufficient, in young children, to examine the posterior portion of the thorax. Doubtless it is more accurate and artistic to explore the whole chest, and this ought to be done in all obscure cases. But when the child is sick and suffering, when it is irritated and exasperated by the presence of a stranger, or by coercion, and still more, when it is weak and exhausted by long or violent illness, it becomes of the greatest importance to shorten, as much as possible, the time occupied in the examination. For these reasons, it is well to be aware of the fact that, in nearly all inflammatory diseases of the lungs, the morbid changes affect first and most severely the posterior surfaces of those organs. This is thought to depend on the fact that the child passes so large a portion of its time in the recumbent position as to cause the fluids of the body to gravitate towards the dependent parts of the lungs, and thus to determine the beginnings of inflammatory action in that direction. Certain it is, be the explanation what it may, that it is rare to find the anterior surface of the lungs affected either with bronchitis, pneumonia, or pleurisy, the posterior surface remaining healthy. When, therefore, upon auscultation and percussion, no signs of disease are met with over the dorsum of the thorax, we may feel pretty well satisfied that the lungs are healthy. Nevertheless, in all doubtful cases, the examination ought to be extended to the whole chest, in order to make what was, before this has been done, only a strong probability, a certainty. Whenever, also, it is important to ascertain the precise amount of disease in any serious or long-continued sickness, the front as well as the back part of the chest must be examined.

The *respiratory sounds* are not of the same character precisely in the child as in the adult, and of this the physician ought to be aware. In children the vesicular murmur is stronger than in the adult, so that it assumes somewhat of a blowing or bronchial sound. It was in consequence of this peculiarity that Laennec gave it the name of puerile respiration, which, though a mark of health in early life, is, at the period of maturity, an indication of a morbid change in some portion of the pulmonary structures. It ought to be remarked, however, that in infants under two, and particularly in those under one year, the vesicular murmur is, in ordinary respiration, weaker than in adults; owing, no doubt, to the fact that the inspirations are short and imperfect, not distending the lungs to their full capacity. When, however, from any cause, a sigh, a sudden disturbance, or the act of crying, a full and complete inspiration takes place, so as to dilate thoroughly the pulmonary structure, the murmur becomes at once loud and strong, or, in other words, *puerile*, as in older children.

The murmurs of inspiration and expiration bear the same relation to each other as in the adult; the expiration being much shorter and feebler than the inspiration, though, at the same time, it, like the inspiration, is

louder than in the adult. In some instances, however, and especially over the posterior, inferior, and lateral regions of the thorax, no sound whatever is heard during the accomplishment of the expiration. This absence of sound during expiration is the more apt to be met with in proportion as the child is younger.

When a young child is made to breathe forcibly and rapidly, the respiratory sounds assume certain characters, even in perfect health, which might mislead an inexperienced observer. The inspiration is short, loud, and hard, so as to assume somewhat of a blowing character, resembling not a little the sound of bronchial respiration. At the same time, the expiration becomes louder also, and longer, which two circumstances, rude or even blowing inspiration, with loud and somewhat prolonged expiration, may very well deceive a young or careless practitioner.

The respiration is most clear and characteristic over the anterior lateral, and posterior inferior regions of the thorax. Over the origin of the larger bronchia, that is to say, in the interscapular region, the respiration is very strong, so as to resemble very closely bronchial blowing. Here, also, the expiration is often very marked; it is sometimes heard as long, or even longer than the inspiration. Over the scapulas, the sound of respiration is always feebler than elsewhere, except in the præcordial region from the interposition of the scapulas and of thick muscles between the ear and the lung.

Percussion yields a much louder and more sonorous sound in children over two years of age than in adults,—a circumstance always occurring coincidently with the presence of puerile respiration, and dependent on the fact that the function of respiration is, at that age, very active, and the lungs therefore filled to their utmost capacity with air. In infants under two years of age, the sonorousness varies to a considerable extent in the same child. When the respiration is, as it usually is, gentle and easy, the inspirations being rather feeble and incomplete, the amount of air contained in the lungs will be somewhat deficient in comparison with what their cells might contain, and the sound yielded upon percussion will necessarily be rather dull and insonorous. When, on the contrary, the respiratory process is quick, active, and energetic, from any cause, so as to give rise to the auscultatory phenomenon called puerile respiration, the percussion will be loudly sonorous, as it is in the later periods of childhood, owing to the thorough dilatation of all the air-cells, and the consequent presence in the thoracic cavity of a large amount of air.

The sonorousness of the thorax is different in different parts in children, as in adults. In front, the percussion is most sonorous from just beneath the clavicle on the right side down to one or two inches below the nipple, where it gradually becomes dull, owing to the position of the liver. On the left side the sonorousness is modified by the presence of the heart in the manner already mentioned. Below the præcordial region we again have pulmonary resonance down to the sixth or seventh ribs, below which is heard the tympanitic sound of the stomach.

Behind, the sound is dull above the spine of the scapula, and considerably so over the scapula beneath its spine. Over the interscapular

space it is clear and strong, and more so in the lower than in the upper half. Beneath the inferior angle of the scapula, likewise, it is clear and full, until we approach the inferior margin of the thorax, where it is dulled, even above the lower edge of the lungs, by the presence beneath of the liver on one side and of the spleen on the other. Over the right side the dulness begins a little higher than over the left, in consequence of the greater bulk of the liver than of the spleen.

The lateral regions are very resonant in their upper portions, but become dull as we approach the liver on the right side and the spleen on the left. On the left side the pulmonary sound is often entirely eclipsed by a tympanitic resonance occasioned by the presence of gas in the stomach.

In practicing percussion in children it is necessary to strike gently, because, from the great natural sonorousness of the chest in early life, any considerable force would bring out so much sound as to prevent the recognition of a degree of dulness which might readily be perceived by the use of more gentle blows. It is necessary always to compare the two sides together, as in adults, since this often leads to the detection of a degree of impaired resonance which might be otherwise inappreciable. Yet, and the physician ought to be well aware of this, the comparison of the two sides is not quite so useful in young as in mature subjects, because of the fact that the diseases in which the differential comparison is most important, pneumonia and pleurisy, are more frequently double than in adults. It becomes, for the same reason, very important to compare the upper and lower portions of the thorax behind, since we may assure ourselves of the existence of dulness below, of which we were before doubtful, by the fact that the sound is less sonorous in that region than above; which is, as already stated, the very opposite of the healthy condition.

EXAMINATION OF THE ABDOMEN.—It is often very important to ascertain, by palpation, the form, size, and degree of tension of the abdomen, the presence or absence of effusions within its cavity, and the condition of the organs which it contains; to learn by percussion the degree of resonance which it affords; and lastly, to find by pressure whether it be unnaturally tender to the touch or not. By a careful inquiry into these various points, and a proper comparison between them and the rational symptoms presented by the patient, we shall be able to discover the existence of tumors, of hypertrophied organs, of unusual developments of gas in the intestines, of dropsical effusions, of enlarged and hardened mesenteric glands, of gurgling, and of soreness on pressure caused by inflammation of some of the contents of the cavity. The examination should be made, if possible, whilst the child is still and composed. It is best, therefore, to perform it before auscultation and percussion, in children who are old enough or amiable enough to be willingly quiet, since the length of the examination of the thorax often wearies out their patience, and they refuse to submit to further inspection; whilst, in infants and in children who obstinately resist the examination, it matters little at what particular period it is attempted, since it must be done at last in the midst of cries and general agitation. It is, at all times, a difficult and not very useful examination, unless the patient consents to it freely and without fear. It is very

necessary, therefore, to resort to every means to obtain this quiet consent. In children over a year old, this condition is to be obtained only during deep sleep, during the act of nursing, or, when the patient is awake, by so pleasing and attracting its attention by toys, by soothing voice and manners, as to cause it to forget what is passing. The reasons why the examination is useless, unless made during a state of calm, are very obvious. In the first place, the contractions of the abdominal muscles give to the walls of the abdomen such a degree of hardness and rigidity, that it is impossible to learn anything in regard to the state of the parts within, except merely what can be learned by percussion; and, in the second place, no acuteness of perception will enable us to distinguish between the cries of anger and fright, and those that may proceed from pain occasioned by pressure.

M. Valleix recommends a plan in the case of young infants, by which tenderness on pressure may very generally be recognized. It is as follows: He carries the child, carefully sustained in the arms, suddenly before a bright light, either that which pours in at a large window during daylight, or that of a bright artificial light at night. The infant, whose greatest pleasure consists in gazing at a bright light, almost always ceases to scream and becomes perfectly quiet while thus attracted. Seizing this opportunity, the physician should pass his hand under the clothes, and applying it directly over the cutaneous surface, he may first learn, by a rapid palpation, the general characters of the abdomen, and then ascertain by sudden and decided pressure, whether it be abnormally sensitive. If the pressure gives pain, the infant will cry out at the moment, while, at the same time, a sudden contraction of the countenance will assist to show the perception of some painful sensation. Should the infant, on the contrary, continue to gaze fixedly at the light, without noticing the manœuvres of the physician, it is fair to conclude that there is no inflammatory tenderness present.

EXAMINATION OF THE MOUTH AND FAUCES.—In all obscure attacks of sickness occurring in young children, and even in those who have attained to the faculty of speech, the physician ought to be most careful to inspect the condition of the mouth and fauces, since not a few cases of fever which seem at first view inexplicable, are at once made plain by this simple exploration. We were once called to see a child three years of age, who had been sick three days with fever, thought by intelligent and educated parents to depend on gastric derangement. A single look into the throat showed it to be completely clogged with pseudo-membranous exudation, whilst a slight hiss in the inspiration, and a husky voice, declared that the same fatal product was just entering the larynx. The time for successful action had slipped by; the patient died two days after in the agonies of slow croup. On another occasion we were called to take charge of two children in one family who had been ailing several days with feverish symptoms, loss of appetite, languor, and some complaint of sore throat. In both we found the fauces covered with plastic deposit, and both died a few days after of membranous croup. Some years ago we attended a child between five and six years old, for a period of four days, with

irregular fever, some vomiting, total anorexia, languor, indisposition to play, and rare complaints of pain in the *chin and neck*, that were not mentioned to us by the attendants, so that all the time we had the idea that the attack was one of gastric embarrassment. Greatly to our amazement and consternation, the mother informed us on the fifth day that she had seen something white in the throat, and upon examination we found both tonsils covered with whitish exudation. Happily the exudation was still confined to these glands, and we were able by appropriate treatment to prevent its further extension.

In croup, also, in whatever form it may make its attack, the fauces ought to be closely watched, in order to know by the presence or absence of false membrane, the probability or improbability of the case being one of the membranous kind. In scarlatina and measles, especially in the former, the throat ought to be examined each day, to ascertain its precise condition, and particularly to learn whether there be present any disposition to membranous, ulcerative, or gangrenous angina.

In young infants also, the mouth requires a thorough examination from time to time in all their ailments, and especially in diseases of their digestive organs, since they are liable to thrush, to aphthæ, and, in chronic and debilitating maladies, to gangræna oris. In teething children, the act of dentition requires that the mouth should be inspected occasionally in order to ascertain the state of that process, and to detect the existence of the form of stomatitis called ulcerative, which generally occurs between the ages of one and five or six years.

The mouth can be readily examined by pressing upon the chin with force sufficient to cause the child to separate the jaws. In the young infant this very generally produces crying, during which the mouth is widely opened, and the state of the cheeks, lips, gums, and tongue can be perfectly well seen. In an older child, who refuses to open the mouth, or to keep it open, the handle of a smooth silver spoon is the best instrument to employ by which to effect our purpose.

The throat cannot be well seen at any age, except by depressing the base of the tongue, which is best done by means of a spoon-handle, as above directed. When a child refuses obstinately to open the mouth, and resists with violent struggles, it should be taken on the lap of a strong assistant, with the back of its trunk resting against the chest of the assistant, whose arms should restrain, by being crossed over the body and limbs of the child, its more vehement movements. Another assistant must hold the head of the child steady, whilst the physician obliges it to open the mouth, either by closing the nostrils with the fingers, or by slowly and gently, but firmly, insinuating the handle of the spoon between the teeth. After the spoon has once been passed over the tongue there is seldom any difficulty in obtaining a good view of the fauces.

The *introduction of the finger into the mouth* is of some use as a diagnostic means in the case of infants. It informs us of the temperature of that cavity, of the state of its secretions, and consequently, of its dryness or humidity, and of the disposition and ability of the infant to suck. When an infant is in good health, it will almost always seize the finger, when

this is placed in the mouth, and suck vigorously for some instants. It will do the same when it is only ailing with some slight malady, and in the early stage of more dangerous diseases. But, in severe and threatening illness, the infant either refuses to suck upon the finger at all, or does so only for an instant. When the mouth is irritated or inflamed, as in the various forms of stomatitis, the child will open the mouth and cry, and make no attempt whatever as suction. In stupor, and especially in coma, but little attention is paid to the finger, the infant being generally unconscious of its presence.

By watching the child when put to the breast, we may acquire nearly the same information as that just referred to, except that the child would naturally make a greater effort to seize the nipple than the finger, and would therefore nurse, even though the act of so doing were painful, under circumstances in which it might refuse to grasp the finger at all. The refusal to nurse, or the nursing but little at a time, may depend on other causes, however, than sore mouth. It often depends on some anginose inflammation. When this is the case, it may be suspected from the peculiar gulping manner in which the child swallows, and from the fact that swallowing often causes fits of coughing. It is caused also by dyspnœa. An infant laboring under severe oppression from pneumonia, bronchitis, or any other cause, never sucks well and steadily, but rather by fits and starts. The nipple is seized often with avidity, and two or three swallows are made in quick succession; then follows a pause to regain the breath, and then again the effort of deglutition. In a few cases attended with very great dyspnœa that we have seen, the patients have been able to swallow only once or twice without pausing, and even then with very great difficulty.

MANNER OF TAKING DRINKS.—The remarks just made as to the inferences to be drawn from the manner in which the infant sucks, will apply also to the mode in which both infants and older children drink. A young child drinks continuously, without stopping to breathe. If, however, it have any disorder which accelerates the respiration, it will, after drinking a few mouthfuls, cease, jerk its head away from the cup or spoon, breathe irregularly and hurriedly, and cough. These symptoms ought to call attention to the respiratory organs. So, if a child, whose breathing is not oppressed, nevertheless drinks with difficulty, slowly, at intervals, and apparently with pain, there is reason to suspect some impediment in the pharynx, and the fauces ought thereupon to be carefully examined.

We may learn also from the manner of drinking whether the child is thirsty or not. When it drinks often and with avidity, and yet has a dry mouth, it is evident that there is very great thirst.

VOMITING AND THE DISCHARGES BY STOOL.—The physician should never think his examination of a sick child concluded until he has inquired as to the occurrence of vomiting, and as to the state of the discharges by stool. Not only, indeed, should he inquire as to these symptoms, but he ought by all means to inspect personally the appearance of

the matters ejected. This is especially important in regard to the dejections, since no description of a mother or nurse, however intelligent, can impart to the physician the precise and accurate idea of the state of those discharges which even a very rapid inspection would give him.

Vomiting is of very frequent occurrence in infancy and childhood. Owing to the fact that the stomach is much less curved in its shape than in the adult, and that the œsophagus enters the organ close to its left extremity, vomiting and regurgitation take place with great readiness, and are, therefore, very common symptoms in the diseases of early life.

The young practitioner must beware lest he regard all kinds of vomiting in the infant as the result of disease. The nursing child is very apt to vomit, even when in the most perfect health, especially if it be suckled at an abundant breast. This kind of vomiting, however, may be readily distinguished from that which depends on some morbid state of the health, by the circumstance of the infant's ejecting nothing but the milk which it has swallowed, either just as it was drawn from the mother, or slightly curdled, and by the fact that it suffers no inconvenience whatever from the act,—neither any violent effort, languor, paleness, nor faintness. And yes we have known a young practitioner to prescribe antacids and absorbents to correct this kind of vomiting, which is most plainly an act of nature kindly intended to rid the infant of any excess of food it may have imbibed.

In older children also, vomiting not unfrequently occurs as a consequence of overdistension of the stomach with food. When, therefore, after vomiting, a child seems relieved and comfortable, when any unpleasant symptoms that may have existed prior to it moderate or disappear afterwards, it is fair to conclude that the act has been beneficial, and wrong to regard it as the signal of a necessity for giving medicine, or for regarding the child as a patient, except inasmuch as to watch lest it be sick as an after-consequence of having had the digestive power overtasked.

Frequently repeated vomiting, attended with retching and effort and with paleness and exhaustion, or with fever, always indicates some considerable derangement of the health. It is impossible to ascertain the precise cause of such vomiting, except by a proper consideration and comparison of all the symptoms the child may present. The cause may be in the stomach itself, consisting of an inflamed state of the organ, or it may be a simple indigestion without any inflammatory condition whatever; it may be that the cause lies in the intestine, being some inflammation, functional disease, or obstruction of that part; it may be pneumonia or pleurisy; it may be the approach of some of the eruptive fevers; or last, and most serious of all, the cause may be some commencing lesion of the brain, which, though as yet determining no proper cerebral symptoms, shall perhaps be destined, by its inevitable progress, to end the patient's life. The detection of the particular causative condition, in any of these forms of vomiting, can be arrived at only by a careful study of the whole constitution of the patient, both through the rational symptoms

that may be present, and by a thorough examination of the different systems of the body by means of the physical methods of diagnosis.

The rule to examine with his own eyes the napkins or cloths of the child, ought never to be forgotten by the practitioner, where there is any reason to suppose that the alimentary functions are at all deranged. The number of the stools in the twenty-four hours ought also to be ascertained, not loosely and carelessly, but precisely and with certainty. Without a close attention to these two precautions, it is impossible for the physician to obtain really useful and exact ideas in regard to the nature of the disorder he is called upon to treat, or to judge of the degree of severity of the attack.

We shall not attempt to consider in this place either the various unnatural appearances of the matters vomited, or passed by stool, the amount of those substances, or the frequency with which the discharges take place, since those various circumstances can be treated of in the manner they require, only when we come to study separately the diseases of which they form a part.

We shall here conclude our remarks upon the methods to be pursued in the clinical explorations of the diseases of children. We have only to add the wish that those who shall honor them with their perusal, may find them of some real assistance in their subsequent studies of the affections of early life. They are intended, of course, chiefly for the student and young practitioner; but we cannot help hoping that they may possibly prove useful to some who have spent a long time in the profession, but who have never, perchance, given any particular attention to the best modes of investigating the diseases of infants and children.

CLASS I.

DISEASES OF THE RESPIRATORY ORGANS.

CHAPTER I.

DISEASES OF THE UPPER AIR-PASSAGES.

SECTION I.

DISEASES OF THE NASAL PASSAGES.

ARTICLE I.

CORYZA.

DEFINITION; SYNONYMS; FORMS; FREQUENCY.—Coryza is an inflammation of the mucous membrane lining the nasal passages. It is called in common language, cold in the head, or snuffles.

We shall describe three forms of the disease,—the simple or mild, the severe, and the chronic. The severe form includes purulent and pseudo-membranous coryza. Simple coryza is very common at all ages; it occurs frequently as a distinct disorder, but still more frequently in connection with laryngitis, bronchitis, pneumonia, measles, scarlet fever, etc. Purulent and pseudo-membranous coryza rarely occur idiopathically, the affections upon which they are most frequently dependent being diphtheria, measles, and scarlet fever. A full account of this complication will be found in the chapters devoted to these various affections.

Chronic coryza occurs most frequently in connection with scrofula or hereditary syphilis. It also results from the persistence of the purulent form which has originated in the course of some specific fever.

CAUSES.—The only clearly evident cause of simple primary coryza, in most cases, is chilling of the body. Insufficient dress,—a very common error in this country,—too low a temperature of the nursery, and exposure to bad weather, may often be discovered to have been the causes of the attack. The other extreme of keeping the nursery overheated tends equally to the development of coryza and of more serious catarrhal affections, because the child becomes so relaxed and sensitive as to be unable to bear the slightest exposure.

As already stated, acute purulent coryza usually occurs in connection

with some general disorder, though we have occasionally met with it as an independent affection, for which no satisfactory cause could be assigned. The cases of MM. Rilliet and Barthez coincided generally with primary or secondary purulent or pseudo-membranous angina. From the account given by Underwood of coryza maligna, there can be little doubt that it was epidemic when observed by himself and Denman. The latter author states that in connection with the coryza there was a general fulness of the throat and neck externally; that the tonsils were tumefied, and of a dark-red color, with ash-colored specks, and in some cases, with extensive ulcerations; and that some of the children swallowed with difficulty; all of which symptoms clearly point to severe concomitant angina. There can therefore be little doubt but that in reality these were cases of nasal diphtheria.

ANATOMICAL LESIONS.—The Schneiderian mucous membrane is found reddened uniformly, or in points, rough, thickened, and sometimes softened. When pseudo-membrane is present, it exists either in fragments, or lines the whole extent of the nasal passages, and is mixed with mucus or mucopurulent fluid, in greater or less quantity.

SYMPTOMS.—The symptoms of simple coryza are sneezing, dryness of the nose at first, soon followed by discharge, which is very small in quantity in the beginning, and more abundant afterwards, and more or less disturbance of the respiration. It is only in young infants that this form of coryza is a disorder of any consequence in itself. In older children it never injures the health seriously by its own action; it is of importance merely as the sign that a cold has been taken, and ought to be regarded as a hint for more careful nursery management in the future. But, in infants at the breast, and very young children, it assumes much greater importance from the very considerable obstacle it opposes to the act of respiration. At this early age coryza becomes a serious and even dangerous disease. If primary, it causes great distress and disturbance to the child, interrupting its sleep, interfering with the act of nursing, and, in some instances, so impeding the function of respiration, as to bring on slight, and more rarely, dangerous asphyctic symptoms. It may, undoubtedly, occasion in weak and debilitated children, more or less extensive collapse of the lungs, an accident which will explain the imperfect performance of the hematosic function in some cases, where the only evident disease is this apparently insignificant one of coryza.

When simple coryza exists in connection with bronchitis and pneumonia, it adds to the severity of those diseases. In children over three or four years old, and particularly in those who are vigorous, it seldom gives any serious trouble. But in young infants, and in weakly children of any age, its influence upon the symptoms is often very marked. The effort to breathe through the nasal passages, when they are partially or wholly occluded by the inflammatory swelling of their lining mucous membrane, or by abundant and viscid secretions, fatigues and wears away the strength of the child, exhausts its energies, and renders it less able to resist the pressure of the sickness. But not only this;—as in primary coryza, the entrance of air into the lungs is impeded, and the

hematotic function is thereby interfered with, while at the same time, the existence of an obstacle to the full inspiratory movement, in addition to that which exists in the lungs themselves from bronchial or pneumonic disease, cannot but assist in the production of that collapse of the pulmonary tissue, which coincides so often with the bronchitis and pneumonia of young children, and especially with the former.

The reason why coryza causes so much difficulty in young children is, that they persist in the effort to breathe through the nose in spite of the obstruction of the nasal passages. They seem to do this instinctively, not apparently having the power to carry on the act of respiration through the mouth, or but for short periods only at a time. The constant struggle to force the air through the nose, and the necessarily smaller quantity that reaches the lungs, are undoubtedly the two chief causes of the symptoms above described as occurring in the coryza of children.

SEVERE CORYZA begins with sneezing and stoppage of the nostrils, soon after which the discharge, which is the pathognomonic symptom of the disease, makes its appearance. This consists of serous or mucous fluid in greater or less abundance, usually of a yellowish color, and which, at first thin and without odor, becomes afterwards thicker and often purulent, with a peculiar, unpleasant, but not fetid odor. In other cases, on the contrary, and especially when pseudo-membranous exudation is present, the discharge is thin, and often contains small granular particles, which seem to be the detritus of the false membrane, while at other times it is ichorous or even bloody. When false membrane is present, it can often be seen, upon examination of the nostrils in a strong light, to cover the mucous membrane in the form of thin adherent layers of a yellowish-white color. The *alæ nasi*, and sometimes the whole extremity of the nose, are red and swollen, and the skin, which is tense and shining, presents an erysipelatous appearance. The upper lip is generally reddened, irritated, swollen, and sometimes excoriated by the nasal secretion.

The *respiration* is generally difficult, nasal, and snoring. When the nasal passages are nearly or quite filled with the secretions, the child being no longer able to breathe through them as in health, is compelled to keep the mouth open. This is exceedingly inconvenient to children of all ages, as it causes dryness and stiffness of that cavity, and of the tongue and throat, and in very young infants, who instinctively respire almost exclusively through the nostrils, it is attended with such violent efforts as to be a chief or perhaps sole cause of the fatal termination of some cases. In one instance that we saw, the child was seized with attacks of suffocative breathing, which threatened fatal asphyxia, whenever the passages became much impeded. Under these circumstances the cleansing of the passages with a brush would afford complete relief, and for a time the little thing would appear to be quite well. Finally, however, death occurred in one of the attacks of dyspnœa, from sudden serous effusion into the lungs. The difficulty of respiration is greater, as we have stated, in proportion as the child is younger, and depends on the physiological fact already referred to, that at a very early age, respiration is performed almost solely through the nostrils, and that the child seems incapable of keeping the mouth open,

in order to compensate for their closure. We have never observed *cough* except in cases accompanied by angina. Slight *epistaxis* occurs sometimes in cases of the pseudo-membranous form. Infants refuse the breast when the nasal passages are much clogged, or suckle with great difficulty and at long intervals.

The character of the *general symptoms* depends much more upon the accompanying disease, in older children, than on the coryza itself, and it is unnecessary therefore to dwell upon them. In two cases observed by ourselves, the principal symptoms in one unaccompanied by angina were restlessness, weakness, emaciation, dry, harsh, and wrinkled skin, and violent attacks of dyspnoea; and in the other case, in which angina was present, there were added to these, fever and somnolence. The *duration*, as observed by ourselves, in cases occurring in infants, has been between two and three weeks. In other cases, which occurred in older children, the duration of the attack depended on the form and degree of the attendant angina. As will be stated, when treating of diphtheria, the complication of pseudo-membranous coryza in that disease is of very unfavorable significance; and such cases, if severe, may terminate fatally in two or three days. But it is evident that this result depends more on the specific blood-disease than on the local inflammation of the nasal passages. In some cases it became chronic, and was accompanied by ulceration of the nasal passages.

The *prognosis* must depend on the age of the child, and the form of the attack. Simple coryza is never dangerous except in very young infants, and rarely in them. When, however, it occurs in a delicate infant, and is accompanied with either sufficient turgescence of the nasal mucous membrane, or with enough viscid secretion, to cause a nearly complete occlusion of the nasal passages, the effort to breathe through the nose, and the diminished quantity of air that reaches the lungs, will sometimes give rise to great and dangerous exhaustion, or to partial or fatal asphyxia, with collapse of lung-tissue. In older children this form of the disease is scarcely ever more than an annoyance.

When simple coryza occurs in connection with other diseases, whether thoracic inflammation, angina, or measles, it always adds, and sometimes seriously, to the difficulties of the patient, since the effort to breathe through the obstructed air-passages tends to exhaust the life-forces, while at the same time a certain amount of the blood in the lungs, which ought to be exposed at each inhalation to the inspired air, is deprived of this necessary contact by the fact that less than the natural quantity of air is drawn through the nasal passages at each expansion of the chest.

The purulent and pseudo-membranous forms of coryza are always dangerous, whether they occur alone or as a part of other diseases. When they occur in connection with diphtheria, or in the course of scarlet fever, the prognosis will of course depend very much on that of those diseases.

CHRONIC CORYZA.—Under this title we shall describe as succinctly as possible a form of inflammation of the Schneiderian membrane, of which we see a good many examples. It is characterized rather by swelling and

thickening of the mucous membrane, as far as this can be seen, and by an accumulation of scabs and crusts, causing obstruction to the passage of the air, than by a discharge. The secretions are, in fact, not much increased in quantity beyond their natural amount, but they consist of very thick mucus, or they are purulent in character.

This form of the disease may be met with at any age, from a few weeks old up to puberty. Its principal *cause* has always seemed to be some faulty state of the general health, some constitutional dyscrasia. Like the keratitis and chronic otorrhœa of children, it makes its appearance without any evident exciting cause whatever, or it follows an acute attack of catarrhal inflammation from cold, or an attack of measles, scarlatina, or epidemic angina. On one occasion, we met with it in three out of a family of four children. Though it is unquestionably very apt to occur in scrofulous children, its presence is not necessarily a sign that the patient is of scrofulous habit, since we have seen it in families in which there was no taint of that disease, and have known a good many of those affected by it to recover perfectly, and show no subsequent symptoms of the scrofulous or tuberculous cachexia. Its chief efficient cause appears to be a low state of the general health, the blood being more or less markedly anæmic, and the nutrition of the body imperfect. In addition to the above conditions, it must also be borne in mind, as a fact of the utmost importance, that this form of coryza occurs frequently as a symptom of constitutional syphilis.

The chief *symptoms* of this form of disease are of a local character. The breathing is at all times more or less nasal and embarrassed. Even in the waking state, the child will sometimes attract attention by the noisy and slightly oppressed character of its respiration, while when asleep the obstruction of the passage of air through the nasal passages will be so great as to give rise to symptoms which, though not alarming, are most annoying to those around. The obstruction to the passage of air through the nasal passages produces snoring or hissing sounds, which are sometimes so noisy as seriously to disturb those sleeping in the same apartment. This obstruction also obliges the child to make much greater muscular efforts than in the healthy state, to supply the thorax fully with air, so that the sleep, instead of being quiet and easy as in health, is broken and disturbed by the unusual play of the muscles, and by the disordered internal sensations caused by the reaction upon the nervous centres of a circulating fluid less thoroughly decarbonized than it should be. The child tosses and rolls, sighs and moans, or it cries out in its sleep, or it wakes suddenly and frequently.

When the nasal passages are examined by a full light, they will be seen to be obstructed in two ways: by a thickened and injected state of the mucous membrane, and by the presence in them of scabs, or of more or less inspissated masses of mucus or muco-pus. The mucous membrane is also redder and more highly vascular than natural, and sometimes exhibits an appearance in some points as though excoriated or slightly eroded. There is seldom, indeed rarely, and considerable amount of fluid secretion, as in acute coryza; the secretions are so much more viscid than usual that

they desiccate in the passages and form scabs and crusts. Not unfrequently the surfaces become so irritable as to bleed very easily. The act of blowing the nose, a rude touch, or a blow, will cause a considerable discharge of blood, and this is often the symptom for which the practitioner is particularly consulted. The voice of the child is usually characteristic; it is nasal; and when the obstruction is considerable this becomes a marked symptom.

The general appearance of the patient almost always shows a deteriorated state of the general health; his color is too pale; the skin is muddy; the expression is languid; the tissues are more flabby and flaccid than they ought to be; and the movements are less brisk and prompt than in full health. Such patients awake from their sleep less refreshed than is natural; their appetite is often capricious and poor; and the digestive and nutritive functions are impaired. The tongue is often flabby in its texture, pale, and more or less furred, the bowels are irregular, and the discharges often scanty, and of an unhealthy color and smell, or there are alternations of diarrhœa and constipation.

In addition, when the case is connected with constitutional syphilis, some of the other evidences of this disease may usually be detected; though we have, like West, met with cases where the coryza has been the only sign of the constitutional taint.

The *duration* of this form of coryza is very indefinite. Under the most patient treatment, it often lasts for many months, and even when cured is very apt to return with or without apparent exciting causes, so that we have known it to last for several years.

TREATMENT.—Simple coryza requires no treatment in children over two years of age, except attention to hygienic conditions. Young children may often be preserved from attacks of spasmodic laryngitis and bronchitis, by calling the attention of the mother to the strong tendency which exists during infancy and childhood to the extension of disease, and advising, in cases of coryza, that the child should be secluded in the house, or else very warmly clothed if sent out.

In young infants, even the mildest coryza gives trouble, by obstructing the full freedom of the respiratory act, by interfering with the suckling, and by the restless and broken sleep which it induces. In such cases, all the treatment required is to keep the child warm, and to clear the nasal passages, and at the same time lubricate them by the occasional introduction of a camel's hair pencil, charged with diluted glycerin, cosmoline, or sweet oil.

When the coryza is more severe, so as to interfere a good deal with the respiration, it is necessary to make use of the brush frequently, to administer a warm foot-bath once or twice a day, and to give a few drops of syrup of ipecacuanha, with sweet spirits of nitre, every two, three or four hours. It is our habit to give quinia in such cases, even in young infants, and the result has satisfied us that the attack is modified and curtailed thereby. The amount should be half a grain twice or thrice daily for a child under a year old, and double that amount twice or three times daily for a child of two or three years. It is usually well received if suspended in a deli-

cately made syrup of liquorice, although it will be found highly convenient to use it in the form of suppositories, of very small size, for administration to young children. In such cases, the late Dr. Charles D. Meigs was in the habit, for many years past, of directing a flannel cap to be put upon the child, and kept there for two or three days;—a simple, and often most effectual mode of treatment. The cap should be removed after two or three days, so soon as the coryza is relieved, as otherwise the child is apt to become so much accustomed to it as to take fresh cold when it is removed.

In infants laboring under purulent or pseudo-membranous coryza, the indications of treatment are to remove the secretions as they collect, and to subdue the inflammation of the mucous membrane by which they are produced. It must never be forgotten, moreover, that this form of the disease is very frequently dependent upon some general specific disorder, of which it may indeed be one of the earliest symptoms, as in diphtheria. It need not be said that in such cases the general treatment is of even more importance than the local. The first indication may be fulfilled by means of a brush made of long camel's-hair, by throwing water or lime-water from a small syringe into the nasal passages, or when the discharges are thin and fluid, by blowing strongly into the nostrils, whilst the tongue is depressed by a finger introduced into the mouth, so as to allow the secretions to pass out of the posterior nares into the fauces.

The second indication is to be fulfilled chiefly by the application of solutions of alum, nitrate of silver, sulphate of zinc or copper, and by insufflations of different substances in powder. The best application is probably the solution of nitrate of silver, which may be made of the strength of five or ten grains to the ounce, or stronger, to be made use of several times a day, with a brush. We have also employed injections consisting of solutions of alum, of from three to six grains to the ounce. It is recommended by MM. Rilliet and Barthez to make insufflations of powdered gum and alum, or of gum and calomel in equal parts, several times a day. There is, however, it seems to us, an objection to this method of treatment, especially in infants,—which is, that the powders would necessarily tend to increase the obstruction which already exists, to breathing through the nose. It has been proposed also to apply a few leeches to the mastoid process, or over the frontal sinuses; but it seems to us that this could scarcely ever be advisable.

In the form of the disease accompanied with angina, an essential part of the treatment must be that of the throat affection. This will be considered in another place.

The treatment of chronic coryza must be two-fold: *general* and *local*. The most important points to be attended to in connection with the general treatment are the clothing; the diet, and the administration of tonics and alteratives. The *clothing* ought to be warm during the cold seasons of the year. Flannel, as a general rule, ought to be insisted upon. The arms and neck must be covered, and the legs should never be exposed, after the very mistaken fashion amongst many persons of the present day. The diet ought to be strengthening and nutritious. Fresh meats, milk, bread,

and good butter, and the plainer vegetables, ought to be urged upon the child. If necessary, some authority must be made use of by the parents to induce the patient to take a sufficient quantity of these plain, but nutritious articles of food. Pastry, cakes, candies, nuts, hot bread, sweet-meats, and all such rich, but not really substantial diet, should be forbidden to as great an extent as possible.

Of the tonics to be given, the best are the preparations of iron and cod-liver oil. Of the former, we prefer commonly the syrup of the iodide of iron, from three to five drops, at four or five years of age, three times a day, in half a teaspoonful or teaspoonful of sarsaparilla syrup. Or the *Pulv. Ferri* of the *Pharmacopœia* may be given, either in the form of powder, mixed with dry sugar, in pill, or in the shape of the chocolate lozenge. From half a grain to a grain, three times a day, is the proper dose from three or four years to six or seven. The carbonate of iron may be given, if it is preferred for any cause. Either of these preparations of iron, or any other that may be chosen, should be combined with a grain of quinine, three times a day, whenever the appetite is poor, and when the digestive process seems to be slow and feeble. Or the child may be made to take half a teaspoonful of the fluid extract of cinchona, mixed with an equal quantity of syrup of ginger, half an hour before the meals, while the iron is given alone soon after the meals. When the attack is particularly obstinate, and when also, it occurs in a subject who either inherits or exhibits signs of the tuberculous or scrofulous diathesis, the best remedy is cod-liver oil, which should be given in doses of from half a teaspoonful to a teaspoonful two hours after each meal. In cases of syphilitic nature, in addition to the above regimen and tonic remedies, we should administer the iodide of potassium, associated in obstinate cases with minute doses of bichloride of mercury.

The *local treatment* must consist in the use of means intended to keep the passages clean and free from scabs and incrustations, and in the employment of astringent and alterative applications. When the patient will submit, the nasal passages should be cleansed by means of a syringe once or twice a day, with tepid water, or milk and water, or with a weak solution of alum in water. The latter may be made in the proportion of from two to four grains to the ounce. If the discharges are offensive, the lotion used for injection should consist of the solution of chlorinated soda, one, two, or three drachms in two ounces of water. After the use of the syringe, and more or less frequently through the day, according to the disposition to dryness of the surfaces, these should be lubricated with some oleaginous application. One of the best is glycerin, or glycerin rubbed up with cold cream ($\frac{f\text{ʒj}}$ of the former to $\frac{\text{ʒj}}$ of the latter); or sweet oil, or oil of sweet almonds, may be used. These applications are best made by means of a camel's-hair brush. In older children, the nasal douche may be occasionally used with excellent results. There is little or no danger from its use, provided that the child can be made to breathe properly through the mouth all the while that the flow continues; that the liquid is used moderately warm; and that the vessel from which it flows is placed but a little above the level of the child's head.

Amongst the astringent applications, the best, are weak solutions (gr. ij to v to water fʒj) of the nitrate of silver, which should be used only once a day, or solutions of the sulphate or acetate of zinc with wine of opium. From two to five grains of either preparation, with a drachm of wine of opium, to an ounce of water, make a proper application. This may be applied twice a day. One of the best means that we know of, however, after the use of the alum or soda injection through the day, is to apply the following ointment at night: R. Ungt. Hydrarg. Nitrat., ʒss.; Ext. Belladonnæ, gr. x; Axungia, ʒss.—M. This has succeeded admirably well in several cases in which we have used it. It should be applied, after being completely softened by a gentle heat, on a camel's hair pencil, care being taken to apply it thoroughly to the surface of the mucous membrane itself, and not merely to the outside of the hardened scabs.

The following case well illustrates the severe form of chronic coryza. It was in all probability of syphilitic nature, though circumstances rendered it impossible to determine this question.

CASE.—The subject of this case, a male, was born after an easy, natural labor, and appeared strong and well, with the exception of a little discharge of blood from the nose soon after birth and slight coryza, the latter of which continued until the child was five weeks old, when it became aggravated, and one of us was requested to visit the infant. It was small and puny; the skin was harsh, dry, and wrinkled, so that the child looked like a little old woman. It was very weak, and had constant secretions from the nostrils of thick, dark-colored pus. When the discharge collected in sufficient quantity to obstruct the passages, the respiration became exceedingly difficult, as the infant seemed incapable of breathing through the mouth. At such moments it seemed as though the child must die of asphyxia. If the nostrils were cleared by any means, by syringing, by the use of a brush, or by blowing into them in the manner already described, the respiration would become easy and natural, until the discharge collected again, when the same scene recurred. During the paroxysms arising from the closure of the nasal passages, the child was entirely unable to take the breast, but after being relieved, had no difficulty whatever; the mouth was either kept shut, or if open, the tongue was observed to be pressed spasmodically against the roof of the mouth, so that it was impossible for more than a very small amount of air to pass over it; the respiration was labored, and accompanied by a loud snoring or nasal sound. There was no other marked symptom, except a nearly constant flatulent distension of the epigastric region. On the day before death, the infant seemed better, appeared to have gained flesh, and looked more intelligent, so that the mother was greatly encouraged; but the next day it was seized during one of the paroxysms of suffocation, which did not seem to be worse than many preceding ones, with copious discharges of bloody and frothy serum from the mouth and nose, and died in about three-quarters of an hour.

At the post-mortem examination we were not allowed to examine the nasal passages or throat. The stomach and bowels were healthy, but much distended with gas. The peritoneum was healthy, but contained a considerable amount of clear yellowish serum. There was serous effusion in both pleural cavities, but no traces of inflammation. The lungs were healthy, with the exception of some ecchymosed points, and general infiltration with sanguineous frothy serum. The trachea and bronchia were natural. The heart was larger than usual, but healthy in other respects.

SECTION II.

DISEASES OF THE LARYNX.

GENERAL REMARKS.

THERE has been much confusion amongst writers on the diseases of children, until within a few years past, in regard to the diseases of the larynx, each one differing from the other in his opinions as to the nature of the several disorders of that organ, and of course as to their classification and symptoms. From later and more rigid observation it has become clear, however, it appears to us, that there are but three diseases of the larynx which deserve to be considered as separate and distinct affections. These are simple erythematous or catarrhal inflammation of the larynx, unattended with spasm of the glottis, or, as that symptom has been emphatically named, laryngismus; simple catarrhal inflammation of the larynx, attended with laryngismus, and called most properly spasmodic simple laryngitis, or more commonly simple, false, spasmodic, or catarrhal croup; and lastly, pseudo-membranous inflammation of the larynx, properly named pseudo-membranous laryngitis, and more commonly called true or membranous croup. There is, moreover, another disease, of which one of the most marked symptoms is spasm of the glottis, or laryngismus, attended with a whoop or stridor, which is now known by the name of laryngismus stridulus, but which is called also Kopp's or thymic asthma, spasm of the glottis, and croup-like convulsion. This disease has often been confounded with the above-named affections of the larynx under the common title of croup, or has been supposed to constitute a distinct disease of the larynx; whereas now it is well known that the laryngismus whence its name was taken, is but one of many symptoms that mark the dependence of the disease upon disordered action of the reflex portion of the general nervous system.

We are well aware, also, that some most competent observers describe a purely spasmodic affection of the larynx, under the title of spasmodic croup, which they believe to be entirely independent of laryngeal inflammation, and to consist in a mere momentary contraction of the sphincter muscle of the larynx, produced by the sympathies which that part holds with other parts of the body, and especially with the digestive apparatus. As we have never, however, in what has now become a very considerable experience in the diseases of children, met with a case of spasmodic croup unconnected with more or less evident catarrhal inflammation of the larynx, we are not disposed to risk increasing the confusion already attending this subject, by making additional and more minute varieties of these affections than those above named. We are quite willing to acknowledge that, in some cases of simple spasmodic croup, the amount of catarrhal inflammation of the larynx is slight, and that the symptoms of digestive disorder are very strongly marked, but in not a single instance of croup that has come under our notice, have we ever had reason to suppose that the croupal symptoms were dependent solely on simple spasm of

the glottis (caused by some distant irritation), unattended with inflammation of the laryngeal mucous membrane. In all such cases that we have met with, it has seemed to us that the condition of gastric, intestinal, or bilious disorders, might be explained in one of two ways. Either the disorder of the digestive function has rendered the child unusually susceptible to cold, by having diminished its power of resistance to the weather; or, the derangement of the bodily functions caused by the cold has weakened, amongst others, the digestive system, and thus brought about various symptoms of gastric or intestinal disturbance, or more commonly of indigestion.

ARTICLE I.

SIMPLE LARYNGITIS WITHOUT SPASM.

DEFINITION; FREQUENCY.—This disease consists of simple erythematous or catarrhal inflammation of the mucous membrane of the larynx, unattended with spasmodic closure of the organ. It is sometimes attended with ulceration, but is unaccompanied by exudation of false membrane. The frequency of the disease, during infancy and childhood, is very considerable; so much so, that not a winter passes without our meeting with a good many well-marked cases.

PREDISPOSING CAUSES.—The disease occurs at all periods of childhood, but is much more frequent under than over five years of age. As to the influence of the seasons, it may be stated that it is by far the most common in the fall, winter, and spring months.

The only *exciting* causes of the disease which appear to have been ascertained with any certainty, are the action of cold, the positive influence of which cannot be questioned; the inspiration of irritating substances, such as gases, smoke, powders floating in the air, etc.; and violent efforts of crying. MM. Rilliet and Barthez state that they have twice known erythematous and ulcerative laryngitis to follow long-continued and violent crying; and M. Billard also cites this as a cause. We are acquainted with one case in which a slight attack of the disease appeared to have been brought on solely by loud and obstinate screaming.

The disease is very apt to occur in the course of other maladies, and particularly of measles, small-pox, scarlet fever, bronchitis, and pneumonia.

ANATOMICAL LESIONS.—The anatomical alterations may consist of simple inflammation of the mucous membrane, with its various effects, or of the same changes in connection with ulceration. The latter class of lesions is almost always confined to secondary cases. In the former class, the mucous membrane varies in color between a deep rose and violet red, which may be either uniform or only in patches. In severer cases, the tissue is at the same time softened or roughened, and sometimes thickened. When redness, softening, and thickening are present, the disease is generally confined to certain parts, and commonly to the epiglottis, and internal portions of the vocal cords; but when redness alone exists, it

usually affects the whole of the larynx, and sometimes extends to the trachea. In cases attended with ulcerations, these alterations exist in connection with those already described. The ulcerations are generally small, few in number, very superficial, linear in shape, and are almost always found upon the vocal cords. They are so slight often as to escape observation, unless a very careful examination be made; and this, perhaps, explains the circumstance of so few persons having met with them in the simple acute disease. Not unfrequently a certain amount of pharyngeal irritation is present at the same time, with or without some degree of tonsillitis.

SYMPTOMS; COURSE; DURATION.—The attack generally begins with an alteration of the *voice* or *cry*. In infants the change in the cry alone exists, so that to detect the disease, it is necessary to hear the child cry. In older children the same alteration of the cry is present, but there is in addition a change of the voice, consisting of various degrees of hoarseness. These symptoms may be so slight as to be observed in the cry only when it is strong and forcible, and in the voice so as to strike only the ear of one accustomed to be with the child; or they may be so marked as to be heard in the faintest cry that is uttered, and to be evident in the voice in a moment to the most careless observer; or there may be complete aphonia. They are often intermittent in this form, and are generally most marked in the after-part of the day and during the night. Simultaneously with this symptom, or very soon after, *cough* occurs. This is generally hoarse and rough, and early in the attack, dry; at a later period it usually becomes loose, and as this change occurs, loses its character of hoarseness. The frequency of the cough is variable, but usually moderate; as a general rule it is most frequent in the evening, and early in the morning, particularly in infants and young children. The disease is almost always preceded and attended with some coryza, which, in the early stage, is marked by sneezing and slight incrustations about the nostrils, and at a later period, by mucous and sero-mucous discharges. The *respiration* remains natural, except that it is sometimes nasal, and sometimes a little accelerated. There is rarely any fever, or it is slight, and occurs only at night. There is no pain in the larynx. In some cases, the hoarseness of the cry, voice, or cough scarcely exists, or is but slightly marked, and the only symptom is a dry, hard, teasing, and paroxysmal cough, which, from its sound, evidently proceeds from the larynx, and resembles very much that produced by the tickling of a foreign body in the throat.

The symptoms of this disease, instead of being of the mild character just described, may be much more severe. The cough is more frequent, hoarse, troublesome, and painful, from the scraping and tearing sensations it occasions in the larynx. The voice is more affected, becoming from husky more and more hoarse, though it is very unusual for it to become weak and whispering, as in membranous and severe spasmodic croup. The respiration is decidedly accelerated, giving rise to slight dyspnoea, and there is more or less fever, which is most marked usually in the after-part of the day and in the night. The pulse is more frequent than in health, rising to 120 or 130 in the minute; the skin is hot and dry; the

child is thirsty, restless, and uncomfortable. After a few days usually, the cough becomes loose and easy, and ceases to be painful; the voice loses its hoarse tone gradually, the fever disappears, the appetite and gayety return, and the child regains its usual health.

When the laryngeal inflammation becomes violent in this disorder so as to be attended with considerable swelling of the mucous membrane, the case nearly always, according to our experience, assumes the nature of grave spasmodic laryngitis. To our article upon this latter affection, spasmodic croup, we must refer the reader for further information on this point.

In nearly all the cases of this form of laryngitis that have come under our observation, we have found, upon examining the fauces, more or less decided inflammation of the tonsils, soft palate, and pharynx. In cases following a rather chronic course, from two to four or six weeks, which are rarely accompanied by fever or hoarseness, except at the invasion, and sometimes in the evening, the pharyngeal mucous membrane presented a roughened, thickened appearance, and the tonsils and uvula were more or less enlarged and tumefied.

There is a form of obstinate, troublesome cough, to which children are subject, and of which we have met with a good many examples, that must be noticed here. It depends evidently upon chronic inflammation, with thickening of the mucous membrane lining the upper portion of the larynx. This can be determined by laryngoscopic examination in the case of children of suitable age; but may be ascertained with confidence from the tone and character of the cough, from its occasional association with hoarseness of the voice, from its being almost invariably coincident with thickening and granulation of the pharyngeal mucous membrane, and from the fact that the most careful physical examination of the chest fails to reveal any disease whatever of the lungs. The cough is harsh, rough, and so to speak, tearing in its character. It often sounds, especially towards evening and in the early part of the night, croupal in its tone. It is usually very frequent, not so much, however, during the day, as in the evening and night. It is very generally increased by the horizontal position, so that when the child is put to bed, he will begin to cough violently and almost incessantly, and will continue to do so for one, two, and even three or four hours. The cough is so constant and so severe as to cause the greatest disturbance not only to the patient, who will toss and turn in bed in the most restless manner, but to the mother or attendants, who are excessively annoyed, and sometimes alarmed, by its constancy and obstinacy. Children who become subject to this species of cough, often have repeated attacks during the cold seasons of the year, the slightest exposure sometimes bringing them on. Each attack may last from a few days to several weeks. In one case we knew it to last, without once entirely ceasing, three months, and in another it lasted, with imperfect suspensions of a few days, during the same length of time. Both these cases occurred in hearty boys, one in the second, and the other in the third year of life, and yet both were vigorous and healthy children, as time has shown. In many other instances, we have known it to last two, three, and four weeks, proving all that time most troublesome and rebellious to treat-

ment. During the day, the child generally seems perfectly well, or at most merely a little pale and languid, and he coughs but moderately, but as soon as night comes on, and especially when he is put to bed, the cough begins and goes on for hours, as stated above, unless some remedy, and particularly some opiate, be given to check it. It is most annoying to the practitioner, for he finds that his usual remedies act merely as palliatives. They check and modify, perhaps overcome it for a time, but the next change in the weather, and especially the least exposure to cold and damp, start it afresh, and he has to resort again to the same round of treatment to subdue it. To the members of the family also it gives great anxiety. At first, they fear it must run into croup, which, however, it very seldom does, and then, finding how difficult it is to cure, and how often it recurs, they can scarcely be persuaded that it does not depend on some serious disease of the lungs.

The principal *cause* of this form of chronic laryngeal irritation is, so far as we have been able to ascertain, an unusual susceptibility of the laryngeal mucous membrane, sometimes the result of a congenital idiosyncrasy, and at other times the result of influences coming into action after birth, and especially of improper dress. We have generally met with it in children dressed upon the hardening system so much in vogue with many of our most highly educated citizens. The low frock, leaving the neck and upper half of the chest exposed to the air, the bare arms and bare legs, persevered in through our cold autumns, winters, and springs, have certainly, in most cases, been the cause of this troublesome and chronic cough.

Our experience since the publication of the earlier editions of this work fully confirms the truth of these remarks upon the style of clothing just referred to. We certainly do not see so many cases of obstinate laryngeal cough as we formerly did, for the simple reason that but few of the families we take care of, adhere to the old-fashioned system of leaving their children half naked.

The *duration* of the disease varies according to its form and the circumstances under which it occurs. When primary, it lasts usually from a few days to one or two weeks, but when it becomes chronic, as we have known to happen in a good many instances, it has lasted from two to four or six weeks, and even two or three months. The duration of secondary cases depends, of course, upon that of the disease during which they occur.

DIAGNOSIS.—The diagnosis of simple laryngitis is very easy. The hoarseness of the cry, voice, and cough, the redness of the mucous membrane of the pharynx, and the absence of general symptoms, will distinguish it from any other affection. In somewhat severer cases of this form, in which the cough is more frequent and harassing, the general symptoms more strongly marked, and the respiration somewhat hurried and oppressed, the attack may at first view present the appearances of bronchitis or pneumonia. The absence of the physical signs of these affections will show at once, by negative evidence, the true nature of the case.

In some cases in which there is little or no hoarseness of the voice or

cough, the symptoms strongly resemble the early stage of hooping-cough. We have met with quite numerous instances in which it was difficult not to believe, for two and three weeks, that the attack was really one of that disease. In one of these the resemblance was so close, that for several days there was a distinct hoop during the fit of coughing, with vomiting at the close of the paroxysm. The grounds for deciding that the case alluded to was not one of pertussis, were, that the attacks came on like laryngitis, after measles, and that the paroxysms occurred only at night. In the other cases a correct diagnosis was arrived at only by attention to the state of the fauces, which are almost always more or less inflamed and thickened in laryngitis, whilst they are not so in pertussis, and by watching the progress of the sickness.

PROGNOSIS.—The prognosis is always favorable in the mild form of the disease. We have never met with a fatal case. In the graver form of the disease, which will be more fully considered in the next chapter, the prognosis must be more guarded, although in our own experience no case of laryngitis without false membrane has proved fatal. It is, however, unquestionable that such a result may occur, so that in grave cases the greatest anxiety is justified.

TREATMENT.—The treatment of the milder cases of this form of laryngitis ought to be very simple. Children under four or five years old ought to be confined for the first few days to the house, unless the weather be dry and not intensely cold. In mild weather they may be sent out for a short time in the middle of the day. When the patient is five or over, he may continue to go out during the day, unless the weather be very bad. Much must depend upon the peculiarities of the child's own constitution. These can only be learned by observation on the part of the mother. Some children bear going out with such attacks perfectly well; others, if sent out with this simple laryngitis, are almost certain to have spasmodic croup or bronchitis more or less severely. When there is any febrile movement in the case, no matter how slight, the child ought to be kept quiet, and confined to the house. Attention to this point, therefore, careful management of the clothing, slight reduction of the diet if there be any fever, a warm foot-bath at night of simple water, or of water containing a little mustard, the application of some slightly stimulating liniment to the front of the neck and throat twice a day, and the occasional internal administration of some gentle expectorant and anodyne dose, constitute all that is necessary in the great majority of cases of this kind. The best internal remedies are a few drops of syrup of ipecacuanha, with paregoric, laudanum, or solution of morphia, given every evening as the child is put to bed, or occasionally through the day also, if the cough is troublesome. A combination of syrup of senega with that of ipecacuanha, will often be found very serviceable.

Without pretending that it is essential here, any more than in the case of coryza (see page 58), it is our habit to give quinia in cases of catarrhal laryngitis; and we believe that it will be found to hasten the resolution of the attack. It is of considerable importance that every acute sickness in young children, of however trifling a character, should be treated so as

to shorten its duration, and so as to leave the forces and tone of the system as little impaired as possible.

The treatment of the chronic laryngeal cough, unattended by fever or any severe constitutional symptoms, described above, requires some special remarks. In the first place, we have to state we have seldom succeeded in curing it until we had obtained from the parents their consent (often obtained with great difficulty) to a proper dress for the child. Expectorants, nauseants, opiates, antispasmodics, counter-irritants, and local applications, have nearly always failed to procure more than temporary alleviation, until the child has been dressed warmly. We have cured on several occasions, this kind of cough, after many ineffectual trials with the above remedies, only by insisting upon a mode of dress which covers the neck, arms, and lower extremities. A merino or soft flannel shirt, with long sleeves and high neck, long merino stockings, and thick muslin or canton-flannel drawers, have done more in such cases to effect a cure than all other means. This style of dress has removed the cause, the constant chilling of the body, and then the usual therapeutic measures will, no doubt, assist in overcoming the local changes which constitute the disease.

The best therapeutic measures to be adopted in such cases are the application, once a day, of a solution of nitrate of silver, of from five to twenty grains to the ounce, low down into the pharynx and chink of the glottis, by means of a small sponge-mop on a bent whalebone. After several applications have been made daily, they should be made only once in two or three days. The strength of the solution is to be determined by the condition of the pharyngeal mucous membrane, as we may assume this to mark, in some measure, the state of the contiguous tissue of the glottis. When the mucous membrane of the fauces is covered with large, protuberant follicles, when the tissue between the follicles is thickened, relaxed, spongy-looking, and when the color of the membrane is dark-red, the stronger solutions are the best; when, on the contrary, the mucous membrane is not roughened or thickened materially, when the follicles are small, when the color of the tissues is bright-red, it is best to use only the five-grain, or even a weaker solution. The most useful internal treatment in our hands has been the exhibition, three times a day, of a fluid-drachm of one of the following mixtures, diluted with a little water:

R. Potass. Carbonat.,	℥j
Tinct. Opii,	gtt. xxiv, vel xlvij .
Syrup. Senegæ,	f℥ij
Syrup. Tolutani,	f℥vj
Aq. Fluvial,	f℥ij
Ft. sol.	
R. Ammonii Bromidi,	gr. lx, vel xcv
Ammonii Muriatis, vel Potassæ Chloratis,	gr. xlvij
Tr. Opii Deodoratæ	gtt. xxiv, vel xlvij
Syr. Scillæ,	f℥ij
Elix. Calisayæ,	f℥j
Aquæ,	q. s. ad f℥ij.
Ft. sol.	

One of the most troublesome cases of cough we ever met with, occurred a few years since in a fine, intelligent, but not robust boy, four years old. He was seized with a hard, obstinate cough, which, in a few days, became really terrible from its almost incessant repetition for many hours at a time. The cough was dry, tickling, choking, repeated with nearly every breath, and so incessant as to drive the parents—and we may add the doctor—almost frantic. The little fellow at last found out, instinctively, that, by placing himself on the front of the body on two pillows, with the chin hanging over the edge of the upper one, he coughed less frequently, and with less violence, than in any other position. Discovering that the uvula was very much lengthened from relaxation and elongation of its mucous membrane, we touched the lower, sharp extremity with the lunar caustic stick twice a day. At the same time, the following mixture, which he had often used to control general nervous irritability in children, was prescribed; and this, with the lunar caustic application, finally controlled the cough. It was as follows:

R. Vin. Antimonii,	gtt. xlviii
Ext. Valerianæ Fl.,	
Tr. Opii Camph., āā	f ʒ ij
Syrup. Simp.,	f ʒ ss
Aquæ,	f ʒ j.—M.

Sig. A teaspoonful every hour or two when the fits of coughing set in.

After a few days, when the irritability of the fauces was somewhat subdued, the elongated portion of the mucous membrane of the uvula was cut off close to the muscle, and there was no renewal of the cough afterwards.

When the cough is very harassing at night, from two to four drops of laudanum, with from ten to twenty drops of syrup of ipecacuanha, or two grains of Dover's powder, given once or twice in the evening, have answered better than any other means. When the patient presents an anæmic appearance, or other symptoms marking a general deterioration of the health, iron, and especially the syrup of the iodide of iron, given three times a day, had assisted in removing the cough, and especially in lessening the extreme susceptibility of the system to changes of weather. The same good result has also followed the use of emulsion of cod-liver oil with lacto-phosphate of lime or with wild cherry bark. The diet ought to be light, but strengthening. Good fresh meat, with simple nutritious vegetables for dinner, and bread and milk morning and evening, constitute the most proper diet. In bad weather, during the cold seasons of the year, the child should be confined to the house.

ARTICLE II.

SPASMODIC SIMPLE LARYNGITIS, OR SPASMODIC OR FALSE CROUP.

DEFINITION; SYNONYMS; FREQUENCY; FORMS.—Spasmodic laryngitis is a disease of the larynx almost peculiar to children, consisting of simple catarrhal inflammation, without pseudo-membranous exudation, of the mucous membrane of that organ, attended with spasmodic contraction of the glottis, or laryngismus, occasioning violent attacks of threatened suffocation.

It is the disease commonly called in this country croup, or, since the distinction between it and pseudo-membranous laryngitis or true croup has been more generally recognized, spasmodic croup. It is known also by the names of false or pseudo-croup. We prefer the term spasmodic laryngitis, because it is expressive of the essential characters of the disease. It is the stridulous laryngitis of MM. Guersent and Valleix; the stridulous angina of M. Bretonneau; the acute asthma of infancy of Millar; and the spasmodic croup of Wichmann, Michaelis, and Double. It is not the laryngismus stridulus described by the English authors, Kerr, Ley, and Marsh, which is the same as the thymic or Kopp's asthma of the Germans, and spasm of the glottis of the French. It is called by Dr. Wood, in his work on the practice of medicine, catarrhal croup.

Spasmodic laryngitis is one of the most *frequent* of the winter diseases occurring during childhood in this country. It is so common in this city, that almost all mothers who have had any experience in sickness, keep some remedy for it in their houses, which they are in the habit of resorting to upon their own judgment.

We shall describe two *forms* or degrees of this disease, the *mild* and the *severe*. Without this distinction it would be impossible to give an accurate account of the disorder, since the two forms differ so much in aspect as to render them almost as much unlike as though they were two distinct affections. Moreover, the mild form differs so widely from membranous or true croup in its course and symptomatology, that the distinction between the two is readily made out, whilst the severe form, on the contrary, resembles true croup so much as to demand often very nice powers of observation to distinguish them, and yet the distinction is one of vast consequence to the patient, since the prognosis and treatment are widely different in the two diseases.

PREDISPOSING CAUSES.—The disease is much more common at some *ages* than others. It occurs most frequently during the period of the first dentition, being more common in the second year of life, which is the time of greatest activity of the first dentition, than at any other age, though it is often met with also in the third and fourth years. In the fifth year it still occurs occasionally, in the sixth and seventh it becomes rare, and after the seventh we have seen it but a few times.

It is said to be more frequent in boys than girls, and this seems borne

out by our own experience (since of 106 cases, 62 occurred in boys, and 44 in girls).

Spasmodic croup occurs usually as a *sporadic* disease, but is said by some authors to prevail at times as an *epidemic*. We have never had any reason to suppose that it was strictly an epidemic like membranous croup, which appears to a considerable extent in some years, and in others is scarcely seen. We believe rather that the unusual prevalence of spasmodic laryngitis at certain periods, in comparison with others, depends on the fact that certain states of the weather or season predispose or excite to it in a greater degree than usual, and thus occasion a large number of children to be attacked with it.

It is generally believed to be hereditary in certain families, and of this we ourselves have no doubt. We are acquainted with one family in this city, in which the children for three generations were extremely liable to it; with another, in which the grandmother and grandchildren were frequently attacked; and with a third, in which the father and children showed the same predisposition in the most marked manner. The idea is, moreover, entertained by many people in this community.

The natural constitution of the child does not seem to have much influence upon the liability to the disease; it occurs indifferently in the weak and strong. We have no doubt, however, that there are certain transient conditions of the health which do affect the liability to it, since it has long been remarked that disturbances of the digestive functions frequently invite it, and since we have often ourselves found it most apt to attack those who are liable to it, when they happen to be laboring under gastric catarrh or indigestions. It is common during cold and rare in warm weather.

EXCITING CAUSES; COLD.—By far the most frequent exciting cause is the action of cold; either the passage from a warm into a cold atmosphere, or prolonged exposure to cold and damp. It has been known on several occasions to follow long-continued crying, doubtless from inflammatory action set up in the larynx, as a consequence of the excessive determination of blood to that part during the act of crying. We were assured, some time since, by a very intelligent woman, that her little daughter had, at the age of two years, a well-marked attack of croup, after a severe and long-continued fit of crying from some contrariety.

ANATOMICAL LESIONS.—Mild cases of spasmodic laryngitis are so rarely fatal, as to leave us in some doubt as to the character of the anatomical lesions, or whether there are indeed any perceptible alterations of the tissues. We have never ourselves met with a fatal case of this form, and are therefore unable to give any personal account of the condition of the larynx, though we have never doubted, from the nature of the symptoms, the hoarseness, the dry cough, which afterwards becomes loose, and the whole aspect of the disease, that the anatomical condition of the affected organ must be one of slight catarrhal inflammation. In some cases, however, that have been examined, a little mucus in the larynx, and slight redness have been found, while in others no change has been detected. Dr. Wood (*Treat. on the Prac. of Med.*, vol. i., p. 779) accounts for this absence of morbid appearances in the following plausible manner: "In

some rare instances, no signs of disease are discovered in the mucous membrane, and the patient has probably died of spasm, consequent upon high vascular irritation or congestion, the marks of which disappear with life."

Cases of severe spasmodic croup have occasionally proved fatal, and the anatomical alterations of this form of the disease have therefore been well ascertained. These alterations consist of either simple catarrhal inflammation of the laryngeal mucous membrane, or of inflammation attended with ulceration. When the inflammation is simple, the membrane is changed in color, either uniformly or in spots, to a deep rose or dark-red tint. This may be the only alteration, or the tissues may be found also softened, or roughened and thickened. When the redness, thickening, and softening, are all present, these appearances are usually confined to certain parts, and particularly to the epiglottis and vocal cords, but when redness alone is present, it generally affects the whole of the larynx, and may extend to the trachea. To the alterations just described are sometimes added, as was stated above, ulcerations. These are commonly small, few in number, of a linear shape, and are usually seated upon the vocal cords. They are so slight as to escape observation, unless carefully looked for.

SYMPTOMS ; DURATION.—The invasion of the *mild form* of spasmodic croup is generally very sudden, for though it is often, probably in a large majority of cases, preceded for a few hours or a day or two by slight coryza, hoarseness, and cough, these symptoms are seldom noticed at the time, and the child is not supposed to be sick until seized with the paroxysm of suffocation, which is pathognomonic of the disease. This occurs in much the larger number of cases during the night, and very generally wakes the child from sleep. Of sixty-four cases observed by ourselves, in which the time of the attack was noted, it occurred in the night in sixty-two, whilst in two it came on in the afternoon. The period of the night at which it takes place is very irregular, but it is much more apt to be before than after midnight, as is shown by the fact that of forty-two cases in which this circumstance was ascertained, the attack was before midnight in thirty, and after in twelve. This agrees very closely with the statement of MM. Rilliet and Barthez, that it has been observed most frequently at eleven in the evening. The duration of the paroxysms varies considerably, and depends a good deal upon the treatment employed. They may last from a few minutes to several hours, but are seldom shorter than from half an hour to an hour. The number of the attacks also varies. In some cases there is but one, though very generally there are several. When the attack occurs early in the night, it is very apt to recur again towards morning, and, unless means of prevention are used, on the following night also, and even, though this happens much more rarely, on the third night. As a general rule, the first attack is the most severe.

When the paroxysm comes on, the child is wakened from sleep by the sudden occurrence of symptoms apparently of the most alarming and dangerous character. These consist of loud, sonorous, and barking cough ; of prolonged and labored inspiration, accompanied by a shrill and piercing sound, to which the term stridulous is applied ; of rapid and irregular

respiration, amounting often to violent dyspnœa, or seemingly impending suffocation; the child, alarmed and terrified at its condition, and at the fright of those around, its countenance expressive of the utmost anxiety, cries violently between the attacks of coughing, and begs to be taken on the lap, or sits up or tosses itself upon the bed, struggling apparently with the disease, which seems for the moment to threaten its very existence. The voice and cry are hoarse, and sometimes almost extinguished during the height of the paroxysms, but become distinctly audible, and often nearly natural, in the intervals between them; differing in this respect from pseudomembranous croup, in which they remain permanently hoarse or whispering. We have never heard, in this disease, the whispering voice and the short smothered cough of true croup. The face, head, and neck, are at first deeply flushed, and as the paroxysm becomes more violent, assume a dark livid tint, which afterwards passes into a deadly paleness, if the attack be long continued. These changes in the coloration depend upon the arrest of the respiratory function and a consequent partial asphyxia. The pulse is frequent during the paroxysm, and the skin sometimes heated. After a longer or shorter period, generally from half an hour to an hour, the respiration becomes more tranquil; the stridulous sound disappears entirely, unless the child be disturbed and made to cry, when it again becomes distinct; the cough is less frequent and less boisterous, and the child generally falls asleep. The attack is very apt to recur towards morning, as has been stated, and if not then, the following night. The patient often seems perfectly well the day after the first paroxysm, with the exception, perhaps, of slight cough. This is no reason, however, for supposing that the disease will not return in the course of the second night, which is almost sure to happen, unless measures be taken to prevent it. The cough generally continues for a day or two, but soon loses the peculiar character expressed by the term croupal; it becomes less frequent and more loose, and the child is commonly well again in two or three days. Sometimes, however, the cough lasts for several days, becoming gradually less frequent, until at last it ceases entirely.

There is very little fever in mild cases, for though the pulse is accelerated and the skin warm during the paroxysm, these symptoms disappear very soon after that is over. In more severe cases, on the contrary, there may be considerable fever, the pulse becoming frequent and full, and the skin hot. The febrile movement is most apt to occur after the first paroxysm, as a consequence, apparently, of the slight catarrh which remains after the attack.

In the few fatal cases on record, the paroxysms have generally become more frequent and more violent by degrees, and death has occurred from suffocation. In other instances, death has been the result of prostration, which itself has probably depended on imperfect hæmatosis.

Recurrences of the disease are very common, children sometimes having several attacks in a single winter. This is not the case in true croup. We have known but two children to have a second attack of that disease.

The *severe form* of spasmodic laryngitis may begin as such or result from an aggravation of the mild form; or the case may commence as one

of simple laryngitis without spasm of the glottis, and as the intensity and extent of the laryngeal inflammation increase, it may assume all the features of the form under consideration. Whatever be the mode of onset of the case, this form of the disease sets in with hoarse, frequent cough, difficult respiration, restlessness, and more or less violent fever, symptoms which almost always become severe for the first time at night, and usually between early evening and midnight; though, in some few cases, they make their first appearance during daylight, and this is very much more apt to happen in this than in the mild form of spasmodic croup. During the night the symptoms increase in severity; the respiration is frequent and difficult, and, after a time, attended with the stridulous sound in inspiration and expiration caused by narrowing of the glottis; the cough is hoarse, dry, and croupal, and unattended with expectoration; the voice becomes hoarse, and fever sets in, the pulse becoming full and frequent, the skin hot and dry, and the face flushed. These symptoms persist, with greater or less severity, throughout the night, while from time to time, they increase to such an extent as to seem to threaten suffocation, resembling then exactly the paroxysms described as occurring in the mild form of the disease. They usually subside, however, very decidedly towards morning, the breathing becoming easier, the stridulous sound less loud, or ceasing altogether, the fever diminishing, and the patient becoming in all respects much more comfortable. This amelioration of the child's condition often continues until the after-part of the day or till evening, when the same train of symptoms reappears. In other cases the disease scarcely subsides at all for two, three, or four days, but continues throughout the day and night to exhibit the same symptoms as have been described above. In cases of this kind, which are not rare, the disease assumes many of the alarming and dangerous characters of pseudo-membranous laryngitis or true croup, and it becomes very difficult often to distinguish between the two. If no favorable change take place, the dyspnœa becomes so violent as to threaten suffocation; the cough is rare and short; the voice is reduced to a mere whisper; the pulse becomes small, extremely rapid and thready; the countenance, at first livid and congested, assumes a pale, cadaveric appearance; the features are contracted; the child becomes comatose or delirious, and death may occur from slow asphyxia, or sometimes in an attack of general convulsions.

In favorable cases, on the contrary, the dyspnœa, and especially the stridulous sound, diminish; the cough becomes loose, less hoarse, and loses its croupal character; expectoration of mucous sputa takes place in older children, whilst in younger, the loose gurgling sound produced by the discharge of the sputa into the fauces, is heard at the termination of each cough; the voice becomes clearer and stronger; the fever diminishes; the child regains its spirits and disposition to be amused; and soon all dangerous symptoms have disappeared, and the recovery is established.

In nearly all the cases that have come under our observation, we have found, upon examining the fauces, more or less decided inflammation of the tonsils, soft palate, and pharynx.

The *duration* of the *severe form* of spasmodic croup depends on the vio-

lence of the attack, and on the mode of treatment. When the treatment is begun at an early period, the disease is much sooner overcome than when allowed to run on for some time without remedies. In cases of moderate severity, the violence of the symptoms usually subsides after thirty-six or forty-eight hours. In more violent cases, on the contrary, the symptoms seldom subside definitively before the third, fourth, and not unfrequently the fifth day. In no case that has come under our observation, has the disease continued to present dangerous symptoms after the fifth day, unless, as not unfrequently happens, the inflammation spreads to the bronchia or tissue of the lungs, producing bronchitis or pneumonia. But even after the signs of severe laryngeal inflammation have disappeared, there almost always remains for several days longer, some cough and huskiness of the voice, showing that the mucous membrane of the larynx has not yet regained completely its healthy condition. The disease is said to have proved fatal in twenty-four hours.

NATURE OF THE DISEASE.—Although by the older writers, spasmodic simple laryngitis was confounded with membranous laryngitis, and this error continued to confuse the minds of medical men until a recent period, there is no longer any doubt as to the totally distinct character of these two affections. The comparative fatality of the two diseases alone is sufficient to establish a wide difference between them. Thus, of 35 cases of the pseudo-membranous form that we have seen, 16 died; while of considerably more than 200 cases of the spasmodic form that we have seen, not one has been fatal. M. Guersent states that of ten cases of the former disease, scarcely two escape; while of upwards of a hundred of the latter that he has seen, not a single one was fatal. (*Dict. de Med.*, t. ix., p. 365.)

The different effects of treatment in the two affections also point to a wide difference in their nature. True croup is almost inevitably fatal, unless attacked at an early period by energetic remedies, while the mild spasmodic form seldom resists the exhibition of an emetic, a warm bath, or of nauseating doses of ipecacuanha; and the severe form, though of a most threatening appearance, almost always yields to prompt treatment. When we add to these circumstances, the differences in the anatomical alterations in the two diseases, the difference in the mode of invasion, in the cough, voice, cry, fever, duration of the attack, and state of the constitution, all of which will be carefully described in the remarks on diagnosis, it is impossible to resist the conclusion that they are two distinct disorders.

We believe, therefore, that mild spasmodic laryngitis is a disease consisting in slight catarrhal inflammation of the mucous membrane of the larynx, attended with violent spasmodic contraction of that organ, or, as that condition has been called, laryngismus. The spasm of the laryngeal sphincter seems to be the result of a disordered action of the excito-motor innervation of the part, the irritant, which is productive of the morbid innervation, being, in all probability, the inflammation of the laryngeal mucous membrane which has been already stated to constitute one element of the malady. The nervous element predominates in the early part of the attack, but towards the conclusion, the spasmodic symptoms disappear entirely, and we have left only those which depend on the local tissue-changes.

In severe cases of the disease we have the same element of laryngeal spasm, or laryngismus, coincident with, and produced by, a much more intense and dangerous inflammation of the mucous membrane of the part than exists in the mild form.

DIAGNOSIS.—Unquestionably the disease with which spasmodic laryngitis is most likely to be confounded is pseudo-membranous laryngitis, or true croup. There is very little difficulty, however, in distinguishing the mild form of spasmodic croup from true croup, whilst in regard to the severe form, it may be safely stated, that the distinction cannot, in some cases, be made with positive certainty, except by watching the course of the sickness.

Mild cases of spasmodic croup may be distinguished from membranous croup by a comparison of the different symptoms as they arise. The most important of these are: the invasion, in one sudden and almost invariably in the evening or night, in the other slow and creeping, the paroxysm first occurring indifferently day or night; the cough, in one hoarse and boisterous, in the other hoarse and frequent at first, but rare and smothered towards the end; the voice, in one hoarse, but never scarcely whispering, and if so, only during the height of the paroxysm, in the other hoarse at first, and soon permanently whispering or entirely lost; the cry, in one hoarse and stridulous only at the moment of the paroxysm, in the other permanently so; the respiration, in one stridulous and difficult only during the paroxysm, and in the interval perfectly natural, in the other, at first natural, becoming by degrees permanently stridulous, and attended by the most violent dyspnœa, with remarkable prolongation of the expiration, and even with recession of the base of the thorax in inspiration; the fever, in one very slight and generally observed only during the nocturnal paroxysm, in the other much more considerable and permanent; and lastly, the duration, in one seldom more than two or three days, in the other rarely less than six, and very often eight or ten days. M. Trousseau states that the *hoarse-sounding croupal* cough is not a sign of the presence of exudation in the larynx, but rather of its absence; but, “when the cough, croupal at first, becomes less and less frequent, and ends with being nearly insonorous with suffocation, there is true croup, that is to say, with plastic exudation in the larynx.” This is precisely our own experience. The rare, insonorous cough of M. Trousseau is the condition which we have expressed by the term smothered.

In order to render the diagnosis still clearer, we add the following table, which is altered from one given by MM. Rilliet and Barthez:

MILD SPASMODIC LARYNGITIS.

Begins with coryza and hoarse cough, or more frequently with a sudden attack of suffocation in the night. Fauces natural, or merely slight redness, as in simple angina.

PSEUDO-MEMBRANOUS LARYNGITIS.

In epidemic form, begins as pseudo-membranous angina. In sporadic form, invasion of slight hoarseness for a day or two. There is fever, increase of the hoarseness, with hoarse, croupal cough; in most of the cases, pharyngeal exudation, and a little later, paroxysms of suffocation.

MILD SPASMODIC LARYNGITIS.

After the paroxysm, the child seems well, the fever disappears, or is very slight. Voice natural, or only slightly hoarse; not whispering.

If the paroxysm returns, it is during the following night, and it is less severe, the hoarseness disappears; the cough becomes loose and catarrhal.

Duration seldom more than three days.

Very rarely fatal.

PSEUDO-MEMBRANOUS LARYNGITIS.

The fever continues; stridulous respiration; prolonged and difficult expiration; recession of base of thorax during inspiration; cough hoarse and smothered; voice hoarse and whispering.

The dyspnoea and suffocation increase the voice and cough are smothered; or extinguished; stridulous respiration persists.

Duration seldom less than five or six days. The hoarseness continues for several weeks.

Fatal in the majority of the cases.

The only real difficulty in the diagnosis is the distinction between the grave form and pseudo-membranous laryngitis or true croup unconnected with angina; and this, it would appear from all evidence, cannot in some cases be made with absolute certainty. The only certain and indubitable sign by which to distinguish them is the presence of false membranes in the expectoration. The existence of this symptom is proof positive of pseudo-membranous disease, but its absence is no proof that the case must be one of simple inflammation; for, even though the membrane has been exuded in large quantities within the larynx, it is not always thrown off by the effort of coughing or vomiting. To show the difficulty of the diagnosis, we will cite the case quoted by M. Valleix (*loc. cit.*, t. i., p. 211) from M. Hache, of a child supposed to be laboring under true croup, who was sent to the Children's Hospital in Paris, in order to have the operation of tracheotomy performed. The absence of false membrane in the expectoration, and a slight remainder of clearness in the voice, occasioned the suspension of the operation. The child died, and no pseudo-membrane whatever was found in the larynx. The only lesions were moderate redness of the mucous membrane, without tumefaction, and without narrowing of the glottis; so that the fatal termination must be ascribed to spasmodic constriction of the glottis, or to tumefaction of that part, which had disappeared after death.

Nevertheless, though the diagnosis is difficult, it can almost always be made out with certainty by attention to the following points. The pseudo-membranous form of the disease is usually preceded or accompanied by the presence of false membranes in the fauces, which is not the case in spasmodic simple laryngitis; the symptoms of invasion of the former disease are less acute than those of the latter, the fever being less violent, and the restlessness and irritability less marked, than is usual in the simple affection, in which the general symptoms are decided from the first. The hoarseness of the voice and cough follow a different course in the two diseases; the progress of these symptoms being slow and gradual in the membranous, and much more rapid in the severe spasmodic form. The fever is marked throughout the attack in the severe spasmodic disease, whilst in the other form it seldom reaches a high degree of intensity. Albuminuria is present in a considerable proportion of cases of pseudo-

membranous croup, while it is habitually absent, or at most very rarely present, in cases of simple laryngitis, even of the most severe type. Lastly, the presence of portions of false membrane in the expectoration, in connection with the laryngeal symptoms, affords positive evidence of the existence of true croup.

Of the characters just enumerated as likely to aid us in distinguishing between severe spasmodic and true or membranous croup, we wish to call the reader's attention in greater detail to two,—the condition of the voice, and the stridulous respiration. The former is, we have no doubt, much the most important single symptom. In membranous croup, the voice begins by being hoarse, but soon becomes weak, so that after the disease has lasted three or four days, it changes from hoarse to whispering; it becomes, in fact, suppressed. In severe spasmodic croup, the voice is hoarse at first, and becomes more so as the disease goes on, but it very rarely becomes whispering as in true croup, but almost always retains a good volume, so that when urged the child can speak out loudly. Now this is never the case in the membranous disease, for, as the fibrinous exudation is deposited on the vocal cords and in the ventricles of the larynx, it suspends almost entirely the functions of those parts, and the voice is more or less completely suppressed. The remarks just made in regard to the voice will apply also to the *cry*, which should be carefully studied in young infants.

The second very important symptom is the stridor. This is, as might be expected, more marked in all its features in true than in false croup, since in the former it depends on a permanent and considerable obstacle to the passage of the air through the larynx. That tube is, in fact, completely coated over upon its internal surface with a more or less thick false membrane, which reduces materially its calibre, and impedes to a greater extent the passage of air, than does the more inflammatory tumescence and swelling of the mucous membrane of the organ in severe spasmodic croup. On this account, therefore, the stridor in the respiration is louder, shriller, more persistent, more marked in the expiration, and attended with greater effort of the respiratory muscles to overcome the obstacles to the passage of the air in membranous than in severe spasmodic croup. We may add that there is something very peculiar in the cough in true croup. When the membrane has come to cover the interior of the larynx, the cough is very distinctive; it has a sound which we can describe only by saying that it always reminds us of the sneezing of a young kitten. This we have never heard in catarrhal croup, no matter how severe.

To conclude, there is in membranous croup a slow, steady, and unrelenting progression of the symptoms, which is not observed in the spasmodic disease. From hour to hour, from day to day, we can perceive, so to speak, from the gradual and steady march of the disease, that a foreign body in the form of a fibrinous moulding is being spread slowly over the cavity of the larynx. In severe spasmodic croup, on the contrary, the course of the symptoms is less regular; paroxysms of suffocation occur as in true croup, but when these are over, the child is often quite comfort-

able; the symptoms indicating a much less considerable permanent mechanical obstruction than in the other affection.

Spasmodic laryngitis has been mistaken also for laryngismus stridulus. The manner in which it is to be distinguished, will be described in the article on that disease.

PROGNOSIS.—Spasmodic catarrhal laryngitis is very rarely a fatal disease. Of its two forms, there can be no doubt that the severe is much more dangerous than the mild, since in the former the patient labors under acute inflammation of the larynx, as well as under spasm of that organ; whilst, in the latter, the amount of inflammation is so very slight as to be of little or no consequence, were it not associated with the laryngismus, which gives to the disorder its most characteristic features.

Of 109 cases of the disease of which we have kept an accurate record, none proved fatal, though 23 of these were of the grave form. We may state, also, that we have seen at least 150 more cases, of which we have no written account, in none of which was there a fatal termination. We have, therefore, never seen a case of croup without false membrane prove fatal. That it does sometimes end unfavorably, however, cannot for a moment be questioned. There are various examples of the kind scattered through the medical journals. MM. Rilliet and Barthez quote in proof of this, two cases from the work of Jurine, in one of which an autopsy was made, and no false membrane discovered. Copland (*loc. cit.*) remarks, that in the few cases of the more purely spasmodic forms that he has had an opportunity of examining, an adhesive glairy fluid, with patches of vascularity on the epiglottis and larynx, and a similar fluid in the large bronchi, were the only alterations observed.

Great imminence of danger in any case is shown by a high intensity of the stridulous sound, especially as heard in the expiration; by great severity of the dyspnoea or suffocation; by permanently whispering voice; by lividity or extreme paleness of the face; by smallness and rapidity of the pulse; by coldness of the extremities; and by delirium or convulsions.

In giving our own experience in regard to the treatment of this disease, we shall first speak exclusively of the mild, and then of the severe form, since the measures proper and necessary in the one, are very different from those called for in the other.

TREATMENT OF THE MILD FORM—*Emetics*.—The great majority of cases will recover perfectly well under the use of emetics employed alone, or in combination with warm baths and revulsives. Of late years we have often succeeded in warding off the slight attack, where there has been good reason to expect it, by the administration of an opiate with syrup of ipecacuanha, at bedtime (early in the evening). At two years of age, two or three drops of laudanum, with ten to twenty drops (according to the gastric susceptibility) of syrup of ipecacuanha; at three or four years, four drops of laudanum with twenty of the ipecacuanha, are about the proper doses. Even when the child has had one attack early in the night, the use of the opiate is most successful, after vomiting, in

preventing the usual return towards morning. If the physician is not called until the day after the first attack, this treatment is excellent in the evening of the second day. In cases attended with violent dyspnœa, hoarse cough, and loud stridulous respiration, the emetic should be given until it produces a full effect. In milder case, in which there is merely loud croupal cough, with an occasional stridulous sound, nauseating doses alone will generally suffice. The most suitable emetic is, as a general rule, ipecacuanha. The best preparation for children is the syrup, of which from twenty to thirty drops may be given to those two years of age, to be repeated every ten or twenty minutes until vomiting is produced, or until the paroxysm is relieved. In very sudden cases, the Syrupus Scillæ Compositus, which is more active in its effects in consequence of the tartar emetic which it contains, might be preferable; about twenty drops of this may be given, and repeated every ten or fifteen minutes, until vomiting or the resolution of the paroxysm is obtained; but, in its employment, care should always be observed not to continue it for too long a time, lest it produce the injurious effects of tartar emetic. Of late years we have almost entirely abandoned the use of the latter emetic, as we succeed perfectly well with the ipecacuanha, and dislike more and more the antimonial preparation in children. When the dyspnœa is very urgent, or when other means fail to produce emesis, we have found nothing so effectual as powdered alum, in doses of a teaspoonful mixed with honey or molasses. (See *Treatment of Pseudo-membranous Laryngitis*.)

A simple and good method of treating the paroxysm is that recommended by Dr. Charles D. Meigs, in the paper referred to. It is to direct a small teaspoonful of powdered ipecacuanha to be diffused in a wine-glassful of water, of which mixture doses of a teaspoonful are to be given every ten, fifteen, or twenty minutes, according to the urgency of the symptoms. This is a plan of treatment often resorted to by parents in this community, where the disease is so common and so well understood, that there are few mothers who have several children, and who have had some little experience, who do not know how to treat a nocturnal attack of mild spasmodic laryngitis.

A very simple and efficient mode of treating the paroxysm, which was first recommended by Graves, consists in gently pressing a sponge soaked in warm water under the chin and to the front of the neck. This may be repeated every ten or fifteen minutes, and under its influence the croupy symptoms will often promptly subside without the use of an emetic.

After the paroxysm is relieved, it is a good plan to direct five or ten drops of the syrup of ipecacuanha to be given every two or three hours during the following day; or, if the child seems perfectly well in the morning, we may begin with these doses in the middle of the day, and continue them until bedtime. By this method the recurrence of the paroxysm during the second night may, we think, often be prevented, and the cough is rendered free and loose much sooner than when the disorder is left to pursue its natural course. Moreover, the child ought to be kept in the house during the next two or three days, or until the cough is thoroughly loose and

easy. If the child be at all a delicate one, or one in whom the disorder is prone to be obstinate, there is no plan so good as to make it sit or lie quietly in bed, sufficiently covered, with a large abundance of playthings, or with a kind nurse to read to and amuse it for two or three days.

Baths.—The warm bath is a very prompt and useful remedy in this disease. In all very violent cases, it ought to be resorted to immediately. It should be used also whenever the emetic fails to relieve the urgency of the symptoms, and in cases attended with much disturbance of the circulation. The temperature of the water ought to be about 95° Fahrenheit, when the child is first immersed, to be raised gradually by the addition of hot water, to 100° or 102°. The child may remain in the bath from ten to twenty minutes.

Revulsives.—The only revulsives that it can be necessary to employ are mustard foot-baths, or mustard poultices applied to the interscapular space, and even these are often needless if the emetic be given. Blisters, which are recommended by some of the French writers, can only be proper in rare cases of the grave form.

Purgatives are required when constipation is present, or when there is so much fever on the second or third day, as to show a considerable amount of laryngeal inflammation. Under the latter circumstances some mild remedy of this class, such as castor oil, may be resorted to with a view to its evacuant effect. We have never had occasion to employ any of the *mercurials*, and believe them to be unnecessary.

Opium is exceedingly beneficial when the emetic, nauseant, or warm bath has failed to relieve entirely, and when a troublesome croupal cough continues after the spasm has been overcome. Laudanum, paregoric, or solution of morphia, in combination with syrup of ipecacuanha, or Dover's powder alone, are the most suitable preparations. It is a very good plan to give the child a moderately full dose of the opiate, with ipecacuanha, after the violence of the paroxysm has subsided. It puts the child to sleep, promotes perspiration, softens the cough, and tends to prevent the return of the spasm. Repeated once or twice early in the second night after the first attack, we believe it often assists materially to avert the recurring nocturnal paroxysm.

TREATMENT OF THE SEVERE FORM.—This form of spasmodic laryngitis requires more active measures than the mild form of the disease.

In some of the former editions of the work, *bloodletting* was recommended when the disorder occurred in robust and vigorous children, and a record was given of the employment of venesection in seven out of twenty-three cases, all of which recovered. Since that report we have learned that depletion is less necessary than we formerly supposed, and as we can still say that we have never yet seen a fatal case of spasmodic croup, either simple or severe, it is fair to conclude that the disease can be safely managed without a resort to this more violent measure. Still, it is but proper to state, that should a case occur to us in a strong and healthy child, in which the breathing should become so much obstructed as to cause deep and alarming venous stasis, and in which these symptoms re-

sisted the more simple means we now employ, we should not hesitate again, as in former years, to employ venesection to the extent of four ounces at the age of four or five years, or a leeching to the same amount.

Our favorite remedies of late years have been *emetics*, *opiates*, *antispasmodics*, and *salines*. Among the combinations that we use most frequently and with the best results, may be mentioned the following:

R. Potass. Citrat.,	3i.
Syrupi Ipecacuanhæ,	f3ij.
Tr. Opii Deodorat.,	gtt. xij.
Syrupi Simp.,	f3ij.
Aquæ,	f3iss.—M.

Dose for a child two years old, a teaspoonful every two hours.

Or,

R. Ammonii Chloridi,	gr. xxiv.
Ammonii Bromidi,	3i.
Syrupi Ipecacuanhæ,	f3iss.
Syrupi Zingiberis,	f3ijss.
Aquæ,	f3iss.—M.

Dose for a child two years old, a teaspoonful every four hours.

Another combination that we have used with excellent effect, especially in cases where examination of the throat (which should be made every day without fail through the earlier stages of these cases) shows redness and swelling of the tonsils, is the following:

R. Potassii Chloratis,		
Ammonii Chloridi, āā	gr. xxiv.
Mist. Glycyrrhizæ Comp.,		
Syrupi Simp., āā	f3i.—M.

Dose for a child two years old, a teaspoonful every three or four hours.

In older children, both the saline and opiate must be suitably increased. In all these cases an emetic ought to be given once, or two or three times in twenty-four hours, when the dyspnœa and stridor become very severe; and in about an hour after its operation, the saline dose should be resumed. Of course, if decided drowsiness supervene from the opiate, the doses must be given at longer intervals. The emetic treatment is not so essential as in true croup, where it is so useful in causing the rejection of the false membrane which obstructs the larynx. Yet it is exceedingly useful, and often indispensable, in assisting to expel the viscid mucus secreted within the larynx, and in relaxing, for a time at least, the spasmodic constriction of the glottis, which plays a most important part in the production of the distressing dyspnœa and suffocation of the disease. They act probably also by lessening immediately, or through their action on the circulatory and nervous systems, the inflammation of the larynx. For their choice and mode of administration, the reader is referred to the article on true croup.

The cough sometimes assumes, especially in children of nervous type, a spasmodic character, resembling not infrequently whooping-cough. It is apt, in such cases, to be very frequent, particularly in the evening and

early part of the night. Here the combination of belladonna and alum which we employ in true whooping-cough is often most beneficial. The formula, for children of two or three years, is as follows:

R. Ext. Belladon.,	gr. i.
Pulv. Aluminis,	3i.
Syr. Acaciæ,							
Syr. Zingiberis,							
Aquæ, āā	f3i.—M.

Dose. A teaspoonful morning, noon, and evening, and once in the night, if necessary. At one year of age the belladonna should be reduced one-half.

Another excellent combination when the cough is frequent and harassing, is the following:

R. Tr. Belladon.,	gtt. iv.
Tr. Opii Camph.,	gtt. xlvij.
Pulv. Aluminis,	gr. vi.
Syr. Acaciæ,	f3ss.
Aquæ,	f3iss.—M.

Dose at six months, a teaspoonful every two or three hours.

The mother should be told to look at the pupils of the child after three or four doses have been given, and should they be at all dilated, to suspend the medicine for some hours, and then use it again. In children of two or three years of age a teaspoonful and a half may be given.

Purgatives are required merely to keep the bowels soluble; they should be repeated as may be necessary throughout the disease. If the bowels are moved every day or every other day spontaneously, there is no use in giving them at all. The most suitable are castor oil, rhubarb, or magnesia, in small doses; or an enema may be given from time to time if the child does not resist its exhibition.

Expectorants are useful after the violence of the disease has been moderated by more energetic remedies. They may consist of small doses of ipecacuanha, of antimonial wine and sweet spirits of nitre, of decoction of senega, snakeroot, or of the citrate or carbonate of potash.

Opiates and antispasmodics are necessary, and are serviceable, as has already been stated, in calming excessive restlessness, and in allaying the violence of the suffocative attacks, which depend, in good part, on spasm of the glottis. The most suitable are Dover's powder or some other preparation of opium, or small doses of belladonna, or hyoscyamus.

Belladonna would seem, from its power to relax the sphincters, and from its excellent effects in whooping-cough, to be indicated in this disease, but we have succeeded so well with opium that we have not often used it. Probably a combination of the two would be found beneficial.

Since the introduction of the bromide salts, the combination of the bromide of potassium, or of ammonium, with an opiate and a saline expectorant has been found very advantageous.

Counter-irritants.—During a paroxysm of dyspnoea in grave, as in mild cases of spasmodic laryngitis, some relief may be obtained from the application of mustard plasters between the shoulders, or over the sternum, and

of a sponge wet with hot water over the larynx. But as there is in these cases a more decided inflammation of the mucous membrane of the larynx and pharynx, it is desirable to use continuous mild counter-irritants for several days. Blisters are of doubtful propriety in any case. The application which we most frequently use is tincture of iodine, diluted with an equal amount of alcohol, which may be painted once or twice a day behind the angles of the lower jaw and all over the larynx, care being taken not to cause too much irritation of the skin. It is well also that a thin layer of new cotton or wool should be kept over the larynx.

A warm *bath* at 97° or 98°, employed once or twice a day, and continued for a period of ten or fifteen minutes, often assists greatly in lessening the sufferings of the child, in calming restlessness, and in moderating the heat of the skin, and violence of the circulation, when the latter symptoms are strongly marked. The same effects may often be obtained, though in less degree, by the use of warm foot-baths, with or without a little mustard in them.

HYGIENIC TREATMENT.—In either form of the disease the child should be placed for the time in a warm room, and warmly clothed. If old enough, it should be kept as much as possible in bed during the paroxysm. If so young as to prefer the lap of the nurse, it should be clothed in a long loose wrapper in addition to its usual night-dress. It is very important to confine the child during the whole term of the acute period in bed, if it is over three or four years old, and in the crib or lap if it be younger. Even after the cessation of the acute condition, it ought to be kept in one room for a few days, in order to make sure of the convalescence. The diet must be simple and of easy digestion, so long as there is any disposition to the recurrence of the disease. It may consist of preparations of milk, of bread, rice, or of thin chicken or mutton-water. Meat and most vegetables had better be avoided until the convalescence is fairly established.

PROPHYLACTIC TREATMENT.—It is certain that much may be done by a wise attention to physical education, to prevent attacks of the disease in children who show a liability to it. We would strongly recommend, with this view, attention to the following advice given by M. Guersent, who says (*loc. cit.*, p. 381): “It is possible, to a certain extent, to prevent attacks of pseudo-croup, if we fortify the constitutions of children, by exposing them well-clothed to a dry and elastic atmosphere, particularly if they can be kept in constant movement. But of all the precautions which have been found unquestionably advantageous, that which seems most useful is to make them sleep in well-ventilated, dry, carefully closed chambers, having a southern exposure, and always without fire. We have several times been convinced of the utility of this habit in families, the children of which were subject to this kind of catarrh.”

There can be no doubt that the style of dress used for children in this country must occasion many and repeated attacks of croup which might just as well have been avoided. The custom is to dress children between the ages of one and four or five years in such a way as to expose to the air the whole of the neck and the upper half of the thorax (for the dresses are made so low and loose at the shoulders as to leave the upper part of

the chest virtually uncovered). The arms are left bare, as are also the legs from the knee, or above the knee, to the ankle, so that very nearly half of the cutaneous surface is without covering, and this too, in the very same rooms and temperature in which sit the parents with the body and limbs warmly clothed to resist our climate, at all seasons changeable and uncertain, and, in the winter, very cold. We are perfectly well convinced that this faulty and unreasonable system of dress, which is chosen because it is fashionable, or in order to *harden* the child, who, however, invariably puts on warm clothing when it comes to years of discretion, will explain in part the enormously greater frequency in children than in adults, of the various diseases of the air-passages and lungs produced by cold.

One of the most important means of prevention, therefore, is the adoption of a suitable dress. In winter this should consist of one that shall cover the body completely. If the child be at all delicate, it ought to wear next to the skin a woollen jacket with long sleeves, and covering the chest to the neck. Over this should be put a long-sleeved stout muslin dress, or one of some light woollen material, made in the same style. In young children, the stockings ought to be of wool, and should reach to the knees; in older ones, they may be shorter, but the legs should be covered with drawers made of canton-flannel, of thick cotton stuff, or of light woollen flannel. To show the influence of dress, Dr. Eberle mentions the fact that in the country, and especially amongst the Germans, who cover the neck and breast, croup is a very rare disease. During a practice of six years amongst that class of people, he met with only one case of the disease.

When the liability to the disease continues after the completion of the first dentition, we have found the daily use of the cold bath, followed by brisk rubbing, so as to insure perfect reaction, in connection always with warm clothing, most useful in preventing the attacks. The bath must be commenced with in the summer, and persevered in during the following winter. The water, after the cold weather begins, should be drawn in the evening, allowed to stand all night in a room in which there is a fire though the day, and made use of on the following day. Prepared in this way, we have found the water in the morning at a temperature of between 50° and 60° F. The child ought to be kept in the water only half a minute or a minute, then well rubbed, and dressed immediately.

When the child is pale, weak, and feeble, and unable to bear exposure to the outer air, it may generally be restored to much better health by careful attention to diet, and by the steady and long-continued use of some tonic remedy. The diet ought to consist of bread and milk, and of meat and the simpler vegetables, as potatoes and rice. The tonics most generally suitable are quinine or iron. Of the quinine a grain may be given in pill or solution, twice or three times a day; while at dinner or lunch, or at both, the child should be made to drink from a dessert to a tablespoonful of port wine, mixed with water. This method ought to be steadily persevered in for from three to six weeks or longer. If quinine be objectionable for any reason, iron must be substituted. The best preparations are the iodide or the reduced iron.

ARTICLE III.

PSEUDO-MEMBRANOUS LARYNGITIS, OR MEMBRANOUS OR TRUE CROUP.

DEFINITION AND SYNONYMS.—Pseudo-membranous laryngitis is an acute inflammation of the mucous membrane of the larynx, attended with the exudation of false membrane.

It is the croup of the French writers, while, in this country, it is called by the various names of slow, creeping, true, membranous, or inflammatory. The term given above seems most suitable, as expressive of the real nature and seat of the disease, and we shall, therefore, make use of it in contra-distinction to that of spasmodic laryngitis or spasmodic or false croup, which is a much more common and less dangerous affection.

NATURE AND RELATIONS.—Of recent years the questions of the essential nature of membranous croup and its relations with diphtheria have been actively discussed, and it has appeared that there are marked differences in the opinions held by the best authorities. In the present state of the discussion, it is improper to attempt any dogmatic assertions on the points at issue, but it seems desirable to restate fully the views we have long held and the considerations on which they are based.

In the first place, there should no longer be tolerated the confusion that has grown up in regard to the very terms employed. It should be unanimously resolved that the terms croup and diphtheria shall hereafter be used as expressing either clinical conditions or anatomical processes. But at present, while many understand by these words definite diseases, others (especially of the German school) apply them to certain anatomical conditions wherever and in whatever clinical relations found. Thus, while one employs "diphtheria" to indicate a specific zymotic blood-disease, and "croup" to indicate acute laryngeal obstruction occurring with febrile symptoms, another will speak of diphtheria and croup of any mucous membrane according to the peculiar anatomical conditions present. We have always urged the adoption of the former clinical definitions, as the only ones that can possibly aid us in reaching clear and intelligible views on this important question; and we trust that their employment in this sense may soon become universal.

By many authorities, true croup is regarded as an idiopathic primary inflammation, presenting the unusual result of pseudo-membranous exudation, and differing thus from diphtheric croup, which is a mere complication in the course of a constitutional disease, depending upon the extension of the false membrane from the fauces into the larynx.

The considerations upon which this distinction has been based may be enumerated as follows: 1, that there are positive differences in the morbid processes present in the two diseases; 2, that croup is a disease peculiar to childhood, commencing in the larynx, and though it may pass down into the trachea, never passes upwards into the pharynx; 3, that it is not attended with enlargement of the cervical glands; 4, that it is a local, non-contagious disease, of a sthenic inflammatory type, without any special alteration of the blood crisis; 5, that it does not present the complications

of diphtheria, such as albuminuria and pseudo-membranous exudation on abraded surfaces, nor its characteristic paralytic sequelæ.

1. So far as the mere anatomical conditions are concerned, it is now generally conceded that there is no essential difference between primary membranous croup, and membranous croup occurring in the course of diphtheria. We shall enter more minutely into details when treating of the morbid anatomy of diphtheria, but it is important to allude here to the various points of difference which have been supposed to exist between croup and diphtheria in this respect. Isambert stated that ulceration of the mucous membrane of the larynx existed in diphtheritic croup alone, but West has met with similar ulceration in cases of primary croup, though somewhat less frequently than in the secondary diphtheritic form.

It has also been attempted to establish a distinction between the morbid process in croup and in diphtheria upon the greater intensity of the lesion in the latter case, associated with more swelling and a more intense congestion of the mucous membranes; but from careful observation of the numerous grades of severity of the diphtheritic process, we are convinced that this difference in degree is not constant, and cannot be made the basis of a radical division of the two diseases.

So too the supposed anatomical differences in the structure of the two kinds of pseudo-membranes were formerly regarded as significant of an essential difference between the two diseases. One by one, however, these hypothetical distinctions, whether chemical or histological, have been abandoned; and the highest authorities of all countries are agreed that the differences between the two are merely in degree, and are to be regarded as due to the different anatomical structure of the pharyngeal and laryngeal mucous membrane.

The Report of the Committee of the Royal Med.-Chir. Society of London on the Relations of Membranous Croup and Diphtheria (*Trans.*, vol. lxii, 9, 1879, p. 80), states that "the testimony of English observers does not seem to supply any anatomical basis for the separation of diphtheritic from croupous products." Of recent German writers, Wagner (*General Pathology*, Amer. ed., 1876, p. 265-266), who considers the pseudo-membranes as the result of a peculiar transformation of the epithelial cells, states "in the greater number of fatal cases of laryngeal croup, diphtheritic exudation is found on the soft palate;" and again "between the croupous and croupous-diphtheritic exudations there is every possible transition, whilst sometimes the epithelial change, sometimes the infiltration of mucous membranes, preponderates." These expressions, croupous and croupous-diphtheritic, it is to be remembered, are used by Wagner in a purely anatomical sense. Rindfleisch also (*Path. Histology*, *Syd. Society Trans.*, vol. i, p. 422, etc.) holds that the morbid process which leads to pseudo-membranous formation is identical in the pharynx, larynx, and trachea, and that the properties of the false membrane, particularly the histological quality of the securing fibrin, and the firmness with which the membrane adheres to the mucous surface, vary with its place of origin, and find their explanation in the normal structure of the affected part.

Cornil and Ranvier, the most recent and authoritative among French

writers on morbid anatomy, assume as established the anatomical identity of croup and diphtheria; and Shakespeare and Simes, well-known American pathologists, who have translated the work of Cornil and Ranvier (Philadelphia, 1880), hold the same views. We may add that in the repeated examinations which we have ourselves made of the false membranes of primary croup and of pharyngeal diphtheria, we have discovered no differences that were not to be accounted for by the peculiarities of the normal structure of the parts.

If, therefore, it must now be admitted, that the anatomical argument in favor of an essential distinction between croup and diphtheria has been refuted, it remains for us to consider the several points of clinical difference.

2. It is undoubtedly true that the primary sporadic form of membranous croup occurs more exclusively in children than does the more fully developed form of diphtheria; but it must be remembered that in childhood there is a peculiar tendency to acute affections of the larynx, and that this part is consequently especially liable to become involved in the cause of diphtheria, and also that a comparatively trifling amount of membranous exudation in a child's larynx will produce grave symptoms of obstruction. Moreover, it has for many years been our decided opinion that in the vast majority of cases of so-called membranous croup, the disease had really begun with some membranous exudation in the fauces, which has too often been overlooked. Unquestionably, the exudation occurs primarily in the larynx in some cases; but as far as can be determined from existing statistics, this does not occur in more than from 10 to 15 per cent. of the cases.

Two cases observed by us in private practice will show how easily membranous croup might be assumed to be idiopathic, when, in truth, it is dependent upon diphtheria: We were called to see a child, four years of age, on Saturday morning, for a croup which had developed the night before. From the severity and steady advance of the laryngeal symptoms, and especially from the tone of the cough, which bore the curious resemblance to the sneeze of a kitten we have referred to, we suspected membranous laryngitis. When first questioned as to the antecedents, the mother insisted that the child had been quite well up to the moment of the invasion of croup the night before. But, after some consideration, she stated that the child had not been quite as well on the previous Monday and Tuesday, and that on Tuesday she had seen a few whitish specks on the child's throat, but had thought nothing of them. On careful examination by ourselves there was not a sign of exudation on the throat. It had entirely disappeared. This child had a violent attack of true membranous croup, and narrowly escaped death.

On another occasion, a boy, two and a half years old, had a well-marked and sharp attack of diphtheria of the tonsils and fauces. Rest in bed and proper treatment for four days dissipated the disease, and the child was allowed to get up. At this time two other boys in the same family, of four and six years of age, were seized with severe diphtheria, marked by high fever, loss of strength, a severe inflammation, with copious exudation on both tonsils and pharynx. They were quite ill, but recovered without any extension to the larynx. At one of our morning visits to these two cases the first child, the one who was supposed to have recovered, ran into the room. We were surprised to hear him give a loud and distinct croupy cough, and to observe that his voice was very husky. There was not a sign of disease in the fauces. We had him put to bed at once, and resorted to the chlorate of potash and tincture of iron, which had seemed to cure the first attack. The croup advanced rapidly in spite of

all that we could do. Early in the morning of the third day afterwards, he was so ill that we told his father he would die, but that there remained the chance of recovery from tracheotomy, though we could scarcely recommend it. He wished everything done. A surgeon was called, who advocated the operation. It was performed with much difficulty. There was some unavoidable trouble in the introduction of the canula, and the child died on the table.

In these two cases there were none of the malignant symptoms of diphtheria present, no fetor, nor any considerable external glandular swelling, and when the larynx was invaded the faucial disorder had entirely disappeared. We believe that many cases of diphtheria are very mild, so much so that only careful inspection of the fauces reveals the true key to the slight constitutional disturbance, and yet, even in these mild cases, the membrane sometimes invades the larynx and brings on the most alarming croup.

3. Much stress has been laid on the fact, that in croup the cervical lymphatic glands are not usually affected, but the cause of this is apparent when it is remembered that the lymphatics of the larynx and trachea communicate only with the single lymphatic gland below the greater horn of the hyoid bone, and with the small gland at the side of the trachea. When on the other hand the pharynx is involved, the close connection between its rich lymphatic supply and the numerous glands below the angle of the jaw, makes enlargement of these glands a prominent symptom from an early date.

4. The assertion that croup is a local non-contagious inflammatory disease of sthenic type, while diphtheria is a specific zymotic disease of an adynamic type, can only be briefly noticed here. Undoubtedly there is a wide difference between the constitutional symptoms of the grave septic form of diphtheria and those of croup. Undoubtedly also in cases of severe diphtheria, especially of epidemic form, where its infectious and contagious characters are pronounced, and the constitutional symptoms are of a low septic type, the pseudo-membrane is likely to be extensive and persistent in the pharynx. When the primary blood-poisoning is intense, death often occurs before the membrane could extend to the larynx. But in many cases, it is our belief that the gravity of the general symptoms of pharyngeal diphtheria, is due to a *secondary* infection of the system from the local disease through the medium of the lymphatics, whose abundance we have above noted. On the other hand, we see cases of pharyngeal diphtheria, both of epidemic and sporadic form, where the constitutional infection, either primary or secondary, is but slight, and where the general symptoms are no more adynamic in character than those we have seen in membranous croup. Especially is this the case in those instances where the pharyngeal exudation has been very slight and transient, and the larynx has been speedily invaded with the development of croup. But although the danger of secondary infection from the exudation in the larynx is less than when the pharynx is involved, the symptoms of prostration in croup are often marked. Of course, when the primary blood-poisoning is intense, and yet the exudation has formed first in the larynx, as it has done with unusual frequency in certain epidemics of diphtheria, the constitutional depression has been marked from the first.

We have recently met with three separate instances where, among

children of a family, one was seized with membranous croup, presenting the symptoms and course described in the primary idiopathic form; while one or more of the other children were affected with pharyngeal diphtheria, running into the laryngeal form in one case and causing death. Here the cause, the type of the disease, and the character of the general symptoms were similar.

5. Contrary to what has often been asserted, albuminuria is frequently present in croup. It is true that it is much less frequent than in pharyngeal diphtheria (47.4 per cent. as against 85 per cent. of the cases analyzed by the Committee of the Med.-Chir. Society); but this is only what would naturally be expected from the greater tendency to secondary systemic infection when the pharynx is involved. In regard to the paralytic sequelæ, it is clear that no inferences can be drawn, owing to the small number who recover from croup, and the small percentage of all cases of diphtheria in which paralysis occurs. Mackenzie, indeed, states (*Diphtheria*, 1879, p. 83) that paralysis has been occasionally met with in those that have survived an attack of croup.

We have thus briefly discussed the various points that have been urged as showing an essential difference between croup and diphtheria. The question is a vitally important one; and, in view of the eminent authorities who do not yet admit their identity, it must be regarded as still unsettled. It has therefore seemed best to treat of membranous croup as a special disease, apart from the brief notice of it we have given in the article on diphtheria.

Our personal experience constrains us, however, to state that the differences between the two forms of membranous croup above enumerated, have not seemed to us sufficient to establish their essential diversity; and that it is our decided opinion that the vast majority, at least, of the cases of so-called pseudo-membranous laryngitis or membranous croup, are in reality instances of laryngeal diphtheria, where the faucial deposit has been slight or possibly absent.

We are led to this conviction, especially by the repeated observation of cases in private practice, such as those recorded on page 87, where we have been summoned upon the first symptoms of indisposition, and have found a trifling amount of membranous exudation on the fauces, which, in a day or two, had disappeared, while the symptoms of croup supervened. We wish, therefore, to impress deeply on the mind of the reader the absolute necessity of immediately and repeatedly examining the throat, whenever the child is taken sick, with even the most trifling croupy symptoms; since, if any membranous exudation be detected on the tonsils or pharynx, the case must be regarded as probably one of membranous croup, a most guarded prognosis accordingly be given, and the most careful treatment be immediately instituted.

FREQUENCY.—The mortality from this disease is in all years considerable, as will be seen from the subjoined table:

Years.	Mortality from Croup.	Mortality from Diphtheria.	Total Mortality less Stillborn.
1846,	111	0	5,944
1847,	121	0	6,881
1848,	177	0	7,268
1849,	130	0	8,989
1850,	151	0	8,034
1851,	180	0	8,374
1852,	208	0	9,745
1853,	303	0	9,184
1854,	304	0	11,280
1855,	265	0	9,906
1856,	268	0	11,720
1857,	256	0	10,331
1858,	292	0	10,162
1859,	312	0	9,084
1860,	354	307	10,849
1861,	304	502	13,838
1862,	258	325	14,386
1863,	443	434	15,045
1864,	455	357	16,794
1865,	350	260	16,453
1866,	239	192	16,005
1867,	185	118	13,153
1868,	206	118	13,949
1869,	237	182	13,428
1870,	316	172	15,317
1871,	264	145	15,485
1872,	296	150	18,987
1873,	200	110	15,224
1874,	199	179	16,238
1875,	428	656	17,805
1876,	386	708	18,892
1877,	338	458	16,004
1878,	388	464	15,743
1879,	291	321	15,473

It is difficult to estimate the number of deaths due to primary membranous croup since diphtheria has made its appearance in the mortality lists of the city, as many cases of secondary diphtheritic croup have unquestionably been returned as mere pseudo-membranous laryngitis.

True croup is, however, rare in comparison with false croup, since while we have seen but 40 cases of pseudo-membranous laryngitis, we have met with upwards of 300 of the catarrhal form. In the following remarks, and in those on the causes of croup, we refer the reader also to the table in the article on diphtheria, showing the comparative monthly and annual mortalities from these two diseases.

From a glance at the accompanying table, it will be seen that since the prevalence of diphtheria, the mortality from croup has not increased disproportionately to the increase in general mortality. Moreover, no change whatever has occurred in the type of this disease during the past ten years, for the experience of one of us for a number of years before the term diphtheria came into use and appeared in the mortality returns of this city, enables us to attest the fact that pseudo-membranous laryngitis, both

of the primary and of the more grave diphtheric form, occurred then precisely as it does now.

PREDISPOSING CAUSES—AGE.—The disease is far most frequent between the close of the first and fifth years. Thus of 2136 fatal cases reported in this city during the seven years from 1862–68, 301 were under 1 year of age; 571 between 1 and 2 years; 951 between 2 and 5 years; or 1522 between 1 and 5 years; and 236 between 5 and 10 years; leaving but 77 cases as occurring after the latter period of life.

Of 38 cases that we have seen, 30 occurred between 2 and 7 years of age; while of the remaining 7, 1 occurred at the age of 18 months, 1 at that of 19 months, 1 at $7\frac{1}{2}$ years, 2 at 11 years, and 1 each at $11\frac{1}{2}$ and $12\frac{1}{2}$ years.

SEX cannot be said to exercise any decided influence upon the frequency of the disease. Thus of the above 2136 cases, 1115 occurred in males, 1021 in females.

CONSTITUTION.—A feeble and delicate constitution is thought by some to be a powerful predisposing cause, but this is at least very doubtful.

Of the 40 cases referred to, of which we have preserved notes, 29 occurred in healthy vigorous children, while the remaining 11 occurred in children who, though neither very weak nor very sickly, presented a rather delicate appearance.

SEASON exerts a very powerful influence upon the development of croup. Thus the mortality from it reaches its maximum during the months of November, December, and January, during which quarter about four times as many deaths occur from croup as during the months of June, July, and August. It is, however, comparatively frequent from October to March, inclusive.

The relation between the mortality from croup and the temperature appears to be a definite and quite constant one, since, as will be seen by referring to the table in the article on diphtheria, with the single exception of February, the mean monthly temperature and the mean monthly mortality from croup vary in inverse ratio throughout the entire year.

The fact that croup occupies a relation to temperature so much more definite than that held by diphtheria, may be due solely to the special tendency to laryngeal irritation that exists during inclement weather. The following interesting table is condensed from the report of the Medico-Chirurgical Committee (*op. cit.*, p. 10).

	Laryngeal, with Faucial Exudation.	Laryngeal only.
January,	6	5
February,	4	5
March,	10	7
April,	8	5
May,	5	3
June,	12	6
July,	9	3
August,	13	1
September,	10	4
October,	8	4
November,	12	5
December,	4	1
Total,	101	49

The *exciting causes* are but little understood. It is known that membranous exudation may follow the application of irritating agents to the laryngeal mucous membrane, but this would account for only very rare instances of croup. Exposure to cold and sudden changes of temperature have been frequently assigned as causes, but careful examination tends to disprove their influence. In none of the cases that we have seen could the exciting cause be even suspected. It seems to us, therefore, altogether probable that it originates from the influences that cause diphtheria, and that the action of such agents as cold and wet is limited to determining the localization of the exudation in the larynx. It is but improbable also that the existence of some predisposing individual peculiarity may be assumed.

SECOND ATTACKS of membranous croup, though rare, are mentioned as occurring by several authors; and, in our remarks on tracheotomy, we quote from Millard an allusion to five cases, in each of which the operation was twice successfully performed for successive attacks of this disease.

We have ourselves met with two instances in which second attacks occurred. One was a girl, who had her first attack at the age of $11\frac{1}{2}$ years, and her second at the age of $12\frac{1}{2}$, and recovered from both without the operation. The second patient was a boy, who had his first attack, a very severe one, but from which he recovered without tracheotomy, at the age of $5\frac{1}{2}$ years; and his second attack, which is fully detailed at the end of the article on tracheotomy (Case 1), at the age of $7\frac{1}{2}$ years.

ANATOMICAL LESIONS.—The false membrane may cover the whole mucous membrane of the larynx, and extend into the pharynx, trachea, and bronchi; or it may be confined to the larynx, either forming a complete lining to the cavity of that organ, or consisting merely of patches of various sizes, with intervals of mucous membrane destitute of exudation.

It is, in the first place, important to ascertain the proportion of cases in which the deposit extends into the bronchi, and those in which it remains limited to the larynx, or larynx and trachea, as the determination of this point has some bearing upon the question of the propriety of the operation of tracheotomy. It appears from a table given by M. Guersent (*Dict. de Médecine*, t. ix, p. 346), containing the results of cases collected by M. Hussenot from various sources, and of autopsies made by M. Bretonneau, numbering in all 171, that in 78 the membrane did not extend beyond the trachea, and that in 42 it invaded the bronchi, and in 30 the condition of the bronchi was not mentioned; and in 21 there were no false membranes; so that of 120 cases, in which the extent of the false membrane was accurately noted, it was confined to the larynx and trachea in 78, and extended into the bronchi only in 42; or in about one-third of the cases. This proportion is the same that Millard gives (*De la Trachéotomie dans le cas de Croup, Thèse de Paris*, 1858), in his masterly memoir upon croup, after an analysis of a large series of cases. Our own experience, based upon 15 cases in which we ascertained with exactitude (by autopsy or by tracheotomy) the extent of the membrane, would indi-

cate that it passed into the bronchi in a large proportion of cases; since in 7 of these 15 cases the exudation extended beyond the trachea. It is to be borne in mind, however, that the cases upon which these calculations are based have very frequently resulted fatally, and presented extensive formation of pseudo-membrane in the bronchi; and it is probable that it really exists there in other instances, but to a much less extent, so that recovery takes place, and renders it impossible to determine accurately the extent of exudation.

The proportion of cases in which the pharynx is implicated is also important, since it affects the diagnosis of the disease, and indeed bears upon the question of the identity or non-identity of pseudo-membranous laryngitis and diphtheritic croup.

We have already referred to this important point, and would here merely repeat our belief, that in the vast majority of cases of membranous croup the disease has begun with exudation on the pharynx, though perhaps only to a very trifling extent. It is evident that most of the statistics published as bearing on this point, are not really applicable. Some of them, as those of Bretonneau and Guersent, only show the proportion of primary membranous laryngitis to diphtheria in general; which from these and other sources may be computed as not more than three per cent. Other sets of statistics, and this remark applies to most that have been published, only show the proportion of cases of croup where the fauces were free from exudation at a variable period after the inception of the disease; and when the cases have been brought to hospitals, it has usually only been after the laryngeal symptoms have become pronounced; by which time, as our experience in private practice has shown conclusively, the faucial deposit, which was frequent at first, may have entirely disappeared. Mackenzie (*op. cit.*, p. 82) states, without giving any figures in support, that croup originates in the larynx or trachea only in 10 or 12 per cent. of the cases. Our own experience, in cases seen at the very outset, would give 16 per cent. as the proportion: thus, in 33 cases observed by ourselves, in which the condition of the throat was recorded, the croup followed membranous angina in 23 cases; in 5 the disease began in the larynx, but was attended later with small deposits upon the tonsils; and in 5 only was there no deposit on the throat at any time.

The fauces and pharynx do not present any constant alterations in cases of croup. Frequently, however, the mucous membrane is red and swollen, and there may be patches of membranous exudation upon the tonsils, velum, half-arches, or on the pharynx. These patches are usually thin, whitish, and may not persist more than twenty-four to forty-eight hours, disappearing and being succeeded by similar formations in some other part of the throat.

We believe, indeed, that such patches of exudation will be found in a large proportion of cases during the first two or three days of the attack; and that they are not more frequently observed, chiefly because the symptoms are usually so slight during this stage, that either no medical attendant is summoned, or his attention is not attracted to the throat.

The most important and characteristic morbid appearances are, however, to be found below the glottis, and consist in the presence of pseudo-mem-

branous exudation, and of certain alterations in the respiratory mucous membrane.

The false membrane may be limited to the larynx, or the larynx and trachea; or it may extend over these parts and into the branches of the bronchi, even to the third and fourth division. In the larynx, trachea, and even the primitive bronchi, it may appear merely as patches of various sizes, with intervening spaces of vascular mucous membrane; but in the smaller air-passages it usually takes the form of complete tubes lining the bronchus. In some cases, such tubular casts may be formed continuously from the larynx down to the minute bronchioles, completely lining the air-passages. It is undoubted, that in the more sthenic idiopathic form of membranous laryngitis, the membrane is more apt to extend deeply into the ramifications of the bronchi, than when it occurs as a complication of diphtheria.

The false membrane is commonly of a yellowish-white color, and from a fifth of a line to a line in thickness. Its consistence is generally considerable, and it is usually somewhat elastic; indeed the more white and fibrous varieties possess a degree of firmness and toughness that renders it difficult to tear the membrane, or tease it out with needles. It is an almost invariable rule, that the membrane lining the upper part of the air-passages is more white and firm than that found in the smaller bronchi; so that it frequently happens, that, on drawing out the firm white tubular membrane lining the larynx, trachea, and primary bronchi, it is seen to terminate in branches which grow progressively softer, more yellow and purulent as they become smaller and smaller.

The free surface of the pseudo-membrane is usually covered with puriform mucus, while the attached surface is adherent with various degrees of force to the mucous membrane beneath. The strength and closeness of these adhesions are often proportionate to the firmness and toughness of the false membrane itself. In the larynx and trachea it is often necessary to employ a good deal of force to separate the exudation from the mucous membrane, and innumerable little fibres are seen passing from one to the other, as though they were processes of exudation dipping into the minute orifices of the mucous follicles. On the other hand, the adhesion between the exudation and mucous membrane is rarely close in the smaller bronchi, or in cases where the pseudo-membrane in the larynx and trachea is less firm and consistent.

These false membranes consist, according to Hasse, mainly of fibrin blended with mucus in various proportions (*Path. Anat.*, Syden. Soc. ed., p. 278). On microscopic examination, they present a more or less close fibrous basis, consisting of interwoven fine fibrils, with imbedded cells in varying number; these cells presenting the ordinary appearances of exudation corpuscles, being round, granular, and containing from one to three small nuclei. The action of various chemical reagents upon them will be found detailed in the article on diphtheria.

The mucous membrane beneath the exudation presents various shades of redness, or it is purplish, or even ecchymosed and blackish. It is also swollen, and may be slightly softened or friable, and has a dull excori-

ated appearance, though actual ulceration very rarely exists. West mentions the occurrence of small aphthous ulcers about the edges of the rima glottidis and the arytenoid cartilages as a frequent lesion in idiopathic croup; but the same lesion has been observed in the diphtheritic form of the disease. There is also vascularity, though usually to a less marked degree, of the bronchial mucous membrane at the points where no exudation exists.

The lungs present some abnormal condition in the great majority of cases. Bronchitis and pneumonia are frequent complications of the disease; and in addition there is often collapse of larger or smaller portions of lung-tissue from occlusion of some bronchus by the pseudo-membrane. In other instances, or frequently in conjunction with collapse of portions of the lungs, the violent respiratory efforts induce either vesicular or even interstitial emphysema, especially of the anterior borders of the lungs.

The morbid appearances found in cases where the croup has followed diphtheritic angina, will be fully described under the head of this latter disease.

In the secondary croup of measles, the appearances are very similar to those observed in primary cases, while in that of scarlet fever the exudation differs in being less consistent and less uniformly spread over the diseased part. In the last-named malady, the membrane is thinner and less adherent, and, in some cases, puriform, soft, and of a grayish color. It is usually poor in fibrin, and prone to decomposition. The mucous membrane is generally discolored and softened.

SYMPTOMS.—In the majority of cases, the development of the symptoms characteristic of croup, is preceded for a few days by the ordinary symptoms of catarrh and slight sore throat. The child is feverish and drowsy; there is cough, which may possess a slight croupy character at some period of the twenty-four hours, but more frequently seems like an ordinary catarrhal cough; coryza is very rarely present, but there is slight soreness behind the angles of the jaws, and the fauces are seen to be reddened, and probably small, thin patches of pseudo-membrane may be visible on the tonsils or fauces. This early stage lasts a variable time, usually from one to three or four days, and is more or less gradually succeeded by the symptoms indicative of laryngeal obstruction.

When, on the other hand, the disease begins in the larynx, the invasion is marked by hoarseness of the voice, and hoarse, croupal cough, which often continue for one, two, or three days, until the disease has made considerable progress, before the parents deem it necessary to send for a physician. In a case that came under the observation of one of ourselves, the child was playing about the room at a time when he had hoarse, whispering voice, and cough, and stridulous respiration. In another we were not called until the evening of the third day, though the child had had stridulous cough and respiration for two nights; but, as he always seemed better in the morning, it was not thought necessary to send for a physician until after he had become violently ill. In a third case there was hoarseness of the voice and slight croupal cough during the afternoon of one day and the ensuing night, and the next morning fully developed croup,

with fibrinous patches on each tonsil. These symptoms are not generally accompanied by fever at first. The appetite is usually unimpaired, the thirst scarcely augmented, and the child, though somewhat dull and languid, is disposed to be amused at times. In other and severer cases, on the contrary, the disease becomes aggravated much more rapidly, and may soon lead to a fatal termination.

The change of the *voice* is the first symptom observed in the cases which begin in the larynx. It has always been described to us as hoarse, like that which is heard in an ordinary cold. As the disease progresses, the voice becomes more and more hoarse and difficult, until at length it is reduced to a mere whisper. The grade of the hoarseness varies, however, to a very great degree in the same case, the diversities depending probably upon the amount of the spasm of the larynx at the moment, and upon the state of the exudation. We have several times observed the voice to become much stronger and clearer after the operation of an emetic, in consequence, no doubt, of its relaxing effect upon the glottis. The *cough* is peculiar. At first slightly hoarse, it becomes, as the case goes on, very hoarse and hollow, and then short and smothered. It is variable in frequency, and is apt to occur in paroxysms, which are often very troublesome from their frequent recurrence. Towards the termination of the disease in fatal cases, or whenever the case is very severe, it is altogether different in character from what it was at the beginning, becoming short, instantaneous, and smothered, so that it might very well be called whispering. As the disease progresses, it is accompanied by stridulous respiration, in which a hoarse, rough, hissing, or crowing sound is produced by the rush of the air through the constricted larynx. This sound is usually heard at first only during forced inspirations, and is therefore noticed first during the long inspiration which precedes coughing. Next it is heard during the violent respiratory movements which accompany the act of crying; and as the larynx becomes more and more clogged with the exudation, it occurs during both inspiration and expiration, in every act of respiration, and is so loud as to be heard over the whole room, or even in adjoining rooms.

The *respiration* is natural in the early part of the attack, but as the voice and cough assume their characteristic features, and the stridulous sound is established, it grows more frequent, rising to 28, 32, 40, and 48 in the minute. At first easy and natural, it becomes, during the height of the symptoms, and especially in fatal cases, the most frightful dyspnœa we have seen in any disease. Every movement of inspiration requires the whole force of the inspiratory muscles to lift the walls of the chest, and enable the air to find its way through the narrow and obstructed glottis; each expiration, instead of being short and easy, as in health, and in nearly all other diseased conditions, requires a slow and laborious contraction of the expiratory muscles to expel from the lungs the air which they contain, and which hisses through the larynx with a sound nearly as loud as that produced during inspiration. The dyspnœa just described is for the most part constant, but exhibits paroxysmal aggravations from time to time.

When a paroxysm of dyspnœa occurs, the expression of the child is that of the most terrible anxiety, or of the wildest terror. In some instances, the face becomes deeply red, then blue, livid, and finally pale and white, and for a moment life may seem extinct. In other cases in which the dyspnœa is constant, the face is of a dusky red color, the expression anxious and haggard, and the child either lies on its side with the head thrown far backwards in a state of somnolence, or constantly changes its position from restlessness without noticing anything around it.

Jacobi (*Amer. Jour. of Obstet.*, May, 1868, pp. 13-65) lays particular stress upon the fact that in membranous croup the dyspnœa exists both in inspiration and expiration, whereas in spasmodic catarrhal croup it is chiefly present in inspiration, and is due, he thinks, to paralysis of the crico-arytenoid muscles from œdema and infiltration, so that the vocal cords are brought into contact during inspiration.

There is one further peculiarity about the dyspnœa of membranous croup to which we would direct especial notice, since we regard it as of the utmost importance. This consists in the occurrence, in certain cases, of a deep sulcus around the base of the chest, and of recession of the lower part of the sternum and the epigastrium during the act of inspiration.

These phenomena are, perhaps, partly due to the violent action of the diaphragm, but undoubtedly their chief cause is the atmospheric pressure, which acts here, as it has been clearly shown by Jenner to act also in rickets, to produce the deformities of the thorax characteristic of that disease. The normal relation which exists between the firmness and resistance of the thoracic walls, the power and rapidity of contraction of the diaphragm, the elasticity of the lungs, and the size of the orifice of the larynx, are here disturbed by the greater or less degree of occlusion of the larynx by membranous exudation. The calibre of the larynx being thus diminished, so that the air enters the lungs but slowly, and the diaphragm contracting violently, there will necessarily be recession of the softer parts of the chest-walls at each inspiration.

The persistence of these phenomena during inspiration for even a short time is, we believe, in the highest degree characteristic of the presence of false membranes in the larynx; and when, despite the use of emetics, this form of respiration continues, it constitutes one of the strongest indications for the performance of tracheotomy.

There is no *expectoration* early in the disease, or it consists of yellowish viscous mucus. At a later period there is usually expectoration of false membrane, sometimes in the form of a complete tube, or, much more frequently, of small, irregular fragments, mixed with mucus, or with the matters ejected from the stomach by vomiting. To detect the membrane, the substances expectorated or vomited ought to be placed in water, when the former detaches itself from the mucus and other matters, and is easily recognized. It is not voided in all cases in which it is known to be present in the larynx.

Thus of the thirty-five cases observed by ourselves, it was expelled by vomiting or coughing in twelve; in twenty-one none was rejected, though its presence in each case was proved by the character of the symptoms,

by its existence in the fauces, by autopsy, or by the operation of tracheotomy; in one there was expectoration of masses of viscid, yellowish fibrin, though none of membrane; and in one there was no positive evidence of its existence. M. Valleix (*Guide du Méd. Prat.*, t. i, p. 330) states that of fifty-one cases, in which the symptoms were very carefully observed, no traces of the exudation could be discovered either in the expectoration or in the matters rejected by vomiting in twenty-six, though its existence was proved by post-mortem examination.

In the severe cases of true croup that have come under our notice, auscultation has been of little or no aid. In fact the chest-sounds has been, in most cases, so completely masked by the loud shrillness of the laryngeal stridor, that we have been unable to judge with any satisfaction to ourselves of the condition of the lungs. It has been impossible to determine whether the inability to detect the natural respiratory murmur depended on the small volume of air that found its way through the obstructed larynx, or on the fact that all sound was masked by the stridor. This is particularly unfortunate, since, were it not for this circumstance, we might be able to judge by auscultation of the extent to which the bronchi have been invaded by the false membrane,—a matter very important to determine when the question of tracheotomy comes to be mooted in any case.

In cases in which the laryngeal obstruction is not very great, and the stridulous sound consequently less loud, we may auscult the chest to some profit. The vesicular murmur is then either natural, or altered according to the state of the lung. This question will be found referred to more fully in our remarks on the indications for the operation of tracheotomy.

There is a slight *febrile movement* at the onset, or a day or two after the appearance of the earliest symptoms. When the disease is fully established, the fever is sometimes violent. The pulse rises to 130, 140, 160, or even higher; it is generally regular and strong at first, but as the case progresses, becomes small, feeble, and very rapid. In one of the paroxysms that we witnessed, it became so rapid that it could not be counted, and at last ceased to beat at either wrist for a few instants. The heat and dryness of the skin are very moderate at first, but increase as the disease reaches its maximum, to diminish afterwards gradually, and in fatal cases, to be replaced by coldness, with copious clammy perspirations. The strength is not diminished at first, but as the disease progresses, becomes more or less so in proportion to the violence and duration of the case. The *digestive organs* are but little disturbed by the influence of the disease, with the exception of diminution or loss of appetite, and moderate thirst, during the violent period. Spontaneous vomiting or diarrhoea are rare, though both sometimes occur. The tongue is moist, and generally covered with a yellowish-white fur. Pain in front of the larynx has been noticed by several authors. We have ourselves observed it in but one case.

Tumefaction of the submaxillary glands, which is a frequent symptom of pseudo-membranous angina, ought always to be sought for, and when present adds additional support to the diagnosis.

The *mode of recovery* in favorable cases is different in different instances. In some it is sudden, taking place rapidly and steadily after the expectoration of a tubular-shaped membrane. The rejection of the deposit in this form is, however, a rare event, and is not always followed by recovery. We have seen in this city three distinct tubules of false membrane, which were thrown from the larynx of the same child at intervals of two days each. The first was the largest, and came evidently from the whole length of the larynx and trachea; the second was somewhat shorter, and the third not more than half so long as the first. The child was greatly relieved for some hours on each occasion of the rejection of a tubule, but then became more oppressed as the exudation again collected. It sank from exhaustion after the third came away.

As a general rule, the recovery is slow and gradual. After free vomiting, after the expectoration of fragments of false membrane mixed with mucus, or, as happened to ourselves in two cases, after the expectoration of masses of tough yellowish fibrin, or lastly, after the rejection of mucoid and frothy sputa only, the symptoms gradually ameliorate; the stridulous respiration slowly subsides, and at last disappears; the cough, which was short, hoarse, and smothered, becomes louder, stronger, less hoarse, and what is still more favorable, loose; the aphonia moderates, but very slowly; the fever disappears; appetite and gayety return; and, after a variable length of time, the child enters into full convalescence. The hoarseness of voice very generally continues for several days after all the other symptoms have lost their dangerous character, and sometimes lasts for weeks. In one case the voice was still weak and hoarse on the tenth day, and in another during the seventh week. (*See a paper on Croup, by J. F. Meigs, M.D., Am. Med. Jour. Med. Sci., April, 1847.*)

DURATION.—Death has been known to occur on the first, second, and third days, but such cases are rare. The duration of the disease may be stated at from three to thirteen days, as its most common term. The cases seen by ourselves lasted from five to fourteen days.

DIAGNOSIS.—There can be no difficulty in recognizing the presence of pseudo-membranous laryngitis, when the development of the symptoms of laryngeal obstruction has been preceded for several days by diphtheritic sore throat.

When, however, the disease seems to begin in the larynx, and especially when there is no exudation whatever in the fauces, the diagnosis becomes more embarrassing, since under these circumstances there are two other laryngeal affections with which true croup may be confounded, to wit: false croup or spasmodic catarrhal laryngitis, and laryngismus stridulus. The mode of distinguishing between these different disorders has been carefully described in the remarks on diagnosis under the head of the former disease. We wish in this place merely to call the attention of the reader, and particularly of the young practitioner, to the extreme importance of the differential diagnosis between the disease now under consideration and false or spasmodic croup, since the former is one of the most dangerous and frightful disorders to which children are subject, demanding vigorous treatment from the start, at which period only is medical treatment likely

to be successful; whilst the latter, though of a much more threatening aspect at the beginning, is in fact a mild and safe disease in comparison, and one rarely requiring other than very simple treatment.

In this connection we would urge again the extreme importance of a careful examination of the throat in every case where there are even the most trifling croupy symptoms present, since if membranous exudation be present either on the pharynx or tonsils, there is great danger that the laryngeal symptoms are due to an extension of the false membrane.

PROGNOSIS.—Pseudo-membranous laryngitis is a very fatal disease. In its sporadic form it is decidedly less dangerous than when it occurs in the course of epidemic diphtheria, owing to an extension of the exudation from the fauces into the larynx; but it still ought, at all times and in all shapes, to arouse the utmost caution of the practitioner.

MM. Rilliet and Barthez state that its common termination is in death. M. Valleix says that, "to speak in general terms, it is fatal when not treated energetically." M. Guersent (*loc. cit.*, p. 365), after a careful consideration of the statements of various authors, says: "In fact, true croup is one of the most dangerous of all diseases; it is generally fatal." He adds that he has seen at least 100 cases of spasmodic croup without a single death, while of 10 children attacked with true croup, it is scarcely possible to save two.

We have ourselves seen upwards of 200 cases of spasmodic or false croup, all of which without exception recovered, while of the 35 cases of true croup, of which we have preserved careful notes, 16 died.

The danger is great in proportion as the child is younger and more feeble, and in proportion to the rapidity of the case and the degree of the dyspnœa. The most unfavorable symptoms are: loud stridulous sound, heard both in the inspiration and expiration; laborious and prolonged expiration; recession of the base of the thorax during inspiration; whispering voice or complete aphonia; congestion of the face and neck; somnolence; weak, rapid, and irregular pulse; cold extremities; and cold, clammy perspirations. The favorable symptoms are: expectoration of false membranes; diminution of the stridulous respiration; the change from whispering to hoarseness or to clearness of the voice; looseness of the cough; moderation of the fever; improvement of the temper and moral state; and amelioration of the general condition.

The case should not, however, be abandoned as hopeless until life is actually extinct. An instance has been elsewhere put on record by one of us (*see paper by Dr. J. F. Meigs, loc. cit.*) of the recovery of a child after momentary suspension of animation from asphyxia on two occasions, though these attacks were followed by a dreadful illness of two days.

TREATMENT.—We are desirous, at the beginning of our remarks upon the treatment of this disease, to express the opinion that none is *likely* to succeed unless it be applied early in the case, and by this we mean in the course of the first or, at the latest, second day. And not only should it be commenced early, but the most active remedies ought to be applied at this period, in their full force. The very moment there is good reason to suppose that a case will prove to be one of membranous croup, the most

energetic means ought to be brought to bear upon it, and if this be done from the first, or even second day, we cannot but hope that a considerably larger proportion of recoveries may take place than has heretofore been thought possible.

In the study of the treatment, it will be necessary to rely chiefly upon the works that have been published since the distinction between the two forms of croup has been correctly drawn, for it is impossible to place much dependence on the assertions of previous writers, inasmuch as their opinions in regard to the effects of treatment must have been formed from indiscriminate experience in two very opposite diseases. It is only necessary to recollect the enormous difference in the mortality of the two affections, as shown by our own experience and the statistics quoted from Guersent, to be convinced that the success of any plan of treatment in the one is no fair argument for its probable success in the other. The most important objects to be held in view in the treatment, are the following: to prevent, if this be at all possible, the formation of false membrane; after its production, to cause its dissolution, or render it less adherent; to provoke its expectoration; to prevent its reproduction after it is once expelled; to subdue the inflammatory condition which exists; to allay the painful symptoms; and in every way to support the system.

Bloodletting.—Some authors still award to bloodletting a high place in importance amongst the medical means in our possession, and it was formerly regarded by many in this country as an indispensable agent in the cure. Moreover, there are not a few who believe that, when promptly and boldly resorted to, it will seldom fail in arresting the disease.

The more careful and extended study which this question has received during the past few years, however, has led many observers to doubt the efficacy of venesection in arresting the course of this inflammation, or preventing the formation of membranous exudation.

In those cases where croup supervenes in the course of epidemic diphtheria, there can be no doubt that bloodletting is entirely contra-indicated; and the same remark may be made of those sporadic cases of pseudo-membranous laryngitis, where the onset of the disease is slow, and its course gradual, and unattended by high febrile reaction. Indeed, the more wide experience we have ourselves had in the treatment of this disease during late years, has convinced us that bloodletting is, to say the least, unnecessary, excepting perhaps in cases where the disease occurs suddenly in vigorous children, and is attended at an early period of the attack by violent febrile action and especially marked suffocative symptoms. Under such circumstances, and such only, it may be advisable to resort to a moderate general venesection, principally for the mechanical relief thus afforded to the acute and intense venous stasis caused by the obstructed respiration.

For all the other indications, however, for which bleeding was formerly recommended in croup, namely, for the reduction of the fever and inflammation, and for the arrest of the exudative process, we prefer resorting to the other remedies hereafter mentioned.

Emetics.—Emetics are recommended by all writers, and are generally acknowledged to be amongst the most, if not the most, efficient of all the means employed. M. Valleix (*op. cit.*, t. i, p. 358) has demonstrated their importance more fully than any other writer. He states that of fifty-three cases of the disease, tartar emetic and ipecacuanha were chiefly relied on in thirty-one, of which fifteen were cured; whilst of the twenty-two others, in which they were parsimoniously given, but a single one recovered. He gives other facts in regard to these cases which are highly interesting and important. Thus, of the thirty-one cases treated with powerful emetics, false membrane was rejected during the efforts of vomiting in twenty-six; and of these, fifteen, or nearly three-fifths, recovered. In the five others of the thirty-one, on the contrary, no membrane was expelled, and they all terminated fatally. Again, of the twenty-two cases in which emetics formed but a secondary part of the treatment, two rejected false membrane, and of these one recovered; whilst of the twenty others in which no false membrane was expelled, not one escaped.

Our own experience in regard to emetics has been as follows: They were administered frequently and in full doses in thirteen of the twenty-one cases which began with angina, of which we have preserved notes; in six they were employed to a moderate extent, and in two not at all. Of the thirteen cases in which they were freely administered, eleven recovered; but, as in one of these life was saved only by tracheotomy, the success cannot be attributed to the emetics. Of the eight cases in which the emetic plan was not pushed, all but one ended fatally. False membrane was rejected in eight out of the thirteen cases above referred to. In one of the eight cases the quantity rejected was very small, and this was the case in which the child was ultimately saved only by operation.

Of thirteen cases in which the disease began in the larynx, emetics were energetically used, and frequently employed, in eight. Of the eight, five recovered. In four of the eight cases, fragments of false membrane were rejected, and in a fifth, a mass of viscid, yellowish fibrin (this case was marked as one of unquestionable membranous croup by patches of false membrane on the tonsils). Of these five, four recovered. In three of the eight, no false membrane was rejected, and of these two died. In five of the thirteen cases they were not freely used, being employed in two only as a secondary means; in one other only at the very termination of the attack, as we were not called to the case until the tenth day, the patient having been under homœopathic treatment before; and in the remaining two cases they were not employed at all. Tracheotomy was performed in four of these five cases, but in only one was a successful result obtained.

It is indeed true that there were peculiarities about the age and the type of the disease in the above groups of cases which may modify to some extent the conclusions which seem inevitable; but the statements and facts above given are quite sufficient to show that emetics exert a most powerful and beneficial influence on the disease, and that they ought, therefore, to form a principal and essential part of the treatment.

The emetics generally employed in Europe and in this country are tartar

emetic and ipecacuanha, which are given in the usual doses to produce full vomiting. We would, however, strongly discountenance the employment of tartar emetic as an emetic, under any circumstances, in children; and, at least in the disease under consideration, we do not like ipecacuanha as an emetic so well as one which, so far as we know, was first recommended by the late Dr. Charles D. Meigs. We refer to the Alumen of the Pharmacopœia.

In an article published by him in the *Medical Examiner* (vol. i; p. 414, 1838), he says he has been "accustomed to make use of an emetic, which, so far as I can learn, is very little employed, but which, from the certainty and the speediness of its operation, ought to be more generally admitted into the list of available medicines for this particular case at least. I have been familiar with its effects for more than twenty years, and my confidence in it increases rather than diminishes by time." He adds, "I think that I have never given more than two doses without causing very full vomiting; but I have often given large quantities of antimonial wine and ipecacuanha, without succeeding in exciting the efforts of the stomach."

The alum is given in powder, in the dose of a teaspoonful, mixed in honey or syrup, or in syrup of ipecacuanha, to be repeated every ten or fifteen minutes until it operates. It is not generally necessary to give a second dose, as one operates in the majority of cases very soon after being taken. We have known it to fail to produce vomiting only in two instances, both of which were fatal cases. In one the disease had gone so far before we were called, that no remedy had any effect upon the stomach. In the other, it was administered several times with full success, but lost its effect at last, as had happened also in regard to antimony and ipecacuanha. The reasons for which we prefer alum to antimony, or ipecacuanha alone, are the following: Antimony, when resorted to as frequently in the disease as we are of opinion that emetics ought to be, is too violent in its action; it prostrates many children to a dangerous degree, and is, we fear, in some cases, itself one cause of death. It acts injuriously upon the gastro-intestinal mucous membrane when used in large quantities and for any considerable length of time. Again, it is very apt to lose its effect, and to fail to produce vomiting. Ipecacuanha is a much safer remedy than tartar emetic, but its operation is often too mild, and it not unfrequently fails to produce any effect after it has been used several times. The advantages of the alum are that it is certain and rapid in its action, and that it operates without producing exhaustion or prostration beyond that which always follows the mere act of vomiting. It does not tend, like antimony, and in a less degree ipecacuanha, to produce adynamia of the nervous system; an effect which, in some constitutions or states of the constitution, or when it has been exhibited frequently, is often attended with injurious or even dangerous consequences. We have given alum in the dose above mentioned every four or five hours, for two or three days, without observing any bad effects to result from it. The alum was given in all the cases that we have seen, in which emetics were used, and was usually the only one employed when it was found to produce full vomit-

ing. In one of the cases accompanied by violent angina, ipecacuanha was substituted because of its smaller bulk. We have already said that it failed to produce vomiting only in two instances. It was the emetic employed in the nine cases in which fragments of false membrane were rejected, and in that in which the yellow viscid fibrin was expelled. Although it did not occasion the rejection of membrane in the other cases, it operated most speedily and efficiently.

Sulphate of copper has been highly recommended by several writers for its emetic operation, and, by some of the German physicians, as exerting a specific influence upon the disease in addition to its emetic effect. As an emetic, it may be given to a child two or three years old, in the dose of from half a grain to a grain every fifteen minutes, until it operates. To obtain its specific action it is continued afterwards in doses of a quarter of a grain every two hours.

We have also employed, with very good results, sulphate of zinc dissolved in syrup of ipecacuanha, in the proportion of 2 or 4 grains to the fluid ounce. Of this, a teaspoonful may be given to a child two or three years old, and repeated every fifteen minutes until it operates. This combination appears, like that of alum and ipecacuanha, to possess the double advantage of mild action without the production of any subsequent depression.

In the third edition of this work we referred to the use of the yellow sulphate of mercury (*Hydrargyri Sulphas Flava*) as an emetic in croup, as recommended by Dr. Hubbard, of Maine. Our own experience with this remedy has been limited, and not very decided. In the *American Journal of Obstetrics*, for May, 1870, Dr. Fordyce Barker, of New York, speaks in the highest terms of praise of its emetic effects in this disease. He always commences the treatment by a dose of from three to five grains, according to the age of the child, which may be repeated if it do not act, which he states very rarely occurs, in fifteen minutes. This he follows up with the use of *veratrum viride*, and states that the treatment has been successful in every case of true croup in which he has employed it. Undoubtedly this high testimony in its behalf justifies a further trial of turpeth mineral in croup, though we confess to a suspicion that not a few of the cases in whose *incipient* stage he has administered this drug so successfully would have proved to be instances of the severe catarrhal and not of the true membranous form.

We conclude these protracted remarks upon emetics with the statement that from what we have read, and from personal experience, we are induced to regard them as the most important remedies we have to oppose to this fearful malady. The emetic, whatever it may be, ought to be given three or four times in the twenty-four hours, and in severe cases, once in every four or five hours. The exact periods and frequency of the administration must be determined by the stage and urgency of the symptoms, and by the constitution and present strength of the patient.

Mercury.—This powerful drug was first employed freely in the treatment of membranous croup in America, and has subsequently been extensively used by English and European physicians. Calomel is the prep-

aration almost always preferred, and many authors still recommend the administration of this remedy, in larger or smaller doses, in the earliest stage of the attack.

During late years, our increased dislike of the administration of mercury to children in large and frequently repeated doses, and the constant observation that even its free use does not appear to arrest the course of true croup, or prevent the formation of membranous exudation, have led us to abandon entirely its employment in this disease.

At the same time we believe there has been found, in the free administration of the alkalies, an agency far less injurious than mercury, and equally powerful, if not more so, in promoting the separation and discharge of the exudation, and preventing its reproduction.

The internal remedies, then, upon which, after emetics, we rely most surely, are various alkaline salts, the use of which, in large doses, has been of late years highly recommended both at home and abroad. Those which we are most in the habit of employing are the chlorate and citrate of potash, which should be given in full and frequently repeated doses, as, for example, two or three grains every two hours to a child of four years old. We are also in the habit of combining with the chlorate of potash, tincture of the chloride of iron, in doses of three to five drops, at the same age.

Antispasmodics are undoubtedly useful in some cases, when there is much laryngeal spasm.

Opium is, however, the best remedy that can be employed for this condition, since it constitutes an important element in the treatment, by alleviating pain and restlessness, at the same time that it relieves the laryngismus, and thus diminishes the asphyctic symptoms. We would consequently recommend the use of some of the preparations of opium, as the tinct. opii deodorata, in such doses and at such intervals as will maintain a gentle opiate impression. In this, as in many other diseases of children, it is better not to prescribe the opium in combination with the other remedies that may be administered, but to either give it separately, or better still, to add it to the dose of the other medicines at the time of administration, so that the amount of the dose of opium and the frequency of its repetition may be modified constantly in accordance with the condition of the child.

Revulsives often prove useful in allaying restlessness, and moderating the violence of the suffocative attacks. Sinapisms and mustard poultices, applied upon various parts of the cutaneous surface, and mustard foot-baths, are amongst the best. The warm bath is often highly beneficial in the same way. We do not think it desirable ever to employ blisters in this disease.

LOCAL TREATMENT.—In those cases, and, as we have seen, they constitute the large majority of all cases of true croup, where the exudation appears in the fauces or on the tonsils before it involves the larynx, local applications to the throat are undoubtedly of importance.

The objects of such applications are here, as in diphtheritic angina, to promote the separation of the false membrane, and to prevent its repro-

duction. To fulfill the first of these indications, many authorities recommend astringent and caustic appliances, which cause the pseudo-membrane to contract and shrink, and thus tend to promote its separation; while others direct the use of those agents which exert a solvent action upon the exudation.

In the former class, the most advisable are, alum; tannic acid; solutions of nitrate of silver; the astringent salts of iron, especially the tincture of the chloride and the perchloride; dilute mineral acids and carbolic acid.

Of these applications, those which we prefer are a solution of nitrate of silver, in the proportion of 5 to 20 grains to $\text{f}\overline{\text{3}}\text{j}$ of distilled water; and tincture of the chloride of iron, in the proportion of $\text{f}\overline{\text{3}}\text{ss.}$ to $\text{f}\overline{\text{3}}\text{ij}$ to the $\text{f}\overline{\text{3}}\text{j}$ of water.

The second group comprises chiefly solutions of various salines, as the carbonate of potash, bicarbonate of soda, chlorate of potash, and lime-water.

If any of the astringent or caustic solutions are employed, we would recommend their application only to the patches of exudation in the fauces, since we regard it as highly doubtful whether they actually possess the power of preventing the formation of membranous exudation when applied to the surrounding mucous membrane. Still more should we doubt the efficacy or advantage of introducing such solutions, and especially the more powerful ones, into the larynx; either by pressing a soft sponge saturated with the solution upon the chink of the glottis, or by passing the sponge directly into the cavity of the larynx,* as recommended by Dr. Horace Green. (*Observ. on the Path. of Croup*, etc., New York, 1852). The practicability of this proceeding is undoubted, and a certain number of cases are on record in which it seems to have been used with success; but we have never resorted to the treatment ourselves.

In cases occurring in older children, who can be induced to inhale the vapor from an atomizer, or to allow a hand-ball atomizer to be used, the various astringent and solvent solutions above mentioned can be applied most satisfactorily in this manner; and, when this is practicable, we would prefer the use of lime-water or one of the alkaline solutions.

We attach so much importance to this remedy, and have found it to be followed by so much relief and comfort that we are in the habit of directing the inhalation of vapor of lime-water for five or ten minutes at least every two hours.

In order to obtain the advantage which undoubtedly follows the inhalation merely of the watery vapor, we are in the habit of causing the child to inhale the vapor from slaking lime for a few minutes in every hour, by covering the patient's body with a thick cloth, and holding a vessel containing the slaking lime a short distance below its mouth under the covering. It is doubtful, however, whether any appreciable amount of lime is carried up by the vapor so as to give the additional advantage of its solvent action upon the exudation.

The reader is referred for more detailed discussion of this question of

local applications in the treatment of croup, to the remarks upon treatment in the article on diphtheria.

HYGIENIC TREATMENT.—The child ought to be warmly clothed and confined to bed. The temperature of the room should be kept equable, and about 70° F.; the air should also be frequently changed, so as to preserve it constantly pure and fresh.

Owing to the loss of appetite and the pain caused by deglutition, it is often very difficult to induce the little patients to take food, so that this important element in the management of the case requires the utmost tact and attention. During the early part of the illness, the food should consist of light animal broths, beef tea, and preparations of milk. Later in the case, when the febrile action subsides, or if any symptoms of exhaustion and prostration appear, a small amount of wine and water, of wine whey, or of weak milk punch should be given.

Ice, given in small pieces to be held in the mouth, should be used very freely, as it relieves the parching thirst, and at the same time appears to act favorably upon the inflamed mucous membrane.

SUMMARY OF THE TREATMENT.—The general plan of treating this disease should, therefore, in our opinion, be somewhat as follows: The child should be confined strictly to bed. The food should be light, digestible, but nourishing, and, upon the earliest approach of exhaustion, a stimulus should be administered. In the early part of the attack we advise the use of revulsives, with mild counter-irritants; topical applications to the fauces if there is any membranous exudation visible, and the internal administration of citrate of potash, with ipecac. and small doses of opium, or of chlorate of potash with tr. ferri chloridi. So soon as the symptoms positively indicate the presence of false membrane in the larynx we resort to emetics, as directed in our remarks upon those remedies. During the whole treatment we also recommend frequent inhalations of the vapor of lime-water, or some other alkaline solution. And finally, after employing these means faithfully but without securing the discharge of the false membrane, while, on the other hand, the symptoms of laryngeal obstruction steadily progress, and the respiration grows more and more difficult, we must consider the propriety of resorting to the operation of tracheotomy, a proceeding which, as will be seen from the ensuing remarks, we approve of under the above circumstances.

TRACHEOTOMY.—The operation of tracheotomy would be apt to suggest itself to a medical man, on his witnessing the closing symptoms of croup, as the very means most likely to afford to the patient relief from the dreadful sufferings under which it labors, and as a possible rescue from impending death. It has accordingly been often resorted to in different parts of the world, at various stages of the disease, but with results that have led to very different conclusions.

In England, for example, the operation was almost universally condemned and abandoned about ten years ago; and in a former edition of this work we presented the unfavorable opinions of the most eminent English authorities.

It was a matter of very great surprise, at that time, that the results of the operation in the hands of English surgeons should differ so widely

from those obtained by the French physicians in similar cases; and, as there was no good ground for believing that sufficient difference existed between the croup of Paris and London, to explain the difference of success in the two cities, it is probable that the great disparity resulted, in part, from the operation being performed in France at an earlier stage of the disease, and in part also from the more careful after-treatment which the patients received.

Within the past few years, however, the operation has been more favorably regarded by English surgeons, and the statistics published show that the proportion of success now obtained does not fall far short of that claimed by French operators.

Thus in a paper read before the Royal Med.-Chir. Soc., in 1857, by Dr. Fuller, it is stated that up to that time 22 cases of tracheotomy in croup had been recorded in England, and that life had been saved in 8 of these, or in 1 out of every $2\frac{3}{4}$ cases.

In the statistical report of English hospitals from 1854-59 it appears that the operation had been performed in 15 cases with 4 recoveries, or 1 in every $3\frac{3}{4}$ cases. Still further, from the statistics published by individual operators in England, since 1858, though it is not to be presumed that we have met with all the cases recorded, it appears that tracheotomy has been resorted to in 63 cases, with successful results in 24, showing a success of 1 in $2\frac{5}{8}$.

When it is borne in mind also that in each of these instances the operation was postponed to the last suffocative stage, and that without exception the operators believe that the proportion of success would have been increased by its somewhat earlier performance, it becomes evident that tracheotomy has occupied a fair position in England among the legitimate operations of surgery.

It is thus advocated by Fergusson in the last edition of his *Practical Surgery*; and Dr. West, in 1859, speaks of it in these terms: "In spite of the unfavorable issue of the few cases in which I have either directed or sanctioned the performance of tracheotomy in croup, I am so far from being opposed to the operation, that my chief anxiety is to make out the indications which may justify me in having more timely recourse to it in future."

In Germany, also, the operation, if not generally practiced, is regarded as fully justifiable, and recommended and successfully performed by many of the most eminent authorities.

The statistics of the results there obtained, borrowed from Fock¹ and Voss,² show that of 50 cases operated on in the last stage, 24 terminated favorably, giving a success of 1 in $2\frac{1}{2}$, or 48 per cent. Steiner has also recently published (*Jahrb. f. Kinderheilk.*, No. 1, 1868) the results of the operation in 52 cases (33 boys and 19 girls), which show a recovery of 18, or 34.6 per cent. of those operated upon; and in an article upon diph-

¹ Report on Tracheotomy, Brit. and For. Med.-Chir. Rev., July, 1860, from Deutsche Klinik, 1860.

² New York Journal of Medicine, January, 1860.

theria and tracheotomy by Güterbock (*Arch. d. Heilkunde*, 1867, No. 6) 100 cases, operated on in Berlin, are reported, with 33 recoveries.

It is, however, in France that the operation first obtained, and has since firmly held, the position of a proper and legitimate method of treatment under certain circumstances of the disease. M. Bretonneau, of Tours, was the first who practiced it with sufficient success in France to give it some vogue. Since that time it has been recommended and performed by many different surgeons and physicians in that country, and particularly, as is well known, by M. Trousseau, who has been undoubtedly the most ardent and persevering, as well as the most experienced advocate of the operation. In one of his later publications upon this subject (*Arch. Gen. de Med.*, Mars, 1855, p. 259), he thus boldly advocates it: "For my part, I am quite determined not to allow myself to be discouraged, but to preach tracheotomy with the greater conviction in proportion as its success increases, and did this proportion remain what it was ten years since, I should still proclaim the necessity of the operation, nor cease to say that it becomes a *duty*, a duty as imperative as the ligature of the carotid artery after a wound of that vessel, though death follows the operation as often, certainly, as recovery."

M. Guersent (*Dict. de Med.*, t. ix, p. 376) recommends the operation when the usual therapeutical methods have failed, "as the only means that offers a remaining chance." He adds (p. 377) that he is certain it does not add to the danger of the disease. MM. Rilliet and Barthez (*Mal. des Enfants*, 2ème ed., t. i, p. 337) say that "the utility of tracheotomy in the treatment of croup cannot at this day be denied; numerous cases of children snatched from a certain and imminent death, reply victoriously to any doubts that may be raised as to the truth of this assertion." The authors of the *Comp. de Med. Prat.* (t. ii, p. 587) remark that of late years, "the successful operations have been numerous enough to dispel the unfortunate prejudices which tracheotomy has hitherto inspired." M. Valleix (*Guide du Méd. Prat.*, t. i, p. 388) says that the reported recoveries are "now too numerous to allow any one to think of opposing the operation except by statistics." MM. Hardy and Behier (*Trait de Path. Int.*, 1850, t. ii, p. 496), in speaking of the contest in regard to the propriety of the operation, say, "But the question seems now to be definitely settled; the operation has succeeded in fact in a little more than one-fourth of the cases in which it has been performed, and, in presence of these results, it may be said to become the duty of the physician to have recourse to it whenever, notwithstanding an appropriate treatment, the general and local symptoms indicate the extension of the false membrane."

M. Bouchut (*Trait. des Mal. des Nouv.-nés*, 2ème ed., p. 316) says, that when medical means have failed, and the disease has produced a "state tending toward asphyxia, in which an attack of suffocation might cause the death of the child, there should be no hesitation; a new route must be artificially opened to the external air; tracheotomy must be performed."

At the time most of the above expressions were written, a comparatively small number of operations had been placed upon record in France, but they were sufficiently numerous to show conclusively that, if the

operation were carefully performed, and the after-treatment skilfully conducted, from 25 to 33 per cent. of the cases would recover. This excellent result is to be in great part attributed to the improvements introduced by Trousseau, and subsequently by other operators, both in the mode of performing the operation, and in the after-treatment of the cases.

Since the publication of the last edition of this work the operation has continued to be so frequently performed in France, that we cannot find space to quote the results obtained by individual operators. The aggregate of their reports, however, as collected by Roger and Sée, Chaillou, Barthez, etc., yield a result of about one recovery in four in a series of over 500 cases.

The proportion of recoveries has varied considerably in different years in accordance with the type of the epidemic; in some years, as 1858, falling as low as 1 in 6.9, or even 1 in 9, as shown by the statistics of the St. Eugénie Hospital for 1876, as quoted by Bergeron, while in other years it has risen even higher than 1 in 3.

It is further to be remembered that these French statistics are chiefly derived from the reports of the Hôpital des Enfants in Paris, and refer, therefore, to a poor class of patients, who have in many instances been subjected to improper and debilitating treatment before reaching the hospital, and who are exposed to unfavorable hygienic conditions while in the institution. When these unfavorable circumstances are allowed their full weight, it must be conceded that the operation of tracheotomy has achieved a considerable share of success in France, and has fully justified the eloquent and enthusiastic advocacy of Trousseau.

In America, tracheotomy has been resorted to but rarely until within the past few years. The statistics which have been lately published, however, fully suffice to show that, in the hands of American physicians, it has been very nearly, if not altogether, as successful as it has been abroad. Dr. H. H. Smith (*Oper. Surg.*, 2d ed., vol. i, p. 473) gives the results of 26 operations performed in this country, of which 9 recovered. Dr. Gay (*Boston Med. and Surg. Jour.*, Jan. 27, 1859, et al.) reports 13 operations, with 7 cures and 6 deaths; and other operators in Boston have performed the operation in all 15 times, with 7 cures and 8 deaths. But by far the most extensive statistics have recently been published by Dr. A. Jacobi, of New York (*Amer. Jour. of Obstet.*, May, 1868, pp. 13 to 65), derived exclusively from the practice of physicians in that city.

The following table shows the results obtained:

Operator.	No. of cases.	No. of cures.	Percentage of success.
Jacobi,	67	13	19½
L. Voss,	43	10	23½
E. Krackowizer,	55	16	29
W. Von Roth,	48	11	23
Total,	213	50	23½

In this city the operation has been as yet but seldom resorted to, and with but moderate success, owing to the fact that in nearly every instance it has been postponed until the child was almost moribund. The follow-

ing tables embraces certainly the great majority of the operations that have been performed; for a knowledge of which we are to a great extent indebted to the courtesy of the operators, since but few of them have as yet been placed on record:

Name of operator.	No. of cases.	No of cures.
Physick,	2	0
Goddard,	2	0
Page,	1	0
J. Pancoast,	6	3
R. J. Levis,	19	3
T. H. Bache,	1	0
A. Hewson,	1	0
H. Lenox Hodge,	9	2
J. H. Packard,	5	1
T. J. Morton,	4	2
Goodman,	3	1
Drysdale,	12	5
Nancrede,	4	1
Cohen,	10	1
Total,	79	19, or 24.05 per cent.

Finally, to sum up the statistics given above, although even this aggregate does not include by any means all recorded cases, Jacobi states (*loc. cit.*), that out of 1024 operations of tracheotomy, performed in various parts of the world, but principally in Europe, 220 or 21.48 per cent. recovered.¹

It is evident, therefore, that wherever this operation has been practiced in true croup, a considerable proportion of cures has been effected; but in order to form a clear opinion as to the real merits of the operation, it is necessary to have some idea as to the number of subjects that might have recovered without resort to it.

This is very easily arrived at in this country, since we believe that it is never performed here except as an ultimate means of relief, when the patient is manifestly in great danger of death, or absolutely moribund.

In regard to the French operations, it is not so clear whether some of the patients, who recovered after the operation, might not have been so fortunate without it, particularly as M. Trousseau formerly recommended that it should be performed so soon as we can be certain that the larynx contains false membranes. But then it is generally understood that he was not called to many of the cases upon which he operated until all other means had failed, and the child had fallen into an apparently hopeless condition. To elucidate this matter, we shall quote the statements made by M. Valleix, one of the most accurate and impartial of writers. M. Valleix (*loc. cit.*, pp. 388-9) tells us that he collected together 54 cases of undeniable, well-marked true croup, treated without the operation, and found that 17 had been cured. Then, examining what had occurred in regard to the operation, he found, as M. Bricheteau had done before, that nearly 1 in 3 had recovered, a success almost precisely the same as had

¹See also Kröulein, *Medical Times and Gazette*, March 30, 1878; and *New York Medical Record*, July 7, 1877.

taken place in the cases treated by medical means alone. "But," he goes on to remark, "there is a consideration of very great importance, one which gives an altogether different value to tracheotomy, to wit, that in the immense majority of instances, the operation was performed under the most discouraging circumstances, and only when all other methods of treatment had proved useless, and the severity of the symptoms, and the near approach of asphyxia, indicated impending death. . . . So that it follows that tracheotomy should be regarded, in connection with croup, as a genuine medical victory, the honor of which belongs to M. Bretonneau, and all preconceived views should fall before the actual facts." We have here the evidence of a most competent witness, living on the spot, to convince us that the operation is not resorted to in France, at least generally, early in the disease, but is performed only as a last resource, when the chance for the patient from the efforts of nature, or from medical means, is almost *nil*. How, then, can we resist the conviction that tracheotomy does afford a sufficient probability of success, after other means have failed and death is fast approaching, to render a recourse to it at least justifiable, if not almost compulsory?

The second point to be examined in discussing the propriety of the operation is, whether it be in itself dangerous.

From the opinions expressed by authors upon this subject, it appears that the only serious danger attendant upon the operation is the occurrence of hemorrhage. When performed for the removal of foreign bodies from the air-passages, the patients almost always recover if the foreign body do but escape. M. Ollivier (*Art Larynx, Corps Etrangers, Dict. de Méd.*) says that the success of the operation is, so to speak, certain, when it is performed early. Liston disapproves of the operation in croup, but states that it is not attended with much danger. Skey regards it as an operation of some difficulty and danger, from the irregularity in the distribution of the vessels, and the existence of numerous veins which may bleed profusely. M. Boyer does not regard it as dangerous, and states that the only danger is from the occurrence of venous hemorrhage into the trachea, and not from the amount of blood lost. Chelius says that it is dangerous below the cricoid cartilage from anastomosis of the thyroid arteries, from the presence of venous plexuses, and sometimes from a deep thyroid artery, Velpeau speaks of the venous hemorrhage as alone dangerous. Trousseau states that he has performed it more than 200 times, and has met with but a single fatal accident in all of these. Dr. Pancoast, of this city, who has operated in more than 6 cases of croup, and a number of times for the removal of foreign bodies in the air-passages, has never met with any serious difficulty in the performance of the operation, nor with any accident which he could suppose might have affected the life of the patient. Dr. H. H. Smith (*op. cit.*, p. 474), when commenting upon the great disparity of the mortality after tracheotomy, when performed for removal of foreign bodies, and when for the relief of croup, remarks that it is very evident that the dangers which ensue upon incising a healthy trachea are comparatively slight, and that the great mortality which has attended the operation when performed for the relief of croup, must be due to some other cause than the mere incision of the windpipe.

If, then, it is the uniform testimony of those experienced in the matter that the operation is in itself alone but slightly dangerous to life, so that its performance adds but little to the danger of the patient; if it affords immediate relief to the suffocation which threatens to be soon fatal, and at least gives additional time, during which the gravity of the disease may subside; if, further, as we think has been most conclusively shown by the statistics quoted, it has unquestionably saved the lives of a considerable number of those upon whom it has been performed, it is difficult to avoid the conclusion that it is our imperative duty to resort to the operation under certain circumstances.

That some who have been operated upon might have recovered without it, is highly probable; but the uncertainty as to the absolute necessity of resorting to it in any individual case is not even so great, probably, as that which exists in regard to many other surgical operations, and to many medical applications.

Our own plan, then, is to try faithfully all medical means; and, being satisfied of their powerlessness and of the certainty of a fatal issue to the case without the performance of tracheotomy, to inform the parents of the inability of mere medical means to afford relief, and to propose the operation to them, setting before them the great probability of its not averting death, but still strongly pointing out the fact that it does not add to the danger of the case, but gives so much additional chance for life that about 1 in every 4 operated upon recovers.

Should they throw the whole responsibility upon us, we should, without hesitation, advise the operation. Our grounds for so doing are very simple, and have been before indicated. The operation does assuredly frequently save life. It is not in itself attended with any great danger. It cannot increase the danger of the patient's position, but certainly gives an additional chance of escape from the disease; and lastly, it mitigates, in a remarkable manner, the sufferings of the patient. On several occasions, indeed, we have been told by the parents, after the death of their child, that they were very glad it had been performed, since, at all events, it had removed the frightful gaspings and strugglings for breath which had previously convulsed the whole frame of the poor little sufferer, and had rendered its last hours easy and tranquil.

If we decide that tracheotomy is justifiable, it becomes all-important to determine the period of the disease at which we should have recourse to it.

M. Trousseau formerly laid down the rule that it was to be performed as soon as it was certain that false membranes had formed in the larynx. He fixed upon this as the proper moment, because he believed that death was, under these circumstances, almost inevitable without the operation.

This opinion is, however, readily proved to be untenable. We have already learned from M. Valleix that of 54 perfectly well-marked cases collected by himself, treated medically (without the operation), 17, or about one-third, recovered. If we add to this, that of 35 cases seen by ourselves 15 recovered without the operation, it becomes very clear that the mere presence of the exudation in the larynx is not sufficient warrant for a resort to the operation.

Accordingly, most authorities advise that we should wait until medical means have been fairly tried. Thus, MM. Rilliet and Barthez (*op. cit.*, t. i, p. 340), in discussing the period at which the operation ought to be performed, conclude that it should not be resorted to until the means that have succeeded in other cases have been fairly tried, and it has become evident that they must fail. They advise the practitioner not to wait, however, too long a time, but to operate even early should the patient suffer a paroxysm of suffocation so severe as to make it probable that another might prove fatal. So, too, Mr. James Spence, in a valuable paper on tracheotomy (*Edin. Med. Jour.*, Feb., 1860), states, as the result of his large experience, "that if, in a case of croup, all measures have been actively tried, if the hard ringing cough has become suppressed, and the respiration is evidently imperfect, as shown by the contracted and depressed appearance of the cartilages of the ribs, and the occasional severe paroxysms of dyspnœa, the operation is fully warranted. When the paroxysms become more and more frequent, and when the dyspnœa is rather persistent than paroxysmal, with turgid or pale lividity, the operation is the little sufferer's only chance for life."

The same course is, we believe, universally pursued in this country, and, as the reader will recollect, corresponds precisely with the advice given in our remarks on the medical treatment of true croup.

The prime indication for the performance of the operation is, then, the degree of laryngeal obstruction as shown by the characters of the respiration, the cry, and cough.

It should, however, be carefully borne in mind that great dyspnœa, or even asphyxia, when intermitting, do not so imperatively claim operative interference, since cases where the dyspnœa is of this character may recover without the operation.

When, however, despite the use of all medical means, and especially the repeated administration of emetics, the dyspnœa grows steadily and progressively greater; when there is marked hissing laryngeal stridor, and, at each inspiratory effort, recession of the base of the thorax; when, in addition, the voice is whispering or suppressed, and the cough short, smothered, and muffled, the operation should, we think, be unhesitatingly performed.

In thus defining the conditions under which tracheotomy is called for in croup, it is clear that we are not to be influenced at all by the mere period of the disease as measured by time, but that, whenever the above symptoms are present, the operation is indicated.

There can be no doubt, however, of the far greater success of the operation when performed in the early period of the attack, before the patient's strength is materially impaired; and it is, therefore, highly desirable that the indications which render its performance necessary should be appreciated so soon as they appear.

A still further argument in favor of the timely performance of the operation is adduced by Dr. George Johnson (*British Med. Jour.*, Jan. 15th, 1870), who dwells upon its value at an early stage, when the indications are present, on account of the danger of œdema of the lung from

venous congestion, and of the coagulation of the blood in the pulmonary artery.

Trousseau, also, in his last publication upon this subject (*Clin. Méd.*, 2d ed., t. i, p. 450), speaks as follows: "I wrote in 1834, and repeated in 1851; so long as tracheotomy was not a trusty weapon in my hands, I said, we should operate as late as possible; but now that I can number many successes, I say, we should operate as early as possible. In removing from this assertion whatever may seem too absolute, I still affirm it, by saying, *that the chances of the success of the operation are so much the greater in proportion as it has been the earlier performed.*"

Notwithstanding this, however, should we be called to a case where the last stage of asphyxia has been reached, it is still not too late to perform the operation. Thus, in one of the cases that occurred in our own practice and which ended favorably, this condition was fully developed, and the bluish skin, drowsiness, and insensibility to pain, showed that the patient had already sunk into very advanced asphyxia.

Perhaps we cannot do better in closing our remarks upon this point than to quote the concise and forcible axiom laid down by Archambault: "We should never operate too late; it is never too late to operate, so long as death is not actually present."

There are, however, certain conditions which have been thought by many authorities to contraindicate the performance of the operation, even under the circumstances above described. The first of these is the very early age of the patient, and it has been advised to refuse the operation in all cases occurring under the age of two years. It is unquestionably true, as might be expected, that age exercises a most powerful influence upon the prognosis after the operation, owing partly to the difficulty in performing it on account of the narrowness of the trachea and the shortness of the neck, but chiefly to the deficiency of vital power, and to the difficulty of nourishing the infant afterwards. Notwithstanding these influences, which render the prognosis so unfavorable in tracheotomy before the age of two years, there are so many successful cases on record that the most tender age can no longer be regarded as a positive contraindication. The following list embraces the names of the operators and the age of the infants in the cases which have been successful at a very early age:

Baizeau,	at 10 months.	Vigla,	at 17 months.
"	" 15 "	Potain,	" 18 "
" (in the hands of his		Moutard-Martin,	" 18 "
colleague),	" 15 "	Trousseau,	" 13 "
Isambert	" 16 "	Barthez,	" 13 "
Archambault,	" 13 "	"	" 7 "
"	" 18 "	Maslieurat Lagemand,	" 23 "
Roger,	" 19 "		

In adults, on the other hand, tracheotomy in croup is less successful than in children, probably because, as Trousseau suggests, the form and size of the larynx allow the pseudo-membrane to extend deeply into the bronchi before producing the symptoms of croup.

There is another condition which, it is thought by many, ought to constitute an insuperable obstacle to the operation, and the possible existence of which, in any case, is one of the most serious objections that has been brought against its performance. The condition to which we allude is the presence of pseudo-membranous exudation in the bronchi.

The existence of this condition must greatly lessen the chances of a successful operation, but that it renders success impossible, as have been supposed, cannot be admitted. MM. Rilliet and Barthez (*op. cit.*, 2ème ed., t. i, p. 338) say: "It has been said that one contraindication was the presence of false membrane in the bronchi. But, besides the fact that the symptoms denoting its presence are uncertain, we cannot see in this a positive objection to the operation. Recovery has been known to occur, in effect, after the rejection of bronchial false membranes, and we were ourselves witnesses of a remarkable example of this kind. And is there any better mode of facilitating the escape of foreign bodies than by opening to them a passage below the larynx? Under such circumstances, we must expect, to be sure, a greater mortality than under more favorable conditions. This opinion is, moreover, that of M. Bretonneau." Numerous cases are indeed on record, and we have ourselves met with such, where, after the operation, large membranous casts of the trachea and bronchi, which could certainly never have escaped through the larynx, have been discharged through the tracheal opening, and their escape followed by complete recovery.

It appears evident, therefore, that if in such cases, when death is even more surely imminent than in those instances when the exudation does not extend below the larynx, tracheotomy affords even a very slight additional chance of recovery, it should be performed despite the fact that the child will in all probability die.

But, apart from this consideration, it must be borne in mind that statistics prove that the false membrane extends below the larynx in about one-third of all cases, and still further, that there are no means by which we can with certainty determine in any individual case whether such extension has taken place or not.

It was at one time thought that auscultation might afford the desired information, but more careful observation has shown that it is not to be depended upon. As already said, in most cases the laryngeal stridor is so loud as to mask all chest-sounds, and even when this does not happen, we have frequently observed that no definite and reliable information is to be gained from physical examination. The following cases may be quoted, out of the number on record, besides several that we have ourselves seen, as proving this statement. MM. De La Berge and Monneret (*Comp. de Méd. Prat.*, t. ii, p. 587) mention a case in which they could not believe that the bronchi contained false membranes, as the vesicular murmur was extremely pure and was heard everywhere; and yet, during the operation, a false membrane was drawn out, which represented the trachea and the division of the principal bronchi. The child died in 15 hours.

The late Professor William Pepper, of this city, reported 2 fatal cases (*Summary of Trans. Coll. Phys.*, vol. iii, No. iii, p. 106), in one of which

"distinct vesicular murmur could be heard throughout the lungs, marked only occasionally by sibilant and sonorous râles," a few hours before tracheotomy was performed. The child died 20 hours after the operation, and the exudation was found to implicate the larynx, trachea, the large bronchi, and even some of the smaller ramifications. In the other case, the state of the respiration was carefully examined the day before death, and not the least respiratory murmur could be heard over any part of the chest, and yet, in this instance, the exudation was confined strictly to the larynx; not a vestige of false membrane was to be found either in the trachea or bronchi.

In a case recently attended by us, where tracheotomy had been performed, so that all laryngeal stridor was absent, auscultation, eight hours before death, revealed quite strong respiratory murmur, much obscured by snoring bronchial râles. The antero-lateral parts of the chest were alone auscultated. Death occurred somewhat suddenly from the lodgment of a very large tubular membrane from the left bronchus in the trachea; and at the autopsy there was a tubular membrane found extending throughout the trachea, and through the right bronchus to its third divisions. The left lung was collapsed and congested; the right one distended and emphysematous.

Since, then, we can learn little or nothing from auscultation, or any other means, as to the presence of false membrane in the bronchi, the question becomes one of expediency, so far as this contraindication is concerned, whether to leave two-thirds of the patients, many of whom could certainly be saved by the operation, to perish without an effort to save them, because one-third *must* probably die; or to perform the operation, with very little prospect of success in one-third, for the sake of the chance of saving many of the remaining two-thirds who must otherwise perish.

The presence of pneumonia is also universally recognized as greatly lessening the chances of recovery after tracheotomy. It must be borne in mind, in regard to this point, that pneumonia is frequently overlooked, and indeed that it frequently cannot be recognized on account of the loud tracheal râles which hide all auscultatory sounds; while, on the other hand, its presence may be simulated by the occurrence of collapse of some portion of the lung, owing to occlusion of the bronchus leading to it. Millard suggests that the degree of dyspnœa may be of service as indicating the presence or absence of pneumonic complication. Thus he has found that in croup not thus complicated the rate of respiration is from 32 to 48, while, when pneumonia is present, it rises above 50. It is probable, also, that by a careful study of the temperature, the occurrence of pneumonia may be suspected by a marked elevation of several degrees. Pneumonia of one lung is not, according to Guersent, a contraindication, nor is even double pneumonia regarded by some operators as absolutely interdicting the operation, though at the same time we are not aware of a single instance in which it has been successfully performed where this condition was unquestionably present.

Another condition in which tracheotomy is thought by many to be contraindicated, is when membranous croup occurs as a secondary affection,

during the course of some constitutional disease other than diphtheria, as for instance, scarlatina, measles, or pertussis. Such cases were regarded even by Trousseau as absolutely unfit for operative treatment. Still, that this contraindication, although of the greatest weight, does not entirely forbid tracheotomy, is shown by a case of croup following scarlatina, in which Dr. Voss operated, and the child survived 31 days, the tracheal wound being nearly closed. Millard, also, in his excellent essay on tracheotomy (*De la Trachéotomie dans le Cas de Croup*, Paris, 1858), records 3 cases of croup secondary to measles, successfully treated by operation. He regards croup occurring in the course of pertussis as far less unfavorable, since the violent cough favors the expectoration of the false membranes.

There remains, finally, one condition to be indicated in which the operation is, in the almost unanimous opinion of authorities upon this question, absolutely contraindicated. We refer to the cases of profound general diphtheritic infection, where the danger of the child depends upon the constitutional disease, even more than upon the laryngeal obstruction, where the blood is gravely altered, and the well-known tendency exists to the formation of pseudo-membranes upon all abrasions or wounds, so that in all probability the operation would merely serve to invite the extension of the exudation.

Trousseau opposes the operation under such conditions, in the following words: "If the diphtheritic infection have profoundly attacked the constitution; if the skin, and especially the nasal passages, are occupied by the specific inflammation; if a frequent pulse, delirium, and prostration show the system to be deeply poisoned, and if the danger is rather from the general condition than from the local lesion of the larynx and trachea, *the operation ought never to be attempted*, for it is invariably followed by death."

Even under this most unfavorable of all conditions, however, there are not wanting some operators of wide experience, who still recommend the operation; thus Jacobi (*loc. cit.*) asserts that whenever the indication of suffocative dyspnœa, steadily increasing and not relieved by emetics, exists, he would operate despite any complications, general diphtheria, or anything else, and uses this powerful language: "Seeing a person suspended by the neck and being strangled, we should hardly investigate the propriety of cutting the rope from the point of view that the sufferer might be or is affected at the same time with tuberculosis, cancer, or diabetes."

After a careful review of the entire question, we believe that the facts upon record justify the following conclusions: that the condition of success which excels all others is the predominance of the characters of asphyxia; that when these are so marked that death is imminent, the operation is justifiable despite any complications which may coexist, save perhaps the presence of grave general diphtheritic infection; and finally that, when no such contraindication is present, and the dyspnœa is continuous and increasing despite all other treatment, the operation is positively indicated, and it becomes the duty of the practitioner to recommend its performance, and, if the decision be entrusted to him, to unhesitatingly assume the responsibility of operating.

We have already indicated with sufficient clearness the influence which the age of the patient, the period of the disease, and the character of the epidemic exert upon the results of tracheotomy. But we would again allude to the marked manner in which the result is modified by the character of the previous treatment, and to the fact that its success is very much interfered with by the earlier employment of any debilitating measures, such as were, until lately, but too frequently adopted.

We have more than once been asked by the parents of children, upon whom tracheotomy was about to be performed, or who had actually undergone it, what influence would be exerted by the effects of the operation, should it be successful, upon a subsequent attack of croup; and since, as has already been seen from the cases quoted by us from our own experience, second attacks of croup are not very rare, it is interesting to know, that so far the statistics which bear upon this question tend to show that a previous attack of croup cured by tracheotomy is a favorable condition for its performance in a subsequent attack. Thus of 5 cases, collected by Millard, in which the operation was performed for the second time, every one recovered. The second operation was uniformly found much easier, on account of the cicatrix of the former incision serving as a guide, and also on account of the slight amount of the hemorrhage.

MODE OF PERFORMING THE OPERATION.—Tracheotomy being an operation which all physicians, whether experienced or not in the use of surgical instruments, are liable to be called upon to perform at a moment's notice, no apology is needed for the introduction here of the details of its performance. The following account is in great part borrowed from the pages of that most experienced tracheotomist, Trousseau,¹ and from a very complete and practical discussion of the operation by F. Howard Marsh, Esq.²

The child should be carefully wrapped up, so as to avoid all exposure to cold; and if an anæsthetic is to be employed, should be allowed to sit or lie in any position he may choose during its administration, as the constrained position necessary during the operation tends to increase the difficulty of breathing. He should then be placed upon a table, furnished with a thin mattress, and a folded pillow or roll of cloth should be placed under the shoulders and back of the neck, so as to put the skin of the throat upon the stretch, and render the trachea prominent. If the operation is performed during the day, the table should be drawn close to the window, and the patient's face directed *toward* it, so that a full light may fall upon the throat; if, however, it be at night, and there is not sufficient gaslight, a special assistant must be intrusted with the duty of holding the candles or lamp. An assistant is also needed to stand behind the patient and hold the head securely; and another, whose duty it shall be to draw aside the successive layers of tissue and the bloodvessels with a hook, and to sponge the wound from time to time.

The instruments needed are a sharp-pointed, slightly curved bistoury; a blunt-pointed bistoury; two flexible hooks; a dilator to stretch the

¹ Clin. Méd., 2ème ed., tom. i. p. 414 et seq.

² St. Barth. Hosp. Rep., vol. iii, p. 331 et seq.

incision in the trachea, so as to favor the introduction of the canula, and made like a pair of curved dressing-forceps, with a little spur projecting backwards, so as to catch in the tissues and prevent its displacement; and finally a canula. The size and form of this canula are matters of great importance, and of late years several marked improvements have been effected in them. The calibre of the canula should, as first clearly directed by Trousseau, be as large as possible without interfering with its easy introduction into the trachea, and its curve should be that of a quarter of a circle.

In regard to this very important question of the size of the canula, we are indebted to Mr. Marsh (*loc. cit.*) for a series of observations, which appear to indicate that a tube somewhat smaller than that recommended by Trousseau, Fuller, and others, may be equally efficient and yet less irritating. By a series of careful measurements of the respective diameters of the trachea and cricoid cartilage, he established the fact that the latter diameter is almost invariably less than that of the trachea, to an extent varying from $\frac{1}{40}$ th to $\frac{6}{40}$ ths of an inch. If, therefore, as his measurements show, the diameters of the trachea are as follows; during the first two years of life, $\frac{3}{40}$ ths of an inch; in the third year, $\frac{11}{40}$ ths; in the fourth and to the seventh, $\frac{14}{40}$ ths; in the eighth and ninth, $\frac{15}{40}$ ths; and in the tenth, $\frac{16}{40}$ ths; it will be seen that a canula having a diameter of $\frac{19}{40}$ ths of an inch will answer for children between the ages of 1 and 4 years; one of $\frac{11}{40}$ ths for children between 5 and 8 years; and one of $\frac{13}{40}$ ths for children between 9 and 12 years old.

It may be added, that after the 12th year the diameters of the cricoid cartilage and trachea increase so rapidly, that the canula now usually made for adults, with a diameter of $\frac{14}{40}$ ths of an inch, is rather small for children between 14 and 16 years old.

The length of the canula should be sufficient to cause it to reach from $\frac{1}{2}$ to 1 inch below the inferior angle of the wound in the trachea.

The canula must also be double, the outer tube having a broad collar in front, with holes through which the band which passes around the neck and secures the canula in position may be passed and tied. It should also be furnished with a key, which plays easily in a notch on the upper part of the inner tube. This inner tube must so fit the larger one as to be readily removed and replaced, being secured in position by the little key above mentioned.

In some canulas a still further improvement is introduced by having the outer tube and collar merely yoked together by means of two arches on the collar, which receive small outjutting bars at the sides of the upper extremity of the outer tube, so that this can shift its position according to any pressure it may receive.

There is also a canula recommended by Fuller, called the "bivalve canula," the outer portion of which is not a tube, but consists of two narrow lateral blades, which are easily compressed by the finger and thumb into the form of a thin wedge, and expand again when the pressure is removed. This instrument supersedes the need of any dilator, and has the great advantage of being readily introduced. It is evident, however,

that it must produce much more irritation while in position than a tubular canula, and in addition, when the inner tube has been removed, as is frequently required, its reintroduction causes pain and irritation, from the constriction of the mucous membrane, which has bulged inwards between the blades of the outer portion. Mr. Marsh, therefore, advises that when there is any difficulty in introducing the canula at the time of the operation, a Fuller's tube should be used, but that this should be exchanged on the second day for one whose outer portion is tubular.

Although it is almost the universal practice to introduce a canula at the time of operation, its use has been objected to by several good authorities, as apt to cause inflammation and ulceration of the trachea, and to favor the development of pulmonary complications; and several plans have been suggested for the separation of the edges of the tracheal wound. Thus Mr. Adams, of the London Hospital, recommends the introduction of a strong metallic wire speculum, such as is frequently used in operations on the eyes, and Dr. Pancoast, of this city, employs a pair of blunt leaden hooks.

In addition to the instruments already enumerated, some operators, following the practice of the Dublin surgeons, use a hook or tenaculum to fix the trachea, while the incision is being made through its rings. This proceeding has certain advantages, especially when it is designed to excise a portion of the trachea, or in case of venous hemorrhage, as the trachea can be raised above the pool of effused blood and speedily opened, which will usually check the bleeding. It is also of service in young children, because the trachea is then so pliable and yielding, that, unless the hook be used, its anterior wall may be easily driven in front of the point of the scalpel, till it is nearly or quite in contact with the posterior one, in which case the latter also may be wounded. Trousseau, Millard, and others, however, strongly objected to this practice, believing it to be dangerous to so fix the trachea and oppose the movements connected with the performance of the function of respiration which is already so much impaired. Our own observation would go to show that, while the advantages to be gained from fixing the trachea are undoubted, especially in young children, the dangers have been somewhat exaggerated.

It has been recommended by several high authorities—Lawrence, Carmichael, G. H. Porter, Brainard, Fergusson—to excise a small piece of the walls of the trachea. By some this has been adopted with the view of dispensing with the use of a canula, but it is claimed that, even when one is employed, this practice renders its introduction more easy; that the tube fits the oval opening thus made much more accurately than a mere slit, produces less pressure upon the edges, and consequently is not so apt to cause caries of the tracheal rings. It seems never to be followed by narrowing of the trachea after the canula has been removed, as might be apprehended.

This practice is followed by Dr. Pancoast, of this city, who, in the case he describes, excised an elliptical piece about one-third of an inch long and two-tenths of an inch broad, from the front part of the third, fourth, and fifth rings of the trachea. As already said, he does not employ either

a canula or dilator, but holds apart the edges of the wound made in the soft parts over the trachea by means of a piece of thick leaden wire, bent so as to form hooks at either end. The wire is of such a length as to fit accurately around the neck when the hooked ends are placed within the edges of the incision, and thus keep up just sufficient traction in opposite directions to maintain the wound open.

In regard to the operation itself, almost all who have had much experience in it direct that it must be performed with great deliberation and care.

The incision through the skin should be made precisely in the median line of the neck, and should extend from the cricoid cartilage to a little above the sternum. The slight white fibrous line which marks the interspace between the sterno-hyoid and sterno-thyroid muscles should then be followed as a guide for the next incision, and the muscular masses drawn aside by hooks.

The trachea is now exposed with the isthmus of the thyroid gland, and occasionally, large thyroidean veins lying upon it, and great care must be observed to avoid wounding these on account of the troublesome hemorrhage which is apt to follow. A still further reason for this caution is the occasional existence of an anomalous distribution of arteries, by which a branch of considerable size, or even the innominate artery itself, passes over the trachea directly in the course of the wound. Any blood-vessels may be drawn aside by hooks, and the isthmus of the thyroid gland may either be treated in the same way, or if it cannot be drawn away far enough to allow a sufficient incision of the trachea, may be ligated in two places and divided between (Brainard, of Chicago), although, when possible, this had better be avoided. The trachea, having been thus carefully exposed, should be punctured just below the cricoid cartilage, and the probe-pointed bistoury being introduced, and its edge guarded by the nail of the index finger of the left hand, the opening should be enlarged downwards to the extent of two or three tracheal rings.

It usually happens that there is some hemorrhage during these incisions; but if it be venous and moderate in amount, the opening of the trachea should not be deferred, as the re-establishment of respiration will usually speedily check it.

So soon as the trachea is incised, the dilator should be instantly introduced with the blades closed; and so soon as in position these should be moderately opened. Air now enters readily, and there is a discharge of mucus, fragments of false membrane, and blood, through the opening. The canula should then be introduced upon the dilator as a guide, its entrance being evinced by the increased facility of respiration, and the escape of mucus and blood through its calibre. A guard of india-rubber or a disk of waxed cloth should then be placed between the guard of the external tube and the skin, to prevent any irritation or chafing, and the canula may be fastened in position by a tape passed around the neck.

Should blood bubble up by the side of the canula, as Géraldes observes, the wound in the trachea has been made too large, so that the blood gains entrance during inspiration, and a larger canula should be at once substituted.

It occasionally happens, as in a case related by Trousseau, that the trachea is lined by a false membrane, which is partly detached and pressed forward by the end of the canula, so that it completely occludes the opening, and thus even increases the asphyxia. When this occurs, the canula should be withdrawn, and an attempt made to seize the false membrane with forceps and withdraw it.

When the operation has been a laborious one, emphysema of the neck may be met with, sometimes extending to a considerable distance, and causing great disfigurement or even seriously complicating the course of the case. It results from a want of parallelism between the cutaneous and tracheal wounds, or from marked disproportion between the size of the tracheal wound and that of the canula, or, as occasionally may happen, from the escape of the canula from the tracheal wound. It has also happened that the inflamed and thickened mucous membrane is stretched over and driven before the point of the scalpel, and so escapes a sufficient division.

It has not been customary to use anæsthetics in the performance of tracheotomy. Fock, however, advises the use of chloroform, and states that he has never, even in extreme dyspnœa, found any ill effects to result from its employment. At first the dyspnœa is increased by the inhalation, but anæsthesia is speedily established, and then the breathing becomes much calmer than before. Dr. Voss, who has also employed it, reports equally favorably of its effects; and Mr. Marsh, who has seen it administered in at least twenty cases, believes that, when carefully and slowly given, it is most beneficial. It must be remembered, however, that, owing to the asphyxia, the sensibility of the child is usually much blunted, so that, even without anæsthesia, the operation has appeared to us to cause but slight pain, and has been borne by the little patients with scarcely any struggling.

AFTER-TREATMENT.—Immediately after the successful performance of the operation, and the satisfactory adjustment of the canula, an almost incredible change occurs in all the symptoms of the patient. The wild restlessness of the little sufferer, with the agonized, appealing glances at those surrounding the bedside, and the frantic clutching at the throat as though to tear it open to admit air, the lividity of the surface, the noisy, hissing stridor of the respiration, all vanish as though by magic. Very frequently the child falls into a placid sleep, the skin and lips regain their normal color, and the breathing becomes regular, full, and nearly as silent as in health. This calm is not, however, to be of long duration; there are frightful dangers still to be undergone, from which nothing but the most assiduous care and skilful treatment can enable the patient to escape with life.

It may, in fact, be asserted that the much greater proportion of success which has of late years attended this operation is to be attributed chiefly to the more judicious after-treatment which patients receive. Indeed, Trousseau has most truly said, with regard to the importance of this portion of the management of the cases, that tracheotomy, badly performed, but well treated afterwards, will end favorably in a third of all cases;

whereas, tracheotomy excellently executed, but badly treated afterwards, will almost invariably be followed by a fatal termination.

It might, consequently, have been added to the contraindications already enumerated, that, unless we can secure constant and skilful attendance upon the case after the performance of the operation, there can be but little hope of obtaining a favorable result.

Wherever it is in any way possible, the constant presence, by day and night, of a physician or student of medicine, should be secured for four or five days after the operation. When this is utterly impossible, all of those engaged in nursing the case should be carefully instructed how to act in the event of any emergency, so that the child shall never be without the presence of some one competent and ready to render the prompt assistance which is frequently necessary to avert instant death. The details of the attention necessary will be given a little further on.

One of the first points to which careful attention must be paid, is to give to the air to be inspired through the canula as much as possible the temperature and degree of moisture that the air attains by its normal passage through the mouth and nose. Various means have been recommended to secure this object; thus a piece of loose coarse sponge, wetted with tepid water, and enveloped in a piece of gauze, may be applied over the canula; or, as directed by Trousseau, "the neck of the child may be surrounded by a cravat of knitted wool, or a large piece of muslin or gauze, so that the patient expires into this thick tissue, and inspires the air impregnated by the warm watery vapor which the expiration has just furnished."

This was the only means adopted by Trousseau; but we may, in addition, by the aid of a spirit-lamp, keep shallow dishes of water evaporating in the room, and at the same time employ a thermometer to regulate the temperature of the chamber, which should be uniformly kept at from 70° to 72° F., though the air should be changed frequently, so that it may be pure and fresh.

By careful attendance to this clear but long-neglected indication, we not only prevent the rapid drying of the mucus in the canula and trachea, but, as Trousseau asserted, avoid to a great extent the occurrence of pneumonia or bronchitis as sequelæ of the operation.

In regard to the treatment of the wound itself, we have already alluded to the advantage of placing a piece of lint spread with cerate, or a caoutchouc ring, beneath the collar of the canula to prevent any irritation of the skin. No sutures should be introduced into the skin incision, as the efforts during coughing will soon tear them out. Trousseau strongly advised that the edges of the wound should be cauterized daily for the first three or four days, with solid nitrate of silver, in order to prevent the formation of diphtheritic deposit.

It very soon becomes necessary, despite every care to render the inspired air moist, to cleanse the inner tube of the coating of viscid, partly dried mucus which collects on its interior, and to effect this, the inner tube should be removed as frequently as is necessary. The frequency with which this withdrawal is required varies in different cases, but it may be stated as a

general rule, that it should be performed from four to twelve times in twenty-four hours.

When the tube is clear, the respiration is almost noiseless, and hence the supervention of noisy breathing is usually the indication of some obstruction in the inner tube, which should immediately be withdrawn and cleaned.

The drying of the mucus in its interior may be partially prevented by dropping, every half hour, a few minims of tepid water into the mouth of the canula, and by smearing the inner surface of the tube with pure glycerin every two or three hours. Some years ago, Barthez¹ recommended instillations of tepid solutions of chlorate of soda through the canula after tracheotomy, in the hope of effecting the softening of the false membranes, and their more rapid and complete expulsion. Although he was inclined to attribute a beneficial effect to the practice at the time, it appears to have since fallen into disfavor even with its originator.

We have ourselves employed lime-water in several cases, and always with obvious relief. We were induced to use it from its well-known solvent action upon pseudo-membranous exudation, and have generally employed it by atomizing warm lime-water through the canula every few hours, or so often as the breathing becomes noisy and labored, despite the removal and cleaning of the inner tube, from the collection of viscid mucus or pseudo-membrane below the end of the canula. The atomization has been continued for a moment or two, and has usually excited cough, while at the same time it softened the viscid mucus and enabled the child to reject it through the tube. So great, indeed, is the relief at times thus afforded, that in one case the little patient asked frequently that the use of the atomizer should be repeated. In all probability it does good, partly by its mechanical action in exciting cough, partly by the softening effect of the watery spray, but partly also, we are inclined to believe, by the action of the lime upon the mucus and pseudo-membranes. We are also in the habit of directing that the child shall breathe, for a few minutes in every hour, the steam from slaking lime, though in all probability this does not contain an appreciable amount of the lime itself.

Recently Dr. Böcker, of Berlin (*Deutsche Klinik*, July 8th, 1876), has reported the very favorable results he has obtained by means of frequently repeated inhalations, after tracheotomy, of the spray of lime-water solution, of chlorate of potash, or dilute lactic acid. Dr. Burrer, who has adopted the same practice, reports eight cases in which these frequent inhalations formed part of the after-treatment; of these eight cases only two died, while of ten cases, tracheotomies performed by the same operator before his adoption of this mode of treatment, seven were fatal.

It occasionally happens, however, that the breathing becomes noisy and obstructed, and remains so even after the withdrawal of the inner tube and the use of the atomizer. The cause of the obstruction, then, probably consists in the presence, near the end of the canula, either of a collection of dried mucus, or of a piece of false membrane, too large to escape through the canula. If, under these circumstances, a paroxysm of dyspnoea should

¹ Bull. Gén. de Thér., May 30th, 1858.

ensue, the strings securing the canula should be instantly cut and the outer tube withdrawn. If this be followed by the rejection of false membrane and a return of quiet respiration, the canula may be returned; but if there is reason to fear that the trachea contains false membranes too large to escape through the tube, it is better to allow it to remain out permanently.

Millard (*loc. cit.*) recommends that the external tube should always be removed at the end of twenty-four hours after the operation, when the track of the wound is usually patulous, being lined by plastic lymph, and after waiting a few minutes for the rejection of false membranes, and cauterizing the wound, be again introduced.

In those cases which progress favorably, it soon becomes necessary to decide at what date the canula shall be finally removed. It is evident that this should be accomplished so soon as possible, as the tube acts the part of a more or less irritating foreign body in the neighborhood of delicate and important structures, and yet it is only in rare cases that the patient can endure its removal before the sixth or seventh day.

At the end of the fifth day, therefore, the experiment may be tried of plugging the mouth of the canula with a little roll of wool, to learn in what degree the larynx has become patulous. Should the child be unable to take a single respiration, the experiment may be deferred for several days, but should breathing be performed through the mouth for several minutes, the measure may be repeated daily, in order to gradually accustom the larynx to a resumption of its function.

About the seventh or eighth day the tube may be removed for an hour or two; and, if its abstraction be well borne, it may be finally withdrawn the following day, and the wound closed by bringing its edges together with adhesive plaster. It is very necessary to observe the caution, that the canula must never be removed unless some one competent to replace it is at hand. It occasionally happens, however, that the larynx remains impervious for a much longer time, and cases are recorded in which it has been impossible to remove the canula for fifteen, twenty-five, forty-four (Trousseau), or even one hundred and twenty-six (Fock) days; or even for months or years. The causes which thus delay the period at which the tube can be removed, are summed up by Mr. Marsh (*loc. cit.*), as follows:

1. Obstruction of the larynx by false membranes, which have been known to linger in its cavity for at least fourteen days after the operation.
2. A chronic inflammation and thickening of the mucous membrane of the larynx, which may remain after the acute disease has passed off.
3. A narrowing or complete obliteration of the passage of the larynx, by the growth of granulations above and around the canula.
4. An impairment or complete loss of those functions of the muscles of the larynx which regulate the admission of air through the rima glottidis.
5. Adhesions of the opposed surfaces of the vocal cords.

After the removal of the tube, the wound heals, either by contracting from the circumference toward the centre, when air escapes until the very last day; or the tracheal wound first closes, and the cicatrization then advances externally. The average time occupied by this process of cica-

trization is about one month, though it may be completed in two weeks, or be protracted for two months.

Among the results which have been known to follow the prolonged stay of the canula in the trachea, are necrosis of the tracheal cartilages, and ulceration about the wound, or of the trachea around the canula, which in several cases, has been followed by fatal hemorrhage. Suppuration among the deeper structures of the neck, even extending into the anterior mediastinum, has been noticed in a few instances, when the deep-seated tissues of the neck had been much disturbed.

GENERAL TREATMENT.—Having carefully discussed the management of the canula, and the treatment of the tracheal wound, it remains to say a few words in regard to the general treatment of the patient after the operation.

The most essential point to be secured is, unquestionably, the proper alimentation of the child. It is, however, frequently very difficult to induce it to partake even of the most tempting food. We should endeavor to persuade it to take, as before the operation, nourishing animal broths, beef-tea, milk, custard, chocolate, wine-whey, or weak milk-punch. If, however, these are refused, and the child expresses a desire for any other digestible article of food—as the breast-meat of fowl, finely minced, or the soft portions of oysters, or eggs—the taste should be gratified. Occasionally ice-cream will be taken willingly, when other food is refused; or, when both wine-whey and milk-punch are rejected, iced wine and water, or brandy and water will be relished. Unfortunately, however, it not rarely happens that, owing partly to the soreness of the throat and partly, undoubtedly, to the pain caused by the canula during the movements of the trachea in deglutition, the little patient utterly refuses to swallow more than a mere sip of iced water. Under such circumstances, so serious a complication is abstinence, that Trousseau recommends that it should be forced to take a little food. “Do not fear,” he says, “to employ intimidation. In such cases I have often—assuming an apparent severity, the expression of which I have exaggerated—forced the child to eat, and so have prepared the way for a recovery, which without this seemed to me impossible.”

Even by this means, however, it may be impossible to secure the administration of a sufficient amount of nourishment, and we would then advise the use of nutritious enemata, consisting either of the yolk of one egg beaten up in an ounce of milk, or of one ounce of beef-tea, and given about every four hours. If they appear to irritate the rectum, and are not retained, one or two drops of laudanum may be added to each enema.

In comparatively rare cases there exists, in addition to this unwillingness to eat, a positive difficulty in swallowing liquids. This results from the inaction of the vocal cords and epiglottis, which allow the fluid to pass through the glottis into the trachea and bronchi, causing violent cough and escaping through the artificial opening. The child is so alarmed by this that it sometimes refuses all nourishment, and can only be supported by nutritious enemata. Under these circumstances, Trousseau advised that

all liquid aliment should be interdicted, and that the food of the child should consist of very thick soups, vermicelli boiled in milk or broth, hard eggs, eggs very much cooked in milk, and rare-cooked meat, in rather large morsels. If the thirst becomes ardent, he allows pure cold water, taking care to give it either some length of time after, or immediately before, the meals, in order to avoid vomiting. This difficulty in swallowing rarely begins until three or four days after the operation, and does not usually last beyond the tenth or twelfth day. Sometimes, as M. Archambault has suggested, the child is enabled to swallow with ease by closing the canula with the finger at the moment of deglutition, but at other times this fails entirely.

In many cases the difficulty in inducing the child to swallow, after the operation, is so great that all medication must be suspended, excepting the administration of small doses of opium, by the mouth or by enema, which we would advise to be continued.

Whenever it is practicable to give remedies, however—without interfering with the ability and willingness to take food—it is very important to bear in mind that despite the very great relief which the operation may have afforded, it has by no means put a stop to the disease, but has simply afforded the system another chance to overcome and cast off the constitutional affection.

Of course, the use of emetics must be suspended, and so if, on any theoretical ground, any depressing remedies have been employed, they should be discontinued. But we should recommend under such circumstances, that the use of the combination of chlorate of potash, tincture of chloride of iron, and sulphate of quinia, should be persisted in.

We subjoin the histories of two cases of true croup, which have lately occurred in our practice, and which will serve to illustrate clinically the remarks that we have made upon this disease.

In both cases tracheotomy was performed by Dr. H. Lenox Hodge, in one instance with complete success, but in the other with a fatal result.

The first and successful case was under the care of Dr. R. Bolling, of Chestnut Hill, and was seen in consultation by Dr. J. F. Meigs, and, subsequently to the performance of the operation by Dr. Hodge, by Drs. Edward Rhoads and William Pepper.

The second case was visited by both of us from the first; and was attended, after the performance of the operation by Dr. Hodge, with the most zealous and skilful care, by Drs. Wharton Sinkler and M. Longstreth.

Angina with Membranous Exudation on Tonsils—Membranous Laryngitis: Tracheotomy at end of second day—Complete Recovery.

CASE I.—F. W., æt. 7½ years, a delicate child, who at the age of four years had suffered from a severe attack of true croup, from which he recovered without the operation. On December 23d, 1868, he was noticed to have the symptoms of an ordinary cold in the head, with slight sore throat, some dysphagia, and laryngeal cough; he was visited and prescribed for by Dr. R. Bolling. On December 24th the cough persisted, and there was slight coryza and redness of fauces, but without any membranous deposit or any croupy symptoms.

He was ordered small doses of Kermes mineral, Dover's powder, and nitrate of potash, and counter-irritation to the throat.

At 5 A.M., December 25th, the child, who had gone to sleep quietly, woke in a frightful paroxysm of dyspnœa, gasping, clutching at its throat, and with oppressed whispering voice.

Emetics of alum were given, and produced free emesis, but without the rejection of any false membrane, nor was any yet visible in the fauces. The powders were continued.

The dyspnœa persisted and grew steadily worse; the voice remained suppressed. Membranous exudation was noticed in the evening on the tonsils, and during the following night the obstruction to respiration became so intense that, after consultation with Dr. Meigs, tracheotomy was performed by Dr. H. Lenox Hodge. The trachea was opened just below the isthmus of the thyroid gland. No false membrane could be seen at the level of the opening, nor was any rejected.

A few hours later it became necessary to remove the canula, cut an oval piece from the trachea, and replace the tube, during which proceeding artificial respiration had to be maintained.

The neck was surrounded by gauze. Nutritious enemata, with small doses of laudanum, were given every two or three hours. Cream and brandy were given by mouth, and the attempt was made to give quinia and iron, but the child absolutely refused to take it. The breathing became somewhat easier, but at 9 A.M., December 26th, it was 66, and the pulse 160.

The internal tube was frequently removed and cleansed of very thick viscid mucus, which rapidly collected in it; and the other treatment was continued. In the afternoon it became evident that the internal tube was entirely too small, and it was therefore abandoned and the external one alone retained. Warm lime-water was now atomized down the tube every two hours, and on the first occasion of its use was followed by the rejection of a large piece of thick, dark-gray, glue-like false membrane. This was followed by marked relief of the dyspnœa.

The child was kept gently under the influence of opium; and was nourished by enemata of beef tea, $\text{f}\overline{3}\text{ij}$; brandy, $\text{f}\overline{3}\text{j}$; tr. opii, gtt. iv, given every three hours; which were retained unless they provoked a fecal discharge, which happened two or three times. The urine was passed freely, and was not albuminous.

The respirations were conducted solely through the tube and once during sleep fell as low as 32.

December 27th.—Still refused to eat, and the bowel also became somewhat irritable, so that several of the enemata were rejected. The respirations varied from 35 to 48; the pulse from 132 to 140. The treatment was continued; the atomization of lime-water through the tube being repeated every three hours. Towards the close of each interval the face became flushed, and the child grew restless, throwing the arms about excitedly, at times leaping up in bed, and turning around so as to lean forward on the pillows and bury his face in his hand, or else looking around with an appealing expression. The atomization was always followed by cough, and the rejection of pieces of false membrane and thick puriform matter. Towards evening he began to swallow some food.

December 28th.—The tissues of the neck had become so much infiltrated and swollen, that the canula was no longer long enough to reach from the cutaneous surface into the trachea; it was in this way pushed forward till it obstructed the tracheal opening and caused great dyspnœa. It was consequently removed, and the child, though much exhausted, sank into a gentle refreshing sleep, with quiet regular breathing. The tube was not replaced, the breathing being readily performed through the wound. There was still marked indisposition to take food, and for a few times he was forced to swallow by holding his nose and pouring beef tea down his throat; this was not, however, continued, as the effort exhausted him very much, owing to his most violent resistance. The discharge from the trachea through the wound was quite fluid, purulent, and very fetid. A solution of carbolic acid, gtt. x, in Oss. of tepid water, was

atomized through the wound; and the atmosphere of the room was kept impregnated by atomizing a stronger solution about the chamber.

During the day he swallowed more food; gr. $\frac{1}{4}$ of opium was given twice; his circulation and respiration improved.

December 29th.—Condition still improving. Respiration 28, quite full and deep, without râles; pulse had fallen steadily from 114 to 84, and was more full and strong. The color of surface was better. Respiration carried on partly through the mouth.

December 30th.—The child passed a very comfortable day. The respirations were about 24: the pulse 78 to 86, soft, full, and strong; the capillary circulation good. Food was taken much better, the child eating a croquette made soft with cream, the soft part of several oysters, a small piece of breast of partridge cut fine and rubbed up with butter and salt, and drinking sherry wine and water, and rich chocolate. The discharge had lost to a great extent its offensive character. Took gr. ss. of opium at night, and slept five hours quietly.

December 31st.—The child's condition was better in every way. The wound was contracting, the edges of the tracheal opening white and clean, and granulations beginning to project over it. The cough was stronger and more laryngeal, and the voice stronger and clearer, though still whispering. A good deal of the discharge was now raised into the mouth and expectorated. The gauze with which the wound had been covered was changed for a piece of patent lint, to encourage the larynx to gradually resume its functions.

From this time the case steadily improved. The matter expectorated grew more and more mucoid, thin, whitish, and scanty, and finally expectoration ceased almost entirely. The general symptoms rapidly improved, though he remained weak and nervous for six weeks. The external wound was covered with patent lint, at first of one, then of several thicknesses, and he gradually regained the power of breathing through the larynx, and of speaking. The larynx seemed quite clear after January 2d, 1869, eight days after the operation. The wound granulated from the bottom outwards, and was entirely cicatrized by the end of six weeks; by which time he was about the house, and had returned to his studies to occupy his mind, as he was very fretful and nervous.

November 1st, 1869.—F. W. remains perfectly well, and is indeed enjoying more robust health than for several years before this attack of membranous croup.

Angina with membranous patches on tonsils; Membranous Laryngitis; Tracheotomy on tenth day; Death on thirteenth day (fifty-eight hours after operation). Autopsy.—False membrane extending from tracheal wound to third division of bronchi; right lung emphysematous; left lung collapsed; blood dark and fluid.

CASE 2.—K. B., girl, æt. 6 years and 1 month; rather tall for her age. Her parents are healthy; but she herself had suffered much from spasmodic asthma during infancy and first dentition. Since then she has enjoyed good health. On the morning of Tuesday, October 5th, 1869, she appeared unwell with a little croupy cough, which passed off in the middle of the day, and she was allowed to play in the square for a couple of hours. On Friday, 8th, her cough was worse, but still she seemed so bright that she was allowed to play about the room; but in the afternoon she complained of sore throat, and Dr. J. F. Meigs was called and found small patches of exudation on the tonsils.

R. Potass. Chlorat., gr. ij.
Tr. Ferri Chloridi, gtt. v.
Every third hour in a teaspoonful of syrup and water.

During the night, violent dyspnoea, with noisy gasping breathing, came on, for which emetics were employed with some relief.

On Saturday, 9th, there were patches of membranous exudation on the fauces and tonsils. The cervical lymphatics were only slightly enlarged. The breathing was

difficult and stridulous; the voice feeble, small, usually whispering, but when raised by an effort was rather piping and shrill; the cough was short and smothered. There was no coryza. Treatment continued, and inhalations of the vapor from slaking lime ordered every hour.

On Sunday, 10th.—The child was restless, with at times jactitation; face flushed, and expression anxious; respiration labored; inspiration imperfect, with shrill stridor; expiration prolonged and stridulous. No expectoration. Membrane still visible in fauces. No albumen present in urine. Pulse frequent, skin hot and moist. Treatment continued, and cloths wrung out from hot water applied to the throat.

On Monday, 11th.—Condition about the same; the degree of dyspnoea varying from time to time with degree of spasm, but the breathing still continuously labored and stridulous. Treatment continued, and frictions with turpentine liniment directed to be made to the throat. Marked unwillingness to eat.

Tuesday, 12th.—There was marked improvement in the child's condition. The breathing was easier and less stridulous; the cough less frequent and looser, with a few thick yellowish purulent sputa; the voice was raised with less difficulty, and was clearer and stronger. There was, however, the same faucial pain and obstinate indisposition to eat. The fauces were still red and swollen, and a small thin patch of exudation was visible on one of the tonsils. The treatment was continued, and the child also took a little port wine and water and beef tea, and had nutritious enemata of egg given every four hours.

Wednesday, 13th.—The condition of the fauces was better, the breathing easier, and the voice more clear. The skin was still heated, pulse frequent, and there was still indisposition to eat.

During the ensuing night the breathing again became more oppressed and tighter, with some return of stridor. The voice also became suppressed and whispering. The circulation was somewhat obstructed, the face becoming flushed, and the lips rather dark.

Thursday, 14th.—These symptoms were aggravated, and in the evening there was marked jactitation and restlessness. The respirations were 40 in the minute, and stridulous, with prolonged expiration; and, during the inspiratory effort, with violent action with the external respiratory muscles, elevation of the shoulders, and recession of the base of the chest and of the epigastrium. There was also complaint of pain at the epigastrium. The cough was infrequent, short, smothered, and muffled. The eyes were anxious, staring and prominent, with large pupils. The pulse was frequent, 140, and small.

During the night there was a steady aggravation of all these symptoms. The respirations rose to 46, and became extremely obstructed, the recession of the base of the chest and at the epigastrium being unusually marked during inspiration. The voice was whispering and almost suppressed; the expression strained, appealing, and anxious; the face deeply flushed and the lips livid. There was the same complaint of constant pain at the epigastrium.

Friday, 15th, at 7½ A.M., respirations 36, pulse 136. Tracheotomy was performed by Dr. H. Lenox Hodge, the trachea being opened just below the isthmus of the thyroid gland, and a small oval piece excised from its walls. A good deal of venous hemorrhage occurred during the operation, but stopped immediately after the trachea was opened, and the tube adjusted.

No anæsthetic was used, but the child made no resistance, and evidently was slightly benumbed from asphyxia. Soon after the operation the respirations grew more easy, a large piece of false membrane was thrown off through the opening, the flush disappeared from the face, and the features became composed and placid.

Very soon after the operation the respirations fell to 28, and throughout the day remained easy and regular. The pulse fell to about 120. The child slept well, but would eat but little, and still had enemata of beef tea, f 3j, q. t. h., given it. The air of the room was kept pure, but warm and moist. The canula was covered with folds of gauze moistened with lime-water, and the wound was covered with a piece of

greased linen, so as to protect it from the canula. The inner tube was removed every hour, cleaned, anointed with glycerin, and returned. Warm lime-water was atomized through the tube every three hours, and always produced strong coughing, with the expectoration of thick purulent matter, and occasionally of flakes of tough white false membrane. All medication was suspended, save the administration of gtt. ij or iij of Tr. Opii Deodorata sufficiently often to keep the child gently under its influence. During the ensuing night the internal tube was removed, owing to the difficulty in expelling the thick viscid mucus.

Saturday, 16th.—The respirations were 24; pulse 116. During the day the child took more beef-tea and wine and water, but still had nutritious enemata given. There was great thirst, and she still complained of pain in swallowing. There was no coryza, and very slight, if any, enlargement of the cervical lymphatics.

Towards evening, breathing again became obstructed, evidently from accumulation of mucus or pseudo-membrane below the end of the tube, which was consequently removed. Its removal was followed by the discharge of several large pieces of false membrane through the wound, the edges of which were well consolidated by lymph. The breathing quickly became noiseless, easy, and tranquil again. During the following night the child enjoyed some refreshing sleep, and took more nourishment. The atomization of lime-water through the tracheal opening was repeated about every two hours, and with such great relief that she several times asked for it herself by signs, as it each time provoked cough, and caused the expulsion of thick purulent matter, dried mucus, and shreds and flakes of false membrane. Urine was discharged freely, and contained no albumen.

Sunday, 17th.—In the morning she appeared quite comfortable. The voice was still whispering, but the cough seemed looser, and she expelled purulent matter more freely through the opening; no false membrane was discharged. The thirst was still great, but the child took beef-tea more freely. The bowels had for several days been opened two or three times daily. Respirations 20–25; quite full, without recession of the base of the chest. Pulse 130–136, of rather better volume. Hands fairly warm, though at times there was a little tendency to coolness. The wound was evidently contracting. Slight emphysema of the base of the neck, which caused complaints of pain about the neck and shoulders.

At 4 P.M. it was observed that the breathing was again becoming obstructed, and that there was recession of the base of the thorax during inspiration. Lime-water was freely atomized through the opening, but without causing any discharge of membrane. The difficulty of respiration increased until 5½ P.M., when suddenly symptoms of asphyxia appeared. Prolonged efforts at artificial respiration were made, and life was thus maintained for a short time, but no essential relief was afforded, and death soon followed, on the thirteenth day of the disease and fifty-eight hours after the operation.

The chest was frequently auscultated throughout the course of the case. Before the operation it was impossible to isolate any respiratory murmur, owing to the loud snoring, whistling, and cooing tracheal and bronchial râles. After the operation, and still more after the final removal of the canula, a faint respiratory murmur could be detected, mingled with the above râles. On the morning before death only was there an obscure flapping sound transmitted to the ear with the tracheal and bronchial râles, but even then it was indeterminate in character.

During the efforts at artificial respiration, a long tubular false membrane was ejected. It had evidently been the immediate cause of death.

AUTOPSY, twenty-four hours after death. *Brain* not examined.

Thorax and Air-Passages.—The wound in the neck looked well, without pseudo-membranous exudation. The larynx itself could not be examined. A long false membrane, extending from the tracheal opening down through the right bronchus to the third or fourth division, lay loose in the trachea, having been detached from the mucous membrane. In the bronchi it was still slightly attached, but separated readily on traction. It was firm, very tough, and white, and in the upper part of the

trachea, at least one line thick. Below the bifurcation it was tubular for the rest of its course; and in its terminal portions grew softer and more yellowish. There was also, in the trachea, a large patch ($1\frac{1}{2}$ inches long by $\frac{1}{2}$ inch wide) of false membrane, of dull white color, and tightly adherent. Upon raising it, numerous little delicate fibrous prolongations were seen attaching it to the mucous membrane. Beneath this patch the mucous membrane was deeply reddened, dry, excoriated-looking, and slightly roughened by minute elevations. There was no enlargement of the mucous follicles. The vascularity of the mucous membrane diminished in the lower part of the trachea, and was but slightly marked in the secondary divisions of the bronchi. No ulceration was seen at any point. There was no pseudo-membrane in the left bronchus or any of its branches, and the mucous membrane here was less reddened than on the right side. In all probability the false membrane removed immediately after death had come from the left bronchus. The right lung was largely distended, the posterior border dark and congested, but the rest of the organ pale and emphysematous. The left lung was dark, purplish, non-crepitant, collapsed, and yielded on section an abundant flow of dark, airless, bloody serum. No pleurisy or pleural effusion.

Heart.—The left ventricle was very firmly contracted and empty and the tissue of its walls hard, tough, and florid red. The walls of the right ventricle were relaxed, and the cavity filled with fluid dark blood, without any clots. No excess of pericardial effusion.

The liver and kidneys were gorged with dark blood.

CHAPTER II.

DISEASES OF THE LUNGS AND PLEURA.

GENERAL REMARKS.

It would be difficult, perhaps, to overestimate the importance to the medical practitioner of a thorough knowledge of the different diseases of the lungs and pleura, as they occur in children. The diseases of the respiratory organs—and much the most frequent of them are pneumonia and bronchitis—cause, according to West, very nearly one-third of all the deaths under five years of age in England; while not above one child in four dies under that age from disease of the nervous system, and not above one in seven from those of the digestive system. In this country, it would seem, from the bills of mortality, that a larger proportion of children die of diseases of the digestive than of the respiratory system. But, while this is true, there can be no doubt that the diseases of the latter system are deserving of our utmost attention, since not only are they of constant occurrence and of fatal tendency, as idiopathic affections, but since, also, they frequently appear as complications in the course of other diseases, adding greatly thereby to their severity and danger. In measles, for instance, by far the most frequent cause of danger is the occurrence of some inflammation of the lungs or pleura. In scarlatina and typhoid fever, bronchitis and pneumonia are very common accidents, and recent researches have shown that in whooping-cough, and in all states of great debility and prostration, a certain change in the condition of the pulmonary tissue, to which the term collapse has been applied, is very apt to occur.

The morbid condition of the lung last referred to, that of collapse, is one that has been well understood only within a few years past, and yet it is so important, in a practical point of view, as to excite a feeling of surprise that it had not been discovered before.

ARTICLE I.

ATELECTASIS PULMONUM, OR IMPERFECT EXPANSION OF THE LUNG.

THE title of atelectasis pulmonum, from *ατελής*, imperfect, and *εχτασις*, expansion, was first employed by Dr. Edward Jörg, to designate a condition of the lungs observed by him in new-born children, a condition in which larger or smaller portions of those organs had never been penetrated by air. The respiration of the infant had, in such cases, been only imperfectly established at birth, and some parts of the pulmonary tissue had consequently never undergone expansion under the distending influence of the inspiratory act; these undilated parts continued in the foetal state.

In addition to this *congenital* form of imperfect expansion of the lung-tissue, this condition is met with at all ages of life, though with especial frequency in young children, as the consequence of a *collapse* of portions of the once-expanded lung, or in other words, of their return to the foetal or unexpanded state. To this latter form of imperfect expansion, the terms post-natal atelectasis, collapse, and foetal condition have been given. Before the discovery of its real nature was made, it had often been described also under the well-known names of carnification and lobular pneumonia. We shall designate it by the title of collapse or post-natal atelectasis, while under that of congenital atelectasis pulmonum, we shall describe the congenital variety of imperfect expansion.

CONGENITAL ATELECTASIS.

ANATOMICAL APPEARANCES.—In congenital atelectasis the parts of the lung most frequently affected are the posterior portion and lower edge of the inferior lobes, the middle lobe of the right lung, and the lower edge of the upper lobes. In some instances that we have examined, the greater part of the lower lobes of both lungs, whilst in others, still larger portions of these organs have been found to present this condition. The imperfectly expanded portions of the lungs are of a dark-red or purplish color, and are diminished in size, so as to be depressed below the level of the healthy parts. They are solid to the touch, and yet they have not lost their cohesive properties, as they are neither friable, easily torn, nor readily penetrable by the finger; their cut surface is perfectly smooth; they do not crepitate under the finger, and no air-bubbles are seen in the fluid squeezed out by pressure; they sink when thrown into water. They, in fact, resemble exactly the foetal lung. The most convincing proof of the real nature of this condition is obtained by the inflation of the lung. When this is done, the depressed, hard, and dark-

colored portions—unless the subject from whom the specimen has been taken may have lived long enough to have allowed the different tissues of the lung to become adherent—rise to their natural level, become elastic, soft, and crepitating, and change, under the influence of the entering air, from a dark and livid tint, to a rosy or pink color of healthy pulmonary tissue. In recent cases, this inflation is performed with great ease and with perfect success; while in other instances, in which the child has lived for some weeks or months, the distension is either effected only by strong efforts, or in a very imperfect manner, or it may fail entirely, owing to some permanent change having taken place in the tissues of the unexpanded portions. In a case that occurred to ourselves, the subject of which died, at the age of fourteen months, of acute pleurisy of the right side, after having presented, at birth and throughout its short life, many of the symptoms of atelectasis, the inferior two-thirds of the lower lobe of the left lung exhibited in the greatest perfection all the atelectasial characteristics. The whole of the unexpanded part was distended by means of inflation with a blowpipe, but only after repeated and powerful expiratory efforts; and Dr. E. Wallace, who made the examination, assured us that he was obliged to use a degree of force much greater than he ever employed to inflate healthy adult lungs.

In some cases there are found small patches of vesicular emphysema associated with the areas of pulmonary collapse. If, in consequence of commencing post-mortem decomposition, there has been any development of gas in the tissue of the lungs, it is seen, by the aid of a lens, in the form of irregular air-bubbles scattered through the interstitial tissue, which are easily distinguished from the minute shining air-bubbles, crowded together in regular arrangement, which are seen in lungs which have been inflated.—Bouchut, *Jour. f. Kinderkrankheiten*, 1863, 3-4, p. 263.

In most cases the foramen ovale and the ductus arteriosus are found to be still open, or the latter has but partially closed.

The *causes* of congenital atelectasis have not been satisfactorily ascertained. The conditions that are probably the most frequent causes are: original debility of the infant, from any cause that has interfered with its proper development in utero, as feeble health on the part of the mother during pregnancy, or multiple pregnancy; and acquired debility, brought about by the fact of the infant's being exposed at birth to unfavorable hygienic influences, and particularly to those which interfere with the proper performance of the respiratory act, as cold, a vitiated and close atmosphere, and the use of too heavy or tight clothing. A very hurried and rapid labor has been thought to cause, in some instances, this imperfect expansion of the lung-substance. In a case that occurred to one of us (see *Am. Jour. Med. Sc.*, Jan., 1852, p. 83), the only explanation of the condition which seemed at all plausible was that the placenta had been separated from the uterus at too early a period of the labor, in consequence of the violent and rapid character of the latter, so that the child was for a short time before birth cut off entirely from its connection with the mother—a time sufficient so to lower its vital forces as to bring on a condition resembling syncope, and to deprive it of the muscular strength

necessary on entering the world, to produce a full expansion of the thoracic cavity, and so, of course, to effect a dilatation of all parts of the lungs.

In addition to this, congenital collapse of the lungs may result from the air-passages of the child becoming obstructed with mucus or fluid in consequence of the umbilical cord being ruptured during labor, and an inspiration thus becoming necessary, before the head is free from the liquor amnii or the secretions of the mother's passages. Finally the want of expansion has been in some cases found to be dependent on pressure upon the medulla oblongata, implicating the roots of the pneumogastric nerves, resulting from inflammatory exudation or from effusion of blood owing to injuries incurred during delivery.

SYMPTOMS.—The symptoms depending on congenital atelectasis vary a good deal in different cases. There are some, however, which exist in most instances. These are the following: the child comes into the world feeble and weak, and instead of crying vigorously and loudly the moment or very soon after it is born, it fails to cry at all, or the cry is low and weak, or it is whimpering or wailing; the color, instead of being brick-red or dark-red, is pale and whitish, leaden, or livid; the muscular movements, which in healthy children are strong and vigorous, are in these languid and slow, or there are none or scarcely any, the limbs being relaxed and motionless. If the breathing is observed, it is found to be short, and imperfect, and it is evident that the thorax is but imperfectly dilated at each movement of respiration. When these symptoms exist in a very marked degree, the infant either dies soon in a state of asphyxia, or the muscular force slowly increasing, the respiration gradually improves, and the child is, after a longer or shorter time, either out of danger, or it falls into the same state as that of one in whom the symptoms have been from the first less severe. Under the latter circumstances, the infant continues feeble and weak. It breathes shortly, rapidly, and imperfectly, but often without any appearance of effort. The cry is rare, and when heard is low and feeble, or there is with each respiration a constant plaintive moan, which is very characteristic, and strongly expressive of exhaustion. The color continues pale and whitish, or it is bluish, and the temperature of the extremities is lower than natural. The child sleeps the greater part of the time, and is unable to nurse, or nurses very feebly, but can swallow when fluid is poured into the mouth. In such cases as these, the infant does not necessarily die, but will often recover when properly treated. In favorable cases, the symptoms just enumerated may last from a few hours to a day or two, or even a few weeks, without much change; then, under the influence of correct hygienic and medical treatment, they will often begin to improve. The color becomes less pale or less bluish; the muscular movements are somewhat stronger; the child begins to cry, and in a louder tone; the act of swallowing is easier and more perfect, or the infant is able to suck when applied to the breast, at first feebly and only for a moment, and then more strongly; the respiration becomes slower, fuller, and more natural, and gradually the dangerous symptoms disappear.

In unfavorable cases, on the contrary, the respiration fails to improve,

but becomes more and more short, quick, and imperfect; the temperature of the body falls; the color of the surface changes, becoming leaden, bluish, or even livid, the change showing itself first in the neighborhood of the mouth, and in the hands and feet, and extending gradually to the rest of the body; the difficulty in swallowing becomes greater, and very generally some spasmodic twitchings begin to show themselves about the muscles of the face. The respiration is very often attended with slight wheezing or rattling, and the convulsive movements returning frequently, and becoming more violent and more general, the child dies in convulsions, or it sinks very slowly and gradually, without convulsions, as though in a state of syncope. According to Steffen (*Klinik, d. Kinderkr.*, 1865, 1 Bd., p. 50), thrombosis of the cerebral sinuses has been found after death under such conditions.

There is another symptom of imperfect expansion of the lungs in newborn and very young infants, which ought not to be passed unnoticed. It is one mentioned by Dr. George A. Rees, of London, in an essay on this subject (London, 1850), and is of much diagnostic value, although not a pathognomonic symptom of this condition, as regarded by him. It is an altered movement of the ribs in respiration. During the inspiratory effort the ribs are seen to move inwards towards the mesial line of the trunk, instead of outwards as in ordinary respiration, thus diminishing instead of increasing the transverse diameter of the thorax. The explanation of the altered movement is as follows: when the diaphragm descends, the lung ought to expand in such a way as to fill up the increased space produced in the thoracic cavity by the descent of that great muscle. Instead of this being the case, however, the lung is collapsed and inexpansive, and cannot enlarge sufficiently to fill up the space alluded to, so that there would remain a vacuum in the chest were it not that the thoracic walls are driven inwards by the pressure of the atmosphere upon their outer surface. In a case that we saw ourselves in a child fourteen months old, who had presented symptoms of atelectasis from birth, and in whom we found after death very extensive collapse, this symptom was very marked. The base of the thorax was indented on both sides by a deep gutter or depression, which remained depressed and unchanged during the inspiratory movements, or which, indeed, rather became more distinctly visible during those motions, so that the chest presented the curious spectacle of dilatation or expansion in its upper parts, during inspiration, and of contraction or collapse at its base.

In regard to this interesting sign the reader is referred to our article on rickets, where is mentioned the explanation given of it by Jenner, in connection with the latter disease.

SYMPTOMS OF COLLAPSE IN THE EARLY WEEKS OF LIFE.—Before taking up the regular consideration of post-natal collapse, as it occurs at all ages of childhood, we wish to refer, for a moment, to that condition as it appears in the first few weeks of life, in infants who have exhibited no sign of it whatever, perhaps, at the moment of birth. We desire to do this now, because the symptoms to which it gives rise resemble much more those of congenital atelectasis, than those of collapse in children over a few

months old. And let it be remarked that these symptoms are very different and much more severe and threatening than those of collapse at later periods. They are in fact those of cyanosis, and in some instance are as strongly marked as those observed in the worst cases of that condition, caused by malformation of the heart or great vessels. The cyanosis and other symptoms of disordered circulation evidently depend on the obstacle offered by the collapsed and condensed portions of lung-tissue to the discharge of blood from the right side of the heart. Though this obstacle to the venous circulation is doubtless the chief cause of the symptoms in these cases, we cannot but think ourselves, that the great difference between the symptoms of congenital atelectasis, as well as of post-natal atelectasis occurring in the first few weeks of life, and the collapse of later periods, must be explained in part, at least, by the fact that the fœtal openings, the foramen ovale and ductus arteriosus, and especially the former, are still patulous, or in such a condition that they may be reopened under pressure, and so allow a portion of the contents of the overloaded and congested right side of the heart to pass into the left auricle, thence into the left ventricle and aorta, and so to the whole body.

In this form of atelectasis, the child may have been born perfectly healthy, or only weaker than usual, or it may have had some difficulty in establishing the respiration, which, however, has afterwards been effected in the most complete manner. Some days, or even weeks after birth, from a cause disturbing the function of respiration, portions of the lung may collapse, and give rise to the different symptoms of that condition in the manner above described. The most important of these symptoms are difficulty of breathing, consisting either in an increased or diminished rate of that function, diminution of the muscular power, cyanotic hue of the skin, and slight or severe spasmodic phenomena. In a case that occurred to one of ourselves (see *Am. Journ. Med. Sc., loc. cit.*):

A child, who had exhibited at birth, and for five days after, every appearance of fine health, was observed on the sixth day to cry rather violently in the morning. At one o'clock in the day he began to moan, and appeared distressed; at two he ceased to moan, became bluish, and seemed to lose his breath. He was placed in a bath, in which the blueness passed off, but the breathing continued irregular and uneven. He soon became blue again, and breathed slowly and irregularly, but had no spasm. At about four o'clock another paroxysm occurred, in which the whole surface became first bluish, and then dark, while at the same time, the trunk and limbs became stiff and rigid under the influence of tonic muscular spasm, and the respiration was slow and imperfect. After the attack had lasted for some moments, the blueness and spasmodic phenomena disappeared, but the child remained in a state of stupefaction. There were two slight paroxysms of convulsive stiffening between this and evening, and later in the evening there was still some blueness, with irregular and short respiration. During the night the breathing was short and uneven, and attended with moaning, but on the following day the symptoms had disappeared entirely, and there was no return.

In another case the symptoms of collapse did not appear until the twenty-fifth day after birth. The infant had been hearty and strong at birth, and had established its respiration fully and completely. Between the birth, however, and the time of the attack, circumstances connected with the lactation had caused the development of diarrhœa and thrush, which had debilitated the child a good deal. On the day of

the attack, frequent sneezing, with stuffing of the head, and some cough, seemed to show the existence of catarrh, and on the same day the child was unfortunately exposed, owing to the accidental opening of one of the gas-burners, to the inhalation of some gas. Late in the evening, a slight whistling or stridulous sound was heard in the breathing, the skin became suddenly a little bluish, and a slight convulsion followed. During the night there were frequent and strong convulsive seizures, always preceded and followed by deep blueness of the mouth, hands, and feet, and it was noticed that the least disturbance, as lifting or nursing, or changing the position, always brought them on. The next morning the attacks continued, but with diminished violence, under the effects of treatment, and they ceased after the middle of the day. The color of the skin had now changed; it had become rosy red, instead of pale or blue, and the hands and feet, which had been cold, were now warm and natural. There was no return after this.

In a third case, a female infant, who had been perfectly well at birth and up to the moment of this attack, was put suddenly into a bath by the nurse on the eighth day, directly after its waking from sleep. The child, who was not thoroughly waked up, seemed greatly terrified, and began to scream most violently. Instead of removing the infant from the water, the nurse persisted in holding it immersed for some minutes, when it became deeply blue, and partially convulsed; it frothed at the mouth and nose, seemed to be suffocating for breath, and appeared to be dying. These symptoms continued for three-quarters of an hour, when they gradually passed away, and it fell into a heavy sleep. When we saw the infant, soon after this, the only signs of disorder that remained consisted of an unusual paleness, drowsiness, and an expression of feebleness. Some three hours later it waked, nursed, and from that time seemed quite well.

In a fourth case, a child born apparently well, with the exception of its having had a rather frequent respiration, and who nursed very well on the second and third day, was attacked on the fourth day with blueness, moaning, short and panting respiration, and then with slight convulsive symptoms. It was unable to nurse, and though kept perfectly still, and fed from time to time with small quantities of milk and brandy, became gradually more deeply blue, had paroxysms of very slow respiration and circulation, with general convulsive seizures, and died at the end of twenty-four hours.

In a fifth case the symptoms, occasioned by a sudden attack of collapse, resembled so closely those of narcotism, as to lead two physicians to suspect that the nurse had given too large a quantity of opium. A female infant, in perfect health at birth, and nursed by the mother, continued well to the end of the fourth week. The child had then an attack of colic, and cried violently and obstinately for the greater part of three days. The nurse had given some small doses of Dewees's carminative. Towards the end of the third day of the crying, the baby was put into its usual warm bath. Directly after the bath, the child was observed to be curiously pale and white, as though it were fainting. Soon after this the color became livid, the respiration was hurried and distressed, and the child seemed to be dying. A physician was called in from the street, and found the child pulseless and very ill. When we arrived, the infant was pale, not blue, the pulse was frequent and feeble, the breathing short and weak, the pupils contracted to a very small size, and the eyes motionless. At first we thought that the child was narcotized. We had it placed gently at the breast, and the nipple inserted into the mouth, but in a moment the head was thrown forcibly backwards, and the body stiffened by a convulsive movement. The face was distorted, the breathing became labored and irregular, and the whole body assumed a livid color. The child was removed from the breast, and laid upon its right side, upon an inclined plane of pillows. Soon the dark color passed away, and the respiration became easy, though still short and feeble. There was no sign of coryza, no faucial rattle, and no evidence of bronchitis. A few drops of brandy, in a teaspoonful of breast milk, were inserted into the mouth from a teaspoon. When this was done, the head was again drawn backwards, the face was contorted, and a slight convulsive movement was produced. A wetted rag was now laid gently between the lips, and the attendants told not to

disturb the infant at all. In a few hours the symptoms subsided. When the accoucheur arrived, he also suspected that the child was narcotized. A nipple shield, with a tube and mouthpiece, was applied over the nipple, and the mouthpiece inserted into the mouth. Through this the child suckled safely for a few moments at a time. On the following day the child was much better, and able to nurse freely through the tube, without being disturbed from the position on the right side on the pillows. In three days the baby was quite well, and is now (1881) over four years old, and perfectly well. We believe that we have seen a case of sudden collapse of a larger portion of the lungs, with extensive and almost fatal obstruction of the right side of the heart, determined by prolonged and violent screaming. The whole amount of laudanum taken amounted to only a drop and a half, given in three doses, distributed over a period of twenty-four hours. For full details of this case, the reader is referred to the journal in which it was published.

("Case of Collapse of the Lung and Cyanosis in a Young Infant, produced by Violent Crying, in which the Symptoms were such as to cause a Suspicion of Opium-poisoning, with Remarks on the Nature and Treatment of Temporary Cyanosis from Post-natal Collapse of the Lung," by J. Forsyth Meigs, M. D., *American Journal of Obstetrics*, and *Diseases of Women and Children*, vol. xii, No. 1, January, 1879, New York.)

DIAGNOSIS.—There can be no difficulty in detecting the nature of the case when the imperfect expansion exists from birth, and when the physician is present at that event.

When, however, collapse of the lung-tissue continues after birth, and the physician is called upon to determine, at the age of some days, weeks, or even months, the cause of the feeble health and puny growth of the child, or to explain those sudden attacks of collapse in very young infants who had previously well established, to all appearances, the respiration, the diagnosis becomes more difficult. In the former class of cases, attention to the following points will usually, however, enable us to make a correct diagnosis. The previous history is particularly important, since, in all such cases, it will be found that the infant was either stillborn and resuscitated with more or less difficulty, or that it was born weak and feeble, and that the respiration had not been established as thoroughly and completely as it ought to have been. Dr. Rees states that certainly half of the cases of this form, in his own practice, occurred in twins, and that they were all born in a more or less completely asphyxiated condition. The present symptoms are also very important. The feeble appearance of the child and its puny growth, in connection with its past history, and the absence, as ascertained by careful examination of the case, of other morbid conditions to explain the general ill-health, ought to direct the attention of the physician to the true nature of the disease; and if we add to these considerations the local thoracic symptoms, the short, rapid, and imperfect breathing, with, perhaps, the altered movement of the ribs, the indentation instead of expansion during inspiration, mentioned above, the absence of fever, and the existence of the physical signs of more or less extensive solidification of the pulmonary tissue, without those of pneumonia, there will seldom be any difficulty in forming a correct diagnosis.

The cases described under the head of collapse, in the early weeks of life, may be readily understood from the simple fact that the symptoms cannot be satisfactorily explained by referring them to any other condition

than that of collapse of portions of the lung, with impeded and deranged circulation.

PROGNOSIS.—The condition of imperfect expansion of the lungs in a new-born child does not necessarily cause it to die immediately or very soon after birth. The fate of the child will depend very much upon the cause of the atelectasis, upon its degree of innate strength and vigor, and upon the kind of hygienic conditions to which it may be consigned. When the child is well developed, and not enfeebled by any fault in the mother's health during the pregnancy, but merely by some momentary condition that has occurred during the labor, there is every reason to hope that proper hygienic and medical treatment may restore it to health. The danger is greatest in those who continue weak and feeble, in spite of the proper measures of care and treatment, for some days or weeks after birth. We have a record of ten examples of this condition in new-born children, in nine of which the symptoms persisted during a period varying between six hours and five days. Of these, seven lived, while three died in from twenty-one hours to three days.

The prognosis of the second class of cases—those in which collapse occurs suddenly a few days or weeks after birth, and after the apparently complete establishment of respiration—will vary, of course, with the violence of the symptoms. Of five cases of this kind that came under our observation, recovery took place in three in spite of the most dangerous and alarming symptoms, while in two death occurred in a period of about twenty-four hours.

In cases where the collapse of the lungs has been extensive, and has in part persisted without proving fatal, serious organic changes in the heart have been found to follow, both by F. Weber and Steffen. The long-continued obstruction to the pulmonary circulation prevents the closure of the ductus arteriosus, and subsequently causes hypertrophy with dilatation of the right side of the heart, a patulous state of the foramen ovale, and at last eccentric hypertrophy of the left auricle and ventricle. Undoubtedly, in most cases, death occurs before these changes in the heart are induced, but it is important to be aware that imperfect expansion of the lungs may thus serve to develop serious cardiac disease of a form likely to be attended with cyanosis.

When the imperfect expansion depends upon the presence of accumulations of mucus in the air-passages, well-directed efforts usually succeed in effecting the removal of the obstruction, and the establishment of free inspiration. In those cases, finally, where there is pressure upon the pneumogastric nerves near their origin, a fatal result must always follow.

TREATMENT.—The treatment of congenital atelectasis must be directed to the removal of its probable cause. If this is suspected to be obstruction of the air-passages by collections of mucus, the infant's mouth should be cleansed, and vomiting provoked by tickling the fauces. In addition to this, all the measures calculated to stimulate respiration should be employed.

When the imperfect expansion appears to depend merely on the weakness of the infant, the treatment resolves itself almost entirely into the

employment of such means as tend to invigorate the general health of the child, and to promote the activity of the respiratory act. In a recent case, one dating from birth, in which the function has always been imperfect, and in which there are present great feebleness, drowsiness, and paleness or blueness, the room in which the infant is placed should be kept up to a temperature of 70° or 75° , and the child should be abundantly covered with warm clothing. Perfect quiet, or at least very gentle motion, is very important, and when there is any disposition to deep blueness or to convulsive movements, attention to this point is essential. It is in such cases, and in those in which these symptoms come on a few days or weeks after birth, that the position recommended by the late Dr. C. D. Meigs, for the treatment of cyanosis neonatorum, was found by him so useful. This position is one upon the right side, with the head and shoulders raised at an angle of 45° . It is obtained by arranging pillows in such a way as to form a plane inclined at that angle. Upon this the infant is placed, and orders are given that it is not to be moved at all, if possible, or only with the greatest care and gentleness, for twenty-four or forty-eight hours. There can be no doubt that this position and the attendant repose have, in many cases recorded by Dr. C. D. Meigs, and in several that we have seen ourselves, been of very great use in controlling the symptoms. Its good effects in cyanosis were supposed by him to depend on the fact that the septum auricularum becomes horizontal in this position of the body, so that the blood in the right auricle must rise against gravity, in order to pass through the foramen ovale, while at the same time the valve of that opening is disposed to fall down by its own weight, and close the foramen, and is, moreover, pressed downwards by any blood that may enter the left auricle from the pulmonary veins. This explanation will apply, of course, only to those cases of atelectasis accompanied by very extensive and deep blueness or purple color of the surface, in which we may suppose that so much of the pulmonary tissue is solidified, as to produce a degree of obstruction to the passage of blood from the right side of the heart into the lungs, sufficient to overload the right ventricle and auricle, until the latter pours a portion of its contents into the left auricle, thus causing admixture of the two kinds of blood. In a large majority of the cases of atelectasis, however, this explanation of the benefit resulting from the treatment referred to, cannot be received, as there is no reason to suppose that in them the slight cyanotic symptoms present indicate anything more than the existence of a moderate degree of fulness of the right side of the heart, unattended by any escape of blood from the right into the left auricle. In such cases the position on the right side is useful, because it is the one most favorable to a full and easy performance of the respiratory and circulatory functions. It leaves the left side free and unembarrassed, so that the heart can act with the greatest possible freedom, while the partial elevation of the head and shoulders renders the movements of the chest more easy and complete than when the body is lying on a horizontal surface. The perfect quiescence which constitutes a part of the treatment is also very important, as in many recent and particularly in cyanotic cases, the symptoms are

greatly aggravated, and convulsive attacks often brought on by moving the child, especially if this be done suddenly or rudely.

Various means are also recommended for rousing the force of respiration, as by compelling the infant to cry, by frictions of the surface, by plunging the body alternately into warm and cold water, or by allowing a stream of cold water to fall on the nape of the neck with the view of exciting the respiratory nerve-centres.

Attempts may also be made to produce full inflation by gentle mouth-to-mouth respiration. The other modes of attempting to accomplish this are either (as by compressing the thorax at regular intervals) of but little value, or (as in case of Hüter's proposal to inflate the lungs after catheterization of the larynx) highly objectionable.

Perhaps the most important point of all in the treatment of this affection, especially when the symptoms tend to become persistent, is the mode of nutrition of the child. If possible, the infant should always have a good breast of milk, and if unable to suck, the milk ought to be drawn by means of a breast-pump, and given to the child in small quantities from a spoon. About two or three teaspoonfuls may be given at first every half hour or hour, and the quantity gradually increased until the child gains strength enough to be put to the breast. If breast-milk cannot be procured, cow's milk and water may be substituted, in the proportion of one part of the former to two or three of the latter. The only medicines to be given are, at first, while the child is still very young and weak, mild stimulants, of which the best, in our opinion, is fine old brandy. Of this about five drops may be given each time that the milk is taken; or, we may make use of from three to five drop doses of the aromatic spirit of hartshorn, or of proper quantities of wine-whey.

When the symptoms of congenital atelectasis tend to persist for several weeks or months, or when we first see the patient some time after birth, the chief points to be attended to in the treatment are, as before, the mode of nutrition, which ought to be by nursing, and the use of gentle stimulants and tonics. Brandy, wine, or Huxham's tincture of bark, are the best stimulants; whilst quinine, in the dose of a quarter or half a grain, three times a day; or iron in the form of Quevenne's powder, or in that of the iodide, are the best tonics.

Vogel (*Dis. of Children*, Amer. ed., 1870, p. 55) speaks very highly of the advantage derived from the cautious application of electricity to the pectoral muscles.

COLLAPSE OF THE LUNG, OR POST-NATAL ATELECTASIS.

GENERAL REMARKS.—By collapse of the lung is meant the return of that organ to its foetal or unexpanded state. It is in fact a condition of atelectasis or imperfect expansion of its vesicular structure. The terms collapse or post-natal atelectasis are employed to contradistinguish it from congenital atelectasis, the former being applied to imperfect expansion as it occurs in lung-tissue after previous expansion, and the latter, as stated in the preceding article, to the same condition as it exists in children who have never expanded certain portions of the pulmonary substance.

The true nature of collapse of the lung was never understood, and its great practical importance never appreciated, until since the year 1844, when MM. Legendre and Bailly published, in the *Archives Générales de Médecine*, their researches on the subject. Since then various observers have repeated the investigations of those gentlemen, and thrown new light upon the matter. Among the most important of the later writers on this subject, we may mention Dr. Charles West of London, MM. Hardy and Behier of Paris, Dr. W. T. Gairdner of Edinburgh, and MM. Rilliet and Barthez, in the second edition of their work.

This discovery in pathology was one of very great value, not merely because it renders our knowledge of the morbid conditions of the lungs more exact and philosophical than it ever was previously, but because it explains certain anatomical changes in the pulmonary structures, often before noticed and described, but never satisfactorily accounted for; and still more, because it points to methods of treatment much more rational and much more successful than those employed under the influence of former ideas as to the nature of the lesions alluded to. The most important result of the new views is the disclosure of the fact that several lesions met with after death, which were formerly thought to depend on inflammation of the affected tissue, are in reality the consequences of collapse or obliteration of the vesicular structure of the lung, and not of inflammation, as was at one time supposed. The lesions alluded to are those which have been hitherto described under the names of lobular pneumonia and carnification.

The peculiar character of the lesions met with in many of the supposed cases of pneumonia, had often attracted attention and been commented upon, before their real nature came to be understood. The points of difference between these alterations and those of true pneumonia were particularly noticed by MM. Denis, De La Berge, Rufz, Rilliet and Barthez, Dr. Gerhard, and Dr. West. In fact, M. Rufz, and MM. Rilliet and Barthez, both approached very near the truth in regard to these lesions, each comparing them, but the former at an earlier period than the latter, to the condition of the lung of a fœtus that has never breathed. The latter writers, in the article on pneumonia in their first edition, have described a condition of the lung which differed so much from ordinary pneumonia as to create a great difficulty in their minds as to its true nature, and to it they applied the term carnification. They were on the very verge of detecting its real character; they did in fact suggest its real character, but were so possessed with the idea that it must be the result of some inflammatory action as to neglect to pursue their own suggestion, but endeavored to explain the condition on the ground that it was "one mode of termination of pneumonia, or else chronic pneumonia." The following passage, quoted from their work (1ère ed., t. i, p. 74), will show how closely they approached the truth: "The first idea that enters the mind on examining this tissue (carnification) is, that it resembles the lung of a fœtus that has not breathed; we should feel inclined to say that the pulmonary vesicles had not yet been dilated under the influence of the thoracic expansion, and had not, therefore, admitted air into their interior;

or, rather, it would seem as though they had been obliterated by some attack of disease, perhaps inflammation, without, however, remaining engorged, and after having lost the power of dilatation."

In the second edition of their great work, MM. Rilliet and Barthez adopt, in great measure, the views of MM. Legendre and Bailly, and of Dr. Gairdner, not only in regard to carnification, but also in regard to the yet more important lesion hitherto generally called lobular pneumonia.

But it is not only the condition of the lung called carnification that has been shown to consist, not in inflammation, but in a collapse of the pulmonary tissue. A much more important consequence of the recent researches has been the discovery that a very large majority of cases of so-called lobular pneumonia, generalized lobular pneumonia, and pseudolobar pneumonia of different writers, are also the results of collapse of the lung, variously combined with bronchitic inflammation and congestion of the pulmonary tissue. The latter discovery has lessened very much the importance of pneumonia as a disease of early life, while it has augmented in the same proportion that of bronchitis, for it has shown that a very large number of cases, formerly regarded as true inflammation of the parenchyma of the lung, are, in fact, cases of bronchitis combined with collapse of the pulmonary tissue.

Now that the nature of collapse of the lung, in connection with bronchitis, and sometimes, also, with true pneumonic inflammation or congestion, has been made known, a number of symptoms occurring in the pulmonary affections of children, which formerly seemed obscure and irregular, have become easily explicable. It had been often observed that many of the supposed pneumonias of children did not present the same symptoms, pursue the same course, nor require the same treatment as the pneumonia of adults, or as some cases of the disease in children. In a great many of the supposed cases there was an unusually large amount of bronchial inflammation, the general symptoms were much less acute than was to be expected in a parenchymatous inflammation, and, what was most singular of all, the physical signs of solidification of the lungs were very variable and uncertain, there being present on one day the signs of simple bronchitis, while on the same day or the following, and over the same region of the thorax, these would be associated with or masked by the signs of induration of the lung; and again, in a day or two, the symptoms indicative of condensation might disappear, to be succeeded yet again by those of simple bronchial inflammation. The effects of treatment seemed also to point clearly to a radical difference between the lobular or bronchopneumonia of children, and the acute phlegmasial disease of adults. It was found, in fact, that depletory measures were seldom borne well in the lobular pneumonia of children, while in the pneumonia of the adult, and in some acute cases occurring in early life, which presented the same general symptoms and the same physical signs as pneumonia in the adult, antiphlogistics, as is well known, are amongst the most successful remedies that can be made use of.

ANATOMICAL LESIONS.—Collapse of the lung (post-natal) occurs in two different forms, the *diffused*, and the limited or *lobular*. The only differ-

ence between the two forms is in the number of lobules affected, and their mode of distribution. In the diffused variety, a large number of adjoining lobules collapse, and give a condensed and solid appearance to larger or smaller portions of the lung, most frequently to the edges merely of one of the lobes, but at others to the greater part or the whole of a lobe, or even the major part of a lung. In the lobular variety, on the contrary, single lobules or clusters of lobules become collapsed in different parts of a lobe or lung, and the affected portions take the form of irregular hardened patches or tumors, situated upon the surface, or disseminated through the interior of the pulmonary texture. In the former kind of collapse, the appearance of the altered portion of the lung is somewhat that of lobar pneumonia, and it is to these cases that the terms generalized lobular, pseudo-lobar, carnification, and splenization, have been applied; while in the latter kind, the isolated and distinct condensed portions have been described by the term lobular pneumonia.

The peculiar or fundamental characters of collapsed pulmonary tissue are the same in both varieties. We will mention them as succinctly as possible, and then compare them with those of pneumonia, for the reason that it is with the lesions of that disease that those of collapse have been so frequently confounded.

Collapsed lung is generally of a dark violet color, but it may be much darker in tint, and even black, when it is much engorged with blood. Its consistence is always changed; the condensation may amount merely to slight hardening, with a diminution of the crepitation, or it may be very dense with an entire absence of crepitation, in which case portions thrown into water sink rapidly. Though more or less hardened, the tissue still retains a certain degree of flaccidity and suppleness. When cut into, the surface is seen to be smooth and uniform, having somewhat the appearance of muscle, and presenting no granulations. Pressure or scraping causes the exudation of more or less semi-transparent bloody serosity. Close examination shows that the organic elements of the tissue, the vessels, bronchi, cellular tissue, etc., can still be distinctly traced. Lastly, inflation of the lung distends the condensed parts, and gives to them again, more or less completely, their natural physiological characters.

MM. Rilliet and Barthez, in their second edition, treat, at considerable length, of congestion of the lung as a very constant accompaniment, and as a very important element in the state of collapse. They regard this congestion as being connected nearly always with bronchitic inflammation, and as being not merely a passive state, but as exhibiting phenomena, in most instances, which prove it to be in some degree an active condition. They say (*op. cit.*, t. i., p. 428): "We readily acknowledge that a state of debility, prolonged dorsal decubitus, and the obstruction to the circulation thus occasioned, facilitate the production of this condition, and give to it the appearance of a simple passive congestion. But we believe that there exists, moreover (frequently, if not always), a really active and even inflammatory movement." They regard this opinion as proved chiefly by the fact that they have found the texture of the affected parts to be somewhat softened, as shown by the facility with which they

are torn by the finger or by scraping with a scalpel; by the swollen and turgid condition the tissues exhibit; by the quantity of sanguineous or sero-sanguineous liquid which escapes on pressure; and by the presence of a serous exudation around the pulmonary vesicles, while the interior of the vesicles appears to be healthy. The last-mentioned condition they found upon their own observation, and upon a microscopic examination made by M. Lebert.

The color is different in the two diseases, being in collapse purple or livid, and in pneumonia brownish-red or fallow-red. In pneumonia the pleura covering the hepatized portions is often covered with false membrane, showing thereby the inflammatory nature of the disease; in collapse this is rarely the case, and only when there is some accidental concomitant pneumonia. The density of the lung in the two conditions is of a different kind; in pneumonia it is hard to the touch, and unyielding; in collapse it always retains a certain degree of flaccidity and softness, like that of muscular tissue. In pneumonia the diseased part is turgid and swollen, so that it projects above the common level of the surrounding surface; in collapse, on the contrary, it is shrunken and depressed below the neighboring parts. In pneumonia the effect of the inflammatory process on the tissues is very strongly marked, and produces changes in them very different from those occasioned by mere collapse. In the former disease the cohesive properties of the pulmonary structure are very much lessened, so that the inflamed parts are readily penetrated by the finger, and are easily torn. In simple collapse, on the contrary, the diseased part is as firm and resisting, or even more so, than in health; whilst in collapse occurring in bronchitis and attended with congestion, though the cohesion of the tissues is somewhat lessened, it is never nearly so much so as in pneumonia. In the true hepatization of pneumonia, a cut surface always presents a granular aspect, while in collapse, on the contrary, it is smooth and even. On scraping a cut surface it is found that in the former state a plastic, fibrinous matter, of a yellowish, orange, or gray color, comes off on the knife, while in collapse only some semi-transparent bloody serosity is scraped off. In the former, the anatomical arrangement of the lobules cannot be seen, as the inflammation attacks indifferently the lobules themselves, the interlobular septa, and parts of neighboring lobules; but, in the latter, the alteration can always be seen to be more or less regularly confined to the lobules, the cellular interstices between the lobules remaining more or less apparent; so that in pneumonia the alteration is not bounded at all by the outlines of the lobules, while in collapse the alteration always affects, more or less, the lobular form. To conclude, the effects of inflation are altogether different in the two conditions. M. Legendre (*Recherches Anatom.-Path. et. Clin. sur quelques, Mal. de l' Enfance*, p. 164) states that air can never be made to penetrate by inflation into a completely hepatized lung. Neither in hepatization of the lobar form, nor in true partial hepatization, has he ever been able, even with the utmost effort, to force air into the inflamed tissues. After repeated trials, the tissue remained compact and friable, and sank as rapidly as before when thrown into water. In the foetal state,

on the contrary, the slightest effort sufficed to fill and distend the collapsed air cells, and to give to the altered portion its natural appearance, excepting that it became more red in consequence of the oxygenation of blood contained in the capillaries. Dr. Gairdner (*Pathol. Anat. of Bronchitis*, etc., Edinburgh, 1850, pp. 13, 14) remarks that, though this test "is very useful in demonstrating the nature of the lesion, in a favorable case, to one not familiar with its character, I do not believe it to be applicable to the determination of the presence or absence of pneumonia in those mixed cases in which alone there is any difficulty." He has observed, in fact, that partially pneumonic lung may be inflated when the affection is recent and combined, as it frequently is, with bronchitic collapse, while in the latter lesion, in its purest forms, complete inflation is often very difficult or impossible after the collapsed state has been of some duration.

The part of the lung in which collapse is most frequently met with depends somewhat on the form of the alteration. In the diffused variety, it may affect a more or less considerable portion of either or both lungs, but is most common at their posterior part. The lobular variety is most common on the anterior edges, but may, like the diffused, occur in any other part. As a general rule, the alteration is most frequent at the periphery of the organ, where its edges are thin, as along the margins of the lobes, in the languette of the upper lobes, and at the bases of both lungs. The parts just named are those most distant from the primary air-passages; they are those in which the inspired air would arrive last, and with the least force of impulsion.

CAUSES.—It has been generally acknowledged that there are two principal causes by which to explain the production of collapse of the lung. These are the presence in the bronchi of some condition which acts as an impediment to the ready passage of the inspired air, and a want of power in the muscular apparatus by which the function of respiration is carried on. To these Dr. Gairdner adds another,—the inability to cough and expectorate, and thus remove the obstructing mucus; but this is, in fact, included in the preceding.

The most important of the above-mentioned causes is evidently the deficient respiratory power, since this is noticed and insisted upon by all observers. It has been found, in fact, that collapse seldom occurs to any considerable extent except in children who are exhausted and debilitated. The debility may be congenital, it may be the result of wearing diseases, as diarrhœa, hooping-cough, measles, typhoid fever, etc., or it may depend on exposure to unwholesome and enfeebling hygienic conditions. It is easy to understand that a child who is either born weak and feeble, or who becomes so in after years from any of the causes just alluded to, must lose, with the general decay of the strength of the body, some portion of the muscular power by which alone a complete and efficient dilatation of the thoracic cavity can be accomplished, and that, when this is the case, the inspirations must be short and imperfect, and that portions of the lung most distant from the primary air-passages, not being reached by the inspired air, will remain in an unexpanded or collapsed state. If we add to this state of feeble respiratory power, the presence of secretions in the

air-tubes, whether these be the consequence of bronchial inflammation, as they are in the immense majority of cases, or as Dr. Gairdner suggests they may sometimes be, the mere natural secretion of these tubes, accumulated for the want of power to throw them off, it becomes very easy to comprehend the mode of production of collapse, in at least some of the examples.

Whether a simple deficiency of inspiratory force alone, without obstructing mucus in the bronchi, will give rise to collapse, is a somewhat mooted point. Dr. West agrees with MM. Legendre and Bailly, in the opinion that it is often due to the inspiratory power having been inadequate to overcome that natural elasticity of the lung which opposes a full dilatation of the organ. Dr. Gairdner (*loc. cit.*, p. 33) cannot "see reason to believe with Dr. West, that mere debility, apart from any obstruction in the tubes, is a sufficient cause for collapse in the child." He remarks, and with strong show of reason, that the very fact of the lesion being usually more or less lobular, or partial in its distribution, appears to indicate special circumstances of a local kind, as having a marked influence on the production of this affection. What is of most consequence, however, to the physician, as an important practical truth, is the fact stated by several observers, and adverted to by Dr. Gairdner himself, that in some cases no signs whatever of obstructive bronchitis or of bronchial accumulation can be discovered during life. Before leaving this point, we desire to call attention to the opinion of Hasse (*Pathol. Anat.*, Syd. Soc. ed., p. 253), that, though this partial introduction of air might be deemed at variance with the laws of respiration, inasmuch as the atmospheric pressure must necessarily distend the entire lung equally, not to the exclusion of a lobe, and still less to that of a lobule, the objection falls to the ground when it is considered that the operation of these laws is the result of previous muscular action. He refers to the fact that in pleurisy one-half of the thorax, and in partial pleurisy certain portions of that cavity, do not share at all in the movements of the remainder. "We need, therefore," he says, "be at no loss to understand how defective breathing may originate in a merely partial activity of the intercostal or other respiratory muscles."

Dr. Gairdner, as already stated, considers as one of the causes of collapse, an inability to cough and expectorate, and thus to remove the obstructing mucus. The views which he expresses on this point are very interesting, and also, we think, very important. He states that Laennec supposed the expiratory force of respiration to be weaker than the inspiratory, while in fact the experiments of Hutchinson and Mendelssohn, to which he refers, prove that though *ordinary* inspiration is more of a muscular act than *ordinary* expiration, yet the residual effective force for overcoming adventitious obstruction is very considerably greater in expiration. "The *forced* or *muscular* expiratory act is, in fact, about one-third more powerful, as measured by its effect upon a pressure-gauge, than the extreme force of inspiration; and it is this force which is thrown into action when obstruction in the tubes is to be overcome." In the act of coughing, the air in the vesicles is brought to bear upon the obstructing substance within the bronchi, at a maximum amount of outward pressure, and with the

additional mechanical advantage of a sudden impulse, so that the practical efficiency of the expiration in forcing air through obstructions must be far greater than that of inspiration. It is clear, therefore, that if the secretions in the air-passages be so abundant or so viscid as to interfere materially with the entrance and exit of air, they must necessarily occasion collapse, either partial or total, of the parts beyond them; since not only does the air enter with difficulty, but being expelled with greater force and in larger quantity than it can be drawn in, the amount remaining in the vesicular structure must gradually diminish. This effect of obstruction will be still more remarkable when the muscular force of respiration is diminished by debility of the patient, for then the inability of the inspiratory act to replace the air driven out by expiration, will be yet more marked than when the muscular powers of the body retain their full force.

There is still another mechanical condition which tends to produce collapse from obstruction, to which Dr. Gairdner refers. This condition is to be found in the form of the bronchial tubes. These tubes are a series of gradually diminishing cylinders, and if a plug of any kind, but especially one closely adapted to the shape of the cylinders, and possessing considerable tenacity, be lodged in any portion of such a cylinder, it will move with much more difficulty towards the smaller end, and in doing so will close up the tapering tube much more tightly against the passage of air, than when moved in the opposite direction into a wider space. From this arrangement of the parts, it will happen that at every expiration a portion of air will be expelled, which, in inspiration, is not restored, owing in part to the comparative weakness of the inspiratory force, and in part to the valvular action of the plug. "If cough supervene, the plug may be entirely dislodged from its position, or expectorated, the air, of course, returning freely into the obstructed part; but if the expiratory force is only sufficient to slightly displace the plug, so as to allow of the outward passage of air, the inspiration will again bring it back to its former position, and the repetition of this process must, after a time, end in perfect collapse of the portion of lung usually fed with air by the obstructed bronchus."

We have been thus particular in our consideration of the causes of collapse, because we are convinced, from personal observation, that it is a subject of very great importance in practice. Many times in the last few years, we have met with cases of bronchitis, either primary or secondary, in weak and debilitated children, in which the general and local symptoms have pointed clearly to the existence of collapse of the lung, and in which, moreover, the good effects of a sustaining and even stimulating treatment have shown the great utility of an acquaintance with the nature of this affection, and its proper remedies.

SYMPTOMS.—As collapse of the lung occurs almost always in connection with bronchitis, though sometimes, also, after, or concomitantly with pneumonia, it is clear that the symptoms which reveal its existence must be mingled, in a greater or less degree, with those of the two diseases just named. It is true, nevertheless, that it sometimes occurs unassociated with more than very slight evidences of any other disease of the lung. Cases of the latter kind have been usually observed only in children dying in

states of utter exhaustion, in whom the muscular power of respiration has been so greatly weakened as to prevent a dilatation of the thoracic cavity sufficient to carry air into the deeper parts of the lung. In such instances, the symptoms of collapse do not show themselves until a very short time before death, and they consist in the sudden appearance of very rapid and oppressed breathing, with little or no cough, in more or less extensive dulness on percussion over different parts of the chest, but most frequently the inferior dorsal regions, and in feeble or suppressed respiratory murmur, or more frequently a distant and imperfect bronchial respiration. In some cases, however, in which there is very little bronchial complication, as shown by the rarity and small amount of the catarrhal râles, the symptoms of collapse continue with more or less irregularity, as to situation and extent, for periods of several weeks, or even months. But here, also, as in the cases previously referred to, the general debility and low health of the child are strongly marked, and are, with slight variations persistent. As an instance of this kind of collapse, we may cite the following case, which occurred in our practice:

A boy, between three and four months old, who, at birth, and up to the time of this attack, had presented every appearance of strong and vigorous health, was seized, on the 3d of October, 1849, with symptoms of a somewhat irregular and anomalous character, but which we soon suspected to be the signs of an intermittent fever. We were induced in part to make this diagnosis, from the fact of having attended the mother during her gestation of this child, in a severe attack of intermittent fever. At the beginning of the sickness, there was some little coryza, but no cough whatever. On the 3d of October, after the coryza had lasted for a few days, he became worse, and we were sent for. During the following six days he had one or two attacks each day of coldness of the extremities, followed by violent fever, and ending sometimes with perspiration. He was exceedingly fretful, screamed a great deal, was at times drowsy and dull, and vomited occasionally. The stools were regular and perfectly natural. The breathing was rapid and short nearly all the time, but there was no cough whatever. On the seventh day, the respiration was 100, and irregular. The child was pale, weak, drowsy, and entirely without cough. Percussion revealed nothing, and no râles could be heard. On the eighth day, the breathing was 96, and a slight, dry cough was heard two or three times. When roused up, the intelligence of the child seemed perfect. On the ninth day, the breathing was 63, and the pulse 120. There was rather more cough, though still very little, and there was a slight return of the coryza, of which there had been none for several days before. Neither auscultation nor percussion revealed any decided change in the lungs. On the eleventh day, the paroxysms of chilliness, followed by fever, were still noticeable, though there was no clearly marked periodicity in the returns. When without fever the breathing was 54; during the fever it was 67. Auscultation revealed nothing decided. Percussion showed dulness beneath the right clavicle. By the seventeenth day, the intermittent nature of the disorder was more decidedly marked, and under a few doses of quinine the symptoms had improved, so that the breathing fell to 30 during sleep. The cough was a little more frequent, though still very slight, and it was loose. The coryza, also, was more considerable, the nasal discharge being quite abundant.

After this the case went on badly, owing, we think, in great measure, to the circumstance of the quinine being abandoned in consequence of the opposition made by the parents to its administration, an opposition which we allowed to influence us more than was proper. During November and December, the child remained weak, pale, languid, and with uncertain appetite, sometimes refusing the breast for a whole day at a time. The quinine was suspended at first on account of the great improve-

ment which had taken place in the symptoms, and though resumed afterwards, was given in such small quantities, and for so short a time, for the reasons just mentioned, as to be of no service. In December the child was very ill. It looked badly, having a pale, waxy face, and a dull, languid expression, though without any want of intelligence; it emaciated moderately, and had occasional vomiting; the stools were natural. At this time also it took the mother's breast with some difficulty, and refused artificial food altogether. Occasionally during this month there was observed a slight blueness around the mouth, and also about the hands and feet. Late in the month it was attacked with thrush in a slight degree, which lasted several days. In the first week of January, finding that it was fast sinking from refusing the mother's breast and artificial food, a wet-nurse was procured, and for a few days it seemed to improve a little, but this did not last. It grew weaker and thinner, the thrush returned, it had now a good deal of loose cough, the abdomen became somewhat contracted and felt hard and doughy, and the breathing was very rapid, though not greatly oppressed. The child died at last on the 26th of January, having for ten days before that event looked wretchedly languid and haggard, and having presented for three days before, slight diarrhoea, loose, frequent cough, entire loss of appetite, thrush, drowsiness, and finally coma.

At the autopsy there were found some fibrinous exudation, and a few adhesions over the lower half of the left lung. The lower two-thirds of the left and the lower half of the right lung were dark-colored, more dense than usual, not friable, and exhibited no granulations on a cut surface. These portions were in fact collapsed. The upper lobes were spongy, crepitant, and healthy. Not a tubercle was found. The foramen ovale presented an oval-shaped opening, of the size of a goose-quill. The abdominal organs were healthy.

When, as indeed most usually happens, collapse occurs in the course of bronchitis, it is associated of course with the symptoms of that disease. The bronchitic symptoms have lasted in their usual form for several days, having been marked by sonorous, sibilant, and subcrepitant râles, when suddenly, or in the space of a few hours, the breathing becomes much worse, the pulse rises in frequency but becomes small and feeble, and certain changes take place in the physical signs which are very important. The subcrepitant râle continues to be heard, but it is associated now with prolonged expiration, and a little later with bronchial respiration, which, however, is of a different kind from the bronchial respiration of pneumonia, being distant and smothered, instead of near and metallic, as in that disease. The percussion becomes, at the same time, dull and obscure, but scarcely ever to the same extent as in pneumonia. The general symptoms are those of exhaustion, rather than of high reaction. The surface is pale or slightly bluish, the skin is either natural in temperature, slightly warmer than usual, or cool, the strength is very much reduced, and the child appears more seriously ill, and particularly more oppressed than the amount of bronchitis present would seem to explain.

As an example of collapse occurring in the course of bronchitis, we will give the following case:

A girl between two and three months old, healthy when born and up to the time of this sickness, saving that she was rather paler and smaller than most robust infants, was seized with coryza and slight cough, and after a few days with the symptoms of a mild bronchitis. For two days there was frequent cough, some little fever, quick but not oppressed breathing, occasional sibilant and mucous râles, perfect ability to nurse, and very moderate restlessness or fretfulness. On the third day, without any apparent reason, the symptoms became suddenly very alarming. The breathing became ex-

tremely rapid and most violently oppressed, so that the movements of the chest at each respiration were heaving and laborious, the shoulders being lifted high at each respiration, the outer angles of the mouth drawn downwards, and the *alæ nasi* widely dilated. There were at the same time abundant subcrepitant, intermingled with dry râles over the dorsum of the chest. There was marked constriction of the base of the chest with each inspiration. The cough was frequent and racking, and occurred in paroxysms. The child was still and quiet, pale, had a haggard and exhausted look, was unable to nurse at all, and its surface was cool and white, especially that of the extremities. These symptoms continued with very little modification for twenty-four hours, when, under the use of brandy administered every hour in milk drawn from the mother, of the spirit of Mindererus and paregoric, perfect quiet, and the assiduous employment of mild revellents, they began to moderate, and at the end of another twenty-four hours the constriction at the base of the thorax during inspiration had disappeared, the breathing was easy and gentle, the extremities had become warm, the child nursed eagerly and abundantly, and, with the exception of a slight catarrh, which lasted a few days longer, it was well.

Collapse depending on bronchitic inflammation, in debilitated children, may sometimes last a considerable length of time. In one case, indeed, that we saw a few years since, and of which an account was published (see *Am. Jour. Med. Sci.* for January, 1852, p. 98), the symptoms, owing probably to the fact that the bronchitis causing the collapse was an accompaniment of whooping-cough, continued with slight variations in degree for a period of about three months, after which the child entirely recovered.

DIAGNOSIS.—The diagnosis of collapse of the lung must always be more or less uncertain where it is of the lobular form, for the reason that the collapsed lobules being disseminated irregularly through the pulmonary tissue, afford no physical sign by which we can detect their condition. The presence of this form ought, however, to be suspected whenever, in a chronic disease, and especially in the course of a catarrhal attack occurring in a feeble and debilitated child, the breathing becomes excessively quick and labored, the skin pale and cool, when the base of the thorax presents a depression instead of an expansion during inspiration, and especially, when these symptoms occur without there being a sufficiently severe and extensive bronchitis to explain their existence.

In cases of collapse affecting a considerable or the greater part of a lobe, the diagnosis, though still perhaps rather uncertain, is much more clear and positive than in the lobular form. In the latter form we are obliged to depend, indeed, almost exclusively upon the rational symptoms, the physical signs being either very slight or entirely null. In collapse of considerable portions of the lung-tissue, we have, on the contrary, some very useful physical signs. These are, the existence of dulness, greater or less, on percussion; feeble respiratory murmur; prolonged expiratory sound, and sometimes bronchial respiration; which, when they occur in connection with, and in the course of bronchitis, are usually quite sufficient to render the diagnosis easy.

The only diseases with which collapse of the lung, presenting the physical signs just mentioned, could be confounded, are pneumonia and pleurisy. From both of these it is usually distinguishable by the absence in collapse, or the slight severity, of the reactional symptoms, by the paleness or blue-

ness and coolness of the surface, by the absence of acute pain, by the greater severity in collapse of the bronchitic symptoms, and by the fact that it rarely occurs except in enfeebled, broken-down subjects, or in those laboring under severe bronchitis. The character of the physical signs, moreover, is different. Though we have dulness on percussion in collapse, it is not so absolute as that either of pleurisy with large effusion, or that of confirmed pneumonia. The bronchial respiration, too, is in collapse different from that of pneumonia. It is muffled and distant, instead of being clear, metallic, and close under the ear, as in pneumonia; and is heard, too, much more in the expiration than in inspiration. In collapse there is also heard, unlike either pneumonia or pleurisy, the sibilant and sonorous dry râles, and the mucous or subcrepitant râles of bronchitis. To add to these differences, it is proper to say that, in cases of pneumonia and pleurisy, the course of the disease is much more regular, and the special symptoms so well marked as to leave no doubt as to the real nature of the attack.

PROGNOSIS.—The prognosis of collapse must depend, in great measure, on two circumstances,—the amount of bronchitis which accompanies it, and the constitutional state of the child. When it occurs during the course of extensive bronchitis, as shown by a great abundance and extent of the bronchitic râles, it must add greatly to the danger of that disease; and if, at the same time, the child be weak and debilitated, either from causes long previously in action, or from the severity of the present attack, the risk to life is very great indeed. Collapse is dangerous, also, but far from necessarily fatal, in subjects in whom its chief cause has been simple debility. The possibility and the probability of recovery will depend on the hygienic conditions to which the child is exposed, the degree of vital strength it is likely to inherit from its parents, the extent of the collapse, as indicated by the severity of the thoracic symptoms, both rational and physical, and the effects of treatment. When the subject can be placed under favorable hygienic conditions, when it has inherited from its parents a good and vigorous hold on life, and when the symptoms of collapse are not very violent, a proper and rational treatment will in all probability save it, while, under opposite circumstances, the chance of recovery would be very small, if there were any.

TREATMENT.—The treatment of collapse, or post-natal atelectasis, must vary somewhat in different cases. One general rule will apply, however, to all; that is, to employ a sustaining and strengthening system of medication, to the exclusion of all exhausting means.

In cases which are entirely, or almost entirely, independent of bronchitis, the most important measures to be attended to are the regulation of the temperature in which the child is kept, of the clothing, and of the diet, the use of mild stimulants and of tonics, and the external employment of revellents. The child ought to be kept in a warm, even temperature of from 70° to 75° ; it should be clothed in soft flannels, and its diet ought to be nourishing and strengthening. If at the breast, we should be sure that the milk is of a good quality, and that the nurse has an abundant flow. If weaned recently, it ought to have, if possible, a wet-nurse, and so also

if it be supposed that the mother has too little milk, or that this is not perfectly healthy. If permanently weaned, the diet should be so arranged as to give to the child what is at the same time easy of digestion and nutritious. In a severe case, coming on suddenly, the most suitable internal remedies are brandy, in small doses, frequently repeated; quinia in full doses, or Huxham's tincture of bark, the spiritus Mindereri, the aromatic spirit of hartshorn, or carbonate of ammonia given in emulsion. In slower and more chronic cases, we must depend on a well-selected and nutritious diet (and food ought to be given almost by force, or at least it should be urged strongly on the child), on warm clothing, and on the use internally of brandy, quinine, the citrate of iron and quinine, reduced iron, the iodide of iron, Huxham's tincture of bark, or some such remedy. In sudden cases, the best revellents are the following; mustard weakened by admixture of flour or Indian meal, and applied once in three or four hours; a plaster made of suet or simple cerate, grated over with nutmeg; or liniments composed of ammonia, spirits of turpentine, or oil of amber, mixed with sweet oil. In chronic cases, the Burgundy pitch, or compound Galbanum plaster, made somewhat weaker than that used for adults, should be applied over the front and back of the chest; or we may rub the thorax twice a day with any ordinary ammonia liniment, made, if necessary, rather more irritating than usual by the addition of some oil of monarda. The daily use of a gentle emetic of ipecacuanha has been recommended, and supposed to prove useful, by emptying the bronchi of their secretions, and also by the fact that its operation induces several deep inspirations, and in that manner promotes the better performance of the respiratory act. We have never employed the emetic except in cases accompanied with a good deal of bronchitis and consequent accumulation of mucus in the air-tubes, and not then when the prostration was very great. In fact, the operation of any emetic is sometimes productive of so much exhaustion of the strength as to cause us to hesitate in prescribing a remedy of that class; though we can fully understand that the act of vomiting, if not followed by too much prostration, could scarcely fail to prove beneficial in collapse, by the strong efforts at breathing which it gives rise to, and also by the succussions it must impart to the lungs through the medium of the diaphragm.

In cases of collapse occurring in the course of, or towards the termination of severe bronchitis, the treatment must resemble a good deal that which we have just described as proper for the same condition, when it exists unassociated, or associated only to a slight extent, with that disease. When the symptoms of imperfect expansion appear towards the termination of, or after the patient has partially recovered from bronchitis, and when of course the strength is more or less reduced by the severity of the previous acute sickness, and also perhaps by the necessary measures of the treatment, the case ought to be managed very much in the same way as has just been recommended for those in which the collapse was caused chiefly by exhaustion, and less by the presence of obstructing secretions in the bronchi. Nourishing, but very light and digestible food; mild

stimulants, as small quantities of brandy or wine-whey; the bitter tinctures, iron, or quinine, with counter-irritants to the surface of the chest, warm clothing, and repose, constitute the necessary and most reasonable remedies. When, on the contrary, the atelectasial condition supervenes in the midst of extensive and severe bronchitis, we are called upon to treat at the same moment two morbid states, one consisting of active inflammation, and another of want of power in the muscles of respiration to force the atmospheric air through the secretions which are obstructing the air-passages. Under these circumstances, there is almost always associated with the bronchitis, as we shall find when we come to treat of that disease, more or less intense congestion of the collapsed portions of the lungs. We must employ, therefore, such remedies as tend to modify the inflammation of the bronchial mucous membrane, and diminish thereby the amount of secretion poured into the air-passages; such as may serve to expel mechanically those secretions; and those which shall unload the congested lung of its excess of blood, always taking care, in our selection of the agents to accomplish these ends, to choose those which are the least perturbative and exhausting. To moderate the inflammation of the bronchial mucous membrane, and with a view also to unload the congested parts of the lung, we may apply a few dry cups, or rely on counter-irritation, the best mode of effecting which is by the repeated application of mustard poultices, consisting of one-third mustard to two-thirds Indian meal or flour, and by mustard foot-baths. These poultices ought to be applied first to the dorsum and then to the front of the chest, once in every three or four hours, and they should be made large enough to cover a considerable portion of the thoracic walls. Counter-irritation, assiduously made use of, is we believe one of the most, if not the most effectual means of treatment in the case. Emetics ought to be given twice a day, or even three times, if they do not reduce the strength too much. The best are those which operate with the least subsequent prostration, such as ipecacuanha or alum. When they are found to exhaust much, and to increase thereby the labor of breathing, their use must be suspended.

After emetics, or when these cannot be used, the remedies from which we have obtained the greatest benefit are carbonate of ammonia or liq. ammon. acetat., and seneka, either in decoction or syrup, combined sometimes with small quantities of opium. To a child two years old we should give one or one and a half grains of carbonate of ammonia or twenty drops of the acetate of ammonia solution, with ten of the syrup of seneka, or with a teaspoonful of decoction of seneka, every two hours. When the cough is paroxysmal, painful, and harassing, about ten drops of paregoric, half a drop or a drop of laudanum, or from four to six drops of solution of morphia, may be added to each of the above doses. The opiate ought to be continued until the cough and restlessness diminish, and then be suspended. In all these cases, there should be no hesitation in giving small quantities of brandy or wine-whey, whenever the symptoms of prostration are so marked as to indicate danger; and these stimuli are urgently called for when the pulse is very rapid and small, when the skin is cool or pale

and bluish, and when the general aspect of the patient, and the convulsive and labored character of the breathing, show that the muscular strength of the child is scarcely sufficient to carry on the function of respiration.

ARTICLE II.

PNEUMONIA.

DEFINITION; SYNONYMS; FREQUENCY; FORMS.—The term pneumonia is now by universal consent, applied only to inflammation of the parenchymatous structure of the lungs. It is often called, in this country, catarrh-fever, lung-fever, or inflammation of the lungs.

It is one of the most frequent, and therefore, one of the most important of the acute diseases of childhood. Dr. West, in a paper on the pneumonia of children (*Brit. and For. Med. Rev.*, April 1843), informs us that the English tables of mortality show pneumonia to be the cause of a larger number of deaths in childhood than any other disease, with the exception of the exanthemata. From the third report of the Registrar-General, he quotes the facts that of all the deaths in the metropolitan districts under fifteen years of age, 13.6 per cent. were from pneumonia, 13 per cent. from convulsions, and 5.4 per cent. from hydrocephalus. He obtained nearly similar results from an examination of the returns from Manchester, Liverpool, and Birmingham.

In this city it appears from the bills of mortality that the deaths from this disease are strikingly below the percentage calculated by Dr. West. Thus, during the ten years ending with 1879, the total mortality from all causes (excluding still-born children) was, at all ages, 166,942; under fifteen years of age, 76,063; and under five years, 66,613. The total mortality from pneumonia during the same period was, at all ages, 9501, or 5.68 per cent. of the entire mortality; under fifteen years, 4140, or 5.44 per cent. of the mortality under that age; and under five years, 3795, or 5.69 per cent. of the mortality under that age. During the same series of years, the total mortality from bronchitis was, at all ages, 2556, or 1.53 per cent. of the entire mortality; under the age of fifteen years, 1774, or 2.33 per cent. of the mortality under that age; and under the age of five years, 1731, or 2.59 per cent. of the mortality during the first five years of life.

Any one who will study with attention the various doctrines in regard to pneumonia and bronchitis that have been set forth in the different works on the diseases of children, will most assuredly acknowledge that there are few diseases about which there has prevailed so much diversity of opinion as to the real nature of the lesions forming the essential anatomical characters of the disorder, and as a consequence of this, so much doubt as to the proper mode of classifying and describing them. From the time of the appearance of the works of M. Valleix, M. Barrier, Dr. Gerhard, and especially that of MM. Rilliet and Barthez, up to the mo-

ment of publication of the essay of MM. Legendre and Bailly (referred to in the article on atelectasis), it was commonly believed that inflammation of the parenchyma of the lung exhibited in children very different characters in the majority of the cases, from those which marked the pneumonia of the adult. Two principal forms of the disease were therefore described by most writers,—the *lobular* and the *lobar*. The former was supposed to be almost peculiar to children, and to occur only on rare occasions in adults; the latter was held to resemble, in almost every respect, the pulmonic inflammation of the adult. Moreover, lobular pneumonia was generally believed to be by far the most common form assumed by the inflammation in children under five years of age, whilst lobar pneumonia was thought to be comparatively rare under the age mentioned. Besides these two chief varieties of pneumonia, two others have been described under the names of *vesicular* and *marginal* pneumonia, while to yet another MM. Rilliet and Barthez applied the title of *carnification*.

The researches of MM. Legendre and Bailly, published in 1844, caused a great revolution in the views of medical observers and writers. These authors first pointed out (as stated in the article on atelectasis) that a very large proportion of the cases previously described under the titles of lobular pneumonia, generalized lobular pneumonia, pseudo-lobar pneumonia, marginal pneumonia, and the carnification of MM. Rilliet and Barthez, were in fact cases of bronchitis variously associated with congestion and collapse of the tissue of the lung. Since the publication of these views, the whole subject has been repeatedly investigated, and we believe that a general agreement now exists in regard to most of the important pathological questions connected with collapse of the lung and the different forms of pneumonia. We refer any one who desires to study the progress of medical opinion on these important subjects to the able essay, *On the Pathological Anatomy of Bronchitis and the Diseases of the Lung connected with Bronchial Obstruction*, by Dr. W. T. Gairdner, of Edinburgh, and to the elaborate article in the second edition of the work of Rilliet and Barthez. We shall ourselves adopt the division of pneumonia into the *lobar* or *croupous* and the *catarrhal* forms. The former of these is sufficiently understood as corresponding with the same form in the adult, while the latter corresponds with the form which we, in previous editions, described under the term *partial*, and which has also been named *lobular*. Despite the fact that formerly many cases of bronchitis with lobular collapse were regarded as cases of lobular pneumonia, it is probable that the catarrhal form must still be regarded as of at least equally frequent occurrence in children as the lobar. In infants under the age of two years, pneumonia is especially apt to assume the catarrhal form. Rindfleisch, however, exceeds the truth in asserting that in children under the age of five, hardly any other form of pulmonary inflammation occurs (*Pathological Histology*, Syd. Soc. ed., vol. ii, p. 14).

PREDISPOSING CAUSES.—It is generally believed that pneumonia is most apt to occur in the course of other affections. This is certainly true in regard to both forms of the disease as it prevails in hospitals, and probably

amongst the poorer classes of society also. MM. Rilliet and Barthez state that of two hundred and forty-five cases observed by themselves, only fifty-eight, or a little less than one-fourth, occurred in children previously in good health. The proportion of cases in which lobar pneumonia occurs in private practice as a secondary affection is much smaller, since of fifty-two well-marked cases, observed by ourselves, in which this point was noted, only seven were secondary. On the other hand catarrhal pneumonia is usually a secondary lesion, and frequently occurs as the result of an extension of severe bronchitis.

Age forms a strong predisposing influence. Of the two hundred and forty-five cases above quoted, one hundred and seventy-two occurred under five years of age. Dr. West (*loc. cit.*) says that during the first five years of life, the cases of pneumonia were in the proportion of 10.3 per cent. to the total of diseases, while in the succeeding five years they were in the proportion only of 1.3 per cent. The mortality bills of this city exhibit the same marked excess in the proportion of deaths from pneumonia under five years of age, as compared with the ensuing years. We have already seen that the proportion during the first five years of life is 5.69 per cent. of the entire mortality under that age; while during the ensuing ten years the deaths from pneumonia form but 3.64 per cent. of the total mortality during that period of life. These statements do not agree with our own experience in private practice, since of fifty-seven cases that we have seen in which this point was noted, thirty occurred under five, and twenty-six between five and eleven years of age, showing that the frequency in the first five and the subsequent six years of life is very nearly the same. Lobar pneumonia is much less frequent in private practice in the first two than in the succeeding years of life.

Sex.—A larger number of cases occur in boys than in girls. The excess is probably not more, however, than may be accounted for by the preponderance of male over female children. Of fifty-five cases of the lobar form in which we have noted the sex, thirty occurred in boys, and twenty-five in girls.

Constitution.—It is doubtful whether constitution has much or any influence upon the liability to the disease. Dr. West says that, according to his experience, weak health is not a predisposing cause. We are convinced that lobar pneumonia attacks strong and vigorous children more frequently than those of more delicate constitution. In children of feeble health and weak stamina, the very same causes which produce pneumonia in the robust, give rise to bronchitis, or perhaps to catarrhal pneumonia.

Season.—The disease is most prevalent during the winter and early spring months, as will be seen from the accompanying table, in which is shown the mean monthly mortality in Philadelphia, for the seven years ending 1870, from this disease, as well as from bronchitis. From this it will be seen that in December, January, February, and March (and the same is very nearly true of April also), the deaths from these diseases are three times as numerous as in August.

Months.	Mean monthly mortality for 7 years, from Pneumonia and Bronchitis.	Mean total mortality from all causes (including still-born) for 7 years.	Mean monthly percentage for 7 years from Pneumonia and Bronchitis.	Mean monthly temperature (F.) for 7 years.
January, .	Pneumonia, . 46 Bronchitis, . 6.29	1296.71	3.54 0.48	30.87°
February, .	Pneumonia, . 45.57 Bronchitis, . 8.43	1206.71	3.76 0.69	33.89°
March, .	Pneumonia, . 48.57 Bronchitis, . 7.14	1344.29	3.61 0.53	40.85°
April, .	Pneumonia, . 41.57 Bronchitis, . 5.71	1281.14	3.24 0.44	52.27°
May, .	Pneumonia, . 34.76 Bronchitis, . 6.57	1234.29	2.81 0.53	62.77°
June, .	Pneumonia, . 26.14 Bronchitis, . 5.00	1178.14	2.21 0.42	71.97°
July, .	Pneumonia, . 24.14 Bronchitis, . 4.00	1837.00	1.31 0.21	77.71°
August, .	Pneumonia, . 20.14 Bronchitis, . 3.00	1825.43	1.10 0.16	76.62°
September, .	Pneumonia, . 14.57 Bronchitis, . 5.29	1215.43	1.19 0.43	68.31°
October, .	Pneumonia, . 22.43 Bronchitis, . 5.43	1218.14	1.84 0.44	56.30°
November, .	Pneumonia, . 28.86 Bronchitis, . 5.71	1052.14	2.74 0.54	46.68°
December, .	Pneumonia, . 39.57 Bronchitis, . 8.14	1191.00	3.38 0.58	34.74°

We have also placed in parallel columns the mean percentage of mortality from these two diseases, and the mean monthly temperatures, in order to show the marked correspondence between the coldness of the weather and the frequency of pneumonia and bronchitis. It is evident, however, that there is another element besides the mere temperature, in determining their frequency, since, in both February and March, more deaths occurred from these causes than in January, although this latter is the coldest month of the year. The additional element is undoubtedly to be found partly in the sudden atmospheric changes, and damp raw days which are so frequent, in both February and March, in our latitude, and partly in the impaired vitality found in many children, as the result of the intense cold of the preceding months.

Previous Diseases.—It is apt to occur as a complication of all the diseases of children, and most frequently in measles, pertussis, typhoid fever, enteritis, and bilious remittent fever. We have already stated that the catarrhal form is very frequently consequent upon severe bronchitis.

EXCITING CAUSES.—The continued action of some of the predisposing causes must be regarded as the exciting cause in the majority of the cases. External violence, as a severe fall, or a blow upon the chest, will sometimes act as an exciting cause. The action of *cold* is almost always alleged to be the immediate cause of the attack. M. Grisolle states that it is impossible to determine the exciting cause in more than one-fourth of the cases, and that in nearly all of these it is cold.

ANATOMICAL LESIONS.—*Lobar* pneumonia in the child is marked by the same physical characters as in the adult. The three stages of the in-

flammation—engorgement, red hepatization, and gray hepatization—exhibit the same alterations of the tissues as in adult life. Moreover, the three stages occur with about the same frequency in early as in later life. Dr. West (*loc. cit.*, 2d ed., p. 189) shows that the third stage occurs very nearly as often in children as in adults, he having met with it in the former in the proportion of sixty-eight per cent., while M. Grisolle found it in seventy-two per cent. of the latter. The chief difference in the disease, as it exists at the two ages, consists in the more frequent coexistence of all three of the stages in the young subject.

In the first stage, or that of engorgement, the affected portion of lung is distended, so that it does not collapse as much as the healthy portions, when the thorax is opened. It is heavier than usual, so that it sinks somewhat in water; it is of a brownish-red color; it pits upon pressure, and crepitates less than healthy lung, the crepitation being observable only here and there. The natural degree of cohesion between the tissues is somewhat diminished, so that the diseased portion is much less tough and elastic, and more soft and friable than it ought to be. When cut into, a large quantity of frothy and more or less deeply-tinged sanguineous fluid escapes.

In the second stage, or that of red hepatization, the lung is increased in volume, so that it continues to fill the side of the chest after that cavity is opened; it is dense and hard, has ceased entirely to crepitate, from the fact of having become completely impermeable to air, and sinks rapidly when thrown into water. Externally, the diseased portion is of a deep-red color, while internally the same color is observed, but often of such different shades as to give to a cut surface a marbled aspect. The cohesion between the tissues is, in this stage, much less strong than in health or in the first stage of the disease; the finger penetrates the lung with some ease, and the texture can be crushed between the finger and thumb. When cut into, there escapes a non-aerated and reddish fluid, which is much less abundant than in the first stage. The most important feature of red hepatization is, however, the granular character of the incised surface. This granular appearance is produced by the presence of numerous minute flat granular elevations, which are the air-vesicles distended with the plastic lymph which has been exuded within them. It is best seen by examining a torn surface of the lung. The most recent investigations tend to show that while some part of the exudation which distends the vesicles may be due to the multiplication of the epithelial cells lining their walls, the greatest portion is derived directly from the blood, and either escapes through the walls of the vessels without rupture or, to a small extent, is associated with rupture of the capillary walls. The exudation itself is seen, on microscopic examination, to have a finely-fibrillated structure inclosing a number of red and colorless corpuscles. (Rindfleisch.)

In the third stage, or that of gray hepatization, the lung continues to exhibit the same volume, density, impermeability to air, and consequent total absence of crepitation, as in the second; but the process of softening has made still further progress, so that a portion of the lung may be squeezed with the greatest ease between the finger and thumb into a pulp.

The color has now changed from a deep-red to a dirty light-gray, or a pale straw-yellow. When incised, the surface still presents a granular appearance, but the granules are more flat and irregular. The diseased portions are now infiltrated with a puruloid fluid, which escapes in considerable quantities in the form of a yellowish-gray liquid, whenever the lung is cut into.

Occasionally instead of local pneumonia involving an entire lobe or a considerable portion of one continuously, it presents itself in the form of several scattered patches, irregular in form and imperfectly circumscribed, but which present the three stages above described, of engorgement, red hepatization, and gray hepatization.

On the other hand, in catarrhal pneumonia there is not the same process of free exudation into the air-vesicles, but instead there is a rapid multiplication of the alveolar epithelial cells which soon fill the vesicles. The accumulated cells are mixed with some sero-mucous secretion, but there is found little or no trace of fibrillation, such as is present in the exudation of the lobar or croupous form. The catarrhal inflammation affects separate lobules, and then gives rise to nodules scattered over and through the lung. These vary in size, from that of a hemp-seed, if a single lobule be affected, even to that of a pigeon's egg, if a large number of affected lobules coalesce. At first they present a reddish or reddish-gray appearance; but later, owing to fatty changes in the proliferated cells, the color changes to a grayish or yellowish-gray. This change may at times occur in the centre only of the nodule, or over its whole extent, or again on its circumference. If the case is destined to end favorably, the cells, after having undergone complete fatty change, are absorbed, or else, an excess of serum being present, may be discharged into the bronchioles and expectorated. The alveolar wall returns to its original condition and a perfect cure is effected. Or, on the other hand, the changes in the alveolar wall persist, the contents of the alveoli undergo cheesy or caseous metamorphosis, and the case passes into a chronic and highly unfavorable stage.

Abscesses are not very uncommon in the lobar pneumonia of children. They occur as a result of the third stage of the disease, so that in the same lung may be observed the first, second, and third stages of the inflammation and abscesses. The cavities of the abscesses are generally circular, sometimes oval, and they measure from half a line to an inch or more in diameter. Sometimes the abscess is multilocular, each of the purulent cavities being partially separated from its neighbor by a wall of hepatized tissue. They are found in various parts of the lung, but seem disposed, generally, to approach the surface of the organ. When the latter event happens, adhesive inflammation between the pulmonary and costal pleura usually takes place; but should this fail to occur, the abscess may rupture into the pleural sac, and produce pneumothorax. MM. Rilliet and Barthez met with two examples in their autopsies in which this accident had occurred, and they report another case in which it occurred during life, and in which the child recovered. We have met with three cases of pneumothorax ourselves, produced in the same way. One occurred in a boy eleven years old, during an attack of secondary pneumonia complicating

a severe bilious remittent fever. The patient recovered entirely after a most violent illness. The two others occurred in very young children, and proved fatal.

We are desirous, before closing our remarks on the anatomical lesions of the disease under consideration, of drawing attention to the subject of simple non-inflammatory congestion of the lung, for the reason that the latter has no doubt, especially when associated with collapse of the pulmonary tissue, been frequently mistaken for pneumonia.

Congestion of the lung occurs either in the lobular or lobar form, the distinction between the two being the same as that between lobular and diffused or lobar collapse. When lobular, the lung presents, generally along the posterior edge of the organ, disseminated lobules, distinctly circumscribed by the interlobular cellular septa, which are rather protuberant than depressed, more friable, and of a lighter purple color than collapsed lobules, and which afford, when squeezed, a considerable quantity of frothless bloody fluid. In very young infants, the congestive disposition often assumes the lobar or diffused form, and is supposed by M. Legendre to have frequently been taken for pneumonia. In this variety of congestion, the affected portion of the lung is increased in size, and is distended and gorged with fluids. The color of the congested part varies from a light to a dark purple, or almost blackish tint. The cohesion of the lung is also variable, the differences depending on the degree of the congestion. When this latter is very great, the part is very friable, while it is much less so under the opposite condition of things. Though the lung is harder in this state than natural, it still retains a certain degree of flaccidity which does not exist in true hepatization. Pressure causes an abundant exudation of blood and serosity from a cut surface, and the latter, instead of being granulated, as is always the case in hepatization, is smooth and even. Neither does the lung exhibit any granulations when it is torn. Lastly, inflation distends all the vesicles, and gives to the condensed parts their natural lightness and their rosy color, though, be it remarked, the development of the affected parts under the operation is not complete and entire, as in collapse, in consequence, no doubt, of the large amount of blood they contain.

Inflation of the lung after death has been much employed of late, as any one who has read the previous remarks on atelectasis must have seen, as a means of distinguishing between pneumonia and collapse. It was there stated that, whilst inflation distended and restored more or less completely to their natural condition parts of the lung that were merely collapsed, it failed almost entirely to have any effect on parts of the lung affected with true pneumonia. It is easy to understand why inflation should fail to exert much effect on inflamed lung, at least when the disease has reached the state of hepatization. In fact the alveoli are distended either with the croupous exudation of the lobar form, or the accumulated epithelial cells of the catarrhal form, so that it becomes impossible to force the air into the midst of the agglutinated structures. In the first stage of pneumonia, that of congestion, inflation will distend in some

degree the affected portions, but, in the second and third stages, not even the strongest force has any affect on the impermeable vesicles.

Lobar pneumonia is stated by most authorities to be generally confined to one lung, and such has been our own experience in regard to it, since of 58 cases in which its location was carefully determined, it was unilateral in 54, and double only in 4. It is much more common on the right than left side, according to most writers. In the 54 cases just referred to, the disease was seated 31 times on the right side, and 23 times on the left. It attacks the lower lobe much more frequently than the upper, though pneumonia of the upper lobe is much more frequently met with in children than in adults. Of 51 cases in which this point was determined, the upper lobe was the part affected in 20, while in 31 the base of the lung was the seat of the disease. Of the 20 cases of inflammation of the upper lobe, in 13 it was seated on the right, and in 7 on the left side. Of 31 cases occurring in the lower lobes, 15 were on the right, and 16 on the left side. In the 4 cases of double pneumonia, the inflammation attacked the lower lobes of both lungs in one; in one the postero-inferior part of both upper lobes was especially involved; while in the two others it attacked first the base of the left lung, and afterwards the summit of the right.

The statements just made as to the seat of the pneumonic inflammation, in the cases that have come under our own observation, do not, we are well aware, agree exactly with the experience of other observers. Dr. West, for instance, found (*loc. cit.*, p. 190) that double pneumonias preponderated greatly, in early life, over those wherein only one lung suffered. This, it will be observed, is widely different from the result of our experience, and it is also directly opposed to that of MM. Rilliet and Barthez, Ruz, and Barrier. M. Barrier, in fact, cites (*Mal. de l'Enfance*, t. i, p. 286) 144 cases of lobar pneumonia as having been observed by the authors just mentioned, and by himself, and of these only 15 were double. Our results in regard to the frequency of double lobar pneumonia agree, therefore, with those of the authors last mentioned, but they differ as to the relative frequency with which the two lungs are attacked. Thus, in our cases, the inflammation occurred with nearly equal frequency in either lung, whilst of 129 cases of unilateral pneumonia observed by the above authors, 84 were seated in the right, and 45 in the left lung. These writers state, as most others do, that lobar pneumonia of the lower lobe is more common than that of the upper lobe. This tallies with our observations, but, as it seems to be a general opinion in the profession, that inflammation of the summit of the lung is rare in comparison with that of the base, we wish to call attention again to the fact stated above, that of 51 cases, in which we ascertained accurately the seat of the disease, it was in the upper lobe in 20, and in the lower in 31.

In regard to catarrhal pneumonia, the statement of West above quoted is certainly true, and it is frequently found that both lungs are involved, and in some cases that there is a more or less symmetrical disposition of the affected lobules.

In the lobar pneumonia of children, as in that of adults, bronchitis does

not usually exist to any very considerable extent, though we have observed a few instances where acute catarrh of the upper air-passages has been followed by the development of extensive croupous pneumonia. It is true that before the recognition of the exact nature of collapse of the lung, the association of bronchitis and lobar pneumonia was thought to be more frequent, because a considerable number of cases of bronchitis with extensive collapse of lung-tissues were regarded as instances of true lobar pneumonia.

But, on the other hand, although the catarrhal form may occur without any associated bronchitis, it is undoubted that bronchitis does very frequently precede or accompany its development. When bronchitis is present it varies from simple increased vascularity with augmented mucous secretion, to intense congestion with purulent or pseudo-membranous secretion.

Pleurisy is a frequent complication, as it is found to exist in about half the cases.

Emphysema is another common complication. It generally occupies the upper part of the lung, or its free edge, and is found most strongly developed in the lung which presents the greatest amount of inflammation, or in both when both are inflamed. Its degree depends on the extent of the pulmonary inflammation and bronchitis, and the severity of the dyspnœa. The vesicular form is much more frequent than the interlobular.

SYMPTOMS; SKETCH OF THE DISEASE; COURSE.—In order to present a faithful account of the disease in its different forms, a general sketch of the symptoms will first be given, after which the most important ones will be considered separately under the head of particular symptoms, so that the reader may first obtain a notion of the course of the disease, and then become intimately acquainted with its details and peculiarities by reference to the remarks on each particular symptom.

LOBAR OR CROUPOUS FORM.—True lobar pneumonia, with well-marked hepatization, is not, according to our experience, a common affection in young infants in private practice, since out of fifty-two cases of the disease that we have met with in children, in which we have noted this point, only three occurred in infants within the first, and four in the second year. Of the three cases within the year, one occurred in a child six weeks old, and the other in one seven months.

In *new-born children, and those still at the breast*, pneumonia is apt to be of the catarrhal form, and very generally begins with more or less marked symptoms of bronchitis, though in some instances it commences suddenly, as it does in adults, without any previous sign whatever of bronchial inflammation. When it occurs during an attack of bronchitis, the symptoms which belong to the pneumonic inflammation will, of course, have been preceded by those which depend on the disease of the bronchial mucous membrane. In these cases, the development of the pneumonia will be indicated by an aggravation of the general symptoms, by an increase of fever with elevation of the temperature, by an increase of the oppression, by the fact that the cough and breathing both become more painful than before,

and in some cases by the occurrence of the physical signs peculiar to lobar pneumonia. In these latter cases, which are very rare, the character of the physical signs and the course of the case are very much such as are described below.

When lobar pneumonia appears as a primary affection in young children, without preceding bronchitis, as sometimes undoubtedly happens, though much less frequently than in children over five years of age, and especially than in adults, the attack is usually sudden. In a child of fourteen months old, we have known the attack to be ushered in by a convulsion, which, with infants, is not rarely the equivalent of a chill. Usually the first symptoms observed are restlessness, peevishness, disposition to cry, a diminished appetite for the breast, and feverishness. These symptoms are most marked in the evening and night. From the very first, or by the second day at least, cough is heard, and careful examination of the breathing will show that it is somewhat hurried. The cough is dry, short, and hacking, at first, and not very frequent, but it soon becomes louder, fuller, more straining, and especially it becomes painful. The fact that it is painful may always be ascertained by watching the motions of the child, its cry, and the expression of the face. We can always perceive, even in an infant, a disposition to restrain the cough, to smother it, a struggle to make it short and sudden, when it causes sharp pain. At the moment of the cough, too, a marked expression of pain, a sudden grimace or twisting of the features, may always be observed, which is accompanied or followed instantly by a loud, sharp cry, or a spell of crying. This grimace of pain, with the accompanying cry, we have never observed in simple bronchitis, but only in pneumonia and pleurisy. We have occasionally seen these symptoms so decidedly marked that they could not fail to have drawn any one's attention; as, for example, in an infant six weeks old, who died of a most violent and extensive pleuro-pneumonia, and again in a child thirteen months old, who died of pleurisy resulting in the formation of pure pus in the pleural sac. The nature and extent of the lesions were ascertained, in both cases, by examination after death. In another case of acute plastic pleurisy, developed during a scarlatinous albuminuria, the expression of anguish in the eye, and in the contracted features of the child, presented one of the most painful scenes we have ever beheld in the sick-room. The presence of pain in the side is shown also by the fact that full inspirations, caused by changing the position of the child, and those which occur during fits of crying, occasion a sudden arrest or stoppage, so to speak, of the act of inspiration, which gives to the crying, and often also to the breathing, a sobbing character, while across the countenance passes at the same moment the expression of pain already referred to. The breathing, which is only slightly disturbed at first, soon becomes frequent and attended with more or less effort, and gives rise to an unusual play of the nostrils, a symptom which ought always to attract attention to the respiratory system as the seat of disorder. It interferes also with the act of nursing, so that whether the child takes the breast less frequently than usual, from want of appetite, or seeks it with greater avidity than common, from thirst, the act of suck-

ing is attended with difficulty. The infant seizes the breast for a few instants, then lets go in order to breathe more easily, and seizes it again; or it drops the nipple suddenly and begins to cry, as though the act of sucking were painful from the necessity it begets of taking occasionally a fuller and deeper inspiration than usual. As a general rule, the bowels are torpid, while vomiting, which is rather unusual in older children, is quite common in young infants.

When the disease is once established, whether it have been preceded by bronchitic symptoms, or occur as a primary affection, the symptoms are generally well marked, so as to leave but little difficulty in the recognition of the disorder. The child now loses all gayety and cheerfulness, and becomes either dull and listless, or very restless, peevish, and troublesome. Young infants generally lie quietly on the bed, or in the lap, merely fretting and crying when they cough, or when they are moved for any purpose, while children of several months old, and those in the second year, are usually very cross and restless, crying and screaming when anything is done for them, and insisting upon being frequently moved from the cradle or bed to the lap, or from the lap to the cradle. As a general rule they are contented only upon the lap, always crying to get back when they are removed from it to the cradle or crib. In some instances, however, they, like young infants, are quiet and dull, being content to lie still when placed in a comfortable position, and crying only after coughing, for the breast or drink, or when disturbed.

A febrile reaction now displays itself in full force. The skin becomes hot and dry, and the pulse frequent, rising to 150 and 160, or higher, in infants, and to 140 and 150, or even 160, in those of several months old. The temperature rises very quickly, so that by the close of the first twenty-four or thirty-six hours it may reach 104° , 105° , or even 106° . The dyspnoea becomes more and more evident. The respiration rises to 60, 70, 80, or even higher. In a case of pleuro-pneumonia at six weeks of age, we counted it at 128. The breathing is at the same time more or less labored and difficult, the *alæ nasi* being seen to dilate spasmodically at each inspiration, while the motions of the chest, and especially those of the abdomen, are much stronger and more active than in healthful respiration. The cough is now more frequent than before, evidently painful, and usually dry, though sometimes a slight degree of looseness may be detected in the sound which it occasions.

Percussion now reveals manifest dulness over the seat of disease, which is usually the base, though not at all unfrequently the upper region of one side. When the disease is double, which is oftener the case, as already stated, in children than in adults, though not so often as has been supposed by some, the percussion will be dull of course over the affected region on each side. Together with the dulness of sound on percussion, and sometimes when this is faintly marked, there is an evident diminution of the elasticity of the walls of the chest, and this becomes, therefore, an important symptom, especially when dulness on percussion is not well-marked. The dulness on percussion is not, indeed, so marked a symptom in infants

as in adults, from the fact that the natural resonance of the chest is so much greater in the former than the latter.

Auscultation reveals over the diseased part distinct and abundant fine subcrepitant râles; but the crepitant râle or fine crepitation, which is the pathognomonic sign of pneumonia in adults, and which in them is rarely wanting, is absent in young children, or is heard only when they make deep and free inspirations. It is most apt to be heard in young children during the deep inspirations which they make just before crying, or during the act of crying. It is, therefore, much less constant, less strongly marked, and more fugitive, in children than in adults, and is, in the former, replaced in good measure by a fine subcrepitant râle. In connection with these symptoms we always have more or less well-marked bronchial respiration. This may be pure, which is rarely the case; it may be, as usually happens, associated with crepitant or subcrepitant râles, or it may be heard only in the expiration. In children who are old enough to talk, there is increased vocal resonance and fremitus; and in infants, we can detect and draw important conclusions from an undue transmission of the resonance and fremitus of the cry or cough.

The symptoms above described show that the inflammation has reached the second stage, or that of red hepatization. After attaining this point, the disease usually remains stationary for a few days, and then either subsides, in favorable cases, by the resolution of the inflammation, or in unfavorable cases, terminates fatally in this stage, or else passes into the third stage, and causes death by a more or less extensive suppuration of the lung. In favorable cases, which are said to be rare in very young infants, but more common in those several months old and in the second year of life, the severity of the symptoms gradually diminishes. The fever subsides, the pulse becoming less frequent, and the skin cooler and less dry; the breathing becomes easier and slower, and is attended with less pain; the cough grows looser, less frequent, less difficult, and ceases to be painful; the child begins to nurse without pain and with greater ease and facility; the restlessness and fretfulness, or the somnolence, when that has been a marked symptom, diminish, and the child becomes more placid, and sleeps quietly and tranquilly. The chest is now less dull than before on percussion; the bronchial respiration begins to diminish in intensity, and is very much masked by the subcrepitant râle, which becomes more and more evident, until at last it takes the place entirely of the bronchial breathing. The symptoms continuing to amend, the physical signs of the disease cease at length to be perceptible, the cough grows more and more loose and rare, the countenance becomes natural, the fever ceases, and convalescence is fully established.

In unfavorable cases, death may occur rather suddenly in the second stage, without any very decided change in the physical signs, from exhaustion or from the supervention of collapse of portions of the lung-tissue. In these cases, the breathing becomes more and more rapid and labored, or it becomes slower than before; the moist râles increase in abundance and extent, while the percussion often remains about the same; the difficulty of sucking increases, so that the child, when put to the breast, attempts to

draw but two or three times and then lets go exhausted and distressed, or it makes no effort whatever; the cough becomes less frequent, but is still painful and difficult; the skin grows pale and white, excepting about the face, hands, and feet, where it often assumes a bluish or cyanotic hue; the extremities, and often the face too, become cool; the child becomes exceedingly restless, and then dull and perfectly quiet or comatose, and death at last occurs from asphyxia. In another class of cases, which, however, are much more rare in very young children than in older ones, the disease passes into the third stage, or that of suppuration, so called. In such cases the febrile symptoms continue much longer than in those just now described; the pulse becomes, and continues for several days together, very frequent and jerking; the skin retains its heat and dryness, though it is often pale at the same time; the child is usually excessively irritable and distressed; the breathing is rapid and oppressed, and often very irregular and uneven; the dulness on percussion extends; the bronchial respiration becomes more distinct and is heard over larger surfaces, and is accompanied with less of the subcrepitant and crepitant râles; the cough is paroxysmal, painful, and often very harassing; the appetite is lost, and the sleep uneasy and often broken. These symptoms continue for several days, or a week or two, when they assume the same characters they exhibit in more rapidly fatal examples; that is to say, asphyctic phenomena develop themselves, and the child dies exhausted and comatose or perhaps convulsed, or after presenting for some hours, or a day, more or less severe spasmodic affections of different muscles or of the extremities.

The *lobar pneumonia of children over two years*, and especially of those over five years of age, exhibits most of the symptoms that characterize the same disease in adults. The chief differences to be noticed at these two periods of life, are the greater predominance of bronchitis in children, particularly in those under five or six years of age, which gives to the physical signs some peculiar features not observed in adults; the frequent absence of expectoration, and when it is present, certain differences between it and that of adults; certain peculiarities in the character and seat of the side-pain; and the existence in many instances of more marked and more dangerous nervous symptoms.

The mode of onset is very different in different subjects. Generally, the attack begins with violent fever, increased frequency of breathing, more or less pain in the side, and short, dry cough. In such cases there is no difficulty in perceiving that the disease consists of some form of thoracic inflammation. But, in other instances, instead of this open and frank development, the disease comes on with symptoms which might well mislead any but a very attentive and competent physician, as to the true nature of the case. The most common cause of obscurity is a predominance of the nervous symptoms, which often gives to the case very much the aspect of a meningeal inflammation. In an example that occurred to one of ourselves, a boy between six and seven years old was seized, after a short exposure during a ride on a raw and cold day, with violent fever, pain in both ears, severe frontal headache, and great sensibility to light when exposed to it. He was, at the same time, very drowsy, sleeping nearly the

whole day, but he could be roused when loudly and vehemently spoken to so as to answer a few questions and manifest great irritability, and, what was extremely suspicious of disease of the brain, when taken with the other symptoms, he vomited frequently. On the second day, the headache was very severe, the sensibility to light continued excessive, and he still vomited frequently, rejecting even water. The bowels were freely moved. There was up to this time no full cough, but only an occasional and slight hacking, that scarcely attracted attention. The respiration was accelerated, but there was no dyspnoea. No pneumonia could be detected, though carefully sought after. On the third day, the breathing was still more frequent, but not at all laborious; the vomiting continued, but the other nervous phenomena had lost some of their intensity, and auscultation revealed well-marked bronchial respiration before and behind, over the summit of the right lung, while over the same regions the percussion was dull. We have met with several cases in which the onset of pneumonia was attended with nervous symptoms that made the diagnosis difficult and obscure.

In other cases the onset of the disease is marked by symptoms of gastro-intestinal irritation, or by such a degree of fever and disturbance of the nervous system, with absence of evident local phenomena, as to render the nature of the attack obscure and uncertain. In one, for instance, occurring in a boy between four and five years old, and six weeks after recovery from measles, the attack began suddenly with violent fever, great restlessness and distress, vomiting, and distention of the abdomen. The case appeared to be one of gastro-intestinal disorder, as there was nothing to call attention to the thorax. On the second day, the symptoms were much worse, the skin being hot and dry, and the pulse one hundred and sixty in the minute, and jerking. The child was drowsy and heavy; it was difficult to make him answer questions, and his answers were confused and unintelligible; his movements were tremulous and uncertain. The tongue was dryish and very thickly coated, and he complained confusedly of pain in the abdomen, which was much distended, and sonorous on percussion. There was no sign of respiratory disease, except quickening of the breathing, and a very slight cough, scarcely to be noticed. At this moment, however, when scarlet fever was apprehended from the great frequency of the respiration, the drowsiness, and the tremulous character of the muscular movements, auscultation and percussion revealed the true nature of the sickness in the shape of a lobar pneumonia of the lower lobe of the left lung.

In a majority of the cases, however, instead of the obscure and deceptive onset we have just described, pneumonia begins with fever, acceleration of the respiration, pain in the side, and short, dry cough. In some instances the disease supervenes upon catarrh or bronchitis. The child ceases to play, refuses to be amused, and is either irritable and cross, or lies listlessly upon the bed, or, if still quite young, insists upon being kept upon the lap. In some few cases, in very young children, convulsions occur. The appetite is lost, or else very much diminished; the thirst is acute, and when the disease is once established, more urgent than in almost any other affec-

tion. Vomiting is quite common, especially in young children, but diarrhœa is rare, the bowels being generally more torpid than usual. From the first day often, and almost always by the second, we can perceive either crepitant or subcrepitant râles, and sometimes bronchial respiration, confined usually to one side, and more frequent below than above, though, be it remarked, not at all rare over the latter part.

As the case proceeds, the fever increases, the bronchial respiration becomes more distinct and is heard over a larger extent of surface, whilst the râles diminish in abundance. The skin is now very hot and dry, so as to impart a burning sensation to the hand; the pulse augments in frequency, seldom counting less than 140 in the minute, often mounting to 160, and in severe cases, and in young children, even to 170, and becoming full and hard; the respiration becomes more and more accelerated, until it rises to 40 or 50, and in a great many cases to 60, 70, or even 80, while it often becomes at the same time oppressed, and, when full inspirations are made, painful; the cough is frequent, dry, or almost dry, and painful at first, but after a few days begins to be moist, and, in children over six or seven years of age, is not unfrequently attended with an expectoration of rusty or sanguinolent sputa; the thirst continues intense, the appetite is null, and the child is very restless and irritable, or drowsy and inattentive. About the fourth or fifth day, as a general rule, the disease has attained its height, the febrile and local symptoms being then most marked and the extent of the inflammation greatest, as shown by the physical signs.

At this stage of the disease the bronchial respiration is generally strongly marked, being clear and distinct, audible both in inspiration and expiration, and accompanied by bronchophony and increased resonance of the cry. The dulness on percussion is also very evident, the change from the natural sound being easily perceptible on a comparison of the two sides.

The symptoms generally remain stationary at this point for one or two days, and then begin to subside. The heat of skin diminishes and perspiration often appears; the pulse falls in frequency and force; the respiration becomes slower, easier, and full inspirations can be taken without pain; the *alæ nasi* no longer dilate; the cough becomes quite loose and ceases to be painful; the thirst is less acute; the child loses some of its irritability and restlessness, and if it have been soporose and dull, becomes more wakeful and observant; the flushing of the face disappears, while the expression is more natural. On auscultation, the bronchial respiration is found to have lost some of its intensity; it has become more distant, or it is heard only in the expiration, and is mingled with, or in part replaced by, crepitant or abundant subcrepitant râles. The dulness on percussion is less marked. A little later the fever ceases entirely, the respiration reassumes its natural rate, the appetite returns, the thirst disappears, the cough subsides very much, and the child begins to be interested in its toys or occupations. About the tenth or fifteenth day, and in some cases rather earlier, convalescence is fairly established, though auscultation may still reveal some prolongation of the expiratory sound and diffuse resonance of the voice.

In unfavorable cases death seldom occurs early in the disease, but usually at some distance of time from the invasion, and in consequence, no doubt, of the transition of the inflammation into the third or suppurative stage. In such cases the disease has usually pursued the course just described up to the period of resolution; but, instead of resolution and convalescence taking place, the fever continues, though perhaps with diminished violence, the skin being less intensely hot, and the pulse less full and active, while it remains quite as frequent. The breathing is sometimes less frequent than before, but it is often more laborious, and very generally it becomes irregular, and is easily hurried under exertion. The cough varies very much, being sometimes almost suppressed, and in other cases very troublesome; it is almost always loose. The strength diminishes, the voice becoming weak and feeble, and the muscular movements tremulous and languid; the face looks pale, haggard, and sunken; the child is sometimes very restless, tossing about from time to time on the bed or lap, with a quick, short, and evidently feeble movement, or it is dull and soporose, awakening only when spoken to, but showing then by its fretfulness and peevishness that its intelligence is retained. While these symptoms are present, the extent over which the bronchial respiration is heard has generally augmented, showing the gradual extension of the hepatization, while outside of the part where the respiration is blowing, and sometimes over the same part, and intermingled with that sound, are heard more or less copious subcrepitant and mucous râles. This condition seldom lasts more than two or three days, at the end of which time the child dies in a state of coma, or after one or more convulsive seizures, which are the result of a gradually increasing asphyxia.

In other cases, again, the termination is more gradual. The child, after presenting many of the above symptoms, may seem to improve somewhat. The fever may diminish, the appetite return to some extent, the respiration become easier, the restlessness subside, and the child becomes more cheerful again; but the face continues pale, emaciation makes progress, the appetite fails again, the pulse remains frequent, diarrhœa comes on, the cough becomes more troublesome, thrush often attacks the mouth, the strength decays continually, and, after some weeks perhaps of struggling, the child dies in a state of great emaciation and debility.

CATARRHAL FORM.—The symptoms of catarrhal pneumonia are much more obscure and uncertain than those of the lobar form of the disease. Owing to the fact that the inflamed patches of the lung are disseminated or scattered through healthy portions of the organ, the signs afforded by physical examination are either very imperfect, or entirely masked by the sounds produced in the healthy texture. We are forced, therefore, to depend much more in this than in the lobar form, on the rational symptoms, in determining the nature of the sickness. The rational symptoms of catarrhal pneumonia are nearly the same as those of the lobar form. The chief differences between the two are in regard to the pain, the dyspnœa, and, when there is expectoration, the amount of the sputa. The febrile and nervous symptoms, and the disturbances of the digestive system, are the same in the two forms, the only difference being in their degree of

severity. In the lobar variety they are usually more acute and severe than in the catarrhal. The mere degree of temperature attained does not differ materially in the two forms, and it is not unusual to find a temperature of 103° on the second day of an attack of the catarrhal form. We have seen, however, that in the croupous form hyperpyrexia (105° to 106°) may occasionally be noted thus early in the attack. Roger (*loc. cit.*) notes, also, that while in lobar pneumonia the high temperature is sustained for six or seven days until defervescence occurs, in the catarrhal form the course of the temperature is marked by a succession of irregular remissions and exacerbations. The local symptoms present important differences which should be noted. In the form under consideration, the pain is either wanting entirely or is much less acute than in the lobar form. When the inflamed patches are few in number, and they are seated in the central parts of the lung, there is entire absence of pain; but when they are more numerous and superficial, pain is complained of, but it is usually diffuse, of slight intensity, changeable, and felt only during cough, or during full inspiration. It makes its appearance commonly on the first day, and very seldom after the third. Cough is rarely wanting. It usually marks the onset of the sickness, is extremely variable as to its frequency and severity, and is not acutely painful, as in the lobar form, unless the inflamed patches be superficial. There is seldom any considerable amount of expectoration, and in some cases none; when there is any it is small in quantity, and it may or may not be characteristic. In one case, however, that came under our observation, in which we had every reason to believe, from the nature of the rational symptoms, and from the absence of physical signs, that the disease was catarrhal pneumonia, there was a rare expectoration of thick, viscous mucus, streaked with blood. The respiration is accelerated, and when the lesion is at all extensive, there is dyspnœa, the degree of these symptoms being determined by the extent and number of the inflamed patches.

The physical signs are not, as above stated, very significant. The percussion is natural, the amount of tissue consolidated being insufficient to affect the sonorousness of the chest. According to Eustace Smith, a general want of healthy pulmonary resonance can be detected over the back by means of broad percussion, striking with three fingers upon three fingers placed on the chest-wall as pleximeters. Auscultation affords no signs of the pneumonic inflammation when the number of affected patches is small; when they are more numerous it is of some, but not of very great utility. Crepitant râles are sometimes heard here and there over circumscribed points of the thorax, and, disseminated in the same way, there is also heard in some instances rude respiration, prolonged expiratory murmur, and bronchial respiration. When, as often happens, this form of the disease co-exists with bronchitis, it will be entirely concealed by the dry and moist râles of the latter affection. If a number of affected lobules coalesce and form a superficial patch of some size, we may have distinct signs of consolidation over a circumscribed area. This is, however, rare.

The *duration* of lobar pneumonia has been fixed with considerable accuracy by the observations of various persons. As a general rule, the

disease reaches its highest point of severity in about four or five days, then remains stationary for one or two days, and diminishes regularly until between the tenth and fifteenth day, when convalescence is established. In our own practice, the longest duration in 23 unmixed cases, in which the period was accurately noted, was 17 days, and the shortest 5. The duration of the 23 cases was as follows: In 1 case, 17 days; in 3 cases, 14 days; in 1, 11; in 4, 10; in 5, 9; in 3, 8; in 2, 7; in 2, 6; and in 2, 5 days. One case lasted 33 days, but it was accompanied and followed by bronchitis.

It is difficult to assign any definite duration for the catarrhal form. In favorable cases, judicious treatment will often be followed by convalescence in from a week to ten days. But in many instances the symptoms persist longer than this; and not unfrequently the case shows a strong tendency to pass into a chronic form.

PARTICULAR SYMPTOMS; PHYSICAL SIGNS.—In order to practice auscultation and percussion in a young child, it should be placed, by the mother, in a sitting posture on her knee, while the physician, kneeling on the floor, or sitting on a low chair, makes the examination he deems necessary. If the child be old enough to take notice, it should be attracted and amused by some toy or glittering object. Even, however, should it cry violently, much valuable information is to be obtained by the examination, for we can ascertain the presence or absence of râles and their characters during the deep inspirations between the cries, and can observe resonance of the cry and cough, and practice percussion.

The physical signs of *lobar pneumonia* are crepitant or subcrepitant râles, feeble respiratory murmur, bronchial respiration, bronchophony, exaggerated resonance of the cry and cough, and dulness on percussion. They are, in fact, the same in the great majority of cases as in adults. Under five years of age, this form often begins with subcrepitant râles, while after that period the earliest auscultatory signs are crepitant râles, and feeble respiration. The bronchial respiration makes its appearance soon after the subcrepitant or crepitant râles, is heard first in the expiration, and then in both inspiration and expiration, and is accompanied by bronchophony, resonance of the cry and cough, and dulness on percussion. Bronchial respiration was present in 46 of 57 cases of lobar pneumonia observed by ourselves; crepitant râles were present in 31, and subcrepitant in 10.

These alterations of the auscultatory phenomena are confined to one side, in the great majority of cases, and are best observed over the postero-inferior portion of the lung. MM. Rilliet and Barthez state that they have never known the bronchial respiration to disappear, in favorable cases, before the fifth day, and in the majority not before the seventh, eighth, or ninth; while, in fatal cases, it continued to the moment of death. Its persistence is always a highly unfavorable symptom in very young children, whilst in those who are older, as in adults, it sometimes remains for several days or weeks, though the general symptoms have entirely disappeared. In *catarrhal pneumonia*, on the other hand, as we have already stated, the physical signs are often present on both sides, and are much less definite.

RATIONAL SYMPTOMS.—*Cough* may be said to be invariably present. It is dry at first, and not very frequent, but in one or two days becomes more frequent, often very troublesome, and from dry and harsh, becomes more or less humid and loose. It continues until the disease moderates, lasting generally from nine to sixteen days. In fatal cases it usually persists to the last. In infants it is not very frequent, occurs in short paroxysms, and in fatal cases often ceases one or two days before death. MM. Rilliet and Barthez remark that in pneumonia of the upper lobes it has a peculiar character. It is little, short, smothered, as it were; or piercing, teasing, or slightly hoarse. We will merely add that cough is sometimes scarcely noticeable in cases which simulate hydrocephalus, during the early part of the attack. In a case already referred to, in which the symptoms bore for several days very much the aspect of a meningeal attack, there was no full cough whatever during the first two days; on the third day, though auscultation and percussion showed the existence of pneumonia of the upper lobe of the right lung, the child coughed only three or four times, and it was not until the sixth day that it became at all frequent. In three other cases the cough was so slight in the early stages of the disease, during the continuance of the cerebral symptoms, as not to have been noticed unless particularly inquired after. Later in the attack, after three, four, or five days, and as the cerebral symptoms moderated, the cough became frequent and loose, and the pneumonic symptoms pursued their regular course.

Expectoration is almost invariably absent under five years of age. MM. Rilliet and Barthez, and Dr. Gerhard, have never observed rust-colored sputa under the age mentioned. In older children there is sometimes, though not very often, voluntary expectoration. Even in them, however, the sputa seldom present the characteristic rust-color and viscosity observed in adults, but consist simply of mucus tinged with blood, or of whitish, brownish, viscous, or non-viscous phlegm. We once, however, saw a child three and a half years old, voluntarily expectorate viscid mucus, tinged copiously with blood. Sanguinolent expectoration was noticed in five of the fifty-seven cases seen by ourselves (not including the one just spoken of). In three the sputa were of the characteristic rusty color, in one they were composed of mucus streaked with blood, and in another portions of mucus streaked with blood were rejected by coughing, and some also by vomiting. The age of the five subjects, just alluded to, was in each case between five and nine years. In another case (not included amongst the five), in a girl seven years old, affected with lobar pneumonia supervening upon pertussis, there was a free expectoration of tenacious mucus, sometimes streaked or dotted with blood, sometimes brownish, and sometimes rust-colored.

M. Valleix mentions a whitish or sanguinolent viscous froth, as sometimes escaping from the mouth of new-born children laboring under the disease, and Bouchut has also noticed in a single case, a little reddish sanguinolent froth, situated on the edge of the lips of an infant with pneumonia. We have never met with this symptom, but know of one case of a child within the month, who, during an attack of pneumonia, vomited

mucus tinged with blood. The child died, and presented the lesions of pneumonia. The nipples of the mother were perfectly healthy, so that the blood could not have been sucked by the child from them, but must have consisted of the sputa which had been swallowed after being coughed into the fauces.

It is scarcely necessary to say that the absence of expectoration is only seeming, for children undoubtedly cough the sputa into the fauces, whence, instead of being rejected, as by the adult, they pass into the stomach.

Thoracic Pain.—It is impossible to ascertain the presence of this symptom with positive certainty prior to the age at which children talk, and very often not for some time after, as they refuse or do not know how to describe their sensations. And yet, even in infants, the presence or absence of the stitch in breathing, and of pain in coughing, may be inferred, almost with certainty, by watching the gestures and expression of the child, and the cries which accompany a full inspiration and the act of coughing. In effect, the deep inspirations induced by moving the child, those which take place during vomiting and gaping, and those also which occur in the act of coughing, cause the child to cry out suddenly and sharply, and give at the same moment an expression of acute suffering to the countenance, which can be referred to nothing else than the causes just mentioned, and which reveals almost as plainly as words the painfulness of a deep inspiration and of the act of coughing. In older children, we have several times known the pain to be most intense, causing bitter and repeated complaints, with crying, fretting, and evident acute suffering. The seat of pain, as complained of by children who talk, ought also to be noticed, since the account given by them might well mislead an unwary and inexperienced physician. It is quite common, in fact, for them to refer the pain to the false ribs, to one of the flanks, to the abdomen, and even to the hip.

The *respiration* is always quickened, except where the constitution of the patient has been greatly deteriorated by long and severe illness or other causes, under which circumstances it may remain at the normal rate, or be very slightly accelerated. This symptom usually dates from the invasion, soon after which the breathing rises as high as 40, 50, and 60 in the minute in older children, and from 60 to 80 in the younger. It sometimes becomes excessively rapid, reaching, as it did in a case of pleuropneumonia in an infant six weeks old under our charge, 128. In favorable cases, the acceleration subsides usually about the seventh, eighth, or ninth day. In most of the cases the breathing is even and regular, while in others it is short, abdominal, uneven, and jerking. When the dyspnoea is very great in a young child, the nostrils dilate widely, the mouth remains open, and its angles are drawn downwards and outwards; the last of these symptoms is almost a fatal one. Sometimes the rhythm of the function is changed, so that it begins with a sudden, active, and moaning expiration, followed by the inspiration, after which comes the interval of rest. MM. Rilliet and Barthez state that unequal, jerking respiration, occurs almost exclusively in cases of inflammation of the upper lobes.

Physiognomy.—The face is almost invariably flushed. The color, at

first scarlet, becomes after a day or two deeper and darker, and in severe cases assumes a livid-red tint. We have noticed in very severe pneumonia, in addition to the deep-red tint, a peculiar glazed appearance of the skin, which looks as though it had been varnished, while the edges of the flush are distinct and abrupt. The lips are generally deeply colored, simultaneously with the face. The flush commonly subsides about the same time, or a little before the diminution in the rate of the respiration. In fatal cases the face is apt to lose its color, and becomes pale and sallow, as the unfavorable symptoms become more and more marked. We have noted extreme pallor of the face in very severe cases occurring in infants, and, although indicative of great danger, a favorable result has followed in some instances.

The expression of the face is one of anxiety and oppression in the early stage; in very severe cases, or those about to terminate unfavorably, the features become drawn and contracted.

Fever exists in all the idiopathic cases. The pulse, at all ages, is rarely under 130 from the first to the sixth or seventh day; in the youngest children it rises as high as 140, 160, and even 180; while in those who are older, it is seldom above 140. In favorable cases, it diminishes about the fifth, sixth, or seventh day. In fatal cases, it is apt to diminish at the same period, but soon becomes more frequent and continues so to the end.

The range of *temperature* in pneumonia is higher than in any other inflammatory disease of children. This is true of the disease in both of its forms, lobar and catarrhal. The highest temperature we have ourselves recorded for pneumonia is 106° , while Roger, in his latest contribution to this subject (*Recherches Cliniques sur les Mal. de l'Enfance*, 1872, p. 356), states that 105.8° has been observed by him in 2 out of 47 cases of pneumonia. In two-thirds of the entire number the mercury reached or exceeded 104° ; and the mean of the highest temperatures in all the cases was 103.9° . Such high degrees are more apt to be found in children over than under two years of age. The maximum temperature reached in any case would not seem to be much influenced by the seat or extent of the inflammation.

In lobar pneumonia the course of the temperature is regular and characteristic. The accession of fever is often very sudden, and the mercury may rapidly rise to its maximum point, reaching 104° , 105° , or even 105.8° , within twelve hours from the onset. After the first abrupt rise, it is sustained nearly at the same point, with moderate morning remissions and evening exacerbations (the variation usually not exceeding one or one and a half degree) until defervescence, which is usually rapid, or even abrupt, occurs. In the catarrhal form, on the other hand, the initial rise of temperature is less abrupt, and the course of the fever is marked by the occurrence of irregular remissions and exacerbations, which Roger attributes to the development of successive patches of pneumonia.

The ratio between the temperature, pulse, and respiration may be quite closely preserved, and the elevated degree of febrile heat be associated with marked acceleration of the pulse and breathing. Thus Roger found

that the mean furnished by 47 cases was: temperature, 103.94° ; pulse, 133; respiration, 52. It is not, however, at all rare in both forms of the disease for the pulse-respiration ratio to vary from the normal 1 to 4 to 1 to 2, or even 1 to 1.5. In the lobar form, it will frequently be found that, following the defervescence, the temperature falls below the normal point, as to 98° or 97° , for a day or two.

The *nervous system* shows more or less marked symptoms of disorder. There is restlessness, peevishness, and irritability during the day, and these increase towards evening. As the night advances, the child becomes still more restless; infants will not sleep except in the arms, and wake crying or fretting every few minutes or hours; older children sleep uneasily, talk in their sleep, or start and cry out, and are often delirious. In some instances, the irritability is most distressing, both to the child and to those around. The child is constantly fretting and whining; it wants its play-things, but will not touch them; food, but rejects it; and slaps and scolds at everything about it. Convulsions sometimes occur at the invasion. They last an uncertain length of time, and are usually followed by insensibility, from which the child wakes with fever, accelerated respiration, and cough indicating the true seat of disease to be the lungs, and not the brain, as might at first be supposed. We have met with but four cases, all of the lobar form, attended with convulsions. One occurred in a boy between ten and eleven years of age, on the second day of the disease. The attack was induced more, however, by an unwholesome meal taken on the first day of his sickness, than by the mere effect of the local inflammation. In a second case, which occurred in a boy between five and six years old, there were two convulsive seizures, a violent one on the first day of the pneumonia, and a slighter one a few days later. In a third case, which occurred in a boy aged two years, the pneumonia occurred in the course of intermittent fever; there were three marked convulsions, but the child subsequently recovered perfectly. In the fourth case, to which reference has already been made, a child of 14 months old, who was improving from a sharp attack of catarrh of the upper air-passage, was seized with croupous pneumonia, of two-thirds of the right lung, ushered in by repeated convulsions, a temperature of 106° on the second day, and with intense fever, with unconsciousness and tonic carpo-pedal spasm persisting for ten days; notwithstanding complete recovery followed. The headache is sometimes very severe; in a few instances we have known it to be so violent as to constitute the most prominent symptom of the case. On one occasion, indeed, it was so intense, and so much complained of, during the first two days of the fever, as to withdraw our attention from the true seat of disease, and it was not until the third day that we discovered the existence of pneumonia. The cough was in this, as in other instances, in which the nervous symptoms were strongly marked, so slight as to escape notice.

Digestive Organs.—Complete anorexia is generally present from the first; the thirst is intense, greater indeed than in almost any other affection of childhood. The tongue is moist, as a general rule, and covered with a whitish or yellowish fur. Vomiting and diarrhoea occur at the

invasion of about half the cases in hospitals; in private practice, vomiting often occurs, but diarrhœa much less frequently.

Urine.—The amount of urine is materially lessened in acute lobar pneumonia, the extent of the reduction being from one-third to one-half (Parkes).

During the height of the disease the urea is increased, and with it, as in most febrile diseases, the uric acid. Simon and Redtenbacher first called attention to the fact that the chloride of sodium is diminished or entirely absent during the early period and at the commencement of hepatization, and reappears during, or rather after resolution; and further researches have fully confirmed this observation, since very few exceptional cases have yet been recorded. The disappearance does not depend upon the reduced diet, since Howitz and Parkes both state that even when chloride of sodium is administered, none can be detected subsequently in the urine. And the fact that it is in reality retained in the system, is further shown by the very excessive excretion during convalescence. According to Beale's observations, the exudation in the lung is very rich in chloride of sodium; and it has been found that as this salt disappears from the urine, it appears in the sputa, and in turn as it returns in the urine, it disappears from the sputa.

It is true that more extended observation has shown that the chloride of sodium is absent or deficient in many other affections, both febrile and inflammatory; but still, although not pathognomonic of pneumonia, this sign is an aid in its diagnosis, and probably serves to distinguish it from collapse of the lung or from tuberculous consolidation.

One more condition of the urine in pneumonia, although as yet, so far as we know, only noticed in adults, deserves attention. We allude to the presence of albumen, which has been noticed by several observers, as Finger, Becquerel, Parkes, and Heller, in almost 45 per cent. of their cases; though others, as well as ourselves, have found it much more rarely.

The period of its occurrence is variable; according to Heller and Parkes, it appears at the time when the chlorides are most deficient, as hepatization advances. The fatality is much increased in cases where albuminuria is present; the combined record of the observers above referred to, yielding a mortality of almost 50 per cent. of such cases; whilst the mortality in cases without albuminous urine was only 14 per cent. According to Parkes, renal cylinders are very common in the albuminous urine of pneumonia; and a little blood is also frequently present, but is usually out of all proportion to the albumen.

DIAGNOSIS.—The *lobar* pneumonia of children is most liable to be confounded with bronchitis, pleurisy, and meningitis. There is little probability, however, that lobar pneumonia would be mistaken for bronchitis by any but a careless or incompetent observer; for the presence, in the former, of subcrepitant, and very often of crepitant râles, of bronchial respiration, bronchophony, resonance of the cry and cough, and dull or flat percussion, confined to one side, would easily distinguish it from bronchitis, which is marked by dry and moist râles over both sides of the

chest, and by a normal condition of the percussion. It is difficult and often impossible, as already stated, to detect the existence of catarrhal pneumonia, or at least to make the diagnosis with absolute certainty. The cause of the difficulty, as before explained, lies in the fact that it presents, in a great many instances, no clear physical signs. When the number of inflamed lobules scattered through the healthy texture of the lung is small, and especially when they are deeply seated, no alteration whatever of the natural respiratory sounds can be perceived, and we are obliged to depend entirely upon the rational symptoms,—the accelerated breathing, oppression, pain, cough, fever, and the absence of the physical signs of other pulmonary inflammation. A careful study of the temperature may here be of service. We have seen that in pneumonia the temperature usually rises quickly to 104° or 105° , while in bronchitis, it rarely attains even the lower of these, and often does not exceed 101° or 102° . Sometimes the presence of the characteristic sputa of pneumonia will, in older children, make the diagnosis clear. When the inflamed lobules are situated near the surface of the lung, we may, in some instances, detect crepitant or fine subcrepitant râles, and bronchial respiration, over circumscribed portions of the lung, and there would be, under such circumstances, no hesitation as to the diagnosis.

It has been stated that pneumonia might be confounded with pleurisy. This could not happen in regard to the catarrhal form, as the slighter degree of the pain, the extent of the râles, the moderate bronchial respiration, and the absence of dulness on percussion in this disease, would prevent such a mistake. The distinction between pleurisy and the lobar form is more difficult, but may generally be made out by attention to the fact that pleurisy is rare under six years of age; by the greater severity of the pain, the less abrupt and extreme elevation of temperature, the absence of râles and presence of friction-sound, the effect of change of position on the sounds yielded by percussion, the shorter duration and greater mildness of the general symptoms, the entire absence or small amount of expectoration, and by the continued dryness of the cough in pleurisy; and lastly, by the disposition on the part of pleurisy to become chronic, while lobar pneumonia nearly always runs an acute course.

Lobar pneumonia in children not unfrequently simulates, in its early stage, an attack of meningitis, constituting a form of the disease sometimes called *cerebral pneumonia*. Vomiting, constipation, extreme irritability or restlessness, and complaints of headache, occur in both; while the absence of thoracic symptoms to draw attention to the true seat of the disease in pneumonia, may readily mislead. The cough in the early stage of pneumonia is sometimes very slight, and not being observed by the attendants, is not reported to the physician. The frequency of the respiration is overlooked, or, if noticed, as ascribed to the fever, which is supposed to depend on the cerebral inflammation. In pneumonia, however, the vomiting is not usually very frequent, nor very obstinate, nor are the bowels so much constipated as in acute hydrocephalus. These variations from the ordinary symptoms of the latter disease, minute though they be, ought to attract the notice of the physician, and lead him to examine the

case more carefully, when, in all probability, the physical signs would immediately reveal the pneumonia. We may mention, in illustration, that we attended a boy six years old, who, for three days, suffered from violent fever and excruciating headache, which last was the only symptom complained of. There was neither cough, expectoration, nor any marked acceleration of the respiration. After three days the headache moderated, and he had slight pain in his side; on examination, we found him laboring under well-marked lobar pneumonia. In April, 1847, we were called to see a boy nineteen months old, who had been taken sick with slight fever, a little hoarse cough, and mild pharyngitis. After remaining in this condition for five days, he began to be drowsy and very irritable, the surface became pale, and the extremities rather cooler than natural. From the sixth to the tenth day there was great somnolence, the child sleeping nearly all the time; when waked from sleep, he was always exceedingly irritable and cross, scarcely opening his eyes, and then shutting them again immediately to avoid the light, which was evidently painful. During this time he took scarcely any food, but little drink, and vomited several times freely; the bowels were moved without medicine; the surface remained very pale, and the extremities often cool; the pulse was frequent and small, the respiration perfectly regular, for which reason it attracted no attention, and there was *no cough* whatever. Under these circumstances, we hesitated between regarding the case as one of meningitis, or of hydrocephaloid disease, as described by Dr. M. Hall. We took the latter view, however, and treated it with small quantities of brandy, cold to the head, and the frequent employment of mustard pediluvia. From the eleventh day the child began to improve; it would open its eyes from time to time, and look round for a few moments; the face began to show a slight degree of color, and the palms of the hands, which had been white and transparent, exhibited a tinge of the natural pink hue which they have in children. Observing about this time that the respiration was accelerated, though perfectly free and regular, and without cough, we counted it, and were astonished to find it 80 in the minute. We now examined the chest carefully, and finding slight dulness on percussion with bronchial respiration, over the inferior half of the left side behind, immediately understood the nature of the case; it was one of latent pneumonia, simulating hydrocephalus. The child was now treated for pneumonia, and after an illness of twenty-seven days longer, recovered perfectly. As the case progressed, the rational signs of pneumonia were more and more apparent, the cough becoming frequent and painful, and after a time loose, while the cerebral symptoms gradually disappeared.

In addition to these cases, we have met with several others which during the early stage resembled very closely the invasion of cerebral disease. One of these has already been referred to in the account of the symptoms of the disease. Two others occurred in children within the year, and one in a child between one and two years old. Attention, however, to the rate of the respiration and the physical signs, and the presence of slight cough, revealed, in two of them, after a little hesitation, the true character of the attacks. The third case, which occurred in one of the children within the

year, was unattended by any cough during the first few days, and was, therefore, very obscure, until our attention was attracted by an acceleration of the respiration, when the physical signs, and, at a later period, the cough, explained the real nature of the attack. We may remark, in addition, that in all these cases, the absence of constipation, the infrequency and short duration of the vomiting, and some clearness of the intelligence when the child was fairly roused, though but for a few moments, from its state of somnolence, were other reasons for doubting the attacks to be meningitis.

We have dwelt at length upon the danger of making this serious mistake in diagnosis, in the hope that our remarks will aid in impressing upon the mind of our readers the great importance, which has indeed been alluded to on several previous occasions, of making a careful examination of the chest by auscultation and percussion in every case of acute disease in children, even though the symptoms do not appear to indicate any affection of the heart or lungs.

Dr. West states that pneumonia is often overlooked in teething children, in whom the cough is called a tooth-cough, whilst the diarrhœa, which frequently occurs, and becomes the prominent symptom, is supposed to depend upon dentition, and is alone attended to. The diarrhœa is obstinate, and when, at last, the cough attracts attention, it is ascribed to phthisis, and the physician is astonished to find at the autopsy purulent infiltration of the lungs, but no tubercles, and no disease of the intestines. The diagnosis is to be correctly made, under such circumstances, only by careful physical examination.

The disease with which catarrhal pneumonia is most apt to be confounded is bronchitis. The two are frequently associated, and, owing to the fact that the physical signs of consolidation of lung-tissue cannot always be detected in the catarrhal form, our diagnosis must sometimes be based solely upon the general symptoms. If we find, however, that with or without previously-existing bronchitis the cough has suddenly assumed the pneumonic character, the temperature rapidly risen, and the degree of dyspnœa suddenly increased, there is strong reason to believe in the super-vention of catarrhal pneumonia. If we find, in addition to this, impaired resonance at various points, with imperfect bronchial breathing and perhaps subcrepitant râles at these spots, our diagnosis would be confirmed. We must also bear in mind the resemblance which exists between catarrhal pneumonia and lobular collapse of the lung. The points of resemblance and the fact that both are apt to appear during the course of a bronchitis have already been alluded to. In lobular collapse, however, although the dyspnœa may be extreme, the symptoms do not indicate any increase in the inflammatory action, but, on the other hand, the temperature is normal, or even lower than natural.

PROGNOSIS.—It may be stated in general terms that lobar pneumonia is the more dangerous in proportion as the child in whom it occurs is younger; and that the secondary, consecutive, or intercurrent form of the disease is much more dangerous than the primary. It is usually supposed to be almost necessarily fatal in new-born children, and to be still very dan-

gerous up to the sixth year of age. There has been so much confusion, however, in regard to atelectasis of the lung and true pneumonia until within a few years past, that it is scarcely possible to trust to former statistics upon this point. From six years of age up to fifteen, the disease is generally curable when of the primary form; when of the secondary form the result is much more doubtful, and will depend in great measure, of course, on the nature of the disorder during or after which it occurs.

MM. Rilliet and Barthez (*loc. cit.*, p. 535) state that they lost about one-eighth of their cases in private practice. Of these, the youngest was a year old, the oldest, three years old. To quote their own words: "Some evidently died of accidents caused by the medication (poisoning by tartar emetic); one was the victim of a relapse, due to faulty hygienic care; and others died of cerebral pneumonia of the upper lobe; they were undergoing the process of dentition." In the hospital, they lost a seventh of their patients. The subjects under five years of age died of cerebral, gangrenous, or intestinal complications. Those over five years of age died, some because they were scrofulous or feeble, the inflammation though lobar, being double; and the others, in consequence of the inflammation having become complicated with pleurisy, scarlet fever, or meningitis. They add, that in the hospital, six-sevenths of the patients attacked with secondary pneumonia died.

In 1862, however, Barthez stated, in a memoir to the French Academy (*Med. Times and Gaz.*, May 10th, 1862), that during the previous seven years, having abandoned the use of depletion in the pneumonia of children, he had treated 212 cases, with a loss of but two patients.

The results of our experience, which, it ought to be remarked, has been acquired chiefly in private practice amongst the easy classes of society, have been as follows: Of 66 cases of well-marked lobar pneumonia, only 2 were fatal. Of these two, one occurred in an infant six weeks old, and was accompanied with extensive and violent pleurisy, and the other occurred in a child between two and three years old, lasted thirty-three days, and was attended with considerable bronchitic inflammation.

In addition to these, we have seen a certain number of cases which are not included in the statistics of our own experience, since some of them were only seen, and, perhaps, but a single time, in consultation, while others occurred among the children in the large public institutions of this city. A far larger proportion of these latter cases proved fatal.

We may conclude, therefore, that pneumonia under two years of age is always dangerous, and much more so when secondary than when primary; that primary pneumonia, between the ages of two and five years, will, if treated judiciously, terminate favorably in the great majority of cases in private practice; and that when the disease attacks children between six and fifteen years of age, the termination is nearly always in health.

The following are some of the most unfavorable symptoms of the disease: convulsions; small, weak pulse; extreme rapidity of the respiration; persistence of the bronchial respiration in young children; incomplete resolution of the disease within the ordinary period; excessive and obsti-

nate diarrhœa; severe cerebral symptoms; great emaciation; greatly altered physiognomy; excessive irritability; and a yellowish tint of the skin. M. Trousseau regards as an unfavorable symptom the occurrence of swelling of the veins of the hands, which he supposes to depend on an obstruction to the function of hæmatosis.

The prognosis in catarrhal pneumonia is much more grave than might be supposed. Supervening, as it so frequently does, in very young children already exhausted by a previous attack of bronchitis, there is danger that the violence of the attack may cause it to prove fatal in the acute stage. The symptoms which would indicate great danger are high temperature, greatly altered pulse-respiration ratio, with extreme frequency of breathing, lividity of face, and the evidences of serious interference with aeration of the blood.

When the attack is less violent, the chances for future recovery are naturally much greater; but, it must be remembered that this form of pneumonia frequently runs on into a chronic stage, and, by the persistence of the changes in the alveolar walls, and the cheesy metamorphosis of the accumulated epithelial cells in the alveoli, leads to the development of chronic pulmonary phthisis. The prognosis therefore should always be a guarded one.

TREATMENT.—When one of the former editions of this work was published, a great change had begun to take place in medical opinion as to the proper treatment of disease, and especially of acute disease. In that edition this change of opinion was referred to, and its effect upon our own convictions and methods of procedure freely acknowledged. Since that period this revolution, as it might be called, has continued to make progress, until, at the present moment, no one can candidly express his own views without referring to it. In view of these facts, we shall not hesitate to write at some length on the treatment of pneumonia, in order that our readers, and especially the younger members of the profession, may be able to comprehend not only the changes that have taken place, but some of their causes.

There is another consideration which has been forced upon us by time and experience, which makes us unwilling to dismiss the treatment of so important a disease in a few words, and this is, that the method of cure to be followed in individual cases must be determined not alone by the simple fact that the patient has an inflammatory exudation in the lung-tissue, but, in large measure, by the state of the general vitality of the subject. What folly, for instance, to suppose that we can safely apply the same therapeutic measures to a case of pneumonia in a child just issuing out of severe measles, to one in the midst of a dangerous typhoid fever, to another in the spasms of whooping-cough, or, finally, to one who was yesterday in consummate health, with every function, up to the moment of the attack, in the finest possible working order. To be sure, this is putting the case in very strong terms, but they are not too decided to make our meaning clear.

Moreover, we think there is a tendency, in some of the later works on diseases of children, and in some, too, of the general treatises on the prac-

tice of medicine, to lengthened scientific descriptions of anatomical changes, symptoms, diagnosis, etc., and to a corresponding diminution of the space devoted to therapeutics. This error, as we think it (not to be wondered at, perhaps, when we consider the relative difficulty of writing on these different subjects), we desire to avoid, and, indeed, we have found it impossible to state our opinions on the subject except at some length.

Bloodletting.—Twenty years ago depletion formed an almost inevitable item in the treatment of pneumonia, but, within the last eight or ten years, the views of most observers have undergone a more or less radical change in regard to its utility and necessity. Some have abandoned it altogether, whilst others employ it still to a moderate extent. In order that the younger practitioner may see the changes which have taken place in this respect, we shall quote the views of some of the more important authorities, and then give our own.

Dr. Charles West (4th Am. ed., from 5th English ed., page 285), writes as follows: "I cannot forget the good results which I saw years ago from the abstraction of blood at the outset of an attack of pneumonia in previously healthy children." He, however, does not advise depletion when small crepitation has become generally diffused, still less when dulness or bronchial breathing is perceptible. He gives no statistics as to his own results whatever. Dr. J. Lewis Smith, of New York, in his work, does not even mention bloodletting. Dr. Eustace Smith (*Medical Times and Gazette*, May 3d, 1873, page 460), says he has never drawn blood from a child suffering from pneumonia, and that he has never met with a case in which such a method of treatment has appeared to him to be in the slightest degree desirable. Dr. Thomas Hillier, of London, says of bloodletting that it "is now for the most part discarded. I have never had occasion to resort to it." He says further, however, that cases might occur where it would be proper to recommend it. Such conditions would be, the second day of the disease, a large extent of inflammation of the lung-tissue, full and bounding pulse, great pain and dyspnœa, and a temperature of 105° or more. If these conditions existed in a previously healthy child, he would think it wise to take a few ounces of blood from the arm. We have already referred to the communication from M. Barthez to the Academy of Medicine of Paris, in April, 1862, intended to vindicate the expectant treatment of pneumonia in early life. In this paper it is stated that of 212 cases of lobar pneumonia, occurring between the ages of two and fifteen, in the course of seven years, at the Hôpital St. Eugénie, only 2 had a fatal termination, although no approach to active treatment was adopted in more than a sixth of the number. Dr. J. Hughes Bennett gives, in *The Practitioner*, for May, 1869, the results of the restorative treatment of pneumonia in 153 cases. Of these, 129 were simple and 24 complicated cases. Of the 129 simple or uncomplicated cases, of which 35 were double, all recovered. Among the 24 complicated cases there were 5 deaths, making of the whole number a mortality of 1 in 30 $\frac{2}{3}$ cases. Dr. Bennett's cases all occurred in adults, but the results are useful to us as showing the effects of this kind of treatment.

In a former edition of this work it was stated that we had treated 50

cases of well-marked lobar pneumonia, with two deaths, in private practice. Full notes of only 46 of these cases were kept. Of the 46 cases, 39 were primary or uncomplicated, and 7 secondary or complicated. The 2 fatal cases occurred, one at six weeks old, and this was attended with very severe pleuritic inflammation, and the other between two and three years old; the latter case lasted 33 days, and was attended with considerable bronchitic inflammation. Depletion was employed in 16 of the 39 primary, and in 2 of the 7 secondary cases. It is proper to state that depletion was not employed in either of the fatal cases.

How difficult is the task of estimating the comparative value of different plans of treatment in any given disease! MM. Rilliet and Barthez lost one-eighth of their cases of pneumonia in private practice, and one-seventh in hospital. We lost one-twenty-fifth of ours in private practice, a result very nearly as good as Dr. Bennett's, though ours were all in children under 15 years of age, and of 37, whose ages were recorded, 19 were under 5 years (2 in the first year, 3 in the second, 5 in the third, 4 in the fourth, 5 in the fifth). Dr. Bennett condemns bleeding almost wholly; we took blood in 16 of 39 primary, and in 2 of 7 complicated cases, and did not deplete at all in the 2 fatal cases. M. Barthez reports 212 cases, treated by the expectant method, with only 2 deaths, or less than one in a hundred; and these cases, too, in children between 2 and 15 years of age, in hospital practice. These last statistics are the most surprising we have seen. We have been unable to find the original memoir of M. Barthez, but have seen the report made to the Academy of Medicine, by M. Blache (*Bulletin de l'Acad. Imp. de Médecine*, t. xxx, p. 21), on the memoir, in which it is stated that "the author has taken care to eliminate the lobular or generalized pneumonias, the pseudo-lobular pneumonias, broncho-pneumonias, and catarrhal pneumonias; he has also thrown aside the lobar congestions which occur in the course of low fevers, and the secondary lobar hepatizations; that is to say, those which occur in the course of any well-determined disease, and particularly tuberculosis." We cannot help thinking that the elimination of so many forms of pneumonia, must be a chief reason for the very great success of the plan of treatment used.

This much, however, has been plainly established by the observations and experience of late years, that the old plan of bleeding, as a rule of absolute practice, merely because of the existence of pneumonia, and especially the Sangrado system of bleeding *coup sur coup*, was a gross mistake, and one which did great harm. But we do not think it has been proved that the restorative or expectant system, to the exclusion of bloodletting under any circumstances, is always and inevitably the right one. We have been led to think that bloodletting was not the only cause of the heavy mortality under the old systems of treatment, but that the use of such agents as antimony, ipecacuanha, and perhaps calomel, in large and frequently administered, and long-continued doses (and particularly the antimony) by their action upon the stomach, in destroying all power to take and digest food, and by the general prostration which their action (especially antimony) upon the nervous system occasioned, were answer-

able for a large share of the fatal results of those days. We doubt, in fact, whether depletion, used in anything like moderation, is not safer for the patient than the continued use, for two or three days, of nauseants and depressants, more particularly of antimony. But of the action of antimony upon children, we shall speak more at length hereafter.

Our own opinion, after the enlarged experience of later years, is, that depletion should not be used save in exceptional cases. When the pneumonia is pursuing a regular and safe course, it is best to trust to the simple means to be spoken of hereafter, and to follow a mild expectant method. Where the physician doubts as to its propriety, and especially when he is young and inexperienced, it is safest to abstain from it entirely, or to employ it only in a very moderate degree. But there is a certain class of cases, in which we believe that local depletion, by cups and leeches, is not only allowable but most useful. When the subject is vigorous and strong, with a fine sanguification; when the temperature is very high; the pulse strong and full; the muscular force good, and the side-pain and cough very severe, we think that the local abstraction of from two to four ounces of blood, at the age of three or four years, has great power to relieve all these symptoms. Again, when the dyspnœa is very great; when the heart pulsates with great force, whilst the pulse is small and feeble, showing that the right heart is overloaded, and the arteries comparatively empty, in consequence of obstruction to the passage of blood through the lungs; and when the child is tolerably vigorous, and not reduced by previous illness, a moderate venesection is often of more use, and of more efficacy in palliating these conditions than any treatment we know of. The quantity to be taken should seldom be over four ounces, at the ages of from three years and upwards. We venture upon these statements the more boldly when we find such men as Chambers and Niemeyer, and even Bennett, giving the same advice. Dr. Bennett (*loc. cit.*) lays down amongst his axioms the following: "Small bloodlettings, of from six to eight ounces, may be used in extreme cases, more especially in double pneumonia and broncho-pneumonia, as a palliative to relieve tension of the bloodvessels and congestion of the right heart and lungs." Niemeyer (*Textbook of Pract. Med.*, Amer. ed., vol. i, p. 184), says, pithily: "Highly as I prize venesection, however, in certain emergencies which may arise in the disease, I had rather that any one, dear to me, and sick of pneumonia, were in the hands of a homœopath, than in the hands of a physician who thinks that he carries the issue of the malady upon the point of his lancet." He recommends venesection in three conditions: 1. When the pneumonia has attacked a vigorous and hitherto healthy subject, is of recent occurrence, the temperature being higher than 105° F., and the frequency of the pulse rating at more than 120 beats a minute. "Here danger threatens from the violence of the fever, and free venesection will reduce the temperature and lessen the frequency of the pulse. In those who are already debilitated and anæmic, bleeding increases the danger of exhaustion. Should the fever be moderate, bloodletting is not indicated, even in healthy and vigorous individuals." 2. "When collateral œdema, in the portions of the lung unaffected by pneumonia, is causing danger to life, the pressure of the

blood is reduced by bleeding, and by prevention of further transudation of serum into the vesicles, insufficiency of the lung, and carbonic acid poisoning are averted. Whenever the great frequency of respiration, in the commencement of pneumonia, cannot be traced to fever, pain, and to the extent of the pneumonic process alone, *as soon as a serous, foamy expectoration appears, together with a respiration of forty or fifty breaths a minute*, and when the rattle in the chest does not cease for awhile after the patient has coughed, we ought at once to practice a copious venesection, in order to reduce the mass of blood, and to moderate the collateral pressure. The third indication for bleeding arises upon the appearance of the symptoms of pressure upon the brain, not headache and delirium, but a state of stupor or transient paralysis." We have made this long quotation because the authority is so high, and because we have nowhere found such clear and concise statements upon this most important point of practice.

Antimony.—In a former edition of this work it was stated that tartar emetic, in the dose recommended by some of the highest authorities of the day, had been found by us a very dangerous drug. Time has but confirmed this opinion. At the time we were in the habit of administering it in doses of a forty-fifth or sixtieth of a grain every hour or two hours. This was at a time when Rilliet and Barthez used it in doses of from two to four grains dissolved in four ounces of water, in twenty-four hours, for very young children, and for those who were older six grains in the same space of time. They continued it for two, three, or four days, and advised its suspension should it give rise to excessive vomiting or severe diarrhœa. Dr. West at that time gave it in doses of one-eighth of a grain, at the age of two years, every ten minutes, until vomiting was produced; to be continued every hour or two afterwards for a period of twenty-four or thirty-six hours. Dr. West had reduced the doses, and the time of continuing it, one-half, between the time referred to and the date of his essay on pneumonia, published in 1843.

The doses used by us, as mentioned above, may seem to some who have not employed them ludicrously small, but we soon found that even they were quite frequently, in certain constitutions, more than could be given with safety. Antimony, even in those small quantities, sometimes caused a very peculiar general prostration. Perhaps without any vomiting whatever, or with only a rare effort at that act, the patient would refuse all nourishment, become very pale and weak, grow limp and motionless, take on a haggard and pinched expression of face, pass into a state in which it would pay no attention to what was going on around, be very peevish and irritable when disturbed, get a very frequent and feeble pulse, and look to an experienced eye as though a very little deeper degree of such prostration might end fatally. After seeing this condition a few times, and finding that the withdrawal of the drug and the use of small doses of brandy (ten to twenty drops in water or milk) every hour or two hours, was followed by rapid improvement, we learned the greatest caution in the use of the remedy. Of late years we never use tartar emetic at all, but give, not unfrequently, in strong and vigorous children, with high febrile heat and rapid circulation, small doses of the precipitated sulphuret of antimony,

always watching its effects carefully, and withdrawing it at once should the above symptoms make their appearance. The formula found most useful and safest is the following:

R. Antimon. Sulphurat., gr. j.
 Pulv. Doveri, gr. iij.
 Sacch. Alb., gr. xij.

M. et. div. in chart. no. xii. One to be given every two, three, or four hours.

To infants under two years of age it is best to give no antimony at all.

Mercury.—In former years, calomel was given freely in nearly all cases of pneumonia, and its excessive use undoubtedly did much harm. But it is in our judgment an error to proceed to the opposite extreme, and to forbid altogether its use in this disease. It should not, however, be given excepting to meet some clear indication; and we will mention the conditions under which, of later years, we chiefly prescribe it. When, for instance, especially in the early stage, there are signs of gastro-hepatic congestion, such as a flabby tongue with whitish fur, fulness of the hypochondriac regions, anorexia and perhaps nausea, a few small doses of calomel, or of blue mass, followed by a mild saline laxative will be followed by relief. Again, when in the stage of consolidation, there are high fever, extreme gastric irritability, and marked nervous symptoms, we believe that we have seen positive benefit from its continued use in very small doses (as for a child of 15 or 18 months, gr. $\frac{1}{4}$ every three or four hours) associated with the use of quinia in full doses by suppository. It is at times impossible to administer any of the alkalies, and still more so quinia, by the mouth without provoking vomiting and interfering seriously with alimentation; and, under such circumstances we have seen the gastric irritability allayed, the power of retaining food restored, and probably the resolution of the exudation favored by the use of a very gentle course of calomel as above mentioned.

Salines.—*Citrate of potash*, either in the form of the neutral mixture or dissolved simply in water with a little sugar, is one of the best febrifuges that can be used. In doses of two and a half grains to children over three or four years old, and half a grain to a grain for younger children and infants, every two hours, it is an excellent remedy. It may be given alone or combined with small doses of syrup of ipecacuanha and opium. Spirit of nitrous ether may be added when the urine is scanty or when the ipecacuanha cannot be borne.

The solution of *acetate of ammonia*, either alone or combined with the spirit of nitrous ether, is useful when the child is feeble, and when the stomach or bowels are irritable, in which case the citrate of potash sometimes offends the stomach and acts upon the bowels. The dose of this remedy may be from twenty or thirty drops to half a drachm or a drachm, according to the age, in sweetened water, or some aromatic water, every two hours.

Quinia is unquestionably a remedy of great value in both forms of the pneumonia of children. When given in full doses, it diminishes the intense febrile heat and the great rapidity of pulse; and, at the same time, is be-

lieved by many observers to possess a tendency to check the extension of the exudative process. It is usually perfectly well accepted by the stomach, and does not interfere with the power of taking food; while, on the other hand, by its tonic influence, it must be of service in sustaining the system until the necessary stages of this exhausting disease have been passed. We are in the habit of giving about one grain three times daily to a child of from twelve to eighteen months, and one and a half grains three to five times daily at the age of three to five years. It may be administered conveniently by diffusing it in a delicate syrup of liquorice. In those cases, however, where much irritation of the stomach exists, it is better not to give it by the mouth until the stomach is thoroughly quieted, but to use it in rather fuller doses in the form of suppositories, which should be made of diminutive size.

Ipecacuanha is preferable to antimony in all conditions except those referred to above. In infants under two years of age, in children of highly nervous temperament, or of feeble and delicate constitutions, in most cases of the secondary form, and in all mild cases, it is much safer than the other drug. The most convenient preparation is the syrup, of which ten drops may be given every two hours at four years of age, five drops between one and three years, and from one to three drops to infants of two or three months. It is often useful to combine the spirit of nitrous ether with it, and, when the stomach is irritable, or the patient very restless and irritable, to add small doses of opium. When the patient is much oppressed by the presence of secretions in the bronchi, and not too much prostrated, an emetic is often very useful. *Ipecacuanha* is the most suitable remedy for this purpose, as it produces less exhaustion and depression than any other, except, perhaps, alum.

Muriate of ammonia has of late years been very largely employed in the acute pulmonary affections both of adults and children. It has seemed to us to possess the power of hastening the softening and resolution of the exudation, and, when there is expectoration, of rendering it freer and less viscous. The best period for administering it is, in the lobar form, after the hepatization is clearly established and the attack has reached its full development; or, in the catarrhal form, after the acute symptoms have somewhat subsided. It may then be given, associated with the febrifuge employed, or else dissolved in a little syrup of Tolu, or syrup of wild cherry bark and water. The proper dose is one grain for children under two years of age, and two to three for those between two and five years, given every six, five, or four hours, according to its effect and the way in which it is tolerated by the stomach.

If the resolution of the exudation does not progress rapidly, and especially if symptoms of exhaustion make their appearance, the carbonate of ammonia may be substituted, in about the same doses, for the muriate.

Laxatives.—A mild laxative dose, with or without a previous small dose of a mercurial, is useful at the beginning of the attack, when the child is constipated, and when the abdomen is tumid and hard. A teaspoonful of castor oil, or two teaspoonfuls of simple syrup of rhubarb, will answer every purpose. After this period, even the mildest laxatives should be

used with the greatest care, and only when demanded by some clear indication. The food taken is almost exclusively liquid, and of this the amount is rarely very large. If, therefore, the bowels are moved spontaneously every two or three days, or, if in case more decided constipation exists, an enema will provoke a satisfactory movement every third day, there is no occasion whatever for the internal use of any laxative. They should be used only in case stools cannot be otherwise secured, or in case there is evidence of irritation from accumulations of undigested food or of morbid secretions. The mildest laxative alone should be ordered, since even a single dose of a powerful or irritating purgative may do irreparable harm by disturbing the stomach, or exciting diarrhœa, and thus inducing greater irritation and exhaustion of the system.

External Applications.—MM. Rilliet and Barthez were of opinion that neither blisters, Burgundy pitch, nor tartar-emetic plasters, exerted the least influence upon any one of the symptoms of pneumonia, but that, on the contrary, they increased the fever. Dr. West gave up the use of blisters entirely, in consequence of the irritation and fever they occasioned, and because of the disposition to sloughing which he observed to follow their use amongst the poor. At one time we thought we had observed great benefit from the use of a blister when other means had failed to produce some moderation of the symptoms after four or five days. If they are used at all, it ought to be with great care, especially in very young or feeble children, whose nutrition is depraved. In children of less than two or three years old, a blister should never remain on the skin longer than two hours. As a general rule, the mother should be told positively to remove it at the end of one hour and a half, even though the surface be still unchanged. A warm bread and milk poultice is then to be used as a dressing, and this rarely fails to cause vesication in a few hours. Employed in this way, we have had but once the misfortune to see a blistered surface slough, and this occurred in a child whose skin had been very much irritated by frictions with amber oil and ammonia.

Since the spring of 1845, however, when we were led to make frequent use of mustard poultices and foot-baths in the treatment of the bronchitis and pneumonia of measles, we have rarely employed blisters, but have preferred the employment several times a day of the remedies just indicated. Two parts of Indian meal and one of mustard, for young children, and for those who are older, equal parts of each, are to be mixed with warm water, and spread thickly like a poultice on a piece of flannel or rag five or six inches square. This is to be covered with fine muslin, linen or gauze, and applied first over the back and then over the front of the thorax. It may remain from fifteen to forty minutes, or until the child cries or complains, or until the skin is reddened. The mustard foot-baths may be employed at the same time with the poultices. These applications are useful whenever the oppression is very great, and, when resorted to in the evening, they often allay irritability and dispose the child to sleep. The number of applications to be made in a day must depend on the urgency of the symptoms. We have employed them from once a day to every two or three hours.

The use of hot linseed-meal poultices, so highly recommended by Dr. Chambers, of London, we have found apparently of much service. The poultices should be spread quite thick on a cloth or flannel as broad as the circumference of the thorax, and deep enough to cover the whole chest, from the collar-bones to the hypochondria. The size of the poultice will be determined by the amount of lung-tissue, though we usually have them spread large enough to cover the whole affected side, even if but a portion of the lung be involved. They should be changed several times during the day, and care should be taken that a fresh hot poultice is ready for application before the one in place has been removed. In order to keep them in place, it is often necessary to have a tape stitched on in front, and a tape behind, which can be tied over the shoulder in the manner of a shoulder-strap. The use of poultices in this way has several disadvantages, and there is much to be said in favor of substituting for them a layer of uncarded wool or cotton large enough to cover the entire thorax, and kept in place by stitching it to the inside of the merino or flannel shirt.

Tonics and stimulants are to be resorted to in cases which manifest undoubted signs of debility. When, therefore, the attack occurs in a feeble child; in secondary cases; when the inflammation remains unresolved after the use of other remedies, and when extensive bronchial respiration persists, though the fever has moderated; or when, in any case, during the acute stage, the child falls into a typhoid state, as shown by pallor of the surface, frequent, uneven pulse, dry tongue, prostration of muscular power, and either incessant jactitation or the listless quiet of exhaustion; attention must be paid to the state of the constitution even more than to the local disease. The vital forces must be sustained and strengthened in order to give time and power to carry on the operations necessary for the removal of the local obstruction. To effect this purpose, we must depend upon the use of food, alcoholic stimuli, and certain tonics. The *food* most suitable for such a condition is milk, animal broths, soft-boiled eggs, and perhaps small quantities of raw or slightly cooked meats. The best stimulants are brandy, given either in milk or in water, as the child will best take it, and wine and water, or wine-whey. The amount of brandy to be given may be stated as 3, 4, or 5 teaspoonfuls in the course of 24 hours at 4 years of age. We have above indicated the conditions that call for the use of alcohol in the pneumonias of young children. Many cases do well without it at any stage, and it should not be given as a matter of mere routine. But when clearly indicated it will usually be found to be well borne and to afford decided relief. The best *tonic* to give in conjunction with the alcoholic stimulus is, as has already been stated, quinia, which should be administered in the quantity of from gr. iv to gr. vj in 24 hours, given in divided doses. When the exhaustion is marked, especially when associated with great embarrassment of respiration and copious viscid secretion from the bronchial tubes, we should recommend the use of the muriate or carbonate of ammonia, either of which may be given in mucilage, in doses of gr. ij to iij every 3 or 4 hours, at 4 or 5 years of age.

Opium is constantly of great service in the treatment of pneumonia. It should always be used when the patient suffers much, either from the side-pain or from cough, whether this be harassing and exhausting from its mere frequency and persistence, or from its effect in developing the stitch; when there is painful jactitation, an unusual degree of distress and malaise, or marked tendency to morbid vigilance. When the fever is very high, the pulse vibrating, the nerves on a rack, opium is of the greatest advantage. The mere comfort it gives is a good warrant for its use, but it has long seemed to us to aid in shortening the duration and lessening the severity of the constitutional disturbance.

The choice of the preparation, and the doses and times of administration, must vary in different cases. When used early in the case, to act upon the circulation and allay general irritability, it is best to give it with the febrifuge every two or three hours. When used to control cough, it can be added to the sulphurated antimony in the form of Dover's powder, as already suggested, or to the syrup of ipecacuanha and spirit of nitrous ether, in a liquid form; or when the cough is particularly troublesome at night, as often happens, it can be given with more advantage in a single dose, or two doses in the evening. The preparations we have found most useful are, laudanum, especially the *tr. opii deodorata*, paregoric, solution of morphia, or Dover's powder. Under six months of age, half a drop of laudanum, from five to ten drops of paregoric, or two or three drops of the solution of morphia, may be given, and repeated twice or three times in the twenty-four hours, according to the effects. From the age of six months to the end of the second year, these doses may be doubled. In the third and fourth years, two drops of laudanum, ten to twenty of paregoric, five to ten of the solution of morphia, may be used several times a day. Where the remedy is given every two or three hours, we have found one drop of laudanum quite enough at the ages last mentioned. When the dose is given only at night, from three to five drops of laudanum, ten to fifteen of the solution of morphia, and thirty to fifty of paregoric are sufficient as a general rule. After the age of four or five years, the doses must be increased in proportion to the age. In very young children, the doses given at first should always be watched with a good deal of care, and never carried to such a quantity, or continued long enough, to induce constant and heavy drowsiness or stupor. In some instances of very nervous and hyperæsthetic children, in whom there is determined, by the violence of the reaction, a degree of irritability of the nervous centres tending to the tetanic state, the doses must be much larger than those mentioned; but here the physician should see the patient himself at least twice, and sometimes three times in the day, to watch and regulate by the dose the exact action of the drug. We have occasionally seen the cough most harassing and exhausting in its effect, occurring almost with every breath, and lasting from twelve to twenty-four hours. Under such circumstances, a mixture like the following has proved most beneficial in our hands:

R. Tr. Opii Deodorat.,	gtt. xxxij.
Vin. Antimon.,	gtt. xxxij.
Ext. Valerian, Fl.,	fʒij.
Syrup. Simp.,	fʒij.
Aquæ,	fʒiss.—M.

Dose. A teaspoonful every hour or two hours, at the age of four years and upwards, until the cough is controlled.

Paregoric, in the proportion of two drachms to half an ounce, in place of the laudanum, in the above mixture, sometimes proves more soothing and comforting.

GENERAL MANAGEMENT.—Since the reign of restorative medicine has set in, the general management of the patient has received a degree of attention which it had never attracted before. Under the expectant plan it constitutes, indeed, the chief portion of the treatment. The most important points to be attended to under this head are the diet, drinks, clothing, air, and state of repose.

The patient ought not to be allowed to go entirely without food even in the early days of the disease, neither should there be any effort made to stuff the child with large quantities of nourishment. The appetite is nearly always in great measure abolished, at first, and food is unwillingly taken except in very small quantities. A nursing child must not be allowed to nurse as heartily as usual. If it attempts to do so, it is probably from thirst and not from hunger. Water, therefore, should be offered to it from time to time, and the breast be allowed only every three or four hours for short periods. Weaned children should have only milk, always reduced by the addition of half or one-third water, and pure water ought to be given frequently. The thirst in this disease is intense, and the physician should himself see that the patient has water freely. We have seen the most violent and obstinate screaming, and painful restlessness, quieted at once by a copious draught of cold water. In children over two and three years of age, milk and water is still the best food; but when this is refused, thin chicken or beef tea may be given in doses of a wineglassful or a gill every four hours. After three or four days have passed by, the administration of food is a very important part of the treatment. The child should now be induced, by persuasion and even gentle force, to take a little food at least three or four times in the twenty-four hours. As the severity of the symptoms subsides, the food ought to be increased in quantity.

The clothing ought to be such as to keep the body comfortably warm. In winter, which is the season when the disease almost always occurs, thin and soft flannels ought to be worn, and, when the child is very restless, either in the bed or on the lap, a sack made high in the neck, with the sleeves to the wrists, buttoning in front, and consisting of a soft and pliable woollen stuff, ought to be put over the bed-dress.

The room ought to be, if possible, a large one with a high ceiling, well ventilated, warmed by an open fire, and kept at a temperature of 65° to 68°. If the child is very young and delicate, a temperature of 70° is not too high, if only the ventilation be good.

The bed or crib is the proper place for a child with pneumonia. The lap of the mother or nurse is a poor substitute for an even, elastic, and steady mattress. We have long endeavored to keep our little patients in bed. A very young infant must, of course, often be taken up to be nursed, soothed, or cleansed, but, as soon as possible, it ought to be replaced in the crib. Children a year or two old can generally, with good management, be kept the greater part of the time in bed. Those of three and four years old and upwards ought always to be confined to the bed. A little firmness on the part of the mother will almost always accomplish this end, and it is a highly important one, and well worth even a quarrel at the beginning of the sickness. We have seen a child three years old kept by a weak and over-tender mother and grandmother nursed on the lap for three weeks, until they were exhausted and demoralized, and the child had œdematous feet from their dependent position during so long a time.

Repose and quiet of mind and body, as complete as can be attained, are things of great value, and to secure them a good bed and a cheerful and resolute manner on the part of the nurse are as important for the child as for the adult. It is only in bed, too, that an even temperature and an avoidance of draught can be fully secured. A direction given by some of the French writers, and by Dr. Gerhard, is not to allow very young children to lie for too long a time in one position in bed, or in the nurse's arms, as it is apt to produce a stasis of blood in the dependent portions of the lungs, and thus to maintain or increase the disease. Dr. West recommends, whenever the inflammation has reached an advanced stage, or involved a considerable extent of the lungs, that the patient be moved with great care and gentleness, lest, as he has often seen occur, convulsions be produced.

ARTICLE III.

BRONCHITIS.

DEFINITION; SYNONYMS; FREQUENCY; FORMS.—The term bronchitis is now universally employed to express inflammation of the mucous membrane of the bronchi; frequently it is called catarrh, and catarrhal fever.

It has been stated, under the head of Pneumonia, that many of the cases known amongst us by the popular term catarrhal fever, are in fact, cases of pneumonia. We shall on account of this misapplication of names endeavor to draw the distinction between bronchitis and pneumonia with great care.

Since bronchitis and pneumonia have been more carefully distinguished in the mortality returns of this city, bronchitis is found to be the cause of a much smaller proportion of deaths than would have formerly appeared.

Thus, during the ten years ending with 1879, the total mortality from all causes (excluding still-born children) was, at all ages, 166,942; under fifteen years of age, 76,063; and under five years, 66,613. The mortality from bronchitis during this period was, at all ages, 2556, or 1.53 per cent. of the entire mortality; under the age of fifteen years, 1774, or 2.33 per cent. of the mortality under that age; and under five years, 1731, or 2.59 per cent. of the mortality under that age.

It is, however, one of the most frequent of the diseases of childhood, especially during the winter and early spring months. It is said to be more common as a secondary than as an idiopathic disease. Of 115 cases observed by MM. Rilliet and Barthez, only 21 were idiopathic. Of 123 cases, however, that we have recorded, 76 were primary, and the remaining 47 secondary. The diseases during the course of which it is most apt to occur, are pertussis and measles.

We shall describe three forms of the disease: 1, *acute bronchitis* of moderate severity; 2, *capillary bronchitis*, or acute suffocative catarrh; 3, *sub-acute* or *chronic bronchitis*.

CAUSES.—Amongst the *predisposing* causes of the disease, *age* is one of the most important. MM. Rilliet and Barthez suppose it to be much more common in children over than in those under five years of age. Of one hundred and fifteen cases observed by them, thirty-seven occurred between the ages of one and five years, and seventy-eight between six and fifteen years of age. It is scarcely fair, however, to compare a period of nine years with one of only four, as is done in the above statement. Of one hundred and twenty cases that we have seen in private practice, in which the age was noted, fifty-four occurred between birth and two years of age; thirty-nine between two and four years; twelve between four and six; six between six and ten; and three between ten and fifteen. Of eighty-one cases under four years of age, of which we have kept an accurate record, eleven occurred in the first half of the first year of life, twenty in the second half, making thirty-one for the first year; twenty-one occurred in the second year of life, nineteen in the third, and ten only in the fourth; showing that the liability is greatest in the first year of life, and particularly in the last half of that year, that it continues very strong in the second and third years, being nearly equal in each of these, and that it then suddenly diminishes. It would seem also that the simple acute and the acute suffocative forms are most common under six years of age, while the secondary cases occur more frequently after that age.

As to the influence of *sex* on the liability to the disease, it would appear from our experience to be rather more common in girls than boys, since of ninety-nine cases in which this point was noted, fifty-four occurred in girls and forty-five in boys. The fact of its being more frequently a *secondary* than a *primary* affection has already been noticed, though this has not been true of our experience. The diseases in which the largest number of cases occur are measles, pertussis, and typhoid fever. The secondary cases are most common, of course, during the prevalence of the disease whose progress they complicate, while the primary cases are most common in the cold months of the year, and especially in the autumn and spring. The

reader is referred to the table in the article on pneumonia for a full exhibition of the effects of season and temperature upon the frequency of this disease. Bronchitis is sometimes *epidemic* amongst children as it is amongst adults. It is important also to be aware that there is a strong tendency to attacks of bronchitis in rickety children.

The only *exciting causes* whose effects in the production of the disease seem clearly proved are sudden transitions from a warm to a cold atmosphere, and sometimes the contrary change; prolonged exposure to cold, particularly when combined with moisture; and the inspiration of irritating gases. We believe ourselves, from what we have seen in this city during the last thirty years, that the most fruitful cause of bronchitis, and also of pneumonia, croup, and angina in early life, is the style of dress almost universally used for young children. The dress is entirely insufficient. It consists usually of a small flannel shirt, cut very low in the neck, scarcely covering the shoulders, and without sleeves; of a flannel petticoat, a muslin petticoat, and an outer dress made in nearly every case of cotton. The dress, like the flannel shirt, is cut low in the neck, is without sleeves, and fits very loosely about the chest, so that not only are the whole neck, the shoulders, and the arms exposed to the air, but, in consequence of the looseness of the dress about the neck, it is fair to say that the upper half of the thorax is also without covering. In the infant, from birth to the age of six or eight months, the dress is made long, —a wise provision so far as it goes; but from the time the skirts are shortened, up to the age of four or five years in boys, when happily the time for boys' clothes arrives, and throughout childhood in girls, the trunk of the body and the arms are dressed, or rather left undressed, as above described. But not only are the neck, breast, and arms left bare, but in many children the greater part of the legs also is kept uncovered, or at least, short stockings, scarcely rising above the ankles, and muslin or sometimes Canton flannel drawers, not reaching or scarcely reaching to the knees, leave exposed to the air a large proportion of the cutaneous surface of the lower extremities. Now, in this dress, the child passes the day in a house, the sitting-rooms of which are heated usually to 68° or 70° , but in which the entries, and sometimes the parlors, are frequently at a temperature of 60° , 50° , or even lower, as we ourselves have tested with the thermometer. And not only are the entries and parlors, and indeed all the rooms, saving the one or two in constant use, frequently at the temperature just mentioned, but the air of the nursery itself is often allowed, through the negligence of the servants, and especially early in the morning, to fall to 60° or 58° , or possibly lower still.

That the style of clothing is not correct, is proved by the simple facts that children who are dressed nearly the same in summer as in winter suffer scarcely at all from colds in the summer season, when the thermometer seldom ranges below 76° , and is usually above that point; and also by the fact that adults who have been driven by long and almost forgotten experience to wear clothing twice or three times as warm as that which they put upon their children. How constantly do we see the strong and fully-developed man comfortably enveloped in a warm, long-sleeved flannel

shirt, woollen or thick cotton drawers, and cloth pantaloons, vest, and coat, in the same room and in the same temperature with the little—often puny, pale, and half-naked—child. But it is almost impossible to make people understand that children need as much clothing as themselves. They always insist upon it that, as the child passes the greater part of the day in the house, it cannot require as much clothing as the adult who is obliged to go out and face the weather; forgetting or refusing to see that the former wears less than half, or probably not more than one-fourth, as much covering as the latter, and that the adult, when in the house, and in the same rooms as the child, finds his one-half or three-fourths warmer clothing not at all superabundant or oppressive. It is true, we are happy to believe, that since these statements were written for the early editions of this work a great change for the better has taken place, at least in Philadelphia, in the manner of clothing young children. Most families now dress the young much more wisely than they did twenty or twenty odd years ago, and we feel sure that we see less acute and dangerous lung diseases in early life, in the easy classes of society, than we did formerly.

We have repeatedly had patients to get well of chronic catarrhal and laryngeal coughs, and to cease to have, as before, frequent recurrences of these disorders, under the simple treatment of a long-sleeved and high-necked merino or flannel shirt; long woollen stockings, and stout Canton flannel drawers coming down below the knees; and that, too, after the most patient and assiduous, and sometimes over-assiduous trials of drugs, diet, and confinement to the house had entirely failed of any permanent good effects. The fact is, that though there are some few children who can bear the dress above described without injury, there are a great many more who, while they wear it, either suffer all winter long from frequently repeated attacks of cold, in the shape of croup, chronic laryngeal irritation with cough, chronic pharyngitis, bronchitis, acute or chronic, or more rarely pneumonia; or, if they escape these direct effects, resulting from the constant and rapid waste of their caloric, they are rendered more pale, thin, and delicate-looking than they would be were their vital forces husbanded by warm clothing, instead of being wasted in the constant struggle to keep up the heat of the uncovered body at the natural point.

ANATOMICAL LESIONS.—We shall describe, first, the lesions met with in cases in which the disease is confined to the large bronchi, the inflammation not extending into the capillary tubes; and next, those observed in cases in which the disease has attacked the capillary bronchi. The former are those which constitute the form designated under the title of acute ordinary bronchitis of moderate severity, while the latter are those to which the term capillary has been applied. Patients seldom die of the first-named variety of the disease alone; but as it often occurs as an accidental complication, or a more or less essential part of different severe and frequently fatal diseases, the morbid alterations which characterize it have been very thoroughly studied and ascertained.

The morbid alterations of *acute ordinary bronchitis* always exist in both lungs, and are confined to the larger bronchi, ceasing on a line with the smaller tubes and the capillary divisions. The most constant alteration

is redness of the bronchial mucous membrane, caused by injection of the minute vessels of that and the subjacent tissues, and varying in shade from a rosy to a bright-red or brownish tint. The mucous membrane is sometimes softened, a change which can be ascertained only in the largest tubes, and it sometimes presents a thickened, unequal, and rough appearance. Ulcerations are very rare. The inflamed bronchi contain a more or less abundant viscid, transparent, or opaque yellowish mucus.

In *capillary bronchitis* the alterations of the mucous membrane of the capillary tubes do not always reveal the existence of the disease. That membrane is sometimes pale in the minute ramifications, and exhibits morbid changes only in those of medium size. The alterations of the membrane consist in redness, which is made up either of a number of fine points, seated in the membrane itself, or of arborizations seated both in the membrane and the cellular tissue beneath; it sometimes presents a granulated appearance, and it may be more or less thickened, and its consistence diminished. The bronchi are usually filled and almost obliterated from the secondary divisions to the final ramifications, by a substance of a yellowish-white or yellow color, non-aerated, and composed of a thick muco-pus. Portions of false membranes are sometimes, not as a rule, but exceptionally, found mixed with the secretions just described, while in other instances false membranes alone are present in certain tubes. The false membrane may exist in the form of patches, or it may constitute a lining to the whole extent of the bronchial ramifications. It is usually soft and but slightly adherent, and the mucous membrane beneath is either very pale, and of its usual consistence, or red, softened, and rough. The different kinds of secretion are commonly most abundant in the bronchi of the inferior lobes.

In a good many of the cases another lesion, dilatation of the bronchi, is also found upon examination. This alteration evidently occurs under the influence of the inflammation; it may affect either the length of the air-tubes, or only their extremities. In the former condition the tube continues of the same size, or becomes gradually larger from one of its early subdivisions until it reaches the surface of the lung. In the latter condition a section of the lung presents an areolar appearance, from the presence of a multitude of little rounded cavities, communicating with each other and with the bronchi, of which they seem to be a continuation. These cavities are generally central, though they are sometimes found upon the surface of the lung, in which case they are formed of the pleura, lined by the thinned membranes of the dilated bronchus.

The fact of these cavities being true dilatations of the bronchi, has been called in question by Dr. Gairdner (*loc. cit.*, p. 76), who believes, on the contrary, "that almost all the so-called bronchial dilatations, and all of those presenting the abrupt, sacculated character here alluded to, are in fact the result of *ulcerative excavations* of the lung communicating with the bronchi." He supposes them to be the result of the expansion of certain small cavities, frequently met with in the bronchitis of children, and to be described directly under the title of vacuoles or bronchial abscesses, either by ulceration or by the act of inspiration.

In addition to the lesions already described as existing in bronchitis, there is another one, not unfrequently met with, to which we shall call attention, that to which the French writers apply the term *vacuoles*, and which Dr. Gairdner designates as *bronchial abscess*. The latter author states that in the centre of the collapsed lobules of a lung affected with acute bronchitis, there are found, not unfrequently, small collections of pus, varying in size from that of a hemp-seed to double or treble that volume. "These small abscesses present, on section, an appearance so much like that of softening tubercles, as to be very readily mistaken by many persons for these bodies; and the resemblance is all the greater on account of the peculiar limited form of the condensation by which they are generally surrounded, which, when felt by the touch from the exterior of the lung, is exceedingly deceptive. In their interior, however, these little abscesses contain, in the recent state, a very fluid pus; moreover, they are often met with as acute lesions produced by a few days of illness, and without a trace of tubercle in any other organ." When the pus is scraped or pressed out of these abscesses, in their recent form, they are found to be lined with a fine villous membrane, while in other instances they are not abruptly limited, but the pus appears to lie in contact with the surrounding pulmonary tissue. The bronchi leading to the part of the lung thus affected, are found, when incised, to be much inflamed, their mucous membrane being vascular, thickened, and covered with pus; and some of them can be observed to communicate with the purulent collections, the mucous membrane having been, at the point of communication, destroyed by ulceration, and either stopping short abruptly, or becoming gradually incorporated with the false membrane lining the abscess. Sometimes these abscesses or vacuoles communicate not only with the bronchi, but also with each other, without difficulty; sometimes, according to Dr. Gairdner, they break into one another and form more considerable excavations, but, more commonly, they remain of limited size, preserving perfectly the direction and relations of the bronchial tubes. They occur both in the diffused and lobular form of condensation from collapse of the lung, and both forms may sometimes be seen in the same lung.

The alteration just now described has excited a good deal of discussion amongst medical writers, and has been very differently accounted for. MM. Rilliet and Barthez regard it as a simple terminal dilatation of the bronchi, while MM. Barrier, Legendre, and Bailly, consider it to depend on a purulent breaking down of the vesicles of one or more lobules. MM. Hardy and Behier look upon it as a lesion of a complex nature, partaking both of dilatation of the bronchi and of pulmonary emphysema. Dr. Gairdner, as already mentioned, describes them as abscesses, and states that they "unquestionably arise from the accumulation of pus primarily in the extreme bronchial tubes of the collapsed lobules." This view, which is closely similar to that of MM. Barrier, Legendre, and Bailly, is, it appears to us, much the most reasonable that has been adduced.

MM. Rilliet and Barthez, in their second edition, as has already been stated, in the article on post-natal collapse, describe at great length a state

of *congestion* of the lung-tissue, as a most important element in the anatomical alterations of the bronchitic diseases. This congestion usually assumes one of two forms: it may be distinctly lobular, consisting then of disseminated patches, or, as more generally happens, large numbers of contiguous lobules are affected, when it takes the form of generalized lobular congestion. These congested portions of the lung are almost always attended with more or less well-marked collapse of the vesicles, so that there are associated together the conditions of congestion and collapse. It is this combination of bronchitis with congestion and collapse, which was formerly described by them under the titles of lobular and generalized lobular pneumonia. The alteration to which the term *carnification* has been applied, and which not unfrequently coexists with bronchitis, they regard as different from the above, and as consisting in a simple collapse of the lung-tissue, without the active or passive congestion which exists in the first form. The principal causes of this condition are, according to them, debility and catarrh. The signs of catarrhal inflammation are, they state, scarcely ever absent. In only four out of thirty-one cases did they fail to discover them. We have dwelt, in our article on pneumonia, on the lesions of catarrhal pneumonia, or lobular pneumonia, as it is now generally recognized; and it is probable that at least in some of the cases described as above by Rilliet and Barthez the condition has really been of that form of inflammation of the pulmonary tissue.

The parenchyma of the lung presents, in bronchitis, different appearances in different cases. It is supple, crepitant, and of a rose-gray color, but does not collapse, especially the anterior portions, when the thorax is opened, as does healthy lung. This imperfect collapse depends either on the fact that the thick mucus and muco-pus which fill and obstruct the bronchi prevent the contained air from being expelled by the natural elasticity of the lung, or, when no secretions exist to produce this effect, on the loss of the natural elasticity of the organ. Another cause is the existence of vesicular emphysema, a lesion observed to a greater or less extent in nearly all the cases, and affecting usually the summit of the lung, its anterior edge, and also its posterior or lateral edge. In a large number of cases, and particularly in those occurring in young children and in weakly and debilitated subjects of all ages, the tissue surrounding the diseased bronchi exhibits the condition which has already been fully described in the article on atelectasis, under the title of collapse of the lung. The extent and mode of distribution of this lesion, its peculiar and distinguishing characters, its causes and mode of production, and the method of treating it, have been carefully discussed in the article just referred to, and we shall make no further allusion to it, in this place, except to beg the reader, who is not already fully acquainted with it in all its bearings, not to suppose himself master of the subject of bronchitis until he has also fully studied that of collapse, as the two go together so constantly, and the latter is practically so important, especially in children, as to make it essential for him to understand both.

The lesions just described as characteristic of acute bronchitis are also met with in the *chronic* form of the disease. The dilatation of the air-

tubes, however, presents different features. The calibre of the enlarged tube is often much greater, its walls are whitish and uneven, and beneath the mucous lining may be seen hypertrophied transverse fibres. The mucous membrane itself remains smooth and polished, while the tissues beneath are thickened and hypertrophied.

SYMPTOMS; COURSE OF THE DISEASE; DURATION.—Acute simple bronchitis exhibits very different degrees of severity in different cases, being in some extremely mild and benign, and in others so much more severe as to border closely on the capillary form of the disease. In its mildest form, it occasions merely slight cough and stuffing, a few mucous râles over the larger bronchi, with a total absence of dyspnœa, or of decided fever. In cases rather more severe than this, it begins with a moderately frequent cough, which, dry at first, soon becomes loose, and is neither paroxysmal nor painful. The expression of the face remains natural, with the exception of an appearance of slight languor. The pulse and respiration are but slightly accelerated; the external phenomena of the latter, an important means of diagnosis in infants, remains natural; it occurs without jerking, the rhythm continues even and regular, and there is no violent action of the *alæ nasi*. The percussion is not modified. Auscultation reveals in very young children a mixture of mucous and sibilant râles on both sides, which come and go, and are of short duration; in older children, the moist râles predominate, and commonly last several days. These sounds are seated in the larger bronchi. The temper of the child is not much changed; the appetite is not entirely lost; there is neither vomiting nor diarrhœa; and the fever is usually slight. The disease remains nearly stationary, or increases for a variable length of time, after which the cough becomes looser, and in children over five years of age is sometimes attended with expectoration of frothy or yellowish mucous sputa, whilst under that age there is no expectoration. The fever and other symptoms, with the exception of the cough, now subside; the cough remains some days longer.

In attacks still more severe than this, the symptoms resemble very much those just now described, but they are all more intense. The cough is tighter, more frequent, harassing, and especially it is more painful, as shown by the fact that the child cries and complains, and that a marked expression of pain passes over the face at the instant of coughing. There is more fever, the skin being hot and dry, and the pulse more frequent, rising often to 130 or 140, and in one case to 156. The respiration is hurried, and, though not attended with the same labor and anxieties as in the capillary variety, it is evidently oppressed; it counted in three cases 60, 60, and 62. The temperature is considerably elevated, but not so much so as in pneumonia, rarely rising above 102° or 102.5° . There is more restlessness, fretfulness, and general distress; the appetite is greatly diminished or lost, and infants nurse with less avidity than usual, or refuse to nurse at all for several hours together. In cases of this kind, the physical signs are more decided than in those of milder degree, there being a greater abundance of mucous and dry râles, and generally some subcrepitant râles, and they are heard over a larger extent of surface, usually over the lower half, two-thirds, or even the whole dorsum of the chest. The symptoms are almost

always most marked and severe in the after-part of the day and night. Very often the patient will be comparatively easy and comfortable in the morning, but as the day goes on, he becomes more feverish, restless, and fretful; the cough grows more troublesome, more frequent, and tighter; the breathing is quicker and more oppressed; the face is more flushed; the sleep is broken and disturbed, and the child may appear through the night quite ill, and yet, as morning approaches, the symptoms moderate, the skin often softens and becomes moist, and the whole aspect of the case shows a great amelioration in the manifestations of the disease.

According to Handfield Jones, this almost invariable tendency to aggravation of catarrhal disorders during the night is due to a lowering of the nerve-power, the vaso-motor nerves partaking of the general debility, and thus allowing dilatation of the arteries, and causing increased hyperæmia of the affected parts with more abundant exudation.

The *duration* of this form of bronchitis is very uncertain; the idiopathic cases last usually from four to seven or eight days, though they may last from sixteen to twenty-five; the duration of the secondary cases depends, in great measure, on the nature of the diseases during which they occur.

In any of these different degrees of acute simple bronchitis, the patient is liable, especially if it be a weak and debilitated child, or a young infant, to sudden and alarming aggravations of the symptoms. The breathing becomes suddenly either greatly increased in frequency, or excessively labored and oppressed; the surface becomes pale, the expression dull and languid, or distressed; the child is drowsy and inattentive, or uneasy and restless; the hands and feet are cool; the act of sucking is performed with difficulty, or the child refuses the breast entirely, and it is evident that, from some sudden change in the condition of the lungs, the act of respiration and the aeration of the blood are very seriously interfered with. If this sudden aggravation of the symptoms be unattended with a corresponding increase of the febrile phenomena, as marked by greater heat of skin and augmented action of the circulation, it is altogether probable that it depends on a collapse of larger or smaller portions of the pulmonary texture, and if, on examination, we discover dulness on percussion, distant bronchial respiration, and cessation or greatly diminished abundance of the bronchitic râles, over parts of the chest where a few hours or a day before there had existed all the physical signs of bronchitis, there can be no longer any doubt as to the cause of the suddenly increased severity of the symptoms—it must be owing to collapse.

It must, however, be carefully borne in mind that it is in the course of bronchitis that catarrhal pneumonia is most apt to occur. The symptoms which indicate this accident have been detailed in the article on pneumonia (page 172), the most prominent being the rapid increase in dyspnœa, the sudden and marked elevation of temperature, the change in the character of cough, the evidences of pain in the chest, and the absence, in most instances, of any positive physical signs of consolidation of lung-tissue. If these symptoms be contrasted with those above stated, as indicating the occurrence of collapse of the lung, it will be seen that with care any error in diagnosis may be avoided.

Capillary bronchitis, or *acute suffocative catarrh*, may succeed to the form just described, or appear as an idiopathic affection. Under either condition the general symptoms are more threatening than in the preceding form, and the disease soon assumes all the appearances of great severity. The child is very uneasy and restless, constantly changing its position, moving about in the crib or bed, or insisting upon being changed from the bed to the lap, or from the lap to the bed. In one case that came under our charge the oppression was very great, and the only position in which the child was at all satisfied was resting on the mother's arms, with the front of its chest applied against her breast, and the head hanging over her shoulder. The expression of the face is anxious and disturbed, and its color usually pale or slightly bluish. The temper is irritable or subdued; the child hates to be disturbed, and generally chooses its own position. The respiration is very much accelerated, running up in a very short time to 60, 70, or 80, and is usually more or less irregular, and evidently laborious and difficult. The cough is very frequent, troublesome, and evidently painful; it occurs in short paroxysms usually, with or without stridulous sound, is at first dry, and after a few days is accompanied, in older children, by whitish or yellowish expectoration. In some instances, the sputa consist of mucus tinged with blood, or of pure blood even, and still more rarely of mucus mingled with small shreds of false membrane. The appetite is entirely lost; the tongue is usually moist and furred white; there is acute thirst, and yet, in severe cases, though the presence of acute thirst is evident from the manner of the child, only very small quantities of water are taken, from the impossibility of suspending the respiration long enough to allow of more being swallowed; the drink is gulped rapidly, suddenly, and with great difficulty, and after a time is refused almost entirely from this cause. In children old enough to talk, the speech is short and abrupt; the patient dislikes to speak, from the fact that the effort obliges him to suspend momentarily the act of breathing. Fever sets in from an early period; the skin is hot and dry, and the face is flushed at first, though it soon becomes pale in most cases, from the approach of an asphyctic state. The pulse becomes frequent, rising soon after the onset to 130, 140, 150, or higher; it is full and hard early in the attack. The resonance on percussion is not modified. Auscultation reveals at first sibilant râles mixed with some mucous râles; but soon a fine subcrepitant râle is heard over all the lower parts of both lungs behind, and approaching sometimes, over the bases of the lungs, the character of crepitus. After a time the subcrepitant râle is heard over the whole, or nearly the whole, dorsum of the chest, and to a greater or less extent, though not so well marked as behind, over the anterior regions of the thorax. This râle is audible at first both in inspiration and expiration, and is very distinct, but at a later period it is heard only in the inspiration, or there is substituted for it a mucous râle, while the subcrepitant râle is now heard only in the forced inspirations during coughing or crying. These râles are fugitive and irregular, disappearing or changing from one to the other after fits of coughing.

Should the case not take a favorable turn, which change would be indi-

cated by a moderation in the symptoms just detailed, and especially by easier and fuller respiration, with diminution of the amount of the subcrepitant râles, and return of the natural respiratory murmur over some parts of the chest, the symptoms look still more alarming. The oppression becomes excessive: fits of dyspnœa occur, in which the child is extremely restless and distressed, tossing itself about on the bed; the respiration runs up to 80, 90, or more, in the minute, and is attended with violent action of the alæ nasi; the pulse grows more and more frequent, rising to 150 or 180, and it loses force and volume; and the face assumes a whitish or slightly bluish tint, looks puffed, and is sometimes covered with perspiration. As the fatal termination approaches more nearly, the pulse becomes small, thready, and irregular; the respiration is uneven, irregular, stertorous, and often slower than before; the cough is smothered and less frequent; the restlessness generally diminishes, and the child sinks into quiet, and often becomes comatose; the paroxysms of suffocation are less frequently renewed, and less violent, and death occurs in a state of quiet insensibility, or is preceded by partial or general convulsive movements.

The *duration* of this form may be stated to be, on the average, between five and eight days. It may, however, end fatally in a much shorter time. In an example that we saw, in a child four months and a half old, death occurred in twenty-six hours from the onset. Dr. Eberle states that it seldom lasts longer than two or three days, and that in very young infants death sometimes occurs on the first day. M. Bouchut gives as the duration in children at the breast, from two days to a week. Dr. West mentions a case that proved fatal in less than forty-eight hours. In the favorable cases that we have seen the duration was seven, eight, and ten days.

Subacute and chronic bronchitis, generally follows one of the acute forms of the disease. The character and severity of the symptoms vary very much in different cases. We have known some children to present for several months together, in the winter season, slight bronchitic symptoms, consisting in wheezing and somewhat accelerated breathing, cough, more or less frequent; occasional feverishness, especially at night; some diminution of appetite and loss of flesh; and sibilant and sonorous with mucous râles, heard here and there, but still without severe symptoms during the greater part of the time. Children laboring under this kind of bronchitic irritation are liable to, and generally have, from time to time, more or less sharp attacks of acute bronchitis, in which they present the usual symptoms of that form of the disease. These attacks are very apt to occur coincidentally with changes in the weather, and in some patients the liability to them is so great, from the excessive susceptibility of the system to the weather, that no care will prevent them. In some instances, we are very sure that an aggravation of the symptoms of the chronic form constantly occurs whenever the child is about cutting additional teeth, whilst in the intervals between the appearance of the successive teeth, the child remains comparatively well. We believe that the cause of the aggravation, at the moment of cutting the teeth, is to be looked for, not only in the act of dentition itself, but in the circumstance that the liability to cold is greatly increased at that particular moment, probably because the forces of the system are

so weakened by the effort of the dentition as to lessen the power of resistance against the disturbing influence of a changing, and particularly of a falling temperature.

Cases of the mild kind of chronic bronchitis that we have just been describing, usually get well under proper medical, and especially under proper hygienic means, after several weeks or two or three months; while in other instances the disorder continues, in spite of every precaution, throughout the winter and spring, and only ceases as the warm summer months arrive. We have known the same disposition to show itself again in the following winter. In other instances again, the frequent attacks of severe bronchitis, together with the effect of a constant slight bronchitic inflammation, ends in the production of an emphysematous state of parts of the lung, and the child exhibits more or less marked asthmatic symptoms, which show themselves whenever a slight increase of the bronchitis occurs, and whenever the digestive system is deranged by imprudence in diet or other causes. It is particularly in such cases as these that the bronchial affection is apt to be associated with rickets, and we should, therefore, always carefully search for the evidences of this latter disease.

In other examples of chronic bronchitis the symptoms are much more severe. These cases almost always follow an acute attack of the disease. The frequency of the respiration and the attacks of dyspnoea persist; the cough is loose and paroxysmal, and the pulse frequent and small; evening exacerbations of fever take place, and the face and sometimes the rest of the surface are often covered with perspiration. Auscultation reveals tubal blowing with mucous or loud sonorous râles, which seem to indicate the presence of dilatation of the bronchi. Emaciation makes rapid progress, the face is pale and blanched, the eyes sunken, the nostrils are covered with mucous or bloody crusts, and the lips ulcerated. Strength diminishes progressively; the appetite is lost, and the thirst acute; colliquative diarrhoea appears; and after twenty, forty, or more days, the child perishes in the last stage of marasmus. This form of bronchitis often simulates phthisis very closely, and may last for a long time, even several years. It rarely occurs under the age of five years. The expectoration consists of purulent or pseudo-membranous secretions in variable quantity.

PARTICULAR SYMPTOMS—PHYSICAL SIGNS.—The *dry râles* are amongst the most frequent physical signs in bronchitis. They may be sibilant or sonorous; they seldom exist alone, but are accompanied with mucous râles, and diminish as the latter become more abundant. As the dry râles cease to be heard, they are replaced by mucous or subcrepitant râles, or by feebleness of the respiratory murmur. The sibilant râle is often heard over the whole thorax, though it may be confined to the posterior portions. It is not restricted to cases of inflammation of the larger bronchi only, but is also present in capillary bronchitis.

Moist Râles.—Mucous and subcrepitant râles do not exist in all cases without exception, as they may be absent in such as are very mild. They may generally be heard over both sides behind, more rarely over the whole of the chest, and almost always both in inspiration and expiration. They

are generally persistent, but are sometimes suspended for a moment and replaced by sibilant râles or feeble respiratory sound.

Feeble respiratory murmur is sometimes observed. It is not permanent, occurs during the interruptions of the subcrepitant or sonorous râles, and does not occupy the whole extent of the thorax, but is limited; it is intermittent, and is not accompanied by diminished resonance.

When dilatation of the bronchi exists to a considerable extent it gives rise to bronchial or even cavernous respiration, and to bronchial resonance of the voice, cry, and cough. The bronchial respiration differs from that of pneumonia by its tone, and by its intermitting character. The *percussion* is generally sonorous.

It has already been stated in the account of the symptoms that it happens not infrequently in severe bronchitis, and also in mild bronchitis occurring in debilitated children, that the respiratory sound suddenly becomes feeble, or even entirely suppressed, over parts of the lung, while in other instances a distant and imperfectly marked bronchial respiration takes the place of the natural vesicular murmur. These changes are heard either over small disseminated points of the lung, or over large surfaces; they are associated with more or less evident dulness on percussion, and what particularly characterizes them, they are very fugitive, being present at one examination, and absent perhaps at the next. The appearance of these changes in the phenomena afforded by auscultation depends on the occurrence of diffused or lobular collapse of the tissue of the lungs.

The physical signs above described are not invariably present in bronchitis. Cases do occur, though they are very rare, in which auscultation fails to reveal the characteristic signs of the disease.

RATIONAL SYMPTOMS.—The rational symptoms are of the utmost importance in informing us of the degree of severity of the attack.

Cough generally exists from the beginning, being in mild cases more or less frequent, and either dry or loose, while in severe cases it is frequent or very frequent, at first dry and then moist, and very rarely hoarse. In acute capillary bronchitis, the cough has a peculiar character. From the first day it occurs in short paroxysms, lasting from a quarter to half a minute. The paroxysms vary greatly in violence, occur at irregular intervals, and generally continue without interruption to the fatal termination, though they are sometimes replaced by simple loose cough a few days before that event. The cough is rarely painful, so long as the inflammation remains simple. Expectoration is never present in very young children. When it occurs in those over five years of age, it consists, in the mild form, of a sero-mucous or of a frothy and yellowish mucous liquid. In general bronchitis it is sero-mucous at first, becoming after a few days yellowish and more or less viscous; it is sometimes nummular and sometimes amorphous.

In the capillary form, as already mentioned, the sputa consist of mucous tinged with blood, or of pure blood even, and in some rare cases there are mixed with the mucus, shreds of false membrane, which may present the form of casts of the minute ramifications of the bronchial tubes.

The *respiration* varies in its characters according to the extent and vio-

lence of the disease. In mild cases, it is not much increased in frequency, being generally between 28 and 40 in the minute. In more violent cases, and particularly when the disease implicates the smaller bronchi, it becomes very frequent. The acceleration is slight in the beginning, but increases regularly as the case progresses; thus it may be 30 at first, and rise afterwards to 50, 60, 80, and even 90. When not very much quickened, it remains even and regular; when more so, it becomes somewhat laborious, and the movements of the chest are full and ample; in severe cases, attended with much dyspnœa, it is often irregular, or assumes the characters to which M. Bouchut has applied the term *expiratory*, that is, the order of the movements is inverted, each respiration beginning with the expiration, leaving a pause between the inspiration and expiration, instead of between the expiration and inspiration. In chronic bronchitis with copious purulent or pseudo-membranous expectoration, the dyspnœa is generally habitual.

Fever.—The fever is slight in mild cases, the pulse rising very little above its natural standard. The heat is not great, and the febrile movement usually subsides before the termination of the disease. In the grave or capillary form, on the contrary, the pulse is always frequent, and continues to increase in rapidity as the disease advances. It varies between 104, 120, 160, and in very violent cases, rises as high as 200. Early in the attack, it is vibrating, rather full and regular, whilst in fatal cases, it always becomes small, irregular, trembling, and unequal. The skin is generally hot in proportion to the activity of the pulse, except towards the termination, when the extremities often become cool. The temperature does not rise so rapidly nor reach so high a point as in pneumonia. Thus Roger gives as the highest temperature observed by himself in bronchitis 102.2° ; while the average in his cases of the acute febrile form was 100.9° . The skin is almost always dry. In very young children it is often pale and cold, and covered with perspiration from the beginning.

The *expression* of the face is unchanged in mild cases, but when the disease is violent and extensive, becomes deeply altered after a few days. The eyes are then surrounded with bluish rings, and the expression is uneasy, anxious, and sometimes, but less frequently, exhibits an appearance of profound exhaustion. The anxiety of the countenance increases with the oppression; the *alæ nasi* are dilated, the nostrils dry or incrustated, and the lips and face, which are extremely pale or momentarily congested, assume a purple tint, particularly after the paroxysms of cough.

The *decubitus* is indifferent at first, but as the disease progresses, the child lies with its thorax more or less elevated, or is restless and constantly changing its position.

In dangerous cases there is great *distress* and *restlessness* after the first few days, or even from the beginning. In some instances the irritability and peevishness are excessive and uncontrollable, while in others there is heaviness and somnolence, especially towards the termination of fatal cases. Some of the disorders of the nervous system just mentioned are present in all the grave cases.

Digestive Organs.—There is moderate *thirst* and incomplete *anorexia*

when the disease is mild, but when severe, the thirst is generally acute, and the appetite entirely lost. The state of the bowels varies. The *tongue* and *abdomen* present no special characters in idiopathic cases.

Urine.—The great majority of recorded observations of the condition of this excretion in bronchitis, relate to the disease as occurring in the adult. The following summary is taken from Parkes: the condition of the urine in bronchitis varies greatly with the grade of the disease; in the grave forms, it resembles that of pneumonia, the urea being increased, and the chloride of sodium at times entirely absent. The urine has also been quite frequently found to be temporarily albuminous in such cases.

DIAGNOSIS.—The mild form of bronchitis, in which the inflammation is confined to the larger bronchi, is not likely to be mistaken for anything but the early stage of hooping-cough. The diagnosis can be made only by attention to the different characters of the cough, which is more spasmodic and paroxysmal in pertussis, by the absence of fever in that disease, and by the development of the peculiar symptoms of each, as the case progresses.

The diagnosis between bronchitis and pneumonia is seldom difficult, except when the latter is grafted on the former, or in cases of partial pneumonia attended with bronchitis. In well-marked cases of the two diseases, there can be no difficulty. The restriction of the physical signs to one side alone of the chest in pneumonia, the peculiar crepitus of that disease, or when this is not heard, the fineness of the subcrepitant râles, limited to the upper or lower regions of one lung, the bronchial respiration and bronchophony, the dulness on percussion over the seat of disease, the greater sharpness and severity and the different location of the pain, the more acute character of the febrile reaction, as marked by the pulse, skin, and thirst, the more abrupt and higher elevation of temperature, and the kind of expectoration, when there is any, will always enable us to distinguish the two with almost absolute certainty. In cases, however, in which the two are combined, the diagnosis is not so easy, but even here the presence of dulness on percussion, and of crepitant or fine subcrepitant râles, or, when these are absent, of pure metallic bronchial respiration with bronchophony, over limited portions of the lung, will generally render the matter clear.

The *sudden* supervention of dulness on percussion over large portions of one of the lobes of a lung, or over disseminated patches, with feeble or absent respiratory sound, or with muffled and distant bronchial respiration, generally indicates the occurrence of collapse in the part of the lung over which these signs exist; and when these symptoms show themselves without any increase in the severity of the febrile reaction, but rather with a diminution, there is every reason to suppose that they depend, not upon inflammatory condensation of the parenchyma of the lung, but upon simple collapse, from the presence of obstructive secretions in the bronchi.

Dr. Gairdner (*loc. cit.*, p. 6) has called attention to a difference in the character of the dyspnœa in the two diseases, which is, we think, of considerable importance, and which we have often remarked ourselves. In

bronchitis, of any considerable severity, the respiration is always evidently labored; it is performed only with the aid of all the accessory muscles of respiration, and in really severe cases it is extremely laborious, the inspiration being long-drawn, exhausting, and inadequate. The dyspnœa of *pure* pneumonia is, on the other hand, quite different. It is merely an "acceleration of the respiration, without any of the heaving or straining inspiration observed in bronchitis, or in cases where the two diseases are combined." Dr. Gairdner states that he has repeatedly seen patients affected with a great extent of pneumonia of both lungs, in whom the extreme lividity and rapid respiration, numbering fifty or sixty in the minute, showed infallibly the amount to which the function of the lung was interfered with, who, nevertheless, lay quietly in bed, breathing without any of the violent effort, or disposition to assume the erect posture, so constantly accompanying the more dangerous forms of bronchitis. In children these differences are even more marked than in adults.

Chronic bronchitis may be mistaken for tuberculosis of the lungs or of the bronchial glands. The distinction can be made only by careful study of the history of the case, and of the phenomena afforded by auscultation and percussion, which are detailed in our article on tuberculosis.

It is also important that we should not overlook the evidence, of rachitis, which, as before stated, very often exist in children who are predisposed to attacks of bronchitis.

PROGNOSIS.—Bronchitis is rarely a fatal disease, so long as it remains confined to the larger bronchi, constituting the acute simple form, of moderate severity. Capillary bronchitis is, on the contrary, a very dangerous affection at all times and at all ages. Even ordinary, simple bronchitis, however, may prove fatal in young infants, and in debilitated children of all ages, from the supervention of collapse of portions of the pulmonary tissue; and it is necessary, therefore, that the prognosis given should always be guarded, when the disease occurs under either of these two conditions. The prognosis differs also in the primary and secondary forms of the disease, since, as might be expected, the danger is much greater in the latter than in the former variety.

We have met with a large number of cases of bronchitis, out of which we have kept more or less copious notes of 123. Of these, 108 were mild, and 15 capillary. Of the 108 mild cases, 65 were primary, all of which recovered; and 43 secondary, of which 2 died. Of the 15 capillary cases, 11 were primary, of which 1 died, and 4 secondary, of which 2 died. Of the whole number, 123 in all, 5 proved fatal. The danger from the disease depends very much also upon the hygienic conditions in which the patients are placed. In hospitals and amongst the poor it is much more dangerous than in private practice amongst the easy classes* of society. This is shown by the fact that all the cases of the capillary form observed by MM. Rilliet and Barthez and Fauvel, in hospital practice, proved fatal, while of 15 cases seen by ourselves, in private practice, under the most favorable hygienic conditions, only 3 died.

The symptoms indicating great danger are, increase of the dyspnœa;

extreme anxiety, small and irregular pulse, coolness or coldness of the skin with clammy sweats, much jactitation, and delirium, drowsiness, or coma. With such symptoms the danger is greater and the fatal termination more imminent in proportion as the child is younger, less robust, and its constitution exhausted by preceding or coincident disease.

TREATMENT.—The acute simple form of bronchitis is frequently so mild as to need no other treatment than careful attention to the hygienic condition of the patient, and the administration of some simple febrifuge and expectorant. The child ought to be confined to one room, in a mild and uniform temperature, and should be kept quiet until the development of the symptoms shows what is to be the type of the attack. The degree of repose of the body necessary will depend on the presence or absence of fever. We believe that the practice of keeping the body quiet in all febrile disorders, is one of the most important therapeutic means we have. It is long since one of us, having seen his father insist upon putting children to bed for a feverish cold, began to follow the same practice. Time and experience have made even more clear to us the wisdom of the practice, especially in regard to very young children.

So soon as the attack of bronchitis becomes severe enough to cause fever, whether the fever be continuous or occur only in the afternoon and night, the patient ought to be confined to the lap or bed. Suckling children, and those under three years of age, must be allowed to lie on the lap at times, but even they may be taught, very early, to rest quietly in the crib the greater part of the twenty-four hours. Children over three years old can almost always be taught to stay in their beds by a little management and authority, if only the parent is resolute. If not very sick, they should have a large pillow put up against the head of the crib or bed, and against this they should be placed in a sitting posture, with the bedclothes arranged over the lap; and, in cool or cold weather, with a light flannel sack over the night-dress. Here they ought to be kept all day, allowed to change their position as they wish, and they should be kept as cheerful and happy as possible with toys, books, pictures, readings, tale-telling, or what not. Under such circumstances, a new and interesting toy will often do more good by far than any drug in the *materia medica*. We have often been surprised, and delighted, too, to find a bronchitis which had been hanging over a young child for several days or a week, getting gradually worse, day by day, under the trotting-about system, begin to mend from the day the child was put to bed, and disappear in two or three days, and that, too, without any change in the other remedies.

The clothing ought to be warm, and yet not sufficient to produce free perspiration, as this, by sudden exposure and evaporation, often induces chilliness. The diet must be simple, and may consist of any of the milk preparations, with or without bread, or bread and butter. Light soups in the middle of the day, or roast potatoes or apples, with bread, may generally be allowed.

As for medicines, in this mild form they are of comparatively little consequence, if the above measures be carried out. In the after part of the

day, when fever sets in, we may prescribe a febrifuge of citrate of potash, such as the following, for children of two to four years old:

R. Potass. Citrat.,	3j.
Syrup. Ipecac.,	f 3j vel 3ij.
Tr. Opii Camph.,	f 3j vel 3ij.
Syrup. Simp.,	f 3 ss.
Aquæ, q. s. ad	f 3 iij.—M.
Dose. A teaspoonful every two or three hours.	

This should be given until the child sleeps, and occasionally in the night if there be cough and restlessness. At six months of age, the following may be used in the same manner:

R. Syrup. Ipecac.,	
Tr. Opii Camph., āā	f 3 ss.
Spts. Æther. Nitros., vel	
Liq. Ammon. Acetat.,	f 3 ij.
Syrup. Simp.,	f 3 v.
Aquæ,	f 3 iij.—M.
Dose. A teaspoonful every two hours.	

If the fever is very slight, and the cough only moderately severe, it is often well to use no drug through the day, but to give in the evening, two hours before bedtime, and again at bedtime, some simple expectorant and anodyne. Thus at two or three months of age, three to five drops of syrup of ipecacuanha with five of paregoric, or half a drop to a drop of laudanum; at one or two years, ten drops of the syrup with ten to twenty of paregoric, or two of laudanum; at five to ten years, ten to twenty drops of the syrup, with twenty to thirty of paregoric or four or five of laudanum. The laudanum is often better than paregoric, as it produces a more decided and lasting impression on the nervous system, and appears to extend its useful control over the symptoms further into the following day.

In this very mild form there is no necessity for giving active purgatives. If the bowels are moved once in the day, or once in two days, it is best not to interfere with them. If, however, the patient be constipated, a little simple syrup of rhubarb, a teaspoonful of castor oil, or an enema, will be quite sufficient. A warm foot-bath, in the evening, containing salt, or better, mustard, will often assist in moderating the cough and promoting quiet sleep.

When in this acute form the symptoms assume greater severity, when signs of reaction are prominent, the dyspnœa considerable, and the cough frequent and harassing, it was formerly quite the custom to employ depletion. In a former edition of this work, it was stated that the abstraction of a few ounces of blood by leeching or cupping was allowable under these circumstances, but that a large majority of such cases would do perfectly well without bloodletting of any kind. We now believe that such practice is unnecessary in any of this class of cases. Attention to hygienic measures is, however, even more important than in the milder cases. Confinement to the bed ought to be a positive rule in such cases. If the bowels are not freely moved, a dose of castor oil, rhubarb, or magnesia

should be given, and the patient then put upon the use of one of the febrifuge mixtures recommended above.

If, as the case progresses, the bronchial secretions become very abundant and the dyspnœa severe, the proper remedy is an emetic. This may be ipecacuanha, either in powder or syrup, or a teaspoonful of powdered alum, to be repeated if necessary, in ten or fifteen minutes. The latter substance is, as we have stated under the head of croup, a very certain, efficient, and safe emetic.

Great benefit may be obtained in all forms of bronchitis, from the more or less frequent application of mustard poultices to the front or back of the thorax, and from mustard foot-baths.

The mercurial preparations, so much recommended by many of the English and by some of our own writers, are, in our opinion, very seldom, if ever, necessary in this, or indeed in any of the forms of bronchitis in children. It may be, however, that the occurrence of gastric disturbance with coated tongue, anorexia, and a torpid state of the bowels, may, in some cases, call for the administration of a single dose of blue mass, followed by a mild saline laxative.

MM. Rilliet and Barthez recommend, when the cough and sibilant râles persist after the disappearance of the febrile symptoms, the use of small doses of the flowers of sulphur. We have ourselves known this remedy to prove of service in such cases. About four grains may be given every three hours to a child four years old.

The treatment of the *grave acute* or *capillary form* of this disease brings up again the question of bloodletting. We, like all the rest of the world, have abandoned the practice as a rule, but we think that when, in a case of the kind now under consideration, the age being over two years, the oppression is very great, the right heart laboring, as shown by a congested surface and a throbbing cardiac impulse at the base and left edge of the sternum, and the strength not too much reduced, the abstraction of from two to four ounces of blood from the interscapular space by cups or leeches, would be a useful and legitimate practice. We venture to give this advice from our own past experience, and from the views taught quite lately as to the effect and value of depletion in relieving the over-distended right heart, produced by an obstacle to the pulmonic circulation.

There is no occasion for repeating here what has been said, under the head of pneumonia, in regard to tartar-emetic. But if the temperature be very high, and the pulse full and strong, we believe that the small doses of sulphurated antimony (gr. $\frac{1}{12}$ th we then recommended, in combination with Dover's powder, every two or three hours), are very useful in moderating the inflammatory symptoms. Should this be followed by nausea or vomiting with exhaustion, they must be suspended at once. The physician, and especially the young and inexperienced one, ought to know that the susceptibility to the action of all antimonials is singularly different in different individuals. We have seen a hearty adult woman thrown into a most violent, and for a time alarming choleraic condition, by two doses of $\frac{1}{12}$ th of a grain of tartar-emetic each. We once saw a fine hearty boy, five years of age, vomit violently, grow pale, weak, and faint away,

from two teaspoonfuls of the *mel. scillæ compositum*, containing in the two doses the fourth of a grain of tartar-emetic. And even twelfths of a grain of the sulphurated antimony will sometimes cause a degree of nausea and prostration in young children which ought not to be kept up, though we never saw it occasion such effects as those just mentioned as following the use of tartar-emetic. When, therefore, the sulphurated antimony acts with any undue violence, it ought to be stopped, and we should substitute the citrate of potash mixture proposed for the mild form of bronchitis.

In connection with one of these internal remedies, counter-irritation to the surface of the chest will be found of very great service. Indeed, we doubt very much whether it is not the most important part of the treatment. It may be obtained by applications of dry cups to the back of the chest, or if this be inconvenient or objected to for any cause, by the use of mustard poultices. The poultice ought to be about the size of the hand, or one-half larger, and it should be made of one part mustard to two of Indian meal or flour. It is to be mixed with warm water, covered with book muslin or cambric, and applied first to the dorsum of the chest; after having reddened at that point, it should be shifted to the front of the thorax. The time necessary for each contact is usually from ten to fifteen or twenty minutes. These applications ought to be renewed once in four hours, when the symptoms are only severe, but when these are urgent they should be made every two hours. We are in the habit of depending very much also on mustard foot-baths. When the oppression is severe, and especially when there is any coolness of the extremities, the use of a foot-bath simultaneously with the mustard poultice will often assist very much in relieving the breathing.

In very young infants, antimony ought not to be employed, in our opinion, and in these, therefore, we need some other remedy. In them *ipecacuanha* is much safer than antimony, and it is quite active enough. The best preparation is the syrup, of which from three to five drops may be given every two hours to infants six months old. In older children, also, in whom we have been obliged to suspend the antimony, and in those in whom its use has been contraindicated by delicacy of constitution or by feeble health, the *ipecacuanha* is preferable. The doses must vary with the age. At five years, about ten drops every two hours, in combination with the same quantity of spirit of nitrous ether, is a proper dose. When the child presents a pale surface and a languid expression, and particularly when the skin is very slightly warmer than usual, or cool, the following prescription has proved a most useful one in our hands:

R. Liq. Ammon. Acetat.,	f ʒss.
Syrup. Ipecac.,	f ʒj.
Liq. Morph. Sulphat.,	gtt. xl.
Syrup. Acaciæ,	f ʒj.
Aquæ,	f ʒjss.—M.

The dose of this is a teaspoonful for a child two years old, to be repeated every two hours. Should there be any nausea present, the syrup of ipe-

cacuanha ought to be reduced to half the quantity; and if there be any drowsiness, the morphia must be left out.

In very severe cases of the disease, in which the dyspnœa is excessive, the pulse rapid and small, the skin cool and pale, the jactitation very great, and when there are present extensive mucous and subcrepitant râles, the treatment generally recommended is the frequent employment of emetics, and the French authors usually prefer tartar-emetic. For our own part, we would not venture to administer, under such circumstances, so powerful a remedy, and especially so potent a sedative, as antimony, one that we have so often known to cause alarming and dangerous prostration in children laboring under much slighter disorders than suffocative bronchitis. If any emetic be given, it ought to be one of milder action and less perturbing influence than tartar-emetic, and we should choose, therefore, either ipecacuanha or alum. The plan of treatment we prefer, however, is to make assiduous use of counter-irritants, and to give internally the spirit of Mindererus and a weak decoction of seneka; or we may combine with the decoction of seneka, in a suitable form, small doses of the muriate or carbonate of ammonia. Depletion is, in these cases, entirely contra-indicated; we may, however, with advantage apply a few small dry cups to the dorsum of the chest in the interscapular space, or over the lower lobes of the lungs.

In the bronchitis of children it often becomes proper and necessary to make use of *stimulants*. In the suffocative form, when the symptoms assume the character described in the last paragraph, small doses of brandy or wine-whey may be administered alternately with the spirit of Mindereus, with great advantage. In milder cases, also, when a sudden increase of the dyspnœa occurs, especially in feeble and debilitated subjects, and when we may suppose, from the character of the rational and physical signs, that collapse of portions of the lung has taken place, it is best to abandon for the time all nauseating remedies, and to make use simply of brandy in doses of from five to twenty drops every half-hour or hour, or wine-whey in dessert or tablespoonful doses, and of counter-irritants, with very light fluid nourishment.

In cases where there is such marked debility, tonics are very useful, and good results may be obtained from the administration of quinia, which was strongly recommended a few years ago, in the form of capillary bronchitis occurring in tropical climates, by Dr. Cameron (*London Lancet*, November 9th, 1861).

In cases of this kind, we have used with great advantage of late years small doses of quinia, prepared as follows:

R. Quiniæ Sulphat.,	gr. vj.
Acid. Sulph. Dil.,	gtt. xij.
Syrup. Simp.,	f 3 ss
Aquæ,	f 3 ijss.—M.

Dose. A teaspoonful every two hours, to children two or three years old.

In older children the proportion of quinia to the dose ought to be doubled.

If this should sicken, as it will sometimes do by the disgust its bitterness produces, and the consequent resistance to the doses, it is best to lay it aside after two or three trials, and to administer the quinia in the form of powder mixed with a little extract of liquorice and sugar, or to substitute the following :

R. Elix. Cinchon. Flav.,	f3ij.
Curaçoa,	f3ij.
Acid. Sulph. Dil.,	℥xij.
Aquæ,	f3 ijss.—M.

Dose. A teaspoonful every two hours.

Here, as well as in other conditions calling for the use of quinia, but where it is difficult to administer it by the mouth, we may give it with good effect in the form of suppositories made with cocoa butter, as small as possible, and containing one or one and a half grains of quinia each.

The child ought to be laid on an inclined plane of pillows, and, with the exception of turning it gently towards one side or the other, from time to time, it should be kept perfectly quiet. These directions are particularly important in very young children, as it is in them that debility and exhaustion of the muscular forces are apt to bring about the state of collapse just referred to.

As an example of the kind of case in which stimulants are useful, and to show also the dangerous effects which antimony sometimes produces, we will quote the following :

"A girl between seven and eight years old, was attacked while in good health with severe bronchitis. On the second day, when we were called, she was very much oppressed, the skin was hot and dry, the pulse rapid, and the surface pale. We ordered a cupping to the amount of four ounces, with some dry cups besides, over the back, and two drops of antimonial wine with ten drops of sweet spirit of nitre to be given every two hours. On the third day a blister was applied over the sternum. On the fourth day we found the child in the afternoon very pale, dozing or tossing about on the bed, and sometimes rising up on her hands and knees with a bewildered look; she was inattentive, so that it was almost impossible to catch her eye; the eyes were sunken, and the countenance was distressed and anxious; she moaned constantly and looked very ill; the skin was still hot; there was neither vomiting nor purging. The respiration was very much oppressed, and she coughed a good deal, though not so much as before. We suspended the antimony at once, and gave a teaspoonful of brandy in water, directing it to be repeated in three-quarters of an hour; after the second dose a teaspoonful was to be given in a wineglassful of milk and water every two hours throughout the night. On the following morning, the child looked better; she was less pale, and the eyes were not so excavated. The breathing was better. She was still very drowsy, but often waked partially with screaming and affright, and when awake took very little notice. The milk and brandy were continued every two hours. On the afternoon of this day, all the unpleasant symptoms had disappeared; there remained only those indicative of a slight bronchitis, and she was soon quite well. Now it seems to us exceedingly clear that, had the antimony been continued in this case, on account of the hot, dry skin, oppressed breathing, frequent cough, and from the absence of vomiting and purging, the child would have died."

The most important points in the treatment of *chronic cases*, are to insist

upon a rigorous and persevering regulation of the hygienic conditions of the patient, and to make use of tonic, balsamic, and expectant remedies. The child should be carefully and warmly clothed, and, when at home, kept in dry, well-ventilated, and, if possible, airy rooms, at a uniform temperature. The bedroom of such a child ought to be heated in winter by a wood-stove, or open wood-fire, if that is sufficient to keep up a proper temperature. In our cold winters we have found no plan so good as a well-managed wood-stove. Coal fires cannot be lowered or extinguished at night, as they ought to be, and often keep up, through the day, too high a temperature. They are unmanageable.

These, indeed, constitute the truly important part of the treatment, for without them, there is but little chance that drugs of any or all kinds, diet, or any other measures, will be of any real service. The dress and temperature ought to be the first things attended to, and after them, and as a secondary matter, certain medical substances will assist in removing the disease. The child ought to be taken as often as possible into the air in fine weather, and only in fine weather. The diet should be selected with a strict view to the improvement of the strength and vigor of the constitution; the food may consist, if the child be of proper age, of light meats, of potatoes and rice, as the only vegetables, and unless there is some contra-indicating circumstance, of a small quantity of wine with the midday meal. The best wine is port, of which one or two tablespoonfuls may be given in a considerable quantity of water.

Tonics must be administered throughout the course of the disease, or until the appetite and strength shall have improved to such an extent as to make them no longer necessary. The best are quinine, in a dose of a grain morning and evening, to be continued for several weeks; or, when the child is thin and anæmic, small doses of arsenic with iron, as recommended in the article on eczema, and cod-liver oil, in doses of half a teaspoonful to a teaspoonful, three times a day after meals, either pure or in some carefully-made emulsion, will often greatly assist in curing these chronic forms of catarrh.

In one case of chronic bronchitis, which came under our care, the patient recovered under careful regulation of the hygiene, and the use of a decoction of seneka, prepared by boiling a drachm, each, of seneka and liquorice-roots, in a pint of water, to half a pint. The decoction was strained, and a large teaspoonful given three times a day. The remedy was continued during a period of two months; under its use the child grew fat and strong, and recovered entirely from the disease.

Other remedies, proposed by different authors, are the various resinous preparations, the balsams of tolu and copaiba, benzoin, and the sulphurous mineral waters. In cases of long standing, where mucous râles persist throughout the lower part of the lungs, showing an abundant morbid secretion, tannic acid has been found, by several good authorities, of much service. While these means are employed, it is recommended, also, to make use of counter-irritants. If any are used, they ought to be such as will not produce too much inflammation of the skin; as, for instance, weak Burgundy pitch plasters, daily frictions with hartshorn and sweet oil, a

simple diachylon plaster, or very mild pustulation with croton oil, or a mixture of croton oil and iodine, such as the following:

R, Ol. Tiglii,	fʒj.
Ether. Sulph.,	fʒj.
Tr. Iodinii,	.
Alcoholis, āā	fʒiij.—M.
S. Locally.	

ARTICLE IV.

EMPHYSEMA.

EMPHYSEMA of the lungs is of quite frequent occurrence in children. It is much more generally met with in an acute form, developed during the progress of some pulmonary disease, than in the chronic form which it so often assumes in the adult. It is probable, however, that in many cases of asthma in childhood, there is an emphysematous condition of the lungs which has been gradually developed at an early period of infancy, in consequence of the respiratory embarrassment attending rachitic disease of the thorax. There is probably in such children a congenital delicacy and weakness of the lung-tissue, and subsequently, if the constitution is re-established, and the deformity of the thorax removed, as it frequently will be, there may be a restoration, to some extent, of the elasticity of the pulmonary tissue, with a corresponding decrease in the evidences of emphysema. It is also highly probable, judging from the frequency with which, in fatal cases of acute pulmonary disease in young children, more or less marked lesions of emphysema of the lungs are found, that this condition is frequently developed to a certain degree in the course of such cases which recover, and that subsequently the lung-tissue regains its normal state.

ANATOMICAL APPEARANCES.—The term emphysema of the lungs, is usually employed to include two conditions essentially dissimilar, and to only one of which it is in reality applicable. One of these is *vesicular* emphysema, which is dependent on dilatation or coalescence of the pulmonary air-cells, without any escape of air into the connective tissue of the lung, and which would, therefore, be more correctly called *rarefaction of lung-tissue*. The use of the term vesicular emphysema is, however, so universal and long-established, that it does not seem desirable to discard it. The other variety is *interlobular* or true emphysema, in which the air escapes from some point into the connective tissue of the lung, and dissects its way between the lobules and under the pleura.

In vesicular emphysema the portions of lung usually most affected are the apex and the anterior border; it may, indeed, be limited to these parts, or may be present, in varying degrees of intensity, along the base and even over the entire surface of the organ. Usually it is present in both lungs simultaneously, though often much more highly developed on one side than the other.

The dilatation of the vesicles causes marked enlargement of the part affected, and when both lungs are seriously involved, they project for-

wards, occupying the mediastinal space, with their anterior borders closely approaching each other. The emphysematous portions do not collapse when the thorax is opened; they are pale, dry, and bloodless, and, when pressed with the finger, afford a soft, doughy feeling, with but an imperfect sense of crepitation. On examining the surface carefully, the dilated vesicles are clearly visible, forming clear, usually round spaces as large as a pin's head or a millet-seed. The effect of this distension upon the surrounding viscera and upon the shape of the thorax are the same in kind, though not so great in degree, as are met with in the adult. The distended anterior portion of the left lung covers more of the heart than normal, and tends to depress this organ downwards and to the right. In the same way when extensive emphysema of the right lung is present, the liver is usually depressed. If both lungs are affected with marked and diffuse emphysema, the thorax is considerably distended, the curve of the ribs is increased, while they are elevated so that their course becomes more horizontal, and the thorax becomes shorter, deeper in its antero-posterior diameter, and more rounded.

The other variety of emphysema—really the only one which strictly merits the name in its usual intention—is the *interlobular*. Here the air makes its escape from a rupture of some air-vesicle or minute bronchiole into the connective tissue of the lung, and then readily makes its way along the bronchial tubes between the lobules so as to reach the surface of the lung. Here it presents itself in the form of minute bubbles of air, of rounded or elongated form, easily recognized by their paleness and transparency, usually arranged in irregular, curving and branching lines, and which can be proved to be in the interstices of the lobules by the fact that they can readily be pressed by the finger from one place to another, or forced to coalesce. When these little bubbles are thickly crowded together they produce an appearance well compared by Rokitansky to froth. Associated with them are often found larger bullæ, where the air has separated the pleura from the surface of the lung; these form flattish, convex prominences above the surrounding surface, and are freely movable. It will be understood that in interlobular emphysema of the lungs, the size of the organ is comparatively little affected, and consequently that little or no influence is exerted by it upon adjacent viscera or upon the shape of the thorax. This condition is comparatively rare, and is not usually associated with marked vesicular emphysema; indeed, the anatomical relations of the two forms are not clearly understood. In places where the pleura is stripped off from the lung over a considerable space, the membrane may be ruptured and air escape into the pleural cavity, constituting pneumothorax, examples of which accident will be found in our article on this latter affection. In other cases, the air makes its way along beneath the pleura to the root of the lung, or by penetrating into the substance of the organ, and following up the divisions of the bronchi, it reaches the same point. It may then pass into the mediastinal spaces, where the loose connective tissue becomes highly emphysematous, so as to present numerous large vesicles with delicate walls, altogether resembling the appearances seen in animals in the slaughter-house. From the mediastinum the air

readily passes upwards into the connective tissue of the neck, where it may first produce a crepitant swelling in the suprasternal, supraclavicular, or inframaxillary regions; and may even extend thence over the surface of the trunk and extremities so as to produce general emphysema.

In the following interesting case, which has already appeared in print,¹ the subcutaneous emphysema did not extend below the clavicle. The minute perforation on the anterior surface of the upper lobe was perhaps due to the inflation of the lungs at the time of the examination, or may have occurred just before death. It would certainly have led to pneumothorax, had it been earlier present:

CASE.—Acute Miliary Tuberculosis: Cough and Dyspnœa: Cervical Emphysema—Interlobular Emphysema—Interlobular Emphysema with Perforation of the Pleura: Emphysema of Mediastinum and Neck.—John F. was born of a stout, hearty young woman, 17 years of age, who nursed him; and he seemed to thrive until eight days before his death, which took place January 24th, 1868, at the age of four months. The symptoms during his sickness were dyspnœa, occasional dry, hacking cough, and anorexia. A few days before his death subcutaneous emphysema made its appearance over the lower part of the neck in front, spreading over both sides, and altering the entire contour of the neck, but not descending below the clavicles. The post-mortem examination was made fifteen hours after death.

The head was not examined. The subcutaneous emphysema persisted as above described.

On removing the sternum, the mediastinal spaces were found much distended with air, the meshes of the connective tissue is some spots forming vesicles more than one inch in diameter, and suggesting forcibly the appearances often seen in animals in the slaughter-house. The emphysema extended up along the trachea and larynx, and to a considerable distance on either side of the neck. There was not a trace of decomposition of the tissues. The lungs collapsed but slightly; the posterior portions were deeply congested, purplish, and almost non-crepitant, but expanded almost fully on inflation. There was neither pneumothorax nor pleuritic effusion or adhesions. The larynx, trachea, and lungs were removed, and inflated under water, when air was found to escape from the right lung in two places—on the anterior face of the upper lobe, and on the inner surface of the apex. On examining the rupture of the anterior surface of the upper lobe, the opening was found to be very small, and to be seated in the midst of a spot where the pleura was separated from the lung so as to form a large vesicle. There were other smaller pearl-like vesicles studding the surface of the lung. The apex was the seat of numerous miliary tubercles, both in the substance of the lung and immediately beneath the pleura. At one point on its inner aspect there was such a sub-pleural deposit, half an inch in diameter, which had undergone cheesy change, and in the centre of which there was an ulcerated opening in the pleura. The escape of air through this perforation was prevented by the close apposition of a tuberculous bronchial gland, about half an inch in diameter, which lay immediately on the right bronchus. The other bronchial glands, especially on the right side, were also tuberculous. The left lung presented no perforation of the pleura. At several points, especially along the anterior edge of the lung, there were large emphysematous bullæ, one inch long by half an inch wide, and in the neighborhood were numerous smaller vesicles of the same nature. On incising the lung near these, smaller clusters of gray miliary tubercles were found. Miliary tubercles were also found on the peritoneal investment of the liver and spleen, and in the substance of these organs and of the mesenteric glands. There were small irregular ulcers in the lower part of the ileum, and numerous small yellowish submucous deposits in the cæcum.

¹ Pepper, On some Cases of Emphysema of the Neck. Philadelphia Medical Times, August 1st, 1872.

Causes.—Although the vesicular and interlobular forms of emphysema are anatomically quite distinct, they may advantageously be considered in connection with each other as regards the mode of their development.

Age.—Vesicular emphysema, though a frequent sequel of acute thoracic diseases in children, cannot be regarded as a disease of childhood in the same sense as the interlobular form. It is true that the delicacy of the walls of the air-vesicles during early life would seem to favor the occurrence of dilatation, but experience shows that it does not favor the development of emphysema nearly so strongly as does the gradual degeneration and weakening of the walls of the air-vesicles which comes on in advanced years.

Interlobular emphysema, on the other hand, is much more frequent in children, and reaches degrees of severity which are scarcely found in later life. So, too, the occurrence of subcutaneous emphysema, in consequence of the rupture of some minute bronchiole or air-vesicle, with the production of sub-pleural and then mediastinal emphysema, is an accident almost limited to early childhood, since of the recorded cases (about 25) in which it has occurred, four-fifths (20) have been observed in young children. Of these 20 cases of "general emphysema in children," to employ Roger's term¹ (of which 19 were collected by him, and 1 subsequently published by ourselves²), 6 occurred under the age of 2 years, 10 between 2 and 4, and only 4 between 10 and 15 years of age.

Previous Diseases.—In children, emphysema occurs as a sequel to some other disease, pulmonary or laryngeal. The affections which most strongly predispose to it are hooping-cough, the bronchitis of measles, simple bronchitis, pneumonia, and pseudo-membranous croup. Of all these, hooping-cough is by far the most fruitful cause. It will be observed that the diseases named present the common symptom of severe cough, often attended with impediment to the escape of air, either from spasm of the air-passages, or accumulation of secretion in the bronchi, or mechanical obstruction of the larynx by false membrane.

Mechanism or Mode of Production.—The way in which pulmonary emphysema is developed has been made the subject of frequent and conflicting speculation. Of the two chief theories which have been advanced in explanation, one (the *inspiratory*) regards the over-distension of the vesicles as the result of the excessive operation of the forces concerned in inspiration; the other (the *expiratory*) explains it as caused by violent but impeded expiratory efforts. The inspiratory theory is still upheld by some eminent writers, but clinical observation is leading to its abandonment. In its original form as advanced by Laennec, it was based upon the erroneous notion that the forces of inspiration are greater than those of expiration, and that consequently emphysema might result from mere excessive inflation of the lungs. This has, however, been universally abandoned as of general application, since the discovery of the important fact that in *forcible* breathing the power of expiration is considerably (at least one-third) greater than that of inspiration; though it is probable that in

¹ Henri Roger, Archives de Médecine, 5ème ed., tome xx, pp. 129, 288, 403.

² W. Pepper (*loc. ante cit.*).

some morbid conditions of the pulmonary tissues, violent inspiration may of itself be capable of producing emphysematous distension of the air-vesicles.

The form of the inspiratory theory, which is still retained by some authorities, is based upon modifications introduced by Dr. William Gairdner, and is an expansion of the idea that if certain portions of the lungs are, from collapse or other cause, incapable of expansion, the atmospheric pressure will determine excessive dilatation of the remaining portions, in order to prevent the occurrence of a vacuum as the thoracic walls expand. There are, however, such grave objections to this theory, which, it will be observed, rests upon the supposition that the expansion of the thorax and the amount of air inspired remain at the normal point, although portions of the lungs are collapsed or otherwise rendered unable to expand, that we are strongly inclined to regard the expiratory theory as the only one capable of general clinical application. We owe to Sir William Jenner chiefly the satisfactory refutation of the principal argument which was formerly brought against this latter theory, that "the expiratory act is mechanically incapable of producing distension of the lung, or of any part of it. The act of expiration tends entirely towards emptying the air-vesicles by the uniform pressure of the external parietes of the thorax upon the whole pulmonary surface; and even when the air-vesicles are maintained at their maximum or normal state of fulness by a closed glottis, any further distension of them is as much out of the question as would be the further distension of a bladder, blown up and tied at its neck, by hydrostatic or equalized pressure applied to its entire external surface" (Gairdner). A little consideration of the anatomical relations of the lungs to the thorax shows the falsity of this argument. The different portions of the lungs are in contact with surfaces and tissues of very different degrees of resisting power, and while the entire postero-lateral portions are supported by the unyielding ribs, the apices are covered only by soft tissues, and the anterior borders of the lungs are supported externally by the comparatively yielding costal cartilages, while centrally they are able to encroach considerably upon the tissues of the mediastinal spaces. In ordinary free expiration the air is forced out of the lungs by a pressure so moderate and gradual that even the weakest parts of the thoracic walls are sufficiently firm to maintain it. But when the expiratory efforts become more violent, the air is pressed with great force from the central, basic, and lateral portions by the ascent of the diaphragm and the compression of the thorax, while the outward current from the apices and anterior margins is comparatively feeble. If, therefore, from any cause, the normal relation between the volume of the expiratory current of air and the calibre of the large bronchi be disturbed, the portion of air which cannot escape will be driven violently into the apices and anterior margins, not only overcoming the outward current of air proceeding from those portions of the lungs, but producing an excessive degree of distension of their air-cells. The strongest possible confirmation of the truth of this view is to be found in the fact that emphysema, both of the vesicular and interlobular form, is found to

be developed in the various parts of the lungs in precise correspondence with the degree in which they lack firm external support.

There are two ways in which a disturbance of the above relations may be effected; either by an obstruction in the air-passages, which prevents the free escape of the air, or by the expiratory act being so sudden and violent that the volume of air hurriedly forced from the air-vesicles is too great to pass freely through the primary bronchi. Instances of this latter condition are familiar to all in violent fits of coughing, during which, even when there is no obstruction in the air-passages, the degree of distension of the apices may be appreciated by the bulging of the supra-clavicular tissues. The full pulmonary resonance, which is elicited by percussion of this bulging, proves conclusively that it is due to distension of the apex; and it is therefore easily understood how the repeated operation of such a cause may gradually lead to the development of vesicular emphysema, or how in an abrupt, violent, and prolonged expiratory effort, attending a fit of coughing, there may be a rupture of some minute bronchiole or air-vesicle, followed by interlobular emphysema. Undoubtedly, also, the mechanical effects of such over-distension will be greatly enhanced by morbid conditions of the lung-tissue which weaken its elasticity, such as are present in severe bronchitis, especially when associated with constitutional diseases. Far more frequently, too, there is associated some cause of partial obstruction to the escape of air, such as the spasmodic contraction of the air-passages in whooping-cough, the presence of layers of false membrane in the larynx or trachea, or thickening of the bronchial mucous membrane with plugs of viscid tenacious mucus in the tubes.

It is very possible, also, that in some cases interlobular emphysema may be caused by the implication of a minute bronchiole in the progress of the softening of some spot of diseased tissue, so that the air might find entrance to the interstitial connective tissue without any mechanical cause of over-distension and rupture of air-vesicles. Thus in the case reported above (p. 220) there was certainly a very close connection between the position of the patches of tuberculous deposit and the bullæ of sub-pleural emphysema, so much so that we cannot doubt that the escape of air was in some way favored by their presence. In another case, also, where interlobular and sub-pleural emphysema, followed by pneumothorax, occurred, and which is reported at length in our article on this latter affection, it seemed to us that probably the softening of superficial circumscribed patches of pneumonia had opened into minute bronchioles, and thus allowed the escape of air.

SYMPTOMS.—We have already seen that in young children emphysema occurs usually in an acute form in connection with some acute disease of the lungs. Although, therefore, its presence may be suspected in such cases where violent paroxysms of cough have occurred, associated with prolonged, severe dyspnœa, there are scarcely any physical signs by which its existence can be determined. The percussion-resonance will continue clear, or even become somewhat exaggerated, and this fact of the absence of any dulness (due to pneumonia, collapse, or pleural effusion), in a case of whooping-cough or bronchitis, when severe cough has occurred with un-

usually extreme dyspnœa, is of diagnostic value. The respiratory murmur undergoes no immediate change in its character, and it is not possible, owing to the violent and rapid respiratory efforts of the child, to detect any diminution in the force of the murmur. Expiration in such cases is, however, often already prolonged and laborious.

The development of acute vesicular or interlobular emphysema, then, is suspected rather on account of the character of the disease from which the child is suffering than from any distinct substantive symptoms of these conditions. Exception is to be made, however, of the rare cases, in which suddenly, in the course of an acute pulmonary disease, a swelling is noticed at some part of the neck, or in the subclavicular space, which on palpation is found to crepitate. This may be regarded as, in all probability, connected with extensive interlobular emphysema. The other chief cause of such subcutaneous emphysema is perforation of the larynx or trachea, and the previous symptoms will enable us to exclude this rare condition without difficulty.

In other cases, however, vesicular emphysema in children assumes the chronic form, more usually found in adults, and will then be attended with the well-known symptoms of this affection. At times, it occurs evidently as a sequel to some acute pulmonary disease, in the course of which it has been developed in the manner above described, and after the original disease has passed away, it persists, either owing to original weakness of the lung-tissue, or to the extreme degree of the dilatation of the air-vesicles. At other times, it is met with as a purely chronic affection, which may begin in early childhood, and gradually increase until the disease is fully developed. In such patients there is probably some congenital weakness and tendency to degeneration of the pulmonary tissues. It is not unfrequently found that there are also evidences of rachitic disease of the ribs in such cases.

Children with chronic emphysema present various degrees of habitual dyspnœa, which is always readily increased by exertion. They are very subject to attacks of bronchitis, during which the breathing is much embarrassed and wheezing, the chest is full of sonorous and sibilant râles, and the cough occurs in severe paroxysms without much expectoration. During these attacks, not unfrequently the child suffers at night from violent paroxysms of spasmodic asthma. Indeed, it may happen that attacks of asthma will be induced in emphysematous children by the most trifling causes, such as changes of weather, indigestion, and the like. The attacks of bronchitis vary greatly in different cases in their relation to season and temperature; in some they occur almost exclusively during the damp, cold weather of fall and winter, while in others they are most frequent and severe during summer and spring, and the child finds more relief during cold weather.

The cough varies much in its intensity and character. During the attacks of bronchitis it is usually very severe, occurring in long spells, at first with a little mucous expectoration, and later, as the attack passes over, with more abundant muco-purulent sputa. In the intervals of the attacks it may continue as an occasional dry and rather wheezing cough,

or it may be more troublesome on account of a certain degree of chronic bronchitis being always present, or finally it may altogether subside. After the disease has lasted a considerable time, however, cough may become persistent, occurring most severely at certain periods of the day, and attended with a considerable quantity of muco-purulent expectoration. In such cases, when emphysema is conjoined with chronic bronchitis, the suspicion is apt to arise that the child is suffering from phthisis, and the positive determination of the diagnosis may indeed be attended with some difficulty. The reader is also referred to the remarks made in this connection on the subject of chronic bronchitis (see p. 210).

In young children under the age of 5 or 6 years, emphysema rarely reaches so great a degree, or persists for so long a time as to induce marked changes in the shape of the thorax, or to seriously affect their nutrition. In children somewhat older, however, when the disease is more severe and chronic, it may be attended with most of the symptoms familiar in the adult. The appearance of such children is apt to be frail and delicate, their muscular system develops slowly, and they become so readily fatigued and out of breath that they avoid play or much exercise. The shape of the thorax becomes gradually altered; the shoulders grew high and rounded, and the chest is prominent and distended in its upper part, while owing to imperfect expansion of the lower lobes, there may be perceptible retraction of the base of the thorax in front, or even a marked depression around the entire base of the chest. Of course, this is likely to occur to a more marked degree if the emphysema is associated with rickets.

The *physical signs* vary greatly with the extent and degree of the emphysema. In cases where it is limited to small areas of the lungs, scarcely any physical signs can be detected; but in partial and more severe forms, the following phenomena can be observed: The respiratory movements are restricted, especially in the way of expansion; and during inspiration the movement is chiefly one of elevation effected by overaction of the upper respiratory muscles, and attended with an evident deepening of the depression around the base of the chest. The percussion-resonance is very full and clear, or even tympanitic, though owing to the marked resonance normal in children, it is difficult to determine the degree of its exaggeration. There may be associated some impairment of resonance over the retracted base of the thorax, and especially posteriorly, where there may be congestion of the lung with accumulation of secretion in the air-passages, due to the coëxisting bronchitis. The respiratory murmur is weakened, though rarely to the degree noticed in adults; the expiratory murmur is decidedly prolonged and frequently wheezing. Both inspiration and expiration are apt to be accompanied with sonorous and sibilant râles. These, and especially the sonorous râles, are most markedly developed over the posterior parts of the lungs, near the larger bronchi. In some cases, moist râles may also be heard over the postero-inferior parts of both lungs, owing to the presence of an unusually large quantity of secretion in the smaller bronchial tubes. During one of the acute aggravations of the bronchitis, attended with nervous asthma, to which we have above al-

The use of this was followed by improvement in appetite and strength, and by a marked diminution in the frequency and severity of the asthmatic attacks. As, however, the cough continued quite severe, with abundant loose mucous râles throughout the chest, the treatment was changed to the following:

R. Syr. Phosphat. Comp., f℥ij.
 Elix. Calisayæ,
 Aquæ, āā f℥j.—M.
 Dose. Two teaspoonfuls in water before meals.

And to relieve the cough:

R. Ammonii Muriatis, gr. lxxij.
 Syr. Senegæ, f℥ss.
 Tr. Hyoscyami, f℥iss.
 Ext. Pruni Virg. Fluid., f℥ss.
 Syr. Zingiberis, q. s. ad f℥ij.—M.

Dose. A teaspoonful in water three or four times in twenty-four hours, according to the severity of the cough.

Under the use of these remedies, her improvement was rapid and continuous, and for the past year and a half there has been scarcely any tendency to bronchitic attacks. Her frame has developed, and the shape of the thorax has improved so as to confirm the evidence of the physical signs that the emphysema is gradually disappearing.

DIAGNOSIS.—In regard to the detection of the acute form of emphysema at the time of its occurrence, we have already shown that there are no signs sufficiently distinctive, and that if any unusual severity or persistency of dyspnœa rouses the suspicion, we may only assume its presence in consequence of the great frequency with which it is developed in certain diseases.

This is even more true with regard to the interlobular than the vesicular form, except when the sudden appearance of subcutaneous emphysema proves its existence.

If, however, emphysema becomes firmly established, and persists, it becomes attended with the well-marked symptoms already described, and there are then scarcely any conditions with which it can be confounded. In almost all cases, there is associated bronchitis, either in the form of repeated acute attacks, or, more frequently, of a chronic form of varying degrees of intensity. It is therefore necessary to guard against overlooking the evidences of emphysema, and considering such cases as simple forms of bronchitis. In cases where emphysema of long standing is accompanied by severe and chronic bronchitis, it may be confounded with phthisis. Apart, however, from the fact that chronic phthisis is rare in childhood, a careful study of the history of the case, and of the physical signs, will enable us to avoid this error.

PROGNOSIS.—Only in extreme cases of emphysema in children is the prognosis so unfavorable as in this condition in adults. In many cases where there can be no reasonable doubt that it exists to a considerable extent, the lung-tissue regains its contractility soon after the exciting cause of the emphysema has been removed, and all evidences of its existence gradually disappear. Even in more protracted cases, when the disease persists for some years, and leads to deformity of the thorax, there is

ground for hope that, under the influence of the developing constitution and frame, and sustained judicious treatment, considerable relief will be obtained. This is equally true in regard to the tendency to attacks of nervous asthma, which is so frequently associated with emphysema, as it is true of its other symptoms. Children who, at an early age, suffer severely with such attacks even from the slightest causes, are frequently seen to entirely outgrow the distressing tendency, and to bear any change of climate or vicissitude of weather without inconvenience.

The prognosis of mediastinal and subcutaneous emphysema dependent upon the interlobular form, is of course controlled entirely by the nature of the primary disease. It undoubtedly of itself aggravates the dyspnœa caused by the original pulmonary disease, but still very rarely reaches so extreme a degree as to endanger life. It must be borne in mind, however, that it is very frequently associated with pre-existing lesions of the lungs, which, either from their character or their extent, are almost necessarily fatal. Thus, of the 20 cases before referred to as on record, in only 4 has this accident been followed by recovery. It must, therefore, be regarded merely as a serious complication, but one which would not justify an altogether unfavorable prognosis in an otherwise curable condition.

TREATMENT.—The treatment of cases of acute pulmonary disease, in the course of which it is suspected that emphysema has occurred, cannot be much modified on account of this complication. As it is, however, closely dependent, in such cases, upon the frequent and violent cough, the most important indication is to allay this by suitable antispasmodics and sedatives. At the same time, as the increased dyspnœa which is caused by the development of the emphysema must seriously add to the exhaustion of the patient, the utmost care must be exerted to sustain the vital powers, and to discard every depressing element from the treatment.

The same remarks, which are used above with special reference to vesicular emphysema, apply with equal force to the interlobular form when it becomes complicated with mediastinal and subcutaneous emphysema. The only chance of recovery in such cases, is to be found in sedulously supporting the system until the primary disease (if of a curable nature) has passed away, and the effused air has been gradually absorbed. This absorption may be, to some extent, hastened by gentle frictions on the emphysematous parts with the hand. In cases where the extent of the external emphysema is such as to threaten life, minute punctures may be made in the distended skin, and the escape of the air favored by gentle pressure with the hands towards the point of puncture.

The most important field for medical treatment is, however, to be found in those cases where emphysema, whether acute or not in its incipency, has passed into a chronic or persistent form. The indications for treatment which here present themselves, are mainly to relieve the chronic bronchitis, which is almost invariably associated with emphysema, if it has not been its chief cause; to eradicate any rachitic tendency which often coëxists, and, so far as possible, to counteract its results; to guard against and relieve the acute attacks of bronchitis often accompanied with nervous asthma, from which such children suffer; and finally to favor, so

far as lies in our power, the restoration of the dilated lung-tissue to its normal condition.

There are several considerations of a general character, which will be found to have an important bearing upon these requirements.

Change of climate, when it can be judiciously made, is often attended with excellent results, particularly as regards the bronchitic irritation and the attacks of asthma. We have known children who suffered most severely from these conditions, which were aggravated by any trivial causes so long as they remained in their native place, but who on removal to other climates, received marked relief and gradually outgrew the disease. It is not at all necessary that this change should be to a distant spot; often the most convenient, dry, elevated, inland locality will answer excellently.

The clothing of such children should be carefully studied and regulated. Without being so heavy as to oppress, it must be at all times warm enough to thoroughly protect; and at the same time there should be a suit of flannel or silk (consisting of an undershirt with long sleeves, and long drawers coming down to the ankles), of varying thickness to suit the season, worn throughout the year, to protect the surface of the body from the chilling effects of sudden vicissitudes of temperature.

As further means to secure activity of the circulation and function of the skin, the use of salt baths (of a temperature to suit the season), and followed by brisk rubbing with a coarse towel, are to be recommended.

In no condition of the system is the use of gymnastic exercise more to be insisted upon. We should select those exercises with light dumb-bells or Indian clubs, which will tend to strengthen the muscles of respiration, expand the lower portions of the chest, which, as we have already pointed out, are apt to be retracted, with some incurvation of the ribs, from the co-existence of rickets at an earlier age. As children in whom emphysema assumes the chronic form are usually over the age of 6 or 7 years, such exercises can be readily carried out.

Very recently several forms of apparatus have been devised for enabling the respiration of air of different degrees of condensation or rarefaction to be employed with accuracy. The one which we have ourselves employed is known as Waldenburg's apparatus, having been designed by this lamented physician, who also was chiefly concerned in introducing this new method of treatment. In the case of emphysema, the patient is caused to expire into a receiver containing rarefied air. It is evident that this will exert an increased suction power on the air in the lungs, and thus will favor the emptying of the air-vesicles and the contraction of the chest to its normal limits. Undoubtedly in children, this method, which we have used with good results in adults, should be employed with caution; but it is especially in young persons, when the elasticity of the ribs and of the lung-tissue is greater, that the most positive effect may be hoped for in causing a restoration of the chest to its normal size.

The selection of the diet should be made with care. It will often be found that all the symptoms are aggravated by any digestive disturbance, and we have seen, as is indeed frequently the case in emphysema of adults, violent paroxysms of nervous asthma induced by indigestion. As

the digestion in such children is apt to be weak, this point requires the greater care.

As regards medication, the most important remedies are such as will affect the constitution favorably.

We should recommend the use of cod-liver oil in properly graduated doses, and, especially where there are evidences of rachitic disease, the compound syrup of the phosphate or the lacto-phosphate of lime may be advantageously associated in the form of emulsion. If iron is not thus administered in the form of the phosphate, the oil may be given alone, and iron should be taken separately in some other combination, as in the following:

R. Potassii Bromidi,	3ij.
Potassii Iodidi,	gr. xlvij.
Syr. Ferri Iodidi,	f3ij.
Syr. Tolutani,	f3vj.
Aquæ,	f3ij.—M.

Dose. A teaspoonful thrice daily in a little water.

The dose here directed is for a child of about eight years of age.

We have also found the prolonged use of arsenic for its constitutional effects of much value in some cases.

Cough should be relieved, so far as possible, without the use of opiates and nauseating expectorants. Among the drugs which we have found most useful in controlling it, as well as relieving the chronic bronchitis upon which it usually depends, are the iodide of potassium, which may be advantageously combined with the potassium bromide, as in the above prescription, the bromide of ammonium, and the muriate of ammonia. If the cough be very troublesome, especially at night, tincture of hyoscyamus and minute doses of morphia may be occasionally associated. It may, however, become necessary to substitute for these, or to combine with them, other alterative and stimulant expectorants, such as seneka or copaiba.

The use of quinia, and, at times, of strychnia with it, is indicated throughout a large part of the treatment for the useful influence of these drugs upon the digestion and general nutrition, and especially upon the tonicity of the muscular system.

The acute attacks of nervous asthma which are apt to occur from time to time must be relieved at the moment by the prompt use of relaxing emetics, hot mustard-water foot-baths, the inhalation of ether or of the smoke of stramonium cigarettes. The frequency of these distressing attacks will, however, be greatly influenced by the persistent employment of the general treatment above sketched.

Finally, it must not be forgotten that despite the obstinacy and severity of the symptoms of emphysema in some cases, and the positive alterations of the shape of the thorax, there is always reason to hope that if we can succeed in removing the element of chronic bronchitis, and in favoring the expansion of the lower lobes of the lungs, so as in these ways to relieve

the strain upon the upper portions of the organs, the distended vesicles will gradually regain their elasticity, and as the thorax enlarges with advancing years, all symptoms of the disease will pass away.

ARTICLE V.

PLEURISY.

DEFINITION; FREQUENCY; FORMS.—Pleurisy consists in inflammation of the pleural serous membrane.

Idiopathic pleurisy is a comparatively rare disease under five years of age, and especially in the first and second years of life. After the age of five years it becomes more frequent. Secondary pleurisy, on the contrary, or that which occurs in the course of other diseases, is common at all ages. M. Bouchut met with it in 23 out of 68 autopsies of new-born and suckling children. Of the 23, 9 accompanied acute pneumonia, 6 tubercular pneumonia, 5 entero-colitis, and 3 different other diseases. This form of the affection is apt to be overlooked during life, being masked by the concomitant disease.

We shall describe two forms of the disease, the *acute* and *chronic*.

PREDISPOSING CAUSES.—As to the influence of *age*, it has already been stated that idiopathic pleurisy is rare between birth and five years of age. It is certainly rare, during those years, in comparison with pneumonia, and especially with bronchitis, for we find we have met with twice as many cases of pneumonia, and seven times as many cases of bronchitis, as of pleurisy under the age of 5 years.

Pleurisy occurs more frequently in boys than in girls. Of our own cases, three times as many were in boys as in girls. The idiopathic form is most apt to occur in vigorous and hearty subjects, while the chronic and cachectic forms attack those who are feeble and delicate. It is often, as already stated, a secondary affection, occurring particularly during pneumonia, and, after that disease, during rheumatism, scarlet fever, and Bright's disease. *Season* is another predisposing cause. It is most common during winter and spring, especially the latter.

The *exciting causes* are very obscure in most cases. The only ones which seem to have been ascertained with any certainty, are exposure to cold and sudden changes of weather. It has been said to follow external violence. In one of the cases that came under our own observation, the child had struck the affected side severely against a pointed stick on the day of the attack.

ANATOMICAL LESIONS.—In some rare cases of very slight inflammation, with serous effusion, the pleura may retain its normal appearance, but usually it presents the characteristic lesions of inflammation. These are more or less minute and abundant injection and punctation, and spots or patches of an ecchymotic appearance, observable particularly at the points where the formation of false membrane has taken place. Another change

produced in the pleura by inflammation is the loss of its natural polish, which is replaced by a more or less granular and rough appearance. In chronic cases it becomes whitish or opaline in color, and thickened. It is very rarely softened.

In addition to the lesions of the pleura itself there are various diseased products of secretion which require notice. These may be either solid or liquid. The solid products are the false membranes which exist so generally in all serous inflammations. They are found both upon the costal and pulmonic pleura. In their recent state they are of variable size and thickness, being in some cases very soft and deposited in small points; in others, more extensive, but thin, like paper; and in others again thicker (one or two lines in thickness), firmer, and separable into several layers. The outer layers are yellow, elastic, and soft, while the inner ones are red, more resisting, and marked with vascular arborization. When examined some time after their formation, the false membranes are found to have been converted into cellular adhesions, which may be either very loose, or they may fasten the lung tightly to the costal pleura. The adhesions are generally, however, thin, transparent, and in the form of loose bridles. After a length of time, the false membranes present the appearances of true serous tissue, and like that, are susceptible of inflammation.

The fluid found in the pleural cavity usually consists of transparent or turbid serum, holding albuminous flocculi in suspension. Not rarely, however, there is an admixture of pus with the serum, or the effusion consists of pure pus. Indeed it would seem that the tendency of severe acute idiopathic pleurisy to produce purulent effusion is greater in young children than in adults. The liquid generally occupies the lowest portion of the thoracic cavity, but is sometimes circumscribed at various heights, or between the lobes of the lung by abnormal adhesions, or by some part of the lung which has been rendered incompressible by inflammation.

The lung presents various alterations from its healthy condition. It is pressed backwards towards its root to a greater or less extent. The tissue of the organ is generally found in one of two conditions: either hard, not crepitating, impenetrable to the finger, and presenting a smooth surface when cut into, a state of things which has been expressed by the term *car-nification*, and which is a mechanical effect of pressure; or else the lower lobe, which is in contact with the fluid, is large, heavy, fleshy, rather hard, not so easily penetrable by the finger as in simple hepatization, yielding under pressure only a small quantity of blood, and but slightly retracted towards the spinal column. The latter condition depends in all probability on an effusion which has occurred after or coincidently with partial hepatization.

In some cases, in which the effusion is but small, or where it has been absorbed, the lung is found to be elastic and crepitating. Whatever the amount of fusion may be, the lung can expand to its normal size if the fluid be absorbed, unless it has been too firmly bound down by false adhesions.

Pleurisy, whether complicated with pulmonic disease or not, is much the most frequently confined to one side. In idiopathic cases, it is said to be

more common on the right than left side; when it accompanies pneumonia, to be, on the contrary, more common on the left than right.

SYMPTOMS.—In describing the symptoms, we shall treat first of the physical, and then of the rational signs, and of the course of the disease. The *physical signs* are exceedingly important, as they often constitute, especially in young children, the only means of recognizing the disease. The *pleural friction-sound* is less important than some other physical signs, as it is by no means constantly heard in children under five years of age, and only during the absorption of the fluid, as a general rule, in those above that age. *Bronchial respiration* may commonly be detected from an early period in the attack. It is, however, to be carefully distinguished from the true, sniffling, superficial bronchial breathing due to pneumonic consolidation of the lung. In pleurisy, with effusion, the bronchial respiration is evidently transmitted from the bronchi of the compressed lung and from the trachea. Indeed, it will sometimes be found that, in a case where distinct bronchial breathing is heard over the seat of the effusion during noisy, hurried breathing, it will be lost and replaced by a feeble respiratory murmur, if the child breathes quietly and without making any noise in the throat. At first the bronchial respiration is usually heard during inspiration, but afterwards it exists both during inspiration and expiration, or in the former alone. In a majority of the cases it is heard over the posterior portion of the thorax, and upon one side only. At first it is audible over nearly the whole height of the affected side, while later in the disease it can be perceived only at the inferior angle of the scapula or in the interscapular space. Its duration is variable; it may disappear in a few days, or last for a much longer time. This sign is almost always present at all ages in acute cases, but is often absent in those which are slow and tedious. In suckling children it is not constant, but intermits occasionally, so that it may be heard at one and not at the next examination. As the effusion subsides in favorable cases it is replaced by feeble vesicular breathing, with or without friction-sounds; and later by pure respiration. *Ægophony* can rarely be detected in children less than two years old. Under that age there is heard instead of it resonance of the cry, especially in the region beneath and on a line with the spine of the scapula. It is intermitting, like the bronchial respiration. In children over two years old, ægophony can often be distinguished by careful examination, but never, of course, unless the quantity of effusion is considerable. It is heard at an early period of the attack, and chiefly in acute cases, and must be sought for in the lower portion of the interscapular space, and the inferior dorsal region. It coexists almost invariably with bronchial respiration, lasts but a short time, disappearing after one, two, three, or four days, and it is intermitting. In older children it is sometimes replaced by diffused resonance of the voice, as it is by resonance of the cry in infants. In a case that occurred to ourselves, in a girl between six and seven years old, and in which the disease became chronic, the voice was not purely ægophonic, but reedy and quavering, from the fifth to the tenth day. After that date the effusion became so great that all sound was suppressed.

Feebleness or absence of the respiratory murmur seldom exists at the be-

ginning of acute cases, but in the subacute or chronic form is generally present when the case is first seen. In the latter class of cases feeble respiration is noticed first over the inferior portion of the dorsal region, but as the effusion increases, it is heard also in the upper and anterior regions, and becomes more and more marked, until at length no sound whatever is audible; the respiratory murmur is suppressed. In acute cases, on the contrary, the absence of the respiratory sound is observed at variable periods of the attack; when noticed soon after the invasion, it is generally coincident with distant and transmitted bronchial respiration, which, heard at first over the whole or the inferior three-fourths of the dorsal region, becomes afterwards perceptible only in the interscapular space, or at the inferior angle of the scapula, while the respiration is feeble or absent over the lower portions of the lung. In acute cases the feeble respiration remains limited to the dorsal region, and disappears after a few days,—in from five to eight, according to our experience; while in chronic cases it extends over a large surface, and continues for several weeks, or even months.

Percussion.—This means of diagnosis is very important in all cases of the disease accompanied by effusion of liquid, unless the quantity be exceedingly small. When, on the contrary, the inflammation results merely in the production of thin false membranes, percussion furnishes no useful information.

Percussion is of no assistance, however, at the moment of invasion, as it is not until the period at which effusion takes place that the resonance of the thorax begins to be altered. In acute cases, the resonance is generally duller than natural, though seldom entirely dull, on the second, third, or fourth day. As the effusion augments, the dulness increases over the region occupied by the fluid, until at length all resonance ceases, and the sound is perfectly flat. The degree of dulness can be properly appreciated only by comparing the two sides together. The degree, extent, and duration of this sign will depend, of course, upon that of the effusion. In children, as in adults, the sounds afforded by percussion vary with the position of the patient, which influence, of course, the situation of the fluid in the pleural cavity. When the effusion has become sufficiently large to compress the lung to a considerable degree, the percussion note over the upper part of the lung, and especially beneath the clavicle, assumes a peculiar wooden tympanitic character. This character is very marked in most cases, and is in itself alone sufficient to arouse the attention of the physician. We have several times been led to a correct diagnosis by this one physical sign alone. Observing its presence ought to lead to a thorough examination of the whole chest, when there is generally little difficulty in coming to a correct conclusion.

In regard to the physical signs of pleuro-pneumonia, it may be stated that when a pleuritic effusion takes place in a child laboring under pneumonia, it happens, as a general rule, that the bronchial respiration occasioned by the inflammation of the lung increases in intensity, though in some few cases it is diminished or suppressed. MM. Rilliet and Barthez lay down the following principle:—“That *when a pleuritic effusion occurs*

in a child affected with hepatization of the inferior portion of the lung, all the abnormal sounds which were perceptible over the diseased point are considerably exaggerated, and the sonority disappears."

Inspection of the thorax affords no assistance at the invasion of the disease, nor generally in acute cases which last but a short time, and in which the amount of effusion is small. When, however, the effusion is large, it may be observed, upon close examination, that the movements of the affected side during respiration are more limited than those of the opposite one, and that the intercostal spaces are more projecting than natural, in consequence of distension by the fluid within. At the same time *mensuration* will show that the side on which the effusion exists is larger than the other. The difference may amount to one-third or two-thirds of an inch. In acute cases, in which the quantity of liquid is small, mensuration will of course show no difference.

Palpation is an important means of diagnosis, especially in making the distinction between pneumonia and pleurisy. In the former disease, the vibration of the thoracic walls during either crying or speaking, is augmented; whilst in the latter it is diminished, or when the effusion is considerable, ceases altogether. This sign is important, both in infants and older children.

Another very important sign, which should always be looked for, is *displacement of the heart* towards the healthy side by the pressure of the effusion. Thus in large effusions on the left side we have found the heart pulsating below the right nipple. The value of this sign is particularly great in children, as it can be determined with accuracy even if their agitation or timidity renders other means of physical exploration difficult.

RATIONAL SYMPTOMS ; COURSE ; DURATION.—Acute pleurisy is comparatively rarely met with, as already stated, in children under six years of age, except as a secondary affection. In idiopathic cases it begins with severe pain in the side, cough, some difficulty of respiration, increased frequency of the pulse, loss of appetite, thirst, bilious vomiting, sometimes headache, and in rare instances delirium. The *pain* in the side or *stitch* is almost always present in acute cases, occurring in healthy children, while in those which are slight, or which occur in weak and debilitated subjects, or very young children, it very often cannot be detected. Sometimes, however, its existence may be ascertained in very young children by tenderness of the side shown during the act of percussion. When present in young children, it can always be detected by watching the face of the child and observing its gestures during the act of coughing, and during full inspirations, as in those made in crying, after sudden movement, or in the act of gaping. In an infant of thirteen months old, who was attacked with pleurisy of the right side, with purulent effusion, and which ended fatally within a month, only the blindest observer could fail to see that every act of coughing was acutely painful, for the child uttered each time a short, sudden cry, which was hushed as soon as given, while at the same moment there passed across the face an expression, amounting almost to a grimace, of suffering, which was unmistakable. The pain is aggravated by coughing, by full inspirations, by change of position, and by percussion.

The seat of pain is almost always in front, but it may extend irregularly over the whole of one side to the arm, or it may be confined to the false ribs, or less frequently to the neighborhood of the nipple; it generally lasts from three to six days, though it sometimes continues longer. This symptom was complained of in most of the cases that we have seen. In some it was very acute or severe for one or two days, while in others it was slight, not well defined, and very transitory. In one, the child said there was no pain, but a sensation of weakness in the side when she coughed. In another, the pain was severe for a few hours, but was relieved by a sinapism, and was not felt again, though the attack resulted in a very large effusion on that side. In a third it lasted a week, and in a fourth only two days, though in both the effusion was extensive, and required several weeks for its absorption. In a fifth case it continued for five days. In the last, the effusion was very slight. It was aggravated in all these cases by coughing, by the act of respiration, especially when this was deep, and by motion.

Cough exists in nearly all idiopathic cases and generally from the onset, though sometimes not before the second or third day. Usually frequent and dry, it commonly retains these characters, in acute cases, for four or six days, and then diminishes rapidly. In more tedious cases it continues for a longer time, but moderates in violence after some days. In secondary cases it has no special characters. It was present at all but one of the cases seen by ourselves. Its character varied very much. In some it was frequent, teasing, and very painful. In others it was rare, scarcely troublesome, and only slightly painful. In all it was very dry, this constituting one of its most marked features, and giving it a very different character from the cough of bronchitis, and also, though somewhat less distinctively, from that of pneumonia. It continued almost entirely dry throughout the disease, except in a case which became complicated after a time with slight bronchial inflammation, and, in that, it became loose. There is generally no *expectoration*; if any, it consists of a small amount of whitish, frothy, sero-mucous fluid.

The *respiration* is usually accelerated in acute cases, but remains natural in other respects; the dyspnœa, however, is slight, as a general rule, compared with that of pneumonia. The difficulty of breathing is commonly in proportion to the earliness of the age, and to the extent and rapidity with which the effusion takes place. In the acute cases that have come under our own observation, the breathing was usually about 36, 38, 40, and 48, but in one case it rose to 68 for a single day. It was not labored, and appeared to be difficult only from the fact of its being more or less painful. In a case of double pleurisy, it was most laborious, and dreadfully painful, as was also the cough. In the cases attended with but slight pain, there was no dyspnœa. It usually subsided after two or three days, when large effusion took place, converting the case into the chronic form.

The *fever* is not usually very great, and seldom lasts more than a few days, or a week. In some few cases, however, that we have seen, the febrile reaction has been very high. In one, in a child between three and

four years old, the pulse rose to 172 on the first day, though the respiration was but 36; the skin was very hot and dry, and there was very great drowsiness and inattention. In other cases the pulse was 140, 128, and 124. The acceleration of the pulse usually lasts three or four days, after which it falls, so that by the end of a week it is seldom over 70, 80, or 90. The *heat* of skin is not very great in most instances, and generally subsides rapidly and disappears after a few days. Thus during the first few days of the attack, the temperature may rise to 103° or 104° ; but it soon falls, and, during the remainder of the case, usually fluctuates between 100.5° and 102° . When the case is complicated with pneumonia, the elevation of temperature is even more marked and persists for a longer time. In acute secondary attacks, the febrile symptoms are more marked, as a general rule, than as has just been described, because of the existence of the concurrent disease.

The *countenance* presents no particular characters, except that an expression of pain passes across it occasionally when the child coughs, or takes a deep breath. It is seldom deeply flushed as in pneumonia. The *alæ nasi* are dilated only during the continuance of the difficulty of respiration.

The *decubitus* is generally dorsal or indifferent. In two cases observed by us, in which the effusion was large, the number of inspirations was always from three to five greater when the child laid on the sound, than when on the affected side.

Headache is often present during the first few days, in children over six years of age, and is sometimes very severe.

Convulsions may occur at the onset in very young children. The *strength* is not usually much diminished, except during the acute period. The *appetite* is diminished and the *thirst* acute, but neither of these symptoms is so marked as in pneumonia. The *tongue* is usually moist, and sometimes covered with a coat of whitish fur; the *abdomen* is natural.

Bilious vomiting occurs in a considerable proportion of cases. The *stools* are generally regular, or there is some constipation.

Urine.—The urine in pleurisy has the so-called febrile characters, but usually not in any degree approaching to the urine in pneumonia, the water being less diminished, and the urea less increased. In cases where there is rapid effusion into the pleura, the chlorides are lessened or almost wanting; and reappear as the effusion is absorbed. Albumen is scarcely ever present.

Course of the Disease.—The cough, pain, fever, and difficulty of breathing continue for several days, after which all but the cough generally disappear, while that commonly persists in a mild form. In acute cases, the appetite now begins to return, the thirst moderates, and auscultation reveals only feebleness of the respiratory murmur, with slight dulness on percussion. The general symptoms cease soon after this, and the patient is entirely convalescent in from one to three weeks, though feeble respiration, with or without friction-sounds, and diminished resonance sometimes persist for a longer period.

CHRONIC PLEURISY may follow the acute form, or occur as an idiopathic disease. In the former case, the acute symptoms diminish after a variable length of time, but the fever does not cease entirely, and often recurs towards evening and assumes a hectic type. In the latter case there is usually a very moderate degree of fever at first, which soon subsides and then disappears, or there is none at all; the pain is generally, though not always, vague, uncertain, and attracts but little notice. In one case that we attended, the cough was frequent, rather dry, and very painful for the first few days, after which it became looser and ceased entirely, though the inferior two-thirds of the right side were filled with effusion for a period of two weeks afterwards. In a second, in which the whole of the left side was occupied by the effusion, there was no cough whatever. In a third, there was a very slight, infrequent cough during the first day, but after that, though the effusion occupied the right side up to the spine of the scapula, there was none through the day, and merely a little hacking at night. In a fourth, in a girl between four and five years old, there was considerable fever during the first week, but literally no local symptoms whatever, so that the case was mistaken for one of bilious fever by another physician. When it came under our notice, some obscurity in the symptoms led us to examine the chest, where we found an effusion occupying the lower third of the right side. The fever was now diminishing, and soon disappeared, but the effusion increased, without pain, and with only an occasional cough, until it filled up three-fourths of the side. It then stopped, and after several days began to recede. At the end of about six weeks the child was quite well again, and continues so to this time, about three years. In a fifth case, in a boy, between five and six years old, the attack was extremely obscure. There was very slight fever, almost no cough, indeed none except upon some exertion being made, and then scarcely noticeable, and no severe pain. In fact, the child complained of no pain whatever, but upon being asked, referred to an uneasy sensation in the inferior lateral region of one side. The tongue was coated, and the symptoms were rather those of some bilious derangement, than of anything more serious. It was not until after four or five days of attendance, that a careful examination of the chest showed the existence of a slight effusion on the right side. This gradually increased until it reached nearly up to the clavicle, and then slowly disappeared again.

The above are mentioned as illustrations, chosen from the very considerable number of cases we have met with, of the great irregularities in the mode of development and general symptoms of chronic pleurisy, and of the danger of errors of diagnosis in consequence. In fact, it is only by the formation of an unvarying habit of making a careful physical exploration of the chest in every case coming under our care, that we can avoid the risk of occasionally overlooking lateral pleural effusions.

The respiration is somewhat accelerated in all cases, and when the effusion is very large, and especially when it is purulent and attended with violent hectic fever, it is sometimes excessively labored and difficult. In some of the cases that we have seen, however, even when the effusion has been very large, the breathing has not been difficult. In one case it was

between 40 and 50 during the first two days, after which it fell, as the effusion took place, to 30. In a second it was 45 at first: at the end of a week it was 38; at the end of the third week, as the effusion was being absorbed, it had fallen to 28, soon after which the recovery was completed. In a third it was so slightly disturbed that we did not at first suspect any disease of the chest. On the fourteenth day, the effusion reaching then nearly to the spine of the scapula, the breathing varied between 40 and 28 during sleep, but during the waking state there was no visible oppression.

The effusion is usually large, excepting in the comparatively rare cases where it is circumscribed by adhesions. The side is evidently enlarged, the increase of size being visible to the eye, and readily ascertainable by mensuration. In addition, the tissues of the affected side seem tense, and the intercostal depressions are obliterated. The heart is pushed by the effusion towards the healthy side, so that the apex-beat is considerably displaced; and, if the right side be affected, the liver is pushed downwards so that its border can be detected by percussion one-half inch or more below the margin of the ribs. On careful inspection it will be seen that, while the respiratory movements of the healthy side are free and even much exaggerated, the side of the thorax on which the effusion exists is almost motionless during respiration. Percussion yields marked dulness over the seat of the effusion. If the pleural sac is not entirely full up to the clavicle, it will be found that, on changing the position of the patient's body, the upper line of percussion-dulness varies its position on account of the gravitation of the liquid. Percussion over the lung above the level of the effusion develops the peculiar pseudo-tympanitic note that has already been described. Quite frequently, if the effusion be large, a slight sense of fluctuation may be obtained by palpation of the intercostal spaces.

Upon auscultation, the respiratory murmur is often suppressed. At other times, a distant and transmitted bronchial breathing may be heard over the seat of the effusion. This can, however, be distinguished from the clear, sniffling, superficial bronchial breathing of lobar pneumonia.

In these cases of chronic pleurisy in children, the effusion is very apt to be purulent, constituting empyema or pyothorax. There are then, in addition to the symptoms and signs of pleural effusion, the evidences of marked hectic fever. The child emaciates, grows pale, and has fever in the latter part of the day, followed by night-sweats. In some cases where the effusion has been at first serous, and later becomes purulent, the development of empyema is clearly indicated by a return of elevated temperature, which now persists with very marked morning remissions and evening exacerbations.

The course and modes of termination of chronic pleurisy differ much in different cases, chiefly in accordance with the character of the fluid. In a large proportion of cases where the effusion is serous, absorption gradually takes place under the influence of treatment in from one to five months, and the patient recovers with a contraction of the side, which eventually disappears as the compressed lung expands. Where, on the contrary, the effusion is purulent, absorption is impossible, and unless paracentesis of

the chest is performed and the pus drawn off, it will be spontaneously evacuated by ulceration through some point of the chest-wall or by an opening into the lung, or the child must die, worn out by the interference with breathing and by the persistent hectic fever. When an external opening forms, it is most frequently in front and in the third or fourth intercostal space. We have, however, met with cases where the opening occurred low down in front or on the postero-lateral aspect of the chest. When an opening forms into the lung, the occurrence is usually announced by a sudden and copious expectoration of pus. The following abstract from our record of a case may be given, as showing the course of empyema when a spontaneous cure takes place by the evacuation of the fluid through an opening in the walls of the chest. This case occurred in a very hearty boy, of between four and five years of age. He was taken sick in the country, with what was supposed to be an attack of typhoid fever. After many weeks of violent illness, an abscess showed itself in the neighborhood of the left nipple. This, at the end of two months, discharged, and the patient began to improve. At the end of three months, he was brought to town, and we saw him. We found a fistulous orifice, discharging occasionally considerable quantities of pus, just below and inside of the left nipple. The left side was very much contracted, and the lung was retracted into the upper part of the chest. He was put upon cod-liver oil, wine, and nutritious food, and gradually improved. He was soon removed to the country, and we did not see him again, but have since heard that he had entirely regained his health.

DIAGNOSIS.—Pleurisy may be confounded with pneumonia or hydrothorax. The latter term is used to imply mere passive serous effusion into the pleural sac, such as occurs in connection with heart disease, or in the course of Bright's disease. The fact that an effusion has resulted from pleurisy will be determined by the history of the case, the acute attack, the pain, and the fever; by the effusion being limited to one side of the chest; and by the absence of the symptoms of those affections which lead to hydrothorax.

The distinction between acute pleurisy and lobar pneumonia is more difficult than that between pleurisy and hydrothorax, and in some instances is subject to considerable doubt. It may generally be arrived at, however, by attention to the differences laid down in the following table, which is taken from the *Bibliothèque du Médecin Practicien*:

ACUTE IDIOPATHIC PLEURISY.

Frequent after six years of age; rare under that age.

Begins with dry cough, sharp thoracic pain, bronchial and metallic respiration during inspiration, either on the first day or later, and more rarely with obscurity of the respiratory sound.

Modification of the physical signs by change of position.

ACUTE IDIOPATHIC PNEUMONIA.

Frequent after six years of age; more rare under that age, but much less so than pleurisy.

Begins with cough, slight thoracic pain, and crepitant or subcrepitant rhonchus; at a later period there is bronchial respiration during the expiration, and bronchophony.

No modification under like circumstances.

ACUTE IDIOPATHIC PLEURISY.

Fever and acceleration of the respiration usually moderate. Rapid diminution of these symptoms from the fourth to the seventh day.

Expectoration absent or very slight.

No rhonchi.

Absence of vibration of the thoracic parietes during speaking or crying.

Course of the disease irregular; rapid disappearance in some cases, prolonged duration in others. The bronchial respiration is substituted or masked by feeble respiration.

In some cases, especially in young children, where the onset of pleurisy is very sudden and acute, the general febrile disturbance may entirely mask the local symptoms, and lead to the belief that some one of the exanthemata is about to develop itself.

Thus we have met with cases where, in the midst of full health, the child has been seized with violent fever; extreme restlessness alternating with stupor; repeated vomiting; great frequency of pulse; acceleration of respiration; but with little or no cough and no complaint of pain in the side. In one instance of this kind, the heat of skin, rapidity of pulse, and frequency of the vomiting were so marked that for twenty-four hours we suspected the approach of scarlet fever, and not until the second day were we able to satisfy ourselves of the nature of the attack by observing that the act of respiration was evidently painful, and by detecting the physical signs of plastic pleurisy over the right apex posteriorly.

In two cases, one at the age of 3 months, the other at $1\frac{1}{2}$ years, we have observed most excessive and almost tetaniform reflex irritability, so that the slightest movement of the child's body, or the attempt to examine the chest, would provoke violent startings and spasmodic contractions of the entire body. In both of these cases a fatal result followed, and post-mortem examination revealed the presence of localized empyema.

The chronic form of pleurisy with extensive effusion may be easily distinguished by the history of the case, by the physical signs which we have carefully detailed, and by attention to the character of the general symptoms.

PROGNOSIS.—Acute idiopathic pleurisy is rarely a fatal disease in healthy children. When, however, it occurs in infants under a year old the mortality is much greater. The danger is also of course greatly increased when the pleurisy is double, or when it is complicated either with pericarditis or with pneumonia. Of 5 cases of primary pleuro-pneumonia, observed by Rilliet and Barthez, 2 died; while of 10 secondary cases, 8 died. A fatal result may follow also when the inflammation is very violent, and

ACUTE IDIOPATHIC PNEUMONIA.

Fever violent; considerable acceleration of the respiration. Diminution of these symptoms less marked, less rapid, and not before the sixth or ninth day.

Expectoration mucous; sometimes sanguineous; very rarely rust-colored.

Rhonchi preceding, following, and often accompanying the bronchial respiration.

Augmentation of vocal resonance very sensible in older children, and in a less degree in all.

Course of the disease regular; steadily increasing in most cases, and then diminishing from the sixth or ninth day. Bronchial respiration more disseminated.

leads to the rapid formation of a large collection of pus; or when the purulent effusion is comparatively small, but is circumscribed by adhesions in such a position that it is difficult to define its precise locality, or to evacuate it successfully. We would specially indicate collections between the under surface of the lung and the diaphragm, or between the pericardial sac and the inner surface of the left lung, as having proved themselves dangerous in our own experience.

Chronic pleurisy is generally a serious, and sometimes a fatal disease, though since the more frequent and more skilful use of paracentesis many cases are cured which would formerly have proved fatal.

TREATMENT.—The *hygienic treatment* in this, as indeed in all the diseases of children, is of the utmost importance, and ought to be regulated by the practitioner himself. In all forms of the disease, the child should be carefully protected from cold, and in the acute form, kept at rest, and if possible, in bed. The diet must be very strict, and should consist for a few days of the preparations of milk. After the fever has disappeared, bread and milk, vegetable soup with a few oysters boiled in it to make it agreeable, and gradually rice, potatoes, and at last small quantities of meat, may be allowed. In the chronic form the diet ought to be nutritious, but regulated with equal care as to quantity and material. In that form the patient should be taken into the air if the weather be mild and dry, and in winter the chamber ought to be well aired from time to time.

Bloodletting.—In acute cases occurring in vigorous children over five years of age, marked by intense fever and pain in the side, and which are seen soon after the onset, local depletion may be employed. The amount of blood to be taken should not exceed two or three ounces; and this should be withdrawn by small cups applied over the seat of inflammation. In younger children, as well as in all whose constitutions are not robust, it is better to limit ourselves to the use of a few dry cups.

Depletion in any form ought to be avoided in most of the secondary cases, unless the symptoms are very acute and the child strong and vigorous; also in all chronic cases, after the febrile symptoms have been dissipated, and in feeble, delicate children.

Antimonials—Febrifuges—Opiates.—A moderate use of the antimonials is of great service in the acute stage of the disease. Small doses of antimonial wine and sweet spirit of nitre, or fractional doses of sulphurated antimony, as recommended in the article on pneumonia, will generally cause the fever, dyspnoea, and cough to subside rapidly. Large doses are unnecessary in any case, and are liable to be injurious in all.

In cases in which antimonials ought not to be used, as where they are opposed by some idiosyncrasy, in children of low vital force, and in the secondary form of the disease, we have found a citrate of potash mixture, containing ipecacuanha and opium, and digitalis, when the heart is much excited, very useful. The quantity of opium must be proportioned to the pain. When this is severe, the doses must be full. The good effects of this remedy in serious inflammations are now generally acknowledged. At two years of age, one drop of laudanum in the above mixture, every two hours; or half a grain of Dover's powder, with the twelfth of a grain

of sulphurated antimony, every two hours, until a decidedly tranquillizing effect is obtained, may be used. When positive drowsiness has been brought about, the doses ought to be given at longer intervals—every three or four hours.

Mercury.—In former years mercury was constantly employed in conjunction with bloodletting. We have, however, long opposed its use as unnecessary in acute cases, but have stated that there was high authority for employing it in cases of the acute form tending towards the chronic, and in confirming chronic cases; adding, however, that we had rarely found it necessary even in these. We find, now, that Dr. West, of London, still speaks highly of it. He says (*loc. cit.*, p. 303): “After depletion, our chief reliance is to be placed on calomel, which should be freely given in combination with opium or Dover’s powder; and an attack of pleurisy thus treated will often be cut off in thirty-six or forty-eight hours.” Dr. J. Lewis Smith, of New York (*loc. cit.* p. 279), does not even mention mercury in his remarks on treatment. Dr. Thomas Hillier (*Diseases of Children*, Amer. ed., 1869, p. 87) says: “Formerly I gave mercury to all cases of primary pleurisy, but this practice I have discontinued, except in the form of an aperient. Instead of it, salines, such as acetate of ammonia, nitrate of potash or soda, the citrate of potash, and nitrous ether, are given.”

The experience we have had, since we last wrote, has not at all increased our faith in this remedy in pleurisy. We believe that as time goes on, and knowledge grows, there is reason to think that the good effects formerly ascribed to calomel in such a variety of diseases, were largely due to the medicines given with it, and particularly the opium (without which it was not often used), the ipecacuanha, the salines, and even the antimonials.

The remedies employed by ourselves, after the disappearance of the acute symptoms, when the effusion has taken place, and especially if there seems any tendency for the case to pass into the chronic form, are either iodide of potassium in syrup of sarsaparilla, according to the following formula:

R. Potass. Iodidi,	gr. xvj ad xxxij.
Syrup. Sarsap. Comp.,						
Aquæ, āā,	f ℥j.—M.
Dose. A teaspoonful three times a day;						

or the syrup of the iodide of iron, of which from thirty to sixty drops should be substituted for the iodide of potassium in such a mixture as the above. The iodide of potassium is preferable in the early stage, and may have the acetate of potassa associated with it. After a time, and especially in anæmic and delicate patients, the iodide of iron should be substituted. Under this treatment, combined with the application of a Burgundy pitch plaster to the side, or some other form of counter-irritant, the effusion has usually disappeared in from two to eight weeks, though diuretics may have failed to make any impression on the cases.

Diuretics are highly recommended in the treatment of cases in which effusion has taken place. Those chiefly employed are squills, digitalis, and nitre. The squill is given alone, or in combination with digitalis, and by some with calomel, or with both. The dose of the powder of squill or digitalis, is about a quarter of a grain every two or three hours. The squill may be used also in the form of syrup or oxymel, and the digitalis in tincture. These two substances may be employed in the following formula :

R. Acet. Scillæ,	℥ij.
Tinct. Digitalis,	gtt. xxx.
Aquæ,	℥iv.—M.

Of this a teaspoonful is to be given three or four times a day to children two years old.

Purgatives ought to be used during the acute stage of pleurisy to an extent sufficient to keep the bowels soluble, and to act as mild evacuants. In chronic cases, on the contrary, they are particularly recommended as evacuants, in order to deplete the bloodvessels, and thus hasten the absorption of the effusion. So far as our experience goes, this treatment is unnecessary, as diuretics and alterative tonics are generally sufficient, without a resort to violent remedies, which must irritate the intestinal mucous membrane, always extremely susceptible in children, to a dangerous degree.

Tonics are often necessary in chronic, and sometimes, after the febrile symptoms have subsided, in acute cases occurring in feeble and delicate children. The most suitable are quinine, in the dose of a grain morning and evening, small quantities of port wine, and the preparations of iron.

External Remedies.—Blisters are very generally employed, in the acute form, to relieve pain and dyspnœa, and, in chronic, to hasten the absorption of the effused liquid. We did not apply them in the cases under our charge, having succeeded very well without ; but we would not hesitate to make use of a small one, applied for a period not longer than two hours, if the pain and oppression persisted. In chronic pleurisy the application of a large Burgundy pitch plaster, made rather weaker than what is used for adults, and large enough to cover nearly the whole side, would be preferable to blisters. We are also in the habit of painting the chest-wall, over the seat of the effusion, with dilute tincture of iodine, every day, or as frequently as the irritability of the skin will permit. The following mixture is of about the proper strength for a young child :

R. Tr. Iodinii,	℥ijj.
Chloroformi,	℥j.
Alcoholis,	℥iv.—M.

Paracentesis.—Of late years, the operation of paracentesis, in cases of pleurisy, both acute and chronic, has been performed so frequently, and with such encouraging results, that it may now be considered to occupy an assured position among the remedies for certain conditions of this dis-

ease. It appears desirable, therefore, to discuss somewhat in detail the circumstances in which it is applicable, the indications which call for it, and to a certain extent the mode of its performance. In doing this, we shall avail ourselves freely of the admirable and exhaustive discussion of this operation by the lamented Trousseau (*Clinique Médicale*. tom. i, pp. 619–698), to whose practice and teaching it was in great part due that paracentesis thoracis was first generally recognized as a justifiable operation for the relief of excessive pleuritic effusions.

In acute pleurisy he recommended the operation more frequently than most authorities consider necessary. Whenever, indeed, the effusion becomes so excessive as to almost entirely fill the pleural sac on the affected side, displacing the adjacent viscera seriously, whether the patient presents intense dyspnœa or not, he advises its performance. The reasons urged by him for this practice were, that although ordinary cases of acute pleurisy almost invariably recover, yet when such excessive effusion exists, it may prove fatal in more than one way. It has not very rarely happened that, from the obstruction to respiration, conjoined with the embarrassment of the heart's action due to its twisting and dislocation, death has occurred suddenly; and we have met with the records of several cases in children which had this unfortunate and unexpected termination.

Again, in these cases of excessive serous effusion, if the fluid be not removed either by absorption or paracentesis, there is great danger that the case will be converted into one of empyema, not from the actual conversion of the serum into pus, but from the altered condition of the secretion from the pleural surface.

But even when the fluid does not become thus converted into pus, but remains clear and serous, absorption is very slow, and the patient may perish from exhaustion and hectic fever. During the long time necessarily occupied in the absorption of the fluid also, the pleurisy really becomes less and less curable, since the lung contracts such close and dense adhesions as prevent it from ever fully expanding again. Finally, if any tuberculous diathesis exists, the long course of the pleuritic attack favors very greatly the development of phthisis.

The chief objections which have been urged against the performance of paracentesis in these acute cases are that the effusion will form again rapidly, requiring repeated punctures and exhausting the patient; that the operation prolongs the duration of the case; and that there is danger of converting the serous effusion into a purulent one.

In regard to the first of these, however, experience has shown that in many cases a single puncture is sufficient, and that even when the fluid does reaccumulate, it is rarely to such an extent as to demand a repetition of the operation,

There is, again, no reason for supposing that the puncture, if properly performed, can in any way tend to prolong the case. In regard to the last objection, the cases recorded sufficiently show that if care be taken to prevent the admission of air, there is not much reason to apprehend the conversion of a serous into a purulent collection, unless the constitutional condition is so impaired that in all probability the case would have passed

into one of extensive empyema, had the operation not been performed at all. Indeed, it is proved by the direct experiments of Nysten and Hewson, that air injected into the pleural cavity does not harm, in the least, the serous membrane.

Since the recent introduction of the greatly improved apparatus for performing paracentesis, also, this source of danger is to a great extent removed. By means of Bowditch's instrument, or, better still, by one of Dieulafoy's aspirators, the effusion can be withdrawn through a canula so fine that its puncture scarcely creates the slightest irritation, and at the same time with entire exclusion of air. In this manner paracentesis has been performed repeatedly of late years, even in the acute stage of pleurisy, without being followed by any of the unfavorable results formerly so much dreaded.

In view of the various risks incurred in cases of excessive hydrothorax, Trousseau thus sums up his remarks upon the operation: "Whenever auscultation and percussion reveal the presence of a very large effusion, whether its formation has been attended with acute symptoms or not, which interferes seriously with respiration, even though dyspnœa is not marked; and when this effusion tends to increase, despite the active employment of local and general remedies for nine or ten days, the operation is indicated." He especially directs attention to the fact that the mere amount of dyspnœa must not be taken as a guide, since this may be absent, although there are at the same time evidences of grave interference with the oxidation of the blood. If, however, during the existence of such an effusion, spells of suffocative dyspnœa should ensue, or syncopal attacks, the operation is urgently called for.

In the *London Hospital Reports* for 1865, these views are warmly advocated and powerfully supported by Dr. Fraser, who believes that the operation should be more generally employed than at present.

We have already alluded to the fact that occasionally an extensive effusion will remain serous for a long time, but in the majority of cases, and especially in children, it sooner or later becomes transformed into pus. Indeed, so frequently does this occur, that West expresses his conviction that in every case of idiopathic pleurisy in childhood, in which fluid is poured out in considerable quantity, the effusion is either originally purulent, or becomes so very speedily. In these unfortunate cases, where there is little or no disposition to absorption, where marked hectic fever and exhausting night-sweats soon set in and rapidly debilitate the patient, and where the most favorable result that can be hoped for is that the pus will either evacuate itself externally, or open into the lung and be expectorated, the operation of paracentesis should undoubtedly be performed.

It is evident, indeed, that paracentesis must here have many advantages, since in cases where the pus discharges externally spontaneously, it is almost invariably about the fourth intercostal space, and outside of the nipple, at a point therefore which renders it impossible for the pus to freely evacuate itself, and which thus tends to keep open the fistula for a very long time. Again, it not rarely happens in these cases that the fistula does not lead directly into the pleural cavity, but that the pus has

burrowed in the thoracic walls, leading to denudation and necrosis of the ribs or sternum.

The termination by the establishment of pulmonary fistula, and the evacuation of the pus through the bronchial tubes, is a comparatively favorable one, but yet the case is apt to be more tedious, and certainly the lung-tissue must be much more seriously affected than when a free exit is given to the matter by the operation of paracentesis. In these cases, of course, the effusion will almost certainly form again, and either require repeated punctures, or a fistula will be established, through which pus will discharge almost daily.

In addition to the advantages afforded by relieving the system of this source of irritation, and giving the lung a chance to expand, paracentesis enables us also to introduce medicated fluids into the thorax, and thus to modify the diseased pleural surface. We will detail below the injections which appear to us most useful for this purpose.

Although, even under the most favorable circumstances, empyema is a most dangerous and not rarely fatal affection, numbers of cases are on record in which life has been undoubtedly saved by a recourse to this operation, and it has been noticed that the proportion of success is much greater in cases of children than of adults. Thus, out of 46 cases in childhood, 13 of which occurred in West's practice, no less than 35 terminated favorably, there being one death in every 4 cases.

In a recent paper by M. Guinier, of Montpellier (*Bull. de l'Acad. de Méd.*, t. xxx, p. 645; *Bien. Retrospect of New Syd. Soc.*, 1865-6, p. 152), the particulars of 31 cases from different authors are recorded. The patients were of all ages up to 14 years; as many as 16, however, were in their 7th, 8th, or 9th year. In one of his own cases, a rapid recovery was effected in a case of extensive sero-purulent effusion in a nursing child one year old. The mortality was about 1 in 6; and in no instance does the operation appear to have done any harm, but, on the other hand, seems to have relieved suffering and retarded death even in the fatal cases.

One reason of this greater success in early life possibly is, that the much greater mobility of the chest-walls in children allows a rapid contraction of the thorax to occur after the pus has been withdrawn, so that the chest-wall comes in contact with the lung, which, in such cases, is always bound down by dense and strong adhesions; whereas, in adults, the more unyielding character of the thorax maintains a space between the two layers of pleura for a much longer time. On the other hand, it must be evident that this same greater mobility of the chest-walls will enable an excessive pleural effusion to be tolerated more readily, and with less injurious effect upon the thoracic organs, than can occur in the comparatively rigid adult chest.

The great deformity of the thorax which ensues upon empyema in childhood is rarely permanent, but as the lung slowly expands, the thoracic walls gradually regain their normal shape, the depression of the shoulder disappears, and, in the course of a few years at the farthest, scarcely any trace of distortion or contraction remains.

Our own opinion in regard to the propriety of this operation, and the

indications for its performance, is as follows: In ordinary cases of pleurisy in children, with moderate effusion, it is unnecessary. When the effusion is very extensive, and causes marked displacement of the heart, distension of the affected side, and severe disturbance of breathing, the question of operating should always be raised, and if, after consultation with the parents, it is determined upon in case of necessity, all preparations for its performance should be made, and we should hold ourselves in readiness to perform it immediately on the appearance of urgent symptoms. Still so long as there is no reason to dread that the case is passing into the stage of empyema, we should recommend a faithful trial for several weeks of the internal remedies, especially digitalis and iodide of potassium, and of the local use of repeated applications of dilute tincture of iodine. In many cases where the effusion has been thus extensive and of quite long standing, we have thus obtained speedy and complete cures, without deformity of the thorax. If, however, positive reduction in the amount of effusion did not soon begin to show itself, we would unhesitatingly operate. Finally, in all cases where the symptoms lead us to conclude that the effusion is more or less purulent, the duty of immediate operation is an imperative one.

In regard to the performance of the operation itself, it may be said to present no difficulty whatever. All the sources of difficulty, and particularly the entrance of air, have been obviated by the improved means of operating lately introduced.

The procedure originally recommended by Trousseau is as follows: The patient being placed near the edge of the bed in a semi-recumbent posture, his body steadied by an assistant, a small incision is made through the skin in the sixth or seventh intercostal space, a little outside of the line of the external border of the pectoralis major. An ordinary trocar, the canula of which is protected by a valve of goldbeater's skin, thin gutta-percha, or a piece of animal membrane of any kind, is then placed in this wound and thrust boldly into the pleural cavity, the precaution being taken of grasping the instrument so that not more than one inch shall be free, to avoid all possibility of wounding the lung.

It is preferable, we think, if a simple trocar and canula be used, that a piece of narrow india-rubber tubing should be attached to the end of the canula, and that the trocar should be passed through from the outside of the tube close to the canula, so that after the puncture into the chest has been made, the trocar may be withdrawn, when the little hole in the elastic tube will close and prevent any entrance of air. The free end of the india-rubber tubing should be carried under the surface of some water placed in a vessel intended to receive the effusion as it escapes. Thus we can simply but surely effect the withdrawal of the fluid without permitting the entrance of air.

These methods of operating have been almost entirely abandoned since the invention of Bowditch's syringe, and better still, of Dieulafoy's aspirator. The latter instrument consists of a syringe, to be attached to the canula after its introduction through the chest-wall and the withdrawal of the trocar, and so constructed that by turning a valve, a vacuum is created

in the barrel when the piston is drawn out. The valve being again turned, the fluid is sucked from the chest into the barrel of the syringe until this latter is filled. A still further turn of the valve establishes a communication with a lateral outlet; the piston is pushed home and the syringe emptied. The vacuum is then renewed, and so the operation is continued until the effusion has been withdrawn so far as desirable. Another improvement in the details of this operation, which has a great influence upon the degree of irritation caused by it, is the use of a very small canula for making the puncture.

It is necessary that the thrust given to the trocar should be fearless and quick, since if it be pushed in a hesitating way, the point may push before it the layers of false membrane which probably coat the pleura, and the effusion will not be reached. Should this accident occur, an attempt may be made to break through the false membrane by a probe introduced through the canula, or a second puncture must be made in a different place.

Different opinions exist in regard to the advisability of withdrawing the entire effusion at once, but experience has, we believe, shown that no unfortunate results need be apprehended from so doing. The last portions of fluid which escape are apt to be stained with blood, probably from rupture of the delicate new-formed vessels of the false adhesions.

The dressing of the wound should be as simple as possible, consisting merely of closing the incision by a piece of adhesive plaster, over which a pledget of lint may be secured by a bandage around the thorax.

Almost the only unpleasant symptom which follows the removal of the fluid is spasmodic cough, which often comes on in severe, and at times painful paroxysms. Syncope is scarcely ever noticed, if the patient be kept in a state of absolute rest after the operation. The internal remedies, especially the diuretics, should be continued, and Trousseau recommends, what we have also found useful, that the side should be painted with tincture of iodine.

When we have reason to believe that the effusion is purulent, which as we have already remarked, is very frequently the case in childhood, there are some points of difference in the operation. Thus we can have no hope that the effusion will not form again, and either require a second operation, or, as frequently happens, cause the cicatrix of the first puncture to reopen. Again, before the case is brought to a successful termination, it is often necessary to employ some medicated injections to alter the character and secretion of the pleural surfaces.

It is doubtful, therefore, whether the admission of a small quantity of air is very objectionable, although West believes that it almost always converts the previous healthy pus into a highly offensive sero-purulent discharge. The ill effects of this can be overcome by the injections to be recommended below; but, on the other hand, care must be taken not to admit so much air as would interfere with the expansion of the lung. It is advisable on the whole, however, to perform the first puncture with the same care, and to employ the same dressing as in the case of serous effusion. We have indeed some evidence to show that, if the operation be

performed with all the details of the antiseptic or Listerian method, it may be possible to lessen the tendency to reaccumulation of the pus. But if a second puncture is required, or if the first one reopens, the wound should be enlarged so as to admit a good-sized canula, which should be allowed to remain. This canula should be of silver, curved so that its extremity may not come in contact with the gradually expanding lung; and its shield should be furnished with a ring of caoutchouc, placed between the instrument and the skin, to prevent excoriation.

After the pus has been withdrawn, the pleural cavity may be washed out through the canula with tepid water, and then there may be injected about an ounce of a mixture of 1 part of tincture of iodine to from 4 to 7 parts of tepid water, effected by the aid of a little iodide of potassium.

The canula should then be closed by a cork, and not disturbed for twenty-four hours, when the accumulated pus should be withdrawn, and a second injection practiced. In the first injections it is better probably to allow the iodine solution to run out again; but after we have assured ourselves of its effect, it may be allowed to remain. Throughout the continuance of the treatment the pus should be allowed to escape at least once every day, though as the secretion diminishes the iodine injections may be practiced only at longer intervals, as once in two, three, or four days. The effects of these injections are usually very beneficial; they correct the fetor of the discharge, diminish its amount, and never, so far as we are aware, are productive of pain or increased inflammation. In cases where they appear to have lost their good effects, other agents may be substituted, as weak solutions of carbolic acid, chlorinated soda, or aromatic wine.

In cases which terminate favorably, the discharge diminishes gradually, though often very slowly, the chest contracts, and finally there is nothing left but a fistula, which for a short time discharges a few drops of serous pus before healing. As an example of the tolerance to this treatment shown even by young children, and of the good results finally obtained in many desperate cases, we would refer the reader to the extraordinary case recorded at length in Trousseau's *Clinique Médicale* (t. i, pp. 650-52), where, in a boy of 6 years, the canula was allowed to remain for eleven months, during which time medicated injections were constantly employed. The amount of pus discharged in all is estimated by Trousseau as not less than 80 pounds, and yet perfect recovery finally ensued, and at the date of the report the child's health was excellent.

Another method of operating in cases of empyema, which we have repeatedly performed with entire success, consists in passing a large curved needle with a stout handle, armed with fine rubber drainage-tube, into the chest through the soft tissues in an intercostal space, and bringing it out through the next interspace above. The needle is then unthreaded and withdrawn; the drainage-tube remains; the spot is covered with a poultice or a wad of oakum; discharge occurs freely through the tube, and it is easy to conduct any subsequent treatment, such as above recommended.

The following case is abstracted from the hospital record, as illustrating the excellent results of this mode of treatment:

Empyema of right side of 8 months' duration; paracentesis; introduction of a drainage-tube, followed by entire recovery.—E. A., a healthy boy of 12 years of age, was attacked in February, 1875, with severe pleurisy on right side. The acute symptoms subsided in 7 weeks, but left him weak and short of breath. On admission to the Hospital of the University of Pennsylvania, the physical signs indicated an effusion on right side reaching up to clavicle, and it was evident from general symptoms that it was purulent in character. On October 13th, 1875, the day of admission, he was tapped by Dr. Pepper in the seventh interspace on line of anterior border of right axilla, and f3xxx pus were withdrawn. A considerable amount still remained, and on October 22d, the effusion having re-formed, a fine rubber drainage-tube was introduced around the seventh rib by means of a large curved needle. A wad of oakum was applied over this and secured in position by a bandage. The amount of discharge was at first very large, but steadily diminished. The chest contracted, the lung expanded partially, the heart returned to its normal position, and in February, 1876, he was sent away cured, the discharge having ceased, and the opening closed after the withdrawal of the drainage-tube. The general treatment consisted of cod-liver oil, quinia, syrup of the iodide of iron, nutritious diet.

In October, 1876, one year after this operation, he was carefully examined. The chest was found to have returned to its healthy symmetrical state, and all traces of the previous disease had disappeared.

During the course of such cases, every attention must be paid to sustaining the child's nutrition by abundant nourishing food, stimulus, if needful, bitter tonics, iron, and cod-liver oil.

We subjoin the following case to illustrate the remarks we have made upon the treatment of pleurisy, and to show the importance of faithfully employing suitable internal remedies before resorting to paracentesis, in cases where the effusion is serous and not so excessive as to seriously embarrass respiration:

Case of chronic pleurisy of the left side, beginning with acute symptoms; extensive effusion, with displacement of the heart to the right of sternum; recovery.—The subject of the case was a boy four years old, of delicate stature and appearance, but enjoying good health. We saw him first at 1 P.M. on February 12th. He was perfectly well the day previous, slept soundly during the night, and rose apparently in good health in the morning. He ate his usual breakfast, but complained afterwards of feeling unwell. Soon after this he complained of headache, of soreness and weakness in the knees in going upstairs, and then of violent pain in the left side beneath the armpit.

At the time of our visit, he was in bed, in the following condition: pulse 130, full and strong; skin warm and moist; headache; sharp, severe pain at the præcordia, extending backwards under the armpit, and aggravated by motion, crying, and by deep inspirations; respiration quick and jerking. No cough at all; absolutely none. Abdomen natural; neither vomiting nor diarrhœa. Tongue slightly furred and moist. Action of heart violent; impulse strong and felt over a large space; sounds loud and strong; to the left, and beneath the nipple, a soft murmur with the second sound. Percussion dull over a larger space than natural.

Behind, percussion dull over whole of left side; natural on right side. Respiration natural on the right side; feeble and indistinct, without bronchial sound, on the left.

Ordered a teaspoonful, each, of extract of senna and syrup of rhubarb, to be given immediately; to have a warm bath in the evening, and to take one of the following powders every two or three hours, beginning in the evening:

R. Pulv. Opii et Ipecac.,	gr. iij.
Potass. Niträt.,	gr. vj.
M., et div. in chart. no. vi.							

February 13. Passed a restless night. Better to-day. Pulse 130, softer; skin moist. Impulse of heart less violent. Pain not so severe. Respiration still quick, and when the child is excited or irritated it becomes jerking, while at other times it is quiet. Physical signs as before, except that the murmur, with the second sound of the heart, is no longer heard. Ordered three ounces of blood to be drawn by leeches from the left side; powders to be continued so as to allay restlessness and pain.

February 14. He had a better night. Pulse less frequent. Respiration 30, and without jerking; no cough at all; makes no complaints of pain. The appetite is returning.

February 15. Better in all respects; no fever nor pain; no cough. Physical signs as before.

The case went on until the 27th of March, when we paid our last visit, making the whole duration of the case over six weeks. During the last two weeks of February there were no acute symptoms. The fever had disappeared entirely. The respiration continued, however, from 28 to 30 during all that time. The effusion occupied nearly the whole of the left side, which was manifestly larger than the right, and the intercostal spaces were protruded. Behind there was total flatness on percussion, from the spine of the scapula downwards, and in front from a short distance beneath the clavicle. The respiratory murmur was absent in the lower three-fourths of the dorsal region, and feeble above. In front respiration was heard only above and just beneath the clavicle. In the course of this period the heart was gradually forced over to the right side of the sternum, so that at last its impulse was felt, not to the left, but to the right of that bone. The cardiac sounds were loudest and most distinct in the same region. The displacement was so remarkable that the mother discovered it herself, as we had avoided telling her to save her from anxiety. The new position of the heart did not seem to produce any inconvenience in addition to that occasioned by the pleuritic effusion. During the last two weeks of March the child was kept in bed; his diet was milk and bread; a large Burgundy pitch plaster was kept on the side, and he took internally vinegar of squill and tincture of digitalis.

Finding that the effusion remained stationary under this treatment, we prescribed a grain of iodide of potassium, three times a day, in a teaspoonful of compound syrup of sarsaparilla. The diet was changed at the same time. He was allowed small quantities of meat every day, and was taken from bed and placed in a chair by the window. Under this treatment he gradually improved, so that by the 27th of March, when we paid our last visit, the effusion had in great measure disappeared, and he was able to play about the room all day. The side was slightly contracted; the respiration was pure and vesicular, but rather more feeble than on the left side; the heart had returned to its natural position.

We examined this child six years later, and found him to be in excellent health. Excepting a slight contraction of the left side, there was no perceptible difference between it and the right one.

ARTICLE VI.

PNEUMOTHORAX.

In this condition there is an accumulation of air in one or both pleural cavities. The source of this air is either from without, when there is an opening through the chest-wall; or from the bronchial tubes, when there is perforation of the pulmonary pleura. There are a certain number of cases recorded, in which it is supposed by the authors that a secretion of

gas has occurred from the pleural surface, or that it has been directly developed from the decomposition of some inflammatory effusion in the pleural cavity; but the evidence upon which the possibility of such occurrences rests is insufficient, and for clinical purposes, at least, it may be assumed that where pneumothorax exists there has invariably been some communication established between the pleural cavity and the atmospheric air. It is therefore to be regarded not so much as a distinct disease as a complication of many other pathological conditions. There are peculiarities, both as to the cause and symptoms of this condition as it occurs in childhood, which render a separate account of it desirable. It is, however, certainly comparatively infrequent in children, owing in part to the rarity of the injuries and wounds, which often cause traumatic pneumothorax in adults, and in part to the fact that the diseases, especially empyema and tuberculosis with the formation of vomicae, which are the most frequent causes of it in adults, are either less frequently attended with this complication in children, or are of comparatively rare occurrence.

ANATOMICAL APPEARANCES.—Pneumothorax may be found to exist on both sides, but as a mere pathological condition which, of course, must have produced death immediately. It is nearly always limited to one pleural sac, and before the thorax is opened the affected side is observed to be distended, with prominent intercostal spaces. The percussion-phenomena, which will be hereafter described, persist, and it is sometimes possible by rapidly moving the body, while the ear is placed in contact with the chest, to develop a succussion-splash. If a small opening be made through an intercostal space, the compressed air will often be heard escaping with a hissing sound, and occasionally the current has so much force as to extinguish a lighted candle held near the opening. The air which escapes is usually, but not always, of offensive odor, in consequence of being tainted by the decomposition of the pleuritic effusion which is apt to coëxist. If the entrance of air has followed a penetrating wound of the chest, or a compound fracture of the ribs, the familiar appearance of these lesions will be found. More frequently it has depended upon perforation of the pulmonary pleura, in consequence of some morbid action, and we may then detect the spot of perforation, and study its characters, by filling the chest with water and blowing through a tube into the trachea, when a stream of air-bubbles will be seen to rise through the fluid from the point of aperture, unless this has been obstructed by layers of false membrane. The lung itself will be found more or less extensively collapsed, according to the nature of the lesion which has caused the perforation. It can rarely be inflated completely in consequence of the free escape of air through the pleural opening. The adjacent movable viscera are displaced by the pressure of the gaseous collection, even to a greater extent than in many cases of hydrothorax. The position of the perforation in the pleura varies, but is most frequently found at some part of the middle lobes, or the adjacent parts of the upper and lower lobes. We have, however, found two points of rupture in one case, both seated near the apex. The opening itself is usually rounded, or occasionally lenticular; in size it varies from one to three lines in diameter. The edges of the pleura are

thin, and often softened and discolored. There may be but a single point of perforation, or several may coëxist, either grouped closely together over an abscess of considerable size, or scattered over the surface of the lung (as in the case on the following page), each opening corresponding to a distinct abscess.

The condition of the lung varies exceedingly, and, of course, presents the appearances proper to the lesion which has caused the pneumothorax. Thus there will be found in about the following order of frequency, the appearances, elsewhere described in their appropriate places, of tuberculosis of the lungs (either in the form of softening sub-pleural miliary formations, or of small superficial vomicæ); of small superficial abscesses resulting from lobular pneumonia; of circumscribed apoplexy or gangrene of the lung; or of vesicular and interlobular emphysema with sub-pleural bullæ. In most cases, there are evidences of pleurisy associated, and the pleural cavity contains a variable quantity of fluid, either turbid or bloody serum, or ichorous pus; and, at the same time, the surfaces of the pleura may present patches or layers of organized lymph. Of course, these appearances will be most marked in cases where the perforation of the pleura has resulted in consequence of a previous empyema. In other cases, the fluids found in the pleural sac have in part escaped from the lung-tissue through the perforation, and are in part due to the pleuritic inflammation superinduced. The irritation caused by the mere admission of air into a healthy pleural sac is not always sufficient to excite inflammation, and thus in rare cases, where pneumothorax results from the rupture of an emphysematous bulla, the pleura may present no inflammatory exudation whatever. But in the great majority of cases, either from the fact that pleurisy coëxists, or that there is an escape of pus from the lung at the time of perforation, the pleura presents the appearances above described. It occasionally happens that, owing to the previous existence of pleuritic adhesions over part of the lung, the escaping air is circumscribed, and produces only a local pneumothorax. In such cases, of course, all the alterations of the thorax, as well as the attendant physical signs, are limited to the seat of the gaseous collection, and may, indeed, be associated with the evidences of chronic pleurisy, with retraction of the remaining parts of the chest. In still other cases, although the pleura has been perforated, the escape of air is entirely prevented by the existence of adhesions of the pleura over the point of rupture; or, as in the interesting case reported on page 226, by the close apposition of enlarged bronchial glands.

The following case, which, owing to the absence of any clinical history, possesses chiefly an anatomical interest, may be given, as showing the usual conditions of a pneumothorax dependent upon pneumonia:

CASE.—Partial Suppurative Pneumonia: Superficial Abscesses, with Sub-pleural Emphysema: Perforation of the Pleura: Pneumothorax: Miliary Tuberculosis.—Mary McC., aged 13 months, died February 11th, 1868, after a short illness, during which the most marked symptoms were dyspnœa and cough, with occasional vomiting. At the autopsy, there were all the physical signs present of pneumothorax of the left side. The *right* lung was found congested and partially collapsed, but admitted of complete

inflation. In the posterior part of the fissure between the upper and middle lobes, the upper lobe presented a separation of the pleura from the lung to the extent of half an inch in diameter. On the apposed portion of the lower lobe there was a similar large bulla. The lung-tissue immediately subjacent to these cavities was consolidated to a distance of an eighth of an inch. On cutting into the bullæ, they were found to be distended with air and dark sanious pus, and their cavities presented minute trabeculæ and septa, consisting of bronchioles and the remains of ruptured air-vesicles. There evidently was no gangrene of the lung-tissue, so that it appeared that these lesions had resulted from a combination of patches of suppurative pneumonia of the superficial layer of the lung with sub-pleural emphysema; and it seems reasonable to conclude from the unusual relations of the emphysematous bullæ, that they were due to the process of softening, which had opened a connection between some of the terminal bronchioles and the sub-pleural connective tissue. There were scattered miliary tubercles in the upper lobe.

The *left* lung presented two similar but larger bullæ (fully one inch in diameter), in exactly the corresponding position between the upper and lower lobes. There was a small perforation of the pleura in the one in the upper lobe. Two other similar but smaller cavities were found on the surface of the lower lobe, in each of which the separated pleura presented a perforation about one-sixth of an inch in diameter. There were traces of localized pleurisy in the neighborhood, but no adhesions; and a considerable pneumothorax had resulted, causing collapse of at least one-half of the lung.

The bronchial glands, spleen, and kidneys contained miliary tubercles. There was no decomposition of the tissues.

Case of Pneumothorax at Nine Months of Age.—We saw, October 27th, 1873, in consultation, a boy nine months old. After careful examination we found that there was dulness on percussion over the lower two-thirds of the left side of the thorax. The expiratory sounds were feeble, and there was also indistinct bronchial breathing. No crepitus or friction-sound could be heard. The right lung was healthy. The pulse was 140, and the respiration 50 to 60. There was much cough, which was evidently painful. We saw the child several times subsequently, and the physical signs continued much the same, but more marked. The case was supposed to be one of plastic pleurisy. Iodide of potash, with citrate of potash and laudanum, were administered; and milk-punch and beef-tea were given as food.

We were called in again on November 6th, as the child had suddenly become worse. The respiration was labored and gasping, 70 to 80, and the pulse 168. The left dorsal region from the scapula downwards was tympanitic, and there was marked amphoric respiration over the same region. The apex-beat of the heart was slightly displaced to the right. Pneumothorax was diagnosed, and the child died at 9 A. M., November 7th.

A post-mortem examination was refused, but we were allowed to puncture the chest with an aspirating needle. A small needle was prepared with a piece of india-rubber tubing attached. The open mouth of the tube was placed in a basin of water, and the needle inserted in the sixth interspace in the line of the axilla. Air escaped freely through the water, the force of the escape being increased at once by pressure on the chest-walls. The needle was withdrawn, and reinserted on the dorsal surface between the eighth and ninth ribs. A syringe was applied, and about three ounces of yellow pus of very fetid odor withdrawn.

The case was, in all probability, one of pleuro-pneumonia, with rupture of a pneumonic abscess into the pleural sac.

CAUSES.—Although, as already stated, pneumothorax is a comparatively rare disease in children, it will be found, when present, to occur most frequently in young children (under the age of 5 years), and especially in those of feeble constitution. The causes which directly lead to

its development vary greatly in their relative frequency, as compared with the causes of pneumothorax in the adult.

Thus we find that the most fruitful cause of pneumothorax in children is unquestionably tuberculosis of the lungs. In adults this condition leads to perforation of the pleura usually only after the production of a vomica; but in children, excavation of the lung-tissue to any extent is rare in tuberculosis, and when it does occur is quite constantly associated with such close adhesions of the neighboring pleural surfaces, as would effectually prevent the escape of any air into the pleural cavity, even in event of a perforation of the walls of the cavity. It is found, therefore, that pneumothorax more frequently results from the softening of small superficial tubercles, which involve the pleura and lead to its softening and perforation.

The next most fruitful cause of pneumothorax in children is probably pneumonia, when it passes on to the stage of suppuration with the formation of a superficial abscess, which seems most likely to happen when the inflammation occurs in a localized and circumscribed form. We have ourselves met with four cases which were due to this cause. It is probable that this unfortunate termination is much more frequent in secondary pneumonias (especially those following such diseases as measles, or, as in one of our cases, severe remittent fever), and in a number of the cases, miliary tubercles have been found associated, as in the instance quoted above by us. In such cases the plastic exudation formed on the pleural surface is often too small to prevent the escape of air after the perforation has occurred.

Gangrene of the lung and the softening of superficial patches of pulmonary apoplexy, are mentioned by several authors, particularly by Rilliet and Barthez, as following next in order of frequency. But, according to our own observation, empyema, with consequent ulceration of the pleura and communication with the bronchi, although not so frequent a cause of pneumothorax in children as in adults, yet furnishes more cases than either of the former rare conditions. Occasionally, also, when the purulent fluid in empyema has discharged itself externally by an ulcerated opening in an intercostal space, air has found entrance to the pleural cavity, and produced a pyopneumothorax.

Finally, pneumothorax has been known to follow the rupture of a subpleural bulla in cases of interlobular emphysema. It is especially in such cases that the collection of air may be found without any coëxisting liquid effusion. It is probable, however, that were life to be prolonged after such an occurrence, some pleural inflammation would be established, and lead to serous effusion.

In most cases the actual perforation of the pleura is the result of the progress of the pulmonary disease which has ultimately involved the serous membrane in its course; but it is probable that the rupture may be at times precipitated by any violent effort, particularly by a fit of coughing or severe vomiting.

SYMPTOMS.—In some cases where the antecedent disease is a very grave one, and the strength of the child is greatly reduced, the supervention of

pneumothorax is with difficulty detected, and death occurs from the sudden increase of obstruction to the respiration, before an opportunity is afforded for careful examination.

The occurrence of the perforation is often marked by an abrupt and decided increase of the dyspnœa which has already existed in consequence of the preceding disease. It will, however, be readily understood that this increase in oppression is not of such constant occurrence in children as in adults, owing to the fact that in the former all acute diseases of the chest are apt to be attended with an extreme degree of dyspnœa. So, too, the sharp lancinating pain usually complained of by adults at the time of the development of pneumothorax may be latent, or only revealed by increased agitation, and more hurried, shallow breathing. In some of their cases, Rilliet and Barthez observed a cough which they considered peculiar, and described as "short, frequent, jerking, painful or convulsive, and sharp or piercing;" and a similar cough has been noticed by other observers.

In cases where death does not occur very quickly, and where a careful examination of the chest can be secured, the physical signs of pneumothorax are much more characteristic than the general symptoms. The affected side is distended, and its intercostal spaces bulge slightly. The respiratory movements are overactive on the opposite side to supplement the marked impairment of motion of the affected one. Percussion over the seat of the pneumothorax gives either merely exaggerated resonance or a tympanitic or amphoric sound. Frequently this morbid resonance will be found associated with dulness upon percussion in some parts of the thorax, owing to the coëxistence of consolidation of the lung or of pleuritic effusion. It may also happen that if the distension of the affected side be extreme, the tympanitic resonance will grow more or less flat, owing to the overtension of the thoracic walls. According to the condition of the lung and the character of the opening in the pleura, the respiratory murmur may be absent, or be present as metallic bronchial breathing, or more frequently as pure amphoric breathing. The vocal fremitus has generally been found decidedly diminished. Metallic tinkling has been detected in several instances; it was observed by Barrier to be most distinct during the effort at coughing. We are not aware that a splashing sound, such as can so frequently be developed in cases of pneumothorax in the adult, by succussion, has yet been observed in children.

The adjacent movable viscera are found to be displaced by the pressure of the gaseous accumulation, especially in left-sided pneumothorax, where the dislocation of the heart to the right is very marked. Of course if the pneumothorax be circumscribed, the above physical signs will be limited to the same spot.

It will thus be seen that the symptoms of this condition in children closely resemble those which it presents in the adult; but that in many cases it is impossible, owing to the great agitation of the child, to fully demonstrate their existence,

COURSE; PROGNOSIS.—The course of pneumothorax in children is usually a rapid one. Occurring, as it does, as a complication of some

serious pre-existing disease of the lung, it so increases the respiratory embarrassment as to generally induce death in from a few hours to a few days.

In rare instances only is life prolonged for a few weeks. The prognosis, although regarded by Rilliet and Barthez as, on the whole, less unfavorable than in the same condition in adults, is still exceedingly grave, both from the serious character of the condition itself, and from the grave nature of the disease (tuberculosis, secondary pneumonia, gangrene of the lung, interlobular emphysema) in whose course it occurs as a complication. Rilliet and Barthez observed one case where recovery ensued after the positive signs of pneumothorax had persisted for twenty days in a boy 3 years of age. They regarded the case as originally one of pneumonia.

We have also observed a case, in a boy 11 years old, during an attack of secondary pneumonia, complicating a severe bilious remittent fever, where complete recovery ensued, though after a most violent illness; and it would indeed seem that, with the exception of the comparatively rare traumatic cases, the prognosis of pneumothorax in children is most favorable when it occurs in this connection.

Steffen (*Klinik d. Kinderkrankheiten*, bd. i, p. 137 *et al.*) expresses this opinion also, and places as the next most favorable variety that which is associated with empyema. Although it might be expected that pneumothorax resulting from the rupture of emphysematous bullæ would be of favorable prognosis, on account of the trifling amount of pleural inflammation which often attends that lesion, the fact is that this condition of the lungs themselves is so serious that a fatal result has followed in all cases so far recorded.

In regard to the *diagnosis*, it is quite true that the occurrence of pneumothorax in the course of one of the thoracic diseases which we have seen it may complicate is apt to be overlooked, either owing to the want of symptoms definite enough to arouse suspicion of the development of some new lesion, or to the difficulty of securing a careful physical examination of the chest. When, however, this physical exploration is made with the frequency and care which are demanded in every case of acute thoracic disease, especially when threatening symptoms exist, the characteristic physical signs will be determined, and can scarcely be attributed to any other than the true cause.

TREATMENT.—The management of pneumothorax must be considered always with reference to the primary disease which it complicates, and its occurrence must not be allowed to interfere with the prosecution of the treatment necessary for this. As it is evident, however, that this additional lesion will still further tax the vital powers, and as the only chance of recovery lies in maintaining life till the cause, if curable, is removed and the air absorbed, we would advise that all remedies capable of reducing the strength or disturbing nutrition should be discarded, and that by every means the system should be sustained. In addition, we should recommend the moderate use of sedatives—either in the form of the preparations of opium or hyoscyamus, associated with bromide of ammonium if the cough be very severe and paroxysmal, to quiet agitation and exces-

sive dyspnœa, and to relieve the cough. Great relief will also be obtained by strapping the affected side with strips of adhesive plaster, overlapping each other, and reaching from the spine to the sternum, so as to restrict the mobility of that side to a great extent, and also to exert a considerable pressure upon it.

If the distension of the affected side and the pressure on the surrounding organs be great, and the evidences of impeded circulation and oxidation of the blood are threatening, recourse may be had to puncture through an intercostal space with a very fine trocar. Although the results of this operation must be regarded as palliative rather than curative in most cases, yet as the paracentesis itself is attended with no danger, its performance is to be recommended whenever the signs of pressure from the accumulation of air in the pleural sac become alarming. It is especially in cases where there is a liquid effusion associated with the gas (constituting a hydro- or pyopneumothorax) that paracentesis will afford most relief. In one case of this kind, Hennig performed paracentesis, evacuating a large amount of purulent liquid and gas, with very great relief of the symptoms of oppression. The child, a boy of 4 years of age, lived four weeks after the operation, and then sank from exhaustion.

ARTICLE VII.

HOOPING-COUGH, OR PERTUSSIS.

DEFINITION; SYNONYMS; FREQUENCY.—Hooping-cough is characterized by a hard, convulsive cough, occurring during expiration, and accompanied by long, shrill, and laborious inspirations, which are called hoops. The cough occurs in paroxysms, which are terminated by the expectoration of tough phlegm, and often by vomiting.

The disease is known by various other names, of which the most common are tussis ferina, chincough, and kincough. The *frequency* of the disease is exceedingly variable, as it occurs both in the sporadic form and as a widely prevailing epidemic. Some idea of its frequency may be gained from the facts that, during the five years from 1844 to 1848 inclusive, there were 390 deaths from it in Philadelphia, under 15 years of age, out of a total mortality of 31,162. During the five years from 1864 to 1868 inclusive, there were, out of a total mortality of 76,354, 543 deaths from hooping-cough; a proportion considerably smaller than that during the first period of five years above mentioned; whilst during the five years from 1874 to 1878 there were only 476 deaths from hooping-cough out of a total mortality of 83,682. The irregularity is even more strikingly seen by comparing single years with each other; thus, while in 1862 there were no less than 208 deaths from this cause, there were but 65 in 1867, and 125 in 1875.

CAUSES; AGE.—It occurs generally in children, and may be met with in the first weeks of life; indeed, Watson, in his lectures, mentions a case

where the mother, during the last week of her pregnancy, lived in a house where the disease was prevalent, and her infant hooped the very day it was born. Of 208 cases in children, in our own private practice, 26 occurred in the first year of life, 147 between the ages of 1 and 7 years, and 35 between 7 and 12 years. To be more explicit, we will state that of 188 cases in which the age was accurately noted, 11 occurred in the first six months of life; 9 between 6 and 12 months; 30 in the second year; 17 in the third, 32 in the fourth, 17 in the fifth, 30 in the sixth, 16 in the seventh, 13 in the eighth, 8 in the ninth, 3 in the tenth, and 1 in the eleventh and twelfth years of life each. Of 130 cases in children, collected by M. Blache, 106 occurred between 1 and 7 years of age, and only 24 between 8 and 14. Of 29 cases observed by MM. Rilliet and Barthez, there were 26 between 1 and 7 years, and 3 between 8 and 12. It is stated by MM. Blache, Rilliet and Barthez, and Valleix, to be most common in girls. Of 208 cases observed by ourselves, 106 occurred in boys, and 102 in girls. Some writers have asserted that certain *constitutions* and *hereditary influences* predispose to the disease. So far as our own experience goes, it has seemed to attack indifferently those who were simultaneously exposed to it. The fact of its being propagated by direct *contagion* is proved beyond doubt by numerous observations. We have rarely known one child in a family to be attacked without its extending to all the others not protected by having had the disease previously. That it often appears also in the form of an *epidemic*, is established by the testimony of many writers, so that at present no doubt is entertained upon this point.

SYMPTOMS.—It is customary to describe three stages of hooping-cough. The first is called the stage of invasion, or the catarrhal stage; the second the stage of increase, or the spasmodic stage; and the third the stage of decline, which is characterized by an amendment of all the symptoms.

First Stage.—The great majority of the cases begin with the ordinary symptoms of simple catarrh. These are coryza, sneezing, slight injection of the conjunctivæ, and dry cough. The cough rarely has any peculiarity in the beginning which will enable us to distinguish it from that of an ordinary cold, though some persons have asserted that they could recognize it. We have often listened with great care to the sound of coughs which parents supposed might be hooping-cough, but were always obliged to confess our inability to determine, until time gave them more decided characters. In addition to the symptoms enumerated, there is generally more languor, lassitude, drowsiness, and irritability, than are commonly present in simple catarrh. In a small proportion of cases the first stage is wanting, and the disease assumes its peculiar features from the first. The duration of this stage is very uncertain, and is ascertained with difficulty. Our own experience would fix it at about two weeks as the average, though it may last undoubtedly a much shorter or longer period. The earliest period at which we have known the distinctive hoop of the disease to be heard was in three days. In another case it was five days. We have also known it to appear at a later period than usual. In a good many instances it has been as late as three weeks, but very rarely later.

Second Stage.—At the beginning of this stage the disease has assumed its peculiar convulsive and paroxysmal character. It consists of violent fits or paroxysms, or as they are often called, kinks of cough, recurring after longer or shorter intervals. Just before the paroxysm the child seems restless, anxious, and irritable, or else keeps perfectly quiet and evidently tries to retard its approach. When it begins, the child, if lying down, rises up suddenly, or if playing about runs to take hold of some fixed object, by which to support itself during the accession. The cough is dry, spasmodic, and sonorous, and occurs in a succession of short, rapid expirations, by which the thorax seems to be emptied of all its air with violent efforts. It is followed by one or two long and deep inspirations, which are accompanied by the peculiar hoop to which the disease owes its name, and which is caused by the drawing of the air rapidly and forcibly through the narrowed glottis, which is spasmodically closed. During the fit the face becomes deeply suffused or even purple, and swollen; the eyes are watery, and the countenance is expressive of great anxiety, and after the fit is over, of fatigue and exhaustion. The latter symptoms are, as M. Valleix remarks, the signs of partial asphyxia, and are the result doubtless of the complete expulsion of air from the thorax, and a consequent partial suspension of the function of hæmatisation. There is almost always an expectoration of colorless ropy fluid, often accompanied by vomiting, at the close of the fit of coughing, and the patients usually appear weak and languid for a short time, after which they return to their play.

In very severe cases there are other symptoms in addition to those just mentioned. *Hæmorrhages* from the mouth, ears, nose, lungs, and beneath the conjunctivæ, are not unusual. We have ourselves seen several instances of epistaxis, one of effusion into the eyelids, a few of extensive subconjunctival ecchymosis, and we are well acquainted with the history of another case, in which there was bleeding both from the nose and ears. In one case, in a girl between and five six years old, that occurred to one of ourselves, in which the paroxysms were violent, the spells were accompanied in the latter half of the fourth and in the fifth week, by a discharge of a good deal of blood from the mouth. This took place particularly during the night-spells, so that in the morning the basin would contain several teaspoonfuls of blood. It was not from the nose. It was bright in color, pure, except that it was intermingled with sero-mucous expectoration, but it was not intimately blended with the sputa, nor was it streaked through the mucus as it sometimes is in the pneumonia of children. On one occasion it was seen to fly from the mouth in a little spirt, as though from a vessel. The child was lively and well at this time, playing about, eating well, strong, not thirsty, without pain, not oppressed between the spells, and sleeping naturally between the paroxysms at night. The only altered physical sign was slight dulness on percussion over the upper part of the right lung behind, with some subcrepitant râles at that point, but without bronchial respiration. After lasting twelve days, it ceased; the child got well gradually, and continues strong and hearty to the present time. In another case, in a girl two years of age, which came under our own observation, a species of syncope, a state of insensibility without convulsive

movements, accompanied by great paleness, occurred after many of the paroxysms.

We have met with general *convulsions* in 12 cases, 5 of which proved fatal. In 2 other cases, both occurring in infants under six months, the paroxysms of cough were accompanied by the most violent struggling and oppression, and by deep blueness of the hands and feet, like that of severe cyanosis.

In some instances, after the paroxysm is apparently over, the child will begin within a few instants to cough again, and may in this way have several fits in such rapid succession as to make an almost continuous paroxysm. It is quite common for this to happen twice, and in one case which we saw, it occurred three times on several occasions. The ordinary *duration* of a paroxysm or kink, is from one-quarter to three-quarters of a minute, though it may last as long as two minutes, or according to some even longer. In a case that occurred to ourselves, one paroxysm lasted the extraordinary period of fifty-five minutes. That it was really a paroxysm of the disease, we are quite sure, as it chanced that we reached the house shortly after it began, and witnessed the greater part of it ourselves. The number of accessions in twenty-four hours is very irregular. It depends chiefly on the stage and violence of the attack. During the height of the disease, we have generally found them to number about 40. In some rare cases, however, they are much more numerous, and amount to 70 or 80. They are generally most frequent in the course of the third or fourth week, after which they remain stationary as to frequency for several days, or for two or three weeks, and then decline gradually. The paroxysms may occur spontaneously, the child being often disturbed from sleep by their sudden occurrence, or they may be excited by various circumstances, such for instance as contrarieties, a fit of crying, change of position, eating, violent exercise, and imitation. We have frequently seen an attack brought on by the sight of another child in a paroxysm of the disease. The duration of the second stage may be stated to be about 30 or 40 days in most cases.

Third Stage.—It is impossible to fix a precise limit from which to date the beginning of this stage. It is generally, however, said to commence from the time when the disease is evidently on the decline. The paroxysms now grow less frequent and less violent, the cough reassumes some of the catarrhal features which it had at first, and gradually loses its peculiar spasmodic character. The child's general health improves, the appetite becomes vigorous, the strength is invigorated, the sleep again becomes sound and tranquil, and the disease disappears. The *duration* of this stage is uncertain, like that of the two others. MM. Rilliet and Barthez state it to be short in uncomplicated cases (ten to fifteen days), and are of opinion that when it has been supposed to have lasted several weeks or months, it has been the result of some complication, as chronic dilatation of the bronchi, tubercular disease, etc. It happens not unfrequently, however, that after the disease has apparently ceased all the distinctive characters of the cough recur, if the child chance to take cold within a few weeks or even longer after its disappearance.

In cases of pertussis unaccompanied by complications of any kind there are no marked *general symptoms*. There is seldom any fever, the appetite continues good, and, with the exception of occasional languor and fatigue and irritability of temper, the child appears to be well.

Urine.—No accurate analyses of the urine in pertussis appear to have been made. Gibb and Johnson, however, state that they have found sugar in variable quantities in almost every case. This question appears well worthy of full investigation, since, if this statement is confirmed, it would link itself in the most interesting manner with the other evidences in this disease of irritation of the pneumogastric nerves, which are at least somewhat concerned in the glycogenic function of the liver.

The *total duration* of the disease, in simple cases, may be set down at from one to three months. We have never known a case to last so short a time as a month, and have rarely found the whole duration much within three months.

COMPLICATIONS.—Though it has happened to us, on several occasions, to meet with children who have been very ill from the violence of the disease under consideration in its uncomplicated condition, we have never known a case to prove fatal except in consequence of some kind of complication. It is exceedingly important, therefore, that the various accidents apt to occur in the course of the disease should be carefully considered.

Convulsions.—This complication is not a rare one, since it occurred in 5 of 29 cases observed by MM. Rilliet and Barthez, and in 12 of 208 observed by ourselves. It is one of the most dangerous accidents liable to occur in the course of the disease. Of the 7 cases reported by the authors quoted (5 of their own, and 2 belonging to M. Papavoine), 6 died. Of our 12 cases, 5 died. In all that we have seen the convulsions were general, extremely violent, and accompanied by insensibility in the fatal cases to the last, and in the favorable ones during from a few minutes to half an hour. In two of the fatal cases the pertussis had lasted nearly two months, and was accompanied by extensive bronchitis. The fatal event took place within twenty-four hours from the supervention of the spasms. The subjects were eight and nine months of age respectively. In the third case, the convulsions came on in the seventh week of the disease, in a child who had been laboring for a number of days under bronchitis. They ended fatally in seven hours. In the fourth they occurred in a child in the second year of its age, at the end of about four weeks, proved fatal in two days, and were caused by bronchitis and collapse of the lung-tissue. In the fifth case they occurred likewise in a child in the second year of life, were attended with violent laryngismus and contraction, and proved fatal in the third week of the disease.

One of the favorable cases occurred in a child five months old, who had been attacked with bronchitis three days before the occurrence of the convulsions, which came on during the height of a severe paroxysm of coughing. The convulsive movements were general, and continued for about half an hour, after which the child was drowsy or irritable for some hours longer. The hooping-cough continued to be severe for two weeks

after this, as many as 42, 46, and 48 paroxysms occurring every day. At last, however, perfect recovery took place. The second favorable case was that of a girl between two and three years old, in whom a convulsion occurred in the third week of the disease, before the paroxysms had become violent, and evidently in consequence of an attack of fever dependent upon dentition. The seizure lasted only a few minutes, was followed by drowsiness for a few hours, but on the following day all the unpleasant symptoms had disappeared. In a third case, in a boy between two and three years old, a violent convulsion occurred at the end of the second week, at the beginning of an attack of pneumonia. The child remained very ill, and nine days afterwards had another convulsion, which was much slighter than the first. After this he gradually recovered. In a fourth case, in a girl between two and three years old, a slight but well-marked convulsion occurred at the onset of an attack of bronchitis, which took place at the beginning of the third week of the whooping-cough. The bronchitis proved to be very severe, but there was no return of the spasm, and the child recovered. In a fifth case, in a boy nine months old, a severe fit occurred in the sixth week, just after the child had been brought home from an expedition to procure his photograph. It lasted fifteen minutes, and was attended with total insensibility and purple discoloration of the face, but in half an hour after the patient was nursing well, and was entirely conscious. There was no return of the convulsions, though the disease was very severe after this attack. In the sixth case, also in a boy nine months old, a slight convulsion occurred during one of the paroxysms in the fifth week, but was not followed by any bad consequences.

Amongst the complications ought to be ranked, we think, though this has not generally been done by writers, an excessive degree of the laryngismus which constitutes one of the natural and essential features of the disease. In some children, in fact, and especially in those of a nervous temperament, and in the anemic and debilitated, and likewise, in certain epidemic types of the disease, this laryngismus assumes a degree of severity which is not only distressing but positively dangerous. In one case that occurred to ourselves, in a child who had suffered many months before from laryngismus and contracture, the occurrence of whooping-cough reproduced the laryngismus, and after a few weeks caused death almost instantaneously, at the beginning of a paroxysm, as the child was sitting upon the floor, where it had been placed only a few moments before to play, it having presented before this no very threatening symptoms. In another case, in which we could detect no other complication, the spasm of the glottis was so very violent, that after a few days the spells were attended with convulsions, and very soon ended fatally. In a third, this symptom was so violent that in many of the spells the child ceased for the time to breathe, seemed to faint, became entirely unconscious, and had to be fanned and carried to an open window to be revived; this patient ultimately recovered. In a great many cases, this symptom, without other complication, has been most distressing, and has required particular treatment.

Collapse of the Lung-tissue.—The recent discoveries in regard to the pathological change in the pulmonary tissue called collapse, and especially a consideration of the causes by which collapse is produced, might well lead us to suppose that pertussis, and especially the bronchitis of pertussis, would be very apt to become associated with collapse. Late researches accordingly show that of all the lesions met with in whooping-cough, this is much the most frequent and important. Dr. Graily Hewitt, of London, in a lecture on the pathology of whooping-cough, read before the Harveian Society of London, in 1855, shows "that the catarrhal inflammation of the bronchial tubes, which occasions whooping-cough, is, in fatal cases, attended almost universally with collapse of the lungs." He states that his observations were made upon nineteen subjects, whose age varied from four years to one month, the average being eighteen months. "In all, the state of the lungs was carefully noted. The chief lesion found after death was collapse of the lung-substance. The following is a statement of the degree to which this pathological condition manifested itself in the different lobes of the two lungs."

"In the *right lung*, portions of the upper lobe were found collapsed in six cases, and in four more to a less degree. The middle lobe was collapsed, wholly or in part, in sixteen cases. The lower lobe was more or less affected with collapse, in eighteen cases. In the *left lung*, the upper lobe presented the same lesion in fifteen cases, the whole of the anterior tongue-like prolongation being in most of the cases affected. The lower lobe was collapsed more or less in eighteen cases. In seven of the cases, the portions collapsed were also congested, in some to a high degree."

"The test of MM. Bailly and Legendre, viz., the inflatability of the portions of the lung thus affected, was used in almost all the cases; and on that and other grounds, it was determined that the particular part of the lung in question was collapsed and not hepatized."

"It will be at once perceived, that the occurrence of collapse was almost universal; all the cases, with the exception of one, in which there was extensive tuberculization of the lungs, presenting a greater or less amount of lung-substance affected in this manner."

We have had but few opportunities of testing this matter for ourselves by post-mortem examinations; but in one case to which we were called in consultation, that of a boy not quite a year old, this lesion was shown, by autopsy, to be present to a great extent. The child had had the disease during three months with considerable severity. He was thought to be doing well, until he was taken one day a long drive into the country. After the ride he seemed very much fatigued, and that night was seized with very great dyspnoea, increased violence of the coughing spells, and after a short time with general convulsions. We saw him on the following day. He was breathing very rapidly and with much effort, there were a great many subcrepitant râles through the chest, the skin was cool, and about the mouth had a cyanotic tint, and he was unconscious. The same symptoms persisted through the day with occasional convulsive seizures, and on the following day he died. At the autopsy, there was found very extensive collapse of both lungs, as proved both by the anatomical ap-

pearances, and by inflation. There was no pneumonia, and very moderate bronchitis.

Bronchitis has always been supposed to be the most frequent complication of hooping-cough, and there can be no doubt that it is one of the most important. The recent discoveries of the existence and nature of collapse have shown, however, that many of the fatal cases, hitherto ascribed to bronchitis, or to bronchitis and pneumonia combined, must have been cases of collapse, so that large allowances must be made for all statistics collected before the discovery of the true nature of the last-named lesion.

There is, as has already been stated, a certain amount of pulmonary catarrh in every case of hooping-cough. This is a normal element of the disease. To constitute a complication there must be a true bronchitis, an inflammation of the bronchial mucous membrane, sufficient to produce the ordinary symptoms of that disease. This exists in a great many cases; MM. Rilliet and Barthez found it to exist either alone or combined with pneumonia in half of the fatal cases. Of the 208 cases observed by ourselves, it existed to a greater or less extent in 42. In 28 of these it was mild or only moderately severe, and of these all but one recovered. In 14 it was severe and very extensive, or else capillary, and of these 6 died. Of the fatal cases, it was in several no doubt attended with collapse of the lung-tissue. In fatal cases, it has often been found accompanied by continuous dilatation of the smaller bronchi.

Pneumonia, according to the authors above quoted, is about as frequent as bronchitis. When, however, the fatal termination took place soon after the beginning of the disease (18th, 26th, or 27th days) it was not generally present. After these periods, on the contrary, it was almost always observed. As these authors, however, include under the title of lobular pneumonia, many cases of bronchitis with collapse, it is clear that a large number of their cases of supposed pneumonia ought to have been ranged under the head of bronchitis. For our own part, we have met with only five well-marked cases of pneumonia. Two of these occurred in girls of seven and nine years old respectively, one in a girl between one and two years of age, a fourth in a boy between two and three years old, and a fifth in a boy in his ninth year. They all recovered. The degree of danger from this complication is in proportion to the earliness of the age at which the disease occurs, and to the extent of the inflammation.

Emphysema undoubtedly follows or accompanies hooping-cough in some cases. In a considerable proportion of fatal cases the lesions of vesicular, and, less frequently, of interlobular emphysema are discovered. These will be found described in full in our article upon the latter affection, where we have also alluded to the rare occurrence of emphysema of the subcutaneous tissue of the neck, and even of the entire body, as a consequence of the free escape of air into the connective tissue of the lung, and thence into the mediastinal spaces. It is therefore probable that, in cases of pertussis which end favorably, but in which the paroxysms of cough have been severe, a less degree of emphysema occurs, which in most instances speedily passes away after the disappearance of the primary affection. Indeed, as nearly all the children whom we have attended with

hooping-cough, continue under our charge, and as only in a very few cases do any symptoms of emphysema persist, we must conclude either that it less frequently attends pertussis than would naturally be supposed, or else that the lung-tissue soon regains its elasticity, and the over-distension of the air-vesicles disappears. In some instances, and especially in those where chronic bronchitis follows the attack of hooping-cough, all the symptoms of emphysema may gradually develop themselves.

Vomiting is a very frequent incident in pertussis, but ought not to be regarded as a complication, unless dependent on some disease of the digestive organs, or symptomatic of cerebral disease. Where it occurs in simple cases, or in those complicated with bronchitis or pneumonia, it has always seemed to us to be advantageous.

Tuberculosis and *scrofula* are not infrequently found to follow hooping-cough, in cases where a marked predisposition to these conditions exists. The tuberculous affection is most apt to take the form of pulmonary or bronchial phthisis. These sequelæ are frequently observed in hospitals, and among the ill-fed and feeble children of the poor, but are comparatively rare among the better classes of society.

DIAGNOSIS.—The diagnosis of pertussis is difficult only during the first stage of the complaint. It is impossible, indeed, to distinguish, during that stage, between it and simple mild laryngitis, or the mild catarrhal attacks which are so common in our climate. After it has once fairly entered upon the second stage, it is scarcely possible to confound it with any other malady. MM. Rilliet and Barthez state, however, that acute bronchitis with paroxysmal cough is not unfrequently mistaken for pertussis, and we recollect perfectly having made this mistake ourselves, in a little girl, five years of age. The cough assumed so exactly the features of pertussis, that after waiting a few days we announced, authoritatively, the presence of pertussis. Only three or four days after this we were forced to take it all back, for the whole thing had disappeared, bronchitis, pertussis, and all. The patient was entirely well. But the mistake need seldom be made, if it be recollected that in acute bronchitis with paroxysmal cough, the invasion is sudden; that there is violent fever, great dyspnoea, and the physical signs of bronchitis; that the hoop is generally wanting, or feebly marked, and that the disease is violent and rapid in its course; all of which circumstances are widely different from what occurs in pertussis.

The same authors assert that tuberculosis of the bronchial glands gives rise to a cough which may be mistaken for pertussis. The following table extracted from their work will show the differences between the two disorders:

PERTUSSIS.	TUBERCULOSIS OF THE BRONCHIAL GLANDS.
Often epidemic, attacking several children at once; transmissible by contagion.	Always sporadic; non-contagious.
Three distinct stages, of which only the second is accompanied by kinks.	No distinct stages.
Kinks attended with hooping, ropy expectoration, and vomiting.	Kinks generally very short, without hooping, ropy expectoration, or vomiting.

PERTUSSIS.

Pure respiration in the intervals between the kinks.

In the intervals between the kinks, respiration and pulse natural, so long as the disease is simple.

Voice natural.

Course generally acute.

TUBERCULOSIS OF THE BRONCHIAL GLANDS.

Physical signs of tuberculosis of the ganglions; but, in certain cases, absence of these signs.

Accessions of asthma in some cases, with the kinks; continuous febrile movement, with evening exacerbations, sweats, progressive emaciation, etc.

Voice sometimes hoarse.

Chronic course.

We would add that we have known the paroxysmal cough attendant upon the development of miliary tubercles in the lungs to simulate whooping-cough so closely as to render it very difficult to distinguish them. In one case where this resemblance was very great, the circumstances rendered the diagnosis additionally difficult. The patient, the eldest of three children, had a perfectly well-marked attack of whooping-cough in his fourth year. Two years later, his little brother and sister contracted the disease in a marked form. Nearly at the same time, the eldest boy, then about six years of age, was attacked with severe paroxysmal cough, recurring in kinks, often inducing vomiting, and occasionally terminating with a sort of hoop. Physical exploration of the chest yielded only negative results. Still our suspicions were aroused by the facts that a previous attack had occurred, that the hoop did not become so perfectly developed as in true whooping-cough, and that there were progressive emaciation and weakness to an unusual degree. In the second month of the cough, the symptoms of tubercular meningitis appeared, death ensued, and, at the autopsy, in addition to tuberculosis of the membranes of the brain, the lungs and spleen were studded throughout with numerous miliary granulations. In the meantime, the two younger children passed successfully through the stages of whooping-cough and entirely recovered.

PROGNOSIS.—Pertussis is rarely a dangerous or fatal disease so long as it remains simple. Of the 208 cases observed by ourselves, 143 were simple, all of which recovered. Nevertheless even the simple disease does something terminate fatally, from the excessive violence of the paroxysms of coughing.

The danger in whooping-cough, which is considerable, depends, therefore, almost entirely on the complications which are so apt to occur, for which reason the physician should watch with the closest attention, in order to prevent their occurrence, and that he may recognize and treat them in their earliest stages. The most dangerous complication is convulsions, and after that bronchitis and pneumonia. So long as the child seems well and lively, and without fever or dyspnoea, in the intervals between the fits, there is nothing to be feared. But if, on the contrary, it becomes languid and irritable, with indisposition to take food, feverishness, and some increase of the rate of respiration, the practitioner should be upon his guard. A very early age and a natural delicacy of constitution, are unfavorable circumstances in the disease. Some form of complication occurred in 65 of the 208 cases observed by ourselves. Of the 65, 12 died.

Five of the 12 fatal cases ended with convulsions. Of these 5 cases, the convulsions were caused by bronchitis and collapse of the lung in 4, the fatal result being the consequence, in fact, of the lung complication. One of the cases was independent, apparently, of disease of the lung (though, as no post-mortem examination was made, this cannot be asserted positively), but seemed to be the result of the violent laryngismus, with contracture and general convulsions, such as will be described in the article on laryngismus stridulus. Two of the cases occurred in children of eight and nine months old, respectively, and proved fatal in twenty-four hours after the setting in of the convulsions. Two others occurred in children in their second year, and the fifth occurred in a boy between three and four years old, and caused death in seven hours.

Of the remaining seven fatal cases, one was the result of collapse of the lungs, supervening suddenly upon a mild bronchitis, in a twin child between two and three months old. The second was caused by tubercular disease of the lungs, in a child between three and four years old, and the remaining five by bronchitis, associated, to a greater or less extent, in all probability, with collapse of the lung. Of the last-mentioned five cases, one occurred in a child between five and six months old, and was rapid in its course; two occurred in children between one and two years old, one being rapid and the other lingering in its course; one occurred in the third year of life, and was attended with severe diarrhœa from teething, as well as with bronchitis and collapse; and the fifth occurred in a child in its fourth year, and was slow and gradual in its course. To sum up, it may be stated that of the 12 fatal cases, 10 were the result of bronchitis and collapse, 1 of tuberculosis of the lungs, and 1 of laryngismus stridulus.

NATURE OF THE DISEASE.—There is no essential anatomical lesion in pertussis, except, perhaps, slight inflammation of the bronchial mucous membrane. In most of the cases, the membrane lining the larger and smaller air-tubes, and very rarely that of the trachea, is reddened and perceptibly thicker than natural, and the tubes contain a considerable quantity of frothy mucus, or a thick, viscid, and tenacious phlegm.

As to the nature of the disease, it seems to us very clear that it ought to be regarded as comprising two elements of morbid action, one of which consists in slight inflammation of the respiratory mucous membrane, and the other of disordered action of the respiratory system of excito-motor nerves. It is neither a pure neurosis nor a pure inflammation, but partakes of the characters of both, and much more of the former than of the latter. The authors of the *Compendium de Médecine Pratique* (t. ii, p. 526) regard it as a neurosis, on the following grounds; “1. In the greater number of cases the respiratory apparatus presents no kind of alteration, or else the lesions are so multiplied or variable that they are surely not the real origin of the disease. 2. The clearly remittent course of the symptoms, and the total absence of fever, unless some complication is present, are not observed in ordinary or even specific inflammation. 3. The cessation or sudden return of the paroxysms, under the influence of moral emotions or change of place, belong to a disorder of innervation, and not to inflammation, which commonly passes through certain stages

before it is resolved. 4. The complete return to health, the integrity of all the functions in slight cases, the resistance which it opposes to treatment, the uselessness of antiphlogistics, and the success obtained from narcotics and antispasmodics, are all so many circumstances peculiar to whooping-cough and to many of the neuroses."

It has, however, so many points of resemblance to the various constitutional diseases, as its undoubtedly contagious nature, the facts that it runs a definite course, and that one attack protects the system against a second, that it also probably depends upon a morbid state of the blood, due to the introduction of some specific poison which possesses the peculiar power of irritating the pneumogastric nerves.

TREATMENT OF SIMPLE PERTUSSIS.—Whooping-cough, like all other diseases, varies greatly in its degree of severity. It is sometimes an affair of no consequence scarcely, the patient passing through its stages without suffering, and without any injurious consequences whatever to the general health. We have known a large family of children to pass through the disease without other treatment than attention to a prudent hygiene, and with no other medicine than a few doses of a mild cathartic, given to relieve some uncomfortable gastric symptoms. We have known one child in a family where the disease was prevailing at the time, to have the cough for only five weeks, and to hoop only on two or three occasions, and to lose neither appetite nor spirits for a moment. Such cases evidently need no interference, and a wise physician will, in such, order no drugs. His business will be simply to direct that the child be guarded against cold and against imprudences in diet.

In other instances the disease assumes, from a very early period, or sometimes not until later, a character of a very different kind. Without any complication whatever, the natural symptoms of the disease develop in great intensity. The spells of coughing are very frequent, very violent, and very long-continued. Instead of some twenty spells or less in twenty-four hours, as is the rule in mild and moderate cases, the patient will average two or more every hour, having fifty or sixty spells in the day. The laryngismus, instead of being slight, will be violent and distressing, so that in lieu of three, four, or five hoops in a paroxysm, there may be fourteen or fifteen, and these so shrill, acute, and prolonged, as greatly to exhaust the poor little patient. Or the laryngismus may be so intense as to close for a few seconds the glottis, and arrest entirely the inspiration, giving rise to the most painful attacks of struggling and suffocation possible to behold. Or the vomiting may be so frequent as seriously to interfere with the nutrition of the child, and thus cause threatening and even dangerous debility. In certain families, and in certain epidemic types of the disease, it assumes these severe features, and such cases must take the same rank in this disease that grave cases of scarlet fever, measles, or variola, take in those affections.

Cases of this latter kind imperatively demand treatment, and they are, we are happy to state, susceptible very generally of great and striking alleviation, by the use of proper means,—means, too, which in themselves are very safe.

At one time we were very much disposed, we confess, to avoid all interference so long as we saw no complication in the case, under the supposition that the disease in its simple form was always safe, and might be trusted to the efforts of nature. More enlarged experience has taught us, however, that the very violence of the disease, even in its simple form, was a source of danger; and that, moreover, such severe cases were much more liable than milder ones to complications, while a proper treatment, instituted so soon as the disease began to show these severe characters, has almost always, after a few days' perseverance, brought about and maintained a most evident amelioration of the symptoms, thus keeping within due bounds a development which might otherwise have gone on to a disastrous termination.

Bloodletting.—Depletion is very rarely necessary in simple pertussis. The only cases in which it can be called for are those occurring in sanguine children, where the laryngismus is so extreme and the paroxysms so violent as to lead to great engorgement of the right side of the heart, and even to endanger the brain by over-distension of the veins. Under these circumstances, we might resort to venesection merely for the mechanical relief afforded, as recommended under similar conditions in pneumonia. In such cases only then, a small bleeding, or the application of a few leeches to the temples or behind the ears, may be proper; but even these may generally be safely treated by reduced diet and by a few doses of saline cathartics, without a resort to the more powerful and more permanently exhausting means of depletion. As for the treatment of simple pertussis by repeated venesections, in the hope of curtailing its duration, or under the idea of their being rendered necessary by the violence of the malady, it seems to us forbidden by the present state of medical knowledge, which informs us that the greater number of the cases do not endanger life so long as they remain simple, however violent they appear to be. Of the 143 simple cases treated by ourselves, depletion was not used in any, and all recovered.

Narcotics and Antispasmodics.—Of the various remedies of this class which have been more or less extensively employed, the most important are belladonna, opium, and hydrocyanic acid. Assafœtida and, of recent years, several of the bromide salts have also been much used with apparent success.

Belladonna is highly recommended by several German authors, by MM. Rilliet and Barthez, who state that it is beyond contradiction the one most deserving of confidence, by Trousseau and Pidoux, and by numerous English and American writers. MM. Trousseau and Pidoux employ the following formula:

R. Pulv. Belladonnæ,	gr. iv.
Extract. Opii,	gr. iv.
Extract. Valerianæ,	3 ss.
M. et div. in pil no. xvi.	
Dose. One to four in the course of the day.	

If the child dislike the pilular form, they give it in syrup, according to the following formula:

R. Extract. Belladonnæ, gr. iv.
 Syrup. Opii,
 Syrup. Flor. Aurantii, āā f ʒj.—M.

Of this, from one to eight teaspoonfuls are to be given in twenty-four hours.

We have ourselves used belladonna in a very large number of cases of whooping-cough, and with such unquestionable benefit, that we regard it as one of the most valuable remedies for this disease in our possession.

We have certainly never seen it cut short the course of the disease as it has been asserted to do, but we have almost invariably found it to moderate the laryngismus, shorten the paroxysms and diminish their number, and probably also shorten the duration of the attack. We have not, however, been in the habit of prescribing such large doses of belladonna as those quoted above (gr. $\frac{1}{4}$); but have usually given it in combination with alum or with bromide of ammonium, in the dose of $\frac{1}{4}$ th of a grain of the extract, every four hours, to a child of one year old. The formula which we employ will be found in our remarks upon the use of alum.

Belladonna has also been largely used, especially by Dr. Fuller, in combination with sulphate of zinc, and with excellent results. This latter author states that he has observed a remarkable tolerance of belladonna in children, so that, beginning with quite large doses, the amount may be rapidly, though carefully, increased until the quantity taken exceeds out of all proportion the corresponding doses which will be tolerated by adults. Even when given, however, in the comparatively small doses of $\frac{1}{4}$ th or $\frac{1}{2}$ th of a grain, it is necessary to watch for any symptoms of the toxic action of the drug, so that its administration may be suspended or the amount diminished.

Opium is confessedly a very valuable remedy in the disease, not as a curative, but as a sedative and palliative. When the cough is frequent and fatiguing, especially if the patient have an irritable and nervous constitution, some opiate preparation is of the utmost service in moderating the frequency and violence of the paroxysms, and in allaying irritability and restlessness. It is best given in the evening, and in combination with ipecacuanha.

Hydrocyanic acid has been employed by various observers, and is highly spoken of by some. Its poisonous properties, however, have deterred many, and amongst them ourselves, from resorting to it. Inasmuch as there are other and safer means for conducting the disease to a favorable termination, it seems to us useless to venture upon so potent a preparation as this.

Since the discovery of the powerful antispasmodic properties of the various bromides, they have been much used in the treatment of this disease. The *bromide of ammonium* has been recommended, especially by Gibb and G. Harley, as a pharyngeal and laryngeal anæsthetic, to diminish the spasm of these parts, while, at the same time, the alkali acts by rendering the secretion from the bronchial mucous membrane more free and readily expectorated. The *bromide of potassium* acts in the same way, and is productive, probably, of equally good results. We have used both of these salts frequently, especially in combination with belladonna, and have ob-

served a marked reduction in the severity and number of the paroxysms of cough in many of the cases. We have, also, used *assafœtida* in a number of instances with decided benefit, both in relieving the general restlessness and in moderating the number and severity of the paroxysms. The doses in which we have given it are either two or three grains in pill, or a teaspoonful of the *mistura assafœtida*, three or four times a day to a child of four years old.

Emetics and *Nauseants* are amongst the most important remedies in the treatment of whooping-cough, since they exert a powerful influence upon the disease, and unless carried to excess, are not in themselves likely to be injurious. Some authors recommend the administration of an emetic every day or every other day, while others give them according to the necessity of the case. Believing that frequently repeated emetic doses are unnecessarily severe, and productive of too much fatigue and exhaustion, we have preferred in the simple disease to give only small doses of *ipe-cacuanha* from time to time, so as to moderate the violence of the cough. Tartar-emetic is never necessary, and ought to be avoided, on account of its disposition to irritate and inflame the gastro-intestinal mucous membrane, and because of its exhausting effects on the general economy. The syrup of *ipe-cacuanha* is the preparation we have almost always used. From ten to twenty drops, given three times a day to a child three years old, will very generally moderate the severity of the paroxysms.

Purgatives are necessary in the simple disease only when constipation is present. The mildest ought to be preferred, in order to avoid irritation and exhaustion. Castor oil, magnesia, or syrup of rhubarb are the best.

Particular Remedies.—Of the different specific remedies that have been employed, none have attained and maintained so high a reputation in this city as the *carbonate of potassa*, which, in the form of the cochineal mixture, is constantly used both by physicians and as a domestic remedy. The beneficial effects of this drug are equally recognized abroad, as may be judged from the language of Niemeyer, who, when speaking of its use in whooping-cough (*op. cit.*, vol. i, p. 101), says: "Its effect in shortening the fits of coughing is often surprising." The following formula is the one generally administered:

R. Potass. Carbonat.,	℥j.
Cocci,	℥ss.
Sacch. Alb.,	℥j.
Aquæ Fontis.,	f ℥iv.—M.

Give a dessertspoonful three times a day to a child a year old. Believing the carbonate of potash to be the active agent in the mixture, we have generally left out the cochineal and used the potash alone, dissolving it in equal parts of syrup of gum and water. We have frequently employed this remedy, and believe that it, with alum and belladonna, are the most useful agents we have to keep down the violence of the disease. We have given it in the dose of a grain three or four times in the twenty-four hours, to children one and two years old, for several weeks at a time, without witnessing any injurious effects from it.

Alum was first highly recommended as a remedy in pertussis by Dr. Golding Bird (*Guy's Hospital Reports*, April, 1845). He states that in the second or nervous period of the disease, when "all inflammatory symptoms have subsided, and when, with a cool skin and clean tongue, the little patient is harassed by a copious secretion from the bronchi, the attempt to get rid of which produces the exhausting and characteristic cough, alum will be found to be of much value." He adds, that he "has not yet met with any other remedy which has acted so satisfactorily, or afforded such marked and rapid relief." From reading Dr. Bird's remarks on alum, and prompted by our knowledge of its admirable qualities in the treatment of croup, we were formerly led to make trial of it in the disease under consideration, and we believe we may say that it has exerted a more decided influence in moderating the violence of the disorder than any that we have ever made use of. We have administered it in 139 cases, beginning in the course of the second stage. In nearly all it was beneficial, and in some the effects were strikingly useful, the improvement being more rapid than we had ever seen to result from other remedies, or to occur when the disease has been allowed to pursue its natural course. In a boy, between five and six years of age, who had been coughing violently for two weeks, the paroxysms diminished so much in intensity and frequency, after he had taken the remedy two days, that he was not once disturbed at night (though before he had always been waked several times), and the spells which occurred during the day were much less severe. After continuing the remedy for ten days, the disease had subsided so much that its employment was suspended. Soon after, however, the paroxysms again became severe and troublesome. The alum was resumed, and with the same results as the first. In another family, in which there were three children, all of whom had been taking syrup of ipecacuanha and carbonate of potash for some days, without any good effects, the alum was given and acted as in the case first referred to. The nights were comparatively quiet, and the spells occurring through the day very much moderated. We may repeat that, so far as our experience in the above 139 cases goes, the effects of alum have been more decided and satisfactory than those of any other remedy. We have never known it to produce ill-consequences, either at the time of its administration or subsequently, though we have given it to children from two months to seven years of age, and have continued its use from one to six weeks at a time. It, like all other remedies, sometimes fails, however, to do any good, and when we have found this to be the case, we have substituted belladonna or carbonate of potash, either alone or combined, and it is curious to observe how, in some instances, the latter remedies will succeed when the other fails. Nothing but a trial will show which is the most proper in any individual case. Of late years we have usually given the alum and belladonna together, and have been much pleased with the results. If administered in large doses, alum produces vomiting. It does not constipate, but, on the contrary, is apt to induce diarrhœa when continued for some time. Dr. Bird gives from two to six grains every four hours. His formula is as follows:

R. Aluminis,	gr. xxv.
Ext. Conii,	gr. xij.
Syrup. Rhœados,	f℥ij.
Aquæ Anethi,	f℥ij.—M.

Give a medium-sized spoonful every three hours.

To children under one year, we give from half a grain to a grain, three or four times a day; and to those over that age, two grains every four or six hours. The formula we have employed is the following :

R. Aluminis,	℥ijss.
Syrup. Zingib., Syrup. Acaciæ, Aquæ Fontis, āā	f℥j.—M.

When this is prepared with good syrups, it tastes very much like lemonade, and is not at all unpleasant, so that children take it without difficulty. The dose is a teaspoonful three times a day, or every four or six hours.

As above said, however, we now generally employ a combination of alum and belladonna, and have obtained better results from it than from any single remedy we have ever used. For a child one year old we use the following formula :

R. Ext. Belladonnæ,	gr. j.
Aluminis,	℥ss.
Syr. Zingib., Syr. Acaciæ, Aquæ āā	f℥j.—M.

Dose. A teaspoonful four times in the twenty-four hours : in the morning, at noon, bedtime, and once in the night, if the cough be troublesome.

Among other remedies which have been highly recommended, but which we have never found it necessary to resort to, may be mentioned the following :

Sulphur is much used by some German authorities, who greatly commend its effects both at the beginning and throughout the course of the disease. It may be given in powder diffused in milk or syrup, or in emulsion, in doses of three grains, two or three times a day, to children from two to four years of age.

Subcarbonate of Iron has been successfully employed by Dr. Steyman and by Lombard, of Geneva.

Dilute Nitric Acid, first recommended by Arnoldi, of Montreal, has been highly praised, especially by Gibb.

Conium has also been frequently used, both alone and as an ingredient in formulæ containing some of the other remedies here mentioned, and appears to alleviate the violence of the paroxysms, though to a less marked degree, we believe, than belladonna.

Inhalations.—It was noticed in France, some years ago, that children suffering with whooping-cough, who lived in the neighborhood of gas-works, were rapidly cured; and the plan has been recently tried with success, of sending patients with this disease to inhale the fumes arising during the purification of gas, which contain ammonia, vapor of tar, and several volatile oils. Dr. Bertolles (*British Med. Jour.*, Nov. 5th, 1864) states that

"the register of the gas-works at Ternes, shows that during the previous six months, 901 patients have been subjected to the vapor treatment, of whom 219 were cured, and 122 relieved." M. Commerege (*id. loc.*) has also reported the effects observed in 142 children who were brought under the action of the fumes in the gas-works at St. Maudé, and believes that the treatment produces excellent results in all stages of the disorder. In general, twelve séances, each of which should be of two hours' duration, are required for the cure. We have ourselves known of quite a number of instances among the children of the poorer classes in this city, where patients, suffering with whooping-cough, have been allowed to inhale the fumes from the gas-works, and have experienced positive benefit. Of like nature is the inhalation of medicated solutions by means of the steam atomizer; and Dr. J. Lewis Smith (*Amer. Jour. Med. Sci.*, October, 1879, p. 386) reports that he has obtained good results from the use of carbolic acid by this method. He recommends the following formula:

R. Acidi Carbol.,	3ss.
Potass. Chlorat.,	3ij.
Glycerinæ,	3ij.
Aquæ,	3vj.—M.

Sig. Three times daily, 2 to 5 minutes at each sitting.

Tonics.—In a number of cases that have come under our notice, the patient has grown pale and weak in the course of the disease, and this without any local complication, but from the disturbance of the digestive system that often exists to a greater or less extent, from the great frequency of the vomiting, which prevents them from taking a sufficient amount of nutriment, and from the exhausting effect of the violent muscular exertion undergone during the paroxysms. In such instances, when there has been no fever, or merely a little evening febricula, we have employed tonics with much advantage, and never to the injury of the patient. We have generally made use of Huxham's tincture of bark, either alone, in doses of from ten to twenty drops three times a day, or in connection with the syrup of the iodide of iron, or half a grain of the metallic iron (*Pulv. Ferri*). When the appetite has been very feeble, we have found that quinine, in the dose of a grain three or four times a day, at the age of three or four years, has restored it more rapidly than any other remedy we have used.

LOCAL APPLICATIONS.—Revulsives.—The milder revulsives are useful in certain complications of pertussis, and as palliatives. To make them the chief basis of the treatment, however, which has been done by some, is a mistake. In order to produce a decided impression upon the disease, it would be necessary to resort to the more powerful remedies of this class, such as moxas, issues, tartar-emetic ointment, blisters, etc., the use of which is not warranted by the nature of the disorder.

When the laryngismus has been severe, we have known the use of a belladonna plaster, 2 by 3 inches, applied over the larynx and worn for several days, to afford relief.

Topical applications to the interior of the larynx of solutions of nitrate

of silver have been used by several practitioners, as by Gibb and Eben Watson, and apparently with much benefit. The strength of the solution should vary according to the stage of the disease, being much reduced during the early acute period.

In cases occurring in older children, where the spasmodic irritability of the larynx is extreme, we may employ with advantage the inhalation, by means of the atomizer, of the vapor of water, or of weak alkaline solutions, to which minute quantities of morphia may be added.

Before concluding our remarks upon the treatment of simple hooping-cough, we wish to state that cases of the disease occur not unfrequently of so mild a form, as to need absolutely no treatment other than the proper degree of attention to hygiene; and that others again, more numerous than those just mentioned, will be met with, in which the only treatment necessary is the use, for a few days or weeks, of some mild expectorant and opiate at night to lessen the severity of the paroxysms, or of moderate doses of alum, belladonna, or carbonate of potash.

In infants particularly it is proper to give as little medicine as possible, allowing the disease to go on without interference so long as it progresses safely. In a good many mild cases, small doses of paregoric and syrup of ipecacuanha, constitute the only remedies we have found necessary in the cases of infants. When, however, the paroxysms become numerous and violent, exhausting the strength of the child and distressing its nervous system, we must make use of some remedy to allay the severity of the attacks. We have found the alum and belladonna formula recommended above safe and effectual. At the age of two and three months, we have usually given from half a grain to a grain of the former, combined with $\frac{1}{4}$ th grain of extract of belladonna, three times a day, taking care to suspend it for a day or two if it caused troublesome vomiting or purging, and then resuming it in diminished dose. Or we have made use of one-quarter or one-half grain of carbonate of potash, also combined with the twenty-fourth part of a grain of the extract of belladonna, three or four times a day.

TREATMENT OF THE COMPLICATIONS.—If any of the diseases which have been mentioned as apt to occur during the course of pertussis should arise, the treatment which is proper for them in their idiopathic form must be adopted without regard to the hooping-cough, with the following reservation; that care must be taken not to use means of too powerful and exhausting a nature, or such as have a tendency to irritate the organ with which they come in contact. For, it must be recollected, that after the complication is cured, the patient has still the original disease to go through with, and therefore requires all his strength; and moreover, the various organs of the body are predisposed, by the very fact of the existence of the original malady, to assume diseased action, should any irritation in the shape of a violent remedy be applied to them.

The cases of *bronchitis* which came under our observation were treated in the simplest manner. The children were put to bed, the diet carefully regulated, the bowels gently opened with castor oil or syrup of rhubarb, and small doses of syrup of ipecacuanha or antimonial wine, with sweet

spirit of nitre, were administered every two hours. Mustard poultices were applied once or twice a day to the interscapular space, and mustard foot-baths used every night, or more frequently, if the dyspnœa were considerable. If the bronchial secretions were very profuse, and the cough troublesome, the decoction or syrup of seneka was given in connection with occasional doses of laudanum or paregoric.

The treatment of *collapse of the lung* should be that which is recommended in the article on that subject, modified, of course, as may be rendered necessary, by the existence of the whooping-cough. A mild emetic, if the patient seem strong enough to bear one; counter-irritants, and especially sinapisms or mustard poultices applied to the chest, nutritious food, and mild stimulants, as brandy, wine-whey, tincture of bark, quinine, or aromatic spirit of ammonia, must form the principal means of treatment.

The complication of *pneumonia* should be treated somewhat differently. At the present time we should advise the use either of the combination of sulphurated antimony and Dover's powder, or of one of the alkaline mixtures, recommended in the article on pneumonia, in conjunction with external applications and the use of the foot-bath; and should not resort to bleeding, whether local or general, unless the indications, elsewhere laid down as calling for depletion in pneumonia, should be present in a marked degree. Indeed, in such cases, the early use of moderate stimulation and of full tonic doses of quinia is apt to be indicated on account of the tendency to depression.

When *convulsions* occur they must be treated according to the cause which produces them, and the constitution and present state of the child. If the patient be strong and sanguine, and not exhausted by previous sickness, the treatment should consist of depletion by leeches to the temples, or behind the ears; of cold applications to the head; the warm bath; large doses of bromide of potassium by the mouth or by enema; hydrate of chloral by enema; cathartics or purgative enemata; and revulsives in the form of sinapisms, or of a small blister to the nucha. If, on the contrary, the patient is of delicate constitution, or exhausted by long illness, and especially when the convulsions are the result of extensive collapse of the lungs, occurring spontaneously or supervening upon bronchitis, we must be content to resort to warm baths, revulsives, antispasmodics, anodynes, stimuli, and stimulating enemata.

Of the 12 cases of convulsions that came under our notice, 5 proved fatal. Two of the fatal cases occurred in children who had long been laboring under bronchitis, probably associated with collapse, that had baffled all treatment. Death took place within twenty-four hours from the appearance of the convulsions, which were, in fact, the result of the diseased condition of the lungs. No treatment further than the warm bath and sinapisms, was resorted to. In the third case, the convulsions came on in the seventh week of the disease, in a child who had been laboring for a number of days under severe bronchitis; they ended fatally in seven hours. The treatment employed at the beginning of the fit was a warm bath, an enema, and mustard plasters. After a few hours, solution of morphia with

fluid extract of valerian were given by enema, cold was applied to the head, and a blister to the nucha. In the fourth case, which occurred in a child in the second year of its life, they were caused by bronchitis and collapse, and proved fatal in two days. The treatment consisted in the use of warm baths, counter-irritants, alum, and small doses of brandy. The fifth case likewise occurred in the second year. This was one in which all the symptoms of laryngismus stridulus—prolonged laryngismus, contracture, and general convulsions—were added to those of the primary disease. It was treated with belladonna, opium, assafoetida, and warm baths, but all to no effect.

Of the favorable cases, one occurred in a boy five months of age, on the third day of a severe attack of bronchitis. The child was immediately placed in a warm bath, and large sinapisms applied over the front of the chest and upon the extremities, when the convulsions ceased. After this he was treated with half-grain doses of alum, repeated every three or four hours, mustard foot-baths and poultices, and small doses of wine of opium. On the sixth day of the attack, the third after the convulsive seizure, there having been no return of the convulsions, the bronchitis subsided with copious sweats and cold hands and feet, for which small quantities of brandy and water and wine-whey were used. The recovery was perfect. A second case occurred in a hearty boy nine months old, and seemed to depend on congestion of the brain, brought on by a severe fit of coughing. In this instance a venesection to a small amount was performed, the child was placed in a warm bath, and cold applied to the head. No return of the spasms took place, and the child recovered without difficulty. In another case the convulsion was caused by an attack of fever depending on dentition, and was treated by lancing the gums, by a warm foot-bath, and by the administration of a grain of calomel in a teaspoonful of castor oil. In the fourth case the convulsions were caused by pneumonia, and were managed by treating the pneumonia, except that at the moment of the attack a warm bath and a stimulating enema were made use of. In a fifth the convulsion, which was a short one, occurred at the onset of an attack of bronchitis. No particular treatment beyond what was necessary for that disease was required. In a sixth, in a boy nine months old, the convulsion occurred suddenly, was violent, and lasted fifteen minutes. The cause could not be ascertained. The only treatment used for the convulsion was a warm bath. There was no return. In a seventh case, in a boy nine months old, a slight convulsion occurred during one of the paroxysms in the fifth week. No treatment was necessary, as the attack was very short, and there was no recurrence of the symptoms.

HYGIENIC TREATMENT.—This part of the management of the disease is of the highest importance, for it is by careful attention to its details that the complications which constitute the chief danger of the malady are to be prevented. In a considerable number of cases of pertussis, nothing more need be done than to insist upon strict attention to hygienic rules. The chief indications are, to preserve the child from taking cold, and to prevent indiscretions in diet. The clothing ought to be warm, and during the autumn, winter, and spring, flannel should always be worn next to

the skin. The child ought to be kept in the house during damp weather at all seasons, and whenever, during the winter season, it is intensely cold. The diet should be nutritious, but of easy digestion. All heavy, rich food ought to be absolutely forbidden during the continuance of the malady.

TREATMENT OF THE PAROXYSM.—It often happens that the paroxysms are so violent that the child seems to be in imminent danger of suffocation or of convulsions. This is especially true of infants. In six cases that we have seen, in infants under six months old, the kinks lasted so long, and the spasm of the larynx was so unyielding, that the children struggled as though laboring under tetanus; the countenance was disturbed and anxious; the face and hands, at first flushed, became purple from deep congestion; and on some occasions the breathing was suspended for several seconds, so that life *seemed* for the time in the greatest danger. The difficulty in these cases depends on the spasmodic closure of the glottis, which is sometimes, no doubt, completely shut. We have never known these alarming symptoms of asphyxia to occur when the hoop has been clear and distinct, for when that is present, the larynx cannot be very tightly closed.

When the symptoms above described occur in a child several years of age, the patient should be raised and supported in the sitting posture; when in an infant, the child ought to be held lightly in the arms, so that it may take any position which instinct prompts it to. At the same time, cold water ought to be sprinkled from the fingers upon the face, the child should be gently fanned, or, if the weather be warm, taken to the open window; and if there be time, it is well to put the feet into mustard-water. It has been recommended on such occasions to apply compresses dipped into cold water to the sternum. We would propose the trial of a means which the late Dr. C. D. Meigs found very successful in arresting tonic spasm of the respiratory muscles, in a case of laryngismus stridulus. This is the sudden application of a piece of ice wrapped in linen to the epigastrium. When the laryngeal spasm is very intense and obstinate, a belladonna plaster, as before recommended, or a small blister to the front of the neck, may be useful in controlling it.

M. Bell speaks very highly of the results obtained by sprinkling a little ether on the clothes of the patient at the onset of the paroxysm; and Dr. Churchill (*Diseases of Childhood*, p. 223), who has tried ether in 12 or 14 cases, and chloroform in 6, regards it as a valuable addition to our remedies. He directs that about half a drachm of sulphuric ether should be sprinkled on the nurse's hand and held before the child's nose and mouth at the commencement of a fit of coughing. In only one or two cases no benefit accrued, while in others great mitigation of the spasm, and in three or four almost complete relief followed when the ether was thus applied. We should certainly recommend a trial of this procedure, making use, however, from preference, exclusively of the sulphuric ether.

CLASS II.

DISEASES OF THE CIRCULATORY ORGANS

ARTICLE I.

CYANOSIS.

SYNONYMS; DEFINITION.—This peculiar condition, known under the various names of *Morbus Cœruleus*, or the blue disease, and *Cyanosis*, may be defined as a permanent state of lividity or blueness of the skin, depending upon numerous malformations or derangements of the heart and great vessels.

In a comparatively slight degree, this condition attends many of the chronic organic diseases of the circulatory organs, and is also transiently present in the course of some acute diseases; but under neither of these circumstances does the lividity merit consideration as a separate affection, being merely due to the imperfect oxygenation of the blood.

There is, however, one form of cyanosis which we have occasionally met with that merits a special reference. In these cases, the blueness of surface has appeared from three or four days to as many weeks after birth, has been intense in its degree, and associated with marked disturbance of respiration, and yet, under proper treatment, the infants have usually recovered. We believe that the cause of such cyanosis is to be found in atelectasis of the lungs, which acts partly by causing general venous congestion, and partly perhaps by obstructing the flow of blood through the pulmonary artery, so that the right cavities of the heart become over-distended, and there results an admixture of venous and arterial blood through the still unclosed foramen ovale.

Before attempting to explain the peculiar blue color in cases of true cyanosis, it will be convenient to allude to the various lesions which have been found present in such cases.

MORBID ANATOMY.—The blood in cyanosis is dark, and contains an excess of carbonic acid; it has also lost, to a great extent, its coagulability. The only organs beside those of circulation which present lesions, with any constancy, are the lungs.

Dr. J. Lewis Smith (*Dis. of Infancy and Childhood*, 1869, pp. 578–599), who has studied this disease with great care, and collected all the cases of it upon record, finds the condition of the lungs recorded with more or less minuteness in 110 out of 191 cases. In 26 cases there was tuberculosis, either confined to the lungs, or chiefly exhibited in these organs; in 35

cases the lungs were of small size, either from compression by effusion in the pleural sacs or pericardium, or sometimes, apparently, from the persistence of the foetal state over a greater or less portion of the organ. In 35 cases the lungs presented a dark color, owing either to atelectasis or to engorgement and congestion. In 9 there was emphysema in a part of the lungs; in 2, pneumonia; in 2 the color was pale; in 1, a bright crimson; in 1, the lungs were larger than natural; in 1, the right lung was absent; and in 17, these organs were recorded as healthy.

There is also found, in a large proportion of cases, venous congestion of the brain, liver, or kidneys. By far the most marked and important lesions, however, are those of the heart and great vessels, which are, excepting in extremely rare instances, the essential seat of the disease. The number of these lesions already recorded is considerable, as will be seen from the subjoined table, borrowed from Smith, which shows their character and relative frequency.

1. Pulmonary artery absent, rudimentary, impervious, or partially obstructed,	97
2. Right auriculo-ventricular orifice impervious or contracted,	5
3. Orifice of the pulmonary artery and the right auriculo-ventricular aperture impervious or contracted,	6
4. Right ventricle divided into two cavities by a supernumerary septum,	11
5. One auricle and one ventricle,	12
6. Two auricles and one ventricle,	4
7. A single auriculo-ventricular opening; inter-auricular and inter-ventricular septa incomplete,	1
8. Mitral orifice closed or contracted,	3
9. Aorta absent, rudimentary, impervious, or partially obstructed,	3
10. Aortic and the left auriculo-ventricular orifices impervious or contracted,	1
11. Aorta and pulmonary artery transposed,	14
12. The cavæ entering the left auricle,	1
13. Pulmonary veins opening into the right auricle, or into the cavæ or azygos veins,	2
14. Aorta impervious or contracted above its point of union with the ductus arteriosus; pulmonary artery wholly or in part supplying blood to the descending aorta through the ductus arteriosus	2
	<hr/>
	162

It is evident from a glance at this table, that the vast majority of the above lesions must occur before the full development of the heart is attained; and that consequently, in nearly every instance, cyanosis is a congenital affection. But further than this it will be observed, that in the first four groups in Smith's table, or in 119 out of 162 cases, the lesions affect the right side of the heart, and are precisely of the kind that we know are caused by inflammation of the endocardium. Bearing in mind then the well-ascertained law, that endocarditis occurring during foetal life, almost exclusively attacks the right side of the heart, we can readily understand how such lesions could be produced by an attack of inflammation affecting either the valves of the pulmonary artery, or the tricuspid valves, or some part of the lining membrane of the right ventricle. Should

such an attack of endocarditis occur after the development of the cavities and septa of the heart, and the closure of the foramen ovale and ductus arteriosus, and lead to occlusion of the orifice of the pulmonary artery, it would of course be impossible for life to be sustained. But where such a lesion is produced while the inter-auricular and inter-ventricular septa are still imperfect, and the ductus arteriosus patulous, so much compensation may be effected that life can often be prolonged for many years. Thus, it is evident, that the first effect of the closure of the orifice of the pulmonary artery, at such an early period, will be to cause a large portion of the blood from the right ventricle to pass directly through the opening remaining in the inter-ventricular septum into the left ventricle. Usually this opening is not free enough to relieve the right ventricle entirely, and there is consequently pressure exerted backwards on the blood entering from the right auricle, which forces part of it through the foramen ovale into the left auricle, and thus still further relieves the fulness of the right cavities. As there is no outward current through the pulmonary artery, owing to the occlusion of its orifice, blood flows back into this vessel from the aorta through the patulous ductus arteriosus, and thus supplies the lungs. At the same time the bronchial arteries become much enlarged, and, in some rare cases, blood has been able to reach the lungs through abnormal branches from the internal mammary or intercostal arteries. In cases of cyanosis which prove fatal very soon after birth, the most diverse and inexplicable lesions, as before enumerated, may be found; but in those instances where life is prolonged, the heart is usually found to present the associated lesions aboved described: contraction or occlusion of the orifice of the pulmonary artery, imperfect inter-ventricular septum, and patulous foramen ovale and ductus arteriosus. In such cases, when the compensatory communications between the right and left side of the circulation are free, life may be prolonged for many years.

This was very nearly the condition found in the following case,¹ the opportunity of examining and describing which we owe to the courtesy of Dr. C. H. Thomas.

"The patient was a young man, *æt.* 22 years, who had been markedly cyanotic from infancy, and was poorly developed. He was unable to maintain a proper temperature. He suffered constantly from slight dyspnoea, with occasional exacerbations. At the autopsy there was marked congestion of the abdominal viscera, and the gall-bladder was packed with gallstones. Both lungs contained numerous yellow miliary tubercles.

"The heart was rounded. The cavities of the ventricles were not much enlarged, nor was there any hypertrophy of the walls of the left ventricle. The walls of the right ventricle were, however, decidedly thickened, though not equalling those of the left. The septum ventriculorum was disproportionately thick, and terminated about one-third of an inch below the level of the origin of the aorta in a smooth, rounded edge, over which the endocardium was thickened. The septum also seemed inside of its normal position, so as almost to bisect the aortic orifice. The aorta, which was slightly dilated but quite healthy, thus communicated freely with both ventricles. The origin of the pulmonary artery was very much obstructed, owing to coalescence and contraction of its valves. The ductus arteriosus was, unfortunately, not preserved,

¹ See Descriptive Catalogue of Path. Museum of Penna. Hosp., No. 1501, p. 84, by William Pepper, M. D., 1869.

but, owing to the large size of the pulmonary artery beyond the seat of obstruction, it had in all probability remained patulous. The foramen ovale was closed."

In this instance, the orifice of the pulmonary artery not being entirely closed, the opening in the inter-ventricular septum had been large enough to allow the right ventricle to relieve itself in that way, and consequently the foramen ovale had closed.

When writing of atelectasis pulmonum (p. 141) we called attention to the fact that in some cases, where the state of imperfect expansion persisted, the continued obstruction to the pulmonary circulation was followed by the same train of lesions, viz., patulous ductus arteriosus and foramen ovale, and hypertrophy with dilatation of the right side of the heart, as are consequent upon obstruction of the orifice of the pulmonary artery, and which, as in the latter case, might be attended with cyanosis.

The mere persistence of the foramen ovale scarcely be regarded as a cause of a permanent cyanosis. It is quite possible that during the early days of extra-uterine life, a certain amount of cyanosis might exist owing to the admixture of venous and arterial blood allowed by this opening, but after the forces of the circulation become equalized, it is quite certain that the valve of the foramen may remain unattached, or may even be somewhat insufficient to close the opening, and yet no cyanosis present.

As an illustrative case of one of the rarer forms of cyanosis, and one which bears in the most interesting manner upon the theory of its production, we abstract the following from a more full account published in the *Proceedings of the Pathological Society of Philadelphia*:¹

The child was a well-developed male, born at full term. No discoloration was noticed at birth, but on the twelfth day, as the grandmother was preparing to wash it, it had a convulsion, and from that time presented coldness of the extremities, gradually increasing lividity, feeble and rapid pulse, and moaning and sighing respiration.

During inspiration, the sternum and upper parts of the chest were elevated, but the lateral regions remained unexpanded, and there was marked recession of the base of the thorax. The percussion-resonance was diminished on both sides, but especially on the right. The vesicular murmur was puerile, except over the right side, where it was feeble. The cardiac sounds were decidedly louder at the right scapula than over the left. The cardiac impulse at the left nipple was very indistinct, and the sounds there feeble but natural. On pressing two fingers lightly to the left of the ensiform cartilage, close to the costal cartilages, a very distinct and quite vigorous impulse could be felt, one much more distinct than at the nipple. At this point, a distinct blowing sound attended the systole of the heart. The diagnosis made at the time was: atelectasis of both lungs, of the right greater than of the left; dilatation with hypertrophy of the right ventricle; obstruction of the pulmonary artery, and open foramen ovale. Death occurred on the forty-fourth day after birth.

At the autopsy, the body was very small and thin. The thorax was flattened laterally, and contracted at the base. There was marked collapse of the lower lobes of both lungs, and especially of the right. The heart was one-half too large, and full, rounded, and distended with soft black clots. The walls of the right ventricle were very thick, and its cavity quite small; it presented the appearance we usually associate with the left ventricle. The walls of the left ventricle were thinner than those of the right, and its cavity was much more capacious than that of the right.

¹Transposition of the Arteries. Dr. J. F. Meigs. Proc. of Path. Soc., vol. ii, p. 37; and Am. Jour. of Med. Sciences, vol. xi, 1860, 415.

The right auricle was dilated and considerably larger than the left. The foramen ovale presented an opening at its lower aspect of about 2 or 3 lines in diameter. The orifices of the venæ cavæ appeared smaller than usual.

The aorta and pulmonary artery were transposed. The aorta arose from the right ventricle in the usual position of the pulmonary artery; the pulmonary artery arose from the left ventricle, and passing under the arch of the aorta, gave to the latter, just beyond the left subclavian, the ductus arteriosus, which was quite pervious and of considerable size. The valves of the heart were healthy and not transposed. The pulmonary artery was of the natural size, and presented no obstruction at its point of origin. After giving off to the aorta the ductus arteriosus, it divided as usual into two pulmonary branches, which soon subdivided into others.

The aorta was of full size and presented nothing unusual. It gave off at its arch the innominate artery, and then the left primitive carotid and the left subclavian. Just beyond the latter it received, from the pulmonary artery, the ductus arteriosus.

The pericardium was normal in all respects.

Theories as to the Production of Cyanosis.—In the vast majority of cases the malformation which causes cyanosis is of such a character as to allow admixture of the venous and arterial blood, and, at the same time, to interfere more or less with the circulation of this mixed fluid. Ever since the time of Morgagni, authors upon this subject have been divided in opinion as to whether the coloration of the skin were due exclusively to one or the other of these causes: obstruction to the cardiac circulation and consequent venous congestion, or intermingling of the venous and arterial blood.

In regard to the first of these causes, although it has numbered among its advocates Morgagni, Louis, and Stillé, it cannot be considered competent to fully explain all the cases and peculiarities of cyanosis, although such central obstruction will unquestionably aid in its production.

Nor can the second theory be held exclusively sufficient, since not only are there cases met with where cyanosis is present and yet no admixture of venous and arterial blood is possible, but also, on the other hand, where a considerable degree of admixture exists without the production of cyanosis.

It seems necessary, therefore, as Smith has clearly pointed out, that any theory which pretends to embrace all the elements of this complex condition, should embody a reference to the fact that the essential defect in cyanosis is a want of arterialization of the blood.

SYMPTOMS.—Even in cases where cyanosis is due to congenital organic lesions, the peculiar symptoms are not always present until some time after birth.

Thus, in 138 of the cases of cyanosis collected by Smith, the time at which lividity was first noticed is stated as follows:

In 97 it was within the first week, and often within a few hours after birth.

In 3 at 2 weeks.

" 1 " 3 "

" 2 " 1 month.

" 7 from 1 to 2 months.

" 5 " 2 " 6 "

" 5 " 6 " 12 "

" 3 " 1 " 2 years.

In 6 from 2 to 5 years.

" 1 " 5 " 10 "

" 6 " 10 " 20 "

" 1 " 20 " 40 "

" 1 over 40 years.

—

41

Dr. Smith adds, "that in these 41 cases, in which blueness did not occur till after the age of one week, if the patient were less than two years old when it commenced, there was frequently no obvious exciting cause; but above this age, with three exceptions, such a cause is known to have been present. It is interesting to observe how trivial the exciting cause frequently is" (an acute attack of sickness, an attack of convulsions, difficult parturition, a fall, or even a severe blow), "and equally interesting to note how long patients have enjoyed good health, not having the least lividity, although the anatomical vice, to which the final development of cyanosis was due, had existed from birth."

The most characteristic symptom of cyanosis is the lividity of the surface, which varies in different cases from mere duskiness to a deep purplish tint. This color also varies in degree in different parts of the body, being most marked in the distant and especially the dependent portions, upon the mucous membranes, and wherever the capillary vessels are abundant, as on the face. Its degree varies, finally, in the same case with the condition of the circulation. In slight cases, when the patient is quiet and the circulation tranquil, the discoloration of the surface may be imperceptible, but upon any exertion, and especially in the more severe cases, the lividity becomes much intensified. In some instances, such as that narrated by us below, there may be actual ecchymoses of the surface, as in purpura.

The state of the general nutrition is much impaired, and the subjects of this disease are usually stunted and poorly developed. In many instances the generative system appears even more imperfectly developed than the rest of the economy. The temperature of the body is always reduced, and exposure to cold is very poorly borne.

In a good many cases the thorax presents the deformity so often found in association with rickets, known as the "pigeon-breast." It usually happens, also, that the ends of the fingers and toes become bulbous. Disturbances of the circulation and respiration are of frequent occurrence. Thus there is often some abnormal bruit heard in the cardiac region, due to the abnormal condition of the heart. The pulse may be regular and of fair volume, but more frequently is small, irregular, or intermittent, and palpitation is very readily induced by exertion. The disturbance of respiration usually corresponds in degree with the embarrassment of the circulation.

While the patient remains quiet his breathing may be easy and regular, but usually any sudden movement or exertion or emotion is sufficient to induce a paroxysm of dyspnœa, during which the lividity of surface becomes much deeper. In infants these paroxysms not rarely terminate in convulsions. Headache is frequently complained of, and is very apt to be caused by whatever disorders the circulation. During the paroxysms of palpitation, pain is often complained of in the region of the heart, but is rarely persistent.

Owing to the extreme venous stasis frequently present, there is a tendency to passive hemorrhages in cyanosis, which expresses itself by bleeding from the nose, mouth, stomach, or rectum, or under the skin. Edema of the lower extremities is often met with as a temporary condition after long

standing on the feet; it is also apt to appear and to invade the rest of the body towards the close of the case, when the circulation becomes more enfeebled.

MODES OF DEATH.—Many cyanotic patients die from the effect of some intercurrent acute disease, as hooping-cough or one of the exanthemata, all of which are very badly borne in this condition.

The theory that venous congestion is opposed to the development of tuberculosis, was applied by Rokitansky to this affection; but without any sufficient ground, since, as we have seen already, tuberculosis was found as the cause of death in no less than 26 of the cases collected by Dr. Smith.

In other cases death occurs suddenly, either during an attack of convulsions or a paroxysm of dyspnoea.

In severe cases of cyanosis life is rarely prolonged more than a few years; but in less marked cases the patients may even attain middle age. In 186 cases collected by Dr. Smith, the age at death was as follows:

In 17 under age of one week.	54	In 21 from 5 years to 10 years.
" 10 from 1 week to 1 month.		" 41 " 10 " " 20 "
" 12 " 1 month to 3 months.		" 20 " 20 " " 40 "
" 11 " 3 months to 6 "		" 4 over 40 years.
667 — " 17 " 6 " " 12 "		—
" 12 " 1 year to 2 years.		186
" 21 " 2 years to 5 "		

So that in 67, or more than one-third, death occurred before the close of the first year; in 121, or more than three-fifths, before the age of 10 years; only 24 survived the age of 20 years, and 4 the age of 40 years.

We subjoin the history of a case of cyanosis which we had under observation for several years, in which the symptoms of this peculiar condition were extremely well marked.

J. W., æt. 16 years and 7 months, has been cyanotic since infancy, but for the past few years, at least, has enjoyed fair health. At present there is marked lividity of the lips and of the nose, especially at the extremity, which seems somewhat increased in size. His hands habitually appear as though stained with solution of carmine, the skin being uniformly livid over the whole hands, but becoming lighter colored on the forearms. Pressure partly removes the lividity, which returns slowly after the withdrawal of the pressure. At times there have been little ecchymoses of the surface, followed by the effusion of serum under the epidermis, and the formation of superficial excoriations, which have left small cicatrices. Only a few of these have appeared on the hands.

These vascular disturbances are even more marked in the lower extremities. The feet are continually deeply livid; and over their surface and the ankles very numerous ecchymotic spots have appeared, which underwent the same changes as those on the hands, and have left shining cicatricial spots, of a deep blackish-red color, from the deposit of pigment, and surrounded by a dark brownish stain. There has also been a good deal of oedema of the feet lately. All of these conditions have been improved by the use of tight-laced stockings. The skin of both the hands and feet is rather soft and moist. The last phalanges, both of the fingers and toes, are markedly clavate and hypertrophied. Firm pressure upon them reduces their size; but, upon withdrawal of the pressure, the blood slowly returns, and they regain their former size. The temperature of the body is always low, and he suffers extremely from exposure to cold.

He suffers somewhat from dyspnœa, even upon slight exertion, but less so than formerly. He is also troubled with cough during the winter months. There is marked deformity of the thorax, the first and second pieces of the sternum uniting at an obtuse angle, and the cartilages of the third, fourth, fifth, and sixth ribs forming a marked prominence on either side of the sternum; the ensiform cartilage is considerably depressed.

The apex-beat of the heart is in the fifth costal interspace, and just inside of the vertical line of the nipple. The heart's action is regular, and at present there is no abnormal cardiac murmur, though two years ago there was a distinct soft systolic bruit. The pulse in the standing posture is 114, in the sitting 108. He has occasional attacks of epistaxis, and suffers quite frequently from attacks of gastric disturbance attended with severe headache.

The above note was taken in December, 1869. Since then this case has terminated fatally, and a post-mortem examination proved the existence of marked congenital malformation of the heart, of the character already fully described, originating evidently in obstruction of the pulmonary orifice.

TREATMENT.—In the form of cyanosis which we have described as depending on collapse of the lungs, the child should be placed in the position below recommended as rendering the heart's action most free; the temperature of the body should be carefully maintained, and a few doses of brandy in water or breast-milk should be given at intervals. There is evidently but little good that can be done by mere medication in cyanosis depending on malformations of the heart. When the heart's action is feeble and irregular, digitalis, iron, and quinia may be administered. During the paroxysms of palpitation and dyspnœa, the best remedies are diffusible stimulants, such as Hoffman's anodyne, spirit of chloroform, ammonia, and brandy; and derivatives, such as sinapisms to the chest, or hot mustard foot-baths. In cases where the digestion is markedly feeble, the use of vegetable tonics is indicated.

By far the most important part of the treatment, however, is a strict attention to the hygienic conditions of the patient. He should, so far as may be practicable, avoid all excitement and active exertion; his diet should be digestible and nutritious, his clothing should be warm, and, in addition, he should carefully avoid all exposure to severe cold.

In cases where the venous congestion of the legs is marked and attended with œdema or with enlargement of the veins, laced stockings should be worn.

It sometimes happens that cyanotic patients find that certain positions afford them peculiar ease and comfort. Believing that in cases where the lividity appeared soon after birth (cyanosis neonatorum) it was due to a patulous condition of the foramen ovale, the late Dr. Charles D. Meigs was led to recommend (*Diseases of Children*, 1850, p. 92) that such children should be placed upon a pillow, on the right side, the head and trunk being inclined upwards about 30° or 45°. The object of this position was "to bring the septum of the auricles into a horizontal position, so that the blood in the left auricle might press the valve of Botalli down upon the foramen ovale."

In a certain number of cases the adoption of this recommendation has

undoubtedly seemed to relieve the lividity, so that it is perhaps desirable that all cyanotic infants should be placed in this position; though from a glance at the anomalies in the formation of the heart which frequently attend cyanosis, it is evident that in most cases it could furnish no material relief. We are ourselves inclined to attribute the relief afforded by this position, not to any influence upon the foramen ovale, but to the fact that the heart's action is far most free and unincumbered when the child is placed upon the right side, with the trunk somewhat elevated.

ARTICLE II.

DISEASES OF THE HEART.

As we are prevented, by the limits of this work, from giving any detailed account of many of the affections which merely occur in childhood in common with adult life, we propose in this article to offer only a few practical remarks upon the differences presented by diseases of the heart occurring at these two periods of life.

Apart from those congenital malformations of the heart, already discussed in the preceding article, the diseases of this organ most frequently met with in childhood are pericarditis, and acute and chronic endocarditis, with valvular disease.

The most frequent causes of these affections are rheumatism, the peculiar alterations of the blood present in scarlatina, rubeola, and diphtheria, and extension of inflammation from the adjacent tissues, in cases of pleurisy or pneumonia. Of these well-recognized causes, rheumatism is by far the most frequent; for, although young children are comparatively rarely the subjects of this disease, it is followed by some cardiac complication in a larger proportion of cases in childhood than in after years. This fact will be more fully referred to in our remarks upon rheumatism, where we dwell upon the importance of recognizing this marked tendency, and of watching most critically for the appearance of any symptom indicating that the heart has become involved. This extreme watchfulness is the more necessary, because it frequently happens in young children, that for several days before the development of any local articular trouble, there may exist marked rheumatic fever, with serious inflammation of the membranes of the heart.

In a few instances an acute cardiac affection cannot be traced to any of the causes above mentioned, but appears to occur idiopathically, without exposure to any recognizable exciting cause.

So, too, in some cases of chronic valvular disease, and especially, it has seemed to us, of contraction and thickening of the mitral valve, the lesion cannot even be traced to any acute attack of endocarditis, but seems more akin to a fibroid degeneration, whose cause and early symptoms have been obscure and entirely overlooked.

Possibly, in some of these interesting cases, the real starting-point of the disease may have been an attack of endocarditis in fœtal life, which partially spoiled the valve, and set on foot degenerative changes, which slowly increased until they produced fatal symptoms.

ACUTE PERICARDITIS may occur at any period after birth. In very young infants it has been observed in conjunction with peritonitis, and was apparently due to erysipelas; while in other cases no cause could be assigned for its occurrence. The symptoms are, however, so vague and difficult to appreciate at this tender age, that the lesion is rarely recognized until after death. The infant is evidently in pain; the features are pinched and shrunken, the skin hot at first, and the pulse and respiration greatly accelerated. The physical signs can, however, rarely be satisfactorily determined, partly because death usually occurs before the lesions reach any considerable degree of development.

In older children the physical signs are often obscured by the coëxistence of some inflammatory condition of the lungs or pleura, and the existence of pericarditis can only be surmised by the presence of a degree of disturbance of the circulation and respiration out of all proportion to the amount of lung trouble.

When, however, pericarditis occurs without any such complication, it may be often recognized by the seat of pain; the existence of great dyspnoea, amounting at times to orthopnoea; the great frequency of the pulse, which is often small, and even irregular; the disturbance of circulation, as shown by lividity of the lips and face; and, finally, by auscultation and percussion, which reveal at first merely a friction-sound, and later, when effusion has occurred, distant and feeble heart-sounds, with an increased area of cardiac dulness. Other cases, however, occur which present but few of these objective signs, and it is only by the most careful physical examination that the disease can be detected.

When severe, pericarditis in children usually proves fatal. After death the same anatomical lesions are found as after pericarditis in adult life. The membrane is, in the first stage, reddened, injected, dryish, and slightly roughened; while later it is still injected and even ecchymosed, thickened, softened, and covered with patches or uniform layers of whitish or yellowish white lymph, the surfaces of which are usually flocculent or irregularly roughened. The pericardial sac contains a variable quantity of turbid, or, at times, bloody serum; or, in secondary cases, a sero-purulent fluid.

In cases where recovery takes place, the results of the previous inflammation are found, after death has occurred from some other cause, in the form of more or less extensive adhesion of the two layers of the pericardium, or merely of thickening and opacity of that membrane. It is seldom that fatal cases of pericarditis are uncomplicated. The most frequent complication is endocarditis, and occasionally broncho-pneumonia and pleurisy.

TREATMENT.—In idiopathic cases, if the disease be recognized in the early stage, we should advise local depletion over the præcordia by three or four leeches or cut-cups in a child of five years of age, followed by the application of warm mush-poultices, the depletion being repeated if indi-

cated; the internal use of large doses of acetate of potash and iodide of potassium, associated with doses of *veratrum viride* of appropriate strength, to quiet the excessive vascular excitement; and the careful administration of nutritious diet and small amounts of stimulus, if the powers of the circulation seem likely to yield to the influence of the disease.

In the very rare instances where the disease becomes chronic, and the effusion remains unabsorbed, the treatment should consist in the repeated application of small blisters over the *præcordia*, and the internal use of iodide of potassium, iodide of iron, with tonics and nutritious diet. In still more rare cases, the effect of chronic pericarditis is to induce extremely thick layers of lymph enveloping the heart. In one instance, which we lately saw in consultation with Dr. C. H. Thomas,¹ where pericarditis of several years' duration existed, the sac was obliterated by a layer of succulent, almost gelatinous lymph, fully one-half inch in thickness. In such a case, it is difficult to distinguish the lesion from hypertrophy of the heart. It might be done, however, by the absence of valvular murmur, and the feebleness of the apex-beat and of the heart-sounds. If, moreover, we have observed or can get an accurate account of the original attack, its character and the future course of the case would be of great value.

ENDOCARDITIS.—In many cases, acute endocarditis in children occurs in conjunction with pericarditis, although it also occurs frequently as an independent affection. It is due to the same series of causes, also, as have been already enumerated when speaking of this latter disease; of these undoubtedly rheumatism, scarlatina, and rubeola are far the most frequent. And as it is of far more frequent occurrence than pericarditis, and productive of even more serious results, it is necessary that we should, if possible, be more upon the alert to detect the very earliest symptoms of its presence. Since recent observation has established the occurrence of acute affections of the heart in the course of some other specific diseases, as stated in our remarks on pericarditis, it is an important rule to use the same care in repeatedly examining the heart in these cases also.

In severe cases, whether occurring idiopathically, or as a complication or sequel of some other disease, there is violent disturbance of the circulation, with great dyspnoea, and short, dry cough, without any of the physical signs of pulmonary disease. The child is extremely restless, and, upon auscultation, an abnormal bruit is heard attending the heart's action. The valvular murmur, it must be remembered, is not harsh and strong as in some cases of chronic valvular disease, but may be so gentle and soft as to be heard with difficulty.

In most cases,² the mitral valve is chiefly affected in acute endocarditis, and the murmur detected on auscultation is heard over the body of the heart and to the left of this organ, and often has its seat of greatest intensity near the apex. We have most frequently observed the murmur to be systolic in time, attending and more or less obscuring the first sound of

¹ See report in Transactions of Philadelphia Pathological Society, vol. vi, 1876-77.

² See table of Dr. Sansom, Medical Times and Gazette, 1879, vol. ii, p. 361.

the heart, in such acute cases. Of course, this indicates the existence of some imperfection in the closure of the mitral valves, allowing more or less regurgitation of blood into the auricle with each contraction of the ventricle. Occasionally a double murmur, attending both the systole and diastole, and indicating roughness as well as insufficiency of the mitral valve is heard. In more rare instances, we have found the aortic valve to be the seat of acute endocarditis, as shown by the presence of a single or double blowing murmur over the base of the heart, and transmitted most strongly upwards over the upper part of the sternum to the second right costal cartilage.

But more frequently the acute symptoms are not so marked or characteristic as this, and, when ensuing, for instance, in the course of acute rheumatism, may consist merely in a little increase of the heat of the skin, frequency of the pulse and restlessness, with or without vague complaints of pain about the præcordia.

Absolutely the only way of recognizing such cases is by auscultation, and consequently we would urge the immense importance of carefully auscultating the heart *daily*, not only in every case of acute rheumatism in a child, but also in every case where anomalous febrile symptoms, with acceleration of pulse, are present, and particularly if there be general soreness, or even resistance to motion.

In very severe attacks of acute endocarditis, death may occur early; but more commonly the disease is less severe, and the urgent symptoms subside, leaving, however, in but too many cases, organic valvular disease.

When death occurs during the acute stage, the endocardium is found injected, reddened, softened, and readily detached from the muscular wall. The lesions are most marked on the left side of the heart, and especially on the endocardium covering the mitral valve, where, in addition to the above mentioned appearances, there are usually patches or rows of minute granular vegetations, which form a fine beaded line along the free border of the valves; or, in other cases, delicate fringe-like processes which hang from the leaflets. We have alluded at some length in our article on chorea, to the theory which has been framed to explain the frequent occurrence of this latter disease in connection with rheumatism, by the separation of minute fragments of such vegetations, and their impaction in some of the vessels of the brain.

The treatment of acute endocarditis should be the same as that recommended for acute pericarditis.

CHRONIC VALVULAR DISEASES.

There are certain general remarks which we desire to make in connection with these affections, which are applicable to them, without reference to the particular valve diseased; in addition to which, we will call attention to the diagnostic signs and special features of the diseases of each set of valves.

CAUSES; FREQUENCY.—In very many cases, heart disease in young children is recognized for the first time when such marked lesions exist

as to convince us that the disease has already been of some considerable duration. Undoubtedly this is partly because the acute symptoms of the early stage have been entirely overlooked. This is particularly the case when the case is rheumatic in its origin. We are convinced that acute rheumatism is often overlooked in young children, and also that endocarditis occurring in the course of such attacks not rarely escapes detection. It is, therefore, very difficult to say in what proportion of cases in young children, valvular diseases have been of acute origin. In our own experience they have, with the exception of contraction of the mitral valve, almost aniversally followed an attack of endocarditis. In the case of mitral contraction, however, it is quite often impossible to trace the disease to any acute attack. It would appear, therefore, either that, contrary to the usual rule in early life, this lesion is often the result of a slow degenerative, fibroid change, or else that, in some cases, it may arise in fœtal life. This latter view does not seem at all impossible when we remember how slow is the development of this lesion, and for how long a time it may remain latent. Thus, Dr. H. Barth (*New York Medical Record*, 1879, p. 292), reports a very interesting case of fœtal endocarditis detected before birth, and in which the autopsy verified the diagnosis.

As to the relative frequency with which the different sets of valves are affected, lesions of the mitral valve undoubtedly preponderate largely. We have, it is true, met with extreme aortic disease in quite young children, marked by all the physical signs that are familiar as occurring in the adult; but such cases have been rare compared to those in which the mitral valve was the seat of the disease.

ANATOMICAL APPEARANCES.—The lesions which are found in chronic valvular diseases do not differ from those which are found in the adult, nor do they present characteristic differences dependent upon the mode of their origin. It is, however, probably true that in those cases which have followed acute endocarditis, it is more usual to find numerous and large vegetations upon the valves, than where the lesion has been chronic and of gradual development from the start. The lesions which are found usually, are vegetations or calcareous incrustations on the valves, or there may be thickening, contraction, and coalescence of the valves and their chordæ tendineæ, either of which conditions may be attended with contraction of the orifices of the heart, and obstruction to the passage of blood. On the other hand, the contraction of the valves may be in such a direction as to render them insufficient to close the orifice, and thus allow regurgitation. The effect of these lesions upon the walls and cavities of the heart will vary with their degree and suddenness of development. Usually they are followed by dilatation of the cavities involved, and by thickening or hypertrophy of their walls, which has usually seemed to us more constant and to bear a larger proportion to the dilatation than in adults.

SYMPTOMS.—The general symptoms during the early stages of chronic valvular disease, are often extremely slight, consisting merely of some interference with the general development of the body; a little palpi-

tion of the heart, and dyspnœa on exertion; occasional præcordial distress, and perhaps slight prominence of the cardiac region.

The vague character of these symptoms accounts for the fact that, after the subsidence of the acute symptoms of endocarditis, when the disease has begun in that way, such cases are very often neglected, and receive no proper care until the occurrence of dyspnœa, cough, or dropsy, gives warning only in time to recognize that incurable or even fatal lesions have been developed.

We make these remarks especially to call attention to the insidious mode of approach of many cases of the chronic valvular disease of the heart in children; and to impress upon our readers the important practical rule that, whenever, in the investigation of a child suffering with obscure ill health, we learn of the previous occurrence of acute rheumatism, or any of the general infectious diseases, or find mentioned among the symptoms any irregularities of the circulation or action of the heart, careful physical exploration of the heart should immediately be practiced.

The special symptoms which attend the diseases of the different valves, may be briefly described as follows:

Diseases of the Aortic Valves.—These affections are, as already said, comparatively rare in children. The blowing murmur which attends them is usually strong and distinct. If the lesion causes obstruction of the aortic orifice, the murmur will attend the first sound; if there be regurgitation through the valve, it will attend or take the place of the second sound. In many cases the lesion causes both obstruction and insufficiency, and there is therefore a double murmur. In either case the murmur will be heard extending from the base of the heart upward and across the sternum to the second right costal cartilage, as well as downward along that bone to the xiphoid cartilage. It is also transmitted into the arteries. The murmur is often so loud that, especially in cases of regurgitation, it may be heard down over the body of the heart to the apex; and also to a varying distance on either side of the sternum over the upper part of the chest. Occasionally also a thrill may be felt over the upper piece of the sternum, in the second intercostal space at either the right or left edge of the sternum, or at the supra-sternal notch.

The action of the heart is regular, and may not be accelerated, though exertion readily excites palpitation. The apex-beat is quick and strong, and is found after a time below and to the left of its normal position. The area of cardiac percussion dulness also becomes moderately increased in consequence of gradual hypertrophy of the walls of the left ventricle.

The pulse is small, quick, and in cases of regurgitation, jerking and unsustained, or receding.

In severe cases, there are marked evidences of interference with the arterial circulation. The surface is pale, and shows the insufficient amount of blood which passes through the arterial capillaries.

The respiration is usually but little disturbed, excepting in consequence of unusual exertion, so long as the lesion is limited to the aortic valve, and the walls of the left ventricle undergo sufficient compensatory hypertrophy to overcome the obstruction to the circulation.

The *prognosis* in aortic disease of moderate severity has not seemed to us unfavorable so far as regards prolongation of life. Thus, for example, we treated a girl of $9\frac{1}{2}$ years, who had a violent attack of acute articular rheumatism with endocarditis. This was followed by a double aortic murmur, which persists to the present time, although she has grown up, married, and has one child. Her health is delicate, and she has very moderate dyspnœa on exertion. We have frequently observed this same tolerance of serious aortic lesions for a number of years. We have never met with a case in which sudden death occurred in the course of aortic regurgitation, as so frequently happens in adults.

Mitral Obstruction.—This interesting form of cardiac lesion would merit a more full description here than any other valvular disease, because its symptoms are somewhat peculiar, and more especially because it is of such comparative frequency in childhood.

Its origin, as we have already remarked, is usually insidious, and it is frequently impossible to gain any history of acute disease in cases where marked mitral obstruction is detected.

The general symptoms which first attract attention to the heart are rarely noticed before the age of 7 or 10 years; and we may then learn that during previous years the child has seemed as active and playful as usual, or that he has always shown an indisposition to active play or exertion, and has become tired readily. Attention is attracted to the heart by the increased tendency to dyspnœa and palpitation on exertion, and by the readiness with which cough of a bronchial character is contracted on very slight exposure. Occasionally during these attacks of bronchitis with pulmonary congestion, hæmoptysis may have occurred. Examination may now show the existence of prominence of the præcordia; and the area of cardiac dulness is usually increased, though not to a marked extent. Frequently a thrill can be felt over the præcordia, and careful examination will show it to occur just before the apex-beat. We have known this thrill to begin distinctly about the base of the heart, and to extend quickly down towards the apex, terminating as the apex-beat was noticed. On auscultation, a murmur, usually of a low, hoarse, or churning character, is heard, which presents these additional peculiarities; it is generally distinctly *presystolic* or *auriculo-systolic* in time, occurring, that is, in the long period of silence preceding the first sound; its relation to the phenomena of the cardiac action can usually be determined without difficulty by observing that it follows the second sound, and that it stops just before, or else runs into, the time of the first sound and of the pulse of the carotid artery. This murmur, also, although usually quite strong, is, as a rule, remarkably localized in comparison to other valvular murmurs; its seat of greatest intensity is at or near the apex, and it loses force rapidly on leaving this point in any direction. Attention to the peculiar physical signs above given, as well as to the general symptoms, will generally render the diagnosis clear.

The prognosis, as regards prolongation of life and maintenance of comfort, is comparatively favorable; as regards improvement in the organic condition of the heart, it is of course entirely the reverse. We have under

our care at present a number of children, of ages varying from 5 to 16 years, who present the typical symptoms of mitral obstruction, but of whom a fair proportion, by care in the manner of living, enjoy entire comfort. Usually, however, the frequent recurrence of pulmonary congestion injures more and more seriously the equilibrium of the heart's circulation and the efficiency of the right ventricle, and eventually grave symptoms of failure of cardiac power, with general venous stasis, appear, and increase until a fatal result occurs.

Mitral Regurgitation.—This, which is the most frequent form of cardiac disease in young children, depends upon inflammatory alterations in the mitral valve, usually resulting from acute endocarditis, and which render it insufficient to close that orifice during the systole of the left ventricle. In this condition, as in the last, the pulmonary circulation is apt to be disturbed from time to time, and therefore the early general symptoms which attract attention to the thoracic organs are usually shortness of breath on exertion, liability to cough, and palpitation of the heart. Of course, where we are in attendance upon a case of rheumatism, for instance, when the acute cardiac inflammation occurs, the fact will be recognized by the symptoms detailed under the head of acute endocarditis. But unfortunately it often happens that this acute stage is quite overlooked, and we would therefore again urge the importance of a careful physical examination of the heart in every case where a child is brought to us complaining of vague symptoms of embarrassed breathing, though no suspicion has ever been raised as to the existence of heart disease. Sometimes, indeed, much more marked general symptoms will have appeared, as, for example, severe dyspnoea on exertion, pulmonary congestion with cough and moist or dry râles over the posterior parts of the lungs, palpitation of the heart, lividity of the lips and fingers, and even œdema of the feet.

On physical exploration we often find prominence of the præcordia, with signs of more considerable hypertrophy and dilatation than in cases of mitral obstruction. The impulse is extended and too forcible, or may even be heaving; it is rarely attended with any thrill. On auscultation a blowing murmur, which varies very greatly in different cases in its force and character, will be heard accompanying or replacing the first sound of the heart. This murmur is heard at the base, and is transmitted most strongly towards the apex, where it often has its point of greatest intensity. It is also strongly transmitted to the left of the apex, being well heard in the infra-axillary space on the level of the apex-beat, and frequently, also, on the dorsum of the left chest, at the angle of the scapula. The only other form of valvular disease with which it is possible to confound this is mitral obstruction; but attention to the evident points of difference noted above will render the diagnosis easy in most cases.

The prognosis varies extremely in different cases, depending upon the extent and rapidity of development of the lesion; the completeness with which the disturbance of the circulation is compensated by the hypertrophy and increased power of the walls of the left ventricle; and the

vigor of the system and the preservation of the tone and nutrition of the muscular fibre of the heart. This form of heart disease illustrates more clearly than any other, the more favorable prognosis which may be made in many cases of organic valvular disease in children, as compared with the same condition in adults. This depends partly upon the fact that when the lesion is not extensive, and when the patient is placed under favorable circumstances, the heart accommodates itself in its growth to the defective state of the valves, and overcomes the impediment to the circulation by acquiring increased propulsive force.

Not only, however, are the valvular lesions in childhood thus partly compensated by hypertrophy of the walls of the heart, but there is also an undoubted tendency, in some favorable cases, for the valvular lesions, both mitral and aortic, themselves to diminish. Thus among the following cases, which we have selected from a large number of records collected in our practice, there will be found several where positive abnormal bruits, due to organic valvular disease, have gradually disappeared in the course of years.

Acute articular rheumatism; endocarditis; recovery; murmur persistent but diminishing.—H. S., a boy, æt. 12 years, had a severe attack of acute articular rheumatism in April, 1869, with swelling, redness, and pain of joints; a systolic murmur appeared at the apex without any pericarditis. He recovered, under the use of alkalis and opium. In November, 1869, seven months after the attack, he seemed perfectly well; had no dyspnoea except on violent exertion. The murmur at the apex was still audible, but less marked than three months ago, when he was last examined.

Acute endocarditis (rheumatic?); marked improvement in general symptoms, but persistent murmur.—B. H., a girl, at age of 4 years suffered from an ordinary catarrh, when we detected a loud, high-pitched murmur at the apex, and, on inquiry, learned that, when 2½ years old, she had a violent inflammation of the chest, supposed to be catarrhal fever. At present, at the age of 12 years, she is in excellent health, without any of the rational signs of cardiac trouble, but she still has a well-marked, rather prolonged, high-pitched, systolic murmur at the apex.

Repeated attacks of rheumatism with severe mitral disease; improvement in general symptoms and force of the murmur.—L. S., a girl, was subject to attacks of rheumatism from very early age, and has presented symptoms of cardiac disease from infancy. At age of 13, there was a strong systolic murmur heard over base and toward apex. She suffered much from violent palpitation, pain in præcordia, headache, and habitual dyspnoea, much increased on exertion. At age of 18, there is still a systolic mitral murmur, but of much less intensity than formerly. Her general health is excellent, and she has but little dyspnoea or palpitation at any time. The heart's action is still readily excited; the impulse strong, but without thrill; there is marked increase in the area of cardiac dulness, but no positive prominence of the præcordia.

Acute rheumatic endocarditis, chronic mitral disease; recovery in five years.—F. R., a girl, at the age of 6 years, was attacked with slight rheumatic fever, without any articular symptoms. In a few days a distinct, but not loud, rather low-pitched systolic murmur was heard at the apex. The treatment consisted of rest in bed, quinia, and Dover's powders. After ten days all the acute symptoms disappeared, but the murmur continued. She regained her health, but for two years the murmur could be detected, but then gradually diminished; and now, five years after the first attack, no murmur can be detected, the first sound at the apex being merely a little prolonged. Her general health is excellent.

Acute rheumatic endocarditis; valvular disease, gradually recovering in course of two

years.—M. B., a girl, at the age of 7 years had fever of a type that made us suspect pneumonia or pleurisy, but without cough, pain in the chest, or any of the physical signs of pulmonary disease. On the third day, there was complaint of pain in one groin, but with no other articular symptoms; rheumatism being suspected, a careful examination detected a roughish systolic murmur at the apex. She was leeches at the præcordia, confined strictly to bed, and had Dover's powders given her. The fever subsided, but the murmur continued for two years, gradually growing faint, and finally disappearing.

It is, however, only when the general nutrition of the patient is good, so that the tonicity of the heart's tissue is preserved; and when all exposure and exertion, which could overtax the energies of the crippled organ, are carefully avoided, that such compensation and gradual recovery are possible.

For in cases where the vigor of the heart's action fails, and degenerative changes occur in its muscular tissue, the tonicity of the walls soon diminishes, and allows the development of passive dilatation of the cavities. In this condition it is not long before the most grave symptoms of embarrassed circulation appear, and the case passes more or less rapidly through the stages common to fatal organic disease of the heart.

The following case may be quoted as a full illustration of the latter remarks, in regard to the effect of exposure and exertion in inducing a fatal result in cases which otherwise might have gradually improved.

Repeated attacks of acute rheumatism in early childhood; valvular disease and hypertrophy; gradual improvement; exposure to hardships of army life; rapid aggravation of symptoms, and death.—W. D., male, as a young child suffered from repeated attacks of acute articular rheumatism with cardiac complication. At the age of 9 years, Dr. William Gerhard pronounced him to be suffering from valvular disease and hypertrophy of the heart.

His condition was gradually improving, and he had so few symptoms of cardiac disease that, at the age of 18 years, he was able to enter the infantry service. At the end of one year, however, he was discharged for disability, and when seen by us in July, 1864, presented the following symptoms: bulging of præcordia; marked extension of the cardiac impulse, which was heaving and powerful; marked increase in the area of cardiac dulness from the presence of pericardial effusion; and strong systolic mitral murmur. He had lost flesh; the surface was sallow and lips livid; there was frequent cough with occasional hæmoptysis and epistaxis; the liver was enlarged, and there was frequently œdema of the feet.

Towards the close of the year, the heart's action grew more labored and feeble, the pulse thready and frequent, the entire body became anasarcaous, and considerable ascites appeared. He suffered from constant orthopnoea and frequent cough, with bloody expectoration. The skin of the legs subsequently became gangrenous in parts, and he died December 28th.

At the autopsy, the heart was found enormously enlarged, extending over to the right of the sternum. The pericardium was firmly adherent throughout its extent, and in places was $\frac{1}{2}$ inch thick; there were several cartilaginous plates in the substance of the investing pericardium.

The heart measured $9\frac{1}{2}$ inches from apex to base, and 6 inches across at the base of the ventricles; the walls of the left ventricle were $1\frac{1}{2}$ inches thick; the auricles were enormously dilated with very thin walls. The aortic and pulmonary valves were healthy and apparently sufficient; the tricuspid valves were also healthy, but probably insufficient. The mitral valves had entirely disappeared, from shrivelling

and contraction, and there merely remained a very thick fibrous ring, studded with calcareous masses, bounding the auriculo-ventricular opening.

The muscular tissue of the heart presented an incipient state of fatty degeneration.

The liver was enormously enlarged, reaching nearly to the umbilicus, and presented intense nutmeg congestion.

The kidneys were large and congested; and the spleen was three times its normal size.

There are, moreover, other dangers attendant on organic disease of the heart in addition to those above referred to as resulting from progressive failure of cardiac power. Embolism, especially of the spleen and kidneys, is quite frequent; and very important cases have been recorded by Gee and Cheadle (*Medical Times and Gazette*, November 17th, 1877), in which in consequence of the local irritation caused by the embolism and the resulting infarction, and of the septicæmia from absorption of the disintegrating tissues at the affected point, a prolonged and decided hectic fever (constituting, in fact, chronic pyæmia) was maintained.

TREATMENT.—Having spoken somewhat in detail of the symptoms and prognosis of the different forms of valvular disease in children, it remains to make some general remarks upon their treatment. In the management of such cases, as in adult life, the most important point to be attended to is the careful regulation of the mode of life. The child should be warmly clothed, and carefully protected from any exposure which might induce rheumatism or congestive attacks; all violent exertion of body or mind should also be avoided, and, so far as possible, all sudden emotions, as fright or anger. On the other hand, care should be taken that proper gymnastic and outdoor exercise should be regularly taken in such ways as to invigorate the frame and strengthen the muscular system, without producing too much exhaustion. The diet should be nutritious and digestible, and if the appetite should fail, and the child appear weakly and pale, vegetable tonics, with iron, should be administered.

The appearance of symptoms of pulmonary congestion or of catarrh, should attract immediate attention, and lead us to employ counter-irritation and suitable expectorants to relieve the lungs.

In cases where the heart's action is excited, and too frequent and powerful, while evidences of excessive hypertrophy begin to show themselves, we should employ cautiously *veratrum viride* or *aconite* to control it. When, on the other hand, any of the cavities of the heart are subjected to overstrain from valvular obstruction or insufficiency, and the heart is acting irregularly and inefficiently, the greatest benefit will be obtained from the use of *digitalis*. Indeed, in many instances we have observed, under the prolonged use of this drug, very great permanent improvement, gradually showing itself both in the action of the heart and in the general symptoms.

Severe paroxysms of palpitation, should they occur, require the use of antispasmodics, diffusible stimuli, and revulsives, just as are indicated under the same circumstances in the adult. Should the attack not subside promptly, recourse should be had to *digitalis*, which may be freely administered, and will be found to afford marked relief.

In cases of rheumatic origin especially, we have thought that good results, in regard to the progress of the organic changes in the heart by the prolonged use of iodide and bromide of potassium, given with due regard to the danger of developing an anæmic state of the blood by the uninterrupted administration of these drugs for a long time.

On the whole, as we have already said, there is reason to be somewhat hopeful in the treatment of chronic valvular disease of moderate severity in young children, bearing in mind the wonderful power which the growing heart possesses of compensating such lesions, so long as by careful attention to hygiene and medical treatment we are able to preserve the tone and nutrition of its muscular tissue.

CLASS III.

DISEASES OF THE DIGESTIVE ORGANS.

CHAPTER I.

DISEASES OF THE MOUTH AND THROAT.

WE shall consider the diseases of the mouth in the following order:

1. Simple or erythematous stomatitis.
2. Follicular stomatitis, or aphthæ.
3. Ulcerative, or ulcero-membranous stomatitis.
4. Gangrene of the mouth.
5. Thrush, or stomatitis with curd-like exudation.
6. Affections of the tonsils.
7. Simple, or erythematous pharyngitis.
8. Retropharyngeal abscess.

In the early editions of this work, we described pseudo-membranous pharyngitis in this place, but further observation and research have clearly established the fact that this is but a local manifestation of a constitutional affection, diphtheria; and we have accordingly given a full account of this whole subject, under this latter name, in the section on constitutional diseases.

Before entering on the consideration of these separate affections, most of which are of frequent occurrence during early infancy, it has seemed best to us to devote a special chapter to the diet of children during the nursing age, instead of the desultory statements that have appeared in previous editions of this work.

ARTICLE I.

FOOD.

EXPERIENCE has shown us that not only the present health of the child, but also its power to resist what may be called the unpreventable diseases of early age, and often its chances of success in the struggle of life, depend largely on the success or failure of the diet provided for it.

We have deemed it best to place this chapter at the head of the section

devoted to diseases of the digestive organs, for the reason that food has much to do in the causation of several of these affections, and because we believe that without a proper knowledge of the diet suitable for infancy, the physician might as well abandon the field, since it is certain that no medical potions can stay or hinder the evil born of, or maintained by, an improper food.

We shall restrict our remarks to the food which is proper during the first two or two and a half years of life, which include the nursing age and the first dentition.

At the very outset of this subject we renew our oft-repeated opinion, that the only food which can satisfy perfectly the demands of the child upon its mother, relations, or the public, is woman's milk, either that of the mother or of a wet nurse. Could this be provided for all children, there would be no need for this chapter. We think a child has a right to this food, if it can be obtained for it. We have met with so many women, and men, too, objecting to wet nurses, that we wish to state the matter strongly. For ourselves, to deny that woman's milk is better for infants than the milk of any animal, or than any other product of the animal or vegetable kingdom, is like denying that two and two make four, or like asserting that the intelligence of man is above the intelligence that created man.

But circumstances constantly occur under which the child must be fed on artificial food, wholly or in part. The parents cannot afford, or they cannot find, a wet-nurse, and children at the public charge cannot always be supplied with nurses; or the child must be fed in part to save the mother; or, lastly, it is weaned early and must have artificial food. So that, however much we may regret the necessity, it is a fact that we are forced to supply artificial food to large numbers of young children.

Experience has demonstrated that the best substitute for woman's milk is the milk of some one of the mammal class of animals. The attempt to hand-feed children on any of the farinaceous substances alone has proved so disastrous that it is astonishing to find any physician of the present day sanctioning it. And yet we have known it to be done quite frequently, and never otherwise than with failure. Either the child has died, or has come to be so feeble or ill that the physician who directed it, or some one called in his place, or the parents, have changed the diet.

The milk generally employed is that of the cow, goat, or ass. It is usually conceded that the milk of the ass most nearly resembles that of woman,—and this milk is a good deal used in Europe,—with us it is so rare that we have never known it to be used.

Goat's milk, also, is employed in Europe, particularly among the farming classes of parts of France, and in Switzerland. It is used to some extent in this country, especially amongst the poorer inhabitants in the suburbs of our cities. We have known it to be employed, and have, ourselves, ordered it in several instances. It has answered in a few cases very well, but its peculiar and disagreeable odor, the difficulty in obtaining it pure and fresh from the poor, who alone keep goats, are great objections to it. In a case where it seems to be necessary we advise the

purchase, if possible, by the family needing it, of a goat for the special use of the child. It is much the safest plan on the whole.

Time and experience have taught us, and most of the profession, that in this country, and especially in our large cities, we must depend on the milk of the cow, and our remarks on artificial food will, therefore, be limited to this form of milk.

In choosing cow's milk the first thing to be thought of is its purity and freshness. In small towns and in the country there ought to be no trouble in obtaining it fresh, but in large cities this is often very difficult. Still, with money at command, and with due care and diligence, it can generally be procured. Our own plan has always been to find a milkman who brings milk from his own farm, or who at least employs the man who delivers it. We never have, and we never shall, so long as we can help ourselves, take milk from the middlemen who buy it of anybody and everybody. Moreover, the person who has charge of the child should always, if possible, know the milkman personally, and know exactly where he comes from, and what manner of man he may be. An honest farmer or dairyman who pastures and feeds his own cows on a healthy farm is the man to be employed. If the character of the milkman is not a sufficient guarantee, or if from any accident the milk must be changed, or if any doubt arise as to its quality, there are some simple methods of examination, which can be made use of by any one of ordinary intelligence, which will reveal most of the gross deceptions apt to be practiced by milk venders. Of these methods we shall treat a little further on.

After good milk has been obtained, it is of the utmost importance that it should be preserved pure at home. The vessels in which it is to be placed must be kept scrupulously clean, and they must not be exposed to foul or stale air, or odors of any kind. They should be kept in a cool, sweet cellar, away from meat or vegetable supplies for the family, or in a special ice-chest intended for the milk alone. It is an established fact, as may be seen in the article on the causes of entero-colitis, that milk has a special tendency to the absorption of the microscopic organisms which go to make up many of the so-called filth-matters, and that when thus contaminated it may cause, by its ingestion, the filth diseases. We repeat, therefore, that the mother at the house cannot be too cautious in having the milk brought to her in clean vessels, and then in causing it to be preserved in the mode and with the care above specified.

A good specimen of cow's milk is slightly acid or neutral; it must have a certain average proportion of cream, and must be of a certain average density.

The following is the average composition of good cow's milk, as given by Dr. Stephen P. Sharples, S.B., chemist, inspector of milk for the city of Boston, in an essay on the adulteration of food. (See Buck's *Hygiene*, vol. ii, p. 366.)

Average Composition of Pure Milk.

Specific gravity,	1.030 +
Cream, per cent., by measure,	8 per cent. +
	Per cent. by Weight.
Sugar,	4.40
Caseine,	4.30
Ash,60
	<hr/>
Solids, not fat,	9.30
Fat,	3.20
	<hr/>
Total solids,	12.50
Water,	87.50
	<hr/>
	100.00

Dr. Sharples gives this analysis as a standard below which pure milk should not fall. "Milk," he says, "can easily be kept up to this standard by proper food and care of the cow. Any falling below it is suspicious." He also states that the Society of Analysts of England has adopted the following slightly lower standard:

Solids, not fat,	9.00
Fat,	2.50
	<hr/>
Total solids,	11.50

This he thinks too low, remarking that it does not give the public a fair chance. The plan he found best in practice (in Boston) was to call all milk falling below the first standard adulterated, but not to prosecute the milkman unless it fell below the Society's standard. The New York Board of Health, it is stated, relies almost entirely on the lactometer, but in Massachusetts, Rhode Island, and perhaps other States, an analysis is required.

It is singular how authorities differ in their estimates of the proportion of cream in milk. Thus Dr. Parkes states that it ought to be from 6 to 11 per cent. Dr. Edward Smith, of London (*Foods*, Am. ed., New York, 1873, p. 313), gives it at 10 to 12 per cent., and states that at the Liverpool Workhouse, they adopt a standard of 10 per cent., and pay a half-penny per gallon for each degree in excess, and deduct a like amount for each degree in defect. The *Maison Rustique*, a French work of high authority on agriculture, states (tome iii, p. 61) that the milk of cows of good race, well kept, furnishes 15 per cent. of cream. The milk supplied by the milkman employed by one of us (his herds contain only good ordinary cows) showed a number of times 15 per cent. of cream, once 19 per cent., and again 10 per cent. At one time the cream fell to 8 per cent. We made a complaint, and were told that the sudden return of a number of his customers at the end of the summer season found him with an insufficient herd, and that he was obliged to purchase some milk. The low proportion has not occurred since. Of three specimens bought at hazard of different dealers, the proportion of cream was 7, 6, and 14 per cent. respectively. We have concluded that good milk in Philadelphia ought to

furnish 10 per cent. of cream, and upon this standard have based our modes of preparing and using it for children.

We will state, further, that the milk of the Alderney cow yields from 30 to 40 per cent. of cream. This milk is sometimes used by the wealthier classes of citizens, who possess country-seats and keep their own dairies. We are opposed to its use as a rule, believing that it contains too much fat, too little caseine, and that it is too unlike human milk.

It is highly important, as we said above, to have some ready and simple means of estimating the quality of milk, and, if the vendors knew that their customers had such means, and used them, they would be more cautious about adulteration, and the dishonest would soon be weeded out. Such an examination, not perfectly accurate, but of great value for household purposes, can be made by the use of litmus and turmeric paper, to determine the acidity or alkalinity of the fluid, by an instrument for measuring the proportion of cream, and by taking the specific gravity. Litmus-paper turns red when touched by an acid; a very weak acid will do this. If a good specimen of this paper (which can be procured of the apothecary) turns faintly red when dipped into milk, the milk is properly acid; if turned bright red, the milk is too acid. When no change is produced in the paper, the milk is either neutral (which is sometimes the case with healthy milk), or it is alkaline. To determine whether it be alkaline or not, turmeric paper, which is turned brown by alkaline solutions, must be used. If the specimen is found to be alkaline in a marked degree, either the cow is, in all probability, diseased, or some alkali has been added to the milk.

It is a curious fact that Dr. Parkes (*Manual of Practical Hygiene*, 2d ed., London, 1866) is almost alone in stating that healthy cow's milk is either faintly acid or alkaline. Most authorities assert that it is alkaline. In order to determine this matter for ourselves, we tested the milk of thirty-one fine cows, fed on the finest pasturage in the neighborhood of this city. This was done by taking the milk, just as it was drawn from each animal, at the milking-house, and testing at once with litmus-paper. In all, the paper was turned red more or less distinctly. Dr. John Ashhurst, Jr., of this city, tested for us, with both litmus and turmeric paper, the milk of nine fine Durham, and of four Alderney cows, belonging to his father, all on rich pasturage. He found that in one Durham and one Alderney, the milk appeared to be almost neutral, but in all the rest more or less acid. In testing the milk of another Durham, litmus-paper was reddened, whilst the turmeric was also slightly changed. He supposed the latter condition to be due to a greasy condition of the milk, owing to the fact that the cow was in the latter period of a long lactation.

The instrument for estimating the cream is called a *creamometer*. A glass vessel, such as can be bought of the apothecary, or at a shop for the sale of chemical apparatus, divided into hundredths, is all that is necessary. We use a vessel tall and narrow, having a foot like a wineglass, and a ground-glass stopper. This is divided into a hundred cubic centimeters. The vessel is filled with fresh milk to the upper mark, and allowed to stand for twenty-four hours in a cool place, away from any

currents of air. At the end of that time the cream will have risen to the top, and the proportion is read off in hundredths or percentage, on the scale.

The specific gravity of milk can be taken with the specific gravity bottle, or, what is better for household purposes, and quite sufficiently accurate, with what is called a *lactometer*. This is nothing more than a common hydrometer with special marks on the stem. These instruments, also, can be bought in the shops, or an ordinary hydrometer may be used. Dr. S. P. Sharples (*loc. cit.*) says: "The one which has been found most convenient is a simple spindle about fifteen centimeters long. The stem of this is graduated from 0° to 40° , 0° representing pure water, 40° representing the specific gravity 1040. This range is sufficient for all uses, and the instrument is readily carried in the pocket, and is so short that it floats in an ordinary quart measure. With this instrument any milk that stands above 33° is pretty sure to be skimmed, while that which falls below 29° is equally sure to be watered. The advantages of this instrument over that in common use are that no standard is assumed on the instrument itself, and its finding is merely a plain statement of facts."

The two most common frauds in milk are the selling of skimmed for pure milk, and the addition of water to increase its bulk and so augment the profits of the salesman. The creamometer will show the proportion of cream in any given specimen, and the lactometer gives the specific gravity, and so declares the amount of water present. But, let it be remembered that the specific gravity does not show the amount of cream. The specific gravity of good milk is stated somewhat differently by different authors. Parkes (*loc. cit.*, 241) says: "The specific gravity varies from 1026 to 1035. A very large quantity of cream lowers it, and after the cream is removed the specific gravity may rise. The average specific gravity of unskimmed milk may be taken as 1030 at 60° F., and the range is nearly 4° above and below the mean." The *Journal of Food, Water, and Air, in Relation to Public Health*, London, edited by Dr. A. H. Hassell (No. 1, November, 1871, p. 3), says: "A genuine milk of good quality should be white, opaque, of a sweet taste, have a specific gravity of about 1030, but not unfrequently ranging from 1032 to 1027, and should yield from 7 to 10 percentage, by measure, of cream, the average being $8\frac{1}{2}$ per cent." Dr. Parkes gives (*loc. cit.*, p. 242) the following table, which we reproduce for the guidance of our readers. He says: "The addition of water is best detected by the specific gravity. No doubt the method is not perfect, but its rate of application strongly recommends it. The following table shows the specific gravity at 60° , with the addition of different quantities of water, as determined by several experiments:

				Sp. gr.	Sp. gr.
Original specific gravity, . .				1030.5	1026
9 milk and 1 water, . .				1027	1023
$8\frac{1}{2}$	"	$1\frac{1}{2}$	"	1025	. .
8	"	2	"	1024	1019
7	"	3	"	1021	1017.5
6	"	4	"	1018	1016
5	"	5	"	1015	. .

We found that the specific gravity of a specimen of excellent milk, as ascertained by the hydrometer, was 1028. When to this milk was added one-fourth part of water, the specific gravity fell to 1024, and, when a half had been added, it fell to 1020. In another specimen, the specific gravity, obtained in the same way, was 1030 at a temperature of 64° F. When, to that specimen, one-half water was added, it fell to 1020.

By these three simple methods of examination, the acidity or alkalinity of the milk, the proportion of cream, and the proportion of water, can be determined. If the milk is either strongly acid or alkaline, it is not to be trusted. If it be strongly acid, it has undergone the acid fermentation, and is not fit for use. If it be strongly alkaline, it has either been adulterated by the addition of an alkali, probably, according to Dr. Parkes, carbonate of soda, to prevent or arrest the lactic acid fermentation, or it may have been taken from a diseased cow. Dr. Parkes suggests the latter probability in a doubtful way. Dr. J. F. Simon, of Berlin (*Animal Chemistry with Reference to the Physiology and Pathology of Man*, vol. ii, p. 67), states, that he analyzed milk drawn from the teat of a cow having vaccinia, and found it strongly alkaline, and showing with the microscope mucus and pus corpuscles, while that drawn from a healthy teat had a mild acid reaction, and contained no pus or mucus corpuscles. He also states (page 68), that Herberger has analyzed the milk of cows suffering from the grease, and found it to contain an increased quantity of the alkaline salts, in the first stage; in the second stage it was thick and viscid, and had, besides, an unpleasant and putrid taste and smell. In both stages, the presence of carbonate of ammonia (an ingredient never before observed in the milk) was detected.

The mother will often wish to preserve milk, especially in our hot summer weather, or for a few days, when on a journey. The best preservative in hot weather, for the day, is of course a good ice-chest. Dr. Parkes says that when boiled, "the bottle quite filled, and at once corked up and well sealed, the milk lessens in bulk, and a vacuum is formed above. It will keep thus for some time. A little sugar aids the preservation. If the heat is carried in a close vessel to 250° Fahr., the milk is preserved for a very long time, even for years; the butter may separate, but this is of no consequence;" or, if a little carbonate of soda and sugar are added, without boiling, he says it will keep for ten days or a fortnight. Cooley, in his *Cyclopædia of Practical Receipts*, states, that the addition of ten to twelve grains of carbonate, or bicarbonate of soda, to each pint of milk, will preserve it for eight or ten days in temperate weather, and adds that this addition is harmless, and, indeed, is advantageous to dyspeptic patients. The method of boiling, proposed by Dr. Parkes, is the one now so much used for preserving fruits fresh.

We have to consider, next, the subject of artificial or hand-feeding. Our remarks will include the new-born, the early weaned, and the period of the first dentition. Any one who has observed the results of artificial feeding of young infants as exhibited in the statistics of foundling hospitals abroad, wards for foundlings in our own almshouses, or hospitals for children; or who has watched for years, as we have, the comparative success

of natural and artificial feeding, even in the houses of the educated and wealthy, will confess the primary importance of this subject. It covers, moreover, very extensive ground, and exhibits surprising differences of opinion amongst high authorities. We shall follow our usual plan of laying before the reader what is largely the result of our own experience and observation.

In prescribing an artificial food to be made of cow's milk, three points demand special attention: 1, The mode of preparing the milk at different ages; 2, the quantity of food to be given each day; 3, the number of meals into which this quantity should be divided.

We shall not attempt to go deeply into the mysteries of the organic chemistry of milk, but, in order that the reader may follow us in our statements, we will lay before him what we believe to be the most correct analysis of human milk. In order, however, that he may understand the difficulty of the subject, and see that, after all, experience must be our chief guide, we will say that the analysis we select is not the one most quoted, and most relied upon, but is that of O. Henri and Chevallier, which is quoted by Dr. Letheby in his *Lectures on Food* (page 131). We find that Dr. Edward Smith follows the analysis of Vernois and Becquerel, published in 1853. Professor Kehrner (*German Clinical Lectures*, Syd. Soc. Ed., 2d series, p. 364) quotes an analysis from Gorup Besanez, which, however, is exactly the same as that given by Dr. Edward Smith, and as those made by Vernois and Becquerel. Believing that this analysis showed too much caseine and too little sugar, and perplexed by the uncertainty of the whole subject, we requested Dr. Arthur V. Meigs, of Philadelphia, to make some fresh analyses of both human and cow's milk, in order, if possible, to clear away some doubts we have had. He has not finished his examinations, but allowed us to publish the following statement of some of the conclusions he has reached. He says: "The question whether young infants that have to be artificially fed should be given pure cow's milk, may be answered in the negative, for two reasons: First, experience teaches that those fed upon a diluted and properly mixed milk are more apt to thrive than those given it pure; and, second, a comparison of human and cow's milk shows that the two are very different."

"The analyses of human milk made thus far may be divided into two classes, and we may take as types of these the analysis of O. Henri and Chevallier, and that of Vernois and Becquerel. I give the two mentioned, and add one of my own of good ordinary cow's milk for comparison.

	V. & B.	H. & C.	Cow's Milk.
Fats,	2.666	3.55	3.544
Caseine,	3.924	1.52	4.405
Sugar,	4.364	6.50	4.278
Salts,138	.45	.764
Total solids,	11.092	12.02	12.991
Water,	88.908	87.95	87.009
	100.000		100.000

"On comparing the two analyses of human milk quoted, it will be perceived that the greatest difference between them is in the percentage of caseine; it may be further seen that in the analysis giving the low estimate of caseine a large estimate of sugar is given, and, in the other, exactly the opposite is the case: the amount of caseine is large, and that of sugar is (comparatively) small. If the total caseine and sugar amounts in the two analyses are compared, it is seen that the sums are nearly the same (8.02 per cent. in one, and 8.28 in the other). I am convinced from experiments of my own that the analysis of Vernois and Becquerel is wrong, that their method classed as caseine a considerable amount of the sugar, and that the other analysis is much nearer the truth. My own experiments proved conclusively that no specimen of human milk I have analyzed contained so much as two per cent. of caseine, whereas Vernois and Becquerel place it at nearly four. Most authors agree that human and cow's milk are very different, that the amount of caseine in human milk is less, and that of sugar greater than in cow's milk, and yet if the analysis of Vernois and Becquerel is compared with the analysis of cow's milk I have placed beside it, it will be perceived that the amounts of caseine and sugar in the two kinds of milk are almost identical. If we accept this analysis as correct, we must give up the old doctrine that human milk contains less caseine and more sugar, and confess that the two are alike in the percentage of the proximate principles contained, except that the cow's milk is richer in salts.

"I wish to be clearly understood as believing that the figures of Vernois and Becquerel and of their followers are totally wrong, and that deductions from their results are largely accountable for the fact that at the present day so many young infants are fed upon cow's milk, pure, which contains an amount of caseine their stomachs are unfit to digest; and I further believe that if physicians in general could be taught to know that fresh cow's milk properly watered, with cream and sugar added in due proportions, is more like human milk than any other food at the present time known, many thousands of infants who now die would live to be men and women.

"My own experiments have not yet been published, as they are still incomplete. I hope, however, soon to have them in such form that they may be offered to the profession."

We believe these statements to be very near the truth. They agree with the results derived from experience, and with the ocular appearance of the two milks—the human and that of the cow. They show why breast-milk is so thin-looking and watery, when compared with cow's milk by the naked eye. The large excess of caseine in cow's milk (quite the double, and sometimes more) gives to this fluid a thicker and richer appearance, and forces upon the infant, when it is taken pure, the effort to digest twice the amount of caseine in the same bulk of water as in human milk. This very effort, we believe, deranges the digestive functions of the child. The appetite is impaired, and often the child takes so little food as to lose much of the water the system absolutely needs. We have cause to think that this loss of water may be one of the principal causes of the

deranged health, from which children fed on pure cow's milk so often suffer. A curious empirical fact, tending to bring out the same conclusions, to wit, that cow's milk ought to be diluted, and that water is one of the most essential elements in infants' food, is derived from the methods in which condensed milk is used. We shall have more to say on this point when we take up the subject of condensed milk. We will now state simply that in Philadelphia those who use condensed milk employ it in such proportion, generally, that the mixture represents only 1 part of fresh milk to 2 parts of water, while some employ it in even weaker proportions.

Cow's milk should never be given pure to young infants, including in this term children under six months, and the rule ought to be the more stringent the nearer we get to the new-born. This is our conviction. We still believe that the old rule of 2 parts of water to 1 of milk, is the proper one during the first and second months, and also for older children when they have been suddenly weaned, and are placed for the first time on artificial food. We know that some recent writers, and some physicians of this city, use, or try to use, undiluted milk for the youngest children, whilst others give it half-and-half, or 2 parts milk to 1 of water. Some children are said to do well on pure milk. We can only say that we have not yet met with them, and we still believe that when milk is of full average richness (containing 10 per cent. cream, and having a specific gravity of 1030), the old rule of 2 parts milk to 1 of water, is the safest. We expressed these same opinions in the first edition of this work, some thirty-three years since, and all our experience, reading, and cogitations have but confirmed us in them.

At the age of six weeks to two months, the proportion of water may be increased to one-half, but the change must be made with circumspection. If the infant be delicate and colicky; if the stools show small undigested portions of milk, instead of being smooth and homogeneous; and if they are whitish in color, instead of yellow, as they ought to be, the change had better be deferred for a time. When it is determined upon, it ought to be made gradually; one meal per day at first, then two, and so on until the change is effected. At the age of five or six months, the proportion may be increased to 2 parts milk and 1 water, this change, like the previous one, being made gradually, and with care. In the second year of life the milk may be given pure, though, even at this age we have met with a good many instances in which the constant addition of a fourth or a third water has rendered the food more digestible, and productive of better results. So long as a child, even six months old or older, thrives well on milk thus diluted, there can be no valid objection, especially during the hot season, to the practice.

We take up next the consideration of the amount of food necessary each day at different ages. This is a matter of primary importance, and yet it is treated in most medical works only in a cursory way. It is clear that the child's appetite, its spontaneity in taking food, will afford often the best criterions as to the amount that ought to be given. This rule of appetite is, however, much more reliable in nursing than in hand-fed children,

since the former live under more natural conditions. A nursing child, when it takes too much, regurgitates the excess with but little trouble to itself. For some reason this is not so much the case with hand-fed children. They do not regurgitate an excess of food so constantly as does the nursing, and when they do, the act has more of the appearance, and, probably, more the effect, of regular vomiting.

The physician ought, therefore, to know accurately the amount of daily food necessary for hand-fed children. He ought to be able to answer with precision the question of the mother or nurse as to how much food is to be used in the twenty-four hours; when, too, the child is feeble or unwell, when it has naturally a small and deficient appetite, there ought to be some fixed standard by which to direct the amount, as well as the nature, of the food.

We can conceive of but one absolutely safe rule by which to determine the amount of food requisite for young children, and this is to find the quantity which nature supplies. It is most curious how few estimates of the quantity of milk furnished by women have been made, and still more curious to see how greatly these estimates differ. To give the reader as clear and satisfactory a view of this matter as possible, we shall quote the best statements we have been able to find, and then give the results of our own observations.

One of the most distinguished recent writers on the diseases of infancy is M. Parrot, of Paris. In his late work (*L'Athrepsie*, Paris, 1877), he quotes Dr. Natalis Guillot ("De la norrice et du nourrisson," *Union Méd.* 1852, p. 61), as having endeavored to ascertain the amount of food taken by infants at the breast. Guillot weighed the child before and after nursing, to determine the amount ingested. He did this only once in the day, and then multiplied the amount by the number of times he supposed the child to nurse. He supposed that a young infant nursed from twenty to thirty times in the day, and assumed twenty-five as the average number. He concluded that a child two days old takes 21 ounces; one five days old, 78 ounces; and one eighteen days, 91 ounces of milk per day. M. Parrot says of these statements: "As we shall soon see, these figures are entirely too large." M. Parrot quotes also an inaugural thesis published in 1864, by M. Bouchard, resident physician in the Maternity Hospital of Paris. M. Bouchard weighed the children at each nursing, the number of which in the day, instead of being twenty-five as Guillot supposed, was but eight or ten. He determined in this way, the average quantity taken each day by children from birth to nine months, to be as follows: First day, 1 ounce; second day, 5 ounces; third day, 14 ounces; fourth day, 17 ounces; after the first month, 20 ounces; after the third month, 23 ounces; after the fourth month, 27 ounces; and from six to nine months, 30 ounces. Of Bouchard's results, M. Parrot says: "These figures are much smaller than those of Natalis Guillot. I accept them entirely, after having proved their exactitude by observations of my own."

M. Parrot makes some statements also in regard to the amount of artificial food taken by children at different ages. We quote from a table obtained by weighing twelve children at the crèche of the hospital, before

and after being fed on cow's milk. They were fed six times in the twenty-four hours. The amounts of food taken and determined in this way were as follows: First day, 5 ounces and 5 drachms; second day (average of three children), 4 ounces and 5 drachms; third day, 5 ounces and 5 drachms; fourth day (average of two children), $7\frac{1}{2}$ ounces; fifth day (average of two children), 7 ounces; eleventh day (two children,) 5 ounces; first month (two children), 7 ounces; second month (two children), 15 ounces; six months, 20 ounces. He says finally: "I think I have shown that $9\frac{1}{2}$ ounces for the first month, 19 ounces for the second, third, fourth, and fifth months, and 25 ounces for the sixth, represent in all cases, an amount of milk sufficient to nourish children raised on the sucking-bottle, with the express condition, that the milk be pure and of good quality, and that, if diluted as some physicians advise, sugar should be added in certain proportion, $7\frac{1}{2}$ drachms in the first month, 10 drachms in the next four months, and $12\frac{1}{2}$ drachms after the sixth month. In my opinion, it is always preferable to give the milk pure." He advises further, that after the sixth month, the ratio shall be increased by from $4\frac{1}{2}$ to $6\frac{1}{2}$ ounces per month, or else, and he prefers this latter plan, that gruels or soups shall be added to the food.

Dr. J. Lewis Smith, of New York (*The Sanitary Care and Treatment of Children and their Diseases*, essays published by the Thomas Wilson Sanitarium of Baltimore, Boston, 1881, pages 295-6), gives some estimates of the amount of milk furnished by the breast. They were obtained by weighing children before and after the act of nursing. In one table of twelve children it is shown that "each of the infants, who were all under the age of five weeks, and all but one under that of twenty days, nursed in the average 12.41 ounces of beast-milk in twenty-four hours, and as the average number of nursings for each during the day was 11, the quantity of milk received at each nursing averaged only a little more than 1 fluid ounce, 1.12." In a second table are given the results of observations on fifteen children from five weeks to two months old. "The average quantity of milk which these infants, who were all well nourished, received in the twenty-four hours, was 24.65 fluid ounces. The quantity received at each nursing was 2.73 fluid ounces in the average."

We shall now give our own observations. In the first edition of this work, in 1848, it was stated that from various inquiries and observations we had been led to believe that a healthy infant of two or three weeks old, would receive from a good nurse and digest well about a pint of food in the twenty-four hours, and that, by the end of the first, and in the second month, the quantity taken would have increased to a pint and a half or a quart. Some of the data upon which these assertions were made were as follows: A woman, attended by one of us in her confinement, had a pint of milk drawn by the nurse daily from the breasts, in addition to what the child took. On asking the nurse how much she supposed the child—a vigorous, hearty boy—might take, she replied that, judging from the frequency and vigor with which he nursed, she supposed he might take as much as was drawn from the breasts. Another patient lost her child at birth, and, desiring to go out as wet-nurse, kept up the flow of her milk

by using a puppy. Six weeks after her confinement a good breast-pump was given her, and she was desired to keep all the milk she could obtain in twenty-four hours. It measured exactly a quart.

It was stated in that edition that careful inquiries were made in regard to this matter of one of the most experienced and intelligent nurses we ever knew. She was desired to answer accurately the two following questions:

1. How much milk do you think a healthy mother gives to her child daily, after the flow is fairly established?

2. What quantity of artificial food do you give in twenty-four hours to infants you are compelled to feed exclusively?

The reply to the first question was that she had often drawn more than a pint from the breasts in the twenty-four hours, in addition to what the child took, and that she had frequently drawn as much as three pints from women who had lost their children. She supposed, therefore, that a hearty child would take, during the first two weeks, at least a pint, and much more afterwards.

To the second question she replied, that she usually gave to hearty children of one, two, and three weeks old, a pint of good milk in twenty-four hours.

Since that time we have had two excellent opportunities for ascertaining the amount of nutriment supplied by nature to young children. A child four months old, who had had a painful and tedious suppuration from an injury to the scapula during birth, and who had not yet recovered, suddenly weaned himself from his mother, who had nursed him in large part, though not wholly, up to this time. The child was fed for a time upon diluted cow's milk and Mellin's food, but, becoming very ill, a wet-nurse was sent for. It was utterly impossible to induce him even to touch the breast. The milk was drawn with a breast-pump and fed to the child from a small sucking-bottle. At this time the wet-nurse's child was two months old. At first only small quantities, 1 and 2 ounces, were taken and retained. Any larger quantity was rejected by vomiting at once. The doses were gradually increased, until, at the end of several days, 36 ounces of the breast milk were consumed daily. Besides this amount, which was drawn by the breast-pump for the sick child, the wet-nurse nursed her own infant several times a day, and, judging from the amount of artificial food the child took, we inferred that it might get from the mother a pint of milk daily. This woman supplied daily, therefore, at the end of the second month of lactation, 3 pints, 48 ounces of milk.

On another occasion a child born of a healthy young woman, was unable, owing to a defective development of the palate, to nurse from the breast. The milk was drawn from her by a breast-pump and fed to the infant from a small sucking-bottle, with unusually large apertures in the mouth-piece. When this child was five and six weeks old it was taking from 18 to 23 ounces of milk daily. The amount obtained by the breast-pump was much larger than this. Accurately measured each day, it was $39\frac{3}{4}$, 41, $33\frac{1}{4}$, 39, $39\frac{1}{4}$, $39\frac{1}{4}$, $31\frac{1}{4}$, $41\frac{1}{2}$, $44\frac{1}{2}$, 35, 40, and $39\frac{1}{4}$ ounces. The largest daily secretion in the fifth and sixth weeks of lactation, was, there-

fore, $44\frac{1}{2}$, and the smallest $31\frac{1}{4}$ ounces. It is reasonable to suppose that, had the child been vigorous, and fit to solicit the flow of milk in the natural method, the mother, who had all the qualities and instincts of maternity in the highest degree, would have had a still larger supply of milk.

In the *Dictionnaire de Médecine*, by Littré and Robin, it is stated under the head of milk, that each breast gives from 25 to 30 grams per hour, or 1440 grams per day, for both. That is about 44 ounces. It is added that Lamperrière (1850) found it to amount to 2144 grams (64 ounces) in some nurses.

We have given now the most reliable estimates we have been able to find of the amount of food supplied by nature to the young child. The differences in the estimates by different authors are certainly very curious. The small amounts stated by MM. Parrot and Bouchard, as compared with those set down by Guillot and Littré, amongst French observers, and with our own are remarkable. The estimates of Dr. Smith are considerably smaller than ours, or those of Guillot, Littré, and Lamperrière, though they were taken with such care that it is difficult to reconcile the discrepancies. We shall assume our own observations as our special guide, for the reason that the milk was on each occasion drawn from the breasts, and accurately measured. Moreover, when we come to consider the quantities of artificial food to be used, we shall find some reason for believing that our own larger estimates as to the amount of food necessary for infants are probably correct.

Before setting forth our own opinions as to the quantities of artificial food proper for young children at different ages, we shall quote the estimates given by M. Parrot upon this point. M. Parrot (*loc. cit.*, p. 435), as has already been stated, advocates the use of pure cow's milk at all ages,—for the newborn as well as for the older child. He ascertained by weighing twelve children of different ages, and chosen from amongst the healthiest in the hospital (*Enfants Assistés*), before and after the use of the sucking-bottle, that the child would take $9\frac{1}{2}$ ounces in the first month; 19 ounces in the second, third, fourth, and fifth months; and 25 ounces in the sixth month. We think these quantities are much too small, not that the amount of pure milk is so deficient, but that the quantity of liquid nutriment is too small. Nature gives much more in bulk, but in a more dilute form, and we believe this is in order to introduce a larger amount of water into the body. We think that the water thus introduced into the organism has its own physiological uses, and that a failure to supply the simple element in sufficient quantity, is a capital error in the attempt to bring children up on artificial food. Moreover, we have frequently known hand-fed children, in our own experience, to consume much larger amounts of food than those given above, and to thrive admirably. Indeed, it has been these hearty-feeding infants who have been the healthiest we have seen. In one case a fine, vigorous boy, twelve weeks old, took in each twenty-four hours a quart and a half pint of good cow's milk mixed with the same amount of water. He was fed at 11 P. M., and again at 6 A. M., and then every two and a half or three hours during the day. Another child, at four months, took two quarts of a food made of milk, cream, arrow-

root-water, and gelatine. A third, eight months old, took, three pints of food per day. One of our patients was in the habit of giving her children (she was forced to wean them very early), at three months of age, a quart of cow's milk mixed with a third water.

As this matter of the quantity of artificial food necessary for the development of the child is a very important one, and as it is a point which has not been very clearly defined by most writers, we have thought it well to lay before our readers the following calculation of what infants may need, from the estimate made by Dr. Parkes as to the amount of food necessary for adults.

According to that author, an adult of average size and activity will, under conditions of moderate exertion, take in twenty-four hours from $\frac{1}{25}$ th to $\frac{1}{20}$ th of his own weight in solid and liquid food. The relative proportion of the so-called solid and liquid food varies greatly, but is usually about 40 ounces avoirdupois of the former, and 60 ounces of water. As, however, all the so-called solid food—bread, meat, etc.—contains a certain amount of water, the actual average amount of water-free food taken by an adult weighing 150 pounds is 23 ounces, or $\frac{1}{105}$ th of the weight of the body; and the amount of water about 75 ounces. Or, in other words, every pound weight of the body receives about 0.15 ounces of water-free food and 0.5 ounces of water in twenty-four hours. This water-free food is composed as follows, according to Moleschott:

	oz. avoirdupois=437.5 grains.
Albuminous substances,	4.587
Fatty "	2.964
Carbohydrates,	14.257
Salts (of all kinds),	1.058
	<hr/>
	22.866

On the basis of these calculations, an infant at birth, the average weight being 7 pounds, would require 1.05 ounces of water-free food; and a child weighing 20 lbs., which is probably the average weight of healthy children of five to six months old, would require 3 ounces.

Assuming the total solid of cow's milk to be 10 per cent., which is rather less than the average as given by Becquerel and Rodier (see composition of healthy milk), it would require to yield an ounce of water-free food rather more than 10 ounces of milk.

Thus on this supposition (*i. e.*, that the total solids of cow's milk of sp. gr. 1026 equal 10 per cent.) one pint imperial (20 oz.) will contain in round numbers,

Caseine,	262 grains.
Fats,	217 "
Lactine,	341 "
Salts,	43 "
	<hr/>
Total,	863 " = very nearly 2 oz. avoirdupois of water-free food.

According to this, therefore, the infant at birth requires little more than

$\frac{1}{2}$ pint imperial of unskimmed cow's milk ; the child at five or six months about $1\frac{1}{2}$ pints imperial.

It is evident that the proportion of fat and water is in great excess in this exclusively milk diet ; but these two principles are required in early infancy in much larger relative amount than at a later period of life. It will also be seen that by diluting the above amounts of cow's milk with one to two parts of water, we obtain, as the proper amount of food for newborn infants, from a pint to a pint and a half ; and for children about five or six months old, from 3 to 4 pints, amounts which correspond closely with the results obtained from examination of the quantity of milk secreted by nursing women.

We resume our consideration of the amounts of artificial food required at the different ages of infancy.

In the first two or three days after birth the child ought to be fed every two hours from early morning until the evening, say six times, and then four times in the night, making ten feedings in all. Each feeding ought to consist of about two tablespoonfuls, or one ounce, making in all ten ounces. From the second or third day to the tenth the feedings may be at the same intervals, but should consist of about three tablespoonfuls, or an ounce and a half, amounting to fifteen ounces, or very nearly a pint. From this time to the end of the first month a vigorous child increases rapidly in appetite and in the power of assimilation, and will be taking generally from a pint and a half to two pints.

It is highly important that the child should have during this early stage an experienced and careful nurse, or, when this cannot be obtained, as amongst the poor or in hospitals, that the physician should lay down the most minute and particular rules for each individual case. For each baby, like each adult, is a law to itself, and the doctor in charge, or the nurse, must, by observation, determine this law as far as may be possible. It is wise to begin with the smaller doses of food, and, after a day or two, to increase with care. The nurse should watch the child closely,—how it eats, whether with appetite and enjoyment ; whether it grows hungry within the proper time, one and a half or two hours after the previous meal ; the condition of the stools ; how it sleeps, and how it behaves when awake. So long as the child is contented, crying only moderately from time to time, when it is soiled or wet, when it is taken up to be changed, or when hungry, it is doing well, and the dose of food may be gradually increased as the appetite grows. The child should never be forced, or persuaded, to take more than it wants, except when the amount consumed in the twenty-four hours is manifestly below the healthy standard for the age. In such cases tonics should be given, or some change made in the food.

We are thus particular, because a young infant once seriously disturbed in its health, by either improper food or by overfeeding, or the opposite, under-feeding or innutrition (the *athrepsia* of Parrot), often falls into a state from which it is very difficult to extricate it. We deem it all-important, therefore, that a newborn child which must be hand-fed shall have the strictest care during the first few days and weeks of its life. We are satisfied that there is no comparison between the results of hand-

feeding in hospitals and amongst the very poor, and in families in easy circumstances, where education gives knowledge and care, and where the child has devoted to it always one person, the mother, and often two, the mother and nurse. Familiar as we are with the details of private practice, and knowing the fact that one young infant, especially if it be a delicate and sickly one, will absorb the whole time of one person, and often wear out her health, we are not surprised at the misery and fatality which so abound in hospitals for foundlings.

In the second month the child will probably still require a meal every two hours or two hours and a half during daylight, and twice or three times in the night, making about eight or nine meals a day. The amount of food at each meal ought to be about 4 ounces (a gill), making 32 ounces in the day. Towards the end of the second and in the third month the rule ought to be, in healthy children, once in three hours during the day, and twice in the night, or about seven meals. These may now amount to 5 or 6 ounces at a time, or from 35 to 42 ounces per day. Some children, as we have shown, are furnished by nature, at the time, with 48 ounces per day.

As the age increases 8 ounces may be given at a time,—five times between six in the morning and ten at night, and once in the night, making five or six meals, and therefore 40 to 48 ounces per day. This amount of food is scarcely greater than in the second and third months, but, by this time, it is much stronger, being composed of milk diluted only a fourth or third, or possibly undiluted, or it may be combined with some farinaceous substance; or probably some animal broth, or bread or cracker, is being taken once or twice besides the milk. It is proper to repeat that the physician must study the appetite of each child. Some, at the age of six and eight months, take with appetite and perfect results two quarts of liquid food in the day, and this is not so rare as we at one time supposed. On the other hand, we think that the child should not be obliged nor coaxed to take more than it fancies, unless the daily *quantum* falls decidedly below the averages given above. In this event, there ought to be no hesitation in coaxing, in gently forcing, the child to take more than it cares to take, and, if the quantity is still too small, the meals ought to be made more frequent again. We have known a number of children so constituted, that even when at a bounteous breast of their own mothers, they would have to be taken into a quiet room in order to be coaxed and enticed to nurse. In children of this type, with careless and deficient appetite, it is the business of the nurse to carry out the general rules of the physician as far as practicable. “*L'appetit vient en mangeant*,” say the French, and we believe there is truth in the saying.

The food must be sweetened, for both chemical analysis and taste show that woman's milk has a larger proportion of sugar than that of the cow. In both milks the variety of sugar is the same, sugar of milk or lactine, and we advise the use of this variety, when it can be obtained, for infant food.

In calculating the amount of sugar to be added to the diluted cow's

milk, we have used the analysis of Henri and Chevalhier (see page 308), which gives the proportion of sugar in human milk at 6.50 per cent. To dilutions of two-thirds water and one-third milk, there should be added about $6\frac{1}{2}$ drachms of the sugar of milk; to dilutions of half and half, the quantity of sugar to be added is $5\frac{1}{2}$ drachms to the pint. If cane-sugar is used, only half the above quantities should be employed.

Many different and more or less complicated preparations of food have been recommended by various authorities. The different feculent substances, so much vaunted and advertised for the use of the public, are of no value in the early months as compared with milk. Milk must be the basis; it is the essential part of the nutriment. To depend altogether on amylaceous food is to sicken and finally starve the child. Yet experience seems to have shown that a small quantity of starchy material combined with the milk does sometimes render the food more digestible. We think the opinion, held by several authorities, that the particles of starch being interposed between the elements of the caseine lessen the tendency of the milk to coagulate into large and hard masses may be a correct one.

The following preparation of food was published in our first edition. We have employed it for many years, and have found it one of the best substitutes for the natural aliment. It is made of prepared gelatine or Russian isinglass, cow's milk, cream, and a very thin arrowroot-water, properly sweetened. The gelatine was introduced originally in imitation of the old German writer, Struve, who maintained that inasmuch as woman used both animal and vegetable food, whilst the cow is herbivorous, it was proper to add some animal material to cow's milk, in order to bring it into closer resemblance to human milk. We have retained it for the simple reason that the food thus prepared has answered so good a purpose. To make this food, a scruple by weight of the isinglass or gelatine, or a portion of gelatine cake, two inches square, is soaked for a short time in half a pint of cold water. The water is then boiled until the gelatine is fully dissolved, about fifteen minutes. A small teaspoonful of arrowroot, mixed into a paste with a little water, is then stirred into the boiling water, after which the milk is added and allowed to boil for a few minutes. At the end of the boiling the cream is added. The proportions of milk are those already laid down: for the youngest children, one-third; and for the older, one-half or two-thirds. Of cream, two tablespoonfuls are added to a pint of the food, so long as this is one-third milk. When the food is half milk, one tablespoonful and a half of cream to the pint is the proper quantity, and when the food becomes two-thirds milk, one tablespoonful is to be added. Of sugar, the proper proportions are those given above.

We have used this food a great deal for over thirty years, as well in children brought up entirely by hand as in those partly suckled, and, on the whole, it has answered better than any other combination we have tried. In a good many cases it has agreed perfectly well with infants who could not, without vomiting, diarrhoea, and colic, take simple milk and water, chicken-water, or, in fact, any other food. In very sickly children it is often well to dilute it for a time, even more than in the proportions

mentioned above. We add further, that we often hear of its being used by other physicians and by families with very positive success.

It is proper to say that, though we recommend, and, on the whole, prefer, arrowroot for very young children, there are several other starchy foods which are of great value in older children, and in some disordered conditions of health. Amongst these are barley, oatmeal, and wheat. Barley or wheat is preferable for children inclined to diarrhœa, and in case of actual diarrhœa; oatmeal is better for costive children. Of barley, a teaspoonful, powdered (Dr. Jacobi recommends that the whole barley be ground in a coffee-grinder at home), should be boiled for fifteen minutes in a gill of the water used for diluting the milk; of oatmeal, the same quantity. Dr. J. Lewis Smith recommends baked wheat-flour, or wheat-flour boiled in a bag four or five hours. This latter preparation is an old favorite in our city in cases of summer diarrhœa. We have frequently employed it, and have found it very useful. The outer portion of the hard cake made by the long boiling is removed, the inner portion is grated down, and the powder boiled in water to a gruel, and mixed with milk in proportion to suit the age. The gruel should not be made too thick; one or two teaspoonfuls for each meal are sufficient.

There is another mode of using cow's milk, mixed with cream, which we have found very useful in the sudden gastro-intestinal disturbances of infants, whether in the cooler seasons from indigestion, or in hot weather during attacks of cholera infantum, diarrhœa, or dysentery. It is made of thin arrowroot-water, lime-water, cream, and milk, in equal proportions. In disorders of this kind we limit the dose of this food, at first, to four tablespoonfuls every two hours. Between the doses of this food we order cool water, with or without a little brandy (a teaspoonful to half a pint of water), according to the state of the patient, to be offered frequently to the child, allowing it to take all that its thirst craves. After one or two days, if the food is retained well, we increase each dose one-half, and when this has been found to be well digested, the dose is made twice the quantity first named. If the case goes on improving, we diminish gradually the proportion of cream and lime-water, increasing that of the milk, until we get back to the child's regular food.

We are well aware that some high authorities oppose the addition of cream to even diluted cow's milk, on the assumption that the fatty element is very indigestible. But experience has convinced us, as we have already declared, that the moderate use of it we recommend has been often most useful. Our desire throughout this long article on food has been to advise the use of an artificial diet as much like the natural food as it can be made. On scientific grounds, therefore, as well as on empirical, we believe it wise to add cream to diluted milk in such proportion as to restore to the fluid its original proportion of fat.

We will merely add that we have met with a few instances in which young children who could not digest, without serious gastro-intestinal disturbance, even very dilute milk and water, were able to digest and to thrive moderately well on cream and water. For further information on this subject, the reader is referred to the essay published by the Thomas Wilson Sanitarium of Baltimore (Boston, 1881, p. 230).

Condensed milk is now much used as a diet for young children. Some medical men prescribe it habitually, use it in their own families, and deem it a more wholesome food than the ordinary cow's milk sold in large cities; some appear to think it a better food than fresh milk. In view of these facts we propose to consider at some length its nature, qualities, and mode of use, in order that our readers may have a correct understanding of what this new article of diet is.

Dr. Edward Smith, of London (*Foods*, American edition, New York, 1873), quotes a report to an American agricultural society (which we have not been able to find) to the effect that American condensed milk is made from fresh cow's milk, of good average quality, by the evaporation of seventy-five per cent. of its water. When thus reduced, white (cane) sugar is added to preserve it. As to the quantity of sugar added we shall speak further on. Thus prepared, condensed milk is of a thick, semi-fluid consistence, and of a syrupy sweetness. It is put up in tin cans, carefully soldered, for preservation, and for ease of transportation. It keeps when closed, it is said, for years, and, even when the can is opened and kept open, it becomes drier and more solid, but does not spoil for some weeks.

There is another form of condensed milk in which the fresh milk is simply condensed by evaporation, no sugar being added. This, it is said, will keep one, or three or four weeks, but it is usually supplied fresh to city customers every three or four days.

The *Journal of Food, Water, and Air, in Relation to Public Health*, edited by Dr. A. H. Hassall, Vol. 1, No. 12, October, 1872, says that whatever be the rule in America the above companies (two English and the Anglo-Swiss Company) take "considerably less than 3 pints of milk to make 1 pound of the sweetened condensed article. We find, further, that the quantity of sugar added is usually about 19 ounces to one gallon of milk, or about 6 ounces only to 1 pound of the sugared milk." Dr. Thomas K. Chambers (*Manual of Diet in Health and Disease*) states that condensed milk is made by driving off by evaporation about six-tenths of the water of fresh milk. To test this point for ourselves we had 2 pints and 3 gills, (44 ounces), of good fresh milk, weighing 2 pounds, 13½ ounces avoirdupois, reduced by evaporation to 9 ounces by weight. To this we added 6 ounces of white sugar, and found that the 15 ounces of sweetened condensed milk, filled not quite full, but very nearly full, one of the tin cans in which the ordinary preparation is sold. This can held 13 ounces, fluid measure, so that the 44 ounces reduced to 9 ounces by weight, with 6 ounces of sugar added, occupied very nearly the bulk of 13 fluid ounces. The original milk had lost in this experiment 80.22 per cent. of its water before the sugar was added.

The normal nutrient principles contained in milk are supposed to be retained in condensed milk, since nothing is taken from it but the water. The journal quoted above says: "Contrasting the analyses given of the several condensed milks with that of normal cow's milk it is obvious that each tin can does really contain, as stated, all the constituents in fair and proper proportion contained in about three pints of normal cow's milk." Mr. J. Alfred Wanklyn (*Milk Analysis*, American edition, New York, 1874),

says; "I have myself examined the principal brands of preserved and condensed milk which are in the London market, and find that the milk which had been condensed, or condensed and preserved, had been charged with its due proportion of fat." We shall assume, therefore, in our remarks upon this subject, that condensed milk, when honestly manufactured, contains all the nutrient constituents of milk in proportion to the amount of evaporation the original milk may have been subjected to.

It contains, however, another ingredient, of which nature has put none in the milk of any animal,—cane-sugar. The journal quoted above states, as we have already said, that from 6 to $6\frac{3}{4}$ ounces of white sugar are added to each 1 pound or 16-ounce tin can of the condensed milk. We had three specimens of condensed milk, the Borden, the Eureka, and the Anglo-Swiss, analyzed for our own purposes. We regret that in these analyses the quantities of sugar of milk and of cane-sugar were possibly not correctly made out. The separation of the two sugars is, we suspect, a very difficult chemical operation. Mr. Wanklyn (*loc. cit.*) does not attempt it. He classes the two sugars together in his analysis of the sweetened condensed, or, what he calls, preserved milk, and gives the percentage of the two as 56.1, or more than one-half of the preparation. In the analyses made for us, the amount of the two sugars combined was 49.1 per cent. in the Borden, 44.7 in the Eureka, and 48.5 in the Anglo-Swiss. The average of the four analyses is 49.6 per cent. If we deduct from this, the amount of milk-sugar which milk reduced three-fourths ought to contain, 17.60 per cent., there would remain, as the average proportion of cane-sugar in condensed milk, 32 per cent. In the analyses made for us, the other nutritive elements, the fat and caseine, are about what they ought to be. The Borden milk, which contained 23.3 per cent. of water, had 11.5 per cent. of fat, and 14.2 per cent. of caseine. In good cow's milk, taking the Boston standard (Sharples, *loc. cit.*), there are 3.20 per cent. of fat, and 4.30 of caseine, which, at the supposed rate of reduction, about three-fourths, gives very closely the amount of these elements which ought to exist in the condensed milk. In the Eureka brand, which contained 29.3 per cent of water, the fat stood at 9.4, and the caseine at 14.0 per cent. We may conclude, therefore, that these two specimens of American condensed milk, like the English, are what they profess to be, good cow's milk condensed and sweetened.

Having shown what condensed milk is, we propose next to consider its advantages and disadvantages.

The mere conveniences which this food offers to the mother of a young child are immense. It saves all bother with the milkman, and a great deal of trouble at home with the servants. It entails no cold vaults, no ice-chests, no care of milk-pans; it is so easily prepared. Even the doctor who prescribes it escapes much annoyance in regard to the choice and management of fresh milk. Those who believe that fresh milk cannot be obtained in large cities resort to it as a matter of conscience. But this is not true of all large cities, and specially not of Philadelphia. When it is true, condensed milk is doubtless better and safer than stale or spoiled milk.

We turn now to its disadvantages. And first is the fact that there is

no reason why fraud may not be practiced in the manufacture of condensed milk, as well as in the preparation for sale of fresh milk. Fraud may be more rare in the former than in the latter case, for the reason that the responsibility, if fraud be detected, is more easily fastened upon one or two manufacturers than upon the many milk vendors. In one respect the housekeeper is safer against fraud in the case of fresh milk than in that of condensed milk. We have shown how this may be done by any intelligent housekeeper in the chapter on food. The analysis of condensed milk, on the contrary, is a difficult problem, and can be made only by the skilled chemist. To show what has happened in the past, we refer the reader to the *First Annual Report of the Board of Health Department of the City of New York* (April 11th, 1870, to April 10th, 1871), New York, 1871. In a report to the board, by the chemist to the board, Dr. C. F. Chandler, upon this very subject, it is stated (p. 314) that a large number of analyses have been made both of ordinary and condensed milk. "The condensed milk is found, with few exceptions, to be made up of skimmed milk entirely or in part. It is thus robbed of its cream, and is therefore deficient in fat (butter)." It is said, further, to be "a notorious fact that most of the condensed milk companies regularly sell cream in the New York market." If this were true in 1871, it may be true again, though the analyses given in the journal above quoted, of foreign preparations, and those made for us, show no material deficiency in the natural milk solids. We will add that Dr. Chandler found the percentage of fat in one specimen he analyzed to be 1.75 instead of 9.50, 10.80, and 11.50, as in the English analyses, or 9.4, 11.5, and 11.15, as in the analyses made for us.

One disadvantage of condensed milk is the large amount of cane-sugar it contains. It is of such syrupy sweetness to the taste that it must be largely diluted to make it agreeable to the palate, and to reduce the proportion of sugar taken in the food made from it. The proportion is so much larger than what long experience has pointed out as the proper amount to be added to a diet made of cow's milk, and so much larger than the difference between the sugar in human milk and cow's milk, that we cannot but look upon it with suspicion, as being so unlike what nature provides for the young child.

We have found, moreover, in examining this subject, that the amount of milk solids is so much smaller in the diet usually made from condensed milk than in one made from fresh cow's milk, that we doubt whether a condensed milk diet can be as good for children over three and four months of age, as the usual diet made from fresh milk. We are well aware that a great many sensible physicians use it largely during the whole nursing age, and point to many fine-looking and apparently healthy children brought up on it. We will, however, lay before our readers the results of our examination, and they can judge for themselves whether there are not good theoretical grounds for our doubts.

To make this matter as clear as possible, we propose to show how it is generally used, the degree to which it is diluted, and then state the amount of cane-sugar and of milk solids in the food so made.

We have found from observation and inquiry, that the physicians who use it most extensively and most successfully, are in the habit of prescribing it in the nursery in the proportion of one heaped teaspoonful in six tablespoonfuls of water. This is a very loose and uncertain rule. We had several heaped teaspoonfuls weighed. In one trial a heaped teaspoonful, what was called a fairly heaped teaspoonful, measured out by an apothecary, weighed 220 grains. In another case a heaped teaspoonful, also measured out by an apothecary, weighed 435 grains. Another heaped teaspoonful, taken by a physician, weighed 334.9 grains. We then asked a child's nurse, one thoroughly accustomed to nursing habits, to take from a can a fairly heaped teaspoonful. This weighed 199 grains. We had a teaspoon, even full, weighed several times; the weight was about 100 grains. We have, therefore, in our calculations, proceeded on the assumption that a fairly heaped teaspoonful contains twice what a teaspoon even full contains,—200 grains. And when we speak, in our remarks upon this subject, of a heaped teaspoonful of condensed milk, we mean, in fact, two teaspoons even full, or 200 grains.

Assuming, as we have already said we should do, that the average amount of cane-sugar in condensed milk is 32 per cent., we find that when 200 grains (two even teaspoonfuls) are diluted with six tablespoonfuls of water, the amount of this sugar is 2.56 grains in each teaspoonful, counting twenty-five teaspoonfuls in the whole mixture. Dr. Edward Smith, of London, in his work on *Foods*, recommends the addition of 4 drachms of milk-sugar, or 2 drachms of cane-sugar, to each pint of food made of two-thirds fresh cow's milk in one-third water. Such a food contains almost precisely 1 grain of cane-sugar in each fluid drachm or teaspoonful. We have advised, in the chapter on food, that to each pint of food for young infants, made of one part fresh cow's milk to two parts water, should be added, to bring the sugar up to the standard of woman's milk, $6\frac{1}{2}$ drachms of milk-sugar or $3\frac{1}{4}$ drachms of cane-sugar. In a diet made after Dr. Smith's rule, there would be 1 grain of cane-sugar to the fluid drachm, and in that recommended by us, 1.52 grains. In a pint of food made from condensed milk in the proportions cited above, there would be very nearly $3\frac{1}{2}$ drachms more cane-sugar than in Dr. Smith's, and nearly $2\frac{1}{4}$ drachms more than what we have learned to believe, from both practical and physiological reasons, to be the proper amount to add to the diet of very young children.

We pass on next to a consideration of the amount of natural milk solids contained in the food as usually made from condensed milk. In these calculations we have taken the analysis of sweetened condensed milk given by Wanklyn, as being, on the whole, the one most likely to be correct.

Wanklyn gives the water in this preparation at 20.5 per cent., the fact at 10.4, the caseine at 11, the ash at 2, and the two sugars, the milk, and the cane, at 56.1 per cent. We find that in a food made of two even teaspoonfuls, or one heaped teaspoonful, weighing 200 grains, in six tablespoonfuls of water, and assuming the weight of a teaspoonful, or fluid drachm of ordinary water to be 54.68 grains, that the percentages are as follows: Water, 89.48; fat, 1.38; caseine, 1.46; ash, .26; and the two

sugars, 7.42 per cent. Such a food represents very closely one part of fresh milk to two parts of water, and is strong enough, with the addition of a little cream for new-born infants.

When, instead of 200 grains, three even teaspoonfuls, or 300 grains, are mixed with the six tablespoonfuls of water, the percentages of milk solids are as follows: Fat, 1.94; caseine, 2.05; ash, .37; the two sugars, 10.44; water, 85.20. This makes a diet of about the strength of half fresh milk and half water, with (calculating the milk-sugar at 4.40 per cent.) 6.04 per cent of cane-sugar.

If, lastly, we mix five even teaspoonfuls, or 500 grains, in the six tablespoonfuls of water, we obtain the following percentages: Fat, 2.86; caseine, 3.03; ash, .55; the two sugars, 15.47; and water, 78.06. In such a mixture the proportions of the fat, caseine, and ash, approximate very closely to those of normal milk, these being, as we have already stated: Fat, 3.20; caseine, 4.30; ash, .60; milk-sugar, 4.40; and water, 87.50. But the sugar is in such large excess, there being 15.47 per cent. instead of 4.40 per cent., that the food would be sickening in taste, cloying to the stomach, and, in all probability, irritating to the digestive apparatus. The amount of cane-sugar in the twenty-six teaspoonfuls of such a mixture would be over 2 drachms (165 grains), or nearly as much as we think necessary for a pint of food made one part milk to two parts water.

In the mixture made of 300 grains in six tablespoonfuls of water, the amount of cane-sugar is a little over a drachm and a half (99 grains).

The large amount of cane-sugar present in a food made of condensed milk and water, of such proportion as to represent fresh milk, must be, it seems to us, a serious objection to it. It may answer very well for very young infants so long as the proportions are those we mentioned first,—one heaped teaspoonful of 200 grains to the six tablespoonfuls of water, in which the mixture represents one part milk to two parts water, with the added cane-sugar. It may answer well enough when three even teaspoonfuls, or 300 grains, are mixed with the six tablespoonfuls of water, representing half milk and half water. Even in such a mixture the amount of cane-sugar is very large, but when we come to the proportions representing fresh milk, 500 grains in six tablespoonfuls of water, the amount of cane-sugar is excessive. It seems impossible, therefore, to make use of condensed milk when the child comes to the age at which pure milk may be used with safety and propriety.

We do not wish to condemn the use of condensed milk for young children, for we know that many excellent physicians use it successfully, and point to numerous children brought up successfully upon it. We desire merely to call the attention of the profession to the above statement of facts. Personally we prefer the old-fashioned mode of using fresh milk, when it can be obtained good, and are of opinion that only long-continued observation and experience can ever demonstrate that the new system is better than the old one.

We shall now give the opinions as to the value of condensed milk as a diet for children, expressed by recent writers on food, and then add some of our own experiences.

Dr. Edward Smith, of London (*Food*, New York, 1873, p. 323), says : "This preparation has been recommended as a food for infants, and it is much liked by them; but it is an error to assume that a given quantity when dissolved in water will yield new milk or be as useful as new milk in feeding infants and young children, and it should never be used as a substitute in such cases whenever new milk can be obtained."

At page 325 he says again: "Without explaining the medical aspect of the question (which would be out of place here), I remark that as a food the addition of nearly two ounces of sugar to the pint of cow's milk greatly lessens its nutritive value, and induces a tendency to starvation of the muscle-forming element. Thus, whilst in natural cow's milk the proportion of nitrogen (flesh-forming) to carbon (fat-forming) is 1 to 12, in the preserved milk it is not much more than one-half, or about 1 to 20. If the object were to feed an animal for the market it would be obtained by this method, but if to make infants into strong muscular men and women, the proportion which nature has provided must be supplied."

Dr. Smith gives also the views of Dr. Daly (*Lond. Lancet*, November 2d, 1872), who, while noting the fact that condensed milk is much liked by children, and that those who are fed upon it grow fat and look very well, yet gives it as the result of his experience that they have not the same degree of resistance and vital power as those who are fed on cow's milk, but sink much more quickly and dangerously under an attack of diarrhœa or any other acute disease.

Dr. Arthur V. Meigs, one of the assistant physicians to the Children's Hospital of this city, informs us that he has been obliged, in most of the cases of cholera infantum, brought to the dispensary during the hot summer months, to change the diet of those fed on condensed milk to fresh milk, as he has found that they rarely do well on the condensed milk diet. His friend and colleague, Dr. Louis Starr, has arrived at the same opinion as to the comparative value in this disease of the two kinds of diet.

Dr. Thomas King Chambers (*Manual of Diet in Health and Disease*, Philadelphia edition, 1873, p. 65) says, of condensed or Swiss milk, that "it certainly is digestible, as is shown by the fact of infants brought up by hand upon it growing fat and apparently strong, a fact of which most of us have ocular proof. Great care should be taken that only the softest water is used for its solution, and precautions taken against its adulteration. As it is a recent invention it is pure enough at present, but extensive use will probably teach ingenious modes of sophistication." Dr. F. W. Pavy (*Treatise on Food and Dietetics*, Philadelphia edition, 1874, p. 194) gives no personal opinion as to its value, but cites, in a foot-note, Dr. Daly's opinion (already quoted) of it.

We have employed condensed milk a few times, and have had charge of cases in which it had been ordered by other physicians. In one instance, a very feeble infant of six months old, who, when we first saw it, had had frequent indigestions and convulsions on fresh milk, did very well on one heaped teaspoonful of condensed milk mixed with four tablespoonfuls of fresh water and two tablespoonfuls of lime-water, with ten drops of wine of pepsin after each feeding, and a mixture of soda, two and a half grains,

sweet tincture of rhubarb five drops, and paregoric two drops, three times a day. We tried fresh milk several times, but it did not answer. When the child reached the age of fourteen months, it had become reasonably healthy, and we ordered some weak beef tea twice a day, and the substitution of one tablespoonful of fresh cream in place of one of the tablespoonfuls of lime-water. Eventually fresh milk was substituted, and the child has grown into a fine healthy boy. In another case, one of a pair of twins was brought to us at the age of four and a half months—a miserable little, pale, feeble, and undergrown infant. Fresh milk had been tried, but had caused indigestion and diarrhœa time and again. We ordered one heaped teaspoonful of condensed milk in five tablespoonfuls of water and two tablespoonfuls of lime-water. On this the child was much more comfortable, and grew slowly. The food was now increased in strength. Two heaped teaspoonfuls of the condensed milk were added to ten tablespoonfuls of water and two tablespoonsful of lime-water every two or three hours. At the age of eight months the child had grown somewhat, was in more comfortable health, but was still very small, white, and puny. We now ordered one tablespoonful of cream to be substituted for one tablespoonful of the plain water, and also two tablespoonfuls of beef tea made by pouring two tablespoonfuls of hot, not boiling, water, on one teaspoonful of Valentine's meat extract. The child did well on this food for some months, when it was gradually changed to fresh milk, and the ordinary food of older children. The other twin, at this time, looked well, was well grown, on condensed milk food, made by adding three teaspoonfuls of the milk to ten tablespoonfuls of simple water and three of lime-water. Both the children are now (June, 1881) living in very good health, though their mother has died of rapid phthisis.

We add a few cases that have come under our personal observation, to show that the use of condensed milk as a food for children is not yet regulated as it ought to be, by a system of rules based on its composition and nutritive value. These cases show, it appears to us, that when used thus carelessly and irregularly, it may give rise to dangerous disturbances of health.

CASE I.—Called to see a child, six months old, in consultation. The mother had had scarcely any milk at first, and this little soon disappeared. The child was put on a food made of condensed milk, 1 teaspoonful to 6 or 7 tablespoonfuls of water. On this diet it did very well, it was said, at first. Six weeks before we saw the patient, it was removed from the city to the seashore. Three weeks afterwards it had some diarrhœa. A physician was sent for, who reduced the food to 1 teaspoonful of condensed milk in a teacupful of water. We measured the cup and found that it held 16 tablespoonfuls of water. We found, by calculation, that a mixture of 1 heaped teaspoonful (200 grains) of condensed milk in 16 tablespoonfuls of water, contained the milk-solids in the following proportions: Sugar, 3.03 per cent.; fat, .56 per cent.; caseine, .60 per cent.; ash, .11 per cent. The water was at 95.70 per cent. The proportion of the solids in fresh milk are: Sugar, 4.40; fat, 3.20; caseine, 4.30; ash, .60; and water, 87.50 per cent. So that the above mixture represents about one part fresh milk to six of water.

During ten days before we saw the child, it had had vomiting and diarrhœa, and had lost flesh, and had become very weak. It was brought to the city, August 24th, 1878, and the family physician sent for on the 26th. Thinking the child very ill, he desired

a consultation, and we met him in the afternoon of that day. The child looked very ill. It was thin, pallid, distressed, and had had, in the morning, a slight spasmodic seizure. It had three stools in the previous twenty-four hours, dark in color from bismuth, and consisting of much fluid, which had run through the napkins, and some thick, gruel-like, flocculent matter in the centre, without any special odor. The child was taking bismuth and pepsin, and for food, a tablespoonful of chicken tea every hour, and a dessertspoonful of whiskey and water (2 teaspoonfuls to the gill) every hour. We recommended 1 teaspoonful of brandy in a half pint of cold water, to be given from a sucking-bottle (as the child had not learned to drink from a cup). Of this the child was to be allowed to take as much as it desired and could retain. The chicken tea was ordered in a double quantity, two tablespoonfuls every two hours, and the alternate two hours four tablespoonfuls of a food made of equal quantities of thin arrowroot-water, lime-water, cream, and fresh cow's milk. The following prescription was ordered: *R.* Liq. morph. sulphat., f ʒ ss.; acid. sulph. dil., gtt. xxx; elix. curaçoa, syrup. simp., āā f ʒ ij; aquæ, f ʒ iss.—*M.* A teaspoonful every four hours; and the alternate four hours, a powder of 3 grains each of bismuth and saccharated pepsin.

On the following day we found that, owing to some mistake, a teaspoonful of condensed milk in 6 tablespoonfuls of water had been given every two hours instead of the food we had proposed. This had been vomited each time that it was taken. The brandy and water, and chicken tea, had been retained. After this the food proposed above was given regularly, and was retained. It was soon increased in quantity, and the child recovered rapidly.

This case was one, we believe, of innutrition. The dangerous symptoms appeared soon after the amazing reduction in the quality of the food, and they disappeared immediately after the stronger food was resorted to.

CASE II.—Called in consultation to see a child fifteen months of age, who had been ill for a few days. There had been vomiting, a moderate diarrhœa, singular and prolonged fits of coldness, lasting four and five hours, followed by paroxysms of fever, with pulse running to 180. There was present a curious and excessive general restlessness, with jactitations and violent startings, of a tetaniform character. The head was retracted. The patient had been living on a daily diet composed of 4 tablespoonfuls of a farinaceous substance, and 4 teaspoonfuls of condensed milk in 2 quarts of water, which food was taken eagerly. The child grew under this system moderately well, and seemed to be doing well before the illness, except that he had been restless and fidgety, had not slept well, and had passed very large amounts of urine. We recommended an increase of the milk, cold water, and brandy, small doses of opium, and quinia in suppository. The treatment was of no avail. The patient died the next day.

This case may have been one of intermittent fever in a badly nourished subject, but it looks like one of the cases described by M. Parrot in his work on athrepsia, under the title of tetaniform eclampsia, as occurring in young children affected with thrush, and therefore badly nourished.

CASE III.—Called in consultation to see a child, a little over one year old, who had been fed on a diet composed of 4 teaspoonfuls of the unsweetened condensed milk in a half pint of water. We do not know the strength of this milk, but believe it to be much less reduced than the regular sweetened preparation. About six meals were taken daily, containing 24 teaspoonfuls of the milk. The patient had had, for some time, very restless nights. Two days before we were called he had been very ill. The symptoms when we saw him were peculiar. There was intense general irritability, jactitation, startings, so that the case had a tetaniform look; there were attacks, in the night, of singular nervous dyspnœa; the pulse ran to 150 and 180; there was very moderate elevation of temperature. We saw the patient but once, and, regarding it

as one of innutrition, advised the food to be doubled in strength, beef tea to be given three or four times a day, weak brandy and water, and small doses of opium, until the extreme nervous agitation was controlled. The child recovered rapidly.

We believe this case to have been one of violent nervous disturbance, caused by faulty sanguification,—itself the result of deficient and defective nutriment.

CASE IV.—In the spring of 1877 we saw a girl, six months old, whose mother had in part nursed and in part fed the child on a diet composed of fresh cow's milk, water, and sugar of milk. It grew moderately well, and looked well, but was excessively restless and wakeful at night. In July it had a cold, and was put upon a food (not advised by us) made of 1 teaspoonful of condensed milk in a half pint (15 or 16 tablespoonfuls) of water. After this the child became much more tranquil, and slept well at night. In August it was given 1 heaped teaspoonful of the condensed milk in 11 tablespoonfuls of water. The child lived for some months on condensed milk. It grew large, very fat, and became very quiet, indeed, quite sluggish. We advised the mother to use fresh milk again. This was done gradually, and the child is now in good health.

We met with another case in which the physician had ordered 1 heaped teaspoonful in 16 tablespoonfuls of water. The mother afterwards increased the food to 1 teaspoonful in 12 tablespoonfuls of water. We saw a child, five weeks old, who was taking, by order of the accoucheur, 1 even teaspoonful in 27 teaspoonfuls of water. Another child, five weeks old, was taking 2 teaspoons, moderately full, in a sucking-bottle which held 13 tablespoonfuls of water. A physician, a very careful and intelligent one, told us that in the case of his eldest child, the breast failed. They tried different kinds of food, amongst others fresh cow's milk, variously diluted, but the child vomited and got on badly. At six months of age, he began the use of condensed milk, giving 1 heaped teaspoonful in a half pint of water. We have said that 200 grains of condensed milk (1 fairly heaped teaspoonful) represents a mixture of one part fresh milk to two parts water. The above proportion (1 heaped teaspoonful in 14 to 16 tablespoonfuls of water) represents one part milk to six of water (see Case I). On this food the child lived for several months. It was small and delicate-looking, but well. We met with another child, seven months old, large and hearty-looking, whose mother told us that she fed him on condensed milk, 3 full teaspoonfuls to 12 tablespoonfuls of water.

We have cited these cases and facts in order that the reader may see how irregular are the rules for the use of this food. We have made calculations that 1 heaped teaspoonful, or, better still, 2 teaspoons, even full (about 200 grains), in 6 tablespoonfuls of water, is the proportion which has seemed to answer best in the hands of those who use it most. To make it much weaker than this would certainly tend to starve the child. Even in this proportion the amount of milk-solids is insufficient for children over three or four months old, and we suspect indeed that the children who are brought up on it live largely on the cane-sugar which it contains.

ARTICLE II.

SIMPLE OR ERYTHEMATOUS STOMATITIS.

DEFINITION; FREQUENCY.—This form of stomatitis consists of simple diffuse inflammation of the mucous membrane of the mouth, unattended by vesicular or pustular productions, by ulcerations, or by membranous exudation. It is a disease of infrequent occurrence, except in the forming stage of other kinds of stomatitis, and of little importance, seldom requiring the attention of the physician.

The *causes* of the disease are the introduction of irritating substances, such as hot drinks, and acrid or caustic preparations, into the mouth; difficult dentition; and probably sympathy with disordered states of the stomach. It occurs not unfrequently as a secondary affection, particularly in the course of measles, scarlet fever, and small-pox.

The *symptoms* of erythematous stomatitis are more or less vivid redness of the mucous membrane, sometimes diffused, and at others punctated or disposed in patches; slight swelling of the same tissue; heat; and tenderness to the touch, and also in the act of sucking or eating. The child is generally fretful and restless, and either loses its appetite, or refuses to nurse or take food freely, on account of the tenderness of the mouth. There are seldom any general symptoms except in secondary cases, in which they are those of the primary affection.

The *treatment* is very simple. It consists in the use of some demulcent wash, as gum-water, sassafras-pith mucilage, a little honey put on the tongue occasionally, and if the inflammation be at all considerable, in the application of some astringent preparation. This may consist of honey and borax, two or three parts of the former to one of the latter, or of the following wash, recommended by M. Bouchut:

R. Mel Rosæ,	f 3j.
Aluminis,	3ss.
Aquæ destillat,	f 3 ss.—M.

The application of any of the washes recommended is best made by means of a thick and soft camel's hair pencil; or it may be done with a soft rag, which should be dipped in the wash, and then conveyed into the mouth on the point of the finger. The remedy ought to be used several times a day.

If signs of gastric or intestinal disorder are present, they should be attended to.

ARTICLE III.

APHTHÆ.

DEFINITION; SYNONYMS; FREQUENCY; FORMS.—The term aphthæ ought to be restricted to the vascular and ulcerous form of disease of the buccal mucous membrane, in which that tissue is covered with an eruption of vesicles, which break and are followed by small rounded ulcerations. Under this title writers formerly confounded the affection we are now considering with ulcerative stomatitis and thrush. It has been called by Billard follicular stomatitis, and by several other writers vesicular stomatitis.

The *frequency* of the disease is very considerable. We shall describe two *forms*, the *discrete* and *confluent*.

CAUSES.—The only causes which seem to have been ascertained with any degree of certainty, are early age and the process of dentition; the contact of irritating substances, particularly stimulating and acrid articles of food, with the mucous membrane of the mouth; and the existence of some morbid irritation of the digestive tube, especially of the stomach. The confluent form is often connected with severe general disease of the constitution.

SYMPTOMS; DURATION.—Aphthæ begin in the form of small red elevations, having little white points upon their centres, which consist of the epithelium of the mucous membrane raised into vesicles. The vesicles are small in size, oval or roundish in shape, and of a white or pearl color. They soon break and allow the fluid which they contained to escape, after which there remains a little rounded ulcer, with excavated and more or less thickened edges, and surrounded almost always by a red circle of inflammation. The bottom of the ulcers is usually of a grayish color.¹ There is seldom any diffuse inflammation of the mucous membrane in this disease. The *number* of aphthæ varies in the two forms. In the discrete variety there are but few, whilst in the confluent form, they are, of course, much more numerous. They generally appear first on the internal surfaces of the lips and gums, and then on the inside of the cheeks, edges of the tongue, and soft palate.

The discrete form is generally accompanied by symptoms of slight disorder of the digestive organs, consisting of thirst, acid eructations or vomiting, imperfect digestion, and a little constipation or diarrhœa. The confluent form, which is much more rare, especially in very young infants, usually coincides, as has already been stated, with severe general or local disease.

¹ The grayish or yellowish-gray secretion, on the base of the apthous ulcers has been closely studied by Dr. J. Worms (Glasgow Med. Jour., July, 1864), who states that both microscopical examination and chemical tests invariably show its sebaceous nature. It is his opinion, therefore, that aphthæ are the acne of the mucous membranes; in support of which, it will be remembered, that they are found most frequently where the muciparous glands are most abundant.

The *duration* of aphthæ is different in the two varieties of the affection. The discrete form generally pursues a rapid course, lasting, from the beginning to the time of cicatrization, between four and seven days. Sometimes, however, when the vesicles are formed successively, one after the other, the disease lasts much longer. The confluent variety pursues a much slower course, and is much more difficult of cure.

DIAGNOSIS AND PROGNOSIS.—The diagnosis of discrete aphthæ is not at all difficult, in consequence of their being isolated and succeeded by small and limited ulcerations. The confluent form, on the contrary, may be confounded with ulcerative or ulcero-membranous stomatitis, and with thrush. From the first-mentioned disease it may be distinguished, however, by attention to the circumstances that that affection begins by small white patches, and not by vesicles, as do aphthæ; that the ulcerations which follow the patches are covered with true pseudo-membrane; and that the white patches just spoken of appear first upon the gums, whilst aphthæ generally begin upon the posterior surface of the inferior lip, and upon the tongue. From thrush it is to be distinguished by the fact that that disease commences by white points, which are not vesicular, and which, running together, form a creamy exudation; by the absence or very small number of ulcerations; and by the presence of the peculiar fungus of thrush.

Discrete aphthæ constitute a very mild disorder. Recovery always occurs without much difficulty. The confluent disease is more serious, because its progress is much slower, its cure more difficult, and because it is often connected, as has been stated, with some other severe disease.

TREATMENT.—Aphthæ, particularly the discrete variety, require in general, very simple treatment. The means to be employed are *general* and *topical*.

The *discrete variety* usually requires only topical remedies, regulation of the diet, and when there are marked symptoms of gastric derangement, the exhibition of some mild emetic, or of a laxative dose. The *local treatment* should consist of applications of demulcent preparations, as the mucilages of slippery elm, sassafras pith, flaxseed, marsh-mallow root, quince-seed, etc., which are to be used pure when there is no pain, or with the addition of a few drops of laudanum or wine of opium, when the mouth is sore and tender; the aphthæ ought to be touched occasionally with the mixture of borax and honey, or the aluminous preparation recommended for simple stomatitis. The application must be made several times a day with a camel's-hair pencil, a pencil made of charpie or cotton, or with a soft rag covering the finger. When the ulcers which follow the vesicles fail to cicatrize rapidly under the above applications, or when they are numerous and painful, their cure may be very much hastened and the pain quickly relieved, by touching them very lightly with a stick of nitrate of silver, or a piece of alum, sharpened to a point; or we may employ a pencil dipped into a strong solution of nitrate of silver, or into a mixture of one part of muriatic acid to two of honey. Light applications, daily or on alternate days, with a solution of iodoform in ether, 40 to 60 grains to the ounce, lessen sensitiveness and promote the healing of the ulcers. Ether

itself has been highly recommended as a local application by Dr. J. Worms, who, as already stated, has observed the fatty nature of the deposit in aphthous ulcers.

The *general treatment* of discrete aphthæ need consist of nothing more than the use of a simple, unirritating diet in most of the cases. If, however, the digestive apparatus is deranged, the case must be treated according to the symptoms; by antacids or a gentle emetic, when the tongue is foul and the secretions acid; and by the use of a mild laxative, as castor oil, magnesia, or rhubarb, when there is constipation. When diarrhœa is present, we should resort first to a small dose of castor oil or syrup of rhubarb, with the addition of half a drop to two drops of laudanum, according to the age of the child, and afterwards to astringents and opiates, as will be recommended in the article on simple diarrhœa.

The treatment of *confluent aphthæ* must depend on their cause. The local treatment is the same as that for the discrete variety, except that mild cauterization should be resorted to at an earlier period. When they seem to depend upon a general morbid condition of the constitution, as congenital debility, a scorbutic diathesis, or upon chronic affections of the digestive organs, they must be treated in the first place by properly regulated and nutritious diet, and by the exhibition of tonics and gentle stimulants, particularly iron, quinine, and small quantities of very fine old brandy; and in the second case, in the manner which will be recommended for chronic derangements of the stomach and bowels, when we come to treat of the diseases of those organs.

ARTICLE IV.

ULCERATIVE OR ULCERO-MEMBRANOUS STOMATITIS.

DEFINITION; SYNONYMS; FREQUENCY.—This form of sore mouth is characterized by a secretion upon the mucous membrane of a plastic exudation in thick, yellowish, adherent patches, and by inflammation, erosion, or ulceration of the subjacent tissues. It is the same disease as the aphtha gangrenosa, and, we believe, the cancrum oris also of Underwood; the ulceration of the mouth of Dewees and Eberle; the stomatite couenneuse, and the ulcerative and pseudo-membranous forms of the stomatite gangreneuse of M. Valleix; the stomatite pseudo-membraneuse or diphtheritique of some writers; and the stomatite ulcero-membraneuse of MM. Rilliet and Barthez. It is the disease described under the title of gangrenous sore mouth by Dr. B. H. Coates (*North American Surgical and Medical Journal*, vol. ii, 1826), with the exception of a few cases which were what we shall treat of as gangrene of the mouth.

Of the different titles given above, we prefer that of ulcero-membranous stomatitis, as most expressive of the distinctive features of the disease. This form of stomatitis is not very frequent in private practice, but some-

times prevails extensively in hospitals, and other public institutions for children, where it often assumes an epidemic character.

CAUSES.—The *predisposing causes* are epidemic influence, of the existence of which we believe there is no doubt; according to some observers, contagion, which, however, has not as yet been positively shown; and bad hygienic conditions as to cleanliness, ventilation, food, clothing, and habitation. That it is epidemic, we have no doubt from our own experience, since we are rarely called to a case without soon meeting with others, while we sometimes pass several months without seeing a single example of the disease. We have also known it to be endemic in a household, having on one occasion met with seven cases in two families of children residing under one roof, on two other occasions with three cases, and on several others with two. It is most frequent between the ages of five and ten years, though it may attack all ages, and is more common in boys than girls. It occurs occasionally during the convalescence from severe diseases, as pneumonia, the eruptive fevers, typhoid fever, entero-colitis, and other affections of children.

The *exciting causes* of sporadic cases are unknown, with the exception, perhaps, of the presence of a carious tooth in the mouth, and fracture or necrosis of the maxillary bones.

SYMPTOMS; COURSE; DURATION.—The disease begins with slight pain and uneasy sensations in the gums, which then become swollen, red, bleeding when touched, and are soon after covered with a grayish pultaceous exudation of varying thickness. The exudation extends from the gums to the internal surface of the lips and cheeks, and sometimes, but more rarely, to the soft palate, and even to the pharynx and nasal passages. The plastic deposit occurs in the form of small, and slightly projecting, yellowish patches, which approach each other, unite, and form bands of pseudo-membrane, somewhat uneven upon the surface, and adhering with considerable force to the tissue beneath. When the exudation is detached, the mucous membrane is found to be of a red or purple color, bleeding, and excoriated or ulcerated. The ulcerations which exist under the false membrane are of various depths, of a grayish, livid, or blackish color, with swelled, softened, and livid red, or bleeding edges. Those which are formed upon the inside of the lips are rounded in shape, whilst those seated in the angle between the lips and gums are usually elongated. In mild cases of this affection, the local symptoms, though perfectly characteristic, are less severe than those just now described. The ulcerations are often few in number, amounting to four, five, or six upon the tongue, to a few scattered over the inner surface of the lips, and to some upon the gums, and especially about the necks of the teeth. The other symptoms are the same as those above mentioned, with the exception that they are milder in degree.

When the disease is mild, and when it is properly treated, the false membranes become detached, leaving the mucous tissue merely excoriated, in which case it soon regains its natural condition; or else the ulcers which exist beneath rapidly become healthy and cicatrize. In violent cases and in those badly treated, the inflammation, on the contrary, persists; the

pseudo-membranes increase in thickness, or if detached, are formed anew; the ulcerations become deeper, the disease extends, and the case lasts an indefinite period of time.

Other symptoms, besides those we have mentioned, characterize the disease.

The *breath* is always more or less fetid, and in bad cases, almost gangrenous. The *salivary* and *submaxillary* glands are generally more or less swollen, hard, and painful, and according to some authors, the surrounding cellular tissue is in the same condition, though this is denied by others. The *movements* of the lower jaw are stiff and painful in severe cases. *Deglutition* is not affected unless the disease extends to the pharynx. In violent cases there is usually a copious *discharge* of fetid, watery saliva, or of bloody serum, which flows from the mouth during sleep. When the ulcerations are deep and large, the tissues beneath are more or less swollen; the swelling, however, rarely assumes the hard, resisting, circumscribed characters, with the tense, smooth, hot, and shining appearance of the skin, which exists in true gangrene of the mouth. In most of the cases there is a moderate but decided *febrile reaction*, especially at the invasion. This usually subsides or disappears after two or three days, though it sometimes increases if the disease becomes extensive.

The disease begins, as already stated, on the gums, and unless limited to these parts, as sometimes happens, extends to the lips and cheeks. In many of the cases it attacks only one side of the mouth, and this is more frequently the left than the right.

The *course* of the disease is usually rapid in epidemic cases, and in those which are properly treated. Where badly treated, on the contrary, it may last from one to several months, or terminate in gangrene of the mouth.

DIAGNOSIS; PROGNOSIS.—The *diagnosis* is, as a general rule, very easy, if proper attention be paid to the characteristic features of the disease. It has, as already stated, been very often confounded with gangrene of the mouth. The method of distinguishing between the two will be given in full in the article on that disease. From thrush it is to be distinguished in the manner which will be pointed out when that disease comes under consideration.

The *prognosis* is favorable in the great majority of cases. Sporadic cases probably always terminate favorably. The epidemic disease, though rarely fatal, sometimes proves so from its extension to the pharynx and larynx, or from its termination in gangrene of the mouth. We have seen a large number of cases in private practice, and have never as yet known one to become gangrenous or to prove fatal. Of upwards of 120 cases of this kind, observed by Dr. Coates at the Philadelphia Children's Asylum, in a period of three months, all but one recovered (*loc. cit.*, p. 21). The cases which occur in the course of other diseases are not dangerous in themselves, but are so as being the sign of a great severity of the primary affection.

TREATMENT.—The treatment may be divided into *general*, and *local* or *topical*. The *general treatment* should consist in most of the cases in at-

tention to the diet, which ought, in healthy and vigorous children, to be simple and unirritating, and in those who are weak and debilitated, nutritious and digestible. If the bowels are costive, or the child feverish and uncomfortable, a laxative dose may be given with advantage; or some simple diaphoretic, as nitre and water, or the neutral mixture, may be used through the day, and a warm foot-bath or an immersion-bath given in the evening. When the constitution is feeble, and the child weak or anæmic, tonic remedies are indicated. The best is probably quinine, or one of the ferruginous preparations; or the compound infusion of gentian, with addition of Huxham's tincture of bark, may be resorted to. The best internal remedy, however, and indeed the only one of any kind that is necessary in most cases, is the chlorate of potash, which possesses a stimulant and alterative action upon the mucous membrane. This is spoken of in the highest terms by Dr. West, of London, who regards it almost as a specific. We have used it now for many years past in a very large number of cases, and have seldom found it necessary to employ any other means, excepting some mild cathartic dose where the bowels have been constipated, and a wash of borax or alum in honey of roses, or borax in simple honey. The symptoms have begun to amend in every case in from three to four or five days, and recovery has taken place in about a week or a little more. The dose is from two to three grains every four hours for a child three years of age, and four and five grains for one of nine or ten years. Mr. Hutchinson (*Med. Times and Gaz.*, 1856), who believes also that this salt is almost a specific in this affection, recommends it in larger doses than the above, giving as much as five grains, thrice daily, to an infant of one year old. We have usually prescribed it in the dose of two grains four times a day, in a mixture of syrup of ginger and water, for children three or four years old.

Much discussion has taken place of late in regard to the injurious results of large doses of this salt administered to children, but we have certainly never seen any bad effects from its use, continued for from a week to ten days, in the amounts above recommended.

Prior to the discovery of the efficacy of the chlorate of potash in this affection, the *local treatment* constituted the only effectual and reliable means of removing it, and the most violent and painful applications were thought necessary and were made use of. Strong solutions of nitrate of silver, and pure or diluted muriatic acid, were frequently employed in severe cases. Now, however, these caustic substances may probably be entirely dispensed with, except in cases that show a tendency to assume the form of gangrene of the mouth. In ordinary cases the only local applications that need be used, and these are not essential when the child resists very much, are demulcent washes to keep the mouth clean, to be employed in the manner recommended in the article on aphthæ, and some mild astringent wash. This may consist of borax and honey, or borax and sugar, in the proportion of two or three parts of the former to one of the latter, or what is in our opinion preferable to either of these, of a drachm of borax rubbed up with an ounce of honey of roses.

Should the disease resist the treatment by the chlorate of potash and

the simple washes just now recommended, we may employ with advantage the ethereal solution of iodoform, as recommended for aphthæ. In cases which prove obstinate, a solution of sulphate of copper of from 3 to 10 grains to the ounce may be used.

MM. Rilliet and Barthez recommended very highly the plan pursued by M. Boneau at the Children's Hospital. This is to cleanse the mouth first, and then to apply dry chloride of lime (*calx chlorinata* of the *Pharmacopœia*) to the diseased surfaces. The application is made by means of a piece of rolled paper, or a stiff pencil, which is to be moistened and then dipped into the powder so that some may adhere, or with the finger. The surfaces are to be gently rubbed with the powder, and after a few moments' contact, washed clean with pure water. This is to be done twice a day, until the ulcerations assume a clean, healthy appearance, after which the following mouth-wash is to be employed :

R. Mucil. G. Acac.,	f℥j.
Syr. Cort. Aurant.,	f℥ss.
Calc. Chlorinat.,	℥j.—M.

The chief danger from the disease depends on the circumstance that it sometimes terminates in gangrene of the mouth, to be presently described. Any disposition to such a termination should be carefully watched, and the proper preventive means, consisting of local stimulating or caustic applications, with the internal use of stimulants and tonics, be at once resorted to.

ARTICLE V.

GANGRENE OF THE MOUTH.

DEFINITION ; SYNONYMS ; FREQUENCY.—Gangrene of the mouth is an affection which occurs chiefly in children of debilitated constitution, and especially as a sequel of some of the eruptive fevers. It begins generally by ulceration of the mucous membrane of the cheek, which after a longer or shorter time, runs into gangrene, and extends rapidly to the gums; after a few days, if the disease be not arrested, the central tissues of the cheek become thickened and indurated, an eschar forms upon the integument, and spreads in depth and width, until at last the cheek may be perforated, the whole side of the face and jaws destroyed, the teeth loosened, and the maxillary bones exposed and necrosed. It is known by a great variety of names: *gangrænopsis*, *cancrum oris*, *gangræna oris*, *canker of the mouth*, *gangrenous erosion of the cheeks of Underwood*; *necrosis infantilis*, *gangrenous stomatitis*, etc. It is a *frequent* disease in the hospitals for children in Europe, and a not uncommon one in institutions of the same kind in this country. It sometimes prevails endemically in hospitals. It is a rare disease in private practice, and we have as yet met with but few cases, excepting in public institutions.

PREDISPOSING CAUSES.—The disease is nearly, but not exclusively confined to the period of childhood. It is most common between the ages of three and six years; is very rare, but does sometimes occur in infants; and is of nearly equal frequency, probably, in the two sexes. Unfavorable *hygienic conditions* constitute a strong predisposing cause. Children living in hospitals or any crowded institution; those whose parents are poor or in want, and whose constitutions have been greatly deteriorated by long illness, by the tubercular diathesis, or by acute diseases, are particularly apt to be attacked. It almost always follows upon some previous acute or chronic disease, particularly measles, or some other acute exanthem; pneumonia; entero-colitis; whooping-cough; long-continued malarial fever, etc. MM. Guersant and Blache say (*Dic. de Med.*, t. 28, p. 601), “The existence of some anterior disease is a necessary condition of gangrene of the mouth; we have never known it, nor has M. Baron, to occur as an idiopathic affection.” It has been affirmed by some persons to be *contagious*, but this is exceedingly doubtful. The fact of its occurring sometimes in an epidemic form has already been referred to. It has been known also to prevail as an epidemic.

The *exciting causes* can rarely be ascertained with any certainty. The only one which seems to have been proved to exist in some instances is the exhibition of large doses of the mercurial preparations, and even this is questioned by some very good authorities.

ANATOMICAL LESIONS.—Upon examination after death, it is found that the *integument* surrounding the mortified spot soon runs into putrefaction. The lip or cheek in which the disease is seated is swelled, hardened, tense, and shining, of a purple or greenish color, and presents a deep, circumscribed engorgement. On the most prominent part of the swelling there often exists a rounded or oval, and distinctly limited eschar, of variable size, from a third of an inch to an inch, or even more, in diameter. In some instances the cutaneous slough is much larger, and extends irregularly to different parts of the face, to the chin, neck, eyelids, and even to the neighborhood of the ear, so as to occupy the whole of one side. Under these circumstances the tumefaction is neither so considerable, nor so regular, as when the slough is smaller. The eschar is always black, and generally dry and parchment-like, and extends a third or two-thirds of a line in depth, or quite through the integument. The tissues beneath the skin are not generally implicated, though in some cases the eschar is detached and there is a perforation of the cheek through which may be seen the alveolar processes.

The *mucous membrane* of the mouth is always affected with mortification. The disease may be limited, so as to exist in the form of an elongated ulceration, of a dark grayish color, situated in the fold where the mucous membrane is reflected from the cheek to the lower jaw; or, in a larger proportion of cases, it is seated on the internal surface of the cheek, opposite the interval between the alveolar processes. Sometimes the disease is much more extensive, and occupies all or a part of the internal surface of the cheek. In such instances the whole thickness of the mucous tissue is destroyed, and it presents upon its surface a blackish or brownish pulta-

ceous slough, almost liquid in consistence, which may be scraped off with a scalpel, leaving beneath loose shreds of mucous membrane, without any trace of organization. The gums frequently participate in the disease, and are converted into shreds, or completely destroyed.

The *maxillary bones* are sometimes, in severe cases, when the disease has extended to the gums, exposed, blackened, and even necrosed. The *teeth* are very often uncovered and loosened, and not unfrequently some are lost. The *tissues* between the skin and mucous membrane are found either hardened and infiltrated, or sphacelated to a greater or less extent. In the least severe cases, the fatty cellular tissue, and the muscular structures of the cheek are infiltrated with serum, but preserve their organization. When the disease is more aggravated, the gangrene extends to these tissues also, and always to those adjoining the mucous membrane first; so that the cellular structure beneath that membrane, and then the muscles, are infiltrated with a sanious fluid, and either in a state of sphacelus or tending thereto, whilst some of the adipose tissue beneath the skin is still merely infiltrated. In yet worse cases, the sloughs formed on the two surfaces of the cheek come into contact, and if their separation from the sound parts has taken place, a perforation is the consequence.

The condition of the *bloodvessels* in the midst of the diseased parts has been carefully examined by MM. Rilliet and Barthez. These authors state that when the tissues of the cheek are merely infiltrated, the vessels remain healthy, permeable, and their parietes are scarcely or very slightly thickened. When the vessels run along the edge of the slough, they are still permeable, but their walls are thickened, and begin to assume the appearances of the mortified tissues. Lastly, when they traverse the centre of the eschar, they can still be traced out, but their canals are found obliterated by coagula, in the whole extent of the mortified parts; or else the coagula occupy the vessels at their points of entrance into and exit from the slough, while between these points their walls are thickened, tend to assume the color and softness of the putrefied tissues, and their canals are filled with pultaceous gangrenous matter. The writers quoted do not suppose that the obliteration of the vessels is the cause of the sphacelus, since that change occurs only after the death of the surrounding tissues has already taken place.

The disease very rarely occurs on both sides of the mouth at once, though this does occasionally happen.

The *submaxillary glands* are nearly always in their natural condition, but in rare instances are softened and engorged.

Gangrene of the mouth never, or very rarely, indeed, exists without lesions of other organs. Of these the most frequent are acute pulmonary affections, and after them, acute or chronic diseases of the gastro-intestinal tract, and then malarial fevers, pleurisy, pneumothorax, peritonitis, and nephritis.

SYMPTOMS ; COURSE ; DURATION.—The following account of the symptoms of the disease is taken chiefly from the work of MM. Rilliet and Barthez. Gangrene of the mouth generally begins during the course or convalescence of some acute or chronic disease, by ulceration, aphthæ, or

phlyctenæ of the mucous membrane, and, in rare instances, by œdema of the substance of the cheek. At the same time the face is pale, and usually continues so throughout the disease; the nostrils and eyelids are often incrustated, and the latter infiltrated or sunken, and surrounded by bluish circles; the lips are swelled and covered with scabs, or dry. The breath of the child is fetid from the beginning, and, as the disease progresses, becomes gangrenous. There is but little fever at first, unless the case be accompanied by some acute disease; the pulse is commonly frequent and small in the beginning, rising gradually from 80 or 90 to 100 or 120, and becoming insensible towards the end. In cases occurring in the course of other diseases, the pulse rises sometimes to 120 or 140, and is larger and fuller. The child is generally languid and quiet at first, or more rarely cross and peevish. The strength may be either lost entirely, merely diminished, or the patient may retain a sufficient amount of force to sit up and observe what is going on around, and even to leave the bed the day before death. Half the children observed by MM. Rilliet and Barthez, in whom this symptom was noted, sat up in bed until within a few days of the fatal termination. In most cases but little complaint is made of pain in the mouth, though in some it is said to be severe.

The ulceration already spoken of as forming the first symptom of the disease is generally of a grayish color, and resembles very closely that which exists in the ulcero-membranous form of stomatitis. It may be seated either on the gums, in the fold formed by the junction of the cheek or lip with the gum, or on the inside of the cheek, opposite the space between the alveolar processes. It may present a gangrenous appearance from the first day, or not until after two or three days; or lastly, it may pass through the stages characteristic of ulcerative stomatitis, and terminate in the affection under consideration. Dr. B. H. Coates (*loc. cit.*) describes, under the title of gangrenous sore mouth of children, the ulcero-membranous form of stomatitis, and a few cases of gangrene, and states that three or four children out of 120 affected with ulcerated gums "suffered small spots of mortification, and one by the delay arising from the tardy report of a nurse, suffered necrosis in a portion of an alveolus."

The ulcerations just described assume the following appearances as the gangrenous nature of the malady develops itself. They become grayish, and then dark in color, bleed easily when touched, and are covered with pultaceous sloughs, exhaling a characteristic fetid odor. The gangrene extends to the neighboring parts, from the gum to the cheek, or from the cheek to the gum, and implicates at last the whole side of the mouth, or of the lower lip. At the same time the affected cheek or lip undergoes a circumscribed infiltration, which is at first rather soft, but becomes afterwards firmer, and forms at last a hard and rounded knot or tumor in the centre of the cheek, which is now tense, shining as though smeared with oil, and pale, or marbled with purple spots, while the slough on the inside is of a brownish color, more extended in size, and sometimes surrounded by a dark ring. The hard tumor of the cheek just described usually appears between the first and third days after the sphacelation of the mucous mem-

brane, though in some instances not until a later period. It is formed, as stated in the account of the anatomical lesions, by engorgement of the cellular and adipose tissues. The child, at this stage, is still able to sit up in bed and take notice, or shows evident signs of weakness and depression; the face is swelled and destitute of expression on the affected side; a bloody or dark-colored saliva runs from the mouth, which is partially open; the appetite is not entirely lost in all cases, the patient still demanding and taking food; vomiting is rare, but diarrhœa is almost always present; the thirst is generally intense; the skin is warm and feverish, natural, or too cool, and almost always dry, the differences depending probably more upon the concomitant disease than upon the mouth affection. The respiration is natural or altered according to the nature of the primary disease, which is, as already stated, in a large proportion of the cases, a pulmonary affection. The intelligence is generally undisturbed, though in some rare cases there is insomnia, delirium, or piercing cries.

If the disease continues to progress, as it almost always does when it has reached the stage we are describing, there appears in many, but not all the cases (8 of the 21 observed by MM. Rilliet and Barthez), a slough or eschar upon the most prominent and discolored part of the swelling of the integument of the cheek or lower lip. This generally makes its appearance between the third and sixth days of the disease, but in other cases, as early as the second, or not before the twelfth, or even later. The skin, at the point where the eschar is about to form, becomes purple, and then black; sometimes a phlyctena makes its appearance, which is very soon converted into a small, dry, black slough. This, if not limited by a process of separation from the living tissues, becomes larger and larger by the extension of the sphacelation, until it may, as already stated, embrace the whole side of the face. In grave and fatal cases, the gangrene sometimes extends to all the tissues of the cheek, and meeting at last, the disease which had commenced on the inside of the mouth, occasions a perforation, through which may be seen the teeth, alveolar processes, and the whole interior of the buccal cavity. In such instances as these, several of which we have seen in the Pennsylvania and Philadelphia Hospitals, the appearance presented by the child is, as may well be imagined, of the most pitiable kind. Even under these circumstances, however, with the cheek perforated, the edges of the opening irregular and covered with shreds of dead tissue, the gums destroyed, the teeth loosened, and the maxillary bones exposed, blackened, and perhaps necrosed, with a dark and fetid sanies flowing from the mouth or perforation, and a putrefactive smell infecting the air around, the child may retain, in some instances, its strength, so as to sit up in bed, ask for food, and drink with avidity. In other cases, on the contrary, the patient at this stage is exhausted to the last degree, and refuses both food and drink. During the closing stage of the disease there is generally profuse diarrhœa, rapid emaciation, dry skin, small, rapid pulse, and at last death in a state of utter prostration.

In favorable cases the recovery may take place in the early stage, before the integument becomes involved, and while the gangrene is limited to the

mucous membrane, or at a later period, after the slough has separated. In the first instance the child generally recovers without deformity, though we saw one case in which necrosis of about an inch of the front of the inferior maxilla took place without any loss of the soft parts. When the child recovers after the formation of the cutaneous slough, a very rare event, the gangrene ceases to extend, the eschar separates and is cast off, the edges of the opening assume the appearances of a healthy ulcer, and after a length of time approach each other and cicatrize, leaving generally a large, uneven, discolored scar, like that of a burn, which remains through life a horrid deformity.

The *duration* of the disease varies according to its termination. When this is unfavorable, which happens in much the larger proportion of cases, death usually occurs about the end of the first, or in the course of the second week, though it has been known to occur at a later period. In favorable cases the duration is commonly longer, particularly if a cutaneous eschar has been produced, as the separation of the slough and cicatrization of the ulcer which remain require a tedious and slow process on the part of nature.

Complications are very apt to arise in the course of the disease. The most frequent is pneumonia. MM. Guersant and Blache state that it exists in nine-tenths of the cases; MM. Rilliet and Barthez found it in 19 out of 21; of the 19, it began in 8 during the progress of the gangrene, and apparently under the influence of the latter, whilst in the remaining cases it existed before, and acted perhaps as a predisposing cause to the affection of the mouth. Another and more dangerous complication is the occurrence of gangrene in other parts of the body, particularly the soft palate, pharynx, œsophagus, anus, and more frequently the vulva and lungs.

DIAGNOSIS.—Some authors have described as identical affections, under the title of gangrenous stomatitis, the disease under consideration and the one already treated of as ulcero-membranous stomatitis. This has been done particularly by M. Taupin, who is followed in his description by M. Valleix (*Guide du Méd. Prat.*, t. iv.) It seems clear to us, moreover, that Dr. B. H. Coates, in his very valuable paper on the "gangrenous sore mouth of children" (*loc. cit.*) mingles in his description the two diseases referred to. It seems clear, however, that the differences between them as to frequency, symptoms, course, amenability to treatment, and termination, which are fully pointed out in the diagnostic table below, fully warrant us in regarding them as different and distinct diseases.

The diagnosis of gangrene of the mouth is, in most cases, very easy. The ulceration of the mucous membrane, followed by gangrene; the deep-seated induration of the cheek, at first pale on the outside, then dark-colored, and terminating after a time in a characteristic slough; the course of the malady, and the nature of the general symptoms, will generally prevent any difficulty in the recognition of the disease.

From stomatitis it may be distinguished by attention to the points laid down in the following table taken from MM. Rilliet and Barthez:

STOMATITIS.

Begins by ulceration or by pseudo-membranous plastic deposit.

Odor very fetid and sometimes gangrenous.

But little extension of the local lesion, which always retains the same appearances.

But little swelling of the cheek or lips, or simply œdema of those parts, without deep-seated induration, tension, or unctuous appearance.

Salivation rarely so considerable as to flow from the mouth; when present, sometimes sanguinolent; never mixed with shreds of gangrenous tissue.

Never an eschar on the exterior.

Never complete perforation of the soft parts; denudation of the bones never occurs; loss of the teeth very rare.

Course of the disease slow when left to itself; recovery rapid under the influence of treatment.

GANGRENE.

Begins by ulceration, which is sometimes gangrenous from the first, or by œdema of the cheek.

Odor always gangrenous.

Considerable and rapid extension; the tissues assume a peculiar dark grayish tint.

Extensive swelling and œdema of the cheek, with deep-seated induration, tension, unctuous appearance, purple spots.

Salivation abundant; constant escape of fluid, at first sanguinolent, afterwards putrefactive.

Often an eschar upon the cheek or lips.

Perforation of the soft parts frequent; denudation of the bones constant; loosening of the teeth constant, and their loss frequent.

Course rapid, and termination fatal, as a rule, when the disease is left to itself, and in spite of all treatment.

Gangrene of the mouth may be confounded with malignant pustule. The method of diagnosis has been drawn by M. Baron in the following words; "Malignant pustule always begins on the exterior; affects the epidermis first, and extends successively to the corpus mucosum, chorion, and subjacent parts; whilst on the contrary, the gangrene under consideration attacks the mucous membrane first, then the muscles, and lastly the skin."

PROGNOSIS.—The prognosis of true gangrene of the mouth is exceedingly unfavorable. The great majority of the subjects die in spite of all that can be done. Dr. Coates (*loc. cit.*, p. 14) says that a black spot on the outer surface of the swelling "has always been in my own experience the immediate harbinger of death. It is proper to state, however, that I have heard it said that cases had recovered in this city, in which the gangrene had produced a hole through the cheek." MM. Rilliet and Barthez state that "death is the ordinary termination of gangrene of the mouth, though there are instances of recovery on record." Of 29 cases analyzed by them, only 3 recovered. MM. Guersant and Blache (*loc. cit.*, p. 596) state that unless arrested in the formative stage, it ends fatally almost constantly in from five to ten days, and frequently before perforation has taken place. Of 36 cases observed by M. Taupin in the Children's Hospital at Paris, not one escaped (Guersant and Blache, *loc. cit.*, p. 597). The authors of the *Compendium de Médecine Pratique* say of this disease (t. i., p. 632), "Death is the almost inevitable termination." Dr. Marshall Hall (*Edin. Med. and Surg. Journ.*, xiv., p. 547) reports six cases of the disease, two of which followed measles, one repeated attacks of pneumonia, one fever, (type not mentioned), one worm fever, and one typhus fever. All but one,

the case occurring in the course of typhus fever, in a girl, twelve years old, died. This girl recovered, with, however, falling in of the right cheek, "a frightful chasm" on the left side of the mouth, and caries of a portion of the alveolar process, palate-bone, and second molar tooth. Recoveries sometimes occur, however, as in the case mentioned by Dr. Hall, after perforation, but nearly always with terrible deformities, with adhesions of the walls of the mouth to the jaws, with incurable fistulæ, etc.

The prognosis is more favorable in private practice than in hospitals. The favorable circumstances in any case are: good hygienic conditions; vigorous constitution of the child; the absence of dangerous concomitant disease; the continuance of appetite and strength; and a disposition to limitation and separation of the slough. Unfavorable symptoms are: weak and debilitated constitution of the patient; severe coëxistent disease; prostration of the strength; and extension of the sloughing process. Death may also occur from hemorrhage in consequence of the separation of the slough, as in a case quoted from Hueter by Bouchut.

TREATMENT.—The reader need but refer to the remarks on prognosis to be assured that no treatment as yet discovered promises much success. We would call attention also to the following statement: that the remarks about to be made apply only to true gangrene of the mouth, and not to all the cases described by some writers under the title of gangrenous sore mouth or even that of gangrene of the mouth, since, as already stated, they confound together true gangrene and ulcero-membranous stomatitis.

The treatment is divided into *local* and *general*. The *local treatment* recommended by the French writers, consists in cauterization of the sloughing parts with one of the mineral acids, with nitrate of silver, or with the actual cautery. This is the plan proposed by MM. Billard, Baron, Guersant and Blache, Barrier, Rilliet and Barthez, Bouchut, and Valleix. The authors of the *Bibliothèque de Médecin Practicien* remark, however, that nearly all the patients subjected to cauterization die, and that of the small number saved, there are as many who had not been subjected to that treatment, as there are of those to whom it had been fully applied. They wonder, therefore, that recent authors continue to repose the same confidence in it as did their predecessors. "For us," they say, "we are of opinion that cauterization exerts but slight influence, if it have any at all, and it is greatly to be desired that the zeal of practitioners might discover some more efficaceous remedy" (*loc. cit.*, t. v, p. 551).

It is very important to make use of the caustic application as early after the beginning of the sphacelus as possible, for if it be allowed to spread to any considerable depth or extent, there is scarcely a hope of arresting it by any means. MM. Guersant and Blache recommend pure nitric, sulphuric, or muriatic acid; MM. Rilliet and Barthez propose the acid nitrate of mercury, or muriatic, sulphuric, or acetic acid; M. Valleix proposes the treatment employed by M. Taupin, which is to remove the pseudo-membrane and a part or the whole of the gangrenous eschar with scissors, to make some scarifications upon the healthy parts, to apply pure muriatic acid, and after the separation of the slough, to make use of dry chloride of lime (*calx chlorinata*). The acid most generally employed is the muriatic.

The local treatment proposed by MM. Rilliet and Barthez is the following: As soon as the ulcerations assume a gangrenous appearance, to touch them with a brush or sponge dipped into acid nitrate of mercury, or pure muriatic acid, the brush to remain in contact with the sloughs for a few instants, and then to be applied rapidly around and on the parts beyond them. After this cauterization, an application is to be made of dry chloride of lime (in the manner recommended in the article on ulceromembranous stomatitis), which is to be left in contact with the sloughs for a few minutes, when the mouth must be thoroughly washed with a strong jet of water from a syringe. The cauterization and use of the chloride of lime are to be resorted to twice a day, and the mouth washed three or four times in the interval with large injections of simple water, barley-water mixed with honey of roses, or better still, with a strong decoction of cinchona. If the case goes on favorably, and the sloughs separate, the cauterizations are to be suspended, and the chloride of lime alone employed. If, on the contrary, a slough forms on the outside of the cheek, a crucial incision must be made into it, and a brush charged with the same caustics introduced between the cuts; powdered cinchona is then placed in the openings, and retained there by a piece of diachylon plaster or by pledgets of charpie, dipped in a solution of soda. This treatment is to be continued until the slough separates, when the edge of the wound, and all the diseased parts that can be reached, are to be cauterized.

In applying escharotics to the mouth, certain general precautions are required, of which it is necessary to give some account. When they are used upon the inside of the cheek, a spoon must be introduced into the mouth, with the concavity directed towards the alveolar processes, in order to preserve the teeth and tongue from being touched. When the application is made upon the gums, the cheek should be drawn to one side by an assistant, and the tongue pushed out of the way with the finger, or a spoon. If the acid happen to touch the teeth or tongue, it must be instantly washed off. The mouth ought always to be thoroughly cleansed with water immediately after the cauterization, to remove any superabundance of acid.

The kind of brush most suitable for the application of the mineral acids is one made of charpie, strongly tied to a solid handle. The sponge-mop which is sometimes used, is made by fastening a small piece of fine sponge to the end of a stick.

MM. Guersant and Blache recommend that the acid be applied to the slough every hour, until the sphacelus ceases to extend. They state that this plan is sometimes advantageous when the gangrene is confined to the gums only, but that it is generally powerless when the disease has extended to the cheek, or has implicated the deep-seated tissues. Under the latter circumstances, and when the inefficacy of caustics has been shown by trial, they propose the use of the actual cautery, as recommended by M. Baron, and other distinguished practitioners, and which, they add, has afforded them some brilliant results in very bad cases.

M. Barrier advises that we should accurately expose the diseased parts by crucial incisions, and apply the escharotic to all the parts forming the

limits of the gangrene, in such a way that the tissues already disposed to slough shall be thoroughly cauterized, while those a little beyond are so in a less degree.

In applying these powerful caustics, several authorities recommend the administration of an anæsthetic.

The English writers, and those of our own country, seem rather less disposed than the French to make use of powerful escharotics, and lay more stress upon the general treatment. Underwood, following Dease, of Dublin, advises that "the parts should be washed and likewise injected with muriatic acid, in chamomile or sage tea, and afterwards dressed with the acid-mixed with the honey of roses, and over all a carrot poultice." Dr. Symonds (*Lib. of Pract. Med.*, vol. iii, p. 23) directs the cheek to be frequently rubbed with a stimulating embrocation of camphorated oil and ammonia, on the first appearance of the swelling, and in the intervals to be kept moist with a tepid lotion containing muriate of ammonia and alcohol. On the slightest appearance of an eschar upon the interior of the mouth, it is to be touched with solid nitrate of silver, or strong muriatic acid. If sloughing have already commenced, the nitrate of silver lotion is said to be the best application. The mouth is to be frequently washed or syringed with a solution of chloride of soda, and when mortification has taken place, we are to endeavor to prevent it from spreading, by carrot or fermenting poultices. Maunsel and Evanson say that the early application of muriatic acid, undiluted or mixed with one or two parts of honey, is the only efficient application in these forms of gangrene. Dr. Fleming (*Dublin Hosp. Gaz.*, May 1st, 1855) recommends the application of a concentrated solution of nitrate of copper, to the sloughing surfaces, and also paints the circumference of the disease and the surrounding cheek with collodion, which, he believes, acts favorably upon the capillary circulation of the part. Dr. Gerhard (*Lib. of Pract. Med.*, vol. iii, Am. ed., p. 24) says "The best local applications are the nitrate of silver, if the slough be small in extent; if much larger, the best escharotic is the muriated tincture of iron, applied in the undiluted state; after the progress of the disease is arrested, the ulcer will improve rapidly under an astringent stimulant, such as the tincture of myrrh, or the aromatic wine of the French Pharmacopœia." Dr. Dunglison (*Prac. of Med.*, vol. i, p. 36) recommends the application with a brush, of a mixture of equal parts of creasote and alcohol, after incisions have been made through the gangrenous parts. Dr. Condie (*op. cit.*, 6th ed., p. 174) states that he has found a strong solution of sulphate of copper (thirty grains to the ounce of water), applied very carefully twice a day, to the full extent of the gangrenous ulceration, by far the most successful lotion.

We have, ourselves, lately employed carbolic acid in two severe cases. The pure acid was carefully applied to the sloughing ulcer on the inside of the cheek, and subsequently a solution of one part of the acid in fifty of water, was frequently employed to wash out the mouth. The application of the undiluted acid seemed to have a beneficial effect, by checking the progress of the sloughing, and completely destroying the putridity of the dead tissue which had not as yet separated. One of the cases recovered

quickly, without perforation of the cheek ; but in the other death occurred, with symptoms of profound adynamia, though there was little, if any, extension of the gangrene.

It seems to us very clear, after the study of the treatment recommended by the different writers quoted above, that the most important part of the local management of the disease is the early application of some escharotic substance to the ulcerations, or to the mortifying parts ; the best is probably pure muriatic acid. This should be made use of twice or three times a day, observing the precaution to wash the mouth with water, immediately afterwards, by means of a syringe. Later in the disease, when it has extended to the skin, the use of escharotics, or of the actual cautery, is still recommended by many writers, but opposed by others. We confess we should be inclined to prefer, at this stage, the use of muriated tincture of iron, as recommended by Dr. Gerhard, of carbolic acid as used by ourselves, of strong lotions of sulphate of copper, of solutions of nitrate of silver of moderate strength, or of the dressings of muriatic acid and honey of roses, as proposed by Underwood, in connection with carrot and fermenting poultices, as recommended by Underwood and Symonds. Throughout the course of the disease the mouth ought to be frequently cleansed by washing or injecting with solution of chlorinated soda, mixed with eight parts of water, or with a dilute solution of carbolic acid, which corrects at the same time the terrible fetor of the disease.

The importance of these measures can scarcely be over-estimated, since the presence of gangrenous tissue about the oval cavity must lead to the introduction of the poisonous results of putrefaction into the system, both by the fetid discharges which partly flow down the œsophagus, and still more by the contamination of the inspired air. Indeed, it seems quite possible, as urged by Dr. Keiller (*Edin. Med. Jour.*, April, 1862), that in cases of unchecked gangrene of the mouth, death occurs, in a great measure, from secondary blood-poisoning, resulting from the continued and unavoidable inhalation of air poisoned by emanations from the gangrenous sloughs. It is evident, therefore, that local applications, both of caustics and antiseptic lotions, must be of great service, by arresting the sloughing and correcting or checking the foul discharges.

GENERAL TREATMENT.—All writers recommend the use of tonics, stimulants, and nutritious diet, unless the presence of high fever, or the state of the digestive organs, seem to contraindicate their employment. From our own personal experience in the treatment of this affection ; from a consideration of what we have seen most successful in other forms of gangrene, as that following accidents and surgical operations in deteriorated constitutions ; from what proved effectual in the case of idiopathic gangrene of the vulva, in a child ten years of age, which came under our charge ; and from what is necessary in that analogous condition of the constitution which accompanies typhoid and cachectic diseases, we are induced to believe that the general treatment must be of at least as great importance as the local, and that the steady and persevering use of tonics, stimulants, and of the most strengthening diet, should always be insisted on from the earliest period, whether fever be present or not. The quantity of stimu-

lants, and the amount of food, ought, it seems to us, to be measured only by the capacity of the digestive organs to receive and assimilate them. Of tonics, the best are quinine and muriated tincture of iron, which may be given in syrup, in doses of a grain of the former with three drops of the latter, four or five time a day, to a child three or four years old. The most suitable stimulants are good brandy or wine given in considerable quantities, and, if the stomach receive it well, carbonate of ammonia, or better still, the aromatic spirit of harts-horn. The diet must consist of milk made into punch with brandy, wine-whey, the yolk of eggs beaten up with wine, strong soups and beef tea, animal jellies, and, if the child wish it, tender meat finely minced.

The room in which the child is placed ought to be large, if possible, and to all events thoroughly ventilated.

ARTICLE VI.

THRUSH.

DEFINITION; SYNONYMS; FREQUENCY; FORMS.—The term *thrush* is applied to a disease long supposed to be a purely local affection, characterized by the deposit on the mucous membrane of the mouth of a whitish or grayish-yellow exudation, of a cheesy consistence, through which ran a parasitic fungus, called by Robin, *oidium albicans*; but of late years it is known that the mouth affection is merely a localization of a widespread constitutional disorder of a special and serious nature. It is the disease described under the title of aphtha or thrush, by Underwood and Eberle; of aphtha, by Dewees; of erythematic stomatitis, with curd-like exudation, by Dr. Condie; and of aphtha lactantium, aphtha lactamen, and aphtha infantilis, by the older writers. It is the muguet of the French.

The *frequency* of the disease is very great in hospitals for young children, especially in foundling hospitals, and in the wards of almshouses devoted to foundlings. It occurs amongst the children of the poor and illiterate, and is very rare, according to our experience, in the middle and upper classes.

It occurs in two forms, the *mild* and the *grave*. In the first, which is met with occasionally in the easy classes of society, the constitutional disturbance, without which the disease probably never exists, may be so slight as to require an experienced eye to detect it. In the second form, the disease is common and very fatal in foundling hospitals, not rare in the neglected, half-starved children of the lowest classes of society, and is occasionally, but very rarely, according to our experience, met with in the easy classes of society.

CAUSES.—The central cause of thrush lies, we think, in a condition of health in which the general vitality is slowly ebbing away under what amounts, virtually, to an inanition, innutrition, denutrition, or what M.

Parrot, of the old Foundling Hospital of Paris (*Clinique des Nouveau-Nés. L'Athrepsie*, par J. Parrot, Professeur a la Faculté de Médecine de Paris, Médecin de l'Hospice des Enfants-Assistés, Paris, 1872), calls athrepsia. M. Parrot derives this new word from *a priv.*, and *θρεψις*, nutrition. He insists, however, that the disorder is not one of inanition, but rather some fault in the evolution or developmental process. The disease occurs in adults, or at least we believe it to be the same, at the close of phthisis and chronic catarrhal pneumonia, in prolonged cases of cancerous disease, and in fatal chronic diarrhœa. It occurs in children deprived of their natural aliment, and virtually starving on some artificial food, often badly chosen, badly prepared, and carelessly or imperfectly administered, as most happens in large foundling hospitals, in the nursery wards of almshouses, where one nurse has charge of three or four or six children; whereas, in families of easy fortune, and often in those of the industrious poor, one hand-fed baby absorbs almost all the time of one woman, and when such a child falls ill, we know that it can be better taken care of by the hands of two than of one woman. It is rarely met with even in the mild, and in the grave form is almost unknown, in suckled children. We have seen, in all our experience, but one case of grave thrush in a child suckled by its own mother, and that was many years since. We shall refer to it again. M. Valleix, whose researches on this subject are amongst the most valuable we have, and whose knowledge was gained in that great school of experience, the old Foundling Hospital of Paris, now the Hospice des Enfants-Assistés, declares (*loc. cit.*, p. 60): "On the other hand, I have never known a child who had been suckled exclusively during the early months of life to have the disease." MM. Trousseau and Delpech, in a valuable paper on the disease (*Journ. de Méd. de MM. Beau et Trousseau*, Janv., Fev., Av., et Mai, 1845), say: "We should be justified, therefore, in asserting that we have never known an infant to die of thrush who had been suckled at a healthy breast, or whose health has not been dangerously compromised by other causes."

M. Parrot (*loc. cit.*, p. 37) refers to the unfavorable atmosphere of hospitals as one of the causes. He adverts to this nosocomial influence, "which," he says, "sometimes acts with a disastrous intensity. It is the agent of unknown nature, which develops in lying-in hospitals and in hospitals or homes for young infants (*crèches*), even when the wards are large and well ventilated." But we still believe that the main cause of the disease is to be looked for in an unhealthy constitutional condition, brought about by the absence of the natural and the substitution of some improper and unhealthy artificial food. M. Parrot himself says (p. 382): "Vicious *ingesta* are, in effect, the most frequent and powerful of all causes." To put this very important matter in the strongest light, we make the following additional quotations. Underwood says: "A principal remote cause of this disease seems to be indigestion, whether produced by bad milk or other unwholesome food, or by the weakness of the stomach." Dewees remarks, "Children fed much upon farinaceous substances are especially exposed to the attacks of this disease, particularly when their food is sweetened with brown sugar or molasses." Dr. Eberle says: "Unwhole-

some and indigestible nourishment, and over-distension of the stomach during the early stages of infancy, almost inevitably lead to the occurrence of aphtha (thrush). Bad and old milk, and thick farinaceous preparations, sweetened with brown sugar or molasses, almost inevitably lead to the occurrence of aphtha (thrush)."

When we come to speak of the nature of the disease, we shall dwell at some length upon a possible element in its causation which has suggested itself to us within a few years past, and which may strike the reader, at first view, as one of very doubtful probability. We refer to the absence of sufficient supply of water in much of the artificial food employed for very young children.

The disease occurs at all *ages*, but is by far most common in the first two months of life. M. Parrot, who has had the enormous experience of the great Parisian hospital, gives us no figures whatever, but says, in speaking of the influence of sex, that he has seen no proof of its having any, and then adds: "In my opinion, as regards the etiology, these are but the newborn, not boys or girls." *Deranged health* from any cause, *deficient ventilation*, invite the disease. The congenital feebleness of *premature* children renders them specially liable to it. *Season* exerts a considerable influence, as M. Valleix found that more than half the cases occurred during the three warmest months of the year.

CONTAGION.—The question of the contagiousness of thrush has been often discussed. We have seen so little of the severe forms of the disease, that our opinion is not worth much, but what we have seen has never aroused in us even a suspicion that it was contagious. M. Parrot cites a few cases (*loc. cit.*, p. 80) which seem to point to a possible infection of the nipple of the nurse, and from thence to the mouth of a second child. But his own opinion against its contagiousness is very positive. He says (p. 83): "I must declare that, at the hospital of the *Enfantes-Assistés*, where I have studied for several years, and where muguet (thrush) is truly endemic, I have never yet met with a well-established case of contagion. And this leads me to believe that it does not often occur." Other authorities, MM. Billard, Baron, Blache, Guersant, and Grisolle, several of whom saw the disease in the Paris hospital, deny its contagiousness. Another writer, M. Seux, says: "My personal experience has not convinced me that the breast of the nurse can be infected with thrush by the mouth of the child. I have watched perseveringly and regularly, the relations with their nurses, of more than 1600 infants affected with thrush, and I have never known the disease to develop on the breast of the women." But M. Seux believes that the child may be infected by the nurse, though he disbelieves in the growth of the fungus on the breast. He supposes that the nipple, without being itself diseased, may be the vehicle between an infected and a healthy child. (*L'Athrepsie*, foot-note, p. 80.)

ANATOMICAL LESIONS.—The characteristic deposit is found upon the mucous membrane of the mouth, pharynx, œsophagus, and, in rare cases, of the stomach and intestines. The question as to the extension of the deposit to the gastric mucous membrane has been much discussed, and the highest authorities have been almost equally divided upon it. This dis-

agreement has arisen solely from the want of microscopical examination, which enables the observer to distinguish readily between true thrush and other appearances of the gastric mucous membrane which closely resemble it. The most conclusive demonstration of its occurrence upon the mucous membrane of the stomach has been furnished in a valuable article on this subject, by M. J. Parrot (*Arch. de Physiologie Norm. et Path.*, Nos. 4 and 5, 1869), and in his work on athrepsia. He has also determined its occasional presence in the large intestine. But the very doubts expressed by observers prior to Parrot, show that it must be very rare. M. Parrot does not state how many times he has seen it in the stomach, but he does refer (*L'Athrepsie*, p. 230) to two cases in which he verified by the microscope the presence of the spores and tubes of this growth, in the large intestine.

It is a curious fact, and a very important one, insisted on by MM. Trousseau and Delpech, and other observers, that the false membrane never extends to the nasal or air-passages; and they call attention to the singular difference in this respect between the affection under consideration and diphtheritic inflammation, which attacks almost exclusively the nostrils, pharynx, larynx, and bronchi. M. Parrot, nevertheless, asserts that it extends quite frequently to the pharynx, and more rarely to the glottis. He mentions (*loc. cit.*, p. 235) eleven cases in which he found the growth on the glottis. It was confined entirely to the inferior vocal cords. He also declares that he has met with one case in which the growth had extended to a pulmonary infundibulum in the apex of the right lung.

Lesions of the digestive mucous membrane are met with in nearly all the cases. M. Valleix states that softening of the gastric mucous membrane is almost constant, and that it is often accompanied by redness and thickening. The authors cited above are of opinion that the gastric lesions have been greatly exaggerated, and assert them to be much the same as exist in other diseases foreign to the digestive apparatus. Various morbid alterations of the mucous membrane of the intestines exist, they state, in nearly all fatal cases. This fact is acknowledged as well by MM. Trousseau and Delpech, who deny the invariable connection of these alterations with thrush, as by M. Valleix, who asserts the connection almost without reserve. The best account of the lesion of the digestive apparatus is given by M. Parrot, in his work on athrepsia. He describes also the lesions of the nervous system and blood, and we shall proceed to quote some of his most important statements.

In the stomach he found gelatiniform *softening*, which he ascribes to the action of the gastric juice on the tissues. He found, also, *ulceration*, and a *diphtheroidal* process. The ulceration appeared in two forms. In one, small isolated spots of a sepia tint, or black in color, are seen scattered through a layer of grayish mucus covering the walls of the organs. These dark spots, on examination, are found to be either depressions in the walls, or true ulcers so small as to be scarcely distinguishable, up to a line in diameter. The second variety is met with in premature children affected with *œdema neonatorum*. On the internal wall of the stomach, whether covered with a layer of mucus or not, are seen small, lenticular spots, not more than half a line in diameter usually, and sometimes much smaller,

of a citron-yellow color, slightly depressed in the centre, or distinctly ulcerated. In this form the mucous membrane was more injected than in the first, and the ulceration was surrounded generally by a red line. He refers to a third form of ulceration, infinitely more rare, in which only one or two are found, but these much larger, deeper, and with projecting and congested edges.

The diphtheroidal lesion also occurs in two forms. In one it resembles closely the ordinary diphtheritic process, occurring in very small points, scarcely half a line in diameter, or in layers of from half an inch to an inch square. In the second the exudation is less compact, less adherent, and of a greenish color, resembling the false membrane of pericarditis.

In regard to the intestinal tract, he says: "We rarely meet with any lesion of the intestine in athrepsia. Indeed, and this is well worthy of attention, it may be stated that of all the organs concerned in the act of digestion, it is the one least frequently affected. This is surprising when we reflect how common diarrhœa is in the disease, and how general is the opinion amongst writers that the intestinal flux is the result of enteritis. In the immense majority of cases, the intestine retains its normal color and thickness; the mucous membrane is grayish in color, and looks as though it had been washed; in the colon, it is not unusual to meet with some of the closed follicles more projecting than natural." He adds that congestion of the different coats is common; that he has occasionally found the mucous membrane injected and thickened; and that, in two instances, he met with the lesions observed in the stomach.

In the liver, though he thinks the organ may play an important part in athrepsia, "it has been impossible for me," he says, "up to this time, to find any particular change." There is little fat in its cells, and, in the chronic forms of the disease, all the fatty matter has disappeared.

He describes an inflammation of the middle ear, asserting that it always exists in the new-born dying of athrepsia. This portion of the ear contains, at first, a thick and flocculent serosity; later, its lining membrane is injected, and the cavity fills, by degrees, with a greenish or greenish-yellow mucus. Still later, the mucous membrane, thickened and softened, incloses a mass of greenish, creamy, and sometimes ropy pus. The membrane of the ossicles is attacked as well as that of the cavity. The eustachian tube is always healthy. The membrana tympani, though softened and friable, is rarely perforated. These lesions, nearly always more marked on the right than left side, exist in some subjects only in the first stage. He believes that the condition may be recovered from, without leaving any notable disturbance in the sense of hearing.

In the *brain* he found fatty degeneration, softening, and hemorrhage. Fatty degeneration occurs in two forms, the diffuse, and the circumscribed or insular. It exists also in the form of small points, irregularly rounded, whitish, and more opaque in the centre than at the periphery, and rarely of a yellowish hue, on different parts of the arachnoid membrane, but especially so near the longitudinal fissure, and in the cerebellum, to the right and left of the median line. Hemorrhage in the brain is a common lesion. He met with it in 34 cases, and observed it in five different localities: in

the cavity of the arachnoid; in the subarachnoid or pia mater region; in the nerve-tissues proper; in the walls of the lateral ventricles, under the ependyma, and in the lateral ventricles. The chief cause of these hemorrhages appears to be the altered blood crasis, which determines congestion, and venous obstruction.

The changes in the *lungs* consist of fatty degeneration of the alveoli, emphysema, and the softening consecutive to thrombosis of the pulmonary artery.

The *kidneys* exhibit fatty degeneration of the tubules, venous thrombosis, and uratic infarctus. By the latter term is meant a condition in which the tubes of Bellini are choked with a deposit in the form of cylinders, fractured at various points. The opaque matter forming the cylinders never invades the interior of the cells. Examined by themselves, this matter consists of elongated masses, of irregular and bosselated contours, similar to stalactites, and formed of a number of spherules of unequal size, perfectly regular, and held together simply by juxtaposition. M. Parrot disagrees with Virchow, who regards these bodies as being composed of urate of ammonia, and declares them to consist of urate of soda.

Of all the anatomical changes, none are more constant and important than those of the *blood*. They make their appearance at the outset, become aggravated each day, and act a considerable part in causing the functional disturbances observed during life, and the various lesions of tissue found after death. "Blood," he says, "drawn in the acute disease from the hand or foot, during life, by a small incision or prick, is of the color of deep lees of wine, and often blackish. If a drop be received on a glass slide, far from diffusing itself rapidly, it preserves its globular form, which is an indication of concentration and viscosity." It is said that, in acute cases, the proportion of red globules is increased. In chronic forms, on the contrary, the blood is watery, of light color, spreads rapidly on a glass slide, and the microscope shows a loss of red globules, which is the more decided as the fatal termination approaches. He is not certain as to the number of the white corpuscles, but is of opinion that their number is increased as the disease progresses, both in rapid and chronic cases. After death in the acute form, the blood found in the heart and veins is very dark in color, almost syrupy, and occasionally formed into soft and friable coagulated masses. In chronic cases it presents nothing particular. One of the important properties of the blood is that of coagulating during life. The thromboses thus formed have their exclusive seat in the veins. They are found only in the sinuses of the dura mater, the veins of the brain and its membranes, in those of the kidneys, and in the pulmonary artery. It is quite unusual to find this condition elsewhere, but it is occasionally seen in the inferior vena cava, and, in such cases, has its point of departure in the emulgent veins.

In severe cases of thrush a certain amount of erythematous inflammation is often found upon the skin of the buttocks and thighs, and ulcerations sometimes exist upon the inner ankles. Before leaving this part of our subject, we may remark that, in the few cases we have met with in private

practice, no ulcerations existed upon the malleoli, and the erythema we observed was only in the neighborhood of the anus.

Symptoms.—We shall first describe the characters of the exudation, and then proceed to the consideration of certain general and local phenomena which exist to a greater or less extent in both forms of the disease.

The mucous membrane of the mouth is often somewhat red, dry, and tender for a longer or shorter time (generally from one to three days), before the appearance of the exudation, and at the same time the papillæ of the tongue swell and become protuberant. Next the exudation shows itself in the form of small, whitish points, sometimes on the tongue first, and in other cases on the inside of the lips, whence it extends to the cheeks in idiopathic mild cases, and to the roof of the mouth, soft palate, pharynx, and œsophagus, in the grave, symptomatic form. The points of false membrane first deposited rapidly increase in size and thickness, so that in from one to three or four days, they assume the form of large patches, or a continuous membrane, which covers the whole or a considerable portion of the cavity of the mouth. When the exudation is recent, it is thin, and its surface smooth; when, on the contrary, it has been longer deposited, it becomes thicker, and its surface is rough. It is at first of a milk-white or pearly hue, but when undisturbed assumes a grayish or yellowish color. It is soft in consistence, breaking down under the finger like cheese, and presenting no traces of organization to the naked eye. It adheres to the mucous membrane with considerable tenacity at first, but becomes looser after awhile, and is detached spontaneously at last without any lesion of the tissue beneath.

The foregoing description applies to the exudation as it appears to the unassisted eye. We shall next give an account of the characters it presents, when subjected to microscopical examination, and in so doing shall quote the language of Berg, who first discovered that thrush essentially depended upon the presence of a peculiar parasitic fungus, to which Robin has given the name of *oidium albicans*. Dr. Berg (*loc. cit.*) states, that the white coating of the exudation consists of epithelium, thickened by the swelling of its constituent cells; from the epithelium there springs a parasitic fungus in greater or less quantity, so that the chief portion of a patch of aphthæ (thrush) is composed either of epithelium or else of the parasitic growth. Under a magnifying power of from 200 to 300 diameters, an aphthous crust is seen to consist of epithelial cells, with a more or less interwoven coat of fibres, and a variable number of spherical or oval cells, without any sign of exudation corpuscles, but only a small quantity of molecular albuminous deposit. “We can often trace the successive development of these cells from a spherical one of the smallest size, to an oval cell, and thence to a filament; and we have no doubt ourselves that the smaller cells are sporules out of whose development the larger oval cells are formed, and finally, the filaments in the same manner as has been observed in other fungoid growths of this nature.” Numerous projecting fibrils are observed in the circumference of an aphthous crust when submitted to the microscope; but these are rendered infinitely more clear by a weak solution of potash, which dissolves the albumen,

and renders the cells of the epithelium transparent, while, at the same time, it diminishes their intimate cohesion, and the network of vegetable fibres is more plainly seen. "These fibres are cylindrical, with sharply defined dark edges, and their centres are transparent in transmitted light; they are generally equal in thickness, but at times they are, as it were, knotted together, and divided by distinct walls of separation In their interior, these fibrils often exhibit nucleated cells; occasionally these are very numerous, and of small size, but at times they are larger. In this course the fibrils divide into numerous branches, whose diameter is not less than that of the original stem, and I have occasionally observed these ramifications to increase in thickness, at their free extremity, and to terminate in a club-shaped end with a species of cell. From the sides of the fibrils spring numerous sporules, forming a point of departure for new ramifications Careful investigation has shown us that these cells are placed upon the sides of the fibrils, and in particular that they are congregated around the terminations of the latter. It must, therefore, be admitted that the cells and the fibrils are both constituent parts of one and the same organization. When this growth vegetates undisturbed, its fibrils penetrate between the layers of the epithelial cells, but do not extend deeper than the inferior layer, though they spread laterally in every direction. On the free surface of the epithelium, the ramifications rise above the surface, exhibiting at the same time an abundant fructification, which gives a yellowish hue to the exterior."

M. Parrot, in describing the appearances of thrush upon the gastric mucous membrane, states that the disease presented itself in the form of small prominent rounded masses, of yellowish color, and either isolated or in groups. These were adherent to the mucous membrane, nearly all umbilicated, and upon pressure the central depression became filled with a cheesy-looking material. On microscopic examination of sections, the spores and filaments of the muguet were found infiltrating the tissue, and as it were planted there, at times scattered in small numbers, at others accumulated in large masses, and holding between them many oil-drops and some débris of the mucous membrane. The muscular coat of the stomach was not involved, but in some instances the spores and filaments penetrated the mucous membrane, and extended to the submucous space. In other cases the mucous membrane was only superficially involved.

The reader is referred for a more full account of the cryptogamic theory of thrush to the interesting review of Berg's work above quoted, and to Bouchut's work on the diseases of new-born children; and for a complete description of the *oidium albicans* to the work of Robin, *Histoire Naturelle des Végétaux Parasites*, Paris, 1853; the works of Parrot above quoted; and the article on thrush in Vogel's work on the *Diseases of Children* (Amer. ed., 1870, p. 99).

Symptoms of the Mild Form of Thrush.—This form is the one most frequently met with in private practice. It is mild in all its characters, and often presents no other symptoms than those connected with the mouth. These are *heat* and *dryness*, with *tenderness* of that part. The tenderness is shown by the child's crying and jerking the head backwards when the

finger is introduced into the mouth, whereas, in health, the infant will almost always seize the finger and suck it with considerable force. It is shown, also, by the refusal to take the breast, or by the difficulty with which this is done, the child occasionally letting the nipple drop with a cry of pain, then seizing it again, and again dropping it with fretting or screaming. In most of the cases there are various signs of disorder of the alimentary tract, which are, however, seldom severe. They consist of slight diarrhœa, the stools being at first yellow, and afterwards green and acid; of occasional vomiting, of attacks of colicky pain, and sometimes of feverishness. To show how frequent is the occurrence of diarrhœa in thrush, and to prove also that it is not a necessary accompaniment of the disease, as has been supposed by some persons, we will quote the fact mentioned by Dr. Berg, that of 115 cases, in only 29 did the stools retain the normal yellow color throughout the whole course of the disease; while in the remaining 86, green evacuations appeared simultaneously with the invasion, or supervened at a later period. We may cite also the cases reported by MM. Trousseau and Delpech, of which only 14 out of 58 presented neither gastric nor intestinal complications.

The amount of exudation is generally small in this form, and it rarely extends behind the soft palate. The *duration* is usually between four and nineteen days, the average being about eight or twelve. Even in mild cases an improper diet, which does not agree with the child, is nearly always the cause of the attacks. If this be changed in time, to one which agrees with the patient, and which satisfies the needs of its constitution, the termination is almost always favorable. If, on the contrary, the real cause be not appreciated, and the unwholesome food be persevered in, the case is apt soon to pass into the grave form.

Grave Form.—It is under this form that the disease is most apt to occur in public institutions for children, and particularly in foundling hospitals. That it sometimes occurs, also, in private practice, will not be doubted, we think, by any who will read with care the descriptions of the disease given by Underwood, Dewees, and Eberle. We have ourselves met with two fatal cases in private practice, which presented all the symptoms described by M. Valleix as characterizing those observed by him in the Foundling Hospital at Paris, with the single exception of the ulcerations upon the internal malleoli. They were both children of parents who had every comfort at their command. One died at the age of four weeks, in consequence of the attempt to rear it on artificial diet. The other perished when six weeks old, apparently from some unhealthy condition of the mother's milk, which seems the more probable from the fact that the same mother had previously lost two children under precisely similar circumstances; all the children of this person were born vigorous and hearty, and did well for a short time, but soon after their birth, the nipples of the mother became dreadfully excoriated, the digestive organs of the infant began to give way, and death finally occurred with all the symptoms of fully developed thrush.

The most important *symptoms* of the grave form are the buccal *exudation*, various *abdominal* symptoms, particularly diarrhœa, vomiting, and

colic, and more or less marked *fever*. The order of succession of the symptoms of severe thrush is not always the same. In most of the cases the first symptom observed is, probably, diarrhœa, which is soon followed by fever, and in a few days by the appearance of the false membrane in the mouth. In a smaller number of instances the buccal exudation is the first symptom observed. The characters of the exudation are much the same as those observed in the mild form of the disease, except that the membrane is thicker, covers a larger portion of the mouth, and generally extends to the pharynx and œsophagus. In addition to the plastic deposit, there sometimes exist, especially in very bad cases, ulcerations upon the roof of the mouth, frænum linguæ, and gums. These are generally few in number, and either confined to the mucous tissue, or they may extend to the fibrous texture beneath; the surface upon which they rest is generally softened in consistence; their edges are irregular, soft, and of a whitish or reddish color. The heat of the mouth is not generally increased, except in very severe cases; the mouth is moist at first, but afterwards becomes very dry, and, from the refusal to suck the finger when it is introduced between the lips, and the difficulty with which the acts of suckling or feeding are performed, is evidently tender and painful.

The symptoms depending on the enteritic affection, are *tenderness* of the abdomen, *diarrhœa*, *vomiting*, and *fever*. The abdomen is usually distended by flatulent collections in the bowels, and is more or less painful to the touch, particularly in the right iliac fossa and epigastrium, and in severe cases over its whole extent. At the same time the child evidently suffers from colicky pains, as shown by restlessness, by uneasy, twisting movements of the trunk, by kicking of the limbs, and by crying, particularly just before or at the moment of the evacuations. The appetite is diminished and the thirst increased. The child desires to nurse or take the bottle more frequently, but is sooner satisfied than is natural, and often, from commencing soreness of the mouth, drops the nipple or bottle soon after taking it. The quantity of urine diminishes notably, and the deeper stain on the napkins shows that its color is darker than before. The amount of urea in it increases very markedly, according to M. Parrot, becoming three or four times greater than natural, and, a little later, uric acid and urates make their appearance. Albumen also is found very frequently. In rapid and violent cases it appears early, and continues to the end. In chronic forms it does not appear until the emaciation is marked, and it is apt to disappear or diminish as the fatal event approaches. It is found even in cases destined to recover, though in such cases only in very small quantity. Diarrhœa comes on. The stools at first are natural in color, but soon become greenish. They are often excessively fetid, contain more or less considerable quantities of undigested caseine, which are whitish, or pale yellow in tint, and intermingled with fluid portions that run through the napkins into the clothes. As the case progresses, they become more and more liquid and numerous, and almost invariably of a bright-green color, and very acid. The green color of the discharges, and their highly acid condition, is noticed by all observers. *Vomiting* occurs in many of the cases, but is less frequent than

diarrhœa. In some instances it is very obstinate and distressing, causing the rejection of whatever alimentary substances the child may take. Under these circumstances it has often been observed to coincide with the presence of a great deal of exudation upon the base of the tongue and soft palate, which has been supposed to act as its exciting cause. In other instances it is not so frequent, and as the matters ejected consist of greenish or yellowish bile, while, at the same time, the epigastrium is very sensible to pressure, this form of vomiting has been thought to depend upon gastritis.

In most cases some fever develops in the early stage. The temperature, however, is very irregular, and the thermometer seldom shows a rise of more than 3° or 4° F. The pulse becomes frequent, running up to 120, 140, or 160. In severe cases, and especially in those approaching a fatal termination, the temperature sinks below the normal, falling as low as 94° or 95° F. The feverish condition is often marked by restlessness and fretting, and often by loud and frequent crying. When the exudation extends into the pharynx or glottis, the cry becomes hoarse and indistinct.

There are two other symptoms which occur in the course of thrush, about which some discussion has arisen. These are, the appearance of an *erythematous redness* about the anus, and upon the buttocks, genitals, and upper parts of the thighs, and *ulcerations* upon the internal malleoli. The erythema is stated by M. Valleix to precede the other symptoms in the greater number of instances, whilst MM. Trousseau and Delpech deny the correctness of the assertion, and observed it to follow the diarrhœa in the majority of their cases. It seems to us that the latter authors are correct in ascribing the erythema to the irritation produced by the contact of the urine with the skin, which is predisposed, by the cachectic state of the constitution, to take on inflammation from causes which would not affect it in a healthy subject. The erythema is sometimes followed by papules, vesicles, blebs, and ulcerations, all of which probably depend upon the cause just referred to. The malleolar ulcerations are ascribed to the friction of the ankles against each other, a cause sufficient to produce such an effect in a broken-down, diseased constitution, though insufficient in a healthy one. We may mention that we have seen the erythema frequently in private practice, but never the malleolar ulcerations.

During the acute period of the disease, the *strength* of the child is not much diminished, but as the case approaches its termination, if no favorable change takes place, the patient becomes weak and exhausted; the face assumes a pale and sallow look; the features are sharp and defined, and the eyes dull and surrounded by bluish circles. At the same time the whole body becomes emaciated, the skin loses its elasticity, and hangs in folds or wrinkles upon the limbs, and the surface assumes a dark and dingy hue. As the fatal termination approaches, all restlessness ceases, and the child lies profoundly still, or only moves the mouth from time to time, or utters a faint cry; the diarrhœa diminishes, and the vomiting generally ceases; the pulse becomes very rapid and weak, the extremities cold, and death occurs in the midst of profound quiet, or after a few slight convulsive movements. The *duration* of this form of the disease is very

uncertain. It is often less than that of the mild form, since many children die in the first five days after the appearance of the exudation. In other cases it is much longer, from a few weeks to two months. Relapses are not uncommon.

Before closing our remarks upon the symptoms, it is proper to state that the disease sometimes occurs at the termination of acute local affections, as pneumonia, bronchitis, or pleurisy, under which circumstances there will be, in addition to the symptoms peculiar to thrush, those of the malady which preceded it.

NATURE OF THE DISEASE.—Repeated microscopic examinations have so uniformly confirmed the statements of Gruby and Berg, that it is no longer doubtful that a peculiar parasite, *oidium albicans*, is a constant element in the exudation of thrush. It is, however, far from being so well determined what relation this growth bears to the disease; since, while one class of authorities consider it the essential and sole cause of the other local and general symptoms, another regard it merely as an epi-phenomenon, the spores of the parasite finding a suitable nidus for development on the already diseased mucous membrane.

We believe thrush to be a constitutional state, in which the local symptom from which the disease has derived its English name, is merely one of the phenomena of a deep and wide-spread perversion of the general health.

We believe the real cause of thrush to be, in the vast majority of cases, the attempt to bring up the child on other than its natural food. M. Val-leix says, "I have never known a child who had been suckled exclusively during the early months of life to have the disease." MM. Trousseau and Delpech say, "We should be justified, therefore, in asserting that we have never known an infant to die of thrush who had been suckled at a healthy breast, or whose health had not been dangerously complicated by other causes." M. Parrot, who is so much opposed to the use of the word inanition as the cause of thrush, as to have invented a new one, *athrepsia*, writes (p. 382), "Vicious *ingesta* are in effect the most frequent and most powerful cause;" again (p. 383), he says that it is unusual for the digestive disorders (of *athrepsia*) to be caused by insufficient alimentation, and, "contrary to the general opinion, I believe that, in most of the cases referred to inanition, it has been an unwholesome food which has made the child ill. It is because cow's milk, or some other food, even less well adapted to the digestive organs of the new-born child, has been substituted for the breast." He writes, a little further on: "So, the term *inanition* is not well chosen to express the condition of the patients we are now studying. It is not, I repeat, the quantity which does the harm, but the quality." At page 411, he states that "the disease has for its constant point of departure a vitiated digestion, followed by an insufficient alimentation; step by step this extends to the whole organism. At the outset acquisition diminishes, then ceases. So the proteine and fatty tissues are themselves burned. To live, the individual consumes himself, and the term of existence alone is the limit of the *autophagia*." And when we, also, can state that we have seen but one fatal case of thrush in a suckled child, it is surely plain that the disease must consist in some perversion of health

determined by the food which has been substituted for the natural aliment. It is, in truth, a form of inanition,—not direct, from deprivation of all food, but indirect, concealed, but none the less an inanition. If the children ask us for bread, and we give them stones, they must die. Some of the artificial foods used are no better than stones.

In discussing the nature of the disease, there is one feature which has forced itself upon our attention of late years, which we think may be of great importance. It is particularly since we have studied M. Parrot's masterly description of the disease, as seen by him in the Foundling Hospital of Paris, that this consideration has grown upon us.

It is impossible to read his descriptions without being impressed with the fact, that there is a singular absence of water in the economy. Both in life and after death, there is constant evidence that the body is too dry. Can it be that the deficiency of water in the food may have an effect in producing so grave a disease as thrush or athrepsia? We will first quote, from M. Parrot, some passages bearing upon the curious diminution of water in the body, and then refer to the food he employs, in order to see whether there may possibly be some connection between the two. The matter is a very important one practically, as it bears strongly upon the question how best we can prepare cow's milk for the artificial food of new-born and very young children.

At page 59, M. Parrot writes: "The emaciation is considerable, and exhibits something quite special to it, for it bears more upon the fluids than solids. The whole organism suffers from aridity, and one might say that the tissues were dried up. Hence arise some features readily perceived by the eye and hand. The flesh has a peculiar sense to the touch; when pressed upon it feels like congealed suet or like wood." At page 407, he says: "In order for the chemical phenomena of nutrition to be carried on, it is necessary that the interior tissues '*milieu interieur*' be humid, that their plasma be abundant. If the physico-chemical composition of the blood be sensibly changed, vitality is disturbed and may be arrested. Have I not repeated to you, even to satiety, that these unfavorable conditions are precisely those of athrepsia? All goes to prove a change in the nutrient fluid. The diminution of its water is shown by the shrunken body, the arid and withered skin, the depressed fontanelle, the dry mucous membrane. The impoverishment of its plasma, and, if we may so speak, its concentration, declare themselves in the cyanosis and in the relative increase of the red globules."

Another condition of the disease during life, which shows a loss of water in the system, is the great diminution or even suppression of the urinary discharge, the presence of a large excess of urea, of uric acid, and of urates in the urine. After death a peculiar dryness of the tissues is observed. The blood is diminished in quantity, and it is often inspissated, as it were. It is disposed to coagulate and form thromboses. One of the most peculiar post-mortem appearances observed, and one which points most clearly to a want of water in the economy, is the presence of saline concretions in the tubules of the kidneys, which we referred to in the paragraph on Anatomical Lèsions.

We will pass on to a consideration of the food employed at the Enfants-Assistés Hospital, to see whether this food may not be deficient in the simple but vital element of water, and so explain in part, at least, one of the many difficulties which beset us in the search for a safe artificial food for infants. The very simplicity of the element water may make us thoughtless in regard to its great importance. If a man can really live for forty days, as is now asserted, on water and air alone, we must be very careful to supply to the feeble and silent infant, all the water it needs, and which it can neither ask nor make a sign for.

M. Parrot, as we shall state in the chapter on food, advocates the use of pure cow's milk, of good quality, for new-born children. He is opposed to its dilution with water. He believes, from careful investigation (see chapter on Food) that the proper daily quantities of this food are $9\frac{1}{2}$ ounces in the first month; 19 ounces in the second, third, fourth, and fifth months; and 25 ounces in the sixth month. We, on the contrary, believe that from 16 to 24 ounces of food are required in the first month, from 32 to 48 ounces in the second, third, and fourth months, and that, not very unfrequently, hearty, hand-fed children take in the fourth month, and afterwards, as much as 64 ounces of food per day.

We never give cow's milk pure in the first month, but always dilute it with two-thirds water, so that were M. Parrot's allowance of pure milk for the first month $9\frac{1}{2}$ ounces, diluted two-thirds, it would make $28\frac{1}{2}$ ounces of food, or very nearly the same in bulk that we give.

We present these thoughts upon the nature of thrush, and especially those upon the possibility that a deficiency of water in the artificial food supplied, may be one of its causes, with some diffidence to the reader. We well know how easy it is to be run away with by an idea, but we have been long convinced, both from scientific and empirical considerations, that the use of pure cow's milk, at least for new-born infants and those under two or three months, is a dangerous practice. And as we are now disposed to believe that a chief error in this practice, is the fact that it gives too little water to the infant, for its active physiological tissue metamorphoses, we deem it wise to call attention to the subject.

DIAGNOSIS.—The diagnosis of thrush is rarely difficult. Aphthæ differ from it in their vesicular nature during the formative stage, in the ulcerations which follow the vesicles, and in the absence of false membranes. From ulcero-membranous stomatitis it may be distinguished, by the formation in that disease of false membrane in layers from the beginning; by the presence of ulcerations; by the spongy, bleeding state of the gums; by the fetid breath; by the absence of the abdominal symptoms which exist in thrush; and by the microscopic appearances of the deposit.

PROGNOSIS.—The prognosis must depend, in great measure, upon the circumstances under which the disease occurs. In private practice, and whenever the patients are suckled by their own mothers, or by healthy nurses, it is as a rule a mild affection. But in foundling hospitals, on the contrary, where the children are mostly brought up by hand, it is one of the most fatal maladies to which children are subject. The prognosis

varies according to the form of the disease. The mild form is rarely fatal, while the grave form is fatal in the great majority of cases.

To show the frightful severity of the disease under certain circumstances, we may mention that of 140 cases which occurred in the wards of M. Barron, at the Foundling Hospital of Paris, only 29 recovered; while of 22 cases observed by M. Valleix, in the same hospital, but 2 recovered (Valleix, *loc. cit.*, p. 74). Again, M. Bouchut states that of 42 cases observed by himself, at the Necker Hospital, 14 were of the idiopathic (mild) form, all of which terminated favorably; and 28 of the grave or symptomatic form, of which 20 died, and 8 left the hospital still laboring under the disease. Of the 20 fatal cases, 12 presented the lesions of chronic entero-colitis, 4 of acute entero-colitis, 8 of pneumonia, and 1 of hydrocephalus.

It is a curious fact, and one, we fear, of bad omen, that M. Parrot says not a word about prognosis, or about the fatality of the disease he describes so well. It may be stated in conclusion that the danger is greatest in private practice, when the child is fed on artificial food, and depends upon the manner in which this food happens to suit the particular child. When the disease appears the food ought, as a general rule, to be changed. After such change the prognosis must depend on the fact whether the new food suit better than the previous one. If it do, the disease will probably soon begin to amend, and the prognosis at once becomes more favorable. If the child is being nursed the state of the nurse should be carefully investigated, and if her health be at all in a dubious state she should be treated medicinally or dietetically, or the nurse should be changed. It would rarely be wise to wean the child at such a moment, unless the necessity is unmistakable.

TREATMENT.—If it be true that thrush is the result of a slow, starving process, determined not by deprivation of food, but by the use of food unfit and incompetent to develop new-born and very young children, it is clear that the primary indication of treatment must be to find a proper food. To discover such a food in each particular case, and apply it, constitutes the chief duty of the physician.

But he has other duties besides this. He should examine into the general hygienic surroundings of the child, and do all that he can to insure it fresh air, cleanliness, and constant attention to its wants. A young hand-fed child ought to have one woman to attend to it, and she should be taught by the physician how to exactly prepare the food, how much to administer at each feeding, and how often, day and night, the feedings are to be repeated. One of the causes of the shocking mortality of new-born children in hospitals is the fact that the number of nurses is insufficient. One woman will have the care of three, four, or more children. It is simply impossible for her to take full and proper care of so many.

In addition to the care as to the food and general hygiene of the patient, much may often be done in the early stage of thrush, before it has reached the grave stage, by a proper use of remedial agents.

If a suckling child be seized with the disease, the health of the nurse ought to be carefully investigated. If this be found deranged, perhaps

by over-fatigue, by worry, by dyspepsia, everything should be done that is possible to remove the cause. The milk should be examined with the microscope, and if any possible fault be found in it, the nurse ought to be changed. If this cannot be, it becomes a question whether the child had best be weaned and put upon a proper artificial food. We confess that we ourselves have such a dread of weaning that we never recommend it until at the very last extremity, and we believe that a really unwholesome milk is a rare thing in the breast of a woman having the signs and appearance of average fair health. In case the mother or nurse have too little milk for the child, we think it far better to continue the nursing, and to feed the child in part. The food must be carefully selected, and may be administered alternately with the nursing.

When the child attacked with thrush is already being fed artificially, as is the case in the great majority of instances, the first question to be solved is whether the food being used is the best in the particular case? For the answer to this question we must refer the reader to the chapter on food, where the whole matter is carefully treated of in detail. We recommend the food composed of cow's milk, cream, milk of sugar, and arrow-root water, as these are laid down for the different ages of infancy. If this have been tried already, or being tried, should fail to suit the case, we advise the diet made of equal parts of milk, cream, lime-water, and plain water. Should this not answer, the cow's milk may be diluted for a short time, with three instead of two parts of water, or pure cream, diluted with six or eight parts of water may be given. It is in such cases as these that condensed milk seems sometimes to succeed. If it be used it ought to be given in the proportion of one heaped teaspoonful to six tablespoonfuls of water, which is the strength of one part fresh milk to two parts water. Sometimes the use of one tablespoonful of lime-water in place of one of the tablespoonfuls of plain water makes it more digestible.

The exact doses and the strength of the food ought to be determined by the physician for each particular case. Nothing but close and careful observation will reveal what is best for each individual child.

When no food can be found to suit the case, and when the child is too weak to suck, a plan which has succeeded with us in a few instances has been to bring a wet-nurse to the house, to have her breast drawn by a breast-pump, and feed it to the child from a spoon or sucking-bottle. It should be given in stated and moderate doses for a time, increasing the dose carefully, as it is found to suit the child.

Besides the food, we believe that in thrush, as in other diseases of the digestive and nutritive functions, it is highly important to administer water to the infant. Two, four, six, or more teaspoonfuls should be offered the baby regularly, half-way between the doses of food, and the child ought to be allowed to take all it wants. It is almost always well to add brandy to the water, certainly in a severe case, and in one attended with considerable diarrhœa or vomiting. Half a teaspoonful of it to a gill of water is the proportion we generally direct, and we give all the child will take with pleasure.

As to the medical treatment, we believe that the measures found most

useful in indigestion, dyspepsia, and diarrhœa, are the proper ones. When the stools are not watery, but in part pasty from undigested milk, and fetid, some mild laxative ought to be given. We prefer the spiced syrup of rhubarb, half a teaspoonful to a teaspoonful once or twice a day. It is well, as a general rule, to combine a little opium with this dose, a quarter or half a drop of laudanum, or three or four drops of paregoric. Or half a teaspoonful of castor oil may be used. After the laxative has been given once or twice, the following mixture, which we have used a great deal in digestive disorders of infants, may be ordered:

R. Sodii Bicarb.,	3 ss.
Tr. Opii Camph.,	gtt. xl.
Tr. Rhei Dulc.,	gtt. lxxx.
Syr. Symp.,	f 3 ij.
Aq. Menthæ Pip.,	f 3 xiv.—M.

Dose. A teaspoonful three times a day.

When the diarrhœa is more severe, and the stools watery and green, the following mixture will be found useful:

R. Sodii Bicarb.,	3j.
Tr. Opii Camph.,	f 3 ss.
Tr. Krameriæ,	f 3j.
Syr. Simp.,	f 3 ijss.
Aq. Menthæ Pip.,	f 3 ijss.—M.

Dose. A teaspoonful three or four times a day.

Or a weak chalk mixture, with paregoric and rhatany may be employed.

It is often well to use pepsin in such cases to strengthen, if possible, the digestion. Ten drops of the wine of pepsin may be given with each meal, or half a grain of the saccharated pepsin three times a day.

Local Treatment.—This should be simple and of a soothing character. A solution of chlorate of potash, five grains to the ounce, one of borax, ten or twelve grains to the ounce, should be pencilled over the mouth, two or three times a day, or applied very gently by means of a soft rag wrapped over the finger. The practice of rubbing the tender and morbid surface with a rag held in the fingers of an awkward and heavy-handed nurse is very injurious. Nor do we approve of one of the favorite applications of the nursery, powdered borax and sugar. It is often applied too copiously, and we have known it to collect between the lip and gum, and cause severe irritation. One of the best applications, we think, is a solution of nitrate of silver, half a grain or a grain to the ounce of distilled water.

In case of erythema or ulceration, the most scrupulous cleanliness is necessary. The erythematous surface may be dusted with powdered starch, rice, or lycopodium. The ulceration may be dressed with benzoated oxide of zinc ointment. When the ulcerations are deep M. Parrot recommends dusting them very lightly with iodoform, and dressing with charpie covered with cerate or cucumber ointment.

ARTICLE VII.

AFFECTIONS OF THE TONSILS.

1. ACUTE INFLAMMATION OF THE TONSILS.—TONSILLITIS.—This painful affection (known also as angina or cynanche tonsillaris), occurs in childhood less frequently, but with the same symptoms as after the age of puberty. We have occasionally met with severe cases of it in children under the age of 5 years.

SYMPTOMS.—When the attack is sudden, there are marked fever, restlessness or heaviness, and complaints of severe pain on deglutition. We have known the fever to reach 104° by the close of the second day, with a pulse rate of 140, in a child of highly nervous temperament. If the child be old enough to answer questions, the pain will be found to radiate from the fauces towards the ear, and to be increased by opening the mouth. Painful enlargement of one or both tonsil glands will be found by pressing the finger below the angle of the lower jaw. On examining the fauces, there is marked redness of the half arches and posterior border of the soft palate; the affected tonsil projects from its bed as a rounded, deep-red body, which may extend even beyond the median line; and if, as less frequently happens, both tonsils are severely inflamed at the same time, they may even meet and entirely occlude the isthmus of the fauces. The surface of the gland often presents small yellowish points which closely resemble patches of false membrane, although careful inspection will show that they are beneath the mucous membrane, and are really only the distended follicles of the gland. Deglutition is so painful, especially for liquids, such as milk or water, that the little patients will at times utterly refuse to swallow.

COURSE AND DURATION.—The disease lasts from 3 to 7 days, and terminates in different ways. It very rarely proves fatal, and only does so by obstructing breathing, and at the same time so seriously interfering with nutrition that the child's strength fails. In most cases the result is favorable, and the termination is either by suppuration or gradual resolution of the enlarged gland. When suppuration occurs, the symptoms have gone on becoming more and more aggravated until they reach their height, and the case seems attended with great danger, when suddenly, after an effort at vomiting, or spontaneously, the tonsillar abscess bursts, a gush of pus occurs from the mouth, and prompt relief is afforded. Occasionally the occurrence of suppuration is marked by a chill, or some decided change in the febrile movement.

More frequently in children, however, the tonsil does not suppurate, but gradually becomes smaller; the redness subsides, and the distended follicles disappear. There is a strong tendency, especially after this latter mode of termination, for the tonsil to pass into a state of moderate chronic enlargement.

DIAGNOSIS.—The conditions with which acute tonsillitis in children is most apt to be confounded, are diphtheria and scarlatinous angina.

From diphtheria it may be told by the more acute and sthenic character of the symptoms; by the slight swelling of the glands at the angles of the jaw, or by the absence of enlargement of any but the tonsil glands, and by the local appearances, particularly the absence of pseudo-membranous exudation.

From the angina of scarlatina it may be distinguished by the less frequent pulse and lower temperature, but chiefly by the absence of eruption, since, as already stated, the fever and pulse may quickly become so high that in some cases the diagnosis cannot be positively determined until the time at which the eruption of scarlatina makes its appearance has passed.

TREATMENT.—So long as the child is able to swallow, quinia may be given in full doses, to diminish the fever, and perhaps diminish the liability to suppuration. It may be given combined as follows:

R. Quiniæ Sulph.,	gr. xvij.
Tr. Ferri Chloridi,	gtt. xlvij.
Potassæ Chloratis,	gr. xxx.
Syr. Zinziberis,	f℥j.
Aquæ,	f℥ij.—M.

Dose. Two teaspoonfuls four times a day, for a child 5 to 7 years old.

If, however, the inflammation be very acute, suppuration will occur in all probability, despite our efforts.

In very young children, it is so difficult to make any local applications to the tonsils, that it is very doubtful whether any good effect they may produce is not more than over-balanced by the fatigue and annoyance they cause. Where, however, they can be made without so much opposition, we would recommend the daily use of the following solution:

R. Potassæ Chloratis,	℥j.
Tr. Ferri Chloridi,	f℥ss.
Glycerinæ	f℥iss.—M.

applied by a brush to the tonsils. Relief will also be obtained from the inhalation of steam or of vaporized warm lime-water. It has seemed to us positively useful to make repeated external applications, according to the tolerance of the skin, of compound tincture of iodine over the post-maxillary triangles. Warm, slightly sedative embrocations may also be applied to the neck.

It is doubtful whether poultices, or any such applications, hasten suppuration sufficiently to make up for the annoyance they cause the child. Even if the occurrence of suppuration be suspected, it is usually impossible to obtain so full a view of the parts as to enable an incision to be made to evacuate the pus. As, however, the abscess will discharge spontaneously in nearly all cases, it is only when the symptoms of obstruction of the throat are very urgent, that it is desirable to insist upon such an examination.

Guaiacum seems to exert a peculiar local action in this affection of the tonsils; and we are in the habit of prescribing the troches of guaiacum,

especially as made by Hancock, of Baltimore, even for young children. It is generally possible to induce them to take a fragment of one of these lozenges every hour or two; and, when herpetic inflammation of the follicles is marked in proportion to the deeper interstitial inflammation of the gland, their use seems to allay pain and to hasten recovery.

The treatment for the chronic enlargement which sometimes remains after an acute tonsillitis, will be considered in the next section.

2. CHRONIC ENLARGEMENT (HYPERTROPHY) OF THE TONSILS—CAUSES.—The tonsils are in young children much more subject to this affection than to acute inflammation. The enlargement may begin during the first year of life, but usually does not become sufficient to attract attention until the second or third year. Most frequently it has no connection whatever with previous acute inflammation of the part, but is chronic and indolent from the beginning. It is often observed that several children of the same family will suffer from this condition, and it is in fact associated in many cases with rickets or with scrofula. We have, however, observed marked and enduring enlargement in children of apparently sound constitution. According to West, the irritation of the latter period of first dentition may be the exciting cause in some of these cases.

We have already alluded to the fact that occasionally, especially in somewhat older children, it has been an acute attack of tonsillitis—either simple or diphtheritic—which has induced the state of chronic enlargement.

ANATOMICAL APPEARANCES.—Both tonsils usually share in the enlargement, though not always to an equal degree. They project into the fauces from either side, forming pale red tumors of rounded form, with a surface that may either be smooth and glistening, or rough and irregular from the rupture of numerous distended follicles. They impart a sense of elastic firmness to the finger when pressed. The exact anatomical condition is in part an enlargement of the follicles of the gland, associated with thickening of the fibro-cellular stroma. The term hypertrophy, commonly applied to this condition, must therefore be regarded as indicating merely the increase in the size of the gland.

SYMPTOMS.—There can be no doubt but that many symptoms have been attributed to the influence of enlarged tonsils which are in reality dependent upon entirely different causes.

The results which are constantly observed are loud snoring during sleep, snuffling, and a thick voice. There is also often a tendency to acute catarrhal attacks, during which the enlargement of the tonsils increases, and the interference with the breathing and voice is much increased. Indeed, in some unusually severe cases the respiration is constantly labored, and the child is annoyed by a frequent dry hacking cough. The pressure of the enlarged glands upon the mouths of the Eustachian tubes may produce tinnitus and hardness of hearing. The most serious results which are, by many authors, attributed to enlargement of the tonsils are alterations in the nose and upper jaw, and the production of the chicken-breast deformity of the thorax. In consequence of the obstruction of the nasal passages caused by the upward pressure of the soft palate, the form of the anterior nares may be somewhat altered and contracted, but we are rather

inclined to refer the small size of the features and the ill-developed upper jaw to the rickety cachexia which is so frequent a cause of enlargement of the tonsils. So, too, the narrowing of the isthmus of the fauces must tend to make inspiration difficult, and thus to prevent full expansion of the chest, but we can hardly imagine that such obstruction could produce marked chicken-breasted deformity of the thorax, if it were not for the fact that in such patients there is usually a high degree of rickets coëxisting. It must be borne in mind that precisely this deformity of the thorax is frequently met with in cases of rickets where there is no enlargement of the tonsils. The condition of these glands and the changes in the jaws and chest-walls must then, we think, be regarded as results of a common cause. So, too, it is probable that the sudden suffocative attacks, which have been described as occasionally attending chronic enlargement of the tonsils have been spells of laryngismus stridulus, dependent upon rachitic disease of the bones of the skull.

PROGNOSIS.—It will be readily seen, therefore, that although this condition of the tonsils is obstinate, and yields slowly, if at all, to treatment, it is rarely of itself followed by any serious consequences. In very many cases it gradually subsides after the patient reaches puberty, while in others treatment is successful in reducing the enlargement. We have, however, known it to persist most obstinately for many years, even after partial ablation and prolonged treatment.

TREATMENT.—The frequent association of enlargement of the tonsils with a rachitic or strumous diathesis must be borne in mind, and if there is any evidence of the existence of such a constitutional taint, the appropriate treatment must be adopted. Even where no decided evidence can be found, it seems desirable to administer such alterative tonics as the iodide of iron, or of the compound syrup of the phosphates of iron and alkalis. The prolonged use of cod-liver oil, with iron and arsenic, has also proved of service.

Counter-irritation by the daily application of dilute tincture of iodine, or compound iodine ointment, behind and below the angle of the jaw, may be used, and sometimes appears to favor the reduction of the swelling.

Local applications to the enlarged glands are of much service in some cases, but to do good must be steadily persisted in, in conjunction with proper internal remedies. Those which have on the whole appeared most useful to us have been Lugol's solution of iodine diluted with two to four parts of water, and nitrate of silver in the form of rather strong solution, as gr. x to the fluidounce.

We have also found it of material service in hastening the reduction of the enlargement to whiten the surface of the tonsils once in three or four days by a light application of solid lunar caustic.

Recently, the use of injections of ergot and iodine as a means of causing the resolution of enlarged glands has been frequently recommended. We have found positive benefit from their employment in cases where the children were old enough to permit this little operation to be performed without too much alarm. The injections should be made into the substance of the hypertrophied tonsil, by means of an ordinary hypodermic syringe; and may

be repeated about once a week, or according to the effect produced. The amount injected should not exceed a few drops, 3 to 5, and should be introduced very gently, so as to avoid pain as far as possible. The fluid used may be either a simple solution of iodine diluted with water, or a dilute solution of ergotin, 48 to 96 grains to the ounce.

Careful attention to diet, and particularly to the proper and sufficient clothing of the child must be insisted on, so as to avoid, as far as possible, the repeated acute attacks of slight tonsillitis which are apt to occur. Under the persistent employment of the general and local means above recommended, we have usually found that the hypertrophy of the tonsils has diminished towards the age of puberty. In some instances, however, we must confess that all forms of treatment, general as well as local, have proved unavailing. We must then resort to excision of the enlarged glands, if the symptoms caused by their presence are sufficiently urgent to render it advisable.

The excision of the tonsil (or rather of the prominent portion of it, for the entire gland rarely needs removal) is an operation attended, in skilful hands, with little difficulty and no danger. It may be readily performed with a Fahnestock's or Physick's tonsillotome, or, as many operators prefer, by raising the gland from its bed with a special kind of forceps, and then slicing it off with a bistoury.

The symptoms which would lead us to advise the early removal of the tonsils are frequent irritative cough, much interference with hearing or with the tones of the voice, or coëxisting rachitic deformity of the chest.

ARTICLE VIII.

SIMPLE OR ERYTHEMATOUS PHARYNGITIS.

DEFINITION ; FREQUENCY.—Simple pharyngitis consists of an erythematous inflammation of the pharynx, tonsils, and soft palate, unaccompanied by ulceration, deposit of false membrane, or gangrene. It is very frequent both as an idiopathic and secondary disease. We constantly meet with it in children of all ages during the cool months of the year.

CAUSES.—It may occur at all ages, and is equally common in the two sexes. The diseases in the course of which it is most apt to occur as a secondary affection, are scarlet fever and measles, and next, pneumonia and bronchitis. It is often an accompaniment of simple laryngitis. The idiopathic form is most common in this city in the late winter and early spring months. It is said to prevail sometimes in an epidemic form.

The *exciting causes* of the disease are not always easily detected. In most instances, however, we believe that exposure to cold is the cause of the attack.

ANATOMICAL LESIONS.—In mild cases the alterations of texture observed during life, and in a few instances after death, the patient having

died of some other disease, consist of greater or less redness, swelling, softening, and a rough or granular and sometimes œdematous condition of the mucous membrane covering the soft palate, tonsils, and pharynx. The uvula and tonsils are generally tumefied, and the crypts of the latter filled with mucous or purulent fluid of a yellowish color. In one very severe case which proved fatal, MM. Rilliet and Barthez found the tonsils very red, soft, only slightly swelled, and infiltrated with pus; the pharynx was covered with a thick layer of bloody mucus; the mucous membrane of the throat was of a dark red color, thickened, and granular, but not softened. The submaxillary glands were of a grayish color, enlarged and soft.

SYMPTOMS.—Simple pharyngitis of moderate severity begins with restlessness, irritability, fever, slight cough, and in some instances, pain in the throat, which is complained of by older children, and betrayed in those who are very young by the refusal to nurse or take food, because of the difficulty of swallowing. The *face* is generally flushed, sometimes very deeply so. Young children are often drowsy, but from irritability and fever refuse to sleep except on the lap. The *fever* is marked by acceleration of the pulse, which rises to 100, 110, or more in children over five years of age, and to 120, 130, or 140 in those under that age, and by unusual warmth or even heat of the skin. At the same time the *respiration* is generally more frequent than natural, but almost always regular; in cases attended with high fever, we have counted the breathing at 42 and 50. *Auscultation* reveals pure vesicular murmur or slight sibilant rhonchus. The *voice* is clear, or, in rather severer cases, obscured and nasal, and in some instances, speaking is painful and difficult. *Cough* is a frequent symptom. It has been present in a great majority of the cases observed by ourselves. In some of these it was harsh and croupal, so that the children seemed threatened with croup. The croupal sound seldom lasted over one night, after which the cough was merely hoarse, and gradually became loose towards the termination of the attack. In the remaining cases it was rare and dry in the beginning, and looser and more frequent as the disease progressed. *Pain* is a frequent, but far from constant symptom at the outset of the disease. It generally exists during deglutition. When present it is shown in infants, as stated, by their refusing the breast, or nursing only at long intervals, and with difficulty; while in older children it is complained of. It is not, however, a constant symptom, as we have often seen children of one, two, and three years old, with severe angina productive of violent fever, who swallowed fluids and soft solids without a sign of pain. Throughout the acute period of the disease there is generally considerable *thirst*; the *appetite* is diminished or entirely suppressed; the *stools* are usually natural, or there is slight constipation.

The throat should always be examined when there is the least reason to suspect that an attack of sickness depends upon inflammation of that part, and whenever a child has been seized suddenly with fever, particularly in cold weather, and there is nothing more evident by which to explain the illness. To examine this part well, the tongue must be strongly depressed

with a small tongue-depressor or with the handle of a spoon, which should be carried back to the base of the tongue. This may be done in the youngest infant.

The *appearances* presented by the throat are as follows: The soft palate, uvula, tonsils, and generally the pharynx also, are more or less reddened and swollen, and the mucous membrane commonly looks rough and granular. The fauces are often filled with frothy mucus, and in severe cases coated all over with mucous or purulent secretions, which sometimes line the inflamed surfaces in such a way as to resemble false membranes. They are to be distinguished only by careful examination, and by removing a small portion on a pencil or sponge-mop, in order to ascertain their real nature. We have seen the mild form of inflammation in a child ten days old, in one eight weeks, another three months, and a fourth nine months old.

Dr. Wertheimer (*Jour. für Kinderkrankheiten*, Band xxxii) calls attention to a variety of angina, which he calls *œdematous*, and which is specially characterized by serous infiltration of the submucous tissue of the pharynx, the mucous membrane itself being pale and smooth, and soft and sticky to the touch.

The *submaxillary glands* and neighboring cellular tissue are sometimes swollen, in consequence of the extension of the inflammation to them. This is often evident to the eye, but it is more correctly judged of by the touch. At the same time the glands are usually somewhat painful to the touch. The amount of swelling is slight in very mild cases, or there may be none at all. In severer cases it is much more considerable.

The *breath* is said to be often fetid. We have not met with this character in the simple disease. *Expectoration* is rarely present. We have never noticed it under six years of age. Slight *nervous symptoms* occur in nearly all the cases, consisting, as already stated, of restlessness and irritability in mild attacks, and of insomnia or drowsiness, with starting and twitching, in those which are more severe.

The fever generally occurs at first only in the latter part of the day and during the night, often becoming intense at that time, with restlessness and starting, and subsiding or disappearing entirely towards morning, to recur again the next afternoon or evening. Children not unfrequently play about all the early part of the day, and are attacked with the symptoms just mentioned as night comes on. The disease generally pursues this course for three or four days, and then passes away entirely or, if it lasts beyond that time, the fever becomes continued, and the attack runs on for seven, eight, or ten days.

In *grave* cases of simple angina, the disease begins with *vomiting*, *fever*, and severe *nervous symptoms*, in the shape of excessive restlessness, or somnolence, and occasionally convulsions. The fever is violent, the pulse being very frequent and full, and the skin hot and flushed. The intense heat and flushing of the skin, which in sanguine children sometimes affects the greater part of the surface of the body, together with the activity of the circulation, not unfrequently make the onset of the disease resemble very closely that of scarlet fever. Four cases of this kind that have come under

our notice presented severe nervous symptoms at the invasion. In a girl between two and three years old, they consisted of wildness and ecstatic expression of the face, and trembling uncertain movements of the limbs, which would probably have terminated in convulsions, but for the timely interposition of a warm bath. In the three others, general convulsions occurred. Two of the subjects in which convulsions took place were between five and six years old, and one between three and four. In two the convulsions occurred at the onset, and in a third on the second day. The convulsive movements lasted from ten to twenty minutes, and were followed by somnolence for a few hours in two, and by stupor for a day in the third. It should be stated, however, that two of these subjects were predisposed by constitution and temperament to spasmodic attacks, as one had had a fit previously from a similar cause, and the other two from difficult dentition. The third had never suffered from any symptoms of the kind, and did not appear predisposed to them.

The *tongue* is generally dry and coated with a thick whitish fur in grave cases; the *respiration* is quick, loud, and nasal; and the *voice* guttural or nasal, and difficult. There is usually extreme *thirst*, and not unfrequently delirium. The throat is commonly violently inflamed, of a deep-red color, and coated over with mucous or purulent secretions. The submaxillary regions are often swollen, and the deglutition sometimes, though not always, difficult. When the disease proves fatal, the different symptoms soon reach their height, and death may occur in two or three days. We have never, however, known simple pharyngitis to terminate fatally. The *duration* of the grave cases is variable. In the four that we have noted, it was between three and eight days.

Secondary pharyngitis, which, as has been stated, is a very frequent disease, will be treated of in the articles on the various diseases in the course of which it occurs.

DIAGNOSIS.—The diagnosis of simple pharyngitis is not always without difficulty, as there are no local symptoms in two-thirds of the cases at the invasion, nor in some instances at any period of the attack. The physician and attendants, therefore, are often deceived as to the real cause of the violent fever which has so suddenly made its appearance, and are disposed to refer it to any but the true one.

It has happened to us several times in cases of children attacked with simple angina, to suspect pneumonia from the sudden occurrence of high fever, rapid respiration, slight, dry cough, and the absence of pain in the throat, difficulty of deglutition, or other symptoms, to call our attention to the real seat of disease. The diagnosis is to be corrected only by the absence of the physical signs of pneumonia, and the consequent necessity of finding some other cause of the sickness. Angina may be mistaken also for indigestion, which is one of the most frequent causes of sudden fever in childhood, and is accompanied, like severe angina, by vomiting. The distinction between the two is to be made by careful inquiry as to the history of the attack, by examination of the matters ejected from the stomach, and by inspection of the throat. Severe cases, particularly when ushered

in by convulsions, may be mistaken for disorder of the nervous system dependent upon dentition. The only method of ascertaining the truth is again the inspection of the throat. Cases of this kind might also be mistaken for the beginning of scarlet fever. Time only, and the development or absence of the symptoms peculiar to the latter disease, could enable us to determine the diagnosis.

The diagnosis between simple and pseudo-membranous pharyngitis will be given under the head of diphtheria.

PROGNOSIS.—Simple pharyngitis of moderate severity is very rarely, if ever, a fatal disease. Severe or grave erythematous pharyngitis, on the contrary, is often a dangerous malady. The four cases that have come under our care, however, all recovered. The unfavorable symptoms in such cases are: very violent fever, greatly altered physiognomy, difficult respiration, choked and guttural voice, excessive jactitation, delirium, convulsions, and coma.

TREATMENT.—Mild cases of simple angina need but little treatment. The child ought to be confined to a warm room in all cases, and kept in bed, or on the lap, if it have fever. The diet must be restricted to milk preparations and bread, so long as the fever continues. The therapeutical part of the treatment may consist in the use of some mild evacuant, as one or two teaspoonfuls of castor oil, half a teaspoonful or a teaspoonful of magnesia, a small quantity of syrup of rhubarb, or what is sufficient in many cases, a simple enema. At the same time we may give, if the frequency of pulse, heat of skin, and restlessness be considerable, a few doses of spirit of nitrous ether, or spiritus Mindereri, alone, or combined with about half a drop of tincture of aconite root, or from one to four drops of antimonial wine, according to the age. At the same time, it is well to give a moderate amount of quinia, which, owing to the pain in deglutition, we are in the habit of giving in the form of very small suppositories, containing one or one and a half grains of quinia, to be repeated from two to five times in twenty-four hours, according to the age and the degree of fever. A warm bath, if the child is not afraid of it, is an admirable remedy when there is much excitement of the circulation; or a foot-bath, containing salt or mustard, may be used. Frictions over the throat and neck are often very advantageous; they may be made with hartshorn and sweet oil, with or without the addition of laudanum, or a small quantity of spirit of turpentine may be applied upon the skin, so as to produce slight counter-irritation. When there is much pain and difficulty of deglutition, the case is best treated by the use of nitrate of silver in solution (5 or 10 grains to the ounce), or of powdered alum, applied by means of a large throat-brush.

In the *severe form* of the disease the treatment must be much more active than in mild cases. When the fever is very high, and threatening nervous symptoms are present, the most speedy means of controlling them is a warm bath, continued for fifteen or twenty minutes. If the effects of this should be slight or transitory, one or two leeches may be applied behind the angles of the jaw, unless the fright and consequent resistance on the

part of the child are so great as to render their application objectionable. Some evacuant dose should be given early in the attack ; it may consist of castor oil, magnesia, epsom salts dissolved in lemonade, fluid extract of senna, or infusion of senna and manna. The quantity must be sufficient to produce several copious stools, and should it fail to operate in three or four hours, and the fever continue, it is always well to assist it by means of a purgative emema. Two hours after the exhibition of the cathartic, it will be proper to resort to small doses of sulphuret of antimony with Dover's powder, repeated every hour and a half or two hours, in the manner recommended in the article on pneumonia. If the secretions into the fauces be very abundant and tenacious, so as to impede respiration, the best means of getting rid of them is by an emetic of ipecacuanha or alum. If they collect again, the throat ought to be cleansed from time to time with a small sponge-mop. The inflamed surfaces should be touched two or three times a day with a solution of nitrate of silver (from five to ten grains to the ounce). The late Dr. C. D. Meigs was in the habit of employing with much benefit, in the severe angina of children, whether idiopathic or secondary, a wash made according to the following formula ;

R. Cupri Sulphat.,	
Quiniæ Sulphat., āā,	gr. vj.
Aquæ Destillatæ,	fʒj.—M.

This is applied in the same way as the lunar caustic solution, and we have frequently seen it produce most excellent effects.

The four grave cases observed by ourselves recovered under very simple treatment. This consisted in the use of the warm bath, of doses of castor oil to move the bowels freely on the first day, and of syrup of rhubarb or enemata afterwards to keep them soluble ; of doses of antimonial wine and nitre every two hours in such quantity as to avoid sickness ; of mustard foot-baths ; stimulating frictions to the outside of the throat ; applications of lunar caustic solution to the throat internally, three or four times a day ; and of rigid diet. In one case the warm bath was used three times in a single day, because of the extreme restlessness and heat of the skin, and was productive each time of much benefit.

ARTICLE IX.

RETROPHARYNGEAL ABSCESS.

THIS name is applied to collections of pus formed in the cellular tissue between the posterior wall of the pharynx and the vertebral column. More frequently the abscess is seated quite high up on the level of the

glottis, though cases are recorded where it occupied a lower position behind the œsophagus.

CAUSES.—Retropharyngeal abscess occurs idiopathically, or as a sequel to some of the specific fevers, or, more frequently, in connection with caries of the cervical vertebræ. In one of our own cases, it followed directly upon a long sleigh ride, and was due evidently to the severe chilling of the body. Although it cannot be regarded as a disease peculiar to childhood, it is far more frequent in the first ten years of life than during any subsequent decade.

SYMPTOMS.—The early symptoms are irregular and not characteristic. In cases where the abscess is connected with caries of the cervical vertebræ, the symptoms of this latter condition have preceded. In all cases, however, the first indications which lead to a suspicion of the existence of a post-pharyngeal abscess are gradually increasing difficulty of swallowing and of respiration, which is attended with a loud, stertorous sound, unlike the stridulous breathing of croup. There is also marked stiffness of the neck, and the head is rigidly retracted. Any effort to bend the head forward it followed by urgent increase of the dyspnœa, and the same result has been noticed to follow when the recumbent position was assumed. There is an appearance of fulness on one or both sides of the neck behind the angle of the lower jaw. Of course the child presents a high degree of restlessness and distress, which increases until the interference with breathing and swallowing may prove fatal from combined exhaustion and asphyxia. In the presence of such symptoms as the above, a careful examination of the pharynx, both by direct inspection, if possible, and by the finger, should immediately be made. The mouth is usually filled with mucus, and the swelling of the posterior wall of the pharynx may frequently be seen projecting forward so as to constrict the pharyngeal space, and obstruct more or less the opening of the glottis. The finger, if carried back over the root of the tongue, comes in contact with a rounded swelling, which is in the early stages firm and somewhat elastic, and later becomes fluctuating. When the abscess is fully formed, the most prominent point may appear yellowish. Occasionally, in the course of caries of the cervical vertebræ, perforation of the posterior wall of the pharynx occurs without being preceded by any such severe symptoms as have just been described as due to post-pharyngeal abscess. We have thus known the expectoration of purulent matter with small fragments of carious bone, to occur in such cases without any previous symptoms of marked obstruction in swallowing or breathing.

DIAGNOSIS.—The recognition of this affection is often difficult, and it is only by bearing in mind the possibility of its occurrence, and making careful examination with the finger, that we can avoid overlooking its existence, in cases where the symptoms are not clearly pronounced. In all cases, therefore, where difficulty in swallowing is superadded to dyspnœa, such an examination should be made. The affection with which it is most likely to be confounded is membranous croup, but the absence of the peculiar croupy cough and stridulous breathing, and the existence of dysphagia,

retraction of the head, with immobility of the neck, fulness at the angle of the lower jaw, and, finally, the detection of the swelling at the back part of the throat, will render the diagnosis easy.

PROGNOSIS.—The termination is always doubtful. When, however, the existence of the abscess is early recognized, and it is promptly evacuated so soon as fully formed, recovery frequently ensues. Even when connected with caries of the vertebræ, the prognosis, although of course unfavorable from the nature of the primary disease, is not necessarily fatal. In the case already referred to as having followed directly upon long exposure to severe cold, the child was very ill, with all the characteristic symptoms of this affection, for a week, after which the abscess burst spontaneously, and the child recovered.

TREATMENT.—The approach of a post-pharyngeal abscess can rarely be detected so early as to enable any preventive treatment to be adopted with success. Indeed, but little could be expected from the use of mild counter-irritants, or absorbent applications to the throat. In older children, if recognized before suppuration has occurred, some benefit might be derived from the use of one or two leeches to the angles of the jaw, or of a blister to the back of the neck. The use of small pieces of ice held in the mouth will also be found to afford relief. The main indication, however, is to watch for the occurrence of suppuration, and then to make as early an incision as possible. When the seat of the abscess is high up this may be done by an ordinary sharp-pointed bistoury, whose blade is guarded up to near the point by being wrapped with adhesive plaster. When the abscess is lower down it can sometimes be more safely reached and evacuated by a trocar and canula. As the opening should be small, there is danger of its closing with a re-accumulation of pus; and it is therefore advisable, as recommended by West, to press with the finger upon the sac of the abscess occasionally for a day or two. In cases dependent upon caries of the vertebræ, it is better to postpone opening the abscess until urgent symptoms are produced by it. Here also it is necessary to employ the other means of treatment suitable for that condition, and especially the use of some mechanical contrivance by which the weight of the head can be supported, and thus relief be afforded to the cervical spine. During the course of the disease every effort must be made to sustain the strength of the patient. If the interference with swallowing be extreme, nutritious enemata should be used until the abscess can be evacuated. In addition, we must use opiates in sufficient amount to quiet the excessive pain and restlessness.

CHAPTER II.

DISEASES OF STOMACH AND INTESTINES.

GENERAL REMARKS.

IN our division of these diseases, we shall treat first of Indigestion, using this term to signify morbid conditions of the digestive function, which we suppose to be the result of functional disorder, or of mild, acute, or chronic catarrh of the stomach. Under the title of Gastritis we shall describe the much more rare and dangerous form of disease, in which there is acute inflammation of one or more of the coats of the stomach, and which is seldom met with except as the consequence of the application of some direct irritant to the organ.

We shall then describe Simple Diarrhœa, in which we suppose the intestinal disorder to be either merely functional, or one of slight catarrhal inflammation of the mucous membrane. Next, under the title of Enterocolitis or Inflammatory Diarrhœa, we shall treat of that form of diarrhœa which is now by many writers styled acute or chronic catarrh of the intestinal mucous membrane, and the chronic forms of which we believe to be of the same nature as the disease designated by most of the observers whose experience was gathered in the vast field of the late war, chronic diarrhœa. We shall pass on then to Cholera Infantum, limiting this term to cases in which the disease is of a true choleraic type; and lastly, we shall consider Dysentery. We have also added separate articles on the diseases of the Cæcum and Appendix Vermiformis, and upon Intussusception.

SECTION I.

FUNCTIONAL DISEASES OR MILD CATARRH OF THE STOMACH AND INTESTINES.

ARTICLE I.

INDIGESTION.

DEFINITION; FREQUENCY; FORMS.—By the term indigestion, we mean that condition of the stomach in which its function of digestion is disturbed or suspended, independent of inflammation or other disease of the organ, appreciable by our senses; or in which there have been found after death, in the few opportunities that have been met with to make such an investigation, the lesions which are now usually designated as mild gastric catarrh. The only anatomical alterations found in such cases, are

reddening of the mucous membrane in spots by a fine injection, relaxation of its tissue, and the presence of a layer of tough mucus. It is a very frequent affection during the whole period of childhood, and is one of great importance on this account, and from the fact of its laying the constitution open, by the debility and cachexia which it produces, to various secondary affections. In our description of the disease, we shall distinguish between the forms which occur during infancy, and after the completion of the first dentition.

CAUSES.—The principal causes of indigestion in infants are an unhealthy state of the milk of the nurse, the use of artificial diet, and lastly, an impaired condition of the digestive function, which disables the stomach from digesting even healthful aliment.

The milk of the nurse may be too old for the child, for it has been found that a breast several months old sometimes, though not always, disagrees with a young infant, in consequence, no doubt, of the milk being somewhat thicker and richer at that time than immediately after parturition. The breast-glands may continue to secrete colostrum for weeks or even months after parturition, and when this is the case the child is almost sure to suffer from indigestion and diarrhœa. The milk may be unwholesome because the nurse is in bad health, or because her diet is not properly regulated. That the diet of the nurse affects her milk, we have no doubt, though it has been denied by some persons.

We have known several children to suffer from indigestion, attended with vomiting, acid secretions, colic, and diarrhœa, in consequence of the nurse having indulged in a very rich diet, and particularly in vegetables and fruits. We do not mean to assert that all nursing-women should abstain from fruits, or even live on a very simple diet, for we have known some who could make use of the richest food, and eat abundantly of all kinds of vegetables and fruits, without the least injury to the child. But there are others who cannot do so without occasioning indigestion in their infants, because, probably, their children are unusually susceptible to the action of the materials absorbed from that kind of food. Again, it is clearly proved by recorded cases and by the opinions of various authorities, that the milk of the nurse is affected by her moral condition. Children have been known to suffer greatly, and even to die, from taking the milk of a nurse who had just before undergone a fit of violent anger. The depressing moral emotions, as anxiety, grief, fear, and despair, are well known to affect the milk secretion in such a way as sometimes to occasion indigestion.

The use of artificial diet for young infants, or as the expression is, "bringing up on hand or the bottle," is, we believe, by far the most frequent cause of indigestion during infancy. Very many children with whom this is attempted die of indigestions, chronic diarrhœa, gastritis, entero-colitis, cholera infantum, and thrush. Very few escape frequent attacks of one or other of the diseases just named. Much depends, no doubt, on the selection and preparation of the food. It may be stated as a well established fact, that a diet consisting wholly or in a great part of farinaceous substances, very rarely fails to disagree with the child, and to produce indigestion and other disorders of the digestive system, which

often prove fatal; while one in which cow's or goat's milk enters as the principal ingredient, though inferior to the natural aliment, and often productive of indigestion, is far less injurious than the one before spoken of.

A third cause of indigestion was stated to be the absence or loss of the digestive power of the stomach, independent of the nature of the food. This is a condition similar to the dyspepsia of the adult. It may be congenital or may result from causes brought into action after birth. It often remains as a consequence of previous indigestions from improper or excessive feeding. It exists during the invasion, course, and convalescence of various diseases. Dentition frequently diminishes or impairs the tone of the digestive function, so that the child is often unable, during that process, to digest aliment which had agreed with it perfectly well at other times.

The causes of indigestion after the completion of the first dentition, are congenital feebleness of the digestive function; a certain want of power of that function, which remains often for years in children reared upon artificial diet, and in those who have been debilitated by frequent attacks of disease of any kind; the habitual use of improper diet; the eating of crude indigestible food; the process of the second dentition; the want of due exercise in the open air; residence in large cities; and undue exercise of the mental faculties in the conduct of the education of the child.

SYMPTOMS.—We shall describe first the symptoms of indigestion as it occurs during infancy, and secondly as it occurs during childhood, or after the completion of the first dentition.

Indigestion during infancy may be advantageously considered under two heads: as occasional or accidental, and as habitual. By the former we mean that which occurs in a healthy infant from a transient cause, such as repletion, or a momentarily unhealthy state of the nurse's milk, from some imprudence on her part as to diet, from some moral cause, or from sickness; and that which depends upon the passing influence of dentition. By habitual indigestion, we mean the form of the affection which is long continued in consequence of a persistence of the cause.

The symptoms of *occasional* or *accidental* indigestion in infants are: paleness and contraction of the face; restlessness and peevishness; moaning and crying, or in some cases, screaming; nausea, shown by excessive paleness, often by very great languor, and by occasional retching, which may either subside without vomiting, or, as more frequently happens, terminate in that act; flatulent distension and hardness of the abdomen, especially in the epigastric region, often accompanied with eructations; and in many of the cases simple diarrhœa. These symptoms usually come on soon after nursing freely, or after a very hearty meal of artificial food, in a child previously in good health. The attack seldom lasts more than a few hours or one or two days. The vomiting which almost always takes place, and which relieves the stomach from the offending cause, very often accomplishes the cure.

Habitual indigestion in infants causes a train of symptoms different from and much more severe than those just described. Of these the most important are: frequent attacks of nausea and vomiting, and of simple

diarrhœa repeated for days, weeks, or months in succession; paleness, or some other unhealthy tint of the cutaneous surface; continual restlessness and discomfort, with fretting or crying, particularly in the latter part of the day and during the evening and night, in place of the natural ease and quiet of a healthy infant; constant fits of the most violent screaming from colic, sometimes lasting for hours; dull and languid expression of the countenance, or else an uneasy, contracted look, like that produced by continued suffering; more or less emaciation; failure of the natural growth in stature and size, so that the child is small and puny for its age; want of calorific power, causing the child to suffer unusually from cold, as shown by frequent coolness of the hands and feet; irregular appetite, which makes it necessary to tempt it by frequent changes of the food, or more or less complete anorexia; and lastly, the various symptoms that indicate an impoverished state of the blood and bad nutrition.

In some cases there are added to the above symptoms, or there follow as a consequence of the indigestion, those of gastritis or entero-colitis, to be hereafter described. Indigestion probably seldom proves fatal in infants, except from the occurrence of some inflammatory complication, as for instance, one of the diseases just named, or acute disease of some other principal organ.

Indigestion in children who have completed the first dentition may, as in the case of infants, be occasional or habitual. *Occasional indigestion* occurs in strong and vigorous, as well as in more delicate subjects. The attack generally begins, within a few hours or a day after the child has eaten some indigestible substance, with languor and chilliness in older children, and with languor and peevishness in those who are younger; after which there is headache, pain in the stomach in most of the cases, and very often a disposition to somnolence. If the child is attacked with vomiting soon after the appearance of these symptoms, and ejects the offending material, it will often seem perfectly well from that time. If, however, this does not take place, fever, sometimes of a violent character, is almost certain to make its appearance. The pulse becomes very frequent, rising to 120, 130, 160 or over; and being full and resisting; the skin becomes flushed, dry, and very hot; the appearance of the tongue is not generally changed early in the attack; there is considerable thirst; the child is restless and uneasy, tossing from side to side, or it lies in an uneasy sleep, attended with frequent starting and jerking of the limbs or crying out; the abdomen is natural, or hard and distended over the epigastric region. When the symptoms just described make their appearance suddenly, by which we mean in the course of a few hours, in a child two, three, four, or five years old, after it has eaten some indigestible substance, there is reason to fear an attack of convulsions. The probability of the occurrence of this accident is great in proportion to the earliness of the child's age, and the impressibility of its nervous system. The attack is particularly to be apprehended, and should be carefully guarded against, whenever the fever is violent, especially if the pulse runs very high, when there are urgent complaints of headache, when the restlessness and agitation are very great, or when there is somnolence, with frequent startings or

twitchings of the muscles. Convulsions sometimes occur without any previous warning, or after such slight signs of disorder as would fail to produce uneasiness in the parents or attendants.

The symptoms produced by occasional indigestion generally continue until nature relieves the stomach by vomiting or diarrhœa, or until the remedies proper in the case, the most important of which are evacuants, have been administered. It happens not unfrequently, that symptoms of gastric or intestinal disorder remain for some days after the violence of the attack has subsided, and in some instances the disturbance is so great as to occasion gastritis, entero-colitis, or dysentery.

Habitual indigestion in children who have completed the first dentition is not at all an uncommon affection. It is a condition analogous to the dyspepsia of the adult. The symptoms of this form are the following: The general appearance of the child is delicate, as shown by a pallid or sallow tint of the skin, instead of the ruddy complexion of health, by thinness and want of proper development of the limbs and trunk, and by softness and flaccidity of the muscular tissues. There is an habitual air of languor and listlessness, with absence of the usual gayety and disposition to play natural to the age, and the child often complains of being tired. The appetite is feeble or uncertain, being sometimes absent, and at other times too great; or it is peculiar, there being a willingness to eat of dainties, but a refusal of food of a simple character. The tongue presents nothing peculiar. It is, however, more frequently somewhat furred than clean and natural. The temper is usually irritable and uncertain. The child rarely sleeps well; on the contrary, the nights are restless and much disturbed, the sleep being broken and interrupted by turning and rolling, by moaning or crying out, and by grinding of the teeth. These latter symptoms, together with picking at the nose, which is a frequent accompaniment, are almost always referred by the parents and nurses to worms, and it is often impossible to convince them to the contrary, even though frequent and violent doses of vermifuges have failed to show the existence of entozoa. The state of the bowels is uncertain. In some instances they are very much constipated, requiring frequent doses of laxatives, or careful regulation of the diet, to keep them soluble; in other they are inclined to be loose, and when this happens the stools are often lenteric. In others, again, constipation and diarrhœa alternate. The abdomen is usually natural, or somewhat enlarged from flatulent distension; complaints of pain are not uncommon. This form of indigestion, like dyspepsia in the adult, is generally a very chronic affection, seldom lasting less than several weeks or months, and sometimes persisting for years.

DIAGNOSIS.—The occasional indigestion of infants is not likely to be mistaken for any other complaint. The suddenness of the attack, the character and quantity of the matters ejected from the stomach, the absence of symptoms indicating the invasion of any other disorder, the short duration of the symptoms, and the rapid recovery, all render the true nature of the case very clear. That which occurs in older children, on the contrary, is not so easy of diagnosis. In many cases the invasion is

not unlike that of scarlet fever. The vomiting, the rapidity of the pulse, the great heat of the skin, and in some cases a certain suffusion of the integument dependent on the activity of the circulation, all render the case doubtful for some hours, or for a day, after which time the difficulty ceases, from the development of the symptoms peculiar to the disorder. We believe that, quite frequently, cases of simple sore throat from cold, or mild forms of diphtheria, or severe ones in the early stage, are mistaken for indigestion. They are referred to and explained in popular language as "bilious attacks." The lassitude, the occasional vomiting, the want of appetite, the more or less decided febrile movement, are explained on the theory of gastric disorder. So true is this that we have formed the habit ourselves, and recommend it to others, of always looking into the throat when a child who cannot speak, or is too young to describe his own sensations, exhibits a train of symptoms pointing to digestive disturbance. This is the only way, often, of making a correct diagnosis. The absence of any apparent pain or difficulty in swallowing goes with us for nothing. We deem it wise to look into the throat. The diagnosis of indigestion accompanied by convulsions will be considered in the article on the latter affection.

The habitual indigestion of infants is not likely to be confounded with any other disease. The absence of fever, of tenderness of the abdomen on pressure, or other acute symptoms, all indicate the dependence of the disorder on functional distress of the stomach. The same remarks apply to this form of the disease occurring in older children. Nevertheless, the practitioner should never neglect to make a careful examination, both of the physical and rational signs, of all the important organs of the body, as it sometimes happens that latent disease of some one of them is the cause of the gastric difficulty.

PROGNOSIS.—The prognosis of occasional indigestion is nearly always favorable. It is rarely a dangerous disorder, unless accompanied by convulsions or some other sign of violent disturbance of the nervous system. Under the latter circumstances the prognosis should be very cautious, as the termination is not unfrequently fatal in consequence of injury done to the nervous centres. It should be recollected also that this form of indigestion sometimes becomes the exciting cause of inflammation of the stomach or intestines, in which event the prognosis will be that of those diseases.

Habitual indigestion in infants is a serious complaint, and ought always to awaken the solicitude both of the physician and parents; for though a simple functional disease of the stomach is probably not often fatal, it is exceedingly apt to prove so by the introduction of gastritis, chronic enteritis, entero-colitis, or thrush, or by its laying the system open to other diseases, and rendering it less able to withstand them should they happen to occur. In older children it is not, according to our experience, so dangerous a malady. We have never, as yet, seen it terminate fatally.

TREATMENT.—The treatment of *occasional indigestion* in infants ought to be very simple. The child has generally relieved itself by vomiting before the physician is called. If, however, it continues pale and languid, with vomiting or retching, after the stomach seems to have been emptied,

the proper plan is to make use of remedies to calm the irritability of that organ. This can almost always be accomplished by giving a teaspoonful every ten or fifteen minutes of a mixture of lime-water and milk, consisting of one-third milk to two-thirds lime-water, or of equal proportions of each, or the same doses of a mixture consisting of equal parts of lime-water and cinnamon-water. At the same time a small mustard-plaster, weakened with wheat flour, or flannels wrung out of hot brandy and water, may be applied to the epigastrium, or a warm Indian mush poultice, in a flannel bag, laid over the whole abdomen. Should these means fail to relieve the sickness, from half a drop to a drop of laudanum, or ten drops of paregoric, may be administered, and repeated, if necessary, in two hours. The child generally recovers its usual health after the sickness has entirely ceased. If, however, it remain fretful and uneasy, if it cry much as though in pain, it is probable that a portion of aliment has passed, in a partially or wholly undigested state, into the intestines. The suspicion will be confirmed if the abdomen is found, upon palpation and percussion, to be swelled, hard, and resonant from flatulent collections in the bowels. Under these circumstances, a laxative ought to be given. The best dose is half a teaspoonful or a teaspoonful of castor oil, a teaspoonful of simple or spiced syrup of rhubarb, or, if there have been evidences of an acid state of the stomach, about a quarter of a teaspoonful of the best magnesia. If, however, the tongue is coated and the stomach irritable, it is better to allay this, and at the same time to promote secretion from the intestines, by the administration of the following powders:

R. Hydrargyri Chloridi Mitis, gr. $\frac{1}{2}$,
 Bismuthi Subnitratiss, gr. x.
 Sodii Bicarb., gr. v.

M. et. div. in chart. No. x.

Dose. One every two hours for a child two years old.

The *occasional indigestion* of older children demands a different and more energetic treatment. After ascertaining that the child had eaten something indigestible, we should inquire whether there has been vomiting. If there has been none, or if only slight, it will be proper to give an emetic immediately. The best one under the circumstances is ipecacuanha. It may be given either in powder or syrup. The dose is familiar to every one. If the ipecacuanha be not at hand, we may use a teaspoonful of powdered alum in honey or molasses, to be repeated, if necessary, in fifteen minutes. Alum is even less apt to fail than ipecacuanha. If the child continue unwell after the operation of the emetic, which is often the case, and particularly if the fever be considerable, a purgative should be given as soon as the stomach will bear it. The best dose is castor oil, which is the most speedy and least irritating. It may be given in orange-juice, which forms an excellent vehicle, or, if the child is old enough, in the froth of beer or porter. A teaspoonful is generally enough. If the oil cannot be taken, we may give infusion of senna and manna, the fluid extract of senna mixed with spiced syrup of rhubarb, syrup of rhubarb alone, magnesia, to be followed by lemonade, salts and magnesia, or

the former alone, or, lastly, a Seidlitz powder. If the fever continue, and the cathartic fail to operate in four or six hours, a purgative enema ought to be given to hasten its effect. A bath at about 96° or 97° will almost always be found useful in these cases. The child should be kept in the bath from eight to twelve or fifteen minutes. The only circumstances which form an objection to this remedy are the facts of the patient being so irritable, or so fearful of the water, as to make it necessary to contend with him in order to succeed in using it. When this is the case, it had better not be employed, and sponging with tepid water and spirit should be substituted. If the child complains of pain in the stomach, the application of a warm mush-poultice over the epigastrium or whole abdomen will be found of much service.

When in this form of indigestion the febrile reaction is violent, as it often is, and particularly when there are signs of great disturbance of the nervous system, consisting of excessive agitation, complaints of severe headache, drowsiness, moaning or crying out in the sleep, or twitching and jerking of the muscles, the physician should beware of a convulsive attack. In such cases as these, the patient ought to take a purgative dose of calomel (from two to three grains), or a dessertspoonful of castor oil, have a warm bath at once, and soon after an injection. The remedies ought to be prompt and energetic, for the case is pressing. A convulsion is always a dangerous event in childhood, and should be prevented if possible. If calomel has been given, a cathartic dose ought to be administered about two hours afterwards, in order to insure an action upon the bowels, and to carry the calomel out of the system. After the administration of the evacuant, bromide of potassium alone, or in combination with small doses of opium are invaluable. At four to five years of age, two and a half grains of the bromide, with one or two minims of laudanum, given every hour or two hours, of two, three, four, or more doses, or until the nervous phenomena are controlled, or sleep is induced, make the proper dose. In several cases in which the nervous symptoms have been very urgent, and where convulsions have occurred, we have known small doses of chloral hydrate, two grains at the age above mentioned, given two or three times, of singular efficacy in calming the threatening erethism of the nerve-centres. The diet should be absolute during the violent stages of the attack, and the usual diet is to be resumed only by degrees. The drinks may be plain water, or gum-water, taken cold.

It not unfrequently happens that occasional indigestion is followed by gastritis or enteritis, or by habitual indigestion lasting for weeks or even months. These different sequelæ must be treated according to the plan proper for each.

The *habitual indigestion* of both infants and older children requires a very different treatment from the occasional or accidental form. In both the indications are nearly the same. The most important are very careful regulation of the diet in all its details, the use of tonics and stimulants to restore tone and vigor to the digestive function, the employment of remedies to correct the state of the bowels, whether they be relaxed or consti-

pated, and attention to securing the child proper exercise, exposure to the air, and suitable clothing.

If the symptoms of the disorder occur in a child at the breast, the milk of the nurse should be carefully examined, in order to ascertain whether it be good. If found to possess any unhealthy qualities, the nurse ought to be changed at once. Attention to this point alone will almost certainly cure the child. It needs no other remedy.

If the patient is fed wholly or in part, it is essential to regulate the diet to suit the state of the digestive function. Milk ought in all cases to form the basis of the food, unless it has been proven by patient trial to be absolutely repugnant to the stomach. We have often found that infants who had been thought quite incapable of digesting cow's milk, could do so very readily when it was very much weakened with water. The usual proportion for an infant of a few months old, are half and half, or two parts milk for one of water. When these are found to disagree, it is well to try three, or even four or five parts of water to one of milk, and if the stomach digest this, as it often will, the proportion of milk may be slowly and cautiously increased to the usual standard. If we conclude that milk cannot be digested by the child, it is best to try cream. Of this one part to three or four of water may be given. Some infants of six or eight months old, it may be remarked, who cannot digest more than very small quantities of milk, will take and digest well very delicate broths made of chicken or mutton, or small quantities of the lightest meats, as mutton, chicken, or very tender beef, mixed up extremely fine, and given by teaspoonfuls.

In cases of this kind we have found the diet consisting of gelatin, milk, cream, and arrowroot, prepared in the manner directed in the article on food, to suit better than anything else. We have met with a number of children, whom it was necessary to feed to the amount of a pint or a pint and a half a day, in addition to their being nursed occasionally, who could take neither milk and water, cream and water, milk and arrowroot, oatmeal gruel, rice gruel, nor indeed anything that was tried, without vomiting, colic, and severe diarrhoea, who digested perfectly well and thrived admirably upon the preparation alluded to. We have used it during many years, and have recommended it for a great many children, and do not hesitate to say that it agrees with a larger number than any diet we have employed or seen employed.

The diet of older children laboring under chronic weakness of the digestive function is as important as that of infants. Two chief ends should always be borne in mind in selecting it, digestibility and nutritiousness. The former is all-important, for without it, the stomach, constantly irritated by improper food, has no chance of regaining its tone, while the latter is necessary in order to sustain the strength of the child, and allow it to carry on its growth. We have generally found it most prudent, and often really necessary to specify as to the substances to be given at each meal. The morning and evening meal ought to consist of bread and milk, mush and milk, or of milk, warm water and sugar (called in this country children's or cambric tea), and bread and butter, and nothing else in most

of the cases. It is sometimes proper to allow a soft-boiled egg, particularly if the child be very fond of it. The dinner ought to consist of light broths containing rice, with bread or toast, or of the plain meats, as mutton, beef, chicken, turkey, birds, or fine game. No vegetable ought to be allowed in most of the cases except rice, as all others, even the potato, are very apt to disagree. We believe that the potato is more digestible when roasted than when boiled. If the child require anything between breakfast and dinner, it may have what is allowed at breakfast, or dry bread and nothing else. There are various articles of diet which should be absolutely forbidden, amongst which are hot and sweet cakes, and hot bread of all kinds; sausages, not unfrequently given to children in this country; corn-beef, ham, veal, pork, goose, ducks, fish; all manner of dessert, excepting rice-pudding, or curds-and-whey, often called junket; sweetmeats, candies, fruits, except some of our finest summer ones; and to conclude, everything which long observation and experience have shown to be unsuitable to a dyspeptic stomach.

It is sometimes very difficult to find anything to agree well with the child. In one case of a child three years old that came under our observation, neither milk, bread, nor meat could be taken. The caseine of milk seemed to be absolutely indigestible, as it would be rejected from the stomach many hours, or even a day or two, after the milk had been taken, in the form of masses of dry, fibrous cheese, of an oblong shape, nearly or quite as large as a peach-stone. After trying various articles, we found that the child digested raw oysters, soda-biscuit, and rennet-whey, and upon these articles alone she lived for two weeks, at the end of which time she had improved so much as to be able to take the white meat of chicken very finely minced. She gradually regained her previous health.

After regulating the diet, such remedies as tend to invigorate the digestive functions ought to be prescribed. The most important of these are the vegetable and mineral tonics, and mild stimulants. We have found quinine, iron, and small quantities of port wine or brandy, to succeed better than anything else. To a child under two years old, from a quarter to half a grain of quinine, and to one over that age, a grain may be given three times a day, and continued for two, three, or four weeks. It is most readily given to young children diffused, without being dissolved, in a mixture of equal parts of syrup of gum and ginger, or in syrup of red oranges, or, what is probably the best of all, elixir of liquorice; while to those who are older it may be administered in pill. Of the preparations of iron we prefer the wine, the syrup of the iodide, or the ferrum redactum. The wine of iron is best given in Dr Erasmus Wilson's formula, consisting of syrup of tolu and caraway water. At the age of six months, from 10 to 15 drops; at two years, 20 to 30 drops, should be given three times a day. Of the syrup of the iodide, 1 drop for infants, 2 to 4 drops for older children, are used three times a day. Of the metallic iron, $\frac{1}{4}$ of a grain for infants, and $\frac{1}{2}$ a grain for older children, is the proper dose, three times a day. It may be mixed with sugar and dropped upon the tongue, or made into a lozenge with chocolate. When

there is any suspicion of a scrofulous taint in the child's constitution, or when it is disposed to have chronic irritations, excoriations, or ulcerations of the nostrils, otorrhœa, or papules or pustules about the eyelids or other parts of the body, it is useful to give the wine of iron mixture, with from $\frac{1}{2}$ a minim to 1 minim, according to the age, of Fowler's solution of arsenic added to each dose, three times a day, directly after food. Under these circumstances, and particularly when the dyspeptic condition is accompanied with frequent nausea or occasional vomiting, with frontal headache, and with constipation, seeming to indicate a disposition to tubercular deposit in the system, we have found cod-liver oil the most efficient of all the remedies that we have tried. It has often removed with great rapidity the dyspeptic symptoms, invigorated the general health, and, in fact, restored the patient to health. The dose is from half a teaspoonful to a teaspoonful twice or three times a day, at the age of six or eight years. It is best taken in a small quantity of malt liquor, or floating on strong mint-water, or syrup of ginger. In very young children, and in older ones also, when the latter refuse to take it in the ordinary methods, the following formula for its administration will be found one of the best:

R. Ol. Jec. Aselli,	f℥ss.
Pulv. Acacia,	q. s.
Ol. Cinnamomi, vel Ol. Gaultheriæ,	gtt. vj.
Sacch. Alb.,	q. s.
Aq. Cinnamomi,	q. s. ad f℥ij.—M.

Dose—A dessertspoonful three times a day, after eating.

The recent introduction of the use of pepsin in the treatment of disorders which, like the one under consideration, are characterized by a want of digestive power, is a valuable improvement in their management. It is nearly always well received by the stomach, and in many cases will enable the child to take and digest the proper amount of suitable food, which before would have caused evidences of gastric embarrassment, with the rejection of a considerable part of the meal in an undigested state by vomiting or stool.

Pepsin may be administered in the form of powder,—the best preparation of which is that now sold under the name of saccharated pepsin,—and the proper dose of which for a young child is two or three grains taken immediately after meals. Or we may use the liquor pepsinæ, which is a solution of this substance in glycerin and water, acidulated with muriatic acid. The proper dose of this latter preparation is from ℥xx to f℥ss., taken diluted with a little water, also directly after meals.

The combination of small doses of muriatic acid is unquestionably of advantage in increasing the digestive power of the stomach. We have thus found the following mixture of much service in the chronic indigestion of children:

R. Acid. Muriatici Dil.,	gtt. xxv.
Liq. Pepsinæ,	
Elix. Calisayæ, aa,	f℥j.—M.

Dose—A half teaspoonful to a teaspoonful, according to the age of the child.

In connection with these remedies, a little port wine or brandy may be allowed twice or three times a day, or at dinner only. To young children, one or two teaspoonfuls of brandy may be given in the course of the day, mixed in water or, better still, in milk; of the port wine, from a teaspoonful to a tablespoonful, according to the age and strength of the patient, may be repeated morning, noon, and night. It may be well for us to say that we do not approve of the daily use of this form of stimulants for children over six or eight years of age. We once knew a boy, ten years of age, to become so fond of his port wine as to purloin it from the pantry. If young children must have such tonics after the age above indicated, we believe the French system of allowing claret, or the German one of allowing light beer, to be the best and safest.

If the bowels are inclined to constipation, they should be kept soluble by laxative enemata, and by the use of rhubarb or aloes; when relaxed, the frequency of the discharges may be controlled by chalk mixture, by anodyne enemata given once or twice a day, by the aromatic syrup of galls (to be described under the head of entero-colitis), or by some of the astringents in common use.

In cases where the evidences of a catarrhal state of the mucous membrane of the stomach and intestine are present, we have frequently found excellent results to follow the administration of small doses of nitrate of silver (gr. $\frac{1}{30}$ to $\frac{1}{15}$) given in solution in thin syrup of acacia two or three times a day.

In all cases of chronic indigestion in children, it ought to be regarded as an essential part of the treatment to secure to the patient a proper amount of exercise in the open air. In summer the child should pass several hours of every day in the air. It ought, indeed, if the heat of the sun can be avoided by proper shade, to pass the whole day in this way. In winter it is, of course, impossible to carry this system to the same extent, but the child should nevertheless be taken out at least once a day; this may be done in the coldest, and even in damp weather, if sufficient clothing be worn. If a child comes back from a walk with warm limbs, and with its cheeks in a glow, there is little danger of cold. The quantity of clothing must depend on the constitution and idiosyncrasy of the patient. Some need twice as much as others. The proper amount is best determined by the temperature and coloration of the surface after a walk.

ARTICLE II.

SIMPLE DIARRHŒA.

UNDER this title we shall describe a mild form of diarrhœa to which children are very subject, in which the pathological condition appears to be one of mere functional disorder, or of very moderate hyperæmia or catarrh of the intestinal mucous membrane. We might, indeed, assume

with some, that the disorder is at all times one of mild catarrh of the bowels, but we deem it best, in a practical point of view, to consider it as being sometimes one of functional disturbance only, since many observers of high authority declare that they meet with cases of even fatal diarrhœa in which no anatomical alterations are found after death, and since we ourselves have met with so many cases in practice which follow a different course in symptomatology, duration, and their effects upon the constitution, from the form of disease which we shall treat of as entero-colitis or inflammatory diarrhœa.

CAUSES.—The causes of the disease during infancy are *unfavorable hygienic conditions*, as the habitation of unwholesome, ill-ventilated, damp, and filthy dwellings, or of contracted and crowded quarters of cities and towns; an *unhealthy state of the milk* of the nurse; the use of *artificial diet* at too early an age, especially that of an improper kind; *cold*; *dentition*; and lastly, great *atmospheric heats*. The most important of these are improper alimentation, by which we mean the use of artificial diet, and particularly one consisting chiefly of farinaceous substances to the exclusion of a proper amount of milk, and dentition. For a fuller account of the influence of these different circumstances on the digestive organs of children, the reader is referred to the remarks on food, the causes of entero-colitis, and to the article on thrush.

The chief causes of the disease after the first dentition are, according to our experience: the habitual use of *improper food*; the *loss* of digestive power, which often follows a severe indigestion, or an attack of some acute disease; the *debility of constitution* which attends sudden and rapid growth; the *want* of proper exercise and exposure to the air; the predisposition which exists in some children from *hereditary* causes; and the disturbing influence of the second dentition.

The system of indiscriminate diet allowed to children in this country is, it seems to us, a fruitful cause of gastric and intestinal complaints. We believe that, as a general rule, children over two and three years of age, are allowed amongst us to eat of the food prepared for the older members of the family. Now, any one who will reflect upon the variety of dishes habitually placed upon an American table, ought not to be surprised to see children permitted a choice amidst such profusion, pale, thin, delicate, exposed to frequent indigestions, attacks of diarrhœa and entero-colitis, to gastric fevers, and the host of minor ills attendant upon feeble digestive powers. We are acquainted with some families in this city, the children of which, from the age of two years, are allowed habitually to breakfast upon hot rolls and butter, hot buckwheat cakes, hot Indian cakes, rice cakes, sausages, salt fish, ham, or dried beef, and coffee or tea; and to dine upon a choice of various meats and a great variety of vegetables, which latter they often prefer to the exclusion of meat, and then to make a rich dessert of pies, puddings, preserves, or fruits; and lastly, to make an evening meal of tea and bread and butter, almost always relished, as the term is, with preserves, stewed fruits, hot cakes of some kind, or with radishes, cucumbers, or some similar dish. Add to such meals as the above, the eating between whiles of all kinds of candies and comfits, which many

children here regularly expect in larger or smaller quantity, cakes both rich and plain, fruits to excess and at all hours, from soon after breakfast to just before going to bed, raisins and almonds, and nuts of various kinds, and the wonder is, not that we are a pale, thin, dyspeptic and anxious-looking race of people, compared with Europeans, but that we have any health at all, when our children are allowed to make use of the indiscriminate and unwholesome diet just described. Such a system undoubtedly occasions frequent attacks of the disease under consideration, and unless the diet be changed early in the attack, it is very apt to become chronic. It has been stated that simple diarrhœa sometimes followed as a consequence of indigestion. We have known such a result to occur in children previously in fine health, and to continue for several weeks or months. In these instances, the disorder appears to depend in good measure on the loss of the digestive power of the stomach. This seems proved by the great influence which the character of the food has upon the malady, which is always aggravated by the use of any articles except those universally acknowledged to be the most digestible, and also by the frequent coëxistence of lientery when the food is not of the lightest kind.

We have several times met with cases which we could ascribe to no other cause than debility and want of power of the digestive organs, dependent upon too rapid growth. That sudden and rapid growth may produce feeble digestion, or, in other words, a dyspeptic state, is, in our opinion, proved by the following consideration. It is attended with loss of appetite, emaciation, paleness, languor, and weakness, and frequent attacks of diarrhœa, or a chronic form of that disorder; all of which symptoms are greatly influenced by the regimen of the child, and are most readily removed by attention to that point, and by the use of tonics and stimulants.

The other causes enumerated need but little comment. We will merely remark that we have several times observed a predisposition to weakness of the digestive organs, transmitted apparently from parent to child. As to the influence of the second dentition, we have no doubt that it is a frequent cause of the complaint, and we believe that it is too little attended to by practitioners.

ANATOMICAL APPEARANCES.—It has already been stated that we look upon this disorder as one of purely functional disturbance in many instances. We are led to take this view by the fact that it is so often unattended by any of the ordinary signs of inflammatory action, and because some very competent observers affirm that they have failed to find in a certain proportion of cases of fatal diarrhœa, any lesions appreciable to the senses. Thus, M. Billard says (*Mal. des Enfants*, p. 392): "Many children at the breast have diarrhœa without enteritis; they lose color, become etiolated, fall into a state of marasmus, and yet at the autopsy not a trace of inflammation of the intestines is found." M. Bertin (*Mal. des Enfants*, 2ème ed., p. 574) states that of fifty-seven cases of gastro-intestinal disease observed by himself, there were four in which not a trace of inflammation, or any other appreciable lesion of the alimentary tract, could be found. MM. Rilliet and Barthez, in their first edition (t. i, p. 491), assert

that is about every twelve children affected with more or less abundant diarrhœa, and in whom we might expect to find colitis, there will be one in whom the gastro-intestinal tract will be found in a state of perfect integrity. They add that this conclusion is deduced from a comparison of nearly three hundred autopsies. We do not find this statement given in their second edition, but we do find there (t. i, p. 693) the following paragraph: "Quite frequently, especially in early infancy, in cases in which the symptoms have pointed to some disease of the gastro-intestinal tube, an autopsy reveals no lesion of the solids, or only changes of minimum importance. The secretions alone are vitiated." One must suppose, therefore, that the class of cases which we describe as simple diarrhœa, are sometimes quite independent of any anatomical changes in the tissues, recognizable by our ordinary methods of examination, or that those changes are so slight and so evanescent as to disappear after death; or that they are those only of the mildest forms of catarrhal inflammation. It is not unlikely, it seems to us, that further and more minute investigation, especially with the microscope, will reveal tissue-changes which are not discoverable by the unassisted senses.

When the anatomical changes, constituting the catarrhal state, are found in children who presented during life the symptoms of simple diarrhœa, they will be such as our described by Niemeyer in the following passage: "Catarrh rarely affects the entire intestinal canal. It is most frequent in the large intestine, less so in ileum, and rarest in the jejunum and duodenum. The anatomical changes left in the cadaver by acute catarrh, are sometimes pale, at others dark redness, swelling, relaxation, and friability of the mucous membrane, which is sometimes diffuse, at others limited to the vicinity of the solitary glands and of Peyer's patches, and a serious infiltration of the submucous tissue. Occasionally, after death, the injection has entirely disappeared, and the mucous membrane appears pale and bloodless. Swelling of the solitary glands and glands of Peyer is an almost constant appearance; they distinctly project above the surface of the mucous membrane. The mesenteric glands also are usually found hyperæmic and somewhat enlarged. The contents of the intestines consist at first of plentiful serous fluid, mixed with detached epithelial and young cells, subsequently of a cloudy mucus, which is adherent to the wall of the intestine, and contains epithelial structures."

The best description that we are acquainted with of the anatomical appearances found in the intestines in fatal cases of diarrhœa, not in children, to be sure, but in adults, is that given by Dr. Woodward in his work on *Camp Diseases* (Philadelphia, 1863). In that work (page 216), under the head of simple diarrhœa, he says that this form of diarrhœa it to be regarded as the result, usually, of irritation of the intestinal mucous membrane, produced by the ingestion of improper food, or other causes mentioned, and as expressing itself in increased secretion throughout the intestinal tract. The irritation, he goes on to say, may even amount to inflammation. Opportunities for post-mortem examination occur but rarely. "They reveal little that bears on the nature of the disease, except congestion of the intestinal vessels of variable intensity." At page

246 will be found a description of the histology of the intestinal lesion in chronic diarrhœa, including the changes observed in specimens but moderately diseased, which latter would probably be the analogue of what we might expect to find in the simple diarrhœa of children we are now describing. We must refer the reader to the work itself, as the passage is too long to be quoted in full here: but we cannot help thinking that Dr. Woodward's descriptions would apply also to the changes induced in children by like causes, and leading to similar forms of disease.

SYMPTOMS.—We shall describe first the symptoms of simple diarrhœa in infants, and afterwards those which characterize the disorder in older children. In infants the appearance of the diarrhœa is usually preceded or accompanied by slight disturbance of the *temper* and *comfort* of the child. There is some degree of *restlessness*, *peevishness*, and disposition to cry; the child sleeps less than usual, and often starts and moans during sleep; all of which symptoms are more marked, as is the case indeed in nearly all the ailments of children, during the night. Though the symptoms described are observed from time to time, and particularly during the night, they are not always present, as the infant will occasionally through the day seem perfectly well and comfortable, with the exception, perhaps, of slight paleness and languor, almost always perceptible upon its countenance. There is no *fever* in these cases, or at least nothing more than unusual warmth of the hands, feet, and abdomen at night. If a marked febrile reaction take place, there would be reason to suspect the existence of some degree of entero-colitis. The *mouth* often becomes, after a few days, a little warmer and less moist than usual; the *tongue* is generally moist and only slightly coated; and the *appetite* is commonly diminished, as shown by the child's nursing with less eagerness and at longer intervals than before. In very mild cases the *stools* are at first, and sometimes throughout the attack, feculent; the only deviation from their ordinary character is that they are more frequent, thinner, more copious than usual, and that the odor is changed so as to become acrid and offensive. In severe cases, they contain less feculent matter, become yet more fluid and sometimes watery, and exhibit small particles of a greenish color scattered through them; or the whole of the discharge is of a deep-green color, and is intermixed with portions of mucus. In many the cases, whitish lumps, evidently consisting of undigested curd, are observed mixed with the other substances upon the napkin. The number of stools varies from two, three, or four, to six or eight in the twenty-four hours. The number last mentioned is seldom exceeded, so long as the diarrhœa remains simple. The *abdomen* is seldom distended or painful to the touch. The *general appearance* of the child almost always shows the effects of the malady upon the constitution after a few days. The countenance becomes paler and thinner; the eyes look somewhat hollow; the edges of the orbits are more defined, and often present a pale-bluish circle; slight emaciation takes place, and the flesh of the child becomes softer and more relaxed than before the attack. The *duration* of the disorder is generally short, as it seldom lasts more than three or four days or a week. It may terminate in complete restoration to health, without having exposed the life of the child

to danger; or, if the causes which gave rise to it continue in action, if the child is of delicate constitution or the treatment not correct, and especially if this is of too perturbing a character, it is very apt to run into enterocolitis and expose the patient to all the dangers of that disease.

In older children (after the first dentition), the disease is much less frequent than in infants, and presents a different train of symptoms. Often it is nothing more than a slight disorder of the bowels, amounting to three, four, or five stools, thinner and more abundant than usual, accompanied by slight colicky pains, and unattended by fever or other signs of sickness, which, after continuing one, two, or three days, ceases, and the child regains its usual health. Some children are particularly liable to these attacks, and suffer from them every few weeks, or after any indiscretion in diet; whilst in others they are rare, let the diet be what it may.

There is another form of simple diarrhœa, however, of which we have seen a number of cases, much more troublesome than the one just described. It occurs in children from two and a half to seven and eight years of age, lasts a considerably longer time, and is much less under the control of remedial measures. This form of the disease has never, in the cases that we have seen, been accompanied by fever, or by any constitutional symptoms rendering it necessary to confine the child either to the bed or house. The only symptoms besides the diarrhœa which we have observed, have been some degree of paleness and moderate emaciation; slight weakness, shown by an indisposition on the part of the child to play with its usual spirit, by an inclination to lie about from time to time through the day on the sofa or floor, and by complaints of "being tired;" irritability of temper and peevishness; irregular appetite; picking of the nose; and restless, disturbed sleep at night, attended with moaning, crying, starting, and grinding of the teeth; all of which symptoms generally convince the mother that the child is suffering from worms. The abdomen is sometimes slightly tumid, but remains natural as to tension, and is not painful on pressure. There is no pain except slight colic in some cases. The stools have generally numbered from three to five, and in a few cases as many as six or eight a day. They are semi-fluid in consistence, often of a very offensive odor, and consist usually of feculent matter, which is sometimes clay-colored, more frequently dark brown, and, in other instances, deep yellow or orange in color. They are often also of a frothy character. In some of the cases that we have seen, there was lientery whenever the aliment was otherwise than of the lightest and most digestible kind. In all, the diarrhœa was evidently greatly influenced by the diet, showing, it appeared to us, a manifest dependence of the malady upon the condition of the stomach, which seemed to have lost to a great degree its digestive power.

The *course* of the disease in this form is variable. In some it lasts a few weeks, and then, under the influence of diet and remedies, ceases, to recur and run the same course after a short period. In others it may last a much longer time in spite of all treatment that we may use. We have known it to thus continue between three and four months, with occasional slight remissions, brought about apparently by remedies which a day or two after would lose their effect.

DIAGNOSIS.—The diagnosis of simple diarrhœa will rarely present any difficulties, since there is nothing with which it could be confounded, except the diarrhœa from tubercular ulceration of the bowels, or enterocolitis. From the former it is to be distinguished by the history of the case, and by the signs of tuberculosis in other parts of the economy; from the latter, by the absence of signs of inflammatory action.

PROGNOSIS.—The prognosis is favorable so long as the disease remains simple. The physician should never forget, however, the disposition which is inherent in it to pass into enterocolitis, nor fail to make the possible occurrence of this transition one element in his prognosis. During infancy it is always more serious than after that period, from the feebler power of resistance on the part of the constitution at that age to disease, which undoubtedly allows this simple affection to prove fatal in some instances, probably from the shock to the nervous system. After infancy it is rarely a dangerous disorder, both because of the greater stamina existing at that age, and from the fact that the disposition to extension of disease is less strong.

TREATMENT.—The *prophylactic management* of simple diarrhœa is the same as that which is proper for enterocolitis, and as that affection will be treated of at considerable length in a future article, we must on account of our limited space refer the reader there for information on this point.

After the disease is established, the treatment must consist first in attention to the *diet, exercise*, and state of the *gums* of the child. In many cases, careful regulation of the diet and exercise, and lancing the gums when they are much distended and vascular from the pressure of the advancing teeth, will suffice to arrest the disorder in a few days, without the necessity of resorting to drugs, which ought certainly to be avoided whenever it is possible to do so. If the child is at the breast, we must ascertain whether the milk of the nurse is good, by inquiry as to its appearance, specific gravity, reaction, and by examination with the microscope, and by reference to her health, diet, temper, etc., all of which circumstances more or less affect the mammary secretion. If we conclude that the milk is good, or that it has been disturbed in its healthy properties only by a transient cause, the child must be continued at the breast, with the precaution, however, of not allowing it to nurse quite so much as usual. An infant suffering from any kind of diarrhœa, had better be restricted entirely to the breast, unless it be clear that the supply of milk is quite insufficient. If we determine that the milk is unhealthy, the nurse must either be changed, or the child weaned; of course the former alternative is infinitely preferable if the child is under a year old, or even under eighteen months, if it seem to have a rather delicate constitution.

If the case occur in a child already weaned, or in one fed partly on artificial diet, the regulation of the kind, preparation, and quantity of aliment is of the utmost consequence. It ought to consist chiefly of milk or cream diluted with water, unless it has been clearly shown by previous trial that these articles do not agree with the child. We prefer as a

general rule, the food made from cow's milk, cream, arrowroot, and gelatine, in the manner described in the chapter on food. The proportions of the milk, cream, and arrowroot must vary with the age and digestive power of the patient. As a general principle, during the existence of diarrhœa, or at least in the early stage of it, and before the strength has been reduced by the disorder, the proportions of cream and milk ought to be somewhat less than in health. Not only so, but the total quantity of food in the day should be diminished, unless the ordinary amount seems to be really necessary for the maintenance of the strength. If it be found, after patient trial, that the child will not take or does not digest this kind of food, we may try arrowroot, rice-water, or barley, with a little cream, or thin gruel or panada, with a small proportion of milk or cream, alternated with very carefully prepared chicken or mutton water. If the child is six or eight months old, it often suits well to allow it a piece of juicy beef or a chicken-bone to suck, or from one to several teaspoonfuls of meat of chicken minced very fine.

For older children with a common attack of simple diarrhœa, the diet should consist for a few days of boiled milk with bread, of gruels made of boiled milk and arrowroot, rice-flower, sago, tapioca, or common wheat-flour, and of small quantities of light broths. Meats are, for the time, improper, and all vegetables, with the exception of rice, yet worse.

In the case of infants it is best to recommend a continuation of the ordinary exercise, unless the weather be cold and damp. Indeed, in good weather, exposure to the air and proper insolation are more important during the existence of this disorder than even during health. The same remarks apply to older children, with the exception that they ought not to be allowed to fatigue themselves, particularly in warm weather, as this tends to aggravate the complaint.

When the disorder occurs in a teething child, the gums ought always to be examined by the physician, and if found swollen, vascular, of a deep-red color, and hot, with the outline of the advancing tooth perceptible, they should be freely incised to the tooth. If, on the contrary, the tooth is too deep to be felt, and yet the gum is red and swelled, we would advise only a slight superficial scarification in order to relieve the tension.

The *therapeutical management* of the disease should be as simple as possible. The fewer drugs we can succeed with in the gastro-intestinal complaints of infants and children, the better. When, however, the diarrhœa continues for some days in spite of attention to the points already mentioned, and earlier if the discharges are either large, frequent, very watery or weakening to the child, we must resort to some of the means which have been found most useful in checking the inordinate action of the bowels. The most important are a careful employment of laxatives, and the use of opiates and astringents. Formerly we generally commenced the treatment by the exhibition of a teaspoonful of castor oil, containing from half a drop to a drop of laudanum for young infants, and two drops for older children; but of late years we have usually preferred the spiced syrup of rhubarb, in a teaspoonful dose, with laudanum, as above recommended. Castor oil sometimes purges more than we like; rhubarb rarely does so.

These doses given for two evenings in succession have oftentimes sufficed to effect the cure. Dr. West recommends very highly in cases of simple diarrhœa, in which the evacuations, though watery, are fecal, and contain little mucus and no blood, small doses of the sulphate of magnesia and tincture of rhubarb. His formula at one year of age is as follows:

R. Magnesii Sulphat.,	3j.
Tinct. Rhei,	f3 ij.
Syr. Zingiber.,	f3 j.
Aquæ Carui,	f3 ix.—M.
Dose—A teaspoonful.	

We often use with excellent effect the sulphate of magnesia, with laudanum, as follows:

R. Magnes. Sulphat.,	3j.
Tr. Opii Deodorat.,	gtt. xij.
Syrupi Simp.,	f3 ss.
Aquæ Menth., vel Cinnamomi,	f3 ijss.—M.

Dose—At one or two years a teaspoonful every two or three hours. For older children, the proportion of the magnesia and laudanum should be doubled.

If the diarrhœa persists after these means have been used for two or three days, or gets rapidly worse, we must resort to some of the astringents. The one most commonly employed is the chalk mixture, which is officinal in our Pharmacopœia. A teaspoonful of this is to be given after each loose evacuation, or three or four times a day. If the case prove obstinate, it will be found useful to add to each dose of the chalk preparation a small quantity of laudanum or paregoric, or some astringent tincture, the best of which is the tincture of krameria. When the chalk mixture fails entirely, powdered crabs' eyes will sometimes succeed; or we may resort to the aromatic syrup of nutgalls. The formulæ and doses for both these remedies will be found in the article on entero-colitis. If the discharges are small and frequent, mixed with mucus and somewhat painful, it is well to use small opiate injections (from one to two drops of laudanum in a tablespoonful of starch-water for young infants, and from three to six drops in double that quantity for older children), or the use of Dover's powder in older children in combination with chalk or sugar of lead, will often succeed in arresting the disease. One of the most valuable astringents in the bowel affections of young children is bismuth, which we are much in the habit of giving in the form of subnitrate, in doses of from two to five grains, according to the age, from three to six times in the course of twenty-four hours. For further and more complete information in regard to astringents, we must refer the reader to the article on entero-colitis, where they will be fully discussed.

The chronic form of simple diarrhœa which we have attempted to describe, occurring in children who have completed the first dentition, has always proved difficult to manage. From the experience we have had, we believe that the best mode of treating it is by proper regulation of the

diet, and by the use of tonics and stimulants, and occasionally of opiates. We were led to adopt this plan in consequence of having failed entirely to control the symptoms by the treatment generally successful in simple diarrhœa, and by the opinion which we came at last to form, that the disease depended in great part on a loss of the digestive power of the stomach and duodenum. The diet must be adapted to the idiosyncrasies of the individual; what we should seek is such a one as will be easily digested by the patient, the materials of which shall not appear in the stools, and one which does not manifestly increase, if it fail to moderate, the frequency of the discharges. The one which we have found to succeed best consists of boiled milk with stale bread for breakfast and tea, and the most tender meats, as very fine beef, mutton, chicken, or birds, with rice, as the only vegetable, for dinner. If the child likes flour or rice pap, it may have either in place of the bread and milk. If it will take none of these, it may have milk, warm water and sugar, with bread; or well boiled mush with milk, or milk toast. Should it refuse the dinner recommended above, we may substitute thin soup, or some of the milk preparations. Raw meat, given in the manner recommended in the article on entero-colitis, should also be tried, and will at times prove very beneficial. Between meals it ought to be allowed nothing but dry bread. All rich food, dessert, fruits, all vegetables except rice, candies and comfits, all kinds of cake and hot bread, in fact everything except the articles which we have mentioned, or similar ones, ought to be rigidly, systematically, and perseveringly forbidden. Until this has been done for many days, or for several weeks, the disease has always, according to our experience, obstinately persisted.

We have already said that we have not found the ordinary remedies for simple diarrhœa to exert much effect upon this form of the disease. On the contrary, the treatment for dyspepsia, that is to say, a simple but nutritious diet, exercise, and the use of tonics and stimulants, has always removed it in a longer or shorter time. The tonics which we have employed are port wine, quinine, and iron. From a dessert to a tablespoonful of port wine was usually given in water three times a day, in connection with iron. The preparations of iron used were Vallet's mass, of which from half a grain to a grain was given in pill three times a day; the solution of iodide of iron in the dose of first one, and then from two to four drops, three times a day, or the solution of the nitrate of iron in the dose of from two to five drops, three times a day, in water, continued for one or two months. We have sometimes combined with each dose of the solution of iron a drop of laudanum, especially if there were pain; or the opiate might be given by injection every evening. The quinine was generally administered alone in the dose of a grain three times a day, for one, two, or three weeks. It has not, however, proved so useful as port wine and iron.

Another tonic which, of late years, we have found very useful in some cases of this kind, is one containing *nux vomica* and compound tincture of gentian, as follows:

R. Tr. Nucis Vomicæ	f℥ss.
Tr. Gentianæ Comp.,	f℥ij.
Syrupi Simp.,	f℥v.
Aquæ,	f℥ij.—M.

Dose.—A teaspoonful three times a day after meals, for children of three or four years of age.

Wine of pepsin, in half teaspoonful doses, three times a day, is also a good remedy in such cases, or we may use the powdered saccharated pepsin, in doses of two to five grains, taken soon after each meal.

We have also found in some obstinate cases excellent results to follow the administration of nitrate of silver (gr. $\frac{1}{10}$ th to gr $\frac{1}{2}$ th) given in solution in syrup of acacia or in pill, according to the age of the child.

In the case attended with all the symptoms usually thought to indicate worms, the use of wormseed oil was followed by the expulsion of several very large lumbricoides. The child did not recover, however, for some weeks afterwards, and not until he had taken port wine and quinine for a considerable period. In other cases in which the verminous symptoms were also strongly marked, and in which the same remedy was given, no worms were expelled.

SECTION II.

DISEASES OF THE STOMACH AND INTESTINES, ATTENDED WITH APPRECIABLE ANATOMICAL LESIONS.

ARTICLE I.

GASTRITIS.

GASTRITIS, in the sense in which the term was used some ten or twenty years since, viz., to express an individual and special inflammatory disease of the stomach, of common occurrence and of supposed great severity and importance in childhood, is now well known to be a rare affection. It is doubtful, indeed, whether it ever forms a special visceral inflammation, except in consequence of the direct application to the organ of some irritant substance, such as the mineral acids or arsenic, or, as Rilliet and Barthez found, in a few instances, certain remedial agents, as tartar-emetic, kermes mineral, and croton oil. In the form of catarrh, acute or chronic, of the mucous membrane, on the other hand, it is doubtless one of the most common affections of childhood, constituting an important element in a great many diseases, and especially in the severe forms of indigestion, in simple and inflammatory diarrhœa, in cholera infantum, and in many of the wasting diseases of childhood, which result from the use of improper artificial diet in infants, and of crude and indigestible articles of food in older children.

We had almost abandoned the plan followed in our former editions, of devoting a special chapter to this subject, but on further consideration, think it will be best to treat of it separately, since, as stated above, cases do occur in practice in which the stomach is the chief, if not the only seat of disease, and which can be properly designated and described only under the title of gastritis.

CAUSES.—It has already been stated that the most violent and typical cases of gastritis, as a distinct disease, are the result of the application to the organ of some special irritant, as the mineral acids, arsenic, boiling water, or of certain remedial agents, and particularly of tartar-emetic, kermes mineral, or croton oil. These latter agents, the drugs just mentioned, cannot produce this effect unless used in large doses, or when continued for too long a time. The quantities of the antimonial preparations formerly administered, were always thought by us to be dangerously large, and we were not at all surprised to find that MM. Rilliet and Barthez, from their experience in former years in the Children's Hospital in Paris, cited them as one of the causes of acute gastritis. In the *Journal für Kinderkrankheiten*, for the years 1859, 1860, and 1861, in the third, fourth, and fifth annual reports of the Public Institute for Children's Diseases of Vienna, by the Director, Dr. Luzinsky, may be found in the third report three cases, in the fourth three cases, and in the fifth two cases of gastritis caused by the accidental drinking of concentrated lye.

The milder forms of gastritis are vastly more common than the ones above referred to. They are generally associated with disturbances of the intestinal tract also, and constitute by far the majority of the cases which come under the observation of the physician. They are caused very generally by improper alimentation; by the same causes, indeed, as those which determine indigestion. In infants, an unhealthy state of the mother's or wet-nurse's milk, the use of too rich a preparation of cow's milk, milk obtained from an unhealthy cow, or a food composed of too large a proportion of farinaceous material, are the most common causes. In older children, an unwholesome meal, as a surfeit of cakes and candies, tough meats, unripe, or an excess of ripe fruits, the swallowing of a quantity of skins of grapes, of orange-peel, of the seeds of oranges, or such like imprudences or accidents, of all which we have seen examples, will sometimes occasion symptoms which we can refer only to acute catarrh of the stomach. In such cases the child may escape any serious consequences if it rejects, by vomiting, the improper food, soon after it has been taken. Or it may have an attack of cholera infantum or cholera morbus, and either recover its usual health in a short time, or pass through a longer or shorter illness, as the result of these disorders; or, lastly, the unhealthy food may be retained for a longer time than usual in the stomach, and acting as a local irritant on the gastric mucous membrane, may set up a true and more or less severe form of the disease we are considering.

ANATOMICAL APPEARANCES.—Death is so rare a consequence of gastritis alone, except in the form produced by the direct application of irritants to the organ (and even in such, recovery appears to be the rule, since all the eight cases referred to as reported by Dr. Luzinsky recovered),

that it is difficult to present a description of the lesions characteristic of this variety of the disease. M. Billard, however (*Mal. des Enfants*, p. 353), gives a case from M. Denis, and one observed by himself. M. Denis found the mucous membrane of a deep-brown color, of a fetid odor, reduced here and there to a state of putrilage, and everywhere easily removed in softened strips. A fluid of the color of lees of wine was found macerating the changed mucous membrane, and this he could ascribe only to gangrene from excessive inflammatory action. The case observed by Billard occurred in a girl three days old, who was brought to the infirmary with a quantity of blackish blood passed into the napkins, and some also vomited. The child died on the following day. The mouth and œsophagus were healthy, but the mucous membrane of the stomach was completely destroyed, not far from the cardiac orifice, over a space as large as a thirty-sous piece. The centre of this space was stained with blackish blood, and its edges, irregularly fringed, were blackened and looked as though they had been burned. Outside of this dark circle, the mucous membrane was thickened, of a violet-red color, and easily reduced to a pulp. The whole surface of the organ was lined with semi-fluid matters, of a bistre color, mixed with sanguinolent striæ, and the mucous membrane beneath these matters was very thin and discolored, especially near the pylorus. The small intestine was stained yellow with bile, and contained fragments of coagulated blood. The large intestine was healthy. The liver was bloodless and pale; the spleen small and but slightly injected. No clue is given as to the cause of this grave lesion.

The gastric lesions belonging to catarrh of that organ are very often met with, as we have already stated, but are almost always associated with changes in the intestinal mucous membrane. They are observed in severe indigestion, in simple and inflammatory diarrhœa, and in cholera infantum. For a full account of the histology of this lesion, we must refer the reader to the essay on Gastritis and Acute Gastric Catarrh, by Dr. Wilson Fox, in the *System of Medicine*, edited by Dr. J. Russell Reynolds. We shall, however, quote the shorter description given by Dr. Niemeyer (*op. cit.*, vol. i, p. 476) of acute gastric catarrh. He says: "We seldom have the opportunity of seeing the remains of acute gastric catarrh in post-mortem examinations; when we do, the gastric mucous membrane is found reddened in spots by a fine injection; its tissue is relaxed, and its surface covered with a layer of tough mucus. But more frequently, especially among children who die with the symptoms of cholera infantum, the autopsy gives negative results, except as to appearances which will be described hereafter. This does not appear strange when we remember that the capillary hyperæmias of other mucous membranes, which we have been able to observe directly during life, leave no trace after death, and that a relaxation and partial loss of epithelium, which we have regarded as the most probable cause of the extensive transudation in cholera infantum, may be very readily overlooked in the dead body, and can very rarely be observed with certainty."

The description of the anatomical appearances in gastritis will not be complete without some reference to a lesion which, some ten or twenty

years since, was thought to be one of great importance in children. This lesion, known by the names of softening or gastro-malacia, was supposed by some to constitute a distinct pathological entity, and to be the result in most cases of inflammatory tissue-changes determined by many different causes. Even then, however, not a few observers believed that the lesion was a post-mortem change, and not the consequence of changes caused by disease during life. This latter opinion has continually gained ground, until now it is generally believed that, when present in a marked degree, it is in fact a cadaveric change. Niemeyer (*op. cit.*, vol. i, p. 476) says that the gastro-malacia or softening of the walls of the stomach, found on autopsy in children, is always a post-mortem appearance, and that "if a child dies who has had vomiting and purging from abnormal fermentation in the stomach, and if there are still fermenting substances left there, the fermentation will not be arrested by the gradual cooling of the body. When the circulation ceases, the stomach can no longer resist the decomposition, which then extends to it also, just as the stomach that has been cut out of an animal and filled with milk, softens if left only for a short time in a warm place. Hence physicians who consider softening of the stomach as a post-mortem appearance, may also predict it with certainty when a child that has died of cholera infantum had eaten milk, or any other easily decomposed substance, shortly before death." We refer the reader, further, to the article on Thrush.

That a certain degree and kind of softening does, however, attend upon catarrhal inflammation of the gastric mucous membrane, as a result of faulty nutrition of the tissues during life, is probably quite as true as that the extensive white softening of one or more of the coats of the organ, not unfrequently met with, is the consequence of a post-mortem change. Thus Dr. Wilson Fox (*loc. cit.*, p. 858) asserts, that the softening of the mucous membrane which accompanies acute catarrh is totally distinct from the post-mortem softenings which are distinguished by the transparency of the tissues. "It rarely exists," he says, "to any marked degree, except in extreme cases, but there is always a certain diminution of resistance to the finger-nail or to the scalpel, which materially assists, when conjoined with opacity and thickening, in distinguishing this condition. Louis's test of the extent to which it can be torn from the submucous tissue is a less available one, and applies rather to the states of post-mortem solution than to this condition."

SYMPTOMS.—It is very difficult to give an accurate account of the symptoms of inflammation of the stomach, for the following reasons: they have not as yet been studied with a sufficient degree of care; gastritis is, as was stated in the early portion of this article, rarely idiopathic, but almost always a secondary affection in the course of other maladies; the symptoms which betray it resemble so closely those of intestinal diseases, as to make it very difficult, if not impossible, to draw a distinction between the two; and lastly, in the great majority of cases, gastric complaints coëxist with intestinal ones.

The most important symptoms are vomiting, diarrhœa, loss of appetite,

thirst, epigastric tenderness, sometimes tension of the abdomen, and slight febrile reaction.

Vomiting is the most important of the different symptoms of gastritis. It is not, however, according to MM. Rilliet and Barthez, invariably present. It was observed by them particularly in cases following the administration of active remedies, while in those which occurred spontaneously it was much less common. It shows itself especially after the taking of food for drink. Sometimes, however, even when the stomach is empty, there will be nausea and retching. In severe cases the vomiting is frequent, and accompanied by violent straining and pain. *Diarrhœa* exists in most cases, whether the attack be one of simple gastritis, or accompanied with enteritis. The *appetite* is generally lost or greatly diminished. *Thirst* is commonly acute, and often intense. The *tongue* is described by some writers as being generally red, and sometimes smooth and glazed. The authors above quoted state, on the contrary, that it presents nothing peculiar in most cases. It was generally moist, only slightly colored, covered with a white or yellow coat of variable thickness, and in some rare instances red on the edges and tip, or gluey, or even dry and harsh. As a general rule, the *abdomen* is normal, according to the same authors, though in some cases there is more or less swelling and tension. According to most writers there is generally tenderness or pressure in the epigastrium. Infants and young children are commonly restless and uneasy, as though in more or less pain, while those who are older complain of burning in the region of the stomach. It is well to remark that MM. Rilliet and Barthez state that tenderness on pressure often exists, not at the epigastrium, but in one of the iliac fossæ, or at the umbilicus, even when the stomach alone is inflamed. The condition of the circulation, and indeed all the symptoms, depend so much upon the nature of the concomitant malady, that it is difficult to ascertain what are their real characters in simple gastritis. Most writers agree that *fever* usually accompanies the disease, and that it is commonly of the remittent type. It is certain, however, from other observations, that it is not always present.

In very violent cases there are added to the symptoms just described, those indicative of an adynamic state of the nervous system: prostration, cool or cold skin, with perspiration; weak, rapid pulse; singultus; sometimes convulsions, and death. The symptoms which have just been detailed as indicating the presence of gastritis, do not generally exist alone. They are much more frequently than not associated with other symptoms, which show the presence of intestinal disease in the form either of simple or inflammatory diarrhœa. That they do sometimes, however, exist alone, and that, too, independently of the action of irritating drugs, or of corrosive poisons, we cannot ourselves doubt, since we have several times seen them follow attacks of simple indigestion. In such cases, we have met with all the symptoms usually supposed to indicate an inflamed state of the gastric mucous membrane,—repeated and obstinate vomiting, epigastric tenderness, entire loss of appetite, and more or less acute fever. We have, to be sure, never seen a post-mortem examination of such a case, for we have never yet known one to prove fatal. Whether we call such an

attack gastritis, acute casarrh of the stomach, or *embarras gastrique*, matters not much. It is the condition which has long been looked upon as indicating an inflammatory state of the gastric mucous membrane, and until we have more positive evidence than has yet been adduced, that inflammation has nothing to do with it, we deem it best to retain the old title.

DIAGNOSIS AND PROGNOSIS.—The *diagnosis* must rest chiefly on the existence and frequency of vomiting, on the presence of epigastric pain or tenderness, of swelling and tension of the abdomen and excessive thirst, and on the absence of other disease which might account for the illness of the child.

The *prognosis* will depend on the severity of the gastric and constitutional symptoms, and on that of the concomitant disease, when the attack is secondary. When there is incessant and obstinate vomiting, so that not even water in small quantities can be retained after several hours of sickness, when the tongue is red and glazed, or dry and brown, and when adynamic symptoms make their appearance, and emaciation makes rapid progress, it is much to be feared that extensive organic change has taken place, and that the case will prove fatal.

TREATMENT.—The two most important points in the treatment are the withdrawal of the causes that may have produced, or may tend to keep up the disease, if these can be detected, and strict attention to diet. Whenever, therefore, the symptoms have made their appearance after the exhibition of powerful drugs, as tartar-emetic, kermes mineral, or cathartics, their use ought to be instantly suspended. The child should be put on the strictest diet. If at the breast, it must be allowed to nurse only at rare intervals, and to take but little at a time. If fed on artificial diet, it should be restricted to barley- or arrowroot-water, to very weak milk and water, or to small quantities of milk diluted with lime-water, in the proportion of a third or a half of the latter. This is one of Dr. Chambers's favorite prescriptions, and is an admirable one. Nothing solid and no rich liquid nourishment ought to be allowed, unless the child is in a state of weakness and debility from previous or concomitant disease, such as to make it absolutely necessary to endeavor to maintain its strength. Billard even recommends that the child be sustained by means of nutritive enemata, while the digestive function is allowed a total rest.

Antiphlogistics are useful and proper when the disease occurs in a strong and healthy child, when it is associated with fever, and when there is nothing in the nature of the accompanying disease, if it be a secondary case, to prevent their employment. The most suitable mode of depletion is by the use of a few leeches, which should be applied to the epigastrium. It is best to take but a very moderate quantity of blood, for fear of exhausting the patient. After the use of the antiphlogistic remedy, a warm bath will be found of great service in moderating the heat of the skin and rendering the child more comfortable. Cool or cold water ought to be offered the child frequently, and it should be allowed to drink as often as it desires, and as much as it can retain. Even though it vomit the water, it should be allowed to repeat the draughts frequently. If the vomiting

be violent and constant, it may be necessary to limit the amount of fluid given each time to one or two ounces; but it ought to be frequently repeated, particularly when the thirst of the patient is very great. Bits of broken ice may also be given frequently. They seem, sometimes, though not often, we think, to allay the nausea better than water, but they do not satisfy the thirst. The addition of a small quantity of brandy to the cold water has, sometimes, a remarkable power of mitigating the gastric distress. A small teaspoonful of brandy to half a tumblerful or to a tumblerful of water, according to the age and present strength of the patient, is the proper dose. As soon as the bleeding from the leech-bites, if leeches have been employed, has ceased, a warm light mush poultice to the epigastrium is a valuable and useful remedy. Some writers recommend the use of blisters to the epigastrium. We should much prefer a warm poultice or the occasional application of a mustard poultice. Opiates are always indicated in these cases. One of the best forms is the following:

R. Liq. Morph. Sulphat.,	.	.	f 3 ss.
Acid. Sulph. Dil.,	.	.	gtt. xxx.
Elix. Curacoæ,	.	.	f 3 ss.
Aquæ,	.	.	f 3 iss.—M.

At a year old give half a teaspoonful, and at two or three years one teaspoonful every hour or two hours. From four to six doses may be used without risk, but the mother or nurse should be warned never to continue an opiate medicine, especially in young infants, if drowsiness or sleep sets in after several doses. Laudanum, or the deodorized laudanum, or paregoric, may also be used. One drop of laudanum, or half a drop of the deodorized, may be given at six months to one year, and repeated in one or two hours, as many as four or six doses being administered if necessary. Of paregoric, ten drops, repeated in the same way, may be used. We are of opinion that opium not only allays nausea, and vomiting, and pain as nothing else will, but that somehow it modifies more favorably what we have to call the inflammatory element of the disease.

When vomiting is frequent and troublesome, it may generally be allayed by the administration of lime-water and milk, given in teaspoonful, dessertspoonful, or tablespoonful quantities every fifteen minutes or half hour; by observing the precaution of allowing the food to be given only in small quantities (from a teaspoonful to a tablespoonful) and at considerable intervals; by the application of warm cataplasms over the abdomen, or a spice-plaster to the epigastrium; or, lastly, by the exhibition of a few drops of laudanum, paregoric, or morphia solution, as just explained, to be repeated if necessary. If the child becomes weak and exhausted, with coolness and abundant moisture upon the limbs, we must resort to the administration of some kind of stimulant. The best stimulant is brandy or whiskey. We prefer the former when it can be had good. Ten drops at six months of age, and twenty at one and two years, should be given every hour or two hours, according to the degree of exhaustion, in one or two teaspoonfuls of lime-water and milk made half and half. When, however, the stomach is excessively irritable, it is wisest, as a rule, to give the

brandy in iced water. If the exhaustion be alarming, the doses of stimuli ought to be doubled. Wine-whey, made of the usual strength and cooled, is sometimes acceptable, and ought in that case to be used. A dessert-spoonful at six months, a tablespoonful at one and two years of age, may be given every half hour or hour. Should it be retained by the stomach, the doses may be increased to one and two ounces and given less frequently. It is a curious fact that the thin chicken-tea referred to before, just touched with salt, will sometimes be taken eagerly by very young children, and retained, when all the milk foods are rejected. A few drops of aromatic spirits of hartshorn, one, two, or three, or from ten to twenty drops of the solution of the acetate of ammonia, in cold water, sweetened, may be tried, though we repeat, we have found nothing so useful as ice, iced water, weak brandy and water, and opium.

ARTICLE II.

ENTERO-COLITIS OR INFLAMMATORY DIARRHŒA.

DEFINITION; FREQUENCY.—By entero-colitis or inflammatory diarrhœa, we mean that form of diarrhœa which presents, during life, in febrile reaction at some period of its course, in marked constitutional disturbances, and in the mucous, muco-purulent, or muco-sanguineous stools, the proofs of inflammatory changes in the intestinal mucous membrane; and which exhibits, after death, the tissue-changes in the small and large intestines which are regarded as the products of inflammation of those organs.

The disease is a very common and fatal one in childhood. Many of the deaths accredited to cholera infantum belong to this disorder. The true choleraic disease is constantly passed through with safety, but is followed by a long, obstinate, exhausting diarrhœa, which is in truth an inflammatory diarrhœa occurring as a sequel to cholera.

A large proportion of the cases of summer diarrhœa are, from the beginning, cases of this kind; or they commence as merely functional disturbances of the intestine, and run, sooner or later, into the disorder we are now considering. It is one of the most important diseases of young children, especially in this country, where our long summer heats, and the filthy condition of many parts of some of our principal cities, give it a degree of prevalence and fatality which raise it to the rank almost of a pestilence.

We believe that most of the cases of diarrhœa in children, no matter what may have been the exciting cause at the start; whether a constantly improper diet, as in hand-fed children; whether ill-judged experiments in new foods by the mother or nurse; whether the accidental use of unwholesome food; whether summer heats, exposure to unhealthy and foul exhalations, crowding, malarial or epidemic causes, dentition, residence in cities, or what not, are prone to end, and nearly certain to end, if they

become chronic, in this disease. This opinion is the result of our experience in private practice, in this city, during many years. It is curious, too, and it is confirmatory of the correctness of this opinion, that in our armies during the late war, diarrhœa, whenever it became chronic, exhibited lesions which are best indicated by the term entero-colitis, if we are to use a name based upon the anatomical lesions of the disorder.

Entero-colitis, then, is undoubtedly one of the most frequent of children's diseases, though it is impossible to determine accurately the mortality it occasions in this city, from the returns as at present made by our physicians.

Thus during the seven years, 1862-1868 inclusive, there were 7273 deaths under five years of age in this city, from the three diseases, cholera infantum, diarrhœa, and dysentery (not to include a comparatively small number returned as due to colic, marasmus, inflammation of the stomach and bowels, aphthæ, etc.). Of these, as will be seen by inspection of the accompanying table¹ (see p. 406), by far the greater proportion, namely, 5963, are recorded as due to cholera infantum. Our extended opportunities of observing the diseases of children in this city have, however, led us to the conviction already expressed in the remarks which preface this article, that the great majority of these cases should in reality be entitled entero-colitis, while the true choleraic disease, to which alone the term cholera infantum should be restricted, is a comparatively infrequent affection.

We may appreciate yet more accurately the importance and frequency of the disease, by reference to the statements of MM. Rilliet and Barthez, who say (1ère., édit., t. i, p. 483), that, taking into consideration all the cases they observed, including tubercular cases, they find that of every two children that die, one presents a more or less serious lesion of the large intestine. They add: "If it be recollected that this holds true particularly in regard to younger children, it will be seen that it is rare for a child to die between two and five years of age, without having either colitis or softening of the large intestine." Bouchut states that entero-colitis is one of the most dangerous affections of children at the breast: "It is the most common of all those incident to that age" (p. 210).

We shall describe two forms of the disease, the *acute* and *chronic*. The acute form is accompanied by active and inflammatory symptoms from the first, and runs its course in a few days or weeks; the chronic form is unaccompanied by acute symptoms, and lasts several weeks or months.

CAUSES.—Much of what we shall say as to the causes of entero-colitis will apply to cholera infantum. The two diseases, together with simple diarrhœa, and some forms of dysentery, are so leagued together in their causation, much of their symptomatology, anatomical changes, and treatment, that they might also be regarded as different forms, stages, or expressions of a single disease. They are, too, largely interchangeable.

¹ We are indebted to the courtesy of Mr. Chambers, the clerk of the Board of Health in this city, for the opportunity of collating portions of this table from the monthly returns of mortality calculated by him.

TABLE SHOWING THE MONTHLY MORTALITY FOR SEVEN YEARS FROM CHOLERA LIFE; COMPARED WITH THE TOTAL MONTHLY MORTALITY

MONTH.	1862.			1863.			1864.			1865.		
	MORTALITY.		Mean temperature.	MORTALITY.		Mean temperature.	MORTALITY.		Mean temperature.	MORTALITY.		Mean temperature.
	Cholera infantum, dysentery, and diarrhoea.	Total.		Cholera infantum, dysentery, and diarrhoea.	Total.		Cholera infantum, dysentery, and diarrhoea.	Total.		Cholera infantum, dysentery, and diarrhoea.	Total.	
Jan.	1 2 4	1314	32.46°	0 3 3	1061	38.25°	1 0 5	1302	33.28°	1 6 3	1373	26.78°
Feb.	1 3 1	1080	32.70°	5 3 2	1122	35.00°	0 3 0	1434	35.97°	2 2 1	1550	32.59°
March.	2 2 3	1204	40.25°	0 2 7	1172	37.26°	4 1 5	1894	40.50°	3 1 2	1868	47.94°
April.	3 1 4	1213	50.61°	2 5 6	1488	49.80°	2 2 6	1377	50.58°	5 3 4	1411	56.46°
May.	9 4 5	1343	63.70°	5 1 6	1060	64.63°	10 5 8	1529	67.20°	10 6 7	1227	63.39°
June.	20 3 8	1002	69.14°	14 2 5	961	68.76°	74 11 14	1245	72.00°	184 10 20	1690	76.73°
July.	300 21 31	1767	75.23°	313 17 38	1859	77.07°	259 24 32	1643	76.08°	364 52 41	1838	77.82°
Aug.	217 19 22	1755	76.70°	464 25 28	2044	79.46°	250 27 31	1956	79.40°	245 42 23	1759	74.74°
Sept.	60 4 9	1037	69.36°	105 15 9	1453	64.73°	28 16 10	1251	65.00°	44 14 7	1040	72.68°
Oct.	15 5 4	1235	58.32°	14 4 5	1104	56.08°	9 8 4	1144	54.75°	15 12 5	1084	54.88°
Nov.	0 2 5	1021	45.20°	5 0 1	1061	47.72°	2 2 1	1212	45.80°	9 8 3	1285	45.35°
Dec.	0 1 2	1124	36.06°	3 2 1	1404	35.41°	2 2 4	1595	36.77°	0 2 3	1044	37.39°
Total.	803			1120			862			1139		

INFANTUM, DYSENTERY, AND DIARRHŒA, DURING THE FIRST FIVE YEARS OF
FROM ALL CAUSES, AND THE MEAN MONTHLY TEMPERATURE.

1866.			1867.			1868.			Mean mortality for seven years from cholera infantum, dys- entery, and diarrhoea.	Mean total mortality for seven years.	Mean temperature for seven years.
MORTALITY.		Mean temperature.	MORTALITY.		Mean temperature.	MORTALITY.		Mean temperature.			
Cholera infantum, dysentery, and diarrhoea.	Total.		Cholera infantum, dysentery, and diarrhoea.	Total.		Cholera infantum, dysentery, and diarrhoea.	Total.				
0 1 4	1402	29.31°	3 0 4	1376	25.89°	2 2 1	1249	30.12°	7½ 2 3½	1296½	30.87°
1 2 0	1156	34.14°	3 2 3	1042	40.21°	0 0 0	1063	26.65°	1½ 2½ 1	1206½	33.89°
3 0 3	1082	40.85°	3 1 2	1094	38.00°	1 1 1	1096	41.12°	2½ 1½ 3½	1344½	40.85°
6 3 4	1034	56.06°	1 2 0	1088	54.13°	9 5 2	1357	48.24°	4 3 3½	1281½	52.27°
8 1 4	1304	61.37°	7 2 3	1260	59.44°	4 1 6	917	59.66°	7½ 2½ 4½	1234½	62.77°
68 2 10	1168	73.04°	38 6 5	980	72.19°	71 8 7	1201	71.99°	67 6 9½	1178½	71.97°
427 21 34	2047	80.37°	423 23 31	1795	76.48°	423 14 32	1900	80.94°	358½ 24½ 34½	1837	77.71°
366 36 41	2401	72.50°	265 26 25	1294	75.10°	327 19 34	1570	78.42°	304½ 27½ 29½	1825½	76.62°
89 15 13	1362	69.42°	88 13 9	1012	68.21°	128 14 24	1353	68.80°	77½ 13 11½	1215½	68.31°
55 3 15	1828	58.35°	24 0 10	1177	57.65°	20 2 3	955	54.08°	152 4½ 6½	1218½	56.30°
6 0 4	1037	48.00°	6 2 5	871	47.79°	3 2 0	878	46.91°	31 2½ 2½	1052½	46.68°
2 2 3	982	33.63°	1 2 2	974	31.78°	1 0 3	1154	32.16°	9 1½ 2½	1191	34.70°
1252			1040			1057					

Nevertheless they do exhibit such different expressions, especially in their march and duration, and, as a consequence, in the treatment proper for each, that we think it best to adhere to our former classification and nomenclature.

The most active causes of the disease are: the heats of summer; residence in large cities, and this includes higher heat than residence in rural districts, with greater density of population and more copious filth emanations; and improper alimentation.

That the heats of summer are a fruitful cause of the disease, a glance at the accompanying table will show better than any words. In July and August, the temperature rising to 70° and 80° F., the deaths run up to 300 and 400 per month, and upwards. In January and December, the temperature being 30° to 40° F., they count from 3 to 5 and from 10 to 15 per month, and most striking of all, not a single death is reported in some of the winter months from diarrhœa, cholera infantum, or dysentery. We might add many more statistical facts, showing the powerful agency of heat, but it is useless.

From the well known fact, however, that those children suffer most who reside in the more filthy and crowded part of a city, whilst the disease is very much less frequent in the open country, and in the cleaner and better ventilated parts of a city, we may safely conclude that it is not heat alone that causes the disease, but that the emanations arising from garbage of various kinds, and the imperfect ventilation of houses built in narrow and crowded streets, have much to do with its causation.

There can be no doubt that improper alimentation may itself produce diarrhœal diseases, for we see them occasionally in cool or cold, as well as in hot seasons. The food most apt to give rise to entero-colitis is the artificial food of hand-fed children. Of the various articles that have been used for this purpose, the kind most apt to produce the effect is one composed exclusively or in considerable proportion of some of the feculent substances, which constitute so large a portion of the diet of children throughout the civilized world. To prove the truth of this assertion, it is only necessary to quote the opinions of those who have most carefully studied the subject. M. Valleix (*Guide du Méd. Prat.*, t. iv, p. 60, 61, and *Bulletin Gén. de Therap.*, article Acute Enteritis of Adults and New-born Children, March, 1845), clearly asserts, that the most frequent cause of muquet, which he believes to be essentially connected with enteritis, is a too exclusively feculent alimentation. In the article last cited, while speaking of the great importance of this cause, he says: "What proves that my assertion is not hypothetical is, first, that all the deaths from enteritis in children that I have seen, occurred in those who had been placed upon this kind of regimen; and, second, that the disease did not occur in any of those observed by me in private practice, for whom I had directed an exclusively milk diet up to four, five, or six months of age." He adds that M. Trousseau had arrived at similar opinions, after studying the same diseases at the Necker Hospital; and that he, on account of the danger of a system of diet disproportioned to the digestive powers, recommended that children be confined almost exclusively to the breast until

after the first dentition is completed. Barrier, speaking of the follicular diacrisis (*op. cit.*, t. ii, p. 40), states that the artificial food given to children at the period of weaning is a frequent cause of the affection, and that of all the different kinds of food habitually employed at that period, feculent substances are the most injurious. We have frequently known enterocolitis to follow the employment of artificial diet, either alone, at the period of weaning, or in children who were partly nursed. Children fed wholly on artificial diet from birth rarely escape, according to our experience, attacks of the disease, which in many prove fatal. It is not merely the quality, but the quantity also, of artificial food that proves injurious to infants. Overfeeding has always been recognized as a fruitful source of bowel complaints in early life. Another cause is the preparation of the food in too thick and rich a manner, thereby overtasking the stomach, intended during the early months to receive only the thin milk supplied by nature. The custom, therefore, of feeding infants on thick oatmeal gruel, with but little or no milk, on what is called cracker victuals (pounded cracker with water and sugar, or milk), on thick bread and milk, on preparations of rice of too solid a nature, or, indeed, on any kind of diet not consisting chiefly of milk, and in which feculent substances enter merely as secondary constituents, may safely be asserted to be the most frequent cause of the disease under consideration.

An unhealthy character of the milk of the nurse is also known to be a cause both of simple diarrhœa and entero-colitis. When the granule cells which exist as a physiological element in the colostrum secreted during the first few days after childbirth, continue to be present after that period, the infant is almost certain to suffer from entero-colitis, and not unfrequently to die, unless weaned or transferred to another nurse. So, also, when the milk departs widely from the normal characters which it should possess, when the nurse is liable to vivid moral emotions of any kind, or when addicted to intemperance, the child is very apt to suffer either from the disease under consideration, or from simple diarrhœa.

Another principal cause is excessive density of the population. In the *Fortieth Report of the Registrar-General of England for 1877* may be found some very important facts bearing upon this point. The figures include the deaths from all causes, but, as it is well known that a large proportion of the deaths are the result of diarrhœal diseases, they are very instructive as to the effects of crowding in cities. The density is calculated upon the proximity of the population in yards. The proximity is given for 593 districts of England and Wales, arranged in seven groups in the order of mortality. The districts of London are excluded. We have space only for a few extracts. In Liverpool, the proximity being seven yards, the number of living out of which one will die annually is 26, and the mean duration of life is 26 years. At the other end of the scale, of 345 districts in which the proximity is 139 yards, the number of living out of which one will die annually is 53, and the mean duration of life is 45 years.

Attention has been drawn of late years to the probability that certain changes, putrescent or fermentative, in the milk used for the food of young

children has a large influence in the causation of diarrhœa. Dr. Thomas B. Curtis, of Boston, (Buck's *Hygiene and Public Health*), says; "Milk, when exposed to atmospheric air, is known to be eminently putrescible. So liable is it to become contaminated by the development of various ferments, that Professor Lister used it as a substitute for Pasteur's solution in his experimental investigations into the subject of fermentation and putrefaction." Dr. Curtis quotes some instructive facts observed by Dr. Baginsky, during an investigation into the causes of infantile diarrhœa in Berlin: "He made a series of comparative experiments for the purpose of ascertaining the degrees of putrescibility of various articles of infant food, comprising woman's milk, cow's milk, Swiss milk, and two kinds of farinaceous food. These, having been previously boiled, were exposed to a continuous temperature of 37° C. (98.6° F.). After twenty-eight hours' exposure to this temperature, the woman's and cow's milk remained almost unchanged; but the Swiss milk, although appearing fresh, and the two farinaceous foods, exhibited bacteria in active motion. The woman's milk was alkaline, the cow's slightly acid, and the farinaceous foods were strongly acid. After a further exposure of eighteen hours, the cow's milk and the Swiss milk were coagulated, and the farinaceous foods in a high state of putrefaction, the woman's milk remaining still alkaline and almost unchanged. The experiments were repeated many times, and always with the same results." Dr. Curtis also quotes Dr. Meissner as asserting that cholera infantum never attacks children raised wholly on the breast, and as being a determined advocate of the bacterial theory of diarrhœal infection. "He expresses his conviction that the agency which, in midsummer, in densely populated districts, occasions fatal diarrhœa, does not reside in animal milk *per se*. The pernicious agent, he says, must be sought for solely in the drawn milk resulting from the access of atmospheric air, and from the imperfect cleansing of the vessels in which the milk is kept and transported, and of the bottles, tubes, and mouth-pieces through which it is administered to infants."

To show the readers the opinions now held by some hygienists, we make another quotation from Dr. Curtis: "It appears probable then, that the poisonous miasmata which are evolved from urban filth under the influence of high temperatures do not exert their universally recognized noxious action upon the infant directly by inhalation, but indirectly through the intermediate instrumentality of putrescive articles of diet. The injurious agent by which the particular form of filth-infection takes place is rotten food taken into the stomach rather than foul air taken into the lungs."

We have quoted these opinions for the benefit of our readers, but are inclined to think that the generalization is too broad. We know that we have seen sudden and violent cholera infantum arise in children feeding at a healthy breast, in the open country, in a perfectly well organized household. There could be no filth-infection here. And we have seen many cases of tedious summer diarrhœa in hand-fed children lasting for weeks, in spite of the fact that the child had been removed to the country (not rarely before the diarrhœa had set in), and under conditions of readily

supplied perfectly fresh cow's milk, where it was very difficult to suspect any fermentative or putrescent change in the milk as possible.

We referred, in the general remarks at the beginning of this chapter, to the resemblance of the chronic diarrhœa of our armies during the late great war, in its mode of causation, symptoms, anatomical lesions, and the effects of treatment, to the chronic form of entero-colitis in childhood.

Any one who will refer to the work of Dr. Woodward, already quoted, or to the essay on *Camp Diarrhœa and Dysentery*, by Dr. S. B. Hunt, in the *United States Sanitary Commission Contributions relating to the Causation and Prevention of Disease, and to Camp Diseases, etc.* (New York, 1867); or to the *Investigations upon the Diseases of the Federal Prisoners confined in Camp Sumter, Andersonville, etc.*, by Joseph Jones, M. D., published in the volume just alluded to; will find ample proof that improper diet, with heat, overcrowding, and want of cleanliness, will give rise to chronic diarrhœa, the essential lesions of which are to be found in radical blood-changes, perverted nutrition, and a localization in the alimentary canal in the form of entero-colitis, very much like the disorder we are describing. Dr. Woodward says, in fact, in speaking of the nature of this affection (chronic diarrhœa), at page 251: "From the account given above of the pathological anatomy of the disease, there can be little doubt that this affection is to be regarded as consisting essentially of a chronic inflammatory process, involving primarily the mucous membrane of the ileum and colon. It may, in fact, be described simply as a chronic ileo-colitis, with a tendency to ulceration." Dr. Hunt (*loc. cit.*, p. 294) says: "The essential fact in the pathology of all these various forms of flux is the same, and autopsies reveal no distinction between cases of diarrhœa and dysentery. They are alike an inflammation of the colon or of the small intestine, or of both, attended by ulceration of the mucous membrane. The solitary follicles of the colon are seen to be enlarged simply, or ruptured, with punched-out ulcerations following. The intestinal wall is thickened and changed in color to a red, brown, black, or greenish hue."

It may seem, at first view, visionary and wild to compare the chronic entero-colitis or inflammatory diarrhœa of childhood to the same disorder in armies and camps; and yet we think there is a most striking analogy between the two as to causation, symptoms, anatomical lesions, pathology, and the results of treatment. The main causes are the same: improper diet; elevated temperatures, the high temperature of the summer season in children, and of the Southern States in the armies; overcrowding, with foul air in camps and cities. The symptoms are very much alike, a most obstinate diarrhœa, with great constitutional suffering and emaciation; the same lesions are present, only less advanced and extensive in most cases of children; and very much the same results follow treatment: as in both diet is found to be more important than drugs, and removal North in the armies, and in children removal from crowded cities or low hot regions of the country to more elevated and cooler tracts, are found necessary. In children, as in armies, if, at the beginning of the attack, the patient is removed from the causes which have produced a simple diarrhœa or a cholera infantum, the case is likely to go no further; but, if

the same causes are continued in operation, the simple diarrhœa passes gradually into the chronic inflammatory form of entero-colitis, and at last the patient recovers only when he is removed to a more favorable locality, when the diet is changed to a more healthy one, or, in the child, when he drags through a long hot summer, and the cooler weather of October or November, and a diminution of the exhalations caused by the summer heats in cities, bring at last, in the course of nature, the change which was essential to his recovery.

After the causes just enumerated, the one which appears to exert the strongest influence is *dentition*. That the evolution of the teeth, though a physiological process, is a powerful predisposing cause of diarrhœa and enteritis, cannot be doubted at the present time. It is one recognized by many of the most able writers and observers of the day, and by most practitioners. MM. Rilliet and Barthez agree with Trousseau in the opinion that the simple diarrhœa so apt to occur in children at the epoch of the first dentition, is often the origin of chronic intestinal lesions which finally reduce them to extreme debility and emaciation. They say that careful investigation will generally show that nearly all the cases of inflammation and softening date either from the epoch of dentition, from the period of weaning, or from the time at which some considerable change in the character of the regimen was made. M. Legendre and M. Barrier (*loc. cit.*) both agree in ascribing very great effect to the influence of dentition in the production of diarrhœa and entero-colitis. The former asserts the diseases referred to, to be much the most frequent between the ages of six or seven months and two or two and a half years, which includes exactly the period occupied in the first dentition, while they are only met with exceptionally after three years of age.

The act of *weaning* is very apt to result in the production either of simple diarrhœa or entero-colitis, in consequence, no doubt, of the irritation set up in the gastro-intestinal surface, by the change of food made at the time. The diarrhœa which occurs at this period was formerly, and is still, not unfrequently, called *weaning-brash*.

Entero-colitis is prone to occur as a secondary affection in many of the acute diseases of children. It is by far the most common in the course of the eruptive fevers, particularly measles, and in that of typhoid fever. It is also a frequent complication of the latter stages of pneumonia.

That children of feeble constitution and lymphatic temperament are more disposed to the disease than others, is sufficiently proved by the evidence of various observers. Lastly, that the incautious and excessive use of perturbing systems of medication, addressed to the digestive tract, often occasions diarrhœa and entero-colitis, is fully proved by the researches of MM. Rilliet and Barthez, and by our own experience.

MORBID ANATOMY.—SEAT OF DISEASE.—It has been already stated, that the alterations in the large intestine are, as a rule, much more frequent and serious than those in the small intestine. It appears from the researches of MM. Rilliet and Barthez, and Legendre, that enteritis rarely exists alone; whilst colitis by itself, or combined with enteritis, is quite frequent. M. Legendre states that inflammation of the small intestines

never occurs without corresponding lesions of the large bowel, while in 28 cases of diarrrhœa he found the large intestine alone diseased in 9. From a table of different intestinal lesions, given by Rilliet and Barthez (*op. cit.*, t. i, p. 692), it appears that they have met with 45 cases of erythematous, pseudo-membranous, ulcerative or pustular enteritis; with 113 of the same forms of colitis; with 90 of follicular enteritis; 64 of follicular colitis; and with 28 of softening of the small, and 35 of softening of the large intestine. Dr. J. Lewis Smith (*op. cit.*, p. 367) offers an analysis of the post-mortem appearances in 82 cases of intestinal inflammation in children. The upper part of the small intestine, embracing the duodenum and jejunum, was found inflamed in 12 cases, while in 51 cases it was free from inflammation and of a pale color. The ileum was inflamed in 49 cases, and the cæcal portion, including the ileo-cæcal valve, was the part in which the inflammation was uniformly most intense, and to which it was often confined; in 13 cases there was no enteritis whatever, and in 16 there was no inflammation of the ileum, so that the ileum was inflamed in all but 3 cases where enteritis was present. On the other hand, in all the cases excepting one, namely, in 81 out of 82 cases, there were lesions indicating inflammation of the mucous membrane of the colon. In 39 the inflammation had affected nearly or quite the entire extent of this portion of the intestine; in 14 it was confined to the descending portion entirely, or almost entirely; in 28 cases, the records state that colitis was present, but its exact location is not mentioned.

We may add, that, in the quite numerous autopsies we have made after death from this disease, we have invariably found the large intestine involved, the inflammatory lesions being in some cases limited to it, while in others they also extended into the small intestine.

It is, therefore clearly established, that in the inflammatory diarrrhœa of children, inflammation of the large is considerably more frequent than that of the small intestine, and much more apt to exist alone. The lower end of the ileum is the portion of the small intestine which presents the most advanced and severe lesions; while in the large intestine the lesions are most marked in the caput coli, sigmoid flexure, and descending colon.

In our description of the lesions of entero-colitis, we shall divide them into those found in the acute and chronic forms of the disease respectively; a division made for the sake of correspondence with the description of the symptoms, although the lesions found in the two stages differ from each other only in extent and degree.

Thus, in the acute stage, the lesions consist of increased vascularity, thickening and softening of the mucous membrane of the intestine, and enlargement of the intestinal follicles; while in the chronic form there is discoloration, thickening, with infiltration and induration of the walls of the intestine, and more or less extensive destruction of the mucous membrane from follicular ulceration.

In the acute stage, the increased vascularity (inflammatory hyperæmia) may present itself as a uniform, more or less intense redness of the mucous membrane; an appearance which may sometimes exist in the duodenum, but far more frequently is observed in the lower end of the ileum and in

the colon. More frequently it takes the form of arborescent congestion, occurring in patches surrounding the enlarged follicles. The peritoneal surface may also be more or less vascular, and quite frequently there are little patches of redness and arborescent vascularity, corresponding to the bases of the inflamed mucous follicles.

The thickening of the mucous membrane usually corresponds to the degree of vascularity, and when the latter is but slight, may be scarcely appreciable; while in other cases, and especially when associated with much enlargement of the mucous follicles and œdema of the submucous tissue, the thickening is highly marked. The inflamed portions of the mucous membrane are also more or less softened, so that they can be detached from the subjacent coats more readily than in health. In some instances the softening is so extreme that it is impossible to raise up the mucous membrane in strips at all. These lesions are all most frequent and marked in the lower part of the ileum, and in the descending part of the colon. In addition to these changes in the color, thickness, and consistence of the mucous membrane, the mucous follicles are prominently enlarged. In the normal state, the isolated follicles of the mucous membrane of the intestine, in young children, appear as minute grayish-white bodies, and present a grayish point, the excretory orifice, which is only visible with the aid of a lens. In the course of entero-colitis, however, the morbid development which they undergo causes them to present the following characters: The isolated glands are enlarged, and seem, therefore, more numerous than in the healthy condition; they appear in the form of lenticular grains seated in the texture of the mucous membrane, sometimes projecting from its surface, sometimes not, and in other instances appearing to be situated beneath it; the excretory orifices of the follicles are often enlarged and tumid, and easily distinguished under the form of a grayish or blackish point in the middle of the gland; in other cases the orifices cannot be distinguished until slight pressure is made upon the crypts, when a drop of turbid mucus may be seen exuding through the open point. The color of the distended follicles is dull white, rosy, or yellowish; they are generally from one-third to two-thirds of a line in diameter. Dr. Horner (*Amer. Jour. Med. Sci.*, Feb., 1829) speaks of them, in this state of development, as resembling "small grains of white sand sparkled over the mucous membrane, and about the size of a millet-seed."

The agminated glands or patches of Peyer are found in the same state of increased development; they are tumefied, and project above the level of the surrounding mucous membrane, and the orifices of the follicles are congested, so as to appear as dark points, giving to the patch a dotted, punctated appearance, which has been compared to the freshly shaven chin.

A little later the enlarged follicles present minute, oval, or round yellowish spots upon their summits, which soften down and allow the contents of the follicles to be discharged. The enlarged orifice of the follicle will then admit a small probe, and may even measure one-half a line in diameter. It leads into a little cavity, which is the follicular sac. The mucous membrane which overhangs this cavity like a fringe is thus undermined

and partly cut off from its vascular supply, so that we may find a process of ulceration advancing in it until the base of the distended follicle is exposed, appearing as a small, oval, or round shallow ulcer.

These various conditions of the follicles may all be seen at the same time in a single portion of intestine. The enlarged patches of Peyer often have the appearance of being ulcerated, but a careful examination will generally show that this is not the case. The appearance depends upon the enlargement of the orifices of the glands, upon unequal tumefaction of the surrounding mucous membrane, and upon the presence, in the patch, of small, irregular grayish points, consisting of pultaceous matter, which makes the patch look more uneven and projecting than usual. If, however, the pultaceous layer be gently rubbed with a piece of linen, it can easily be detached, when the mucous membrane beneath is found red, softened, and thickened, but not ulcerated. In comparatively rare cases, however, there are superficial erosions of the mucous membrane, covering the prominent patch.

The exact date at which the ulceration of the follicles begins, is as yet undetermined, and probably varies greatly in different cases. It frequently happens, however, that death occurs, especially from the super-vention of a choleraic condition, whilst they are still merely in a state of enlargement. When, on the other hand, the disease passes into the chronic form, the lesions which we have above described become more and more extensive. This is especially the case with the lesions in the large intestine, for it is even more true with regard to chronic than acute enterocolitis, that the chief seat of the disease is in the colon.

In chronic entero-colitis, the intestine is often contracted, and the peritoneal surface may present patches of discoloration. The thickening and infiltration have now affected the submucous and muscular coats, and have been followed by induration of the tissues, so that the walls of the intestine are often abnormally rigid. This is especially true with regard to the lower part of the descending colon and the rectum. The mucous membrane is seen to be riddled, not with mere superficial erosions, but with true ulcers, affecting the whole thickness of the membrane. These ulcers, when isolated, are from one to one and a half lines in diameter, oval or circular in shape, and either have sharp-cut edges, as though the piece of mucous membrane had been cut out with a punch, or the mucous membrane bounding them is undermined. Frequently, however, these ulcers coalesce, and at the same time extend in depth, so that large, sinuous, irregular ulcers are formed, with thickened, slate-gray, undermined edges, and having for their base either the submucous or muscular coats, which may be covered with a pultaceous, apparently pseudo-membranous layer, of a grayish-white color. These ulcers surround and include irregular islets of mucous membrane, which are swollen, infiltrated, vascular, and discolored. That the large and deep ulcerations just described, even when most extensive, take their start from the mucous follicles, is proved by the frequent presence amongst them of other ulcerations of more recent date and smaller size, which present all the characters of the follicular ulcer, and show clearly the origin of the larger and more advanced ulcerations.

Occasionally there is a marked deposit of pigment in the bases of the ulcers, and in some cases small coagula of blood have been found adherent to their bases.

We have already had occasion to allude to the marked analogy between the disease under consideration, and the form of camp diarrhoea described by Woodward (*op. cit.*); and one of the most powerful arguments in favor of the essential identity of the two affections, is the perfect correspondence between their anatomical lesions. We present below a summary of the microscopical changes in the intestine during the development of these lesions, as determined by the careful investigations of Dr. Woodward (*op. cit.*, p. 246). In the early stage, attended merely with thickening and softening of the mucous membrane, microscopic examination shows marked multiplication of the connective-tissue cells about the base of the follicles, and soon the tissue is occupied by great groups of small, rounded, or slightly polygonal cells. The delicate layer of muscular tissue immediately beneath the base of the follicles, presents, at first, enlargement and proliferation of its nuclei, whilst later it often ceases to be recognizable, being obscured by the luxuriant cell-growth. In the most intense cases, the cell-growth here described as attained toward the surface of the membrane, may take place throughout its whole thickness, and even involve the subjacent muscular layer.

A similar proliferation takes place in the connective tissue, which lies between the follicles. The epithelial layer, which invests the mucous membrane, and is prolonged into the tubular follicles, either is the seat of rapid cell multiplication, or is exfoliated and replaced by round granular cells from the adjacent connective-tissue cells. The epithelial lining, near the orifice of the follicles, appears to undergo these changes most readily and with the greatest rapidity.

The *closed* follicles also present rapid cell multiplication, which affects the parenchyma of the follicle, as well as the connective tissue of its capsule and the surrounding cellular tissue. Microscopic examination then shows the follicle distended with small, rounded, granular cells, and imbedded in a luxuriant growth of similar cells, which render it almost or quite impossible to draw the line where the follicle terminates and the surrounding connective tissue begins. "Ulceration usually appears to originate in the rupture of one of the closed follicles, and the discharge of its softened contents into the intestinal cavity. This is followed by the liquefaction of the intercellular substance, and the consequent liberation of the broods of minute cells, into which the surrounding connective tissue has been transformed. Hence results one of the punched-out ulcers described above. In the subsequent extension of the ulceration, by which large, irregular, sinuous ulcers are produced, the progress seems to take place chiefly in the submucous connective tissue, the superficial part of the mucous membrane resisting the process until undermined, and its nutritive supply cut off. Hence arises the excavated undermining character of the edges of the ulcers. From the anatomical point of view, it will therefore be perceived that the morbid process, in the cases in which there is no

ulceration, is essentially the same as in those in which ulceration is present. The one lesion is only a later stage of the other."

Not unfrequently there will be found one or more intussusceptions of the ileum. These are usually readily restored, and have evidently occurred during the act of dying. Smith has, however, "in a few instances, found intussusceptions which sustained the weight of two feet or more of intestine without being reduced, and which, from being in their interior more vascular than the contiguous membrane, had probably occurred some hours or days before death, but being sufficiently pervious to allow the food to pass, the symptoms of obstruction were lacking."

The *Mesenteric and Mesocolic Glands* are nearly always enlarged, the most marked enlargement corresponding to the lower end of the ileum and the descending colon. The enlarged glands are of a pink color, and rather more soft and succulent than normal.

Stomach.—In the great majority of cases the stomach is quite healthy; in a few instances, however, there may be found congestion of the mucous membrane, slight enlargement of the mucous follicles, or softening of the mucous membrane, probably cadaveric in most cases.

Liver.—Many authorities, apparently led by the presence of symptoms supposed to indicate disturbance of the function of the liver, have assumed that there is in most cases of entero-colitis some morbid condition of this organ, but extended observation has disproved this view.

Thus Hallowell (*Amer. Journ. Med. Sci.*, July, 1847) found, that in 14 cases, the liver was affected in but 1 case, being simply enlarged; and Smith (*op. cit.*, p, 370) has published the result of 32 post-mortem examinations in regard to this point, which confirm the same conclusion. Thus, he states, "there was no evidence from the post-mortem appearances of the liver in these cases of any congestion, or torpidity, or hyper-activity or perverted secretion. The size of the liver was in some cases very different in those of about the same age, but probably there was no greater difference than usually obtains among glandular organs within the limits of health. In most of the cases the liver was examined microscopically, and the only fact worthy of note observed was the variable amount of fatty matter. Sometimes it was in excess, sometimes in moderate quantity or rather deficient, and sometimes in greater amount in one portion of the organ than in another."

The thoracic viscera present no constant or important lesion, though in a certain proportion of cases there may be found more or less hypostatic congestion with collapse of portions of the lungs.

When death occurs during the acute stage, the brain presents no lesions dependent upon the disease. When the case has been protracted and attended with much wasting of the solids and fluids of the body, the brain also diminishes in size, and there is frequently found marked excess of subarachnoid effusion in cases where the fontanelles have closed; while if these spaces still remain unossified they become markedly depressed. These appearances are, however, purely passive in their character, and depend upon the wasting of the brain.

PATHOLOGY.—The pathology of inflammatory diarrhœa is involved in great obscurity. We are now pretty well acquainted with the physical conditions under which the disorder is most apt to be developed. Early age, the period of dentition, high temperatures, improper food, residence in cities, and especially the crowded occupation of small and ill-ventilated buildings, in narrow courts and alleys, where unhealthy exhalations arise from the decomposition of filth and dirt of all kinds, are the chief conditions which precede the outbreak of the disease. But how these conditions act to produce their effect is still a matter of doubt. To attempt to reason upon a matter so full of difficulty seems almost useless, and yet we shall venture to place before the reader some thoughts we have had upon the subject.

There are two broad generalizations which we think may be safely assumed to be true. 1. An unhealthy food, one incompetent to furnish to the body what it needs for the purpose of nutrition, as farinaceous food or unhealthy milk, is sure to produce the disorder we are considering, no matter how favorable may be the circumstances, in all other respects, in which the child is placed. 2. The best breast-milk in the world, or the most correct artificial diet, will not save a child from this disorder who is located in an ill-ventilated house in a dirty and filthy quarter of a large city during hot weather. Here the heat to which the child is exposed, the heavy air loaded with foul exhalations which it breathes, determines a condition of the health in which the digestive organs can no longer digest properly the food offered them. In both cases the same result is produced. In the first, the stomach cannot change the originally improper character of the food into healthy material. In the second, the diseased and enfeebled organ loses the power to digest even proper food. In both the alimentary canal is filled with the products of an improperly digested food. Whether these unhealthy products in the alimentary canal act as local irritants to the mucous membrane, and thus determine the tissue-changes met with; or whether, as Rilliet and Barthez suppose, some morbid condition of the blood is brought about, which gives rise to the changes in the mucous membrane through a morbid action of the diseased blood on the nervous system, and particularly on the sympathetic nerves, we cannot say. Most probably they act in both ways, and the resultant effects are the consequence of the two trains of diseased action set up, the local and the general.

In either case a constitutional condition is brought about, the essential feature of which is a slow innutrition or inanition. It is altogether probable, moreover, that a condition partaking of the scorbutic must be induced, so that we have, after the disorder has lasted for several days or weeks, the general debility of a slow inanition, and blood-alterations which resemble those of scurvy.

SYMPTOMS; DURATION.—In infants the *acute* form of entero-colitis generally begins with restlessness and fretfulness. The mother observes that the child sleeps less than usual and for shorter periods, and that its sleep is uneasy and broken by sighing and moaning, or by occasional expressions of pain flitting across the face. It takes the breast less frequently,

and is satisfied to nurse for a shorter time, showing thereby an evident diminution of appetite. At the same time it is apt to reject its milk in larger quantities than usual, and this is often observed to have a very acid smell. After these symptoms have lasted a few days, and sometimes without them, the peculiar symptoms of the disease, the diarrhœa and other abdominal symptoms make their appearance, and are accompanied by febrile reaction in most cases.

In older children the acute form may come on suddenly, with diarrhœa, loss of appetite, thirst, sometimes vomiting, abdominal pain and fever from the first; or, as happens very frequently, the case begins with slight diarrhœa, unaccompanied by fever or other signs of sickness, and it is not until after several, or eight, ten, or even more days, that signs of inflammation make their appearance.

After the disease is established, the most important symptoms are the following. The *diarrhœa*, which is the most prominent and characteristic, presents various characters. In order to appreciate this symptom as its importance requires, the practitioner ought always to see the napkins of the child at least once, and often more frequently, in the day. It exists in almost all cases of entero-colitis, in the erythematous and follicular inflammations, and in the ulcerations and softening which accompany or succeed simple inflammation. It is seldom absent, and yet that it is so sometimes, is proved by the facts mentioned by MM. Rilliet and Barthez, who state that they have calculated, from their cases, that it is wanting in about one of every twelve cases of inflammation or softening of the intestine. They add, however, that it is absent only in slight attacks, and is always present when the disease is severe. It varies greatly as to the frequency, abundance, and character of the stools. It varies also in its mode of progress, so that it presents great differences as to all these points from day to day, and at different portions of the same day. We may remark, in general, however, that in proportion to the severity of the inflammation, so is the diarrhœa violent and constant, and that it usually increases as the signs of inflammation become more and more marked. It is rare to have severe diarrhœa when the anatomical lesion is of slight extent, though this does sometimes happen. The *number* of the stools, as has been stated, is exceedingly variable. This depends in a great measure upon the violence of the case; for, while in those which present the symptoms of an inflammation of small extent the stools seldom amount to more than six or eight a day, in those in which the evidences of more extensive and severer inflammation are present, there will be fifteen, twenty, twenty-five, or even more per diem. The *consistence* of the stools may vary between that which characterizes them in a state of health, and that of the thinnest serous fluid. The *materials* of which they are composed consist chiefly of mucus, bile, serum, small portions of feculent matter, portions of undigested caseine or other food, and blood.

After the epoch of the first dentition the disease becomes much more rare, and when it occurs, is generally of a milder character, so that the discharges differ less from their healthy characters. Under these circumstances, they are usually less frequent, not often exceeding six, eight, or

ten in the day, and retaining generally their yellow color or becoming brownish; they are commonly of a semifluid consistence, and may be called bilious. When, on the contrary, more frequent, they become fluid, abundant, mixed with mucus, and are either of a light yellow or brownish, or more rarely, of a greenish color. In some cases there are, in addition to the substances mentioned, pus, which indicates ulceration of the lower portion of the intestine, and fragments of false membrane. Moreover, it is very common in older children to observe traces of blood in the stools, sometimes in considerable quantities. We may remark that we have several times met with stools containing blood in children within the year, but much less frequently than after that age. The presence of blood generally coincides with small and frequent stools, attended with much straining, and often severe pain, and almost always indicates follicular inflammation and ulceration of the large intestine.

The serous fluid alluded to sometimes constitutes the whole of the discharge, so that the napkins are merely wetted through, without any or but a very small quantity of solid matter being left upon them. This kind of stool is very frequent in the cholera infantum of this country. The *odor* of the stools is important. In the beginning, while the discharges still retain some of their natural characters as to color and consistence, it is often very offensive, but as the case goes on, and the greenish color predominates, it is either sour, or becomes very slight. In some violent cases, in which the discharge consists of a watery, dark brown fluid, the odor is fetid.

After diarrhœa, the most important symptoms are those which concern the *form*, *size*, and *tension* of the *abdomen*, and the presence or absence of *pain* or *tenderness* on pressure. In infants the abdomen is more distended than usual; but, according to Bouchut, the tension depends on the muscular effort made by the child to resist the hand of the physician. He says that when it is carefully examined, while the attention of the child is attracted in some other direction, it is found to be soft and supple, and rarely painful to the touch. In older children it is, in many acute cases, but not in all, enlarged, sometimes tense and sonorous, and very generally painful to the touch. The seat of pain is variable, but generally it is in one of the iliac fossæ or at the umbilicus. It is seldom acute, though the child not unfrequently shrinks away and cries out, as though it were excessive, from fear of the examination. It is easy to distinguish when the pain is real and when apparent, by withdrawing the attention of the child, by some device, from the examination, in which case it will cease to notice the palpation; or by touching some other part of the body, when, if the crying and shrinking depend on fear or nervous excitation, they will be as violent as when the abdomen is touched. Pain to the touch is an important symptom, as it is very generally indicative of acute enteritis. *Gurgling* is rare, according to MM. Rilliet and Barthez, in ordinary entero-colitis, though very generally present in typhoid fever.

Vomiting is very common in young infants, and is generally repeated several times a day. In severe and rapid cases it is a very troublesome and alarming symptom. In older children it is much less common, and

is never really violent, except in some of the most acute cases. In them it is confined to the first few days of the attack.

After the diarrhœa is fairly established, young infants are almost always either very irritable, peevish, and restless, or weak, languid, and subdued. Their slumber is short and disturbed, and generally they sleep much less in the twenty-four hours than when in health, unless under the influence of anodynes. Older children are generally somewhat restless and irritable, but much less so than infants. There is seldom any disorder of the intelligence, though in acute cases there is sometimes slight delirium, or headache. *Fever* exists in all acute cases. It is seldom continuous in infants except for the first few days, after which it almost always assumes the remittent type. It is marked by increased frequency of the pulse, which rises to 120 and 140, or in bad cases much higher; by heat of skin, often intense during the exacerbations; by thirst and diminished appetite; and by dryness and heat of the mouth. In older children the pulse is not generally so high as in infants, and in many of the mild cases the fever is very slight or there is none at all. In acute cases, however, it is sometimes continuous, and marked by rapid pulse and great heat of skin.

The *tongue* is generally normal, though sometimes red on the edges and tip in acute cases. It is seldom dry, except during the fever. The *appetite* is almost always lost, and the *thirst* generally increased, though to a less degree than in diseases of the stomach.

The *countenance* presents nothing peculiar, except that the features are, according to MM. Rilliet and Barthez, drawn down towards the inferior portion of the face. Emaciation always takes place as the disease progresses, and in very severe cases occurs with the greatest rapidity, so that in a very few days the child will be reduced from an appearance of vigor and strength to that of the greatest debility. As this occurs the flesh loses its firmness, the skin hangs in folds upon the trunk and limbs, and is dull and dirty in its tint, the eyes become sunken and surrounded with bluish circles, and the whole appearance of the child is that of misery and exhaustion.

In infants, it is very common to meet with erythema of the buttocks and thighs, produced by the contact of the acrid stools and urine with those parts. This symptom is said by Bouchut to exist in five-sixths of the cases. We feel quite sure that it does not exist in so large a proportion of those which occur in private practice, though we have met with it in numerous instances. When severe it is generally accompanied by papules, which ulcerate after a time and form superficial ulcerations upon the skin. These ulcerations sometimes run together, and become of considerable size and depth. In the form of the disease met with in the children's hospitals in Paris, erythema and ulcerations of the heels and internal malleoli are also met with, and constitute a serious complication in the case. They are said to depend on want of cleanliness, and the rubbing together of the feet of the child, unprotected by covering. We have never met with them in private practice,

The *duration* of the disease is stated by the French writers to be generally about fifteen days, at the end of which time convalescence is usually

established. It may be shorter or longer. According to our own experience it is entirely uncertain. Most of the cases that have come under our notice have been rather shorter. The disease subsides gradually. The number of stools diminishes; they become less abundant and more consistent, and return to their natural color and odor; the pain on pressure, and the enlargement and tension of the abdomen disappear; and as this occurs, the fever subsides, the appetite returns, the temper improves, and the child enters into full convalescence.

The *chronic form* of entero-colitis generally follows the acute, though it sometimes presents many characteristic features from the first. It differs from the acute form chiefly in the absence or the much slighter degree of fever and other constitutional symptoms in the early stage. The diarrhœa is less abundant and less frequent. At first the child retains its spirits and many of the signs of health. But gradually its strength fails, the temper becomes irritable, the complexion grows dark, sallow, and unhealthy, the skin becomes dry and harsh, and, in consequence of the emaciation which takes place progressively with the other symptoms, hangs in folds around the shrunken extremities, or is drawn tightly over the joints and other osseous protuberances. The tongue is generally moist and natural, though in some cases red and dry, whilst in others it, together with the lips, partakes of the pallor which pervades all parts of the body. The abdomen is usually distended and sonorous on percussion, and may be painful or not on pressure in different cases, or in the same case at different periods of the disease; its parietes sometimes offer no resistance to the touch, so that the intestinal convolutions may be readily felt by the hand, or even between the fingers; and in some cases we have seen them so thin and relaxed, though the abdomen was more prominent than natural, that the outlines of the intestines, and even their peristaltic movement, were visible upon the exterior. The appetite generally persists in spite of the gravity of the disease, and is sometimes increased. The stools, as has been stated, are not so frequent as in the acute form, seldom numbering over six or ten in the day and night. They consist of the products of an imperfect digestion, and contain not unfrequently the alimentary substances in the state in which they were swallowed, mixed with mucus, serum, pus, and sometimes blood. Their consistence varies constantly, but they are usually semi-fluid. Their odor is seldom natural, and often extremely offensive.

The *course* of the disease is very irregular. Even in the worst and most prolonged cases intermissions or remissions occur, so that the child will often improve greatly for days or weeks, and then suddenly relapse into as bad a condition as ever. In favorable cases these remissions become more and more frequent, and the symptoms gradually improve, until at length the child is restored to health. In fatal cases death is occasioned by the utter deterioration of the general health which finally occurs, and the child perishes, worn out by long illness, or, as more frequently happens, some complication arises which hurries on the fatal event. Thrush is a frequent complication of chronic entero-colitis, and doubtless often hastens the death by the impediment which it occasions to the nursing or feeding of the child. Vomiting has almost always occurred towards the close of

the fatal cases that we have seen, especially in those in which extensive thrush was present.

The *duration* of this form is of course very uncertain. It may last for weeks or months. We have known it to last for two or three months in several cases, and in two others it lasted with occasional intermissions, in one a year, and in the other eighteen months.

DIAGNOSIS.—The diagnosis of acute entero-colitis is not difficult. There is no disease with which it is likely to be confounded. The characteristic features of the malady are the diarrhœa and other abdominal symptoms, and the absence of signs of other disease. The secondary cases are distinguished by the occurrence of the usual symptoms of entero-colitis during the progress of the primary malady.

The chronic form is not likely to be mistaken for any other disorder, unless it be the diarrhœa which occurs in tubercular disease, from which is to be distinguished by the presence in the latter of the signs of tuberculosis of other organs.

PROGNOSIS.—Acute entero-colitis is always a serious disease in infants. The prognosis will depend in great measure on the circumstances under which the affection has been developed. It is much more unfavorable in a child fed on artificial diet, either wholly or in part, than in one who is nursed at a fine breast of milk. It is more unfavorable also in weak and delicate than in robust and vigorous children, and in those of poor people, who live in crowded and unhealthy portions of cities and towns, whose habitations are small, damp, and ill-ventilated, and whose food is coarse and insufficient, or improper, than in those placed in more fortunate and more healthful hygienic conditions. It is a more dangerous disease in summer than in winter. In hospitals for children it is a very fatal disorder, owing to the bad hygienic conditions under which the inmates are placed. In children who have passed through the first dentition, the prognosis is, as a rule, favorable. The disease is seldom dangerous in such cases when it occurs as a primary affection, while, as a secondary affection, on the contrary, it is much more apt to be serious.

The unfavorable symptoms are: great frequency of the stools; collapse; violent vomiting or retching; and dangerous cerebral symptoms, as coma, rigidity of the limbs, paralysis, or convulsions.

TREATMENT.—The *prophylactic treatment* is very important. It includes attention to *habitation, diet, dress, and exercise*. The most frequent causes of entero-colitis are high summer temperatures, residence in an unhealthy locality, and improper diet. A child may have been born of the most healthy parents; may be living, if an infant, on the most healthy food in the world, the milk of a perfectly sound woman, or, if it have been weaned, on the best possible substitute for breast-milk, one selected by the most consummate medical art; and yet, if it be the unfortunate resident of some low, crowded, and unclean part of any of our cities in the summer season, it has but few chances of escaping inflammatory diarrhœa or cholera infantum, to be followed by chronic diarrhœa. Or, a child may be living in the best part of these cities, with every advantage that wealth and the medical art can give, and, if in the period of the first dentition, and the

summer heats be great, it will be only too apt to have some form of the disease we are considering. Under the latter circumstances, its chance of escaping the disease will be vastly greater than under the first named conditions, but the true prophylaxis is, where the parents are so situated as to be able to do that which is best for the child, removal from the city during the hot season (from the early part of June to the last week of September) into some cool and healthy region of country. We have long thought that the best region to spend summer in is a somewhat high and cool part of the country, where the breezes have full sweep, and where the topography is such that water runs off rapidly, or sinks fast into the soil. The seaside, if it be a point where there are no marshes and no malaria, and where the supply of milk and other wholesome food is abundant, is an excellent place. We have seen more remarkable sudden effects from the removal of a dangerously sick child to the seaside, than from a change to the interior; but, nevertheless, for a continued residence of three months, we prefer a high interior locality.

On the other hand, if a child be placed in the most favorable possible condition as to locality, and the diet be a radically bad one, a deficient or unhealthy breast, improper artificial diet, or a foolish allowance on the part of the mother or nurse to the child of a variety of vegetables, of fruits, and especially of berries like currants or gooseberries (and we have known such things), it can scarcely escape the penalty of a fit of illness more or less severe.

A child who is so unfortunate as to get a sharp attack of entero-colitis in June or July, is very apt to continue more or less sick during the rest of the summer, so that the true prophylaxis is to take it away from the city early in June to avoid this danger, and not to return until after the September heats are over.

As the reasons for decisive medical action in any disorder cannot be too strongly demonstrated, and as this subject of removal is a very important one, we think it well to advert here to the results of experience in this matter in the diarrhœa and dysentery of our armies during the late war. Here we have the experience of intelligent army medical officers in vast numbers of cases,—cases, too, so grouped together as to give opportunity for the most accurate observation. In the article by Dr. S. B. Hunt, in the *Sanitary Memoirs of the War of the Rebellion* (*loc. cit.*), will be found a most valuable discussion of the causes and treatment of diarrhœa and dysentery, which no one can read without being impressed with the similarity (saving the ages of the patients) of those diseases to the one we are describing. At page 304, Dr. Hunt says: "But in others the disease progressed, became follicular, and finally ulcerative. In the treatment of these, great difficulty was experienced, from the fact that the patient was still exposed to the causes of this malady; and it came to be a fixed doctrine at Southern and Southwestern stations that confirmed cases had no security for cure except by removal to the North. This soon became a governmental policy, and hospitals were established in New England, along the Lakes, and in the Northwest, to which chronic cases were sent in great numbers. Among patients not thus removed, but treated in

Southern hospitals, much vacillation and irresolution were exhibited in the prescriptions of surgeons, as happens in all diseases, the treatment of which by drugs is usually unsuccessful. To trace the history of an individual case was to find that the prescriber had run the round of all remedies, from opium to astringents, from astringents to quinine, from quinine to bismuth, and from bismuth to *nux vomica*, from *nux vomica* to mercurials, returning almost always to opium as the drug, which at least alleviated, if it did not cure."

The *dress* ought to be suited to the weather. It is best to keep on the child, even in hot weather, a very thin and soft flannel skirt, with short sleeves. This should never be removed. A young infant should wear all summer long a thin and light flannel petticoat. A child a year old may have the flannel petticoat removed for a few days when the temperature rises above 85° or 90° , when it suffers evidently from the heat; but so soon as the temperature falls to 85° or below, the petticoat should be replaced. This happens only for a few days in our summer season, and the change should be made with great care, and only under the supervision of an intelligent and watchful mother or nurse.

Exposure to the open air is another point in the prophylactic treatment which is of great importance. In country houses in the summer, a young infant may get nearly as much air as it needs, but in cities the air of houses is much more dull and stagnant, and the child ought to be carried out into the streets and squares for several hours morning and evening. If possible, it should be taken to drive into the open country. Short excursions, by rail or boat, for the children of the poor, who cannot escape from the city in summer, are very useful in carrying the child safely through the summer. But in all such jaunts after health, the parents should so arrange matters that the child shall be as little fatigued as possible. The best plan is to go in the morning and return in the evening, resting through the middle of the day at some point where the child can take rest and midday sleep, which are quite as important as fresh air. Included in this subject is that of *exercise*. This becomes very important when the child is old enough to walk and run, for then an ignorant or thoughtless woman might think the more exercise the better, whereas it is necessary to watch such children very carefully, since, if they are allowed or enticed to take undue exercise, the resulting fatigue becomes a positive cause of diarrhœa. A child of two or three years old should never be induced to take long and continuous walks; it ought to frisk and play, not walk straight ahead, like a man in training; for that kind of exercise, we have remarked, never suits children well.

It has already been stated that one of the most frequent causes of the malady is the attempt to bring up the child on artificial diet, and particularly on one of an improper kind. It is clear, therefore, that to avoid the disease it is necessary that the child should, if possible, be nursed. If this cannot be done, the diet ought to be wisely selected and regulated in all its details by the physician. The one most proper is evidently that which most closely resembles the natural aliment of the infant. For directions as to

diet, we must refer the reader to the remarks upon diet, where we have discussed this point quite fully.

Diet in the Attack.—After the disease has made its appearance, the diet should be very carefully regulated. This constitutes, in truth, the crucial point in the treatment. If the child is nursing, it ought to be confined entirely to the breast, and should the nurse have a large quantity of milk, and the stools exhibit considerable quantities of undigested caseine, it must be somewhat restricted as to the frequency and length of time it is allowed to nurse; in other words, it must be moderately dieted for two or three days. Should there be the least suspicion that the milk of the nurse is unhealthy, it ought to be examined as before directed, and, if found unhealthy in any respect, a new nurse must be provided. If the disease comes on shortly after weaning, and persists for several days in spite of careful diet and treatment, it is safest to restore the child to the breast. When this cannot be done, we must select that form of artificial diet which seems most suitable. The best is, in our opinion, the cow's milk prepared with the solution of gelatine in the manner already recommended, but made very weak for a few days. We have often found it necessary, under these circumstances, to add four and even more parts of water to the milk, instead of two or equal parts, as is the usual custom.

In older children the diet, for a few days, ought to consist of simple milk and water, or of thin preparations of arrowroot, rice-flour, sago, tapioca, or wheat-flour, made with milk, or milk and water, with small quantities of bread, or, if the child refuse such articles, panada, or light beef-tea, or chicken or mutton-water may be allowed. The quantity of food, whatever it be, must be determined very much by the child's instincts. When the appetite continues, we can seldom go wrong in allowing as much of these simple foods as the patient will take. Still, the physician ought to know accurately the amounts that are given, and if he finds the patient taking a full healthy average, or more, it will be best to restrict the quantity somewhat, and offer water frequently, on the supposition that the little patient is taking its liquid food more from thirst than hunger; or else increase the water of the food, if he have reason to believe that the solid matter is in too large a proportion.

THERAPEUTICAL TREATMENT.—We have found a large number of the mild cases that have come under our notice to recover under very simple treatment. When the patient is an infant at the breast, before the period of dentition, of simple direction not to allow it to nurse as much as usual; the use of a warm bath morning and evening, if the skin be heated and the child restless and fretful; the administration of a small dose of castor oil (half a teaspoonful to a teaspoonful), or of spiced syrup of rhubarb in the same quantity, with a half drop to a drop of laudanum, at the beginning of the attack, to remove any undigested food that may be lying in the bowels, followed in one or two days, if the disorder continues, by some simple astringent remedy, generally suffices to effect a cure. When, on the contrary, the case depends on an unhealthy or insufficient milk, when the child subsists entirely on artificial food, and when the disease coincides with the process of dentition, the attack is kept up and aggra-

vated by these causes, and it is more difficult to obtain a cure. In the former case the diet is, of course, of all importance; in the latter the gums must be carefully examined, and if found to be swelled and inflamed, and the teeth near the surface, they should be freely incised. After these matters have been attended to, the kind of treatment will depend on the character of the general symptoms and the violence of the enteritic disorder.

When the pain is violent, the discharges frequent, painful, and mixed with mucus, muco-pus, or blood, and the abdomen tense and painful to the touch, we employ warm baths, poultices to the abdomen, or warm stupes, and refrigerant medicines. Small doses of the sulphate of magnesia and laudanum are very useful; or we may employ spirit of nitrous ether, or solution of the acetate of ammonia with paregoric or laudanum, or the following mixture:

R. Sodæ Bicarb.,	3ss.
Pil. Hydrarg.,	gr. iij.
Tr. Opii Camph.,	gtt. ℥ vel f3j.
Syrupi Simp.,	f3ij.
Aq. Menth.,	f3xiv.—M.

Dose.—A teaspoonful every three or four hours.

The warm bath, used at a temperature of 95° to 97° , twice or thrice a day, is most excellent. It is a good plan to wrap the child, immediately on being taken out of the bath, in a warm muslin sheet, to put over this a light blanket, and let it lie on the lap or bed for twenty minutes or half an hour. During the past two years, the external use of cold water in febrile diseases has been extended widely, and among the affections in which it has been applied are entero-colitis and cholera infantum. We have ourselves limited the use of water to tepid bathing or cool sponging when the temperature was considerably elevated; but excellent results have been obtained by some observers, both here and in Europe, by the use of cold baths (72° to 78° Fahr.), repeated several times during the day when the febrile temperature rose to 103° or 104° . If the temperature should remain elevated, despite the use of repeated cool spongings and the warm bath, the cool baths may be tried, though we should not recommend them quite so cold as above mentioned.

The hot poultice or stupe recommended above should be covered with oiled silk, secured by a towel pinned around the body, changed every three or four hours, and kept on for the greater part of the day, or for several days.

Calomel has been so highly recommended and so long employed in these cases, that we feel some hesitation in saying how often it has disappointed us. Certainly we have found in many children that it was of no evident use, and in the old-fashioned doses of a grain or half a grain, we think it only adds to the irritation of the bowels. In doses from the $\frac{1}{4}$ th to $\frac{1}{2}$ th of a grain combined with small doses of opium and with chalk or bismuth, it is sometimes useful, especially when there is marked gastric irritability coëxistent. We cannot doubt, however, that much of the benefit

that was formerly attributed to calomel in this disease, has been really due to the opiate and astringent with which it was usually combined.

We still have confidence in, and employ, the mixture of blue mass and soda above recommended. It does not irritate, as we have known the larger doses of calomel to do; but on the contrary, when given for thirty-six or forty-eight hours, under the circumstances mentioned, it is frequently followed by an improvement in both the number and character of the stools.

Before quitting this question of the use of mercurials in diarrhœa, we wish to quote the results at which some of the more modern observers have arrived, with the remark, as we pass on, that our own conclusions were much the same twenty years ago as those expressed above. We shall do this even at the risk of being tedious, for we think the point a very important one. In the first place, we shall quote the opinion of one of the ablest of the United States army surgeons, as to the use of this drug during the late war. The writer (*Outlines of the Camp Diseases of the United States Armies as observed during the Present War*, by J. J. Woodward, M. D., Philadelphia, 1863), in the article on Chronic Diarrhœa, a disorder closely akin in many of its symptoms and anatomical lesions to the entero-colitis of children, says, at page 262: "Among the remedies liberally employed in chronic diarrhœa is one which can only be mentioned with disapprobation. This is the mercurials, which are too frequently administered to gentle salivation in the form of blue pill or calomel, combined with opium and ipecacuanha. The authority of some of the most distinguished American medical writers is in favor of the employment of mercurials in the chronic diarrhœa of civil life; yet when it is remembered that even those modern writers, who most warmly advocate their general employment in the treatment of inflammation, recommend them to be discontinued as injurious whenever the process has gone on to ulceration, it would appear that even sound mercurialists would avoid using them in the form of chronic diarrhœa which is most common in the army."

"Practically, it will be found that although in some cases mercurials may succeed, as much less dangerous remedies would have done, in checking the progress of the disease, yet that in the majority of cases their employment is accompanied by an increase of the debility, the loss of appetite, the anæmia, and the general constitutional symptoms, without any diminution in the frequency of the stools. They are, therefore, to be regarded as dangerous and inefficient, and their use in these cases has been completely abandoned by those surgeons who are most successful in the treatment of the disease."

Dr. T. K. Chambers, of London (*Clinical Medicine*, London, 1864, p. 517), in considering the treatment of diarrhœa in which the stools exhibit the products of acute inflammation, says: "The drugs I have most trust in are calomel, ipecacuanha, and carbonate of soda. Of the first and second equal quantities, and a double quantity of the third, may be made into powders, of which from four to six grains, according to the child's age, may be given every three hours. This is a traditionary powder, but it is

right to say that I have in a good many instances lately left out the calomel, and the case has done just as well, if not better, without it."

Dr. J. L. Smith, of New York (*op. cit.*, p. 379), says nothing whatever about mercurials in his article on the treatment of inflammatory diarrhœa, from which we are led to suppose that he does not use them. He, however, quotes Dr. E. H. Parker as giving, when the condition approaches that of dysentery, a mixture consisting of about ten grains of blue mass rubbed up in two drachms of syrup of rhubarb, to which is added one-half teaspoonful of paregoric, and four ounces of chalk mixture. Of this the dose is a teaspoonful every two or three hours. Dr. Parker says that the "blue mass certainly does not act like the calomel, not producing in purgative doses so great prostration, and in small doses it does not lessen the proportion of fibrin in the blood, as in the case with calomel." Dr. Smith's comment on this is: "I have never used this mixture, having been generally satisfied with the effects of the castor oil mixture."

It is unnecessary to say any more upon the use of mercurials, and especially of calomel. We have quoted enough to show that our own opinions find us in very good company.

We regard *opium* as one of the most valuable remedies we have in the treatment of this disease. In a former edition of this work it was stated that some writers objected to its employment in the early stage as injurious, but that we had not been deterred from using it, except in cases in which drowsiness and a tendency to stupor or coma, point to some cerebral disorder; but that when there has been nothing more than irritability, restlessness, and insomnia, when there was evident pain during the discharges, and when the latter have been very frequent, we had always made use of some of its preparations without hesitation, and certainly without injury, but, on the contrary, with very great benefit. Our longer experience confirms us in this view and practice. The propriety of using large doses of opium in the early stages of cholera infantum may well be questioned, as it has come to be by some of the best observers in Asiatic cholera; but this matter will be considered under the head of that disease. In the disorder under consideration, which is one of an inflammatory catarrhal type, we have never seen the moderate use of opium do anything but good. When the nervous symptoms are very marked, if they be of the kind which denote disturbance of the reflex functions of the nervous system rather than those indicating cerebral disorder, we find nothing which answers our purpose so well as this remedy. When, however, there is unusual quiet, tending towards drowsiness or stupor, with contraction of the pupils, we make use of it only with great caution and in very small doses. We are glad to find that Dr. Stokes also employs opium without hesitation. He says: "It is a remedy that requires caution in its exhibition, but one of great utility." It generally lessens the number of discharges, and very often diminishes the heat of skin and frequency of the circulation, by allaying the irritability of the nervous system, while at the same time it greatly promotes the comfort of the child. We have used it in the form of laudanum or paregoric, given in combination with a laxative early in the case, or by enema, and afterwards in that of the Dover's powder or

powdered opium. For a child under six months old half a drop of laudanum is enough to give by the mouth. Of the Dover's powder about a sixth or eighth of a grain may be administered mixed with two grains of chalk, to be repeated every two or three hours, until three or four doses have been taken, or until the child shows some degree of drowsiness from the action of the opium, after which it ought to be suspended for six or eight hours, and then resumed. Or the opium may be given in the form of laudanum combined with the sulphate of magnesia as recommended above. The old-fashioned castor oil emulsion, in the proportion of one drachm in a two-ounce mixture, with half a drop of the deodorized laudanum to each teaspoonful of the mixture, is often very soothing and beneficial. When there is marked tenesmus, with frequent small evacuations, opium may also be used with great advantage by the rectum, either to the exclusion of any in the mixture, or in addition to that, taking care to graduate the quantity by the degree of drowsiness that may be induced. At one year two drops in one or two teaspoonfuls of water or thin starch-water, may be used two or three times a day. In such cases, suppositories are sometimes retained better than enemata. A twelfth of a grain of powdered opium, made up with cocoa butter, may be given instead of the injection. We have learned to be cautious in the use of opium in substance in children under one year of age, and especially under five or six months, whether in Dover's powder, powdered opium, or in suppositories. The difficulty in securing an accurate subdivision into such small doses as are necessary is the chief reason for this caution, and, whenever possible, we prefer the fluid medicine.

Generally speaking the acute constitutional symptoms either subside or disappear under the above treatment, and very often the diarrhœa also ceases and the child recovers. When, however, the diarrhœa persists, it is necessary to resort to two other classes of remedies, upon which great reliance is placed in the treatment of this affection. These are *astringents* and *absorbents*, of which the most important are prepared chalk, powdered crab's-eyes, bismuth, acetate of lead, rhatany, kino, and catechu. The chalk may be used in the form of the officinal *mistura cretæ*, a teaspoonful of which is given after each loose evacuation, or several times a day. When the case is severe, the efficacy of this remedy is much increased by the addition of tincture of *krameria*, in the proportion of a drachm to two or three ounces of the mixture, of some opiate preparation, or of ten or fifteen drops of the aromatic syrup of galls (to be described presently) to each teaspoonful. Chalk may be used also with great advantage, as stated above, in powder, combined with Dover's powder.

The powdered *crab's eyes*, it has been thought, will sometimes arrest cases in which prepared chalk fails to produce any effect. It is generally employed in mixture. The formula we employ is the following:

R. Ocul. Cancror. Pulv.,	3j.
Pulv. Acaciæ,	3ij.
Sacch. Alb.,	℞j.
Aquæ Fontis, Aquæ Cinnamom., āā	f 3 iss.—M.
A teaspoonful to be given four, five, or six times a day.	

M. Bouchut recommends the following prescription of Hufeland's:

R. Ocul. Cancror. Pulv., gr. x.
 Aquæ Fœniculi, Syr. Rhei, āā fʒss.—M.
 Give a teaspoonful every hour.

Subnitrate of bismuth has been highly recommended, for a number of years past, as a remedy in diarrhœa. Dr. Woodward (*op. cit.*, p. 258) quotes Assistant Surgeon Dr. John B. Trask, U. S. A., as lauding it very highly in the chronic diarrhœa of the armies during the late war, and in California and Oregon, especially in those cases in which there is nausea or other disorder of the stomach. Dr. Woodward states that "he has given it a fair trial, and while he is far from regarding it as specific, believes it to be a most valuable article in both simple, irritative, and in chronic diarrhœa." Dr. Trask prefers to give the whole quantity for the day in a single dose; but Dr. Woodward states that this view does not correspond with the general experience on the subject. It may be given in doses of one to two grains, to children one year old, every two or three hours. It can be administered in powder with sugar alone, or combined with prepared chalk, or in mixture with simple syrup, or ginger or acacia syrup, and some aromatic water. We have employed it quite frequently, but, on the whole, have not found it so effective as we had been led to hope.

Acetate of lead has been highly extolled by many writers in the treatment of the diarrhœas of children. We have had but little experience in its use, and are, therefore, unable to offer an opinion in regard to the influence which it may exert. It may be given in doses of from a sixth to an eighth of a grain, alone, or combined with chalk or Dover's powder, every two hours. *Krameria*, *kino*, and *catechu* may be exhibited alone, in the form of infusion or solution, or they may be given in conjunction with the chalk mixture. We have frequently employed the tincture of *krameria* in the latter way, and believe it adds very much to the efficacy of the remedy. One or two drachms may be added to three ounces of the mixture, and the usual dose given. We have used, with much advantage, either alone or with chalk or crab's-eyes mixture, an aromatic syrup of *galls*, in the dose of from fifteen to forty drops three or four times a day, or, when the discharges are very frequent, every two or three hours. It is prepared according to the following formula:

R. Gallæ Opt. Pulv., ʒss.
 Cinnamomi Pulv., ʒij.
 Zingib. Pulv., ʒss.
 Spt. Vini Gall. Opt., Oss.—M.

Let the ingredients stand in a warm place for two hours, and then burn off the brandy, holding some lumps of sugar in the flames. Strain through blotting-paper.

Nitrate of silver has been highly recommended as a remedy of late years by several writers. It is given both internally and by enema. The modes of administration will be described in the remarks on the treatment of the chronic form of the disease.

Revulsives are often of much service in the treatment of this, as of almost all the diseases of childhood. When there is much restlessness and irritability, with heat of the head and trunk, and coolness of the extremities, it will be found that mustard foot-baths or sinapisms to the extremities, often allay these symptoms, and greatly comfort the little patient. When the abdomen is tense and painful, and the discharges preceded or accompanied by movements or crying indicative of pain, the application of a poultice of mush and mustard from time to time, to be followed by a simple mush poultice, sometimes acts very usefully.

Tonics and stimulants are often necessary in weak and delicate children from an early period in the attack, and in those who are stronger, after the disease has lasted for some time, and the acute symptoms have ceased, and been followed by weakness and exhaustion. The best tonic is, probably, *sulphate of quinine*, in doses of from a quarter of a grain to a grain three times a day, continued for one, two, or three weeks if necessary. Old *brandy* has answered better in our hands as a stimulant than wine, wine-whey, or any of the tinctures. It may be given to the youngest children in doses of from five to ten drops every two hours, or a teaspoonful may be added to a wineglassful of sweetened water, and a teaspoonful given whenever the child will take it. We have been obliged, in several cases, to continue the use of the brandy for three, four, and five weeks. At the time when we are obliged to resort to this class of remedies, it is almost always necessary also to pay attention to the improvement of the diet. The proportion of milk to water ought to be increased, if it has been small heretofore; and we should employ every means to induce the child to take a sufficient quantity without overloading the stomach. At this stage small quantities of animal broths are proper, or the child may be allowed to suck pieces of juicy meat, or to eat very finely minced meat of chicken or mutton. The diet is in fact a most important part of the treatment at this period. Dr. Stokes says of it, that "many children are lost by the practitioner neglecting this point."

Occasionally, indeed quite frequently, vomiting becomes a most troublesome symptom in enterocolitis. When it occurs at rare intervals, and without much distress to the patient, it needs no attention, since it is to be supposed that the physician has already arranged the hygienic and therapeutical treatment to suit the ordinary conditions of the disorder. But when vomiting becomes frequent and violent, so that the child rejects a large proportion of all that is given to it, and when, between the acts of vomiting, the little thing refuses almost everything that is brought to it, all its usual foods, medicines, and sometimes even water, it becomes evident that there must be more or less nausea which causes loathing of food, and the symptom becomes a serious complication which requires special attention and treatment. In such cases, there is no use in forcing food or drugs, which it loathes, upon the child, unless all other means have failed, when, of course, we must attempt to make it take concentrated foods in small doses. The better plan, at first, is to change the diet *in toto*,—to abandon milk and all its preparations for one or two days,—and to give light beef or chicken tea, just touched with salt, or raw beef, or, if this also is refused,

cold extract of beef in one or two tablespoonful or wineglassful quantities, or pieces of juicy and rich beef, very slightly cooked, to be sucked. Or we may try small portions of yolk of egg, hard boiled, or what we have often found was eagerly taken in such conditions, wine-whey, of which we have given, in the second year of life, as much as a tumblerful in twenty-four hours, and this without the slightest effect of undue stimulation, febrile heat, or excitement. Sometimes, when the child persistently refuses its ordinary milk, or vomits it so soon as taken, it will drink willingly, and retain very well, lime-water and milk, in the proportion of one of the former to two or three of the latter, with just enough brandy to change the taste. We know that some medical men object entirely to the use of stimuli in children on two grounds: 1. That alcohol has no remedial power whatever, or that it is positively injurious in all cases. 2. That its use tends to produce a pernicious taste for stimulants and invites the habit of drunkenness. To the first objection we can only reply that our observation and experience have led us to a different conclusion, and that, when employed in certain conditions of the vital powers, which we have carefully endeavored to describe, stimuli are of the highest value as a therapeutic means. To the second we reply that we have never, so far as very careful observation goes, produced a drunkard by any use we have made of them. We agree that physicians ought to be careful not to employ them in any attractive form, as a long-continued remedy, in children over six or eight years of age. When we desire to use any form of alcohol in a chronic case in children over the age mentioned, we give some of the bitter tinctures or elixirs. When brandy is to be used, we always order the oldest and most delicate that can be procured. As to the quantity, this must depend on the age of the patient, the instinct and idiosyncrasy of the child, and the degree of severity of the case. At the age of six months, from ten to fifteen drops may be given every two or three hours in two or three ounces of the lime-water and milk; and at one and two years, from twenty to twenty-five drops in from four to six ounces of the milk food every two, three or four hours. It may be a sign of the old Adam in the little sufferer, but we have often known children to take, for days together, milk with brandy in it, who would not touch the milk without this addition. We cannot but think that in such cases it is an instinct for a useful agent, like the appetite of patients in typhoid or typhus fever, in certain of their phases, for wine or brandy, which disappears when the necessity for it passes away, as has been so well described by Dr. Corrigan, of Dublin, in his able essay on the treatment of Irish typhus.

Under these circumstances, all medicines which disgust the child must be laid aside. A bitter, or nauseous, or gritty dose will, in such states, surely cause vomiting, as, in older persons, under such conditions, does an odor or taste, or even an idea. We have seen a little infant, sick with diarrhœa, who was sitting languidly upon the floor, made to gag and retch by chancing to pick from the floor a piece of softened bread. The impression produced upon the tactile sense of the fingers by the wet and mushy substance caused sickness at once, as the filing of a saw sets the teeth of a delicate nature on edge, and brings water into the mouth. All

offensive and bitter doses must therefore be abandoned. We have often used, in such cases, the following prescription with much benefit:

R. Liq. Morph. Sulphat.,	℥xxxij.
Acid. Sulph. Dil.,	℥xv.
Curaçœ,	f℥ij.
Aquæ,	f℥xiv.—M.

Dose.—A teaspoonful every hour or two hours, at the age of six months to a year.

For older children, the proportions of the opiate and acid must be increased. When the nausea subsides or passes away, or when the child becomes drowsy, the intervals between the doses must be lengthened, and as the symptoms disappear, the other remedies necessary for the diarrhœa may be resumed, and so too of the food. Dr. J. L. Smith, of New York, states that the best remedy he has used for vomiting is the neutral mixture, as follows:

R. Potassii Bicarb.,	gr. xxv.
Acid. Citrici,	gr. xvij.
Aquæ Amygd. Amar.,	f℥j.
Aquæ,	f℥ij.—M.

One teaspoonful to a child from eight to twelve months old, repeated according to the nausea or vomiting.

We have ourselves more frequently directed the freshly prepared effervescing draught, made with lemon juice and bicarbonate of potash, and have found it very useful.

Creasote is often of great value in relieving such nausea. It may be given in the dose of an eighth of a drop every three or four hours at the age of one or two years, and may be administered either in a teaspoonful of lime water, further diluted with a teaspoonful of water or of milk, or in the following form:

R. Sodii Bicarb.,	gr. xxxij.
Creasoti,	gtt. iv.
Pulv. Acaciæ, Sacchar., āā	q. s.
Spt. Lavandulæ Comp.,	f℥ij.
Aquæ.	q. s. ad f℥xiv.—M.

Dose.—A teaspoonful in a little water three or four times daily.

In some cases no remedy will allay the irritability of the stomach so promptly as very minute doses of calomel (gr. $\frac{1}{4}$ every two hours at two or three years of age) placed dry on the tongue.

TREATMENT OF CHRONIC ENTERO-COLITIS.—The management of the hygiene of the patient is more important than any other part of the treatment, in this, as in nearly all the diseases of the digestive organs in children; for cases will often recover when the diet, drinks, and exercise are properly regulated, without the use of any drugs whatever, whereas, most assuredly, but a small proportion of them would terminate favorably under the best and wisest therapeutical medication, were the hygiene of the child neglected. The remarks that have been made as to the diet most proper in the acute form will apply here. If the child have been weaned only a

few weeks before the time at which we are consulted, and the case is at all serious, it is best to advise a wet-nurse. We have several times known cases of the disease which had resisted the most carefully managed artificial diet and therapeutical treatment, recover in a few days after the child had been restored to the breast. It is often, however, impossible to follow this course, from the refusal of the parents to obtain a nurse, or of the child to take the breast of a stranger, and we are obliged to rest content with artificial food. Cow's milk, in some form, makes the best diet under these circumstances. For full information we refer to the chapter on food.

In some of these cases beef-tea or chicken-tea will be taken willingly by the child and retained, when milk preparations are turned from with disgust or rejected by vomiting. Beef-tea is best made after the mode laid down by Dr. Letheby. Equal weights of lean beef, cut into small pieces, and cold water (a pound to a pint), are infused together for half an hour. They are then put into a pipkin, placed near the fire, and allowed to heat gradually, so as to reach the boiling point in fifteen minutes. They are allowed to boil a few minutes—two minutes are enough. The water is then poured off the beef, the beef squeezed, and the water added to the rest. The amount of sediment here is very small and soft, and is to be given with the broth. Salt, of course, is added.

Chicken-tea is made by taking half of a small chicken, or the leg and thigh of a large one, removing the skin, breaking the bones, and simmering in a pint of water down to half a pint. Salt is added. It is quite remarkable with what pleasure and avidity young children will take this thin food.

It sometimes happens that the child will refuse everything that has been mentioned, and yet the prostration and emaciation are such as to make it essential to procure some aliment that it will consent to take. We have, under such circumstances, given small portions of bread and butter, or stale sponge-cake, with weak brandy and water, if the child is old enough to swallow solid food. If the white of an egg be stirred in a small glass of water, the child will usually drink it freely without recognizing the presence of the albumen, and we are thus enabled to administer a considerable amount of nutritious food by giving the whites of two or three eggs in the course of the day. Sometimes the child will eat small quantities of meat, and when this has been the case, we have not hesitated to allow a chicken-bone, with a little meat attached to it, or a piece of ham, or better still, a portion of roast beef, or of the tenderloin of beef-steak, to be held in the hand and sucked; or we may give the white meat of chicken cut up very fine, or torn into the finest shreds. Of the latter about a teaspoonful is sufficient for the first day, given with a little brandy and water. The quantity can be gradually increased afterwards. We have of late years also given small quantities of raw beef in many cases, minced very fine and flavored with salt, or prepared in the manner described below,¹ and have found it to be readily digested and to agree well with

¹ The use of raw meat in the diarrhoea either of infants deprived of their mothers' milk, or of weaned children, was recommended by Weisse, of St. Petersburg, as long

the little patients. There is another article which we have sometimes given when children have been exhausted for want of food, and when they require constant change in order to be tempted to take it. This is the yolk of a *hard-boiled* egg, which has the advantage of being very nutritious if digested, and of not being injurious should it happen to pass into the bowel in the crude state, as it falls into a state of fine powder, which is not irritating to that organ.

The quantity as well as character of the food is of the utmost importance, and should be strictly regulated by the physician, and attended to by the mother or nurse. As a general rule the child may be allowed as much as it wants of proper food, since the appetite is almost always greatly diminished, and it is not likely, therefore, that too much will be taken. If, however, there is a disposition to nausea or vomiting, or if the appetite remain as good as usual, the quantity must be restricted. The difficulty in most cases is to get the patient to take enough, and not to prevent it from taking too much, for we have very often ascertained, upon careful inquiry, that the quantity was entirely too small to support the strength of the constitution. This is a matter of great importance. We believe that the life of the patient often hangs upon the physician's action in such cases. He should know, by the most minute and thorough inquiry, just what the patient is taking each day. A child six months old, as we have shown elsewhere, ought to take from a quart to three pints of liquid food per day, and one of a year old as much or more than this made with a larger proportion of milk, or in connection with some solid food. Now, we have frequently known children with this disease to take not more than two or three gills a day, which is manifestly quite too little to sup-

ago as 1840 (Oppenheim's Journal). Of late years it has been extensively used with excellent results, and is highly praised by Trousseau and other eminent authorities. The administration of the muscular tissue itself appears much more useful than any form of beef essence or soup, probably for this reason among others, that these fluids pass too quickly through the intestinal canal. The best meat for the purpose is the fillet of beef, though fine mutton may also be used. It should be cut very fine, and, according to Trousseau, pounded in a mortar and strained through a sieve or cullender: the pulp, thus separated from the cellular texture of the meat, may be rolled into small balls in salt or powdered sugar.

The quantity upon the first day should not exceed three drachms, given in divided doses; but it may be doubled on the successive days, until young children may take from six to ten ounces a day. Under this regimen the diarrhœa frequently ceases, and the children quickly recover their plumpness and natural spirits.

Trousseau calls attention to the fact that the stools are frequently red and fetid at first, even when the nature and abundance of the diarrhœa have already undergone a favorable change.

In the second article upon this subject (Jour. für Kinderkrankheiten, January and February, 1858) Weisse calls attention to the fact that in many children who had been treated by raw beef, tapeworms have been developed. As these worms were all specimens of *tania solium*, which is not indigenous in St. Petersburg, it is probable, as suggested by Von Siebold, that they had been conveyed in the undeveloped state in the flesh of oxen brought from distant points. We are not aware that this unfortunate consequence has been observed frequently in other localities, and certainly in the quite numerous cases in which we have ourselves administered raw meat to children, no entozoa have been developed.

port life for more than a short time. In such cases, the persevering use of stimulants and tonics, and changing the food until something is discovered that is accepted willingly by the child, makes the essential part of the treatment.

In connection with this most important matter of the food, we will again quote from Dr. S. B. Hunt (*op. cit.*, page 305), to show the results of his experience in the use of foods in chronic inflammatory diarrhœa in the army. For the sake of any non-professional reader, we will state that by albuminoid food Dr. Hunt refers to meat, meat broths, eggs, etc.; and by antiscorbutic food he means tomatoes, fresh fruits, onions, etc. Dr. Hunt says: "The value of drugs was, perhaps, overestimated in this, as in all other diseases of assimilation, and only a careful avoidance of the original causes of the malady, and an equally careful recognition of their continued existence in the system, could secure any degree of success. The scorbutic and malarial taints were almost uniformly present, the former very frequently in as pronounced a form as the latter. The bowels, enfeebled by the inflammatory process, were unable to perform their normal function of the digestion of starches, and the diet, therefore, became necessarily albuminoid. A full nutritious diet of albuminoid and antiscorbutic food assumed the first importance in the treatment. Coupled with this came pure air and absolute cleanliness. And, with these hygienic measures alone, when they could be properly enforced, it was possible to treat chronic diarrhœa and dysentery with a fair degree of success, even in the great heats of a Southern summer." These views confirm what we have said above, that milk, meat, raw or cooked, broths, eggs, gingerbread, tomatoes, bread and butter, and we may add currant-jelly, make the best food for children over two and three years of age. Even in children of eight months and a year or upwards of age, milk and beef or chicken-tea ought to form the chief diet. The starches, such as arrowroot, barley, wheat preparations, etc., do not answer, except in very small quantities cooked in milk. We saw one child, a year old, weaned in August in consequence of the illness of the wet-nurse, whose life was apparently saved in dysentery by Liebig's cold extract of beef, and by its fortunately having developed a strong taste for the sucking of large pieces of rapidly and slightly cooked beef-steak.

The *therapeutical treatment* of the chronic form consists principally in the administration of tonics, astringents, and absorbents. Of these the most important are bismuth, powdered chalk and crabs'-eyes, and the different vegetable astringents, which have already been noticed in the remarks on the acute form. These are to be given in the manner there recommended, and it is therefore unnecessary to repeat what has already been said. In addition to these there are some remedies which are particularly adapted to the chronic form of the disease. Amongst them is *nitrate of silver*. Dr. Eberle (*op. cit.*, p. 251) says he has found its internal administration to produce the happiest effect in a few instances. His prescription was a grain of the nitrate dissolved in an ounce and a half of gum arabic water, with the addition of twenty drops of laudanum. The dose was a teaspoonful three times a day. He adds that he has never "known

the slightest inconvenience to result from the use of this article in chronic mucous inflammation of the bowels, when administered in a mucilaginous solution and in very small doses." It has been much used of late years in France. MM. Trousseau and Pidoux recommend its internal use in the chronic diarrhœas of children occurring during dentition, after bismuth, powdered crabs'-eyes, and diet have failed to effect a cure. Their formula is as follows :

R. Argenti Nitrat.,	gr. $\frac{1}{2}$.
Aquæ Destillat.,	fʒvj.
Syr. Sarsap.,	fʒijss.—M.

To be given in eight or ten doses.

At the same time, they employ an enema composed of a grain of the nitrate in three ounces of distilled water. It is highly recommended also in these cases by Hirsch, of Königsberg. His formula is as follows :

R. Argenti Nitrat. Crystal.,	gr. $\frac{1}{4}$.
Aquæ Destillat.,	fʒij.
Acaciæ Pulv.,	ʒij.
Sacch. Alb.,	ʒij.—M.

A teaspoonful of this mixture to be given every two hours, and an enema, consisting of a quarter of a grain of the salt, with mucilage and a little opium, to be administered (*Ranking's Abst.*, No. VI, p. 61). We have for a number of years past used nitrate of silver so frequently in this disease, and with such excellent results, that we can confidently recommend it. Internally it is best given in solution, in a thin and delicately made mucilage or syrup of acacia, in the dose of gr. $\frac{1}{4}$ to gr. $\frac{1}{10}$, three or four times daily, for a child two or three years of age. Each dose should be given in about two or three fluid drachms of liquid, to which whatever amount of deodorized tincture of opium is considered desirable may be added. The best time for its administration is towards the close of digestion, or about one hour after food has been taken. We have also given it very frequently in the form of enema, in cases where it was apparent that the rectum and the lower part of the large bowel were considerably affected, and under such circumstances the happiest effects may be secured. The dose and mode of administration are stated at the close of the following paragraph.

Dr. Woodward (*op. cit.*, p. 264) says, in his article on the treatment of the chronic diarrhœa, which was a true entero-colitis, that "by far the most valuable local measure is the employment of solutions of the mineral astringents as enemata." He mentions sulphate of copper, nitrate of silver, sulphate of zinc, and acetate of lead, but thinks that the sulphate of copper and nitrate of silver are probably the most efficient. The strength he recommends is of one or two grains to the ounce of water, of which from one to six ounces may be thrown into the rectum two or three times a day. He advises that, when the rectum rejects the injection immediately, twenty to forty drops of laudanum be added to each enema, that the

injection be thrown carefully into the bowel, and the nozzle of the syringe be withdrawn as gently as possible, in order that the fluid may be retained at least for some little time. We quote these statements, not to induce the use, in children, of solutions of one or two grains to the ounce, but to draw attention to one of the means, the ability and advantage of which bore the test of the vast army experience in this most severe and troublesome disease. In children it is best to begin with one or two ounces of a solution of the strength of gr. $\frac{1}{4}$ of the nitrate of silver to an ounce of water or thin mucilage, repeated morning and evening; and, if this gives no pain, or but little, and does not produce the desired benefit, the quantity may be increased, and the proportion may be doubled, or, after two or three trials, brought up to that of a grain to an ounce. It is well to add a suitable amount of deodorized tincture of opium, carefully adapted to the age of the child and to the amount of opiate that is being given by the mouth.

Another excellent remedy in the chronic diarrhœas of children, one from which we have sometimes obtained very satisfactory effects, is the solution of the *nitrate of iron*. It is given in doses of from two to five drops three times a day, in sweetened water, at the age of one or two years.

The following formula is recommended by Dr. Eustace Smith. We have used it ourselves in several chronic cases, and have been much pleased with its effects:

R. Liq. Ferri Pernitrat.,	f 3 ss.
Acid. Nitric. Dil.,	f 3 ss.
Syr. Zingib.,	f 3 j.
Aq. Anethæ.	q. s. ad	f 3 iij.—M.
A teaspoonful every six hours at one year of age.						

We have found a teaspoonful every three or four hours not too much at three and four years of age.

Creasote also has been highly recommended, and we have used it with advantage in cases attended with nausea, flatulent distension of the bowels, or a very fetid state of the discharges. It may be given in the manner perscribed on page 434, or subnitrate of bismuth may be substituted in this mixture for the soda.

Bouchut recommends enemata of from ten to twelve grains of extract of *rhatany*, or six to ten of *tannin*, in about five to seven ounces of some vehicle.

Sulphuric acid has been found very useful in the treatment of this affection, and by some authors, as, for instance, Pollock (*Trans. Amer. Med. Assoc.*, vol. viii, p. 260), has been given in large doses as the sole remedy.

We have never used it in this manner, but for some years past have been in the habit of employing it in the following mixture with excellent results. In cases of diarrhœa, showing a disposition towards dysentery, as often occurs in entero-colitis, and especially when the stomach has been irritable, so as to bear other medicinal substances badly, we have found this combination very beneficial:

R. Acid. Sulph. Arom.,	gtt. xlvijj.
Tinct. Opii,	gtt. xij, vel xxiv.
Syr. Krameriaë,	f 3ss.
Aq. Fluvial.,	f 3ijss.—M.

A teaspoonful every two hours.

It should never be forgotten in the treatment of chronic diarrhœa in children, that the most important point of all is the regulation of the diet and other hygienic conditions. We are fully convinced that we have seen several children saved from death by attention to these points, and by the persevering and careful employment of tonics and stimulants. It often happens, after the disease has lasted for some weeks or months, that the powers of the stomach are almost wholly lost. The child either refuses food or takes so little that the quantity is evidently insufficient to carry on the vital processes, or the greater part of what is taken is rejected by vomiting, or, lastly, much of it passes off through the bowels, and appears in the stools in an undigested state, forming what is called lientery. If this condition of things is allowed to continue, the emaciation and exhaustion make rapid progress, and the case must soon terminate fatally. Under these circumstances all the ingenuity and skill of the physician are required to find articles of diet of a digestible and nutritious kind, which shall, at the same time, wake up and tempt the patient's worn-out and perverted appetite. There is almost always present more or less nausea, which keeps the patient on the sharp edge of vomiting. It is worse than useless for the physician to direct the mother to give, in such a case, doses of an ill-tasting or nauseous medicine. Either they would not be given more than once or twice, or, if persisted in by too believing a mother, they would cause vomiting or retching, and do more harm than good. We must depend chiefly, in such cases, on doses of the oldest and most delicate brandy that can be found, of which from one to two teaspoonfuls may be put into a wineglassful of cold water, and the whole given by teaspoonfuls in the twenty-four hours; or fifteen- to twenty-drop doses of the elixir of Peruvian bark every three or four hours may be used; or solution of pepsin, in half-teaspoonful doses three times a day; or, two or three drops of tincture of nux vomica in sweetened water three times a day, if the bitterness does not cause nausea or increase the loathing. In such cases, wine of iron, in doses of twenty drops to a fourth of a drachm, with syrup of tolu and caraway-water, will sometimes do exceedingly well; or the following, which has sometimes succeeded in our hands:

R. Tr. Ferri Chlorid.,	f 3j.
Acid. Acet. Dil.,	f 3j.
Liq. Ammon. Acetat.,	f 3ij.
Syrupi Simp.,	f 3ss.
Aquæ,	f 3ij.—M.

Dose at four years, a teaspoonful, and under that age, half a teaspoonful, three or four times a day.

In some very obstinate cases, especially where there is any reason to suspect the existence of a malarial element in the case, from half a minim to one minim of Fowler's solution of arsenic, with the wine of iron, three

times a day, has been very serviceable. Whilst this is being done, an occasional dose of anodyne, just enough to tranquillize without stupefying, may be given. If the rectum will retain it, it is better to give it by enema. In some cases we have found the aromatic syrup of galls, given with brandy, to be taken by the child without any difficulty or disgust; and strange to say, we have found occasionally that an emulsion of cod-liver oil made very weak, from two drachms to half an ounce in a three-ounce mixture, flavored with oil of cinnamon or of partridge-berry, and given in teaspoonful, and afterwards in dessertspoonful doses, three times a day, could be taken readily, and with excellent results.

Gentle exercise each day in an easy carriage, or in a baby-carriage, is very useful when properly managed. It is very possible to have too much of it, and this does more harm than good. If the child comes home fagged, it has been injured. If it return a little wearied, and disposed to sleep, it has been benefited. Exposure to the open air, under the shade of trees or in a piazza, through much of the day—taking the daytime nap in this way—is useful. In severe and tedious cases, change of residence, from the interior to the seashore, or, if this have failed, to some considerable altitude, will often cure when nothing else will. In one case in this city, which had lasted with but short intervals for two years, we obtained a perfect cure by persuading the parents to send the child into an elevated part of the country in the month of May, where it was kept until July, after which it was removed to the seaside until the end of August. Nothing was done in the meantime except to regulate the diet most carefully, and to keep the child the greater part of the day in the open air.

ARTICLE III.

CHOLERA INFANTUM.

GENERAL REMARKS.—In the early editions of this work we failed to draw with sufficient clearness the distinction between what we think ought exclusively to be called cholera infantum, and the much more common disorders which are properly styled simple and inflammatory diarrhœa or enterocolitis. In this we did but follow the practice of most American writers, and the custom of the day. Indeed, many physicians amongst us are still in the habit of designating the various intestinal disorders of children so frequent during the summer heats, under the common title of cholera infantum. We believe, on the other hand, that a large majority of the deaths registered in our mortality returns under this name, are the result not of a true choleraic disease, but rather of simple diarrhœa or enterocolitis. We have, however, only too often to contend with a disease in children which deserves the title of cholera, which is the analogue of cholera morbus in the adult, and which is the disease we propose to consider in the present chapter.

DEFINITION; SYNONYMS; FREQUENCY.—We can define cholera infantum only by an enumeration of its most specific characters, and we shall do this very much in the words in which Dr. Aitken describes epidemic cholera. Cholera infantum, as we understand it, is characterized by the occurrence, almost solely during the summer months, in young and generally teething children, who have been previously either healthy or the subjects, for a longer or shorter time, of simple or inflammatory diarrhœa, of sudden muscular debility, occasional nausea, spasmodic griping pains in the bowels, depression of the functions of respiration, and an appearance of faintness; copious purging of thin, serous fluid, or of large watery and fetid evacuations, succeeded by more or less obstinate vomiting, coldness and dampness of a part or of the whole surface of the body, coldness and lividity of the lips and tongue, cold breath, a craving thirst, a feeble rapid pulse, difficult and oppressed respiration, with extreme restlessness, diminished or suppressed urinary secretion, pallor of the entire surface of the body, a sunken and pinched countenance, weakness of the cry or partial aphonia, and collapse, more or less complete, which may prove fatal, or be followed by reaction and speedy recovery, or by a subsequent more or less severe and obstinate simple or inflammatory diarrhœa.

This disease is not so common as simple and inflammatory diarrhœa, most cases of which have been hitherto, as stated above, improperly grouped under the common name of summer complaint. Though rare in Europe, in comparison with its frequency in this country, it is easy to recognize from the descriptions, the identity of some of the cases called by Billard follicular enteritis, by Barrier apyretic and febrile follicular diacrisis, by Rilliet and Barthez, in their second edition, choleriform gastro-intestinal catarrh, and by Copland, the choleric fever of infants, with the true cholera infantum of America.

It is impossible to determine its real frequency amongst us, for the reason that fatal cases of simple diarrhœa and entero-colitis, are so generally included in our mortality returns, with those of the true choleraic disease, under the common title of cholera infantum or summer complaint. That it is a frequent cause of death is shown, however, by the tables of Dr. Emerson (*Am. Jour. Med. Sciences*, vol. i, 1827), wherein it appears that from 1807 to 1827, 3576 deaths from cholera, under five years of age, were returned in this city; of course many of these deaths were from a true choleraic disease. This is the largest number of deaths from any one disease given in the table. The next largest item of mortality is under the head of convulsions, of which it appears that 3192 died in the same period of time. During the five years, from 1876 to 1880 inclusive, there occurred in this city, 37,709 deaths from all causes, under fifteen years of age. Of this total, 4547 died of the so-called cholera infantum, which is the largest number of deaths from any one disease. After cholera infantum the largest number of deaths was caused by convulsions (3464) and by marasmus (3386). We also refer the reader to the table given at pages 406–407, obtained from the Board of Health of this city, exhibiting the mortality under five years of age from cholera infantum, diarrhœa and dysentery.

with the total mortality at all ages, and with the mean temperature of each month.

CAUSES.—In discussing the causes of cholera in children, we meet again the difficulty so often alluded to, viz., the custom in this country of classing in mortality returns, all the deaths from intestinal affections in childhood, under the common title of cholera infantum or summer complaint. Our own experience leads us to the conviction that the causes are the same as those of simple and inflammatory diarrhœa, acting with greater intensity. When that cause, or those causes, whatever they may be, act with moderate force, the result will probably be a simple or inflammatory diarrhœa. When, on the contrary, the causes are intensified in degree, the case will be apt to take the form of choleraic disorder. Thus *heat* is one of the most influential of these causes. So long as the atmospheric temperature is moderate, the resulting disorders will probably take the form of simple or inflammatory diarrhœa. But let the temperature rise to 85° or 95° Fahr., or even higher, as happens occasionally in our summers, and continue at that height for three or four days, and children previously well will be seized with the true choleraic forms of diarrhœa, whilst those who are already suffering with simple or inflammatory diarrhœa, are prone to have these milder diseases assume suddenly the choleraic type.

A glance at the table above referred to, shows most plainly the effect of heat upon the mortality from bowel diseases in children, under five years of age. It will there be seen that, in the two months of July and August, when the mean monthly temperature is between 75° and 80°, the mortality from cholera infantum rises to between two and four hundred, and even over; whilst during the cool months, as January, February, November, and December, when the mean monthly temperature is between 30° and 40° generally, only one, two, three, or none at all, are reported. This table shows also, what we have so frequently remarked upon, that most of the fatal cases of bowel disease, in early life, are classed in the medical returns of this city, under the common title of cholera infantum, whereas, we are sure from our own personal experience, that many of these deaths would be more correctly referred to simple or inflammatory diarrhœa, or enterocolitis. Thus, in the very months when three and four hundred deaths are grouped under the title of cholera infantum, only from fifteen to twenty, or a little over, appear usually under the term diarrhœa.

Diet.—Improper diet is another frequent cause of choleraic disease in hot weather. Sudden weaning, a change in the character of the artificial food, the unfortunate use by accident, or by the carelessness of the nurse, of unwholesome milk, of improper vegetables, or, as not unfrequently happens, of green or unripe or unhealthy fruit, as apples, currants, gooseberries, or blackberries (instances of all of which we have ourselves met with), will sometimes bring on, in a very few hours, the most violent attacks of cholera, or convert a previously mild and comparatively safe diarrhœa into the more violent form of disease we are considering. These results are especially apt to follow such accidents or imprudences in large cities, where the hygienic conditions are always in summer of a kind to

invite the more violent and dangerous forms of intestinal disorder. In fine, the conditions which have been ascertained to be most certain to produce epidemic cholera, when that disease is present in a locality, are those which develop cholera in children.

To put before the reader the conditions most certain to cause cholera in children, we cannot do better than to quote from the *Report on Epidemic Cholera to the Citizens' Association of New York*, in 1865, the localizing causes of cholera.

These are:

1. Decaying organic matters, bone, hide, fat and offal houses, neglected stables, putrescent mud and filth.
2. Bad drainage, local dampness, malaria.
3. Obstructed sewers, filthy streets, gutters, stables, garbage, and cess-pools.
4. Water and beverages in any manner contaminated by putrescent organic matter, particularly by any soakage from privies.
5. Neglected privies and putrefying excrement.
6. Overcrowding and neglect of ventilation.

It is just where these conditions are most rife that choleraic diseases in children are most apt to occur. Amongst the poor, who inhabit the crowded quarters of cities, where the streets and alleys are small and narrow, where heaps of decaying vegetable and organic matters abound, where water is scant, or scantily used, where ventilation, from the manner in which the streets are laid out, and from the crowding together of buildings, is necessarily imperfect, we have the most numerous and the severest forms of the disease. Add to these the small size of the houses, the low ceilings, the small and few windows, and the interior arrangement of the rooms, which is such that a thorough draught is unattainable, and we need not wonder at the prevalence of the disease. It is amongst the poor, too, that the food is often of necessity, as well as from ignorance and recklessness, of the most improper kind, and not unfrequently insufficient in quantity.

But not only the poor, in their unhappy lot, suffer from this disease. The children of the rich, with all the advantages of the most wholesome hygienic appurtenances which ease and knowledge can supply, are apt to contract it if they remain in town during the hot summer months. So well is this known, that most families in easy circumstances leave the city for the seaside or the interior, so long as their children are young, remaining absent usually from the middle of June to the middle or end of September. It is nevertheless true that, whilst all the residents in our cities during the summer season are liable to see their young children suffer from this disease, those who are so fortunate as to occupy large and airy houses in the best and cleanest quarters, and who follow a wise system of hygiene as to diet, water, dress, and exposure to fresh air, escape with much more certainty the disease than those who are compelled by the necessities of their position to submit to the unhealthy conditions mentioned above. For further information, and especially for certain opin-

ions in regard to the part that unwholesome milk may play, the reader is referred to the article on the causes of entero-colitis.

Dentition.—We believe this also to be a most powerful predisposing cause of the disease, and yet it is less influential than age, for vital statistics show that it is about twice as fatal in the first year as in the second, though the process of dentition is certainly more active and continuous in the second than in the first year. We have rarely observed it before the beginning of the process of dentition, and it is certainly very rare after its completion.

Age, as has just been stated, exerts a strong influence in the production of the disease. In the tables of Dr. Emerson, the cases of cholera infantum and cholera morbus are included under the one head of cholera, but as all cases of the disease under five years of age are called cholera infantum, the want of the distinction does not make the statements less useful to us. From them it appears that there were 2122 deaths in the first year, 1186 in the second, and only 268 between the second and fifth. Between five and ten years, only 52 cases are noted, and these would of course be entitled cholera morbus. In the five years, from 1844 to 1848 inclusive, of 1611 deaths from cholera infantum under fifteen years of age, 969 occurred in the first year of life, 529 in the second, 103 between two and five years, and only 10 after that age.

Sex.—There are no large tables of reference, by which to ascertain the exact proportion in which the disease occurs in the opposite sexes. It would appear, however, from our own experience, to be much more common in males than females, since of 77 cases of which we have kept a record, 48 occurred in boys, and only 29 in girls.

Constitution.—This disease is most apt to occur in feeble, delicate children, and in those of nervous, irritable temperament.

Hereditary Predisposition.—Our own observation leads us to believe that the disease is apt to occur in certain families. It would seem probable that this peculiarity must depend on the fact that the constitutions of some families are particularly disposed to disorders of the digestive apparatus. We are acquainted with one family in this city, in which eight out of ten children suffered more or less from the disease. Again, of these children four have grown up, married, and have children. Two of these families have each lost a child from the disease; in a third, the two children of the family have been exceedingly ill with it; while in the fourth some of the children have been sick, though not to the same degree. Again, we have attended two children in a family, one not quite two years, and the other three months and a half old, who have both been very sick with the disease. The elder child was ill the summer before in the same way. The mother of these children was herself very ill with the disease on several occasions during her infancy, as was also her brother.

ANATOMICAL LESIONS AND PATHOLOGY.—It will be readily understood that it is far from an easy task to define precisely what are the essential lesions in true cholera infantum, as we have described it. Having been confounded so long with ordinary inflammatory diarrhœa, the lesions usually attributed to it are precisely those we have detailed in our article

on the latter affection. In those cases again where the true choleraic disease appears during the course of inflammatory diarrhœa, it is of course difficult to determine to which affection the lesions presented after death are in reality due. We must, therefore, seek for the true and proper lesions of cholera infantum in the comparatively rare cases in which this affection has appeared in the midst of good health, and has proved fatal during the acute stage. With this restriction then it appears that the only anatomical changes which can be regarded as constant and essential to the disease, are enlargement of the mucous follicles, and, to a less degree, of the glands of Peyer; and softening, and in some cases erythematous inflammation of the mucous membrane.

There can be little doubt that the appearances thus indicated depend upon the presence of an early stage of inflammation of the tissues of the intestinal walls, and of the mucous follicles. This view is supported by the similarity between these lesions and those found in cases of enterocolitis, proving fatal during the early stage, as well as by the fact that where the child survives the choleraic stage, and ultimately dies after a continuance of diarrhœa for some days, or even several weeks, the lesions are found to have developed into those ordinarily found in primary enterocolitis.

It is, however, necessary to consider briefly what additional element is present, in this form of disease, which impresses upon it such peculiar and fatal features; or, in other words, what is the pathology of the collapse which characterizes cholera infantum.

It is a matter of much regret, that as yet we are wanting in careful microscopical examinations of the condition of the epithelium of the mucous membrane, and of the characters of the evacuations. We should anticipate, however, from the evident similarity between cholera infantum and sporadic cholera, or cholera morbus in the adult, that in the former as in the latter disease, such examination would reveal rapid proliferation and exfoliation of the cells of the mucous membrane.

In regard to the explanation of these lesions, we would refer the reader to the remarks upon the pathology of enterocolitis, where we have expressed our belief that the causes of these affections (heat, noxious emanations, unwholesome food), act in a complicated manner, by inducing a state of malnutrition in which the tissues are prone to undergo inflammatory changes, by loading the blood with noxious substances, which may irritate the glands which excrete them, and finally by interfering with digestion, so that the contents of the intestinal canal undergo changes which render them highly irritating.

We repeat that we recognize in cholera infantum the presence of the general alteration of nutrition, and the change in the entire blood mass, as well as the local irritant action of the morbid contents of the intestines. But it is in the highest degree interesting and significant of the importance of this last element in the causation, that symptoms altogether indistinguishable from those of cholera collapse, may be produced by agencies acting directly and solely upon the coats of the stomach and intestines.

Attention was directed to these analogous conditions by Sedgwick, in a

highly valuable article, "*On some Analogies of Cholera, in which suppression of urine is not accompanied by symptoms of uræmic poisoning*" (*Med.-Chir. Trans.*, vol. li, p. 1, 1868), in which he collected many such examples. Among the causes which are clearly established as capable of producing such an analogous condition, are poisonous doses of corrosive sublimate, arsenic, some of the mineral acids, especially nitric acid; and also of certain drastic purgatives, especially croton oil. In these cases the peculiar symptoms produced, which are uniformly described by accurate observers as most closely analogous to those of cholera collapse, are due exclusively to the direct irritant action of the substance upon the gastro-intestinal mucous membrane.

The same effects have frequently been observed to follow the eating or drinking of poisonous animal matters, such as tainted or simply unwholesome meat or fish, and milk which has undergone some injurious, but as yet unknown change, decomposing vegetables, and some of the poisonous fungi. In this last group of cases, the local irritant action of the substances swallowed must certainly be regarded as the principal cause in the production of the symptoms, although it is quite possible that the ingestion of such putrid animal or vegetable substances should also cause an altered condition of the blood.

In like manner, there are numerous morbid conditions of the intestines, or their peritoneal covering (as perforation with subsequent peritonitis, peritonitis, from extension of inflammation, intestinal obstruction), which may be attended with symptoms closely analogous to those of cholera collapse.

We will also quote from Rilliet and Barthez the following passage in regard to the remarkable memoirs upon *Inanition*, by Dr. Chossat, of Geneva, which show the analogy which exists between the results of experimental inanition and the chief symptoms of cholera infantum. "This is seen especially: (1.) In the diminution of temperature, which, conjoined with the loss of weight, is in inanition, as in cholera infantum, one of the principal causes of death. (2.) In the stupor which follows the jactitation as the temperature falls. (3.) In the colliquative diarrhœa during the last few days of life, the severity of which is proportioned to the rapidity of the fatal termination, and to the increase of the algidity.

It is not within the scope of the present work to discuss, critically, the various theories which have been advanced to explain the *modus operandi* of such causes in producing a state of collapse analogous to that of cholera, as well as the pathology of true cholera collapse.

It is, however, evident that the mere drain of fluid from the alimentary canal, although it undoubtedly has much influence upon the course of the disease, cannot be regarded as the efficient cause of collapse, since in many cases profound collapse occurs with comparatively scanty discharges.

So too we must regard Dr. Johnson's hypothesis (*Medico-Chir. Trans.*, vol. l, 1867, p. 103, *et seq.*), that the symptoms of collapse are due to a spasm of the minute branches of the pulmonary artery, caused by the specific alteration of the blood in cholera, as based upon insufficient arguments. Thus, in the first place, we have cited instances above where

symptoms altogether similar to those of cholera collapse, are produced under circumstances in which it is impossible even to suspect the existence of a poisoned state of the blood. Again, there is neither any clinical nor anatomical evidence to show that the contraction of the pulmonary artery is relatively greater than that of the rest of the arterial system; or again, that such contraction precedes the other signs of collapse.

In an earlier edition of this work, we quoted the opinion of Rilliet and Barthez in regard to the implication of the sympathetic nervous system in cholera infantum, and since that time we have been led to regard this more and more strongly as the essential cause of the collapse which characterizes this and other choleraic conditions.

The passage extracted from the admirable work of Rilliet and Barthez was as follows: "The disease we have just described is, in our opinion, a catarrh which has localized itself upon the digestive tract and the great sympathetic nerve. It is, of all forms of the catarrhal affection, that which most clearly justifies the idea of a poisoning. It proves also that anatomical differences alone will not suffice to establish a separation between the various species of the disease.

"Its catarrhal nature is demonstrated by the causes, which are those of all catarrhs (improper alimentation, epidemic influence, etc.); by the analogy of the symptoms; by the gradual passage of the mild into the grave forms, through intermediate cases; and lastly, by the fact that simple intestinal catarrh is often but the prodrome of choleraiform enteritis.

"Reasoning from the simple fact that the disease is catarrhal, we admit the existence of a modification of the whole economy, and of some alteration of the blood.

"A study of the anatomico-pathological descriptions of the disease, and especially the observation of cases, demonstrates that the gastro-intestinal tract of children dying of this affection may be found in four different conditions:

"*a.* Either the stomach is softened without any lesion of the digestive tube.

"*b.* Or the stomach is softened, at the same time that the mucous membrane of the intestines, and especially its follicular apparatus, is diseased.

"*c.* Or the stomach is healthy, whilst the follicular apparatus or the mucous membrane are diseased.

"*d.* Or, lastly, the gastro-intestinal tract fails to exhibit any lesions appreciable by our senses in the present state of our knowledge, or it presents alterations too insignificant to explain the gravity of the symptoms."

. . . . "Up to this point the disease resembles all other catarrhs, but what gives to it a special type is the abundance of the serous secretion and the disturbance of the great sympathetic nerve.

"The serous secretion, which seems to be produced by perspiration (analogous to that of the respiratory passages and of the skin), rather than by a follicular secretion, shows, perhaps, that the elimination of morbid matter is accomplished by other organs than the follicles; and we ought perhaps, to see in this a proof that the matters to be eliminated are not the same as in simple catarrh. On all these points we are compelled to remain in doubt, we content ourselves with stating the fact.

"The functional derangements of the trisplanchnic nerve play an important part in the disease; under this point of view it differs from the mild form, in which the innervation is normal, and from the cerebral form, in which it is especially the cerebro-spinal apparatus that is sympathetically affected. The proof of a disturbance of the ganglionic nervous system, rests upon the following physiological and nosological considerations:

"The disease exists at the age and in the physiological condition (dentition), in which functional derangements of the nervous system without lesions of organs are most frequent; it is often complicated with those very disorders of the general innervation, as is proved by certain profound changes in the functions of nutrition, circulation, and calorification, which the amount of material waste will not always account for. We occasionally observe the same symptoms of nervous sideration, and particularly the extreme smallness of the pulse, and the algid phenomena, to arise in certain of the most violent attacks of spontaneous peritonitis. Now these phenomena, which cannot always be referred to the intensity of the pain, and which do not exist in inflammations of the other serous membranes, no matter what the rapidity of their course, are only to be explained by the fact that the disease, seated in the abdomen, envelops the ganglia of the great sympathetic nerve."

Since the date at which this was written, our knowledge of the functions of the sympathetic nerve, especially with regard to its power of regulating the calibre of the arteries, by inducing contraction or allowing relaxation of their muscular coat, has been much advanced; and we are fully prepared to understand how the symptoms of cholera collapse might be explained upon the supposition of a wide-spread powerful irritation of the fibres of the sympathetic nerve, so richly distributed to the coats of the vessels throughout the alimentary canal, and which have such intimate relations with the nervous supply of the whole arterial system, as well as of the heart and lungs.

Thus we can most readily explain in this way the small, thready pulse; the cold, pale, and shrunken skin; the asphyxia and coldness of the breath; the diminution in the formation of urea and in the secretion of urine.

The above views of the pathology of choleraic collapse have been of late ably supported by Sedgwick (*loc. cit.*) and Dr. Horace Jaeffreson (*Edin. Med. Jour.*, December, 1866, p. 520).

At the same time the probability is that the vaso-motor nerves of the intestinal walls themselves are paralyzed, from exhaustion of their excitability, so that dilatation of the vessels occurs with profuse discharge of serum.

So far as experimental research can be made available in deciding questions involving such deep-seated and delicate parts, the results entirely confirm the explanation given above. Thus Moreau¹ has found that, after

¹Comp. Rend. de l'Acad. des Sciences, t. lxvi, p. 554, 1868, in Medical Times and Gaz., April 11th, 1868, p. 397.

section of the branches of the sympathetic nerve supplying the intestines, a copious secretion of alkaline serous fluid takes place into the bowel.

SYMPTOMS.—Restricting, as we now do, the term cholera infantum to cases which have a truly choleraic character, we shall have a smaller ground to go over than we had in our early editions.

The *invasion* of the choleraic symptoms is sudden. The child may have been quite well previously, or may have been the subject for an indefinite length of time—days or weeks—of simple or inflammatory diarrhœa, when, from exposure to high summer heats (85° to 95° Fabr.) in a city, or more rarely, in the country; from being allowed to take some unwholesome article of food; from the effort of cutting teeth; or perhaps from having been chilled by night air, or by a sudden change of the weather from hot to cool; the choleraic disorder breaks out, with almost simultaneous vomiting and purging. The diarrhœa is, from the beginning, violent. The stools are usually frequent, consisting almost entirely of a thin fluid, which runs through the napkins and wets the clothes of the child. Sometimes the discharges are not very frequent, but each one may be so large as to wet not only the napkins and clothes of the child, but to run through to the lap or bed on which the patient lies. The chief and important characters of the stools in true cholera infantum, as in cholera of the adult, are their fluidity and quantity. These two characters, more than the vomiting or the nature of the discharges in any other respect, are the special signs of the disease, and by the degree in which they are present do we recognize the disease, and usually determine its severity. The fluid thus rendered by stool may be of different characters. It may be an almost colorless liquid, merely wetting the napkins and clothing, as though they had been dipped into a bucket of water, or saturated with the pale urine of a healthy infant; or they may consist of the same watery fluid, holding in suspension small and soft flocculi of fecal matter of a yellowish or greenish color, or small detached portions of mucus, which are left upon the napkins as the watery fluid drains through them. When the stools are of this kind they are usually almost inodorous. In other cases they are still very watery, but the fluid is yellowish or brownish in color, contains rather a larger amount of thin feculent matter, and has a most offensive odor,—an odor which is peculiar for its extreme fetidity, a fetidity so great that we have known it to cause vomiting in those exposed to it, and so adhesive as to render it necessary to change at once all the clothing and bed-linen of the child, and even then the fœtor may cling to the body of the patient, after repeated washings. This odor we have seldom met with except in the choleraic form of summer diarrhœa. The number of the stools varies greatly. We have known as many as twelve to be passed in as many hours. In other cases they are not so frequent, but the quantity at each time may be so great as to drain the body of its fluids at a more rapid rate than many more evacuations of an ordinary size. Eight, twelve, fifteen, or more than twenty stools in twenty-four hours are not rare. In one fatal case, in a child between one and two years old, there were between twenty-five and thirty stools during the second night of the attack, in a space of twelve hours.

Simultaneously with, or soon after the diarrhœa sets in, there is vomiting. The matters vomited consist at first of the ordinary contents of the stomach, food, and the gastric liquids, Soon these matters consist of the water or medicines that may be taken, and of a serous or sero-mucous fluid mixed with small portions of bilious matter. Sometimes they are tinted green, as so often happens in the gastro-intestinal affections of children. The vomiting may or may not be very frequent. It is frequently one of the severest elements of the disease, causing everything taken to be rejected almost as soon as swallowed, or assuming the form of repeated and exhausting retching, even when the stomach is quite empty. In connection with these symptoms there is rapid loss of strength. The child is listless and still between the evacuations and vomiting, or tosses and moans with the jactitation of severe illness. The appetite is lost, but thirst is extreme, and constitutes one of the marked phenomena of the disease. Water and ice are seized upon with the greatest avidity, and taken almost incessantly, if allowed, though rejected a few moments afterwards.

The abdomen is flaccid or retracted, not tender to the touch usually, and its walls inelastic, so that they can be readily pinched up into folds. The tongue, moist at first, with a thin white fur upon it, becomes pasty or dryish after a time, and is sometimes protruded from time to time between the lips.

The pulse runs up from the first, rising soon to 130, 140, and 150, and being usually small in volume, whilst the temperature remains for a time normal, rises slightly above the natural point, or, in some few cases, becomes quite high. The urine diminishes in all these cases, and in very severe ones, ceases to flow, or flows only in the smallest quantities. As in true cholera, the degree of suppression of this function is in proportion to the severity of the choleraic discharges. The respiration, natural at first, soon becomes, if the case goes on unfavorably, irregular, unequal, and anxious. The temper is irritable at the beginning, the child being restless, peevish, disposed to fret and cry at the least contradiction or disturbance. The sleep is restless and disturbed, especially at night. The child wakes frequently, and almost always with crying. When asleep, the eyes are often but half closed, and the brow contracted and frowning. The countenance soon becomes anxious and distressed. In sudden and severe attacks, it is languid and subdued, pale and contracted.

If the disease is not soon checked, signs of collapse make their appearance, and become more and more marked. The body becomes cool and then cold, the pulse grows smaller, thready, and very rapid; the features are drawn; the nose is sharp and thin; the eyes shrink within the orbits; the cheeks become sunken; the patient passes into a still, quiet, and drowsy state; the vomiting may cease, but the diarrhœa usually persists; the child falls into a comatose or semi-comatose state, and death occurs quietly in this condition, or it may be preceded by slight convulsive movements. According to the researches of Roger (*op. cit.*, p. 399), the reduction in the temperature of the axilla never approaches, in these cases of sporadic cholera, that which is found in cases of the true epidemic form occurring in children. Some very violent cases run their course in a day,

a day and a half, or two or three days. We, ourselves, do not recollect to have seen any case terminate sooner than in three days and a half.

In favorable cases, after one, two, or three days, the diarrhœa ceases to be so violent; the stools grow less frequent, smaller in quantity, thicker in consistence, containing a better concocted fecal matter, and regaining a more natural odor. The vomiting and thirst gradually subside; food is again taken and retained; the circulation falls, and the child, though weak and thin, and the subject for some days of a simple diarrhœa, may regain its health in great measure, at the end of a week or ten days. More frequently, however, the disease assumes the form of a more obstinate simple, or inflammatory diarrhœa, which may last for several weeks, to take on again, perhaps, from a recurrence of the exciting causes, the choleraic form, or to persist in one of the former shapes until the return of cool weather.

Such is a picture of the disorder to which we think the name of cholera infantum ought to be restricted. If physicians could agree to limit the title to this true choleraic disease, our mortality returns would soon show the comparative frequency of death from this disorder, and from those more tedious and chronic diseases which have already been treated of under the designation of simple and inflammatory diarrhœa or enterocolitis.

The *duration* of cholera infantum, as we restrict the term, is seldom more than two, three, or four days. It may prove fatal in a much shorter time. Dr. Eberle (*Dis. of Children*, p. 285) says it sometimes runs on to a fatal termination in five or six hours. Dr. J. Lewis Smith (*op. cit.*, p. 392) reports a case in a child sixteen months old, which ended fatally in less than one day; a second, at seven months, after a sickness of about one day; and a third, at twenty months, in thirty-six hours. We do not recollect, in our own experience, which has been chiefly in private practice, a shorter case than one of three days and a half. In favorable cases the diarrhœa usually persists, as already stated, for several days after the disappearance of the choleraic phenomena, and very frequently runs on into a simple or inflammatory diarrhœa, which follows the law of these disorders.

DIAGNOSIS.—The diagnosis of cholera infantum requires no particular elucidation. The season at which it is most prevalent; the profuse, serous, or at least fluid evacuations; the frequent and severe vomiting; the early exhaustion of muscular strength; the rapid pulse, with absence of, or a very moderate febrile heat; the threatening or the actual supervention of collapse, marked by cool or cold surface, pinched and anxious countenance, shrivelled skin, sighing or irregular respiration, rapid and feeble or extinguished pulse, diminished or suppressed urinary secretion; with, finally, the still and limp body, and drowsy or comatose brain, all mark a disorder which is readily recognized after being once seen, or which may be distinguished by any intelligent person who has never yet met with such a case, if only the progression of the symptoms be carefully inquired into, and correlated with the present condition.

PROGNOSIS.—Cholera infantum, as we restrict the use of the term, is, of

course, always a dangerous disease. Collapse, which either threatens all who are attacked by it, or actually supervenes to a greater or less degree, is well known by all physicians to be one of the most formidable morbid conditions to which the body is liable. The degree of danger in any individual case must depend chiefly upon the ability of the physician to arrest, and of the patient to resist, this state. The probability of its supervision depends very much upon the hygienic condition in which the child is placed, upon the age of the patient, the stage of the process of dentition, the present state of health, the innate vigor of the constitutional force, and also, we may say, upon the period of the disease and the degree of wisdom with which medical means are applied. Children placed in favorable hygienic conditions in the country, or in the healthier parts of cities, in large and well-ventilated rooms, and who have been fed upon proper diet, and who have, therefore, been attacked by the disease whilst in previous fair health, are much more apt to escape collapse, or to recover from it after it has made its appearance in a more or less marked degree, than those who are placed in conditions the opposite of those we have enumerated. Early age, recent weaning, improper artificial diet, unwholesome hygienic surroundings, and feeble vital powers from any cause, either inherent or acquired, are amongst the most unfavorable conditions. Still, we should never despair until the last moment, since we have seen some most surprising recoveries from apparently desperate conditions in this disease.

The prognosis may be stated in general terms to be unfavorable in proportion to the frequency and violence of the vomiting, the number of the stools, the severity of the fever, and the more or less marked character of the collapse. When the discharges consist merely of serous fluid, and are copious and frequent; when they consist of small quantities of deep-green matter, mixed with much mucus or with blood; when accompanied by straining; when they number from fifteen to twenty-five in the day; when they are very fetid; and when, with these symptoms, the abdomen is tense and tympanitic, the countenance pinched, the expression languid, the extremities cool, the pulse rapid and small, and the child irritable and restless, or, on the other hand, very still and subdued, the prognosis is exceedingly bad. If, after the symptoms just enumerated, drowsiness or stupor and then coma, convulsions, rigidity, or paralysis make their appearance, there is scarcely a hope left.

The favorable symptoms in any case are, diminution of the fever; equal temperature of the whole surface; cessation of vomiting; decrease in the number of the stools, and a return to their natural color, consistence and odor; quiet, tranquil sleep; return of appetite; and lastly, a restoration of the natural temper and gayety of the child.

PROPHYLACTIC TREATMENT.—The danger to which teething children are exposed from residence in our American cities during the hot months of the year, are now so well understood that most families who can afford it remove to the country during the warm season, and by this course very generally avoid the disease. It is undoubtedly the best plan that can be adopted, and very commonly succeeds. When this cannot be done, however, the prophylactic treatment consists in a most careful attention to

diet, dress, thorough ventilation of the dwelling, and exposure to the open air. If possible, the child should be kept at the breast until it has passed through its second summer, as there is but little danger from the disease after that period. If the weaning must take place prior to that age, it ought to be accomplished before the hot weather begins, as a change from the breast to artificial food during the warm season is very apt to bring on the disease. If the child is weaned, the diet must be strictly attended to. Up to the age of ten months or a year, the food should consist almost wholly of milk containing arrowroot, rice, oatmeal, or some farinaceous substance in small quantity. A little plain chicken or mutton water, with rice boiled in it, or a piece of beef or chicken to suck, may be given occasionally, but all vegetables and fruit should be strictly forbidden. After the age of ten months, some light soup and small portions of mutton, chicken, or very tender beef, minced very fine, may be given every day in addition to the milk food, which must still form the major part of the child's nutriment. Fruit of all kinds, all vegetables except rice and potatoes, and the latter are doubtful, ought to be carefully avoided until after the hot season has passed entirely away, or until the child has its full set of teeth. We have found the food prepared with gelatine, in the manner described, to answer better than anything else for a large number of children to whom we have prescribed it. For details in regard to this essential matter we refer the reader to the article on food.

The dress ought to be arranged according to the heat of the day. We have not rarely known young children to be kept clothed all summer in this city in thick flannel jackets, and petticoats, and woolen socks. This is certainly too much for the hot days which so frequently occur in July, August, and early in September, and is often, we believe, very injurious. A light gauze flannel shirt is the only woolen garment that need be worn during the warm season. On hot days, a child should have only this, a muslin petticoat and frock, and the lightest possible socks, or none at all. If, as constantly happens in our climate, a cool day comes, there should be added to these a light flannel petticoat.

It is of the utmost importance that children should pass as large a portion of the day as possible in the open air. In the country this is easily managed, and parents almost always contrive to accomplish it; but in a city, many people seem to think it of less importance, or their servants are occupied with other things, and it is neglected. It is, nevertheless, a matter of the greatest consequence; the child ought to be kept in the air by the nurse for several hours in the morning and evening, either in the garden attached to the house, if there be one, at the front door, walking in shady streets or public squares, or, better still, making short excursions into the neighboring country, taking care, however, to avoid the intense heat of the sun during the middle of the day.

We believe that with constant and wise attention to these points, viz., diet, dress, careful ventilation of the house and bedroom, exposure to the air, and exercise, much may be done towards preventing the disease even in families obliged to remain in the city during the summer.

As stated in the account of the symptoms, the choleraic disease often

supervenes in children who have already been the subjects of simple or inflammatory diarrhœa. When, therefore, a child in the city has diarrhœa, if it do not yield readily to treatment, and especially if the stools begin to be thin and watery, with any marked tendency to exhaustion, it ought to be regarded as being threatened with cholera. In such an event, the best prophylaxis in the world is instant removal to some high country locality or the seaside.

TREATMENT OF THE ATTACK.—Regarding this disease as a truly choleraic one, we shall follow, in the consideration of its treatment, the plan adopted by some of the more recent writers on Asiatic cholera; and shall accordingly divide our discussion of this subject into the treatment appropriate for the three stages of *evacuation*, *collapse*, and *reaction*.

Every young child who is attacked with diarrhœa, whether simple or inflammatory, in the summer season, ought to be regarded as liable to cholera, and should be carefully watched to prevent the development of this disease. For the proper treatment of such conditions, the reader is referred to the article on those affections.

Should a child, either previously well, or the subject of diarrhœa of the ordinary form, be attacked with sudden, profuse, frequent, and watery discharges, and especially, should these be associated with vomiting, with spasmodic intestinal pain, and with any appearance of general exhaustion, it ought to be presumed to be in the early or *evacuation stage* of cholera infantum, or in what is the analogue of the evacuation stage of epidemic cholera. Under these circumstances, it has been a prevalent practice here to give a cathartic, castor oil, calomel, or rhubarb. We think the practice wrong, unless there be positive evidence that the attack has followed directly upon the use of some unwholesome article of diet. If it be found that the child has certainly eaten some such food, green apples, currants, gooseberries, or articles of this kind, and that these have not come away in the discharges, it is right to give first a moderate purgative. We prefer castor oil or syrup of rhubarb, half a teaspoonful of the former, or a teaspoonful of the latter, with two drops of laudanum at the age of one year, or a teaspoonful of castor oil, or two of the syrup of rhubarb, with four drops of laudanum, or two drops of chlorodyne,¹ at two or three years of age. Two hours after this dose, if the stools continue frequent and watery, we use the chalk mixture, with tincture of krameria and laudanum or paregoric (at easpoonful of the chalk mixture, with ten to fifteen drops of the krameria, and one drop of laudanum, or five of paregoric) every two hours at the age of one year. Thirty drops of the syrup of nutgalls (see article on entero-colitis), with an opiate every two hours, is often very useful. We believe that the great object is to arrest the watery discharges by stool. If the above means fail, laudanum by injection, two drops at one year, and double the dose at two years, every two or three

¹ The preparation which we prescribe under the name of chlorodyne is not Dr. J. Collis Browne's, but is made by Messrs. Bullock & Crenshaw, of this city. It contains one grain of morphia to the fluid drachm; but, as it does not drop less than 120 to the fluid drachm, the dose for an adult is 10 to 15 drops. It is a very elegant preparation, and has proved most efficient in our hands.

hours, may be tried in addition to the above treatment. The quantity of opium to be used must depend on its action. Children, like adults, bear very different amounts. As soon as positive drowsiness appears, or the pupils become contracted much below their natural size, the doses must be suspended or diminished, or the intervals between them lengthened. Of course, if the stools lessen in frequency, quantity, or fluidity, the same reduction in the amount of the opium ought to be made.

When vomiting is severe and frequent, and the above remedies are rejected, we may use the one proposed in the article on inflammatory diarrhoea, consisting of solution of morphia, dilute sulphuric acid, and curaçoa-cordial. This, or some similar remedy, is at times very successful. It is nineteen years since one of us saw a child nine months old, in deep collapse from a most violent attack of cholera infantum, who rejected its mother's milk as though from the action of an emetic, whose stomach was only made worse by calomel, but who began to improve very soon upon doses consisting of two drops of aromatic sulphuric acid, and five drops of solution of morphia, in a teaspoonful of iced water, every hour. Since then we have frequently used the mixture above recommended in such cases, and we think, on the whole, with more control over the vomiting than anything we have tried. In other cases, minute doses of calomel and bismuth, or nitrate of silver, as already recommended (page 438), will allay gastric irritability and afford relief.

The experience gained by careful and lengthened observation in the treatment of the evacuation stage of Asiatic cholera, may well be applied to the affection under consideration, so much alike are they. Dr. Good-eve (*loc. cit.*, p. 177) gives first a full dose of opium (he says that calomel was generally combined with it in India, and though he does not "know that the calomel does good, it does no harm"), to an adult two grains, and half an hour afterwards he begins with an astringent, in his own practice, usually the following mixture :

R. Plumbi Acetat., gr. xxx.
 Acid. Acet., ℥x.
 Aq. Destillat., fʒvj.—M.
 One ounce or half an ounce every half hour or hour.

At the end of an hour from the administration of the first dose of opium, if the purging persisted, he gave one grain of opium and continued the astringent. A small teaspoonful, or two-thirds of an ordinary teaspoonful of this solution would contain about half a grain of the acetate of lead, and this might safely be given to a child a year old for several doses. We have not used this remedy ourselves, but it comes from a source which commends itself to us, and we shall not hesitate to use it when the occasion presents itself. As soon as the frequency of the discharges is arrested, the doses should be given at longer intervals, and when the peculiar serous character of the stools has disappeared, this remedy ought to be suspended, and some more simple one substituted, in order to avoid the possibility of producing the toxic action of lead.

If, in spite of the treatment, the stage of *collapse* should set in, other

methods of treatment must be adopted. Here the stools are usually in great measure arrested, or they are few in number and small in amount. The object to be sought after is to produce reaction, or rather to favor the efforts of nature to bring about this change. It is now generally acknowledged by men of experience, that the old plan of pouring in large doses of opium and alcohol is a great mistake. But little is absorbed by the stomach whilst the body is in this condition, and not unfrequently the patient is injured, perhaps fatally, by the sudden absorption of these substances, when the stomach begins to absorb after reaction has taken place. The opium may cause dangerous or fatal stupor, or may increase or keep up the tendency to suspension of the urinary function, and thus promote one of the great dangers of the disease, uræmic intoxication. The alcohol, if it has been used in large quantities, would also tend to clog the nervous centres, to cause gastric or gastro-intestinal catarrh, and to heighten beyond a safe point the febrile movement which is so apt to accompany the reaction stage. Opium, therefore, should be avoided during collapse, or given only in the smallest doses. Alcohol, though it should never be given in large doses, and recklessly, as has so often been done, may be used in small quantities, especially if it be found by close watching that it promotes the force and volume of the pulse. Ten or fifteen drop doses of old and delicate brandy, in a teaspoonful or tablespoonful of ice-water, ought to be given every hour or two hours, at one year of age. During collapse the stomach is still often very irritable, and yet the thirst continues intense. We are glad to find that such men as Drs. Maclean and Goodeve recommend the free use of ice and water under these circumstances. Our own practice, for years past, has been to allow ice and cold water, almost without limit, to children in this condition, and we are much pleased to know that such, too, is the practice of these gentlemen. We never could understand the wisdom of refusing water to patients who were suffering the horrid thirst produced by the immense losses of the water of the body by serous purging. The degree of thirst for water (a natural and not a secondary diseased instinct, like that of the drunkard for alcohol) must be the safest guide we can have as to the need of the body for water, and as such, it ought always, it seems to us, to be gratified, unless under very rare and most peculiar conditions. We give water and ice, even though the child vomits from time to time, believing and hoping that some will be absorbed to take the place in the tissues of that which has been drained off through the intestines. This point in the treatment we regard as so important, and one, we think, so much misunderstood by the public and by some medical men, that we make the following quotation from a note of Professor Maclean's to Dr. Aitken (*Aitken's Practice*, vol. i, foot-note, p. 663); "Urgent thirst is one of the most distressing symptoms in cholera; there is incessant craving for cold water, doubtless instinctive, to correct the inspissated condition of the blood, due to the so rapid escape of the *liquor sanguinis*. It was formerly the practice to withhold water—a practice as cruel as it is mischievous. Water in abundance, pure and cold, should be given to the patient, and he should be encouraged to drink it, even should a large portion of it be rejected by the stomach; and when

the purging has ceased, some may with advantage be thrown into the bowel from time to time." The use of water by enema, when the diarrhœa is checked, is a point which ought not to be neglected, especially if the stomach continues weak and irritable. A gill of tepid water may be used at a time, thrown slowly and gently into the bowel, in the case of a child one or two years old. If this is retained well, the same quantity may be repeated in one or two hours.

Whilst the collapse lasts, but little food can be taken. It is seldom retained if used in any quantity, and the stomach has lost, in great measure, its digestive power. The only food we have found at all available has been thin chicken tea, Liebig's cold extract of beef, or weak wine-whey, given in two or three teaspoonful doses, every half hour or hour. It is worse than useless to attempt more than this, as not only is it not retained, but it evidently tends to keep up the nausea and vomiting, and thus retard the natural effort at reaction. As to remedies in this condition, we doubt whether anything better can be done than to use water, as just advised, small doses of brandy, and, if they can be borne, the acid and morphia mixture recommended above, small quantities of the liquor ammoniæ acetatis, ten to twenty drops, in cold water, every hour, at one year of age. There is, however, a remedy which has obtained a great reputation amongst the English army surgeons in India, for the promotion of reaction in the collapse stage of epidemic cholera, which we have used ourselves with advantage in adults, but not in children, though we propose trying it when we next have a good opportunity. It is spoken of highly by Dr. Maclean. The formula is as follows:

R. Ol. Anisi, Ol. Cajuput., Ol. Junip., āā,	f3ss.
Ætheris,	f3ss.
Liq. Acid. Halleri,	f3ss.
Tinct. Cinnamomi,	f3 ij.—M.

The dose for an adult is ten drops every quarter of an hour, in a tablespoonful of water.

An opiate may be given with the first and second doses, but should not be continued, for the reasons already given. The liq. acid. Halleri consists of one part of concentrated sulphuric acid to three parts of rectified spirit. The dose of this mixture for a child a year old, ought, we think, to be about one or two drops in a teaspoonful of water, given, as above stated, every quarter of an hour. So much is this valued in India, according to Dr. Maclean, that it is always ordered to be kept in store in the "medical field companion" of armies on the march.

It must not be supposed that all children seized with choleraic diarrhœa are necessarily to pass through the collapse stage in all its terrors. On the contrary, many, when judiciously treated early in the disorder, escape collapse altogether, and yet they have had none the less the true choleraic disease. Others suffer more profuse and exhausting losses of water by the discharges, or their vital power of resisting disease is less, and they pass into more or less deep collapse; or hang, as we have seen them, on the very edge of that condition, for one or two days, and then emerge from

the danger, without having done more than cause the experienced physician the grave anxiety which such suspense must and ought to create. During these doubtful moments of the attack, the child should be kept as quiet and still as possible. He should be made to lie in a constantly horizontal position, on a smooth and easy mattress, in the crib, or on a large and roomy bed, and as little as may be on the lap, which is uneven and unsteady, and which must give his weak and exhausted muscles more work to do than they would have on the more solid and even bed. If, however, the nature of the child be such that he clings to the mother's or nurse's lap as his only safety, or if he have been taught (a most ill-judged lesson) to prefer the lap to any other position, we must yield to him, rather than cause fretting or unhappiness, when his very life may hang upon the avoidance of all disturbing influences. In this case it is well to place him upon as firm a pillow as can be found, and let him be held on this in the lap. It is important to move him, when this becomes necessary, as slowly and gently as possible, always keeping the body on a horizontal plane, to avoid the tendency to the syncopal state, which sudden movements, and especially the sitting or erect position, are apt to produce. When the tendency to cooling of the body shows itself, and this is usually first noticeable in the hands and feet, ears and nose, he should be kept wrapped in warm, dry, and soft flannels or blankets. Flannels heated at the fire, thus supplying dry artificial heat, are of great use here. Bottles or tins filled with hot water, ought to be placed at the feet, under the blanket. A warm, soft, and light poultice of Indian meal or flaxseed, with a little mustard incorporated with it, may be placed over the abdomen, or three or four thicknesses of flannel, wrung out of hot water and whiskey, may be laid over the lowest part of the thorax and over the abdomen, and covered with oiled silk, to retain their heat and prevent the wetting of the clothes. Whilst artificial heat is thus made use of, fresh air must not be excluded. On the contrary, as these cases almost always occur in the hottest summer weather, the largest supply of fresh air that can be obtained must be admitted. Warm baths, which were proper and useful during the early stage, especially when fever was present, we have not found useful in these cases. The fatigue and irritation caused by the disturbance of undressing and dressing the child, have seemed to us to do more harm than any good derived from the heat of the water compensated for.

When the case takes a favorable turn, and the *reaction stage* begins, it is usually best to do nothing more than supply food and water carefully, and keep the body quiet and tranquil. The food may be cautiously and slowly increased in quantity, if the stomach has become settled. Tablespoonfuls of thin chicken-tea, just flavored with salt, or of Liebig's cold extract of beef, or of light beef-tea, or of a mixture of wine-whey with two or three parts of thin arrowroot decoction (a teaspoonful to a pint), may be given every half hour or hour. If these are retained several times, and the child shows some little anxiety for food, the same materials may be given in wineglassful quantities. At the same time, water and ice ought to be allowed from time to time, as the thirst may call for them. On the second or third day of the reaction, we may give, if the child shows a desire for

it, a little milk and water and lime-water, one part of milk to one or two of water, with one of lime-water, commencing with not more than two or three ounces of the mixture at each feeding. The milk ought certainly to be very much diluted for the first three or four days after it is allowed. We have used with success the food made of equal parts of milk, cream, lime-water, and plain water, as described in the chapter on food. When the child has been carried thus far safely, we may gradually return to its former habits of feeding, allowing meat to suck, a little bread, and so on, if it is old enough for such habits.

As to drugs during the reaction stage, they are not necessary if everything goes on well. If, however, the fever run high, we may use small doses of the spirit of nitrous ether, as ten drops, in iced water, every two hours at one year, or twenty drops of the solution of acetate of ammonia, in the same manner, at the same age. If, as often happens, the urinary secretion remains scanty, water, in such quantities as the stomach takes willingly, makes the best diuretic; or we may use the spirit of nitrous ether, as just recommended, with a grain of acetate of potash and half a drop to a drop of tincture of digitalis, every two hours, for a day or two.

When reaction is thus successfully brought about, the child may either improve rapidly and regain its previous health, or simple or inflammatory diarrhœa may set in, and pursue the usual course of those disorders. In the latter event, the child, if the attack of cholera have occurred in the city, ought certainly to be removed to the country if possible, since it is only too apt to have a recurrence of the choleraic disease if kept in town, or to suffer, at least, a tedious and more or less dangerous attack of the simpler form of diarrhœa. For the proper treatment of either of these sequences to cholera, the reader is referred to the articles on those diseases, with the warning, however, that all such patients ought to be treated with every minute care as to hygienic and therapeutic measures that experience and art have taught us, since the health has been so rudely shaken by the sickness already endured.

We have now laid before the reader, to the best of our ability, what we think is the best method of treating cholera in children; but, before quitting the subject entirely, we wish to make a few remarks upon points not referred to in the above account.

Attention to the *state of the gums* should never be neglected in teething children. Our experience leads us to believe most implicitly that the process of dentition, or at least that and other concomitant constitutional conditions, are constant predisposing causes of gastro-intestinal disorders in early life, and that the active hyperæmic state, or positive acute inflammatory condition, which often attends upon the near approach of teeth to the surface of the gum, may become an exciting cause of acute digestive diseases, such as cholera. We think it is always well, therefore, to examine into the state of the mouth in a choleraic child as in other infantile disorders; and if the teeth are felt distinctly through the gums, and the gums be found swollen, tense, hot, and highly vascular, to cut them freely once. If, on the contrary, the gums are firm, not hot, not redder than

usual, and the edges of the teeth cannot be felt, it is foolish to cut them.

Baths.—In the early stage of cholera, before collapse has begun, and whilst the child is still reasonably strong, and particularly when there is marked febrile heat and dryness of the body, we think that the use of the warm or hot bath, or of sponging with hot water and spirit, are excellent measures. The bath may be used twice, or even three times a day if the child does not resist and scream. The temperature should be 95° to 98° , and the child may be kept in the water from five to ten minutes. It is an excellent plan to wrap the child, directly on lifting it from the bath, in a heated muslin sheet, and to apply over this a blanket, and keep it thus enveloped on the lap, for half an hour or more if it is comfortable and disposed to rest. If the child be somewhat weak, whiskey, added to the water, renders the bath more useful and safe. When the use of a bath alarms or annoys, so as to cause violent agitation, it is best to substitute sponging with hot water and whiskey or vinegar, under a light blanket, two or three times a day.

Antiphlogistics.—It may appear to many, in these modern times, a mere waste of words for us to state that we are opposed to bloodletting, in any form or any stage of cholera infantum. But if any such will take the trouble to look over the works of writers of ten and twenty years back, he will find reason to think that if this be our opinion, it ought to be expressed. When one of ourselves began to practice, in 1841, it was quite the custom to take blood for the nervous symptoms which are present in the early stage, and still more for the comatose phenomena at the close. This was done on the theory that these symptoms were the result of congestion or inflammation of the brain, whereas now they are looked upon as the results of exhaustion, of the altered conditions of the blood, or of uræmia.

Calomel.—The opinion was expressed in a former edition of this work, that the doses of calomel usually recommended were too large for young children, and were apt to aggravate the existing irritation of the digestive mucous membrane; and that such doses of a remedy acknowledged to be a powerful sedative, could not be proper in a disease which constantly tended towards exhaustion and collapse. It was also stated that the small doses which we did recommend had been declared by some critics to be entirely too small, and that to this we could only reply that the larger and more careful, and, we hoped, the wiser our observation had been in the last few years, the more thoroughly convinced were we that the larger doses, such as were formerly recommended and used by nearly all writers and practitioners, were not only unnecessarily large, but most seriously objectionable. We went on to say that the indiscriminate use of this remedy, in nearly all cases of the gastro-intestinal diseases of childhood, became with some, we believed, a mere routine habit,—that they never tried what might be accomplished without it, but went on pushing the drug in constant doses, when the case, if trusted to simpler means, or even left to the efforts of nature, would often do much better, we had learned to believe, than when these delicate organs were made the receptacle of

doses that could not but tend to keep up the nausea, vomiting, and diarrhœa, which forms so important a part of the morbid phenomena. The experience we have had since that time has but confirmed us in these opinions. Indeed we have so often been disappointed in obtaining any good effects from the drug, and have so often had reason to think that, instead of allaying nausea and vomiting, it increased them, and added to the exhaustion which is one of the dangers always to be contended against, that we have virtually abandoned its use in this affection.

ARTICLE IV.

DYSENTERY.

It seems to us unnecessary to make more than a few remarks on dysentery, since we have already spoken of the morbid conditions of the large intestine, in our article on entero-colitis. Dysentery, however, differs from this latter affection by the fact that it frequently occurs in an epidemic form, and that there is a tendency to more rapid and extensive ulceration of the mucous membrane of the rectum and colon. It is an acute febrile disease, characterized by frequent evacuations, attended with more or less severe pain and straining, and consisting of muco-sanguinolent or sanguineous substances, which are due to ulcerative inflammation of the rectum and colon.

The *causes* of dysentery are but little understood, beyond the mere fact that it occurs as an endemic in some regions of country, and as an epidemic over large districts. It is frequent, also, as a sporadic disease, and in this form seems to depend upon the same causes as those already cited as productive of entero-colitis. Like cholera infantum, it appears to be more common in boys than girls, since of 41 cases of which we have kept notes, in which the sex is mentioned, 28 occurred in boys, and only 13 in girls. It is most frequent in the second and third years of life. Of 40 cases in which the age was noted, 1 occurred in the first year of life, 16 in the second, 8 in the third, 3 in the fourth, 3 in the fifth, 1 in the sixth, 3 in the seventh, 3 in the eighth, and only 2 from the eighth to the end of the eleventh year. It may be either idiopathic or secondary. As a secondary affection it is most apt to follow measles and variola. We have often known dysenteric stools to occur in the course of cholera infantum, and in a considerable number of cases such as we have described under the title of entero-colitis.

The *anatomical lesions* are confined chiefly to the large intestine, and are the same as those described under the head of entero-colitis, except that they are of a graver character. The mucous membrane is commonly found thickened, swelled, red, and softened; the submucous tissue sometimes presents ecchymosed points; the follicles are often diseased, their orifices being enlarged and ulcerated, as described under entero-colitis.

In grave cases, particularly those occurring under an epidemic influence, there are usually more or less extensive ulcerations, which may implicate only the mucous, or extend to the muscular or even the peritoneal coat. In such instances, pseudo-membranous exudations are often formed, sometimes in large quantity, and often covering the ulcerations. The intestine contains sanguinolent mucus, or at times a brownish or greenish material which is evidently the result of a gangrenous condition of the mucous membrane, pus, and lastly false membranes. In some rare cases perforation has been known to take place.

SYMPTOMS.—The symptoms are much the same as those already described as existing in entero-colitis, excepting that the local symptoms are more severe, and the presence of blood in the stools constant. The disease often begins as a *diarrhœa*. The stools at first contain feculent materials, but after a time become very thin, small in quantity, and consist chiefly of mucus mixed with blood. The blood may be black and in considerable quantity, or of a dark rosy red color, or like the washings of flesh; it is mixed with greenish or yellowish substances, whitish mucus, fragments of false membrane, or purulent fluid. In young children there is evidently *pain*, from the restlessness, moving of the limbs, and crying about the time of the evacuations, while in those who are older, there is true *tenesmus*, like that observed in adults, and severe pain at the anus. The number of stools varies according to the severity of the case. There may be only four, eight, or ten in the day, or many more. We have quite frequently known as many as 30 and 40 to be voided in the twenty-four hours, and in fatal cases the dejections sometimes number three or four in an hour, while between the discharges the child often suffers from most violent and painful tenesmus.

The *abdomen* is generally distended, tympanitic, warmer than natural, and painful.

In mild cases there is usually no *fever*, or very little, while in severe attacks there is high fever during the first few days, marked by frequent pulse, hot dry skin, followed after a time, unless a favorable change takes place, by coolness of the surface, contraction of the countenance, hollow, sunken expression of the eye, rapid emaciation, and death.

It is useless to give a longer detail of the symptoms, as they are the same as those already described in the article on entero-colitis.

The *diagnosis* presents no difficulties. The frequency of the discharges, the pain in the course of the colon and in the anus, the tenesmus, the character of the evacuations, and the febrile reaction, all make the disease easy of recognition.

The *prognosis* is favorable in mild cases, unattended with much fever, or very frequent discharges. When, on the contrary, there is violent fever in the beginning, followed by disposition to coolness and collapse: when the stools are exceedingly frequent, and attended with severe pain and almost constant straining; and when they consist of nothing but mucus, mixed with considerable quantities of blood, or with pulse or false membranes, the prognosis is very unfavorable. Of 38 cases, the termination of which we have recorded, 4 proved fatal.

TREATMENT.—The treatment of dysentery in children is often very unsatisfactory. The mere variety of the remedies recommended by different writers and practitioners marks the uncertainty of the effects obtained from drugs. Mild cases so generally get well under any treatment that all methods have had their supporters and advocates, while grave cases, and especially those occurring under the influence of severe epidemic visitations, are so difficult of treatment, and often so little under the evident control of medical means, as to leave the careful observer in great doubt as to what he ought to set down as the evident result of his own action in the case, and what as the results of the effort of nature to cure the disease.

Mild cases, in which the fever is not very high, the number of stools not great, and the pain and distress moderate, require little else than *rest* in bed, a light and unirritating *diet*, and the use of *opium* in small quantities either internally or by injection. When there is reason to suspect the presence of unwholesome food in the stomach, or of unhealthy secretions in the intestines, it is necessary to give in the beginning small doses of some mild *cathartic*. The one generally preferred is castor oil, which may be given either simple, in the dose of a small teaspoonful containing one to four drops of laudanum according to the age, or in the form of emulsion. The latter is the mode of employing it usually chosen. A drachm of oil should be rubbed up with a scruple of gum, a little sugar, from two to eight drops of laudanum, according to the age of the child, and seven drachms of some aromatic water. The dose is a teaspoonful every three or four hours. If the case continue to improve under the emulsion it may be continued for a couple of days, but should the stools become more and more frequent, and the pain and tenesmus increase, it must be suspended after one or two days, and laudanum enemata, with or without the internal use of absorbents and astringents, substituted. The injections ought to consist of four or five drops of laudanum at two years of age, and of ten drops at five or six years, suspended in from half an ounce to an ounce of some mucilage, or thin farinaceous fluid, or simply mixed in a tablespoonful of tepid water, which is perhaps the best plan of all. The injections may be given every four or six hours if necessary, or they may be made use of only at night, while small doses of Dover's powder are administered every three or four hours through the day.

If the signs of rectal inflammation continue marked, it will be well to add to the injections nitrate of silver as recommended on the next page, or in smaller doses as recommended in chronic entero-colitis (p. 438).

The internal remedies that we depend upon chiefly are subnitrate of bismuth with small doses of Dover's powder or of opium alone; prepared chalk given in emulsion with an astringent, as *krameriaë*, and with a suitable amount of opium according to the amount given by enema; or acetate of lead.

The *diet* in these cases should consist of arrowroot, sago, tapioca, or some such food, made into thin pap with milk and water; and the quantity allowed ought to be very moderate. Rest in bed, in the cradle, or in

the lap, is essential. The child must not be allowed to run about, to be on the floor, or to use exertion of any kind.

In very severe cases of dysentery the treatment is, as above stated, difficult and uncertain, owing to the dangerous character of the disease, and to the fact that so many different methods have been recommended by different writers.

In the early stage of a *severe case*, whilst the febrile reaction is high and the strength of the patient still unsubdued, depletion by leeches is strongly approved of by many able practitioners. For our own part we have not resorted to it as a general rule, from the fact that we have so often found, the strength of the child to fail rapidly under the disease itself. In a few of our cases, however, where the pain was very severe and the fever high, and where there was marked soreness of the abdomen, the application of a few leeches around the anus has been followed by manifest benefit. All occasional *warm bath* is also very soothing and useful in such cases.

The internal remedies most commonly depended upon are castor oil in emulsion with laudanum, mercury, sugar of lead, opium, nitrate of silver, spirit of turpentine, and astringents. The castor oil emulsion, prepared as mentioned above, is useful in the early part of the attack, but ceases to be so, according to our experience, after the first twenty-four or forty-eight hours. Whichever astringent or alterative remedy is now selected, all agree as to the propriety of continuing the use of opium, and the very fact that it is so universally employed points it out as one of the most reliable and valuable means we have at our command. It is certainly the one upon which we most depend ourselves. It may be given either alone or in connection with other substances. Where injections can be retained it is best given in that way. About five drops of laudanum at two years of age, or ten drops at four or five years, may be given in a tablespoonful of any bland vehicle every four hours. When the rectum rejects the enema as soon as administered, the opium should be given either by the mouth, or in the form of laudanum or solution of morphia, or in that of Dover's powder; or in the form of suppository. We should indeed strongly recommend the administration of opium in this latter form in such cases, since we unquestionably obtain a certain beneficial local action, in addition to its constitutional effect through its absorption. The amount of opium should be about the one-eighth of a grain at two years of age, which, together with any other remedy, such as acetate of lead, if it be desired, should be incorporated with butter of cocoa, a most bland and soothing substance, which dissolves readily at the temperature of the body. When made of this substance, and of proper shape and sufficiently small, the suppository can be introduced without pain, and will usually be retained. It should of course be repeated at intervals, depending upon the effect produced. Opium is almost always employed in connection with some other remedy, and particularly with calomel, acetate of lead, or nitrate of silver.

There is much difference of opinion as to the value of mercurials in severe cases of dysentery. Calomel is the form that is most commonly prescribed, and many excellent authorities strongly recommend its use in combination with small doses of opium and ipecacuanha. We use it not rarely

ourselves, but chiefly in those cases where the heavily coated tongue, the irritable stomach, and the tumid abdomen indicate that the mucous membrane of the upper part of the alimentary canal is also involved in the affection. The best mode of giving it, according to our own experience, is in small doses frequently repeated; as, for instance, from gr. $\frac{1}{20}$ to gr. $\frac{1}{12}$ every two or three hours at the age of two or three years. This may be combined with gr. $\frac{1}{20}$ to gr. $\frac{1}{12}$ powdered opium, or with half a grain or a grain of Dover's powder; or two grains of subnitrate of bismuth may be given with each dose of the calomel, while the opium is given by the rectum in the form of enema or of suppository.

Acetate of lead is also much relied upon, and we have ourselves obtained excellent effects from its use in some instances. It is difficult to define precisely in what cases it is preferable to calomel or nitrate of silver. It has seemed to us to produce the best results in cases where the abdominal pain was severe and not limited to the region of the lower bowel, and where the discharge was frequent and not composed merely of mucus, with more or less admixture of blood, from the rectum. The dose is from one-third of a grain to a grain every two or three hours at two or three years of age.

The two remedies which have been of more positive efficacy in our own practice than any others, with the exception of opium, are the nitrate of silver and the solution of the nitrate of iron. The former we have used both internally and by injection, the latter only by injection. For an account of the mode in which these remedies are employed by different authorities, the reader is referred to the remarks on chronic entero-colitis. We have employed nitrate of silver in sixteen cases of dysentery. These were all severe attacks, and some of them most violent. Of the sixteen cases, three died. The remedy was given by the mouth alone in seven cases, by injection alone in five, and by the mouth and by injection both in four. It has proved most beneficial in its effects, in our hands, when given by the mouth, though its influence over the disease has always been less immediate than when used by injection, but it has been more permanent. The dose in which we have used it has varied with the age of the child, and with the severity of the symptoms. For children two years old we have usually employed from one grain to one and a half grains, and for those of five or six years or upwards, two grains dissolved in two ounces of vehicle, consisting of an ounce each of syrup of gum arabic and distilled water. The dose is a teaspoonful every two or three hours. It is well, as a general rule, to add from four to sixteen drops of laudanum, according to the age of the subject, to the mixture. For use by injection we have commonly employed for each enema two grains for young children, and four grains for older ones, dissolved in four ounces of distilled water. The injections are to be repeated twice or three times a day. After the nitrate of silver enema has come away, it is a good plan to throw into the bowel a laudanum and starch injection.

We have made use of the solution of nitrate of iron, to which allusion was made above, only as an injection in acute dysentery. We have employed it in eight cases, and are quite sure that it was of essential service in six, while in two it appeared to irritate, probably because the quantity

given was too large. Our mode of exhibiting it is to mix from ten to twelve drops in four ounces of tepid water for each injection. The injections were given twice or three times a day, and they were followed, as soon as they had returned, by a laudanum injection. On two occasions, the nitrate of iron injection remained in the bowel for several hours before being rejected, and thus restrained for that time the stools, which had previously been very frequent, and attended with much tenesmus.

When the stools continue very frequent in spite of the use of opium in some of its many forms, when sugar of lead and nitrate of silver have been employed without controlling the frequency of the discharges, we have sometimes found the mixture of aromatic sulphuric acid, laudanum, and syrup of rhatany, before recommended, very beneficial. When the stools, in addition to their dysenteric characters, have been watery, and greenish in color, the chalk mixture, with laudanum and tincture of rhatany, kino, or catechu, repeated every two hours, with occasional laudanum enemata, has been very useful.

The *hygienic management* of dysentery should be precisely the same as that which was suggested as proper for entero-colitis.

ARTICLE V.

DISEASES OF THE CÆCUM AND APPENDIX CÆCI—TYPHLITIS AND PERITYPHLITIS.

SYNONYMS ; DEFINITION.—The diseases of the cæcum and of its vermiform appendix are so important and frequent, and present so many peculiarities, as to demand a separate and detailed consideration. In approaching their discussion, it is necessary to bear in mind several important points in which the cæcum differs from the rest of the large intestine. Thus its peritoneal investment is deficient over the posterior part, which is generally quite firmly attached to the right iliac fossa by connective tissue, containing a small proportion of fat. Its anatomical relations moreover indicate that the semi-feculent materials passing from the ileum are destined to be retained in the cæcum to undergo some important action. The ileum at its lower portion rarely has a calibre greater than one-third that of the cæcum, a circumstance which must materially retard the progress of the contents of the latter, and a further detention is caused by the ileo-cæcal valve, which prevents all reflux, and by the position of the cæcum, which compels it to force onwards its contents in opposition to gravity. The view that the cæcum is the seat of an important part of the digestive process, either in the appropriation of any remaining nutritious elements of the semi-feculent chyme, the absorption of its watery parts, or the elimination of some excrementitious matter from the system, receives confirmation from the very rich vascular and glandular supply of the walls of this part of the intestine.

In addition to this, the cœcum has opening into it, usually at its lower and back part, the appendix vermiformis, a narrow, elongated, glandular process, varying from three to six inches in length, and having an average diameter about equal to that of a goose-quill, although its calibre is quite small. It is usually directed upwards and inwards behind the cœcum, and lies coiled upon itself. Its function appears to be the secretion of a viscid ropy mucus.

We thus see in the anatomical and physiological relations of the cœcum strong predisposing causes of many morbid conditions. Among these the most frequent are distension and impaction of its calibre by hardened fæces; the lodgment of a foreign body or intestinal concretion in one of its pouches or in the appendix, an accident which often excites violent and destructive inflammatory action; and finally, localized inflammation of one or all of the coats of the cœcum or the vermiform appendix.

The last condition has received the names of typhlo-enteritis, from *τυφλος*, blind, and *εντερον*, intestine; typhlitis; and cœcitis, from the Latin word *cæcum*, also signifying blind.

The pericœcal connective tissue is also occasionally the seat of inflammatory action, constituting a condition known as perityphlitis.

SEAT AND CHARACTER.—Clinical experience and the researches of pathological anatomy fully justify us in recognizing the above-mentioned morbid conditions, but the question as to their relative frequency and importance is still far from being settled.

By some authorities the diseases of the cœcum are regarded as secondary to morbid affections of the appendix, the latter consisting generally in the presence of foreign bodies, or of hardened, inspissated mucus, which act as the focus and exciting cause of the inflammation of the cœcum.

It is probable, however, in regard to the simple form of typhlitis, that both the cœcum and its appendix are subject to a peculiar localized inflammation, involving all their coats, and due to the temporary arrest of some foreign substance or intestinal concretion in their cavity, or to the action of the causes to be hereafter considered. It is indeed possible that the inflammation excited by the presence of a foreign body may subside, whilst the cause still remains arrested in the appendix or one of the pouches of the cœcum; but experience would lead us to infer, that, when once inflammatory action has been excited, so long as the foreign substance which has caused it remains in contact with the mucous membrane, the tendency is usually to produce ulceration and preforation of the coats of the bowel.

We find the same discrepancy of opinion in regard to those causes attended with perforation of some portion of the cœcum, and the formation of an abscess in the iliac region. Dupuytren, who was the first to call attention to the pathology of these *iliac abscesses*, attributed them to suppurative inflammation of the pericœcal connective tissue, produced in many cases by extension of inflammation from the coats of the cœcum, and held that the perforation of the bowel often found in connection was a secondary phenomenon, and was in fact the *mode* by which the abscess was discharged. Inflammation and suppuration of the pericœcal tissue

does indeed occur as an idiopathic affection, or from extension of inflammation from the cœcum, but it is of extremely rare occurrence; and there can be no doubt that nearly all cases of iliac abscess are due to perforative ulceration of either the cœcum or appendix. As Bouchut suggests, one proof that most cases of non-puerperal iliac abscess are thus due to perforation of the cœcum or appendix, is afforded by their almost constant occurrence on the right side. Thus of fifty-seven non-puerperal iliac abscesses collected by Grissolle, nine only were on the left side; while of twenty-six puerperal ones, fifteen were on that side.

It is necessary, however, to carry this question one step further, and to determine, if possible, the relative frequency of perforation of the cœcum and of the appendix. It has been supposed, as by Ferrall, that ulceration of the cœcum is in most cases the starting-point in the development of the lesions. But, while we are in possession of a sufficient number of recorded cases, 12 of which we have collected, where post-mortem examination has proved the abscess to have originated in perforation of the cœcum, there is good reason to believe that perforation of the intestine is much more frequently found associated with disease of the appendix than with ulceration of the cœcum itself.

CAUSES.—In addition to the anatomical peculiarities of the cœcum and appendix which must be regarded as predisposing causes of these affections, there are other conditions which exert an unquestionable influence.

The *strumous diathesis* has been regarded as a predisposing cause of diseases of the cœcum and appendix. It does not appear, however, that inflammation of these parts is more frequent in strumous subjects, but merely that it has a greater tendency in such patients to run on to ulceration and perforation of the bowel.

Age.—The greater irritability and proneness to inflammation which the intestinal canal presents in early life, appears to have its effect upon the development of typhlitis, since a considerable majority of reported cases have occurred under the age of 25 years. This is particularly true of the milder attacks, which are not attended with ulceration. Thus, of 42 cases of typhlitis at all ages, which recovered without perforation of the bowel, 32 occurred at or under the age of 25; 10 only were in older persons. Of these 42, 17 occurred in our own practice, and 13 of them were in children whose ages were as follows: 2 under 6 years; 6 between 6 and 12 years; 5 between 12 and 15 years. Finally, 19 of the 42 cases occurred at or under the age of 15 years. This does not appear to hold true, however, with regard to perforative ulceration of the cœcum and appendix.

We have not met with any case of perforation of the cœcum occurring during childhood, but of 25 cases collected from different sources, 13 occurred after the age of 25; 12 at or under that age. Of these 25 cases, 12 only were verified by post-mortem examination, of which 3 were under 15 years of age, 2 between 15 and 25 years, and 5 above 25 years.

Of perforation of the appendix vermiformis, we have met with 3 cases in children, aged respectively $4\frac{1}{2}$, 8, and 11 years. Of 25 other cases, collected from various sources, in which the age is stated, 9 were above, 16 below 30 years of age. Of these 16, 3 only were under 15 years of

age, so that, including our own 3 cases, we find 6 cases occurring under 15 years, 13 between 15 and 30, and 9 above 30 years of age.

Sex.—The influence of sex has been very variously stated by different observers. It appears, however, that males are somewhat more prone to all these forms of disease than females. Thus of 43 cases of typhlitis, which recovered without perforation of the bowel, 27 were in males; 16 only in females. Of 13 of these 43 cases, which we observed in children, 8 were males and 5 females.

Of 25 cases of perforation of the cæcum, 13 occurred in males; 12 in females.

The sex is stated in 27 of 32 cases of perforation of the appendix. Of these, 21 were males; 6 only were females. Of 6 cases occurring under 15 years of age, the sex is stated in 5, 4 of which were males.

Occupation.—Various occupations, especially those involving sedentary habits, have been supposed to predispose to these affections, as also the practice among females of wearing tight corsets. Experience, however, has not verified these suppositions.

Constipation.—A constipated state of the bowels undoubtedly predisposes to these affections by favoring the production of a distended and impacted condition of the cæcum, even if the presence of the hardened fecal matter does not prove the exciting cause of some cases of typhlitis. Rokitansky considers this cause so important that he has given the name typhlitis stercoralis to one form of inflammation of the cæcum.

EXCITING CAUSES.—Cold and Exposure.—The action of these ordinary exciting causes has been denied by some observers on account of the frequent absence of a chill or rigor at the inception of the attack, and the development of the local before the general symptoms. It cannot, however, be doubted that typhlitis may be idiopathic, and arise from the ordinary exciting causes; and, indeed, our recent experience indicates that these influences play a much larger part in the production of this disease than is commonly assumed, though the cases are comparatively rare.

Food.—In several instances the attack appears to have been brought on by the use of indigestible or irritating articles of diet, among which may be especially mentioned unripe acescent fruits. It has been said that the use of oatmeal, which favors the formation of intestinal concretions, is also liable to be followed by this disease. It does not, however, appear that typhlitis is any less frequent in countries where wheaten bread is used, than in those where oatmeal forms a chief part of the food.

Blows or Exertion.—There are a few cases recorded in which a blow upon the abdomen, or a sudden violent strain, appears to have been the immediate cause of an attack of typhlitis; and we have met with several instances ourselves, where the attack could be traced distinctly to such a cause.

Foreign Bodies and Intestinal Concretions.—This class, comprising very various substances, certainly forms an important and frequent cause of diseases of the cæcum and appendix.

We cannot be positive as to the amount of influence they exert in the milder and more tractable cases of simple typhlitis, though it is quite

probable that many of these are caused by the temporary arrest of some foreign substance in the appendix, or one of the pouches of the cœcum. Thus, in a case reported by Dr. Wynn Williams (*Lancet*, January 25th, 1862), in a male adult, three months after a well marked acute attack of typhlitis, which yielded to judicious treatment, a large intestinal concretion, having a plum-stone for a nucleus, was passed by the rectum. They are, however, the efficient cause of a large majority of all the cases of perforative ulceration of the cœcum and its appendix.

The diseases of this latter part, however, are far more uniformly dependent upon the presence of foreign bodies even than in cases of the cœcum; almost three-fourths of all recorded cases of perforation of the appendix having been due to this cause. In 6 cases occurring in children, some extraneous substance was found in the appendix in each one; in 2 a foreign body was present; and in each of the other 4, an intestinal concretion.

Many of these bodies are true intestinal concretions, having for their nucleus merely a nodule of hardened fæces or inspissated mucus. They vary considerably in size, the majority of them being about the size of a cherry-stone or date-stone, though Habershon mentions having seen one as large as a hen's egg. They are also of very varying consistence, according to Volz, as quoted by Hanbury Smith, constituting three varieties: the soft, resembling excrement in appearance and odor, and having a nucleus of hardened fecal matter; the semi-hard, of a grayish-brown color, consisting of shining concrete layers, with a nucleus which is not a foreign body; and the stony, which are of a greyish-white or earthy color, and have a surface from which may be detached delicate scales, or which is smooth, shining, yellowish-white, or brown and studded with calcareous projections.

Many of these concretions consist of carbonate and phosphate of lime, united with inspissated mucus. Copland also mentions one which consisted of cholesterin.

In addition to these, however, numerous foreign bodies have been found in connection with the cœcum or appendix, either free or forming the nucleus of an intestinal concretion. Among these may be mentioned grape-seeds, cherry-stones, date-stones, pins, bristles, fragments of glass, biliary calculi, and balls of worms, either ascarides or lumbricoids.

It may not be amiss to remark here, that some intestinal concretions resemble, to a marked degree, the seeds or stones of different fruits, particularly of the cherry, date, and plum; and there is no doubt that many of the bodies found in the cœcum or the appendix, and reported as cherry-stones or date-stones, have been in reality intestinal concretions.

Whatever be the nature and origin of these bodies, it is probable that in many cases some morbid condition of the mucous membrane of the cœcum or appendix precedes their formation or lodgment, and the development of the grave symptoms which often follow.

As Habershon justly remarks, the ordinary calibre of the appendix is so extremely small and so thoroughly lubricated, that it must be very rare for any extraneous substance to become impacted in it so long as it re-

mains healthy. A further argument in favor of this view is the fact that the presence of these concretions is attended by the most varying results, since very large and irritating bodies have been occasionally found occupying the cavity of the appendix without having produced any symptoms during life, or any inflammation of its surface; while, on the other hand, minute concretions of semi-solid consistence, and apparently unirritating in character, have frequently been observed to act as the foci of the most serious and destructive inflammatory action.

ANATOMICAL APPEARANCES.—In the simple forms of typhlitis, the mucous membrane of the cæcum presents the usual appearances of inflammation; the peritoneal investment is also involved, and besides injection and opacity of this membrane, there are adhesions formed between folds of the intestines.

When, however, ulceration is present, as often results from the presence of foreign bodies, or in strumous subjects, it is a matter of the utmost importance which portion of the cæcum is involved, since, as such ulcers have a strong tendency to perforate the coats of the bowel, if they occur on the anterior part of the cæcum, which has a peritoneal investment, there is the greatest danger of an escape of the contents of the bowel into the peritoneal sac, and the development of rapidly fatal periitonitis. Thus, of 10 fatal cases of perforation of the cæcum, in which the seat of the perforation was determined by post-mortem examination, the anterior wall was involved in 6 instances. If, on the other hand, the ulcer be seated on the posterior part of the cæcum, where it is attached to the iliac fossa by connective tissue, and devoid of a peritoneal covering, perforation is not directly followed by any such unfortunate results. Inflammation is excited in the pericæcal connective tissue, suppuration ensues, and the resulting abscess follows one of several courses, precisely as in idiopathic suppuration of the pericæcal tissue. Thus it may reopen into the bowel; may burrow along the sheath of the psoas muscle, and point below Poupart's ligament; or it may discharge in the lumbar region, or at any point along the crest of the ilium.

In one case the iliac artery was opened, leading to speedy death from hæmorrhage.

Occasionally these abscesses discharge themselves by more than one avenue, as, for instance, through the bowel and in the groin or iliac region simultaneously. When, as occasionally happens, the inflammation or the cæcum passes into a chronic form and the ulcerative process ceases, the adhesions of the cæcum to the iliac fossa become preternaturally dense, the cæcum itself is contracted, its coats thickened, and the mucous membrane almost entirely destroyed, or converted into a retiform and trabecular fibroid tissue. Rokitansky has found in such cases the cæcum converted into a slate-colored capsule, with dense parietes, of the size of a walnut or a pigeon's egg.

The *appendix vermiformis* may be the seat of catarrhal inflammation, associated with inflammation of its peritoneal covering. Death does not result from this condition, but the pathological appearances are probably analogous to those found in all cases of localized sero-enteritis.

When, however, the appendix has been the seat of ulceration, and death has resulted before perforation has occurred, its cavity is found distended with pus, its mucous membrane deeply ulcerated, and in nearly every instance, a foreign body or an intestinal concretion is present.

The ulceration of the appendix varies in its position and extent, at times being seated at the free extremity, at others occupying the lower third of the appendix, which is perhaps the more frequent seat. In regard to its size, the ulcer and the subsequent perforation may be either very small, or else may involve almost the entire circumference of the appendix.

Under favorable circumstances, especially if the foreign body is discharged, the ulceration ceases, and the appendix becomes converted into a ligamentous cord, its calibre being entirely obliterated.

When perforation of the appendix occurs, the results vary according to the degree of local peritonitis which has been excited. If the appendix has become strongly adherent at the point where perforation is about to take place, this accident may not be followed by the development of general peritonitis. The points to which the appendix generally becomes adherent are the cœcum, the anterior abdominal wall, and the right iliac fossa. In the first case, the circumscribed abscess which follows the perforation of the appendix will discharge itself through the cœcum by effecting a perforation of its wall from without inwards, and this is the most favorable termination possible. When, however, the appendix has become adherent to the abdominal wall or iliac fossa, the resulting abscess will follow the course, already described, of abscess from perforation of the cœcum.

It is in this connection that the various abnormal positions which the appendix may assume, are of importance, as determining the position in which the abscess will point.

Unfortunately, however, the adhesions are rarely strong enough to circumscribe the purulent matters escaping from the appendix, so that these generally find their way into the peritoneal cavity, and excite general peritonitis.

We subjoin the histories of 3 fatal cases of perforation of the appendix from intestinal concretions, occurring in children, in all of which some local peritonitis with adhesions had occurred, but had not sufficed to prevent the above unfortunate termination.

CASE 1. Intestinal concretion in the appendix cœci, causing perforation and fatal peritonitis.—T. D. S., a healthy, well-grown boy, 11 years of age, rose on the morning of December 25th, 1860, apparently quite well. Soon afterwards, however, he complained of pain in the right iliac and lumbar regions, was chilly, and returned to bed. A dose of castor oil was given him. In the course of the day fever came on.

Next day he was feverish, with a pulse of 132, a hot and dry skin, and a moderately furred tongue. The pain still continued, with tenderness and slight distension of the abdomen on the right side; there was no vomiting. His bowels had been acted upon three times by the oil. Leeches and a poultice locally, and a mixture of blue pill with rhubarb syrup internally, were ordered.

On the 27th and 28th, the symptoms were much the same, except that the tenderness and distension increased. The pain was aggravated by coughing, by a full inspiration, and by motion, especially of the right leg. The bowels were slightly moved

by the mixture; no vomiting as yet. His fever continued, but the pulse fell to 108, and his skin was somewhat cooler.

On the 29th he was worse. All his symptoms were aggravated, and vomiting set in; his bowels became confined. Small doses of calomel and opium were given, enemata of various kinds were tried, and rhubarb syrup with a little fluid extract of rhubarb was perseveringly employed, but without effect. The abdomen now became greatly distended, exceedingly sonorous, and painful; the stomach grew more and more irritable, rejecting from time to time, towards the last, with a sudden spasmodic effort, everything that was taken by the mouth. The bowels were completely obstructed, so that repeated injections of various kinds elicited no discharges, even of flatus. The urine continued to be secreted to the last; and there was at times, in spite of the nausea and vomiting, quite a strong desire for milk and bread.

During the last few days wine-whey and beef-tea were given in small quantities; and opium by enema and by the mouth was used to allay pain. On the third day of the treatment a blister four inches square was applied over the seat of tenderness; but neither this nor any of the other remedies employed seemed to exert the least effect upon the course of the disease.

Death took place on the eighth day, January 1st, 1861.

The autopsy was made by Dr. Packard, twenty-four hours after death. Body large, muscular, and well-formed; rigor mortis well pronounced. Abdomen only examined.

On making the usual section, several coils of small intestine, very greatly distended with gas, and markedly injected, with flakes of lymph here and there over the surface, at some points gluing the adjacent coils together, were seen concealing the rest of the abdominal viscera. After some search, the colon was found, very much contracted, except at the cæcum. The ileum was in like manner contracted, the narrowing beginning at about the end of the jejunum, which formed the distended coils above mentioned. No cause was assignable for the constriction at this point; but a little lymph was thrown out here, and it may have been that the bowel had been twisted.

The appendix vermiformis was bound down by peritoneal adhesions. Within it, near its origin, was a mass as large as a small bean, but perfectly oval. Just beyond this mass, at what seemed to have been its position, was an ulcer extending all round the tube, and of a gangrenous aspect. At the distal end of this ulcer was a perforation, by which matter had found an exit into the peritoneal cavity. The rest of the tube looked as if it had been distended by the pus before the opening was formed. After its escape from the appendix, the matter seemed to have caused a circumscribed peritonitis, in addition to the general one already indicated. The adhesions bounding this peritonitis had extended up to the liver, the convex surface of which was hollowed to a slight depth in an oval shape, the depression being lined by false membrane. The whole quantity of the pus was perhaps $f\frac{3}{4}$ iv.

The liver was pale in patches, but was not degenerated. Rather too large a number of oil-drops existed in a dark, inflamed portion of its substance, just beneath the depression above mentioned; but even here the quantity was not great. The mesenteric glands were swollen and injected over the surface. No other lesions were observed.

CASE 2. Intestinal concretion in the appendix cæci, causing perforation and fatal peritonitis.—C. B., æt. $4\frac{1}{2}$ years, was taken sick with slight fever, pain in the abdomen, some vomiting, constipation, and inflation of the abdomen. With these symptoms there was marked tenderness in the right iliac fossa. After three days the bowels were well opened, and the fever subsided; the abdomen, however, continued inflated, and a small but distinct tumor had appeared just inside of the right anterior superior spinous process of the ilium.

He continued to improve, and was apparently much better, but was strictly confined to bed, when on the ninth day, at $3\frac{1}{2}$ p. m., he was seized with severe abdominal pains; symptoms of collapse rapidly appeared, and he died at 2 a. m. the following morning.

At the autopsy an intestinal concretion of the shape and size of a date-stone was found in the appendix. The end of the appendix was perforated, and had become attached to the anterior wall of the abdomen, where a small abscess had formed in the cellular tissue between the peritoneum and the abdominal muscles, evidently seeking an outlet through the abdominal parietes. The wall of this had unfortunately ruptured into the peritoneal sac, and death had resulted in a few hours from general peritonitis.

CASE 3. R. P., a healthy girl, aged $7\frac{1}{2}$ years, died at the end of the second week of a well marked attack of perforative disease of the appendix veriformis.

At the autopsy a large, rounded intestinal concretion was found in the appendix cœci, which was perforated, allowing an escape of matter into the peritoneal cavity. There was marked general peritonitis, with the formation of a large quantity of pus.

SYMPTOMS.—Mere *distension* of the cœcum by hardened fæces, without acute inflammation of its coats, may be attended with constipation, some vomiting, and the presence of a somewhat sensitive tumor in the cœcal region. According to Copland, when the distension by accumulated matters is great, it may, from rising high in the abdomen and pressing upon the nerves, vessels, and ducts in its vicinity, occasion numbness and œdema of the right lower extremity, retraction of the right testicle, and derangement of the urinary secretion, so as to be mistaken for disease of the kidney.

Inflammation of the mucous membrane only of the cœcum, is generally attended with a moderate degree of fever, slight pain and tenderness in the right iliac fossa, and some diarrhœa, with mucous, offensive stools. This condition is not unfrequently chronic, and evinces its presence by no very positive symptoms, unless adjacent parts have become involved in the inflammation, or an acute attack of typhlitis supervene.

TYPHLITIS, or inflammation of all the coats of the cœcum or appendix, usually appears suddenly during full health, or it may be preceded by slight intestinal derangement, such as diarrhœa or constipation.

Pain.—The earliest and most marked symptom is generally pain in the region of the cœcum, which appears suddenly, becomes fixed and constant, rarely remitting, and is greatly increased by a deep inspiration or by coughing.

This pain is attended from the very first with such exquisite tenderness on pressure in the right iliac region, that the weight of the bedclothes cannot be borne, and the patient shrinks from the lightest touch. To relieve this pain the patient lies toward the right side, with the thighs flexed upon the pelvis, and any attempt to draw the right leg down causes agonizing suffering. These local symptoms are usually confined to the right iliac fossa, though the entire peritoneum may become somewhat involved, and the symptoms of general peritonitis develop themselves.

Fulness of Tumor.—Owing to the distended state of the bowel itself, and to the adhesions formed between folds of the intestines, or in some rare cases to an inflammatory effusion behind the cœcum in the iliac fossa, there is marked fulness, or even a well defined tumor in the right iliac region. Frequently there will be merely fulness during the first few days of an attack, and then a distinct tumor will be developed. In 14 of 42 cases of acute typhlitis, recovering without perforation of the bowel, a distinct

tumor was present. In most of the other cases the condition of the cæcal region is described as one of fulness or distension. Of these 42 cases, 19 occurred in children under 15 years of age, in only 3 of which a distinct tumor is recorded to have been observed.

Constipation.—The bowels are almost invariably constipated; in many cases very obstinately so. This constipation is frequently associated with quite severe tormina and tenesmus, and if the cæcum be much distended, there may be pain shooting down the right thigh, or numbness and even œdema of this part, together with retraction of the right testicle.

It is important to observe here, that in most cases, when once the constipation is relieved, and free feculent stools procured, the most threatening symptoms of the attack rapidly subside,

Vomiting nearly always attends in children; it was present in all of our 13 cases. It is never stercoraceous, and indeed is rarely troublesome unless the constipation is marked, or perturbing treatment has been adopted in the beginning of the attack.

Fever.—The attack is not usually ushered in by any chill or rigor; but marked febrile symptoms soon appear, the pulse becomes accelerated, the skin hot, the tongue furred, and the thirst extreme. These symptoms usually subside under appropriate treatment after a variable time, generally from four to twelve days; the bowels are opened freely, the pain and tenderness diminish, and the fulness in the right iliac region gradually disappears.

This description of symptoms applies to acute inflammation both of the cæcum and appendix, as there are no well-recognized differences in the symptoms of these two conditions. The only probable points of difference are, that in inflammation of the appendix the pain is more acute, that the thorough evacuation of the bowels is not followed by the same prompt and complete relief.

PERFORATION OF THE CÆCUM.—When, however, perforative ulceration is progressing, the symptoms follow a different course. The constipation may be relieved and the vomiting cease, but the local symptoms persist, until the rupture of the bowel leads either to speedily fatal peritonitis, or to the effusion of fecal matter mixed with the products of inflammation into the pericæcal tissue. When this latter event occurs, the constitutional symptoms soon indicate the occurrence of suppuration, and hectic irritation, with rigors or marked chills succeeded by drenching sweats, colliquative diarrhœa, rapid prostration and emaciation, with a dry brownish tongue and feeble running pulse, soon appear. Despite the desperate character of these symptoms, however, recovery may take place if the abscess points externally in the way already described, and does not open into the peritoneal cavity. It is necessary to be aware that the approach of a fecal abscess to the surface is not attended with the appearances which usually accompany the pointing of an abscess. Thus, instead of the skin becoming tense, prominent, and reddish, with a distinct sense of fluctuation present, the surface becomes doughy and dark-colored, and upon palpation a distinct sense of emphysematous crepitation is often obtained. Upon incising such a point, a discharge of fetid gas and grumous matter follows the punc-

ture, and this peculiarity has more than once led surgeons to believe that they had opened a knuckle of intestine.

PERFORATIVE ULCERATION OF THE APPENDIX.—The symptoms of this disastrous condition closely resemble those of perforation of the anterior part of the cœcum. They are, however, often even more acute, the pain is sudden and violent, and a distinct tumor is more uniformly present; while, on the other hand, the symptoms of obstruction of the intestine are not so well developed. Constipation and vomiting are not constant in the early stage, and at a later period spontaneous diarrhœa may appear, but without any favorable result. The perforation of this part is, as already said, far more apt to be followed by general peritonitis; and, indeed, so far as we know, there is but one well authenticated case on record of recovery after this accident, which was published by one of us in the *Proceedings of the Pathological Society of Philadelphia*. (See *Amer. Jour. Med. Sciences*, vol. liv., July, 1867, p. 145.)

PERITYPHLITIS, or inflammation of the pericœcal tissue, when it does occur independently of typhlitis, is ushered in by pain, with deep-seated tenderness in the right iliac region. There is also some fulness of this part, but not the formation of a distinct tumor, as may frequently be detected in typhlitis. There are usually colicky pains in the abdomen, with either constipation or diarrhœa, and with a moderate degree of febrile excitement. This disease, when judiciously treated, frequently seems to terminate in resolution; when, however, suppuration occurs, the symptoms will approximate those given above, and the abscess which forms may discharge itself externally, into the bowel, or into the peritoneal cavity.

DURATION.—Many attacks of acute typhlitis, when promptly and judiciously treated, yield on the second or third day; though the case is often prolonged to the ninth or twelfth day, and, in violent attacks, it may be many weeks before all local tenderness in the cœcal region passes away, and the function of the bowel is again completely restored. It should be carefully borne in mind also, that after the first attack, there is a distinct tendency to relapses, or to recurrences of typhlitis from slight causes. In our experience, this has been more marked in cases occurring after the age of fifteen years than in children; and in several instances we have seen six series of four, six, or even ten mild attacks recurring under more and more slight provocation, until at length the disease assumed what must be called a chronic form.

When perforation of the cœcum occurs, the after-duration of the case depends entirely upon the point of perforation. If the ulcer have penetrated the anterior wall, general peritonitis is usually excited, and death results in less than forty-eight hours. But if, on the other hand, the posterior wall be perforated, a fecal fistula may be formed, and continue open for very many years. The duration of perforative ulceration of the appendix varies considerably. In three cases in children, observed by ourselves, the duration was respectively seven, nine, and fourteen days, with a mean of ten days.

In eleven cases, at all ages, in which duration is distinctly stated,

the mean duration was nine days, the extremes being two and a half and twenty-nine days.

Bamberger, however, gives the duration of seven cases, occurring at various ages, at from twenty to fifty days, with a mean of thirty-one days. It is probable, however, that this last mean is rarely attained in cases occurring in children.

PROGNOSIS.—Nearly all cases of simple acute typhlitis, without perforation of the bowel, recover under proper treatment. Indeed, there are no cases on record of acute typhlitis proving fatal, in which post-mortem examination did not show the existence of perforation of the cœcum or appendix.

When the cœcum has become the seat of chronic inflammation, however, death may result, either from the sudden development of the acute peritonitis, without perforation of the bowel, or from such contraction of the cœcum as finally to lead to obstruction of the intestine.

When perforation of the cœcum does not prove speedily fatal from peritonitis, but leads to the formation of an abscess in the iliac fossa, the prognosis of the case depends, in a considerable degree, upon the course taken by this abscess. Dupuytren regarded the reopening into the bowel as the safest termination of an iliac abscess, and the opening upon the surface of the body as almost universally fatal. Further experience has confirmed the truth of the first portion of his opinion, but has also established the fact, that almost one-half of the abscesses opening externally recover.

Perforation of the appendix vermiformis is invariably fatal, so far as our experience goes, if we except the case before referred to, where, in an old man about whose past history nothing could be learned, we found the appendix converted into a solid fibrous cord, with a small opening, near the free extremity, leading to its centre.

DIAGNOSIS.—The general diagnosis of most of these conditions is not attended with much difficulty. We have already mentioned that simple excessive distension and impaction of the cœcum is sometimes attended with severe pain, some tenderness, constipation, and even vomiting, and that these symptoms are relieved upon free action of the bowels being secured. We do not have here, however, the sudden attack occurring in a state of perfect health, as in typhlitis, nor the marked febrile symptoms, nor are the local signs in the right iliac fossa, and especially the peculiar, exquisite sensitiveness, nearly so well developed.

Inflammatory disease, in connection with the right ovary, with local peritonitis, is unquestionably sometimes mistaken for typhlitis. The local symptoms in the former affection are, however, lower down in the abdomen than is usual in typhlitis; there is not the well-defined tumor nor the obstinate constipation; and, in addition, there is generally the history of some menstrual trouble, or the attack occurs in immediate connection with the period of menstruation.

Pain in the course of the last dorsal nerve may arise from spine disease, or, in the course of the genito-crural nerve, from the passage of a renal calculus, and, according to Habershon, be confounded with cœcal disease.

It is evident, however, that most of the characteristic symptoms of typhlitis would be absent, whilst a careful investigation of the case would probably educe more symptoms of the existing trouble.

The diagnosis of typhlitis from intussusception, an affection which presents many features of resemblance, will be fully considered in the article devoted to this latter disease.

Ulceration of the cœcum or appendix may be suspected, if the violent pain and the exquisite tenderness persist in the right iliac region, after the other symptoms of an acute attack of cœcal disease, especially the vomiting and constipation, have been overcome. Ulceration of the cœcum is much more apt to have been preceded by bowel complaint for some time; it is also much more rare than ulceration of the appendix.

In cases where we are consulted only after perforation has taken place, with the production of a fecal abscess, we must endeavor, by obtaining a most accurate history of the case, to establish the presence or absence of symptoms of inflammation of the cœcum at the beginning. And further, care must be taken to exclude the following conditions, all of which may at times simulate iliac abscess, namely: psoas abscess, or abscess connected with caries of the pelvic bones; abscesses in the walls of the abdomen, with local peritonitis, resulting from blows; suppuration originating in connection with the right kidney or its envelope; and finally, some cases of disease of the right hip-joint.

The differential diagnosis of these affections of the cœcum and appendix from one another is as yet scarcely possible. The following general remarks contain, perhaps, all that can be surely advanced:

Simple inflammation of the appendix presents symptoms of even greater acuteness and severity than those of simple cœcitis, and which do not subside so promptly after the bowels have been freely acted upon.

In ulcerative disease, both of the cœcum and appendix, the symptoms also persist after the constipation and vomiting have yielded.

Ulceration of the cœcum, however, is rare, and is apt to be preceded by symptoms of bowel complaint. Whilst ulceration of the appendix, on the other hand, is often terribly acute, advancing from a state of apparent perfect health to perforation and death in forty-eight hours; it is also much more frequently attended with a distinct tumor in the right iliac region.

TREATMENT.—The indications for treatment in the acute stage of typhlitis are clearly to reduce the local inflammation of the peritoneum and intestine, to relieve the pain and tenderness, and to secure free and natural action of the bowels. At the same time, all perturbing and strongly reducing treatment is forbidden, by the knowledge that the attack is frequently caused by an irritating foreign body; and that, in a certain number of cases, perforation will occur, in which event the only hope of recovery often rests upon the adhesions which have been formed during the early stage, and upon the vigor of the constitution to resist a prolonged and exhausting process of suppuration.

Depletion.—The local abstraction of a few ounces of blood by the application of leeches to the cœcal region, should be practiced in acute cases.

This measure, while it does not seriously reduce the strength of the patient, relieves the pain and tenderness, and probably facilitates the action of the internal remedies employed. Beyond this degree, however, depletion is injurious, or, at least unnecessary.

Purgatives.—The experience of all observers agrees in condemning the use of powerful, irritating purgatives at any stage of typhlitis. In the early stage, they aggravate the pain and inflammation, increase or establish vomiting, and frequently fail entirely in their object; while, on the contrary, the constipation which will resist the strongest, most drastic purgatives, will quickly yield to mild, saline, or vegetable laxatives.

It is a good plan to combine a small amount of opium with the laxative; since, so far from counteracting its operation, it appears, by allaying the intense sensitiveness of the bowel, to promote its painless and thorough action.

Burne recommends highly the following laxative draught, the dose of which is arranged for an adult :

R. Sodæ Sulphatis, ʒj.
 Tr. Opii, gtt. v.
 Inf. Sennæ, fʒj.—M.
 S.—Repeat every four hours until the bowels are freely moved.

We have ourselves been led by experience to rely upon the combination of comp. ext. colocynth with opium, given in small and frequently repeated doses. Thus, for a child of from five to eight years, the following pill may be prescribed :

R. Pulv. Opii, gr. ij or iij.
 Ext. Colocynth. Comp., gr. xij to xvij.
 Ft. mas. et div. in pil. No. xxiv.

S.—One every three or four hours until free action of the bowels is secured.

Enemata.—The action of these laxatives may be furthered by the administration of large enemata, which may consist either entirely of tepid water, or of water containing a small proportion of some stimulating or laxative substance, such as soap, molasses, or castor oil. In cases where the irritability of the stomach precludes the administration of laxatives by the mouth, enemata become especially important, and at times their use will be followed by the most happy results, the irritating contents of the cæcum being brought away, with almost immediate relief to the most threatening symptoms.

Mercury.—It is difficult to support the practice of giving this drug in typhlitis. In the early stage, indeed, when it may be supposed that the intestinal canal contains irritating ingesta and secretions, a small dose of calomel or blue pill may be administered; and, in a large number of the successful cases on record, this was done. It is not, however, at all necessary. Beyond this, the further use of mercury appears to us injurious since, if it be given until any constitutional effects are produced, it must have a tendency to prevent the formation of those strong adhesions which constitute the sole chance of recovery in case of perforation of the appendix or the anterior wall of the cæcum.

Opium.—We have already mentioned the way in which opium is most advantageously given in this affection, in combination with the laxative employed. Its use is absolutely called for, and the violence of the local symptoms, the pain and exquisite tenderness, form the best guide as to the amount required.

Poultices and Counter-irritants.—In case even the local abstraction of blood appears undesirable, resort should be had to the frequent application of mustard plasters or turpentine stupes to the cœcal region. Hot fomentations or light poultices, to which some sedative substance may be added, should be kept constantly applied to the abdomen.

Vomiting when present, should be allayed by counter-irritation, by swallowing small fragments of ice, by carbonated drinks, hydrocyanic acid, or any other suitable remedy.

The diet during the early stage should be fluid and unirritating in character.

When the persistence of the symptoms leads us to apprehend the presence of ulceration, either of the cœcum or appendix, all depletory and perturbing treatment should be abandoned, and we should limit our efforts to the relief of pain, by the use of opium and the continued application of poultices; to regulating the functions of the intestinal canal, and to the sustentation of our patient's strength.

If perforation has occurred, without the speedy development of general peritonitis, our attention should be mainly directed to supporting the system during the long and exhausting process of suppuration which must ensue. For this purpose a generous, though digestible diet, with as much stimulus as appears necessary, should be enjoined; and resort may also be had to the various tonics, as quinia or the preparations of bark. If a tumor forms, and it becomes evident that the abscess is tending to discharge externally, its approach to the surface should be encouraged by poulticing; and the moment an emphysematous condition of the skin is detected at any point, a free incision should be made, and the discharge of matter furthered by the introduction of a sponge-tent or a pledget of lint, and the application of a poultice.

In those unfortunate cases where the perforation of the bowel has been followed by general peritonitis, all treatment is unavailing. Our main reliance must, however, be placed upon the exhibition of opium, and the use of counter-irritation.

ARTICLE VI.

INTUSSUSCEPTION.

DEFINITION; SYNONYMS; FORMS; FREQUENCY.—Obstruction of the intestinal canal, from one or another of the numerous causes capable of producing it, is an accident liable to occur at all periods of life. But the variety of it which forms the subject of this article is of rare occurrence

excepting in early childhood. It has been called ileus, volvulus, miserere mei; but is best known under the descriptive names of intussusception or invagination of the intestines. It consists in the passage or introduction of one portion of intestine within another, as a small tube might slide into a large one, or, to borrow a familiar illustration, as the end of a glove finger may be pushed back upon itself into the glove. This simple invagination, however, is not the only element present, for in order that the symptoms of intussusception should be produced, it is necessary that the included portion of bowel should be so incarcerated and constricted as to give rise to more or less complete intestinal obstruction. This has led to a very just division of intussusceptions into such as are slight, unattended by inflammation, or spasmodic; and such as are grave, or attended by inflammation and incarceration. The slight form of invagination is found very frequently at autopsies of children who have died of other diseases, and in whom during life there was no symptom of disturbed function of the alimentary canal. It is in all probability produced in the death agony.

M. Louis states that the greater part of 300 children dying during the period of dentition at the Salpêtrière, had 2, 3, or even 4 volvuli without inflammation.

Baillie, Cheyne, and Billard speak of such intussusceptions, as being frequently found at the autopsies of children; and Burns, as quoted by Gorham,¹ gives the results of the autopsies of 50 children who had died from diarrhœa, in every one of which they were found. This species of invagination in children occurs almost exclusively in the small intestine; the invaginated part is usually of no considerable length; and the very slightest traction suffices to restore it.

The grave form, on the other hand, differs from this alike in the very positive symptoms by which its presence is announced, in the condition of the parts involved, and in the part of the bowel affected; and as the form first mentioned scarcely deserves to be called a disease, it is to the latter alone that the following remarks are addressed.

Frequency.—Although numerous well authenticated cases of intussusception occurring in adults are on record, statistics prove that it is relatively much more frequent during the first four years of life. Thus of 100 cases given by Duchaussoy² in which the age is mentioned, there were 31 under 4 years of age, 6 between 4 and 10 years, and 63 adults. Smith's³ tables go to show that "this complaint is rare under the age of 3 months, and that the period of greatest frequency is from the third to the sixth month of life, the maximum number being at the fourth month." Thus there were 11, of the 50 cases collected by him, at the age of 4 months, or 21 in all between 3 and 6 months inclusive; 8 from 6 months to 1 year; and only 18 between the ages of 1 and 12 years.

¹ Guy's Hosp. Reports, 1st series, vol. iii., 1838, p. 330.

² Duchaussoy, Mém. de l'Acad. de Méd., vol. xxiv., p. 97 (New Syd. Soc. Year-Book, 1863, p. 294).

³ Smith, Statistics of Intussusception in Children (Am. Jour. Med. Sci., vol., xliii., 1862, p. 17).

We must, however, call attention to the rarity of this disease at any age among us; for although, in the course of a very extensive practice among children in this city, we have met with several well marked illustrations of the various forms and terminations of intussusception, it has been a rare occurrence in our experience.

ANATOMICAL APPEARANCES.—Intussusceptions, anatomically considered, may be divided into descending or progressive, and ascending or retrograde, according to the direction with the invaginated portion takes; and into central or lateral, according as the entire intestine, or but one wall, is invaginated. Lateral invaginations, however, are exceedingly rare, occurring but twice in 137 cases collected by Duchaussoy.

Excepting when invagination occurs as a complication of some other affection, it is almost invariably of the descending form. Thus, of Duchaussoy's 137 cases, only 16 were retrograde, all of them being complicated; and Haven gives but 3 instances of ascending intussusception out of 59 cases.

It is a matter of considerable importance to determine what is the most frequent seat of intussusception in children. Rilliet and Barthez¹ declare that in infants the small intestine is hardly ever the seat of intussusception, but that ordinarily it is the lower end of the ileum which is invaginated into the large intestine. The reasons for this are found in the anatomical conditions of the intestines in infancy: the adhesions of the cæcum to the right iliac fossa being much more limited and less powerful than in later life; and the muscular coat of the cæcum being but slightly developed in childhood, a circumstance which must also tend to favor the passage of the lower end of the ileum through the valve.

The statistics of Duchaussoy and Smith confirm this opinion; as of 31 cases of simple descending intussusception in children under 4 years of age, collected by the former, the large intestine alone, or both the large and small, formed the intussusception in all but 4 cases; and Smith states that he has found no exception to Rilliet's remark, as regards early infancy. In children above the age of 2 years, fatal invagination in the small intestines may occur in rare cases. In a few cases also, the ileum has preserved its normal relations to the ileo-cæcal valve, the cæcum being the first part inverted, and drawing after it the lower end of the ileum.

An intussusception, then, is made up of three folds of intestine: 1st, The inner, or contained part, which in descending intussusceptions is always in the natural direction; 2d, The middle, which is a reflection of the inner, and passes in a direction contrary to the intussusception; and 3d, The outer, containing part or sheath, which is in its natural position, and in the direction of the intussusception. We find, therefore, the mucous membrane of the middle and outer parts in apposition; and the peritoneal investment of the middle and inner parts in contact.

The amount of intestine invaginated and the condition of the parts depend, in great measure, upon the duration of the case. If death takes place early, only a small portion of the ileum may have passed the valve;

¹ Mal. des Enfants, 2ème ed., tom. i, p. 806.

but as the case progresses, the tenesmus or the active peristaltic action of the outer part, bring down more and more of the ileum with its accompanying mesentery, until finally, the constriction of the ileo-cæcal valve preventing the descent of any more of the ileum, the cæcum is inverted and forced into the ascending colon. This in turn may be invaginated in the descending colon and rectum, until not unfrequently a portion of the invaginated intestine protrudes from the anus. In rare cases, the whole invaginated mass descends into the intestine below, thus forming a double intussusception of great thickness. It has occurred, in a few rare cases, that the amount of constriction was so slight that the intestine remains pervious to a certain extent; so that life has been protracted for many weeks, and death has finally ensued only from exhaustion. But ordinarily the parts are in the following condition: the intestine above the point of constriction is distended with gaseous and fecal contents, and more or less discolored from congestion of its walls. It is rare, however, to find any evidences of enteritis either here or in the intestine below the intussusception, which is generally pale and contracted. The invaginated portion itself, at the upper part, where it seems to plunge into the containing portion of the intestine, presents a series of concentric circular folds. The walls of the bowel thus incarcerated are thickened and infiltrated; their serous investment either deeply injected or discolored by congestion and ecchymosis, so as to be of a deep blackish-red color; and frequently evidences of local peritonitis are present. The mucous membrane in cases of short duration may be merely thickened and injected, but more frequently it is turgid from congestion, ecchymosed in points, and shows the effects of violent inflammation by its unequal roughened surface, presenting either ulcerations or greyish false membranes. The capillaries of the constricted portion become greatly distended, so that, especially in young children, in whom the vascular rete of the intestines is remarkably rich, whilst the tissues are delicate and yielding, they frequently rupture, filling the invaginated intestine with blood, and producing bloody discharges.

If the case is protracted and the powers of life sufficient, when treatment has not sufficed to reduce the intussusception, nature endeavors to effect a cure by eliminating the invaginated portion. The incarcerated bowel becomes gangrenous, a line of separation forms, union and cicatrization take place between the part of the bowel above the intussusception and the upper part of the containing intestine, and the invaginated portion is discharged per anum. This process of elimination is extremely rare in infants; but it is stated by Rilliet to be the ordinary method of cure in children in their second infancy. In 59 cases reported by Haven,¹ of all ages, discharge of the intestine per anum took place 12 times, with recovery in all but two cases. The average length of intestine passed in these cases was $23\frac{1}{2}$ inches; in the two fatal cases, the portions passed was respectively 39 and 44 inches long. The earliest age at which we have met with this process of cure is at 13 months in a case reported by M. Marage.

¹ Haven on Intestinal Obstruction, Amer. Med. Sci., vol. xxx, 1855, p. 351.

In the report of the *Proceedings of the Pathological Society of London*, vol. xiii., a specimen is described by Dr. Hare, where this process had taken place. The patient was a female 41 years of age, and her death resulted from tubercular disease three months subsequently to the passage of the sphacelated bowel, "which was $6\frac{1}{2}$ inches in length, of a very dark purplish-gray color; it formed a perfect cylinder, but the intestine was turned inside out, the exterior of the specimen, as voided, being the mucous membrane, and the interior of the cylinder being the peritoneal covering of the intestine."

At the autopsy, at the point where the invaginated portion had been separated, about fifteen inches above the cæcum, the line of union was found running obliquely across the intestine, "but the union was so perfect that it could scarcely be detected except by holding up the intestine between the eye and the light, when the thinness of the intestine clearly pointed out the line or seam where the union had taken place. Exactly at the point of union the intestine was notably narrower than natural; but the intestine above this point was a little dilated."

We have recently had an opportunity, through the courtesy of Professor Alfred Stillé, of studying a specimen in which a similar process of cure had been effected. The patient was an adult, who died of some chronic disease, and no history could be obtained of the occurrence of the attack of intestinal obstruction, or of the discharge of the sphacelated portion of bowel from the anus. The specimen, however, presented appearances which left no doubt that invagination of a portion of the ileum had occurred, that the invaginated portion had sloughed away, and that union had taken place between the intestine, just above the intussusception, and the upper part of the sheath, so as to preserve the continuity of the bowel. The external surface presented a marked constriction encircling the intestine due to the entrance of the upper part of the bowel into the sheath. There was a layer of organized lymph investing the peritoneum at the line of junction, and firmly uniting the two serous surfaces. Upon laying open this part of the ileum, a narrow rim of indurated tissue, evidently the altered intestinal wall, projected downwards into the intestine from the line of constriction, and formed, as it were, a perforated diaphragm across the calibre of the bowel.

We thus see that even when the slough is cast off, and the patient recovers from the intussusception, the cure is not always permanent, since in a small proportion of cases there may be serious contraction of the bowel, caused by the ensuing cicatrization.

In addition to the modes of recovery already adverted to, namely, the reduction of the intussusception either by the movements of the bowel itself or by the remedial measures adopted, and the elimination of the invaginated portion, there is still a third mode possible, in which the intestine remains invaginated, but by agglutination of the outer folds becomes pervious, and undergoes such atrophy and contraction as not to interfere materially with the functions of the bowel. Rilliet and Barthez, as well as other Continental authors, speak of this as of occasional occurrence, but we have not found any well authenticated cases recorded.

There are few morbid changes found in intussusception excepting those pertaining to the intestines. It is, however, worthy of mention, that in some cases the invaginated mass appears to produce serious compression of the large vessels of the abdomen.

CAUSES: Age.—We have already given the statistics which prove that intussusception is relatively very much more frequent during the first four years of life, the period of maximum frequency being between the third and sixth months. It is very rare before the age of three months. All forms of invagination, however, do not occur with equal frequency at these various ages. During early infancy, for the anatomical reasons already assigned, the almost invariable seat of the invagination is the lower end of the ileum and the upper part of the large intestine; while, after the age of two years, invagination of the small intestine alone, though still very rare, may occur.

Sex.—All statistics agree in giving a majority of males over females, at least in the proportion of 2 to 1; while in some tables the proportion is as high as 7 to 1; thus Rilliet and Barthez collected 25 cases, of which 22 were boys.

Previous Condition.—In by far the majority of cases, intussusception in the infant occurs as an idiopathic affection, appearing during perfect health. In children over one or two years of age, however, it is much more apt to be preceded by some disturbance of the alimentary canal, as constipation, diarrhœa, dysentery, or even by symptoms of imperfect obstruction of the intestines.

Intussusception may also occur during the course of other diseases, as in a case quoted by Rilliet from Legoupil, where the invagination appeared during the progress of variola; the child, $4\frac{1}{2}$ years old, recovered.

Exciting Causes.—*External violence*, as blows upon the abdomen, or sudden jerking of the child's body, as in tossing it in the arms, are assigned as the probable exciting cause of a certain number of cases. It has been supposed, also, that violent fits of coughing or screaming, or strong straining at stool, have produced invaginations, especially in very young children.

Improper alimentation and sudden changes of diet appear to act quite frequently as efficient causes; thus in a case reported by Gorham, occurring in a healthy infant of four months old, the only assignable cause was the administration of panada for three days preceding the attack. It is, however, frequently impossible to assign any plausible reason for the sudden production of severe intussusceptions.

Granting, however, the presence of any of these causes, the question still remains as to the exact mechanism of the invagination. According to Gorham, "it is necessary to the production of an intussusception that there should be either: 1st, A contraction of the part to be intussuscepted; or 2d, A dilatation of that part which is to be the outer fold; or 3d, A natural and sudden inequality of calibre of some portion of the intestinal tube. The first of these conditions may be produced by spasm; the second by flatus; whilst the third is always present at the termination of the ileum in the cœcum." It is at this point, accordingly, that intussusception

most frequently occurs, and, from the anatomical arrangement of the parts making it very difficult for restitution to occur, puts on its most dangerous and fatal characters.

The invagination having once begun, its increase and persistence are probably due to the active peristaltic action of the outer fold, aided by the spasmodic contractions of the diaphragm and abdominal muscles, causing the powerful tenesmus so frequently observed.

There is one more question in regard to the etiology of this affection, about which various opinions have been expressed; whether, namely, enteritis holds the relation of cause or effect to intussusception. Rilliet and Barthez appear to us to have given it its true importance in stating that it sometimes plays one part and sometimes the other. We have already seen that, though in many cases intussusception occurs suddenly in full health, there are a sufficient number of instances where the attack has been preceded by symptoms of intestinal irritation or inflammation, to make it clear that at times enteritis acts as a predisposing or determining cause. And, on the other hand, the pathological anatomy of the disease, showing the inflammation of the bowel to be limited to the immediate vicinity of the invagination, and to be the more intense as the constriction is tighter, proves that enteritis frequently appears as a result of intussusception. This becomes especially evident in those cases where the disease has been caused by external violence, and where after death the above conditions have been noticed.

SYMPTOMS; DURATION; TERMINATIONS.—The principal symptoms of intussusception are furnished by the gastro-intestinal apparatus; and towards the termination of unfavorable cases, by the nervous system. We have seen that a considerable difference exists in the seat of the invagination at different periods of childhood, and in examining the symptoms we find a corresponding disparity, according as the intussusception occurs in the first infancy, under the age of two years, or in the second infancy, between the second and sixth year. These points of difference will be mentioned as each symptom is discussed.

The most important and characteristic symptoms are: vomiting, constipation, and bloody discharge from the anus; abdominal pain, tenesmus, and protrusion of the intestine, the presence of a tumor in the abdomen, and tympany.

Vomiting is an almost constant symptom, being present in about 95 per cent. of the cases. Very rarely the gastric disturbance amounts only to nausea, but nearly always vomiting sets in early in the attack and persists, despite all treatment, until either the invagination is relieved, when it promptly ceases; or until the approach of death. Quite frequently it ceases a day or two before the fatal event occurs. The matters vomited at first consist of the ingesta, the stomach rejecting everything taken into it; soon, however, they become mixed with mucus and bile. In very young children it is rare for stercoraceous vomiting to occur, but in those who are above two years of age it may occasionally be present. In Smith's 50 cases it occurred in three at the respective ages of 3, 6, and 11 years.

The condition of the bowels is generally one of obstinate *constipation*, so far as the passage of fecal matters is concerned. It is not unusual for one natural abundant stool to occur after the intussusception begins, but this is succeeded by constipation. It is only in those very rare cases where the invaginated portion remains pervious, that a small amount of fecal matter finds its way into the stools.

The discharges which, however, do take place almost invariably in intussusception in children are due to the rupture of the capillaries of the constricted bowel, and consist of blood mixed in varying proportions with mucus and serum. It is rare for the blood to be so deficient that the discharges resemble the gelatinoid mucous discharges of dysentery, merely streaked and tinged with blood, whilst, at times, the blood is in such excess as to appear pure, and to constitute a true intestinal hemorrhage. This symptom, the true value of which was first recognized by Gorham and Clarke,¹ is of more uniform occurrence in children under two years, on account of the greater ease with which the intestinal capillaries give way in infancy. Thus of 26 children under one year of age, bloody evacuations occurred in 23, usually several times in the twenty-four hours; in 2 of the 26 there is no record of this symptom, and in 1 only is it recorded as absent. In case No. 2, of Mr. Gorham's table, a child of 3½ months passed within a few hours more than a teacupful of fluid blood. In older children, on the other hand, bloody discharges occur less frequently; thus Smith records 18 cases of invagination between one and two years, in only 6 of which it is stated that there were bloody motions.

We have already mentioned the various ways in which recovery takes place, and when elimination of the invaginated portion is about to occur which is almost exclusively limited to cases occurring in the second infancy, the stools become highly fetid, contain more or less blood, are blackish or brownish in color, and are soon accompanied by the discharge of the slough. The interval elapsing between the inception of the attack and the discharge of the portion of bowel varies considerably in different cases, but seems to be less in childhood than in adult age. Thomson states that in adults the elimination takes place in the majority of cases within thirty days; and in one of his cases it occurred as early as the sixth day. In children the interval rarely exceeds twelve days; and the average of all recorded observations would seem to fix about nine days as the usual time.

Abdominal pain is among the earliest and most constant symptoms at all ages. During the early part of the attack, it appears in paroxysms; and may be detected even in the youngest children, by the violent paroxysmal screaming, and contortions of the limbs and trunk. At the commencement, the abdomen is generally relaxed, supple, and indolent; and this condition may remain until death, perhaps because the constriction in some cases is not complete and allows the passage of gas. But, after a few days, there is apt to be more or less continuous pain and soreness on pressure in the part of the abdomen corresponding to the invagination, due to the local enteritis and peritonitis. This may or may not be accompanied

¹ London Lancet, January, 1838.

by tympany and diffuse tenderness of the abdomen ; but, as a general rule, intussusception in very young children is not attended by the great distension and marked symptoms of general peritonitis which frequently appear in intestinal obstruction in adults. In children over two years of age, the abdominal symptoms are more apt to indicate peritonitis. In a considerable proportion of cases, tenesmus occurs and adds much to the suffering. It does not appear so early as the abdominal pain, and generally ceases a few days before death.

Tumor.—It would appear natural that when a considerable intussusception has taken place, the knot formed at the point of obstruction should be readily detected through the abdominal walls. And yet the cases on record show that this tumor is recognizable in not more than two or three out of every ten cases. When it can be detected, it is generally found in the left iliac region, varying in size from a walnut to a large goose-egg, and giving the sensation of a solid, but doughy and compressible mass. It is ordinarily quite movable, and percussion elicits a dull note over its position.

Another symptom depending upon the displacement of the intestine, to which considerable importance has been attached in the diagnosis of invagination in the adult, is a depression of the abdomen at a point corresponding to the displaced intestine, and a fulness at the corresponding point on the opposite side. Experience has shown, however, that but little value can be attached to this sign in young children, on account of its great rarity.

We have seen that the presence of a tumor in the abdomen is far from an invariable sign of intussusception, and the same remark applies to the *protrusion of the invaginated bowel from the anus*, a symptom to which very different diagnostic value has been attached by different authors. It is stated by some to be hardly ever present, but we have found it recorded particularly in six of Smith's cases, the same number in which an abdominal tumor was present in the same series ; and in three other cases, although no tumor protruded from the anus, the invaginated mass was readily felt by examination per rectum.

When the bowel protrudes, it forms an oblong tumor, at times even two inches in length, much congested from the constriction, and smeared with blood and mucus.

When we pass from these positively diagnostic symptoms, we find little elsewhere characteristic of the disease. The tongue is normal until inflammatory action sets in, when it often becomes dry and brown ; the appetite is impaired or absent, and the thirst is generally but moderate. Rilliet and Barthez call attention to the importance of this last symptom in a diagnostic point of view, as well as to the fact that the emaciation is usually not so marked as in other acute diseases of equal duration and severity.

The amount of febrile action is generally slight in infancy ; the surface, cool at first, may at times become hot, or is alternately hot and cold, and as death approaches remains continuously cold. The pulse soon be-

comes frequent, though small and feeble. There is no marked disturbance of respiration.

In older children there is apt to be more febrile action, the skin being hot until late in the attack, and the pulse frequent and more full. The physiognomy of the little patient is greatly altered from the commencement of the attack. The eyes are dull and languid, sunken in their orbits, and surrounded by discolored areolæ; the countenance is expressive of the most profound prostration, so as to have elicited a comparison to the physiognomy of cholera patients.

Almost all cases, at whatever age, present symptoms of marked disturbance of the nervous system, as great restlessness, indescribable malaise, sharp cries, and, toward the close of the case, profound prostration. But in infancy, in addition to these symptoms, the case is more apt to present an attack of convulsions, either as one of the earliest symptoms, or toward death, alternating with coma.

Duration.—It is necessary to distinguish here between cases occurring during extreme infancy, when we cannot hope for elimination to take place, and those in more advanced childhood. In early infancy, when the attack is about to take a favorable turn, the symptoms usually yield in from two to four days, owing to reduction of the invagination. In fatal cases, death occurs within five days, as the rule. In some cases, however, where the constriction was not complete, life has been prolonged even for six weeks.

In second infancy, where the constriction is complete, and the result fatal, death occurs within seven or eight days in the vast majority of cases. But when elimination is to result, the case is more protracted, and complete recovery is postponed to the third week. Thus, in 7 cases out of Smith's statistics, which resulted favorably by sloughing, the ages were 5, 6, 6, 9, 11, 12, and 12 years respectively; and the separation of the invaginated portion took place between the ninth and twelfth days, with an average of nine and a half days. After the discharge of this, which is soon followed by the fetid, brownish-black stools already described, the symptoms rapidly disappear, and in one or two weeks the cure is complete; so that, if we can carry a patient, advanced beyond the first infancy, through the first week of the attack without too much exhaustion, we may each day look for the discharge of the invaginated bowel, the restoration of the function of the intestines, and ultimate recovery.

Terminations.—We have already described the favorable modes of termination, namely, by the subsidence of the intussusception, either spontaneously or as the result of treatment; by restoration of the calibre of the bowel by sloughing of the invaginated bowel, and union and cicatrization of the divided edges; and finally, by agglutination of the outer layers of the invaginated portion with subsequent thinning and atrophy, thus rendering the intestine pervious, although the intussusception remains.

In those cases in which death takes place very early, as on the first or second day, it is frequently produced by cerebral congestion or an attack of convulsions. In the majority of cases, however, it occurs somewhat later, and is preceded by a state of collapse. Even in those cases where

the constriction is not at first complete, and where there are daily feculent evacuations for a time, death is apt to occur from exhaustion, or from the invagination becoming more extensive and symptoms of complete obstruction arising.

PROGNOSIS.—A single glance at the character of the lesion and the accompanying phenomena, suffices to assure us of the grave nature of intussusception, and of the impotence of all ordinary methods of treatment against it. In young infants, indeed, where the strength of the system cannot be expected to hold out until elimination occurs, intussusception is almost invariably fatal. In a single instance only has recovery by elimination been noticed so early as the end of the first year. In a few cases, where the symptoms were well developed and threatening, they have subsided and the infant has recovered, apparently from spontaneous reduction of the invagination.

We must not, however, forget that during the early stage of this affection the diagnosis is somewhat doubtful, since young children frequently present symptoms of obstructed and loaded intestine, such as a distended, hard abdomen, constant unnatural straining, with evident suffering, and yet are entirely relieved after the administration and operation of laxatives.

Not to refer now to the recent cases of successful abdominal section, a considerable number of cases of cure of undoubted intussusception, by means of inflation, have also been reported even at this early age; so that, when treatment is instituted soon after the appearance of the symptoms, the case is not absolutely hopeless. In older children, that is to say above three years of age, the prognosis is much less unfavorable, since treatment offers a certain amount of hope, and there is always the prospect of the occurrence of elimination of the invaginated bowel, if the strength of the patient has been sustained during the first week.

Even after elimination has taken place, however, the prognosis should still be somewhat guarded, as the slightest indiscretion in diet may, either by the development of flatulence or by the escape of irritating, undigested particles into the intestine, cause a rupture of the recently formed cicatrix and speedy death.

DIAGNOSIS.—Intussusception has been, until recently, regarded by all authors as an affection of obscure and doubtful diagnosis. With the light, however, which has been thrown upon this subject by the labors of Clarke, Gorham, Smith, and especially Rilliet, the diagnosis in the great majority of cases can be made with precision. It is true, however, as conceded by Rilliet, that "very rarely in early infancy, more frequently than later, there are certain cases of invagination impossible to distinguish from other forms of intestinal obstruction; and that at all periods of childhood the diagnosis presents many difficulties."

With what diseases, then, could we confound this affection, occurring, as we have seen, suddenly in perfect health; attended by obstinate, though rarely fecal vomiting; by marked constipation, but with frequent bloody discharges; by paroxysmal abdominal pain and tenesmus; by the presence of a tumor, generally in the left iliac region; by the protrusion of the in-

vaginated bowel from the anus; and by profound prostration and disturbance of the nervous system? It is to be remembered, indeed, that this group of symptoms, so characteristic when viewed together, are rarely all present; and that with the exception of the vomiting, constipation, and bloody discharges, there is no single symptom which is not more frequently absent than present. There are, nevertheless, a sufficient number present in nearly every case to enable us to form a diagnosis.

The diseases which may most readily be confounded with intussusception are, 1st, impaction of the intestine with hardened feces; 2d, typhlitis or perityphlitis; 3d, cholera infantum; 4th, dysentery; 5th, intestinal hemorrhage; 6th, the various forms of internal strangulation; 7th, peritonitis.

1st. When an accumulation of fecal matter takes place in either the cœcum or sigmoid flexure, the case may present many symptoms similar to those of intussusception. There is frequently such gastric and intestinal irritation as to lead to occasional vomiting and paroxysmal abdominal pain; the bowels are constipated, and there is frequent and strong tenesmus, so as often to cause protrusion of the bowel. In addition to these symptoms, a well defined tumor is present in one or the other iliac fossa.

These cases, however, often have presented symptoms of intestinal disturbance for some time previous to the attack; the vomiting is rarely so constant as in intussusception; the tumor is quite painless and has a peculiar doughy consistence; bloody discharges from the bowels are very rare; and we do not notice the profound prostration which exists in well established invagination. During the early stage of the case, however, the diagnosis is doubtful; and when we have reason to suspect the presence of fecal accumulations, we must await the result of the administration of laxatives and laxative enemata, before deciding upon the nature of the case.

2d. Inflammation of the cœcum, appendix vermiformis, or of the pericœcal connective tissue, is attended with fulness or a well defined tumor in the right iliac fossa, with vomiting, constipation, and occasionally tenesmus, with distension of the abdomen and pain radiating from the right iliac region.

There is, however, a marked degree of fever, and the symptoms of local peritonitis appear early in the case; the patient assumes a characteristic position, with the thighs flexed upon the pelvis, and the right iliac fossa is the seat of exquisite tenderness, so that the slightest pressure cannot be tolerated. The vomiting and constipation are not so marked and obstinate, and excepting in those cases which have been preceded by dysenteric symptoms, there are no bloody discharges, and as we have remarked above, the tumor or fulness is in the right iliac fossa; whereas when this sign is present in intussusception, it usually occupies the left iliac region.

3d. In cholera infantum, the vomiting is often incessant; the stools are frequent, with painful tenesmus; the abdominal pain paroxysmal, and occasionally the intestine protrudes from the anus. It is almost impossible, however, to mistake this affection for intussusception, if we remember that

it is almost always accompanied by fever, with insatiate thirst, and prompt and extreme emaciation; that the abdomen is without tumor, and rarely distended until towards the close of the case, and that the stools, instead of being bloody, are large and fluid.

4th. Dysentery frequently offers a close resemblance to intussusception so far as the characters of the stools are concerned, as they are often small and bloody, or muco-sanguinolent. But we do not see in dysentery the sudden inception, the rapid progress, the obstinate vomiting, the moist tongue and moderate thirst, which characterize intussusception.

5th. We have seen that occasionally the amount of blood passed by stool in intussusception is very great, and constitutes a true intestinal hemorrhage; thus in the case reported by Marwick,¹ it amounted to a large teacupful of pure blood.

Intestinal hemorrhage is a very rare occurrence during childhood, but has been noticed in children in connection with polypus of the rectum, especially by Mr. Bryant; in typhoid fever, or the hemorrhagic form of some others of the exanthemata, and in the course of purpura. The absence of the other symptoms of intussusception, however, and the presence of the local or general symptoms peculiar to these various conditions, will serve to render the diagnosis easy.

6th. Other forms of internal strangulation, such as those produced by a diverticulum from the intestine compressing it, by the adhesion of the vermiform appendix so as to constrict the bowel, or by a contraction of the calibre of the bowel, produce symptoms so identical with those of intussusception in second infancy, when the affection more nearly resembles intestinal obstruction in the adult, as to render diagnosis impossible. The presence of an abdominal tumor, the occurrence of bloody stools, or the protrusion of the constricted bowel from the anus, would be the only diagnostic signs.

7th. Peritonitis, when diffuse, presents a few symptoms in common with intussusception; as the vomiting, constipation, abdominal pain and tenderness; and when the inflammation of the peritoneum is localized, there is in addition a well defined sensitive tumor, which soon appears as the result of the inflammatory action. The diagnosis here rests upon the greater frequency of the vomiting in intussusception, the more obstinate constipation with bloody discharges from the bowels; the paroxysmal nature of the abdominal pain, with less tenderness; the less degree of fever, the moist tongue, slight thirst, quiet respiration, and only moderately accelerated pulse.

TREATMENT.—There is no special plan of treatment for intussusception deserving the name of *preventive*, owing to our ignorance of any symptoms which can be definitely regarded as the precursors of the invagination. The fact, however, that various derangements of digestion, such as pain upon going to stool, diarrhœa, or constipation alternating with diarrhœa, have been occasionally noticed to precede the attack, should be an additional motive to urge us to meet these symptoms by the most assiduous

¹ London Lancet, July, 1846.

attention to the hygiene of the child, and to the regulation of its alimentary functions.

The *curative* treatment may be divided into three classes : the medical, mechanical, and surgical treatment.

Medical Treatment.—*Depletion* is strongly contra-indicated by the tender age of the patients, and by the necessity of preserving the vital powers; since elimination, which affords the principal chance of recovery, does not occur until after the eighth day. In order, however, to relieve the engorgement at the point of constriction, without reducing the strength of the patient, it is advisable to apply a few leeches or cups to the abdomen, and preferably to the right iliac region, unless a tumor can be detected, when, of course, they should be applied over its seat.

Purgatives were formerly strongly advocated by most authors; the one most generally advised being quicksilver, which was given with a view of overcoming the obstruction by its great weight and fluidity. The use of this agent is now, however, universally reprobated.

In regard to other and less mechanical purgatives, there is still some difference of opinion.

During the early stage of the attack, before the symptoms of intussusception are very positively developed, we should advise the administration of a mild but thorough laxative, such as castor oil, in conjunction with large laxative enemata. If, however, at the end of twenty-four or forty-eight hours, the administration of these remedies, aided by the local depletion, has failed to produce an evacuation from the upper bowel, these measures should be abandoned, and recourse be had to means of calming pain and nervous disturbance, and to the sustentation of our patient. Among the remedies best calculated to allay the pain, the tenesmus, and the nervous irritability are : opium, in doses proportionate to the intensity of the pain; warm anodyne poultices applied to the abdomen, and warm baths carefully given. These latter are especially serviceable when the symptoms of nervous disturbance are marked, even amounting, as they occasionally do, to general convulsions.

In endeavoring to sustain the child's strength, attention must be paid to the vomiting, which is generally so severe as to prevent any nourishment being retained. The remedies of most service against this are counter-irritants to the epigastrium, opium, hydrocyanic acid, carbonated water, small pieces of ice kept constantly in the mouth or swallowed whole.

Nutritious enemata may also be tried, but are rarely retained.

The *mechanical* treatment consists in the injection of fluids or air into the bowel in such quantities as to distend it; and in the introduction of a large sound, with the view of pushing up the invaginated portion of intestine. The fluids generally used have been either tepid water or warm gruel, injected forcibly into the bowel, until the sudden cessation of resistance informs us of the reduction of the intestine. We have already seen that the seat of intussusception in the child is almost invariably the lower end of the ileum, which passes into the cœcum and is there constricted; and, when we reflect that it has been frequently demonstrated that if fluid be forcibly injected into the large bowel, the ileo-cœcal valve will rupture

before any fluid is allowed to pass into the ileum, it is evident that we can in this way exert a most powerful pressure upon the invaginated intestine. Experience shows that this procedure is frequently successful, even in cases where all medicinal treatment has proved unavailing; and there are now a sufficient number of such cases on record to render a resort to it proper. The fluid may be introduced by an ordinary syringe, or better by a Bowditch's syringe, the limbs being held together so as to prevent as far as possible any reflux. It has been recently suggested by Simon that hydrostatic pressure might be employed to force fluid into the bowel. For this purpose a glass funnel attached to a long india-rubber tube terminating in an olive-shaped plug is used. The plug is inserted in the anus, and the funnel is held on a level with the body, and water poured in until it is filled. The funnel is then gradually elevated, and more and more water poured in to replace that which is forced by the hydrostatic pressure into the bowel. Owing to the gradual and uniform increase in pressure thus brought about, extreme distension of the entire colon can thus be produced. We have recently employed this mode of treatment with most gratifying success in a very severe case in an infant of 6 months of age.

Air, also, both on account of its great elasticity and mobility, as well as the great facility of its introduction in sufficient quantity, is to be highly recommended. Indeed, inflation was advised by Hippocrates as a remedy in intussusception, but until within the past forty years does not seem to have been much practiced. Two cases of obstruction of the bowels, occurring in adults, successfully treated by inflation, are reported in the *American Journal of Medical Sciences*, for 1833; one by Dr. Janeway, of New York; the other, which, however, was transcribed from the *Glasgow Medical Journal*, for 1831, by Dr. King. The following year, in the *Boston Medical and Surgical Journal*, December 15th, 1834, Dr. J. Wood published a case, also in an adult, where death seemed imminent, but where the obstruction was readily overcome by inflation, and the patient recovered. Since then, this remedy has been frequently employed in intussusception in children, and with such good results, that it may fairly be said that the prognosis of this affection is less grave since the introduction of this remedial measure. To obtain the best results, inflation should be employed early in the case, before any considerable amount of adhesive inflammation has taken place between the sheath and the contained intestine. The air is readily introduced by a pair of ordinary bellows; the nozzle being inserted well into the rectum, and inflation continued until the obstruction yields. The return of the invaginated intestine is sometimes attended by a clearly audible sound, a species of crack, but it never gives any pain, and has generally seemed to afford relief. The complete restoration of the calibre of the intestine is proved by the copious feculent stools which frequently come away soon after the inflation.

A third mechanical means for restoring the displaced intestine has been recommended by Dr. Nissen, and consists in pushing up the invaginated portion by means of an œsophageal sound protected by a sponge. This proceeding would probably be readily accomplished, if the intussusception occurred far down in the large intestine; but it would appear very

difficult to replace in this way an invagination as high up as the ileo-cæcal valve. Dr. Nissen, however (in the *Journal de Canstatt*, quoted by Rilliet and Barthez), gives two cases in which he succeeded in pushing up the intestine into the ascending colon, with complete relief of the symptoms of obstruction. There are also a few other cases of cure, by this means, upon record in medical literature.

The *surgical* treatment consists in the performance of the operation of gastrotomy, finding the invaginated portion of bowel and reducing it by gentle traction. We had already expressed ourselves in favor of this operation under certain circumstances, while there still existed much diversity of opinion on the subject and many authors condemned it. Their disapproval was based upon the grounds of the great difficulty of ascertaining the exact position of the intussusception; the difficulty of restoring the invaginated intestine even if found; and finally upon the dangers of the operation.

We have seen, however, that in the majority of cases the invaginated mass will be found in the neighborhood of the left iliac fossa; the lower end of the ileum having traversed the cæcum, ascending and transverse colon, and these parts being successively inverted; that in a certain proportion of cases a tumor is readily detectable; and further, that some idea as to the seat of obstruction may be obtained from the distance to which enemata appear to penetrate. So that in a considerable proportion of the cases we have the means of localizing the point of constriction with a certain amount of definiteness.

In regard to the difficulty of reducing the invaginated parts, authors differ greatly. It has been remarked, that even if the equivocal and uncertain nature of the symptoms of volvulus were not sufficient to deter us from undertaking the operation, the state of the invaginated parts would entirely banish all thoughts of such an imprudent attempt; since the different folds of intestine become so agglutinated to each other than they can hardly be withdrawn, even after death.

Rilliet and Barthez (*loc. cit.*), however, conclude from their anatomical researches, that in the majority of cases the disengagement of the intestines is very easily accomplished; and accordingly they declare that, "after employing medical treatment during three or four days, and after having made several attempts at inflation, we should not hesitate to perform gastrotomy."

The great danger of the operation is, of course, apparent, but should hardly be considered an objection, when we consider the fatal nature of this affection. Nor have the results of operation been such as to destroy hope. In addition to several successful operations previously recorded, the only 3 cases out of the 57 collected by Haven, in which gastrotomy was performed, terminated favorably. More recently, also, the operation has been performed several times by different operators (J. Hutchinson, Howard Marsh, Legge, Sands, and others), and with such encouraging results as to fully justify us in repeating our former advice in regard to its performance.

To sum up our remarks upon this subject: after having tried for two

or three days the medical and mechanical means recommended without success, we must forbear and decide whether to trust the case to nature, with the hope of elimination of the invaginated bowel occurring, or to resort to gastrotomy. And in this decision, the circumstances of each case must be taken into account; for if the case has not yet progressed so far that adhesive inflammation has certainly taken place, and if we are able to detect the exact seat of constriction by the presence of a tumor, the operation certainly has strong arguments in its favor, and should not be hastily rejected.

In those cases which have been trusted to nature, and when elimination has fortunately occurred, we must treat the child, during this crisis, with the utmost care. The diet must be rigidly regulated, and the child kept in absolute repose. Nor must we relax these precautions for several weeks, and allow either indigestible food, or too large a meal of even the most digestible articles; since death has been several times known to follow this imprudence, from a rupture of the imperfectly formed cicatrix.

CLASS IV.

DISEASES OF THE NERVOUS SYSTEM.

GENERAL REMARKS.

It is a very common opinion, both in and out of the medical profession, that this class of diseases occasions a much larger number of deaths in childhood than any other. Indeed, it was formerly supposed by many persons that, whatever the primary disease might be, nearly all children who died, died, as it was said, by the brain. The careful study of mortality statistics and the advance of pathological knowledge have effectually disposed of this idea, and have shown that in a large proportion of fatal cases where nervous symptoms have been prominent towards the close, these phenomena were merely the result of functional derangement sympathetic with the primary disease, or due to the circulation in the blood of some specific poison.

Before beginning the consideration of the particular diseases of this class, we are desirous of stating that we shall be compelled, on account of our limited space, to devote attention chiefly to those which are most important from their frequency or severity, avoiding or merely alluding to those which are of less consequence, or which occur in childhood merely in common with adult life.

In our earlier editions we divided this subject into two classes, one containing all the diseases attended with and dependent upon some appreciable alteration of the nervous centres, the second containing those in which no such alteration exists. We have since discarded that arrangement, principally on account of the minute researches of histologists during the past few years, which have all gone to prove the existence of positive and definite tissue changes in many diseases previously regarded as purely functional.

ARTICLE I.

TUBERCULAR MENINGITIS.

DEFINITION ; SYNONYMS ; FREQUENCY.—This disease is characterized by violent cerebral symptoms, dependent upon the existence of tubercular granulations in the pia mater, as the essential anatomical lesion ; accompanied in the great majority of cases, by coincident inflammation of that membrane, by softening of the central parts of the brain, by effusions of serum into the ventricles, and in many instances by tubercular deposits in

other organs. Formerly tubercular meningitis, simple acute meningitis independent of tuberculization, and simple dropsical effusion within the cavity of the cranium independent of inflammation, were confounded together under the single term of acute hydrocephalus or water on the brain. It has been shown, however, that a large majority of the cases of acute hydrocephalus of authors are, in fact, cases of tubercular meningitis, and more recent researches have further shown that most of the remaining cases are in reality due to the altered condition of the blood, called uræmia, and are independent either of any material lesion of the brain or of the presence of an excess of serous fluid in its cavities.

The term acute hydrocephalus ought to be therefore restricted to the single condition of sudden serous effusion in or around the brain, independent of any inflammation; a condition which only occurs in connection with the causes of general dropsy, and especially with renal disease, and is, indeed, merely the most rare form of internal dropsy, and, as such, not to be regarded as a separate disease. A description of the symptoms of this condition will be found in our remarks upon the renal complication of scarlatina.

There can be no doubt that tubercular meningitis is of rather frequent occurrence, though it is difficult to obtain statistics which will enable us to form anything like an accurate idea upon this point. M. Barrier (*loc. cit.*, t. i, pp. 34, 36) states that during the period in which his observations were carried on at the Children's Hospital in Paris, there occurred 576 medical cases of all kinds. In this number there were only 10 cases of tubercular meningitis, whilst there were 83 of pneumonia, 48 of pleurisy, 24 of typhoid fever, 48 of measles, etc., etc., showing the first-named disease to be much less frequent than many other affections. We may also form some idea of its frequency in proportion to other diseases, by a reference to the work of MM. Rilliet and Barthez (1ère édit.), who report 33 cases of tubercular meningitis, against somewhat over 245 of pneumonia, 174 of bronchitis, 111 of typhoid fever, 167 of measles, and 87 of scarlet fever. We are of opinion that it is not of frequent occurrence amongst the easier classes of this city, since we have met with less than 60 cases in private practice in the course of thirty-five years. We observe it more frequently, however, in our large children's hospitals, and from what we have been told by other practitioners, it seems probable that it is much more common among the destitute classes, and particularly the blacks, who crowd the southern parts of the city, and who suffer to a great extent from tubercular and scrofulous diseases. It is, however, impossible to obtain accurate information in regard to the frequency of the disease in this city, in comparison with other affections of the brain, from a reference to the bills of mortality. Thus during the year 1874, with a total mortality of 16,254, there were 8349 deaths among minors; of these, 143 are recorded as from hydrocephalus (tubercular meningitis), 382 as from cephalitis, 211 as from congestion of the brain, 83 as from brain disease undefined, and 654 from convulsions. It cannot be doubted that a considerable number of cases of tubercular meningitis are included under these latter vague headings.

PREDISPOSING CAUSES.—MM. Rilliet and Barthez (2ème édit., t. iii, p. 511) state that the disease is very rare in the first year of life; that it becomes notably more frequent in the second year, but that it is between two and seven years of age that it occurs with the greatest frequency. After this, it diminishes, they say, rapidly from eight to ten, and especially from eleven to fifteen years of age. The influence of sex has not been determined, but it appears probable that boys are somewhat more subject to it than girls. It has been clearly shown by the observation of various writers that the disease usually attacks delicate children, and especially those born of parents who are either themselves laboring under tuberculosis, or in whose families that diathesis has existed to a greater or less extent. Of the 31 cases that have come under our own observation in which we have preserved complete notes of the disease, in 20, one of the parents either had phthisis at the time, or died of it subsequently; in 3, one or the other parent came of a tuberculous family, though in these both parents were living at the time in seeming good health; in 4, no trace of tuberculosis could be found in the parents or in their families, and in 4 the history of the parents or of their families could not be traced out. It is not uncommon for several children in a family to die of tubercular meningitis. Under these circumstances, it has nearly always been ascertained that the parents, or some of the immediate relations, have either died of tuberculous or scrofulous disease, or shown unequivocal signs of one of those diatheses. Thus, 4 of the 20 cases mentioned above occurred in two families, in one of which the father is since dead of phthisis, and in the other the mother has long been ailing with inactive tubercle of the lungs, and slow caries of a bone, in all probability of tuberculous origin. It may follow other diseases, and has been observed particularly after measles and other fevers, and after the suppression of eruptions.

M. Barrier (*op. cit.*, t. ii, p. 379) explains, and we think with good show of reason, the causes of the disposition on the part of the tubercular diathesis in children to localize itself in the brain, as well as the disproportionate violence and extent of the inflammatory action in comparison with the degree of the tubercular lesion, by the physiological conditions of the nervous system in early life, which are those of great functional energy and nutritive activity. The affection, though much more frequent in childhood, is by no means peculiar to that period of life, and we have met with, in addition to the cases above referred to, a number of cases occurring in the adult, and presenting the same general clinical symptoms and anatomical lesions.

As to the *exciting causes*, nothing positive is known. The disease has been supposed to be brought into action by falls and blows upon the head, by violent moral emotions, and by exposure to the sun. These causes, however, are all of doubtful influence.

Recent pathological investigations have established the fact that, in many cases, the development of true miliary tuberculosis of the cerebral membranes, or of other tissues, is connected with the previous existence of foci of cheesy degeneration, as in an enlarged lymphatic gland, a patch of unabsorbed pneumonia exudation, or otherwise. Undoubtedly such a con-

dition exerts its power of infecting the general system, and leading to the development of tuberculosis, especially when there exists a hereditary predisposition to that disease. In a number of instances, we have been able to trace the origin of tubercular meningitis to this cause.

ANATOMICAL LESIONS.—The tubercles which constitute the essential anatomical element of the disease are very rarely found upon the free surface of the arachnoid, but almost invariably beneath that tissue, or in the meshes of the pia mater. They usually appear as more or less opaque gray granulations, the so-called *miliary tubercles*, and may generally be seen through the arachnoid, scattered about in the shape of small, rounded, or flattened bodies, of grayish or yellowish-gray color, and varying in size from two-fifths to four-fifths of a line. When the finger is passed over the arachnoid above them, they may be usually felt as little granular bodies. Their size, however, varies very much, and they are in some cases so small and so closely resemble in color the surrounding parts, that it requires a careful search to detect them. They vary also greatly in number, being in some cases thickly scattered over a considerable extent of the pia mater, while in other cases but two or three can be discovered on each hemisphere.

Frequently they can be detected with most ease upon the processes of pia mater which dip down between the convolutions, so that if we fail to find any granulations upon the surface, we should always strip off the pia mater and carefully examine these processes. Upon a careful examination of the arrangement of the miliary tubercles, it will often be observed that they are clustered about the small arterioles of the pia mater, and evidently follow in their distribution the branches of these vessels.

These granulations are not found upon all portions of the brain equally in cases of tubercular meningitis. On the contrary, they are rarely present upon its convexity or lateral aspects, while they are uniformly present at the base, and especially about the optic chiasm and the fissures of Sylvius.

Upon *microscopic examination* of one of these granulations, its issue is seen to be composed of numerous oval cells, with a single nucleus, though there are also some larger cells mixed with these which contain several nuclei. In many instances, as has been observed by Cornil,¹ Hayem,² Bastian,³ and ourselves,⁴ the tuberculous granulation will be seen to envelop a small arteriole, whose calibre is obstructed at the point of its development. There is also marked proliferation of the cells of the perivascular sheath of the vessel for a varying distance on either side of the granulation, and it is highly probable that it is from these cells that the granulation has been developed.

We think it probable that some of the granulations may also be devel-

¹ Arch. de Phys. Norm. et Path., 1868, p. 98.

² Études sur les Diverses Formes d'Encephalite, Paris, 1869.

³ Edin. Medical Journal, 1867, p. 875.

⁴ Trans. of Biological and Micros. Section of Acad. of Nat. Sci. of Phila., 1869.

oped from the cells of the connective tissue which holds together the vessels of the pia mater.

These miliary tubercles precede the occurrence of the inflammatory changes in the meninges described below, and sometimes it happens, in very acute cases, that the only lesions discoverable consist of a few gray granulations scattered in the meshes of the pia mater. It is not probable, however, that they exist any great length of time without giving rise to meningitis, since they are usually found associated with more or less abundant inflammatory exudation, which surrounds and often conceals them. The chief seat of this inflammation, as of the tubercular deposition, is the pia mater; the arachnoid membrane being, as a general rule, affected only to a slight extent. That membrane sometimes, however, contains a very small quantity of clear or turbid serum in its cavity. Its surface is often dry and viscid, and in some instances its whole tissue is opaque and thickened. But it is chiefly in the pia mater that are found the evidences of severe inflammation. In order to detect these changes, it is necessary to examine the membrane not merely upon the surface of the brain, but to tear it off, so as to bring into view the portions which dip in between the convolutions, and which often exhibit the greatest amount of morbid alteration. The inflammatory lesions vary between mere vascular injection, infiltration with clear, turbid, or gelatinous liquid, and abundant formation of lymph. When the inflammation has gone beyond mere sanguine injection, it is marked by infiltration of the membrane with turbid, whitish, or sanguinolent serum, with pus, or with whitish or yellowish lymph. These products are, like the tubercular granulations which they imbed and often conceal, most abundant at the base of the brain, about the peduncles of the cerebrum, the optic chiasm, and in the fissures of Sylvius; like these granulations also, the products of the inflammation are most marked along the track of the bloodvessels. In this respect the disease differs from simple meningitis, in which the results of inflammation are usually more abundant and well marked upon the convexity than at the base. The pia mater, which, in a healthy brain, can be readily detached from the surface of that organ, becomes, in cases of meningitis, particularly in those which are violent, more or less adherent, so that in tearing it off portions of the cineritious substance, which is itself softened, come with it. The proper tissue of the membrane is thickened and indurated, the degree of thickening depending on the amount of infiltration.

After the changes in the pia mater, the most important anatomical feature is effusion within the ventricles. This was formerly thought to be the essential lesion of the disease, but recent researches have shown that it is absent in some instances which have followed in all respects the ordinary course of the malady. According to M. Barrier, effusion cannot be supposed to exist unless the ventricles contain from one and a half to two ounces of fluid, whilst Rilliet and Barthez assert that the normal quantity is a few grammes (about a drachm). The quantity in this disease is very variable; sometimes there are only a few drops or a teaspoonful, while in other instances it amounts to three ounces and a half, or much more. It may be so large as greatly to distend the ventricles, rupture the soft com-

missure of the thalami, and even the septum lucidum, diminish considerably the thickness of the hemispheres, and flatten the convolutions against each other. In such cases the effused fluid passes through the membrane of the ventricle and infiltrates into and softens the substance of the brain, so that the latter becomes almost of the consistence of thick cream. The characters of the fluid vary in different cases. It is white, perfectly limpid and transparent, or may be turbid, either from being secreted in that condition or from holding in suspension albuminous or purulent flocculi, or portions of the broken-down walls of the cavity. In some rare instances it is sero-sanguinolent. Rilliet and Barthez remark that the effusion which coincides with tubercular meningitis is different from that which accompanies tubercles of the substance of the brain. In the former it takes place rapidly, is turbid, exists in smaller quantity, and constitutes the condition formerly called acute hydrocephalus. In the latter it is secreted slowly and in considerable quantity, dilates the walls of the cranium, and constitutes one form of chronic hydrocephalus.

The brain itself presents various morbid alterations. The whole organ often seems enlarged, so that the dura mater appears distended, and when the latter is cut into, the cerebral substance protrudes in the form of a hernia. At the same time the convolutions are observed to be pressed against each other, and the anfractuositities seem to have disappeared. The compression of the brain depends either upon the distending action of the ventricular effusion, or upon sanguine turgescence of the organ. In most cases, but not in all, there is evident congestion of the cerebral substance, shown by a more or less abundant dotted redness, and sometimes by a general rosy tint of the medullary, and vivid redness of the cortical portion. Softening of the substance of the brain is of common occurrence in connection with the other lesions. We have already spoken of the softening of the walls of the ventricles where there is much effusion, and which in some cases appears to result from the macerating influence of the fluid. In many other cases, however, microscopic examination of the softened brain-tissues shows the effects of inflammation in the presence of numerous granule-cells, free granular matter, and a disintegrated condition of the nerve fibrils. In addition to this, as figured by Rindfleisch (*Syd. Soc. edit.*, vol. ii, p. 312), the proper vessels of the cortical substance of the brain frequently present tubercular degeneration of their walls. The lining membrane of the ventricles also presents abnormal appearances in a majority of cases. In some these consist merely in injection with loss of polish and transparency: in others, however, by viewing the surface sideways, we can detect a very finely granular condition, as though the membrane had been sprinkled with fine sand. Löschner (*Aus dem Franz Joseph Kinder-spitale*, 1860, Prague), has found this appearance to be due to a proliferation of the cells of the ependyma, the minute granulations consisting of rounded nucleated cells. In Dr. West's minute analysis of 61 autopsies of tubercular meningitis, also, the lining membrane of the ventricles presented evidences of inflammation in a large proportion of the cases. We have also referred, very cursorily, to the softening which exists under the

inflamed portions of the membranes, and which occasions adhesion of the pia mater to the brain beneath. In the latter cases the softening may be either red or white, and does not penetrate more than a line, and often less, in depth.

In addition to the changes already described tubercles of the brain itself may be occasionally met with, having no connection with the meninges. These are found in various parts of the organ, and differ greatly in size, varying generally between that of a millet-seed and hazel-nut, but reaching sometimes the volume of a pigeon's or hen's egg, or even that of half the fist.

We have but few words to say in regard to the lesions of other organs. It is undoubtedly true that in the vast majority of cases tubercles are found in other parts of the body. Of all the cases of tubercular disease observed by Rilliet and Barthez, amounting to 312, in only one was the deposit confined to the meninges (*op. cit.*, 1ère édit., t. iii, note, p. 49). M. Valleix (*op. cit.*, t. ix, pp. 196, 197), states, that in all the cases, without exception, of tuberculosis of the meninges in adults, tubercles exist also in the lungs, and that the same is true, in the vast majority of cases, in regard to children. According to Henoch (*Centralblatt Zeit. f. Kinderk.*, May 1, 1879), limitation of the eruption of tubercle to the pia mater or the brain-substance, to the exclusion of other organs, is very rare in childhood, and when it is reported, give rise to suspicion that the examination has not been thoroughly made. The organs in which the deposit is most apt to exist are the bronchial glands, lungs, mesenteric glands, pleura, spinal cord, and peritoneum.

Another very frequent lesion is softening of the stomach. This may affect only the mucous or all the coats, so that a slight degree of force will suffice to tear the organ. Dr. Gerhard (*Am. Jour. Med. Sci.*, vol. xiv, 1834) states, that lesions of the stomach existed in six of the ten cases detailed by him, and in four-fifths of others not detailed.

Before quitting this subject, we would call the attention of the reader to the fact mentioned by M. Valleix (*op. cit.*, t. ix, p. 214) that all the symptoms about to be described as constituting the disease under consideration, with the exception of paralysis, may depend on simple tuberculosis of the meninges. Several cases have been cited, in fact, in which the only lesion found after death consisted of granulations in the pia mater. No traces of inflammation were observed. It is clear, therefore, that the evidences of the disease, or symptoms, depend not merely on inflammation caused by the tubercular deposits, but on the presence of that morbid production. The paralysis, which is one of the important symptoms, depends partly upon the inflammatory changes in the brain-tissue itself, and partly upon the pressure exerted on the structures at the base of the brain by the exudation which forms there.

SYMPTOMS; COURSE; DURATION.—The disease has been divided by authors into different stages, founded on the predominance of certain symptoms at particular periods of its course.

These divisions are all imperfect and unsatisfactory, because the disease

is in fact a continuous one, and for this reason some writers have avoided attempting any classification of the symptoms. We can, however, obtain a more faithful picture of the disorder by adopting the division made by M. Valleix, which, though arbitrary and imperfect, because of the want of a natural line of demarcation, seems warranted by the very great differences in the character of the symptoms at an early and late period of the affection. We shall therefore describe first the invasion of the malady, and then two stages or periods of the symptoms after the disease is confirmed.

The *invasion* of the disease may be either insidious or sudden. In a large majority of the cases, the onset is preceded by a well marked prodromic period. The length of this period varies greatly in different subjects. Its duration is stated by MM. Rilliet and Barthez to be, as a general rule, between fifteen days and three months, scarcely ever less, and rarely more. During this period, the symptoms presented by the child are those which are usually held to be indicative of a failure in the general health. The nutritive functions especially show disorder. The appetite diminishes, or becomes capricious, there are alternations of constipation and diarrhœa, the body grows thin, the color pales, the gayety of childhood disappears, and the patient becomes listless, apathetic, and complains of being tired and weak; or he is irritable and peevish, or too mild and gentle; study and exercise both become distasteful, and there is a degree of weakness and debility, which, though slight at first, becomes at length so evident as to arouse the attention of the parents, or those who have charge of the child. If, as not rarely happens, there is a development of miliary tubercles in the lungs also, there may be troublesome, dry, spasmodic cough before the appearance of cerebral symptoms. Besides these symptoms, there is often very great restlessness at night. The only pain complained of is headache, and sometimes abdominal pain. The headache is, in subjects old enough to notice and describe their sensations, often a prominent symptom. It is not constant, but occurs at intervals, and is sometimes severe, and its returns frequent. Fever is not generally present until after the more positive symptoms have fairly begun, and when present is generally slight and fugacious. The emaciation and loss of strength are seldom present to such a degree, in the prodromic stage, as to confine the child to the house. On the contrary, he continues to amuse himself at times, and to walk as usual.

The following is a rapid and summary account of the mode of invasion in some of the cases that have come under our own observation:

In one case, in a girl six years of age, the invasion was preceded during three months by occasional cough, and irregular attacks of fever, by progressive emaciation, paleness, languor alternating with extreme irritability, disinclination to take exercise, and during the latter part of the time by partial lameness, and in fact by all the signs of general tubercular disease. In another, which occurred in a boy eight years of age, it was preceded for several months by frequent complaints of intense headache, especially after taking active exercise, and by unusual languor, but no other symp-

toms. The boy was sent to a boarding-school apparently well, and was suddenly attacked there. In five cases the meningeal symptoms were developed in connection with those of phthisis, whilst in an eighth they followed a state of general weak health, with dyspeptic symptoms, which had lasted for several months. In a ninth case, a violent convulsion, seemingly dependent on a fit of indigestion, was followed during four months by irregular and diminished appetite, by some loss of strength and flesh, and by frequent attacks of severe headache, and at the end of that time by the symptoms which denote inflammation of the membranes. In a tenth, after some months of gradual thinning and general debility, a convulsion occurred, also from indigestion apparently. This was recovered from, but a few days afterwards the symptoms of meningitis showed themselves, and followed their usual course. In an eleventh case, occurring in a girl ten years of age, there was a mild, almost continuous fever, lasting four weeks, and resembling most closely typhoid fever, except that there was no diarrhœa and only a very few doubtful rose spots, when severe frontal headache, vomiting, slow and intermittent pulse, with drowsiness, declared the invasion of tubercular meningitis. In a twelfth, a girl three years old, born of healthy living parents, presented for four days the signs of gastric catarrh, with, however, unusual irritability of temper alternating with a suspicious quiet. On the fifth day, there was just preceptible strabismus, after which the case went on in the usual way to a fatal result. In a thirteenth, in a girl five years old, of healthy living parents, but with tuberculous grandparents on the father's side, the general health failed slowly, with loss of appetite and flesh for one month. Then there set in lassitude, desire to lie about, with the most petulant irritability on the slightest disturbance, occasional vomiting, constipation, loathing of food, and gradual conversion of drowsiness into coma, and so on to the end. In a fourteenth, a case of general miliary tuberculosis, to which allusion has already been made in the article on Hooping-cough, there was a hard, spasmodic cough for some weeks, possibly connected with enlargement of the bronchial glands, which were subsequently found to be tuberculous, and succeeded by an irregular febrile state simulating typhoid fever, and lasting some days before the appearance of cerebral symptoms. In the remaining cases that we have seen, the invasion was preceded by much less decided prodromic symptoms.

After the different phenomena above described as characteristic of the prodromic stage have continued during a variable length of time, the disease enters into activity, a change which is ushered in by three important symptoms: *headache*, *vomiting*, and *constipation*, to which is added, in a large majority of the cases, slight *acceleration* of the circulation. At the same time the *intelligence* remains perfect, the *strength* is not greatly diminished, the *appetite* is not entirely lost, and the *thirst* is moderate.

First Stage.—The headache, vomiting, and constipation persist and become more marked. *Headache* is a nearly invariable symptom in children old enough to describe their sensations, and is therefore very important. In infants its presence is to be inferred when the child carries its hands

frequently to various parts of the head, and presses strongly against it, and when the head is constantly rolled from side to side. It is generally frontal, and is usually referred to a point just over one or both brows. In other cases it extends over the whole head. It is commonly severe, so that the child when old enough complains of it spontaneously. In the case of a girl seven years old, whom we saw, it was so severe that she cried frequently and bitterly, begged to have the doctor sent for, and submitted willingly to any remedy suggested with a view to its relief. It is thought that the acute, shrill cry of the disease, to which the term *hydrencephalic* has been applied, depends on the acuteness of this pain. It usually lasts throughout the first stage, and ceases only as the delirium and coma of the second stage come on. *Vomiting* is also a nearly constant symptom. Of 80 cases collected from different sources by M. Barrier, it was absent only in 15, or less than one-fifth. This symptom generally makes its appearance on the first day, rarely later than the second or third, and lasts two or three days, and sometimes longer. In one case that we saw, it lasted eleven days, though it was but slight on the tenth and eleventh. The matters ejected from the stomach consist of the ingesta, and of mucus and bile in various proportions. It is commonly repeated two or three times a day. *Constipation* is even more important as a symptom than the one last named. Of 87 cases it was absent only in 7, according to Barrier. MM. Rilliet and Barthez state, however, that it exists at the beginning only in about three-fourths of the cases. Where there is diarrhœa instead of constipation, at the invasion, as sometimes happens, the former symptom almost always depends on tubercular disease of the intestine. Even under these circumstances, however, the diarrhœa is sometimes arrested, and constipation substituted under the influence of the cerebral disease. The constipation generally persists obstinately for several days, and then gives way under the influence of purgative medication, or is replaced spontaneously by diarrhœa with involuntary stools towards the termination of the case.

In connection with the three important symptoms just described, there are others, which, though less characteristic, are of much assistance in forming the diagnosis. The child is *dull* and *sad*, or *excited* and *irritable* by turns; he shuns the *light*, or closes the eyelids and contracts the brows when it is thrown upon the face; his hearing becomes painfully acute, so that sudden and jarring sounds distress and irritate him; the *sleep* is restless and disturbed, and accompanied by grinding of the teeth; and he utters from time to time, both sleeping and waking, the peculiar shrill, sharp, and sudden scream, which seems to depend upon internal pain, probably headache, and which has been called by Coindet the *hydrencephalic cry*. In young children, those who have not yet learned to put their sensations into words, a peculiar, apparently causeless, obstinate peevishness and positive ill-temper, shown by sudden, sharp crying at any disturbance, as even the kindness of a father or mother, especially when this alternates with sluggishness or drowsiness, and when there is no evident disease of a painful or exhausting kind to explain such a state, ought to arouse the fears of the physician as to the possible inception of this disease,

even when there is as yet no vomiting or distinct signs of headache to call attention to the brain. The general as well as the special sensibility is sometimes but not by any means always, exaggerated at this time. Rilliet and Barthez met with exaltation of this function only in four of their patients. The *intellectual faculties* remain undisturbed in the majority of the cases during the first few days, and this fact, which is so contrary to what might be expected, is one of the utmost importance in the judgment of the case. We remember being asked by a little girl seven years old, to whom we have already referred, "why it was that she saw double; why she saw two mothers and two doctors?" At the time when she first asked the question there was no perceptible strabismus, but on the following day we thought we could detect a deviation of one of the eyes from its proper axis, and on the third day the deviation was very marked, though the poor child still wondered why she saw two objects instead of one. In another case in a boy five years old, there was no disorder of the intelligence until the eleventh day, when there was slight delirium alternating with somnolence; yet it was clear from the first that the attack would prove one of tubercular meningitis, from the co-existence of violent frontal headache, obstinate vomiting, constipation, slow and irregular pulse, and the absence of other local or general symptoms. In only a fifth of the cases observed by MM. Rilliet and Barthez was there perversion of the intellectual faculties at the invasion. Let us observe, moreover, that even when children present some of these disorders early in the attack, they generally consist only of slight delirium, dulness of the intelligence, slowness and hesitation in answering questions, disposition to somnolence, excessive irritability and peevishness of temper, and what is more important and characteristic than any of these, perhaps, of a certain expression of the countenance, and particularly of the look, which is expressive of astonishment or of the utmost indifference. The look is, in fact, fixed or staring, like that of one in a mild ecstasy. Even when these symptoms exist, however, at an early period, they not unfrequently alternate with the most perfect clearness of the faculties, so that the physician in private practice, who sees his patient only at long intervals, and for a few moments at a time, should never venture to disbelieve, without due consideration, the account of the mother or nurse as to their occasional presence during his absence, even though never observable during his visit. We knew this to happen in regard to two boys of eight and ten years of age respectively, whose mothers constantly insisted to the physician in attendance that during his absence the children occasionally presented slight delirium, and a wild uncertain expression of the countenance, which made them fear that the brain might be affected. As the children's intelligence was perfect, however, whenever the doctor saw them, he determined that the mothers were fanciful through over-anxiety, and ascribed the sickness to a bilious disorder of the stomach. After a few days the cases developed themselves, and the boys died with every symptom of tubercular disease of the brain.

When disorders of intelligence do not occur in the early days of the attack, they usually make their appearance about or soon after the fifth day.

In this disease, as in acute simple meningitis, the obstruction to the return of venous blood through the sinuses produces in both eyes, but especially in the one corresponding to the hemisphere where the inflammation is most intense, congestion and œdema of the optic papilla and surrounding tissue (Bouchut's peripapillary congestion); tortuosities and varicosities of the retinal veins; and occasionally thrombosis or rupture of these vessels, causing minute hemorrhages in the retina. In some cases the size of the globe is increased, owing to hydrophthalmia. In addition to this, the characteristic appearances of optic neuritis often become visible; thus in a series of observations by Dr. Garlick (*Med. Chir. Trans.*, vol. lxij, p. 441) the ophthalmoscope disclosed changes in the optic disks of about 80 per cent. of the children who died of tubercular meningitis. These lesions are indeed more frequent in this disease than in simple meningitis, since the inflammation and resulting exudation are more apt here to involve the base of the brain, and cause a greater degree of obstruction to the circulation.

Enlarged experience has convinced us of the high value of these retinal lesions in establishing the differential diagnosis of tubercular meningitis. It is true that they cannot be regarded as pathognomonic of the existence of this disease, and are therefore only valuable as confirmatory of the general symptoms, still in certain cases the development of the ocular lesions before the appearance of the more characteristic symptoms enable the diagnosis to be made at an earlier date than would be possible without ophthalmoscopic examination; while in others where the general symptoms leave it for the time doubtful whether the case is one of typhoid fever or of tubercular meningitis, the use of the ophthalmoscope renders invaluable aid. As the disks vary physiologically in different individuals, and even in the same person the two are often not alike, progressive change is better evidence, therefore, than can be obtained from a single observation.

During the first stage the *coloration* of the face ought to be noticed. It is generally paler than natural, though from time to time a sudden flush of redness may be seen to pass over it. The condition of the *senses* is natural, except that the acuteness of the eye, ear, and sometimes that of touch, are exalted, so that the child avoids the light, starts at sudden or loud sounds, and cries when it is touched or moved. The *respiration* becomes unequal and irregular, and is interrupted by sighing or yawning.

Convulsions rarely occur in the first stage. MM. Rilliet and Barthez conclude that meningitis without complication of tuberculous disease of the cerebral substance, never begins with convulsions. In one of the cases that came under our charge, a severe and prolonged convulsive seizure did occur, however, on the very first day of the attack of the disease. The subject of the case was a boy between four and five years old. The death took place on the eighteenth day, and the autopsy showed no tubercular disease of the cerebral substance. It is proper to state, however, that the child had eaten on the morning of the day that he was attacked, a most unwholesome meal, and it is very possible, as we in fact supposed at the time, that the convulsions were caused by the presence in the stomach of undigested food. When they do occur in tubercular meningitis, they may

be limited to the extremities, upper lip, eyeballs, or they may be general. Sometimes the child dies in a convulsion. They are generally much less important as a symptom, according to M. Valleix, than in simple acute meningitis.

The *tongue* remains moist; the *appetite* is not entirely lost; *thirst* is moderate; the *constipation* continues, unless removed by treatment; the abdomen becomes *retracted*, so that its walls approach closely to the spinal column, and allow us to feel the pulsations of the aorta without using more than very slight pressure. The latter symptom comes on gradually, and is generally well marked by the sixth day or a little later. MM. Rilliet and Barthez regard it as a very important sign, and state that they have observed it almost exclusively in cerebral affections. They think it depends not upon contraction of the abdominal muscles, but upon retraction of the intestines. We can corroborate by our own experience, the evidence of the above authorities as to the value of this symptom. It has been very marked in most of the cases that we have seen.

The state of the *circulation* is of the utmost importance in forming the diagnosis. So true indeed is this, that Dr. Whytt, of Edinburgh, whose description of acute hydrocephalus, published in 1768, has been most highly commended by all recent writers as a singular instance of accurate observation, makes three stages of the disease, each of which is characterized by the state of the pulse. In the early part of the attack the pulse is accelerated, rising to 110, 120, or, according to Whytt, in a few cases to 130 or even 140. At the same time it is neither full nor tense, as a general rule, but rather soft and compressible. This condition of the pulse changes, as we shall find, in the middle period of the disease, and again shortly before the fatal termination. The *heat* of the skin is usually moderate and sometimes quite natural, at this time, as might be supposed from the state of the circulation. Frequently the temperature will not during this period exceed 100° to 100.5° , though occasionally marked and rapid changes in it are observed. It is especially to be noted that the temperature does not follow the regular mode of development so characteristic of typhoid fever.

Second Stage.—This stage begins about the time the more marked nervous symptoms show themselves. The headache generally subsides or ceases at the beginning of this period and gives place to *delirium*. This occurs usually somewhere between the sixth and twelfth days. The delirium which occurs has been generally supposed to be always mild and calm. MM. Rilliet and Barthez state, however, that in one-third of their cases it was intense, and accompanied with cries, agitation, and frequent changes of position. In most of the cases, however, it is mild, and is manifested in older children by their muttering unintelligible words, by inattention to what is going on around them, by an expression of wildness and astonishment, and by their giving hesitating answers to questions. In children under two years of age there is no proper delirium. There is, however, an analogous condition, which is characterized by disorder of the two faculties of attention and perception. The delirium seldom lasts more than two or three days, and generally alternates with somnolence, so

that the child is either dozing and sleeping, talking in its sleep, or frequently waking with loud cries, and restlessness. The *general sensibility*, which may have been exaggerated in the early period of the disease, is diminished in the early part of the second stage, or about the seventh day, and completely abolished towards the end. The face in the second stage is almost always pale, or pale and flushed alternately. During this stage, and especially during the latter part of it, it is very common to see sudden alterations in the color of the face. Sometimes without any apparent cause, but more frequently from disturbances of any kind, as from pain, or from external influences acting upon the child, such as moving it, or the administration of food or medicine, the face becomes suffused of a more or less deep pinkish or scarlet tint, the color beginning faintly at first and gradually deepening and expanding until it covers the whole face and forehead, and then as gradually fading away. It is during this stage also that another symptom, which we have often noticed, and to which M. Trousseau has called attention, may usually be observed. M. Trousseau refers to it as a red line or spot remaining upon the skin of the forehead or abdomen when the finger has been drawn across it, and has given to it the name of "*tache méningitique*" or "*tache cérébrale*." We had often remarked, before knowing that M. Trousseau had drawn attention to this phenomenon, that the slightest pressure with the finger on any part of the face or forehead, caused the appearance at the point of pressure of a spot of a peculiar pink or rose color, which, like the flush above referred to began faintly, became more or less deep in tint, remained a few moments, and then as gradually faded away. This symptom is undoubtedly due to the extreme modification of the innervation of the minute bloodvessels of the skin. There is no doubt that it is nearly always present in cases of tubercular meningitis, and thus may be said to possess a diagnostic importance. We have, however, so frequently met with the same phenomenon in cases of typhoid fever, and more rarely in cerebral pneumonia, that we must warn against it being regarded as a pathognomonic symptom of tubercular meningitis. Occasionally contractions pass over the features, giving rise to grimaces, after which the countenance resumes its expression of indifference and stupor. The eyelids are generally only partially closed, and between them the globes of the eyes can be seen to oscillate and move in various directions, as though by some automatic force.

As the case progresses, the *nervous symptoms* become more and more marked; somnolence gradually deepens into coma; the delirium becomes less and less frequent; and the child no longer observes what is going on, nor answers questions. As the somnolence and coma increase, various lesions of motility make their appearance, consisting, in order of frequency, of paralysis, which is generally partial; contraction with rigidity of the limbs; stiffness of the muscles of the back of the neck, causing retraction of the head; stiffness of the trunk; spasmodic closure of the jaws; carphologia; subsultus tendinum, and convulsions. The paralysis is almost always partial and of very limited extent, affecting, for instance, the jaw, the orbicularis muscles of the eyelids, the levator of the upper eyelid, the tongue,

or one side of the face. It is very rare to see one of the limbs paralyzed. Contraction with rigidity of the muscles is an important symptom, but is not always present. When it exists it generally appears at an advanced period of the attack, commonly between the seventh and thirteenth days and is usually partial. It may affect either the extremities, back of the neck, trunk, or inferior maxilla. It is seldom permanent, but after lasting one or two days, disappears, to reappear at a later period. The carphologia, subsultus, and chewing motion of the under jaw generally occur only a few days before death, and last but a few days.

The *decubitus*, in the early part of the second stage, is generally lateral, with the thighs flexed upon the pelvis, the legs upon the thighs, the arms applied against the thorax, the elbows bent, and the hands placed in front, the *decubitus* called by the French "*en chien de fusil*," or *gun-hammer*. At this time the child will still occasionally move its position with facility, showing that strength is not by any means entirely lost. At a still later period the *decubitus* is dorsal. In the latter part of the first and early part of the second stage, the *pulse*, which we have ascertained to be accelerated at the invasion, falls to the natural standard, or becomes slow, and at the same time irregular. From 110 or 120, as it was, it now sinks to 90, 80, 60, or as happened in one instance to M. Guersant, to 48 in the minute. Coincidentally with this change it almost always becomes irregular. The irregularity affects both its force and frequency, so that a strong pulsation may be followed by a feeble one, or the rhythm may be regularly or irregularly intermittent. The irregularity varies greatly at different periods of the day, or within short spaces of time, so that the pulse is found to be very slow at one moment and much more frequent the next. On this account it is necessary to examine it on different occasions. Slowness and irregularity of the circulation are important as a means of diagnosis, since it has very rarely been met with as a permanent condition, except in the tuberculo-inflammatory affections of the brain and its appendages. Towards the termination of the disease, generally speaking two or three days before death, the pulse rises again in frequency, so that it counts at first 112 or 120, and gradually increases to 140, 160, or even 200 the day before, or that on which death takes place. Simultaneously with this change it also becomes extremely feeble and small, and often ceases to be perceptible at the wrist on the last day.

The *respiratory movements* also show marked irregularities. During the early stage of the disease they are frequent, though we think rarely so much as to preserve the normal ratio to the pulse. But during the stage we are now considering, the breathing becomes unequal and irregular, and deep sighing respirations alternate with quick superficial ones. We have called attention lately¹ to a very peculiar modification of respiration, which has been frequently observed by ourselves and others in the exudative stage of tubercular meningitis. The following description may serve to give an idea of its general character: "The breathing is from time

¹ Remarks on Cheyné-Stokes Respiration, especially in connection with Tubercular Meningitis, by Dr. William Pepper, Phila. Med. Times, May 27th, 1876.

to time interrupted by periods of apœna of varying length (five to thirty seconds), between which occur a series of respiratory acts, which begin by very feeble and barely perceptible movements, and gradually grow fuller and stronger until they reach a climax, when they occasionally end by a long-drawn sigh, or more commonly, pass through a descending scale of movements, each growing more and more feeble until they end with barely perceptible respirations, such as marked the beginning of the series. This period of respiration, which also occupies from five to thirty seconds, is followed by a second period of complete apnœa, which is in turn succeeded by a group of respirations similar to the first." (*loc. cit.*) This disturbance of breathing, which is known as Cheyne-Stokes or tidal respiration, depends, as does also the interference with the heart's action, upon the pressure of the exudation upon the pneumogastric nerves; and, as they are not found in the affections which may simulate tubercular meningitis, must be regarded as possessing considerable diagnostic value.

The *heat* of skin, which has fallen with the reduction in the frequency of the pulse, generally increases with its acceleration. This is not invariable, however, since in some cases the temperature remains but moderately elevated, about 101° or 102° , until death; and in others an algid condition precedes death, in which the temperature falls as low as 79.4° . (*Reynolds's Syst. of Medicine*, vol. ii, p. 379, art. Tuberc. Meningitis). On the other hand, in some cases the temperature increases irregularly as the fatal result approaches, and may attain an extreme height. Thus Roger (*op. cit.*, p. 323) has observed on the day of death in an attack of tubercular meningitis, a temperature of 108.5° F; and Satterthwaite (*New York Medical Record*, May 8th, 1880), reports a temperature of 110° a few minutes before death in a case of this disease. During the last few days the surface is often covered with an abundant perspiration; the tongue becomes dry; the teeth and gums are fuliginous; the exhaustion increases; the respiration becomes stertorous, unequal, difficult, and anxious, and at the very last attended with great dyspnœa; and the urine and stools are discharged involuntarily. Death finally occurs in this condition, or is hastened by an attack of convulsions. In some cases it is most lingering. In one instance we expected the death of a young child in this disease every day for eight in succession.

The *duration* of tubercular meningitis is exceedingly variable in different cases. As a general rule it lasts between eleven and twenty days, though it may continue a considerably longer time. Rilliet and Barthez have never known death to occur before the seventh day.

DIAGNOSIS.—The disease with which tuberculosis of the meninges is most likely to be confounded are simple meningitis and typhoid fever. It might also be confounded, though this is much less probable, with the cerebral symptoms which complicate the exanthemata and some local diseases, especially pneumonia, and to which symptoms, as a group, M. Barrier has applied the term pseudo-meningitis.

The diagnosis between tubercular and simple meningitis will be best understood from the following synoptical table, based upon the one contained in the last edition of the work of MM. Rilliet and Barthez.

SIMPLE ACUTE MENINGITIS.

I. The subjects of acute simple meningitis are usually robust and well-developed, and present no trace of either internal or external tubercular disease. Born of healthy parents.

II. The disease may prevail epidemically.

III. *Condition Prior to Invasion.*—The disease begins in the midst of the most blooming health, or, if secondary, it occurs in the course of, during the convalescence from, some acute non-tubercular disease, or it follows an external cause.

IV. *Mode of Invasion.*—Violent convulsions attended with intense febrile movement, and with very hurried respiration in young infants; or very acute frontal headache, accompanied by fever, bilious vomiting, and towards the end of the first, or in the course of the second day, at the latest, excessive restlessness, preceded or not by somnolence; most violent delirium; formidable ataxia.

V. *Symptoms.*—*Very intense* headache, *obstinate* vomiting, *moderate* constipation, *violent* fever, *high* delirium.

VI. From the beginning, the aspect of a grave disease of ataxic form.

VII. Course rapid, aggravation progressive and continuous; convulsion after convulsion, or else violent delirium, extreme agitation, violent fever, etc.

Duration.—Disease of short duration, ending sometimes in 24 or 36 hours, but lasting generally from three to six days, and seldom more.

REGULAR TUBERCULAR MENINGITIS.

I. Subjects of tubercular meningitis delicate, puny, exhibiting often precocious intelligence and sensibility. Have sometimes had, in infancy, enlarged glands or chronic cutaneous eruptions; the parents, or brothers and sisters, often present the signs of tubercular disease.

II. Disease always sporadic.

III. *Condition Prior to Invasion.*—For some months or weeks the patients grow languid, lose their strength, become pale, emaciate; their temper changes, they are dull, they lose appetite, the digestion is deranged, etc. Absence of prodromic symptoms is rare.

IV. *Mode of Invasion.*—Never with convulsions at the onset; the change from the prodromic to the acute stage sometimes imperceptible. It takes place by a progressive increase of the symptoms before mentioned, and by the setting in of headache; in other cases, the acute stage is better marked by headache, vomiting, and constipation; generally, the intelligence remains clear; no ataxia. In the rare cases in which there is ataxia at the onset of the acute symptoms, the prodromic stage, above described, has been observable, or the meningitis has occurred in the course of advanced phthisis. In cases in which no prodromes exist, the meningitis begins with vomiting, constipation, moderate headache, and slight febrile movement; ataxia, if it is to appear, occurs later, and a mistake is impossible.

V. *Symptoms.*—Not *very intense* headache, vomiting *less frequent*, *very obstinate* constipation, *very moderate* fever, slowness and irregularity of the pulse, delirium usually mild.

VI. Invasion insidious, with the aspect of a mild disease.

VII. Course slow, preservation of the intelligence to an advanced period, fever slight, and some slowness and irregularity of the pulse, sighing, changing color of the face, eye dull or ecstatic, etc.

Duration.—Always much longer in the regular form.

We will remark in regard to this table, which is, in most respects, admirable, that we have never met with more intense and persistent headache than we have in some cases of the disease under consideration. In some of our cases this has been a most prominent and striking symptom.

Before quitting the subject of the diagnosis of these two affections, it is desirable to state for the information of the reader, that some of the highest authorities acknowledge it to be sometimes nearly or quite impossible to distinguish between them.

From typhoid fever, tubercular meningitis is to be distinguished by the antecedent history of the patient, which often reveals the existence of a tubercular diathesis in the latter affection; by the symptoms of the invasion, which in meningitis consist of severe and persistent headache, frequent vomiting, and constipation, whilst in typhoid fever the headache is less severe and less persistent, the vomiting much less frequent, and the constipation replaced by diarrhœa, or at least by an unusual susceptibility to the action of laxatives; by the different characters of the febrile movement, which, in typhoid fever, is more marked, and attended with a frequent, full, and regular pulse, while in meningitis it is less marked and is accompanied after a few days by slowness and irregularity of the pulse, and by irregularity of respiration; lastly, in meningitis, the constipation is usually marked, the abdomen is retracted, and there are various important and characteristic lesions of motility, and the special senses; in typhoid fever there is diarrhœa, the abdomen is distended and meteoric, there are characteristic rose-colored spots, whilst there are no considerable lesions, either of motility, or of the special senses. Much assistance in the diagnosis can also be obtained by a careful study of the course and changes of the temperature in the two diseases. In tubercular meningitis, instead of the gradual progress in development, with moderate evening exacerbations, which is so characteristic of typhoid fever, the temperature presents great and irregular variations; it is specially marked by a period of reduction, even to 97° or 96° (Roger), corresponding to the middle stage, and then by a final rise, which may continue increasing until the last day of life, or may be replaced by an algid state, with great lowering of the heat of the surface. In doubtful cases, the use of the ophthalmoscope will often be of great value. We have already mentioned the retinal changes which are frequently seen in tubercular meningitis, whilst in typhoid fever no lesions will be detected.

Although the above general remarks apply to the majority of cases, it must not be imagined, however, that the diagnosis between the two affections is always easy or even possible in the early stages. This is largely due to the fact that the typhoid fever of children presents as many irregularities and departures from the typical course which it more frequently follows in the adult. Thus it is not very rare to have epistaxis, diarrhœa, and even the peculiar eruption absent in typhoid fever in young children, and if, when this occurs, the *tache cérébrale* be developed by drawing the nail over the skin, it is evident that it will be difficult to decide whether the

irregular fever, with cerebral disturbance, be the result of the one or the other of these affections. The symptoms of most value in such obscure cases, are irregularity and slowing of the pulse, with unequal and irregular respirations; and strabismus, diplopia, or changes in the optic nerves or retina. When these appear, as they usually do in the second stage of tubercular meningitis, the diagnosis can scarcely possess any further difficulty.

It is unnecessary to do more than allude to the possibility of confounding the disease with the exanthemata, or with local diseases accompanied by cerebral symptoms, and particularly with pneumonia in very young children. The resemblance of pneumonia of the apex of the lung in the early stage of tubercular meningitis, has been referred to in the article on pneumonia. The diagnosis must be made by careful consideration of the symptoms peculiar to each, and in the case of a local disease, by accurate physical examination of all the important organs of the body.

Occasionally, also, cases are met with where, in connection with gastro-hepatic disturbance, there is probably some cerebral congestion, and which may simulate the early stage of tubercular meningitis. For instance, we were called to see a boy eight years old who had been suffering for two weeks with violent frontal headache, frequent vomiting, constipation, slight fever, and somnolence. We feared that the case might prove to be one of tubercular meningitis. However, a large dose of calomel, followed by castor oil and free leeching to the temples, relieved him in two days perfectly, and he has remained well ever since, though this was a number of years ago.

PROGNOSIS.—M. Barrier, in speaking of the prognosis of this affection, says: "The gravity of tubercular meningitis is not surpassed by that of any other disease. Thoracic and abdominal phthisis, though almost constantly fatal, pursue a slower course, and last a longer time. We may even allow as proved, that in a small number of cases, they are susceptible of cure, or may remain stationary for months or years. Unfortunately it is not so in regard to tubercular meningitis." MM. Rilliet and Barthez, in their second edition, do not express the same entire hopelessness as to recovery from the disease, that they did in their first. They say, amongst other conclusions (*op. cit.*, t. iii, p. 510), that there are on record incontestable examples of the complete disappearance of the symptoms, but remark, that such cures have occurred in the first stage, or in the first half of the second stage, after seven or eight days of sickness, rarely later, and after alternations of amelioration and aggravation. They state also that, in excessively rare instances, a return to health has been obtained even in the course of the third stage, after many weeks of illness. They are of opinion that the disease often returns and proves fatal in from one to five years and a half after the recovery. The cause of the relapse is to be found in the fact that the local lesion remains, and that the diathesis has not been eradicated. M. Valleix is of opinion that after having acquired the conviction that a case is really one of tuberculosis of the meninges, we should regard the patient as lost; "for the exception that I have men-

tioned (a case belonging to M. Rilliet, then unpublished), even did no doubt as to the exactness of the diagnosis remain, ought not, standing by itself, to impart to us any real security." M. Guersant (*Dict. de Méd.*, t. xix, p. 403), seems to think it possible that the disease may sometimes terminate favorably in the very early stage, but adds that "such cases are always more or less doubtful, and seem to us to belong rather, for the most part, to simple meningitis." During the second period (that of slowness and irregularity of the pulse), he has scarcely seen one child in a hundred survive, and even then they perished at a later period of the disease, or of phthisis pulmonalis. Of those arrived at the third stage (marked by renewed frequency of the pulse, coma, and lesions of motility and sensibility), he has never seen any recovery, even momentarily. Dr. George B. Wood (*Prac. of Med.*, vol. ii, p. 365), states that he has "never seen a well-marked case of tuberculous meningitis and favorably." Dr. Robert Whytt (*Works of Robert Whytt, published by his son*, quarto, Edinburgh, 1768, p. 745), says: "I freely own, that I have never been so lucky as to cure one patient who had those symptoms which with certainty denote this disease; and I suspect that those who imagine they have been more successful have mistaken another distemper for this."

In the quarterly abstract furnished by Dr. A. Wiltshire in the *Brit. and Foreign Med.-Chir. Rev.*, for April, 1876, at page 465, it is stated that Dr. Clifford Allbutt has known of two cases of recovery from tubercular meningitis, in which the diagnosis made with the aid of the ophthalmoscope was verified some years later at the autopsies of the patients. And at the same place is quoted another case by Dr. Rinteln (*Berl. Klin. Wochenschr.*, No. 21, 1876, p. 287), where recovery occurred after all the recognized symptoms of tubercular meningitis had been present, and where this diagnosis with a consequent fatal prognosis was made by all the physicians who saw the patient.

Our own experience coincides with the mass of evidence given above as to the almost hopeless fatality of the disease. All the undoubted cases that we have seen have proved fatal. A case, however, came under our observation, in 1850, which might, perhaps, be classed as a recovery from tuberculosis of the meninges, though not from tubercular meningitis, since there were no well-marked signs of inflammation of the membranes of the brain, though there was every reason to suppose that the symptoms depended on the deposit of tubercles in those membranes. The case was as follows:

A girl between four and five years old, whose mother was then laboring under tubercular disease of the apex of one lung (which has since proved fatal), and who had lost several brothers and sisters with consumption, had had nearly constant cough during the winter of 1849-50. During the months of April, May, and June, of 1850, she had exhibited all the signs of induration over the upper two or three inches of the right lung, before and behind,—marked dulness on percussion and bronchial respiration, but no râles. For these symptoms she had been treated with cod-liver oil, iodide of iron, opium for the cough, and good diet. From the middle of June she complained frequently of headache, had occasional vomiting without any gastric derangement, and was much disposed to be constipated. She had no appetite, grew thin, and was very languid, listless, and weak. On the 27th of June the mother thought she observed

some squinting. On the 29th we found that the child had lost all power over the right muscles of the right eye, so that when she looked towards the right hand, she squinted dreadfully. She was dull and heavy, and vomited two or three times a day. The pulse was 62 to 75 or 80; there was a slight hitch in its beat, but no decided intermittence. The child said that she sometimes saw two things instead of one. From this time until July 7th, she continued in much the same state. On July 1st, finding that the eyes were quite yellow, and that the child was constipated, we ordered half a grain of calomel morning and evening. After three doses she was purged. This relieved her a good deal, there being less headache, more appetite, and an improvement in color afterwards. But still there was every day some vomiting, complaints of headache, and more or less listlessness and heaviness in the morning, while in the afternoon she would brighten up and seem better. The intelligence continued perfect; the temper was rather irritable, but not very much so.

The treatment after the 29th of June was calomel, given as above stated, from time to time, to keep the bowels soluble; cod-liver oil, a teaspoonful twice or three times a day, as the child would take it; mustard foot-baths every day or two; and meat, bread, and ice cream for diet. On the 5th of July we ordered half a grain of iodide of potassium, three times a day, in addition to the oil.

On the 11th of July, she was taken, by our direction, to the seaside, where the use of the oil and of the iodide of potassium was to be continued.

On the 7th of August she was brought back from the seaside, and we saw her on the 8th. We were astonished to see how well she looked. The strabismus had entirely disappeared. We were told that it had begun to diminish two weeks after her arrival at the sea, and had then gradually disappeared. She had grown somewhat, though not very much, stouter. Her whole appearance was very much improved. The coloration of the body, the expression of the face, were both much better; she was much stronger, running about, in fact, all day; she ate well, and with the exception of a little cough, and a rather delicate frame, looked very well. Except one day, she was well all the time at the seashore. On that day she was feverish, had much headache and vomiting, and laid abed. The cod-liver oil and iodide of potassium were ordered to be continued.

The child remained pretty well throughout the winter of 1850-51. There was no return of either the strabismus or the vomiting. She was thin, pale and delicate-looking, coughed occasionally, and the solidification of the apex of the lung continued, but she was not confined to the house. Late in the winter she went south with her mother, and there, after having become quite stout and healthy during their travels, died of dysentery in April or May. The mother died in 1852, of phthisis, with large cavities in both lungs.

Another case, in which the early symptoms of the disease were well marked, and in which recovery took place, will be detailed in the remarks on prophylactic treatment.

In the following case, also, the diagnosis was of tubercular meningitis in the early stage:

The patient was a boy eight years of age, whose father had died a few years before of phthisis, the younger brother died of tuberculous meningitis, and the sister of whooping-cough, with the lungs filled with miliary tubercles, as ascertained by a post-mortem examination. The child, after having had fair health previously, was seized, towards the end of March, 1865, with frontal headache, very slight fever, occasional vomiting, constipation, hesitating pulse, languor, willingness to lie abed, and a tendency to somnolence. He was treated with rest, milk and beef tea in alternate doses, and mustard foot-baths morning and evening; the bowels were kept moderately open, and he took tincture of the chloride of iron in combination with dilute acetic acid and solution of

the acetate of ammonia every three hours. Under this treatment he improved, and in ten days had quite recovered. His mother removed from this city to Washington, where he died on the 30th of June of the same year, after an illness of twenty-one days, of what was called water on the brain.

In another case to which we were called in consultation, a boy whose mother had died a few years before of diabetes mellitus, and whose father's family was tuberculous, presented a series of symptoms which we could explain only as the result of slow thickening of the membranes at the base of the brain, in all probability the result of a tubercular deposit. This child had, for several weeks, violent frontal headache, constipation, loss of flesh, lassitude, a peculiar one-sided or lateral gait in walking, strabismus, and great impairment of vision, so that he could see a small object only by bringing it almost in contact with the face. There was scarcely any disturbance of the circulation, and only slight febrile heat at night. He was treated at first with rest, nutritious food, minute doses of bichloride of mercury in combination with iodide of potassium, three times a day, and then, when he began to improve, with tincture of the chloride of iron and cod-liver oil for a long period. He finally recovered his health, grew stout and strong, but has remained ever since so blind that he reads with great difficulty, but manages to pick his way through a room or the street, with only occasional stumbling. The illness occurred several years ago, and he is still living in good general health at this time.

Are we then to abandon all hope of deriving any good from medical means in the disease under consideration? To this most serious question we ought clearly to respond in the negative. The grounds for entertaining hope are first, the evidence of M. Guersant that he has seen cases which appeared to be tubercular meningitis recover in the first stage. Let it be supposed, even, that they were cases of simple inflammation. But they were undistinguishable from the tubercular disease by one of the most celebrated of modern physicians. Surely, therefore, it may happen to men of inferior skill to meet with the same difficulty, or, if we may so speak, to make the same mistake, if a mistake was made. It is said by M. Valleix that M. Rufz, after determining at the autopsy, that a case which he had witnessed was one of simple meningitis, asserted that it would have been impossible to distinguish it from the tubercular disease during life. Again, M. Rilliet has, according to M. Valleix, seen one case of recovery from what he believed to be the tubercular affection, and MM. Rilliet and Barthez, in their second edition, as above quoted, assert its occasional curability. To these authorities must be added the valuable evidence of Dr. Clifford Allbutt, already quoted; and it would be possible to cite still further instances, if it were necessary. We know of the occurrence of a case in this city, under the charge of one of our friends, than whom we believe no one can be more competent to make a correct diagnosis, in which, after the child had presented in regular order all the early symptoms of the disease, and had arrived at the last and most hopeless stage, perfect recovery, to his utter amazement, gradually took place. This child, when our friend last heard of it, three months afterwards, was in all respects strong and hearty. No doubt the probabilities are that the case was one

of simple meningitis, but who could have known this at the time?—and should it not deter us from abandoning all hope, and, as a consequence, all active treatment, when we seem to have under our hands a case of this dreadful malady? Our own cases, given above, also go to prove that the disease is sometimes curable in its early stages.

It is important, in tubercular meningitis, to avoid making a positive prognosis as to the period at which death will occur, notwithstanding that the patient may present every mark of an immediately fatal termination. We have already adverted slightly to this subject. On one occasion we expected the death of a patient with this malady for three days in succession, and on another, we visited a child for a week, during every day of which it seemed as though existence could not endure until the next. It had during this time profound coma, subsultus tendinum, and enlarged pupils; the eyelids were half open, the eyes constantly oscillating, or else rigidly distorted, and both corneas dimmed and slightly eroded, from constant exposure to air and light. Convulsions occurred from time to time, the pulse was variable, and at times exceedingly frequent, and indeed everything threatened a speedy termination. MM. Rilliet and Barthez say, "Often have we inscribed upon our notes *death imminent*, and been astonished the next day to find still alive, children to whom we had allowed scarcely two hours of life."

The symptoms which most positively indicate the near approach of death are, livid color of the face, sweats occurring about the face, glassy expression of the eye, dry and incrustated nostrils, and especially a very rapid pulse, and the various nervous symptoms mentioned, as carphologia, subsultus tendinum, and general convulsions.

TREATMENT.—In the early editions of this work we took the ground that it was proper in the early stage of the disease to employ bloodletting. Further experience and knowledge compel us to retract this opinion. We believe now, that abstraction of blood should not be resorted to unless when the diagnosis between this disease and simple meningitis is very uncertain. Where there are no marked signs of active inflammation, where, from the family history, from the absence of marked fever, and the peculiar state of the pulse, we have every reason to believe that the low-typed inflammation present is the result of the presence of tubercle, we deem it safest to avoid all lowering measures. The case is so critical, so almost hopeless from its very nature, that we prefer a treatment based on the theory of promoting a retrogression of the tuberculous deposit. The only measures which, in an experience of over thirty years, we have found to delay and, in the cases referred to in the article on prognosis, to cure the disease in part, have been the following: quiet of body and mind, obtained by means of rest in or on the bed, in a pleasant room, with attendants who know how to soothe and still the child. We always insist upon a nutritious diet; and one consisting mainly of milk and cream, or the two mixed, with beef-tea, bread and butter, if the patient will take it, or milk-toast, in moderate quantities, every three or four hours, a soft-boiled egg, or the yelk of a hard-boiled egg, once or twice a day, is what we usually endeavor to get the patient to take. A mustard foot-bath two or three

times a day, is always safe, and we think useful and tranquillizing. The bowels should be moved gently once a day, or every two days, by means of an enema or some simple laxative, as simple syrup of rhubarb. Active purgation we have found of no use. As remedies, we prefer the following :

R. Tr. Ferri Chloridi,	f ʒj.
Acid. Acet. Dil.,	f ʒj.
Liq. Ammon. Acetat.,	f ʒij.
Syrupi Simp.,	f ʒj.
Aquæ,	f ʒ ijss.—M.
A teaspoonful at five years of age, every three or four hours.	

In connection with this, we give half a teaspoonful of cod-liver oil in emulsion three times a day. Calomel we have abandoned of late years entirely, as it has utterly failed in our hands to do any good.

Iodine has been very much employed as a remedy in this disease, both in the forms of Lugol's solution and iodide of potassium. Perhaps the strongest argument which exists in its favor is the benefit which often follows its employment in other scrofulous and tuberculous diseases; through there are several cases in which it is asserted to have been successfully used in tubercular meningitis. Iodine itself is comparatively little used. M. Rilliet (*op. cit.*, t. iii, p. 308, 1847) states that it has entirely failed in his hands in the tubercular form of the disease; the only influence which it seemed to exert was to cause the immediate suspension of the coma. This was its effect also in a case in which we employed it, that of a girl seven years old, to whom we gave two drops of Lugol's solution three times a day, from the thirteenth to the twentieth day, when she died. The day before her death she seemed to improve somewhat, and we were in hopes that it had been of some service. The amelioration did not continue, however, and we are now disposed to believe that the change was one of those which often take place naturally in the disease.

Iodide of potassium was recommended more than twenty years ago by Roeser (*Hufeland's Journal*, April, 1840), as a remedy of special power in this disease. It has since then been very widely employed, and there are quite a number of cases in which it is asserted that its administration has been followed by successful results.

Dr. West (*op. cit.*, 4th Amer. ed., p. 97) thinks that he has seen good from its employment, "and that in one instance of what seemed to be advanced tubercular hydrocephalus, under the care of my friend and former colleague, Dr. Jenner, recovery took place under its employment."

Niemeyer (*op. cit.*, vol. ii, p. 218) speaks as follows of its use: "On the strength of two successful cases, opposed, it is true, by a large number of unsuccessful ones, I recommended large doses of iodide of potassium, continued for a long time."

Dr. J. Lewis Smith (*op. cit.*, p. 145) also recommends its use throughout the entire course of the disease, beginning as early as possible in the premonitory period.

Successful cases of its administration are also reported by Drs. Bourrose de Lafore, Coldstream (*Edin. Med. Jour.*, Dec. 1859), and Carson (*Med. Times and Gaz.*, March 5th, 1857).

We have ourselves frequently administered it, either alone or in combination with small doses of bichloride of mercury, but have not yet been fortunate enough to arrest the progress of any case when once the second stage has been fully developed. In a few cases, however, the use of the following combination :

R. Potass. Iodidi,	3j.
Hydrarg. Chloridi Corrosivi,	gr. j.
Syrupi Simp.,	f3j.
Aquæ,	f3ij.—M.
Dose, a teaspoonful three times a day at five years of age,	

has seemed to delay the march of the disease, in one some weeks, and in another, the one already mentioned, it seemed to have a positive effect in promoting the absorption of the exudation upon the membranes at the base of the brain. It is improbable, also, that in *all* of the reported cases, errors of diagnosis were made, and simple meningitis taken for the tubercular from ; so that there is no remedy from which so much benefit may be hoped for in this almost hopeless disease, as iodide of potassium in full doses, and it should therefore be faithfully tried whenever opportunity offers. We have been in the habit of giving it in doses of one or two grains every three or four hours, to children two years of age. It has, however, been given to the extent of a drachm in the course of a single day to children of that age. It ought to be begun with early in the case, and continued in connection with counter-irritation and cold to the head. We must remark, however, that it sometimes irritates the bowels too much, causing diarrhœa ; and here the dose ought to be greatly reduced, or the remedy withdrawn.

Bromide of potassium, on account of its undoubted power in cases of active cerebral congestion with great nervous excitement, may be advantageously associated with the iodide, and we are in the habit of combining from three to five grains of the former with the dose of the latter above recommended.

The treatment which has just been described is that which we have been led by our convictions as to the nature of the disease, and by our personal experience of different plans, to adopt as the most reasonable and the best. It is proper to state, however, that we have never seen it, nor any other method, of any avail after the disease has passed into the latter part of the second stage—when coma, dilatation of the pupils, marked strabismus, paralytic or convulsive phenomena, show the presence of inflammatory exudation under the membranes, and of serous effusion into the lateral ventricles, or the peculiar lesions of the substance of the brain which exist at that period of the malady. It is also proper to add that other means have been recommended by high authorities, and to these we shall now devote some remarks.

We have already stated that *calomel* has not succeeded in our hands, so that we have abandoned its use. We deem it right, however, to lay before the reader the opinions of others upon this point. Thus, it is highly recommended by many of the English writers on acute hydrocephalus, and is asserted to have effected cures when it has been pushed to such an extent as to produce salivation. But little dependence, however, can be placed on these assertions, as in all probability the reported recoveries occurred in cases of simple meningitis. The French writers speak of having used it in very large quantities without any success. It was given to many of the patients of MM. Rilliet and Barthez, in the quantity of from six to ten, increased to twenty grains, in twenty-four hours, in connection with frictions with mercurial ointment, of which two drachms and a half were used at first, and the quantity afterwards doubled and trebled. They state that salivation did not occur in any of the cases, though fetor of the breath and inflammation of the gums were of frequent occurrence. Calomel may be given, as has been remarked, in purgative doses, at the beginning, and for the purpose of procuring its specific effects. With the latter view the dose may be from a quarter of a grain to a grain, every hour or two hours. Mercurial inunction, in conjunction with the internal administration of the remedy, has been highly recommended by several writers as an efficient means of procuring the full effect of the drug upon the constitution. About a drachm of the ointment is to be rubbed into the insides of the arms and thighs morning and evening, and the quantity gradually increased if no effect is produced. For our part, we will merely state that we have never known calomel given in large quantities, in order to procure salivation, of the least benefit in the disease. On the contrary, we cannot but think that the violent irritation of the digestive mucous membrane which it has determined, whenever we have used it largely, and the inflamed, irritated condition of the mouth which it caused in one case, must have been a serious aggravation of the state of disease under which the constitution was laboring. Mercury is well known to be an injurious and dangerous remedy in the tubercular diseases of adults, having for its effect to increase the dyscrasia of the constitution, which already exists, and thereby to hasten the progress of the malady. Why it should have a different effect in children is difficult to understand. It may be said, to be sure, that in the disease we are considering, it is given to overcome the inflammatory element of the malady, which, for the time, constitutes the danger of the case, and also to allow the patient the chance of its beneficial operation should the disease happen to be one of simple meningitis. In support of the views just expressed, we will quote the opinion of Dr. John Abercrombie (*Diseases of the Brain and Spinal Cord*, Philad. ed., 1831, pp. 173-6): "Mercury has been strongly recommended in that class of cases which terminates by hydrocephalus, but its reputation seems to stand upon very doubtful grounds. In many cases, especially during the first or more active stage, the indiscriminate employment of mercury must be injurious. . . . In the preceding observations, I shall perhaps be considered as having attached too little importance to mercury

in the treatment of this class of diseases, particularly in the treatment of hydrocephalus; but in doing so, I have stated simply what is the result of an extensive observation, . . . and I confess, the result of my observations is, that when mercury is useful in affections of the brain, it is chiefly as a purgative."

It was recommended by Sir B. Brodie, to employ mercurial inunction as especially applicable in using mercury for children. He advised that a drachm or more of the ointment be spread upon one end of a flannel roller, which is to be applied, not very tightly, around the knee; repeating the application daily. "The motions of the child produce the necessary friction; and the cuticle being thin, the mercury easily enters the system." The editors of the journal in which this communication is made (*Braith. Retrospect. of Med.*, vol. iv, 1846, p. 147, from *Quart. Med. Rev.*, July, 1846, p. 169), state that they tried this plan in a case of acute hydrocephalus, in which some of the most urgent and fatal symptoms were present, "such as very dilated pupils, constant convulsions, hemiplegia, and more or less stertorous breathing; in short, so violent were the symptoms, that we considered the case perfectly hopeless; but on reflecting on Sir Benjamin's method, we ordered strong mercurial ointment to be smeared on each leg every twelve hours, and covered with a stocking made to tie tightly above the knees. The symptoms soon began to abate, and by following this up with small doses of iodide of potassium, frequently repeated (gr. i, every three or four hours), the head symptoms vanished."

"In a second case, the same set of symptoms were approaching, but were stopped by the same mode of treatment."

When the convulsive symptoms are violent and distressing, they may often be moderated by the use of a warm bath, which must be carefully given, and by the administration of some of the *antispasmodics*. We prefer for this purpose the fluid extract of valerian, of which from ten to twenty drops may be exhibited every two or three hours to young children, and a larger dose to those who are older. Bromide of potassium, to which reference has already been made, has also been recommended on account of its peculiar sedative action, and M. Bazin (*Gaz. de Hôpitaux*, 1865) narrates a case in which large doses of this remedy were successful in checking the progress of tubercular meningitis, in a lad who presented at the same time the symptoms of pulmonary tuberculosis.

Ergot, which is believed to possess the power of reducing cerebral and spinal hyperæmia, has of late years been recommended in large doses in this disease. Dr. V. P. Gibney, of New York, in a communication to the Academy of Medicine (*N. Y. Med. Record*, 1877, p. 709), reports at length a case of a boy aged 11 years, which certainly presented many of the characteristic symptoms of tubercular meningitis, where recovery followed the use of ergot in large doses (Ext. Ergotæ Fl. fʒss. to fʒj, or Ext. Ergotæ, gr. ijss. every three hours), continued for thirty days. No other remedy was employed to which any share in the result could be attributed. We have used this remedy in several cases, but without any appreciable influence upon the fatal course of the disease.

As a general rule, *narcotics* of all kinds are to be avoided, from their effect of increasing the constipation, and exciting more or less the cerebral circulation. When, however, neither antiphlogistics, evacuants, nor cold or warm applications relieve the sufferings of the child, it would be proper to employ small laudanum poultices or opium plasters upon the forehead or temples, or we may use morphia by the endermic method.

Counter-irritation in different forms has been employed, and apparently with success, though it has failed in our hands. Blisters to the nucha, behind the ears, or over the whole scalp, have been used. At one time, in this city, it was a common practice to cover the scalp with a blister, but it was found to fail so constantly, and was so painful a sight to the relatives of the child, that it has been very much abandoned. Surely, if it had succeeded in any considerable proportion of the cases, it would have been received as a boon, however revolting to the sight. We have, ourselves, in past years, blistered the nucha, the back of the ears, and the temples in a number of cases, but have always failed to obtain any evident good from them. Within a few years it has been claimed that pustulating the whole sinciput with croton oil has been of great service. The last case of tuberculous meningitis we saw, occurred in an adult, and here we had nearly the whole of the crown of the head shaved, and pustulated with the oil, but it was of no use whatever.

Cold applications to the head have been very much used. We have employed them ourselves and still use them whenever the head is hot, or when their use relieves the headache or soothes the patient, but we confess that they have not seemed to us of much use except as palliatives. They may consist of cloths wet with cold water, of affusions with cold water, or, as has been proposed by M. Guersant, of irrigation as employed in surgery. M. Guersant prefers this mode of applying cold to any other, believing it to be the most convenient and comfortable to the child, and from its continuous action, the most efficacious. To make use of it, the hair is to be shaved or closely cut, and the child placed upon a mattress without a pillow, and with its head near the edge of the bed. The head is then covered with compresses of soft rag, or, better still, of patent lint, while under it is placed a piece of oiled silk or india-rubber cloth, so arranged as to keep the thorax from being wet, and doubled into a gutter above to convey the water off into a vessel placed on the floor. A bucket or basin filled with fresh, cool water, is placed near the head of the bed, and from this a siphon made of lint or lamp-wick is so arranged as to convey a stream of water upon the compresses covering the head. If the heat of the whole body falls so much as to threaten collapse after the irrigation has been continued for some time, the stream of water should be stopped, and compresses, merely wet with water not quite so cool, kept on the head. The latter precaution is necessary in order to prevent injurious reaction from the sudden and total removal of so powerful a sedative as irrigation proves to be.

Some practitioners prefer the use of ice in a bladder. This seems, however, too severe a remedy to be long continued, and we should therefore rather use only cloths wet with iced-water, or irrigation. Dr. Abercrombie

is of opinion that the application of cold is by far the most powerful local remedy that we have. M. Gendrin recommends cool or cold affusions over the whole surface, the temperature to be proportioned to the heat of the skin. When there is but little heat of head, only a slight febrile movement, and the headache is not relieved by cold applications, Guersant recommends the substitution of warm poultices to the scalp, in the place of irrigation or cold applications.

The treatment described in the preceding pages, is that which is proper for cases of the disease occurring in subject previously in good health, or evincing but few signs of the tubercular cachexia. When, on the contrary, it occurs in children with extensive tubercular affections of other organs, by which they are already weakened and exhausted, the treatment must of course be modified to meet the circumstance of the case. It ought to consist chiefly of cold applications, and of an early use of cod-liver oil, of iodine, or of the iodide of iron. We should recollect that experience has long since shown the weakness of our art in such cases, and for that reason avoid such a degree of interference as might possibly abridge the little span of life allowed the patient by this relentless malady.

PROPHYLACTIC TREATMENT.—It must be evident that the prophylactic treatment is of special importance in a disease so little amenable to curative means as the one under consideration. When, therefore, there is reason to suspect a tendency to tubercular meningitis in a child, either from the fact that other children in the family have perished with it, or from a bad state of the general health, and frequent complaints of headache, it becomes proper and necessary to regulate both the moral and physical education with a view to its prevention. For this end the hygienic management of the child ought to be such as is best calculated to prevent the formation or development of tubercles in the constitution. During infancy, such a child should be nursed, if this be possible, by a strong, hearty woman, with an abundant flow of milk. If the mother is not possessed of these qualities, if there be, indeed, the least doubt upon the point, she ought without hesitation to give up the pleasure of nursing the child herself, and procure for it a wet-nurse of the kind described. This alone will, in all probability, often make the difference between a vigorous and a fragile constitution. When the time for weaning arrives, the change ought to be made with the greatest care and circumspection. During and for some time after weaning, the diet must consist principally of milk preparations and bread, and of small quantities of light broths, or of meat very finely cut up. As the child grows older, the meals ought to be arranged at regular hours, and should consist of four in the day. The principal food must be bread and milk, well chosen, well cooked meats, and rice and potatoes as almost the only vegetables. After the first dentition is completed, a moderate use of ripe and wholesome fruits may be allowed, but always with care, in order to avoid injury to the digestive organs, and also so as not to mar the appetite for some wholesome and nutritious food. Coffee and tea ought to be forbidden at all times; since, as we have often observed, when the palate of a child is taught, by habit, to become accustomed to these more highly sapid substances, it is very apt

to abandon the use of milk, which ought to constitute a large proportion of its food, at least up to the age of twelve or fifteen years. In no circumstance of life is the old saying, "where ignorance is bliss, 'tis folly to be wise," a better rule of action than in regard to the diet of our children. The child should not taste improper articles of food, so that it may escape the torment of desiring what is improper.

After diet the most important points in the treatment are air and clothing. The child should inhabit, if possible, a large, dry, well ventilated room, which ought to be kept as cool as possible in summer, and moderately warm in winter. Not a day should be allowed to pass, unless the weather is totally unfit, without the child's being sent out for several hours into the open air, and we believe that it is much better for it to walk than drive, unless the weather be very hot. The clothing ought to be suitable to the season, cool in summer and warm in winter. In our country there is a great inclination to *harden* children by dressing them very slightly in cold weather; so that they frequently suffer from catarrh, pneumonia, and spasmodic croup brought on by improper exposure. This cannot but be wrong in a child who shows the least evidence of tendency to tubercular affections.

For our own part we are fully convinced from what experience we have had of the diseases of children, that by far the most certain and effectual means of preventing the development of a tubercular, or indeed any other cachexia in a child, is to have it brought up in the open country, or in some healthy village, until the epoch of puberty has passed by safely. A very good plan for parents whose occupations compel them to live in cities or large towns, is to have their residence a few miles in the country, and to come to town every day. Children brought up in this way have a far better chance of obtaining strong and vigorous constitutions, than those reared entirely in the close and confined dwellings and streets of crowded cities.

When a child, who, from the health of its parents, or from its own appearance, may be suspected of having any tubercular or scrofulous taint in its system, becomes subject to frequent attacks of apparently causeless headache, and especially when such headaches are associated with a constipated habit of body and with occasional vomiting, it ought to be looked upon as threatened with tubercular disease of the brain. Under these circumstances we would advise, in addition to the measures just now recommended as to diet, dress, exercise in the open air, and a residence in the country, that it be put at once upon the use of cod-liver oil, iodide of iron, and mild laxatives, and that these be persevered in for several weeks or months, until in fact the strength and general health are restored and the headaches cease. When the appetite is poor, and the digestion is imperfect, in such a case we may use with advantage, beside the above remedies, solution of pepsin, a teaspoonful three times a day with the meals, or tincture of *nux vomica*, three or four drops in a mixture of syrup and compound tincture of gentian, or in a teaspoonful of elixir of cinchona three times a day. If the child is of an age to be going on with its education this should for the time cease, or be carried on in such a way as to

avoid all excitement or fatigue. A case occurred to us in the course of the year 1852, which showed, we think, very clearly the utility of these measures.

A boy between seven and eight years old, whose mother had died of well marked phthisis a few months before he was put under our charge, had been losing flesh and strength, and suffering from occasional headache for some time before we were called to see him. We found him in bed complaining of severe frontal headache; so severe at times, and usually in the after part of the day, as to cause great distress, with crying. The intelligence was perfectly natural. The child was rather dull and listless, from suffering and from weakness, but not from any want of a healthful state of the mental operations. There was no sign whatever of spasmodic or paralytic affection. In the morning the skin was cool and natural, but in the afternoon it became warm and dry, but not very hot. The pulse was 62 to 68, and though not actually irregular, it was halting or hesitating. There was occasional, but not frequent, unprovoked vomiting, and he complained often of sick stomach, even when he did not vomit. The bowels were very much constipated, and had been a good deal so for some weeks previous to his falling actually sick. There was no cough, no sore throat, and no soreness about the abdomen. The tongue was moist, soft, slightly furred, and not red nor gashed. The urinary secretion was healthy. Physical examination showed the lungs and heart to be without disease.

The treatment during the first week was small doses of calomel and rhubarb, half a grain of the former to two of the latter, given for a day, and followed by syrup of rhubarb and fluid extract of senna, until the bowels were copiously evacuated. After this the bowels were kept soluble by the administration every day, or every other day, of doses of Seltzer powder, sufficient to produce the effect. Blisters were applied behind the ears. In the after-part of the day, when the head and body became heated, cooling applications were made to the head, and the feet were put into mustard-water, once, twice, or three times. Two grains of iodide of potassium were ordered to be given three times a day. The diet was to be light but nutritious. It was to consist of bread and milk and a soft-boiled egg in the morning, oysters or light meats with rice for dinner, and milk with bread in the evening. Of these he was to have any reasonable quantity that he might desire. Under this treatment he improved slowly, with occasional drawbacks for a week, when the iodide of iron was substituted for the iodide of potassium. The bowels continued very costive, requiring daily doses of the Seltzer powder; the headaches diminished in frequency, duration, and severity; the pulse went up to 72 and 78, and became more free and even; the appetite had improved, but the child remained still very weak, pallid, and quite emaciated. After another week, as he continued to mend, and the stomach had become stronger, cod-liver oil was ordered in addition to the iron; a teaspoonful was to be taken three times a day in a wineglassful of table-beer. As he gained strength, the amount and kind of food was increased. He was, indeed, encouraged to eat heartily of plain and digestible substances.

He now improved gradually in health. The headaches subsided, and finally ceased; the bowels became soluble; the appetite grew hearty and strong, and all feeling of nausea disappeared; he regained his strength, flesh, and color, so that at the end of two months we saw him looking quite fat and well. The iodide of iron and cod-liver oil were, however, to be continued for a month longer. He is now (1869) a young man in very good health. He has passed several years in Germany pursuing a scientific education, and has returned lately to this country, and is about to marry.

If in any child whose hereditary tendencies or whose physical characteristics are such as to make us fear a predisposition to tuberculosis, there should also be evidences of marked nervous irritability or precocious mental development, it is desirable to use every means to prevent a continuance

of such undue cerebral activity, which might tend to induce tuberculosis of that organ or of its membranes. We must, however, be satisfied for the most part with a careful attention to all the details of sound hygiene, in addition to which, however, the following special points deserve mention, viz., to keep the head cool by not allowing it to be very warmly covered, and by keeping the hair short; to keep the extremities warm; and to avoid stimulating the intellectual faculties to any considerable extent by education, until after eight or ten years of age. The long-continued employment of a powerful derivative from the brain, as a seton in the neck, seems to us to be attended with too many serious objections to be at all desirable. If finally in such children there should be extensive eruptions on the scalp, it may be safe to undertake their cure by suitable internal remedies and mild external applications, rather than to try by powerful local treatment to rapidly remove the affection of the skin. We still mention this caution, despite the fact that the classical investigations of Hebra have shown that for the most part the danger of inducing internal disease by quickly curing cutaneous eruptions is a purely imaginary one.

ARTICLE II.

SIMPLE MENINGITIS.

DEFINITION; SYNONYMS; FREQUENCY.—By this term is understood inflammation of the membranes of the brain, independent of tuberculosis of those tissues, or of other organs of the economy.

The disease was for a long time confounded with tubercular meningitis under the titles of water on the brain, dropsy of the brain, and acute hydrocephalus. It has also been called arachnitis; and more rarely phrenitis.

Its frequency is much less than that of tubercular meningitis. West (*op. cit.*, 4th Amer. ed., p. 100) states that he has seen seven cases of fatal acute meningitis, in five of which post-mortem examination was made and confirmed the diagnosis. Vogel (*op. cit.*, p. 359) speaks of it as being much rarer than the tubercular form, and states that it is no more frequent in children than in adults. It appears that MM. Rilliet and Barthez, during their researches, met with only five cases of this disease, while they report thirty-three of tubercular meningitis. Bouchut states that he has met with two cases of simple meningitis to six of tubercular disease, whilst Barrier reports only four of the former in nearly thirty autopsies of meningitis. He states, however, that he has met with three cases of recovery, all of which he believes to have been instances of the simple form. Our own experience agrees with that of Bierbaum (*Die Meningitis Simplex*, Leipzig, 1866) in showing that it chiefly attacks infants under two years of age.

CAUSES.—The causes of simple meningitis are not very clearly ascertained. M. Rilliet, who published a very valuable paper on this affection

(*Arch. Gen. de Méd.*, t. xii., 1846), divided it into two forms, the convulsive and phrenitic, the former of which he believed to be most common under two, and the latter between five and fifteen years of age. Subsequent experience has confirmed this view. From the fact that the disease is most frequent in the first and ninth years of life, Rilliet concluded the process of dentition has something to do in its production. It appears also to be more frequent in boys than girls, and in robust than in weak constitutions. Exposure to extremes of temperature predisposes to attacks of acute meningitis; and, in particular, continued exposure to the direct rays of the sun has been known to act as an immediate cause.

Guersant has known it to follow such exposure in several instances, particularly in young infants; MM. Rilliet and Barthez report a case of the same kind, and Rilliet (*loc. cit.*) another. Other causes cited by authors are injuries upon the head, such as blows, falls, and wounds. It also occurs as a consequence of extension of inflammation to the membranes of the brain, and usually from the internal ear in cases of otorrhœa.

The disease seems to have sometimes occurred in an epidemic form. There is reason to believe, however, when we consider the purely sporadic nature of its recognized causes, that the reported epidemics have for the most part been of cerebro-spinal meningitis, a disease which we shall treat of in its appropriate place, among the acute specific febrile affections.

ANATOMICAL LESIONS.—The dura mater is generally much injected, and its sinuses, together with the large cerebral veins, contain coagulated or semi-coagulated blood, sometimes in large quantities. On opening the dura mater, the whole, or nearly the whole of the convex surface of both hemispheres, or in some instances of one only, are found to be covered with a yellowish or greenish-yellow layer, which consists of fluid or concrete pus, or of false membranes. These deposits exist also on the internal surfaces of the hemispheres, on the upper surfaces of the cerebellum, and often also at the base of the brain, though in some cases the latter presents none whatever. The inflammatory products are seated in the pia mater, and sometimes in the cavity of the arachnoid membrane, but in much smaller quantity than in the tissue beneath that membrane.

The *arachnoid* membrane, which covers the brain, seldom participates in the inflammation, but remains smooth and transparent. Its cavity, however, sometimes contains inflammatory products, which, when death occurs early in the attack, consist of a small quantity of pure pus, or of larger quantities of a turbid, yellowish serosity, consisting of serum and pus mixed together. When death has occurred later in the disease,—after five, six, or seven days,—the pus is mixed with lymph, or else true false membranes are found. The *pia mater* is observed to contain fluid or semi-fluid pus when death occurs before the fourth or fifth day; while in less acute cases there are patches or large layers of lymph, which sometimes dip into the anfractuositities, and give to the membrane under consideration a swelled and thickened appearance. These appearances are more marked on the superior and lateral, than on the inferior surface of the brain. Where the deposits exist the membrane presents a vivid injection, which is more marked in proportion as death has taken place earlier in the disease. The

pia mater is generally easily detached from the cerebral substance, particularly when the fatal termination has occurred early. The *substance* of the brain is firm, and but slightly colored, in rapid cases. When the course of the disease has been slower, the cineritious portion is generally of a bright rose color, and the medullary substance abundantly dotted with red, showing that the inflammation has involved the superficial layer of the brain. In the latter class of cases the surface of the convolutions is usually softened, and the pia mater adherent. In very young children the whole brain is sometimes soft.

The *ventricles* do not, as a general rule, contain transparent serum, except at a very early age, when serous effusion takes place with great facility. They, often, however, contain one or two teaspoonfuls, and rarely more than one or two tablespoonfuls, of pus or purulent serum. The serous membrane of the ventricles and the plexus choroides exhibit signs of inflammation in some instances. They are of a bright red color, uneven, rough, and very much softened, in children who die early; and pale, opaque, slightly thickened, and rough, in those who die at a later period.

The central parts of the brain often retain their firmness, but are sometimes softer than natural, or even diffuent. This softening is particularly apt to exist in very young children, in connection with large effusion into the ventricles; though it also occurs in those who are older, and in whom there is only slight effusion of pus or purulent serum. In the former case it is probably due to the macerating effect of the effusion, while in the latter it is more likely to be owing to inflammation.

In some cases, and especially those of the epidemic form of the disease, the membranes of the spinal cord are found to present the same inflammatory appearances which have been described as existing in the cerebral meninges. These cases are, therefore, more correctly designated by the name cerebro-spinal meningitis.

The other organs are healthy except in secondary cases. Tubercles, which so constantly exist in various other organs in tuberculosis of the meninges, are never found, according to M. Rilliet, in this form of meningitis. This author believes himself entitled from his researches to formulate the following law of pathological anatomy; "That general meningitis and meningitis of the convexity of the brain occur only in non-tuberculous children, whilst meningitis of the base of the brain without inflammation of the lining membrane of the ventricles, belongs exclusively to tuberculous children." (*Op. cit.*, t. iii, 1846, p. 408.)

This law cannot, however, be adopted without exception, since we have already seen, when speaking of tubercular meningitis, that there are, in a large proportion of such cases, evidences of inflammation of the lining membrane of the ventricles.

SYMPTOMS.—The following account of the symptoms of the disease is taken chiefly from the paper of M. Rilliet. That author describes two forms of the affection, the *convulsive* and *phrenitic*; the former of which is characterized by a predominance of convulsive phenomena, and the latter by disorders of the intelligence.

The disease may also be idiopathic or secondary, simple or complicated, sporadic or epidemic.

The *convulsive form* generally occurs in children under two years of age. The disease usually begins suddenly or after a restless night, with a violent and prolonged attack of *convulsions*, oftener general than partial, and is accompanied by violent *fever*, and sometimes by considerable quickness of *respiration*. The existence of *headache* cannot be ascertained at this early age. *Vomiting* is often absent, and the *bowels* generally continue regular in this form, though they are sometimes constipated. After awhile the convulsions cease, and the child remains for the time in a state of quiet, somnolence, or coma, when they return with renewed violence. The returns of the convulsions generally take place at intervals of one or two hours or more. In the intervals between the crises the child is restless or drowsy, or in a state of partial *stupor*, attended with tremulous movements of the extremities; there is *strabismus*, *contraction of the pupils*, *trismus*, and sometimes *hemiplegia*. The skin retains its warmth, the pulse is accelerated, irregular, and unequal; the face is pale; the stools are spontaneous or easily procured by remedies. It is unusual to see the child regain its consciousness so as to recognize objects in the intervals between the convulsions, or after the appearance of coma and other cerebral symptoms. Death occurs during coma or in a violent attack of convulsions. This form seldom lasts more than four days.

Occasionally this form begins in a different manner. The convulsions, though they still predominate, do not occur until later in the disease, and the whole course of the affection is slower. Such cases begin with a violent febrile movement, lasting several days, and accompanied by acceleration or unevenness of the respiration, or by almost constant drowsiness, preceded or followed by agitation, screaming, staring expression of the eyes, and dilatation of the pupils; vomiting and constipation are sometimes present, at others absent. After a time, however, convulsions make their appearance, and the case follows the course already described. The duration of this form may be the same as that of the first, or it may last about two weeks.

The *phrenitic form* of simple meningitis generally begins suddenly with fever, which is sometimes preceded by a chill; the skin is warm and dry, and the pulse, in idiopathic cases, full and accelerated. In secondary cases the pulse has been found slow and irregular; in all it becomes irregular, small, and very rapid the day before death. Simultaneously with the fever there is frontal headache, which is often so violent as to draw cries from the child, and, according to M. Rilliet, is more severe than either in tubercular meningitis or typhoid fever. It is also more constant, and lasts generally one, two, or three days, until the appearance of restlessness, delirium, or coma. At the same time there is great sensibility to light and noise, and abundant vomiting of bilious matter. The latter symptom is one of the earliest; it generally ceases after a few days, but sometimes continues to the very end. Constipation exists in some cases, but is much less constant and more easily overcome than in the tubercular disease. The appetite is lost, and the thirst very acute. The abdomen is

flattened and retracted, especially towards the end, while in secondary cases of this form, and in very young children, it retains its usual shape.

About the end of the first day, generally, or, in rare instances, after two or three days, appear various disorders of the intelligence. The first symptom of this kind is observable in the expression of the face, which becomes a little wild or wandering, and sometimes grimacing. Soon afterwards occur restlessness, which is sometimes extreme, and, in succession, delirium, somnolence, and, later in the attack, coma. The restlessness and somnolence often alternate early in the case, though the former generally predominates and soon passes into delirium, which is usually violent. When in this condition the child seldom recognizes any one, and either refuses to answer questions, or answers incoherently. In connection with the disorders of intelligence there exist also trismus, grinding of the teeth, subsultus tendinum, partial convulsive movements, stiffening of the extremities or trunk, retraction of the head, strabismus, contraction first and then dilatation of the pupils, and in some cases violent convulsions, followed by deep coma. Death sometimes occurs at this period. In other instances, the disease continues longer, and other symptoms declare themselves. Vomiting generally ceases; constipation increases; the abdomen is retracted; headache is no longer complained of; the fever continues, but the pulse becomes irregular; the respiration is uneven and irregular, being sometimes more and at other times less frequent than natural; the face is distorted and extremely pale, or there may be a purple flush on the cheeks; an erythematous streak (*tâche meningitique* of Trousseau) may often be observed after drawing the finger lightly over the skin; the restlessness is excessive, and accompanied by subsultus, carphologia, or partial convulsive movements; the delirium, at first so violent as to make it necessary sometimes to hold the child in bed, subsides into a state of coma and collapse, in which general sensibility is obtunded, and special sensibility extinguished; the respiration becomes stertorous, and at length asphyxia, coma, or a severe attack of convulsions terminates the scene.

The course of the disease is generally continuous. In very rare cases, however, occasional remissions occur, so that the child recovers its intelligence for a short time, and recognizes persons around. The duration has varied between a day and a half and nine days.

When, on the other hand, a case of either form tends towards recovery, the graver symptoms gradually subside. Convulsive movements or cerebral excitement lessen and are replaced by moderate stupor, from which the child can be at least partially aroused. More or less complete paralysis of one side or of one member, irregularities in the pupils, with or without strabismus, tonic muscular contractions, affections of the special senses, as deafness or impaired vision, may persist for a number of days or even for several weeks. Gradually the capacity for taking food returns, the above-mentioned nervous phenomena disappear, intelligence increases, and convalescence passes slowly into recovery, which may be complete or may be marred by the persistence of some defect of special sense.

We have observed several cases where severe and fully developed meningitis, undoubtedly of the simple non-tubercular form, has terminated favorably as above described.

DIAGNOSIS.—The convulsive form may be confounded with the essential or symptomatic, and with the sympathetic convulsions of children. The mistake may generally be avoided by attention to the following points. In essential convulsions, the attacks are usually less violent, seldom last more than a few moments, occur from some evident cause, and do not recur often. When they have ceased, the child generally soon regains its consciousness and health, or exhibits slight drowsiness, or derangement of movement for a short time only. In such cases the respiration is not permanently accelerated, as in convulsive meningitis; the pulse, if it has been increased in frequency, soon falls to the natural standard, and special sensibility remains undisturbed.

It is to be distinguished from sympathetic convulsions by the characters just described, aided by a reference to the disease which may have caused the attack of eclampsia, and which may be one of the eruptive fevers, enteritis, indigestion, pneumonia, or any other acute affection. In some instances, however, the distinction cannot be made except by attention to the progress of the attack.

The phrenitic form may be confounded with tubercular meningitis, with congestion of the brain, or with the early stage of the eruptive fevers. The distinction between it and tubercular meningitis has already been considered under the head of the latter disease.

It is sometimes difficult, as pointed out by Rilliet, to distinguish between simple meningitis and cerebral congestion and partial encephalitis.

In congestion, however, there is not the same intense headache, the febrile movement is not so marked, vomiting is usually absent, and the development of delirium or coma, or of convulsive or paralytic symptoms is more sudden or even instantaneous.

Partial encephalitis is even more rare in children than simple meningitis. It may be distinguished from the latter by the less severity of the headache, by the less marked delirium, by the comparative infrequency of vomiting, and by the less activity of the febrile movement, and the more gradual course of the case in encephalitis.

Severe or malignant scarlatina in its stage of invasion may stimulate meningitis, but can be readily distinguished by attention to the remarkable elevation of temperature and rapidity of the pulse, to the heavily coated tongue, and to the decided throat symptoms. Typhoid fever may also resemble simple meningitis in some respects, but can be recognized by its more gradual onset, by the course of the febrile action, by the tendency to bronchial irritation and to abdominal symptoms, and by the characteristic eruption. It may happen, however, that meningitis develops during the course of typhoid fever, in which case it is extremely difficult or impossible to determine its existence, unless an ophthalmoscopic examination can be made and reveals changes in the fundus of the eye.

PROGNOSIS.—The prognosis of simple meningitis is very grave, but much less hopeless than in the tubercular form. M. Rilliet (*loc. cit.*) cites several instances of recovery, but states that death is much the most frequent termination.

TREATMENT.—It must be evident, it seems to us, that but little depen-

dence can or ought to be placed on any but prompt and powerful *antiphlogistic* treatment. *Depletion*, therefore, *mercury*, *cold* applications to the head, *laxatives*, *counter-irritants*, and the most rigid diet, ought to be employed from as early a period as possible.

If there is any case, excepting those in which venesection is indicated for the relief of mechanical engorgement of the right heart, when general bleeding would appear to be preferable to local depletion, acute simple meningitis in a vigorous child may be cited as such. We are not prepared, therefore, to discountenance its performance even in very young children, although in our own practice we have relied upon the application of leeches to the temples or behind the ears. We may remark that MM. Rilliet and Barthez object to the application of leeches to the head, and propose that they should be placed rather about the anus or on the inferior extremities. The quantity of blood to be drawn must depend upon the age and constitution of the subject, and the violence of the attack, in some measure. It should always, however, be large, as much or more, we think, than is necessary in any other of the acute affections of childhood. In a child two years old, of good constitution, from two to four ounces would not be too much at first, and should the symptoms not moderate in six or eight hours, as much more may be taken. We are disposed to believe that in such a disease as this, bleeding is by far the most powerful remedy, and it is perhaps the only one which offers us any real chance of success, at least in those rapid cases in which extensive layers of pus and false membranes are found on the surface of the brain, in the pia mater, or in the subarachnoid tissue, in from two days and a half to three or four days after the commencement of the disease.

The application of *cold* to the head constitutes another most efficient remedy in inflammations of the brain and its membranes. These means may consist of a bladder containing water and pounded ice, which is perhaps the most convenient and powerful, of cloths wrung out of iced or very cold water, to be constantly renewed, of cold affusions upon the head, or lastly, of irrigation as recommended by M. Guersant, and described in the article on tubercular meningitis. *Purgatives* are often employed very freely in this disease, with the view of producing a decided revulsion upon the gastro-intestinal mucous membrane. We would, however, advise caution in their administration, lest excessive irritation of that membrane be produced, with consequent derangement of digestion. If there is reason to suspect the presence of undigested or irritating matters in the alimentary canal, a dose of castor oil, citrate of magnesia, or rhubarb may be given. But the method we prefer is to give calomel in small and frequently repeated doses until the bowels are moved, and afterwards to continue it in still smaller doses, given at longer intervals, in order to secure its specific influence on the inflammation.

Some writers also recommend very highly the use of mercurial inunction. Vogel (*op. cit.*, p. 361) states that a mercurial treatment is decidedly effectual, and adds that the only two children he has seen recover from this disease were treated exclusively with mercury, internally and externally.

Iodide of Potassium should unquestionably, in our opinion, be given in full doses as soon as the calomel is suspended, or from the beginning in case this latter remedy is not employed. The evidence in its favor, as tending to induce absorption of the exudation, seems very strong; and in our own successful cases it was given after the first few days, during which calomel was administered, and continued until convalescence was fully established.

Ergot, as already mentioned in the article on tubercular meningitis, has recently been recommended strongly in both that and the simple form of the disease; and certainly, in the latter, its power of lessening hyperæmia of the intracranial vessels indicates its use in conjunction with iodide of potassium.

Bromide of Potassium and bromide of sodium are the most valuable sedatives in the acute stage of this disease when marked symptoms of cerebral excitement or of a convulsive tendency exist. The doses should be full and frequently repeated; as, for instance, at one year of age, from 3 to 5 grains every two or three hours.

Counter-irritants are useful as adjuvants to the more powerful remedies already indicated. During the first day or two they should consist chiefly of sinapisms and mustard poultices, applied from time to time to the trunk and extremities. Authorities differ somewhat as to the effect of blisters, and as to the time at which they ought to be applied. M. Valleix (*op. cit.*, t. ix, p. 187) opposes their employment in this affection as often injurious and still more frequently useless. We think the advice given by Dr. Abercrombie, as to their employment, is probably the most prudent. This is, not to apply them in the early stage, but to wait until the active symptoms of the disease have been subdued. They may be applied to the head itself, to the nucha, or to the extremities. We believe that we have seen them most useful when applied to the neck and inside of the calves of the legs. Nevertheless, there is his authority in favor of their good effects when applied upon the head itself.

M. Rilliet (*loc cit.*) recommends a vigorous revulsion upon the scalp when the disease has followed the suppression of an eruption. He proposes with this view the employment of pustulation by croton oil, and relates a case of recovery which followed this treatment under a most unfavorable train of symptoms. To make use of it the head must be first shaved; from fifteen to twenty drops of the oil are then to be rubbed over the scalp with a glove four or six times a day. Before making the friction, the eyes of the patient must be covered with a band to prevent the introduction of any of the oil into them, as this would be apt to occasion severe ophthalmia. In the case reported by him, a considerable number of pustules were produced in twenty-four hours, and in a few more hours the eruption was general, so that the head was covered with a kind of cap of a fine yellow color.

ARTICLE III.

CEREBRAL CONGESTION.

CEREBRAL CONGESTION implies a condition in which the bloodvessels of the brain contain an excessive quantity of blood. It occurs under the two forms of *active* congestion, where there is an increase in the amount of arterial blood, and of *passive* congestion, where the quantity of venous blood is excessive.

While it cannot be doubted that the amount of blood circulating in the brain is thus liable to vary, and that such variations are of frequent occurrence, and are attended with very important symptoms, it is difficult to determine the relative frequency of cerebral congestion at different periods of life. Some authors of large experience assert that it is much more frequent in infancy and childhood than at any later period of life; and this opinion is supported by the extreme mobility of the circulation in early life, and by the frequency in childhood of the symptoms that are usually attributed to congestion of the brain. Without desiring to dissent strongly from this opinion, it is important to remember that, in the first place, as asserted long ago by Rilliet and Barthez (1ère edit., t. i. p. 649), the anatomical appearances of cerebral congestion are occasionally found in children dying of different diseases without having presented any cerebral symptoms; and, again, that the symptoms of congestion of the brain may, in our opinion, be due sometimes to mere excitement and undue rapidity of the circulation, and in other cases to the irritation of the brain, caused by the circulation through it of blood vitiated by the poison of some of the acute specific diseases.

CAUSES.—Active congestion may occur during the process of dentition, or may result from exposure to the sun, from falls or blows on the head, or from excessive excitement or fatigue in children who are predisposed to the affection. The cerebral symptoms occurring at the outset of the eruptive fevers have been, by West and others, attributed to the development of intense cerebral congestion. We feel, however, that the nervous symptoms just alluded to ought, most frequently at least, to be regarded as the result of the presence in the nervous centres of a diseased and vitiated blood, rather than of congestion. That congestion does not always produce them is shown by the statement of Rilliet and Barthez (*op. cit.*, t. ii, p. 620) in regard to the cerebral symptoms of scarlet fever, “that a more or less sanguine congestion (of the cerebro-spinal apparatus) is the only alteration *generally, but not always found*, and sometimes the congestion is not more marked than in other diseases in which there had been no cerebral symptoms.”

In reference to the cerebral symptoms which less frequently occur in inflammations of important organs, as, for instance, pneumonia or enterocolitis, we are not yet in possession of sufficient facts to determine whether they result from reflex irritation, from the elevated temperature and acceleration of circulation, or from changes in the quantity of blood circu-

lating through the brain. Probably all three of these elements take part in varying proportion in the production of such symptoms; we do not, therefore, feel at liberty to regard them, in the majority of cases, as actually depending on cerebral congestion.

Passive congestion results from such causes as offer an impediment to the reflux of the venous blood from the brain. Among them may be mentioned the mechanical objection caused by the pressure of an enlarged thymus gland, or of enlarged cervical or bronchial glands, or by the partial or complete occlusion of a large vein or sinus from the formation of a fibrinous concretion (thrombosis) in its cavity, or from the pressure of a tumor upon its walls. Passive congestion occurs also in affections which, like whooping-cough, are attended with violent paroxysms of cough, during which the return of venous blood from the brain is greatly impeded. Finally it often appears that the state of feeble, languid circulation, depending upon want of pure air or of sufficient and nourishing food, strongly predisposes to, or actually induces, this form of cerebral congestion.

SYMPTOMS.—Recognizing that cerebral congestion may occur at the onset or during the course of various acute diseases, it is evident that the symptoms due to the congestion itself must frequently be complicated by those of the primary affection. We may, however, give the following brief sketch of the symptoms which may be attributed to the two forms of cerebral congestion.

The *active* form usually appears suddenly, though it may be preceded for a few days by a state of indisposition, with irritability and peevishness, some fever, and a disordered state of the bowels generally, but not always consisting of constipation. The chief symptoms of the attack are great heat of the head, and complaints, in older children, of headache, intolerance of light and sound, nervousness, with startings during sleep and twitchings of the muscles. The pulse is frequent, the carotids throb, and if the skull be still unossified the anterior fontanelle is tense and prominent, or the brain is seen and felt to pulsate forcibly through it. If these symptoms be not relieved by appropriate treatment, or, in some cases, without any premonitory stage, the child may pass into a state of more or less profound stupor or coma; or, on the other hand, an attack of partial or general convulsions may occur.

In *passive* congestion the symptoms are apt to be less suddenly developed, but when marked they resemble in many respects those which we have described as indicative of the active form. There is, however, less febrile excitement, and the force of the arterial pulsation and the prominence and tension of the fontanelle are notably less. Still there are usually present great irritability, restless, disturbed sleep, muscular twitchings, or even convulsions; or, on the other hand, deepening indifference to surrounding objects, resulting in profound stupor.

The *duration* and *termination* of these symptoms are very variable. If the anatomical condition present has been only one of great vascular distension, and appropriate treatment is promptly employed, the threatening symptoms subside in the course of two or three days. If, however, the

congestion has been so extreme or has lasted so long as to lead to serous effusions or even to minute extravasations of blood, the symptoms may continue to deepen in gravity until the fatal result occurs, or else the attack subsides but leaves behind it some evidence of injury to the brain-tissue in the form of more or less lasting paralysis. Occasionally, however, complete recovery takes place, contrary to all expectation, after the grave symptoms above described have continued but slightly modified for days or even weeks.

The *diagnosis* of cerebral congestion must be based upon the symptoms above detailed, as well as upon the general considerations with which we prefaced this article. The reader is also referred to the remarks made under the head of diagnosis in the article on simple meningitis. It is, of course, essential not only that the existence of cerebral congestion should be recognized, but also that the form in which it presents itself should be determined, as this has a most important bearing upon the treatment to be employed.

TREATMENT.—Active cerebral congestion is to be treated like the first stage of simple meningitis, with cathartics and purgative enemata, calomel, cold applications to the head, baths, revulsives, full doses of bromide of potassium, low diet, and confinement to a cool dark chamber. We desire to lay particular emphasis upon the employment of bromide of potassium, as its power of lessening active congestion of the nerve-centres is established by very positive evidence. Ergot or belladonna, which exert a similar action, may be associated with it. If the symptoms do not promptly yield to these measures, and there is no special contra-indication, we should recommend the use of local depletion by leeches or wet cups to the temples or some part of the head.

In the treatment of the *passive* form, particular attention must be given to the relief or removal of the primary cause. Unfortunately, however, in many instances this is not possible. The urgent symptoms themselves must be promptly treated by cold applications to the head, by active revulsion, by strict attention to the diet and the state of the bowels. If great danger exists, and the nature of the cause and the condition of the child justify it, mild local depletion may be cautiously employed. If, on the other hand, the case be associated with enfeebled nutrition, it may be necessary to employ quinia, ammonia, as nourishing a diet as can be digested, and small amounts of stimulus.

If the congestion terminate in extravasation, the treatment for this condition and the paralytic symptoms which may result must be such as is recommended under the head of cerebral hemorrhage.

ARTICLE IV.

CEREBRAL HEMORRHAGE.

WE shall consider hemorrhage of the brain under two heads, that of the substance, and that of the membranes; the former is usually designated as cerebral, and the latter as meningeal apoplexy. Both these forms of hemorrhage are of rare occurrence in childhood compared with other diseases of the brain, and with their frequency during adult life and old age. Of the two kinds, that of the meninges is the more common.

DEFINITION; FREQUENCY; FORMS.—By cerebral apoplexy or hemorrhage is understood an effusion of blood into the substance of the brain. By meningeal apoplexy or hemorrhage is understood an effusion of blood between the dura matter and cranium, into the cavity of the arachnoid membrane, beneath the arachnoid, or in the meshes of the pia mater. Cerebral hemorrhage is a rare affection in childhood, while meningeal apoplexy is of more frequent occurrence. Rilliet and Barthez met with only eight cases of the former in their extensive experience, and with eighteen cases of the latter. We have ourselves met with three cases of hemorrhage into the substance of the brain, and with several of meningeal apoplexy.

Hemorrhage into the substance of the brain occurs in two different forms: one in which the effused blood is contained in a cavity caused by a laceration of the tissue of the organ, and designated apoplexy in a cavity; and the other in which the blood is effused in a multitude of little points of different sizes, and designated capillary apoplexy.

In addition to this, as in a remarkable case published by Dr. Dulles (*Philada. Med. Times*, July 22d, 1876), the hemorrhage may occur into the ventricles of the brain, completely filling these cavities.

In meningeal hemorrhage the blood may, as we have stated, be effused between the dura mater and the bone. This form, however, is very rare, so rare, indeed, that several writers deny its existence. It is proved, however, to have occurred, by a case reported by MM. Rilliet and Barthez, which is the only one they have met with. In by far the most common form of the disease, the blood escapes into the cavity of the arachnoid membrane, though in rare instances it is effused beneath or in the meshes of the pia mater.

CAUSES.—The causes of intracranial hemorrhage, which appear to be much the same in both forms of the affection, are rather obscure. In new-born infants, hemorrhage may ensue upon the intense congestion of the vessels of the brain and its meninges occurring during severe and prolonged labors. Again it is common to find more or less intracranial hemorrhage, either cerebral or meningeal, in children who have died from tetanus neonatorum; a result to be explained by the intense passive congestion occurring during the stage of tetanic rigidity.

The causes of cerebral hemorrhage are very obscure, so much so, indeed, that some writers have not attempted to ascertain them. They ap-

pear to be the same in both forms of the affection. Amongst the ascribed causes are the sudden disappearance of eruptions of the scalp, observed in two cases by MM. Rilliet and Barthez, in one of which this effect is stated to have been produced suddenly by medical treatment, while in the other it followed the application of poultices to a favous eruption upon the same part. This cause must, however, it appears to us, be regarded as purely illusory. The disease is stated by M. Legendre to have followed in one case a violent fit of anger. It is said also to have been produced by various causes which acted as impediments to the circulation. The obstacle may be situated within or exterior to the cranium. To the first class belong cases in which the sinuses and large venous trunks of the head have been found obstructed by coagula of blood, or by the pressure of tumors, generally of a tubercular nature; to the latter, those in which there is intense engorgement of the superior cava produced, as in prolonged paroxysms of hooping-cough, or in obstructive cardiac disease, or where there is compression of this vessel by enlarged and tubercular bronchial glands. Another cause is thought to be the existence of confirmed cachexia and general debility from any diseased condition whatever, in which the blood having become thin and lost its plasticity, escapes from the vessels with great facility. This last condition is one which almost always exists in connection with the causes cited as acting through the agency of obstruction to the circulation, and tends of course to augment their dangerous effects. Occasionally, also, aneurisms of the cerebral arteries, especially of the middle cerebrals, occur at an early age (several cases are on record at the age of 14 years), and by the rupture of the sac give rise to excessive and rapidly fatal cerebral hemorrhage.

We have met with one case of extensive hemorrhage into the left corpus striatum and adjoining tissue in a boy of 13, evidently connected with advanced granular degeneration of the kidneys, resulting from a previous attack of scarlatina.

In some instances the hemorrhage occurs in the healthiest and most vigorous constitutions, and cannot be accounted for in any way.

It appears that meningeal apoplexy is most frequently met with in very young children, according to MM. Rilliet and Barthez, between the ages of one and two and a half years, whilst M. Legendre did not meet with a single case after three years of age in 248 autopsies. Cerebral and ventricular hemorrhage, on the contrary, are much more common after three years of age than before, which is just the reverse of the law in regard to meningeal effusion.

ANATOMICAL LESIONS.—The description of the lesions of hemorrhage into the substance of the brain need not detain us long, for they are much the same as those observed in the adult. When the blood is effused into cavities (apoplexy in cavities), the latter are usually small in size, seldom exceeding from one to two-thirds of an inch in diameter, though in rare cases they have been found much larger. The cavity is formed by a laceration of the substance of the brain, and is filled with soft, dark coagula, or sometimes with fluid blood; the walls of the cavity consist sometimes of the substance of the brain, which may be of a rosy color and natural con-

sistence, or yellowish and softened, while in other instances they are formed of more or less numerous points of capillary apoplexy. The capillary form of effusion occurs in the shape of a number of points, scarcely so large as the head of a small pin, and of a dark or brownish color, which contrasts strongly with that of the cerebral tissue. Those points evidently consist of true coagula, which are sometimes surrounded by small yellowish areolæ. The substance of the brain around the effusion is either white, firm, and perfectly healthy, or softened, and of a whitish, reddish, or yellowish color. The capillary effusions are generally limited within a space of from a third of an inch to an inch and a half in size, but they have been found scattered over a large portion of the hemispheres.

Both forms of hemorrhage are much more common in the cerebrum than cerebellum, and occur more frequently on the left than right side. In addition to the sanguine effusion there is generally considerable congestion of the pia mater, of the venous sinuses, or of the substance of the brain itself.

In describing the lesions of meningeal apoplexy, we shall confine our remarks to the effusion which occurs into the cavity of the arachnoid, this being, as we have already remarked, by far the most frequent form of the disease.

The appearances presented by the cavity of the arachnoid into which the effusion has taken place vary greatly in different cases, according to the age of the child, the quantity of the hemorrhage, and the period of time which may have elapsed between the accident and the death of the patient. It is very uncommon to find pure liquid blood, though this has been met with. In the case published by Dr. Dulles (*loc. cit.*), which occurred at the age of six months, the lateral ventricles were filled completely with firm and partly organized clots; a large firm clot filled the third ventricle. The surrounding brain tissue was deeply stained, but was not lacerated. In most instances, there is a bloody serum mixed with thin, reddish coagula, contained in a soft and very delicate membrane lining the internal surface of the arachnoid. Sometimes the effusion is thin, limpid, and more or less yellowish in color, while at other times it is thick and brownish, or chocolate-colored. In some rare cases it is perfectly transparent and colorless. The fluid, in whatever state it exists, appears to be the result of transformations undergone by the effused blood. The solid portion of the blood or clot is found either in the condition of more or less recent coagula, or changed into false membranes, which sometimes resemble very closely the arachnoid itself, and sometimes a true fibrous membrane. The coagula are found in the form of thin membranes, varying between one or two lines in thickness, and an inch and a half or two inches in size. They are thickest generally in the centre, where they measure between a fifth of a line and two lines, and are brownish or greenish in color, and of variable consistence, according to their age. These coagula may exist upon any portion of the brain, but, according to MM. Rilliet and Barthez, are most frequently met with upon its convex surface.

The coagula just referred to undergo in some instances a curious change,

of which we shall give a short description. In the course of time the fibrinous portions of the blood are deposited upon the internal surfaces of the cavity of the arachnoid, in the form of a new membrane. When death occurs soon after the onset of the attack, the parietal layer of the arachnoid is found to be completely lined with this membraniform production, whilst the visceral or cerebral layer is covered by it only in certain points. When the case has lasted a longer time, on the contrary, the visceral as well as the parietal layer of the arachnoid may be covered with the new production, and when this happens there is formed a true sac or cyst, destitute of opening, which lines the whole interior of the arachnoid, and contains within its cavity bloody serum and coagula. At first this new membrane is reddish in color, elastic, and of a stronger texture than might be supposed from its apparent thinness and softness. Its thickness is generally about a tenth of a line. At a later period the walls of the cyst become so thin and transparent that they have been mistaken for the arachnoid itself. They differ, however, from the latter in being rather less transparent and thin, and particularly in the circumstance of presenting numerous vascular arborizations. When death occurs at this stage, which M. Legendre (whose description we chiefly follow) calls the second period, or that of complete organization of the cyst, the external surface of the latter is found to adhere intimately to the parietal portion of the arachnoid membrane, by very delicate cellular tissue, though not with so much force but that it may be detached by traction. The internal portion of the new membrane, on the contrary, which is lubricated by the serosity of the arachnoid tissue, is very slightly adherent to the layer of that membrane covering the brain.

So long as the cyst formed by the new membrane, or, as it is called by MM. Rilliet and Barthez, the *pseudo-arachnoid* membrane, contains an amount of fluid sufficient to keep its surfaces separated, its cavity is single. When, on the contrary, the walls of the cyst have come into contact, either because of the partial absorption of the contained fluid, or because the fluid has accumulated at the lowest points, or wherever there is the least resistance, the cavity becomes multilocular in consequence of the cohesion of its walls at certain points.

The size of the cyst varies exceedingly. Sometimes it covers the greater part of the convex surface of one hemisphere, sometimes the whole, while in other instances it extends to the base, forming in that case a nearly complete shell for the whole brain. The quantity of fluid varies in different cases. Sometimes it amounts only to a few large spoonfuls; in others, to one or two, or eight or nine ounces; in one case observed by MM. Rilliet and Barthez there was upwards of a pint on each side, or more than a quart in all. In most instances the hemorrhage occurs into both halves of the arachnoid membrane, so that there is a cyst for each hemisphere. More rarely it occurs only on one side.

In the second stage, and when the effusion is very large, which rarely happens except in young children, and prior to ossification of the fontanelles or sutures, the lesion constitutes a form of chronic external hydrocephalus, and the symptoms are such as will be detailed under the head

of this latter disease. The vault of the cranium is enlarged by the unnatural prominence of the frontal and parietal bones; the sutures are more open than usual, and the anterior fontanelle is distended and protuberant. When the effusion occurs thus early in life before complete ossification of the skull, the brain does not appear compressed or flattened, as it does when the disease occurs at a later period.

The visceral portion of the arachnoid is often thickened, opaque, and more resisting than natural. The pia mater is frequently infiltrated with a good deal of serosity, which sometimes has a gelatinous appearance. When death has occurred in the first stage of the disease, the brain usually presents signs of hyperæmia. The veins on the surface of the hemispheres are enlarged, the cortical substance is of a bright rose-gray color, and the medullary portion is dotted over with drops of blood. Sometimes the cellular substance beneath the arachnoid is slightly infiltrated with serosity, at other times not. The ventricles contain a very small quantity of fluid.

The exact anatomical cause of the cerebral hemorrhage in children is still subject to some doubt. It appears probable that it usually results from intense determination of blood to the head, or from extreme passive congestion, which lead to the rupture of vessels so minute as to escape notice, or possibly in some cases to the transudation of blood through the capillary walls without actual rupture. We are not aware that any careful microscopic examination has yet been made of the condition of the walls of the vessels in such cases. In some rare instances, however, as in one witnessed by M. Legendre, the effusion is the result of the rupture of a vessel of some size. In the case observed by him, death took place in twelve hours from the attack, and the left hemisphere was found covered with a layer of coagulated blood, which had escaped from a ruptured vein. (*Biblioth. du Méd. Prat.*, t. vi, p. 192.)

SYMPTOMS; DURATION.—The symptoms of *hemorrhage into the substance of the brain* in the child are, as a general rule, extremely obscure and uncertain, though in some few cases that have been observed they were as characteristic as those which occur in adults. In obscure cases the chief symptoms that have been noticed were restlessness, delirium, headache, violent fever, grinding of the teeth, and, after a time, complete abolition of the intelligence, fixity of the eyes, invariable dilatation of the pupils, stertorous respiration, and general insensibility. Of three cases reported by M. Valleix (*Clinique des Mal. des Enf.*), the nature of the disorder was easily diagnosticated in one by the existence of complete hemiplegia, while in the two others the only marked symptom was entire immobility. The only certain symptom of the disease, therefore, would be a sudden attack of hemiplegia, either as the primary symptom, or following coma or convulsions, and lasting for at least several days. An attack of general paralysis would not be by any means so certain, as this may exist in several other diseases of childhood.

In a case which came under our charge, we believe the attack to have been one of apoplexy of this kind.

CASE.—A girl, two years and a half old, apparently in the enjoyment of excellent health, was suddenly, and without ascertainable cause, attacked with violent general convulsions and entire insensibility, which lasted with very slight remissions of the convulsive movements, but without any return of consciousness, for twelve hours. At the end of that time the convulsions ceased entirely, and she very soon regained her consciousness, remaining merely peevish and languid. She was, however, completely hemiplegic on the left side, so that she could neither rise in bed nor turn towards the right side. The paralysis diminished rapidly, but regularly, so that at the end of three days she could sit up in bed, and in a few weeks was perfectly well. This child remained well, with the exception of rather unusual excitability, and some peevishness of temper, for three years, when she died of scarlet fever. No autopsy could be made.

The obscurity which exists in these cases will be clearly understood by any one who will read two examples given by Dr. West (*loc. cit.*, p. 1062).

With a short quotation from the work of MM. Rilliet and Barthez we shall pass on to the subject of meningeal apoplexy. These authors remark (*op. cit.*, t. ii, p. 54), in speaking of this affection, that “cerebral symptoms have been observed to exist, but of so unusual a character, and so different from what have been assigned by writers to apoplexy, that they could not lead to a diagnosis of the disease.”

We shall describe the symptoms of the *meningeal form* of hemorrhage under two heads: first, as they present themselves in the acute, and, second, as they occur in the chronic or second stage of the affection.

Unfortunately the symptoms of the acute or first stage are not much more certain and distinct than those of cerebral hemorrhage. The disease may begin with fever and some convulsive movements, or, as happened in a case reported by M. Valleix, with violent general convulsions. Vomiting sometimes occurs at the beginning, but is usually very slight. It is difficult to know whether headache exists or not at the early age at which this disease commonly occurs. The convulsive movements generally affect particularly the eyes, and are followed by some degree of strabismus. The appetite is lost from the first; the thirst is moderate; there is no constipation. Soon after the symptoms just described, permanent contractions of the hands and feet appear, which are followed by attacks of tonic or clonic convulsions, during which sensibility and intelligence are abolished. Between the attacks of convulsions there is somnolence, which though slight at first, becomes more marked as the case goes on. The attacks of convulsions become more and more frequent as the case progresses, until at last they are nearly constant. The tonic convulsions affect both the limbs and trunk, but particularly the former, whilst the clonic spasms occupy sometimes one side of the body, sometimes the upper extremity alone, and at other times the whole body, but even then are usually stronger on one side than on the other. Paralysis is rarely noticed in the disease; it occurred only in one out of nine cases observed by M. Legendre, and in one out of seventeen observed by MM. Rilliet and Barthez.

Dr. West remarks (p. 1061): “The absence of paralytic symptoms, however, is not the sole cause of the obscurity of these cases, but the indi-

cations of cerebral disturbance, by which they are attended, vary greatly in kind as in degree. The sudden occurrence of violent convulsions and their frequent return, alternating with spasmodic contraction of the fingers and toes in the intervals, appear to be the most frequent indications of the effusion of blood upon the surface of the brain. I need not say, however, that such symptoms, taken alone, would by no means justify you in inferring that an effusion had taken place." Dr. West adverts particularly to the fact that apoplexy in the child is especially apt to occur in those who are weakly and feeble, and gives to this form of the disease the appellation of the cachectic form of cerebral hemorrhage.

It must be remembered that in cases of valvular disease of the heart, embolism of one of the cerebral arteries may occur from the detachment of a fragment of a vegetation. We have observed this accident in a young girl, eight years of age, where the symptoms which marked the occurrence of the embolism were brief unconsciousness, followed by complete left-sided sudden hemiplegia. The diagnosis may be made in these rare cases by the detection of the physical signs of organic disease of the heart, and by the less severe cerebral symptoms which, as a rule, attend it. The hemiplegia which follows embolism may be partial or complete. In our cases it persisted, with contraction of the paralyzed members, until death.

The chronic form presents most of the symptoms which exist in acquired chronic hydrocephalus from serous effusion into the ventricles. The cranium is very large in proportion to the face; the sutures are not ossified; there is strabismus, with dilatation of the pupils; the sense of sight is generally but not always retained; the face loses its expression; if the child was old enough at the moment of the attack to show signs of intelligence, the latter are found to diminish rather than increase, and sometimes they are lost entirely, as the size of the head augments; and the child is apt to utter loud cries, particularly during the night. The cutaneous sensibility is in general neither lost nor diminished. The power of motion usually remains, though it was entirely lost in one case. The appetite and thirst persist.

The *duration* of cerebral apoplexy is very irregular. In one case quoted by MM. Rilliet and Barthez, it was a quarter of an hour; in another, an hour; in a third, forty-eight days; and in one reported by M. Valleix, in a very young infant, recovery was nearly perfect in a little less than two months, when the child was seized with pneumonia and died.

The duration of meningeal apoplexy is also irregular. According to M. Legendre, all the recent cases seen by him in the Children's Hospital, died in from eight to twelve days, apparently rather from intercurrent diseases than from the primary affection itself, whilst cases occurring in subjects placed in better hygienic conditions, and not attacked with intercurrent affections, passed into the second or hydrocephalic stage of the disease. The second stage lasted, according to the same author, in the four cases which he witnessed, from eight to thirty months, and then death was the result, not of cerebral symptoms, but of complications affecting the thoracic organs.

DIAGNOSIS.—The diagnosis of cerebral hemorrhage is, as we have already stated, very difficult, unless hemiplegia exist. When the case commences, as it often does, with convulsions or with inflammatory symptoms, it is often impossible to distinguish it from acute or tubercular disease of the brain.

The diagnosis of meningeal hemorrhage is also very often extremely difficult. Not unfrequently it occurs in the course of other diseases, and is then entirely latent. In acute, primary cases, the most important and distinctive symptoms are the early age of the subjects, between one and three years generally; the violent fever from the commencement, marked by full, frequent, and *regular* pulse; the absence of constipation; the frequency of the convulsive attacks, and particularly the permanent contraction with rigidity of the feet and hands.

The diagnosis between the form of hydrocephalus which follows meningeal apoplexy, and ventricular serous hydrocephalus, is exceedingly obscure. The only circumstances which seem to have any real value are the acute commencement of the disease with the symptoms above detailed, and the early age of the patient. MM. Rilliet and Barthez state that they have never known a child of two years old, or younger, to die of ventricular serous hydrocephalus from tumors, whether tubercular or not, of the brain; in all such cases the effusion has been the result of a hemorrhage.

PROGNOSIS.—The prognosis of both forms of the disease is very grave, but it is impossible to ascertain it with any certainty, so long as the symptomatology of the two affections is so obscure as we have found it to be. That cerebral hemorrhage is susceptible of cure, however, is proved by the case reported by M. Valleix, already referred to, in which the child had nearly recovered, when it was seized with another disease which destroyed it. Recovery from meningeal apoplexy is certainly extremely rare; we believe, however, that we have met with at least one case in which this affection terminated favorably.

TREATMENT.—The treatment must depend on the diagnosis and the special character of the symptoms in each case. In a sudden and severe attack, occurring in a strong and hearty child, in which the symptoms of congestion of the brain are strongly marked, and where we are not yet certain that actual hemorrhage has taken place, we should immediately resort to a general or local bloodletting. It was formerly customary to employ venesection in all such cases, but we believe that equal relief can be obtained by freely cupping or leeching the back of the neck.

When, however, we have every reason to believe that blood has been effused, either in the membranes or into the substance of the brain, it is evident that bloodletting can produce but little effect, and that only in reducing the general fulness of the cerebral vessels. In such cases we should certainly limit ourselves to the application of a few cut cups or leeches to the nucha, if any blood at all is to be withdrawn.

It must be further remarked, however, that in many cases of cerebral or meningeal apoplexy, depletion in any form is entirely contra-indicated; since, as has already been stated, the effusion of blood occurs frequently

in feeble and weakly children, and either in the course of some acute or chronic disease, or as a consequence of previous diseases which have exhausted the forces of the constitution and induced a state of dyscrasia and diffuence of the blood. In such cases as these it is clear that the only chance of recovery must depend upon maintaining the system in perfect rest, avoiding any perturbation or depressing measures, and endeavoring to support the vital powers till reaction occurs, and an opportunity for the absorption of the effused blood is secured.

If there is undue heat of the head, cold applications should be immediately made to it, either by wet cloths, the ice bladder, or by cold affusion. At the same time, if there is reason to suspect the presence of undigested or irritating matters in the alimentary canal, a moderately active purgative dose should be administered.

Counter-irritants are always useful adjuvants to the remedies already mentioned. They should consist at first of mustard plasters applied to the extremities, and shifted from place to place. When the severe symptoms do not yield after some hours, it may be well to apply a blister to the nape of the neck.

The diet must be very strict, and should consist only of barley or arrow-root-water, for a few days.

The temperature of the room should be kept cool; and the child should be placed with the head and trunk somewhat elevated, and kept profoundly quiet.

For the paralysis which follows apoplexy in children, we believe that the most important, and indeed the only treatment necessary, is attention to the general health of the patient, in order to give to nature time and opportunity to effect the absorption of the clot which has been thrown out into the substance of the brain, or into the cavity of the arachnoid membrane. This process may, however, be aided and hastened by the prolonged administration of iodide of potassium with the iodide of iron. In cases of meningeal apoplexy, when the disease assumes the chronic form, occasioning the kind of hydrocephalus we have described, there is little more to be done than to attend to the general health of the child, and to endeavor to promote absorption of the fluid by the internal administration of diuretics, and the preparations of iodine. It has been proposed also to get rid of the fluid by tapping, as has been done in congenital hydrocephalus, and it is indeed in cases of the form we are now considering, when the fluid is entirely external to the brain, and where no malformation or organic disease of the brain exists, that this operation has been found most successful. (See treatment of chronic hydrocephalus.)

ARTICLE V.

CHRONIC HYDROCEPHALUS.

THIS term is applied to an affection characterized by an excessive accumulation of serous fluid, either within the ventricles of the brain or the sac of the arachnoid.

The names *internal* and *external* have also been applied to it, in accordance with the position of the fluid: the former being given to those cases where the ventricles are the seat of the morbid collection, and the latter indicating that the fluid has accumulated in the cavity of the arachnoid and consequently surrounds the exterior of the brain. Chronic hydrocephalus may either be *congenital* or *acquired*, the latter variety presenting the most interest in a practical point of view, since congenital hydrocephalus is usually associated with some malformation of the brain which renders extra-uterine life almost impossible.

In either form it is a comparatively rare disease in this city. Our more recent experience, however, has furnished us with numerous opportunities for studying its symptoms, treatment and pathology.

MORBID APPEARANCES.—There are indeed few diseases in which it is of more importance to correctly establish the exact nature of the morbid process and the resulting lesions, since, as we shall see in a later part of this discussion, questions of the utmost practical value hinge upon the determination.

Internal Hydrocephalus.—In this condition the amount of fluid is often very large, and varies from half a pint or a pint, to even as much as a gallon. Trousseau mentions a case where the fluid weighed 30 pounds, and Frank one in which it weighed 50 pounds. The formation of this accumulation being gradual, the cavities of the brain accommodate themselves to it, the ventricles become distended, and the communications between their cavities are all enlarged; and occasionally the septum lucidum is perforated. This distension is usually most marked in the lateral ventricles. The hemispheres of the brain yield to the pressure of the increasing collection in the ventricles; their convolutions are unfolded and flattened, so that the interval between them is only marked by a sinuous shallow groove, and the hemispheres are so thinned out as to form a layer not exceeding a few lines in thickness. It is not unusual, however, even when the distension of the brain has proceeded to this extreme degree, to be able to trace the cineritious and white layers, preserving their normal relations. The consistence of the expanded brain-substance varies in different cases; usually, however, it remains normal, or is even increased, though in some cases it has been found so soft as to tear upon the slightest traction. The structures at the base of the brain present the same changes in consistence.

One of the most important questions in this relation, as bearing upon the causation of the affection, concerns the condition of the lining membrane of the ventricles.

The analogy of all other serous membranes would lead us to infer that in those cases where no mechanical obstruction to the circulation exists, such as a tubercular tumor pressing upon the sinuses of the brain, we should look for the cause of the serous accumulation in a morbid state of the lining membrane of the ventricles. This view is fully confirmed by the study of fatal cases of internal hydrocephalus, since in many cases this membrane is found much thickened, and either softened or roughened and granular. The granular condition of the membrane presents many degrees:

in some cases it is merely a slight irregularity of the surface, while in others there is an unevenness as marked as that of shagreen, or even a formation of granules, which, at times, measure one-third of an inch in diameter, or even become distinctly pedunculated.

Occasionally, a false membrane is found lining one or both ventricles, as the result of the chronic inflammation of the lining membrane of these cavities.

Even when the symptoms of hydrocephalus have not appeared until some time after birth, the brain may be found to present positive evidences of congenital malformation, in the retarded development of some of the structures at its base.

The veins of Galen and sinuses of the dura mater are usually found in a healthy state, with their calibres quite free; a fact which is of importance in considering the mode of production of internal hydrocephalus.

In *external* hydrocephalus, the collection of fluid occurs in the sac of the arachnoid, or in a pseudo-cyst resulting from the transformation of a blood-clot, as described in our remarks on meningeal apoplexy; the brain is separated from the cranial vault and compressed against the base of the skull, as the lung is forced back against the spinal column by the fluid of hydrothorax.

The superior cerebral veins, passing from the surface of the brain to the longitudinal sinus, traverse the fluid, and at times are so much stretched as to raise the surface of the brain into points.

Excepting in cases, however, where the disease is congenital and coincident with some original malformation of the brain, there is no absolute diminution in the size of this organ.

The character of the fluid varies considerably in different cases, and probably depends to a great extent upon the cause.

In an analysis by Spengler of the fluid evacuated in a case of hydrocephalus by puncture, the fluid was clear and colorless; specific gravity 1010, of acid reaction, and contained no albumen. It also contained chlorides and phosphates of soda and potassa, but no sulphates. It appears, therefore, in such cases as this, that the fluid is not the result of inflammation, but rather due to a passive dropsy. It is, we believe, especially in cases of external hydrocephalus, where the fluid results from the transformation of a sanguineous effusion, that it possesses these characters.

On the other hand, the fluid frequently contains a large amount of organic matter, and closely resembles the effusion in pleurisy or pericarditis. Thus, in a case reported by Battersby, which was tapped eight times, the fluid always contained varying, and sometimes very large, proportions of albumen.

Causes of Internal Hydrocephalus.—The opinions of the highest authorities and most experienced observers still differ widely upon this important point.

We have alluded to the fact that not unfrequently the brain is found to present evidences of congenital malformation, and this fact has led to the opinion that internal hydrocephalus is almost invariably the effect of arrested development of the brain.

Rilliet and Barthez place the effusion in this affection in the class of passive dropsies, and express their belief that most frequently the cause of internal hydrocephalus is to be found in compression of the veins of Galen or ventricular veins, caused by the development of a tumor in the cranial cavity, and usually in the lobes of the cerebrum.

The unfavorable influence which either of these views would have upon the prognosis and treatment of this disease is of course evident.

On the other hand, however, the opinion is advanced that the starting-point of internal hydrocephalus is, in fact, a morbid condition of the lining membrane of the ventricles.

We have briefly described the appearances of this membrane which have now been observed in numerous well authenticated cases of internal hydrocephalus, and which plainly indicate the pre-existence of a chronic inflammation so that we are led to believe that in a certain number of cases, at least, the effusion is due to a slow inflammatory action in the lining membrane of the ventricles. Those cases in which these appearances have been found associated with retarded development of the brain, may be readily explained upon the supposition that the inflammation has been excited at a more or less advanced period of intra-uterine life, and that the resulting effusion has so compressed the structures at the base of the brain as to prevent their normal development. We may add that many eminent authorities, as Trousseau, now adhere to this view.

In a very interesting case of this form which we have lately had under observation, the mother, a very intelligent and healthy woman, had quite a severe fall about the fourth month of pregnancy. In addition to this, however, she lost a favorite brother-in-law from a violent attack of cerebrospinal meningitis about the same time. She nursed him constantly during his illness, and was very deeply impressed with the unnatural appearance of his face and head, which had been shaved. This became so fixed an impression that more than once, between the time of his death and the birth of her child, she said it would not be strange if something were to prove wrong about the baby's head. We attributed the hydrocephalus which did actually develop in her infant to inflammatory changes, perhaps induced by the fall; but the powerful maternal impression above mentioned certainly constituted a curious coincidence.

In cases, however, where the effusion into the ventricles depends upon the development of a tumor in the cranial cavity, the growth will usually be found to occupy the cerebral lobes in such a manner as to compress the veins of Galen, which pass along the under surface of the corpus callosum, and are indeed the only true ventricular veins.

The causes of *external hydrocephalus* are perhaps less obscure and uncertain than those of the internal form.

In some cases, the effusion in the sac of the arachnoid is evidently due to a rupture of some portion of a brain distended by accumulation of fluid in the ventricles, and hence is merely a sequel of internal hydrocephalus.

According to the able investigations of Legendre, and Rilliet and Barthez, one of the most frequent causes of external hydrocephalus is hemor-

rhage into the arachnoid space; the effused blood undergoing changes which result in the presence of large quantities of clear fluid, as described at length in our remarks on meningeal apoplexy. We have alluded to the fact that in many cases of external hydrocephalus the diminution in size of the brain is comparative rather than real; but there are instances where this form of the disease is found associated with malformation of the brain, which appears as a small, misshapen mass, pressed against the anterior part of the base of the skull. In such cases, it appears as though the fluid were poured out to fill up the vacuum between the skull and atrophied brain. It is also possible that these conditions may be produced by the occurrence of hemorrhage into the arachnoid space during intra-uterine life, and before the brain had attained its normal development.

SYMPTOMS; PHYSICAL APPEARANCE.—The unusual size of the head is one of the most striking symptoms of hydrocephalus. In many cases associated with atrophy or retarded development of the bones of the face and the rest of the body, this enlargement appears even more monstrous than it in reality is. The diameters of the cranium are, however, very much enlarged; cases being on record in which at the age of a few weeks the circumference of the head has been twenty-three inches, or even more.

The increase in the size of the head is not, however, invariably the earliest sign of the disease, being frequently preceded by marked symptoms of nervous disturbance, or of impaired nutrition.

The bones of the cranial vault which contribute to this enlargement are the frontal, the parietals, the occipital, and the squamous portion of the temporals. When the disease makes its appearance before the ossification of the sutures and fontanelles has been completed, the gradual increase of the fluid separates these bones more and more widely. The occipital bone thus is pushed backwards, the parietals outwards and backwards, the frontal upwards and forwards. The increase of the size of the head is thus effected by the widening of the sagittal and coronal sutures, and by enlargement of the anterior fontanelle.

The displacement of the frontal bone gives rise to a marked prominence of the forehead, which overhangs the diminutive features; while at the same time the pressure of the fluid depresses its orbital plate into an oblique position, contracts the orbital space, and gives rise to the characteristic appearance of the eye, the globe being prominent but directed downwards so as to be buried below the lower eyelid, which conceals almost the entire cornea.

The membrane which covers in the enlarged sutures is often distended and prominent, or remains on the normal level. A distinct sense of fluctuation is readily obtained by palpating one of these spaces, and in some cases, principally in young infants, and where the collection is very large, the head is absolutely translucent. When life is prolonged, and the disease arrested, the ossification of the cranial vault is effected by the development of numerous supernumerary bones, or ossa triquetra, in the membranous spaces. These little bones are consequently found in the

largest numbers in the coronal and sagittal sutures, where the deficiency is greatest and most wide. When, on the other hand, the disease does not begin until the sutures have united and the fontanelles ossified, it is rare for the head to attain any very large size. In a few cases, however, occurring in children of even nine years of age, the sutures have reopened under the continuous pressure, and the bones have been found separated as much as half an inch.

More usually, however, in such cases, the pressure seems to expend itself in thinning the cranial bones, which become reduced to mere shells of light, fragile compact bone. Occasionally, so far from inducing thinning of the bones, actual hypertrophy occurs, and the bones of the cranial vault acquire an unusual thickness, and at the same time are dense and indurated.

The early symptoms of the disease vary much. When it is congenital, there are nearly always evidences of cerebral disturbance either from the date of birth, or appearing within a few days. These symptoms are occasionally slight, consisting merely in an unnatural expression, with oscillation of the eyes or strabismus; or, on the other hand, there may be attacks of convulsions frequently repeated.

These symptoms speedily become associated with enlargement of the head and the characteristic alteration of physiognomy. When the disease is strictly acquired, the early symptoms are even more varied. In one set of cases they are those of hemorrhage into the arachnoid; in another the evidences of inflammation of the serous lining of the ventricles, of more or less acute character, are present; whilst in numerous cases the only symptoms which precede the enlargement of the head are those of failing nutrition.

Usually the aspect of children suffering with this affection is tranquil, or they may even present a certain unnatural gravity and apathy of expression.

Cerebral Symptoms.—At times the intelligence of the child, though perhaps poorly developed, remains intact, and there is no marked cerebral disturbance.

In other cases, however, the advance of the disease is attended with a gradual failure of the intelligence, and impairment of the special senses, and especially of vision.

In addition to the displacement of the globes of the eyes and alterations in the pupils already mentioned, the accumulation of fluid rapidly causes obstruction to the return of venous blood through the sinuses, so that even at an early stage ophthalmoscopic examination shows marked changes in the fundus of the eyes. These consist in increase in the number and size of the veins of the retina, with later serous infiltration or even atrophy of the optic papilla.

The nervous symptoms are at times much more marked; and there may be frequently recurring convulsive attacks, or, as West mentions having seen in several cases, spasmodic attacks of difficult breathing, with a crowing sound in inspiration (*laryngismus stridulus*).

According to Rilliet and Barthez, the common sensibility of the surface

is often impaired; and there may be more or less complete paralysis, or contraction with rigidity of the extremities.

It is, of course, difficult to estimate the amount of suffering experienced by the little patients; ordinarily it does not appear great, and indeed in some cases it has seemed chiefly due to the opposition offered by the cranial walls to the distension of the brain.

In one case of MM. Rilliet and Barthez, the development of acute pain coincided with the ossification of the fontanelles.

The *general condition* of children suffering with chronic hydrocephalus varies greatly.

In some cases they preserve their appetite and digestion, and appear well nourished and strong to a late period in the attack; but more frequently they present marked evidences of impairment of nutrition.

The appetite may indeed remain, but the child loses both flesh and strength; the bowels are irregular; usually constipated, but alternating with temporary attacks of diarrhœa.

In the majority of cases, perhaps, these symptoms are not sufficiently pronounced to establish the character of the attack, until the increasing size of the head becomes manifest, and the child acquires the distinctive physiognomy of hydrocephalus. Even after marked enlargement of the head has occurred, however, the advance of the case is far from being uniform. In almost every instance there are pauses of the most variable frequency and duration, during which the child seems free from pain, improves in general condition, and the development of the head is temporarily arrested.

Death is frequently directly induced by some intercurrent affection, wholly unconnected with the disease of the brain; while, in other cases, it immediately follows a violent attack of convulsions, or is preceded by symptoms of an acute exacerbation of the cerebral disorder. In some cases, also, the patients sink into a condition of atrophy, and die worn out by the protracted suffering and malnutrition.

DIAGNOSIS.—During the early stage of the disorder, if the nervous symptoms are slight, consisting merely in occasional attacks of heat of the head, attended with pulsation or tension of the anterior fontanelle, and restlessness, and crying, the diagnosis must remain uncertain. After the enlargement of the head has progressed to any considerable degree, the expression of the little patient, taken in conjunction with the other symptoms, is usually perfectly characteristic and conclusive.

The morbid condition with which it is most likely to be confounded, is rickets of the skull. In fact, in some cases, the enlargement of the head, which results from these two affections, is quite identical. Usually, however, this is not the case; and the hypertrophy of the rachitic bones takes place irregularly, so that the skull acquires a square instead of a rounded form; the orbital plates of the frontal bones are not displaced; so that, although the forehead may be large and overhanging, the axes of the eyes are not disturbed; the fontanelles are not widely open, prominent, or distended; and, finally, of course, fluctuation on palpation is never present. In addition to this, the evidences of rickets in other portions of the body,

and the peculiar symptoms of that affection, as detailed in the article devoted to its consideration, nearly always enable the diagnosis to be readily made.

We have already mentioned the changes which ophthalmoscopic examination shows in the retina in this disease, and as a similar examination reveals no lesion whatever in cases of rachitic enlargement of the head, it is evident that the use of the ophthalmoscope may be of material aid in establishing the diagnosis between these affections, which is, despite all the points of distinction above referred to, obscure and difficult in some few cases.

In doubtful cases assistance may possibly also be derived from cerebral auscultation; the presence of a bruit over the anterior fontanelle being thought by some authors to be a valuable indication of the rachitic nature of the enlargement of the skull. The significance of this cephalic bruit is, however, so much disputed, that it is at present impossible to assign any definite value to it.

The only other pathological condition with which chronic hydrocephalus is apt to be confounded, is hypertrophy of the brain, an extremely rare affection, due to an increase of the interstitial connective tissue of the brain, the so-called neuroglia.

In hypertrophy of the brain, however, the symptoms do not usually appear so early as in chronic hydrocephalus, nor is the cerebral disturbance so marked as in the latter affection. The enlargement of the head, also, which is the most characteristic feature of both conditions, is not so great in hypertrophy of the brain, and, instead of being uniform and assuming a rounded form as in hydrocephalus, occurs especially at the occiput. There is, further, no depression of the orbital plates of the frontal bones in hypertrophy of the brain, so that the axes of the eyes are not disturbed, and the globes are not displaced in the way we have already described as so characteristic of hydrocephalus.

Finally, the sutures are not so widely open, nor the fontanelles tense and prominent as in hydrocephalus; and, of course, the fluctuation which can be detected on palpation in some cases of this latter disease is never present.

PROGNOSIS.—Chronic hydrocephalus still ranks among the most fatal diseases; so much so that Rilliet and Barthez—who, however, attribute its production usually to the presence of a tumor in the brain—express their belief that it is invariably fatal. Indeed, it must be borne in mind that in many cases treatment must necessarily fail from the coëxistence of some extensive congenital malformation of the brain. We should suspect the presence of this complication when there is serious disturbance of the nervous system, such as paralysis, or frequent and apparently causeless convulsions. Unfortunately, however, these hopeless cases cannot always be distinguished.

While the prospect of a complete cure is very slight, it must be remembered that even a high degree of hydrocephalus is not incompatible with considerable prolongation of life. Thus cases have been known to attain the age of 15, 20 or even 25 years, with the maintenance of a fair degree of mental and bodily power.

The prognosis in cases of external hydrocephalus, especially when of acute origin, is much less unfavorable than when the effusion takes place into the ventricles.

Whatever be the seat of the effusion, however, and the size of the head, the case must not be regarded as hopeless and beyond reach of remedial measures, so long as the functions of the brain are well performed, since there are well authenticated cases of complete recovery from chronic hydrocephalus, even when congenital.

In a case we have recently had under observation, the symptoms of hydrocephalus, apparently of congenital origin, advanced rapidly up to the age of three and one-half years, at which time the enlargement of the head was arrested, the mental and physical powers of the child developed regularly, and at the age of seven years it seemed probable that complete recovery would result.

TREATMENT.—It must be sufficiently evident, from the previous consideration of this affection, that there are numerous cases in which all treatment must prove unavailing, from the serious organic disease of the brain which accompanies it. Under any circumstances, however, the nature of the treatment and its efficiency will be much influenced by the early stage at which it is instituted.

In regard to the utility of various special remedies, also, there is the greatest diversity of opinion; and, indeed, there is no plan of treatment which possesses so much evidence in its favor as that originally proposed by Professor Gölis, of Vienna.

If the disease be in its incipience, and the constitution and hereditary tendencies of the child free from taint, this distinguished physician recommends that the head should be shaved, and one or two drachms of dilute mild mercurial ointment rubbed daily into its scalp. While this treatment is being carried out, the head should be constantly protected by a flannel cap. At the same time, calomel should be given in doses of one-sixth to one-fourth of a grain twice daily, unless it irritate the bowels, when the inunction alone should be continued.

If after pursuing this treatment, conjoined with the most careful attention to diet and all hygienic precautions, for five or six weeks, there is marked improvement in the condition of the child, the mercurials may be gradually discontinued.

The iodide of potassium has been highly recommended as a substitute for the mercurials above mentioned, and several cases of apparent recovery under its use are on record. It should be given in large doses, and for a considerable length of time. Trousseau, who recommends its use, joins to its internal administration the external application to the head of lotions containing iodine.

Should the disease remain uninfluenced at the end of this time, it is proper to add to the treatment diuretics and counter-irritants, in the form of issues in the back of the neck, which may be kept open for several weeks. Dr. West recommends the frequent application of blisters as a substitute for the use of issues.

During the employment of this or any other mode of treatment, it will

be occasionally necessary to have recourse to antiphlogistic remedies, to subdue the exacerbations of heat and restlessness which occur more or less frequently, and threaten the development of an acute inflammatory condition. Nor should we fail to pay attention to the proper performance of all the functions; to the maintenance of the appetite and digestion by the use of tonics; and in case of the existence of a scrofulous diathesis, to the administration of cod-liver oil, iodide of iron, etc.

When, despite the most careful employment of well directed measures, the disease is clearly advancing, it is worse than useless to persist in any plan of treatment which annoys or absolutely pains the doomed child; our only endeavor should then be to subdue any intercurrent disorder which might hasten the fatal result.

More than twenty-five years ago, the use of compression of the head, to prevent its yielding to the accumulating fluid, was urged by Barnard, and experience has shown it to be a valuable adjunct to other treatment, though it is inapplicable while any acute symptoms are present, and according to West is best adapted to cases of external hydrocephalus succeeding to hemorrhage into the arachnoid space.

M. Trousseau recommends the following mode of applying this pressure: Strips of adhesive plaster, about one-third inch wide, are passed from each mastoid process to the outer part of the orbit of the opposite side; from the nape of the neck along the longitudinal sinus to the root of the nose; across the whole head, intersecting at the vertex; and finally are kept securely in position by a strip passed thrice around the head, the ends of the previous strips being turned up over the first coil of this strip, and secured by the succeeding turns.

It becomes necessary to loosen these strips instantly, if any symptoms of compression of the brain develop themselves, since the increasing pressure of the accumulating fluid may produce irreparable injury to the base of the brain, or even, as happened to M. Trousseau, detach the ethmoidal bone from its connections.

The unfavorable results of all strictly medicinal treatment, impelled physicians, at an early date, to resort to active surgical interference in chronic hydrocephalus, by puncturing the cranium and evacuating the fluid.

The operation should be performed with a delicate trocar and canula, the puncture being made in the coronal suture, about an inch or an inch and a half from the longitudinal sinus,—and in a majority of cases, no evil consequences appear to follow the operation itself. Much difference of opinion still exists, however, as to its curative influence. From a rigid analysis of 56 reported cases in which this operation has been performed, Dr. West came to the conclusion that in only 4 had a permanent cure been effected. Other successful cases have been since reported, so that the operation must be recognized as at least a justifiable one in certain cases.

In a case of internal hydrocephalus upon which we operated recently, death followed the operation in less than 48 hours, and was unquestionably hastened by our interference. Examination showed, however, that the case was an utterly hopeless one, since the structures at the base of the brain were becoming disorganized by the pressure of the liquid, and

in addition the child, less than 3 years old, was the victim of miliary tuberculosis. On the other hand, we have recently met with a case of external hydrocephalus where a permanent cure was effected by the operation. In this interesting case, the operation was repeated three times, at intervals of 3 and 6 weeks respectively; 16, 8, and 6 ounces of limpid serum were removed; subsequently no tendency to reproduction of the effusion showed itself; the fontanelles closed rapidly; every function developed normally; and at the age of 4 years, over 3 years since the last operation, although the head is abnormally large, the disease is evidently arrested and the child is in excellent health.

The circumstances favorable to its performance are, therefore, when the hydrocephalus is external; or when internal, is due to previous inflammation of the lining membrane of the ventricles; when there is no reason to believe that the disease is congenital, and attended with arrested development of the brain; when, though the head may be very large and increasing in size, the cerebral functions are not seriously impaired; and finally, when the nutrition of the child is still good.

In cases of internal hydrocephalus the operation should never be performed until the treatment previously recommended has been faithfully tried without any influence on the advance of the disease. It involves great risks with but slight prospects of success. In external hydrocephalus, on the other hand, the operation holds out much more prospect of success, and is to be recommended when the symptoms of the disease do not yield satisfactorily to less radical methods of treatment.

Brainard, of Chicago, recommended the injection of solutions of iodine into the cranial cavity, after puncture and evacuation of the fluid. He employed this in at least two cases, without the development of any severe symptoms as a direct consequence of the treatment.

One of the cases died at the end of eight months; the other, at the date of the report, only thirty-five days after the operation, had shown no unfavorable symptoms.

He advises the use of an aqueous solution of iodine, in the proportion of one-third of a grain with one grain of iodide of potassium, to $\text{f}\overline{\text{3}}\text{j}$ of distilled water; of this from $\text{f}\overline{\text{3}}\text{j}$ to $\text{f}\overline{\text{3}}\text{j}$ may be injected; the strength of the solution and the amount injected being increased at subsequent punctures.

In one of his cases, twenty-one injections were practiced in the course of seven months.

Injections of this strength are usually followed by no symptoms of inflammation whatever; and this exemption has led to the employment of much stronger solutions.

Thus Dr. Tournesko, of Bucharest (quoted by Bouchut), injected $\text{f}\overline{\text{3}}\text{ij}$ of tr. iodine in $\text{f}\overline{\text{3}}\text{v}$ distilled water, immediately after having drawn off by puncture $\text{f}\overline{\text{3}}\text{xxiv}$ of serum. The operation was followed by slight febrile excitement; but, at the expiration of fifteen days, the child seemed in excellent health, the circumference of the head having diminished from $56\frac{1}{2}$ to 43 centimeters.

We have had no experience with the use of such injections in hydrocephalus, and should regard them as admissible only in cases of the internal form.

ARTICLE VI.

GENERAL CONVULSIONS, OR ECLAMPSIA.

GENERAL REMARKS.—The word convulsions is a generic term applied to different forms of spasmodic disease, very dissimilar from each other in many of their characters.

Writers make different classifications of convulsions according to their peculiar notions in regard to the nature and causes of these disorders. The best division is, it seems to us, one which arranges them according to their supposed causes, making three classes, *idiopathic* or *essential*, *sympathetic*, and *symptomatic* convulsions. The first two classes are unaccompanied by appreciable lesions of the nervous centres, while the third is called symptomatic, because it includes cases of convulsions which are the sign or symptom of an appreciable lesion of the cerebro-spinal axis, as for instance, those which occur in the course of meningitis, tubercular disease, hydrocephalus, apoplexy, etc. In idiopathic or essential convulsions, the cause of the attack acts directly upon the nervous centres, while in those to which the term sympathetic is applied, the cause lies in the influence or effect upon the brain or spinal marrow of disease of some other organ; to the latter class belong the convulsions which occur in the course of pneumonia, bronchitis, the eruptive fevers, etc.

We shall not pretend to give an accurate account of symptomatic convulsions in this article, as they have already been treated of under the head of the different organic diseases of the brain in the course of which they occur. We shall refer to them in the present article only so far as may be necessary to elucidate the pathology, diagnosis, prognosis, and treatment of idiopathic and sympathetic convulsions.

There is a form of eclampsia occurring in children which we shall describe separately, as it differs in many of its characters from ordinary convulsions. This is the disease known by the names of spasm of the glottis, thymic or Kopp's asthma, laryngismus stridulus, and eclampsia with suffocation.

DEFINITION; SYNONYMS; FREQUENCY.—By the term *convulsion* is meant a paroxysm of variable duration, usually attended with unconsciousness, and followed by stupor, and characterized by a primary involuntary tonic contraction followed by irregular clonic spasms of the affected muscles.

In general convulsions, to which the above definition especially applies, the entire system of voluntary muscles is usually affected; though, as will be described hereafter, the attack may be a complete and genuine one of eclampsia, and yet the convulsive movements be limited in their extent to a single group of muscles, or even a single muscle.

The only synonyms which it is necessary to mention are *epilepsia puerilis*, *insultus epilepticus*, and *eclampsia*. The latter term, eclampsia, is, we believe, preferable to any other, and we would gladly introduce it instead of convulsions, which is too general a term to express the form of disease under consideration.

The frequency of eclampsia is very great. During the two years 1879-80, 1316 children under fifteen years of age died in this city of convulsions; whilst, during the same time, 1695 died of infantile cholera, and 781 of pneumonia. It must be recollected, however, that a very large number of these cases of eclampsia ought, beyond doubt, to have been returned under other titles, as many of them must have been a mere result of organic disease of the cerebro-spinal axis, or of other acute local or general diseases.

PREDISPOSING CAUSES.—Essential and sympathetic convulsions are much the most frequent before the age of seven years, which is the case also in regard to symptomatic convulsions, though the latter often occur after the age mentioned. We have ourselves met with not less than 200 cases of convulsions, though we have preserved records of but 96. Of these the age is noted in 91 cases, 19 of which occurred in the first year, 26 in the second, 20 in the third and fourth, 23 between the fourth and ninth, and 3 between the ninth and thirteenth years of life. Dr. West (*op. cit.*, p. 42) states that according to the Fifth and Eighth Reports of the Registrar-General, the deaths from diseases of the nervous system in London, under one year of age, bore a proportion of 30.5 per cent. to the deaths from all causes; from the first to the third year, the proportion was 18.5 per cent.; from the third to the fifth year it was 17.6 per cent.; from the fifth to the tenth year, it was 15.1 per cent.; whilst from the tenth to the fifteenth year it was only 10.6 per cent., and the total above fifteen years was but 10.4 per cent. Again, to show the very great influence of age upon the predisposition to convulsions, Dr. West states that, within the first year, the deaths from convulsions constituted 73.3 per cent. of the total mortality from diseases of the nervous system; between the first and third years, the proportional mortality from convulsions to the total mortality from affections of the nervous system, was 24.9 per cent.; between the third and fifth years, it was 17.8 per cent.; between the fifth and tenth years, it was 9.9 per cent.; while between the tenth and fifteenth years it had fallen to 2.4 per cent.; and above fifteen years it was but 0.8 per cent.

It is generally stated that convulsions are more common in girls than boys. MM. Rilliet and Barthez found this to be the case in their private practice, whilst in the hospital, sympathetic and symptomatic convulsions were most frequent in boys. According to our experience, they have been almost equally frequent in the two sexes, since of 92 cases that we have seen in which the sex was recorded, 47 occurred in boys, and 45 in girls.

It has been generally supposed that a delicate and nervous constitution is a powerful predisposing cause to convulsive attacks. This has been denied, however, by several recent writers, whose observation is very careful and accurate. We are disposed to believe that it is not so much a feeble or delicate constitution that predisposes to convulsions, as it is one characterized by a highly susceptible, irritable, and nervous temperament, which often exists, in our opinion, in connection with a healthy and vigorous physical organization. Of 96 children in whom we have seen convulsive attacks, and in whom this point was noted, these occurred more than once in 13. Of the 13, nine presented every appearance of strong and vigorous health, with the exception that when laboring under any kind of sickness,

as dentition, indigestion, the fever accompanying simple angina, in two the invasion of measles, and in one that of erysipelas, they immediately became extremely restless and irritable, or heavy and drowsy, and at a very early period, and sometimes with very little warning, were seized with convulsions. In one, a well developed infant in its first year, the convulsions occurred every month or six weeks, without any appreciable cause. Three of the 13 were delicate; one was puny and feeble until after the completion of the first dentition, when it grew strong and hearty; one had had an apoplectic attack when an infant which had caused partial loss of power of one side; and the third was very weak at birth, then grew stronger, and died in its second year of hydrocephalus following scarlet fever. The number of convulsions varied in the different subjects. In 1 there were five different attacks, in another four, in 4 there were three, and in 5, two.

In two the attacks were very numerous, recurring frequently, and from very slight causes, or without any appreciable cause. They all recovered but two, and are still living. Of the 11 now living at various ages, all but one are free from anything like epilepsy, and that one, though liable during three years to attacks of an epileptiform character, became gradually less and less subject to the seizures, and has now been for several years perfectly well in all respects.

We have another patient, a boy, whose case is not included amongst the above, now five years old, who has had ten different attacks of convulsions. These attacks were all produced by some disturbance of his health. Several of them have occurred at the outset of a febrile reaction caused by a simple catarrh of the upper air-passages,—the convulsions ushering in the catarrh just as they sometimes do an attack of measles or scarlet fever. On other occasions, the seizure has evidently been the result of a febrile movement caused by indigestion or gastric irritation. After having had nine different attacks, he remained free from them for a whole year, and then had the tenth at the very beginning of a catarrh of the larynx, fauces, and nasal passages. This child has never as yet exhibited any symptom whatever of disease, either acute or chronic, of the cerebro-spinal axis, and as the convulsions have always been connected with a febrile movement, there is every reason to hope that they are not epileptic. Another patient, likewise not included amongst the above, a girl now five years old, has also had frequent attacks, but as they are of short duration, always coincident with the fever of catarrh, or digestive disorder, and on one occasion that of measles, and as between the seizures her health is excellent, there is but little reason to fear epilepsy.

It is generally believed that the predisposition to convulsions is sometimes hereditary. We have remarked in regard to this point, that several children in the same family sometimes suffer from the disease, and that the nervous temperament to which we alluded above, appeared in some instances to have been inherited by the child from its parents.

In one family that we attend, out of six children, all but one have had attacks of convulsions; one of these children had but one attack, and that was at the age of ten years, and was caused by a fit of indigestion occurring during convalescence from pneumonia. The other four children had

each several attacks, occasioned always by the febrile movement resulting from some of the numerous disorders of infancy. In none of these has there been any reason to suppose that the attacks were settling into epilepsy.

Some very interesting evidence confirmatory of this view has been furnished by Dr. Robert P. Harris, of this city, in an article read before the Philadelphia Obstetrical Society (see *Amer. Jour. of Obstet.*, vol. ii, No. 2, August, 1869).

His record embraces 38 cases of eclampsia, 37 of which occurred in 13 families, in which, collectively, there were 55 children who lived long enough after birth to prove their liability or exemption; 4 having died too early to determine whether they were subject to convulsions or not.

All of the individuals included in the statistics were descendants of the first, second, or third generations, of two pairs of ancestors; of the present rising generation (the second) there are 31 members, only one of whom has as yet married; twenty of the 31 have had convulsions.

The *exciting causes* of convulsions are exceedingly numerous and dissimilar. Amongst the causes of essential convulsions are cited vivid moral emotions, violent pain, high temperature, exposure with the head uncovered to the sun, and sudden exposure to cold. In many cases, however, the exciting cause cannot be detected. The exciting cause of sympathetic convulsions may be almost any of the diseases incident to childhood. Amongst them we will cite as the most frequent, hooping-cough, pneumonia, catarrh, scarlatina, measles, violent fever from any cause, dentition, and indigestion.

It will be observed that many of the causes here assigned for convulsions are also regarded as inducing cerebral congestion, either of the active or passive form. It is accordingly held by some authors that the way in which such influences act is to cause congestion, and that the condition of cerebral congestion is the cause of the convulsions. In our article on congestion of the brain will be found a brief statement of the considerations which lead us to doubt the propriety, in many cases, of ascribing to that condition the convulsive and other nervous symptoms which occur so frequently in connection with the acute local or general diseases of childhood. In regard to some of the other causes above mentioned, however, there can be no doubt that they are likely to induce extreme cerebral congestion. We have seen that in both the active and passive forms of such congestion, convulsions are of frequent occurrence. It is proper then to say that congestion of the brain, of either form and however produced, is among the frequent causes of eclampsia in children.

Of 96 cases of convulsions, of which we have preserved notes, we have regarded only 4 as essential, while 70 were sympathetic, and 22 symptomatic. Of the 4 essential cases, we could not detect the exciting cause in any. Of the 70 sympathetic cases, it was scarlet fever in 12; pertussis in 9; indigestion in 13; pneumonia in 3; the fever of simple angina in 6; cholera infantum and bronchitis, each 3; dysentery, 4; measles and dentition, each 6; enteritis, the fever and irritation caused by a burn upon the back, and the onset of erysipelas, each 1; an overdose of castor oil (3vj)

given to a young child with a slight cholera; 1; and lastly, fecal accumulations in the large intestine, 1.

SYMPTOMS.—*Prodromic Symptoms.*—It has been asserted by some writers that most attacks of convulsions in children are preceded by prodromic symptoms, which indicate to the experienced eye their approach. This does not agree exactly with our own experience, at least in regard to the essential and sympathetic forms, since of the cases of the former variety, well marked prodromes did not occur in any, and of 64 cases of the latter, in which the early symptoms were noted, strongly marked precursory phenomena occurred only in 8. We do not mean to say that there were no symptoms in the other 56 cases which might have indicated to an experienced eye the probability of an approaching attack of convulsions, but merely that there were none that were strikingly characteristic, none which pointed out clearly and decisively that such a crisis was close at hand. In many of the 56, there were symptoms that might be regarded as indicating, with various degrees of probability, the approach of the convulsive seizure; but, inasmuch as they were such as constantly exist in children not predisposed by temperament or constitution to eclampsia, without the development of the disease, they scarcely deserve to be called precursory symptoms.

The precursory symptoms of idiopathic and sympathetic convulsions are, therefore, difficult to describe because of their variable and uncertain character. They consist in general, however, of whatever indicates a highly disordered condition of the nervous system. The most marked symptoms are unusual drowsiness, excessive irritability, a peculiar physiognomical expression, general tremors, and the drawing of the thumbs into the palms of the hands, or rigid flexion of the toes. The drowsiness which precedes an attack of eclampsia, is almost always accompanied with some restlessness. The sleep is light and easily disturbed; the child moves and turns, or starts and moans; often it seems to have frightful dreams, and will scream out or wake suddenly bewildered and terrified, and when roused is generally exceedingly irritable, crying violently or fretting at the slightest contrariety, or without cause. The face, and particularly the eyes, often exhibit a peculiar expression, altogether different from their usual appearance. The expression which has most struck us, and which we have seen on several occasions, is a fixed and staring look, lasting but for an instant, as though the child were looking intently at some object, while in fact it is gazing at vacancy; at the same time the expression is entirely without meaning. The child seems, in fact, for a moment, to be in a state of ecstasy. In some instances a sardonic smile is seen to pass over the countenance just before the attack. The tremors or tremblings alluded to above, occur both in the sleeping and waking state, but particularly in the former. Flexion of the thumbs and toes has been noticed by different observers, but is, we believe, a sign rather of the approach of symptomatic, than of essential or sympathetic convulsions.

The precursory symptoms of symptomatic convulsions will depend on the nature of the disease in the course of which they occur. Not unfrequently the convulsions occur at the very onset of the disease of the brain

or spinal marrow, when of course there will be no prodromic symptoms whatever. According to Dr. Marshall Hall (*Diseases of the Nervous System*, p. 149), the first and most frequent sign showing that the excito-motory system is becoming complicated in diseases of the brain is vomiting, after which come strabismus, a contracted state of the muscles of the thumbs or fingers, or some unequivocal spasmodic or convulsive affection of the respiratory muscles, or of the muscles of the limbs.

Symptoms of the Attack.—With or without the precursory symptoms just described, the convulsion itself usually begins suddenly. The child often utters a cry; loses consciousness and is seized with powerful tonic contraction of the voluntary muscles; the eyes are for a moment fixed and staring, and then drawn obliquely upward under the upper lid, so that the white portions of the balls alone are visible for an instant between the partially open lids; the trunk is rigid and stiff, the thorax immovable, the respiration suspended by rigid spasm of the respiratory muscles; the face, for a moment pale, usually becomes livid and congested, and the veins of the neck are distended.

This state of tonic spasm is followed quickly by the stage of clonic spasm, in which involuntary and most irregular convulsive movements occur. The eyes are rarely fixed in one position, but are constantly agitated in various directions, from side to side, or upwards and downwards; very often there is the most violent strabismus; the eyelids are sometimes open, at others shut; the pupils may be contracted or dilated. The muscles of the face next enter into contraction, and occasion the most hideous contortions of the features. The mouth is distorted into various shapes, the lips are often covered with a whitish or sanguinolent froth, and the jaws tightly clinched together by tonic spasms, or agitated by convulsive movements, so as to produce grinding of the teeth. The trunk of the body is also sometimes variously contorted by clonic convulsions. The head is usually strongly retracted upon the trunk, but in other instances is drawn to one side, or violently rotated. The muscles about the front of the neck enter into action, and alternately elevate and depress the larynx; the tongue, when it can be seen, is observed to be moved in different directions, and is sometimes caught between the teeth and severely bitten. The extremities, particularly the superior, are more violently convulsed than any other parts. The fingers are drawn into the palms of the hands, the forearms are flexed and extended upon the arms by short, rapid, and generally rythmical movements, the hand is quickly pronated and supinated upon the arm, or finally the whole upper extremity is twisted and distorted into various positions, which it is impossible to describe. The inferior extremities undergo similar movements, but almost always in a less degree than the upper. The respiration during the attack is irregular, sometimes suspended by rigid spasm of the respiratory muscles, and sometimes accelerated. A spasmodic contraction of the larynx, producing noisy inspirations, has been noticed by several writers. We shall find when we come to consider the nature of this disease, that Dr. Hall was of opinion that a more or less complete closure of the larynx is the most important feature of the convulsive crisis. The face is often livid and deeply congested, especially when the respiration is embarrassed; the head is hot, whilst the

extremities are cold; the pulse becomes large and full, or frequent and small, and sometimes cannot be counted in consequence of the contractions of the muscles of the forearm. The face is not always, however, congested. We have sometimes seen it perfectly white, while the convulsions were severe, and the child profoundly insensible. The action of the heart is tumultuous, and sometimes irregular or intermittent. When the attack is very violent, the urine and fæces are occasionally discharged involuntarily, but these are rare symptoms. Deglutition is seldom impossible even in the severest fit. In severe, and especially in long-continued attacks, consciousness, and general and special sensibility, are all abolished. In milder cases, though consciousness is destroyed, some of the special senses still respond to irritants, whilst in still slighter cases, the intelligence also is more or less preserved.

As the termination of the attack approaches, the convulsive movements become more and more feeble, until they finally cease entirely, and the child falls into a state of deep sleep, or of more or less profound stupor.

Convulsions are not always, as we have just described them, general. They may be circumscribed or partial, affecting one side of the body more than the other, or one side alone, or a single arm, or in some cases, indeed, only a single muscle, as the biceps. Sometimes they implicate the eyes only. The inferior extremities are rarely affected alone. Of the partial convulsions the most frequent are those in which some parts of the face and upper extremities are attacked. In this form of the disease, the disorders of the circulation and respiration, the congested tint of the face, the froth upon the lips, and the derangements of intelligence and sensibility, are much less strongly marked than in general attacks.

In still other cases, which have been by various authors grouped together under the objectionable title of "inward convulsions," the spasm affects chiefly the muscles of respiration; at times being limited to the muscles of the larynx, and constituting the affection we shall describe in a special article under the name of laryngismus stridulus; at others affecting principally the diaphragm and the thoracic and abdominal muscles of respiration.

The *duration* of an attack of eclampsia concerns both the length of the convulsive crisis and the continuance of the disposition to renewals of the crisis. Both of these are very uncertain. We have known the attack to last in all its violence eight hours and a half in one case, and twelve in another, and it is said to have lasted much longer in some instances. When the spasmodic movements continue during a long period, they are almost always interrupted by remissions. As a general rule, the duration is much shorter than the periods above mentioned—from a few minutes to half an hour. When the attacks cease and recur, as they often do, several times a day, they leave the patient during the intervals in a state of more or less perfect consciousness or somnolence, restlessness or delirium, or finally of coma. The period during which the disposition to recurrence continues, depends principally upon the cause of the convulsions. If this continue in action, they will be apt to return until it is removed.

Idiopathic and sympathetic convulsions generally consist of a single attack, though there are sometimes several, which occur at intervals of

some hours, or of one or two days. Sympathetic convulsions usually occur either at the beginning or termination of the disease which they complicate, and much less frequently during its middle period. Of 46 cases of this form observed by ourselves, complicating measles, scarlet fever, erysipelas, pneumonia, bronchitis, cholera infantum, simple angina, and dysentery, in which the period was carefully ascertained, they occurred at the invasion alone in 25, at the termination alone in 15, at the middle period alone in 3, and at the invasion and termination both in 3. It is curious to remark, that of the 25 cases that occurred only at the invasion of the disease, all but 7 recovered; that the 3 occurring in the middle period alone, also recovered; that of the 3 occurring both at the invasion and termination, 2 died; and that all of those which occurred at the termination alone, proved fatal.

MM. Rilliet and Barthez state that half the cases of symptomatic convulsions observed by them occurred at the commencement of the encephalic disease. This form seldom consists of a single crisis; the attacks, on the contrary, are repeated from time to time. The authors just quoted state that whenever the convulsive attacks have recurred repeatedly within a period of a few days, they have proved symptomatic of disease of the brain.

NATURE OF THE DISEASE.—One of the most important contributions which have been made towards a plausible and satisfactory explanation of the pathology of convulsions in children, was afforded us in the writings of Dr. Marshall Hall; and, although more advanced knowledge of the physiology of the nervous system has shown that the part of the cerebro-spinal axis involved in the production of convulsions is not limited, as he supposed, to the *true spinal system*, his theory of excito-motor action furnishes the most ready explanation of very many cases of eclampsia. Dr. Hall says (*Diseases and Derangements of the Nervous System*, p. 145): "That the whole *class* of convulsive diseases consists of affections of the true spinal system, there is no longer any doubt. But these diseases do not all *originate* in this system." All convulsive disorders are, according to this doctrine, affections of the true spinal or excito-motory system. The causes of these disorders may be of incident origin, acting upon excitor nerves; of centric origin, seated in the brain or spinal marrow; or of reflex origin, acting upon reflex or motor nerves. They are called, therefore, according to their causes, central or centric, when they depend on disease of the nervous centres; centripetal when they are excited through excitor nerves; and centrifugal when they depend on disease of the motor nerves.

Dr. Hall, as is well known, ascribed great importance to the condition of the glottis in convulsions. He says (p. 323), in speaking of epilepsy: "The second symptom is a forcible closure of the *larynx* and *expiratory efforts*, which suffuse the countenance and probably congest the brain with venous blood." At page 327 he says: "A spasmodic affection of the larynx has obviously much to do in this disease (epilepsy, as well as in the crowing inspiration or croup-like convulsion of infants; so much, indeed, that I doubt whether *convulsion* would occur without closure of this organ." In describing the croup-like convulsion or laryngismus stridulus

(p. 180), he says: "I must repeat the observation that the respiration is actually arrested by the closure of the larynx; and there are forcible expiratory efforts only or principally in the actual convulsion." In a later publication, Dr. Hall says: "Without closure of the larynx, extreme laryngismus, and the consequent congestion of the nervous centres, there could, I believe, be no convulsion! This closure of the larynx must be complete in the affection under consideration (laryngismus stridulus), as in all others, before convulsions *can* take place." (*Braith. Ret. from Lancet*, June 12th, 1847, p. 609.)

It is, however, evident that the obstruction to respiration exists not only in the larynx, but in the thorax, the muscles of which are rigidly contracted. Nor can we at present admit that this spasm of the muscles of respiration is more than coincident with the other phenomena of the convulsive attack; and, indeed, there are reasons for believing that the accumulation of venous blood in the nervous centres which follows the obstruction of respiration, so far from causing the convulsion, has a tendency to arrest it, and to induce a state of coma.

It is, however, easy to comprehend the mode of production of sympathetic convulsions by reference to these doctrines. They evidently depend upon morbid impressions conveyed to the cerebro-spinal axis through the excitor nerves having their origin in the diseased organs, probably conjoined with a state of undue reflex excitability of certain parts of the nervous centres. Thus it is easy to understand why inflammation of the parenchyma of the lung in pneumonia, of the bronchial mucous membrane in bronchitis, of the mucous membrane of the bowel in entero-colitis or dysentery, or the pharynx in angina; why the pressure of a tooth upon an inflamed gum during dentition, the presence of a foreign body, as newspaper (in one of our own cases), or crude food, in the stomach, or fecal, or lenteric accumulations in the intestine, should produce a degree of irritation in excitor nerves, sufficient, when transmitted to the sensori-motor ganglia, to occasion the convulsions we have been considering.

It is more difficult to explain the mode in which continued fevers, measles, scarlatina, etc., give rise to convulsions. To us, however, their occurrence is explicable by the morbid effect produced upon the nervous centres by the blood, which is known to be more or less changed in these affections from its healthful condition, and also by the mere fact of the existence of fever; for we have met with a number of children in our own practice, who are almost certain to have a convulsive seizure, whenever the circulation becomes greatly excited in force and frequency by the existence of fever, no matter what be its cause.

The explanation of the production of idiopathic or essential convulsions is not always so easy, because we are sometimes unable to detect any cause, either centric, centripetal, or centrifugal, to account for the excitation of the nervous system. It seems probable, however, that they must depend, like those of the sympathetic form, upon some unhealthful, and therefore irritating condition, acting upon the excito-motory system of nerves. The cause may be so slight as to escape the notice of the physician, and yet sufficient to produce a convulsive crisis in a child predisposed to eclampsia. It may be an unnoticed dentition, some undigested food in contact with

the stomach or intestines, or accumulations of unhealthy fecal substances, or of vitiated secretions, in the intestines. When convulsions have followed a vivid mental emotion, as passion or vexation, they are evidently a result of the influence of that condition upon the nervous centres. Acute pain, which is said to have occasioned essential eclampsia, as well as exposure to violent heat or severe cold, must produce their effects through their action upon incident excitor nerves. There is also in all probability, in most children who suffer with convulsions, a state of preternatural mobility and increased reflex excitability of certain parts of the cerebro-spinal axis, which predisposes to disorderly nervous action, even upon trifling causes. There can be no doubt that this irritability of the nervous system is frequently inherited, though it may be acquired in the course of chronic debilitating diseases. Although we have described these convulsions under the title of essential and sympathetic, we do not mean to assert that they are absolutely independent of any material changes in the nervous centres, but merely that, up to the present time, no appreciable lesions have been detected as their causes. It is indeed true that, in a certain number of instances, after death from eclampsia, there are found engorgement of the vessels of the membranes and of the substance of the brain, serous effusion into the cavity of the arachnoid or the lateral ventricles, or even actual cerebral hemorrhage. In a certain proportion of cases, undoubtedly, cerebral congestion acts as the direct cause of eclampsia, as we have stated in our article on the former subject. But in the majority of cases, the lesions above mentioned cannot be considered as the *causes* of the convulsive attack, but on the other hand must be regarded as the direct *result* of the convulsion, and due to the intense vascular engorgement caused by the spasm of the respiratory muscles and the consequent arrest of the venous circulation. And indeed it is the danger of the occurrence of such lesions which imparts much of the gravity to the prognosis in all severe attacks of eclampsia in young children.

All symptomatic convulsions belong, of course, to the class of centric diseases. These need no further remarks.

DIAGNOSIS.—There are two important points to be considered in treating of the diagnosis of eclampsia: the diseases with which it may be confounded, and the causes which may have produced the convulsions, or, in other words, their distinction into essential, sympathetic, and symptomatic.

The only disease with which eclampsia is likely to be confounded is epilepsy; the mistake could only be made when the former is violent, and when it is accompanied and followed by unconsciousness. In epilepsy, however, the invasion is more sudden, the convulsions are accompanied with greater rigidity, there is always frothing at the mouth, the duration of the crisis is shorter, and it is generally followed by more marked stupor. If the convulsive attack have occurred under the influence of an appreciable cause, if the parents are not epileptic, and if the child is very impressionable, it is probably eclampsia. Again, the younger the patient, the more likely is the case to be one of eclampsia; whilst if the child is approaching towards puberty, if the attacks are frequently repeated, and yet not dependent on fever, and if they are followed by complete restoration to health in the interval, the disease is much more likely to be epilepsy.

The diagnosis of the form of the attack, whether idiopathic, sympathetic, or symptomatic, is exceedingly important, as upon this must depend in great measure the prognosis and treatment. It is often very difficult, and sometimes impossible, to determine at the moment to which class the convulsions belong. The most difficult points in the diagnosis are the following: first, when a child previously in good health, is suddenly seized with the disease, to determine whether it is essential; whether it is sympathetic and occasioned by disease which, up to this instant, has been latent, or by the invasion of some one of the acute local diseases, or of one of the continued fevers; or lastly, whether it is symptomatic, marking the invasion of a disease of the cerebro-spinal axis: second, when the convulsion occurs in the course of a disease not primarily implicating the nervous centres, to determine whether it is merely sympathetic of that disease, or whether it is symptomatic of an intercurrent affection of the brain or spinal marrow.

It is impossible, for want of space, to treat of all these points in detail. The enumeration of them, however, will be useful in calling the attention of the reader to their importance.

An essential convulsion is only to be distinguished by careful study of the antecedent history and present condition of the patient. If, after a thorough examination of all the organs, no diseased point can be detected, and if the child recover perfectly from the convulsion, we must conclude that the case has been an idiopathic one, in which the cause is beyond our reach. We are disposed to believe, however, as has already been stated, that in most of such cases there has been a source of irritation in some of the organs of the body, which has acted as the excitant to the excito-motory system, and which, if we could but detect it, would warrant us in classing the case amongst sympathetic convulsions; and on this account a searching physical examination should be made in every case, as a matter of course.

The sympathetic and symptomatic forms of eclampsia are to be diagnosed by the same careful attention to the antecedent history and present condition of the child. If the latter is teething at the time of the fit, and there is no other cause to explain the attack, and should there be nothing in the consecutive symptoms to render such an explanation inadmissible, we may refer it to that condition. We may remark merely, that, as a general rule, eclampsia depending entirely upon the irritation of dentition, is seldom either violent or long-continued, and that the return to consciousness and health is speedy. The probable dependence of the attack upon indigestion is to be ascertained by the absence of other causes, and by our learning upon inquiry that the child had eaten of some indigestible substance within a few hours or a day or two before the attack. Its dependence on intestinal accumulations is to be arrived at by the same negative or exclusive method, and by learning that the patient is usually, or has been of late, of a constipated habit.

When the attack occurs in the course of some other disease, as pneumonia, catarrh, enteritis, pertussis, scarlatina, or measles, it is almost certainly sympathetic. It may possibly, however, be indicative of an intercurrent attack of cerebral disease. This can be determined only by

attention to the consecutive phenomena. If the attack be short, and soon followed by complete restoration to consciousness, it is in all probability sympathetic. If, on the contrary, the convulsive crisis be long and severe, if the recovery from it be slow and imperfect, if it be followed by violent agitation, somnolence, or coma, or by some persistent lesion of motility, there is every reason to fear an attack of disease of the brain.

Sympathetic convulsions, occurring at the invasion of different local or general diseases, are to be distinguished only by observation of the symptoms that follow the crisis, which will be those belonging to the particular malady whose approach has caused the attack of eclampsia.

Symptomatic eclampsia is characterized by various signs of encephalic disorder, which soon follow the convulsive attack. The most important of these are severe and continued headache; diminution or exaltation of general or special sensibility; dilatation or contraction of the pupils; irregular movements of the eyes; flexion or stiffness of some of the limbs, or of the fingers or thumbs; disordered intelligence; or the symptoms which have already been described in the articles upon the diseases of the brain.

PROGNOSIS.—The prognosis of essential convulsions must depend on the nature of the cause and the violence of the attack. When the cause has been slight, or one which soon ceases to act, or can be readily removed, the prognosis is much more favorable than under opposite conditions. If the convulsive crisis is short and of moderate severity, if the pulse and respiration are but slightly disturbed, if there be but little congestion of the face, and no stertor, there is every reason to hope a successful issue in the case. Of the three cases of this class that we have seen, two recovered and one died.

Sympathetic is more dangerous than essential eclampsia, but much less so than symptomatic. The prognosis will depend chiefly on the nature of the disease which it complicates, and on the stage of that disease at which it occurs. Thus, in scarlatina, convulsions, especially when they occur in the first few days of the disease, are almost always fatal, in measles much less so, and in other diseases in various proportions. They are very apt to terminate unfavorably when they occur after the malady which they complicate has been in progress for several days. This is a remark made by various authors, and we have already stated that of 46 cases of this form in which we carefully ascertained the period of their occurrence, 25 appeared at the invasion, of which all but 7 ended favorably; 3 at the middle period, which all recovered; 3 both at the invasion and at a later period, 2 of which were fatal; and 15 after the cases had been progressing for a considerable time, all of which proved fatal. In addition to these important elements for making the prognosis, we must consider, also, the duration and degree of violence of the paroxysm, the state of the patient after the fit as to its cerebro-spinal functions, and lastly the age and constitution of the child.

The prognosis of symptomatic convulsions must depend very much upon that of the disease of which they are the symptom. It may be stated, as a general rule, that, like those of a sympathetic class, they are less dangerous when they occur at the beginning than at a later period of the dis-

ease. They are always, however, very dangerous. Of 22 cases that we have seen, 19 were fatal.

It frequently happens, however, that although life is not destroyed by the convulsions, certain grave sequelæ remain, among which the most frequent are paralysis, disorders of the mind, and defects of speech or vision. These symptoms are, it is true, far most frequent in cases of symptomatic convulsions, where they depend upon the same lesion of the brain which occasioned the fit. They may, however, succeed convulsions which we are still obliged to call essential, although very possibly there is some minute alteration or defect in a part of the nervous centres, which our means of observation do not suffice to detect. In such cases these sequelæ probably depend upon some lesion of the brain, such as cerebral hemorrhage, which has occurred as a result of the convulsion.

Hemiplegia, which is the form of paralysis which most frequently follows eclampsia, is most apt to occur when the fit has been limited to one side of the body; it is often temporary, and passes away in a few days, though it may remain persistently. Dr. J. Hughlings Jackson suggests that, in the absence of evident disease in such cases of unilateral convulsion and palsy in children, the symptoms may depend upon the plugging of very small vessels in the brain.

TREATMENT.—We shall confine our remarks upon the treatment of eclampsia to the essential and sympathetic forms of the disease, having already treated of that of the symptomatic form in the articles upon the cerebral diseases which give rise to it.

We think that the treatment of eclampsia in children may be simplified if we pay attention to two distinct conditions of disorder, which appear to exist in every case. These are the condition of morbid irritation or derangement of the excito-motory system of nerves, and the cause which occasions that derangement. The condition of irritation or disease of the cerebro-spinal axis exists in all cases, and is always the same, differing only in degree and extent; whilst the morbid cause of that irritation differs in each case, being in one dentition, in another pain, in another constipation, in others pneumonia or indigestion, pleurisy, catarrh, or angina, scarlet fever, measles, fright, or other violent emotions. If this view of the subject be correct, it is clear that in treating a case of convulsions we have to attend to the two morbid conditions referred to, and we shall be careful, therefore, in the course of our remarks, to treat of the remedies most proper for the removal of the cause, whatever it may be, which acts as the irritant to the nervous centres; and of those calculated to subdue or allay the deranged condition of those centres and the effects of that derangement.

There are some general rules to be followed in the treatment of convulsions which apply to all cases, and of these we shall first speak. They are, to place the child in a large *well-ventilated* room, if such can be procured; if it have been seized in a little close room, where the atmosphere is dense and impure, removal to another room, or exposure to fresh air before an open window, has sometimes sufficed to terminate the crisis. At the same time the clothes of the child should be loosened, in order to prevent all constriction, and, if necessary, taken off, to allow of a careful examination

of the whole body. We believe it is a good rule always to place the child, no matter what be the cause of the convulsion, if it be a severe one, in a *warm bath* (96° or 97° F.). This has frequently proved an efficient remedy, according to our experience. It is easily procured in most cases, and we are quite confident that we have never known it to do harm, though we have used it in almost every case. The patient should be kept in the bath some ten, fifteen or twenty minutes, or until the convulsive movements cease; when taken out it is most convenient, and at the same time useful, to envelop it in a small, light blanket, or flannel, for a short time, before the clothes are readjusted.

In cases where the attack of eclampsia is limited to a single convulsive seizure, we rarely have an opportunity of instituting any treatment for the paroxysm itself, since it is usually over before we reach the patient. In such instances, bloodletting is unnecessary. If, however, the opportunity offers, and if the convulsion occur in a strong and vigorous subject; if it be violent, and accompanied by a deep red, or yet more by a livid flush of the face, and distension of the veins of the head and neck; if it last more than a few minutes, or is repeated after short intervals of quiet, we would, without hesitation, recommend the use of *bloodletting*. The detraction of blood is called for, in our opinion, for the same reasons as in puerperal convulsions, and indeed in every violent convulsive attack,—to save the nervous centres from the effects of the paroxysm, which are, in all severe cases, excessive congestion, and, in some, fatal effusions. These instances are, however, comparatively rare, and in the great majority of cases we would not advise depletion in any form. This is particularly true, for instance, in those where the convulsion depends upon an anæmic condition, and in which depletion is contra-indicated by a naturally feeble, or by a debilitated state of the constitution; those in which it is clearly unnecessary, from the slight severity or short duration of the attack; or those which occur in the course of other diseases, and particularly at their termination, and in which a resort to it is rendered evidently improper by the circumstances of the concomitant affection.

During the convulsion it is usually impossible to make the child swallow anything, and when there is but a single attack, and no reason exists for attributing it to the presence of irritating or undigested matters in the stomach, it is not necessary to administer an emetic after the attack. But if the convulsion is prolonged, or a tendency to its recurrence is manifest, we believe that the use of an emetic is very desirable, even when no gastric irritation exists. The act of vomiting alone is often sufficient to break up a paroxysm of convulsions which has resisted various other means. This we learned first from the advice of an old and experienced practitioner, who was in the habit of employing emetics in all cases of eclampsia of children, and we have seen it tested on numerous occasions. The emetic which we prefer in this condition is *ipecaquanha*.

Cold applications to the head will be found proper and useful in nearly all cases of eclampsia which are of any considerable violence. Their use would be improper, however, when the surface is pale, the features contracted, and the pulse small and feeble; but whenever the skin, especially that

of the head, is deeply colored and turgid, and the pulse full and strong, they ought to be employed from the beginning. While the child is in the bath, its head may be wrapped in a cloth wet with ice-water; or, after it has been removed, cold water may be poured from pitchers or a tea-kettle upon the same part. If the latter is done, enough should be employed to prevent the sudden reaction which inevitably takes place when but a small quantity is used. During the subsequent treatment of the case, the cold applications ought to be continued so long as the head remains unnaturally heated.

The administration of a *purgative* dose is proper and useful in most cases of convulsions; particularly when it is found upon inquiry that the child has been constipated prior to the attack; when it is suspected that the bowels may contain crude food or some foreign body; when it is desirable to produce an evacuant effect in a strong plethoric child, or a derivative action from the brain, and when the attack is attended with violent determination of blood towards the organ. The best purgative in severe cases occurring in hearty children is *calomel*. It is advantageous because of its easy administration, its speedy operation, and the powerful sedative influence which it exerts upon the whole economy. The dose should be from two to four grains, according to the age. It ought to be followed in one or two hours by a dose of castor oil or rhubarb. The best of all is castor oil, if it can be given. When the attack is slight or the patient weak and delicate, castor oil is particularly applicable, as it operates with so little irritation to the intestine, or we may employ a mixture of castor oil and spiced syrup of rhubarb. Whatever the remedy may be, it should be given only in such quantity as to produce complete evacuation of the bowels and a moderate derivation upon these organs, without the risk of occasioning a degree of irritation sufficient to increase the disturbance of the nervous system already existing.

In many, indeed, in most cases of eclampsia it will be found that purgative *enemata* are of great service. They may be administered immediately before or after the bath, and not unfrequently have the effect of stopping the paroxysm. They must consist of water holding in suspension or solution castile soap, common salt, molasses, castor oil, sweet oil, or spirit of turpentine. If the first fails to operate in ten or fifteen minutes, another or even a third ought to be given.

Revulsives are of the utmost importance in the treatment of convulsions. They should be employed from the very first, or immediately after the use of the bath. In slight attacks, they alone are often sufficient to suspend the paroxysm, or at least the fit often ceases under their use. Mustard is the most useful and convenient form of application in the great majority of cases. It may be used either in the form of sinapisms, which are to be shifted from place to place, or in that of the foot-bath. When sinapisms are used, they should always be covered with gauze or fine muslin, to avoid the danger of leaving any of the mustard upon the skin after they are taken off. We once saw very bad ulcerations upon the feet of a child from the neglect of this precaution. In the hurry and bustle of the moment, the feet were not washed when the plasters were removed, and the mustard

that remained produced vesications which ulcerated. In obstinate attacks, the revulsives ought to be reapplied from time to time, taking care to shift their position in order to avoid vesication.

Antispasmodics are very valuable remedies in eclampsia, but as they are somewhat slow in acting, we should first resort to the means already detailed. We should then give full doses of one of the antispasmodics recommended below, and should continue its use so long as may be thought desirable after the attack has passed over. They should also be used as a means of prevention in children threatened with eclampsia.

The bromides of potassium and of sodium are the most powerful and reliable remedies of this class in the treatment of almost all forms of convulsion in children. They may be given in the dose of three to five grains, three or four times a day, from the age of six months to one year, with an additional grain for every year.

Next to the bromides, the ones most highly recommended are chloral hydrate, valerian, oxide of zinc, assafoetida, and camphor. Chloral hydrate, given as an enema, as originally recommended by Polaillon (*Union Médicale*, March 23d, 1876), has proved successful in numerous instances in checking the attack. Caution is to be observed in using large doses; those we have ourselves administered have been of from three to five grains, at from two to three years, dissolved in from three or five drachms of thin mucilage of acacia. Valerian is best given in the form of the fluid extract, of which from ten to twenty drops may be administered in water, to a child two years old, every half hour or hour, until several doses have been exhibited, after which it ought to be suspended for awhile or given in smaller quantity. Assafoetida is best given in the form of emulsion, half a teaspoonful being administered by the mouth, or one or two teaspoonfuls mixed with a little warm water may be thrown from time to time into the rectum. M. Brachet (*Traité Prat. des Convulsions dans l'Enfance*, 2ème éd., pp. 102-3) highly recommends the oxide of zinc in combination with extract of hyoscyamus, to the amount of at least two grains of the former and four of the latter in twenty-four hours, divided into four, eight, or twelve doses. A dose was given every two or three hours, and when the symptoms were very violent, the first two or three were repeated at much shorter intervals.

Opium is a remedy which requires much care and discrimination in its employment, but which, in certain conditions of the disease, is of the greatest service. It should not generally be given while there remains any evidence of considerable determination of blood to the brain, but when this condition does not exist, or after it has been removed by bloodletting and revulsives, opium proves very useful in allaying irritability and restlessness, which themselves seem to keep up a disposition to a return or continuance of the convulsive phenomena. Somnolence also, and still more, coma, likewise contra-indicate the use of opium. Dr. Eberle thinks he has seen much advantage from frictions over the spinal region with a mixture of equal parts of oil of amber, laudanum, and spirit of camphor, particularly in very young infants.

Chloroform has been highly recommended in the essential convulsions of

children, by Sir J. Y. Simpson (*Edin. Med. Jour.*, June, 1858), and is favorably spoken of by both West and Trousseau. It should be used when the fits are violent and recur frequently, and do not yield to any of the remedies previously mentioned. By careful administration, anæsthesia may be prolonged for many hours, though, as Simpson recommends, it should be allowed to partially pass off every two or three hours for the purpose of feeding the child. It has been employed in numerous cases with the effect of arresting the convulsions, and in no instance has any unfavorable result been observed to follow its use, although we should ourselves prefer *sulphuric ether* for the production and maintenance of anæsthesia in such cases.

Ice.—In a case of severe convulsions in a new-born infant recorded by Dr. Ellwood Wilson (*Phila. Med. Times*, Nov. 1st, 1873, p. 65), the introduction of a small conical piece of ice into the rectum immediately arrested the paroxysm each time.

We shall here conclude our remarks upon the general treatment of eclampsia, and proceed to make some observations on the conduct to be pursued under particular circumstances.

It is always highly important for the direction of the treatment, to discover the cause of the attack. This is sometimes very easy, while in other instances it is exceedingly difficult, and not unfrequently impossible. If the attack occur in the course of some acute disease, as pneumonia, catarrh, angina, enteritis, or dysentery, or during the progress of one of the eruptive diseases, the diagnosis of the case is, as a general rule, very easy. If, on the contrary, it occurs at the commencement of one of these affections, the diagnosis will be much more difficult, unless indeed the symptoms of the concomitant disease have already declared themselves, or should do so very soon after the convulsion. The treatment in such cases should be that laid down in our general remarks, modified, however, by the requirements of the particular disease during the course of which the eclampsia occurs.

When the attack occurs suddenly in a child previously in good health, or who had been merely slightly ailing for a few hours, the detection of the cause is still more difficult. The most probable causes under such circumstances are, however, dentition, indigestion, intestinal disorder, or the approach of an acute general or local disease. It is easy to determine by inquiry of the attendants, and by examination of the mouth, whether the child is teething or not. As a general rule, the convulsions which depend solely on the process of dentition, are slight, and last but a short time. In all the instances that we have seen, in which such was the only cause to be detected, the attack was of this nature. The treatment in such instances is to lance the gums, if they are swollen and inflamed over the advancing tooth; to use warm baths, and to administer purgative and then antispasmodic enemata. These simple means will seldom fail when eclampsia depends on the process of dentition alone. But when, on the contrary, there is present indigestion, intestinal accumulations, or enteritis, as often happens during dentition, the case becomes more serious, and requires, in addition to the treatment above described, one directed to the particular coëxisting morbid condition.

The existence of indigestion as the cause of the attack, can be discovered only by ascertaining with great care the diet of the child during the previous days. If it appear that something of an indigestible nature has been eaten within a short time, and if, at the same time, it be impossible to detect any more evident or probable cause for the attack, we should have a right to conclude that it depends upon indigestion. Under these circumstances the proper treatment is the immediate use of the warm bath, and the earliest possible administration of an emetic of ipecacuanha. The operation of the emetic may often be hastened by tickling the fauces with a quill.

The presence of intestinal accumulations as the cause of the paroxysms may be inferred, when it is found upon inquiry that the patient has been constipated for some days, or that the stools have been scanty and hard, or scanty and very offensive; when the abdomen is distended and hard, and the distension is ascertained by palpation and percussion, not to be merely tympany; and, lastly, when there is no more evident cause for the attack. In such a case the particular treatment is the use of purgatives and enemata, in addition to the other means detailed.

If the child presents the symptoms of dyspepsia and malnutrition, associated with anæmia, and the convulsions recur during a long period, the most scrupulous care must be taken to secure a nutritious digestible diet, combined with the use of tonics and iron.

The dependence of the attack on the approach or commencement of some acute general or local disease, can be inferred only from a very careful examination of the antecedent and present phenomena of the case. One of these may be suspected as the cause when we can account for the occurrence of the convulsion on no more reasonable supposition; when neither dentition, indigestion, nor intestinal irritation exists. It is scarcely likely that a convulsion could be occasioned by any of the acute thoracic or abdominal affections, unless the disease had already gone far enough to allow a careful examination of the different radical and physical symptoms, to determine its existence. Perhaps the most difficult cases to diagnose, are those which occur at the beginning of the eruptive fevers. Even here, however, a careful search for the prodromic symptoms of the disease, a watchful observance of the condition of the patient in and after the paroxysms, will generally lead to a correct opinion within a few hours, or after a day, and sometimes at the moment of the attack. Of the eruptive diseases, scarlet fever is much the most apt to be accompanied by convulsions at the onset, and in that disease the remarkable rapidity and activity of the pulse, the state of the fauces, the heat of skin, and early appearance of the eruption, will generally enable us to understand the cause of the convulsion at a very early period.

The treatment of sympathetic eclampsia depending on acute thoracic or abdominal disease, should be that which is proper for the particular malady which they complicate, with the addition of warm baths, revulsives and antispasmodics. The management of the convulsions which complicate the eruptive fevers, will be specially treated of in the articles on those maladies.

ARTICLE VII.

LARYNGISMUS STRIDULUS.

DEFINITION ; SYNONYMS ; FREQUENCY.—Laryngismus stridulus belongs to the class of neuroses. It is characterized by crowing inspirations, or by momentary suspension of the act of respiration; these attacks occur suddenly, and at irregular intervals, are of short duration, cease suddenly, and are unaccompanied by cough, or other signs of irritation of the larynx. If the disease progress, it becomes associated with other convulsive symptoms, as strabismus, distortion of the face, carpopedal spasms, or general convulsions.

It is "the peculiar species of convulsion" of Dr. John Clarke; the inward fits of Underwood; the spasm of the glottis of Marsh, West, Vogel, and some of the French writers; the laryngismus stridulus of Good; the croup-like convulsion of Dr. Marshall Hall; child-crowing; one form of the internal convulsion of MM. Trousseau and Pidoux, of MM. Rilliet and Barthez, and of J. L. Smith; and the thymic asthma of some of the German authors. It is described by Eberle under the title of carpopedal spasms.

The frequency of the disease seems to vary in different countries. In France it would appear to be somewhat rare. MM. Rilliet and Barthez (2ème édit.) speak of having seen nine cases. At the time of publication of their first edition, they had met with only one case, and then stated that they were acquainted with only one other, published by M. Constant in the *Bulletin de Thérapeutique*. In Germany, on the contrary, it would seem to be a rather frequent disease. In England it cannot be very infrequent, since Merriman says it is by no means uncommon. Copeland (*Stridulous Laryngic Suffocation in Children, Dict. of Prac. Med.*) speaks of numerous cases that he has seen, and states that he has had as many as three under treatment at the same time. Ley speaks of having met with considerably above twenty cases. Dr. Marshall Hall remarks that "within the short space of one month, I have seen five cases of croup-like convulsion." Dr. Charles West (4th edit., p. 162) mentions thirty-seven cases of which he has preserved some record. The statements of more recent English writers indicate that it continues to be of quite frequent occurrence.

We do not think it is a common disease in America, though it is certainly not extremely rare, since we have either seen ourselves or heard of the occurrence of a comparatively large number of cases.

PREDISPOSING CAUSES.—Age.—It is generally acknowledged that the disease occurs most frequently during the period of the first dentition, though it has been known to occur as late as six or seven years of age. Of 30 cases selected indifferently from our practice and from authors in which the age is given, 13 were six months or less of age, 11 between six months and one year, 4 between one and two years of age, 1 of two, and 1 of four years of age; so that of the 30, 24 were under one year. It is

evident, therefore, so far as these cases go, that the majority occur within the first, and very few after the second year.

Of the 37 cases mentioned by Dr. West, 31 occurred in children between six months and two years of age. All the cases seen by MM. Rilliet and Barthez were in children under two years old. Those authors state that the seven subjects observed by M. Hérard were more than two years of age, and that 2 of them were between three and four years old. From the statements made by the authors in general, it would seem to be most frequent between the ages of three weeks and eighteen months. It has been known, however, in one very rare instance, to occur as late as seven years of age.

Sex.—It is most frequent in the male sex. Of 50 cases (45 from authors, and 5 by ourselves), 39 occurred in boys, and 11 in girls. MM. Rilliet and Barthez state that of 16 cases observed by themselves and by M. Hérard, 12 occurred in boys, and 4 in girls; of 183 cases collected by M. Lorent, in which the sex was noted, 125 occurred in boys, and 58 in girls.

Constitution.—It seems established that it sometimes occurs in the most healthy and vigorous subjects, being then probably dependent upon reflex nervous irritation. It is, however, far most frequently met with in children who are delicate and feeble, and especially in those of scrofulous or rickety constitutions. The very frequent association of rachitis with laryngismus has been more and more prominently developed during the past few years. Some high authorities, since the publication of Elsässer's researches, in 1843, have even asserted that this connection is a constant one, and that laryngismus is essentially dependent upon craniotabes or rachitic disease of the skull. There are certain cases in our own experience, and others who are reported by careful observers, which do not allow us at present to admit that this connection is an invariable one, but there can be no doubt that in the great majority of cases laryngismus occurs in rachitic children, and particularly in those who have craniotabes, or "soft spots" in the occiput. It not unfrequently attacks several children in a family. Ley quotes four instances from other writers, in which three children in each family had the disease, and in one all three died. He states that his own experience fully confirms this fact.

MM. Rilliet and Barthez (2ème édit., note, t. ii, p. 527) state that Davies and Henrich have met with four, and Torgord five children of the same family affected with the disease. They quote from Reid the curious fact that Powell saw one family of thirteen children, not one of which escaped the disease.

Amongst the causes of the disease, in addition to those already mentioned, must not be forgotten *dentition* and *improper food*. These two are, indeed, probably the most influential of all in the production of the complaint. The age at which it occurs most frequently, the last half of the first, and the first half of the second year, the very period during which the process of dentition is most active, would alone go far to show that this must constitute one of its most powerful predisposing, if not exciting, causes. The opinions of writers on this point are also conclusive as to the great influence of this vital process. Improper food, and especially early

weaning, and the attempt to bring the child up by hand, is clearly a potent predisposing cause of the disease. This has been clearly shown in the cases that have come under our own observation, and especially in one in which contraction with rigidity followed the symptoms of laryngismus. The details of this case will be found appended to the article on contracture. Dr. James Reid, in an excellent work on the disease (see *Brit. and For. Med. Chirurg. Rev.*, July, 1849, p. 163), gives the following conclusions as to its ætiology: "1. That for the occurrence of this complaint, the cerebro-spinal system is required to be in a peculiarly excitable state, which then acts as a predisposing cause. The period of teething is the most likely to produce this condition. 2. That during this irritable state of the nervous centres, the two most frequent (and in the majority of instances the combined) causes are the improper description of food which is administered to the infant, and the impure and irritating atmosphere which it breathes." It must not be forgotten that, while in some cases these causes act in producing laryngismus by reflex irritation from the gums or mucous membrane of the alimentary canal upon a weak and over-sensitive nervous system, in other cases, the laryngismus is essentially connected with rickets, which has been induced by improper feeding.

NATURE AND EXCITING CAUSES ; FORMS.—Much difference of opinion has prevailed in regard to the nature and exciting causes of laryngismus stridulus since the disease has attracted the particular notice of the profession. Kopp and other German authors originally ascribed it to compression of the trachea by an enlarged thymus gland; and Ley supposed it to depend on compression of the pneumogastric nerves by enlarged cervical and bronchial glands. It has become a generally accepted opinion, however, that laryngismus is to be regarded as a neurosis, and to be classed with other partial and incomplete convulsive affections. There are various ways in which the attacks may be excited, supposing the predisposition to exist. Marshall Hall considered it as due to reflex irritation, a view that probably holds true in a certain proportion of cases. Many recent authors, as already stated, are disposed to regard it as dependent upon the direct irritation of the brain, due to the existence of craniotabes. But it is evident that if it is regarded, as we are disposed to do, as really one form of internal convulsions in children, a wider view of its nature and pathology must be entertained.

Before examining in detail the different opinions that have been prominently advanced, we will refer to the anatomical appearances of the malady.

The mucous membrane of the air-passages, as a general rule, is found perfectly healthy, presenting neither redness, inflammatory swelling, œdema, nor accidental products of any kind. The lungs are usually of the natural color and density, and crepitant. M. Hérard (*Bib. du Méd. Prat.*, t. v, pp. 319, 320) observed that in several autopsies made by himself, they always presented one marked change from their natural condition, however, which was a very high degree of emphysema, more general and strongly marked than in any other disease. This alteration is believed to depend, as it does in whooping-cough, upon the impediment to

respiration which exists during the disease. MM. Rilliet and Barthez state, however, that emphysema was not present in any of their autopsies.

The heart and great vessels of the thorax often, but not always, contained more blood than usual, as in asphyxia.

M. Hérard states that he has made very minute researches in regard to the condition of the nervous system, examining the brain and spinal marrow, the pneumogastric, recurrent, and diaphragmatic nerves, and those of the extremities even, to their terminations, without, however, finding important lesions in any case. He excepts only serous effusion in small quantity, and evidently consecutive, in the ventricles and particularly in the membranes of the brain, and slight venous congestion of the same kind. The tissues of the brain and spinal marrow retained their ordinary consistence, and presented neither redness nor softening.

The condition of the pneumogastric nerves has, however, been variously reported by different authors, some having found them softened, others indurated.

In some cases tuberculosis of the lungs or bronchial glands has been observed. But as these, as well as all the other lesions mentioned, are not constant, they cannot be regarded as characteristic. In many instances more or less marked evidences of rickets are discovered upon the bones of the cranium, the ribs, or the long bones of the extremities.

We will now examine as succinctly as possible the different opinions which have been advocated in regard to the causes of laryngismus stridulus. These may be classed, it seems to us, under four heads. 1. Enlargement of the thymus gland. 2. Enlargement of the cervical and bronchial glands. 3. Organic disease of the cerebro-spinal axis. 4. That which regards it as a simple neurosis, without appreciable anatomical alterations.

1. *Enlargement of the Thymus Gland.*—That the disease is in some cases coincident with, if not dependent upon, this condition, is proved by the observations of Kopp, Hirsch, Haugsted, Kyll, and others. Hasse (*Pathol. Anat.*, Syden. Soc. Ed., p. 384) says there can be little doubt that it sometimes depends upon this cause.

It appears to us, however, that it has been clearly shown by M. Hérard (*loc. cit.*, pp. 320, 321), that the disease is entirely independent of any alteration of the thymus. That observer found that in six children between two and four years old, dying of the affection, the gland weighed between half a drachm and a drachm in five, and four drachms and two scruples in the sixth. These cases alone show that the size of the gland varies greatly in different subjects attacked with the disease. M. Hérard has examined the gland, with a view to the elucidation of this point, in sixty children dying with various diseases, between two and four years of age (the age of those who had died of the disease under consideration). In fifty he found that it presented the same arrangement, color, density, and weight, as in those who had perished with laryngismus stridulus.

All of these subjects exhibited the same aspect; they were pale, thin, and most of them exhausted by diarrhœa. In ten of the sixty the gland was much more voluminous, weighing from two to two and a half or five drachms, and in one instance an ounce and a quarter. The ten subjects upon which these observations were made died of different diseases, croup,

acute laryngitis, asthma, meningitis, and varioloid. All exhibited the appearances of strong and vigorous health; the one which presented the largest gland was very fat, and so robust, that, though only twenty-two months old, he looked to be three or four years. It appears to result therefore from these researches, that the gland is liable to great variations of size, and that its size bears a very exact proportion to the force of the child, being small in those who are slightly developed, or emaciated by chronic disease, and voluminous in those who are vigorously constituted, or who have died of acute diseases.

That the disease does not depend, at least in all cases, on this cause, is shown also by Haugsted (*Arch. de Méd.*, t. xxxiii, 1833, p. 111), who reports the case of a girl, seven years old, in whom the gland weighed five ounces, and measured four inches long, and one and a half in thickness, without its occasioning the least difficulty of breathing of any kind. That it occurs in children in whom the gland is very small, is shown also by Caspari and Pagenstecher (quoted by Hasse, *loc. cit.*).

2. *Enlargement of the Cervical and Bronchial Glands.*—This condition as a cause of the disease, so strongly advocated by Dr. Ley, and adopted upon his authority by Kyll and Hasse, would seem from certain facts and arguments to be of doubtful agency.

Thus, Mr. Wakely (quoted by Kerr) states that "he possesses more than one case of tubercular affection in children, where the pneumogastric nerve has been completely flattened by the pressure of tubercles, without giving rise to any remarkable disturbance of the function of respiration." Dr. Hall doubts the correctness of this explanation of the phenomena of the disease, and says that if the contiguity of enlarged glands with the pneumogastric nerve have any affect, it is by their action upon it as an incident excitor, and not as a motor or muscular nerve.

3. *Organic Disease of the Cerebro-spinal Axis.*—That it may depend on this cause is proved by a case mentioned by Dr. Coley (*On Infants and Children*, Bell's edition, p. 226), who states that in a fatal instance which occurred in his own family, the only morbid appearance found on dissection was a large exostosis growing on the inner surface of the occiput, which compressed the cerebellum and produced chronic inflammation of the dura mater. No disease was discoverable either in the cervical or thoracic glands. Dr. Kyll (*Arch. Gén. de Méd.*, t. xiv, 1837, p. 94) quotes a case from Dr. Corrigan, of Dublin, which had lasted three months, in spite of calomel, emetics, and antispasmodics. Attention was called by chance to the spinal column, when it was discovered that pressure over the third and fourth cervical vertebræ was very painful, and produced loud cries from the child. Two applications of four leeches, at an interval of two days, to that point, removed all the symptoms, and the child recovered perfectly.

Dr. M. Hall (*Diseases and Derangements of the Nervous System*, 1841, p. 99) states that the crowing inspiration may arise from affections of the centre of the excito-motory system. He quotes a case related to him by Mr. Evans, of Hampstead, of spina bifida, in which "there was a croup-like convulsion whenever the little patient turned so as to press upon the

tumor." He states, moreover, that he found induration of the medulla oblongata in one case of the disease.

Dr. West has also noticed occasional attacks of laryngismus stridulus in chronic hydrocephalus, occurring even before much enlargement of the head had appeared.

We have already stated that, in many cases of laryngismus, the patients will be found to be rachitic, and that some have explained this connection by supposing a direct irritation of the brain due to craniotabes; but it seems probable that in most cases the true connection is to be found in the fact that the derangement of general nutrition associated with rickets induces a state of irritability of the nervous centres, which allows the production of convulsions by slight direct or reflex irritations.

4. *That it is a Neurosis.*—We have seen that in very few cases of laryngismus there is actual organic disease of the brain or spinal cord. It is necessary, therefore, to regard it as most frequently a purely spasmodic nervous affection dependent upon irritation of certain parts of the nervous system which are directly or indirectly connected with the muscles of the glottis. Almost all recent authorities concur in the main with this opinion.

That it is not always, however, a neurosis, is also shown by the cases quoted under the first head from Drs. Hall and Coley, and by those in which the disease is accompanied from the first by symptoms of inflammation or congestion of the brain.

It has now been shown that the causes of the disease are exceedingly variable and uncertain, and that any opinion which asserts its dependence on one invariable and constant cause is untenable. We must, therefore, seek some explanation which shall reconcile, as far as possible, the facts related above, and harmonize the various opinions expressed by the authors quoted.

It seems to us that the explanation given by Dr. Hall (*loc. cit.*) is the only one which accounts satisfactorily for the phenomena of the disease, and reconciles the contradictory accounts of its nature and causes brought forward. Dr. Hall regards it as an affection of the excito-motory or true spinal system of nerves, producing in mild cases partial closure of the glottis, and difficult inspirations, while in more severe cases the spasmodic disposition extends to other parts of the body,—to the eyeballs, and to the flexors of the fingers and toes. We have already alluded to his theory that in very violent attacks of laryngismus, where the glottis is entirely shut, the suspension of respiration produces congestion of the nervous centres and general convulsions. As already stated, however, this theory has not been accepted, and we regard the occasional occurrence of general convulsions in connection with laryngismus stridulus, as one proof that this latter affection is merely a partial and imperfectly developed convulsion.

The causes may be either centric, seated in the nervous centres, or centripetal, in the excitor or incident nerves. In the great majority of cases, the causes are centripetal, consisting of various morbid conditions situated at the peripheral extremities of the nerves, which become causes in consequence of the irritation they establish in the nerve-extremities; this irri

tation is transmitted to the nervous centres, and thence reflected through the various efferent or motor nerves to the different portions of the muscular apparatus affected in the disease, the larynx, face, extremities, and lastly, in severe cases, the whole body. The principal causes of this class are dental irritation, occurring during dentition; gastric irritation, arising from excessive or improper food; intestinal irritation, from constipation, intestinal disorder or catharsis; and perhaps the pressure of an enlarged thymus or enlarged cervical or bronchial glands.

The centric class of causes includes such as are seated in the nervous centres. These are much less common than the former class, and give rise to a vastly more dangerous and intractable form of the disease. Foremost among them, according to recent observations, must be placed the development of "soft spots" in the occipital bone in connection with rickets, which allows pressure upon the back of the head to induce irritation of the brain. Indeed, the more this subject is investigated the closer and more frequent does the connection appear to be between laryngismus and rickets. There are also different morbid conditions of the brain and spinal marrow, as inflammation, congestion, and effusion, which appear to have occasionally proved the cause of laryngismus. That such causes sometimes produce the disease is shown by the cases of exostosis already quoted from Coley, that of spinal irritation from Kyll, that of Dr. Hall, in which he found induration of the medulla oblongata, and the one of spina bifida reported to Dr. Hall by Mr. Evans. In the latter case the tumor was seated on the loins. Mr. E. proposed to treat it by compression, but on making the attempt found that it was followed immediately "by the affection described by Dr. J. Clarke" (*Hall, loc. cit.*, p. 144). Other centric causes, which have been ascribed in some rare instances, are passion, vexation, fright, contradiction, etc.

This theory of the nature of the disease likewise accounts for the varying character of the convulsive symptoms. The laryngeal spasm, from which the disease derives its name, does not constitute the whole malady; it is only one of the symptoms, though the principal one, and that by which it is particularly characterized. The other convulsive phenomena, which generally occur only in severe attacks, or after the disease has continued for some time, are distortion of the face, strabismus, carpopedal spasms, and general convulsions. The hydrocephalic symptoms which occur towards the termination of such cases, and the serous effusion within the cranium found after death, are, it ought to be recollected, often the consequences of the congestion of the brain and asphyxia, which take place during the more or less complete closure of the larynx.

SYMPTOMS; COURSE; DURATION.—Laryngismus stridulus begins suddenly with a paroxysm of difficult respiration. The larynx is contracted spasmodically, and the entrance of air into the lungs is either prevented or impeded. In most cases the closure of the larynx is only partial, and the respiratory movement continues, but is accompanied by prolonged and difficult inspirations, which give rise to the crowing or stridulous sound, whence the disease derives its name. The crowing sound is generally heard several times in each paroxysm, owing to the repeated but only partially successful attempts at inspiration; while in very violent cases it

occurs only at the beginning and end of the accession, the respiration being entirely suspended in the middle period. At the same time the child presents an appearance of great distress. The body is thrown forcibly backwards, the eyes are fixed and staring, the nostrils dilated, and the whole countenance indicative of great anxiety. If the paroxysm continues many seconds, the face becomes bluish, the extremities cold, and the fingers and toes contracted. After a few seconds, or a minute, or even longer, the spasms of the larynx ceases; a loud, full inspiration takes place; a fit of crying generally follows, and the child either very soon regains its usual spirits, or, if the paroxysms have been very severe, seems weak, languid and drowsy, and returns more slowly to its ordinary condition. Between the paroxysms the child may seem perfectly well so far as concerns the character of the respiration, but it almost always exhibits the symptoms of some derangement of the general health, or, in other words, of the morbid condition which is the ultimate cause of the laryngeal spasm.

The paroxysms are most apt to occur during sleep, or as the child is waking. They occur spontaneously, and are brought on by fretting or crying, coughing, fright, contrarieties, deglutition, by the sudden application of cold, and other sudden impressions. At the commencement of the disease they occur at rare intervals, and often attract little notice; but, as the case progresses, they become more frequent, and may amount to twenty or thirty in the day, according to Kerr. They sometimes cease entirely for some weeks, or even months, and then recommence. In a case attended by one of ourselves (reported in the *Am. Jour. Med. Sci.*, April, 1847, p. 287), the attacks lasted eighteen days, occurring sometimes two or three times in an hour, and sometimes much less frequently. The child then recovered entirely for a period of seven months, when the disease returned, and after continuing for five days, caused the death of the child in one of the paroxysms.

If the disease continues to progress, it almost always becomes associated with other spasmodic symptoms. The thumbs are drawn tightly into the palms of the hands, and the fingers clasped over them, which gives to the back of the hands a swelled and tumid look. At the same time the toes are strongly fixed under the feet, and the insteps look swelled like the backs of the hands. Sometimes the hands are bent on the forearms, and the forearms on the arms. There is often distortion of the face. In severe cases, or when the disease has continued for a considerable period, epileptiform convulsions make their appearance, and generally prove fatal.

The disease is apyretic in a large majority of cases. When fever arises it almost always depends on the condition which has occasioned the disordered action of the excito-motory system, or on some accidental complication. The pulse during the paroxysm is small, corded, rapid, and sometimes imperceptible. In the intervals it is natural or nearly so.

Death may occur very early in the disease, or after some weeks, months, or, according to Kyll, years. Vogel states (*op. cit.*, p. 272) that "sometimes even the very first attack terminates in death, and a seemingly perfectly healthy child may be carried off in a few seconds." In a case quoted

by MM. Rilliet and Barthez, death took place at the end of three weeks, and in another in twenty months.

The *duration* is very uncertain. It generally, however, lasts several months. In one of our own cases it lasted eighteen days, then ceased for seven months, returned, and proved fatal in five days. In another case, the attacks of spasms returned from time to time, during a period of three weeks. In another case, the notes of which were obligingly furnished us by our friend Dr. Benedict, and which we shall append to this article, it lasted, in connection with contracture, four months and a half, and was followed by perfect recovery.

Other Forms of Internal Convulsions.—We have for the sake of clearness, limited ourselves so far in the present article, to cases where the spasm is confined to the muscles of the larynx, when the attack might be called one of *laryngeal convulsion*.

In other cases, however, the spasm may effect, either solely, or in conjunction with the larynx, the diaphragm, and the respiratory muscles of the abdomen and chest, constituting what is termed by some authors "*internal convulsions*." The most common form of internal convulsion as described by Trousseau, "is characterized by rolling upwards of the eye-balls, by an almost complete loss of consciousness, by extreme difficulty or impossibility of deglutition, by irregular respiration, at times barely preceptible, or free, deep, and blowing, indicating that the diaphragm and the respiratory muscles of the abdomen and chest are especially affected."

These internal convulsions may be associated with partial or even general convulsions of the face and extremities; more frequently, however, they are accompanied by more or less general *tonic* muscular contraction.

In most cases, as indicated in the passage quoted from Trousseau, the muscles of the pharynx are involved, and there is marked dysphagia or utter inability to swallow.

In some instances, also, the frequency, irregularity, and smallness of the pulse, and the irregular and tumultuous character of the action of the heart, indicate, as pointed out by Rilliet and Barthez (*op. cit.*, t. i, p. 510), that the organs of circulation probably share in the convulsion.

The degree in which the larynx participates in the attack varies much in different cases; at times there is no obstacle whatever to the entrance or exit of air through its cavity, at others, the spasm of its muscles is so extreme that the passage of air is entirely obstructed; whilst in still other cases, of which the one communicated to us by the late Prof. Pepper, and quoted at the end of this article, in an example, respiration is difficult and accompanied by a stridulous noise.

The above description applies to those cases of internal convulsions where the convulsion is complete, and presents both the primary tonic contraction and the subsequent clonic spasms of the respiratory muscles.

But in other cases, the attack consists merely of a sudden tonic spasm of the diaphragm and respiratory muscles of the abdomen and chest, followed by a sudden and complete relaxation. The entire suspension of the respiration during the spasm would of course rapidly induce fatal asphyxia,

but fortunately the attacks, as we have met with them, have usually been so brief as not to cause and dangerous symptoms.

These attacks are popularly known in this country, and were described in the earlier editions of this work, under the title of "*Holding-breath Spells.*"

We have met with a considerable number of well-marked cases of the affection, and believe it to be of quite common occurrence. It seldom happens that the physician is consulted in regard to it, as those who have charge of children in whom it occurs, almost always ascribe it to temper, and think it of but little moment. It appears to be the result of a sudden spasm of all the respiratory muscles, so that the child ceases for the time to breathe, from which circumstances, no doubt, it has received its name of "holding-breath spell." There is no stridulous sound, nor hoarseness of the cry, nor indeed sound of any kind. The face is contracted and bluish, the base of the thorax retracted and immovable, and the limbs violently agitated at first, and then stiff; after a few seconds, or perhaps a minute in severe cases, the spasm yields, the child instantly makes a full inspiration, unattended with stridulous sound, and generally bursts into a loud fit of crying, which lasts for a few moments, after which the child seems perfectly well, or else the attack is followed by excessive paleness, with languor or prostration, lasting half an hour or even longer. The attacks recur with variable frequency; there may be several in a day, or but one, or they may occur only at intervals of several days. The most frequent cause of the paroxysms is contradiction. They are determined also by fright, pain, and crying. They never occur spontaneously, and never during sleep, so far as we know. It is to be distinguished from laryngismus stridulus by the absence of the crowing sound, by its not occurring spontaneously or during sleep, and by the absence of carpopedal or other spasmodic symptoms. It is, we believe, a spasmodic affection of respiration, analogous to though not exactly similar to laryngismus stridulus. We have never met with it except during the period of the first dentition, and always in children of nervous temperament. The cases that we have met with all recovered, and in one only did the life of the child seem to be at all endangered. In this instance the paroxysms had recurred very frequently for eleven months, and on two occasions were terminated by slight spasmodic movements of the limbs, lasting only for a few instants, and unaccompanied by insensibility or other dangerous symptoms. After these attacks the child was removed to the country, where he recovered perfectly.

DIAGNOSIS.—The only disease with which laryngismus stridulus is likely to be confounded is spasmodic laryngitis, or false croup. From this it may readily be distinguished by the absence of catarrhal symptoms, or fever; by the fact that the paroxysms occur indifferently in the day or night, and that they are much more frequent; by the duration of the paroxysms, which last only a few seconds, or more rarely a minute; by the absence of cough or hoarseness of the voice, even during the height of the paroxysm; by the occurrence of tonic muscular spasms, and convulsions; and, finally, by the chronic course of the malady; the converse of all of which symptoms exist in spasmodic croup.

PROGNOSIS.—The prognosis of laryngismus stridulus is always serious, since even the mildest cases may terminate fatally in any one of the paroxysms. It is, however, far from being so dangerous a disease as has been supposed by some writers, and amongst others M. Valleix, who states that it is almost always fatal (*Guide du Méd. Prat.*, t. i, p. 564). Of 56 cases collected from Pagenstecher, Hachman, Ley, Kopp, Hall, Constant, Rilliet and Barthez, Kyll, and 5 from our own observation, making 61 in all, 4 died of intercurrent or consecutive disease, while of the remaining 57, 32 were cured, and 25, or about 43 per cent., died of the malady itself.

MM. Rilliet and Barthez quote from M. Lorent, the translator of Dr. Reid's work, the statement, that of 289 cases collected from various writers, 115, or rather more than 39 per cent., proved fatal.

The prognosis given by the physician ought to depend in great measure upon the cause of the malady. When it depends on difficult dentition, improper diet, or gastro-intestinal disease, whether or not connected, as they very frequently are under these circumstances, with rickets, the case will in all probability terminate favorably if the proper treatment can be, and is, brought to bear against those morbid conditions; while if it occur under the influence of a centric cause, or of enlargement of the cervical or bronchial glands, the prognosis becomes much more unpromising.

TREATMENT.—If the views taken of the nature of the disease in the above remarks be correct, it must be evident that for the treatment to offer any considerable chance of success, it must be directed not merely to the removal of the spasm of the larynx, which is only a symptom and not the whole disease, but to the remedying of the deeper-seated cause of the disordered functional action of the excito-motory system of nerves. In this connection it is especially important to search for the symptoms of rickets, which we have seen to be so often the primary underlying cause of the attacks.

When the disease seems to immediately depend upon difficult dentition, the gums ought to be carefully watched, and freely scarified, so soon as there is the least heat or swelling over the advanced teeth. Dr. Marshall Hall deems the use of the gum-lancet one of the most important means of treatment we are possessed of, and recommends that the gums should be fully divided, "not once, or occasionally, but *twice* or even *thrice* daily."

In another place, he says: "We should lance the gums *freely* and *deeply*, over a great part of their extent, *daily*, or *even twice a day*, and apply a sponge with warm water, so as to encourage the flow of blood." He even recommends that, in very urgent cases, the *lateral* as well as the more prominent portions of the gum should be scarified. Lancing of the gums is undoubtedly a most important point in the treatment of this and other diseases of childhood, connected with dentition. We have long been convinced, however, from personal observation, that a resort to this operation, merely because the child is passing through the period of dentition, is at least useless. We have never found it to do any good, unless the teeth are near enough to the surface to produce manifest swelling, attended with heat and soreness of the gums. So long as the gum is hard, insensible, not turgid, and of its natural color, and the mouth not hot, cutting has done no good.

When the disease depends on gastric irritation, the result of an unhealthy milk or of artificial diet, or when there are evidences that these morbid influences have induced rickets, our attention must be directed principally to the removal of these conditions. A wet-nurse ought to be procured at once if one can be obtained, and if the child will nurse. If this cannot be done, the diet must be carefully regulated by the physician. Ass's milk or goat's milk ought to be used if they can be procured; if not we would recommend the gelatin diet, prepared as recommended at page 318. The proportion of the ingredients must be regulated by the condition of the stomach. If the digestive power be very weak, the proportion of milk must be only a fourth, or even a sixth for a few days, while the amount of cream must bear its usual ratio to the milk.

When the child is thin and pale, and the stomach evidently weak and dyspeptic, it is well to resort to small quantities of stimulants, and to tonics in proper doses. The best stimulant is fine old brandy, of which from ten to twenty drops may be given three or four times a day, or every two or three hours. Or we may administer the aromatic spirit of harts-horn in connection with, or without the brandy; of this about ten or fifteen drops should be given four or five times a day, or alternately with the brandy. Of tonics, the most suitable, it seems to us, are quinine, in the dose of a quarter to half of a grain, three or four times a day, or the citrate of iron and quinine, in the dose of half a grain, given in the same way. Another very excellent stimulant and tonic is Huxham's tincture of bark, of which about five to fifteen drops may be prescribed in the place of brandy. This kind of treatment will scarcely fail to stimulate the digestive power of the stomach to greater activity after a few days, and of course to improve the nutritive functions and the strength of the patient. In addition to this we would recommend the persistent use of the remedies which, as cod-liver oil and the compound syrup of the phosphates, are most beneficial in the treatment of rickets. The reader is referred to the article on the latter subject for more detailed discussion of this point.

When the disease is associated with marked intestinal irritation, we must inquire carefully into its nature and causes. It may be connected with constipation, diarrhœa, or with an unhealthy state of the contents of the bowels. It is often dependent on the presence of crude or imperfectly digested food in the alimentary canal, and when this is the case, the only proper method of treatment is to attend to the state of the digestive function, and to discover and employ a proper diet. The bowels are quite frequently very torpid, and the stools, when obtained by medicine, are often found to be very offensive, light-colored, and pasty, conditions generally resulting from imperfect action of the liver. Under these circumstances, small doses of mercurials, or taraxacum, should be resorted to in combination with or followed by light aperients, as castor oil or rhubarb. One of the very best cathartic remedies, when this combination of symptoms is present, is Chaussier's mixture of castor oil and aromatic syrup of rhubarb, consisting of three parts of the former rubbed up with five parts of the latter. The dose is a teaspoonful every two or three hours, until the bowels are well evacuated. It is gentle in its action, and yet very efficient, gives no pain, and is easily taken. If a mercurial be desired,

about two or three grains of blue mass, one or two grains of calomel, or four grains of the mercury with chalk, may be incorporated into an ounce of the mixture. When diarrhœa is present, it must be treated according to its causes, as recommended in the articles on simple diarrhœa and entero-colitis. When, on the contrary, constipation is a marked symptom, this is to be treated by regulation of the diet, by the daily use of warm water enemata (particularly recommended by Dr. M. Hall), or, if these do not answer, by the exhibition of small doses of the mildest aperients.

Dr. Hall states that by strict attention to the dentition process, and to gastric and intestinal irritation in the *dawn* of the disease, he has succeeded in curing all the cases he has seen but one, and in that he found induration of the medulla oblongata.

By those who suppose the disease to depend on enlargement of the thymic, cervical, or bronchial glands, it has been proposed to endeavor to procure a reduction of the hypertrophy of those glands by frequent applications of leeches, by the use of exutories upon the thorax, by the employment of strong purgative medicines, and by the administration of mercury, digitalis, and iodine. In a case apparently connected with enlargement of the bronchial or cervical glands, we should prefer to direct our treatment to the invigoration of the general health by attention to diet, by the use of tonics, and by proper exposure to fresh air, whilst we should employ internally, cod-liver oil, iron, iodide of potassium, the preparations of iodine, and antispasmodics.

When the disease depends on a centric cause, this must be treated, if it can be detected, according to its nature.

Antispasmodics.—Whatever be the causes of laryngismus stridulus, it is undoubtedly proper, whilst our chief efforts are directed towards their removal or mitigation, to make use of antispasmodics in order to moderate the spasmodic symptoms which are but the expression of those causes. The remedies of this class most highly recommended are bromides of potassium and ammonium, belladonna, valerian, musk, assafœtida, oxide of zinc, and small doses of ipecacuanha. One of the bromide salts mentioned should be given in full doses, and may be combined with any of the other antispasmodics. As stated in our remarks on the use of remedies of this class in whooping-cough, there is reason to believe that the bromide of ammonium possesses greater power than the other bromide salts in relieving the spasmodic affections of the larynx. The oxide of zinc which, as stated in the article on eclampsia, is so highly recommended by Brachet and others, may be given alone or combined with extract of hyoscyamus. M. Brachet always combines the oxide of zinc with extract of hyoscyamus, and gives at least two grains of the former with four of the latter, in divided doses, in the twenty-four hours. He states that he has never given more than ten grains of each in the period mentioned. Of the fluid extract of valerian, about a teaspoonful, or even more, might be given in the twenty-four hours, to a child one or two years old. It should be mixed with water, of course.

It must never be forgotten, however, that remedies of this class are

to be employed only as palliatives and adjuvants, and not as curative agents.

Iron.—Of all the remedies to be employed, after attending in the strictest manner to the removal of the exciting causes of the disease, there is none of such almost universal applicability as iron or its preparations. The patient is almost invariably, owing to the faulty state of the digestive and nutritive functions, more or less anæmic, a condition imperatively demanding iron; and as this remedy rarely conflicts with the other means indicated, it should be given probably in all, or nearly all the cases. The metallic iron in powder or in lozenges, in doses of half a grain or a grain three times a day, or the syrup of the iodide of iron in doses of from two to four drops three times a day, in a mixture of syrup and cinnamon water, are the best preparations, and they should be continued, as a general rule, throughout the treatment of the case.

We have already referred to the great value of the prolonged use of remedies which improve the general nutrition, and particularly counteract the rachitic diathesis so often present in cases of laryngismus. Among these may be mentioned cod-liver oil, arsenic, and the alkaline phosphates.

Treatment during the Paroxysm.—When the child is attacked with a paroxysm of difficult breathing, it should be lifted at once into a sitting posture, if it be reclining, and fanned, or carried to an open window, if the weather be not too cold. At the same time cold water should be sprinkled upon the face, and if the attack be violent, we may resort to what is recommended by Dr. Hugh Ley and Dr. Hall, tickling of the fauces to produce nausea or vomiting, or irritation of the nostrils with a feather, so as to occasion gasping respiration. In a case which occurred to the late Dr. C. D. Meigs, accompanied with severe general convulsions, he found that the suspension of the respiration could very generally be broken in upon, and the paroxysm sometimes averted, by the application of a piece of ice, wrapped in a cloth, to the epigastrium and lower part of the sternum.

Dr. Edmunds (*Méd. Times and Gaz.*, March 12th, 1864) also found that the application of one of Chapman's ice-bags to the spine, did more than anything else to keep off the paroxysms in an obstinate case of laryngismus.

If there is marked determination of blood to the head during the attack, it will be proper to apply cloths wet with cold water, or to bathe the head with a cooling alcoholic lotion.

In cases, especially of the more general form of internal convulsions, where the attacks are so frequently repeated and severe as to threaten life, we would recommend the induction of partial anæsthesia by either ether or chloroform, as advised in the article on eclampsia.

Removal to the Country.—When the disease persists in spite of the means above recommended, and especially when it depends on dentition or digestive irritation, change of air will often produce a wonderful effect, and should always be tried.

The following cases are reported in full, as illustrating the peculiarities and treatment of this curious affection :

CASE.—“The subject of this case was a boy, born in July. He was a large, hearty child, and remained well until January of the following year, when his mother's milk failed, and he was placed upon artificial diet. From this time to May following, his diet was cream and water, barley-water, oatmeal, arrowroot, pounded crackers boiled with water, and gum-water, all of which were tried in turn, being prepared and administered with the greatest caution as to time and quantity. A wet-nurse was tried, but the child refused the breast entirely.

“On the 27th of January he was attacked with diarrhœa, which lasted one week. This was followed by constipation, the stools being white, firm, tenacious, and offensive. The constipation continued up to July, when it was replaced by diarrhœa.

“February 4th. On this day, the child being seven months old, was first observed a spasm of the larynx, producing a shrill, croupal whistle, or *ooh, ooh*, during two or three successive respirations, and followed by a cessation of breathing for some seconds, long enough to dash water in his face, carry him to the window, pat him on the back, etc. These spells occurred during the sleeping and waking state, and especially during crying or laughing, and continued almost daily and often many times a day and night until June, when he was taken into the country.

“Simultaneously with the laryngeal spasm, appeared contraction of the upper extremities, the thumbs being drawn tightly into the palms of the hands, the fingers flexed over the thumbs, and the hands bent on the forearms. The backs of the hands were swollen, and the skin looked tight and polished.

“For a few days in the middle of February there was a subsidence of all the symptoms, with decided improvement in every respect.

“On the 25th of the same month occurred a return of all the symptoms, with extension of the spasm to the feet, the toes being bent under the feet, the insteps much swelled and having a polished appearance. At the same time there were occasional spasmodic movements of the muscles of the face, arms, and body, resembling those of chorea. This condition continued with occasional relaxations up to the 11th of June.

“The stomach was exceedingly delicate, rejecting the most carefully selected nourishment, and at times refusing all food. The child became pale, thin, and timid; was disturbed by the slightest noise, and shunned the light as painful.

“He was removed to the country on the 11th of June. There his health was gradually restored. The appetite improved, the spasm of the larynx and contractions of the extremities gradually relaxed, and the thumbs were at last liberated, the skin under them having taken on the appearance of mucous membrane. There was no return of the disease after the middle of June, although the child had a severe attack of diarrhœa in July, after which he got perfectly well, and has remained so up to the present time (twelve months subsequently). The first tooth made its appearance in September, and he now has fourteen, and has cut them all without the least accident. During the last eight months he has been remarkably fat and hearty.

“I am not aware that any medicine had any effect in removing the disease. Calomel, in large and small doses, antispasmodics of all kinds, frictions over the spine, blisters to the back of the head, alteratives, laxatives, etc., were persevered in without benefit. On removing him to the country, and feeding him on milk warm from the cow, at first diluted, and afterwards pure, an improvement was speedily observed.”

The above case, which was communicated to us by Dr. Benedict, was probably associated with rachitis; unfortunately no record is made of the condition of the occiput. The result illustrates most strikingly the good effects of removal from the city to the country, and the adoption of a more healthy diet.

CASE.—The following case is one that occurred to one of us. We extract the account of it from a paper on croup by Dr. J. F. Meigs (*Am. Jour. Med. Sci.*, April, 1847).

The patient was a girl, five months of age. I saw the child on the 28th of March, 1844. The first attack occurred the day before I was called, but as the mother supposed it to be a matter of little consequence, she did not send for me until the next day. The child was well grown, and except a rather too great paleness, looked strong and healthy. It was playful and good-humored, nursed freely, had no fever, and between the paroxysms presented the appearance of perfect health. The crowing fits occurred frequently in the course of the day and night, sometimes two or three times in an hour, or not so often. They often waked the little thing suddenly from tranquil sleep. They consisted of a succession of long and difficult inspirations, accompanied by a peculiar whistling or crowing sound, such as might be supposed to depend on the passage of air through a narrow aperture. During the attack the face assumed an expression of great anxiety, the respiratory muscles contracted with violence, and there seemed to be for the time imminent danger of suffocation. After several seconds or a minute the shrillness of the sound diminished, the struggling subsided, and soon the respiration became perfectly natural, and the child seemed well. The paroxysms were usually followed by fits of crying, which, however, were easily pacified.

The paroxysms gradually diminished in frequency and violence, and ceased entirely after the 13th of April. The treatment consisted simply in careful attention to the general health, and in frequent use of warm baths and mild nauseants.

The child remained perfectly well, with the exception of a slight attack of cholera infantum, until the following November, seven months after, when the disorder recurred. Several paroxysms occurred between the 12th and 17th of the month; but as they were slight and unattended by other symptoms of illness, the mother was not alarmed, and paid but little attention to them. On the 17th of the same month, the child was sitting on the floor amusing itself with some playthings. There were no persons in the room except young children. They saw the little thing stoop forward suddenly, as though in play, and did not therefore regard it immediately. As it remained in that position, however, they went to it, took it up, and found it dead. It had perished suddenly, no doubt in one of the paroxysms of laryngismus.

An autopsy was made, in which the larynx and thoracic organs were examined, but nothing was found to explain the cause of the disease or the sudden death.

In the following interesting case, communicated to us by the late Prof. William Pepper, the attack consisted of persistent laryngismus stridulus, accompanied by frequently recurring internal convulsions affecting the diaphragm and other respiratory muscles, and by tonic contraction of the muscles of the arms.

CASE.—A boy, aged four months, remarkably healthy and well-developed, after suffering a few days with slight catarrhal symptoms, was suddenly seized with a peculiar stridulous crowing respiration.

I saw the child about half an hour from the commencement of the attack, and found it with a pulse of 140, pale face, and livid lips. The pupils were contracted, and the hands firmly clenched; the crowing sound was very loud, and attended every act of inspiration. At times the respiration and circulation would be entirely suspended for many seconds, followed by great lividity of the surface and coldness of the extremities.

Eight or ten leeches were applied behind the ears, the feet placed in warm water, and a dose of castor oil administered, to be followed by saline enemata.

Four hours from the commencement of the attack, all the symptoms were greatly aggravated; the wrists and fingers were firmly flexed, these spasms coinciding with the arrest of the circulation and respiration; there was now perfect insensibility. The

child was placed in a warm bath, cold water was applied to the head, and a sinapism along the spine, without, however, affording any relief to the crowing inspiration, or other spasmodic symptoms.

At the suggestion of Dr. C. D. Meigs, the child was now placed on its right side, with the shoulders elevated; this position to be maintained at least six hours. At the end of that time the child was in no respect improved, and accordingly, at the suggestion of Dr. M., six leeches were applied over the cardiac region; fʒj of lac. assafoetid. was thrown into the rectum, and a blister applied to the back of the neck.

The child expired at midnight, about ten hours from the commencement of the attack, the crowing respiration, with more or less asphyxia, having persisted throughout.

Autopsy, thirty-six hours after Death.—Mucous membrane of the larynx injected, but in other respects natural. Thymus gland three and a half inches long, two and a half wide, and at its upper part three-quarters of an inch thick, its weight was 620 grains, or 10 drachms and 1 scruple. Lower lobes of both lungs greatly congested. Heart natural. The brain, unfortunately, could not be examined.

It will be observed that, in the above case, the laryngismus and other spasmodic symptoms appeared after slight catarrhal symptoms had existed for a few days; and it may be possible that the irritation of the mucous membrane acted as the exciting cause of the convulsive attack, although the absence of a careful post-mortem examination renders it impossible to say positively that no lesion of the nervous centres existed.

ARTICLE VIII.

CONTRACTION WITH RIGIDITY.

THIS is the disease called by the French *contracture*. We shall treat of it as idiopathic contraction with rigidity. It has been little studied until of recent years; since then a number of cases have been placed on record, especially by French writers. We have ourselves met with but one well marked example of it in an independent form. This case, of which we shall give a sketch at the end of this article, and the one of laryngismus stridulus communicated to us by Dr. Benedict, and appended to the article on that disease, furnish very good examples of contraction coëxisting with the former affection. We have also seen two other cases in which the contraction was decided, but in which it lasted but a short time.

The disease is evidently one of the forms of excito-motor disturbance, which present themselves under such a variety of shapes during infancy and childhood. Though it generally exists as an idiopathic and distinct malady, it is in other cases associated with, or follows laryngismus stridulus or spasm of the glottis, and in others again is combined with attacks of general convulsions.

DEFINITION.—By idiopathic contraction with rigidity (*contracture* of the French writers) is meant the involuntary tonic contraction of different flexor muscles of the extremities, particularly those of the fingers and

toes, but sometimes of the forearms and arms also, existing independently of any appreciable organic disease of the cerebro-spinal axis. It has been described by different English writers in connection with laryngismus stridulus, under the title of "carpopedal spasms," "cerebral spasmodic croup," "croup-like convulsions," etc., etc. We believe, however, that it will be useful to describe it separately from that disorder, for though of the same nature, and sometimes associated with it, it often exists as an independent affection.

CAUSES.—It is most common between the ages of one and three years. It is much oftener sympathetic than essential, and its most frequent causes are dentition, disordered states of the digestive function dependent upon improper alimentation, anæmia and its accompanying nervous excitability, brought about by digestive and nutritive derangements, pneumonia, bronchitis, masturbation and other forms of irritation of the genitals, and unfavorable hygienic conditions. In some few cases, the disease is truly essential, since no pathological cause for it whatever can be detected. It is merely necessary to say that it is also often symptomatic of disease of the brain, but of that form of the affection nothing will be said in the present article.

NATURE OF THE DISEASE.—It appears to consist in a functional derangement of the motor tract of the cerebro-spinal axis, occurring without any cause that can be detected, or determined by the existence of some irritation affecting incident excitor nerves. We once saw a child two years of age, who, after a restless, uneasy night, presented in the morning tonic contraction of the flexors of all the toes of both feet, so that the insteps were swollen, and looked smooth and polished. There was no other sign of sickness except peevishness. Learning on inquiry that the bowels had been somewhat constipated for several days, and that the materials of the scanty stools which had been discharged were dark-colored and very offensive, we ordered a dose of castor oil containing two grains of calomel. The contraction continued unyielding until six o'clock in the afternoon, when a very copious, dark-colored, viscid, and offensive stool occurred, and the contraction immediately ceased. Here the cause of the contraction was evidently an accumulation of unhealthy fecal matter in the intestine, which, by irritating certain sensitive fibres of the excitomotor system, caused a reflex motor action that gave rise to permanent muscular contractions. In other cases the disturbance of the excitomotor system depends on the reflex irritation occasioned by the process of dentition, by indigestion, by diarrhœa, pneumonia, pleurisy, etc. In other instances, again, to which the term essential must be applied, it seems to depend simply on general debility and anæmia, which are well known to be productive of functional disease of the nervous system.

SYMPTOMS; COURSE; DURATION.—The disease rarely attacks children previously in good health, but generally those already suffering from some disorder of the general health, or a severe local affection. When sympathetic, the first symptom noted is the contraction which constitutes the disease. When essential, on the contrary, the onset is sometimes marked

by various nervous symptoms, such as giddiness, headache, or somnolence, which soon pass off, leaving the simple contraction with rigidity as the only morbid condition. In most cases, however, the attack begins with the muscular contraction, which generally affects the superior extremities first, and gradually extends to the inferior.

When the disease is fully developed, the thumbs are drawn down into the palms of the hands, and the fingers, strongly flexed at the metacarpophalangeal articulations, cover and conceal the thumbs. At the same time that the metacarpophalangeal articulations are flexed, the phalanges themselves remain extended and the fingers are separated from each other. The contraction generally affects the wrist-joints also, so that the hands are strongly flexed upon the forearms, and in some rare cases the latter upon the arms. The disorder usually affects the inferior extremities likewise, the toes being in a state of tonic flexion or extension, the foot rigidly extended upon the leg, and its point sometimes drawn inwards. The spasm very rarely extends to the knees.

Children old enough to describe their sensation generally complain of stiffness in the affected parts, with more or less severe pains darting along the course of the nerves. The contracted muscles are hard and rigid to the touch, and sometimes enlarged so as to appear in strong relief under the skin. In slight cases the contractions can be overcome by very moderate force and without pain, whilst in those which are more severe, the attempt to overcome the contraction is productive of acute pain in the rigid parts. The backs of the hands and the insteps present a swollen appearance, and the skin over these points is smooth and polished. In the case communicated by Dr. Benedict, appended to the article on laryngismus stridulus, and likewise in our own case, the skin under the thumbs had assumed the appearance of mucous membrane, from the long and close confinement of the member.

In addition to the symptoms already enumerated as characteristic of the malady, there are others which require attention. The child is of course unable to walk or perform any prehensile movement. The intelligence and senses always remain perfect in simple, uncomplicated cases. The nervous system shows signs of disorder in the form of restlessness or languor, and irritability, with crying and peevishness. In the great majority of instances, these are the only nervous symptoms, though in some there are general or partial convulsions, strabismus, and diminution of sensibility. Of these the most frequent are convulsions, which generally come on a few days after the attack, or precede the fatal termination. In the case of Dr. Benedict, referred to above, there were occasional choreic movements of the face, arms, and body. The simple disease is unaccompanied by any febrile movement, and the organic functions go on naturally. In the sympathetic form, on the contrary, we have the various symptoms of the disease which acts as the cause of the contraction, whether that be abdominal or thoracic. The most common train of symptoms, in young children, is the same as that which accompanies gastric or intestinal derangement, morbid dentition, etc. The *course* and *duration* of the disease

are very irregular and uncertain. When once developed it may last from weeks to months, either slowly increasing in severity, or remaining stationary for a length of time. As a general rule, after it has lasted for some time, it becomes intermittent, sometimes diminishing or even disappearing entirely for a period, then reappearing or increasing, to subside or cease again, and so changing without regularity or evident cause, until at last recovery gradually takes place, or death occurs from the concomitant disease, or in a paroxysm of convulsions.

DIAGNOSIS.—The only difficulty in the diagnosis of idiopathic contraction is to distinguish it from symptomatic contraction, or that which depends upon cerebral or spinal disease. The kinds of cerebral disease which most frequently occasion contraction are tubercle of the brain, and meningeal hemorrhage. The distinction can generally be made with considerable facility, however, by attention to the various disorders of intelligence and sensibility, to the fever, constipation, vomiting and different modes of invasion and progress which characterize the symptomatic form. The following table, taken from MM. Rilliet and Barthez, will assist in the diagnosis.

SYMPTOMATIC CONTRACTION.	ESSENTIAL CONTRACTION.
Cerebral symptoms, special functional disorders (convulsions, strabismus, dilatation of the pupils, etc.), preceding or accompanying the contraction.	Similar cerebral symptoms, but only in exceptional cases, sometimes accompanying, but scarcely ever preceding the contraction.
In many cases irregularity of the pulse,	No irregularity of the pulse.
Generally partial, and commencing usually in the elbows and knees, and in a single extremity.	Binary, commencing in the fingers and toes.
Almost always permanent.	Remarkably intermittent.

PROGNOSIS.—The prognosis must depend on the cause of the malady. The contraction itself has no influence whatever on the termination. The fatal termination has always resulted from the anterior or concomitant disease. Six cases observed by M. Barrier all recovered. The case communicated to us by Dr. Benedict, which was connected with laryngismus stridulus, and one very severe one that occurred in our own practice, also terminated favorably. The prognosis is favorable, therefore, when the attack occurs in a child of naturally good constitution, and when the cause of the disease is not a permanent or incurable one. The possibility of the occurrence of fatal convulsions should always lead us to make a guarded prognosis.

TREATMENT.—The treatment must depend on the circumstances under which the disease has made its appearance. When it occurs in the course of an acute local affection, the treatment must of course be that which is proper for the concomitant disorder. When it depends on dentition, or on gastric or intestinal derangement induced by improper diet, the treatment is the same precisely as that recommended for laryngismus stridulus dependent on the same causes.

It may be stated that, as a general rule, all violent remedies, as bleeding, calomel, except in very minute doses as an alterative, drastic cathartics, and blisters, can scarcely fail to be injurious, unless manifestly necessary in the treatment of the concomitant affection.

It is proper in almost all cases to combine with the treatment already recommended, the employment of antispasmodic remedies, particularly when the contractions persist after the removal of the primary disease.

The best remedies of this class are the warm bath, used every day; belladonna; conium; bromide of potassium; the fluid extract of valerian; assafoetida, and camphor. We would further recommend the use of remedies calculated to improve and invigorate the nutrition, and particularly cod-liver oil and iron. The diet ought generally to be nutritious and strengthening, particularly when the patient is weak and delicate.

In conclusion we may state that the treatment should be very much the same as that proposed for laryngismus stridulus, and we therefore refer the reader to that subject for more detailed information.

CASE BY DR. J. F. MEIGS.

CASE.—The subject of this case was a girl nine months old. The parents were healthy persons, but the mother, owing to some indiosyncrasy, had made but a poor nurse for the preceding child, and I had strongly advised her, therefore, at the birth of this one, to give it a wet-nurse. This was not done, however, and it was found necessary to feed the infant a great deal from its birth. During the early months of its life it had some slight attacks of disorder of the digestive system, but being taken to the country for several months in the summer, it there improved very much. On being brought back to town I saw it, and found it pretty well developed, but very pale, and, on the whole, delicate looking. It was still nursed by the mother, but not to any very considerable extent, as it was obliged to be fed several times each day. The food consisted of different farinaceous substancee made with cow's milk.

On the left forearm of the child there was situated a congenital aneurism by anastomosis, which had grown, by the age of nine months, to be as large as a five-cent piece. It was deemed necessary to remove this tumor, and accordingly, on the 11th of January, 1852, a surgeon tried it with a needle and double ligature. The child bore the operation very well, and soon quieted, and was cheerful and ate well until the evening of the 15th, when it was attacked with fever, which lasted all night, and was accompanied with a good deal of cough and some gurgling in the fauces. On the following morning, at about 7½ o'clock, it had a slight convulsive seizure, lasting a few moments, and marked by stiffening of the body, and a staring expression of the eyes. In the middle of the day, it was seized again, and during that and the next day (17th), up to 10 P. M., it had twenty-four convulsions. These lasted from three to eight minutes each; they were general, and consisted of flexions of the limbs, working of the face, and were attended with unconsciousness. There was no opisthotonos during the attacks, no extensions of the limbs, and no contraction of the jaw. Between the seizures, the child nursed perfectly well, sucked the finger, had no stiffness of the lower jaw, and was perfectly conscious. There was during these two days, some fever, as the skin was too warm, and the pulse between 161 and 180. The respiration was more frequent than natural, there was a good deal of cough, some catarrhal râles in the chest, and also some gurgling in the fauces. The stools were scanty, pasty, and white. There was a well marked but rather faint rash on the limbs and trunk, like erythema or mild scarlet fever, and the lymphatic glands on both sides of the lower jaw were somewhat swelled, and quite hard. The treatment directed was one-sixth of a grain of calomel every two hours; two drops of solution of morphia with five of fluid extract of valerian, to be given also every two hours; warm im-

mersion baths, and mustard foot-bath. On the second day, blisters were applied behind the ears.

On the 19th, the child was better. There was no convulsion; she noticed well, smiled a little, nursed heartily, and took some arrowroot-water.

During all this time the tumor in the arm was not at all inflamed. It was neither red, sore to the touch, nor swelled. It was suppurating slightly. Under the idea that the convulsions might depend in part on the operation, and in order to promote suppuration, a warm poultice was kept constantly applied over the tumor.

The child continued better, with the exception of slight angina and severe cough, until the morning of the 22d, when it waked early, crying violently as though in severe pain, and I found the fingers of both hands strongly flexed at the metacarpal articulations over the thumbs, which were themselves drawn into the palms of the hands. The phalanges, though bent, as just stated, at the metacarpal articulations, were stiffly extended at the phalangeal articulations, and at the same time separated from each other. The hands were flexed at the wrists. The toes were flexed, and the feet stiffly cramped at the ankles, and the insteps, as also the backs of the hands looked swollen and cushiony. Any attempt to open the hands was painful and caused crying. The pulse was frequent and small, the skin pale, and very slightly too warm; the intelligence was perfect. The jaw was open, and the act of sucking was performed, but with some difficulty. On the previous day the bowels had been opened three times, and on this day once; the stools were scanty, pasty, and white. At 9 A.M. I ordered two drops of solution of morphia, five of the fluid extract of valerian, and twenty of milk of assafœtida, to be given every two hours.

4 P.M.—Same state, except that the contraction is stronger. There is more heat of skin, much crying, and a restless, distressed motion of the head. At 4½ o'clock, two drops of laudanum were given with assafœtida. A teaspoonful of the following mixture was ordered every hour.

R. Pil. Hydrarg.,	gr. iij.
Ol. Ricini,	fʒ iij.
Syr. Rhei. Aromat.,	fʒ v.—M.

10 P.M.—Has taken three doses of the mixture and had one large, whitish, pasty, stool. Much easier. Has slept a good deal. Contractions not so strong, as the hands can be opened more easily, and with very little pain. Skin soft, of natural temperature, and moist. Ordered one or two more doses of the mixture, and a repetition of the laudanum and assafœtida, in case of restlessness. During all this time the tumor has not separated. A process of ulceration is going on around the ligatures, but there is no inflammation of any consequence; the arm is not swollen, and there is neither redness nor soreness to the touch.

January 30th.—The contracture diminished very much for two days, and then returned, so that during the 27th, and 28th, and 29th it was very marked, the forearms being flexed on the arms, and the hands strongly flexed on the forearms. The feet also were very stiff, and strongly flexed. The head was occasionally but not constantly retracted upon the trunk. The child evidently suffered very much, as it cried constantly and was very restless, except when under the influence of anodynes or antispasmodics. The bowels were sluggish, but had been kept open by the oil and rhubarb mixture. The dejections were generally whitish and pasty, but occasionally there was a healthy yellow stool. On the 28th the following mixture was ordered:

R. Ext. Valerian. Fl.,	fʒ j.
Sp. Ætheris Comp.,	fʒ ss.
Liq. Morph. Sulph.,	gtt. lx.
Syr. Tolutani.	fʒ vj.
Aquæ,	fʒ iij.—M.

A teaspoonful to be given every hour or two, when there is much suffering or restlessness.

On the evening of the 29th the ligatures were removed, as they had become entirely loose, though without cutting off the tumor. The diseased point was not much inflamed, nor was it tender.

The child is still nursed and fed. Since the 29th it has had goat's instead of cow's milk. On the evening of the 30th the patient was more tranquil, the expression was more placid and open, and the contracture not quite so strong.

Up to February 7th, there was no decided change in the symptoms. They continued quite as severe as before. The dyspeptic symptoms, the torpid state of the bowels, the want of appetite, and the white, pasty state of the evacuations were never relieved, except momentarily, by means of cathartics. On the 7th a wet-nurse was procured, but only after the most persevering and urgent solicitation and argument on our part, I having long been convinced that the cause of the contracture lay in the disordered state of the digestive functions, produced and kept up by artificial diet, and perhaps by an unhealthy state of the mother's milk. The parents, however, had always thought that the operation had been the cause of the convulsive disease, and for a length of time would not consent to a wet-nurse.

After the child had been suckled by the wet-nurse for two days, the stools, which, since the beginning of the sickness, now twenty-three days, and to a greater or less extent since birth, had been very unhealthy, became yellow, homogeneous, and natural in character; while the bowels, instead of being obstinately constipated, so as to require large doses of cathartic medicine, were moved spontaneously two or three times a day.

On the 10th we noticed strong divergent strabismus, and the child looked very badly. The left leg was drawn up, whilst the right was stiffened. The left arm was more used than the right, the left hand being carried often to the mouth, while this was never done with the right. It was difficult to measure the degree of the intelligence, but the child occasionally looked at and evidently noticed objects, but during most of the time it was dull and inattentive.

On the 13th there was an evident improvement, the previous night having been very good. The face was improved in color and expression, and was not quite so thin. The contraction was about the same.

14th.—Some diminution of the contraction, the forearm being a little extended upon the arm, and the wrists, though still very rigid, not quite so much drawn. The child looks better; she nurses a great deal, taking all that the mother, and most also of what the wet-nurse, a hearty woman, has.

February 20th.—Doing very well up to last night, when she became more restless, cried a great deal, rolled the head on the pillow, and had slight retractions of the whole trunk of the body. Occasionally she ceased to cry, scarcely breathed, and the eyes were rolled upwards and fixed for several seconds. She looked pale and pinched again, and refused to nurse. Had one whitish, curdy stool.

21st.—Better; more quiet; nurses well. The boring with the head has ceased, and also the retractions of the trunk. One healthy stool.

22d.—Much better; nurses well; one healthy stool. The contraction of the right arm is yielding, and that of the forearm on the arm is gone on both sides. The left wrist is straight; the right one is yielding very much, though it is still somewhat bent. The fingers of the right hand, though still bent, have relaxed very much; those of the left hand are still very much bent, but are less rigid than before. The integument of the palms of both hands has become, in the flexures, whitish, soft, moist, mucus-like, and has an offensive odor. To-day and yesterday the child uses the arms, touches and reaches out for articles; she is much more intelligent, and looks at and observes objects; she now holds her head up, and likes to be carried about sitting up in the arms of the nurse, which before she could not do at all. She is gaining flesh; the color of the surface is improving; the ears have become pink and pretty.

A fresh assafoetida plaster was applied upon the back yesterday.

March 1st.—Continues to do well. The right hand is to-day almost natural, being opened and shut, and used to grasp with, though it still looks a little stiff. Left hand much better; she opens and shuts the forefinger, and grasps and holds toys with it, but the other fingers are still much contracted. The movements of the arms are quite easy and natural. There is no bending of the hands at the wrists, except, perhaps, very slightly in the left extremity. The feet are natural, except a slight stiffness. She now nurses very well, and is growing fat. She is larger, in fact, than before the sickness. The intelligence is improving rapidly, as she notices, smiles occasionally, and distinguishes, her attendants think, between persons. The bowels are regular without medicine. She has taken no remedy of any kind for three days past.

March 11th.—Almost entirely recovered. There is still a slight but only very slight flexion of the fingers of the left hand. General health excellent.

March 29th.—The patient is now perfectly well, except that she uses the forefinger of either hand rather better than all together, so that in grasping and holding an object, she is more apt to seize it with the forefinger than with all. Still she can and does grasp with all, when the object is large, and no one, unless very observant, would notice the peculiarity just described. Embonpoint very good; complexion clear and healthy; sleeps sound; bowels in excellent condition. Intelligence perfect; smiles and laughs a great deal, and distinguishes between persons; takes a great deal of notice. She is about equal in intelligence to a child of eight months old. Does not attempt to speak.

April 10th, 1852.—I was sent for to-day. The child had not been well for three days, having had three or four thin and greenish stools a day, with whitish specks in them. She was fretful and did not sleep well, and had a good deal of loose catarrhal cough and some acceleration of the breathing. I found her in the morning, after a restless night, quite feverish, hot and dry, with frequent respiration, and with some catarrhal wheezing in the chest. She had coughed a good deal, and her mother had found her hands showing some signs of spasm, the forefingers being extended as though pointing, and separated from the other fingers, which were flexed, with the thumbs also, into the palms of the hands.

There is some degree of laryngismus, as on waking from sleep the breathing is labored, difficult, partially suspended, and accompanied with a slight crowing, or rather choking sound, while at the same time the face becomes pale and the mouth bluish. Bowels open three times yesterday, the stools being mucous, greenish, and containing small lumps of undigested caseine.

Ordered a quarter of a grain of mercury with chalk diffused in a teaspoonful of syrup of jalap to be given every two hours.

At 1 p. m. there was a slight general spasm, with stiffening of the limbs and retraction of the head, lasting, however, only a few moments. This occurred again in the afternoon. The dose of the mercury and jalap was reduced one-half in the middle of the day, as the quantity first ordered was found to cause sickness and vomiting.

Evening.—Rather better. No fever; some moisture of the skin; spasm of the hands very much relaxed. The diminished dose of mercury and jalap was well borne.

11th.—Rather better. Some fever still, with cough, gurgling in the throat, and distinct enlargement and hardening of the lymphatic glands at the angles of the jaw on both sides. There is still some contraction of the hands. Bowels open freely twice last night, and the stools better, being of a pale yellow color, and more homogeneous. The jalap and mercury to be suspended.

In the course of the day there were two slight general spasms, with laryngismus. The latter occurred several times during the waking state, but was not severe. Ordered three drops of syrup of ipecac., with four of sweet spirit of nitre, to be given every two hours.

12th.—Much better. Contraction of hands almost gone; very slight feverishness;

cough less frequent and looser; respiration easy. No spasm to-day. Stools more healthy, yellow, homogeneous, and of natural quantity.

13th.—Continues better. Contraction slight. Cough diminishing very much.

14th.—Rather pale, dull, and languid. Has had several attacks of laryngismus, one of which was quite severe, being attended with deep blueness about the mouth, and some of the face also. Does not nurse so well as formerly. The hands exhibit decided flexion of the third, fourth, and fifth fingers at the metacarpo-phalangeal articulations, with stiffened extension of the other phalangeal articulations. Thumbs slightly drawn into the palms, and the forefingers rather extended. Bowels natural. Ordered fifteen drops of brandy, and a very small pinch of the Quevenne's metallic iron in powder, three times a day.

15th.—Condition about the same. On the 22d of March, the first wet-nurse, under whose charge the child had improved so rapidly, was changed, on account of some objection to her personal appearance, and another one procured in her place. This one was a healthy-looking woman, with milk enough, but she was red-haired, irritable, and excessively high-tempered, and the child has been losing ground ever since her arrival. Under the idea that her milk did not suit the child, a third nurse was by my advice obtained to-day (15th), a calm, placid, fat, and comfortable-looking woman, with an abundant supply of milk of ten months old.

17th.—The child has improved very much. She is fatter already, has a contented, tranquil expression, takes more than she did from the previous nurse, and rejects much less of the milk. The stools are now regular, occurring twice daily without aid, and of a natural appearance. The sleep of the child is better now than it has been at any time since the first wet-nurse was dismissed. The attacks of laryngismus are already much less frequent, and less severe. The hands are very nearly in a natural condition. The child is less nervous, not starting now as formerly at sounds.

To continue the brandy and iron.

From this period the child continued to improve regularly in health. She was removed to the country during the summer months, and when brought back in the autumn, was entirely well, with the exception that she was less forward in walking than most children, but not more so than might have been expected in one who had been dangerously ill for so long a time. Her intelligence was good in all respects.

February 5th, 1853.—We have seen this child to-day, and find her in very good health, except that she is rather smaller in size than is usual at her present age. She has been weaned now for about six weeks, and eats heartily and digests well most ordinary food, as milk, meat, potatoes, etc. The weaning was borne very well, except that the appetite was rather deficient and capricious for about a week after the departure of the nurse. She can stand up when placed in the erect position, and can walk feebly when well supported, but not alone, nor can she rise up from a sitting posture. Her intelligence is, in all respects, perfect, but she does not talk as yet. There is no vestige of her former spasmodic symptoms, when she is in good health; but any little turn of sickness reproduces some contraction of one leg, and a slight flexion of the hands.

Some months after this, the child was unfortunately seized with whooping-cough. She did well for several weeks, but one day, being seized with a fit of coughing while seated upon the floor playing, died instantly, doubtless from asphyxia, caused by complete closure of the glottis by spasm. This is the only case of whooping-cough that we have ever known to prove suddenly fatal in this way. There is every reason to suppose that the fatal suspension of respiration was caused by the unnatural excitability of the sphincter muscle of the glottis, left by the previous attack of laryngismus stridulus.

ARTICLE IX.

TETANUS NASCENTIUM.

DEFINITION; SYNONYMS; PERIOD OF OCCURRENCE; FREQUENCY.—Tetanus nascentium is a most fatal affection, occurring principally during the first two weeks after birth, usually running an acute course, and characterized by a more or less general tonic contraction of the voluntary muscles, with paroxysmal exacerbations, and usually without any period of complete relaxation until the close of the malady.

From this definition it will be seen that the affection does not differ in its essential nature from tetanus as it occurs in adults; though there are so many peculiarities in its causes and symptoms as to demand a special discussion. This disease has also been described under the names of trismus nascentium or neonatorum, in accordance with the prominence and frequency of contraction of the muscles of the lower jaw; but as the spasm is rarely limited to these muscles, but usually involves the other muscles of the face and those of the extremities, the more comprehensive name of tetanus seems more appropriate. It most frequently makes its appearance between the third and tenth days after birth, although there are cases on record in which it set in fifteen hours after birth (West), and others where it did not manifest itself until the twelfth or fifteenth day.

CAUSES.—The causes which have been assigned for the production of tetanus nascentium are very numerous; they may, however, be generally divided into the groups of *general* and *local*. Among the local causes, the various morbid conditions of the umbilicus and umbilical vessels hold the most prominent place. These are, however, far from being constantly present, and yet the weight of evidence is at present in favor of regarding diseases of the umbilicus, and more especially of the umbilical arteries, as occasional causes of tetanus nascentium.

In other cases, the disease has been attributed to some blow or accidental injury which the infant had received. It is, however, still a vexed question as to how much influence should be ascribed to those purely mechanical impressions in the production of this affection. One of the most powerful efforts yet made to establish their importance was by Dr. Marion Sims,¹ who published a series of articles to prove that "trismus nascentium is a disease of centric origin depending on a mechanical pressure exerted on the medulla oblongata, and its nerves; and that this pressure is the result, most generally, of an inward displacement of the occipital bone." This displacement is physiological during the paturient state, but its persistence after birth is dependent, according to his theory, chiefly upon the improper position in which infants are allowed to lie, resting upon their occiput for days together.

Further experience, however, has not confirmed this view, nor justified the admission of injury to the cranial bones into the list of common

¹ Amer. Jour. of Med. Sci., April, 1846, p. 363; July, 1848, p. 59; and October 1848, p. 355.

causes; and yet there are a few cases on record in which tetanus undoubtedly appears to have been developed from this source.

GENERAL CAUSES.—Vicissitudes of temperature appear to favor the development of tetanus, since it is frequent in many countries where a high temperature during the day is succeeded by great cold during the night. In the same way, exposure of the infant to wet and cold, as by putting damp clothes upon it, may be productive of the disease. The most frequent and well established cause of tetanus nascentium, however, is a vitiated state of the atmosphere; whether engendered by a filthy condition of the bedding or house, or by imperfect ventilation; and it is to this that we must attribute the frequency of the affection in such dissimilar localities as the Western Hebrides, Iceland and the neighboring islands, Minorca (see Cleghorn, *Observ. on Epidemical Diseases of Minorca*, London, 1768, p. 81), and some of the Southern States of America, where it was formerly not at all unusual for 50 per cent of all infants born to perish during the first two weeks from this cause alone. It was formerly supposed that certain localities, pre-eminent among which are those just mentioned, were peculiarly favorable to the development of this disease, but it is probable that no predisposition exists excepting the fluctuations of the climate and the filthy habits of the people.

The very great importance of filth and deficient ventilation as a cause of tetanus nascentium is, however, most forcibly shown by the great reduction in the frequency of this disease in large lying-in asylums, effected by the introduction of more thorough ventilation and a greater regard to cleanliness. This was conclusively demonstrated in the Dublin Lying-in Asylum towards the close of the last century. Previously to the year 1782, of 17,650 infants born alive in the asylum, 2944, or almost one-sixth, had died within the first fortnight, and in almost every one of these the cause of death was tetanus nascentium. During the next seven years, after Dr. Clarke had simply introduced a much more complete system of ventilation in the wards, of 8033 children born, only 419 in all died, or about 1 in 19, or $5\frac{1}{3}$ th per cent.

Our comparative immunity in this part of America, even among the poor in our cities, is probably due to the greater degree of cleanliness in their houses, and to the improved construction of our hospitals and asylums. In New York, however, according to Dr. Smith,¹ there are more deaths from tetanus during the first year of life than at all other ages together.

The mortality returns of this city indicate that tetanus, although comparatively frequent among infants, is much less so than in New York.

Thus during the 5 years from 1876 to 1880 inclusive, the returns show a total mortality (less still-born), at all ages, of 83,823, and under 1 year of 19,514. During this period there were 246 deaths from tetanus at all ages; 95 of which were during the first year of life, and 151 after that age. Thus the proportion of deaths from tetanus to those from all causes was, after the age of one year, as 1 to 425, and during the first year of life, as 1 to 205.

¹ Amer. Jour. of Med. Sci., July and October, 1865; and op. cit., p. 168.

During this same period the number of deaths in Philadelphia amounted to 93,207.

PATHOLOGICAL APPEARANCES.—We have already alluded to the morbid conditions of the umbilical vessels or umbilicus occasionally found in tetanus nascentium; it is evident, however, that if these lesions have any connection with the disease, they merely act as exciting causes.

The only characteristic lesions of this affection are presented by the nervous system.

The brain and its meninges are frequently found intensely congested, though this is not so uniformly present as a similar condition of the spinal cord; according to numerous observers, however, it is more frequently present than absent. In some cases, this congestion has led to an actual effusion of blood, either between the skull and dura mater, into the arachnoid cavity, or into the ventricles. In some cases, instead of hemorrhage, there has been found serous effusion into the ventricles or into the sub-arachnoid space, accompanied with a diminution of consistence of the cerebral substance, as reported by Matuszynski.

The morbid appearances found in connection with the spinal cord are the same in character as the above, but more constant and even more marked. The vessels of the spinal meninges and of the substance of the cord are intensely congested, and there is frequently effusion of blood into the cavity of the arachnoid.

The value of these appearances was formerly under-estimated from a suspicion that they might be partly, at least, due to the mere gravitation of the blood after death. This suspicion has, however, been entirely removed by the observations of Weber of Kiel, and Finckh of Stuttgart, who placed the bodies of infants dying with tetanus in various positions before examining them, and yet invariably found the above-mentioned conditions.

There is, however, a further source of doubt as to the significance of these lesions. We have already seen, in speaking of eclampsia, an affection in which no appreciable material lesion has as yet been detected, that, in a certain proportion of cases, congestion, serous effusion, or actual hemorrhage might be present not as causes but as effects, and due merely to the intense venous engorgement caused by the embarrassment of the respiration and venous circulation during the convulsion. It is, indeed, it seems to us, highly probable that a similar interpretation may be placed, in many cases at least, upon the morbid appearances above mentioned as being found after death from tetanus nascentium.

We have thus enumerated the lesions of the nervous system which are readily discoverable in many fatal cases of tetanus; and yet these lesions are, it will be observed, almost without exception concerned merely with the vascular supply of the brain and spinal cord, and we are as yet without any accurate investigations into the condition of the nervous tissue itself. Within the past few years the wonderful advances of microscopical science, as applied to pathological anatomy, have revealed structural changes in the nervous system in connection with more than one disease, whose pathology has heretofore been utterly obscure, and it is not too much

to hope that at no distant period the question of the presence of any definite structural change in the brain or spinal cord in cases of tetanus nascentium will be positively settled. In connection with this suggestion, especially in consideration of the analogy between this disease and tetanus in the adult, we append the results of the investigation of Rokitansky and Demme upon the microscopical appearances in the spinal cord in fatal cases of this latter affection.¹

1. The constant anatomical character of tetanus appears to be proliferation of the connective tissue (of the cord); the most striking peculiarity of this lesion is the extent over which it is found.

2. The product is a viscous mass, abounding in nuclei; it remains at this stage of development in both acute and chronic cases, never progressing to the formation of fibres.

3. This change is found almost exclusively in the white medullary substance; the gray matter seems to suffer only secondarily, and then from compression rather than interstitial deposit.

4. The proliferation is not always followed by corresponding swelling of the white matter; it can often be recognized only by means of the microscope.

5. It was principally found in the medulla oblongata, the crura cerebri, the inferior peduncles of the cerebellum, and in the greater part of the spinal cord.

6. This lesion of the connective tissue appears to be due to long-continued or repeated congestions.

7. The period at which it occurs probably varies in different cases.

These observations, which were originally published about 1860, have been confirmed in all essential particulars by Wagner (*Syd. Soc. Year-Book*, 1862, p. 219); and still later by J. Lockhart Clarke, who published in the *Med.-Chir. Trans.*, vol. xlviii., the results of the microscopic examination of the spinal cord in six cases of tetanus, in all of which structural lesions were discovered; and by Dr. Dickinson (*Med.-Chir. Trans.*, vol. li. p. 265).

SYMPTOMS.—There are rarely any premonitory symptoms of the attack, but the onset and development of the disease are usually gradual. The earliest symptom noticed is, in most cases, difficulty in nursing; the infant appearing anxious to nurse and eagerly pressing its mouth against the nipple, but being unable to fully take it into the mouth or to suck, from a rigid condition of the masseter muscles. At the same time it utters a whimpering, whining, unnatural cry.

The tonic muscular contraction very rarely remains limited to the masseters, but soon invades the other muscles of the face, and those of the trunk and extremities.

The expression of the face thus produced is indicative of great suffering; though it is impossible to say how truly this represents the sensations of the patient.

The face is drawn into wrinkles and furrows, and has a strange appear-

¹ Schmidt's Jahrb., vol. iii. (in New Syd. Soc. Year-Book, 1864, p. 232).

ance of age. The condition of the mouth, however, is most characteristic; the jaws are firmly fixed, the lips slightly separated and pressed firmly against the gums, and the angles of the mouth drawn backwards and downwards in the well known risus sardonicus.

During this time, the other voluntary muscles gradually become rigid. At first, their contraction can be overcome by the use of a moderate degree of force, but in the course of twelve or twenty-four hours the period of maximum rigidity is attained. The head is drawn backwards, and firmly fixed; the arms are flexed, and the hands clenched, with the thumbs drawn across the palms. The thighs may be flexed upon the pelvis, or the legs crossed; the great toes are usually adducted and separated from the rest, which are flexed.

The contraction of the dorsal muscles frequently produces opisthotonos; and the entire body is at times rendered so rigid that it can be raised, without bending, by placing a hand under the heels and head. This extreme degree of spasm of all the voluntary muscles may never be developed in some cases; or, when present, it often is not so persistent. When the infant is quiet or sleeping, there is usually a certain degree of relaxation. It is a marked peculiarity of the affection, however, that exacerbations of the tonic spasm are produced by the slightest exciting causes, as an effort at deglutition, a sudden noise, a puff of air, the most delicate touch, or even the alighting of a fly upon the surface. During these paroxysms or clonic spasms, the muscular rigidity and contraction attain their greatest height, and produce the most painful distortion of the face and limbs. The fit, according to West, may be ushered in by a screech. During its continuance, there is a serious interruption of respiration and circulation; the surface becomes livid, and epistaxis may occur. It is during this condition, too, that hemorrhages into the brain or spinal cord, or their meninges, may result.

These paroxysms recur at irregular intervals, but usually in fatal cases, occur with increasing frequency until either the child expires suddenly during one of the fits, or passes into a state of coma.

The pulse does not present any characteristic change; in some cases it has been found accelerated, but in others has continued normal, or has even fallen below the healthy rate.

The condition of the bowels is not uniform. Diarrhœa is frequently present, but is probably due to irritation of the bowels from the irritating nature of the ingesta, or to some accidental cause; particularly as the bowels are occasionally constipated in well marked cases.

The appetite generally appears to continue, but we have already alluded to the fact that any attempts to feed the child bring on violent spasms, which expel the greater part of the food taken into the mouth. Owing principally to this obstacle to the nourishment of the infant, the emaciation is more rapid and marked in this than in almost any other affection of infancy.

This state of the pupils in tetanus nascentium has not been noted with sufficient frequency or accuracy to allow any deductions to be drawn with

regard to it. Smith has been the pupils contracted in the last stage of the disease.

PROGNOSIS.—The majority of authors state that they have never met with a case of recovery from fully established tetanus nascentium.

Dr. Smith has, however, collected 8 cases of recovery, in the histories of which he calls attention to two important peculiarities: that the children were all about a week old when the initiatory symptoms appeared, and that there were fluctuations in the symptoms of the disease. The only circumstances, then, which would lead us to form a less gloomy prognosis than usual are the late appearance of the disease, and the mildness and intermittent character of the symptoms.

Dr. Hüttenbrenner (quoted in *Boston Med. and Surg. Jour.*, Feb. 12th, 1874) has lately published the results of more recent clinical experience in regard to this disease, from which it appears that although the prognosis is very unfavorable, it must not be considered absolutely fatal.

The *diagnosis* of this affection presents no difficulties, being readily made by attention to the persistent muscular contraction, the inability to suck or to take food, and the exacerbations which are produced by the slightest causes.

DURATION.—In fatal cases the duration rarely exceeds forty-eight or seventy-two hours, and death frequently occurs during the first day. There are instances, however, in which its course has been prolonged to the sixth, or even the ninth day; and Smith refers to two remarkable fatal cases, recorded by Underwood and Elsässer, in one of which the duration was six weeks, and in the other thirty-one days.

Dr. Wells has reported (*Brit. Med. Jour.*, Dec. 21st, 1861) the following case of chronic trismus: The child died at the age of one year, having been, from its birth, in a state of tonic spasm or trismus; it was always restless, and appeared ill nourished, though there was no reason for this. All treatment was unavailing. It was suggested that the child's state might proceed from irritation due to the mother's milk; and the child was weaned, but without benefit. At the post-mortem examination there was found a considerable opalescent effusion over the surface of the brain; the cerebellum was harder than usual, and on being cut into presented a homogeneous appearance. The *arbor vitæ* was entirely wanting.

In favorable cases the duration varies from a few days to one month, or even more.

In the 8 favorable cases collected by Smith, the duration was, in 1 case, two days; in 1, a few days; in 1, fourteen days; in 2, fifteen days; in 1, twenty-eight days; in 1, thirty-one days; and in the remaining case about five weeks.

PREVENTION AND TREATMENT.—It is fortunate that we can by wise hygienic measures do much to prevent the occurrence of a disease of such fatality, and in which, when once fully developed, treatment is so unavailing. We have already alluded to the vast diminution in the number of deaths from this disease, which followed the introduction of free ventilation and cleanliness into the wards of the Dublin Lying-in-Hospital. Nor are

the good effects of this practice limited to public institutions, but it has been found that wherever the disease has prevailed to any extent, as on the Southern plantations, its progress can be arrested by insisting upon the observance of cleanliness in bedding and clothing, of mother and child; by cleaning, disinfecting, and freely ventilating the houses; by care in dressing the umbilical cord; and, finally, by attention to the food of the infant, and the condition of its bowels.

Even when the disease has made its appearance these same measures should be carried out with equal care, since by removing all possible causes, so far as we are acquainted with them, we may mitigate the severity of the attack.

In addition to the removal of the causes, the strictest quiet should be enjoined, and all care employed to avoid exciting the violent paroxysms, which are so readily induced.

It would be well, in addition, to examine the occipital region, to discover if the occipital bone be unnaturally depressed, since in one or two cases this has appeared to act as the exciting cause of the attack. If such depression be found, the position of the child should be varied by placing it on its side, in accordance with the recommendation of Dr. Sims.

The application of leeches to the nape of the neck or along the spine, appears indicated in the early stage of the disease. Dr. West advises the practice, though he has had no experience in its use. Collins, however, states he has tried frequent leeching along the spinal column without the least benefit.

Purgatives are only useful to the extent of maintaining regular action of the bowels.

The remedies which have been most highly recommended as directly curative are ether and chloroform, and various narcotics and antispasmodics, as opium, hydrate of chloral, belladonna, aconite, cannabis indica, conium, woorara, tobacco, and assafoetida.

Anæsthetics have been employed frequently in tetanus of the adult, and occasionally in the affection under discussion. Despite, however, the great expectations which were entertained in regard to their utility, their action cannot be considered directly curative. They relieve suffering, however, and by temporarily allaying the spasmodic contraction of the muscles, enable us to administer food or remedies, and thus prolong life, and give time for other agents to act. "So long, therefore, as the patient is able to take food and to obtain periods of comparative quiet, the use of anæsthetic inhalations is not desirable. Great advantages may, however, be obtained from them if he be unable to open the jaw sufficiently to permit of taking food, or if the tetanic spasms are without remission. Ether appears to have stronger facts in its recommendation than chloroform." (J. Hughlings Jackson, and Hutchinson's Report on Tetanus, *Med. Times and Gaz.*, April 6th, 1861.)

The evidence in regard to the superior efficacy of any particular narcotic is highly conflicting. Opium has, until recently, been the one usually relied upon, and several recoveries have occurred under its use.

Of late years, however, various other narcotics have been employed, especially in traumatic tetanus in the adult. Thus belladonna and its alkaloid atropia have been used, the latter hypodermically, with occasional good results. If the sulphate of atropia is used hypodermically in infants, the first dose should not exceed the $\frac{1}{200}$ th or $\frac{1}{150}$ th of a grain, so that its effects may be tested carefully. One-half grain of the salt may be dissolved in a fluid ounce of water, and four to six drops injected under the skin along the spine.

The various preparations of cannabis indica have also been extensively used. Dr. Gaillard reports two cases of recovery from tetanus nascentium under this treatment; in one of which the infant, aged eight days, took as much as f $\overline{3}$ ss. of tincture of cannabis indica in a single day—being equivalent to about eleven grains of the pure extract. This quantity, however, appears excessive.

Woorara has been given in twenty-two cases, according to Demme, with eight cures. It has been recommended by Harley, Spencer Wells, Broca, Vella, Chassaignac, and others. The dose in which this poisonous substance has been given, is from one-eighth to one-half of a grain to an *adult*. The great objection, however, to both this remedy and cannabis indica, is the great want of uniformity in the strength of their preparations, which necessitates the utmost caution in their use.

More recently still, numerous cases of tetanus in the adult have been treated with the various preparations of conium, and with its alkaloid conia, and also with hydrate of chloral, and the results have been of a decidedly encouraging character. Hüttenbrenner (*loc. cit.*) especially recommends hydrate of chloral, which, according to his observations, is preferable to all other remedies in this disease.

Physostigma, or the Calabar bean, has rapidly acquired a very high reputation in the treatment of traumatic tetanus, and although we are not aware of any cases of the disease under consideration in which this remedy has been used, there is no doubt as to the propriety of employing it in tetanus nascentium. The dose for an infant would be about two drops, repeated at short intervals, of a tincture containing in one pint the virtues of two ounces of the bean.

Among the antispasmodics most frequently used, are assafœtida and tobacco, either given internally or by enema, or added to a warm bath. There is no very positive evidence, however, of their efficiency in this disease.

Baths, either of warm water or vapor, should be repeatedly given; they tend to act favorably as sedatives, by relaxing the muscular spasm, and, in addition, excite the action of the skin.

The free use of large doses of quinine, usually in combination with one of the narcotics above mentioned, appears to be serviceable in traumatic tetanus, by reducing the frequency of the pulse and mitigating the tendency to spasm, so that the induction of cinchonism in tetanus nascentium is a measure worthy of a fair trial.

The application of ice to the spine has been highly recommended in tetanus in adults, and is reported to have been used with success in sev-

eral cases. The condition of the bloodvessels of the cord and its membranes, in fatal cases of tetanus nascentium, would certainly appear to indicate its use in this affection also.

Whichever of the above plans of treatment may be adopted, it must never be forgotten that one of the principal dangers and most frequent causes of death in this disease, is the obstacle offered to the nourishment of the infant. We must pay attention, therefore, to the administration of milk, meat-broth, and alcoholic stimuli in small quantities, but frequently repeated; and if the rigidity of the jaw and the occurrence of spasms upon every attempt at deglutition, prevent the child from taking food, we should have recourse to anæsthetics to relax the spasmodic muscular contraction, and enable us to get nourishment into the stomach.

ARTICLE X.

CHOREA.

DEFINITION; SYNONYMS; FREQUENCY.—Chorea is a non-febrile, convulsive disease, characterized by irregular and imperfectly co-ordinated, but not completely involuntary contractions, of different parts of the muscular system, and particularly of the muscles of the face and of the extremities.

It is also called St. Vitus's dance, chorea sancti viti, choreomania, epilepsia saltatoria, and by various other names.

It is evidently impossible at present to determine the frequency of chorea, as it rarely proves fatal, and consequently scarcely figures in the mortality reports. It must, however, be quite frequent, since it rarely happens to us not to have several cases under treatment at any one time, either in private practice or in some public institution. M. Rufz states (*Dict. de Méd.*, t. vii. p. 544) that of 32,976 children admitted into the Children's Hospital of Paris in ten years, only 189 were affected with chorea, or 1 in 377.

PREDISPOSING CAUSES.—*Age.*—Chorea very rarely occurs during infancy. According to M. Rufz, it is seldom met with between one and six years of age, since of 189 cases, in only ten did it occur within that period; while between six and ten years of age it is much more common (61 in 189 cases); and between ten and fifteen years still more so (118 in 189).

M. Sée, in a valuable essay on chorea (*Mém. de l'Acad. Nat. de Médecine*, t. xv, p. 373), and the relations of rheumatism and diseases of the heart with nervous and convulsive diseases, states (page 448), that of 531 cases of chorea treated in the Children's Hospital at Paris, during a period of twenty-two years, 28 were under six years, 218 between six and ten years, and 235 between six and fifteen years of age. M. Sée concludes, after carefully sifting the facts, that the true maximum of frequency is comprised between six and eleven years of age, and that it corresponds especially to the tenth year. Under six years of age it becomes more and more rare as we approach the moment of birth. MM. Simon and Con-

stant, however, met with it in nursing children of twelve, six, and four months of age.

The statistics furnished by Hillier¹ confirm these statements in every detail. Thus, of 422 cases treated as out-patients at the Children's Hospital in London (where no patients over twelve years are received), the numbers at different ages were as follows:

From 3 months to 6 months, .	3	From 6 years to 7 years,	48
" 6 " 12 " .	5	" 7 " 8 " .	51
" 12 " 18 " .	2	" 8 " 9 " .	58
" 18 " 2 years .	4	" 9 " 10 " .	80
" 2 years 3 " .	6	" 10 " 12 " .	104
" 3 " 4 " .	11		—
" 4 " 5 " .	20		422
" 5 " 6 " .	30		

Of 1984 cases of all kinds treated in the wards of the Children's Hospital of this city during the period of twenty years ending with the close of 1875, there were 65 cases of chorea. The ages of these were as follows:

Under 5 years,	2	At 11 years,	8
At 6 years,	7	" 12 "	9
" 7 "	5	" 13 "	4
" 8 "	6	" 14 "	2
" 9 "	7	" 15 "	1
" 10 "	14		

Sex.—It is much more frequent in girls than boys. Of the 531 cases cited by M. Sée, 393 occurred in girls, and only 138 in boys. This is the same result as that attained, M. Sée remarks, by Reeves, Good, etc.,—131 girls in 186 cases. This accords entirely with our own experience, and in a very interesting statistical report by Dr. George S. Gerhard, based on 80 cases observed in this city (*Amer. Jour. Med. Sciences*, July, 1876, p. 99), the number of female patients just doubles that of the males, 53 to 27. Of the 65 cases occurring in the Children's Hospital, tabulated above, 38 were in girls, 27 in boys.

This excess of females over males obtains in chorea of every grade, from the mildest to the most rapidly fatal cases.

Rapid growth and the *second dentition* probably act, in a considerable degree, as predisposing causes of the disease. Particular attention is drawn to these conditions by MM. Rilliet and Barthez, and the precise age at which it is most frequent (between six and eleven years) would seem to show that they exert a very positive influence. The general deterioration of the health, resulting in anæmia, and the exaggerated nervous susceptibility, so often observed at these periods, are probably the immediate causes of the frequency of the disease at this epoch of life.

Drs. Gerhard (*loc. cit.*) and S. Weir Mitchell report that they have observed that chorea occurs more frequently and in a more severe form in the spring than at any other season; and also that relapses of the disease are most apt to take place at that time. They think this is probably at-

¹Diseases of Children (Amer. ed., 1868, p. 234).

tributable to the condition of weakness of the system which exists in the spring.

An altered and anæmic state of the blood has also been supposed, as by Ogle¹ and Barnes,² to be the efficient and exciting cause of the affection. Rilliet and Barthez,³ also, when speaking of rheumatism as a cause of chorea, say that, "while admitting the existence of rheumatic chorea, it must not be forgotten that the disease is frequently of a different nature, and that we meet in authors with incontestable examples of chorea consecutive to chronic diseases that have produced a debilitated condition of the economy, . . . as chlorosis, anæmia, and tuberculosis."

Constitution does not seem to exert much influence in its production, though it is generally thought to be most apt to occur in children of delicate, excitable, and nervous temperament. The belief in *hereditary* predisposition seems to be unfounded save in rare cases. The disease appears to commence more frequently in spring and summer than in winter, and yet it is scarcely known in tropical climates.

Rheumatism, however, is probably the condition in connection with chorea occurs more frequently than with any other. The evidence of many observers of experience is decided upon this point. M. Sée (*loc. cit.*) asserts, after much examination of this subject, that one-half the cases of the chorea are dependent upon the rheumatic poison. Thus of 109 cases of rheumatism admitted into the Hôpital des Enfants, he found that 61 were complicated with chorea. Trousseau⁴ also states that in his experience rheumatism was undoubtedly the most marked cause of chorea. M. Henri Roger⁵ asserts their connection even more strongly, and states that "the coincidence of chorea and rheumatism is so common a fact that it ought to be regarded as a pathological law, just as much as the coincidence of heart disease and rheumatism."

In England, also, this connection between rheumatism and chorea, both of the mild and severe or fatal form, is positively stated by numerous authorities. Thus in 104 cases of the list collected by Dr. Hughes,⁶ "where special inquiries were made respecting rheumatic and heart affections, there were only 15 in which the patients were both free from cardiac murmur, and had not suffered from a previous attack of rheumatism." Hillier (*op. cit.*, p. 236) "believes there is a very close connection between these diseases." West (*op. cit.*, 4th Am. ed., p. 188) says: "Be the exact relation then what it may, it does seem that rheumatism, or the rheumatic diathesis, is a very powerful predisposing cause of chorea." Dr. H. M. Tuckwell, in a valuable article⁷ on the pathology of chorea, strongly upholds their frequent connection, and cites 17 cases of his own, in 11 of

¹ Brit. and For. Med.-Chir. Rev., Jan. and April, 1868, pp. 208, 465.

² Chorea in Pregnancy, Proc. of Obstet. Soc. of London, vol. x, 1868, p. 147.

³ *Op. cit.*, 2ème ed., t. ii, pp. 565-598.

⁴ Clin. Med., 2ème ed., t. ii, pp. 160-198.

⁵ Arch. Gén. de Méd., 1866, vol. ii, p. 641; and 1867, vol. i, p. 54; and Gaz. Méd. de Paris, March 7th, 1868.

⁶ Guy's Hospital Rep., 2d series, vol. iv, 1846.

⁷ St. Barth. Hosp. Rep., vol. v, 1869, pp. 86-105.

which the previous occurrence of rheumatism was allowed, while it was denied only in 6.

Dr. Chambers found in his books, that out of 33 cases of chorea in 6 the affection either began during rheumatic fever, or followed immediately after it, or else rheumatic fever succeeded to the chorea. In 80 cases of non-fatal chorea recorded by Ogle,¹ it appears that in 8 cases rheumatic fever had existed.

On the other hand, several German authors of high authority do not attach so much importance to the causative influence of rheumatism in chorea. Thus Romberg² states that he has not frequently observed their connection; and Vogel³ states that, "although it must be acknowledged that chorea may succeed to acute rheumatism, still the frequency of the occurrence has been very much over-estimated."

Steiner⁴ also states that out of 252 cases of chorea the disease ensued during the decline of acute articular rheumatism in but 4 cases; of 3 fatal cases, however, reported by him, 1 was complicated with rheumatic heart disease.

We must also allude to the argument of Vogel (*op. cit.*, p. 399), that if there were any actual connection between these diseases, then more girls than boys ought to suffer from rheumatism; for it is well known that the former are predominantly subject to chorea. "Just the reverse happens to be the case in rheumatism, which notoriously attacks more boys than girls." We have already quoted extensive statistics, which prove the truth of the first of Vogel's statements; but we are by no means convinced that the latter is correct, and that rheumatism is more frequent in boys than in girls. On the contrary, the statistics quoted by Tuckwell (*loc. cit.*, p. 102) go to show that the reverse even may be the case. Thus during sixteen years there were admitted to the Children's Hospital in London, 478 patients with rheumatism, 252 of whom were females, and 226 males.

We are not aware of the existence of any accurate statistics of the disease in this country in regard to this point, excepting those of Gerhard (*loc. cit.*), in whose 30 cases rheumatism was assigned as the cause in only 4.

The great weight of evidence, however, which has been accumulated in favor of such a connection, together with the decided results of our own observation, appears to us to leave no doubt that in a considerable proportion of cases, chorea is in some way connected with the previous occurrence of rheumatism. We shall have occasion to call attention to the obscurity which frequently attends the manifestations of rheumatism in young children; and it is, therefore, highly probable that in not a few cases of chorea where, on inquiry, the parents deny the previous occurrence of rheumatism, the truly rheumatic nature of some acute febrile attack, with which the child may have suffered months before, has been entirely overlooked.

We will postpone, until we come to discuss the nature of this affection, the consideration of the manner in which rheumatism disposes to chorea,

¹ Brit. and For. Med.-Chir. Rev., April, 1868, p. 1490.

² Dis. of Nerv. Syst. (Syd. Soc), 1853, vol. ii, p. 57.

³ *Op. cit.*, p. 399.

⁴ Preg. Vjrschr. xcix (xxv, 3), p. 43, 1868; in Schmidt's Jahrb., Bd. 142, No. 4, 1869, p. 26.

whether by directly causing centric lesions, as of the spinal meninges; or by inducing a state of anæmia, impaired nutrition, and preternatural mobility of the nervous system; or whether the choreic movements are in some way connected with cardiac disease, which so frequently attends rheumatism in the young.

Syphilitic disease of the nervous centres as a cause of chorea is so rare that, after an extended search, Alison (*Amer. Jour. Med. Sci.*, July, 1877, p. 75) has been able to find only two cases recorded. Two others, however, have been seen by himself.

EXCITING CAUSES.—Of many exciting causes that have been mentioned by different writers, the one most frequent and most clearly proven is the influence of terror. It was assigned as a cause in 31 out of 56 cases collected by Duffosse and Bird, in 34 out of 100 cases collected by Hughes, in 25 out of 128 by Sée, in 9 out of 31 by Peacock, in 9 out of 38 by Hillier, and in 7 out of 30 by Gerhard. Besides this are cited imitation, blows and falls upon the head, fits of violent anger, contrarieties, prolonged excessive mental effort in young subjects, masturbation, the difficult establishment of the menstrual function in girls, or suppression of that function, the sudden drying up of ulcers or eruptions, and, in females after puberty, pregnancy, which indeed is a well ascertained and most important cause.

In a very interesting case reported by Packard (*Amer. Jour. Med. Sciences*, April, 1870, p. 347), a child of 11 was attacked with very violent and persistent chorea following severe irritation of the ulnar side of the matrix of the right thumb-nail due to a large splinter of wood. The chorea persisted, despite judicious treatment and a residence at the seashore, until the irritated filaments of the ulnar nerve were excised, after which speedy and permanent improvement ensued.

Chorea has also been observed in the course of, or as a sequel to, various acute diseases, as pneumonia, the eruptive, typhoid, and intermittent fevers, and affections of the gastro-intestinal tube.

Dr. S. Weir Mitchell (*Amer. Jour. Med. Sciences*, Oct., 1874, p. 342) has called attention, under the name of "post paralytic chorea," to the disorderly movements of choreic nature which are frequently seen to follow paralysis either in the adult or in children. He has pointed out that it is not the ordinary infantile palsy, but rather the hemiplegia of cerebral origin, which is apt to be thus followed by chorea; and that the younger the child the more likely are these choreic sequelæ to ensue.

ANATOMICAL LESIONS.—It would appear that as yet we are unacquainted with any truly characteristic lesion in chorea. In many of the recorded autopsies, it is stated that no lesion either of the cerebro-spinal axis or any other viscus was present. As, however, most of these autopsies were made before the improved methods of microscopic examination of the nervous system were introduced, they cannot be regarded as conclusive upon this point. In many cases, also, the examination of other viscera has been too superficial to have led to the detection of minute but positive and important lesions. Upon the whole, therefore, it may be fairly said, that it is chiefly the examinations which have been made during the past few years

which are of real value, and that there is still need of numerous accurate autopsies before we can consider ourselves justified in speaking of the true lesions in chorea. According to Dr. Octavius Sturges (*Brit. Med. Jour.*, Aug. 23d, 1879), the affection in its simple and uncomplicated form is not due to any lesion which is demonstrable anatomically; and that its symptoms are not otherwise to be explained than by reference to the general character of disturbed muscular movement, when the source of disturbance is, directly or indirectly, a mental impression.

It is evident that the determination of this question presents great difficulties, apart from the fact that fatal cases of chorea are comparatively rare, and that it requires an amount of skill and patient labor, rarely at command, to make the examination with the requisite minuteness. One of these difficulties consists in the fact that, although chorea may exist as a special, individual affection, there are numerous other cases of nervous disease which are of very varied nature, but which are attended with irregular muscular movements truly choreic in character.

We think it highly probable, therefore, that all cases of so-called chorea will never be found to be invariably associated with any one anatomical lesion.

Thus, passing to the actual results of post-mortem examination, we find a number of lesions recorded which evidently refer to cases of organic disease of the nervous centres, which were merely attended with choreoid symptoms.

Among these are enlargement of the odontoid process, effusions into the arachnoid, tumors in the substance of the brain, abscess in the cerebellum, bony plates upon the spinal meninges, and many other entirely disconnected lesions.

On the other hand there are cases on record, in which careful examination has failed entirely to detect any material lesion, either of the nervous centres or of the other viscera, and in which the choreic movements were probably of a reflex character.

Of late years, however, since this question has been subjected to more frequent and critical examination, there are certain lesions which have been found so frequently after death in fatal cases of true chorea, that they must be regarded as possessing some definite connection with the disease. These lesions consist in certain morbid conditions of the heart, and of the nervous centres.

In regard to the lesions of the *heart*, M. Sée (*loc. cit.*, p. 390) states, after a careful examination of eighty-four autopsies, that "in most of the cases, and especially in those most strongly attested, chorea is the result of the rheumatic diathesis, and that it reveals itself by plastic inflammations of the cardiac membranes, of the pleura, and of the peritoneum, with or without articular rheumatism."

Bright, Copland, Todd,¹ Kirkes,² Nairne,³ Begbie,⁴ were also among the first to call attention to the frequency of rheumatic endocarditis in connection with chorea. In an interesting article on "Maniacal Chorea,"⁵ Tuck-

¹ Lumleian Lectures, 1849.

³ London Jour. of Med., 1851.

⁵ Brit. and For. Med.-Chir. Rev., Oct., 1867.

² Medical Gazette, 1850.

⁴ Edin. Med. Jour., 1852.

well gives an analysis of the lesions in 34 fatal cases of chorea collected by himself. In 25 of these the endocardium was found diseased, the presence of warty vegetations on the valves being especially alluded to in 20. Of the remaining 9, no mention is made of the heart in 5, and it is reported as healthy only in 4. The pericardium was found diseased only in 8 of the 34 cases.

In Ogle's fatal cases (*loc. cit.*, pp. 208 and 507), there were in 11 out of 17 instances more or less fibrinous deposit or granulations upon the valves or some part of the endocardium. In two cases only was the pericardium diseased. In the 14 fatal cases collected by Hughes (*loc. cit.*), vegetations were found on the valves of the heart in not less than 11.

The results of careful auscultation, during life, come to support those of post-mortem examination.

Hillier states (*op. cit.*, p. 236) that, "of 37 cases in my note-books there was probably organic disease of the heart in 25, and in 4 others there was evidence of functional derangement, whilst in 8 only was there no sign of cardiac disturbance."

Jules Simon writes from a large experience, and says: "I have been almost always able to detect well-marked evidence of cardiac affection in chorea, in the shape of organic murmurs, hypertrophy of the heart, etc."¹

In our own experience, evidences of rheumatic heart disease have very frequently been present in cases of chorea; and also in cases which have come under our care for organic disease of the heart, there has frequently been a history of previous attacks of chorea.

It is sufficiently evident, therefore, that in a large proportion of cases of chorea, some morbid condition of the endocardium is present. The particular lesion which has been usually found, consists of fine bead-like vegetations, which either fringe the border of the mitral valve, or are seated upon the auricular surface of its leaflets.

These vegetations are in most cases readily detached from the valve, by lightly brushing them with the tip of the finger, or with a camel's-hair brush; and it has been supposed by some observers, as Ogle and Barnes, that they consisted merely of the fibrin of the blood, deposited in the agony of dissolution. We believe, however, both from the previous occurrence of valvular murmurs in cases where such vegetations have been found, as well as from a careful study of the anatomical descriptions of their appearances, and the occasional presence of the positive results of embolism, that these vegetations are produced by a process of endocarditis.

We will, however, discuss the question of their connection with chorea, when we come to speak of the nature of that disease.

In regard to the condition of the nervous system in fatal cases of chorea, there is at times no lesion appreciable, even on microscopic examination, while on the other hand there is not unfrequently marked disease, either of the nervous tissue or of the meninges.

Thus, in the 14 fatal cases collected by Hughes, the *brain* was healthy in 4, only congested in 3 cases; there was softening of the brain, with or

¹ Nouv. Dict. de Méd. et de Chir. Prat., Art. Chorée (quoted by Tuckwell, St. Barth. Hosp. Rep., *loc. cit.*, p. 101).

without opacity of the membranes and serous effusions in 6, and in the seventh with opacity and congestion of the dura mater.

In 11 of the 35 fatal cases collected by Tuckwell, the brain was found softened, and in 9 only is it reported as healthy. In the 16 fatal cases reported by Ogle, the brain was healthy in 6, much congested in 8, softened in but 1, and anæmic in 1 also.

It appears, therefore, that in a notable proportion of the cases upon record, positive organic disease of the brain, and especially in the form of softening has been discovered. In a few instances embolism, or occlusion of the vessels by fibrinous masses, has been observed, either in the carotid artery (Ogle), or in the minute arterial branches leading to patches of softened brain-tissue (Tuckwell). We need, however, a large series of careful observations to determine more positively how frequently lesions of the brain occur, and especially in what proportion of cases embolism is present.

The *spinal cord* has also been found softened with or without opacity and thickening of its membranes, though in a much smaller number of cases, probably in part because it has not been so frequently examined in such cases as the brain.

Of the 16 fatal cases reported by Ogle, its tissue was congested in 5; there was slight softening in 2; in 1 the upper dorsal region of the cord was completely broken down and almost diffuent. In 2 cases the cord was examined by Mr. J. Lockhart Clarke, who found in one (*loc. cit.*, p. 221) that "in the lower part of the dorsal region, at the ninth dorsal nerves, the anterior columns were swollen, and formed a convex protuberance of considerable size. In a transverse section of the cord carried through this part, and examined under the microscope, it was very evident that extensive morbid changes had been going on, the white substance had been softened, . . . and in two or three places there were circumscribed effusions of blood, surrounded by granular exudations which had probably occurred before the effusions." Similar appearances were discovered in the lower dorsal region in the other case (*loc. cit.*, p. 507).

In a case already referred to, observed by Tuckwell, of rapidly fatal maniacal chorea in a lad of seventeen years of age, in addition to several patches of embolic softening of the brain, there was marked softening of the spinal cord in the middle dorsal region.

In 3 fatal cases reported by Steiner (*loc. cit.*), there was increase in the connective tissue of the spinal cord; serous effusion in the spinal canal; and congestion or effusion of blood in the membranes at the exit of the nerves.

Finally in the cases where embolism of the brain was observed by Tuckwell, there was also minute embolism of the *kidneys*.

In a case of fatal chorea reported by Monckton,¹ embolism of one brachial artery occurred, and, after death, large vegetations were found on the aortic valves.

We will have occasion to refer again to these various anatomical appearances when speaking of the nature of chorea.

¹ Brit. Med. Jour., 1866, No. 305.

SYMPTOMS ; COURSE ; DURATION.—The disease may be general or partial: in the first case, it affects all the limbs, the face, and some of the muscles of the trunk; in the second it implicates only one side, the upper extremities, a single member, or a certain group of muscles. It happens not rarely that the choreic movements are limited to one side of the body: thus in 80 cases of non-fatal chorea reported by Ogle (*loc. cit.*, p. 488), the right side alone was affected in 24, whilst the left alone was affected in 20; and in 25 both sides were affected, though in some instances one or the other side was more involved than the opposite one. Of the 30 cases reported by Gerhard (*loc. cit.*), no less than 15 were strictly unilateral, the choreic movements being confined to the right side in 10 instances, and in 5 to the left. In a large majority of the recorded cases of unilateral chorea the right side was the affected one. It occasionally happens, as noted by Russell (*Med. Times and Gazette*, 1868 and 1869) and Gerhard (*loc. cit.*), that a chorea, which begins as unilateral, may subsequently invade the opposite side and become general. Of 7 cases that we have seen, in which this point was noted, it was general in 4, and confined entirely to the right side in 1, and to the left in 2. We shall describe first the prodromes of the disease, then the invasion, and afterwards the symptoms as they exist in fully developed cases.

Prodromic Symptoms.—It is doubtful whether there are, as a general rule, any well marked prodromic symptoms. The only ones that have been mentioned with any authority are irritability and peevishness of the temper, an unusual degree of impressibility, languor, debility, disturbance of the organic functions, exhibited by deranged appetite and an irregular state of the bowels, and, after a time, a certain quickness and irregularity of the movements, which mark the commencement of the characteristic symptoms of the malady.

Invasion.—The onset of the disease is, as already stated, either sudden or gradual, so that there may be several days or more before it reaches any considerable degree of severity, or it may, particularly when the case has been of a sudden and energetic nature, reach its height in a few hours. In most cases, however, it begins with some unusual and singular movements in one of the upper extremities. The choreic movements are often observed first in the fingers, and at the same time, or soon after, in the face. Sooner or later they increase in severity, and extend to the other arm, to the legs, and to the tongue, and the disease is fully developed.

Symptoms of Confirmed General Chorea.—When the disease has become fully confirmed the movements are exceedingly diversified and irregular. The limbs are agitated by involuntary contractions of the muscles into every attitude possible for them to assume. The fingers are opened and shut, brought together or separated, without any regularity. The hands are flexed and extended upon the forearms, or pronated and supinated, whilst the forearms are flexed or extended upon the arms, and the arms moved at the shoulders into every imaginable position. Such are the irregularity and rapidity of the motions that it is often with great difficulty that the patient can seize anything with the hands, and when once the object is obtained, he frequently cannot do with it what he wishes.

This imperfect control over the hands and arms sometimes prevents the patient from carrying food and drink to the mouth, excepting with the utmost difficulty, and may make it necessary to feed the child.

The inferior extremities are affected in the same way as the arms. Walking is always more or less difficult, and in some severe cases impracticable. The patient totters from side to side, or walks rapidly a short distance, and then suddenly stops. Sometimes the progress is accomplished in a zigzag direction, and at others by fits and starts as it were, whilst in others again the walk is rapid and sudden, almost a run. The child often falls while walking or running, either from meeting a slight obstacle, or in consequence of the irregular and imperfect muscular action. In some instances standing is impossible, the knees bending suddenly under the weight of the body. It was no doubt the peculiar irregular and dancing movements of the inferior extremities during the attempts to walk and stand, that gave to the disease its original name of St. Vitus's dance.

The convulsive movements of the face and head are not less singular than those of the limbs. The face is distorted into all kinds of expressions, so that it assumes by turns that of the most opposite emotions,—sadness, terror, joy, or grief. The mouth is opened and shut, or its corners drawn apart, with the greatest irregularity; the tongue is occasionally protruded between the teeth, and sometimes moved rapidly in the mouth, so as to cause a clacking sound; the lower jaw is depressed and elevated, or moved in a lateral direction, and with such violence perhaps as to injure the tongue or teeth. In consequence of the irregular motions of the tongue and mouth, articulation becomes difficult, and the child either stutters, or speaks slowly and badly, or can pronounce only monosyllables. In a case that occurred to one of ourselves, the movements of the mouth and tongue were so violent and uncontrollable that the patient, a boy nine years old, lost for three weeks all power of speech. He was at the same time unable to open or shut the mouth at will, or to swallow at the proper moment, so that in the act of feeding him, which became necessary from his entire want of control over the arms, the food was constantly spilled and spluttered about as though by an idiot. The act of mastication also was quite impossible, so that he could take nothing but fluids for a number of weeks. In another case also that occurred to one of ourselves, in a girl between eight and nine years of age, and which moreover was a relapse, the patient exhibited the same inability to feed herself, and the same difficulty in regard to mastication, so that she had to be nourished for several weeks on soft food. The speech was likewise greatly affected, it being very difficult to understand her muffled, thick, and indistinct utterance.

Whilst the face and limbs are contorted as above described, the head is moved rapidly from side to side, or backwards and forwards, or undergoes constant rotation, and, in some instances, as in two that came under our own notice, all power over the muscles of the back of the neck is lost, and the head falls from side to side, or forwards, as in an infant. In severe cases the choreic movements affect the trunk also, so that the

patient cannot lie upon a bed, but rolls and twists about the floor with such violence as to bruise and excoriate the skin. Deglutition is sometimes slightly embarrassed, and the child is obliged to swallow with great rapidity; in some few cases a peculiar loud cry, like that which occurs in hysteria, dependent apparently upon spasm of the larynx, has also been observed. The muscles of the external and internal respiratory apparatus are rarely affected, though Romberg narrates three remarkable instances, in which dyspnœa, loud whistling respiration, spasmodic contractions of the glottis, or hiccough, were present. Occasionally irregular action and palpitation of the heart are observed, and have been attributed to chorea of its muscular structure.

In some cases, also, the sphincters of the bladder or rectum are partially paralyzed. Retention of urine has been noticed in a few cases; and, on the other hand, the late Professor William Pepper mentions having known incontinence of urine to alternate with chorea of the external muscles.

The disease is unaccompanied by pain unless it be attended with some complication, and what is very singular and remarkable, the constant and often very violent muscular contractions do not seem to occasion fatigue.

There is, however, frequently evidence of a want of muscular power, which may merely amount to an unusual susceptibility to fatigue on voluntary exertion; or complete paralysis may be present, especially in the form of hemiplegia, in cases of unilateral chorea. This latter is by far the most frequent form of palsy in choreic patients, according to our own observation. It occurred in no less than 17 of Gerhard's 80 cases—in 10 times on the right side, and in 7 upon the left.

The general symptoms require some attention. The choreic movements are almost always increased by emotion, as terror, anger, contrarieties, and by the consciousness of being observed. Sleep generally suspends them entirely. In very bad cases they are said to produce insomnia, or to wake the child frequently in the night. The intelligence is rarely affected, except in very severe and long-continued attacks; though some authors appear to have met with frequent instances of impairment or perversion of the intellectual faculties. It is said that idiocy is apt to occur in cases which last for a number of years, but it is probable that in such cases the choreic movements have been associated with some organic lesion of the nerve-centres. The temper is often irritable and capricious. General and special sensibility commonly remain natural; though in some cases, impairment of general sensibility of the parts most convulsed, even amounting to anæsthesia, is noticed. In simple, uncomplicated attacks, the pulse, as a rule, remains natural; the appetite is preserved; there is no unusual thirst, and the bowels continue regular.

The *urine* has at times been observed to be of unusually high specific gravity, and to contain an excess of urates and oxalates. These conditions do not, however, appear to be at all constant or characteristic.

In a considerable proportion of cases of chorea (see statistics on page 611), a bruit is heard on ausculting the heart, usually of low pitch, and not very great intensity. In some cases this is undoubtedly due to the

vegetations so frequently found on the valves of the heart in this disease, but in others it appears to be rather due to the anæmic state of the blood; and in those cases where palpitation exists, it may be due to the irregular contractions of the walls of the heart. It has also been noticed that these murmurs in chorea are often transitory, and even intermitting.

The *course* of the disease is acute or chronic. In a large majority of cases it is acute, the symptoms becoming more and more violent until they reach their height, when they remain stationary for a time, and then subside and disappear under the influence of treatment, or in the natural course of the malady. It has been frequently noticed that when an acute febrile or inflammatory disease is developed during the course of chorea, the spasmodic movements are very apt to diminish or entirely cease for the time. In fatal cases the symptoms are constantly aggravated; the movements become so violent as to make it necessary to secure the child in bed, or in a strait-jacket; the patients, deprived of sleep, become feeble and emaciated; the respiration becomes difficult; intelligence is abolished; the pupils are contracted; and the child dies.

The *duration* is irregular, varying in acute cases between one and three months. The average duration is probably about six or nine weeks. In very slight attacks it may be much less. The duration of chronic cases is from months to years. In fatal cases the duration is sometimes very short. In one it was only nine, and in another twenty-seven days. The local forms of the disease are often peculiarly intractable, and last many years.

Relapses.—Relapses are quite common and are said by Trousseau to be shorter than the original attack. We have, however, in a few cases, observed that the relapse was much worse than the first attack. In one case in particular, the relapse was one of the most violent and prolonged attacks that we have seen. MM. Rilliet and Barthez state they occurred in six out of nineteen cases seen by them. The relapses in these cases occurred once, twice, and three times. M. Sée (*loc. cit.*, p. 408) says that it is not uncommon, after some weeks of respite, or several months of apparent recovery, to see the disease reappear with renewed intensity, and be thus repeated twice, thrice, and even seven times in succession. Out of four patients, at least one, he states, remains thus under the influence of the disease. Of 158 cases he counted 37 relapses, of which 17 were arrested after the second attack; 13 suffered a third, and 6 a fourth attack; and, lastly, one had seven distinct seizures, each one of which was separated from the following by a well marked interval. In 46 of Ogle's cases in which this point was noted, previous attacks had occurred in 25; in 5 of which there had been 2 previous attacks, and in one no less than 7. According to Gerhard, relapses, like the primary attacks, occur most frequently in spring.

NATURE OF CHOREA.—In considering the essential nature of chorea, it is evident that there are two points of importance to be determined, namely, the precise portion of the nervous system involved, and the nature of the morbid change in this part.

Before alluding to the views which have been entertained in regard to

the first of these questions, we would refer to the very great irregularity which exists in different cases in the extent and distribution of the choreic movements. Thus it frequently happens that the disease is strictly confined to one or the other side of the body, or it may be entirely symmetrical. In other cases the muscles of the head and neck may almost or quite escape, while both legs and one or both arms are affected. Or, on the other hand, the choreic movements may first appear and remain most severe in the muscles of the face, mouth, and tongue. It seems probable to us, therefore, that there is no one special portion of the motor centres which is exclusively the seat of lesion in all cases of chorea. In the great majority of cases, however, the symptoms are so far uniform that the muscles of the face and tongue, as well as those of the extremities, are affected, and the only peculiarity is that the irregular movements may be confined to one or the other side, a circumstance susceptible of ready explanation.

Marshall Hall considered chorea as an affection of the true spinal system, and possibly in some cases where the choreic movements are limited to the extremities and symmetrical, this supposition may be correct.

In the vast majority of cases, however, it is undoubtedly necessary to locate the seat of disturbance in chorea at a higher point in the cerebro-spinal axis, one above the decussation of the anterior pyramids, and probably in or near the corpora striata. Among the arguments which lead to this view, many of which have been advanced by J. Hughlings Jackson¹ and Broadbent,² who strongly uphold it, may be stated the following: That the muscles of the face are very frequently affected by the choreic movements; that in the great majority of cases the movements cease during sleep; that the affection is frequently limited to one side of the face and body, and that the spasmodic movements not rarely terminate in complete hemiplegia. In a footnote (*loc cit.*, p. 93) Tuckwell says: "It is just to Dr. Todd's memory to add, that he long ago (*Lancet*, 1843, vol. ii, p. 463) showed that the choreic phenomena cannot be explained by the hypothesis which refers them to irritation of the spinal cord." He says: "The hemiplegic tendency is utterly inexplicable according to that view. The affection of one-half the body would alone refer to some point above the decussation of the pyramids as the seat of irritation." The supposition of Carpenter and others that the cerebellum is the seat of the disturbance in chorea, was based upon the view that that organ possessed the chief power of co-ordinating muscular movements. Recent researches into the functions of the cerebellum, as well as the arguments which have been adduced above, render this supposition untenable.

The further question now remains as to the condition into which the affected part of the motor centres is brought, in order to produce the phenomena of chorea. And it is especially in regard to this point that the investigations of Jackson and Broadbent, above referred to, are of so much value. These pathologists, and particularly the latter, have called attention to the fact that the choreic phenomena are symptomatic merely of the

¹ Reynolds's Syst. of Med., Art. Chorea, vol. ii, p. 127, footnote; and *Med. Times and Gaz.*, March 6th, 1869.

² *British Med. Jour.*, 1869.

seat of the disease, and that the only essential condition of their production is an impairment of vigor and instability of the sensori-motor ganglia, a condition which may probably be induced in different ways.

We are now prepared to consider the manner in which the various causes of chorea may be supposed to act.

We have already seen that in a certain number of cases chorea is independent of any appreciable lesions of the nervous system. In some of these cases it is possible that the impaired nutrition of the motor centre may result from an altered and anæmic state of the blood: and, indeed, it appears to us quite as reasonable to explain a certain class of cases of chorea in this manner, as to apply the same explanation to analogous cases of paralysis.

It is probable, also, that in another group of cases, chorea may be *reflex* in character, and depend upon a different degree of that peculiar action upon the motor centres which produces reflex paralysis, whether by exhausting their excitability or by causing a reflex spasm of their vessels. The view is maintained by Broadbent (*loc. cit.*) as well as by Radcliffe,¹ who states that irregular choreic movements may be produced not only by injury of certain parts of the nervous system, but by injury of certain nerves at a distance from the nervous centres, the portions of the cerebro-spinal axis which are concerned in the development of such movements, being affected by reflex action.

It is probable that if this mode of production be admitted, it will serve to explain a large number of cases of chorea, both where the source of irritation is at a distance (as in cases of pregnancy, or where there are worms in the intestinal canal, or, as in the case already quoted from Packard, where a splinter was lodged in the matrix under a finger-nail) and where it is seated in immediate connection with the nervous centres. As instances of the latter kind, may be suggested such conditions as thickening of the meninges of the brain or spinal cord, and the presence of bony spiculæ developed in the meninges.

Finally, we must admit as a cause of chorea, primary alterations of the tissue of the sensori-motor ganglia and adjacent parts; the degree of disease not being so great as to abolish entirely their function and produce paralysis, but only sufficient (as for instance would be secured by an early stage of softening) to weaken it and render it unstable.

It will be seen from the foregoing remarks that we deem it impossible, at least in the present state of our knowledge upon the subject, to consider the cause and mode of production essentially the same in all cases of chorea, and that we are disposed to admit the existence of cases due to mere anæmia and impaired nutrition, or to an altered state of the blood; of cases due to reflex irritation, in both of which classes of cases, some minute and as yet inappreciable lesion may exist; as well as of cases which are due to primary material alterations of the sensori-motor ganglia.

We have already, in considering the causes and anatomical appearances of chorea, had occasion to dwell upon the connection which exists between

¹ Reynolds's Syst. of Med., Art. Chorea, vol. ii, p. 126.

it and rheumatism, and before leaving the present subject it is desirable to refer to the various explanations which have been offered of this circumstance. Among these, the most important and interesting is that of Kirkes,¹ who, noticing the frequent presence of vegetation upon the valves of the heart in fatal cases of chorea, was led to suggest that very small fragments of fibrin might be detached from the valves, and entering the circulation cause temporary obstruction of the minute capillaries of the nervous centres, producing irritation and impaired nutrition. This theory, which attributes the production of chorea to embolism, has been accepted by J. Hughlings Jackson (*loc. cit.*), by Savory,² by Tuckwell (*loc. cit.*), and in part, at least, by Broadbent (*loc. cit.*).

It is supported strongly by the facts that continued observation of cases of chorea has shown even more clearly the very frequent existence of cardiac murmurs during life, and of vegetations upon the valves after death; that complete paralysis, usually in the form of hemiplegia, frequently follows the choreic movements; that in many fatal cases there is found just such cerebral softening as follows embolism; and, finally, that in a few cases, already referred to, the existence of embolism has been actually demonstrated.

It is also to be borne in mind that recent researches have shown that endocarditis with the production of fine vegetations on the margins of the cardiac valves, is not a complication of rheumatism alone, but occurs in connection with scarlatina, diphtheria, and some other acute specific diseases. If then the theory is finally substantiated which would explain some cases of cholera as the result of minute embolisms, it is possible that it may be found that such is the mode of production of a part of the post-scarlatinal or post-diphtheritic choreas.

There have, however, been numerous objections advanced against this theory, the most powerful of which are urged by Barnes (*loc. cit.*) and Ogle (*loc. cit.*). Thus it has been objected that, on the supposition of numerous minute fragments of fibrin circulating in the blood and becoming impacted in the minute capillaries, it would be difficult to explain the fact that chorea is so frequently unilateral, or even localized in a single group of muscles. It must be remembered, however, in answer to this, not only that in some cases of fatal chorea embolism of single large arterial branches has been found, but that the number of minute fragments of fibrin detached from the heart's valves may be very small, and that it is quite supposable that they should in some instances nearly all pass into the innominate, or the left carotid artery, and thus be chiefly distributed to one side of the brain. It may be mentioned also in this connection, that it is especially in these cases of unilateral chorea that the affection is succeeded by paralysis, such as might readily follow in case of embolism.

Again, it has been objected that if chorea be invariably dependent upon embolism, the results of this accident must be of a very transient and trifling character, since in so great a majority of cases the disease termi-

¹ Med. Times and Gaz., 1863, vol. i, pp. 636 and 662.

² St. Barth. Hosp. Rep., vol. i, 1865, p. 107.

nates in complete and permanent recovery. The weight of this objection must be admitted, and yet Tuckwell fairly remarks in answer to it, that the "mere fact of recovery is not enough to condemn the notion of embolism. On the other hand, the very frequent presence of a cardiac murmur, even in the milder attacks of chorea which recover, would rather dispose me to look for the same exciting cause in the mild as in the severe cases, viz., embolism." It is evident also that if the supposed embolus were minute, and therefore obstructed only a very small vessel, a collateral circulation might soon be established and restore the nutrition of the area affected.

Another objection advanced by Ogle (*loc. cit.*, p. 232) is, that in other cases of capillary embolism the symptoms produced are not those of chorea, but rather of pyæmia or of gangrene. It is quite evident, however, that those symptoms alluded to (which are met with for instance in ulcerative endocarditis) are due, as remarked by Savory and Tuckwell, not to the mere capillary embolism, but to the concomitant septic condition of the blood. In this connection, reference may be made to the elaborate experiments of Panum as to the results of embolism (*Arch. f. Path. Anat.*, xxv, 308, 433; *Syd. Soc. Year-Book*, 1863, p. 211), in which he demonstrates that embolism of the vessels of the brain and medulla oblongata is followed by tetanic symptoms.

This extremely interesting question cannot be considered as definitely settled; there is still needed a series of careful examinations in regard to the various points under discussion. It appears to us, however, conclusively shown that, in a certain number of cases, the peculiar irritation and impaired nutrition of the sensori-motor ganglia, which leads to the development of the choreic phenomena, are due to embolism of the vessels supplying these parts. We have, however, already expressed our opinion that, at present at least, there must be admitted two other classes of cases of chorea, due primarily to alterations in the blood and to reflex irritation respectively. It is quite possible, therefore, that in some instances rheumatism induces chorea indirectly, either by causing anæmia and impaired nervous vigor, or by causing inflammatory lesions, as of the spinal meninges or sheaths of spinal nerves, which may serve as the foci of reflex irritation.

DIAGNOSIS.—The diagnosis of chorea cannot be attended with any difficulty, and we shall therefore make no remarks upon it.

PROGNOSIS.—Idiopathic simple chorea in young children is rarely a fatal disease. Nevertheless, even under these circumstances, it sometimes terminates fatally, and usually from exhaustion. Thus MM. Ruz, Legendre, and Rilliet and Barthez have each met with an instance. M. Sée (*loc. cit.*, p. 406) states that of 158 cases, 4 passed into the chronic condition, and 9 proved fatal. Dr. Copland states that he has met with 3 or 4 fatal cases, that Dr. Prichard has recorded 4, and that Dr. Brown refers to 3 in his practice; but he does not inform us whether they were idiopathic, complicated, or symptomatic. We have already referred to the list of 14 fatal cases, of which the autopsies were reported by Dr. Hughes (*loc. cit.*); and to the 34 additional fatal cases collected by Tuckwell (*loc. cit.*). Dr. J. W.

Ogle has (*Brit. and For. Med.-Chir. Rev.*, January and April, 1868) published the details of 19 more fatal cases; and from the same source we find that the mortality from this disease in Great Britain during 23 years was 1255.

It is quite possible, however, that many cases of organic disease of the nervous system merely attended with irregular choreoid muscular movements have been included in these reports. On the other hand, out of 84,332 deaths at all ages occurring in this city during seven consecutive years, but 3 are attributed to chorea. It must not, however, we think, be positively inferred from this that severe and fatal chorea has been really so rare among us; since, during the same time, there are reported in addition to the deaths from convulsions, 79 deaths from cramps, a vague and most improper term, which, in all probability, includes a certain proportion of cases of chorea.

In regard to any special rules in prognosis to be deduced from a study of the fatal cases, it may be observed that their average age is considerably greater than that of ordinary mild chorea. Thus in 17 out of Ogle's 19 fatal cases, but two were under the age of ten; the average being 15 $\frac{1}{4}$ ths years.

So too in 32 of Tuckwell's 34 fatal cases, 21 were at or above the age of fourteen, and 6 of this 21 were at or above the age of twenty.

The duration of the case scarcely seems to have a direct bearing upon its fatality. It is true that in cases which have passed into the chronic form and persisted for several months, the prospect of being able to effect an entire cure diminishes, but still such patients may live very many years, and ultimately die only from some intercurrent disease. And, on the other hand, death has been known to occur as early as the end of the first week. Of course the existence of any serious complication, and perhaps especially of marked cardiac disease from previous rheumatic attacks, renders the prognosis unfavorable.

Of 58 cases treated in the Children's Hospital of this city, the average duration of treatment was 39 days. Of this number, 42 were discharged cured, 13 much improved or almost well, and 3 without any improvement.

In conclusion, whenever, in a case of chorea, the convulsive movements become incessant, and the respiration embarrassed, and still more when subsultus tendinum takes the place of the choreic movements, a fatal termination is greatly to be apprehended.

TREATMENT.—Many different plans of treatment, and a great variety of drugs have been proposed for the cure of the disease under consideration. These facts alone may serve to teach us that the effects of treatment are not clearly appreciated, and also, when taken in connection with the circumstance that fatal cases are rare, that the disease tends naturally to recovery in a good proportion of the cases. This feature of the natural history of the disease is shown also by the evidence given by Dr. Bardsley, who mentions that in the Manchester Infirmary, notwithstanding the variety of treatment adopted by successive practitioners, an incurable case had not presented itself in the course of thirty-three years. (*Tweedie's Lib. Pract. Med.*, Am. ed., vol. ii, p. 46.)

The only rules to be laid down for its treatment are those which apply to all the convulsive affections depending on functional disorder of the nervous system, and on disordered states of the general health, connected with a faulty condition of the functions of digestion and assimilation. These are attention to the general health, and especially a careful regulation of the diet and other hygienic conditions of the patient, the removal of any local derangement or disease that may exert an unhealthy influence upon the nervous system, the use of tonics and iron, and the employment of such remedies as have been found to exert a controlling effect upon spasmodic and convulsive affections generally, and upon this disease in particular.

We shall consider, under different heads, the various means that have been recommended, endeavoring in the course of our remarks to distinguish the cases to which each remedy is best adapted.

Purgatives.—It was formerly the custom to rely largely on the use of purgatives in the treatment of chorea. In our own practice this never succeeded well, and for many years we have used laxatives, of the milder and less irritant class, only when called for by the existence of decided constipation. When the discharges from the bowels are clay-covered, or dark and offensive, when the mouth is pasty, the tongue loaded with a thick yellowish fur, and the breath heavy, it is proper to employ a mercurial.

Antispasmodics are amongst the most important remedies we have to oppose to the disease. The weight of evidence seems to show, indeed, that they, in conjunction with a moderate use of laxatives, of tonics, especially ferruginous tonics, and of certain particular remedies, and with careful regulation of the hygienic conditions of the patient, ought to constitute the treatment in the great majority of cases. Of the various remedies of this class that have been employed, those which have exerted the most beneficial influence are the root of the cimicifuga or black snakeroot, valerian, assafoetida, oxide of zinc, camphor, bromide of potassium, conium, and calabar bean.

Cimicifuga was first introduced into use by Dr. Jesse Young, and is now extensively employed and much relied upon. Dr. Wood (*Pract. of Med.*, vol. ii, p. 755), says: "I have in repeated instances found it of itself adequate to the cure of the disease." We have employed it ourselves quite frequently in primary cases, and in two cases of relapse. In several of the former the children recovered entirely under its use; in some, however, it failed to do any good, and recovery took place under the use of iron, arsenic, the sea-bath, and in the course of time. In the two relapsed cases, the patients recovered finally under the use of the cimicifuga, iron, cod-liver oil, and good diet. One of the cases that recovered under its use was among the worst we have ever met with. It was that of a boy of nine years, in whom the disease went so far as to destroy all power of locomotion. The child was unable even to stand. At the same time, the movements of the lips, cheeks, and tongue were so violent and irregular, and so little under the control of the will, that the power of speech was lost entirely for a period of four or five weeks. The choreic spasm appeared

to affect even the muscles of deglutition, so that the act of swallowing was often difficult and uncertain. Mastication also was impossible, and the child was unable to carry anything to its mouth, rendering it necessary to feed him, as one would a baby, with soft solids and fluids. During some two months, the muscles at the back of the neck were so weakened that the head could not be lifted from the pillow or held direct, but fell from side to side or forwards like that of an infant. The condition of the child was altogether one of the most complete and distressing helplessness. During the first month of the case it was treated with active cathartics, chiefly very large doses of cream of tartar and jalap, and with iron, but as the symptoms became worse and worse the cathartics were abandoned, except so far as to maintain, by the occasional use of rhubarb and senna, a soluble state of the bowels, which were very much disposed to constipation. The patient was now put upon the use of decoction of cimicifuga, of which he began with four ounces, soon increased to half a pint per day, made in the proportion of half an ounce to the pint. The iron was continued. Under this treatment he very soon began to amend, and in two weeks showed a very decided improvement. Cod-liver oil was now added to the iron and cimicifuga, and in six weeks he was in great measure restored to health, and in the end recovered completely. In another case almost as bad as this, the patient finally recovered under the same treatment.

The cimicifuga is given in powder, tincture, decoction, or fluid extract, and should be continued for several weeks in gradually increasing doses, until some visible effect is produced, as nausea, headache, vertigo, or disordered vision. The usual doses are from half a drachm to a drachm of the powder, from one to two ounces of the officinal decoction, and one or two drachms of a saturated tincture, given three times a days. For our own part we prefer the decoction, of which we give to children of eight or nine years old, from four ounces to half a pint a day, made in the proportion of half an ounce of the root to a pint of boiling water. Prepared in this way, it is not a disagreeable drink, and is usually taken without much objection.

The bromide of potassium in full doses has in some cases in our experience proved of marked benefit. We have used it especially in those cases which were connected with rheumatism as a cause, and have then frequently administered it in combination with the iodide of potassium and the iodide of iron.

We have not used valerian extensively in chorea, but from the evidence adduced in its favor there can be no doubt that it often exerts a very beneficial effect upon the disease. It may be given in the form of powder, infusion, or fluid extract. The dose of the powder is from twelve to eighteen grains in the day, to commence with, to be rapidly increased to several drachms, as the stomach becomes accustomed to it. It may be given in honey or preserve-syrup. We should prefer the fluid extract, of which half a teaspoonful may be given to a child eight or ten years old three times a day, and the quantity gradually increased. The oil of valerian is employed by some practitioners. Oxide of zinc is given

in doses of a grain every three hours to children eight years old, and is much relied upon by some practitioners. *Assafœtida* is recommended both by English and French writers. It is best given in pill, on account of the nauseous taste of the mixture. Two three-grain pills may be given to a child of four or six years of age, three times a day. Dr. Bardsley gave it by injection, in combination with laudanum, every evening, after using musk and camphor during the day.

Conium maculatum, given in the form of the *succus conii*, has been highly recommended by Dr. John Harley (*The Old Vegetable Neurotics*, London, 1869) in the treatment of chorea; and a certain number of cases have already been reported of its successful administration. Dr. Harley prescribes the succus in the doses of 20 or 30 drops for a child of six months old; a drachm for one over two years old; and from one to two drachms at ten years of age. In explaining the use of doses so large as these, he insists upon the fact "that hemlock given in doses which fall far short of producing its proper physiological action, is useless for the treatment of the diseases to which it is adapted." We have not had any extended experience ourselves with this drug in chorea, except of the chronic form in older subjects, but from our observation of its use in other conditions we should strongly advise beginning with much smaller doses, and gradually increasing as they are found to be tolerated.

Bouchut gives (*Bull. de Ther.*, April 15th, 1875, quoted in *Medical Times and Gazette*, June 5th, 1875) the results of numerous trials of eserine, the active principle of calabar bean, in chorea. It may be given either hypodermically or by the stomach: in the former way it is more energetic, and must be given in doses of $\frac{1}{25}$ th to $\frac{1}{15}$ th of a grain (3 to 5 milligrammes) three times a day; by the stomach, one-half as much more may be given. Its action is temporary, lasting two or three hours. It often causes, in the above doses, some uneasiness and restlessness, and occasionally nausea. It arrests the choreic movements during its action, and gradually effects a permanent modification in them, so that, according to Bouchut, the cure is effected in ten days on an average.

Narcotics have been recommended by some writers. Those which are most employed are opium, belladonna, stramonium, and cannabis indica. Substances of this class are seldom, however, made the basis of treatment. Opium is useful in some cases in which the agitation is very great, so that the sleep of the child is much disturbed, but it is seldom necessary except as an adjuvant to other means; and the remark applies equally to other remedies of this class.

Arsenic.—There is no remedy in regard to whose curative action in chorea testimony is more unanimous. Romberg and Begbie speak of it as curing the affection in as short a time and with even greater certainty than any other remedy; and Trousseau also testifies to its good effects, but adds that it has the disadvantage of being difficult of administration, owing to its irritant properties. Gerhard (*loc. cit.*) also speaks of it as having proved of marked benefit in his hands. Dr. Radcliffe, after meeting with the same difficulty in maintaining the use of full doses of this remedy for any length of time, tried with marked success the hypodermic injec-

tion of Fowler's solution. He was first led to employ this in cases of chronic local chorea in adults, where the injection of doses of Fowler's solution, varying from five to fourteen minims, produced a speedy cure. He also employed it successfully in two cases of general chorea, the duration being twenty-eight and thirty-two days respectively. This method of giving arsenic in obstinate cases of chorea has subsequently been tried in a sufficient number of instances to establish its positive value.

The usual manner in which we have administered it is in the form of Fowler's solution, given in the ordinary doses, and immediately after eating, and steadily persisted in until some of its constitutional effects are produced. By carefully watching for these, and immediately reducing the dose until the signs of irritation have passed away, and then again cautiously increasing it, we have usually been able to administer it without serious inconvenience, and with excellent results in a large proportion of cases. This preparation may also be advantageously combined with the bitter wine of iron. If, however, there is any individual peculiarity that makes it impossible to continue the use of ordinary full doses of arsenic; or if the choreic symptoms do not yield to this or the other remedies we have mentioned, we should recommend the use of arsenic hypodermically in larger doses.

Strychnia.—Trousseau recommends more highly than any other plan of treatment, the use of sulphate of strychnia in gradually increasing doses, until the extreme limit of tolerance is reached. He begins by giving gr. $\frac{1}{28}$ th twice or thrice daily, to children between five and ten years old, and cautiously increases this dose until it reaches about gr. $\frac{3}{4}$ th in twenty-four hours. The results obtained by this treatment in Trousseau's hands certainly appear good, but the risk attending it, and the care demanded to prevent accidents are so great, that we should prefer some of the equally successful and less dangerous methods. It appears, however, that other observers, as West, have obtained good results from its use in doses much smaller than those recommended by Trousseau, not exceeding gr. $\frac{1}{4}$ th thrice daily, for children of eight or ten years of age.

Stimuli.—The well known views of Dr. Radcliffe upon the pathology of spasmodic affections, have led him to recommend the free use of alcoholic drinks, to the point of obtaining their decidedly sedative action on the economy, as the foundation of a rational treatment in chorea.

Without being prepared to adopt this as a regular plan of treatment for ordinary cases of the disease, we should certainly be disposed to administer alcoholic stimuli whenever the symptoms indicated the approach of nervous exhaustion.

Tonics.—Whenever the disease occurs in debilitated and anæmic individuals, remedies of this class are evidently necessary, and prove of great efficacy. The ferruginous preparations are those most clearly indicated under the circumstances; and, indeed, there are many authorities, as Watson, Elliotson, and others, who consider the preparations of iron sufficient, of themselves, to cure almost all cases of chorea. Any of them may be selected. The best are the subcarbonate, Vallet's pills, the syrup of the iodide, and the pure metallic iron (ferrum per hydrogen). Quinine is also

recommended when the patient is feeble and weak. It may be given alone or in combination with iron. The citrate of iron and quinine would form a very good prescription under the circumstances mentioned. Cod-liver oil is an admirable remedy when the child is thin and weak, and especially when there is cause to suspect any tubercular predisposition.

A great variety of remedies besides those we have mentioned have been employed, and have more or less evidence in their favor. Amongst them are sulphate of zinc, nitrate of silver, subnitrate of bismuth, iodine, and a host of others which it is useless to enumerate. The sulphate of zinc has undoubtedly proved efficacious in some instances. About two grains may be given at first three times a day, and gradually increased to six or eight if the stomach bears the remedy well.

External Remedies.—The cold plunge and shower-bath as well as cold affusions to the nape of the neck and along the spine have been frequently employed as adjuncts to the internal treatment, and are of unquestionable value in many instances. The cases in which they are used should, however, be selected. They ought not to be employed unless followed by full reaction, nor unless the child is willing to take them. When the use of the baths terrifies or shocks the patient greatly, it cannot be proper. A warm or tepid bath used once a day, or every second day, would always be useful in promoting the general health, when the cold bath is not borne well. Ether spray has been recommended by Lubelski (*Gaz. Hebdomadaire*, April 19th, 1867) as an application along the spine, and a number of cases in which its use was successful have been placed on record.

Sulphurous baths have been recommended and employed with much success by M. Bandelocque, of Paris. A rapid and definite cure was obtained in 58 out of 65 cases. Thirty drachms of sulphuret of potassium are added to each bath, which is employed for at least one hour daily, at a temperature of 91°. Generally amelioration occurs after the second or third bath, but sometimes not until after twelve or fifteen days, a mean of twenty-two days having served for the cure of fifty out of fifty-seven cases. Where the cure is retarded, it ordinarily depends upon the patient's powers being lowered by other remedies or insufficient diet, upon irritation of the skin induced by the bath, or upon acute irritation of the internal serous membranes,—circumstances contra-indicating the baths while they continue. The conjunction of other remedies retards rather than aids the cure. Deducting the cases in which the bath was improperly used under the above circumstances, there remain but nine true failures in eighty-one cases, these being almost all recent or rheumatic choreas. (See on Chorea, *Ranking's Abstract*, No. 16, p. 51.)

Counter-irritation to the spine, in all its shapes, from pustulation with tartar-emetic, issues, and blisters, down to frictions with coarse towels, has been proposed and employed in the treatment. The use of any but the milder remedies of this class is unnecessarily harsh and cruel, except when the disease is evidently dependent upon an affection of the brain or spinal marrow. The great majority of cases will recover perfectly well without a resort to such violent means, and they ought therefore to be avoided.

Electricity has been resorted to, and apparently with good effects in

some instances, and it might therefore be tried when other and simpler means fail, or in conjunction with these means. In cases where the spasmodic movements are constant and persistent despite the use of internal remedies, the inhalation of anæsthetics has been tried, but with uncertain results.

In violent cases, it is of course desirable to confine the patient to bed; and it may be necessary to have padded sides made for it to prevent him from dashing himself out of bed in his uncontrollable and violent movements. In such cases it may even become necessary to employ padded splints, or to envelop the body with bandages carefully applied over layers of wadding, so as to secure the legs together, and to confine the arms by the sides.

Gymnastic Exercises.—M. Sée (*loc. cit.*, p. 481) says that this method is one of the best that have been employed. He states that it was recommended by Darwin, and then by Mason Good, and was first employed by Louvet Lamarre in one case, after which it fell into oblivion until some of the physicians at the Children's Hospital, at Paris, and amongst others, MM. Bouneau, Baudelocque, Guersant, and Blache, "struck, no doubt, like myself, with the good effects of gymnastics in scrofula and other cachectic diseases, and taught especially by the effects of musculation on the general health, conceived the idea of applying this treatment to nervous diseases, and particularly to chorea, which, besides the perturbation of the nervous system, is so often attended with disorders of nutrition and of the functions of organic life. To put a stop to this state of languor, to re-establish at the same time the equilibrium of the movements, which are rather irregular than convulsive, to endeavor, in fine, by regulating the contractions, to break up their vitiated habit,—this is the triple object sought to be attained by gymnastics. Be it theory or empiricism, success crowned these previsions, and proved the utility of the new treatment, of which we are about to study the methods and its consequences." M. Sée says, that to commence the treatment, we must prescribe first simple and cadenced movements, and exercise at the same time the larynx by means of singing. "To place the child in a vertical position, make it flex and extend the knees, touch the ground, stretch out and bend the arms, harmonizing at the same time these various movements by regulated singing,—such are the first means by which to replace the contractions under the power of the will. This end will be so much the more rapidly attained, as the attention of the patient is the less distracted, its intelligence the less changed, and its temper the less capricious; so also is it often impossible to succeed unless we first obtain control over the patient by kindness and gentleness."

"After reaching this point, we may attempt walking, regulated to a slow or quick step, running, jumping, hanging by the arms, or other more complicated movements, always graduating them to the degree of the disease, watching them most carefully, and repeating them daily without prolonging them beyond fifteen or twenty-five minutes, in order to avoid muscular fatigue and palpitation of the heart, which occur sometimes when the exercises are too long continued."

"With these precautions, and no matter how severe the symptoms, we may, after a few lessons, and sometimes after the first, and at latest after the fifth or sixth, perceive a manifest change in the abnormal mobility, which is usually so rapid that we are generally able to decide, after the first eight days, as to the efficacy of the treatment. When, after this length of time, the patient can neither stand erect, walk in a straight line, nor hang by the arms, there is reason to fear that the method will fail; it is at least certain that it will be tedious and difficult."

In *Ranking's Abstract* (*loc. cit.*, p. 50) may be found the following statements in regard to the treatment by gymnastic exercises:

They were first employed under the guidance of M. Laisné, gymnastic professor of the Polytechnic School, their effects being tried first on scrofulous children. "Commencing with simple movements of the legs and arms, accompanied by appropriate songs, the children's progress was so rapid that they were soon able to employ the orthopædic ladder, the parallel bars, and other machinery, in succession. By the twentieth lesson they were exercised in wrestling, and afterwards in running, special exercises being devised for the lame. From the first lesson the children became fired with emulation, and movements which seemed impossible were soon executed with ease and pleasure. A marked amelioration was speedily observed, their countenances becoming animated, their flesh firm, their voices stronger, their appetite keener and more regular; glandular swellings, which had long resisted all treatment, were resolved, and fistulous sores, that had been open for years, closed up. The lessons, one hour each, were given three times a week; and in the intervals the children amused themselves by repeating such of them as did not require machinery. The treatment, at first applied to scrofulous children, was, as stated above, extended to those laboring under nervous affections, partial paralysis, rickets, and especially chorea. Since 1847 ninety-five children suffering from chorea, sometimes so obstinate as to have resisted the most various treatment, have been cured by this means alone, or in conjunction with others, and no accident has resulted from the employment of the exercises. The movements are graduated according to the severity of the case, and they are repeated daily, but not for more than from fifteen to twenty-five minutes, so as not to induce fatigue or palpitation. "Improvement is sometimes seen after the first lesson, and at latest after the fifth or sixth; so that at the end of a week we can judge whether the means are likely to prove efficacious, and if manifest improvement has not then taken place, it is doubtful whether the cure will be thus effected, or if it is, it will be so only after a long time. The worst as well as the slightest cases have reaped equal benefit, the cure in the favorable ones only requiring a mean of twenty-nine days, and old or relapsed chorea being more amenable than recent. Dr. Sée has found that when other remedies are conjoined with the gymnastics, the proportion of cures is less, and the period of their attainment later; and he recommends no other adjunct to be employed than good diet." (Dr. Sée on Chorea, *loc. cit.*, No. 16, p. 50.)

HYGIENIC TREATMENT.—The management of the hygiene of the patient

is quite as important as any other part of the treatment. The diet should be arranged to suit the particular condition of the individual, and with a view to procure and maintain the most healthy possible state of the digestive apparatus. It should always be light and easily digestible, in order that neither the stomach nor bowels may be oppressed and deranged by the products of an imperfect digestion. When the stomach is weak and dyspeptic, the food ought to consist for some days chiefly of preparations of milk and bread, whilst in the meantime, a tonic remedy is administered internally, in order to invigorate the power of that organ. As the digestive function becomes stronger, the child ought, as a general rule, to be put upon the kind of diet most likely to promote the general health and vigor of body. It ought to consist of bread, milk, plain wholesome meats, and simple vegetables. Coffee and tea, and all other nervous stimulants, had better be avoided. The meats ought to be mutton, beef, or poultry. There are few vegetables, besides rice, potatoes, and tomatoes, which are suitable under the circumstances. All candies, preserves, unripe, coarse, or dried fruit, hot bread and cakes, except the very simplest, ought to be withheld.

Of dress we need merely say that it must be suited to the season. Exercise, or at least exposure to fresh air and insolation, are of the utmost consequence. When the disease is so violent as to prevent the child from walking, it ought to be taken to drive as often as possible. In cases which seem connected with a debilitated and anæmic condition of the constitution, removal to the country, and particularly to the seaside, will often effect a cure with great rapidity. Whenever, indeed, a patient inhabiting a large city or town can be conveniently taken to the seaside in the summer, it ought to be done, for the change is useful not only at the time, but it lessens, also, by strengthening and invigorating the constitution for the future, the danger of a relapse.

ARTICLE XI.

ATROPHIC INFANTILE PARALYSIS, OR POLIOMYELITIS ANTERIOR.

PARALYSIS occurs in the young child in almost, if not quite, all the forms observed in the adult. Many of these are, however, rare in childhood; whilst, on the other hand, there is one form which, although it is occasionally observed in the adult, occurs with such peculiar frequency in young children as to have received the name of infantile palsy. It is characterized by total or partial loss of power over one or several groups of muscles, usually without impairment of sensation, occurring suddenly as a rule, and often followed by atrophy of the palsied muscles, and consequent deformities.

HISTORY AND SYNONYMS.—Occasional allusions to infantile paralysis may be met with in medical writings even as far back as the latter part of the last century, but of such a vague and indefinite nature that the full

recognition and accurate description of this peculiar affection cannot be said to date further back than the writings of Kennedy¹ and Heine,² in 1836 and 1840 respectively. Since the publication of Heine's classical memoir, however, a number of observers have studied the disease with much attention and success. The vague and discordant views which have been held in regard to its cause and nature, have led to the employment of many names by which to designate it. Thus it has been called by Heine infantile spinal paralysis, and Meyer and others follow him in the use of this term; by Gull³ it was called paralysis during dentition; by Rilliet and Barthez,⁴ Vogt,⁵ Eulenberg,⁶ Valleix,⁷ Brunniche,⁸ Laborde,⁹ and Niemeyer,¹⁰ essential paralysis of children; by Duchenne,¹¹ who is followed by Echeverria,¹² fatty atrophic paralysis of infancy; by Reynolds,¹³ paralysis with wasting of the muscles; by Bouchut,¹⁴ myogenic paralysis; by Hammond,¹⁵ organic infantile paralysis; and it has also been called idiopathic and congestive infantile paralysis.

The names above enumerated appear to us to be either vague and inaccurate, as the terms essential and idiopathic; or to neglect one of the most striking features of the disease, the muscular atrophy, as the term infantile spinal paralysis does; or to convey a partial or even erroneous theory of the pathology of the disease, as the names congestive and myogenic respectively do. The terms organic and fatty atrophic paralysis also seem to us defective, since the first is equally applicable to cases of palsy due to organic disease of the brain, while the second is based upon the fatty degeneration of the affected muscles, which, however, occurs only in a portion of the cases of infantile paralysis.

In former editions we employed the term *atrophic infantile paralysis*, but with the statement that a term would doubtless be introduced, em-

¹ Kennedy, Observations on Apoplexy and Paralysis of New-born Infants: Dublin Jour. Med. Sci., 1836; and Dublin Med. Press, 1841; and Dublin Quart. Jour. of Med., 1850, and Nov., 1861.

² Heine, Beobach. ü. Lähmungszustände der unteren Extremitäten und deren Behandlung, Stuttgart, 1840; and Spinale Kinder Lähmung, Stuttgart, 1860; see Canstatt's Jahr., vol. iii., p. 70, 1860, and Med. Times and Gaz., London, 1863.

³ Gull, On Paralysis during Dentition, Guy's Hosp. Rep., 2d ser., vol. viii., pt. 1, 1852, p. 81.

⁴ Rilliet and Barthez, Traité des Mal. des Enfants, ed. 2ème, 1854, t. ii., p. 545.

⁵ Vogt, Essential Paralysis of Children, Berne, 1858, p. 86; New York Journal of Med., Jan., 1859, p. 117.

⁶ Eulenberg, On Essential Paralysis of Children, Virch. Arch., 1859, 177; and Schmidt's Jahrb., vol. 107, p. 55.

⁷ Valleix, Guide du Medecine Prat., ed. 4ème, 1860, t. i., p. 759.

⁸ Brunniche, ü. d. sogenannt. Essentiellen Lähmungen bei Kleinen Kindern., Journ. f. Kind., 1861.

⁹ Laborde, De la Paralysie (dite Essentielle) de l'Enfance, Paris, 1862, p. 122.

¹⁰ Niemeyer, Pract. Medicine (Amer. ed.), N. Y., 1869, vol. ii., p. 338.

¹¹ Duchenne, De l'Electrization Localisée, Paris, 1861, p. 275.

¹² Echeverria, Atrophic Fatty Palsy in Infancy, Amer. Med. Times, July 13, 1861.

¹³ Reynolds, Lancet, vol. ii., July 11, 1868, p. 35.

¹⁴ Bouchut, Des Maladies des Nouveaux-nés, ed. 4ème, Paris, 1862, p. 122.

¹⁵ Hammond, Organic Infantile Paralysis, N. Y. Med. Jour., Dec., 1865, p. 168; and Journ. of Psych. Med., vol. i., 1867, p. 49, and vol. ii., 1868, p. 531.

bodying a recognition of the seat and character of the anatomical lesion of this form of palsy, which would supplant all others. Since the localization of this lesion in the gray matter of the anterior columns of the spinal cord, the name of *poliomyelitis*¹ *anterior* has been rapidly coming into general use.

CAUSES.—The etiology of this affection is very obscure, doubtless partly owing to the fact that, as the paralysis occurs when the spinal system is extremely impressible, the causes which induce it are trivial and usually entirely overlooked. Age is the only influence which can be said to have a positive action in its production, since the great majority of cases occur between the ages of six months and two years, during the period of primary dentition. By several of the early observers especially, the disease was on this account attributed solely to dental irritation, but more careful observation shows that in most cases no such direct connection can be traced; and it is probable that early age and dentition only act indirectly by inducing a remarkably susceptible condition of the entire spinal system.

Sex appears to have no influence whatever upon its production; and the disease is almost as frequent among the children of the wealthy as among the ill-fed and ill-tended children of the poor. In some few cases, where the loss of power is sudden, the exciting cause seems to be the direct exposure to the local action of cold, as from sitting upon a stone step (West), or lying on the damp ground (Hammond).

Atrophic infantile paralysis is usually primary, and occurs in the midst of good health; but it has also been observed in a secondary form, appearing during the convalescence from measles, scarlatina, or typhoid fever, or during rheumatism and chorea.

In one of the cases following chorea, which are recorded by Kennedy (*loc. cit.*), it is positively stated that there was a distinct cardiac murmur, due to organic valvular disease; and it may be suggested that the essential cause of the paralysis was embolism of some of the spinal arteries, as observed by Panum.²

MODE OF ATTACK; INITIATORY SYMPTOMS.—There is considerable variety in the mode in which this disease makes its appearance. In some cases the paralysis is the first symptom observed, and is found to have almost immediately attained its full extent, without any recognizable cause or premonitory symptom. Thus the child may have appeared perfectly well when put to bed in the evening, and yet on the following morning, there may be more or less complete loss of power over the lower extremities. But in the great majority of cases, especially the more severe ones, the attack is preceded by quite marked constitutional disturbance. This may consist merely of fever, appearing without evident cause and lasting from a few hours to a week or more, unattended by any

¹ Πολιός, gray.

² Ueber den Tod durch Embolie (Bibliothek für Läger, 1856), quoted by Jaccoud (*op. cit.*, p. 297), and Arch. f. Path. Anat., xxv, 308, 443, 1863, in Yearbook of N. Syd. Soc., 1863, p. 210.

gastro-intestinal disturbance. Or, during this period, the child may also complain of pain in the back, or there may be tenderness on pressure, especially in the lumbar region; there is frequently slight dulness of the mind; and finally, in comparatively rare cases, one or more, convulsions may occur. It is the rule, however, for no marked symptoms of cerebral disturbance to be present at any period of the disease. There are rarely any symptoms connected with the parts about to become paralyzed, though in an interesting case recorded by Kennedy (*loc. cit.*), there was spasm of the muscles subsequently affected.

The disease usually makes its appearance during health, but it is probable that many of the cases of paralysis occurring during convalescence from the various exanthemata properly belong to this variety.

Whether preceded by initiatory symptoms or not, the development of the paralysis is generally sudden, and it is only in rare cases that it is partial at first and increases gradually. Indeed it usually happens that when first observed the paralysis is at its maximum, both as regards the number of muscles affected and the degree of the loss of power, and that there soon occurs a diminution in its extent, so that only some of the parts first affected remain palsied.

The form of the paralysis clearly indicates its spinal origin. Complete hemiplegia is scarcely ever observed, though in a few cases the arm and leg of the same side, or even all four extremities, have been palsied. Most frequently the disease takes the form of incomplete paraplegia; though occasionally the paralysis affects single groups of muscles or even individual muscles.

According to Mr. Adams,¹ the groups of muscles most frequently affected are: 1. The muscles of the anterior parts of the leg, forming the extensors of the toes and the flexors of the foot; 2. The extensors and supinators of the hand, these muscles being always affected together; and 3. The extensors of the leg, and with them generally the muscles of the foot, as in the first group. When single muscles are affected, the most likely to suffer are these; 1. The extensor longus digitorum pedis; 2. The tibialis anticus; 3. The deltoid; and 4. The sterno-mastoid.

The bladder and rectum are scarcely ever involved. In very rare cases (5 out of over 1500)², also, the abdominal muscles are palsied, giving rise to marked protrusions of portions of the abdomen. The muscles of the back are more frequently involved.

The degree of the paralysis varies as much as its extent; usually complete at first, in some cases it soon becomes partial or even slight; while in others the loss of power remains absolutely complete. The paralyzed muscles are perfectly relaxed, so that the affected parts can have all their normal movements impressed upon them without difficulty, and fall in a lifeless manner if left unsupported. The special senses are unimpaired; and general sensibility is usually only blunted for a time. Occasionally it is not affected at all, or, as stated by West, there may even be hyperæsthesia for a variable time.

¹ Adams, On Club Foot, London, 1866.

² Birdsall, Jour. Nerv. and Ment. Dis., vol. viii, July, 1881.

The paralyzed muscles are rarely the seat either of painful subjective sensations or of tenderness on pressure ; though in some cases severe pain may be present in the affected parts.

Reflex movements are, as a rule, abolished in those parts where there is complete loss of voluntary motion ; though Laborde (*loc. cit.*) has shown that they may occasionally be preserved even in the first stage of the paralysis.

During the early stage we are at present considering, the electro-muscular contractility usually remains intact, and the muscles respond both to the induced and direct current.

The constitutional disturbances which we have described as preceding the paralysis may persist for a variable time after its development, or disappear quickly, leaving no other symptoms present but those connected with the paralyzed parts.

The following case may be quoted as an illustration of this form of paralysis.

A male child, æt. thirteen months, was brought for treatment by its mother, an intelligent woman with several healthy children. The following history of the case was obtained : The little boy had walked at the age of nine months, and always seemed a vigorous, bright child ; he had also cut eight teeth, without much irritation. About September 10th, 1868, after no particular exposure, he became fretful and feverish, with occasional vomiting ; and after three days it was noticed that right-sided hemiplegia had developed itself. The paralysis of the arm was never complete, while the leg had entirely lost all power of motion. This loss of power had not become complete suddenly, but, at first partial, had gradually increased. There was no tendency to coma and no evidence of any acute pain. The febrile symptoms soon disappeared ; the arm regained the power of motion in a few days, but the leg remained palsied. It also soon grew remarkably cold, and when seen on October 1st, three weeks after the attack, the temperature was decidedly lower than that of its fellow. Sensation was impaired, but had never been abolished. There had been no paralysis of either bladder or rectum. At the time of the examination the child seemed bright and lively, though rather pale. There was no tenderness along the spine, nor in the leg. No reflex movements were developed in the paralyzed leg by tickling the sole of the foot. Neither atrophy nor deformity had as yet occurred.

The subsequent course of the disease varies greatly in different instances. In one set of cases, though the paralysis may be quite extensive and complete at first, the symptoms gradually subside, the paralysis disappears, and complete recovery ensues in from four to six weeks. These cases correspond exactly to the form of paralysis originally described by Kennedy (*loc. cit.*) under the name of "Temporary Infantile Paralysis," and, as we shall see hereafter, in all probability depend upon mere congestion of the spinal cord.

In the other set of cases, on the contrary, the loss of power persists, and after it has continued for a time, varying from one to several months, is followed by marked and more or less rapid atrophy of the affected muscles. The circulation in the paralyzed parts becomes feeble, the subcutaneous veins are smaller, and Heine, and Rilliet and Barthez each cite a case of paralysis of the arm in which it was almost impossible to detect the radial pulse. The temperature of the affected part becomes

perceptibly lower, the fall amounting, according to Hammond,¹ to from 5 to 8, or even 10 degrees, as tested by a galvanometer. The muscles themselves undergo marked atrophy, frequently accompanied by fatty degeneration; and their reflex motility and electro-muscular contractility disappear. It is important to notice, however, that long after muscular contractions fail to be produced by the induced current, they may frequently be excited by the use of a direct current of low tension, slowly interrupted.

The mere wasting of the muscles is not, however, the only cause of the great difference in size between the healthy and paralyzed members. The nutrition of the whole limb is affected, and the growth and development of all its tissues arrested, so that the paralyzed member becomes smaller in all its dimensions than its fellow. Rilliet and Barthez cite an example which they observed, to show to how remarkable a degree this conjoined atrophy and arrest of development may progress. The patient was a young girl who was seized with instantaneous paralysis of the right lower extremity; and the following measurements show the degree of inequality which was produced by four years' continuance of the paralysis and arrest of development.

	Right leg.	Left leg.
1. From the great trochanter to the external malleolus,	49 cent.	54 cent. 5 mill.
2. From the patella to the malleolus,	29 "	32 "
3. Length of foot from heel to great toe,	14 " 3 mill.	18 "

Five months previously, the following diminution in thickness of the limbs was noticed: at three fingers' breadth above the patella, on left side, 20 centimetres, 16 on right; at the middle of the thigh, on left side, 29 centimetres, and 22 on right. The height of the child was 116 centimetres.

This wasting and palsy of the muscles is associated with relaxation of the ligaments, and the combination of these causes induces many of the deformities observed in childhood. When the paralysis affects one side of the body chiefly, it indirectly leads to various lateral curvatures in the spinal column, probably from a want of symmetrical action in the muscles of the two sides.

In cases of paralysis of the arms, the relaxation of the ligaments about the shoulder-joint and the atrophy of the deltoid allow the head of the humerus to drop out of the glenoid cavity, so as to produce even complete dislocation, with apparent elongation of the paralyzed limb to the extent of three-fourths of an inch (West).

As the muscles of the lower extremities are far most frequently affected in this form of paralysis, we usually find the resulting deformities involving the feet and legs, where they constitute the greater proportion of all cases of club-foot. According to Adams (*loc. cit.*), "these deformities occur in the following order of frequency: 1, Talipes equinus; 2, equino-varus; 3, equino-valgus; 4, calcaneus or calcaneo-valgus; and 5, talipes varus. When both feet are affected, equino-varus of one foot is generally found with equino-valgus of the other."

In addition to the influence which the actual wasting of the limb and

¹ Dis. of Nervous System, 1871, p. 690.

the arrest of its development exert, Adams believes that the great cause of such deformities is the "adapted atrophy" of Paget, the changes which ensue in consequence of the mechanical relations of the foot to the leg. Although, however, it is true that paralysis of a group of muscles does not excite active contraction in their opponents, it appears that in the efforts of the child to move the part, the non-paralyzed muscles must gain control over the limb, and aid at least in producing the various characteristic distortions.

During the development of this atrophic stage, the general sensibility of the affected parts is usually normal, and the general health, intelligence, and nutrition of the patient unimpaired.

DURATION.—As will be inferred from our description of the course of this affection, the entire duration and that of its different stages varies greatly in different cases. In some, which have hence had the name "temporary" infantile paralysis bestowed upon them, the loss of power rapidly diminishes, and complete recovery follows in from a few days to a few weeks; while, in other cases, the paralysis persists until atrophy ensues, and the limb may remain crippled and useless throughout life. The period which elapses before atrophy commences, and the rapidity with which it advances, also vary extremely, even in apparently similar cases. Thus the palsied muscles may begin to atrophy within four or five weeks, though more frequently this change cannot be noticed for several months. Different muscles also atrophy with very different rapidity, the deltoid and tibialis anticus appearing to waste more rapidly than any other muscles of the body; and, in different cases, the same groups of muscles show equal variety in this respect, a few weeks serving in some instances for as much wasting to occur as would require months to produce in other cases.

PROGNOSIS.—The great uncertainty of the progress and duration of atrophic infantile paralysis renders it highly desirable to ascertain, if possible, the conditions which determine its result. Of itself, it is never fatal; but, unfortunately, our prognosis is limited, in the early stage of the disease, to this assertion, for the duration and course of the case are not influenced, in any constant and reliable way, either by the age of the patient, the extent of the paralysis or the parts affected, or the initiatory symptoms. It may perhaps be stated that, in general, cases which are ushered in by high fever, especially if associated with convulsions, and in which the paralysis is extensive, will prove severe and tedious. But there are too many exceptions to every particular of this statement for it to be regarded as a general rule of much positive value in prognosis.

When paralysis has lasted three or four weeks, we are able to determine with much accuracy the approach of atrophy by the condition of the electro-muscular contractility; for it has been frequently observed that those muscles which lose their power of responding to the interrupted current, soon begin to waste.

After the occurrence of atrophy, also, much valuable aid in prognosis is gained from the use of electricity.

We may here mention the interesting and highly important observation, first made in connection with this disease by Hammond (*loc. cit.*) and J.

Netten Radcliffe,¹ that in many cases where the atrophied muscles have lost entirely their power of reacting to the most powerful induced electrical currents, they will still react vigorously to a direct (galvanic) current of low tension and slowly interrupted. The importance of this discovery, in the treatment of the disease, can scarcely be overrated; and it has also enabled this point to be established in the prognosis, that whenever muscular contractions can be excited by either induced or direct currents, no matter how far advanced the atrophy of the muscles, the restoration of their power can certainly be accomplished; though it would appear from a case successfully treated by Hammond, that even when such contractions are not at first produced, the prognosis is not absolutely unfavorable. The still more curious, and as yet inexplicable observation has also been frequently made, that as the muscles regain their power of voluntary motion, their susceptibility to the direct galvanic current is apt to diminish, but, on the other hand, their normal reaction to the induced current returns.

The prognosis will also be materially influenced, especially when the atrophic stage has begun, by the condition in which the tissue of the palsied muscles is found, as in cases where advanced fatty degeneration is present, it is far more unlikely that they will ever regain their power. In order to ascertain this point, Duchenne has devised a small trocar,² called by him "emporte-pièce," by which small pieces of muscle can be extracted, and subsequently submitted to microscopic examination.

It is evident, finally, that the duration and result will depend, to a great extent, upon the period at which treatment is instituted. In those cases where the paralysis has been allowed to continue until marked atrophy has ensued, and the electro-muscular contractility is almost lost, although the prognosis may still be favorable as regards the ultimate cure, it must be carefully guarded as to the duration, since the treatment will probably require to be steadily pursued for many weeks, or even months.

MORBID ANATOMY AND PATHOLOGY.—It appears desirable to introduce the consideration of the anatomical appearances at this point, in order to facilitate the subsequent discussion of the pathology and diagnosis of the disease.

In regard to the changes which take place in the atrophied muscles, the brief yet complete summary given by Hillier³ may be quoted:

"1. The transverse striæ become less apparent and separated by wider spaces, which are filled with opaque granules, which are not dissolved by ether, but are sensibly acted on by acetic acid.

"2. The transverse striæ disappear, and there is an abundant appearance of granular substance.

"3. There remain but slight traces of longitudinal fibres, filled with granules, with a larger quantity of connective tissue between the bundles.

"4. The granules have disappeared, and empty transparent tubes of

¹ See footnote to page 665, vol. ii, Reynolds's System of Medicine.

² These trocars are manufactured by Tiemann, of New York. Dr. Hammond has published (*Jour. of Psych. Med.*, July, 1867) a description of their form and mode of use, illustrated by a woodcut.

³ T. Hillier, *Diseases of Children*, Philadelphia, 1868, p. 255.

myolemma with a few scanty granules on their walls remain, with more connective tissue and some elastic fibres.

"5. In some cases, fat globules take the place of the granular matter in the muscular fibres, and in the cellular tissue between the bundles of muscular fibre. This change is not universally present in cases even when atrophy has proceeded to an extreme degree."

The last conclusion stated here, which has been confirmed by other observers, shows that perhaps the most frequent change which occurs, is a simple atrophy of the muscles, with a granular but non-fatty degeneration, and conclusively shows the inaccuracy of the name proposed by Duchenne for the disease (namely, fatty atrophic paralysis of infants).

In approaching the question of the lesions of the nervous centres in this affection, which have now been definitely determined, it is necessary to refer to the general question of the existence of so-called essential, purely neurotic paralyses. In one form of paralysis, the reflex, it is true that as yet no material lesion has been detected, and that the most plausible explanation of the loss of power in such cases is simply the exhaustion of the functional activity of the spinal cord, owing to the prolonged irritation of some of the peripheral nerves. And it must be borne in mind that the form of infantile paralysis under consideration was formerly by some regarded as a reflex paralysis depending on dental irritation. Apart, however, from the fact, that the symptoms much more closely resemble those due to spinal congestion than those seen in reflex paralysis, it is to be remembered that the disease is by no means limited to the period of dentition, and that all local signs of dental irritation are frequently absent at the time of the appearance of the paralysis. With the exception, then, of reflex paralysis, it may be asserted with confidence that all other forms of spinal paralysis are associated with some material lesion of this nervous trunk. It is to be remembered that it is only a few years since the beautiful researches of J. Lockhart Clarke have shown that positive structural changes, in both nerve-cells and nerve-fibrils, may be detected by microscopic examination in spinal cords, which present no alteration apparent to the naked eye. In rejecting the evidence of all post-mortem examinations of the spinal cord, made before the introduction of Clarke's method, as incomplete and inconclusive, we find that in all those diseases formerly classed as pure neuroses (such as tetanus and chorea), which have been subjected to this latter mode of examination, positive demonstrable lesions have at least occasionally been detected.

Among this class of diseases, so long considered as purely functional neuroses, atrophic infantile paralysis has always, until lately, occupied a prominent position, as is evinced by the large number of authors who have described it under the terms "essential," or "idiopathic."

It is indeed difficult to secure opportunities of examining the state of the spinal cord in this affection, owing to the fact that the disease is scarcely ever, if at all, fatal of itself; so that the arguments in opposition to the view of its functional nature, will be in part drawn from the close analogy of its symptoms to those of certain spinal diseases which are well known to be attended with positive lesions of the nervous tissue. Thus, in its

mode of appearance, and in the character of the paralysis, there is so perfect a resemblance to the onset and symptoms of congestion of the spinal cord, as to leave little room for doubt that this is the condition at first present in many cases of atrophic infantile paralysis. In both this affection and spinal congestion, the paralysis may appear quite abruptly, or be preceded by pains in the back and fever; in both, the paralysis is usually paraplegic, the loss of power only partial, and the affected muscles are relaxed; in both, general sensibility is but slightly impaired, the bladder and rectum are not involved, and there are no disturbances of the cerebrum or special senses; in both, finally, recovery usually follows, if proper treatment be promptly instituted.

In those cases where the paralysis disappears within a few days or weeks, it has been supposed by various authors that the nature of the disease is entirely different from that of atrophic infantile paralysis; but it appears to us highly unnecessary to complicate the question by such a supposition; since the temporary character of the paralysis is readily accounted for by supposing that the spinal congestion which produced it was slight and transient.

It is quite possible also that in other cases the loss of power caused by more severe spinal congestion should persist until atrophy of the affected muscles ensued, and rendered the case more protracted.

Indeed, some of the authors who most forcibly support the view of the pathology of this affection which we have given above, as Dr. C. B. Radcliffe (*loc. cit.*), hold that the lesion of the cord does not advance beyond this stage of congestion. The evidence in support of this opinion is principally found in the result of post-mortem examinations, as those reported by Rilliet and Barthez, Fliess and Adams, where no lesions of the cord were detected. But in none of these cases does it appear that the careful and skilful microscopic examination, which is now recognized as necessary to detect some lesions of the nervous tissue, was performed; so that we may feel at liberty to doubt the complete accuracy of these autopsies. On the other hand, it certainly seems entirely consistent to suppose that in certain cases, where the congestion is unusually marked and prolonged, or where it is repeated, that a process of subacute inflammation should be excited, resulting in the permanent structural change.

The usual change which takes place in the spinal cord, under such circumstances, is that described under the name "sclerosis," in which there is marked proliferation of the connective-tissue elements of the cord, with swelling and consequent pressure upon the nerve-tubules. In the subsequent development of the new-formed connective tissue, it undergoes contraction, and induces atrophy of the compressed nerve-tubules. The portions of the spinal cord where this lesion exists, may either be atrophied or retain their normal size, shape, and external appearance, but on transverse section, though the tissue is firm, certain parts of the white substance are seen to present a grayish, translucent appearance, differing noticeably, in well marked cases of the lesion, from the opaque whiteness of the surrounding healthy tissue. In other instances, however, the change in color cannot be detected, and it is only by microscopic examination that we can

discover the increase in the connective tissue of the cord, and the atrophy of the nerve-tubules.

This view of the nature of the lesions in atrophic infantile paralysis was forcibly urged by Heine, in the last edition of his classical monograph on this subject (*op. cit.*), who based it merely upon an analysis of the symptoms, and it has since been adopted by Jaccoud (*loc. cit.*). It does not rest, however, solely upon such reasoning, for there have been a limited but rapidly increasing number of autopsies made in which the lesions of sclerosis above described have been actually observed.

Heine quotes three post-mortem examinations in support of this theory. One of these, quoted from Longet, was of a girl of eight years, with club-foot on the right side, following an attack of paralysis, who died of variola; and at the autopsy the muscles and nerves of the right leg were atrophied, and the anterior roots of the spinal nerves which make up the right sciatic nerve, were scarcely one-quarter the size of the corresponding roots on the left side.

In the second case, quoted from Hutin, the subject was forty-five years old, had been paraplegic from the age of seven years, and had considerable deformity of the lower members; at the autopsy, after death from dysentery, there was atrophy of the lower part of the spinal cord.

The third observation quoted by Heine, has been quoted more fully from the original source (*Trans. de la Soc. Med. de Berlin*, Dec. 7th, 1862), by Jaccoud (*op. cit.*, p. 450). It was the autopsy of a child with paralytic club-foot, reported by Berend and Remak, where the "spinal arachnoid was found thickened by inflammatory product, and exercising such pressure upon the cord, that when the false membranes were cut, the nervous tissue immediately protruded through the incision."

Berend also reported (*id. loc.*) another observation upon a child four years old, who died paraplegic with contraction of the legs and feet. The autopsy was preformed by Recklinghausen, who found tubercles in the cord.

Hammond reports (*Jour. of Psych. Med.*, vol. i, p. 51) a case where the paralysis affected the left leg, and had lasted four years, in which he found, upon post-mortem examination, a cicatrix, partly filled with clot, in the lower part of the dorsal region, in the left anterior column. Recently, however, the opportunities for careful study of the lesions in atrophic infantile paralysis have multiplied, and have been seized by numerous able observers, especially in France, where the first demonstration of the true characteristic morbid changes in this disease was effected. The earliest cases placed upon record in which this lesion was accurately described were by Cornil (*loc. cit.*), in 1863; by Laborde (*loc. cit.*), in 1864; by Prevost (*loc. cit.*), in 1866; J. Lockhart Clarke (*loc. cit.*), in 1868; Charcot and Joffroy (*loc. cit.*), in 1870; Parrot and Joffroy,¹ in 1870; Roger and Damaschino,² in 1871; Dujardin-Beaumetz,³ in 1872; Petitfils,⁴ in 1873; and

¹ Arch. de Physiologie, tom. iii, 1870, p. 135.

² Gaz. Méd. de Paris, 1871.

³ De la myélite àigue, Paris, 1872.

⁴ Considérations sur l'atrophie des cellules motrices, Paris, 1873.

numerous other observers have confirmed their results, so that the morbid anatomy of atrophic infantile palsy may be regarded as clearly and fully determined.

The lesions occupy the antero-lateral columns, and especially the anterior horns of gray matter. There is atrophy of the nerve-fibres in the anterior and lateral columns, which varies in amount in different cases, and is associated with a varying degree of hypertrophy of the interstitial connective tissue (sclerotic). These parts are more translucent than natural, and often present a very appreciable grayish rose tint to the naked eye. The consistence of the affected tracts is diminished, and upon microscopical examination there may be observed a marked proliferation of the elements of the connective tissue, the cells and nuclei being dispersed in the midst of a finely granular substance, in which there are fibrils of extreme tenuity. In the parts which are most affected the nerve tubules are either lost altogether, or they present a varicose appearance, while the other portions of the spinal column preserve a perfect integrity.

But the most characteristic changes are found in the anterior horns of gray matter, where there is invariably atrophy of the ganglion nerve-cells and of their processes, so that in some instances the anterior group of cells has entirely disappeared from atrophy. In other cases the remains of the cells are found atrophied, misshapen, and with granular degeneration of their contents. The other elements of the gray tissue are usually changed also; there is proliferation of the nuclei of the neuroglia, and occasionally increase in the delicate fibrils of this connective tissue. In some cases the walls of the vessels in the affected parts are found thickened, with proliferation of their nuclei. These changes have been so prominent in some cases as to have led to the opinion (Damaschino,¹ Duchenne) that they constituted the primary and essential lesion. This, however, does not seem probable. It will be seen, therefore, that the name poliomyelitis anterior is, as we have already stated, strictly appropriate.

The progress of anatomical investigation has thus at last developed the true pathology of this affection. It is possible that in some cases the lesion of the antero-lateral columns may be the result of hæmorrhage into the substance of the cord, or of pressure from thickening of the meninges; but in the vast majority of cases the morbid process is one of slow subacute inflammatory, sclerotic change, with atrophy of the nerve-tubules in the antero-lateral columns and anterior horns of gray matter, and especially with atrophy and destruction of the anterior groups of ganglion nerve-cells.

DIAGNOSIS.—There is but little danger of overlooking the nature of those cases where the paralysis appears quite suddenly in the midst of apparent good health, excepting in cases occurring in young children who have not yet learned to walk, and where the loss of power is limited to the lower extremities. In such instances the paralysis may be entirely overlooked by the parents or nurse for some time. So also in cases preceded by constitutional disturbance, as there is nothing whatever characteristic in these premonitory symptoms, it is quite possible to fail to rec-

¹ Damaschino and Royer, *Gaz. Med.*, 1871, p. 457.

ognize the presence of paralysis. It is well, therefore, whenever a child between six months and three years of age presents feverish symptoms for which no apparent cause exists, to ascertain carefully whether there is any loss of power of its extremities.

The diseases with which atrophic infantile paralysis is most likely to be confounded, are other forms of paralysis of cerebral or spinal origin, and progressive muscular atrophy.

In paralysis due to hemorrhage into the substance of the brain (see page 546), the case is more apt to be ushered in by delirium or convulsions, followed by more or less marked coma, while in atrophic infantile paralysis there is either entire absence of cerebral symptoms, or at most a single convulsion occurs. Cerebral paralysis is usually hemiplegic, while in the form of spinal paralysis we are considering, paraplegia is more common, or the loss of power may be limited to one leg or to a single group of muscles. In those comparatively rare cases where the paralysis is at first hemiplegic, the arm usually soon regains its power of motion, leaving the leg paralyzed; while the reverse of this occurs in cerebral hemiplegia, where the leg usually improves much more rapidly than the arm. In cerebral paralysis, also, the affected muscles are frequently rigid instead of being relaxed; and there is not the tendency to atrophy and deformity, the loss of electro-muscular contractility, nor the lowering of the temperature of the affected part, which are observed in atrophic infantile paralysis.

In cases of meningeal apoplexy, where the hemorrhage has occurred *upon* the surface of the brain, the symptoms are still more distinct. Thus (see page 546) there are usually repeated convulsive seizures, with somnolence during the intervals; paralysis is rare and partial, while strabismus and tonic contraction of the hands and feet are very common.

In acute inflammation of the spinal cord, or myelitis, the loss of power is complete, and there is also more marked loss of sensation and paralysis of the rectum and bladder, with alkaline urine; though there is here as well as in atrophic infantile paralysis, diminution of reflex excitability and electro-muscular contractility, and wasting of the paralyzed muscles. The symptoms first mentioned, the more grave character of the case, and the tendency of the paralysis to increase rather than decrease, suffice to distinguish myelitis from the affection under consideration.

Progressive muscular atrophy, of very rare occurrence in children, may be distinguished by its gradually progressive course; and by the preservation of the temperature of the affected parts, of the power of motion, and of electro-muscular contractility, until atrophy has far advanced. There is usually a quivering of the atrophied muscles in this disease, due to fibrillar contraction, which is entirely wanting in atrophic infantile paralysis.

We have already expressed our belief that some of the cases where the loss of power is very temporary, are really instances of reflex paralysis, and in such some source of peripheral irritation can usually be detected.

West alludes to the fact that in those cases where the affection is limited to one leg, and attended by hyperæsthesia and painful sensations, the dis-

ease may be mistaken for coxalgia, though the diagnosis may readily be made by attending to the slow course, the absence of paralysis, the fixed pain in the knee-joint, and the marked increase of suffering caused by forcing the head of the femur against the acetabulum, which characterize hip-disease.

TREATMENT.—The treatment of atrophic infantile paralysis may be divided into that adapted to the early stage and that directed against the second stage or period of atrophy.

In the first instance we must endeavor to discover and remove any exciting cause of the paralysis that may exist. If symptoms of morbid dentition have preceded, and the appearance of the gums indicate it, they should be lanced; or if gastro-intestinal disturbance is present, or the presence of worms is suspected, laxatives should be administered. Tepid baths are also recommended, as tending to allay irritation and reduce feverishness.

When, however, no local irritation can be detected to render it possible that the case is one of reflex paralysis, we should direct our remedies towards relieving the spinal congestion, which we believe to exist in cases of true atrophic infantile paralysis. Counter-irritation should be applied along the spine, and may be effected by producing a narrow blister, or preferably by the use of sinapisms or stimulating liniments, containing croton oil, ammonia, or turpentine.

Local abstraction of blood by means of cups or leeches applied along the spine has been recommended by Fliess; and we should certainly advise its employment, especially in those cases where there is considerable febrile disturbance and pain in the back.

There are also certain remedies from which we have obtained excellent results in the treatment of spinal congestion in the adult, and should, therefore, recommend their employment in the early stage of this affection.

These are ergot, which may be given in the form of fluid extract, beginning with doses of 5 to 10 minims for a child of two years old; and belladonna, which may be given either in the form of tincture, or an aqueous solution of the extract. Iodide of potassium may also be given in combination with one or the other of these, in doses of gr. j or ij for a child of two years old, in the hope of preventing the development of any inflammatory changes in the cord.

In addition to these remedial measures, the child should be absolutely confined to bed.

If, despite the use of these agents, the paralysis persists, the temperature begins to fall, and the muscles to atrophy, every means must be adopted to promote the general nutrition of the child so as to favorably influence indirectly the changes in the spinal cord; and, at the same time, local treatment must be instituted to promote the circulation and nutrition of the paralyzed parts.

Among the internal remedies, iron is one of the most suitable, and may be given in any eligible form. The pyrophosphate is perhaps especially indicated on account of the phosphoric acid with which the iron is combined.

The various preparations of *nux vomica* or its alkaloid *strychnia* are also very valuable after the acute stage has passed. Heine advises the use of *tr. nucis vomicæ* in combination with camphor and pyrethrum; while West recommends the alcoholic extract of *nux vomica*. *Strychnia*, which is more frequently employed than the preparations of *nux vomica* itself, is usually given in the form of solution. Hillier has also used it hypodermically, but without marked benefit.

The doses of these powerful drugs, which are recommended by some authors, especially Heine, appear to us too large to be safely administered.

We should recommend beginning with a dose of at most gtt. ij of the tincture, or gr. $\frac{1}{20}$ th of the alcoholic extract of *nux vomica*, or gr. $\frac{1}{80}$ th of sulphate of *strychnia*, for a child of two years old; the amount being increased steadily but cautiously so long as no unpleasant symptoms are produced by it.

Local means must also be employed for inducing increased circulation in the affected parts. For this purpose, the stimulating liniments already mentioned, or moist heat, may be applied. Passive motion and kneading the muscles, also aid in improving their nutrition and contractile power.

Electricity, however, certainly ranks first among the local means for restoring the contractile power of the paralyzed muscles. It is true that several authorities have asserted that they derived no good results from its employment, but since the introduction of localized electricity (faradization), as developed by the researches of Duchenne, and of the use of the constant current, the most marked benefit has been obtained at all stages of this form of paralysis.

If the induced current be used, it must be carefully isolated and limited to the affected muscles, by means of wet sponges fastened to the electrodes. In those cases where the muscles refuse to respond to an induced current even of considerable power, the direct current, slowly interrupted (the labile current of Remak), will be found to induce contractions, excepting where the muscular tissue is far advanced in fatty degeneration. In all such cases then, this direct current should be employed. We have already alluded to the fact, that as the palsied muscles regain their power under the use of the direct current, they respond to it less and less strongly, while the induced current is found to again have the power of exciting muscular contractions. When this period in the treatment of the case arrives it is desirable to substitute the use of the induced current.

In order that the use of electricity, in either form, may be productive of the excellent results it is capable of yielding, it must be applied thoroughly to each of the paralyzed muscles three or four times weekly, and this treatment pursued for months, until the muscles regain both their size and contractile power.¹ The value of this mode of treatment is, indeed

¹For a full description of the best forms of electrical batteries for medical purposes, the reader is referred to some of the manuals on medical electricity, as Meyer, Tibbits, etc.

The best batteries in the American market are made by Flemming & Talbot, of Philadelphia, or by the American Galvano-faradic Manufacturing Company of New York.

so great "that so long as muscular contraction can be induced, recovery is merely a matter of time, but if no action of the paralyzed muscles can be brought about, the prognosis must be unfavorable, though even here there is some hope." (Hammond, Radcliffe.) One of the earliest symptoms of improving nutrition is an elevation in the temperature of the part, which may readily be detected by the galvanometer, as before mentioned.

In addition, however, to the local and general measures above recommended, there is another kind of treatment scarcely less important, which should be employed in conjunction with them.

This consists in the use of such mechanical apparatus and gymnastic exercises as shall tend to bring the affected muscles into play, and to obviate the deformities of the atrophic period. The greater part of our knowledge upon this subject is due to the admirable and extensive observations of Heine, who had the superintendence of a large orthopædic institute, and most carefully studied the effects of these agents upon cases of paralysis which have progressed to the stage of atrophy and deformity. But it is by no means to this advanced stage alone that such measures are adapted, for it is a matter of the highest importance, that from a very early period of the paralysis, the little patients should be subjected to this treatment.

If the legs be affected, it is not surprising that the child, who has, perhaps, gained but imperfect use of its limbs, and is making its first essays in walking when the paralysis appears, should feel such a sense of insecurity, even when the power of motion has returned to a considerable extent, that it will refuse to make any renewed efforts to walk. And the parents, finding all their attempts to persuade or compel it to do so unavailing and distressing to the child, are apt to desist, waiting until increased power of movement returns; a delay which is too often followed by all the steps of the atrophic period.

To supply the indispensable exercise of the muscles, and in a form attractive to the little patients, numerous mechanical contrivances have been resorted to.

While the legs are still almost powerless, some form of baby-jumper at the same time delights the child and effectually exercises its limbs. When the power of motion has returned to a somewhat greater extent, we gain the same results even more completely by the use of the go-cart or velocipede, a frame or a chair upon wheels, the motive power being furnished by the alternate pressure of the rider's feet upon a pair of treadles which are connected with the wheels by cranks. This imparts such a sense of security and so much pleasure, that the child can readily be encouraged to take enough exercise to preserve the play of the articulations, and to aid in developing muscular power.

Dr. West makes a single objection to the use of the go-cart; that it encourages the tendency to lean very much forward in walking, which always exists until after the little patients have learned to walk pretty well; he, therefore, advises that, after the child has gained some facility in the use of the go-cart, a jacket should be worn, supplied with a stout

strap before and behind, so that the attendant can conveniently hold them and support the child's weight more or less completely, thus enabling it to walk without being thrown forward as when stepping in a go-cart.

In children of from five to seven years even, the use of crutches is soon acquired, and it is desirable, so soon as possible, to abandon the other contrivances spoken of, and trust the child to its own exertions to walk with a pair of crutches.

When the paralysis affects the arms, precisely the same principle should guide us, and every form of persuasion, of stratagem, and contrivance, must be used to induce the child to exercise the crippled member. Trundling a hoop, or raising a weight by means of a cord passing over a pulley, furnish good exercise to the arm; or we may encourage the little one to use a contrivance, also called a velocipede, in which the wheels are turned by handles, instead of treadles, attached to the cranks.

In addition to these forms of exercise, however, it is often found necessary to employ splints of different kinds, such as Stromeyer's, which enables the angle of the splint to be changed without removal from the limb, and various modes of extension to counteract the tendency which exists to contraction of the paralyzed part. In some cases, indeed, all means are powerless to avoid this consequence, and we are obliged to resort to the section of the tendons of the contracted muscles and subsequent extension, though tenotomy should not be performed until time has been allowed to show the extent of permanent paralysis, and until the conjoined use of electricity and orthopædic apparatus has proved insufficient to restore the limb to its shape.

It may readily be surmised that this orthopædic plan of treatment is one requiring the utmost patience and persistence, and the most loving persuasion and encouragement; for, indeed, it must be pursued, in face of all apparent failure, for months and years. Nor must we be satisfied during this period with these efforts we are making to restore the power of the muscles; but careful attention must be paid to the nutrition and general health of the child, and we must continue the use of the warm douche, in conjunction with the persistent use of electricity, of stimulating frictions, and of every remedy calculated to promote the general nutrition of the child.

ARTICLE XII.

FACIAL PARALYSIS.

PARALYSIS of the muscles supplied with motor power by the facial nerve, is frequently met with as a temporary condition in infants who have been delivered by forceps, as a result of the pressure of the blade of the instrument upon the nerve as it emerges from the cranium. It is by no means rare, however, during childhood, and either appears suddenly after exposure to cold, when it is possibly due to pressure caused by

congestion and swelling of the tissues around the stylo-mastoid foramen; or more gradually, when it is usually due to pressure from an enlarged gland, or to disease of the petrous portion of the temporal bone.

The symptoms of this affection are so striking that no difficulty can exist as to its diagnosis. The eye upon the affected side remains open; the power of knitting the forehead and of raising the eyebrow is lost; the angle of the nose and mouth on the same side hang down. The tears trickle over the cheek, and the conjunctiva frequently becomes injected or inflamed; saliva dribbles from the mouth, portions of food collect between the teeth and paralyzed cheek, and there is inability to whistle, spit, or distend the cheeks with air. During the acts of laughing or crying, the face becomes distorted, owing to the immobility of the paralyzed side, while the antagonistic muscles act strongly and draw the features towards the sound side.

In addition to these symptoms, which are common to all cases of facial palsy, there are others which depend upon the point at which the lesion involves the trunk of the facial nerve. Thus if the nerve be paralyzed between its point of emergence from beside the pons and the point where it gives off its petrosal branches (soon after entering the Fallopian canal), there will also be paralysis of one side of the soft palate, greater acuteness of hearing on one side, and loss of the sense of taste on one-half the anterior part of the tongue. This latter symptom is due to the implication of the chorda tympani branch. If, therefore, the seat of the lesion is in the Fallopian canal between the points of origin of the petrosal branch and the chorda tympani, the palate will not be paralyzed, but the sense of taste will be lost.

It is usually true that if the nerve be paralyzed before the origin of the stapedius branch, the hearing becomes more acute; but in children the cause of the palsy is so often necrosis of the petrous portion of the temporal bone associated with disease of the internal ear, that there is frequently deafness with purulent otorrhœa. Finally, if the point of paralysis be near the stylo-mastoid foramen and below the chorda tympani, none of the above symptoms will be present, and there will only be the palsy of the external muscles already indicated.

The possibility of mistaking simple facial paralysis for hemiplegia from cerebral disease must be borne in mind, though attention to the symptoms of the case will prevent any error in diagnosis. Thus in hemiplegia of cerebral origin, the paralysis is usually ushered in by convulsions and coma; the frontalis and orbicularis oculi muscles are not paralyzed; the sense of taste is not affected, but, on the other hand, the masseters, temporals, and pterygoids, supplied by the fifth nerve, occasionally are paralyzed, and the tongue is protruded towards the paralyzed side; and, finally, there is loss of power in the arm and leg on the same side.

Webber (*Chicago Jour. of Nervous and Mental Disease*, July, 1876, p. 363) records several interesting cases where the facial palsy appeared after convulsions, and while the paralysis of the arm and leg were very transient so as to have passed away before the case came under observation, the loss of power of the muscles of the face was persistent, and was associated with

impairment of electro-muscular contractility. In these unusual cases the author thinks the lesion was in the brain, affecting the centre of innervation for the facial muscles, which the researches of Hitzig and Ferrier tend to locate in the lower part of the central ascending convolution.

The prognosis of cases of facial palsy must evidently depend upon the cause. When the paralysis is due simply to exposure to cold, a cure may be expected, though the affection is often very tedious, the paralysis at times persisting for months. But when, on the other hand, it depends upon disease of the temporal bone, the prognosis is usually unfavorable.

The treatment must also be modified according to the cause of the attack.

In simple acute cases, the application of hot fomentations to the part, or of one or two leeches near the stylo-mastoid foramen, should always be directed, and is often productive of good results. Later in the affection, if the paralysis persists, small blisters should be repeatedly applied near the point of exit of the nerve.

Electricity is here also of very great service, and the same curious observation, which was mentioned in atrophic infantile paralysis, as to the power of the direct current to excite muscular contractions when the muscles have ceased entirely to respond to an induced current, has been frequently made in this affection.

In addition to these local remedies, the internal use of strychnia, iron, or iodide of potassium, is often followed by benefit. In cases where there is reason to suspect that disease of the bone, or scrofulous enlargement of the cervical glands, are the cause of the paralysis, the patient should be put upon the use of iodide of iron or cod-liver oil.

ARTICLE XIII.

PROGRESSIVE MUSCULAR SCLEROSIS, OR PSEUDO-HYPERTROPHIC MUSCULAR PARALYSIS.

DEFINITION.—This curious affection is characterized by progressive loss of power, which first appears in certain groups of muscles, and advances until nearly all the muscles of the body may be involved, while at the same time the affected muscles increase in size and firmness owing to excessive hypertrophy (sclerotic) of their inter-fibrillar connective tissue. The muscular fibres usually present changes themselves, and at a later stage there is a process of fatty degeneration or accumulation in the newly formed interstitial tissue.

HISTORY; SYNONYMS AND FREQUENCY.—True progressive muscular atrophy is extremely rare in young children; and among the cases which have been described, as by Meryon, a certain number seem to belong to the disease now under consideration. The merit of having first clearly recognized and described the distinctive features of this latter affection

certainly belongs to Duchenne, whose first observations were published more than twelve years ago. Since then cases have been reported in rapid succession until the number now upon record probably exceeds 150. The disease cannot, therefore, be regarded as a very rare one. We have ourselves had an opportunity of carefully studying seven cases, including the one of which a full account was published in 1871.¹

Various names have already been applied to the affection. It was originally called "hypertrophic paraplegia of infancy," by Duchenne, but he has since substituted the terms, paralysis with muscular sclerosis (*paralysie myosclérosique*), or muscular paralysis with apparent hypertrophy (*paralysie musculaire pseudo-hypertrophique*). It has also been called "lipomatosis luxurians musculorum progressiva" (Heller); lipomatous muscular atrophy (Seidel); progressive muscular paralysis, as a result of hypertrophy of the interstitial fatty tissue (Niemeyer); fatty muscular hypertrophy (Bergeron and Lutz); pseudo-hypertrophic spinal paralysis (Hammond); and, finally, progressive muscular sclerosis (Jaccoud and others). We much prefer this latter term, since it expresses the true pathological process which is present, and at the same time does not tend to confound this disease with any of the forms of true paralysis, from which it is, in reality, clearly distinguished by the facts that its essential feature is a progressive change in the structure of the muscle, and that the loss of power is dependent upon the change in the muscular tissue, and is not primary, as in all true palsies.

CAUSES.—The essential causes of progressive muscular sclerosis are unknown. There are, however, some influences which exert marked control over its occurrence. One of the most important of these is *early age*, since in a very large majority of cases the disease begins in childhood, and has even appeared in some cases to be congenital (Niemeyer). Although, however, it must be distinctly classed among the affections of childhood, it has been shown (Benedikt, Lutz, and Laycock) to occasionally occur in adult life. *Sex* also exerts a powerful influence: of 45 cases collected by Estrazulas (*loc. cit.*), in which this point was noted, it occurred only 7 times in females.

The curious fact has also been observed, that several children in the same family are apt to be affected, probably indicating some hereditary tendency. Eulenburg² is consequently inclined to regard the disease as dependent upon some congenitally defective formation of the central nervous system, probably in the cells of the gray substance of the spinal cord. Instances are on record where four brothers were affected (Meryon); and in another two brothers (Eulenburg); and in still another by the son of the latter author, in which the affection first showed itself in three sisters successively in the eighth year of their age.

SYMPTOMS.—The disease either begins in early infancy, and is first manifested at the time the child should begin to walk, or it makes its appearance some years after the power of walking has been acquired.

¹ Clinical Lecture on a case of Progressive Muscular Sclerosis, by Prof. William Pepper, M. D., Philadelphia Med. Times, June 15th and July 1st, 1871.

² Virchow's Archiv, liii, 361.

The disease usually affects first the muscles of the legs, and advances upwards; in Niemeyer's case, on the other hand, it began in the gluteal muscles, and subsequently affected all the muscles of the lower extremities. The early symptoms are, therefore, connected with walking, and it is observed either that the child does not begin to walk until very late, and then walks imperfectly, or that, having walked well for several years, he begins to be readily tired by standing or walking, and soon presents peculiarities in his gait. In a few instances, pains in the limbs have been complained of in the early stage. When the disease is fully established, though before it has advanced far, the mode of walking and standing are quite characteristic. The patients find that, without some support, these operations become more and more difficult and painful, and that they are subject to frequent falls. In order to maintain their equilibrium while standing or walking, the lower dorsal and lumbar spine is arched forwards, while the upper part of the spine, the shoulders and head are bent backwards, frequently to so great an extent that their point of equilibrium falls behind the pelvis, thus producing the deformity known as "ensellure" or "saddle-back." The legs are widely separated, and in walking the body is inclined laterally towards the leg which rests on the ground, thus producing a characteristic balancing of the body during progression, while the arms are swung about, and the legs are advanced by jerks, describing a small arc.

While this impairment of strength and power of progression is developing, the affected muscles undergo remarkable changes. For a time they may be noticed merely to cease developing and increasing in size, or, more rarely, as in the case reported by one of ourselves (*loc. cit.*), they may present a well marked stage of atrophy. After the stage of muscular weakness has lasted for a variable time, from a few months to two or even three years, whether or not there has been any noticeable atrophy of the affected muscles, a progressive enlargement of them makes its appearance. This usually affects the gastrocnemii first, then the glutei, the lumbar muscles of the spine, of the trunk, and finally the muscles of the arms, and even of the face and tongue. In five of the recorded cases, the heart has been hypertrophied. In one of these, reported by Dr. B. W. Foster, quoted by Poore (*loc. cit.*), the heart was normal when first examined, but three years later, and without apparent cause, it was found to be enlarged.

The above order is not invariably followed, and in by no means every case is the affection of the muscles so universal. The apparent hypertrophy may occur in nearly all the muscles which have shown weakness, but in general, according to Duchenne, it does not, and may even be limited to a very small number of them. The same observer (*loc. cit.*) thus describes the appearance of the muscles after this consecutive enlargement has occurred.

. . . "The hypertrophied muscles are firm and elastic; they become very hard while they contract, and show all the relief or projection which properly belongs to their contracted state; they then appear to form a hernial protrusion through the integument, which is very thin; moreover, their

great size shows off the apparent smallness and delicacy of the joints at the knee, ankle, etc."

When this pseudo-hypertrophy is marked, and affects many muscles, it gives a most curious appearance to the children. Niemeyer speaks of his patient as looking "as if he had the body and head of a weak child on the hips and thighs of a strong man;" and J. Lockhart Clarke, in describing one of Duchenne's patients, says; "He looked like a little Hercules. Every visible muscle of the body, except the pectorals, was enormously developed; his head, even, appeared swollen, and the temporal muscles stood out like convex shells. Yet, when the poor boy attempted to walk, he labored to get along, presenting the most grotesque appearance; and when laid on the ground, he was wholly unable to rise by his own unaided efforts."¹

Dr. Mitchell (*loc. cit.*) calls attention to the fact, however, which we have also observed, that the enlargement of the calves is lower down than would be the case in excessively developed, but well formed limbs.

The marked enlargement of the muscles of the calves is often attended with forced extension of the feet, producing double pes-equinus or equinovarus. In the case reported by Estrazulas (*loc. cit.*), there was also marked enlargement and retraction of the posterior muscles of the thigh, with atrophy of the extensor group, so that there was forced flexion of the legs, rendering the boy unable to stand at all. Knoll² also describes such contractions in the enlarged muscles, but they are not usually present. According to Berger,³ fibrillar contractions are of constant occurrence in the affected muscles; this does not accord with our own observations, nor with many of the reported descriptions of the disease. In one of the cases reported by Gerhard (*loc. cit.*) there was constant tremor of the flexors of the legs and feet, and of some of the muscles of the forearms.

The electrical condition of the affected muscles is peculiar. Frequently the results, when tested with faradic currents, are different from those obtained with galvanism. The results also vary at different stages of the same case. Usually the muscular contractility, as tested by faradization, is impaired in all the affected muscles, those which are hypertrophied, however, contracting more actively than those which are atrophied. The galvano-contractility is also slightly impaired. The electro-muscular contractility has been found unimpaired in the earlier stages of the disease; but later it diminishes, the muscles continuing, however, to respond actively to galvanism after they have partly lost their power of responding to faradization.

Electro-muscular sensibility has been found normal or impaired in different cases; in one of our patients it was diminished to faradization, but remained acute to galvanism.

The skin over the affected parts often presents a marbled or mottled appearance. In one case that we have seen (described by Mitchell, *loc. cit.*), the mottling "consists of spaces of pallid skin surrounded by quite regular

¹ Trans. of London Path. Soc., vol. xix, 1868, p. 6.

² Wien. Med. Jahrb., 1872; and in Syd. Soc. Bienn. Retrospect, 1871-72, p. 71.

³ Deut. Arch. f. Klin. Med., March, 1872, Bd. ix; Hft. 4, 5, p. 363.

circles of congestion, which affect an irregular polygonal shape." The skin is usually thin and delicate, and can be easily lifted from the muscles. Disorders of the cutaneous sensibility have not been usually found, but Berger (*loc. cit.*) describes violent neuralgic pains and formication, followed at a later stage by anæsthesia. We have already alluded to the pains in the limbs occasionally complained of in the early stages of the disease.

The temperature of the parts is lowered. This can be distinguished by the hand, and has been found, on careful thermometric study, by Mitchell,¹ to be as follows: Left axilla, 97.5° ; right axilla, 97° ; perineum, 94.5° ; right calf, 91.5° ; left calf, 91° ; and Estrazulas, in the case observed by him, reports the temperature in both axillæ 98° ; on right calf, $91\frac{3}{8}^{\circ}$; and on left calf, 91° .

There is usually an entire want of disturbance of the general health.

The appetite remains good until a late period, digestion is well performed, and the action of the bowels is regular. Neither the rectum nor the urinary bladder become paralyzed. There is frequently an entire want of cerebral symptoms, and the mind may be clear until the close of the case. In several instances, however, the patients have been of feeble intelligence, or even idiotic; and in the case above reported, it will be remembered that the disease was complicated with epileptiform convulsions.

COURSE AND DURATION.—As will be inferred from the foregoing description, the duration of this disease is very considerable, varying from five to fifteen years, or even more. It may occupy several years in reaching its full development, and may then remain at this stage for several years, or even until a tolerably advanced period of youth, but finally it is succeeded by a stage in which the loss of power becomes more complete and extensive, involving the upper extremities and muscles of respiration, and confining the patients to the recumbent position. During this final stage there is a rapid decrease in the size of the hypertrophied muscles, and the limbs may even come to present an appearance of great atrophy.

Death usually occurs before adult age from sheer prostration or from some intercurrent affection of the respiratory organs.

PROGNOSIS.—The course of this disease is steadily progressive, and, despite the various plans of treatment adopted, usually leads to a fatal result. In one case, however, recovery took place, and in one other there was some improvement. In the case we have here reported, there seemed to be some temporary improvement under treatment.

DIAGNOSIS.—The diseases from which it is most important to distinguish progressive muscular sclerosis are atrophic infantile paralysis and progressive muscular atrophy. In infantile paralysis, however, the suddenness of attack, frequently associated with fever or with some cerebral disturbance, as convulsions; the occurrence of complete and more or less extensive paralysis; the gradual disappearance of the paralysis in some parts, while in others it remains permanent; the diminution and ultimate loss of electro-

¹ This case was re-examined by Gerhard (*loc. cit.*, p. 31) at a later period of its development with the following results: Right deltoid, 92° ; left deltoid, $92\frac{1}{2}^{\circ}$; right thigh (inner side), $94\frac{1}{8}^{\circ}$; left thigh (inner side), $94\frac{1}{8}^{\circ}$; right calf, 90° ; left calf, $90\frac{3}{8}^{\circ}$.

muscular contractility; the occurrence at a later period of fatty degeneration and atrophy of the affected muscles, with arrest in the development of the bones and marked deformities; and the entire absence of any secondary enlargement of the part involved, constitute a series of distinctive features so clear and decisive as to render the differential diagnosis easy and certain.

A disease from which it is much more important to carefully distinguish progressive sclerosis of the muscles is progressive muscular atrophy occurring in childhood. The especial importance of the relations of these two diseases depends on the fact that both are alike diseases of nutrition of the muscles, thus constituting a group quite distinct from all the forms of true paralysis. In both the disease begins—usually without any apparent cause—insidiously, and progresses slowly but surely. In both the loss of motor power is secondary to the changes in the muscular tissue; in both the muscular degeneration and consequent loss of power almost invariably progress steadily to a fatal result. These two diseases, then, stand related to each as being alike caused by disturbance of the trophic nervous system, but they are at the same time most positively separated from each other by marked differences in their course and symptoms.

Thus, in progressive muscular atrophy, the disease nearly always begins in the upper extremities, and invades subsequently the trunk and lower extremities. Indeed, Duchenne has pointed out that when this disease appears in childhood, which is quite rare, it usually begins in the face, where it produces atrophy of the orbicularis oris and the zygomatici, and does not extend to the trunk and extremities until after a period varying from two to three years. It then follows the same descending course seen in cases occurring in adults. The atrophy usually affects the muscles irregularly, so that various deformities and vicious positions of the parts involved are developed. Microscopic examination shows a progressive fatty degeneration and atrophy of the muscular fibrils, and in proportion as this increases there is loss of power and of electro-muscular contractility. One further symptom of high diagnostic value is the frequent occurrence of fibrillar contractions in the affected muscles, which, although stated by Berger to be of constant occurrence in progressive muscular sclerosis, has not been found so by ourselves or other observers. Finally, the muscles which have progressively atrophied never undergo any secondary enlargement, nor does microscopic examination reveal any lesion of the interfibrillar connective tissue. In all these particulars, then, progressive muscular atrophy differs widely from progressive muscular sclerosis, which is almost exclusively a disease of childhood, beginning in the muscles of the lower extremities and advancing upwards, producing a peculiar mode of standing and walking, and in which the affected muscles, with or without a previous stage of atrophy, undergo remarkable enlargement, usually without fibrillar contractions, and with preservation of electro-muscular contractility till a comparatively late period of the disease. The results of microscopic examination, also, as detailed in the next paragraph on the morbid anatomy, are entirely different from those observed in progressive atrophy.

MORBID ANATOMY AND NATURE.—There is still an urgent need of careful, skilfully conducted microscopic examinations of the nerve centres in this disease. The examinations which have been made up to the present have not yielded uniform results. In two of them, Cohnheim's (where, however, the microscopic study was not conducted with the requisite care and thoroughness) and Charcot's,¹ no lesions were found in the spinal cord; while in the cases reported by Müller,² Barth,³ and Lockhart Clarke,⁴ positive lesions of the cord were discovered, chiefly affecting the anterior columns of gray matter and the large nerve-cells which exist there. Eulenburg had already suggested that the pathological origin of this affection would be found in some defective formation or disease of these parts; and in the first publication made on this subject by one of ourselves, when Cohnheim's imperfectly studied case was the only one on record, we stated that analogy with other diseases of the nutrition of the muscles supported this suggestion. Hammond (*op. cit.*, p. 500) "feels warranted in at least provisionally accepting the view, that the anterior tract of gray matter is the seat of lesion in pseudo-hypertrophic paralysis." Charcot, on the other hand, contends that the anatomical cause of this affection is not seated in the spinal cord. In this view, he agrees with many other authorities. Duchenne ascribes it to a paralysis of the vaso-motor nerves; and Berger (*loc. cit.*), who assumes the existence of trophic nerves, attributes the disease to some disturbance of their function. In a case recently reported by Brigidi (*Lo Sperimentale*, March, 1878; *N. Y. Med. Record*, May 25th, 1878) the sympathetic ganglia presented marked alterations: the nerve-cells were atrophied and pigmented, the connective tissues hyperplastic, and the nerve-fibres altered. Others again regard it as a primitive muscular lesion; as Gowers, who asserts that it is not a disease of the spinal cord, but attributes it to "a congenital nutritive and formative weakness of the striated muscle substance." It is evident, therefore, that further careful examinations of the spinal cord in this disease are necessary before a definite conclusion can be reached on this point of vital importance.

The condition of the affected muscles themselves has been very carefully studied during life, on small fragments removed by Duchenne's trocar⁵ (*emporte-pièce*), and the results confirmed by examination after death.

¹ Sur l'état anatomique des muscles et de la moelle épinière dans un cas de paralysie pseudo-hypertrophique, *Arch. de Phys.*, March, 1872, p. 228.

² Beiträge z. Path. Anat. u. Phys. d. menschlichen Rückenmarks, Hft. ii, Leipzig, 1870.

³ Beiträge z. Kenntniss d. atrophie musculorum lipomatosa; *Arch. d. Heilkund.*, Leipzig, 1871, p. 120.

⁴ *Medico-Chir. Trans.*, vol. xlvii, 1875, p. 247.

⁵ This useful little instrument is shaped like a trocar. The blade is, however, a hollow cylinder, composed of two parts, one of which, bearing the point, is fixed, while the other can be withdrawn a little by sliding a movable button on the handle. The trocar is introduced close into the substance of the muscle, the button withdrawn, so as to open the cylinder and allow a fragment of muscle to project into it; the button is then pushed forward, cutting off and securing the little morsel of tissue.

When examined by the naked eye, their color is altered, and the muscles present either a uniform pale or yellowish appearance, or are marked with stripes of yellow or yellowish-white; on section they shine with a dull, greasy lustre.

The results of microscopic examinations vary somewhat at different periods of the disease. The changes affect both the muscular fibrils, and even more markedly, the inter-fibrillar connective tissue.

In the early stage, Berger asserts (*loc. cit.*) that he found in two cases an absence of change in the interstitial tissue, and a marked hypertrophy of the fibrils themselves. This enlargement has not, however, been constantly observed. The fact of its occurrence, and of its persistence in some fibrils even in a comparatively advanced period of the disease, is confirmed by the observation of Leyden,¹ Estrazulas (*loc. cit.*), Knoll (*loc. cit.*), and ourselves. In the later stages, many of the fibrils are pale and small, being occasionally reduced, according to Cohnheim, to $\frac{1}{3}$ th their normal diameter; in some places empty sheaths of sarcolemma are seen. Many of these fibrils, though altered in size, present no other morbid condition, either fatty or granular. Knoll observed in some of the border fibres a tendency to split into two; and Martini² describes a peculiar process of fission or division of some of the atrophied fibres. In the cases recorded by Meryon (*loc. cit.*), which were probably of this form of disease, a granular degeneration of the muscular fibres with rupture of the sarcolemma was observed.

The most marked change is, however, in the condition of the interstitial tissue. It is not known definitely whether this precedes all change in the muscular fibrils themselves. But at least by the time that enlargement of the muscular masses can readily be detected, there is usually, despite the two observations of Berger, marked proliferation of its nuclei and hyperplasia of the fibrils. This continues to increase until at places the muscular fibres are separated by broad tracts of wavy fibrous tissue, interspersed with fine nuclei. At a later period this is associated with increasing interstitial fatty accumulation and degeneration, which advances with varying rapidity, even leading in some cases to such extreme accumulation of fat as to be visible to the unaided eye as yellowish streaks. It is probable that the muscular fibres may temporarily share the exaggerated nutrition of the surrounding connective tissue, but later, as this interstitial tissue accumulates, the fibrils are subjected to severe pressure, and undergo atrophy in many instances. The entire process, therefore, seems to be of a strictly sclerotic character, so far as the inter-fibrillar connective tissue is concerned, but associated with an irregular and as yet undetermined stage of true hypertrophy of the muscular fibrils themselves.

TREATMENT.—The results of treatment in progressive muscular sclerosis have so far been highly unsatisfactory. The internal remedies from which most benefit may be expected are those which tend to improve nutrition, and especially to improve the tone of nutrition of the nerve-centres. Among these, cod-liver oil, iron, the compound syrup of the phosphates, and arsenic, may be specially mentioned. In the case reported by our-

¹ Berl. Klin. Wochenschr., 1866.

² Centralblatt, 1871, 641; in Syd. Soc. Bienn. Retrospect, 1871-72, p. 70.

selves, where there was the complication of epileptiform convulsions, benefit was derived from a course of bromide of potassium.

The remedy, however, from which most good is to be expected, is electricity. This has been used by Duchenne with great benefit, in the form of faradization of the affected muscles. It is asserted by Benedikt, that good results have been attained in three cases by the use of the direct current, the copper pole being placed over the lower cervical ganglion, and the zinc pole along the side of the lumbar vertebræ, by means of a broad metal plate. Others have, however, tried this mode of treatment for a long time without any success. As, however, no more plausible mode of treatment has yet been suggested, we should be inclined to adopt it in conjunction with direct faradization of the affected muscles, and the use of the internal remedies above recommended.

ARTICLE XIV.

NIGHT TERRORS.

THE night terror of children is a condition, in some respects, analogous to the nightmare of the adult. It is not quite the same, however, for in nightmare the subject is relieved of the symptoms so soon as he awakes from the sleep in which the dream has occurred, while in night terror the symptoms continue for some time after the patient has been roused sufficiently to utter cries, and to exhibit in the expression of the countenance, and in the movements, an accordance with the painful idea which occupies the mind. It is, perhaps, more akin to somnambulism than nightmare. But we will cite some of the cases we have met with, and then endeavor to explain the pathology of the condition, and to describe its proper treatment.

Several years since, we had the charge of a family of five children, all of whom were more than usually intelligent, and all were liable in early childhood to frequent attacks of night terror. One of these children died early of serous effusion into the ventricles of the brain, occurring after a severe attack of diarrhœa. The eldest child, a son, who became quite a distinguished student and lawyer, was specially liable to this condition. He would wake partially from sound sleep, screaming, struggling, and exhibiting all the signs of a violent terror. This condition lasted for several minutes after the screams and struggles had begun, during which time it was evident that he was under the influence of some terrifying idea. The only thing to be done at the time was to hold him gently in the arms, and endeavor to rouse him into full wakefulness by soothing words and caresses. During his early years he was unable to explain the character of the idea which caused the distress. At a later period he recollected that the terrifying thought was always one connected with some object of vast size. Many of the attacks, he said, arose from his seeing in his sleep an elephant

in the nursery, which, from being small in size at first, expanded before his eyes to such a size that he was being crushed between it and the walls.

In another case, we were called to see a young child, one of a large family, of nervous and rather timid type, which, whilst playing on the floor of the nursery one morning, was bitten in the foot by a tame parrot as it hopped about the room. The child, though not seriously bitten, was very much alarmed at the time. On the following night, at midnight, it started suddenly from a sound sleep, shrieking, "Take the parrot away, take the parrot away!" and was so alarmed and terrified, and struggled so violently and so long with the idea, that the mother, a very sensible and experienced person, became quite alarmed lest it should have a convulsion, and was on the point of sending for us. At last, by carrying it about and soothing it gently, it was fully waked and the terror passed away for the time. This scene was repeated for several nights afterwards at about the same hour, but with diminishing violence, until the impression faded away and disappeared. At the time of the occurrence the child was perfectly well, and when we visited it on the following day and for several days afterwards, there was no disturbance of the health requiring medical interference. The attacks were evidently the result of the vivid impression made upon the child in the daytime, which, by the play of memory during sleep, reproduced to the child all the pain and terror endured at the moment of the occurrence.

We were sent for on another occasion to see a little girl of six years of age, who had alarmed the mother greatly during the previous night by partially waking from sound sleep, screaming, "Take the white dog away, oh, take the white dog away!" with such terror in the countenance and eye, such agitation and struggling of the limbs, that the mother feared it must have a fit. It was many moments before it could be roused sufficiently from the delirious terror to recognize those around it, and to know that there was no dog by its side. The terror was repeated for several nights at about the same time, and then gradually passed away. In this case the child had been walking in the street with the mother, on the day before the first dream occurred, and had been suddenly shocked and terrified by a large white dog brushing sharply against her, as it was careering along the street. There was no disturbance of the health at the time nor afterwards. As in the last case, the attack was evidently the result of the reproduction in a dream of the fright it had had in the street.

The following case was still more curious. It is related very much in the language of the mother. The child, a girl two years and two months old, was the daughter of a very bright and highly educated mother, and of a father of unusual intelligence and force of character. The child was herself very precocious and active-minded. The parents were both very healthy persons. On the night of September 13th, 1873, the child awoke partially in a great fright, screaming, trembling, and covered with cold perspiration. Nothing calmed her for some time. She repeated over and over again, "Teeth bite you, teeth bite you!" Finally, at the end of an hour, she was roused from the state of half sleep, the sobs and crying ceased, and the attack ended, but at dawn of the same night there was

another seizure which was not quite so violent. During the attack she begged to have her head covered, evidently fancying that this protected her from some threatening object. During the following day she manifested great fear of the large slop jar in the nursery, and could not be induced to approach it. When her father took it into his hands to explain to her that it was "only a jar," she screamed most pitifully and seemed in abject terror. Being a child of peculiarly vivid imagination, it was supposed that she had, by some freak of fancy, come to imagine the top or opening of the jar, to be an open mouth armed with teeth, ready to rend and tear her. She conceived, also, a great terror of the carved ornaments on the head-board of her crib, and, indeed, these fears became so marked and so great, that she was moved into another room, where the furniture was plainer and unvarnished, so that the shadows and reflections might not feed her disturbed fancy. In December she was seized with typhoid fever, from which she did not recover until February, 1874. In that month her mother had another child, a boy. When she was taken into the chamber to see her little brother, she had another nervous attack, in which she fancied that an animal was biting her mother and that it might bite her also. The nervous seizures at night continued to recur at intervals of about five weeks, generally after violent exercise or hearty eating, until May, 1875, when a severe attack of scarlet fever seemed to eradicate them in great measure. At the present time, she has a night terror only when her imagination has been, in some way, excited, or her sympathies overtaxed.

With one more case, which shows the curious features of this sleep intoxication in a very marked manner, we shall end the citation of cases, and pass on to a consideration of the pathology and treatment of the condition.

A gentleman, whose eldest son was rather remarkable for his intellectual development, was roused from a sound sleep in the early morning by a touch on the shoulder. On waking he saw this boy, then about twelve years old, standing by the bedside, dressed in his wrapper, his legs bare, and with his bare feet in slippers. The boy said: "Father, there are robbers in the house, they are downstairs now, but I do not think there is any danger, for I have been downstairs and have locked the doors, but I thought I had best tell you." The father thus waked suddenly from sleep, thought at first that the child was insane, but seeing some peculiar fixity in his look and manner, it flashed into his mind that the child was acting under the influence of a dream. He touched him sharply, and said loudly, "My son, you are sleeping; wake up, wake up." The boy drew a long, deep breath, gaped, waked, and said, "Why so I am," and walked off to bed again in the most natural manner.

PATHOLOGY.—The best explanation of the conditions which exist in this curious disorder of sleep is, we think, to be found in the works of writers on forensic medicine. Thus, Dr. Johann L. Caspar (*Forensic Medicine, Syd. Soc. Ed.*, vol. iv., p. 273) remarks that "the dreaming state passes quite insensibly into that of somnolence, that middle state betwixt sleeping and waking, in which the connection with the outer world is neither that

of sleep nor waking. The dreaming state is wholly sleep; somnolence is half sleep, half waking. In it the senses are neither quite awake nor quite roused, but are surrounded by a cloud of dream phantasms; the somnolent man sees and hears self-made phantoms instead of real objects; he hears a shot fired, and dreams of it, while it was only a stool that fell. He reasons logically, as is well known to be the case also in dreams, in regard to the impressions supposed to be felt, and may, since muscular action is not prevented by sleep, act in the most illegal manner."

In the *Treatise on Medical Jurisprudence* (Philadelphia, 1855, p. 119), by Wharton and Moreton Stillé, in the article on "Mental Unsoundness as Connected with Sleep," it is stated that, "under this general head may be grouped somnolentia or sleep drunkenness, somnambulism, and nightmare, the two last of which may be joined." Sleep is interrupted, they suppose, by whatever terminates the peculiar condition of the brain upon which sleep depends, by the natural exhaustion of the state of the brain, by vivid and sudden impressions on the senses, and by disagreeable sensations. "Now, in a certain morbid condition of the brain this awakening is not complete, and does not restore the waking state with a full and correct perception of surrounding things, but an intermediate state between sleeping and waking is produced, which resembles intoxication, and is called the *intoxication of sleep* (schlaftrunkenheit). This state admits of action, which is directed by the phantoms of the dream; talking in sleep being very nearly allied to waking, and dreams themselves being midway between sleeping and waking, for in the depths of sleep we no longer become conscious of dreams." In this explanation they differ somewhat from Caspar, who asserts that dreams are "purely phantasmagoric conceptions arising spontaneously in the brain, which continues to act during sleep and during the so-called dreamy waking, without any stimulation produced by the external world through the senses." Wharton and Stillé say further: "It is important to distinguish somnolentia, or sleep-drunkenness, which is a state which in a greater or less extent is incidental to every individual, from somnambulism, which is an abnormal condition incident to very few." The experience of every-day life demonstrates how much the former enters into almost every relation. Children, particularly, sometimes struggle convulsively in the effort to wake up, which often is continued for several minutes. The very exclamations, "wake up," "come to," which are so common in addressing persons in the waking condition, are scarcely necessary in bringing to the mind many recollections of cases where the waking struggles were peculiarly protracted. Of course there are constitutions where this struggle is peculiarly distressing, just as there are constitutions in which the tendency to sleeplessness is equally marked. If we recall to the reader the fact, that it is in the state of somnolentia or sleep-drunkenness, that acts of violence have been committed by persons, as stabbing a friend, shooting a passer-by who sought to wake a sleeping sentinel, and other acts of this kind, for which the unfortunate individual has been tried for his life, all of which unhappy events have been committed in the mental state induced by some dream, which has pursued the patient into the only partial awakening, it will not be difficult to understand the

phenomena we have described as occurring in some of the cases of night terrors above cited.

CAUSES.—In some children it evidently needs but a vivid impression upon the mind in the waking state, to produce in the course of the following night, and sometimes for many nights afterwards, the dream which is to cause all the phenomena of the severest night terror. The child may be in perfect health, and yet the mind shall, in sleep, so act as to reproduce in full or in exaggerated force, the terrors which have been first felt in the waking state, and perhaps whilst the child was in full, happy play. Such were the cases in which the child had been bitten by a parrot whilst playing on the floor, and that in which it had been shocked and terrified, whilst walking the streets, by the large dog brushing against its person. In other cases, children predisposed to this condition by some unusual activity of the brain, have the attacks whenever their health is deranged in any way, as by indigestion, or by febrile disturbances from any cause. We have met with the attacks often in the various greater or lesser perturbations of health which accompany the different diseases of childhood.

TREATMENT.—The only treatment necessary during the attack is for the mother or nurse to take the child into her arms, and endeavor by gentle and soothing means to wake it fully from its half sleep. Gentle movements, caresses, soft words, stroking the head and limbs, indeed, the very conduct which any tender mother would naturally adopt towards a terrified and frightened child, are the proper means to be used in the paroxysm.

If, at the time the child is having these attacks, there is any fault in the health, this should be attended to. The digestive system, especially, ought to be carefully examined; constipation should be relieved; the diet ought to be arranged with great care, so that it may be readily digested, and yet be abundantly nutritive. If the patient is pale, iron ought always to be given for some weeks. If there be any trace of periodic disturbance from latent or open malarial disorders, quinia is of the utmost use and importance. When the attacks recur night after night, we know nothing so useful as the bromide of potassium or sodium, of which from two and one-half to five grains may be given at bedtime for one or more weeks. It is often wise, particularly when the disturbance is of obstinate continuance, to add from one to two or three minims, according to the age, of deodorized laudanum, to the dose of bromide. We have obtained decided advantage in several obstinate cases from the use, each evening at bedtime, of a suppository containing two or three grains each of quinia and assafoetida. The avoidance of all causes of nervous excitement, the cessation of study, and even a change of residence may be required to break up the morbid habit in cases where such attacks recur frequently in children of a very sensitive nervous organization.

CLASS V.

GENERAL DISEASES.

INTRODUCTORY REMARKS.

THIS great class includes a large number of diseases, both acute and chronic, in which the system at large, including the blood, is affected by the morbid process. These diseases are so numerous and there are such marked points of difference between some of them, that they have been subdivided into groups in different ways by various authors. The most striking distinction is based on the mode of their causation. Many of them depend upon the introduction from without into the system of specific poisonous principles which excite directly the peculiar symptoms of the disease. As illustrations may be mentioned small-pox and measles. Of late years the names infectious and zymotic are often applied to this group, which includes diseases unattended with eruptions, as mumps and malaria, as well as the eruptive fevers. On the other hand, a group may be formed where no such specific exciting cause can be shown to exist, but where the disease depends upon a derangement of the ordinary processes of nutrition, either from inherited taints of constitution or from the operation of ordinary morbid agencies. As illustrations may be mentioned rheumatism and congenital syphilis.

It will not be necessary to treat of all the diseases that are included in these two great groups, since some of them, as typhus and relapsing fevers, do not present enough peculiarities as occurring in children to justify a special discussion in this work; while others, as gout, are so rare in childhood as to render it undesirable to include them here. We shall therefore divide general diseases as follows:

Those resulting from derangements of the normal processes of nutrition, including:

Rheumatism,
Scrofula,

Tuberculosis,
Rickets,
Congenital Syphilis.

Those resulting from special morbid agents operating from without, including:

Typhoid Fever,
Variola and Varioloid,
Vaccinia,
Varicella,
Scarlatina,
Rubeola,

Rötheln,
Malaria,
Mumps,
Erysipelas,
Diphtheria,
Epidemic Cerebro-spinal Meningitis.

GENERAL DISEASES RESULTING FROM DERANGEMENTS
OF THE NORMAL PROCESSES OF NUTRITION.

ARTICLE I.

ACUTE RHEUMATISM.

As it is not designed to enter into a full discussion of the numerous affections which merely occur in childhood in common with the other periods of life, we shall present but a brief account of rheumatism in children, alluding particularly to those points in which it differs from the same disease in adults.

The importance of this subject is, we believe, not usually appreciated; and it is not treated of at all in many of the treatises on diseases of children. Rheumatism in children deserves careful consideration, however, not only on account of its frequency and peculiarities, but also on account of its marked tendency to cardiac complications, and of its recently established relation to chorea.

SYMPTOMS.—Acute rheumatism may express itself in the child, as in the adult, by painful inflammation of one or more of the larger joints, usually accompanied by a high grade of febrile action. It is probable, however, that in the majority of cases in children, the fever is not so intense nor the course of the disease so long, as in adults.

The fever, which is one of the most marked symptoms, may precede the development of inflammation of the joints by one or two days, or may coincide with the appearance of pain and swelling. It is generally marked in severe cases, and attended by frequency of the pulse, great heat of the skin, and, usually, copious acid perspirations. The heat of the skin and frequency of the pulse constitute a good index of the severity of the disease, and we may always apprehend a dangerous attack when the temperature rises above 104° .

With this febrile action we find disturbance of the digestive functions; the tongue is heavily coated, the appetite lost, or nausea may be present, and the bowels are sluggish, the evacuations being dark and offensive.

The local phenomena attending this fever depend upon acute inflammation of some of the large joints.

Occasionally the ankles, knee-joints, wrists, elbows, and shoulder-joints will be simultaneously affected; but in far the majority of cases, a few only of these articulations will be involved, and the others become affected subsequently, if indeed they do not escape entirely. But one of the most characteristic features of this specific rheumatic inflammation, though most marked in the chronic form, is its tendency to shift its seat, and we may find the intense pain and heat of one part transferred within twenty-four hours to a distant joint. We can rarely learn from the little patients the character of the pain which causes such bitter complaints; in one mild case, recorded by Rilliet and Barthez, it was compared to frequent light blows given upon the affected joints.

The *heat* of the inflamed part is always much increased, and it is not unusual to find its temperature ranging from 100° to 105° (Aitken).

The *swelling* is generally considerable, so that the shape of the parts may be much changed. When the knee-joint is inflamed, the effusion may raise the patella from its position on the condyles.

The skin over the inflamed joints usually presents a more or less decided blush.

While the articular form of acute rheumatism above described is not rarely met with in children, our own experience shows that it is even more common for it to assume the form of acute continued fever, with more or less severe general soreness, a slight development or even a complete absence of joint affections, and a very marked tendency to inflammation of the cardiac serous membranes. It requires care and thorough familiarity with the peculiarities of infantile rheumatism to avoid overlooking the true nature of such cases. The soreness may be extreme, so as to cause cries on every motion; or it may be moderate and localized, so that it escapes detection unless carefully inquired after. We have seen cases where severe and apparently causeless fever existed, with decided complaints of vaguely localized pain about the epigastrium, and with complaints of indistinct soreness about the limbs or back; but where examination showed fully developed endocarditis, and the subsequent progress of the case demonstrated the rheumatic nature of the entire morbid process. These observations have impressed on us most forcibly the necessity of examining every case of acute febrile disease in young children with special reference to the possibility of its being one of acute rheumatism without apparent articular inflammation.

Occasionally when the joints are not markedly implicated, but the muscular or tendinous tissues are more specially attacked, the case assumes a subacute character, is attended with a lower grade of fever, and runs a more protracted and irregular course. It is essential to note that in these cases, contrary to what we find the rule in the analogous form of rheumatism in adults, there is a strong tendency to cardiac complications. Meynet and Heischsprung have called special attention to this subacute fibrous rheumatism in children, and have added the important contribution to our practical knowledge of it that there is a peculiar tendency to relapses. They confirm the statement made above as to the danger of cardiac inflammation.

DURATION.—The duration of acute rheumatism varies exceedingly. According to Rilliet and Barthez, it follows a much more rapid course in children than in adults, occasionally yielding at the end of six days, and nearly always before the fifteenth day. We have, however, seen the rheumatic fever last twenty-one days, and before convalescence was fully entered upon, six weeks had elapsed.

There is a marked tendency to relapses and second attacks in rheumatism, at whatever age it occurs; and we frequently meet with children of twelve or fifteen years of age who have passed through three or four acute attacks of this disease.

CAUSES.—*Age.*—Early infancy appears to protect, to a certain extent,

against this affection, but we are confident that, in some one of its irregular forms, it occurs at that tender age more frequently than is usually recognized. Rilliet and Barthez allude to a case occurring at the age of seven months; and Jacobi (*Amer. Clin. Lect.*, vol. i, no. ii, p. 35) refers to a case at the age of nine weeks, reported by Staeger.

We have ourselves observed one case of acute articular rheumatism in the second year, and several others between the close of the second and fifth years. Of its imperfectly developed forms, we have met with a number of cases in very young children, and are satisfied that the nature of these attacks is often overlooked.

The influence which *sex* exercises upon the frequency of rheumatism in childhood seems still undetermined. It is usually stated that boys are far more liable to the disease than girls, but in our own experience it has been more frequent in girls; and from the register of the Children's Hospital in London (quoted by Tuckwell, *St. Barth. Hosp. Rep.*, vol. v, 1868, p. 102), it appears that of 478 cases of rheumatism treated during sixteen years, 252 were in females, and only 226 in males. The marked difference in this respect between rheumatism in children and in adults may probably be explained by the fact that the two sexes are exposed to the exciting causes much more equally in childhood than in later life.

Cold and Dampness.—Of external causes, the most prominent undoubtedly are, sudden vicissitudes of temperature, especially when joined with dampness of the atmosphere, whereas the mere degree of coldness exercises but little influence upon its development. Of course the action of damp and cold is markedly increased by insufficient clothing.

Complications.—We have already alluded to the occurrence of chorea and cerebral symptoms in connection with rheumatism (see article on chorea), and the most important and frequent of all these complications, the various inflammations of the membranes of the heart, have been treated of under the head of diseases of that organ.

PROGNOSIS.—Uncomplicated rheumatism in childhood, though at times severe, is scarcely ever fatal. When complicated, however, with endo- or pericarditis, the gravity of the prognosis must depend upon the extent and severity of the inflammation. For although, even when the heart is seriously involved, the child frequently survives the acute symptoms, it too often bears with it the seeds of premature death, in an organic disease of that organ.

DIAGNOSIS.—The diagnosis of acute rheumatism, after the appearance of the articular symptoms, can hardly present any difficulty. When, however, marked rheumatic fever, accompanied merely by vague pains, precedes by several days the development of any local symptoms, the diagnosis must remain uncertain, or we may be led to regard as rheumatism one of those cases of phlegmon of the deep tissues of the extremities, such as is alluded to in the introductory essay of this work. In addition to this, we must be careful to distinguish the articular affections occurring in pyæmia, or those supervening upon attacks of small-pox and scarlet fever, which are probably also of pyæmic nature. The diagnosis in these cases must be chiefly established by attention to the general symptoms and the patient's

history; to the occurrence of repeated chills or irregular febrile paroxysms, the diarrhœa, the greater degree of prostration and more rapid emaciation, and the more frequent fatality. The joints involved in these latter affections present large collections of creamy pus, and the articular cartilages are discolored, or eroded and destroyed in patches.

Finally, Rilliet and Barthez cite a case (from *Jour. Hebdomadaire*, t. ii, p. 260) of hemorrhage under the periosteum of the clavicles, which simulated rheumatic inflammation of the sterno-clavicular joints, but which could be distinguished by ordinary attention to the general symptoms, in case of the occurrence of such a rare condition.

From our own experience we should think that, during the early stage of rheumatic fever, the affections with which it might be most readily confounded, are pleurisy and pneumonia, and typhoid fever.

The absence of the physical signs of the two former affections, and of the diarrhœa and delirium of the latter, should, we think, lead the physician to suspect the rheumatic nature of the attack. And if, in addition, there should be any fixed pain about the limbs, or unusual soreness and pain on being moved, or if any sign of cardiac inflammation be detected, this suspicion would be confirmed. Thus in a case seen by us, where at first the height of the fever and the great thirst led us to suspect the existence of pneumonia or pleurisy—of which, however, no physical signs could be detected—the occurrence, on the third day, of complaints of pain in the right groin, led to a more careful examination of the heart, where the presence of a soft, faint mitral murmur, declared the nature of the attack.

TREATMENT.—The indications for treatment presented by acute rheumatism have been universally recognized as uniform, but the measures adopted to meet them embrace almost all known remedies.

The prominent indications are :

1. To aid in the elimination of the rheumatic poison, which has set up the specific inflammations and fever.
2. To relieve pain.
3. To guard assiduously against all complications, and to aid convalescence by suitable nourishment and tonics.

Among the remedies which appear to be most productive of benefit, are *alkalies*, especially the bicarbonate of soda or potash and the acetate of potash, as recommended by Garrod; and we have ourselves employed these in the majority of our cases with considerable satisfaction. The formula which we are in the habit of using is the following :

R. Potass. Acetat.,	3j.
Potass. Bicarb.,	3j.
Tr. Opii Deodor.,	gtt. xxiv.
vel Tr. Opii Camph.,	f 3 ij.
Syr. Zingiberis,	f 3 j.
Aquæ,	q. s. ad f 3 iij.

Ft. sol. S.—A teaspoonful every two or three hours, at four or five years of age.

When the fever is very marked, nitrate of potash, in carefully graduated doses, may be substituted for the bicarbonate in the above mixture. We

have usually employed quinia in full doses in addition, frequently giving it in the form of very small suppositories to avoid the risk of irritating the stomach by a multiplicity of doses. Aconite or digitalis may be used instead of or in addition to the nitrate of potash, especially if the action of the heart is much excited. Their administration is, as will be seen, imperative in case of cardiac complication. The bromide of ammonium has been recommended of late as of value in acute rheumatism. We have not been sufficiently pleased with its effects, however, to lead us to substitute it for the alkalies above mentioned, excepting in cases where a high degree of nervous restlessness with sleeplessness exists.

Salicylic acid and salicylate of soda have established themselves in our estimation as of positive value in certain cases of acute rheumatism, although we are well aware that their action is not uniformly favorable, owing presumably to some unrecognized differences between apparently similar cases. We have of recent years used salicylate of soda in a number of instances of acute rheumatism, both with and without marked articular inflammation, and have found it serviceable in a majority of them in relieving the fever, pain, and local lesions. As in the case of adults, however, if benefit does not follow its use in the course of 3 or 4 days, it is better to stop it and substitute the alkaline mixture.

The iodide of potassium is most serviceable in subacute cases affecting the muscular or tendinous tissues, such as we have above described, or in somewhat chronic cases of the articular form. We can fully indorse, moreover, the statements of Rilliet and Barthez, that more benefit is to be derived from large doses of this salt than from any other drug in the inflammatory complications of rheumatism (endo- and pericarditis, and pleurisy).

Iron, particularly in the form of Basham's solution of the peracetate of iron, should be given so soon as the intensity of the fever has mitigated. The necessity for this remedy is but too often seen in the sallow, anæmic appearance of convalescents from rheumatism, which proves the rapid and extreme disintegration of the red corpuscles of the blood during an acute attack of this disease.

When the acute symptoms have subsided, the alkalies may be diminished and withdrawn, and *quinia* in the dose of one grain every four hours, at the age of five years, may be given in connection with opium.

The following formula is one we frequently use for the administration of these remedies in this and other conditions:

R. Quiniæ Sulph.,	gr. xxiv.
Liq. Morph. Sulph.,	fʒ ij.
Acid. Sulph., Dil.,	gtt. xxx.
Curaçoa,	fʒ ij.
Syrupi,	fʒ vj.
Aquæ,	q. s. ad fʒ iij.

Ft. sol. S.—A teaspoonful every four hours, at four or five years of age.

To fulfil the second indication, the mitigation of pain, opium must be given in proportion to the severity of the suffering. It is best given in small doses at short intervals, and by administering it in combination with

ipecacuanha, as in the form of Dover's powder, we derive the double benefit of a sedative and diaphoretic action. We have already given the formula by which we usually direct it in this disease.

In addition to the other remedies, particular attention must be paid to the condition of the bowels, and if constipation exists, as is very frequent, mild saline laxatives or laxative enemata should be administered as frequently as required. Anything like purgation, however, should be avoided, on account of the excruciating suffering often produced by the movements necessary to have a stool. We desire, however, to call attention to the fact that young children with this disease may persist in lying in one fixed position for even several days, dreading to be touched; so that there is added to the inevitable pain of the disease, the distress occasioned by the long-continued contact of single points of the opposing articulating surfaces. Under these circumstances it is wise, and greatly promotes the comfort of the patient, to gently change the angle of the limbs by arranging pillows so as to support them and alter their direction.

In regard to the last indication—the prevention of complications—the most important means is the avoidance of all exposure of the patient to damp or to changes of temperature. In the fulfilment of this, the greatest care must be paid to the temperature of the sick-room, to the clothing of the patient, and to the mode of conducting all our examinations. Dr. Chambers, in his admirable lectures upon this subject (*Clinical Lectures*, American edition, pp. 156, 177, etc.), dwells with special force upon this point, and enjoins the exclusive use of blankets and flannels for the bedding and clothing of patients with rheumatism, and gives the following summary of his observations of the effects of this precaution alone in the treatment of nearly two hundred cases of rheumatism: "That bedding in blankets reduces from sixteen to four, or by three-fourths, the risk of inflammation of the heart, diminishes the intensity of the inflammation when it does occur, and diminishes still further the danger of death by that or any other lesion."

The importance of confinement to bed in this disease is difficult to overestimate; the inflamed condition of the joints absolutely demands it, and the tendency to cardiac inflammation warns us to save the heart all unnecessary exertion, which strict attention, as above recommended, to the equable warmth of the surface, effects better than any other means.

As to the diet in this affection, we must be guided by the acuteness of the symptoms and the condition of the patient. If the fever be marked, and the child vigorous, a diet chiefly consisting of milk and water is best suited to the early part of the attack, but so soon as the febrile stage has passed off, or when the patient is of feeble constitution, we may give soft-boiled eggs, and meat broths, with advantage; and frequently we will find concentrated nourishment and a moderate amount of stimulus required towards the close of the case.

LOCAL TREATMENT.—As severe arthritis is much more rare in rheumatism in children than in adults, it is less frequently necessary to employ systematic local treatment for the relief of the inflamed joints.

Local depletion is rarely justifiable; and if the swelling and congestion

are severe, relief may usually be obtained from the local application of cold wet compresses. More commonly we apply tincture of iodine freely over the affected joints, and envelop them in raw cotton held in place by a light bandage. If the pain is great, a small and mild blister may be used with advantage; or the joints may be bathed with a sedative liniment, such as the following:

R. Tr. Opii Deodoratæ,	
Tr. Aconiti Radicis,	aa f3ij.
Lin. Chloroformi,	f3iss.
Ol. Olivæ,	q. s. ad f3iv.
M. et ft. lin.,	

and then enveloped in bats of wool and covered with oiled silk.

It is important to pay attention to the position of the affected parts. The joint should be slightly flexed and carefully supported on small down pillows or rolls of raw cotton. From time to time the angle at which the joint is flexed should be changed very gently. At a later stage, when the acute inflammation has subsided, the absorption of any thickening or exudation that remains may be hastened by friction with stimulating liniments, by the continued use of iodine, and by gentle uniform pressure by a skilfully applied bandage or by a plaster-of-Paris dressing.

COMPLICATIONS.—In those cases where, despite our precautions, the membranes of the heart are threatened with inflammation, as evinced by sudden pain in the cardiac region, frequency of pulse, and oppression—even before the development of any murmurs—we should lose no time in employing local depletion by leeches or cups, abstracting as much blood as the urgency of the symptoms and the vigor of the constitution justify.

If, for any cause, local depletion should appear contra-indicated, the immediate application of a blister is to be recommended.

After the removal of the cups or leeches, or blister, warm mush-poultices should be applied steadily over the whole præcordial region.

It is our custom to order immediately the iodide of potassium in combination with the acetate of potash. The dose of the iodide must be carefully graduated to suit the age and susceptibility of the child; but usually one grain every four hours may be safely ordered at three years of age, and this may cautiously be increased if the symptoms are urgent. Digitalis should be given at the same time in full doses, as two or three drops of the tincture every four hours at three years of age; its effects being of course carefully watched at short intervals. We have already expressed, when speaking of diseases of the heart, our sense of the importance of maintaining careful observation and judicious treatment in such cases after the acute symptoms have subsided, since it is sometimes possible to secure complete removal of organic lesions occurring at such an early age.

ARTICLE II.

SCROFULA.

It does not seem appropriate, in a work whose chief character is designed to be practical, to enter upon a full discussion of the important pathological questions connected with the subject of scrofula, particularly in regard to its relations to simple chronic inflammation on the one hand, and to tuberculosis on the other. Indeed, in some respects these questions may be said to be still in such an unsettled state that no definite position in regard to them can be assumed with confidence. We propose, therefore, to confine our remarks at present chiefly to a description of the most marked manifestations of scrofula as generally recognized, and to a discussion of the appropriate treatment.

DEFINITION; CHARACTERS.—The term scrofula is of very long standing. It appears to have been originally applied to a peculiar cachectic state of the system in which there is a special tendency to enlargement of the lymphatic glands. Subsequently it has been employed in so many and such varied senses as to make it difficult in many cases to decide in which way it is meant to be understood. We ourselves would be understood to employ it much in the old sense, to indicate a peculiar constitutional condition in which there is a "*vulnerable*" or irritable state of the lymphatics, which renders them liable to become enlarged from trifling causes, and at the same time indisposed to healthy reparative action; and which is also apt to manifest itself by various obstinate chronic inflammations of the skin, mucous or synovial membranes, or bones.

Scrofula is undoubtedly closely associated with tuberculosis. It very often happens that the children of tuberculous parents are scrofulous. And again we frequently observe that patients who have suffered with some chronic scrofulous affection become the subjects of tuberculosis, even of the most acute miliary form. So also there is a stage, that of yellow cheesy degeneration, in which it is not possible to distinguish between products of a scrofulous and of a tuberculous character. Still, however, we do not regard these two cachexiæ as identical, and enough points of difference can be indicated to fully support this opinion. Tuberculosis, it is true, often follows scrofulous affections, just as it follows any other condition attended with the formation of cheesy deposits, which may infect the system and give rise to acute miliary tuberculosis. On the other hand, it is not common for tuberculous subjects to develop any manifestations of scrofula; and, as West points out, we frequently see whole families which display one or the other diathesis in its most intense form, and yet perfectly uncomplicated. Scrofula, moreover, is, far more markedly than tuberculosis, a disease of early life. The most common and characteristic of its manifestations also are very different from those of the latter disease; it affects the bones, the skin and adjacent mucous membranes, the glands, the synovial membranes in preference to the serous membranes, the lungs, the solid abdominal organs, and the alimentary and respiratory mucous membranes. These differences in the leading pathological tenden-

cies of these two great cachexiæ, as well as the many points of difference in the physical peculiarities of children who are liable to tuberculosis or scrofula, are clearly and forcibly pointed out by Jenner in a clinical lecture published in *The Medical Times and Gazette*, 1860, p. 259.

CAUSES.—Scrofula is, we think, in many cases undoubtedly due to inherited predisposition. As in the case of other cachexiæ, the actual disease is not transmitted from parent to offspring, but merely so strong a tendency to its development that in some cases no care or favorable hygienic influences will overcome it. Not only do we meet with scrofula in the children of parents who themselves have been scrofulous, but also in cases where a feeble and vitiated constitution has been inherited from parents affected with tuberculosis or constitutional syphilis. In other cases, it is undoubtedly acquired after birth, appearing in children born to parents of sound constitution. The causes which tend to thus develop it act by impairing the nutrition, and include such influences as insufficient, improper food, protracted exposure to damp, cold, and especially to vitiated atmospheres, attacks of certain diseases, which like measles, typhoid fever, and chronic malaria, exercise a remarkably injurious action upon nutrition.

SYMPTOMS.—Although by no means all scrofulous children present the same physical peculiarities, there are yet certain features so commonly met with in such subjects as to have led to their recognition as forming together the symptoms of a *scrofulous diathesis*. Thus, as a rule, such children are heavy and lethargic in mind, and of phlegmatic temperament, with dull expression, and thick, opaque skin. The features are apt to be coarse, especially the lips and nose; the lymphatic glands are perceptible to the touch; the abdomen is apt to be full and large; and the bones are large, with coarse, thick ends.

There is nothing peculiar or pathognomonic about the special manifestations of scrofula. Almost all of them may also appear as simple idiopathic affections due to some definite exciting cause, in children of entirely sound constitution. That which characterizes these same affections when they occur in what we term the scrofulous form, are the trivial causes which excite them, the inveterate obstinacy with which they persist, and their association with other analogous phenomena in the same subject. There is also in some cases a certain order of succession of the manifestations of scrofula which has even led to the division of its course into three stages, corresponding somewhat to the classic phases of constitutional syphilis. Thus in the earliest stage, the lymphatic glands and skin are chiefly affected; subsequently affections of the mucous membranes and cellular tissue make their appearance; and in the final and most aggravated form the bones and viscera suffer. We cannot affirm, however, that this division and order of succession of the manifestations of scrofula is by any means constant or even marked in many cases.

Most of these manifestations appear as chronic inflammation of the part affected. At times such inflammation seems to arise spontaneously, while more frequently some more or less trivial exciting cause can be assigned. Thus the scrofulous enlargement of any group of glands is apt to be preceded by irritation of the area whose lymphatics pass to the affected glands,

as, for instance, enlargement of the cervical lymphatics follows eruptions on the scalp or behind the ears, or attacks of sore throat.

Among the most frequent affections which are generally classed as scrofulous may be mentioned, without any reference to their frequency of occurrence, enlargement of the superficial lymphatic glands, cutaneous eruptions, especially of the vesicular and pustular varieties, small subcutaneous abscesses, chronic, inflammation of the mucous membranes which are continuous with the external skin, as of the conjunctiva, the membrane of the external auditory meatus, that of the nose, and of the vulva and vagina, chronic effusions in the synovial membranes, chronic ostitis with caries.

The reader is referred for more detailed accounts of these numerous local scrofulous affections to the special works which treat of the diseases of the skin or organs of special sense, or to general treatises upon surgery. Our own purpose is of necessity limited to a discussion of the general symptoms and treatment of the scrofulous cachexia, rather than of its numberless local manifestations. In the most advanced and severe forms of scrofula, lesions of various internal viscera may be developed. Among the most frequent and clearly marked in their nature, of these, are caseous bronchitis and pneumonia, and albuminoid degeneration of the abdominal viscera, the liver, spleen, and kidneys. Bronchitis and pneumonia at times appear in forms which entitle them to be regarded as scrofulous from the first. At other times they apparently originate as acute inflammatory affections, but which owing to the strong scrofulous diathesis of the patient, pass into a chronic form characterized by the low grade of the morbid products developed, by the obstinate and intractable course the affections run, and by the marked tendency to the occurrence of caseous degeneration and destructive changes in the diseased parts. In this condition a sudden development of miliary tuberculosis not rarely occurs, either in the adjacent portions of the diseased organ or throughout the other parts of the system. The exact nature of the primary changes in such cases is, at the present moment, one of the most unsettled and disputed points in pathology. The reader will find a tolerably full description of the lesions and symptoms under the head of pulmonary phthisis.

The exact relation of albuminoid degeneration of the viscera to scrofula is also somewhat uncertain. Although one of the most frequent of the unfavorable sequelæ of scrofulous affections, it cannot itself be regarded as scrofulous in nature, since it makes its appearance in connection with other cachectic states of the system. The attempt of Dickinson to associate it with the changes in the blood and tissues caused by prolonged suppuration (which so often occurs in scrofulous disease of the bones, joints, or glands), has not been altogether successful. Although it is undoubtedly true that in many cases where albuminoid degeneration has been developed there has been previous prolonged suppuration, there are many exceptions where the visceral lesions have apparently been induced directly in connection with the scrofulous or other cachexia. The occurrence of this sequel must always be anticipated with anxiety in protracted and severe cases of scrofula. Although usually involving, simultaneously or

in rapid succession, the various abdominal organs, the liver, spleen, kidneys, and gastro-intestinal canal, it may present a marked localization, for an indefinite time, in any of these parts.

When one of the above solid organs is affected with advanced albuminoid degeneration, it is found enlarged, though still preserving its original shape; the peritoneal capsule is unchanged; and on section the tissue presents a homogeneous, waxy, or lardaceous appearance, which is associated, when the section is examined by transmitted light, with abnormal translucence. The intimate nature of the change consists in an infiltration of the organ with a peculiar structureless albuminoid neoplasm or exudation. This first affects the walls of the arterioles, and later the glandular cells of the organ.

When the kidneys are involved, there is usually œdema, which appears early and increases rapidly; the urine is abundant, clear, with but slight reduction in its specific gravity, contains a large amount of albumen, and deposits numerous hyaline tube-casts. Albuminoid disease of the liver and spleen usually coexists. These organs are markedly enlarged, as can readily be detected by palpation and percussion. There is usually abdominal dropsy, with distension of the subcutaneous veins of the abdominal walls; and frequently there is also albuminuria and diarrhœa from coexisting disease of the kidneys and intestine. We have much less frequently observed marked albuminoid disease of the gastro-intestinal canal than of the solid abdominal organs, as above described. When it occurs, the walls of the stomach or intestine are thickened and present a peculiar homogeneous, glistening, and infiltrated appearance. The same microscopic changes are found as already described. The lesion of the mucous membrane is usually attended with chronic diarrhœa, and, if the stomach is also seriously involved, frequent and obstinate vomiting. Hemorrhages from the bowels have been observed, but much less frequently than in the same condition in the adult. The general symptoms which mark the later stages of fatal cases of scrofula, especially when these serious visceral lesions have been developed, are expressive of the most profound anæmia and malnutrition.

DIAGNOSIS.—The recognition of the existence of scrofula depends, not so much upon the presence of any special symptom or local affection, as upon the general marks of the scrofulous diathesis, the existence of hereditary tendency, or of some of the well-known exciting causes; the spontaneity and order of evolution of the phenomena; and finally their intractable resistance to the ordinary remedies, and the marked benefit which is often found to follow the use of special anti-scrofulous treatment.

PROGNOSIS.—The prognosis in cases of scrofula must of course depend upon the intensity of the diathesis, the gravity of the local manifestations, and the hygienic surroundings of the child. When the general health is fair, and the only scrofulous affections present are superficial, although the case is likely to prove obstinate and tedious, complete recovery can often be insured. It must never be forgotten, however, that such children are liable to the recurrence of scrofulous disease in some other form, and even to the development of the grave visceral lesions we have above al-

luded to. In the latter and more advanced stages of the cachexia, when serious disease of the osseous and glandular tissues exists, the prognosis becomes in the highest degree unfavorable.

TREATMENT.—A great variety of local treatment—both medicinal and operative—is required for the various local scrofulous affections. We shall not, of course, attempt even to refer to these, but shall merely allude to the general principles that we think of prime importance; that, in the first place, all such affections should be cured as promptly as possible, and also that, in their treatment, the essential value of proper hygiene and constitutional remedies should never be forgotten.

The *preventive* treatment is of the greatest value; but it merely consists in the employment, with special and continued care, in the case of any child who probably possesses a scrofulous diathesis, of all those precautions as to diet, dress, exercise, and residence, which sound hygiene would dictate. In children born of scrofulous or tuberculous parents, a wet-nurse should be secured even if the mother is able to suckle them; and under no circumstances should the attempt be made to rear them on artificial food. Later, when the child has been weaned, the diet should be of the most nutritious and digestible character, especially containing a large proportion of well selected animal food. The utmost care should also be exerted as to the dress, in order that it may be adapted to the season and sufficiently warm to prevent the child from contracting any of the catarrhal attacks, to which there is so great a liability in the scrofulous diathesis. Outdoor exercise in fair weather and gymnastic exercises indoors, when it is unfit for the child to be exposed to the weather, must be enjoined. As a general rule, it may be said that the child should be encouraged to spend as much time out of doors as possible, when the weather is fine, dry, and sunny. If the circumstances of the parents admit of it the residence of the child should be chosen in an elevated, dry, and comparatively open part of the city, and for several months in each year it should be taken to the sea-shore, or to some elevated inland locality. While at the sea-shore, sea-bathing should be regularly followed, and throughout the rest of the year brine-baths, made with either bay-salt, or rock-salt, may be used daily. All forms of catarrhal inflammation, as angina, conjunctivitis, enteritis, and the like, should receive prompt and careful attention, and be cured as soon as possible, since there is danger, if they are allowed to continue, not only of their becoming chronic and extremely obstinate, but also of troublesome glandular enlargements being induced by the protracted irritation. After any of the local manifestations of scrofula which we have above enumerated have made their appearance, the above hygienic management must be sedulously persisted in. There are also various medicinal substances which exercise a beneficial effect by their alterative and tonic action upon the general nutrition. Among these the best are cod-liver oil, various preparations of iodine and of iron. Cod-liver oil may be used alone, or combined with the compound syrup of the phosphates of the alkalies and iron.

The preparations of iodine most frequently used, and which we have been led to prefer, are the compound tincture or solution of iodine, in the

dose of from two to four drops three times a day, and the iodide of potassium, either alone, given in solution, as follows :

R. Potassii Iodidi, gr. xxiv.
Decoct. Sarsaparillæ Comp., f℥iv.

Ft. sol. Dose, a dessertspoonful to a tablespoonful thrice daily, at three to five years of age.

Or in combination with the iodide of iron, as follows :

R. Potassii Iodidi, gr. xlvij.
Syr. Ferri Iodidi, f℥ij.
Syr. Zingiberis, f℥x.
Aquæ, f℥iss.

Ft. sol. Dose, a teaspoonful thrice daily in water, at five years of age.

It is probable that the above is the best mode in which iron can be administered, though it is often desirable to give it in association with quinia or some other vegetable bitter, in order to stimulate the appetite and digestion.

Mercury, despite its powerful absorbent action, is not to be recommended for the treatment of scrofulous enlargement of the glands, in any form in which it is likely to produce its characteristic effect upon the blood. We are satisfied, however, that in some very obstinate cases which resist all other modes of treatment, minute doses of the bichloride or biniodide may be employed without risk, and with much advantage.

Arsenic deservedly occupies a high place among the internal remedies in scrofula. It may be given in combination with iron or quinia; or in some cases will be found of service in the form of small doses of Donovan's solution, the liquor hydrargyri et arsenici iodidi of the U. S. Pharmacopœia.

When circumstances permit, the use of certain mineral waters, particularly if the child can have the advantage of a temporary change of residence to the locality of the spring, is often attended with marked benefit. The waters which prove most useful are the sulphurated and iodo-bromated.

ARTICLE III.

TUBERCULOSIS.

THIS subject has received from many authors upon diseases of children, far less attention than it merits, under the idea that it is merely a repetition, upon a small scale, of the same disease in the adult, and not possessed of any individual characteristics. In fact, however, tuberculosis in childhood is an affection possessing characters and presenting symptoms entirely special, and differing from its manifestation in adult life both in causes, locality, and clinical history.

CAUSES.—The causes which exert most manifest influence in its produc-

tion are hereditary tendency, and all those debilitating agencies which act directly or indirectly upon nutrition. Of these latter causes, early weaning is the most prominent. Thus, we have met with a case where a healthy woman, the mother of several vigorous children, all of whom she had nursed, gave birth to one which she was unable to suckle, and this child, after pining for some months, died of an attack of tubercular meningitis. A bad quality of the nurse's milk, or improper artificial food after weaning, also exert a powerful influence in the production of tuberculosis; and not unfrequently its development has been traced to repeated attacks of indigestion or diarrhœa.

It has also a tendency to develop itself after certain acute affections, especially in children predisposed by hereditary influence. Of these diseases, rubeola, pertussis, typhoid fever, and, according to Greenhow, variola, are most frequently followed by tuberculosis.

There is still some difference of opinion in regard to the rôle which pneumonia plays in the development of tuberculosis. When the two coëxist, the inflammation is by some regarded as a secondary affection, induced by the deposit of tubercle in the lung; while by others it is held that, amongst predisposed children, it is the pneumonia which causes the development of tuberculosis of the lung. We believe that pneumonia occupies each of these relations in a certain number of cases; but reliable statistics upon this point are still too scanty to determine the exact proportion.

Of recent years the influence exerted by foci of cheesy degeneration in the production of general tuberculosis has been established by careful clinical and experimental observation. In childhood the most frequent seat of such foci is in connection with the lymphatic glands, though they may also occur in connection with disease of the bones or with chronic ulcers.

ANATOMICAL APPEARANCES.—The most frequent seats of tubercular deposit in the child are the brain, constituting tubercular meningitis, which has already been treated of at length; the bronchial glands, the lungs, and the mesenteric glands and peritoneum. It is, however, one of the distinguishing features of tuberculosis in the young subject, that it is apt to involve several viscera simultaneously, while not unfrequently the lungs remain free. Thus, in 312 children in whom Rilliet and Barthez found a deposit of tubercle in one or more of the viscera, the lungs were healthy in 47; while in 123 similar instances in the adult, Louis only found one such exception.

Locality.—In *bronchial phthisis*, which generally accompanies pulmonary phthisis, but also exists as a separate affection (though according to Bouchut, this is a rare occurrence), the glands are much enlarged and inclose tubercular matter, frequently in large proportion. This is especially marked in those glands which lie along the trachea and around its bifurcation, and, when many of them are involved and adherent to each other, they form masses varying in size from a hen's egg to a large apple. The deposit, which in by far the majority of cases exists as infiltrated tubercle, does not usually soften, though cases are recorded where such softening has occurred, and the fluid has been discharged through an opening into

a bronchus. Obsolescence and calcification, however, are quite common terminations of bronchial tubercles; and when the lungs do not become involved in the morbid process, a cure may be effected by these transformations. Calcified tubercle may be eliminated through a communication between the gland and one of the air-passages; and a few cases are also reported where the œsophagus, trachea, and even the pulmonary artery have been perforated in this manner. Most of these tuberculous glands are inclosed in a distinct and dense capsule, which may attain the thickness of one or two lines, and is usually quite vascular. This fibrous capsule is due to the hypertrophy of the originally delicate cellular investment of the gland.

Pulmonary Phthisis.—The anatomical characters of tuberculosis of the lungs in children present several peculiarities, as distinguished from the same disease in adults. Thus gray granulations and crude miliary tubercles frequently exist in the lungs, independently of each other and of any other form of tubercular deposit. In the adult, Louis discovered miliary tubercles unassociated with gray granulations only in 2 out of 123 cases, or in 1.6 per cent.; and gray granulations alone in but 5 more, or 4 per cent.; while in the child, Rilliet and Barthez found miliary tubercles without gray granulations in 107 out of 265 cases, or in 40.4 per cent.; and gray granulations alone in 36 instances, or in 13 per cent.; and the observations of West, "which are based on 102 cases, yield 20 instances of the presence of miliary tubercles alone, and 17 of the presence of gray granulations alone in the tissue of the lungs."

The great frequency with which the so-called yellow infiltrated tubercle is observed in early life constitutes another anatomical peculiarity, Rilliet and Barthez, and West, having found it in from 23 to 33 per cent. of their cases. This condition rarely exists as an isolated state, but is found in conjunction with gray granulations and crude yellow tubercles, and not unfrequently also with advanced tuberculization of the bronchial glands.

The rare occurrence of cavities in the lungs is a most striking peculiarity of phthisis in children. It is probably no exaggeration to say that in adults, cavities are found in the lungs in 90 out of every 100 cases of tuberculosis; whilst out of 265 cases of tuberculosis of the lungs in children that came under the notice of Rilliet and Barthez, only 77, or 29 per cent., presented cavities; they existed only in 23.5 per cent. of West's cases, and Bouchut found them in but three out of 36 cases.

Occasionally the cavities resemble the vomicæ found in the lungs of adults, and this occurs with more frequency as we advance beyond the age of six years. In other cases, the excavation is produced by the softening of very small tuberculous deposits, distinct, though in close proximity, which form small vacuoles, communicating with each other and with the neighboring bronchial tubes. All three of M. Bouchut's cases appear to have been of this form.

In addition to these two varieties of tuberculous cavities, there is still a third, produced by the simultaneous softening of considerable portions of a lung affected with yellow infiltration. This action, which is most com-

monly met with in very early life, and in cases which progress with great rapidity, pervades the whole of the tissue affected, instead of producing a central cavity. Cavities of this kind sometimes form very quickly, and involve large portions of lung, the whole of one lobe even being converted into a mere sac, with thin walls.

There is another form of excavation occasionally noticed, which is not a true pulmonary vomica, but the result of the softening and evacuation of a tuberculous pulmonary gland. The diagnosis, however, may be rendered easy by reflecting that a pulmonary cavity of such small dimensions is hardly ever solitary, unless it proceeds from the softening of tubercular infiltration, whilst the deposit of tubercle which takes place in the neighborhood of a diseased pulmonary gland is always in the form of distinct deposits, not of tubercular infiltration (West).

The last anatomical peculiarity, already alluded to, of pulmonary phthisis in children, is its frequent complication with tubercular deposit in the bronchial glands.

Peritoneum.—Tubercular deposit on the peritoneum rarely or never occurs without the presence of a similar disease in some other parts of the economy. It may be either general or partial in its disposition, though it is far more frequently the latter. The deposit varies also in its character, appearing generally in the form of yellow granulations or of miliary tubercles, either isolated or united into small masses. Gray granulations, however, are also of quite frequent occurrence.

The relation which the tubercles bear to the peritoneum is not uniform, though they are more frequently found deposited on its surface than beneath it. In 86 cases examined by Rilliet and Barthez, the seat was as follows: intra-peritoneal in 40; extra-peritoneal in 22; both intra- and extra-peritoneal in 14; in the other 10 cases the exact seat was doubtful.

When the deposit involves the entire extent of the serous membrane, we find the anterior parietes of the abdomen adherent to the subjacent structures, and the viscera so matted together and adherent, as to form an almost inseparable mass. More frequently, however, the tuberculization is partial, and even limited to the vicinity of a single organ. The peritoneum investing the diaphragm, especially that portion which is in contact with the liver or spleen, or the adjacent parietal peritoneum, is very often affected; and as tubercles rarely fail to be deposited in the peritoneum covering these viscera, we find them firmly adhering to the diaphragm or abdominal wall.

In some cases the omentum is the chief seat of the disease, and may either present numerous gray granulations scattered through its folds, or may be thickened or matted together from a kind of grayish tubercular infiltration, due to the coalescence of innumerable minute gray granulations. It is more rare to find the tuberculization limited to the intestines, merely causing adhesion of the adjoining coils.

In examining the adhesions which are almost universally found to exist between the various organs and portions of peritoneum affected, we find them to present two elements. In the first place, the tubercular deposits on the adjoining surfaces gradually coalesce as they increase in size, and

finally unite the surfaces by more or less extensive patches of tubercular matter. And again, at the same time, the subacute inflammation caused by their presence leads to the formation of cellular and fibrous adhesions as in cases of simple peritonitis. This is well seen in cases where some coils of the intestine present tubercular adhesions to each other, forming masses which can only be separated by rupturing the walls of the bowel, while between other coils the adhesions merely consist of delicate and easily lacerated cellular bands.

It is a well established fact that the tubercular granulations on the surface of the peritoneum have no tendency to perforate this membrane; but that the perforations which are occasionally found, especially in the walls of the intestines, are due to the development and softening of the sub-peritoneal tubercles, which always tend to penetrate into its cavity. This same law holds elsewhere, and it is on this account that the adhesions which so constantly form between tuberculous membranes are of such great value in preventing the escape of foreign matters into the serous cavities. In the intestines this action can be traced even further, and when tubercles exist under both layers of the peritoneum at a point of adhesion between two folds of intestine, as softening advances the layers of peritoneum are destroyed, and the little collection of tuberculous pus remains confined only by the inner coats of the two layers of bowel. Sooner or later these also break down, the softened tubercle is discharged into the bowel, and a direct communication established between distant parts of the intestinal canal, as between a fold of the ileum and the ascending or descending colon. This perforation, then, is not caused by tuberculous ulceration of the mucous membrane; nor does this latter affection bear any fixed relation to the degree of tuberculization of the peritonæum.

There is generally some deposit of tubercle in the mesenteric glands in these cases; and when the splenic portion of the peritoneum is involved, we frequently find an abundant deposit in this organ.

Tuberculization of the *mesenteric glands*, or *tabes mesenterica*, offers few anatomical features in addition to those present in bronchial phthisis. It is, moreover, far from being a frequent form of the disease, for although, according to Rilliet and Barthez, some tubercle is found in these glands in one-half of all tuberculous subjects, it exists in considerable quantity only in one out of every sixteen of the whole number. The deposit generally appears as infiltrated tubercle, though not unfrequently miliary tubercles are present. The glands attain a size varying from that of an almond to a pigeon's egg, and occasionally, from the aggregation of several enlarged glands, a mass is formed double the size of the child's fist.

The capsule which surrounds them is usually more delicate and less vascular than the same structure in tuberculous bronchial glands. The tubercular deposit here, as elsewhere, is liable to undergo calcification or softening, the latter process being more frequently met with.

Owing both to the yielding nature of the abdominal walls, which do not resist the forward growth of the mesenteric glands, and to the mobility of the adjacent viscera, we never see the same degree of compression exerted on surrounding structures, as is noticed in tuberculization of the bronchial glands.

Occasionally, however, adhesions may form between a tuberculous mesenteric gland and a fold of the intestine, and ultimately result in perforation of the bowel.

In thus describing these various lesions as being all tuberculous in their essential nature, we have purposely employed this term in the somewhat inaccurate and vague sense which was assigned to it until within the past few years.

Recognizing, as we distinctly do, but one elementary form of tuberculous deposit, the gray granulation or miliary tubercle, which may, it is true, undergo cheesy degeneration, it is evident that many of the cases in which extensive and uniform cheesy deposits are found, rather depend upon scrofulous inflammation of the part than upon true tuberculous formation. It is comparatively rare to meet with such cheesy deposits in the lungs in children, while, as already described, they occur very frequently both in the bronchial and mesenteric glands. And, therefore, we are disposed to believe that in many cases of so-called bronchial or mesenteric phthisis, the enlargement and degeneration of the glands are really due to an inflammatory process of a low and unhealthy type, excited by the previous occurrence of attacks of bronchitis or enteritis, and leading to the formation of a cacoplastic lymph, which soon undergoes cheesy degeneration.

It is in this way, doubtless, that the comparatively numerous cases are to be explained in which such deposits soften and are evacuated, or undergo partial absorption and calcification, and where ultimately the child's health is restored. We have preferred, however, in the present edition, for practical purposes, to group the descriptions of these various conditions under one common head, being unwilling to separate them until more extended study shall have more clearly demonstrated the degree of resemblance which exists between true tuberculous matter and such cacoplastic inflammatory formations.

SYMPTOMS.—The symptoms of tuberculosis in children may be studied under the forms of bronchial phthisis; acute and chronic pulmonary phthisis; and tuberculization of the peritoneum and mesenteric glands.

Bronchial Phthisis.—In addition to the general symptoms of tuberculosis, which will be fully given under the head of pulmonary phthisis, the most marked symptoms of bronchial phthisis are those due to the mechanical effect of the enlarged and hardened glands upon the surrounding tissues. Our knowledge of the functions of the lymphatic glands is as yet so inaccurate that we are entirely unable to appreciate the symptoms of disordered action which are probably present in cases of extensive disease of these organs.

Bronchial phthisis occurs in its most marked form between the ages of two and six years; and in many cases appears to be developed after some severe attack of bronchitis, either accompanying measles or arising without apparent cause.

The cough which, in the early stage, is hacking and not very troublesome, soon acquires severity, and becomes intermittent, recurring in paroxysms like those of pertussis.

The respiration becomes habitually labored and oppressed, with a prolonged wheezing sound, as in asthmatic cases.

The veins of the neck are often greatly distended, the distension becoming extreme during the violent paroxysms of coughing; the face becomes puffy and œdematous, a condition occasionally extending to the upper extremities; and, as West points out, the superficial vessels of the thorax become enlarged, just as those of the abdomen do in cases of cirrhosis of the liver. The obstruction to the return of blood from the superior vena cava is further shown by the occurrence of epistaxis, or even of hemorrhage into the arachnoid; and the compression of the pulmonary tissue occasionally produces hæmoptysis and œdema of the lungs. Dr. Jenner has seen hydrothorax produced from compression of the vena azygos.

The œsophagus does not always escape the encroachment of the glands, but may be so compressed as to produce dysphagia.

It is hardly necessary to say that so long as the tubercular deposit remains small, it may exist without causing any symptoms, and it is only when several glands become infiltrated with tubercle, enlarged and firm, that they give rise either to the symptoms already enumerated, or to the physical signs below alluded to.

Physical Signs.—In estimating the value of these, it is necessary to constantly bear in mind the fact that the enlarged and tuberculous bronchial glands, while they still surround the trachea and bronchi, also come into contact with the spinal column, or, in a few cases, with the sternum. From their solidity, and the consequent readiness with which they are thrown into vibration, they transmit directly to the ear, and seem to exaggerate many respiratory sounds, which are in reality produced at a distance from the thoracic walls, and which are either entirely normal or dependent upon a small amount of disease.

It is also due to these relations, that the signs, both of auscultation and percussion, of bronchial phthisis are best detected at the summit of the lungs posteriorly, or at the level of the vertebræ with which the enlarged glands come into contact.

Our knowledge of these important considerations is chiefly due to the investigations of Rilliet and Barthez.

Percussion.—In the young child in health there is a diminution in resonance over the manubrium of the sternum, owing to the remains of the thymus glands: but, in some cases of marked bronchial phthisis this dullness extends both downwards and laterally to a varying but perceptible degree, owing to the projection of the enlarged glands into the anterior mediastinum.

More generally, however, as we have said, the tuberculous glands are in contact with the spinal column, so that we find dullness on percussion in the inter-scapular space as a pretty constant and characteristic symptom.

According to Dr. Jenner, it is common to have a cracked-pot sound on percussing the cartilages of the upper three ribs on one or both sides. This is due to the fact that the enlarged glands accompanying the bronchial tubes frequently extend under the anterior margin of the lungs, so that,

in percussing, the air-containing lung is compressed between the solid mass of glands behind and the in-driven parietes in front, and the air is forced out suddenly from the healthy layer of lung, producing the chinking sound.

Auscultation often reveals true tubular breathing over the upper part of the sternum, extending almost to the base of the heart. In those cases where a large bronchial tube is compressed or occluded, we, of course, find an enfeebled or extinct respiratory murmur over the corresponding lung segment.

Occasionally the enlarged glands compress the superior vena cava, and give rise to a permanent venous hum; or a systolic murmur, having its seat of greatest intensity at the second left interspace, may be produced by similar compression of the pulmonary artery.

There is one characteristic, however, of this form of phthisis, which is especially dwelt upon by Dr. West, and which it is well to bear in mind, to avoid being misled. This is the frequent occurrence of great fluctuations in the condition of the patient; so that, even when the rapid breathing, frequent cough, emaciation, and loss of strength would betoken a speedily fatal issue, a pause will occur in the progress of the disease, during which the diminution of any bronchitic complication, with partial disappearance of the dyspnœa and cough, and the return of flesh and strength to the little patient, all tend to awaken delusive hopes. In the great majority of cases, this respite is but brief, and the disease again resumes its onward course; but there are well authenticated cases on record in which the gravest symptoms have gradually disappeared, and the child has ultimately regained fair health. In these cases, the tuberculous deposit may either have undergone cretaceous degeneration, or having softened and formed an opening into a bronchus, have been expectorated.

The characteristics of bronchial phthisis, which we have been considering, are thus summed up by West:

“1. The frequent development of its symptoms out of one or more attacks of bronchitis.

“2. The peculiar paroxysmal cough which attends it, resembling that of incipient pertussis.

“3. The great and frequent fluctuations in the patient's condition, and the occasional and apparently causeless aggravation both of the cough and dyspnœa.”

Symptoms of Pulmonary Phthisis.—Valuable as are the general symptoms of tuberculosis in the adult, it is in the young child peculiarly that they reach their highest importance, owing either to the absence or the difficulty of appreciation of many symptoms which aid greatly in the diagnosis of phthisis in adult life.

It is necessary, therefore, to examine with the greatest care the child's hereditary tendencies, its past history, and its appearance and physical development. Thus it is in cases of inherited tuberculosis that we see its characteristic features most strongly marked, in the tall, slim frame; the firm bones, with small and yielding cartilages; the delicate diaphanous complexion; the fine, silky hair; the active, often precocious intelligence;

the ease with which the general health is affected by slight causes, and the peculiar proneness to catch cold on the least exposure. By careful attention to these and other similar points, as much, or often more valuable information can be obtained in the phthisis of children, than from the most careful investigation of the physical signs.

In enumerating the symptoms, it is unnecessary to detail those which exist in common with pulmonary phthisis in the adult, save to point out any particular in which they may differ as seen in the young subject.

The *cough* varies much, in accordance with the varying amount of bronchial irritation, being at one time scarcely troublesome, or so aggravated and accompanied with such violent dyspnœa, from some intercurrent attack of bronchitis, as to threaten immediate death.

It not unfrequently has a somewhat paroxysmal character from the accompanying tuberculization of the bronchial glands. One of the most marked peculiarities of the cough in the phthisis of children is the entire absence of expectoration, since the secretions are either retained in the bronchial tubes, or, if raised into the pharynx, are swallowed without any effort at expulsion.

Hæmoptysis very rarely occurs in the early stage or during the progress of the disease; and when it occurs as the cause of sudden death, is due to the complication with bronchial phthisis, rather than to the rupture of a bloodvessel in a pulmonary vomica.

The *temperature* of the body is, as a rule, higher than normal, although it presents fluctuations on different days, and even at different hours of the same day; at times being normal, and again rising as high as 102°, or more. The greatest elevation of the temperature is generally noticed at night, and is usually accompanied by flushing of one or both cheeks; but it is rare to find the colliquative night-sweats which prove so exhausting to adults.

The *pulse* is always accelerated, and becomes very frequent as the temperature rises.

The *appetite* is capricious, the tongue furred, and the digestion imperfect; the bowels alternate from a state of constipation to diarrhœa, and the stools are unhealthy in appearance, being generally putty-like or clay-colored. Naturally, with this disturbed state of the primæ viæ, nutrition is seriously impaired, and the child steadily loses flesh and strength. Indeed, in very many cases, the little patient presents merely the symptoms of impaired nutrition, becomes languid and drooping, and loses appetite, strength and flesh, for many weeks before the development of cough reveals the lungs as the seat of the disease.

Physical Signs.—We have already remarked the fact that in the investigation of phthisis, the physical signs are of much less value in the case of children than in adults. This arises not only from peculiarities of the physical and moral organization in childhood, but also from the mode in which the tuberculous deposit takes place. Thus, as a rule, the deposit of tubercle in the lungs of children is more generally diffused and uniform; so that we lose to a great degree the advantages derived in adults from a comparison of the results of auscultation and percussion in one part, with

those obtained in other. For the same reason, we are also deprived of those signs which, in the adult, are developed in a single point : as, for instance, the *coarse breathing*, which is of so much diagnostic importance as one of the earliest signs of the deposit of tubercle at the apex of the lungs.

Another source of difficulty and error lies in the fact already alluded to, that the bronchial glands, when enlarged by the deposit of tubercle, as so constantly happens in conjunction with the pulmonary phthisis of children, are brought into contact with the thoracic walls, and transmit many sounds with intensified force. It is thus that prolongation of the expiratory sound beneath the clavicle, and jerking respiration, lose much of the importance they have as signs of the early stage of phthisis in adults. For although, when heard in children, they should always be regarded as probable evidence of phthisis, they have frequently been noticed in cases whose progress shows the tubercular deposit to have been, at the most, trifling. In the same way, caution must be used not to mistake the blowing sound, mixed with moist râles, which is thus transmitted from a compressed bronchial tube containing mucus, for a large tubercular vomica. The only way in which this mistake, and the consequent too unfavorable prognosis, can be avoided, is by comparing daily the results of auscultation and percussion, and noticing whether they remain exactly the same, or whether, while the dulness on percussion over the enlarged glands persists, the results of auscultation vary from day to day with the varying amount of compression of the bronchus and the nature of its contents.

A still further source of difficulty results from the loss of all the information which is derived, in older persons, from the vocal resonance and its alterations ; although it is occasionally possible to draw reliable conclusions from the resonance and fremitus of the cry or cough. And again, owing to the excitability of children, patient and prolonged observation is required, both as to the situation, degree, extent, and duration of any inequality of breathing, before any conclusion can be drawn from it.

Finally, the extreme resonance of the thorax in early life tends to vitiate the results of percussion by preventing the recognition of fine variations of sonority, such as are readily detected in more advanced life.

We have thus far been considering the symptoms of pulmonary phthisis in its usual moderately acute form, but it is necessary to be aware that in some cases it deviates from this course, being at one time extremely rapid, and at another very chronic in its progress.

In the acute cases, we often find that there has been a previous deposit of tubercle or of caseous inflammatory exudation in different parts of the economy, though to so small an extent as scarcely to have interfered with nutrition or the performance of the functions, or to have attracted the least attention.

In such a state of the system, death may be produced in a few days or weeks by an acute development of tubercle. When this occurs in the lungs, it is not unfrequently attended by inflammation of the pulmonary parenchyma, constituting tuberculous pneumonia : and, whatever may be the view entertained as to the relation between the inflammation and the tubercular deposit, the recognition of this latter element is of the greatest importance, from its bearing on the treatment to be adopted.

In *tuberculous pneumonia*, in addition to the hereditary tendency and past history of the child, we rarely find the same heat of skin or vascular excitement as in pure pneumonia. The degree of oppression of the chest is also, from the beginning, out of proportion to the catarrhal or bronchial symptoms with which the case sets in. And auscultation reveals both that the amount of inflamed lung tissue is not sufficient to account for the dyspnoea, and that the râles developed are of the subcrepitant and mucous varieties, rather than the true fine crepitant râle of uncomplicated pneumonia.

In the *chronic form* of phthisis alluded to, the symptoms may be prolonged during several years. They consist of progressive emaciation, chronic cough, with or without expectoration according to the age of the patient, and the physical signs of more or less advanced tubercular deposit. In favorable cases, it is not unusual for some degree of temporary improvement to occur in the general symptoms, and in some rare cases the child slowly regains good health, and the physical signs gradually diminish, leaving merely some dulness and feeble respiration at points where positive signs of advanced pulmonary disease previously existed. It is needless to add that in such cases the nature of the morbid condition present has probably been of a chronic inflammatory rather than of a truly tuberculous character,

Symptoms of Tuberculous Peritonitis.—The peritoneum may either become implicated late in the course of general tuberculosis, or it may be the first structure involved. Apart, however, from the general symptoms of the tuberculous cachexia which in some cases precede its appearance, there are few symptoms of much diagnostic value during its early stage. Thus the child retains its appetite and spirit; does not lose flesh rapidly; and only complains of occasional and apparently causeless abdominal pain. This condition does not, however, last long; the nutrition soon fails, the appetite becomes capricious, the bowels irregular, the colicky pains more frequent and severe, and the abdomen acquires an abnormal size and appearance. These symptoms, however, merit a more detailed allusion. The tongue rarely indicates, either by dryness or furring, any serious disturbance of the digestive functions. The bowels are almost invariably loose, or alternations of constipation and diarrhoea present themselves, the stools usually being unhealthy in appearance. This condition frequently appears to depend upon inflammation or tuberculous ulceration of the intestines. Vomiting is not usually present; it is rarely spontaneous, and merely consists in the occasional rejection of alimentary matters.

Pain in the abdomen has been mentioned as one of the earliest symptoms. It is rarely constant or confined to the seat of the lesion, but is rather shifting, intermittent, and colicky in its nature, recurring with greater or less frequency. There is also tenderness on pressure over the abdomen, which becomes especially marked during the latter stages of the disease, though in some cases the abdomen remains indolent throughout. At a variable period after the appearance of the preceding symptoms, and sometimes simultaneously with the occurrence of colicky pains, the abdomen undergoes a marked modification in its size and shape. It becomes

tense and large, and assumes an oval or globular form, the depressions and fossæ being all effaced. It generally retains its tympanitic note upon percussion; and in proportion as the distension increases, the sound may become more and more tympanitic. It is not, however, rare to note that careful and gentle percussion gives irregular areas of impaired resonance, due to the presence of layers of lymph coating the visceral or parietal peritoneum at those points. The tension often varies without any apparent cause; and when it is much diminished, an imperfect sense of fluctuation may be obtained by flapping the sides of the abdomen. This sign is rarely due to any ascites being present, but is beyond doubt rightly explained by Rilliet and Barthez as due to the transmission of the impulse of the hand by the agglutinated intestinal mass.

It is only in very exceptional cases that even the most careful percussion or palpation will detect any inequalities in the abdominal walls, due to the presence of large tuberculous patches. In every case in which the last-named observers detected any abdominal tumor, the omentum was found to be the chief seat of the tuberculous deposit.

After this condition of the abdomen has persisted some time, the distended skin desquamates, and assumes a rough and dirty appearance; and the cutaneous veins of the abdomen become prominent and dilated, owing to the obstruction to the abdominal circulation. Deep inspirations are apt to cause pain owing to the descent of the diaphragm, and the breathing becomes shallow and thoracic in type from this cause as well as from the distension of the abdomen.

As the case progresses, and the general symptoms assume more gravity, these local symptoms become more pronounced. Like all forms of tuberculosis in children, however, the advance of this disease is rarely uniform, and intermissions and fluctuations in the symptoms are often noticed. Toward the close of life, all the symptoms usually undergo aggravation, and the remissions become more and more rare and brief.

Death is either produced by the advance of tuberculous disease in the lungs, or by tubercular meningitis; or the little patient sinks from sheer exhaustion under the persistent diarrhœa and the repeated accessions of peritoneal disease.

Symptoms of Tuberculosis of the Mesenteric Glands.—The symptoms of this condition are even less positive and diagnostic than those of tuberculous peritonitis. So long as the glands remain only moderately enlarged, buried as they are beneath the small intestine, it is impossible to detect their presence, especially as the absorption of chyle may not be materially interfered with.

We have already mentioned, moreover, the comparative rarity of symptoms due to the pressure of the enlarged glands upon neighboring structures, such as perforation or compression of the intestines, and dilatation of the cutaneous veins, or œdema.

The modifications of the size and shape of the abdomen occasionally furnish useful information. It is rarely so large and tense as in tuberculous peritonitis, and its shape is rather globular than oval.

There is scarcely any tenderness on pressure over the abdomen, unless there is some accompanying peritonitis.

The only really pathognomonic symptom, indeed, is the detection of the enlarged glands by palpation. This, however, is far from being possible in all cases, even when the bulk of the glands is very considerable, as they are frequently covered and concealed by the intestines.

It is, in fact, only in those cases where the abdomen is supple and relaxed, that we can establish the presence of the tumor, which is usually lobulated, varying in size from a hen's egg to a large orange, and seated in the neighborhood of the umbilicus.

The digestive system here also presents more or less marked disturbances; the bowels in particular being loose, a condition generally due to the existence of tuberculous ulceration of the intestine. It is probable that in most cases the disease of the intestinal mucous membrane is primary, and leads to irritative hyperplasia of the mesenteric glands, which ultimately undergo caseation and become the seat of tuberculous formation.

The general symptoms which accompany tuberculization of the mesenteric glands alone, are often not so marked as when other organs are affected; in fact, MM. Rilliet and Barthez assert that they have not met with a case in which this affection, isolated from all others, has produced any considerable emaciation. This does not, however, correspond with our own observation, since we have met with cases where the interference with general health and the attendant emaciation were very marked.

DURATION.—The duration of tuberculosis in children, as might have been expected, varies considerably according to the position and surroundings of the patients. In large hospitals, where the children have not the advantage of the best hygienic influences, the majority of cases terminate in from 3 to 7 months, though occasionally protracted to upwards of 2 years. In private practice, on the other hand, many cases of chronic phthisis are met with, in which the disease continues 3, 4, or even 5 years before producing death. It is extremely difficult to assign any probable duration for either tuberculous peritonitis or tabes mesenterica, as they can rarely be diagnosed during the early stages of their development.

DIAGNOSIS.—The danger in regard to the diagnosis of phthisis in children is not so much of entirely overlooking the nature of the disease, as of over-estimating its amount. We have already given the reasons why the physical signs of pulmonary and bronchial tuberculosis in children are less reliable and more difficult to appreciate than in adults. A proper attention to the hereditary tendencies and individual history of the child; a close scrutiny of its physical conformation and development, with an intelligent interpretation of the physical signs, will, however, generally suffice to prevent any serious error.

In the earlier stages of the more acute forms of phthisis, the disease with which it is most apt to be confounded is remittent fever; from which it may be distinguished by the history of malarial exposure, by the definite commencement of the case, and by the very marked exacerbations which

occur towards night, attended with high fever, great heat of skin, and considerable delirium. In its more chronic forms, the diagnosis of pulmonary phthisis from chronic bronchitis is often attended with the greatest difficulty. In fact, the physical signs of the two conditions are frequently so entirely analogous, that it is only by the general symptoms of tuberculosis, the greater amount of hectic irritation, the more rapid emaciation, and the frequent supervention of tubercular deposit in other organs, that a diagnosis can be established.

We have already dwelt upon the value of abnormal development of the abdomen as a symptom of tubercular peritonitis. There are, however, many cases of simple functional derangement of the intestines, in which no suspicion of tuberculous deposit can be entertained, where this symptom is also noticed. It is due to this circumstance that tubercular disease of the peritoneum and mesenteric glands was formerly considered of such frequent occurrence. A careful regard, however, to the age of the patient (for simple distension of the abdomen occurs generally in infancy, whilst tuberculous peritonitis is most frequent after the age of 3 years), and to the effects of simple remedies, will usually remove any doubt.

Ascites from other causes than peritonitis is not very rare in young children; but may be distinguished by the distinct fluctuation on palpation, by the symmetrical arrangement of dulness on percussion which occupies the dependent parts of the abdomen and which is greatly influenced by changes in the position of the body. In peritonitis, on the other hand, fluctuation when present is rarely so distinct and general, and a gentle percussion will often reveal irregularly distributed areas of relative dulness, alternating with tympanitic resonance, which are only to a moderate extent influenced by changes in the position of the child's body. The character of the breathing should be carefully studied, since, in consequence of the distension of the abdomen and the pain caused by any downward pressure of the diaphragm, it assumes in the highest degree the thoracic type.

Whilst it is usually possible, by attention to the above conditions, to determine the existence of peritonitis in children, it must be borne in mind that the disease is not always of tuberculous nature, but that subacute idiopathic peritonitis occurs in children, and may terminate favorably under suitable treatment. We have met with some most interesting cases of this character, and Kersch of Prague (quoted in *London Medical Record*, October 16th, 1876) has published an instructive article on the subject.

The cases which are most apt to be confounded with *tabes mesenterica* are those in which abdominal tumors, due to some other cause, are present. Thus, in extensive tubercular deposit in the omentum, we may have, in addition to the general symptoms of tuberculosis, a well-defined tumor about the middle of the abdomen. The greater degree of tenderness of the abdomen, and the mobility in this case, may, however, serve to distinguish it. Again, it is not rare to find in cases of digestive derangement, where irregular action of the bowels with more or less pain may have been present, a distinct and only slightly movable tumor in the abdomen, due to the impaction of the intestine with hardened feces.

A careful consideration, however, of the position of these masses, which is generally in one or the other iliac fossa; their entire painlessness and doughy character upon palpation, and their complete disappearance after the administration of laxatives and enemata, will reveal their true nature.

PROGNOSIS.—The very name of tuberculosis has grown, with only too much reason, to be almost synonymous with impending, unavoidable death. And yet, while pulmonary phthisis shows the same fatal tendency in childhood as in adult life, the prognosis is somewhat less gloomy. For not only does well-directed treatment occasionally render the morbid deposits in the lungs in some cases of phthisis, whose existence has been proved by the symptoms of the incipient stage, inert and obsolescent; but in rare cases, where the deposit has advanced to softening and destruction of lung-tissue, a cure has been slowly effected by the evacuation of the softened tubercle and the gradual cicatrization of the cavity.

In tuberculization of the bronchial and mesenteric glands, moreover, numerous cases have been noticed where the glands have undergone complete calcification, and the progress of the disease has been arrested.

While, therefore, the prognosis must ever be grave and unfavorable, we must bear in mind the possibility of recovery when the hereditary tendency of the child is not too strongly pronounced, and the actual tuberculous deposit not extensive or rapidly progressing.

MODES OF DEATH.—Having thus spoken briefly of the prognosis, a few words will suffice to call attention to the various modes in which phthisis brings about a fatal issue in children.

In the majority of cases, death occurs from sheer exhaustion of the powers of life, from impaired nutrition and perverted functions. In a few instances of bronchial phthisis, death is suddenly caused by copious hemorrhage, owing to the perforation of one of the pulmonary blood-vessels.

The immediate cause of death is frequently found in an intercurrent attack of bronchitis, pneumonia, or peritonitis: while, in other cases, the cerebral symptoms which precede the fatal event, show that the membranes of the brain have become the seat of tuberculous deposit.

It is not unusual, moreover, whether the original seat of the tuberculous deposit have been in the abdomen or thorax, for marked abdominal symptoms to be developed towards the close of the case; the tuberculous ulceration of the intestines serving to maintain an uncontrollable and exhausting diarrhœa.

TREATMENT.—*Prophylactic.*—In children whose parents are tuberculous, and who in early life give evidence of delicate health, the prophylaxis becomes most important. The infant should be kept at the mother's breast up to the age of fifteen or eighteen months; but in case the mother be herself tuberculous, on no account should she be allowed to nurse the child, for whom a healthy wet-nurse should be immediately procured. By attention to this precaution, we have succeeded in raising children of tuberculous mothers who had suckled their previous children and had lost them all in early life from tuberculous disease.

As the child advances in age, every caution should be paid to its food

and clothing, to securing sufficient exercise in the open air, and free ventilation in its sleeping apartment. When the circumstances of the parents permit it, it is of the greatest consequence that the child should enjoy the benefits of a country life, in some healthy, invigorating atmosphere, for four or six months out of every year.

The child should further be guarded sedulously from the ailments incident to early life, and especially from hooping-cough and measles; and the slightest disturbance of either the respiratory or digestive functions should receive prompt and careful treatment; nor should we be tempted to discontinue these efforts, even if positive signs of tuberculous deposit appear; for the possibility of these deposits in childhood becoming latent or being evacuated, and the general health re-established, should never be lost sight of.

Curative.—Little need be said of the treatment of fully developed tuberculosis in children, since the same indications present themselves as in adults, and call for the same remedies. The most essential points in the treatment are attention to all hygienic conditions, careful regulation of the diet, and the administration of remedies calculated to improve nutrition and primary assimilation.

It is indeed impossible to over-estimate the importance of maintaining the appetite and powers of digestion; and if these show any sign of failing, we should resort to some of the bitter vegetable tonics, of which, perhaps, the combination of tincture of *nux vomica*, gtt. ij to v, with the compound tincture of gentian, ℥ xv to xxx, according to the age of the child, is most desirable. On the other hand, if we find reason to believe that any remedy we are administering disturbs the nutrition of the child, disgusts it, lessens its appetite, or rouses violent opposition at every dose, it should be instantly abandoned as producing the very effect we most desire to avoid.

The child should be strongly encouraged to take nourishing food at regular intervals, and so soon as any of the articles of its diet become unattractive, other preparations of similar nature should be substituted. Milk should enter largely into the diet, and ought to be taken at least every morning and evening. Tender, finely divided meat should be eaten at the midday meal, in such quantities as the digestion will easily bear. If marked signs of debility present themselves, a few drachms of good brandy may be taken at intervals through the day, with advantage.

When the stomach does not reject it, there are few remedies whose action is more beneficial than cod-liver oil, given in the dose of a teaspoonful or even less, three times a day. In many instances, children soon become accustomed to the taste of this substance, and even grow to relish it almost as a luxury, and to take it eagerly; in some cases, however, the taste is so unpleasant that the children refuse to take it, and it is, therefore, advisable in such instances to prescribe it in the combination which we have already recommended, at least during the first few weeks of its administration.

In those cases where it is impossible to administer cod-liver oil internally, very good results may often be obtained here, as well as under

similar circumstances in other wasting diseases in children, by the use of the oil by inunction.

Iron and its various preparations are strongly indicated, and we can generally find some of the milder forms which will be readily tolerated.

In those cases where there is considerable implication of the lymphatic glands, the syrup of the iodide of iron appears especially useful, and this may be well given alternately or in conjunction with iodide of potassium.

Sea-bathing is strongly recommended, especially in the tuberculization of the glandular system; or when this is not attainable, baths in which some tonic drug has been mixed may be used.

In tuberculous deposit in the peritoneum or mesenteric glands, the diet must be regulated with peculiar care; the most bland, unirritating, and digestible food being selected. If, however, despite our precautions, diarrhoea should make its appearance, the various astringents in combination with opium should be given freely. The pain in the abdomen, which is frequently so severe in these forms of tuberculosis, may be relieved by the application of sinapisms, or of warm anodyne poultices, or by gentle friction with a sedative liniment.

When the symptoms of any intercurrent inflammation in the diseased organ present themselves, we must limit our treatment to the application of a few cups or leeches over the part, and the administration of a less stimulating diet, with some mild febrifuge. When the peritoneum is involved in the tubercular deposit, and we have reason to fear an accession of inflammation of that membrane, there is urgent necessity for the use of topical depletion in moderation; but we must, at the same time, bear in mind the cachectic nature of the disease, and refrain from the adoption of any depressing plan of treatment.

ARTICLE IV.

RICKETS.

DEFINITION; SYNONYMS; FREQUENCY.—Rickets is a constitutional disease peculiar to childhood, which first manifests itself by various disturbances of nutrition, and later by a specific alteration in the bones.

The disease has been known under a vast variety of names in many different languages;¹ almost the only terms by which it is designated by English or American authors, however, are rickets and rachitis.

An idea of the vast importance and frequency of this disease may be gained from the statements of some of the recent writers upon this subject. Thus Sir W. Jenner, whose lectures upon this subject² present a most original, philosophical, and lifelike description of the disease, speaks

¹ For Synonym, see Art. Rickets in Reynolds's Syst. of Med., vol. i, p. 768.

² Med. Times and Gazette, 1860.

of it as "without question the most common, the most important, and in its effects the most fatal of the diseases which exclusively affect children." Hillier, at the close of an excellent chapter upon rickets (*op. cit.*), presents a table showing the proportion borne by the number of cases of this disease to the total number of out-patients treated at the Hospital for Sick Children, London, from which we calculate that of 128,656 children treated during thirteen years (1854-66), not less than 8419, or 6.5 per cent., were rachitic; and in some years the proportion of such patients rose as high as 9 per cent.

The statistics furnished by other English writers, as Gee (*loc. cit.*), Merei, and Ritchie, support the view that in all classes of English society a notable proportion of the children are rachitic. In the same way the highest German authorities, as Ritter von Rittershain and Henoch, state that the proportion of the children treated at public institutions in that country, who are found to be rachitic, is not less than 30 per cent.

Of late years the attention of observers in this country has been more forcibly attracted to this subject, and, as a consequence, the number of cases in which the early and less prominent symptoms of rickets are now recognized is rapidly increasing. In a paper on this subject by Parry,¹ which we regard as the most valuable contribution to the literature of rachitis which has been made on this side of the Atlantic, the writer states that he has been "irresistibly forced to the conclusion that rachitis is scarcely less frequent in Philadelphia than it is in the large cities of Great Britain and the continent of Europe." We must add that, although, judging from our own experience, the above statement is an over-estimate, the number of cases in which we meet with the early, or even the more grave symptoms of rickets, is quite large both in private practice and in connection with public institutions.

The fact that during the past twelve years the mortality returns of this city contain but two deaths reported as from rickets, is of little importance, since so rarely is it assigned as a cause of death even in Great Britain, that the Registrar-General has not found it necessary to devote a column of his tables of mortality to the disease. "The secondary diseases," as Hillier says, "are recognized, such as bronchitis, collapse of the lungs, atrophy, measles, whooping-cough, or convulsions, but the primary disease, which renders these secondary diseases fatal, is ignored."

We shall limit ourselves to an account of the causes, general symptoms, and treatment of the disease, with a brief description of the anatomical changes in the bones, and the deformities which result, referring the reader who desires more minute knowledge on these latter points, to any of the elaborate memoirs published on this disease.

CAUSES.—*Age.*—Rickets is essentially a disease of childhood, and indeed may make its appearance during early infancy. There are also a few cases on record which show that it may, although rarely, occur in the fœtus before birth. It frequently may be detected during the first six months of extra-uterine life. Gee has noticed positive beading of the ribs at the third and fourth weeks, and Parry at the sixth week after birth.

¹ Amer. Journ. Med. Sciences, Jan. 1872, p. 17.

The age at which it ceases to be frequent for rachitis to begin is variously estimated. We have observed a number of cases where the earliest symptoms were detected during the second year; and we should be inclined to assign as the limits of its most frequent occurrence the second or third month to the close of the second year. It grows rarer after this latter date, and many high authorities unite in saying that it never comes on after the completion of the first dentition in a child hitherto perfectly healthy. Considerable difference of opinion exists upon the question whether rickets is hereditary or not; but there seems no evidence to show that it ever is so, in the sense, for example, in which infantile syphilis is hereditary. There can, however, be no doubt as to the great influence exercised by the health of the parents upon the development of the disease.

It is stated by some authors that too early marriages, or marriages between relatives, and chronic tuberculosis or constitutional syphilis of the father, predispose to it. These causes are, however, of doubtful power; and certainly are inoperative as compared with the very positive influence exercised by the condition of the mother. Thus, it is well ascertained, that whatever tends to induce debility and anæmia in the mother, as too frequent pregnancies or prolonged lactation, renders it probable that her next born children will be rickety. Thus, Jenner states that it is very uncommon for the first, or the two or three first born children, to be free from any sign of rickets, and yet for every subsequent child to be rickety; which he explains by the fact, "that among the poor the parents are generally worse fed, worse clothed, and worse lodged, the larger the number of their children; and among the rich and poor alike, the larger the number of children, the more has the mother's constitutional strength been taxed, and the more likely is she to have lost in general power." (*Loc. cit.*)

In addition to the tendency derived from the mother, there are numerous causes acting directly upon the child, which strongly predispose to the disease. These will be found to be nearly the same as those which favor the development of tuberculosis. Thus, premature weaning, and the substitution of improper food for the mother's milk; or, on the other hand, the continuance of suckling long after the proper period for weaning, and after the mother's milk has deteriorated in quality and become insufficient and unwholesome; or the use of indigestible, or of poor, scanty, and innutritious food at any period during early childhood, are all potent causes of rickets. So, too, many of the acute and chronic diseases of children, which impair assimilation and nutrition, as entero-colitis; and all such depressing influences as impure water, foul air, poor ventilation, small, damp, and dirty habitations, may be classed among the predisposing causes.

The marked alterations and deformities of the bones, which are so characteristic of rickets, are not developed until after a more or less marked cachectic state of system has persisted for a time varying from a few weeks to several months.

During this initiatory stage, the most marked symptoms are connected with the digestive system. The appetite may remain good or grow capricious; and the bowels are irregular, though for the most part of the time there is diarrhœa, with stools which are at first greenish and mucous, subsequently serous, watery, of a brownish or slate color, and horribly offen-

sive. If this chronic intestinal catarrh be but slightly marked, the child may retain a good deal of its fat, though frequently there is extreme emaciation.

The head is frequently bathed in profuse perspiration, which occurs especially during sleep, but also after any exertion, or even while the child is lying quiet. The skin of the trunk and extremities is hot and dry, and even the lightest covering seems oppressive to the little patient; so that there is a tendency to get rid of all the bed-clothing at night.

Another symptom which makes its appearance in a certain proportion of cases, but not so constantly as the digestive disturbances, local sweatings, and restlessness at night, is general soreness and tenderness of the body, with pain on movement; when this is marked, the child dreads to be moved or even touched, cries if its limbs be pressed firmly, and will lie almost motionless for hours. According to Parry (*loc. cit.*), this symptom is associated with the commencing bone changes, so that it properly belongs to the early part of the second stage.

If the disease begins before the completion of primary dentition, the development of the teeth is always impeded, and they are not only cut late, but either decay or fall very early from their sockets. The urine does not present any constant alteration, but in a certain proportion of cases the amount is increased, and there is an excess of the phosphatic salts, while in other instances excess of some free acid, said to be usually lactic, has been detected. The mental condition in rickets has been variously described; some authors regarding the intelligence as precocious, owing probably to the isolation of the patient from other children, and his constant association with his elders; while others assert that there is an actual deficiency in intellectual capacity and power. At a somewhat later period of the disease, the child acquires a peculiar staid and sedate aspect, which, when associated with the unusual breadth and squareness of the face, imparts a strange expression of age.

According to Roger and Rilliet, a blowing murmur may frequently be heard over the anterior fontanelle in this disease, synchronous with the arterial pulse. As, however, this murmur is to be heard in other conditions, and is often absent in cases of rickets, it cannot be considered as a sign of any positive value. The causes which appear to intensify it, are the anæmic state of the blood and the patency of the anterior fontanelle; yet Hillier states that he has found it present in thirteen, and absent in twenty-nine rickety children whose fontanelles were open. According to Jurasz, this murmur originates either in the carotid canal or the foramen spinosum, and is without diagnostic value. He never found it prior to the third month or after the sixth year; but between these ages found it in twenty-eight out of sixty-eight cases, though not constant nor always in the same place.

The phenomena above described, when present in the same case, may certainly be regarded as positively indicative of the existence of this initiatory stage of rickets, but they are by no means invariably all present, so that it is often impossible to determine the approach of the next stage in which the characteristic lesions and deformities of the bones make their appearance.

They are not, moreover, limited to the stage of invasion, but continue, with more or less severity, for a varying time after the bone changes have begun. The length of this stage of invasion is exceedingly irregular, and the earliest physical sign of bone change may occur after it has lasted a few weeks, or may be deferred for several months after the peculiar prodromic symptoms have been marked.

Stage of Deformity.—After the initiatory stage has lasted for a varying time, bead-like swellings begin to be noticed at the line of junction of the ribs and costal cartilages, which is usually regarded as the earliest lesion of the bones, and of the epiphyses and shafts of the long bones of the upper and lower extremities, giving in these latter places, as at the ankles and wrists, a peculiar knobby double-jointed appearance. With this, there is such a degree of softening of the bones, that they yield readily to pressure.

Early in this stage the presence of craniotabes, or “soft spots” in the occipital bone, may often be detected. Indeed, in some instances this appears to be the first recognizable bone lesion.

If the disease reaches this stage before the child has begun to walk, there may be no deformity of the lower extremities whatever; but in cases where the little patient has already been walking about, the femora bend so that they become markedly convex forwards; the tibiae bend in the same forward direction, while the knees may be bent inwards, thus giving to the legs a series of curvatures. The forward curvature of the femora may indeed be produced before the child walks, simply by the weight of the legs and feet, which hang pendant from the knee-joints as the child sits in its mother’s lap or on a chair.

The bones of the upper extremities also share in these deformities; thus the humeri bend at the point of insertion of the deltoids, from the weight of the arms when raised by the action of these muscles; and both the humeri and the bones of the forearms become bent, from the pressure which the child makes on its open palms to assist itself in sitting up.

The clavicles are very constantly deformed, and present a double curvature; one curve being forwards and somewhat upwards, and seated just outside of the attachment of the sterno-cleido-mastoid muscle, the other being backwards, and seated about half an inch from the acromio-clavicular articulation.

By far the most important deformities, however, are those presented by the head, spine, thorax, and pelvis. The peculiarities by which the head in rickets is distinguished are thus described by Jenner:

1st. By the length of time the anterior fontanelle remains open. In the healthy child, it closes completely before the expiration of the second year. In the rickety child, it is often widely open at that period.

2d. By thickening of the bones. This is usually most perceptible just outside the sutures, the situation of the sutures being indicated by deep furrows.

3d. By the relative length of the antero-posterior diameter of the head.

4th. By the height, squareness, and projection of the forehead. The first two of these peculiarities of the rickety head are the result of the

affection of the bones; the last two are due chiefly to disease of the cerebrum.

Besides this thickening of the edges of the cranial bones, there are spots, irregularly distributed, where the bones are so thinned and softened that they yield to the pressure of the fingers; and, indeed, in some cases the thinning is so extreme that the pericranium and dura mater come in contact. These "soft spots,"¹ which constitute the condition known as *craniotabes*, were first observed by Elsässer.¹

The nature and mode of their production has been a matter of much discussion. By some authorities their rachitic nature has been denied, but there seems to us no valid reason for doubting their essential connection with the rachitic alterations of the bones. They are usually limited to the occipital region, but may rarely be present over the other cranial bones.

They are never observable save in those parts of the bones which are developed from membrane. At first, the spots affected are the seat merely of softening, with perhaps some thickening; then thinning of the bone occurs, and subsequently the entire thickness of the occipital bone is often removed, causing perforations. These vary in number from one or two to as many as twenty-five or thirty. In order to detect them, the skull should be carefully examined by fixing the head between the hands, and then pressing carefully over the upper part of the occipital region and the posterior portions of the parietal bones. The diseased spots are felt to be soft and easily depressed, and "impart the sensation of an orifice in the bone, closed by parchment." It is necessary to use much caution and gentleness in making this examination, since any undue pressure may produce severe nervous symptoms, even convulsions, according to Niemeyer. It is difficult to account for the production of these spots, but the most probable explanation is that they are dependent upon the prolonged pressure upon the softening bone, caused by the head resting on the pillow on one side, and by the counter-pressure of the brain on the skull on the other.

The curvature of the spine varies according as the child is able or unable to walk. In the latter case, there is a posterior curvature of the spine, beginning at the first dorsal, and extending to the last lumbar vertebra; while if the child is able to walk, this posterior curvature is limited to the dorsal region, but is combined with an anterior curvature in the lumbar region. The cervical anterior curve is increased, and consequently the face is directed upwards, and the head falls backwards, and being unsupported, owing to the muscular debility, sways loosely from side to side. Jenner points out that these curvatures may readily be distinguished from angular curvature, by the fact that the weight of the legs will usually remove them if the child be held by the upper part of the trunk, especially if the physician at the same time raises the lower limbs with one hand, and places the other on the curved spine.

The thorax is subject to deformities, which in a practical sense exceed all others in importance, owing to the serious interference which they occasion with the action of the heart and lungs.

In the first place, owing to the curvature of the spine, the ribs are flat-

¹ Der weiche Hinterkopf, Stuttgart, 1843.

tened laterally, and run forwards more horizontally, so that the lateral diameter of the chest is greatly diminished, while the sternum is carried forwards, and thus the antero-posterior diameter of the thorax is increased. In addition, there is a marked groove on either side of the sternum, extending from the first to the ninth or tenth ribs, along the line of junction of the ribs with their cartilages. These grooves are produced by the bending of the ribs where the dorsal and lateral portions unite; from which point they pass forwards and inwards to unite with their cartilages, which curve outwards before uniting with the sternum.

The curvatures and deformities which have been described before this are chiefly due to the action of muscles or the weight of dependent parts; but the production of the last-described deformities of the thorax is attributed by Jenner chiefly to the atmospheric pressure, which, during inspiration, causes recession of the most yielding part of the thoracic walls, *i. e.*, the softened ribs at the line of junction with their cartilage. In consequence of the support which the liver, heart, and spleen furnish to the ribs corresponding to their position, the groove extends further down on the left than on the right side, but is deeper over the fifth and sixth ribs on the right than on the left side.

The pelvis is frequently affected in rickets, and the deformities which result, on account of the great interference they cause in childbirth in the female, rank next in importance to those of the thorax. The rickety pelvis is characterized by a shortening of the antero-posterior diameter, so that the upper strait assumes an oval form, or is at times heart-shaped. In extreme instances the sides also approximate, and give to the pelvis a triangular shape. It is evident that the form will be influenced by a number of conditions; as the stage of ossification, and the direction in which the pelvis is compressed by the spine from above, and the thigh-bones from below.

Partly in consequence of the diminished capacity of the thorax and pelvis, partly in consequence of the weakness of the abdominal muscles, the flatulent distension of the intestines, and the enlargement of the liver and spleen which are frequently present, the abdomen is unusually prominent in rickety children.

During the development of the alterations in the bones the general symptoms before described persist; the digestion is enfeebled, and the stools liquid and fetid; the emaciation and debility increase; the respiration is more or less embarrassed by the deformities of the thorax; the pulse is quick, small, and irritable; the skin hot, excepting on the head and neck, where it is still frequently bathed in sweat; and the general tenderness of the body is aggravated.

In cases where the disease approaches a favorable termination, the earliest signs of improvement consist in a decrease in the emaciation, debility, and suffering; the stools become more healthy, and the febrile symptoms, if any have been present, disappear.

During this stage of early convalescence, when the children attempt to leave the bed and walk about, holding on to the chairs, there is great danger of increased curvature and even of partial fractures of the bones of the lower extremities.

When, on the other hand, death occurs during the course of rickets, it is rarely from the intensity of the cachexia (which explains the apparent anomaly of so fatal a disease being scarcely represented in the mortality returns), but from the supervention of some secondary disease. Among these, the following are enumerated by Jenner as the most frequent causes of death:

1. Catarrh and bronchitis, which are rendered far more dangerous from the mechanical interference with respiration caused by the deformed thorax.

2. Albuminoid (?) infiltration of various organs, especially of the liver, spleen, and lymphatic glands. As will be seen by the remarks in the section on morbid anatomy, recent researches make it probable that the enlargement of these organs in rickets differs from ordinary albuminoid change. This peculiar form of degeneration is not unfrequently developed during the course of rickets; it manifests itself by increased emaciation, extreme pallor, occasional œdema and albuminuria, and enlargement of the affected organs.

3. Laryngismus stridulus, which, according to Jenner, is essentially connected rather with the nervous irritability due to rickets than with the tardy and difficult dentition which is itself but another expression of the constitutional disease.

4. Chronic hydrocephalus.

5. Convulsions, depending, like the laryngismus stridulus, upon the heightened irritability of the nervous system.

6. Persistent and severe diarrhœa, which is probably due in many cases to albuminoid degeneration of the intestinal mucous membrane.

DURATION; PROGNOSIS.—The duration of rickets varies so greatly, that the disease may be said to present an acute and chronic form.

When the diathesis is marked, the hygienic conditions of the child very unfavorable, and the disease makes its appearance at an early age, its course is often very rapid, and death usually follows. When, on the other hand, the disease does not begin till late in the second or third year, and when the surroundings of the child are more favorable, recovery usually occurs, although the disease may last for several years.

An unfavorable prognosis may be made, then, when the disease begins in very early infancy; when it is attended with marked constitutional disturbances; when the deformities of the head and thorax are rapidly and extremely developed; when any of the secondary morbid conditions above enumerated have supervened. When, on the other hand, the reverse of these conditions obtains, recovery may be expected, though often only after prolonged illness.

DIAGNOSIS.—It is only during the initiatory stage of rickets, that the true nature of the attack is likely to be mistaken. But during this period the disease may be confounded either with chronic entero-colitis, or with tuberculosis of the peritoneum and intestinal canal. Careful attention to the peculiar symptoms of rickets, especially the sweating of the head, the general soreness and tenderness of the body, and the retardation of dentition, will, however, lead to a correct diagnosis, even before the swelling of

the sternal ends of the ribs and of the epiphysial lines of the long bones, and the projection of the sternum, remove all doubt as to the nature of the case.

MORBID ANATOMY.—The essential lesions in rickets consist of the changes in the bones, though there are also certain lesions of the viscera which are frequently met with.

The long bones affected by rickets, in addition to the deformities already described, are clumsy, and present marked swellings at the line of their junction with the epiphyses. This enlargement is due to excessive development of the spongy tissue in the extremity of the bone and the epiphysis, and to marked proliferation of the epiphysial cartilage. The fact that the epiphyses widen instead of elongating, is due to the pressure of the superimposed parts upon the soft proliferating layers, causing them to bulge laterally.

The deposition of calcareous granular particles at the line of ossification is also wanting, and the cartilage cells calcify before the matrix begins to ossify, and are converted into bone cells.

There is thus excessive formation of the structures which precede or form the nidus for ossification, while there is at the same time retardation or incomplete performance of that process.

At the same time, the diaphyses present rarefaction of their tissue, not owing to undue softening and removal of old bone, but simply to the fact that, while the old layers of bone are consumed by the normally progressive formation of medullary cavities, the new layers which are produced are soft and do not ossify.

The medullary space may reach the line of ossification, or even project beyond it into the proliferating epiphysial cartilage.

The periosteum of rickety bones is usually thickened and highly vascular.

The bones themselves become so soft that they may be bent in any direction or even cut with a knife without difficulty.

Upon section the spongy tissue and the enlarged areolæ are found filled with a crimson pulp, containing blood globules, a large amount of free fat in some cases, and very many round, faintly granular cells, with one or two nuclei. According to Hillier and Parry the reaction of rachitic bones is alkaline or neutral.

The softening of the bones is fully accounted for by the diminution in the proportion of their calcareous salts. Thus Jenner states as the mean of the analyses of several observers, that the bones of healthy children yield about thirty-seven parts of organic and sixty-three of inorganic matters; whereas those of rickety children yield about seventy-nine parts of organic to twenty-one parts of inorganic matters. In addition to this, it would appear that the organic matters themselves undergo change, since it has been found by several experimenters that the bones in advanced rickets yield neither chondrin nor gelatin on boiling.

The thickening of the flat bone is caused by the formation of new osseous layers from the thickened and vascular periosteum, which are especially formed at or near the growing margins of the bones, thus accounting for the thickened ridges near the sutures of the cranial bones.

The thickening of the bones of the skull may reach a very high degree, a thickness of $\frac{7}{8}$ in. having been quite frequently observed. There are also frequently found on the skull evidences of craniotabes in the form of round or oval perforations of the bone, which have been observable during life as "soft spots." These perforations are most constant and frequent in the occipital bone, and are also found in the parietal bones, or wherever the skull has been subjected to pressure. They are surrounded by thickened bone, and are produced by the wasting and resorption, under the influence of pressure, of the young and unossified layers of sub-periosteal formation, while the absorption of the inner vitreous table, which keeps pace with the growth of the brain, proceeds at its usual rate. In number they vary from one to twenty or thirty.

In addition to these changes in the bones, which are the constant and essential lesions in rickets, there are certain lesions of the viscera frequently met with, which depend partly upon the deformities of the skeleton and partly upon the general cachexia. Thus in consequence of the peculiar deformity of the thorax, the anterior borders of the lungs become highly emphysematous, while the band of lung tissue corresponding to the deep groove at the sternal end of the ribs is compressed and collapsed.

This peculiar and constantly present strip of collapsed lung, is due to the recession of the corresponding part of the ribs during inspiration; but frequently there is also found extensive collapse of the postero-inferior parts of the lungs from the ordinary causes, bronchitis and impeded respiration. Jenner has also called attention to the frequent presence in rickets of white spots upon the pericardium, near the apex of the heart. These spots thus correspond to the depressed part of the fifth left rib, and are in all probability due to the friction of the heart against this hard knuckle of bone.

We have before alluded to the enlargement of the liver and spleen which appears in some severe cases of rickets. This was formerly regarded as due to albuminoid degeneration, but recent study of such organs has made it probable that the alteration is a peculiar and specific one.

The differences between this change and albuminoid (amyloid, of Virchow) degeneration were first pointed out by Jenner,¹ who showed that in the rickety enlargement, the organs present no reaction with iodine, and that in the spleen there is an absence of the peculiar sago-like transformation of the Malpighian corpuscles. Dr. W. H. Dickinson has more recently examined this subject with care, and has confirmed the view that the change in rickets differs both from albuminoid degeneration and from the peculiar enlargement of the spleen and lymphatic glands known as Hodgkin's disease. The *liver* in rickets undergoes an increase of size evenly throughout its whole bulk; it becomes pale, containing little blood, and is less friable than in health, hard, dense, and elastic. The acini are yellowish and are surrounded by a pinkish or grayish line, due to increase

¹ Medico-Chir. Trans., vol. lii, 1869, p. 359.

of the interlobular connective tissue. There is not, however, any bacony translucency as in albuminoid degeneration. The spleen is even more markedly enlarged than the liver, so that its weight may increase from one ounce to half a pound. The organ presents a resilient hardness which in extreme cases was compared by Bright to the consistence of a half-ripe apple. The color is generally a deep-red or purple, besprinkled with smooth white spots, which are enlarged Malpighian corpuscles. The trabeculæ are much thickened, and there may be also morbid hyperplasia of the cellular contents of the meshes, the corpuscles being much crowded together. The above change is described by Dickinson, as due not to the presence of any formation foreign to the structure of these organs, but to an irregularity of growth which alters the natural proportions of their tissues. The epithelial and corpuscular element is generally increased, while in the liver the capsule of Glisson, and in the spleen the trabecular tissue, is abnormally developed. There would appear also to be a deficiency of earthy salts in these organs.

In cases where death is directly due to any secondary disease, as bronchitis, intestinal catarrh, or chronic hydrocephalus, there will of course be found, in addition, the lesions common to such affections.

PATHOLOGY.—The description which has been given of the symptoms of rickets, clearly establishes the fact that it is a constitutional disease, in the same sense that scrofula and tuberculosis are; and we are consequently to regard the lesions of the bones as merely a local manifestation of the general cachexia. We are unable, however, to advance beyond this point, since we are ignorant, not only of the essential nature of the vice of nutrition, but equally so of the specific cause of the changes in the bones.

The result of chemical analysis has led to the theory that the disease essentially consists in a deficiency of the calcareous salts of the bones; and the attempt has been made to explain this deficiency by supposing an excess of lactic acid in the primæ viæ and blood, which holds the calcareous salts in solution, and prevents them from being deposited in the bones. Apart from the purely hypothetical nature of this supposition, and its entire inadequacy to explain many of the most serious symptoms of rickets, it is to be borne in mind that the excess of free acid in the urine is far from being constant, and that the changes in the bones are characterized not merely by a deficient deposit of the calcareous salts, but by their abnormal position, and by all the evidences of an active vital process.

Again, the marked vascularity of the bone and periosteum, the rapid proliferation of cells, and the pain and constitutional irritation which attend the disease, have induced others to regard the process as an inflammatory one. But this view is controverted as well by the etiology and clinical history of the disease as by its constant anatomical results.

We can only assume that, in consequence either of a special predisposition on the part of the tissues themselves, or of an abnormal quantity of some stimulus which normally excites the cartilage cells to undergo division, and the periosteum to form new layers of tissue opposed to the

surface of the bone, there is a morbid activity of these processes resulting in an excessive production of preparatory or intermediate structures, which can only become ossified in an imperfect, irregular, and slow manner.

The pathological process is thus seen to consist, for the most part, in a morbid acceleration of the changes which precede the normal formation and growth of bone.

TREATMENT.—In cases where there is reason to anticipate the development of rickets, as where the previous children of the mother have become rickety, the utmost attention must be paid to the feeding and hygiene of the young infant. If careful examination of the mother's milk proves that it is unsuitable in quality, a wet-nurse should be immediately provided, or if that be unattainable, the child should be fed upon carefully selected cow's milk, or upon one of the substitutes for human milk described in the article on food.

So, too, after the disease has made its appearance, the most appropriate, nutritious, and digestible diet must be selected, care being taken that it shall contain a large proportion of animal food.

The teeth of rickety children are so defective that, when they begin to take solid food, it is highly necessary to insure its complete mastication, and in cases where the condition of the teeth renders this impossible, the meat should be chopped finely and bruised in a mortar.

The child should be suitably and warmly dressed, and be taken freely into the sunlight and open air. The use of salt-water baths, followed by active friction of the skin, is also to be recommended.

During the early stage, when there is marked constitutional irritation and pain, the remedies used to relieve these symptoms should be alkaline mixtures, such as the effervescing draught or neutral mixture, or magnesia (Copland), conjoined with sedatives and tonics. Under no circumstances should any depressing plan of treatment be adopted.

If the digestion be much impaired and diarrhœa is present, the use of vegetable tonics, or wine of iron, with mild astringents and antacids, is indicated.

The remedy, however, from which most benefit is usually derived is cod-liver oil, and it should consequently be given, in conjunction with iron and vegetable tonics, and a small amount of some generous wine, so soon as the nature of the attack is recognized, and persevered with for months, or until the disease is overcome.

The efficacy of cod-liver oil in the treatment of this disease is, indeed, so remarkable that all other remedies formerly used have been supplanted by it. Vogel asserts (*op. cit.*, p. 534) that "rachitis may be cured by the use of cod-liver oil alone, even if the circumstances are in other respects unfavorable." Rickety children usually tolerate the oil well, and even become so fond of it that they will willingly take large doses. In some cases, however, it disagrees with the stomach and is obstinately refused by the children; and when this happens, so important is the introduction of the oil into the system, that we should recommend its use by inunction. It very rarely happens, however, that the difficulty in its administration cannot be overcome by having the oil prepared in the form of an emul-

sion, either according to the formula recommended on page 386, or in combination with the lacto-phosphate of lime.

There can be no doubt that when rickets is recognized in its early stages, and a suitable medicinal and hygienic treatment promptly instituted, it is usually curable in a comparatively short time. When, however, the diathesis is strong and the case overlooked until softening of the bones has occurred, and deformities begin to appear, the treatment must be persisted in for many months or even years. In such cases, unfortunately, there is only too great probability of the deformities increasing and becoming permanent, even if death does not ensue from some intercurrent or super-induced disease.

In order to guard against deformities, the little patient should lie upon a firm, smooth mattress, and high pillows should be forbidden. Niemeyer recommends that small children should be carried out in a basket; while larger ones should be drawn about in a carriage provided with a mattress. Sitting up for any length of time, or attempts at walking, should be prohibited until the bones have grown firm and inflexible.

It is not advisable, especially during the earlier stages of the disease, to employ any mechanical contrivances to prevent or relieve deformities. During convalescence, however, attempts may be made to control the deformities by means of leather or pasteboard splints.

In the treatment of any intercurrent affections it must be remembered that we have to do with a condition of malnutrition and enfeebled vitality, so that all remedies of a depressing character must be scrupulously avoided.

ARTICLE XIII.

CONGENITAL SYPHILIS.

INFANTILE syphilis may be either inherited or acquired subsequent to birth. As, however, the characters of the latter form do not differ materially from those of acquired syphilis in the adult, we shall limit our description to hereditary syphilis.

Careful clinical observation appears to have clearly demonstrated the following facts with regard to the transmission of syphilis, in addition to the direct contagiousness of both the primary and secondary manifestations :

That the embryo in utero may be infected, if either of the parents have constitutional syphilis at the period of conception, no matter whether the disease be latent, or if secondary or tertiary symptoms are present. That if both parents are syphilitic the child will more surely suffer from the disease, and in a more severe form. That if the mother, though healthy at the time of conception, contract syphilis during the first six or seven months of pregnancy, the child will probably be infected. That when the mother infects the embryo, the disease is probably more severe than when the

father alone is syphilitic, and thus such embryos usually perish, and are prematurely cast off by abortion, so that the great majority of children with congenital syphilis have inherited it from their father. While the last statement is almost universally admitted, there are some authors, as Hutchinson,¹ who do not admit the greater severity of the disease when the mother is the source of contagion. Finally, that a syphilitic father may infect the ovum without contaminating the mother's system, though the mother may subsequently herself be infected by the embryo.

In very many cases, though unfortunately not in all, the infected embryo perishes, and abortion follows. When, however, such infants are born living, they usually present no trace of syphilitic disease at birth, but may appear well nourished and healthy. Occasionally, however, children have been observed who presented, at the time of birth, copper-colored blotches upon the skin, condylomata, or mucous patches.

In the majority of cases the first symptoms of the disease appear between the fifteenth and thirtieth days after birth, though in many instances also during the second month. Thus of 158 cases collected from various sources by Diday,² the disease showed itself—

During 1st month in	86
" 2d "	45
" 3d "	15
At 4th month in	7
" 5th "	1
" 6th "	1
" 8th "	1
" 1 year,	1
" 2 years,	1

So that 131 children out of 158, or 83 per cent., presented evident symptoms of syphilis before the end of the second month.

Among the earliest evidences of the disease are the signs of failing nutrition. The infant, who has grown well, and has been plump and apparently vigorous for a few weeks, begins to emaciate, the features become pinched, the skin assumes a dry, sallow, shrivelled appearance, and presents patches of yellowish-brown discoloration, especially on the prominent parts of the face; the voice becomes feeble, whimpering, and plaintive, and the infant soon acquires a remarkable expression of premature old age.

The appearance of the skin has been most minutely described by Trousseau,³ West,⁴ Diday, and others, and is in a high degree characteristic of the disease.

In addition, however, to these general symptoms of malnutrition, there soon appear the signs of constitutional syphilis, familiarly met with in the adult, as well as some which are peculiar to the disease in infancy.

These symptoms now to be described belong partly to the secondary and

¹ Art. Constitutional Syphilis, in Reynolds's Syst. of Med., vol. i, pp. 297 and 315.

² Infantile Syphilis (Syd. Soc.), 1850.

³ Clin. Méd., 3ème ed., 1865, t. iii, p. 291.

⁴ Dis. of Children, (4th Am. ed.), 1866, p. 577.

partly to the tertiary stage, for it is a peculiarity of infantile syphilis that the evolution of the symptoms does not follow so orderly a course as in syphilis of the adult. The symptoms most frequently met with are certain affections of the skin and mucous membrane. The former may manifest themselves before the latter, simultaneous with them, or later; not rarely, however, the skin is the first tissue attacked. The cutaneous eruptions do not appear in any fixed order of succession, but are subject to marked variations. They may appear in a macular, papular, pustular, or bullous form, and thus produce roseola, erythema, mucous patches, acne, impetigo, ecthyma, and pemphigus. Certain of these eruptions manifest themselves sooner and occur more frequently than others; and some preserve their original form throughout, whilst others frequently combine. They may invade the whole surface of the skin, but generally have certain places of election, and a particular manner of grouping. Some of them differ considerably in appearance from those of acquired syphilis, and present features which distinguish them from non-specific eruptions. They are contagious; their color, in the majority of cases, is of a peculiar coppery hue, or yellowish-red, and varies in different stages; they are rarely attended with itching or smarting; they are annular in shape, and are prone to relapses.

Roseola is generally one of the first manifestations of constitutional syphilis, and is characterized by spots or patches of a bright or brownish-red color. The spots occur usually upon the abdomen, the inner surface of the thighs, or the lower part of the thorax. They are irregularly rounded, circumscribed, and vary in size from a finger-nail to the palm of the hand. They rarely disappear upon pressure, and finally fade away as dark-gray stains. Syphilitic roseola has occasionally been mistaken for simple roseola, measles, and scarlatina. This need not happen, however, if we remember that in scarlatina the eruption is most marked on the neck and upper part of the chest, is of an intense red color, disappears upon pressure, is punctiform, and has an accompanying angina that is more severe. In measles the eruption occupies the face and is crescentic in shape; besides there are coincident catarrhal phenomena, such as coryza and bronchitis, which are absent in syphilitic roseola. In simple roseola the eruption is of shorter duration, disappears upon pressure, itches, and does not leave behind it any of the dull-gray stains before mentioned. At times the syphilitic roseola is so extensive that the whole of the lower portion of the body is covered by a sheet of erythema. Again, the erythema may attack the palms of the hands and the soles of the feet, the skin peeling off in thin, dry flakes. Intertrigo and simple erythema are apt to be confounded with this eruption, but they appear in situations and forms, and from the influence of causes which distinguish them at once from a venereal exanthem.

Not infrequently the erythematous spots assume a papulated form, very slightly prominent, of the size of a finger-nail, and with a curved border. This maculo-papular eruption is considered by some authorities¹ as the most frequent syphiloderm occurring in the infant.

¹ D. Hring, *Treatise on Skin Diseases*, Philadelphia, 1881, p. 496.

Papules appear as both dry and moist lesions. The dry papule occurs less frequently than the moist kind, is broad and flat, with a glazed surface, and usually presents a superficial desquamation. At first it is of a red color, but afterwards assumes a tawny hue. It occurs most usually on the upper half of the body. The predominance of the moist papule, or "mucous patch," is to be found in the fact¹ that papules occupy by choice skin that is "thin, moist, and exposed to constant friction," and that just as soon as formed they are "macerated by the normal moisture of the part affected, which approaches the character of the mucous membranes."

They are slightly elevated, of varying size, have an ashy-white or diphtheritic color, and are covered with a thick glistening secretion. Mucous patches are almost invariably present in cases of congenital syphilis. They affect especially the region around the umbilicus, vulva, scrotum, anus, axilla, and corners of the mouth. Occasionally they appear between the toes and fingers, behind the ears, and about the alæ of the nose. Wherever moisture, warmth, and friction are present, there they most commonly are seated. When not treated they increase in extent, but do not become deeper.

Rhagades or fissures sometimes form in the skin, in the flexures of the joints, particularly in those of the fingers and toes, and may assume the character of moist papules. They also occur at the junction of the skin and mucous membranes, as on the lips, and at the verge of the anus. These rhagades bleed upon any stretching of the parts, and by their laceration so much pain is caused, that when the mouth is affected, the child dreads to smile, talk, or suckle; and when they are seated on the anus, defecation is attended with extreme suffering. Dry, scaly, or squamous eruptions are quite rare in infantile syphilis.

Syphilitic pustular eruptions, in new-born infants, may appear in the form of acne, impetigo, or ecthyma, and are peculiar in that they do not belong to any particular period of the disease. A form of acne, attended with the semblance of indurated pustules which leave little depressed cicatrices, is not infrequently met with. It is observed principally on the back, buttocks, shoulders, and chest.

Impetigo attacks the face as numerous pustules, which soon coalesce. These burst, and the pus drying the children are covered with an unsightly and horrible mask. The chest and neck may rarely be invaded, but the first eruption never oversteps its original situation. The diagnosis between syphilitic and simple impetigo is easily made out when we remember that in the latter there is commonly an eruption of like nature on the scalp, which is absent in the former, but has no ulceration under the crusts, and none of the many other symptoms that are present in syphilis.

Ecthyma seldom appears in the early stage of the disease. It occurs on the legs and buttocks as dark colored patches. These soon become converted into pustules in which the pus is mixed with blood. Subsequently there is ulceration and loss of substance. It is readily distinguished from ordinary ecthyma, which only occurs in adults and old people. Of the

¹ Diday. op. cit.

bullous form of eruption pemphigus is the most characteristic. It is also the first eruption to appear, not rarely being present at birth, and never, according to Niemeyer,¹ commencing later than the end of the first week. It usually appears first on the palms and soles of the feet, and may afterwards spread to various parts of the surface. It begins as small round spots, of reddish color, which become converted in a day or two into bullæ filled with turbid fluid. These burst, leaving irritable excoriations, and are succeeded by fresh crops of similar vesicles. The early appearance of pemphigus is of most fatal import, though in some cases recovery gradually occurs in the course of a few weeks.

Next to the cutaneous eruptions the affections of the mucous membranes are the most frequent. Thus coryza, of a serious and most obstinate form, is one of the most constant symptoms met with, and presents here all the characters fully described in our article upon that subject. The nasal mucous membrane is so much swollen that breathing and nursing are seriously interfered with. There is a profuse discharge from the nostrils, either of a thin, irritating fluid, which flows over the lip and excoriates it, or of a thicker pus, which tends to concrete and form thick, discolored crusts. The obstruction to respiration, and the accumulation of secretion in the nasal cavities, give rise to a peculiar snorting or snuffing quite characteristic of the disease.

There is apt to be, at the same time, a superficial diffuse inflammation of the mucous membrane of the mouth and throat, which may extend into the larynx, causing, in conjunction with the coryza, great alteration in the cry or voice, which is hoarse, and has been under such conditions compared by West to the sound of a child's penny trumpet.

Despite the severity and obstinacy of the coryza, there comparatively rarely occurs any ulceration of the mucous membrane, or necrosis of the nasal bones, or of the hard palate. In a few cases, however, we have observed depression of the bridge of the nose in consequence of the destruction of the nasal bones and perforations of the septum between the nostrils or of the hard palate, and West records a case in which there was necrosis of the hard palate in a young infant.

When mucous patches occur on mucous membranes they are seen most frequently in the mouth and at the anus. They may form in all parts of the mouth, but generally occupy either the furrow which unites the gums and lips, or the cheek, edges and tip of the tongue, soft palate, roof of the mouth, tonsils, and half arches. They present essentially the same features as those occurring on the skin, but are less prominent, and ulcerate much more rapidly. Condylomata are prominent mucous patches, which are either hard and warty, or fungus-like granulations, according as they occupy exposed surfaces or moist clefts. They are most frequent at the orifice of the mouth and anus, though they may also form elsewhere upon the skin. In consequence, probably, of the softening and ulceration of these growths, large, sinuous, irregular ulcers may form in such positions, extending for some distance into the surrounding skin.

¹ Textbook of Pract. Med. (Am. Trans.), 1869, vol. ii, p. 700.

Stomatitis and aphthæ may be confounded with mucous patches occurring in the mouth; but according to Duhring,¹ the ulcerations of stomatitis can be distinguished by their gray color, and by the redness and swelling of the surrounding parts. Aphthæ are seated upon an inflammatory base, are circular, isolated, and have distinct margins and areolæ. They also occur in successive crops, and are generally attended with derangement of the stomach.

In a few cases, iritis occurs; and so too the deeper seated tissues of the globe, as the vitreous humor, retina, or choroid, may become inflamed.

Death very frequently ensues before the end of the first year, either in consequence of the severity of the coryza and the inability to nourish the little patient, or in consequence of the profound cachexia and anæmia, or the development of some of the visceral lesions, to be hereafter described. When, however, owing to judicious treatment, or the comparatively slight development of the early symptoms, the child survives, the disease frequently subsides about the end of the first year; but often, after remaining latent for a variable time, reappears in the form of tertiary symptoms. According to Hutchinson, this tertiary epoch may begin at any period after the fifth year, but is commonly delayed till at or near the period of puberty. In addition to the traces which may remain of the earlier symptoms, such as little pits and scars upon the skin, alterations in the form of the nose from long-standing nasal obstruction, or actual disease of the nasal bones, there are several very characteristic symptoms amongst the later manifestations.

Among these is a peculiar alteration of the permanent incisor teeth, first described by Mr. Jonathan Hutchinson. Although we are not altogether disposed to attach the overpowering weight which Mr. Hutchinson does to the evidence furnished by this alteration of the teeth, of the existence of inherited syphilis, there is no doubt that it is an important sign, and we, therefore, quote in full his description of it (*loc. cit.*, p. 317):

"In these patients (those suffering with inherited syphilis), it is very common to find all the incisor teeth dwarfed and malformed. Sometimes the canines are affected also. These teeth are narrow and rounded, and peg-like; their edges are jagged and notched. Owing to their smallness, their sides do not touch, and interspaces are left. It is, however, the upper central incisors which are the most reliable for purposes of diagnosis. When the other teeth are affected these very rarely escape, and very often they are malformed when all the others are of fairly good shape. The characteristic malformation of the upper central incisors consists in a dwarfing of the tooth, which is usually both narrow and short, and in the atrophy of its middle lobe. This atrophy leaves a single broad notch (vertical) in the edge of the tooth, and sometimes from this notch a shallow furrow passes upwards on both the anterior and posterior surface nearly to the gum. This notching is usually symmetrical. It may vary much in degree in different cases; sometimes the teeth diverge, and at others they slant towards each other. In a few rare cases, only one of the upper central

¹ Duhring, *op. cit.*, p. 498.

incisors is malformed, the other being of natural shape and size. It is only in the permanent set that such peculiarities are to be observed; the first set are liable to premature decay, but are not malformed."

Another valuable symptom of inherited syphilis at this stage, and one which never occurs in acquired syphilis, is a peculiar form of keratitis, or inflammation of the cornea, which has been termed *interstitial* or *syphilitic*. It also is usually symmetrical, and is attended by opacity of the cornea from the formation of lymph in their substance. The inflammation usually subsides in a few weeks or months, leaving slight cloudy opacities here and there in the substance of the cornea.

Occasionally also there are symptoms indicative of grave visceral disease. The liver and spleen may be found enlarged and firm, and in such cases ascites is not rare. So, too, affections of the nervous system, usually limited to a single pair of cerebral nerves, as the auditory, and causing deafness, or the optic, and causing amaurosis, are met with in some instances.

Even at this stage marked disease of the bones is rare, though nodes quite frequently form upon the long bones; and Parrot calls attention to the fact that, in young infants affected with congenital syphilis, even when all other signs may be wanting, tumefaction of the bones will often be found.

These are more clearly marked on the inner surface of the tibiæ, on the lower part of the shaft of the humerus, and on the cranial bones. These latter form rounded bumps, and may not appear before the seventh or eighth or even twelfth month. More rarely, in very young infants (two weeks to two or three months), there may be found one or two fusiform nodosities in the continuity of long bones, which are due to semi-osseous callus around the seat of an undetected fracture.

In some few cases, the disease breaks out in the form of destructive lupus, which is apt to be associated with serious disease of the bony tissue.

MORBID ANATOMY.—The principal lesions found in the victims of inherited syphilis are in connection with the liver and lungs; more rarely other organs, as the brain or thymus gland, present evidences of disease. The liver is at times enlarged, rounded, and indurated, apparently the result of diffuse subacute hepatitis, or of infiltration of the organ with the peculiar albuminoid substance, called "amyloid" by Virchow. It is comparatively rare in children to find gummy tumors developed in the substance of the liver, with thickening and cicatricial puckering of the capsule, as are so often met with in visceral syphilis in adult life.

In the lungs, gummy tumors of various sizes form, and usually present cheesy degeneration of their central portion; and there is at times also a form of consolidation, called by Virchow "white hepatization," which depends upon chronic catarrhal pneumonia, with infarction of the air-vesicles with epithelial cells, in a state of partial cheesy degeneration.

More rarely, gummy tumors have been found in the substance of the brain. The thymus gland is occasionally the seat of suppurative inflammation, so that, on section, abscesses may be detected in the substance of the organ. Of course in cases where periostitis, with the formation of nodes, has been present, the ordinary appearances of such lesions will be observed.

DIAGNOSIS.—During the presence of the early symptoms the diagnosis is usually made with ease, by observing the presence of pemphigus soon after birth; of other eruptions, with copper-colored discoloration of the skin, appearing a few weeks later; of condylomata and rhagades; and of coryza, stomatitis, and laryngitis. The general symptoms are also peculiar, especially the physiognomy and the discoloration of the skin. And we should, in addition, endeavor to confirm our suspicion by obtaining a clear history of the parents' condition at the time of conception.

During the later periods of the disease, at or after the period of puberty, the diagnosis is no less important, and far more obscure. We must now rely upon the history of the case, upon the condition of other children of the same family, upon the detection of traces of the earlier symptoms, upon the presence of the peculiar alteration of teeth described by Hutchinson, of interstitial keratitis, of nodes, or of a symmetrical affection of some of the cranial nerves.

In deciding between the inherited or acquired nature of any case, the points which will aid us are the existence of primary disease of the mother at the time of delivery (which is rare, and can hardly be discovered even if it have been present); the existence of secondary contagious symptoms on either the mother or the nurse who suckled the infant; the presence of notched incisor teeth or of interstitial keratitis, which are peculiar to the inherited form; and the symmetrical distribution of all the secondary and tertiary manifestations, which is asserted by Hutchinson to be also an attribute of inherited as distinguished from acquired syphilis.

PROGNOSIS.—The most unfavorable conditions in inherited syphilis are the infection of both parents; the appearance of the disease soon after birth, especially in the form of pemphigus; and the occurrence of rapid and extreme emaciation. On the other hand, if the father alone has secondary symptoms, and those of a mild character; if the disease do not make its appearance till the third or fourth week; if the general nutrition is not greatly impaired, and if proper treatment can be immediately instituted, the prognosis is favorable, at least as regards preservation of life.

TREATMENT.—If the previous children of a mother have proved syphilitic, it is well to subject her to a mild mercurial course during her pregnancy.

In the treatment of the infant, every care must be paid to support its strength by the most nutritious diet, if it is unable to suckle the mother. It is, however, improper to employ a wet-nurse, on account of the danger of her being infected by the child.

In regard to medicinal treatment, the use of mercury is universally recommended during the presence of marked symptoms. The mercurial may be given either in the form of hydrarg. cum creta; calomel; or bichloride of mercury, in solution in some aromatic water or syrup; or, finally, it may be introduced into the system in the form of mercurial ointment by inunction. The most convenient mode of introducing it in the latter form is by smearing a flannel roller with mercurial ointment, and binding it around the child, whose movements cause its speedy absorption.

The dose of the mercurial should be small, and it is to be continued steadily though with caution, so as to avoid producing salivation, until a decided improvement in the symptoms manifests itself. During its administration it will frequently have to be temporarily discontinued, on account of gastro-intestinal irritation.

So soon as the mercury is stopped, we should order the iodide of potassium or iodide of iron, either one or both together being employed, according to the toleration of the stomach.

We should also recommend the use of cod-liver oil, and some preparation of cinchona, from an early period in the case; and even when the child suckles, a certain amount of Liebig's cold extract of meat, or of raw beef scraped finely and given as directed at page 435, should be administered.

The best local application to the sores is black-wash, though the condylomata usually require to be touched occasionally with solid nitrate of silver.

CLASS VII.

GENERAL DISEASES RESULTING FROM SPECIAL MORBID AGENTS OPERATING FROM WITHOUT.

ARTICLE I.

TYPHOID FEVER.

It is only of late years that the frequent occurrence of typhoid fever in young children has been fully recognized by medical authors. From the date of the publication of the classical work of Louis on this disease, until the year 1839, it appears to have been the almost universal belief that it was an affection limited to adult life; and with the exception of a few brief and vague descriptions, which evidently referred to this disease, though other names were used to designate it, medical literature contained no account of typhoid fever as it occurs in childhood. In the latter part of 1839, however, Rilliet (*Thèse de la Faculté*, 1840; and *Maladies des Enfants*, t. ii, pp. 663-739) and Taupin (*Journal des Connaissances Méd.-Chirurgicales*) published separate and independent memoirs on this subject; and since that time the occurrence of typhoid fever in children has been frequently observed and very carefully studied.

The fact that it was so long overlooked, is undoubtedly to be explained, in great part, by certain peculiarities which the disease presents in children, which caused its real nature to be mistaken, and led to the application of other names.

Of these names, that of infantile remittent fever was the most frequently used, and though this term was made to include a number of other diseases, and although remittent fever does occur in children, there can now be no doubt that a large proportion of the causes so styled were in reality cases of typhoid fever.

CAUSES.—*Age*.—Typhoid fever has been observed during the first year of life, but is rare under the age of two years. We have, however, met with well marked instances of it at the age of eighteen or twenty months. It is comparatively frequent between the ages of three and eight years, and it attains its maximum of frequency in childhood between the ages of eight and eleven years.

Sex.—The statistics of most authorities show a preponderance, more or less marked, of cases occurring in boys. In some series of cases this disparity has been remarkable (three to one); but, notwithstanding, it is probable that in a very extensive series the difference would be comparatively trifling.

Contagion: Epidemic Influence.—If typhoid fever be at all contagious, it is so in the slightest degree. On the other hand, it is well known that the dejecta from patients with this disease possess the power of producing it in those who are exposed to their emanations, or who drink fluids which have been allowed to become in any way tinted by them. The noxious vapors from foul sewers, drains, or cesspools are also frequently the cause of typhoid fever. It is, moreover, subject both to epidemic and endemic influences in a marked degree; and it is owing to the varying action of these causes that it presents the wide variety, in type and severity, which will be described.

ANATOMICAL APPEARANCES.—These are strictly analogous to those found in the adult. When death occurs early in the attack, the agminate glands of the ileum are found swollen, prominent, injected; the alteration being most marked in those nearest to the ileo-cæcal valve. Later, however, these glands ulcerate, the softening beginning either on the surface, and extending more and more deeply, or beginning in the deeper portion of the patch, so that the superficial layer may be thrown off as a slough.

These ulcers thus destroy the mucous membrane, and present the sub-mucous or muscular coat for their base; or, in some instances, the ulcerative process may extend through the muscular and even through the peritoneal coats. We have known the most violent general peritonitis to be excited early in an attack of typhoid fever, in a girl 4 years of age, by the simple extension of inflammation, without the occurrence of perforation.

The glands of Peyer are much less fully developed in the child than in the adult, so that it is probable that, as Jacobi suggests, the greater mildness of typhoid fever in childhood may be due to the fact, noted also by Rilliet (*loc. cit.*), that the ulcers are more slow in forming, smaller, and less numerous and deep.

The solitary glands are, in the early stage of the disease, prominent, and may be distended with a serous or more thick and yellowish secretion, so as to resemble vesicles or even pustules. Later in the attack their mucous covering is destroyed, and small, round, or oval ulcers, with everted edges, remain. These ulcers are also most numerous in the lower part of the ileum, though in some cases they are met with quite abundantly in the large intestine.

The mesenteric glands are enlarged, softened, and strongly injected, the change corresponding in intensity to that of the agminate glands in the ileum, and being most marked in those glands which are nearest the ileo-cæcal valve. Usually the swelling of these glands subsides without suppuration occurring, but occasionally this ensues, and the gland is converted into an abscess with thin walls.

The cicatrization of the intestinal ulcers appears usually to occur rapidly; thus Rilliet has seen the process completed by the thirtieth day, though this is probably sooner than it is entirely finished in the majority of cases.

Ulcers of other mucous surfaces, as of the pharynx and larynx, are

more rarely met with in children than in adults. The spleen is nearly always considerably enlarged and softened.

The blood in severe cases is dark and uncoagulable, and the lining membrane of the heart and large vessels is stained by imbibition. In some cases quite firm coagula are met with in the cavities of the heart.

Even in cases where the most violent nervous symptoms have been present, the brain rarely presents any more positive lesion than mere congestion of the vessels of its membranes and substance, with at times some subarachnoid effusion. Of course, in cases where death has resulted in consequence of some complication, the lesions of the intercurrent disease will be found.

SYMPTOMS.—The general course of typhoid fever is much the same in children as in adults.

It presents also the same wide variety in its type and degree of severity, depending upon the predominance and excessive development of some one of the elements of the disease; and it would be easy, therefore, to divide the disease into a great number of forms, according to the prominence of each functional disturbance; but as our object is merely to give a practical description of the disease as met with in children, we will, in considering its course, give a brief sketch of an ordinary case, and then dwell in detail upon certain symptoms which require special notice, as presenting special peculiarities in childhood.

In the majority of cases, the attack is preceded for some days by slight prodromes; the child, who ordinarily may enjoy robust health, appears languid, and is easily tired, and indisposed to play; he loses his appetite, is restless during sleep, and possibly may complain of colicky pain in the abdomen, perhaps attended with slight looseness of the bowels. After this state of vague indisposition has lasted from three or four to eight or ten days, more decided symptoms manifest themselves, and the attack may be said to fairly begin.

More or less febrile action now appears; but this is rarely continuous, and for the ensuing five or six days there are distinct and marked remissions, usually in the morning, but sometimes so marked and prolonged that it is only towards night that the skin becomes heated, the pulse frequent, and the child grows restless, while, during the day, he has merely appeared somewhat dull and languid. The loss of appetite continues, and becomes more complete, though thirst is marked; vomiting is apt to follow eating, and is sometimes frequent and spontaneous; the tongue presents a moist, whitish-yellow fur in the centre. The bowels either continue loose, or now become so for the first time; the abdomen becomes somewhat large and tympanitic, and slight tenderness may be present in the right iliac region towards the close of the first week.

The strength is rapidly lost, and the child, after the first few days, shows no desire to leave the bed. The respirations are somewhat hurried, and are often accompanied by sonorous râles and slight dry cough. The pulse is accelerated, but rarely rises at this stage above 110.

The expression grows dull and listless, unless temporarily excited during delirium, and the child takes but little notice of surrounding persons or

objects. During the night there may even now be a tendency to more marked cerebral disturbance, and the little patient grows very restless, utters sharp, shrill cries, or talks unmeaningly.

About the end of the first week the characteristic eruption appears, first on the upper part of the abdomen, in the form of small, oval spots, scarcely, if at all, elevated above the surface, of a light rose color, disappearing on very slight pressure, and quite rapidly returning.

During the second week, the symptoms become more severe. The fever is more continuous, and the temperature ranges in different cases from 102° to 105° ; it may still present, however, decided morning remissions, and it is not rare for profuse warm perspiration to occur, without having any critical value whatever. The pulse becomes more frequent, 120, 140, even 160, and at the same time smaller and of less force. The respirations are also more hurried, and, when the pulmonary complication is marked, may be very rapid and shallow, and the cough frequent and annoying; in such cases, auscultation reveals, especially over the postero-inferior part of the lungs, abundant mucous or subcrepitant râles. The vomiting ceases, and the child will usually take the liquid food offered it; the tongue becomes more heavily furred, and may be dry and brownish in the centre, though it often remains moist and yellowish-white throughout. Thirst is apt to diminish, owing to dulness of the perceptions, but the child will frequently drink greedily of cold water if offered to it. The diarrhœa persists, however, and the stools are ochre-yellow and fluid; the belly is more tympanitic, and may be extremely distended. The discharges, both of urine and fæces, are often involuntary, and the child does not even appear conscious of them. The urine is high-colored and scanty. The eruption continues and becomes more abundant, the spots which appeared passing away and being followed by successive crops. Sudamina are also frequently present, especially when sweating occurs. The mind becomes more and more dull, though it is nearly always possible to rouse the child by speaking loudly to it; delirium is usually present, especially in the night, and manifests itself in young children by restlessness, sharp, unmeaning cries, and a wild expression of the face, and in older ones by muttering, or even by attempts to leave the bed. Irregular muscular movements such as floccitatio and subsultus, are rarely noticed; though at times these, and even spasmodic rigidity of the trunk or limbs, or convulsions, may be present.

We have thus sketched the course of what is, perhaps, the most common form of typhoid fever in children, where the disease begins gradually, and either remains mild throughout, or assumes a more grave character during the second week.

In a certain number of cases, however, the onset of the disease is far more sudden and violent, and the severity of the attack is manifested from its earliest period. In this form, the prodromes are brief, or almost entirely absent; and there may be in older children an initial chill, or the only symptoms present are marked debility, languor, and drowsiness.

During even the first two or three days, however, there is apt also to be frequent vomiting, severe headache, or marked hebetude, and high fever,

which usually presents the same marked morning remissions and evening exacerbations as in the milder form. The sleep is restless and disturbed, and the child either utters sharp cries, or, if older, talks incoherently. The pulse and respiration are much accelerated, and the temperature of the surface rapidly rises, till, by the end of the first week, it may reach 103° or 105° . The cerebral disturbance may mask the presence of any abdominal pain; and as it is not unusual for the bowels to be quiet for the first few days, the case may closely simulate some acute cerebral disorder. By the end of the first week the disease is developed in its full severity. The fever is more nearly continuous, the morning remissions being comparatively slight, and the skin remains constantly dry and hot. There is deep stupor, from which the child is roused only with much difficulty, and which occasionally alternates at night with restlessness, jactitation, and noisy delirium. The pulse is very frequent and feeble, and the breathing accelerated, and usually accompanied with bronchial râles. The vomiting ceases, but the abdomen becomes tympanitic, and there is more or less abundant diarrhœa; the stools are often passed quite involuntarily, and the urine is either retained or dribbles away unconsciously. Epistaxis occurs in a large proportion of cases, and about this time the characteristic rose-colored eruption makes its appearance. During the second week, all of the symptoms become more grave, and the patient may succumb to the violence of the disease, or remain for a week or ten days plunged in profound stupor, with subsultus and marked muscular tremor; with the lips and teeth coated with sordes, the tongue tremulous, dry, and coated with brown crusts, the abdomen tympanitic, and the stools frequent, thin, and passed involuntarily; with the pulse running, feeble, from 130 to 160 in the minute; the respirations shallow, imperfect, and attended with subcrepitant râles, indicating passive congestion of the lungs; with the urine retained, dark-colored, and even albuminous; and yet gradually emerge from this apparently hopeless condition to enter upon convalescence about the close of the third week.

In favorable cases, between the fifteenth and twenty-first day, the grave symptoms begin to abate. The child's expression becomes more natural, and often the earliest sign of approaching convalescence will be the appearance of a smile of recognition, or of pleasure at the consciousness of improvement. The fur upon the tongue becomes looser, moister, and begins to separate, and the appetite slowly returns; the distension of the abdomen diminishes, and the stools are again passed consciously and voluntarily, and gradually assume a healthy appearance. Restlessness and delirium disappear, and the sleep becomes quiet and refreshing; the fever subsides, and the temperature falls, and again shows a marked difference between the morning and evening. The child thus passes into a state of convalescence, which, when not disturbed by complications, is quite rapid, though attended with marked emaciation, extreme debility, and feebleness of digestive power, with a tendency to intestinal disturbances. In some rare cases, at times without assignable cause, at others from improper exposure or exertion, or indiscretions in diet, the patient suffers a relapse, the original symptoms reappear, and a second fully developed attack of typhoid

fever, attended with marked nervous symptoms, characteristic eruption, and diarrhœa, may ensue.

In very severe cases, on the contrary, and especially when a fatal result is to follow, the condition of the patient grows more and more grave after the end of the second week, unless, as at times happens, death has occurred sooner from the violence or malignancy of the attack. The nervous symptoms become more marked, and the child sinks into a deeper stupor, even approaching true coma, or the stupor is interrupted by violent agitation, with cries or efforts to leave the bed, or by muscular twitchings, picking at the bed-clothes, or even general convulsions. The pulse is very rapid and small; the respirations hurried and noisy, and physical examination frequently reveals the existence of extensive bronchitis or hypostatic pneumonia. Vomiting is rarely present, but hiccough may be frequent and distressing; the belly is enormously distended, the stools frequent, involuntary, and at times bloody. Bedsores form on points subjected to pressure, and death ensues amid profound stupor and with signs of extreme pulmonary obstruction.

At other times death occurs not so much from the extreme violence of the disease itself as from the development of some one of the complications which will be mentioned hereafter.

SPECIAL SYMPTOMS.—Although, as has been seen, the general course of typhoid fever is much the same in children as in adults, there are a few symptoms which require more detailed notice, as presenting peculiarities which impress special features upon the disease as it occurs in childhood.

Prodromes.—In children, as in adults, typhoid fever is nearly always preceded by a marked prodromic stage, and the passage from the state of health to the fully developed disease is usually very gradual. The duration of these prodromes varies from three or four to ten days, being least in the more severe cases.

Fever.—Condition of Skin.—We have already remarked that, in the early stage, there are apt to be very marked remissions in the febrile action, lasting even throughout a considerable part of the day; the exacerbations of the fever usually occurring towards evening. West states that in some few instances two distinct remissions and exacerbations may be noticed in the course of every twenty-four hours. It is this feature which gained for the disease the name of infantile remittent fever, and caused it to be ranked formerly with the malarial diseases. Towards the middle of the second week, however, the remissions become much less marked; the temperature, which in some cases reaches 104° or 105° , merely presenting a somewhat marked fall in the morning. In general terms it may be said that the law of the accession of febrile temperature in typhoid fever in children, although in general correspondence with that found in adults, presents occasional marked interruptions. Thus, in some instances, the onset is so abrupt that a temperature of 104.5° has been noted by Roger on the first day, while it will be remembered as a law laid down by Wunderlich for the typhoid fever of adults, that the disease is never attended by a temperature of 104° so early as the second day. The skin is hot and dry as a general rule, but

sweats are more apt to occur during the height of the disease than they are in adults ; they are not, however, of any prognostic value.

Digestive Symptoms.—Among the earliest and most important symptoms are various disturbances of the digestive functions. The appetite rapidly fails, and is often lost before the attack fairly begins. Thirst is, however, marked until dulness of the mind appears, after which it also may be entirely absent, though the child will usually drink if cold water be offered to it. The tongue is already furred, usually being covered throughout the course of the disease by a thick yellowish-white coat, which may remain moist and loose, or, in very grave cases, become dry and brownish. Sordes are not often observed. Vomiting, which is perhaps not more frequently met with in the early stage in children than in adults, may be very frequent and persist until far into the second week. In the majority of cases, diarrhœa is either present or the bowels are peculiarly sensitive to the action of laxatives. In some case, however, and especially those where vomiting is marked, constipation of a quite obstinate form is present. The conjunction of these two symptoms, in connection with the cerebral symptoms present, may cause the case to strongly resemble the first stage of tubercular meningitis ; the doubt may, however, be usually resolved by careful examination, as will be more fully alluded to under the head of diagnosis.

The stools, when diarrhœa exists, are ochre-colored, fluid, and, on standing, deposit a sediment of shreds of mucous membrane, epithelium, and partially digested food. Mucus is rarely present ; but blood, in varying amount, may be mixed with the fecal matter. When the amount is large, it is usually due to the ulcerative process in the intestine having opened a vessel of considerable size, and then constitutes a very grave complication.

In young children it is difficult to establish the existence of abdominal pain, but when they are capable of describing their sensations, colicky pain is frequently complained of in the early stages ; and even in the youngest children, pressure in the right iliac region may often be seen to be painful.

Tympany is usually present at some time during the attack, especially when there is diarrhœa. Even when the bowels are confined, however, the abdomen is never retracted. Rilliet states that, in some grave cases, he observed such great tympany that the abdominal walls were thin enough to allow the outlines of the convolutions of the intestine to be clearly seen.

Enlargement of the spleen nearly always exists, but frequently to so slight a degree that it cannot be readily detected either by palpation or percussion, and even when considerably enlarged, it is apt to be entirely hidden by the distension of the abdomen. On the other hand, we have repeatedly found this organ so much enlarged as to be distinctly perceptible on careful palpation.

The *urine* presents the ordinary febrile conditions, being scanty, high-colored, and of high specific gravity ; the pigment is increased, and the chlorides much diminished.

The stools are, as we have already said, often involuntary during the height of grave cases, after the beginning of the second week. Until this

time, however, and throughout the entire course of more mild cases, the child is conscious of the desire, and can control the passage, or even wishes to be taken from the bed for the purpose.

The urine is also, though more rarely, discharged involuntarily; in rare cases, which may ultimately recover, retention of urine is present, and is of grave import. Rilliet never observed this symptom, but we have seen it more than once, and especially in a boy aged five years, who required catheterization for several days successively, but who finally recovered.

Respiratory and Circulatory Symptoms.—Even during the first week there is usually more or less dry cough, with sonorous and sibilant râles over the posterior part of the lungs. Indeed, in some cases, we have known the cough and signs of catarrhal inflammation to be so marked in the first days of the disease as to cause the attack to be regarded as one of severe acute bronchitis. Later in the disease, and owing merely to the passive hypostatic congestion of the lungs, and the accumulation of mucus in the bronchial tubes, the cough is apt to grow more frequent and troublesome, the respiration is hurried and oppressed, and auscultation reveals moist and dry râles throughout both lungs. When pneumonia or bronchitis supervene, these symptoms of respiratory obstruction increase to a marked degree. Extreme rapidity of breathing, with alterations in its character and rhythm, are also met with, however, in cases where the pulmonary obstruction seems moderate, but where the nervous system is profoundly disturbed.

The *pulse* is accelerated from the very first, and during the height of the disease rises to 120, 140, or even 180, according to the age of the child. In grave cases it may become extremely small, feeble, and compressible, but scarcely ever is intermittent or irregular.

The *eruption* of typhoid fever in children presents precisely the same appearances as in the adult; it usually appears first on the upper part of the abdomen, and often presents several successive crops. It is, however, more frequently absent entirely, and presents even greater irregularities, as to the date of its appearance, in them than in adults. The abundance of the eruption certainly bears no relation whatever to the severity of the attack; and in a varying proportion of cases (7 in 30, Hillier), the most careful daily examination fails to detect the characteristic spots at any period of the case. The eruption makes its appearance in a large majority of cases between the sixth and twelfth days, but the first spot has been observed so late as the twenty-fifth day (Hillier), or the twenty-ninth (Rilliet).

Sudamina are frequently present in large numbers at any time after the ninth day.

It is very important to be aware that in some cases, owing to the peculiar state of the cutaneous circulation, a marked reddish streak will be produced if the finger be somewhat firmly drawn across the skin. This sign, which we have described under the name of "*tâche meningitique*," in our article on tubercular meningitis, does not therefore possess the high degree of diagnostic value accorded to it by Trousseau and others, which would make it of much use in doubtful cases.

Epistaxis is very rarely abundant, but is met with in a majority of cases at some period after the third day.

Nervous Symptoms.—In none of the symptoms of this disease is such variety observed as in those furnished by the nervous system.

In mild cases, consciousness is retained throughout the attack; the expression of the face is stupid and heavy; the child is dull and disposed to doze during the day, but becomes feverish and restless towards night, and sleeps uneasily and wakes frequently.

In more severe cases, the nervous symptoms soon become prominent. The face assumes an almost characteristic expression; the eyes are dull and vacant, or bright and excited during temporary delirium; the cheeks present a circumscribed flush; the lips are dry and parched; and the features remain almost motionless.

Headache is sometimes complained of, and without doubt exists in many cases when the child is too young to call attention to it. It is especially observed in the early part of the attack, when there may be some hebetude and deafness present, and, according to Dr. Jenner, ceases upon the appearance of delirium.

This latter symptom rarely appears in marked degree before the second week, but then may become violent, the child crying out loudly, or muttering incoherently, and struggling violently to leave its bed. The delirium is rarely continuous, but is more marked during the night, being replaced during the day by more or less profound stupor, which, however, rarely amounts to actual coma.

Subsultus and carphologia, as well as muscular rigidity, are comparatively rarely observed in children, and only in very grave cases. Convulsions, even of a general and violent character, are met with in a very small proportion of cases; they may occur in the early stages of cases which subsequently recover, or as one of the final phenomena in fatal cases. They are, however, at whatever stage they present themselves, of very great import. In a case mentioned by West, the convulsions recurred on two successive days at the middle of the third week of the fever, and were succeeded by hemiplegia, which continued, though gradually diminishing, for four days. The child was unconscious even before their occurrence, and continued so for several days, though he eventually recovered.

As a general rule, the course of typhoid fever is much less apt to be attended by any *complication* in children than in adults; there are, however, some which occur with considerable frequency.

We have already stated that cough and signs of slight bronchitis are frequent in the early stage. In a considerable number of cases these symptoms become aggravated as the case progresses, and there may be a development of general bronchitis or even pneumonia; more frequently, however, the condition of the lungs is rather one of hypostatic congestion than of true inflammation. These complications when present in a marked degree, protract the case and add greatly to its danger. Pleurisy is comparatively rare.

Perforation of the ileum, from ulceration of Peyer's patches, is more rare in children than in adults; but when present gives rise to the same

symptoms, and leads to an equally rapidly fatal result. In some cases, its occurrence is announced by an attack of convulsions (Rilliet). We have already alluded to the fact that we have known violent general peritonitis to be excited by extension of inflammation without the occurrence of actual perforation of the bowel.

Intestinal hemorrhage, on the other hand, is comparatively frequent; thus Hillier observed it four times out of thirty in which the stools were carefully examined. It is usually of grave significance, but is at times seen in mild cases, which recover readily.

Earache is not infrequently observed after the height of the disease; in some cases it is followed by abundant purulent discharge.

Inflammation of the parotid gland is much less frequent than in adults, as is also phlegmasia alba dolens, of which, however, there are instances on record.

There is very little tendency to the formation of bedsores in children, and with care in the management of the patient, they will scarcely ever occur. In some epidemics, gangrene of other parts, as of the vulva or cheek, have been observed in a few instances. Angina, and occasionally pseudo-membranous laryngitis, have also been noticed.

We have seen that the urine is at times albuminous, and in these cases there is undoubtedly an intense congestion of the kidneys, which in very rare instances eventuates in Bright's disease. Œdema is not usually present, even when there is albuminuria, though Rilliet records two cases where anasarca, accompanied by albuminous urine, appeared on the fifth day, and lasted about a week. When œdema appears late in the course of the disease, it is probably to be rather attributed to a watery state of the blood and the debility of the circulation.

We have already seen that the febrile movement in typhoid fever, in children, presents such marked remissions, as to have led many observers to apply the name infantile remittent fever to the disease. We must bear in mind, however, that it is far from being rare for a true malarial element to be present, complicating the case, and constituting it a typho-malarial fever.

During the height of the disease, it is rare for any of the other eruptive fevers to make their appearance; but during convalescence, variola, rubella, and scarlatina have all been occasionally observed to appear, and run through their regular course.

Tuberculosis is by some regarded as one of the most frequent of the sequelæ of typhoid fever in childhood; and in some cases, indeed, it appears as though the extreme debility of constitution induced by the disease favored the development of tubercle in children with hereditary predisposition. In other cases it is probable that the production of tuberculosis after typhoid fever depends upon the infection of the system by the products of cheesy metamorphosis derived from some of the mesenteric glands, which, instead of returning to their normal state, have undergone this form of degeneration. It is probable, however, that in some cases also the early stage of acute tuberculosis has been mistaken for typhoid fever, with which,

as will be more clearly pointed out, it possesses some strong features of resemblance.

Various disturbances of the nervous system may occur as sequelæ of typhoid fever in childhood. Among these may be mentioned paralysis, either in the forms of paraplegia or limited to a single nerve-trunk, chorea, and locomotor ataxia.

CONVALESCENCE.—The convalescence is, as in adults, tedious and uncertain. The child often remains for many weeks in a condition of great debility, and with such extreme nervous exhaustion, that hydrencephaloid symptoms may even be present.

The digestive system also manifests this debility in a most marked degree, and it requires the greatest tact and care to encourage the child to eat, and at the same time to regulate the diet, since the slightest indiscretion will serve to excite serious symptoms. Not rarely death ensues many weeks after the termination of the disease itself, in a state of intense emaciation, the child being worn out by persistent diarrhœa, which resists all change of diet and treatment.

We have already alluded to the fact, that occasionally relapses have been observed, either without cause or following some trifling indiscretion, in which the symptoms of the fully developed disease have reappeared and gone through their regular course.

DURATION.—The duration of the fever varies according to the severity of the case. Even in the mildest forms it rarely begins to subside before the end of the second week, while much more frequently it is protracted until from the twentieth to the twenty-third day. In many cases, indeed, convalescence cannot be said to be fairly entered upon before the end of the fourth week.

PROGNOSIS AND MORTALITY.—The symptoms and conditions which indicate a favorable or unfavorable termination to the case are the same as present themselves in the adult, and may be readily gathered from the foregoing description. The mortality of typhoid fever is, however, decidedly less in children than in adults, partly owing to the comparative rarity of dangerous complications, and partly to the fact that the disease is usually of a less severe type. In mild cases, death scarcely ever occurs; and even in the more severe forms, the mortality is only from 5 to 10 per cent., under favorable hygienic circumstances.

DIAGNOSIS.—We have already stated that, partly owing to the imperfect recognition of typhoid fever, and partly to the various names which were loosely applied to this disease as occurring in children, it was formerly frequently confounded with other affections.

There are, however, several diseases from which it is not always easy, even with our improved knowledge of its peculiar symptoms, to distinguish it.

Thus, in some cases of gastro-enteritis, such as are not rare among children, and especially when the disease assumes a typhoid form, the resemblance to typhoid fever is so great as to have led Rilliet and Barthez to assert that it is impossible to make a differential diagnosis.

It should be borne in mind, however, that typhoid fever may often be

traced to epidemic or endemic influence, and occasionally to contagion; that it is very rarely possible to assign any direct exciting cause for the attack; and that it especially attacks children over five years of age, comparatively rarely those between two and five years, and very rarely those under the former age. Its onset is usually more gradual; the vomiting and diarrhœa are rarely so marked; the fever is more intense, the loss of strength greater and more rapid; while the marked dulness alternating with delirium during the night, the occurrence of the characteristic eruption and of epistaxis, and the more fixed duration, form a group of symptoms which should serve, when present, to clearly distinguish these two diseases.

In some cases, as already stated, the pulmonary complication, either in the form of diffuse bronchitis or of pneumonia, appears so early and causes such marked symptoms as tend to conceal those of the constitutional disease, and render care necessary to avoid overlooking it entirely.

On the other hand, it occasionally happens, and more frequently in children than in adults, that cases of pneumonia assume a typhoid condition, and present very many of the general symptoms of typhoid fever. It will, however, usually be sufficient in cases of this kind to pay careful attention to the early symptoms and mode of development of the disease, as well as to the existence or absence of the characteristic symptoms of typhoid fever, such as diarrhœa, tympany, epistaxis, rose-colored eruption, to avoid any error in diagnosis.

In some cases of acute, general tuberculosis, in which the deposit affects the brain, lungs, and intestinal canal, the symptoms may closely resemble those of typhoid fever. This form of tubercular disease may develop itself in the midst of seeming good health, the child losing strength and spirits; fever of a remittent type soon appearing; with vomiting, diarrhœa, tympanitic abdomen, and dry, furred tongue; and dulness of mind during the day, alternating with delirium at night. At the same time there is cough and rapidity of respiration, though the deposit in the lungs may be too slight and uniformly diffused to reveal itself by any positive physical signs.

In some such cases, indeed, it is only possible to form a probable diagnosis, based upon the age and previous history of the child; for acute general tuberculosis appears even at the earliest ages, and especially in children who have an hereditary tendency to tubercular disease, or who are delicate and frail, or have lately passed through an attack of some one of the eruptive fevers, or of hooping-cough; and upon the absence of eruption and the greater duration of the case.

Usually, however, there is a sufficient ground for diagnosis furnished by the special symptom, even early in the course of the case. Thus, in typhoid fever the vomiting in the early stage is rarely frequent or obstinate, and only follows eating; and, though the bowels may be constipated for a day or two, diarrhœa soon makes its appearance, and the abdomen begins early to enlarge. In acute tuberculosis, on the other hand, the vomiting in the early stage is usually both frequent and obstinate, and occurs entirely causelessly; whilst the bowels are in most cases constipated, and the abdomen retracted until a much later period in the case, when the

disease of the mucous membrane excites diarrhœa. The approach of fever in the tubercular disease is more slow, its course less regular, and its degree less intense, as a general rule, than in typhoid fever.

The nervous symptoms in the early stage of the two affections may be almost identical, but before long, in cases of tuberculosis, some of the unmistakable signs of tubercular meningitis, such as strabismus or partial paralysis, usually appear. Epistaxis is rare in tuberculosis, and, of course, the characteristic eruption of typhoid fever is absent, though it must be borne in mind that this is not constant in the latter disease. In doubtful cases, ophthalmoscopic examination should never be omitted, since it will frequently reveal lesions of the optic nerve or retina in meningitis which are absent in typhoid fever. And, finally, through the pulmonary disease may in some cases of the tubercular affection be slight, and not reveal itself by positive physical signs, most important aid is often derived from a careful exploration of the chest. We have already alluded to the fact that although the production of a reddish streak by drawing the finger over the skin is frequently observed in tubercular meningitis, the occasional occurrence of the same sign in typhoid fever deprives it of much of its diagnostic value.

TREATMENT.—Typhoid fever in childhood requires the same general plan of treatment as in adults. In mild cases little else is required than strict attention to all hygienic precautions, and a supporting, but fluid and digestible, diet. Whatever complications ensue, should of course be treated appropriately. There are, however, a few indications in regard to which it may be well to speak more in detail.

When the fever is high, febrifuges, such as *liq. ammoniæ acetatis* and *sp. ætheris nitrosi*, should be given; to which a little *syr. ipecac.* may be added, if the cough be troublesome. The surface of the body should be sponged daily or several times a day with tepid water, to which a little vinegar may be added; or the child may be carefully lifted for a few minutes every day or every other day into a bath of about 65° to 75°.

If there is much gastric irritability in the early stage, food should be given in very small quantities, and should be of the lightest character, as milk with lime-water or weak beef extract; counter-irritation may be employed in the form of mustard-plasters to the epigastrium; or, if there be reason to think that the stomach contains undigested, irritating food, an emetic of *ipecacuanha* may be given. In cases where marked symptoms of gastro-hepatic disturbance occur at the beginning of the attack, a few small doses of calomel with bismuth, or of blue mass, followed by a very gentle laxative, will be followed by relief to these symptoms. If the bowels are constipated, very small doses of some mild laxative, as castor oil or *syr. rhei aromat.*, should be given during the first week; but when spontaneous diarrhœa is present, it should, unless it becomes excessive, not be interfered with. When, however, the stools exceed three or four daily, chalk mixture, with some vegetable astringent and opium, or small doses of opium and acetate of lead, or of nitrate of silver and opium, or of paregoric alone, may be administered.

In ordinary cases, the nervous symptoms scarcely require any especial attention. When, however, they become marked, it will often suffice to

apply wet cloths to the head, and to administer warm mustard foot-baths to allay the agitation. In cases where delirium becomes extreme, with great nervous agitation, the above remedies should still be used, but, in addition, small doses of chloroform with camphor-water, or even of opium, should be given, and will often produce the happiest effect. Dr. West speaks highly in such cases of the combination of opium and tartar-emetic, recommended by Graves in the treatment of the head symptoms of typhus fever. It will be found, also, that chloral, in doses of five grains at three to five years, or of bromide of potassium, in doses of seven to ten grains at the same age, repeated according to the urgency of the symptoms and the effect produced, will often prove successful in affording relief.

In regard to the occurrence of complications, we have already alluded to the remedies by which diarrhœa is to be checked if it becomes excessive. When the symptoms of pulmonary obstruction become marked, frequent counter-irritation by mustard or turpentine should be applied to the chest, and stimulating expectorants, as carbonate or muriate of ammonia, administered internally.

Hemorrhage from the bowels and peritonitis from perforation on extension of inflammation, must be treated exactly as in the adult, the one by astringents, either vegetable or mineral, or by ergot and moderate doses of opium; the other by the free use of opium.

Of the special remedies which are recommended in this disease, we may allude to the treatment by means of mineral acids, especially the muriatic and nitro-muriatic, which is highly praised by some authorities, and to that by nitrate of silver as recommended from extensive experience by ourselves. This latter remedy is best given to young children in the form of solution in a thin syrup of acacia; and the dose at the age of 3 to 5 years should be from gr. $\frac{1}{24}$ to gr. $\frac{1}{12}$ three or four times daily, to which may be added from one-half to two drops of deodorized laudanum.

Quinia is necessary in many cases as tonic when adynamic symptoms begin to appear, and is of service in cases attended with high temperature, when given in full doses with a view of reducing the excessive heat of the body. In some cases, also, where the remittent character of the fever is marked, and where there is a suspicion that the case is complicated with a malarial element, it should be administered in full antiperiodic doses in the earliest stages.

Opium is rarely necessary in the early part of the disease, unless it be required to check diarrhœa; but when, in the latter part of the second or third week, the delirium becomes extreme, and the child sleeps but little, the night being spent in violent restless agitation, with loud screaming, opium should be fearlessly given until quiet sleep is produced.

The oil of turpentine is to be administered under the same conditions which call for its use in adults.

Stimulants are by no means necessary in all cases of typhoid fever in children. Expecting in the very mildest, however, it is prudent to administer them in small quantities after the middle of the second week. When, however, the condition of the child calls for their freer use, as shown by the frequent feeble pulse, rapid labored breathing, dry, brownish tongue, dulness alternating with noisy delirium, and other

marked nervous symptoms, they should be given to the extent of fʒiij to fʒvj of sherry wine or even of brandy, to children of six years old. It will be found in these cases that even such large amounts of stimulants as the above are very well borne by children.

The food should be given in small quantities, and frequently repeated. It should throughout the entire course of the disease be exclusively fluid, consisting of milk, chicken-water, or the various animal broths. It is rarely difficult to regulate the diet of children suffering with this disease, since their entire loss of appetite renders them indifferent to all food, and they will usually take whatever is offered to them.

During convalescence, the utmost care must be exercised, both in regard to food and exercise. Solid food should be permitted very gradually, and with much caution; beginning with the lightest and most digestible forms, and watching the manner in which each article is digested.

In those cases where the child remains a long time in a condition of extreme debility, with impaired power of digestion, the bitter tonics and iron should be given, if the stomach will tolerate them. Sea, or cold-water bathing, change of residence, and the utmost attention to all hygienic rules, are also to be recommended. When there is any reason to dread the development of tubercular disease, this treatment must be carried out with the greatest assiduity; and, if the child can digest it, cod-liver oil may be given with advantage.

ARTICLE II.

VARIOLA, OR SMALL-POX.

DEFINITION; FREQUENCY; FORMS.—Variola is an epidemic and contagious disease, characterized by an initial fever, lasting from three to four days, and followed by an eruption, at first papular, then vesicular, and afterwards pustular; the eruption attains maturity in from six to nine days, after which the pustules are converted by desiccation into scabs, which fall off between the fifteenth and twenty-fifth days.

The *frequency* of the disease varies greatly in different years, because of its epidemic nature. It is far less common in childhood amongst the middle and upper classes of the community, than either measles or scarlatina, in consequence, no doubt, of the attention paid to vaccination. During the early months of 1865, one of us had the opportunity of studying a severe epidemic which occurred in portions of this city, and we have published elsewhere¹ an analysis of thirty cases in children under fifteen years of age, observed at that time. Apart from these cases, however, we had met with but two cases of the disease under fifteen years of age, during the fifteen years preceding 1871–72, whilst during the same period we had met with 263 of scarlatina, and upwards of 314 of measles. In the last-mentioned years, a severe epidemic occurred in this city, when we again

¹ Amer. Jour. of Med. Sci., October, 1869, p. 322.

saw numerous cases at all ages. It prevails to a greater extent amongst the poor and destitute classes, who neglect the attention to vaccination necessary to preserve children from the disease.

We abstract from the article already referred to, the following table, showing the entire annual mortality from variola in Philadelphia, together with the relative mortality during the early years of life, for the twenty-four years ending 1873; to which we have added the figures for the subsequent years up to 1879:

					MORTALITY FROM VARIOLA.			
					Under, 1 year.	Between 1 and 2 years.	Between 2 and 5 years.	Total of all ages.
1848,	21	13	17	100
1849,	25	20	34	152
1850,	13	8	4	40
1851,	40	30	54	216
1852,	89	54	100	427
1853,	22	9	9	57
1854,	12	4	6	49
1855,	57	39	85	275
1856,	86	44	88	390
1857,	19	17	11	65
1858,	1	2	1	7
1859,	0	0	1	2
1860,	14	10	16	57
1861,	159	105	200	758
1862,	52	44	66	264
1863,	33	24	28	171
1864,	57	31	61	260
1865,	104	50	112	524
1866,	32	17	27	144
1867,	16	4	11	48
1868,	0	0	0	1
1869,	1	3	0	6
1870,	3	0	3	9
1871,	203	112	292	1879
1872,	347	188	446	2585
1873,	10	5	3	39
1874,	2	0	0	15
1875,	8	4	14	54
1876,	57	35	102	407
1877,	17	11	39	155
1878,	0	0	0	0
1879,	0	0	2	6
Total,	1500	883	1832	9162

An inspection of this table undoubtedly establishes the fact that whenever the contagious principle of variola, favored by some peculiar epidemic influence, is introduced into this community, it finds a large number of unprotected subjects who fall ready victims to its attack.

We shall, in our description of variola, refer to several forms which it may assume. These are merely degrees of severity of the same disease—types given to each case by several causes. Chief amongst these is the presence or absence in the person attacked, of the protective power of the

vaccine disease, next is the type of the epidemic prevailing at the time, and last we must place the inexplicable and utterly uncertain influence of individual constitution. According to the degree of reaction of the variolous poison in the system of the patient, shall we have *distinct* or *discrete*, *confluent* or *hemorrhagic* small-pox; or that form modified by vaccination, inoculation, or previous natural small-pox, called *varioid*.

We shall also describe the *complications* of the disease.

CAUSES.—The principal causes of variola are contagion and epidemic influence.

It is not clearly ascertained at what period of its course the disease first acquires the property of infectiousness. Some assert that is not until after suppuration is established. This is, however to say the least, doubtful, and it is best, therefore, to take every precaution that may be necessary to prevent the extension of the disease, from the moment that its real nature becomes apparent. There can be no doubt that the body may still impart the disease after death, and that clothes worn by the patient retain the contagious principle, unless freely exposed to the air, for days, months, and, it is said, even for years. It is also capable of infecting furniture or letters, and may thus propagate the disease at any distance, and for an indefinite period, by fomites.

One attack protects the constitution, in the great majority of cases, against subsequent contagion. When persons who have once had the disease contract it again, it almost always assumes a much milder and less dangerous form.

In the report of the Municipal Hospital of Philadelphia, made to the Board of Health of Philadelphia, for the year 1872, Dr. Wm. M. Welch, the physician in charge, states that out of the whole number of cases (2377) of variola admitted during the violent epidemic of 1871–2, 15 were said to have had a previous attack of the disease. Of these 15 cases, those which could not show a single scar as the result, he should classify as of doubtful authenticity; those which exhibited only a few scars, as of probable authenticity; and those which exhibited well-marked pitting as authentic. To the first class belonged 7 cases, of which 3 died; to the second class belonged 3 cases, all of which recovered; to the third class belonged 5 cases, all of which recovered, and in all of which the eruption was very light, so much so in one as to be characterized as doubtful.

The period of *incubation*, or the time elapsing between the reception of the poison and the onset of the malady, varies generally between nine and twelve days. It may, however, be seven or fifteen days.

SYMPTOMS; COURSE; DURATION.—We shall describe three stages of the disease: 1. That of the initial or eruptive fever; 2. That of the progress and maturation of the eruption; 3. That of decline or desiccation. In addition to these, some writers make another stage, that of incubation, which includes the period between the introduction of the poison into the system and the appearance of the first symptoms. This stage is seldom marked by symptoms sufficiently characteristic to enable us to detect the approaching disease, and in many instances is probably entirely unnoticed by the patient.

The *first stage*, or that of initial fever, commences generally in children

with pains in the bones and loins, and sometimes with rigors or chilliness, accompanied with headache, and soon followed by fever. Nausea and vomiting often exist from the first, or come on soon after the appearance of fever and headache. At the same time there is loss of appetite, thirst, and more or less obstinate constipation. The tongue is red at the point and edges. One of the characteristic symptoms of this stage is pain in the loins, which generally dates from the first or second day, and which, though varying much in degree, is usually severe. The patients often complain also of abdominal pains, which seem to be colicky, and are referred either to the epigastric or umbilical region.

Fever and headache are the most constant of all the initial symptoms. The chilliness and rigors which frequently exist in adults are not easily ascertained in the cases of children, and are therefore much less important. The fever varies greatly as to degree; the heat of skin is generally considerable, the temperature rising to 104° or 105° , and may be accompanied either with dryness or moisture. The pulse is commonly full and frequent, rising to 120, 140, or 160 beats, according to the severity of the case and the age of the child. The headache is usually frontal and often very severe. In some cases there are strongly marked cerebral symptoms, consisting of excessive restlessness and irritability, insomnia or somnolence, delirium, and even convulsions.

The various symptoms just enumerated continue up to the moment when the eruption begins to make its appearance, which happens generally in the course of the third day, though it may occur as early as the second, or as late as the fifth, sixth, or even seventh. In severe and confluent attacks the eruption, as a general rule, begins earlier than in mild and discrete cases.

Second Stage, or that of Eruption.—In the great majority of cases, the specific eruption makes its appearance in the course of the third day from the beginning of the fever. This is the law of the disease. Before, however, describing it, we must state that not rarely a more or less extensive roseolous rash precedes the specific eruption. So well known is this that it has been called *roseola variolosa*. It looks so like measles as to make a correct diagnosis difficult, since nothing could reveal its true character unless it were known that the subject had been exposed to variolous infection, in which event the unusual severity of the constitutional phenomena, compared with those generally attendant upon roseola, might well lead the practitioner to defer his opinion. This roseola occurs in all forms of small-pox. Dr. Welch thinks he has seen it most frequently and in greatest quantity in cases of mild varioloid.

The specific eruption appears, then, on the third day, in the form of small, isolated, and rounded red specks, which soon become projecting and solid, or in other words are converted into papules. The papules are from a third to two-thirds of a line in diameter, of a more or less vivid red color, which disappears under pressure, to return immediately when the pressure is removed. They are also hard, and feel almost like shot imbedded in the derm. The eruption shows itself first on the face, and generally about the chin and mouth, and then extends to the rest of the face, to the neck, trunk, limbs, feet, and hands. It sometimes happens, par-

ticularly in very young children, that the eruption appears first about the genital organs, whilst in other cases it is first observed on the lower part of the loins, or upon the thighs. The papules increase gradually in size and prominence for one, two, or three days, and, as a general rule, some time in the course of the second day of the eruption begin to change into vesicles. This change takes place by the formation on the top of each papule of a little transparent elevation of the cuticle, beneath which is deposited a drop of serosity. The conversion of the papules into vesicles occurs first on the face, and then on the neck, trunk, and extremities. The vesicles are at first smaller than the papules, and acuminate in shape, but as they grow larger, become gradually flattened and depressed in the centre; after a time they cover the whole papule, and before long exceed it in size. As these changes take place the fluid they contain loses its transparency, becomes opaline, and by degrees the vesicles are transformed into pustules, and thus the third stage of the eruption or that of suppuration begins.

The pocks are more or less numerous, according to the extent and severity of the eruption. When scattered over the surface so as not to touch at their edges, the disease is said to be *distinct* or *discrete*; when, on the contrary, so numerous as to come into contact and run together, it is called *confluent*. Of these two varieties, the latter is necessarily more severe and dangerous than the former, in consequence of the greater extent of tegumentary surface inflamed. During the various changes the vesicles undergo, they are surrounded by small, inflamed areolæ, which differ in appearance according to the number of the vesicles. In cases of the discrete form, in which the eruption is sparse, so that the pocks are widely separated, the areolæ fade gradually into the natural color of the skin, at the distance of a third or two-thirds of a line or more from the base of the vesicles, whilst in those in which the eruption is more abundant, they run together, so that the spaces between the pocks are of a more or less bright-red color. In confluent attacks again, the areolæ are more or less imperfect, according to the manner in which the vesicles are grouped together.

The change of the vesicles into pustules takes place generally from the fourth to the sixth day of the eruption. During this process the fluid of the pocks becomes more and more opaque, whitish, and at length assumes a yellowish color, being in fact converted from serum into pus. At the same time the pocks become larger, begin to distend, and, as they approach complete maturation, gradually lose their umbilicated shape and become convex on the surface. The formation of the pustules follows the same course as did the vesicles, beginning on the face and extending thence to the neck, trunk, and extremities. The areolæ that have just been described as existing during the vesicular stage of the disease, continue during the early part of the stage of pustulation, but decline towards its termination, assuming as they disappear a purple tint. The number of pustules is in proportion, of course, to that of the vesicles, but from the increase in size of the pocks during the changes from papules into vesicles and pustules, the eruption, when at its height, seems to be greatly more extensive than would have seemed probable at the beginning

of the first stage. As a general rule the pocks are most numerous on the face, and after that part on the neck and limbs. On the trunk the eruption is always much less abundant than on other parts of the body, and even when confluent in the highest degree on the face and neck, it is generally so only in patches on the limbs, while it is discrete on the thorax and abdomen.

Simultaneously with the eruption upon the skin, there occurs one also upon the mucous membranes, particularly those of the mouth, nasal passages, fauces, eyelids, and sometimes of the prepuce and vulva. It begins with more or less vivid redness of the membrane, which is followed by the production of little elevations, the real nature of which, whether papular or vesicular, seems not to be clearly determined. About the second or third day these red elevations assume the appearance of small, whitish, rounded, and umbilicated pseudo-membranous points, which last generally about five days, and are then detached, leaving usually a little ulceration or erosion, which heals without leaving a cicatrix.

A short time after the appearance of the pustules in the mouth and throat, a true inflammation of the mucous membrane of those parts takes place. When the gums are inflamed they become swollen, red, and spongy, and are dotted over with white pseudo-membranous points of a rounded shape. Sometimes the velum pendulum, and more rarely the tongue, present the same white points, with redness and injection of the membrane between. In most of the cases there is also partial or general inflammation of the pharynx, which occurs subsequently to the formation of the variolous pustules. The existence of this inflammation is denoted by more or less severe sore throat, attended with difficulty of swallowing, and with swelling and tenderness of the submaxillary glands. When the mucous eruption extends to the larynx, as often happens, there is pain in that part; the voice becomes hoarse or whispering, and there is a hoarse, laryngeal, smothered cough. The pharyngo-laryngitis just described occurs generally between the third and sixth days of the eruption, and ceases about the eighth or thirteenth. In some instances it does not exist at all or only to a slight extent.

During the eruption there is more or less inflammation and swelling of the subcutaneous cellular tissue, the degree of which depends on the extent of the eruption. The skin becomes tense, red, shining, and elastic under the finger, and more or less hot and painful. The swelling is greatest upon the face, where it commences about the fourth or fifth day of the eruption, and goes on increasing for five or six days, occasioning much pain, stiffness, and inconvenience to the child. The swelling diminishes when desiccation begins, and ceases entirely as the latter is accomplished.

It is important to study carefully the general symptoms of the second stage. The fever which existed during the initial stage sometimes continues during the first day or two of the eruption. When, however, the papules are fully thrown out, the fever subsides or disappears entirely, so that the pulse falls from 100, 120, or 140 beats, to 100, 80, 76, or 74, and the heat of skin diminishes at the same time. The child remains without fever usually throughout the vesicular period of the eruption, that is to

say, until the fourth, fifth, or sixth day; during which time the appetite sometimes returns, sleep is tranquil and quiet, and the patient is in most respects well and comfortable.

About the fifth or sixth day of the eruption, at which time the maturation of the pustules is nearly completed on the face, and that process is commencing on the extremities, a new fever, to which the technical term *secondary fever* is applied, makes its appearance. The pulse rises again to 88, 100, 120, and 140, and becomes strong, hard, and full, whilst the skin is hot and dry. After continuing for some days the secondary fever diminishes after the suppuration is fully established, and disappears about the time that desiccation is nearly completed on the face, and has commenced upon the limbs. It ceases generally, therefore, about the ninth or eleventh day, having lasted between four and six days. This attack of fever is evidently the consequence of the suppurative stage of the disease, or of the conversion of the vesicles into pustules.

Towards the termination of the second stage, at the very height of the disease, when the pustules begin to break and discharge their contents, the patient exhales a peculiar, disagreeable, and fetid odor, which is characteristic of the disease.

The *third* or *declining stage* is that of the desiccation or drying of the pustules, and their desquamation. The desiccation commences generally between the sixth and ninth days, and terminates between the tenth and fourteenth. The formation of the crusts begins upon the face and extends thence to the neck and limbs. It does not reach the limbs usually until about two or four days after it has commenced on the face. The mode in which the drying of the pustules takes place is not the same in all. In some a dark point is formed in the centre, which gradually extends and converts the whole pustule into a hard crust, in others the whole surface dries at the same time; while in others again, the epidermis gives way and allows the contained fluid to escape, which then hardens into yellowish, irregular crusts, which become brown before they fall off. Some of the pustules, particularly those upon the arms and legs, do not form scabs at all, but shrink away from the absorption of their fluid, leaving behind nothing but pellicles of cuticle, which fall off by desquamation.

The desquamation or falling of the crusts begins from the eleventh to the sixteenth, and ends somewhere between the nineteenth, twenty-fifth, and even fortieth days of the eruption. When the scabs fall off, the appearances presented by the skin beneath vary in different cases. In some a true ulceration and loss of substance of the derm has taken place, which gives all the characters of a suppurating ulcer when desquamation has begun early in the disease; when that process occurs at a later period, the ulcer is found to be dry and cicatrized. In both these forms of desquamation, the cicatrices form little pits or depressions, which remain during life. In other instances, the fall of the scabs leaves red and excoriated surfaces which are on a level with the surrounding skin, but which soon dry, leaving blotches of a reddish-brown color, that do not disappear entirely for months. No cicatrices remain when desquamation takes place in this manner. In a third series of cases the crusts do not fall until the surface beneath has completely cicatrized, and the only traces left behind

are more or less deeply tinted reddish spots, with occasional slight furfureaceous exfoliation of the cuticle, all of which disappear entirely after a time without leaving pits or cicatrices.

To conclude the account of the symptoms of the disease, we have a few words to say in regard to the condition of some of the important organs throughout the course of the malady.

The *tongue* presents no appearance peculiar to the disease, other than the eruption already described. It is generally moist, more or less furred, and either pale or red in color. The *abdomen* usually remains soft and undistended, though in some instances it is slightly tumid and hard, with occasional pains in the epigastric, umbilical, or iliac regions; in simple cases, the latter symptoms rarely last more than a short time, and when otherwise they are almost always the sign of some complication. The *constipation* which exists during the initiatory stage generally continues throughout the disease, though in some instances a slight diarrhœa occurs about the end of the first or second week, after which the bowels regain their natural condition. If severe diarrhœa should make its appearance, it is almost always the sign of a dangerous complication. The *nausea and vomiting*, which are so often present during the initial stage, cease after the appearance of the eruption, and recur only in rare cases, or in consequence of some complication. The *appetite* is almost always lost during the course of the disease, though it sometimes returns in the period between the termination of the initial and the commencement of the secondary fever; *thirst* is acute as a general rule, and more or less so according to the violence of the fever.

The *urine* presents, during the course of the disease, the ordinary febrile characters of lessened quantity and heightened color. The urea, uric acid, and pigment are increased, and the chlorides much diminished. Albuminuria is occasionally present at the height of the disease; it is, however, temporary, and apparently not of very grave import. Casts of the renal tubules are also present in some cases. The frequency with which this condition exists probably varies in different epidemics, since we have detected it but rarely in our cases, while Parkes states that it is present in about 30 per cent. of all cases. After the subsidence of the secondary fever, the urine frequently becomes very abundant, of pale color, and of low sp. gr. Thus, in one of our cases, in a girl aged 18 years, the daily amount of urine passed from the tenth to the thirteenth day of the eruption was f̄3clx, or ten pints, of sp. gr. 1001, as clear as spring-water, containing no albumen, but with a fair proportion of chlorides. In another case, in a young man, aged 20 years, the patient also passed, on the eighteenth day, f̄3clx of crystal-clear urine, of sp. gr. 1007, without albumen but containing abundant chlorides. In a third case, in a boy aged 13 years, the amount, on the twelfth day of the eruption, was f̄3xlv.

The *strength* of the child is not, as a general rule, greatly diminished, except in severe and dangerous cases. *Restlessness, irritability, crying, and delirium*, which are of such frequent occurrence in the febrile diseases of children, are not usually very strongly marked in regular cases of variola. They exist, but it is to a moderate extent only.

We pass on now to the *confluent* forms of the disease.

It is not possible to predict from the character of the initial fever what is to be the type of eruptions which is to follow, since in discrete variola, and even in varioloid, the precursory fever and other symptoms, often run alarmingly high, while, on the other hand, a case destined to be confluent, or even hemorrhagic, does not always exhibit violent phenomena at the onset. As the time for the eruption approaches in confluent cases, the skin usually gives evidence of active inflammation of its deeper structures. It becomes thickened, swollen, hard, dark in tint, and as the eruption advances, the countless number of papules and vesicles, which cover all parts of the body, increase the violence of this inflammatory action, and give rise to an earlier appearance of the secondary fever, which is marked by higher temperature, more active delirium, and greater disturbance of the circulation than in discrete or moderate small-pox. As the vesicles form upon the papules, they so crowd the surface that their edges run together, thus making the confluence, and no portions of natural skin remain on which to form the areolæ, which, therefore, are absent. As the pustules follow the vesicles they do not develop well, but remain flattish and sluggish, with a whitish, ill-concocted pus on some parts of the body, particularly the face and backs of the hands. They run together into large flat blebs or bullæ, of several inches or more in extent. Sometimes portions of the loosened cuticle are rubbed off by the movements in bed, or by scratching. The parts thus denuded look raw, and exude a sanious fluid.

In severe confluent cases the eruption extends to the mucous membrane of the nose, mouth, fauces, eyelids, and perhaps to the prepuce or vulva, as in the distinct form, but with very different severity and consequences. The inflammation produced by the eruption causes enlargement of the tongue, swelling of the fauces, pain, and often great difficulty of swallowing. The rawness and soreness of the passages, and an abundant and usually dark-colored viscid secretion, which clogs and clings to the parts, cause great distress, and add to the exhaustion of the patient. At the same time the laryngeal catarrh causes cough, hoarseness, partial or total loss of voice, and difficulty of breathing. Thus, as in violent anginose scarlet fever, and in some cases of diphtheria, the supply of air to the lungs is so diminished by the various causes of obstruction (swelling, collections of viscid phlegm, and spasm of the glottis) that the blood does not receive its due amount of oxygen, a venous stasis is established, the skin becomes dark-brown or purplish from capillary stagnation, and the patient dies, sometimes in great distress, though at others with very little apparent suffering, in a state of asphyxia and exhaustion.

In some cases the heart presents evidences of disease: the sound becomes feeble and obscure, the impulse weak, that the action of the heart irregular and intermittent. These symptoms, to which special attention has been called by Desnos and Huchard,¹ are dependent upon grave inflammatory changes, either in the endo- or pericardium, or frequently in the muscular tissue of the heart. In cases where this latter lesion is present they have occasionally observed a want of agreement in frequency between the contractions of the heart and the radial pulse; and also, but as a much more

¹ Des complications cardiaques dans la variole, Paris, 1871.

constant sign, a murmur at the apex of the heart, soft, deep, diffuse, and inconstant, which differs in its character from the murmur which attends endocarditis of the valves. Undoubtedly in many cases of variola where death occurs suddenly, with signs of failure of cardiac power and pulmonary engorgement, the fatal event will be found to depend on the development of one of these cardiac lesions, and especially of myocarditis.

There is a form of confluent small-pox called superficial confluent, in which, though the eruption is really confluent, it runs through the stages of maturation, desiccation, and desquamation so rapidly that the constitution is not greatly tried, and the patient recovers without difficulty.

Even in the severe form, the patient may, if his constitution be good, pass safely through the disease.

The *hemorrhagic*, *malignant*, or *petechial* form is happily rare. We had rarely seen it until the epidemic of 1871-72 showed it to us in all its terrible power. Our forefathers knew all about it. We, of the generation which has risen since the introduction of vaccination, had read of it, but took little heed of what the variolous poison might do when it exerted its malignant forces. In this form the patient is weak and feeble from the beginning. The surface assumes a singular reddish hue as the eruption comes out. The vesicles when they form upon the papules, instead of filling with lymph, and then pus, contain only a thin, sanguinolent liquid; they mature very imperfectly, or rather not at all, not acuminating but remaining flattish, or irregular in shape, and flabby. While the eruption is struggling along in this irregular mode, the vessels of the cutaneous tissues become gorged and partially stagnant, so as to give to the surface dark-red, brown, blue, or purplish, and livid tints. Extravasations take place amongst and beneath the eruptive points, the cuticle forming the bloody pocks breaks, blood exudes, and forms dark scabs, and the patient is so changed from his natural aspect that we may comprehend how in the olden time, people who had not the consolation which vaccination gives, may have been driven from the bed and even from the house of the sufferer in hopeless terror. Such cases look no longer human. The swollen face, purple or black, the dark or crimson-red eyeball, with the whitish cornea sunken into a pit formed by the projection of the blood-colored and œdematous conjunctival membrane, the eyelids thick and stiff and imperfectly closing, the gross body, changed from all its natural bright to blackish tints, the cuticle dissected from the skin by bloody exudations, which weep and stain the clothing and bed-linen. Such is the variola nigra or black small-pox of the old writers, and well does it deserve its name.

VARIOLOID, OR MODIFIED SMALL-POX.—This is a term now usually applied to the modified form of the disease, as it occurs in individuals who have been vaccinated, or who have already had the natural or inoculated disease.

Dr. Welch's rule is a very good one,—“to classify as variola all unvaccinated cases, no matter how mild, all malignant cases, and all the vaccinated cases in which the eruption does not reach maturity until after the sixth or seventh day from its first appearance.” The true point of

distinction here, when any uncertainty as to vaccination exists (and this is not rare amongst the poor), is the time of maturation of the eruption. This, in varioloid, ought to be matured and in the decline by the sixth or seventh day.

The initial symptoms of varioloid are of the same general kind as those of natural small-pox, differing merely in degree. But the physician ought to know that, in a few cases of even very mild varioloid, while the eruption is destined to be sparse, to consist of but few pocks, and to run through its stages in five or six days, the initial fever may be very high, and the attendant phenomena of pain, nervous disturbances, loss of strength, etc., very marked. We once saw a girl nine years old, who was ill for three days with very high temperature, delirium, stupor, prostration, violent headache, and rapid pulse, so that her case looked very threatening and left the diagnosis in great doubt. On the third day a moderately abundant variolous eruption came out, when all the unpleasant symptoms rapidly abated and disappeared. The eruption ran through its stages in six days, and the patient recovered without a pit. She had been well and carefully vaccinated in infancy.

These severe initial symptoms are rare, however, in children as compared with adults. Usually the attack begins with slight fever, headache, languor, and sometimes constipation, which are followed, in two or three days, by the eruption. The vomiting, lumbar pains, and different nervous symptoms which exist in regular variola, are not often present, or, if so, in a very slight degree. The eruption consists of papules like those of true small-pox, but usually they are few in number, and entirely discrete in their arrangement. The initial fever and other symptoms subside completely upon the appearance of the eruption, and the child often seems perfectly well.

The progress and character of the eruption are very similar to those of the regular form of the disease, with the exception that the changes are more rapidly effected, and, as a consequence, the duration of the attack is rendered much shorter. The papules are converted into vesicles at a much earlier period—as early as the first or second day. The vesicles soon assume a whitish, opaline appearance, become umbilicated, and in the course of the second or third day begin to change into pustules. The suppurative stage of the eruption, or maturation, is seldom accompanied by any marked secondary fever, as in the regular disease. When the fever does occur, it is generally very moderate, consisting merely in slight acceleration of the pulse and a little increased heat of skin, and in one or two days it disappears entirely. The pustules do not fill usually so well as in regular variola, and not unfrequently their contents are rather sero-purulent than purulent in the proper sense of the term. The third stage occurs earlier and goes through its period more rapidly than in true small-pox; desiccation soon takes place, is speedily finished, and the falling of the scabs, which begins as early as the eighth day of the eruption, is usually completed about the twelfth or fourteenth. After desquamation is completed, the only traces of the disease left are reddish spots or blotches,

which disappear after a time without leaving cicatrices. The whole duration of the attack is generally from ten to twenty days.

Varioloid may be so mild that the patient never goes to bed. Some malaise, a little loss of appetite, the appearance on the skin of half a dozen papules, which soon become umbilicated vesicles, and then rapidly form scabs, constitute the whole history of some cases. Here it is that a correct diagnosis is invaluable to the family. To the patient it is of no consequence. He is safe, but he may inoculate any or all of those who have not been properly protected.

COMPLICATIONS.—The most frequent and important complications of variola in children, are inflammations of the mucous membrane of the lower half of the intestinal tube, conjunctivitis, otitis, and different hemorrhages. In a smaller number of cases, attacks of bronchitis, pneumonia, anasarca, articular inflammations, subcutaneous abscesses, simple and pseudo-membranous coryza, angina, and laryngitis, and other eruptive diseases, occur at different periods of the malady.

It is impossible for us, for want of space, to attempt a description of the various symptoms of the different complications just enumerated. Having mentioned the possibility and probability of their occurrence, we must leave the reader with the advice always to suspect the existence or approach of some one of them, when the symptoms, in any case, differ much from those which have been described as characteristic of the regular form.

ANATOMICAL LESIONS.—The characteristic lesions of small-pox are a certain deteriorated state of the blood, congestion of the internal organs, and the inflammation of the skin and mucous membranes constituting the eruption. The blood is found to be entirely liquid and uncoagulable, and of a dark color; or if coagula exist, they are small, soft, and very dark in color. The exceptions to this rule are those in which some acute and severe inflammation exists, under which circumstances the dissolved state of the blood is less marked, and fully formed coagula are more abundant. The congestion referred to affects almost the whole system. The muscles are firm and of a deep-red color; the membranes of the brain are strongly injected, the sinuses are filled with blood, and the cerebral substance presents numerous red points or dots. The vessels of the lungs contain a large quantity of blood, and the liver, spleen, and kidneys are all deeply congested.

The condition of the mucous membranes is important. The pharynx, larynx, and trachea present an eruption, or simple inflammation without eruption. The eruption exists under the aspect of small, circular, thin, and whitish pseudo-membranous points, scattered over the mucous tissue, and slightly adherent to it, beneath which that tissue is often observed to be red and inflamed. At a more advanced degree, and in severer cases, the false membranes have disappeared, and in their places we find circular ulcerations, which are either superficial, or they penetrate the tissue of the mucous coat and rest upon the muscular, or even pierce that and reach to the cartilaginous tissue beneath. In addition to these lesions are found inflammation of the mucous tissue with its consequences, redness, soften-

ing, thickening, and extensive deposits of false membrane, quite distinct from the appearances above described as characteristic of the eruption upon these tissues.

It has been a contested point whether a true vesicular or pustular eruption ever exists upon the mucous lining of the stomach and intestines. The general opinion appears now to be, however, that the changes observed in these organs cannot be ascribed to the formation either of vesicles or pustules. The appearances that have led some observers to consider them as the result of a proper eruption, are the following: The follicles at the commencement and termination of the small intestines, and in rarer cases, of the large intestines also, present an abnormal degree of development, appearing in the form of small hemispherical or pointed, and sometimes flattened projections, on which there often exists a dark, and sometimes depressed central point. At the same time Peyer's glands are often enlarged, more projecting than usual, softened, and red.

According to the valuable researches of Desnos and Huchard (*loc. cit.*), the heart and pericardium present marked lesions in a considerable proportion of cases of confluent variola. These changes were rare in cases of the discrete form, and were not detected in any case of varioloid. The lesions may consist solely of endocarditis, or pericarditis, or these may be associated. These inflammations present the ordinary morbid products, and are not attended with the development of pustules. In other cases the muscular walls of the heart are affected with an acute myocarditis, which is marked at first by a granular state of the muscular fibres, which soon passes into fatty degeneration.

The *anatomy of the variolous pock* is important and interesting. When a vesicle is opened soon after its formation, it is found to contain nothing but a little serosity, which is perfectly limpid and alkaline, while the skin beneath is red, softened, and moist. The umbilicated character depends on a filiform adhesion between the centre of the pock and the surface of the skin beneath. This adhesion is broken, when, at a later period, the pustule becomes globose in shape. The vesicle is also subdivided into several chambers by delicate radiating partitions, so that a single puncture will not discharge the entire contents. About the period of the conversion of the vesicles into pustules, or very soon after the formation of the latter, the cavity of the pock will be found to contain a false membrane, which is of an opaque white color, soft and friable in its texture, and seated upon the derm in small isolated points. After a time these points enlarge, and meeting, unite and form a soft pseudo-membranous disk, uneven upon its surface, and which either fills the pock completely, or is covered at first with serosity and afterwards with pus. This false membrane is secreted originally by the true skin. At a somewhat later period it forms an adhesion to the inner surface of the cuticle, while still later in the progress of the pock, it becomes detached from the cuticle, and remains loose and free in the cavity of the pustule, surrounded by the fluid contents of the latter.

DIAGNOSIS.—The diagnosis of this disease in all its forms ought to be made as early as possible, in order that the persons in contact with the

patient, whether from necessity or by accident, may be vaccinated or re-vaccinated. It is well known that exposure to the mildest varioloid may produce in the unprotected any form of small-pox, from discrete to malignant, according to the constitution of the subject and the type of epidemic prevailing. Therefore the only safety after exposure is in the vaccine disease, and, therefore, the lives of the exposed hang upon the knowledge and action of the physician in charge, a responsibility from which he cannot escape either in the estimate of the public or in his own consciousness.

Dr. Welch concludes from his observations that "vaccination performed at a period less than seven days previous to the appearance of the eruption (small-pox) will not modify the disease," but that when performed, "seven days previous (it) will almost always modify the disease to the extent of rendering it harmless."

Dr. Masson (article on Small-pox, in Reynolds's *System of Medicine*, vol. i, p. 477) says that to be effective vaccination should have gone on to the stage of areola before there is any illness from small-pox. "It has before been stated that when small-pox has been taken into the system there is twelve days freedom from illness, generally, forty-eight hours' illness, and then the disease begins to appear on the skin. The areola of vaccination is not fully formed until the ninth or tenth day of the progress of the vaccine vesicles on those who have never been vaccinated before, so that unless there has been time for the areola to be formed after the vaccination, before the illness produced by small-pox begins, the vaccination will not be of the least benefit." He gives an example: "Suppose an unvaccinated person to inhale the germ of a variola on a Monday; if he be vaccinated as late as the following Wednesday, the vaccination will be in time to prevent the small-pox being developed; if it be put off until Thursday, the small-pox will appear, but will be modified; if the vaccination be delayed until Friday, it will be of no use, it will not have had time to reach the stage of areola, the index of safety, before the illness of small-pox begins. This we have seen over and over again, and know it to be the exact state of the question. Revaccination will have effect two days later than vaccination will have that is performed for the first time, because revaccinated cases reach the stage of areola two or three days sooner than in those persons vaccinated for the first time.

It is plain, therefore, that the diagnosis ought to be made as early as possible. Can it be made in the initial stage? Not, we think, with any certainty. Except in a time of general epidemic prevalence, cases of small-pox are almost unknown, and varioloid is very rare amongst children, and the medical man thinks of anything but varioloid or small-pox to explain a fever attended with vomiting, anorexia, restlessness, or drowsiness in the infant, and the same symptoms with headache and general soreness in the older child. The initial fever has no characteristic phenomena. When the disease is epidemic, the initial fever, as it has been described, may arouse suspicion, and the attendant physician may dare to announce the probable approach of the dreaded disease, and examine all the exposed persons as to their being fully protected. But this course is justified only by the presence of the epidemic. Not until the eruption

begins to appear can the diagnosis be made with certainty ; and however easy it may be for old and experienced physicians to make it then, we desire to caution the younger and more inexperienced as to the possibility of mistake.

The important points to bear in mind are the following ; 1. The prodromic stage, whether of mere ailing and lassitude, such as may not send the patient to bed, or violent fever with nervous symptoms and the different signs which declare a severe disease, which lasts two days, and on the third of which, as the law, the eruption makes its appearance. 2. The eruption appears first on the face and about the upper part of the neck, and consists of hard, distinct, shotty papules, seated, in mild cases, on a nodosal skin. 3. As the eruption appears, the fever diminishes. These three points kept steadily in view will usually prevent any mistake.

The eruption of measles shows itself on the third day of fever, as in small-pox, and occasionally appears in distinct points, which give it a suspicious likeness to that disease. But the attendant catarrhal conditions, the coryza, cough, and conjunctival catarrh, with the fact that the fever increases as the eruption comes out, instead of diminishing as in variola, ought alone to decide between the two. Moreover, a careful study of the eruption ought to enable us to decide. In variolous disease the papules are small, hard, very distinct one from another ; in measles the papular character is not well marked, the stigmata are larger, broader, flatter, and much less hard and shotty to the touch, and very soon they run together and assume irregularly crescentic outlines. By the second day of the disease there is rarely any difficulty.

Varicella, which from its name, one would think, ought to resemble variola closely, has rarely given us any trouble. The prodromic stage of varicella never lasts over a day ; it often consists of but a restless night, and sometimes the first thing to attract the attention of the mother or nurse is the eruption. When the prodromic stage does exist, it consists merely of some lassitude or irritability, loss of appetite, and slight fever. The eruption shows itself at once upon the face and front and back of the body. So much is this the case that we always have the child undressed in order to get a good view of the body. If, on inspection, a number, three or four or a dozen, or very many rounded, projecting, globose vesicles are to be seen, consisting of a thin and transparent layer of the cuticle, filled often to bursting with a limpid serum, there ought to be no difficulty in the diagnosis. Such an eruption, appearing with scarcely a prodrome, or merely a slight ailing of twelve or twenty-four hours, cannot be small-pox or varioloid.

In very mild varioloid, where the eruption counts three or four or half a dozen vesicles, and where the prodromes are very mild, it is not always easy to be quite secure in one's opinion, and a careless or inexperienced person might very well fail to detect the true nature of the disorder. But even here careful inquiry will generally show that the health has been disturbed for two days, at least by altered temper, lassitude, lessened appetite, and one or two restless nights. These prodromes, when followed by a few hard, distinct papules, which become on the second day vesicles, and

then umbilicated pustules, to dry up on the fourth, fifth, or sixth, can be nothing but variolous in their nature.

Again, in severe cases of small-pox itself, embarrassments sometimes occur. We once saw an infant, five weeks old, who had never been out of the mother's room, seized in the midst of perfect health, with violent fever, vomiting, loathing of the breast, and heavy stupor. On the second day of the illness the whole cutaneous surface began to redden; soon the tint became bright red, not unlike some scarlet fevers, but of a more crimson-red; the skin was swollen, tight, hard, and, so to speak, shining. On the third day innumerable hard and distinct papules formed upon this evidently acutely inflamed skin, and on the following day the child died comatose. The child had not been vaccinated, and there were at the time a few cases of varioloid and small-pox in the city. Even in such cases, however, where a deep roseolous or erythematous efflorescence precedes and masks the variolous eruption, the violence of the prodromic symptoms, so unlike the mild phenomena which precede ordinary roseola or erythema, and particularly the intensity of the coloration and the hard and swollen condition of the skin, indicating active inflammatory states of its deeper layers, will go far to prepare an experienced eye for what is coming.

PROGNOSIS.—The fatality of small-pox varies greatly in different epidemics. The result is also markedly influenced by age. It is particularly fatal in infants under one year of age. Of the whole number of cases, 2377, admitted into the Municipal Hospital of this city in 1871-2, there were 35 children under one year of age. Of these 26, or 74.28 per cent., died. Between the ages of 1 and 15 years there were 291 cases, of which 95, or 32.64 per cent., died. The mortality was therefore nearly three-fourths of the whole number under one year, and very nearly a third of those between 1 and 15 years of age.

Of a series of 23 cases that we have met with, 5 were fatal. All of these were under 5, and 3 under 1 year of age.

The amount of the eruption governs the prognosis to a great degree. As the number of pocks is abundant or otherwise,—as the case is a discrete, moderately full, semi-confluent, or confluent one,—so is the danger. Cases of full confluence are almost as fatal as malignant scarlatina. Few children escape in the confluent form. A moderately full eruption, and of course a discrete one, is favorable. The hemorrhagic form is, almost without exception, fatal. Varioloid rarely kills. Under 15 years of age we have never seen a fatal case of it. In one case only have we known it to be dangerous.

The favorable symptoms in any case of variola are the occurrence of the disease in children previously in good health and over one year of age; the absence of any violent nervous symptoms during the initial stage; a proper duration of the first stage; and the subsidence of the fever after the appearance of the eruption. When, in addition to these circumstances, the secondary fever is not too violent, and no complication arises, there is but little doubt that the patient will recover.

The unfavorable symptoms are the occurrence of the disease at a very

early age ; the existence of severe nervous symptoms during the first stage ; the occurrence of a thick and abundant eruption upon the face indicating a probably confluent case ; continuation of the fever after the appearance of the eruption, or a merely slight subsidence of it ; delirium and other nervous symptoms during the secondary fever ; and any irregularity in the appearance of the eruption, as paleness instead of the usual red color, a livid or purplish color of the pustules, imperfect development of the pocks, or their sudden shrinking without diminution of the general symptoms. The occurrence of the signs which mark the hemorrhagic form, as petechiæ and local hemorrhages, stamp the case as almost necessarily fatal. It is scarcely necessary to say that many of these symptoms are indicative of the existence or threatened production of some complication, upon the nature of which must depend, after all, in great measure our prognosis. The complications most apt to occur have already been considered in a previous section.

TREATMENT.—The treatment must be regulated by the type of the case under charge. It will vary, therefore, from a mere quiet expectancy throughout, to the vigorous use of such means as moderate fever, abate nervous agitation, and allay suffering in the early stages, with the peremptory exhibition of stimulants, tonics, and nutritious foods, in the period of eruption and maturation.

In the varioloid of children over eight or ten years of age, during the initial fever, rest in bed, light diet, and the use of sweet spirit of nitre, in iced lemonade, often suffice. Should there be much restlessness, insomnia, or pain, solution of citrate of potash, with small doses of laudanum or paregoric, may be given. When the eruption appears, if it be slight, and the fever disappears, nothing more is necessary than to keep the diet moderate and seclude the patient in one room, for the sake of others, until the crusts have fallen. If the eruption be more copious, enough to cause a good deal of irritation and restlessness, a warm bath at night, especially with some bran added to it, and the application through the day of an ointment of glycerin and cold cream, with a mild opiate at night, are sufficient.

In the variola of unvaccinated children, the treatment must also depend on the type of the symptoms. In the initial stage, when the fever is high, the child must be confined to the breast, if it is still nursing, and, if weaned, it is to be kept upon a proper mixture of milk and water, with lime-water, and upon chicken or beef-tea, for food. Cold water must be frequently offered to the child at all ages, and it should be allowed to take all it desires. A tepid bath morning and evening, or even three times a day, if the child does not resist, is very soothing, and tends to reduce the heat. Spongings with tepid or cool water, from time to time, according to the degree of heat, and the effects of the application, may be used, if the bath terrifies or fails to reduce the fever.

Diaphoretics, and especially the citrate of potash, with sweet spirit of nitre, and very small proportions of laudanum, should be administered in this stage. Or the spirit of Mindererus may be given,—twenty to thirty drops, with ten drops of nitre and five of paregoric, in a tablespoonful of

iced water, every two hours, to children of six months to two years. For older children the doses must be enlarged.

When there is, as often happens, great agitation of the nervous system, as shown by jactitation, insomnia, and mild or active delirium, some remedy should be given to control these symptoms. If the citrate of potash and opium fail to relieve these conditions, the best remedy is bromide of potassium, one to two and a half grains, with one to two drops of deodorized laudanum, at the age of one to three years, every two hours until rest is obtained, or until three or four doses have been given. After the age of four years the proportion of the bromide may be doubled.

When great heat and swelling of the skin, severe headache, and signs of congestion of the lungs or brain, exist, cold applications to the head, with hot mustard foot-baths, may be used with the diaphoretics. If, in older children, the headache or pain in the loins be very severe, a few dry cups or a sinapism may be applied to the back of the neck or loins.

If the bowels are not moved spontaneously, a moderate laxative ought to be used, as syrup of rhubarb or castor oil, or an enema may be ordered. Purging with large doses of cathartics must be avoided at all ages.

In the eruptive stage the treatment must vary with the type of the eruption and the constitutional peculiarities of the patient. It may be laid down as a rule that, the more copious the eruption, the more carefully should the strength be husbanded, and the vitality supported, to enable the patient to pass through the long and exhaustive process of maturation and desiccation necessary to a cure.

If the eruption come out slowly and tardily, and the extremities be cool, even though the body is hot, hot mustard foot-baths, or warm baths, with hot drinks, as milk and water, hot broths, and small quantities of brandy, ought to be employed, and are often very useful.

If the eruption be discrete and moderate in amount, nothing but rest in bed, simple sustaining foods, and some local remedy to allay cutaneous irritation, as an ointment or an occasional warm bath, will be necessary until the secondary fever appears. When this arrives, the same means, in the form of diaphoretics, anodynes, and nervous sedatives, should be used as in the initial stage. In the stage of maturation the strength must be sustained by a diet adapted to each particular case. If the patient be feeble, and therefore much reduced by even a moderate eruption, he must have brandy added to his milk, or wine-whey, from time to time, increased doses of beef or chicken soup, if he can take them, and, if old enough, eggs, or egg-nog. Quinia and muriated tincture of iron should be used as in confluent cases, of which we shall speak directly.

In the semi-confluent and confluent cases all must be done to sustain the strength and vitality. From an early period of the eruptive stage, alcohol, quinia, and iron must be employed. From twenty to thirty drops of brandy, in a wineglassful of milk, may be given every two hours, and two or three tablespoonfuls of thin beef-tea, every alternate two hours, at the age of two or three years. After the age of five, these quantities may be doubled. To infants, brandy, in doses of ten to twenty drops, may be

given every two hours in breast-milk, or in warm water and sugar. Quinia, in doses of half a grain, at a year old, and one grain at four and five years, with or without muriated tincture of iron, ought to be administered every four hours. It is best to choose the four-hour interval, because of the difficulty there is in giving frequent doses to children. If the stomach will not retain the iron and quinia mixed together, the quinia may be used in suppository, two grains every four hours, and the tincture of iron in doses of two to five drops, according to the age, every two hours, in syrup of ginger, or in combination with dilute acetic acid and solution of acetate of ammonia, as proposed in the article on scarlet fever. On account of the well-known frequency of cardiac complications in such cases, digitalis may be added to the treatment if the symptoms indicate marked failure of the heart's action.

The condition of the pharynx and larynx present in confluent small-pox, as described in the article on symptoms, constitutes one of the great difficulties of the disease. The patient suffers so much in the act of swallowing, the respiration is so interfered with when he attempts to drink or eat, that it ends in his taking but little, and, at last, almost nothing. Here ice should be given, iced flaxseed-tea and iced brandy and water, or frozen beef-tea. A solution of chlorate of potash may also be tried. Lemonade may be used, and a warm poultice to the throat is to be recommended. Still we must persevere, as small quantities are better than nothing, and we may employ nutritive injections of beef-tea, of milk, or of egg and milk.

If the patient survives the stage of eruption, we must continue the tonics, stimulants, and nutritious food through the decline of the disease. During the latter period something must be done to allay the itching, burning, and irritation of the skin. If the patient is not too weak, a flaxseed or bran bath is very soothing, or we may use lime-water and sweet-oil liniment, or glycerin and cold cream ointment, applied with a large camel's-hair brush frequently.

In hemorrhagic small-pox, which is almost always fatal, we know nothing better to recommend than the treatment just advised for the confluent form, to which turpentine in full doses may be added, on account of both its stimulant and hæmostatic properties.

Treatment of Complications.—If complications occur in the course of the disease, they must be treated always with a full consideration of the primary importance of the general disorder. The angina and laryngitis of confluent cases can scarcely be looked upon as complications. They belong to the disease. We have already alluded to their treatment, and may refer the reader to what has been said of the same series of symptoms in scarlet fever. In pleurisy or pneumonia we can do nothing better than persevere with the measures most proper to combat fever. Pain may make it necessary to use opium in full doses. Counter-irritation is not to be thought of because of the eruption, and even cataplasms, which are so useful in ordinary pleurisy and pneumonia, are objectionable here.

The treatment of the conjunctivitis which so often threatens, and sometimes occasions great or irreparable injury to the eye, is very important.

Niemeyer says that much may be done to prevent the development of a severe eruption on the conjunctiva by the assiduous employment of cold-water compresses, or, still better, by compresses moistened with a weak solution of corrosive sublimate, one of one grain to six ounces of distilled water. When ulcerations occur upon the cornea, they ought to be touched, if this be practicable, with solid nitrate of silver sharpened to a point, or with a fine camel's-hair pencil which has been moistened and rubbed over the nitrate of silver crystal to insure a caustic solution. When it is impossible to apply the solid caustic or the brush, we must resort to some collyrium. This may consist of a solution of nitrate of silver, a grain to the ounce, or of one or two grains of sulphate of zinc, with twenty or thirty drops of wine of opium, dissolved in an ounce of rose-water, two or three drops of either of which may be introduced into the eye, morning and evening. An excellent collyrium is one composed of twelve grains of borate of soda, one grain of sulphate of zinc, a drachm of camphor-water, to seven drachms of distilled water.

Catarrh of the intestine must be treated by the most careful attention to the diet, by emollient and anodyne injections, and by the internal administration of astringents, and small doses of opiates. When the diarrhoea is severe, and the stools mucous and bloody, we may use with advantage the nitrate of silver by enema, as recommended in the article on enterocolitis.

The treatment of the convalescence is important. The same rules apply here as in other infantile and children's diseases.

Ventilation and Disinfectants.—It is even more important in this disease than in others, for the physician to see to it himself that the rooms occupied by the patient, and the house of which they form a part, shall be well ventilated, and that so soon as the eruption becomes purulent, and its exhalations more or less fetid, proper disinfectants shall be applied. This is necessary, not only for the good of the patient, but also for the safety and comfort of the other inmates of the house. The best ventilation in winter is that procured by an open fire, or, if this cannot be had, by a stove. If the room can be warmed only by a furnace, the windows must be very carefully opened from time to time, so as to supply fresh air, and yet avoid currents flowing over the patient. In summer, of course, the windows must be open.

Among the best disinfectants is Labarraque's solution. If this cannot be had, or if more than one be desired, chloride of lime in saucers, wetted, or a mixture of equal parts of impure sulphate of iron and of chloride of lime, wetted, and placed in saucers, in the entries and passages of the house, are very efficient. Solutions of carbolic acid, or permanganate of potassa, chloral, and other disinfectants may be substituted for the above if more convenient.

Before terminating our remarks upon the subject of small-pox, it will be proper to give some account of the treatment of the eruption which has been recommended and practiced, with a view to prevent the scarring and disfiguration which so often result from the ravages of the disease. Of the different means that have been employed with this view, there are two

which are chiefly relied upon at present. One is to cauterize the pustules with nitrate of silver, and the other to make a mercurial application upon the part where it is desirable to cause the abortion of the eruption. The cauterization has been performed in two modes; by the application of the caustic to each pustule separately, or to masses of the eruption without puncturing the cuticle. It appears, however, that the first-named method is much the most preferable. To succeed perfectly, it is necessary to touch the derm forming the base of the pustule; so that the best plan is to remove or lift up a portion of the top of the vesicle with a lancet, and then to introduce into its interior the sharpened point of a stick of caustic. This operation is certainly successful only when performed on the first or second day of the eruption, through MM. Rilliet and Barthez have known it to answer as late as the third and fourth, or even fifth day. The process of cauterization is productive of acute pain, but does not increase the local inflammation, according to the authors just quoted, at least when applied to a small number of the pocks. They state that when applied to the pustules seated upon the edges of the eyelids, it is almost incredible to behold how great is the diminution of the œdema of those parts in a single day. The conclusion of these gentlemen is, that individual cauterization of the pustules with nitrate of silver does certainly cause them, as well as the surrounding tumefaction, to abort, and prevents them from leaving cicatrices.

This plan is, however, manifestly inapplicable to any but cases of the discrete form, where the vesicles are not very numerous.

The other method which has been employed to cause the abortion of the pustules and thus prevent disfiguration, is, as has been stated, the application of some one of the mercurial preparations. The effects of this treatment are said to be an almost certain arrest of the development of the eruption, when it is used from the first or second, or not after the third day; the vesicles and pustules remaining small and isolated, and not assuming, or else soon losing the umbilicated character. When applied early, while there are as yet but few vesicles formed, it prevents the development of new ones, and diminishes the accompanying swelling and soreness. When the application is removed on the seventh or eighth day, it is found that desiccation has occurred imperfectly, the surface presenting small soft scabs, or little whitish, soft elevations, consisting of the pseudo-membranous substance situated between the true skin and the new epidermis, the old cuticle having generally peeled off with the plaster. In some places a light rose-colored surface alone remains.

In regard to the success of this treatment in preventing disfiguration, we may quote the statement of MM. Rilliet and Barthez, that none of the patients upon whom they saw it tried presented any cicatrices, though several had had confluent small-pox, which pursued its usual course on the parts not covered by the application. Dr. Stewardson, of this city, made a considerable number of trials of this treatment at the Small-pox Hospital of this city in 1841-42. He gave his conclusions in the following words (*Am. Jour. Med. Sci.*, January, 1843, pp. 86-7): "From these experiments, it seems pretty evident that the mercurial plaster has a decided

influence upon the small-pox pustules, preventing more or less completely their perfect maturation, and diminishing the concomitant swelling and soreness, the process of desiccation being completed without the formation of thick scabs, and the resulting cicatrices less marked than when the process of suppuration was left to pursue its natural course. . . . That, by its use, pitting may be entirely prevented, or the mortality from small-pox materially lessened, seems to me very doubtful, although had all the precautions above mentioned been taken, it is not improbable that the effects would have been still more decided."

The use of the mercurial application is attended with some inconvenience. In the first place it is difficult to keep it accurately applied, particularly in children, in consequence of the unpleasant sensations it occasions. In the second place, it not very unfrequently, according to MM. Rilliet and Barthez, produces an eruption of hydrargyriasis, or mercurial roseola, in about eight or fourteen days after the variolous eruption, or four or ten after the application of the remedy. M. Rayer, however, states this effect to be a rare one.

Dr. Stewardson says that he thinks no apprehension need be felt as to constitutional affection from the mercury, for scarcely ever were the gums even touched. One of ourselves, however, when in Paris, in 1840, saw this effect produced in a young girl at the Children's Hospital.

The method of its application is different in different hands. The French generally employ the *emplastrum de Vigo cum mercurio*. Dr. Stewardson prefers the strong mercurial ointment, either pure or rubbed down with an equal bulk of lard, spread upon a piece of thick muslin. The muslin is to be cut into the shape of a mask, with apertures for the eyes, nose, and mouth. It is secured upon the face by means of strings attached to its margin and tied across the back of the head and neck. It is important always for the success of the measure, that the application should be kept in close contact with the skin. To insure this, he employed a separate piece of muslin for the nose, which is the part most difficult to fit. With the same view, the French authors recommend that the plaster should be cut in pieces to suit the different portions of the face, making one for the forehead, and others for the cheeks, sides and back of the nose, and upper and lower lips. Any spaces that may remain are to be covered with other portions of the plaster, and the whole secured with strips of diachylon. On account of the difficulty of applying the mercurial plaster, the following ointment was compounded by the apothecary of the Children's Hospital at Paris, and has been found to answer very well:

R. Mercurial Ointment,	24 parts.
Yellow Wax,	10 parts.
Black Pitch,	6 parts.—Mix.

The application ought to be confined to the face, as that is the part which it is most important to save from disfiguration, and as it is better not to use it upon a larger surface than necessary, lest it might occasion the mercurial roseola, or possibly salivation. As a general rule, four or

five days are sufficient, according to Guersant and Blache, to leave it in contact with the skin, in order to avoid the bad effects just referred to.

The object sought in these applications being, to a great extent, to protect the vesicles from contact with the atmosphere, it has been advised to paint a saturated solution of gutta-percha in chloroform, over the neck and face, so soon as the papular eruption is fully out. This plan was tried in five of our own cases (*loc. cit.*, p. 345), two of which were discrete and three confluent, and with very satisfactory results.

To conclude this matter we will add that Niemeyer states that Skoda prefers compresses moistened with solution of corrosive sublimate (gr. ij-iv to water \bar{z} vj) to mercurial plaster, which induces an injurious elevation of temperature. He also says that Hebra rejects both mercurial plaster and solution of corrosive sublimate, as well as collodion, and touching the individual pocks with nitrate of silver, and that he has come to this decision from the observation in his wards, that the pocks do not leave cicatrices any oftener since he has ceased to employ these remedies than when he used them. He (Hebra) applies only cold-water compresses, which while the skin is tense, relieve the patient, although they do not protect the skin from destruction.

ARTICLE III.

VACCINIA.

DEFINITION ; SYNONYMS ; HISTORY.—The vaccine disease is an affection produced by the inoculation of the virus of variola, modified by passing through the system of the cow.

The proofs which exist as to the truly variolous nature of the vaccine disease in the cow, are altogether incontestable ; so that we must regard the vaccine disease in the human subject merely as a remarkably modified form of variola.

It is susceptible of propagation from individual to individual by inoculation, but is contagious in no other way, and it possesses the invaluable quality of protecting, with very great, though not with absolute certainty, those through whom it has passed, against small-pox.

Besides the name given above, it is known by the titles of cow-pox, kine-pock, vaccina, and vaccinia.

Some knowledge of the nature of the vaccine disease, and of its power to protect the human constitution against small-pox, has been found to have existed in different parts of the world, but there can be no doubt that we owe to the genius and patient research of Dr. Jenner the inestimable blessing of vaccination, since it was by him that its marvellous virtue was demonstrated and proclaimed to the world. Dr. Jenner learned, at an early period of his life, that there existed a popular belief in Gloucestershire, England, that persons who had contracted a peculiar vesicular disease from the udder of the cow, were thereby protected from the attack of

small-pox. Becoming convinced by a long course of patient observation, that this belief was founded in fact, he determined at last to try whether the disease might not be transmitted from one person to another, and thus increase immeasurably the utility of this wonderful protective means. On the 14th of May, 1796, accordingly, he vaccinated a child eight years old with matter taken from the hands of a milker who had received the disease from the cow. The experiment succeeded perfectly, the child having received and passed through the disorder in the most satisfactory manner. On the 1st of July following, this child was inoculated with variolous matter, and resisted the contagion entirely, as Dr. Jenner had expected. It was not, however, until two years later, in 1798, after additional experiments, that the results of his researches were published to the world. From this time the belief in the utility of vaccination and its application in practice spread rapidly throughout England. In 1799 it was introduced into this country; in 1800 it reached France, and in the course of a very few years extended to all civilized nations.

SYMPTOMS ; COURSE.—It is very important for the physician to be thoroughly acquainted with the appearances presented by the vaccine disease in its various stages, since he is to judge by those appearances whether the subject has had the disease in such perfection as to derive all the benefit from its protective power which it is possible for it to impart.

The first effect of the puncture by which the virus is introduced into the tissues, is to produce a very slight redness at the point where the operation is performed. This redness usually disappears within twenty-four hours, and there is left merely a little mark or scab at the point of insertion. On the third day after the operation we first begin to perceive the specific effects of the virus, in the shape of a small, hardened point at the seat of the wound, surrounded by a faint, erythematous redness. Over this hardened point, which grows gradually larger, the cuticle is elevated on the fifth day into a vesicle, by a thin, transparent, and pearl-colored serous exudation. This vesicle soon becomes umbilicated, so that by the following day, the sixth, the depression in the centre, constituting the umbilicated character, is generally perfectly manifest, and at the same time the vesicle is surrounded by a very narrow ring of inflammation. The vesicle continues to increase in size, until on the eight or ninth day it has reached its highest degree of development. At this stage the vesicle or pock is large, usually about one-third of an inch in diameter, and it projects very considerably above the general surface. Its shape is circular, as a general rule, though not unfrequently it is oval, this depending apparently upon the mode in which the puncture has been made. The color of the pock is dull white or pearly, or sometimes it has a yellowish tint. The quantity of fluid contained in the cavity of the vesicle differs, of course, according to its size. The structure of the pock is found, upon careful examination at this time, to be cellular, the number of cells amounting commonly to eight or ten; very often there is a small, dark-colored scab on the very centre of the vesicle, even at this period, though in other instances this is absent, the surface of the vesicle being formed exclusively of thin and transparent cuticle. The scab just alluded to has

seemed to us to consist of the little incrustation, formed at the point where we had introduced the virus by the drying up of the minute quantity of blood escaping after the puncture, and of the dissolved virus which had not been absorbed. We have often noticed that when the small scab just alluded to has been rubbed off the arm on the second day, the vesicle has presented no scab as early as the eighth day. On the eighth day the little ring of redness at the base of the pock, which has hitherto been very small and narrow, begins to enlarge so as to form the areola. This increases during the ninth and tenth days, forming a brilliant scarlet or dark-red inflammatory circle of about two inches in diameter, and constituting one of the most strongly marked features of the vaccine disease. The color of the ring is most intense at the edge of the vesicle, and then fades gradually to its outermost boundary. On the ninth and tenth days, in connection with the areola, the skin and cellular tissue on which the vesicle is seated, and that for a short distance beyond the margin of the latter, become hardened and tumefied, forming a solid knot or lump in the derm, like the base of a furuncle. The inflammation which causes the areola is often so intense as to occasion the production of vesicles, which are almost always discoverable with the aid of a lens, and are sometimes distinctly visible to the naked eye. On the tenth day the disease is usually at its height, and it is then, of course, that all its peculiar characteristics are most strongly marked. At this time the child, when of an age to describe its sensations, will often complain of heat, itching, and pain in the inflamed spot; the arm is heavy and not willingly moved, or it is moved with care and caution; there is, in a good many instances, some irritation and swelling of the axillary glands, and very frequently a decided febrile reaction may be noticed. In other cases, on the contrary, none of these symptoms will be present. The child is gay and cheerful, its movements are free, quick, and unembarrassed, and it seems in all respects to be in its ordinary condition of health.

From the tenth day the disease begins to subside. The areola fades so as to have nearly disappeared by the fourteenth day; the fluid contained in the vesicle is gradually converted into pus, and the cellular structure of the pock is broken down so as to form, by the thirteenth day, but a single cavity, in which the pus is contained; the process of desiccation is going on rapidly during this time, so that about the fourteenth day the vesicle has disappeared, and in its place there is a firm, hard scab, of the shape and size of the vesicle. This scab continues to harden for some days longer, and at the same time contracts somewhat in size and grows darker in color, until at last it is of a very dark-brown or mahogany tint. It separates gradually from the tissues beneath, the separation beginning at the circumference, and falls off usually about the eighteenth or twenty-first day, leaving beneath a small ulcer, which soon heals, or else a cicatrix of the shape and size of the pock. The cicatrix is at first of a deep-red or purple color, but fades gradually, until it becomes much whiter than the surrounding skin. The scar left by the vaccine disease is very characteristic, and is often, though not by any means invariably, indelible. To be at all depended on as a mark of the disease, the scar should be small,

circular, of a smooth and somewhat shining appearance, and it should exhibit radiations and little depressions or pits. The depressions are supposed to have been caused by the cells constituting the pock in its early period.

There is rarely more than a very slight constitutional disturbance attendant upon the course of this disease. About the eighth day, a decided febrile reaction, attended with some unusual warmth of the surface, restlessness at night, and fretfulness of the temper, is often observed. In a few instances we have noticed distinct disturbance of the health about the third and fourth days; amounting only, however, to unusual irritability and discomfort through the day, and to wakefulness or disturbed sleep at night.

IRREGULARITIES AND ANOMALIES.—We have now described the regular course of a vaccination—that which it pursues in a large majority of the cases. Certain variations from the above standard or typical course are frequently, however, met with, and require some notice. These variations may consist merely in the degree of severity of the local and general symptoms, or in the appearances presented by the pock, without affecting at all the validity of the disease; or they may concern the duration of the phenomena; or, lastly they may be such as to call in question the validity of the disease, leaving us in some doubt as to whether it has protected the constitution against variolous attacks or not.

The severity of the local inflammation occasioned by the vaccination, and that of the general symptoms also, varies often to a considerable extent. In some instances, and especially when the virus employed has been procured recently from the cow, the specific inflammation proves very severe. We have seen the arm intensely red, and very considerably swollen, from the shoulder to an inch below the elbow, while at the same time the axillary glands were tumefied and tender, and the child very feverish and uncomfortable. This happened in three children, in all of whom we had employed the same virus; which, as we afterwards learned of the person from whom we obtained it, had been taken quite recently from the cow. It produced the same violent inflammation, moreover, in several other subjects in whom it was employed. This, however, is not to be regarded as by any means a usual occurrence when bovine virus has been employed, since now that we very frequently vaccinate with lymph directly from the cow, it is rarely that we observe any severe inflammation.

If the vesicle happens to be broken by accident soon after its formation, its appearances during the subsequent progress of the disorder will often be very different from those exhibited in subjects in whom no such accident occurs. The vesicle loses a portion of its contents; it becomes conoidal and irregular in shape, instead of being circular and umbilicated; it does not exhibit the pearly white and diaphanous color which belongs to it, but is yellowish and opaque; the areola is often premature and irregular in shape, and the scab is frequently small, uneven on the edges, and falls off at an unusually early period.

Occasionally there is observed in the course of cow-pox a papular erup-

tion over the body of the child. This occurs usually between the ninth and twelfth days.

It is quite common for the disease to be retarded in its progress. The delay generally takes place in the appearance of the vesicle, this not showing itself until the sixth or eighth day, or, in some rare instances, not until the sixteenth, or even the twentieth, or forty-sixth day. The longest retardation that we have met with has been seven days. In this kind of retardation, the disease usually runs through its regular and natural phases after the vesicle has once made its appearance. In another kind of retardation the delay occurs in the vesicular and pustular stages of the affection, the papule appearing at the ordinary time, but the disease not reaching its height or maturity until the eleventh or twelfth day.

The forms of variation from the ordinary course of cow-pox just described, do not seem to be connected with any diminution in the protective power of the disease.

It sometimes happens that the operation of vaccination gives rise to a disease totally unlike the true vaccine disease, one which does not protect against small-pox, and which has therefore been called *spurious vaccine disease*.

It was formerly the custom to describe quite a variety of appearances as indicating with greater or less probability a spurious disease. Of late years, however, it is generally admitted that the spurious pock is of much less frequent occurrence than was at one time supposed, and that, when it does occur, its characters are so marked as to make it easy of recognition. In fact, it happens in a very large majority of cases, that the vaccination either fails entirely, the puncture being productive of no other results than those which would naturally flow from a slight wound of the skin, or else that it is followed by a true and easily recognized vaccine pock.

When, however, the operation is followed immediately or within a day or two days by inflammation, and the appearance of a pustule, without the previous production of a vesicle; when this pustule is irregular in shape, yellow in color, acuminate, easily broken, and terminating in a soft, yellowish, ragged-looking crust, which falls off upon the fifth, sixth, or seventh day, there is assuredly reason enough to call the vaccination spurious, and it becomes the imperative duty of the practitioner to regard it as such until subsequent and repeated trials with other and fresh virus, have proved the child to be protected.

DIAGNOSIS.—There can be no difficulty whatever in distinguishing the vaccine disease when it occurs in its regular form. The successive phases through which the eruption passes, and the particular appearances which it presents in each stage, are so unlike all other diseases, except, indeed, small-pox, as to render it very easy of recognition.

Sometimes, however, there is a little difficulty in determining whether the eruption is spurious or regular. But this rarely happens except under circumstances in which we should expect some modification in the phenomena of the disease, to wit, when its course is interfered with by the effects of a previous vaccination, or of an attack of variola. The irregularities arising from these causes are such as might be anticipated, and

will be described in the article on revaccination. Whenever the disease fails, in any important respect, to exhibit the perfect attributes of a well-marked pock, both as regards its time of development, its changes, and its particular appearances at each stage, in a child not previously vaccinated, nor having had small-pox, the only wise and prudent plan to follow is to repeat the operation a few weeks after the doubtful one, so as to test thereby the protective power of the first.

PROTECTIVE POWERS.—Though vaccination in infancy has not proved a sovereign protection against small-pox, as was at first hoped and expected, the security it does afford, when properly used, against one of the most loathsome and dangerous of disease, is so nearly perfect that the thought of its benevolent power ought to rouse every feeling of thankfulness of which the human heart is capable. It has come to pass within a few years, here and there in the world, and we know this was the case in Philadelphia, that some persons have begun to question the real value of vaccination. Such persons always seemed to us the most crotchety and foolish of mankind, and since small-pox exhibited its powers here, as it did in the epidemic of 1871-2, we imagine those very persons are quite ready once again to thank Providence for its great boon, and to do true homage to the great discoverers of vaccination.

As to the protective power of vaccination, we have had abundant proof, in our own experience alone, to satisfy us that this is complete when it is properly applied. We have never seen life lost or the face disfigured, during forty years of experience, in any one who had been well vaccinated in infancy, and then *successfully* revaccinated at puberty. We had never seen a fatal case of small-pox in a subject under 43 years of age, who had been well vaccinated in infancy, until the late epidemic, though we had seen two who had it severely enough to pock-mark them. We knew that such cases occurred, but none had occurred in our own practice; and our experience in the late terrible epidemic has but confirmed our faith in the powers of vaccination. During its prevalence we saw no severe nor dangerous varioloid or variola in children under 10 and 12 years of age. It was not except among those over 15 and 20 years of age that we began to see and hear of dangerous cases of the disease; and after successful revaccination, even in those most exposed, we saw not a case even of varioloid, much less of severe variola. We could, had we the space, cite particular instances in our own practice in proof of the absolute protection afforded by revaccination, but deem it best to give some facts illustrative of this power from the hospital experiences of the late epidemic.

In the report made to the Board of Health, of this city, by Dr. Welch, of the Municipal Hospital, during the epidemic of 1871-2, are some facts which show most strikingly the power of vaccination. At page 9 are given the following cases:

“Case I. Child, *æt.* two years; vaccinated in infancy; two good cicatrices; came in with mother, who had small-pox; sixteen days in hospital; no disease.

“No. VI. Infant, *æt.* 10 months; not vaccinated; admitted February 10th, along with its mother, who had varioloid, and from whose

breast it was nursing; vaccinated same day. February 16th.—Two convulsions. 17th.—Perfectly well again; vaccination taking well; fourteen (14) days in hospital; no disease. (This child returned to the hospital with measles.)

“No. IX. Child, æt. 7 years; vaccinated six months ago; fair cicatrix; eleven (11) days in hospital; no disease.

“No. X. Child, æt. 8 years; vaccinated six months ago; fair cicatrix; eleven (11) days in hospital; no disease.”

At page 12 Dr. Welch states another very interesting fact, which coincides with the experience of the London Small-pox Hospital. He says: “In this connection we might add that the physician in charge, his two assistants, the matron,—who has been connected with the hospital for twenty-four years,—the chief male nurse, and a number of others employed at the hospital during the epidemic, were protected only by vaccination and revaccination. Indeed, not a single person connected with the hospital, who had been revaccinated, contracted the disease; while, on the other hand, some three or four of the nurses, who had been affected by small-pox previously, took the disease a second time.”

How any one can read such facts as these, and they might be indefinitely increased, and yet refuse a child the boon of vaccination, is beyond our comprehension.

In former editions of this work we endeavored to show the necessity and propriety of revaccination. Hereafter we shall advocate revaccination in all cases, no matter how perfect the first vaccination may be stated to have been, or how perfect the cicatrix or cicatrices. At the age of fifteen, or as soon afterwards as possible, all young persons ought to be revaccinated. There should be no waiting for an epidemic or for direct exposure to infection. The operation ought to be preformed as regularly as the primary vaccination.

There is now a host of evidence on this point, but that which is given by Dr. Welch, in the report just quoted, of facts demonstrated by the late epidemic in this city will be sufficient.

“With reference to the practical efficacy of revaccination,” he says, “the hospital record shows as follows: Among 2377 cases of small-pox admitted during the epidemic, only 36 are said to have been revaccinated, of which 4 died. But by subjecting these cases to a careful analysis, we find as follows: Seventeen (17) were revaccinated at a distant period, some as far back as thirty-one (31) years; five (5) had not been revaccinated until after exposure; seven (7) were said to have been successfully revaccinated, but were unable to exhibit any cicatrices as the result; sixteen (16) bore upon their arms very poor and uncharacteristic scars, some of which, indeed, were scarcely visible; five (5) presented fair cicatrices; and only three (3) were able to show good cicatrices.

“Of the four (4) who died, two (2) occurred among those without cicatrices, one among those revaccinated after exposure, and one among those showing poor and uncharacteristic scars.

“All the cases which bore upon their arms unmistakable evidence of successful revaccination, suffered from the mildest form possible of the dis-

ease. Indeed, three (3) of these cases exhibited an eruption of doubtful character, and have therefore been recorded as cases of varioloid (?). The eruption on three (3) others did not advance beyond the papular stage, and on seven (7) it was barely vesicular."

It is unnecessary to add anything more as to the protective power of the vaccine disease against small-pox. Those who are not convinced by such facts as these, would not believe one though he rose from the dead.

PERIOD OF PERFORMANCE.—The period usually chosen for the performance of this operation, is soon after the age of three months. If, however, the infant be exposed to the contagion of variola, it is necessary to perform it immediately, even upon the first day of life; and in such cases the protective power is as perfect, and the local or constitutional irritation little greater, than when the operation has been deferred to the usual time.

SUSCEPTIBILITY TO THE DISEASE.—The susceptibility to the vaccine disease varies greatly in different persons and different families, and is modified to a greater or less extent by the existence of other diseases in the individual at the moment of the operation. In some it is said never to be received, no matter how frequently or how carefully the virus may be inserted. In others it is received with difficulty, requiring several repetitions of the operation before it can be made to take; whilst in yet another class of subjects, the smallest amount of virus, when inserted in a careless and imperfect manner even, will produce the disease with the greatest certainty. Nevertheless a large majority of children take the disease after a single operation, if this be preformed with ordinary care and nicety. No explanation of the different susceptibilities of individuals to the disease can be given. The same difference is known to exist in regard to other contagious and even epidemic diseases, as measles, scarlatina, pertussis, variola itself, typhoid fever, and cholera.

The susceptibility varies also in the same person at different times, without its being possible to ascribe this fact to any evident cause, since the child may appear on both occasions to be in the same condition as to health and other circumstances likely to influence its susceptibility to the contagion. Thus, we knew a child a few months old to be vaccinated four times, twice by the late Dr. C. D. Meigs and twice by one of ourselves, each operation following rapidly the preceding one, without success, though the virus was known to be good from its having succeeded in other subjects, and though it was changed each time. The child appeared to be in perfect health. There was no eruption of any kind upon its surface, nor and other condition that could explain its insusceptibility. After the fourth operation, the attempt was suspended for about four months, then renewed, and with instant and entire success. In another case, the varying susceptibility of the same individual to the disease was still more strikingly exemplified. An infant, a few months old, was vaccinated four times in succession from the scab without success. It was then vaccinated with fresh lymph taken from the arm of an infant who was undergoing the disease. This also failed. A few weeks after this, the operation was

again performed with the dried scab, and this time with perfect success. This same experience has more recently occurred to us in a case where vaccination was performed four times at short intervals with fresh bovine virus without success, but on a fifth attempt complete success was obtained.

Certain eruptions existing previously upon the surface, have seemed to us to prevent the reception of a vaccination. The eczematous and impetiginous diseases of infancy and childhood have certainly had this effect in our experience, though M. Taupin (*Dict. de Médecine*, t. xxx., p. 406) is of the contrary opinion; he having found that the disease has been merely retarded when the operation was performed during the initial stage of the eruptive fevers, whilst its course was suspended even entirely when any of these affections occurred in a child already vaccinated, to be resumed again after the cure of the eruptive fever.

There is another circumstance concerning the supposed effects of other diseases on the vaccine affection, to which it will be well to draw attention. We are sure there are few practitioners, having any considerable amount of business, but must have been annoyed, and injured perhaps in their reputations, by the notion so prevalent in the community that vaccination may impart to children other diseases. This prejudice exists particularly in regard to the chronic cutaneous eruptions of infancy and childhood, so that we have frequently had parents to insist to us that the impetiginous or eczematous disease under which their child might be laboring, has been caused by the vaccination, performed perhaps recently, or even months before. M. Taupin, quoted by MM. Guersant and Blache (*Dict. de Méd.*, t. xxx., p. 414), vaccinated a large number of children at the Children's Hospital in Paris, with virus taken from subjects affected with itch, scarlatina, measles, varicella, varioloid and variola, rachitis, scrofula, tuberculosis, chronic eruptions of the scalp, darts, etc., without communicating to the patient any of these affections, either those of acknowledged contagious or non-contagious nature. A very curious case illustrative of this point is mentioned by Dr. Gregory in his *Lectures on the Eruptive Fevers* (Am. ed., New York, p. 270). "A child, who had been exposed to the infection of small-pox, was vaccinated. Both diseases advanced. A lancet charged with lymph from the vaccine vesicle produced cow-pox. Another lancet charged with matter from a *variolous* pustule, *formed within the vaccine areola*, communicated small-pox." We mention the result of these experiments in order to show how little foundation there is for the popular notion above alluded to, and to give to the practitioner an argument with which to defend himself against the unjust accusations of those who may assert his vaccination to have been the cause of any disorder that may have followed upon it. Not that we would ourselves employ virus taken from a child suffering from disease of any kind whatsoever, since this is, to say the least, unnecessary, and ought to be avoided. Indeed, we have never employed a vaccine crust taken from a child who was not apparently in perfect health. The smallest amount of cutaneous eruption upon a child has always been sufficient reason with us to reject

the virus afforded by such a patient, and as this *must* be the safest plan to adopt, it is of course the proper one.

The still more serious charge has, of recent years, been made against vaccination, that it may be the means of transmitting constitutional syphilis. And there are well-authenticated cases in which the operation has undoubtedly been followed by this terrible result. In every instance, however, so far as we are aware, in which the exact mode of the vaccination could be ascertained, it has been found either that the child from whom the virus was obtained, presented at the time evidences of constitutional syphilis, or that the virus had been impure, being mixed with blood or pus, which may have been the medium of infection. There is, indeed, no evidence whatever to show that the lymph or crust derived from a typical vaccine eruption, in an apparently healthy child, can possibly be the means of transmitting any constitutional disease. It is more prudent, however, that if the lymph be used, it should not be taken after the eighth day of the existence of the vesicles; and that in obtaining it, all hemorrhage should be avoided.

Mr. Jonathan Hutchinson (*Med.-Chirurg. Transactions*, for 1871) gives two series of cases which show the possibility of communicating syphilis by means of vaccination. At page 322 he states his belief that the blood is the source of the contamination. He says: "There can, I think, be little doubt that in this instance it was the blood, and not the vaccine lymph, which was the source of contamination." At page 325 he quotes, from a previous report, the following, amongst other conclusions: "That the blood of a child suffering from inherited syphilis can, if inoculated, transmit the disease with great certainty.

"That it is quite possible for vaccine lymph and blood to be transferred at the same time, and for each to produce its specific results, the effects of the syphilitic inoculation occurring subsequently to those of vaccination.

"That it is quite possible to vaccinate successfully from a syphilitic infant in the stage of the utmost potency as regards its blood, without communicating syphilis."

In regard to this most important point we have two statements to make; that we have never had occasion to suspect even that we have been the unfortunate instruments in communicating this disease in our own practice, and that we have always used the dried scab. Is this happy exemption from such an accident the result of care in selecting the virus, or does it depend on our constant use of the dried scab? Is not the danger of having blood intermixed with the lymph much greater, when the vesicle is opened by the surgeon on the eighth day, than when the lymph is left to dry and form a scab in the natural mode?

Still as the danger of communicating syphilis by badly selected vaccine virus does exist, it is important to be aware that recently there has been introduced into the American market a supply of lymph directly taken from the cow. This is furnished in the form of quill slips, one end of which is charged with the lymph: and they can be constantly obtained fresh in our larger cities, as the supply is replenished every day or two.

Whenever it is impossible to obtain perfectly satisfactory humanized virus, either lymph or crust, the bovine virus above mentioned should invariably be used. Indeed, of late we have been more and more in the habit of employing it on account of its convenience and reliability. The more severe local inflammation which was formerly thought to attend the use of lymph directly or only a few removes from the cow, is not found to follow the employment of these slips to any objectionable degree.

OPERATION.—Under this head we shall consider several important points: the relative value of the dried scab and fresh lymph; the question as to whether it is best to raise more than one vesicle by more than one insertion of the virus; and the various modes of performing the operation.

In this city it has been the custom for many years past to use the dried scab, and to raise, as a rule, but one vesicle. After an experience, extending, in the case of one of us, over forty years, during which we have never used anything but the crust, and have rarely made more than one insertion, we can aver that we have never known any one to die of small-pox who had been successfully vaccinated and then successfully revaccinated by this method. We have seen a good many mild varioloids, in subjects that had not been revaccinated, from the ages of twelve and fifteen upwards, but only in two cases have we known the disease to be severe enough to pock-mark the patient. We know of but one death from small-pox in our own circle of patients. This occurred in a gentleman 43 years of age, who was originally vaccinated by the late Dr. C. D. Meigs, and who was never revaccinated until four or five days after he had been exposed directly to the small-pox infection. The operation came too late. Though the puncture took, he died of hemorrhagic small-pox of a virulent form.

In using the crust we have always taken great care to select only those from the most healthy children. Any blemish upon the skin, any shadow of doubt as to the perfection of the vaccine disease, ought always to cause the rejection of the crust.

The scab is less certain to take the first time than the fresh lymph, but it can always be made to take by perseverance, and we confess that it is hard for us to understand why the vaccine disease, if it be perfect in all its stages and phenomena, is not as much a vaccine disease when it springs from the crust as when it proceeds from fresh lymph, and therefore as competent to affect the economy through which it passes according to its natural law.

If the crust is to be used, it ought to be as fresh as possible, to insure its taking at the first operation. When the physician is obliged to keep it for several weeks, he should preserve it in some close receptacle, as between glasses, in tin-foil, oiled paper, between two pieces of wax, or in hermetically closed glass vials.

If the fresh lymph is preferred, the children to be vaccinated should be collected together about the vaccinifer (the child from whom it is to be taken) on the eighth day of the disease. The vesicle must be very carefully opened, so as to avoid wounding the true derm, and thereby causing

any effusion of blood, and the lymph conveyed on a lancet directly from arm to arm.

It is proper to say that this is the mode of vaccinating usually preferred in Europe as the most certain and successful.

Though we have stated that, in our own practice in this city, one thoroughly characteristic vesicle at the primary vaccination, and one again at the vaccination, has been entirely successful in securing complete protection against small-pox, the opinion is held abroad that more than one vesicle gives greater security, in the event of small-pox attacking the vaccinated, than a single one. This opinion, which is based upon very numerous observations in England and Germany, is so strong and positive that we think it best to advise hereafter that at least two insertions, so as to raise two vesicles, shall be made in this country. Any one who wishes to study this question may refer to an excellent article on Vaccination, by Dr. Edward Cator Seaton, in *Reynolds's System of Medicine*, vol. i, page 483, where the whole subject is fully discussed. At page 499 Dr. Seaton insists that it is the duty of the physician to produce four or five genuine good-sized vesicles.¹

It is proper to put before the reader this opinion of so able an authority as Dr. Seaton, so that any one who feels bound by such authority may follow his rule. For ourselves, we can only repeat that thus far in our own experience, one thoroughly good primary vaccination, and a second characteristic vesicle obtained at the revaccination, have been entirely successful and sufficient. In obedience, however, to the facts collected in England and Germany, we shall, hereafter, as stated above, advise the raising of at least two good vesicles at each vaccination. It makes but little difference whether the two be raised on one arm, or one on each. For the convenience of handling the child, we think it will be best to make the two insertions on one arm.

We think it the duty of the physician who vaccinates a child always to see to it himself that the result is a perfect vaccine disease. This matter

¹ The protective power of vaccination, as well as the influence exerted by the perfection and the number of the insertions, as shown by the cicatrices, is remarkably well exhibited in the following table quoted by Dr. Seaton (Art. Vaccination, in *Reynolds's Syst. of Med.*, vol. i, p. 499), from Mr. Marson. The table is based upon 15,000 cases. Of these it was found that the unvaccinated died at the rate of 37 per cent., and the vaccinated at the rate of only $6\frac{1}{2}$ per cent.

Classification of Patients affected with Small-pox.	Number of Deaths per cent. in each class respectively.
1. Unvaccinated,	37.
2. Stated to have been vaccinated, but having no cicatrix,	23.57
3. Vaccinated :	
a. Having one vaccine cicatrix,	7.73
b. Having two vaccine cicatrices,	4.70
c. Having three vaccine cicatrices,	1.95
d. Having four or more vaccine cicatrices,	0.55
a. Having well-marked cicatrices,	2.52
β. Having badly-marked cicatrices,	8.82
4. Having previously had small-pox,	19.

is too important to be trusted to any inexperienced person. The physician has not done his duty who trusts to anything but his own eye as to the genuineness of the vesicle which results from his operation. He should examine it himself on the eighth or ninth day of the disease. The special characters of the disease have already been fully described.

It would be well, too, that physicians in charge of families should examine the cicatrices which follow vaccination, and if they fail to present the characters which belong to successful operations, he ought to repeat the vaccination. Dr. Welch, whose experience in this matter was large, says that a good cicatrix is one "with a well-defined margin, slightly excavated, and reticulated or honeycombed." What he classifies as a fair cicatrix presents the same characteristics, but to a less marked degree, and poor ones are those "pointed out as the result of vaccination, but which are so indistinct or uncharacteristic as to make it difficult, and sometimes even impossible, to recognize them as vaccine scars." In case any practitioner should meet with the latter in a family he may be attending, he ought, we think, to urge upon the parents the necessity of repeating the operation at once.

REVACCINATION.—We think few physicians or laymen who watched the violent epidemic of small-pox which prevailed in this city during 1871 and 1872, can doubt as to the necessity of revaccination. So convinced are we by what we saw during that epidemic of this necessity, that we shall hereafter advocate the repetition of the operation at the age of puberty as a matter of domestic habit and law, a matter to be attended to by the heads of families with the same regularity and care that is now universally bestowed by all educated and careful people upon the vaccination of infants. Each child of a family ought to be subjected to this operation at or about the age of fifteen, and we think the family physician ought to bestow the same care upon this as upon the primary vaccination. One trial, without result, we hold to be of no more use than it would be in an unvaccinated child. The trial should be made again and again until a result is obtained. We have ourselves of late years repeated it twice, three times, and, in one instance, seven times, before we succeeded in obtaining a vesicle. Once the vesicle obtained, with a good areola, we believe the subject is safe for many years, probably for the lifetime.

The characters of the vaccine disease produced by a revaccination are not always the same as those obtained at the primary vaccination, especially when the time between the two operations is only that extending from birth to puberty. We have seen at later periods of life, at thirty and forty years of age, for instance, as perfect specimens of the vaccine disease from a secondary vaccination as we have ever seen in the infant. Not a feature has been wanting. The exact phases of the disease, the papule, the vesicle, the precise duration as to time, the areola, the constitutional disturbance, and the resulting cicatrix, have all been perfect in every point.

It is difficult to escape the conviction, that in such cases as these just mentioned the protective power of the primary vaccination had been entirely obliterated, and such, indeed, is and has been the opinion of many. A careful observation has shown, however, that this is not correct, and

that, to use the words of Dr. Seaton (*loc. cit.*, p. 511), we cannot "draw from the local phenomena of revaccination any inferences whatever as to the state in which the revaccinated persons were as to liability to small-pox. Jenner himself, indeed pointed this out in his first treatise, and showed that the natural cow-pox might be induced again and again in persons who, being protected against variola by their first attack of cow-pox, could not be variolated either by inoculation or by exposure, as well as that cow-pox might be made to take on those who had had small-pox." A table, given by Dr. Seaton to show the results of revaccination in the Würtemberg army in 1831-35, and in the English army in 1861, shows conclusively that revaccination was nearly as successful in producing a perfect vaccine disease in those who bore the marks of previous small-pox, and in those who had good cicatrices of previous vaccinations, as in those who bore no marks of previous vaccination or small-pox.

These facts overthrow the prevalent notion held by the public at large and by many physicians, that a successful revaccination is a sure sign that the subject had lost the protection afforded by the previous vaccination. They also overthrow the idea that it is necessary to revaccinate every few years in order to renew the protective power of the vaccine disease. One good primary vaccination doubtless affords full protection throughout life in many, but it does not in all, and since it is impossible to determine which are the protected and which the unprotected, it is necessary to revaccinate all. But one successful revaccination is probably all that is required. Should, however, any one who has been thus successfully revaccinated be exposed directly to the infection of small-pox many years afterwards, it might be well to repeat the operation once again. The fashion, however, of being revaccinated every few years, which some persons indulge in and some physicians assent to only too readily, is simply a work of foolish supererogation not unattended with risk, since vaccine punctures, though made in the most legitimate way, will occasionally cause severe and even dangerous sores.

We have already said that revaccination at puberty rarely produces a vaccine disease of typical character. Still more is this true of children under puberty. At that early age the disease usually begins earlier after the puncture than in the primary form, reaches its height by the fifth or sixth day, and then declines. The vesicle is apt to be acuminate rather than umbilicated, the areola is irregular in outline, narrower, paler and is usually hard. The scab is small and imperfect, looking more like one formed by the desiccation of pus than like that formed from true vaccine lymph, and is often complete by the eighth day, and soon falls. There is often a good deal of constitutional irritation caused by revaccination, more even than in the primary disease, and there is also much local irritation in the form of itching and pain. Nevertheless, these appearances are invaluable as showing that the lymph employed has affected the constitution of the patient. Without some such response to the revaccination, we hold the operation to have been useless, and always repeat it, as has already been stated.

To impress upon all the power and value of revaccination, we will quote

some facts given by Dr. Seaton (*loc. cit.*, p. 509): "Heine found that in five years there occurred among 14,384 revaccinated soldiers in Würtemberg only one instance of varioloid, and in 30,000 revaccinated persons in civil practice only two cases of varioloid, though during these years small-pox had prevailed in 344 localities, producing 1674 cases of modified and unmodified small-pox among the not revaccinated, and in part not vaccinated, population of 363,298 persons in those places in which it had prevailed. In the Prussian army, since the introduction of systematic revaccination of all, the annual deaths from small-pox (which at one time were 104) have not averaged more than 2; and on analysis of 40 fatal cases that occurred in twenty years, it appeared that only 4 were in persons who were said to have been successfully revaccinated."

He also cites Mr. Marson's statement, to the effect that in "thirty years no nurse or servant at the Small-pox Hospital has taken small-pox, he having taken care always to revaccinate them on their coming to live in the hospital; and further, that when a large number of work-people were employed for several months about the hospital, most of whom consented to be revaccinated, two only were attacked by small-pox, but they were amongst the few who were not revaccinated."

With a few words on the mode of performing the operation of vaccination, we shall bring this article to a close.

Different methods of inserting the vaccine virus have been employed by different practitioners. The two methods most frequently resorted to are those by incision and puncture. The former consists in making a superficial incision of several lines in length into the skin, in such a way as to cause a very slight effusion of blood. Into this is introduced a small quantity of a dried vaccine scab reduced to a fine powder, or a piece of fine thread wet with the vaccine fluid, or with water holding in suspension a portion of dried virus. Over the wound is then placed a piece of isinglass plaster, which is secured by a bandage. This is to be removed after two or three days, and the disease allowed to pursue its regular course. The operation by puncture is performed by introducing horizontally beneath the skin a needle or lancet charged with the virus, and then withdrawing it in such a way as to leave the virus in the wound. Of these two modes the latter is the one now most frequently adopted, the former having been found to occasion, not unfrequently, a spurious disease, and to be of very difficult application in the cases of children. For our own part we have used for some years past a method that we have found much the most convenient in children, and which rarely fails when it is carefully performed. We take a common thumb lancet, which should not be too sharp. Holding the arm of the child with our left hand, and stretching the skin between the forefinger and thumb, whilst the under part of the arm is grasped by the second finger placed beneath the first, we lay the lancet flat upon the skin, and using the point, remove, by a repeated and very gentle rubbing movement, the cuticle, until the surface of the derm is laid bare, so as to allow of a perceptible, and merely perceptible oozing of blood, or, in other words, so as to expose a living surface. This surface should be about as large as a small-sized bird-shot, and it should not

bleed, but merely show that the vascular part of the derm has been reached and slightly exposed. On this surface the vaccine fluid or dissolved scab is to be placed in quantity sufficient to cover it, and the nurse should be told to leave the arm bare and untouched for twenty minutes, or until the applied fluid has dried into a little scab, when no further precautions are necessary. This mode of operating may at first seem tedious and painful. We can only say that when performed gently and gradually, it causes so little pain that we have often practiced it upon sleeping children without waking them.

If the quills of fresh bovine lymph are used, an abrasion of the cuticle is made in the ordinary manner, and the end of the quill charged with lymph is very slightly moistened and then rubbed on the abraded spot until the virus is thoroughly removed.

The place usually selected for the operation is, as every one knows, on the arm close to the insertion of the deltoid muscle. This is the best place as a general rule, and particularly in girls, whose parents often object to having the insertion made below this, lest the scar should be visible in after years, when the arm is uncovered. In boys we often select the radial edge of the forearm some two inches below the elbow, since in this place the pock is least apt to be injured in the act of dressing the child, or of lifting it about.

ARTICLE IV.

VARICELLA.

DEFINITIONS; SYNONYMS; FORMS.—Varicella is a contagious eruptive disease of benign nature, characterized by more or less numerous transparent vesicles following rapidly upon small red elevations. The eruption is usually preceded by slight initial symptoms lasting from one to two days, and it terminates by the desiccation of the vesicles about the fifth or eighth day after their appearance.

It is known also in English by the names of chicken-pox, swine-pox, and crystalli.

Several different forms of the disease have been described by different writers under the titles of lenticular, conoidal, and globular; but inasmuch as these varieties are of no real importance in practice, we shall merely advert to them casually in our account of the eruption.

CAUSES.—Varicella is propagated in two ways: by *contagion*, and by *epidemic* influence. That it is contagious there can be no doubt, since nearly all observers agree upon this point. In our own experience we have seldom known any child, who had not had the disorder previously, to escape it when once it has entered a household. It rarely attacks any but children. Its epidemic nature is shown by the fact that in some seasons it is scarcely seen, whilst in others it prevails extensively over large districts of country, and attacks many children in the great towns and cities of those districts. Varicella occurs only once in the same individual.

Considerable discussion has taken place at various times as to the real nature of varicella, some asserting that the disorder is merely one of the varieties of modified small-pox, while others maintain as strenuously that it is an independent and specific disease. The weight of authority, however, seems to be clearly in favor of the last-mentioned opinion, and we have no hesitation in avowing this to be the conclusion to which our own reading and experience have brought us. When we consider, indeed, that varicella is, unlike either variola or varioloid, incommunicable by inoculation,¹ that it attacks indifferently the vaccinated and unvaccinated, that its course is entirely unaffected by previous vaccination, and that the vaccine disease is readily taken, and passes through its regular phases after varicella, we do not see how we can refuse to believe that the latter is something entirely independent of small-pox, and therefore a distinct and peculiar malady.

SYMPTOMS ; COURSE ; DURATION.—The eruption is usually, but not always preceded by prodromic symptoms. These seldom last more than one, or at most two days, and consist at the very beginning of slight chilliness, or of a chill even, which is followed by a more or less marked febrile reaction. In some instances there is vomiting, but this is rare, and when it does occur, slight. When fever exists it is marked by headache, accelerated pulse, slight warmth of the surface, pain in the back and limbs, languor, indisposition to play, some unusual irritability of temper, diminution or loss of appetite, and unusual thirst. These symptoms may be present, and yet in so mild a shape that the child shall show no disposition to abandon its ordinary habits of activity and play, while in other cases again, there are literally no initiatory symptoms whatever, and the appearance of the eruption is the first declaration of the presence of the malady. Even when constitutional symptoms are present, they usually disappear by the third day.

The eruption appears in the form of small papular spots, of a deep-red color, and irregularly circular shape, which generally show themselves first on the front and back of the trunk, and extend very soon to the face, and a little later, to the extremities. We have known a child to go to bed at night with slight headache and fever, and present a well-marked though not yet abundant varicellous rash upon the upper part of the trunk, and on the face, on the following morning. These papules exhibit,

¹ Steiner of Prague (quoted in Medical and Surgical Reporter, July 17th, 1875, p. 57) has lately inoculated varicella in several cases, with the following results :

1. The contents of varicella vesicles are inoculable: of 10 cases of inoculation, 8 were successful and 2 failed.
2. After the successful inoculation of varicella, varicella and not variola or varioloid was invariably produced.
3. The stage of inoculation in all the successful cases was eight days.
4. In 4 cases there were no prodromes; in 4 cases there were prodromes of four days' duration.

Vaccination has no influence on the production of the exanthem; of the 8 cases of successful inoculation, 5 had been vaccinated, 3 had not.

Varicella does not protect against variola; in one case a child died of confluent variola fourteen days after convalescence from an attack of varicella.

in the course of a very few hours, small vesicles in their centres; indeed, according to some observers, the eruption is vesicular from the very beginning. On the second day the papules are in great measure converted into vesicles, which may be either small and acuminate, constituting the *lenticular* form of the disease, or they may be larger and of a more globular shape, constituting the *conoidal* and *globular* or *globose* forms of Willan and Bateman. We deem it unnecessary, as above stated, to describe different varieties of varicella, since this is useless for any practical purposes, and because we constantly see upon the same subject vesicles of very different shape and size. When fully completed, the vesicles are often of very considerable size—two or three lines in diameter; they contain a transparent fluid, which is either entirely colorless or of a faint orange tint, and some of them are surrounded by a small ring of inflammation. On the third day, the eruption continues in nearly the same state as on the second, except that the fluid contained within the vesicles assumes a yellowish appearance, owing to its passage from the serous into the purulent condition. On the fourth day, the process of desiccation begins and goes on rapidly, the vesicles that have not been broken by accident, or torn by the fingers of the child in its efforts to appease the itching which they give rise to, assuming a shrivelled and shrunken appearance at their margins. As this process goes on, the vesicles are gradually converted into light brownish scabs, so that by the sixth day they are nearly all dried up.

The scabs are usually thin; they dry from the circumference to the centre, and between the eighth and ninth days fall off, leaving behind faint red spots, not depressed below the general surface, and which soon disappear.

The eruption is generally accompanied, as was stated above, by a sensation of heat and itching in the vesicles, which causes the child to rub and scratch them in such a way as often to break those which he can reach, and thus prevent them from passing through the regular periods of maturation and desiccation.

DIAGNOSIS.—There is but one disease with which varicella could be confounded, and that is variola in some of its shapes. With regular small-pox such a mistake could scarcely happen even to the inexperienced. With varioloid, on the contrary, there might be some difficulty, and yet, if it is borne in mind that in varioloid the initiatory fever is much more severe, lasting three days instead of twelve or thirty-six hours, that the eruption appears first on the face and extends very slowly to the trunk and extremities, and that the conversion from the papular into the vesicular condition is much more gradual than in chicken-pox, we think no serious difficulty can ever occur in making the distinction between the two affections.

PROGNOSIS.—The prognosis is always favorable. The only real trouble that we have ever known to occur has been from catarrh or pneumonia contracted by imprudent exposure during the convalescence.

TREATMENT.—In a large majority of the cases, varicella requires no treatment beyond attention to diet for the first two or three days, and the avoidance of cold during the convalescence. When the constitutional symptoms are marked, the fever and headache being considerable, a dose

of some mild cathartic, a little sweet spirit of nitre in cold lemonade or orangeade, rest in bed, and one or two foot-baths, will be all that is necessary to reduce these symptoms and make the patient comfortable.

ARTICLE V.

SCARLET FEVER OR SCARLATINA.

DEFINITION; FREQUENCY; FORMS.—Scarlet fever is an epidemic and contagious eruptive fever, characterized by a scarlet rash, which appears on the first or second day of the disease, and ends usually about the sixth or seventh, or in rare cases so late as the tenth; by simultaneous inflammation of the tonsils, and of the mucous membrane of the mouth and pharynx; and by desquamation.

The frequency of the disease is exceedingly variable in different years, owing to its epidemic nature. This may be readily seen by a glance at the following table, which gives the annual mortality for the past sixty years in this city, from scarlatina and measles:

Scarlatina. Measles.			Scarlatina. Measles.			Scarlatina. Measles.		
1809	3	0	1833	61	1	1857	704	66
1810	2	1	1834	83	7	1858	241	28
1811	3	2	1835	305	248	1859	232	51
1812	1	20	1836	240	4	1860	591	15
1813	0	1	1837	205	49	1861	1190	74
1814	0	9	1838	134	123	1862	461	109
1815	0	7	1839	225	136	1863	275	82
1816	0	2	1840	244	2	1864	349	90
1817	0	0	1841	83	119	1865	624	54
1818	1	0	1842	220	24	1866	491	221
1819	2	108	1843	395	1	1867	367	83
1820	31	47	1844	269	3	1868	224	108
1821	13	0	1845	199	90	1869	799	85
1822	9	0	1846	221	6	1870	956	48
1823	11	156	1847	344	77	1871	262	41
1824	9	102	1848	172	99	1872	174	136
1825	9	38	1849	242	27	1873	319	30
1826	4	101	1850	440	72	1874	461	117
1827	1	9	1851	391	17	1875	1032	12
1828	0	58	1852	434	90	1876	328	53
1829	9	53	1853	388	14	1877	379	69
1830	40	7	1854	162	62	1878	554	12
1831	200	23	1855	163	24	1879	336	8
1832	307	118	1856	992	141			

It will be noticed that for five successive years 1813–17 inclusive, not a single death from scarlatina is reported; and that during twenty years, 1809–28 inclusive, only 99 deaths occurred from this cause; while in the single years 1856 and 1861, 992 and 1190 deaths respectively are reported.

During the entire series of sixty years, there have been 18,616 deaths from scarlatina returned.

Hillier states, that during the eighteen years from 1848 to 1866, the deaths from scarlatina in London amounted to 52,461.

It is impossible to estimate the actual relative frequency of scarlatina and measles, owing to the absence of any returns of non-fatal cases. It is evident, however, from the above table that, although the mortality from measles is also very variable, and thus may for a short time exceed that from scarlatina, in a long series of years the latter disease is far the more fatal. Thus the number of deaths from measles in this city, during the past sixty years, amounts to but 2279.

MM. Guersant and Blache (*Dict de Méd.*, t. 28, p. 173) state that it is less frequent than measles or variola. They added together the cases of the eruptive fevers collected in 1838 and 1839, by MM. Roger, Rilliet and Barthez, and Barrier, in the Children's Hospital at Paris, and found that there had only been 157 of scarlet fever; whilst there were 267 of measles, and 213 of variola and varioloid.

The *forms* of the disease generally enumerated are the *simple*, *anginose*, and *malignant*. Authors differ widely in their descriptions of these three forms. Many of the English authors include in the simple form only the cases in which there is no affection of the fauces, while the anginose form includes all in which there is any throat affection whatever. M. Rayer, on the contrary, describes under the head of the simple form the cases in which the throat affection is mild, while he considers the anginose form to be that in which a pseudo-membranous angina occurs. Again, the descriptions of the malignant form are vague and uncertain, some including under this term only the rapidly fatal cases in which cerebral symptoms are present, while others include those also which are rendered malignant by the occurrence of pseudo-membranous angina.

We believe this division of scarlet fever into distinct forms and varieties to be, for several reasons, a faulty arrangement. It is not, it appears to us, in the first place, consonant with the nature of the disease. Scarlet fever is, in fact, with all its degrees of severity, and apparent differences, a single and distinct fever, produced by one cause, determining similar effects, howsoever much they may vary in degree, and requiring no more than does typhoid fever to be divided into the variety of different forms, which it has been customary to ascribe to it. Again, the above mode of division is not, we are sure, a good one for practical purposes. It is impossible, indeed, as we have often found it, to refer many cases we meet with in practice, clearly and satisfactorily, to any one of the forms of the disease described in books. The simple form of some of the English writers, or that in which there is no anginose affection, has no existence whatever, so far as we have been able to discover. We believe that inflammation of the mucous membrane of the fauces constitutes an essential element of the disease, for we have never yet seen a case of scarlatina in which it was not present to a greater or less extent. It is often very slight, so slight, indeed, as to be unaccompanied by any evidence of pain in the part, but in all that we have examined, it has been decided and

obvious. This supposed form of the disease does not, therefore, in our opinion, exist.

The two other forms usually described, the anginose and malignant, are also of little value practically, since we have found that in all severe or grave cases, in which the patient did not die with violent nervous symptoms under the first shock of the scarlatinous poison, there has been developed a severe and dangerous anginose inflammation about the third or fourth day; so that it is fair to say that we cannot imagine any malignant case, lasting over the third or fourth day, which is not anginose, nor any severe anginose case, which might not also be styled, from its dangerous character, malignant. We have found it impossible, in our experience, to draw the distinction clearly and indubitably between the anginose and malignant varieties, because all severe cases partake more or less of the features of both.

Feeling this difficulty of describing the disease according to the mode that had before that time been generally followed, and believing it also to be insufficient for practical purposes, we were led to attempt, in the first edition of this work, a different arrangement.

We made, accordingly, two forms or degrees of the disease, which we designated by the terms *regular* and *grave*. In the first form or degree we included all the cases in which the angina was simple and the eruption regular in all respects; in which there was no predominance of one set of symptoms over another, but in which all held a due relation to each other.

In this form were embraced all the cases of scarlatina simplex of writers, and many of those of scarlatina anginosa of the English authors. In the second form we included the cases which departed from the regular course of the disease, and which were rendered dangerous by the occurrence of severe symptoms not belonging in the same degree to the simple affection. This form we subdivided into two varieties, the *grave anginose*, which contained all the cases accompanied by pseudo-membranous, ulcerative, or gangrenous angina; and the *grave cerebral*, which comprised all those marked by the early occurrence of dangerous cerebral symptoms. The grave form comprehended, therefore, some of the cases of scarlatina anginosa, and all those of scarlatina maligna of writers, dividing, however, those in which a pseudo-membranous, ulcerative, or gangrenous angina determined the type of the attack, from those in which the cerebral or nervous symptoms gave to the case its stamp.

More extended observation and more patient reflection have taught us that this division also is incorrect,—that it does not afford a good classification for the purposes of description, and that it is defective as a guide in practice.

We adopted, therefore, in the third edition, and shall follow in the present one, a different method of considering the disease, one which we believe to be more consistent with its nature, more suitable for the purpose of description, and much more likely to prove useful in practice. We shall follow the same arrangement in regard to scarlet fever as that now generally employed for typhoid fever. We shall consider it as a single and distinct disease, and not as made up of a number of uncertain

and imperfectly separated forms of varieties, since these so run into each other, as to make it absolutely impossible to draw the line clearly and palpably between them. The only division we shall make is into *mild* and *grave* cases, since the only real difference between the cases is a difference in the degree of severity they exhibit.

CAUSES.—It has been abundantly proven by long and repeated observation that scarlatina is propagated by two causes,—*contagion* and *epidemic* influence. Of these two modes of propagation, we have not the least doubt ourselves that the latter is by far the most active. It is only necessary to look over the results afforded by the tables of mortality for this city, as quoted in the early part of this article, and to observe that in some years the disease caused a heavy mortality, in others a very small one, and that in others again not a single death from it is reported, to be convinced that it is of a highly *epidemic* nature.

The *contagious* character of scarlatina has been doubted by some few persons, but seems to us clearly proved by the evidence adduced by various writers. Our own experience also convinces us that it is a contagious disease, though much less so, we think, than either small-pox, measles, hooping-cough, or chicken-pox. We have quite frequently, indeed, known children exposed directly and for a considerable length of time to the infection to escape entirely, while it is extremely rare for us to meet with children, unprotected by previous attacks, who can resist the contagion of measles, hooping-cough, or varicella. Thus Dr. C. E. Billington (New York *Med. Record*, March 23d, 1878, p. 221) reports that in 26 families with 90 children, who were all exposed to the contagion of scarlatina, 43 had the disease and 47 escaped. He justly says that if such a result had occurred while any prophylactic was being used, false conclusions might readily have been drawn. But, though we believe it to be much less highly contagious than has been generally supposed, and than the other contagious diseases just named, we are also well convinced, as was stated above, that it is propagated to a considerable extent by a direct contagion. We have, in a number of instances, known one child in a family to contract the disease from direct exposure to it, or from the epidemic constitution of the atmosphere, and a second, third, and even a fourth, to take the disease from the first, in five, seven, or nine days after the latter had fallen sick. In other instances, on the contrary, it would seem that either several children in one family contract the disease nearly simultaneously from the epidemic influence, or else that the period of incubation is sometimes very short. For example, during the winter season, a child five months old, who had never been out of the house, was seized with it. On the second day after the eruption appeared on this child, her sister, between four and five years old, fell sick, and on the third day another sister, the only remaining child, between two and three years of age. In the first of these cases it must have been contracted through the epidemic influence which was at that time prevalent in the city, since the child had in no way been directly exposed to it. In the other two, we must either suppose the cause to have been the same, or else that the period of incubation was only two and three days in the respective cases.

The period of incubation is shorter than in other contagious eruptive diseases. It may be stated to vary between twenty-four hours and two or three weeks. MM. Guersant and Blache are of opinion that in the majority of cases, it is from three to seven days. MM. Rilliet and Barthez found that of 38 cases in which the time was recorded, it was between 2 and 7 days in 16, between 8 and 13 in 15, and 15 and 40 in 8 cases. Our own observation would fix it at from 9 to 15 days in the majority of cases.

Occasionally, however, it is very short; thus Trousseau mentions a case in which the evidence is almost conclusive that the period of incubation was less than twenty-four hours. Murchison also states that this latent period varies from a few minutes to five days, rarely, if ever, exceeding six days.

It is impossible to state with any certainty the length of time during which the power of imparting the contagion continues in the patient. M. Cazenave (*Abrégé Prat. des Mal. de la Peau*, p. 54) states that it lasts throughout the period of desquamation, and that it would even seem to be most active at that time.

Whatever may be the duration of this period, it is certain that the virus may attach itself to clothing, bedding, or furniture, and that the disease may thus be transmitted by one who is not himself attacked. We also learn from some remarkable instances, as for example, from a case related by Richardson in "*The Asclepiad*," that when the virus is thus attached to fomites, it may retain its activity for many months.

In regard to the essential nature of the poison, it appears probable, in the first place, that it is contained in the secretions of the skin and fauces.

The distance to which it may be carried by the air does not appear to exceed a few feet, and in those cases where prompt isolation does not prevent the communication of the disease, the virus has either been previously imbibed or is carried by fomites. It is probably of material nature, and is admitted to the system either through the skin, the respiratory, or, perhaps, the gastric mucous membrane.

As we have seen, it retains its activity for a long time; but is rendered inert by a temperature somewhat below 212° F.

Scarlatina is stated to be also inoculable, by the blood, the secretion from the fauces, and the fluid from the miliary vesicles which occasionally form on the skin. The resulting disease appears in some instances to have been favorably modified, but the operation has been comparatively rarely practiced.

The epidemics of scarlet fever vary exceedingly in their extent and violence. During the years 1842 and 1843, the disease prevailed very extensively in this city, and assumed a malignant type, so that in a considerable number of families, two, three, and even four children, died within a very short period.

During the winter of 1856-57, and throughout the spring of 1857, we had one of the most prevalent epidemics that ever visited this city, and yet the proportion of deaths to the whole number of cases in our own practice and that of our friends, was such as to seem to show that the type of the epidemic was mild.

The disease prevails at all seasons, but is most frequent in the spring and summer, and next in the autumn. It rarely occurs more than once in the same individual, but that it does so sometimes, is proved by facts brought forward by different authors. It has been asserted that second attacks of scarlet fever occur in the same person not more than one in a thousand cases. Of the truth of this assertion we are, however, very doubtful, since it has occurred to us to see no less than three examples of second attacks in our own experience. We attended in this city one child with perfectly well-marked scarlet fever, attested by subsequent anasarca, who had had the disease two years previously under the care of the late Prof. C. D. Meigs. In the winter of 1852, we attended two children in one family with the disease, one of whom died, and both of whom had had the disease four years and a half before. They were attended in the first attack by one of ourselves, and as it chanced, owing to our absence from town during one day, they were seen also by one of our friends, who made no exception whatever to the diagnosis of scarlet fever. The only doubt as to these cases having been veritable examples of double attacks of the disease, must rest of course upon the diagnosis. In the first example, the diagnosis was made by Prof. Meigs in the first attack, and by one of ourselves in the second. In the two latter it was made by one of ourselves in both, accidentally confirmed in the first attack, in both children, by the opinion of a competent professional friend. The first attacks in the latter cases were both mild, but well marked; the second were both severe, and one proved fatal on the sixth day. We have not the least doubt ourselves that all of the three were cases of true scarlet fever. If they were not, the two latter must have been cases of roseola, so closely resembling scarlatina as to oblige us to confess ourselves incompetent to distinguish between the two diseases. What adds to the certainty that the two which came under our own observation were examples of scarlet fever, is the fact that they occurred simultaneously with a third case in the same family. Now, roseola is not apt, so far as we know, to occur epidemically in a household. Most of the cases of that disease that we have seen, have been solitary ones. Again in the spring of 1857, one of us saw a well-marked attack of the disease in a boy nearly four years old, who had had it one year before, under the charge of a perfectly competent practitioner.

Dr. Richardson (*loc. cit.*) asserts that he has known the disease to occur twice in the same patient, and also states that he himself has suffered from it three times.

Age.—MM. Rilliet and Barthez state that it is most common from six to ten years of age. Of 251 cases that we have seen, in which the age was noted, 64 occurred under 3 years of age, 78 between 3 and 5 years, 51 between 5 and 7, 47 between 7 and 10, and 11 between 10 and 15. From this it would appear to be more common in the first five years than between the ages of five and ten, since of the 251 cases, 142 occurred in the former, and only 98 in the latter period. By uniting the statistical tables of Dr. Emerson with those of Dr. Condie (*Dis. of Child.*, 2d ed., note, p. 86), we obtain the deaths from scarlatina in this city at different ages for a period of thirty years. These tables show clearly that the disease is

most common between the ages of one and five years. The total mortality from scarlatina under ten years, during the time stated, was 2171, of which 132 were under one year of age, 411 between 1 and 2, 1130 between 2 and 5, and 510 between 5 and 10.

Of 148,829 cases collected by Dr. Murchison from the death returns of Great Britain, 9999, or about 7 per cent., were under 1 year; 30,974, or 20 per cent., under 2 years; 95,070, or 64 per cent., under 5 years; 38,591, or 26 per cent., between 5 and 10; and but 13,168, or about 9 per cent., at all ages above 10.

This agrees quite closely with the averages calculated from the extensive statistics collected by Dr Richardson, which show the following percentage at different ages :

Under 5 years,	67.63
From 5 to 10,	24.33
“ 10 to 20,	5.52
“ 20 to 40,	1.73
“ 40 upwards,	0.66

Out of 12,962 deaths under 5 years, 1289, or 9.9 per cent., were under 1 year; 2874, or 22 per cent., between 1 and 2; so that 4163, or 31.4 per cent., were under 2 years.

The earliest age at which we have seen it perfectly well marked, was twenty-one days. We saw it once also in a child five months of age, and twice at the age of six months. It is not nearly so common in the first year of life as it is afterwards. The largest number of cases occur, according to our experience, in the third, fourth, and fifth years of life.

The influence of *sex* seems not to have been determined with certainty. Dr. Tweedie (*Cyclop. of Prac. Med.*, art. Scarlatina) says it is most common in girls. MM. Rilliet and Barthez, on the contrary, state it to be more common in boys. Of 262 cases under 15 years of age that we have seen, in which the sex was noted, 133 occurred in males, and 129 in females. The truth is, probably, that under puberty it attacks the two sexes with about equal frequency, while after that age it is most common in females.

It occasionally happens, that patients, both adults and children, who have undergone surgical operations, are attacked with a scarlatinous rash, with mild constitutional symptoms (Hillier, Gee). The disease, according to these authorities, is true scarlatina; and its occurrence at that time probably depends upon the system being in an unusually favorable condition for the reception of the virus.

SYMPTOMS ; COURSE ; DURATION.—As has already been stated, we intend, in our description of the symptoms of scarlet fever, to depart from the ordinary mode of arrangement of the subject. We shall discard the old division of the disease into three forms or degrees, scarlatina simplex, anginosa, and maligna, and substitute, for reasons already given, the simple division into mild and grave cases. We shall class as mild cases those which pursue an even and regular course, without being accompanied by dangerous or malignant symptoms, in which there occur neither violent

nervous, nor threatening anginose symptoms ; while among the grave cases we shall place those in which there occur severe nervous symptoms, in the form of delirium, coma, or convulsions, dangerous symptoms in the form of diphtheritic, ulcerative, or gangrenous inflammation of the mucous membrane of the fauces, and finally, those in which the general symptoms assume a low and typhoid character. When it seems convenient, we shall follow the usual division of the course of the disease into the three stages of invasion, eruption, and desquamation.

Mild Cases—Stage of Invasion.—The following description of the symptoms of scarlet fever in its mild form is drawn partly from books, but much more from our own observation of several hundred mild cases of the disease, of 213 of which we have kept a faithful record, and, when there was anything peculiar or important, full notes.

The onset of mild cases of scarlet fever is generally sudden. A child is well, or so slightly ailing, that no change from its usual condition is noticed at the time, though some slight signs of indisposition may be recalled afterwards, and on the following day, or often within twelve hours or less, the symptoms of the disease become marked and characteristic. In a large majority of the cases that we have seen, the eruption was already visible at our first visit. Frequently the patient has been to bed well in the evening, and, becoming restless and feverish in the night, is found on the following morning with fever, sore throat, and very considerable eruption ; or, as happened in one of our cases, a child gets up in the morning apparently well, breakfasts as usual, goes to church, and falling sick there, comes home and, a few hours later, shows the eruption over the neck and upper part of the trunk, and has fever and sore throat. In another case, a boy between seven and eight years old was perfectly well in the morning. At 2 P.M. his mother, a most sensible and accurate person, observed him playing in the garden, and remarked upon his healthy looks. Fifteen minutes after this he felt sick at the stomach ; he came into the house and went up to the nursery, looking pale and pinched, with a cold skin, and nearly fainted in the nurse's arms. He had then in the course of an hour three copious and watery stools, each one accompanied with vomiting. We saw him one hour after this, dozing, very pale, with pinched features, sunken and half-closed eyes, cool surface, and with the pulse at 128, and rather feeble. There was no eruption. At 6 P.M. we found him with a hot and dry skin, with the tongue heavily coated, the fauces swollen and showing flecks of exudation upon the tonsils, a pulse at 128, and with a well-marked scarlatinous eruption coming out abundantly. The case pursued a very regular course, without dangerous or malignant symptoms of any kind.

But the invasion, though sudden in nearly all cases, is not always so precipitate as we have just described. When we come to analyze the early symptoms, we find that the first one observed in most of the cases is fever, marked by considerable acceleration of the pulse and heat of skin. In some few cases the fever is preceded by the ordinary prodromes of febrile diseases, languor, lassitude, pains in the back and limbs, and slight rigors. Simultaneously with the fever there is in nearly all cases more or less sore-

ness of the throat. Dr. Billington (*loc. cit.*) thinks that the precedence of angina to every other symptom is invariable. He describes it as differing from the appearance of catarrhal angina. It consists in diffuse redness, sometimes at first punctate on one or both half-arches, then extending around them and involving the uvula, extending also to the tonsils, which become reddened and more or less enlarged. The posterior wall of the pharynx is little if at all affected. In all that we have examined, even those in which no pain was complained of, there has been redness, or redness with swelling of the fauces. In a majority of the cases vomiting occurs, or if not vomiting, some degree of nausea. There is complete anorexia; the thirst is acute; the bowels are usually in their natural condition, or slightly constipated. The child is quiet and dull, or else restless and irritable, and sometimes there is delirium; the face is generally flushed, and the eyes often slightly injected. The duration of these symptoms is irregular. They are said to last generally about a day, but they may continue either a shorter or longer period. We are very sure, from our own observation, as we have already stated, that these premonitory symptoms rarely precede the eruption more than twelve hours, and very often the time is less, so that the eruption may be the first symptom noticed.

Stage of Eruption.—The eruption generally appears first on the face and neck, whence it extends rapidly over the whole surface. It continues to increase in extent and intensity, so as to reach its maximum about the third or fourth day. It appears first in minute dark-red points, dotted upon a rose-colored surface, forming patches of irregular shape, of considerable size, level with the skin, disappearing under pressure, divided at first by portions of healthy skin, but running rapidly together, and giving to large portions of the surface a uniform scarlet color. The eruption is not generally equally diffused over the body, but is more marked upon one portion than another. It is often most intense on the back, and is there of a deeper color than elsewhere, not unfrequently assuming a purple hue. It is generally very well marked on the abdomen and thighs, and about the articulations, and assumes in those regions a particularly bright tint.

It does not always cover the whole surface, but in some very mild cases, and, as we shall find when treating of the grave cases, in them, also, it may occur only in patches of moderate extent upon different portions of the body, leaving us at times in doubt as to the real nature of the rash.

The surface of the eruption is smooth and even to the touch, unless, as not unfrequently happens, it is accompanied by the development of miliary vesicles, or crops of minute pimples or pustules. A certain degree of roughness is sometimes occasioned also by enlargement of the papillæ of the skin in various parts of the body, particularly on the extensor surface of the limbs; but this is evidently independent of the characteristic eruption. The skin upon some parts of the body, especially the face, hands, and feet, often presents a swollen appearance, rendering the movements somewhat stiff. There is in most cases a feeling of burning, irritation, and itching in the skin, the latter of which symptoms increases as the malady progresses.

If the nail be drawn firmly over the skin where the eruption exists, a white line is produced, which lasts for a short time and then passes away; if the pressure be more firm, a central red line with a white streak on either side is developed. This was originally pointed out by Bouchut as pathognomonic of scarlatina, the peculiarity, according to him, consisting in the great duration of the white line so caused. It does not appear, however, to have any positive value in distinguishing this affection from many forms of erythema.

The eruption generally reaches its height about the fourth day, and then remains stationary for one, or less frequently for two days, after which it begins to decline. Its decline is marked by a diminution in the intensity of the color, which, from scarlet, becomes red, then rose colored, and growing paler and paler, finally disappears entirely about the sixth, seventh, or eighth day. In some very mild cases, however, the whole duration of the eruption is not over two or three days, and in such the color it imparts to the skin is never very bright nor very deep, nor is it accompanied by intense heat, or by much irritation or itching.

The symptoms which precede the eruption do not subside on its appearance, but persist or are augmented. The febrile movement continues unabated; the pulse is full, strong, and frequent, running up very soon after the onset to 120, 140, 150, and often to 160. This frequency of the pulse is, in fact, one of the most marked symptoms of the disease. We have rarely, even in very mild cases, found it less than 140, and in not a few it has been in the first few days, and in children of four or six years old, as high as 168 or 170. Occasionally, however, it has been lower, and in a case that occurred to one of us, in a boy five years old, it was 96 on the second day, and only 88 on the third, though there was still a good deal of rash upon the skin. The skin is burning hot and dry, as a general rule, and loses its usual softness and suppleness. The expression of the face is usually natural. The eye is often animated, and slightly injected. The respiration is generally easy and natural, though sometimes, when the fever is violent, it becomes quickened. The auscultation and percussion signs are natural, unless some complication exists. There is often a rather frequent cough, which is dry, and evidently depends on the guttural inflammation, and not on any bronchial or pulmonary affection; it exists during the early period of the eruption, and declines with the inflammation of the fauces. The voice is seldom altered beyond having a nasal sound, so long as the disease continues simple and regular. If the voice become hoarse or whispering, it indicates an extension of the inflammation from the pharynx to the larynx. The anorexia continues until the eruption begins to decline, and the thirst is acute up to the same period, when it moderates. At first the dorsum of the tongue is covered with a whitish or yellowish-white fur of variable thickness, while its tip and edges are of a deep-red color. After two or three days, and during the course of the eruption, the coating just described disappears from the tongue, and its whole surface assumes a deep-red tint and a shining appearance, which makes it look like raw flesh. At the same time it is often much diminished in size from contraction of its tissues, and its papillæ be-

come enlarged and projecting; this condition generally lasts from six to ten days, after which it returns to its natural state; it is commonly moist throughout the attack. Vomiting is rarely troublesome in mild cases, though it often occurs; the bowels continue nearly in their natural condition; in some few cases slight diarrhœa occurs, but more frequently there is very moderate constipation. The abdomen is natural in most of the cases; sometimes, however, there are slight distension and pain for a few days, which coincide generally with slight enlargement of the liver, or more rarely of the spleen.

The urine during this stage usually presents the ordinary febrile characters; it is diminished in quantity, often of high color, though the pigment is not necessarily increased. The urea is not increased, which Ringer regards as indicating that the kidneys are affected from the beginning of the attack. The chlorides are always more or less diminished. The phosphoric acid, according to Dr. Gee, is about normal for the first three or four days; it then diminishes, and remains for a few days at a half or a third of its normal amount. Uric acid appears to be retained during the pyrexia, and excreted in excess so soon as it begins to subside. According to Holder's examination of 17 cases, there is bile pigment present during the first six days.

Early in the second, or even in the first stage, the fauces present the signs of inflammatory action; the pharynx is reddened, and in some instances swollen; the tonsils enlarge and become red; the submaxillary and lymphatic glands are somewhat tumefied and tender to the touch, and when the case is at all severe, deglutition is generally painful, and in some instances extremely so. The absence of complaints of sore throat in a child, or the fact of its swallowing without hesitation or apparent difficulty, is no proof that angina does not exist, since we have always found upon examination in a good light much greater redness than natural, and in many instances redness and swelling combined. As the eruption progresses, and the tongue loses its coat and becomes red, the inflammation of the pharynx usually augments; the redness becomes deeper, and the tonsils are more swollen and painful, and, in a good many, but not by any means all the cases, are dotted over with small white spots, or with thin, whitish, and soft false membranes. The throat affection, however, is rarely severe enough to constitute a serious danger in mild scarlatina, while in many of the malignant cases it is a frequent cause of a fatal termination. During the eruption, the nostrils are either dry and incrustated, or there is some coryza. The strength of the child is reduced for the time, but there are no signs of prostration, and the decubitus is indifferent. There is almost always more or less disorder of the nervous system, sometimes amounting only to headache and restlessness, while in other instances there is great irritability, wakefulness, and occasional mild delirium, especially at night.

Stage of Decline and Desquamation.—The eruption reaches its height, as already stated, about the third or fourth day, then remains stationary for one or two days, and afterwards declines gradually, so that no traces are left on the sixth, usually, or at most, in rare cases, on the ninth or tenth

day. In some very mild attacks, the whole duration of the eruption is not over two or three days. By the third day it has disappeared entirely. Such cases are not, however, very common. The other symptoms, both general and local, decline with the eruption; the pulse loses its frequency, and falls to the natural standard; the heat of the surface first subsides and then disappears, but the skin remains somewhat harsh; the redness and swelling of the tonsils and pharynx diminish; the spots of false membrane, if these be present, are thrown off; the deglutition becomes easy if it have been difficult, and soon all signs of throat affection vanish; the tongue cleans off, becomes reddish and glossy, and after a time returns to its natural state.

At the time of subsidence of the symptoms *desquamation* begins. It dates, therefore, in most cases from about the sixth day, though it may be either earlier or later. According to Hillier, the date of commencement varies from the sixth to the twenty-fifth day. It commences in most of the cases on the face and neck, though in a few instances it appears first on the abdomen. It then extends gradually over the body and becomes general. About the thorax and abdomen it occurs in the form of minute points, like those which result from the desiccation of sudamina; on the face it is in the form of thin light scales or squamæ, while on the extremities large flakes of the epidermis become separated from the derm, and are removed by the child, or rubbed off by his movements in bed. The whole process usually occupies some ten or twelve days, but may be prolonged into the third week, or even until the middle of the second month. It is generally accompanied by roughness and dryness, and some itching and irritation of the skin. Not unfrequently the surface beneath the exfoliation is left tender and irritable for some time afterwards.

During this period, dating from the sixth or eighth day, the urine becomes abundant, pale, of neutral or faintly acid reaction, and according to Gee, deficient in phosphoric acid.

It has been stated by some observers that albumen quite frequently appears in the urine during desquamation, and much more rarely during the stage of eruption, in short being unaccompanied by the signs of dropsy or of any special constitutional disorder. The proportion of cases in which this is said to occur varies from twenty-five or thirty to ninety per cent. in different epidemics. In such cases the albumen may disappear without causing any evil consequence, or it may continue or recur until, after a variable length of time, renal catarrh is developed.

Temperature.—The fiery redness of the surface, and the pungent character of the heat, have led to much exaggeration in the description of the pyrexia in this disease. The range of temperature is indeed from 100° to 105° , and only in rare cases does it reach 106° .

In 30 cases reported by Rinder (*Med. Times and Gaz.*, Feb. 15th, 1862), the temperature remained at the same point throughout the day in the more severe attacks; in slighter ones it fell in the morning and rose during the day, being most frequently at its highest point between 2 and 8 P. M. When the morning remission was marked, it indicated the approach of a favorable termination. The first decided fall of temperature, coinciding

with a diminution in the eruption, occurred in the majority of cases on the fifth day, or if not on this day, it was deferred usually until the tenth or fifteenth. In these latter cases, however, a fall of varying extent had occurred on the preceding fifth days. After the marked fall on the fifth, tenth, or fifteenth day, the temperature remains from 99° to 101° for a variable time, coinciding, when persistent, with continuance of the angina, or some of the other lesions of the disease. If at any time after the complete fall of the temperature there is any considerable elevation again, it indicates the development of some sequel, either an affection of the kidneys, throat, or one of the serous membranes. It is thus seen that the temperature in scarlatina tends to form arcs or cycles, usually of five days' duration.

Before quitting this part of our subject we must remark that, though the above is a correct description of the usual symptoms of mild cases of this disease, the reader would be greatly deceived should he expect always, and invariably, to find this exact train of morbid phenomena. On the contrary, the mild and the grave cases both vary so much, that it is impossible to describe them accurately by one or two portraits. Taking the above sketch as a standard of the mild cases, the observer will find that many fall short of it in all their features, while others deepen gradually in their shades, so to speak, until they pass into the grave type. Those that are milder in their type than the above sketch may be so to such an extent as to make it very difficult, and in some cases impossible, to determine positively whether the child has had the disease or not. Indeed, we doubt not ourselves that children sometimes have the disease so slightly that it is not discovered by either physician or parents, and, being protected by the attack, are in after-life classed amongst those insusceptible to the disease. Our grounds for this assertion are the facts that we have seen some cases so very mild that, but for the existence of the disease in other members of the family, they might have passed unobserved; and that in one instance we saw a patient, who had had the eruption for three days, and yet who was so slightly sick that he was sent from school, to which he had gone, for us to see. It was a well-marked case, and the child had no troublesome symptoms afterwards, notwithstanding the exposure he had undergone.

A still more remarkable case occurred to one of us two years since, which shows clearly how a child may pass through scarlet fever without its being recognized by the family.

CASE.—The mother of one of the families we attended consulted us about one of her sons, a sturdy boy of seventeen years of age. She stated that he had not been well for two days; that he had severe sore throat, was restless and feverish at night, could eat nothing, and complained of debility, but, as he was passing his examination at the High School of this city, he had refused obstinately to remain at home. It was agreed that he should call at our office the next morning, on his way to the school. When he arrived, which he did on foot, as usual, we found him covered with a copious, dark-red, perfectly characteristic scarlatinous eruption; his throat was very red, swollen, and quite painful; the pulse was over 120, active and full, and the skin hot. He was, of course, ordered home, there to remain until perfectly well. He recovered and had no drawback. The distance from this boy's home to his school was not less than a mile and a half, and he had walked this distance twice a day.

Grave Cases.—The following description of the symptoms of grave cases of scarlet fever is, like that which has just been given of the mild cases, drawn partly from books, and partly from our own observation of the disease. We have had the opportunity of carefully observing a very large number of grave cases of scarlatina, and we have preserved a more or less complete record of 61 such cases. We shall include under this division of the subject, as already stated, most of the cases usually classed by writers under the title of scarlatina anginosa, and all those generally described under the title of scarlatina maligna.

The symptoms which mark the *invasion* of grave cases of scarlet fever, though sufficiently alike in all to show the unity of the disease, differ very materially as to their degree of severity in different cases. In one set (rather less than a third, or 18 in 61, of our cases) they are most violent and dangerous, or, indeed, appalling in their character. From the first, they declare the imminent danger of the attack. In the second set (rather more than two-thirds, or 43 in 61, of our cases) they may be either evidently severe and dangerous, though not appalling, as in the first, or they may be much milder, more like those which mark the invasion of mild cases; but even under these circumstances they soon put on their grave and dangerous character.

The first set of cases, or those in which the symptoms are the most severe of all, and which include most of the malignant cases ordinarily styled ataxic, usually begin with nervous symptoms. The onset is in some instantaneous.

In one, the little patient, a girl two years old, whose brother and sister had been sick for some days with scarlatina, was put to bed in the evening in her usual health, which was strong and vigorous. She slept quietly through the night, but was found by the mother the next morning in a state of drowsiness, violent fever, and covered with a deep-red scarlatinous rash. She soon became comatose, and died on the third day. In another case, a boy eleven months old was a little fretful in the afternoon, but was put to bed in the evening as usual and went to sleep. About ten o'clock the nurse heard a rustling in the bed, and on going to it found him in a violent general convulsion. The next morning he was covered with a scarlet rash, which became deeper and deeper as the disease went on. On the second day he was nearly insensible, and had frequent attacks of convulsions; on the third day he had retraction of the neck, with spasmodic twitchings, and at the end of that day died in a state of coma. In a third case, a boy six years old, whose sister had been sick for a week with a mild attack, went to bed well. At three o'clock in the morning, he was seized with vomiting and purging, paleness and coolness of the skin, and great exhaustion. At nine o'clock he was drowsy and dull, the skin was pale and cool, and the pulse extremely rapid; the vomiting and purging had ceased; at 12 M. he was comatose and had a convulsion. From this time he continued comatose until he died at 6 P. M. of the same day, after an illness of fifteen hours. In a fourth instance, the invasion was that of croup; after a few hours coma and convulsions developed; patches of eruption then appeared on the trunk, and death occurred in twenty-four hours from the beginning. The subject of this case, a boy five years old, was thought to be so well in the afternoon of the day he was taken sick, that he had been sent out to visit a relation, and while there fell sick. In the fifth case the onset was sudden, with violent fever, drowsiness, deep suffusion of the skin, and in a few hours insensibility, general convulsions, and death in thirty-six hours. In a sixth, in a boy four years old, the attack came on with vomiting, pallor, drowsiness, and then a scarlet rash; after a few days, coryza and otorrhœa occurred;

the tongue and lips became cracked and dry; in the second week the child was comatose, with occasional attacks of extreme jactitation and the most violent hydrocephalic cries, which condition lasted ten days. After this came diarrhœa, extreme emaciation, loss of speech, and entire deafness. Gradually, however, the fever disappeared, the tongue cleaned off, and intelligence very slowly returned; in the sixth week convalescence was firmly established, and the child recovered perfectly with the exception of his hearing, which remained very dull in consequence of the perforation of both membranæ tympanorum. In a seventh, a girl eight years old, whose brother was then sick in the house with the disease, was in the morning well. At breakfast, she said she felt sick and soon went to bed. At 5 P.M. of that day she was attacked with a general convulsion which lasted about fifteen minutes. The pulse, immediately after the convulsion, was 150. At 11 P.M. she had another convulsion. Through that night she was very restless and wandering. On the morning of the second day there was a third convulsion, which, however, was very short. The pulse was now 160, small, and feeble. The patient was very heavy and dull, answering questions slowly and with great difficulty, and during part of the day she was comatose. On the third day she was better, the pulse having fallen to 152, and she was less dull, though she still continued very heavy and inattentive unless aroused by persevering efforts. The limbs were cool, while the head and trunk were hot. The eruption was thick on the trunk and upper part of the extremities; elsewhere it was scanty. Wherever it existed, it was of a deep-red or purplish color, and the capillary circulation was sluggish and imperfect. On the fourth day her intellectual condition continued better, but the extremities were still cold, and the lymphatic glands and subcutaneous tissues about the lower jaw and neck had begun to swell. On the fifth day, the swelling had become very great; the stupor had returned; a profuse and disgusting coryza and otorrhœa had set in; and the edges of the eyelids were inflamed and sore. On the sixth day the discharges from the mucous membranes of the head were very copious, and consisted of a thick, offensive, purulent fluid intermixed with dull whitish grumous particles. The patient was now comatose or very restless; she swallowed with great difficulty; the swelling under the lower jaw and about the throat was enormous; the pulse was rapid and small; the eruption was very dark in tint; the cutaneous circulation was slow; the extremities were cold, and death occurred about the middle of this day. In another case, the subject of which was a girl between three and four years old, the attack began with severe inflammation of the throat, causing great difficulty in swallowing. The rash on the first day was very extensive and of a deep-red color. The child was drowsy and heavy, or else delirious. On the second day she was comatose, and had strabismus and automatic movements of the limbs. On the third day the coma continued, and there were automatic movements of the extensor muscles, with retraction of the head. The eruption continued vivid, but was of a dark-red color. Death occurred in the middle of the fourth day, in a state of coma, without convulsions. In still another case, a boy, between eight and nine years old, was attacked suddenly, while in good health, with vomiting, sore throat, and high fever. Twelve hours after the onset, he had a severe convulsion, which lasted fifteen minutes. He soon recovered from this, however, and remained perfectly intelligent. On the second day the rash was moderate; there was violent fever, and the child was heavy, but, when roused, still intelligent. Early in this day a severe fit occurred. This was most violent, as severe as the worst epileptic convulsion. It lasted one hour and three-quarters. The pulse, after this, was 145. On the third and fourth days, the symptoms improved very much, the pulse having fallen to 125 and 132, but he continued drowsy and heavy. The eruption came out most abundantly. The fauces were very much inflamed, and somewhat ulcerated, and the external lymphatic glands were enlarged, but still the swallowing was not difficult. On the fifth he was not so well, being more restless and heavy alternately. There had now come on much difficulty in breathing, and some croupal sound. The latter symptom increased through the day, until the dyspnœa became very great. Deglutition now became excessively difficult; the external swelling increased; attacks of suffocation attended with the most

painful and distressing jactitation came on, and were renewed more and more frequently; and death occurred by asphyxia about the middle of the sixth day. In a tenth case, in a girl five months old, convulsions occurred on the second day. These were followed by coma lasting several days, and by enormous swelling of the lymphatic glands and subcutaneous tissues on the left side of the neck, and by a less degree of swelling on the right side of the neck. The glands of both sides suppurated and were opened, and the child finally recovered perfectly. In an eleventh case, in a boy seven years old, an attack of general convulsions took place on the third day, after which there were delirium and coma alternately for several days, with coryza, angina, and offensive otorrhœa, lasting in all six weeks. The child recovered, but remained deaf.

In this form of the disease, therefore, the symptoms are of the most virulent character. The onset is sudden. The child passes within a few hours from a state of apparent health, into one of the most extreme danger. Most of the cases begin with violent fever, and great depression of the strength. The pulse soon becomes very rapid (140, 150, 180), or so frequent that it cannot be counted, and it is at the same time small and often irregular. The skin is dry and burning hot in some parts, in others cool or even cold. There is generally nausea or vomiting, and these may be violent and constant. These are accompanied in some cases, but in our experience, only in the severest of all, by colliquative diarrhœa and meteorism. Delirium often exists from the first, or else there is drowsiness and dulness of intelligence, verging gradually into coma. In the most violent cases, the stupor or coma alternate with convulsions, which may cause a fatal termination in eighteen, twenty-four, or thirty-six hours.

When a case of this kind lasts over three, or even two days, the violence of the nervous symptoms almost always subsides; the convulsions cease to recur; the delirium is less violent; the coma gives way to drowsiness, or the patient becomes again quite intelligent and observant; the pulse often falls in frequency, and the heat of skin may diminish, and the eruption assume a more favorable appearance. All the symptoms seem, indeed, to be more promising, and very often both the physician and friends are greatly elated by the improvement in the patient's condition. Nor are these hopes always illusory, since children do recover occasionally even in cases that have exhibited the most threatening and malignant appearance at the moment of invasion. It happens, unfortunately, however, in a large majority of such attacks, that the improvement which takes place on the third or fourth day is only momentary. The nervous symptoms subside, but new phenomena make their appearance in the shape of severe inflammation, membranous deposit upon, and perhaps ulceration of the fauces, with extensive swelling and induration of the lymphatic glands and subcutaneous tissues about the angles of the lower jaw, and under the chin and throat. In connection with the throat affection which develops itself in this way, it is very common to have abundant purulent or membranous coryza, and often also otorrhœa. The symptoms assume, in fact, the features of the cases usually described under the title of scarlatina anginosa. As we shall, however, describe them directly in our account of the second set of grave cases, it is unnecessary to pursue the description at the present moment. We will state, however, before pro-

ceeding further, that the anginose and general symptoms which occur in cases beginning with violent nervous phenomena, and especially with convulsions, are nearly always of the most dangerous and malignant character, and usually end fatally in two, three, or four days after their appearance.

The eruption in this class of cases varies according to the violence of the attack. In the severest one that we saw, that which proved fatal in fifteen hours, no eruption whatever was perceived, and we only knew it to be scarlatina by the general character of the symptoms, and by the fact that a sister of the boy had been sick in the same house with the disease for a week. In the case which terminated in twenty-four hours, the eruption showed itself in the form of scarlet patches about the face and upper parts of the body, twelve hours after the onset. In a third case the eruption was moderate, but perfectly well marked and general. In the other thirteen cases, which lasted, with one exception, not less than three days, the eruption was entirely characteristic. It covered the whole surface, was at first scarlet in color, soon ran into a deep red, and then became violet or purplish. The exceptional case was one which lasted thirty-six hours, and proved fatal in that time. In this also, the eruption was well marked and extensive. M. Guérin (*Arch. de Méd.*, t. i, p. 292, 1842), in his account of the acute malignant form which he witnessed, states that the eruption was nearly constant. In all our cases it occurred within twenty-four hours from the invasion, while in those of M. Guérin, it appeared within twenty-four or forty-eight hours, or, as more frequently happened, not until the fourth or fifth day.

If no favorable change take place in these severe cases, and if they do not prove fatal at once, the patient grows weaker and weaker; the delirium continues, or is replaced by coma; subsultus tendinum, rigidity of the limbs, spasmodic twitchings or general convulsions, make their appearance; the eruption becomes more and more livid; the pulse grows smaller, more frequent, and irregular; the respiration is excessively embarrassed; deglutition becomes impossible; and the patient dies in from three to seven or nine days. In some few instances, the child struggles on for several weeks, and dies in a state of utter exhaustion, or having a constitution of great powers of endurance, at last surmounts the disease and recovers.

The invasion of grave cases is not always, as we have stated above, so violent as in those which have just been described. In rather more than two-thirds (43) of the 61 grave cases of which we have preserved notes, the onset was less threatening than in the other third, though the symptoms were severe and dangerous in most of these also, and when not so at the very start, very soon assumed the serious characters which make it necessary to class the cases in which they occurred as grave. The chief difference between the symptoms that mark the onset of grave cases of this kind, and of those in which the symptoms are still more violent, which latter we have thus far been describing, lies in the character of the nervous phenomena—in the latter most severe, threatening, and dangerous, consisting

of stupor, coma, or convulsions, and in the former, merely excessive agitation, restlessness, heaviness, or stupor.

In one well-marked case of the kind now under consideration, the patient, a boy between seven and eight years old, was attacked in the evening with headache, fever, and vomiting. On the following morning a faint rash was perceptible, which, by the afternoon of that day, was distinct, though not very full. The case now rapidly assumed unpleasant features. The pulse rose to 150. There was much drowsiness and delirium, and on the fourth day constant picking at the bedclothes and at the fingers. In another case, in a boy between four and five years old, the first sign of sickness was slight languor after dinner, which was followed by fever in the evening, and the development in the course of the night of a scarlatinous rash. On the following day there was some pain in the throat, with redness; the pulse was 140, and the skin hot and dry; there were no nervous symptoms, except slight drowsiness. On the third day the pulse was 136, the rash was well out, and there were no unpleasant symptoms whatever. From this time, however, the symptoms gradually grew worse; the throat affection increasing, the cervical lymphatic glands becoming very much swelled, and the child growing more uneasy and restless, though retaining perfectly its intelligence. By the sixth day, the grave character of the case was fully developed, the eruption being intense, and of a deep brick-red, verging towards a purple color. There was at the same time very great drowsiness, abundant discharges from the nasal passages of thick sero-mucous and purulent fluids, membranous exudation in the fauces, with gurgling and great difficulty in swallowing, and an utter loss of appetite. In a third case, a boy between one and two years old, was a little fretful in the morning, and was seized in the evening with vomiting and fever, and very considerable restlessness. On the next day he was covered with a scarlet rash from head to foot, and the skin was fiery hot. The pulse was 160, regular, not large. The child was very drowsy, dozing nearly all the time, but quite intelligent when aroused. The fauces were intensely red and rough, and the tonsils much swollen; there was very little external swelling. On the third day he was still very drowsy, and, when roused, less observant than before, though he still recognized persons. The pulse was 168, small, difficult to count, very hard, and corded. The skin, especially that of the limbs, was scarlet, very hot, and dry; the cutaneous capillary circulation was good. After this the symptoms grew rapidly worse; the pulse continued at from 148 to 168 on the fourth and fifth days, and on the sixth rose to 172, at which it stood a few hours before death. On the fourth and fifth days, he was still very heavy and drowsy, and so much so on the former as to take no notice whatever except when moved. On the fifth day, an abundant sero-mucous discharge took place from the nostrils; the cervical lymphatic glands, which had begun to swell before, now increased in size; there was some loud faucial gurgling, and the swallowing became difficult. On the morning of the sixth day, some of the symptoms improved so much as to flatter very greatly some of his attendants, who were unacquainted with the treacherous character of the disease. He roused up from his state of stupor, and noticed several things that were shown him, even taking them into his hand; but the breathing continued bad, the lymphatic glands were swelling rapidly, and had already become very large, so that they formed great projections on either side of the neck. The pulse was 155, and small. In the middle of the day the breathing became difficult, from the internal and external swelling, and from the collection in the fauces of thick and vivid phlegm. The surface had now become pale. The tumefaction about the neck was immense. Down the front of the neck and along its sides to the clavicles, a kind of œdematous swelling of great size had come on, and was rapidly increasing. The pulse was 160, small and feeble. The legs and arms were of a dark, congested tint. Deglutition was excessively difficult. In the evening the pulse was 172, and death took place just before midnight, with slight convulsive movements.

The mode of invasion is different, therefore, in different examples of the kind of grave cases now under consideration. In some it is even milder

than in any of those that have just been detailed; and it is not until the third, fourth, or fifth day, or even later, that the severity of the attack shows itself fully and unmistakably.

After the disease is once established, it will be found upon examination that the fauces are of a deeper red color, and that they are more swollen, than in mild cases. At the same time there is more difficulty and pain in deglutition, these are complained of by older children, and are shown in those who are younger by their refusal to swallow, by their crying upon making the attempt, and in some instances, especially at a later period of the sickness, by a positive inability to perform the movement. In nearly all these cases, false membrane is formed upon the mucous membrane of the throat. This is never, or very rarely, present on the first day of the attack. In most cases it is not found until the second or third, and often not before the fifth or sixth day. MM. Rilliet and Barthez state that they have known it not to appear until the tenth and eleventh days. It appears first in small, thin, whitish, yellowish, or ash-colored points or patches, on one or both tonsils, or on the soft palate only, where it remains limited, or from whence it extends to the pharynx, which it may cover in whole or in part. The patches are of variable thickness and consistence, and adhere sometimes very slightly, and sometimes with considerable tenacity to the mucous membrane beneath. They may remain for a day, and then be thrown off not to be again produced; or they may form in several successive crops, until the case is terminated; or, as most frequently happens, they last three or four days or more, and are then detached. The mucous membrane upon which they are seated is found in various conditions. It may present the redness and swelling indicative of severe inflammation, or it may be softened, ulcerated, and, according to MM. Guersant and Blache, gangrenous, though as a general rule, what have been supposed to be sloughs are in fact portions of altered false membrane. There is more or less fetor of the breath, sometimes amounting to a gangrenous odor, after the appearance of the pseudo-membrane. The severity of the symptoms is in proportion to the extent and thickness of the false membrane.

We have already seen that it is not uncommon to find ulcerations beneath the false membranes. In other cases of this kind the throat affection assumes very great violence without the presence of any exudation whatever. In some the mucous membrane is of a deep-red or even purplish hue, its consistence is softened, and it is swollen and covered with a layer of grayish or sanious pus. The tonsils are enlarged, infiltrated with pus, and softened. In other cases, in addition to the redness and softening, ulcerations are present. These may be superficial, amounting only to erosions, or they may extend through the mucous, and even submucous tissue to the muscles beneath. They are seated generally in the pharynx, but may exist also on the tonsils, and in some rare cases they extend into the larynx. In still more malignant attacks of the disease we find evidences of gangrene of the pharynx. It is important to distinguish between those in which the pseudo-membrane becomes so changed as to assume the appearance of sloughs, and those in which the tissues of the pharynx are

really gangrenous. The former constitute by far the greater number of the cases which have been generally regarded as instances of gangrene of the throat. That gangrene of these tissues does actually occur in some few cases, is proved, however, by the evidence of Dr. Tweedie, who says (*op. cit.*, p. 650) that in malignant scarlatina "the membrane of the pharynx is sometimes of a dark, livid color, and occasionally in a sloughing state," and by that of MM. Guersant and Blache, who state that they met with several instances of gangrene of the pharynx in the pseudo-membranous angina which prevailed in 1841.

An almost constant accompaniment of cases of this kind is inflammation and swelling of the submaxillary lymphatic glands and surrounding cellular tissue. The tumefaction is generally confined at first to the glands beneath the jaw, which become painful to the touch. After a short time it extends to the parts behind the angle of the jaw, and beneath that bone, until at last the sides of the neck and the throat are greatly swollen, so as to interfere with, or even prevent in large measure, the opening of the mouth, and by the pressure exerted on the internal parts of the throat, to add to the difficulty of deglutition which already exists. In some cases the pressure is so considerable as to embarrass the respiration of the child. This swelling has been generally supposed to depend on inflammation of the parotid glands; but MM. Bretonneau, Guersant and Blache, and Rilliet and Barthez, all state that parotitis is of exceedingly rare occurrence, and that the swelling in question depends nearly always on the causes just described. The last-named writers state, moreover, that the tumefaction of the cellular tissue is often of the nature of active œdema. The swelling of the cervical lymphatic glands, and of the cellular tissue of the sides of the neck, and that under the throat and chin, seldom takes place to any considerable extent, according to our experience, prior to the third or fourth day. During the first two or three days the chief symptoms are the fever, the eruption, and the nervous phenomena, which latter consist in this class of cases, of either excessive agitation and restlessness, or of drowsiness or stupor. Very often, after a child has seemed to be very ill for two, three, or four days, from the violence of the febrile reaction and the severity of the nervous symptoms, it will appear to improve very decidedly on the third or fourth day, and thus lift up the hopes of those interested in it. It is just at this time, however, that the throat affection is apt to set in severely, and it rarely fails to come in children who have presented violent symptoms during the first three days. The enlargement generally disappears, in favorable cases, in from three to twelve days, by resolution, while in others it terminates by suppuration of the glands and surrounding parts.

In the form of the disease we are now considering it is common to observe violent *coryza*, which may be either purulent or pseudo-membranous. It may appear from the very first, or not for several days after the eruption has commenced. The discharge is yellowish, granular, thin at first, and afterwards thick; it contains often flakes and shreds of exudation, and becomes sometimes very offensive, and highly acrid, so as to excoriate the upper lip. It often flows in surprising quantities, and gen-

erally continues up to the moment of death, or until all the symptoms have moderated.

Otorrhœa is another symptom of this form. It is apt to occur simultaneously with coryza. The discharge is at first thin and watery, like that from the nostrils, but becomes thicker as the case advances. The quantity is extremely variable. In some cases we have known it to fill the meatus and concha of each ear, and then to flow out and make large stains upon the pillow, or to collect very rapidly after being wiped away. It is, like coryza, an unfavorable symptom, as it is a mark of the grave form of the disease, and because, if the child recovers, it is very apt to result in deafness, which is but too often permanent.

These symptoms, coryza and otorrhœa, sometimes exist also in mild cases, but they do not then assume the peculiar characters which they present in grave cases. The discharges are much less abundant, and the mucus or pus is healthy, and scarcely offensive to the smell; they last but a short time, and are very rarely accompanied at the time or followed by more than a slight degree of deafness.

The eruption is generally stated to appear later than in mild cases, and often to be less vivid and less extensive. It is also said to occupy only portions and not the whole of the body, to occur in irregular patches, or to appear and disappear alternately. This has not been the case in the instances which we have seen. In all but two of the forty-one, the eruption occurred early, generally within twenty-four hours from the onset. It was of a deep brick-red or livid color, and covered the whole surface. In one of the exceptional cases it did not take place until the seventh day, when it appeared in patches on the wrists and knees. On the eighth day, it extended to the rest of the extremities and abdomen, and on the ninth was general and of a rather dark hue. In the second exceptional case the eruption did not appear until the second day. It then came out over the whole trunk, and to a moderate extent upon the limbs also. In this, as in the previous one, it was dark in its tint. In three other cases it was quite moderate in amount, but general and well marked.

The *general symptoms* are more severe in grave than in mild cases. It sometimes happens that for one or two days, or even longer, the case promises to be mild, but then suddenly assumes the threatening features of the form under consideration. The fever is usually intense, the pulse being full and strong, and rising very soon after the onset to 140, 150, or 170; the skin is very hot and dry; there is more restlessness and irritability than in the mild form, and after one, two, or three days, appears a strong disposition to delirium and stupor, not unfrequently merging into coma. The respiration is accelerated, and in many instances, owing to the throat affection, labored and difficult. In most of the cases, a loud gurgling, which is very characteristic, is heard in the throat, particularly when the child is asleep or dozing. This depends in part upon the collection of viscid and tenacious secretions in the fauces,—which sometimes embarrass the respiration so much as to make it necessary to remove them with a mop,—and in part upon the existence of the coryza of which we have spoken. The coryza is a symptom of very serious consequence at

all ages, but especially in young children. There is generally some cough, which may be frequent and troublesome, though not usually so unless there be a disposition to laryngeal complication. The voice is hoarse, guttural, and sometimes whispering. When the cough is very frequent, and still more, when it becomes hoarse and croupal, in connection with hoarse or whispering voice, or aphonia, there is great reason to fear the extension of the exudation into the larynx, which constitutes an almost necessarily fatal accident. The face is deeply flushed at first, and the expression anxious. If no improvement take place, the case assumes in four or five days, or even less, a still more threatening aspect. The pulse becomes very rapid and small; the restlessness and delirium pass into drowsiness or coma; the tongue becomes brown and dry; the teeth are covered with sordes; the lips are dry, cracked and bleeding; diarrhœa is apt to occur; and the patient dies in from three to ten days, in a well-marked typhoid condition. In other instances, on the contrary, the case runs on from week to week, and at last, after an illness of four, five, or six weeks, the child either dies, or recovers after all chances for life seem to have been lost.

In order to show in their natural connection, the different symptoms that have just been described, we will cite the following abstract of three of our cases:

The first occurred in a boy between seven and eight years old. On the fourth day of the attack the pulse was 150, and the fauces presented flakes of false membrane. The fauces were very much swollen, and deglutition became difficult; faucial gurgling came on, and the throat was filled with viscid and tenacious secretions. The nasal passages now became occluded by constant discharges, at first mucous and then mucopurulent, with admixture of membranous flakes. From the fifth to the ninth day there was an excessive fetor from the nose and mouth. The lymphatic glands just beneath the ear swelled very greatly, so as to extend much beyond the line of the inferior maxilla. The tongue and lips became dry and cracked, the teeth were covered with sordes, and the angles of the eyelids inflamed, and then ulcerated. On the sixth, seventh, eighth, and ninth days, there were taken away from the mouth and throat of the child, with a mop, hard and most offensive masses of dried-up mucus and incrustated epithelium, enveloped in thick, gluey, dark-colored mucus. These masses stuck to the fauces, tongue, and lips so tenaciously, that they could be removed only by means of a mop, the boy himself being quite unable to detach them. On the seventh, eighth and ninth days, though the cervical lymphatic glands were very much swollen, the patient was better. The pulse came down gradually from 152 to 132, 128, and 112, and the swallowing improved so much that the child could take liquids with less convulsive effort, and could drink continuously. The drowsiness diminished, and the delirium ceased. On the eighth day a slight erythematous redness appeared on the bridge of the nose, and extended towards the malar bones. The skin of the face and eyelids became somewhat swollen and puffed by an œdematous effusion. On the ninth day the pulse was down to 104, and the skin was nearly natural as to temperature. The swelling was very great on both sides of the neck, and the glands on the right side were red on the surface, very hard, and quite painful. The swallowing was much easier for drinks, but as yet no solid, not even the very softest kind, could be taken. On the fourteenth day from the onset we opened a very large abscess on the left side of the neck, which discharged abundantly a healthy and laudable pus. On the fifteenth day we opened a still larger abscess on the right side, and after this, perfect recovery took place.

In another example, which has been alluded to already, occurring in a boy between

four and five years old, the gravity of the case did not show itself clearly until the sixth day. On the evening of that day the pulse was 128, the skin very hot and dry, and there was an intense eruption of a brick-red color. There was, at the same time, great drowsiness, and utter loss of appetite. Deglutition was difficult, and there was a loud faucial gurgling during sleep. There was now also a considerable amount of membranous exudation in the fauces. During the seventh and eighth days, the boy continued very sick. He was drowsy, almost comatose; the eyes were half open and the conjunctivæ minutely injected; there was an abundant coryza, the discharges being composed of offensive mucous and sero-mucous fluid, with an admixture of pus and of flocculent or grumous particles, the latter consisting evidently of broken-down membranous exudation. There was no otorrhœa. The pulse rose from 120 to 128. During the night of the seventh day the anginose affection was so severe that the child could swallow nothing from 10 P. M. to 3 A. M.; fluids poured into the mouth ran out again in part, and were in part returned through the nostrils. On the tenth day there was still no decided improvement, except that the pulse had fallen to 112. The coryza continued as before; the fauces were covered thickly with whitish exudation; the deglutition was a little easier. The drowsiness continued, as the child dozed nearly all the time, merely rousing from time to time to take drinks, and then, in spite of all solicitation, sinking into sleep again. The abdomen was tympanitic. The urine was rather free, more so than it had been before, and it was also clearer and of a lighter color. By the twelfth day there was a decided improvement; the pulse had fallen to 106, and the child was not quite so heavy. The act of swallowing was easier, and the fauces showed less of the plastic exudation, but they were still very much coated with tenacious mucus. On the thirteenth and fourteenth days the patient continued to mend. The pulse fell to 98 and 92; the fauces had become clear of the exudation, and presented instead an excoriated and ulcerated appearance. The secretions into the fauces were less viscid and less copious. The coryza had diminished, and the discharges had become first muco-purulent and then mucous. The drowsiness had diminished, so that he woke spontaneously and began to ask for his toys. He now began to demand food, but refused to eat when things were brought to him. On the fifteenth day he was extremely irritable, screaming most violently from the slightest causes. On the sixteenth day the pulse was 92, and the skin nearly natural as to temperature. He was now exceedingly emaciated and very weak. The orifices of the nasal passages were very much irritated and incrustated, but there was scarcely any coryza. The tongue was clean, pink in color, and moist, the thirst not too great, and there was a little appetite. The temper was improving. From this time forward the child improved steadily but slowly, so that he sat up for the first time on the twenty-seventh day. He was as much emaciated at that time as after violent typhoid fever.

The reader must not, however, suppose that all grave cases present throughout their whole course, symptoms so dangerous as those which marked the two examples that have just been detailed. In some, on the contrary, the symptoms, though of such a character as to deserve and require the title of grave, are of a much milder kind. In order to make this part of our description of the disease as clear as may be, we will relate the following as an example of a grave case in which the symptoms, though severe, were neither malignant, nor at any one time very dangerous to life:

A girl between seven and eight years old was well at breakfast. In the course of the morning she complained of sore throat, and of not feeling well, and at 4 P. M., when we saw her, was quite feverish, with a frequent pulse and hot skin, and showed already a well marked but rather faint scarlet rash upon the trunk of the body, and about the elbows. On the following day the trunk and upper parts of the limbs were covered thickly with an intense eruption, of a bright scarlet color. The fauces were

very red, somewhat roughened, and a good deal swollen. The only nervous symptom present was severe frontal headache. There was no unusual agitation, no drowsiness, and nothing like convulsive movement. On the evening of this day, the pulse had run up to 168, and was rather full, but not hard. The skin was exceedingly hot and burning; during the night there was great restlessness, and the child was wakeful and occasionally delirious. On the third day the symptoms continued much the same, except that the pulse was down in the morning to 152, that the rash had extended to the hands and feet, and that some small spots of whitish exudation were now visible on each tonsil. On the night of this day the fever again increased very much, and the child was again delirious. On the fourth day the pulse was 148; the exudation had increased so much as to cover a good portion of both tonsils, and it had extended also in a slight degree to the posterior wall of the pharynx. There was now a considerable enlargement of the lymphatic glands situated at the angle of the jaw on the left side, and a smaller one on the right side. Deglutition was somewhat painful, and a little difficult, but not seriously so. The case continued in much the same way until the seventh day, when the pulse had fallen to 132, and the eruption had faded very much on the trunk of the body, and, to a considerable extent, upon the limbs also. The fauces now exhibited the false membrane over the whole of both tonsils, over the half-arches, the sides of the uvula, and upon the upper portion of the posterior wall of the pharynx. Instead of being whitish and clean-looking as at first, however, the false membranes now looked exactly like sloughing portions of the mucous membrane. They were of a dirty-brown color, softened, and seemed to be detaching themselves like sloughs from the tissues beneath. On the ninth day the patient was much better, the pulse having fallen to 116; the eruption had almost wholly disappeared; the heat of skin was very much reduced; the dark-colored portions of false membrane had disappeared from the fauces, leaving the mucous membrane beneath red, excoriated, and in parts ulcerated. On the thirteenth day the child was convalescent, the pulse having fallen to 96, the heat of skin having disappeared and the throat being nearly well. The appetite had returned, the temper was serene and cheerful, and the patient was, in fact, well, with the exception of weakness, and some remaining soreness of the throat.

Laryngitis has been supposed by some persons to be of frequent occurrence in the course of the disease, while others assert that it rarely, if ever, occurs. M. Bretonneau has never met with it. M. Rayer says he does not know that the exudation has ever been found in the larynx or trachea. Tweedie (*Cyclop. Pract. Med.*, Art. Scarlatina, p. 640) states that in the dissections he has made he has not seen an instance of the membranous exudation extending into the larynx. That it does sometimes occur, is proved nevertheless, beyond a doubt, by the evidence of MM. Guersant and Blache, Rilliet and Barthez, and others, and by our own observation. MM. Rilliet and Barthez report three cases in which it was found in the larynx after death. These gentlemen state, however, that they have never observed the peculiar symptoms of croup. This does not accord with our own experience; for in several cases that we have seen, all the peculiar symptoms of the malady were present during life.

The subject of one of these cases was a boy two years of age. A few days after the invasion of the disease, a severe and extensive pseudo-membranous angina was developed. This was soon followed by all the symptoms of croup: hoarse cough, stridulous respiration, weak, feeble cry, dyspnoea, and whispering voice, which lasted about five days, when the angina and croupal symptoms both diminished very much, and the child seemed in a fair way to recover; suddenly, however, extensive tumefaction of one side of the neck took place, and he died in twenty-four hours. Unfortunately no

examination could be made. In another case, in a child between six and seven years old, who had a most violent attack of the disease, severe croupal symptoms set in on the eighth day. They consisted of harsh, croupal cough, stridulous respiration, and great difficulty in swallowing, and the act of swallowing occasioned much harsh cough and strangling. The symptoms continued on the ninth day, after which they moderated, and the child finally recovered entirely. In a third case, also a violent one, in a boy between eight and nine years old, and in which general convulsions occurred on the first and second days, the symptoms had improved a good deal on the third or fourth day. On the fifth day he was not so well, being more restless and heavy, and having much difficulty in breathing, with some croupiness of sound. These symptoms increased rapidly until they gave rise to most violent fits of suffocation, and caused a fatal termination on the sixth day. In a fourth case, in a child nine months old, death occurred on the thirteenth day from laryngitis, occurring in connection with membranous angina. The fatal termination was preceded by hard, dry, and croupal cough, stridulous respiration, and great difficulty of deglutition. In a fifth, in a child under a year old, croupal symptoms made their appearance on the sixth day, the fauces being at that time covered with membranous exudation, and they caused or assisted to cause a fatal termination on the eighth day. In yet another case, the subject of which was between one and two years old, a grave attack of scarlet fever was entirely recovered from. At the end of the second week the child was seized, owing to improper exposure in a cold house, against which the parents had been properly warned, with anasarca. This also was recovered from, and again the parents were warned against improper exposure. On the very day after our last visit, however, the child was taken down stairs into a room with the windows open, and this on a mild day in the month of February. The child was seized now with diphtheritic angina, and died, after a few days, of croup. This was in the fourth week from the outset of the scarlet fever. In a seventh case, severe from the beginning, the patient recovered so as to be apparently out of danger, but owing to the room being very cold from the fact that it was large, with wide rattling windows down to the floor, and from the fire being too small the child took cold, and, at the end of the third week, was seized with severe croup, which had many of the features of membranous croup, but which was, in all probability, spasmodic croup, dependent on ulcerative laryngitis. The case continued seven days, during which time the patient was violently ill, but finally, after a most dangerous struggle, it ended favorably.

The symptoms which indicate a disposition to implication of the larynx are frequent, hoarse, and croupal cough, hoarse and whispering voice or cry, aphonia, and dyspnoea with stridulous respiration.

The *duration* of grave cases of scarlet fever is very uncertain. In some the disease runs its course with frightful rapidity, destroying life within a few hours or days. In others, though the symptoms of the early stage may seem to be as violent as in those where death occurs in a very short space of time, the patient either lingers for several days or two or three weeks, and then dies, worn out by the violence or malignancy of the attack, or else, after a most dangerous and apparently desperate illness, he finally struggles through and recovers.

In the most violent of the grave cases, those which we described first as forming a separate group, 18 in number, of which 13 proved fatal, the duration in the fatal cases was between eighteen hours and six days. Of the 13, 2 proved fatal in eighteen hours, 1 in twenty-four hours, 2 in thirty-six hours, 4 in three days, 1 in four days, 1 in five days, and 2 in six days. Of the 5 favorable cases, 1 lasted three weeks, 1 four weeks, 2 six weeks, and 1 two months.

Of the less violent of the grave cases, 43 in number, 15 died, and 28 recovered. Of the 15 fatal cases, 1 died in four days, 2 in five days, 2 in seven days, 3 in eight days, 2 in thirteen days, 1 in fourteen days, 1 in fifteen days, 2 in four weeks, and 1 in five weeks. Of the 28 favorable cases, the duration of the shortest was seven days. The remainder lasted from twelve days to six weeks, the most common period being between three and four weeks.

COMPLICATIONS AND SEQUELÆ.—*Dropsy*.—This is the most frequent and important sequela of the disease. In the vast majority of cases, when dropsy appears as a sequel to scarlatina, the urine will be found to present all the characters present in acute Bright's disease; and yet there are some high authorities (Simon, Becquerel, Philippe, Rayer) who assert that marked dropsy may occur without the slightest albuminuria. It is possible that some of these cases may be explained on the supposition that the urine has only been occasionally examined, and that albumen may have been temporarily present and overlooked; but it seems undeniable, that, in some instances also, dropsy may appear without any abnormal condition of the urine whatever. It is probable that, in these cases, it depends upon an anæmic state of the blood, developed during the course of the disease. We have never met with dropsy following scarlet fever in which we did not find albumen, and, in only too many instances, it is not only discoverable, but it is in much larger proportion than in most renal diseases unconnected with scarlet fever.

The frequency with which dropsy is developed varies greatly in different epidemics and in different forms of scarlatina. It occurred in a fifth of the cases of MM. Rilliet and Barthez, and in 31 of the 274, or in about a ninth, of those observed by ourselves of which we have kept notes. It occurs generally in the course of the second or third week of the disease, and during the process of desquamation. It is thought to follow cases of moderate severity much more frequently than those of a grave character. Dr. Tweedie states that it has never been observed to succeed a malignant attack. This does not, however, accord with our own experience, since of the 31 examples that we have seen, 8 occurred in grave cases of the disease. Still it may be said on the whole, that the susceptibility to renal disease bears an inverse proportion to the activity and complete development of the scarlatina. The effusion may attack any one of the cavities, or the cellular tissue of the body, or all at once. The most common form in which it appears is anasarca, after which the most frequent are, in the order in which they are mentioned, œdema of the lung, hydrothorax, ascites, and hydropericardium.

The *exciting cause* of the dropsy is generally believed to be cold, contracted usually by exposure to air and moisture at too early a period. We have rarely known it to occur when the patient has been confined to the chamber or house until after the twenty-first or twenty-eighth day; while, on the other hand, we have seen it follow immediately upon a ride in cool weather on the fourteenth day, the child having been convalescent for several days before. We have known it to occur also when the child has been allowed to run through the house exposed to draughts from open

doors and windows. We have been able, in a number of instances, to trace it directly and obviously to cold. Thus, in one very marked example, a boy between six and seven years old had had a mild attack of the disease, and was so entirely recovered that we ceased our visits on the tenth day, leaving strict injunctions with the mother as to the necessity of confining the child to the house for at least ten days longer. On the fourteenth day he was allowed to sit for fifteen minutes, late in the afternoon of a cool April day, on the marble front-door step. He was seized that night with fever and vomiting, had anasarca next day, and, during an illness of two weeks, had dropsy of the pericardium, effusion into the right pleural sac, ascites, and some signs of uræmia. In another case, a boy eleven years old had recovered entirely of a mild attack. He slept in a room heated by a stove. On the nineteenth day, the weather being cold, he got up early in the morning to light the fire, which had gone out accidentally. He was attacked that day with bronchitis, and was, on the following day, anasarcaous. In another instance, anasarca was produced at the end of the third week, the child being quite well previously, by his being taken into a cold room to sleep. We could cite other instances of the same kind, but these are enough. It is sufficient to say that in a large majority of the cases that we have seen, it has manifestly and obviously followed improper exposure during the second or third week. In a few cases, however, it has come on without any imprudence whatever, and we have been entirely unable to ascertain the cause. It has been doubted by some whether the action of cold will cause dropsy, unless the urine have been already albuminous. We have no doubt on this point ourselves. We have too frequently seen children who were, unless all signs fail, entirely convalescent, attacked by acute renal catarrh directly after exposure to cold, to have any doubts as to the sequence of events.

We are in the habit now of always directing the mother or nurse to keep the patient confined to the chamber for four weeks from the onset of the disease, or, if it be allowed to run through the house, to take care to have it well clothed, and to keep the windows and doors carefully closed should the weather be cold or cloudy. This rule is one of the most important of all in the care of the disease. It ought to be insisted upon in all and every case occurring in the cool season of the year.

The question was formerly much discussed, whether the condition of the kidney which accompanies scarlatinal dropsy was one of the forms of Bright's disease. Dr. Johnson suggested that it was a peculiar affection of these organs, characterized by a desquamation of the epithelium of the tubules, for which he proposed the name of desquamative nephritis. Recent observations have, however, shown that there is in reality nothing specific in the lesion, but that it is identical with other cases of renal catarrh or tubal nephritis, to use the excellent name bestowed by Dickinson, occurring from whatsoever cause. Indeed, it may be said that in almost 75 per cent. of all cases of chronic renal disease in children, the cause of the affection has been scarlatina, and the form of the lesion is that which we have above mentioned.

Various causes have been assigned for the frequent development of tubal

nephritis in the course of scarlatina. Thus it has been supposed that the affection of the kidneys resulted from inaction of the skin, owing to the intense congestion which attends the eruption; but clinical experience shows that it is precisely in cases where the affection of the skin is most intense that the kidneys are least disposed to disease. It would rather appear that when the action of the virus is not fully determined to the surface, violent congestion of the kidneys is established, which, especially when the patient is exposed to the action of cold, may result in the development of tubal nephritis.

Morbid Anatomy of the Kidneys.—When death occurs in the acute stage of the renal disease, the kidneys are found enlarged and very heavy. The surface is smooth and injected; on section, the organ drips with blood; the Malpighian bodies are congested, and appear as red dots; and the vessels of the cortex and cones are gorged with blood. The tubules are distended with granular epithelium, granular matter, or fibrinous plugs. The cortex appears coarse-grained, and presents intermingled dots or streaks of red and buff color. In the more chronic form of the disease, the kidney is also much enlarged and very heavy, its surface smooth and pale, or dotted with congested stellate vessels. The capsule is not thickened, and is readily removed. On section, very little blood escapes; the cones retain their pinkish or red color; while the cortex is coarse-grained, thickened, and of a peculiar opaque white color. The Malpighian bodies may be distended, owing to obstruction to the escape of the blood. The principal lesion, however, is still found within the tubules, which are stuffed with epithelial cells, or with granular matter resulting from their disintegration; occasionally, clear fibrinous plugs are also seen occupying their calibre. It frequently happens that the epithelium undergoes fatty degeneration, and when this is marked, the cortex acquires a yellowish tint. According to Dickinson, there is less tendency to this change in tubal nephritis following scarlatina than when it follows other causes; a circumstance which he thinks may possibly account for the comparatively curable nature of scarlatinal dropsy.

The *dropsical symptoms* usually show themselves in the third or fourth week of the disease, and are generally preceded for a few days by albuminuria. In most of the cases that we have seen they occurred in the third week, but they sometimes appear at the end of the second, and sometimes not until the fourth week. In one case they showed themselves first on the thirtieth day, after the child had been exposed to too cool a temperature in an insufficiently warmed room. They occur, therefore, as a general rule, during the stage of desquamation. The attack is sometimes very sudden, but in most instances it is slow and gradual. The effusion is not commonly the first symptom observed. On the contrary, the dropsy is almost always preceded for one or two days by the signs of a more or less considerable constitutional disturbance. The patient has usually passed safely through the eruptive stage of the fever, and has been considered for several days as convalescent, for, as we have already remarked, the dropsical affection is more rare after grave than after mild cases. The child has perhaps been running about the house, or it has

even been out, the parents supposing, unless warned by the physician, from the disappearance of the fever and other symptoms of illness, and from the return of appetite and gayety, that complete recovery has taken place. We have seen a few cases, however, in which, without any suspicion of exposure or negligence, for the children had not been out of bed, much less out of the room, the renal disease made its appearance in a most treacherous way. In one family, a brother and sister, six and eight years old respectively, occupied separate beds in a large, thoroughly well-ventilated chamber. They had had the disease decidedly, though in a mild form. The boy was the elder of the two and robust. The girl was small, delicate in appearance. After convalescence was well established, the boy was allowed to be up and about the room; the girl was kept in bed, because of her supposed delicacy. After two days the boy lost some of his vivacity and appetite; his urine was examined, and was found to contain a notable quantity of albumen. He was put in bed again at once. A few days later, the girl, who had not been out of bed, and who seemed quite well, showed some slight signs of indisposition. Her urine was examined, and found to be in the same state as her brother's. These children were at no time in any danger, and yet the urine of both remained unhealthy for a year, showing albumen and tube-casts in gradually diminishing amounts at each examination. They both recovered and are living now (1881), ten years after the illness, in excellent health.

Generally, however, it happens that after some exposure the child becomes drooping, languid, and irritable, or uneasy, peevish, and restless. Simultaneously with or very soon after these symptoms, fever sets in; the skin becomes dry and heated, and there is usually an elevation of the temperature to the extent of 4° or 5° ; the pulse is frequent and hard, or it is frequent and jerking; the appetite is diminished or lost, and there is more or less thirst; the bowels are generally constipated; the urine is usually diminished; and there is not unfrequently some nausea or vomiting, and complaints of headache.

The symptoms which precede the appearance of the effusion are not always, however, so marked, and in other instances are scarcely noticeable, and yet the effusion may take place suddenly, and, affecting the subcutaneous cellular tissue and different internal organs simultaneously, may cause a fatal termination with frightful rapidity.

The effusion usually commences in the face and may be very slight, leaving us in doubt even whether there is really any, or it may be very large and disfiguring. The swelling is most marked about the eyelids, which look puffed, and it may be confined entirely to them, or, at least, it may be only in them that we can feel sure of its existence. From the face it extends to the hands and feet, and either remains limited to these parts, or spreads over the whole surface, and gradually or rapidly to the internal organs. The skin over the parts in which the effusion has taken place is firm, hard, and elastic to the touch; it does not generally pit, at least not in the early stage, and it is of a dull white color.

In very mild cases the constitutional disturbance is usually but slight, and the effusion may be so small as to leave us in doubt as to the cause

of the sickness. Generally, however, we have been able to determine the cause of the fever by a careful examination of the face, and particularly of the eyelids, which look a little swelled and distended, and by the presence of a slight puffiness or cushiony appearance of the backs of the hands and feet. In such cases the general symptoms usually pass away after a few days; the urinary secretion, which had been diminished in quantity and of a deeper color than natural, becomes again healthy; the anasarca disappears, and the child returns to its ordinary condition. In more severe cases the general symptoms are all more marked; the anasarca is more extensive and the swelling more considerable; the child, if old enough to describe its sensations, may complain of pain in the back, though we believe this to be rare, and the lumbar region may be tender to the touch; the urine exhibits much more marked changes in its character; but still, unless some important internal cavity be attacked, the symptoms diminish after a week or ten days, and the child recovers gradually. In still more violent cases, the amount of the effusion is very large indeed, the face is disfigured by the swelling, the limbs are largely distended, the cellular tissue of the trunk of the body is infiltrated, the quantity of urine discharged is very small or the secretion is arrested entirely for one or several days, and the fever is high. If the disease be not removed, the effusion may extend to the internal organs; to the lung, producing œdema of that organ, to the pleural sac, causing hydrothorax, to the pericardium, to the peritoneal cavity, or to the brain. Death may occur in these violent cases from asphyxia occasioned by œdema of the lung, by hydrothorax, or by the obstacle to the circulation caused by the presence of the effusion in the pericardium; from hydrocephalus, or, finally, the patient may sink into a comatose state like that which often precedes the fatal termination of Bright's disease in the adult, and due, like that, to uræmia.

It sometimes happens, as was stated above, that death occurs with very great rapidity. MM. Guersant and Blache have known it to end fatally in twelve, fourteen, and thirty-six hours. In a case that came under our own observation in consultation, a child between one and two years old, who had had a very mild attack of scarlet fever, was seized suddenly towards the end of the third week, after it was supposed to be quite well, and after exposure to draughts of cold air in the lower room of a small house, with vomiting, and shortly afterwards with convulsions and coma, which terminated fatally in thirty-six hours. In another case, in a boy between thirteen and fourteen years old, who had had a mild but well-marked attack, and who had convalesced, and been out of bed for a few days, fever with a slight headache, and diminution of the urine, came on at the end of the second week. After two days of slight ailment, without any signs of anasarca, he suddenly, without any warning, fell into violent convulsions, which were repeated frequently, with lulls of imperfect consciousness, for a few hours. After twelve hours he became completely comatose, with occasional convulsive seizures, and died at the end of eight hours more.

According to Gee (*loc. cit.*), uræmic convulsions and coma are not fre-

quent in the course of scarlatinal dropsy, nor are they of such fatal import as in acute Bright's disease in the adult. We have, however, seen quite a number of convulsions, and advise the practitioner to be very careful as to his prognosis. It is true, probably, that they are less fatal than in adults, but so also in acute renal catarrh (not scarlatinous) from exposure to cold.

The symptoms which mark the occurrence of internal effusion will depend of course upon the part attacked. In one case they will be those of œdema of the lung, in another of hydrothorax, and in another of hydro-pericardium or ascites.

Urine.—The particular condition of the urinary function is next to be described. It has already been stated that the amount of urine secreted is less than natural during the early period of the dropsical attack. But, at the same time, the patient generally voids the secretion more frequently than usual. There is in fact micturition, occasioned no doubt by the irritating character of the urine, which causes the bladder to contract and expel that fluid so soon as even a small quantity collects. The diminution in the amount of the secretion is usually a very marked symptom. It is sometimes almost, or even entirely suppressed for a considerable period. In one case that occurred to one of ourselves, in a boy between one and two years old, there was no discharge whatever for a period of thirty-six hours. During this time there was no distension of the bladder, as we ascertained this point by careful palpation and percussion. In another case, which occurred in a girl between three and four years old, and who was nursed by her grandmother, one of the most accurate, reliable, and experienced nurses in the city, we were assured that there was no discharge whatever of urine for five days in succession. During the suppression there was no accumulation in the bladder. On the contrary, the hypogastric region was flat, depressible, and sonorous on percussion. The patient was very ill during all this time. She was feverish and passed nearly the whole time in a semi-comatose state, but could be roused with much effort, so as to show some intelligence; she rejected by vomiting almost everything that was given her, and complained when aroused of severe headache. She had no convulsions nor any convulsive movements, and finally recovered as the kidneys regained gradually their secretory function. In many other cases that have come under our observation, especially those which we have seen in later years, when we have watched this symptom more carefully, the diminution of the urine had been very great, so much so as to constitute a marked and important symptom.

In mild cases, when the diminution is not very marked, the urine is of a deeper color than natural, but retains its transparency when first voided. It is apt, however, to become turbid on cooling, and to deposit a more or less abundant precipitate of urates. Its reaction is acid; its specific gravity increased in proportion to the concentration; and urea and the chlorides are much diminished. Albumen is present, and microscopic examination shows epithelial or hyaline casts of the renal tubules, renal epithelium, and blood-globules. In more severe cases, the urine is very much diminished in quantity, the color is either a very dark red, or has a

blackish or brownish tint, or is like smoke or soot; the specific gravity is very high; the amount of albumen large, and the precipitate contains many casts and blood-globules, mixed with abundant urates.

The amounts of albumen and blood bear no definite relation to each other; in some cases, the albumen may be abundant without any blood being present; while in other cases, with a large precipitate of blood globules, the urine may contain but a moderate amount of albumen.

Basham calls attention to the occasional development of a bluish-green, and subsequently greenish-black color, on the addition of nitric acid to the heated urine, as a sign of very grave augury, being associated with extensive and advanced renal disease.

The duration of this stage of diminution of urine varies greatly in different instances, and is, to a certain extent, indicative of the future progress of the case. It is succeeded by a stage in which the urine becomes abundant, even exceeding the normal amount, the specific gravity falls, and the urea and chlorides return to the normal figure, but albumen is still present, the smoky color is apt to persist, and the precipitate which forms on standing contains renal epithelium, blood-globules, and granular or epithelial casts.

In favorable cases, the smokiness and albumen now gradually disappear, the urine often continuing for a little while to be secreted in excessive quantity; but in other cases, and unfortunately they are but too frequent, the albumen persists, and the urine assumes the characters indicative of chronic Bright's disease.

The form which the dropsy takes varies greatly in different cases, and seems to depend on inappreciable causes. Of the 29 cases that we have met with, in which its distribution was noted, anasarca alone was present in 22. In 1, there was extensive anasarca, hydrothorax of the right side, hydropericardium, and ascites. In 5, grave cerebral symptoms, probably uræmic in character, were present; and in 4 of these anasarca also existed. In 1 there were also hydrocephaloid symptoms, but of much less violent form.

Recent researches have established the fact that most of the cases formerly regarded as acute hydrocephalus are in reality due to the poisoned state of the blood, the so-called uræmia, so familiar to all that it is merely necessary to allude to it in this place.

The degree of danger to be apprehended from this dropsical complication depends upon the form which it assumes. M. Cazenave (*loc. cit.*, p. 52) says that there is no danger from it so long as it remains confined to the subcutaneous cellular tissue; and this is probably true. When, however, it attacks the serous cavities, or becomes associated with cerebral symptoms, due to the retention of urea and other excrementitious matters in the system, it is exceedingly dangerous. Of the 29 cases that we have had under charge of which we have preserved notes, 6 were fatal. Of the 22 cases in which the effusion was anasarca alone, but 1 was fatal. All of the 5 in which well-marked cerebral symptoms, due to uræmia, occurred in connection with anasarca, ended fatally. In one other case, which ended favorably, there were mild uræmic symptoms present. In the

case above adverted to, in which hydrothorax, hydropericardium, and ascites were added to the anasarca, the patient recovered after a long and severe illness.

In addition to the cases of dropsy and uræmia just referred to, and which all occurred in our own practice, we have seen quite numerous examples of scarlatinous dropsy with uræmic symptoms in consultation or in hospital practice. In one instance, uræmic symptoms came on very suddenly in a young child, and proved fatal in thirty-six hours, while in other cases, they have run a more gradual course, either ending fatally or terminating favorably after a severe illness of from several days to several weeks. In one case where recovery followed, the patient, a girl between three and four years old, was in a semi-comatose state for a week, with fever, excessive irritability of the stomach, and complaints of headache. For a period of five days the urine was entirely suppressed, not a drop having been voided during all that time, at least with the knowledge of the nurse, who was a most accurate and competent person. It would seem to be much more dangerous in the Parisian hospitals than in private practice in this country, since MM. Guersant and Blache speak of having seen it prove fatal in twelve, fourteen, and thirty-six hours, after one or two weeks, or even two or three months; and MM. Rilliet and Barthez refer to it as often proving fatal.

Diarrhœa is not an uncommon accident in the disease. It generally depends on simple functional derangement of the bowels. In some cases, however, it is so severe or long-continued as to constitute a serious complication. Under these circumstances, it depends on follicular enterocolitis, or slight erythematous inflammation of the intestinal mucous membrane.

In some cases, chronic angina remains after the subsidence of the disease; so, too, coryza may persist, even taking the form of ozæna.

Otorrhœa is not an infrequent sequel, and when following angina, and due to the extension of inflammation up the Eustachian tube, may be associated with permanent deafness, necrosis of the temporal bone, facial palsy, and even abscess of the brain.

Occasionally during the desquamative period, a painful swelling of the joints appears, attended with a renewal of the fever and, frequently, with sweating. This form of rheumatism is in all probability of a pyæmic character, and connected with the imperfect elimination of excrementitious substances, owing to the state of the various emunctories. In rather rare cases, the inflammation of the joint runs on to fatal suppuration. We have seen two fatal cases of this form of rheumatism. One occurred in our own practice in a girl five years old, and of good constitution seemingly. The type of scarlet fever was severe, but not malignant. The outbreak of rheumatism took place in the third week, after convalescence had seemed to have begun. It was impossible to trace the exciting cause of the rheumatic attack. There had been no imprudence that we could discover. In the second case, the patient, a boy three years old, previously healthy, had a severe scarlet fever with very high temperature, but without dangerous nervous phenomena. In the middle of the second

week, rheumatism of most of the joints, with acute pain, tense swelling and high heat, appeared. We saw the case in consultation. The child suffered intensely and died in three days.

Bronchitis and pneumonia are rare. *Inflammation of the serous membranes* is more common, occasioning in some cases the dropsical effusions which have already been treated of. It is in most cases connected either with renal disease or with the form of rheumatism above described. The pleura is more frequently affected than any other of the serous membranes; and not rarely the effusion becomes purulent.

Inflammation of the investing or lining membrane of the heart also occasionally occurs. Thus, of 39 cases of endo- or pericarditis mentioned by Dr. West, 6 could be traced to an attack of scarlatina.

Peritonitis is much more rare, and the effusion here also is especially apt to be purulent.

Scarlatina may be coincident with variola or measles. We have never seen it in connection with the former, but in two cases which came under our observation it was complicated with measles.

Diphtheria has also been observed not very rarely, usually appearing during convalescence. In a considerable number of cases, scarlatina has been noticed in the course of typhoid fever.

In some rare cases, as in the one detailed under the head of prognosis, more or less complete paralysis ensues during the convalescence from scarlatina.

Tuberculosis is not nearly so apt to be developed after scarlatina as after either rubeola or typhoid fever.

ANATOMICAL LESIONS.—The eruption sometimes disappears entirely after death, and on other occasions assumes a deep livid or purple appearance. The epidermis is generally loosened upon the integument, so as to be peeled off with great facility. The most important lesions, and those which seem to belong to the nature of the disease independent of complications, are the altered condition of the blood, and congestions of different parts of the body, particularly the brain, serous membranes, kidneys, spleen, glands of Peyer, and intestinal follicles. We have already alluded to the fact that, even when the cerebral symptoms have been most severe, and we might expect to find evidences of violent inflammation of the brain, nothing is observed after death, in the majority of cases, but congestion of the large veins and sinuses of the brain, of the pia mater, or of the cerebral substance. There is rarely any unnatural amount of serous effusion into the ventricles, or meshes of the pia mater; and it is evident that the symptoms have been due entirely to the vitiated condition of the blood. Nevertheless, effusions within the cranium may exist, in some few cases, as has been already stated in the remarks upon hydrocephalus.

The respiratory organs are usually healthy, with the exception of congestion and serous engorgement.

According to the researches of Fenwick, Fox, and Murchison, it appears that the entire gastro-intestinal mucous membrane is affected in many cases of this disease. There is congestion of the subepithelial layers, with excessive formation and subsequent desquamation of the epithelium.

The gastric tubules are greatly distended and obstructed by cells mixed with granular and fatty matters, and casts of their calibres are frequently found in the matters vomited or in the contents of the stomach after death.

The condition of the skin resembles this closely, the rete mucosum being thickened, with a formation of numerous round nucleated cells, and the sudoriferous glands being often obstructed by the rapidly formed cells.

The glands of Brunner and Pyfer are not unfrequently enlarged, and they are sometimes reddened or softened. In a smaller number of cases the mesenteric glands are slightly inflamed and increased in size, and the spleen is redder than usual or softened. These lesions have no necessary relation to the form of the disease, since they are often absent in typhoid cases, and present in those of a different type.

According to the observations of Dr. Klein, in twenty-three cases of scarlet fever dying from the second to the forty-first day (*Med. Times and Gaz.*, May 5th, 1877, p. 487), the kidneys, in the first week of the disease, showed an increase in the number of nuclei in the Malpighian bodies; hyaline degeneration of the intima, and multiplication of the nuclei in the muscular coat of the minute arteries; swelling and increase of the nuclei of the epithelium, and a granular appearance of the tubules and Malpighian bodies. After the first week, the changes noticeable were infiltration around the tubules, and tubal nephritis,—the tubules being crowded with hyaline cylinders and the epithelium presenting fatty degeneration.

The heart occasionally presents the results of inflammation of its lining or investing membrane; and in some cases its cavities contain firm white ante-mortem clots.

The blood exhibits very different appearances in different cases. It is viscid or serous, dark-colored or light, and fluid or coagulated, the clots being of variable color and density. The proportion of its constituent elements is changed. The fibrin maintains its usual relation to the mass of the fluid (3 parts in 1000), or it is very slightly augmented, while the quantity of the globules is increased to 136 or 146 parts, according to M. Andral, instead of 127, in 1000 parts. This increase in the proportion of fibrin may be in part the cause of the fibrinous depositions which occasionally are found in the cavities of the heart, and appear to have been instrumental in causing death.

In an article on "The Pathology of Scarlatina, and the Relation between Enteric and Scarlet Fevers" (*Med.-Chirurg. Trans.*, vol. lv, p. 103), Dr. John Harley, of London, reports thirty-six cases of scarlet fever, to show that the anatomical lesions of that disease are the same as those of typhoid fever in its early period, and that not unfrequently scarlatina, when long continued, passes into enteric fever. After describing these lesions, he says (p. 125): "From this view *our general conclusion as to the connection of scarlet fever and enteric fever is inevitable, viz., that the pathological changes accompanying an attack of scarlatina, include all those of the first stage of enteric fever, and are so far identical with them.* And it follows, therefore, that the transition from the former disease to the latter is nothing more than a natural pathological sequence, readily determined by

any cause which may increase the intestinal irritation." The italics are Dr. Harley's.

We have, on a few rare occasions, known cases of scarlet fever in our private practice, where the disease has been prolonged beyond its usual period, to assume some of the phenomena of typhoid fever, but this occurrence has been so infrequent that we doubt whether it ought to be regarded as the development of a pathological law connecting the two affections. That typhoid fever may attack a child just recovering from scarlet fever is as probable as that measles and scarlet fever may directly follow each other, or even coëxist at the same moment. Of both of these accidental coincidences, we have seen a few well-marked examples. Several of the cases described by Dr. Harley, in which typhoid fever certainly followed scarlet fever, occurred in patients admitted to hospitals. Five occurred in the London Fever Hospital, and in some of these the attacks of enteric fever began after full convalescence from scarlet fever; in one on the 28th day of convalescence; in a second on the 31st day; in another on the 37th day; in another on the 32d day; and again on the 56th and 32d days. We would ask whether in such cases the sequent typhoid fever ought not to be explained as the result of fever poison imbibed during residence in the wards of a fever hospital?

One very interesting fact observed by Dr. Harley, is the frequency with which he found fibrinous clots in the heart and great vessels "during a pyrexial state, at any period of the disease. This," he states, "is the commonest cause of death during the early stage of scarlatina; it is indicated during life by the reduction, often sudden, of a full pulse of about 120, to a dribble of 150 or 160 almost imperceptible impulses. The failure of the heart's action is commonly attended with orthopnoea and delirium, from obstruction to the pulmonary and cerebral circulations. On opening the body before it has lost a degree of temperature, and while the hot blood is still fluid, the right heart will be found distended, partly with dark fluid blood which coagulates on exposure; and partly, sometimes chiefly, by a large, firm, white, bifid clot continuous through the auriculo-ventricular opening. Each portion is interlaced with and firmly adherent to the tendinous cords and outstanding muscular bands of the cavity in which it lies, and sends outwards a rope-like continuation, the one into the pulmonary artery, and the other into the superior cava. These processes not only occupy a large portion of the area of these tubes, but extend with their branches upwards into the cranial cavity, and outwards into the lungs, whence they may often be withdrawn in ramifications up to the eighth degree, and eight or nine inches long.

"The left heart was generally empty and firmly contracted; in one case (1) each cavity was occupied by a large fibrinous clot, that in the ventricular spreading into the brachio-cephalic vessels of the arch of the aorta, and that in the auricle sending large ramifying branches into the pulmonary veins. In another case (12) the auricle was distended with dark softly-clotted blood."

We desire to call attention to these facts, since we doubt not that they explain many of the cases of early death in this disease, in which all

medical treatment has proved so futile, and also on account of the great interest of these observations in connection with similar results which will be mentioned in the article on diphtheria.

DIAGNOSIS.—It is impossible to distinguish scarlatina from the other eruptive fevers by the symptoms which precede the eruption. The only signs upon which a diagnosis at that time might be grounded, are great frequency of pulse, which is characteristic of this disease, some soreness or redness of the fauces, and the prevalence of the disease in the community. But these are all exceedingly fallacious, and the physician should be content to wait for the eruption before he ventures to speak with certainty. After the eruption has come out, it can scarcely be mistaken for anything else, except it be roseola.

From *measles* it may be distinguished by the differences in the prodromes, course, and eruption of the two affections. The prodromic stage of scarlatina rarely lasts more than twenty-four hours, and is very often much less; that of measles, on the contrary, is almost always from two to three days; in scarlatina the rash appears suddenly and often spreads over the whole body in a single day; in measles it appears on the face first, and extends gradually to the rest of the surface, seldom reaching the hands and feet before the end of the second day; the eruption of measles occurs first in distinct papules, which coalesce and form patches of an irregular crescentic shape, while that of scarlatina is in the form of innumerable minute dots or punctations, placed so closely together as to give to large portions of the surface a uniform color, like that produced by blushing. The color of the two eruptions is different, that of measles being dark like raspberry juice, and that of scarlatina of a more or less bright scarlet tint. The presence of catarrhal symptoms in measles, and their absence in scarlet fever; the absence of angina in the former disease, or its very slight character, and the severity of the throat affection in scarlatina; and lastly, the greater severity of the febrile symptoms, particularly the frequency of the pulse and the heat of skin in scarlatina, are other points of difference which will assist in making the diagnosis, rarely, it seems to us, difficult, still more certain. A very great frequency of the pulse is one of the most unfailing symptoms of the early stage of scarlet fever. It almost always runs up to 140, 150, or 160, in young children, within the first twelve or twenty-four hours, and to 120, 130, 140, or higher, in those who are older. Nevertheless, this, like all other symptoms, is sometimes wanting. We have lately seen a boy, between five and six years old, with a marked but very safe attack of the disease, whose pulse ranged between 80 and 90 throughout the sickness. This was, however, the only case we have ever met with, in which the pulse remained so little disturbed.

It is sometimes very difficult to determine with precision between erythema and scarlet fever. By the eruption alone, we believe it to be often impossible. We have seen quite a number of cases, in which the eruption of erythema resembled so closely that of scarlet fever, that we should have been obliged to confess our inability to make the distinction, had it not been for the other symptoms, and particularly the frequency of the circu-

lation, the heat of the skin, and the throat symptoms. The most important differential symptoms are the tint of the eruption, which in erythema is dark-red, in scarlet fever bright-red or scarlet; the characters of the patches of eruption, which are more regular in shape, but of much smaller size in erythema than in scarlet fever; the total absence or very slight degree of anginose inflammation in erythema; and, what is decidedly the most important of all, the very much slighter degree of febrile reaction in erythema, in which the pulse, instead of being doubled in frequency as it is in scarlet fever, is scarcely above its natural rate, and in which the heat of skin is but little above the standard of health. Moreover, erythema is generally of shorter duration, and is a milder affection, and therefore accompanied by far less fever and general disturbance of the constitution.

Diphtheria occasionally resembles scarlatina to so great an extent, as to have even led some observers to consider them identical. Thus, there is in diphtheria a pseudo-membranous angina, with swelling of the cervical glands, and at times albuminuria, and even an erythematous rash. We will elsewhere (see article on diphtheria) give at length the differential diagnosis between these affections, and will here merely call attention to the fact that the rash is a rare exception in diphtheria, and is a mere uniform erythematous redness; that even when albuminuria is present, the urine does not present the other characters noted in scarlatina; and that the condition of the fauces in the two diseases is somewhat different. There is, further, a wide difference in the sequelæ of the diseases; and, finally, they do not exercise any protective power whatever against each other.

There is a form of disease known as rubeola notha, epidemic roseola, rosalia (Richardson), rötheln, in which there are some of the symptoms of both measles and scarlatina; the eruption appearing on the second or third day, at first resembling that of measles, but becoming soon more like that of scarlatina. Coryza and angina may both be present, and there is subsequent desquamation. Some authorities regard this as a union of the poisons of measles and scarlatina, while others consider it a separate disease, because epidemics of it occur when neither measles nor scarlatina are prevailing. Previous attacks of these latter do not protect against it. In an extensive epidemic in the lower part of this city, which appeared to be of this nature, not a single case, of the numbers which came under our observation, was followed by any of the sequelæ of either measles or scarlatina. We believe this to be a specific disease. A very prevalent and wide-spread epidemic, the first thoroughly marked one we have met with, occurred in Philadelphia in the winter and spring of 1880-1881. It attacked several children in a family, and sometimes all. The eruption resembled measles much more than scarlet fever. There was some moderate fever and lassitude for one or two days, distinct but inconsiderable angina, and slight catarrh of the eyes and nose. One of the most characteristic conditions of the disorder was slight swelling of some of the cervical lymphatic glands. This was generally, but not always present. The swelling was very moderate, seldom great enough to attract

the eye, but to be found by careful touching. The glands were not larger than cherry-stones usually, not at all numerous, some three or four or more, and situated in the region of the sides of the nucha, behind the ear, or at the sides of the neck. They were often so small as to be discovered only by careful manipulation, and were not painful to the touch. In no case did the symptoms look for a moment like a full or severe case of scarlet fever, and in none did we see any dangerous conditions during the attack or subsequently.

PROGNOSIS.—It is impossible to obtain a useful average mortality of scarlet fever, since the disease varies so greatly in different epidemics, and under different hygienic conditions, that the results obtained during one period are inapplicable to cases observed at another. This is proven by the experience of almost every physician, and by the evidence of many writers. It is proven, also, by the following facts: M. Guérin (*loc. cit.*, p. 283) states that the mortality in the epidemic observed by him was about 1 in 12; of 99 cases, 8 died. MM. Rilliet and Barthez lost a little more than half their cases; of 87, the total, 46 were fatal. These cases, let it be remarked, however, occurred in the Hospital for Children in Paris, which will account for the heavy fatality. The degree, however, to which the mortality may vary in the same place and under the same plan of treatment, is shown by the fact, mentioned by Hillier, that in the course of eleven years the annual mortality from scarlet fever in the London Fever Hospital, varied from 2.5 per cent. to 16.5 per cent.; and in the Hospital for Sick Children in London, from 9 to 31 per cent. Of the 274 cases that we have observed, 31, or rather more than one-ninth, were fatal. Of the 274 cases, 104 occurred between 1849 and 1853, and in these the mortality was much smaller than in those which occurred prior to that year. Of 104, 11 were fatal, or about one in nine and a half. Seventy-eight cases occurred between 1853 and the spring of 1857. Of these 78, only 4, or 1 in 12, were fatal. Of 81 cases observed previous to 1849, 13, or about 1 in 6, proved fatal. Of 11 cases occurring in 1872-3, 3 proved fatal. The mortality met with by ourselves in private practice has greatly varied, therefore, in different series of years. In one series it was 1 in 6, in another 1 in $9\frac{1}{2}$, and in a third 1 in 12. Lastly, to show the influence of the epidemic type upon the mortality still more clearly, we may state that of the last series of cases observed, 78 in number, 43 occurred during the epidemic which lasted from the summer of 1856 to the spring of 1857, and of these only 3, or 1 in 14, died.

The prognosis must be based, therefore, in part on the character of the epidemic prevailing at the time. It must depend, also, on the nature of the case. Mild and regular cases are rarely fatal. Of 206 mild cases that have been under our care, only three proved fatal. One of these would probably not have so terminated had it not been for the imprudence of the nurse. This was, in fact, the case of a young child who had recovered from the eruptive stage of the disease, but whom the nurse carried out of the room in the second week, notwithstanding express directions to the contrary. The child took cold and was seized with catarrh and slight anasarca; on the fifteenth day uræmic symptoms set in, and it

died on the seventeenth day, comatose, and with convulsive movements of different parts of the body. The second case was that of the boy thirteen years old, already described, who died with sudden hydrocephaloid symptoms, at the end of the second week. The third fatal case occurred in a girl between eight and nine years old, who died suddenly at the end of six weeks. The patient had convalesced sufficiently to have been out several times, but remained very hydræmic and weak. After being much fatigued one afternoon by playing with some little friends, she was seized next day with vomiting, and soon after with great difficulty of breathing and extremely rapid and feeble action of the heart. These symptoms increased on the following day. The dyspnœa was most severe, and was attended with cyanotic color of the hands and face, and with cold colliquative sweats. The lungs were free, there was no cough, and auscultation revealed no pericardial lesion. Death occurred suddenly at the end of a day and a half. No post-mortem was made, owing to circumstances that could not be controlled. Our own opinion was, and is, that the death was caused by a coagulum in the heart.

Grave cases of scarlet fever are always, on the contrary, exceedingly dangerous: thus of 61 cases of this kind that we have had under charge, 28, or nearly a half, were fatal. In order to render the description of the symptoms of this class of cases more clear, we divided them into two groups: one, in which the onset of the disease is instantaneous and most violent, being characterized by excessive disturbance of the nervous system, taking the form usually of convulsions, but sometimes only of profound coma; and a second, in which the symptoms of the onset, though severe enough usually from the first to mark the character of the case as grave, are less violent than in the first group, and especially not marked by the occurrence of convulsive phenomena. Of 18 cases belonging to the first group, 13 died; while of 43 belonging to the second, 15 died. Violent nervous symptoms occurring early in scarlet fever augur, therefore, great danger to the patient, since of 18 cases in which they were present, 13 died, whilst of 43 in which they were more moderate, though still marked and severe, only 15 died.

The character of the nervous symptoms is, therefore, all-important in the determination of the prognosis, as the probable termination of the case is to be foretold more certainly by a just appreciation of these particular phenomena of the disease than by any other means. Excessive jactitation or irritability, delirium, coma, and the hydrocephalic cries, are all unfavorable symptoms, but not in the same degree as those connected with the locomotor apparatus. MM. Rilliet and Barthez state that they have seen recoveries take place in cases in which the intelligence of the patient had been very much disordered, while of those who "*during the first fifteen days of scarlatina*, were taken with convulsions, convulsive movements, contractions, in a word, any symptoms affecting the locomotor apparatus, all, without exception, died." This does not accord exactly with our own experience, through nearly enough so to show how exceedingly dangerous are the symptoms just enumerated when they occur early in the disease. General convulsions occurred on the first day of the disease in 9 of the 61

grave cases observed by ourselves, and of these not one terminated fortunately; in 4 they occurred on the second day, and of these 3 recovered and 1 died; in one they occurred on the ninth day, and this patient also recovered; in another case there were no general convulsions, but on the first day there were automatic motions, with involuntary extensor motions of the arms and fingers, and on the second day strabismus, with a continuation of the automatic motions. This case proved fatal. Of the 15 cases, therefore, in which marked disturbances of the muscular system occurred, only 4 ended favorably. Of 10 subjects in which the convulsive phenomena occurred on the first day of the disease, not one escaped. Of 5 subjects in whom these symptoms appeared on or after the second day, 4 escaped. One of the favorable cases occurred in a boy seven years old, who had a general convulsion, lasting several minutes, on the second day of the attack; this was followed by delirium and coma alternately, but no return of the convulsions. The case was a most violent one, and lasted six weeks, leaving the child at the termination very deaf, but otherwise in good health. The second case occurred in a child five months old. The convulsive symptoms appeared on the ninth day, and consisted of strabismus, spasmodic retraction of the head, and occasional slight spasms of the limbs. They alternated with coma, and disappeared on the tenth day, until the seventeenth and eighteenth, when the strabismus reappeared. The child recovered perfectly. The third was that of a very healthy and vigorous boy between eight and nine years old, who, on the second day of an attack which had begun like a severe cholera morbus, had, twice, fits of insensibility, with stiffening of the extensor muscles of the fingers, rigid contractions of the flexors of the arms, and spasms of the eyeballs. This case proved afterwards very violent, so that the patient nearly died on the fifth day, with asphyctic symptoms, caused by very great swelling of the tonsils and fauces, and enormous enlargement of the external cervical lymphatic glands, complicated moreover with extensive acute œdema about the chin and front of the neck. These symptoms were followed again by diphtheritic deposit covering the whole of the pharynx. He finally, however, recovered perfectly. The fourth case was that of a boy between five and six years old, who, on the second day, had an attack of general convulsions, which were repeated frequently on the third day. This patient continued very ill for several days, and when, at last, he began to improve somewhat in the middle of the second week, it was found that he had lost entirely the power of speech, and all control over nearly the whole of the locomotor apparatus of the body. He could neither lift his head nor turn it; the legs were immovable, the hands perfectly helpless. The only motion that remained was a jerking, apparently almost automatic, movement of the arms upon the shoulders, and the forearms upon the arms. But even these were most irregular, and badly co-ordinated. He was very much in the condition of a new-born child. It was difficult to ascertain what the condition of his senses was; but after a short time we were able to satisfy ourselves that he saw and heard, and only after many weeks was he able to hold a very light object in his fingers, then to move his head from side to side, and at a still later period to hold it up. At the end of

about two months he could sit in a chair, when placed in it, but could not sit on the floor unsupported. At the end of three months he was learning to walk by being held up by the arms. He had never spoken a word. The only approach to anything like articulation was the ability to hum a low gentle musical note; his intellectual faculties, so far as we could judge by the signs he made, were awakening. At the end of two months he could speak intelligibly some three or four words. When we last heard this patient he was a man of over twenty years of age, with thick speech, slow mind, irritable and unreasonable temper; in fact, of very low mental development. The fifth case was that of a male infant, nine months old, who, on the second day, had severe general convulsions, followed by very deep drowsiness. The eruption became intense, and, on the third day, the convulsive symptoms recurred from time to time, but with less violence. On the fourth day he seemed somewhat better, but on the fifth very severe anginose symptoms set in, and he died.

Again, in 20 of the 61 grave case, severe and more or less prolonged delirium or coma occurred, and of these 14 died. We may conclude, therefore, that convulsive symptoms appearing early in scarlet fever indicate a highly dangerous and, in all probability, a fatal attack; while severe, and especially prolonged delirium or coma, are also extremely unfavorable symptoms, but somewhat less so than are those of a convulsive character.

Other unfavorable symptoms are: extremely frequent or very violent pulse; intense heat or unnatural coolness of the skin; persistently elevated temperature after deflorescence; deficiency or sudden disappearance of the eruption; a livid or purple tint of the eruption; slow and imperfect capillary circulation, as ascertained by pressure; the appearance of petechiæ; ecchymoses, or hemorrhages; violent vomiting and colliquative diarrhœa; great violence of the throat affection, as shown by tumefaction, abundant pseudo-membranous exudation, or disposition to ulceration and sloughing; and lastly, severe coryza or otorrhœa. A disposition to a typhoid state, indicated by dulness of the intelligence, dusky hue of the skin, frequent and feeble pulse, dry, brown tongue, sordes on the teeth, meteorism, and disposition to diarrhœa, is always dangerous.

When, on the contrary, the fever is moderate, the cerebral symptoms absent or very slight, and the eruptions regular, and of a bright tint; when there is no disposition to typhoid symptoms; when the throat affection is mild, and the disease pursues a regular, uniform course, we have every reason to expect a favorable termination in a large majority of the cases.

In addition to these remarks it may be said that neither age, sex, nor social position influences the prognosis. A delicate constitution does not seem to predispose to a violent attack of scarlatina, and, indeed, many of the most malignant cases occur in robust children; but, on the other hand, it has been noticed that in certain families there exists a strong tendency for the disease to assume a grave and fatal form.

TREATMENT.—*Hygienic Treatment.*—In all cases of the disease, whether of the mild or grave kind, the strictest attention should be paid to the

hygienic condition of the patient. The room in which the child is placed ought to be, if possible, large, and at all events well ventilated. The temperature in winter should be carefully attended to. We usually direct it to be kept at from 68° to 70° F., during the early stages of the disease, unless the fever is violent and the child complains of heat, in which case it may be allowed to fall to 66° , or 62° . The clothing ought to be moderate, not enough to increase the heat of the skin, nor yet so little as to endanger chilliness. During the latter stages of the disease, when the fever has subsided, and particularly when the heat of the skin has fallen, the temperature of the chamber ought to be kept, as a general rule, at from 68° to 70° , and, when the child is pale, weak, and chilly, it may be maintained with great propriety at 72° .

One of the most important points in the treatment of scarlet fever is, undoubtedly, the management of the patient during the convalescence, and especially during the desquamative period. It is during this period that the child is liable, as we have already shown in our account of the different complications and sequelæ of the disease, to dropsy, which is the most frequent, and at the same time the most dangerous accident to which the patient is exposed. There can be no doubt, we think, from the opinions expressed by various writers, and also from our own experience, that the most common cause of this accident is exposure to cold. Chilling of the body, no matter how produced, is exceedingly apt, when it occurs within three, or, more rarely, four weeks from the invasion of scarlet fever, to be followed by a more or less marked attack of some form of dropsy. It is true, we are well aware, that dropsical effusions sometimes take place in subjects who have been guarded in the most careful possible manner, and in whom there has been no evident exposure to cold; but it is also true, that a much larger number of those who have been thus guarded escape than of those who are not thus taken care of. We have, therefore, no doubt whatever, that it is most wise and prudent to confine the patient to well-warmed rooms, or at least to the house, for twenty-one or twenty-eight days from the outset of the disease. The fact that the attack has been a slight one only makes it the more necessary to carry out this regulation, as it has been found by experience that dropsy occurs more frequently after mild than after severe attacks. M. Legendre (*Recherches Anat. Pathol.*, p. 311) is of opinion that the patient ought not to be allowed to leave the house until the skin, completely deprived of the old epidermis, shall have regained its suppleness, its smooth and polished appearance, and all its functions. When, therefore, after a mild case, the desquamation is completely terminated in three weeks, the patient, he thinks, may be allowed to go out. But, on the contrary, this period would be too short by one-half, if the eruption had been very intense, as the desquamation is, in such cases, scarcely finished on the hands and feet at that time. Our own opinion, as already stated, is, that in the cool seasons of the year, the patient ought to be restricted to the house during full four weeks.

TREATMENT OF MILD CASES.—Mild cases, those in which the eruption

is moderate, the temperature but little above the normal point, even though the pulse be very frequent, in which neither delirium, stupor, nor unnatural jactitation betray threatening conditions of the nerve-centres, need but the simplest treatment. The child must be confined to a comfortable, well-ventilated room, and cooling drinks, as cold water, lemonade, or orangeade, should be allowed, and indeed they ought to be recommended, and the nurse should be made to understand that she is not to wait until a young child calls out for a drink, but that she is to offer it frequently. Young children, or at least some, seem not to know when they are thirsty, or hot, or cold; they have not yet learned to express their sensations in words, and a wise nurse or physician will think for them.

In all cases in which there is heat of skin and frequent pulse, and these conditions attend all but a very small fraction of the whole number, the patient ought to be kept in bed whilst the fever lasts, and for two days afterwards. This point, so important in all fevers, is too often neglected. Cases so treated are apt to be shorter in their duration, milder in their symptoms, and less likely to be followed by any of the troublesome sequelæ so prone to occur, as inflammations of the cervical glands, of the ear, or of the kidneys.

In many mild cases no drug whatever is needed. If the bowels are positively costive, that is to say, if they are not moved every second or third day spontaneously, a simple enema, a dose of syrup of rhubarb, a baked apple, or stewed prunes, will suffice. If the temperature is high, the pulse active, and the patient restless and suffering, sweet spirits of nitre, solution of the acetate of ammonia, or two or three grains of the citrate of potash, with from a half to one drop of deodorized tincture of opium, every two or three hours, according to the age, will usually lessen the heat and promote quiet. We wish to repeat, however, our opinion that in a great many cases of this type, no drugs whatever are necessary. The time is fast coming when even the vulgar and illiterate will no longer quarrel with the physician because he gives no drugs, not even in infinitesimal doses, and the time has come when the wise and educated trust the intelligent physician, so that he need no longer give placebos in order to earn his fee.

The diet should be, for the first five or six days, in great measure, liquid. Milk, with or without some farinaceous substance, to suit the tastes or habits of the patient, or with bread and butter, and beef or chicken soup, with rice or bread, are sufficient. After five or six days, when no severer symptoms have made their appearance, and the disease is on the decline, light meats, eggs, stewed fruits, or potatoes, may be added.

Baths, tepid or warm, spongings with tepid or warm water, cloths wetted with cold or tepid water, applied to the forehead, may be used, according to the judgment of the physician. They are not necessary agents, but in certain cases, when the heat of the skin tends upwards, when the patient is restless from nervous irritation, and particularly if the child is in the habit of being bathed, they may be used with much advantage and comfort.

The throat, in mild cases, rarely needs any treatment. If, however, the patient complain of pain, if there be some uneasiness in swallowing, or if decided patches of exudation make their appearance on the tonsils or pharynx, it will be well to let the child gargle, if it be old enough, with solution of chlorate of potash or alum, or with flaxseed tea. An excellent gargle is one made of a wineglassful of table claret, two wineglassfuls of water, and forty grains of chlorate of potash. If the patient is too young to gargle, some chlorate of potash or alum can be mixed with powdered sugar, and a small pinch placed upon the tongue every two or three hours. One part of the chlorate or alum may be rubbed up with five or six parts of the sugar. There is no necessity for the application of strong agents of any kind to the throat. Even though patches of exudation of considerable size appear upon the fauces, they will disappear spontaneously in all cases of the kind we are discussing. We object to the forcible application of medicinal solutions to the throat in young children, unless they are absolutely necessary to clear the passages of obstructing viscid and offensive secretions. In children of a certain type,—those of sensitive nerves and strong wills, in whom fear of pain on the one hand, and will to resist on the other, form a combination which prompts the child to resist such an operation to the last,—even though we might hope some benefit from the application, the irritation and exhaustion caused by the struggle, and the agitation kept up by its expected renewal, will do more harm, we think, than the treatment can compensate for.

Inunction, as one of the means of treatment in scarlet fever, is now so well known that we suppose nearly all physicians use it. For our part, we order the ointment for external use just as regularly nowadays as we do cold drinks and proper food.

It was first proposed and strongly urged, we believe, upon the profession, by Dr. Schneeman, a German physician. Dr. Schneeman makes use of bacon fat. He takes a piece about as large as the hand, still covered with its rind, in order to obtain a firm grasp upon it. On the soft side of the piece slits are made in various directions in order to allow the oozing out of the fat. The patient is to be rubbed with this, as soon as we are aware of the nature of the case, from head to foot, excepting the face and scalp, every morning and evening. The rubbing is to be so performed that the skin may be regularly, but not too quickly, saturated with the fat. During the process only the part being rubbed is to be uncovered, or the whole can be done under the bedclothes. (*Ranking's Abst.*, No. 12, p. 26.)

For our own part, we used the bacon fat but twice, soon finding how disagreeable an application it was, and not believing that the salt it contained could do any good whatever. We now always employ an ointment made by rubbing together a drachm of pure glycerin with an ounce of cold cream (*ungt. aq. rosæ*). We have seen children smeared from head to foot with lard, and, what is worse, with goose-grease, with their clothing saturated, their pillows and sheets a mass of discolored grease, most offensive to the eye and nostrils. This is quite unnecessary. Our own method is to explain to the mother or nurse, that she must take a little of the oint-

ment above recommended in the palm of the hand, and with this rub gently the various parts of the surface, first one limb, then another, and then the body. The ointment should be rubbed in with gentle pressure, and it is well, we think, to knead and squeeze lightly the various portions of the body being anointed, as is done in the *massage* of the French, or by the "rubber" of the English. These manipulations assist, we think, the capillary circulation, which is often a good deal impeded. After applying a moderate amount of the ointment, until the skin is well softened and oiled, any excess of the material should be wiped off with a soft towel or handkerchief. In this way the anointing is thoroughly accomplished, and yet the clothing and bed linen are not so soaked and saturated with the oleaginous substance, as to be disagreeable to the patient, nurse, or mother. Cosmoline can be used in place of the ointment above mentioned, and is equally as efficacious and valuable.

There can be no doubt, at the present time, that the employment of inunction in scarlet fever has proven a most useful addition to our former means of treatment. In our hands it has had the effect of allaying, in all cases, the violent irritation caused by the intense heat and inflammation of the skin. In nearly all cases, it sensibly diminishes the frequency of the pulse, and in many this effect is very strongly marked. It removes, of course, the dryness and harshness of the skin, keeping it, instead, soft and moist. It lessens or even removes the burning, irritation, and itching caused by the eruption. By these effects, to wit, lowering of the pulse, and alleviation of the external heat, dryness, itching, and irritation, it cannot but, and evidently does modify and diminish, most happily, the injurious effects of the disease upon the constitution at large. So great is the comfort it gives to the patient that we have several times had young children, still untaught to speak, to make signs and motions, at shorter or longer intervals, showing their desire to have the application renewed. The frequency of the application must depend upon the case. When the eruption is intense, the skin very hot, and the febrile symptoms marked, they should be made every two or four hours, or even oftener. In milder cases they need to be repeated only three or four times in the twenty-four hours.

TREATMENT OF GRAVE CASES.—The most dangerous cases of this disease are those of the type described at page 786, in which the attack is sudden, and in which disorders of the nervous system in the form of convulsions, tremors or rigidity, retraction of the head, delirium, stupor, or coma, appear within a few hours of the onset. When this type of the disease attacks very young children, they, so far as we have seen, nearly always die in sixteen, twenty-four, or forty-eight hours. Older children have more chance of escape, but, even in them, the danger is extreme.

We have seen everything tried in these cases, from depletion by bleeding and leeching, many years since, to expectancy, and must confess that we have little faith in the power of human agency to contend against this particular array of symptoms. Depletion is no longer, we believe, thought of by any, and there is often no time for the action of drugs.

It is in such cases that the use of water at different temperatures, applied in the form of baths, affusions, packings, ablutions, and ice, has been recommended, and has seemed in some cases to do good. We shall give a rapid sketch of the opinions of those who have used this means, and then state our own views.

Dr. J. Currie, of Edinburgh, was the one who first and most prominently brought before the profession the use of cold water. It must be observed, however, that Dr. Currie limits its uses to cases to which he applies the term *anginose*, many of which, we doubt not from his description, ought to be classed as mild cases. He mentions another class of cases which he thinks ought rather to be called "*purpurata*," characterized by "extreme feebleness and rapidity of the pulse, and great fetor of the breath. . . . The heat does not rise much above the standard of health. Great debility, oppression, headache, pain in the back, vomiting, and sometimes purging, accompany its rapid progress; the patient sinks into the low delirium, and expires on the second, third, or fourth day. . . . The cold affusion is scarcely applicable to it, and the tepid affusion makes little impression upon it. In my experience, indeed, all remedies have been equally unsuccessful. It outstrips in rapidity, and it equals in fatality, the purple confluent small-pox, to which it may be compared." (*Currie's Med. Reports*, Philada. ed., p. 277.) It is clear, therefore, that Dr. Currie, when he speaks of nearly invariable success in upward of one hundred and fifty cases (p. 286), had to do, not with the malignant, or, at least, not with the most malignant forms, for which we are seeking a remedy, but with cases of a mild form, or at most with those of the severe *anginose* type. Indeed, at page 294, we find the following remarks: "It has come to my knowledge, that in two cases of *scarlatina*, of the most malignant nature, the patients have been taken out of bed; under the low delirium, with the skin cool and moist, and the pulse scarcely perceptible. In this state, supported by the attendants, several gallons of perfectly cold water were madly poured over them, on the supposed authority of this work! I need scarcely add that the effects were almost immediately fatal." We have been induced to enter thus much into detail in regard to the use of cold affusions, because of the intrinsic importance of the subject, and because of the remarks upon it in the work of MM. *Rilliet and Barthez*, who bring forward Currie's success as a strong argument in favor of their employment in that form of the disease in which cerebral symptoms predominate. Currie does not recommend them, however, except in cases in which the reaction is full and strong, as indicated by very great heat of skin, scarlet eruption, and rapid, but not feeble pulse. In the famous cases of his own two children, it is evident that the attacks were not malignant, for the skin was very hot (108° and 109° F.), and no mention is made either of stupor or delirium, much less of convulsive phenomena. Dr. George Gregory, of London, whose opinions upon all matters connected with the eruptive fevers are of course worthy of great weight, says (*Lect. on the Eruptive Fevers*, edited by Dr. Bulkley, New York, p. 190), in relation to the use of cold affusion: "Sanctioned

by my uncle, the late Dr. Gregory, of Edinburgh, this plan has been amply tried in all parts of the world, but it has not realized the expectations of its proposer.

"The truth is that the cold affusion is applicable only to a small number of cases. It is adapted for young people with high anginose inflammation and a burning hot skin, without plethora, without depression of nervous energy; but it is inapplicable to the scarlatina of adults, accompanied with coma, phrenitis, or marked debility. It is wholly unfit for cases of cynanche maligna. It answers its purpose very well for the first day or two, but it is often impossible to continue its use. Lastly, it seems to increase the disposition to dropsy."

Dr. Currie's method of using water was by affusion. The child is undressed and placed, erect or sitting, in a tub, while four or five gallons of water, at from 60° to 70° F., are poured over the head and body. The good effects of the remedy are said to be an immediate reduction of the heat, a diminution in the rapidity of the pulse, which, in one of Dr. Gregory's children, fell in half an hour after the cold affusion from 160 to 120, a disposition to sleep and quiet, and, according to Dr. Gregory, a seeming arrest of the throat affection. These good effects of the affusions are transient, however, as the heat of skin and rapidity of the circulation return in the course of one or two hours. For this reason it is necessary to repeat them frequently, once in two or three hours at least, in order to render the effects permanent. Currie used fourteen affusions for one of his own children, and twelve for another, in thirty-two hours. These were not, however, all cold. Gregory used for his child five "good sousings," to use his own words, in twenty-four hours.

MM. Rilliet and Barthez give, in the following words, the conclusions of Henke in regard to the use of cold affusions: 1. Cold affusions are not adapted for a *general* method of treatment. 2. The slight, or simply inflammatory forms, do not all demand so energetic a treatment. 3. Their employment must be reserved for cases in which the disease is epidemic, and accompanied by intense heat and dryness of the skin, with smallness and acceleration of the pulse, and for those in which the cerebral symptoms are very violent and characterized by great restlessness, alternating with drowsiness, commencing from an early period of the disease. Scarlet fever under these circumstances is so dangerous, they say, and so often mortal, that recourse ought to be had to all curative means, and in children the cold affusions are much more strongly indicated than bleeding (*op. cit.*, vol. ii, p. 653).

Dr. Hiram Corson, of Montgomery County, in this State, has, so far as we know, used cold externally more boldly than any one in this country. He began this treatment in 1844, and, in a report made by him to the Pennsylvania State Medical Society, "On the External Application of Ice to the Throat as a Remedy in Scarlet Fever and Diphtheria" (see *Transact. of the Med. Soc. of the State of Pennsylvania for the year 1864*), declares his unabated faith in the excellence and safety of the treatment. He advises, in cases attended with convulsions, the pouring of cold water from a height

of a few feet on the head for several minutes at a time,—this to be repeated every fifteen or twenty minutes until relief is obtained. At page 467, he says: “Hundreds of times have I had patients brought to the side of the bed and cold water poured freely over the head, until the stupid, almost comatose child, was yelling, and kicking, and striking to get rid of the falling water; and this I have repeated whenever the symptoms called for its repetition.” He prefers in these cases the cold affusion to ice. He also applied pieces of ice wrapped in cloths to the neck, when the anginose symptoms were severe, and, when the temperature was very high, washed the whole body with iced water, until the heat was reduced.

Dr. Corson, in this article, speaks with the greatest possible confidence of his treatment, and when others evince some dubitation as to the invariable success of the cold treatment, avers that they had used it imperfectly or with timidity. It is most unfortunate that his paper deals altogether in general assertions. At page 458, he says: “And now, after twenty years of experience in the use of it, and after treating scores and scores of cases, I am most happy to say that I have never seen the least injury produced by it, but, on the contrary, regard it as the means, above all others, of comfort and safety to the patient.” He does not refer to a single fatal case during the twenty years he has been using this system. At page 453, however, he speaks of having “during the whole winter, in about one hundred cases, continued the treatment in degrees apportioned to their mildness or severity, and without the loss of a single patient thus treated.”

Nevertheless his experience is valuable, for it shows that, in some cases at least, the use of means which reduce rapidly the heightened temperature of the body in scarlet fever, acts as favorably as it has been found to do in the hyperpyrexia of sunstroke, rheumatic fever, and in continued fevers.

Hillier (*Dis. of Children*, p. 326) states that he has employed cold affusions with good effects in a few malignant cases. He used water from 70° to 75° F., wrapping the child immediately after the affusion in dry blankets. He adds that “in cases of collapse with cold extremities, it would not be prudent to resort to the operation.”

Trousseau (*Clin. Med.*, Syd. Soc. ed., vol. ii) recommends cool or cold affusions when dangerous ataxic nervous symptoms make their appearance. At page 198, he says: “I declare to you that I have never resorted to the employment of cold affusions without obtaining beneficial results. I am far from pretending that all my patients recovered; like my colleagues, I have lost the greater number, but even those who died experienced a temporary relief from suffering, and the affusion, so far from proving injurious to them, always moderated the symptoms, and also seemed always to retard the fatal termination.” At page 206, he states that he does not use them indiscriminately in all cases, but only “to subdue serious nervous complications—formidable ataxic symptoms.”

Dr. Gee (Reynolds's *Syst. of Med.*, vol. i) speaks of the cold affusion as being sometimes useful in the malignant form of the disease, attended with delirium, diarrhoea, vomiting, full pulse, and great heat of skin. He adds, however (p. 355), that in the “primary adynamic form, all treat-

ment will be baffled. The cold affusion is the only means which has seemed to me to be of even momentary benefit."

We shall now refer to our own experience in the employment of external cold. We never use it to its full extent except in really dangerous cases. So long as the case is mild or moderated, or even severe, if there be no cerebral, and especially no locomotor disturbances, we deem it unnecessary, and rest content with more simple means; or we use simply ablutions with tepid or cool water, with cold applications to the head, so long as they are agreeable, and until the temperature is reduced. But, when the temperature rises very high (105°), or, as Currie asserts in one case, to 112° , and Dr. Woodman (*Wunderlich on Medical Thermometry*, Syd. Soc. ed., p. 204, footnote) to 115° F., with nervous symptoms, the danger is extreme, and we have used, and shall use hereafter, means to reduce the heat. In one case we made repeated affusions upon the head with water at 70° , pouring at one time seven bucketfuls upon the part. This was a case attended with coma, strabismus, and spasmodic retraction of the head. In addition to the affusions, cloths dipped into iced water were kept applied the greater part of the time. These means, especially the affusions, were evidently advantageous, and the child recovered.

We have made use of lotions with cool water (70° to 72°) in three grave cases. In two they were evidently useful; in one they did no good, and were perhaps injurious, as we believe now that the case might have been better treated with prolonged warm baths at a temperature of 92° to 95° , cold to the head, and internal stimulation.

The latter case was one in which the patient had two convulsions on the first day, and one on the second. The pulse rose at once to between 160 and 170; the head and trunk were very hot, whilst the extremities were cool; the child was either excessively dull or comatose. Cloths wet with iced water were kept constantly upon the head and the body, and occasionally the limbs were sponged with cold water. The internal remedies consisted of carbonate of ammonia and milk punch. The patient improved decidedly on the third day, so that the pulse came down to 152, the intelligence returned, though the child was still very drowsy and heavy, and the case looked quite promising. On the fourth and fifth days, the throat affection came on; the neck and throat swelled enormously, the cervical lymphatic glands became very large, the nasal passages discharged streams of offensive grumous pus, the ears ran copiously, the fauces became pseudo-membranous, the deglutition grew worse and worse, until at last it was impracticable, and the child died on the middle of the sixth day, a mass of the most disgusting and offensive disease. One of the grave cases in which the cool applications proved useful, occurred in a hearty, vigorous girl, twelve years of age. On the third day of the attack, the symptoms were as follows: the pulse was between 160 and 170, small and quick; skin intensely hot; eruption very copious and of a dark-red color tending to violet; capillary circulation slow and languid; tongue black, and covered with a hard, dry crust; teeth and lips dry, and covered with dark incrustations. There was very great agitation and restlessness, with constant moaning and complaining, the total insomnia. Under these circumstances, we directed the nurse to sponge the head and extremities of the patient with water of the temperature of the room (68° to 70°). As the water became heated by contact with the skin, small pieces of ice were put into the basin so as to keep the temperature at the point mentioned. At the end of four hours, the washing having been continued all the time, we found the patient decidedly more comfortable. The pulse had fallen to 140, and increased in volume; the heat of skin was much reduced; the color of the eruption had improved,

having become much more scarlet in tint; the capillary circulation was more active; the agitation and restlessness had very much moderated, and the child had slept somewhat at short intervals. This treatment, in conjunction with the internal administration of the solution of chlorinated soda, and small doses of oil of turpentine, was continued for several days, the sponging being used whenever the heat and restlessness were great, and the pulse very rapid. The child convalesced about the end of the third week, but was unfortunately seized with uræmic symptoms on the twenty-fifth day, and died in twenty-three hours, after the most frightful convulsions we ever saw.

Since the publication of the last edition of this work we have seen but few cases of scarlet fever, and our experience as to the exact value of cold has not been much increased. It was used by our advice in the following case, of the most violent type, to which we were called in consultation:

CASE.—The patient was a girl, two years old, who, seized in the morning of one day with vomiting, fever, and restlessness, had, during the following night, high fever, violent jactitation, and moaning. She refused all food. Next day, at 9 A. M., when we saw her, she was dangerously ill. She knew no one, paid no attention to father or mother, tossed incessantly about the bed, or in their arms, so that it was almost impossible to hold her; and at times had rigid contractions of the muscles, like those in tetanus. The features were drawn and rigid; the pulse running up to 180, very feeble and small; the skin very hot, but without any eruption. The latter fact might have thrown some doubt on the diagnosis; but the character of the symptoms, the rapid fatality, and the fact that a few days afterwards two children were seized with distinct scarlet fever in the house opposite, left no doubt in our own mind. The prognosis was as bad as it could be, and so we announced, but added that external cold ought to be tried. The body temperature was very high, and we directed basins of water with ice in it, as used by Dr. Corson, to be prepared. Towels wrung out of this water were kept on the head, and the body and limbs sponged with the same until the heat fell, when the washings were suspended temporarily, to be renewed when the heat rose again. The treatment was carried out very correctly, as there was a medical man present all the time, but it was of no use whatever. The child died at 12 M. of that day, in a little over twenty-four hours from the onset.

This case was not a fair test of the value of the treatment. The cold was applied too late to show clearly what may be its power. But we confess that its total failure, though used within twenty-four hours of the inception of the disease, is a melancholy proof of the extreme danger of such cases.

A second case, which occurred about the same time, also shows the violence of this form of the disease.

CASE.—A very healthy girl, within a few days of two years old, whose sister and two of whose cousins had been ill with scarlatina in the same house for some ten days, was seized at six in the morning with vomiting. She then slept for a time and vomited again. At 9 A. M., she had a convulsion, which lasted, with short lulls, during which she was comatose, until 6 P. M., when one of us saw her in consultation. She was then very hot, covered with a copious, dark, dingy eruption, and insensible. Despairing of any other treatment, we advised cold externally, and arranged for its use by a physician, with the thermometer as a guide. Before the treatment could be commenced, the child became again convulsed and died.

After thus stating the conditions under which we think external cold may be properly used, we must protest against its indiscriminate use in all cases of dangerous scarlet fever. If the reader will glance back at page

815, he will see what Currie thought of the rash use of his cold affusions, and what Dr. Gregory also states of the effects of cold.

When the body, instead of being hot, is cool—when a dingy and stagnant capillary circulation shows a feeble and struggling heart, it would be most dangerous to use cold. Here the warm or tepid bath or affusion should be used, or warm mustard foot-baths may be resorted to every two or three hours. If, even whilst the body and limbs are cool, the head is hot, it would be proper to apply cold by cloths or affusion to that part, whilst the body is immersed in warm water or wrapped in proper coverings.

The true guide as to the propriety or impropriety of using cold is, none can doubt now, to be found in the thermometer. The method followed by Drs. Wilson Fox and H. Weber in the hyperpyrexia of rheumatic fever, is the one which we propose to use ourselves, and to recommend to others. It is the only scientific one, and therefore the one which can be accurately described and followed. If errors occur, they can be definitely stated and afterwards avoided. If successful, the exact means which led to success can be ascertained and communicated in precise language. Dr. Fox first published his cases in the *London Lancet*, and then presented them in a separate form as an essay "On the Treatment of Hyperpyrexia, as Illustrated in Acute Articular Rheumatism, by Means of the External Application of Cold;" Macmillan & Co., London, 1871. Dr. H. Weber's case is to be found in the *Transactions of the Clinical Society of London*, vol. v., p. 136, under the title of "A Case of Hyperpyrexia (Heatstroke) in Rheumatic Fever Successfully Treated by Cool Baths and Affusions."

The first point to be determined is the degree of febrile temperature dangerous to life, to prevent or reduce which we must resort to the application of cold externally. Of course there is but one certain guide to the temperature of the human body—the thermometer. A practiced hand may be relied on to a certain extent; but no hand, however experienced, can give the certainty of the thermometer. Inasmuch, too, as the state of hyperpyrexia is always attended by concomitant phenomena of a peculiar kind, these, to the experienced physician, will assist in guiding him in his treatment. These phenomena constitute the group called ataxic or adynamic nervous symptoms. The patient is usually delirious and restless or comatose, and not unfrequently has local or general convulsive movements; the pulse is frequent and feeble, and sometimes so small as to be felt with difficulty, and the capillary circulation is sluggish and congested; the respiration is usually hurried and embarrassed, so that the patient is readily judged to be in extreme danger.

According to Dr. Fox, a temperature in rheumatic fever which rises suddenly from 103° or 104° to 107° , 108° , or 109° , has usually proved fatal within a very short time after the latter temperature (109°) has been reached. He, however, saved one patient, by external cold, in whom it reached 110° in the rectum. Dr. H. Weber thinks that until the cold treatment was used, a temperature of 108° had been nearly always fatal. Dr. Fox asks the question, After what degree of temperature attained by the human body in febrile states is recovery naturally possible without

medical interference? He states that the highest recorded temperature he knows of after which recovery has taken place (with the exception of relapsing fever) was in a case of tubercular pneumonia, in which it rose suddenly from 105° to 108° , and then fell as suddenly to 104° . He refers, of course, to cases not treated by cold, since, as stated above, he himself saved a case in which the temperature had reached 110° in the rectum.

It must not be forgotten that the axillary temperature is lower than that of the mouth, under the tongue, and this less than that of the vagina or unloaded rectum. Wunderlich gives the averages in the adult as follows: The axilla, 98.6° F.; the mouth, 98.78° to 98.96° ; and the vagina or unloaded rectum, 99.14° to 99.5° F. In children the temperature is more variable, but does not differ very greatly from that of the adult. M. Roger gives 98.97° as the average axillary temperature between 4 months and 6 years, and 99.15° between 6 and 14 years. Dr. Finlayson, in 21 children under 6 years, found the morning temperature in the rectum to be 99.41° F.

Such being the normal temperatures, we will now give those which have been observed in scarlet fever. Wunderlich (*Med. Thermometry*, Syd. Soc. ed., p. 348) says that the height reached by the temperature in scarlet fever is almost always above 104° F., very commonly over 104.9° , while in cases which terminate favorably it seldom exceeds 105.8° F. The translator of Wunderlich (Dr. Woodman, footnote, p. 221) gives the noon temperature of typical non-malignant scarlatina in a good many cases at 105° , 104° , 103° , and 102° , on the first, second, third, and fourth days. In a note at page 204, he states that he has put on record (*Med. Mirror* for February, 1865) some fatal cases of scarlet fever in which the temperature reached 115° . "The observations were made with one of Negretti and Zambra's thermometers, divided into fifths, which had been recently compared with a standard."

From these facts we may assume that a temperature of 105° F. in scarlet fever is not necessarily very dangerous to life, but that from the moment it tends to rise above this point, the patient enters into a very dangerous period.

If, with a temperature of 105° , there appear any of the nervous phenomena so often alluded to, delirium, drowsiness, coma, vomiting or purging, and especially any locomotor disturbances, the time has come for the use of external cold; and should the temperature continue to rise after it has reached 105° , the cold treatment ought to be resorted to, even though these nervous phenomena have not shown themselves, since they will be almost certain to appear should the temperature go on rising.

And next as to the best mode of applying cold. It does not matter much how this is done, if only it be so managed as to reduce with certainty the heat of the body towards the normal point. Dr. Fox used baths at different temperatures, and in one case applied ice to the chest and along the spine in an ice-bag, whilst he reduced the temperature of the bath rapidly from 96° to 66° . At other moments in the same case he used the ice-bag applied to the spine for several hours at a time, and on

still other occasions employed the cold pack, wrapping his patient in a sheet wrung out of ordinary cold water (probably 60°). Dr. H. Weber placed his patient (a boy of 16) in a bath at 71° F., keeping him there the first time thirty minutes, when the temperature under the tongue had fallen from 108.2° to 101.8° . Some hours afterwards, when the temperature had risen to 105.8° , the patient was again put into a bath at 72° , and water poured over the back of the head and neck. In twenty-five minutes the temperature fell to 101° .

Dr. Fox says, at page 34: "I believe, however, that the bath may be altogether dispensed with, and that for the future it will be sufficient to place a Mackintosh sheet under the patients, so arranged that the water may escape into a receptacle, and to pour cold water over them from time to time."

Dr. Corson uses affusions of cold water over the head of the child, which is held over a tub, as the most powerful means in cases of coma or convulsions, and, when the temperature is very high, washes the whole body with iced water, or even rubs it with ice.

The most convenient mode, it appears to us, in children, will be either the one proposed by Dr. Fox, the Mackintosh sheet on the bed, and affusions of cold water; or a bath-tub or common large wash-tub, containing water at a temperature proportioned to the heat of the body, 80° to 70° , with affusions of cold water upon the head, or the application of towels, wrung out of cold or iced water, to the head. So soon as the thermometer, held in the rectum or under the tongue, shows that the temperature has fallen to 101° or 102° , it will be best to remove the patient to bed between two blankets. Not unfrequently, as the temperature falls, the patient becomes partially conscious, grows pale, and shivers. When these signs appear it is time to cease, at least temporarily, the use of the cold.

The physician, when he first uses this mode of treatment, should know that the temperature is apt to continue to fall, even after the use of the cold has been suspended. Thus in one of Dr. Fox's patients it fell from 103° to 99.4° after the removal from the bath. The danger to be apprehended from these continued falls in the temperature is not so great as might be supposed. Thus, Dr. Fox says that it may be doubted whether, in future cases, any external warmth may be necessary to prevent too great a fall of temperature. "Even severe collapse produced by cold has been shown by F. Weber's, Bartel's and Ziemssen's observations on the pneumonia of children to be less dangerous than it at first appears." And Dr. H. Weber says, "Although the duration of the bath will be influenced in some degree by the temperature of the water, we must be entirely guided by the condition of the patient while in the bath; the heat of the blood ought to be reduced, if possible, to almost its normal degree, and the nerve-centres ought to be reduced to a more healthy condition."

After the heat has been once reduced by the cold to near its normal state, the patient must be carefully watched by means of the thermometer, and if the temperature rises again, the cold should be reapplied. This may have to be done several times a day at first, and less frequently afterwards, if the treatment prove successful. It is not always necessary to resort to the

bath for every rise of the heat. Cold to the head, affusions upon the head alone, or the application of an ice-bag to the spine, may suffice to keep the temperature within safe limits.

While the cold is being used to reduce the temperature, we may employ certain internal remedies with advantage. If the patient is not very much exhausted, we may make use of the antiseptic salts of Polli, to which reference will presently be made. But if the exhaustion be very great, if the circulation is rapid, feeble, and uncertain, with a dusky and congested skin, we should use brandy with milk, beef or chicken-tea, and wine-whey. Dr. Fox used in the two cases which recovered (adults) very large quantities of brandy, from twelve to eighteen, and even thirty-three, ounces in twenty-four hours. He gave also large amounts of beef-tea, two to three pints, and as much milk as three and four pints. In such cases quinia and carbonate of ammonia would also be proper means until the vitality is restored.

In a former edition of this work it was stated that we had used the hyposulphite of soda or magnesia in 11 cases, of which 2 were malignant in type, 3 grave, and 6 moderate. All these recovered. It was then said that so small an experience was of little weight in determining their value. Since that time we have used the same salts in most of the cases that we have seen. Only 4 of these could be called grave. They were not of the convulsive form, but exhibited high fever, severe anginose symptoms, tedious duration, and copious desquamation. They were severe, but not malignant cases, and they all did well. In two cases of the malignant form, with profound adynamic nervous symptoms from the very onset, they were also freely used, but without effect. On the whole, we think these salts deserve a further trial. The soda or magnesia salt ought to be selected according to the state of the bowels. When these are constipated, the magnesia is to be used. In the contrary case, or when the bowels are relaxed, we use the soda salt. The dose of either is five or ten grains, every two hours according to the age. They are best given in solution in water, with a little ginger syrup.

There is a class of cases which, though they do not exhibit the extreme severity of those we have just considered, well deserve the name of grave. The temperature is high, the pulse rapid, the nervous system shows disturbance by extreme agitation or by drowsiness; there may be muscular starting, or tremors, or a single slight convulsion; the eruption is very abundant, and vivid or dark in tint, and the anginose symptoms are marked and severe. Such cases are dangerous; seldom last less than two or three weeks, and require all the care of the practitioner and nurse.

In this second grade of the grave form the temperature ought to be reduced, using the thermometer as a guide, by the careful use of the cool or tepid bath, or of cool or tepid ablutions, and by the use of cold water or ice to the head. Internally, the hyposulphite salts or an alkaline febrifuge ought to be administered for the first few days. In addition to this, we may employ with advantage full doses of quinia or salicylic acid; both of which, and especially the latter, possess the property of greatly reducing the degree of febrile heat.

The patient often, indeed generally, in this class of cases, sinks after a few days into a low ataxic condition. Here the best remedies, we think, are chlorate of potash with muriated tincture of iron, quinia with muriated tincture of iron, or muriated tincture of iron with solution of acetate of ammonia and dilute acetic acid. Hillier is of the opinion that carbonate of ammonia is the best remedy in such cases, and Trousseau also advises it strongly. These medicines may be given in the following manner:

R. Potass. Chlorat.,	3j.
Tr. Ferri Chloridi,	f 3 ss.
Syr. Zingiberis,	f 3 vij.
Aquæ,	f 3 ij.—M.

Dose—A teaspoonful every two hours at five years of age, and under that age, half a teaspoonful.

R. Quiniæ Sulphat.,	gr. xij.
Tr. Ferri Chloridi,	f 3 ss.
Syr. Zingib.,	f 3 v.
Syr. Simp.,	f 3 ij.
Aquæ,	f 3 ij.—M.

Dose—A teaspoonful every two hours at five years, and under that age, half the quantity.

R. Tr. Ferri Chloridi,	f 3 ss.
Acid. Acet. Dil.,	f 3 j.
Liq. Ammon. Acetat.,	f 3 vj.
Syr. Simp.,	f 3 ss.
Aquæ,	f 3 jss.—M.

Dose—A teaspoonful every two hours at five years of age, and under that age, half the quantity.

The dose of carbonate of ammonia is from one to two grains every two hours at five years, given in a mixture of syrup and gum, or in milk and water.

In all severe cases attention to the diet is highly important. At first only milk and broths should be allowed. After a few days bread or some other farinaceous substance may be added. Until the fever has disappeared in great measure, no other diet ought to be permitted. The cases are very uncertain in their course.

Under certain conditions alcohol must be used as has already been said. When the tongue is dry, the skin harsh, the pulse rapid and feeble, the cardiac impulse weak, the muscular force reduced, we use brandy or whiskey, or wine in the form of wine-whey, or mixed with water. The choice between these agents must depend on the degree of vital, and especially of circulatory, prostration present, and the fancy of the patient. We use brandy generally, giving it in milk or water. From ten to twenty drops at the age of one or two years; from twenty to forty drops between two and five years of age; and after these ages from half a teaspoonful to a teaspoonful every two or four hours, or three or four times a day, according to the effects produced upon the pulse and nervous symptoms, are the doses we have found best. When wine-whey is preferred, one or two teaspoonfuls every hour or two hours, at two years of age, and a table-spoonful at the age of four or five and upwards, may be given.

Treatment of the Angina.—The angina is seldom troublesome before the third day. It never becomes, we think, a source of danger in itself, in the cases destined to end fatally on the first, second, or third day. But, when the disease begins with grave nervous symptoms, and the patient survives these, the throat almost always exhibits, on the third or fourth day, the conditions which have already been described, and which partake so much of the character of severe diphtheria. The general treatment ought to be steadily preserved in,—that by the hyposulphites, or by the muriated tincture of iron, with or without cholorate of potash and quinia, as has been described. The local treatment should consist, in the early stage, of cold applications, if the constitutional state of the patient will allow of it. When the circulation is active, and the temperature of the body high, there need be no fear as to the use of cold. When, on the contrary, the heat is not high, it should be used with caution, and if the temperature falls rapidly under its use, it must be abandoned, or used only from time to time, and with great care. When the temperature continues rather low, warm poultices, inclosed in portions of thin, soft flannel, and secured by a light cravat round the neck, may be tried. They may prove comforting to the patient.

When the cold is to be used, pieces of ice wrapped in flannel, and applied behind the angle of the jaw, or cloths wetted with iced water, may be employed. We have used the cold several times, and in two cases with marked benefit. In one case, which we shall relate, the effects were most striking.

CASE.—The patient was a boy between eight and nine years of age, who had had slight convulsive movements and delirium on the first day, violent jactitation and unconsciousness on the second and third days, with very active pulse, profuse dark eruption, and very high temperature. There appeared on the third day threatening anginose symptoms. On the fourth these had increased, and, by the night of the fifth day, had reached such a height as to make us almost despair of the child's life. The violent pharyngeal inflammation was attended with excessive swelling of the tonsils, and with œdematous infiltration of the submucous tissue, while externally the cervical glands were enormously enlarged and hard as paving-stones, and the subcutaneous tissues of the front and lateral regions of the neck packed and hard with acute œdema. The general symptoms were most threatening. Owing to the swelling of all the parts composing the neck, the respiration was so interfered with as to cause the development of dangerous asphyctic symptoms. The pulse, which for the first three days had been running at 168, had fallen on the fourth to 140, and on the fifth to 128; the skin was hot and dry, and the face had assumed a dark, bluish tint; there was almost constant muttering delirium and a degree of tossing and violent jactitation painful to witness. The swallowing was so much impeded that it was with great difficulty that the patient could take the thinnest liquids. Up to this time the case had been treated with inunctions, cold drinks, and a febrifuge containing spirit of Mindererus and sweet spirit of nitre. In the midst of these threatening symptoms, and when we had almost lost hope, the late Dr. Charles D. Meigs, who saw the case with us, proposed the withdrawal of all drugs, and the use of cold applications externally, and stimuli. Accordingly a large towel was wrung out of iced water and wrapped around the neck, and weak wine and water was given as often as the child could take it. The cloth was dipped afresh into the water every few minutes. This treatment was commenced about 1 A. M., and carried on steadily all night. At 9 A. M. it was evident that the symptoms had somewhat improved, and by the afternoon of that day the patient was

greatly better. The improvement consisted principally in a moderation of the pharyngeal swelling, so that both respiration and deglutition were much easier. The dark color of the face had lessened; the pulse had risen in frequency, and was stronger; and the delirium and excessive jactitation had almost disappeared. On the day after this the external cervical swelling continued very much the same, except that the oedema had notably diminished. The pharyngeal swelling had disappeared, the tonsils having regained their natural size, but the whole pharynx was covered with a thick mould of white exudation. The cold application, which of late had been used more sparingly, was now discontinued; the fauces were touched with a solution of nitrate of silver of ten grains to the ounce; broths, milk, wine-whey, and wine and water were given for nourishment, and the patient recovered at the end of the fifth week, after having had a large suppuration just above the inner end of the left clavicle.

It was at one time very much the custom to make various applications to the fauces. Nitrate of silver, pure, or in strong solution, muriatic acid, or capsicum, were deemed necessary and useful. They have been very much abandoned, and we think wisely. The agitation and terror caused by them in some children, and the violent resistance they often make, exhaust the patient, and we therefore avoid them whenever we can. Sometimes, however, and especially in young children, viscous secretions collect in the fauces in such quantity as to cause serious annoyance to the child and embarrass the respiration; they ought, therefore, to be removed by means of a sponge-mop or camel's-hair brush. This point in the treatment is a very important one, especially in young children. We believe that we have rescued more than one patient, by going three or four times a day, to make use ourselves of means by which to remove from the fauces, the viscid, glue-like secretions, the purulent fluids, and the masses of pseudo-membranous exudation which collect in and occlude those passages, and which the child often cannot, by any effort of its own, get rid of. The best mode of effecting this object is by the use of mops, made of sponge or rags, fastened to a stick or whalebone, or by the injection from a small syringe, or elastic bottle, of detergent washes or gargles into the throat, the mouth being held open and the tongue depressed by the handle of a spoon. One of the best washes or injections is made of a strong decoction of green tea containing alum; or we may employ sage-tea and alum; or honey of roses and borax mixed with water; or lime-water; or what is highly recommended by Dr. Watson as efficacious, a solution of common salt. For local application by means of a pencil or mop, the following mixture is one of the best of the many we have tried:

R. Acid. Carbolicæ Crys.,	gr. x.
Liq. Ferri Subsulphat.,	fʒj.
Glycerinæ,	fʒj.—M.

To be used two, three, or four times a day.

Muriated tincture of iron, one part to five or six of water, or to one of glycerin and five of water, is another excellent local application. In older children, gargles of salt and water, alum and water, chlorate of potash, in claret and water, or plain water, may, and ought to be used, when possible. In many cases, even in young children, it is possible to secure the inhalation of fluids vaporized by the steam atomizer. It will

be found that the frequent inhalation of lime-water exerts a very favorable effect of the condition of the throat. When coryza is present, the nasal passages should be cleansed by means of camel's-hair brushes, or by the injection of some of the mild washes just referred to, and then freely anointed with sweet oil or some mild ointment, or they may be touched with the wash used for the throat.

Diarrhœa, when present, probably depends on congestion and desquamation of the intestinal mucous membrane, and should be treated with bland demulcent drinks, and absorbent antacids, especially chalk mixture.

Rheumatism is to be treated by opiates to allay the pain, and the swollen joints should be enveloped in bats of cotton. If suppuration occur, either in connection with the rheumatic inflammation of the joints, or involving the glands or cellular tissue, and indicating a pyæmic tendency, large doses of quinia with stimulants must be given. The abscesses which may form should be opened so soon as fluctuation can be detected.

For the *otorrhœa* which sometimes occurs, it is seldom necessary to do more during the violence of the attack, than to cleanse the ears twice or three times a day, by syringing with warm water and castile soap, or with a weak solution of alum. After the violence of the attack has subsided, this complication should be treated as in idiopathic cases.

Treatment of Dropsy.—It is our habit, when directing the general treatment of a case of scarlet fever, always to explain to the mother or nurse, or both, that the most frequent and dangerous sequel to be apprehended in this disease is dropsy, that this is even more apt to follow mild than grave cases, and that it usually occurs in the third or fourth week of the disorder, though it does occur, on very rare occasions, at a still later period. We also assert our belief that this consequence, or complication, or sequela, is apt to be produced by cold, and that subjects confined to bed through the third and fourth weeks, and those rigidly secluded in a warm room for four weeks from the onset of the disease, no matter how mild the case, are much less prone to dropsy than they who leave their beds or rooms at an early period to take the usual chances to which children are exposed. We know well that Hebra ridicules the stupidity of English physicians in ascribing so many disorders to cold. But, whilst we believe that in England, and amongst ourselves, the word "cold" is often used both by medical men and by the public as a scapegoat to bear the weight of our ignorance as to the real cause of disease, we also believe, most emphatically, that chilling of the body, if it be continued for any length of time, is very apt to be followed by some disturbance of the health. We are quite sure that we have on several occasions, traced a relation of cause and effect between exposure to cold in the third or fourth week of scarlet fever, and a rapidly sequent dropsy. Several such cases are mentioned in the section on the symptoms and causes of dropsy at page 796. We always, therefore, urge upon the mother or nurse not to allow the child, in the autumn, winter or spring months, no matter how mild may have been the disease, to leave a well-warmed and well-ventilated room for four weeks, and, if there be any sign whatever of ailing health at the end of the fourth week,

to continue the seclusion for one or two more weeks. One of the annoyances in a private medical career is the contention one has with people in their hurry to get children, who have been sick, out of the nursery. In summer weather, this may be all very well, but in the cool and cold months it is not very well, as the bills of mortality and the experience of any older physician, or experienced mother, will show. Why a child should lie abed for one or two months for a broken bone, without fear for his general health, and yet be regarded as a suffering martyr, because some tyrannical doctor insists upon his remaining a few more days or weeks in a comfortable chamber, with all the household at his feet, to escape a disease like acute Bright's disease, passes our comprehension.

Before leaving this subject we wish to say that we have known dropsy to follow scarlet fever in only one or two cases, in which the child had not left the bed. One of these cases has been fully described in the remarks upon dropsy.

It is important to recognize the renal disease early. The mother ought to be warned to send for the physician again, if he have resigned the charge of the case as being convalescent, should there be any delay or irregularity in the convalescence, and especially should she observe an unusual scantiness in the quantity of urine discharged, a dark and especially a brownish or blackish tint of this fluid, fulness of the eyelids, swelling of the cervical glands, or, indeed, any departure from a regularly progressive return to health.

In all cases of scarlatinous dropsy, the patient ought to be put to bed at once, and kept there throughout the acute period of the disease. The diet must be restricted to fluids. Milk and animal broths or farinaceous preparations alone ought to be allowed. The patient should be encouraged to drink freely and often of water, lemonade, or orangeade, or sweet spirit of nitre and water. A hot bath, used once or twice a day, is, we think, one of the most important of all remedies in the early stage. It is best used in the following mode: A portable bath-tub should be brought into the bedroom, if possible. This can always be done in the cases of young children. The water must be warm— 96° to 98° or 100° . The patient ought to be fully immersed, and kept in the water from ten to thirty minutes, the time being regulated by the degree of willingness of the child to remain, and by the effect of the bath on the system at large, as shown by the countenance and circulation. A soft cotton sheet is to be heated at the fire, and in this, when the patient is removed from the bath, he is to be carefully wrapped. Over this is to be put a light blanket, and thus wrapped in the two coverings, the child is to be laid in bed, or held in the arms, for half an hour or an hour. By this procedure sweating is generally induced. When this is over, the sheet and blanket may be removed and the child dressed in warm bedclothes again. The bath, carefully used in this way, once a day in slight cases, and twice or even three times in severe ones, has proved in our hands a potent means of cure.

In mild cases, without fever, the bowels ought to be soluble, no medicine being needed if they are removed spontaneously. If they are not, a little syrup of rhubarb or Rochelle salts will be all-sufficient. If the

amount of urine is scanty, a diuretic ought to be used. The following combination is excellent :

R. Potass. Bitart.,	3j.
Spt. Junip. Comp.,	f3 ij.
Spt. Æther. Nitros.,	f3j.
Tr. Digitalis,	℥ xv.
Syr. Simp.,	f3 v.
Aquæ,	f3 ij.—M.

Give a teaspoonful every two hours, at two, three, or four years of age.

In more severe cases, when vomiting, fever, anorexia, restlessness, rapid anasarca, scanty and dark-colored urine with blood, blood and granular casts, and a large proportion of albumen, all demonstrate a serious and extensive catarrh of the renal tubules, it is proper to use dry cupping to the loins, or in subjects of vigorous constitution, uninjured by the previous scarlet fever, we may take three or four ounces of blood from the loins by wet cups. If the cupping cannot be used, hot cataplasms of Indian mush or flaxseed or bags of hot sand should be applied from time to time over the loins. The bowels ought to be kept open by rhubarb, Rochelle salts, or Seidlitz powder. A febrifuge and diuretic, such as the following, must be used :

R. Potass. Acetat.,	3j.
Tr. Digitalis,	f3 ss.
Syr. Scilæ,	f3j vel f3ij.
Syr. Zingib.,	f3 v.
Aquæ,	q. s. ad f3ij.—M.

Give a teaspoonful every two or three hours to children two or three years old. For those above that age the proportion of the active ingredients should be doubled.

We desire to call the attention of the reader, especially if he be a young physician, and therefore disposed to trust overmuch to mere drugs, to the necessity of supplying to a child needing diuretics an abundant amount of water. The diuretic does but stimulate the action of the kidney, and to enable it to do this, the supply of water by injection should be copious. A child laboring under renal catarrh following scarlet fever, rarely, perhaps never, takes too much water. The fact that the patient vomits water is no proof that his system does not need that liquid. He vomits sometimes for hours everything given him, and this is one of the dangers of the disease. Like the traveller who has been deprived of water for several days, and who vomits repeatedly the first supplies he obtains, so the child, in the condition we are considering, must not be deprived of the element essential to its safety, because it rejects a portion of what is allowed it. Water should be offered in various forms,—lemonade, orangeade, weak brandy and water, claret and water, water in which the white of an egg has been incorporated by careful and prolonged beating (the albuminous water of the French), very thin chicken-tea touched with salt, thin milk and water with a little tea, Liebig's cold beef extract made from raw beef, thin cocoa or chocolate, or, indeed, anything not plainly injurious which the fancy, tastes, or habits, of the patient may indicate, may and ought to be tried. As to the quantity of water there should be no stint,—as much and as

often as the child can take and retain, is our rule. It will not drink plain water after its thirst is assuaged, any more than will the adult man, or a horse, until the supply in the body is exhausted, and more is never needed for the physiological uses of the body.

By these means the case should be treated for several days, until the fever subsides and the patient is no longer in present danger. When the fever is over, it is best to continue the above mixture, or something of the same kind, at longer intervals, and to give also one of the preparations of iron. Our own favorite is the mixture of muriated tincture of iron with acetic acid and spirit of Mindererus, or the simple tincture itself, in doses of from two and a half to five drops, according to the age, three or four times a day. If from any cause the tincture cannot be taken, wine of iron, or ferrated elixir of Calisaya bark, may be substituted, in half-drachm or drachm doses.

As the fever disappears, the food ought to be increased. This is, of course, more important than drugs, and ought to be strictly attended to by the physician himself at each visit.

In some of these cases the most perverse irritability of the stomach exists for several days, so that the patient may almost or actually die from exhaustion. In such cases it is worse than folly to give drugs which are resisted with loathing, and vomited the instant they enter the stomach. Something must be chosen which at least does not clearly cause vomiting by its smell or taste. A mixture of wine of iron with syrup of Tolu and some aromatic water, chocolate lozenges with iron, or powdered metallic iron with white sugar, can often be taken even in these cases. A weak cream of tartar lemonade, flavored with lemon-juice, sweet spirit of nitre in lemonade, watermelon-seed tea, and such remedies may be used. In regard to details, as to the best method of feeding in such cases—on which, we desire to say, much more depends than upon drugs—we must refer the reader to the remarks on diet in obstinate sick stomach in chronic diarrhœa. We venture to hope that we have seen lives saved in cases of this kind by constant attention to little details of food and medicine, which must have been lost by any less constant care. In one instance the child was almost comatose, with total suppression of urine for five days, and the stomach so irritable that no remedy scarcely could be borne. Finally, under small doses of watermelon-seed tea, given frequently, mustard foot-baths, blisters behind the ears, and feeding with lime-water, milk and brandy, wine-whey, chicken-tea, and such preparations, the patient recovered. When the stomach refuses everything, the patient may be fed for a few days by the rectum, unless there be diarrhœa or unusual irritability of the intestine. One or two ounces of beef-tea, of chicken tea, of extract of raw beef, of lime-water and milk, of water with raw egg incorporated in it, or of plain water, may be given every two or three hours, and continued as long as they are retained or needed.

PROPHYLACTIC TREATMENT.—It was formerly asserted that *belladonna*, used by persons exposed to the contagion of scarlet fever, had the power of imparting perfect or nearly perfect immunity from its attack. The

evident difficulty of determining a question such as this, in reference to a disease so uncertain and irregular in its mode of extension as scarlatina, long maintained a certain degree of doubt as to the possibility of the truth of this most unlikely assertion. We believe, however, that by common consent, all belief in the supposed efficacy of belladonna for this purpose has now been abandoned.

Dr. Brakenridge (*Medical Times and Gazette*, July 24th, 1875, p. 92) has given sulpho-carbolate of sodium in doses varying, according to age, from five to thirty grains three or four times a day, to those exposed to the poisons of scarlet fever, diphtheria, and measles; and, although not feeling that his observations have proved the power of this treatment to prevent attacks of these diseases, he is inclined to attribute some prophylactic power to it.

In order to purify articles which have been exposed to scarlatina, they should be either put in boiling water or exposed to a temperature of over 200°, as we have seen that a temperature somewhat below the boiling-point of water destroys the activity of the virus.

ARTICLE VI.

MEASLES, RUBEOLA OR MORBILLI.

DEFINITION ; FREQUENCY ; FORMS.—Measles are an epidemic and contagious exanthem, characterized by catarrhal symptoms, continued fever, and an eruption, generally on the fourth day, of a crimson rash, in the form of stigmatized dots, like fleabites, slightly elevated, which coalesce into irregular circles or crescents. It ends about the seventh day by desquamation.

The frequency of the disease is very irregular in different years because of its epidemic nature. Thus, according to the mortality tables of the Board of Health, there have been 2279 deaths from measles in this city during the sixty years ending with 1870. In five of these years, as will be seen by a reference to the table at page 769, there is not a death recorded from this cause, while, on the other hand, the annual mortality exceeds 100 in eleven years, and 200 in two. During the same period, the deaths from scarlatina in this city, as already stated, amounted to 13,016.

Measles are probably a more common though a less fatal disease, and attack a larger number of persons than scarlet fever: thus, during a period of fifteen years, during which we noted every case that occurred in our practice, we met with 314 cases of the former to 263 of the latter.

We shall describe two *forms* of the disease: the *regular* or *rubeola vulgaris*; and the *malignant* or *rubeola maligna*. We shall afterwards treat of its *irregularities* and *complications*.

CAUSES.—A chief cause of the disease is *epidemic* influence. Of this there can be no doubt, as it is proved by the evidence of all observers.

Contagion.—That it is a contagious disease is universally admitted. The

contagious quality is thought to begin with the primary fever, and to continue up to the period of desquamation, though some authorities believe that it is also contagious during the stage of incubation. That the disease is contagious prior to the appearance of the eruption seems to be proved by observations like the following: A child living in Philadelphia visited a relative in the country, and returned to the city the same day. On the next day the child in the country showed the measles eruption, and twelve days afterwards, the one in the city sickened with the same disease. The precise period at which the contagious property disappears is not, however, known. The disease may be carried in fomites. It has been propagated also by inoculation with the blood taken from a patient, and with serum obtained from the vesicles which sometimes accompany the eruption.

The period of incubation is difficult to determine, but is usually stated as from five or six, to twenty days, or even longer. In the great majority of cases, however, the eruption appears in from twelve to fifteen days after exposure to the contagion, thus making the duration of the period of incubation from nine to twelve days. Thus, in twelve cases where we were able to determine with precision the interval between the exposure to contagion and the appearance of the rash, it was ten days in 1 case, eleven in 1, twelve in 3, thirteen in 5, fourteen in 1, and fifteen in 1. In 108 cases observed by M. Girard, of Marseilles (quoted in *Med. Times and Gaz.*, Aug. 21st, 1869, p. 225), the eruption appeared as late as the sixteenth day only in 3 cases; in all the others it was developed on the thirteenth or fourteenth day, never before the thirteenth, and never after the sixteenth.

MM. Rilliet and Barthez conclude that measles are more frequent, less contagious, and have longer incubative and prodromic stages than scarlet fever.

The same authors are of opinion that variola is somewhat more rare, rather more contagious, and that its period of incubation and its prodromic stage are a little shorter than those of measles.

Measles, like other contagious diseases, rarely occur a second time in the same individual. We have, however, met with undoubted cases of second attacks; although unquestionably in a large proportion of the numerous cases in which we have been told that two attacks had occurred, one of them had been not of true measles, but of r  theln or roseola.

Age.—We find by uniting Dr. Emerson's tables with some given by Dr. Condie (*Dis. of Child.*, note, p. 100), that the disease appears to be most frequent between the age of one and two years, for while 395 deaths occurred in the second year, only 468 occurred between two and five years of age. This does not agree, however, with our own experience, since of 280 cases of the disease that have come under our own observation, in which the age was accurately recorded, only 36 occurred in the second year, while 84 occurred between the end of the second and the end of the fifth year. This discrepancy depends probably, in part at least, on the greater mortality of the disease during the earliest years of life, which would of course give a large number of deaths for those attacked in the second, than for those in the third, fourth, and fifth years. The cases that

have come under our own observation occurred as follows. They are stated in their order of frequency. In the sixth year, 37; in the second, 36; in the seventh, 35; in the fifth, 34; in the fourth, 30, in the eighth, 27; in the first, 19; in the ninth, 11; and then in the eleventh, tenth, thirteenth, twelfth, and fifteenth.

Sex.—It appears to be more common in the male than in the female sex. Of 290 cases that we have seen, in which the sex was noted, 156 occurred in males, and 134 in females.

Fungous Origin.—In 1862 Dr. Salisbury, of Ohio (*Am. Jour. of Med. Sciences*, July and October, 1862), published two elaborate articles, in which he attributed measles to the action of the fungus developed on damp, mouldy straw. He reported the results of numerous cases in which this fungus had been inoculated, with the production of a modified form of rubeola, which, however, protects the system against a future attack of true measles; and also instances where measles had broken out in camps where damp straw was used for bedding.

A complete examination of this question, embodying the evidence of Dr. Woodward (*Camp Diseases of the U. S. Armies*, Philadelphia, 1863), and the experiments of Dr. C. E. Smith, and one of ourselves, will be found in a paper by Dr. H. C. Wood, Jr., in *The American Journal of the Medical Sciences*, October, 1868, p. 342.

The results of the inoculation of nearly 50 cases, prove that in nearly every instance, the introduction of the straw fungus into the system is entirely without effect; and that in the few cases where any symptoms have followed, they have not been those of true rubeola, nor have they protected the system from an attack of genuine measles.

In regard to the occurrence of camp measles also Dr. Woodward remarks, that it prevailed almost exclusively in regiments raised in the rural districts, while those from cities and towns were more or less completely exempt; and that the inevitable inference from this, confirmed by personal inquiry, is that the recruits from the country had generally escaped the disease before their enlistment, while those from towns had usually suffered from it at some previous period; a condition of things entirely at variance with the idea that the straw fungus is the veritable cause of measles.

SYMPTOMS; COURSE; DURATION.—*Regular Form of the Disease.*—*Stage of Invasion.*—Measles begin with languor, irritability, sometimes chilliness, anorexia, aching in the back and limbs, fever, thirst, headache, and various signs of irritation of the mucous membrane of the eyes, nose, fauces and larynx.

The chilliness or horripilations which are mentioned by almost all writers are difficult to appreciate in children. We have seldom known the child itself to complain of them, but upon inquiry of the mother or nurse, have sometimes been told that they had observed some coolness of the hands or feet, or a disposition to keep near the fire, and a desire for additional clothing. These, therefore, are not important symptoms. Neither is the aching in the back and limbs, as it is seldom complained of, and can be ascertained in the older only by close questioning, or suspected in the younger by their complaining when they are moved. Fever

is rarely absent, but is often very moderate in degree. It almost always comes on with, or very soon after the other prodromes, but in rare cases does not begin until the second day. It is almost invariably continued, after it once begins, except that it remits somewhat about daylight and in the early part of the morning, to become exacerbated again in the after-part of the day. Its intensity increases, and the remissions become less distinct and shorter, as the time for the appearance of the eruption approaches. The pulse is increased in frequency, force, and volume, but rarely attains the same rapidity as in scarlet fever. At the same time the skin becomes warm and dry, the face is generally flushed, and there is considerable restlessness and irritability at first, often passing into quiet and drowsiness as the eruptive point approaches. The fever is accompanied by thirst, partial or complete anorexia, and generally by headache, which is frontal, and often complained of by children old enough to give an account of their sensations. The symptoms do not always follow this regular and orderly course. We have known a number of cases in which the approach of the disease was not even suspected until a copious rash made its appearance, so latent were the usual prodromes. In one case, a boy nine years old gave a party to his little friends. He danced himself all the evening, and next morning, when the windows were opened, was found covered with a copious measles rash. All the children of the family, six in number, and several of the guests, broke out with the disease in the usual time.

Vomiting occurs sometimes, but not as a general rule. The catarrhal symptoms commence with, or may precede the fever. They constitute the most characteristic early symptoms of the disease, and indeed the only ones by which we are able to distinguish it with an certainty in the first stage. They consist of irritation and redness of the conjunctivæ, especially that of the eyelids, lachrymation, suffusion of the eyes, sensibility to light, stuffing of the nose, coryza, sneezing, slight soreness of the throat, cough, some constriction of the thorax, and slight dyspnœa. The state of the eyes and nose are very important as signs of the disease. The above symptoms are not always present in the same degree, being very strongly marked in some instances, in others less so, and in some rare cases, absent. They are important, because there are few cases of ordinary cold in which they are present to the same extent, or if so, the accompanying general symptoms are slight compared with those of measles. We have rarely known the faucial affection severe enough to elicit complaints, and never to produce difficulty of deglutition. It consists generally only of slight redness of the tonsils, soft palate, and pharynx, which is most strongly marked about the time that the eruption makes its appearance. The cough usually appears on the first day. Infrequent and slight at first, it becomes more troublesome as the case progresses, until it assumes, on the third or fourth day, a character which is peculiar, and which may often lead to a suspicion as to the true nature of the attack. It is laryngeal, hard, dry, rather hoarse, and occurs generally in short paroxysms. Expectoration, if present at all, is slight, and consists of a clear, viscid mucus. At the same time the voice is often hoarse.

The tongue is usually white and somewhat furred; the bowels remain in their natural condition, or there may be slight constipation or diarrhœa. Constipation is most frequent, according to our own experience. The drowsiness, to which we have already alluded, often exists during the first stage. We have noticed it in a great many cases. The child, if undisturbed, sleeps quietly for many hours, or for the greater part of one or two days, waking only from time to time to ask for drink, and then sinking off to sleep again. So common is this symptom that old nurses have a saying,—“The child is sleeping for the measles.” The symptom is not alarming, unless it be connected with others which indicate local disease, or unless it pass into coma, or alternate with marked delirium. Other nervous symptoms which sometimes occur, especially when the fever is violent, are restlessness, irritability, occasionally delirium at night, and, in very rare cases, convulsions. Of 167 cases observed by Rilliet and Barthez, the latter symptom appeared in the first stage only in one, and was then confined to the eyeballs. We have met with convulsions in 5 out of 314 cases, at the beginning of the eruption, and in one, of which we shall not now speak, at the close of the eruption. In one of the cases the convulsions occurred on the first day, in a boy five years of age, of nervous temperament, and who had had several convulsive attacks during the process of dentition. The convulsions were general, but slight; they lasted only a short time, and were not followed by any bad consequences. In the second case the sickness began with fever, drowsiness, tremulous movements of the hands, delirium, and in a few hours a slight general convulsion. On the second day there were two attacks of convulsions, both, however, slight. The other symptoms continued as before. On the third day the child was better, the fever having diminished, and the nervous symptoms in great measure disappeared. On the fourth, fifth, and sixth days, the fever returned, and on the middle of the sixth day, a full measles rash made its appearance. There was no recurrence of the nervous symptoms, and the case ended favorably. The third case occurred in a boy between seven and eight years old, of nervous and impressible temperament. The convulsive seizure took place just as the rash was coming out; it was very slight, and lasted not more than one or two minutes. In the fourth case, in a boy in the second year of life, who had already had three convulsive attacks from other causes, showing thereby a manifest predisposition to that kind of disorder, the convulsions occurred as in the previous case, just at the coming out of the eruption. In this case also the convulsions were slight, lasting only a few minutes. In neither of these two cases were the convulsions followed by dangerous symptoms. In the fifth and last case, the convulsions, as in the two preceding examples, occurred just as the rash was appearing; they were very slight, and were followed by no serious consequences. The subject of this case was a girl between seven and eight years old, who had previously had an attack of convulsions produced by a severe febrile reaction occasioned by simple angina, and another attack, caused by indigestion.

MM. Guersant and Blache (*Dict. de Méd.*, t. 27, p. 658) mention another initial symptom, which has sometimes enabled them to recognize the ap-

proach of measles before the eruption. This is a peculiar redness, a rose-colored punctation, of the roof of the mouth, soft palate, and uvula, differing from that of scarlatina. We have observed this symptom ourselves in quite a number of cases, and, as it not unfrequently appears twenty-four hours before the cutaneous eruption has come out, we think that it is of some value as a sign in the early stages.

M. Girard (*loc. cit.*) states that the early diagnosis may be aided by the fact, that a red papule appears near the free border of the velum palati several days before the appearance of the eruption.

The *duration* of the initial stage is generally from two to three days. In a large majority of the cases that we have seen, the eruption has begun to appear in the course of the third day. This stage may, however, last only one or two days, or be prolonged to five, six, or seven, and according to Guersant and Blache (*loc. cit.*, p. 659), it lasted in one case, with all the characteristic symptoms, fifteen days. In a case that occurred to one of ourselves, the subject of which was a girl between one and two years old, the eruption, owing no doubt to the presence of severe general bronchitis, did not make its appearance until the ninth day of the sickness, and even then came out slowly and with much difficulty. The disease was known to be approaching from the fact that another child in the house had just recovered from an attack. In another case, in a girl between twelve and thirteen years of age, the eruption began on the fourth day of the sickness, but was so faint and indistinct that we could not, until the sixth day, feel sure that it was a measles rash. Even after this, the eruption continued pale and insufficient until the seventh day of the eruption, when it was out fully and completely.

Second Stage, or that of Eruption.—The eruption generally appears some time in the course of the third or fourth day, showing itself first on the chin or cheeks, or some other part of the face, and extending gradually to the neck and trunk, and finally to the extremities. It is often completed in from twenty-four to forty-eight hours. It begins in the form of distinct spots, not unlike fleabites, of a more or less bright rose or crimson color, verging sometimes towards a deep red, of a roundish shape, with irregular edges, and of different sizes, varying between half a line and three lines in diameter. When fully formed they constitute true papules, which are felt to be slightly elevated and firm to the touch, with broad flat summits. When pressed upon, their color disappears, to return rapidly when the pressure is removed. Distinct and scanty at first, the spots or stigmata soon become more numerous, and arrange themselves into clusters of an irregularly crescentic or semilunar shape. The number of these clusters and the consequent general tint of the skin, depend upon the amount and intensity of the eruption. In very mild cases, or when the eruption is imperfect, the clusters of papules are few in number, and they are separated by large portions of healthy skin. In severe cases, on the contrary, the patches are so numerous, and coalesce so closely, that the skin assumes a general deep-red tint. Occasionally in these severe cases minute vesicles form on the summit of the papules. Yet it ought to be remarked that it can be observed on close examination that the papules

never run completely into each other, so as to form a continuous red surface, unless it be over very small spaces and on certain parts of the surface, more particularly the face.

The fever does not diminish when the eruption makes its appearance, so that the highest temperature is usually attained soon after the full development of the rash. The skin retains its heat; the irritation of the eyes continues and is sometimes very severe; the nostrils are dry and incrustated, or there is coryza, and in some few cases epistaxis. The face is at the same time flushed, independently of the eruption, the red color of the skin being observable in the intervals between the papules, and it looks swollen and turgid, from tumefaction of the cheeks and particularly of the eyelids. The cough continues, and is loud, hoarse, and frequent in most cases, but in others short, scarcely hoarse, and but slightly marked. The voice is usually but not always a little hoarse. The respiration is slightly quickened in regular cases, but generally very little beyond the natural rate. The pulse is accelerated, though to a less degree than in scarlatina; its frequency is usually found to be in direct proportion to the height of the temperature. The tongue is covered with a yellowish or whitish fur in its middle, while the edges and tip are clean and red. It remains moist and soft unless some complication occurs. The tonsils, soft palate, and pharynx present considerable redness, without tumefaction. The abdomen commonly remains natural, though in some few cases there is slight soreness over its whole extent or in the iliac fossæ. Slight diarrhœa often occurs at this time. It seldom lasts more than from one to three days. In other cases the stools are natural, or there may be moderate constipation. The anorexia and thirst continue up to the stage of decline. About the time of the appearance of the rash there is often considerable restlessness, anxiety, starting, and twitching in sleep, slight delirium, and in children old enough to describe their sensations, complaints of headache. The strength of the patient is not decidedly affected in most of the cases.

The urine during this stage is scanty, of dark-yellow color, and not rarely contains a trace of albumen.

Stage of Decline and Desquamation.—The disease having reached its height in the course of the fifth or sixth day, the second of the eruption, it remains nearly stationary for one or two days longer, and begins to subside about the seventh or eighth of the disease, or third or fourth of the eruption. The eruption fades first on the face and neck, and has often very much or wholly subsided on those parts while it is still vivid on the extremities. The papules lose some of their color, become less prominent, diminish in size, and when pressed upon do not disappear entirely as they did at first, but leave a dull or yellowish stain behind. A little later they assume a dirty yellow or copperish tint, which does not disappear under pressure, showing that a slight ecchymosis has taken place into the substance of the derm. These stains continue a variable length of time, and are finally removed by absorption. As the eruption disappears, a slight furfuraceous desquamation takes place in a considerable number of the cases, but not by any means in all. This begins usually about the face,

and may either be limited to that part, or extend to other portions of the body. It is seldom general, however, and is often scarcely noticeable. It occurs between the eighth and eleventh days of the disease, or fourth and seventh of the rash.

From the moment the eruption passes its highest point of intensity, and begins to decline, the other symptoms do the same. The pulse lessens in frequency, and regains its ordinary characters. The heat of skin passes away, often with considerable perspiration, but sometimes with gentle moisture only. The various catarrhal symptoms subside; the cough is less frequent, loses its hoarseness, becomes softer, and generally ceases. The expectoration, if present, now becomes more copious and thinner, and presents nummular masses of muco-purulent matter floating in a clear, watery fluid. The tongue cleans off; appetite returns; thirst ceases; the restlessness and irritability disappear; and the child returns to its ordinary condition of health. The young physician must not expect to meet with all the phenomena we have cited as making up the regular form of measles. We have seen, as in scarlet fever and in most acute diseases, cases so very mild as to render the diagnosis very difficult, not to say impossible. In one instance, a second child in a family where a well-marked case of measles had occurred, had, twelve days afterwards, some slight languor, a hint of coryza, and six or eight faint stigmata on the face. She was put to bed for two days and then returned to her usual health. We believe this to have been a very mild attack of measles, and regard it as belonging to the same category of disorders as walking typhoid fever.

Temperature.—According to the observations of Ringer (*Reynolds's Syst. of Med.*, vol. i, art. Measles), the highest temperature reached in ordinary cases is about 103° F. From the observations of Roger (*op cit.*, p. 298) this would appear higher than is usually attained, the mean of his records having been only 101.5° F. If it rises above 102.5° it indicates a severe, if it continues below this, a mild attack. The temperature presents the diurnal variations usual in fevers, until the close of the disease, when it suddenly declines. The duration of measles, measured by the temperature, varies considerably; the decline of the fever occurring in some cases on the fourth day, in others not until the eighth or tenth day.

IRREGULARITIES OF THE DISEASE.—Under this term we shall describe only the anomalous symptoms of the disease, which occur independently of complications. Those which are produced by the latter causes will be fully treated of when we come to consider the subject of the complications.

In some cases, the symptoms of the prodromic stage are so slight that they pass almost unobserved, and the child is scarcely thought to be sick until the rash makes its appearance. In others, owing to some peculiarity of the temperament, or to the state of the constitution at the time, they are much more severe than usual, or some one symptom may be in excess. In one case that came under our own observation, in a girl seven years old, the nausea and vomiting were very distressing, and were accompanied by the most intense frontal headache. She complained precisely as children generally do with tubercular meningitis, and was, moreover, extremely restless, and at night delirious. Nevertheless, the eruption came out on

the fourth day, and was perfectly regular in its characters and course; the unpleasant symptoms ceased from that moment, and the patient recovered without any further bad symptoms. We have already spoken of five cases accompanied by general convulsions at the commencement of the first stage. The course of the disease in the subsequent stages was regular in all respects. In two other cases, in girls, sisters, seven and nine years old respectively, of highly nervous temperament, the headache in the first stage was so intense as to require the application of leeches for its relief; yet the disease was regular in its other characters.

The eruption presents various irregularities which ought to be noticed. It has already been stated that the amount of the rash varies according to the severity of the case, although in other respects regular. Sometimes the papules are comparatively small in size and few in number, and consequently, the clusters in which they are arranged have considerable spaces of healthy skin between. When this is the case, the stigmata are usually rough, lighter in color, and from this circumstances and the fact that the spaces between the clusters are large, the general tint of the skin is much less deep than in severer cases, in which the opposite of these characters prevails. In some of the mildest cases, the amount of eruption upon the extremities has been very small, and after forming, it has very rapidly, in the space of a night, faded to such a degree as to seem almost a retrocession. But as this sudden disappearance has not been accompanied or followed by dangerous symptoms, it is clear that it was dependent simply on the mildness of the attack. In such instances the general symptoms have always been slight, and the whole duration of the sickness shorter by two or three days than in severer cases. At times the order of appearance of the eruption is reversed, and the papules appear first on the trunk, thence spreading to the face.

We have already described the dull yellowish stains which remain after the papules have faded. These stains sometimes assume, in malignant cases, a livid or purplish hue, from the occurrence of passive hemorrhage into the tissue of the derm. They may, however, assume a dark and purpureous appearance, without any malignant or dangerous symptoms whatever. This happened in a family in which one of us attended seven cases of the disease in 1845. In three of them (boys of 10, 5 and 1 year old, respectively), the eruption, which was copious and regular in all, became in a single night, at the moment of decline, of a dark-brown or light-purple hue. The spots did not appear at all under pressure, and were evidently formed by true ecchymoses. The general symptoms were all favorable. The only peculiarity to be observed was that the fever had disappeared very suddenly, and that the extremities were slightly cooler than natural. The convalescence was as usual, except that the ecchymotic spots disappeared very slowly and gradually. We have, since the above-named period, seen a great many similar cases, but in none have the symptoms been attended or followed by any evil consequences.

Several authors describe a form of measles without eruption. They state that during the epidemic prevalence of the disease, some children present all the catarrhal and febrile symptoms, without the eruption, and

that they are protected against future attacks. The last assertion, at least, must be very difficult to prove. For our own part, we have never met with such cases, and should we ever seem to do so, would certainly not call them measles, lest by so doing the parents might be induced, on future occasions, to expose the child unnecessarily to the disease, when, should any evil consequences follow, they might justly question the wisdom of the physician's advice.

Willan and other authors have described another variety of the disease, to which is applied the term *rubeola sine catarrho*, or measles without catarrhal symptoms. Such cases are said to present no catarrhal symptoms whatever, and little or no febrile reaction. They are stated, moreover, to occur generally during the epidemic prevalent of measles. Most authors agree that this form does not protect the constitution against the true disease, and some regard it only as an eruption resembling measles, dependent upon gastric disorder. Our own opinion is that such cases, of which we have seen a considerable number, are nothing more than examples of roseola. The entire absence of catarrhal symptoms and of fever, of their very slight character, the short duration of the cases, and the little constitutional disturbance exhibited by the patient, all serve to convince us that they cannot be attacks of true measles. We recollect three such cases in particular, which, had they been accompanied by cough and fever, we should certainly have called measles. They all occurred in infants. The rash was preceded for two or three days by feverishness, uneasiness, restlessness during sleep, and slight diarrhoea, after which the eruption suddenly made its appearance and covered the whole integument within twenty-four hours. There were no catarrhal symptoms whatever. At the same time the febrile symptoms disappeared, and the children seemed quite well. The eruption never lasted over forty-eight hours, and disappeared without leaving a trace behind. They were, no doubt, cases of roseola.

RUBEOLA MALIGNA.—This form may occur either as an epidemic or sporadic affection. Generally, however, it prevails as an epidemic, and depends upon some peculiarity which it is impossible to understand. The few sporadic cases which are met with may be traced generally to some vicious state of the constitution of the individual, or to the unfavorable hygienic conditions in which he is placed. The symptoms assume ataxic, or adynamic characters, which give to the case the features of the typhoid type of the disease. They may make their appearance in the prodromic, or, as happens more frequently, not before the eruptive stage. When they begin in the first stage, the case is marked by great frequency and feebleness of the pulse; by prostration; by unusual dyspnoea and oppression; and especially by greater violence of the nervous symptoms, as delirium or stupor. Sometimes, even in this stage, petechiæ make their appearance, and there is lividity and soreness of the fauces, with discharges of dark blood from the nostrils, and sometimes profuse and exhausting diarrhoea or dysenteric discharges. When the time for the eruption to appear arrives, this comes out slowly and imperfectly, or irregularly, and generally assumes a livid, purplish, or blackish color, owing to the passive ex-

udation of blood into the papules, and hence the name sometimes given to such cases, of *Rubeola Nigra*, or *black measles*.

This form of the disease assumes, in fact, many of the features of purpura hæmorrhagica. The patient may die of exhaustion, of congestion of some important organ, as the brain or lungs, of the diarrhœa or dysentery which sometimes complicate the disease, or finally of the hemorrhages which occur in consequence of the dissolved and fluid state of the blood; or he may, after a severe struggle with the disease, recover his health.

COMPLICATIONS AND SEQUELÆ.—MM. Rilliet and Barthéz begin their chapter on the complications of this disease with the following excellent remarks: “*Rubeola* manifests itself by an inflammation or inflammatory fluxion upon the skin and mucous membranes. The regular course of the disease depends upon the conservation of a due equilibrium between these two kinds of fluxions. That which is seated in the skin ought in general to predominate. If the equilibrium be destroyed by any cause whatever, whether accidental or inherent to the disease, and should the predominance of the inflammation take place in the mucous membranes, there will result a phlegmasia of some one of those tissues.

“It is easy to foresee, by attention to these circumstances, that the inflammatory complications of measles will be most apt to fall upon the mucous membranes, and that broncho-pneumonia, pharyngo-laryngitis, and intestinal inflammation will be the most frequent of all.”

Bronchitis and Pneumonia.—These constitute by far the most frequent and important complications of measles. In 167 cases, MM. Rilliet and Barthéz met with 24 cases of bronchitis, 7 of pneumonia without bronchitis, and 58 of lobular broncho-pneumonia. This statement shows how very large a proportion of the cases of measles occurring in the Children’s Hospital at Paris, became complicated in the course of the attack. The proportion in private practice is much smaller, since in 314 cases, we have met with only 24 of bronchitis, and 6 of lobar pneumonia. These are, however, in private practice, according to our experience, much the most important of the complications likely to occur. Of six deaths which occurred in the 314 cases that we have seen, 3 were caused by bronchitis.

The time at which these different complications make their appearance is important. They may occur during the initial stage, early in the eruptive stage, during the decline of the eruption, or after the eruption. The most common period for their occurrence is the initial stage. It is difficult or impossible to ascertain their causes in a great many cases. In some instances they depend evidently upon cold. Age has some influence upon their production, as bronchitis is most apt to occur in young children, whilst lobar pneumonia attacks those who are older.

The *physical signs* of these affections are the same as when they exist in the idiopathic form. The rational signs are increase of cough, which instead of being merely laryngeal, becomes deeper and either pneumonic or catarrhal; and dyspnœa, which is sometimes excessive, the number of respirations mounting to 40, 50, and, in severe cases, to 60 and 80. The pulse is more frequent than in regular measles, and in very bad cases becomes rapid and small; the skin is hot and dry; the face is pale and

anxious in severe cases, in which the eruption does not appear; and the child is generally restless and irritable, with broken irregular sleep, or, in the most violent cases, it is dull and soporose. In two of the fatal cases that came under our observation, convulsions occurred. It should be remarked, however, that in one, the patient, a boy only nine months old, was laboring under an attack of whooping-cough, and that it was in one of the paroxysms of that malady that death took place. In the other case, that of a boy eighteen months old, the convulsions occurred first on the day of eruption, and then ceased, to recur again the third day afterwards. The bronchitis dated from before the appearance of the eruption, and was no doubt the cause of the convulsions and death.

When a pulmonary complication begins in the prodromic stage, it almost always modifies the eruption in some manner, either retarding or rendering it irregular or imperfect. When it dates from the second stage it may cause a partial or complete retrocession of the eruption. We have known the eruption to be retarded several days, so as not to come out until the fifth, sixth, or even ninth. When the rash does appear, whether at the usual period or later, it is evidently with difficulty. It is pale and scanty, or abundant on one part of the body, and scanty on another, or it appears and disappears alternately. At length it either comes out fully, and the threatening symptoms pass away, or the eruption lasts the usual, or nearly the usual length of time, in its pale and imperfect condition, and the child recovers slowly and gradually from the complication, which has become the most important part of the sickness; or, in fatal cases, the symptoms grow worse and worse, and the child dies after a few days, or a longer time, according as the inflammation assumes the acute or chronic type.

Whenever it is observed in a case of measles, that there is more drowsiness or irritability than usual, or that the pulse is more frequent or stronger than it ought to be, it becomes important to ascertain carefully the state of the respiration. If this be accelerated, the thorax ought to be examined with strict attention, by auscultation and percussion, to discover whether there be not some pulmonic inflammation at work, likely to convert the disorder from a mild one, as it almost always is when uncomplicated, into one dangerous to life, which it will assuredly become should any pulmonic complication be allowed to steal unawares upon the patient.

The prognosis of the pulmonic complications of measles would appear to be very unfavorable in hospitals for children, since Rilliet and Barthez state that scarcely one patient in four or five recovered. Of the 30 cases that we have seen, we have already stated that 3 died of bronchitis, and if we recollect that one of these was complicated also with pertussis and morbid dentition, it will be seen that the prognosis is, as might be expected, vastly more favorable in private than in hospital practice.

There is, however, a tendency, especially marked in delicate, strumous children, for the inflammation of the bronchial mucous membrane to become chronic, in which case the cough may persist for years, at times intermitting, but returning after the slightest exposure, and particularly in cold, damp seasons of the year.

Laryngitis is a common complication of the disease. The authors just

quoted met with it in 35 of their 167 cases. It occurred in 8 of the 314 cases that came under our observation. It is often accompanied by pharyngitis.

Autopsies show that the laryngitis may be slight, severe, or accompanied with pseudo-membranous exudation. The inflammation may be simple, consisting merely of different degrees of redness, or of redness with thickening and softening of the mucous membrane; it may be more intense and accompanied by ulcerations or erosions; or, lastly, it may be associated with an exudation of false membrane.

The symptoms of this complication will depend upon the form the inflammation assumes. It is unnecessary to describe them here, as they are the same as those of the idiopathic affection, which has already been fully treated of.

The occurrence of laryngitis exerts but little influence on the rash, particularly as it almost always appears during the decline of the latter. It is seldom fatal, unless it assumes the pseudo-membranous form. The eight cases that came under our observation were attacks of the simple disease, and they all recovered.

Inflammation of the Intestines.—According to Rilliet and Barthez, lesions of the intestinal mucous membrane are the most frequent complications, after pulmonary affections. About a third of their cases presented at the autopsy erythematous inflammation of the mucous membrane; a fifth offered follicular entero-colitis, a seventh ulcerative inflammation, and a fourth softening. Some presented several of the lesions united, and in a few no lesion was found, though the symptoms of entero-colitis had existed during life. We give these data from the above authors, not because they apply to private practice, but merely in order to show what are the tendencies to the disease, when disposed from unfavorable hygienic conditions to take on complications. We have met with only seven instances of intestinal inflammation in the 314 cases that have come under our own observation. Four of these occurred in the same family, in children of seven, five, three, and one year old respectively. They were cases of entero-colitis, accompanied in two with dysenteric symptoms, and all made their appearance towards the close of the disease. The three remaining cases were attacks of dysentery, one of which was very severe, the stools amounting to twenty in the day, while the other two were much less so.

The intestinal complications may appear during the initial stage, or on the day of eruption, but if not at one of these periods, they are most apt to occur during the decline of the rash. The slight cases, constituting the common diarrhœa of the disease, generally begin early, whilst the grave cases usually date from a later period of the disease. The *causes* of these complications seem to be various exciting agents acting upon a mucous membrane predisposed, by the nature of the disease, to inflammatory action. These agents are said to be, generally, improper food, giving rise to indigestions; and the too early use of purgative remedies, and laxatives. In the cases observed by ourselves it was impossible to detect the causes.

The *symptoms* are more or less abundant diarrhœa, and in some, but not all the cases, tenderness with tumidity and tension of the abdomen. This

complication does not exert much influence upon the measles, which usually pursue their regular course. Sometimes, however, it occasions an aggravation of the febrile symptoms, and, when of a grave character, may no doubt interfere with the regular progress of the eruptive disease.

According to Rilliet and Barthez, this complication was very seldom the only, or even chief cause of a fatal termination. Scarcely five or six of all that they observed could be considered as of that kind. It increases very much, however, the danger of the pulmonic attacks, for the latter are much less serious, so long as they exist alone, while so soon as intestinal inflammation is added to them, they become almost necessarily fatal. The seven cases that we met with recovered under simple treatment.

In a considerable number of cases, a slight diarrhœa, to which we have already referred as a common event in measles, occurred, but only in the seven above mentioned did it amount to a serious complication.

In one case that came under our observation, in a girl between five and six years old, fatal cerebral symptoms, due either to congestion of the brain or uræmia, occurred just as the rash was disappearing. There was no evident cause whatever for this accident. There had been no imprudence either as to diet or exposure. The child was, however, of a tubercular family, the mother having at this very time tubercular disease of the lungs. The eruption had come out well and properly, and continued to do so on the second day without any irregular or threatening symptoms. On the third day of the eruption this began to decline, and the child had an attack of spontaneous vomiting, but continued through the day cheerful and pleasant. On the night of that day she was restless and feverish, and wanted much drink. On the fourth day she was drowsy and heavy, and complained of her head. We saw her first in the evening of this day. She was then very dull and heavy, scarcely answering questions, and protruding the tongue slowly and after much urging. She had some little, but not a troublesome cough. Careful examination revealed no disease of the thoracic organs. The respiration was natural, and the pulse full and very frequent. On the morning of the fifth day the patient was comatose, neither answering questions nor protruding her tongue. In the course of the day there were some irregular convulsive movements. In the evening the right arm was rigidly flexed at the elbow, and the left one stiffly extended. The patient died that night. No autopsy could be obtained.

In another case death occurred from sudden *effusion of serum* into the internal cavities, caused apparently by the existence of an excessively hydræmic state of the blood, possibly connected with albuminuria, which had been allowed to come on gradually, without any attempt on the part of the parents to seek a remedy during the slow approach of this condition of the circulating fluid.

CASE.—The patient was a boy in the second year of his age, who had a phthisical mother. The attack of measles took place in the last week of January, 1852, and was regular, and not, according to the account of the parents, we not having seen the child, at all severe or dangerous in any respect. After the attack was over, however, and though he was running about the house as before, he continued to look more and more pale and sickly until the evening of February 25th, when suddenly after 11 P. M. he

was seized with fever, and became very restless. On the following day, at 9 A. M., we saw him. He was then extremely pallid, and very drowsy and heavy; the breathing was rapid and oppressed, the pulse very frequent, and the skin hot and dry. He was evidently dropsical, as both the face and hands, and the feet also, were puffed, smooth, and doughy. The bowels had not been opened the previous night. In the evening the pulse was 170; the skin was still hot, and the breathing very rapid and much oppressed. There was scarcely any cough. The percussion was dull over too large a space in the præcordial region; the cardiac impulse was obscure, and the sounds indistinct and muffled; there was no bellows-murmur. The percussion was dull over the inferior dorsal regions. No râle whatever was heard. The child died on the following morning at 3½ o'clock. Ten minutes before his death he asked for a drink, lifted himself up in bed, drank freely, looked around intelligently, and then lay down and died. At the autopsy the subcutaneous cellular tissue was found to be infiltrated with serum. On puncturing the right pleural sac, there was an immediate escape of a clear, straw-yellow serum. There was considerable effusion in the left pleura also, but less than in the right. The pericardium contained at least two ounces of serum, so that it was pushed off to a considerable extent from the heart. There was a slight pleuritic adhesion of the upper lobe of the right lung to the ribs. This was, however, evidently of an ancient date. There was no other inflammation of the pleuræ, and none of the pericardium. Both lungs contained tubercles, which were not very numerous, but in the upper lobes of considerable size. There was no pneumonia, but both lungs were somewhat congested. The heart was larger than usual. In the right auricle there was a rather large, and white, but soft concretion, and a smaller one in the right ventricle. The left cavities presented no concretions. The valves were healthy.

There are several other disorders which sometimes complicate or follow measles, but as we have already given as much space to this subject as the limits of the work will allow, we shall be satisfied with a simple enumeration of them. They are otitis, ophthalmia, hemorrhages, gangrene of the cheek or vulva, anasarca, and different cerebral symptoms. We will merely add that measles appear to possess a special tendency to develop tubercular disease in the system, and that it is necessary, therefore, to treat a child showing any predisposition to that diathesis, or one born of tubercular parents, with particular caution, both at the time of the disease and during the convalescence. It is not uncommon for measles to be conjoined with other eruptive diseases. We have known it to coëxist with scarlatina in two instances, and Dr. G. B. Wood has met with a fatal case of the same nature. It may be associated likewise with variola or with erysipelas; of the latter we have met with one instance. We will mention here that of the whole 314 cases of measles that we have observed, 257 were simple and 57 complicated. The complications were as follows: bronchitis, 24; pneumonia, 6; laryngitis, slight or severe, 8; dysentery, 7; pertussis, 7; scarlatina, 2; convulsions in the early stage of the disease, 5, and in the latter stage, 3; keratitis, 2; intermitten fever, 1; erysipelas, 1; meningitis, 1; congestion of the brain, 1; serous effusion into the internal cavities, 1. It ought to be observed, however, that in the above enumeration several cases are referred to twice, and one, a case in which pertussis, bronchitis, and convulsions occurred, three times.

ANATOMICAL LESIONS.—It is difficult to ascertain what are the characteristic lesions of measles, because of the fact that most of the fatal cases prove so in consequence of some complication. Some few fatal

cases, however, of the regular form and some in which the complication was so slight as not to be likely to change the morbid appearances much, have led to the following conclusions.

The lesions present in measles are the following: general congestion of different organs, which are colored red from the imbibition of blood and sometimes softened. The congestion affects the mucous membranes particularly, and imparts to them a reddish or slightly blackish color. In some of the cases there is morbid development of the intestinal follicles. The most important lesion, however, is that of the blood, which presents the appearances common to the class of pyrexia. These are a normal proportion or diminution of the fibrinous, with increase of the globular elements of the blood. Dr. Copland (*Dict. Prac. Med.*, vol. ii, p. 819) gives the appearance in a few fatal cases of malignant measles. They were, softening of the tissues and the facility with which they were torn; the presence, in some of the cases, of a turbid or sanguineous serous fluid in the serous cavities; general congestion of the lungs; dark appearance, and livid or purple ecchymoses of the bronchial mucous surface, of the fauces, stomach, and cæcum; engorgement with dark and semifluid blood of the veins and sinuses of the brain, and of the auricles and large veins; and finally a livid and mottled appearance of some parts of the body, with petechia of a dark color.

DIAGNOSIS.—It is impossible to diagnosticate measles in the first stage with any considerable certainty. The existence of the disease may be suspected in that period from the appearance of the eyes, from the coryza and sneezing, the frequent, hoarse, scraping cough, and the fever, headache, and thirst. If, in connection with these symptoms, it happens that an epidemic of measles be prevailing at the time, or that the child has been exposed to the contagion of the disease, the inference becomes still more plausible. Nevertheless, any opinion upon this point ought to be given with much reservation.

We have already alluded to the opinion of some authorities, that the diagnosis in the early stage is aided by the presence of punctated redness of the roof of the mouth, or of a red papule on the velum palati. We have met with this symptom so often that we have formed the habit of looking for it in doubtful cases. It is often present twenty hours before the cutaneous rash appears. When, therefore, this punctated eruption on the hard and soft palate is discovered, in a child in whom lachrymation and catarrh of the upper air-passages, with fever, suggest the probable approach of measles, the probability that this case is one of that disease is very much augmented, though no cutaneous rash whatever may yet be visible.

After the eruption has come out fully, it is not likely to be mistaken for any other disease, unless it be roseola or rōtheln, the rash of both of which sometimes resembles that of measles very closely. It may be distinguished, however, from the former by attention to the concomitant symptoms, by the entire absence or very slight degree of fever, the more rapid evolution of the rash, and the absence of the peculiar catarrhal symptoms. From rōtheln it is not so easily distinguished. We have known rōtheln to be

pronounced true measles by experienced men on several occasions. The diagnosis may be made, we think, by the shorter prodromes, the much less marked laryngeal catarrh, even when the ophthalmic symptoms are quite decided, by the more rapid appearance and darker tint of the eruption, by the very moderate degree of fever, and particularly by the presence in most cases of r  theln, of slight enlargement of one or several of the cervical lymphatic glands, and specially of those on the back of the neck.

In the very early stage of the eruption, measles may be confounded with variola. A careful attention, however, to the size and shape of the papules, which in measles are much larger, flatter, less elevated, softer, and without the shotty feel peculiar to the papules in variola, and the presence of the catarrhal symptoms, will usually suffice to distinguish them, even in the earliest stage. In measles also the general symptoms persist, or even become aggravated after the appearance of the eruption, instead of abruptly subsiding as they do in variola. A little later, the appearance of vesicles on some of the papules about the face in variola, will show the difference still more strongly. The distinction between measles and scarlatina has already been drawn in the description of the latter disease. It rests chiefly on the much shorter duration of the prodromic stage, the greater violence of the anginose symptoms, the absence of the peculiar catarrhal symptoms, and the more rapid evolution of the eruption in scarlet fever; and lastly, on the differences in the two eruptions, observable especially at their first appearance.

The eruption of typhus fever appears nearly at the same time as that of measles, and in their earliest stage the two eruptions often resemble each other closely. In typhus, however, there is an entire absence of the characteristic catarrhal symptoms. The spots are less elevated; are isolated and round, instead of coalescing to form crescentic patches; do not appear first on the face, but on the trunk or wrists (Ringer); more frequently become petechial, and last a much longer time.

When measles are conjoined with some other eruption, the diagnosis is to be made out by a careful study of the initial symptoms, and of the eruption on different parts of the body, for we can generally find well-marked patches of the rash peculiar to each on some portions of the surface. In one of the cases of measles and scarlatina that we saw, the latter disease was developed first. The eruption made its appearance in the usual form; on the second day of the eruption, the child was seized with hard, hoarse, laryngeal cough, and with redness of the eyes and lachrymation. These symptoms continued three days, at the end of which time the scarlatinous rash had disappeared from the face, but remained visible upon the trunk and extremities. Characteristic measles papules now made their appearance on the face, and pursued their regular course, while on the trunk and extremities the measles eruption was never well defined, being mixed with and disguised, as it were, by that of the scarlatina. In the other case, the measles appeared first and went on regularly until the eruption was declining and the general symptoms moderating, when suddenly the fever, heat of skin, restlessness, and irritability returned,

and the child was very soon covered with the punctated scarlet rash of scarlatina.

PROGNOSIS.—The prognosis of simple, uncomplicated measles is very favorable; the cases almost always recover without difficulty. This is shown to be true by the following facts; Rilliet and Barthez report 36 cases of simple measles, of which all but one recovered. Of 257 cases that we have seen, all terminated favorably. When, on the contrary, complications occur, the disease always becomes more or less dangerous, the degree of danger depending on the nature of the intercurrent affection, and on the hygienic conditions in which the patient is placed. Thus of 131 cases observed by the above authors, in which some form of complication occurred, 89 or about two-thirds proved fatal, while of the 53 complicated cases that we have seen, only 6 were fatal. It must be recollected that the cases of the French observers all occurred under the unfavorable hygienic conditions of a large hospital, in children of bad constitution from congenital or acquired causes, whilst ours were observed in private practice, where the hygienic conditions are favorable in the same degree as they are unfavorable in hospitals.

The six fatal cases that came under our observation, proved so from the circumstances we are about to mention. The first occurred in a child nine months old, who was laboring under pertussis when attacked with measles. Bronchitis supervened upon the measles, and proved fatal by convulsions, which came on during a paroxysm of whooping-cough, two weeks after the disappearance of the rubeola. The second case was that of a boy, eighteen months old, who was prescribed for by an apothecary from behind his counter, until we saw him. The eruption made its appearance imperfectly, we were told, and with a convulsion. After this he was very restless, and had rapid and difficult respiration and much cough. On the morning of the fourth day of the eruption, this had almost entirely disappeared, and the child was again attacked with convulsions. We saw him shortly after this for the first time, and found him comatose, with convulsive movements of the limbs, extreme dyspnoea, and all the symptoms of extensive bronchitis of both lungs. He died thirty-six hours from this, as was to be expected. The third was a case of pneumonia in a child between one and two years of age, in which the inflammation came on as the eruption was fading, and proved fatal, in spite of all that could be done, on the eleventh day. The fourth occurred in a boy between four and five years old, who appeared to recover perfectly from the measles, but was attacked in ten days with meningitis, and died. The fifth was the case of congestion of the brain, already detailed in the remarks upon complications, as proving fatal shortly after the decline of the rash. The sixth was that of sudden dropsical effusion into the internal cavities, also described in the remarks upon complications.

To conclude, we may state that the prognosis is always highly favorable under the following circumstances: when the disease is primary; when the initial stage is of the proper duration; when the eruption begins upon the face and extends gradually to the rest of the body; when the febrile movement is moderate; when the eruption, after increasing for one, two,

or three days, gradually decreases; when the cough and other concomitant symptoms diminish with the fever; when the cutaneous surface, after the fading of the rash, assumes a natural color, and is neither flushed nor pale; when the appetite returns, the disposition to be amused and take notice continues, and lastly when the sleep is natural.

On the contrary, the prognosis becomes unfavorable under the following circumstances: when the initial stage lasts longer than usual, and when it is accompanied by violent symptoms of any kind, as extreme jactitation, irritability, dyspnœa, much stupor, coma, or convulsions; when the eruption is irregular in its appearance or course; when the fever does not disappear with the eruption, whether it remains violent or assumes the form of hectic; when, after the eruption, the face continues deeply flushed or becomes very pale; when the cough, dyspnœa, or diarrhœa persist: and lastly, when the child remains weak, languid, dispirited, or irritable.

It may be stated, in conclusion, that the prognosis of measles is always favorable in proportion to the health of the child at the time of the invasion, and the regularity with which the disease passes through its different phases; while it becomes unfavorable, though far less so in private practice amongst people in easy circumstances, than in hospitals, or amongst the poor and wretched, whenever it attacks a child already laboring under some disease, and when it becomes complicated with any other malady, either local or general.

TREATMENT OF THE REGULAR, SIMPLE FORM.—This form requires, in a large majority of the cases, little other treatment than strict attention to the hygienic condition of the patient, the use of simple diaphoretics, of a simple laxative when there is positive constipation, and the palliation of any of the symptoms that may chance to become somewhat more troublesome than usual.

The child ought to be confined to bed in a large, well-ventilated chamber, the light in which should be somewhat softened. Every precaution should be observed to prevent chilling of the body, while at the same time it is nearly, if not quite as important, to avoid overheating the patient, either by excessive clothing, or by keeping the temperature of the room too high. In winter it is well to direct the temperature to be maintained at between 65° and 70° F., night and day. If this be done, the child is not apt to take cold, even though it be uncovered at times, and yet the warmth is not oppressive. We have often remarked that this temperature is just what it ought to be when the room is well ventilated, either by means of an open fire place, or by communication with adjoining rooms; but when, on the contrary, the room is heated by a furnace-flue, and not ventilated at all, or very imperfectly, the same temperature, as indicated by the thermometer, becomes close and oppressive. Under such circumstances, a door into an adjoining room, or if this cannot be, one into the entry, ought to be kept more or less open, with a screen of some kind between it and the child, in order to secure a good ventilation, which is assuredly of the very highest importance, and yet to prevent by the screen a current of cool air from chilling the patient. Miss Florence Nightingale

remarks that doors are made to be shut and windows to open. There is much in this saying, and when the nurse is intelligent and observant, we much prefer to shut the door and open a window. In our winter temperatures in this city this must be done very carefully. One of the sashes raised an inch, or one or two inches, will make a large difference in the temperature and vitality of the air of the sick-room.

The *diet* during the febrile period must be light. It may consist of milk and water, of arrowroot, sago, or tapioca, prepared with milk or water; or of crackers soaked in water, with salt, or some similar food. When the eruption and fever have in great measure disappeared, some light broth, either vegetable or animal, with dry toast or bread, plain boiled rice, or a roasted potato, may be added; and after all the symptoms have ceased, the usual diet can be gradually resumed. The drinks may consist of simple water, of lemonade, orangeade, gum-water, or flaxseed tea, with the addition of a little sweet nitre; or of weak infusions of balm, sweet marjoram, or saffron, or cascarilla with a few drops of hydrochloric or nitric acids. They may be given in any reasonable quantity, at the temperature of the room.

Cool or cold water is the best drink the patient can have, and he should have as much of it during the febrile period as he desires. It is a mistake to allow very large draughts of cold water to be taken at once. We saw a boy, nine years old, attacked with violent epigastric pain and partial retrocession of the eruption directly after swallowing suddenly a half pint of iced water. The unpleasant symptoms passed off in a few hours, and he had no difficulty afterwards. But we have never known anything but good come of the use of cool, and even of iced water, in frequent, *small* amounts so as to satisfy the sense of thirst.

The patient should not be permitted to leave the room until a few days after the entire disappearance of the disease. This precaution is necessary for all, but particularly for the delicate, and in the cold weather of these latitudes. He should be kept in the house until he has regained in some degree his usual health, and then sent out with due precautions.

Medical Treatment.—Many cases of measles—the mild, the moderate, the uncomplicated—need no other treatment than that just laid down in the paragraph on the hygiene of the disease. So long as the case goes on regularly, so long as the symptoms are moderate and such as to cause but little suffering, there is no necessity for drugs, or, at the most, a simple diaphoretic, as sweet spirit of nitre or the solution of acetate of ammonia, with a little paregoric or laudanum once or twice in the evening, will be all that ought to be given.

The child does not require, and therefore ought not be made to take as a mere routine, *cathartics*. If the bowels are known to be costive, and not to have been moved for two or three days, a teaspoonful of dessertspoonful of castor oil, or, better still, a dessertspoonful to a tablespoonful of simple syrup of rhubarb, or a simple enema, will answer every purpose. We are sure that active purging is unnecessary, and apt to do harm.

When the case is a very decided one, and the eruption extensive and deep in color, the fever runs high, and the patient often suffers greatly

from fever-pains, and from the violence and frequency of the cough. Here medical treatment is necessary, since it lessens suffering, diminishes the violence of the symptoms, and so promotes the safety of the patient. In infants, under these conditions, we order five drops of sweet spirit of nitre, two or three of syrup of ipecacuanha, and two of paregoric, in a teaspoonful of sweetened water every two hours, at the age of six months. At one and two years, we double the proportions of the active ingredients. Should even these small doses of ipecacuanha cause any sickness of stomach, we lay that drug aside. One of the best combinations is the following:

R. Potass. Citrat.,	3j.
Spt. Etheris Nit.,	f 3 ij.
Tr. Opii Deodorat.,	℥xij vel xxiv.
Syr. Simp.,	f 3 vj.
Aquæ,	f 3 ij.—M.

Dose.—A teaspoonful every two or three hours, at five years of age.

In younger children, from two to five years, the same formula may be used, except that the laudanum should be reduced to six minims. When the cough is very dry, scraping, and, as it sometimes is, incessant, there should be added to the above mixture syrup of ipecacuanha, in the proportion of five or ten drops to every teaspoonful, according to the age of the child; and there may and ought to be given from time to time, if the patient be not too drowsy from the effects of the fever or the mixture, an extra dose of opium. We prefer on the whole the deodorized laudanum. Of this two drops in a teaspoonful of water may be given two or three times a day, or, better still, once or twice in the evening, to children over five years of age. From one to five years of age, one or two drops are enough. In some few children paregoric may answer better, but this rarely happens. When this is used, ten to twenty drops at five years, five or ten at one year, and from half a teaspoonful to a teaspoonful over five years, may be used instead of the laudanum.

Depletion, except that which comes of the above treatment, is unnecessary. We did, in past years, use depletion in 2 cases out of 257 regular cases of which we kept notes. In one, a venesection to four ounces was used in a boy seven years old, on account of the great violence of the febrile movement; and in the second, leeches were applied to the temples for an intense headache in a girl nine years old. For many years past we have used no general bleeding, but might be tempted to use leeches in a case of the same kind as that just mentioned, in which the pain in the head was something quite out of the usual way. Instead of venesection we should make use of a warm bath continued for fifteen to twenty minutes. If the temperature of the body be very high, it may be reduced by careful sponging with tepid or cool water.

Sometimes, when the cough is very troublesome, a mustard foot-bath used every three or four hours, and a mild liniment, as one composed of sweet oil and spirit of hartshorn, or of chloroform, camphor, and soap liniment, rubbed gently upon the front of the neck and over the upper part of the sternum, will assist materially in palliating this symptom.

When the conjunctival inflammation is acute and painful, it may be relieved by lotions with simple warm water, milk and water, or sassafras-pith mucilage, alone or mixed with rose-water. If the headache be very violent it can generally be relieved by the use of a laxative, by the occasional use of a mustard foot-bath, or of a sinapism to the nucha, and by the application of cold to the head.

If, at any time during the course of the case, symptoms of exhaustion appear, the most nourishing and concentrated food, with alcoholic stimulants in graduated doses, should be promptly resorted to.

The *malignant form* of the disease must be treated chiefly with stimulants and tonics. The most useful are wine and brandy, quinia, ammonia, capsicum, etc. Camphor and opium would be proper, were the case attended with severe nervous symptoms. The diet ought to be nutritious and digestible, and may consist of milk and bread, light broths, and beef-tea or essence of beef.

When local inflammations occur, they may be treated by a few dry cups, or by means of counter-irritants, of which the most suitable are mustard, spirit of turpentine, or ammonia. Blisters ought to be avoided, as they are very apt to occasion dangerous and even fatal sloughing.

TREATMENT OF THE COMPLICATIONS.—*Bronchitis, Pneumonia.*—The mode of treatment of these complications must depend upon the stage at which they are developed, and upon the age and constitution of the subject. When they occur during the first stage, one of the most important points in the treatment is to endeavor to favor the appearance of the eruption, and when in the second stage, and the eruption has retroceded wholly or in part, the same indication applies with equal force. When they appear during the third stage, they are to be treated without any regard to the eruption, but always with reference to the fact that the patient has just passed through an acute febrile disease, which must have weakened in some degree the vital powers.

It may be stated in general terms, that the treatment proper for these local inflammations when they occur as primary affections, is proper also, with some reservations, under the circumstances we are now considering.

Thus even local depletion should be employed only with the greatest care, and, indeed, we should recommend in preference the application of dry cups, or of sinapisms.

Purgatives should also be used with caution, on account of the disposition to gastro-intestinal irritation which is always present in this disease. Our own practice is to employ moderate counter-irritation, in conjunction with minute doses of sulphuretted antimony and Dover's powder, or a mixture containing citrate of potash and syrup of ipecacuanha. When in these cases the skin is at all coolish, or bathed with too considerable a perspiration, we have found the liquor ammoniæ acetatis a very useful remedy.

It is universally acknowledged that it is exceedingly important to assist nature in throwing out the rash, whenever these complications either prevent its formation, or cause its retrocession. The true mode of doing this is to cure or alleviate the internal inflammation, which is the cause of the

difficulty. To attain this end the above plan of treatment ought to be instituted at once. At the same time, we may greatly assist the appearance of the eruption by a persevering employment of counter-irritants. The best of these is, we believe, mustard, and in some cases a warm bath. The mustard may be used in the form of plasters, poultices, or baths. Our own plan in moderately severe cases, is to apply a mustard poultice to the interscapular space, and to make use of a mustard foot-bath, two or three times a day, while in severe and urgent attacks we direct the cataplasm and bath to be renewed every two or three hours, taking care, however, to apply the former alternately to the front and back of the chest, in order to avoid all possibility of too violent an action upon the skin; the feet and limbs also ought to be carefully watched, to avoid the same danger. We have had occasion to observe the great efficacy of this unremitting employment of revulsives, in several severe cases of bronchitis in young children. In some we have depended solely upon this treatment, and the use of small doses of ipecacuanha and spiritus Mindereri. In one particularly, which occurred in a child eight months old, the attack came on in the first stage. On the fourth, fifth, and sixth days, the dyspnœa was excessive, the respiration running up to 70 and 80; the pulse was frequent and small; the skin pale and rather cool; and the irritability and restlessness very great. For a period of twenty-four hours, we used the poultices and foot-baths every two hours regularly, and gave internally the spiritus Mindereri at the same intervals. Nothing else was done. On the sixth day, when one of the poultices was removed from the interscapular space, the integument beneath was observed to be covered with the measly stigma, whilst there were none as yet on any other part of the surface. From this time the eruption came out freely, and the child recovered rapidly.

The warm bath may be used under the same circumstances. It should be given with great care, the child being wrapped in a warm blanket the moment it is removed from the water, to prevent the least sensation of chilliness. It may either lie for a short time in the blanket, or be wiped dry beneath it, and then dressed.

In some of the cases of bronchitis, there has been profuse secretion attended with extensive subcrepitant and mucous râles. In such instances we have found the internal use of the syrup or infusion of polygala seneka, with an occasional revulsive, very effectual.

The *diarrhœa* which occurs so frequently seldom requires any treatment. Indeed, unless it indicates evident entero-colitis, or is accompanied by frequent mucous or bloody stools, and by pain and tenesmus, it is better not to interfere with it beyond paying strict attention to the diet. When attended, however, with the symptoms just mentioned, it must be treated by astringents, by opium and ipecacuanha, and by the application of poultices to the abdomen. The seven cases that occurred to ourselves recovered under the use of laudanum enemata, given twice or three times a day, the strictest diet, and small doses of Dover's powder.

Laryngitis, as it occurs in most of the cases, needs but little treatment beyond careful avoidance of cold, the use of some mild nauseant, and re-

vulsives to the neck. It is very seldom of a dangerous character. When, however, it assumes the character of pseudo-membranous croup, it must be treated with all activity, in the manner described in the article on that disease. In only two of the eight cases, we have seen, did it appear at all threatening, and both of these recovered under the use of emetics and moderate leeching of the throat.

The *cerebral symptoms* which sometimes occur, must be treated differently in different periods of the disease. In the early stage, when they last but a short time and do not recur, they require nothing more than a warm bath and the use of revulsives. If they continue to recur, or are followed by stupor or other cerebral symptoms, more energetic treatment becomes necessary. If the child is strong and hearty we may apply dry cups to the back of the neck or temples, and resort to purgatives, revulsives, and cold applications to the head. When the symptoms are violent, and when the heat is intense, it has been proposed to use cold lotions in the manner recommended in scarlatina. The evidence upon this point is not very conclusive, and as we have never used them, nor seen them used, nor indeed seen any necessity for a resort to them, we can offer no opinion in regard to their value.

We have met with five cases of convulsions in the first stage. One occurred in a boy five years old; the convulsions were slight, lasted not more than ten or fifteen minutes, and were followed by no bad symptoms. The intelligence of the child returned very soon afterwards. The only remedy used was a warm bath. The other cases have already been described.

When convulsions occur in the second or third stages, it is very important to ascertain whether they are not the result of some local disease. Two of the three cases that came under our notice accompanied violent attacks of bronchitis. The third was caused by congestion of the brain. Here the treatment must be directed against the local disease, if that can be detected. When, on the contrary, the convulsions seem to depend on nervous irritation, they may be treated with baths, revulsives, purgatives, and the careful administration of opium, as recommended by Sydenham, Copland, Rilliet and Barthez, and other authors; or of bromide of potassium, chloral, camphor, assafœtida, musk, or hyoscyamus. If accompanied by intense heat and great dryness of the skin, without local complications, cold or tepid lotions may also be tried.

The treatment suitable when any of the complications or sequelaë become chronic, will be found in the articles devoted to the respective diseases. Bearing in mind the tendency to the development of scrofula or tuberculosis after this disease, the most careful attention should be paid to all hygienic measures; and alteratives and tonics, as syr. ferri iodidi, cod-liver oil, and quinia, should be administered.

ARTICLE VII.

RÖTHELN.

DEFINITION ; SYNONYMS ; FREQUENCY.—Rötheln is a contagious eruptive disease of benign nature, occurring epidemically, and bearing a very close resemblance to mild forms of measles and scarlatina. The eruption is seldom preceded by any marked premonitory symptoms, but appears suddenly, lasts generally about four days, and usually fades away without any desquamation. This describes the ordinary mild form of the disease, but we have frequently observed febrile symptoms of twenty-four hours' duration to precede the eruption, and a marked furfuraceous desquamation to follow.

It has frequently been designated German measles; and it is probable that many cases described as rubeola sine catarrhis have been really cases of rötheln.

It affects all ages, but especially childhood. According to the observations of J. Lewis Smith in a series of 96 cases, studied by him in two epidemics in New York in 1873 and 1880, it occurred in 33 under five years of age, in 41 between five and ten years, and in 22 between ten and forty-two years.

It is probable that limited epidemics of rötheln have been of frequent occurrence in this country without being recognized. As far back as 1866, we had the opportunity of observing closely more than fifty cases that occurred in the practice of the late Dr. George Pepper, then one of the district physicians of the Philadelphia Dispensary. The disease was limited to a small area of one of the poorest quarters of the city, and the cases which occurred almost exclusively in young children were quite severe, though all terminated favorably. From that year until 1880 we met with rötheln but rarely; but in the latter year a very widespread epidemic prevailed in various parts of America, and we saw in this city a large number of cases of a mild type in children and adolescents. As these pages are passing through the press, we have again met with a few scattered cases. Dr. Honan, Sr., in 1845, reported some cases occurring in Boston, and Drs. Cotting and Howard, in 1853 and 1871, also described several cases seen by them (*Boston Med. and Surg. Journal*, March 15th, 1878).

J. Lewis Smith has given an account of an epidemic of it in New York in 1873-74 (*Sanitarium*, July, 1874), and Dr. Forrest of one in Charleston, S. C., in 1880 (*Amer. Journ. Med. Sci.*, April, 1881.) A number of cases occurred in Philadelphia in 1875, and Drs. Duhring and Hays have described (*Philada. Med. Times*, March 26th, 1881) numerous cases seen by them in 1881.

NATURE.—Rötheln presents points of marked resemblance to both measles and scarlatina, and in certain cases the symptoms may be so closely like those of either of these latter affections as to render a differential diagnosis very difficult. It is not strange, therefore, that considerable discussion should have taken place as to its real nature, some asserting that it is a mild form of measles, while others have held that it is a poorly developed

scarlet fever, It seems to us however, that the weight of evidence indorses so strongly the view that rōtheln is an independent and distinct disease as to leave no room for further discussion. Apart from the peculiarities of its symptoms, the following considerations establish its essential distinction from both measles and scarlatina: An attack of rōtheln affords no immunity from either of the latter diseases, nor do they afford any protection against it; it occurs epidemically at times when neither measles nor scarlatina are prevalent, and all of the cases exhibit such distinctive characters as to prove conclusively that they represent an independent zymotic disease.

SYMPTOMS; CAUSE; DURATION.—The eruption may appear suddenly in the midst of apparent health, constituting the first evidence of the disease. But in a number of cases, especially in older children, and adolescents, we have noted the occurrence for twenty-four or even thirty-six hours before the appearance of the eruption, of feverishness, headache, and pain in the back and limbs, nausea, and, less frequently, soreness of the throat with short, dry, hacking cough.

The eruption appears in the form of an erythema, and, as well described by Duhring (*loc. cit.*), is "multiform, more or less confluent, disseminated, ill-defined, pale-red or rosy, punctate, and small split-pea sized, faintly defined macules." It occupies the face, neck, chest, and back, and sometimes the arms and thighs. It resembles the eruption of measles, but it is macular, not papular, less distinct, the spots are round, not oval, and there is no tendency to a concentric arrangement. The color of the eruption occasionally looks like that of scarlet fever, but it is much paler than that met with in that disease. Moreover, in discriminating, reliance is to be placed upon the non-prevalence of scarlet fever and the mildness of the constitutional symptoms. The eruption usually lasts four days and disappears rather quickly. In some cases no desquamation occurs, but in our own experience a fine branny or furfuraceous exfoliation has been usual.

It is generally accompanied by a slight suffusion or even a fine injection of the eyes, by mild coryza and redness of the fauces, and usually by engorgement of the cervical and post-cervical glands. Any or all of these symptoms may be absent. The digestive system is not materially disturbed. The pulse is slightly increased in frequency, and the temperature rises from one-half to two degrees. Rōtheln rarely lasts more than five days, and the prognosis is always favorable.

DIAGNOSIS.—Enough has been said to indicate the mode of differentiating rōtheln from measles and scarlatina, the only disease with which it is likely to be confounded. It may not, however, be amiss to recapitulate briefly. The child who is taken sick may or may not have already passed through an attack of measles or of scarlatina, or of both. It may be known that it has been exposed to the contagion of rōtheln or that this disease is prevalent. The prodromoes, if such exist, are suggestive of scarlatina, but the rise of temperature and the acceleration of pulse are scarcely sufficient to accord with that suspicion. When within twenty-four or thirty-six hours the eruption appears, with, it is true, suffusion of the eyes and

slight coryza, but without distinct bronchial catarrh, the idea of measles may be dismissed. When further, the mild character of the angina, the continuance of only moderate fever and pulse-rate, the absence of nervous symptoms, and the disseminated macular character of the eruption are carefully considered, the question of scarlatina may be eliminated; and the diagnosis of rötheln is established by positive symptoms as well as by occlusion.

TREATMENT.—Beyond strict confinement to bed and a restricted diet, only simple remedies are required in the treatment of rötheln. We are in the habit of giving moderate doses of quinia, internally or by enema or suppository; and appropriate doses of aconite combined with sweet spirit of nitre, or effervescing draught in case of nausea. As the fever subsides and the eruption fades we advise the inunction of the surface as recommended in scarlatina.

ARTICLE VIII.

MALARIAL FEVER.

THE propriety of introducing a chapter upon malarial fever in the present work, is shown not only by the fact that malarial disease is extremely frequent in children, but also because it presents, as it occurs in them, so many peculiarities as to frequently lead to the true nature of such attacks being overlooked.

CAUSES; FREQUENCY.—There are cases upon record in which malarial disease appears to have been contracted in utero, and where immediately after the birth of the infant it has presented unmistakable evidences of the disease. We have ourselves met with several such cases, where the symptoms, and the prompt effect of quinine, left no doubt as to the diagnosis. At all periods of childhood, even from the age of a few weeks upwards, there can be no doubt that children readily contract malarial disease on exposure to its cause. Indeed, we have met with cases which have shown that the susceptibility of children to malarial poison may be even greater than that of their parents or other adults exposed to the same influences. In children over five or six years old the symptoms of malarial fevers are apt to be almost the same as in adults; the following remarks must therefore be understood to apply especially to those diseases as they present themselves in younger subjects.

SYMPTOMS.—Malaria presents itself in children both in acute and chronic forms. The former occurs both as intermittent and remittent fever. In our article upon typhoid fever we have carefully pointed out the fact, that in children the febrile movement in this latter disease often presents such marked remissions as to have led many authors to confound it with malarial disease, under the name of "Infantile Remittent Fever." But apart from this, true malarial remittent fever occurs in children, and indeed it is a peculiarity of all forms of malarial disease in early life to present a less marked development both of the paroxysms and of the in

missions. Intermittent fever in children may occur in any of the forms met with in adults, still the quotidian is by far the most frequent, the tertian less common, and the quartan decidedly rare. Whichever form may be present, is apt to present several peculiarities. In the first place, the features of the paroxysms are apt to be imperfectly developed. This is particularly true of the cold stage. It is very rarely present as a well-developed chill; in some cases, it seems to be entirely absent, but usually can be detected by careful observation. The child may merely become pale, seem weaker and more languid, or with this there may be distinct coolness of the hands and feet, and blueness of the nails; less frequently is there any discernible rigor, and, as before stated, a fully developed chill is very rare. The cold stage is of short duration, lasting from a few minutes to a quarter of an hour. It is followed by the hot stage, or in some cases the beginning of the attack is marked by the appearance of fever. The degree of this is rarely very high. Sometimes the child, who has been merely drooping during the earlier part of the day, is noticed to grow more dull, to wish to be constantly in bed, or on the lap, and its head and hands grow warm, with perhaps some flushing of the cheeks. Indeed, in some cases, the fever is so slight as to pass unnoticed, unless the attention of the nurse is directed to it by the physician. In other cases the accession of fever is more marked: the skin becomes very hot, and the cheeks brightly flushed; the child is dull and yet restless; there is rapid breathing, and marked acceleration of pulse. In some children, the fever is attended with delirium, and it is not a very rare thing to have it ushered in by a convulsion. This fact of the occasional occurrence of a convulsion, as a substitute for the chill as the initial symptom of the malarial paroxysm, must be borne in mind as of positive diagnostic importance. The fever lasts a very variable time, and rarely terminates abruptly, as in the case of adults by a sudden defervescence with profuse sweating. Indeed, in many cases, the child seems somewhat feverish during the entire twenty-four hours, but on careful observation is found to present increase of heat at some period of the day, and this is often preceded or followed by a short period during which the child is pale and languid, with cool moist brow and hands. Added to this irregularity in the symptoms and duration of the paroxysms, is the further source of difficulty, that the accession of fever occurs at very irregular hours. In children of even five years of age, it may occur at the ordinary time towards noon, but in younger children it may appear much later in the day, or even, as we have several times seen, late in the night.

There are a few other symptoms to be mentioned in connection with the paroxysms. We have already alluded to the occurrence of convulsions ushering in the hot stage. Frequently the child will vomit whatever food was in the stomach at the time of the attack. The urine that is passed during the paroxysm is scanty and high-colored, while not long after the subsidence of the fever, there is apt to be a quite free discharge of limpid urine. Between the paroxysms, if no complication exists, the child may appear merely listless, with scanty appetite. Quite frequently, however, the disease is attended with some more marked disturbance, either of res-

piration or digestion. The complications which we have ourselves most frequently observed have been gastro-intestinal catarrh, bronchitis and pneumonia. In cases where the latter has been present, the seat of the inflammation has occasionally been the apex of the lung.

The *chronic form* of malaria reveals itself in children in the same way as in adults. No well-marked paroxysms may occur, but the patient has a sallow, cachetic, or anæmic appearance, which of itself is quite characteristic. There is more or less emaciation from interference with nutrition, as the appetite is poor or capricious, and the action of the liver and bowels sluggish and insufficient. Enlargement of the spleen frequently follows, and we have met with well-marked examples of ague-cake in very young children. The blood becomes very poor and watery, and this, added to the obstruction to the circulation through the liver and spleen, in advanced cases may lead to ascites or œdema. We are not aware that the marked development of pigment-granules in the blood, which has been so often observed in the adult, has yet been detected in children suffering with chronic malaria. In some very severe and protracted cases, granular degeneration of the kidneys with albuminuria, and finally uræmia, has seemed to follow in quite young children. Some of the manifestations of malaria which are quite common in the adult are very rare in children. This applies especially to the various forms of neuralgia, which, as met with in the adult, are so frequently of malarial origin, while we do not remember to have met with a single case of this character occurring in children.

DIAGNOSIS.—It is our belief that malarial disease in children is often not recognized, and that this is due, not so much to its real difficulty of detection, as to the fact that the frequent occurrence of the different forms of malaria in young children, is not sufficiently borne in mind. Undoubtedly also there are difficulties in its diagnosis, which do not usually exist in adults. These arise, as before said, from the irregularity and imperfect development of the paroxysms. Our own experience has taught us in all cases of irregular febrile action, especially when occurring during the spring or fall, without any discoverable lesion to account for it, to suspect the malarial character of the attack. So, too, in cases where some slight lesion or disturbance of function exists, and yet the child seems too seriously and too obstinately ill for the apparent cause, and presents irregular fever with considerable fluctuations, the idea of the malarial nature of the attack should always be entertained. In some such cases, where it is impossible to reach a definite decision from a study of the symptoms, the diagnosis may be made by the therapeutic test of administering full doses of quinia for several days in succession.

PROGNOSIS.—The result of malarial fever is quite as favorable in children as in adults, when uncomplicated with any serious local inflammation. All of its forms usually yield readily to specific treatment. The chief source of danger lies in the tendency to severe bronchitis or pneumonia. In protracted chronic malaria, the anæmic and cachectic symptoms have seemed to us to yield to treatment even more rapidly than in the case of adults.

TREATMENT.—Children, even at a very early age, bear full doses of quinia very well. The amount which we have usually found necessary to arrest an attack of intermittent fever is three grains daily for children of one year of age or under, and one grain additional for each succeeding year, though we have given as much as five grains by the mouth in the course of the day to children of ten months, and without the slightest ill effect. It may be administered in the form of powders containing one-half grain, mixed with an equal amount of sugar and powdered extract of liquorice, repeated as necessary, and given at such times as to bring the system thoroughly under the influence of the drug before the hour at which the accession of fever has been noticed. Some children, however, will not take the powders without difficulty or nausea, and the quinia may then be given merely suspended in syrup of liquorice, or in the following combination:

R. Quiniæ Sulph., gr. xxiv.
 Acid. Sulph. Diluti, gtt. xxx. /
 Syr. Zingiberis, Syr. Simplicis, Aquæ, āā. f℥j.

Ft. sol.—Dose, a teaspoonful three or four times a day, according to age.

If, however, the stomach rejects it in all of these forms, as we have known it to do, we have found the administration by enema of two grains of quinia in a tablespoonful of starch-water, three times a day, equally successful. We may also resort to the use of suppositories, which when neatly made with butter of cacao and of small size are perfectly well tolerated, as a rule, even by very young infants. A small amount of dilute sulphuric acid, about one-half of a drop to each grain, should be added to the quinia in either of these modes of administration to favor its solubility and absorption.

In ordinary acute cases no other treatment is really required. It may be well to give a few doses of some saline febrifuge during each day, until the fever is entirely subdued, and of course any special disturbance of function must be relieved by appropriate remedies. The treatment of pulmonary complications must be subordinate to that of the general disease. All depleting or perturbing treatment must be avoided, and it will generally be found that with the aid of mild counter-irritation, the local symptoms will begin to improve after the malarial fever has been subdued by quinia. It is necessary to keep up the action of quinia for some time after the paroxysms are broken, because the tendency of the disease to recur is fully as great in children as in adults. We are in the habit of thus continuing it for three or four weeks in diminished doses, giving, however, on each septenary period, dating from the arrest of the paroxysms, the full antiperiodic dose, suited to the age of the patient. At the same time the child should take suitable doses of iron and arsenic, which may be conveniently given in the following form:

R. Liq. Potassæ Arsenitis, f℥j.
 Vini Ferri Amari, f℥ij.—M.

Dose.—From a half to a whole teaspoonful thrice daily in water after meals.

In chronic malaria we must persist in the use of quinia, iron, and arsenic

for a considerable period. At the same time careful attention must be paid to securing the best possible hygienic influences for the child. When practicable, a change of climate should be secured by a journey to the mountains or to the sea-shore. The patient should be warmly dressed, and carefully guarded against all exposure to damp or cold. The diet should be carefully selected, and every error of digestion promptly corrected. Even after the child is apparently restored to health, it should not be allowed to return to the locality where it contracted the disease, and for several successive springs and autumns should take a short course of quinia and arsenic. In the treatment of enlargement of the spleen, which frequently occurs in chronic malaria, we have obtained excellent results from the use of hypodermic injections of ergotin into the subcutaneous tissue of the abdominal wall.

ARTICLE IX.

MUMPS.

DEFINITION; SYNONYMS; FREQUENCY.—Mumps is an acute febrile specific disease, contagious and epidemic; occurring but once in an individual; attended by an inflammation of the parotid and sometimes of the submaxillary glands, with a tendency to metastasis to the testicles in males and to the mammae, vulva, or ovaries in females; and almost invariably resulting in recovery.

Some authors, as Niemeyer, object to classifying mumps with constitutional diseases; but the fact that it undoubtedly possesses the features enumerated in the above definition, and which, in the present state of our knowledge, must be regarded as specifically characteristic of that class of affections, seems to us to fully entitle it to be included with the other general diseases.

Mumps is known under a variety of names in every language. The other terms usually employed to designate it by English and American authors are cynanche parotideæ, parotitis, parotiditis, and inflammation of the parotid.

It will be impossible to obtain any definite idea as to the frequency of this affection, until the system has been introduced of registering not merely deaths but all cases of disease, since mumps is scarcely ever fatal. Its frequency is, however, known to vary very widely in different years, owing to epidemic influences; so that while in certain years we do not meet with a single case, in others we are called to see a considerable number.

CAUSES.—Nothing is known in regard to the essential nature of the cause of mumps. The disease is, however, unquestionably contagious, and it quite rarely happens that one member of a family sickens with mumps, without some of the other children being attacked.

Mumps rarely occurs as a *sporadic* affection, but appears, as already

stated, in *epidemics* of varying extent and severity, at times being limited to a single locality or even a single institution, and at others affecting large cities or districts.

Season appears to exert a powerful influence upon the development and activity of the specific poison of mumps, since the epidemics occur nearly always in the spring or autumn. According to Vogel, it is said to be endemic on the damp coasts of Holland, England, and France.

Age also exerts an unquestionable influence, by modifying the susceptibility to the contagion of mumps. Thus the disease is far more common between the ages of seven and fifteen years; whilst it is almost unknown before the end of the first year, comparatively rare between the ages of one and five years, and, on the other hand, quite rare in adults.

Although it appears certain, however, that the susceptibility to the contagion of mumps diminishes with each succeeding year after the age of fifteen, we must in great part explain the rarity of the disease in adult life, by the fact that a large proportion of people have had it in childhood, and are thus protected against a second attack.

Second attacks of mumps are indeed of extreme rarity.

ANATOMICAL APPEARANCES.—Opportunities very rarely occur for the examination of the parotid glands in mumps, since the disease is scarcely ever fatal. Virchow,¹ who has shown that, in cases of symptomatic secondary parotitis, the affection starts in the gland-ducts, maintains that the idiopathic form occupies the same seat. Bamberger,² on the other hand, states that the whole gland appears enlarged and reddened, with its tissues swollen and flaccid, owing to an interstitial exudation of lymph. The softness and indolent character of the swelling, however, the fact that it usually extends beyond the borders of the gland, and its usually rapid and complete subsidence, all induce us to believe rather that there is slight catarrh of the ducts, with mere oedema of the interstitial and surrounding connective tissue.

It is only in rare and very severe cases that there is sufficient lymph effused to undergo organization and lead to persistent increase in the size of the gland, or to so compress the ducts as to induce atrophy of the true gland-tissue. In even more rare cases it is said that suppuration may occur. In the secondary form, on the other hand, such as is seen in connection with the various specific fevers, the occurrence of suppuration is frequent.

SYMPTOMS.—In some cases the attack of mumps is preceded for a day or two by slight prodromes, consisting of restlessness, feverishness, loss of appetite or vomiting; in excitable children, symptoms of nervous disturbance may occur. More frequently, however, the local symptoms appear simultaneously with the fever, and we have generally found positive swelling of the parotid gland upon our first visit to the child.

The earliest local symptom is often pain, complained of under the ear,

¹ Quoted by Niemeyer (op. cit., vol. i, p. 436).

² Quoted by Vogel (op. cit., p. 113).

and increased by pressure and by all movements of the jaw, as in mastication. There is also stiffness felt in opening the mouth. The swelling appears first immediately beneath the ear; the depression between the mastoid process and the ramus of the jaw quickly becomes filled, and the swelling rapidly extends on to the cheek and neck. At first the swelling is flat, indurated, and presents the outlines of the parotid glands; but it soon becomes prominent, the most marked projection usually being observed immediately anterior to the lobe of the ear, and extends beyond the limits of the affected gland. The central part of the swelling corresponding to the parotid, remains firm, indurated, and more or less elastic, while at the periphery it is softer and often pits on pressure. The degree of enlargement varies much in different cases, being at times moderate and confined to the parotid region, while in other cases it extends over a large part of the neck and face, and may be so great as, especially when both glands are affected simultaneously, to give to the head and neck a pyramidal shape.

Quite frequently the submaxillary glands are involved, and the swelling consequently extends along the base of the jaw; in more rare cases, the enlargement is most marked in this region, or, indeed, the submaxillary glands may be almost exclusively the seat of the affection.

It seems important to call especial attention to this latter class of cases, since when the parotid swelling is absent and the submaxillary glands alone are involved the true character of the disease is apt to be overlooked. The fact that such irregular cases are true mumps is shown conclusively by the prevalence of an epidemic of mumps, by the occurrence of ordinary typical cases in their immediate connection, and by their power of communicating the disease to unprotected persons who may come in contact with them.

The skin over the seat of enlargement is at times scarcely altered in color, or may present more or less marked redness. There is usually only very moderate tenderness on pressure. The pain suffered during the attack varies greatly; in some cases it is merely a marked sense of tension and pressure, while in other instances it has been complained of as constant and severe, and extending even to the chest and shoulders. The movements of the head are impaired, and those of the jaw are impeded to such an extent that the mouth can only be slightly opened, and mastication is performed imperfectly and with great difficulty.

Usually the swelling increases for from three to five days, remains at its acme for a day or two, and then rapidly subsides, so that in about ten days the face has regained its natural appearance.

Mumps usually involves both parotids, though they rarely become affected simultaneously; the left gland is said to be most frequently the first inflamed, and subsequently, in twenty-four or forty-eight hours, or even when the swelling has disappeared from the side first affected, the opposite gland becomes enlarged. Occasionally the enlargement does not undergo complete resolution, and a circumscribed, painless, hard swelling remains for a variable time in the parotid region. In very rare cases suppuration is said to have occurred. The salivary secretion is variously

affected, and may be either diminished or excessive, or remain unaltered. Occasionally the external swelling is associated with enlargement of the tonsils and œdema of the submucous tissue of the pharynx. In such cases the difficulty of deglutition is much increased, and there may even be marked obstruction to respiration.

GENERAL SYMPTOMS.—Usually the constitutional disturbance in mumps is but slight and subsides even before the swelling of the parotid gland. Until the disease reaches its height, however, there is fever, with heat and dryness of the skin; the pulse and respiration are accelerated, the appetite impaired or lost, and the thirst usually extreme. There may also be, especially in nervous children, marked restlessness, sleeplessness from the pain and discomfort caused by the great swelling of the neck and face, and even mild delirium at night. As already mentioned, however, these febrile symptoms usually disappear about the fifth or sixth day.

One of the most curious features in parotitis is the tendency which occasionally exhibits itself to metastasis. The parts which are liable to be thus secondarily inflamed are the testicles and scrotum in males, and the mammæ, the vulva, and the ovaries in females. The most frequent of these metastatic inflammations in mumps is the affection of the testicle, which is much more common in men than in boys, is usually seated upon the same side with the enlarged parotid, and is attended with enlargement of the body of the testicle, serous effusion into the tunica vaginalis, and œdematous swelling of the scrotum. The swelling of the parotid ordinarily subsides when any of these metastatic affections appear, but occasionally the two inflammations continue together, a circumstance which shows, as Niemeyer pointed out, that they are in reality due to a common cause, and that no true transference of inflammation takes place from one point to the other. In some instances the swelling of the parotid subsides a variable time before the development of the metastatic affection, and, during the interval, alarming symptoms of depression and cerebral disturbance have been noticed, and at times referred to a metastasis to the membranes of the brain. There is, however, no actual meningitis present, and upon the redevelopment of the external swelling these nervous symptoms disappear.

PROGNOSIS ; DURATION ; COURSE ; TERMINATION.—Idiopathic parotitis or mumps almost invariably terminates favorably. The duration of the case varies from four or five days in very mild cases, to ten or twelve in severe ones. As already stated, the inflammation usually terminates in complete and rapid resolution. In some cases, however, a large amount of lymph is formed in the interstitial tissue of the gland, undergoes partial organization, and causes a hard, painless swelling, which persists for some time. In some epidemics, suppurative degeneration of the gland has been noticed, and the abscess which formed has either opened outwardly or into the external auditory meatus. We have known persistent hydrocele to follow the inflammation of the testicle occurring during an attack of mumps.

DIAGNOSIS.—The acute febrile character of the affection, and the pecu-

liar seat and shape of the swelling, always serve to render the disease readily recognizable.

TREATMENT.—As mumps almost invariably runs a favorable course, the treatment should be of a mild and expectant character.

The child should be strictly confined to bed; the diet should be fluid, partly on account of the great difficulty in mastication, light and digestible, consisting chiefly of preparations of milk and light animal broths. The only internal remedies required are febrifuges, such as spirit of nitrous ether and solution of acetate of ammonia, with a free supply of water and acidulated drinks; occasional laxatives; and, if there is sleeplessness, small doses of Dover's powder or some other anodyne.

Jaborandi has been asserted by Testa (*Il Morgagni*, July, 1878, quoted in *Brit. Med. Jour.*, August 23d, 1879) to be a most efficient remedy in this affection, and even to possess the power of aborting it if administered in time. We have no experience, however, to offer on this interesting point.

Local applications appear to have little or no influence upon the course of the swelling. The only ones to be recommended are warm, light poultices, or light water-dressings, covered with oiled silk, which do not annoy the child, and tend to favor resolution. If the induration be marked and extensive, so as to threaten suppuration, it has been advised to apply a few leeches behind the angle of the jaw. If it should become evident that suppuration has occurred, the abscess should be opened immediately, and the discharge favored by the application of poultices, in order to prevent further destruction of the gland or perforation of the external auditory meatus. In cases where induration and enlargement of the gland persist, absorbent applications, such as inunctions of iodine or mercury, should be made over the tumor.

In cases where alarming symptoms of depression and cerebral disturbance make their appearance after the sudden subsidence of the parotid swelling, the effort may be made to redevelop the external inflammation by stimulating applications to the surface, and by the internal administration of nervous and diffusible stimulants, such as ammonia, musk, or brandy.

After the acute symptoms of the attack have subsided, and the child has fully entered upon convalescence, we would caution against allowing it to leave bed too soon, since we have occasionally observed such premature exposure to be followed by marked febrile sequelæ. Thus in one case, occurring in an adult, there was marked fever lasting for a week; in another case, in a child, there was high fever for ten days; and in a third case, also in a child, there was a most obstinate and violent vomiting for four days, so violent, indeed, that we feared lest some renal complication might have been developed. On examination, however, the urine was found to be entirely normal.

ARTICLE X.

ERYSIPELAS.

DEFINITION; FORMS; FREQUENCY.—Erysipelas is a specific, acute, febrile, non-contagious exanthem, characterized by a deep red rash, attended with heat and swelling of the skin, sometimes with inflammation of the subjacent cellular tissue, and terminating generally in resolution, but sometimes in suppuration or gangrene. The disease is very variable as to its extent, and has the peculiarity of spreading from place to place, the part first attacked recovering, whilst the neighboring surface is becoming affected.

The disease, as it occurs in children over six months of age, presents the same characters as in adults, and requires therefore no particular attention in this work. In younger children, on the contrary, and especially in the new-born infant, it is different in several particulars from that of older children or adults, and this we shall attempt to describe. The form which occurs in new-born infants, has been technically named *erysipelas neonatorum*.

Erysipelas is a rare disease in private practice, particularly amongst families in easy circumstances. In lying-in and foundling hospitals, on the contrary, it is of frequent occurrence, and it is not uncommon in hospitals for children and in the children of the poor. We have ourselves met with but four cases of erysipelas in children under six months of age, whilst we have met with seven in older children.

CAUSES.—The erysipelas of young children almost always starts from some previously existing cutaneous inflammation, the most frequent seats of which are the umbilicus during the process of separation of the cord, the irritated folds of the skin existing in erythema intertrigo, the inflammation accompanying the vaccine disease, and that which exists in the eczematous and impetiginous eruptions of the scalp, ears, and face. In a large majority of the cases observed in new-born children, the disease begins upon the abdomen, and generally at the umbilicus. In those which occur in children at the breast, it may show itself at any of the points above mentioned.

The disease occasionally follows vaccination. We have ourselves met with three instances, in two of which the erysipelas broke out about the eighth day; and in the third on the tenth day. In none of these cases could there be any doubt as to the purity of the vaccine virus used. In two, the disease extended over the greater part of the cutaneous surface, lasting three weeks, but terminating favorably in both cases. In the third case, it extended over the whole of the vaccinated arm, then attacked the upper part of the trunk, the face, and the right arm, and terminated fatally in the second week.

But, though erysipelas commonly starts from, and may at first view seem to be produced by these different local irritations, it is impossible to suppose that they can be anything more than the exciting agencies or causes, which bring into action a disease of which the seeds already exist

in the economy. We must, therefore, in order to understand the real mode of causation of erysipelas, seek for the conditions that give rise to this predisposition to the malady, without which the above-mentioned exciting causes would rest without effect. These conditions are either a general epidemic constitution of the air, affecting certain districts of country, and acting more or less upon all classes of the community, but with especial force upon the destitute and miserable; or else a local epidemic constitution, such as that often occasioned by the unfavorable hygienic conditions of hospitals, and particularly of lying-in and foundling hospitals, or that not unfrequently determined by the same causes in the crowded and miserable habitations of the poorer classes of the inhabitants of large towns and cities.

SYMPTOMS.—Infantile erysipelas is not generally preceded by any constitutional symptoms. The appearance of the eruption is usually the first sign of the disease. So soon, however, as the eruption appears, or very soon after, the child is attacked with fever, marked by frequent pulse, heat and dryness of the skin, restlessness and insomnia, and thirst. In the form of the disease which occurs in very young infants and in hospitals, or amongst the lower classes of the population, the eruption almost always begins upon the abdomen, and very generally at the umbilicus, whence it extends to the rest of the trunk, to the genital parts, and sometimes to the inferior extremities. Even under the circumstances just mentioned, however, the eruption sometimes commences upon the face or upon the limbs. In children over two weeks of age, and in those observed in private practice, the disease may begin upon any part of the surface. It very often commences in the neighborhood of a vaccine pock, in a patch of erythema intertrigo, whether this be seated on the neck or about the pelvis, or it may appear first upon the face or upon one of the extremities, without any apparent exciting cause, and extend thence with greater or less rapidity to other parts of the body.

The form of the disease which recurs in very young infants, and which is by far more frequent in lying-in and foundling hospitals than under any other circumstances, begins almost always, at least when of a severe type, on the abdomen. It attacks hearty as well as more delicate children, and is generally very rapid in its progress. The erysipelatous surface is at first of a bright-red and shining appearance, but soon assumes a purplish hue, and as this occurs, becomes exceedingly hard to the touch, and somewhat, though not very much swollen. As the case goes on, unless resolution, which is a rare event, should take place, or death occur at an early period, the purple color deepens into livid, vesications occur, the cellular tissue is destroyed, and in many instances extensive gangrene takes place, so that the scrotum has been seen to "become black and slough away, leaving the testicles bare, and hanging loose by the cords." (Maunsel and Evanson.) In a case that occurred to one of ourselves in private practice, the disease began on the ninth day at the umbilicus, and involved the soft tissues of the anterior wall of the thorax and abdomen. The skin sloughed in several places, exposing the muscles; and at one point, just below the epigastrium, perforation of the abdominal wall occurred. Death followed

on the fifteenth day of the disease. In this form of infantile erysipelas, examination after death almost always discloses severe and extensive peritoneal inflammation, a condition which cannot fail, of course, to add greatly to the danger of the disease.

But infantile erysipelas does not always exhibit these violent characters, though whenever it occurs in infants under a year old it must be regarded as a very dangerous affection. When it attacks children over two weeks or a month old, it usually starts, as has been stated, from the neighborhood of a vaccine pock, from the inflamed surfaces of intertrigo or those of eczematous or impetiginous eruptions, or it begins without evident cause, as in adults, on the face, or on some part of the extremities. It appears first in the shape of a bright-red inflammation of the skin. After a short time the erysipelatous surface becomes tense, shining, very hot, slightly swollen, and painful to the touch. Pressure causes the color to disappear, but this rapidly returns when the pressure is removed. Coincidentally with the appearance of the cutaneous redness the child is seized with fever, restlessness, and severe thirst. From the spot first attacked the disease extends rapidly to the neighboring surfaces, from the neck and arms, to the head and trunk, and from the groin or genital parts to the rest of the trunk and to the inferior extremities. When it begins upon the face, it extends to the scalp, and may thence travel over the whole surface, or it may remain limited, as it often does in adults, to the head alone. In one case that we saw, in an infant three weeks old, in which it began upon the face, it extended gradually over the whole cutaneous surface, and yet the child recovered. In another, two months old, it began upon the bridge of the nose, and from thence extended over the whole head, but did not reach the trunk or limbs. In a third case, a vaccinated arm was attacked with erysipelas on the eighth day of the vaccination. The disease extended down to the fingers, and upwards to the shoulder. From the shoulder it spread gradually over the whole trunk, and down the whole length of both lower extremities. As it was subsiding on the feet, it appeared on the arm opposite the one first attacked, and then attacked the corresponding side of the head, where it ceased. The child finally recovered after an illness of three weeks.

As the peculiar inflammation spreads to the neighboring surfaces, the parts first attacked lose their red color and swelling, and undergo a process of desquamation. In some instances, the inflammation has caused suppuration of the subcutaneous cellular tissue, so that even when the greater part of the surface first attacked has ceased to present the peculiar character of the erysipelatous inflammation, there remain behind abscesses of greater or less extent. Thus, in one of the cases that came under our own notice, when the erysipelas had left the head and thorax, and was confined to the pelvis and inferior extremities, there were two abscesses on the scalp, and one over the right pectoral muscle, whilst all the skin between the abscesses had regained its natural appearance, with the exception of the desquamative process, which was going on as usual. In another, but rarer set of cases, the inflammation sometimes returns to the parts over which it has already passed.

The swelling which accompanies this disease is usually of an œdematous nature,—the œdema being most marked in the hands and feet, and upon the face, whilst upon the trunk it is much less considerable.

The general symptoms consist at first, as already stated, of those indicating a strong febrile reaction. If the case goes on favorably these symptoms continue until the disorder terminates. But when the disease is severe, and especially when it ends in vesication, in extensive destruction of the cellular tissue, or in gangrene, the general symptoms are much more violent, marking thereby the gravity of the attack. The face and lips become pale, and the tongue and mouth dry. The child is in a state of constant agitation at first, and expresses its uneasiness and suffering by incessant moaning or crying, but after a time, it becomes heavy and drowsy from exhaustion. The pulse is very frequent and feeble; diarrhœa and vomiting make their appearance, and the child dies at last in a state of profound debility; or convulsions occur towards the last, and terminate the case, as they so often do in the diseases of infancy and childhood.

The *duration* of erysipelas in children is extremely uncertain, and depends very much upon its form. In that which occurs in the new-born child, or within one or two weeks after birth, it sometimes proves fatal within seven days according to Canstatt (*Handbuch der Med Klinik*, 2d ed., vol. ii, p. 264). M. Bouchut (*Mal. des Enf. Nouv.-Nés*, p. 532) gives as an approximation to the ordinary duration of infantile erysipelas, between four and five weeks, and states that this is also the result arrived at by M. Trousseau. In one of the cases alluded to by us, in which the disease extended over the whole cutaneous surface, the duration was four, while in another it was three weeks; in the one in which the eruption was limited to the head, the duration was a week. In the seven remaining cases, the disease was limited to the nose and eyelids, or the face and scalp, and lasted from three to ten days.

DIAGNOSIS.—The diagnosis is very easy. The peculiar shade of the red color, the presence of decided though moderate tumefaction of the affected part, the severity of the general symptoms, and the characteristic erratic mode of extension from surface to surface, all assist to render the diagnosis very clear to those who have a proper amount of medical knowledge.

PROGNOSIS.—Erysipelas is always a dangerous disease in young children. The precise degree of danger in individual cases will depend chiefly on two circumstances: first, the age of the subject; and second, the hygienic conditions under which the disease occurs. It is exceedingly dangerous in new-born infants, so much so indeed that M. Bouchut declares that they all die (*loc. cit.*, p. 532). This is in all probability almost strictly true of the cases which occur in infants only a few days old, particularly when they take place in lying-in hospitals, or even in private practice, during the prevalence of an epidemic of puerperal fever. The disease is always very dangerous in hospitals, even in infants over two weeks old. Yet it would appear not to be so grave as represented by M. Bouchut, who thinks that very few indeed have been cured even at that age; for, of thirty cases in infants between one day and a year old observed by Bil-

lard at the Foundling's Hospital of Paris, sixteen, or only one more than half, proved fatal. Schwebel reports 54 deaths in 86 cases (Meissner, *Kinderkrankheiten*, 3d ed., vol. i, p. 372).

In private practice, erysipelas, as it occurs in children between two weeks and a few years old, is a dangerous malady, but yet it is far from being so in the same degree as in the new-born infant, and in hospitals. We have already stated that we have seen four cases in young infants; one nine days old, in whom the disease proved fatal in fifteen days; one three weeks old, in whom the disease lasted four weeks, and travelled over the whole cutaneous surface; another ten weeks old, in whom also it travelled over the greater part of the cutaneous surface; and a fourth two months old, in whom it remained limited to the head. These last three recovered. Again, we have seen seven cases of erysipelas of the face or head in children between seven months and twelve years old, and these also ended favorably. It must be recollected, however, to account for these recoveries; that they all occurred in robust children, and under the most favorable hygienic conditions met with in private practice. To conclude MM. Rilliet and Barthez report nine cases of erysipelas of the face in children, all of whom, with three exceptions, were over five years of age. Five of the nine cases were idiopathic; in four the disease complicated other affections. All of the spontaneous and one of the complicated cases recovered. The two others, both of which occurred in subjects laboring under measles attended with pneumonia, proved fatal.

TREATMENT.—The treatment of erysipelas in new-born infants, especially when the subjects of the disease are the inmates of a hospital, and when it occurs coincidently with a puerperal fever epidemic, is, as may be learned from the almost certain fatality of the disorder, exceedingly hopeless. M. Trousseau (Barrier, *Traité Prat. des Mal. de l'Enfance*, t, ii, p. 560) has made trial unsuccessfully of emollients in every form, of fomentations, lotions, baths, and of ointments containing sulphate of iron. "I have tried," he says, "surrounding the whole body and limbs with blisters in the form of strips; the erysipelas has passed over the obstacle. I have applied without success blisters upon the surfaces already invaded by the inflammation. I have obtained no advantage from mercurial ointment or from baths containing corrosive sublimate." He even tried the application of the actual cautery in points where the disease was beginning, but without effect. So, too, with methodical compression.

Underwood says that "upon the complaint being first noticed in the British Lying-in Hospital, various means were made use of without success; the progress of the inflammation has seemed, indeed to be checked for awhile by saturnine fomentations and poultices, applied on the very first appearance of the inflammation; but it soon spread, and a gangrene presently came on; or where matter has been formed, the tender infant has sunk under the discharge." He adds that he then proposed bark, to which sometimes a little confectio aromatica was added, and that from that period several cases recovered. After this, linen compresses, wrung out of camphorated spirit, were applied in the place of the saturnine solution, and proved successful in several instances in checking the inflammation.

"Nevertheless, the greater number of infants attacked with this disorder sink under its violence, and many of them in a very few days." (*Treat. on the Dis. of Children*, Am. ed., by Dr. Bell, from the 9th Eng. ed., p. 103.) In a note to the above, Dr. M. Hall stated that fomentations of extract of poppies diffused in warm water, and poultices consisting of the same fluid and crumbs of bread, proved beneficial in many instances. Dewees recommended the application of a blister, when the erysipelas is so situated as to allow the whole surface of inflammation and a portion of the neighboring healthy surface to be covered by the plaster. When this cannot be done, he preferred the use of the strong mercurial ointment, which must be applied over the whole of the eruption, and partly upon the healthy skin, and renewed as often as the part becomes dry.

It is very difficult amidst the variety of advice given by different writers, and especially when we reflect upon the great mortality of the disease under every kind of treatment, to determine which to select. For our own part, we should prefer the use of cooling emollient applications during the first part of the attack, whilst the skin is of a bright-red color, hot, and shining. When the circulation becomes languid, and the color of the eruption is disposed to deepen from red to purple, we should suspend the use of the emollient applications, and employ instead the lotion of camphorated spirit recommended by Underwood; the camphorated tincture of soap, which we have known to be of great service in the erysipelatous inflammations occurring in patients of broken-down constitution, and which is to be applied three or four times a day by means of a soft sponge; or lastly, we would make trial of Kentish's ointment, a remedy found of great service by the late Dr. Charles D. Meigs, in the erysipelas of children (*North Amer. Med. and Surg. Jour.*, vol. vi, p. 77). This ointment he prepared by rendering basilicon ointment soft (not fluid) with spirit of turpentine. It is rubbed upon the inflamed part with the fingers, the anointing being "repeated often enough to keep the part always very thinly covered." The internal treatment should consist in attention to the state of the bowels, which are to be kept soluble by the mildest laxatives, without being purged, and in a resort to tonic and stimulating remedies upon the very first approach of symptoms indicating exhaustion. The best remedies of this class are proper diet, wine whey, small quantities of brandy, and bark in connection with minute doses of carbonate of ammonia.

In addition to these, the tincture of the chloride of iron, whose remarkable and almost specific influence upon the course of erysipelas in more advanced life is so well established, should be given in large doses, proportioned to the tender age of the patient. Thus we may give two or three drops every three hours to an infant of a month old, as in the following formula:

R. Tr. Ferri Chloridi,	f ʒ ss.
Acid. Acetici Dil.,	f ʒ ss.
Liq. Ammoniae Acetat.,	f ʒj.
Syr. Simp.,	f ʒ ss.
Aquæ,	q. s. ad f ʒ iij.—M.

Dose.—A teaspoonful every three hours.

When the inflammation has gone on to the production of subcutaneous suppuration, it becomes still more important to sustain the forces of the constitution, by giving the infant a healthy and abundant breast of milk, and by the internal use of brandy in small quantities, of bark, or better still, of quinia in combination with small doses of carbonate of ammonia. At the same time the suppurating surfaces must be well fomented, and dressed with warm poultices, and when necessary, laid open by careful incisions, observing the precaution to cause as small a loss of blood as possible. If the case occur in a hospital, or in a child placed in unfavorable hygienic conditions, let the following statement of M. Barrier (*loc. cit.*, t. iii, p. 562) be borne in mind: "However much the life of an infant be threatened by erysipelas, if we can but persuade a wet-nurse to take charge of it, the pure air of the country is often seen to replace most advantageously all other therapeutical resources."

As the proceeding remarks have been restricted to the form of the disease which occurs in infants under two weeks of age, we have now to make some observations on the cases which occur in older children.

The disease is still, even at this latter age, a very dangerous one, though much less so, certainly, than in the new-born child. We have been deterred from the use of depletion in any form by two reasons,—the fear of exhaustion, which is so apt to occur in the disease, and the apprehension lest the leech-bites or cup-marks, in the case of local depletion, might prove new foci of the erysipelatous inflammation. The only internal remedies necessary in the beginning, are such laxatives as may be required to keep the bowels soluble when they are constipated, such as shall correct acidity or diarrhœa when either is present, and those which promote an open state of the skin, and a free discharge of the urinary secretion. For the latter purpose we know none better than the solution of the acetate of ammonia, and the sweet spirit of nitre, about twenty or thirty drops of the former, with five of the latter, in sweetened water, to be repeated every two or three hours. The tincture of the chloride of iron should also be given, in the combination before recommended, in large doses, as three to six drops, every three hours, at the age of one or two years. Should the attack be attended by any symptoms of prostration, or at a later period of the disease, when the child begins to emaciate and grow feeble, its strength must be carefully supported by the use of proper diet, and of stimulants and tonics. The only proper diet for nursing children is, of course, breast-milk; for those who have been weaned, the diet should consist of preparations of milk, light animal broths, or beef-tea. The best stimulants are five or ten drops of brandy, five drops of aromatic spirit of hartshorn, or a quarter or sixth of a grain of carbonate of ammonia, in weak syrup of ginger, to be administered four or five times a day, or more frequently, when the forces of the child are greatly prostrated. The proper tonic is half a grain of quinia, in some suitable vehicle, every three or four hours.

The best local treatment is, in our opinion, cooling or tepid emollient applications, as slippery elm bark, marsh-mallow, or flaxseed tea, during the first few days, whilst the reaction is marked, and the temperature of the body high. Somewhat later, when the strength begins to be reduced,

and the color of the eruption to deepen, we should make use either of mercurial ointment, which is highly recommended by some, or of Kentish's ointment, or camphorated tincture of soap, to which attention has already been called. We would here propose the trial of an ointment which we have found not only soothing and comforting to the child, but also of manifest curative efficacy in the violent cutaneous inflammation of scarlatina. It consists of one ounce of fresh cold cream, rubbed up with a drachm of glycerin. It should be smeared over the inflamed surface several times a day, and need not interfere with the use of emollient applications. In scarlatina it has been most useful in reducing the burning heat of the eruption, and in softening the harsh and distended skin, and by these effects has aided greatly in moderating the severity of the general, and especially of the nervous symptoms. Cosmoline may also be used for the same purpose and with the same good effect.

In children over two or three years of age, erysipelas must be treated on the same principles as in adults, by light but nourishing diet, and rest in bed, by the occasional use of laxatives, of full doses of the tincture of chloride of iron, and of febrifuges, and by the external application of emollient infusions, so long as the symptoms remain acute and the strength unreduced. But when, after a time, the fever begins to subside, or the child begins to show signs of debility and a tendency towards the typhoid condition, we must endeavor to maintain the life-actions in a proper degree of energy by a more nourishing and abundant diet, by the prudent administration of bark or of quinia, and even by the use of brandy and ammonia, should the strength of the patient be disposed to give way suddenly or rapidly. Under these circumstances, moreover, the best local application will be either Kentish's ointment, or the camphorated tincture of soap.

ARTICLE XI.

DIPHTHERIA.

DEFINITION ; SYNONYMS ; HISTORY ; FREQUENCY.—Diphtheria is an acute febrile, moderately contagious, and infectious asthenic blood disorder, occurring both endemically and epidemically ; without characteristic eruption, and distinguished by a disposition to the formation of false membranes upon inflamed mucous surfaces, especially in the fauces, or upon abrasions of the cutaneous surface.

It is the disease called by the older writers, *angina maligna* or *gangrenosa* ; *cynanche maligna* ; *garotillo* ; *angina suffocativa*, under which name

it was described by Dr. Samuel Bard, of New York, in one of the best of the early essays upon this subject (*Trans. Amer. Philos. Soc.*, vol. i).

It is, indeed, thought probable, that the history of this affection can be traced back to a period beyond the time of Hippocrates; but unquestionably the writings of Aretæus, who flourished in the second century of the Christian era, contain a distinct description of this malignant sore throat. He describes it under the names of *ulcus Syriacum* and *malum Ægyptiacum*.

From this period, there is quite frequent mention of the disease in the works of medical writers; the earliest account of its appearance in modern times being given by Hecker, who describes an epidemic of it that prevailed in Holland, in 1337.

About the middle of the last century, it prevailed in Paris, where it was described by MM. Malonin and Chomel; and in some parts of England, where it was studied and described by Fothergill, though it is now doubted whether the disease to which he refers was not more nearly allied to scarlatinous angina.

The first full description of this affection published in this country, was the paper, already referred to, by Dr. Bard, based upon an epidemic which appeared in 1771; the views advanced in which have been universally recognized, even to the present day, as most clear and just.

From that time, the complaint seems to have attracted but little attention, until its occurrence at Tours, in 1818, and subsequent years, called forth the treatise of Bretonneau in 1826, in which he gave the first precise notion of the disease, and bestowed the name diphtherite upon it.

Since then it has occurred frequently epidemically in France; in 1857 it appeared almost simultaneously in England, and in the extreme western part of our own country, and from that time has occurred in the form of epidemics of greater or less extent and severity, in the most varied climates and seasons, in almost all known parts of the globe.

Diphtheria, the name by which this epidemic pseudo-membranous angina is commonly designated, is a synonym of the word diphtherite, originally used by Bretonneau in his treatise on this subject.

Διφθέρα and *Διφθέρης* both mean "the prepared skin of an animal;" and *Διφθερίτης* and *Διφθερίας* signify alike, "that which is covered with a skin or membrane."

No cases of death from diphtheria in Philadelphia are reported in the annual lists of mortality, published by the Board of Health, until the year 1860. In the preceding report, however, Dr. Jewell mentions that several severe cases had occurred, some of which had proved fatal. One of us can, however, assert from his personal experience, that well-marked cases of diphtheria were of not rare occurrence in this city for a number of years before that time, but were reported under other names, and usually as either croup or angina.

It is probable, however, that the disease did not prevail at all extensively previously to its great outbreak in 1860, as may be seen by a reference to the number of deaths from croup and scarlatina, returned for the years preceding and subsequent to that date.

TOTAL NUMBER OF DEATHS FROM

	Scarlatina.	Croup.	Diphtheria.
1855,	163	265	
1856,	992	268	
1857,	704	256	
1858,	241	292	
1859,	232	312	
1860,	206	354	307
1861,	329	304	502
1862,	461	258	325
1863,	275	444	434
1864,	349	455	357

The total number of deaths from scarlatina, from 1855 to 1859 inclusive, was 2332; from 1860 to 1864 inclusive, 1620, or 712 less than in the previous period.

The total number of deaths from croup from 1855 to 1859 inclusive, were 1393; from 1860 to 1864 inclusive, 1815, or 422 more than in the previous five years. And, further, during the latter five years, 1860 to 1864, the deaths from diphtheria amount to 1925.

CAUSES.—Diphtheria occurs in both endemic and epidemic forms; and the various outbreaks vary widely in gravity of type, and in the extent of territory involved. No less surely established is its contagious and infectious nature. Until recently, doubts were frequently expressed as to whether diphtheria is really contagious, but the evidence accumulated is sufficient to show that it is unquestionably so, although, as in other zymotic diseases, the activity and virulence of its contagious principle varies greatly in different cases.

The infectious nature of diphtheria was clearly recognized by Bretonneau, and many incontestable cases are on record to show its transmissibility by the direct contact of the diphtheritic exudation with an absorbing surface. Thus, the disease has, in repeated instances, been acquired by physicians in attendance on cases of diphtheria, by the entrance of fragments of exudation to the lip or mouth, while making local applications to the pharynx or while sucking the wound during the performance of tracheotomy.

Apart from these well-ascertained properties, nothing is as yet positively known with regard to the general conditions which favor its production; and it appears to have prevailed with equal severity in healthy and unhealthy situations; in damp marshy districts and in dry hilly regions; in the crowded filthy houses of great cities, and in sparsely populated villages; in the depth of winter, and in the intense heat of summer.

Nor can it yet be positively asserted (although it is probably true with regard to diphtheria, as in the case of other zymotic diseases), that children of feeble constitution and those subjected to bad hygienic conditions, or debilitated by severe illness, are particularly exposed to it, especially in the sporadic form.

Of late years, we have been inclining to the opinion, although as yet

more positive evidence is needed, that the prevalence of diphtheria and especially its virulence, are favored by defects of drainage and by contamination of the air and drinking-water.

The effect of local causes, of a depressing character, upon the production of diphtheria, was investigated by Dr. Ballard, in regard to 57 fatal cases. Inquiries at the houses where the 57 deaths had occurred, showed that in 24 instances the houses were damp, and that defective drains or some similar cause gave rise to offensive smells; in four houses the inmates were overcrowded, and the ventilation deficient; in 8 cases the drinking-water was foul, or there was some noxious accumulation; and in 25 cases nothing whatever could be discovered amiss in the hygienic condition of the houses.

As an instance of bad sanitary conditions which would certainly induce or favor an outbreak of diphtheria, the following is instructive:

On March 12th, 1878, the mother and the eldest daughter of a family, comprising father, mother and six children, and living in a very healthfully situated cottage in Newport, R. I., visited for a short time a house where two days subsequently three persons had mild diphtheria. On the evening of the 12th, the oldest daughter was attacked, and within fifteen days all the children were seized with malignant diphtheria and died. The father had a severe attack, the mother a mild one, and both recovered. A careful examination of the premises by Colonel G. E. Waring, Jr., the well-known sanitary engineer, revealed a foul condition of the water-closet, a very unhealthy arrangement of the waste-pipe of the sink, and a break in the trap of the water-closet through which fecal matter had escaped and had accumulated in large quantity under the floor of the scullery.

It is certain, however, that occasionally diphtheria appears in a sporadic form, and isolated cases occur which can be attributed to no known cause whatever. Attention has lately been called by Mr. W. H. Power (*Med. Times and Gaz.*, Jan. 18, 1879, pp. 66 and 75) to the possibility of contaminated milk serving as a means of promoting the spread of diphtheria.

We subjoin a table of the mortality from croup and diphtheria in this community during the seven years from 1874 to 1880 inclusive; upon which we base, to a great extent, the remarks which follow as to the causation of the latter disease.

MORTALITY TABLE OF CROUP AND DIPHTHERIA DURING SEVEN YEARS, FROM 1874-80, INCLUSIVE.

MONTH.	1874.		1875.		1876.		1877.		1878.		1879.		1880.		Mean of these Seven Years.		Mean Temperature for Seven Years.	Mean Fall of Rain during Seven Years.
	Croup.	Diphtheria.	Croup.	Diphtheria.	Croup.	Diphtheria.	Croup.	Diphtheria.	Croup.	Diphtheria.	Croup.	Diphtheria.	Croup.	Diphtheria.	Croup.	Diphtheria.		
January, .	22	9	57	41	42	82	41	46	53	57	51	46	40	35	43.7	45.1	33.21	2.818
February,	16	12	39	27	57	56	19	32	37	32	41	38	28	30	33.8	32.4	33.68	2.398
March, . .	20	11	38	48	31	64	33	42	53	46	28	37	32	31	33.6	39.8	39.68	3.438
April, . .	19	7	24	47	40	74	28	22	22	27	12	31	29	16	24.8	32.0	49.21	3.802
May, . .	16	9	22	46	27	53	18	41	20	26	12	21	24	30	19.8	32.3	62.32	2.101
June, . .	8	12	38	65	19	70	12	34	21	28	8	14	13	22	17.0	35.0	71.82	3.822
July, . .	5	8	20	45	17	36	8	22	7	30	10	18	19	22	12.3	25.8	76.32	4.690
August, .	5	3	14	57	18	31	11	25	11	18	8	12	10	8	11.0	22.0	73.18	4.251
September,	5	9	33	42	23	48	22	37	18	42	9	16	13	24	17.5	31.1	66.10	3.318
October, .	18	15	43	76	34	62	38	42	36	52	18	22	27	34	30.5	43.3	55.97	2.301
November,	20	38	47	65	42	51	53	44	61	47	50	36	23	40	42.3	45.8	43.30	3.641
December,	45	48	53	97	36	81	55	71	49	59	44	30	45	31	46.7	59.5	34.05	2.858
Total,	199	181	428	656	386	708	338	458	388	464	291	321	303	323

Season.—As we have already remarked, the influence of season upon the prevalence of diphtheria is comparatively slight, and there are numerous records of epidemics occurring in the summer, as well as in the winter months. Thus it will be seen from the accompanying table that it was only in July and August that the mean mortality from diphtheria fell decidedly below the average of the rest of the year, while even in those months a notable proportion of deaths occurred from this disease. During the seven years taken as a basis of this calculation, the highest mortality was in the months of October, November, December, and January.

It is true that croup exhibits the influence of season and temperature more markedly than does diphtheria, and yet a study of this table will show that, even in regard to croup, this influence has been generally overestimated. Thus the fluctuation in the mortality from diphtheria and croup will be seen to correspond closely; the highest point being reached by both in the same month, December, and the lowest by both in August. The difference between the minimum and maximum mortality of the year is much greater in the case of croup than in that of diphtheria, being as 1 to 4.25 in the former, and as 1 to 2.7 in the latter.

Sex appears to have absolutely no influence upon the frequency of diphtheria, since of 3111 fatal cases occurring in this city during the above seven years, 1457 were males, and 1654 females.

Age, on the other hand, unquestionably exerts a very strong predisposing influence, a large majority of all recorded cases occurring between the ages of one and eight years.

Of the 3111 cases in our table, 212 occurred under the age of one year; 489 between one and two years; 1353 between two and five years, and 836 between five and ten years. Although the liability thus diminishes, in an uncertain ratio, with advancing years, no age is exempt from it. By reference to the influence of age upon the frequency of true primary croup, it will be seen that the maximum of its frequency is also attained between the ages of one and five years. We would also call attention to the much greater frequency with which diphtheria occurs in later life than croup; since of 2333 deaths from croup, but 22 were over ten years of age; while of 3111 deaths from diphtheria, not less than 221 occurred after that period. Of course it is evident that the above statistics not only prove that diphtheria is much more frequent during the first decade of life than at any subsequent period, but also that it is much more fatal then.

NATURE.—In his earliest writings upon this subject, Bretonneau attached little importance to the constitutional symptoms attending diphtheria, and upheld the view that it was essentially a local affection; and though he subsequently somewhat modified his views, he yet only admitted that the constitution becomes involved secondarily.

It is indeed true that the epidemics which have occurred during the past twenty-five years seem to have been attended by far more grave constitutional symptoms than were present in the cases upon which Bretonneau's memoir was founded. Moreover, the development of our knowledge of zymotic diseases has advanced rapidly during that period; so that it may be stated to be the almost universally adopted view that diphtheria is a

blood disease, dependent upon the admission to the system of some specific morbid principle.

The chief arguments in favor of its being a constitutional disease, are its epidemic and contagious nature; the continued febrile action, of asthenic type, which attends its course; the marked alteration of the blood mass in color and consistence; the tendency to pseudo-membranous exudation on mucous membranes, or abrasions of the skin; the occurrence of albuminuria; and, finally, the frequent development of paralytic sequelæ, showing the presence of some morbid agent, acting especially upon the nervous system.

It must be admitted, however, that considerable plausibility attaches to the later view of Bretonneau, which has been strongly advocated by Bouchut.¹ He divides diphtheria into false, or non-infecting, which is mere pseudo-membranous angina; and the true, or infecting, which involves the entire system, by means of the absorption of septic substances from the pharynx. In this respect it resembles pyæmia, and produces swelling of the lymphatics, alteration of the blood, albuminuria, and even metastatic deposits.

Recently² the results obtained by Drs. H. C. Wood and Henry F. Formad, of the University of Pennsylvania, working at the suggestion and under the auspices of the National Board of Health, from an investigation of diphtheria occurring epidemically, sporadically, and from artificial causes, tend strongly to support the above view.

PATHOLOGICAL ANATOMY.—False Membranes.—We have already dwelt upon the fact, that the pseudo-membranous exudation can no longer be regarded as the essential and most important element in diphtheria; it is, however, one of the most constant and striking phenomena, and in certain cases, where it extends into the larynx, becomes the effective cause of death.

When fully developed, the pseudo-membranous deposit has the ordinary appearance of a fibro-plastic membrane, as more fully described below. Its appearance is preceded by swelling and infiltration of the mucous membrane, and by some submucous exudation of a viscid, sero-mucous liquid. The membrane itself appears in the form of points of grayish-white or slightly yellowish tint, which, at first isolated and circumscribed, soon coalesce.

This pellicle is more dense and thick at its centre than towards the edges, and soon after its formation, the exudation continuing beneath it, and coalescing with it, it gains in thickness by the apposition of an under layer; until, when the membrane is fully developed, it may consist of several layers, and appear imbricated.

At this period its adhesions are so strong that, if it be detached from its connections, slight hemorrhage will follow, or numerous minute bloody points may be seen upon the subjacent mucous membrane.

The appearance of the opaque points has been attributed to the coagu-

¹ Mal. des Enfants, 4ème éd., pp. 907-923.

² Supplement No. 7, *National Board of Health Bulletin*, and *Phila. Med. Times*, Oct. 22, 1881, p. 33.

lation of fibrin in the clear sero-mucous fluid; but according to the most recent researches the exudation is almost exclusively composed of cells. The microscopic appearances which are constant are the ordinary elements of corpuscular lymphatic leucocytes, granular cells, and free fatty granules, more or less abundant and closely interlacing fibrillæ, mixed with epithelial cells of various shapes and sizes. It appears, therefore, that the cellular elements are derived in large part from the cells of the superficial layers of the mucous membrane, and an interesting discussion has been maintained between Wagner,¹ who insists upon a specific alteration of these layers, and Boldygreu, Steudener, and others as to the precise character of these changes. Rindfleisch² says: "The false membrane is undeniably produced by the secretion of young elements upon the irritated mucous surface, followed by their gradual stiffening, sclerosis, glassy swelling, or whatever term we may choose to apply to their degeneration. Accordingly, the false membrane occupies the precise position which belongs of right to the epithelium; the degeneration in question taking the place of the normal evolution of epithelial elements." In addition to this must be clearly admitted the coagulation of a liquid rich in fibrin, and the escape from the distended vessels of white blood corpuscles, which become fixed by this process of coagulation.

It is necessary to allude here to the very important question of the relations of minute parasitic organisms to the exudation and to the general symptoms of diphtheria. This is the more important on account of the remarkable results recently published by Wood and Formad (*loc. cit.*). It has long been known through the observations of Vogel, Laycock, Wade, Oerbel, Letzerick, and others, that a form of fungus is often found in the exudation and secretions in cases of diphtheria, but it also appears that a fungus is present in numerous diseased conditions of the mouth and fauces. Wide differences of opinion have existed as to whether this fungus was the cause of diphtheria, or whether it was merely an accidental development, due to the fact that the spores found a favorable nidus in the diseased secretions. It appears from the observations of Wood and Formad, that the micrococci found in diphtheria do not differ essentially from those found in ordinary sore throat; but that they are the same organism in a state of higher reproductive activity. They do not directly cause diphtheria, nor do they, by entering the blood, directly cause the symptoms of septicæmia. But it is possible that they exert upon the diphtheritic exudation somewhat the same action that the yeast-plant does upon sugar, and cause the production of a septic poison which if absorbed will induce the symptoms of constitutional diphtheria. It would certainly appear from the observations and experiments of Wood and Formad, that in simple sporadic diphtheria the micrococci do not multiply so actively, nor develop a powerful septic poison, and that, if death occurs from intercurrent croup, no micrococci will be found in the blood or internal organs, while in grave epidemic cases, with marked septi-

¹ Manual of General Pathology, New York, 1876, p. 265.

² Pathological Histology, New Syd. Soc., ed. 1872, p. 425.

cæmia, very numerous micrococci were invariably found in these localities, being especially numerous in the spleen and liver.

Color.—The color of the pseudo-membrane varies at different stages, and somewhat according to its seat.

In the fauces, the deposit is often whitish at first, but soon acquires a yellow tint; though in some cases it is quite gray, and produces the appearance of extensive sloughs on the fauces and pharynx. In severe cases, there is usually a bloody sanious fluid effused, which imbues the pseudo-membrane, discolours it, and promotes its decomposition, so that it forms dark-colored shreddy patches, exhaling a fetid, gangrenous odor.

It is essential to bear in mind that these appearances of the fauces in diphtheria are usually due to decomposition of the false membrane alone; and that if this be removed, the mucous membrane will generally be found merely raw, excoriated, and oozing blood.

It is, however, true that in certain epidemics the rule has been for serious lesions of the mucous membrane, involving even its entire thickness, to occur.

In milder cases, where the disappearance of the false membrane can be studied, it is never seen to separate all at once, leaving in its place a cicatrized surface, but the pellicle gradually diminishes in thickness and extent. When the pseudo-membrane extends into the larynx, it is more apt to remain whitish throughout its course there than in the fauces.

Consistence.—The consistence of these deposits varies considerably. In cases of ordinary severity, where the symptoms are not of a very adynamic type, the pseudo-membrane is often quite firm, tenacious, and elastic; while in grave asthenic cases, with severe inflammation of the throat, the deposit is apt to be much less firm, or even quite pultaceous.

It has been attempted to base upon these conditions and corresponding microscopic appearances, a division of diphtheritic pseudo-membranes into two classes, answering to the well-known division of inflammatory lymph into the fibrinous and the corpuscular.

Chemical Characters.—The false membranes contract and shrivel when treated with alcohol; mineral acids, such as sulphuric, muriatic, nitric, or chromic; strong solutions of nitrate of silver; or solutions of the per-salts of iron.

On the other hand, they soften more or less quickly when treated with alkaline solutions, as of potassa, soda, lime, or ammonia; or of chlorate of potash, chlorate of soda, bromide of potassium; or with glycerin and various other agents. Recently, pepsin and lactic acid have also been announced as powerful solvents. These various chemical properties are constantly turned to account in the treatment of diphtheria, in guiding our selection of the most appropriate local applications.

Condition of the subjacent Mucous Membrane.—Even before the appearance of the slightest exudation, the mucous membrane of the fauces is often seen to be red and somewhat swollen. After the pseudo-membrane is fully developed, it is of course impossible, without forcibly detaching it, to gain any idea of the condition of the mucous membrane beneath, and unquestionably very many of the descriptions given of extensive gan-

grenous ulceration of the fauces and pharynx, have referred merely to the changes in the pseudo-membrane due to its decomposition and the imbibition of sanious fluid.

In the vast majority of cases, the subjacent mucous membrane is not truly ulcerated, but is merely much congested and swollen, with an exco-riated and roughened appearance from removal of its epithelium, and occasionally presents spots of ecchymosis.

At times it is whitish, opaque, or unnaturally pale; while in other cases it is purplish or otherwise discolored. When the deposit is raised up, especially if it be of the firmer variety, it is often seen to be attached to the surface beneath by numerous small filaments, as though processes of the deposit passed into the mucous follicles.

Although these may be considered as the most usual conditions of the mucous membrane, it is undoubtedly true that in some cases extensive and deep ulceration, and even gangrene occur, exposing the muscular tissue of the pharynx, or even producing the destruction by sloughing of an entire tonsil gland.

The accident occurs much more frequently in some epidemics than others, as may be readily seen by a comparison of the accounts given by different authors of the anatomical lesions noticed in the epidemics they have respectively studied.

The submucous tissue is often œdematous, infiltrated with bloody serum, or is the seat of an interstitial exudation of lymph. In some cases the œsophagus and the muscular tissue around the fauces and pharynx are congested and infiltrated.

When croup ensues, the mucous membrane of the larynx and trachea is more or less swollen and congested, and, according to West, presents distinct erosion of its surface, with small ulcers about the edges of the glottis, in a larger proportion of cases than ulceration is met with in the fauces. M. Isambert¹ suggested that this condition might serve to distinguish diphtheritic from idiopathic croup; but West has met with precisely similar ulceration of the mucous membrane of the larynx in cases of primary croup, and is disposed to regard its presence or absence as mainly dependent on the rate of progress of the disease towards a fatal termination.²

SEAT OF THE EXUDATION.—The pseudo-membranous deposit is usually first seen upon the tonsils and soft palate, and in some cases is limited to these parts throughout the whole course of the case.

Frequently, however, the exudation spreads and coats the pharynx more or less extensively, or extends into the posterior nares, or downwards through the larynx into the trachea and bronchi, or more rarely into the œsophagus.

It is rare for any exudation to occur on the mucous membrane lining the cheeks, or upon the gums, though according to some authors, as Hutchinson,³ Trousseau, and Bouchut, ulcerative stomatitis is in reality buccal

¹ Arch. Gén. de Méd., March and April, 1857.

² Diseases of Children, 4th Am. ed., 1866, p. 356.

³ Med. Times and Gaz., March 19th, 1859.

diphtheria. The epiglottis is at times covered with a pseudo-membranous deposit, so as to become swollen, rigid, and almost immovable, and hence partially obstructing, without being able to protect, the entrance into the larynx.

The tendency for the exudation to extend into the nasal passages varies much in different epidemics, and when present, almost always betokens the great gravity of the case.

According to Bretonneau, the exudation occasionally begins in the nares and extends thence in so insidious a manner as readily to escape detection.

We will discuss more fully the questions relating to the extension of the exudation into the larynx under the head of diphtheritic croup.

The diphtheritic pseudo-membrane is not, however, limited to these mucous surfaces, but is occasionally seen, and especially in very severe cases, to form upon the mucous membrane of the vulva or of the anus.

It is, moreover, a most significant fact in regard to this affection, that any portion of the external cutaneous surface which has been denuded of epidermis, may become the seat of this deposit, and that in some cases the pseudo-membranous formation is even limited to the skin, constituting the so-called external or cutaneous diphtheria. So far, however, from the attending constitutional symptoms being less severe in the external than in the ordinary form, the tendency to deposit upon the cutaneous surface usually presents itself in cases of a typhoid adynamic type.

It appears, indeed, that this pseudo-membrane may occur at any point of the body to which the atmospheric air has access; but it has never been noticed on parts which are removed from its influence.

Notwithstanding these apparently distinctive features of the diphtheritic deposit, it is impossible by mere ocular or microscopic examination to distinguish it from the pseudo-membranous deposit in cases of ordinary scarlatinous angina.

It is more, therefore, in the peculiar constitutional disturbance that we must look for the specific nature of diphtheria, than in the presence and characters of the false membranes.

Bouchut and Dubrisay¹ have estimated the proportion of blood-corpuscles in diphtheria, employing the method of Hayem. They have found that the number of white corpuscles is considerably increased, while that of the red globules is diminished. The increase of white globules varies in direct proportion with the severity of the disease.

We have already alluded to the fact that in fatal cases of grave epidemic diphtheria, numerous micrococci have been found by Wood and Formad in the blood, and even developing within the white corpuscles.

The *Submaxillary Glands* are almost always enlarged, though they rarely acquire the enormous size and peculiar brawny induration so often noticed in scarlatina. It is, moreover, very rare for this condition to terminate in suppuration of the gland.

The *Heart* has been found, by Hillier², in a state of fatty degeneration in two cases, and by Bristowe (*id loc.*) in one; all of which were rather

¹ Arch. Gén. de Méd., October, 1877, p. 484.

² Diseases of Children, Am. ed., 1868, p. 154.

chronic. We have observed marked granular degeneration of the cardiac fibre in several instances, however, where the disease had been of a violent and rapidly fatal form. In some instances where symptoms of endocarditis were present during life, the auriculo-ventricular valves have been found in an incipient stage of inflammation. (Bridger¹.) We have also observed in several cases, in at least one of which a valvular murmur was heard during life, incipient inflammation of the mitral valve, with rows of minute, delicate, beadlike vegetations fringing the free borders of the leaflets. In other cases, pericarditis has been developed during the course of diphtheria. We desire to call particular attention to these cardiac complications, on account of their great importance as influencing not only the prognosis of the attack itself, but possibly also the subsequent development of chronic cardiac disease.

Heart-clots of large size and firm consistence, evidently of ante-mortem formation, are also found in a certain number of cases where death has been preceded by peculiar signs of circulatory embarrassment.

The *Lungs* are not rarely found inflamed and consolidated to a greater or less extent. In other cases the exudation is found penetrating deeply into their structure, filling the smaller bronchial tubes, and the lung itself is in parts collapsed or carnified.

Bouchut speaks of having seen small apoplectiform patches, similar to those which precede the so-called metastatic abscesses in pyæmia.

The *Kidneys* are at times quite healthy; in other cases, however, they have been found congested, and the renal epithelium granular and detached, so as to distend the tubules, which also contain fibrinous casts (inclosing granules of hæmatin, blood-corpuscles, or a few altered epithelial cells). (Hillier (*loc. cit.*), Greenhow,² etc.)

The *gastro-intestinal canal* presents no lesions of importance; in a few cases enlargement of the solitary glands of the lower part of the ileum has been noted.

Secondary Form.—When diphtheria appears in the secondary form, the mucous membrane is more violently inflamed. It is of a deep red color, rough, and very much thickened and softened. The tonsils are large and soft, uneven, and often infiltrated with pus. In addition, the mucous membrane is far more frequently and seriously ulcerated in this form than in the primary. False membranes are almost always present, generally on different portions of the fauces, and more rarely over their whole extent. They are generally rather soft and thin, of a whitish, grayish, or yellow color, dispersed in fragments and easily torn.

The inflamed parts are usually bathed in a purulent fluid. The sub-maxillary glands are large, red and soft; and, in addition, there may be found various lesions of other organs, due to the primary disease, in the course of which the diphtheritic angina has been developed.

SYMPTOMS.—Diphtheria occurs both in a sporadic and epidemic form; it also presents itself either as a primary or secondary affection. The symptoms of this latter form are, however, so involved with the symptoms

¹ Med. Times and Gaz., Jan. 1864, p. 201; and Brit. Med. Jour., Oct. 22d, 1864.

² On Diphtheria, New York, 1861, p. 160.

of the disease in the course of which it is developed, that it seems desirable to consider it in connection with them severally.

It would appear that, under the influence of widespread epidemic influences, the symptoms of diphtheria have of late years presented a higher degree of severity. Owing to the fact, however, that, until the description of diphtheria by Bretonneau, it was confounded with anginose scarlatina, with ulcerated sore throat, etc., it is extremely difficult to form a correct comparison between the disease as we are now familiar with it, and as it undoubtedly occurred in former years.

In a strictly systematic discussion it might be well to divide diphtheria into a mild form, which would include most sporadic cases and many of the epidemic ones, and a severe form, under which head would be comprised all cases distinguished by a high degree of constitutional disturbance. For practical purposes, however, it is sufficient to give a description of the ordinary course of the disease, dwelling upon some of the most important symptoms, and alluding to the chief peculiarities which at times present themselves.

LOCAL SYMPTOMS.—*Examination of the Throat.*—The onset of diphtheria is often very insidious; so that our attention may not be called to the throat by any complaint of the patient, even when a considerable amount of exudation is already present.

If the throat be examined, however, on the first day of the disease, the exudation may often be found even at that time, though it is sometimes not found before the second day. The fauces generally present more or less swelling and redness prior to the appearance of the false membrane, which almost always shows itself first on one of the tonsils only, in the form of whitish or opaline spots, like coagulated mucus, which soon run together and extend over the whole gland, and then to the soft palate and pharynx, though it sometimes remains limited to the tonsils and soft palate. A little later in the attack the plastic deposit exists in the form of layers of greater or less extent; it has lost its transparency, become firmer in consistence, thicker, and changes from a white to a yellowish-white or lardaceous, and sometimes grayish color.

The breath in this case is offensive, but not fetid; and there is but little salivation.

When, in favorable cases, the disease is left to pursue its natural course, the pseudo-membrane becomes thinner, assumes a grayish tint, and falls off about the sixth or seventh day. When, on the contrary, topical remedies are applied to the throat, the membrane is often detached after one, two, or three days, but may be reproduced several times before the conclusion of the case.

In some unfavorable cases, on the contrary, even though the exudation may disappear more or less completely from the pharynx, it extends downwards into the larynx, and we have true croup developed, which but too often proves fatal in spite of all remedies.

In more violent cases, the pseudo-membrane, about the time that it begins to be detached, assumes a grayish or blackish color, and hangs in shreds from the surfaces to which it was attached. The fauces, under these cir-

cumstances, present a gangrenous aspect, the mucous membrane having an appearance as though it were falling off in sloughs; the breath is extremely fetid, and there is more or less abundant salivation, or in some cases an expulsion of sanguinolent fluid.

There can be no doubt that it was from a misconception of such cases as these, that the titles of gangrenous and putrid sore throat arose.

As the exudation disappears from the pharynx, the swelling of the parts affected gradually subsides. The mucous membrane, from which the plastic matter has just fallen, is more or less injected and red; the tonsils and soft palate are sometimes found to be reduced below their natural size.

Even when the throat affection is very severe, there is not often so much difficulty in opening the jaws nor in deglutition as is met with in scarlatina.

The *submaxillary glands* are almost always enlarged and slightly painful to the touch, about three or four days after the appearance of the pseudo-membrane. The enlargement is usually greatest on the side where the inflammation of the fauces is most intense. The surrounding cellular tissue shares in the inflammation, so that the swelling is often very great, and impedes the movements of the jaw; it is rarely, however, save in very bad cases, so hard and painful as the corresponding swelling in scarlatina.

Pain and Difficulty in Deglutition.—There is sometimes no complaint of pain in the throat, although, even at the outset, swallowing is usually somewhat difficult and painful, and pressure behind the angles of the jaw causes a moderate degree of suffering. In some cases, especially of the sporadic sthenic form, the earliest symptom may be excessive pain on swallowing.

As the pseudo-membranous exudation increases, and the submaxillary glands become swollen and tender, deglutition becomes more difficult and painful, and, at times, attempts to swallow fluids are followed by cough and the return of the fluid through the nostrils.

In cases where the false membranes decompose and acquire a gangrenous aspect, and typhoid symptoms are present, the pain and difficulty in swallowing, if they have existed, are apt to disappear.

VARIETIES DEPENDING UPON EXTENSION OF THE EXUDATION.—1. *Croupal Variety.*—It would be a matter of much interest to determine in what proportion of cases this complication may be anticipated, and whether there be any definite and constant relation between the amount or character of the exudation in the pharynx and its extension to the larynx. As yet, however, no general conclusions can be arrived at in regard to any of these points. The frequency of its occurrence varies much in different epidemics, the proportion varying from one or two per cent. to as high as fifty per cent. of all the cases. Indeed, as in an epidemic referred to by Trousseau, the disease may, in almost every instance, assume a primary laryngeal form.

As might be expected from the considerations presented under the head of croup, this complication occurs more frequently and is much more fatal in children than in adults.

It is a well-recognized fact that true diphtheritic croup is nearly always preceded or accompanied by pseudo-membranous exudation in the fauces or pharynx, but the amount of deposit in these latter places may be extremely small, and yet be followed by extensive exudation in the air-passages; while, on the other hand, there is often copious deposit upon the pharynx in cases where the larynx does not become invaded.

No case, indeed, is free from the chance of this complication; it constitutes the chief source of danger in the mild variety, and yet is occasionally met with as the immediate cause of death in the most malignant attacks.

The pseudo-membrane is quite frequently found, in cases where the air-passages have become involved, to extend through the larynx and trachea, as far down as the tertiary bronchi, or in some instances, even to their finest divisions.

In this respect diphtheritic croup does not differ from primary croup, unless it be, indeed, that it seems to be more frequent in the former for the exudation to extend to the smaller bronchial tubes.

We have seen that there is no essential difference in the condition of the mucous membrane beneath the deposit in the two affections; and that they are equally liable to be associated with inflammatory conditions of the lungs.

Unless, therefore, the more highly corpuscular character of the exudation in diphtheria constitutes a ground of distinction between these two forms of croup, it seems difficult to establish a diagnosis between them on merely anatomical grounds.

When diphtheritic croup is secondary, appearing in the course of measles, scarlatina, or other general disorders, the conditions found after death in the larynx are much the same as in primary diphtheritic croup. The mucous membrane here, however, as in the fauces, is usually more intensely inflamed, and is more frequently ulcerated.

The possibility of the occurrence of croup should never be lost sight of, and every case should be treated as though it tended to invade the larynx. It is especially important to detect the very earliest signs of the approaching danger, since its onset is frequently extremely insidious.

If violent cough is excited by attempts to swallow liquids, it usually indicates that the epiglottis is inflamed, and the seat of pseudo-membranous exudation, which impedes its movements and thus allows the fluid to pass into the larynx. The extension of the exudation to the larynx is indicated by the cough acquiring a rough croupy sound, though it often has not the loud clangor of ordinary croup; the respiration becoming sibilant, and the voice weak and hoarse.

When the false membrane in the larynx is fully developed, the voice is almost or quite extinct, and the cough, losing its croupy character, becomes stifled and less frequent. The respiration is now peculiar; there is constantly a certain degree of dyspnœa, as shown by the frequent labored breathing, but there are, in addition, paroxysms of suffocation, induced by spasm of the laryngeal muscles, during which the dyspnœa is frightful, and

attended with tossing of the whole body and the most violent efforts at inspiration.

Death may occur during one of these paroxysms; but usually they subside and are followed by intervals of comparative ease, soon interrupted by the recurrence of the same alarming phenomena.

The intervals become more and more brief, and finally the patient sinks into a comatose condition, and dies with all the symptoms of asphyxia. If, during the violent efforts at respiration which attend these paroxysms of dyspnœa, or owing to the action of remedies, portions of the exudation are dislodged and coughed up, the most urgent symptoms are often immediately relieved. It is, however, but a deceitful repose, for in most cases the pseudo-membrane re-forms, and the recurrence of the croupy voice and sibilant respiration announce that the danger of suffocation is again imminent.

In favorable cases, however, either when the membrane does not re-form, or when it is dislodged as often as formed, recovery may occur; the paroxysms of dyspnœa recur at lengthening intervals, and finally disappear; the cough becomes gradually more soft, and fragments of pseudo-membrane, mixed with muco-purulent fluid, are discharged; the voice returns, and the capillary circulation becomes re-established.

In some cases, where the exudation has extended through the larynx and trachea deeply into the minute bronchial tubes, there is an absence of marked croupal symptoms, and death occurs slowly, after extreme dyspnœa and oppression of the chest, with all the symptoms of deficient aeration of the blood. These cases occur more frequently in adults than in children on account of the larger size of the larynx in the former.

The reader is referred for a more full account of this condition to the article on pseudo-membranous laryngitis.

2. *Nasal Variety*.—We have already mentioned that Bretonneau states that the disease occasionally begins at the nares, and extends thence in a most insidious manner. More frequently, however, the affection of the nares is consequent upon an extension of the exudation from the pharynx.

This complication is second in gravity only to the occurrence of croup. It impedes still further the already obstructed respiration, is attended with a foul acrid discharge from the nostrils, and, in addition, experience has shown that it is usually a sign of great malignancy in the case. According to Trousseau, the result is almost always fatal, the blood-poisoning being marked, as shown by the great alteration in the physical properties of the blood, the proneness to hemorrhage, the waxy pallor of the skin, and the ultimate fatal termination by syncope.

The detection of this complication in its incipient stage is therefore of the highest importance, and Bretonneau (5th memoir, *Syd. Soc. Trans.*, pp. 196, 197) has laid down the most minute directions for its recognition at this stage. If the patient present any evidences of disease of these passages, as a slight snuffling or coryza, during the prevalence of diphtheria, the finger should be placed behind the angle of the lower jaw, below the lobe of the ear, and thence passed down the side of the neck, and if

swelling of the cervical glands be noticed, it renders it probable that there is false membrane in the nares.

If, further, the upper lip be found reddened exclusively under one nostril, and that on the side of the glandular swelling, or if the swelling exists on both sides, but unequally, and if the lip is correspondingly reddened, the probability that there is nasal diphtheria is converted into a certainty, since ordinary coryza, acting equally on both nostrils, produces equal redness of both sides of the upper lip.

3. *Cutaneous Diphtheria*.—It is one of the characters of diphtheria which entitles it to be regarded as a blood disease, that different and distant parts are apt to become affected simultaneously or consecutively with the peculiar inflammation and exudation. We find, indeed, that in many cases of diphtheria there is a tendency to the formation of pseudo-membrane upon any portion of skin denuded of its epidermis.

This tendency varies greatly in different epidemics; according to our experience it is of rare occurrence in this city. It was, however, noticed by Bard nearly a century ago, and has been made the subject of special study by Bretonneau and Trousseau.¹ The pseudo-membrane forms upon any blistered surface; upon leech-bites; upon excoriations; in fissures, as behind the ears, or at the angles of the mouth; or on the outlets of the vagina and rectum.

The part that is to be the seat of pseudo-membranous deposit becomes surrounded by an erysipelatous redness; it is painful, exudes an abundant fetid serous fluid, and soon becomes covered with a grayish false membrane. This deposit gains in thickness from beneath; and, at the same time, extends in every direction, by the development of vesicles in the neighborhood, the bases of which become the seat of diphtheritic deposit.

The layers of membrane, bathed in the fetid serous fluid, soon change color, decompose, become horribly offensive, and impart the appearance of true gangrene.

Trousseau has observed this cutaneous exudation in cases where no affection of the throat existed, and has clearly established the identity of these various forms of diphtheria by facts collected in an epidemic in the neighborhood of Orleans, where the disease in some persons presented its ordinary features, while in others the exudation occurred on the vulva, on blistered surfaces, on the hairy scalp affected with favus, or upon ulcers.

The constitutional symptoms which accompany cutaneous diphtheria are usually extremely grave and adynamic.

GENERAL SYMPTOMS.—In the mild form of this disease the invasion is often highly insidious; there is usually fever, but the strength and appetite are not much disturbed at first. There is at the same time, in some, but not all cases, pain in the throat, which may or may not be accompanied by difficulty of deglutition. Both these symptoms are, however, often very slight, or they may be entirely wanting, a fact with which the practitioner should be well acquainted, as this absence of local symptoms by which to explain the cause of the sickness, gives to the disease, in some

¹ On Cutaneous Diphtheria, Arch. Gén. de Méd., 1830 (et loc. ante cit.).

instances, a remarkably insidious character which may well mislead. In one fatal case, at three years of age, that came under our notice, there were neither complaints of pain, nor difficulty of swallowing, so that the parents had not the least suspicion of the throat being the seat of disease, though we found it violently inflamed, and covered with deposits of thick false membrane in points. On another occasion, we were called to see two children who had been sick for four days with slight fever, languor, and loss of appetite, but who were not thought to be seriously ill. We found them laboring under extensive pseudo-membranous angina, with the early symptoms of croup. They both died a few days later of croup. The symptoms, prior to the development of the croup, had been so mild in both cases as to cause no alarm, and yet the anginose disease had evidently been progressing insidiously for several days. We attended, a few years since, for three days in succession, a boy who was attacked suddenly with vomiting and slight fever, loss of appetite and languor, and whom we supposed to be suffering from mere gastric irritation. His only local symptom was pain in the chin, and this was not reported to us until afterwards. The mother chanced to look into his throat, and, finding there some whitish spots, sent us word. We found him with very considerable membranous exudation, which was fortunately prevented from extending into the larynx by proper treatment. Quite frequently have we been called to see children attacked with croup, and on finding the fauces thickly covered with exudation, have been told that the patient has been ailing for near a week before with languor, slight peevishness, loss of appetite, and some little pain in the throat. To this point, the strangely insidious character of the anginose symptoms in the early stage of many cases, we cannot too strongly invite the attention of the reader. It is one of the very greatest importance, since at that time, above all others, ought the case to be placed under proper treatment.

It is to this class of cases that, owing to the trifling character of the constitutional symptoms, the name of diphtheroid sore throat is sometimes applied, though inaccurately, since it is calculated to create doubt as to their essentially diphtheritic nature.

It has been, on the other hand, stated, that, during epidemics of diphtheria, cases occur which present the usual general symptoms, with some difficulty of swallowing and swelling of the cervical glands, but in which no pseudo-membrane is formed, the fauces being merely of a dark-red color, with swelling and elongation of the uvula, and sometimes tumefaction of the tonsils.

Such cases are rarely fatal, and, as a rule, yield readily to the ordinary treatment for diphtheria.

In addition to these mild cases, in which the chief danger is from the extension of the exudation into the larynx, the disease in many instances, and especially under the influence of epidemic causes, assumes a grave form, in which the danger depends not upon an accidental extension of inflammation, but upon the essential alteration of the blood, and the condition of the entire system.

In these cases also the onset may be insidious, though it is often pre-

ceded for a short time by general malaise, indisposition to play on the part of children, and to exertion on that of adults, and slight swelling of the cervical glands, and pain on deglutition.

Whether these prodromes have been present or not, a more or less marked chill ushers in the febrile action, which is often quite intense for a few days; so that, when the throat affection is decided, a doubt may exist for a short time whether the approaching attack is one of scarlatina or diphtheria. The fever, however, soon subsides almost completely, sometimes indeed leaving the surface pale and cooler than natural. The pulse may remain frequent, but is weak and compressible; and the general symptoms are all characteristic of deficient vital force.

There is not usually any marked mental disturbance after the second day, the child being intelligent, though dull and indisposed to pay attention to anything.

There are but few symptoms of digestive disorder; the appetite, which is often retained for the first day or two, soon diminishes, and the child often becomes unwilling to take any food, partly from the pain caused by the efforts to swallow, partly from complete anorexia. There is rarely any vomiting, unless provoked by remedies; and the bowels, though usually torpid, occasionally incline to be loose. The urine is rather scanty, quite frequently albuminous, and upon microscopic examination is found to contain renal epithelium and casts from the renal tubules. This symptom will be again and more fully alluded to among the complications.

At the same time, the submaxillary glands enlarge, and the fauces assume the appearances we have already described. There is a great increase in the secretion of saliva, which often dribbles quite profusely from the mouth, and is apt to be offensive, though rarely fetid. In many cases there is in addition a discharge from the nostril, which becomes acrid and offensive when there are false membranes in the nasal passage.

The voice is commonly obscured and nasal, or somewhat hoarse, even when the larynx is not involved.

Cough sometimes exists, and may have a slightly ringing spasmodic character, due to mere irritation of the larynx, though it usually resembles in sound that produced by the action of hawking, rather than a common cough.

In a very small proportion of the cases, an eruption, resembling that of scarlatina, appears at irregular periods in the course of the disease. It appears, however, that this eruption lacks the punctated appearance of the scarlatinous rash; does not appear at any fixed day of the disease; is irregular in its progress, and is not followed by desquamation.

The reports of it are, however, scarcely numerous or accurate enough to enable us to say positively that intermingled cases of scarlatina have not been mistaken for diphtheria, or that the two poisons may not have been acting jointly.

The further course of these cases varies widely. If the result is to be unfavorable, the depression and loss of strength increase rapidly; the surface grows pale or sallow, and is below the natural temperature; the pulse becomes exceedingly frequent and feeble; the fauces assume a gangrenous

appearance from decomposition of the false membrane; the swelling of the cervical glands increases, and the patient often refuses to make the effort to swallow, though deglutition is still generally possible; there is a constant fetid discharge from the mouth and nostrils; the breath is horribly offensive; and death ensues amid the most profound prostration. Or, at a much earlier period of the disease, the fatal event may be precipitated by the extension of the exudation to the larynx.

We must also allude to the occasional occurrence of sudden death, even in cases not of the gravest type. The dreadful accident appears to result from paralytic failure of the heart's action, or, less frequently, from the sudden formation of a heart-clot; and the fact that it may occur, should call for the most careful attention to the avoidance of all exertion on the part of the patient.

If, on the other hand, the case tends towards recovery, the false membranes become detached and thrown off, the strength improves, the pulse becomes fuller and stronger, and the appetite returns. Even in advanced convalescence, however, there is serious danger, as will be seen more fully hereafter, of the occurrence of troublesome or even fatal sequelæ.

In a still more severe group of cases than those above sketched, the symptoms are of the most asthenic or malignant type.

In these cases the anginose affection, though it may be severe, rarely attracts much attention. The pseudo-membranes in the fauces are soft and pulpy, and, when examined microscopically, highly corpuscular and granular; they soon decompose, and become discolored by the blood which exudes from the mucous membrane. There is, moreover, a strong disposition for the exudation to extend to the posterior nares, or to appear on various portions of the external cutaneous surface. The breath and the discharge from the mouth and nostrils are indescribably fetid. In some cases true ulceration, and even gangrene, of the fauces occurs. There is, however, less pain complained of, and less indisposition to swallow than in many lighter cases, owing probably to the depression of the nervous centres from the poisoned state of the blood. There may be high fever during the first few days, but this soon disappears, and is replaced by a deadly pallor of surface; extremely feeble, running pulse; and at times low muttering delirium.

Passive hemorrhages from the nostrils, mouth, rectum, or other mucous passages, are of frequent occurrence.

The result in these cases of profound diphtheritic infection is almost invariably fatal; death resulting quietly from pure exhaustion, without the development of any complications.

The *duration* of diphtheria varies considerably. Ordinary cases recover in about seven, eight, or nine days, whilst more severe attacks are often protracted until the end of the second week.

It is impossible, however, to say that the disease has actually run its course in this time, since there are sequelæ which may appear during advanced convalescence, and retard the recovery even for many weeks.

On the other hand, in fatal cases, death may occur from croup, as early as the end of the second day; though usually the larynx does not become

implicated under five or six days, and this accident may occur so late as the twelfth or fourteenth day of the attack.

In extremely malignant cases, death may also occur during the first few days. On the whole, however, it may be said that the majority of deaths from all causes occur in the period between the sixth and twelfth days. When death results from one of the sequelæ, either disease of the kidneys or paralysis, it may be deferred for weeks, or even for several months.

PROGNOSIS.—In cases of ordinary severity, when the patient is seen early, and the disease remains limited to the pharynx, the result is usually favorable; though no case, not even the mildest, is free from danger, either of extension into the larynx or bronchial tubes, of exhaustion, or of the supervention of some complication, such as endocarditis, or the formation of heart-clots. If, on the contrary, the exudation extends to the nasal passages, the prognosis is more unfavorable; and when the larynx becomes implicated, the prognosis is exceedingly grave; if the disposition to the production of false membrane spread to the skin, rectum, or vulva, the prognosis is also very grave, and death generally occurs in a state of profound adynamia.

If any other signs of unusual malignancy are present, such as abnormal slowness, or great frequency and smallness of pulse; marked prostration with pallor and coolness of the surface; great tumefaction of the cervical glands; abundant pseudo-membranes, pultaceous and rapidly decomposing; hemorrhages from various mucous surfaces; acrid, fetid discharges from the mouth or nostrils; intense and persisting albuminuria, with diminution of the amount of urea excreted; the prognosis is, of course, much more unfavorable.

It must be remembered, however, that no one of these symptoms, nor even any combination of them, is necessarily of fatal import; that cases are often rescued apparently from inevitably impending death; and that, however threatening the symptoms may be, it is our duty, in this disease even more than in many others, to persevere to the very latest moment in the judicious application of suitable remedies.

It is as yet impossible to arrive at any plausible estimate of the average mortality of diphtheria, so widely does the proportion vary in different epidemics. Neither sex nor temperament appears to have any influence upon the result; but extreme youth undoubtedly renders the prognosis much more grave.

The prognosis in the secondary form of diphtheria is also more unfavorable than in the primary.

DIAGNOSIS.—We have already sufficiently dwelt upon the general symptoms and local signs which enable us to detect diphtheria, in every instance, after the disease has fully developed itself.

In examining the fauces in the early stage of the affection, it is well to remember that in simple angina, the crypts of the tonsil-glands occasionally become so distended by their secretion as to present the appearance of small, round, and slightly elevated whitish patches, which might readily impose upon a hasty observer for pseudo-membranous deposits.

In regard to the value of the peculiarities upon which a differential diagnosis between diphtheritic croup and idiopathic primary membranous croup is so frequently based, we have fully expressed our opinion in the article on the latter disease, to which we would refer the reader.

Diagnosis from Scarlatina.—The great resemblance which at times exists between the anginose symptoms of scarlatina and diphtheria has led some authors to suggest that they are identical diseases, and the following further points of resemblance have been adduced: the two affections prevail frequently simultaneously in the same region, and even in the same family; in certain cases of diphtheria, a rash, very similar to that of scarlatina, is said to appear; and the urine, in diphtheria, is frequently albuminous. That this similarity is, however, more apparent than real, is evident from the following considerations:

1. Although in some epidemics of diphtheria a rash is said to have been occasionally noticed, its occurrence is at most the rare exception, instead of the almost invariable rule, as in scarlatina; it differs, too, from that of scarlatina, in appearing at irregular periods, in being partial, appearing suddenly in patches, not deepening gradually in intensity, and in being of a uniform erythematous redness, without the punctated appearance peculiar to the scarlatinous eruption.

2. The albuminuria of diphtheria presents these distinctive features as compared with that of scarlatina, that there is not always any diminution in the amount, nor any constant change in the character of the urine when it is present; that it occurs in the early part of the attack, and increases as the disease approaches its height, or may disappear suddenly, even in the early part of its course; that although usually noticed in severe cases (and probably a very unfavorable symptom), there seems to be no necessary connection between the urine becoming non-albuminous and the disease assuming a milder type.

3. There is a wide difference in the sequelæ which succeed the two affections; dropsy scarcely ever following diphtheria, while various paralytic phenomena, which are rarely noticed after scarlatina, are of frequent occurrence. It is very much more common, also, to have suppuration of the glands of the neck after scarlatina.

4. In the same way, endocarditis, though it has recently been noticed in a few cases of diphtheria, is much more frequent in scarlatina.

One of the most positive proofs of the essential difference of these two affections is the fact, attested by universal experience, that they exercise no protective power whatever against each other, and that individuals whose systems are protected against a second attack of scarlatina, are fully as likely to contract diphtheria as those who have never suffered with either of these diseases.

It may also be added that second attacks of scarlatina are very rare, while they seem to be much more common in diphtheria.

It seems evident to us, therefore, that in the present state of our information upon this subject, scarlatina and diphtheria must be regarded as entirely distinct affections, although presenting quite numerous points of singular resemblance.

COMPLICATIONS AND SEQUELÆ. — *Albuminuria*. — We have already briefly alluded to the peculiarities of the albuminuria of diphtheria, but the importance of the symptom merits a more full discussion.

The occasional presence of albumen in the urine in cases of diphtheria was first noticed by Mr. Wade in 1857, who also found associated with the albumen, tube-casts and renal epithelium. It was shortly afterwards recognized by MM. Bouchut and Empis¹ in thirteen out of fifteen cases; and since then has been found, in a varying proportion of the cases, by many observers in different epidemics.

The character of the urine when it contains albumen is not constant, but usually it is quite pellucid, of acid reaction, and apparently free from any deposit; although, on standing, both tube-casts and epithelium may settle to the bottom. The quantity also varies considerably, Hillier having found it much diminished, while, according to West and Wade, it frequently remains normal.

The amount of urea excreted is usually increased in diphtheria, and, according to Sanderson, the presence of albumen and tube-casts in the urine is not necessarily associated with any interference in its elimination, but this does not agree with the examination of others, who have found a diminution of the solid excreta when albumen was present.

The quantity of albumen varies much, being at times a mere trace, and again being present in large amount. The kinds of tube-casts noticed by Wade, and which are the ones usually found, were small, waxy casts; casts of a similar size, but granular, probably from commencing disintegration, and ordinary epithelial casts, and fibrinous flakes.

Albuminuria in diphtheria occurs at various stages of the disorder, in some cases even during the first few days. It not rarely comes on insidiously, and may manifest its presence by no peculiar constitutional symptoms. There can be, however, little doubt of the grave import of its appearance, though as yet its exact significance has not been accurately defined.

It is indeed true, that it has been found in large quantities in cases which have presented a mild character throughout (Sanderson); but on the other hand, Bouchut and Empis regard it as a highly unfavorable sign, coinciding with very great gravity of the disease; and Wade believes that the quantity of albumen is usually in direct proportion to the retention of effete material, and that indications of impairment of the renal function are almost constantly precursors of an unfavorable termination.

Hillier (*loc. cit.*), examined 38 very severe cases in regard to this point, and found albumen present in 33, 32 of which proved fatal, while of the 5 free from albuminuria, all recovered.

The albumen appeared in 1 case on the fourth day, in 3 on the fifth day, in 2 on the seventh day, in 5 on the ninth, and in 1 each on the thirteenth and nineteenth days. Usually the albumen disappears from the urine as the severity of the symptoms diminishes, but Bouchut has

¹ De l'Album. dans les Mal. Couenneuses, Compt. Rendus, 1859.

known it to persist after convalescence, and finally produce, as in Bright's disease, anasarca and hydrothorax.

Heart-clot.—The formation of coagula in the cavities of the heart during life has been noticed in many conditions of the system; and this terrible, because almost necessarily fatal accident, is now always dreaded in the course of several diseases, of which diphtheria is eminently one.

There have even been epidemics of an unknown nature, but where the only discoverable lesions have been enormous fibrinous concretions in the heart. Such epidemics have been recorded by Huxham, Chisholm, and recently by Armand.¹

The symptoms mentioned by these authors as significant of this accident are pain at the pit of the stomach; difficulty in respiration; extreme anxiety and restlessness; anxious expression and depression of spirits; slight, dry, and rather spasmodic cough; the face being at times livid, and the surface dry and inclining to be cool, with coldness of the extremities. The pulse was small and irregular, and, in some of Armand's cases, an abnormal murmur was detected in the heart; there was usually considerable dullness over the cardiac region; the respiratory murmur remained pure and quite full, and the chest normally resonant.

According to Robinson, the first observation of sudden death in diphtheria from the formation of heart-clot was made by Dr. Werner, of Linz, in Austria, in 1842; and the second by Winkler, in 1852.

In England, Dr. Richardson² appears to have been the first to call attention to the difference between these symptoms of embarrassed circulation and those of obstructed respiration, as met with in diphtheritic croup.

His account of the symptoms of the former condition agrees closely with that given above as to the coolness and almost marbly pallor of the surface; the moderate lividity of the face; the constant restlessness and intense anxiety; the feeble, quick, and irregular action of the heart, with a muffled character of the sounds, and in some cases an abnormal murmur. He also calls attention to a peculiar prominence of the anterior part of the thorax in very young children, which he believes to be strictly diagnostic of fibrinous obstruction.

In obstruction of respiration, on the other hand, the surface becomes livid, the veins turgid, and the muscles are often convulsed; the heart-sounds are clear, though feeble, and the breathing is the first to stop at death, instead of the circulation, as in the other case.

In three cases occurring in the practice of one of ourselves,³ in which we were able to diagnosticate the condition, death took place on the twenty-first, twenty-fifth, and twenty-eighth days respectively. In each case the local symptoms had given way and almost disappeared, and the children seemed to have entered upon convalescence, when slight but steadily increasing signs of circulatory embarrassment became perceptible, and after

¹ Des Concretions Fibrineuses et Polypiformes du Cœur, 1857.

² Med. Times and Gaz., March 8, 1856; British Med. Jour., Feb. 16, and April 7, 1860.

³ Dr. J. F. Meigs, Am. Jour. Med. Science, April, 1864, vol. xlvii, p. 305.

a few days' battling against the constantly increasing obstruction, each of the little patients died as though worn out by the unequal struggle.

In no case was there any evidence of any other organ being implicated; one of the cases was, however, complicated with albuminuria.

The pulse was not noted to be over one hundred; the cardiac sounds were unattended with murmur, but confused, indistinct, and seeming as though reduplicated.

There was no marked paralysis, but in one case partial paralysis, and in another marked muscular debility.

At the autopsy, in each case, the right side of the heart was full of clots, which were either dark-colored, with whitish spots, or yellowish-white throughout, quite firm, and adherent to the endocardium, and appeared to have been forming for several days. In one case, a clot in the left ventricle presented at its lower extremity a broken, irregular, uneven, and frayed or granulated appearance, as though the disintegrating process by which thrombi are broken up, had commenced in it. In none of the cases were there any evidences of endocarditis. The same accident has been observed during the past ten years by Dr. Barry,¹ Mr. H. Smith,² and Mr. C. R. Thompson,³ and many others, and in a valuable thesis published recently⁴ by Dr. Beverley Robinson (now of New York), ten cases are fully described, in at least five of which the ante-mortem formation of clots occurred.

The symptoms which he deduces, from a careful analysis of his own and the other recorded cases, as indicative of this condition are: coolness of the extremities, pallor of the face, prostration, anxiety, agitation, and peculiar intense dyspnœa; associated with a feeble pulse, dull, weak, and veiled heart-sounds, and frequently with the signs of emphysema of the lungs.

Most of the cases have occurred in young subjects, and the clot has formed late in the course of the disease, or even after convalescence has begun. The cause of this deposition of fibrin is not very apparent; in our article on this subject, already referred to, it was suggested that the coagulation might depend upon some peculiar change in the tissue of the endocardium, analogous to that which gives rise to the diphtheritic exudation on mucous surfaces.

No such alteration has, however, as yet been detected, and Dr. Richardson, to whom the profession is so much indebted for his investigations upon the coagulation of the blood, attributes it, in this case, to a deficiency of the volatile agent which retains the fibrin in solution, together with an actual increase in the amount of the fibrin of the blood, this combination producing the most favorable condition possible for fibrinous deposition.

Endocarditis.—Although in the above cases no lesion of the endocardium has been found, inflammation of this membrane has, as already

¹ British Med. Jour., 1858.

² Med. Times and Gaz., Dec. 17, 1859.

³ Med. Times and Gaz., Jan. 7, 1860.

⁴ De la Thrombose Cardiaque dans la Diphthérie. Paris. 1872.

stated (page 884), been quite frequently noticed in diphtheria. It has usually appeared late in the course of the disease, and has been attended with pain in the præcordia, frequent pulse, hurried respiration, an anxious countenance, with in some cases a systolic murmur. In fatal cases, there was found a roughened, reddened, thickened appearance of the valves, as if due to interstitial deposit. In some cases, also, a granular or fatty degeneration of the muscular fibres of the heart has been observed, as by Bristowe, Hillier, Robinson, ourselves, and others.

Paralysis.—One of the most frequent and important, and certainly the most peculiar of the sequelæ of diphtheria, is the occurrence of paralysis. It originally attracted the attention of MM. Trousseau, Lasèque, and Faure, under the form of difficulty of deglutition, and a nasal character of the voice; but since then it has been observed in the most varied forms and degrees, affecting both general and special sensation and the power of motion. In most cases, every trace of the primary disease has disappeared before any paralysis is noticed; the patient sleeps, eats, and digests well, yet many cases emaciate, and there is often marked pallor of the surface. In many instances, also, especially in children, there is great irascibility or irritability of temper.

Most frequently a nasal character of voice and regurgitation of liquids through the nose are the first symptoms to call attention to the disease, though these may be preceded by some slight difficulty in articulation, or by alteration of the sense of taste at the back of the tongue. On examining the fauces the soft palate is found hanging relaxed, and, if it be pricked, there is no contraction of it, nor does it give the patient pain.

At times but one side is paralyzed, and the uvula is drawn towards the sound side. The affection may extend no further than the fauces, and soon disappear; or may advance, the eye usually becoming next affected, following the throat affection, and preceding any paralysis of the limbs. The impairment of vision is rarely of long duration, lasting from a few days to two months, and is of every grade, from mere inability to read fine print to perfect blindness.

Greenhow has noticed that the pupils become dilated, and act sluggishly under the influence of light, for a day or two before the sight becomes sensibly impaired, and may remain so for a time after sight has been regained. He has also observed that patients who were unable to read with unassisted sight could do so with the aid of convex glasses, so that he attributes the impairment of sight to paralysis of the ciliary muscle and temporary loss of the adjusting power.

In addition to this want of accommodation, however, depending on paralysis of the ciliary muscle, Bouchut believes that there is in many cases, and especially in those who have had albuminuria, a serous infiltration* of the fundus of the eye, due to the anæmic condition of the blood, and which may impair the nutrition of the optic nerve, and even lead to its atrophy.

The following case, which came under our observation recently, at the clinic at the University of Pennsylvania, affords an interesting illustration

of the peculiarities of this form of paralysis. The ophthalmic examination was made by Dr. S. D. Risley, who has kindly placed the results at our disposal.

CASE.—Emma W., *æt.* 7 years, suffered with an attack of sore throat, the nature of which was not recognized by the physician in attendance. It was quite severe, was accompanied by marked swelling of the glands at the angles of the jaw, and compelled her to be confined to bed for a week or ten days. Soon after the convalescence began, it was noticed that her voice became altered, and that she occasionally regurgitated fluids which she attempted to swallow. Her general health improved, however, and in a few days she returned to school, which she was soon obliged to quit in consequence of rapidly increasing inability to read, on account of the print seeming blurred and "the letters running together."

Two weeks later, or about five weeks from the time of the first attack, examination of the eyes showed the fundus of both eyes entirely healthy. *O. D.*, $V = \frac{20}{xx}$. Acuteness of vision, as determined by Snellen's types, normal, and she can read Jr. No. 14 at 2½'. *O. S.*, $V = \frac{20}{xxx}$, and she reads Jr. No. 16 at 2'. *O. D.* emmetropic; *O. S.* hypermetropic = $\frac{1}{4}$ 8. With glasses $\times \frac{1}{12}$ (convex glasses with 12'' focus) *O. D.* reads Jr. No. 1 at 12'', *O. S.* at 14''. The pupils react promptly to light.

She was directed to wear $+\frac{1}{12}$ glasses for near work, and strychniæ sulph. gr. $\frac{1}{60}$ was ordered four times daily. This was in a few days increased to five times a day, and its use was followed by prompt improvement, so that in less than two weeks the power of accommodation was entirely restored.

Deafness may follow this amaurosis; then the lower limbs become affected, the patient becoming paraplegic, and next the upper extremities, then the muscles of the alimentary canal and bladder, causing impaction of the rectum with fæces and retention of urine, or the sphincters of these organs alone may be involved, and lead to involuntary discharges. Finally, the muscles of the trunk, including those of respiration, may become paralyzed, and in some very rare cases even the muscles of the heart are involved. It is stated that the paralysis of the extremities is never strictly unilateral. The paralysis is rarely confined to loss of motion, but, in a majority of cases, sensation is either much modified or lost; and indeed in some instances there has been no loss of motion, the sensory nerves alone being affected. In other cases the sensibility has been found exalted, or there has been in the same case hyperæsthesia in the upper, with anæsthesia in the lower extremities.

The paralysis, whether it be of motion or sensation, is progressive and gradual, even in the same set of muscles, and usually involves one limb before it extends to other parts. The mind, though often feeble and dull, acts correctly in most cases.

During the continuance of these phenomena, the appetite may remain good and digestion easy; but there are often marked evidences of the continuance of some morbid action in the economy. The surface is of an earthy, sallow hue, calorification is often imperfect, and the circulation is much depressed, the pulse being small, weak, and much reduced in frequency.

In some cases, indeed, the affection runs on to a fatal issue, usually consequent upon a failure of one of the vital functions of circulation or res-

piration. M. Faura has given a vivid picture of these sequelæ in their worst form, when the patient, paralyzed, indescribably prostrated, with imperfect speech and power of deglutition, impaired vision, imbecility of mind, œdema, and even gangrene of the extremities, finally dies in some fainting fit, or passes away almost imperceptibly.

The result of diphtheritic paralysis is, however, favorable in a large majority of cases; thus of 77 cases collected by Dr. Reynolds, but 9 were fatal. We have ourselves never met with a fatal result in a single instance, although a large number of cases, some of them of very severe character, have come under our observation. The duration is, however, more uncertain, varying from one or two weeks to several months, the mean duration being about a month.

It is as yet impossible to advance any satisfactory explanation of the cause of these grave paralytic sequelæ. They occur probably in one-fourth of all cases, in greater or less degree, and are noticed with at least equal frequency after mild as after severe attacks.

At first, indeed, the faucial paralysis was attributed to some such local cause as inflammation of the sheath of the nerves supplying these parts, and Greenhow still contends that the nerve affections bear some proportion to the local severity of the attack, the paralysis and anæsthesia being more complete on that side of the fauces which has been most severely affected by the primary disease; but we have been able to satisfy ourselves that this does not occur with any uniformity. Charcot and Vulpian have, however, demonstrated in a case of paralysis of the palate, lesions both of the palatine nerves and muscles; and Dejerino (*Gaz. Méd.*, 1877, No. 33), reports that he has found in three cases of diphtheritic palsy, signs of parenchymatous neuritis of the anterior roots of the corresponding nerves. It is difficult to determine whether such lesions are of constant occurrence in the ordinary cases which rapidly recover. It seems improbable, also, that in cases where widespread paralytic symptoms are present, which subsequently entirely disappear, any serious lesion of the nerve-trunks or of the muscles could have existed.

Nor in the occurrence of albuminuria necessary for the development of paralysis, since the urine is often quite normal throughout the entire course of cases, which are nevertheless followed by marked palsy.

The most plausible view we can entertain of the nature of these nerve affections, is that they are the direct effect of the diphtheritic poison, which while modifying the blood crisis, and so acting on the system at large, has an especial tendency to the nervous system; while at the same time, some of the local forms of the paralysis may be associated with lesions of the nerves and muscles of the part affected.

Ataxic Form.—In some cases, which are comparatively rare, the nerve affection does not constitute actual paralysis, but takes the form of locomotor ataxia. In such cases, the muscular force in the affected parts, usually the lower extremities, is not materially diminished, so that the patient can move them forcibly when he is lying down; but there is such a degree of incoördination in the motions communicated to them, that combined movements, even in the supine position, may become impossible.

It is, however, especially in walking, that this loss of coördinating power manifests itself; the gait becomes irregular, the patient falls if the eyes are closed, and the case presents all the characteristics peculiar to well-marked locomotor ataxia.

The first instance of this diphtheritic ataxia appears to have been observed by Jaccoud¹ in 1861; it was soon after noticed by Eisenmann;² and more recently a well-marked case has been reported by Dr. Gray,³ in a boy nine years old, following an apparently mild case of diphtheria. It is evident, also, as pointed out by Jaccoud, that a certain number of the cases which have been reported under the name of diphtheritic paralysis, have in reality been examples of locomotor ataxia, the paralysis having been only apparent. We have met with two well-marked examples ourselves, in both of which entire recovery followed. This diphtheritic ataxia is in all probability due to the same unknown morbid condition or dyscrasia, which causes the actual paralytic symptoms which are more frequently observed as sequelæ of diphtheria. It usually yields to the treatment recommended for the latter conditions, though in Gray's case death occurred, apparently from rapid loss of nervous power, seven weeks after the appearance of the nervous symptoms.

TREATMENT.—The treatment may be usefully considered under the two heads of local and general. Of late years, the importance of the latter has been more and more recognized as supreme, and, indeed, the utility of all local treatment has even been questioned on the ground that the throat affection is merely a local evidence of the constitutional disease, and that the disease rarely kills save by involving organs beyond the influence of such agents. We have, however, no doubt as to the very great importance of proper local treatment; although, on the other hand, we are not prepared to say, with some eminent authorities, as Trousseau, that topical applications are the most successful and important remedies in diphtheria.

The great objects to be held in view in the local treatment, are to favor the separation of the pseudo-membranes, and to prevent their extension from the fauces into the larynx and nasal passages.

LOCAL TREATMENT.—The most important of the local remedies are included in the lists of astringents and caustics.

Of these, nitrate of silver has properly been used more than any other substance for many years past, and is highly recommended by MM. Bretonneau, Valleix, Grisolle, Rilliet and Barthez, Trousseau, West, and many others.

It is employed both in solution and substance. The latter form is, however, open to the objections, that if the extent of the false membranes be at all considerable, the solid caustic can seldom be applied to more than a small portion of it, and that it is attended with the risk of slipping from the porte-caustic into the pharynx, and thence passing into the

¹ Les Paraplégies et l'Ataxie du Mouvement, p. 631, Paris, 1864.

² Die Bewegungs-Ataxie, Wien, 1863.

³ London Med. Times and Gazette, February 6th, 1869, p. 141.

stomach.¹ The solution is therefore generally preferred. M. Bretonneau advises its employment in the proportion of half an ounce of the salt to an ounce and a half of water; and West employs a solution of the strength of a drachm to an ounce.

We have usually made use ourselves of a solution of ten or twenty grains to the ounce, and have found it abundantly strong. It may be applied either by means of a piece of sponge fastened upon a proper handle, which is the best method, or a camel's-hair pencil, nearly as large as the end of the little finger. The application should be made once, twice, or even three times in the course of twenty-four hours.

Hydrochloric acid is also frequently employed, either pure or diluted with from one to ten parts of honey; the more dilute forms being used in the case of children.

It possesses the great advantage over the other mineral acids, that its caustic action does not extend much from the point of application, but is open to the objection of causing a white plastic exudation on any part of the mucous surface, not covered with false membrane, with which it may come in contact, which may lead the physician into error.

When the limits of the pseudo-membrane can be seen in the pharynx, following M. Bretonneau's advice, the acid may be used more concentrated, and the sponge, after being dipped into the acid and squeezed so as to be merely moistened, should be carried rapidly into the pharynx, and withdrawn after lightly cauterizing the surface.

When, on the contrary, the limits of the membrane cannot be seen, the acid should be more diluted, and leaving more of it upon the sponge, this should be passed down over the epiglottis and then pressed against the base of the tongue, by raising strongly the handle to which it is tied, in order to express a few drops upon the mucous membrane of the larynx. The cauterization is to be performed once or twice a day, according to the necessity of the case. For children under ten years of age, the sponge ought to be about half as large as a pigeon's egg. It is to be fastened to a piece of flexible whalebone, by making a crucial incision into it, introducing into this the end of the whalebone, and securing it with good sealing-wax, which is not acted upon by the acid as any ligature would be. When about to be used, the whalebone is warmed and curved into such a shape as will allow it to pass into the pharynx without touching the roof of the mouth. M. Valleix proposes that the sponge should be fastened to the whalebone with waxed thread, and that this should be covered with sealing-wax, to preserve it from the action of the acid. This would certainly be safer than the mere wax alone.

Applications of powdered alum, tannic acid, and chlorinated lime, are recommended by writers of high authority. In slight cases, in which the disease shows but little disposition to extend, such applications may answer very well; but when the attack is threatening, and especially when

¹ Dr. Geddings recommends, when it is desirable to use the solid nitrate, to reduce it to powder, and to roll the sponge probang, previously moistened with mucilage of acacia and squeezed, in the powder until a sufficient quantity adheres, and to apply it thus prepared to the diseased parts.

the exudation is spreading, we should neglect these minor remedies, and resort at once either to nitrate of silver, dilute muriatic acid, or the tincture of the chloride of iron. If, however, these powders are employed, they may be applied by means of a throat brush, or by causing a sufficient quantity to adhere to the forefinger of the right hand, and conveying it upon this to the diseased surfaces.

The astringent and caustic preparations of iron have lately been introduced in the treatment of this affection with much benefit. They cause the pseudo-membranes to contract and shrivel, and thus favor their separation, while, at the same time, they modify the action of the mucous membrane, and also tend, as does the sol. sodæ chlor., to correct the fetor arising from the putrefaction of the false membranes, and to prevent poisoning of the system by absorption.

The tr. ferri chloridi and the ferri perchloridum are among the best preparations, and may be applied either pure or diluted, several times in the course of twenty-four hours. Monsel's salt, in powder, has also been highly recommended by Beardsley, of Connecticut, and possesses the same mode of action, though somewhat more escharotic.

Carbolic acid, diluted with glycerin and water, applied by a mop to the throat, appears to possess almost equal virtue in causing the separation of the pseudo-membranes, and preventing their re-formation.

Various applications have also been recommended from the fact that they exercise a direct solvent power over the pseudo-membranes, and thus promote their removal. Among those which have been thus recommended are solutions of lime, potassa, and soda; solution of chlorinated lime; of chlorate of potash or soda; of permanganate of potash; of bromide of potassium; of pepsin; and of dilute lactic acid.

Dr. Jacobi (*Amer. Jour. of Obstet.*, May, 1868, pp. 13-65), has published an analysis of the relative value of these solvent applications. According to him, lime-water requires four to ten hours to thoroughly liquefy soft diphtheritic exudation; while for firm pseudo-membranes, it requires from thirty to seventy-two hours. Potash and soda, and their salts, act more slowly; and the one other application which he recommends as equally rapid in its action is a solution of bromine gr. j, bromide of potassium gr. j, in fʒvj of water.

We have carefully tested the latter solutions, as well as those mentioned above, and from the results of repeated tests, have concluded that lime-water is the most powerful in its solvent action upon pseudo-membranous exudations. We have frequently found, when fragments of firm white exudation have been placed in lime-water at a temperature even lower than that of the buccal cavity, that the exterior began in a very short time (half an hour) to undergo disintegration, and that the whole fragment was reduced in a few hours to a granular putrilage. It is, however, undoubtedly true that this effect will be produced with very different rapidity upon different specimens of pseudo-membrane.

There is no real difficulty in making use of any of these applications, if the children be properly managed. One or two assistants must hold the patient in such a way that the head shall be thrown backwards, and

the hands and feet secured. The physician must depress the tongue with the handle of a spoon held in the left hand, while he holds in the right the pencil or sponge-mop. If the child refuses to open the mouth, it can generally be made to do so by holding the nose in order to force it to breathe through the mouth. If this fail, all that is necessary is to press the handle of the spoon against the teeth, when the patient will soon become too much fatigued to offer further resistance.

Of late years, considerable difference of opinion has been expressed as to the importance of topical treatment in diphtheria; but in the light of the recent observations, especially of Wood and Formad, as to the secondary character of the blood-poisoning, in many cases it is clear that suitable local applications must be of positive and real value.

Gargles.—When the patient is sufficiently old and intelligent to be able to use gargles thoroughly, any of the substances which have been recommended as local applications may be thus used, being of course largely diluted. Thus tr. ferri chlor., hydrochloric acid, sol. sodæ chlorinatæ, in the proportion of fʒj or fʒij to fʒvj, or chlorate of potash in strong solution, may be used as gargles with much advantage in some cases.

These solutions may also be very efficiently applied to the throat in a finely divided condition, by means of the steam or hand-ball atomizer, a mode of application which is peculiarly useful in cases where the pseudo-membrane has extended into the larynx. As it is often impossible to employ gargles, and as we attach very great importance to the frequent use of mild solvent applications, particularly lime-water, we would strongly recommend the use of the steam atomizer for this purpose.

A very convenient and ready application, and one from which we have obtained marked advantage in several cases, especially where the exudation had extended into the larynx, is by covering the patient's head with a sheet, and introducing a vessel containing slaking lime, so that the steam may be freely inhaled. It is probable that the chief benefit is here derived from the warm watery vapor; though a small quantity of lime, in the form of impalpable powder, probably gains entrance to the fauces and air-passages.¹

Ice.—In a rather early stage of the disease, if there is much heat and engorgement about the throat, cold, wet compresses may afford temporary

¹ Bouchut has lately strenuously advised active cauterization of the fauces, or ablation of the tonsils, not only for the purpose of removing the exudation which appears on them, which he considers the localization of the disease, but also of facilitating respiration.

According to him, the operation of ablation has now been performed fifteen times, five by himself, and ten by MM. Domere, Symyan, Speckahn, and Paillot, with successful results in each case, no false membrane reappearing.

Despite this favorable report, however, the procedure appears to us objectionable, regarding, as we do, the importance of the local condition as secondary to that of the alteration of the blood. The operation must further cause the greatest alarm and most powerful resistance on the part of young children, and it seems highly improbable that a large proportion of cases should be attended with the same fortunate exemption from a recurrence of pseudo-membranous formation, as occurred in Bouchut's cases.

relief; and great benefit is often obtained in cases where there is much swelling, and inflammation of the fauces and pharynx, by the free internal use of ice, allowing the patient to hold small pieces of it almost constantly in the mouth.

Other external applications may also be employed to reduce the swelling of the cervical and submaxillary glands, render deglutition more easy, and relieve suffering; and, in this way, the persistent use of poultices or spongio-piline fomentations is of service.

It is essential to remember, however, that all blisters or irritating applications capable of destroying the epidermis, must be carefully avoided, owing to the tendency, already alluded to, of the pseudo-membranous deposit to occur on such abrasions.

When the nasal fossæ have become implicated from extension of the pseudo-membrane, one of the dilute solutions recommended as gargles should be injected frequently through the nostrils, or the desired effect may be even more thoroughly secured by the use of the same fluid through a Thudichum's nasal douche.

GENERAL TREATMENT.—Whatever differences of opinion may exist in regard to the relative merits of the various local applications we have enumerated, all high authorities are now agreed as to the general character of the constitutional treatment which should be adopted.

Some years ago, before opportunities had been presented for studying diphtheria in its epidemic form, as it has since occurred, it was customary to employ moderate depletion early in the attack, if the patient was vigorous and strong, and to follow this by the use of mercury and antiphlogistics, with a view of subduing the febrile excitement, and causing the dissolution and absorption of the pseudo-membrane.

With the increase of knowledge, however, of the true pathology and natural history of the disease, which has been gained of late years, all depleting and antiphlogistic plans of treatment have been, by common consent, abandoned as indefensible either in theory or practice, and all efforts are directed to promoting the nutrition of the patient and supporting the strength of the system, as indicated by the marked tendency to prostration, the feeble pulse, and the manifest deterioration of the blood.

It is probable that those cases in which bloodletting and the administration of mercurials were adopted with such apparent benefit, were either erroneously considered diphtheritic, or that the disease, when occurring sporadically, as it formerly did, was of a far more sthenic type than it has presented of late years.

It must be added, however that in the sporadic cases still frequently met with, when no grave epidemic influence is prevailing, the use of alkalies, as soda, combined with small doses of calomel, has been found very successful.

Regarding diphtheria as a constitutional affection, depending upon a peculiar alteration of the blood, we must admit that we are in possession of no remedy which in any respect merits the name of a specific in its treatment.

Among the best internal remedies, however, are the various preparations of chlorine, iron, and bark, which may be given singly, or, preferably, in combination.

Thus there are no remedies of more uniform and marked advantage than sulphate of quinia and tincture of the chloride of iron, given in full doses at short intervals. Some good observers prefer the liquor ferri chloridi to the tincture, and administer it successfully in the dose of gtt. j or iss every two hours for a child two years old. Hydrochloric acid or chloric ether may be added to these tonics, and this combination is strongly recommended by West and other high authorities.

The Sanitary Commission, in London, reported very strongly in favor of a mixture containing tincture of the chloride of iron, with chlorate of potash, chloric ether, and hydrochloric acid, sweetened with syrup; full doses being employed according to the age of the patient, and frequently repeated. This combination has been by Gibb, rendered still more stimulating by the addition of muriate of ammonia.

Oil of turpentine has been recommended (Dr. Perrey, *Med. Times and Gaz.*, March 5th, 1859) in large doses, both for its stimulating effect, and from its tendency to promote the absorption of lymph in adynamic states of the system, where mercury cannot be given.

Chlorate of potash, given in Huxham's tincture of bark, has been vaunted as almost specific in the treatment of diphtheria; but, as remarked by West, it unquestionably fails to produce here those excellent effects which are obtained from its use in ulcerative stomatitis.

Permanganate of potash, which has been so extensively used of late years in zymotic diseases, has been used both locally and internally in this affection, but apparently without any very positive advantage.

Mercurials.—Recently Dr. G. A. Linn¹ reported remarkable results from the use of large doses of bichloride of mercury in grave cases of diphtheria. He found that even so large a dose as gr. $\frac{1}{10}$ every three hours was well borne by children of one year old, and asserts that from his experience it prevents the spread of the membrane or the development of blood poisoning, and acts as much as a specific in diphtheria as quinia does in intermittent fever. These bold assertions have been corroborated by several good observers. We have not used this remedy sufficiently to authorize an expression of opinion, but a truly remarkable case, occurring in the practice of Dr. T. J. Yarrow of Philadelphia, and seen by us in consultation,² where this remedy was used in the above manner with excellent results, convinces us that further cautious trials should be made in this direction. The same may be said for the treatment by enormous doses of calomel, which has been advocated by some good observers as producing specific curative effects. It is difficult to define the cases in which it might be justifiable to try either of these modes of treatment, but it seems to us that it would chiefly be in cases where a continued tendency to the formation of

¹ Trans. Penna. State Med. Society, p. 886, 1879.

² Address on Medicine, by William Pepper, M. D., Trans. Amer. Med. Association, 1881.

pseudo-membrane showed itself while as yet no extreme degree of blood poisoning had occurred.

Emetics; Purgatives.—Emetics are useful when the exudation shows a disposition to extend into the larynx, or when there is much difficulty of breathing from tumefaction of the fauces, or from accumulation of the pseudo-membranous deposits. We would recommend under these circumstances the use of alum or ipecacuanha, as recommended in the article on pseudo-membranous laryngitis; the emetic being repeated in six or twelve hours, if the same indication should continue or recur.

A purgative dose is useful at the commencement of the disease, merely as an evacuant. After that period only such laxatives need to be employed as may suffice to keep the bowels soluble.

Stimulants.—In the milder forms of diphtheria, where no complications exist, the cases usually terminate favorably without the use of any stimulants; but there are many cases, on the other hand, characterized by pallor of surface, marked weakness of the circulation and tendency to prostration, great enlargement of the cervical glands, and extensive disease of the throat, where the pseudo-membranes rapidly decompose and assume a gangrenous appearance, and the urine is frequently albuminous, in which stimulants, freely administered, are positively required.

In cases where such adynamic symptoms are present, we should begin early in the attack with the administration of the weaker stimuli, and employ the stronger forms as the disease advances and the strength of the system succumbs more and more.

Food.—In no disease should more sedulous care be paid to securing to the patient a proper amount of suitable nourishment; and, indeed, in the absence of any remedy which can be looked upon as essential or specific, we must assign, perhaps, the most important part in the treatment of diphtheria to food and stimulants. It is at least certain that where these cannot be administered in proper quantity, all other treatment is unavailing, and hence it is our duty, upon finding that the pain and fatigue experienced by the child when forced to take frequent doses of medicine make it utterly unwilling to take food, to abandon all strictly medicinal treatment, and trust to sustaining the powers of the system by the free use of stimulants and concentrated food.

In cases where mechanical obstruction exists, or where all efforts at voluntary deglutition are obstinately resisted from fear of the great pain caused by the act, nutritious and stimulating enemata must be immediately resorted to. These may consist of beef-tea, eggs beaten up in milk, brandy in the form of milk-punch, and, further, may be medicated by the addition of quinia. They should be given every three or four hours, in rather small quantity, and not so concentrated as to irritate the bowel. When thus administered it is quite possible to sustain life for several days, until food can again be introduced into the stomach.

In addition to the local and general treatment above recommended, the patient should be rigorously confined to bed during the whole treatment, and for at least ten days after the disappearance of the exudation. This caution is given, not only on account of the danger of that most fatal ac-

cident, the formation of a heart-clot, but because we have twice known the exudation to reappear when the patient had been allowed to leave the bed at too early a period; and in one of these the exudation extended into the larynx on the occasion of the second attack, in spite of all that could be done, and life was saved only by the operation of tracheotomy.

The most scrupulous cleanliness of the person and surroundings of the patient should be preserved; free and uninterrupted ventilation secured; and on account of the positive, though perhaps slight, contagiousness of diphtheria, it is wise to practice separation of the well children in the family from the sick.

The treatment required in those cases where the pseudo-membrane extends into the larynx, and especially the discussion of the indications for the operation of tracheotomy, will be found in detail in the article on pseudo-membranous laryngitis.

Treatment of Paralysis.—We have already stated that the prognosis in diphtheritic paralysis is usually favorable, the symptoms often disappearing in the course of time without treatment. The cure may, however, be much hastened by a persistence in the administration of iron and quinia, to which strychnia should be added in full doses.

Nitrate of silver has also been employed in full doses with apparent benefit.

The paralyzed muscles should be faradized daily; and, when accessible, sea-bathing or sulphur baths may be employed with advantage.

In those cases where the muscles of deglutition are especially affected, and the nutrition of the patient is suffering from his inability to swallow sufficient food, it is desirable to resort to the use of nutritious enemata.

Treatment of Heart-clots.—Under the supposition that the blood is hyperinotic in the latter stage of diphtheria, the various salines, especially the vegetable ones, such as the citrates and acetates, and ammonia, given either as the carbonate or in the liquid form, have been recommended by Richardson.

When, however, the symptoms indicate that deposition of fibrin has absolutely occurred, it is probable that nothing can be done in the way of curative treatment. Alkalies may be given internally, the vapor of ammonia inhaled, alkaline solutions injected into the veins, but there is little reason to hope that any effect upon the clot can be produced.

In one of the cases reported by us (*loc. cit.*), the clot presented at one extremity a granular, partially disintegrated condition, as though its removal had begun by interstitial action, and the mechanical effects of the blood current; and it is possible that by supporting the powers of nature the removal of the clot might be effected in this way. Indeed, there are cases on record (quoted by Robinson, *loc. cit.*) in which the symptoms have most clearly demonstrated the existence of a clot in the cavities of the heart, where recovery has still occurred.

ARTICLE XII.

EPIDEMIC CEREBRO-SPINAL MENINGITIS.

DEFINITION; SYNONYMS; HISTORY; FREQUENCY.—Among the large number of names which have been applied to this disease, the only additional ones which call for mention are cerebro-spinal fever and spotted fever. The latter has been much used in this country, but as it is based upon a symptom of occasional occurrence only, is evidently inadmissible. The name we have adopted is the one now almost universally accepted.

Epidemic cerebro-spinal meningitis is an acute specific febrile affection, occurring in epidemics of widespread or local character, but not propagated by contagion; characterized by alterations of the blood and by inflammatory changes in the membranes of the brain and spinal cord, and running a most irregular course both as regards symptoms and duration.

	Epidemic Cerebro-spinal Meningitis.	Typhus Fever.	Typhoid Fever.	Diphtheria.
1860	0	14	68	214
1861	0	17	148	502
1862	0	37	654	325
1863	49	131	486	434
1864	384	335	648	357
1865	192	334	773	460
1866	92	96	381	192
1867	102	138	367	119
1868	54	108	395	119
1869	36	49	373	182
1870	36	69	409	172
1871	44	37	313	145
1872	128	35	369	150
1873	246	31	382	110
1874	82	26	461	179
1875	83	21	419	652
1876	84	27	761	708
1877	56	15	542	458
1878	90	9	404	464
1879	62	1	344	321

Although it is highly probable that epidemics of varying extent occurred previous to the present century, it appears that their true nature was not recognized, and the first distinct account of epidemic meningitis

was published in 1805 by Vieusseux. Since that time, however, it has appeared more or less frequently in almost every country of the globe. It began its course in the United States in 1806, and numerous epidemics of it occurred between that time and the year 1816; again between 1823 and 1830; again between 1842 and 1850, and again between 1856 and the present time. The last epidemic began in Philadelphia in 1863, and returned annually for some years, reaching its height in 1866-67. Since then, however, as will be seen from the table, it has continued to furnish annually a considerable number of deaths.

It was made the subject of several valuable memoirs, among which may be mentioned those of Gerhard,¹ Githens,² Levick,³ and particularly the admirable treatise of Stillé.⁴

Owing to the employment of various names for this disease in the mortality reports, it is difficult to estimate the actual mortality occasioned in this city by it, but a good idea may be obtained from an inspection of the table, which presents the annual mortality for twenty years from epidemic cerebro-spinal meningitis, typhus and typhoid fevers, and diphtheria.

CAUSES.—The disease we are considering has occurred "in all portions of the temperate zone inhabited by European races and their descendants; in all sorts of localities, among all ranks and conditions of society, at all ages, and in both sexes; and it is, therefore, in the strongest sense of the word a pandemic disease." (Stillé, *op. cit.*, p. 94.) While its epidemic character is so strongly marked, there is no reliable evidence as to its being contagious. A few authors, however, even of such recent date as Bristowe, assert their belief in its contagiousness.

Although occurring in both sexes and at all ages, it is more frequent in females, and a large majority of the cases are among minors. The disease has undoubtedly been more frequent and virulent among over-crowded and filthy populations, but there is no reason to think that these conditions exert any more than a general depressing influence.

ANATOMICAL LESIONS.—The chief lesions in this disease are found in connection with the blood and the nervous centres.

Blood drawn from a vein during life usually presents its normal characters or else those indicative of inflammatory action, excepting in cases of the most malignant and rapidly fatal type. On the other hand, in post-mortem examinations, the blood is most frequently found to be dark, and either altogether fluid, or with only small dark and soft clots, in the cavities of the heart and the larger bloodvessels. These alterations are evidently what should be expected in a disease presenting the double condition of a specific blood poison and a distinct and serious local inflammation, which two elements vary greatly in their relative preponderance in different epidemics, and even in different cases of the same epidemic; no definite chemical changes have been ascertained. In cases where the

¹ Amer. Jour. Med. Sci., July, 1863, p. 105.

² Ibid., July, 1867, p. 17.

³ Trans. of Amer. Med. Assoc., xvii, p. 311.

⁴ Epidemic Meningitis, Phila., 1867, p. 178.

crasis of the blood is greatly impaired, microscopic examination has frequently shown an absence of the ordinary mode of arrangement of the red corpuscles in rouleaux, and a crenated appearance of the corpuscles themselves.

The lesions of the nervous centres are chiefly seen in the meninges of the brain and spinal cord. In the early stage, there is extreme engorgement of the vessels of the meninges, with a loss of the normal translucency of the pia mater and arachnoid. Later, an exudation occurs which at first may be serous, but soon becomes sero-purulent or entirely composed of thick creamy pus, with a varying proportion of lymph. The amount of this exudation varies greatly in different cases; at times being quite scanty, while in other instances several ounces of pus are present.

The exudation usually occupies the subarachnoid space, and is at times associated with effusion into the ventricles of the brain. The convexity and the base of the brain are both involved, though the exudation is as a rule more abundant over the latter, and especially about the optic chiasm, the fissures of Sylvius, and the base of the cerebellum and under surface of the pons. One or more of the cranial nerves are completely imbedded in the exudation.

The spinal meninges present the same general changes, the vessels being congested, the pia mater infiltrated with sero-purulent fluid, and the subarachnoid space occupied by a more or less extensive exudation. These lesions are usually most marked about the medulla and again at the lower part of the cord. It is probable that the tendency of the exudation to accumulate at the latter points is partly, at least, due to gravity.

The substance of the brain is usually vascular and more or less softened; and the spinal cord presents the same changes, though less constantly and usually to a less degree. In some cases, however, the softening of the cord is extreme, and Hirsch has recorded a case in which the central canal of the cord was distended with pus.

In some cases where death occurred very rapidly, within twenty-four or forty-eight hours, the cerebro-spinal lesions have been found wanting, not having yet been developed to an appreciable extent.

There are no characteristic lesions of any other organ.

SYMPTOMS ; COURSE.—In some cases the attack is preceded by the usual prodromes of acute specific fever; but in many instances it occurs without warning in the midst of perfect health. Usually the actual outbreak is marked by a distinct rigor, followed by fever, prostration, vomiting, intense headache, and pains in the back and limbs. It is soon evident that the patient is seriously ill. He grows restless and tosses about, at times with slightly spasmodic twitchings of the muscles. The intellect often remains clear; but delirium or a disposition to heaviness and dozing may be present. The temperature is rarely very high at first, and the pulse ranges from 90 to 120, according to the age of the child. As early as the second day we may observe that the head is retracted, and this increases in its degree and soon becomes associated with a tendency to opisthotonos. The other nervous symptoms increase in severity; headache persists, and

is even so severe as to elicit screams of pain; the pupils are contracted; there is delirium, with excessive restlessness or even convulsions. Extreme cutaneous and muscular hyperæsthesia are very common symptoms. Vomiting is apt to continue; the bowels are constipated, the abdomen retracted, and the urine is usually retained. The pulse grows more frequent and small, the respiration is greatly accelerated, and the temperature increases to perhaps 103° or 104° F. Herpetic eruptions about the mouth are common; and in a varying proportion of cases a purpuric eruption appears from the second to the fourth day. In some instances the course of the symptoms presents marked fluctuations, which may assume the form of distinct quotidian or tertian remissions. In unfavorable cases, stupor supervenes, interrupted by more or less marked spasmodic movements; respiration and circulation become more and more impaired, and the patient dies in profound coma. In favorable cases, on the other hand, the nervous jactitation and delirium diminish, the tetantic muscular spasms relax, the urine is passed voluntarily, the pulse and respiration gradually become uniform, and the patient enters upon convalescence.

Death may occur even in a few hours from general paralysis, from the overwhelming effect of a violent and universal meningitis. It may occur in the course of one or two days, in malignant cases with extremely marked blood lesions. In cases of the regular form, death may occur from the fourth to the eighth day, from the effects of the cerebro-spinal lesion; but after convalescence has commenced, the supervention of some complication or sequel may prove fatal, even after the lapse of weeks or months.

Before proceeding to discuss a few of the principal symptoms more in detail, the following case of severe type, which occurred in our practice, may be cited in illustration of the general sketch above given:

CASE.—David R., æt. 12 years, in good health, was attacked on June 8th, without apparent cause, with violent headache, languor, and occasional vomiting. Cerebro-spinal meningitis was prevalent as an epidemic at the time (1864).

On June 9th (second day), at noon, headache continued intense, the fever increased in severity, and he twice vomited a greenish liquid. The bowels were constipated and urine scanty. He lay in a drowsy condition; the eyes slightly injected, but pupils of normal size; the nose was pinched, the mouth closed. There was no rigidity of the muscles of the neck or of other parts. There was marked jactitation, with mild delirium. The skin was dry and slightly pungent; respiration 48, quick and sighing; pulse 105. Over the whole body, but especially on the extremities, there were small petechiæ, of livid color. Abdomen meteoric. Ordered brandy $f\text{℥ss}$. every half hour, a saline laxative, and the following:

R. Morphiæ Sulph.,	gr. $\frac{1}{4}$
Acid. Sulph. Aromat.,	℥v.
Elix. Cinchonæ,	$f\text{℥ss}$.
Ft. Sol.—To be taken every two hours.					

In the evening, his mind was more clear; but rigidity and soreness of muscles of back of neck had appeared, with tendency to opisthotonos. Pulse 120, irritable and quick. Vomiting continued. The bark mixture was changed for one containing quinia, morphia, and dilute sulphuric acid. Iced champagne and beef-tea were given in small quantities. Counter-irritation by capsicum and whisky.

In the night, at two o'clock, vomiting continued frequent and uncontrollable. Pulse

was very feeble, almost-thread-like, and very irregular, varying from 100 to 120 within a few minutes' time. Temperature equally variable, surface being alternately burning hot and chilly. Quite conscious, and complained of intense headache; extreme restlessness continued. Frequent thin, dark stools.

June 10th.—In morning, pulse 90, varying from minute to minute; respiration less hurried. Surface warm. He passed urine and feces. Ordered suppositories of gr. ss. opium and gr. ij quinia every five or six hours; a blister behind each ear; f $\frac{3}{4}$ ss. brandy every hour, and concentrated nourishment.

About five hours later, stupor supervened and continued until night, interrupted by fits of violent delirium. Slight convergent strabismus. Petechiæ continued, and a herpetic eruption appeared about the mouth. Temperature reduced in extremities. Ordered nutritious enemata; blisters 4x4 to back of neck; small doses of opium by rectum.

June 11th.—More quiet; no more active delirium. Bowels opened more freely. Urine free, acid reaction, sp. gr. 1.032, with a very heavy deposit of whitish urates. Nourishment and medicines retained by stomach. Pulse 85 to 105, with more volume. Consciousness partially regained; complained of frontal headache and muscular soreness. No opisthotonos. Eyes slightly injected. Continued opium and quinia; beef tea; iced champagne; milk with lime water.

June 12th.—Condition about the same. Some quiet sleep. Skin of pleasant temperature. Herpes abundant about mouth. Urine quite free, and light-colored. Pulse about 100; respiration more full. Treatment continued.

June 13.—Pulse 72 to 100. Consciousness perfect, but complained of intense frontal pain. Temperature almost normal. Petechiæ disappearing; herpetic eruptions on face still marked. Passed urine freely; sp. gr. 1.011; no albumen; bowels costive; quinia discontinued on account of headache. Ordered blue mass, gr. j, every two hours for four doses, followed by a laxative enema, which acted freely, and gave much relief.

June 14th.—Restless, and complained of his head. Still some retraction of the head, and muscular soreness. Respiration 35; pulse 100 to 116. Stomach retentive. Tongue cleaning, but dry and cracked. Passed urine copiously, very light-colored and watery. Suppositories of quinia and opium resumed; stimulus and nourishment as before.

June 15th.—Doing well. Has emaciated very rapidly. Pulse 110, falling during the day to 96; respiration less sighing and labored. Decubitus more natural, and movements more easy and free. Opiates suspended. Takes quinia sulph., gr. j; acid. muriat. dil., gtt. x; q. q. h. Stimulus and food as before.

June 16th.—Doing well. Petechiæ disappearing; herpes around the mouth better. Tongue moist, but very sore, with ulcerated cracks covered with pultaceous crusts. Pulse 100, of good volume. Expression natural. No signs of spinal irritation.

June 19th.—Ninth day of disease. Convalescent.

June 20th.—Rapidly regaining strength. Sleeps well, and eats with great appetite. No headache; expression bright and natural. Tongue has healed, and herpes around mouth nearly gone. Bowels regular; urine free and normal. Continues quinia and mineral acid. The subsequent course of the case was that of rapid restoration to health without any sequelæ.

In entering upon a study of the special symptoms, it will first be observed that the mode of onset of the disease is peculiar, and is characterized by its suddenness and by the early appearance of grave nervous symptoms without a high grade of fever.

The symptoms furnished by the nervous system merit the closest study. So important are they that many authors have been led to regard the whole affection as due to the meningeal inflammation, losing sight of the

coëxistent lesion of the blood. The most marked of them is headache, which, as already stated, appears early, is usually sharp and lancinating in character, and violent in the extreme. It is accompanied with pain along the spine, especially in the cervical region, which is much increased by pressure or by motion. Neuralgic pains are also very common, and at times are extremely severe: they chiefly affect the extremities and the abdomen. From an early period of the case, there is apt to be marked and painful hyperæsthesia of the skin, and also of the muscles; so that handling the patient causes complaints or even cries of pain. Later this may be followed by more or less marked anæsthesia. They are frequently also various forms of muscular spasm. At times a high group of muscles will be affected with spasm, either tonic or clonic. The most frequent instances of this are the retraction of the head from rigid spasm of the cervical muscles, and the tendency to opisthotonos; both of these conditions are very constant, and may be present in a high degree. Trismus is not very infrequent; while in other cases, the muscles of the extremities are affected with spasm; or again, the tetanoid symptoms may assume the form of general epileptiform convulsions, which are much more frequent in children than in adults.

Paralysis in various forms, hemiplegia, paraplegia, but much more frequently as affecting a single cranial nerve (facial, abducens, or oculo-motor) has been observed, but as a rule is not met with until the later stages of the disease. In some cases, marked subsultus is present. In addition to these symptoms, there is rapid and marked debility, frequently associated with vertigo on rising into the sitting posture. Disturbances of the intellect are also more frequent in children than in adults. At first there are great restlessness, jactitation, and wandering delirium, which may persist or be replaced by more or less profound stupor; perhaps occasionally interrupted by active delirium. In fatal cases, this stupor deepens into coma as the end approaches.

There is frequently loss of control over the bladder and rectum, attended with retention or with involuntary discharges.

The organs of *special sense* furnish important symptoms. There is often a uniform, diffused redness of the conjunctiva. The pupils are usually contracted at first, but later may be dilated or unequal. Not rarely blindness follows; due either to keratitis, to exudative inflammation of the retina or choroid, or to purulent exudation into the chambers of the eye. Squinting frequently results from paralysis of one of the motor nerves of the eye as already stated. In like manner, deafness often results from suppurative inflammation of the internal ear, or from inflammation of the auditory nerve.

The *expression* of the patient varies much at different periods of the disease. In the early stage, the face is often pale, with pinched nose and sunken features. This peculiar facies, which was noted in the case above narrated, is regarded by Hirsch and others as very characteristic.

The *pulse* and *temperature* are less uniformly affected than in any other of the acute specific febrile diseases. At first the pulse may not be much

accelerated, but later it may be very rapid, or again, may present remarkable variations at short intervals. The course of the fever is very irregular; in most cases the temperature does not rise above 103° or 104° , and not infrequently is much lower throughout the whole course of the attack.

The disturbance of respiration is marked but very irregular. In the early stages it corresponds with the condition of the pulse; but later it presents irregularities due to the pressure of exudation on the pneumogastric nerve, and may then assume the peculiar form known as *ascending and descending breathing*, which is so frequently seen in the exudative stages of tubercular meningitis.

The *digestive system* presents few constant symptoms. Vomiting, however, is nearly always present in the early stage, and may be frequent and uncontrollable. It is often unattended by any nausea, and there evidently depends an irritation at the base of the brain. We have already referred to the severe abdominal neuralgic pain sometimes complained of. The bowels are usually constipated, and the abdomen retracted. In the later stage, it is not rare for involuntary discharges to occur.

The *urine* presents the ordinary febrile characters, and in addition is at times albuminous or even bloody. There is frequently retention, requiring the use of the catheter, though we think this less frequent in children than in adults.

We have already alluded to the occurrence of *eruptions* on the skin. They are not constant, and in some epidemics have been rarely noticed. Still they constitute very important symptoms in the majority of cases; and indeed the occurrence of a petechial eruption has been so marked a feature in most American epidemics as to render the objectional name, "spotted fever," the popular title for this affection.

Among these eruptions, groups of herpes are frequently seen on the face, especially about the mouth, while in more rare instances, erythema and urticaria have been observed. But by far the most frequent and important is the petechial eruption, which appears early in the case, usually in the form of small spots. The proportion of cases in which such an eruption is present varies greatly in different epidemics. Stillé concludes (*op. cit.*, p. 64), that, taking the whole of the cases of epidemic meningitis in Europe and America, it did not occur in more than 10 per cent. On the other hand, of 96 cases recorded by Dr. Githens (*loc. cit.*), 36 had marked petechial eruptions; and of the cases we have ourselves observed, the proportion has been even greater.

The *duration* of this disease is extremely variable. We have seen it prove fatal in the adult from collapse within forty-eight hours, with all the evidences of profound blood alterations, and before the characteristic lesions of the meninges had passed beyond the first stage. Death has been known to follow even in a few hours, with symptoms of general paralysis.

In cases of ordinary severity which terminate favorably, the duration of the acute attack may be stated as from five to fourteen days, but complete convalescence may be postponed for weeks or months on account of the occurrence of some of the sequelæ so frequent after this affection. On the other hand, death may occur at any period from the first day, as above

stated, to as late as months after the original attack. Hirsch states that its duration is between a few hours and several months.

Convalescence may be prompt, satisfactory, and complete; or it may be irregular and protracted, in consequence of the persistence of some symptoms or the development of some of the sequelæ. Among them may be mentioned blindness and deafness due to causes already stated. Some neuralgic pains, dependent upon irritation of the posterior roots of spinal nerves, tonic or clonic spasm of certain muscles, and more rarely paralysis of a single muscle or of one or more members. It will be observed that these sequelæ are, for the most part, the results of the inflammation of the cerebro-spinal meninges.

The *mortality* varies greatly in different epidemics. According to Hirsch it has varied from 20 to 75 per cent. As we have met with it in this city, the mortality was about 33 per cent.

The *prognosis* should always be guarded in epidemic cerebro-spinal meningitis. Even when the threatening symptoms of the acute attack begin to subside, it is impossible to predict that there will not be left behind some sequel which may protract convalescence indefinitely or induce a lingering and painful death.

The *diagnosis* of this disease is not difficult. It could scarcely be confounded with typhoid fever; but in case of the coëxistence of an epidemic of typhus fever, it is important to note the differential marks by which this latter may be distinguished from epidemic meningitis. Thus the headache of typhus fever is less violent and sharp, and the delirium less active and marked. In typhus, also, the face presents a dusky flush, and the conjunctiva is injected; while in meningitis the face may be pale at first, and the conjunctivæ are of a uniform pinkish color. The rapidity of pulse and elevation of temperature are much more marked in typhus, and follow a much more regular course. The eruption of typhus appears on the third or fourth day; while in meningitis the occurrence of petechiæ is not at all constant, and when present they may appear as early as the first or second day. On the other hand, the herpetic eruptions so frequent in meningitis are wanting in typhus. Vomiting is much more frequent in meningitis than in typhus. Retraction of the head, opisthotonos, muscular spasms, neuralgic pains, cutaneous hyperæsthesia, irregularity of the pulse and respiration also, which are such marked symptoms of meningitis, are not characteristic of typhus. We may add that although typhus occurs in childhood, it is comparatively rare; while, as we have already seen, epidemic meningitis occurs more frequently among children than at any other age.

We would call attention also to the possibility of confounding mild cases, without eruption, for rheumatism of the cervical and dorsal muscles. We knew this error to be committed in one case, with fatal results.

TREATMENT.—In the treatment of this affection it is necessary to bear in mind its real nature. There can be no doubt, however, that, in the majority of cases at all amenable to treatment, the meningeal lesions must be the chief object of our medication.

The use of cathartics is to be adopted with great caution. A mild saline

laxative may be administered, and its action aided by an enema, if marked constipation exists at the onset. But it is to be remembered that but little food can usually be retained, and also that the constipation is the result of the cerebral irritation.

Opium administered freely, so as to quiet the intense pain and nervous restlessness, is to be recommended. Its use will also at times control the cerebral vomiting. It is best given by suppository or enema, or in the form of morphia by hypodermic injection.

Bromide of potassium, as a nervous sedative and antispasmodic, may be employed with advantage.

Ergot has also been highly recommended on account of its well-established power of influencing congestion of the cerebro-spinal vessels; and certainly this would appear sufficient reason for the employment of this remedy alone or in conjunction with *belladonna*.

Our own experience agrees with that of most American physicians, in regard to the beneficial effects of quinine, given in large doses, either by the mouth, or, if vomiting be persistent, in the form of suppository.

Local depletion by leeches or cut cups to the back of the neck and along the spine should be employed, unless the evidence of blood dyscrasia forbid it. After its employment, or instead of it, in case it be deemed inadmissible, the repeated application of dry cups along the spine is to be recommended.

Food should be very carefully administered in small quantities, and if necessary, nutritious enemata should be resorted to.

In case of extreme debility, of depression of the circulation, or of tendency to collapse, alcoholic stimulus should be freely employed, and the same is true, when, later in the course of the case, the typhoid state is markedly developed. It may also be necessary to apply external warmth to maintain the temperature of the body.

For the relief of the sequelæ which result from the imperfect absorption of the exudation, and the consequent irritation or pressure upon the roots of the cranial or spinal nerves, evincing itself by neuralgic pains, muscular spasms, contractions or paralysis, the necessary treatment comprises continued counter-irritation to the spine; the internal administration of iodide of potassium, nitrate of silver, or bichloride of mercury; and the use of one or the other form of electricity.

CLASS VII.

DISEASES OF THE SKIN.

INTRODUCTORY REMARKS.

IT would be worse than useless, in a work like the present, to attempt a full description of all the diseases of the skin to which children are subject. Such a course would compel us to devote to more important matters than the affections of the skin, a much smaller proportion of space than they require and deserve. We shall therefore select only those cutaneous diseases occurring in early life, which are most important either from their frequency, or because they present in children some particular aspect or peculiarities, which make it necessary that they should be studied separately from the same affections in adults. Moreover, we shall treat of each one as it comes before us with greater or less copiousness of detail, according to its respective consequence to the medical practitioner, eschewing carefully any useless detail in regard to the more unimportant ones, but endeavoring anxiously to describe with accuracy the history, diagnosis, and treatment of such as demand a greater degree of consideration.

The progress in scientific dermatology has been so rapid during recent years that it may be hoped that ere long a strictly satisfactory classification, with definite significations attached to each name, will be formed and accepted universally. For the present, we are led to accept the classification adopted by the American Dermatological Association in 1878. According to this, there are nine classes of skin diseases, as follows :

1. Disorders of the sweat glands, including hyperidrosis, etc. ; and of the sebaceous glands, including seborrhœa, etc.
2. Inflammations, including erythematous, vesicular, bullous, papular, pustular, squamous, and phlegmonous forms.
3. Hemorrhages, as purpura.
4. Hypertrophies, as affecting the pigment, the epiderm, papillæ, hair or nails, or the connective tissue of the corium.
5. Atrophies, as affecting the same individual elements.
6. New growths, either of connective tissue, of vessels, or of granulation tissue.
7. Ulcers.
8. Neuroses, including hyperæsthesia and anæsthesia.
9. Parasitic affections of vegetable or animal origin.

In beginning the study of any case of skin disease, it is essential to form a correct diagnosis, so that the case may be referred to the class where it

belongs. There are certain elementary lesions which, taken in connection with other considerations, have a good deal of value in determining the character of any given eruption. A simple enumeration of these will suffice here, as follows, *Maculæ*, or stains; *hyperæmia*, or redness; *pomphi*, or wheals; *papulæ*, or pimples; *vesiculæ*, or little bladders; *bullæ*, or blebs, bladders of a larger size; *pustulæ*, or pustules; *squamæ*, or scales; and *tuberculæ*, or small solid lumps in the skin.

As the eruption in many forms of skin diseases passes through various stages and presents different appearances at different times, it is important to study not one patch of eruption only, but as many points as exist, so as to determine the character of the latest developments of the disease.

The two following practical rules, quoted from Tilbury Fox's *Epitome of Skin Diseases* (Phila., 1879, p. 16), express in forcible terms the true diagnostic method: "All diseased places, or as many as possible, should be carefully examined, and not one only, or one here and there, for the simple reason that the eruption may be at very different stages of development, and therefore present diverse aspects, in different localities, upon the same patient.

"When in any given case the earlier stages are not present so as to be recognizable, careful inquiry should be made by interrogation of the patient, as to the changes that have occurred before the disease came under observation, with the view of determining its nature." A complete diagnosis available as a guide to successful treatment, must include, not only the determination of the primary and essential form of the eruption and its proper classification, and the detection of any complication or coëxisting eruption, but should also embrace the recognition of the causes, predisposing and exciting; and of any conditions, local or constitutional, that may modify the character or course of the disease. It is more and more clearly demonstrated that there is, in reality, nothing special or peculiar in the pathology of skin disease, but that here, as elsewhere, we have to deal with the ordinary pathological changes merely modified in their expression by the anatomical peculiarities of the tissue affected.

It will be found that the large majority of important skin diseases among children come under the class of Inflammations or else of Parasitic Affections. It will be found, also, that with them, even more than at a later age, these affections are influenced or caused by imperfect digestion or assimilation, or by unfavorable climatic or hygienic conditions; and that they are frequently modified by inherited constitutional taints, most commonly of a scrofulous nature.

As already stated, we shall not consider all the classes of skin affections systematically, but shall devote attention only to those that present unusual importance in childhood owing to frequency of occurrence or peculiarity of manifestation.

CHAPTER I.

ERYTHEMATOUS AFFECTIONS.

THE chief feature of these is the presence of hyperæmia, mainly affecting the papillary layer, with or without some slight consequent effusion of serum, swelling of the cells of the rete mucosum, rarely vesiculation, but subsequent desquamation.

These comprise erythema, roseola, and urticaria.

ARTICLE I.

ERYTHEMA.

DEFINITION; FREQUENCY; FORMS.—Erythema is a superficial inflammation of the skin, occurring in patches of irregular form and varying extent. In some cases there is mere hyperæmia, causing superficial redness of a rosy or deep-red hue, but without any swelling whatever; while in other cases there is more or less exudation, so that the patches of hyperæmia also present papules, vesicles, or tubercles. It occurs either as an idiopathic affection, from the action of local causes; or else as a symptomatic one, in connection with some systemic disturbance.

The causes that give rise to idiopathic erythema are various, including extremes of heat and cold; the action of chemical or mechanical irritants, etc. Of these, the only one that is specially operative in children is the chafing, caused by contact of the natural folds of the skin with one another, producing what is known as erythema intertrigo, or simply intertrigo.

Symptomatic erythema is observed in connection either with the febrile stage of acute local diseases, or with specific constitutional disorders. So varied are the manifestations of symptomatic erythema, under these different conditions, that a considerable number of forms have been described and named. Hebra adopts the term erythema multiforme to include such varieties as are called by many writers papulosum, tuberculosum, nodosum, fugax, marginatum, circinatum, iris, etc. Of these we shall describe only erythema fugax and nodosum as being the forms most commonly met with in children.

ERYTHEMA INTERTRIGO.—This form of erythema was for a long time, and is still by some known by the single name of intertrigo. It occurs on the portions of the body exposed to friction by the contact of opposite surfaces, and to irritation from the passage over, or retention upon them, of the urinary secretion or the fecal discharges. The most common seats of it are, therefore, in the folds of the skin about the neck, in the axillæ, the groins, about the anus, in the cleft of the nates, and on the inside of the thighs.

As it appears in the creases of the skin about the neck, or in the axillæ, it may be a mere red blush lasting a few days, and then disappearing; or,

after presenting this appearance for a short time, the inflammation may become much more intense, and pass into a true dermatitis, occasioning an excoriated condition of the surfaces attended with the discharge of a serous or a sero-purulent fluid; or, lastly, the inflammation may run into veritable ulceration, giving rise to extensive and very painful ulcers occupying the depth of the crease, presenting abrupt and jagged edges, and discharging very considerable quantities of pus. In one child, two months of age, of delicate constitution, and imperfectly supplied with food, we saw the last-described form of the disease occupying at the same time the groins, the axillæ, and the folds of the neck. The attack lasted two weeks, and very nearly proved fatal from the violent suffering it caused. In another child, not quite a year old, who was teething, it presented these characters in the neck and axillæ, while in the groins it was much less severe, the latter parts being merely excoriated.

Infants attacked with severe diarrhœa, with dysentery, or entero-colitis, and especially with that form of entero-colitis which so generally accompanies thrush, are very apt to have an erythema of the nates, genital parts, and the internal surfaces of the thighs. So common, indeed, is this occurrence that M. Valleix regards erythema of these parts as an almost constant accompaniment and even precursor of thrush. For our own part we have very often met with it in cases of diarrhœa in infants, even in those of very moderate severity, but we have never seen it precede the appearance of the intestinal disorder.

This form of erythema begins as a simple redness of the skin about the anus, between the buttocks, about the genital parts, and over the inside of the upper parts of the thighs. In a mild case of diarrhœa, and in a child properly cleansed after each evacuation by stool or urine, it will go no further than this; but in a severe attack of inflammatory diarrhœa, attended with frequent acid stools, and in a case in which proper cleanliness is not attended to, the long-continued contact of the discharges and soiled napkins will often cause the erythema to assume very distressing features. The redness extends in such instances along the legs to the feet; small papules, more or less numerous, make their appearance upon the inflamed skin; these are converted into pustules and then into ulcerations, and if the case goes on unchecked, the ulcerations become larger, run together, and present raw, deep red, and bleeding surfaces, sometimes of considerable size. Very often the ulcerations present a grayish plastic exudation upon their surfaces. When these conditions present themselves, the case has passed into an exudative form, and is properly to be regarded as an *eczema papulosum* or *pustulosum*. After cicatrization there remain, at the points where the ulcerations had existed, reddish and copper-colored spots, which do not disappear for a considerable length of time. This form of erythema rarely ceases entirely until the diarrhœa which has occasioned it has itself been cured.

ERYTHEMA FUGAX.—This form of erythema occurs chiefly as a symptomatic affection, in the course of various acute internal inflammations, and especially those which occur during dentition. It may occur during high febrile reaction brought on by any cause, especially in children having

an active cutaneous circulation. We have observed it several times in the local inflammations accompanied with great disturbance of the circulation, and particularly in cases of severe catarrh occurring during dentition, and in attacks of severe simple angina. In these cases it appeared in the form of a bright red rash, resembling very much a mild scarlatinous eruption. It was seated upon the upper part of the front of the thorax, and upon the outer surfaces of the arms. The red flush disappeared readily under pressure, and flashed back the moment the pressure was removed. There was no swelling whatever attending it, and the color was never so bright as that of a severe scarlatina, nor so deep as that of erysipelas or roseola. It lasted only a few hours or half a day, and then disappeared without desquamation.

The chief point of interest in regard to this form of erythema, as it has come under our notice, has been the diagnosis between it and scarlet fever. This is to be made out only by recollecting that it has made its appearance in the course of another disease, while the child is already suffering under some kind of sickness, which is not generally the case with scarlatina; by the less scarlet tint of the eruption, its more superficial character, and more limited extent; and lastly, by its short duration.

ERYTHEMA NODOSUM is an acute inflammatory affection of the skin, characterized by the formation of rounded or oval, variously sized, more or less elevated nodes. It is uncommon after the age of twenty years, and generally occurs in feeble and delicate children. We have never met with it under five years of age. Duhring (*Dis. of Skin*, 12th ed., p. 145) refers to the form which Uffelmann and Oehme have described as occurring in the younger members of tuberculous families. It may develop itself upon different parts of the body, but occurs in by far the greater part of the cases on the fore part of the legs, or over the anterior edge of the tibia. We have only twice seen it elsewhere, and then it was situated upon the outer surfaces of the arms and forearms. It is preceded usually for several days by general indisposition, by lassitude, thirst, loss of appetite, and some feverishness. It appears in the form of red spots of an oval shape, somewhat elevated in the centre, and which increase gradually in size. After a short time these patches become decidedly elevated above the surrounding surface, and in passing the hand over them they give the sensation of nodosities. They increase gradually in size, so as to measure from a few lines to an inch or an inch and a half long, by half an inch or an inch broad, when they present the appearance of reddish tumors, somewhat painful to the touch, and having an obscure feeling of fluctuation, as though about to suppurate. This, however, they never do, but after a short time they diminish in size, their red color changes into a bluish or livid tint, they soften, and finally disappear entirely in about twelve or twenty days. As a rule they do not appear at once, but come out at intervals in the form of crops. Rheumatic pains occasionally precede and attend the attack, and according to Fox, chorea sometimes occurs in connection with it. It has been supposed by Hebra that the nodes are due to an inflammation of the lymphatic vessels, while by others (Bohn) they have been attributed to minute embolisms of the cutaneous vessels; but we agree with Duhring

(*op. cit.*, p. 146), that its nature is still involved in uncertainty. We have met with five well-marked cases of this disease. Three occurred in girls between six and twelve years of age, and two in boys of the same age. They all appeared to depend on derangement of the digestive function, attended with a somewhat impoverished state of the blood, and general debility.

DIAGNOSIS.—The only disorders with which erythema could be confounded are erysipelas, roseola, or scarlatina, and this could happen only in regard to the erythema fugax. From erysipelas it may be distinguished by the superficial character of the eruption, the absence of swelling and of smarting and burning pain, and by the slighter severity and much shorter duration of the symptoms in erythema. Another important feature is the peculiar, abrupt, well-defined, and slightly elevated margin which marks the edge of the erysipelatous rash, and which does not exist with the same distinctness in erythema. Lastly, the singular regularity observed by erysipelas in its gradual extension from place to place, is altogether unlike the march of erythema, which shows itself suddenly, or in a few hours, over large surfaces, and, after lasting some hours or a few days, quickly disappears.

In roseola the peculiar deep-rose tint of the rash will serve to distinguish between it and the lighter red tint of erythema.

The mild character of the general symptoms, and the absence of throat affection in erythema, will prevent any one who is careful from mistaking the disease for scarlatina.

Erythema intertrigo cannot be mistaken for any other disease, and if the course and peculiar local character of erythema nodosum be borne in mind, it also may be easily recognized. The only thing with which the latter might be confounded is phlegmonous erysipelas, but if the mild character of the general symptoms in erythema nodosum, the distinctly circumscribed form of the tumors, and the fact that the disease never terminates by suppuration, are recollected, there need be no difficulty in making the diagnosis.

PROGNOSIS.—Erythema is a very mild disorder in a large majority of the cases. The only conditions under which it proves serious are when in the form of intertrigo, it attacks children laboring under chronic entero-colitis, or those affected with severe thrush connected with gastro-intestinal inflammation, when it cannot fail to increase the sufferings and danger of the patient; or, when it implicates, as we have seen it do in two instances, extensive portions of the cutaneous surface, involving the folds of the neck, armpits, groins, and genital organs, and this, too, without any other signs of disorder of the digestive apparatus than those showing functional derangement. In one of these cases the extent and depth of the ulcerations were so great, and the resulting suffering and constitutional distress so severe, as to have very nearly destroyed the life of the infant, who was but two months old at the time of the attack.

Erythema nodosum would almost certainly excite some uneasiness in the mind of a practitioner unacquainted with its real nature and probable course, and not only so, but it would prove tedious and difficult of cure,

unless treated in the proper way. When managed correctly, however, it almost always gets well without any difficulty.

TREATMENT.—Ordinary mild cases of erythema intertrigo require no other measures than attention to strict cleanliness. The irritated parts must be carefully washed two or three times a day, and if the nates, genital parts, and thighs are concerned, the washing must be repeated after each evacuation of urine or stool. After this the parts should be dusted with fine starch, with the powder of chalk or lycopodium, or with calomel, which, in our hands, has answered best of all, or else be well anointed with some mild ointment, the best of which is, in our opinion, Goulard's cerate. The washing ought to be performed with a fine soft sponge and warm water. The sponge is far better than the cloth generally employed, because, with the former, the cleansing can be effected by pressure, whilst with the latter it is necessary to use a kind of wiping or rubbing process, which cannot fail to irritate the inflamed and tender surfaces.

When the surfaces have become excoriated or ulcerated, attention to cleanliness is as important as ever. The application of the drying powders generally employed by the public becomes, under these circumstances, insufficient, and often rather injurious, except, indeed, in cases in which the excoriation is very slight; here the lycopodium powder, or very fine starch or magnesia will sometimes answer a good purpose. When the excoriation is severe, and when ulceration is present, we have never obtained any good effects from powdering; on the contrary, it has often proved injurious, and is at least troublesome and annoying from the incrusting of the powder about the ulcer. We prefer, therefore, when ulceration is present, to dress the part with simple cerate, Goulard's cerate, Turner's cerate, or with ointment of oxide of zinc. The ointment should be applied on a fine rag greased on one side, the rag being doubled and interposed in such a way between the opposite surfaces of inflammation as to be accurately applied to the whole extent of the disease, and thus prevent all friction or even contact of the opposite sides. These compresses ought to be changed three or four times a day, and all the discharges gently but carefully washed off by *pressure* with the sponge between each change of dressing.

Whilst this topical treatment is being carried out, constant attention must be paid to the state of the digestive function. It is scarcely necessary to apply this remark to cases occurring in the course of thrush or entero-colitis; but there is another class of cases that we have met with, in which, though the intertrigo is severe and obstinate, lasting as much as two, three, or four weeks, the signs of gastro-intestinal disorder are so slight as to pass unnoticed unless carefully inquired into. Thus they may consist merely in the fact that a child has a few more stools per day than usual, or that the stools are more liquid than they should be, or that they exhibit marks of derangement of the digestive process by the appearance in them of imperfectly digested curd of milk, or by their green color and sour smell. Whatever be the character of the derangement of this function, as shown by the general appearance of the child, its appetite, degree of thirst, or the appearances presented by the stools, we should always endeavor to rectify

the disorder, and if the attempt prove successful, we shall often see the intertrigo vanish at once, while before it had resisted all the means employed for its cure.

Erythema fugax requires no special treatment. The disorder which has occasioned it is the point to which our attention must be directed, and not the eruption, which is a mere consequence.

Erythema nodosum occurs generally, as already stated, in feeble children, and is usually accompanied with constipation or unhealthy stools, and slight febrile reaction. The proper treatment is a laxative at the beginning of the attack, and again in the course of the disorder, if necessary; rest in bed, or on a sofa, which is very important; and, after the operation of the laxative, the administration of tonics, and the use of a light but strengthening diet. The best tonic, as a general rule, is quinia. If this is not liked, or if there be anything in the case to contra-indicate its employment, we may substitute the compound tincture of bark, in the dose of fifteen or twenty drops, three times a day. If the child is pale and anæmic, iron is the proper remedy. It should be given in connection with the tincture of bark, or with small doses of brandy, when the appetite is poor, and the strength and spirits of the child much below their natural level.

Topical remedies are not necessary as a general rule. When, however, the local symptoms are severe, or there is much heat or pain in the tumors, they should be kept covered with compresses moistened with some kind of mucilage, or with lead-water and laudanum.

ARTICLE II.

ROSEOLA.

DEFINITIONS; SYNONYMS; FREQUENCY; FORMS.—Roseola is a non-contagious erythematous affection characterized by hyperæmic patches, of a rosy color and of irregular size and shape, which are unaccompanied by elevations or papules, and the appearance of which is preceded and accompanied by febrile symptoms.

It is often called in this country scarlet-rash, and under that title, erroneously supposed to constitute a very mild form of scarlatina. It is sometimes called also French measles, and rubeola sine catarrho. By some writers, as Duhring, roseola is included under the general term of Erythema.

Roseola is of rather *frequent* occurrence amongst children, though more rare than either measles or scarlet fever.

There are three *forms* of the disease met with in children, roseola æstiva, roseola autumnalis, and roseola annulata. As the two former, however, present no differences of any importance, we shall describe them under

one head, whilst the latter, quite unlike the other two, requires that we should describe it apart.

CAUSES.—Roseola may occur at all *ages* of infancy and childhood, and at any *season*, but is most common in summer and autumn. It has been known to prevail as an *epidemic*, but is not *contagious*. It may attack the same individual on several different occasions, one attack not protecting from repetitions. The variolous eruptions are sometimes preceded by roseola, and in some children it makes its appearance on the ninth or tenth day of the vaccine disease. Of the various causes that we have known to produce it, the most frequent is certainly derangement of the digestive function during the first dentition. It is said also to be occasioned by sudden changes of temperature, by violent exercise, and by the use of cold drinks while the body is heated and moist with perspiration; causes which strongly indicate that the nervous system is closely connected with its production.

SYMPTOMS.—Young children who have been suffering for a few days with disorder of the digestive function, often exhibit a slight roseolous eruption, lasting twenty-four or thirty-six hours, and then disappearing. The eruption in this mild form of the disease appears suddenly, often in the course of a single night, covering the trunk or even the whole surface with numerous patches, nearly circular in shape, or in irregular, broad, and waving lines, situated close together, and yet distinct, and of a light rose color. In another, and rather more violent form, occurring especially during dentition, the eruption appears after vomiting, fever, diarrhœa, and slight nervous symptoms, or possibly after slight convulsions, with the characters above mentioned, except that the rash is deeper in color, greater in extent, and that it lasts generally a longer time—two, three, or four days. Again, in a yet more marked form, which frequently, but by no means exclusively, occurs in warm weather, when it is styled *roseola æstiva* and *autumnalis*, the eruption is symptomatic of a more definite constitutional disturbance. It begins with more or less chilliness, alternating with heat, with loss of strength and spirits, with headache, restlessness, sometimes mild delirium, and even, it is said, though we have never seen them, with slight convulsive phenomena. At the same time there is some febrile reaction, marked by accelerated pulse, heat and dryness of the skin, thirst and loss of appetite; the digestive function is shown to be deranged by the presence either of constipation or diarrhœa. After these symptoms have continued for two, three, four, or even six or seven days, the eruption appears first upon the face and neck, whence it extends in twenty-four or forty-eight hours to the rest of the body. The rash resembles very closely, in some cases exactly, that of measles; but the catarrhal symptoms are absent. It is in the form of irregularly circular and rather large patches, at first of a red, but soon changing to a deep rose color, and separated from each other by portions of healthy skin. The eruption is sometimes accompanied by itching, and sometimes by stinging pain, and the febrile symptoms generally continue, though moderated in degree, after the appearance of the rash; while in other instances the fever disappears entirely from that moment. The rash lasts between one and two or three

days, as a general rule, and fades away gradually until it has entirely disappeared. In some cases it comes and goes alternately for a week after its first appearance.

ROSEOLA ANNULATA is a curious and rather rare form of the disorder, from the singular and beautiful appearance of the bright rose-colored rings which constitute the eruption.

This variety of roseola appears in the form of rosy rings, or circles, whose centres retain the natural color of the skin. The favorite seats of the eruption are the abdomen, loins, buttocks, or thighs, or it may cover the greater part of the body. In one case that we saw, the eruption covered the face, neck, and trunk. In another it was seated upon the face, trunk, and upper extremities. The rings are at first not more than one or two lines in diameter, but they enlarge gradually until their centres measure as much as half an inch in diameter. In some instances two or three rings surround one another, the skin in the intervals between them still retaining, however, its natural appearance. The disease is, when accompanied by symptoms of reaction, usually of short duration. The cases which occurred to ourselves lasted only three days, and were accompanied by decided febrile symptoms, together with signs of digestive derangement. It sometimes assumes a chronic form, the eruption fading in color in the morning, and increasing again and causing heat of skin, in the evening.

DIAGNOSIS.—Roseola æstiva might be readily mistaken by a careless observer for measles or scarlatina, and especially for the former. We have no doubt whatever that cases of roseola are often regarded, under the title of scarlet-rash, as examples of a very mild form of scarlatina, a misapprehension which will explain some at least of the supposed instances of second attacks of scarlet fever in the same individual. This is a mistake, however, that ought not occur, and need not, if the following characters of the two diseases are properly understood. The rash in scarlatina is, in the first place, of a much brighter tint, and it is more persistent and more uniformly spread over the surface than in roseola. When we come to analyze the characters of the two eruptions, there are other distinctions between them which assist greatly in making the diagnosis. In scarlatina, the eruption is composed of very large patches, or it is absolutely uniform, and evenly distributed over large surfaces, as over the whole trunk, or over the flexor or extensor aspects of the limbs. It is seen to be composed, too, when minutely examined, of an aggregation of very minute red points, which are dotted so closely together as to present the appearance of a general scarlet blush. In roseola, on the contrary, the rash is composed of irregularly circular, crescentic, or waving patches, with portions of skin between of a natural or nearly natural color. The patches, moreover, are of a different tint from that of scarlatina, being of a deep rose, instead of a bright red or scarlet color, and they cannot, upon close examination, be resolved into the minute dotted points which make up the scarlatinous eruption. When we add to these circumstances the facts; that in roseola there is no faucial inflammation, that the pulse has not the great frequency almost invariably present even in very slight cases of scarlet fever, that

all the general symptoms are much less strongly marked, that no desquamation takes place in roseola, and that the duration of the attack is much shorter, we think we have points of difference between the two, quite numerous and marked enough, to render the differential diagnosis easy to a careful observer.

It has always seemed to us impossible to distinguish with certainty between roseola and measles by the eruption alone, and we find that MM. Rilliet and Barthez are also of this opinion (*Mal. des Enfants*, t. i, p. 732). We are told by writers that in roseola the patches composing the eruption are more distinct, larger, paler, and more irregular in shape than in measles, and that they are separated by intervals of healthy skin; but we are quite satisfied that, in some cases witnessed by ourselves, these differences were not sufficient to distinguish them. The diagnosis is to be made by attention to the following points: by the absence of catarrhal symptoms in roseola, by the slighter severity of all the general symptoms, and by the much shorter duration and greater irregularity of the initial phenomena, which latter seldom last in roseola more than one or two days, and consist of symptoms of gastro-intestinal derangement, whilst in measles they last three and almost always four full days, and consist of very strongly marked catarrhal or respiratory symptoms, with very slight signs of gastro-intestinal derangement.

Roseola annulata is so peculiar and characteristic in all its appearances as to prevent its being mistaken for any other disease that we are acquainted with.

PROGNOSIS.—Roseola is never dangerous to life. If it ever seems to be so, it must be in consequence of its occurring in connection with severe internal disease.

TREATMENT.—The only treatment necessary in roseola is attention to diet; the correction by that means, or, if necessary, by a mild laxative, by some antacid preparation, or by a mercurial dose, of the gastric or intestinal disorder; rest in bed, or seclusion in a chamber with a properly regulated temperature; and the use of mild diaphoretics and cooling demulcent drinks.

ARTICLE III.

URTICARIA.

DEFINITIONS; SYNONYMS; FREQUENCY; FORMS.—Urticaria is an erythematous affection, characterized by hard elevations upon the skin, of uncertain size and shape, and of a reddish or whitish color, or, more frequently, partly red and partly white; the eruption is generally of short duration, is almost always accompanied with intense heat, and violent itching and burning.

The affection may be idiopathic or symptomatic. In the first case, the wheals seem to exist as the sole disease present, though signs of gastro-

intestinal disease may exist ; in the latter case, it is preceded and accompanied by more marked digestive or systemic disturbance, or else it occurs either during or after some specific disease, as scarlatina, measles or whooping-cough ; or finally it is secondary to some other skin disease, as scabies.

Its most common title is that of *nettle-rash*. The mild, discrete form of the disease is generally called in the nursery *hives*. It is sometimes described under the name of *essera*. It is of very frequent occurrence amongst children in a mild type. We have seldom seen in early life the abundant and severe eruption covering the greater part of the surface, which is met with in adults.

Urticaria occurs both in an acute and chronic form, the latter being rare in children. The typical and most common variety of the acute form in the adult is *urticaria febrilis*, which includes many of the unnecessary subdivisions that have been made. It is also not uncommon in childhood, though the form which is most frequently met with there is urticaria papulosa, often called lichen urticatus. Among the subdivisions of chronic urticaria, those known as *evanida* and *tuberosa* appear to occur only in adults. We have never met with an example of either earlier than the twentieth year.

CAUSES.—Children possessing a fine and delicate skin, especially when they are at the same time endowed with a highly nervous temperament, are particularly predisposed to attacks of urticaria. Very slight disturbances of the gastric functions, a very warm day, or excessive clothing, will cause an attack in such subjects ; while in many others the disease is never seen under any circumstances. Tilbury Fox (*op. cit.*, p. 126) dwells upon the important part that mal-hygiene, uncleanness, and bad air, play in the production of urticaria in children of the lower classes. Amongst the most frequent causes may be mentioned the functional disorders of the digestive apparatus which occur in the spring and summer seasons, the influence of dentition, derangement of the gastric functions from the use of improper food, and lastly, the ingestion of certain articles of diet which have been proved by long experience to be apt to occasion attacks of the disease. Of the articles last referred to, those which most frequently produce this effect are crabs, the eggs of particular kinds of fish, certain crayfish, and some kinds of smoked, dried or salted fish.

Certain external irritant and poisons to the skin are capable of producing urticaria in a marked degree ; thus we have seen it in children caused by the stinging nettle, mosquitoes, and bedbugs.

As to the intimate pathology of the affection, it seems evident that the vaso-motor nerves play an important part in the production of the wheals and large papules characteristic of urticaria. The process appears to be an acute hyperæmia, seated for the most part in the papillary layer of the skin, and leading to a sudden and intense œdema of the affected spots. The irritant, whether applied internally or externally, acts on the unduly sensitive nerves of the skin and causes spasm of the vessels and muscular fibres, which is followed by paralytic dilatation with rapid effusion. The pressure caused by the interstitial œdema empties the bloodvessels at the

centre of the wheal, making it very pale in color, while there is a reddish congested areola.

SYMPTOMS.—The most common form of this disease met with in children is known as *urticaria papulosa*, because the wheals are rather small and assume the appearance of papules. It has also been called *lichen urticatus*. It is often idiopathic, dependent on mal-hygiene or cutaneous irritation, and unassociated with fever or marked signs of disorder of the general health. It is most commonly met with among children of the lower classes.

The eruption consists of large inflamed papules, which are irregular in shape, being either rounded or oblong, projecting most in the centre, and which appears suddenly, without any or with only slight prodromic symptoms. The papules are of a bright red color, excepting in their projecting central portions, where they are whitish or of a very pale red tint. The eruption is accompanied with a smarting and burning pain, and with the most violent and annoying itching, which the child endeavors to allay by frequent and often rude scratching, in consequence of which the summits of the papules are often torn and present little crusts of dried blood. It is very fugacious in its character, appearing suddenly, lasting for a few hours or several days, and then disappearing entirely, or recurring again after a short time in the same or in new places. It terminates finally, after from a few days to several weeks, by resolution or by a slight furfuraceous desquamation. The most common seats of the eruption in children and the face, about the buttocks, or upon the thighs, or upper part of the arms.

This is the form of the disease we have met with in infants, and in children under two and three years of age. It is, as already stated, of very slight consequence, being merely annoying and never dangerous. In young infants it occasions sometimes much crying and irritability, which can be explained only by the discovery of the eruption.

The *urticaria febrilis* is usually, but not always, preceded for a few hours or two or three days, by feverishness, and by more or less marked signs of gastric disorder, such as nausea, chilliness, headache, and languor. In other instances the fever and the rash occur at the same time. The eruption begins with a sense of itching, and with heat and burning of the skin, and soon after there appear on the shoulders, loins, inside of the arms, and about the thighs and knees, reddish and solid elevations, irregular in outline, but generally roundish or oblong. The latter shape is the one the elevations most frequently assume, and it is from the resemblance which they bear in this form to the marks left by the stripes from a rod or whip-lash, that they are often called wheals. The elevations project a good deal above the surrounding surface, forming knots or ridges; their size is variable; they have hardened edges; they are reddish in color, except over the central and most projecting part, which is generally, and always when the swelling is considerable, whitish in its tint; and they are surrounded by a narrow areola of a bright red or scarlet color. The amount of the eruption is very uncertain, the elevations being sometimes separated by considerable intervals of healthy skin, while in severe cases they are

extremely numerous, and from their confluent character in such attacks, give to the part upon which they are seated, a nearly uniform red color, and occasion at the same time a very decided puffing and swelling of the skin.

The eruption, when at all considerable in degree, is attended with violent itching and burning. The former is often so severe and troublesome as to occasion the most distressing irritation to the patient, precluding all comfort and quiet. It is increased by heat, and especially by that of the bed. The patches of eruption which appear first do not continue throughout the disease, but, after lasting from a few minutes to a few hours, fade away, and are replaced by new and successive crops. During the attack, the patient is usually more or less feverish, and he suffers from languor, loss of appetite, furred tongue, and the usual signs of gastric derangement. The symptoms subside gradually, so that, after a period varying from two or three days to a week, the disorder has entirely disappeared, leaving behind no traces, except, in a few instances, a slight desquamation.

When this form of urticaria follows the ingestion of certain articles of food, the eruption usually appears within a very few hours after the meal, being preceded and accompanied by nausea or vomiting, pain and distress in the epigastric region, giddiness, headache, and feverishness.

In cases where the cause persists, and especially among children of the poorest classes, who live in squalor, or at least under any unfavorable hygienic conditions, the disease may assume a chronic form. The individual wheals are most persistent, and the rude scratching and absence of cleanliness aggravate the eruption and modify its character.

DIAGNOSIS.—There can be no difficulty in recognizing a case of urticaria. The peculiar characters of the eruption, and especially the size, shape, and color of the solid elevations of which the patches consist, the violent itching and burning which accompany it, and its fugacious character, render it unlike any other cutaneous disease, and ought to prevent any mistake as to its nature.

PROGNOSIS.—Urticaria is probably never dangerous in children. If it be accompanied by symptoms of a threatening or alarming character, these are dependent rather upon the gastric disorder, which is the cause of the urticaria, than upon the latter affection itself. We have never known it to be more than troublesome and annoying.

TREATMENT.—There are but two really important indications for the treatment of this disease: to attend to the state of the digestive functions; and to allay, by proper means, the distressing irritation occasioned by the itching and burning of the eruption.

In the mild form of urticaria, called in the nursery "hives," and in scientific language, urticaria papulosa, the only treatment necessary is careful regulation of the diet, and the use of means proper to correct any evident derangement of the digestive functions. The food should be light and digestible, but at the same time nourishing. Milk, bread, light meats, and the plainest vegetables, form the proper diet for children over three years of age. Under that age, milk preparations, bread, and in those over a year old, light broths, ought to constitute the diet. In a large

majority of such cases, no drug whatever ought to be given. The only ones likely ever to be required are occasional mild laxatives or gentle mercurials, when constipation is present; and some of the antacids, as very small quantities of magnesia or carbonate of soda, or lime-water and milk, when the stomach is acid. To allay the itching and consequent restlessness of the child, the patches of eruption should be well and frequently dusted with toasted rye or wheat flour, which are often very successful. Washing the eruption with salt and water, or with brandy or whiskey diluted with water, when the cuticle is not broken, is sometimes very soothing, and, when the patches are of small extent, may be frequently repeated. Various lotions may be used with advantage; as, for instance, one composed of a drachm each of carbonate of ammonia and acetate of lead, and eight ounces of rose-water; or of carbolic acid with water, a drachm to the pint; or of benzoic acid and borax, of each five grains, to a pint of water; or of corrosive sublimate, two to four grains to a pint of water.

In the urticaria febrilis the treatment must depend upon the cause of the attack. When it follows upon the eating of some unwholesome food, we must rid the stomach of the offending substance by an emetic, unless nature has already caused its rejection by spontaneous vomiting. When this end has been accomplished, it will be proper to give some kind of cathartic medicine, and the best is castor oil, as the mildest and most certain, in order to insure the discharge of the whole of the aliment which has been causing the mischief: or small doses of blue pill; or hydrargyrum cum cretâ, with rhubarb, where there are present any signs of hepatic derangement. After this the only treatment necessary will be the use of cooling and demulcent drinks, containing perhaps a little sweet spirit of nitre; rest in bed, or at least seclusion in the house, for a few days; and careful regulation of the diet. The latter ought to be very light during the continuance of the eruption, consisting merely of milk and bread, or of some kind of gruel or plain broth; after the cessation of the disease, it should be augmented only with due care and quite gradually. To allay the itching and burning of the eruption, and the general distress of the child, the best remedy is a warm bath carefully administered. This may be repeated in six or eight hours if necessary, and there may be added to the bath an alkali, or some starch or bran.

As the children who are subject to urticaria often present evidences of defective nutrition or impaired nervous tone, tonics should be administered after the acute symptoms have subsided. Quinia with small doses of a mineral acid, is one of the most valuable, especially in cases where a tendency to recurrence of the eruption is manifest. If the disease is at all chronic, arsenic and iron are frequently of service,—provided that no undue irritability of the stomach exists to contra-indicate their use.

CHAPTER II.

VESICULAR OR CATARRHAL INFLAMMATIONS OF THE SKIN.

THESE are characterized by hyperæmia, with serous effusion into the corium, together with the escape of leucocytes into the same tissue, tending to give rise to sero-purulent discharge and crusting, though sometimes terminating in simple desquamation.

They comprise eczema and herpes.

ARTICLE I.

ECZEMA.

THE term eczema is no longer restricted to a disease characterized by the formation of vesicles, but embraces all the numerous affections which present redness of the skin, frequently punctated, itching, infiltration, and exudation on the surface, with the formation of crusts. So far, indeed, from vesicles being characteristic of it, it may be said, and especially in regard to eczema in children, that its rarest form is that which is attended solely with their formation. The elementary lesions which may be present at the beginning of the attack, are either erythema, papules, vesicles, or pustules, and the disease is divided accordingly into eczema erythematosum; eczema papulosum, which embraces eczema lichenoides, and eczema prurigiosum; eczema vesiculosum, the typical eczema of Willan, one of the rarest of all its varieties; eczema pustulosum, or impetiginoides, which includes impetigo; and eczema squamosum, which is usually of the chronic form, and resembles, in many cases, psoriasis. It is indeed called psoriasis by Dr. Wilson, who gives the name "*alphos*" to that scaly disease, which is still, by most authorities, and especially by Hebra, designated as psoriasis.

It not unfrequently happens, also, that the various elementary lesions enumerated above may be present at the same time on a patch of eczematous eruption, so that a case which has begun as eczema erythematosum, or vesiculosum, may present the development of papules or pustules, or thick scabs, and thus become converted into the pustular or squamous form. This tendency for the blending of several elementary lesions in the same eruption, and especially for the conversion of the eruption into the pustular form, is very markedly seen in cases of eczema of children.

Eczema is also divided, according to its course, duration, and stage, into acute and chronic.

Eczema shows, moreover, an especial tendency to attack certain parts of the surface, and presents various peculiarities in the different localities; in children, it frequently occurs on the scalp and face, though it extends over the entire surface of the body far more frequently in them than in adults.

The special forms of eczema which will be here described, are simple

acute eczema; eczema of the scalp, and of the face; eczema pustulosum, or impetigo; eczema papulosum; and chronic eczema, or eczema squamosum.

CAUSES.—Eczema is by far the most common of all diseases of the skin in this country, as shown by the statistics collected by Duhring (*op. cit.*, p. 165), and it is not improbable that climatic influence may account for something in its production. There are also certain constitutional or general causes that predispose to the development of eczema; thus it seems children of scrofulous or tuberculous diathesis are specially liable to it. All influences that impair nutrition and lessen the vital resistance of the tissues must be regarded as favoring the occurrence of eczema. It is generally conceded that the nervous system exerts a very powerful, though not altogether demonstrated, control over the nutrition of the skin, and it is probable that such depressing causes as specially impair and irritate the nervous system predispose most strongly to this affection. Exposure to bad hygienic conditions, or want of cleanliness, insufficient or improper food, crowded or ill-ventilated habitations, extreme and continued heat, and sudden atmospheric changes, may be mentioned as among the most powerful of such influences. Irregularities or indiscretions in diet, alteration in the quality of the mother's milk, and unsuitable artificial food, operate so frequently and so potently as to demand special mention. They predispose to eczema, as they do to many other affections, but they may also serve as exciting causes, either by causing the entrance of imperfectly digested and, therefore, irritating substances into the blood, or possibly by the reflex influence of the irritated gastro-intestinal mucous membrane. So, too, the relation of dentition to eczema demands special mention, since the majority of cases of this affection in children occur during either the first or the early part of the second dentition. Undoubtedly the disturbing influence which this process exerts on the nervous system and general nutrition of many children so powerfully predisposes to eczema, that the most trifling exciting causes suffice to develop the eruption, which is subsequently maintained by the irritated and enfeebled state of the system. It is a mistake, however, to assume that dentition acts also as the direct and exciting cause of eczema, excepting, perhaps, in cases where the irritation of the nervous system is unusually severe and prolonged, so that the innervation of the skin is seriously perverted. It is highly important to recognize this truth, as it is altogether too much the custom to refer skin diseases, as well as other affections, in teething children to the irritation of dentition alone, without searching carefully for the presence of some of the other causes that produce such disorders at other periods of life.

Vaccination occasionally serves to develop eczema in children who are predisposed to it.

Many local causes act in the same way, but it is only necessary to mention among them the direct action of excessive heat or of extreme cold, the inordinate use of water in bathing, especially if conjoined with the use of alkaline soap.

PATHOLOGY.—In order to fully appreciate the clinical features of

eczema, it is desirable to consider the anatomical changes that occur in the affected portions of the skin. These have been carefully studied by Neumann (*Lehrb. d. Hautkrankheiten*, Wien, 1873) and by Biesiadecki. In the acute stage of the disease there is, in the first place, extreme congestion of the capillaries, which soon becomes associated with effusion of serum, chiefly into the papillary layer. According to the intensity of the morbid action, probably, the effusion is either serous or mixed with a varying amount of plastic material. And there is also a varying amount of migration of white blood-corpuscles from the vessels. At the same time, the papillæ are decidedly enlarged from serous effusion and from proliferation of the cells. There is also a development of spindle-shaped connective tissue corpuscles, extending up into the rete mucosum, over and between the affected papillæ. It occasionally happens that if both the serous effusion and cell development are very slight, they may subside without leading to any further changes or to any discharge. When this cell-infiltration is considerable, the eczema assumes the papular form; while, when the cells of the more superficial layers are greatly swollen and distended, and the free liquid effusion is at the same time abundant, the epidermis is raised, so that vesicles of varying size are formed. When these vesicles rupture, even the deeper layers of the corium may be exposed, and a discharge occurs of viscid, yellowish, serous liquid, which stains and stiffens linen, and, when exposed to the air, dries and forms yellowish crusts.

In chronic eczema, the further stages of these lesions are observed. The layers of the skin are indurated, thickened, and infiltrated, with imperfectly developed cells. The papillæ are remarkably enlarged and prominent. Both the bloodvessels and lymphatics have been found enlarged, but in other cases, owing to proliferation of the cells in the sheath of the bloodvessels, and to the hyperplasia of the intervening tissue, the channels are greatly narrowed. There is almost constantly an excessive deposit of pigment, especially along the course of the affected vessels.

As regards the essential cause of the above phenomena, we have already expressed an opinion that in eczema, as in many other inflammatory states, there is a faulty state of the nervous supply, together with the local morbid condition of the vessels and cells of the affected parts.

SYMPTOMS.—We have already alluded to the fact, that in the eczema of young children, as indeed is true, to a less degree, of the disease at all ages, we constantly meet with the most varied forms of eruption in the same case; and have the opportunity of watching the development of papules, vesicles, or pustules, until a case which has begun as one of erythematous eczema presents the characters of the papular, vesicular, and ultimately of the pustular form. The predominance of one or the other of these typical forms of eczema is determined by the temperament and general condition of the child, and the grade of inflammatory action present.

Eczema simplex, or *vesiculosum*, may occur on any part of the body, but in children is most frequent on the face and arms. The eruption appears, without any precursory symptoms, as an erythematous patch, which is red

and itchy, and may present slightly raised pimples, and being rubbed and scratched, soon presents the formation of numerous, closely aggregated, exceedingly minute vesicles, containing a transparent limpid serum. After a short time the contained fluid becomes turbid and then milky, and is either absorbed, while the vesicles shrivel up and disappear by a slight desquamation, or else the fluid escapes by the rupture of the vesicles, and little thin scales follow, which are detached before long from the surface beneath. The eruption is attended with more or less itching and smarting, but does not generally give rise to constitutional symptoms. The vesicles are generally renewed by successive crops, so that, though the case may terminate in from two to three weeks, it is apt to continue for two or more months.

Eczema papulosum, or Lichen simplex.—In the same way as the above form of eczema is characterized by the formation of vesicles, there are other cases where the eruption principally consists of small reddish papules, associated with erythematous patches. The papules in infantile papular eczema are small, from the size of a small to a large pin-head, and may either be firm and acuminate, or softer and more rounded or flat. They may be either discrete or confluent. The papules either continue as such throughout their course, or some of them undergo a transformation into vesicles. It is this clinical fact that, while true papules predominate, imperfectly formed papules, half-developed vesicles, or even typical vesicles may appear, which proves that the vesicular and papular varieties of eczema are really manifestations of the same morbid process. Duhring states (*op. cit.*, p. 162) that papular eczema attacks by preference the arms, trunk, and thighs, especially the flexor surfaces. It is an obstinate form of the disease, and, as the itching is severe, the papules are often torn by the scratching, so that their summits are covered by small blood crusts.

Eczema pustulosum or impetiginosum.—Under this head we will describe the affection usually styled impetigo, and formerly classed among the pustular diseases of the skin, but which possesses peculiarities which have induced many dermatologists to transfer it to the group of eczematous affections.

It may be described as a form of eczema characterized by the production of pyodermic sero-pustules, containing a thin purulent fluid, which either break and discharge, or dry up and form thin amber-colored or more thick yellowish-brown crusts.

The eruption usually begins as a reddened patch, studded with slightly raised pimples. As the inflammation increases, the cuticle is often raised into more or less well-defined vesicles, or the surface becomes excoriated, and there is a discharge of turbid or whitish-yellow secretion; the skin now becomes infiltrated, and numerous rather small pustules, containing a light colored pus, form on the red swollen surface. Not infrequently there are vesicles on the same patch, surrounding its margin. These pustules are usually broken by scratching or by friction against the clothes, and their contents dry up, forming amber-colored or brownish crusts. Frequently, also, blood is mingled with the discharges, and the crusts become dark-colored, or at times positively black.

The crusts separate in a few days, the time varying according to their firmness and thickness, and leave the surface reddened, but without any permanent scar. Frequently, however, the disease passes into a chronic form; the eruption retreats to certain seats, as the scalp, or the flexures of the joints, where the skin remains somewhat infiltrated, while the cuticle is rough, scaly, and constantly desquamates, either in the form of a fine furfuraceous exfoliation, or of scales of considerable size.

There are in reality but two specific varieties of this form of eczema, *impetigo figurata* and *impetigo sparsa*, so named from the manner in which the pustules forming the eruption are arranged. When the former variety occurs on the trunk or limbs, it usually presents a large eruptive surface. On the arms, we have seen it extend from the shoulders to the hands, and, as a general rule, it has been most severe on the outer portions of the limbs. On the trunk and legs, it has usually affected surfaces of much less considerable size, and has commonly appeared in a patch of an irregularly oval shape, and of four, five, or six inches in diameter.

Impetigo sparsa is quite a common affection in children of all ages. Here the pustules, instead of being confluent or grouped closely together, appear singly or in small clusters. It most frequently appears on the face and scalp, but is also met with on the extremities, being not infrequent, according to Wilson, on the hands and feet.

The great frequency and severity of *eczema pustulosum* as it appears on the scalp or face in young children, makes it desirable to give a brief description of these two local forms.

Eczema capitis is often met with in infants at the breast during the first dentition, and at later periods of childhood in those who are scrofulous, or who are placed in unfavorable hygienic conditions. It may be confined to a small portion of the scalp, or it may cover the head, and extend to the face and neck; or again, it may be limited entirely to the latter localities, when it constitutes eczema of the face. In both cases, the eruption is very apt to run into the pustular form, constituting the disease known as *impetigo capitis*.

When mild in its features, it consists of an eruption of numerous small vesicles or sero-pustules, spread over certain portions of the scalp, to which it may remain limited; or it may cover the face at the same time, or it may attack alone the forehead, temples, and, perhaps, portions of the cheeks. It is attended, under these circumstances, with very slight redness and heat of the integument. The sero-pustules discharge their fluid contents and form thin crusts, which gradually fall off, leaving slightly reddened or excoriated surfaces, which soon disappear, or are followed by fresh crops of eruption, destined to pass through the same changes as the preceding ones.

In more severe cases the disease may be confined either to the scalp or face, or it may, as stated above, exist upon both simultaneously. The eruption presents different appearances in these two situations.

When seated on the scalp, it is often called by the English, *milky crust* or *milk-crust*, *crusta lactea*, *tinea lactea*, and *porrigo larvalis*; and by the French, *croûte de lait* and *gourme*.

On the scalp, as already said, the eruption may be either partial or gen-

eral. It may consist at first of disseminated minute vesicles, which break, and form thin lamellated crusts, of a yellowish or brownish color; or of pustules, yellowish-white in color, and of small size, seated on an inflamed base. The surface affected is at first small, but the eruption gradually extends to surrounding parts. It is attended with great heat and itching; and, as the disease advances, the scalp becomes very much inflamed, red, tense, swollen, and painful. The eruption is now more completely pustular, and as the pustules open or are torn by the uncontrollable scratching, they discharge an abundant thin sero-pus, or even a thick and viscid fluid, which glues the hairs together, and hardens into uneven brownish-yellow crusts. If the scalp is not kept clean by constant washing or by emollient applications, the crusts increase rapidly in thickness by successive discharges of fluid from the pustular surface beneath, until at length the whole of the diseased part is covered with thick, heavy, rough, and adherent crusts, of a brownish or yellowish-white color, or at times of a positive black from the admixture of blood which oozes from the inflamed surface, torn by the nails of the little sufferer.

When neglected, the crusts become more and more thick, and from the heat of the head and exposure to the air, they undergo partial decomposition, and exhale a fetid, sickening odor, of the most disgusting kind. Among the children of the poor and destitute, lice often form in abundance, and add to the repulsive character of the disease. At first, the crusts are somewhat soft and moist, from the percolation through them of the fluid exuded beneath; but as they become more abundant and thicker, their outer surface becomes dry and sometimes very friable. The secretion from the inflamed surfaces often makes its way under the crusted mass above, and, flowing down over the forehead and behind the ears, irritates the parts that were before healthy, and thus extends the disease.

When the crusts are removed by any means, the surface of the eruption is found to be red, shining, wet, and discharging an abundant purulent or sero-purulent fluid, which escapes from minute excoriated points, dotted thickly over the inflamed scalp. The scalp is at the same time tumefied, tender to the touch, and abscesses may form beneath it. The lymphatic glands, as the occipital, submental, or cervical, are frequently enlarged, and at times suppurate.

When the disease has lasted a considerable length of time, it tends to assume a chronic form. The inflammatory action extends to the hair-follicles, and often occasions partial loss of hair over larger or smaller surfaces. This kind of alopecia is not, however, permanent. The hair-bulbs are not destroyed, but merely inflamed, so that the hair grows again after the cure of the disease. The tissues of the scalp remain thickened, but the amount of the secretion diminishes; and the painful irritation and itching are less troublesome. Under these circumstances, the crusts are less thick and massive; they become lighter, thinner, and are more easily detached. The epidermis is dry, uneven, and rough, and there is a continual desquamation of fine furfuraceous particles, constituting a form of pityriasis capitis, or of epithelial scales of various sizes, resembling a case of psoriasis.

On the *face* (*Eczema faciei*), the disease usually shows itself first on the forehead and cheeks, to which parts it may remain limited, or whence it may extend to the lips, chin, ears, and neck. The nose and eyelids are seldom attacked, though we have occasionally seen the upper eyelids slightly affected.

The disease begins by the appearance of minute vesicles or sero-pustules on a patch of reddened and slightly swollen skin; there is also excessive pruritus. When the eruption is scanty, and rather vesicular, and the degree of inflammation slight, the cuticle breaks, and there is a discharge of a thin, turbid, serous fluid, which dries into delicate scales, or thin lamellated crusts.

When the accompanying inflammation is more severe, however, the eruption is more truly pustular, the pustules being numerous and rather large, and the discharge copious, so that when the formation of crusts is not interfered with by topical applications, or by the scratching of the child, large portions of the affected surface become covered with thick yellowish, brownish, or brownish-red crusts, which present the general appearance of a mass of incrustation, broken by cracks and fissures into portions of very irregular size and shape.

In the milder cases, when the scales drop off, the skin may appear reddened and moist, or may seem to be covered with a very delicate, shining epidermis, which is perfectly dry or presents tiny drops of serum or minute cracks. In the more severe cases, if the crusts are detached from any cause, the skin beneath appears red, swollen, inflamed, and wetted with a more or less abundant sero-purulent fluid, sometimes mixed with blood, that oozes from numerous small points on the excoriated and inflamed surface. The eruption is attended with severe itching and smarting, to relieve which the child often tears the affected surface with the nails, so as frequently to remove the crusts, wound the skin beneath, and cause more or less bleeding from the part.

In this more severe form, when the discharge forms a thick discolored scab covering the scalp or face like a mask, the disease has received the names of *porrigo* or *impetigo larvalis*, which are less accurate than *eczema larvale*. It corresponds to the *impetigo figurata*, as met with in other parts of the body.

When the eruption is more scanty and developed in small groups on the scalp alone, the discharge is less copious, and soon concretes into dry, friable, brownish crusts of irregular shape, some of which are very adherent, matting together a larger or smaller number of hairs, while others are broken into small and dry fragments, which have been compared to particles of mortar dispersed among the hair. Many of the pustules in this variety are formed at the base of the hairs, so that these particles of crust, being pierced by the hairs, have somewhat the appearance of a string of rude beads. This form of the disease has been known as *tinea*, or *impetigo*, or *porrigo granulata*; but for the sake of uniformity, it might be styled *eczema granulatum*. It corresponds to the *impetigo sparsa*, as met with on other parts of the body.

Eczema larvale, whether confined to the scalp or face or existing on both parts at once, causes, when it exists in the acute form, much distress and annoyance to the child. The heat and tension of the part, and particularly the itching, occasion much restlessness and irritability; they make the child cross and peevish, disturb its sleep, and sometimes cause slight febrile attacks, which debilitate and injure the health. Indeed, when the disease has lasted a considerable time, it often induces extreme anæmia and impairs severely the general nutrition of the child. In other cases, however, the general health remains perfect,—all the functions of the body going on well, notwithstanding the local distress and irritation. The lymphatic glands situated behind and in front of the ear, and those on the back and front of the neck often inflame, enlarge, are frequently hard and painful to the touch, and in a few instances suppurate, though the latter occurrence is not frequent.

The *duration* of eczema larvale is very variable in different cases. Mild cases, and particularly those in which the eruption is confined to a limited extent, often get well, or are readily cured in two or three months. When, on the contrary, the disease is severe and extensive, the duration is much longer, seldom less, according to our experience, than several months or even one or two years. In most cases, however, the intensity of the disease varies from time to time, so that at one period it may seem to be subsiding rapidly, or it may even disappear almost, or be very greatly ameliorated, only to break out again with renewed violence under the influence of some exciting cause, as the cutting of new teeth, some change in the weather or season, or some alteration in the health of the child which cannot be explained. This affection is, as already stated, almost entirely confined to the age of dentition. The disease often begins some months before the appearance of the first teeth, and though it generally ceases or is cured before the termination of dentition, we have known it to run on unchecked three months after the conclusion of that process and then to be removed only by medical treatment.

Eczema granulatum is comparatively a slight disorder, and is usually much more under the control of remedies and of much shorter duration.

There is a local variety of eczema which requires a brief allusion. It is known as eczema tarsi and affects the edges of the eyelids, especially in strumous children, in whom it is often associated with strumous ophthalmia. It is attended with the formation of pustules at the openings of the hair-follicles, itching, thickening of the eyelids from infiltration, the formation of crusts, and a tendency to adhesion of the edges of the lids together, especially in the morning after they have been in contact during sleep. If not cured by appropriate treatment, it frequently leads to distortion of the hairs, which assume abnormal directions in their growth, and to inversion or eversion of the lids.

Eczema Chronicum.—Eczema infantile, if left to itself, has no natural tendency to cure, but usually becomes chronic, as in the adult.

When the disease passes into this form, no matter what may have been the original type of the eczema, the eruption gradually assumes uniform and characteristic appearances. The various forms which have already

been described are then to be regarded as varieties of acute eczema, while the chronic form is common to them all, and represents the condition into which all the acute varieties may merge.

The skin in chronic eczema is either very much inflamed and thickened, presenting excoriations with deep cracks and fissures, which pour out an abundant ichorous secretion, or, more frequently, the inflammation is less severe, there being much less heat, redness, and infiltration of the skin, fewer excoriations and cracks, and a smaller amount of effusion. The affected surface is, in these cases, dry and parched, and constantly throws off a fine furfuraceous desquamation, as in pityriasis, or scales of dried cuticle of various sizes, as in psoriasis.

This form is most common on the scalp, behind the ears, about the neck and upper part of the trunk, and in the flexures of the joints. It usually lasts for months, and is difficult of cure. It is attended with severe itching, which is sometimes so troublesome as to occasion the most distressing and uncontrollable restlessness at night.

Not rarely also, on the application of any exciting cause, the eruption will spread from the spots where it has been lurking in the chronic form, and invade more or less of the surface, assuming all the appearances of acute eczema.

It is very important that the correct principles of studying cutaneous diseases should be carefully applied in forming the diagnosis of an affection of such protean character as eczema.

It must be borne in mind that its characteristic symptoms, which are present in varying proportion in nearly every case, are redness and infiltration of the skin, with some œdematous swelling; itching of a peculiarly intense character; and liquid exudation on the surface, with the formation of crusts.

Acute eczema may occasionally suggest scarlatina, but the absence of high febrile action, of sore throat, will prevent mistakes; and very soon the eruption will present its characteristic appearances.

From erysipelas, with which it may much more readily be confounded, eczema may be distinguished by the less degree of constitutional disturbance, and by the absence of the great thickening, with distinctly elevated margin of the erysipelatous patch, and of the peculiar mode of spreading of the latter eruption.

Eczema simplex when seated on the hands and between the fingers, may be mistaken for scabies. The distinction can, however, be made by attention to the following points: the vesicles of eczema are flattened and aggregated; in scabies they are acuminate, isolated, and entirely distinct. There will also be frequently found, in scabies, vesicles on the hips where the hands of the nurse from whom the child has caught the disease has been placed to support it. In scabies, also, the vesicles present little red lines, running off from their margins, and marking the course taken by the acarus; and lastly, in that disease careful search will almost always enable us to detect the insect or its ova, which are infallibly characteristic of the disease.

From sudamina, with which eczema vesiculosum might perhaps be con-

founded, the latter disease may be distinguished by the facts that the vesicles constituting sudamina are much larger, that they are discrete and scattered, that they are associated nearly always with profuse perspiration, and that they are unaccompanied by an inflammatory state of the skin or by itching.

Psoriasis is often confounded with eczema in its chronic scaly forms. The history of the case will often solve the question by showing that discharge occurred at an early period of the eczema. The scales in psoriasis are larger, more purely epithelial, silvery, and imbricated, while in eczema they are thin, yellowish, and scanty.

Pityriasis rubra may in like manner be distinguished from eczema by the abundant large, whitish, papery epithelial scales, and by the absence of discharge or of marked infiltration of the skin.

Eczema impetiginosum, especially when affecting the scalp, might possibly be mistaken for favus, from which, however, it may readily be distinguished by the facts that, in the latter disease, the pustules are imbedded in the epidermis, and that the crusts present a peculiar bright yellow color, and are of an umbilicated or cup-like shape. Favus is also followed by incurable alopecia and is contagious, and microscopic examination will detect the peculiar fungus, the *achorion*, upon which it depends, in all of which circumstance it differs entirely from eczema.

Tinea carcinata is occasionally mistaken for the squamous variety of eczema. But the history of *tinea* will often show its contagious nature; the patches of eruption are circular and sharply defined; and the microscope will reveal the presence of the peculiar fungus.

PROGNOSIS.—Eczema infantile is rarely dangerous to life, though it sometimes occasions much distress to the health by the suffering, irritation, and especially by the loss of sleep, which it entails. In one instance, however, that came under our observation, of very severe eczema larvale combined with extensive impetigo figurata, in a child a few months old, the disease ended fatally some weeks after the child has been put under the charge of a homœopathic practitioner.

In the prognosis given by the physician, especially in the instance of extensive eczema pustulosum, he should never forget to refer to its probable long duration, and to its disposition to return even after an apparent cure has been effected. It often lasts, in this way, for many months, and sometimes for one or two years or even longer. This difficulty of cure, and obstinate tendency to recur, are often owing to its dependence on some constitutional disturbance, or upon derangement of the digestive system. It ought, therefore, to be looked upon as an expression of a general disorder, and its cure will at times be found to depend upon the removal of the constitutional fault.

It is on this account that the opinion has long been popularly entertained, that extensive eczema should not be treated by severe local remedies, since, if suddenly arrested by such means, the disease might fall with all the greater severity upon parts more important to life.

If proper attention be devoted, however, to the removal of any underlying constitutional disturbance, there can be no danger in using suitable

local remedies to effect as rapid a cure as possible. In a general way, therefore, eczema may be said to be always curable. But in forming our opinion as to the probable duration of the disease, we must carefully estimate the general and local conditions that may prolong its cause.

TREATMENT.—The remarks which have been already made in connection with the causes and constitutional character of many skin diseases, will readily suggest the indications which are to be followed in treatment. It is necessary to remove the constitutional disturbance which may be the essential cause of the affection, to allay the local distress, and to promote the healthy vigorous nutrition of the skin. Our own observation has convinced us that the most rapid and certain cures can only be effected by a judicious combination of general and local remedies, either of which, however, may, under special circumstances, assume peculiar and paramount importance.

The general treatment of eczema must depend on the state of health of the patient at the time, on the extent and activity of the eruption, and on its acute or chronic character.

In mild cases, which show but little disposition to extend and are not attended by much irritation, regulation of the child's diet, and the use of the same simple bland applications, will be sufficient.

When the disease is more extensive and attended with much irritation, it is necessary to examine carefully into the state of the digestive function, and if this be in any way disordered, to endeavor to restore it to a more healthful condition.

When the child is teething, the gums ought to be examined, and, if found swollen or inflamed, they should be lanced as often as necessary. The diet must be properly regulated, the food being changed if that which has been previously taken is found not to be well and completely digested.

Constipation, if it be present, must be overcome by altering the diet, or by the administration of rhubarb, small doses of magnesia, Rochelle salts, or sulphur. Purgatives have been strongly recommended by some writers in the treatment of eczema putulosum, but we should discountenance their use, save in the form of very gentle laxatives when absolutely required, since in no other form of eczema is an early resort to tonic and nutrient treatment so strongly demanded as in this.

If there are evidences of acidity of the stomach, it is well to employ some of the various preparations of the alkalies.

So also when diarrhœa is present, it should be treated by attention to the diet; and by the administration of a weak castor-oil emulsion, containing small quantities of laudanum, when the stools are feculent, but small, frequent, and attended with griping; when they are thin and watery, greenish, and composed in part of mucus, the following prescription will often prove very useful:

R.	Tr. Krameriæ,	f ʒj.
	Tr. Opii,	gtt. vj.
	Sodii Bicarb.,	ʒj.
	Syr. Zingiberis,	f ʒ vij.
	Aquæ.	f ʒ ij.—M.

S.—A teaspoonful two or three times a day, for children of one and two years old.

When the eruption has persisted for some time, and tends to become, or has actually become, chronic, resort must be had to remedies which are capable of modifying the constitutional condition of the child. In many such cases, the child shows evidences of impaired nutrition, and is weak and debilitated; so that the remedies clearly indicated are those which will tend to invigorate the general health and aid in the restoration of power.

The remedy which extensive experience has led us to regard as the most useful in all suitable cases of chronic eczema is *arsenic*. It is essential, however, that there shall be no undue irritability of the gastro-intestinal mucous membrane, or it will inevitably disagree. The preparation of arsenic which is best adapted for administration to children is Fowler's solution; which we are in the habit of giving in combination with iron, as in the following formula:

R. Liq. Potassii Arsenitis,	℥ xvj ad xxxij.
Vin. Ferri Amari,		
Syr. Tolutani, āā,	f℥ j.
Aq. Carui,	f℥ ij.—M.

Dose.—A teaspoonful thrice daily, directly after food, for an infant from six months to a year old.

We have never known any serious inconvenience to follow the administration of this remedy, the only annoying symptoms occasionally produced being slight gastric irritation and diarrhœa, and a little puffiness of the eyelids. By giving it immediately after taking food and properly diluting it, it rarely causes any gastric irritation, and even should it do so, the symptoms rapidly disappear if the remedy be temporarily suspended, or given in a smaller dose or less frequently. The mother or attendant, should, therefore, be carefully instructed to instantly suspend its administration upon the appearance of any disturbance of digestion. The puffiness of the eyelids, which is one of the earliest and most characteristic symptoms of the physiological action of arsenic, is of no alarming importance, and the remedy need not be instantly suspended on account of its appearance; though it is more prudent, at least, to reduce the dose and frequency of administration, and to watch carefully for the occurrence of any further signs of the over-action of the drug.

The period of continuance of this treatment must depend upon the state of the eruption, and the manner in which the arsenic is tolerated; if necessary, however, and if it causes no gastric irritation, it may be continued for many weeks or months.

In cases which persist despite local treatment and the internal administration of arsenic, we have frequently found the use of cod-liver oil followed by marked benefit. It may be given combined with the arsenic, or, if the stomach will not tolerate it in an undisguised form, in the form of an emulsion with aromatics, as already recommended at page 386.

In cases attended with marked anæmia and debility of constitution associated with a scrofulous tendency, we have obtained good results from the administration of the syrup of the iodide of iron. This may be given in

combination with the compound syrup of sarsaparilla, in the dose of from gtt. ij to gtt. v of the former, diffused in from a quarter to a half teaspoonful of the latter, three times a day, for children of one or two years of age.

Where the tongue is heavily coated, and the bowels constipated, with whitish or clay-colored stools, minute doses of blue pill or calomel in combination with bicarbonate of soda, may be given from time to time with manifest advantage.

The diet should be nutritious and strengthening, but at the same time, light and of easy digestion.

If the appetite is weak and capricious, tonic remedies, as compound tincture of cinchona bark, or quinia, in combination with the ferruginous preparation employed, ought to be administered.

In rare cases, when the patient is of full habit, of gross development, and of florid complexion, the diet must be somewhat restricted, and a moderate use of cathartic remedies, as small doses of saline laxatives, of sulphur, of blue pill, or of extract of taraxacum, resorted to.

LOCAL TREATMENT.—In all cases of eczema, the use of local remedies, carefully adapted to the stage and form of the disease is a matter of the greatest importance. The affected parts must be critically examined, all possible sources of local irritation removed, and minute directions given as to the exact manner in which the steps of the local treatment are to be carried out.

In the first place, whenever crusts or scales cover the affected surfaces they should be removed very gently, by first softening and loosening them by oily applications or by poulticing, and then by using tepid water with or without soap. In all stages of the disease the inflamed parts must be carefully excluded from the air; but the character of the local application must vary in accordance with the precise condition of the individual case.

In the acute stage, attended with great local irritation, the indications are to soothe the inflamed surface, and to relieve the engorgement by mild astringents. Great relief may often be obtained from the use of compresses repeatedly wet with water, either cool or hot, or with some emollient decoction, as of marshmallow, poppyheads, bran or flaxseed, sassafras pith, or slippery elm bark. Weak lotions, as of one drachm of bicarbonate of soda, or of half a drachm of borax with a little morphia to a pint of water, or of diluted lead-water or carbolic acid $\frac{1}{2}$ to 1 drachm to a pint of water, may be found more soothing in other cases. Such applications may be retained upon the part for several hours at a time or throughout the day as they may be found to suit the eruption, though care must be taken not to macerate the skin by too prolonged use of lotions. At night they may be replaced by a mild ointment or by a protective powder. These latter often given marked relief. The surface may be powdered from time to time with finely powdered lycopodium, carbonate of zinc, or with the following:

R. Pulv. Amyli,	3vj.
Pulv. Zinci Oxidi,	3ss to 3j.
Pulv. Camphoræ,	gr. xv.—M.

If powders are used, they must be very carefully removed at least once in twenty-four hours, so as to prevent the formation of hard crusts of dried discharges mixed with the powder.

But in a large proportion of cases, ointments do more good than powders or lotions. To secure this good result, they must be applied carefully and systematically. The affected part should be washed with tepid water, with or without soap, and dried by gently pressing a wad of absorbent cotton or a very soft sponge upon the surface, and then it should be covered closely by strips of old linen spread with the ointment. Among the best purely soothing salves are cucumber ointment, cosmoline, and vaseline. We prefer, however, the ointment of the oxide of zinc to any other, selecting the simple ointment when the irritation is very acute, and the benzoated ointment when the most acute stage has somewhat subsided. This may be rendered more sedative by the addition of a little camphor, as in the following formula from Duhring:

R.	Pulv. Camphoræ,	.	.	.	ʒj.
	Pulv. Zinci Oxidi,	.	.	.	ʒij.
	Glycerinæ,	.	.	.	fʒss.
	Adipis Benzoat.,	.	.	.	ʒvj.
	M. et ft. ung.				

Dr. White, of Boston, treats acute eczema very successfully by applying the lotio nigra several times daily to the inflamed surface, and then gently rubbing on oxide of zinc ointment. Ointments containing lead are also of great value. The most generally applicable formula is the following, recommended by Hebra: fʒxv of olive oil and ʒij to ʒvj of litharge are boiled together to a good consistence, and then fʒij of oil of lavender are to be added.

Subnitrate of bismuth in the proportion of 20 to 40 grains to an ounce of simple ointment or cosmoline, makes a very useful application. It must be remembered that in young children it will be better to reduce the strength of the above formula by about one-half, so as to adapt these to the extreme sensibility of the skin.

It is impossible to draw the line accurately between the acute cases for which such soothing or gently astringent applications are required, and cases of a subacute or chronic character which will tolerate a more powerful local treatment. In many cases, the applications already recommended will suffice from the beginning to the end of a case, but at times it becomes necessary to resort to those of a more stimulating character. The amount of local heat and irritation; the character of the discharge; the color of the infiltrated skin; and the duration of the case; will assist greatly in deciding this occasional difficult question.

When the patches of eruption are small, ointment containing mercury will often cause rapid recovery. We have used with much satisfaction a preparation of one part of ung. hydrarg. nitratis with three or four parts of simple cerate or cosmoline.

The following may also be recommended :

R. Hydrargyri Protiodidi,	gr. xij.
Camphoræ	gr. v.
Axungia,	℥j.
M. et ft. ung.	S. Apply twice daily.

Or,

R. Hydrargyri Chloridi Mitis,	℥j.
Camphoræ,	gr. v.
Glycerinæ,	f℥j.
Ung. Aquæ Rosæ,	℥j.
M. et ft. ung.	

Or, weak solutions of bichloride of mercury, gr. ss. to f℥j, as Van Swieten's liquor :

R. Hydrarg. Chloridi Corros.,	gr. xviii.
Alcoholis,	f℥iij.
Aquæ Destillat.,	f℥xxix.
Ft. sol.	

These lotions may be applied on pledgets of lint wetted with them, or, if such prolonged applications prove irritating, they may be used by merely washing the part with them for a quarter of an hour each time.

Applications of tar are applicable to chronic eczema, whether of limited or considerable extent, so that they must rank as the most useful of all local applications in these affections. Mere irritation is no contra-indication, for in fact they are the best anti-pruriginous remedies; but as long as the morbid process presents any of the signs of the acute stage, no form of tar is to be recommended.

In proportion as the case presents the features of the chronic form will these applications prove of value. They may be used in the form of ointments, as the officinal tar ointment, suitably diluted; or instead of *common tar*, the *oil of cade* (oleum cadinum) may be used as follows :

R. Olei Cadini,	f℥ss to f℥j.
Cerati Simplicis,	℥j.
Ol. Amygdal. Amar.,	gtt. v.
M. et ft. ung.	

For application to the scalp, as is so often required in children, Duhring advises a fluid preparation of tar instead of an ointment :

R. Picis Liquidæ,	f℥j.
Glycerinæ,	f℥j.
Alcoholis,	f℥vj.
Ol. Amygdal. Amar.,	gtt. xv.
Ft. sol.	S. To be diluted suitably and rubbed firmly into the skin.

Particular attention must be paid to the way in which tarry applications, liquid or solid, as well as other stimulating applications used in chronic skin diseases, are employed. They must not be merely smeared over the surface, but a small quantity being taken on a piece of sponge or flannel it should be firmly and patiently worked into the skin.

Especially when much thickening of the epidermic layers and infiltration of the skin exist, various alkaline substances may be combined advantageously with the tarry preparation. Thus one part of tar may be added, three parts of Hebra's spiritus saponatus kalinus, the formula for which is given below, and may be applied as there directed. In such chronic cases, when the patches of disease are circumscribed, we may use with great care a lotion of tar, alcohol, and sapo viridis in equal parts; or the following prescription of Bulkley:

R. Picis Liquidæ,	f3ij.
Potassæ Causticæ,	3j.
Aq. Destillatæ,	f3v.—M.

S. "Liq. Picis Alkalinus." To be used very largely diluted with water, as from one fluid drachm to the pint up to one part in ten of water, according to the extent of the disease and the susceptibility of the surface.

Carbolic acid, which produces an action analogous to that of the tarry preparation, is a very valuable application, and may be used either in the form of a lotion or in the proportion of 5 to 10 drops to an ounce of cosmoline, simple cerate, or benzoated oxide of zinc ointment. So also is diluted fluid extract of *grindelia robusta*. Alkaline preparations are very useful without the addition of any tarry substances, especially for the removal of circumscribed infiltrations and epidermic thickenings in chronic eczema. The remedy from which we have obtained the most beneficial effects in such cases, in the spiritus saponatus kalinus of Hebra, to which we have referred above, and which is prepared by dissolving soft (potash) soaps in alcohol, filtering the solution, and scenting it with oil of lavender or any aromatic spirit:

R. Saponis Mollis,	3ij.
Alcoholis,	f3j.
Sp. Lavandulæ,	gtt. xx.
M. et colque.	

In the use of this application it is essential, as directed by Hebra, that the soap should be finely rubbed into the eruptive patch by means of a piece of flannel or brush, till the accumulated masses of epidermis are removed, and a little blood is seen to ooze from the red base which has thus been exposed. The treatment is not as painful as would be supposed; and it should be followed immediately by the thorough application of an ointment of oxide of zinc or of litharge (see formulæ).

Other ointments and lotions containing alkaline substances are also recommended, such as carbonate of potash, gr. xx to xxx to 3j of lard; or caustic potash, gr. ij to gr. x to f3j of water. This latter application is especially useful in cases where the eruption is confined to limited patches, and is attended with much chronic infiltration of the skin. If the stronger forms of the solution are used, they should, after being applied quickly by means of a brush, be washed off by a large brush wetted with pure water.

In cases of eczema tarsi, attended with infiltration of the eyelids, McCall Anderson recommends that the eyelashes should be extracted, the eyelids

everted, and a solution of caustic potash, gr. v or x to fʒj, applied and quickly washed off by a large brush. Care should be observed in case the edges of the eyelids are adherent in the morning, not to separate them rudely, but to moisten them with tepid water or milk and water, so as to soften the crusts. Afterwards an application of citrine ointment, diluted with about two parts of lard, should be made along the edges of the lids night and morning.

ARTICLE II.

HERPES.

DEFINITION; VARIETIES; FREQUENCY.—Herpes is a non-contagious cutaneous disease, characterized by an eruption of vesicles assembled in groups on inflamed surfaces, of irregular size and shape, which are separated from each other by perfectly healthy portions of skin. The disease is usually acute in its course, seldom lasting more than two or three weeks, but it is not, as a general rule, accompanied by any severe constitutional symptoms. The separate vesicles composing the eruption last about ten days, and then disappear by the absorption of their contents, by the drying up of the contained fluid without rupture of the vesicles, or by the rupture of the vesicles, the escape of the fluid, and the formation of thin, brownish, or yellowish scabs.

There are several different varieties of herpes, which have been well divided by Mr. Wilson into two groups, the *phlyctenoid* and *circinate*. The phlyctenoid group is characterized by the irregularity of form exhibited by the eruption, and includes the variety called herpes phlyctenodes, and the local forms, called, according to their seat, labialis, nasalis, palpebralis, auricularis, præputialis, and pudendalis; whilst the circinate group is characterized by the arrangement of the vesicles in circles, and includes the herpes zoster and iris. Of these different varieties we shall describe, as of importance in children, only the *phlyctenodes*, *zoster*, and *iris*. Herpes circinatus, formerly included in this group, will be found described in the article on tinea.

HERPES is quite a frequent disease in children, though one rarely of any considerable importance.

CAUSES.—The causes of herpes are often obscure and uncertain, and in many cases entirely inappreciable. The disease is most common in persons who possess a delicate and irritable skin. The most frequent and most clearly ascertained cause is some disturbance of the digestive functions; and when there exist, in connection with this condition, irritations or inflammations of the respiratory mucous membrane, it is especially apt to be developed. Herpes phlyctenodes often follows exposure to a hot sun, while herpes labialis is frequently caused by exposure to a cold wind, especially when this occurs immediately after leaving a heated room. The latter variety also frequently accompanies coryza, angina, and stomatitis;

it also appears quite frequently in the course or at the termination of typhoid or intermittent fevers, pneumonia, etc.

The usual exciting causes of the disease are irregularities in diet, exposure of the body while in a heated state to cold and damp, local irritants, malarial disease, and bilious disorders of all kinds.

The cause of herpes zoster is peculiar. The eruption appears, in nearly if not quite all cases, to be dependent upon a morbidly irritable state of some nerve-trunk, which may be the result of simple or rheumatic inflammation, of pressure, of mechanical irritation, etc.

Bärensprung, who was one of the first to recognize the dependence of herpes zoster on morbid conditions of the nerve-trunks, has not only actually detected inflammatory lesions of the intercostal nerves in cases of zoster pectoralis, but has also shown that there is a primary lesion of the corresponding spinal ganglia. These views have received such repeated confirmation, that they may be accepted as representing the true pathology of this interesting affection.

HERPES PHLYCTENODES.—This variety of herpes, unlike the other forms of the disease, may appear upon any part of the cutaneous surface, and does not assume a determinate shape. It may appear, indeed, upon several parts at the same time. It is usually, however, met with upon the upper parts of the body, and particularly the cheeks, neck, chest, and arms. It is rare to observe it on the lower extremities.

We believe it to be a rare affection amongst the children of families in easy circumstances. The only examples that we have seen have been the result of poisoning by the different kinds of *Toxicodendron*.

SYMPTOMS.—The eruption appears in the form of vesicles, usually of very small size, looking like mere points, or attaining sometimes the size of a pea, which are seated in groups or clusters on inflamed patches of the skin, varying in size from that of a dollar to that of the palm of the hand. Sensations of heat, smarting, and itching are often felt in the part where the eruption is about to show itself; and within a day, usually, after these symptoms have been observed, or without them, the disease makes its appearance, exhibiting one or more red and inflamed surfaces, of an irregular or rounded shape, dotted over with projecting, globular vesicles, which are hard, resisting, and, on the first day, transparent, but which become, in the course of a day or two, turbid or lactescent. The red color of the eruptive patch generally extends a short distance beyond the vesicles; the integument between the different patches retains, however, in all cases, its healthy color and character. A sense of smarting and itching accompanies, as well as precedes, the eruption. On the second day of the eruption, the number of vesicles gradually increases, and they become full and distended. About the third or fourth day, the vesicles have become very turbid, and they begin to shrink. About the seventh or eighth day, they are usually transformed, by the drying up of their contents, into thin, brownish crusts, which fall off by desquamation about the tenth or twelfth day. There also remains, for a few days after the disappearance of the eruption, some redness of the surface, which subsides little by little.

This variety of herpes is never accompanied by constitutional symptoms

of any severity. A very slight febrile reaction, some languor, loss of appetite, and thirst may precede the appearance of the eruption for a few days, and continue for a short time after the nature of the case has declared itself.

HERPES LABIALIS.—This is the most frequent of all the varieties of the disease. The eruption, as the name implies, is seated on the lips. Usually it occurs upon the line of junction of the mucous membrane with the integument; but it may affect either the former or the latter alone. Though generally confined strictly to the lips, the eruption, in some instances, extends to the cheeks, chin, or alæ of the nose.

The disease begins generally with redness, heat, smarting, and painful tension of the portion of the lip upon which the eruption is about to appear. After a few hours, or a day, vesicles begin to show themselves upon the inflamed spot, and there is then observed a red, swollen, and shining point, upon which is seated a group of vesicles. The tumefaction and redness commonly extend some distance beyond the vesicles. The latter develop themselves rapidly until five or six small, rounded vesicles, filled with a transparent fluid, are seen. The vesicles remain solitary, or several may unite together to form one of considerable size. After the complete development of the eruption, the burning pain which existed at first commonly subsides. The contents of the vesicles soon become turbid and lactescent, and are converted, by the third or fourth day, from a serous into a sero-purulent fluid, at which time, also, the accompanying redness and swelling have, in great measure, disappeared. Soon after this, brownish crusts, are formed by the drying up of the fluid of the vesicles, and these drop off usually about the seventh or eighth day. A slight redness remains for a short time at the point of eruption, and then disappears entirely.

HERPES ZOSTER.—This variety of herpes, known also by the names of *zona* and *shingles*, is of a rather rare occurrence in children. It derives its name, which signifies a girdle or belt, from the fact that when it attacks the trunk the eruption often surrounds one-half of the body in the form of a belt, of varying width. It attacks various parts of the body, especially the trunk and the face, though it may also appear on the extremities. It always follows the distribution of some nerve-trunk, and, as a rule, is confined to one side of the body.

When it occurs at the base of the thorax (*zoster pectoralis*), the course of the eruption is determined by that of the adjacent intercostal nerve, so that it usually extends from the median line in front to the same point behind. In *zoster abdominalis* the course is much the same. About the head it may appear in the course of the supraorbital nerve, extending over the brow into the scalp (*zoster capitis*); or else in the course of the lower branches of the trifacial, involving the cheek, and extending downwards towards the neck (*zoster faciei*). It may also occur about the shoulder or back of the neck and extend down the arm (*zoster brachialis*), or on the thigh (*zoster femoralis*). When the eruption appears in the form of a belt, the zone is not composed of a continuous line of vesicles, but is made up of distinct patches of eruption, all following the same general direction,

but divided from each other by portions of healthy integument. The eruptive patches may be very closely approximated, or they may be separated by considerable spaces of skin untouched by the disease.

The disease is acute in its character, lasting as a general rule, from one to three or four weeks.

An attack of zoster is usually preceded for several days by smarting and burning, and by severe neuralgic pains in the part that is to be affected. The eruption then appears in the form of irregular patches of a vivid red color, more or less widely separated from each other, and grouped so as to form a racemose or belt-like form according to the distribution of the affected nerve. Soon after the appearance of the inflamed patches, numerous small white projections can be seen, by careful examination, upon the red surfaces: these increase rapidly in size, and are soon converted into distinct transparent vesicles. The vesicles augment in size, and arrive, in the course of three or four days, at their fullest development, when they are about as large as small or large peas, or, in some few instances, much larger, and containing a clear yellowish fluid. At this stage of the eruption the red surface upon which each group of vesicles is seated extends a slight distance beyond the patch, thus forming a kind of areola.

After remaining in this state for four or five days, each group of vesicles begins to subside. The redness of the inflamed patch diminishes; the vesicles shrink, and become shrivelled; their contents, which were transparent at first, become opaque and puriform, and finally they dry up and form small, dark-brown scabs, which fall off about the tenth or twelfth day, leaving behind reddish spots, which disappear little by little.

The constitutional symptoms of herpes zoster consist usually of slight feverishness, languor, and the signs of gastro-intestinal irritation. The local symptoms are pungent and burning pain at the beginning of the eruption, and more or less severe tension, and sometimes acute pain, in the part upon which the disease is seated, which latter lasts, in some instances, throughout the course of the disorder, or even for some considerable time after it has disappeared. This neuralgic pain, which is dependent on the implication of a nerve-trunk, varies much in intensity, being at times slight, and at others very intense.

HERPES CIRCINATUS.—This variety of herpes has been called also ring herpes, herpetic ringworm, and vesicular ringworm; it will be found described under the name *tinea circinata* in the article on parasitic diseases of the skin.

HERPES IRIS.—This is a very rare variety of herpes, and one that we have met with in children in but a few instances, although according to Duhring (*op. cit.*, p. 222) it is comparatively frequent in children and young people. It begins with small red spots, which are soon surrounded by four or five rings of different shades of redness. About the second day of the eruption, the central red spots present in their centres one or more vesicles, and on the third and fourth days, vesicles of very minute size generally appear on the outer concentric rings. After two or three days, the fluid contained in the central group of vesicles, which was transparent

at first, becomes turbid, and about the fifth or sixth of the eruption, it is absorbed, and the disease terminates by a slight desquamation. All the colors of the rainbow, subdued in tone, may usually be observed at one time or another in the course of the disease, the red, yellow, and violet shades predominating (Duhring). The vesicles formed on the outer ring undergo the same changes as those described as occurring on the central ones. In some instances, the vesicles open, and their contents escaping, form small, thin, and brownish scales, which fall off in ten or twelve days.

Herpes iris may attack any part of the body, but is most frequently developed upon the face, hands, fingers and neck.

According to some dermatologists, as McCall Anderson, herpes iris is a parasitic disease and merely a form of *tinea versicolor*. This view, however, does not appear to us to be correct, as this affection seems, on the contrary, to have the closest analogies with *erythema vesiculosum*.

DIAGNOSIS.—The diagnosis of herpes is seldom attended with any difficulty. The small size of the vesicles, their globular shape, their number, their aggregation upon distinct patches of inflamed integument, and the slight degree of constitutional disturbance attendant upon the disease, all render the eruption unlike any other cutaneous affection, and therefore easy of recognition.

Herpes phlyctenodes might possibly be confounded with pemphigus. The recollection that the eruption in pemphigus consists of distinct bullæ, much larger, of course, than the vesicles of herpes, while that of herpes phlyctenodes consists of numerous vesicles, much smaller than the bullæ of pemphigus, and closely dotted over isolated red patches, will always serve to distinguish the two affections. It might be mistaken also for eczema, when the vesicles of the latter are disposed, as sometimes, though rarely, happens, in groups. The distinction may be made, however, by attention to the facts that the eczematous vesicles are redder, less elevated, scarcely transparent, and that, though arranged in groups, they are confluent, whilst in herpes they are always distinct.

Herpes labialis is not likely to be mistaken for any other eruption. Herpes zoster may always be distinguished by the peculiar forms assumed by the eruption, and by its arrangement in the course of some nerve tract, and by the neuralgic pains which attend it.

There is but one disease with which herpes iris is likely to be confounded,—*roseola annulata*. The entire absence of vesicles in the latter affection will always, however, enable us to make the distinction.

PROGNOSIS.—The *prognosis* of herpes is always favorable. It is never in itself a dangerous disease, though zoster often causes much suffering, and is moreover usually the expression of a considerable disturbance of the general health.

TREATMENT.—The different varieties of herpes seldom require more than the mildest treatment. In all, attention should be paid to the general health. The diet must be regulated according to the state of the digestive function. When constipation is present, especially if there be some febrile reaction at the same time, gentle laxatives ought to be ad-

ministered, such as sulphur, magnesia, syrup of rhubarb and magnesia, or castor oil. If the skin be sallow, the tongue heavily coated, the breath foul, and the stools scanty and light-colored, or very offensive, small doses of blue pill in combination with rhubarb, or followed by rhubarb and magnesia, would be the most appropriate remedy. Excessive or frequently repeated doses of any purgative ought to be avoided, as the debility and gastro-intestinal irritation that so often follow such practice, are more injurious than the original disease.

The local treatment of herpes is important, and is, indeed, in many cases, all that is necessary.

Herpes phlyctenodes requires nothing more than mucilaginous lotions, an occasional warm bath, or the frequent moistening of the eruption with a liniment made of equal parts of lime water and sweet oil. Herpes labialis, if it demand local treatment at all, may be relieved by the use of any mild lip salve; a very good ointment is one composed of equal parts of Goulard's cerate and simple cerate, with a few drops of glycerin. Mr. Wilson recommends the following ointment:

R. Unguent. Flor. Sambuci, ℥j.
 Liq. Plumbi, fʒj.—M.

During the early stage of herpes zoster, the local treatment should be such as will tend to allay inflammation and relieve pain. These results may be obtained by applying compresses moistened with some kind of mucilage, such as barley-water, or decoction of flaxseed or slippery-elm bark, or with simple cold water, or with weak lead-water and laudanum. The application of a dusting-powder of starch, camphor, and morphia often affords relief. When the eruption is followed by excoriations or ulcerations, and the pain is severe, the latter may be allayed by the use of an ointment consisting of equal parts of Goulard's cerate and lard, either alone, or containing two or three grains of opium, or half a drachm of the watery extract of opium. Duhring speaks highly of the application of flexible collodion, with morphia in the strength of ten grains to the ounce. Underwood recommends, when the discharge has subsided, and the scabs have formed and become adherent, that they should be anointed twice daily with the ung. hydrarg. ammoniat. It is, however, necessary to also employ some internal treatment to relieve the neuralgic pains, which are so prominent a symptom. For this purpose, opium, or preferably the hypodermic injection of morphia, must frequently be employed. We have also found that the combination of iron, quinia, arsenious acid, and belladonna has afforded marked relief in some cases. Phosphide of zinc has been highly recommended by Ashburton Thompson and Bulkley in doses of one-third of a grain for an adult. The employment of a galvanic current applied along the tract of the affected nerve, has proved very beneficial.

Herpes iris seldom requires any treatment. If any be determined on, it should consist of alkaline lotions, or of water rendered slightly astringent by the addition of alum, or sulphate of zinc.

ARTICLE III.

MILIARIA.

MILIARIA is an acute inflammatory disorder of the sweat glands, characterized by numerous minute papules or vesicles, attended by prickling, tingling, and burning sensations.

In many cases both papules and vesicles are present, although usually one or the other will predominate. It is on this account that one of its forms, *miliaria papulosa*, has long been regarded as a form of lichen, under the name of *lichen tropicus*, or *prickly heat*. This affection is a very common one at all ages of childhood, from early infancy upwards, in this city, and in most of our Middle and Southern States. It is, as already stated in the definition, which is taken from Duhring, essentially an inflammatory affection of the sweat glands, and differs, therefore, entirely from true lichen. In some cases the congestion and exudation about the ducts lead to papules for the most part; while, in others, there is a greater tendency to vesiculation. On the whole, the disposition is to the latter lesion. There can be no hesitation, however, in view of the essential pathology of the disease, in assigning it to a place among the vesicular affections.

The chief *cause* of prickly heat is the action on the skin of a high temperature, aided, no doubt, by the disturbances of the digestive function so apt to coincide with extreme heat. It is especially common upon the sudden development of unusually hot weather. Very warm weather, and particularly the contact of thick rough flannels with the skin, are apt to develop the eruption.

SYMPTOMS.—Prickly heat, or lichen papulosa, or tropicus, appears suddenly in the form of numerous minute papules, few of them being larger than a pin's head, scattered more or less thickly over the affected surface. The pimples are of a red color, which are more or less bright in tint, according to the extent and intensity of the eruption. Usually vesicles, or vesico-papules, are seen here and there upon the affected surface.

The skin between the papules retains its natural appearance when the eruption is but slight or moderate; but when this is copious and severe, it assumes a faint reddish appearance, owing no doubt to the activity of the circulation in the part.

The eruption is most abundant on the parts covered by the dress, or rubbed by the edges of the dress, particularly about the neck, upper part of the chest, and on the arms and legs. We have sometimes seen it covering the greater part of the body. It is always attended with more or less itching, burning, and prickling, which, in older children, causes much fretfulness and scratching, and, in those who are younger, restlessness, worrying, and more or less disturbance of the sleep. The disorder usually remains stationary for several days, and then disappears gradually without desquamation or other change in the skin; or, it subsides and increases, or disappears and returns, with the rising and falling of the temperature, or without any very evident cause, until at last it ceases, not to appear again.

When the eruption lasts many days, it is almost always accompanied by a slight scaly desquamation of the tops of the pimples.

The vesicular form of miliaria is rather rare in children. It is characterized by the sudden development of numerous minute discrete vesicles, occurring in large patches on a congested or slightly inflamed skin; running an acute course, drying up in a few days, and ending in slight desquamation.

The *diagnosis* of prickly heat is never difficult. Its sudden occurrence during hot weather, the character of the papules, their minuteness and abundance, and the entire absence of constitutional disturbance, will always render it easy of recognition.

TREATMENT.—Prickly heat needs no treatment, except when very abundant, and when it annoys the child by the heat and itching it occasions. Under these circumstances, the skin should be dusted with rye-meal, or anointed two or three times a day with some mild ointment, as, for instance, one consisting of glycerin and cold cream or lard, or the benzoated zinc ointment; or, the child may be bathed once or twice a day in warm water, containing bran, slippery-elm, or some other mucilaginous substance. Alkaline baths or lotions may also be used with benefit. If any predisposing cause exists, as a warm apartment, too heavy clothing, or injudicious diet, it must be corrected. Small doses of quinia with one of the mineral acids will often have a tendency to lessen sweating and thus afford some relief.

CHAPTER III.

BULLOUS INFLAMMATION OF THE SKIN.

THE distinguishing feature of this form of inflammation is the formation of blebs or bullæ of considerable size. We include pemphigus and rupia under this heading, though the latter might with equal propriety have been treated of in the chapter on syphilitic diseases of the skin.

ARTICLE I.

PEMPHIGUS.

DEFINITION; SYNONYMS; VARIETIES; FREQUENCY.—Pemphigus is an acute or chronic inflammatory disease, characterized by the presence on one or several parts of the body of more or less numerous bullæ of considerable size, nearly always isolated, resting upon circular or oval erythematous patches, about as large or somewhat larger than the bases of the bullæ themselves. The bullæ form in the course of a few hours, and contain at first a limpid serum, which soon becomes reddish or turbid; they terminate by desiccation and the formation of thin crusts, or by rup-

ture and the escape of their contents, when there remains behind a superficial ulceration.

The two most clearly marked varieties are pemphigus vulgaris and foliaceus. The former occurs both as an acute and chronic affection, but it is only the acute form which occurs with special frequency in children, and we shall therefore give a detailed account of this alone.

Pemphigus is not unfrequently met with in young children who become the inmates of hospitals, almshouses, and foundling hospitals, and amongst the poor and destitute classes of large cities. Under such unfavorable conditions, it occasionally assumes what must be termed an epidemic form. Still it cannot be said to be a frequent disease.

CAUSES.—The causes of pemphigus are often obscure or entirely inappreciable. It is usually supposed, however, to depend, in children, upon the influence of the act of dentition, on disturbances of the gastro-intestinal tract brought about by improper food or overfeeding, and on general disorder of the nervous system. The so-called syphilitic pemphigus, which is one of the most frequent eruptions in congenital syphilis, and is not rarely present at birth in such cases, is not a true pemphigus, but has been described as a bullous syphiloderm.

SYMPTOMS.—Acute pemphigus may be confined to a very small portion of the cutaneous surface, or it may affect several regions of the body at once. It is usually attended with symptoms of constitutional disturbance, which, especially in very young infants, may be slight, consisting merely of general uneasiness, languor, and some acceleration of the pulse; or they may be severe, exhibiting in such cases a dry and burning skin, frequent pulse, thirst, and loss of appetite.

After the above constitutional symptoms have lasted one, two, or three days, the eruption makes its appearance in the form of small circular red spots, which increase in size, and soon exhibit a bleb or bulla rising in the middle or over the whole of the red spot. The vesicle commonly appears a few hours after the red patch, and consists of an elevation of the cuticle by an effusion of serum beneath it. The bulla rapidly distends by the increase of the serous effusion, until it attains the size of a pea, a hazelnut, or a large walnut. It is of a circular or oval form, and may be confined to the centre of the erythematous surface on which it rests, being surrounded in such cases by a more or less wide red line of inflammation, or it may occupy the whole or nearly the whole of the red patch, under which circumstances it entirely conceals the latter, or is surrounded by a very narrow red ring. The color of the areolæ around the bullæ is very bright during the first day of the eruption, while the integument between remains perfectly healthy.

The fluid contained in the bullæ soon becomes turbid; the bullæ become wrinkled, and usually burst after one or two days, and are replaced by thin yellowish or brownish scabs. The crusts begin to form before the redness of the integument has disappeared. In some instances the bullæ do not break, but their contained fluid becomes yellowish in color, and then turbid; it diminishes by absorption, and, at the end of about a week, dries into a thin dark-colored scab. The crusts usually fall off in the course of

two or three weeks, leaving the skin beneath of a reddish color, but in other respects healthy. The whole duration of the disease is commonly from one to three weeks, the time in each case varying with the mode of the eruption; when all the bullæ appear simultaneously, seldom lasting more than one or two weeks; while in cases in which they appear at successive periods, lasting three or even four weeks.

When pemphigus occurs in children who are cachectic, ill-fed, and surrounded by poor hygienic conditions, and especially when it appears in an epidemic form in badly-managed public institutions, it may assume a much more grave form, known as pemphigus gangrenosus or cachecticus. The eruption affects the neck, chest, abdomen, scrotum, hands, and feet. It begins as purplish or livid spots, raised slightly above the level of the skin, upon which bullæ soon form, of irregular shape, flattened on the top, and surrounded by purplish areolæ. The fluid they contain soon becomes fetid, turbid, and dark-colored, or almost black. If the bullæ burst, deep unhealthy ulcers are exposed, with a dark, shreddy, or gangrenous surface, secreting a fetid sanies. The constitutional symptoms are severe, indicating blood-poisoning and rapid collapse of the vital powers; death often occurs as early as the tenth or twelfth day. In cases where the dyscrasia is not so intense, the case is much prolonged; successive crops of such bullæ appearing until the child dies, worn out by suffering and exhausting discharge, or else enters gradually into a slow and difficult convalescence.

Chronic pemphigus is the usual form of the disease in adults, in whom it may run a course extending over many years. In pemphigus foliaceus, also, the affection is essentially a chronic one, and occurs only in adults. It differs from pemphigus vulgaris chronicus chiefly in the imperfect development of the bullæ, which are flaccid and only partly filled with fluid. This dries rapidly into thin whitish flakes, and as the bullæ are very numerous and large the whole surface may present a red scalded appearance, with loose, shreddy, and flaky epidermis.

DIAGNOSIS.—The diagnosis of pemphigus acutus is seldom difficult. The large isolated bullæ, seated on inflamed patches of the integument, filled with transparent serum, and followed by thin lamellated scabs, are unlike any other kind of eruption. The mode of distinguishing it from the bullous form of syphiloderm has been considered in the article on the latter subject.

PROGNOSIS.—Acute pemphigus is rarely dangerous when it exists without complications. When, however, it is very extensive, and accompanied with severe constitutional symptoms, and particularly when it exists in connection with other diseases, or occurs in a child whose health has been broken down by unwholesome hygienic influences, it may assume a dangerous character, and the prognosis should, therefore, always be guarded under such circumstances. The gangrenous form of pemphigus, especially when occurring under bad hygienic circumstances, is a very fatal disease.

TREATMENT.—Simple acute pemphigus may require no other treatment than attention to diet, and regulation of the digestive function. When constipation is present, this should be overcome by means of simple enemata, or by the use of some mild laxative, as manna, spiced syrup of

rhubarb, or very small doses of castor oil. If the discharges be too frequent, they should be restrained by the use of opium, in doses proportioned to the age of the child. In young infants, it will often be found that the gastro-intestinal secretions are of an acid and irritating character. This condition may be treated with small doses of paregoric or laudanum, combined with lime or magnesia water, or with soda. The diet must be managed according to the state of health of the child. For an infant, a good breast of milk is, of course, the best treatment in the world. For older children, the diet ought to be light and unirritating, but, at the same time, nourishing and strengthening.

When, however, the child shows signs of debility during the progress of the disease, and also when the eruption tends to assume a chronic course, the treatment ought to be tonic and invigorating. It should consist in the use of a nutritious diet, and in the exhibition of tonics, as Huxham's tincture of bark, quinia, arsenic, cod-liver oil, or in the use of wine-whey, or small quantities of brandy.

In the cachectic or gangrenous variety, the treatment must be supporting and stimulating in a high degree. Full doses of quinia with mineral acids, as sulphurous or muriatic, and chlorate of potassa, together with milk punch of suitable strength and carefully regulated diet, are to be recommended.

The local treatment should consist, in the early stage, of an occasional warm bath. When the bullæ have fully formed, they ought to be punctured, and the fluid gently pressed out, care being taken not to remove the cuticle, as this forms the best possible dressing for the inflamed integument. If much irritation be present, as is sometimes the case, relief may be gained from the use of water dressings by means of cloths, or from lotions of diluted fluid extract of *grindelia robusta*, or diluted lead water. When the bullæ have been followed by excoriations, these may be treated with lotions, as dilute *lotio nigra* or a weak solution of nitrate of silver or of sulphate of copper; or with ointments, as the benzoated oxide of zinc ointment, or one of equal parts of Goulard's cerate and cosmoline. An ointment made from the leaves of the *scrophularia nodosa* was found to be the most useful application in gangrenous pemphigus, by Dr. Whitley Stokes, who saw much of the disease among poor, ill-fed children in Ireland.

ARTICLE II.

RUPIA.

DEFINITION; VARIETIES.—Rupia is an eruptive disease, characterized in its early stage by distinct, somewhat flattened bullæ, of more or less considerable size, containing at first a serous, and then a purulent or

blackish fluid; at a later period the disease exhibits very thick scabs, and still later, ulcerations.

There are two varieties of this eruption: *rupia simplex*, and *rupia prominens*. *Rupia escharotica*, formerly included as a variety of this affection, will be found described in the article on pemphigus, under the title of pemphigus gangrenosus.

CAUSES.—*Rupia* is most apt to occur in weakly, badly nourished, and scrofulous children, and seems to depend, therefore, upon that state of debility and exhaustion of the general health which results from exposure to unfavorable hygienic conditions, which follows exhausting diseases, or which exists as a consequence of some hereditary taint. It also occurs in connection with congenital syphilis; and is then described (Duhring) as a large flat pustular or ecthymatous syphiloderm.

SYMPTOMS.—*Rupia simplex* begins almost always on the inferior extremities, or more rarely on the trunk or arms, without previous inflammation, in the form of small, flattened bullæ of about three or four lines in diameter. The bullæ contain at first a serous and transparent fluid, which soon becomes thicker, and is converted into pus. At an early period they shrink and become wrinkled, their contained fluid hardens and is converted into rough, brownish scabs, which are always thicker at the centre than on the edges, and which leave beneath, after their fall, superficial ulcerations. These ulcerations either soon cicatrize and disappear, or are covered by fresh scales. After the fall of the final scabs, there yet remain, for some time longer, dark-brown or livid spots, which gradually fade and disappear.

Rupia prominens exhibits the same general characters as the preceding variety, but with more marked and peculiar features. The eruption commences with a circumscribed inflammation of the skin, on which inflamed spot soon appears a bulla filled with yellowish serum, or sometimes with a blackish fluid, which rapidly hardens into a brownish or blackish wrinkled crust. The crust is surrounded by an erythematous areola, formed by the extension of the cutaneous inflammation beyond the circumference of the scab. Upon this areola a fresh elevation of the cuticle, by purulent deposit, often takes place, which, by its desiccation, adds to the size of the crust. This successive increase at the margin of the scab enlarges it in breadth, and at the same time raises the height of its centre, so as to give it a peculiar and characteristic appearance, and causes it to resemble very closely the shell of a limpet or oyster. The scabs thus formed usually adhere to the surface beneath with much tenacity, and remain attached for a variable, and, as a general rule, considerable length of time. When at length they fall off, or are removed, there are left beneath ulcers of variable depth and extent, which are either covered by fresh crusts, or, as more frequently happens, remain open, presenting a foul surface of a livid red color, with thickened edges. The ulcers are difficult to heal, and, after cicatrization, leave livid or purplish stains, which often remain for months. The number of bullæ is usually small, there being generally one at its height, and one or two about to appear, or on the decline.

DIAGNOSIS.—Rupia is likely to be confounded only with pemphigus and ecthyma. Pemphigus is to be distinguished from rupia by the larger size and greater distension and prominence of its bullæ; by the fact that the contained fluid of the latter is serous and transparent in pemphigus, instead of being turbid and sanguinolent, as in rupia; by the different character of the crusts, which, in pemphigus, are thin and lamellated, while in rupia they are thick and rugous; and, lastly, by the deep and unhealthy-looking ulcerations that follow rupia.

Ecthyma is unlike rupia in being a pustular disease from the first. Moreover, the pustules of ecthyma are surrounded by a highly inflamed areola, which is not the case in rupia, while the crusts in the former disease differ from those in the latter, in being smaller, harder, more irregular, and more adherent.

PROGNOSIS.—Rupia simplex and prominens, though tedious and slow of cure, seldom prove fatal; if any danger accompany the disease, it arises rather from the enfeebled and disordered state of the general health under the influence of which it is produced, than from any injury caused by the eruption.

TREATMENT.—The most important point in the treatment is to attend to the hygienic state of the patient. When the child is living in an unhealthy house, or a close and confined room, it should be removed, if possible, to a more salubrious position, or to a larger and well-ventilated room. The diet ought to be such as to invigorate the strength, and promote the nutrition of the body. For an infant who is fed upon artificial food, or who is suckling a nurse of doubtful health, the best remedy in the world is a fresh and full breast of milk. If a nurse cannot be procured, the diet must be most carefully regulated in accordance with the principles already detailed in full in the article on thrush, at page 361. While the diet is thus attended to, it is necessary to watch the state of the digestive organs, and if there be either constipation or diarrhœa, these must be overcome by suitable remedies. Tonics and stimulants are always advisable in this disease, and may consist either of brandy or wine, given alone, or in connection with Huxham's tincture of bark, extract of cinchona, small doses of quinia, iron, cod-liver oil, or any other remedy of this kind that may be preferred. In cases of syphilitic origin specific treatment should be resorted to as detailed on page 713.

Rupia simplex and prominens are to be locally treated in the early stage by opening the bullæ so soon as they form, and covering them with dry lint and a light bandage, or with the water-dressing. The ulcerations that follow the bullæ may be treated with Goulard's ointment, applied on pieces of fenestrated lint, and by washing occasionally with lime-water, or with weak solutions of alum, copper, zinc, or nitrate of silver. At a later period of the disease, when the ulcerations are covered with the characteristic thick crusts, these are first to be removed by means of poultices of bread and water, or flaxseed meal, and the surfaces beneath them treated with the applications recommended above. When the ulcerations are very obstinate and difficult to heal, they should be modified by occa-

sional touchings with nitrate of silver, either pure or in strong solution, or with dilute nitric or muriatic acid.

Billard recommends that the ulcerations should be dusted with powdered alum or cream of tartar, and Rayer also speaks very highly of the last-named application.

CHAPTER IV.

PUSTULAR OR SUPPURATIVE INFLAMMATION OF THE SKIN.

THIS is characterized by the development of pustules, superficial and painless, or deeply-seated and painful. A considerable number of affections are included under this heading, only two of which require detailed consideration from us. Impetigo has already been treated of among the vesicular affections, under the name of *eczema pustulosum*; but there is one form, *impetigo contagiosa*, which properly belongs here. In addition to this, *ecthyma*, *sycosis non-parasitica*, and, by many authorities, *acne* and *acne rosacea* are also included. But only the first of these, *ecthyma*, is a disease of childhood, and this alone will therefore be described.

ARTICLE I.

ECTHYMA.

DEFINITION; SYNONYMS; VARIETIES.—*Ecthyma* is an eruption characterized by prominent, rounded, and usually discrete pustules of considerable size, with hard and inflamed bases. The pustules, which are sometimes termed *phlyzacious*, are followed by thick, brownish crusts, which leave on their fall a reddish mark, or more rarely a superficial ulcer or a true cicatrix.

Ecthyma occurs both in an acute and chronic form. The variety known as *ecthyma infantile* is usually of an essentially chronic character.

CAUSES.—*Ecthyma* is especially an affection of impoverished systems. It rarely occurs except in those who have become predisposed by improper or insufficient nourishment, lack of fresh air and sunlight, and similar depressing influences. It is met with in feeble, cachectic children, and in those whose health has been broken down by exhausting diseases, and particularly by disorders of the gastro-intestinal apparatus. It may be provoked in such subjects by the application of an irritant to the skin, by scratching, or by the presence of other eruptions, particularly that of small-pox, measles, scarlet fever, or scabies.

SYMPTOMS.—Acute *ecthyma* is rare in children. It may be preceded by symptoms of mild febrile disturbance. The eruption occurs most frequently on the extremities and neck, and more rarely on the trunk of the body. It appears in the form of small, red, and circumscribed spots, projecting above the surface of the skin, hard to the touch, and accompanied by smarting and often severe pain, and by soreness on pressure. The centre

of the spots is soon elevated into a pustule, filled with a purulent fluid. The size of the pustules varies, but is usually about that of half a pea. Each pustule is generally surrounded by a hard base of a bright red color, constituting an areola, while, in some instances, the whole of the red elevation is covered by the pustular formation. The pustule remains unchanged usually for three or four days, and more rarely for a week, and is then converted, by the drying up of the effused fluid, into a thinish brown scab, which drops off after a few days, and leaves a congested purple spot that remains for some time longer. In other instances, the pustule breaks and leaves a small ulceration which terminates with a slight cicatrix.

Even when ecthyma begins as an acute affection, the eruption is apt to appear in successive crops; and in the vast majority of cases, especially in children, it tends to pass into the chronic form. This is particularly the case when it is connected with some chronic disorder of the digestive or respiratory apparatus. The eruption in chronic ecthyma is similar to that of the acute form. The pustules may, however, be more variable in size, some being as large as a sixpence. They are circular in form, and surrounded by an areola of a red or purplish tint; the fluid which they contain is generally not very thick, and is of a dark and sanguinolent appearance; they terminate by the formation of a dark and adherent crust, by absorption of the contained fluid and a kind of desquamation, or by a bloody excoriation, or true ulceration, which are followed by a deep stain upon the skin or a true cicatrix.

DIAGNOSIS.—Ecthyma is more likely to be confounded with rupia, the bullous syphiloderm, than with any other disease. The pustular character of ecthyma from the very beginning, will, however, almost always enable us to distinguish it from the broad and distended bullæ of rupia, filled with sero-purulent fluid; and the difference between the two becomes still more marked, when we recollect the hard and inflamed bases on which the pustules of ecthyma rest, and the shapeless crusts and superficial excoriations of that disease, instead of the projecting, rugous, and imbricated scabs, and deep ulcerations of rupia. Ecthyma is not at all likely to be mistaken for the small and numerous pustules of eczema pustulosum, for those of impetigo contagiosa, or the umbilicated ones of small-pox.

PROGNOSIS.—Ecthyma is never a dangerous disease in itself. If any danger accompany it, it arises rather from the enfeebled and disordered state of the general health under the influence of which it is produced, than from any injury caused by the eruption. The prognosis must depend, therefore, upon the state of the general health existing during the attack of the disease.

TREATMENT.—In both varieties of ecthyma, attention to the general health of the patient constitutes the most important point in the treatment. In the acute form, mild laxatives, small doses of some alterative, as the hydrargyrum cum cretâ or sulphur, the use of a nutritious and wholesome, and especially of an unstimulating diet, and the local application of mucilaginous infusions, or of a mild and cooling ointment, as Goulard's cerate,

Turner's cerate, or the carrot, cucumber, or elder-flower ointments, with occasional warm bathing, are all that the case demands. In chronic ecthyma, the great deterioration of the general health usually requires close attention. As this deterioration depends usually upon the exposure of the child to unwholesome hygienic influences, and a consequent unhealthy state of the digestive and nutritive functions, it is of primary importance that these should be early attended to. The patient ought to be placed in a healthy and well-ventilated apartment; the clothing must be regulated according to the age of the child, and the season of the year; and, what is most important of all, the diet ought to be such as is digestible, suitable to the age, and, at the same time, nourishing and strengthening. The internal remedies must consist of tonics in all cases, and, when the digestive power and general strength are reduced much below the normal standard, of stimulants. The best stimulant is old and pure brandy, either given mixed with water, three or four times a day, or combined with the food. The best tonics are, in most cases, some preparation of iron, and the one we prefer is the iodide. . . . ; cod-liver oil, in emulsion with the lactophosphate of lime; quinia, and the mineral acids. While these remedies are being employed, or prior to their administration, the gastro-intestinal functions ought to be carefully regulated by the use of mild laxatives when the bowels are constipated, or by some kind of astringent when they are loose and disordered.

The external or local treatment must consist in the use of mild demulcent applications, or of soothing or cooling ointments, during the pustular stage of the eruption. When crusts have formed, they should be removed by poulticing, and a more or less stimulating ointment applied to the surface. When unhealthy excoriations or ulcerations follow the pustules, these may be brought into good condition by the employment of weak solutions of nitrate of silver or sulphate of zinc, or of a very weak lotion of nitric or muriatic acid.

ARTICLE II.

IMPETIGO CONTAGIOSA.

THE affection that was formerly called impetigo, has been already described as the pustular variety of eczema (*E. pustulosum*, see p. 936).

Impetigo contagiosa is a very different affection. It is, as defined by Duhring, an acute, inflammatory, contagious disease, characterized by the formation of superficial, flat, discrete, roundish vesico-pustules, the size of a split pea or finger-nail, which pass into granular, straw-colored crusts.

SYMPTOMS.—The appearance of the eruption is apt to be preceded by some mild febrile disturbance. The eruption occurs usually on the face, and about the head; but also on the arms and other parts of the surface. Small isolated vesicles form, with very slight surrounding redness, and if not broken by scratching, they enlarge, in the course of 5 or 6 days, into flal bullæ, as large as a sixpence, frequently with a distinct central de-

pression. Their contents grow turbid and purulent, and soon begin to dry up and form crusts. These are flat, rather thin, straw-colored, and granular-looking. Beneath them there is an erythematous base. The disease is auto-inoculable, and may be spread from place to place by means of its secretion. In this way the mucous membrane of the eye and nose may become implicated.

CAUSES AND PATHOLOGY.—The fact that the disease is both auto-inoculable, inoculable to others, and contagious, would be readily intelligible if it could be shown that it is of a parasitic nature. But thus far, although elements of a vegetable fungus are occasionally demonstrable (Kohn and Giffard) in the crusts it has not yet been shown that they exist in the liquid of the vesico-pustules (T. Fox), nor that they have any definite relation with the disease.

It is almost exclusively a disease of childhood. Mal-hygiene probably predisposes to it, but it may occur in children who have been well cared for. It has been noticed to follow vaccination in so many instances as to give rise to a suspicion (Duhring, *op. cit.*, p. 279) of some connection between them, at least in some cases.

DIAGNOSIS.—The affections with which impetigo contagiosa may most readily be confounded are eczema pustulosum and varicella. From the former it may be distinguished by the history of the case, the mode of development of the eruption, its inoculable and contagious character, and the features of the pustules. These are flat, isolated, itch but little, and are followed by superficial, flat crusts, which Fox well describes (*op. cit.*, p. 225) as seeming “stuck on.”

From varicella it may be distinguished by the fact that the pustules of impetigo contagiosa are larger, and that the crusts are totally different from those of chicken-pox. The eruption in this latter affection is much more copious, as a rule, develops much more simultaneously, and is more uniformly distributed over the various regions of the body.

PROGNOSIS.—As soon as the nature of the case has been recognized, it is readily curable; so that the prognosis is always favorable.

TREATMENT.—If signs of impoverished nutrition are present, careful attention to diet and hygiene, and the administration of tonic remedies, such as cod-liver oil, iron, and quinia are required. Strict cleanliness, and the prevention of the extension of the disease by scratching, are always to be insisted on. The local treatment needed is of the simplest character. The crusts, if any have formed, are to be removed by poulticing; any pustules that form to be punctured, and their pus removed by absorbent cotton; and every spot of eruption is to be dressed with the benzoated oxide of zinc ointment, to an ounce of which five grains of ammonio-chloride of mercury had better be added.

CHAPTER V.

PAPULAR OR PLASTIC INFLAMMATION OF THE SKIN.

THE essential feature of this class of skin affections is the formation of plastic lymph in the papillary, or sometimes in the deeper dermic layer. considerable confusion formerly existed in regard to the diseases that should be included under this heading. Willan recognized three, namely, strophulus, lichen, and prurigo. But a careful examination of the so-called varieties of strophulus has shown that in reality they are not all papular affections, but that widely different conditions have been grouped together under this title. We shall briefly describe lichen as the type of this class, referring especially to the only variety, *L. scrofulosum*, which is frequent in children. It will be remembered that the so-called lichen tropicus has already been carefully described in its proper place as a variety of urticaria. A short account will also be given of the lichenoid form of strophulus, and of prurigo.

ARTICLE I.

LICHEN.

THIS is a chronic inflammatory disease, characterized by the appearance of small papules, about the size of a millet-seed, either pale-red, slightly yellowish, or of the color of the surrounding skin. At first they may be isolated, but later are apt to become closely grouped together. Itching is often present, though not a constant symptom. The papules present no other change but the occasional formation of thin, delicate scales on their summits. The skin is dry and becomes harsh and thickened. The cause of the disease is essentially chronic, the papules lasting a considerable time, and new ones forming as the old ones slowly disappear.

Such being a general definition of lichen, some authors describe a variety termed *lichen simplex*. Fox (*op. cit.*, p. 139) admits its rarity while recognizing its existence. We have never met with a case of it in young children; and many authorities regard it as only the papulous form of eczema.

Lichen ruber, although also a rare affection, is one of the distinctive forms of papular disease of the skin. It does not occur in childhood, and we shall therefore limit ourselves to a mere definition of it. It runs a chronic course, and is characterized by the formation of papules, varying in size from a pin-head to a pea, and either flat (*l. ruber planus*) or pointed (*l. ruber acuminatus*). The papules are of a dull crimson red color, and may be either discrete or confluent, in which latter case the integument becomes much infiltrated and greatly thickened. When discrete the individual papules rise abruptly from the surrounding healthy skin, and are angular in outline instead of round as other papules. In the latter stages, considerable desquamation may be present. The disease is stubborn and essentially chronic in its course, but still usually yields to persistent treatment. In very extensive cases, a fatal termination may occur.

Lichen scrofulosus, on the other hand, requires careful study, as it most commonly occurs before the age of puberty.

The papules are always minute, about the size of pin heads or millet-seeds, and are pale-red or yellowish in color. They show a disposition to become grouped in patches with curved borders. The papules are found to be developed in connection with the hair-follicles, the new-formed exudation cells occurring both in and around the follicle. The eruption is usually limited to the trunk, being rare on the face or extremities. After the papules have existed for some time, their summits are apt to be covered with minute scales, and the skin becomes dry, harsh, and yellowish. It is important to note that there is very little or no itching. The course of the disease is essentially chronic, and it may last for years, although it yields quite promptly to appropriate treatment.

CAUSE.—As its name indicates, this affection is one of the manifestations of the scrofulous diathesis, and is apt to present itself in children or young persons who exhibit other marks of scrofula, as enlarged glands, ulcers, or necroses. From the accounts of it given by Hebra, Kaposi, and Kohn, it would seem to be a common enough affection in Austria, but it certainly is a very rare disease in this part of America, a statement confirmed by Duhring, who has not met with a single case.

TREATMENT.—The treatment that has been found successful is the free administration, internally, of cod-liver oil, together with its daily use by means of thorough inunction. The use of arsenic or of iodide of iron in conjunction is to be recommended.

LICHEN STROPHULUS, OR STROPHULUS.—Various opinions are entertained with regard to the true character and position of the affections grouped by Willan under the above heading. According to the older view, it was essentially the same as lichen, and its various subdivisions corresponded in the infant with those of the latter disease in the adult. But more careful study has shown that several of the subdivisions, made by Willan, really belong under forms of skin disease, are entirely and essentially different from lichen; and consequently the whole subject of strophulus has been discarded by some authors. We must recognize, therefore, that it is not to be admitted in this place, except with distinct limitations, but still, as some of its subdivisions appear to us to deserve to be regarded as lichenoid, and as they are the only truly papular affections occurring in young infants, it seems better to retain the familiar name for them.

We understand by strophulus, then, an inflammatory disease of the skin, of rather acute character, usually affecting infants at the breast, and distinguished by a more or less extensive, and sometimes a general, eruption of pale or vivid-red color, accompanied by more or less irritation and itching.

VARIETIES AND SYMPTOMS.—The *strophulus intertinctus*, or *red gum*, consists of an eruption of prominent pimples of a vivid red color, scattered here and there over different parts of the whole of the body, and intermingled with small erythematous patches. The eruption remains upon the skin for some time, the papules disappearing and reappearing in successive crops, for a week or two, or more, until they terminate by

desquamation. It is most common upon the cheeks, backs of the hands, and forearms.

In *strophulus confertus*, the papules are much smaller, more closely aggregated, much more numerous, and more confluent, than in the first variety, and they constitute a more severe eruption. It may be distributed over the whole surface, but is more commonly limited to a single spot, or to several regions, as the face, breast, or arms. The eruption is less vivid, but more lasting than that of the *strophulus intertinctus*, and usually reaches its height in twelve or fourteen days, and then subsides.

In *strophulus volaticus*, the papules, which are very ephemeral, are of a vivid color, and are disposed in small, not very numerous, circular groups, scattered over the surface of the body, but met with most frequently on the cheeks and arms. The ephemeral character, which is its only distinguishing feature, does not justify us in regarding *strophulus volaticus* as a distinct variety; and it possesses characters which show that it is closely related to urticaria.

The two remaining varieties, *strophulus albidus*, or *white gum*, and *strophulus candidus*, are both characterized by whitish instead of red papules. In the former, the papules are white, minute in size, and surrounded by an areola of a faint red color; they appear usually on the face, neck, and breast, and continue for some length of time. According to Tilbury Fox, the term *strophulus* is a misnomer as applied to this form, which he regards as a disease of the sebaceous glands. In *strophulus candidus*, which may more correctly be regarded as a form of urticaria, the papules are much larger, broader, more hard and tense, and are unaccompanied by any redness. They last usually about a week. This eruption is most common during dentition.

CAUSES AND PATHOLOGY.—The causes of *strophulus* are various disturbances of the digestive apparatus, aggravated in older infants by the irritation of the system due to dentition. Tilbury Fox (*op. cit.*, p. 155), regards *strophulus* as the result of hyperæmia of the sweat glands, and inclines to consider it as due to excessive clothing, to overheated rooms, and changes of weather. This view is based upon the observation that, while the characteristic papules of lichen are due to exudation into the papillary layer of the derma, the papules of *strophulus* may be seen clearly with a powerful glass to be seated at the sweat follicles. If this observation be correct, it would render it improper to retain *strophulus* among the truly papular affections.

DIAGNOSIS.—There is no difficulty in distinguishing *strophulus*, as it is the only papular eruption to which infants are subject. The absence of general symptoms and the extreme mildness of the disease are amongst its chief characters. It must be remembered that we only regard such papular eruptions as are unassociated with exudation or eczematous patches elsewhere on the surface as true instances of *strophulus*, since papules in all respects resembling those of this disease are to be frequently observed in cases of *eczema papulosum*.

PROGNOSIS.—The eruption is never attended with any danger. If severe

symptoms happen to coincide with it, they must depend on some other causes than the cutaneous affection.

TREATMENT.—As a general rule, strophulus needs no treatment whatever. In infants within the month, the irritation of the skin, if it be such as to disturb the comfort of the child, may be allayed by the use of the tepid bath, and by dusting by some mild powder, or by anointing with cold cream, glycerin and cold cream, simple cerate, or cocoa-butter. When any marked disturbance of the digestive apparatus is present, this should be attended to by the administration of mild laxatives, and of tonics, with some preparation of iron, as the tartrate or superphosphate.

In older children, in whom the disease appears to be associated with dentition, the local means spoken of above may be employed, while, at the same time, the gums should be lanced, if necessary, and any gastrointestinal disturbance removed by appropriate treatment.

ARTICLE II.

PRURIGO.

DEFINITION; FREQUENCY.—Prurigo is a chronic inflammatory disease characterized by an eruption, more or less extensive, of isolated papules, about the size of a small split pea, and pale red or of the color of the surrounding skin. They are developed usually on the extensor surfaces of the limbs, and give rise to the most violent and distressing itching, a symptom which constitutes one of the most marked features of the disease. Wilson includes it among the nervous affections of the skin, and attributes it usually to nervous debility, with an impaired state of the nutrition and innervation of the skin.

Prurigo is a rare disease in this city amongst the children of the middle and upper classes, since we have seldom met with it. In Europe, it is described as occurring in the children of the poor, though it is much less common than the eruptive diseases already treated of. Doubtless it occurs in this country also, but with the exception of a case reported by Wigglesworth, of Boston (*Amer. Jour. of Syph. and Derm.*, vol. iv, 1873, p. 21), we have not found any original account of it in the works of American writers. Duhring (*op. cit.*, p. 252) states that the disease is extremely rare and almost unknown in the United States.

CAUSES.—The only well-ascertained causes of the disease are the unfavorable hygienic conditions which exist amongst the destitute classes of society,—damp and ill-ventilated dwellings, unwholesome food, especially the use of salted meats and fish, and want of cleanliness as to person and clothes.

SYMPTOMS.—The papules of prurigo are small, but slightly prominent, and attended with moderate itching, constituting the *prurigo mitis*; or they are larger, more projecting, and attended with the most violent pruritus, forming the *prurigo ferox formicans* papule. The papules are usually

red or of the color of the skin, except when they have been torn by the nails, and are generally seated upon the outer surfaces of the limbs, and the upper part of the trunk.

When the itching is severe, the tearing of the papules by the nails causes the escape of a small drop of blood from the tops of many of them. The blood dries and forms so many small black crusts crowning the summits of the papules, a peculiarity which constitutes one of the most distinctive features of the disease. The papules terminate by absorption or by a slight desquamation. After the disease has lasted some time, the skin acquires, partly from the constant and violent scratching, a peculiar thickened and harsh character, which is most marked on the lower extremities.

The *duration* of the eruption is very uncertain. In acute cases, when properly treated, it may end in a few weeks, though it often, and indeed more generally, lasts for several months or years, or even through life.

DIAGNOSIS.—The only diseases with which prurigo is likely to be confounded are lichen or pruritis. It may be distinguished, however, generally with ease, by the facts that the papules of prurigo are larger, less numerous, and more extended, than those of strophulus or lichen; that in the latter diseases the papules are never crowned by the small black crusts of prurigo, and they are never attended with the same violent itching as the former.

From pruritus, it is to be distinguished by the absence in the former affection of papules, as well as of thickening and roughness of the skin. The regions affected in pruritus are also quite different from those most frequently involved in prurigo.

PROGNOSIS.—Prurigo is never, perhaps, a dangerous disease, though usually a very troublesome one from the severe irritation which attends it, from its not unfrequently obstinate resistance to treatment, and its disposition to relapse. According to Duhring, it is perhaps curable in children, but scarcely so when it has lasted until adult life.

TREATMENT.—The internal treatment of prurigo in children should consist in the use of sulphur, given alone or in the form of the compound liquid powder, if there is much constipation; of demulcent drinks, and of such remedies as may be rendered necessary by any disordered state of the digestive function. The diet must be carefully regulated. It ought to be nourishing and sustaining, but at the same time light and easy of digestion. In obstinate cases, recourse must be had to the administration of arsenic, cod-liver oil, iron, and other powerful nutrient and alterative remedies.

In addition to the internal treatment, simple warm-water baths, or emollient baths of flaxseed, bran, slippery-elm, or marsh-mallow, should be made use of in the early stage of the disorder. At a later period, alkaline baths, containing from three to eight ounces of carbonate of potash to each bath, according to the age, are recommended by Cazenave and Schedel. To allay the cutaneous irritation, mild ointments are often found useful. Billard employed with success, in a child six months old, inunctions with the oil of sweet almonds. Soaps or lotions, containing juniper tar or carbolic acid, are excellent anti-pruriginous applications; and relief will fre-

quently be obtained from the application of a dilute solution of chlorinated soda. When the case is obstinate, resisting emollient and alkaline baths, sulphurous baths must be made use of.

CHAPTER VI.

SQUAMOUS INFLAMMATION OF THE SKIN.

THE affections of this class are characterized by inflammatory hyperæmia of the derma, and hyperplastic growth of the cuticle forming scales or squamæ, and with a varying amount of secondary thickening.

There are two affections embraced under this heading of which, as they are of rare occurrence in children, only a brief description will be given.

ARTICLE I.

PSORIASIS.

PSORIASIS is usually chronic in its course, and is characterized by slightly elevated hyperæmic patches, of varying size and shape, which are covered with abundant, dry, silvery-white scales. If the scales are removed, which can be done readily, the cutis is seen to be inflamed, rough, and dry, or with merely a little blood exuding from mechanical irritation. It is essentially a disease of the upper layers of the corium and the papillæ, with hyperæmia, cell-proliferation, and with a remarkably copious formation of epidermic cells. Any part of the surface may be affected, but the disease shows marked preference for the extensor surface of the limbs and for the scalp.

Itching is usually present, and may be quite severe, especially in the early stages of the disease. It appears in several forms, depending chiefly on the size, form, and distribution of the patches of eruption. Psoriasis guttata is the name applied to the disease when it occurs as small, reddish, rounded elevations, more raised at the centre than at the circumference, and varying in size from a pin-head (also called *p. punctata*) to a large pea, and which soon become covered with fine, minute, whitish scales. Psoriasis circinata and gyrata are also described, in consequence of the wing-like or curving forms assumed by the eruption. When the patches are large, and irregular in shape, and cover a large amount of surface, the name of *p. diffusa* is applied.

CAUSES.—Psoriasis is a very rare disease in children. It seems not to occur in infants; but we have met with several well-marked cases, chiefly of *p. guttata*, in young children. In its more extensive and inveterately chronic forms it rarely if ever appears much before puberty. The most common period for its manifestation, according to Duhring, is

at about the age of twenty. It may also be inherited. The other predisposing causes are obscure. The syphilitic taint has no influence on the production of the true psoriasis. It affects both sexes, and all classes; but in children especially it seems to be associated with nutritive weakness, and perhaps with some special defeat of assimilation.

DIAGNOSIS.—The special localities affected, as the knees, elbows, and the extensor surfaces of the extremities; the absence of any stage of discharge; the abundant, silvery-white imbricated scales; and the rough, red, readily-bleeding surface beneath, render the diagnosis of psoriasis from eczema an easy matter. It is at times more difficult to distinguish psoriasis from the papulo-squamous syphiloderm, but attention to the following points will usually enable a diagnosis to be established. Psoriasis is more apt to be symmetrical, and to occupy certain localities, above mentioned. The patches of eruption in psoriasis are larger, occur simultaneously in more widely separated parts of the body, are more apt to be the seat of itching, and present much more copious formation of scales. The detection of any other evidences of inherited syphilis would of course aid greatly. In doubtful cases, the test of specific treatment should be resorted to.

PROGNOSIS.—Psoriasis is one of the most chronic and intractable of the inflammatory diseases of the skin; but we think it is more amenable to treatment in children than in adults.

TREATMENT.—In psoriasis it is especially necessary to pay strict and equal attention to the constitutional and to the local treatment. In all cases, the condition of digestion must be carefully examined, and any disorder that it may present should be removed as rapidly and effectually as possible by the proper remedies. Arsenic is the most valuable remedy we have in psoriasis on account of its peculiar tonic and alterative action. For children it is best prescribed in the form of the ferro-arsenical mixture:

R. Liq. Potassæ Arsenitis, f ʒj.
Vini Ferri, q. s. ad f ʒ iij.—M.

S.—Fifteen to sixty minims, according to age of child, three times daily in water immediately after meals.

In other cases, the following may be substituted with advantage:

R. Hydrargyri Bichloridi, gr. j.
Liq. Arsenici Chloridi, f ʒ ij.
Tr. Ferri Chloridi, f ʒ vj.—M.

S.—Four to eight drops in a wineglassful of water through a glass tube, three times a day after eating.

At the same time, cod-liver oil may be given with benefit, if it is well digested.

LOCAL TREATMENT.—In all cases, the crusts must be first removed by suitable means. If then the case is recent, only mild applications should be used, such as compresses wet with warm water; or inunctions

with oil of sweet almonds or olive oil ; or with soothing ointment or lotions. Tilbury Fox recommends that in extensive psoriasis, with irritability of the skin, the child should be soaked in an alkaline bath, containing 2 ounces of bicarbonate of soda, and 2 pounds of clarified size, every night for fifteen minutes, and then be thoroughly anointed. But in many instances, the disease has already reached a more chronic stage, requiring more stimulating applications. The particular mode of local treatment we have found most successful in such cases is by the use of alkaline tarry applications. Reference may be made to the brief description of Hebra's method in our remarks on chronic eczema (see page 947). Ointments of common tar or of oil of cade, or solutions of these substances in alcohol, are very valuable applications. Carbolic acid as a lotion or ointment may also be used, though less useful than tarry preparations in our experience. Various mercurial preparations, as dilute citrine ointment; an ointment of the protiodide, in the proportion of 10 or 15 grains to the ounce of simple cerate or cosmoline; solutions of the bichloride, 3 to 5 grains to an ounce of glycerin and water; are all successful in many cases, but should chiefly be used where the patches of eruption are quite limited in extent. Where much thickening of the skin exists, the use of caustic potash, in weak solution, or added in suitable proportion to some of the above recommended preparations, is desirable. Chrysophanic acid and pyrogallie acid, used in the form of ointment, in the proportion (for children) of 5 to 15 grains to the ounce, are among the latest remedies suggested for the treatment of psoriasis. They are applicable chiefly to cases of circumscribed eruption, when they often give excellent results even in chronic and very obstinate cases. In the use of any of these stimulating preparations in psoriasis, it is very important to begin with weak ointments or solutions, to use *very small quantities*, and to rub them thoroughly into the surface, and if very undue irritation is caused, to immediately resort to soothing or milder applications.

ARTICLE II.

PITYRIASIS RUBRA.

THIS very rare form of disease does not, we believe, occur often in young children. It usually affects the whole surface, and is attended with deep-red coloration of the skin, due to hyperæmia, and with abundant desquamation in the form of large, thin, whitish scales. There is little if any infiltration of the skin, and no discharge occurs at any time. There is excessive sensitiveness to damp and to changes of weather; but the patient suffers little, as a rule, from itching. As already stated, the entire surface, including the palms of the hands and the soles of the feet, is usually affected, and the nails are not rarely shed. This form of the disease occurs in anæmic and enfeebled subjects, and is a chronic and stubborn affection. Careful attention to its peculiarities will prevent its

being confounded with general eczema squamosum, psoriasis, lichen ruber, or pemphigus foliaceus. The internal treatment should consist of cod-liver oil, with iron, quinia, or mineral acids. Locally, the use of mild soothing alterative applications is to be recommended. Fox has found very good results follow from keeping the patient wrapped up in olive oil.

There seem also to occur analogous conditions of more limited extent in children, to which the name of pityriasis simplex may be appropriate. Thus an affection of the scalp, with innumerable small, thin, whitish, furfuraceous scales, but without any inflammation or infiltration of the scalp, is occasionally met with. Sometimes this is connected with irritation of the sebaceous follicles, and is really a seborrhœa. But in other cases it is due to a simple hyperplasia and rapid desquamation of the epidermic layer. It is a condition of small importance, and requires merely strict cleanliness and the use of mild lotions or ointments; as weak solutions of bichloride of mercury, benzoated oxide of zinc ointment, dilute citrine ointment, etc. If the child's nutrition is impoverished, careful attention to diet and hygiene, and the internal use of iron and arsenic, are desirable.

CHAPTER VII.

HYPERTROPHIES.

THIS group includes a considerable number of affections of the skin, since each of the anatomical elements of this tissue (pigment, epidermis, papillæ, corium), as well as its appendages (hair and nails) are liable to be affected by hypertrophy. The only two of this class of diseases, however, which occur with greater frequency or with unusual features in childhood, so as to demand consideration here, are *ichthyosis*, due to a hypertrophy of the epidermis; and *sclerema*, due to a hypertrophy of the corium.

ARTICLE I.

ICHTHYOSIS.

ICHTHYOSIS is a chronic disease of the skin, usually affecting the whole surface, characterized by dryness, harshness, desquamation, and more or less papillary hypertrophy. It is observed in the two forms of *ichthyosis simplex* and *hystrix*. *Ichthyosis simplex* varies much in its degree of development. When there is merely a dry, harsh, ill-nourished condition of the skin, with slight furfuraceous exfoliation, it is termed *xeroderma*. But in its more usual, fully developed form, the *simplex* is characterized by a high degree of dryness and harshness of the skin, together with an extensive production of variously sized and shaped fish-like scales. The skin has a dirty, unwashed look; and the scales are white and silvery, or

at times yellowish or greenish. The scales correspond in size and shape with the spaces between the normal lines and furrows of the skin. The entire surface is usually affected, but the face and the flexures of the joints suffer least, while the elbows and fronts of the knees are specially involved.

In ichthyosis hystrix, the skin may also be uniformly affected, or else there may be scattered patches of various sizes occurring on any part of the body. These patches are hard, rough, elevated, and of a yellowish, brownish, or greenish color, often resembling dried mud. They are made up of thickened epidermis and enormously hypertrophied, hard or even horny papillæ.

ETIOLOGY.—The disease is often called a congenital one, and although this is not strictly true, since it does not make its appearance until towards the close of the first or second year of life, it is probable that the tendency to the disease exists from the time of birth. Occasionally several members of the same generation of a family may be affected; in other instances, the disease appears hereditary; but for the most part it occurs without ascertainable cause in isolated individuals.

PATHOLOGY.—The papillæ and the epiderm are chiefly affected. The papillæ are enlarged or elongated, and are infiltrated with cells. The mucous and horny layers are both thickened with accumulations of heaped up cuticular cells. The sebaceous glands are frequently atrophied. Fox states that the inorganic salts in the skin, especially those of lime and iron, and silicic acid, are increased in quantity.

The milder form, xeroderma, may be confounded with other harsh, ill-nourished conditions of the skin; but the well-developed disease in either of its forms can be recognized without difficulty.

TREATMENT.—The internal use of alteratives or nutrients, as arsenic, cod-liver oil, or iodide of iron, would seem to be indicated, and some authors have found them beneficial. But our own experience agrees with that of Duhring, and others, who state that external treatment alone is of service. The general plan that will be found most useful is to favor softening of the hypertrophied layers of epidermis by warm baths, or by vapor or alkaline baths; and then their removal by kneading or friction of the skin, with or without the aid of soft soap or some analogous alkaline application. This process is to be followed by the inunction of the whole surface, once or twice daily, with olive oil, cocoa oil, or vaseline; and after several days of such rubbing, the bath may be repeated. In this way the accumulations of hardened epiderm are prevented, and the skin is kept comparatively soft and pliable; but no treatment has yet been found that rapidly cures the disorder, which continues throughout life.

ARTICLE II.

SCLEREMA.

DEFINITION; SYNONYMS; CAUSE; FREQUENCY.—This peculiar affection, characterized by induration of the skin and subcutaneous tissue, with or without œdema, has been described by numerous writers, almost each one of whom has given a distinct name and theory for the disease. Among these names the most appropriate are sclerema, scleriosis, scleroderma, induration of the cellular tissue, or chorionitis.

It is an affection not altogether peculiar to infants, though it is rare to find it well developed after the first few months of life. There are, however, a sufficient number of cases in adults on record to establish the fact of its occasional occurrence at all ages. It must be a very rare disease in this country, even among infants, and especially in private practice, as we have met with but three well marked cases in adults, and but one instance occurring in childhood, which was the case in which imperfect induration of the skin was developed in connection with atelectasis pulmonum, to which allusion is made in our article upon the latter affection.

In the large foundling hospitals in Europe, however, where so many causes exist to depress the vitality of the infants, it is of very frequent occurrence. Under such circumstances, moreover, it generally develops itself within the first twelve or fourteen days after birth.

The most varied causes were formerly assigned for this disease, before the researches of Bailly and Legendre appeared to point out sclerema as one of the results of imperfect expansion of the lungs. As we have already remarked, it is seldom observed among the children of the upper classes of society, so that all those conditions which depress the strength of the child, as insufficient or unhealthy nourishment, imperfect clothing, cold, especially when associated with moisture, may be considered as the predisposing causes of sclerema. The influence of dampness and cold in developing this affection is shown by the fact, deduced from numerous statistics, that twice as many children are attacked during the cold and wet months of the year as at other times, although there are cases recorded as occurring in the hottest months.

Authors still differ in regard to the relation between atelectasis and sclerema. West¹ accepts the results of the researches of Bailly and Legendre, and considers it a result of the imperfect expansion of the lungs. Bouchut,² on the other hand, regards the changes found in the lungs as the result rather than the cause of the induration of the skin.

Letourneau³ agrees with West in regarding sclerema as a condition depending primarily on congenital weakness, imperfect expansion of the lungs, and defective hæmotosis. According to his view it is a slow asphyxia,

¹ Diseases of Children (3d Amer. ed.), p. 238.

² Diseases of Infancy (Bird's trans.).

³ Letourneau, Sclerema and Œdema. Paris, 1858 (Canstat. Jahrb., iv, 456).

the body becoming gradually cooled down and the child remaining in a state of organic torpor until death occurs.

SYMPTOMS.—The disease presents some variety of symptoms according as it occurs in early infancy or in more advanced life.

In infants the induration of the skin appears within two weeks after birth, either with or without a preceding febrile condition for a day or two. It invades successively the feet, hands, limbs, the back, the face, and finally involves the entire surface of the body. At this early age, the skin retains its reddish tint in the affected parts; whilst later in life, the surface assumes a dull, slightly yellowish aspect. The skin becomes hard, is with difficulty pinched up, and instead of thinning, remains thick and wax-like. The parts appear somewhat swollen, though never to any great extent, and pressure with the finger scarcely leaves an impression on their surface. Occasionally the induration is associated with an effusion of serum beneath the skin, and when this exists, the surface is much more readily indented.

It is this occurrence of œdema which has led some observers to consider sclerema as a form of anasarca; it is probable, however, that its presence is merely a result of the obstructed cutaneous circulation, and that it does not, in reality, constitute an essential element of the disease. The skin is also quite frequently jaundiced. The children usually preserve the power of moving the affected parts, and there is no loss of cutaneous sensibility. The temperature of the body, and especially of the indurated portions, rapidly decreases, so that from 100° it may fall to 90° , 80° , and even, in some exceptional cases recorded by Roger, to 73° and 70° .

The little patients appear to suffer much pain during this disease. They utter a sharp, abrupt, isolated, but very frequently repeated cry, quite characteristic of the affection, and occasionally they present nervous symptoms, such as twitching of the hands or more general convulsive movements.

The strength fails rapidly, and they soon become too weak to suck. The pulse is feeble, though not much accelerated, unless some complication has ensued. The appetite fails, and the bowels are constipated, unless there is entero-colitis, which occurs in a few cases. This condition is naturally attended with great emaciation, as we find in Elsässer's¹ cases, where the average loss of weight was three-fourths of a pound, the extreme being six ounces and two pounds.

The respirations are imperfect, and, after a short time, cough makes its appearance and continues throughout the case, indicating the occurrence of either pneumonia or collapse of the lungs, which are by far the most frequent complications, even if the state of atelectasis be not regarded as an efficient cause of sclerema. The disease, however, is not always so general and severe as above described; occasionally it occurs in limited portions of the body, and without any very alarming symptoms.

In later life the disease is more frequently thus limited to small portions of the body; the symptoms follow a more chronic course, and are somewhat

¹ Sclerema, Arch. Gén., N. S., t. i., 1853, p. 531.

amenable to treatment. Rilliet and Barthez¹ describe an acute and chronic form, and mention the following symptoms as distinguishing the disease in the adult: the severe epigastric pain associated with violent palpitations, the less acute progress of the case, and the more frequent implication of the serous membranes.

In one well-marked case, occurring at the age of thirty-five years, which one of us had the opportunity of observing during its entire course, these symptoms were very prominent.

PROGNOSIS.—In infants, when the induration is at all general, the disease almost invariably terminates fatally in from two to six days. Under favorable circumstances, however, and when the induration is limited, resolution may occur, and the case terminate favorably; though it requires from fifteen days to a month to effect the cure.

The fatal result is either caused by the gradual exhaustion of the vital powers, or by the supervention of one of the complications already mentioned, by far the most usual of which are lesions of the lungs.

In later life, when the disease tends to recovery, a long time may be consumed before the induration completely disappears. Rilliet and Barthez report a case, occurring in a girl aged eleven years, which lasted two years, although it was at no time very general or accompanied by very severe symptoms.

Of 53 cases reported by Elsässer, all but four proved fatal, either from the sclerema itself, or from some incidental disease.

DIAGNOSIS.—The absence of any lesion of the internal organs, together with the perfectly characteristic appearances of the induration, render an error of diagnosis almost impossible.

ANATOMICAL APPEARANCES.—The induration of the surface persists after death, and on incising the part, a turbid fluid, resembling that of anasarca, often flows out. The subcutaneous tissue is also indurated, and the fat is found in the form of solid granules. This layer, which varies from one-half a line to three lines in thickness, is sometimes followed by a gelatinous one.

The fluid which is contained in the meshes of the tissues has been subjected to analysis by several observers, but with conflicting results: Chevreul and Breschet reporting that it contained a plastic matter, spontaneously coagulable on contact with the air, which they were inclined to regard as characteristic of the disease; whilst Billard, on repeating this observation with fluid derived from an ordinary case of anasarca, found it to possess the same property. This subject, therefore, of much importance in regard to the pathology of sclerema, requires to be more fully investigated.

Different observers are not agreed as to the condition of the corium and subcutaneous tissue. In sclerema adultorum the essential element in the changes of the skin appears to consist in a morbid increase of the connective tissue, associated with a marked development of lymphoid cells, by multiplication of the cells in the perivascular sheaths of the minute bloodves-

¹ Op. cit., t. ii., p. 106.

sels of the derm and subcutaneous tissue. This condition was pointed out by Rasmussen (translated in *Edin. Med. Journ.*, vol. xiii, part i, pp. 200 and 318), and accords with our own investigations. In sclerema in children, Jenks¹ and Löschner² have observed marked increase in the connective tissue of the corium; but this condition has not been found by other observers.

The indurated tissue is traversed by numerous vessels, permeable, and for the most part gorged with dark blood. Bouchut believes that the cutaneous capillaries are in great measure obliterated in the indurated parts, and that the œdema which occasionally coëxists with sclerema is due to this obliteration; founding his opinion upon an unsuccessful attempt to inject the skin of a limb affected with sclerema, although the injecting fluid freely entered all the deeper tissues. The observations of Elsässer, however, render this view doubtful, since in 49 cases, he failed to find this condition.

Apart from these morbid changes in the skin and subcutaneous tissue, there is no lesion characteristic of sclerema. In a large number of cases, however, the lungs present some abnormal condition. According to Bouchut they are often gorged with blood, and here and there contain patches of lobar pneumonia; conditions which he regards rather as the result than the cause of sclerema.

Elsässer found lobular pneumonia present in one-tenth of his cases; and in one-third of them, portions of the lungs were impermeable to air. We have already stated that West, following the researches of Bailly and Legendre on atelectasis, believes that sclerema is one of the results of this persistence of the foetal condition of the lung, not differing in its essential nature from œdema following pulmonary obstruction. The occurrence of undoubted cases of sclerema in the adult, and the frequent absence of atelectasis in well-marked cases of sclerema in infants, appear, however, to render this view untenable.

The entire venous system and the cavities of the heart are distended with dark fluid blood; but the heart presents no constant condition to which could be attributed the production of the disease. The jaundice which has been mentioned as occasionally existing, is not found to be associated with any abnormal condition of the liver, excepting congestion. Entero-colitis is a rather frequent complication of sclerema, and has been regarded as influencing its development; but this view has long since been abandoned. Elsässer found intestinal lesions and hyperæmia of the abdominal viscera quite commonly; and in eight of his cases, peritonitis was present.

TREATMENT.—The preventive treatment of sclerema consists in attention to all the hygienic conditions of the young infant.

The curative treatment implies the removal of all the causes, and the application of remedies calculated to restore the force of the circulation, and the function of the skin. *Warmth* stands foremost as a curative measure, and recourse may be had to warm baths or hot vapor-baths, and

¹ Amer. Journ. of Obstetrics, May, 1871, p. 129.

² Prager, Vierteljahrschrift, 1868.

to frictions with hot oil; hot sand or bran-bags may be applied to the surface, and the temperature of the room should be carefully regulated.

The child should be nourished with breast-milk; and stimulants, such as wine whey, should be freely given. Cordial and aromatic draughts are also recommended, which may be formed of any of the diffusible stimulants.

As there is reason to believe that some relation exists between sclerema and atelectasis pulmonum, we should, in addition, resort to all those means especially adapted to remove this condition, for a full account of which we refer the reader to the article on collapse of the lungs.

The same plan of treatment is advisable in cases in adult life.

By these means we may hope to arrest, and even cure this strange affection, when it has not involved any considerable portion of the surface.

CHAPTER VIII.

SECTION I.

PARASITIC SKIN DISEASES.

GENERAL REMARKS.—The diseases now regarded by many authorities as due to the presence of a vegetable parasite upon the skin are as follows:

1. Tinea Favosa or Favus, Parasite: Achorion Schœnleinii.
2. Tinea Trico-phytina, $\left\{ \begin{array}{l} \text{Tinea Tonsurans (Ringworm of scalp),} \\ \text{Tinea Circinata (Ringworm of body),} \\ \text{Tinea Sycosis (Ringworm of beard),} \end{array} \right\}$ Parasite: Tricophyton.
3. Tinea Versicolor (Chloasma, Wilson), . . . Parasite: Microsporon Farfur.
4. Tinea Decalvans (Alopecia Areata), . . . " Microsporon Audouini.

There are also diseases of the skin due to the presence of animal parasites, namely scabies or itch: and pediculosis due to the presence of lice. The latter of these does not require description here.

There are several questions in regard to the vegetable parasitic affections upon which doubts still exist, and which are of so much importance as to demand a brief examination.

In the first place, it can scarcely be doubted by any one familiar with the use of the microscope, and who has taken the trouble to examine the subject, that parasitic fungi are found with remarkable constancy in the eruptions of these diseases. The opinion advanced by Wilson (*Br. and For. Med.-Chir. Rev.*, 1864; and *Diseases of the Skin*, 7th Amer. ed., p. 614), that the structures found in these cases, are due to a peculiar "granular" degeneration of the normal elements of the part, owing to which they lose their power of developing into healthy epithelial structures, but retain their power of proliferation, appears to us opposed to all sound reason and accurate observation.

In addition, however, to the evidence furnished by the chemical and microscopical examination of the growths in question, their fungous nature is shown by the facts that they can be cultivated after removal from the body, and that the diseases with which they are associated are contagious and can be communicated by inoculation to healthy persons, or even to some of the lower animals.

In searching for these growths, the scraping from the surface of the diseased spot, or the hairs which traverse it, may be taken for examination; but before subjecting them to microscopic study, they should be treated with dilute acetic acid to render them more translucent, and subsequently with a little sulphuric ether to remove the fatty granules which often obscure the fungus.

The structures which the fungi affect are the hairs with their follicles, and the epidermis.

The special alterations which the hairs undergo will be detailed under the head of the different diseases; the fungus gains entrance to the follicle, penetrates the bulb of the hair, insinuates itself between its longitudinal fibres, thus splitting it up and rendering it brittle. In the epidermis the fungus is said at first usually to appear beneath the superficial layer, until, by its development, it causes such irritation as leads to the exfoliation of this layer, when it reaches the surface and then multiplies rapidly.

The objection which has been based upon this fact, that the growth cannot be a parasitic one, does not seem to us of much force, since it is easy to account for the introduction of such extremely minute bodies as the spores of these fungi beneath the superficial layer of the cuticle.

Admitting then the presence of these parasitic growths, a more interesting question arises in regard to the relation which exists between them and the diseases with which they are associated; whether, that is, they are essential to, and actually the causes of the respective diseases, or are merely accidental, and are present only because they find a suitable nidus for development in the diseased skin. Opinions are at variance upon this question, but there are at least two considerations which render it probable that the fungi are essential rather than accidental productions. The first of these is, that they are present in the early stages of the disease, before any considerable inflammatory change has occurred, and that in proportion as suppuration ensues they diminish in abundance. And, secondly, that, as already stated, they are capable of transmission to perfectly healthy persons by inoculation.

There can, however, be no doubt that the development of the fungus, under ordinary circumstances, is greatly favored by the constitutional condition of the patient and the state of the cutaneous surface. Thus it is especially in children of a delicate or strumous constitution, that these various diseases are most frequently met with; and when, in addition, personal filthiness with inattention to properly combing and cleansing the hair, and changing the clothing, are combined, the spores find the most favorable conditions possible for their rapid development.

There remains the further question, upon which authorities are still di-

vided, whether there are various fungi concerned in the production of these diseases, or whether the apparently different species are merely different stages of a single fungus. For the sake of greater ease of reference and comparison, we will here give a brief description of their characteristic appearances.

Fungus of Tinea Favosa.—In the earliest stage of development of the favus crust, it is still covered by the superficial layer of epidermis; but later, when this is ruptured, it presents an envelope of a sulphur-yellow color, which on microscopic examination shows a homogeneous or finely granular substance. The interior, of a pale white color, is the true favus matter, and consists of the sporules, thalli, and mycelia of a fungus named the *achorion schoenleinii*, in honor of Schoenlein, who first fully described it.

The sporules are of a rounded, or more frequently of an oval form, and have well-marked edges, and a homogeneous and slightly opalescent interior. Their average diameter is about $\frac{1}{3000}$ th of an inch. Many of these sporules are seen to be grouped together, while some are more elongated and present a contraction in the middle; others are nearly triangular in form, with rounded angles; others, yet more elongated, are marked with several contortions. Some sporules, completely formed, seem to have a double enveloping membrane, and others present in their interiors something like a nucleus.

There are also present numerous diaphragmated tubes, formed by the development and confluence of the sporules, which are either simple or present ramifying branches. These tubes vary in diameter from $\frac{1}{4000}$ th to $\frac{1}{1500}$ th of an inch, and are either empty or have granular contents. Amongst the sporules and mycelia, especially towards the circumference of the cups, may be seen a considerable number of molecular granules, which are probably imperfectly developed sporules.

Fungus of Tinea Tricophytina.—The next parasite, the *tricophyton*, is that which produces *tinea tonsurans*, *tinea circinata*, and *tinea sycosis*.

The microscopic characteristics of this parasite, as first described by Malmsten, in 1845, and since confirmed by numerous observers, are very numerous rounded or oval sporules, about $\frac{1}{7000}$ th of an inch in diameter, which are isolated or united together into chains, and a comparatively small number of mycelial thread.

Again, the parasite, which by many observers is believed to cause *tinea versicolor*, is the *microsporon furfur*, discovered by Eichstädt, in 1846. This fungus presents numerous rounded spores, and long tubes. The spores are about $\frac{1}{5000}$ th of an inch in diameter, and are frequently collected together in large clusters, like bunches of grapes (Anderson). Some of the tubes observed are simple, and others jointed.

In regard to the parasitic nature of alopecia areata, there is great doubt, and even so warm a supporter of the fungous origin of the other diseases we have mentioned as Dr. Anderson, does not allow it.

Numerous observations have been made which go to show the existence of a very wide range of variation as regards form in these fungi; and have led some observers to assert not only the identity of these particular

forms, but indeed to refer all varieties of epiphytic fungi to some one central type.

The evidence upon which this view rests, mainly drawn from the results obtained from germination of the various fungi, and from the study of their transitional forms, cannot at present be considered conclusive; and further investigation of the question is demanded.

It is, however, thought by some high authorities, that no doubt can be entertained in regard to the identity at least of the parasites which produce the various forms of tinea, including the achorion of favus, the microsporon furfur of tinea versicolor, and the tricophyton of the various varieties of ringworm. The most complete exposition of the arguments upon which this view is based, will be found in Dr. Tilbury Fox's admirable treatise on skin diseases of parasitic origin (London, 1863).

On the other hand, some eminent dermatologists believe that the fungi which produce these diseases are essentially distinct. The arguments upon which they base this opinion may be briefly expressed as follows, in the language of Dr. Anderson (*loc. cit.*, p. 170).

That in all cases of successful inoculation with the achorion, tricophyton, and microsporon furfur, the same parasitic disease has been produced as that from which the parasite was taken. That of the innumerable cases occurring in the human subject, illustrative of the contagious nature of favus, tinea tonsurans, and tinea versicolor, there is no authentic case in which one of these diseases gave rise to one of the others.

That the difference in the appearance of the eruption, when fully developed, is so very striking as to lead to the belief that they are produced by separate parasites.

That there is no authentic record of the transition of one of these diseases into one of the others.

That the microscopic differences between the three fungi are in many cases sufficient to base a correct diagnosis upon.

That of the numerous instances on record of the transmission of tinea favosa, and tinea tricophytina, from the lower animals by contagion or inoculation, favus has always given rise to favus, and tinea tricophytina to tinea tricophytina.

We regard then the parasitic nature of these affections as undoubted, but more extended observation is necessary before the relations of their respective fungi can be determined.

ARTICLE I.

TINEA FAVOSA OR FAVUS.

FAVUS is a parasitic disease of the scalp, long confounded by different writers with other and very dissimilar affections of that part. In consequence of this confusion it has received a great variety of names, of which the most generally known are porrigo and tinea. In adopting the above

title, we follow the example of Erasmus Wilson and other recent authorities, amongst the English, and of MM. Rilliet and Barthez, Gilbert and Rayer, amongst the French.

DEFINITION; SYNONYMS; VARIETIES; FREQUENCY.—Favus is a specific contagious eruption of the scalp, characterized by inflammation of the hair-follicles dependent upon the presence of a peculiar fungus, the *achorion schœnleinii*. It is distinguished at first by small yellow pustules, countersunk in the skin; these are soon converted into yellow cup-like crusts, which adhere often for a very long period. It usually causes permanent loss of hair at the affected part.

The disease is described by most of the former English writers under the title of *porrigo*, but as several other eruptions have been included under the same name, we think it best to follow the example of Mr. E. Wilson, and call it *favus*. By MM. Bielt and Cazenave it is designated, after Willan, *porrigo favosa* and *porrigo scutulata*. MM. Rayer and Gilbert, as above mentioned, give it the name of *favus*.

There are two varieties of *favus*, the *favus dispersus*, the *porrigo favosa* of most writers, and the *favus confertus*, the *porrigo scutulata* of many observers.

The disease is much less frequent than *eczema* of the scalp, but is nevertheless constantly met with amongst the crowded populations of Europe. In this country it is more rare, and amongst the middle and upper classes, at least of this city, is almost unknown, since we have never met with a case of it in our own private practice, though we have occasionally seen it in the hospitals here.

CAUSES.—The only well-ascertained exciting cause of *favus* is generally thought to be *contagion*, a quality of the disease acknowledged by most observers, though denied by Mr. E. Wilson, who considers its cause a debility of nutritive vitality, allied with *struma*. It may be propagated by direct contact of the disease with a healthy skin, or by means of combs, brushes, or other articles of the toilet; and it is also probable that the spores may be carried by the atmosphere so as to communicate it by infection. It has been frequently propagated by direct inoculation,—by Remak, Bennett, Hebra, Bazin, Gruby, Köbner, etc.

Favus is also said to be met with in the lower animals, and especially amongst mice and cats; and cases are on record which render it highly probable that it may be communicated from them to the human subject.

It occurs at all seasons, attacks either sex indifferently, and is met with at all ages, but is especially frequent in children and young people, and, indeed, when met with in adults, is usually found to have commenced in early life, and to have persisted for years. Certain conditions act as predisposing causes in its production, and may alone, perhaps, give rise to its development. These conditions are unhealthy hygienic influences, as unwholesome and insufficient food, poverty, filth, and the living in low, damp, and ill-ventilated dwellings. It is met with most frequently in persons of feeble, lymphatic, and especially in those of scrofulous constitution, though, be it remarked, it occurs also in persons of strong and vigorous health.

Among those who believe in its truly parasitic nature, there are some, as Devergie, who believe that it may be spontaneously generated, the parasite originating in the body of the affected person. One of the facts upon which the theory is based is the asserted occasional cure of the disease by internal remedies, but we believe that these can only relieve it by fortifying the system, and so removing the conditions which favored the development of the parasite.

SYMPTOMS.—FAVUS DISPERSUS, OR PORRIGO FAVOSA.—This variety begins with very small pustules of a peculiar straw-yellow color, which exhibit from the first the special character of not being raised at all above the level of the skin. Directly after their formation, the yellowish matter which they contain begins to concrete, and there can be perceived from this early period a central depression in the crusts, which becomes more marked as these augment in size, so that at the end of five or six days it is perfectly evident. Each pustule, and of course each crust is, as a general rule, traversed by a hair. The favus crust is a very remarkable feature of the disease, and it is in itself a pathognomonic symptom. As it increases in size, which it does gradually until it reaches in some instances a diameter of half an inch, the central depression above spoken of becomes more and more distinct, and the crust assumes, from this circumstance, the shape of a cup with an inverted edge. Their structure is made up of a series of concentric layers, or layers or rings, compactly arranged one upon the other. This cup-like form with the concentric arrangement of layers, the peculiar straw-yellow color, and the fact that each crust is usually pierced by a hair, are the distinguishing characters of the disease.

The pustules are usually isolated at first, though they may be arranged in groups of irregular size. When numerous, the crusts, by their gradual enlargement, touch at their edges, and blend into larger or smaller patches of irregular shape, but still presenting many little depressions corresponding to the first formed pustules. In rare cases, the disease is so extensive as to form a kind of mask covering the whole scalp.

When the disease is not interfered with by treatment, the crusts remain adherent for a long time,—for months or even years; they become also paler in color than they were at first, and so dry and pulverulent, as to break very readily when rubbed or touched. They become, moreover, thicker and more massive, and lose their first regular cup-like form, from the disappearance of their depressions, and from the irregular and uneven shape given to their edges and surfaces, by the breaking which they undergo. When the case runs on in this way, the head exhales a most unpleasant odor, which has been compared to that of mice or the urine of a cat; McCall Anderson has, however, noticed a very similar odor in cases of eczema impetiginoides of the scalp. In some instances, where the disease is grossly neglected amongst the very poor, pediculi form in abundance amidst the crusts, and add to the disgusting appearance of the disorder.

When the crusts have been removed by any means, the surface of the scalp is seen to be red, moist, and to present slight erosions or even ulcerations. The crusts are reproduced only by the eruption of new pustules.

An invariable and unfortunate sequel to the favus disease is a more or less extensive loss of the hair. The hairs become loose from a very early period of the disease, and can be pulled out with great ease. As the case goes on they fall out, and the scalp is left smooth, shining, uneven, and deprived of hair. On these spots the hair seldom grows again, and if it does, it comes out thin, woolly, and with every appearance of weakness and unhealthfulness.

Though the usual and favorite seat of favus is the scalp, it is met with occasionally on the forehead, temples, chin, and eyebrows, and, in still rarer instances, on the shoulders, elbows, forearms, on the upper and outer parts of the legs and thighs, on the scrotum, and even on the nails. The nails are also liable to be invaded, probably in consequence of the parasite gaining entrance during the act of scratching. The affected nails become thickened, yellowish and opaque, and brittle. Even in such cases, however, it has frequently existed first on the scalp, and extended thence to the other parts, though it may sometimes begin upon the trunk or limbs in consequence of a direct application to them of the contagious element.

FAVUS CONFERTUS, OR PORRIGO SCUTULATA.—In this variety of favus the pustules are arranged so as to form circles or rings upon the forehead or scalp, instead of being dispersed irregularly over the scalp, as in the preceding variety. The disease begins with red, circular patches, attended with a good deal of itching, upon which, after a short time, appear small yellow pustules, that seem to be sunken in the skin. The pustules are more numerous on the circumference than at the centre of the red patch or disk; or the latter increase in size by the extension of the disease to the follicles just beyond its outer edge. The pustules are exactly like those of favus dispersus, except that their yellow color is of a lighter tint. They desiccate very rapidly, and form crusts which are very thin at first, never very thick, and of an irregular shape.

When the disks are very numerous, either originally, or by propagation of the disease from part to part, they meet at their borders, blend together and give to the scalp the appearance of an extensive and irregular crust, presenting at its circumference curved lines, marking the segments of circles, of which the whole is composed. The crust has sometimes covered the whole scalp, excepting merely a small border at its circumference, where may still exist some scanty remains of the hair.

When the crusts are removed, the surface beneath is found to be red and tumid, according to Wilson, and to present numerous yellow points. Cazenave and Schedel state that when the crusts fall, they leave exposed a large, uneven, furfuraceous patch, upon which new favus pustules do not appear often for a long time. The hair is in great measure destroyed over the diseased surfaces, though not so completely, it is said, as in the other variety.

Favus is not, in either variety, attended with constitutional symptoms.

The only marked local symptom complained of is the itching, which is always greatly aggravated by want of cleanliness.

NATURE OF FAVUS.—We have already, in our general remarks, intro-

ductory to this class of skin diseases, given the arguments which prove their parasitic nature.

Mr. E. Wilson, alone among dermatologists of note, persists in regarding favus and the others, as due to mere alterations in the nutrition of the skin dependent upon constitutional nutritive debility; and he refers the characteristic fungous elements revealed by microscopic examination, merely to a peculiar granular degeneration of the epithelial elements.

We refer the reader, for a more full discussion of this question, to the works already quoted, merely adding here that, in our opinion, the results of microscopic examination, the results of inoculation of the parasite in man, as well as in the lower animals and plants, the undoubtedly contagious nature of the disease, and finally the astonishing and never-failing success of the local treatment when properly carried out, conclusively show its parasitic nature. The reader is also referred to the remarks introductory to this chapter for a full description of the parasite, the *achorion schœnleinii*, which is the essential cause of favus; as well as for the differences which distinguish it from the parasites which are found in the various forms of tinea.

DIAGNOSIS.—The diagnosis of favus rarely presents any difficulties. The peculiar pustules which exist at first—small, yellow, on a level with or below the surface of the scalp, and the crusts which so soon follow these, saffron-yellow in color, dry, and cup-shaped, will mark a case of favus *dispersus* from every other disease. In favus *confertus* the same characters exist, but the crusts and pustules are arranged on circular erythematous disks, instead of being isolated or dispersed as in favus *dispersus*.

From impetigo of the scalp, which is the only disease with which it is at all probable that it would be confounded, it may readily be distinguished by an examination of the primary characters of the two disorders. This primary character can always be found by searching at the outer edges of the diseased surface. In favus the pustule is small, depressed, and contains very little fluid, while in impetigo it is large, globular, and projecting. The crusts are very different: in the former dry, as though dusted with sulphur, cup-shaped, depressed, and usually traversed by a hair; in the latter, rugous, irregular in shape, not cupped, resting above the skin, and generally somewhat moist and soft. The peculiar odor which is present in cases of favus may be of assistance, although a very similar odor has been observed in cases of different characters. The microscopic examination of the hair or crusts in favus also shows the presence of the *achorion schœnleinii*, which is never met with in impetigo. Lastly, the alopecia which so constantly results from favus, does not occur in impetigo.

PROGNOSIS.—Favus is a serious disease because of its usually long duration, the difficulty often experienced in effecting its cure, and because of the loss of hair which it occasions.

TREATMENT.—The treatment of favus should be both general and local, for though some writers, and particularly Cazenave and Schedel, state that it must be altogether external, and that in spite of numerous trials they do not feel authorized to propose any internal means (*Malad. de la Peau*, 4ème éd., p. 326); others, as Wilson, Bennett, and Neligan, recom-

mend constitutional remedies as of very great importance in assisting the cure.

The *general* treatment must be such as may seem called for by the state of health of the individual patient. When, as so often happens, the disease occurs in a scrofulous person, cod-liver oil, iodide of potassium, nourishing food, air, and exercise, are of the utmost importance. When the health of the patient is feeble and broken from the want of wholesome and abundant food, from insufficient clothing, or from residence in a vitiated, close, and confined air, the removal of these conditions, which undoubtedly act as predisposing causes in the production of the disease, cannot but aid in its cure. Dr. Neligan (*Dublin Quart. Journ. of Med. Sci.*, vol. vi, p. 56) recommends very highly the use of the iodide of arsenic as a constitutional remedy. It must be given in doses carefully graduated to the age of the patient (one-eighth of a grain being the proper dose for an adult), and should any symptoms of its irritative action ensue, its use must be immediately suspended for a few days, and a purgative be administered.

The *local* treatment of favus is undoubtedly that upon which we must chiefly rely, since the essential element in the treatment must always be the destruction of the parasite.

The mere application of remedies adapted for this purpose, called parasiticides, is, however, rarely of itself sufficient, since they cannot penetrate to the hair-follicles, and it is, therefore, directed by most authors of experience in the treatment of this disease, that the hairs must be removed from the affected parts before the application can be efficiently and successfully made. Before doing this, the crusts must be removed. Some recommend for this purpose poultices, but these are condemned by Wilson as clumsy, and by Lebert as causing the extension of the disease by the softened sporules which spread to the surrounding surfaces and propagate the disorder. This objection does not, however, appear valid, and their use is countenanced by many good authorities. Wilson recommends their removal by means of a local vapor-bath, applied through the medium of a caoutchouc cap, or, if this is not at hand, by laying a piece of folded lint, wetted in a solution of subcarbonate of soda or potash, upon the head, and covering it with an oiled silk or gum elastic cap, which should include the entire scalp. M. Lebert insists upon the necessity of removing the favi (not the pustular crusts which accompany the specific vegetable growth) in their dry state, by means of small spatulas, needles, or some kind of instrument. The epidermis is readily detached from around the favus, and this latter, which adheres but slightly to the skin, is then easily removed. M. Lebert states that this is so easily done, that he has been able to teach his ward-attendants to remove them without pain to the patients. Hebra uses applications of alcohol, which cause the crusts to shrink and thus lose their attachments, when they are readily removed.

After the crusts have been gotten rid of, the scalp should be well washed with soap and water in order to remove any favus sporules that may have escaped and become free, and the hair should then be cut short. Various applications are then recommended, before proceeding to depilation, as

tending to allay the irritability of the scalp and to render the hair less friable; among these are oil of cade (Bazin) and almond-oil (Anderson), which may be applied for a few days before depilation is begun. There are various methods which have been adopted for the extraction, but the best is undoubtedly to employ a small pair of forceps with square ends, and fine but not sharp teeth, so as to enable the operator to catch the delicate and brittle hairs surely without breaking them. The hairs must be extracted singly, and so soon as a little space has been cleaned, the parasiticide remedy should be applied so as to secure its entrance to the follicle. A single depilation is frequently not sufficient, but it is easy to distinguish, by the appearance of the surface and the growing hairs, those parts where the disease has been eradicated. This process is at first somewhat tedious both to operator and patient, but by practice a degree of skill is acquired which enables the physician or trained nurse to remove the hair rapidly and with very little discomfort to the patient.

So soon as a clean surface has been thus obtained, some application intended to destroy the vitality of the vegetable growth ought to be made use of. One of the best for this purpose is a solution of corrosive sublimate, the strength of which, according to Lebert, ought to be, when employed in lotion, from two to four grains to the ounce, and, when used as a fomentation, weaker. This is also McCall Anderson's favorite application. Dr. Bennett (*Ranking's Half-Yearly Abstract*, No. xii, 1850, Am. ed., p. 73), employs, to fulfil this indication, cod-liver oil. The head is kept constantly smeared with the oil, and covered with an oiled silk cap. This application is, however, merely palliative, and, so soon as it is intermitted, the disease reappears.

There are various other remedies that have been applied to the diseased scalp empirically, either to "modify the state of the skin," to "excite the disordered follicles to healthy action," or, lastly, to "destroy the vitality of the fungus, and, by altering the nature of the soil on which it flourished, to prevent its reproduction." Without attempting to define the mode in which any of these various substances may produce their effect, we deem it best to mention as succinctly as possible those which have the strongest testimony in their favor.

Mr. E. Wilson, who it will be remembered does not believe in its parasitic nature, is less favorable to strong applications than he was formerly. Those he now prefers are the ceratum tigllii, containing from ten to thirty drops of the oil to the ounce; the unguentum hydrargyri nitratis, diluted one-half; the unguentum hydrargyri nitrico-oxidi, diluted in similar proportion; the compound sulphur ointment, and some others.

Dr. Bennett's application of cod-liver oil has been referred to above. This, in connection with the constitutional treatment for scrofula, is said to have cured, on an average, in six weeks.

MM. Cazenave and Schedel recommend alkaline and sulphurous applications, and acidulated lotions. They speak very favorably of, and give much the highest place, amongst the substances to be used in friction, to the iodide of sulphur. This remedy was originally made use of by Bielt, and employed by him with much success. Its efficacy is attested also by

Lebert. It is used in the form of an ointment, consisting of from a scruple to half a drachm of the drug to an ounce of lard, which is to be applied morning and evening to the diseased surfaces by gentle friction.

Applications of hyposulphite of soda, in proportion of \mathfrak{zj} to $\mathfrak{f}\mathfrak{zj}$ of water, or of sulphurous acid lotions, are highly recommended. Among the parasitocides most valued in France, are oil of cade and turpeth mineral, which latter may be employed in the proportion of \mathfrak{zj} to $\mathfrak{f}\mathfrak{zj}$ of glycerin of starch, which is perhaps the best excipient for the various parasitocides.

Ointments and lotions containing carbolic acid have been much employed of late, but apparently not with entire success.

Dr. Fuller recommends the ablution of the head twice a day by means of soft soap, and the inunction of an application composed of equal parts of unguentum hydrargyri ammonio-chloridi and unguentum picis liquidæ. He states that a cure may usually be effected by this plan in from two to four weeks.

When the disease affects other parts of the body, the treatment must be similar to that above recommended; depilation is, however, unnecessary, and a cure is usually more promptly obtained. When the nails are invaded, they should be cut and scraped, and the parasiticide application should be rubbed into and beneath the free border of the nail.

Under any plan of treatment, a complete cure is rarely obtained in less than from four to eighteen weeks; the disease is extremely obstinate, and there is a strong tendency to the redevelopment of the parasite after the cessation of the local treatment, until it be completely eradicated. By persevering in the plan above recommended, however, this can invariably be effected, and a perfect cure obtained, with the exception of patches of baldness, which but too frequently follow, from the destruction of the hair-follicles.

ARTICLE II.

TINEA.

WE have already, in our general remarks introductory to this chapter, stated our belief that the various forms of tinea or ringworm are contagious diseases, and due to the presence of a peculiar fungus, the trichophyton.

The ordinary varieties of tinea which are described, are tinea tonsurans, or ringworm of the scalp; tinea circinata, or ringworm of the general surface; and tinea sycosis, or ringworm of the beard. With the latter form, of course, we are not at present concerned, nor are its relations to the two other varieties indisputable, since opinions are still divided as to its contagious and parasitic nature.

There is, however, abundant reason for believing the essential identity

of *tinea tonsurans* and *tinea circinata*. In addition to the results of microscopic examination, which reveal the presence of the same fungus in both, there is the strongest clinical testimony to the same effect. Thus it constantly happens that patches of the two varieties will be observed upon the same patient, and there are innumerable instances on record to prove that they give rise to each other.

These diseases were formerly described by some authors under the generic name of *porrigo*; by others under that of *herpes*. Wilson, in his last edition, employs the term *trichinosis* to designate the group.

CAUSES.—The peculiar parasite, the *trichophyton*, is the essential cause of the disease; and the mode of its propagation is chiefly by contagion. Mr. Wilson believes the cause of the disease to be imperfect nutrition; but it is quite certain that the only way in which a scrofulous or debilitated constitution can influence the production of the disease, is by favoring the more ready growth of the parasite. In like manner, filthiness of every kind may be said to be a predisposing cause.

The influence of these is, however, trifling, and we have frequently met with the disease among families living in easy or very affluent circumstances, the children of which were perfectly well lodged, well clothed, and well fed, and to whom every attention required by the nicest cleanliness was given. The means by which the affection is communicated are such as brushes, combs, caps, etc., or by the direct contact of the diseased surfaces.

One of us has but lately had an opportunity of studying, on a large scale, these affections and the mode of their transmission, at a large Children's Home in this city. There were a considerable number of children, about twenty in all, affected with the disease in a severe form; by strict isolation, by the utmost care in preventing any use of their combs, brushes, caps, or clothing, by the other children, by covering the entire scalp with an oiled-silk cap, whenever they mingled with their comrades, the disease was prevented from spreading. It was, however, frequently observed, that in the children who suffered with *tinea tonsurans* of the scalp, patches of *tinea circinata* would appear either on the neck or face, or on some part which could be brought in contact with the affected surface; and its highly contagious nature was unhesitatingly believed by all the attendants, who had indeed themselves furnished the strongest evidence possible of it, by each and all contracting the disease repeatedly from handling the children in dressing them, or in making applications to the affected parts.

Age exercises a marked influence upon the production of these diseases, *tinea tonsurans* being confined to childhood and early youth, most commonly occurring between the ages of three and twelve years; though *tinea circinata* may be met with at any age.

TINEA TONSURANS.—SYMPTOMS.—The disease most frequently begins with little erythematous patches, which soon become covered with furfureous scales, and which increase circumferentially while they heal in the centre, leaving the skin more or less furfuraceous. Occasionally there may be a crop of minute vesicles on the patch, which are soon followed

by desquamation. When fully established, the disease appears in the form of furfuraceous patches of oval or circular shape, which are at first not more than $\frac{1}{8}$ th or $\frac{1}{4}$ th of an inch in size, but which increase gradually until they attain a diameter of one or two inches, and seldom more. The diseased surface is slightly thickened, elevated, of a grayish, bluish, or slate color, and covered with fine dry scales, which are very easily rubbed off, and are quickly renewed after being removed by any cause.

The hairs are altered from the very first. In the early stage, the apertures of the follicles of the diseased hairs are generally more or less prominent or papillated, and the hairs are unnaturally brittle, dull, and dry, and are bent on themselves and twisted, so as not to lie smooth, and the roots are somewhat matted together by the furfuraceous scales. A little later they break off at a short distance from the diseased surface, leaving the circular patches partially deprived of hair. The broken hairs are uneven in length, and otherwise altered in appearance, being bent and twisted, and having become lighter in color than the original hairs, so as to assume somewhat the look of bundles of tow. The enlarged follicles also dot the surface, giving it the appearance of cutis anserina, or the skin of a plucked fowl. The epidermis and the stumps of the broken hairs now become covered with a characteristic grayish-white powder, consisting of the sporules of the trichophyton, the peculiar parasite; and, on examining the hairs, the same fungus will be found penetrating into the bulbs and shafts between the separated fibres, and causing here and there, by its accumulation, swellings or bulgings of the shaft.

The disease is unattended by any local sensations, excepting a moderate degree of itching.

If the disease persists and the degree of inflammation increases, there may be a good deal of infiltration of the scalp, and the surface becomes tumid, and dotted with enlarged orifices of hair-follicles, or there may be an eruption of vesicles or pustules, which dry and form scaly, yellowish crusts.

DIAGNOSIS.—This disease is easily distinguished from other eruptions of the scalp. The appearances it presents when fully developed, are utterly unlike those of favus or eczema impetiginodes capitis. In favus, the peculiar cup-shaped crusts and the presence of the spores of the achorion, are sufficient to prevent mistakes; while in eczema, the eruption is sero-pustular, with the formation of yellowish or brownish yellow crusts; the patches are not circular, the hairs are healthy, the itching is extreme, and finally the disease is not contagious; in all of which particulars it differs entirely from the eruption of ringworm.

Pityriasis capitis does not occur in circular patches, but affects the whole scalp; it is not parasitic nor contagious, and does not lead to so much alteration of the hairs.

Occasionally tinea tonsurans, either from the irritation of scratching, or some other cause, may be associated with eczema impetiginodes, which to a great extent obscures the former disease, though a careful search will usually detect some of the characteristic broken stumps of hairs, loaded with the parasitic growth.

PROGNOSIS.—Ringworm of the scalp is entirely devoid of danger, but is an exceedingly troublesome disease, as it is apt to spread to other children, and is often very difficult to cure. Its duration is very indefinite, and it not rarely results in patches of permanent baldness.

TINEA CIRCINATA, as we have already said, frequently occurs in connection with *tinea tonsurans*, appearing on the neck or face; though it occurs also as an independent disease on any part of the body, and in patients of every age.

It begins as a little rose-colored, slightly elevated spot, which soon becomes the seat of a slight furfuraceous desquamation; and extends circumferentially, healing in the centre, until it forms a large slightly elevated erythematous ring, inclosing a portion of sound skin.

In other cases, minute vesicles form on the reddened inflamed ring. They follow the usual course of development, being at first transparent, then turbid, and finally drying into small thin scales.

The size of the patch varies greatly, being in some instances small, not larger than a shilling, and in others presenting a diameter of two or three inches. When small, the redness covers the whole of the patch, but is much fainter in the centre than at the circumference; when large, the centre regains the natural color of the skin. Usually the ring is exactly circular, but at times it assumes an oval shape.

If any hairs have been growing on the affected spot, they become brittle and changed, as before described. There are usually several such circles present, and in some cases they are formed in great numbers. The only symptoms accompanying the eruption are slight pricking, smarting, and itching in the part.

Occasionally the parasitic growth invades the nails, which then become opaque, whitish, thickened, and brittle.

DIAGNOSIS.—There are but few diseases with which there is any danger of confounding *tinea circinata*. It is distinguished from *erythema marginatum* by the greater elevation of the marginal ring, by the presence of the parasite, and by its contagious nature; and the last two peculiarities serve to distinguish it from *psoriasis circinata*.

According to McCall Anderson, and some other dermatologists, *herpes iris* is merely a form of this affection.

TREATMENT.—The cases of *tinea tonsurans* that have come under our charge, have proved in many instances very rebellious to treatment.

Strict attention should always be paid to cleanliness and hygienic rules; and, if the disease be associated with any impairment of the constitution, cod-liver oil, iron, in the form of the syrup of the iodide in syrup of sarsaparilla, arsenic, and bitter tonics, should be administered.

The local treatment is, however, the most essential. Where the disease occurs on a part covered with hair, depilation is advised by some authorities, and it would in all probability facilitate and hasten the cure.

Among the local applications which have proved most useful to us have been sulpho-alkaline lotions, composed of \mathfrak{zj} of subcarbonate of potash and \mathfrak{zij} of sulphur, to a pint of water, applied by washing with a sponge several times a day; strong solutions of sulphite of soda; and an ointment

consisting of \mathfrak{zj} of muriate of ammonia, mixed in an ounce of sulphur ointment, applied first at night by inunction, and after a time on rags.

Alkaline remedies have also been much used by other observers, who recommend washing the scalp every morning with a lotion composed of gr. xxx or xl of carbonate of potassa or borax to a pint of water, and applying in the evening an ointment containing \mathfrak{ij} of tannic acid to \mathfrak{zj} of lard.

Much more stimulating applications are, however, highly recommended, and often prove very serviceable. Thus, Mr. Wilson advises a single application of the acetum cantharidis, or the stronger acetic acid; and Devergie recommends a solution of nitrate of silver, \mathfrak{zj} to $\mathfrak{f}\mathfrak{zj}$ of water.

Various mercurial applications are also advised, as solutions of corrosive sublimate, the citrine ointment, or the following, recommended by Jenner:

R. Hydrargyri Ammonio-Chloridi,	.	.	.	gr. xx.
Ung. Sulphuris,	.	.	.	\mathfrak{z} iv.

Tarry applications may also be employed in obstinate cases, in the form of lotions, ointments, or soaps, containing tar, or oil of cade.

Nayler speaks highly of a plan used by Mr. Coster, who saturates the part with the following mixture:

R. Iodinii,	.	.	.	\mathfrak{z} ij.
Ol. Picis,	.	.	.	$\mathfrak{f}\mathfrak{z}$ j.

This solution is to be rubbed in firmly with a piece of sponge on the end of a piece of wood or whalebone. It is allowed to dry on the part, and left until the cuticle and the black crust separate at the end of a week or ten days.

In cases where many patches are present over the body, it is advisable to employ mercurial or sulphur vapor-baths.

It must not, however, be forgotten that these varieties of tinea are among the most obstinate disorders to which children are subject. The most faithful trial may be made with the remedies recommended above, for a long time, without success, and it is often necessary to persevere in their use for months; conjoining the treatment with a change of diet, and, when possible, with a change of residence, before the affection will be entirely and permanently cured.

CASES.—The following cases may be taken as types of the aggravated form of tinea, after it has persisted a long time and become complicated with secondary eruptions of eczema or pityriasis. It will be noticed that in the following records, all of the patients are stated to have been markedly scrofulous; but this circumstance must not have too much importance attached to it, since in the Home where these cases occurred, almost every one of the children presented unquestionable marks of the strumous diathesis. There can be no doubt, however, that this condition of constitution strongly favored the development of the disease, rendered it more severe and obstinate, and also favored the occurrence of the secondary inflammatory eruptions.

CASE I.—George T., æt. five years, scrofulous, admitted to the Home in 1864, with bleeding piles. *Tinea tonsurans* appeared two months after admission, and persisted with various fluctuations for eighteen months, when it became complicated with eczema impetiginodes. Applications of tar and corrosive sublimate had been chiefly relied on.

November 30th, 1866. Scalp covered with grayish-yellow crusts, one-fourth inch thick, in places running together or forming isolated lumps. A few spots of *tinea circinata* on face and neck. The scalp is reddish, and there is very little discharge from it. The hairs are sparse and broken. The cervical glands are much enlarged on both sides. On removing the crusts and examining the base, numerous exudation corpuscles and some spores of *trichophyton* were found; the epithelium not very granular.

Poulticed to remove the crusts. Ordered iodide of iron and potassium internally.

December 4th. Scalp quite clean from crusts, but remains reddish, with here and there bald patches. Numerous spores of *trichophyton* found in the hairs and among the epidermic cells.

Solution of sodæ sulphas ($\frac{3}{j}$ to Oss water), applied morning and evening, and kept on during the whole time by means of folds of linen saturated in the solution, and covered with an oil-silk cap.

December 16th. Much improved. Scalp cleaner, and less red. Some flat, thin, whitish scales over surface. Hairs more free from *trichophyton*, but numerous spores can still be seen by scraping moist surface beneath the thin crusts.

Treatment continued, with ultimate success.

CASE II.—William L., æt. 5 years; hereditary tendency to tuberculosis; scrofulous; cervical glands enlarged on both sides; admitted in 1865, and has had *tinea* ever since, many forms of treatment having been tried, but none with more than temporary success.

November 30th, 1866. The scalp is reddened and wax-like from infiltration, with patches of baldness. In places where the eruption is oldest it is covered with whitish scales; elsewhere, there are scattered or confluent grayish-yellow or yellow crusts. Discharge of pale, thin, fetid pus.

On examining the surface beneath the crusts, numerous pus-cells and spores of *trichophyton*, often aggregated together, are found. The hairs have lost their normal appearance entirely, are bent, and where they emerge from the scalp, the shaft is swollen, with bulging outline. The shafts are covered with spores of *trichophyton*, and their longitudinal fibres separated by collections of the fungus. Some of the bulbs remain healthy, others are broken and apparently converted into masses of fungous spores.

Ordered poultices to remove crusts; iodide of iron and potassium internally.

December 4th. Scalp clean, with exception of minute white scales; shows bald glazed patches, with light, short, thin hairs. Ordered same application of sulphite of soda as used in previous case.

December 15th. Greatly improved; the large bald patches still covered with minute shining white scales, but few *trichophyta* to be seen.

CASE III.—Charles L., æt. 4 years, admitted in May, 1866; hereditary tendency to tuberculosis; cervical glands slightly enlarged. *Tinea* soon appeared on the face and scalp, and in early part of November, thin flat grayish-yellow crusts formed over vertex, the rest of the scalp being covered with minute whitish scales.

November 30th, 1866. Ordered poultices to remove crusts; iodide of iron and potassium internally.

December 4th. Scalp comparatively clean. Patches of baldness, especially over parietal protuberances, with straggling, short, light-colored hairs. Abundant spores of *trichophyton* found. The hair-shafts much involved, collections of the parasite existing between the longitudinal fibres. The bulbs are also diseased, and seem to have become affected just below the exit of the hair; the bulb first becoming swollen at this point and then its sheath having become destroyed, so that the fungus forms a bed surrounding the shaft.

Ordered same application of solution of sulphite of soda.

December 16th. Very much improved; scalp smooth and clean, excepting above

the ears, where there is on each side a collection of thin yellowish crusts. The hairs passing through these had numerous pus-cells adherent to their shafts, but the hair-bulbs seemed healthy, and no spores of trichophyton could be found on any of them.

ARTICLE III.

ALOPECIA AREATA.

THIS affection, which is also known by the names of *area* and *tinea decalvans*, is characterized by the loss of hair in circumscribed patches of round or oval shape. It is by no means a rare disease, and is much more common in children than in adults; thus of 42 cases cited by Hutchinson, 28 were under fifteen, 14 above that age.

CAUSE.—There is still much doubt as to the essential cause and nature of alopecia areata, although it appears to us that the view which attributes it to a perversion or suspension of innervation is the correct one.

Gruby is said to have discovered in 1843 a fungus in it, which has been called the *microsporon audouini*, and some dermatologists of high authority accept the view of its parasitic nature. It cannot, however, be said to be demonstrated, since the parasite is very rarely found; so that Anderson, who has made numerous microscopic examinations, has never succeeded in detecting it.

Wilson, Dubring, and many others, consider it as due to suspended innervation, as a kind of paresis.

The disease appears to be, at least in some instances, propagated by contagion; though it certainly possesses this property to a much less degree than either of the forms of ringworm.

SYMPTOMS.—The disease is limited to the scalp in children; though in adults it may attack any hairy part. In some cases, the first intimation of the existence of the disease is the sudden discovery of a bald spot, but in others, though less frequently, there is slight itching, with redness and branny desquamation of the affected spots.

The bulbs of the hairs then atrophy, and become tapering instead of being rounded and club-shaped; the hairs themselves become dry, lustreless, and brittle, with a fibrous fracture, and rapidly fall out, leaving bald patches.

These patches vary in size from one-half inch to an inch or even more in diameter, and there may be but a single one present, or they may be numerous, in which case they often coalesce, forming large patches of irregular shape; when the patches are single they usually assume a round or oval form.

The denuded portion of scalp is peculiar in appearance, being very white and polished, and thinner than the surrounding healthy scalp; the sensibility of the affected surface is also frequently impaired.

DIAGNOSIS.—There can be no difficulty in recognizing the fully devel-

oped disease, excepting in the comparatively rare cases when it is combined with other skin diseases, as eczema or pityriasis.

PROGNOSIS.—The only danger attendant upon alopecia areata is that of deformity, which is, in some cases, very great, depending of course upon the extent of the disease and the stage at which it is brought under treatment.

If the patches are small, the scalp not materially atrophied, and the orifices of the hair-follicles still visible on the bald patches, there is good reason to hope that steady persistence in treatment will effect a cure. In the majority of instances, it may be said that recovery occurs after a length of time varying from several weeks to several months.

TREATMENT.—Those who regard this as a parasitic affection, advise the removal of the hairs immediately surrounding the patch, and the application of some of the stimulating parasiticides recommended in the article on tinea.

The majority of authors, however, content themselves with the application merely of such stimulating lotions and ointments as will increase the nutrition of the affected spots, and favor the renewed growth of hair.

Among the ointments which are most highly recommended are those containing the red iodide, the nitrate, the ammonio-chloride of mercury; some form of sulphur; or tar, iodine, or cantharides.

Hillier recommends, as the treatment he has found most useful, the application at long intervals of acetum cantharidis to the bald patches; painting them every other day with tincture of iodine, washing the head twice a week with soap and cold water, and applying a wash (consisting of one pint of rum, one ounce of tinct. cantharidis, one-half ounce of spt. ammoniæ aromat., and ten ounces of water) to the parts of the head which are not bald, twice a week.

The effect of this local treatment will be much increased by the internal administration of arsenic and iron.

Alcohol, carbolic acid, ammonia, and capsicum have also all been highly recommended as local applications in the treatment of this affection; they should of course be employed in a sufficiently dilute form.

ARTICLE IV.

SCABIES.

DEFINITIONS; SYNONYMS; FREQUENCY.—Scabies is a contagious affection of the skin, characterized by the formation of papules, vesicles, or pustules; the vesicles being pointed, generally discrete, and usually presenting small red lines, of one or several lines in length, running off from them. The eruption is attended with severe itching, and is caused partly by the presence in the skin of a small insect, called the *acarus scabiei*, and partly by the scratching which the intolerable itching provokes.

CAUSES.—Itch is a contagious malady, and is in all probability caused only by contact, either immediately with some person laboring under the disease, or with articles of clothing worn by an infected individual.

It is much more frequently met with amongst the poor and destitute, whose habits are uncleanly, who live close packed together in small and inconvenient houses, and in whom, therefore, the means of communication are more abundant than amongst the easy classes of society, whose habits, and, consequently, liability to contact, are the opposite of those just named. It is, however, comparatively rare in the United States, and particularly in this city; thus Duhring met with but 12 cases out of 2472 consecutive cases of skin diseases.

The disease usually appears in children in from four to five days after exposure to the contagion. In healthy, sanguine children, it often shows itself within a shorter time—after two days—while in those who are feeble and weakly, the period of incubation may be even longer than four or five days.

SYMPTOMS.—The first symptoms of itch appear in the part to which the cause, a contagious contact, may have been applied. In infants at the breast, it is usually first developed on the hips and thighs, as it is those parts which are most constantly in contact with the nurses who carry the child, and from whom young children generally receive the infection. In older children, the disease commonly appears first on the wrists and between the fingers, and extends thence more or less quickly to the flexures of the elbows, and to the axillæ and abdomen. It rarely or never attacks the face in adults; but in children, even this part is not, according to M. Richard, exempt. (*Trait. Prac. des Mal. des Enfants*, p. 590.)

The disease is always attended with severe itching, which, in infants, causes uneasiness and fretfulness, and, in older children, violent scratching. The itching is increased by the heat of the bed-coverings, and is, therefore, most troublesome at night. The eruption appears in the form of more or less numerous vesicles, which are small, discrete, acuminate, and transparent at the top. The vesicles are at first of a faint rose color, and they contain a viscid transparent serum. Their number is variable, being sometimes very abundant, and at others sparse. They either open spontaneously, or are soon broken by the fingers or clothes, and are followed by small, thin, slightly adherent scabs. In some instances the action of the nails causes slight effusions of blood, which dry into small bloody scabs, like those of prurigo, thus embarrassing to a certain extent the diagnosis of the disease. Sometimes, particularly when the inflammation attendant upon the eruption, or that caused by scratching, is marked, there are, intermingled with the psoric vesicles, pustules of impetigo, or perhaps papules of lichen, which tend, like the sanguine crusts just alluded to, to render the diagnosis difficult. Indeed, it is more strictly correct to describe the eruption of scabies as multiform instead of vesicular, as was formerly done.

When a recent vesicle is carefully examined, there may generally be observed running off from it, in a straight, curved, or zigzag direction, a whitish or reddish line, like that produced by the scratch of a pin. This line marks the course of the fecundated female of the *acarus scabiei* in its

burrowings under the epidermis, and is called the *cuniculus*, or burrow. It varies in length from one or two, to five or six lines. At the point where it terminates opposite to the vesicle, there is usually to be seen a small rounded projection, deeper in color than the rest of the *cuniculus*, beneath which lies the insect. The *acarus* can often be found at this spot, and removed, by carefully introducing horizontally under the epidermis the point of a small needle, and by manipulating so as to take off a small layer of the epidermis. The insect clings to the point of the needle, and can then be extracted from its lodgment. This furrow is the certain diagnostic mark of scabies.

The number and extent of the vesicles vary greatly in different subjects. In some they are confined to limited surfaces, while in others, and particularly in robust, sanguine children, and in those who are neglected and imperfectly cleansed, they extend to many different parts, or over the greater part of the body.

Itch occasions in children much irritability and suffering, and when neglected may injure seriously the general health, and cause emaciation and debility.

The *acarus scabiei* is an arachnoid insect, varying, according to Mr. Wilson's measurements, between $\frac{1}{17}$ and $\frac{1}{7}$ of an inch in length, and between $\frac{1}{33}$ and $\frac{1}{9}$ of an inch in breadth. It is of a whitish and shining color, when examined with the naked eye, of a globular form, and is provided with eight legs, four anterior and four posterior. A most accurate and minute account of the structure of the insect is given by Mr. Wilson in his work on diseases of the skin (7th Amer. ed., p. 739). Besides the female, which is formed as before stated, at the extremity, the *cuniculus* contains a varying number of ova, rarely more than twelve or fourteen; there are in addition numerous little oval or round blackish spots, which are supposed to be excrement. These ova are about $\frac{1}{30}$ of an inch broad, and $\frac{1}{10}$ of an inch in length, though their size varies according to their age. After the escape of the *acarus* the shell appears shriveled, with two slits in it.

DIAGNOSIS.—The most characteristic marks of itch are the presence of the *cuniculi* and of the insect which causes the disease. If the *acarus* or its ova can be extracted from the skin, there will remain, of course, no doubt; and if the *cuniculi* be distinct and numerous, the diagnosis becomes almost as certain as when the insect itself is obtained. Before endeavoring to detect the *cuniculi*, it is always advisable to make the patient wash the part thoroughly.

In doubtful cases, it has been recommended by Gull and Hilton Fagge to search for the ova in the crusts or the thickened and undermined cuticle in the neighborhood of the vesicles. In order to detect these, a small piece of the crust should be boiled in a solution of caustic soda, 3ss to fʒj of water, until it is in great part dissolved; the fluid should then be allowed to settle, the supernatant part decanted and the deposit examined, which will, in cases of true scabies, be generally found to contain larvæ, ova, or egg-shells.

When, on the contrary, the insect cannot be found, and when the cuniculi are absent or not distinct, the diagnosis becomes more uncertain. The diseases with which it is most likely to be confounded are eczema simplex, prurigo, and lichen simplex. From the former it may usually be distinguished with certainty by attention to the following points: in eczema the vesicles are flattened, or globular, scarcely raised above the surface, and they are collected together in clusters; in itch they are acuminate, elevated, and either entirely distinct, or much less confluent than in eczema; in eczema there is a sensation rather of pricking than itching, whilst in itch the sense of itching is severe and distressing; and lastly, itch is communicable by contact, whilst eczema is never contagious.

Prurigo begins with papules, which always remain such. The scabs in prurigo are small and black, consisting of coagulated blood, caused to exude by the rubbing off of the top of the papule; while in scabies the scabs are more like thin, yellowish, and friable scales. The seat of the two eruptions is different. Prurigo is developed upon the back, the shoulders, and upon the extensor surfaces of the limb; while itch appears first about the thighs and buttocks, between the fingers, or about the flexures of the joints. Lastly, prurigo is never, itch always, contagious.

Lichen simplex is a papular disease, in which the papules are closely agglomerated, while in scabies the papules, if present, are conjoined with vesicles or pustules, and are discrete. Lichen sometimes affects the hands, and might then be mistaken for itch; but in the former the eruption affects the dorsal surface of the hands, while in the latter it appears in the interspaces of the fingers. Lichen is never attended, as itch always is, by severe pruritus. Attention to these points of difference will almost always render the diagnosis of the two diseases very easy and certain.

When, as often happens, scabies is intermingled with other eruptions of the pustular, papular, or vesicular kind, the diagnosis can be arrived at with certainty, only by careful attention to the cuniculi, or by the detection of the insect. When neither of these characteristic conditions are present to mark the true nature of the disease, there will always remain some doubt as to the diagnosis.

Under these circumstances, however, it is advisable to treat the case as one of scabies, since the specific remedies for this affection will not be injurious, even if they do not speedily cure the eruption.

PROGNOSIS.—Itch is a mild disease, which never disturbs the health seriously.

TREATMENT.—If the inflammation produced by scratching be very severe, it may be necessary to allay it by emollient applications, though this rarely happens.

In children, as in adults, the best treatment of itch is the use of sulphur by inunction. The ungt. sulphuris of the U. S. Pharmacopœia, consisting of one part of sulphur to two of lard, should be well rubbed into the skin before a fire, morning and evening, for two days. The child should be kept in a flannel gown, and in bed, during this treatment. On

the morning of the third day, the skin may be washed clean with soap and water, or by immersion in a warm bath. This plan rarely fails to effect a cure. Should it happen, however, to fail, the treatment must be repeated. Before the application of this or any of the other ointments, the surface should be well scrubbed with soap and hot water, so as to cleanse and soften the skin.

It also increases the effect of the sulphur, to conjoin with it some alkaline substance, as in the various sulpho-alkaline ointments and lotions, of which the following are among the best:

UNG. SULPHURIS CUM POTASSA (WILSON).

R. Sulphuris Sublimata,	℥j.
Potassæ Carbonatis,	℥ij.
Unguenti Benzoati,	℥v.
Olei Anthemidis,	f℥ss.—M.

Tilbury Fox's Formula.

R. Sulphuris Sublimati,	℥ss.
Hydrargyri Ammonio-chloridi,	gr. iv.
Creasoti,	gtt. iv.
Ol. Anthemidis,	gtt. x.
Adipis,	℥j.—M.

Helmerich's Formula.

R. Sulphuris Sublimati,	℥ij.
Potassæ Carbonatis,	℥j.
Adipis Preparati,	℥ viij.—M.

Vlemingxz's Formula.

R. Calcis Vivi,	℥ij.
Sulphuris Sublimati,	℥iv.
Aquæ Fontanæ,	f℥xx.

Boil in an iron vessel, and stir with a wooden spatula to a perfect union.

These are all quoted in the proportions directed for adults, which are much too active to be applied to the delicate skin of children; they should therefore be diluted one-half at least.

Anderson recommends the use of oil of cade or tar, combined with the sulpho-alkaline ointments.

As the use of the sulphur ointment is sometimes objected to in private families, an account of its disagreeable odor, various substitutes have been recommended. Mr. Wilson states that he found camphor dissolved in oil, in the proportion of one drachm to the ounce, answer every purpose of eradicating the disease; and Dr. Coley (*Prac. Treat. on Dis. of Children*, Phil. ed., 101) speaks highly of an ointment composed of one drachm of iodide of potassium to one ounce and a half of lard, of which a little is to be applied all over the body, except the head and face, every night.

Ointments containing carbolic acid or petroleum are also used with good effect.

The use of stavesacre and hellebore has lately been revived, and apparently with good success; and Anderson highly recommends an ointment made by melting together one part of liquid styrax with two of lard.

The disease rarely requires any constitutional treatment. If, however, any complication exist, or the general health be deranged in any way, such measures as may be necessary for the removal of either of these conditions should be employed, in connection with those proper for the specific disease.

CLASS VIII.

WORMS IN THE ALIMENTARY CANAL.

GENERAL REMARKS.

THERE are five different species of worms found in the alimentary canal. These are the *Ascaris lumbricoides*, or round-worm; *Ascaris vermicularis*, thread-worm, seat-worm, or, as it is popularly called, ascarides; *Tricocephalus dispar*, or long thread-worm; *Tænia solium*, and *Tænia mediocanellata*, the most common varieties of tape-worm; and the *Bothriocephalus latus*, *tænia lata*, or broad tape-worm.

We shall give a short description of each of the intestinal entozoa, in order that they may be readily distinguished, but will treat of the causes, symptoms, and treatment only of the first two, inasmuch as the *tænia*s very rarely exist during infancy or childhood, and the *tricocephalus* is much less frequent than the round and seat-worms, and gives rise to symptoms of the same kind as the former.

DESCRIPTION.—The *Ascaris lumbricoides*, or, as it is commonly called, *lumbricoides*, *lumbricus*, or round-worm, is shaped not unlike the common earth-worm, having a cylindrical body, which is attenuated towards either extremity, but particularly the anterior. It varies in length generally between six and twelve inches, and is usually about two or three lines in thickness. The young worm, about an inch and a half long, is rarely met with. The head of the animal is at the smallest extremity, and may be distinguished by a circular depression, around which may be seen three tubercles. When recently voided, the worms are somewhat transparent, so that the viscera may sometimes be seen through the parietes. The integument is marked by circular fibres, and by four lines extending at equal distances from the head to the tail, the former of which indicate the course of the muscles, while the latter indicate that of the vessels and nerves.

The color of the worm is whitish, yellowish, or more or less deep rosy in tint, according to the nature of the aliment they contain; they are, as already stated, somewhat transparent when first voided. The alimentary canal, which may be distinguished by its brownish color, terminates by a transverse opening or anus, situated on the inferior surface of the animal, just in front of its posterior extremity.

The two sexes are in different individuals. The male may be known by its tail, which is shortly curved, while that of the female is straighter and thicker. The genitals of the male consist of a double penis, which may sometimes be seen to protrude just in front of the caudal extremity; those

of the female may be distinguished by the vulva, seated at a constricted point of the body, about a third of the distance from the head to the tail. The male is smaller and much less abundant than the female.

The *Ascaris* or *Oxyuris vermicularis*, thread-worm, seat-worm, or maw-worm, is the smallest of the intestinal worms, and is generally distinguished in popular language by the title of ascarides. The sexes are in separate individuals.

The male is generally about two lines in length; its body is elastic, of a whitish color, very slender, and looks not unlike a piece of cotton thread, whence one of its names was derived. The female is larger than the male, reaching a length of four or five lines. The anterior part of the body is of the same shape in both sexes. It is obtuse, and surrounded by a transparent membrane, through which may be seen a straight tube, forming a kind of bladder, which is the œsophagus, and which terminates in a globular stomach. The head is provided with three tubercles, as in the lumbricoides. The intestinal tube in the male continues the whole length of the body, which becomes somewhat thicker towards the end, and is arranged into a spiral shape at the tail. The body of the female is shaped like that of the male as far back as the stomach, and increases in size in the first third of its length, after which it diminishes, and becomes so small at the end as to be seen with difficulty by the naked eye.

The *Tricocephalus dispar*, or long thread-worm, is generally about an inch and a half or two inches long, and consists, as it were, of two portions, of which the anterior, constituting about two-thirds of the length, is exceedingly slender, scarcely thicker than a horse-hair, while the posterior third swells out suddenly so as to become much thicker and larger. The sexes are in different individuals. The worm is provided with an alimentary canal, which, commencing at an orbicular mouth placed in the small extremity, runs through the animal to the anus, placed at the caudal extremity. The male is smaller than the female, and is usually found convoluted. This worm is met with chiefly in the cœcum and colon, particularly the former. It usually exists in very small numbers, and sometimes but a single one is found. The symptoms which it occasions are the same as those produced by the lumbricoides.

The *Tænia solium*, common or long tape-worm, as well as the *Tænia lata*, are of rare occurrence in children. Of 206 cases observed by M. Wavrouch, only 22 occurred in subjects under fifteen years of age, and of them the youngest was three years and a half old (*Bib. du Méd. Prat.*, t. v., p. 626). These worms have, however, been met with at an earlier age, but as they are rare, we deem it unnecessary to do more than describe their appearance, in order that the reader may be able to distinguish between them and the varieties which generally exist in children, the *Ascaris lumbricoides* and *vermicularis*. For a full account of the symptoms produced by the two varieties of the *tænia*, and their treatment, the reader is referred to any of the standard works on the practice of medicine.

The *Tænia solium* is usually of a whitish color, flat in form, and varying in length from five to ten feet, its ordinary length, to twenty feet, or even

more. It is uneven in shape, being thick and broader behind, and measuring three or four lines at its widest part, while it tapers gradually towards the anterior extremity, where it becomes slender and thread-like. The head is globose and very minute, being about $\frac{1}{30}$ th of an inch in diameter. It has a projecting papilla in the centre, furnished with a double circle of hooklets. There are also four projecting suckorial disks placed at equal distances around the head. The neck is delicate and thread-like, but on microscopic examination presents transverse wrinkles at a short distance from the head, and soon merges into the distinctly jointed body. This is composed of numerous segments, which at first are small, and broader than they are long, but lower down increase more rapidly in length and in breadth. The largest joints measure about one-fourth of an inch wide by half an inch long. Each joint contains both male and female sexual apparatus, opening by a common aperture on the side.

The *Tænia mediocanellata* was formerly confounded with the *tænia solium*. It attains, however, a greater length, its joints are longer and broader, and its head is about three times as thick. The four suckers are present, but there is no central projecting papilla, nor any hooklets.

The *Bothriocephalus latus*, *Tænia lata*, or *broad tape-worm*, is long and flat like the preceding variety, but it is generally thinner and broader, measuring from four to ten lines in breadth. It attains even a greater length than the common tape-worm. It is usually of a dirty-white color, and rather less opaque than the *tænia solium*. It is distinguished also from the other *tænia*, by the shape of the segments, which are broader than they are long; by the form of the head, which is small, elongated, without spines, and divided into two lobes by a longitudinal fossa on each side; and by having, instead of the four mouths of the *tænia solium*, a single minute pore in the centre between the fossa, or else two pores, one at the extremity of each lobe.

The *frequency* of intestinal worms, and their *importance* as a cause of disease, have certainly been, and are still by many physicians, and especially by the public, very greatly exaggerated. There can be no doubt that they do, when they exist in large quantities, and particularly in certain countries, give rise to great disturbances of the digestive organs, and even occasion death; but such instances are, it seems to us, extremely rare, in this city, at least. We are quite sure that we have never as yet met with a case, in our own experience, in which life was at all seriously endangered by their existence,—though we have seen numerous instances in which slight disorders of the digestive apparatus, and various nervous symptoms, generally of very moderate severity, have disappeared after the administration of anthelmintics, sometimes followed, and in an equal number of cases probably, not followed, by the expulsion of worms.

To show the truth of the above remarks, as to the importance of worms as a cause of disease, we make the following quotations: Dr. Rush (*Med. Inquiries and Observations*, vol. 1, p. 205) remarks: "When we consider how universally worms are found in all young animals, and how frequently they exist in the human body, without producing disease of any kind, it

is natural to conclude that they serve some useful and necessary purposes in the animal economy." M. Guersant says (*Dict. de Méd.*, t. xxx, 669): "It has always been the custom to assign to entozoa much too important an influence upon the diseases of childhood. In proportion as this part of pathology is perfected, it becomes evident that the greater number of children dying after having discharged worms, or even while having them still, are affected with acute or chronic diseases, which leave after death incontestable traces of their effects, and which are of themselves necessarily fatal." M. Barrier (*Mal de l' Enf.*, t. ii, p. 100) quotes M. Trousseau as making the following remarks: "For sixteen years we have not met with a single child who has presented any verminous symptoms; never or almost never does a child born and reared in Paris discharge worms, while just the contrary is true as to the provinces. . . . Young children, to be sure, are sometimes met with in our hospitals, who discharge worms, but they are those who have been born in the country, and have lived in the capital only for a short time." Dr. Condie (*Dis. of Child.*, 2d ed., p. 226), remarks: "Worms are a very common occurrence in the intestines of children, and may unquestionably, under certain circumstances, become a cause of severe irritation; but much less frequently than is generally supposed."

We believe we may conclude, therefore, that though these parasites are of very common occurrence, and productive of grave disorders in some countries, they are rarely met with in quantities sufficient to do serious injury to the health, in other places, as for instance Paris, and probably in this country, or at least in the northern parts of it.

That intestinal worms do, however, not unfrequently in some countries, and occasionally in all, produce dangerous and even fatal disturbances of the health, cannot be doubted after careful perusal of the evidence brought forward by different authorities. M. Guersant, amongst others, remarks (*loc. cit.*, p. 670): "It is nevertheless incontestable, that the development of these animals in the gastro-intestinal and abdominal cavities does sometimes give rise to very varied morbid phenomena, which are in some instances grave enough to cause death." Nevertheless, we are disposed to believe, as stated above, that fatal or even dangerous results from the existence of these parasites are of rare occurrence in this city, and probably throughout our Northern States. Dr. Dewees, however, mentions several cases in which they produced alarming symptoms, and one in particular (*Dis. of Child.*, p. 492), in which the subject, a child twenty months old, was extremely emaciated, and whose abdomen was "enormously distended, and semi-transparent," who recovered rapidly after ninety-six lumbricoides, from six to ten inches long each, had been expelled under the use of pink-root in infusion.

ARTICLE I.

ASCARIS LUMBRICOIDES.

THE *description* of this worm has already been given at page 1003.

CAUSES.—Under this head we shall not pretend to consider the question of the origin of worms, but only the causes which predispose to their production or favor their growth.

Age has no doubt a considerable influence upon the predisposition to lumbricoides. According to M. Guersant (*loc. cit.*, p. 685), infants at the breast under sixth months of age are very rarely affected with them. Instances occasionally occur, but are altogether exceptions to the general rule. Above six months of age, they begin to be met with, but still very rarely, so that scarcely one or two will be found in several hundred children of a very early age; while from three to ten years of age they will be observed in about a twentieth, or in some seasons perhaps in a larger proportion. M. Valleix states that he has never met with them in newborn children. Dr. Dewees says (*loc. cit.*, p. 481), that he has never seen worms in children under ten months old; and in only two instances at that age. We do not recollect ourselves ever to have seen them in subjects younger than eighteen months, and very rarely in those under three or four years.

There can be little doubt that the disposition to worms is *hereditary* in some families. It is generally believed that the species under consideration is more common in *girls* than *boys*; that it is most common in children of *lymphatic* and *scrofulous* constitutions; and that a too exclusively vegetable or milk *diet*, and an abuse of *fruits*, strongly predispose to their production. The *habitation* of a cold and damp, or warm and damp climate, and the *seasons* of summer and autumn, are supposed by many also to favor their production and growth. It is a general belief, and we should suppose from personal experience, a well-founded one, that a feeble and disordered state of the digestive function from any cause, often acts as a predisposing cause of worms, and particularly of lumbricoides.

SEATS.—The small intestine is, in a very large majority of the cases, the seat of the ascaris lumbricoides. They are met with, however, in other parts of the digestive tube, particularly the stomach and large intestine, and more rarely in the œsophagus or pharynx. In some instances they are found to have migrated to other organs, as to the liver, gall-bladder, and in still rarer cases they have passed into the peritoneal cavity, bladder, larynx, trachea, bronchi, and even into the nasal passages and frontal sinuses. They have also been met with occasionally in the walls of the abdomen, forming verminous abscesses, whence they have escaped on the opening of the abscess.

The *number* of ascarides is exceedingly variable; there may be only two or three, ten or twenty, or several hundred. When very numerous, they are apt to be rolled or twisted into knots or balls, which have been seen as

large as the fist, so as to block up completely the canal of the intestine. In a case cited by Rilliet and Barthéz, from M. Daquin, the duodenum was so filled with worms as to be distended, and to have acquired a considerably larger size than natural, while at the same time it was hard and elastic. The jejunum, ileum, and cœcum were filled, so that it seemed as though the worms must have been pushed in by force. They were found also, but in smaller quantity, in the colon. Dr. Condie (*loc. cit.*, p. 230) states that he has known one hundred and twenty lumbricoides to be voided in a single day by a child five years old. It ought, however, to be remarked, that the instances in which such large numbers are met with are altogether exceptional, especially in our Northern States. We have never ourselves known more than six, eight, or ten to be expelled within a few days' time, and very generally there have not been more than three, four, or five.

ANATOMICAL LESIONS.—When the number of lumbricoides is small, the mucous membrane has been found in a state of perfect health, while, on the contrary, when they are numerous, and especially when collected together into knots, the membrane has presented a fine injection like that which exists in erythematous enteritis; in some very rare instances on record, in which the quantity of worms has been very great, the mucous membrane has been found deeply injected, thickened, granulated, and, in a small proportion of cases, softened, and even eroded. Not unfrequently the intestine presents all the characters of well-marked enteritis, or enterocolitis, though the number of worms may be very small. In such cases, it is reasonable to suppose that the inflammatory affection has been an accidental complication of the verminous disorder.

Much discussion has arisen in regard to the manner in which *perforation* of the intestine, as an accompaniment of worms, takes place. It is necessary to suppose, in subjects in whom worms are found in the peritoneal cavity, or in abscesses formed in the abdominal parieties, that perforation of the bowel has taken place, and yet in some instances no trace of the openings is left, no inflammation of the serous membrane is met with, nor has there been any escape of the contents of the digestive canal into the abdominal cavity. In others, however, and much the most numerous cases, it is evident from the anatomical appearances, that the perforation has taken place in consequence of previous ulceration of the coats of the bowel, and that the worms have escaped with the other contents of the intestine. It is in regard to the former class, therefore, that discussion has principally taken place; some asserting that the parasite itself makes the opening, by an active process, while others deny the possibility of this occurrence, and maintain a previous ulceration or softening in all cases. Amongst those who advocate the possibility of perforation independent of previous change in the intestinal coats by disease, are MM. Mondière and Charcelay, the former of whom has examined the subject with a great deal of care, quoted by Rilliet and Barthéz; Rilliet and Barthéz themselves; the authors of the *Biblioth. du Méd. Prat.*, and M. Guersant; while amongst those opposed to this opinion may be cited, MM. Cruveilhier, Barrier, Dr. Arthur Farre, who greatly doubts the possibility of the accident, and Dr. Condie. We confess ourselves inclined to believe, from facts stated by different authors,

and from the history of two cases which occurred to M. Guersant in 1841, at the Children's Hospital of Paris (*loc. cit.*, p. 680), that worms may in some instances cause a perforation independently of previous disease of the coats of the intestine. In one of these, two lumbrici were found engaged in an opening in the appendix veriformis, half the bodies of the animals being in the appendix and half in the peritoneal sac; while in the other, an opening of the same kind as in the previous case was found in the appendix, and though the three worms which were found lying in the abdominal cavity might have escaped through an ulcerated perforation of the colon, it is not the less true that the opening in the appendix presented the same characters exactly as in the first case, in which the animals were, as the author remarks, "taken in the act." In both instances, the perforation of the appendix was at the extremity of that canal, and in the form of a narrow opening of a conical shape; the membranes were smooth, thinned, and the edges of the orifice sloped off from within outwards; no trace of ulceration was perceptible. On the other hand, we have met with a fatal case of intestinal perforation, dependent on extensive ulceration of the bowel, in which a lumbricoid worm was found lying loosely half-way out through the opening. In this case it was evident that the presence of the worm was purely accidental.

In regard to the *verminous abscesses* already referred to, we shall make but few remarks, referring the reader to more extensive treatises for fuller information. These abscesses have been, in very rare instances, met with in the pharynx and nasal passages, but much more frequently they exist in the abdomen. The latter may be of two kind, *stercoraceous* and *non-stercoraceous*. In the former, the abscess, which forms upon some portion of the walls of the abdomen, gives issue not only to the worm or worms, and pus, but also to fecal and even alimentary substances, and leaves behind a fistula connecting with the cavity of the intestine, which may cicatrize after a short time, or remain open during life. In the other form of abscess, the opening through the coats of the intestine has been closed immediately after the passage of the worm, so that the abscess gives issue only to the animal and pus after which it heals up without giving rise to a fistula.

The verminous abscesses are said to be found generally about the inguinal and umbilical regions; to occur most frequently between the ages of seven and fourteen years, and not to be, as a general rule, very dangerous to life.

SYMPTOMS INDICATIVE OF THE PRESENCE OF WORMS.—We believe it is almost universally acknowledged by later writers, that there is no single symptom, nor group of symptoms, other than the expulsion of the worms, and their detection, which indicates with certainty their existence in the alimentary tract. This is the expressed opinion, amongst others, of MM. Guersant, Rilliet and Barthez, Barrier, Valleix, and Drs. Eberle and Condie, and it is also the opinion which we have ourselves been led to form from our experience amongst children.

Another point worthy of remark is, that even though one or several worms may have been expelled, it is not always fair to conclude that the

symptoms under which the child labors, are the result of the presence of others of these animals, as there may be no more in the bowels, or they may be so few in number as not to produce injurious effects; while, on the contrary, various disorders of the alimentary tract, as chronic indigestion, simple diarrhœa, and inflammatory diseases of the gastro-intestinal mucous membrane, may and do exist simultaneously with, and yet independently of, the presence of these parasites.

The *symptoms* generally enumerated as indicative of the presence of worms are the following: The child presents various signs of disturbed health. The stomach is more or less deranged, as shown by furred tongue, eructations, variable appetite, which is sometimes diminished, and sometimes increased, thirst, acid or heavy breath, and nausea. The abdomen may be enlarged or retracted, generally the former, and is often more or less hard and painful to the touch; the condition of the bowels varies in different cases, as they are sometimes costive, and sometimes affected with diarrhœa. According to M. Guersant, the stools often contain glairy substances, and are sometimes streaked with blood and of a yellowish-green color; the patient often suffers from colic, which may be either dull or acute, though more generally the latter, and which is generally felt at the umbilical region. Children affected with lumbricoides are said to present a peculiar physiognomy; the face is usually paler than natural, and sometimes has a leaden tint; the eyes are surrounded by bluish rings, and have at the same time a dull and languid expression; the inferior eyelids are often swelled and puffy; the sclerotic coat of the eye assumes a bilious tint; the nostrils are said to be sometimes swollen, and the child complains much of irritation and itching of those parts, and is constantly picking at them with the fingers. In some instances epistaxis takes place. The child is generally pale and thin, indolent and languid, or irritable and unhappy. The sleep is almost almost disturbed. This indeed is, it seems to us, one of the most important signs both of worms and of chronic functional disorders of the stomach and bowels. The nights are almost always restless, the patient either waking often to drink, or waking in fright and alarm from dreams, or else constantly tossing and turning in sleep, moaning, or grinding the teeth.

Other symptoms mentioned by different observers, and by some very much depended upon, are acceleration with irregularity of the pulse, and dilatation, especially unequal dilatation, of the pupils. We might cite also strabismus, and occasionally cough.

In children in whom the number of lumbricoides is very large, the constitution suffers to a dangerous degree. The symptoms above enumerated are very marked, and at the same time the child is very pale or sallow, emaciated, weak, and without appetite; the abdomen is hard and tumid; the nervous symptoms are severe, and some of the symptoms which we shall describe presently, under the head of disorders, occasioned by worms, are also observed.

It should be remarked, however, again, that all or any of the symptoms just described may exist independently of the presence of worms, the only certain sign of which is their expulsion from the patient.

MORBID EFFECTS OCCASIONED BY WORMS.—MM. Rilliet and Barthez divided the accidents or effects produced by the existence of lumbricoides into two groups: those which result from the mechanical influence of the entozoa, as their accumulation or displacement; and those which appear to be the consequences of a purely sympathetic action on the different systems of the body, and particularly the nervous system.

MECHANICAL EFFECTS.—Under this head are included perforation and hemorrhage of the intestine, enteritis, abscesses, and the symptoms determined by the displacement or migration of the worms into the ductus communis choledochus, the liver, or the air-passages.

Of perforation and abscesses, we have already treated under the head of anatomical lesions. Hemorrhage is a very rare event, but it occurred in one instance cited by MM. Rilliet and Barthez, and Guersant, from M. Charcelay, in consequence of the rupture of an arteriole in a small rounded ulceration in the duodenum, apparently occasioned by the presence of a large number of lumbrici. Enteritis, as an effect of the presence of worms, has also been referred to under the head of the anatomical lesions. In many instances it is, no doubt, a mere accidental complication, in no way connected with the presence of entozoa; probably this is true of a large majority of the cases. When, however, the number of the parasites is very great, and particularly when they are collected into large or firm knots and bundles, they may, no doubt, occasion, by their mechanical irritation, inflammation, thickening, softening, and even destruction of the mucous tissue, as in cases cited by M. Guersant, from MM. Bretonneau and Charcelay, and in one which occurred to himself. It should be remarked, however, that the cases on record in which ulcerations evidently depend upon the presence of worms, are, so to speak, infinitely few in comparison with those in which no such alteration existed, or in which it was evidently independent of any influence exerted by the worms.

EFFECTS CAUSED BY THE DISPLACEMENT OR MIGRATION OF WORMS.—Lumbricoides have been found, as we have already seen, in the walls of the abdomen, giving rise to abscesses. They have been discovered, also, in the vermiform appendix, in the ductus communis choledochus, in the gall-bladder, in the hepatic ducts, in the substance of the liver, forming abscesses, and in the pancreatic duct. The symptoms occasioned by the latter class of cases are very obscure. In one instance, M. Guersant supposed that an attack of convulsions depended upon the presence of worms in the common duct.

More numerous examples are on record, in which violent dyspnœa and cough, and fatal asphyxia, have occurred in consequence of the pressure of lumbricoides which had passed into the œsophagus, or from their introduction into the larynx, trachea, or bronchi. The symptoms occasioned by these accidents are a sudden attack of dyspnœa, anxiety, agitation, threatened suffocation, dry, spasmodic cough, acute painful cries, pain in the larynx or trachea, and, unless relief be obtained in a few hours, death. This kind of attack may depend on the rising of a worm or bundle of worms into the œsophagus, causing pressure on the larynx

and trachea, as in the case reported by M. Tonnelles, in which the symptoms disappeared after the expulsion of a large number of worms; or else it may be due to reflex spasm of the œsophagus or larynx dependent upon the irritation transmitted from the intestine which is excited by the presence of these parasites. One of us has met with an instance of this kind. It occurred in a boy fifteen years old, presenting every mark of strong and vigorous health, but who, for three or four weeks before we were consulted in regard to him, had been subject to sudden and apparently causeless attacks of suffocation, which seized him without the least warning. When the attack came on, he would for some instants cease to breathe, or breathe with much difficulty. He always seemed to suffer from the greatest anxiety; the countenance became altered and distressed; he was unable to speak, but made signs for water, and when able to swallow a mouthful, which was always exceedingly difficult, was at once relieved. His mother told us that he always appeared to be in the greatest distress, so that, on several occasions, she feared for his life. Striking him violently on the back, which she, when present, always did, sometimes relieved him, but generally the difficulty continued until he could swallow a little fluid of some kind. These attacks were unattended at the time by cough, nor was there the least sign of disorder of the respiratory system in the intervals between them. Suspecting that the difficulty must depend on the rising of a worm or worms into the œsophagus, or upon sympathetic irritation from the presence of these parasites in the stomach, and learning that he had been troubled with worms some years previously, we gave him wormseed oil, which caused the expulsion of a few large lumbricoides, after which he had no return of the symptoms.

The attacks of dyspnoea may depend also, as already stated, on the introduction of worms into the air-passages. Under these circumstances death is very apt to be the result. In one instance, however, reported by M. Arronsson, after the difficulty had lasted two hours, the patient, a little girl eight years old, after violent efforts at coughing, threw up a living lumbricus.

We have next to consider the *sympathetic effects*, and particularly the *nervous symptoms*, occasioned by worms. We may include amongst the nervous symptoms produced by worms the headache, languor, irritability, restless and disturbed sleep, and grinding of the teeth, so frequently observed. These, however, are of but slight importance in comparison with certain other disorders of the nervous system, which do undoubtedly occur sometimes, though we should suppose *very rarely*, in proportion to the whole number of subjects affected with the parasites. The disorders to which we allude are partial or general convulsions, chorea, hysteria, and catalepsy, which are the most frequent, though, as so often stated already, extremely rare in comparison with the number of cases in which the presence of the worms produces no such effects. Other disorders cited by the authors of the *Bib. du Méd. Prat.*, with cases to prove their reality, are insanity, paralysis, coma, palpitations, strabismus, cough, hyperæsthesia of the skin, amaurosis, and aphonia.

DIAGNOSIS.—It has already been stated that there are no certain signs of the presence of worms in an individual except their expulsion. The symptoms which have seemed to us most strongly to indicate their presence are, a chronic disordered state of the digestive apparatus, producing irregular appetite, which is sometimes good and at others bad; slight emaciation; paleness or unhealthy tint of the complexion; languid expression of the face; some irritability of the temper, or a want of the gayety and activity of disposition natural to childhood; picking at the nose; often some tumidity of the abdomen, which may be at the same time either hard or merely tympanitic; and, what seems to us more important than any that we have named, very restless and broken sleep at night, with frequent grinding of the teeth.

M. Valleix remarks that, in a case presenting nervous symptoms simulating disease of the brain, we may suspect the existence of worms, if we learn upon inquiry that the symptoms of marked intestinal disorder, the various signs cited above as indicative of the presence of worms, and different derangements of digestion, had preceded for some time the appearance of the nervous symptoms; chiefly for the reason that, in most diseases of the brain, the alimentary tract is, at the invasion, in a state of integrity, with the exception of sympathetic vomiting. If we can learn, upon inquiry, that the child has discharged worms on some previous occasion, the probability of the dependence of the symptoms upon them becomes still stronger.

It is sometimes difficult to determine positively whether certain substances discharged at stool are fragments of worms, or whether they are portions of imperfectly digested aliment, or foreign bodies. The things which most resemble lumbricoides, are the remains of tendons, ligaments, vessels, fibres of plants, etc. To make the distinction with certainty, the doubtful substance ought to be placed in water, so that it may be thoroughly cleansed, after which it must be carefully examined as to its structure, arrangement, consistence, etc., with the eye, and with the microscope, if necessary. M. Guersant has suggested a very easy method of ascertaining whether the substance be animal or vegetable, which is to subject it to heat after it has been carefully washed, when the odor will at once inform us of its real nature.

PROGNOSIS.—It is no doubt a very rare event, at least in the northern parts of our country, for life to be endangered by the presence of worms. We have never, ourselves, met with an instance in which the general health was more than moderately disturbed by this cause. That verminous affections are sometimes, however, dangerous to life in this city, is shown by three cases related by Dr. Dewees, in which very severe and threatening symptoms were instantly relieved upon the expulsion of lumbrici after the exhibition of vermifuges.

Worms become dangerous to life when they migrate from their original seat to neighboring and important organs, particularly the air-passages and liver. The prognosis is unfavorable also when they accumulate in very large numbers, and give rise to the different nervous symptoms above described.

TREATMENT.—Before commencing our remarks upon the particular remedies employed for the destruction and expulsion of worms from the alimentary canal, we would call the attention of the reader to the fact that most of the recognized anthelmintics are more or less irritating to the gastro-intestinal mucous membrane, and some of them to the nervous system also, producing, in overdoses, severe and even dangerous nervous symptoms. It is evident, therefore, that remedies of this class ought not to be exhibited unless they are manifestly called for, and not at all when symptoms of severe gastro-intestinal irritation, and particularly of inflammation, are present, unless there be the very strongest reasons for supposing that those symptoms depend upon accumulations of worms. We are quite sure that we have, in a considerable number of instances, met with children whose digestive organs had been injured, and in whom slight functional derangement had been converted into severe indigestion, and even inflammatory disorder, by the too frequent or long-continued use, or the administration in excessive quantities, of different vermifuges and of various quack nostrums, which are sold to an amazing extent in this city, and all over the country.

As the diagnosis of worms is always doubtful, it is best never to risk the administration of any of the irritating vermifuges, unless convinced by the previous expulsion of worms, that they are almost certainly present; and indeed, we ourselves rarely give any other remedy than small quantities of the *wormseed oil* in slight, and especially in doubtful cases, unless this has already been tried and failed. For our own experience, we believe that this remedy is all-sufficient in a large majority of the cases that occur in this city; as these are almost always of a mild character, and, as it not only produces the expulsion of the parasites when they exist, but also acts beneficially upon the forms of digestive irritation which simulate so closely the symptoms produced by worms. We are persuaded, indeed, that of all the cases that have come under our notice, in which it seemed probable that worms might be present, none were expelled in nearly half, and yet the signs of disturbed health have passed away under the use of the remedy. The oil of wormseed may be given in doses of four drops to children of two years of age, and of six or ten to those above that age, three times a day for three days, to be followed on the morning of the fourth day by a moderately active, but not irritating cathartic dose, the best of which is castor oil or syrup of rhubarb. The objection to the remedy is its nauseous taste and smell; these, however, may be partially disguised by making it into a mixture with yolk of egg, powdered gum, and syrup of ginger. Some children take it very well dropped upon a lump of white sugar, while others take it best mixed with common brown sugar. If one course of the oil, as it is called, fail to relieve the symptoms, another should be administered. It ought to be recollected that, when given in large doses, the wormseed oil is irritating to the digestive mucous membrane, and produces dangerous nervous symptoms. We know of one case, in which a girl six or seven years of age was made exceedingly ill and suffered for years afterwards, from the effects of a teaspoonful of the oil given by mis-

take. The following is a very good formula for the administration of this remedy :

R. Ol. Chenopodii.	gtt. lx. vel f 3j.
Pulv. Acaciæ,	3 ij.
Syr. Simplicis,	f 3j.
Aq. Cinnamomi,	f 3 ij.—M.

Give a dessertspoonful three times a day, for three days, and repeat after several days.

The wormseed may be given also in powder, in the dose of from twenty to forty grains.

The remedies most frequently employed in this country besides the wormseed, are pink-root or spigelia, oil of turpentine, calomel, and the bristles of cowhage.

We believe that the *pink-root* is more depended upon by us than any other single remedy. It is given either in substance or infusion. The dose of the powder is from ten to twenty grains for a child three or four years old, to be repeated every morning and evening for several days, and followed by an active cathartic. The powder is seldom used, however, as the drug is almost always given in infusion. The best and safest mode of administering it is in combination with cathartic substances. Thus, half an ounce each of pink-root and senna may be infused for a few hours in a pint of boiling water, and a tablespoonful given two or three times a day to children two or three years old, for three, four, or five days, when it should be suspended for a time, and resumed, if necessary. A preparation much used in this city under the title of worm-tea, and which we have ourselves given with very good success, consists of the spigelia mixed with senna, manna, and savine, in different proportions, made into an infusion and sweetened with brown sugar. Dr. G. B. Wood (*Pract. of Med.*, vol. i., p. 623) recommended the following formula :

R. Sennæ, Spigeliæ,	āā 3 ss.
Magnesiæ Sulphat.,	3 ij.
Mannæ,	3 j.
Fœniculi,	3j.
Aquæ Fervent.,	0j.

These are to be macerated for four hours in a covered vessel, and a tablespoonful given to a child two years old once or twice a day, or every other day, so as to procure two or three evacuations in the twenty-four hours. The remedy is continued for a few days, or for one or two weeks, if necessary, and if it do not debilitate the child.

The fluid extract of spigelia and senna has been introduced as a more convenient and acceptable mode of administering this vermifuge with a cathartic. The dose for a child is from thirty minims to a teaspoonful, according to the age.

The *spirit of turpentine* is highly recommended as an efficient remedy for worms by several authorities, and particularly by Dr. Joseph Klapp and Dr. Condie, of this city. Dr. Condie states that it is the article from

which he has derived the most decidedly beneficial effects, and remarks that it may be given when there exists considerable irritation of the alimentary canal, or even subacute inflammation, without any fear of its increasing either. He gives the rectified spirit in sweetened milk, in molasses, or in the following mixture:

R. Mucil. Acaciæ,	f ℥ ij.
Sacch. Alb.,	℥ x.
Spir. Æther. Nitr.,	f ℥ iij.
Ol. Terebinth.,	f ℥ iij.
Magnes. Calcinat.,	℥ j.
Aquæ Menthæ,	f ℥ j.—M.

Of this mixture a dessertspoonful is given every three hours.

We have used the spirit of turpentine but seldom, on account of its extremely disagreeable taste, having always succeeded perfectly well with the wormseed oil, or with infusion of pink-root with cathartics.

Calomel also is highly thought of by many persons as a vermifuge, and, no doubt, when used in combination with or followed by cathartics, or given in full purgative doses, it is very effectual. We can only repeat what we have already said on several occasions, that it is a remedy which, from the powerful influence it exerts upon the constitution, ought not to be given except when really called for; and, as we can almost always succeed in curing verminous affections by milder drugs, we see no occasion for resorting to this, except in rare cases. When used it is given alone in considerable doses, and followed by some cathartic, or in combination with rhubarb and jalap, or jalap, or scammony.

The bristles or down of *cowhage* are also used by some practitioners, no doubt sometimes with success. We have never used them, and can give no opinion, therefore, from personal experience, as to their efficacy. They are administered by making them into an electuary with honey, syrup, or molasses, a teaspoonful of which is given every morning for three days, and then followed by an active cathartic.

The following *electuary*, recommended by Bremser, is very much employed in Europe, and is highly spoken of by Dr. Eberle:

R. Semin. Santonicæ (semen-contræ of the French writers),	
Semin. Tanaceti contus.,	āā ℥ ss.
Valerian. pulv.,	℥ ij.
Jalapæ pulv.,	℥ jss.—ij.
Potass. Sulphat.,	℥ jss.—ij.
Oxymel. Scillæ,	q. s.—ut ft.*
	Electuarium.

A teaspoonful of this is given morning and evening for three or four days, when the dejections generally become more copious and liquid. If they do not produce this effect, Bremser advises that the dose be increased. Dr. Eberle gave it for six or seven days, and says it does far less good when it produces frequent and watery evacuations, than when it causes only three or four consistent stools a day. This preparation has a very dis-

agreeable taste, and children sometimes refuse to take it on that account. When this is the case it may be made into pills.

MM. Rilliet and Barthez recommend the following *syrup*, which was proposed and highly thought of by M. Cruveilhier:

R. Follicul. Sennæ, Rhei, Semin. Santonicæ, Artem. Abrotan.,
Helminthocort, Tanaceti, Artemis Pontic, āā ʒj.

To be infused in half a pint of cold water, strained, and made into a syrup with sugar, of which a tablespoonful is to be given every morning for three days. .

Of late years, santonin, the active principle derived from the European wormseed, has been much employed, and with very good success. The remedy may be given in doses of from one-fourth to one-half of a grain for a child two years old, combined or followed by a dose of castor oil or senna. It is also prepared in the form of sugar-coated dragées, which renders it quite acceptable to children.

The *empyreumatic oil* of Chabert is also highly spoken of by some European authorities. It is made by mixing one part of the empyreumatic oil or fetid spirit of hartshorn, with three parts of spirit of turpentine, and allowing them to digest for four days. The mixture is then put into a glass retort and distilled in a sand-bath until three-fourths of the whole have passed over into the receiver. The product should be kept in small and tightly-closed vials. The dose is about fifteen or twenty drops, three or four times a day, for children between two and seven years old. This is recommended highly by Bremser and other authorities. The great objection to it is its exceedingly nauseous taste. Dr. Eberle speaks in very favorable terms of a strong decoction of *helminthocorton* or Corsican moss, which he has found "not only valuable as a vermifuge, but particularly so as a corrective of that deranged and debilitated condition of the alimentary canal favoring the production of worms." An ounce of helminthocorton, with a drachm of valerian, are to be boiled in a pint of water down to a gill, and a teaspoonful of the decoction given morning, noon, and evening. It is particularly beneficial in cases attended with the usual symptoms of worms, connected with want of appetite and mucous diarrhœa, and arising from debility of the digestive organs, and a vitiated condition of the intestinal secretions.

Kameela, the reddish-brown powder which clothes the capsules of the *Rottlera tinctoria*, has been of late highly recommended, not only in cases of tænia, but of *ascaris lumbricoides*. The dose for children is about gr. v, repeated till it has acted on the bowels.

In all cases of deranged health supposed, either from the nature of the symptoms, or proved by the previous expulsion of worms, to depend on the presence of these animals in the alimentary canal, it is exceedingly important to attend to the *hygienic treatment* of the child, and in some instances to administer *tonics* and *stimulants*. In not a few cases that have come under our own notice, in which many of the symptoms supposed to indicate the presence of worms have been extremely well marked, we have succeeded in removing them all without a resort to any vermifuge, by the treatment proper for the chronic indigestion or dyspepsia of children. The method

of treatment to be employed in such cases has already been laid down in the article on digestion, to which the reader is referred for full information. It should consist chiefly in strict attention to exercise and diet, and in the use of tonics, as quinia and iron, and small quantities of fine port wine.

Whenever any complication exists in connection with worms, the treatment must be modified according to its nature. If it consist in inflammation of any part of the alimentary tract, the inflammation ought to be attended to first, and the verminous disorder for the time let alone. If the inflammation be very slight, or if the symptoms indicate only severe irritation rather than positive inflammatory action, we may exhibit the milder and least injurious vermifuges, as very small doses of wormseed oil, which we have never known to do harm, the decoction of helminthocorton and valerian, recommended by Dr. Eberle, or, according to Dr. Condie, the spirit of turpentine. If the verminous affection coëxist with any of the acute local inflammations of the thorax, the former ought to be, as a general rule, let alone, until the latter has been relieved by appropriate treatment. In doubtful cases, in which it is impossible to ascertain with certainty whether the symptoms depend on worms, or upon a simple dyspeptic condition of the digestive organs, it is most prudent to give only the simplest and least irritating vermifuges, to regulate the hygienic conditions of the patient, and afterwards to resort to tonics, if necessary.

Various writers, and particularly M. Guersant, advise that we should forbid, in verminous cases, the use of relaxing food, especially of milk preparations, fruits, and of fatty and farinaceous substances; and that, after the expulsion of the worms, we should direct a tonic and strengthening regimen. The diet should consist of boiled and roasted meats, of wine, and of bitters. The author just quoted, states that a change of food alone will often suffice to procure the expulsion of worms. He says (*Dict. de Méd.*, t. xxx, p. 689), "I have met with children who had been tormented with ascarides lumbricoides while residing in the country and living upon milk and fruits, and who, upon being brought to the city, and put upon the use of broths and soups, passed considerable quantities of worms, and after that got entirely rid of them."

Occasionally our opinion is asked with reference to worms of other varieties, which are reported to have been passed from the rectum of children. Thus, tapering elongated pieces of coagulated casein may be mistaken for worms.

So, too, we have seen a specimen, submitted to us by Dr. Bussey, of Buena Vista, Texas, and said to have been passed by a boy there, of male Gordicus aquaticus, or horse-hair worm. This is a nematoid worm of chestnut-brown color, a foot in length, a little more than one-half line in breadth, with a bifid caudal extremity. It grows in stagnant water, and thus may readily have been swallowed and passed per anum.

ARTICLE II.

ASCARIS VERMICULARIS.

THE *description* of this worm has already been given at page 1004.

SEAT.—The ascaris vermicularis is found almost exclusively in the large intestine, and in a large majority of the cases is confined to the rectum. It is said to have been found in the vagina in the female, having no doubt passed from the rectum into that canal.

The *causes* which determine the presence of this worm are not at all understood.

SYMPTOMS.—The characteristic, and often the only symptom indicative of their presence, is violent itching about the anus, which is sometimes almost insupportable, and which is generally most troublesome and most apt to occur at night when the child is in bed. In consequence of this, the sleep is much disturbed, and the child grows peevish and irritable, and may suffer considerable impairment of general health. In some instances they give rise to acute and violent pain in the region of the anus, and sometimes to tenesmus and mucous or bloody stools. When the last-named severe symptoms exist, the worms may occasion dangerous nervous disorders, and even give rise to general convulsions. The worms not unfrequently escape from the rectum and are found upon the bed-clothes, or upon the clothes which the child has worn through the day. Sometimes they are discharged in considerable numbers, and are found, in that case, either mixed with the fæces, or with mucus, or collected into balls or knots.

The *diagnosis* of the seat-worm, like that of the lumbricoides, cannot be regarded as positive, unless some have been expelled, or unless they can be seen by examination of the rectum. This can generally be done when they are present in any number, by pressing the nates apart so as to open the anus and bring the folds of the mucous coat of the bowel into view. The only other symptom which indicates their presence with any certainty, is the existence of severe itching about the anus, not to be explained upon any more reasonable supposition.

PROGNOSIS.—These worms do not, as a general rule, produce the same disturbances of the general health as lumbricoides, and in not a few instances are entirely innocuous, with the exception of the pain and inconvenience they occasion.

They are, however, exceedingly troublesome, because of the difficulty of removing them entirely by any treatment. No matter how many are discharged, some almost always remain concealed in the folds of the mucous membrane, and as they are propagated with great rapidity, the same train of symptoms is very apt to return soon after they may have been seemingly dislodged.

TREATMENT.—It has been found by long experience that the common vermifuges, given by the mouth, exert much less influence in causing the expulsion of these worms than of the lumbricoides. For this reason enemata are generally resorted to in the treatment, instead of remedies

given by the mouth. Dr. Dewees, however, recommends the elixir proprietatis (tinct. aloes et myrrhæ), in small and often-repeated doses, continued for some time, and followed by enemata of lime-water, camphor, or aloes. He gave twenty drops of the elixir three times a day, in a little sweetened milk, to children from two to four years old, and thirty drops to those between five and seven years.

The plan we have generally resorted to has been to give small doses of the wormseed oil, as directed in the article on lumbricoides, and to direct an injection of from four to six grains of powdered aloes, suspended in a gill of warm milk, for children four years old, to be repeated once in three, four, or five days, according to the necessity of the case.

Lime-water by injection is recommended by several different authorities. It may be given of its ordinary strength, or mixed with an equal quantity of warm milk, or flaxseed mucilage. Other enemata recommended are spirit of turpentine in milk, a teaspoonful of the former to a gill of the latter; decoction of helminthocorton; a strong infusion of quassia (℥ij to Oj) affords a most efficient and harmless injection; an injection made by infusing two drachms of fresh garlic-cloves in three ounces and a half of boiling water, and adding to the infusion, after it has been poured off, a scruple of assafoetida rubbed up with the yolk of an egg; a solution of from six to twelve grains of sulphuret of potassium in half a pint of water; injections of sweet oil, or of lard beaten up with water until it becomes fluid, and even of cold water. The last two mentioned substances have the advantage of calming the itching and irritation of the rectum almost immediately. Enemata of a solution of nitrate of silver, in the proportion of two to four grains to the ounce of water, have been recommended by Schultz (*Deutsche Klinik*, quoted in *Med. Times and Gaz.*, 1858), who asserts that two, or at most three, of these injections suffice to effect a cure. Again, it has been recommended to pass a bougie smeared with mercurial ointment into the rectum. We should much prefer a method of using this ointment which succeeded in the hands of M. Cruveilhier in a very severe case. This was to place a little of the ointment on the anus, by which course the patient was entirely relieved after a few days. In a very obstinate case in an adult, we succeeded in entirely destroying the worms by the daily use of suppositories, made unusually long, and impregnated with carbolic acid. M. Valleix states that he has obtained the same results by causing the anus to be anointed with the following preparation, a small quantity of which was introduced at the same time into the inferior extremity of the intestine:

R. Hydrarg. Chlor. Mitis,	.	.	.	℥iv.
Axung.,	.	.	.	℥vj.—M.

Dr. Wood states that a dose of sulphur taken every morning before breakfast has been found very useful.

The diet and general health ought always to be strictly inquired after, and attended to by the physician. For information upon these points the reader is referred to the remarks upon hygienic treatment in the last article.

INDEX.

- ABDOMEN**, condition of, in cholera infantum, 451
 in gastritis, 401
 in entero-colitis, 420
 in tuberculous peritonitis, 689
 in tuberculosis of mesenteric glands, 689
 in typhoid fever, 721
 in worms, 1009
 examination of, and signs from, 46, 47
Abscess of leg, simulating rheumatism, 32
 of lung following pneumonia, 162
 bronchial, in bronchitis, 200
 iliac, in disease of cæcum and appendix, 468
 retropharyngeal, 373
 following erysipelas, 868
Absorbents in entero-colitis, 431
Acarus scabiei, description of, 999
Achorion, Schœnleinii, description of, 982
Acids in local treatment of gangrene of the mouth, 343
 carbolic, in gangrene of the mouth, 343
 in diphtheria, 903
 in favus, 989
 in scabies, 1001
 muriatic, in gangrene of the mouth, 343
 in diphtheria, 902
 in typhoid fever, 728
 sulphuric, in chronic entero-colitis, 439
Acne, in congenital syphilis, 709
Ægophony, in pleurisy, 233
Affusion of cold water, in scarlatina, 815
 of warm water, in scarlatina, 815
Air, as an injection in intussusception, 494
Albuminoid degeneration of viscera in scrofula, 675
Albuminuria, in croup, 89
 in diphtheria, 894
 in pneumonia, 179
 in scarlatina, 729
 in scarlatinous dropsy, 794
 in scrofula, 676
 in typhoid fever, 724
 in variola, 736
Alkalies, in membranous croup, 105
 in rheumatism, 669
 in thrush, 363
 local use of in diphtheria, 903
 in skin diseases, 988, 992, 994
Alopecia areata, article on, 996-997
 frequency of, 996
 fungous nature of, doubtful, 996
 contagion as cause of, 996
Alopecia areata, symptoms of, 996
 condition of hair in, 996
 baldness following, 996
 diagnosis of, 996
 prognosis in, 996
 treatment of, 997
Alphos, 933
Alum as an emetic in true croup, 103
 in whooping-cough, 274
Anæsthetics, use of during tracheotomy, 123
 in eclampsia, 574
 in tetanus, 608
Analysis of condensed milk, 321
 of cow's milk, 303
 of human milk, 308
 of bones in rickets, 702
 of fluid in hydrocephalus, 550
 of fluid in sclerema, 978
Anatomical lesions, in albuminoid degeneration of the viscera, 675
 in ascaris lumbricoides, 1008
 in atelectasis pulmonum, 134
 in bronchitis, 198
 in bronchial phthisis, 683
 in cerebral congestion, 537
 hemorrhage, 542
 in cholera infantum, 445-449
 in chorea, 614
 in cæcum and appendix, diseases of, 472
 in collapse of the lung, 145.
 in convulsions, 567
 in contraction with rigidity, 594
 in coryza, 53
 in croup, 92
 in cyanosis, 281
 in diarrhœa, 389
 in diphtheria, 879-884
 in dysentery, 462
 in eclampsia, 567
 in eczema, 935
 in emphysema, 218
 in enlargement of the tonsils, 366
 in endocarditis, acute, 292
 in entero-colitis, 412
 in gangrene of the mouth, 337
 in gastritis, 398
 in hemorrhage, cerebral, 542
 in whooping-cough, 269
 in hydrocephalus, 549
 in intussusception, 483
 in laryngismus stridulus, 579
 in laryngitis, simple, 62
 in laryngitis, simp. spasmodic, 70
 in laryngitis, pseudo-membranous, 92

- Anatomical lesions, in measles, 845
 in meningitis, simple, 530
 tubercular, 501
 epidemic cerebro-spinal, 910
 in mumps, 862
 in night terrors, 661
 in paralysis, atrophic infantile, 641
 pseudo-hypertrophic, 658
 facial, 651
 in pericarditis, 290
 in pharyngitis, 368
 in pleurisy, 231
 in pneumonia, 160-162
 in pneumothorax, 253
 in phthisis pulmonum, 680
 in rickets, 702
 in rōtheln, 851
 in scarlatina, 802
 in sclerema, 978
 in stomatitis gangrenosa, 337
 in syphilis, congenital, 712
 in tetanus nascentium, 604
 in thrush, 349
 in tuberculosis, 679
 in typhoid fever, 716
 in valvular disease of heart, 293
 in variola, 740
- Angina; *see* Pharyngitis and Tonsillitis.
 in diphtheria, 885
 in scarlatina, 813
- Antimony in catarrhal croup, 80
 in capillary bronchitis, 213
 in children, excessive action of, 188
 in pleurisy, 242
 in pneumonia, 188
 in pseudo-membranous laryngitis, 103
- Antispasmodics in chorea, 627
 in eclampsia, 574
 in laryngismus stridulus, 589
 in laryngitis, simp. spasmodic, 81
 in tetanus, 609
- Aortic valve, diseases of, 293
- Appendix cœci, catarrhal inflammation of, 472
 perforative ulceration of 477
 article on diseases of cœcum and appendix, 467
 age as cause, 469
 sex as cause, 470
 intestinal concretions as cause, 470
 anatomical lesions of, 472
 cases of, 473
 symptoms of, 475
 duration of, 477
 prognosis in, 478
 diagnosis of, 478
 treatment of, 479
- Arsenic in chorea, 629
 in scrofula, 678
 in malarial fever, 860
 in eczematous affections, 944
- Artificial food (*see* Food, Diet, and Milk).
- Ascaris lumbricoides*, article on, 1007
 description of, 1004
 synonyms of, 1004
 early age as cause of, 1007
 disposition to hereditary, 1007
- Ascaris lumbricoides*, seat of, 1007
 number of, 1007
 anatomical lesions in, 1008
 condition of mucous membrane in, 1008
 perforation of intestine by, 1008
 hemorrhage from bowel in, 1009
 verminous abscesses in, 1009
 no diagnostic symptoms of, 1009
 digestive disturbances caused by, 1010
 restlessness caused by, 1010
 peculiar physiognomy caused by, 1010
 mechanical effects of, 1010
 effects caused by displacement of, 1011
 dyspnœa and cough caused by, 1011
 nervous symptoms caused by, 1011
 diagnosis of, 1012
 prognosis in, 1012
 treatment of, 1013
 caution in use of vermifuges in, 1013
 wormseed in cases of, 1014
 pinkroot in cases of, 1015
 turpentine in cases of, 1015
 calomel in cases of, 1016
 santonin in cases of, 1016
 kameela in cases of, 1017
 general treatment in cases of, 1017
 treatment of complications in cases of, 1018
 diet in cases of, 1018
- Ascaris vermicularis*, article on, 1019
 description of, 1004
 synonyms of, 1019
 seat of, 1019
 causes of, 1019
 symptoms of, 1019
 diagnosis of, 1019
 prognosis in, 1019
 treatment of, 1019
 enemata in treatment of, 1019
 ointments in treatment of, 1019
- Assafœtida in chorea, 627
- Astringents in cholera infantum, 456
 in entero-colitis, 420, 437
 local use of, in diphtheria, 902
- Ataxia, locomotor, after diphtheria, 900
- Atelectasis pulmonum, and collapse of the lung, 134-143
 peculiarity of respiration in, 40, 41
 forms of, 134
 congenital, anatomical appearances in, 134
 causes of, 135,
 symptoms of, 136
 in early weeks of life, symptoms of (*see* Collapse), 137
 case of, 138
 diagnosis of, 140
 prognosis in, 141
 treatment of, 141
 effects of position in, 142
 post-natal, 143
 as cause of sclerema, 976
- Athrepsia, 316, 348
- Atmospheric pressure as cause of deformity in rickets, 700

- Atrophy, muscular (*see* Infantile Paralysis), 634
- Auscultation, of heart, 36
- of lungs, 42-45
- best position of child in, 43
- in true croup, negative results of, 98, 116
- in pneumonia, 168, 174
- in bronchitis, 206
- in pleurisy, 233
- of heart in chorea, 615
- in bronchial phthisis, 685
- in pulmonary phthisis, tuberculosis, 687
- of head in rickets, 697
- Baldness, following alopecia areata, 996
- Baths, in treatment of cholera infantum, 461
- in treatment of skin diseases, 954-971
- cold, as prophylactic in catarrhal laryngitis, 84
- in treatment of chorea, 631
- hot, in grave cases of scarlatina, 812, 828
- in treatment of scarlatinous dropsy, 828
- in treatment of tetanus, 609
- warm, in treatment of catarrhal laryngitis, 84
- in eclampsia, 572
- in rubeola, 851
- in scarlatina, 828
- in tetanus, 609
- in variola, 746
- Becquerel, pulse in children, 34
- Belladonna, in spasmodic laryngitis, 82
- in whooping-cough, 271
- in infantile paralysis, 647
- in tetanus, 609
- as a prophylactic in scarlatina, 830
- Bennett, J. Hughes, restorative treatment of pneumonia, 183
- on bleeding in treatment of pneumonia, 185
- Berg, fungous nature of thrush, 353
- Billard, cry and pulse in children, 33
- Bismuth, in eczema, 946
- in entero-colitis, 431
- Blebs (*see* Bullæ), 956
- Bleeding, in bronchitis, 212
- in cerebral hemorrhage, 547
- in eclampsia, 572
- in gastritis, 402
- in whooping-cough, 271
- in intussusception, 494
- in laryngitis, spasmodic, 80
- pseudo-membranous, 101
- in meningitis, simple, 535
- tubercular, 520
- epidemic cerebro-spinal, 917
- in paralysis, atrophic infantile, 647
- in pleurisy, 242
- in pneumonia, 185
- in rubeola, 851
- in typhlitis, 479
- Blisters, mode of using in children, 191
- Blisters, in pericarditis, 291
- in pleurisy, 244
- in pneumonia, 191
- Blood, condition of, in malarial fever, 859
- in measles, 846
- in scarlatina, 803
- in typhoid fever, 717
- in variola, 740
- Bloodvessels of skin in sclerema, 979
- Bones, alteration of, in rickets, 698
- diseases of, in congenital syphilis, 712
- Bothriocephalus, 1005
- Bouchut, pulse in children, 35
- expectoration in pneumonia, 175
- Brain, condition of, in cerebral congestion, 537, 541
- in cerebral hemorrhage, 541
- in chorea, 617
- in congenital syphilis, 712
- in meningitis, simple, 530
- tubercular, 503
- in scarlatina, 802
- in tetanus, 604
- in typhoid fever, 717
- Bretonneau, on nasal diphtheria, 888
- Bronchi, dilatation of, in capillary bronchitis, 199
- in chronic bronchitis, 206
- physical signs of, 207
- Bronchial abscess, in bronchitis, 200
- glands, tuberculous of, 679, 683
- phthisis (*see* Tuberculosis of Bronchial Glands).
- Bronchitis, connection of, with atelectasis, 145
- in typhoid fever, 722
- in whooping-cough, 266
- in measles, 841, 852
- in rickets, 701
- in scrofula, 675
- effects of temperature and season on mortality of, 196
- article on, 195-218
- definition of, 195
- synonyms of, 195
- frequency and mortality of, 196
- forms of, 196
- predisposing causes of: age, sex, season, clothing, 196
- exciting causes of, 197
- anatomical alterations in:
- acute ordinary form, 198
- capillary form, 199
- bronchial abscess, 200
- dilatation of bronchi, 199
- condition of lung tissue in, 201
- lesions in chronic form, 201
- symptoms of simple acute form, 202
- aggravation of symptoms at night, 203
- duration of simple acute form, 203
- danger of collapse of lung in, 203
- symptoms of capillary form, 204
- duration of capillary form, 205
- symptoms and course of chronic form, 205
- physical signs of, 206
- cough in, 207

- Bronchitis, sputa in capillary form of, 207
 peculiar cough in capillary form of, 207
 respiration and pulse in, 207
 temperature in, 208
 decubitus in, 208
 digestive organs in, 208
 expression in, 208
 urine in, 209
 diagnosis of, 209
 peculiarity of dyspnoea in, 209
 prognosis in, 210
 treatment of, 211
 importance of confinement to bed in, 211
 bleeding in, 212
 emetics in, 213
 antimony in, 213
 ipecacuanha in, 214
 external applications in, 214
 use of stimulants in, 215
 use of quinia in, 215
 treatment of chronic form, 216
- Bullæ, chapter on, 956
- Bullous inflammation of the skin, 956
- Calomel (*see* Mercury), use in spasmodic croup, 80
 in membranous croup, 104
 use in dysentery, 465
 in entero-colitis, 427
 in cholera infantum, 461
 in tubercular meningitis, 523
 in simple meningitis, 535
 in chronic hydrocephalus, 556
 in eczematous affections, 945
 as a vermifuge, 1016
- Cannabis indica in tetanus, 608
- Calabar bean in chorea, 629
- Canula for tracheotomy, details of size and form, 120
- Cane sugar in condensed milk, 322
- Capillary bronchitis (*see* Bronchitis), 204
- Carbolic acid (*see* ACID).
- Carmo-pedal spasms, 595
 in laryngismus stridulus, 584
- Caseine in human milk, 309
- Cases, illustrative of cerebral pneumonia, 178
 cœcum and appendix, diseases of, 473
 collapse of the lungs, 138, 151
 contraction with rigidity, 597
 coryza, chronic, 60
 cyanosis, 283, 287
 croup in scarlatina, 790-793
 emphysema, 220, 226
 heart clot in diphtheria, 896
 hemorrhage, cerebral, 545
 heart, valvular diseases of, 297
 intussusception, 485
 laryngitis, pseudo-membranous, 128
 laryngismus stridulus, 591
 meningitis, epidemic cerebro-spinal, 902
 tubercular, 517, 528
 milk, condensed, use of, 326
 paralysis, atrophic infantile, 638
- Cases illustrative of paralysis, following diphtheria, 899
 pleurisy, chronic, 251
 pneumothorax, 254
 protective power of vaccination, 756
 scarlatina, 781, 782, 786
 tinea, 994
 valvular diseases of the heart, 297
- Catarrh of stomach and intestine (*see* Indigestion), 376
 of stomach, 399
 in measles, 834
- Catarrhal inflammation of the skin, 933
- Cauterization of variolous pock to prevent pitting, 749
 in treatment of diphtheria, 901
- Cavities, tuberculous, 680
- Cerebral congestion, article on, 537-539
 symptoms of, 537
 forms and frequency of, 537
 duration of, 538
 prognosis in, 538
 diagnosis of, 539
 treatment of, 539
- Cerebral form of pneumonia, 178
- Cerebral hemorrhage, article on, 540-548
 definition and frequency of, 540
 forms of, cerebral and meningeal, 540
 causes of, 540
 anatomical lesions of, 541
 of the meningeal form, 542
 transformation of the clot in, and formation of pseudo-cyst, 543
 symptoms of cerebral form, 544
 case of, 545
 meningeal form, 545
 chronic hydrocephalus following meningeal form, 546
 duration of, 546
 diagnosis of cerebral form, 547
 of meningeal form, 547
 prognosis in, 547
 treatment of, 547
 depletion in, 547
 cold and counter-irritation in, 548
 treatment of paralysis following, 548
 chronic hydrocephalus following, 548
- Cerebral symptoms, in pneumonia, 178
 in cholera infantum, 451
 in intussusception, 490
 in tubercular meningitis, 506, 511
 in simple meningitis, 532
 in cerebral congestion, 538
 in cerebral hemorrhage, 545
 in chronic hydrocephalus, 553
 in laryngismus stridulus, 584
 in contraction with rigidity, 594
 in chorea, 620
 absence of, in atrophic infantile paralysis, 636
 in mumps, 864
 in scarlatina, 779, 786
 in measles, 837, 844
 in variola, 732
 in typhoid fever, 718, 720, 723
 caused by worms, 974, 981

- Chambers' analysis of condensed milk, 320
 Chemical characters of false membranes, 881
 Chenopodium, oil of, in treatment of worms, 1015
 Chicken-pox (*see* Varicella.)
 Chicken-breast in rickets, 700
 Chloral in eclampsia, 574
 in tetanus, 580
 Chloroform in eclampsia, 574
 in laryngismus, 590
 Cholera Infantum, article on, 441-462
 general remarks on, 441
 definition and synonyms of, 442
 frequency of, 442
 causes of, 443
 great heat as a cause of, 443
 improper diet as a cause of, 443
 hygienic conditions favorable to, 444
 anatomical appearances and pathology of, 445
 symptoms of, 450
 character of stools in, 450
 of vomiting in, 451
 course and duration of, 452
 diagnosis of, 452
 prognosis in, 452
 prophylactic treatment in, 453
 treatment of stage of evacuation, 455
 of stage of collapse, 456
 importance of free supply of water in, 457
 mistura indica in, 458
 importance of rest in, 459
 treatment of stage of reaction, 459
 importance of attending to state of gums in 460,
 use of baths in, 461
 use of calomel in, 461
 Chorea, article on, 610-634
 definition and synonyms of, 610
 frequency of, 610
 early age as predisposing cause of, 610
 other predisposing causes of, 611
 rheumatism as cause of, 612, 668
 fear and other exciting causes of, 614
 anatomical lesions in, 614
 lesions of heart in, 615
 brain in, 617
 spinal cord in, 617
 portions of body affected in, 618
 prodromic symptoms of, 618
 symptoms of invasion of, 618
 of the confirmed disease, 618
 respiratory muscles and heart at times affected, 620
 paralysis of sphincters in, 620
 loss of voluntary power in, 619
 general symptoms in, 620
 condition of urine in, 620
 cardiac murmurs in, 620
 course of, 621
 effects of acute intercurrent disease upon, 621
 duration of, 621
 frequency of relapses in, 621
 Chorea, nature of, 621
 probable seat of lesion in, 622
 alterations of blood as cause of, 623
 reflex irritation as cause of, 623
 embolism as cause of, 292, 624
 mode of action of rheumatisms as cause of, 624
 diagnosis of, 625
 prognosis in, 625
 statistics of mortality in, 626
 unfavorable symptoms in, 626
 duration of, 626
 treatment of, 626
 use of purgatives in, 627
 antispasmodics in, 627
 cimicifuga in, 627
 bromides in, 628
 conium in, 629
 physostigma in, 629
 arsenic in, 629
 strychnia in, 630
 stimuli in, 630
 tonics in, 630
 baths in, 631
 counter-irritation in, 631
 electricity in, 631
 gymnastic exercises in, 632
 hygienic treatment of, 633
 Chronic bronchitis, 200
 pleurisy, 233
 Cimicifuga racemosa in chorea, 627
 Circulatory organs, diseases of, 276
 Clark, J. L., state of spinal cord in tetanus, 579
 in chorea, 591
 Clinical examination of children, 17
 Club-foot in infantile paralysis, 625
 in progressive muscular sclerosis, 655
 Cod-liver oil in habitual indigestion, 362
 in rickets, 875
 in tuberculosis, 862
 in congenital syphilis, 881
 in eczematous affections, 944
 in tuberculosis, 693
 in scrofula, 677
 in rickets, 705
 Climate, in tetanus, 603
 in scrofula, 677
 Cœcum and appendix cœci, article on diseases of, 467-481
 (*See also* Typhlitis, Perityphlitis, and Appendix.)
 synonyms and definition of, 467.
 seat and character of, 468
 causes of, 469
 intestinal concretions and foreign bodies as causes of, 470
 anatomical lesions in, 472
 illustrative cases of, 473
 symptoms of, 475
 duration of, 477
 prognosis in, 478
 diagnosis of, 478
 treatment of, 479
 typhlitis, 475
 perforation of cœcum, 476

- Cæcum and appendix cæci, perforative ulceration of, 477
perityphlitis, 476
- Cæcum, symptoms of fecal distension of, 475
perforative ulceration of, 477
inflammation of (*see* Typhlitis).
- Cold, as cause of dropsy after scarlatina, 794
applications in tubercular meningitis, 525
in simple meningitis, 535
in cerebral hemorrhage, 548
in eclampsia, 572
in laryngismus stridulus, 590
in tetanus, 609
in scarlatina, 820
- Collapse of lung, in bronchitis, 203
in whooping-cough, 265
in rickets, 703
article on, 134-155
in early weeks of life, symptoms of, 137
cyanosis in, 138
cases of, 134, 138
diagnosis of, 140
prognosis in, 141
treatment of, 142
of post-natal atelectasis, 143-155
general remarks on the pathology of, 143
identity of lobular pneumonia with, 144-145
anatomical lesions in, 145
congestion of lung accompanying, 146
difference between condition of lung in, and in pneumonia, 147
portions of lung affected in, 148
causes of and explanation of mode of production, 148
symptoms of, 150
cases of, 151-152
diagnosis of, 153
prognosis in, 154
treatment of, 154
use of emetics in, 155
treatment of, when combined with bronchitis, 155
- Coloration of skin, significance of changes of, 22
in infants, 30
of face, in pneumonia, 176
in tubercular meningitis, 511
- Compression of head in hydrocephalus, 557
- Concretions, intestinal, 470
- Condensed milk (*see* Milk).
- Condylomata in congenital syphilis, 710
- Congenital syphilis (*see* Syphilis).
- Congestion of the brain (*see* Cerebral Congestion), 537
of the lungs, not inflammatory, 163
in bronchitis, 201
in typhoid fever, 722
- Conium in tetanus, 609
in chorea, 629
- Conjunctivitis, in variola, 747
- Constipation, as cause of diseases of the cæcum, 470
in intussusception, 488
in tubercular meningitis, 507-509
- Contagion of alopecia areata, 996
of diphtheria, 875
of favus, 984
of whooping-cough, 260
of mumps, 861
of röteln, 855
of rubeola, 831
of scabies, 998
of scarlatina, 772
of thrush, 349
of tinea, 991
of typhoid fever, 716
- Contraction with rigidity, article on, 593-601
a rare affection, 593
definition of, 593
causes of, 594
nature of, one of the forms of eclampsia, 594
symptoms of, 594
carpo-pedal spasms in, 595
diagnosis of, from symptomatic contraction, 596
prognosis in, 596
treatment of, 597
- Contracture (*see* Contraction).
- Convulsions, general, or Eclampsia, article on, 559-576
general remarks on; forms of, 559
definition, synonyms, frequency, 559
predisposing causes of, 560
most frequent before age of seven years, 560
nervous temperament as a predisposing cause of, 560
hereditary nature of, 561
exciting causes of, 562
frequency of different forms of, 562
prodromic symptoms of, 563
symptoms of the attack, 564
partial, varieties of, 565
general, duration of, 565
nature of, 566
M. Hall's views on spasm of the larynx, 566
centric and eccentric causes of, 567
no lesion as yet detected in, 567
diagnosis of, from epilepsy, 568
the form of convulsion, 569
prognosis in, 570
treatment of, 571
importance of discovering cause of attack, 571
treatment of attack, 572
bleeding in, 572
emetics in, 573
purgatives in, 573
antispasmodics and opium in, 574
chloroform in, 574
internal, definition of, 575
symptoms of paroxysm, 585

- Convulsions, general, degree of laryngismus present, 585
 incomplete, or holding-breath spells, 586
 rarely dangerous, 586
 in pneumonia, 178
 in whooping-cough, 262, 278
 in tubercular meningitis, 509
 in simple meningitis, 532
 in meningeal apoplexy, 545
 in scarlatina, 784
 in measles, 835
 uræmic, in scarlatina, 798
 in rickets, 701
 in initial stage of measles, 835
 in later stages of measles, 845, 849
 in typhoid fever, 720
 in worms, 1012, 1019
 Cooley, method of preserving milk, 307
 Coruill and Ranvier, false membrane in diphtheritic croup, 86
 Corson, cold affusions in scarlatina, 816
 Coryza, definitions, synonyms, forms, frequency, 52
 causes of, 52
 anatomical lesions in, 53
 symptoms of mild form, 53
 of severe form, 54
 epistaxis in, 54
 duration of, 55
 prognosis in, 55
 in the course of other diseases, 55
 chronic symptoms and duration of, 55
 treatment of acute, 57
 local, of acute, 58
 of chronic, 59
 case of chronic, 60
 in congenital syphilis, 710
 in scarlatina, 788
 in measles, 834
 Cough, in bronchial phthisis, 683
 in bronchitis, 207
 in croup, 95
 in whooping-cough, 261
 in laryngitis, chronic, 64
 simple, 63
 spasmodic simple, 73, 75
 in measles, 837
 in phthisis, 686
 in pleurisy, 235
 in pneumonia, 175
 in typhoid fever, 722
 in worms, 1011
 Countenance, alterations of, 21
 Counter-irritation in bronchitis, 214
 in tubercular meningitis, 525
 in simple meningitis, 536
 in chorea, 631
 in pulmonary complications of measles, 853
 Country residence, importance of in summer, 424, 454
 in tuberculosis, 693
 Cowhage as a vermifuge, 1016
 Cow-pox (*see* Vaccine Disease).
 Coxalgia, simulated by infantile paralysis, 647
 Cracked-pot sound in bronchial phthisis, 684
 Craniotabes, 698
 Cream, proportion of in cow's milk, and mode of determining, 304
 in human milk, 309
 Creamometer, 305
 Croup, diphtheritic, 882
 relations of to pseudo-membranous laryngitis, 85, 89, 882
 false, spasmodic, or catarrhal (*see* Spasmodic simple laryngitis), 69
 true or membranous (*see* Pseudo-membranous laryngitis), 85
 secondary, in scarlatina, 792
 tracheotomy in, 107
 Crust, vaccine, characters of, 761
 Crusta lactea (*see* Eczema capitis), 937
 Crûte de lait, 937
 Cry, characters of the, 24
 peculiar in tubercular meningitis, 507
 alterations of in simple laryngitis, 63
 peculiar in sclerema, 977
 Crystalli (*see* Varicella).
 Cuniculus in scabies, 999
 Currie, cold affusions in scarlatina, 816
 Cutaneous diseases, 918
 not transmitted by vaccination, 759
 Cutaneous diphtheria, 883, 889
 surface, signs from, 22, 30
 Cyanosis, in collapse of the lung, 138
 article on, 281
 definition of, 281
 anatomical appearances in, 281
 illustrated cases of, 283
 theories of mode of production of, 285
 symptoms of, 285
 date of appearance of lividity in, 286
 modes of death in, 286
 duration of life in, 287
 treatment of the form due to atelectasis, 288
 of paroxysms of dyspnoea, 288
 hygienic treatment of, 288
 effect of position on, 289
 neonatorum, Professor Meigs on treatment of, 142
 Cynanche parotidea (*see* Mumps).
 Cynanche maligna (*see* Diphtheria), 873
 tonsillaris (*see* Tonsillitis), 364
 Cyst, pseudo-, in arachnoid, in meningeal apoplexy, 543
 Deafness after diphtheria, 899
 Decubitus of children, 28
 in different diseases, 29
 in tubercular meningitis, 512
 Deformities in chorea, 639
 in rickets, 706
 Deglutition, difficulty of in bronchial phthisis, 684
 in diphtheria, 886
 in retropharyngeal abscess, 374
 in scarlatina, 788
 Dentition as cause of entero-colitis, 412
 of cholera infantum, 445
 of eclampsia, 562

- Dentition as cause of laryngismus stridulus, 578
 of infantile paralysis, 636
 impeded in rickets, 697
- Depilation in favus, 989
 in tinea, 994
- Desiccation in variola, 735
- Desquamation in scarlatina, 779
 in measles, 837
 in small-pox, 735
 in erysipelas, 869
- Development, degree of as aid in diagnosis, 27
- Diagnosis in children, difficulties of, 18
 general method of, 19
 of alopecia areata, 996
 of aphthæ from ulcero-membranous stomatitis, 989
 of ascaris lumbricoides, 1009, 1012
 vermicularis, 1019
 of atelectasis from pneumonia and pleurisy, 153
 of bronchitis from pneumonia, 209
 from whooping-cough, 209
 of cholera infantum, 452
 of chorea, 625
 of cœcum and appendix, diseases of, 478
 of collapse of the lung, 153
 of contraction with rigidity, 596
 of congestion, cerebral, 539
 of coryza, 57
 of diarrhœa, 393
 of diphtheria, 893
 of dysentery, 463
 of ecthyma, 963
 of eczema, 958
 of eclampsia, 568
 of emphysema, 227
 of entero-colitis, 423
 of erysipelas, 869
 of erythema, 921
 of favus, 987
 of gastritis, 402
 of hemorrhage, cerebral, 547
 of herpes, 953
 of whooping-cough, 267
 of hydrocephalus, chronic, 554
 of impetigo, 965
 of indigestion, 380
 of intussusception, 491
 of laryngitis, simple, without spasm, 65
 simple spasmodic, 75
 of laryngitis, pseudo-membranous, 99
 catarrhal from true croup, 75, 99
 of laryngismus stridulus, 586
 of lichen strophulus, 968
 of malarial fever, 859
 of meningitis, simple, 534
 tubercular, 513
 epidemic cerebro-spinal, 917
 of miliaria, 956
 of mumps, 864
 of night terrors, 662
 of paralysis, atrophic infantile, 645
 facial, 651
- Diagnosis of paralysis, pseudo-hypertrophic muscular, 656
 of pemphigus, 958
 of pharyngitis, simple, 371
 of pleurisy, 240
 of pneumonia, 179-182
 of pneumothorax, 258
 of progressive muscular sclerosis, 656
 of prurigo, 970
 of psoriasis, 972
 of rheumatism, 668
 of rickets, 701
 of roseola, 727
 of rôtheln, 856
 of rubeola, 846
 of rupia, 961
 of scabies, 999
 of scarlatina, 805
 of sclerema, 978
 of scrofula, 676
 of stomatitis, ulcerative, 334
 of syphilis, congenital, 713
 of tetanus nascentium, 607
 of thrush, 360
 of tinea tonsurans, 992
 of tinea circinata, 993
 of tonsillitis, 364
 of tuberculosis, 690
 of mesenteric glands, 690
 of peritoneum, 690
 of typhoid fever, 725
 of urticaria, 901
 of vaccinia, 755
 of variola, 741
 of varicella, 767
- Diaphoretics (*see* Formulæ).
 hot bath as, in scarlatinous dropsy, 812
- Diarrhœa in thrush, 355, 356
 simple or catarrhal, article on, 387-397
 nature of, 387
 causes of, 388
 improper diet as a cause of, 388
 anatomical lesions in, 389
 symptoms of, 391
 course of, 392
 diagnosis of, 393
 prognosis in, 393
 treatment of, 393
 of chronic form, 395
 inflammatory (*see* Entero-colitis), 404
 in rickets, 701
 in tuberculous peritonitis, 688
 in scarlatina, 801
 in measles, 853
 in variola, 747
 in typhoid fever, 716, 720, 726
- Diday, on infantile syphilis, 707
- Diet (*see also* Food) in pneumonia, 192
 after tracheotomy, 127
 in chronic bronchitis, 211
 in pleurisy, 242
 in thrush, 362
 after premature weaning, 362
 in indigestion, 384
 improper, as cause of indigestion, 377
 as cause of diarrhœa, 388
 proper in diarrhœa, 393

- Diet in gastritis, 402
 in acute entero-colitis, 426, 435, 440
 in chronic entero-colitis, 436
 improper, as cause of cholera infantum, 443, 454, 458
 suitable for children, 301-328
 in tubercular meningitis, 525
 in laryngismus stridulus, 588
 in diphtheria, 907
 in rheumatism, 671
 in rickets, 705
 in tuberculosis, 693
 in congenital syphilis, 713
 in scarlatina, 812, 824
 in rubeola, 850
 in variola, 746
 in typhoid fever, 729
 in scrofula, 677
 in erysipelas, 871
 as cause of urticaria, 929
 in urticaria, 931
 in rupia, 960
 in ecthyma, 964
 in ascaris vermicularis, 1018
- Digestive organs, diseases of, 301-497
 disturbances in early stage of rickets, 696
 in tuberculous peritonitis, 688
 in tuberculosis of mesenteric glands, 690
 in variola, 736
 in typhoid fever, 721
 in erythema intertrigo, 921
 caused by worms, 1010
- Digitalis in heart disease, 299
 in scarlatinous dropsy, 829
- Dilatation of bronchi, 200
- Diphtheria, article on, 873-908
 definition and synonyms of, 973
 history of, 873
 statistics of frequency of, 874
 causes of, 875
 epidemic, contagious, and infectious nature of, 875
 influence of season upon, 878
 table showing monthly mortality of, 878
 influence of age upon, less than in croup, 878
 nature of; a constitutional disease, 878
 pathological anatomy of, 879-884
 development of false membranes in, 879
 color and consistence of false membranes, 881
 microscopic anatomy of false membranes, 881
 parasites in the exudation of, 880
 chemical characters of false membranes in, 881
 condition of mucous membrane in, 881
 lesions in croup following, 882
 seat of exudation in, 882
 exudation on skin in, 883
 condition of submaxillary glands in, 883
 fatty degeneration of heart in, 883
- Diphtheria, condition of kidneys in, 884
 lesions in secondary form, 884
 forms of, 884
 symptoms of, 885
 condition of throat in, 885
 difficulty in deglutition not constant, 886
 danger of exudation extending to larynx, 886
 symptoms of croup in, 887
 nasal variety of, 888
 cutaneous symptoms of, 889
 invasion often insidious, 889
 general symptoms of mild form, 890
 of severe form, 890
 urine in, 891
 eruption in, 891
 course in fatal cases, 891
 malignant symptoms in, 892
 duration of, 892
 prognosis in, 893
 diagnosis of, 893
 from scarlatina, 894
 albuminuria in, 895
 heart-clot in, 896
 Richardson's account of symptoms of, 896
 Robinson on, 896
 cases of, 896
 endocarditis in, 879
 paralysis following, 898
 order of muscles affected in, 898
 motion and sensation both affected, 899
 result usually favorable, 899
 explanation of, 900
 locomotor ataxia following, 900
 treatment of, 901
 local applications to throat in, 901
 solvents for the false membranes in, 902
 use of gargles in, 904
 local use of ice in, 904
 external applications in, 905
 injections in nasal form of, 905
 general treatment of, 905
 mercurials in, 906
 emetics and purgatives in, 907
 necessity for supporting remedies in, 907
 stimulants in, 907
 diet in, 907
 necessity for absolute rest, 907
 treatment of the paralysis after, 908
 of heart-clots, 908
- Diuretics (*see* Formulæ), in scarlatinous dropsy, 829
- Dress, suitable for children, 84, 426, 454
- Drinks, manner of taking, as a diagnostic sign, 49
- Dropsy, after scarlatina (*see* article on Scarlatina), 794
 treatment of, 760-763
 after measles, 776
- Duchenne, on progressive paralysis, 630
- Dysentery, article on, 462-469
 definition of, 462

- Dysentery, causes of, 462
 anatomical lesions in, 462
 symptoms of, 463
 diagnosis of, 463
 prognosis in, 463
 treatment of, 464
- Earache, violent crying in, 25
- Eclampsia (*see* Convulsions), 559
- Ecthyma, article on, 962-964
 definition, synonyms, varieties, 962
 causes of, 962
 symptoms of, 962
 diagnosis of, from rupia, 961, 963
 prognosis in, 963
 general treatment of, 963
 local treatment of, 964
 in syphilis, 709
- Eczema, article on, 933-949
 definition of, 933
 elementary lesions in, 933
 eruption frequently mixed in, 933
 seats of eruption in, 933
 forms of, 933
 causes of, 934
 anatomical lesions in, 935
eczema simplex or *vesiculosum*.
 symptoms of, 945
eczema papulosum.
 symptoms of, 936
eczema pustulosum or *impetiginoides*.
 symptoms of, 936
eczema capitis.
 symptoms of, 937
 chronic form of, 938
 condition of scalp in, 938
eczema faciei.
 symptoms of, 939
eczema larvale.
 symptoms of, 940
 duration of, 940
eczema granulosum, 940
eczema tarsi.
 symptoms of, 940
eczema chronicum.
 common to all varieties, 940
 symptoms of, 941
 seats of, 941
 diagnosis of, from scarlatina, 941
 from erysipelas, 941
 of *eczema simplex*, from scabies, 941
 of *eczema vesiculosum*, from sudamina, 941
 of *eczema chronicum*, from psoriasis, 942
 of *eczema impetiginosum*, from favus, 942
 of *eczema squamosa*, from *tinea circinata*, 942
 prognosis in, 942
 treatment of, 943
 principles of, 943
 general treatment in, 943
 attention to digestive symptoms in, 943
 use of arsenic in, 944
 of cod-liver oil in, 944
- Eczema, use of iron in, 944
 of calomel in, 945
 local treatment of, 945
 mode of removing crusts in, 945
 cool and emollient applications in, 945
 benzoated oxide of zinc ointment in, 946
 lotions in, 947
 ointments in, 946
 spiritus saponatus kalinus of Hebra in, 948
 solutions of potash in, 949
 tarry applications in, 947
 use of soaps in, 948
 mercurial applications in, 947
- Electrical batteries, 649
- Electricity in chorea, 631
 in infantile paralysis, 648
 in facial paralysis, 652
 in progressive paralysis, 660
 in diphtheritic paralysis, 907
- Electro-muscular contractility in infantile palsy, 616
 in facial palsy, 651
 in progressive palsy, 655
- Embolism in endocarditis as cause of chorea, 292, 624
 as a cause of atrophic infantile paralysis, 636
- Emetics in catarrhal croup, 80
 in membranous croup, 102
 in collapse of the lung, 155
 in bronchitis, 213
 in whooping-cough, 273
 in eclampsia, 572
 in diphtheria, 907
- Emphysema, time and cause of existence, 218
 anatomical appearances, 218
 vesicular, 218
 interlobular, 219
 causing pneumothorax, 256
 case of, 220
 causes of, age, previous disease, etc., 221
 mechanism of production, 221
 symptoms, rational, 333
 physical signs, 224
 case of, 226
 diagnosis of, 227
 prognosis in, 227
 treatment of, 228
 association with pneumonia, 165
- Empyema, symptoms and course of (*see* Pleurisy), 239
 paracentesis in, 250
- Empyreumatic oil of Chabert as a vermifuge, 1017
- Endocarditis, in diphtheria, 897
 in scarlatina, 802
 acute, 291
 symptoms of, 292
 prognosis in, 292
 anatomical appearances in, 292
 embolism in, in connection with chorea, 292

- Endocarditis, acute, treatment of, 292
- Enemata, astringent, in chronic enterocolitis, 438
- in dysentery, 467
- in typhlitis, 480
- in disease of cæcum, 480
- of air and fluids in intussusception, 494
- in eclampsia, 573
- in diphtheria, 973
- in treatment of worms, 1020
- Enteritis in measles, 843
- in variola, 748
- Enterocolitis, article on, 404-441
- definition of, 404
- frequency of, 404
- improper food and intense heat as causes of, 408
- table of mortality in, 406
- analogies to camp diarrhoea, 411
- dentition and weaning as causes of, 412
- anatomical lesions in, 412
- seat of disease in, 412
- condition of intestinal follicles in, 414
- lesions in chronic form, 415
- microscopic changes in intestines, 416
- condition of stomach in, 417
- of liver in, 417
- of mesenteric and mesocolic glands in, 417
- pathology of, 418
- symptoms of in acute form, 418
- condition of stools in acute form, 419
- of abdomen in acute form, 420
- vomiting in acute form, 420
- erythema of buttocks in acute form, 421
- duration of acute form, 421
- symptoms of chronic form, 422
- course and duration of chronic form, 422
- diagnosis of, 423
- prognosis in, 423
- treatment of acute form, 423-434
- prophylactic, 423
- necessity for change of residence, 424
- importance of exercise in open air, 425
- diet in, 426
- therapeutical treatment of, 426
- cold water in treatment of, 427
- use of calomel in, 427
- use of opium in, 429
- use of astringents and absorbents in, 430
- of tonics and stimulants in, 432
- remedies for vomiting in, 432
- treatment of chronic form, 434
- diet in chronic form, 435
- use of raw meat in chronic form, 435
- nitrate of silver in, 437
- astringent enemata in, 438
- creasote in, 439
- nitrate of iron in, 439
- sulphuric acid in, 439
- tonics and stimulants in, 440
- Epidemic nature of diphtheria, 875
- Epidemic nature of mumps, 862
- of roseola, 926
- of rubeola, 832
- of scarlatina, 772
- of typhoid fever, 716
- of variola, 729
- of varicella, 766
- Epistaxis in coryza, 57
- in measles, 837
- in bronchial phthisis, 684
- in typhoid fever, 723
- Ergot, in infantile paralysis, 647
- in tubercular meningitis, 524
- in epidemic cerebro-spinal meningitis, 917
- Eruption in cerebro-spinal meningitis, epidemic, 915
- in diphtheria, 891
- in ecthyma, 962
- in eczema, 935-941
- in erysipelas, 867
- in erythema intertrigo, 920
- fugax, 921
- nodosum, 922
- in favus, 985
- in herpes, 950, 952
- in ichthyosis, 975
- in impetigo contagiosa, 965
- in lichen strophulus, 962
- in measles, 836-840
- in miliaria, 966
- in pemphigus, 957
- in pityriasis, 973
- in prurigo, 969
- in psoriasis, 971
- in roseola, 926
- in rōtheln, 856
- in rupia, 960
- in scabies, 998
- in scarlatina, 777
- in sclerema, 977
- in tinea, 991
- in typhoid fever, 718-722
- in urticaria, 930
- in vaccinia, 752
- in varicella, 767
- in variola, 733
- in varioloid, 739
- Eruptive fevers, 715
- Erysipelas, article on, 866-873
- definition and forms of, 866
- frequency of, 866
- causes of, 866
- following vaccination, 866
- epidemic and endemic nature of, 867
- symptoms of, 867
- starting-point of eruption, 867
- characters in very young infants, 867
- sloughing of skin in, 867
- characters of in older children, 868
- desquamation in, 868
- abscesses following, 868
- febrile symptoms in, 869
- typhoid symptoms in, 869
- duration of, 869
- diagnosis of, 869
- prognosis in, 869

- Erysipelas, treatment of in young infants, 870
 local applications in, 870
 tr. ferri. chlor. in, 871
 treatment of in older children, 872
 stimulants and tonics in, 872
 local applications in, 872
- Erythema about anus in thrush, 822
 of buttocks in entero-colitis, 421, 921
 in congenital syphilis, 908
 in thrush, 357
 article on, 920-925
 definition and forms, 920
- intertrigo.*
 seat of eruption, 920
 character of eruption, 920
 ulceration in, 921
 form of, occurring in connection with diarrhoea, 921
 diagnosis, 923
 prognosis in, 923
 treatment of, 924
 local applications in, 924
 attention to digestive derangement in, 924
- fugax.*
 symptoms of, 921
 diagnosis of from scarlatina, 921
 from erysipelas, 923
 from roseola, 923
 treatment of, 925
- nodosum.*
 symptoms of, 922
 diagnosis of, from phlegmonous erysipelas, 923
 prognosis in, 923
 treatment of, 925
 tonics and stimulants in, 925
 local applications in, 925
- Escharotics, in gangrenous stomatitis, 343
- Eserine, in chorea, 629
- Essay, introductory, 17
- Essential convulsions (*see* Convulsions).
- Essera (*see* Urticaria), 929
- Examination, clinical, in children, 17
 difficulties of, 17-19
 of abdomen, 46
 of the heart, 36
 of lungs, 43
 of mouth and fauces, 47
 of the pulse, 33-35
- Exercise in open air, importance of, 425
- Expectoration, nummular, in measles, 838
 in true croup, 97
 in bronchitis, 207
 in pneumonia, 175
- Expiratory respiration in bronchitis, 209
- External applications (*see* Local applications).
- Facial paralysis (*see* Paralysis).
- Facies (*see* Physiognomy).
- False membrane, in croup, characters of, 92-94
 in secondary croup, 95
 extent of and frequency with which bronchi are invaded, 94
- False membrane in diphtheria, 879
 action of chemical reagents upon, 881
 in capillary bronchitis, 199
 in scarlatina, 786
- Fauces, examination of, 47, 48
 in membranous croup, 93, 100
 in simple pharyngitis, 369
 in diphtheria, 885
 in scarlatina, 777-787
 in measles, 835, 836
- Favus, article on, 983-990
 definition and synonyms, 984
 varieties and frequency of, 984
 description of the fungus which causes it, 984
 contagion as cause of, 984
 occurs in lower animals, 984
 other causes of, 984
- dispersus*, 985
 symptoms of, 985
 condition of skin in, 985
 appearance of crusts, 985
 alopecia, 986
 seat of eruption, 986
 nature of, 986
 diagnosis of, 987
 prognosis in, 987
 general treatment of, 988
 local treatment of, 988
 mode of removing crusts in, 988
 depilation in, 989
 parasitocides in, 990
- confertus.*
 eruption in, 986
 course of, 986
 condition of hairs in, 986
 baldness following, 986
- Fecal accumulation in cæcum, symptoms of, 475
 diagnosis of, 492
 as cause of eclampsia, 576
- Fever, eruptive, 715
 scarlet (*see* Scarlatina), 769
 in variola, subsides on appearance of eruption, 734
 secondary, in variola, 735
 typhoid, 715
 febrile action in, 720
- Finlayson, normal temperature in children, 38
- Follicles of intestine in entero-colitis, 413, 414
- Fomites as means of transmitting scarlatina, 773
 variola, 731
- Food, article on, 301-328
 artificial, 302-307
 importance of a knowledge of, 302
 improper, as cause of indigestion, 377
 of diarrhoea, 388
 of entero-colitis, 408
 of laryngismus stridulus, 578
 milk, *goat's*, 302
 cow's, 303
 plan of obtaining it fresh, 303
 method of preserving it, 303, 307

- Food, milk, reaction of, 303, 305
 composition of, 303
 proportion of cream in, 304, 305
 methods of estimating quality of, 305
 proportion of water in, 306
 specific gravity of, 306
 adulteration of, 306
 its use in artificial or hand-feeding, 307
 principles to be observed in prescribing it, 308
woman's, 308
 only proper food for infants, 302
 composition of, 308
 proper dilution of cow's milk, for infants, 310
 quantity required by infants, 310, 316
 method of ascertaining it, 311, 316
 hygienic care of infants in hand-feeding, 316
 manner of feeding, 317
 substitutes for, 318, 319
 formula for gelatin food, 318
 Merei's food, 318
condensed, 320
 method of preparation, 320
 analysis of, 321
 different kinds of, 321
 advantages and disadvantages of, 321
 how it is, and how it should be used, 323
 quantity to be given, 324
 value of, 325
 illustrative cases of its use, 326, 327
- Foot-baths, in treatment of diseases, 191, 212, 852
- Formulary of receipts recommended.
- Alteratives.*
 Formula for mixture of iodide of potassium and bichloride of mercury, 522
 for iodide of potassium and compound decoction of sarsaparilla, 677
 for iodide of potassium, syrup of iodide of iron, and syrup of ginger, 677
- Antacids and alkalies; laxatives.*
 Formula for mixture of soda, rhubarb and paregoric, 363
 of magnesia, and tr. opii, 395
 of sulphate of magnesia and rhubarb, 395
 of sulphate of magnesia and laudanum, 395
 of soda, blue mass, and paregoric, 427
 of crab's eyes, 431
 for neutral mixture, 434
 of acetate and bicarb. of potash and opium, 669
- Antiseptic.*
 Formula for mixture of chlorinated lime, 336
- Anthelmintics.*
 Formula for mixture of ol. chenopodii, 1015
 of spigelia, magnesia, and manna, 1015
 of ol. terebinth., and magnesia, 1016
 of santonin, 1016
- Antispasmodics.*
 Formula for pills of belladonna, opium, and valerian, 271
 for mixture of belladonna and opium, 272
- Astringents.*
 Formula for mixture of alum and honey of roses, 329
 of sulphate of copper and quinia, 373
 for aromatic syrup of galls, 431
 of morph. sulph., and dil. sulph. acid, 434
 of soda, krameria, and opium, 363
 of nitrate of silver, 438
 solution of nitrate of silver, 438
 of pernitrate of iron and nitric acid, 439
 of aromatic opium and krameria, 440
 of acetate of lead and acetic acid, 456
- Febrifuges and diuretics (see also Alkalies).*
 Formula for mixture of citrate of potash, ipecac., and paregoric, 81
 of ipecac., opium, sp. æther. nitrosi, 212
 of morphia, liq. ammon. acet., 214
 of iodide of potassium and sarsaparilla, 243
 of squill and digitalis, 244
 of carbonate of potash, squill, and opium, 67
 senega and opium, 67
 of acetate of potash, digitalis, and squill, 829
 of bitart. of potash, juniper, and sp. ætheris nitrosi, 829
 of powders of sulphurated antimony and Dover's powder, 189
 of opium, ipecac, and nitrate of potash, 251
- Foods.*
 Formula for gelatin food for children, 318
 for Merei's food for children, 319
 of preparation of raw meat, 435
- Laxatives (see also Antacids).*
 Formula for mixtures of sulphate of soda, senna, and laudanum, 480
 for pill of opium and colocynth, 480
 for blue pill, castor oil, and aromatic syrup of rhubarb, 598
- Local applications.*
 Formula for lotion of carbonate of potash and sulphur, 1001
 of soft soap and alcohol (Hebra's sp. saponatus kalinus), 948
 of glycerin and ung. aq. ros., 59
 for benzoated oxide of zinc ointment (Bell's formula), 946

- Local applications.*
 Formula for mercurial ointment to prevent pitting in variola, 750
 for ointment of nitrate of mercury and belladonna, 60
 of protiodide of mercury, 947
 of calomel and camphor, 947
 of ammon. chl. of mercury with sulphur, 994
 of tar and iodine, 994
 of sulphur and carbonate of potash (Wilson), 1001
 of sulphur and carb. potash (Helmreich), 1001
- Nervous sedatives.*
 Formula for mixture of antimony, valerian, and paregoric, 68
 of antimony, valerian, and laudanum, 194
- Nervous stimulants.*
 Formula for mistura indica, 458
- Specific remedies.*
 Formula for mixture of carbonate of potash in whooping-cough, 273
 of alum and conium in whooping-cough, 275
 of alum in whooping-cough, 275
 and belladonna in whooping-cough, 275
- Tonics.*
 Formula for mixture of cod-liver oil, 386
 of nux vomica and gentian, 397
 of tr. ferri chlor., acetic acid, and sp. Mindereri, 824
 of quinia, morphia, and sulph. acid, 670
 of quinia and dil. sulph. acid, 215
 of elix. cinchona and curaçoa, 216
 of arsenic and bitter wine of iron, 944
- Forster, temperature in new-born children, 37
 Fox, Wilson, on softening of stomach, 400
 on use of cold in treatment of hyperpyrexia, 816
 French measles (*see* Roseola).
 Frequency of diseases (*see* Statistics).
 Fungi in skin diseases (*see* Parasite).
 Fungous origin of measles, 833
- Gairdner, collapse of the lung, 144
 lesions in bronchitis, 199
 difference between dyspnoea of pneumonia and bronchitis, 209
 Galls, aromatic syrup of, 431
 Gangrene, of mouth (*see* Stomatitis gangrenosa).
 of pharynx in diphtheria, 886
 in scarlatina, 787
 of skin in erysipelas, 868
 Gargles in diphtheria, 904
 Gastritis, article on, 397-404
 frequency and nature of, 397
 causes of, 398
 anatomical lesions in, 398
 softening of stomach in, 399
 symptoms of, 400
- Gastritis, diagnosis of, 402
 prognosis in, 402
 treatment of, 402
 Gastromalacia (*see* Softening of stomach),
 Gastrotomy in intussusception, 496
 Gelatin food, formula for, 318
 General diseases, introductory remarks upon, 665
 Gerhard, G. S., on chorea, 611
 German measles, 855
 Gestures, significance of, 30
 Glands, bronchial, tuberculosis of, 679, 683
 cervical, in scarlatina, 788
 intestinal, in scarlatina, 803
 in typhoid fever, 716
 mesenteric, tuberculosis of, 682, 689
 parotid, in mumps (*see* Mumps).
 in scarlatina, 788
 submaxillary, in diphtheria, 883
 enlarged in some cases of mumps, 863
 enlarged in scarlatina, 778
 Glottis, spasm of, in eclampsia, 566
 (*see* Laryngismus stridulus), 577
 Gölis on treatment of hydrocephalus, 556
 Gregory, cold affusions in scarlatina, 816
 Guaiac in tonsillitis, 265
 Gum (*see* Strophulus).
 Gummy tumors in congenital syphilis, 712
 Gums, importance of lancing in cholera infantum, 460
 in laryngismus, 587
 Gutta-percha, solution of, to prevent pitting in variola, 751
 Gymnastic exercises in chorea, 632
 in infantile paralysis, 649
- Hæmoptysis in pulmonary phthisis, 686
 Hair, in favus, 986
 in tinea, 992
 in alopecia areata, 996
 Hall, Marshall, on spasm of glottis in eclampsia, 566
 on nature of laryngismus, 579
 Hammond, electrical condition of muscles in infantile paralysis, 640
 Harley, John, on connection of scarlet and enteric fevers, 803
 Harris, R. P., on hereditary nature of eclampsia, 562
 Hassall, analysis of milk, 306, 320
 Head, peculiarities of, in hydrocephalus, 552
 in rickets, 697-699
 soft spots on, in rickets, 867
 Heart, physical examination of, 36
 sounds of, 36
 diseases of, article on, 289-300
 (*see* Pericarditis and Endocarditis.)
 causes of, 289
 chronic valvular diseases of 292
 causes of, 292
 anatomical appearances in, 293
 symptoms of aortic disease, 293
 prognosis in, 295
 symptoms of mitral obstruction, 295

- Heart, mitral obstruction, prognosis in, 295
 symptoms of mitral regurgitation, 296
 prognosis in, 296
 compensation for affected by growing heart, 297
 tendency to improve by time, 297
 treatment of, 299
 illustrative cases of, 297
 in chorea, 615
 irregular action of, in chorea, 620
 fatty degenerations of, in diphtheria, 883
 inflammation of endocardium in diphtheria, 897
 inflammation of membranes of, in scarlatina, 802
- Heart-clot in diphtheria, 896, 908
 in scarlatina, 804
- Heat, intense, as cause of entero-colitis, 408
 as a cause of cholera infantum, 443
- Heat of surface (*see* Temperature).
- Hebra, on treatment of eczema, 946
- Heine and Chevalier, analysis of milk, 306
- Helminthocorton as a vermifuge, 1017
- Hemiplegia in chorea, 620
 in cerebral hemorrhage, 544
 cerebral, diagnosis of from facial paralysis, 651
- Hemorrhage during paroxysms of whooping cough, 261
 intestinal, in intussusception, etc., 493
 in typhoid fever, 724
- Herpes, article on, 949-954
 definition of, 949
 varieties of, 949
 frequency of, 949
 causes of, 949
 diagnosis of, 953
 prognosis in, 953
 general treatment of, 953
 local treatment of, 954
- phlyctenodes*.
 seat of eruption in, 959
 symptoms of, 950
 diagnosis of from pemphigus, 953
 treatment of, 953
- labialis*.
 seat of eruption in, 951
 symptoms of, 951
 local applications in, 954
- zoster*.
 definition of, 951
 seat of, 951
 character of eruption, 952
 course and duration of, 952
 general symptoms in, 952
 pain in, 952
 diagnosis of, 953
 local applications in, 954
- circinatus* (*see* Tinea circinata).
- iris*.
 symptoms of, 952
 seat of, 952
 parasitic nature of, 953
- Herpes, iris, diagnosis of from roseola annulata, 953
 treatment of, 954
- Hewitt, Grailly, on collapse of the lung in whooping-cough, 265
- Hives (*see* Urticaria), 929
- Hillier, indications for bleeding in pneumonia, 185
 on pathology of infantile paralysis, 641
 albuminuria in diphtheria, 895
- Holding-breath spells (*see* Convulsions, internal), 586
- Whooping-cough, article on, 259-280
 definition, synonyms, frequency, 259
 causes, influence of age, 259
 contagion and epidemic influence, 260
 stages of, 260
 symptoms and duration of first stage, 260
 of second stage, 261
 character of paroxysms, 261
 hemorrhages during this stage, 261
 convulsions during this stage, 262
 duration of paroxysms, 262
 number of paroxysms, 262
 symptoms and duration of third stage, 262
 absence of general symptoms in, 263
 urine in, 263
 total duration of, 263
 convulsions as a complication, 263
 excessive laryngismus as a complication, 264
 collapse of the lung as a complication, 265
 bronchitis as a complication, 266
 pneumonia as a complication, 266
 vomiting in, 267
 emphysema as a sequel, 266
 tuberculosis and scrofula as sequelæ, 267
 diagnosis of, from acute catarrh, 267
 from tuberculosis of bronchial glands, 267
 prognosis in, 268
 nature of, 269
 anatomical lesions, 269
 mortality in, 269
 treatment of simple form, 269
 bloodletting in, 271
 belladonna in, 271
 hydrocyanic acid in, 272
 carbonate of potash in, 273
 alum in, 274
 inhalations in, 275
 local applications in, 276
 treatment of complications of, 277
 of paroxysms of, 280
 hygienic treatment of, 279
- Hunger, crying from, 25
- Hunt, S. B., change of residence in chronic diarrhoea, 424
- Hutchinson, J., on transmission of syphilis by vaccination, 760
 alteration of teeth in congenital syphilis, 711

- Hydrencephalic cry, 507
 Hydrocephalus, acute (*see* Tubercular meningitis).
 chronic, following meningeal apoplexy, 546
 article on, 548-558
 forms of, 548
 anatomical appearance in, 549
 analysis of fluid in, 550
 causes of internal form, 550
 external form, 551
 symptoms of, 552
 enlargement of head in, 552
 cerebral symptoms in, 553
 mode of death in, 554
 diagnosis of, from rickets of the skull, 554
 from hypertrophy of the brain, 555
 prognosis in, 555
 treatment of, 556
 use of mercury in, 556
 compression of head in, 557
 paracentesis in, 557
 injections into cranial cavity in, 558
 Hydrochloric acid in diphtheria, 902
 Hydrocyanic acid in whooping-cough, 272
 Hygienic conditions, influence of unfavorable, 444
 treatment in chorea, 633
 in measles, 849
 in scarlatina, 810
 Hypertrophy of tonsils (*see* Tonsils, chronic enlargement of), 366
 Hyposulphites, use of in scarlatina, 823
 Hypertrophic disease of the skin, 974
 Hypertrophy of the heart, 297

 Ice, local use of, in eclampsia, 575
 in tetanus, 609
 in diphtheria, 904
 in scarlatina, 825
 in angina of scarlatina, 825
 Ichthyosis, symptoms of, 975
 etiology of, 975
 pathology of, 975
 treatment of, 975
 Ileus (*see* Intussusception).
 Impetigo (*see* Eczema impetiginoides).
 Impetigo contagiosa,
 article on, 965-966
 definition of, 965
 symptoms of, 965
 causes and pathology of, 965
 diagnosis of, 965
 prognosis in, 966
 treatment of, 966
 larvalis (*see* Eczema larvale).
 granulata (*see* Eczema granulosum).
 figurata, 937
 sparsa, 937
 Incubation of scarlatina, 772
 of measles, 832
 of small pox, 731
 Indigestion, article on, 376-387
 definition, frequency, forms, 376
 causes of, 377
 Indigestion, symptoms of occasional form, 378
 of habitual form, 378
 diagnosis of, 380
 prognosis in, 381
 treatment of occasional form in infants, 381
 in older children, 382
 of habitual form, 383
 as cause of eclampsia, 562
 Infantile paralysis (*see* Paralysis), 634
 Infantile remittent fever (*see* Typhoid fever).
 syphilis (*see* Syphilis).
 Inflammation, catarrhal, of larynx, without spasms, 62
 with spasms, 69
 of larynx with pseudo-membranous exudation, 85
 of lungs, 157
 of bronchi, 195
 of pleura, 231
 of skin, bullous, 956
 catarrhal, 933
 erythematous, 920
 papular, 966
 pustular, 962
 squamous, 971
 vesicular, 933
 Inflation, effect of, on collapsed lung, and in atelectasis, 134, 147
 impossible in pneumonic lung, 163
 in congestion of lung, 162
 Inhalations in treatment of membranous croup, 106
 of coal-gas in whooping-cough, 275
 Injections (*see* Enemata).
 into cranial cavity in hydrocephalus, 558
 into pleural sac after paracentesis, 250
 in nasal diphtheria, 905
 of air in intussusception, 495
 Inoculability of scarlatina, 773
 Inspection of thorax in pleurisy, 235
 Inspiration, recession of base of thorax in, 41
 Internal convulsions (*see* Convulsions and Laryngismus).
 Intertrigo (*see* Erythema), 921
 Intestines, inflammation of, in measles, 843
 and stomach, general remarks upon diseases of, 376
 Intussusception, article on, 481-497
 definition, synonyms, and forms, 481
 frequency, 482
 anatomical appearances, 483
 divisions of, 483
 most frequent seat of, 483
 pathology of, 483
 modes of termination of, 484
 case of elimination of the invaginated bowel, 485
 causes of, 486
 mode of production, 486
 symptoms of, 487
 duration of, 490

- Intussusception, modes of termination of, 490
 prognosis in, 491
 diagnosis of, 491
 differential diagnosis of, 492
 treatment of, medical, mechanical, and surgical, 493
 use of purgatives in, 494
 injections of air and fluids in, 494
 hydrostatic pressure in treatment of, 495
 gastrotomy in, 496
- Inunction in scarlatina, 813
 of cod-liver oil in rickets, 705
 in tuberculosis, 694
- Invagination (*see* Intussusception), 481
- Invasion of diphtheria often insidious, 889
 of mild cases of scarlatina, 776
 of grave cases of scarlatina, 785
 of measles, 833
- Iodide of potassium (*see* Potassium).
- Iodine in scrofula, 678
 in tubercular meningitis, 521
 as injection in chronic hydrocephalus, 558
- Iron in laryngismus stridulus, 590
 in infantile paralysis, 647
 in rheumatism, 670
 in eczema, 944
 local use of, in diphtheria, 903
 in tuberculosis, 694
 chloride of, with acetic acid and sp. Mindereri, 440
 in diphtheria, 903
 in scarlatinous angina, 826
 in erysipelas, 871
 iodide of, in pleurisy, 243
 in chronic eczema, 944
 nitrate of, in chronic entero-colitis, 434
 enema of, in dysentery, 466
- Isambert, ulceration of larynx in diphtheritic croup, 86
- Itch (*see* Scabies), 997
- Jaborandi, in mumps, 865
- Jaccoud, diphtheritic ataxia, 901
- Jacobi, statistics of tracheotomy in croup in New York, 110, 118
 local applications in diphtheria, 903
- Jenner, recession of base of chest in inspiration, 41
 on rickets, 694, 698, 701
- Joints, condition of, in rheumatism, 666
 affections of, in scarlatina, 801
- Kameela as a vermifuge, 1016
- Keratitis, in congenital syphilis, 712
- Kidneys, condition of, in diphtheria, 884
 in scarlatinous dropsy, 795
 in thrush, 352
- Kine-pock (*see* Vaccine disease).
- Klein, condition of kidneys in scarlatina, 803
- Laborde, sclerosis of spinal cord in infantile palsy, 644
- Lancing gums (*see* Gums).
- Lactometer, use of, 306
- Laryngismus, excessive in whooping-cough, 264
 in eclampsia, 566
 in rickets, 701
 stridulus, article on, 571-593
 definition and synonyms, 577
 frequency of, 577
 predisposing causes of, 577
 nature and exciting causes of, 579
 anatomical appearances in, 579
 enlargement of thymus gland in, 580
 centric and eccentric causes of, 581
 symptoms of paroxysm of, 582
 duration and course of, 584
 danger of sudden death in, 584
 other forms of, 585
 holding-breath spells, 586
 diagnosis of, 586
 prognosis in, 587
 treatment of, 587
 importance of lancing gums in, 587
 attention to diet in, 588
 antispasmodics in, 589
 iron in, 590
 treatment of paroxysms, 590
 change of residence in, 590
 illustrative cases, 591
- Laryngitis, in scarlatina, 792
 in measles, 843, 853
 in small-pox, 734
 chronic, 67
 pseudo-membranous, 85
 definition, synonyms of, 85
 nature and relations to diphtheria, 85-89
 morbid process in pseudo-membranous laryngitis and diphtheria, 86
 age in, 87
 condition of cervical glands in, 88
 type of general symptoms in, 88
 complications in, 89
 frequency of faucial deposit in, 89
 frequency of, 89
 mortality from, 90
 predisposing causes of, 91
 exciting causes of, 92
 second attack of, 92
 anatomical lesions in, 92
 extent and seat of false membranes, 92
 appearance of fauces in, 93
 characters of membrane in, 94
 symptoms of, 95
 of initial stage, 95
 characters of voice and cough, 95
 of respiration, 96
 explanation of recession of base of chest in, 97
 expectoration and rejection of false membrane in, 97

- Laryngitis, pseudo-membranous, negative
 results of auscultation in, 98
 mode of recovery in, 99
 duration of, 99
 diagnosis of, 99
 importance of examination of
 throat in, 100
 prognosis in, 100
 treatment of, 100
 bloodletting in, 101
 emetics in, 102
 antimony in, 103
 mercury in, 104
 alum and ipecacuanha in, 103
 alkalies in, 105
 opium in, 105
 local treatment in, 105
 inhalations in, 106
 hygienic treatment, 107
 summary of the treatment of, 107
 tracheotomy (*see* under that
 head), 107
spasmodic, simple, 69-84
 forms of, 69
 synonyms of, 69
 causes of, 69
 anatomical lesions in, 70
 symptoms of, 71
 duration of, 73
 nature of, 74
 diagnosis of from true croup, 75
 from laryngismus stridulus,
 586
 peculiarities of voice in, 77
 prognosis in, 78
 treatment of the mild form of, 78
 of the severe form of, 78
 hygienic treatment of, 83
 prophylactic treatment of, 83
 style of dress suitable in, 84
simple, without spasms, 62
 definition and frequency of, 62
 causes of, 62
 anatomical lesions in, 62
 symptoms and course of, 63
 duration of, 64
 diagnosis of, 65
 prognosis in, 66
 treatment of, 66
- Larynx, general remarks on diseases of, 61
 Legendre and Bailly, researches on col-
 lapse of lung, 144
- Lichen, article on, 966-968
 definition of, 966
 varieties of, 966
 simplex, ruber, and scrofulosus, 967
 symptoms of, 966
 causes of, 967
 strophulus, varieties and symptoms of,
 967
 causes and pathology of, 968
 diagnosis of, 968
 prognosis in, 968
 treatment of, 969
 intertinctus, confertus, volaticus, al-
 bidus, candidus, 968
 tropicus, 955
- Liver, state of, in entero-colitis, 417
 in rickets, 701
 in congenital syphilis, 712
- Lobular pneumonia, in reality collapse of
 lung, 144
- Local applications in alopecia areata,
 997
 in aphthæ, 331
 in bronchitis, 217
 in chorea, 631
 in congenital syphilis, 713
 in coryza, 59
 in diphtheria, 901
 in cæcum and appendix, diseases of,
 481
 in eclampsia, 575
 in ecthyma, 964
 in eczema, 945
 in erysipelas, 870, 872
 in erythema, 924
 in favus, 988
 in gangrene of the mouth, 343
 in herpes, 954
 in hooping-cough, 276
 in ichthyosis, 975
 in impetigo contagiosa, 966
 in lichen, 967, 969
 in meningitis simple, 535
 tubercular, 525
 in membranous croup, 105
 in miliaria, 956
 in mumps, 865
 in pemphigus, 959, 962
 in pityriasis, 974
 in pleurisy, 244
 in pneumonia, 191
 in prurigo, 971
 in psoriasis, 973
 in rheumatism, 671
 in rupia, 961
 in scabies, 1001
 in scarlatina, 813
 in strophulus, 969
 in tinea, 993
 in thrush, 363
 in ulcero-membranous stomatitis, 335
 in urticaria, 932
 in variola, 749
- Locomotor ataxia following diphtheria,
 900
- Lotions (*see* Formulæ).
 of water in scarlatina, 815-822
 in treatment of eczema, 947
- Lumbricus (*see* Ascaris lumbricoides).
- Lungs, auscultation of, 44
 general remarks on diseases of, 133
 collapse and imperfect expansion of,
 143
 inflammation of, 157
 abscess of, following pneumonia, 162
 congestion of, non-inflammatory, 163
 condition of, in pleurisy, carnification,
 233
 perforation of, 257
 collapse of, in rickets, 703
 tuberculosis of, 680, 685
 percussion of, 43, 44, 45

- Lungs, condition of, in congenital syphilis, 712
 in typhoid fever, 719, 722
 in sclerema, 977
- Mackenzie, paralysis in croup, 89
 anatomical lesions in croup, 93
- Magnesia, hyposulphite of, in scarlatina, 823
- Malarial fever, article on, 857-861
 causes, frequency of, 857
 symptoms of acute, 857
 features of paroxysm imperfectly developed, 858
 of chronic, 859
 characters of blood in, 859
 enlargement of spleen in, 859
 neuralgia rare in, 859
 diagnosis of, 859
 prognosis in, 859
 treatment of, 860
 quinia in, 860
 iron and arsenic in, 860
- Marsh, F. H., on tracheotomy in croup, 119
- Maturative fever in variola, 735
- Maw-worm (*see* *Ascaris vermicularis*).
- Malleolus, ulceration of, in thrush, 357
- Measles (*see* *Rubeola*), 834
 French, 925
 German, 855
- Meigs, Charles D., treatment of coryza, 58
 use of alum as an emetic, 103
 on proper position of body in atelektasis and cyanosis, 142, 288
 treatment of paroxysm of laryngismus, 590
- Meigs, J. F., case of contraction with rigidity, 597
 heart-clot in diphtheria, 896
- Membrane, false (*see* *False membrane*).
- Meningeal apoplexy (*see* *Cerebral hemorrhage*), 540
- Meningitis, simulated by cerebral form of pneumonia, 180
epidemic cerebro-spinal, article on, 908-912
 definition, synonyms, history of, 908
 frequency of, 908
 causes of, 910
 anatomical lesions of, 910
 symptoms and course of, 911
 fever, headache, vomiting in, 911
 case of, 912
 paralysis following, 914
 eruptions in, 915
 duration of, 915
 diagnosis of, 917
 prognosis in, 916
 treatment of, 917
simple, article on, 529-536
 definition, synonyms, frequency, 529
 causes of, 529
 anatomical lesions in, 530
 symptoms of convulsive form, 531
 of phrenitic form, 532
 course and duration of, 533
- Meningitis, diagnosis of, from congestion of brain, 534
 differential diagnosis of, 534
 from tubercular form, 514
 prognosis in, 534
 treatment of, 534
 bleeding in, 535
 calomel in, 535
 cold and counter-irritation in, 535
tubercular, article on, 498-529
 definition, synonyms, and frequency, 498
 predisposing causes of, 500
 exciting causes of, 500
 pathology of, 500
 anatomical lesions in, 501
 microscopical changes in, 501
 division into stages, 504
 mode of invasion, 505
 symptoms of first stage, 506
 hydrocephalic cry, 507
 condition of mind in, 507
 use of ophthalmoscope in, 509
 convulsions in, 509
 circulation in, 510
 symptoms of second stage, 510
 tache méningitique, 511
 nervous symptoms in, 511
 decubitus in, 512
 pulse in, 512
 respiration in, 512
 temperature in, 513
 diagnosis of, from simple meningitis, 513
 from typhoid fever, 515
 cases simulating, 516
 prognosis in, 516
 cases of recovery from, 517
 case of apparent recovery from, 518
 prognosis not absolutely hopeless, 519
 uncertainty of date of death in, 519
 treatment of, 520
 bleeding not to be used, 520
 iodine and iodide of potassium in, 521
 bromide of potassium in, 522
 ergot in, 524
 counter-irritation in, 525
 cold applications in, 525
 calomel and mercury in, 523
 prophylaxis in, 526
 narcotics in, 526
 diet in, 526
 importance of country residence, 527
 illustrative case, 528
- Mental condition in rickets, 697
- Mercurial ointment to prevent pitting in variola, 750
 in congenital syphilis, 713
 applications in eczema, 948
 in favus, 990
 in tinea, 993
- Mercury (*see* *Calomel*).
 in membranous croup, 104
 in pneumonia, 189
 in pleurisy, 242
 in entero-colitis, 403

- Mercury in diseases of cœcum and appendix, 480
 in tubercular meningitis, 523
 in simple meningitis, 535
 in chronic hydrocephalus, 556
 in scrofula, 678
 in congenital syphilis, 713
 in diphtheria, 906
- Merei's food, 318
- Mesenteric glands, tuberculosis of, 682, 689
 condition of entero-colitis, 417
- Metastasis in mumps, 864
- Microscopic examination of false membranes in diphtheria, 879
 changes in tubercular meningitis, 501
 in spinal cord in tetanus, 605
 in chorea, 617
 in infantile paralysis, 642
 in muscle in infantile paralysis, 641
 in progressive paralysis, 658
 in kidneys in scarlatinous dropsy, 795, 796
- Microsporon furfur, 982
- Miliaria, article on, 955-956
 definition of, 955
 frequency of, 955
 cause of, 955
 symptoms of, 955
 duration of, 955
 diagnosis of, 956
 treatment of, 956
- Miliary tubercles, 501
- Milk (*see also* Food and Diet).
- Milk-crust (*see* Eczema capitis), 957
- Mineral acids, use of in typhoid fever, 728
- Mitchell, S. W., observations on chorea, 611, 614
- Mitral valve, disease of, 293
- Morbilli (*see* Rubeola), 834
- Mortality (*see* Statistics).
- Mouth, examination of, 47
 mode of examining, 48
 diseases of (*see* Stomatitis), 301
- Mucous membrane, condition in entero-colitis, 414
 of fauces in diphtheria, 885
 affections of, in congenital syphilis, 709
 of fauces, in scarlatina, 709
 gastro-intestinal, in scarlatina, 803
 eruption on, in small-pox, 734
 condition of, in small-pox, 737
 in cases of ascaris lumbricoides, 1008
- Muguet (*see* Thrush).
- Mumps, article on, 861-865
 definition, synonyms, and frequency, of, 861
 causes of, 861
 anatomical appearances in, 862
 symptoms of, 862
 characters of swelling in, 862
 salivary secretion in, 863
 general symptoms in, 864
 tendency to metastasis in, 864
 prognosis always favorable in, 864
- Mumps, course and duration of, 864
 usually terminates by resolution, 864
 suppuration of parotid in, 864
 diagnosis of, 864
 treatment of, 865
 danger of febrile sequelæ in, 865
- Muriate of ammonia (*see* Ammonia).
- Muriatic acid (*see* Acid).
- Murmur, cardiac, in chorea, 620
 cerebral, in rickets, 697
- Muscle, condition of, in atrophic infantile paralysis, 637, 638
 in progressive paralysis, 654, 658
- Muscles, paralyzed in diphtheria, 898
- Muscular sclerosis, progressive (*see* Paralysis), 652
- Narcotics in tetanus nascentium, 609
- Nasal variety of diphtheria, 888
- Nature of alopecia areata, 996
 of aphthæ, 330
 of cerebral congestion, 537
 of chorea, 621
 of cholera infantum, 445
 of collapse of the lung, 143
 of contraction with rigidity, 594
 of cyanosis, 285
 of cœcum and appendix, diseases of, 468
 of diarrhœa, 387
 of diphtheria, 875, 878
 of eclampsia, 566
 of emphysema, 218
 of entero-colitis, 418
 of erysipelas, 867
 of favus, 987
 of gastritis, 397
 of herpes, 949
 of hooping-cough, 269
 of hydrocephalus, 549
 of ichthyosis, 975
 of laryngitis, spasmodic, 74
 pseudo-membranous and diphtheria, 85-89
 of laryngismus stridulus, 579
 of mumps, 861
 of night terrors, 662
 of paralysis, atrophic infantile, 641
 of parasitic skin diseases, 980
 of pneumonia, 158
 of pneumothorax, 253
 of progressive muscular sclerosis, 658
 of prurigo, 969
 of rickets, 704
 of rœtheln, 855
 of rubeola, 832
 of scabies, 987
 of scarlatina, 773
 of thrush, 358
 of tonsillitis, 364
 of typhoid fever, 716
 of varicella, 766
 of variola, 729
- Nephritis, after scarlatina, 795
- Nervous system, general remarks on diseases of, 498
 symptoms (*see* Cerebral symptoms).

- Nervous system, influence of in eczema, 935
 Nettle-rash (*see* Urticaria), 929
 Neuralgia following malaria, 859
 Neurosis, evidence in favor of whooping-cough being a, 269
 evidence of laryngismus stridulus being a, 579
 infantile paralysis not a, 642
 Niemeyer, indications for bleeding in pneumonia, 187
 on lesions in catarrh of stomach, 400
 Night terrors, article on, 660-664
 definition of, 660
 cases of, 660
 pathology of, 662
 causes of, 664
 treatment of, 664
 Notha, or rubeola sine catarrho, 840
 Obstruction of the intestines (*see* Intussusception), 481
 Occipital bone, depression of, as cause of tetanus, 602
 Œdema of face in bronchial phthisis, 684
 of neck in scarlatina, 788
 in variola, 733
 in typhoid fever, 724
 in erysipelas, 869
 in sclerema, 977
 Ogle, J. W., on chorea, 613
 Oidium albicans (*see* Thrush), 347, 350, 353, 358
 Ointments (*see* Local applications).
 in treatment of seatworms, 1020
 Omentum, tuberculosis of, 681, 689
 Ophthalmia, in variola, 738
 Ophthalmoscope, use of in tubercular meningitis, 509
 Opisthotonos in tetanus nascentium, 606
 Opium in catarrhal croup, 80, 82
 in membranous croup, 105
 in pneumonia, doses and modes of administering, 193
 in pleurisy, 242
 in thrush, 363
 in entero-colitis, 429
 in cholera infantum, 456
 in dysentery, 464
 in diseases of cœcum and appendix, 481
 in eclampsia, 574
 in rheumatism, 670
 in variola, 747
 in typhoid fever, 728
 Otorrhœa, in scarlatina, 789; 801
 Ova of *acarus scabiei*, 999
 Oxyuris vermicularis (*see* *Ascaris vermicularis*), 1004
 Packing, cold, in scarlatina, 815
 Pain, modes of expressing, 22
 in pneumonia, seats and peculiarities of, 167, 176
 in pleurisy, seats and peculiarities of, 236
 abdominal, in intussusception, 488
 in tuberculous peritonitis, 688
 Pain in variola, 730
 in herpes zoster, 953
 Palate, papules on, in measles, 836
 Palpation of thorax in pleurisy, 235
 Pancoast, J., on tracheotomy in croup, 112
 Papules, chapter on, 966
 in small-pox, 732
 Paracentesis, in pleurisy (*see* Pleurisy), 244-252
 case of, 251
 in hydrocephalus, 557
 Paralysis, in tubercular meningitis, 507
 in cerebral hemorrhage, 545
 in chorea, 620
 in diphtheria, 898
 after scarlatina, 802
 in epidemic cerebro-spinal meningitis, 914
 atrophic infantile, article on, 634-650
 history and authors on, 634
 synonyms of, 635
 causes of, 636
 forms of, and muscles affected in, 637
 mode of attack, 636
 reaches its maximum suddenly, 637
 condition of paralyzed muscles in, 638
 illustrative case of, 638
 at times temporary, 638
 at other times followed by atrophy, 638
 temperature lowered in palsied parts, 639
 subsequent deformities in, 639
 duration of, 640
 prognosis in, 640
 electrical condition of muscles as aid in prognosis, 640
 microscopic examination of muscles in, 641
 anatomical lesions in and nature of, 641
 cannot be considered reflex, 642
 primary condition usually one of spinal congestion, 643
 sclerosis of cord in a later stage, 643
 other lesions of cord occasionally met with, 644
 diagnosis of from other forms of paralysis, 645
 from progressive muscular atrophy, 646
 occasionally simulates coxalgia, 647
 use of local treatment to spine in, 647
 ergot, belladonna, and iodide of potassium in acute stage, 647
 iron and strychnia in later stage, 648
 use of electricity in, directions, choice of current, 648
 mechanical contrivances in treatment of, 649

- Paralysis, tenotomy to relieve deformity, 650
 necessity of pursuing treatment for years, 650
facial, article on, 650-652
 causes of, 650
 symptoms of, 651
 diagnosis of from cerebral hemiplegia, 651
 prognosis in, 652
 treatment of, 652
progressive, with apparent hypertrophy of the muscles (*see* Progressive muscular sclerosis), 652-660
- Parasite of thrush (*see* *Oidium albicans*).
 in false membrane of diphtheria, 880
 of favus (*see* *Achorion Schoenleinii*), 982
 of tinea trichophytina (*see* *Trichophyton*), 982
 of tinea versicolor (*see* *Microsporon*), 982
 of alopecia areata, 996
- Parasitic skin diseases, 980-1002
 general remarks on, 980
 varieties of, 980
 nature of, 980
 mode of detecting fungus in, 981
 relation between fungus and the eruption, 981
 description of *achorion Schoenleinii* (*see* *Favus* also), 982
 description of *trichophyton* (*see* *Tinea* also), 982
 description of *microsporon furfur* (*see* *Alopecia areata*), 982
 relation of the various fungi in, 983
- Parasitocides, 989
- Parkes, analysis of milk, 304, etc.
- Parotid gland, condition of in mumps, 863
 suppuration of in mumps, 864
- Parotitis (*see* *Mumps*).
 in typhoid fever, 724
- Paroxysm of whooping-cough, peculiarities of, 261
 treatment of, 280
 of laryngismus stridulus, symptoms of, 583
 treatment of, 590
 of eclampsia, symptoms of, 563
 treatment of, 571
 in tetanus, 606
 in diphtheritic croup, 890
- Parrot, on thrush, 311, 314
- Parry, observations on rickets, 695
- Pathology of catarrhal pneumonia, 162
 of entero-colitis, 418
 of intussusception, 483
 of night terrors, 662
 of rickets, 704
 of diphtheria, 879
- Pelvis, alterations of in rickets, 700
- Pemphigus infantilis in congenital syphilis, 710
 article on, 956-959
 definition and synonyms of, 956
 forms and frequency of, 956
- Pemphigus, causes of, 957
 symptoms of, 957
 duration of, 958
 diagnosis of, 958
 prognosis in, 958
 general treatment of, 958
 local treatment of, 959
- Pepper, Professor, case of laryngismus stridulus, 592
 incontinence of urine in chorea, 620
- Pericarditis, acute, symptoms and difficulty of detection of, 290
 prognosis in, 290
 anatomical appearances in, 290
 treatment of, 290
 chronic, 291
 in scarlatina, 802
- Percussion of heart, 37
 of lungs, mode of performing, etc., 44, 45
 in pneumonia, 167, 174
 in pleurisy, 234
 in bronchial phthisis, 684
 in pulmonary phthisis, 687
- Perforation of cæcum, 476
 of appendix cæci, 477
 of intestine in typhoid fever, 723, 728
 from *ascaris lumbricoides*, 1008
- Peritoneum, tuberculosis of (*see* *Tuberculosis*), 681, 688
- Peritonitis, from perforation of cæcum and appendix, 477
 tuberculosis (*see* *Tuberculosis*).
 in scarlatina, 802
- Perityphlitis, in art, on dis. of cæcum and appendix, 467-481
 definition, 468
 iliac abscess in, 468
 anatomical appearances in, 472
 symptoms of, 477
- Perspiration, tendency to profuse, in rickets, 866
- Pertussis (*see* *Whooping cough*), 259
- Peyer's patches in typhoid fever, 716
- Pharyngitis, simple, 368-373
 definition and frequency, 368
 causes of, 368
 lesions in, 368
 symptoms of, 369
 diagnosis of, 371
 prognosis in, 372
 treatment of, 372
- Phlyzacia (*see* *Ecthyma*), 962
- Phthisis (*see* *Tuberculosis*).
- Physical signs (*see* *Auscultation and Percussion*).
- Physiognomy in diseases, 21
 in pneumonia, 176
 in pleurisy, 235
 in tetanus, 605
 in cases of worms, 1010
- Physostigma in chorea, 629
- Pinkroot as a vermifuge, 1015
- Pitting in variola (*see* under *Variola*), 735
 in varioloid, 739
 treatment to prevent, 748

- Pityriasis, symptoms of, 973
 treatment of, 974
 Pleura, remarks on diseases of, 133
 Pleurisy, article on, 231-252
 definition, frequency, and forms, 231
 predisposing causes of, 231
 exciting causes of, 231
 anatomical lesions in, 231
 symptoms of acute form, 233
 physical signs from auscultation, 233
 percussion, 234
 inspection, 235
 palpation, 235
 rational symptoms, pain, cough, respiration, pulse, 235
 temperature in, 236
 urine in, 237
 course of, 237
 symptoms of chronic form, 238
 of empyema, 239
 diagnosis of, 240
 obscurity in early stage from violence of constitutional symptoms, 241
 prognosis and mortality in, 241
 treatment of, 242-252
 blood-letting in, 242
 antimony in, 242
 mercury in, 243
 diuretics and purgatives in, 244
 external remedies in, 244
 paracentesis, 244-252
 indications for, 245
 objections to, 245
 Trousseau's rule in regard to, 246
 indicated in empyema, 246
 success greater in children, 247
 rules to guide in advising, 248
 mode of performing, 248
 after-treatment, 249
 use of medicated injections through canula, 250
 illustrative case of chronic form, 251
 Pleuro-pneumonia, physical signs in, 234
 mortality in, 241
 Pneumonia, presence of, contraindicating tracheotomy, 117
 lobular, in reality collapse of the lung, 144
 catarrhal, 172
 differential diagnosis of, 182
 prognosis of, 174
 differences between condition of lung in, and in collapse, 147
 in whooping-cough, 266
 in measles, 841, 852
 article on, 157
 definition and synonyms of, 157
 frequency and mortality of, 157
 forms and classification, 158
 lobular, identity of with collapse, 145, 158
 predisposing causes, 158
 age at which most frequent, 159
 table showing influence of season, 160
 relation of mortality from, to the temperature, 160
 exciting causes of, 160
 Pneumonia, anatomical lesions of, 160
 of lobular form, 161
 of partial form, 162
 abscess of lung following, 162
 difference between condition of lung in, and in non-inflammatory congestion, 163
 inflation of lung impossible in, 163
 usually unilateral, 164
 portion of lung involved in, 164
 apex quite frequently the seat of, 164
 not so frequently attended by bronchitis as formerly thought, 165
 association with pleurisy, 165
 with empyema, 165
 general course of, in young children, 165
 in children over two years old, 166
 varieties of mode of onset, simulating other affections, 165
 unfavorable symptoms and modes of death, 168, 173, 183
 general course of the partial form, 172
 duration of, 173
 physical signs of, 174
 cough in, 175
 expectoration in, 175
 thoracic pain in, 176
 state of respiration in, 176
 physiognomy in, 176
 grade of fever in, 167, 177
 rate of pulse in, 167, 171, 178
 nervous symptoms, convulsions, 178
 appetite, vomiting, diarrhoea in, 178
 thirst in, 178
 urine, albuminuria in, 179
 chlorides in, 179
 diagnosis from bronchitis, 179
 from pleurisy, 180, 240
 of catarrhal form, 180
 of cerebral form, 180
 during teething, 182
 from typhoid fever, 726
 prognosis of, 182
 treatment of, 184-195
 question of blood-letting in treatment of, 185
 indications for bleeding in, 187
 use of antimony in, 188
 calomel in, 189
 salines in, 189
 quinia in, 189
 ipecacuanha in, 190
 muriate of ammonia in, 190
 purgatives in, 190
 external applications in, 191
 tonics and stimulants in, 192
 diet in, 192
 use of opium in, 193
 general management of, 194
 diet in, 189
 importance of administration of water in, 194
 confinement to bed, 195
 change of position in, 195
 scrofulous, 675

- Pneumonia, relations of to tuberculosis of the lung, 679, 688
in typhoid fever, 722
- Pneumothorax, from rupture of abscess of the lungs, 256
article on, 252-259
nature of, 253
anatomical appearances in, 253
case of, 254
causes of, 255
 phthisis, pneumonia, gangrene, and emphysema, 256
 symptoms of, 256
 course of, 257
 prognosis in, 258
 diagnosis of, 258
 treatment of, 258
- Pock, anatomy of variolous, 741
of vaccine, 752
- Poliomyelitis, anterior (*see* Infantile Paralysis), 636
- Pompholyx (*see* Pemphigus), 956
- Population, density of, as a cause of enterocolitis, 409
- Porrigo (*see* Tinea), 991
- Porrigo larvalis (*see* Eczema capitis), 937
 granulata (*see* Eczema granulosum), 940
 favosa (*see* Favus), 985
 scutulata (*see* Favus), 986
- Position recommended in atelectasis, 142
- Post-pharyngeal abscess (*see* Retropharyngeal), 373
- Potash, carbonate of, in whooping-cough, 273
and acetate of, in rheumatism, 669
chlorate of, in ulcerative stomatitis, 335
 in diphtheria, 904
 in scarlatina, 824
 caustic solutions of, in eczema, 949
- Potassium, iodide of, in pleurisy, 243
 in chronic valvular disease of the heart, 300
 in simple meningitis, 536
 in tubercular meningitis, 521
 in chronic hydrocephalus, 556
 in infantile paralysis, 647
 in rheumatism, 670
 in scrofula, 678
 in congenital syphilis, 714
 sulphuret of, baths of, in chorea, 631
- Poultices in pneumonia, 191
 in bronchitis, 214
 in eczema, 945
 in favus, 988
- Prickly-heat (*see* Lichen tropicus), 955
- Prognosis in alopecia areata, 997
 in aphthæ, 330
 in ascaris lumbricoides, 1012
 in ascaris vermicularis, 1019
 in atelectasis pulmonum, 141
 in bronchitis, 210
 in chronic enlargement of the tonsils, 367
 in cholera infantum, 452
 in chorea, 625
 in cæcum and appendix, diseases of, 478
 in collapse of the lung, 154
- Prognosis in contraction with rigidity, 596
 in diarrhœa, 393
 in diphtheria, 893
 in dysentery, 463
 in ecthyma, 963
 in eczema, 942
 in entero-colitis, 423
 in eclampsia, 570
 in emphysema, 227
 in erysipelas, 869
 in endocarditis, 292
 in erythema, 923
 in favus, 987
 in gangrene of the mouth, 342
 in gastritis, 402
 in hemorrhage, cerebral, 547
 in herpes, 953
 in whooping-cough, 268
 in hydrocephalus, 555
 in impetigo contagiosa, 966
 in intussusception, 491
 in indigestion, 381
 in laryngitis, simple, 66
 spasmodic, 78
 pseudo-membranous, 100
 in laryngismus stridulus, 587
 in lichen, 968
 in malarial fever, 859
 in meningitis simple, 534
 tubercular, 516
 epidemic cerebro-spinal, 916
 in mumps, 864
 in night terrors, 664
 in paralysis, atrophic infantile, 640
 facial, 652
 pseudo-hypertrophic muscular, 656
 in pemphigus, 958
 in pharyngitis, 372
 in pleurisy, 241
 in pneumonia, 182
 in pneumothorax, 258
 in prurigo, 970
 in psoriasis, 972
 in retropharyngeal abscess, 375
 in rheumatism, 668
 in rickets, 701
 in roseola, 928
 in röteln, 857
 in rubeola, 848
 in rupia, 961
 in scabies, 1000
 in sclerema, 978
 in scarlatina, 807-809
 in stomatitis, gangrenous, 342
 ulcerative, 332
 in scrofula, 676
 in syphilis, congenital, 713
 in tetanus, 607
 in thrush, 367
 in tinea, 993
 in tonsillitis, 364
 in tuberculosis, 692
 in typhoid fever, 725
 in urticaria, 931
 in varicella, 767
 in variola, 744

- Prognosis in varioloid, 740
 in valvular diseases of the heart, 295
- Progressive muscular sclerosis, article on, 652-660
 definition of, 652
 history and synonyms of, 652
 causes of, 653
 symptoms of, 653
 peculiar gait in, 654
 condition of muscles in, 654, 658
 produces club-foot, 655
 electrical condition of muscles in, 655
 appearance of skin in, 655
 temperature in, 656
 course of, 656
 mental condition in, 656
 duration variable, 656
 termination fatal, by affection of respiratory muscles, 656
 diagnosis of, 656
 anatomical appearances in, 658
 treatment of, 659
- Prophylaxis in dropsy of scarlatina, 828
 in scrofula, 677
 in tetanus nascentium, 607
 in tubercular meningitis, 526
 in tuberculosis, 692
- Prurigo, article on, 969-971
 definition of, 969
 frequency of, 969
 causes of, 969
 symptoms of, 969
 duration of, 970
 diagnosis of, from strophulus or lichen, 970
 from scabies, 926
 prognosis in, 970
 treatment of, 970
- Pseudo-hypertrophic muscular paralysis (*see* Progressive muscular sclerosis), 652
- Pseudo-membranous angina (*see* Diphtheria), 873
 in true croup, 85
 laryngitis (*see* Laryngitis).
- Psoriasis, article on, 971
 definition of, 971
 symptoms of, 971
 causes of, 971
 diagnosis of, 972
 prognosis in, 972
 treatment of, 972
 local applications in, 972
- Pulmonary resonance, characters of, in children, 44
 tuberculosis, 684, 687
- Pulse, in children, 33, 34, 35
 rate of, at different ages, 33, 34
 intermittence or irregularity of, 35
 irritability of, 35
 to be examined during sleep, 33
 peculiarities of, in tubercular meningitis, 510, 512
 in mild cases of scarlatina, 778
 in grave cases of scarlatina, 784
 in variola, 734
- Pulse in typhoid fever, 722
- Purgatives (*see* Formulæ) in pleurisy, 244
 in diseases of cæcum, 480
 in intussusception, 494
 in eclampsia, 573
 in chorea, 627
 in rheumatism, 671
 in scarlatina, 811, 827
 in scarlatinous dropsy, 827
 in rubeola, 850
 in variola, 746
 in eczema, 943
 in worms, 1014
- Pustules in variola, 733
 chapter on, 962
- Quinia in pneumonia, 189
 in bronchitis, 215
 in entero-colitis, 432
 in tetanus, 609
 in rheumatism, 670
 in diphtheria, 906
 in malarial fever, 860
 in variola, 746
 in typhoid fever, 728
- Quinsy (*see* Tonsillitis), 364
- Rachitis (*see* Rickets).
- Radcliffe, J. N., electrical condition of muscles in infantile paralysis, 641
- Rashes, 920
- Raw meat, use of, in entero-colitis, 426
- Reaction of cow's and human milk, 303, 305
- Recession of base of chest in membranous croup, 97
- Rees, G. A., recession of base of chest in inspiration, 41
 respiration in collapse of the lung, 137
- Relapse of chorea, 621
 of acute rheumatism, 667
 of typhoid fever, 725
- Remittent Fever (*see* Typhoid fever), 715
- Residence, change of, in laryngismus stridulus, 590
 in unhealthy localities, influence of, 423
 change of, in treatment of entero-colitis, 434
 in treatment of scrofula, 677
 in treatment of tubercular meningitis, 527
 in country, importance of, 424
- Resonance, pulmonary, character in children, 44
- Respiration, general characters of, in children, 39, 40, 41, 43
 rate of, 39
 expiratory, 41
 diagnostic signs from, 41
 peculiar in atelectasis and croup, recession of base of thorax, 41
 puerile, 44
 alteration of in simple laryngitis, 64
 in catarrhal croup, 77
 in true croup, 96
 in pneumonia, 176

- Respiration, alteration in bronchitis, 207
 in pleurisy, 235
 in bronchial phthisis, 684
 in tubercular meningitis, 512
 in typhoid fever, 717, 718, 722
 in sclerema, 977
- Respiratory muscles affected in some cases of chorea, 620
 organs, diseases of, 52
 sounds, 44
- Rest, importance of, in cholera infantum, 459
 in bed, importance of, in rheumatism, 671
- Restlessness in disease, 29
- Retropharyngeal abscess, 373
 definition, causes, symptoms of, 374
 diagnosis, prognosis, treatment of, 374, 375
- Return-cry, alteration of, 27
- Revaccination, 763
- Rhagades in congenital syphilis, 709
- Rheumatism as a cause of heart disease, 289
 of chorea, 612, 624
 in scarlatina, 801, 828
 acute, article on, 666-672
 symptoms of, 666
 temperature in, 666
 condition of joints in, 666
 local symptoms often comparatively slight, 667
 duration and tendency to relapses, 667
 causes of, 667
 influence of sex not yet determined, 668
 chorea as complication of, 668, 612
 heart disease as complication of, 289
 prognosis in, 668
 diagnosis difficult when local symptoms are slight, 668
 treatment of, 669
 alkalies in, 669
 salicylic acid in, 670
 iodide of potash in, 670
 iron and quinia in, 670
 opium in, 670
 importance of strict rest in bed, 671
 diet in, 671
 local applications in, 671
 treatment of complications, 672
- Ribs, beaded in rickets, 698
- Richardson, B. W., heart-clot in diphtheria, 896
- Rickets, article on, 694-706
 definition and frequency of, 694
 causes of, 695
 symptoms of initiatory stage, 696
 digestive disturbances in, 696
 general soreness of body in, 697
 tendency to profuse perspiration in, 697
 dentition impeded in, 697
- Rickets, urine in, 697
 mental condition in, 697
 cerebral blowing murmur in, 697
 stage of deformity, 698
 alterations of long bones in, 698
 of head in, 699
 of spine in, 699
 of thorax in, 699
 of thorax due to atmospheric pressure, 699
 of pelvis in, 700
 general symptoms in later stage, 701
 in favorable cases, 701
 secondary diseases causing death in, 701
 prognosis of duration in, 701
 diagnosis of, 701
 morbid anatomy of bones in, 702
 collapse of lung in, 703
 condition of viscera in, 703
 pathology of, 704
 treatment of, 705
 importance of proper diet in, 705
 cod-liver oil in, 705
 means of avoiding deformities, 706
- Rilliet and Barthez, size of heart by percussion, 37
 diagnosis between true and false croup, 75
 atelectasis and collapse of the lung, 145
 state of vessels in gangrene of the mouth, 338
 diagnosis of gangrene of the mouth from ulcero-membranous stomatitis, 338
 lesions in diarrhoea, 389
 on pathology of cholera infantum, 447
 diagnosis of simple from tubercular meningitis, 514
 diagnosis of simple meningitis from congestion of the brain, 534
 diagnosis of symptomatic from essential contraction, 596
 on convulsions in scarlatina, 808
 cold affusion in scarlatina, 816
- Rindfleisch, characters of false membrane, in croup, 86
- Ringworm (*see* Tinea), 993
- Robinson on heart-clot in diphtheria, 896
- Roger, pulse in children, 33
 respiration in children, 40
- Rokitansky, on changes in spinal cord in tetanus, 605
- Romberg, relation of rheumatism to chorea, 613
- Rosalia, or rubeola notha, 806
- Roseola, epidemic, or rubeola notha, 806
 article on, 925-928
 definition and synonyms, 925
 frequency of, 925
 forms of, 925
 causes: occasionally epidemic, 926
 symptoms of; eruption, 926

- Roseola, duration of, 926
annulata.
 symptoms of, 927
 diagnosis of, from scarlatina, 927
 diagnosis of, from rubeola, 928
 from herpes iris, 928
 from erythema, 923
 prognosis in, 928
 treatment of, 928
 in congenital syphilis, 708
- Rötheln, article on, 855-857
 definition, synonyms, frequency, of 855
 history of, 855
 nature of, 855
 symptoms of, 856
 date of eruption, 856
 characters of eruption, 856
 epidemic nature of, 856
 diagnosis of, 856
 from scarlatina, 856
 from measles, 856
 treatment of, 857
- Round-worm (*see* *Ascaris lumbricoides*), 965
- Rubeola, article on, 831-854
 definition of, 831
 forms of, 831
 frequency of, 831, 769
 epidemic nature of, 832
 contagiousness of, 832
 period of incubation of, 832
 influence of age on frequency of, 832
 straw-fungus as cause of, 833
 symptoms of, 833
 mode of invasion of regular form of, 833
 fever in initial stage of, 834
 catarrhal symptoms in initial stage, 834
 marked drowsiness in initial stage, 835
 convulsions in initial stage of, 835
 red papules on palate in initial stage of, 836
 duration of initial stage of, 836
 date of appearance of eruption in, 836
 characters of eruption in, 836
 symptoms during eruption in, 837
 duration of eruption in, 837
 urine, during eruption in, 837
 symptoms of stage of decline of, 837
 desquamation in, 837
 temperature in, 838
 irregularities of prodromic stage in, 838
 of eruption, 839
 petechial character of eruption without any malignant symptoms, 839
 form of, without eruption, 839
notha or *sine catarrho*, a form of roseola, 840
 malignant form of, 840
 eruption in, 840
 complications and sequelæ of, 845
 bronchitis and pneumonia in, 841
 effect of, upon eruption, 842
 prognosis in bronchitis and pneumonia in, 842
- Rubeola, laryngitis in, 842
 enteritis in, 843
 frequency of, 843
 causes of, 843
 symptoms of, 843
 fatal cerebral symptoms in, 844
 cases of, 844
 serous effusions in, 845
 tendency of, to develop tuberculosis, 845
 coëxisting with variola, scarlatina, or erysipelas, 845
 anatomical lesions in, 845
 diagnosis of, 846
 from roseola and rötheln, 846, 856
 from variola, 847
 from typhus, 847
 prognosis in, 848
 causes of death in, 848
 treatment of, 849
 hygienic treatment of, 849
 diet in, 850
 laxatives and febrifuges in, 850
 depletion in, 851
 treatment of malignant form of, 852
 of pulmonary complications in, 852
 counter-irritation in pulmonary complications in, 853
 treatment of diarrhæa in, 853
 of laryngitis in, 853
 of cerebral symptoms in, 854
- Rufz, statistics of chorea, 610
- Rupia, article on, 959-962
 definition of, 959
 varieties of, 960
 causes of, 960
 symptoms of, 960
 diagnosis of, from pemphigus, 961
 from ecthyma, 961
 prognosis in, 961
 general treatment of, 961
 local treatment of, 961
- simplex*.
 symptoms of, 960
- prominens*.
 symptoms of, 960
- Salicylic acid, in treatment of rheumatism, 670
- Salines, in treatment of membranous croup, 105
 of pneumonia, 189
- Salisbury, on straw fungus as cause of measles, 833
- Salivary secretion in mumps, 863
- Santonin as a vermifuge, 1016
- Scabies, article on, 997-1002
 definition of, 997
 caused by a *carus scabiei*, 998
 symptoms, 998
 seat of eruption in, 998
 local symptoms in, 998
 character of eruption in, 998
 cuniculi in, 999
 mode of detecting *acarus*, 999
 description of the *acarus*, 999

- Scabies, diagnosis of, by finding acarus or its ova, 999
 from eczema simplex, 1000
 from prurigo, 1000
 from lichen, 1000
 prognosis in, 1000
 treatment of, 1000
 applications of sulphur in (*see* Formulæ), 1001
 substitutes for sulphur in, 1001
 tarry applications in, 1001
 carbolic acid in, 1001
 general treatment of, 1002
- Scarlatina, article on, 769-831
 definition of, 769
 frequency of, 769
 forms of, 770
 contagion as a cause of, 772
 period of incubation of, 772
 transmitted by fomites, 773
 inoculability of, 773
 epidemic nature of, 773
 occasional occurrence of second attacks of, 774
 influence of age upon frequency of, 774
 of sex upon, 775
 symptoms of mild cases of, 775-781
 invasion generally sudden in, 776
 occasionally a short prodromic stage, 776
 characters of eruption in, 777
 duration of eruption in, 777
 pulse and fever in stages of eruption, 778
 tongue in stage of eruption, 778
 urine in stage of eruption, 779
 fauces in stage of eruption, 779
 symptoms of decline of, 779
 desquamation in, 780
 duration of mild cases, 780
 temperature in, 780
 no sharp line between mild and grave cases, 781
 illustrative cases, 781
 symptoms of grave cases of, 782
 sudden invasion in ataxic form of, 782
 case of, 784
 general symptoms in ataxic form of, 784
 convulsions in ataxic form of, 784
 delusive improvement in ataxic form of, 784
 eruption in ataxic form of, 785
 fatal symptoms in ataxic form of, 785
 invasion of grave cases sometimes less sudden, 785
 case of, 786
 condition of fauces in grave cases, 787
 pseudo-membrane in, 787
 swelling of submaxillary and cervical glands in grave cases, 788
 coryza and otorrhœa in grave cases, 788
 eruption in grave cases, 789
 general symptoms in grave cases, 789
 cases of this grave form, 790
 laryngitis in grave form, 792
- Scarlatina, duration of grave form, 793
 complications and sequelæ, 794
 dropsy as a sequel of, 794
 preceded by albuminuria, 794
 frequency of, very variable, 794
 period of occurrence, 794
 usually due to cold, 795
 due to tubal nephritis, 795
 condition of kidneys in, 796
 preceded by febrile symptoms, 796
 seat of effusion in, 797
 course and duration of, 798
 mode of death in, 798
 uræmic symptoms in, 798
 urine greatly diminished in, 799
 characters of, in, 799
 prognosis in, 800
 uræmia not necessarily fatal, 800
- diarrhœa as a complication of, 801
 rheumatism during, 801
 inflammation of serous membranes in, 802
 endo- or pericarditis in, 802
 peritonitis in, 802
 complicated with variola, measles, or diphtheria, 802
 paralysis after, 802
 anatomical lesions in, 802
 condition of gastro-intestinal mucous membrane in, 802
 of skin in, 803
 of blood in, 803
 connection of, with enteric fever, 803
 heart-clot in, 804
 diagnosis of, from measles, 805
 from roseola, 805
 from diphtheria, 806
 from rubeola notha, 806
 from erythema fugax, 806
 prognosis in, very variable in different epidemics, 807
 in mild cases, 808
 in grave cases, 808
 grave significance of convulsions in, 809
 unfavorable symptoms in, 810
 favorable symptoms in, 810
 hygienic treatment of, 810
 diet in, 811
 treatment of mild cases, 811
 use of warm baths and affusions in mild cases, 812
 care in use of purgatives in mild cases, 812
 treatment of angina in mild cases, 813
 inunction in, 813
 treatment of grave cases, 814
 cold affusions in grave cases, 815
 general remarks on baths, lotions, and affusions in grave cases, 816
 guide as to using cold in, 820
 best method of applying, 821
 temperature likely to continue to fall after application of, 821
 hyposulphite of soda and magnesia in, 823

- Scarlatina, use of tonics and stimulants in grave cases, 824
 treatment of angina in grave cases, 825
 external use of ice in grave cases, 825
 case of, 826
 importance of removing viscid secretions from throat, 827
 diarrhœa in, 827
 treatment of the rheumatism in, 828
 of otorrhœa, 828
 prophylactic of dropsy, 828
 of mild cases of dropsy, 828
 of severe cases of dropsy, 829
 diuretics in, 829
 hot baths as diaphoretics in dropsy, 828
 treatment of cerebral symptoms in dropsy, 830
 of the later stages of dropsy, 830
 use of belladonna as a prophylactic in, 830
- Schneeman, inunction in scarlatina, 813
- Sclerema, article on, 976-979
 definition and synonyms of, 976
 frequency of, 976
 date of occurrence of, 976
 cause of, 976
 atelectasis as cause of, 976
 symptoms of, 977
 extent of, 977
 condition of skin in, 977
 œdema in, 977
 temperature in, 977
 peculiar cry in, 977
 general symptoms of, 977
 symptoms of, in later life, 977
 prognosis in, 978
 diagnosis of, 978
 anatomical appearances, 978
 condition of skin in, 978
 bloodvessels in, 979
 lungs in, 979
 treatment of, 979
- Sclerosis of spinal cord in infantile paralysis, 642
 progressive muscular, 652
- Scrofula as sequel of hooping-cough, 267
 article on, 673-678
 definition and characters of, 673
 associated with tuberculosis, 673
 causes of, 674
 symptoms of, 674
 stages of, 674
 pneumonia and bronchitis in, 675
 albuminoid degeneration of viscera in, 675
 anatomical appearances in, 676
 diagnosis of, 676
 prognosis in, 676
 treatment of, 677
 preventive of, 677
 cod-liver oil in, 677
 iodine preparations in, 678
 mercury in, 678
 arsenic in, 678
- Screw-driver teeth, 711
- Season, influence upon frequency of true croup, 91, 878
 pneumonia and bronchitis, 159
 diphtheria and croup, 877
- Seaton, E. C., on vaccination, 762
- Seat-worm (*see* *Ascaris vermicularis*), 1007
- Second attacks of scarlatina, 774
- Secondary fever in variola, 735, 746
- Sée, on gymnastic exercises in chorea, 632
- Sensibility affected in diphtheritic paralysis, 899
- Sharples, Stephen P., analysis of milk, 303, 304, 306
- Shingles (*see* *Herpes zoster*), 951
- Simon, reaction of milk, 307
- Sims, Marion, on cause of tetanus nascens, 602
- Silver, nitrate of, in chronic entero-colitis, 437
 in dysentery, 466
 local application of, in diphtheria, 901
 in diphtheritic paralysis, 907
 local use of, in scarlatinous angina, 826
 to prevent pitting in variola, 749
 enema of, in worms, 1020
- Skin, diseases of, introductory remarks on, classification of, and method of arriving at a diagnosis, 918, 919
 examination of, 30, 31, 32
 color of, in infants, 31, 32
 in different diseases, 31, 32
 exudation on, in diphtheria, 889
 in congenital syphilis, 707
 in scarlatina (*see* *Eruption and desquamation*).
 in measles (*see* *Eruption and desquamation*).
 in variola (*see* *Eruption and desquamation*).
 in typhoid fever, 720
 in progressive muscular sclerosis, 656
 gangrene of, in erysipelas, 867
 in sclerema, 977
 in favus, 985
- Sleep, diagnostic signs from, 23
- Small-pox (*see* *Variola*), 729
- Smith, J. Lewis, on lesions in cyanosis, 281
 on symptoms in cyanosis, 286
 state of intestines in entero-colitis, 413
 on liver in entero-colitis, 417
- Smith, E., analysis of milk, 304
- Snuffles, 52
- Soaps, in treatment of eczema, 948
- Soda, hyposulphite of, in scarlatina, 823
 in favus, 990
- Softening of stomach, 399
 of bones in rickets, 702
- Soreness of body in rickets, 701
- Sounds of heart, 36
 respiratory, 44
- Spasm of glottis (*see* *Laryngismus stridulus*), 577
 carpopedal, 584, 595
- Spence, J., on tracheotomy in croup, 114
- Sphincters affected in chorea, 620
- Spigelia as a vermifuge, 1015

- Spinal cord in tetanus, 605
 in chorea, 617
 in infantile paralysis, 642
 column, alteration of, in rickets, 699
 Spiritus saponatus kalinus, 948
 Spleen, enlarged in typhoid fever, 721
 in malarial fever, 859
 Spotted fever, 908
 Sputa, in bronchitis, 207
 in pneumonia, 175
 Squamous inflammation of the skin, article on, 971
 Statistics of frequency and mortality of
 bronchitis, 196
 of croup, pseudo-membranous, 90
 of cholera infantum, 442
 of chorea, 611, 626
 of cœcum and appendix, diseases of, 469
 of diphtheria, as compared to true croup, 90, 878
 of eclampsia, 560
 of entero-colitis, 406
 of hooping-cough, 269
 of intussusception, 482
 of laryngismus stridulus, 577
 of meningitis, epidemic cerebro-spinal, 909
 of meningitis tubercular, 499
 of nervous diseases, 498
 of paracentesis thoracis, 247
 of pleurisy, 241
 of pneumonia, 157, 160
 of rickets, 695
 of rubeola, 769
 of scarlatina, 769, 775
 of tracheotomy in croup, 108
 of thrush, 361
 of typhoid fever, 725, 909
 of tetanus nascentium, 603
 of tuberculosis, 680
 of variola, 730
 of variola after vaccination, 762, 764
 Stimulus, indications for, in pneumonia, 192
 in bronchitis, 215
 use of, in entero-colitis, 432, 440
 in stage of collapse in cholera infantum, 457
 in chorea, 630
 in diphtheria, 907
 in scarlatina, 824
 in measles, 852
 in variola, 746
 in typhoid fever, 728
 in erysipelas, 872
 Stomach and intestines, general remarks on diseases of, 376
 functional disease of (*see* Indigestion), 376
 diseases of, attended with lesions, 397
 condition of, in thrush, 350, 353
 inflammation of (*see* Gastritis), 397
 softening of, 399
 condition in entero-colitis, 417
 eruption on, in variola, 740
 Stomatitis.
 in congenital syphilis, 711
 in small-pox, 734
erythematous.
 article on, 329
 definition and frequency of, 329
 causes of, 329
 symptoms of, 329
 treatment of, 329
follicular.
 article on, 330-332
 definition, synonyms, frequency of, 330
 forms of, 330
 causes of, 330
 symptoms and duration of, 330
 diagnosis of, 330
 prognosis in, 331
 treatment of, 331
ulcerative or ulcero-membranous.
 article on, 332-336
 definitions, synonyms, frequency of, 332
 causes of, 333
 symptoms and course of, 333
 duration of, 334
 diagnosis of, 334
 prognosis in, 334
 treatment of, 334
gangrenous.
 article on, 236-347
 definition, synonyms, frequency of, 336
 causes of, 337
 anatomical lesions of, 337
 symptoms and course of, 338
 duration of, 338
 complications of, 341
 diagnosis of, 341
 prognosis in, 342
 treatment of, 343, 347
 Stools, diagnostic signs from the, 50
 in simple diarrhœa, 391
 in acute entero-colitis, 419
 in chronic entero-colitis, 422
 in cholera infantum, 451
 in dysentery, 463
 bloody, in intussusception, 488
 in typhoid fever, 721
 in case of worms, 1010
 Straw fungus as cause of rubeola, 833
 Strophulus, article on, 967-969
 definition of, 968
 causes of, 968
 varieties and symptoms of, 968
 diagnosis of, 969
 treatment of, 969
 intertinctus or *red gum*, 968
 confertus, 968
 albidus or *white gum*, 968
 Strychnia in chorea, 630
 in infantile paralysis, 648
 in diphtheritic paralysis, 907
 Submaxillary glands, swelling of, in croup, 98
 in diphtheria, 883
 in mumps, 863

- Sucking, signs from mode of, 49
 Sudamina in typhoid fever, 718, 722
 Sugar in urine in whooping-cough, 263
 in woman's milk, 309
 in condensed milk, 322
 Sulphurous applications in favus, 989
 Sulphur in treatment of scabies, 1001
 Summer complaint (*see* Cholera infantum), 441
 Suppurative fever in variola, 735
 Swine-pox (*see* Varicella).
 Sympathetic nerve, affection of, in cholera infantum, 449
 Syphilis, congenital, article on, 706-714
 modes of transmission of, to embryo, 706
 date of appearance of symptoms, 707
 appearance of skin in, 707
 affection of the skin in, 708
 occurrence and varieties of, 708
 coryza and stomatitis in, 710
 alteration of voice in, 710
 course of, 711
 development of tertiary stage, 711
 alteration of teeth in, described by Hutchinson, 711
 interstitial keratitis in, 712
 affection of internal organs in, 712
 anatomical lesions in, 712
 diagnosis of, in early stage, 713
 in later stage, 713
 between inherited and acquired, 713
 prognosis in, 713
 treatment of, 713
 use of mercury in, 713
 Syphilis transmitted by vaccination, 760
 Syphiloderma, 708
 Tabes mesenterica (*see* Tuberculosis of mesenteric glands), 682
 Table of mortality (*see* Statistics).
 Tache méningitique in meningitis, 511
 Tænia solium, 1004
 lata, 1005
 Tape-worm (*see* Tænia), 1004
 Tarry applications in eczema, 947
 in scabies, 1001
 in tinea, 994
 in favus, 990
 Tartar emetic (*see* Antimony).
 Tears, arrest of, in disease, 27
 Teeth, alteration of, in congenital syphilis, 711
 delayed development of, in rickets, 697
 Temperature, normal, at different ages, 37
 tables of observations on, 37, 38
 effect of, on mortality of pneumonia and bronchitis, 160
 of body, low in cyanosis, 286
 in tubercular meningitis, 513
 lowered in infantile paralysis, 639
 in progressive muscular sclerosis, 656
 in acute rheumatism, 666
 influence of, upon membranous croup, 91
 Temperature, effect upon mortality of
 diphtheria and croup, 877, 878
 in pulmonary phthisis, 686
 in scarlatina, 778, 780
 in measles, 838
 in typhoid fever, 717
 high, as cause of miliaria, 955
 in sclerema, 977
 proper for sick-room, 849
 Tenotomy in infantile paralysis, 650
 for deformities in chorea, 650
 Terror as a cause of chorea, 614
 Testicle affected by metastasis in mumps, 864
 Tetanus nascentium, article on, 602-610
 definition and synonyms of, 602
 period of occurrence of, 602
 morbid conditions of umbilicus as cause of, 602
 Sims's view of displacement of occipital bone as cause of, 602
 general causes of, 603
 frequency of, 603
 anatomical lesions in, 604
 microscopical changes in spinal cord in, 605
 symptoms of, 605
 recurrence of paroxysm in, 606
 prognosis in, 607
 diagnosis of, 607
 duration of, occasionally chronic, 607
 prophylaxis in, 607
 treatment of, 608
 anæsthetics in, 608
 narcotics and antispasmodics in, 609
 Thermometer, observations with, in children, 37
 Thirst, as a cause of crying, 25
 Thorax, sonorousness of, in children, 45
 percussion of, 44-46
 auscultation of, 43, 44
 recession of base of, in croup, etc., 41
 alterations in shape of, in rickets, 699
 Thread-worm (*see* Ascaris vermicularis), 1004, 1019
 (*see* Tricocephalus dispar), 1004
 Throat, examination of, 47
 diseases of (*see* Pharyngitis), 370
 Thrush, article on, 347-363
 definition, synonyms of, 347
 frequency of, 347
 predisposing causes of, 347
 exciting causes of, 348
 contagion of, 349
 anatomical lesions of, 349
 description of fungus of, 350
 symptoms of, 353
 nature of, 358
 diagnosis of, 360
 prognosis in, 360
 prophylactic treatment of, 361
 general treatment of, 362
 local treatment of, 363
 Thymus gland, enlarged in laryngismus stridulus, 580
 Tinea lactea (*see* Eczema capitis), 937

- Tinea granulata* (*see* *Eczema granulosum*), 939
decalvans (*see* *Alopecia areata*), 996
trichophytina.
 varieties of, 990
 identity of *t. tonsurans* and *t. circinata*, 990
 synonyms of, 991
 description of its fungus, *trichophyton*, 991
 contagion as cause of, 991
 other causes of, 991
tonsurans.
 eruption in, 991
 character of hairs in, 992
 condition of scalp in, 992
 diagnosis of, 992
 prognosis in, 993
circinata.
 eruption in, 993
 character of hairs in, 993
 diagnosis of, 993
 general treatment in, 993
 local treatment in, 994
 depilation in, 994
 alkaline applications in, 994
 mercurial applications in, 994
 tarry applications in, 994
 cases of, 994
 Tongue in scarlatina, 778
 in typhoid fever, 721
 Tonsils, sloughing of, in diphtheria, 882
 ablation of, in diphtheria, 904
 condition of, in scarlatina, 787
 acute inflammation of (*see* *Tonsillitis*), 364
 chronic enlargement of, 366
 nature and causes of, 366
 anatomical appearances in, 366
 symptoms of, 366
 prognosis in, 367
 treatment of, 367
 Tonsillitis, 364-365
 definition of, 364
 symptoms of, 364
 course and duration of, 364
 prognosis in, 364
 diagnosis of, 364
 treatment of, 365
 Tracheotomy, in membranous croup, 107
 statistics of its performance in different countries, 108
 estimation of its value, 112
 dangers of the operation, 113
 rules to guide in advising, 113
 proper period for performing, 113, 118
 indications for, 114
 influence of period of performance upon result, 114
 contraindications, 115, 118
 age as a contraindication, 115
 successful cases at early age, 115
 extension of false membrane into the bronchi as a contraindication, 116
 negative results of auscultation, 116
 presence of pneumonia as a contraindication, 117
 Tracheotomy, general diphtheria as a contraindication, 118
 contraindicated in secondary forms of croup, 118
 twice successfully performed in same subject, 119
 mode of performing, 119-123
 instruments required in performing, 119
 details about canulas, 120
 substitutes for canulas, 121
 question of fixing trachea in, 121
 excising piece of trachea in, 121
 mode of performing, 122
 emphysema of neck following, 123
 use of anæsthetics during, 123
 after-treatment, 123-127
 great importance of, 124
 modes of rendering inspired air moist, 124
 treatment of wound, 124
 instillation and atomization into trachea after, 124
 directions for cleansing tubes after, 124
 mode of removing canula, 126
 date of removing canula, 126
 causes of delay in removing canula, 126
 results following prolonged stay of canula, 127
 general after-treatment, 127
 importance and manner of feeding patients, 127
 question of medication after, 128
 difficulty of deglutition following, 127
 illustrative cases of, 128
 Treatment of alopecia areata, 997
 of aphthæ, 331
 of *ascaris lumbricoides*, 1013
 of *ascaris vermicularis*, 1019
 of atelectasis pulmonum, 141
 of atrophic infantile paralysis, 647
 of bronchitis, acute, 211
 chronic, 216
 of cerebral congestion, 539
 hemorrhage, 547
 paralysis following, 548
 of cæcum and appendix, diseases of, 479
 of cholera infantum, 453-459
 of chorea, 626
 of collapse of the lung, 154
 of contraction with rigidity, 596
 of congenital syphilis, 713
 of coryza, 57
 of cyanosis, 288
 of diarrhœa, acute, 393
 chronic, 395
 of diphtheria, 901
 of dysentery, 464
 of eclampsia, 571
 of ecthyma, 963
 of eczema, 943
 of emphysema, 228
 of endocarditis, 292
 of entero-colitis, 423-434

- Treatment of epidemic cerebro-spinal meningitis, 917
 of erysipelas, 870
 of erythema fugax, 925
 intertrigo, 925
 nodosum, 925
 of facial paralysis, 652
 of favus, 988
 of gangrene of the mouth, 343
 of gastritis, 402
 of herpes, 953
 of hooping-cough, 269
 of hydrocephalus, 556
 of ichthyosis, 975
 of impetigo contagiosa, 966
 of indigestion, 381
 of laryngeal cough, chronic, 67
 of laryngitis, simple, 66
 spasmodic, 78
 pseudo-membranous, 100
 of laryngismus stridulus, 587
 of lichen scrofulosus, 967
 strophulus, 969
 of malarial fever, 860
 of meningitis, simple, 534
 tubercular, 520
 epidemic cerebro-spinal, 917
 of miliaria, 956
 of mumps, 865
 of night-terrors, 664
 of paralysis, atrophic infantile, 647
 facial, 652
 pseudo-hypertrophic, 659
 of pemphigus, 958
 of pericarditis, 290
 of pharyngitis, 372
 of pityriasis, 974
 of pleurisy, 242-252
 of pneumonia, 184-195
 of pneumothorax, 258
 of progressive muscular sclerosis, 659
 of prurigo, 970
 of psoriasis, 975
 of retropharyngeal abscess, 375
 of roseola, 928
 of r  theln, 857
 of rheumatism, 669
 of rickets, 705
 of rupia, 961
 of scabies, 1000
 of scarlatina, 801-830
 of sclerema, 979
 of sclerosis, progressive muscular, 659
 of scrofula, 677
 of stomatitis, erythematous, 329
 follicular, 331
 ulcerative, 334
 gangrenous, 343
 of strophulus, 969
 of syphilis, congenital, 713
 of tetanus nascentium, 608
 of thrush, 361
 of tinea, 993
 of tonsillitis, 365
 of tonsils, chronic enlargement of, 367
 of tuberculosis, 692
- Treatment of typhoid fever, 727
 of urticaria, 931
 of valvular diseases of the heart, 299
 of varicella, 767
 of variola, 745
 Typhoid fever, diagnosis from tubercular meningitis, 515
 article on, 715-729
 formerly confounded with remittent fever, 715
 causes of, 715
 but slightly contagious, 716
 epidemic nature of, 716
 anatomical appearances in, 716
 condition of intestine in, 716
 condition of blood in, 717
 condition of brain in, 717
 symptoms of ordinary cases, 717
 prodromes in, 717
 marked remissions in febrile action, 717
 eruption in, 718, 722
 symptoms of fully developed attack, 718
 of grave cases, 723
 favorable symptoms in, 719
 unfavorable symptoms in, 718
 condition of skin in, 720
 digestive disturbances in, 721
 character of stool in, 721
 distension of abdomen in, 721
 enlargement of spleen in, 721
 urine in, 721
 respiration in, 722
 pulse in, 722
 nervous symptoms in, 723
 pulmonary complications in, 723
 perforation of intestine in, 723
 intestinal hemorrhage in, 724
 albuminuria and œdema in, 724
 complicated with malaria or one of the eruptive fevers, 724
 tuberculosis as sequel of, 724
 convalescence in, 725
 relapses in, 725
 duration of, 725
 prognosis and mortality in, 725
 diagnosis of, from gastro-enteritis, 725
 from typhoid pneumonia, 726
 from acute tuberculosis, 726
 treatment of febrile symptoms in, 727
 of gastric irritability in, 727
 of intestinal symptoms in, 727
 of nervous symptoms, 728
 of complications in, 728
 use of mineral acids in, 728
 of quinia in, 728
 of opium in, 728
 stimulants in, 728
 diet in, 729
 management of convalescence, 729
 relation of scarlatina to, 803
- Ulceration of internal malleoli in thrush, 357
 of mucous membrane of fauces in diphtheria, 881

- Ulceration of mucous membrane in scarlatina, 787
 in erythema intertrigo, 921
- Umbilicus, morbid states of, as cause of tetanus, 602
- Umbilication of variolous pock, cause of, 740
- Uræmia, in scarlatinous dropsy, 798
- Urine, in bronchitis, 209
 in chorea, 620
 in diphtheria, 191
 in hooping-cough, 263
 in measles, 837
 in pleurisy, 237
 in pneumonia, 179
 in rickets, 697
 in scarlatina, 779, 798, 799
 in typhoid fever, 721
 in variola, 736
- Urticaria, article on, 928-932
 definition and synonyms of, 928
 forms of, 929
 causes of, 929
 symptoms of, 930
 diagnosis of, 931
 prognosis in, 931
 treatment of, 931
 diet in, 932
- Vaccination (*see* under Vaccine disease).
- Vaccine disease, article on, 751-769
- Vaccinia, 751-769
 definition and synonyms of, 751
 history of, 751
 date of appearance and development of pock, 752
 cellular character of pock, 752
 local and general symptoms in, 753
 desiccation and desquamation in, 753
 character of cicatrix after, 753
 irregularities and course of, 754
 severe local symptoms in, 754
 erysipelas following, 866
 appearance of pock retarded, 754
 spurious form of, 755
 diagnosis of, 755
 protective power of against variola, 756
 cases illustrative of, 756
 against death, 757
- Vaccination*, period of performing, 758
 susceptibility to, variable, 758
 effect of cutaneous eruptions on, 759
 alleged transmission of cutaneous diseases by, 759
 of syphilis by, 760
 forms of virus employed, 760
 characters of good vaccine crust, 760
 modes of introducing virus, 761, 765
 advantages of several punctures in, 761
 influence of number and quality of cicatrices on mortality after, 762
- Revaccination*, necessity for, 763
 results of, 763
 period of performance, 764, 765
- Vacuole (*see* Bronchial abscess), 199
- Valerian in chorea, 628
- Valleix, return-cry in infants, 27
 plan of examining abdomen, 47
 expectoration in true croup, 98
 emetics in true croup, 102
 on tracheotomy in croup, 111
 expectoration in pneumonia, 175
- Valvular diseases of heart (*see* Heart).
- Varicella, article on, 766-767
 definition of, 766
 synonyms of, 766
 forms of, 766
 contagious nature of, 766
 epidemic nature of, 766
 essentially distinct from variola, 767
 symptoms of, 767
 eruption in, 767
 diagnosis of, 767
 prognosis in, 767
 treatment of, 767
- Variola, article on, 729-751
 definition of, 729
 frequency of, 729
 table of mortality of, 730
 forms of, 731
 contagious nature of, 731
 epidemic nature of, 731
 transmitted by fomites, 731
regular form of, period of incubation in, 731
 symptoms of initial stage, 732
 pain in the loins in, 732
 symptoms of eruptive stage of, 732
 discrete and confluent forms of, 733
 date of appearance of eruption in, and character of papules in, 733
 development of eruption in, 734
 occurrence of eruption on mucous membranes, 734
 stomatitis in, 734
 laryngitis in, 734
 swelling of subcutaneous tissue in, 734
 subsidence of general symptoms on appearance of eruption, 734
 secondary fever, 735
 date of desiccation, 735
 date and modes of desquamation, 735
 pitting after, 735
 digestive symptoms during, 736
 urine in, 736
 excessive discharge of urine during desiccation, 736
 nervous symptoms in, 736
irregular forms of, confluent symptoms of initial stage, 736
 course of eruption in, 737
 hemorrhagic eruption in, 738
modified (*see* Varioloid), 738
 complications of, 740
 anatomical lesions, 740
 condition of blood in, 740
 mucous membranes in, 740
 anatomy of pock in, 741
 diagnosis of, 741

- Variola, prognosis in, 744
 favorable symptoms in, 744
 unfavorable symptoms in, 744
 treatment of, 745
 mild febrifuges in mild cases, 745
 laxatives in, 746
 remedies for favoring appearance of eruption in, 746
 diet in, 746
 quinia in, 746
 opium in, 746
 stimulants in, 746
 treatment of complications in, 747
 of conjunctivitis in, 747
 ventilation and disinfectants in, 748
 prevention of pitting in, 748
 cauterization of pock with nitrate of silver, 749
 mercurial applications in, 749
 applications of solution of gutta percha in, 751
- Varioloid, definition of, 738
 symptoms of, 739
 course of eruption more rapid in, 739
 absence of secondary fever in, 739
 no pitting after, 739
 duration of, 740
 prognosis in, 740
- Vasomotor nerves, influence of, in urticaria, 929
- Veins, pressure on by bronchial glands, 684
- Venesection (*see* Bleeding).
- Ventilation, importance of, in sick-chamber, 849
- Vermifuges, 1013
- Verminous abscess, 1009
- Vernois and Becquerel, analysis of milk, 308
- Vesicular inflammation of skin, article on, 933
 in variola, 732
- Virus, vaccine, forms of employed, and mode of introduction, 761
- Viscera, peculiar changes of in rickets, 703
- Voice, alteration of, in coryza, 57
 in simple laryngitis, 63
 in false croup, 77
 in true croup, 96
 in diphtheritic paralysis, 898
 in congenital syphilis, 710
- Volvulus (*see* Intussusception), 481
- Vomicæ, rare in tuberculosis of lungs, 680
- Vomiting, diagnostic signs from, 50, 51
 in gastritis, 401
 in entero-colitis, 420
 in entero-colitis, remedies for, 432
 in cholera infantum, 451
 treatment of, 456
- Vomiting in intussusception, 487
 in tubercular meningitis, 507
 in typhoid fever, 721
 in measles, 834
- Wagner, characters of false membrane in croup, 86
- Water, importance of administering in pneumonia, 194
 in cholera infantum, 457
 in entero-colitis, 427
 lessened quantity of in the system a cause of thrush, 359
 importance of, in feeding children, 314
 quantity required by children, 315
 injections of, in intussusception, 494
 external use of, in scarlatina (*see* also Bath and Affusion), 815, etc.
 local use of, in eczema, 945
- Waldenburg, treatment of emphysema, 229
- Wanklyn, analysis of condensed milk, 321
- Wells, case of chronic trismus, 607
- Wertheimer, on oedematous angina, 370
- West, C., on statistics of pneumonia, 159
 on cerebral congestion, 537
 on characteristics of bronchial phthisis, 685
- White gum, 968
- Wilson, Erasmus, description of *acarus scabiei*, 999
- Wood, H. C., origin of diphtheria, 879
- Woodward, J. J., on lesions in diarrhœa, 390
 microscopic changes in intestine in entero-colitis, 411
 use of mercury in diarrhœa, 428
- Woorara in tetanus, 609
- Worms in the alimentary canal, chapter on, 1003-1020
 varieties of, 1003
 description of *ascaris lumbricoides*, 1003
 of *ascaris vermicularis*, 1004
 of *tricocephalus dispar*, 1004
 of *tænia solium*, 1004
 lata, 1005
 frequency of, much exaggerated, 1005
 dangerous symptoms from, 1006
- Wormseed oil as a vermifuge, 1014
- Wunderlich, on temperature in scarlatina 821
- Xeroderma, 974
- Zinc ointment in eczema, 946
- Zona (*see* Herpes zoster), 951

ERRATA.

Page 715, line	i,	instead of vii,	read vi.
" 770, "	i,	" sixty,	" seventy.
" 856, "	xi,	" cause,	" course.
" 857, "	vii,	" occlusion,	" exclusion.

1871
MAY 11
P. 11

Received of the
Hon. Secy of the Navy
the sum of \$100.00

for the purchase of
the sum of \$100.00

for the purchase of
the sum of \$100.00

for the purchase of
the sum of \$100.00

for the purchase of
the sum of \$100.00

for the purchase of
the sum of \$100.00

for the purchase of
the sum of \$100.00

for the purchase of
the sum of \$100.00

for the purchase of
the sum of \$100.00

CATALOGUE

No. 1.

CATALOGUE

OF

MEDICAL, DENTAL, Pharmaceutical & Scientific Publications,

WITH A CLASSIFIED INDEX,

PUBLISHED BY

P. BLAKISTON, SON & CO.,

(SUCCESSORS TO LINDSAY & BLAKISTON)

Booksellers, Publishers and Importers of Medical and Scientific Books,

No. 1012 WALNUT STREET, PHILADELPHIA.

THE FOLLOWING CATALOGUES WILL BE SENT FREE TO ANY ADDRESS,
UPON APPLICATION.

This Catalogue, including all of our own publications.

A Catalogue of Books for Dental Students and Practitioners.


A Catalogue of Books on Chemistry, Pharmacy, The Microscope, Hygiene, Human Health, Sanitary Science, Technological Works, etc.

Students' Catalogue, including the "Quiz-Compendis" and some of the most prominent **Text-books** and **manuals** for medical students.

A Complete Classified Catalogue (32 pages) of all Books on Medicine, Dentistry, Pharmacy and Collateral Branches. English and American.

A Catalogue of Medical and Scientific Periodicals and Physicians' Visiting Lists, giving club rates.

P. Blakiston, Son & Co.'s publications may be had through Booksellers in all the principal cities of the United States and Canada, or any book will be sent, postpaid, by the publishers, upon receipt of price, or will be forwarded by express, C. O. D., upon receiving a remittance of 25 per cent. of the amount ordered, to cover express charges. Money should be remitted by postal note, money order, registered letter, or bank draft.

 All new books received as soon as published. Special facilities for importing books from England, Germany and France.

CLASSIFIED LIST, WITH PRICES,

OF ALL BOOKS PUBLISHED BY

P. BLAKISTON, SON & CO., PHILADELPHIA.

For further information in reference to these Books, send for full descriptive catalogue (No. 1), which will be sent, free, to any address. When the price is not given below, the book is not to be had at present. Cloth binding, unless otherwise specified.

ANÆSTHETICS.

Sansom. Chloroform.	\$1.25
Turnbull. 2d Ed.	1.50
— Cocaine.	.50

ANATOMY.

Handy. Text-book.	3.00
Heath. Practical.	5.00
Holden. Dissections.	5.00
— Landmarks.	—
Morris. On the Joints.	5.50
Potter. Compend of	1.00
— Visceral.	1.00
Wilson. 10th Ed.	6.00

ATLASES AND DIAGRAMS.

Bentley and Trimens. Medical Plants.	75.00
Braune. Of Anatomy.	8.00
Flower. Of Nerves.	3.50
Fox. Of Skin Dis.	20.00
Godlee. Of Anatomy.	20.00
Heath. Operative Surgery.	12.00
Hutchinson. Surgery.	25.00
Jones. Membrana Tympani.	4.00
Marshall's Physiol. Plates.	80.00
Schultze. Obstetrical Plates.	25.00

BRAIN AND INSANITY.

Bucknill and Tuke. Psychological Medicine.	8.00
Gowers. Diagnosis of Diseases of the Brain.	2.00
Mann's Psychological Med.	5.00
Roberts. Surgery of	1.25
Wood. Brain and Overwork	.50

CHEMISTRY.

Allen. Commercial Analysis. Volume 1.	4.50
Bartley. Medical.	2.50
Bloxam's Text-Book.	3.75
— Laboratory.	1.75
Bowman's Practical.	2.00
Frankland. How to Teach.	—
Leffmann's Compend.	1.00
Muter. Med'l and Pharm.	6.00
— Practical and Analy.	2.50
Richter's Inorganic.	2.00
— Organic.	3.00
Stammer. Problems.	.75
Sutton. Volumetric Anal.	5.00
Thompson's Physics.	—
Trimble. Analytical.	1.50
Vacher's Primer of.	.50
Valentin. Quant. Anal.	3.00
Ward's Compend of.	1.00
Watts. (Fowne's) Inorg.	2.25
— (Fowne's) Organ.	2.25
Wolff. Applied Medical Chemistry.	1.50

CHILDREN.

Chavasse. Mental Culture of.	1.00
Day. Diseases of.	3.00
Dillenberger. Women and.	1.50
Ellis. Manual of Dis. of.	3.00
— Mother's book on.	.75
Goodhart and Starr.	3.00
Hale. Care of.	.75
Hillier. Diseases of.	1.25
Meigs. Infant Feeding and Milk Analysis.	1.00
Meigs and Pepper's Treatise.	5.00
Smith. Wasting Diseases of.	3.00
Starr. Digestive Organs of.	—

COMPENDS

And The Quiz-Compend.

Brubaker's Physiology.	\$1.00
Horwitz. Surgery.	1.00
Hughes. Practice. 2 Pts. Ea.	1.00
Landis. Obstetrics.	1.00
Leffmann's Chemistry.	1.00
Mendenhall's Vade Mecum.	—
Potter's Anatomy.	1.00
— Visceral Anatomy	1.00
— Materia Medica.	1.00
Roberts. Mat. Med. and Phar.	2.00
Stewart. Pharmacy.	1.00
Ward's Chemistry.	1.00

DEFORMITIES.

Churchill. Face and Foot.	3.50
Coles. Of Mouth.	4.50
Prince. Orthopædics.	4.50
Reeves.	2.25

DENTISTRY.

Barrett. Dental Surg.	1.00
Coles. Dental Note Book.	1.00
Flagg. Plastics.	4.00
Gorgas. Dental Medicine.	3.00
Harris. Principles and Prac.	6.50
— Dictionary of.	6.50
Heath. Dis. of Jaws.	4.50
Hunter. Mechanical Dent.	1.50
Leber and Rottenstein.	—
— Caries.	1.25
Richardson. Mech. Dent.	4.00
Stocken. Materia Medica.	2.50
Taft. Operative Dentistry.	4.25
— Index of Dental Lit.	—
Tomes. Dental Surgery.	—
— Dental Anatomy.	4.25
White. Mouth and Teeth.	.50

DICTIONARIES.

Cleveland's Pocket Medical.	.75
Cooper's Surgical.	12.00
Harris' Dental.	6.50
Longley's Pronouncing	1.00

DIRECTORY.

Medical, of Philadelphia,	—
Pa., Del. and South N. J.	2.50

EAR.

Burnett. Hearing, etc.	.50
Dalby. Diseases of.	1.50
Jones. Aural Surgery.	2.75
— Membrana Tympani.	4.00
Woakes. Deafness, etc.	1.50
— Catarrh, etc.	—

ELECTRICITY.

Althaus. Medical Electricity.	6.00
Reynolds. Clinical Uses.	1.00

EYE.

Arlt. Diseases of.	2.50
Carter. Eyesight.	1.25
Daguenet. Ophthalmoscopy.	1.50
Fenner. Vision.	3.50
Gowers. Ophthalmoscopy.	6.00
Harlan. Eyesight.	.50
Hartridge. Refraction.	2.00
Higgins. Handbook.	.50
Jones. Sight and Hearing.	.50
Liebreich. Atlas of Ophth.	15.00
Macnamara. Diseases of.	4.00
Morton. Refraction.	1.00
Wolfe. Diseases of.	7.00

FEVERS.

Welch. Enteric Fever.	2.00
-----------------------	------

HEADACHES.

Day. Their Treatment, etc.	\$1.25
Wright. Causes and Cure.	.50

HEALTH AND DOMESTIC MEDICINE.

Bulkley. The Skin.	.50
Burnett. Hearing.	.50
Cohen. Throat and Voice.	.50
Dulles. Emergencies.	.75
Harlan. Eyesight.	.50
Hartshorne. Our Homes.	.50
Hufeland. Long Life.	1.00
Lincoln. Hygiene.	.50
Osgood. Winter.	.50
Packard. Sea Air, etc.	.50
Richardson's Long Life.	.50
Tanner. On Poisons.	.75
White. Mouth and Teeth.	.50
Wilson. Summer.	.50
Wilson's Domestic Hygiene.	1.00
Wood. Brain Work.	.50

HEALTH RESORTS.

Madden. Foreign.	2.50
Packard. Sea Air and Bath'g.	.50
Solly. Colorado Springs.	.25
Wilson. The Ocean as a.	2.50

HEART.

Balfour. Diseases of.	5.00
Fothergill. Diseases of.	3.50
Sansom. Diseases of.	1.25

HISTOLOGY.

See Microscope and Pathology.

HOSPITALS.

Burdett. Cottage Hospitals.	4.50
— Pay Hospitals.	2.25
Domville. Hospital Nursing.	.75

HYGIENE.

Bible Hygiene.	1.00
Frankland. Water Analysis.	1.00
Fox. Water, Air, Food.	4.00
Lincoln. School Hygiene.	.50
Parke's Hygiene. 6th Ed.	3.00
Wilson's Handbook of.	2.75
— Domestic.	1.00
— Drainage.	1.00
— Naval.	3.00

KIDNEY DISEASES.

Edwards. How to Live with	—
Bright's Disease.	.50
Greenhow. Addison's Dis.	3.00
Ralfe. Dis. of Kidney, etc.	2.75
Tyson. Bright's Disease.	3.50

LIVER.

Habershon. Diseases of.	1.50
Harley. Diseases of.	3.00

LUNGS AND CHEST.

See Phy. Diagnosis and Throat.

MARRIAGE.

Ryan. Philosophy of.	1.00
Walker. For Beauty, Health,	1.00

MATERIA MEDICA.

Biddle. 10th Ed.	4.00
Charteris. Manual of.	—
Gorgas. Dental.	3.00
Merrell's Digest.	4.00
Phillips. Vegetable.	—
Potter's Compend of.	1.00
— Handbook of.	—
Roberts' Compend of.	2.00

CLASSIFIED LIST OF P. BLAKISTON, SON & CO.'S PUBLICATIONS.

MEDICAL JURISPRUDENCE.

Abercrombie's Handbook,	\$2.50
Reese's Text-book of	4.00
Woodman and Tidy's Treatise, including Toxicology.	7.50

MICROSCOPE.

Beale. How to Work with.	7.50
— In Medicine.	7.50
Carpenter. The Microscope.	5.50
Lee. Vade Mecum of.	3.00
MacDonald. Examination of Water by.	2.75
Martin. Mounting.	2.75
Wythe. The Microscopist.	3.00

MISCELLANEOUS.

Allen. The Soft Palate.	.50
Beale. Life Theories, etc.	2.00
— Slight Ailments.	1.25
Black. Micro-Organisms.	1.50
Cobbold. Parasites, etc.	5.00
Edwards. Malaria.	.50
— Vaccination.	.50
— Constipation.	—
Gross. Life of Hunter.	1.25
Hardwicke. Med. Educat'n.	3.00
Hare. Tobacco. Paper.	.50
Hodge. Fœticide.	.50
Holden. The Sphygmograph.	2.00
Kane. Opium Habit.	1.25
MacMunn. The Spectroscope.	3.00
Mathias. Legislative Manual.	.50
Sieveling. Life Insurance.	2.00
Smith. Ringworm.	1.00
Smythe. Med'l Heresies.	1.25
Wickes. Sepulture.	1.50

NERVOUS DISEASES.

Buzzard. Ner. Affections.	5.00
Flowers. Atlas of Nerves.	3.50
Gowers. Dis. of Spinal Cord.	1.50
— Epilepsy.	4.00
Granville. Nerve Vibration.	2.00
Page. Injuries of Spine.	3.50
Radcliffe. Epilepsy, Pain, etc.	1.25
Tuke. Hypnotism, etc.	1.75
Wilkes. Nervous Diseases.	6.00

NURSING.

Cullingworth. Manual of.	1.00
— Monthly Nursing.	.50
Domville's Manual.	.75
Lückes. Hospital Sisters.	1.00
Record for the Sick Room.	.25
Temperature Charts.	.50

OBSTETRICS.

Barnes. Obstetric Operations.	3.75
Cazeaux and Tarnier. New Ed. Colored Plates.	—
Gallabin's Manual of.	—
Glisan's Text-book.	4.00
Landis. Compend.	1.00
Meadows. Manual.	2.00
Rigby and Meadow's.	.50
Savage. Female Pelvic Org.	12.00
Schultze. Diagrams.	25.00
Swayne's Aphorisms	1.25

OSTEOLOGY.

Holden's Text-book. 1 Vol.	6.00
----------------------------	------

PATHOLOGY & HISTOLOGY.

Gibbes. Practical	1.50
Gilliam. Essentials of.	2.00
Jones and Sieveling.	5.50
Paget's Surgical Path.	7.00
Piersol. Histology, 40 Plts.	6.50
Rindfleisch. General.	2.00
Virchow. Post-mortems.	1.00
Wilkes and Moxon.	6.00

PHARMACY.

Beasley's Druggists' Rec'ts.	2.25
— Formulary.	2.25
Flückiger. Cinchona Barks.	1.50
Kirby. Pharm. of Remedies.	2.25
Mackenzie. Phar. of Throat.	1.25
Merrell's Digest.	4.00
Oldberg. Unofficial Pharm.	3.50
Piesse. Perfumery.	5.50
Proctor. Practical Pharm.	4.50

Roberts. Compend of.	\$2.00
Stewart's Compend.	1.00
Sweringen's Pharm. Lex.	3.00
Tuson. Veterinary Pharm.	2.50

PHYSICAL DIAGNOSIS.

Barth and Roger.	1.00
Bruen's Handbook. 2d Ed.	1.50
West. Exam. of Chest.	1.75

PHYSIOLOGY.

Beale's Bioplasm.	2.25
— Protoplasm.	—
Brubaker's Compend.	1.00
Fulton's Text-book.	4.00
Kirkes' 11th Edition.	—

Cloth, 4.00; Sheep, 5.00	
Landois. Text-book. 2 Vols.	10.00
Sanderson's Laboratory B'k.	5.00
Tyson's Cell Doctrine.	2.00
Yeo's Student's Manual	4.00

POISONS.

Black. Formation of.	1.50
Reese. Toxicology.	4.00
Tanner. Memoranda of.	.75

PRACTICE.

Aitken. 2 Vols. New Ed.	12.00
Beale. Slight Ailments.	1.25
Charteris. Handbook of.	1.25
Cormack. Clinical Lectures.	5.00
Fagge's Practice. 2 Vols.	—
Fenwick's Outlines of.	1.25
Hughes. Compend of. 2 Pts.	2.00
Roberts. Text-book. 5th Ed.	5.00
Tanner's Index of Diseases.	3.00
Warner's Case Taking.	1.75

PRESCRIPTION BOOKS.

Beasley's 3000 Prescriptions.	2.25
— Receipt Book.	2.25
— Formulary.	2.25
Oldberg's New Prescriptions.	1.25
Pereira's Pocket-book.	1.00
Wythe's Dose and Symptom Book.	1.00

RECTUM AND ANUS.

Allingham. Diseases of.	1.25
Cripps. Diseases of.	4.50

SKIN AND HAIR.

Bulkley. The Skin.	.50
Cobbold. Parasites.	5.00
Fox. Atlas of Skin Dis.	20.00
Van Harlingen. Diagnosis and Treatment of Skin Dis.	1.75
Wilson. Skin and Hair.	1.00

STIMULANTS & NARCOTICS.

Anstie. On.	3.00
Kane. Opium Habit, etc.	1.25
Lizars. On Tobacco.	.50
Miller. On Alcohol.	.50
Parrish. Inebriety.	1.25

STOMACH & INDIGESTION.

Allbutt. Visceral Neuroses.	1.50
Edwards. Constipation	—
Fenwick. Atrophy of.	3.25
Gill. Indigestion.	1.25

SURGERY.

Cooper's Dictionary of.	12.00
Druitt's Handbook of.	—
Gamgee. Wounds and Fractures.	3.50
Heath's Operative.	12.00
— Minor.	—
— Surgical Diagnosis.	1.25
— Diseases of Jaws.	4.50
Horwitz. Compend. 2d Ed.	1.00
Hutchinson's Clinical	25.00
Mears. Practical.	3.75
Pye. Surgical Handicraft.	5.00
Roberts. Surgical Delusions.	.50
Watson's Amputations.	5.50

TECHNOLOGICAL BOOKS.

See also Chemistry.	
Gardner. Brewing, etc.	1.75
— Bleaching & Dyeing.	1.75

Gardner. Acetic Acid, etc.	\$1.75
Overman. Mineralogy.	1.00
Piesse. Perfumery, etc.	5.50
Piggott. On Copper.	1.00
Thompson. Physics.	—

THERAPEUTICS.

Biddle. 10th Ed.	4.00
Cohen. Inhalations.	1.25
Headland. Action of Med.	3.00
Kirby. Selected Remedies.	2.25
Mays. Therap. Forces.	1.25
Ott. Action of Medicines.	2.00
Phillips. Vegetable.	7.50
Potter's Compend.	1.00
Waring's Practical.	3.00

THROAT AND VOICE.

Cohen. Throat and Voice.	.50
— Inhalations.	1.25
Dobell. Winter Cough, etc.	3.50
Greenhow. Bronchitis.	1.25
Holmes. Laryngoscope.	1.00
James. Sore Throat	1.25
Mackenzie. Throat & Nose.	6.00
— Larynx.	1.25
— Hay Fever.	.50
— Pharmacopœia.	1.25
Potter. Defects of Speech.	1.00
Thorowgood. Asthma.	1.50

TRANSACTIONS AND REPORTS.

Penna. Hospital Reports.	1.25
Power and Holmes' Reports.	1.25
Trans. College of Physicians.	3.50
— Amer. Surg. Assoc.	4.00

TUMORS AND CANCER.

Hodge. Note-book for.	.50
Thompson. Of the Bladder.	1.75
Wells. Ovarian and Uterine.	7.00
— Abdominal	1.50

URINE & URINARY ORGANS.

Acton. Repro. Organs.	2.00
Beale. Urin. & Renal Dis.	1.75
— Urin. Deposits. Plates.	2.00
Curling. On the Testes.	5.50
Legg. On Urine.	.75
Marshall and Smith. Urine.	1.00
Thompson. Urinary Organs.	1.25
— Surg. of Urin. Organs.	1.25
— Calculous Dis.	1.00
— Lithotomy.	3.50
— Prostate.	Paper, .75
— Tumors of Bladder.	1.75
— Stricture.	2.00
Tyson. Exam. of Urine.	1.50

VENEREAL DISEASES.

Cooper. Syphilis.	3.50
Durkee. Gonorrhœa.	3.50
Hill and Cooper's Manual.	3.00
Lewin. Syphilis.	1.25

VETERINARY PRACTICE.

Armatage's Pocket-book of.	1.25
Tuson's Vet. Pharmacopœia.	2.50

VISITING LISTS AND ACCOUNT BOOKS.

Lindsay and Blakiston's Regular Edition.	1.00 to 3.00
— Perpetual Edition.	1.25
Watson's Led. and Cash Bk.	4.00

WATER.

Fox. Water, Air, Food.	4.00
Frankland. Analysis of.	1.00
MacDonald. " "	2.75

WOMEN, DISEASES OF.

Byford's Text-book.	5.00
— Uterus.	1.25
Courty. Uterus, Ovaries, etc.	6.00
Dillnberger. and Children.	1.50
Duncan. Sterility.	2.00
Galabin. Diseases of.	3.00
Savage. Surgery of Female Pelvic Organs.	12.00
Tilt. Change of Life.	1.25

New Books—Just Ready.

DISEASES OF THE DIGESTIVE ORGANS IN INFANCY AND CHILDHOOD, with Chapters on the Investigation of Disease, and on the General Management of Children. By LOUIS STARR, M.D., Clinical Professor of Diseases of Children in the Hospital of the University of Pennsylvania; Physician to the Children's Hospital, Philadelphia, etc. Octavo. Handsome Cloth, cut or uncut edges, as may be ordered. \$2.50.

A MANUAL OF MIDWIFERY, for Students and Practitioners. By A. LEWIS GALABIN, M.D., F.R.C.P., Obstetric Physician to Guy's Hospital, London, and Professor of Midwifery in the same Institution. 227 Illustrations. Cloth, \$3.00; Leather, \$3.50.

STUDENT'S MANUAL OF VENEREAL DISEASES. Being a Concise Description of those Affections and of their Treatment. By BERKELEY HILL, M.D., Professor of Clinical Surgery in University College, London; Surgeon to University Hospital and to the Lock Hospital; and ARTHUR COOPER, M.D., Late Surgeon to the Lock Hospital. Fourth Edition. Revised. With over 60 Formulæ. 12mo. Cloth, \$1.00.

This edition has been carefully revised and enlarged, without, however, increasing, to any extent, the number of pages.

CHEMISTRY OF THE CARBON COMPOUNDS, OR ORGANIC CHEMISTRY. By PROF. VICTOR VON RICHTER, University of Breslau. Authorized Translation from the Fourth German Edition, by EDGAR F. SMITH, M.A., PH.D., Prof. of Chemistry in Wittenberg College, Springfield, Ohio, formerly in the Laboratories of the University of Pennsylvania; Member of the Chemical Societies of Berlin and Paris, etc. Illustrated. 12mo, 710 pp. Cloth, \$3.00.

Excellent methods of preparing the most typical bodies will be found distributed through the entire volume. These will enable the student to acquaint himself practically with those portions of organic work which are now receiving so much attention. The reactions of the groups of compounds are complete, the methods of analysis described are the most modern and reliable, so that the volume may be depended on as a trustworthy laboratory guide. Numerous references to the literature of the subject will be found upon almost every page.

"It is equaled by nothing, so far as I know, in the language, in comprehensiveness, and clear statement of the matters one wants to know. I have already found in it—clearly stated and with sufficient fullness—points that might well have required the consultation of much larger works. Its arrangement, distinction of type, etc., add greatly to the convenience of the user. It is well up to the present state of that most important and progressive branch of the science. * * * Remarkably full and complete."—*Professor Wm. Hale Herrick, Pennsylvania State College.*

MATERIA MEDICA AND THERAPEUTICS. TENTH EDITION. For Physicians and Students. By JOHN B. BIDDLE, M.D., late Professor of Materia Medica and General Therapeutics in the Jefferson Medical College, Philadelphia. Revised and Enlarged, with Special Reference to Therapeutics and to the Physiological Action of Medicines. By CLEMENT BIDDLE, M.D., U.S.N., and HENRY MORRIS, M.D., Fellow of the College of Physicians of Philadelphia; Demonstrator of Obstetrics and Gynæcology in Jefferson Medical College, etc. Illustrated. Octavo, 524 pp. Cloth, \$4.00; Leather, \$4.75.

From the Preface.—"The favor extended the ninth edition of Biddle's *Materia Medica* has exhausted a large edition in about two years. This appreciation shown their work has been an additional stimulus to the editors in preparing the tenth edition, and encourages them to hope that the fruit of their present labors may be a continuation of this popularity.

A very prominent advantage of the book, and probably its most popular feature, is its small size and practical character; realizing this, and knowing the value of a concise, practical book to the student, endeavor has been made to decrease rather than increase the number of pages, so, though a great deal of new matter has been added, space has been made for it by striking out that which was either obsolete or useless. * * * Therapeutics—the practical application of remedies—has been given more prominence than ever before, and it will compare favorably with other text-books."

P. BLAKISTON, SON & CO., 1012 Walnut St., Philadelphia.

P. BLAKISTON, SON & CO.'S
Medical and Scientific Publications,

No. 1012 WALNUT ST., PHILADELPHIA.

- ABERCROMBIE. Medical Jurisprudence**, for Medical and Legal Students and Practitioners. By JOHN ABERCROMBIE, M.D. 387 pages. Cloth, \$2.50
- ACTON. The Functions and Disorders of the Reproductive Organs** in Childhood, Youth, Adult Age and Advanced Life, considered in their Physiological, Social and Moral Relations. By WILLIAM ACTON, M.D., M.R.C.S. Sixth Edition. 8vo. Cloth, \$2.00
- AITKEN. Science and Practice of Medicine.** By WILLIAM AITKEN, M.D., F.R.S., Professor of Pathology in the Army Medical School, London. Seventh Edition. Revised throughout. 196 Engravings on Wood, and a Map. 2 vols. 8vo. Cloth, \$12.00; Leather, \$14.00
- ALLBUTT. Visceral Neuroses.** On Neuralgia of the Stomach, and Allied Disorders. By T. CLIFFORD ALLBUTT, M.D., F.R.S., Consulting Physician to the Leeds General Infirmary. 8vo. Cloth, \$1.50
- ALLEN. Commercial Organic Analysis.** A Treatise on the Modes of Assaying the Various Organic Chemicals and Products employed in the Arts, Manufactures, Medicine, etc., with Concise Methods for the Detection of Impurities, Adulterations, etc. Second Edition. Revised and Enlarged. By ALFRED ALLEN, F.C.S.
 Vol. I. Alcohols, Ethers, Vegetable Acids, Starch and its Isomers, etc. \$4.50
 Vol. II. Fixed Oils and Fats, Hydrocarbons and Mineral Oils, Phenols and their Derivatives, Coloring Matters, etc. *In Press.*
 Vol. III. Cyanogen Compounds, Alkaloids, Animal Products, etc. *In Press.*
- ALLEN'S New Method** of Recording the Motions of the Soft Palate. By HARRISON ALLEN, M.D., Professor of Physiology University of Pennsylvania. Cloth, .50
- ALLINGHAM. Diseases of the Rectum.** Fistula, Hæmorrhoids, Painful Ulcer, Stricture, Prolapsus, and other Diseases of the Rectum, their Diagnosis and Treatment. By WILLIAM ALLINGHAM, F.R.C.S. Fourth Edition, Enlarged. Illustrated. 8vo. Paper covers, .75; Cloth, \$1.25
London Edition, thick paper and larger type, \$2.00
- ALTHAUS. Medical Electricity.** Theoretical and Practical. Its Use in the Treatment of Paralysis, Neuralgia, and other Diseases. By JULIUS ALTHAUS, M.D. Third Edition, Enlarged. 246 Illustrations. 8vo. Cloth, \$6.00
- ANSTIE. Stimulants and Narcotics.** With special researches on the Action of Alcohol, Ether and Chloroform on the Vital Organism. By FRANCIS E. ANSTIE, M.D. 8vo. Cloth, \$3.00
- ARLT. Diseases of the Eye.** Clinical Studies on Diseases of the Eye. Including the Conjunctiva, Cornea and Sclerotic, Iris and Ciliary Body. By Dr. FERD. RITTER VON ARLT, University of Vienna. Authorized Translation by LYMAN WARE, M.D., Surgeon to the Illinois Charitable Eye and Ear Infirmary, Chicago. Illustrated. 8vo. Cloth, \$2.50
- ARMATAGE. The Veterinarian's Pocket Remembrancer.** Containing concise directions for the Treatment of Urgent or Rare Cases, embracing Semeiology, Diagnosis, Prognosis, Surgery, Therapeutics, etc. 32mo. Cloth, \$1.25
- BALFOUR. Clinical Lectures on Diseases of the Heart and Aorta.** By G. W. BALFOUR, M.D. Illustrated. Second Edition. Cloth, \$5.00
- BARNES. Lectures on Obstetric Operations**, including the Treatment of Hemorrhage, and forming a Guide to Difficult Labor. By ROBERT BARNES, M.D., F.R.C.P. Fourth Edition. Illustrated. 8vo. Cloth, \$3.75

- BARRETT.** *Dental Surgery* for General Practitioners and Students of Medicine and Dentistry. Extraction of Teeth, etc. By A. W. BARRETT, M.D. Illustrated. Cloth, \$1.00
- BARTH AND ROGER.** *Auscultation and Percussion.* 12mo. Cloth, \$1.00
- BARTLEY.** *Medical Chemistry.* A Text-book for Medical and Pharmaceutical Students. By E. H. BARTLEY, M.D., Associate Professor of Chemistry at the Long Island College Hospital; President of the American Society of Public Analysts; Chief Chemist, Board of Health, of Brooklyn, N. Y. With Illustrations, Glossary and Complete Index. 12mo. Cloth, \$2.50
- BEALE.** *On Slight Ailments; their Nature and Treatment.* By LIONEL S. BEALE, M.D., F.R.S., Professor of Practice, King's Medical College, London. Second Edition. Enlarged and Illustrated. Paper covers, .75; Cloth, \$1.25
Finer Edition, Heavy Paper. Extra Cloth, \$1.75
- Urinary and Renal Diseases* and Calculous Disorders. Hints on Diagnosis and Treatment. Demi-8vo. 356 pages. Cloth, \$1.75
- The Use of the Microscope in Practical Medicine.* For Students and Practitioners, with full directions for examining the various secretions, etc., in the Microscope. Fourth Edition. 500 Illustrations. 8vo. Cloth, \$7.50
- How to Work with the Microscope.* A Complete Manual of Microscopical Manipulation, containing a full description of many new processes of investigation, with directions for examining objects under the highest powers, and for taking photographs of microscopic objects. Fifth Edition. Containing over 400 Illustrations, many of them colored. 8vo. Cloth, \$7.50
- Protoplasm; or Matter and Life.* Sixteen Colored Plates. Part I. Dissentient. Part II. Demonstrative. Part III. Suggestive. *New Ed. Preparing.*
- Bioplasm.* A Contribution to the Physiology of Life, or an Introduction to the Study of Physiology and Medicine, for Students. With numerous Illustrations. Cloth, \$2.25
- Life Theories; Their Influence upon Religious Thought.* Six Colored Plates. Cloth, \$2.00
- On Life and Vital Action in Health and Disease.* 12mo. Cloth, \$2.00
- One Hundred Urinary Deposits,* on eight sheets, for the Hospital, Laboratory, or Surgery. New Edition. 4to. Paper, \$2.00
- BEASLEY'S Book of Prescriptions.** Containing over 3100 Prescriptions, collected from the Practice of the most Eminent Physicians and Surgeons—English, French and American; a Compendious History of the Materia Medica, Lists of the Doses of all Official and Established Preparations, and an Index of Diseases and their Remedies. By HENRY BEASLEY. Sixth Edition. Revised and Enlarged. Cloth, \$2.25
- Druggists' General Receipt Book.** Comprising a copious Veterinary Formulary; Recipes in Patent and Proprietary Medicines, Druggists' Nostrums, etc.; Perfumery and Cosmetics; Beverages, Dietetic Articles and Condiments; Trade Chemicals, Scientific Processes, and an Appendix of Useful Tables. Ninth Edition. Revised. Cloth, \$2.25
- Pocket Formulary** and Synopsis of the British and Foreign Pharmacopœias. Comprising Standard and Approved Formulæ for the Preparations and Compounds Employed in Medical Practice. Eleventh Edition. Cloth, \$2.25
- BENTLEY AND TRIMEN'S Medicinal Plants.** A New Illustrated Work, containing full botanical descriptions, with an account of the properties and uses of the principal plants employed in medicine, especial attention being paid to those which are officinal in the British and United States Pharmacopœias. The plants which supply food and substances required by the sick and convalescent are also included. By R. BENTLEY, F.R.S., Professor of Botany, King's College, London, and H. TRIMEN, M.B., F.H.S., Department of Botany, British Museum. Each species illustrated by a colored plate drawn from nature. In forty-two parts. Eight colored plates in each part.
Price reduced to \$1.50 per part, or the complete work handsomely bound in 4 volumes. Half Morocco, Gilt, \$75.00.

- BIBLE HYGIENE; or Health Hints.** By a physician. Written to impart in a popular and condensed form the elements of Hygiene; showing how varied and important are the Health Hints contained in the Bible, and to prove that the secondary tendency of modern Philosophy runs in a parallel direction with the primary light of the Bible. 12mo. Cloth, \$1.00
- BIDDLE'S Materia Medica and Therapeutics.** Tenth Edition. For the Use of Students and Physicians. By Prof. JOHN B. BIDDLE, M.D., Professor of Materia Medica in Jefferson Medical College, Philadelphia. The Tenth Edition, thoroughly revised, and in many parts rewritten, by his son, CLEMENT BIDDLE, M.D., Assistant Surgeon, U. S. Navy, and HENRY MORRIS, M.D., Demonstrator of Obstetrics in Jefferson Medical College, Fellow of the College of Physicians, of Philadelphia, etc. The Botanical portions have been curtailed or left out, and the other sections, on the Physiological action of Drugs, greatly enlarged. Cloth, \$4.00; Leather, \$4.75
- BLACK. Micro-Organisms.** The Formation of Poisons by Micro-Organisms. A Biological study of the Germ Theory of Disease. By G. V. BLACK, M.D., D.D.S. Cloth, \$1.50
- BLOXAM. Chemistry, Inorganic and Organic.** With Experiments. By CHARLES L. BLOXAM, Professor of Chemistry in King's College, London, and in the Department for Artillery Studies, Woolwich. Fifth Edition. With nearly 300 Engravings. 8vo. Cloth, \$3.75; Leather, \$4.75
- Laboratory Teaching.** Progressive Exercises in Practical Chemistry. Intended for use in the Chemical Laboratory, by those who are commencing the study of Practical Chemistry. 4th Edition. 89 Illus. Cloth, \$1.75
- BOWMAN. Practical Chemistry,** including analysis, with about 100 Illustrations. By Prof. JOHN E. BOWMAN. Eighth English Edition. Revised by Prof. BLOXAM, Professor of Chemistry, King's College, London. Cloth, \$2.00
- BRAUNE. Atlas of Topographical Anatomy.** Thirty-four Full-page Plates, Photographed on Stone, from Plane Sections of Frozen Bodies, with many other illustrations. By WILHELM BRAUNE, Professor of Anatomy at Leipzig. Translated and Edited by EDWARD BELLAMY, F.R.C.S., Lecturer on Anatomy, Charing Cross Hospital, London. 4to. Cloth, \$8.00; Half Morocco, \$10.00
- BRUBAKER. Physiology.** A Compend of Physiology, specially adapted for the use of Students and Physicians. By A. P. BRUBAKER, M.D., Demonstrator of Physiology at Jefferson Medical College, Prof. of Physiology, Penn'a College of Dental Surgery, Philadelphia. Third Edition. Revised, Enlarged and Illustrated. "No. 4, ?Quiz-Compend Series?" 12mo. Cloth, \$1.00
Interleaved for the addition of notes, \$1.25
- BRUEN. Physical Diagnosis.** For Physicians and Students. By EDWARD T. BRUEN, M.D., Asst. Professor of Physical Diagnosis in the University of Pennsylvania. Illustrated by Original Wood Engravings. 12mo. 2d Ed. Cloth, \$1.50
- BUCKNILL AND TUKE'S Manual of Psychological Medicine:** containing the Lunacy Laws, the Nosology, Aetiology, Statistics, Description, Diagnosis, Pathology (including morbid Histology) and Treatment of Insanity. By JOHN CHARLES BUCKNILL, M.D., F.R.S., and DANIEL HACK TUKE, M.D., F.R.C.P. Fourth Edition, much enlarged, with twelve lithographic and numerous other illustrations. 8vo. Cloth, \$8.00
- BULKLEY. The Skin in Health and Disease.** By L. DUNCAN BULKLEY, M.D., Attending Physician at the New York Hospital. Illustrated. Cloth, .50
- BURDETT'S Pay Hospitals and Paying Wards throughout the World.** Facts in support of a rearrangement of the system of Medical Relief. By HENRY C. BURDETT, M.D. 8vo. Cloth, \$2.25
- Cottage Hospitals.** General, Fever and Convalescent; their Progress, Management and Work. Second Edition. Rewritten and Enlarged, with Plans and Illustrations. Crown 8vo. Cloth, \$4.50
- BURNETT. Hearing, and How to Keep It.** By CHAS. H. BURNETT, M.D., Prof. of Diseases of the Ear, at the Philadelphia Polyclinic. Illustrated. Cloth, .50

- BUZZARD. Clinical Lectures on Diseases of the Nervous System.** By THOS. BUZZARD, M.D. Illustrated. Octavo. Cloth, \$5.00
- BYFORD. Diseases of Women.** The Practice of Medicine and Surgery, as applied to the Diseases of Women. By W. H. BYFORD, A.M., M.D., Professor of Obstetrics and the Diseases of Women and Children, in the Chicago Medical College. Third Edition. Revised and Enlarged, much of it rewritten, with numerous additional illustrations. 8vo. Cloth, \$5.00; Leather, \$6.00
- On the Uterus.** The Chronic Inflammation and Displacement of the Unimpregnated Uterus. With Illustrations. Paper, .75; Cloth, \$1.25
- CARPENTER. The Microscope and Its Revelations.** By W. B. CARPENTER, M.D., F.R.S. Sixth Edition. Revised and Enlarged, with over 500 Illustrations and Lithographs. Cloth, \$5.50
- CARTER. Eyesight, Good and Bad.** A Treatise on the Exercise and Preservation of Vision. By ROBERT BRUDENELL CARTER, F.R.C.S. Second Edition, with 50 Illustrations, Test Types, etc. 12mo. Paper, .75; Cloth, \$1.25
- CAZEAUX and TARNIER'S Midwifery.** Seventh Revised and Enlarged Edition. With Colored Plates and numerous other Illustrations. The Theory and Practice of Obstetrics; including the Diseases of Pregnancy and Parturition, Obstetrical Operations, etc. By P. CAZEAUX, Member of the Imperial Academy of Medicine, Adjunct Professor in the Faculty of Medicine in Paris. Remodeled and rearranged, with revisions and additions, by S. TARNIER, M.D., Professor of Obstetrics and Diseases of Women and Children in the Faculty of Medicine of Paris. A New American, from the Eighth French and First Italian Edition. Edited and Enlarged by ROBERT J. HESS, M.D., Physician to the Northern Dispensary, Phila., etc. About 1100 pages quarto, with 12 Full-page Plates (five of which are beautifully colored) and over 175 Wood Engravings. Royal Square Octavo. *Sold by subscription only. Circulars and information will be sent, upon application to the Publishers.*
- CHARTERIS. The Practice of Medicine.** A Handbook. By M. CHARTERIS, M.D., Member of Hospital Staff and Professor in University of Glasgow. With Microscopic and other Illustrations. Cloth, \$1.25
- Materia Medica and Therapeutics.** A Manual for Students. *In Press.*
- CHAVASSE. The Mental Culture and Training of Children.** By PYE HENRY CHAVASSE. 12mo. Cloth, \$1.00
- CHURCHILL. Face and Foot Deformities.** By FRED. CHURCHILL, M.D., Ass't Surgeon to the Victoria Hospital for Sick Children, London. Six Plain and Two Colored Lithographs. 8vo. Cloth, \$3.50
- CLEAVELAND'S Pocket Dictionary.** A Pronouncing Medical Lexicon, containing correct Pronunciation and Definition of terms used in medicine and the collateral sciences, abbreviations used in prescriptions, list of poisons, their antidotes, etc. By C. H. CLEAVELAND, M.D. Thirty-first Edition. 16mo. Cloth, .75; Tucks with Pocket, \$1.00
- COBBOLD. A Treatise on the Entozoa of Man and Animals,** including some account of the Ectozoa. By T. SPENCER COBBOLD, M.D., F.R.S. With 85 Illustrations. 8vo. Cloth, \$5.00
- COHEN on Inhalation,** its Therapeutics and Practice, including a Description of the Apparatus Employed, etc. By J. SOLIS COHEN, M.D. With cases and Illustrations. A New Enlarged Edition. 12mo. Paper, .75; Cloth, \$1.25
- The Throat and Voice.** Illustrated. 12mo. Cloth, .50
- COLES. Deformities of the Mouth,** Congenital and Acquired, with Their Mechanical Treatment. By OAKLEY COLES, M.D., D.D.S. Third Edition. 83 Wood Engravings and 96 Drawings on Stone. 8vo. Cloth, \$4.50
- The Dental Student's Note-Book.** A new Edition. 16mo. Cloth, \$1.00
- COOPER'S Surgical Dictionary.** A Dictionary of Practical Surgery and Encyclopædia of Surgical Science. By SAMUEL COOPER. New Edition. By SAMUEL A. LANE, F.R.C.S., assisted by various eminent Surgeons. 2 vols. Cloth, \$12.00

- COOPER on Syphilis and Pseudo-Syphilis.** By ALFRED COOPER, F.R.C.S., Surgeon to the Lock Hospital, to St. Marks, and to the West London Hospitals. Octavo. Cloth, \$3.50
- CORMACK'S Clinical Studies.** Illustrated by Cases observed in Hospital and Private Practice. By Sir JOHN ROSE CORMACK, M.D., K.B., etc. Illustrated. 2 vols. 1127 pp. Cloth, \$5.00
- COURTY. The Uterus, Ovaries, etc.** A Practical Treatise on Diseases of the Uterus, Ovaries and Fallopian Tubes. By Prof. A. COURTY, of Montpellier, France. Translated from the Third Edition, by his pupil and assistant, AGNES McLAREN, M.D., M.K.Q.C.P.I. With a Preface by J. MATHEWS DUNCAN, M.D., LL.D., F.R.S., Obstetric Physician to Saint Bartholomew's Hospital, London. With 431 Illustrations. 8vo. *Sold by Subscription.* Cloth, \$6.00; Leather, \$7.00
- CRIPPS. Diseases of the Rectum and Anus,** including a portion of the Jacksonian Prize Essay on Cancer. By HARRISON CRIPPS, M.D., Ass't Surgeon to St. Bartholomew's Hospital, London. Lithographic Plates and other Illustrations. Cloth, \$4.50
- CULLINGWORTH. A Manual of Nursing, Medical and Surgical.** By CHARLES J. CULLINGWORTH, M.D., Physician to St. Mary's Hospital, Manchester, England. Second Edition. With 18 Illustrations. 12mo. Cloth, \$1.00
- A Manual for Monthly Nurses.** 32mo. Cloth, .50
- CURLING. On the Diseases of the Testis, Spermatic Cord and Scrotum.** By T. B. CURLING, M.D., F.R.S. Fourth Edition, Enlarged and Illustrated. 8vo. Cloth, \$5.50
- DAGUENET'S Ophthalmoscopy.** A Manual for the Use of Students. By Dr. DAGUENET. Translated from the French, by Dr. C. S. JEAFFERSON, F.R.C.S.E. Illustrated. 12mo. Cloth, \$1.50
- DALBY. The Ear.** The Diseases and Injuries of the Ear. By W. B. DALBY, M.D., Surgeon and Lecturer on Aural Surgery, St. George's Hospital. With Illustrations. 12mo. Cloth, \$1.50
- DAY. Diseases of Children.** A Practical and Systematic Treatise for Practitioners and Students. By Wm. H. DAY, M.D. Second Edition. Rewritten and very much Enlarged. 8vo. 752 pp. Price reduced. Cloth, \$3.00; Sheep, \$4.00
- On Headaches.** The Nature, Causes and Treatment of Headaches. Fourth Edition. Illustrated. 8vo. Paper, .75; Cloth, \$1.25
- DILLNBERGER. On Women and Children.** A Handbook of the Treatment of the Diseases Peculiar to Women and Children. By Dr. EMIL DILLNBERGER. 12mo. Cloth, \$1.50
- DOBELL. On Winter Cough, Catarrh, Bronchitis, Emphysema, Asthma, etc.** By HORACE DOBELL, M.D., Senior Physician to the Royal Hospital for Diseases of the Chest. Third Edition. Octavo. Cloth, \$3.50
- DOMVILLE. Manual for Nurses** and others engaged in attending to the sick. By Ed. J. Domville, M.D. Fifth Ed. With Recipes for Sick-room Cookery, etc. Cloth, .75
- DRUITT'S Modern Surgery.** The Surgeon's Vade Mecum; a Manual of Modern Surgery. By ROBERT DRUITT, F.R.C.S. Twelfth Enlarged Edition, with 369 Illustrations. 864 pp. *In Preparation.*
- DULLES. What to Do First, In Accidents and Poisoning.** By C. W. DULLES, M.D. Second Edition, Enlarged, with new Illustrations. Cloth, .75
- DUNCAN, On Sterility in Women.** By J. MATHEWS DUNCAN, M.D., LL.D., Obstetric Physician to St. Bartholomew's Hospital, etc. Octavo. Cloth, \$2.00
- DURKEE, On Gonorrhœa and Syphilis.** By SILAS DURKEE, M.D. Sixth Edition. Revised and Enlarged, with Portrait and Eight Colored Illustrations. Cloth, \$3.50
- ELLIS. Diseases of Children.** A Practical Manual of the Diseases of Children, with a Formulary. By EDWARD ELLIS, M.D. Late Physician to the Victoria Hospital for Children, London. Fourth Edition, Enlarged. Cloth, \$3.00
- What Every Mother Should Know.** 12mo. Cloth, .75

EDWARDS. Bright's Disease. How a Person Affected with Bright's Disease Ought to Live. By JOS. F. EDWARDS, M.D. 2d Ed. Reduced to Cloth, .50

Malaria: What It Means; How to Escape It; Its Symptoms; When and Where to Look For It. Price Reduced. Cloth, .50

Vaccination and Smallpox. Showing the Reasons in favor of Vaccination, and the Fallacy of the Arguments advanced against it, with Hints on the Management and Care of Smallpox patients. Cloth, .50

FAGGE. The Principles and Practice of Medicine. By C. HILTON FAGGE, M.D., F.R.C.P., F.R.M.C.S., Examiner in Medicine, University of London; Physician to, and Lecturer on Pathology in, Guy's Hospital; Senior Physician to Evelina Hospital for Sick Children, etc. Arranged for the press by PHILIP H. PYE-SMITH, M.D., Lect. on Medicine in Guy's Hospital. Including a section on Cutaneous Affections, by the Editor; Chapter on Cardiac Diseases, by SAMUEL WILKES, M.D., F.R.S., and Complete Indexes by ROBERT EDMUND CARRINGTON. 2 vols. Royal 8vo.

Sold only by subscription. Full information upon application to the Publishers.

FENNER. On Vision. Its Optical Defects, the Adaptation of Spectacles, Defects of Accommodation, etc. By C. S. FENNER, M.D. With Test Types and 74 Illustrations. Second Edition, Revised and Enlarged. 8vo. Cloth, \$3.50

FENWICK'S Outlines of Practice of Medicine. With Formulæ and Illustrations. By SAMUEL FENWICK, M.D., 12mo. Cloth, \$1.25

Atrophy of the Stomach and Its Effect on the Nervous Affections of the Digestive Organs. 8vo. Cloth, \$3.25

FLAGG'S Plastics and Plastic Filling; As Pertaining to the Filling of all Cavities of Decay in Teeth below Medium in Structure, and to Difficult and Inaccessible Cavities in Teeth of all Grades of Structure. By J. FOSTER FLAGG, D.D.S., Professor in the Philadelphia Dental College. 8vo. Second Ed. Cloth, \$4.00

FLOWER'S Diagrams of the Nerves of the Human Body. Exhibiting their Origin, Divisions and Connections, with their Distribution to the various Regions of the Cutaneous Surface, and to all the Muscles. By WILLIAM H. FLOWER, F.R.C.S., F.R.S., Hunterian Professor of Comparative Anatomy, and Conservator of the Museum of the Royal College of Surgeons. Third Edition, thoroughly revised. With six Large Folio Maps or Diagrams. 4to. Cloth, \$3.50

FLÜCKIGER. The Cinchona Barks Pharmacognostically Considered. By Professor FRIEDRICH FLÜCKIGER, of Strasburg. Translated by FREDERICK B. POWER, PH.D., Professor of Materia Medica and Pharmacy, University of Wisconsin. With 8 Lithographic Plates. Royal octavo. Cloth, \$1.50

FOTHERGILL. On the Heart and Its Diseases. With Their Treatment. Including the Gouty Heart. By J. MILNER FOTHERGILL, M.D., Member of the Royal College of Physicians of London. 2d Ed. Re-written. 8vo. Cloth, \$3.50

FOX. Water, Air and Food. Sanitary Examinations of Water, Air and Food. By CORNELIUS B. FOX, M.D. 94 Engravings. 8vo. Cloth, \$4.00

FRANKLAND'S Water Analysis. For Sanitary Purposes, with Hints for the Interpretation of Results. By E. FRANKLAND, M.D., F.R.S. Illustrated. 12mo. Cloth, \$1.00

How to Teach Chemistry. Six Lessons to Science Teachers. Edited by G. G. CHALONER, F.C.S. Illustrated. Second Edition. *In Press.*

GALABIN'S Midwifery. A Manual for Students and Practitioners. By A. LEWIS GALABIN, M.D., F.R.C.P., Obstetric Physician to Guy's Hospital, London, and Professor of Midwifery in the same institution. 227 Illustrations. Cloth, \$3.00; Leather, \$3.50

GAMGEE. Wounds and Fractures. The Treatment of Wounds and Fractures. Clinical Lectures, by SAMPSON GAMGEE, F.R.S.E., Consulting Surgeon to Queen's Hospital, Birmingham. 34 Engravings. Second Edition. 8vo. Cloth, \$3.50

- GARDNER'S TECHNOLOGICAL SERIES.** *The Brewer, Distiller and Wine Manufacturer.* A Handbook for all Interested in the Manufacture and Trade of Alcohol and Its Compounds. Edited by JOHN GARDNER, F.C.S. Illustrated. Cloth, \$1.75
- Bleaching, Dyeing, and Calico Printing.* With Formulæ. Illustrated. \$1.75
- Acetic Acid, Vinegar, Ammonia and Alum.* Illustrated. Cloth, \$1.75
- GIBBES'S Practical Histology and Pathology.** By HENEAGE GIBBES, M.B. 12mo. Third Edition. Cloth, \$1.75
- GILL.** *Indigestion:* What it is; What it Leads to; and a New Method of Treating it. By JOHN BEADNELL GILL, M.D. Third Edition. 12mo. Cloth, \$1.25
- GILLIAM'S Pathology.** The Essentials of Pathology; a Handbook for Students. By D. TOD GILLIAM, M.D., Professor of Physiology, Starling Medical College, Columbus, O. With 47 Illustrations. 12mo. Cloth, \$2.00
- GLISAN'S Modern Midwifery.** A Text-book. By RODNEY GLISAN, M.D., Emeritus Professor of Midwifery and Diseases of Women and Children, in Willamette University, Portland, Oregon. 129 Illus. 8vo. Cloth, \$4.00; Leather, \$5.00
- GODLEE'S Atlas of Anatomy.** Illustrating most of the Ordinary Dissections and many not usually practiced by the Student. With References and an Explanatory Text, and 48 Colored Plates. By RICKMAN JOHN GODLEE, M.D., F.R.C.S. A large Folio Volume, with References, and a Separate Volume of Letter-press. The two Volumes, Atlas and Letter-press, Cloth, \$20.00
- GOODHART and STARR'S Diseases of Children.** The Student's Guide to the Diseases of Children. By J. F. GOODHART, M.D., F.R.C.P., Physician to Evelina Hospital for Children, Demonstrator of Morbid Anatomy at Guy's Hospital. Edited, with notes and additions, by LOUIS STARR, M.D., Clinical Professor of Diseases of Children, in the University of Pennsylvania. Cloth, \$3.00; Leather, \$4.00
- GORGAS'S Dental Medicine.** A Manual of Materia Medica and Therapeutics. By FERDINAND J. S. GORGAS, M.D., D.D.S., Professor of the Principles of Dental Science, Dental Surgery and Dental Mechanism, in the Dental Department of the University of Maryland. Second Edition. Enlarged. 8vo. Cloth, \$3.25
- GOWERS, Spinal Cord.** Diagnosis of Diseases of the Spinal Cord. With Colored Plates and Engravings. Third Edition. Enlarged. By WILLIAM R. GOWERS, M.D., Ass't Prof. Clinical Medicine, University College, London. Cloth, \$1.50
- Ophthalmoscopy.** A Manual and Atlas of Ophthalmoscopy. With 16 Colored Autotype and Lithographic Plates and 26 Wood Cuts, comprising 112 Original Illustrations of the Changes in the Eye in Diseases of the Brain, Kidneys, etc. 8vo. Cloth, \$6.00
- Epilepsy and other Chronic Convulsive Diseases:** Their Causes, Symptoms, and Treatment. 8vo. Cloth, \$4.00
- Diagnosis of Diseases of the Brain.** 8vo. Illustrated. Cloth, \$2.00
- GRANVILLE.** *Nerve Vibration and Excitation as Agents in the Treatment of Functional Disorder and Organic Disease.* By J. MORTIMER GRANVILLE, M.D. Illustrated. 8vo. Cloth, \$2.00
- GROSS'S Biography of John Hunter.** John Hunter and His Pupils. By S. D. GROSS, M.D., Professor of Surgery in Jefferson Medical College, Philadelphia. With a Portrait. 8vo. Paper, .75; Cloth, \$1.25
- GREENHOW.** *Chronic Bronchitis,* especially as connected with Gout, Emphysema, and Diseases of the Heart. By E. HEADLAM GREENHOW, M.D. 12mo. Paper, .75; Cloth, \$1.25
- Addison's Disease.** Illustrated by Plates and Reports of Cases. By E. HEADLAM GREENHOW, M.D. 8vo. Cloth, \$3.00
- HABERSHON.** *On Some Diseases of the Liver.* By S. O. HABERSHON, M.D., F.R.C.P., late Senior Physician to Guy's Hospital. A New Edition. Cloth, \$1.50
- HALE.** *On the Management of Children in Health and Disease.** A Book for Mothers. By AMIE M. HALE, M.D. New Enlarged Edition. 12mo. Cloth, .75

- HANDY'S Text-Book of Anatomy** and Guide to Dissections. For the Use of Students. By W. R. HANDY, M.D. 312 Illustrations. 8vo. Cloth, \$3.00
- HARDWICKE. Medical Education** and Practice in All Parts of the World. Containing Regulations for Graduation at the Various Universities throughout the World. By HERBERT JUNIUS HARDWICKE, M.D., M.R.C.P. 8vo. Cloth, \$3.00
- HARE. Tobacco**, Its Physiological and Pathological Effects. 8vo. Illustrated. The Fiske Fund Prize Dissertation for 1885. Paper Covers, .50
- HARLAN. Eyesight** and How to Care for It. By GEORGE C. HARLAN, M.D., Prof. of Diseases of the Eye, Philadelphia Polyclinic. Illustrated. Cloth, .50
- HARLEY. Diseases of the Liver**, With or Without Jaundice. Diagnosis and Treatment. By GEORGE HARLEY, M.D. With Colored Plates and Numerous Illustrations. 8vo. Price reduced. Cloth, \$3.00; Leather, \$4.00
- HARRIS'S Principles and Practice of Dentistry**. Including Anatomy, Physiology, Pathology, Therapeutics, Dental Surgery and Mechanism. By CHAPIN A. HARRIS, M.D., D.D.S., late President of the Baltimore Dental College, author of "Dictionary of Medical Terminology and Dental Surgery." Eleventh Edition. Revised and Edited by FERDINAND J. S. GORGAS, A.M., M.D., D.D.S., author of "Dental Medicine;" Professor of the Principles of Dental Science, Dental Surgery and Dental Mechanism in the University of Maryland. Two Full-page Plates and 744 Illustrations. 994 pages. 8vo. Cloth, \$6.50; Leather, \$7.50
- Medical and Dental Dictionary**. A Dictionary of Medical Terminology, Dental Surgery, and the Collateral Sciences. Fourth Edition, carefully Revised and Enlarged. By FERDINAND J. S. GORGAS, M.D., D.D.S., Prof. of Dental Surgery in the Baltimore College. 8vo. Cloth, \$6.50; Leather, \$7.50
- HARTRIDGE. Refraction**. The Refraction of the Eye. A Manual for Students. By GUSTAVUS HARTRIDGE, F.R.C.S., Consulting Ophthalmic Surgeon to St. Bartholomew's Hospital; Ass't Surgeon to the Royal Westminster Ophthalmic Hospital, etc. 94 Illustrations and Test Types. Second Edition. Cloth, \$2.00
- HARTSHORNE. Our Homes**. Their Situation, Construction, Drainage, etc. By HENRY HARTSHORNE, M.D. Illustrated. Cloth, .50
- HEADLAND'S Action of Medicines**. On the Action of Medicines in the System. By F. W. HEADLAND, M.D. Ninth American Edition. 8vo. Cloth, \$3.00
- HEATH'S Operative Surgery**. A Course of Operative Surgery, consisting of a Series of Plates, Drawn from Nature by M. Lèveillé, of Paris. With Descriptive Text of Each Operation. By CHRISTOPHER HEATH, F.R.C.S., Holme Professor of Clinical Surgery in University College, London. Quarto. Second Edition. Revised. *Sold by Subscription*. Cloth, \$12.00
- Minor Surgery and Bandaging**. Seventh Edition. Revised and Enlarged. With 115 Illustrations. 12mo. *Preparing*.
- Practical Anatomy**. A Manual of Dissections. Sixth London Edition. 24 Colored Plates, and nearly 300 other Illustrations. Cloth, \$5.00
- Injuries and Diseases of the Jaws**. Third Edition. Revised, with over 150 Illustrations. 8vo. Cloth, \$4.50
- Surgical Diagnosis**. Cloth, \$1.25
- HIGGINS' Ophthalmic Practice**. A Handbook for Students and Practitioners. By CHARLES HIGGINS, F.R.C.S. Ophthalmic Assistant Surgeon at Guy's Hospital. Second Edition. 16mo. Cloth, .50
- HILLIER. Diseases of Children**. A Clinical Treatise. By THOMAS HILLIER, M.D. 8vo. Paper, .75; Cloth, \$1.25
- HILL AND COOPER. Venereal Diseases**. The Student's Manual of Venereal Diseases, being a concise description of those Affections and their Treatment. By BERKELEY HILL, M.D., Professor of Clinical Surgery, University College, and ARTHUR COOPER, M.D., Late House Surgeon to the Lock Hospital, London. 4th Edition. 12mo. Cloth, \$1.00

- HODGE'S Note-book** for Cases of Ovarian Tumors. By H. LENOX HODGE, M.D.
With Diagrams. Paper .50
- HODGE on Foeticide** or Criminal Abortion. By HUGH L. HODGE, M.D.
Paper, .30; Cloth, .50
- HOLDEN'S Anatomy.** A Manual of the Dissections of the Human Body. By LUTHER HOLDEN, F.R.C.S. Fifth Edition. Carefully Revised and Enlarged. Specially concerning the Anatomy of the Nervous System, Organs of Special Sense, etc. By JOHN LANGTON, F.R.C.S., Surgeon to, and Lecturer on Anatomy at, St. Bartholomew's Hospital. 208 Illustrations. 8vo.
Oil Cloth Covers, for the Dissecting Room, \$4.50; Cloth, \$5.00; Leather, \$6.00
- Landmarks.** Medical and Surgical. Third London Edition. Revised and Enlarged. *New Edition Preparing.*
- Human Osteology.** Comprising a Description of the Bones, with Colored Delineations of the Attachments of the Muscles. The General and Microscopical Structure of Bone and its Development. Carefully Revised. By the Author and A DORAN, F.R.C.S., with Lithographic Plates and Numerous Illustrations. Sixth Edition. 8vo. Cloth, \$6.00
- HOLDEN. The Sphygmograph.** Its Physiological and Pathological Indications. By EDGAR HOLDEN, M.D. Illustrated. 8vo. Cloth, \$2.00
- HORWITZ'S Compend of Surgery,** including Minor Surgery, Amputations, Fractures, Dislocations, Surgical Diseases, etc., with Differential Diagnosis and Treatment. By ORVILLE HORWITZ, B.S., M.D. Second Edition, Enlarged. 62 Illustrations. 12mo. Cloth, \$1.00
Interleaved for the addition of notes, \$1.25.
- HUFELAND. Long Life.** Art of Prolonging Life. By C. W. HUFELAND. Edited by ERASMUS WILSON, M.D. 12mo. Cloth, \$1.00
- HUGHES. Compend of the Practice of Medicine.** Second Edition. Revised and Enlarged. By DANIEL E. HUGHES, M.D., Demonstrator of Clinical Medicine at Jefferson Medical College, Philadelphia. In two parts:
PART I.—Continued, Eruptive and Periodical Fevers, Diseases of the Stomach, Intestines, Peritoneum, Biliary Passages, Liver, Kidneys, etc., and General Diseases, etc.
PART II.—Diseases of the Respiratory System, Circulatory System and Nervous System; Diseases of the Blood, etc.
Price of each Part, in Cloth, \$1.00; interleaved for the addition of Notes, \$1.25
Physicians' Edition.—In one volume, including a section on Skin Diseases. Full Morocco, \$2.50
- HUNTER. Mechanical Dentistry.** A Practical Treatise on the Construction of the Various kinds of Artificial Dentures, with Formulæ, Receipts, etc. By CHARLES HUNTER, D.D.S. 100 Illustrations. 12mo. Cloth, \$1.50
- HUTCHINSON'S Clinical Surgery.** Consisting of Plates, Photographs, Woodcuts, Diagrams, etc. Illustrating Surgical Diseases, Symptoms and Accidents; also Operations and other Methods of Treatment. With Descriptive Letter-press. By JONATHAN HUTCHINSON, F.R.C.S., Senior Surgeon to the London Hospital, Surgeon to the Moorfields Ophthalmic Hospital. Imperial 4to. Volume 1. (Ten Parts) bound in cloth, complete in itself, \$25.00. Parts Eleven to Seventeen of Volume 2 Now Ready. Each, \$2.50
- JAMES on Sore Throat.** Its Nature, Varieties and Treatment, including its Connection with other Diseases. By PROSSER JAMES, M.D. Fourth Edition, Revised and Enlarged. With Colored Plates and Numerous Woodcuts. 12mo. Paper .75; Cloth, \$1.25
- JONES' Aural Atlas.** An Atlas of Diseases of the Membrana Tympani. Being a Series of Colored Plates, containing 62 Figures. With Explanatory Text. By H. MACNAUGHTON JONES, M.D., Surgeon to the Cork Ophthalmic and Aural Hospital. 4to. Cloth, \$4.00
- Aural Surgery.** A Practical Handbook on Aural Surgery. Illustrated. Second Edition, Revised and Enlarged, with new Wood Engravings. 12mo. Cloth, \$2.75

- JONES and SIEVEKING'S Pathological Anatomy.** A Manual of Pathological Anatomy. By C. HANDFIELD JONES, M.D. and EDWARD H. SIEVEKING, M.D. A New Enlarged Edition. Edited by J. F. PAYNE, M.D. Illus. Cloth, \$5.50
- JONES. Defects of Sight and Hearing,** their Nature, Causes and Prevention. By T. WHARTON JONES, M.D. Second Edition. 16mo. Cloth, .50
- KANE'S Drugs that Enslave.** The Opium, Morphine, Chloral, and Similar Habits. By H. H. KANE, M.D. With Illustrations. Paper, .75; Cloth, \$1.25
- KIDD'S Laws of Therapeutics;** or, the Science and Art of Medicine. By JOSEPH KIDD, M.D. 12mo. Paper, .75; Cloth, \$1.25
- KIRBY. Selected Remedies.** A Pharmacopœia of Selected Remedies, with Therapeutic Annotations, Notes on Alimentation in Disease, Air, Massage, Electricity and other Supplementary Remedial Agents, and a Clinical Index; arranged as a Handbook for Prescribers. By E. A. KIRBY, M.D. Sixth Edition, Enlarged and Revised. 4to. Cloth, \$2.25
- KIRKE'S Physiology.** A Handbook of Physiology. By KIRKE. Eleventh London Edition, Revised and Enlarged. By W. MORRANT BAKER, M.D. 460 Illustrations. Cloth, \$4.00; Leather, \$5.00
- LANDIS' Compend of Obstetrics;** especially adapted to the Use of Students and Physicians. By HENRY G. LANDIS, M.D., Professor of Obstetrics and Diseases of Women, in Starling Medical College, Columbus, Ohio. Second Edition. Revised and Enlarged. With New Illustrations. Cloth, \$1.00; interleaved for the addition of Notes, \$1.25
- LANDOIS. A Manual of Human Physiology;** including Histology and Microscopical Anatomy, with special reference to the requirements of Practical Medicine. By DR. L. LANDOIS, of the University of Greifswald. Translated from the Fourth German Edition, with additions, by WM. STIRLING, M.D., D.Sc., Professor of the Institutes of Medicine in the University of Aberdeen. With very numerous Illustrations. 8vo. 2 Volumes, Cloth, \$10.00
- LEBER AND ROTTENSTEIN. Dental Caries and Its Causes.** An Investigation into the Influence of Fungi in the Destruction of the Teeth. By Drs. LEBER and ROTTENSTEIN. Illustrated. Paper, .75; Cloth, \$1.25
- LEE. The Microtomist's Vade Mecum.** By ARTHUR BOLLES LEE. A Handbook of the Methods of Microscopical Anatomy. 660 Formulæ, etc. 12mo. Cloth, \$3.00
- LEFFMANN'S Organic and Medical Chemistry.** Including Urine Analysis and the Analysis of Water and Food. By HENRY LEFFMANN, M.D., Demonstrator of Chemistry at Jefferson Medical College, Philadelphia. Quiz-Compend Series. 12mo. Cloth, \$1.00. Interleaved for the addition of Notes, \$1.25
- LEGG on the Urine.** Practical Guide to the Examination of the Urine, for Practitioner and Student. By J. WICKHAM LEGG, M.D. Sixth Edition, Enlarged. Illustrated. 12mo. Cloth, .75
- LEWIN on Syphilis.** The Treatment of Syphilis. By Dr. GEORGE LEWIN, of Berlin. Translated by CARL PROEGLER, M.D., and E. H. GALE, M.D., Surgeons U. S. Army. Illustrated. 12mo. Paper, .75; Cloth, \$1.25
- LIEBREICH'S Atlas of Ophthalmoscopy,** composed of 12 Chromo-Lithographic Plates (containing 59 Figures), with Text. Translated by H. R. SWANZY, M.D. Third Edition. 4to. Boards. \$15.00
- LINCOLN. School and Industrial Hygiene.** By D. F. LINCOLN, M.D. Cloth, .50
- LONGLEY'S Pocket Medical Dictionary** for Students and Physicians. Giving the Correct Definition and Pronunciation of all Words and Terms in General Use in Medicine and the Collateral Sciences, with an Appendix, containing Poisons and their Antidotes, Abbreviations Used in Prescriptions, and a Metric Scale of Doses. By ELIAS LONGLEY. Cloth, \$1.00; Tucks and Pocket, \$1.25
- LIZARS. On Tobacco.** The Use and Abuse of Tobacco. By JOHN LIZARS, M.D. 12mo. Cloth, .50
- LÜCKES. Hospital Sisters and their Duties.** By EVA C. E. LÜCKES, Matron to the London Hospital; Author of "Lectures on Nursing." 12mo. Cloth, \$1.00

MACDONALD'S Microscopical Examinations of Water and Air. A Guide to the Microscopical Examination of Drinking Water, with an Appendix on the Microscopical Examination of Air. By J. D. MACDONALD, M.D. With 25 Lithographic Plates, Reference Tables, etc. Second Ed., Revised. 8vo. Cloth, \$2.75

MACKENZIE on the Throat and Nose. Complete. Including the Pharynx, Larynx, Trachea, Œsophagus, Nasal Cavities, etc., etc. By MORELL MACKENZIE, M.D., Senior Physician to the Hospital for Diseases of the Chest and Throat, Lecturer on Diseases of the Throat at London Hospital Medical College, etc.

Vol. I. Including the Pharynx, Larynx, Trachea, etc. 112 Illustrations.

Vol. II. Including the Œsophagus, Nose, Naso-Pharynx, etc. Illustrated.

The two Volumes, Cloth, \$6.00; Leather, \$7.50

Volume II, sold separately, Cloth, \$3.00; Leather, \$4.00

Author's Edition, issued under his supervision, containing all the original Wood Engravings, and the essay on "Diphtheria, its Causes, Nature, and Treatment," formerly published separately.

The Pharmacopœia of the Hospital for Diseases of the Throat and Nose.

Fourth Edition, Enlarged, Containing 250 Formulæ, with Directions for their Preparation and Use. 16mo. Cloth, \$1.25

Growths in the Larynx. Their History, Causes, Symptoms, etc. With Reports and Analysis of one Hundred Cases. With Colored and other Illustrations. 8vo. Paper, .75; Cloth, \$1.25

Hay Fever; Its Etiology and Treatment. Paper, .50

MAC MUNN. On the Spectroscope in Medicine. By CHAS. A. MAC MUNN, M.D. With 3 Chromo-lithographic Plates of Physiological and Pathological Spectra, and 13 Wood Cuts. 8vo. Cloth, \$3.00

MACNAMARA. On the Eye. A Manual of the Diseases of the Eye. By C. MACNAMARA, M.D. Fourth Edition, Carefully Revised; with Additions and Numerous Colored Plates, Diagrams of Eye, Wood-cuts, and Test Types. Demi 8vo. Cloth, \$4.00

MADDEN. Health Resorts for the Treatment of Chronic Diseases. The result of the author's own observations during several years of health travel in many lands. With remarks on climatology and the use of mineral waters. By T. M. MADDEN, M.D. 8vo. Cloth, \$2.50

MANN'S Manual of Psychological Medicine and Allied Nervous Diseases. Their Diagnosis, Pathology, Prognosis and Treatment, including their Medico-Legal Aspects; with chapter on Expert Testimony, and an abstract of the laws relating to the Insane in all the States of the Union. By EDWARD C. MANN, M.D., member of the New York County Medical Society. With Illustrations of Typical Faces of the Insane, Handwriting of the Insane, and Micro-photographic Sections of the Brain and Spinal Cord. Octavo. Cloth, \$5.00; Leather \$6.00

MARSHALL. Anatomical Plates; or Physiological Diagrams. Life Size (4 by 7 feet.) Beautifully Colored. By JOHN MARSHALL, F.R.S. New Edition. Revised and Improved, Illustrating the Whole Human Body.

The Set, 11 Maps, in Sheets, \$50.00

The Set, 11 Maps, on Canvas, with Rollers, and Varnished, 80.00

An Explanatory Key to the Diagrams, .50

No. 1. The Skeleton and Ligaments. No. 2. The Muscles, Joints, and Animal Mechanics. No. 3. The Viscera in Position—The Structure of the Lungs. No. 4. The Organs of Circulation. No. 5. The Lymphatics or Absorbents. No. 6. The Digestive Organs. No. 7. The Brain and Nerves. No. 8. The Organs of the Senses and Organs of the Voice, Plate 1. No. 9. The Organs of the Senses, Plate 2. No. 10. The Microscopic Structure of the Textures. No. 11. The Microscopic Structure of the Textures.

MARSHALL & SMITH. On the Urine. The Chemical Analysis of the Urine. By JOHN MARSHALL, M.D., and Prof. EDGAR F. SMITH, of the Chemical Laboratories, University of Pennsylvania. Illustrated by Phototype Plates. 12mo. Cloth, \$1.00

MARTIN'S Microscopic Mounting. A Manual. With Notes on the Collection and Examination of Objects. 150 Illustrations. By JOHN H. MARTIN. Second Edition, Enlarged. 8vo. Cloth, \$2.75

- MATTHIAS' Legislative Manual.** Rules for Conducting Business in Meetings of Societies, Legislative Bodies, Town and Ward Meetings, etc. By BENJ. MATTHIAS, A.M. Eighteenth Edition. Cloth, .50
- MAYS' Therapeutic Forces ;** or, The Action of Medicine in the Light of the Doctrine of Conservation of Force. By THOMAS J. MAYS, M.D. Cloth, \$1.25
- MEADOWS' Obstetrics.** A Text-Book of Midwifery. Including the Signs and Symptoms of Pregnancy, Obstetric Operations, Diseases of the Puerperal State, etc. By ALFRED MEADOWS, M.D. Third American, from Fourth London Edition. Revised and Enlarged. With 145 Illustrations. 8vo. Cloth, \$2.00
- MEARS' Practical Surgery.** Including: Part I.—Surgical Dressings; Part II.—Bandaging; Part III—Ligations; Part IV.—Amputations. With 490 Illustrations. By J. EWING MEARS, M.D., Demonstrator of Surgery in Jefferson Medical College, and Professor of Anatomy and Clinical Surgery in the Pennsylvania College of Dental Surgery. Second Edition, Revised. 12mo. 794 pages. Cloth, \$3.75; Sheep, \$4.75
- MEDICAL Directory of Philadelphia,** Pennsylvania, Delaware and Southern half of New Jersey, containing lists of Physicians of *all Schools of Practice*, Dentists, Druggists and Chemists, with information concerning Medical Societies, Colleges and Associations, Hospitals, Asylums, Charities, etc. Published Annually. For 1885 now Ready. 12mo. Full Morocco, Gilt edges, \$2.50
- MEIGS. Milk Analysis and Infant Feeding.** A Practical Treatise on the Examination of Human and Cows' Milk, Cream, Condensed Milk, etc., and Directions as to the Diet of Young Infants. By ARTHUR V. MEIGS, M.D., Physician to the Pennsylvania Hospital, Philadelphia. 12mo. Cloth, \$1.00
- MEIGS and PEPPER on Children.** A Practical Treatise on the Diseases of Children. By J. FORSYTH MEIGS, M.D., Fellow of the College of Physicians of Philadelphia, etc., etc., and WILLIAM PEPPER, M.D., Professor of the Principles and Practice of Medicine in the Medical Department, University of Pennsylvania. Seventh Edition. Cloth, \$5.00; Leather, \$6.00
- MENDENHALL'S Vade Mecum.** The Medical Student's Vade Mecum. A Compend of Anatomy, Physiology, Chemistry, The Practice of Medicine, Surgery, Obstetrics, etc. By GEO. MENDENHALL, M.D. 11th Ed. 224 Illus. 8vo.
- MERRELL'S Digest of Materia Medica.** Forming a Complete Pharmacopœia for the use of Physicians, Pharmacists and Students. By ALBERT MERRELL, M.D. Octavo. Half dark Calf, \$4.00
- MILLER and LIZAR'S Alcohol and Tobacco.** Alcohol. Its Place and Power. By JAMES MILLER, F.R.C.S.; and, Tobacco, Its Use and Abuse. By JOHN LIZARS, M.D. The two essays in one volume. Cloth, \$1.00; Separate, each .50
- MORRIS on the Joints.** The Anatomy of the Joints of Man. Comprising a Description of the Ligaments, Cartilages and Synovial Membranes; of the Articular Parts of Bones, etc. By HENRY MORRIS, F.R.C.S. Illustrated by 44 Large Plates and Numerous Figures, many of which are Colored. 8vo. Cloth, \$5.50
- MORTON on Refraction of the Eye.** Its Diagnosis and the Correction of its Errors. With Chapter on Keratotomy and Test Types. By A. MORTON, M.B. Third Edition, Revised. Cloth, \$1.00
- MUTER'S Chemistry.** An Introduction to Pharmaceutical and Medical Chemistry. Part I.—Theoretical and Descriptive. Part II.—Practical and Analytical. By JOHN MUTER, M.D., President of the Society of Public Analysts. Second Edition, Enlarged and Rearranged. The Two Parts in one volume. 8vo. Cloth, \$6.00
- Part II. Practical and Analytical. Bound Separately. Cloth, \$2.50
- OSGOOD. The Winter and Its Dangers.** By HAMILTON OSGOOD, M.D. Cloth, .50
- OTT'S Action of Medicines.** By ISAAC OTT, M.D., late Demonstrator of Experimental Physiology in the University of Pennsylvania. 22 Illus. Cloth, \$2.00
- OVERMAN'S Practical Mineralogy, Assaying and Mining,** with a Description of the Useful Minerals, etc. By FREDERICK OVERMAN, Mining Engineer. Eleventh Edition. 12mo. Cloth, \$1.00

- PACKARD'S Sea Air and Sea Bathing.** By JOHN H. PACKARD, one of the Physicians to the Pennsylvania Hospital, Philadelphia. Cloth, .50
- PAGE'S Injuries of the Spine and Spinal Cord, without apparent Lesion and Nervous Shock.** In their Surgical and Medico-Legal Aspects. By HERBERT W. PAGE, M.D., F.R.C.S. Second Edition, Revised. Octavo. Cloth, \$3.50
- PAGET'S Lectures on Surgical Pathology.** Delivered at the Royal College of Surgeons. By JAMES PAGET, F.R.S. Third Edition. Edited by WILLIAM TURNER, M.D. With Numerous Illustrations. 8vo. Cloth, \$7.00; Leather, \$8.00
- PARKES' Practical Hygiene.** By EDWARD A. PARKES, M.D. The Sixth Revised and Enlarged Edition. With Many Illustrations. 8vo. Cloth, \$3.00
- PARRISH'S Alcoholic Inebriety.** From a Medical Standpoint, with Illustrative Cases from the Clinical Records of the Author. By JOSEPH PARRISH, M.D., President of the Amer. Assoc. for Cure of Inebriates. Paper, .75; Cloth, \$1.25
- PENNSYLVANIA Hospital Reports.** Edited by a Committee of the Hospital Staff: J. M. DACOSTA, M.D., and WILLIAM HUNT. Containing Original Articles by the Staff. With many other Illustrations. Paper, .75; Cloth, \$1.25
- PEREIRA'S Prescription Book.** Containing Lists of Terms, Phrases, Contractions and Abbreviations used in Prescriptions, Explanatory Notes, Grammatical Construction of Prescriptions, Rules for the Pronunciation of Pharmaceutical Terms. By JONATHAN PEREIRA, M.D. Sixteenth Edition. Cloth, \$1.00; Leather, with tucks and pocket, \$1.25
- PHILLIPS' Materia Medica and Therapeutics.** Vegetable Materia Medica and Therapeutics. By CHAS. D. F. PHILLIPS, M.D., F.R.S., Edin., late Lecturer on Materia Medica and Therapeutics, Westminster Hospital, London. Second Edition, Enlarged and Revised. Cloth, \$7.50
- PHYSICIAN'S VISITING LIST.** Published Annually. Thirty-fourth Year of its Publication.

SIZES AND PRICES.

For 25 Patients weekly.	Tucks, pocket and pencil,	\$1.00
50 " "	" " "	1.25
75 " "	" " "	1.50
100 " "	" " "	2.00
50 " " 2 vols.	{ Jan. to June }	2.50
	{ July to Dec. }	
100 " " 2 vols.	{ Jan. to June }	3.00
	{ July to Dec. }	

INTERLEAVED EDITION.

For 25 Patients weekly, interleaved, tucks, pocket, etc.,	1.25
50 " " " " " "	1.50
50 " " 2 vols. { Jan. to June }	3.00
	{ July to Dec. }	.	.	.	

Perpetual Edition, without Dates and with Special Memorandum Pages					
For 25 Patients, interleaved, tucks, pocket and pencil,	\$1.25
50 " " " " " "	1.50

EXTRA Pencils will be sent, postpaid, for 25 cents per half dozen.

- PIESSE'S Art of Perfumery, or the Methods of Obtaining the Odors of Plants, and Instruction for the Manufacture of Perfumery, Dentrifices, Soap, Scented Powders, Cosmetics, etc.** By G. W. SEPTIMUS PIESSE. Fourth Edition. 366 Illustrations. Cloth, \$5.50
- PIERSOL. Normal Histology.** A Synopsis. Adapted to the course at the University of Penn'a. By GEORGE A. PIERSOL, M.D., Demonstrator of Histology. 40 Photo- and Micrographic Plates, containing 200 Figures. \$6.50
- PIGGOTT on Copper Mining and Copper Ore.** With a full Description of the Principal Copper Mines of the United States, the Art of Mining, etc. By A. SNOWDEN PIGGOTT. 12mo. Cloth, \$1.00

- POTTER.** *A Handbook of Materia Medica, Pharmacy and Therapeutics*, including the Action of Medicines, Special Therapeutics, Pharmacology, etc. By SAMUEL O. L. POTTER, M.A., M.D. *Nearly Ready.*
- Speech and Its Defects.** Considered Physiologically, Pathologically and Remedially; being the Lea Prize Thesis of Jefferson Medical College, 1882. Revised and Corrected. 12mo. Cloth, \$1.00
- Compend of Anatomy.** 63 Illustrations. Third Edition, Revised.
- Compend of Visceral Anatomy.** Illustrated. Second Edition.
- Compend of Materia Medica and Therapeutics**, arranged in accordance with the Sixth Revision U. S. Pharmacopœia. Revised Edition, with Index. Price for each, Cloth, \$1.00; Interleaved for taking Notes, \$1.25
- POWER, HOLMES, ANSTIE and BARNES (Drs.)** Reports on the Progress of Medicine, Surgery, Physiology, Midwifery, Diseases of Women and Children, Materia Medica, Medical Jurisprudence, Ophthalmology, etc. Reported for the New Sydenham Society. 8vo. Paper, .75; Cloth, \$1.25
- PRINCE'S Plastic and Orthopædic Surgery.** By DAVID PRINCE, M.D. Containing a Report on the Condition of, and Advance made in, Plastic and Orthopædic Surgery, etc. Numerous Illustrations. 8vo. Cloth, \$4.50
- PROCTER'S Practical Pharmacy.** Lectures on Practical Pharmacy. With 43 Engravings and 32 Lithographic Fac-simile Prescriptions. By BARNARD S. PROCTER. Second Edition. Cloth, \$4.50
- PYE.** *Surgical Handicraft.* A Manual of Surgical Manipulations, Minor Surgery and other Matters connected with the work of Surgeons, Surgeons' Assistants, etc. By WALTER PYE, M.D., Surgeon to St. Mary's Hospital, London. 208 Illustrations. Cloth, \$5.00
- RADCLIFFE on Epilepsy, Pain, Paralysis, and other Disorders of the Nervous System.** By CHARLES BLAND RADCLIFFE, M.D. Illus. Paper, .75; Cloth, \$1.25
- RALFE.** *Diseases of the Kidney and Urinary Derangements.* By C. H. RALFE, M.D., F.R.C.P., Ass't Physician to the London Hospital. Illustrated. 12mo. Volume 3. *Practical Series.* Cloth, \$2.75
- RECORD for the Sick Room.** Designed for the Use of Nurses and others engaged in caring for the Sick. It consists of Blanks, in which may be recorded the Hour, State of Pulse, Temperature, Respiration, Medicines to be Given, Food Taken, etc., together with a List of Directions for the Nurse to pursue in Emergencies. *Sample Pages Free.* One Copy, .25; Per Dozen, \$2.50
- REESE'S Medical Jurisprudence and Toxicology.** A Text-book for Medical and Legal Practitioners and Students. By JOHN J. REESE, M.D., Editor of Taylor's Jurisprudence, Professor of the Principles and Practice of Medical Jurisprudence, including Toxicology, in the University of Pennsylvania Medical and Law Schools. Crown Octavo. Cloth, \$4.00; Leather, \$5.00
- REEVES.** *Bodily Deformities and their Treatment.* A Handbook of Practical Orthopædics. By H. A. REEVES, M.D., Senior Ass't Surgeon to the London Hospital, Surgeon to the Royal Orthopædic Hospital. 228 Illus. Cloth, \$2.25
- REYNOLDS.** *Electricity.* Lectures on the Clinical Uses of Electricity. By J. RUSSELL REYNOLDS, M.D., F.R.S. Second Edition. 12mo. Cloth, \$1.00
- RICHARDSON.** *Long Life, and How to Reach It.* By J. G. RICHARDSON, Prof. of Hygiene, University of Penna. Cloth, .50
- RICHARDSON'S Mechanical Dentistry.** A Practical Treatise on Mechanical Dentistry. By JOSEPH RICHARDSON, D.D.S. Fourth Edition. With 185 Illustrations. 8vo. Cloth, \$4.00; Leather, \$4.75
- RIGBY'S Obstetric Memoranda.** Fourth Edition, Revised. By ALFRED MEADOWS, M.D. 32mo. Cloth, .50
- RINDFLEISCH'S General Pathology.** A Handbook for Students and Physicians. By Prof. EDWARD RINDFLEISCH, of Wurzburg. Translated by WM. H. MERCUR, M.D., of Pittsburgh, Pa., Edited and Revised by JAMES TYSON, M.D., Professor of Morbid Anatomy and Pathology, University of Pennsylvania. Cloth, \$2.00

- RICHTER'S Inorganic Chemistry.** A Text-book for Students. By Prof. VICTOR VON RICHTER, University of Breslau. Second American, from Fourth German Edition. Authorized Translation by EDGAR F. SMITH, M.A., PH.D., Prof. of Chemistry, Wittenberg College, formerly in the Laboratories of the University of Pennsylvania, Member of the Chemical Societies of Berlin and Paris. With 89 Illustrations and a Colored Plate of Spectra. 12mo. 424 pages. Cloth, \$2.00
- Organic Chemistry.** A Text-book for Students. Authorized translation from the Fourth German Edition, by Prof. Edgar F. Smith. Illus. Cloth, \$3.00
- ROBERTS. Practice of Medicine.** The Theory and Practice of Medicine. By FREDERICK ROBERTS, M.D., Professor of Therapeutics at University College, London. Fifth American Edition, thoroughly revised and enlarged, with New Illustrations. 8vo. Cloth, \$5.00; Leather, \$6.00
Recommended as a text-book at the University of Pennsylvania, Yale and Dartmouth Colleges, University of Michigan, and many other Medical Schools.
- Materia Medica and Pharmacy.** A Compend for Students. 12mo. Cloth, \$2.00
- ROBERTS. Surgical Delusions and Follies.** By JOHN B. ROBERTS, M.D., Professor of Anatomy and Surgery, in the Philadelphia Polyclinic. Paper, .25; Cloth, .50
- The Human Brain.** The Field and Limitation of the Operative Surgery of the Human Brain. Illustrated. 8vo. Cloth, \$1.25
- RYAN'S Philosophy of Marriage,** in Its Social, Moral and Physical Relations, and Diseases of the Urinary Organs. By MICHAEL RYAN, M.D. Member of the Royal College of Physicians, London. 12mo. Cloth, \$1.00
- SANDERSON'S Physiological Laboratory.** A Handbook of the Physiological Laboratory. Being Practical Exercises for Students in Physiology and Histology. By J. BURDON SANDERSON, M.D., E. KLEIN, M.D., MICHAEL FOSTER, M.D., F.R.S., and T. LAUDER BRUNTON, M.D. With over 350 Illustrations and Appropriate Letter-press Explanations and References. One Volume. Cloth, \$5.00
Adopted as a Text-book at Yale College and other Medical Schools.
- SANSOM'S Diseases of the Heart.** Valvular Disease of the Heart. By ARTHUR ERNEST SANSOM, M.D. Illustrated. 12mo. Cloth, \$1.25
- On Chloroform.** Its Action and Administration. Paper, .75; Cloth, \$1.25
- SAVAGE. On the Pelvic Organs.** The Surgery, Surgical Pathology and Surgical Anatomy of the Female Pelvic Organs. In a Series of Colored Plates taken from Nature, with Commentaries, Notes and Cases. By HENRY SAVAGE, M.D., F.R.C.S. New Edition. Issued by arrangement with the Author, from the original Plates. 4to. Cloth, \$12.00
- SCHULTZE'S Obstetrical Plates.** Obstetrical Diagrams. Life Size. By Prof. B. S. SCHULTZE, M.D., of Berlin. Twenty in the Set. Colored. In Sheets, \$15.00; Mounted on Rollers, \$25.00
- SIEVEKING on Life Assurance.** The Medical Adviser in Life Assurance. By E. H. SIEVEKING, M.D. Second Edition. Revised. 12mo. Cloth, \$2.00
- SOLLY'S Colorado Springs and Manitou as Health Resorts.** By S. EDWIN SOLLY, M.D., M.R.C.S., Eng. 12mo. Paper cover, .25
- SMITH on Ringworm.** Its Diagnosis and Treatment. By ALDER SMITH, F.R.C.S. With Illustrations. 12mo. Cloth, \$1.00
- SMITH'S Wasting Diseases of Infants and Children.** By EUSTACE SMITH, M.D., F.R.C.P., Physician to the East London Children's Hospital. Fourth London Edition, Enlarged. 8vo. Cloth, \$3.00
- SMYTHE'S Medical Heresies.** Historically Considered. A Series of Critical Essays on the Origin and Evolution of Sectarian Medicine, embracing a Special Sketch and Review of Homœopathy, Past and Present. By GONZALVO C. SMYTHE, A.M., M.D., Professor of the Principles and Practice of Medicine, College of Physicians and Surgeons, Indianapolis, Indiana. 12mo. Cloth, \$1.25
- STAMMER. Chemical Problems,** with Explanations and Answers. By KARL STAMMER. Translated from the 2d German Edition, by Prof. W. S. HOSKINSON, A.M., Wittenberg College, Springfield, Ohio. 12mo. Cloth. .75

- STARR. The Digestive Organs in Childhood.** The Diseases of the Digestive Organs in Infancy and Childhood. With Chapters on the Investigation of Disease and the Management of Children. By LOUIS STARR, M.D., Clinical Professor of Diseases of Children in the Hospital of the University of Pennsylvania; Physician to the Children's Hospital, Philadelphia. 8vo. Cut or uncut edges. Cloth, \$2.50
- STEWART'S Compend of Pharmacy.** Based upon "Remington's Text-Book of Pharmacy." By F. E. STEWART, MD., PH.G., Quiz Master in Chemistry and Theoretical Pharmacy, Philadelphia College of Pharmacy; Demonstrator and Lecturer in Pharmacology, Medico-Chirurgical College, and in Woman's Medical College. Quiz-Compend Series. Cloth, \$1.00; Interleaved for the addition of notes, \$1.25
- STOCKEN'S Dental Materia Medica.** The Elements of Dental Materia Medica and Therapeutics with Pharmacopœia. By JAMES STOCKEN, D.D.S. Third Edition. 12mo. Cloth, \$2.50
- SUTTON'S Volumetric Analysis.** A Systematic Handbook for the Quantitative Estimation of Chemical Substances by Measure, Applied to Liquids, Solids and Gases. By FRANCIS SUTTON, F.C.S. Fourth Edition, Revised and Enlarged, with Illustrations. 8vo. Cloth, \$5.00
- SWAYNE'S Obstetric Aphorisms,** for the Use of Students commencing Midwifery Practice. By JOSEPH G. SWAYNE, M.D. Eighth Edition. Illus. Cloth, \$1.25
- SWERINGEN'S Druggists' Reference Book.** A Pharmaceutical Lexicon or Dictionary of Pharmaceutical Science. Containing Explanations of the various Subjects and Terms of Pharmacy, with appropriate Selections from the Collateral Sciences. Formulæ for Official, Empirical and Dietetic Preparations, etc. By HIRAM V. SWERINGEN, M.D. 8vo. Cloth, \$3.00; Leather, \$4.00
- TAFT'S Operative Dentistry.** A Practical Treatise on Operative Dentistry. By JONATHAN TAFT, D.D.S. Fourth Revised and Enlarged Edition. Over 100 Illustrations. 8vo. Cloth, \$4.25; Leather, \$5.00
- Index of Dental Periodical Literature.** 8vo. *In Press.*
- TANNER'S Index of Diseases and their Treatment.** By THOS. HAWKES TANNER, M.D., F.R.C.P. Second Edition, Revised and Enlarged. By W. H. BROADBENT, M.D. With Additions. Appendix of Formulæ, etc. 8vo. Cloth, \$3.00
- Memoranda of Poisons and their Antidotes and Tests.** Fifth American, from the Last London Edition. Revised and Enlarged. Cloth, .75
- TEMPERATURE Charts** for Recording Temperature, Respiration, Pulse, Day of Disease, Date, Age, Sex, Occupation, Name, etc. Put up in pads; each .50
- TILT'S Change of Life in Women, in Health and Disease.** A Practical Treatise on the Diseases incidental to Women at the Decline of Life. By EDWARD JOHN TILT, M.D. Fourth London Edition. 8vo. Paper cover, .75; Cloth, \$1.25
- THOMPSON. Lithotomy and Lithotripsy.** Practical Lithotomy and Lithotripsy; or, an Inquiry into the best Modes of Removing Stone from the Bladder. By Sir HENRY THOMPSON, F.R.C.S., Emeritus Professor of Clinical Surgery in University College. Third Edition. With 87 Engravings. 8vo. Cloth, \$3.50
- Urinary Organs.** Diseases of the Urinary Organs. Clinical Lectures. Seventh London Edition, Enlarged, with 73 Illustrations. Cloth, \$1.25
- On the Prostate.** Diseases of the Prostate. Their Pathology and Treatment. Fifth London Edition. 8vo. Illustrated. Cloth, \$1.25
- Calculous Diseases.** The Preventive Treatment of Calculous Disease, and the Use of Solvent Remedies. Second Edition. 16mo. Cloth, \$1.00
- Surgery of the Urinary Organs.** Lectures on some Important Points connected with the Surgery of the Urinary Organs. Illustrated. 8vo. Paper, .75; Cloth, \$1.25
- Tumors of the Bladder.** Their Nature, Symptoms and Surgical Treatment. Preceded by a Consideration of the Best Methods of Diagnosing all Forms of Vesical Disease. Illustrated. 8vo. Cloth, \$1.75
- Stricture of the Urethra and Urinary Fistulæ,** their Pathology and Treatment. Fourth Edition. Illustrated. Cloth, \$2.00

- THOMPSON'S Manual of Physics.** A Student's Manual. By SYLVANUS P. THOMPSON, B.A., D.Sc., F.R.A.S., Professor of Experimental Physics in University College, Bristol, England. *Preparing.*
- THOROWGOOD on Asthma.** Its Forms, Nature and Treatment. By JOHN C. THOROWGOOD, M.D. Second Edition. Cloth, \$1.50
- TOMES' Dental Anatomy.** A Manual of Dental Anatomy, Human and Comparative. By C. S. TOMES, D.D.S. 191 Illustrations. 2d Ed. 12mo. Cloth, \$4.25
- TOMES. Dental Surgery.** A System of Dental Surgery. By JOHN TOMES, F.R.S. Fourth Edition, Revised and Enlarged. By C. S. TOMES, D.D.S. With 263 Illustrations. 12mo. *Preparing.*
- TRANSACTIONS of the College of Physicians of Philadelphia.** New Series. Vols. I, II, III, IV, V, Cloth, each, \$2.50. VI, VII, Cloth, each, \$3.50
- TRANSACTIONS American Surgical Association.** Volumes I and II. Illustrated. Edited by J. EWING MEARS, M.D., Recorder of the Association. Royal 8vo. Price of Vol. I, Cloth, \$3.50; Vol. II, Cloth, \$4.00; Vol. III, Cloth, \$3.50
- TRIMBLE. Practical and Analytical Chemistry.** Being a complete course in Chemical Analysis. By HENRY TRIMBLE, PH.G., Professor of Analytical Chemistry in the Philadelphia College of Pharmacy. Illustrated. 8vo. Cloth, \$1.50
- TUKE on Sleep Walking and Hypnotism.** By D. HACK TUKE, M.D., LL.D., F.R.C.P., Co-Editor of the Journal of Mental Diseases. 8vo. Cloth, \$1.75
- History of the Insane in the British Islands.** Cloth, \$3.50
- TURNBULL'S Artificial Anæsthesia.** The Advantages and Accidents of Artificial Anæsthesia; Its Employment in the Treatment of Disease; Modes of Administration; Considering their Relative Risks; Tests of Purity; Treatment of Asphyxia; Spasms of the Glottis; Syncope, etc. By LAURENCE TURNBULL, M.D., PH. G., Aural Surgeon to Jefferson College Hospital, etc. Second Edition, Revised and Enlarged. With 27 Illustrations of Various Forms of Inhalers, etc., and an appendix of over 70 pages, containing a full account of the new local Anæsthetic, Hydrochlorate of Cocaine. 12mo. Cloth, \$1.50
- Hydrochlorate of Cocaine.** 12mo. Paper, .50
- TUSON. Veterinary Pharmacopœia.** Including the Outlines of Materia Medica and Therapeutics. For the Use of Students and Practitioners of Veterinary Medicine. By RICHARD V. TUSON, F.C.S. Third Edition. 12mo. Cloth, \$2.50
- TYSON. Bright's Disease and Diabetes.** With Especial Reference to Pathology and Therapeutics. By JAMES TYSON, M.D., Professor of Pathology and Morbid Anatomy in the University of Pennsylvania. With Colored Plates and many Wood Engravings. 8vo. Cloth, 3.50
- Guide to the Examination of Urine. Fifth Edition.** For the Use of Physicians and Students. With Colored Plates and Numerous Illustrations Engraved on Wood. Fifth Edition. Enlarged and Revised. 12mo. 249 pages. Cloth, \$1.50
- Cell Doctrine.** Its History and Present State. With a Copious Bibliography of the subject. Illustrated by a Colored Plate and Wood Cuts. Second Edition. 8vo. Cloth, \$2.00
- Rindfleisch's Pathology.** Edited by Prof. TYSON. General Pathology; a Handbook for Students and Physicians. By Prof. EDWARD RINDFLEISCH, of Würzburg. Translated by WM. H. MERCUR, M.D. Edited and Revised by JAMES TYSON, M.D., Professor of Morbid Anatomy and Pathology, University of Pennsylvania. Cloth, \$2.00
- VACHER'S Primer of Chemistry.** With Analysis. By ARTHUR VACHER. Cl., .50
- VALENTIN'S Qualitative Analysis.** A Course of Qualitative Chemical Analysis. By WM. G. VALENTIN, F.C.S. Sixth Edition. Revised and Corrected by W. R. HODGKINSON, PH.D. (Würzburg), Fellow of the Institute of Chemistry, and of the Chemical, Physical and Geological Societies of London; Lecturer on Chemistry in the South Kensington Science Schools. Assisted by H. M. CHAPMAN, Assistant Demonstrator of Chemistry in the Royal School of Mines. Illustrated. Octavo. Cloth, \$3.00

- VIRCHOW'S Post-mortem Examinations.** A Description and Explanation of the Method of Performing them in the Dead House of the Berlin Charité Hospital, with especial reference to Medico-legal Practice. By Prof. VIRCHOW. Translated by Dr. T. P. SMITH. Third Edition, with Additions and New Plates. 12mo. Cloth, \$1.00
- VAN HARLINGEN on Skin Diseases.** A Practical Manual of Diagnosis and Treatment. For Students and Practitioners. By ARTHUR VAN HARLINGEN, M.D., Professor of Diseases of the Skin in the Philadelphia Polyclinic; Consulting Physician to the Philadelphia Dispensary for Skin Diseases. Including Formulæ. Illustrated by two Colored Plates, with a number of figures showing the appearance of various lesions. 12mo. Cloth, \$1.75
- WALKER on Intermarriage;** or, the Mode in which, and the Causes why, Beauty, Health and Intellect result from certain Unions; and Deformity, Disease and Insanity from others. Illustrated. 12mo. Cloth, \$1.00
- WARD'S Compend of Chemistry** for Chemical and Medical Students. By G. MASON WARD, M.D., Demonstrator of Chemistry in Jefferson Medical College, Philadelphia. Containing a Table of Elements and Tables for the Detection of Metals in Solutions of Mixed Substances, etc. 12mo. 2d Edition. Cloth, \$1.00
Interleaved for the addition of Notes, \$1.25
- WARING. Practical Therapeutics.** A Manual for Physicians and Students. By Edward J. Waring, M.D. Fourth Edition. Revised, Rewritten and Rearranged by DUDLEY W. BUXTON, M.D., Assistant to the Professor of Medicine, University College, London. Crown Octavo. Cloth, \$3.00; Leather, \$3.50
- WARNER. Case Taking.** A Manual of Clinical Medicine and Case Taking. By FRANCIS WARNER, M.D. Second Edition. Cloth, \$1.75
- WATSON on Amputations** of the Extremities and Their Complications. By B. A. WATSON, A.M., M.D., Surgeon to the Jersey City Charity Hospital and to Christ's Hospital, Jersey City, N. J.; Member of the American Surgical Association. With over 250 Wood Engravings and two Full-page Colored Plates. Octavo. 770 pages. Cloth, \$5.50
- WATSON'S Physician's Ledger and Cash Book.** Based upon and Designed to be Used with Lindsay and Blakiston's Physician's Visiting List (see page 17). *Sample Pages Free.* Price, for 1000 accounts, Leather, \$6.50; 500 accounts, Leather, \$5.00; 500 accounts, Cloth, \$4.00
- WATTS' Inorganic Chemistry.** A Manual of Chemistry, Physical and Inorganic. (Being the 13th Edition of FOWNE'S PHYSICAL AND INORGANIC CHEMISTRY.) By HENRY WATTS, B.A., F.R.S., Editor of the Journal of the Chemical Society; Author of "A Dictionary of Chemistry," etc. With Colored Plate of Spectra and other Illustrations. 12mo. 595 pages. Cloth, \$2.25
- Organic Chemistry.** Second Edition. By WM. A. TILDEN, D.Sc., F.R.S. (Being the 13th Edition of FOWNE'S ORGANIC CHEMISTRY). Illustrated. 12mo. Cloth, \$2.25
- WELCH'S Enteric Fever.** Its Prevalence and Modifications; Ætiology, Pathology and Treatment. By FRANCIS H. WELCH, F.R.C.S. 8vo. Cloth, \$2.00
- WELLS. Abdominal Tumors.** Their Diagnosis and Surgical Treatment. By T. SPENCER WELLS, M.D., F.R.C.S., Consulting Surgeon to the Samaritan Hospital for Women, etc. Illustrated. 8vo. Cloth, \$1.50
- WEST'S How to Examine the Chest.** A Practical Guide for the Use of Students. By SAMUEL WEST, M.D. Oxon., M.R.C.P., Physician to the City of London Hospital for Diseases of the Chest. Illustrated. 32mo. Cloth, \$1.75
- WHITE. The Mouth and Teeth.** By J. W. WHITE, M.D., D.D.S. Editor of the Dental Cosmos. Illustrated. Cloth, .50
- WICKES' Sepulture.** Its History, Methods and Sanitary Requisites. By STEPHEN WICKES, A.M., M.D. Octavo. Cloth, \$1.50
- WILKES' Pathological Anatomy.** Lectures on Pathological Anatomy. By SAMUEL WILKES, F.R.S. Second Edition, Revised and Enlarged by WALTER MOXON, M.D., F.R.S., Physician to and Lecturer at Guy's Hospital, London. Cloth, \$6.00
- The Nervous System.** Lectures on Diseases of the Nervous System, Delivered at Guy's Hospital, London. New Edition. 8vo. Cloth, \$6.00

- WILSON.** *The Summer and Its Diseases.* By JAMES C. WILSON, M.D. Cloth, .50
- WILSON'S Drainage for Health;** or, Easy Lessons in Sanitary Science, with Illustrations. By JOSEPH WILSON, M.D., Medical Director U. S. N. Cloth, \$1.00
- Naval Hygiene;** or, Human Health and Means for Preventing Disease. With Illustrative Incidents from Naval Experience. Illus. Cloth, \$3.00
- WILSON'S Text-Book of Domestic Hygiene** and Sanitary Information. A Guide to Personal and Domestic Hygiene. By GEORGE WILSON, M.D., Medical Officer of Health. Edited by JOS. G. RICHARDSON, M.D., Professor of Hygiene at the University of Pennsylvania. Cloth, \$1.00
- Handbook of Hygiene** and Sanitary Science. With Illustrations. Sixth Edition, Revised and Enlarged. 8vo. Cloth, \$2.75
- WILSON.** *Human Anatomy.* The Anatomist's Vade-mecum. General and Special. By Prof. ERASMUS WILSON. Edited by GEORGE BUCHANAN, Professor of Clinical Surgery in the University of Glasgow; and HENRY E. CLARK, Lecturer on Anatomy at the Royal Infirmary School of Medicine, Glasgow. Tenth Edition. With 450 Engravings (including 26 Colored Plates). Cloth, \$6.00
- Healthy Skin and Hair.** A Practical Treatise. Their Preservation and Management. Eighth Edition. 12mo. Paper, \$1.00
- WILSON'S The Ocean as a Health Resort.** A Handbook of Practical Information as to Sea Voyages, for the Use of Tourists and Invalids. By WM. S. WILSON, M.D. With a Chart showing the Ocean Routes, and Illustrating the Physical Geography of the Sea. Cloth, \$2.50
- WOAKES.** *Post-Nasal Catarrh* and Diseases of the Nose, causing Deafness. By EDWARD WOAKES, M.D., Senior Aural Surgeon to the London Hospital for Diseases of the Throat and Chest. 26 Illustrations. Cloth, \$1.50
- On Deafness, Giddiness** and Noises in the Head, or the Naso-Pharyngeal aspect of Ear Disease. Third Edition. Illustrated. *In Press.*
- WOLFE on the Eye.** A Practical Treatise on Diseases and Injuries of the Eye. By M. WOLFE, M.D., Senior Surgeon to the Glasgow Ophthalmic Institution, etc. With 10 Colored Plates, and Numerous other Illustrations. Octavo. Cloth, \$7.00
- WOLFF.** *Manual of Applied Medical Chemistry* for Students and Practitioners of Medicine. By LAWRENCE WOLFF, M.D., Demonstrator of Chemistry in Jefferson Medical College, Philadelphia. Cloth, \$1.50
- WOOD.** *Brain Work and Overwork.* By Prof. H. C. WOOD, Clinical Professor of Nervous Diseases, University of Pennsylvania. 12mo. Cloth, .50
- WOODMAN and TIDY.** *Medical Jurisprudence.* Forensic Medicine and Toxicology. By W. BATHURST WOODMAN, M.D., Physician to the London Hospital, and CHARLES MEYMOTT TIDY, F.C.S., Professor of Chemistry and Medical Jurisprudence at the London Hospital. With Chromo-lithographic Plates, representing the Appearance of the Stomach in Poisoning by Arsenic, Corrosive Sublimite, Nitric Acid, Oxalic Acid; the Spectra of Blood and the Microscopic Appearance of Human and other Hairs; and 116 other Illustrations. Large Octavo. *Sold by Subscription.* Cloth, \$7.50; Leather, \$8.50
- WRIGHT on Headaches;** their Causes, Nature and Treatment. By HENRY G. WRIGHT, M.D. 12mo. *Ninth Thousand.* Cloth, .50
- WYTHE on the Microscope.** A Manual of Microscopy and Compendium of the Microscopic Sciences, Micro-mineralogy, Biology, Histology and Practical Medicine, with Index and Glossary and the genera of microscopic plants. By JOSEPH H. WYTHE, A.M., M.D. Fourth Edition. 252 Illus. Cloth, \$3.00; Leather, \$4.00
- Dose and Symptom Book.** The Physician's Pocket Dose and Symptom Book. Containing the Doses and Uses of all the Principal Articles of the Materia Medica, and Original Preparations. Sixteenth Revised Edition. Cloth, \$1.00; Leather, with Tucks and Pocket, \$1.25
- YEO'S Manual of Physiology.** A Text-book for Students of Medicine. By GERALD F. YEO, M.D., F.R.C.S., Professor of Physiology in King's College, London. With over 300 carefully printed Illustrations. A Glossary and Complete Index. Crown Octavo. Cloth, \$4.00; Leather, \$5.00

NEW CHEMICAL BOOKS.

PREPARED ESPECIALLY FOR THE WANTS OF MEDICAL, DENTAL AND PHARMACEUTICAL STUDENTS AND PRACTITIONERS.

A Text-Book of Medical Chemistry.

BY E. H. BARTLEY, M.D.,

Associate Professor of Chemistry at the Long Island College Hospital; President of the American Society of Public Analysts; Chief Chemist, Board of Health, of Brooklyn, N. Y., etc.

Illustrated 12mo. Cloth, \$2.50.

This book, written especially for students and physicians, aims to be a text-book for the one and a work of reference for the other. It is practical and concise, dealing only with those parts of chemistry pertaining to medicine; no time being wasted in long descriptions of substances and theories of interest only to the advanced chemical student.

PART I—Treats of Light, Heat and Electricity, which are described at some length, and explanations made and applied to common phenomena. In the subject of light, only so much is given as will explain the theory and use of the spectroscope. In electricity, the principal aim has been to give such information as is needed for the proper understanding, working and care of the medical battery.

PART II—Theoretical Chemistry. Only such portions of the well established principles of modern chemistry as are necessary to an understanding of the subject are given. It has been deemed best to present all these elementary parts first, that the student may be better able to study any set of isolated facts. These theories are presented in a concise, clear way, in a logical order and in a manner which the author has found specially successful in an experience of over twelve years of teaching.

PART III—Treats of the natural history of the elements, of their principal compounds, with their physiological action and toxicology.

PART IV—Organic bodies commonly used in medicine and pharmacy. The principal organic substances derived from animal life are given a place. In the appendix will be found analyses of the principal secretions and tissues, tables of solubilities and of specific gravities, the metric system, and other useful information.

Applied Medical Chemistry.

Containing a description of the apparatus and methods employed in the practice of Medical Chemistry, the Chemistry of Poisons, Physiological and Pathological Analysis, Urinary and Fecal Analysis, Sanitary Chemistry and the Examination of Medicinal Agents, Foods, etc.

BY LAWRENCE WOLFF, M.D.,

Demonstrator of Chemistry in the Jefferson Medical College; Member of the Philadelphia College of Pharmacy and of the Chemical Section of the Franklin Institute, etc.

Octavo, Cloth, \$1.50.

* * * The object of the author of this book is to furnish the practitioner and student a reliable and simple guide for making analyses and examinations of the various medicinal agents, human excretions, secretions, etc., without elaborate apparatus or expensive processes.

Practical and Analytical Chemistry.

Being a complete course in Chemical Analysis, for pharmaceutical and medical students.

BY HENRY TRIMBLE, Ph.G.,

Professor of Analytical Chemistry in the Philadelphia College of Pharmacy.

Illustrated. 8vo. Cloth, \$1.50.

SUMMARY OF CONTENTS. Part I. Practical—Preparation and Properties of Gases, Preparation of Salts, etc. Part II. Section I—Bases. Group I—Potassium, Sodium, Lithium, Ammonium. Group II—Barium, Strontium, Calcium, Magnesium. Group III—Manganese, Zinc, Cobalt, Nickel. Group IV—Iron, Cerium, Chromium, Aluminium. Group V—Arsenic, Antimony, Tin, Gold, Platinum. Group VI—Mercury (ic), Bismuth, Copper, Cadmium. Group VII—Silver, Mercury (ous), Lead. Section II—Acids. Section III—Detection of Bases and Acids. Section IV—Some of the Reactions and Tests of Purity of the more important Organic Compounds. Part III. Quantitative Chemical Analysis. Section I—Gravimetric Estimation. Section II—Volumetric Estimation. There are also a number of useful Tables.

LEFFMANN'S ORGANIC AND MEDICAL CHEMISTRY. Including Urine Analysis and the Analysis of Water and Food. By HENRY LEFFMANN, M.D., Demonstrator of Chemistry at Jefferson Medical College, Philadelphia. 12mo. Cloth, \$1.00; Interleaved for the addition of Notes, \$1.25

P. BLAKISTON, SON & CO., 1012 Walnut St., Philadelphia.

Reissue, in an Improved Form, 12mo size, neatly bound in Flexible Cloth. Each volume sold separately.

PRICE FIFTY CENTS.

American Health Primers.

EDITED BY W. W. KEEN, M.D.,

Fellow of the College of Physicians of Philadelphia.

This Series of American Health Primers is prepared to diffuse as widely and cheaply as possible, among all classes, a knowledge of the elementary facts of Preventive Medicine, and the bearings and applications of the latest and best researches in every branch of Medical and Hygienic Science. They are intended to teach people the principles of Health, and how to take care of themselves, their children, pupils, employes, etc.

Handsome Cloth Binding, 50 cents, each.

Sent, postpaid, upon receipt of price, or may be obtained from any book store.

HEARING, AND HOW TO KEEP IT. With Illustrations. By CHAS. H. BURNETT, M.D., Aurist to the Presbyterian Hospital, Professor in the Philadelphia Polyclinic.

LONG LIFE, AND HOW TO REACH IT. By J. G. RICHARDSON, M.D., Professor of Hygiene in the University of Pennsylvania.

THE SUMMER AND ITS DISEASES. By JAMES C. WILSON, M.D., Lecturer on Physical Diagnosis in Jefferson Medical College.

EYESIGHT, AND HOW TO CARE FOR IT. With Illustrations. By GEO. C. HARLAN, M.D., Surgeon to the Wills (Eye) Hospital, and to the Eye and Ear Department, Pennsylvania Hospital.

THE THROAT AND THE VOICE. With Illustrations. By J. SOLIS COHEN, M.D., Professor of Diseases of the Throat and Chest in the Philadelphia Polyclinic.

THE WINTER AND ITS DANGERS. By HAMILTON OSGOOD, M.D., of Boston, Editorial Staff Boston *Medical and Surgical Journal*.

THE MOUTH AND THE TEETH. With Illustrations. By J. W. WHITE, M.D., D.D.S., of Philadelphia, Editor of the *Dental Cosmos*.

BRAIN WORK AND OVERWORK. By H. C. WOOD, JR., M.D., Clinical Professor of Nervous Diseases in the University of Pennsylvania.

OUR HOMES. With Illustrations. By HENRY HARTSHORNE, M.D., of Philadelphia, formerly Professor of Hygiene in the University of Pennsylvania.

THE SKIN IN HEALTH AND DISEASE. By L. D. BULKLEY, M.D., of New York, Physician to the Skin Department of the Demilt Dispensary and of the New York Hospital.

SEA AIR AND SEA BATHING. By JOHN H. PACKARD, M.D., of Philadelphia, Surgeon to the Pennsylvania and to St. Joseph's Hospitals.

SCHOOL AND INDUSTRIAL HYGIENE. By D. F. LINCOLN, M.D., of Boston, Chairman Department of Health, American Social Science Association.

"Each volume of the 'American Health Primers' *The Inter-Ocean* has had the pleasure to commend. In their practical teachings, learning, and sound sense, these volumes are worthy of all the compliments they have received. They teach what every man and woman should know, and yet what nine-tenths of the intelligent classes are ignorant of, or at best, have but a smattering knowledge of."—*Chicago Inter-Ocean*.

"The series of American Health Primers deserves hearty commendation. These handbooks of practical suggestion are prepared by men whose professional competence is beyond question, and, for the most part, by those who have made the subject treated the specific study of their lives."

P. BLAKISTON, SON & CO., Medical Publishers and Booksellers,
1012 WALNUT STREET, PHILADELPHIA.

Holden's Manual of Anatomy.

FIFTH EDITION, REVISED AND ENLARGED. 208 ILLUSTRATIONS.

A MANUAL OF THE DISSECTIONS OF THE HUMAN BODY.

By LUTHER HOLDEN, M.D., F.R.C.S., Consulting Surgeon to St. Bartholomew's and the Foundling Hospitals, London, and JOHN LANGTON, F.R.C.S., Surgeon to and Lecturer in St. Bartholomew's Hospital. Fifth Edition. Revised and Enlarged, with many new Illustrations. Octavo.


OIL CLOTH BINDING, \$4.50

Cloth, \$5.00; Leather, \$6.00

*** As Holden's Anatomy is the chief "Dissector" now in use, the publishers have put it in an Oil Cloth Binding. This does not retain the odors of the dissecting room, is not easily soiled, and may be washed without damage.*

DIAGRAM OF AXILLA.

(From Holden's Anatomy.)



1. AXILLARY ARTERY.
2. BRACHIAL ARTERY.
3. THORACICA HUMERALIA ARTERY.
4. SUPERIOR THORACIC ARTERY.
5. SUBSCAPULAR ARTERY.
6. DORSALIS SCAPULÆ ARTERY.
7. POSTERIOR CIRCUMFLEX ARTERY.
8. SUPERIOR PROFUNDA ARTERY.
9. POSTERIOR THORACIC NERVE.
10. LONG SUBSCAPULAR NERVE.
11. MEDIAN NERVE.
12. CEPHALIC VEIN.
13. MUSCULO-CUTANEOUS NERVE.
14. TERES MAJOR.

"No student of anatomy can take up this book without being pleased and instructed. Its diagrams are original, striking and suggestive, giving more at a glance than pages of text description. All this is known to those who are already acquainted with this admirable work; but it is simple justice to its value, as a work for careful study and reference, that these points be emphasized to such as are commencing their studies. The text matches the illustrations in directness of practical application and clearness of detail."—*New York Medical Record*, April 15th, 1885.

BY THE SAME AUTHOR.

HUMAN OSTEOLOGY. Comprising a description of the Bones, with Colored Delineations of the Attachments of the Muscles. The General and Microscopical Structure of Bone and its Development. Carefully Revised. By the Author and A. DORAN, F.R.C.S., with Lithographic Plates and Numerous Illustrations. Sixth Edition. 8vo. Cloth, \$6.00


HEATH'S PRACTICAL ANATOMY. A Manual of Dissections. Sixth London Edition. 24 Colored Plates, and nearly 300 other Illustrations. Cloth, \$5.00

P. BLAKISTON, SON & CO., 1012 Walnut St., Philadelphia.

NEW SURGICAL WORKS JUST PUBLISHED.

Watson on Amputations.

Amputations of the Extremities and Their Complications. By B. A. WATSON, A.M., M.D., Surgeon to the Jersey City Charity Hospital, to St. Francis' and to Christ's Hospital, at Jersey City, N. J.; Fellow of the American Surgical Association; Member of the New York Pathological Society, etc. Two full-page Colored Plates, and two hundred and fifty-five Wood Engravings. Octavo. 762 + xix Pages. Handsomely bound in Cloth, \$5.50.



SPECIMEN OF ILLUSTRATIONS IN WATSON'S AMPUTATIONS.

"This volume is an encyclopædic monograph, containing the important facts, theories and arguments relating to amputations of the extremities, and their complications. The author's aim has been to collect facts on this subject from English, French, German and American literature. He does not lay claim to originality but, of course, introduces, in their proper places, those observations which his own experience has led him to make on the general subject. Under the division of 'Complications,' the general subject of the treatment of wounds is discussed, with an outline of the present views on germs and germicides, for, in the words of the preface, the complications of amputation wounds are essentially the same as those which pertain to any solution of continuity involving the various tissues of the body. A great service has been done the profession by the insertion of a translation of Gaupot's and Spellmann's writings on artificial limbs. This is the fullest exposition of the subject we have yet seen in an American text-book. Too much praise cannot be given to the typographical appearance of the work. The illustrations are marvels of clearness, and do, what is not always the case, elucidate the text." —*Medical Record, New York, June 20th, 1885.*

Pye's Surgical Handicraft.

A Manual of Surgical Manipulations, Minor Surgery, Bandaging, Dressing, etc., etc., for the use of General Practitioners and Students. With special chapters on Aural Surgery, Extraction of Teeth, Anæsthetics, etc. By WALTER PYE, F.R.C.S., Surgeon to St. Mary's Hospital and the Victoria Hospital for Sick Children; Examiner in Surgery at the University of Glasgow. 208 Illustrations. Octavo. Cloth, \$5.00.

Heath's Operative Surgery.

A Course of Operative Surgery, consisting of a Series of Colored Plates, each plate containing Several Figures, Drawn from Nature by the Celebrated Anatomical Artist, M. Lèveillé, of Paris, Engraved on Steel under his immediate superintendence, with Descriptive Text of Each Operation, and numerous Wood Engravings. By CHRISTOPHER HEATH, F.R.C.S., Surgeon to University College Hospital, and Holme Professor of Clinical Surgery in University College, London. One Large Quarto Volume. Second Edition, Revised and Enlarged. *Sold by Subscription.* Full information upon application. Cloth, \$12.00.

P. BLAKISTON, SON & CO., 1012 Walnut St., Philadelphia.

Practical Handbooks

FOR THE PHYSICIAN AND MEDICAL STUDENT.

VAN HARLINGEN ON SKIN DISEASES. A Handbook of the Diagnosis and Treatment of Skin Diseases. By ARTHUR VAN HARLINGEN, M.D., Professor of Diseases of the Skin in the Philadelphia Polyclinic; Consulting Physician to the Philadelphia Dispensary for Skin Diseases, and Dermatologist to the Howard Hospital. With colored plates representing the appearance of various lesions. 12mo. Cloth, \$1.75

* * * This is a complete epitome of skin diseases, arranged in alphabetical order, giving the diagnosis and treatment in a concise, practical way. Many prescriptions are given that have never been published in any text-book, and an article incorporated on Diet. The plates do not represent one or two cases, but are composed of a number of figures, accurately colored, showing the appearance of various lesions, and will be found to give great aid in diagnosing.

"This new handbook is essentially a small encyclopædia. * * * Contains a very complete summary of the present state of Dermatology. * * * We heartily commend it for its brevity, clearness and evidently careful preparation."—*Philadelphia Medical Times*.

"The author shows a proper appreciation of the wants of the general practitioner."—*New York Medical Record*.

"It is concisely and intelligently written, and contains many of the best formulas in use for the various forms of Skin Disease."—*New York Medical Times*.

"This is an excellent little book, in which, for ease of reference, the more common diseases of the skin are arranged in alphabetical order, while many good prescriptions are given, together with clear and sensible directions as to their proper application."—*Boston Medical and Surgical Journal*.

"It is just the kind of book that the general practitioner will find most convenient for reference, and we feel confident that it will be appreciated."—*Southern Practitioner*.

RINDFLEISCH'S PATHOLOGY. The Elements of Pathology. By PROF. EDWARD RINDFLEISCH, University of Würzburg. Authorized translation from the first German edition, by WM. H. MERCUR, M.D. (Univ. of Pa.) Revised by JAMES TYSON, M.D., Professor of Pathology and Morbid Anatomy in the University of Pennsylvania. 12mo. Cloth, \$2.00

Prof. Tyson, in his Preface to the American edition, says:—"A high appreciation of Prof. Rindfleisch's work on Pathological Histology, caused me to make careful examination of these 'Elements' immediately after their publication in the original. From such an examination I became satisfied that the book would fill a niche in the wants of the student, as well as of others who may desire to familiarize themselves with general pathological processes, viewed from the most modern standpoint."

BRUEN'S PHYSICAL DIAGNOSIS. Second Edition. A Pocket-book of Physical Diagnosis of the Heart and Lungs; for the Student and Physician. By EDWARD T. BRUEN, Demonstrator of Clinical Medicine in the University of Pennsylvania; Lecturer on Pathology in the Women's Medical College of Philadelphia; 2d Edition, revised, with new original illustrations. 12mo. Cloth, \$1.50

"We consider the description of the manner and rules governing the art of percussion well given. The subject is always a difficult one for beginners, and requires to be well handled in order to be properly understood."—*American Journal of Medical Sciences*.

WOAKES ON CATARRH AND DISEASES OF THE NOSE CAUSING DEAFNESS. By EDWARD WOAKES, M.D., Senior Aural Surgeon to the London Hospital for Diseases of the Throat and Chest. 29 Illustrations. 12mo. Cloth, \$1.50

"Out of the large number of special works on catarrh, there is none for which we have such an unqualified good opinion. * * * The subject is clearly presented. * * * The line of treatment suggested is rational."—*North Carolina Medical Journal*.

P. BLAKISTON, SON & CO., 1012 Walnut St., Philadelphia.

PRACTICAL HANDBOOKS—Continued.

VON ARLT. DISEASES OF THE EYE. Including those of the Conjunctiva, Cornea, Sclerotic, and of the Iris and Ciliary Body. By DR. FERDINAND RITTER VON ARLT, Professor of Ophthalmology in Vienna. Translated by LYMAN WARE, M.D., Surgeon to the Illinois Charitable Eye and Ear Infirmary; Ophthalmic Surgeon to the Presbyterian Hospital, and to the Protestant Orphan Asylum, Chicago. Illustrated. 8vo. 325 pages. Cloth, \$2.50

"His style is condensed but clear, and his pages contain a vast amount of information, couched in such language that it will be equally instructive to the general practitioner and the specialist."—*Philadelphia Medical and Surgical Reporter*, May 30th, 1885.

TYSON ON THE URINE. A Practical Guide to the Examination of Urine. For the Use of Physicians and Students. With Colored Lithographic Plates and Numerous Illustrations Engraved on Wood. Fourth Edition. 12mo. Cloth, \$1.50

"The practical man will find in this little book all that is absolutely necessary for him to know, in order to utilize fully the data supplied by the urine."—*Chicago Medical Journal*.

GILLIAM'S ESSENTIALS OF PATHOLOGY. The Essentials of Pathology. By D. TOD GILLIAM, M.D., Professor of Physiology, Starling Medical College, Columbus, Ohio. With 47 wood engravings. 12mo. Cloth, \$2.00

"The general practitioner will find in this little 12mo a convenient compendium of the current pathology of the day."—*Chicago Medical Journal and Examiner*.

THE PRACTICAL SERIES.

A NEW VOLUME JUST READY.

* * * The volumes of this series written by well known physicians and surgeons, of large private and hospital experience, recognized authorities on the subjects of which they treat, will embrace the various branches of medicine and surgery. They are of a thoroughly practical character, calculated to meet the requirements of the practitioner, and will present the most recent methods and information in a compact shape and at a low price. Bound uniformly, in a handsome and distinctive cloth binding.

DISEASES OF THE KIDNEYS, AND URINARY DERANGEMENTS. By C. H. RALFE, M.A., M.D., F.R.C.P., Assistant Physician to the London Hospital; late Senior Physician to the Seamen's Hospital, Greenwich. 12mo. With Illustrations. 572 pages. *Just Ready.* Cloth, \$2.75

"The object of this volume is to present the student and practitioner with a clear, concise and systematic account of urinary pathology and therapeutics, based upon the latest ascertained facts, and supported by the best authorities. Throughout, the author has endeavored to put prominently forward the characters upon which the diagnosis of the various renal and urinary diseases is founded, and their treatment indicated."—*Extract from The Preface.*

BODILY DEFORMITIES AND THEIR TREATMENT. A Handbook of Practical Orthopædics. By H. A. REEVES, F.R.C.S., Senior Assistant Surgeon and Teacher of Practical Surgery at the London Hospital; Surgeon to the Royal Orthopædic Hospital, etc. 12mo. 228 Illustrations. 460 pages. Cloth, \$2.25

"From what we have already said, it will be seen that Mr. Reeves has given us a trustworthy guide for the treatment of a very extended class of cases. * * * If the other volumes of the Practical Series are as good as this, we shall be agreeably disappointed."—*American Journal of Medical Sciences*, April, 1885.

"The utility of the work now before us cannot be better recommended to the appreciation of the professional reading public, than by recalling that it is the first of its kind, dealing with orthopædics from a modern standpoint."—*Hospital Gazette and Students' Journal*.

DENTAL SURGERY FOR GENERAL PRACTITIONERS AND STUDENTS IN MEDICINE. By ASHLEY W. BARRETT, M.D., M.R.C.S. ENG., Surgeon-Dentist to, and Lecturer on Dental Surgery and Pathology in the Medical School of, London Hospital. 12mo. Illustrated. Cloth, \$1.00

"Replete with an abundance of practical information of unquestionable utility."—*Hospital Gazette and Students' Journal*.

P. BLAKISTON, SON & CO., 1012 Walnut St., Philadelphia.

GOODHART AND STARR

ON

The Diseases of Children.

A Manual for Students and Physicians. By J. F. GOODHART, M.D., Physician to the Evelina Hospital for Children; Assistant Physician to Guy's Hospital, London. American Edition, Revised and Edited by LOUIS STARR, M.D., Clinical Professor of Diseases of Children in the Hospital of the University of Pennsylvania, and Physician to the Children's Hospital, Philadelphia. Containing many new Prescriptions, a list of over 50 Formulæ, conforming to the U. S. Pharmacopœia, and Directions for making Artificial Human Milk, for the Artificial Digestion of Milk, etc.

Just Ready. Demi-Octavo. 738 Pages. Cloth, \$3.00; Leather, \$4.00.

The NEW YORK MEDICAL RECORD, for May, 1885, says:—

"As it is said of some men, so it might be said of some books, that they are 'born to greatness.' This new volume has, we believe, a mission, particularly in the hands of the younger members of the profession. In these days of prolixity in medical literature, it is refreshing to meet with an author who knows both what to say and when he has said it. The work of Dr. Goodhart (admirably conformed, by Dr. Starr, to meet American requirements) is the nearest approach to clinical teaching without the actual presence of clinical material that we have yet seen. It does not discuss mooted questions of Pathology, but is a terse, straightforward account of the author's experience at the bedside of ailing children. Domestic hygiene is awarded its important place in the therapeutics of pediatrics. The details of management so gratefully read by the young practitioner are fully elucidated. Altogether, the book is one of as great practical working value as we have seen for many months."

From the JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION, June 6th, 1885.

"Nothing that concerns disease as found in childhood seems to have escaped the author's attention. From introduction to the end it is replete with valuable information, and one reads it with the feeling that Dr. Goodhart is writing of what he has seen at the bedside. It need scarcely be added that the revisions and additions by the American editor are of much value, neither too full nor too spare, and very judicious."

From the BOSTON MEDICAL AND SURGICAL JOURNAL, June 4th, 1885.

"This work is written in a very agreeable style, carrying weight, from its simplicity and clearness, and the evidently large and matured experience of the author. It is especially adapted to the needs of the practicing physician rather than for the medical student, as with rare discernment it takes up important points in the details of the disease and deals with them practically, leaving the general typical course of the case to the other numerous writers who have already covered the ground in this class of cases. The type and paper are especially to be commended, and the editor, Dr. Starr, can be said to have offered a very attractive book to the medical profession."

From the LONDON MEDICAL TIMES AND GAZETTE, March 7th, 1885.

"Among the great superfluity of medical books which issue from the press we are occasionally gladdened by the reading of some which not only have an unquestionable *raison d'être*, but also as certainly fulfill their purpose. Such a book, we do not hesitate to say, is that which is now before us; and, after a careful perusal, productive of both pleasure and profit, we can assure Dr. Goodhart that he owes no apology for his work, and that if, as he says, he has repeated tales that have been told before, he has repeated them with 'excellent differences.' The book cannot be abstracted. It must, and we think will, be read by all who are interested in or desire to study its subject. We feel sure, moreover, and this forcibly struck us while reading through the work, that those practitioners who have given even half the thought and study to the subject of the diseases of children that Dr. Goodhart has done, will be able to endorse the bulk of his teaching, and will recognize very truly of their own, unwritten, and sometimes unspoken, thoughts and beliefs; an evidence at once of the great value of the book, and an explanation of the undoubted pleasure that every expert and earnest student will inevitably experience in its perusal."

OTHER WORKS ON DISEASES OF CHILDREN:

DAY. DISEASES OF CHILDREN. A Practical and Systematic Treatise for Practitioners and Students. Second Edition. Rewritten and very much Enlarged. 8vo. 752 pp. Cloth, \$3.00; Sheep, \$4.00

MEIGS AND PEPPER ON CHILDREN. A Practical Treatise on the Diseases of Children. Seventh Edition, thoroughly Revised and Enlarged. Cloth, \$6.00; Leather, \$7.00

P. BLAKISTON, SON & CO., 1012 Walnut St., Philadelphia.

? QUIZ-COMPENDS ?

A NEW SERIES OF PRACTICAL MANUALS FOR THE PHYSICIAN AND STUDENT.

*Compiled in accordance with the latest teachings of prominent lecturers
and the most popular Text-books.*

They form a most complete set of Compendes, containing information nowhere else collected in such a condensed, practical shape. The authors have had large experience as quiz masters and attachés of colleges, with exceptional opportunities for noting the most recent advances in therapeutics, methods of treatment, etc. The arrangement of the subjects, illustrations and types, are all of the most improved form, and the size of the books is such that they may be easily carried in the pocket.

Bound in Cloth, each \$1.00. Interleaved, for the Addition of Notes, \$1.25.

No. 1. Human Anatomy. Third Edition. Illustrated. By SAMUEL O. L. POTTER, M.A., M.D., late A. A. Surgeon U. S. Army. With 63 Illus. 3d Revised Ed.

"To those desiring to post themselves hurriedly for examination, this little book will be useful in refreshing the memory."—*New Orleans Med. and Surg. J.*

Nos. 2 and 3. Practice of Medicine. Second Edition. Especially adapted to the use of Students and Physicians. By DANIEL E. HUGHES, M.D., Demonstrator of Clinical Medicine in Jefferson Med. College, Phila. In two parts.

PART I.—Continued, Eruptive and Periodical Fevers, Diseases of the Stomach, Intestines, Peritoneum, Biliary Passages, Liver, Kidneys, etc. (including Tests for Urine), General Diseases, etc.

PART II.—Diseases of the Respiratory System (including Physical Diagnosis), Circulatory System and Nervous System; Diseases of the Blood, etc.

* These little books can be regarded as a full set of notes upon the Practice of Medicine, containing the Synonyms, Definitions, Causes, Symptoms, Prognosis, Diagnosis, Treatment, etc., of each disease, and including a number of prescriptions hitherto unpublished.

No. 4. Physiology, including Embryology. Third Edition. By ALBERT P. BRUBAKER, M.D., Prof. of Physiology, Penn'a College of Dental Surgery; Demonstrator of Physiology in Jefferson Med. College, Phila. Revised, Enlarged and Illustrated.

"This is a well written little book."—*London Lancet.*

No. 5. Obstetrics. Illustrated. Second Edition. For Physicians and Students. By HENRY G. LANDIS, M.D., Prof. of Obstetrics and Diseases of Women, in Starling Medical College, Columbus. Revised Ed. New Illustrations.

"We have no doubt that many students will find in it a most valuable aid."—*The Amer. J. of Obstetrics.*

No. 6. Materia Medica and Therapeutics. Second Revised Edition. With especial Reference to the Physiological Actions of Drugs. For the use of Medical, Dental and Pharmaceutical Students, and Practitioners. Based on the New Revision (Sixth) of the U. S. Pharmacopœia, and including many unofficial remedies. By

SAMUEL O. L. POTTER, M.A., M.D., late A. A. Surg. U. S. Army. Revised Edition, with Index.

"One of the very best we have ever seen."—*Southern Clinic.*

No. 7. Inorganic Chemistry. New Edition. By G. MASON WARD, M.D., Demonstrator of Chemistry in Jefferson Med. College, Phila. Including Table of Elements and various Analytical Tables. New Ed.

"This neat pocket volume is a brief but excellent compend of inorganic chemistry and simple analysis of the metals."—*Pharmaceutical Record, N. Y.*

No. 8. Visceral Anatomy. Illustrated. By SAMUEL O. L. POTTER, M.A., M.D., late A. A. Surg. U. S. Army. With 40 Illus.

"Worthy our recommendation to students, and a ready reference to the busy practitioner."—*Chicago Med. Times.*

No. 9. Surgery. Second Edition. Illustrated. Including Fractures, Wounds, Dislocations, Sprains, Amputations and other operations; Inflammation, Suppuration, Ulcers, Syphilis, Tumors, Shock, etc. Diseases of the Spine, Ear, Eye, Bladder, Testicles, Anus, and other Surgical Diseases. By ORVILLE HORWITZ, A.M., M.D., Resident Physician Pennsylvania Hospital, Phil'a. Second Edition, Revised and Enlarged. With 62 Illustrations.


"Will prove very useful, both to the student and practitioner."—*Valentine Mott, M.D., Asst. to the Prof. of Surgery, Bellevue Hospital, New York.*

No. 10. Organic Chemistry. Including Medical Chemistry, Urine Analysis, and the Analysis of Water and Food, etc. By HENRY LEFFMANN, M.D., Demonstrator of Chemistry in Jefferson Med. College; Prof. of Chemistry in Penn'a College of Dental Surgery, Philadelphia.

"It is a useful and valuable addition to the series of Quiz-Compendes."—*College and Clinical Record.*

No. 11. Pharmacy. Based upon "Remington's Text-Book of Pharmacy." By F. E. STEWART, M.D., PH.G., Quiz-Master at Philadelphia College of Pharmacy.

Bound in Cloth, each \$1.00. Interleaved, for the Addition of Notes, \$1.25.

 These books are constantly revised to keep up with the latest teachings and discoveries.


P. BLAKISTON, SON & CO., 1012 Walnut St., Philadelphia.

The Physician's Visiting List.

(LINDSAY & BLAKISTON'S.)

PUBLISHED ANNUALLY; NOW IN ITS THIRTY-FIFTH YEAR.

Containing Calendar, List of Poisons and Antidotes, Dose Tables rewritten in accordance with the Sixth Revision of the U. S. Pharmacopœia, Marshall Hall's Ready Method in Asphyxia, Lists of New Remedies, Sylvester's Method for Producing Artificial Respiration, with Illustrations; Diagram for Diagnosing Diseases of Heart and Lungs; a new Table for Calculating the Period of Utero-Gestation, etc.

 The Quality of the Leather used in Binding this List has been again Improved, and a Superior Pencil, with Nickel Tip, manufactured especially for it, has been added.

SIZES AND PRICES.

For 25 Patients weekly.	Tucks, pockets, etc.,	\$1.00
50 " "	" "	1.25
75 " "	" "	1.50
100 " "	" "	2.00
50 " " 2 Vols.	{ Jan. to June }	2.50
	{ July to Dec. }	
100 " " 2 Vols.	{ Jan. to June }	3.00
	{ July to Dec. }	

INTERLEAVED EDITION.

For 25 Patients weekly.	Interleaved, tucks, etc.,	1.25
50 " "	" "	1.50
50 " " 2 Vols.	{ Jan. to June }	3.00
	{ July to Dec. }	

PERPETUAL EDITION, without Dates, can be commenced at any time and used until full, similar in style, contents and arrangements to the above.

For 25 Patients, Interleaved, \$1.25
 " 50 " " 1.50

" For completeness, compactness, and simplicity of arrangement it is excelled by none in the market."—*N. Y. Medical Record.*

" The book is convenient in form, not too bulky, and in every respect the very best Visiting List published."—*Canada Medical and Surgical Journal.*

" After all the trials made, there are none superior to it."—*Gaillard's Medical Journal.*

" It has become Standard."—*Southern Clinic.*

" Regular as the seasons comes this old favorite."—*Michigan Medical News.*

" It is quite convenient for the pocket, and possesses every desirable quality."—*Medical Herald.*

" The most popular Visiting List extant."—*Buffalo Medical and Surgical Journal.*

" We have used it for years, and do not hesitate to pronounce it equal, if not superior, to any."—*Southern Clinic.*

" This Visiting List is too well known to require either description or commendation from us."—*Cincinnati Medical News.*

WATSON'S

Physician's Ledger and Cash Book Combined.

WEEKLY AND MONTHLY.


This Ledger is based upon, and designed to be used in connection with, Lindsay & Blakiston's Physician's Visiting List.


PRICES.

Ledger for 1000 accounts,	Leather, \$6.50
" 500 "	" 5.00
" 500 "	Cloth, 4.00

* * Sample pages of both books sent upon application. Books sent, postage prepaid, upon receipt of full price, or can be obtained through any bookseller.

P. BLAKISTON, SON & CO., 1012 Walnut St., Philadelphia.





RJ
886

2
1
on
it
p
on

