

## **A universal formulary.**

### **Contributors**

Griffith, R. Eglesfeld 1798-1850.  
Harvey Cushing/John Hay Whitney Medical Library

### **Publication/Creation**

Philadelphia : Lea & Blanchard, 1850.

### **Persistent URL**

<https://wellcomecollection.org/works/qww5k9zb>

### **License and attribution**

This material has been provided by This material has been provided by the Harvey Cushing/John Hay Whitney Medical Library at Yale University, through the Medical Heritage Library. The original may be consulted at the Harvey Cushing/John Hay Whitney Medical Library at Yale University. where the originals may be consulted.

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection  
183 Euston Road  
London NW1 2BE UK  
T +44 (0)20 7611 8722  
E [library@wellcomecollection.org](mailto:library@wellcomecollection.org)  
<https://wellcomecollection.org>










*Gift of*  
New Haven Dental  
Association

YALE MEDICAL LIBRARY







UNIVERSAL FORMULARY

BY

R. EGLESFELD GRIFFITH, M. D.

UNIVERSAL TORRENTIAL

BY

W. B. BENTLEY, M.D.



# UNIVERSAL FORMULARY:

CONTAINING THE METHODS OF

PREPARING AND ADMINISTERING

OFFICINAL AND OTHER MEDICINES.


THE WHOLE ADAPTED TO

PHYSICIANS AND PHARMACEUTISTS.

BY


R. EGLESFELD GRIFFITH, M.D.

*Selecta sunt quæ medicum nobilitant.*—LINNÆUS



PHILADELPHIA:  
LEA AND BLANCHARD.  
1850.

Entered according to the Act of Congress, in the year 1850, by  
LEA AND BLANCHARD,  
in the Office of the Clerk of the District Court for the Eastern District of Pennsylvania.



RS125  
850G



PHILADELPHIA :  
T. K. AND P. G. COLLINS, PRINTERS.



TO

GEORGE B. WOOD, M. D.,

PROFESSOR OF MATERIA MEDICA AND PHARMACY IN THE UNIVERSITY OF PENNSYLVANIA, ETC.

AND

FRANKLIN BACHE, M. D.,

PROFESSOR OF CHEMISTRY IN JEFFERSON MEDICAL COLLEGE OF PHILADELPHIA, ETC.

AUTHORS OF

"THE DISPENSATORY OF THE UNITED STATES OF AMERICA,"

THIS WORK

IS RESPECTFULLY DEDICATED,

BY THEIR FRIEND,

R. E. G.

Digitized by the Internet Archive  
in 2012 with funding from  
Open Knowledge Commons and Yale University, Cushing/Whitney Medical Library



## PREFACE.

---

THE design of this work is to present a compendious collection of formulæ and pharmaceutic processes, with such additional information as may render it useful to the physician and apothecary; and the principal aim has been to select materials most generally applicable, and of practical utility. The sources from which they have been derived are very numerous, as will be seen by a reference to the various authorities cited.

The introduction contains tables and observations on the weights and measures employed for pharmaceutical purposes in the United States and in foreign countries, and an explanation or vocabulary of the principal abbreviations and Latin terms used by physicians in writing prescriptions, followed by observations on the management of the sick-room, with rules for the administration of the different classes of medicines.

The formulary is arranged alphabetically, according to the pharmaceutic names adopted in the United States Pharmacopœia; but in each formula, the English appellations for the articles composing it are used, and the quantities of these ingredients are expressed in words, and not in the usual pharmaceutic signs.

These innovations may, and probably will, be objected to by many; but we feel convinced that a change has become requisite, and that fewer mistakes would be committed by physicians in writing prescriptions, both in the names of the ingredients and in the quantities, were they given at length, and in common language, instead of in the abbreviated cabalistic terms now used. In France this change has been made, and it is full time that other countries should follow her example.

In this portion of the work, the compiler has derived much important assistance from several of his friends, and is under great obligation to Mr. Wm. Procter, Jr., for numerous formulæ and many useful suggestions. He, also, is much indebted to Dr. Robert Bridges for his attentive revision of the sheets, as well as for some important corrections.



The various tables which follow, it is hoped, will add to the value of the work. The observations and directions on officinal preparations are given in as concise a form as possible; and are, for the most part, condensed from the excellent edition of *Mohr and Redwood's Pharmacy*, as edited by Mr. Wm. Procter. To this is added a short view of the action of poisons, with the best means of obviating their effects.

To facilitate a reference to the contents of the work, copious indices have been added, not only of the formulæ, but of the diseases for which they have been advised.

Notwithstanding the care with which the sheets have been revised, a few typographical errors have escaped notice; but, so far as observed, they are of but little consequence. Should any important ones be noticed, or any articles of practical utility have been omitted, the author would be pleased to be informed of them, that they may be noticed in a new edition, should one be called for.

In the botanical portion, the author has been obliged to cite, in almost every instance, his "Medical Botany." This has arisen from the fact, that no other work of a general character on medicinal plants has been issued from the American press.

PHILADELPHIA,  
March, 1850.

# TABLE OF CONTENTS.

	PAGE
PREFACE . . . . .	vii
INTRODUCTION . . . . .	9
WEIGHTS AND MEASURES . . . . .	9
Weights of the United States and Great Britain . . . . .	9
Foreign Weights . . . . .	12
Measures . . . . .	16
SPECIFIC GRAVITY . . . . .	20
TEMPERATURES FOR CERTAIN PHARMACEUTICAL OPERATIONS . . . . .	26
HYDROMETRICAL EQUIVALENTS . . . . .	26
SPECIFIC GRAVITIES OF SOME OF THE PREPARATIONS OF THE PHARMACOPEIAS . . . . .	31
RELATION BETWEEN DIFFERENT THERMOMETRICAL SCALES . . . . .	32
EXPLANATION OF PRINCIPAL ABBREVIATIONS USED IN FORMULÆ . . . . .	33
VOCABULARY OF WORDS EMPLOYED IN PRESCRIPTIONS . . . . .	34
OBSERVATIONS ON THE MANAGEMENT OF THE SICK ROOM . . . . .	45
Ventilation of the Sick Room . . . . .	45
Temperature of the Sick Room . . . . .	46
Cleanliness in the Sick Room . . . . .	46
Quiet in the Sick Room . . . . .	47
Examination and Preservation of the Excretions . . . . .	47
Administration of Medicine . . . . .	47
Furniture of a Sick Room . . . . .	48
Proper use of Utensils for Evacuations . . . . .	49
DOSES OF MEDICINES . . . . .	50
Age . . . . .	50
Sex . . . . .	51
Temperament . . . . .	51
Idiosyncrasy . . . . .	51
Habit . . . . .	51
State of the System . . . . .	51
Time of Day . . . . .	52
Intervals between Doses . . . . .	52
RULES FOR ADMINISTRATION OF MEDICINES . . . . .	53
Acids . . . . .	53
Antacids . . . . .	53
Antilithics and Lithontriptics . . . . .	54
Antispasmodics . . . . .	54



	PAGE
Anthelmintics . . . . .	54
Cathartics . . . . .	55
Enemata . . . . .	55
Suppositories . . . . .	56
Demulcents or Emollients . . . . .	56
Diaphoretics . . . . .	57
Diluents . . . . .	57
Diuretics . . . . .	57
Emetics . . . . .	57
Emmenagogues . . . . .	58
Epispastics . . . . .	58
Errhines . . . . .	60
Escharotics . . . . .	60
Expectorants . . . . .	60
Narcotics . . . . .	61
Refrigerants . . . . .	61
Sedatives . . . . .	61
Sialagogues . . . . .	61
Stimulants . . . . .	62
Tonics . . . . .	62
MANAGEMENT OF CONVALESCENCE AND RELAPSES . . . . .	63
UNIVERSAL FORMULARY . . . . .	65
DIETETIC PREPARATIONS NOT INCLUDED AMONG THE PREVIOUS PRESCRIPTIONS . . . . .	417
LIST OF INCOMPATIBLES . . . . .	422
POSOLOGICAL TABLE OF THE MOST IMPORTANT MEDICINES . . . . .	426
TABLE OF PHARMACEUTICAL NAMES WHICH DIFFER IN THE UNITED STATES AND BRITISH PHARMACOPŒIAS . . . . .	433
OFFICINAL PREPARATIONS AND DIRECTIONS . . . . .	446
INTERNAL REMEDIES . . . . .	446
Powders . . . . .	446
Pills and Boluses . . . . .	447
Extracts . . . . .	448
Confections, Conserves, Electuaries . . . . .	449
Pulps . . . . .	450
Syrups . . . . .	450
Mellites or Honeys . . . . .	451
Infusions . . . . .	451
Decoctions . . . . .	451
Tinctures . . . . .	451
Wines . . . . .	453
Vinegars . . . . .	454
Mixtures . . . . .	454
Medicated Waters . . . . .	455
Distilled, Essential, or Volatile Oils . . . . .	456
Fixed Oils and Fats . . . . .	460
Alkaloids . . . . .	461
Spirits . . . . .	461

# TABLE OF CONTENTS.

ix

	PAGE
Troches or Lozenges . . . . .	462
Inhalations . . . . .	462
EXTERNAL REMEDIES . . . . .	462
Baths . . . . .	462
Cold Bath . . . . .	463
Cool Bath . . . . .	463
Temperate Bath . . . . .	463
Tepid Bath . . . . .	463
Warm Bath . . . . .	463
Hot Bath . . . . .	464
Shower Bath . . . . .	464
Local Baths . . . . .	465
Vapor Bath . . . . .	465
Warm Air Bath . . . . .	465
Douches . . . . .	465
Medicated Baths . . . . .	466
Affusion . . . . .	466
Sponging . . . . .	467
Fomentations . . . . .	468
Cataplasms, or Poultices . . . . .	468
Lotions, Liniments, Embrocations . . . . .	469
Vesicatories, or Blisters . . . . .	470
Issues . . . . .	470
Setons . . . . .	471
Ointments . . . . .	471
Cerates . . . . .	471
Plasters . . . . .	472
Fumigations . . . . .	472
BLOOD-LETTING . . . . .	474
General Blood-letting . . . . .	474
Venesection . . . . .	474
Arteriotomy . . . . .	476
Topical Blood-letting . . . . .	476
Cupping . . . . .	476
Leeching . . . . .	477
Scarifications . . . . .	478
POISONS . . . . .	479
INDEX OF DISEASES AND THEIR REMEDIES . . . . .	495
INDEX OF PHARMACEUTICAL AND BOTANICAL NAMES . . . . .	511
GENERAL INDEX . . . . .	519



1	General Index
2	Index of Phrases and Botanical Names
3	Index of Diseases and Their Remedies
4	Poisons
5	Remedies
6	Phrases
7	Botanical Names
8	Diseases
9	Remedies
10	Phrases
11	Botanical Names
12	Diseases
13	Remedies
14	Phrases
15	Botanical Names
16	Diseases
17	Remedies
18	Phrases
19	Botanical Names
20	Diseases
21	Remedies
22	Phrases
23	Botanical Names
24	Diseases
25	Remedies
26	Phrases
27	Botanical Names
28	Diseases
29	Remedies
30	Phrases
31	Botanical Names
32	Diseases
33	Remedies
34	Phrases
35	Botanical Names
36	Diseases
37	Remedies
38	Phrases
39	Botanical Names
40	Diseases
41	Remedies
42	Phrases
43	Botanical Names
44	Diseases
45	Remedies
46	Phrases
47	Botanical Names
48	Diseases
49	Remedies
50	Phrases
51	Botanical Names
52	Diseases
53	Remedies
54	Phrases
55	Botanical Names
56	Diseases
57	Remedies
58	Phrases
59	Botanical Names
60	Diseases
61	Remedies
62	Phrases
63	Botanical Names
64	Diseases
65	Remedies
66	Phrases
67	Botanical Names
68	Diseases
69	Remedies
70	Phrases
71	Botanical Names
72	Diseases
73	Remedies
74	Phrases
75	Botanical Names
76	Diseases
77	Remedies
78	Phrases
79	Botanical Names
80	Diseases
81	Remedies
82	Phrases
83	Botanical Names
84	Diseases
85	Remedies
86	Phrases
87	Botanical Names
88	Diseases
89	Remedies
90	Phrases
91	Botanical Names
92	Diseases
93	Remedies
94	Phrases
95	Botanical Names
96	Diseases
97	Remedies
98	Phrases
99	Botanical Names
100	Diseases



# INTRODUCTION.

BEFORE entering upon the main object of the work, some introductory observations are required, on the forms in which remedial substances are administered, the processes to be pursued in their preparation, and on the weights and measures employed in the proper apportionment of the ingredients, and in the regulation of the doses to be administered.

## WEIGHTS AND MEASURES.

These vary much in different countries, and even in the same country. Thus, in Great Britain and the United States, there are three standards of weight recognized and employed: the Avoirdupois, the Troy, and the Apothecaries.

### WEIGHTS OF THE UNITED STATES AND GREAT BRITAIN.

#### AVOIRDUPOIS WEIGHT.

This is the common standard employed in the selling or buying of such articles as are measured by weight, and is also in use among apothecaries in this country, except in the compounding and dispensing of medicines.

	Equal to troy grains.
1 drachm - - - - -	27.34375
16 = 1 ounce - - - - -	437.5
256 = 16 = 1 pound - - - - -	7000.
3584 = 224 = 14 = 1 stone - - - - -	98000.
28672 = 1792 = 112 = 8 = 1 hundred weight - - - - -	784000.
473440 = 35840 = 2240 = 160 = 20 = 1 ton - - - - -	15680000.

In weighing many articles, what is called the short ton, or 2000 pounds, is employed

#### TROY WEIGHT

Is used in the sale of gold, silver, platina, and precious stones.

1 grain
24 = 1 pennyweight
480 = 20 = 1 ounce
5760 = 240 = 12 = 1 pound.

The following are the relative proportions between the troy and the avoirdupois:—

One pound troy is equivalent to 0.822857 pound avoirdupois, or 13 ounces, 2 drachms, 1 scruple, 9.5 grains.

One ounce troy is equivalent to 1 ounce, 1 drachm, 1 scruple, 6.225 grains avoirdupois.

One pound avoirdupois is equivalent to 1.21577 pound troy, or 1 pound, 2 ounces, 4 drachms, and 2 scruples.

One ounce avoirdupois is equivalent to 7 drachms, 17.5 grains troy.

144 pounds avoirdupois are equivalent to 175 pounds troy.

175 ounces troy are equivalent to 192 ounces avoirdupois.

The following table of equivalents in troy and avoirdupois weight, derived from Redwood's edition of Gray's Supplement to the Pharmacopœia, will be found useful in making these calculations.



## EQUIVALENTS IN TROY AND AVOIRDUPOIS WEIGHT.

Troy grains.	TROY.				AVOIRDUPOIS.		
	lbs.	oz.	drs.	grs.	lbs.	oz.	grs.
60			1				60
120			2				120
240			4				240
437.5			7	17.5		1	
480		1				1	42.5
875		1	6	35		2	
960		2				2	85
1312.5		2	5	52.5		3	
1440.		3				3	127.5
1750.		3	5	10		4	
1920.		4				4	170
2187.5		4	4	27.5		5	
2400		5				5	212.5
2625.0		5	3	45.		6	
2880.		6				6	255
3062.5		6	3	2.5		7	
3360		7				7	297.5
3500		7	2	20.		8	
3840		8				8	340
3937.5		8	1	37.5		9	
4320.		9				9	382.5
4375.		9	0	55.		10	
4800		10				10	425
4812.5		10		12.5		11	
5250		10	7	30.		12	
5280		11				12	30
5687.5		11	6	47.5		13	
5760.	1	12				13	72.5
6125.	1	0	6	5.		14	
6562.5	1	1	5	22.5		15	
7000	1	2	4	40.	1	16	
7680	1	4			1	1	242.5
9600	1	8			1	5	422.5
10500	1	9	7		1	8	
11520	2				1	10	145
14000	2	5	1	20	2		
17280	3				2	7	217.5
21000	3	7	6	0	3		
23040	4				3	4	290
28000	4	10	2	40	4		
28800	5				4	1	362.5
34560	6				4	14	435
35000	6	0	7	20	5		
40320	7				5	12	70
42000	7	3	4	0	6		
46080	8				6	9	142.5
49000	8	6	0	40	7		
51840	9				7	6	215
56000	9	8	5	20	8		
57600	10				8	3	287.5
63000	10	11	2	0	9		

Troy grains.	TROY.				AVOIRDUPOIS.		
	lbs.	oz.	drs.	grs.	lbs.	oz.	grs.
63360	11				9	0	360
69120	12				9	13	432.5
70000	12	1	6	40	10		
74880	13				10	11	67.5
77000	13	4	3	20	11		
80640	14				11	8	140
84000	14	7	0	0	12		
86400	15				12	5	212.5
91000	15	9	4	40	13		
92160	16				13	2	285
97920	17				13	15	357.5
98000	17	0	1	20	14		
103680	18				14	12	430
105000	18	2	6	0	15		
109440	19				15	10	65
112000	19	5	2	40	16		
115200	20				16	7	137.5
119000	20	7	7	20	17		
120960	21				17	4	210
126000	21	10	4	0	18		
126720	22				18	1	282.5
132480	23				18	14	355
133000	23	1	0	40	19		
138240	24				19	11	427.5
140000	24	3	5	20	20		
144000	25				20	9	62.5
147000	25	6	2	0	21		
149760	26				21	6	135
154000	26	8	6	40	22		
155520	27				22	3	207.5
161000	27	11	3	20	23		
161280	28				23	0	280
167040	29				23	13	352.5
168000	29	2	0	0	24		
172800	30				24	10	425
175000	30	4	4	40	25		
178560	31				25	8	59
182000	31	7	1	20	26		
184320	32				26	5	131.5
189000	32	9	6	0	27		
190080	33				27	2	204
195840	34				27	15	276.5
196000	34	0	2	40	28		
201600	35				28	12	149
203000	35	2	7	20	29		
207360	36				29	9	421.5
210000	36	5	4	0	30		
230400	40				32	14	275
280000	48	7	2	40	40		
288000	50				41	2	125
345600	60				49	5	412.5
350000	60	9	1	20	50		
403200	70				57	9	262.5
420000	72	11	0	0	60		



Troy grains.	TROY.				AVOIRDUPOIS.		
	lbs.	oz.	drs.	grs.	lbs.	oz.	grs.
460800	80				65	13	113
490000	85	0	6	40	70		
518400	90				74	0	400.5
560000	97	2	5	20	80		
576000	100				82	4	250.5
630000	109	4	4	0	90		
645120	112				92	2	245
700000	121	6	2	40	100		
784000	136	1	2	40	112		

When applied to the compounding or dispensing of medicines, this standard of weight is known as Apothecaries' weight, and differs from the last in the subdivision of the ounce.

#### APOTHECARIES' WEIGHT.

1 grain  
 20 = 1 scruple  
 60 = 3 = 1 drachm  
 480 = 24 = 8 = 1 ounce  
 3760 = 288 = 96 = 12 = 1 pound.

To designate these divisions, the following marks are generally used in prescriptions: a grain, gr.; a scruple,  $\mathfrak{S}$ ; a drachm,  $\mathfrak{z}$ ; an ounce,  $\mathfrak{℥}$ ; a pound,  $\mathfrak{℔}$ .

#### FOREIGN WEIGHTS.

FRANCE.—Anterior to the French Revolution of 1789, the scale of weight used was the *poids de marc*, the unit of which was the pound of Charlemagne, which was equivalent to 7561 troy grains, and was divided as follows:

#### POIDS DE MARC.

	Troy grains.	Grammes.
1 grain - - - - -	= 0.8203	= 0.0531
24 = 1 scruple - - - - -	= 19.687	= 1.274
72 = 3 = 1 gros - - - - -	= 59.070	= 3.824
576 = 24 = 8 = 1 once - - - - -	= 472.542	= 30.594
4608 = 192 = 64 = 8 = 1 marc - - - - -	= 3780.500	= 244.750
6612 = 288 = 96 = 12 = 1 livre medicinal - - - - -	= 5670.750	= 367.125
9216 = 384 = 128 = 16 = 1 livre marchand or poid de marc - - - - -	= 7561.000	= 489.500

When the decimal system was introduced by the National Assembly, a new series of measures was adopted, termed the metrical, in which the *metre*, or the ten-millionth part of a quarter of the meridian of the earth, is taken as the unit; this is divided into ten parts, each of which is called a *decimetre*, which in turn is divided into ten *centimetres*. A cubic decimetre is assumed as the unit of measures of capacity, and termed a *litre*. The unit of weight is called a *gramme*, and is a cubic centimetre of distilled water at 39.5 Fahr.

#### METRICAL WEIGHT.

	Troy grains.
1 milligramme - - - - -	= .015444
10 = 1 centigramme - - - - -	= .15444
100 = 10 = 1 decigramme - - - - -	= 1.5444
1000 = 100 = 10 = 1 gramme - - - - -	= 15.444
10000 = 1000 = 100 = 10 = 1 decagramme - - - - -	= 154.44
100000 = 10000 = 1000 = 100 = 10 = 1 hectogramme - - - - -	= 1544.4
1000000 = 100000 = 10000 = 1000 = 100 = 10 = 1 kilogram. - - - - -	= 15444.



This system of weights, although adopted by the scientific men of France, was not favorably received by the people generally, who obstinately adhered to the old system of the *poids de marc*; and, although many stringent laws were passed, from time to time, to render the metrical system obligatory, the government in 1812 was forced to make an attempt to amalgamate the two systems, by altering the *livre*, and making it equal to half a kilogramme, assuming this as the unit, and calculating the other divisions from it, according to the old nomenclature. The following table shows the equivalent of the two scales and of avoirdupois:—

## FRENCH WEIGHTS OF 1812.

French weights of 1812.	Metrical weight grammes.	lb.	oz.	dr.	grains.
1 livre =	500	= 1	1	10	11.07
$\frac{1}{2}$ " =	250	=	8	13	5.53
$\frac{1}{4}$ " =	125	=	4	6	16.18
2 once =	62.5	=	2	3	8.9
1 " =	31.25	=	1	1	17.71
$\frac{1}{2}$ " =	15.625	=		8	22.52
2 gros =	7.812	=		4	11.26
1 " =	3.906	=		2	5.63
$\frac{1}{2}$ " =	1.9021	=		1	2.81
1 grain =	0.0542	=			0.90

The adoption of this system was not made obligatory upon the pharmaciens by law until the year 1827; and indeed it appears never to have been generally adopted, the greatest confusion having prevailed with regard to the weights and measures used in the preparation of medicine, as well as in commerce generally, up to the year 1840. In July, 1837, a law was passed which definitively abolished the use of all other weights and measures, excepting those of the metrical or decimal system, from and after August, 1840. The METRICAL WEIGHT, therefore, is now the only one permitted to be used throughout France.

The exact proportion of troy weight to the metrical standard has never been fully and exactly ascertained. In the table previously given, the kilogramme has been considered as equivalent to 15444 grains troy, which is the usual estimation of it. Francoeur, however, estimates the pound troy as equal to 392.9986 grammes; Matthieu, Legendre, and Duborg, as equal to 373.0956 grammes; Chelius and Houschild to 373.243 grammes. But the most correct estimation appears to be that of Lochman, who considers 9216 grains poid de marc as equivalent to 7555 grains troy; the pound troy as equal to 7026.320 grains marc; and the kilogramme as equal to 15434 grains troy. The subjoined tables are calculated on this estimation:—

## VALUE OF TROY WEIGHT IN METRICAL WEIGHT.

One pound	-	-	-	= 373.202	grammes
One ounce	-	-	-	= 31.10017	"
One drachm	-	-	-	= 3.887521	"
One scruple	-	-	-	= 1.295840	"
One grain	-	-	-	= 0.06479201	"

## VALUE OF TROY WEIGHT IN MARC WEIGHT.

One pound	= 12 ounces,	1 drachm,	42.33	grains
One ounce	= 1 ounce		9.53	"
One drachm	=	1 "	1.19	"
One scruple	=		24.40	"
One grain	=		1.2203	"

The other European States differ much in their medicinal weights from the troy standard; but it is superfluous to examine each in detail, the following synopsis and table being sufficient to show the variations.



1. The medicinal pound, except in Turkey, and by the old pouds de mare, is in all of them divided into 12 ounces.

2. In all of them, the ounce is divided into 8 drachms, except in Naples, where it contains 10.

3. In all, the ounce is divided into 3 scruples.

4. But the value of the scruple differs. It is composed of 24 grains at Bologna, Coni, Lucca, Modena, Parma, Rome, and in Spain, France, Portugal, Tuscany, and Sardinia.

Of 20 grains in Great Britain, the United States, Austria, Bavaria, Holland, Poland, Prussia, and Sweden; at Lubeck, Naples, Nuremberg, and Venice.

5. Consequently, the pound consists of

5760 grains in the United States, Great Britain, Austria, Bavaria, Holland, Poland, Prussia, and Sweden; at Lubeck, Nuremberg, Naples, and Venice.

6400 grains in Turkey.

6912 grains in Spain, Portugal, Tuscany, Sardinia, Bologna, Lucca, Modena, Parma, Rome, and Coni.

7200 grains at Naples.

9216 grains in France by the pouds de mare.

The subjoined table shows the differential values between the troy weight and the medicinal weights of the countries of Europe calculated in grains:—

COUNTRIES, &c.	Value of the pound, 5760 grains.	Value of the ounce, 480 grains.	Value of the drachm, 60 grains.	Value of the scruple, 20 grains.	Value of the grain.
Austria <sup>(1)</sup> . . .	5118.09	426.51	53.31	17.77	0.89
Bavaria <sup>(2)</sup> . . .	5971.23	497.60	62.20	20.73	1.04
Bologna . . .	7920.93	660.08	82.51	27.50	1.38
Coni . . . . .	8392.40	699.37	87.42	29.14	1.46
Corte . . . . .	7807.42	650.62	81.33	27.11	1.36
Holland <sup>(3)</sup> . . .	5732.38	477.70	59.71	19.90	1.00
Lubeck . . . . .	5823.60	485.30	60.66	20.22	1.01
Lucca . . . . .	7711.73	642.64	80.33	26.78	1.34
Modena . . . . .	7576.80	631.40	78.92	26.31	1.32
Monticelli <sup>(4)</sup> . .	8334.64	694.55	86.82	28.94	1.45
Naples <sup>(5)</sup> . . . .	8377.11	698.09	87.26	29.09	1.46
Nuremberg <sup>(6)</sup> . .	6007.22	500.60	62.58	20.86	1.04
Parma <sup>(7)</sup> . . . .	7864.55	655.38	81.92	27.31	1.37
Piacenza <sup>(8)</sup> . . .	8124.20	677.02	84.63	28.21	1.41
Poland . . . . .	5996.04	499.67	62.46	20.82	1.04
Portugal . . . . .	7494.62	624.55	78.07	26.02	1.30
Prussia <sup>(9)</sup> . . . .	6128.12	510.68	63.83	21.28	1.06
Rome . . . . .	7607.72	633.98	79.25	26.42	1.32
Spain . . . . .	7475.46	622.96	77.87	25.96	1.30
Sweden . . . . .	6034.48	502.87	62.86	20.95	1.05
Turin . . . . .	7770.71	647.56	80.94	26.98	1.35
Turkey . . . . .	7436.78	619.73	77.47	25.82	1.29
Tuscany . . . . .	7597.21	633.10	79.14	26.38	1.32
Venice . . . . .	7136.23	594.69	74.34	24.78	1.24

(1) This weight is used in Austria Proper, Bohemia, Moravia, Hungary, Transylvania, Tyrol, and the Lombardo-Venetian kingdom, with the exception of the city of Venice.

(2) This weight has been adopted in Greece.

(3) This weight is also used in Belgium.

(4) Monticelli, Castelvetro, and Polesine.

(5) Kingdom of the Two Sicilies.

(6) This weight is no longer used in Nuremberg; this city, being now incorporated in the kingdom of Bavaria, has adopted the Bavarian standard. But the Nuremberg weight is still extensively used in Germany and other countries. In Germany, we find it in the duchies of Saxe-



The value of these various pounds in grammes is thus calculated by Jourdain:—

The pound of 5760 grains.			
Venice	- - - = 301.230	grammes	Piacenza - - - = 317.577 grammes
Prussia	- - - = 350.761	"	Bologna - - - = 325.665 "
Sweden	- - - = 356.227	"	Soragna - - - = 325.800 "
Nuremberg	- - - = 357.843	"	Parma - - - = 328.000 "
Poland	- - - = 358.510	"	Corte - - - = 330.400 "
Bavaria	- - - = 360.000	"	Turin - - - = 331.961 "
Lubeck	- - - = 369.126	"	Lucca - - - = 334.500 "
United States and	} = 373.202	"	Rome - - - = 339.073 "
Great Britain			
Holland	- - - = 375.000	"	Tuscany - - - = 339.542 "
Austria	- - - = 420.009	"	Modena - - - = 340.457 "
The pound of 6400 grains.			Portugal - - - = 344.190 "
Turkey	- - - = 321.317	"	Spain - - - = 345.072 "
The pound of 6912 grains.			The pound of 7200 grains.
Coni	- - - = 307.370	"	Naples - - - = 320.230 "
Monticelli	- - - = 307.370	"	The pound of 9216 grains.
			France - - - = 489.503 "

#### LIQUID MEASURES OF THE UNITED STATES AND GREAT BRITAIN.

The liquid measures employed by the apothecaries in the United States and Ireland, are the wine gallon and its sub-divisions.

Altenburg, and of Anhalt-Bernburg; grand duchy of Baden; city of Bremen; duchies of Brunswick, Saxe-Meiningen, and Saxe-Coburg-Gotha; cities of Frankfurt-on-the-Maine and Hamburg; kingdom of Hanover; electorate of Hesse, grand duchy of Hesse, landgraviate of Hesse-Homburg; principality of Hohenzollern-Sigmaringen; duchy of Nassau; grand duchy of Oldenburg; principalities of Reuss-Plauen and Reuss-Schleitz, of Schwartzburg-Rudolstadt, and Schwartzburg-Sonderhausen; duchy of Sleswick-Holstein; principality of Waldeck; grand duchy of Saxe-Weimar; and kingdom of Wirtemberg.

The Nuremberg standard is also adopted in Denmark, Norway, Finland, Russia, and Switzerland.

(7) Parma, Borgo San-Donnino, Pellegrino, San-Secondo, Fontanellato, and Bussetto.

(8) Piacenza, Fiorenzola, and Carpaneta.

(9) This standard has been adopted in the kingdom of Saxony, the principalities of Lippe-Detmold and Lippe-Schaumburg, and the duchy of Anhalt-Dessau. It differs so little from those of Anhalt-Coethen and the grand duchies of Mecklenburg-Schwerin and Mecklenburg-Strelitz, that the standard of these three territories may be considered identical with that of Prussia.

This table needs hardly any explanation: it will enable the pharmacist—in all cases of foreign formulæ—to calculate the foreign weight in troy weight; thus, *e.g.*—one grain troy being equal to 0.89 grain of Austria—the value of one grain Austrian weight in troy weight may be readily ascertained by a simple rule of three example:—

$$\begin{array}{l} 0.89 : 1 :: 1 \\ 1 \times 1 = \times 1 = 0.100 \\ 0.89)0.100(1.1235 \end{array}$$

89

110

89

210

178

320

267

530

445

65



## WINE MEASURE.

1 minim					
60 =	1 fluidrachm	-	-	=	0.2254 cubic inches.
480 =	8 = 1 fluidounce	-	-	=	1.8047 "
7680 =	128 = 16 = 1 pint	-	-	=	28.875 "
61440 =	1024 = 128 = 8 = 1 gallon	-	-	=	231 "

For a long time, the Royal Colleges of Physicians in England, Scotland, and Ireland declined to recognize the use of measures in preparing and dispensing medicines, on the ground that the varying densities of different fluids rendered it difficult to use one common measure for all without risk of serious errors. But as druggists and medical practitioners constantly employed measures, in defiance of the prohibition, and as the practice, besides being attended with great and obvious convenience, was found to be less fraught with danger than had been conceived, the London College in the first instance, and subsequently the other colleges also, recognized a system of measures of their own, founded on the standard measures of the country. The Colleges of London and Edinburgh, adopting the imperial pint of 1826 as the basis, divide it into twenty parts, called fluidounces; each of which corresponds exactly with an avoirdupois ounce of distilled water at 62° F. and 30° B., and therefore contains 437.5 grains troy. The fluidounce is subdivided into eight parts of 54.6875 grains, termed fluidrachms; and each of these consists of sixty parts, called minims, which therefore amount each in weight to 0.91146 troy grain of distilled water.

The last edition of the Dublin Pharmacopœia having been published immediately before the passing of the Act of 1826, the imperial standard measures could not be adopted in that work as the basis of apothecaries' measure. The College, however, recognized a system adopted in a previous edition of the London Pharmacopœia, and founded on the wine measure, in general use in Britain prior to the legislative alterations of 1826. According to this system, the old wine gallon of 231 cubic inches, or 58317.8 troy grains of distilled water at 62° F. and 30° B., is divided into eight "pounds by measure" or old wine pints, each containing 7289.725 grains of water. The pint is subdivided into sixteen "ounces by measure," containing each 455.6075 grains; the ounce is divided into eight "drachms by measure" of 56.95 grains each; and the drachm is divided into three "scruples by measure," each of which contains nineteen grains troy, and consists of twenty "grains by measure." Care must be taken, therefore, not to confound the Dublin apothecaries' measure with that more recently adopted by the two British Colleges.

## IMPERIAL MEASURE.

1 minim					Grains troy.
60 =	1 fluidrachm			=	0.91
480 =	8 " 1 fluidounce			=	54.7
9600 =	160 " = 20 " = 1 pint			=	437.5 = 1 oz.
76800 =	1280 " = 160 " = 8 " = 1 gallon			=	8750 = 1.25 lb.
				=	70000 = 10 lbs.

## VALUE OF WINE OR APOTHECARIES' MEASURE IN IMPERIAL MEASURE.

Wine Measure.		Apothecaries' Measure.			
		Pints.	Fluidounces.	Fluidrachms.	Minims.
1 gallon	=	6	13	2	23
1 pint	=		16	5	18
1 fluidounce	=		1	0	20
1 fluidrachm	=			1	2½

## VALUE OF IMPERIAL MEASURE IN WINE OR APOTHECARIES' MEASURE.

	Gallon.	Pints.	Fluidounces.	Fluidrachms.	Minims.
1 gallon	1	1	9	5	8
1 pint		1	3	1	58
1 fluidounce				7	41
1 fluidrachm					58



Besides these regular and authorized measures, there are others constantly used in the preparation and administration of medicines, which require notice. These, which have been adopted for convenience, are far from being uniform, but may be used without danger for ordinary purposes. Drs. Wood and Bache estimate the

Teacup	as containing about	four fluidounces.
Wineglass	“ “	two fluidounces.
Tablespoon	“ “	a half a fluidounce.
Teaspoon	“ “	a fluidrachm.

Dr. Christison gives a somewhat different calculation; he considers

A Tumbler	to contain	eight fluidounces.
Breakfast cup	“	do do
Teacup	“	five fluidounces.
Wineglass	“	two fluidounces.
Tablespoon	“	half a fluidounce.
Dessertspoon	“	two fluidrachms.
Teaspoon	“	one fluidrachm.

In both these estimates, the teaspoon is rated too low; they apply very well to the teaspoonful formerly used, but not to the much larger kind now in general use, which approaches the dessertspoon in capacity. Quart and pint bottles are also sometimes employed as measures of capacity for medical purposes; but they are very fallacious guides of quantity, as the so called quart bottle never holds a quart, nor the misnamed pint bottle a pint. The first seldom holds more than twenty-six to twenty-seven fluidounces, and the latter from thirteen to fourteen.

It is still more common to estimate small quantities or doses of medicine by drops, as representing, and equivalent to, minims. This is so convenient, that it is not likely to be abandoned, though nothing can be more erroneous, as the size of the drop of the same fluid varies much with the form of the mouth of the bottle, its size, the mode in which the operation is performed, and still more from the nature of the fluid thus attempted to be measured. Numerous experiments have been made on the subject, but can only be considered as approximate estimates, though sufficiently accurate for practical purposes.

Mr. Alsop, of London, some years since, published the result of some experiments made by him, which show the great variation to which this mode of measuring is liable. The following proves that the number of drops required to measure a fluidrachm are widely different when dropped from a large or a small bottle.

One fluidrachm.		Large bottle.	Small bottle.
Diluted sulphuric acid	- - - -	24 drops	84 drops
Scheele's hydrocyanic acid	- - - -	35 “	60 “
Distilled water	- - - -	31 “	54 “
Solution of ammonia	- - - -	40 “	48 “
Tincture of opium	- - - -	84 “	135 “
Rectified spirit	- - - -	100 “	130 “
Tincture of muriate of iron	- - - -	100 “	150 “

Mr. Durand, of Philadelphia, in an able paper on the subject (*Journ. Phila. Col. of Pharm.*, vol. i. p. 165), says, “The bulk of drops depends not only on the density of the liquids which furnish them, and the cohesion of the constituent particles of that liquid, but also on the shape of the mouth of the vessel from which they are poured. An open vessel with a beak, such as the common graduated measure, affords a larger drop than a bottle with the stopper half drawn out; a mode commonly practiced. That furnished by the dropping tube is still smaller, and is ever liable to vary with the greater or lesser diameter of its extremity. Besides, in every instance, the first drops poured from any vessel are always smaller than those subsequently obtained.”

He goes on to say that the following may be considered as established:—

1. That liquids which contain a small proportion of water afford a small drop; while, on the contrary, liquids containing a large quantity of water furnish a large



drop. For instance, concentrated acids, ethers, rectified alcohol, fixed and essential oils, &c., which contain a very small proportion of water, yield a smaller drop than diluted acids, weak alcohol, wine, &c.

2. That, among liquids containing a large proportion of water, those which are not charged with remedial substances give a larger and heavier drop than those same liquids containing extraneous bodies in solution. Thus, weak alcohol, wine, vinegar, and water furnish a larger and heavier drop than the tinctures prepared from them.

The following table illustrates his remarks:—

TABLE

*Showing the Differences between Minims, Drops, and Grains of various Medicinal Liquid Preparations of the Pharmacopœia of the United States, &c.*

PREPARATIONS.	No. of drops in 20 minims.	No. of min. in 20 drops.	No. of drops in 20 grains.	No. of grains in 20 drops.
Sulphuric acid - - - - -	30	13.3	25	16
Sulphuric ether - - - - -	50	8	60	6
Rectified alcohol - - - - -	46	8.6	57	7.1
Nitric acid - - - - -	28	14.2	22.2	18
Acetic acid (crystallizable) - - - - -	40	10	40	10
Muriatic acid - - - - -	18	22.2	18.1	22
Oil of wormseed (chenopod, anthelminticum) - - - - -	40	10	50	8
of peppermint, aniseed, sweet almond, olive, palma christi - - - - -	40	10	43.5	9
of cloves - - - - -	40	10	36	11
of cinnamon - - - - -	40	10	32	12.5
Copaiba - - - - -	40	10	40	10
Diluted alcohol - - - - -	40	10	42	9.5
Tincture of hydriodate of potassa, cantharides, kino, digitalis, assafetida, sulphuric acid, colchicum, opium, valerian, guaiacum - - - - -	40	10	43	9.3
of valerian, guaiacum (volatile) - - - - -	40	10	50	8
of muriate of iron - - - - -	44	9.1	50	8
Wine, Teneriffe - - - - -	26	15.3	25	16
antimonial - - - - -	24	16.6	26	15.3
of opium (Sydenham's laudanum) - - - - -	26	15.3	29	13.7
of colchicum root and seeds - - - - -	25	16	29	13.7
Vinegar, distilled - - - - -	19	21	20	20
of opium (black drop) } of colchicum } of squill - - - - -	26	15.3	25	16
Water, distilled - - - - -	15	26.6	17.5	24.5
solution of hydrocyanic acid - - - - -	15	26.6	17.5	24.5
sulphuric acid (1 to 7) - - - - -	17	23.5	17	23.5
nitric " " - - - - -	17	23.5	17	23.5
ammonia (strong) - - - - -	18	22.2	18.5	22
" (weak) - - - - -	15	26.6	20	20
hydriodate of potassa - - - - -	18	22.2	20	20
arsenite of potassa - - - - -	19	21	20	20

The only mode to obviate these discrepancies is for the physician to order the administration of small quantities of fluid medicines in minims, and not in drops, which would induce the use of a minim measure in every sick room.



## FRENCH MEASURES OF CAPACITY.

1 millilitre	-	-	-	-	=	16.2318 minims.
10 =	1 centilitre	-	-	-	=	2.7053 fl. dr.
100 =	10 =	1 decilitre	-	-	=	3.3816 fl. ounces
1000 =	100 =	10 =	1 litre	-	=	2.1135 pints.
10000 =	1000 =	100 =	10 =	1 decalitre	-	2.6419 gallons.
100000 =	10000 =	1000 =	100 =	10 =	1 hectolitre	= 26.4190 "
1000000 =	100000 =	10000 =	1000 =	100 =	10 =	1 kilolitre = 264.1900 "

Litres.	Eng. cubic inches.	Imperial pints.	Wine pints.	Troy ounces of water.
1 =	61.028 =	1.7608 =	2.1135 =	31.104
2 =	122.056 =	3.5216 =	4.2270 =	62.208
3 =	183.084 =	5.2822 =	6.3405 =	96.312
4 =	244.112 =	7.0430 =	8.4541 =	128.416
5 =	305.140 =	8.8038 =	10.5476 =	160.520
6 =	366.168 =	10.5646 =	12.6811 =	192.624
7 =	427.196 =	12.3253 =	14.7947 =	224.728
8 =	488.224 =	14.0861 =	16.9082 =	256.832
9 =	549.252 =	15.8469 =	19.0217 =	288.936

The former measure in use before the introduction of the metrical system was the *pint*, and its subdivisions; and it still continues to be employed in some parts. It agrees with the metrical measure as follows:—

1 demi-poisson	=	62.50 grammes	=	0 litre	0 decilitre	1 centilitre.
1 poisson	=	125	=	0 "	1 "	2 "
1 demi-setier	=	250	=	0 "	2 "	5 "
1 chopine	=	500	=	0 "	5 "	0 "
1 pinte	=	1000	=	1 "	0 "	0 "

The subjoined is a list of the principal liquid measures of Europe, with their corresponding value in the metrical measure of France:—

AUSTRIA.—The fundamental unity is the *mass* or *kanne* = 71.3343 Par. cubic inches, or 1.415015 litre. The mass is divided into 3 *seidel*, and 40 mass make one *eimer*.

BADEN.—The fundamental unity is the *mass* = 1.5 decimetre cubic, and accordingly equivalent to 1.5 litre.

BAVARIA.—The fundamental unity is the *mass* = 43 decimal cubic inches of Bavaria, or 1.06921728 litre. The mass is divided into 4 *schoppen*.

DENMARK.—The fundamental unity is the *pott* ( $\frac{1}{32}$ d part of a cubic foot Danish), equivalent to 0.96529 litre. The pott is divided into four *pegel*, and 2 potts make a *kanne*.

GRAND DUCHY OF HESSE.—The fundamental unity is the *mass*, which corresponds exactly to 2 litres. The mass is divided into 4 *schoppen*.

NAPLES.—The unity is the *barile*, equivalent to 43.6216 litres.

PORTUGAL.—The unity is the *almude*, equivalent to 16.451 litres.

PRUSSIA.—The unity is the *quart*. This measure has a capacity of 64 cubic inches of distilled water, at 61.25 Fahr. and 27° 10' barom., = 57.724 cubic inches of Paris, or 1.145 litre.

ROME.—The unity is the *barile*, equivalent to 58.3416 litres.

RUSSIA.—The unity is the *wedro*, which contains 750 cubic inches of distilled water at 50° Fahr. and 30° barom., = 30 pounds Russian. The *wedro* is equivalent to 12.28963047 litres, and one litre is accordingly equivalent to 0.081369410776019 *wedro*. The *wedro* is divided into 10 *stof*, and the *stof* into 10 *tschark*.

SPAIN.—The unity is the *arroba*, equivalent to 16.073 litres.

SWEDEN.—The fundamental unity is the *kanna*, which contains 100 cubic decimal inches of pure water at 62° Fahr., *in vacuo*, equivalent to 6.151951 pounds Swedish. One pound Swedish of pure water at 62° Fahr. *in vacuo*, is equal to 0.4250104 litre;



the *kanna* is consequently equivalent to 2.6146431552904 litres, or, calculated upon the maximum density of water, 2.617341607126 litres, or, in shorter terms, 2.62 litres. The *kanna* is divided into 2 *stop*, the *stop* into 4 *quarter*, and the *quarter* into 4 *jungfrus*.

TURIN.—The unity is the *brenta*, divided into 36 pints, and equivalent to 49.28468 litres.

TUSCANY.—The unity is the *barile*, equivalent to 45.584 litres.

WIRTEMBERG.—The unity is the *mass*, which contains 78.125 cubic inches (Wirtemberg measure), and is equivalent to 1.83705 litre. The *mass* is divided into four *schoppen*.

### SPECIFIC GRAVITY.

The following remarks on Specific Gravity have been taken from Redwood's edition to Gray's Supplement, and comprise in a condensed form much useful information.

The determination of the specific gravity of a body consists in estimating the weight of a given volume of it, as compared with an equal volume of some other body. The bodies usually taken as the standard of comparison are *pure water* for solids and liquids, and *atmospheric air* for gases.

The *specific gravity of a solid* is determined by first weighing it in the ordinary manner with an accurate balance suspended in the air; then attaching a horse-hair, or fine silken thread to the solid body, immersing it in pure distilled water, and weighing it while thus immersed. The weight of the body in air, divided by the difference between its weight in air and its weight in water, will be its specific gravity. Thus a piece of lead is found to weigh 398 grains in air. When immersed in water, its weight is 362.4 grains; and the difference between these two weights, namely, 35.6, is the weight of the volume of water displaced by the lead, or of a volume of water equal to that of the lead. The volume of water being taken as unity, the specific gravity of the lead is found by the following rule of three sum:—

35.6 : 1 : : 398 : 11.176, the specific gravity of the lead.

In taking the specific gravity of a *solid substance lighter than water*, some modification of the process is required: but we have, nevertheless, the same preliminary points to determine; first, the weight of the substance in air; and secondly, the weight of an equal volume of water. This may be illustrated by taking the specific gravity of a piece of wax. The weight of the wax in air is 105.4 grains. On immersing the wax in water, two pressures are exerted—a pressure downwards, equal to the gravity or weight of the wax, and a pressure upwards, equal to the weight of the volume of water displaced by the wax; but the specific gravity of water being greater than that of wax, the upward pressure preponderates, and the wax rises to the surface. Thus, we find that a volume of water equal to that of the wax weighs as much as the wax, and something more. We must ascertain how much more; and this is done in the following manner: Some body heavier than water, and the weight of which in water is known, is attached to the wax, and the two bodies are weighed in water together. A piece of lead may be used for this purpose. The lead, alone, weighs 378 grains in water; with the wax attached to it, the weight in water is 372.4 grains, making a difference of 5.6 grains; and this 5.6 grains is equal to the excess of the upward over the downward pressure on the wax, when immersed in water. Thus, a volume of water equal to that of the wax weighs 5.6 grains more than the wax, or  $105.4 + 5.6 = 111$  grains.

Then,  $111 : 1 : : 105.4 : 0.949$ , the specific gravity of the wax.

It sometimes happens that the solid substance, the specific gravity of which is to be determined, is in *powder*, or in *several small particles*. In such cases, it is found convenient to proceed as in the following method of taking the specific gravity of calomel:—

100 grains of calomel are introduced into a specific gravity bottle, which holds 1000 grains of distilled water; the bottle is filled up with water, and the weight of the contents is found to be 1083.7 grains; deducting the weight of the calomel (100 grains) from this, the remainder (983.7 grains) will be the weight of the water in the



bottle, and the difference (16.3 grains) between this and 1000 grains, the weight of the whole contents of the bottle when filled with distilled water, is the weight of a volume of water equal to the volume of the calomel.

Then,  $16.3 : 1 :: 100 : 6.03$ , the specific gravity of the calomel.

In taking the *specific gravity of substances soluble in water*, other modifications of the process are required. Sometimes the substance may be covered with a thin coating of varnish, so as to protect it from the action of the water. This method answers very well for blue pill, which may be brushed over with a strong tincture of mastic, and then proceeded with as in the case of the lead. In other instances, however, it is necessary to pursue a different course. Thus, any powder that is soluble in water must have its specific gravity taken, in the first instance, with reference to some liquid in which it is not soluble. Spirit of wine, oil of turpentine, or olive oil, may be used in such cases. The process may be illustrated by describing the method of taking the specific gravity of guano in oil of turpentine.

In the first place, the specific gravity of the oil of turpentine is ascertained to be 0.874. Then 100 grains of guano are introduced into a specific gravity bottle, as in the case of the calomel; and the bottle being filled up with oil of turpentine, the weight of the contents is found to be 922.7 grains, from which, deducting 100 grains, the remainder (842.7 grains) will represent the oil not displaced by the guano; and this, deducted from 874 grains, the quantity of oil the bottle is capable of holding, leaves 51.3 grains as the weight of a volume of oil of turpentine equal to that of the guano. Now,  $874 : 51.3 :: 1000 : 58.7$ , the weight of a volume of water equal to that of the guano.

Then,  $58.7 : 1 :: 100 : 1.7$ , the specific gravity of the guano.

The methods by which the *specific gravities of liquids* are usually determined may be divided into two classes:—

1st. Those which consist in filling any suitable vessel with the liquid to be estimated, ascertaining the weight of the contents, and dividing this by the weight of the same volume of water.

2d. Those which consist in displacing a portion of the liquid by some solid body floating in it, and estimating the specific gravity according to the weight and volume of the substance immersed, as compared with its immersion in water.

In the first case, the instruments employed are a specific gravity bottle and an ordinary balance.

In the second case, the instruments used may be comprehended under the general terms of *hydrometers* or *aréometers*. These, however, are distinguished from each other, for there are many varieties of them, by different names, according to the particular purpose for which they are respectively intended, or from some peculiarity in their construction.

The *specific gravity bottle* affords the most accurate means of determining the comparative densities of liquids. It consists, usually, of a globular bottle with a flat bottom and a slender neck, which holds exactly 1000 grains of distilled water at a certain fixed temperature. It is very easy at any time to test the accuracy of one of these bottles by a single experiment; and, having ascertained that the bottle is correctly adjusted with regard to distilled water, the indications afforded with any other liquid will be equally trustworthy. The weight in grains of the quantity of any liquid filling such bottle will indicate its specific gravity.

*Hydrometers*, or *aréometers*, are floating instruments, and their application for the purpose of determining the specific gravities of liquids depends upon the fact that a body immersed in any liquid sustains a pressure from below upwards equal to the weight of the volume of the liquid displaced by such body.

The use of hydrometers for determining the specific gravities of liquids has been traced back to a period about 300 years before Christ; an instrument of this kind being described as the invention of Archimedes, the Sicilian mathematician. It subsequently fell into disuse, but was again brought into notice by Basil Valentine.

There are two kinds of hydrometers, which may be taken as the types of all the different varieties in regard to construction:—

1st. Those which are always immersed in the liquids to be tried, to the same



depth, and to which weights are added to adjust the instrument to the density of any particular liquid. Of this description are Fahrenheit's, Nicholson's, and Guyton de Morveau's hydrometers.

2d. Those which are always used with the same weight, but which sink into the liquids to be tried to different depths, according to the densities of the liquids. These usually have graduated scales attached to their stems. Of this description are the common glass hydrometers, generally including those of Baumé, Cartier, Gay Lussac, Twaddle, Zanetti, &c., and the specific gravity beads.

Sikes's and Dicas's hydrometers combine the principles of both types, having movable weights and graduated scales.

Hydrometers may also be divided into two classes, as follows:—

*First.* Those having a general application, for determining the comparative densities of any liquids.

*Second.* Those intended for special application; as for estimating the comparative strengths of spirits, or the comparative densities of syrups, oils, &c.

Fahrenheit's, Nicholson's, Guyton de Morveau's, and the common glass hydrometers, including Baumé's, Cartier's, Zanetti's, and the specific gravity beads, belong to the first class.

Gay Lussac's, Sikes's, and Dicas's hydrometers, the saccharometer, urinometer, and elaëometer, belong to the second class.

*Fahrenheit's hydrometer* consists of two glass bulbs blown in a glass tube, like a common hydrometer, excepting that the upper bulb is larger, and the stem, which is small, is terminated at the top in a cup or funnel. It has a mark on the middle of the stem, indicating the point at which the instrument is to be made to float, by means of weights put into the cup.

*Nicholson's hydrometer* is a modification of Fahrenheit's. It is made of brass, and consists of a hollow globe, to which is fixed a slender stem surmounted by a cup; on the opposite side of the globe is another cup fixed in a kind of stirrup, and loaded, so that this may always form the lowest point of the instrument when immersed in any liquid. There is a mark on the middle of the upper stem, indicating the point at which the instrument is to be made to float. A certain weight is introduced into the cup, to cause the instrument to sink to the proper mark in distilled water. On immersing the hydrometer into any other liquid, more or less weight will have to be put into the cup, according as such liquid is more or less dense than water. Thus the relative densities of liquids are determined.

This instrument is also applicable for taking the specific gravities of solids. If the solid substance be put into the cup as part of the weight required to sink the hydrometer in distilled water, the weight of the substance in air is ascertained; and if it be then put into the lower cup, immersed in the water, and the instrument again adjusted, its weight in water is ascertained; and from these its specific gravity is calculated.

*Guyton de Morveau's hydrometer* is similar to Fahrenheit's.

*Baumé's hydrometers* are used extensively in this country, as well as in France, and are applicable for all kinds of liquids. There are two distinct instruments: one for liquids lighter than water, and the other for liquids heavier than water. The latter is, for distinction, called the *acidometer* or *saccharometer* (*pèse-acide* or *pèse-sirop*); the former, the *spirit hydrometer* (*pèse-esprit*).

*Baumé's acidometer* is made in the form of the common hydrometers. It consists of a glass tube terminated at the lower end by two bulbs, the lowest bulb being much smaller than the other, and intended to contain the ballast with which the instrument is loaded. The scale is marked on a slip of paper, or of ivory, fixed in the tube, and is adjusted in the following manner: The top of the tube being open, the slip of paper on which the scale is to be marked is put into the stem, and the instrument is then immersed in pure distilled water; quicksilver is now dropped into the lower ball until the instrument sinks so low in the water that only the top of the stem remains above the surface, and a mark is made on the glass denoting exactly the point to which it sinks. The instrument is now taken out of the pure water and put into a solution of fifteen parts of common salt in eighty-five parts of distilled



water, this solution being at the same temperature as the water in which the instrument was previously immersed; the point to which it sinks in this solution is to be marked on the stem as before, and the distance between the two marks being taken with a pair of compasses, and transferred to the slip of paper, the first is made the zero or 0, and the other the 15th degree of the scale. This distance being divided into fifteen equal parts or divisions, each division is called a degree, and the scale is completed by adding as many more degrees as the length of the stem will admit of. This being done, the slip of paper is again introduced into its place, and so fixed that the zero (0) of the scale shall be exactly opposite the first mark made on the glass. The end of the stem is now sealed with the flame of a blowpipe.

*Baumé's spirit hydrometer* is similar in form to the acidometer; but the weight of the instrument, and the scale, are different. In this case, the hydrometer is first immersed, as before, in pure distilled water; but it is made to float, so that the greater part of the stem shall be above the surface of the water. This point is marked, and the instrument is then transferred to a solution of ten parts of common salt in ninety parts of water, when another mark is made. The distance between these marks is made ten degrees of the scale, which are divided with the compasses, and marked on the slip of paper, as in the other case, the floating point in the solution of salt being made the zero, and the degrees carried upwards from this point.

The temperature at which these instruments were originally adjusted by Baumé, was 10° Reaumur, or 12.5 Centigrade; but those made in England are usually adjusted at 60° Fahrenheit. It is sometimes important to be aware of this difference.

*Cartier's hydrometer* is much used in France. It is only applicable for liquids lighter than water. This instrument is a modification of Baumé's spirit hydrometer, the form of the instrument being the same, and the same point being taken as the zero of the scale; but the space which, in Baumé's scale, is divided into 32°, is in Cartier's divided into 30°.

It is becoming the common practice in this country to have the scales of hydrometers marked with the specific gravities intended to be indicated, and this is by far the most convenient kind of hydrometer for general use.

*Twaddle's hydrometers* are much used in Scotland, and occasionally in England. They are made of glass like the common hydrometers, and are sold in sets of six. Each degree on the scale is equal to 0.005 of specific gravity, so that the specific gravity of a liquid is found, with these hydrometers, by multiplying the number of degrees indicated by 5, and adding 1000. Thus, 10° by Twaddle's hydrometer,  $\times 5 + 1000 = 1.050$  specific gravity.

*Zanetti's hydrometers*, which are made at Manchester, are also sold in sets of six. With these the specific gravity is got by adding a cipher to the number of degrees indicated.

*Specific gravity beads*, sometimes called *Lovi's beads*, are hollow sealed globes of glass, about the size of small pistol-bullets. Each bead is a small hydrometer, intended to indicate one fixed density, by its remaining half way between the top and bottom of the liquid into which it is introduced. These beads are sold in sets, each one being marked with the specific gravity it is to indicate at a certain fixed temperature. They are very useful in making mixtures of any required densities, as, for instance, in making test acids.

*Gay Lussac's alcolometre* is frequently employed in France; it is adapted only for estimating the strength of spirits. The instrument is made like a common glass hydrometer, the scale of which is divided into 100 parts or degrees. The lowest division marked 0, at the bottom of the scale, denotes the specific gravity of pure water at a temperature of 15° cent., and the highest division, at the top of the scale, the specific gravity of absolute alcohol at the same temperature. The intermediate degrees indicate the number of volumes of absolute alcohol in 100 volumes of the spirit tried. The instrument is accompanied by a table for correcting the numbers marked on the scale, when it is used at any other temperature than that of 15° cent.

*Sikes's hydrometer* is used exclusively in the collection of the spirit revenue. It consists of a spherical ball or float, and an upper and a lower stem made of brass; the upper stem has ten principal divisions, numbered 1, 2, 3, &c., which are each subdivided into ten parts.



vided into five parts; the lower stem is made conical, and has a pear-shaped loaded bulb at its lower extremity. There are nine movable weights, having the form of circular discs, and numbered 10, 20, 30, and so on to 90. Each of the circular weights is cut into its centre, so that it can be placed on the inferior conical stem, and slid down to the bulb; but, in consequence of the enlargement of the cone, they cannot slip off at the bottom, but must be drawn up to the thin part for this purpose. The instrument is adjusted to strong spirit, specific gravity .825, at 60° Fahr., this being reckoned as standard alcohol. In this spirit, the instrument floats at the first division, 0, or zero, without a weight. In weaker spirit, having a greater density, the hydrometer will not sink so low; and, if the density be much greater, it will be necessary to add one of the weights to cause the entire immersion of the bulb of the instrument. Each weight represents as many principal divisions of the stem as its number indicates: thus, the heaviest weight, marked 90, is equivalent to 90 divisions of the stem, and the instrument with this weight added floats at 0 in distilled water. As each principal division on the stem is divided into five, the instrument has a range of 500 degrees between *standard alcohol*, specific gravity .825, and water. In using this instrument, it is immersed in the spirit, and pressed down by the hand to 0, till the whole divided part of the stem be wet. The force of the hand required to sink it will be a guide in selecting the proper weight. Having taken one of the circular weights, which is necessary for this purpose, it is slipped on the conical stem. The instrument is again immersed, and pressed down as before to 0, and is then allowed to rise and settle at any point of the scale. The eye is then brought to the level of the surface of the spirit, and the part of the stem cut by the surface, *as seen from below*, is marked. The number thus indicated by the stem is added to the number of the weight employed, and with this sum at the side, and the temperature of the spirits at the top, the strength *per cent.* is found in a table which accompanies the hydrometer. The strength is expressed in numbers denoting the excess or deficiency *per cent.* of proof spirit in any sample; and the number itself, having its decimal point removed two places to the left, becomes a factor, whereby the gauged contents of a cask or vessel of such spirit being multiplied, and the product being added to the gauged contents if over proof, or deducted from it if under proof, the result will be the actual quantity of proof spirit contained in such cask or vessel.

*Dicas's hydrometer* is similar in construction to Sikes's; and it is used in a similar manner, with the same result, indicating the relation of the spirit tried to standard proof spirit.

It is the practice in commerce to designate the strength of spirit as so many degrees above or below proof, the government having fixed upon what is called *proof spirit* as the standard in comparison with which the strength of all spirit shall be estimated. The term *proof* is said to have been derived from the ancient practice of trying the strength of spirit by pouring it over gunpowder in a cup, and then setting fire to the spirit; if, when the spirit had burned away, the gunpowder exploded, the spirit was said to be *over proof*: if, on the other hand, the gunpowder failed to ignite, in consequence of the water left from the spirit, it was said to be *under proof*. The weakest spirit capable of firing gunpowder in this way was called *proof spirit*: but it requires a spirit nearly of the strength of what is now called rectified spirit to stand this test. The *standard proof spirit* of the excise is defined by law (56 Geo. III. cap. 140) to be "*that which, at a temperature of 51° by Fahrenheit's thermometer, weighs exactly twelve-thirteenth parts of an equal measure of distilled water.*" This will have a specific gravity of .923 at 51° Fahr., or about .920 at 60° Fahr. The *standard alcohol* of the excise is spirit, the specific gravity of which is .825 at 60° Fahr. By "*spirit 60 degrees over proof*," is understood a spirit, 100 measures of which, added to 60 measures of water, will form *standard proof spirit*, sp. gr. 920. By "*spirit 10 degrees under proof*," is understood a spirit, 100 measures of which, mixed with 10 measures of standard alcohol, sp. gr. .825, will form *standard proof spirit*.

*Saccharometers*, which are hydrometers intended for determining the density of syrups, are usually made and graduated in the same manner as Baumé's acidometers, and differ only from these in being made smaller; but the scale is sometimes graduated to indicate the proportion of sugar in the solution.



The *urinometer* is a small hydrometer, originally suggested by Dr. Prout, for estimating the density of urine. The scale is divided into 60 degrees, the zero being the point at which it floats in distilled water. The numbers on the scale, added to 1000, the assumed sp. gr. of water, give the specific gravities at the respective points; thus, supposing the number cut by the surface of the liquid to be 30, this indicates a sp. gr. 1030. The letters H. S., on the back of the scale, signify *healthy standard*, which ranges from 10° to 20° of the scale.

The *elaiometer* is a very delicate glass hydrometer, intended for testing the purity of olive oil or oil of almonds, by determining their densities. The 0 or zero of the scale is the point at which the instrument floats in *oil of poppy seeds*. The point at which it floats in *pure olive oil* is made the 50th degree, and the space between these two points is divided into 50 equal parts, and numbered accordingly. It floats at 38 or 38½° in *pure oil of almonds*.

The following tables have been drawn up for the purpose of showing the relations between the indications afforded by some of the foregoing instruments.

*Relation between Specific Gravities, and Degrees of Baumé's Hydrometer for Liquids heavier than Water.*

Sp. gr. Baumé.	Sp. gr. Baumé.	Sp. gr. Baumé.	Sp. gr. Baumé.	Sp. gr. Baumé.
1.000 = 0	1.116 = 15	1.264 = 30	1.454 = 45	1.714 = 60
1.007 1	1.125 16	1.275 31	1.470 46	1.736 61
1.014 2	1.134 17	1.286 32	1.485 47	1.758 62
1.022 3	1.143 18	1.298 33	1.501 48	1.779 63
1.029 4	1.152 19	1.309 34	1.526 49	1.801 64
1.036 5	1.161 20	1.321 35	1.532 50	1.823 65
1.044 6	1.171 21	1.334 36	1.549 51	1.847 66
1.052 7	1.180 22	1.346 37	1.566 52	1.872 67
1.060 8	1.190 23	1.359 38	1.583 53	1.897 68
1.067 9	1.199 24	1.372 39	1.601 54	1.921 69
1.075 10	1.210 25	1.384 40	1.618 55	1.946 70
1.083 11	1.221 26	1.398 41	1.637 56	1.974 71
1.091 12	1.231 27	1.412 42	1.656 57	2.002 72
1.100 13	1.242 28	1.426 43	1.676 58	2.031 73
1.108 14	1.252 29	1.440 44	1.695 59	2.059 74
				2.087 75

*Relation between Specific Gravities, and Degrees of Baumé's Hydrometer for Liquids lighter than Water.*

Sp. gr. Baumé.	Sp. gr. Baumé.	Sp. gr. Baumé.	Sp. gr. Baumé.	Sp. gr. Baumé.
1000 = 10	0.936 = 20	0.880 = 30	0.830 = 40	0.785 = 50
0.993 11	0.930 21	0.874 31	0.825 41	0.781 51
0.986 12	0.924 22	0.869 32	0.820 42	0.777 52
0.980 13	0.918 23	0.864 33	0.816 43	0.773 53
0.973 14	0.913 24	0.859 34	0.811 44	0.768 54
0.967 15	0.907 25	0.854 35	0.807 45	0.764 55
0.960 16	0.901 26	0.849 36	0.802 46	0.760 56
0.954 17	0.896 27	0.844 37	0.798 47	0.757 57
0.948 18	0.890 28	0.839 38	0.794 48	0.753 58
0.942 19	0.885 29	0.834 39	0.789 49	0.749 59
				0.745 60



## TEMPERATURES TO BE OBSERVED IN CERTAIN PHARMACEUTICAL OPERATIONS.

In the fermentation of saccharine solutions, the highest temperature should not exceed 86° (Thomson).

The lowest temperature at which they will ferment is 38° (Thomson).

The process of acetous fermentation is best conducted at a temperature of about 86°.

The temperature requisite to coagulate albumen varies with the state of dilution. If the quantity of albumen be so great that the liquid has a slimy aspect, a heat of 145° or 150° suffices, but in a very dilute condition boiling is required (Fownes).

*In the London and U. S. Pharmacopœias:*

When a boiling heat is directed, a temperature is meant of 212° Fahr.

When a gentle heat is directed, a temperature is meant of from 90° to 100°.

The specific gravities of substances ordered in the London Pharmacopœia are to be taken at a temperature of 62°.

A water bath is that by which any substance contained in a proper vessel is exposed either to hot water, or the vapor of boiling water. A sand bath is made of sand, to be gradually heated, in which anything is placed contained in a proper vessel.

Syrups are to be kept in a place where the temperature never exceeds 55°.

Vegetables, shortly after they have been gathered, those excepted which ought to be fresh, are to be lightly strewed, and dried as quickly as possible with a gentle heat (90° to 100°). They are to be kept afterwards in proper vessels, excluded from the access of light and moisture.

*In the Dublin Pharmacopœia:*

By the term superior heat is meant some degree between 200° and 212°.

When a medium heat is directed, a temperature is meant between 100° and 200°.

When an inferior heat is directed, a temperature is meant between 90° and 100°.

In the process of digestion, an inferior heat is to be applied, unless it should be otherwise directed. In the process of maceration, a heat should be applied between 60° and 90°.

*In the Dublin, Edinburgh, and U. S. Pharmacopœias:*

Whenever mention occurs of the specific gravity of any body, its temperature is supposed to be at 60°.

## HYDROMETRICAL EQUIVALENTS.

Sp. gr. at 60° Fahr.	100 parts contain of		1000 parts contain of standard alc. sp. gr. 825	Sikes.	Baumé.	Cartier.	Per cent. of alcohol, sp. gr. 796 by volume. Gay Lussac.
	Alcohol sp. gr. 796.	Water. By weight.					
796	100	0			46.5	43.48	100
797	99.5	.5					99.75
798	99	1			46	43.06	99.50
799	98.67	1.33					99.25
800	98.33	1.67					99
801	98	2					98.75
802	97.67	2.33			45	42.14	98.50
803	97.33	2.67				42	98.28
804	97	3					98.15
805	96.67	3.33					98
806	96.33	3.67					97.80
806.5	96.17	3.83			44	41.22	97.70



Sp. gr. at 60° Fah.	100 parts contain of		1000 parts contain of standard alc. sp. gr. 825.	Sikes.	Baumé.	Cartier.	Per cent. of alcohol, sp. gr. 796 by volume. Gay Lussac.
	Alcohol sp. gr. 796.	Water. By weight.					
807	96	4					97.60
808	95.5	4.5				41	97.40
809	95	5					97.29
809.5	94.89	5.10					97.10
810	94.67	5.33					97
811	94.33	5.67			43	40.34	96.75
812	94	6					96.50
813	93.67	6.33				40	96.25
814	93.33	6.67					96
815	93	7					95.75
816	92.5	7.5			42	39.40	95.50
817	92	8					95.25
818	91.67	8.33					95
818.6	91.5	8.5				39	94.90
819	91.33	8.67					94.75
820	91	9					94.50
821	91.5	9.5			41	38.46	94.25
822	90	10					94
823	89.67	10.33				38	93.75
824	89.33	10.67					93.50
825	89	11	1000	63	O. P. 40	37.55	93.25
826	88.5	11.5	993	62			93
827	88	12	988.5	61.5		37	92.6
828	87.67	12.33	984	61	39.5		92.3
829	87.33	12.67	979.5	60.5	39	36.63	91
830	87	13	975	60			91.7
831	86.5	13.5	970.5	59.5	38.5	36.17	91.35
832	86	14	966	59		36	91
833	85.67	14.33	961.5	58.3			90.65
834	85.33	14.67	957	58	38	35.72	90.3
835	85	15	953	57.5			90
836	84.67	15.33	949	57	37.5	35.26	89.7
837	84.33	15.67	944.5	56.5			89.35
837.6	84.25	15.75	942.5	56.3		35	89.20
838	84	16	940	56			89
839	83.5	16.5	936	55.5	37	34.80	88.75
840	83	17	932	55			88.5
841	82.67	17.33	928	54.5	36.5		88.25
842	82.33	17.67	924	54		34.94	88
843	82	18	920	53.5		34	87.65
844	81.67	18.33	916	53	36	33.88	87.3
845	81.33	18.67	912	52.5			87
846	81	19	908	52			86.7
847	80.5	19.5	903	51	36.5	33.42	86.35
848	80	20	898	50			86
849	79.67	20.33	893	49.5			85.65
850	79.33	20.67	888	49	35	33	85.3
851	79	21	883	48.5			85
852	78.5	21.5	878	48	34.5	32.43	84.7
853	78	22	873	47.5			84.35
854	77.5	22.5	868	47			84
855	77	23	862.5	46.5	34	32.04	83.65
856	76.5	23.5	857	46			83.3



# INTRODUCTION.

Sp. gr. at 60° Fah.	100 parts contain of		1000 parts contain of standard alc. sp. gr. 825.	Sikes.	Baumé.	Cartier.	Per cent. of alcohol, sp. gr. 796 by volume. Gay Lussac.
	Alcohol sp. gr. 796.	Water. By weight.					
857	76	24	853	45.5 O. P.	33.5	31.58	83
858	75.67	24.33	849	45			82.7
859	75.33	24.67	844.5	45			82.35
860	75	25	840	45	33	31.13	82
861	74.67	25.33	836.5	44.5		31	81.7
862	74.33	25.67	833	44			81.3
862.5	74.16	25.84	830.5	43.75	32.5	30.76	80
863	74	26	828	43.5			80.8
864	73.5	26.5	823	43			80.3
865	73	27	818	42.5	32	30.21	79.95
866	72.5	27.5	813	42			79.6
867	72	28	810	41			79.3
867.5	71.83	28.17	808.5	40.5	31.5	29.78	79.15
868	71.67	28.33	807	40			79
869	71.33	28.67	802.5	39.5			78.65
870	71	29	798	39	31	29.29	78.3
871	70.5	29.5	792.5	38.5			78
872	70	30	787	38		29	77.7
873	69.5	30.5	781.5	37	30.5	28.83	77.35
874	69	31	776	36			77
875	68.67	31.33	772	35			76.5
876	68.33	31.67	768	34	30	28.38	76
877	68	32	762.5	33			75.65
877.5	67.75	32.25	759.25	32.5		28	75.5
878	67.5	32.5	757	32			75.3
878.5	67.25	32.75	753.75	31.5	29.5	27.91	75
879	67	33	751.5	31			74.8
880	66.5	33.5	746	30			74.3
881	66	34	742	29.5	29	27.44	74
882	65.5	34.5	738	29			73.7
883	65	35	733.5	28.5		27	73.35
883.5	64.83	35.17	731.25	28.25	28.5	26.99	73.17
884	64.67	35.33	729	28			
885	64.33	35.67	724	27.5			72.5
886	64	36	719	27	28	26.53	72
887	63.67	36.33	714	26			71.5
888	63.33	36.67	709	25			71
889	63	37	704	24.5	27.5	26.07	70.65
890	62.5	37.5	699	24			70.3
891	62	38	694	23			69.8
892	61.5	38.5	689	22	27	25.61	69.3
893	61	39	684.5	21			69
894	60.67	39.33	680	20			68.7
895	60.33	39.67	675.5	19.5			68.35
895.5	60.16	39.84	673.25	19.25	26.5	25.15	68.17
896	60	40	671	19		25	68
897	59.5	40.5	666.5	18			67.65
898	59	41	662	17	26	24.69	67.3
899	58.5	41.5	655.5	16			67
900	58	42	649	15			66.7
900.5	57.75	42.25	647	14.75	25.5	24.23	66.52
901	57.5	42.5	645	14.5			66.35
901.5	57.25	42.25	643	14.25		24	66.17



Sp. gr. at 60° Fah.	100 parts contain of		1000 parts contain of standard alc. sp. gr. 825.	Sikes.	Baumé.	Cartier.	Per cent. of alcohol, sp. gr. 796 by volume. Gay Lussac.
	Alcohol sp. gr. 796.	Water. By weight.					
902	57	43	641	14 O. P.			66
903	56.5	43.5	636	13	25	23.77	65.5
904	56	44	631	12			65
905	55.5	44.5	626	11.5			64.5
906	55	45	621	11	24.5	23.31	64
907	54.5	45.5	616.5	10.5			63.65
908	54	46	612	10		23	63.3
909	53.5	46.5	607	9	24	22.85	62.65
910	53	47	602	8			62.3
911	52.5	47.5	595.5	7.5			61.9
912	52	48	591	7	23.5	22.39	61.5
913	51.67	48.33	586	6			61
914	51.33	48.67	581	5			60.5
915	51	49	576	4	23	21.94	60
916	50.5	49.5	571	3			59.6
917	50	50	560.5	2			59.3
918	49.67	50.33	562	1	22.5	21.48	59
919	49.33	50.67	554	.5			58.5
920	49	51	550	Proof			58
921	48.5	51.5	545	1 U. P.	22	21.02	57.5
922	48	52	540	2			57
923	47.5	52.5	535.5	3	21.5	20.56	56.5
924	47	53	531	4			56
925	46.5	53.5	526	5			55.5
926	46	54	521	6	21	20.10	55
927			515.5	6.5			
928	45	55	510	7			54
929	44.5	55.5	505	8			53.5
929.5	44.25	55.75	502.5	8.5	20.5	19.64	53.25
930	44	56	500	9			53
931	43.67	56.33	495.5	10			52.5
932	43.33	56.67	489	11			52
933	43	57	484	12	20	19.18	51.5
934	42.5	57.5	479	13		19	51
935	42	58	472.5	14			50.5
936	41.5	58.5	468	15	19.5	18.72	50
937	41	59	462	16			49.5
938	40.5	59.5	456	17			49
939	40	60	450	18	19	18.26	48.5
940	39.5	60.5	444	19			48
940.5	39.25	60.75	441	19.5		18	47.63
941	39	61	438	20			47.25
942	38.5	61.5	432	21	18.5	17.80	46.5
943	38	62	426.5	22			45
944	37.5	62.5	421	23			45.5
945	37	63	416	23.5	18	17.35	44.75
946	36.5	63.5	411	24			44
947	36	64	399	25		17	43.5
948	35.5	64.5	397	26	17.5	16.89	43
949	35	65	389.5	27			42.25
950	34.5	65.5	382	28			41.5
951	34	66	376	29.5	17	16.43	40
952	33.5	66.5	370	31			40.5



Sp. gr. at 60° Fah.	100 parts contain of		1000 parts contain of standard alc. sp. gr. 825.	Sikes.	Baumé.	Cartier.	Per cent. of alcohol, sp. gr. 796 by volume. Gay Lussac.
	Alcohol sp. gr. 796.	Water. By weight.					
953	33	67	364	32.5 U. P.			39.75
954	32.5	67.5	358	34			39
955	32	68	352	35	16.5	16.3	38.5
956	31.5	68.5	346	36		16	38
957	31	69	339.5	37.5			37.25
958	30	70	333	39			36.5
959	29.5	70.5	324	40.5	16	15.51	35.75
960	29	71	315	42			35
961	28.5	71.5	307.5	43.5			34.5
962	28	72	300	45	15.5	15	34
963	27	73	292.5	46.5			33
964	26.5	73.5	285	48			32
965	26	74	277.5	49.5	15	14.59	31
966	25.5	74.5	270	51			30
967	25	75	261.5	52.5			29
968	24	76	253	54			28
968.5	23.75	76.25			14.5	14.13	27.5
969	23.5	76.5	244.5	55.5			27
970	23	77	236	57			26
971	22.5	77.5	227	58.5			25
972	22	78	218	60	14	13.67	24
973	21	79	209	62			23
974	20	80	200	64			22
975	19	81	195	66	13.5	13.21	21
976	18.5	81.5	190.5	68			20
977	18	82	183.5	70			19
978	17	83	175	72	13	12.76	18
979	16	84	163	73.5			17
980	15.5	84.5	150	75			16
981	15	85	143	76			15
982	14	86	135	77	12.5	12.30	14
983	13.5	86.5	128	78.5			13
984	13	87	120	80			12
985	12.5	87.5	112	81			11.25
986	12	88	105	82	12	11.84	10.5
987	11	89	98	83.5			9.75
988	10	90	90	85			9
989	9	91	82	87	11.5	11.38	8
990	8	92	75	89			7
991	7	93	67.5	90.5			6.5
992	6	94	60	92			6
993	5.5	94.5	52.5	93.5	11	10.92	5
994	5	95	45	95			4
995	4	96	37.5	95.5			3.5
996	3.5	96.5	30	96	10.5	10.46	3
997	3	97	22.5	97			2
998	2	98	15	98			1
999	1	99	7.5	99			.5
1000	0	100	0	100	10	10	0



SPECIFIC GRAVITIES OF SOME OF THE PREPARATIONS ORDERED  
IN THE PHARMACOPŒIAS.

The London Pharmacopœia directs the Specific Gravity to be taken at a temperature of 62° Fahr. The Edinburgh, Dublin, and U.S. Pharmacopœias at 60° Fahr.

						Sp. gr.
Acetum Destillatum	-	-	-	-	Edin.	1.005
	-	-	-	-	Dublin	1.005
Acidum Aceticum	-	-	-	-	London	1.048
	-	-	-	-	U. S.	1.063
	-	-	-	-	Edin.	to
	-	-	-	-		1.068
	-	-	-	-	Dublin	1.074
	-	-	-	-	Dublin	1.005
Dilutum	-	-	-	-	London, U. S.	1.160
Hydrochloricum	-	-	-	-	Edin.	1.170
	-	-	-	-	Dublin	1.160
	-	-	-	-	Edin.	1.050
Dilutum	-	-	-	-	U. S.	1.046
	-	-	-	-	Dublin	1.080
Hydrocyanicum	-	-	-	-	Dublin	.998
Nitricum	-	-	-	-	London, U. S.	1.500
	-	-	-	-	Edin.	1.500
Dilutum	-	-	-	-	London, U. S.	1.080
	-	-	-	-	Edin.	1.077
	-	-	-	-	Dublin	1.280
Phosphoricum Dilutum	-	-	-	-	London	1.064
Sulphuricum	-	-	-	-	London, U. S.	1.845
	-	-	-	-	Edin.	1.845
	-	-	-	-	Dublin	1.845
Sulphuricum venale	-	-	-	-	Dublin	1.850
Sulphuricum Dilutum	-	-	-	-	London	1.110
	-	-	-	-	Edin. U. S.	1.090
	-	-	-	-	Dublin	1.084
Æther Nitrosus	-	-	-	-	Dublin	0.900
Sulphuricus	-	-	-	-	London, U. S.	0.750
	-	-	-	-	Edin.	0.735
	-	-	-	-	Dublin	0.766
Alcohol	-	-	-	-	London	0.815
	-	-	-	-	U. S.	0.835
	-	-	-	-	Edin.	0.796
	-	-	-	-	Dublin	0.810
Dilutum	-	-	-	-	U. S.	0.935
Aqua Destillata	-	-	-	-	L. E. D.	1.000
Ammoniaë	-	-	-	-	Edin.	0.960
	-	-	-	-	Dublin	0.950
Acetatis	-	-	-	-	Edin.	1.011
	-	-	-	-	Dublin	1.011
Ammoniaë Carbonatis	-	-	-	-	Dublin	1.090
Barytæ Muriatis	-	-	-	-	Dublin	1.230
Calcis Muriatis	-	-	-	-	Dublin	1.202
Aqua Potassæ	-	-	-	-	Edin.	1.072
	-	-	-	-	Dublin	1.080
Carbonatis	-	-	-	-	Dublin	1.320
Potassii Sulphureti	-	-	-	-	Dublin	1.117
Sodæ Carbonatis	-	-	-	-	Dublin	1.024



Liquor Ammoniaë	-	-	-	-	-	London	Sp. gr. 0.960
Fortior	-	-	-	-	-	London, U. S.	0.882
Plumbi Diacetatis	-	-	-	-	-	London	1.260
Potassæ	-	-	-	-	-	London	1.063
Carbonatis	-	-	-	-	-	London	1.473
Oleum Æthereum	-	-	-	-	-	London	1.05
	-	-	-	-	-	U. S.	1.096
Spiritus Æthereus Nitrosus	-	-	-	-	-	Dublin	0.850
Ætheris Nitrici	-	-	-	-	-	London, U. S.	0.834
	-	-	-	-	-	Edin.	0.847
Sulphurici	-	-	-	-	-	Edin.	0.809
Ammoniaë	-	-	-	-	-	London	0.860
	-	-	-	-	-	U. S.	0.831
Aromaticus	-	-	-	-	-	London	0.914
Fœtidus	-	-	-	-	-	London	0.861
Rectificatus	-	-	-	-	-	London	0.838
	-	-	-	-	-	Edin.	0.838
	-	-	-	-	-	Dublin	0.840
Tenuior	-	-	-	-	-	London	0.920
	-	-	-	-	-	Edin.	0.912
	-	-	-	-	-	Dublin	0.919
Tinctura Ferri Sesquichloridi	-	-	-	-	-	London	0.992

### RELATION BETWEEN DIFFERENT THERMOMETRICAL SCALES.

The thermometer always used in this country is that of Fahrenheit; it is also used in parts of Germany.

In this instrument, the range between the freezing and boiling points of water is divided into 180°, and as the greatest possible degree of cold was supposed to be that produced by mixing snow and salt together, it was made the zero. Hence, the freezing point became 32°, and the boiling point 212°.

The Centigrade thermometer places the zero at the freezing point of water, and divides the range between the freezing and boiling points into 100°. This scale has long been used in Sweden, under the title of Celsius's thermometer, and is generally adopted on the Continent.

Reaumur's thermometer, which was formerly used in France, divides the space between the freezing and boiling points of water into 80°, and places the zero at the freezing point. It is now little employed.

Le Lisle's thermometer is used in Russia. The graduation begins at the boiling point, and increases towards the freezing point. The boiling point is marked 0°, and the freezing point 150°.

*To reduce Centigrade degrees to those of Fahrenheit.*

RULE.—Multiply by 9, divide by 5, and add 32.

$$\text{Thus, } 40^{\circ} \times 9 \div 5 + 32 = 104^{\circ} \text{ Fahr.}$$

*To reduce Fahrenheit's degrees to those of Centigrade.*

RULE.—Subtract 32, multiply by 5, and divide by 9.

$$\text{Thus, } 104^{\circ} \text{ Fahr.} - 32 \times 5 \div 9 = 40^{\circ} \text{ Cent.}$$

*To reduce Reaumur's degrees to those of Fahrenheit.*

RULE.—Multiply by 9, divide by 4, and add 32.

$$\text{Thus, } 32^{\circ} \text{ Reaumur.} \times 9 \div 4 + 32 = 104^{\circ} \text{ Fahr.}$$



*To reduce Fahrenheit's degrees to those of Reaumur.*

RULE.—Subtract 32, multiply by 4, and divide by 9.

$$\text{Thus, } 104^{\text{Fahr.}} - 32 \times 4 \div 9 = 32^{\text{Reaum.}}$$

*To reduce Reaumur's degrees to those of Centigrade.*

RULE.—Multiply by 5, and divide by 4.

$$\text{Thus, } 32^{\text{Reaum.}} \times 5 \div 4 = 40^{\text{Cent.}}$$

*To reduce Centigrade degrees to those of Reaumur.*

RULE.—Multiply by 4, and divide by 5.

$$\text{Thus, } 40^{\text{Cent.}} \times 4 \div 5 = 32^{\text{Reaum.}}$$

## EXPLANATION OF THE PRINCIPAL ABBREVIATIONS OCCURRING IN PHARMACEUTICAL FORMULÆ.

R. Recipe—Take.

F. S. A. Fiat secundum artem—Let it be made or prepared according to the rules of the art.

M. Misce—Mix.

M. S. D. Misce, signa, da—Mix the medicine, and deliver it afterwards, with the requisite instruction, to the patient (or nurse) in writing.

M. F. P. Misce fiat pulvis—Mix to form a powder.

M. F. Mixt. Misce fiat mixtura—Mix to form a liquid mixture.

M. F. Pil. Misce fiant pilulæ—Mix to form pills.

Div. Divide—Divide.

Sol. Solve—Dissolve.

Fasc. j. Fasciculus—An armful.

Man. j. Manipulus—A handful, a gripe.

Pugil. j. Pugillus or Pugillum—A pinch.

Cyat. j. Cyathus—A glassful.

Cochl. j. Cochlear or cochleare—A spoonful.

Gutt. Gutta—Drop.

No. 1, 2, 3, &c. The number of pieces or parts.

Ana, or āā. Of each.

P. Ae. Partes æquales—Equal parts.

Q. S. Quantum sufficit—As much as will suffice.

Q. L. Quantum libet—As much as you like.

Q. V. Quantum volueris—As much as you like.

℔. Libra—A pound.

℥. Uncia—An ounce.

ʒ. Drachma—A drachm or dram.

ʒ. Scrupulus—A scruple.

Gr. Granum—A grain.

Pil. Pilula—A pill.

Pot. Potio—A potion.

Pulv. Pulvis—A powder.

Tinc. Tinctura—A tincture.

Ext. Extractum—An extract.

Chart. Chartula—A small paper.

Collyr. Collyrium—An eye water.

Collutor. Collutorium—A mouth wash.

Cong. Congius—A gallon.

O. Octarius—A pint.



f3. Fluiduncia—A fluidounce.  
 f3. Fluidrachma—A fluidrachm.  
 m. Minimum—A minim.  
 Decoct. Decoction—A decoction.  
 Garg. Gargarisma—A gargle.  
 Haust. Haustus—A draught.  
 Infus. Infusum—An infusion.  
 Mass. Massa—A mass.  
 Mist. Mistura—A mixture.  
 Ss. Semis—A half.

The use of the Latin language for prescriptions and for the accompanying directions is universal in Europe, except in France, where, although the prescription itself may be in Latin, the directions are usually given in French; this is also the case in the United States, where few, if any, physicians give the directions for the use and mode of administration of a formula in any other language than English. As, however, it often becomes of importance to refer to the prescriptions of foreign practitioners, the following table, from Mowbray's *Conspectus*, is introduced:—

### A VOCABULARY

*Of the Words most frequently occurring in the Prescriptions of Physicians.*

#### A.

A, āā, or ana, of each ingredient; more promptly expressed by the genitive case plural (singulorum) of each; for instance:  
 R.—Aquæ cinnamon, tinct. rhei, āā ʒij, means  
 Take of cinnamon-water, and tincture of rhubarb, of each two drachms.  
 Abdomen, the belly. Gen.—inis, of the belly. Dat.—ini, to the belly.  
 Absente febre, while the fever is off.  
 Accuratè (adverb), accurately. Accuratè pensi, weighed with the utmost exactness.  
 Accuratè misceantur, mix very completely.  
 Aciditas, sharpness. Ad gratam aciditatem, make it just acid enough to be palatable, and not to sour.  
 Acme, the height of a fever or any other disease.  
 Ad, to. Ad duas vices, at twice taking. Ad libitum, as you like.  
 Adde, add. Addantur, let there be added. Addendus, to be added. Addendo, adding.  
 Admoveatur, let there be applied—antur, plural, when more than one is to be applied.  
 Adstante febre, while the fever is on.  
 Æger, a sick person, a patient. Ægra, a female patient.  
 Aggrediente febre, when the fever is coming on.  
 Aggressus, an attack. Aggressus febris, the attack of a fever.  
 Agitato vasè, shaking the phial.  
 Albus, white.  
 Aliquot, some. Aliquoties, sometimes.  
 Alter, altera, alterum, the other.  
 Alternus, alternate. Alternis horis, every second hour. Alternis diebus, every alternate day.  
 Aluta, leather. Extende super alutam mollem, spread on soft leather.  
 Alvus, the belly, the bowels. Adstrictâ alvo, when costive.  
 Amplus, large. Cochleare amplum, a tablespoonful.  
 Antè, before. Ut antè, as before.  
 Antemeridianus, in the forenoon; any time before twelve o'clock at noon.  
 Apparatus, any sort of preparation, instruments, or, in short, everything that is requisite to be had in readiness for performing any sort of operation. Also, the more delicate term for a bladder and pipe for clysters.  
 Aperiens, opening, gently purging.  
 Applicetur, let there be applied.



Aqua, water. Aquæ, of water. Aqua bulliens, Aqua fervens, boiling water.  
 Armatus, armed. Fistula armata, an apparatus for clysters; a pipe and bladder.  
 Auris, the ear. Auri, to the ear. Aures, ears. Auribus, to the ears.  
 Aut, or.

## B.

Benè, well.  
 Bibat, let him drink.  
 Biduum, two days. Omni biduò vel triduò, every two or three days.  
 Bihorium, the space of two hours. Omni bihorio, every two hours.  
 Bis, twice; bis, terve, twice or thrice.  
 Bullio, to boil; bulliens, boiling. Aquæ bullientis, of boiling water.  
 Bulliant, let them boil.

## C.

Calefactus, made warm.  
 Capiat, let (the patient) take. Capt. cochl. iij magn., three tablespoonfuls to be taken.  
 Catharticus, purging.  
 Cautè, cautiously.  
 Cerevisia, beer. Cerevisia Londinensis, porter.  
 Charta, paper. Charta cærulea, blue paper. Chartula, a little piece of paper. Cola trans chartam, filter through paper.  
 Cibus, food.  
 Circa, about. Circiter, about.  
 Citò, soon, quickly. Citissimè (the superlative degree), as quick as possible.  
 Clausus, clausa, clausum, covered. Vase clauso, in a covered vessel.  
 Cochleare, a spoonful. Cochlearia, spoonfuls. Coch. magnum, amplum, a table-spoonful. Co. infantulorum, coch. modicum, child's spoonful, a dessertspoonful. Co. parvum, a teaspoonful.  
 Coctio, a boiling. Sub finem coctionis, towards the end of the boiling, when almost boiled.  
 Cæruleus, blue. Charta cærul., blue paper. Unguentum cæruleum, mercurial ointment.  
 Cola (imperat. of Colo), strain, to filter.  
 Colatus, strained, filtered. Colato liquori, to the strained liquor.  
 Colatura, a straining. Colaturæ, to the strained liquor.  
 Colatus, let be strained. Colentur, let them be strained.  
 Collum, a neck.  
 Comp., Compositus, compound.  
 Continuo, to continue. Continuantur remedia, go on with the same medicines as last prescribed.  
 Contundo, to bruise. Contusus, bruised.  
 Contusio, a bruise, a contusion.  
 Coquo, to boil. Coque, boil. Coque parùm, boil a little while.  
 Coquantur, let them be boiled.  
 Cor, the heart. Scrobiculus cordis, the pit of the stomach.  
 Cōxa, coxendix, the hip.  
 Cras, to-morrow; c. m., cras mane, early to-morrow morning; c. m. s., cras mane sumendus, to be taken early to-morrow morning.  
 Crastinus (adj.), of to-morrow. In usum crastinum, for to-morrow's use.  
 Cribrum, a sieve. Trans cribrum, through a sieve.  
 Cujus, of which. Cujus capiat, orumat, of which (the patient is to) take.  
 Cum, with.  
 Cyathus, a cup. Cyathus vinarius, a wineglass. Cyath. theat, a teacup full.

## D.

Da, give. Detur, dentur (plural), let be given.  
 De, of. De quo, or quâ, of which. De die, in a day.



Deauratus, gilt. Deaurentur pilulæ, let the pills be gilt.  
 Debilitas, weakness.  
 Decem, ten. Decimus, the tenth.  
 Debitus, due. Ad debitam spissitudinem, to a proper degree of thickness, as to consistence.  
 Decubitus, lying down. Horâ decubitûs, at bedtime.  
 Deglutio, to swallow. Deglutiatur, is swallowed—etur, let be swallowed.  
 Dein, deinde, then; afterwards.  
 Dejectio, a depositing, or putting down, also a going to stool; as, post duas dejectiones alvi, after two motions.  
 Dejicio, to deposit. Donec alvus bis dejiciat, until the patient shall have had two stools.  
 Dexter, the right. Manus dextra, the right hand. Auri dextro, to the right ear.  
 Dictus, spoken of, said.  
 Dies, a day. Die, in a day; as, bis die, twice a day. Diebus, in days; as, tertiis diebus, every third day. Alternis diebus, every alternate day.  
 Dilutum, diluted.  
 Dimidium, the half. Dimidius, a, um (adjective), half.  
 D.P., Directione propria, with its proper direction.  
 Diu, a long time. Tere diu, rub for a long time.  
 Diurnus, long continued. Diurnâ trituratione, by long-continued rubbing, or grinding in the mortar. Diurna coctione, by long boiling.  
 Dolor, pain. Dolores, pains. Doloris, of the pain.  
 Dolens, pained. Parti dolenti, to the pained part.  
 Donec, until. Dum, whilst.  
 Durante dolore, while the pain continues.

## E.

Eadem, eandem, the same. Eodem, in the same.  
 Effunde, pour out.  
 Effervescentia, the effervescence.  
 Effervescet, effervesces.  
 Ejusdem, of the same, the genitive case of idem.  
 Enema, a clyster. Enemata, clysters.  
 Erit, shall be.  
 Evanescio, evanui, to disappear. Evanuerit, shall have disappeared.  
 Exhibe, exhibeatur, give. Exhibendus, to be given.  
 Extendo, to extend, also to spread. Extende super alutam, spread it on leather.  
 Extensus, a, um, spread.

## F.

Fac, make. Fac in pilulas xij, make into twelve pills.  
 Farina, flour. Farina seminis lini, linseed meal.  
 Febris, a fever. Febre durante, while the fever is on.  
 Femur, a thigh.  
 Femoribus, to the thighs. Femoribus internis, to the inner sides of the thighs.  
 Fervens, boiling. Fervidus fervida, hot.  
 Ferventis, of boiling. R aq. ferventis, take of boiling water.  
 Finis, the conclusion. Sub finem coctionis, when almost boiled enough.  
 Fiat, make (the singular number). Fiant, make (plural).  
 Fontana, fountain. Aqua fontana, spring water.  
 Formula, prescription.  
 Fetus, a fomentation.  
 Fuerit, shall have been; as, donec alvus soluta fuerit, until a motion is (or shall have been) procured.

## G.

Gargarisma, a gargle.  
 Gelatina, jelly. Gelatina ribesiorum, currant jelly. Gelatina quovis, any sort of jelly.



Globulus, a little ball. Globuli Gascoigni, Gascoign's ball. Donec globuli evanuerint, until the globules (of quicksilver) totally disappear (so that they cannot be seen even with a microscope).

Gradatim, by slow degrees.

Gratus, grata, gratum, agreeable, pleasant. Ad gratam aciditatem, so as to make it pleasantly acid without being too sour. In quovis grato vehiculo, in any agreeable vehicle.

Gutta, a drop. Guttæ, drops. Guttas, drops.

Guttatim, drop by drop.

## H.

Hac, this. Hac nocte, this night. Hanc, this. Sumat hanc, let him take this.

Hactenus, hitherto, heretofore, up to the present day.

Harum, of these. Harum pilularum sumat tres, of these pills let him or her take three.

Haustus, a draught.

Hebdomada, a week.

Heri, yesterday. Ut heri, as yesterday.

Hesternus, of yesterday. Hesternæ nocte, last night.

Hirudo, a leech. Hirudines, leeches.

His, in these, to these. His adde, add to these.

Hora, an hour. Horæ, of an hour. Horæ (plural), hours.

H.S. (horâ somni), at the hour of rest.

H.S.S. (horâ somni sumendus), to be taken at bedtime. Hora decubitus, at the hour of going to rest, bedtime.

Horâ vespertinâ, in the evening.

Horæ unius spatium, in the space of one hour.

Horæ  $\frac{1}{4}$ , horæ quadrante, quarter of an hour.

Horis intermediis, at intermediate hours, when two medicines are to be taken. Horis intermediis means that one is to be given exactly at midtime from the other: suppose a draught is ordered (to be taken every six hours), and a powder horis intermediis, that is every six hours intermediately, then a draught will be taken at six o'clock and at twelve, and a powder at three and at nine.

Hujusmodi, of this sort, like these.

H. p. n., Haustus purgans noster, a formula of purging draught made according to a practitioner's own private Pharmacopœia, and is prepared so as to keep a long time without spoiling, that we may not have the trouble of preparing it every time a draught is wanted. Mitt. H. p. n.  $\mathfrak{z}$ ij ad ij Vices c. m. s. Mitte Haustus purgantis nostri uncias duas, ad duas Vices cras mane sumendus; send two ounces of our purging draught, to be taken to-morrow morning, at twice, that is half at first, and the remaining half in an hour if the first do not operate.

## I.

Idoneus, proper, appropriate.

Idem, eadem, the same. Ejusdem, of the same, the genitive case of idem.

Imponatur (sing.), let there be put on—nantur (plural).

Imprimis, first.

In, in. In die, in a day.

Indiè (an adverb), every day, daily.

Indicaverit, shows, indicates.

Infusio, infusum, an infusion.

Infunde, infundatur, infuse.

Inter, between.

Injectio, injection.

Inquietudo, restlessness. Ungente inquietudine, if restless.

Injiciatur, throw in, throw up. Injiciatur enema, let a clyster be administered.



Injiciendus, injiciendum, to be administered.

Instar (an adverb), as big as. Sumat molem instar mucis moschatae, the bigness of a nutmeg to be taken.

Intermedius, intermediate.

Internus, the inner side.

## J.

Jam, already. Jampridem, jamdudum, some time ago, heretofore.

Jugulum, the throat.

Jusculum, broth. Jusculum ovillum, mutton broth. Jus bovinum, beef tea.

Juxta, near to.

## L.

Lac, milk. Lactis, of milk. Lacte, in milk.

Laevigatus, levigated.

Lana, flannel. Lana nova, new flannel.

Languor, faintness, lowness. In languoribus, in the fainting fits, when low and faint.

Latus, the side. Lateris, of the side. Lateri, to the side. Latus dolens, the pained side. Lateri dolenti, to the pained side.

Latus, lata, latum (adjective), broad.

Lectus, a bed. In lecto, in bed.

Liber, a book.

Libra, a pound. Libris, libras, pounds.

Linteum, lint; also linen.

Liquesco, to liquefy, to melt. Donec liquescat, till it melts.

Liquidus, liquid. Sedes liquidæ, loose stools. In quovis liquido, in any liquid.

Londinensis, of London. Pharmacopœia Londinensis, the London Dispensatory.

Lumborum, of the loins.

Libet (a verb impersonal), it pleases. Ad libitum, just as you please.

## M.

Magnus, magna, magnum, large. Magnum cochleare, a tablespoonful.

Major, greater, larger. Cochlearia duo majora, two tablespoonfuls.

Maximus, the greatest. Maximâ curâ, with the greatest care.

Maximè, chiefly.

Malleolus, the ankle. Malleolus internus, the inner ankle.

Manè, in the morning. Manè primo; valde mane, very early in the morning.

Manus, a hand. Manu calefacta, with a warm hand.

Massa, a mass. Massa pilularis, a mass fit for forming into pills.

Matutinus, in the morning or forenoon.

Mediocris, middle-sized. Pilulæ mediocres, middle-sized pills. Cochleare mediocre, a dessertspoonful, a pap-spoonful, or a child's spoonful. Mediocris also means indifferent, as to quality.

Medius, middle. Media nocte, in the middle of the night.

Melior, melius, better.

Minatur, minaretur, threatens. Minante, threatening.

Minimus, very small. Cochleare minimum, a teaspoonful.

Minutum,\* a minute.

Misce, mix. Bene misceatur, let it be well mixed.

Mite, send.

Mittatur (singular), let it be sent. Mittantur (plural), let them be sent.

Mittatur sanguis, take blood away; i. e., bleed the patient.

Mistura, mixtura, a mixture.

Mitigatio, mitigation, alleviation. Donec dolor is mitigatio sit, until the pain is easier.

Mitigatus, a, um, mitigated, lessened.

\* The word minutum, for a minute, is very barbarous Latin; we believe there is no such word; but the right Latin for a minute, Sexagesima pars horæ, is as long and as tiresome to write as "Semivitreous oxide of lead" for the simple word "Litharge."



Modicus, middle-sized.

Modus, a manner. Modo præscripto, in the manner directed.

Moles, a mass, a lump, a piece. Sumat molem instar nucis moschatæ, let him (or her) take the bigness or size of a nutmeg.

Mollis, molle, soft.

Molestus, troublesome. Molestor, to trouble, to be troublesome. Molestante dolore, when the pain is troublesome. Molestante tussi, when the cough is troublesome.

Mora, delay. Sine morâ, without delay.

Mos, moris, manner. More solito, in the usual manner; also, in the same manner as I am in the habit of prescribing it to other people.

Mortarium, a mortar. Mortario aheneo, in the brass mortar. Mortario marmoreo, in the marble mortar. Mort. vitreo, in the glass mortar.

## N.

Narthecium, a gallipot.

Nates, the buttocks.

Ne, lest, also, do not; as, ne tradas sine nummo, do not deliver the medicine without the money.

Necnon, also.

Niger, nigra, nigrum, black.

Ni, nisi, unless.

Nihil, nothing.

Nisus, an endeavor, an attempt, a straining, a motion, a straining to vomit, or go to stool.

Nimis, nimium, too much.

Nodulus, a little knot. Nodulo ligati, tied up in a piece of clean rag.

Nomen, nominis, a name. Signetur nomine proprio, write its common name upon the label.

Novem, nine.

Nonus, the ninth.

Novus, nova, novum, new.

Novissimè, very lately, the last of all.

Nocte, at night. Noctes, nights. Nocte maneque, night and morning. Alternis noctibus, every second night.

Nuper, lately. Nuperrime, very lately.

Nucha, the nape of the neck. Nuchæ, to the nape.

N.M., Nux moschata, a nutmeg. Sumat magnitudinem nucis moschatæ, take the bigness of a nutmeg.

## Numeri. Numbers.

1. or j. unus, una, unum, one.

unius, of one.

2. ij. duo, duæ, two.

duorum, of two.

duobus, in two, to two.

3. iij. tres, tria, three.

tribus, in three, to three.

trium, of three.

ter, three times.

4. iv. quatuor, four.

quartus, a, um, fourth.

quater, four times.

5. v. quinque, five.

quintus, fifth.

6. vj. sex, six.

sextus, sixth.

7. or vij. septem, seven.

septimus, seventh.

septimana, or,

7 mana, a week.

8. viij. octo, eight.

octavus, eighth.

9. ix. novem, nine.

nonus, ninth.

10. x. decem, ten.

decimus, tenth.

11. xj. undecim, eleven.

12. xij. duodecim, twelve.

20. xx. viginti, twenty.

24. xxiv. viginti quatuor, twenty-four.



## O.

Obstante, hindering, preventing.

Occasio, occasion, opportunity.

Octo, eight.

Octavus, eighth. Octava quaque horâ, every eighth hour.

Olim, some time ago.

Olla, a pot, a gallipot.

Omnis, all. Omni mane, every morning. Omni horâ, every hour. Omni bihorio, every two hours. Omni nocte, every night. Omni  $\frac{1}{4}$  horâ, Omni quadrante horæ, every quarter of an hour. Tere omnia, rub all together.

Omninò, quite, wholly, entirely.

Optimè, very well, as well as possible.

Optimus, ma, mum, best.

Opus, need, occasion. Si opus sit, or fuerit, if it be necessary.

Ovillum jusculum, mutton broth.

Ovum, an egg. Vitell. ovi, the yolk of an egg. Ovorum, of eggs.

## P.

Pannus, a rag. Pannus linteus, a linen rag. Pannus laneus, Pannus è lanâ, a piece of flannel.

Paroxysmus, a paroxysm, a fit, a convulsion fit.

Pars, a part. Partes, parts. Parti dolenti, to the pained part.

Partitus, parted. Partitus vicibus means that you are not to give a medicine all at once, but divide the dose according to the directions most commonly previously given; for example, if a purging or emetic draught, half or a third (as the case may be) to be taken at first, and the other half or third at a certain distance of time, if the former quantity be not found sufficient to produce the desired effect.

Parvus, little. Cochleare parvum, a teaspoonful.

Parum, a little. Parumper, a little.

Pauculum, pauxillum, paululum, a little.

Pectus, the breast. Pectoris, of the breast.

Pediluvium, a bath for the feet.

Per, by, or through.

Pergo, to go on with. Pergat in usu medicinarum, continue the medicines as before.

Peractus, { completed, perfected, quite done, gone through with; as, Peractâ ope-

Perfectus, { ratione emetici, after the emetic has quite done operating.

Perfricetur, let it be rubbed. Perfricandus, to be rubbed.

Pensus, weighed. Accuratè pensi, exactly weighed.

Perpetuus, perpetual. Fiat perpetuum, keep it open (when it refers to a blister).

Pluvialis, also pluviatilis aqua, rain-water.

Pharmacopœia, the dispensatory.

Pollex, the thumb. Pollex pedis, the great toe.

Pomeridianus, postmeridianus, in the evening or afternoon, time of the day.

P. R. N., Pro re natâ, occasionally, according as circumstances may occur, according as the symptoms may require.

Poculum, a cup.

Pomum, an apple. Pomi, of an apple.

Pone, behind. Pondere, by weight.

Porro, moreover. Post, after. Postea, then, afterwards.

P. M., Post meridiem, afternoon, in the afternoon, after twelve o'clock at noon.

Potus, drink, any kind of beverage.

Postulet, postulent, may require, demand.

Præcipue, especially.

Præparatus, prepared.

Præparo, to prepare. Præparentur, let them be prepared.



Primus, ma, mum, the first. Primò, first of all. Primo mane, very early in the morning.  
 Prior, prius, the former, the first.  
 Priusquam, before that.  
 Pro, for. Pro re natâ, p. r. n., occasionally, &c. Pro ratione, according to, or, in proportion to; as, pro ratione ætatis, according to the age of the patient.  
 Proprius, a, um, proper. S. N. P., Signetur nomine proprio, mark it with its proper direction.  
 Pruritus, an itching. Pruriens; itching. Dolichos pruriens, cowhage.  
 Psora, the itch.  
 Pulvis, a powder. Pulveres, powders. Pulveribus, in powders.  
 Purificatus, a, um, purified.  
 Pyxis, pyxidis, a pill-box, or lozenge-box.

## Q.

Quadrans, quadrantis, quadrante, quarter. Omni quadrante horæ, every quarter of an hour.  
 Quacum, with which.  
 Quamprimum, as soon as possible, without the least delay.  
 Quartus, a, um, the fourth.  
 Quâqua, every. Quâvis (foem), with any.  
 Quater, four times. Quatuor, four. Quarta pars, a fourth part.  
 —que (at the end of any Latin word), and.  
 Quem, quam, quod, which (the accusative case).  
 Qui, quæ, quod, which (the nominative case).  
 Quibus, to which, with which.  
 Quibusdam, to or with some. Cum guttis quibusdam, with a few drops.  
 Quilibet, quaelibet, quodlibet, gen. Cujuslibet, abl. Quolibet, any.  
 Quiescat, goes to rest, is easier.  
 Quinque, five. Quintus, a, um, the fifth. Quinquies, five times.  
 Quinquina (cinchona), Peruvian bark.  
 Q. S.—q. s., Quantum sufficiat, as much as may be sufficient.  
 Quorum, quarum, of which. Quos, quas, which. Quocum, quacum, with which.  
 Quovis, with any.

## R.

Ratio, a reason, also a proportion. Pro ratione ætatis, according to the age of the patient. Pro ratione doloris, according to the urgency of the pain.  
 Raucedo, hoarseness.  
 Recipe, take. Recipe (taken substantively), a prescription.  
 Regio, a region (an anatomical term for certain parts of the body); as, regio epigastrica, the epigastric region; regio lumborum, the region of the loins; appl. emplastr. regioni umbilicali, to the umbilical region or parts in the neighborhood of the navel.  
 Redigo, to reduce. Redige, reduce. Redigetur, may be reduced. Redigatur, let it be reduced.  
 Redactus, a, um, reduced. In pulv. redact., reduced to powder.  
 Refriferit—pl. int—the subjunctive future of refrigescere, to cool.  
 Reliquus, a, um, the rest, the remaining part.  
 Remedium, a remedy. Continuandur remedia, go on with the same medicines as before.  
 Repetatur, let be repeated, repeat.  
 Repetendus, to be repeated.  
 Respondeo, responderit, shall have answered. Donec alvus ad sedes ij vel iij responderit, until two or three stools shall have been procured.  
 Ribes, currants. Gelatina ribesiorum, currant jelly.  
 Ruber, rubra, rubrum, red.



Rubus, idæus, raspberry.  
Retinendus, to be retained, or kept.

## S.

S.V.R., Spiritus vinosus rectificatus, rectified spirit of wine.  
S.V., Ten, or tenuis, proof spirit.  
Sæpe, often; sæpius, oftener; sæpissime, very often.  
Saltem, at least.  
Sanguis, blood—in, of blood. Sanguinis missura, blood-letting.  
Saphena vena, the ankle vein.  
Scapula, the shoulder blade. Inter scapulas, between the shoulders.  
Scilicet, to wit, namely.  
Scrobiculus cordis, the pit of the stomach.  
Secundus, a, um (adjective), second.  
Secundum (preposition), according to.  
S.A., Secundum artem, according to art; that is, you are to use your own ingenuity to do it in the most proper and scientific way.  
Sed, but.  
Sedes, a stool—plural, stools.  
Semi, or semis, half. Semihora, half an hour. Semi drachma, half a drachm.  
Sesqui, one and a half, as sesquihora, an hour and a half. Sesquiuncia, or sescuncia, an ounce and a half. Sesquidrachma, a drachm and a half. Remember well to attend to the difference of these two words, for many young men, by not knowing that sesqui means one and a half, but confounding it with semi, have made bad mistakes.  
Semper, always.  
Semel, once.  
Septem, seven.  
Septimana, a week, seven days. Septimus, seventh.  
Sequens, following.  
Serum lactis, whey. In sero lactis vinoso, in wine whey.  
Serum is also the watery part of the blood which separates from the red part, or crassamentum, on standing until cold.  
Sex, six; sextus, sixth.  
Si, if. Sive, or; whether.  
Signatura, a label or direction.  
Signetur, let it be marked, directed, written upon.  
S.N.P., Signetur nomine proprio, mark it with the name it is usually known by.  
Simul, together; as, terantur simul, let them be rubbed together. Simul ac, at the same time that.  
Sine, without. Sine morâ, without delay.  
Singultus, hiccup.  
Singulus, a, um, each; in singulis, in each; singulorum, of each.  
Sinister, tra, trum, the left. Auri sinistro, to the left ear.  
Satis, thirst; si sitiât, if thirsty.  
Solitus, accustomed.  
Solutus, alone; only.  
Solutus, a, um, dissolved, also loosened; as, donec alvus soluta fuerit, until a stool is procured.  
Solve, dissolve.  
Somnus, sleep. Hora somni, at bed-time.  
Spina, the spine, the backbone; also, a thorn.  
Spissus, thick. Spissitudo, thickness of consistence.  
Statim, directly, immediately.  
Stent, let them stand. Stet, let it stand.  
Sternutatorius pulvis, sneezing powder, snuff.  
Stupa, tow.  
Sub, subter, under. Sub finem coctionis, when the boiling is almost finished. Sub,



prefixed to a word, implies diminution, or a process not completed; also, in many words has the same signification as the termination *ish* in English words, as *sub-niger*, blackish, not quite black. *Subtepidus*, lukewarm. For its meaning when applied to terms of chemistry, as *subcarbonas*, *submurias*, the pupil must study his chemical nomenclature.

*Subactus*, a, um, subdued, dissolved.

*Subitus*, a, um, *Subitaneus*, sudden. *Subito* (adv.), suddenly.

*Subtepidus*, a little warm, lukewarm.

*Subige*, dissolve it, make it unite. *Subdue* quicksilver with lard or balsam of sulphur.

*Subinde*, frequently, now and then.

*Subtilis*, subtile, reduced to fine powder. *Pulv. subtilissimus*, the very finest powder.

*Sumat*, let him take. *Sumatur*, *sumantur*, let it be taken, take.

*Sudor*, sweat.

*Superbibo*, to drink after taking anything; as, chamomile tea or warm water after an emetic; or a cup of water or any liquid medicine to wash down a dose of any sort of pills.

*Superinfundo*, to pour upon.

*Supra*, above. *Supradictus*, above mentioned.

*Syncope*, a fainting fit.

## T.

*Tabellæ*, *tabulæ*, lozenges.

*Talis*, such. *Sumat talem*, let him take such a one as this.

*Talus*, the ankle.

*Tam*, so. *Tamen*, yet.

*Tactus*, the touch.

*Tænia*, the tapeworm.

*Tempus*, *temporis*, time.

*Tempora*, the temples. *Temporibus*, to the temples. *Tempori dextro*, to the right temple. *Tempori sinistro*, to the left temple.

*Temperies*, *temperamentum*, temperament, degree of heat.

*Tenacitas*, tenacity. *Ad debitam tenacitatem*, of a proper degree of tenacity or consistence.

*Teneo*, to hold. *Tenendus*, to be held.

*Tenuis*, weak, thin, small, slender.

*Tepefactus*, warmed, made warm.

*Ter*, three times, thrice. *Ter quaterve*, three or four times.

*Teres*, *teretis*, round, taper; also, *teres* is a name for the long and round worm infesting the human body, *qu. d. vermis teres*.

*Tergum*, the back. *A tergo*, behind.

*Tero*, to rub. *Tere* (imperative), rub. *Terendus*, to be rubbed. *Terantur*, let them be rubbed.

*Tertius*, *tertia*, *tertium*, the third.

*Testacea*, the prepared powders made of oyster-shells, egg-shells, crabs' claws, &c.

*Thorax*, the chest. *Thoracis*, of the chest.

*Thus*, frankincense.

*Tinea capitis*, scald head.

*Torrefactus*, toasted.

*Tres*, *tria*, three; *tribus*, in three, to three.

*Triduum*, three days.

*Trituratio*, a grinding. *Trituratus*, triturated, ground. *Tritus*, ground.

*T.O.*, *Tinctura Opii*, what is commonly called Laudanum.

*T.O.C.*, *Tinctura Opii Camphorata*, Camphorated Tincture of Opium, called, formerly, Paregoric Elixir. This tincture is now called by the London College, in the late reform of the Pharmacopœia, edit. 1809, *Tinctura Camphoræ Composita*.

*Trans*, through. *Cola trans chartam*, filter through filtering-paper.

*Tussis*, a cough. *Tussi molestante*, when the cough is troublesome.



Ultimus, ultima, um, the last. Ultimó præscript, which was last prescribed.  
 Umbilicus, the navel.  
 Unà (an adverb), together.  
 Uncia, an ounce. Unciam cum semisse, an ounce and a half.  
 Undecim, eleven.  
 Unus, una, unum, one. Unius, of one. Uni, to one.  
 Urgente tussi, when the cough is troublesome.  
 Urgeo, to urge, to oppress, to be troublesome or painful.  
 Usus, use. Pergat in usu remediorum, continue the use of the medicine as before.  
 Usque ad, up to, as far as.  
 Ut, as, that, so that; in the same manner as.  
 Utendus, to be used.  
 Utatur, let him make use of.  
 Uterque, utraque, utrumque, both. Utriusque, of both. Utrique, to both, to either.  
 Admoveantur hirundines ij tempori utrique, apply two leeches to each temple.  
 Utriuslibet, of whichever of the two, or more, the patient likes best.

## V.

Vaccinatio, the act of inoculating for the cowpock.  
 Variolæ vaccinae, the cowpock.  
 Vaccinum lac, cow's milk.  
 Valde, very, very much.  
 Valeo, to avail; si non valeat, if it does not answer.  
 Variolæ, the smallpox.  
 Varicella, the chickenpox.  
 Vas, a vessel. Vasis, of a vessel. Vase clauso, in a covered vessel. Agitato vase, shaking the vessel.  
 Ve, vel, or; either: -ve is never at the beginning, but the end, of a word.  
 Vertebrae, the joints of the neck, back, or loins; the vertebrae, altogether, form that column of bones which is called the spine.  
 Verus, true, real, genuine.  
 Vena, vein. Vena saphena, the ankle vein.  
 Venæsectio, bleeding. Fiat venæsectio, bleed him.  
 V. S. B., Venæsectio brachii, bleeding in the arm. Fiat venæsectio in venâ saphena, bleed the patient in the ankle vein—or, it may be understood, to bleed wherever you can find the best vein at the top of the foot to get blood from.  
 Vespere, in the evening.  
 Vespertinus, in the evening, as horâ vi<sup>a</sup>, vespertina, at six o'clock in the evening.  
 Vehiculum, a vehicle; that is, whatever liquid or any other eatable or drinkable we take a medicine in, as barley-water, whey, jelly, or panada, &c.  
 Viginti, twenty. Vicesimus, vigesimus, the twentieth.  
 Vinosus, vinarius, of wine. Cyathus vinarius, a wine-glass.  
 Vices, turns; ad duas vices sumendus, to be taken at twice; that is, half to be taken at first, and the other half in some time after.  
 Vicibus partitis. See Partitus in P.  
 Vice, in the room of. Vix, scarcely, hardly. Ut vix sentiatur, so that it can scarcely be perceivable.  
 Vitellus, the yolk of an egg.  
 V. O. S., Vitello ovi Solut<sup>m</sup>; dissolved in the yolk of an egg.  
 Vitrum, a glass. Vitreus, made of glass.  
 Vires, strength; si vires permittant, if the strength will bear it.  
 Vomitio, a vomiting. Vomitione urgente, when the vomiting is troublesome.  
 Vultus, the countenance.



## OBSERVATIONS ON THE MANAGEMENT OF THE SICK ROOM.

In the treatment of disease, as much depends on the proper management of the patient, or as it is usually called *nursing*, as on the remedial powers of medicines appropriate to the exigencies of the case. How many, in fact, have owed their lives to the sedulous and skillful attention of their nurse or friends, wholly independent of professional assistance! and, on the other hand, what numbers have been sacrificed to a want of knowledge of the proper government of a sick room, in spite of the most judicious treatment on the part of the physician! This most important subject has been strangely overlooked by both lecturers and writers on the practice of medicine. We are acquainted with but two treatises on the subject, in our language, that can be referred to with confidence—the general observations prefixed to “A Practice of Physic,” by the late Dr. Dewees, and “The Domestic Management of the Sick Room,” by Dr. A. T. Thomson, of both of which we shall freely avail ourself in the following pages.

## VENTILATION IN THE SICK ROOM.

Where it is possible, the room in which the patient is confined should be large and lofty, and in all cases provided with a chimney; the upper sashes of the windows should be capable of being let down, which is not always the case; in a word, the room should be of such a construction, as to permit the freest ventilation and renewal of the air without injury to the patient. Fortunately, for the comfort as well as for the benefit of the sick, a great change has taken place as regards their treatment in this respect; at one time, and that not many years since, it was considered essential to their recovery that air should be carefully excluded from their apartments, and every means was taken to accomplish this result; but a more rational mode of treatment is now generally adopted, though physicians occasionally meet with individuals who obstinately adhere to the idea that all access of the outer air is injurious to the sick.

The air of a sick chamber soon becomes impure, and must be changed by the admission of fresh air from without, and not, as is too frequently attempted, by the use of disinfecting agents in the chamber itself; which, although they may mask or destroy offensive odors, only tend to vitiate the air still more. It may be confidently asserted, that no disinfecting or fumigating agent is capable of rendering the air of a sick chamber so pure and salutary as it can be made by proper attention to cleanliness and free ventilation.

This latter can in most cases be accomplished without risk to the patient, by proper management, and should never be neglected. It should be regulated by the season of the year, the state of the atmosphere, and the character of the disease. In winter, in consequence of the draught occasioned by the fire, a renewal of the air of the room takes place to a greater or less degree; but a sick chamber should, if possible, be heated by an open fire, and not by a stove, as the air is thus more rapidly changed, and the unpleasant emanations caused by a large surface of heated iron is obviated. The state of the weather should also influence the mode of ventilation, for it must be obvious that, if it is damp, it would be injurious, if admitted into a sick room directly or in large quantities. The character of the disease should always be considered; as a general rule, all acute affections require more ventilation than those of a chronic nature, except such as are attended with offensive discharges. No precise rules, however, can be laid down on the subject of ventilation as respects the mode in which it is to be attained; this must, in a great measure, be left to the discretion and good sense of the attendants.

Ventilation is especially required in infectious and contagious disorders, as, in close, ill-ventilated apartments, the power of infection is greatly augmented, and is, as it were, concentrated. The infectious miasm is greatly weakened or diluted by a free access of air, and its powers much augmented by a damp, close atmosphere. Next to ventilation, and of equal importance, is the



## TEMPERATURE OF THE SICK ROOM.

There is a very general, but erroneous, opinion among nurses, that a sick person should be kept very warm, to prevent his taking cold, and, consequently, the temperature of the chamber is maintained at a fever heat, to the great inconvenience and positive detriment of the patient. The best general temperature of a sick room is from 60° to 70° F., to be regulated by the thermometer rather than by the sensations of the sick person, as these are oftentimes morbid and deceptive, but at the same time are not to be wholly disregarded; thus, where the temperature is agreeable to the sensations of the attendants, and the patient, notwithstanding, complains of chilliness, it should be increased a few degrees.

In febrile complaints, nothing conduces more to the comfort, and nothing is more proper than a reduced temperature, as it materially aids the physician in relieving the morbid heat of the patient's body, and the increased action of the arterial system. Thus, in a patient in the hot stage of fever, it is surprising to see the almost immediate relief experienced by the admission of cool air into the chamber; a tranquil state ensues, replacing his former jactitation and restlessness; his skin becomes cooler; his respiration, from being hurried and laborious, becomes calm and gentle; his pulse less frequent and active; and, in many cases, a placid and refreshing sleep comes on; or he breaks out into a general perspiration.

Where, on the other hand, the temperature of the room is maintained at a temperature equal to, or nearly equal to that of his body, every morbid symptom will be much aggravated, and his condition rendered still more deplorable. But, whilst a moderately depressed temperature is beneficial in many diseases, it exercises a decidedly injurious effect in others; in affections of the lungs, even a slight change in the thermometer will often cause an attack of cough and an augmentation of the symptoms.

During convalescence, the air of a sick chamber should be frequently renewed, and the temperature kept at about 60° to 65°; but no sudden transitions permitted, as nothing is more essential than a guarded care against extremes and rapid changes of heat and cold.

It is of equal importance that a proper attention should be paid to the temperature of the patient as regards the amount of his covering, for constant errors are committed by nurses in this respect. When a sick person complains of feeling cold, it is always proper to provide him with additional covering: but, if reaction takes place, and heat of skin ensues, it is seldom thought advisable by the attendants to remove any of the now superfluous clothes, "as he might take cold;" the consequence is that a febrile condition is produced, with a dry, hot skin, without a sign of perspiration. If some of the bedclothes be removed, however, this will soon make its appearance to the great relief of the patient. Every nurse should be aware that there is a sweating temperature, and that, when this is transcended, perspiration will not take place, and will cease if it be present. Attention to a simple rule on this point will obviate any difficulty,—to add covering when the patient complains of chilliness, and to remove it when he experiences too much heat.

## CLEANLINESS IN THE SICK ROOM.

Nor is attention to cleanliness of less importance, for, although much is accomplished, in preventing a foul and offensive atmosphere, by proper ventilation, this is not sufficient, and is not always practicable. The observance of this is one of a nurse's most important duties; and she should be careful to remove anything that is capable of emitting an unpleasant smell, or of giving out an injurious gas, as speedily as possible. All evacuations are to be taken from the room at once, and never suffered to taint the air, under any pretence whatever. The bedclothes, as well as the personal clothing of the patient, should be changed as frequently as circumstances will allow, and no dirt suffered to accumulate in any part of the room. All articles used by the patient in taking his food, medicine, &c., should be cleaned as soon after they have been employed as possible. No slop-basin or slop-pail should ever be allowed to remain in a sick chamber.



A patient's face and hands should be frequently wiped with a towel moistened with cold water, or vinegar and water, in all cases in which there is no danger of causing chilliness, and he should be allowed to rinse his mouth and clean his teeth, several times a-day; when he is unable to perform the latter office for himself, it should be done by the nurse. This is very grateful to patients in fever, when the tongue is dry or coated with fur or tenacious mucus.

Whilst every means of ensuring cleanliness in the sick room should be practiced, it is to be accomplished with as little noise and bustle as possible; otherwise it becomes annoying and injurious to the patient. It is not requisite that the room should be swept, nor that the furniture should be dusted, every day. What is done should always be performed in the morning, as the patient is usually better able to bear the little noise and bustle necessarily attendant on these operations, after a night's rest and quiet. When the carpet is swept, it should be sprinkled with moist tea leaves a short time previous, to prevent dust from rising.

#### QUIET IN THE SICK ROOM.

In most diseases, and especially in those attended with fever, nothing is more annoying and distressing to the patient than noise; hence, the utmost quiet should be strictly maintained. All unnecessary conversation should be avoided, as a sick room is an unfit place for gossiping. If conversation be carried on, however, it should be in such a tone of voice that the patient may comprehend it, for nothing is more injudicious or hurtful to a sick person, especially if of a nervous temperament, than the mode usually adopted by the attendants of conversing in whispers, as it leads him to imagine that it refers to something they are afraid or unwilling to communicate to him, and hence, he fatigues himself by endeavoring to ascertain the purport of conversation, or gives way to despondency; on this account, it is better that all talking that is required should be carried on in a moderate tone of voice, and not in whispers, as much less likely to disturb the patient or to excite his attention.

In some cases, especially in those of a nervous character, on the contrary, cheerful conversation is of much benefit to the patient, provided it is not continued so long as to fatigue him.

A great fault in many nurses is being always in a bustle, "putting things to rights," or "fixing the room," thus constantly disturbing the patient, and preventing him from sleeping; nothing is more annoying to a sick person than this bustle or this constant recurrence of noises, which, however insignificant in themselves, render him nervous and impatient, from their repetition. Most of these may be prevented by an attention to order and method on the part of the nurse.

One common source of annoyance to a sick person arises from a frequent opening and shutting of his door, more especially if the lock or hinges are not in good order. In many cases, the door may be kept permanently open, and the patient protected from the light and draught of air by means of a screen. Where this cannot be done, the door should never be opened, except where it is absolutely required, and it is by all means to be avoided whilst the patient is asleep. One means of avoiding a too frequent opening and shutting of the door is to permit no person to enter the room except such as are required to assist in attendance on the patient.

#### EXAMINATION AND PRESERVATION OF THE EXCRETIONS.

This is of much importance in many cases, as it enables the physician to judge with some degree of certainty of the condition of the patient and of the progress of the disease. It should never be trusted to a nurse, as it is very rarely that their account can be depended upon. In all cases where the physician deems it necessary, the nurse should be directed to preserve the excretions for his inspection; but they should never be retained in the sick room.

#### ADMINISTRATION OF MEDICINE, &c.

However skillful the physician may be, and however judicious his treatment, they are rendered perfectly nugatory, if the remedies he orders are not administered accord-



ing to his directions, as the cure of the patient depends in many cases on the regular application of the prescribed remedies. It is, therefore, the duty of the attendant on the sick to follow implicitly the directions of the physician, as well in exactly complying with his orders, as in doing nothing that she has not been ordered to do. At the same time, there are exceptions to this rule, in which a suspension of the remedy, or a deviation from the order of the physician, is not only allowable, but is absolutely required. Thus, from idiosyncrasy or some other cause, the remedy in the doses ordered may have no effect, or may produce one widely different from that intended or expected. In such cases, it is evident that a strict adherence to the directions of the physician would be productive of evil; but he should be immediately apprised of the circumstance.

Nurses, however, more frequently commit errors in the administration of drinks and nourishment than in that of the prescribed medicines. In the former, they are apt to imagine that they are as good, if not better judges than the physician, and, in consequence, thwart the best directed treatment. The nurse should as scrupulously adhere to the directions of the physician as regards drink and nourishment, as with the doses and times of administration of medicines; nor should the exhibition of food or drink be left to her discretion more than that of remedies.

One error is very common, that of giving them too largely or too frequently; this should be guarded against, as far as possible, by the physician prescribing the quantity and quality of food or drink, as well as the times of giving it, with as much precision as he prescribes doses of medicine and the periods of their exhibition. In fact, in the treatment of the sick, as little as possible should be left to the discretion and judgment of the nurse or attendant of the patient.

#### FURNITURE OF A SICK ROOM.

Where a disease is of a serious character, or appears likely to be protracted, it becomes of importance that the chamber he is to occupy, if circumstances will admit of it, should be selected and arranged in such a manner as will most minister to his comfort, and aid his recovery. It should, if possible, be large and airy, with a northern aspect, so as to avoid the glare of the sun; but, in towns, a room exposed to the sun is preferable to one on the street, in which the patient is constantly disturbed by the noise of passing vehicles, &c. As mentioned under the head of ventilation, it should always be provided with an open chimney, that a due circulation of air may take place.

No article of furniture that is not required should be suffered to remain in the room, as the dusting and arranging it will cause unnecessary noise and bustle, annoying to the patient. Two tables are wanted in most cases: one of a small size, to stand by the side of the bed, to hold such articles as are in immediate use, as the medicines he is taking, the spoon or glass in which they are administered, &c.; but nothing should be permitted to remain upon it, except articles frequently wanted. The other table, which should be large, and adapted for the reception of medicines not in immediate use, such as spare spoons and glasses, should contain a drawer, supplied with a roll of old and soft linen, a sponge, scissors, a spatula, a roll of muslin and one of flannel bandage, some lint, some adhesive plaster, a pincushion well supplied with pins; in fact, all such articles as may be needed. It may appear unnecessary to allude to these things; but it has happened to every medical man, to observe that, without such previous preparation, much of his time is wasted in the search for what is wanted.

Where a second table is inconvenient, the top of a chest of drawers can be employed instead, for bottles, glasses, &c.; and one drawer should be appropriated to the reception of the various articles above enumerated; another should contain towels, a free supply of which should always be at hand; but none are ever to be used for the reception of dirtied articles of clothing, which should always be at once removed from the room. The washstand should be provided with additional basins, one of which should be of small size; and a large pitcher of water should be always kept in the room.

If there is a looking-glass in the room, it should be placed in such a position as not to be in view of the patient. A large sofa, or a small additional bed, is constantly required, as in certain cases where the patient is unable to sit up, he finds much comfort



in being removed to it; it is also required when his bed is to be made. An invalid, or easy chair is also productive of much benefit in certain cases; and where it can be commanded, it should be of such a construction as will admit of changes, so as to vary the position of the patient at pleasure.

What is called a French bedstead, without curtains, is the best adapted to the sick room. The four post bedstead, when used, should not be furnished with curtains, as these tend to exclude the air, and to retain smells and a vitiated atmosphere. In all cases, a mattress is preferable to a feather-bed; it may be placed over the latter, when greater softness is required. The pillows should be elastic, but so firm as not to permit the head of the patient to sink in them, as this prevents coolness and a due circulation of air. The bed-coverings should be as light as is consistent with warmth and comfort; hence, Marseilles quilts should not be used, as from their weight they oppress the patient, without affording as much warmth as a light blanket.

The change of bedclothes, and shaking up the bed or mattress, must be regulated by the character of the disease and the condition of the patient. In febrile complaints, it should be done, if the strength of the patient will admit, twice in the twenty-four hours. When there is much restlessness, it will be found advantageous to move him night and morning to a large sofa, or to another bed; this tends to promote sleep, and to insure a proper airing of the beds. When he is thus moved, the clothes of the bed he has left are to be turned down, and fully exposed to the air, which will render so frequent a change of the sheets less necessary than would otherwise be required. When there is only one bed, and in febrile diseases, the sheets which have been used at night are to be replaced by others in the morning, and freely exposed to the air during the day, and again used at night. When the patient is too much debilitated to be moved from bed to bed, he should be merely removed to the other side of his bed, and the one he has left properly arranged by changing the linen, beating up the bed, &c.; but these changes are never to be made whilst he is in a perspiration.

If a fire in the room is required, it should not be made use of to prepare the food or drinks of the patient; this should be done elsewhere, as well to avoid the smell emitted by the heated articles, as the noise and bustle caused in their preparation.

#### PROPER USE OF UTENSILS FOR EVACUATIONS, &c.

As it is of the utmost importance to the patient that he should be spared all unnecessary fatigue, some precautions are to be used to prevent this during his evacuations; and one of the duties of a nurse is to enable him to perform those offices with the least expenditure of strength. When the patient is feeble, he should never be allowed to rise to fulfil the calls of nature, and therefore a bed pan and urinal should always be provided; by means of these, the evacuations can be performed with ease, and without exertion. Many persons, it is true, have an aversion to using them; but this difficulty can generally be overcome by a proper representation of their advantages. One of the objections commonly made is that the use of a bed-pan gives them pain in the back; and this is often the case, where the patient is not properly supported; but can always be prevented by placing a pillow in the hollow of the back, before the pan is used.

In the case of giving drinks or nourishment, much unnecessary fatigue may be spared to the patient. Owing to the form of the vessels usually employed to administer drinks, without the sick person is raised in bed to receive them, much of the fluid is apt to be spilt on the bedclothes, to his great discomfort. Hence, a tumbler, tea-cup, or bowl should never be employed for this purpose, where the patient is too feeble to raise himself without inconvenience. The sick cup, as it is called, should therefore be employed for this purpose, as it spares all fatigue, and prevents any spilling of fluid on the patient or on the clothes.

When the medicine or food is taken in spoonful quantities, and in cases of nauseous medicines to be administered to children, the medicinal spoon will be found very useful. This is a spoon with a hollow handle, with an opening at its extremity; the bowl is provided with a hinged lid, but is open at the apex. When a fluid is poured into it, and the lid closed, the pressure of the air at the opening at the end of the bowl will prevent any of the fluid escaping, as long as the orifice at the end of the



handle is closed, but is projected with some force when this opening is free. By means of this contrivance, medicine or food may be conveniently administered without the patient being obliged to be raised in bed, or in spite of resistance on the part of a child.

### DOSE OF MEDICINES.

The doses of medicinal substances, being specific as regards each other, can only be learnt by experience; but their remedial or active powers do not always increase with an increase of the dose: thus, if a purgative dose of calomel be ten grains, an increased quantity of the medicine will not induce an equivalent increase in the purgation, though it may cause other consequences. The dose in which a medicine is given often determines its specific action; two grains of ipecacuanha will usually quiet the stomach, or act as a diaphoretic, whilst thirty will produce active emesis, and, in like manner, almost every medicine displays different powers, according to the dose in which it is administered.

Dr. Paris states "that powerful doses are disposed to produce local rather than general effects." This is proved by experience, in many cases, but does not hold good in all; thus, many of the active poisons cause general effects in both large and small doses, and their virulent action is in general proportionate to the quantity taken. In most of what are termed the alterative medicines, however, the alterative action is only displayed when they are administered in small, but long-continued doses, and does not take place when they are taken in such quantities as to excite much local action. It is erroneous to suppose that a medicine is devoid of power, or does not exert an influence on the system, because no marked effect immediately follows its exhibition; for, as is observed by Dr. Barlow, "it is very possible that practitioners often err, especially in the treatment of chronic maladies, from requiring an obvious effect from each dose administered; where it is ascertained that a medicine actually possesses inherent powers, the slow and almost imperceptible exercise of these powers should not be despised. There is often more wisdom in seconding the efforts of nature than in superseding them." In fully admitting the truth of this remark, we are far from advocating the doctrine of Hahnemann, or his invisible doses; nor does it, in fact, accord with his views, for he gives a long catalogue of symptoms, caused by the administration of the minutest quantity of the articles of his *materia medica*.

So many circumstances influence the effects of medicines, that it is almost impossible to say what dose will produce identical effects in different individuals; but it has been sufficiently ascertained that, in the generality of patients, we may reasonably expect certain results from certain doses. The circumstances that influence the action of medicines, and the doses in which they are to be given, are

AGE.—This exercises much influence; the young require smaller doses of a medicine to produce the desired effect than an adult; and the aged, although less susceptible to impressions than formerly, are unable to bear any forcible shock on their system. Various formulas or tables have been published for the graduation of doses to the respective ages; but none of them are as generally adopted as that attributed to Gaubius, which is as follows:—

Dose for a person in adult age being	1 drachm or 1,
That for a person from 14 to 21 years would be	2 scruples or $\frac{2}{3}$ .
“ 7 to 14 “	$\frac{1}{2}$ drachm or $\frac{1}{2}$ .
“ 4 to 7 “	1 scruple or $\frac{1}{3}$ .
“ 4 “	15 grains or $\frac{1}{4}$ .
“ 3 “	10 grains or $\frac{1}{6}$ .
“ 2 “	8 grains or $\frac{1}{8}$ .
“ 1 “	5 grains or $\frac{1}{12}$ .

This table, although mainly correct, is often found to be erroneous, when applied to particular cases. Dr. Paris, in his *Pharmacologia*, gives a formula devised by Dr. Young, which is simple, and has been found to be generally correct; this is:—

For children under twelve years, the doses of most medicines must be diminished in the proportion of the age to the age increased by twelve; thus:



At two years to  $\frac{1}{7}$ , viz :

$$\frac{2}{2+12} = \frac{1}{7};$$

At 21 the full dose to be given.

But no scheme can be devised, founded on age alone, to which there are not many exceptions. Thus, children require proportionably larger doses of castor oil or calomel to affect them than adults, whilst, and especially when very young, they are exceedingly susceptible to the action of opium and its preparations.

**SEX.**—It is generally admitted that females require smaller doses of medicines than males, and, as a general rule, this may be correct; but so much depends upon habits of life, temperament, &c., that the exceptions are almost as numerous as the examples.

**TEMPERAMENT** also exercises much influence on the doses of certain articles of the materia medica; an individual of a nervous temperament is unable to bear with impunity doses of opium that scarcely act on the phlegmatic; the sanguineous are readily affected by the action of stimuli, &c.

**IDIOSYNCRASY**, or constitutional peculiarity, has more influence than either sex or temperament on the action of medicines. Some of these anomalies are very striking, and almost incredible, and evidence the necessity of particular inquiries being made by the physician of every patient with whose constitution he is unacquainted. In some cases, opium, or any of its preparations, cannot be administered without producing the most distressing symptoms; in others, the smallest doses of calomel will cause salivation; whilst in others again it may be profusely given without inducing any effect on the salivary glands. In some persons, the mere vicinity to the poison oak (*Rhus radicans*) will bring on an unpleasant inflammation of the skin; whilst on other individuals this plant has no action whatever. It is often found that copaiba will cause a peculiar eruption, and the smell of ipecacuanha induce a distressing dyspnoea. So tartar emetic will be productive of ptyalism in some individuals even when applied to the surface of the body; blisters of cantharides will cause strangury in some patients, &c. In fact, these idiosyncrasies are so numerous that it would require a volume to notice in detail those already recorded.

**HABIT** exercises an equal influence with idiosyncrasy in modifying the action of medicines. The usual consequence of a long-continued use of any remedial or stimulating substance is to diminish the susceptibility of the organs on which it acts to its impression. Hence, to obtain its effects, it must be given in gradually increased quantities. In many cases, however, instead of a diminished, an increased action is caused by repetition of certain substances.

The influence of habit, in diminishing the susceptibility of the system to stimuli, is liable to fewer exceptions as regards vegetable than mineral substances, as may be daily remarked in the use of opium and tobacco. Dr. Christison accounts for this effect of habit in diminishing the influence of vegetable substances, by an increased power of the stomach in decomposing them; but it must also be attributed to a diminished susceptibility in the nervous system to their influence. But mineral medicines appear, in almost every case, to act with increased power by repetition, though the converse has been observed in some cases.

**STATE OF THE SYSTEM.**—It is well known that, in certain deviations from a healthy or normal condition, a remarkable change is often observed in the action of remedies on the system. Thus, in severe pain, opium can be administered in doses that cannot be borne with impunity in a state of health; but this is still more the case in tetanus, in which scruple doses, frequently repeated, have been administered without producing any marked effects. This modifying influence of disease has been noticed with regard to many other remedies, as in mercury, which can be given in large and repeated doses, in a febrile condition, without causing salivation. Where there is a diminished sensibility of a part, or a comatose condition, the pain, or irritation caused by the application of stimulating or vesicating applications, not being felt by the patient, there is a risk of inducing a sloughing and destruction of the part to which they are applied, if care be not taken to remove them in time; so, also, the application of ammonia to the nostrils, to relieve fainting, has caused violent irritation, and even



inflammation of the trachea, the patient being unable, from his condition, of appreciating or expressing his sensations. Dr. Christison notices several fatal cases of bronchitis from this cause.

TIME OF DAY at which medicines are administered also influences their action on the system. As a general rule, no medicine should be given soon after a meal, though, in the case of some of the irritants, as arsenic, corrosive sublimate, iodine, &c., it is often found most expedient to administer them whilst the stomach is filled with food, to avoid their local action on that organ. Dr. Paris observes, under this head: "Evacuating medicines ought to be exhibited late at night, or early in the morning; thus, substances of tardy operation, as calomel, and the resinous purgatives, should be given at night, while saline purgatives, senna, &c. may be given in the morning. It would seem that during sleep the bowels are not so irritable, and, consequently, not so easily acted upon, which allows time for the full solution of the substance." In all cases, the exhibition of purgatives should be so timed that their operation may take place during the day, so as to avoid an undue disturbance of the patient at night. Other directions will be found in a subsequent part of the work.

Numerous other causes exercise much influence on the action of medicines, as climate, the imagination of the patient, the tissue or organ to which they are applied, &c.

Nor should the *variable activity* of a medicine be overlooked; this has occasioned more accidents than is generally supposed. Some remedies, and generally those of a vegetable nature, lose much of their activity by age, and, therefore, require to be given in large doses to produce the desired effect; hence, where a physician has been induced to increase the usual dose of some article, which in a recent or unaltered state is active, and even poisonous, but, from age or other deteriorating cause, has become weakened, serious accidents may occur, if he persist in administering the same doses of a parcel which is fresh, or which is obtained from another apothecary. In such cases, the safest plan is always to commence with the usual dose of the medicine, and to gradually increase it, if this be found necessary.

It should also be borne in mind that certain medicines have the power of accumulating in the system, or, in other words, of not displaying their usual effects when given in small and repeated doses, until a certain quantity has been taken, when, on a sudden, symptoms resembling those caused by an overdose of the article, make their appearance. This has been remarked in relation to mercury, arsenic, and several other of the metallic medicines, as well as iodine, digitalis, &c. In giving remedies of this character, the physician should always be cautious, and not repeat the dose of them too frequently, because no sensible effect is produced at the expected time.

INTERVALS BETWEEN DOSES.—As every medicine produces a specific and definite action on the system, which, in most cases, continues for a certain time, and then ceases, it becomes necessary to renew or maintain the medicinal impression as long as is required to counteract the morbid condition of the system. In some cases, however, as in the case of an emetic or purgative given for the purpose of merely evacuating the stomach and bowels, the single effect produced may be sufficient, and no repetition of the dose is required. In most attacks of disease, on the other hand, the physician is obliged to prescribe medicines in such quantities as to produce a certain impression, and to repeat the doses at such intervals as will keep up the influence of this impression. Where, from a want of knowledge of the time during which the remedy he has administered will continue to exercise its full action on the system, he postpones a repetition of it for too long a period, although a fresh impression is made by each dose, but little real good is the result, and, from a want of continuity in the remedial impression, no progress is made towards a cure. Hence, a physician should be acquainted with the usual duration of the action of the medicine he is administering, and order a repetition of the doses so as to keep the system under its influence. A nurse should, therefore, in all cases, follow his directions implicitly, as regards the periods at which medicines are to be given, for, by inattention to this rule, injustice is done to the physician, and, what is of more importance, the life of the patient is endangered. Where the diffusible stimulants are administered, especially those of an evanescent character, it is of much consequence that their influence should be kept up; and hence a neglect on the part of a nurse in giving



them at the prescribed periods might cause a fatal collapse of the patient. It should be observed, however, that, except in cases where it is necessary thus to maintain the vital powers of a patient, this strict observance of the prescribed intervals between the doses of medicines is not intended to extend to the night, so as to disturb the invalid, as, in most diseases, sleep is one of the best restoratives; but advantage is to be taken of those moments when he is awake, timing the doses as nearly as possible to the designated periods.

**RULES FOR THE ADMINISTRATION OF MEDICINES.**—As has been stated, fluid medicines are usually administered in drops, teaspoonfuls, tablespoonfuls, &c.—an approximative measurement, which, although not precise or accurate, is sufficiently so in most cases, though not suited to medicines of a very active nature. As the generality of medicines are nauseous, it is necessary to modify, as far as possible, their disagreeable taste, which renders them so repugnant to patients, and often causes much difficulty in their administration, especially to children; but this modification must not be of such a character as to alter their properties. As regards young children, much difficulty will be obviated by the use of the medicine-spoon already described; this will also be found useful in cases of insanity, where patients refuse to take medicine or nourishment.

It is often found that patients experience much difficulty in swallowing pills, especially when these are of small size. The usual mode is to place the pill on the tongue, and to take a mouthful of water, when, on making an effort to swallow, both pass down the throat together; this plan is generally successful; but some persons appear to be unable to take a pill in this manner, and gorge themselves with fluid without effecting the deglutition of the pilular body; in such cases, by enveloping the pill in some bread, or some conserve or sweetmeat, so as to form a bolus, the difficulty is obviated.

Mixtures containing insoluble ingredients should always be shaken up before they are administered; otherwise, the first doses will contain but little of the active ingredients, whilst the last portions will, on the other hand, be more powerful than was intended. Where the article administered is of a volatile nature, it should be given the moment it is poured out, and the bottle containing the remainder immediately closed.

As each of the classes of medicinal articles requires the observance of certain rules to insure its full benefit, these will be briefly noticed in an alphabetical order.

**ACIDS.**—These are given to fulfil various indications; some acting as tonics, some as refrigerants, &c. As those most generally used are corrosive, they require to be diluted with some bland fluid, so as not to exercise too powerful a local action on the tissues with which they come in contact; nor should they be kept in a metallic vessel, nor given by means of a metal spoon. As most of them act on the enamel of the teeth in a very rapid and energetic manner, the mouth should be well rinsed with water after the administration of each dose; or they may be taken by being sucked up by means of a small glass tube, thus avoiding any contact with the teeth: this observation applies to many of the acid or super-salts; the usual solution of the sulphate of quinine, which contains free sulphuric acid, is known to act on the teeth very rapidly, and, from a constant repetition of the doses, to injure them in a serious manner; and hence requires to be given with precaution.

**ANTACIDS** are medicines intended to neutralize an undue or morbid acidity of the stomach or intestines. Their action is chemical, and they produce an innoxious compound which takes the place of the morbid fluid that causes the uneasiness or pain. In many cases, they act most efficaciously, when an emetic or cathartic has been previously administered. To obtain a permanent benefit from them, they should be combined with tonics, aromatics, &c.; as, when given alone, the relief afforded by them is but transient.

They are most advantageously administered early in the morning, or some time previous to a meal during the day. When the salt resulting from the union of the antacid and the morbid acid is purgative, the former is better given at bed-time; thus, the combination of the solution of potassa, of the carbonates of potassa and soda, or



of magnesia, forms salts with the gastric acid that act with some energy on the bowels; but this is not the case with chalk mixture or lime water.

The patient should be restricted to light animal food and well boiled vegetables, avoiding sweets, acids, pastry, and, in fact, everything liable to increase the aced condition of the stomach.

ANTILITHICS AND LITHONTRIPTICS are medicines supposed to possess the power of obviating or dissolving urinary calculi. That there are remedial agents capable of removing or relieving that morbid condition of the system on which a lithic diathesis depends, there can be no doubt; but it is very questionable if we possess remedies capable of dissolving calculi existing in the urinary organs. Among the antilithic remedies, those most generally employed are magnesia and its salts, the alkalies, soap, several of the acids, some astringent diuretics, many bitter tonics, &c. In cases of a lithic diathesis, however, much depends on the regulation of the diet of the patient; thus, when there is a superabundant secretion of uric acid, the food should contain as little azote as possible, and, therefore, be mainly of a vegetable nature; but, at the same time, care must be taken that the healthy action of the stomach is not impaired. There should be an avoidance of all indigestible substances, as fresh bread, pastry, salt meats, and even soups of all kinds; nor should malt liquors or acid wines be allowed. Where the white sand deposits or the phosphates occur, an acid system of diet is beneficial, with abstinence from all alkalies, soda water, &c.; whilst lemonade, the acid wines, and fruits in moderation can be allowed with advantage.

ANTISPASMODICS are remedies which have been considered to be endowed with the power of allaying spasm, and of calming or entirely removing irregular actions of the nervous system. As spasmodic action of the muscular apparatus and a morbid condition of the nervous centres may arise from a variety of causes, it has been found that the most opposite modes of treatment have been required; thus, in some cases, the most efficient antispasmodic is the lancet, whilst in others the morbid symptoms are best alleviated by the most active diffusible stimuli. It is not our intention to enter into the discussion of the mooted question of the *modus operandi* of antispasmodics, or whether there exists a class of remedies which are endowed with the power of exerting a direct control over spasmodic action. It is, however, shown by experience that the influence of this class of remedial agents is very limited in their operation and very various in their effects, and that to be efficient much discrimination is requisite on the part of the physician in the selection of the article best calculated to attain the desired effect. In general, what are termed antispasmodics have some of the properties of the narcotics, but differ from them in not causing stupor or insensibility. They are mostly substances possessed of a strong odor, and of a more or less volatile nature; and hence also partake of the properties of stimulants.

ANTHELMINTICS are medicines which are capable of expelling or destroying worms situated in the intestinal canal. Several kinds of worms are apt to infest the human intestines, and often give rise to distressing and even fatal symptoms. The medicines calculated to dislodge them are of various kinds: some act mechanically, as active purgatives, powder of tin, cowhage, &c.; some by being poisonous to the worms, as pink root, pomegranate bark, the vegetable bitters, &c.; and to this class belong, in all probability, the acrid, volatile, and resinous vermifuges, as turpentine, camphor, garlic, &c.; and the mineral remedies, as calomel, the preparations of iron, sulphur, &c.; some seem to act by asphyxiating the worms, as the fatty oils, the mucilages, &c.

Vermifuge medicines should be given in as simple a form as possible, and in as large doses as is compatible with their nature, and the age and condition of the patient; and hence the plan of administering them in the form of lozenges, biscuits, &c., is erroneous. One of the most certain of these remedies is the pink root (*spigelia*), either in the form of simple infusion, or united with senna or savin; in the first case, it should be followed by the administration of a brisk cathartic. In consequence of its sometimes causing temporary loss of sight, tremors, &c., which, although seldom dangerous, are apt to frighten the relations and friends of the patient, it is better, especially in cases of children, to administer the infusion in the evening, so that its effects may not be observed. The worm-seed oil is also another native anthelmintic of great power, but has the objection of having so disagreeable an odor and taste as



to prevent its general use; the least unpleasant mode of giving it is in emulsion with mint or cinnamon water.

Whatever vermifuge is employed, and whatever success attends its administration, the treatment should not be confined to the mere expulsion of the worms; but the condition of the stomach and bowels should be attended to, and their tone rendered healthy by the use of tonics, and a well-directed and generous diet.

CATHARTICS are those medicines which increase in a greater or less degree the peristaltic action of the intestines. Cathartics have been divided into *laxatives*, which are characterized by acting mildly, without stimulating in any great degree the vessels of the intestines, or exciting a general disturbance of the system; and *purgatives*, which, in addition to evacuating the contents of the bowels, also promote secretions from their mucous coat, giving rise to copious watery stools. Some purgatives have a more violent action, occasioning nausea, faintness, pain, tenesmus, &c.; and some, acting in an overdose as acrid poisons, causing great irritation, and even inflammation of the intestines. But there is another and important character in the action of purgatives, which is that different portions of the intestinal canal are most affected by different cathartic substances; thus, calomel acts, in a marked manner, on the upper portions of the tube, as is shown by the increase of the biliary matters in the evacuations; gamboge also manifests its power on the stomach by often causing vomiting, but also, like the other drastics, causes much irritation in the large intestines; aloes, again, exert very little action on the upper part of the canal, but spends its force on the lower portion; in fact, each purgative substance is characterized by a peculiarity of action as regards the part of the intestinal canal affected by it, and the nature of the evacuation it produces; and hence, it is of importance, in prescribing an article of this class, that it be suited to the impression wished to be made, and not, as is too frequently the case, selected at random. No class of remedies has been productive of more injury than the present, from their being resorted to without judgment, and from being persevered in for too long a time; for, independent of the irritation they create and keep up in the bowels, they impair digestion, and too often lay the foundation of fatal organic diseases. Purgative medicines may be exhibited at any period during the day; but in all cases the patient should not be exposed to the influence of cold air; if, however, the circumstances are not urgent, it is better to administer them late at night or early in the morning, so as to prevent the disturbance and loss of rest of the patient, and also to secure their retention by the stomach; when given on an empty stomach, they also act more speedily and effectually than when this viscus is filled with food. To promote their action, and to obviate their griping effect, warm diluents, as chicken water, gruel, &c., should be freely taken after the first discharge. Where they operate too much, this inordinate action is to be checked by laudanum given by the mouth, or in the form of injections.

Many of these medicines are very nauseous, and so repugnant to the patient as to induce nausea and vomiting, and, therefore, require the addition of correctives. The taste of rhubarb is best disguised by being given in milk, and that of castor oil by the froth of porter. The disagreeable taste of infusion of senna is much lessened if made with cold water, or, if made in the usual manner, by the addition of strong coffee, black tea, or by a few grains of cream of tartar. Aloes are rendered more palatable by the addition of extract of liquorice. Magnesia is mixed more perfectly with water or milk, if it is poured on the surface of the fluid, and permitted to sink as it becomes saturated, than by the opposite course. The neutral salts are rendered more palatable by adding a small portion of lemon juice to their solution, &c.

When a cathartic is given in the form of pill, its operation is always less speedy than when administered in powder or mixture; and it should be borne in mind that the drastic purgatives operate more mildly, and even effectually, in combination than when given alone, as is instanced in the compound cathartic pill.

*Enemata or Clysters.*—This mode of purgation is now generally employed in our large towns; but a great prejudice exists among many persons against its use, from a fastidious and mistaken delicacy. In consequence of the improved apparatus now to be procured, the administration of these remedies is attended with much less diffi-



culty and exposure than formerly, when the pipe and bag or common syringe were employed; and, therefore, much of the objection to their use is obviated.

Clysters are of almost indispensable utility when it is necessary to evacuate the bowels as speedily as possible, or when the stomach will not bear the administration of a purgative by the mouth, and also in cases where it is of importance to make a direct application to the lower bowels, as in dysentery, colic, &c. Where it is merely wished to open the bowels, an injection of tepid water will often be found sufficient; where this is not found sufficient, the addition of table salt, sweet oil, or molasses, will, in most cases, induce a full discharge. In all cases, the patient should be directed to retain the injection for as long a time as possible, and not to attempt to empty his bowels immediately after the reception of the medicine. Though clysters seldom can be forced beyond the great curve of the colon, and hence their local action confined to the lower part of the bowels, still, in many cases, their purgative influence extends to the small intestines, and their administration is followed by full and copious evacuations. They are also directed for other purposes than evacuating the bowels, for, from the intimate connection of the rectum with the other pelvic viscera, impressions made upon it are speedily transmitted to the adjoining parts, and hence, remedies can be directed by this channel with much advantage. In irritation of the bladder or uterus, an anodyne injection into the rectum will often afford much relief. In diseases of the lower bowels, also, clysters are of almost indispensable utility, as also in the dislodgement of ascarides seated in the rectum; nor are they less beneficial in those cases of sudden sinking of the powers of life where deglutition is impossible, and yet a prompt stimulating impression is requisite to save the patient; under such circumstances, clysters of some of the diffusible stimuli have proved of the greatest benefit.

*Suppositories.*—These are substances introduced into the rectum to induce a discharge of its contents. Their use is principally confined to the mere object of relieving costiveness, for which purpose a piece of soap of a conical shape, and of about an inch in length, is the most generally employed; but in some cases, articles of a more stimulating character are necessary. In cases of costiveness in children, the introduction of the end of a small roll of paper, moistened with oil, for a few moments into the rectum, will, in many cases, be found sufficient. The objection to their use is the irritation they often create in the rectum. Sometimes suppositories of opium are found useful in those cases in which anodyne enemata are usually employed, and may be used where patients have a repugnance to the use of clysters, or where the soreness of the rectum prevents the introduction of the pipe of a syringe.

In all cases where enemata or suppositories are employed, care must be taken to introduce them with care; otherwise serious injury may be done to the mucous coat of the rectum; instances have occurred where, from a careless insertion of the pipe of a syringe, the gut has been seriously affected, giving rise to fistula, &c. In most cases, the injection for a child under one year is about an ounce, which is to be gradually increased to five or six ounces, as it advances in age; to a youth, from ten to fifteen years old, a pint may be given; whilst an adult requires a pint and a-half to a quart. Where, however, the injection is of an anodyne nature, to restrain discharges or to alleviate pain, the laudanum should be mixed with not more than half an ounce to an ounce of bland mucilage, so as not to excite contraction of the bowels by the stimulus of quantity.

DEMULCENTS OR EMOLLIENTS are medicinal agents which diminish tone or irritation in the tissues to which they are applied, and thereby cause a relaxation of the parts, or sheathe and protect them from the action of substances capable of acting on them injuriously.

The term *Emollients* is generally employed to designate remedies which are applied externally to soften and relax external surfaces, and, by contiguous sympathy, the deeper seated parts; these effects have been referred by some to a physical, and by others to a vital, agency. They consist principally of bland watery mixtures, or of simple water, at a somewhat elevated temperature, of oleaginous substances, or those combined in various forms, as fomentations, poultices, &c., which will be more particularly noticed in a subsequent part of the work.

*Demulcents* are bland substances of various kinds, administered internally, to pre-



vent the action of irritating matters on mucous membranes, or to soothe or lubricate these parts when irritated or inflamed. They are useful in irritation and inflammation of the stomach and bowels, in like conditions of the air-passages and lungs, as well as in affections of the urinary apparatus. When prescribed in cases of catarrh, and other morbid conditions of the respiratory organs, they are usually directed to be taken *ad libitum*; but they should be swallowed slowly, for their main efficacy arises from their lubricating effect on the parts with which they come in contact; and this soothing influence is extended by sympathy to the bronchial membranes and pulmonic structure. They should be taken after a fit of coughing, for, as, by that spasmodic effort, the acrid secretion of the inflamed mucous surface is thrown off, they are enabled to come in direct contact with the irritated part, and thus exercise their soothing influence more efficiently.

DIAPHORETICS are those medicinal agents which promote and increase cutaneous transpiration. These articles are numerous, and of a very heterogeneous character. In general, warm diluents, aided by external heat, are very effectual means of promoting perspiration; whilst, in some cases, this excretion is better promoted by the use of cool liquids, or even by the sudden and temporary application of cold to the surface. They are in almost every case relative agents; and, to produce the desired effect, should be given in certain states of the system, and with certain precautions. Thus, during the exhibition of these remedies, the patient should be confined to bed, and covered with light blankets; his shirt should be of flannel or muslin, as these are good non-conductors of heat. Their action is much promoted by the free use of diluents, given either warm or cold, according to the temperature of the patient's skin. During the continuance of the perspiration, great care should be taken that it is not suddenly checked from any cause. When it is wished to moderate the sweating, the patient may be removed to another bed, covered with a warm blanket, his clothes changed, and his body dried with dry towels, care being taken that he is not exposed to cold air. During the exhibition of diaphoretics, no medicines of a purgative or diuretic character should be administered, as the action of the latter is incompatible with that of the former, besides causing a frequent exposure to cold during their operation.

DILUENTS are watery fluids which increase the fluidity of the blood, and render the secretions and excretions less acrid or viscid. The best diluent is water, which is to be given either alone or variously flavored, according to the taste of the patient or the circumstances of the case. The temperature at which a diluent is given has much influence on its action. Thus, water at 60° or 70° F. merely dilutes; whilst at 45° to 60° it acts somewhat as a tonic, and at 70°, and to as high a temperature as it can be drunk, it is stimulant. Merely tepid water will often act as an emetic, whilst at 90° or 100° it will quiet the stomach and relieve nausea. In a cool or cold state, diluents diminish the temperature of the body and moderate an undue excitement of the pulse, and are hence called *refrigerants* or *temperants*, and principally consist of aqueous solutions of many of the acids, or of some of the neutral salts.

DIURETICS are medicines which increase or promote the secretion of urine: this they do by increasing the quantity of fluid taken into the stomach, or by stimulating the kidneys to increased action. They are always inconstant in their effects, and cannot be relied upon to produce the desired effect in all cases. To obtain their effect, the surface of the patient's body should be kept cool; otherwise, they are apt to operate as diaphoretics. Hence, the patient should remain out of bed, and the medicine be administered during the day-time; or, if confined to bed, the clothing should be as light as is compatible with comfort. The action of diuretics is much aided by a free use of the cooling diluents.

EMETICS are those medicines which invert the natural action of the stomach, so as to cause an ejection of its contents through the mouth, or vomiting. When the patient is plethoric, blood-letting should always precede the administration of an emetic, as it not only promotes its operation, but prevents any injurious rush of blood to the head. When it is required, a full dose of the emetic substance is to be administered; but, in the generality of cases, it is better to give it in divided doses, frequently repeated, until the desired effect is produced. It often happens, when a physician orders an emetic, that he will be asked not to give a strong one, as the patient is too weak, &c.;



but it should be known that, unless the medicine causes full vomiting, it will create more uneasiness and debility than when it acts in the desired manner. When vomiting takes place, it should be encouraged by draughts of tepid water or weak chamomile tea, to be repeated after each discharge; but the use of these fluids is not to be carried too far, because, when the stomach is gorged with liquids, the action of vomiting is often not fully performed, or is wholly checked. When the operation of an emetic is too violent, the best means of checking it is by the administration of laudanum, or the application of fomentations or warm poultices to the epigastrium; if these fail, a spice plaster or a sinapism to the same part will generally be found effectual. If the stomach continues irritable, drink, food, or medicine is to be given in very small quantities. There is always much thirst, which patients are very apt to endeavor to alleviate by freely taking some fluid; but this in most cases being rejected, the evil is only increased; the best plan to relieve it is to frequently wash the mouth with some cool drink, swallowing as little as possible. When the stomach is inactive, and a full dose of an emetic substance does not cause vomiting, it is always unsafe to repeat the dose, as by such a course inflammation may be produced. When the stomach thus resists the action of one article, carried to a due extent, it is better to have recourse to another, or to endeavor to produce emesis by titillating the fauces and throat by means of a feather. Emetics should always be given at the time ordered by the physician, and, as a general rule, on an empty stomach, in the morning; but it is sometimes of benefit to administer them in the evening, as, from the exhaustion produced by their operation, a disposition to sleep is caused.

EMMENAGOGUES are medicines which excite or promote the catamenial discharge; but there are, strictly speaking, few remedies which have this specific power on the uterus, the majority of them acting by their influence on the system generally, or on parts contiguous to the uterus—ergot being almost the only article which exercises a direct power on this organ, and that rather in increasing its expulsive energy than in promoting the menstrual function, though it has been advantageously employed for the latter purpose. To insure the successful administration of this class of remedial agents, the system must be previously prepared for their use by invigorating it, if there are relaxation and debility; and by an opposite course, if there is an undue degree of arterial action. In a majority of cases, the restoration of the discharge is rather attributable to a proper regulation of the system than to any specific power in the emmenagogue administered.

EPISPASTICS are external applications which are capable of inflaming the skin, and causing an effusion of serum beneath the cuticle. These effects can be produced by a variety of means; but that most generally employed is the Spanish fly, or cantharides. The mode of applying a blister, and of dressing the sore it occasions, is very simple; but, without special directions from the physician, many nurses and attendants on the sick are strangely ignorant on the subject. The best preparation of cantharides for blistering purposes is the cerate of Spanish flies, or common blistering plaster; this will be always found effectual, if properly prepared, and of good materials. It is best spread on soft leather, though linen, muslin, or even stout paper will answer. To obtain a prompt action from a blister, it should be spread thick; powdered flies should not be sprinkled on its surface, for, if the plaster is well made, it requires no such addition; added to which, the powder is apt to adhere to the skin, and to give the patient much uneasiness. Before applying a blister, the part on which it is to be placed should be well washed with soap and water, and thoroughly dried, the hair shaved off, and the skin irritated by being rubbed with a coarse towel, or, if this is not sufficient, with warm vinegar, spirits of turpentine, &c. The plaster is to be secured to the part by means of a bandage; or, what is better, by means of strips of adhesive plaster, or by coating the margin of the leather or cloth on which the blister is spread with the adhesive plaster. Whenever either of the last modes of securing a blister is resorted to, it is necessary, to secure its adhesion, that the skin be perfectly dry. The plaster should be in perfect contact with the skin; but should not, as is too often the case, be subjected to much pressure, for this retards vesication, and is a source of uneasiness to the patient.

The period generally required for a blister to draw is twelve hours; but this is liable



to many exceptions: thus, for children, delicate females, and for persons with tender skins, a much shorter time is sufficient. The best plan, when the patient complains of pain and burning, is to lift a corner of the plaster and examine the condition of the skin beneath, and, if it be found vesicated or fully reddened, to remove the blister; or, if the skin is not sufficiently inflamed, to permit the blister to remain till this result is attained. When the skin is sufficiently reddened, but not vesicated, the application of a warm poultice, or of a dressing of basilicon ointment, will cause a copious effusion of serum. In cases of children and irritable persons, or those liable to strangury, this should always be done. To guard against this latter, it has been advised to interpose a piece of gauze, fine muslin, or tissue paper between the skin and the plaster: these should be moistened with oil. It is very doubtful, however, whether this plan is of any real advantage.

In many persons, the action of a blister disposes to sleep; but in very sensitive individuals it causes much irritation; this may be often obviated by the addition of a few grains of extract of hyoseyamus, or of belladonna, to the plaster. When it is requisite to cause a blister as rapidly as possible, the plaster of Spanish flies must be replaced by other means. In such cases, paper, moistened with oil of cantharides, or with the ethereal extract, will be found useful. The application of a pledget, wet with water of ammonia, will also act very rapidly. When a still more rapid effect is desired, the direct application of heat to the part may be resorted to, by means of a disk of polished metal, heated in boiling water, or by applying a piece of wet blotting paper to the skin, and passing a hot flatiron over it.

Some persons have a great dread of blisters, and object to the application of a large one; but, when the full effect of blistering is required, a large one should always be applied, for the pain and irritation caused by a large blister are very little greater than a small one occasions, whilst the benefit is more decided. Still, as the quantity of the discharge is in a great measure equivalent to the size of the vesication, large blisters depress more than small ones; therefore, when they are intended merely as counter-irritants, small blisters are to be preferred. The size and shape of blisters vary greatly, according to the part to which they are to be applied, and the age of the patient. Dr. Dewees gives the following as the general size of blisters:—

For the legs or thighs, from 7 to 8 inches long by 3 to 3½ broad.

For the back, from 7 to 8 inches long by 4 to 4½ broad.

For the thorax, from 8 to 9 inches long by 7 to 8 broad.

For the stomach, from 8 to 9 inches long by 6 to 7 broad.

For the abdomen, from 10 to 11 inches long by 8 to 10 broad, if designed to cover the whole surface.

For the ears, from the peculiarity of the shape, the size cannot well be defined.

For the temples, from an inch to an inch and a half in diameter; they are usually circular.

These are the sizes for adults; for younger persons they must, of course, be proportionably smaller.

The dressing of a blister requires some skill on the part of a nurse; and much pain and inconvenience will be saved to the patient by the exercise of a little dexterity in the operation. Some trouble will be avoided by having everything necessary for the operation in readiness, as the plasters spread, bandages ready, &c. Each of the vesicles is to be carefully snipped, with the points of a pair of sharp scissors, at their most depending part, and the serum evacuated; but when the vesications are very small and numerous, this should not be attempted, as it would occasion unnecessary fatigue to the patient. In such case, the larger vesicles only are to be opened, and the smaller left till a subsequent dressing, when it is probable they will be found much increased in size. The cuticle should not be removed from the surface of the blistered part, except when it is wished to keep up a continued discharge, in which case the cuticle is best removed by the application of a hot poultice. When it is wished to heal the blistered surface, the dressing should be of simple cerate, thinly spread on a linen rag, and renewed twice a-day. When the object is to maintain the discharge for a short time, the dressing should consist of basilicon ointment. When a more permanent impression is desired, some more stimulating application, as savin ointment, must be



used. A blistered surface should seldom be washed, as this often causes chilliness and fatigue, and is generally useless. Some persons become faint when a blister is dressed for the first time: this should not excite alarm, as it is a mere sympathetic effect. The plaster for the dressings should not be larger than the blistered surface, as it causes irritation of the sound skin without aiding the discharge. Blisters sometimes, instead of healing properly, become painful and inflamed, and assume the appearance of an eroding or a phagedenic sore; this is best remedied by the application of a soft bread and milk poultice, having its surface anointed with fresh lard or cold cream. If this does not succeed, a lotion of lime-water and linseed will often be found effectual; oxide of zinc ointment will likewise prove useful in some cases. When a blistered surface causes much itching, and renders the patient restless, washing the irritated surface with an infusion of flaxseed or slippery elm will, in most cases, relieve the uneasiness. One of the most distressing consequences, arising from the application of a blister, is the production of strangury, which invariably takes place in some persons whenever a blister is used; this is remedied by a free exhibition of emollient and mucilaginous diluents, opiates by the mouth and as an enema, external fomentations to the pubic region, &c.; in some cases, a clyster of solution of sulphate of soda will act very efficaciously. To prevent its occurrence, it has been found of benefit to incorporate camphor and opium with the epispastic ointment, say twelve grains of the former and four of the latter, for a blister six inches square. It is said that boiling the flies in water, previous to their formation into an ointment, will deprive them of the property of causing strangury.

ERRHINES are medicines which, snuffed up the nostrils, cause an increased discharge of the nasal mucus, and generally the convulsive action of sneezing. They are now seldom prescribed, though undoubtedly useful in some cases; but they should never be employed in persons disposed to apoplexy, or in those affected with hernia, prolapsus uteri, &c.

ESCHAROTICS are topical agents, which, by a chemical or mechanical action, are capable of eroding or disorganizing the solid tissues of the part to which they are applied: those which act mechanically are actual cauteries, as a heated iron, moxa, &c.; those which act chemically are caustics, as fused potassa, nitrate of silver, and chloride of zinc. In cauterizing with a heated iron, this should be at a white heat, as, at this temperature, it occasions less pain to the patient, from its causing an immediate death of the part to which it is applied. In applying it to any part, the surrounding surface should be protected by some non-conductor of heat, but not by wet paper or cloth, as the sudden extrication of steam will produce a blistered surface around the burn, and will much increase the pain. The hot iron is rarely employed in this country, except to arrest hemorrhage; in leech bites, where the usual means of stopping the bleeding are ineffectual, the application of the end of a heated wire to the wound will, in most cases, be found effectual. Another mode of applying the actual cautery is by means of moxa. This term is used for a small cone or cylinder of some slowly burning substance, applied to the skin, set fire to, and suffered to burn slowly for a length of time proportionate to the effect intended to be produced; this is a very painful operation, and seldom employed in the United States. The chemical cauterants are numerous, and of various degrees of activity. In using the most powerful of them, caustic potassa, some precautions are necessary, owing to its great deliquescence; when it is employed to form an issue, layers of adhesive plaster should be placed on the skin, with a hole of the proper size cut in their centre, through which the caustic can be applied without danger of its spreading. The chloride of zinc, which is an excellent cauterant, is generally employed by forming it into a paste with flour. The chloride of antimony, though very powerful, from being a fluid, is difficult to manage, but has been advantageously used in cases of poisoned wounds.

EXPECTORANTS are medicines which promote the expulsion of mucus or other matters from the air-passages. It is an ill-defined class of remedies, and is extremely uncertain in its action; but we cannot, in this place, examine the propriety of retaining a division of the materia medica, the articles composing which act in an obscure and by no means regular manner, in almost all cases by a remote or secondary influence of a nauseating or sedative character. At the same time, daily experience proves



that much benefit is derived, in diseases of the respiratory organs, from the administration of substances which are usually considered expectorants, in whatever way they may primarily act on the system. No class of remedies has been more productive of injury than that of so-called cough medicines; nor in any one has quackery exercised a more unbounded sway. It is too common for persons to resort to their use in cough and catarrh without consulting a physician, thus often aggravating a trifling irritation until it becomes a serious, and oftentimes a fatal, inflammation; for, as many of the articles composing them, especially those most frequently employed in domestic and empirical practice, are stimulants, much mischief is apt to follow their indiscriminate use. The following rules ought to be observed in the exhibition of expectorants—ought always to be enforced on the patient. The surface of the body should be kept moderately warm, so as to promote a gentle diaphoresis, and nothing done that may interfere with the primary operation of the remedy on the nervous, lymphatic, or muscular systems, by the secondary effect of which the lungs are benefited; hence, the use of any that excite diuresis, and more especially of purging, should be avoided; nor should the patient be allowed to indulge his appetite in food or drink of a stimulating character.

NARCOTICS are medicines which lessen the manifestation of vital phenomena dependent upon the nervous system, especially deadening sensibility and diminishing the motor power: their full operations shown in sleep, or even coma. The most general objects in view in the administration of these articles are the production of sleep, or the alleviation of pain. No medicines are so much modified in their action by idiosyncrasy and habit. This does not extend to them as a class, but to each individually; hence, it is of importance that a physician should ascertain from a patient, before prescribing a narcotic, whether he has been long or frequently in the habit of using it, or whether it causes any unpleasant effects upon him, so that some other article of the class may be substituted. As regards the time at which a narcotic should be given, it depends upon the cause for which it is to be administered. Thus, if prescribed to allay pain, it may be given at any time, and repeated as long as the pain remains; if to procure sleep, it is best given at night, and so late as to prevent the patient from being disturbed before it produces its full effect; otherwise, it is apt to exercise a stimulating action. In many persons, the secondary effect of opium is to cause much nausea and headache after its narcotic action has ceased, which are best relieved by a cup of strong coffee, or by vegetable acids. When, however, this or any other narcotic produces these consequences, its use should be relinquished, and some other substituted.

REFRIGERANTS are remedies which are supposed to diminish the heat of the body without causing a marked diminution of sensibility or nervous energy, and they are thought to do this, whether administered internally or applied externally; but it is very doubtful whether any medicines act as internal refrigerants independently of the coldness of the medium in which they are administered, except in a secondary manner. Diaphoretics act as refrigerants by inducing an increase of superficial evaporation; sedatives, by lessening the force and rapidity of the circulation; purgatives, and especially the saline, by the copious watery evacuations they produce. The great refrigerant is cold water, whether applied topically or given internally. As a topical application, it is often mixed with fluids of greater volatility, that, by a more rapid evaporation, a proportionate abstraction of heat may take place.

SEDATIVES are those remedial agents which diminish muscular tonicity, and the manifestation of nervous influence, even to the production of syncope, or a suspension, for a certain time, of the functions of the systems affected. Some act generally; others more locally. Blood-letting and tartar emetic are examples of the first; hydrocyanic acid, of the second. Most of the articles of this class are capable, if given in too large doses, or too long continued, of producing a depression of the vital powers which may induce death; and hence require to be administered with much care and judgment.

SIALAGOGUES are medicines which induce an increased secretion from the salivary glands. The modes in which the articles composing this class act are various. Some are merely local irritants, as pellitory, horseradish, &c.; and, when chewed,



produce such a local stimulation of the salivary glands as to produce a copious effusion of saliva and mucus. Another and more important section includes such articles as evince their sialagogue powers in a secondary manner, which does not take place until a powerful impression has been made on the system generally; to this belong mercury, gold, &c., and their preparations. In the first of these divisions, their medicinal powers depend wholly on sialagogue effects; hence, they are merely useful as counter-irritants; the local stimulation, and consequent increased secretion produced by their action, often acting advantageously in relieving inflammations and congestions of adjoining parts. The curative value of internal sialagogues, however, is not dependent on the salivary flow they occasion, for though, in many cases, this may be very great, yet it is not essential to their remedial action, and may, in fact, rather be considered as a proof that their full constitutional influence has been attained.

In administering mercury, with a view to the production of its constitutional effect, it should always be recollected that, if it be permitted to exert its sialagogue powers to any great extent, it always causes great distress to the patient, without producing a beneficial effect on the disease; and its operation should, therefore, be closely watched. The condition of the gums should be examined daily, and, as soon as they display evidence of the action of the remedy, it should be discontinued, or given at longer intervals. The constitution of the patient should be sedulously attended to, as mercurials are apt to produce unpleasant consequences in delicate, nervous individuals, and are badly borne by the aged, who have suffered from previous attacks of illness. It is always dangerous to push them to salivation in children, as it frequently produces sloughing of the gums and cheeks. When too great a degree of salivation has been induced by the mercurials, it is of importance to check it as speedily as possible. This is to be attempted by cool air, and by washing the mouth with an infusion of some vegetable astringent, a solution of opium, or one of acetate of lead; this latter is very efficacious, but has the property of rendering the teeth black. Marked benefit has also been derived from a weak solution of chloride of soda; much reliance is placed by some practitioners on the preparations of sulphur, especially the soluble sulphurets, both as a wash in solution, and administered as a powder. An infusion of the smooth sumach (*rhus glabrum*), as well as a strong solution of borax, has likewise been found very useful as a local application. When the inflammation is severe, the administration of cathartics, especially the saline, is requisite, and in some cases blisters and leeches. Where a tendency to slough exists, the application of diluted pyroligneous acid will be found advantageous.

STIMULANTS are remedies which increase the vital activity of the system, or of a single organ, the first being termed *general*, and the latter *local* stimulants. Much discrimination and caution are required in the administration of articles of this class, because, if given when inflammation is present, they are liable to create more mischief than benefit; but they are called for when, on the decline of that condition of an organ, or organs, a state of relaxation or torpidity exists. In this state of things, a gentle stimulation materially assists the functions, and is productive of much benefit.

TONICS are closely allied to stimulants, and some of them act in this manner; but most of them do not cause any sensible excitation of the system, whilst, at the same time, they gradually impart tone and activity to the vital powers. With the exception of iron, most tonics act primarily on the nervous system, and secondarily on the muscular, on which latter their power is most distinctly marked. Iron appears to act in a different manner, by altering or improving the condition of the blood, and consequently increasing the tone of the system. One of the most valuable properties of many articles of this class is that of curing diseases of a periodic type; in what manner, it is difficult to explain, as the laws that govern periodicity are but little understood.

In prescribing the antiperiodic tonics in fevers, as cinchona and its salts, their use should be confined to the intermissions, and, in most cases, in as large doses as the stomach will bear with impunity, so timing the doses that one may be taken a short time before the expected paroxysm. Quinia and its preparations, although eminently antiperiodic, are not possessed of the tonic properties of the bark itself, and, in the



large doses recommended of late, are apt to induce unpleasant affections of the head, as vertigo, ringing in the ears, headache, and mental delusions. In administering arsenic as antiperiodic, its effects should be carefully watched, and be discontinued when it causes constitutional symptoms, as vertigo, and œdema; nor should it be given in debilitated habits, and where the pulse is weak and feeble. It is better, in all cases, to commence its use in small doses, and to increase them until the stomach is somewhat affected, when the quantity is to be diminished, or even discontinued.

#### MANAGEMENT OF CONVALESCENCE AND RELAPSES.

The management of a patient, after the cessation of an attack of disease, and when he is declared convalescent, is of equal importance with the treatment during the continuance of the malady; and yet it is one strangely neglected by the physician, and is too often trusted to nurses and friends. In all recoveries from sickness, the organs, or parts which have been affected, remain for some time in an excitable condition, and liable to recurrence of the disease on the slightest provocation; and it is of much consequence to inquire what causes operate most frequently in thus producing a relapse. It will be found that they generally consist in the indulgence in food, either in too great quantity, or of an improper character, or in the too early or imprudent exposure to the weather, or in over-exertion of any kind.

*Diet.*—The stomach, in common with every other part of the body, suffers from a diminution of its normal vigor and power of digestion, and, as a necessary consequence, when food of too nutritious a character is taken into it, it is unable to digest it properly; hence, part of it remains in an undigested state, oppressing the weakened organ, and is either rejected by vomiting, or causes diarrhœa; or, if the stomach is capable of converting it into nourishment, an undue stimulation of that organ ensues, which will result in fever. This latter state of things is also induced by the administration of stimulating drinks. No animal substance, in any form, should be allowed during the continuance of acute febrile disease, nor immediately after its cessation. When it is permitted, it should be at first as little stimulating as possible, and in small quantities at a time; but is to be gradually increased in quantity and quality as the patient gains strength. An invalid should be very cautious in the use of certain articles, considered by nurses and others as peculiarly nourishing and proper for the sick, namely, animal jellies of all kinds. These are exceedingly indigestible, even in a state of perfect health—added to which, they are, in most cases, rendered highly stimulating by the addition of wine or spices.

*Exercise.*—Another frequent cause of a relapse is over-exertion, from the common, but erroneous, notion that a convalescent will not recover his strength, except he takes as much exercise as he can bear. Properly regulated exertion is highly serviceable; but it should never be carried so far as to produce exhaustion, and should be pursued for some time in doors, before it be attempted in the open air: the latter, at first, should always take place in a carriage, that can be opened or closed at will; the patient may then attempt short walks in the open air; but, in all cases, it is of importance that he is not unduly fatigued, as otherwise injury instead of benefit will be the result. One of the most serious errors, committed with regard to exertion, is that of permitting a convalescent to sit up too frequently, or for too long a time, under the mistaken notion of giving him strength. A patient should never be allowed to sit up longer than is agreeable to his feelings, and never so long as to produce a sense of fatigue.

*Air.*—Great care must be taken that an invalid is not exposed to cold or damp air, especially if his skin be disposed to moisture; to guard him from sudden changes of temperature, he should wear flannel next to his skin, and have his feet well protected by warm stockings. In every convalescence, from whatever disease, country air is far more invigorating than the vitiated atmosphere of towns; but care must be exercised in the selection of the location, as to its freedom from malaria, &c., for it is far preferable that he should remain within the precincts of the city, than expose himself to the influence of causes which act with peculiar activity on a system debilitated by previous disease.







# UNIVERSAL FORMULARY.

## A

### ABSINTHIUM.

#### WORMWOOD.

This name is applied to several species of *Artemisia*, all, however, possessing the same properties. That recognized in the U.S. Pharmacopœia, and most generally employed, is *A. absinthium*, a native of Europe, and commonly cultivated in our gardens.

*Ser. Syst.* Synges. super. *Nat. Syst.* Asteraceæ.  
Linn. Sp. Pl. 1188. Griffith, Med. Bot. 405.

The parts used, are the tops or extremities of the branches. These have a grayish-white color; a soft, silky feel, a fragrant odor, and a very bitter taste. They impart their properties to water or alcohol.

Wormwood has been employed as a stimulating tonic, in various forms of dyspepsia, in amenorrhœa, chronic leucorrhœa, gout, verminose affections, &c. It is given in a variety of forms.

#### INFUSION OF WORMWOOD.

R. Wormwood one ounce  
Boiling water one pint  
Infuse. Dose, one to two fl. ounces.  
*Brande.*

#### EXTRACT OF WORMWOOD.

R. Wormwood one part  
Boiling water six parts  
Boil, express, and evaporate to a proper consistence. Dose, ten to twenty grains.  
Tonic and stomachic. *Giordano.*

#### CLYSTER OF WORMWOOD.

R. Wormwood  
Rue  
Savine, each, three drachms  
Boiling water one pint  
Boil, strain, and add to each half pint  
Castor oil half an ounce  
*Foy.*

#### TINCTURE OF WORMWOOD.

R. Wormwood one part  
Alcohol eight parts  
Digest with a gentle heat for five days, express, and filter. Tonic, stomachic, and

vermifuge. Dose, from one fl. scruple to two fl. drachms. *Guibourt.*

#### ALKALINE TINCTURE OF WORMWOOD.

R. Wormwood  
Tansy  
Centaury  
Bog Bean, each, eight parts  
Alcohol one hundred and twenty parts  
Carbonate of Potassa one part  
Digest for six days, and filter. Dose, one or two fl. drachms. In dyspepsia and flatulent colic. *Hoffmann.*

#### SYRUP OF WORMWOOD.

R. Fresh wormwood at will  
Bruise, without washing, express, coagulate the juice by heat, filter, form a syrup with sugar, nine parts to five of juice.  
*Van Mons.*

#### WINE OF WORMWOOD.

R. Wormwood two parts  
White wine fifty parts  
Digest for four days, express, and filter. A stomachic bitter and anthelmintic. Dose, one fl. ounce. *Taddei.*

#### OIL OF WORMWOOD.

R. Tops of wormwood one part  
Olive oil eight parts  
Heat for two hours on a water-bath, stirring often, express, and filter. As an embrocation to the abdomen, in cardialgia, diarrhœa, &c. *Wurtem Ph.*

#### ESSENTIAL OIL OF WORMWOOD.

R. Tops of wormwood at will  
Water sufficient  
Distil, and collect the oil that floats on the product. Dose, two to four drops several times a-day. *Par. Cod.*

#### COMPOUND SPIRIT OF WORMWOOD.

R. Wormwood four pounds  
Juniper eight ounces



Cinnamon two ounces  
 Angelica root half an ounce  
 Alcohol eighteen pints  
 Macerate for fourteen days, distil twelve pints, pour back and re-distil ten pints.  
*Guibourt.*

## RATAFIA OF WORMWOOD.

R. White sugar  
 Water, each, forty ounces  
 Dissolve, and add  
 Orange-flower water six fl. ounces  
 Previously beaten with white of egg one  
 Then add  
 Compound spirit of wormwood fifty-six fl. ounces  
 Heat on a water-bath, cool, and filter.  
*Guibourt.*

## ACACIA.

## GUM ARABIC.

This gum is derived from several species of *Acacia*, as *A. vera*, *A. senegal*, *A. arabica*, &c.  
*Ser. Syst.* Polygam. monœcia. *Nat. Syst.* Fabaceæ.

Willdenow. Sp. Pl. iv. Griffith. Med. Bot. 270.

In tears or fragments of various sizes, often rounded; colorless or of a yellowish tint, semi-transparent, inodorous, of a slightly sweetish mucilaginous taste. Wholly soluble in water, forming a mucilage. Insoluble in alcohol. Demulcent and nutritive.

## MUCILAGE OF GUM ARABIC.

R. Powdered gum Arabic four ounces  
 Boiling water half a pint  
 Add the water gradually to the gum, and rub together, till the mucilage is formed. Strain. Principally used to suspend insoluble powders or to diffuse oily and resinous substances through aqueous fluids.  
*U. S. Ph.*

## MIXTURE OF GUM ARABIC.

R. Mucilag. of gum Arabic three fl. ounces  
 Sweet almonds ten drachms  
 Sugar five drachms  
 Water two pints  
 Blanch the almonds, beat them to pulp with the sugar and mucilage, add the water gradually, constantly stirring. Strain. *Ed. Ph.* Demulcent and emollient. Used to appease cough and to allay irritation.

## GUM ARABIC MIXTURE.

R. Mucilage of gum Arabic six fl. ounces  
 Oil of sweet almonds three fl. drachms  
 Syrup of saffron one fl. ounce

Solution carbonate ammonia (*Lond.*) half fl. drachm  
 Mix, and make emulsion. A tablespoonful occasionally, in catarrh. *Ainslie.*

## GUM LOZENGES.

R. Gum Arabic four ounces  
 Starch one ounce  
 Sugar one pound  
 Pulverize, mix, and form a consistent paste with sufficient rose water, to form lozenges of convenient size. A useful article to allay irritation of the throat in catarrh.

## PECTORAL PASTE OF GUM ARABIC.

R. Gum Arabic two pounds  
 Dissolve in water, and add  
 Sugar one pound and a half  
 Evaporate to a proper consistence, adding  
 Orange-flower water three drachms  
 Employed to allay irritation, like the last.  
*Foy.*

## PATE DE GUIMAUVE.

R. Gum Arabic powdered two pounds, four ounces  
 White sugar two pounds  
 Whites of eggs twenty  
 Double orange-flower water six fl. ounces  
 Dissolve the gum and sugar in a sufficiency of water, strain, next day decant, evaporate by a gentle heat to consistence of honey, add the whites of eggs and stir briskly, continue the evaporation; when almost complete add the orange-flower water, and continue to stir; pour on a marble slab, and keep in a tin box.  
*Cules.*

## GUM ARABIC PASTE.

R. Gum Arabic one pound  
 Dissolve on a water-bath, in water eight fl. ounces  
 Evaporate to the consistence of honey. Add  
 Whites of egg six  
 Orange-flower water two ounces  
 Beat well together. When sufficiently consistent, run into moulds.  
 Used for the same purposes as last. *Beral.*

## SYRUP OF GUM.

R. Gum Arabic four ounces  
 Sugar one pound  
 Boiling water one pint  
 Dissolve by means of a water-bath, let settle for some time, remove scum, and strain. A good addition to pectoral and other mixtures.  
*U. S. Ph. 1830.*

## COMPOUND SYRUP OF GUM ARABIC.

R. Syrup of gum Arabic four fl. ounces



Sulphate of morphine      one grain  
 Oil of sassafras      one drop  
 Hoffmann's anodyne      half fl. drachm  
 Mix. Known as Jackson's cough syrup,  
 and much used in catarrhs, &c. *Jackson.*

#### LINCTUS OF GUM ARABIC.

R. Gum Arabic      one part  
 Clarified honey      three parts  
 Incorporate the gum with the honey. The  
 dose is a teaspoonful occasionally, to re-  
 lieve irritation of the throat in catarrh.  
*Beral.*

#### WENDT'S MIXTURE OF GUM ARABIC.

R. Mucilage gum Arabic  
 Linden-flower water      equal parts  
 Found useful in doses of a teaspoonful  
 every hour in the diarrhoea of infants.  
*Radius.*

#### TRONCHIN'S LOZENGES OF GUM ARABIC.

R. Powdered gum Arabic      eight ounces  
 Precipitated sulphuret  
 of antimony  
 Anise      each four scruples  
 Extract of liquorice      two ounces  
 Extract of opium      twelve grains  
 White sugar      two pounds  
 Mix, and form lozenges of six grains, one  
 to be used occasionally in catarrh and  
 bronchial affections. *Tronchin.*

### ACETOSELLA.

#### WOOD SORREL.

*Ser. Syst.* Decand. Pentag. *Nat. Syst.* Oxali-  
 daceæ. Lindley.

*Oxalis acetosella.* Linn. Torrey and Gray, Fl.  
 i. 211. Griffith, Med. Bot. 208.

A native of Europe and the United States.  
 The whole plant is used. It is agreeably acid,  
 from its containing a binoxalate of potassa. It  
 is refreshing and cooling, and is said to be useful  
 in scorbutic affections.

#### EXTRACT OF WOOD SORREL.

R. Wood sorrel      one part  
 Bruise in a mortar, express the juice, and  
 pour on the marc  
 Water      three parts  
 Boil for an hour, express, evaporate, add-  
 ing the juice, to the consistence of an ex-  
 tract. Acid and bitter. Dose, a scruple  
 to half a drachm. *Pideret.*

### ACETUM.

#### VINEGAR.

Impure and very dilute acetic acid, contain-  
 ing about five per cent. of real acid. It is the

result of acetous fermentation of any fluid which  
 is susceptible of vinous fermentation. It is prin-  
 cipally obtained in Europe from malt or weak  
 wine, and in this country from cider.

It is employed internally as a refrigerant and  
 diuretic, and is a grateful addition to diluent  
 drinks in febrile affections. It is also sometimes  
 used as a clyster. Externally it is beneficial as  
 a fomentation, and likewise forms a useful addi-  
 tion to gargles and collyria. *U. S. Dis.*

#### VINEGAR MIXTURE.

R. Vinegar      three ounces  
 Honey      two ounces  
 Water      three pints

Mix. As a cooling drink in fevers.

*Ammon.*

R. Vinegar      one ounce  
 Acetic ether      one drachm  
 Syrup of raspberries      three ounces  
 Water      two pints

Mix. As a refreshing drink in fevers.

*Augustin.*

#### MIXTURE OF VINEGAR AND CARDAMOMS.

R. Vinegar      one drachm  
 Comp. tincture of car-  
 damoms  
 Simple syrup, each,      half an ounce  
 Water      ten ounces

Mix. To be taken in such portions as  
 the stomach can bear. In sick headache.

#### VINEGAR PASTILLES.

R. Powdered white sugar      four parts  
 Heat moderately, and add  
 Vinegar      one part  
 Make pastilles. *Neuman.*

#### GARGLE OF VINEGAR.

R. Vinegar      two parts  
 Honey  
 Barley water, each,      eight parts  
 Mix. As a mouth-wash or gargle in in-  
 flamed fauces. *Soubeiran.*

R. Vinegar      two fl. drachms  
 Muriate of ammonia      one drachm  
 Honey      one and a half fl. ounces  
 Water      twelve fl. ounces  
 Mix. As a gargle in inflamed fauces.

*Ratier.*

#### SYRUP OF VINEGAR.

R. Vinegar      eleven fl. ounces  
 Sugar      fourteen ounces  
 Boil together. *Ed. Ph.*  
 Mixed with water forms a pleasant and  
 refreshing drink in febrile complaints.

#### VINEGAR CATAPLASM.

R. Vinegar      one part  
 Flour      three parts



Make cataplasm. An astringent and antiseptic application.

*Beral.*

Oil of cloves, each, three drops  
Mix, and keep in a glass-stoppered bottle.  
*Gray.*

#### DISTILLED VINEGAR.

Distil a gallon of vinegar, by means of a sand-bath, from a glass retort into a glass receiver. Discontinue the process when seven pints shall have been distilled, and keep these for use.

*U. S. Ph.*

### ACIDUM ACETICUM.

#### ACETIC ACID.

R. Acetate of soda in powder one pound  
Sulphuric acid half a pound  
Red oxide of lead one drachm  
Pour the acid into a glass retort, gradually add the acetate of soda, and by means of a sand-bath, at a gentle heat, distil the acetic acid, till the residuum becomes dry. Mix the distilled liquid with the red oxide of lead and again distil, with a moderate heat, to dryness.

*U. S. Ph.*

#### DILUTED ACETIC ACID.

R. Acetic acid half a pint  
Distilled water five pints  
Mix.

*U. S. Ph.*

#### AROMATIC VINEGAR.

R. Acetic acid one pint  
Camphor two ounces  
Oil of lavender one drachm  
Oil of cinnamon twenty drops  
Oil of cloves  
Oil of rosemary, each, thirty drops  
Mix.

*Gray.*

R. Acetic acid one pint  
Oil of cloves one drachm  
Oil of rosemary two scruples  
Oil of bergamot  
Oil of cinnamon, each, half a drachm  
Oil of pimento twenty-four grains  
Oil of lavender one scruple  
Oil of neroli ten drops  
Camphor one ounce  
Alcohol half an ounce

Mix.

*Cooley.*

Aromatic vinegar is used as a pungent and reviving perfume in fainting, &c. As it is corrosive, it should not be allowed to come in contact with the skin or clothes. It is also prepared in the dry way, as follows:

R. Crystallized acetate of soda one drachm  
Sulphuric acid twenty drops  
Oil of lemons

R. Dried rosemary  
Origanum, each, one ounce  
Lavender half an ounce  
Bruised cloves half a drachm  
Acetic acid (1.068) one pint and a-half  
Macerate for seven days, strain, express, and filter.

*Edin. Ph., 1839.*

#### AROMATIC SPIRIT OF VINEGAR.

R. Strong acetic acid one pint  
Camphor two ounces  
Oil of lavender nine minims  
Oil of cinnamon twenty minims  
Oil of cloves half a drachm  
Mix. To be kept in a glass-stoppered bottle.

*Paris Cod.*

#### ACETIC ACID CATAPLASM.

R. Rye meal three parts  
Diluted acetic acid one part  
Mix. Used as an application to ill-conditioned ulcers.

*Taddei.*

#### ACETIC ACID CLYSTER.

R. Barley water six ounces  
Diluted acetic acid one to three ounces  
Mix. The proportion of diluted acid may be varied according to the indication. Has been advised in obstinate constipation, ileus, &c.

*Swediaur.*

#### PROPHYLACTIC VINEGAR.

R. Dried rosemary and sage, each, one ounce  
Dried lavender flowers half an ounce  
Bruised cloves half a drachm  
Distilled vinegar two pounds  
Macerate for seven days, press, and filter.

*Ed. Ph.*

This is intended as a substitute for the complicated formula formerly used, for *Thieves' vinegar*, at one time so celebrated as a preservative against the plague. It has no advantages over it.

#### VINEGAR OF THE FOUR THIEVES.

R. Wormwood  
Roman wormwood  
Rosemary  
Sage  
Mint  
Rue, each, one ounce and a-half  
Lavender flowers two ounces  
Garlic  
Calamus  
Cinnamon  
Cloves  
Nutmegs, each, two drachms  
Distilled vinegar eight pints



Digest by a moderate heat, in a closely stopped matrass, for twelve days, strain, express, and filter, then add one ounce of camphor dissolved in alcohol. *Lewis.*

## ACIDUM ACETICUM EMPYREUMATICUM.

### PYROLIGNEOUS ACID.

Pyroligneous acid is an impure acetic acid, obtained by a destructive distillation of wood. It contains creasote, empyreumatic oil, &c., and has been employed in gangrene to correct the fetor and to promote the separation of sloughs, and also as an application to unhealthy ulcers.

Dispens. U. S. Dunglison, N. R. 13.

### MOUTH WASH.

R. Pyroligneous acid half a fl. ounce  
Cinnamon water four fl. ounces  
Syrup of mulberries two fl. ounces  
Mix. Used as a mouth-wash in cancrum oris. It should be kept in an opaque glass vessel to prevent decomposition. *Phabus.*

### INJECTION OF PYROLIGNEOUS ACID.

R. Pyroligneous acid two fl. drachms  
Distilled water six fl. ounces  
Mix. In purulent discharges from the ear. *Buchanan.*

### CATAPLASM OF PYROLIGNEOUS ACID.

R. Bran half a pound  
Linseed meal one ounce  
Mix well, and add  
Pyroligneous acid q. s.  
To make a cataplasm.  
As an application to foul ulcers. *Reece.*

## ACIDUM ARSENIOSUM.

### WHITE ARSENIC.

Arsenious acid is found in the shops, in the form of semi-transparent or opaque lumps, or in that of a heavy white powder. It has no smell, and its taste is slightly acid. It is volatile at a red heat, giving off white fumes of a garlic-like odor. Extremely poisonous and corrosive, even in minute doses. *U. S. Dis.*

### ARSENICAL POWDER.

R. White Arsenic one part  
Calomel one hundred and ninety-nine parts  
Mix. Recommended in obstinate cases of lepra and cancerous ulcers. *Dupuytren.*

### COMPOUND POWDER OF WHITE ARSENIC.

R. White arsenic six drachms  
Powdered dragon's blood two drachms  
Animal charcoal four scruples

Cinnabar three ounces  
Mix, and triturate well. Has been used in the same description of cases. *Van Mons.*

R. White arsenic one part  
Kino eight parts  
Cinnabar sixteen parts  
Mix, and triturate well. *Ratier.*  
Employed in the French hospitals as a caustic to cancerous sores.

### ARSENICAL OINTMENT.

R. White arsenic one drachm  
Lard  
Spermaceti cerate, each, six drachms  
Melt the cerate and lard by a gentle heat, and add the arsenic; triturate well in a glass mortar till perfectly united. *Soubeyran.*

### ARSENICAL CERATE.

R. Powdered white arsenic one scruple  
Simple cerate one ounce  
Mix the arsenic with the cerate softened by heat, and triturate well. *U. S. Ph.*

### SOLUTION OF ARSENIC.

R. White arsenic one scruple  
Distilled water two fl. ounces  
Dissolve. This and the foregoing ointment have been employed as applications to cancerous sores, but are always dangerous from the poisonous symptoms induced by the absorption of the arsenic.

### ARSENICAL PILLS.

R. White arsenic two grains  
Powdered opium three grains  
White soap eight grains  
Powdered liquorice root q. s.  
Mix well and divide into twenty pills. Each of these pills contains  $\frac{1}{10}$ th of a grain of arsenic. They have been given in doses of one, two, or three times a-day, according to circumstances, in diseases of an intermittent character. *Ellis.*

### ARSENIC AND PEPPER PILLS.

R. White arsenic one grain  
Black pepper twelve grains  
Gum Arabic two grains  
Distilled water q. s.  
Triturate the arsenic and pepper for a long time, add the gum and water, and rub well together. Make twelve pills; each of which will contain  $\frac{1}{12}$ th of a grain of arsenic. *Guibourt.*

R. White arsenic fifty-five grains  
Powdered black pepper nine drachms  
Conserve of roses sufficient



Mix, and make eight hundred pills. One daily in chronic psoriasis.

*Cazenave and Schedel.*

#### PILLS OF ARSENIC AND OPIUM.

R. Arsenious acid two grains  
Powdered opium eight grains  
Soap one scruple

Beat together and divide into twenty-four pills. One to be taken three times a-day. Have been found useful in intermittent fevers, periodical headaches, neuralgia, and lepra.

*A. T. Thomson.*

### ACIDUM BENZOICUM.

#### BENZOIC ACID.

R. Benzoin, any convenient quantity, put it into a glass vessel on a sand-bath, gradually increase the heat, and sublime as long as anything ascends. Wrap the sublimate in bibulous paper, and press, to separate the oil, and again sublime.

*U. S. Ph.*

R. Benzoin five parts  
Quicklime  
Hydrochloric acid, each, one part  
Water two hundred parts

Triturate the benzoin with the lime, boil in 100 parts of water, let stand, and when cold pour off the liquid. Boil residuum in 70 parts water, and pour off liquid when cold. Mix the solutions and evaporate to one half, filter, and gradually add the hydrochloric acid, wash the precipitate with the remainder of the water, dry, and sublime by a slow fire.

*Dub. Ph.*

Benzoic acid is in the form of white, feathery hexagonal crystals, when pure having no odor, but usually with an agreeable smell from the presence of oil. The taste is acrid acid. It is converted into hippuric acid, and voided by the urine when taken by man, and has been recommended in the uric acid diathesis and gout, but does not appear to be of any benefit. It has, however, been serviceable in catarrh of the bladder, and where there is a secretion of granular mucus with phosphates. It is also beneficial as a stimulating expectorant.

#### MIXTURE OF BENZOIC ACID AND COPAIBA.

R. Benzoic acid one drachm  
Balsam copaiba half fl. ounce  
White of egg q. s.  
Camphor water seven fl. ounces

Mix. Dose, two tablespoonfuls three times a-day. Recommended in chronic dysuria and vesical irritation.

*Walker.*

#### POWDER OF BENZOIC ACID AND IPECACUANHA.

R. Benzoic acid two scruples  
Ipecacuanha six grains  
Golden sulphur of antimony three grains  
White sugar one drachm

Mix, and triturate well. Divide into six powders. Dose, one, four times a-day in barley water. Said to be useful in asthenic pneumonia.

*Phœbus.*

#### MIXTURE OF BENZOIC ACID.

R. Benzoic acid  
Sulphur, each, one scruple  
Ipecacuanha six grains  
Honey six ounces  
Syrup of seneka  
Syrup of squills, each, one ounce

Mix. Dose, a teaspoonful, three or four times a-day, in the chronic catarrh of elderly persons.

*St. Marie.*

### ACIDUM BORACICUM.

#### BORACIC ACID.

R. Powdered borax forty parts  
Boiling water one hundred parts  
Dissolve, and add  
Muriatic acid twenty-five parts  
Collect the acid, which crystallizes on cooling, on a filter, drain, wash with cold water, and dry at 234° F. If not pure, dissolve and recrystallize.

*Wackenroder.*

If sulphuric acid be used, as is generally advised, it is almost impossible to free the boracic acid from a trace of it.

#### MIXTURE OF BORACIC ACID.

R. Boracic acid ten grains  
Oil of sweet almonds  
Syrup of lemons, each, one drachm  
Mix. Advised in cardialgia; to be repeated until a cessation of the pain.

*Pierquin.*

#### COLLUTORY WITH BORACIC ACID.

R. Cloves  
Galangal  
Ginger  
Peruvian bark, each, two drachms  
Gum lac one and a half drachms  
Benzoin  
Storax, each, half a drachm  
Alcohol six fl. drachms  
Acetic acid one and a half fl. ounces  
Vinegar five fl. ounces  
Macerate, press, and filter, and add



Boracic acid half a drachm

This has been recommended as a mouth-wash in a scorbutic condition of the gums. When used it is to be diluted with water. *Phoebus.*

## ACIDUM CITRICUM.

### CITRIC ACID.

This exists naturally in the juices of many fruits; to obtain it pure, boiling lemon or lime juice is to be saturated with chalk, when carbonic acid is given off, and an insoluble citrate is formed; this is to be washed and decomposed by boiling with dilute sulphuric acid; insoluble sulphate of lime precipitates, and the citric acid remains in solution and crystallizes on evaporation. To purify, dissolve in water and recrystallize. *Dispens. U. S.*

### ARTIFICIAL LEMON JUICE.

R. Citric acid one ounce  
Distilled water fourteen fl. ounces  
Oil of lemons five drops  
Mix. *Beasley.*

### LOZENGES OF CITRIC ACID.

R. Citric acid three drachms  
White sugar one pound  
Oil of lemons sixteen drops  
Triturate well, and add  
Mucilage of tragacanth q. s.  
Make lozenges of twelve grains each. *Cottureau.*

### SYRUP OF CITRIC ACID.

R. Citric acid five drachms  
Water ten drachms  
Simple syrup two pounds  
Tincture of fresh lemon peel one drachm  
Dissolve the acid in the water, mix with the syrup at a boiling heat, and, when cold, add the tincture. *Soubeyran.*

### DRY LEMONADE.

R. Citric acid two drachms  
White sugar four ounces  
Oil of lemons eight drops  
Mix well. A spoonful to a tumbler of water. *Gray.*

## ACIDUM GALLICUM.

### GALLIC ACID.

It is not certain that this acid exists as such in nature, or whether it arises from the decomposition of tannic acid. It is usually prepared from galls. Many processes have been devised,

R. Decoction of galls at will  
Expose to action of air in a loosely covered vessel for some months; it will grow

mouldy, and become covered with a glutinous pellicle, and gallic acid will be deposited on the sides of the vessel and on the under surface of the pellicle; collect, dissolve, and recrystallize. *Scheele.*

R. Decoction of galls at will  
Sulphuric acid sufficient  
to precipitate, wash the precipitate with and dissolve by aid of heat in diluted sulphuric acid, boil for a few minutes, let cool, and collect the crystals. *Liebig.*

Gallic acid is a powerful astringent, and has been found useful in hemorrhages and fluxes, as well as in checking night sweats in phthisis.

### PILLS OF GALLIC ACID.

R. Gallic acid q. s.  
Extract gentian sufficient to form pills of two to five grains each. One to be given every three or four hours.  
Useful in menorrhagia, hematuria, &c. *Dunghison.*

### INJECTION OF GALLIC ACID.

R. Gallic acid one scruple to one drachm  
Water two pints  
Mix. Found beneficial in leucorrhœa. *Dunghison.*

## ACIDUM HYDROCYANICUM.

### HYDROCYANIC ACID.

R. Ferrocyanuret of potassium two ounces  
Sulphuric acid one ounce and a half  
Distilled water sufficient  
Mix the acid with four fl. ounces of the water, and pour the mixture, when cool, into a glass retort. To this add the ferrocyanuret, previously dissolved in ten fl. ounces of the water. Pour eight fl. ounces of distilled water into a cooled receiver, and having attached the retort, distil on a sand-bath, with a moderate heat, six fl. ounces. Lastly, add to the product five fl. ounces of distilled water, or as much as will render the acid of such strength, that 100 grains will be accurately saturated by 12.7 grains of nitrate of silver. *U. S. Ph.*

### EXTEMPORANEOUS HYDROCYANIC ACID.

R. Cyanuret of silver fifty grains and a half  
Muriatic acid forty-one grains  
Distilled water one fl. ounce  
Mix the acid with the water, and add the cyanuret, and shake in a well-stopped bottle. Let settle, decant, and keep for use. *U. S. Ph.*

This acid should be kept in closely stopped



bottles, protected from the light. It is a transparent, volatile liquid, of a cooling and then somewhat irritating taste, and a peculiar smell. It is the most active poison known, and must be used with extreme caution. It is used as an anodyne and antispasmodic, in many diseases. The dose is from one to six drops mixed with gum water or syrup, always beginning with the smallest quantity and gradually increasing.

#### MIXTURE OF HYDROCYANIC ACID.

R. Medicinal hydrocyanic acid one fl. drachm  
Distilled water one pint  
Sugar one ounce and a-half  
Mix. A deserts spoonful twice a-day, gradually increasing the dose. *Magendie.*

R. Powdered gum Arabic half an ounce  
Water seven fl. ounces and a-half  
Dissolve, and add  
Syrup of tolu half fl. ounce  
Hydrocyanic acid twelve drops  
Mix. A tablespoonful every three hours in the cough of phthisis. *S. G. Morton.*

R. Medicinal hydrocyanic acid half a drachm  
Powdered sugar  
candy one ounce and a-half  
Syrup of red cabbage  
" Mallow, each, two ounces  
" Balsam of tolu  
" Maidenhair, each, one ounce  
" Poppies  
" Cinnamon, each, two drachms  
Mix. A spoonful occasionally. *Pierquin.*

#### SYRUP OF HYDROCYANIC ACID.

R. Medicinal hydrocyanic acid four grains and a-half  
Clarified syrup one ounce  
Mix. Contains  $\frac{2}{15}$  of grain of anhydrous acid. *Magendie.*

#### JULEP OF HYDROCYANIC ACID.

R. Medicinal hydrocyanic acid fifteen drops  
Hoffmann's anodyne two fl. ounces  
Syrup of marsh mallow three fl. ounces  
Mix. A spoonful every two hours. *Pierquin.*

R. Medicinal hydrocyanic acid two to four drops  
Syrup of peppermint one fl. ounce  
Infusion of linden four fl. ounces  
Mix. A spoonful every hour. *Foy.*

#### INJECTION OF HYDROCYANIC ACID.

R. Medicinal hydrocyanic acid one part  
Distilled water four parts  
In gonorrhœa. *Foy.*

#### LOTION OF HYDROCYANIC ACID.

R. Hydrocyanic acid half fl. ounce  
Alcohol one fl. ounce  
Distilled water ten fl. ounces and a half  
Mix. As lotion, in impetigo. *A. T. Thomson.*

R. Medicinal hydrocyanic acid two fl. drachms  
Lettuce water two pints  
Mix. In hepatic affections. *Magendie.*  
R. Hydrocyanic acid half fl. drachm  
Bicarbonate of soda two drachms  
Milk eight fl. ounces  
Mix. In milky scall. *A. T. Thomson.*

#### ACIDUM HYDRIODICUM.

##### HYDRIODIC ACID.

R. Tartaric acid two hundred and sixty-four grains  
Iodide of potassium three hundred and thirty grains  
Dissolve each in one fl. ounce and a half of distilled water. Mix the solutions, shake and let settle, filter, and add distilled water to make up measure of six and a quarter fl. ounces. *Buchanan.*

Has the same properties as iodine; dose, at first a few drops, gradually to half a fl. ounce three times a-day.

#### ACIDUM HYDROSULPHURICUM.

##### SULPHURETTED HYDROGEN.

R. Sulphuret of iron  
Sulphuric acid equal parts  
Pulverize the sulphuret, and gradually pour on it the acid diluted with three times its weight of water, and collect the gas. *Van Mons.*

This gas is an active poison, but has been administered in colica pictonum and mercurial affections. It is, however, much employed externally in cutaneous diseases.

##### POWDER AGAINST SALIVATION.

R. Sulphuret of magnesia one drachm  
Tartaric acid two scruples  
White sugar four scruples  
Mix, and divide into twelve powders. One to be taken every three hours in a little water, during the effervescence. *Radius.*

##### HYDROSULPHURETTED BATH.

R. Sulphuret of potash four ounces  
Water one pint  
Dissolve, and add



Muriatic acid two drachms  
 Pour the whole into the bath.  
 Advised in chronic diseases of the skin,  
 rheumatism, and certain cases of paralysis.  
*Cadet de Gassicourt.*

## ARTIFICIAL SULPHURETTED WATER.

R. Carbonate of soda twelve grains  
 Liquid hydrosulphuric acid two pints  
 Water six pints  
 To be kept in well-closed bottles. Has  
 been recommended in colica pictonum.  
*Foy.*

## HYDROSULPHURETTED LOTION.

R. Sulphuret of potash twenty-four parts  
 Water two hundred and fifty parts  
 Dissolve, and add  
 Sulphuric acid one part  
 Used as a wash in chronic diseases of the  
 skin.  
*Dupuytren.*

## ACIDUM LACTICUM.

## LACTIC ACID.

R. Sugar six pounds  
 Tartaric acid half an ounce  
 Boiling water twenty-six pounds  
 Mix, and set aside for a few days, then add  
 Old stinking cheese eight ounces  
 well diffused in  
 Curdled acid skimmed  
 milk eight pounds  
 Powdered chalk three pounds  
 Place in a warm situation, so as to keep  
 the mixture at a temperature of 86° to 95°. Stir often; in eight or ten days it will  
 solidify into a stiff paste of lactate of lime;  
 now add twenty pounds of boiling water,  
 and half an ounce of caustic lime, boil for  
 half an hour, and filter through linen.  
 Evaporate the liquid to the consistence of  
 syrup, and set aside for four days; remove  
 the lactate of lime, express, agitate with  
 one-tenth of cold water, and express, re-  
 peating this operation two or three times.  
 Now dissolve in twice its weight of boiling  
 water, and for every pound of the lactate,  
 add three and a-half ounces of sulphuric  
 acid diluted with an equal weight of water.  
 Filter the hot liquid through a bag, and  
 boil it with one pound and three-eighths  
 of carbonate of zinc, for every pound of  
 sulphuric acid, for a quarter of an hour.  
 Filter whilst boiling hot, and let stand to  
 crystallize; remove these crystalline crusts  
 of lactate of zinc, and wash them with cold  
 water. Then dissolve them in seven and  
 a-half parts of boiling water, and pass

through the solution a current of sulphu-  
 retted hydrogen, till sulphuret of zinc no  
 longer separates. Filter, boil the liquid,  
 to expel the excess of sulphuretted hydro-  
 gen, and evaporate on a water-bath to the  
 consistence of syrup.  
*Bensch.*

The juice of beets is to be permitted to  
 ferment for two months in a warm place,  
 then evaporate to the consistence of syrup,  
 treat with alcohol, evaporate, dissolve the  
 residuum in water, saturate, filter, concen-  
 trate, and crystallize. Dissolve the crys-  
 tals in water, purify by boiling with animal  
 charcoal, filter whilst hot, treat with baryta  
 and then with sulphuric acid; the lactic  
 acid will be set free, and may be concen-  
 trated in vacuo.  
*Cottureau.*

Lactic acid is in the form of a colorless syrup,  
 inodorous, but of a very sour taste. It attracts  
 moisture from the air. It has been recom-  
 mended by Magendie in dyspepsia, and may,  
 perhaps, be useful in a phosphatic diathesis.

## LOZENGES OF LACTIC ACID.

R. Lactic acid two drachms  
 Powdered white sugar one ounce  
 Gum tragacanth q. s.  
 Oil of vanilla four drops  
 Mix, and form pastilles of half a drachm  
 each, of which six may be taken during  
 the day. They should be kept dry.  
*Magendie.*

## LEMONADE OF LACTIC ACID.

R. Lactic acid one to four drachms  
 Simple syrup two fl. ounces  
 Water two pints  
 Mix. Dose, a cupful several times a-day.  
*Magendie.*

## ACIDUM MURIATICUM.

## MURIATIC ACID.

Prepared by distilling common salt in a glass  
 retort with sulphuric acid and water. It has a  
 density of 1.16, a suffocating odor, an acrid and  
 sour taste. Is very volatile. It gives a curdy-  
 white precipitate with nitrate of silver. It is  
 never given internally except in a diluted form.

## DILUTED MURIATIC ACID.

R. Muriatic acid four fl. ounces  
 Distilled water twelve fl. ounces  
 Mix.  
*U. S. Ph.*

## GASEOUS MURIATIC ACID.

R. Common salt two parts  
 Sulphuric acid three parts  
 Mix. The muriatic acid fumes that arise  
 are considered to be disinfectant, but are  
 much inferior to chlorine.  
*Swediaur.*



## MURIATIC ACID BATH.

R. Muriatic acid           ten fl. ounces  
 Water                   fifty gallons  
 Mix. Found useful in some chronic diseases of the skin. *Soubeiran.*

## MURIATIC ACID GARGLE.

R. Muriatic acid           two fl. drachms  
 Honey                   two fl. ounces  
 Barley water           one pint  
 Mix. Useful in the angina of scarlet fever, and in ulceration of the mouth and throat. *Ratier.*

## MURIATIC ACID LOTION.

R. Muriatic acid           one part  
 Water                   sixteen parts  
 Mix. Advised as a wash for chilblains, and also found beneficial in lepra and other skin diseases. *Foy.*

## MURIATIC ACID PEDILUVIUM.

R. Muriatic acid           two fl. ounces  
 Water                   eight pints  
 Mix. Advised by Scott and others in chronic enlargements of the liver and spleen. *Beral.*

## MURIATIC ACID DRAUGHT.

R. Muriatic acid           ten to twenty drops  
 Barley water           eight ounces  
 Mix. Dose, a tablespoonful three or four times a day. Has been advised in stone in the bladder. *Ellis.*

## MURIATIC ACID LINIMENT.

R. Balsam Peru           one drachm  
 Spermaceti  
 White wax  
 Muriatic acid, each,   two drachms  
 Olive oil               two ounces  
 Water                   six ounces  
 Rub well together. To be applied twice a-day. *Ratier.*

## ACIDUM NITRICUM.

## NITRIC ACID.

Nitric acid of the specific gravity 1.5.

## DILUTED NITRIC ACID.

R. Nitric acid           one fl. ounce  
 Distilled water       nine fl. ounces  
 Mix. *U. S. Ph.*  
 Dose, twenty to forty drops, in sweetened water, three times a-day.

## COLLUTORY OF NITRIC ACID.

R. Nitric acid           one fl. scruple  
 Honey of roses  
 Syrup of mulberries, each, half fl. ounce  
 Mix. To touch obstinate venereal ulcers in the mouth. *Phæbus.*

## FOMENTATION OF NITRIC ACID.

R. Nitric acid           one part  
 Distilled water       ninety-six parts  
 Mix. Used to destroy the fetid smell of foul ulcers; also as a wash in itch. *Soubeiran.*

## LOTION OF NITRIC ACID.

R. Nitric acid           half to one fl. drachm  
 Laudanum           one fl. drachm  
 Rose water           six fl. ounces  
 Mix. To wash venereal ulcers. *Phæbus.*

## NITRIC ACID MIXTURE.

R. Nitric acid           two fl. drachms  
 Raspberry syrup   three fl. ounces  
 Sugar               three ounces  
 Water               two pints  
 A wineglassful, with as much Seltzer water, in fetid breath. *Cadet.*

R. Nitric acid           half fl. drachm  
 Water               one pint and a-half  
 Syrup of cinnamon   one fl. ounce and a-half  
 Mix. In syphilis, chronic hepatitis, &c. A spoonful every two or three hours. *Augustin.*

R. Nitric acid           two fl. scruples  
 Opium               two grains  
 Water               two fl. ounces  
 Syrup of cinnamon   half fl. ounce  
 Mix. A spoonful every hour, in barley water. In dysentery, cholera, &c. *Ammon.*

R. Nitric acid           half fl. drachm  
 Gum Arabic  
 Sugar, each,       three drachms  
 Water               eight fl. ounces  
 Mix. A small tablespoonful in water, as occasion may require. As a tonic, to arrest colliquative sweats, &c. *Ellis.*

## OINTMENT OF NITRIC ACID.

R. Olive oil           one pound  
 Lard               four ounces  
 Nitric acid five fl. drachms and a half  
 Melt the oil and lard together in a glass vessel; when, on cooling, they begin to stiffen, add the acid, and stir. *Dub. Ph.*  
 Used as application to porrigo, psora, &c.

## LINIMENT OF NITRIC ACID.

R. Nitric acid           two fl. ounces  
 Oil of turpentine   three fl. ounces  
 Theriac           three drachms  
 Honey           one ounce  
 Alcohol           six fl. ounces  
 Mix. As a rubefacient, has been used in cholera. *Foy.*



## ACIDUM NITRO-MURIATICUM.

## NITRO-MURIATIC ACID.

R. Nitric acid by measure      one part  
 Muriatic acid      two parts  
 Mix in a refrigerated bottle, and keep in  
 a cool, dark place. *Dub. Ph.*

## NITRO-MURIATIC ACID BATH.

R. Nitro-muriatic acid-bath      six fl. ounces  
 Water      three gallons  
 Mix. Recommended by Scott as a foot-  
 bath in chronic hepatitis.

## ACIDUM OXALICUM.

## OXALIC ACID.

This acid exists in many plants in combina-  
 tion with lime or potash, but is usually obtained  
 by the action of nitric acid on sugar or starch.  
 It is in the form of colorless, transparent, pris-  
 matic crystals, having no odor, but a very acid  
 taste. This acid is an active poison in large  
 doses. It is not used in medicine in this coun-  
 try, but is employed in France.

## LOZENGES OF OXALIC ACID.

R. Powdered oxalic acid      one drachm  
 White sugar      eight ounces  
 Gum tragacanth      two scruples  
 Water of orange peel      five drachms  
 Oil of lemons      eight drops  
 Mix, and form lozenges of ten grains each.  
*Cottureau.*

## ACIDUM PHOSPHORICUM.

## PHOSPHORIC ACID.

R. Bones calcined to whiteness      five parts  
 Sulphuric acid      three parts  
 Water      thirty parts  
 Mix, and boil for an hour, constantly stir-  
 ring, strain, evaporate the fluid to consist-  
 ence of honey, pour on an oiled slab, and  
 when cold break in pieces. *Van Mons.*

This affords not pure phosphoric acid,  
 but an acid phosphate of lime, yet is the  
 formula recognized by the Batavian, Rus-  
 sian, Prussian, Saxon, &c. Pharmacopœias.

R. Phosphorus      one part  
 Nitric acid      eight parts  
 Cut the phosphorus in small pieces, and  
 add it very gradually to the acid, placed  
 on a sand-bath; when all the phosphorus  
 has been changed into an acid, evaporate  
 to the consistence of a syrup. *Guibourt.*

## DILUTED PHOSPHORIC ACID.

R. Phosphorus      one ounce

Dilute nitric acid      four fl. ounces  
 Distilled water      ten fl. ounces

Add the phosphorus gradually to the nitric  
 acid and water in a glass retort, on a sand-  
 bath, apply heat, and distil eight fl.  
 ounces. Pour these back into the retort  
 and again distil eight fl. ounces, which  
 reject. Evaporate the residue in a plati-  
 num crucible to two ounces and six  
 drachms. When cold, add as much dis-  
 tilled water as will make twenty-eight  
 fl. ounces. *London Ph.*

This acid is stated to be useful in nervous  
 disorders, in lithiasis with phosphatic deposits,  
 in morbid ossifications, in diabetes, to allay thirst,  
 &c., and as a local application in caries. The  
 dose of the dilute acid is from ten drops to a  
 fl. drachm in gum water.

## PILLS OF PHOSPHORIC ACID.

R. Phosphoric acid  
 Assafoetida, each,      three drachms  
 Powdered calamus      q. s.  
 Make one hundred and eighty pills. Dose,  
 five to ten, three times a-day.

Recommended in caries. *Phœbus.*

## FOMENTATION OF PHOSPHORIC ACID.

R. Phosphoric acid      one ounce  
 Decoction of chamomile      eight ounces  
 Mix. Employed in caries, and as injec-  
 tion in fistulas. *Augustin.*

## PHOSPHORIC TINCTURE OF MYRRH.

R. Tincture of myrrh      half an ounce  
 Diluted phosphoric acid      thirty drops  
 Mix. Used in caries of the teeth or bones.  
*Augustin.*

## PHOSPHORIC ACID LEMONADE.

R. Diluted phosphoric acid      twenty-four  
 drops  
 Simple syrup      two ounces  
 Water      two pints  
 Mix. Dose, about an ounce to relieve  
 thirst, in diabetes. *Ratier.*

## ACIDUM SULPHURICUM.

## SULPHURIC ACID.

## DILUTE SULPHURIC ACID.

R. Sulphuric acid      one fl. ounce  
 Water      thirteen fl. ounces  
 Add the acid gradually to the water in a  
 glass vessel, and mix them. *U. S. Ph.*

## ELIXIR OF VITRIOL.

R. Sulphuric  
 acid      three and a-half fl. ounces  
 Ginger, bruised      one ounce  
 Cinnamon, bruised, one ounce and a-half



Alcohol two pints  
Drop the acid gradually into the alcohol, digest in a close glass vessel for three days, add the ginger and cinnamon, macerate for a week, and then filter. *U. S. Ph.*

#### SULPHURIC ACID AND ALCOHOL.

R. Sulphuric acid four ounces  
Acetic acid  
Alcohol, each, two pounds  
Mix. Vulnerary, antiseptic, and astringent. Dose, internally, twenty or thirty drops, in an appropriate vehicle. Externally, as a lotion to foul ulcers and contusions, and to arrest bleeding. *Campana.*

#### SULPHURIC ACID AND NITRIC ETHER.

R. Sulphuric acid  
Nitric ether equal parts  
Mix very gradually. Dose, five to ten drops in an ounce of water. Found beneficial in spasmodic attacks, sinking, and hemorrhages. *Vogler.*

#### SULPHURIC ACID LEMONADE.

R. Sulphuric acid one part  
Simple syrup thirty-two parts  
Water one hundred and sixty parts  
Mix. As a cooling drink in hemorrhages, and also found useful as a preventive of the effects of lead. *Beral.*

#### SULPHURIC ACID LINIMENT.

R. Sulphuric acid two drachms  
Olive oil two and a-half ounces  
Oil of turpentine one ounce  
Mix. In gentle frictions to chilblains, where the skin is not broken. *Foy.*

#### SULPHURIC ACID OINTMENT.

R. Olive oil eight parts  
Add gradually, constantly stirring,  
Sulphuric acid five parts  
After standing for twenty-four hours, wash well in tepid water, till it will not redden litmus paper.  
Has been used in itch and paralysis, in frictions. *Brugnatelli.*

### ACIDUM TANNICUM.

#### TANNIC ACID.

R. Powdered galls  
Sulphuric ether of each a sufficient quantity  
Put into a glass adapter, loosely closed at its lower end with carded cotton, sufficient galls to fill about one-half of it, pressing the powder slightly. Then fit the adapter accurately to a receiver, fill it with sul-

phuric ether, and close the upper end slightly with a cork. The liquid which passes into the receiver separates into two portions, the lower being the smaller and denser. Pour on additional ether until the lower stratum of liquid in the receiver no longer increases. Separate this, and evaporate to dryness in a capsule by a moderate heat, and reduce to powder. *U. S. Ph.*

Tannic acid is of a yellowish-white colour, of a powerfully astringent taste, soluble in water, less so in alcohol or ether. It is a strong astringent, and has been successfully used in hemorrhages and other fluxes. The dose is from two to ten grains.

#### PILLS OF TANNIC ACID.

R. Tannic acid six grains  
Gum Arabic twelve grains  
White sugar seventy-two grains  
Triturate well, and add  
Simple syrup sufficient  
to make pills of four grains each. Dose, one to four, morning and evening. *Beral.*

#### PILLS OF TANNIC ACID AND OPIUM.

R. Tannic acid half a drachm  
Extract of opium three-quarters of a grain  
Conserve of roses sufficient  
to make twenty pills.  
One every hour, in uterine hemorrhage. *Dumars.*

#### GARGLE OF TANNIC ACID.

R. Tannic acid one drachm  
Honey of roses two ounces  
Rose water two fl. ounces  
Distilled water eight ounces  
Mix. As a gargle, to arrest mercurial salivation. *Beral.*

#### INJECTION OF TANNIC ACID.

R. Tannic acid half a drachm  
Distilled water eight fl. ounces  
Dissolve. In gleet and leucorrhœa. *Beral.*

#### MIXTURE OF TANNIC ACID.

R. Tannic acid twelve grains  
Syrup of rhatany  
Syrup of gum, each, one ounce  
Camphor water four fl. ounces  
Mix. Six to twelve spoonfuls a-day, in diarrhœa. *Beral.*

#### OINTMENT OF TANNIC ACID.

R. Tannic acid  
Distilled water, each, two drachms  
Dissolve, and rub well with  
Lard twelve drachms



ACIDUM TARTARICUM.

### TARTARIC ACID.

Is prepared from bi-tartrate of potash, by saturating the excess of acid with chalk, and decomposing the insoluble tartrate of lime by means of dilute sulphuric acid, evaporating and crystallizing.

It is cooling and diuretic in doses of a drachm or more.

LOZENGES OF TARTARIC ACID.

R. Powdered tartaric acid	three drachms
White sugar	one pound
Oil of lemons	sixteen drops
Mucilage of tragacanth	sufficient

Make into lozenges of twelve grains.

Used to allay thirst. *Cottureau.*

LEMONADE OF TARTARIC ACID X

R. Tartaric acid	one drachm
White sugar	one ounce
Water	twelve fl. ounces

Dissolve. For an agreeable acidulous  
drink. *Foy.*

ACIDUM VALERIANICUM.

### VALERIANIC ACID.

R. Valerian	forty pounds
Distilled water	three hundred and twenty pints

Distil, collect the oil, continuing the distillation as long as the water is acid. Agitate the oil with milk of lime; nearly saturate the acid water by milk of lime, first adding that already used, and then fresh; and, lastly, add lime water to excess. Concentrate till a pellicle appears, then decompose by nitric acid in a long, narrow flask with a ground stopper, decant the valerianic acid, which floats on the liquid, and distil it with a gentle heat till the distilled fluid ceases to be oily. *P. L. Bonaparte.*

R. Valerian at will  
Boil the root for three or four hours with rather more than its bulk of water, in which an ounce of carbonate of soda is dissolved for every pound of the root, replacing the water as it evaporates. Express strongly; boil the residue twice with the same quantity of water, expressing each time. Mix the liquids, add two fl. drachms of strong sulphuric acid for every pound of the root, and distil till three-fourths of the liquid have passed over. Neutralize this by carbonate of soda, concentrate, decompose the valerianate of soda by sulphuric acid, and separate the free valerianic acid by distillation.

*T. and H. Smith.*

ACONITUM.

ACONITE. MONKSHOOD.

Many species of *Aconitum* are official in the various pharmacopœias, but those recognized by the U. S. Ph. are *A. napellus* and *A. paniculatum*, both natives of Europe.

*Sex. Syst.* Polyand. Trigyn. *Nat. Syst.* Ranunculaceæ.

Linn. Sp. Pl. 751. Griffith, Med. Bot. 90.

The whole plant is possessed of highly deleterious properties, but the root is the most powerful. The leaves are, however, principally made use of. They have a somewhat nauseous odor, and a bitterish, acrid taste, followed by a peculiar tingling and burning of the lips. They owe their properties to the presence of a peculiar principle called *Aconitina*.

COMPOUND POWDER OF ACONITE.

R. Powdered aconite leaves  
Precip. sulph. of  
    antimon., each,                     one grain  
Carbonate of magnesia               one scruple  
Mix. As an anodyne and diaphoretic in  
gout and rheumatism.               *Vogler.*

EXTRACT OF ACONITE.

R. Recent leaves of aconite                      one pound  
Moisten with water, bruise in a stone mortar, express the juice, and evaporate to proper consistence.                      *U. S. Ph.*

R. Recent leaves of  
aconite                      a sufficient quantity.  
Beat to a pulp, express the juice; subject  
the residuum to displacement with rectified  
spirit, as long as the liquid is colored;  
unite the juice and the tincture; filter;  
distil off the spirit, and evaporate to a  
proper consistence on a vapor-bath. *E. Ph.*

Both are uncertain preparations; when good, they should cause a numbness and tingling of the lips, in a few minutes after taking. Dose, one to two grains, to be gradually increased.

ALCOHOLIC EXTRACT OF ACONITE.

R. Monkshood                                one pound  
Diluted alcohol                              two pints

Make a tincture by displacement; distil off the alcohol, and evaporate. *U. S. Ph.*

Dose, one-sixth of a grain.

## PILLS OF EXTRACT OF ACONITE.

R. Alcohol. extract of aconite    one grain  
Powdered liquorice                   twelve grains  
Syrup     sufficient  
to form a consistent mass. Divide into  
six pills. Dose, one, every three or four  
hours. *Turnbull.*

ACONITE PLASTER.

R. A small portion of the alcoholic extract spread over the surface of common adhesive plaster.



This has been highly recommended in neuralgia, over the painful part. *Curtis.*

#### ACONITE OINTMENT.

R. Alcohol. extract of aconite one part  
Lard two parts

Recommended as a friction in neuralgia.  
*Turnbull.*

#### COMPOUND WINE OF ACONITE.

R. Alcohol. extract of aconite one drachm  
Antimonial wine one ounce  
Make a solution. Dose, fifteen or twenty drops every three hours, gradually increasing till some effect is produced.

Found beneficial in chronic rheumatism, toothache, neuralgia, &c. *Richter.*

#### TINCTURE OF ACONITE.

R. Aconite four ounces  
Diluted alcohol two pints  
Macerate for fourteen days, and filter; or prepare by displacement. Dose, five drops three times a-day. The effects should be carefully watched. In rheumatism, neuralgia, &c. It is also employed as an embrocation to the painful spots, by means of a small piece of sponge attached to a handle. *U. S. Ph.*

#### TINCTURE OF ROOT OF ACONITE.

R. Root of aconite, in  
fine powder, sixteen ounces  
Alcohol .835 sufficient  
Mix the powder with a pint and a-half of the alcohol, macerate for two days, put it into a displacer, cover the surface with a piece of filtering paper, and return the fluid until it becomes clear; then add alcohol in small portions, suffering one to disappear before adding another, until the amount displaced amounts to four pints, reduce this by evaporation to two pints; filter, if necessary. *W. Procter, Jr.*

#### AMMONIATED EXTRACT OF ACONITE.

R. Alcohol. extract of aconite one drachm  
Water of ammonia ten drops  
Evaporate gently, to expel excess of ammonia.

#### AMMONIATED ACONITE OINTMENT.

R. Ammon. extract of aconite one drachm  
Lard three drachms  
Rub well together.

Used for the same purposes as the aconite ointment, but more active and pungent. *Turnbull.*

#### MIXTURE OF ACONITE.

R. Tincture of aconite one fl. drachm  
Carbonate of soda one drachm and a-half

Sulphate of magnesia

one ounce and a-half

Water

six fl. ounces

Mix. In gastralgia. A tablespoonful, when the pain is urgent. *Fleming.*

#### ACONITINA.

##### ACONITINE.

R. Aconite root, dried and  
bruised, two pounds  
Rectified spirit three gallons  
Diluted sulphuric acid  
Solution of ammonia  
Purified animal charcoal, each, sufficient

Boil the aconite with a gallon of the spirit, for an hour, in a retort attached to a receiver. Pour off the tincture, and repeat the operation a second and third time. Then express, mix the tinctures, filter, and distil off the spirit. Evaporate the residue to consistence of an extract. Dissolve this in water, and filter. Evaporate the solution with a gentle heat to consistence of syrup. Add to it the diluted acid mixed with distilled water. Drop in the solution of ammonia, and dissolve the precipitated aconitina in diluted sulphuric acid mixed with water. Then add the animal charcoal, occasionally shaking, for a quarter of an hour. Lastly, filter, and, having again dropped in the solution of ammonia, wash the precipitate and dry it. *Lond. Ph.*

Not used internally, but successfully employed externally in neuralgia, gout, and rheumatism.

##### ACONITINE OINTMENT.

R. Aconitine two grains  
Alcohol six drops  
Rub well together, and add  
Lard one drachm  
*Turnbull.*

R. Aconitine sixteen grains  
Olive oil half a drachm  
Lard one ounce  
Incorporate thoroughly. *Turnbull.*

These ointments are employed as frictions in neuralgia, over the painful parts. A small portion, not exceeding the size of a pea, is to be used at a time. The operation to be repeated three or four times a-day.

##### ACONITINE LOTION.

R. Aconitine eight grains  
Alcohol two fl. ounces  
Dissolve. Used by means of a friction sponge over the affected part. Never to be employed where the skin is broken or abraded. *Turnbull.*



ADIANTUM.

MAIDENHAIR.

Two species of this genus, *A. pedatum* and *A. capillus veneris* have been much used in Europe as remedies in pectoral affections. They are both bitter and aromatic, but the former is the most active, and is a native of the United States.  
*Sex. Syst. Cryp. Fil. Nat. Syst. Polypodiaceæ.*

INFUSION OF MAIDENHAIR.

R. Maidenhair three drachms  
Liquorice root two drachms  
Water sufficient  
to obtain two pints of infusion. As a drink in catarrh.

SYRUP OF MAIDENHAIR.

R. Maidenhair four ounces  
Boiling water three pints  
Infuse for two hours, strain, and add  
Sugar four pounds  
Clarify with white of egg, and add  
Maidenhair two ounces  
Digest for two hours, and strain.  
*Cottureau.*

This syrup is known under the name of syrup of capillaire, and is much employed in Europe as a pectoral.

COMPOUND SYRUP OF MAIDENHAIR.

R. Maidenhair five ounces  
Marsh mallow two drachms  
Asparagus root one ounce  
Liquorice root two ounces  
Water two pints  
Boil to a pint and a half, express, strain, and clarify, and add  
Sugar thirty-two ounces  
Make syrup. *Wurtemberg Ph.*

ÆTHER ACETICUS.

ACETIC ETHER.

R. Alcohol (36 degrees) three thousand parts  
Acetic acid (10 degrees) two thousand parts

Sulphuric acid  
(66 degrees) six hundred and twenty-five parts

Mix the alcohol and acetic acid in a glass retort, add gradually the other acid, distil in a sand-bath four thousand parts, rectify this product with a small quantity of carbonate of potassa to obtain three thousand parts.  
*Cottureau.*

This ether is milder, more agreeable and diaphoretic, than the other ethers. It is used in low fevers, spasmodic vomiting, cardialgia, in doses of ten drops to a drachm.

SPIRIT OF ACETIC ETHER.

R. Acetic ether one part  
Alcohol three parts  
Mix. This is employed as a substitute for Hoffmann's anodyne, in about the same doses.  
*Giordano.*

MIXTURE OF ACETIC ETHER.

R. Acetic ether  
Ethereal tincture of valerian  
Tincture of opium equal parts  
Mix. Dose, from ten to thirty drops, in hysteria and hypochondriasis.  
*Radiis.*

ÆTHER HYPONITROSUS.

NITROUS ETHER.

R. Rectified spirit fifteen fl. ounces  
Pure nitric acid (1.500) seven fl. ounces  
Put the spirit with a little clean sand into a two pint matrass, fitted with a cork, through which are passed a safety tube, terminating an inch above the spirit, and another tube leading to a refrigerator. The safety-tube being filled with pure nitric acid, add through it gradually three and a-half fl. ounces of the acid. When the ebullition which ensues is nearly over, add the rest of the acid gradually, half a fl. ounce at a time, waiting till the ebullition which slowly arises has subsided, and cooling the refrigerator with a stream of water. The ether which distils over, being received in a bottle, is to be agitated first with a little milk of lime, till it ceases to redden litmus paper, and then with half its volume of a concentrated solution of muriate of lime.  
*Edin. Ph.*

This ether is stimulant, antispasmodic, carminative, and diuretic. The dose is from ten to forty drops.

SWEET SPIRIT OF NITRE.

R. Hyponitrous ether one part  
Rectified spirit four parts  
Mix. *Edin. Ph.*

R. Nitrate of potassa, in powder, two pounds  
Sulphuric acid a pound and a-half  
Alcohol nine pints and a-half  
Diluted alcohol one pint  
Carbonate of potassa one ounce

Mix the nitrate of potassa and the acid in a glass retort, and, having gradually added the alcohol, digest with a gentle heat for two hours; then raise the heat and distil a gallon. To the distilled liquor add the diluted alcohol and carbonate of potassa, and again distil a gallon.  
*U. S. Ph.*



Diuretic, diaphoretic, and carminative.  
Dose, half a fl. drachm to three fl. drachms.

#### MIXTURE OF SWEET SPIRIT OF NITRE.

R. Sweet spirit of nitre                      one drachm  
Hoffmann's anodyne  
Aromatic ammoniated  
alcohol, each,                      two drachms  
Mint water                      six ounces  
Mix. Dose, a spoonful occasionally.  
*Brera.*

#### MIXTURE OF HYPONITROUS ETHER.

R. Ipecacuanha                      half a drachm  
Boiling water                      sufficient  
To obtain six ounces of infusion, strain  
and add  
Hyponitrous ether                      one drachm  
Extract of juniper berries                      one ounce  
Mix. Dose, a spoonful every two hours,  
in the dropsical swellings consecutive to  
scarlatina.  
*Radius.*

### ÆTHER HYDROCYANICUS.

#### HYDROCYNIC ETHER.

R. Cyanuret of potassium  
Sulpho-vinate of barytes, equal parts  
Mix, and put in a glass retort, distil by a  
moderate heat. The product separates  
into two strata, the lighter of which is im-  
pure hydrocyanic ether; this is to be sepa-  
rated and agitated with four or five times  
its bulk of water at 120° to 140° F., again  
agitated with a little water, decanted, and  
placed in contact with chloride of lime for  
twenty-four hours, and then distilled.

*Magendie.*

This preparation is said to resemble hydro-  
cyanic acid in its therapeutical effects, but is less  
active. Its smell is, however, penetrating and  
offensive. The dose is two to six drops, in mu-  
cilage or emulsion, in obstinate or convulsive  
coughs.

### ÆTHER MURIATICUS.

#### MURIATIC ETHER.

R. Muriatic acid  
Alcohol                      equal parts  
Distil in a Wolf's apparatus, and preserve  
the product contained in the second re-  
ceiver.  
*Cotteneau.*

It has much the same properties as the other  
ethers. Dose, thirty to forty drops.

#### SPIRIT OF MURIATIC ETHER.

R. Muriatic ether  
Alcohol                      equal parts  
Mix.                      *Cotteneau.*

#### MIXTURE OF MURIATIC ETHER.

R. Spirit of muriatic ether                      half a drachm  
Parsley water  
Syrup of rhubarb, each,                      one ounce  
Mix. A teaspoonful every hour for young  
children as a diuretic.  
*Wendt.*

### ÆTHER SULPHURICUS.

#### SULPHURIC ETHER.

This is commonly known under the name of  
ether. It is an energetic diffusible stimulant, and is  
much employed as an excitant and antispasmodic  
in what are termed nervous disorders. It is also  
considered to be anthelmintic. In vapor it has  
been used in small quantities as an inhalation in  
chronic irritations of the lungs, and of late under  
the name of *letheon* to induce unconsciousness  
to pain in surgical operations. The dose of the  
fluid is from half a fl. drachm to two drachms.  
When employed as an inhalation it should be  
purified.

#### RECTIFIED SULPHURIC ETHER.

R. Sulphuric ether                      fourteen fl. ounces  
Potassa                      half an ounce  
Distilled water                      eleven fl. ounces  
Dissolve the potassa in two fl. ounces of  
the water, and add the ether to the solu-  
tion, shaking well; then distil at 120° F.  
twelve ounces of rectified ether. Shake  
the distilled fluid with nine fl. ounces of  
water, and set them by for the water to  
subside, then pour off the ether and keep  
in a well-closed bottle.  
*U. S. Ph.*

#### ETHEREAL OIL.

R. Alcohol                      two parts  
Sulphuric acid                      four parts  
Mix. Distil till a black froth begins to  
arise, remove the retort from the fire, when  
cool, add water to the fluid in the receiver,  
and remove the oil that may float on the  
surface. Agitate this with a solution of  
potassa and separate the ethereal oil.

*Van Mons.*

#### SPIRIT OF SULPHURIC ETHER.

R. Rectified sulphuric ether                      half a pint  
Alcohol                      one pint  
Mix.                      *U. S. Ph.*

#### HOFFMANN'S ANODYNE.

R. Spirit of sulphuric ether                      one pint  
Ethereal oil                      two fluidrachms  
Mix.                      *U. S. Ph.*

#### LOTION OF SULPHURIC ETHER.

R. Sulphuric ether                      two fl. ounces  
Vinegar                      six fl. ounces  
Rose water                      four fl. ounces  
Distilled water                      two pints



Mix. As a lotion to painful inflammatory tumors. *Pierquin.*

SYRUP OF SULPHURIC ETHER.

R. Sulphuric ether one fl. ounce  
Syrup one pint

Mix in a glass vessel, having a stop-cock at the lower part, shake occasionally for a week, and draw off when clear into small bottles. *Par. Cod.*

Dose, half an ounce to an ounce.

R. White sugar twenty-one ounces  
Distilled water six ounces  
Sulphuric ether two ounces

Mix, and pour into a closed flask, agitate for some minutes, and add

Distilled water six ounces  
Agitate occasionally for five days, filter in a covered funnel. *Paton.*

MIXTURE OF SULPHURIC ETHER AND CAMPHOR.

R. Camphor mixture seven fl. ounces  
Sulphuric ether  
Syrup of saffron, each, half fl. ounce  
Mix.

MIXTURE OF SULPHURIC ETHER AND TURPENTINE.

R. Sulphuric ether one fl. drachm  
Oil of turpentine two fl. drachms  
Sugar four drachms  
Water two fl. ounces

Mix. Two fl. drachms every quarter of an hour in poisoning by nux vomica. *Orfila.*

ÆTHER TEREBINTHINATUS.

TEREBINTHINATED ETHER.

R. Alcohol two pounds  
Spirit of turpentine half a pound  
Mix, and add gradually

Nitric acid two pounds  
Distil off one-half at a gentle heat. Dose, from twenty to forty drops, in syrup or mucilage. Used internally and externally in cases of biliary calculi, jaundice, engorgements of the liver, and rheumatism. *Cadet.*

ALETRIS.

STAR GRASS.

*Aletris farinosa.* Linn. Bigelow. Am. Med. Bot. iii. 50. Griffith, Med. Bot. 623.

*Sex. Syst.* Hexandria monogynia. *Nat. Syst.* Hæmodoraceæ. Brown.

A native plant, with an intensely bitter root, which is the officinal portion. The powder is tonic in doses of ten grains.

TINCTURE OF ALETRIS.

R. Aletris root, bruised, three ounces  
Diluted alcohol two pints  
Make tincture by displacement. Dose, half an ounce to an ounce in colic, and in smaller doses in chronic rheumatism.

ALLIUM, U. S.

GARLIC.

Several species of *Allium* have been employed in medicine, but the most important, and that recognized by the Pharmacopœia, is *A. sativum*. Linn. Griffith, Med. Bot. 653. The part used is the bulb, which is composed of several bulb-lets or *cloves* (*spicæ*), surrounded by a dry, white, thin capsular membrane. These cloves have a peculiar odor, and an acrid and pungent taste. The properties depend on the presence of a volatile oil, and are lost on drying.

*Sex. Syst.* Hex. monog. *Nat. Syst.* Liliaceæ.

Garlic is a stimulating expectorant; diuretic and diaphoretic when given internally, and acts as an irritant, or even vesicant, when applied externally. The dose in substance is from half a drachm to a drachm, or more.

SYRUP OF GARLIC.

R. Fresh garlic two ounces  
Distilled vinegar one pint  
Sugar two pounds

Macerate the garlic in the vinegar, in a glass vessel, for four days, express, and permit dregs to subside. Add sugar and form syrup *s. a.* *U. S. Ph.*

As an expectorant in chronic catarrhs, well suited for children. Dose, for a child, about a teaspoonful.

CATAPLASM OF GARLIC.

Bruised cloves of garlic, mixed with common bread and milk poultice, in different proportions. Used as a revulsive, but less active than the mustard cataplasm.

LINIMENT OF GARLIC.

R. Garlic  
Lard equal parts  
To be well rubbed together. Has been recommended as a revulsive in infantile nervous and convulsive disorders.

*Gassicourt.*

CAPILLARY LOTION.

R. Garlic three or four cloves  
Alcohol two pints

Macerate for thirty-six hours, filter, and add of

Burdock eight ounces

The head to be sponged with this every evening, for some weeks. It is said to be efficient to promote the growth of hair. *Phæbus.*



## A L O E .

## A L O E S .

Under this name are included the inspissated juices of several species of aloes, and especially of *A. vulgaris*, *A. socotrina*, and *A. spicata*.

*Sex. Syst.* Hexandria monogynia. *Nat. Syst.* Liliaceae.

Pereira, Mat. Med. ii. 113. Griffith, Med. Bot. 649.

Three varieties of aloes are found in the shops, the Cape, Socotrine, and Hepatic. The second, when genuine, is the most esteemed, but the first is by far the most abundant, and, when good, answers every purpose for which the drug is used.

*Dispens. U. S. art. Aloe.*

## P O W D E R O F A L O E S .

R. Aloes six grains  
White sugar one drachm  
Pulverize very finely. Recommended to be blown in the eye, to remove films and specks on the cornea. *Radius.*

## P O W D E R O F A L O E S A N D C A N E L L A .

R. Aloes one pound  
Canella bark three ounces  
Powder separately, and mix. Dose, five to fifteen grains. A popular emmenagogue, known as *Hiera Picra*.

## C O M P O U N D P O W D E R O F A L O E S .

R. Aloes an ounce and a half  
Guaiacum resin one ounce  
Compound powder of cinnamon half an ounce  
Powder the aloes and resin separately, and then mix them with the compound powder of cinnamon. Dose, ten to twenty grains. Used as a purgative and diaphoretic.

*London Ph.*

## E M M E N A G O G U E P O W D E R .

R. Iron rust six grains  
Powdered aloes two grains  
Magnesia thirty grains  
Mix. For a single dose, to be repeated three times a-day. *Brera.*

## A L O E S P I L L S .

R. Powdered aloes  
Soap, each, an ounce  
Form a mass, to be divided into two hundred and forty pills. Dose, one to three as a laxative, five or more as a purgative.

*U. S. Ph.*

## C O M P O U N D P I L L S O F A L O E S .

R. Powdered aloes half a drachm  
" rhubarb one drachm  
Oil of cloves four drops  
Soap eight grains  
Syrup of rhubarb sufficient  
Rub well together, and form forty pills.

In tardy menstruation, one at night or oftener if required, so as to open bowels but not to purge. *Dewees.*

R. Powdered aloes one ounce  
Extract of gentian half an ounce  
Oil of caraway forty minims  
Beat together till incorporated. *Lond. Ph.*  
Dose, five to fifteen grains. A valuable purgative in habitual costiveness.

## A L O E S A N D A S S A F E T I D A P I L L S .

R. Powdered aloes  
Assafetida  
Soap, each, half an ounce  
Beat with water to form a mass; divide into one hundred and eighty pills. Dose, two to five. Useful in costiveness with flatulency. *U. S. Ph.*

## A L O E S A N D M Y R R H P I L L S .

R. Powdered aloes two ounces  
Powdered myrrh one ounce  
Saffron half an ounce  
Syrup sufficient  
Beat together, to form a mass; divide into four hundred and eighty pills. *U. S. Ph.*  
Dose, from three to six. A well-known cathartic and emmenagogue, under the name of Rufus' pills.

## A L O E S A N D I R O N P I L L S .

R. Aloes two parts  
Sulphate of iron three parts  
Aromatic powder six parts  
Conserve of red roses eight parts  
Pulverize the aloes and sulphate of iron, mix the whole ingredients and beat into a mass, and divide into five grain pills. Dose, one to three. *Edin. Ph.*

A useful emmenagogue in chlorosis and atonic amenorrhœa.

## C H A P M A N ' S A P E R I E N T P I L L S .

R. Aloes sixteen grains  
Powdered rhubarb twenty-four grains  
Mastic twelve grains  
Mix, and make twelve pills. *Chapman.*

## C H A P M A N ' S A N T I - D Y S P E P T I C P I L L S .

R. Aloes half a drachm  
Powdered ipecacuanha ten grains  
Mastic thirty grains  
Oil of fennel eight drops  
Mix, and make twenty pills. *Chapman.*

## M I T C H E L L ' S A P E R I E N T P I L L S .

R. Aloes sixteen grains  
Powdered rhubarb thirty-two grains  
Calomel two grains  
Tartar emetic one grain  
Mix, and make sixteen pills.

*J. K. Mitchell.*



HOOPER'S PILLS.

R. Aloes four hundred parts  
Sulphate of iron two hundred parts  
Extract of black hellebore  
Myrrh  
Soap, each, one hundred parts  
Powdered canella  
Powdered ginger, each, fifty parts  
Break into a mass with water, and divide into pills of two and a-half grains.

Much used as cathartic and emmenagogue.  
*Journ. Phil. Col. Pharm. v. 25.*

ANDERSON'S PILLS.

R. Aloes seven hundred and eighty-seven parts  
Soap one hundred and thirty-one parts

Colocynth  
Gamboge, each, thirty-three parts  
Oil of aniseed sixteen parts  
Reduce the aloes, colocynth, and gamboge to a very fine powder, add oil of aniseed and soap, and beat into a mass with water; divide into three grain pills. A mild purgative. *Journ. Phil. Col. Pharm. v. 25.*

LADY WEBSTER'S PILLS.

R. Aloes six drachms  
Mastich  
Red rose-leaves, each, two drachms  
Syrup of wormwood sufficient  
to form a mass. Divide into three grain pills, used as a laxative in costiveness, dependent on impaired digestion. Dose, one to three. A better form is to substitute powdered rhubarb for the rose-leaves.  
*Cooley.*

MORRISON'S PILLS.

No. 1. R. Aloes  
Cream of tartar equal parts  
Mucilage sufficient  
to form mass.

No. 2. R. Aloes three parts  
Gamboge two parts  
Colocynth one part  
Cream of tartar four parts  
Powder finely, and add  
Simple syrup sufficient  
to form mass. Dose, five to ten grains.  
These purge actively, especially No. 2.  
*Cooley.*

FULLER'S PILLS.

R. Aloes half a drachm  
Senna  
Myrrh, each, one scruple  
Assafetida  
Galbanum, each, ten grains

Saffron

Mace, each, five grains  
Sulphate of iron two scruples  
Simple syrup sufficient  
to form mass. Dose, five to fifteen grains, used as an antispasmodic and aperient.  
*Cooley.*

JAMES' ANALEPTIC PILLS.

R. Antimonial powder  
Resin of guaiacum  
Aloes and myrrh pill equal parts  
Syrup sufficient  
Form a mass, and divide into four grain pills. A diaphoretic purgative. *Cooley.*

PETERS' PILLS.

R. Aloes  
Jalap  
Scammony  
Gamboge, each, two drachms  
Calomel one drachm  
Beat into a mass with alcohol. A powerful purgative. *Cooley.*

SPEEDIMAN'S PILLS.

R. Aloes  
Myrrh  
Rhubarb, each, one ounce  
Extract of chamomile half an ounce  
Beat into a mass with syrup, and divide into four grain pills. A good tonic and stomachic purge. *Burnett.*

SPLENETIC PILLS.

R. Strained aloes  
Gum ammoniac, each, an ounce and a-half  
Myrrh  
Bryony, each, two drachms  
Beat into a mass, and divide into four grain pills. Dose, three to five. Extolled in amenorrhœa and hypochondriasis.  
*Saunders.*

DUCHESNE'S PILLS.

R. Aloes  
Gum ammoniac, each, twenty-four grains  
Myrrh six grains  
Mastich  
Carbonate of potassa, each, two grains  
Saffron one grain  
Simple syrup sufficient  
to make a mass. Dose, from ten to twenty grains. Prescribed in engorgements of the abdominal viscera, supervening on intermittent fevers. *Van Mons.*

ANTICHLOROTIC PILLS.

R. Aloes  
Iron rust, each, half a drachm



Gum ammoniac one drachm  
 Extract of dandelion sufficient  
 to form mass. Divide into pills of three  
 grains each. Dose, three to six, morning  
 and evening, in chlorosis and amenorrhœa.  
*Radius.*

## BICKER'S PILLS.

R. Iron rust two drachms  
 Sulphur  
 Myrrh  
 Aloes, each, one drachm  
 Beef gall sufficient  
 to form mass. Divide into four grain pills.  
 Six, morning and evening. *Radius.*

## WHYTT'S PILLS.

R. Chloride of iron  
 Aloes  
 Extract of horehound, each, half a  
 drachm  
 Assafetida one drachm and a half  
 Form a mass. Divide into two grain pills.  
 Dose, four to five, three times a day, in  
 leucorrhœa and hysteria with constipa-  
 tion. *Radius.*

## BARTHEZ'S PILLS.

R. Aloes half a drachm  
 Myrrh one drachm  
 Musk twenty grains  
 Camphor ten grains  
 Balsam Peru sufficient  
 to form mass. Dose, eight grains, three  
 times a day. Advised in amenorrhœa and  
 chlorosis. *Pierquin.*

## PITSCHAFT'S ECCOPROTIC PILLS.

R. Strained aloes  
 Sulphate of quinine, each, one scruple  
 Mix, and divide into twenty pills. Dose,  
 one from time to time in the evening.  
 Found useful in torpor of the large intes-  
 tines. *Radius.*

## FRANK'S PILLS.

R. Aloes  
 Jalap, each, four parts  
 Rhubarb one part  
 Syrup of wormwood sufficient  
 to form mass. Divide into three grain  
 pills. Dose, one to four during the day.

These are the *Grains de Santé*, so much em-  
 ployed as dinner pills in Europe. *Foy.*

## ALOE AND RHUBARB PILLS.

R. Powdered aloes  
 Rhubarb, each, half a drachm  
 Soap sufficient  
 to form mass. Divide into twenty-five  
 pills. Three or four occasionally in dys-  
 pepsia with costiveness. *Ellis.*

## GRIFFITTS' PILLS.

R. Powdered rhubarb one drachm and  
 a half  
 Sulphate of iron half a drachm  
 Soap two scruples  
 Water sufficient  
 to form mass. Divide into forty pills.

A favorite remedy with the late Dr. S. P.  
 Griffiths, to remove costiveness and impart tone  
 to the bowels. Three or four, to be taken at  
 bed time.

## APERIENT PILLS.

R. Aloes  
 Rhubarb, each, one drachm  
 Ipecacuanha six grains  
 Soap one scruple  
 Form mass with water, and divide into  
 sixty-four pills. Dose, one at bed time,  
 as an aperient; two or three as a purga-  
 tive. *Ellis.*

## PILLS OF ALOES AND BLUE MASS.

R. Blue mass. one scruple  
 Powdered aloes twenty-five grains  
 Incorporate, and divide into fifteen pills.  
 Dose, one every two hours till they ope-  
 rate. Useful in constipation where there is  
 a deficiency of bile. *Ellis.*

## LAXATIVE ELECTUARY.

R. Aloes eight grains  
 Cream of tartar two drachms  
 Honey sufficient  
 to form electuary. For a single dose.  
 Advised in amenorrhœa attributed to ab-  
 dominal engorgement. *Brera.*

## ANTHELMINTIC SUPPOSITORY.

R. Aloes half an ounce  
 Common salt three drachms  
 Flour two ounces  
 Honey sufficient  
 to form a firm paste. As a suppository in  
 cases of ascarides. *Foy.*

## COMPOUND DECOCTION OF ALOES.

R. Extract of liquorice seven drachms  
 Carbonate of potassa one drachm  
 Powdered aloes  
 myrrh  
 Saffron, each, one drachm and a-half  
 Compound tincture of cardamoms  
 seven fl. ounces

Distilled water one pint and a-half  
 Boil down the liquorice, carbonate of po-  
 tassa, aloes, myrrh, and saffron with the  
 water to a pint, strain, and add the com-  
 pound tincture of cardamom. Dose, half  
 a fl. ounce to two fl. ounces. A mild ca-  
 thartic, tonic, antacid, and emmenagogue.  
*Lond. Ph.*



CLAUDER'S ELIXIR.

R. Carbonate of potassa  
Muriate of ammonia, each, one ounce  
Elder-flower water a pint and a-half  
Make a solution, and add  
Strained aloes  
Myrrh, each, one ounce  
Saffron two drachms  
Digest for twenty-four hours, and filter.  
Dose, half a drachm to a drachm. Has  
been recommended in obstructions of the  
abdominal viscera, in amenorrhœa, consti-  
pation, scurvy, &c. *Pideret*

DETERSIVE INJECTION.

R. Strained aloes ten grains  
Muriate of ammonia four grains  
Honey of roses one ounce  
Fennel water six ounces  
Make solution, and filter. Stated to be  
useful in chronic discharges from the ure-  
thra, to be injected three or four times a  
day. *Soubéiran.*

ANTHELMINTIC CLYSTER.

R. Powdered aloes one drachm  
Barley water one pint  
Mix. Very effectual against ascarides in  
the rectum. *Radiis.*

ALOETIC MIXTURE.

R. Strained aloes  
Myrrh, each, one drachm  
Cream of tartar half an ounce  
Myrrh water six ounces  
Make solution, and filter. Dose, a spoon-  
ful night and morning, to provoke hemor-  
rhoids. *Foy.*

WINE OF ALOES.

R. Aloes one ounce  
Cardamom *x*  
Ginger, each, one drachm  
Wine one pint  
Macerate for fourteen days, occasionally  
agitating, then filter. *U. S. Ph.*  
Purgative, in doses of half an ounce to  
two ounces, stomachic and tonic, in doses  
of one to two drachms.

ALKALINE WINE OF ALOES.

R. Aloes  
Myrrh *+*  
Saffron, each, one ounce  
Carbonate of potassa two ounces  
Wine two pints  
Digest for ten days, and filter. Bitter,  
tonic and stimulant. Recommended in  
dyspepsia with pyrosis, in doses of an  
ounce. *Swediaur.*

BALSAMIC WINE OF ALOES.

R. Aloes  
Myrrh  
Olibanum  
Angelica, each, half an ounce  
Balsam Peru one ounce  
Storax two ounces  
Benzoin three ounces  
Flowers of hypericum four handfuls  
Wine four pints  
Macerate for fourteen days, and strain.

TINCTURE OF ALOES.

R. Powdered aloes one ounce  
Liquorice three ounces  
Alcohol half a pint  
Distilled water one pint and a-half  
Macerate fourteen days, and filter. Dose,  
two drachms to an ounce. Purgative and  
stomachic. *U. S. Ph.*

TINCTURE OF ALOES AND MYRRH.

R. Powdered aloes three ounces  
Saffron one ounce  
Tincture of myrrh two pints  
Macerate for fourteen days, and filter.  
*U. S. Ph.*

R. Tincture of aloes three ounces  
Tincture of myrrh four ounces  
Tincture of saffron three ounces  
Mix. *Soubéiran.*

Long celebrated under the name of *elixir pro-  
prietatis*. It is purgative, stomachic, and em-  
menagogue. The dose is half a drachm to a  
drachm. Well suited to cold, torpid habits.

ETHEREAL TINCTURE OF ALOES.

R. Myrrh one ounce and a-half  
Spirit of sulphuric ether one pound  
Digest for four days, and add  
Aloes one ounce and a-half  
Saffron one ounce  
Again digest for four days, and filter.  
*Edin. Ph.*

This tincture has been highly esteemed as  
stomachic, vermifuge, and emmenagogue, in  
doses of half a drachm to a drachm.

BOERHAAVE'S ELIXIR.

R. Aloes  
Myrrh *x*  
Saffron, each, one ounce  
Tartrate of potassa two ounces  
Alcohol fourteen ounces  
Distilled water eight ounces  
Macerate for three days, and filter.  
*Han. Ph.*

This has been highly praised in visceral ob-  
structions. The dose is from half a drachm to  
a drachm.



## STOUGHTON'S ELIXIR.

R. Aloes	
Cascarilla, each,	one drachm
Rhubarb	four drachms
Wormwood	
Germander	
Gentian	
Orange peel, each,	six drachms
Alcohol	two pints

Macerate for four days, and filter. *Foy.*

Stimulant, tonic, and stomachic, in doses of twenty or thirty drops.

## ELIXIR OF GARUS.

R. Saffron	one ounce
Myrrh	two ounces
Aloes	ten ounces
Cinnamon	
Nutmeg, each,	half an ounce
Orange-flower water	one pound
Alcohol	sixteen pounds

Macerate for two days and distil off eight pounds, to which add

Syrup of maidenhair	two pounds
And color with caramel, dissolved in	
Orange-flower water	eight ounces

This preparation can scarcely be considered as a medicine, but is rather an agreeable liqueur. *Soubéiran.*

## OINTMENT OF ALOES.

R. Powdered aloes	two drachms
Lard	one ounce

Triturate well.

Has been recommended as a friction to the abdomen as a vermifuge in children.

*Soubéiran.*

## ALTHÆA.

## MARSH MALLOW.

Several species of Mallow are employed in medicine, but that which is officinal is *A. officinalis*, a herbaceous perennial, with pale purplish flowers, a native of Europe, in moist situations.

*Ser. Syst.* Monadelph. Polyand. *Nat. Syst.* Malvaceæ.

Linn. Sp. Pl. 966. Griffith, Med. Bot. 161.

The parts used are the roots and leaves. These are inodorous, and have a vapid mucilaginous taste. They are much employed in Europe as demulcent and emollient, but their place is supplied in this country by other articles.

## DECOCTION OF MARSH MALLOW.

R. Marsh mallow dried	four ounces
Raisins, stoned,	two ounces
Water	seven pints

Boil down to five pints. Strain, and set aside until the dregs have subsided, and decant. *Dub. Ph.*

A good demulcent drink.

## SYRUP OF MARSH MALLOW.

R. Marsh mallow root	eight ounces
Sugar	two and a-half pounds
Water	four pints

Boil down the water with the root to one half, and press out the cooled liquor. Set by for dregs to subside, then decant, add the sugar, and boil to proper consistence.

*Lond. Ph.*

A demulcent, but not equal to the syrup of gum Arabic.

## MARSH MALLOW PASTE.

R. Marsh mallow root powdered	
Wheat flour, each,	three parts
White sugar	six parts
Mucilage of gum Arabic	two parts

Mix, and form paste. *Beral.*

A good demulcent in irritation of the throat.

## MARSH MALLOW LOZENGES.

R. Powdered mallow root	one part
White sugar	seven parts
Mucilage of tragacanth	sufficient

Form into lozenges of sixteen grains each. *Cottreau.*

## MARSH MALLOW OINTMENT.

R. Flaxseed	
Mallow root, each,	one pound
Water	sufficient

to make a thick mucilage; strain, and add

Lard, melted,	six pounds
---------------	------------

Evaporate, and add

Resin	eight ounces
Turpentine	
Yellow wax, each,	one pound

Melt the whole together. *Beral.*

## POWDER OF MARSH MALLOW.

R. Powdered mallow root	
liquorice root, each,	three ounces
nitre	half an ounce
camphor	one drachm

Mix, and divide into thirty powders; one to be taken three times a-day. *Foy.*

## ALUMINA.

## ALUMINE.

The hydrate of alumine is found naturally in an impure state in the form of different clays, boles, &c., and can be prepared in a pure state by the decomposition of alum.

## HYDRATE OF ALUMINE.

R. Alum	any quantity
Distilled water	sufficient

to dissolve the salt, add gradually a solution of carbonate of potassa, digest at a moderate temperature for a short time,



until a precipitate no longer takes place, wash this carefully, and dry. *Sax. Ph.*

## RUST'S ASTRINGENT.

R. Armenian bole  
Carbonate of magnesia  
White sugar, each, two drachms  
Oil of mace two drops  
Rhubarb one drachm  
Mix, and triturate well. *Radius.*  
Dose, a teaspoonful three times a-day, in chronic gonorrhœa, vaginal and uterine catarrh and slight menorrhagia.

## ELECTUARY FOR THE TEETH.

R. White bole an ounce and a half  
Gum lac half an ounce  
Cinnamon a drachm and a half  
Syrup of pinks sufficient  
to form soft paste. Triturate well together and keep in closed boxes. Used as a dentifrice, but also beneficial in scorbutic gums.

*Swedliaur.*

## ALUMINÆ ACETAS.

## ACETATE OF ALUMINE.

R. Alum a sufficient quantity  
Dissolve in water, precipitate by means of aqua ammonia, filter, and dissolve the moist precipitate in acetic acid, and again filter.

This has been recommended diluted with water in chronic diarrhœa, and mixed with syrup of poppies in slight cases of hæmoptysis.

*Van Mons.*

## ALUMINÆ SULPHAS.

## SULPHATE OF ALUMINE.

R. Alum a sufficient quantity  
Dissolve in water, precipitate by aqua ammonia, filter, and dissolve the moist precipitate in dilute sulphuric acid and again filter; evaporate and crystallize.

## DETERGENT WASH.

R. Sulphate of alumine two drachms  
Water half a pint  
Mix. As a wash to foul ulcers.

*Pennypacker.*

Mr. Gannal has found that a solution of this salt, made with about a pound to the quart of water, will preserve a body fresh for a long time, if injected into the blood-vessels; where it is only wished to preserve the body for a month or six weeks, an enema of one quart or an injection of about a quart into the œsophagus is sufficient.

*Dunglison, N. R.*

## ALUMINÆ ET POTASSÆ SULPHAS.

## ALUM.

Several varieties of alum are known in commerce, all containing more or less sulphate of iron. Most of that used in this country is of domestic manufacture.

Alum is employed both internally and externally as an astringent, in doses of ten grains to a scruple or more.

## BURNT ALUM.

R. Alum any quantity  
Melt in an iron or earthen vessel over a fire, and continue the heat till it ceases to bubble and becomes dry, then rub it into powder. *U. S. Ph.*

Astringent, and a mild escharotic. It is much used to repress the growth of exuberant granulations or proud flesh.

## POWDER OF ALUM.

R. Alum one drachm  
Gum Arabic half a drachm  
Triturate, and divide into four powders, one to be taken every three hours in atonic hemorrhagia. *Radius.*

R. Alum a drachm and a half  
Catechu one drachm  
Armenian bole one ounce  
Triturate well. To arrest hemorrhagia. *Van Mons.*

R. Alum four parts  
Kino one part  
Triturate well. Dose, ten to twenty grains, in hemorrhages or diarrhœa. *Landerer.*

R. Alum thirty grains  
Powdered opium three grains  
Mix, and make six powders, one to be taken every four hours. *A. T. Thomson.*

R. Alum one drachm  
Powdered opium four grains  
" cinnamon one scruple  
Mix and triturate, divide into four powders, one to be taken every four hours, in menorrhagia or in diarrhœa. *Radius.*

R. Alum two drachms  
Powdered opium three grains  
" kino one scruple  
Sugar of milk one drachm  
Mix, triturate, and divide into six powders. One to be taken every three hours, in fluxes. *Phœbus.*

## ESCHAROTIC POWDER OF ALUM.

R. Burnt alum  
Yellow oxide of iron  
Savine, each, equal parts



Triturate well. As an escharotic to venereal excrescences. *Swediaur.*

## ALUM ERRHINE.

R. Alum  
Armenian bole, each, one drachm  
Kino half a drachm  
Red oxide of iron two drachms  
Mix, and triturate. As an errhine in epistaxis. *Radiu.*

## GARGLES OF ALUM.

R. Alum  
Nitrate of potassa, each, three ounces  
Cream of tartar four ounces  
Acetic acid four pounds  
Dissolve, evaporate to dryness, and powder the residuum. Half an ounce, dissolved in eight ounces of water, forms a gargle which has been highly praised in inflammation of the fauces and tonsils. *Wirtem. Ph.*

R. Alum two drachms  
Water four fl. ounces  
Dissolve. Said to be useful where the breath is offensive. *Cavarra.*

## ODONTALGIC SOLUTION OF ALUM.

R. Finely powdered alum two drachms  
Spirit of nitric ether seven fl. drachms  
Dissolve. Stated to be an almost infallible cure for toothache. *Blake.*

## BOLUSES OF ALUM.

R. Alum  
Extract of Peruvian bark  
Nutmeg, each, half a scruple  
Simple syrup sufficient  
to make a bolus. Prescribed in uterine and other hemorrhages, and repeated as may be required. *Ellis.*

R. Alum five grains  
Extract of rhatany eighteen grains  
Conserve of rose hips half a drachm  
Astringent syrup sufficient  
to make ten boluses. Dose, one every three or four hours. *Foy.*

## ALUM PILLS.

R. Alum six grains  
Opium  
Catechu, each, one grain  
Mix, and form into six pills. One to be given every two to four hours. In passive hemorrhages and atonic mucous discharges. *Ellis.*

R. Alum two drachms  
Kino  
Honey of roses, each, one drachm

Make pills, of six grains each. Dose, one to six a day, in passive hemorrhages. *Foy.*

R. Alum  
Extract of Peruvian bark  
Muriate of iron and ammonia  
Aromatic powder, each, a drachm and a half  
Oil of cinnamon twelve drops  
Mix, and make pills of two grains. Dose, four to six, morning and evening, in passive hemorrhages and mucous discharges. *Augustin.*

## ASTRINGENT PILLS OF ALUM.

R. Alum  
Catechu equal parts  
Extract of gentian sufficient  
to make pills of two grains each. Dose, four every three hours. Useful in diarrhoea, and especially in leucorrhœa. *Radiu.*

## PILLS OF ALUM AND BENZOIN.

R. Alum one scruple  
Benzoic acid five grains  
Gum Arabic  
White sugar, each, ten grains  
Mix, with sufficient water to form thirty-five pills, to be taken in two days. Have been thought useful in phthisis pulmonalis. *Augustin.*

## ELECTUARIES OF ALUM.

R. Alum one drachm  
Catechu  
Extract of Peruvian bark, each, two drachms  
Conserve of roses six drachms  
Simple syrup sufficient  
to make an electuary. Dose, one drachm every four hours. In chronic diarrhoea, leucorrhœa, and passive hemorrhages. *St. Marie.*

R. Alum one drachm  
Extract of logwood half an ounce  
Balsam Peru six drops  
Water of sage sufficient  
Make an electuary. Astringent and antiseptic, recommended in sponginess of the gums. *Phœbus.*

## ALUM COLLYRIUM.

R. Alum twelve to twenty grains  
Rose water four fl. ounces  
Dissolve. As an application in chronic ophthalmia. *Ellis.*

## ALUM CURD.

R. Powdered alum half a drachm  
White of egg one



Agitate well till a coagulum is formed.  
To be applied on a rag to inflamed eyes.  
To be removed when it becomes warm.

*Ellis.*

#### SOLUTIONS OF ALUM.

R. Burnt alum  
Sulphate of zinc, each, half an ounce  
Distilled water two pints  
Dissolve.

R. Alum one drachm  
Sulphate of zinc half a drachm  
Borax four grains  
Rose water six ounces

Dissolve. Used as an astringent in bruises, slight hemorrhages, and mucous discharges.

*Cadet de Gassicourt.*

R. Alum half a drachm  
Armenian bole six drachms  
Vinegar

Red wine, each, half an ounce  
Mix the powdered bole to the solution of the alum in the vinegar, and add the wine.  
Used as a local application in epistaxis.

*Swediaur.*

#### GARGLES OF ALUM.

R. Alum one drachm  
Wine one pound  
Tincture of bark four drachms  
myrrh two drachms  
Honey of roses two ounces  
Laudanum one scruple

Mix. As a gargle in scurvy of the gums.

*Foy.*

R. Alum one drachm  
Infusion of red roses  
Barley water, each, three ounces  
Add to the solution  
Honey of roses two ounces

*Ratier.*

R. Powdered oak bark one ounce  
Boiling water a pint and a half  
Evaporate to one pint, filter, and add  
Alum half a drachm  
Brandy two fl. ounces

Used as a gargle in inflammation of the mouth and throat.

*Augustin.*

#### INJECTIONS OF ALUM.

R. Alum one drachm  
Rose water two ounces

Mix. In chronic gonorrhœa.

*Foy.*

R. Alum  
Sulphate of iron, each, twelve grains  
Honey of roses one ounce  
Barley water five fl. ounces

Mix. In chronic gonorrhœa.

*Radius.*

R. Stramonium half an ounce  
Boiling water two pints  
Infuse for half an hour, strain, and add to infusion

Alum two drachms to an ounce  
In cancerous affections of uterus.

*Foy.*

#### ALUM INJECTION.

R. Infusion of flaxseed fifteen fl. ounces  
Alum two drachms  
Tincture of kino one ounce

Mix. As an injection in cauliflower excrescence of the uterus.

*Clarke.*

#### JULEP OF ALUM.

R. Alum two drachms  
Sulphuric acid ten drops.  
Essence of citron six drops  
Syrup of lemon two ounces  
Water three ounces

Mix. A tablespoonful every hour, in lead colic.

*Radius.*

#### ALUM LINIMENT.

R. Alum half an ounce  
Whites of eggs four  
Spirit of camphor two ounces

Mix. As an application to bed sores.

*Augustin.*

#### LOTIONS OF ALUM.

R. Alum  
Sulphate of zinc, each, two drachms  
Plantain water two pints

Mix. As a wash to wounds and bleeding ulcers.

*Cadet de Gassicourt.*

R. Alum three drachms  
Muriate of ammonia one drachm  
Liquid hydrosulphuric acid

Rose water one scruple  
half a pint

Mix. As a wash, in obstinate cutaneous affections.

*Alibert.*

#### ALUM WHEY.

R. Cow's milk one pint  
Powdered alum a drachm and a-half  
Mix, separate the curd, and add to the whey

White sugar one ounce  
Dissolve. In passive hemorrhages. A wineglassful occasionally, to be taken cold.

*Foy.*

#### AROMATIC ALUM WHEY.

R. Powdered alum one drachm  
Sugar of milk half a drachm  
Powdered cinnamon fifteen grains  
Cow's milk a pint and a-half

Boil, and strain after coagulation. To be taken in cupful doses, cold, in hemorrhagia.

*Niemann.*



## ALUM DRAUGHTS.

R. Alum one to two drachms  
 Syrup of gum two ounces  
 Distilled water four fl. ounces  
 Mix. Dose, a wineglassful every six hours,  
 in lead colic and uterine hemorrhages.

*Guibourt.*

R. Alum two drachms  
 Wine four ounces  
 Gum fragacanth  
 Catechu, each, a drachm  
 Water eight ounces  
 Mix. One or two spoonfuls every hour  
 in lead colic.

*Radius.*

## ANGLO-SAXON OINTMENT.

R. Red lead  
 Olive oil  
 White wax, each, one pound  
 Powdered amber  
 Burnt alum  
 Camphor, each, two drachms  
 Heat the oil until it becomes of a reddish-brown color, add the red lead and continue the heat; when the mass has the consistence of a plaster, add the amber, and finally, when it cools, the alum and camphor. Highly spoken of as a dressing to foul ulcers.

*Niemann.*

## OINTMENT FOR CHILBLAINS.

R. Sweet almonds, blanched, half a pound  
 Honey six ounces  
 Camphor  
 Flour of mustard  
 Burnt alum  
 Olibanum, each, two ounces  
 Yolk of eggs three  
 Triturate well. This paste is to be thinned in a little water, and rubbed on the inflamed parts night and morning, and then washed off in tepid water, and afterward the parts well dried with a linen cloth.

*Swediaur.*

## RUST'S OINTMENT.

R. Alum a drachm and a-half  
 Camphor  
 Opium, each, a scruple to half a drachm  
 Balsam Peru one drachm  
 Lead ointment half an ounce  
 Triturate well together. Found useful as an application to frosted limbs.

*Phæbus.*

## PILE OINTMENT.

R. Powdered alum four scruples  
 Simple cerate five ounces and a-half  
 Mix, and triturate well. As an application to painful hemorrhoids.

*Taddei.*

## AMMONIA.

## AMMONIA.

This, under ordinary circumstances, exists as a gas, which is readily taken up by water or alcohol. Water absorbs nearly 800 times its volume of the gas at 32° F., and this solution possessing the properties of the gas is usually employed.

## WATER OF AMMONIA.

R. Pulverized muriate of ammonia  
 Quicklime, each, one pound  
 Distilled water one pint  
 Water nine fl. ounces

Break the lime in pieces, pour the water upon it in a proper vessel, cover this and permit it to rest until the lime slacks and is cold. Mix this with the muriate of ammonia in a mortar, and introduce the mixture into a glass retort. Place the retort on a sand-bath, and adapt to it a receiver connected by means of a glass tube with a quart bottle containing the distilled water. Apply heat, and continue as long as ammonia comes over. Remove the fluid in the bottle, and keep in small well-closed bottles. The *liq. ammon. fort.* has a sp. gr. of 0.880, and the usual *liq. ammoniæ* that of 0.960; one fl. ounce of the former with three of water will form the latter.

*U. S. Ph.*

## LINIMENT OF AMMONIA.

R. Water of ammonia half a fl. ounce  
 Olive oil two fl. ounces  
 Mix. Used as a rubefacient, and counter-irritant.

*U. S. Ph.*

## COMPOUND LINIMENT OF AMMONIA.

R. Stronger water of ammonia five fl. ounces  
 Spirit of camphor two fl. ounces  
 Spirit of rosemary one fl. ounce  
 Mix well. More powerful than the simple liniments.

*Edin. Ph.*

## GRANVILLE'S LOTION.

*(Milder.)*

R. Water of ammonia (.872) four fl. drachms  
 Spirit of rosemary three fl. drachms  
 Spirit of camphor (one  $\frac{3}{4}$  to Oj) one fl. drachm

Mix.

*(Stronger.)*

R. Solution of ammonia five fl. drachms  
 Spirit of rosemary two fl. drachms  
 Spirit of camphor one fl. drachm  
 Mix. As a counter irritant.

*Gray.*

## GARGLE OF AMMONIA.

R. Water of ammonia (.960) half fl. ounce



To be added to a strained decoction of  
 Mallows  
 Dry figs, each, six drachms  
 Cow's milk sixteen ounces  
 Employed as a resolvent in certain cases  
 of angina. *Pringle.*

## INJECTION OF AMMONIA.

R. Water of ammonia one scruple  
 Cow's milk one ounce  
 Mix. An an emmenagogue this injection  
 is to be used three times a-day. It induces  
 a mucous flow, often followed by the ap-  
 pearance of the catamenia. It must be  
 used with great caution, and made less  
 stimulating at first. *Lavagna.*

## MIXTURE OF AMMONIA.

R. Water of ammonia  
 Carbonate of potassa,  
 each, two drachms  
 Cinnamon water four ounces  
 Mix. A spoonful every two or three hours,  
 in cardialgia and acidity of the stomach.  
*Jourdan.*

## MIXTURE OF AMMONIA AND ETHER.

R. Water of ammonia  
 Sulphuric ether, each, two drachms  
 Peppermint water four ounces  
 Syrup of cinnamon one ounce  
 Mix. A spoonful every hour. Said to be  
 very useful in bites of venomous snakes.  
*Radiis.*

## AMMONIACAL SINAPISM.

R. Flour of mustard one ounce and a-half  
 Rye flour half an ounce  
 Water of ammonia sufficient  
 To form a cataplasm.  
 This is more active than the simple mus-  
 tard poultice. *Phœbus.*

## LOTION OF AMMONIA.

R. Water of ammonia  
 Spirit of thyme  
 Spirit of camphor equal parts  
 Mix. As a lotion to the forehead, or ap-  
 plied to the temples, in compresses, in sick-  
 headache. *Swediaur.*

## FETID SPIRIT OF AMMONIA.

R. Ammoniated alcohol two pints  
 Assafetida two ounces  
 Macerate for twelve hours. Then, by a  
 gentle heat, distil off a pint and a-half into  
 a cooled receiver. *Lond. Ph.*

Employed as a stimulant and antispasmodic,  
 in doses of half a drachm to a drachm.

## STEER'S OPODELDOC.

R. Alcohol eight pints

Soap twenty ounces  
 Water of ammonia four fl. ounces  
 Camphor eight ounces  
 Oil of rosemary  
 of monarda, each, one fl. ounce  
 Dissolve the soap in the alcohol with a  
 gentle heat, add the other articles, suffer  
 the impurities to subside, and pour into  
 vials whilst warm. *Phil. Coll. Pharm.*

## LIQUID OPODELDOC.

R. Dry Spanish soap sixty parts  
 Camphor fifteen parts  
 Alcohol five hundred parts  
 Oil of thyme four parts  
 of rosemary three parts  
 Solution of ammonia thirty parts  
 Mix, dissolve, and filter. *Giseke.*

## PLASTER OF AMMONIA.

R. Strong water of ammonia one ounce  
 Camphor two ounces  
 Opium fifteen grains  
 Ammoniac  
 Saffron  
 Gum plaster  
 Galbanum plaster, each, half an ounce  
 Mix well. For corns. A thick layer of  
 this plaster is to be spread on a piece of  
 linen or kid-skin, of the exact size of the  
 corn, as it will blister the sound skin.  
*La Foret.*

## SULPHURETTED AMMONIACAL LINIMENT.

R. Water of ammonia one ounce  
 Camphor half a drachm  
 Sulphuret of potassium four drachms  
 Olive oil six ounces  
 Mix. Recommended as an application in  
 tenia capitis. *Bories.*

TEREBINTHINATE AMMONIACAL LINI-  
MENT.

R. Oil of chamomile  
 of henbane  
 Essence of turpentine,  
 each, two fl. ounces  
 Water of ammonia  
 Tincture of capsicum, each, one ounce  
 Camphor half an ounce  
 Mix well. A very stimulating liniment,  
 found beneficial in asthenic gout and chro-  
 nic rheumatism. *Ammon.*

## AMMONIÆ ACETAS.

## ACETATE OF AMMONIA.

## SOLUTION OF ACETATE OF AMMONIA.

R. Diluted acetic acid one pint



Carbonate of ammonia a sufficient quantity  
Add the carbonate gradually to the acid, stirring constantly until all effervescence ceases. *U. S. Ph.*

A valuable diaphoretic in febrile and inflammatory diseases, when aided by warmth, otherwise acts as a diuretic. Dose, half a fl. ounce to a fl. ounce every three or four hours. It is also a good refrigerant lotion. It is known as the *spirit of mindererus*.

#### CATAPLASM WITH ACETATE OF AMMONIA.

R. Extract of hemlock one ounce  
of henbane half an ounce  
Powder of belladonna one drachm  
Solution of acetate of ammonia sufficient  
to form a cataplasm. Has been found useful in ulcerated cancerous sores. *Radius.*

#### COLLYRIUM OF ACETATE OF AMMONIA.

R. Solution of acetate of ammonia  
Rose water equal parts  
In chronic ophthalmia. *Saunders.*

#### GARGLE OF ACETATE OF AMMONIA.

R. Solution of acetate of ammonia  
Honey of roses, each, one ounce  
Elder water eight ounces  
Mix. Recommended by Wendt in scarlatina anginose with ulcerations. *Phœbus.*

#### MIXTURE OF ACETATE OF AMMONIA.

R. Solution of acetate of ammonia two fl. ounces  
Cinnamon water one fl. ounce  
Antimonial wine one fl. drachm  
Distilled water two fl. ounces  
Mix. A tablespoonful every two or three hours. *Ellis.*

R. Camphor mixture  
Solution of acetate of ammonia, each, half a fl. ounce  
Antimonial wine  
Laudanum, each, twenty drops  
Mix. To be taken at bedtime, or oftener, in which latter case the laudanum to be diminished. *Ellis.*

R. Liquid acetate of ammonia half fl. ounce  
Camphor mixture seven fl. drachms  
Antimonial wine fifteen drops  
Syrup of saffron half a fl. drachm  
Mix. To be taken three times a-day. *Burke.*

### AMMONIÆ ARSENIAS.

#### ARSENATE OF AMMONIA.

R. Arsenic acid one ounce  
Strong water of ammonia sufficient  
to saturate the acid. To be left to evaporate and crystallize spontaneously. *Cottureau.*

#### SOLUTION OF ARSENATE OF AMMONIA.

R. Arseniate of ammonia eight grains  
Water eight ounces  
Angelica water four drachms  
Mix. Has been recommended, in doses of twenty to thirty drops, in obstinate cutaneous affections. *Soubeiran.*

### AMMONIÆ CARBONAS.

#### CARBONATE OF AMMONIA.

R. Muriate of ammonia one pound  
Dried carbonate of lime one pound and a-half  
Pulverize them separately, then mix them thoroughly, and sublime into a cooled receiver. *U. S. Ph.*

#### AROMATIC CARBONATE OF AMMONIA.

R. Carbonate of ammonia one pound  
Oil of lavender three ounces  
verbena half an ounce  
Grind together, and sublime with a gentle heat.

R. Muriate of ammonia one drachm  
Caustic potassa three drachms  
Grind together, and add  
Oil of lemons fifteen drops  
*Cooley.*

R. Carbonate of ammonia half an ounce.  
Oil of peppermint  
cageput, each, twelve drops  
Mix, and keep in closed bottle. *Phœbus.*

R. Powd. carb. ammon. one ounce  
Stronger sol. of ammon. half a fl. ounce  
Oil of rosemary  
Oil of bergamot, each, ten drops  
Mix, and while moist, put in a wide-mouth bottle, which is to be well closed.

All these are used as smelling salts in faintness, &c.

#### POTION OF CARBONATE OF AMMONIA.

R. Carbonate of ammonia ten grains  
Cinnamon water one ounce  
Tincture of allspice half a drachm  
Syrup of saffron half an ounce  
Mix. To be taken before going to bed. *Phœbus.*



R. Comp. tinct. of  
 ammonia two fl. drachms and a-half  
 Camphor water seven fl. ounces  
 Tincture of castor one fl. drachm  
 Simple syrup three fl. drachms  
 Mix. Two spoonfuls in an hysteric attack.  
*Ainslie.*

## PILLS OF CARBONATE OF AMMONIA.

R. Carbonate of ammonia  
 Powdered capsicum  
 cloves  
 mace, each, one scruple  
 Oil of caraway five drops  
 Extract of gentian twelve grains  
 Simple syrup sufficient  
 to form twenty pills. One every two  
 hours in gout in the stomach. *Parrish.*

## MIXTURE OF CARBONATE OF AMMONIA.

R. Carbonate of ammonia  
 Powdered white sugar  
 gum Arabic, each, a drachm  
 and a-half  
 Compound spirits of  
 lavender two fl. drachms  
 Mint water four fl. ounces  
 Mix. A tablespoonful every two or three  
 hours. If required, forty to fifty drops of  
 laudanum may be added to the mixture.  
*Ellis.*

Useful in low states of the system.

## DRAUGHT WITH CARBONATE OF AMMONIA.

R. Valerian one scruple  
 Carbonate of ammonia ten grains  
 Cinnamon water two fl. ounces  
 Mix. To be taken every four hours, in  
 depression of the spirits and nervous head-  
 ache. *Ellis.*

MIXTURE OF CARBONATE OF AMMONIA  
AND GINGER.

R. Powdered ginger fifteen grains  
 Carbonate of ammonia eight grains  
 Spirit of cinnamon two fl. drachms  
 Water one and a-half fl. ounces  
 Mix. To be taken in gout or cramp in  
 the stomach. *Ellis.*

## DROPS OF CARBONATE OF AMMONIA.

R. Carbonate of ammonia twenty parts  
 Oil of lavender one part  
 Alcohol four parts  
 Distil. Dose, thirty or forty drops in  
 nervous headache. *Cottureau.*

## OINTMENT OF CARBONATE OF AMMONIA.

R. Carbonate of ammonia one part  
 Simple cerate eight parts

Mix. Used in frictions to the throat in  
 croup. *Foy.*

## PLASTER OF CARBONATE OF AMMONIA.

R. Carbonate of ammonia  
 Opium, each, fifteen grains.  
 Camphor half a drachm  
 Oil of cajeput twenty drops  
 Galbanum plaster half an ounce

Mix. Said to be useful as an application  
 to the pit of the stomach to arrest vomit-  
 ing in sea-sickness. *Phœbus.*

## CAMPHORATED AMMONIACAL LINIMENT.

R. Liquid carbonate of ammonia four parts  
 Oil of olives sixteen parts  
 Spirit of camphor three parts

Mix. As a revulsive embrocation.  
*Swediaur.*

## AMMONIACAL OINTMENT.

R. Carbonate of ammonia one drachm  
 Rose ointment one ounce  
 Oil of jasmine four drops

Mix. Recommended as a friction, three  
 or four times a-day, in engorgements of  
 the mammæ. *Augustin.*

## GONDRET'S AMMONIACAL OINTMENT.

R. Suet  
 Hog's lard, each, one part  
 Strong water of ammonia two parts  
 Melt the fats and incorporate the ammo-  
 nia. *Soubeiran.*

R. Hog's lard seven drachms  
 Oil of almonds a drachm and a-half  
 Strong water of ammonia five  
 drachms

Melt the lard, mix it with the oil, pour  
 into a wide-mouthed bottle with a glass  
 stopper, add the ammonia, agitate well.  
 Keep in a cool place.

Rubbed on the skin it causes rubefaction, and,  
 if covered with a compress, speedily vesicates.  
 A useful rubefacient and counter-irritant.

*Gondret.*

## COMPOUND TINCTURE OF AMMONIA.

R. Mastic two drachms  
 Alcohol nine fl. drachms  
 Oil of lavender fourteen drops  
 Oil of amber four drops  
 Strong water of ammonia one pint  
 Dissolve the mastic in the alcohol, and fil-  
 ter; add the other ingredients, and agitate  
 them well.

Well known under the name of *eau de luce* as  
 a remedy in bites of venomous reptiles. It is a  
 good antispasmodic, in doses of from ten to forty  
 drops; and also forms a stimulating embroca-  
 tion. *Lond. Ph.*



## AMMONIATED ALCOHOL.

R. Alcohol twenty fl. ounces  
 Quicklime one pound  
 Muriate of ammonia one pound  
 Water nine fl. ounces  
 Prepare as directed for water of ammonia.  
*U. S. Ph.*

## AROMATIC AMMONIATED ALCOHOL.

R. Ammoniated alcohol one pint  
 Oil of rosemary  
 lemons, each, two fl. drachms  
 cloves  
 cinnamon, each, half a  
 fl. drachm

Mix the oils, add the ammoniated alcohol, and pour in as much water as will prevent empyreuma in distilling; distil a pint with a gentle heat.  
*U. S. Ph., 1830.*

## MIXTURE OF AMMONIATED ALCOHOL.

R. Ammoniated alcohol half an ounce  
 Oil of amber one drop  
 mace two drops  
 aniseed  
 cloves  
 cinnamon, each, three drops

Mix. Dose, ten to thirty drops in sweetened gum water, in anorexia and flatus.

*Ammon.*

## ANISATED AMMONIATED ALCOHOL.

R. Alcohol twenty-four parts  
 Oil of aniseed one part  
 Dissolve, and gradually add, constantly stirring  
 Ammoniated alcohol six parts  
 Keep in well-closed bottles.  
*Jourdan.*

## LOTION IN HICCUGH.

R. Anisated ammoniated alcohol  
 one drachm  
 Spirits of lavender two ounces

Recommended by Siebold to be rubbed on the thorax and abdomen of infants to check hiccough.  
*Radius.*

## MIXTURE OF ANISATED AMMONIATED ALCOHOL.

R. Extract of liquorice one drachm  
 Barley water three ounces and a-half  
 Anisated ammoniated alcohol half a  
 drachm  
 Simple syrup half an ounce

Mix. A teaspoonful occasionally in chronic catarrh.  
*Hufeland.*

R. Anisated ammoniated alcohol  
 one drachm  
 Syrup of mallows one ounce  
 Extract of henbane one scruple  
 Mix well. A desertspoonful, three times

a day, in chronic bronchitis, or in the second stages of hooping-cough. *Radius.*

## VICAT'S ANODYNE MIXTURE.

R. Ammoniated alcohol half an ounce  
 Diluted alcohol one ounce  
 Opium two scruples  
 Camphor one scruple  
 Digest for three days, often stirring, and filter. Used on cotton to fill the cavity of carious teeth, and as a lotion to the temples in headache.  
*Spielman.*

## AMMONIÆ CITRAS.

## CITRATE OF AMMONIA.

R. Fresh lemon juice one fl. ounce  
 Carbonate of ammonia sufficient  
 to saturate  
 White sugar one drachm  
 Distilled water four fl. ounces

Mix.

A pleasant saline diaphoretic in febrile disorders, resembling the solution of the acetate of ammonia in its effects. Dose, a tablespoonful, as may be required.  
*Ellis.*

## AMMONIA HYDRIODAS.

## HYDRIODATE OF AMMONIA.

R. Iodine one hundred parts  
 Iron filings thirty parts  
 Water five hundred parts  
 Mix, and heat till the liquid becomes almost colorless; filter, and add carbonate of ammonia, as long as any precipitate takes place; filter, evaporate to dryness, redissolve in four or five times its weight of water; filter, evaporate, and crystallize.  
*Beasley.*

## OINTMENT OF HYDRIODATE OF AMMONIA.

R. Hydriodate of ammonia one scruple  
 Lard one ounce  
 Mix.  
*Beasley.*  
 R. Hydriodate of ammonia one drachm  
 Lard one ounce  
 Mix.  
*Gibert.*

As an application to scrofulous tumors and cutaneous affections.

## AMMONIÆ HYDROSULPHAS.

## HYDROSULPHATE OF AMMONIA.

R. Water of ammonia four fl. ounces  
 Pass hydrosulphuric acid (obtained from the sulphuret of iron and sulphuric acid



diluted with four times its weight of water) through the water of ammonia in a suitable apparatus, to perfect saturation. To be kept in a well-stopped bottle.

*U. S. Ph.*

A powerful sedative, lessening the action of the circulatory system to a great degree, and causing nausea, vomiting, vertigo, &c.

Used in diabetes to check the morbid appetite, by Cruickshank, Rollo, and others. Dose, five or six drops, in a tumbler full of water, three or four times a-day, gradually increasing the dose until some vertigo is induced.

#### SULPHURETTED HYDROSULPHATE OF AMMONIA.

*R.* Quicklime  
Sulphur, each, three parts  
Triturate together, adding water sufficient to form a paste, and incorporate

Sulphate of ammonia seven parts  
dissolved in water, let stand, decant, wash the residuum, rubbing it with a small portion of water, unite the solutions, and filter.

*Van Mons.*

Known as Boyle's fuming liquor, and recommended in venereal complaints, in phthisis, rheumatism, and gout, in doses of three to four drops in water, augmenting the dose until vertigo ensues.

#### WILKINSON'S LINIMENT.

*R.* Boyle's fuming liquor ten grains  
Chalk one scruple  
Flowers of sulphur  
Lard

Tar, each, half an ounce  
Mix well. Advised in chronic diseases of the skin. To be rubbed on a certain extent of the surface at a time.

*Phæbus.*

#### SULPHURETTED SYRUP.

*R.* Boyle's fuming liquor three drachms  
Sulphuret of antimony  
and soda one drachm  
Simple syrup an ounce and a-half  
Mix. Has been prescribed in croup, in the dose of a teaspoonful every hour.

*Swedjaur.*

### AMMONIÆ MURIAS.

#### SAL AMMONIAC.

##### POWDER OF SAL AMMONIAC.

*R.* Sal ammoniac two drachms  
Camphor six grains  
Arnica half a drachm  
White sugar six drachms

Mix. Dose, a teaspoonful, three or four times a day. Said to be efficacious as an emmenagogue, and also in smaller doses in chronic catarrh.

*Radius.*

#### BOLUS OF SAL AMMONIAC.

*R.* Sal ammoniac  
Extract of liquorice, each, one scruple  
Extract of taraxacum sufficient  
to form a bolus. One to be taken every three or four hours, in schirrus of the prostate, bladder, or rectum.

*Radius.*

#### LOTIONS OF SAL AMMONIAC.

*R.* Sal ammoniac two scruples  
Vinegar one ounce  
Water one pint  
Mix.

*Augustin.*

*R.* Sal Ammoniac one ounce  
Vinegar  
Alcohol, each, four ounces

Mix. In strains, bruises, and external inflammation.

*Foy.*

#### FOMENTATION OF SAL AMMONIAC.

*R.* Bistort  
Pomegranate, each, two ounces  
Red wine one pound  
Heat, digest for an hour, strain, and add

Sal ammoniac two drachms  
Useful as a fomentation, in strains and bruises.

*Foy.*

*R.* Sal ammoniac one ounce  
Spirit of rosemary one pound

Mix. Said to be useful in engorgement of the mammae. The breast to be covered with cloths dipped in the fluid.

*Foy.*

#### LINIMENT WITH SAL AMMONIAC.

*R.* Sal ammoniac one ounce  
Camphor half an ounce  
White soap six drachms  
Alcohol two pounds

Digest, and filter. Useful as an application to contusions.

*Niemann.*

#### COLLUTORIUM OF SAL AMMONIAC.

*R.* Pyrethrum two drachms  
Lavender water  
Vinegar, each, two ounces  
Opium two grains  
Sal ammoniac one drachm

Macerate for a few days, and filter.

*Cadet de Gassicourt.*

#### DRAUGHT OF SAL AMMONIAC.

*R.* Sal ammoniac five grains  
Camphor three grains  
Gum Arabic one drachm  
Parsley water four ounces

Mix. To be taken every two hours in paralytic retention of urine.

*Radius.*

#### CATAPLASM WITH SAL AMMONIAC.

*R.* Sal ammoniac  
Henbane, each, half an ounce



Linseed meal      an ounce and a-half  
Boiling water      sufficient  
to make a cataplasm. Advised in an engorgement of milk in the breasts.

*Radiis*

#### WASH OF SAL AMMONIAC.

R. Bran      an ounce and a-half  
Mallow leaves      six drachms  
Water      sufficient  
to make sixteen ounces of solution. Boil, strain, and add to decoction

Sal ammoniac      two drachms  
Tincture of benzoin      half an ounce

*Radiis.*

#### EMULSION OF SAL AMMONIAC.

R. Sweet almonds      one ounce  
Orange-flower water      two ounces  
Rose water      eight ounces  
Sal ammoniac      one drachm  
Tincture of benzoin      two drachms  
Rub down the almonds previously blanched, with the rose water, and add the other articles.

*Phœbus.*

### AMMONIÆ NITRAS.

#### NITRATE OF AMMONIA.

R. Diluted nitric acid      at will  
Carbonate of ammonia      sufficient  
to saturate.

Evaporate by a gentle heat, till a pellicle forms; set aside to crystallize. *Pereira.*

Soluble in two parts of cold water. In doses of one to two scruples, it reduces the frequency of the pulse and heat of skin, and also acts as a diuretic.

#### MIXTURE OF NITRATE OF AMMONIA.

R. Carbonate of ammonia      one scruple  
Nitric acid      sufficient  
to saturate; add  
Syrup of mallows      one ounce  
Distilled water      two ounces

Dose, a deserts spoonful, every two hours.

*Radiis.*

### AMMONIÆ PHOSPHAS.

#### PHOSPHATE OF AMMONIA.

R. Phosphoric acid      at will  
Saturate with

Carbonate of ammonia  
and set by to crystallize. *Niemann.*

Diuretic and discutient. Recommended by Dr. Buckler, of Baltimore, in gout and rheumatism, as a solvent of uric acid calculus, and in diseases dependent on a lithic acid diathesis. Dose, twenty to thirty drops of the saturated solution, three times a-day.

R. Powdered calcined bone      ten pounds  
Sulphuric acid      six pounds

Mix in a stoneware vessel, adding one gallon of water, digest for three or four days, frequently stirring, add a gallon of boiling water, strain, adding more boiling water, till it passes almost tasteless. Concentrate to one gallon, saturate with carbonate of ammonia, filter, concentrate by gentle heat, not above 100° F., and let crystallize. After removing each crop of crystals, add more ammonia, and concentrate. *C. Ellis.*

#### SOLUTION OF PHOSPHATE OF AMMONIA.

R. Phosphate of ammonia      half an ounce  
Water      six fl. ounces

Dissolve. Dose, a tablespoonful three times a-day. *C. Ellis.*

### AMMONIÆ SUCCINAS.

#### SUCCINATE OF AMMONIA.

R. Succinic acid      one part  
Water      four parts

Put in a capsule, and add sufficient water of ammonia, till perfectly saturated, with a slight excess of alkali, filter, evaporate, and crystallize. *Guibourt.*

#### IMPURE SUCCINATE OF AMMONIA.

R. Spirits of hartshorn      at will  
Saturate with a solution of succinic acid, filter, evaporate, and crystallize.

*Bavan Ph.*

#### SPIRIT OF SUCCINATE OF AMMONIA.

R. Spirit of hartshorn      two fl. ounces  
Sal ammoniac      four ounces  
Carbonate of potassa      twelve ounces  
Powdered amber      six ounces  
Oil of lemons      two fl. drachms

Mix the sal ammoniac, oil, and amber, add the carbonate of potassa to three pints of alcohol, mixed with a pint of water, and distil on a sand-bath. Dose, thirty to fifty drops, as an antispasmodic. *Niemann.*

#### MIXTURE OF SUCCINATE OF AMMONIA.

R. Liquid succinate of ammonia  
Elixir of liquorice, each,      two parts  
Wine of opium

Antimonial wine, each,      one part

Mix. Useful in whooping-cough, in doses of ten to forty drops, every two hours.

*Augustin.*

#### SPIRIT OF AMMONIA AND SUCCINIC ACID.

R. Succinic acid      one part  
Hot water      eight parts

Saturate the solution with carbonate of



ammonia, and filter. Antispasmodic, sudorific. *Van Mons.*

## AMMON. SUCCINIC ACID, AND ETHER.

R. Ammon. succinic acid  
Sulphuric ether equal parts  
Mix. Much praised in gout and rheumatism. Dose, twenty to forty drops in sugar and water, two or three times a-day. *St. Marie.*

## MIXTURE OF SUCCINATE OF AMMONIA.

R. Liquid succinate of ammonia five fl. drachms  
Tincture of castor  
Anisated ammon. alcohol, each, three fl. drachms  
Tincture of mace two fl. drachms  
nutmeg one fl. drachm  
Mix. Fifteen to twenty-five drops, three or four times a-day, in hysteria. *Augustin.*

## LINIMENT OF SUCCINATE OF AMMONIA.

R. Liquid succinate of ammonia one fl. ounce  
Camphorated oil two fl. ounces  
Mix. As a liniment to the neck in angina. *St. Marie.*

R. Camphor two scruples  
Oil of turpentine two drachms  
Dissolve, and add  
Soft-soap  $\times$  one ounce  
Ointment of bayberries two drachms  
Succinate of ammonia fifteen grains  
Mix. As a friction in chronic rheumatism, especially sciatica. *Cadet.*

## AMMONIÆ SULPHAS.

## SULPHATE OF AMMONIA.

R. Diluted sulphuric acid at will  
Carbonate of ammonia sufficient  
to saturate; filter, and crystallize. *Soubeiran.*

Laxative and stimulant. Dose, from a scruple to half a drachm.

AMMONIACUM.  $\times$ 

## GUM AMMONIAC.

This gum-resin is found in the shops, either in whitish or yellowish tears, or in yellowish or brownish masses. It has an unpleasant odor, and a nauseous and somewhat bitter taste. Is the product of *Ferula tingitana*, Linn.; and of *Dorema ammoniacum*, Don.

Pereira, Mat. Med. ii. 489. Griffith, Med. Bot. 325, 331.

Its effects on the system are somewhat like those of assafetida, and appear to be most

marked on the bronchial mucous membrane. It also is considered to be emmenagogue. The dose is from six to twelve grains. It is employed externally as a resolvent.

## AMMONIAC PILLS.

R. Ammoniac one drachm  
Syrup of gum sufficient  
to make thirty-six pills. *Soubeiran.*

R. Ammoniac  $\times$   
Myrrh, each, two drachms  
Extract of horehound one drachm  
liquorice three drachms

Mix. Make two grain pills. Dose, four to ten every three hours. *Augustin.*

R. Ammoniac one ounce  
Sagapenum two ounces  
Elixir proprietatis sufficient  
to make four grain pills. As a remedy in obstructed menstruation. *Bories.*

R. Ammoniac two drachms  
Soap one drachm  
Oil of juniper twenty drops  
Mix, and form pills of two grains. *Augustin.*

## KLEIN'S PILLS.

R. Ammoniac  
Extract of centaury, each, half an ounce  
Soap one ounce  
Oil of amber ten drops  
Triturate, and make pills of two grains. *Spielmann.*

## PILLS OF AMMONIAC AND RHUBARB.

R. Ammoniac  
Soap  
Rhubarb, each, two drachms  
Triturate, and make three grain pills.  
Dose, ten, three to four times a-day. *Tissot.*

## COMPOUND AMMONIAC PILL.

R. Ammoniac one drachm  
Blue pill fifteen grains  
Powdered squill six grains  
Simple syrup sufficient  $+$   
Mix, and make sixteen pills. One, three times a-day, in asthmatic cough with hepatic derangement. *Ainslie.*

## DE HAEN'S PILLS.

R. Ammoniac  
Pill. aloes and myrrh, each, one drachm  
Soap  
Extract of conium, each, a drachm  
and a-half  
Triturate, and form into three grain pills.  
Dose, four a day, in chlorosis and amenorrhœa. *St. Marie.*

## AMMONIAC MIXTURE.

R. Ammoniac two drachms



Water half a pint  
 Rub the ammoniac with the water gradually added, until they are thoroughly mixed.  
*U. S. Ph.*

The dose is from one to two tablespoofuls.

#### COMPOUND AMMONIAC MIXTURE.

R. Ammoniac one drachm  
 Distilled water four fl. ounces  
 Make a mixture, and add  
 Oxymel of squills two fl. ounces  
 Paregoric elixir half a fl. ounce

Mix. A teaspoonful every two hours, as a stimulating expectorant. *Ellis.*

R. Ammoniac mixture five fl. ounces  
 Oxymel of squill half fl. ounce  
 Antimonial wine twenty-six minims  
 Distilled vinegar three fl. drachms  
 Mix. A tablespoonful occasionally in cough, or humoral asthma. *Ainslie.*

R. Ammoniac  
 Assafetida  
 Soap  
 Powdered valerian  
 Flowers of arnica, each, two drachms  
 Tartar emetic eighteen grains  
 Mix, and make two grain pills. In doses of twenty or thirty, three times a-day. In amaurosis, arising from abdominal affections. *Richter.*

#### AMMONIAC AND NITRIC ACID MIXTURE.

R. Ammoniac two and a-half scruples  
 Nitric acid two fl. drachms  
 Water eight fl. ounces  
 Add the acid to the water, then gradually rub the gum with the acid compound, in a composition mortar.

A teaspoonful, two or three times a day. It is highly beneficial in chronic catarrh, especially in that of elderly persons. *Ellis.*

#### MIXTURES OF AMMONIAC.

R. Ammoniac mixture three fl. ounces  
 Tincture of castor half a fl. drachm  
 Syrup of tolu half a fl. ounce  
 Laudanum twenty to thirty drops  
 Cinnamon water one fl. ounce  
 Mix. A desertspoonful every three or four hours in troublesome cough. *Meigs.*

R. Gum ammoniac  
 Arabic, each, half an ounce  
 Infusion of hyssop six ounces  
 Syrup of capillaire two ounces  
 Rub the gums with the infusion until a solution is formed.  
 In tablespoonful doses occasionally in catarrhs. *Radius.*

R. Ammoniac mixture four ounces

Wine of ammoniac  
 Paregoric elixir, each, half an ounce  
 Syrup of tolu one ounce  
 Mix. In teaspoonful doses occasionally as an expectorant.

#### AMMONIAC PLASTER.

R. Ammoniac five ounces  
 Vinegar half a pint  
 Dissolve the ammonia in the vinegar and strain, evaporate in an iron vessel on a sand-bath to a proper consistence.  
*U. S. Ph.*

Useful as an application to indolent and scrofulous tumors to promote suppuration.

#### PLASTER OF AMMONIAC WITH MERCURY.

R. Ammoniac one pound  
 Purified mercury three ounces  
 Olive oil one fl. ounce  
 Sulphur eight grains  
 To the heated oil, add the sulphur gradually, stirring constantly, then rub the mercury with the compound, until the globules disappear; lastly, add the ammoniac previously melted, and with a gentle heat rub them well together. *Edin. Ph.*

More active than the simple ammoniac plaster. Used as a discutient to tumid glands and other indolent swellings, especially syphilitic. Sometimes salivates.

#### GUM PLASTER.

R. Litharge plaster four ounces  
 Ammoniac  
 Galbanum  
 Beeswax, each, half an ounce  
 Melt the gum resins together and strain, melt also the plaster and wax; add the two mixtures together and thoroughly incorporate. *Edin. Ph.*

A good digestive and suppurative for indolent tumors.

#### AMMONIAC PLASTER WITH HEMLOCK.

R. Strained ammoniac three ounces  
 Extract of hemlock two drachms  
 Melt, and add  
 Solution subacetate of lead one drachm  
 Stir till cold. *Ph. Chem.*

### AMYGDALA.

#### ALMONDS.

#### AMYGDALA AMARA.

#### BITTER ALMONDS.

#### AMYGDALA DULCIS.

#### SWEET ALMONDS.

The almond-tree, *Amygdalus communis*, is about the size and much resembles the peach-tree in



appearance. It is a native of Persia and Syria, and is extensively cultivated in the southern parts of Europe. There are several varieties, the most important of which are the sweet and the bitter: the former having a sweet, bland kernel; and the latter, one that has the bitter taste of the peach kernel.

*Sex. Syst.* Icosand. monog. *Nat. Syst.* Amygdalaceæ.

Linn. Sp. Pl. 677. Griffith, Med. Bot. 284.

Sweet almonds act as a demulcent; but the bitter have some of the properties of hydrocyanic acid.

#### BUTTER OF ALMONDS.

R. Blanched almonds ten drachms  
White sugar twelve drachms  
Orange-flower water two drachms  
Triturate till reduced to a homogeneous paste. *Guibourt.*

A spoonful, stirred in a tumbler of water, forms an extemporaneous syrup of orgeat, which latter preparation is difficult to preserve.

#### ALMOND PASTE.

R. Blanched almonds eight ounces  
Oil of almonds  
Lemon juice, each, a sufficient quantity  
Diluted alcohol five ounces  
Form a paste. *Taddei.*

R. Blanched almonds  
Rice flower, each, eight ounces  
Bean flower sixteen ounces  
Myrrh two ounces  
Camphor one ounce and a-half  
White soap twenty ounces  
Form a paste. *Spielmann.*

Used as a cosmetic.

#### ALMOND POWDER.

R. Blanched almonds two pounds  
Rice flower  
Orris root, each, four ounces  
Gum benzoin  
Carbonate of potassa  
Spermaceti, each, one ounce  
Oil of rhodium  
lavender  
cloves, each, thirty drops  
Form a powder. Used as a cosmetic.

#### ALMOND EMULSIONS.

R. Blanched almonds one ounce  
Pound in a marble mortar, and gradually add

Water sixteen ounces  
Strain. *Beral.*

R. Blanched almonds two ounces  
White sugar half an ounce  
Pound in a marble mortar, and add  
Boiling water twelve fl. ounces  
Strain, and add  
Orange-flower water two drachms

#### ALMOND CONFECTION.

R. Blanched almonds one ounce  
Powdered gum Arabic one drachm  
sugar half an ounce  
Pound well in a marble mortar till thoroughly mixed. *U. S. Ph.*

#### COMPOUND ALMOND LOTION.

R. Sweet almonds one ounce  
Bitter " four drachms  
Form an emulsion with  
Cherry brandy ten fl. ounces  
Add  
Corrosive sublimate six grains  
Tincture of benzoin six drachms  
Lemon juice four drachms  
Mix. As a lotion for pustular eruptions on the face. The mixture should be shaken before it is used. *Siemerling.*

#### SYRUP OF ORGEAT.

R. Sweet almonds one pound  
Bitter " five ounces  
White sugar six pounds  
Orange-flower water eight ounces  
Water three pints.  
Blanch the almonds, beat them into a paste with some of the water and the sugar, mix this paste with the rest of the water, press, and strain; dissolve the remainder of the sugar in the emulsion by means of a water-bath, add the orange flower-water, and strain. *Paris Cod.*

#### ARTIFICIAL MILK.

R. Blanched almonds No. 2  
Make an emulsion with  
Boiling water four ounces  
Cow's milk six ounces  
White sugar one drachm  
Strain. As a substitute for breast milk for infants. *Rosenstein.*

#### ANODYNE MILK.

R. Blanched almonds four ounces  
Melon seeds  
Squash seed, each, one ounce  
Lettuce seeds  
Poppy seeds, each, one drachm  
Triturate well, adding gradually  
Barley water sufficient  
Strain and express, to be taken in glassful doses, to which are to be added two drachms of sugar. *Pierquin.*

#### MILK OF ROSES.

R. Almonds one ounce  
Rose water five ounces  
Alcohol one ounce  
Soap half a drachm  
Attar of roses two drops



Blanch the almonds, dry them well with a cloth, beat them to a paste, add the soap, triturate well, then add the rose water and alcohol, rub well together, and strain through fine linen. This keeps well.

*Bateman.*

- |                       |               |
|-----------------------|---------------|
| R. Blanched almonds   | eight ounces  |
| Rose water            | three pints   |
| Windsor soap          |               |
| White wax             |               |
| Oil of almonds, each, | half an ounce |
| Rectified spirit      | twelve ounces |
| Oil of bergamot       | half an ounce |
| lavender              | one drachm    |
| Attar of roses        | half a drachm |

Beat the almonds with the rose water, so as to form an emulsion; melt the soap, wax, and almond oil together by a gentle heat, and rub the mixture with the emulsion, strain; dissolve the essential oils in the spirit, and mix with the strained emulsion.

*Redwood.*

#### GOULARD'S LOTION.

- |                       |                   |
|-----------------------|-------------------|
| R. Emulsion of bitter |                   |
| almonds               | two hundred parts |
| Corrosive sublimate   |                   |
| Sal ammoniac, each,   | one part          |

*Foy.*

#### ALMOND CREAM.

- |                   |            |
|-------------------|------------|
| R. Marrow pomatum |            |
| Almond oil, each, | two ounces |
- Melt by a gentle heat, and add  
Oil of jasmine two drachms  
stirring till cold. As an application to the hair.

*Bateman.*

#### OIL OF ALMONDS.

The kernels of almonds are to be rubbed and shaken in a bag to remove the reddish dust that adheres to them, then ground in a mill or pounded in a stone mortar. The marc is to be subjected to pressure in a linen bag and the oil filtered.

#### COLD CREAM.

- |                       |                |
|-----------------------|----------------|
| R. Rose water         |                |
| Oil of almonds, each, | two fl. ounces |
| Spermaceti            | half an ounce  |
| White wax             | one drachm     |
- Melt together the oil, spermaceti, and wax, by means of a water-bath, add the rose water, and stir constantly until the mixture is cold.

*U. S. Ph.*

#### ALMOND EMULSION.

- |                      |             |
|----------------------|-------------|
| R. Almond confection | one ounce   |
| Water                | half a pint |
- Rub the confection with the water gradually added; when thoroughly mixed, strain.

*U. S. Ph.*

A good demulcent, in irritation or inflammation of mucous membranes, and may be freely used.

#### AROMATIC ALMOND EMULSION.

- |                            |                 |
|----------------------------|-----------------|
| R. Blanched almonds        |                 |
| Simple syrup               |                 |
| Orange-flower water, each, | one ounce       |
| Water                      | fourteen ounces |
- Rub down the almonds with a little of the water, incorporate the paste with the remainder of the fluid, and strain.

*Beral.*

#### GREEN LINCTUS.

- |                     |                |
|---------------------|----------------|
| R. Pistachio nuts   | No. 14         |
| Syrup of violets    | one ounce      |
| Oil of almonds      | half an ounce  |
| Gum tragacanth      | sixteen grains |
| Tincture of saffron | twenty drops   |
| Orange-flower water | two drachms    |
| Water               | four ounces    |

Make an emulsion.

*Soubiran.*

Employed as a demulcent cough mixture.

#### EMOLLIENT CLYSTER.

- |                    |             |
|--------------------|-------------|
| R. Oil of almonds  |             |
| Brown sugar, each, | one ounce   |
| Barley water       | half a pint |
- Mix.
- Barley water a sufficient quantity  
Strain. To be used as an anodyne demulcent, in doses of a small tumblerful, adding to each two drachms of white sugar.

*Pierquin.*

#### BITTER ALMOND WATER.

- |                |           |
|----------------|-----------|
| R. Almond cake | one part  |
| Boiling water  | six parts |
- Mix well, and let stand for twelve hours, then distil one part with a moderate heat.

*Guibourt.*

Used as a flavoring ingredient, but with caution, as it contains hydrocyanic acid.

#### DIURETIC WATER.

- |                     |             |
|---------------------|-------------|
| R. Bitter almonds   | four ounces |
| Dried elder flowers | six ounces  |
| Peach kernels       | four ounces |
| Cherry kernels      | two ounces  |
| Wine                | eight pints |
- Digest for two days and distil off one half.

*Palat. Ph.*

Said to act as a diuretic in doses of half an ounce, but requires to be used with much caution.

#### OIL OF BITTER ALMONDS.

- |                   |                       |
|-------------------|-----------------------|
| R. Bitter almonds | twenty pounds         |
| Water             | a sufficient quantity |
- Pound the almonds, express the oil, let the cake macerate in water for twenty-four hours, distil as long as the product is



odorous, separate the oil, redistil the water, and again separate the oil. *Cottureau.*

Very poisonous, especially when fresh; principally employed as a flavoring ingredient, but should always be used with great caution.

LOCKSTADT'S PILLS.

R. Sulphate of quinine three grains  
Aromatic powder ten grains  
Oil of bitter almonds one drop  
Extract of gentian sufficient  
Make ten pills. The whole to be taken at a dose, before an expected paroxysm of intermittent fever. *Phœbus.*

WHITE LINCTUS.

R. Blanched almonds No. 12  
White sugar six drachms  
Gum tragacanth twelve grains  
Orange-flower water one drachm  
Water four ounces  
Make an emulsion *Guibourt.*

Used as a demulcent, in catarrhs and bowel complaints.

R. Blanched almonds four drachms  
White sugar six drachms  
Oil of almonds three drachms  
Orange-flower water two drachms  
Gum tragacanth sixteen grains  
Water four ounces  
Make an emulsion *Giordano.*

MIXTURES OF OIL OF ALMONDS.

R. Oil of almonds  
Simple syrup  
Water equal parts  
*Ratier.*

R. Oil of almonds one ounce  
Gum Arabic  
White sugar, each, half an ounce  
Water eight ounces  
Make an emulsion *Foy.*

R. Oil of almonds  
Gum Arabic  
Orange-flower water, each, half an ounce  
Syrup of mallows one ounce  
Water three ounces  
Make an emulsion. *Paris Ph.*

R. Oil of almonds one ounce  
Yolk of egg No. 1  
Syrup of mallows one ounce  
Orange-flower water two drachms  
Water two ounces  
Make an emulsion. *Guibourt.*

R. Oil of almonds  
Syrup of orgeat, each, seven drachms  
Gum Arabic two drachms  
Bitter almond water half an ounce

Make an emulsion. Dose, a teaspoonful. *Radius.*

R. Oil of sweet almonds  
Syrup of tolu, each, one fl. ounce  
Distilled water six fl. ounces  
Solution subcarbonate potassa sufficient

Make emulsion. A tablespoonful two or three times a-day in catarrh. *Ainslie.*

COSMETIC LINIMENT.

R. Oil of almonds one ounce  
Balsam of Mecca one drachm  
Sub. carb. of potassa half a drachm  
Rose water four fl. ounces  
Rub the balsam with the oil, add the potassa, triturate for ten minutes, and gradually introduce the rose water.

MACASSAR OIL.

R. Oil of almonds at will  
Alkanet root sufficient  
to color  
Oil of cinnamon sufficient  
to scent *Steph. and Church.*

FIXATURE.

R. Oil of almonds one ounce  
White wax one drachm  
Tincture of mastic three drachms  
Oil of bergamot one drachm  
Melt the oil and wax together by a moderate heat, then add the tincture of mastic and oil of bergamot, and strain.  
To fix and retain the hair in form. *Redwood.*

AMYLUM.

STARCH.

MUCILAGE OF STARCH.

R. Starch three drachms  
Water one pint  
Triturate the starch with the water gradually added, then boil for a few minutes. *Edin. Ph.*

Mainly employed as a demulcent injection in irritation of the rectum, &c., and forms a good vehicle for laudanum and other active remedies when given in enemata.

COMPOUND POWDER OF STARCH.

R. Liquorice root one ounce  
Starch  
Gum Arabic  
tragacanth  
Quince seeds  
Poppy seeds, each, two ounces  
Blanched almonds four ounces  
Sugar candy twelve ounces  
Triturate well. *Van Mons.*



Has been advised in spitting of blood, catarrh, and diarrhœa, in doses of half a drachm to a drachm.

#### STARCH LOZENGES.

R. Starch  
Orris root, each, half an ounce  
Extract of liquorice one ounce  
Saffron half an ounce  
White sugar one pound  
Mix, and form into lozenges. *Pideret.*

R. Starch one ounce  
Gum Arabic two ounces  
White sugar one pound  
Benzoic acid half a drachm  
Rose water sufficient  
to form a paste. To be divided into lozenges. *Swedjaur.*

These lozenges are somewhat stimulating, and are best suited to chronic catarrhs.

#### STARCH JELLY.

R. Starch six ounces  
Water six pounds  
Boil till reduced to three pounds, and add  
Wine four ounces  
Lemon juice one ounce  
Simple syrup two ounces  
Mix well, and permit to cool. *Swedjaur.*

A nourishing diet for convalescents.

#### MIXTURE OF STARCH AND SUET.

R. Cow's milk six pounds  
Mutton suet two ounces  
Boil by a gentle heat; constantly stirring, and add  
Starch a spoonful  
Boil a short time, and add  
White sugar a sufficiency  
Recommended in wineglassfuls in dysentery. *Pringle.*

### AMYLI IODIDUM.

#### IODIDE OF STARCH.

R. Iodine twenty-four grains  
Triturate with a little water, and add gradually

Powdered starch one ounce  
Continue the trituration till the mixture assumes a deep and uniform color.

*Buchanan.*

Dose, from half a drachm, to be gradually increased to four drachms. Given in cases requiring the use of iodine.

### ANEMONE.

#### ANEMONE

A genus of perennial herbs with radical leaves. They are all acrid and stimulating. The spe-

cies most generally employed, are the *A. pratensis* and *A. pulsatilla*, both natives of Europe, but it is probable that the *A. patens* of this country has the same properties.

The *A. pratensis* was at one time in much repute in amaurosis and chronic diseases of the skin.

*Sex. Syst.* Polyand. polygyn. *Nat. Syst.* Ranunculaceæ.

Lindley, Med. Flor. 2. Griffith, Med. Bot. 80.

#### EXTRACT OF ANEMONE.

R. Fresh anemone a sufficient quantity  
Pound in a mortar with a little water, express the juice, and evaporate in a water-bath. *Beral.*

The dose is from one grain to four, gradually increasing. It has been advised in doses of half a grain to a grain, in combination with sugar of milk, in whooping cough.

#### ANEMONE PILLS.

R. Powder of anemone one drachm  
Extract of anemone sufficient  
Make sixty pills. Dose, one to five, three times a-day, gradually increasing.

Said to have proved useful in amaurosis.

#### COLLYRIUM OF ANEMONE.

R. Anemone three drachms  
Water sufficient  
to make six ounces of infusion, add,  
Corrosive sublimate one grain  
To be dropped in the eye, in opacity of the cornea and incipient cataract. *Radius.*

#### WATER OF ANEMONE.

R. Fresh anemone four parts  
Water sufficient  
to prevent empyreuma. Distil six parts. *Van Mons.*

This is the best preparation, as it contains the active principle. It is somewhat caustic, and has been recommended to remove freckles on the skin.

### ANGELICA.

#### ANGELICA.

The only species of this genus, that is officinal in our Pharmacopœia, is *A. atro-purpurea*. This possesses much the same properties as the Garden Angelica, and is popularly used in flatulent colic and cardialgia. In Europe, the *A. officinalis* is generally used; and, as both species appear to possess the same properties, they may be indifferently employed.

*Sex. Syst.* Pentand. monog. *Nat. Syst.* Apiaceæ.

Torrey and Gray, Flor. 1, 64. Griffith, Med. Bot. 322.

#### COMPOUND TINCTURE OF ANGELICA.

R. Aniseed  
Angelica seeds, each, half a pound  
Proof spirit one gallon



Water sufficient to prevent empyreuma. Macerate for twenty-four hours, and distil a gallon.

A stimulant, used as a stomachic and carminative. Dose, half a drachm to four drachms. *Dublin Ph.*

#### TINCTURE OF ANGELICA.

R. Angelica root                      one part  
Alcohol                                  eight parts  
Water                                   sixteen parts  
Distil six parts. *Van Mons.*

Employed as a carminative and alexipharmic, in doses of one or two fl. drachms.

#### WATER OF ANGELICA.

R. Angelica root                      one part  
Water                                   six parts  
Distil three parts *Cottureau.*

#### ERRHINE MIXTURE.

R. Water of angelica  
orange flowers  
roses                                  equal parts  
Mix. *Pierquin.*

This has been recommended to destroy the fetid smell in ozæna.

#### CONSERVE OF ANGELICA.

R. Fresh root of Angelica           one part  
Water                                   eight parts  
Macerate for a few days, boil, clarify the decoction, and add  
Sugar                                  one part and a-half  
cook the root in the syrup, and preserve in this state, or dry. *Giordano.*

#### ESSENCE OF ANGELICA.

R. Angelica root                      one part  
Diluted alcohol                      eight parts  
Water                                   sixteen parts  
Distil off six parts. Stomachic and carminative. *Van Mons.*

### ANGUSTURA.

#### ANGUSTURA BARK.

This is recognized by the U. S. Pharmacopœia as the product *Galipea officinalis*, but is also obtained from one or more other species, all natives of South America. It is compact, in flat or rolled, but seldom quilled pieces, of a light gray color externally, and of a yellowish-fawn within. It is very fragile, with a resinous fracture. The odor is faint but disagreeable; the taste is bitter and somewhat aromatic, leaving a sensation of pungency. Its powder is pale yellow.

*Sec. Syst.* Heptand. monog. *Nat. Syst.* Rutaceæ. Hancock, Trans. Med. Soc. 1849. Griffith, Med. Bot. 192.

It is a stimulating tonic, in large doses, acting on the stomach and bowels. It is more used in chronic diarrhœa, and a debilitated condition of the stomach and bowels, than as a febrifuge.

#### INFUSION OF ANGUSTURA.

R. Bruised angustura bark    half an ounce  
Boiling water                      one pint  
Macerate for two hours in covered vessel, and strain. *U. S. Pharm.*

The dose is two fl. ounces, every two to four hours.

#### TINCTURE OF ANGUSTURA.

R. Powdered angustura bark   two ounces  
Proof spirit                          two pints  
Macerate for fourteen days, then filter. *Dub. Pharm.*

This contains the active principles of the bark, and may be given in the dose of one or two fl. drachms.

#### ELECTUARY OF ANGUSTURA.

R. Powdered angustura        half an ounce  
canella                              half a drachm  
Honey                                a sufficient quantity  
Make an electuary. *Foy.*

This has been employed with benefit in chronic diarrhœa and dysentery, in doses of about a drachm.

#### MIXTURE OF ANGUSTURA.

R. Infusion of angustura       six fl. ounces  
Cinnamon water                half fl. ounce  
Laudanum                          twenty drops  
Mix. Three spoonfuls a-day. *Radius.*

### ANISUM.

#### ANISE.

This, the fruit of *Pimpinella anisum*, a perennial plant, cultivated largely in some parts of Europe, and often met with in gardens in this country. The seeds are of a grayish-green color, of an ovate, compressed form, with five pale, narrow ridges. They are aromatic, and have a pleasant, sweetish taste.

*Sec. Syst.* Pentand. digyn. *Nat. Syst.* Apiaceæ. Linn. Sp. Pl. 379. Griffith, Med. Bot. 319.

Anise is an aromatic stimulant, and is used in various forms, in colic and dyspepsia, and to prevent the griping of some purgatives. The oil is officinal, but is much superseded by that of *Illicium anisatum*, which is almost identical.

#### SPIRIT OF ANISEED.

R. Anise, bruised,                      ten ounces  
Proof spirit                              one gallon  
Water                                      two pints  
Mix, and distil a gallon by a gentle fire. *Lond. Pharm.*

#### ANISE LOZENGES.

R. White sugar  
Anise water, each,                  two ounces  
Evaporate to the consistence of honey, and add a mixture of  
White sugar                              four ounces  
Oil of anise                              half a drachm



Pour on a cold marble slab. *Cottreau.*

Used as a carminative and anti-dyspeptic.

#### MIXTURE OF ANISEED.

R. Powdered anise  
fennel  
Extract of liquorice, each, one ounce  
Syrup of mallows twelve ounces  
Mix. *Hanover Pharm.*

Said to be an excellent carminative, especially for children, in doses of about a teaspoonful.

#### TINCTURE OF ANISEED.

R. Spirit of anise sixteen ounces  
Elixir proprietatis one ounce  
Aromatic confection half an ounce  
Pink flowers one drachm  
Filter, after sufficient maceration.

*Wirttemberg Pharm.*

Stomachic, carminative, and pectoral. Dose, one to two drachms.

#### OIL OF ANISE MIXTURE.

R. Oil of anise twelve minims  
Sugar one drachm  
Tincture of ginger two fl. drachms  
Peppermint water six fl. ounces  
Mix. Dose, two or three spoonfuls, as a carminative. *Ainslie.*

### ANTHEMIS.

#### CHAMOMILE.

Several species of *Anthemis* are employed in medicine, but the most important and the only one recognized in the U. S. Pharm., is *A. nobilis*. This is an herbaceous perennial, native of Europe, where it is also extensively cultivated, and is also to be found in gardens in this country. The part used is the flowers; these, as found in the shops, are large, roundish, of a yellowish white color, a peculiar somewhat balsamic odor, and a warm, bitter, aromatic taste.

*Sex. Syst.* Syngen. super. *Nat. Syst.* Astera-  
raceæ.

Linn. Sp. Pl. 1260. Griffith, Med. Bot. 401.

Chamomile is an aromatic bitter tonic, in small doses, but emetic in large ones; the usual mode of administration is in infusion.

#### INFUSION OF CHAMOMILE.

R. Chamomile half an ounce  
Boiling water one pint  
Macerate for ten minutes in a covered vessel, and strain. *U. S. Pharm.*

When cold, it is tonic in doses of one or two fl. ounces, several times a day. As an aid to the operation of emetics, it should be given in a tepid state, and largely.

#### COMPOUND DECOCTION OF CHAMOMILE.

R. Chamomile half an ounce  
Fennel seeds two drachms  
Water one pint

Boil for a short time, and strain.

*Dublin Pharm.*

#### EXTRACT OF CHAMOMILE.

R. Chamomile one pound  
Water one gallon  
Boil down to four pints, filter whilst hot, and evaporate on a water-bath to proper consistence. *Edin. Pharm.*

A mild, bitter tonic, in doses of ten to twenty grains.

#### MIXTURE OF CHAMOMILE.

R. Infusion of chamomile four ounces  
Syrup of orange flowers one ounce  
Ethereal tincture of valerian one drachm  
Mix. *Foy.*

This has proved useful in hysteria, and other nervous disorders, especially where the stomach is disordered. Dose, a dessertspoonful.

R. Extract of chamomile  
Gum Arabic, each, two drachms  
Chamomile water five ounces  
Tamarind pulp  
Syrup of chamomile, each, one ounce  
Mix. *Phæbus.*

Said to be useful in diarrhœa, in tablespoonful doses.

#### SYRUP OF CHAMOMILE.

R. Water of chamomile two pounds  
Syrup of mallows half a pound  
Extract of liquorice one ounce  
*Cadet de Gassicourt.*

Has been recommended in chronic cough and affections of the lungs, in one or two ounces, to be taken during the day.

R. Coarsely powdered chamomile one ounce  
Cold water twelve fl. ounces  
Coarsely powdered sugar twenty ounces

Make an infusion by displacement, of the chamomile and water. Remove the residue, and place the sugar in its place; on this pour the infusion, till entirely dissolved. Dose, a tablespoonful.

*E. Parrish.*

#### COMPOUND CHAMOMILE PILLS.

R. Assafetida one scruple and a-half  
Extract of chamomile one drachm  
Powdered rhubarb one scruple  
Make mass, and divide into thirty pills.  
Three, twice a-day in flatulent dyspepsia. *Ainslie.*

### ANTIMONIUM.

#### ANTIMONY.

A metal of a silvery blue color, very brilliant,



of a lamellar texture, brittle and easily pulverized. Metallic Antimony or Regulus of Antimony, is not officinal in our Pharmacopœia, but is the basis of a great number of very important preparations.

## ANTIMONII CHLORIDUM.

## CHLORIDE OF ANTIMONY.

R. Sulphuret of antimony           one part  
Muriatic acid                       five parts  
Dissolve by aid of a gentle heat, let stand, decant, evaporate, and then distil almost to dryness. *Soubairan.*

Used as a caustic, to destroy fungous flesh and to cauterize poisoned wounds.

## NITRO-MURIATIC OXIDE OF ANTIMONY.

R. Prepared sulphuret of antimony           twenty parts  
Muriatic acid                       one hundred parts  
Nitric acid                       one part  
Digest the sulphuret with the acids, mixed in a glass, avoiding the fumes, with a gradually increasing heat. Boil for an hour, and pour the liquor, when cooled and filtered, into a gallon of water. Wash the precipitate with plenty of water, till the latter no longer reddens litmus paper, then dry the oxide on bibulous paper. *Dub. Ph.*

R. Liquid chloride of antimony   at will  
Distilled water                       sufficient  
Agitate together, wash the precipitate that forms with cold water, and dry by a gentle heat. *Van Mons.*

This preparation, which is known as *Powder of Algaroth*, is of very uncertain action, and is not used, except in making tartar emetic.

## OINTMENT OF CHLORIDE OF ANTIMONY.

R. Liquid chloride of antimony  
Corrosive sublimate, each, one drachm  
Powdered savin                       two drachms  
Lard                                       six drachms  
Mix. To destroy venereal excrescences. *Radiis.*

## ANTIMONII SULPHURETUM.

## SULPHURET OF ANTIMONY.

The sulphuret or crude antimony is now seldom used as a medical agent, but is the basis of almost all the preparations of that metal. Some forms of exhibition are, however, still in favor, on the continent of Europe.

## PILLS OF SULPHURET OF ANTIMONY.

R. Sulphuret of antimony           one ounce  
Guaiacum                               two drachms  
Extract of fumitory a sufficient quantity

Mix, and make pills of two grains each.

*Radiis.*

Much employed in Germany, as an alterative in diseases of the skin.

R. Sulphuret of antimony half a drachm  
Extract of opium five to eight grains  
  guaiac                       two drachms  
Make fifty pills. *Foy.*

Advised in gouty affections, nine pills being given three times a-day.

## LOZENGES OF SULPHURET OF ANTIMONY.

R. White sugar one hundred and ninety-two parts  
Sulphuret of antimony eight parts  
Mucilage of gum Arabic one part  
Mix, and make lozenges of twelve grains, each containing half a grain of the sulphuret. *Beral.*

Much employed in Europe, as an alterative in diseases of the skin, in rheumatism, gout, &c.

## OINTMENT OF SULPHURET OF ANTIMONY.

R. White wax                               four ounces  
Olive oil                                       twelve ounces  
Melt together, and add  
Powdered charcoal                       two ounces  
Precipitated sulphur  
Sulphuret of antimony, each, one ounce  
Stir well until cold. *Radiis.*

Used as an application in tinea capitis.

## PRECIPITATED SULPHURET OF ANTIMONY.

R. Sulphuret of antimony six ounces  
Solution of potash four pints  
Distilled water  
Diluted sulphuric acid, each, a sufficient quantity  
Mix the sulphuret with the solution of potash and twelve pints of the water, boil by a gentle heat for three hours, frequently stirring, and adding more water, to keep up the original quantity. Strain, while hot; and gradually add the sulphuric acid as long as a precipitate is produced, wash well with hot water, dry, and powder. *U. S. Ph.*

## KERMES MINERAL.

R. Crystallized carbonate of soda 128 parts  
Water 1280 parts  
Sulphuret of antimony 6 parts  
Boil the antimony for an hour in the solution of soda, filter the hot liquor into warm earthen pans, cool very slowly, wash the precipitate with cold water, dry at a moderate temperature, and keep in well-closed bottles. *Paris Cod.*



## KERMESINE POWDER.

- R. Kermes mineral two grains  
 Sugar of milk  
 Gum Arabic, each, one drachm  
 Divide into six powders. *Radius.*

Given as a diaphoretic and expectorant, in the dose of one every four hours.

## COMPOUND KERMESINE POWDER.

- R. Kermes mineral half a drachm  
 Ipecacuanha one grain  
 Prepared chalk  
 Gum Arabic, each, twelve grains  
 Mix, and divide into three powders; one to be taken three times a-day, in spoonful of tea. *St. Marie.*

This formula has been much praised in whooping-cough; the proportions are intended for a child of about twelve years of age.

## LOZENGES OF KERMES MINERAL.

- R. Kermes mineral  
 Powdered squills, each, sixty grains  
 Extract of opium one drachm  
 Ipecacuanha two drachms  
 White sugar three ounces  
 Mucilage of tragacanth a sufficient quantity

Mix, and make four hundred lozenges.

*Foy.*

These are expectorant and calming in catarrh. One is to be taken every two hours.

- R. Kermes mineral one drachm  
 Powdered gum Arabic eight ounces  
 Extract of opium twelve grains  
 liquorice two ounces  
 Sugar thirty-two ounces  
 Oil of aniseed six drops  
 Mucilage of tragacanth sufficient

Mix, and form ten grain lozenges. Known as Tronchin's lozenges. Used in catarrh.

*Van Mons.*

- R. Benzoic acid two drachms  
 Sugar thirty-two ounces  
 Powdered orris root four drachms  
 gum Arabic two ounces  
 starch four ounces  
 Water four fl. ounces

Mix, and make fifteen grain lozenges.

*Vandamme.*

## EMULSION WITH KERMES MINERAL.

- R. Kermes mineral ten grains  
 Oxymel of squills one ounce  
 Emulsion gum Arabic four ounces  
 Mix. A spoonful occasionally. *Radius.*

## MIXTURE WITH KERMES MINERAL.

- R. Kermes mineral two grains  
 Gum Arabic one drachm

Syrup four fl. ounces  
 of grapes one fl. ounce

Mix. A spoonful occasionally in catarrh, when expectoration is difficult. *Pierquin.*

## GOLDEN SULPHURET OF ANTIMONY.

- R. Mother water of kermes mineral  
 add gradually  
 Acetic acid  
 till a precipitate is no longer formed, wash this well in cold water, and dry by a gentle heat. *Paris Cod.*

## POWDER OF GOLDEN SULPHURET OF ANTIMONY.

- R. Golden sulph. of antimony  
 Camphor, each, a grain and a-half to two grains  
 Nitrate of potassa a half to one drachm  
 Gum Arabic  
 White sugar, each, one drachm  
 Triturate well, and divide into six powders. *Radius.*

The dose is one, every two hours, in peripneumonia, after a reduction of the primary symptoms.

- R. Golden sulph. of antimony eight grains  
 Opium two grains  
 Powdered Peruvian bark

Chamomile, each, four scruples

Mix, and divide into eight powders; to be taken during the apyrexia. *Radius.*

## MIXTURE OF SULPHURET OF ANTIMONY.

- R. Golden sulph. of antimony six grains  
 Extract of lettuce one scruple  
 Syrup of senega two ounces  
 Mix. A teaspoonful every two hours.

*Phæbus.*

## SULPHURET OF ANTIMONY AND LIME.

- R. Golden sulph. of antimony one part  
 Quicklime three parts  
 Triturate together, and add  
 Boiling water twenty-four parts  
 Dry with a gentle heat, constantly stirring, and keep in a well-closed bottle.

*Batav. Ph.*

Emetic and resolvent. Much praised in gout, rheumatism, scrofula, &c. Dose, one to six grains.

## PILLS OF SULPHURET OF ANTIMONY AND LIME.

- R. Sulphuret of antimony and lime half a drachm  
 Mucilage of tragacanth one drachm  
 Mix, and make sixty pills; five, thrice a-day. *Augustin.*



PLASTER OF SULPHURET OF ANTIMONY  
AND LIME.

R. Yellow wax      one drachm and a-half  
Melt, and mix.

Pitch      one ounce  
and incorporate in the cooling mixture  
Sulphuret of antimony and  
lime      five scruples

As a plaster in sciatica, chronic head-  
ache, &c. It sometimes occasions pustules.

*Radiis.*

## ANTIMONII SULPHAS.

## SULPHATE OF ANTIMONY.

R. Antimony      two parts  
Sulphuric acid      three parts

Heat in an earthen vessel, stirring from  
time to time, leave the mixture on the fire  
till it has assumed a grayish-white color,  
wash carefully, to remove the excess of  
acid, and dry the powder. *Van Mons.*

ANTIMONII ET POTASSÆ  
TARTRAS.

## TARTAR EMETIC.

## POWDER OF TARTAR EMETIC.

R. Tartar emetic      one grain  
Sugar      thirty grains

Mix, and divide into ten powders, one  
every three or four hours, as a diaphoretic  
in fevers, after bleeding. *A. T. Thomson.*

## COMPOUND POWDER OF TARTAR EMETIC.

R. Tartar emetic      half a drachm  
Sulphate of potassa      one ounce  
Powdered liquorice      one ounce and a-  
half

Mix well. Two scruples contain one grain  
of tartar emetic. *Beasley.*

R. Tartar emetic      one grain  
Kermes mineral      two grains  
Powdered orris root      one scruple

Gum Arabic  
Sugar, each,      one drachm  
Mix, and divide into twenty-four powders,  
one every hour, as expectorant. *Cadet.*

POWDER OF TARTAR EMETIC AND  
QUININE.

R. Tartar emetic      three grains  
Sulphate of quinine      ten grains

Mix, and divide into six powders, one,  
every two hours, in apyrexia of obstinate  
intermittents. Both vomits and purges.

*Gala.*

POWDER OF TARTAR EMETIC AND PHOS-  
PHATE OF LIME.

R. Tartar emetic      one grain  
Phosphate of lime      thirty-two grains

Mix. To be divided into four powders;  
one to be taken in the evening, in chronic  
dysury, produced by checked perspiration.  
*Swediaur.*

POWDER OF TARTAR EMETIC AND IPECA-  
CUANHA.

R. Tartar emetic      two grains  
Ipecacuanha      two scruples  
Starch      half a drachm

Mix, and divide into three powders; one  
every quarter of an hour, till vomiting is  
induced. *Radiis.*

## BOLUS OF TARTAR EMETIC.

R. Tartar emetic      six grains  
Powdered Peruvian bark      six drachms  
Extract of juniper      sufficient

Mix, and make sixty boluses. To be taken  
in twenty-four hours in quartan intermit-  
tent. *Laennec.*

## PILLS OF TARTAR EMETIC AND CAMPHOR.

R. Tartar emetic      two grains  
Powdered opium      six grains  
Camphor      thirty-six grains  
Alcohol      three drops  
Conserve of roses      sufficient

Powder the camphor with the alcohol, add  
the other powders, mix and incorporate  
with the conserve, and divide into eleven  
pills. One every fourth hour as a diapho-  
retic. *Ellis.*

## PILLS OF TARTAR EMETIC AND OPIUM.

R. Tartar emetic  
Opium, each,      one grain and a-half  
Powdered tragacanth  
Conserve of roses, each,      sufficient

Mix, and make twenty-five pills. Two,  
night and morning, in chronic pulmonary  
catarrh. *Parrish.*

R. Tartar emetic      twelve grains  
Opium      ten grains

Crumb of bread  
Gum Arabic, each,      sufficient

Mix, and make pills of half a grain. Dose,  
one increased to three in chronic rheuma-  
tism. *Broussais.*

PILLS OF TARTAR EMETIC AND  
GUAIAIACUM.

R. Tartar emetic      one grain  
Powdered guaiacum      half a drachm

Pill of aloes and myrrh      half a drachm  
Molasses      sufficient

Mix, and make sixteen pills. *Beasley.*



## SOLUTION OF TARTAR EMETIC.

R. Tartar emetic four grains  
 Sugar one drachm  
 Distilled water four fl. ounces  
 Dissolve. A tablespoonful every ten or fifteen minutes till it operates. *Ellis.*

## SOLUTION OF TARTAR EMETIC AND SQUILL.

R. Tartar emetic two grains  
 Oxymer of squill half an ounce  
 Water two fl. ounces and a-half  
 Mix. One-half to be taken, and if it does not vomit in fifteen minutes, the remainder. *Augustin.*

## EMULSION OF TARTAR EMETIC.

R. Tartar emetic five grains  
 Camphor half a drachm  
 Blanched almonds  
 Syrup, each, one ounce  
 Water ten fl. ounces  
 Make an emulsion with the almonds and water, mix the camphor rubbed with a few drops of alcohol, add the tartar emetic; when dissolved, add the syrup; emetic and expectorant, according to the dose.

*Van Mons.*

## CLYSTER OF TARTAR EMETIC.

R. Tartar emetic one to two scruples  
 Tepid solution of gum Arabic one pint  
 To overcome obstinate constipation. To be used with much caution. *Chapman.*

R. Tartar emetic three to eight grains  
 Infusion of arnica twelve fl. ounces  
 Dissolve. In apoplexy and cerebral affections. *Foy.*

## LOTION OF TARTAR EMETIC.

R. Tartar emetic one scruple  
 Water one pint  
 Dissolve. In cutaneous affections.

*Pierquin.*

## LOTION OF TARTAR EMETIC AND CORROSIVE SUBLIMATE.

R. Tartar emetic one drachm  
 Corrosive sublimate five grains  
 Compound spirit of lavender one fl. drachm  
 Water one fl. ounce

Mix. Wet the finger with the solution, and rub on part for five or ten minutes; it will cause a pustular eruption in a few hours. *Hannay.*

## LOTION OF TARTAR EMETIC AND CAMPHOR.

R. Tartar emetic one drachm  
 Boiling water one pint  
 Tincture of camphor half fl. ounce

Mix. As a counter-irritant on the chest, in pulmonary complaints. *Augustin.*

## OINTMENT OF TARTAR EMETIC.

R. Tartar emetic one drachm  
 Lard one ounce

Mix. When applied to the skin occasions a pustular eruption. *Hannay.*

R. Tartar emetic two drachms  
 Sugar one drachm  
 Cinnabar five grains  
 Spermaceti ointment nine drachms

Mix. *Jenner.*

## COMPOUND TARTAR EMETIC OINTMENT.

R. Tartar emetic half a drachm  
 Sal ammoniac one drachm  
 Camphor twenty-five grains  
 Musk ten grains  
 Lard one ounce

Mix. As a counter-irritant, in chronic diseases of the liver. *Fabre.*

## PLASTER WITH TARTAR EMETIC.

R. Tartar emetic one part  
 Burgundy pitch seven parts

Mix, and spread. *Beral.*

Or, it may be made by sprinkling tartar emetic on a pitch, or adhesive plaster.

*Ellis.*

It takes some time for it to produce its effect, and the eruption that ensues is very irritating and painful.

R. Resin plaster one ounce  
 Resin four drachms  
 Venice turpentine three drachms  
 Melt together by a gentle heat, and when nearly cold, add

Powdered tartar emetic one drachm  
 To be applied to the nape of the neck in scarlatina in children, also in rheumatism.

*Niemann.*

## AMMONIACAL LINIMENT WITH TARTAR EMETIC.

R. Ammoniacal liniment one fl. ounce  
 Tartar emetic one drachm  
 Mix. To indolent tumors, &c. *Ellis.*

## ANTIMONIAL WINE.

R. Tartar emetic one scruple  
 Wine ten fl. ounces  
 Dissolve. *U. S. Ph.*

Dose, as emetic for children, from thirty drops to a fl. drachm, every fifteen minutes, till it operates.

## COLLYRIUM WITH ANTIMONIAL WINE.

R. Antimonial wine half fl. ounce  
 Rose water four fl. ounces  
 Balsam of Fioraventi two drops



Mix. Recommended in chronic ophthalmia. *Spielmann.*

#### MIXTURE OF ANTIMONIAL WINE AND AMMONIAC.

R. Gum ammoniac one fl. scruple  
Oxymel of squill half fl. ounce  
Antimonial wine one fl. drachm  
Syrup of liquorice one fl. ounce and a-half

Mix. A spoonful every two hours, as expectorant. *Phœbus.*

#### MIXTURE OF ANTIMONIAL WINE AND LAUDANUM.

R. Antimonial wine three fl. drachms  
Laudanum one fl. drachm

Mix. Fifteen drops, every half hour or hour in rheumatic diarrhoea. *Monro.*

#### MIXTURE OF ANTIMONIAL WINE AND BITTER SWEET.

R. Antimonial wine half fl. drachm  
Syrup of marsh mallow three fl. drachms  
Fennel water one fl. ounce and a-half  
Extract of bitter sweet fifteen grains  
Cremor Tartar forty-five grains

Mix. A teaspoonful every hour, in whooping cough in young children. *Phœbus.*

#### SYRUP OF ANTIMONIAL WINE.

R. Antimonial wine  
Sal ammoniac  
Gum ammoniac, each, two drachms  
Oxymel of squill one ounce  
Syrup of mallow half a pound

Mix. In pectoral affections, to facilitate expectoration and to keep the bowels open. A spoonful every hour. *Cadet.*

### PULVIS ANTIMONIALIS.

#### ANTIMONIAL POWDER.

R. Sulphuret of antimony, powdered,  
Horn shavings, equal parts  
Mix, and put them in a red-hot iron pot, and stir constantly till they acquire an ash-gray color, and no vapor arises. Pulverize the product, put it in a crucible with a perforated cover, and expose this to a heat gradually increased to whiteness, which is to be kept up for two hours. When cold, reduce the product to a fine powder. *Ed. Ph.*

Emetic and diaphoretic; in doses of three to eight grains, diaphoretic; in larger, emetic and purgative, but very uncertain in its operation.

#### JAMES' POWDER.

R. Tartar emetic one scruple

Prepared burnt hartshorn

Oxide of antimony, each, five scruples  
Mix, and divide into twenty-one grain powders. Said to be the formula by which the original powder was prepared by Dr. James. (*Vid. Phil. Journ. Pharm., vi. 282*)

#### PILLS OF ANTIMONIAL POWDER AND CALOMEL.

R. Antimonial powder ten grains  
Powdered opium  
Calomel, each, two grains  
Conserve of roses sufficient

Mix, and make four pills. Two to be taken at bed-time in acute rheumatism.

*Ellis.*

### APOCYNUM.

#### DOGSBANE.

Two species of this genus are officinal in the U. S. Pharm. *A. androsæmifolium* and *A. cannabinum*, possessed of much the same properties; these are emetic, cathartic, and sometimes diuretic, the latter is rather the most powerful.

*Sex. Syst.* Pentand. digyn. *Nat. Syst.* Apocynaceæ.

### A. ANDROSÆMIFOLIUM.

#### DOGSBANE.

Bigelow, Med. Bot. ii. 148. Griffith, Med. Bot. 449.

A native of most parts of the U. S. Emetic and diaphoretic. Dose, 40 grains; where it is wished to act on the skin, combined with a grain of opium.

### A. CANNABINUM.

#### INDIAN HEMP.

Griscom, Am. Journ. Med. Sci. xii. 55. Griffith, Med. Bot. 450.

Found in many parts of the United States. Emetic, in doses of fifteen to thirty grains.

#### DECOCTION OF INDIAN HEMP.

R. Root of Indian hemp half an ounce  
Water a pint and a-half  
Boil to a pint. *Griscom.*

This has been found useful in dropsy, in doses of one or two fl. ounces, two or three times a-day, acting as a hydragogue purgative.

#### EXTRACT OF INDIAN HEMP.

R. Powdered root of Indian hemp one pound  
Water one gallon  
Macerate for twenty-four hours, boil down one quart, strain, and evaporate to a proper consistence. Dose, three to five grains.



Acts like the decoction, but is not as efficient.

### ARALIA.

Most of the species of this genus are medicinal, but two only are officinal.

*Sex. Syst.* Pentand. pentag. *Nat. Syst.* Araliaceæ.

#### A. NUDICAULIS.

##### FALSE SARSAPARILLA.

Torrey and Gray, Fl. i. 646. Griffith, Med. Bot. 344.

A mild stimulating diaphoretic, the root has been employed as a substitute for sarsaparilla; best given in infusion.

#### A. SPINOSA.

##### ANGELICA TREE.

Torrey and Gray, Fl. i. 647. Griffith, Med. Bot. 345.

The bark is a stimulating diaphoretic; a decoction has been found useful in rheumatic, syphilitic, and cutaneous affections; as has also a tincture.

### ARCTIUM.

##### BURDOCK.

One species of this genus is officinal in our Pharmacopœia, the *A. lappa*.

*Sex. Syst.* Syngen. æqual. *Nat. Syst.* Asteraceæ. Linn. Sp. Pl. 1143. Griffith, Med. Bot. 411. (Lappa.)

This is a native of Europe, but generally naturalized in the United States. The parts used are the root and seeds. They are diaphoretic and diuretic, especially the latter.

##### INFUSION OF BURDOCK.

R. Burdock root                      one ounce  
Boiling water                      two pints

Infuse for six hours, and strain. *Cottrean.*

Used as a diaphoretic, in rheumatic pains, &c.

##### DECOCTION OF BURDOCK.

R. Burdock root                      two ounces  
Water                      three pints

Boil down to two pints, and strain. *Wood.*

### ARGENTUM.

##### SILVER.

A solid, white, brilliant, very ductile metal. It is wholly soluble in nitric acid. The only officinal preparation in our pharmacopœia is the nitrate; but several others are employed in Europe.

### ARGENTI CHLORIDUM.

##### CHLORIDE OF SILVER.

R. Solution of nitrate of silver      at will  
Add gradually a solution of common salt, wash well the resulting precipitate, and dry by a gentle heat in the shade.

##### POWDER OF CHLORIDE OF SILVER.

R. Chloride of silver                      one grain  
Powdered orris root                      two grains  
Triturate well, and divide into ten parts. *Taddei.*

Used in syphilis, in friction on the tongue.

##### CHLORIDE OF SILVER AND AMMONIA.

R. Liquid ammonia  
Saturate, by the aid of heat, with recently precipitated and well-washed chloride of silver. Filter whilst hot, and crystallize.

##### PILLS OF CHLORIDE OF SILVER AND AMMONIA.

R. Chloride of silver and ammonia                      one grain  
Powdered orris root                      two grains  
Conserve of linden flowers                      sufficient  
Mix, and make fourteen pills. *Serre.*

### ARGENTI CYANIDUM.

##### CYANIDE OF SILVER.

R. Nitrate of silver                      fifteen drachms  
Distilled water                      one pint  
Add to the solution  
Diluted hydrocyanic acid                      one pint  
Wash the precipitate well with distilled water, and dry by a gentle heat.

*U. S. Pharm.*

Used in syphilis, in doses of one-twelfth to one-tenth of a grain.

##### OINTMENT OF CYANIDE OF SILVER.

R. Cyanide of silver                      ten grains  
Lard                      one ounce  
Mix *Serre.*

Employed as an application to ulcerations of the cornea, irritable ulcers, &c.

### ARGENTI IODIDUM.

##### IODIDE OF SILVER.

R. Solution of nitrate of silver  
Add gradually a  
Solution of iodide of potassium  
Wash the precipitate, and dry by a gentle heat. *Patterson.*



ointment of Iodide of Silver.

R. Iodide of silver                   ten grains  
Lard                                   one ounce  
Mix.                                   *Serre.*

Used in the same cases as the ointment of the cyanide.

ARGENTI OXIDUM.

OXIDE OF SILVER.

R. Crystallized nitrate of silver one ounce  
Caustic potash                   seven ounces  
Water                               eighteen ounces

Dissolve the nitrate in two or three ounces of the water, and the potash in the remainder, mix the solutions, stir and collect the precipitate, wash it carefully, and dry.

*Duhamel.*

Used as a substitute for the nitrate, in doses of about half a grain twice a-day, in diseases of the stomach, of undue secretion, &c.

ointment of Oxide of Silver.

R. Oxide of silver                   twenty grains  
Lard                                   one ounce  
Rub well together.                   *Serre.*

Used as an application to irritable ulcers, &c.

ARGENTI NITRAS.

NITRATE OF SILVER.

R. Silver, in small pieces,       one ounce  
Nitric acid                       five fl. drachms  
Distilled water                   two fl. ounces

Mix the acid with the water, and dissolve the silver in the mixture, on a sand-bath, then gradually increase the heat, so as to dry the salt. Melt this in a crucible, and continue the heat till ebullition ceases, then immediately cast into moulds. *U. S. Ph.*

Used as a tonic and antispasmodic, in doses of a fourth of a grain, gradually increased.

SOLUTION OF NITRATE OF SILVER.

R. Nitrate of silver               one drachm  
Distilled water                   one fl. ounce  
Dissolve, and strain. Keep in the dark.

*London Ph.*

PILLS OF NITRATE OF SILVER.

R. Crystallized nitrate of silver one grain  
Crumb of bread                   one drachm  
Make sixteen pills.                   *Guibourt.*

Each pill contains a sixteenth of a grain of the nitrate.

R. Nitrate of silver               ten grains  
Opium                               four grains  
Extract of gentian  
liquorice, each,           a drachm  
and a-half

Make pills of a grain each, these each contain a twentieth of the nitrate. *Brera.*

R. Nitrate of silver               two grains  
Crumb of bread                   sufficient

Mix well, and divide into four pills. One to be taken every six hours.

Advised in chronic epilepsy and other spasmodic disorders. To prevent the blue tinge of the skin, so often produced by a continued use of the salt of silver, eight drops of diluted nitric acid in a fl. ounce of water, should be taken after each pill. *A. T. Thompson.*

MACKENSIE'S SOLUTION.

R. Nitrate of silver               one scruple  
Distilled water                   one ounce

Dissolve. *Mackensie.*

Used to wash the throat and fauces, in affections of those parts.

NITRATE OF SILVER COLLYRIUM.

R. Nitrate of silver               ten grains  
Laudanum                       half a drachm  
Distilled water                   one ounce

Mix. *Foy.*

Employed as a wash in purulent ophthalmia.

ANTI-EPILEPTIC MIXTURE.

R. Nitrate of silver               twelve grains  
Laudanum                       twenty drops  
Mucilage of Gum Arabic       one ounce

Mix. *Radius.*

Given in epilepsy, in doses of twenty drops, twice a-day.

HAIR DYE.

R. Silver                           two drachms  
Steel filings                   half an ounce  
Nitric acid                   an ounce  
Rain water                   eight ounces

Dissolve, and strain. To be applied by means of a close brush. *Bateman.*

INDELIBLE INK.

R. Carbonate of soda           half an ounce  
Distilled water               four ounces

Mix, and make mordant.

R. Nitrate of silver           five scruples  
Gum Arabic                   two drachms  
Sap green                   one scruple  
Distilled water               one fl. ounce

Mix, and make ink.

INDELIBLE INK WITHOUT MORDANT.

R. Nitrate of silver           one ounce  
Crystallized carbonate of  
soda                   one ounce and a-half

Tartaric acid               eight scruples

Strong solution of ammonia

two fl. ounces

Archil                   half fl. ounce

White sugar               four drachms

Powdered gum Arabic twelve drachms



Distilled water sufficient  
Dissolve the nitrate and carbonate in separate portions of the water, mix the solutions, collect and wash the precipitate on a filter, rub it whilst still moist in a porcelain mortar with the tartaric acid, till effervescence has ceased, add the ammonia to dissolve the tartrate of silver, then mix the archil, sugar, and gum, adding as much as will make six fl. ounces.

*Redwood.*

#### OINTMENT OF NITRATE OF SILVER.

R. Nitrate of silver two grains  
Lard two drachms  
Mix well. As an application in acute ophthalmia.

*Velpeau.*

R. Nitrate of silver four parts  
Lard thirty parts  
or, eight parts of the salt to same proportion of lard, or, twelve parts to thirty. Used as application to white-swelling.

*Jobert.*

R. Nitrate of silver one part  
Lard seven and a-half parts  
Mix, to smear bougies, in cure of gonorrhoea.

*Macdonald.*

#### COMPOUND OINTMENT OF NITRATE OF SILVER.

R. Nitrate of silver ten to twenty grains  
Zinc ointment half an ounce  
Balsam of Peru one drachm  
Triturate thoroughly together. To heal venereal ulcers, and to promote cicatrization.

*Fricke.*

R. Nitrate of silver three grains  
Solution of subacetate of lead, five drops  
Lard one drachm

Mix well. In chronic ophthalmia.

*Guthrie.*

### ARMORACIA.

#### HORSERADISH.

This is a species of *Cochlearia*, the *C. armoracia*. A native of many parts of Europe, and much cultivated both there and in this country, for the sake of its roots, so well known as a condiment. The root is the part used in medicine. It has a pungent odor, and a warm, acrid taste.

*Sex. Syst.* Tetrad. silic. *Nat. Syst.* Brassicaceæ.

*Linn. Sp. Pl.* 904. *Griffith, Med. Bot.* 131.

As a remedial agent, it is an active stimulant, and when applied to the skin, acts as a rubefacient.

#### INFUSION OF HORSERADISH.

R. Fresh horseradish, sliced,

Mustard seed, bruised, each, one ounce  
Boiling water one pint  
Macerate for two hours in a covered vessel, and strain.

*U. S. Ph.*

It is sometimes used in paralytic, scorbutic, and dropsical affections, attended with debility. The dose is two fl. ounces, three or four times a-day.

#### COMPOUND SPIRIT OF HORSERADISH.

R. Sliced horseradish  
Dried orange peel, each, twenty ounces  
Bruised nutmeg five drachms  
Proof spirit one gallon  
Water two pints

Mix, and distil a gallon by a gentle heat.

*London Ph.*

Employed as a diuretic, in dropsy with debility. Dose, one to four fl. drachms.

#### MIXTURE OF HORSERADISH.

R. Infusion of horseradish four ounces  
Syrup of cinchona one ounce  
Extract of fumitory one drachm

Mix.

*Foy.*

In spoonful doses, as an antiscorbutic.

#### COMPOUND HORSERADISH CATAPLASM.

R. Bruised horseradish six drachms  
Mustard seed one ounce  
Flaxseed meal two ounces and a-half  
Vinegar sufficient

Mix, and make cataplasm, to be applied to the feet as revulsive in diseases of head.

*Ainslie.*

#### COMPOUND INFUSION OF HORSERADISH.

R. Horseradish one ounce  
Boiling water one pint  
Infuse for an hour in a close vessel; on cooling, add

Simple syrup one ounce and a-half.

A wineglassful, several times a-day, in dropsy, especially when arising from a granular affection of the kidneys.

### ARNICA.

#### WOLFSBANE.

Several species of this genus are medicinal, but one only is officinal, *A. montana*, a native of Europe; but it is likely that the *A. nudicaule* of this country is possessed of the same powers.

*Sex. Syst.* Syngen. super. *Nat. Syst.* Asteraceæ.

*Linn. Sp. Pl.* 1245. *Griffith, Med. Bot.* 407.

Arnica is a stimulant, acting with much energy on the cerebro-spinal system, and is also an active irritant on the stomach and bowels. In Germany, where it is much employed, it is given to fulfil a variety of indications.

#### COMPOUND POWDER OF ARNICA.

R. Powdered arnica root  
serpentaria



sugar, each, two drachms  
Oil of peppermint ten drops  
Mix, and divide into sixteen powders.  
*Augustin.*

Dose, a powder, every two hours, in the diarrhoea complicating typhoid fevers.

## EXTRACT OF ARNICA.

R. Flowers of Arnica one part  
Water eight parts  
Alcohol one part  
Mix, and macerate for three days at a gentle heat, express, distil off the alcohol, and evaporate the residue to a proper consistence.  
*Wurtzburg Ph.*

Dose, ten grains to a scruple, in chronic rheumatism and paralysis.

## INFUSION OF ARNICA.

R. Flowers of arnica one ounce  
Boiling water one pint  
Infuse for half an hour, and strain.  
*Cottureau.*

Used in the same cases as the last.

## COMPOUND INFUSION OF ARNICA.

R. Flowers of arnica one drachm  
chamomile half an ounce  
Peppermint two drachms  
Boiling water nine ounces  
Mix, macerate, and strain.  
*Copenh. Ph.*  
Dose, one ounce.

## FOMENTATION OF ARNICA.

R. Flowers of arnica half an ounce  
Boiling vinegar sufficient  
To obtain six ounces, add  
Carbonate of ammonia two drachms  
*Radiu.*

Used as a warm fomentation in œdema of the scrotum.

## DECOCTION OF ARNICA.

R. Flowers of arnica one ounce  
Water three pints  
Boil till reduced to two pints, strain, and add  
Syrup of ginger two ounces  
*Swed.aur.*

Dose, one to two ounces, every two hours, in aphonia, paralysis, rheumatism, &c.

## FOMENTATION OF ARNICA AND RUE.

R. Flowers of arnica two ounces  
Rue one ounce  
Boiling water sufficient  
To obtain twelve ounces when strained, after one hour of maceration.  
*Radiu.*

Used as a fomentation in contusions.

## TINCTURE OF ARNICA.

R. Flowers of arnica one part

Alcohol sixteen parts  
Digest, express, and filter.  
*Beral.*  
Dose, thirty drops, several times a-day.

## ARSENICUM.

## ARSENIC.

Metallic arsenic is not employed in medicine, but several of its combinations are of much importance.

## ARSENICI IODIDUM.

## IODIDE OF ARSENIC.

R. Arsenious acid sixteen parts  
Iodine one hundred parts  
Heat in a glass alembic, and preserve the sublimation.  
*Magendie.*

R. Arsenious acid three parts  
Iodine ten parts  
Digest in

Water one hundred and nine parts until no odor of iodine is perceptible; decant, and evaporate to dryness, and sublime in a close vessel.  
*Plisson.*

Dose, one-tenth of a grain, gradually increased to a quarter, three times a-day. Given in cancer, and obstinate cutaneous affections.

## SESQUIODIDE OF ARSENIC.

R. Finely powdered metallic arsenic one part  
Iodine five parts  
Triturate together, and introduce into a small flask or retort; place this on a sand-bath, and apply gentle heat till liquefaction is produced. The vessel should be nearly full, to prevent the formation of much iodine vapor, and to enable the operator to bring the fixed mass in contact with every part of it, so as to include any iodine that may have been sublimed. If no iodine odor is perceptible, and the contents assume a reddish yellow color, and crystallize on the sides of the vessel, the operation is complete.  
*W. Procter.*

## IODIDE OF ARSENIC PILLS.

R. Iodide of arsenic one grain  
Extract of conium twenty grains  
Mix, and make ten pills.  
*Thompson.*  
Used in lepra, and obstinate eruptions.

## IODIDE OF ARSENIC OINTMENT.

R. Iodide of arsenic three grains  
Lard one ounce  
Triturate well.  
*Bielt.*

Employed in cases of lepra, and other cutaneous affections, but requires much caution.



LIQUOR ARSENICI ET HYD-  
RARGYRI IODIDI.SOLUTION OF IODIDE OF ARSENIC AND  
MERCURY.

DONOVAN'S SOLUTION.

R. Levigated metallic arsenic six grains,  
·08

Mercury fourteen grains ·82

Iodine forty-nine grains

Alcohol one fl. drachm

Rub together till dry, and has assumed a pale red color. Add eight fl. ounces distilled water, triturate for a short time, pour into a flask, and add half a drachm of hydriodic acid, made with two grains of iodine, and boil for a few moments. When cold, add sufficient distilled water to make up eight fl. ounces; filter. *Donovan.*

R. Sesquiodide of arsenic thirty-six grains

Biniodide of mercury thirty-four grains

Distilled water half a pint

Triturate the two iodides with half an ounce of the water, until they combine and dissolve, then add the rest of the water, and filter. *W. Procter.*

Each fl. drachm contains one-eighth of a grain of arsenic and one-fourth of a grain of protoxide of mercury. This preparation is said to be useful in obstinate cutaneous diseases, as lepra, lupus, &c. The dose is from fifteen minims to half a fl. drachm two or three times a-day. *Dunghison, N. R.*

DRAUGHT OF SOLUTION OF IODIDE OF  
MERCURY AND ARSENIC.R. Solution iodide mercury and arsenic  
two drachmsDistilled water three fl. ounces and a-  
half

Syrup of ginger half fl. ounce

Mix, and divide into four draughts. One to be taken night and morning; not to be given in a metallic spoon. *Donovan.*

## ARSENICI SULPHURETUM.

ORPIMENT.

R. Orpiment one ounce

Quicklime one pound

Starch ten ounces

Water sufficient

to form a soft paste. *Foy.*

This is spread on the part where it is desired to remove the hair, and washed off when it begins to dry. It is the *rusma* of the Turks.

POWDER OF ORPIMENT.

R. Orpiment one ounce

Quicklime twelve ounces

Jasmine powder ten ounces

Powder of palm soap four ounces

Mix well. *Morfit.*

This, when applied, is mixed with a little water.

Both these, although efficient for the purposes designed, are dangerous, and should never be used, except with extreme caution.

LOTION OF ORPIMENT.

R. Orpiment two drachms

Verdigris

Aloes

Myrrh, each, half a drachm

Rose water three ounces

White wine six ounces

Mix. *Van Mons.*

Used as an application to fungous ulcers.

GARGLE OF ORPIMENT.

R. Lanfranc's lotion half an ounce

Barley water six ounces

Honey of roses one ounce

Mix. *Giordano.*

As a gargle, in venereal sore throat.

## ARTEMISIA.

A genus of bitter herbs, several of which are used as medicinal agents to fulfil various indications; such as tonics, (*V. absinthium*,) some as moxas, (*A. moxa*,) and some as anthelmintics.

*Sex. Syst.* Syn. pol. super. *Nat. Syst.* Astéraceæ.

## 1. A. ABROTANUM.

SOUTHERNWOOD.

A native of the south of Europe and the Levant, and generally cultivated in gardens. Its odor is aromatic and peculiar, and its taste bitter. It is used as a tonic, antispasmodic, and vermifuge.

*Linn. Sp. Pl.* 1185. *Griffith, Med. Bot.* 406.

INFUSION OF SOUTHERNWOOD.

R. Southernwood six drachms

Boiling water one pint

Digest for two hours, and strain. Given in hysteria and in dismenorrhœa. *Taddei.*

CLYSTER OF SOUTHERNWOOD.

R. Southernwood half an ounce

Boiling water one pint

Digest for half an hour, and, to six or eight fl. ounces of strained infusion, add

Olive oil one ounce

Said to be effectual in cases of worms, and especially ascarides. *Radius.*



## 2. A. SANTONICA.

## SEMEN CONTRA.

Under this name are included several species, as the *A. santonica*, *contra*, *judaica*, &c. They are principally from Barbary and the Levant. The parts used are the flowers, buds, peduncles, &c. Semen contra is much employed in Europe as an anthelmintic, but is seldom used in this country. The dose in powder is from ten to thirty grains.

## POWDER OF SEMEN CONTRA.

R. Semen contra	
Extract of tansy, each,	six grains
Oxide of iron	four grains
Oil of valerian	one drop.
Mix.	<i>Augustin.</i>

## ELECTUARY OF SEMEN CONTRA.

R. Semen contra	ten grains
Sulphate of iron	four grains
Jalap	
Honey, each,	one scruple
Mix.	<i>Rosenstein.</i>

## ANTHELMINTIC MIXTURE OF SEMEN CONTRA.

R. Infusion of semen contra	four ounces
Syrup of senna	one ounce
Mix.	<i>Foy.</i>

## ARUM.

## INDIAN TURNIP.

Several species of Arum are used in medicine, but the only one officinal in the U. S. Pharm., is *A. triphyllum*; a native plant, with a large, perennial cormus; this is white, fleshy, and extremely acrid in a fresh state, but becomes edible and bland when dried, and kept for any time; and the fecula obtained from it resembles arrowroot, in appearance and uses.

*Ser. Syst.* Mon. polyand. *Nat. Syst.* Araceæ.  
Bigelow, Am. Med. Bot. 1, 52.

The powder of the dried root may be given in doses of ten grains, in an emulsion of gum Arabic, sugar, and water, several times a-day. It has been found useful in chronic catarrh, chronic bronchitis, rheumatism, &c.

## ASARUM.

There are two plants recognized by this name, one by the London College, and the other by the U. S. Pharm.

*Ser. Syst.* Dodecand. monog. *Nat. Syst.* Aristolochiaceæ.

## 1. ASARUM EUROPÆUM.

## ASARABACCA.

A native of Europe, of which the leaves and root are used. The first have a feeble aromatic odor, when rubbed, and a somewhat spicy taste;

the latter has a strong, penetrating odor, and an acrid, bitter, and nauseous taste; this is sometimes used abroad, as a stimulant emetic, and anthelmintic; the leaves, as the basis of various errhine powders.

Linn. Sp. Pl. 633. Griffith, Med. Bot. 527.

## COMPOUND POWDER OF ASARABACCA.

R. Dried leaves of asarabacca	one ounce
Lavender flowers	one drachm
Rub together to a fine powder.	

*Dublin Ph.*

Has been found useful as an errhine, in some cases of headache, toothache, &c. Five or six grains are to be snuffed up the nostrils at night.

## 2. ASARUM CANADENSE.

## WILD GINGER.

Bigelow, Med. Bot. 1, 150. Barton, Veg. Mat. Med., p. 85.

This species is common to most parts of the U. S. The leaves are said by Dr. Barton and Bigelow, not to be emetic, but other authorities state that a teaspoonful of this powder never fails to act on the stomach. Like those of the asarabacca, they are eminently errhine. The root is an aromatic stimulant, with active diaphoretic properties, and may be used as a substitute for serpentaria.

## INFUSION OF WILD GINGER.

R. Root of wild ginger	half an ounce
Boiling water	one pint
Digest for an hour in a covered vessel, and strain.	
In all cases where Virginia snake-root is indicated.	
Dose, one to two fl. ounces.	

## ASCLEPIAS.

*Ser. Syst.* Pentand. digyn. *Nat. Syst.* Asclepiadaceæ.

Many species of this large genus are employed in medicine, and it is probable that the whole of them are endowed with active properties.

## 1. A. TUBEROSA.

## PLEURISY ROOT.

A native of most parts of the United States. The root is large, and irregularly tuberous; the taste of it is bitter, nauseous, and somewhat acrid. It is an active diaphoretic, and, in large doses, purgative. It has attained much popular reputation in the treatment of diseases of the respiratory organs, and more especially of pleurisy. The dose of the powder is from a scruple to a drachm, several times a-day, but is better in infusion or decoction.

## INFUSION OF PLEURISY ROOT.

R. Bruised Pleurisy Root	one ounce
Boiling water	two pints



Digest for two hours. Dose, three or four fl. ounces, warm, to be given every two or three hours, until it operates.

## 2. A. INCARNATA.

### SWAMP SILK WEED.

A native of many parts of the U. S., usually growing in wet places. The root is the officinal portion. It is emetic and cathartic in full doses of half a drachm to a drachm, and expectorant and alterative in small ones.

## 3. A. SYRIACA.

### SILK WEED.

Very common in the U. S. The root appears to have much the same properties as the last, and also some anodyne powers.

## ASPARAGUS.

### ASPARAGUS.

Several species of Asparagus have been used in medicine, but the only one that is officinal, is the *A. officinalis*. This is a native of Europe, and is generally cultivated there and in this country.

*Sex. Syst.* Hexand. monog. *Nat. Syst.* Liliaceæ. *Linn. Sp. Pl.* 448. *Stokes, Bot. Mat. Med.* ii. 252.

The parts used are the roots and young shoots, (turiones.) These are diuretic, aperient, and deobstruent, and have been thought to exercise a sedative influence on the heart. Dr. Wood states (*Am. U. S. Dispens.* 1231) that the berries are more efficient than the shoots.

### EXTRACT OF ASPARAGUS SHOOTS.

R. Clarified juice of asparagus sufficient  
Evaporate by a mild heat to proper consistence.  
*Soubéiran.*

### EXTRACT OF ASPARAGUS ROOTS.

R. Roots of asparagus sufficient  
Bruise them, add sufficient water to cover them, express, strain, and evaporate by a mild heat.  
*Soubéiran.*

These extracts are given in doses of a scruple to a drachm, as diuretics.

### SYRUP OF ASPARAGUS SHOOTS.

R. Juice of asparagus shoots one pint  
White sugar two pounds  
Dissolve by means of a water bath, and strain.  
*Cottureau.*

Advised in pulmonary catarrh, too great action of the heart, in one to four ounces during the day.

### DECOCTION OF ASPARAGUS ROOTS.

R. Asparagus roots one ounce  
Water two pints

Boil, and strain.

*Radius.*

To be taken by cupfuls in dropsies.

### DIURETIC MIXTURE.

R. Asparagus roots one ounce  
Water one pint

Boil for an hour, strain, and add

Acetate of potassa twenty grains  
Honey of squills half an ounce

*Foy.*

Used as the last, and in the same diseases.

## ASSAFÆTIDA.

### ASSAFETIDA.

A gum resin or inspissated juice from an umbelliferous plant, a native of Persia, usually thought to be *Ferula assafetida*, but now shown by Dr. Falconer to be a *Narthex*, which, though similar to *Ferula*, is distinct from it. (*Royle, Mat. Med.* 407.)

*Sex. Syst.* Pentand. digyn. *Nat. Syst.* Apiaceæ. *Royle, Mat. Med.* 407. *Griffith, Med. Bot.* 326.

Assafetida as found in the shops is in masses of a whitish, reddish, or violet hue, composed of adhering tears. Odor, fetid and alliaceous; taste, bitter and somewhat acrid; forms an emulsion with water.

Is stimulant and antispasmodic, and also emmenagogue and anthelmintic. It is used in spasmodic and convulsive diseases, as hysteria, chorea, hooping-cough, flatulent colic, &c.

### ASSAFETIDA PILLS.

R. Assafetida one ounce and a half  
Soap half an ounce

Beat with water into a mass, and divide into two hundred and forty pills; each of which contains three grains of the gum resin.  
*U. S. Ph.*

R. Assafetida one drachm and a-half  
Powdered orris root

Mucilage of gum Arabic, each,

sufficient

Beat together, and divide into thirty pills. One to be given occasionally in the hooping-cough of children.  
*Kapp.*

R. Assafetida a drachm and a-half  
Powdered ginger half a drachm

Syrup sufficient

Mix, and make thirty pills. Three to be taken every three hours. Have been advised in palsy.  
*A. T. Thomson.*

### PILLS OF ASSAFETIDA AND LACTUCARIUM.

R. Assafetida three drachms  
Lactucarium two scruples and a-half

Mix, and divide into eighty pills. Two or three every hour in hooping-cough of adults.  
*Radius.*



## PILLS OF ASSAFETIDA AND IRON.

R. Assafetida  
Sulphate of iron  
Extract of chamomile, each,  
half an ounce

Mix well, and divide into one hundred and eighty pills. Two or more, three or four times a-day, in hypochondria and hysteria. *Syffret.*

R. Black oxide of iron half a drachm  
Assafetida a drachm and a-half  
Oil of tansy ten drops  
Extract of wormwood sufficient

Mix, and divide into ninety pills. Six to be taken three times a-day. Said to be useful against lumbrici and ascarides.

*Phœbus.*

## PILLS OF ASSAFETIDA AND MUSK.

R. Assafetida two drachms  
Musk  
Camphor, each, one drachm  
Ambergris half a drachm

Beat well together, and divide into one hundred pills. Two to be taken three times a-day. Said to have proved useful in angina pectoris, hysteria, &c.

*Sainte Marie.*

## PILLS OF ASSAFETIDA AND OPIUM.

R. Assafetida half an ounce  
Powdered opium  
“ ipecacuanha, each,

four grains  
Oil of peppermint eight drops  
Alcohol sufficient

Beat well together, and divide into one hundred and twenty pills. Ten to be taken three times a-day in chronic ischuria, consecutive on gonorrhœa.

*Radius.*

## ASSAFETIDA MIXTURE.

R. Assafetida two drachms  
Water half a pint  
Rub together, adding the water by degrees, till well mixed. *U. S. Ph.*

R. Assafetida one drachm  
Sugar six drachms  
Rose water five ounces  
Make an emulsion, and add  
Hoffmann's anodyne thirty drops  
Taken in spoonful doses, in spasmodic asthma, and hysteria. *Berles.*

## ASSAFETIDA MIXTURE.

R. Assafetida half a drachm  
Solution of acetate of ammonia,  
half fl. ounce  
Pennyroyal water three fl. ounces

Mix. One or two spoonfuls in whooping-cough. *Miller.*

R. Assafetida mixture  
five and a-half fl. ounces  
Compound spt. of lavender  
half fl. ounce

Aromatic spt. of ammonia  
two fl. drachms

Mix. Three spoonfuls a day in hysteria. *Ainslie.*

## ASSAFETIDA MIXTURE.

R. Assafetida two ounces  
Opium two grains  
Infusion of chamomile four ounces

Mix. As an injection in convulsive attacks. *Augustin.*

R. Assafetida half a drachm  
Yolk of egg one

Rub together, and gradually add  
Water eight ounces

Strain. To form two injections. Said to be useful in the whooping-cough of children. *Soubéiran.*

## MIXTURE OF ASSAFETIDA AND OXYMEL OF SQUILLS.

R. Assafetida one scruple  
Powdered digitalis six grains  
Oxymel of squills two ounces  
Linden-flower water four ounces

Rub well together. Said to be found useful in the dry cough consequent on deranged menstruation. *Pierquin.*

## MIXTURE OF ASSAFETIDA AND TOLU.

R. Assafetida one drachm  
Water four fl. ounces

Make an emulsion and add  
Tincture of balsam of tolu half a fl. ounce

“ opium forty to fifty drops  
Mix well. A teaspoonful every two hours, in whooping-cough, and a dessert-spoonful or more to an adult. *Ellis.*

## TINCTURE OF ASSAFETIDA.

R. Assafetida four ounces  
Alcohol two pints  
Macerate for fourteen days, and filter. *U. S. Ph.*

Has all the properties of the gum resin, but is more stimulating. Dose, a fl. drachm.

## ALKALINE TINCTURE OF ASSAFETIDA.

R. Assafetida  
Carbonate of potassa, each, two ounces  
Diluted alcohol one pint  
Macerate for three days, by a mild heat and filter. Anti-hysteria, &c. Useful in



nervous disorders. Dose, about twenty drops. *Wirtumberg Dis.*

#### AMMONIATED TINCTURE OF ASSAFETIDA.

R. Ammoniated alcohol sixteen ounces  
Assafetida one ounce

Macerate for twenty-four hours, in a closed vessel, and then distil sixteen ounces.

*Van Mons.*

Stimulant and anti-hysterical. Dose, five to sixty drops.

#### TINCTURE OF ASSAFETIDA AND SOOT.

R. Assafetida one drachm  
Wood soot two drachms  
Alcohol three ounces

Digest for six days, and filter. *Guibourt.*

Highly praised in hysteria, flatulence, and convulsions caused by dentition. Dose, ten to thirty drops in an appropriate vehicle.

#### ASSAFETIDA PLASTER.

R. Assafetida  
Lead plaster, each, one pound  
Galbanum  
Yellow wax, each, half a pound  
Alcohol three pints

Dissolve the assafetida and galbanum in the alcohol, by means of a water-bath; strain while hot, and evaporate to consistence of honey, add lead plaster and wax melted together; stir well, and evaporate to due consistence. *U. S. Ph.*

A good application to the stomach, in hysteria with flatulence, and to the chest in whooping-cough.

#### DEWEES' CARMINATIVE.

R. Carbonate of magnesia half a drachm  
Tincture of assafetida sixty drops  
" opium twenty drops  
Sugar one drachm  
Distilled water one fl. ounce  
Mix well. *Deweese.*

Very useful in the flatulent colic, diarrhoea, &c. of infants. Dose, twenty-five drops to a child of two to four weeks old, increasing the dose for those of a more advanced age.

#### PILLS OF ASSAFETIDA AND ALOES.

R. Assafetida  
Powdered aloes  
Soap, each, half an ounce  
Beat with sufficient water to form a mass; divide into one hundred and eighty pills.

*U. S. Ph.*

A good purgative in cases of costiveness with flatulence and debility of digestive organs. Dose, two to five.

#### TINCTURE OF ASSAFETIDA AND CASTOR.

R. Tincture of assafetida  
castor, each, one fl. ounce

Aromatic spirit of ammonia one fl. drachm

Mix. In hysteria, &c. Dose, about a teaspoonful. *Ellis.*

#### COMPOUND ASSAFETIDA MIXTURE.

R. Assafetida one drachm  
Peppermint water three fl. ounces

Make a solution, and add

Ammoniated tincture of valerian two fl. drachms

Tincture of castor three fl. drachms

Sulphuric ether one fl. drachm

Mix. Useful in hysteria, in doses of a tablespoonful, largely diluted, every second hour. *Ellis.*

#### ASSAFETIDA ENEMA.

R. Assafetida one drachm  
Hot water two ounces  
Triturate together. In flatulent colic, and against ascarides. *Foy.*

### ATROPIA.

#### ATROPINE.

Take recently dried root of belladonna, exhaust it by alcohol, 40° Cartier; add to the tincture a quantity of slaked lime equal to one-twentieth of the weight of the root. After twenty-four hours' contact, filter; acidify slightly with sulphuric acid, and again filter. Two-thirds of the alcohol is then removed by distillation. Evaporate the residue to one-twelfth of the weight of the root employed. Carbonate of potassa is then added until the liquid begins to be rendered opaque by a grayish-brown precipitate, carefully avoiding an excess. The liquid is again filtered, carbonate of potassa added, till it ceases to precipitate, and after twenty-four hours the precipitate is collected on a filter and dried. It is then dissolved in concentrated alcohol, treated with animal charcoal and filtered, evaporated, and crystallized. *Soubiran.*

#### SOLUTION OF ATROPIA.

R. Atropia (one, two, or three grains, as the strength may wish to be increased) one grain  
Nitric or acetic acid sufficient  
Water one fl. ounce  
Dissolve.

As an application to the eyes for dilating the pupil.

### AURANTIUM.

#### ORANGE.

This fruit is the product of the *Citrus Aurantium*, a native of Asia, but now cultivated in all warm climates; two varieties occur, the one with bitter, the other with sweet fruit; these are considered distinct by Risso.



*Sex. Syst.* Polydelph. icosand. *Nat. Syst.* Aurantiaceæ.

*Linn. Sp. Pl.* Griffith, *Med. Bot.* 165.

Almost every part is employed in medicine, but the rind of the fruit of the bitter kind only is official in the U. S. Pharm.

## AURANTII CORTEX.

### ORANGE PEEL.

#### POWDER OF ORANGE PEEL AND RHUBARB.

R. Powdered orange peel  
rhubarb

Bitartrate of potassa, each, one ounce  
Mix well. Dose, one or two teaspoonfuls  
a-day. In dyspepsia, with a costive condition of the bowels. *Radius.*

#### CONFECTION OF ORANGE PEEL.

R. Fresh orange peel, grated, one pound  
White sugar three pounds  
Beat together, till thoroughly mixed.  
*U. S. Ph.*

As a vehicle or adjunct to powders.

#### ELECTUARY OF ORANGE PEEL.

R. Orange peel half an ounce  
Willow bark two drachms  
Syrup of orange peel sufficient  
Make an electuary. To be taken in divided doses, during the apyrexia.  
*Phœbus.*

#### TINCTURE OF ORANGE PEEL.

R. Dried orange peel three ounces  
and a-half  
Proof spirit, (Imp. meas.) two pints  
Digest for fourteen days, and filter.  
*Lond. Ph.*

Used principally as an addition to infusions, decoctions, &c.

#### SPIRIT OF ORANGE PEEL.

R. Orange peel three ounces  
Alcohol sixteen ounces  
Water four ounces  
Digest for four days, and distil sixteen parts.  
*Cottureau.*

Used for the same purposes as the last.

#### ELIXIR OF ORANGE PEEL.

R. Orange peel one ounce  
Cascarilla half an ounce  
Water of citron peel  
wormwood  
Alcohol, each, half a pint  
Digest for four days, and filter. *Moscatti.*

Dose, from four to six fl. drachms, as a carminative and stomachic.

### INFUSION OF ORANGE PEEL.

R. Orange peel No. 3  
Boiling water three pints  
Infuse for half an hour.

### COMPOUND INFUSION OF ORANGE PEEL.

R. Dried orange peel half an ounce  
Fresh lemon peel two drachms  
Bruised cloves one drachm  
Boiling distilled water one pint  
Macerate for a quarter of an hour, and strain.  
*Lond. Ph.*

These two preparations are used as stomachics, in doses of one or two fl. ounces.

### SYRUP OF ORANGE PEEL.

R. Bruised orange peel two ounces  
Boiling water one pint  
Macerate for twelve hours, strain, and add  
Sugar two pounds and a-half  
Make syrup. *U. S. Ph.*

Used principally as an agreeable drink mixed with water.

R. Recently dried sweet orange  
peel two ounces  
Powder, and put in a displacement apparatus, and pour on it a mixture of two parts alcohol, and one of water, until six fl. ounces are obtained. Pour this on thirty-two ounces (av.) of coarsely powdered sugar, and spread on paper, until the alcohol has evaporated. Then form it into a syrup, with sixteen ounces of water, merely carrying the heat to the boiling point, strain, and bottle whilst hot.  
*W. Procter.*

## AURANTII FOLIA.

### ORANGE LEAVES.

#### ELECTUARY OF ORANGE LEAVES.

R. Orange leaves  
Valerian, each, half a drachm  
Syrup of orange peel sufficient  
Make an electuary. *Radius.*  
A teaspoonful occasionally as an antispasmodic.

#### JELLY OF ORANGE LEAVES.

R. Orange leaves an ounce and a-half  
Boiling water two pints  
Digest for twenty-four hours, strain, and add  
Powdered salep two drachms  
Boil to a jelly, and add  
Wine  
Syrup of orange flowers, each, one ounce  
In teaspoonful doses, in marasmus.  
*Radius.*



## AURANTII FLORES.

## ORANGE FLOWERS.

## CONFECTION OF ORANGE FLOWERS.

R. Orange flowers                      one part  
Syrup                                      three parts  
Evaporate to proper consistence. *Taddei.*

## PASTILLES OF ORANGE FLOWERS.

R. Sugar  
Orange-flower water, each, two ounces  
Dissolve, and evaporate to consistence of honey; add a mixture of  
Sugar                                      four ounces  
Oil of orange flowers              half a drachm  
Make pastilles. *Cottureau.*

## ORANGE-FLOWER WATER.

R. Orange flowers                      ten parts  
Water                                      thirty parts  
Distil off twenty parts. *Guibourt.*

## SYRUP OF ORANGE FLOWERS.

R. Orange-flower water              one part  
Sugar                                      two parts  
Dissolve in a close vessel, and evaporate to proper consistence. *Soubeiran.*

## OIL OF ORANGE FLOWERS.

R. Orange flowers                      one part  
Water                                      three parts  
Distil, and separate the oil that floats on the product. *Taddei.*

## AURUM.

## GOLD.

Gold is used as a remedial agent in its pure state, but only in an infinitely divided form, in which it is stated that it acts on the system, and to be equally efficacious as its salts, but its action is much milder. It has been prescribed in various forms of syphilitic diseases, in doses of a quarter of a grain to a grain, three or four times a-day.

## POWDER OF GOLD.

R. Gold leaf                              one part  
Sulphate of potassa              eight to ten parts  
Triturate thoroughly, wash out the sulphate of potassa, and sift through fine gauze. *Par. Cod.*

To be applied in friction on the gums and tongue.

R. Dilute solution nit. mur. of gold  
at will  
Solution sulphate of iron              sufficient  
to precipitate; collect this, and dry. *Tromsdorff.*

R. Pulverized gold                      six grains  
Starch                                      sixty-six grains

Mix, and divide into twelve powders. One to be given four times a-day. *Riecke.*

R. Pulverized gold  
Lycopodium, each,              two grains.  
Mix. To be rubbed, during the day, on the tongue. *Riecke.*

## POWDER OF GOLD AND IRON.

R. Pulverized gold                      two scruples  
Clean iron filings                      two scruples  
Powdered gum Arabic              half a drachm  
Mix. To be given at once in water, acidulated with a few drops of diluted sulphuric acid, as an antidote to corrosive sublimate. *Buckler.*

## SYRUP OF GOLD.

R. Pulverized gold                      twenty-four grains  
Syrup                                      one ounce  
Mix. As an application to chancres on the fauces. *Bories.*

## OINTMENT OF GOLD.

R. Pulverized gold                      one drachm  
Lard                                      half an ounce  
Triturate well. As a dressing to venereal ulcers, and also to a surface having the cuticle removed; when the surface becomes dry, substitute the ointment of chloride of gold. *Legrand.*

## AURI CHLORIDUM.

## CHLORIDE OF GOLD.

R. Leaf gold                              one part  
Nitro-muriatic acid                      three parts  
Dissolve by aid of a sand-bath, and evaporate till vapors of chlorine are given off, and set aside to crystallize. Dose, one-twentieth to one-sixteenth of a grain. *Par. Cod.*

R. Pure gold                              one part  
Nitro-muriatic acid, (made with three parts muriatic acid, one part nitric acid, and one of water,)              two parts  
Put the gold in a porcelain capsule, and add the acid; cover with a plate of glass, and place on a salt-water bath, continue heat till nitrous fumes are no longer evolved. Raise the glass by means of a fold of paper, and continue heat, till, on introducing a glass rod, the adhering chloride of gold becomes solid on cooling. Remove from fire, and let crystallize. *Leval.*

## POWDER OF CHLORIDE OF GOLD.

R. Chloride of gold                      one part  
Powdered orris root                      two parts



Mix. In frictions on the gums, in the dose of a fifteenth to a tenth of a grain.

*Foy.*

**BOLUS OF CHLORIDE OF GOLD.**

R. Chloride of gold half a grain  
Extract of aconite twelve grains  
Mix, and make six boluses. One to be taken three times a-day. *Brera.*

**PILLS OF CHLORIDE OF GOLD.**

R. Chloride of gold ten grains  
Powdered liquorice three drachms  
Syrup sufficient  
Mix, and make one hundred and fifty pills. One, thrice a-day. *Ellis.*

**TINCTURE OF CHLORIDE OF GOLD.**

R. Leaf gold one drachm  
Nitro-muriatic acid one ounce  
Dissolve, and add  
Oil of rosemary two ounces  
Alcohol eight ounces  
Mix. Dose, ten drops. *Spielmann.*

**CAUSTIC OF CHLORIDE OF GOLD.**

R. Chloride of gold one ounce  
Nitro-muriatic acid one fl. ounce  
Mix. Used as a caustic to the surface of cancerous ulcers, applied by means of a dossil of lint. *Recamier.*

**COLLYRIUM OF CHLORIDE OF GOLD.**

R. Chloride of gold two grains  
Distilled water six fl. ounces  
Dissolve. As a collyrium in ophthalmia. *Jahn.*

**OINTMENT OF CHLORIDE OF GOLD.**

R. Chloride of gold eight grains  
Lard half an ounce  
Mix. To relieve rheumatic pains. *Legrand.*

**AURI ET SODII CHLORIDUM.**

**CHLORIDE OF GOLD AND SODIUM.**

R. Chloride of gold eighty-five parts  
Chloride of sodium sixteen parts  
Dissolve in a small quantity of water, evaporate to a pellicle, and let crystallize. *Par. Cod.*

R. Leaf gold four parts  
Nitro-muriatic acid sufficient  
to dissolve, evaporate to dryness, add  
Chloride of sodium one part  
Water thirty-two parts  
Evaporate to one-half, and set aside to crystallize. *Figuiet.*

**POWDER OF CHLORIDE OF GOLD AND SODIUM.**

R. Chloride of gold and sodium one part

*Lycopodium* two parts  
Mix. *Cadet.*

R. Chloride of gold and sodium one part  
Sugar of milk sixteen parts  
Mix. *Radius.*

R. Chloride of gold and sodium three parts  
Powdered orris root nine parts  
Mix. Three grains contain about three-fourths of a grain of the chloride, and are used for thirty frictions, for the weakest doses, and three for the strongest. *Legrand.*

**PILLS OF CHLORIDE OF GOLD AND SODIUM.**

R. Chloride of gold and sodium ten grains  
Potato starch four grains  
Gum Arabic  
Distilled water, each, one drachm  
Mix, and make one hundred and twenty pills. Each contains one-twelfth of a grain of chloride. *Soubeiran.*

R. Chloride of gold and sodium four grains  
Extract of bitter sweet one drachm  
aconite ten grains  
Powdered marsh mallow  
Distilled water, each, sufficient  
Mix, and make eighty pills. Dose, three thrice a-day, in dropsy, incontinence of urine, amaurosis, &c. *Groetznar.*

R. Chloride of gold and sodium one grain  
Extract of mezereon one drachm  
Mix, and make sixty pills. *Magendie.*

**LOZENGES OF CHLORIDE OF GOLD AND SODIUM.**

R. Chloride of gold and sodium five grains  
Powdered sugar one ounce  
Mucilage of tragacanth sufficient  
Rub the chloride in a glass mortar with the sugar, add the mucilage, and make sixty lozenges. *Soubeiran.*

**SOLUTION OF CHLORIDE OF GOLD AND SODIUM.**

R. Chloride of gold and sodium two grains  
Distilled water one ounce  
Dissolve. Ten drops, every two hours, in dropsy and incontinence of urine. *Radius.*

**SYRUP OF CHLORIDE OF GOLD AND SODIUM.**

R. Chloride of gold and sodium one grain  
Syrup six ounces  
Dissolve. Dose, from one ounce to two. *Bories.*



COMPOUND SYRUP OF CHLORIDE OF GOLD  
AND SODIUM.

R. Peruvian bark  
Gentian, each, three ounces  
Mezereon one ounce and a-half  
Foxglove two drachms and a-half  
Water one pint and a-half

Boil down to one pint, and pour the boiling decoction on

Bitter orange peel one ounce  
Cloves half an ounce

After a sufficient infusion, strain, and add

Chloride of gold and sodium four grains  
A spoonful, night and morning. *Bories.*

OINTMENT OF CHLORIDE OF GOLD AND  
SODIUM.

R. Chloride of gold and sodium one scruple  
Lard one ounce

Mix well. *Foy.*

R. Chloride of gold and sodium three to  
four grains

Lard half an ounce  
Mix thoroughly. *Groetzner.*

## AURI CYANIDUM.

## CYANIDE OF GOLD.

R. Leaf gold one part  
Nitro-muriatic acid six parts

Dissolve, and evaporate to dryness. Dissolve in

Distilled water eight parts  
Reduce on a water-bath to one-fourth, and add very gradually

Cyanide of potassium half a part  
Distilled water twenty-four parts

Agitate, let rest, and separate the cyanide. *Soubeiran.*

R. Fresh precipitated oxide of gold  
(washed, but not dried) at will

Diluted hydrocyanic acid sufficient

Boil till solution assumes a beautiful yellow tint, evaporate to dryness on a water-bath. *Desfosses.*

Dose, from eighteenth to tenth of a grain.

## POWDER OF CYANIDE OF GOLD.

R. Cyanide of gold one grain  
Lycopodium fifteen grains

Mix, and divide into sixteen powders. One every day, as friction on the gums and tongue. *Radius.*

## PILLS OF CYANIDE OF GOLD.

R. Cyanide of gold one grain  
Powdered liquorice thirty-one grains

Mucilage sufficient  
Mix, and make sixteen pills. *Radius.*

R. Cyanide of gold one grain  
Extract of mezereon three grains  
Powdered marsh mallow sufficient  
Mix, and make five grain pills.

## LOZENGES OF CYANIDE OF GOLD.

R. Cyanide of gold one grain  
Chocolate sufficient  
to make sixteen lozenges. *Soubeiran.*

## SOLUTION OF CYANIDE OF GOLD.

R. Cyanide of gold three grains  
Diluted alcohol eight fl. ounces

A teaspoonful, twice a-day, gradually increasing the dose. *Christien.*

## AURI IODIDUM.

## IODIDE OF GOLD.

R. Solution chloride of gold at will  
Iodide of potassium sufficient

to precipitate, wash the precipitate with alcohol, and dry it. *Par. Cod.*

R. Solution chloride of gold at will  
hydriodate of ammonia sufficient

to precipitate. Wash with alcohol, and dry. *Miellat.*

Used like the other preparations, and in the same doses.

## AURI OXIDUM.

## OXIDE OF GOLD.

R. Pure gold one part  
Mercury six parts

Make an amalgam; triturate with double the weight of sulphur and calceine. *Wurt. Ph.*

R. Leaf gold half an ounce  
Nitre ten ounces

Common salt five ounces  
Alum eight ounces

Triturate to extinction, dissolve in lime-water, precipitate by solution of potassa; wash the precipitate, and calceine. *Spielmann.*

R. Chloride of gold one part  
Distilled water forty parts

Dissolve, and add

Fresh calcined magnesia four parts

Boil; wash the precipitate with distilled water, then with nitric acid diluted with twenty parts of water, and again with water. Dry in the shade. *Par. Cod.*

Dose, from a tenth to three-fourths of a grain.

## POWDER OF OXIDE OF GOLD.

R. Oxide of gold two drachms



Sulphuret of antimony half an ounce  
 Oil of cinnamon eight drops  
 Triturate together. *Spielmann.*  
 R. Oxide of gold four scruples  
 Prepared hartshorn one ounce  
 Triturate together. Used in malignant  
 fevers and small-pox. *Spielmann.*

## PILLS OF OXIDE OF GOLD.

R. Oxide of gold five grains  
 Extract of mezereon two drachms  
 Mix, and make sixty pills. *Magendie.*

## AURUM AMMONIATUM.

## AMMONIATED GOLD.

R. Gold leaf two drachms  
 Nitro-muriatic acid one ounce  
 Dissolve, and add  
 Water of ammonia sufficient  
 to precipitate. Wash and dry, with great  
 caution. *Niemann.*

This is what is termed fulminating gold, and  
 explodes at 400°. It has been used in fevers,  
 &c., as a diaphoretic, but has produced unpleasant  
 consequences.

## PILLS OF AMMONIATED GOLD.

R. Ammoniated gold  
 Calomel  
 Extract of squill, each, half a drachm  
 Powdered rhubarb two drachms  
 Conserve of juniper sufficient.  
 Mix, and make pills of two grains. *Plenck.*

## AURUM STANNO PARATUM.

## PURPLE OF CASSIUS.

R. Chloride of gold one part  
 Distilled water two hundred parts  
 Add to this solution, very gradually,  
 Pure tin one part  
 Nitro-muriatic acid three parts  
 Dissolve without heat, and add  
 Distilled water one hundred parts  
 as long as any precipitate falls. Wash this  
 and dry by a gentle heat. *Par. Cod.*

This is used like the other preparations of gold.

## AVENA SATIVA.

## OAT.

The common oat, although generally cultivated  
 from a very early age, is not known in a wild  
 state.

*Sex. Syst.* Triand. digyn. *Nat. Syst.* Gramin-  
 aceæ.

*Linn. Sp. Pl.* 61. *Griffith, Med. Bot.* 662.

Though, for the most part, used as food for

horses, it contains much nutritive matter, and  
 forms an important article of diet in some parts  
 of Europe. It is also employed in medicine as  
 a bland, nutritious, and somewhat laxative nutri-  
 ment in inflammatory diseases. When merely  
 freed from their husk and coarsely broken, they  
 are called groats.

## GROAT GRUEL.

R. Groats three ounces  
 Wash well in cold water, and then put into  
 Fresh water four pints  
 Boil slowly to one half, strain through a  
 fine sieve. *A. T. Thomson.*

R. Groats five drachms  
 Liquorice root three drachms  
 Boil the groats in sufficient water, to have  
 one quart of decoction, add the liquorice,  
 and after a sufficient infusion, strain.  
*Cottureau.*

## OATMEAL GRUEL.

R. Oatmeal two ounces  
 River water one pint and a-half  
 Rub the meal in a basin, with the back of  
 a spoon, in some of the water, pouring off  
 the fluid after the grosser particles have  
 subsided, but whilst the milkiness remains,  
 repeat this with fresh water. Stir well,  
 and unite the washings, and boil until a  
 soft, thick mucilage is formed.

*A. T. Thomson.*

Both these preparations may be flavored ac-  
 cording to circumstances.

## COMPOUND DECOCTION OF OATMEAL.

R. Groats six ounces  
 Red sanders, chipped, one ounce  
 Chicory root one ounce and a-half  
 Water twelve pints  
 Boil down one-third, and add to the strained  
 decoction

Nitre half an ounce  
 Sugar two ounces

Dissolve. Advised in gout, nephritis, &c.,  
 to be taken hot or tepid, morning and even-  
 ing for some weeks, in doses of six or eight  
 fl. ounces. *Wurt. Ph.*

## FLUMMERY OF OATMEAL.

R. Oatmeal or groats a quart  
 Rub for a considerable time with two  
 quarts of hot water, and let the mixture  
 stand till it becomes sour, then add another  
 quart of hot water, and strain through a  
 hair sieve. Let stand till a white sediment  
 is deposited, decant the fluid, and wash the  
 sediment with cold water. This is now to  
 be boiled with fresh water, till it forms a  
 mucilage, stirring the whole time. It is a  
 very light and somewhat nutritious food,  
 during early convalescence.

*A. T. Thomson.*



## POWDER FOR A CATAPLASM.

R. Linseed meal                      one part  
 Oatmeal                                two parts  
*Dub. Ph.*

## OATMEAL POULTICE.

R. Oatmeal                              sufficient  
 Stir gradually into water kept boiling, until it is sufficiently consistent.

*A. T. Thomson.*

R. Powder for a cataplasm            at will  
 Boiling water                          sufficient  
*Dub. Ph.*

## AZEDARACH.

## AZEDARACH.

The *Melia Azedarach* is a tree, a native of many parts of Asia, and naturalized in the warmer parts of Europe and the United States.

*Ser. Syst.* Hexand. monog. *Nat. Syst.* Meliaceæ.

Linn. Sp. Pl. 550. Griffith, Med. Bot. 179.

The parts used are the bark of the root, the berries, and the leaves. The former only is official in the U. S. Pharm. It is cathartic and emetic, in large doses somewhat narcotic; it is also an efficient anthelmintic.

## DECOCTION OF AZEDARACH.

R. Azedarach                            four ounces  
 Water                                    two pints  
 Boil to a pint, and strain.

Dose, one fl. ounce, every two or three hours, for a child.

## OINTMENT OF AZEDARACH BERRIES.

R. Pulp of berries  
 Lard, each,                              one ounce  
 Rub well together. Said to be useful as an application in tinea capitis.

## B

## BALLOTA LANATA.

## HAIRY HOREHOUND.

Several species of *Ballota* have, at different times, been used in medicine, but this is the only one that appears to possess active properties.

*Ser. Syst.* Didynam. gym. *Nat. Syst.* Lamiacæ.

The whole plant is used, with the exception of the root, and is stated to be an excellent diuretic, and useful in dropsical cases.

## DECOCTION OF HAIRY HOREHOUND.

R. Hairy horehound one ounce and a-half  
 Boiling water                          two pints  
 Boil down to one pint. Dose, a cupful twice a-day, gradually increasing the quantity.  
*Rehmann.*

## DECOCTION OF WILD INDIGO.

R. Root of wild indigo                one ounce  
 Boiling water                          a pint and a-half  
 Boil down to a pint. Dose, half a fl. ounce every four to eight hours.

*Comstock.*

## OINTMENT OF WILD INDIGO.

R. Contused root of wild indigo  
     two ounces  
 Lard    six ounces

Simmer together for an hour, and strain. Has been found beneficial as an application to burns and ulcers.

## BARIUM.

## BARI CHLORIDUM.

## CHLORIDE OF BARIUM.

R. Carbonate of barytes                one pound  
 Muriatic acid                          twelve fl. ounces  
 Water                                      three pints

Mix the acid with the water, and gradually add the barytes; towards close of effervescence, apply a gentle heat, and, when action has ceased, filter, and boil down, for crystals to form. *U. S. Ph.*

## POWDERS OF CHLORIDE OF BARIUM.

R. Chloride of barium                  two drachms  
 Calomel                                    ten grains  
 Sulphuret of antimony                six grains

## BAPTISIA TINCTORIA.

## WILD INDIGO.

A native plant, found in most parts of the country; becoming black when dried.

*Ser. Syst.* Decand. monog. *Nat. Syst.* Fabaceæ.  
 Torrey and Gray, Flor. i. 386. Griffith, Med. Bot. 231.

The root, which is the part used, is emetic and purgative when fresh. Has been considered a valuable antiseptic, and febrifuge, and has been given with advantage in typhus fever, scarlatina, &c., and also used as a wash to foul ulcers, aphthæ, &c. Most employed in decoction.



Mix, and divide into forty-eight powders.  
Dose, two a-day in syrup, in cutaneous affections. *Swed.aur.*

## PILLS OF CHLORIDE OF BARIUM.

R. Chloride of barium  
Extract of liquorice, each,  
half a drachm

Powdered liquorice root

Water, each, sufficient

Mix, and divide into one hundred and twenty pills. *Phœbus.*

Dose, four to eight, three or four times a-day.

R. Chloride of barium one drachm  
Resin of guaiacum half an ounce  
Conserve of fumitory sufficient

Mix, and make one hundred eighty-eight pills. One to be taken morning and evening, increased to two, against tapeworm. *Pierquin.*

## SOLUTION OF CHLORIDE OF BARIUM.

R. Chloride of barium one ounce  
Distilled water three fl. ounces  
Dissolve, and filter. *U. S. Ph.*

Has been used in small doses in cancer and scrofula. Dose, five drops, two or three times a-day, cautiously increasing.

## COLLYRIUM OF CHLORIDE OF BARIUM.

R. Chloride of barium ten grains  
Distilled water one ounce  
Dissolve, filter, and add  
Mucilage of quince seeds two drachms  
Wine of opium half a drachm

*Radius.*

The eyelids are to be washed several times a-day with this, in scrofulous ophthalmia.

## MIXTURE OF CHLORIDE OF BARIUM.

R. Chloride of barium  
iron, each, half a drachm  
Distilled water one ounce  
Dissolve *Augustin.*

Dose, twenty to sixty drops.

R. Chloride of barium  
Extract of cicuta, each, half a drachm  
Distilled water half an ounce  
Antimonial wine one drachm and

a-half

*Phœbus.*

Dose, fifteen to twenty drops every three hours, in chronic orchitis.

R. Chloride of barium one drachm  
Balm water seven fl. drachms  
Antimonial wine one fl. drachm

Dose, twenty to thirty drops four times a-day. *Hufeland.*

R. Chloride of barium four grains  
Distilled water two fl. ounces

Extract of hemlock four grains  
Common emulsion one pound  
Syrup one ounce

Mix. To be taken during the day, in scrofula, and scrofulous phthisis. *Brera.*

## BARI IODIDUM.

## IODIDE OF BARIUM.

R. Iodine one hundred parts  
Iron filings thirty parts  
Water sufficient

Prepare an iodide of iron, add baryta dissolved in twenty parts of water, as long as a precipitate is formed, heat a moment, filter, evaporate, and crystallize.

*Magendie.*

Been used with success in scrofula. Dose,  $\frac{1}{2}$  of a grain, three times a-day, cautiously increasing.

## POWDER OF IODIDE OF BARIUM.

R. Iodide of barium one grain  
Powdered cinnamon  
Sugar, each, four scruples  
Mix well, and divide into eight powders. One to be given two or three times a-day in scrofula. *Radius.*

## OINTMENT OF IODIDE OF BARIUM.

R. Iodide of barium four grains  
Lard one ounce  
Mix. As a friction in scrofulous swellings. *Bielt.*

## BARI SULPHURETUM.

## SULPHURET OF BARIUM.

R. Sulphate of barytes eleven parts  
Charcoal one part  
Oil of turpentine sufficient  
Triturate the barytes with the coal, moisten the mixture with the turpentine, and heat the whole in a crucible to redness; let cool, and preserve. *Van Mons.*

## BARYTA.

## BARYTES.

R. Nitrate of barytes sufficient  
Mix in a platina crucible to a red heat, when the mass has become solid and porous, raise to a white heat, remove from fire, and cool. *Van Mons.*

## SOLUTION OF BARYTES.

R. Barytes one part  
Distilled water twenty parts  
Dissolve. *Van Mons.*



Has been recommended in scrofula, in doses of four to five drops, in some appropriate vehicle.

#### BARYTIC LINIMENT.

R. Solution of barytes                      one part  
Olive oil                                      six parts  
Rub together.                              *Soubieran.*

Advised as an external application in lepra, and other obstinate cutaneous diseases.

#### BARYTÆ ACETAS.

##### ACETATE OF BARYTES.

R. Carbonate of barytes                      at will  
Acetic acid                                      sufficient  
to saturate; filter, evaporate, and crystallize.                      *Van Mons.*

##### SOLUTION OF ACETATE OF BARYTES.

R. Acetate of barytes                      one part  
Distilled water                              nine parts  
Dissolve, and filter. In same cases and doses as the solution of barytes.                      *Hamb. Ph.*

#### BEBEERINA.

##### BEBEERINE.

This is an alkaloid obtained from the bark of a tree, growing in British Guiana. It is said to belong to the genus *Nectandra*, and has been named *N. radxi*. The bark is in flat pieces, smooth, grayish, hard, heavy, and brittle, with but little odor, though of a very bitter taste. Bebeerine is extracted from this bark in the form of a sulphate, by a process similar to that used to obtain sulphate of quinia. In this form it contains both bebeerine and siperina, and is in thin, somewhat glittering scales of a brownish-yellow color, forming a yellow powder, soluble in cold water, but often forming a turbid solution, which is rendered clear by a few drops of diluted sulphuric acid. Pure bebeerine can be obtained from it as follows:—

Decompose by ammonia, wash the precipitate, and whilst moist, triturate with moist hydrated oxide of lead; dry on a water-bath, exhaust with alcohol, and distil off the spirit, treat the residue with ether; on the evaporation of the ether, bebeerine will be left of a bright Canary yellow color, but in powder appears nearly white.

Dose of the sulphate is one to three grains as a tonic, and five grains to a scruple as a febrifuge.

It is not equal to quinine as an antiperiodic, but is a good substitute for that article.

##### PILLS OF SULPHATE OF BEBEERINE.

R. Sulphate of bebeerine                      two drachms  
Conserve of roses                              sufficient  
Mix, and form twenty-four pills—one to three, thrice a-day in intermittent fevers.                      *Christison.*

#### SOLUTION OF SULPHATE OF BEBEERINE.

R. Sulphate of bebeerine                      half a drachm  
Diluted sulphuric acid                      twenty-five minims

Syrup  
Tincture of orange peel, each,                      one ounce

Water                                      four ounces

Mix. A tablespoonful three times a-day.                      *Christison.*

#### BECCABUNGA.

##### BROOKLIME.

Two species of *Veronica* of similar properties are included under this name, the *V. beccabunga* and *V. anagallis*; they are both semi-aquatic plants indigenous to Europe and to this country. *Ser. Syst. Diand. monog. Nat. Syst. Scrophulariaceæ.*

These plants are employed usually in a fresh state, and considered to be antiscorbutic and alterative. They are usually given in infusion made with one or two handfuls of the herb to a pint of boiling water. They have also been employed in conserve, syrup, &c.

##### WATER OF BROOKLIME.

R. Brooklime  
Water, each,                                      two parts  
Distil off one part.                              *Cottureau.*  
Dose, one to four ounces.

##### SYRUP OF BROOKLIME.

R. Clarified juice of brooklime                      one part  
Sugar                                      two parts  
Dissolve, by means of a water-bath.                      *Taddei.*

#### BELLADONNA.

##### BELLADONNA.

This is the *Atropa Belladonna*, an herbaceous plant, native of Europe, and cultivated in some places in the United States.

*Ser. Syst. Pentand. monog. Nat. Syst. Solanaceæ.*

*Linn. Sp. Pl. 260, Griffith, Med. Bot. 486.*

The whole plant is officinal in some Pharmacopœias, but the leaves only are ordered in that of the U. S. This plant is a powerful narcotic, owing to the presence of a peculiar principle called *Atropia*. It has been used in a variety of diseases as an anodyne, antispasmodic, and discutient. The dose of the powdered leaves is one to two grains, daily or twice a-day, gradually increasing.

##### POWDER OF BELLADONNA.

R. Powdered belladonna root                      three grains  
   ipecacuanha                      two grains  
Oxide of zinc                                      six grains  
Sugar                                      one drachm  
Rub together, and divide into six powders. One every two hours in epilepsy.                      *Radius.*



R. Powdered belladonna root two grains  
 ipecacuanha one grain  
 Washed sulphur thirty-two grains  
 Sugar of milk two drachms  
 Rub well together, divide into twenty powders. One every three hours in whooping-cough. *Phœbus.*

R. Powdered belladonna one grain  
 nitrate of potassa twenty-one grains  
 sugar nine grains  
 Make a powder, to be taken at bedtime.

In chronic rheumatism, extensive ulcerations, mania, epilepsy. *A. T. Thomson.*

R. Powdered root of belladonna  
 ipecacuanha, each, six grains  
 liquorice root  
 sugar, each, half a drachm  
 Precipitated sulphur two scruples  
 Oil of anise  
 amber, each, three minims  
 Mix, and make five to twenty powders. *A. T. Thomson.*

R. Powdered leaves of belladonna one to three grains  
 Musk  
 Camphor, each, five grains  
 Sugar thirty grains  
 Triturate well together, and divide into eight powders. *A. T. Thomson.*

POWDER OF BELLADONNA AND RHUBARB.  
 R. Powdered belladonna leaves ten grains  
 rhubarb two scruples  
 Mix, and divide into ten powders. One, two, or three times a-day, in obstructions of the liver and spleen. *Radiis.*

#### OPIATED BELLADONNA LEAVES.

R. Belladonna leaves  
 Water, each, two ounces  
 Extract of opium one drachm  
 Dissolve the opium in the water, wash the leaves with the solution, and dry them by a gentle heat. *Guibourt.*

Said to be very useful in phthisis, the patient to smoke a pinch every morning in a common pipe.

#### FUMIGATION OF BELLADONNA.

R. Powdered belladonna one drachm  
 Boiling infusion of sage one quart  
 The steam to be inhaled. Said to be useful in whooping-cough and phthisis. *Soubéiran.*

#### EXTRACT OF BELLADONNA.

R. Belladonna leaves one pound  
 Bruise in a stone mortar, with the addition of a little water; express the juice, heat to

boiling point, strain, and evaporate to proper consistence. *U. S. Ph.*

Dose, one grain night and morning, to be gradually increased.

#### ALCOHOLIC EXTRACT OF BELLADONNA.

R. Belladonna leaves one pound  
 Diluted alcohol four pints  
 Moisten the leaves with half a pint of alcohol, let stand for twenty-four hours, put in a displacement apparatus, add gradually the remainder of alcohol. When this has penetrated the belladonna, add water occasionally so as to keep the powder covered. Stop the process when the liquid that passes forms a precipitate in what has already passed. Distil off alcohol, and evaporate to proper consistence. *U. S. Ph.*

Dose, half a grain to a grain, to be gradually increased.

#### COMPOUND BELLADONNA PILLS.

R. Extract of belladonna  
 Blue pill  
 Powdered ipecacuanha, each, twelve grains  
 Mix, and make twelve pills. One to be taken morning and evening, in cancerous affections. *Ainslie.*

#### SOLUTION OF EXTRACT OF BELLADONNA.

R. Extract of belladonna three grains  
 Cinnamon water one fl. ounce  
 Make a solution. Dose, for a child under one year, two or three drops twice or thrice a day, and an additional drop for each additional year. *Ellis.*

Said to have proved efficacious as a preventive of scarlatina.

#### SYRUP OF BELLADONNA.

R. Extract of belladonna forty-eight grains  
 Distilled water two ounces  
 Dissolve and filter, then add  
 Syrup three pounds  
 Boil, and strain. *Cottureau.*

Said to be useful in whooping-cough, in doses of one to three drachms.

#### GILLET'S SYRUP OF BELLADONNA.

R. Dried belladonna leaves one ounce  
 Digitalis  
 Common nightshade, each, one drachm  
 Water ten fl. ounces  
 Digest for two days, and filter, then add  
 Syrup two pounds  
*Cadet.*

Advised in whooping-cough, in the dose of a teaspoonful every hour or two.

#### TINCTURE OF BELLADONNA.

R. Belladonna leaves four ounces  
 Diluted alcohol two pints



Macerate for fourteen days, express, and filter. *U. S. Ph.*

The dose is from fifteen to thirty drops.

#### ETHEREAL TINCTURE OF BELLADONNA.

R. Belladonna leaves dried one part  
Sulphuric ether eight parts  
Macerate for eight days, and filter.

*Guibourt.*

#### BELLADONNA MIXTURE.

R. Extract of bella-  
donna eight to twelve grains  
Wine of ipecacuanha one fl. drachm  
Syrup of seneka half fl. ounce  
tolu three and a half fl. ounces

Mix. Advised in catarrh, in the dose of a teaspoonful, three or four times a-day, using also gum water acidulated with lemon-juice. *Ellis.*

R. Extract of belladonna fifteen grains  
Fennel water five fl. ounces  
Alcohol one fl. drachm  
Dissolve. Dose, a drop for every year of the age of a child, not to exceed fifteen; as a preventive of scarlatina. *Maisier.*

#### LINIMENT OF BELLADONNA.

R. Extract of belladonna two scruples  
Sulphuric ether one drachm  
Cherry laurel water two ounces  
As a friction to the abdomen in colica pictonum. *Phæbus.*

#### CLYSTER OF BELLADONNA.

R. Belladonna  
leaves twelve to twenty grains  
Jalap one scruple  
Boiling water half a pint  
Infuse, and strain. *Pitschaft.*

This has been recommended in strangulated hernia, to overcome spasm.

#### PLASTER OF BELLADONNA.

R. Resin plaster three ounces  
Extract of bella-  
donna one ounce and a-half  
Melt plaster by means of water-bath, and add the extract; mix well. *U. S. Ph.*

An efficacious application in neuralgic and rheumatic pains.

#### OINTMENT OF BELLADONNA.

R. Extract of belladonna two drachms  
Distilled water two fl. ounces  
Lard two ounces  
Rub well together. Used by Chaussier as an application to the neck of the uterus in cases of rigidity. *Ratier.*

R. Fresh belladonna leaves one part  
Lard two parts

Simmer together, express, and strain. Used in frictions of one or two drachms, to indolent tumours. *Soubeiran.*

## BENZOINUM.

### BENZOIN.

Is the concrete juice of the *Styrax Benzoïn*, a tree of some size, native of several of the East Indian islands.

*Ser. Syst.* Decand monog. *Nat. Syst.* Styracaceæ.

Dryander, Ph. Tl. lxxvii. 308. Griffith, Med. Bot. 439.

There are several varieties of benzoin, but the best is in whitish tears, united by a reddish-brown connecting medium. It has an aromatic, agreeable odor, and a somewhat acrid taste. It is stimulant and expectorant, and is much used in chronic catarrhs.

### BENZOATED LARD.

R. Benzoin in coarse powder one ounce  
Lard twenty-five ounces  
Heat them together in a water-bath for two or three hours, strain without pressure, and stir while cooling.  
Employed for ointments to be long kept, to prevent them from becoming rancid. *Deschamps.*

### FUMIGATING POWDER.

R. Powdered olibanum two pounds  
benzoin  
storax, each, half a pound  
dried roses  
lavender  
flowers, each, six ounces  
Mix. A small quantity to be thrown on hot coals, to raise a smoke; to be inhaled in whooping-cough, &c. *Dohrn.*

### FUMIGATING PASTILLES.

R. Benzoin ten parts  
Charcoal twenty-four parts  
Nitrate of potassa one part  
Sassafras two parts  
Mucilage of gum Arabic sufficient  
To make pastilles, which are to be conical. *Beral.*

R. Benzoin sixteen parts  
Sandal wood four parts  
Ladanum one part  
Balsam of tolu four parts  
Charcoal forty-eight parts  
Nitrate of potassa two parts  
Mucilage of tragacanth sufficient  
Make conical pastilles. *Foy.*

### POWDER OF BENZOIN.

R. Powdered benzoin  
sugar candy equal parts



Mix. Said to be useful with camphor water in asthma and chronic catarrh. *Pierquin.*

OIL OF BENZOIN.

R. Benzoin at will  
Water a small quantity  
Distil on a sand-bath, and separate the oil that passes over.  
Advised as a friction in neuralgic and rheumatic pains. *Swedliaur.*

COSMETIC WASH OF BENZOIN.

R. Tincture of benzoin three drachms  
Liquid subcarbonate of potassa two drachms  
Rose water eight ounces  
Mix. As a lotion in acne. *Augustin.*

COSMETIC EMULSION OF BENZOIN.

R. Almond paste two drachms  
Rose water eight ounces  
Make an emulsion, and add  
Tincture of benzoin three drachms  
*Schubarth.*

MILK OF ROSES.

R. Tincture of benzoin one drachm  
Rose water one pint  
Mix. *Taddei.*  
R. Tincture of benzoin one drachm  
balsam of Peru twenty drops  
Rose water one pint  
Mix. *Giannini.*

Both these are used as cosmetic washes.

LOTION FOR BURNS.

R. Benzoin six drachms  
Storax four drachms  
Balsam tolu two drachms  
Aloes one drachm  
Alcohol eight fl. ounces

Make a tincture, to be used as a lotion for burns and scalds, before vesication has taken place.

TINCTURE OF BENZOIN.

R. Benzoin one part  
Alcohol eight parts  
Digest for some days, and filter. *Guibourt.*

Used as a stomachic carminative, &c., in doses of ten to twenty drops.

COMPOUND TINCTURE OF BENZOIN.

R. Benzoin three ounces  
Purified storax two ounces  
Balsam tolu one ounce  
Powdered aloes half an ounce  
Alcohol two pints  
Macerate for fourteen days, and filter. *U. S. Ph.*

A stimulating expectorant, and also used as an application to indolent ulcers.

TURLINGTON'S BALSAM.

R. Benzoin twelve ounces  
Liquid storax fourteen ounces  
Balsam of Peru two ounces  
Myrrh  
Aloes, each, one ounce  
Balsam of tolu  
Extract of liquorice, each, four ounces  
Angelica root half an ounce  
Alcohol eight pints  
Digest for ten days, and strain. *Phil. Coll. Pharm.*

BERBERIS.

BARBERRY.

One species of this genus, *B. vulgaris*, has been employed in medicine in Europe. It is a native of Europe, but is naturalized in some parts of the U. S.

*Ser. Syst. Hexand. monog. Nat. Syst. Berberidaceæ.*

Torrey and Gray, Fl. i. 49. Griffith, Med. Bot. 112.

The parts used are principally the berries, which are acidulous, and form a substitute for tamarinds, in the preparation of cooling drinks. The bark of the root is bitter and astringent, and is useful in the treatment of aphthous sore mouth, and was at one time much esteemed in the treatment of jaundice.

LEMONADE OF BARBERIES.

R. Juice of berries one part  
Sweetened water fifteen parts  
Mix. As a refreshing drink in fevers. *Beral.*

BISMUTHUM.

BISMUTH.

BISMUTHI SUBNITRAS.

SUBNITRATE OF BISMUTH.

R. Purified bismuth one ounce  
Nitric acid one fl. ounce and a-half  
Diluted water sufficient  
Mix one fl. ounce of water with the acid, and dissolve the bismuth in the mixture; when action has terminated, pour clear solution into three pints of the water, and permit precipitate to subside; pour off supernatant fluid, wash the precipitate well, and dry it by a gentle heat. *U. S. Ph.*

Used as a tonic and antispasmodic, especially in certain painful affections of the stomach. Dose, five grains, twice or thrice a-day, gradually increasing the quantity.

COMPOUND BISMUTH POWDER.

R. Subnitrate of bismuth twelve grains  
Powdered ipecacuanha two grains  
Carbonate of magnesia two drachms



Mix, and form twelve powders. One, two or three times a-day, in gastrodynia.

*Clarus.*

R. Subnitrate of bismuth eight grains  
Opium one grain  
Sugar two drachms

Mix, and form four powders. *Ammon.*

Much praised by the author, in cholera; one powder to be taken every two hours.

#### PILLS OF SUBNITRATE OF BISMUTH.

R. Subnitrate of bismuth one drachm  
Mucilage of gum Arabic sufficient  
Mix, and make thirty pills. One to be given every two hours in dyspepsia. *Ellis.*

#### SUBNITRATE OF BISMUTH OINTMENT.

R. Subnitrate of bismuth one part  
Lard three parts  
Rub well together. Said to be useful in psora, and other cutaneous eruptions.

*Kerksig.*

### BISTORTA.

#### BISTORT.

This is the root of the *Polygonum Bistorta*, a native of Europe, and also found in the northern parts of this country, growing in wet situations.

*Sex. Syst.* Octand. trigyn. *Nat. Syst.* Polygonaceæ.

Linn. Sp. Pl. 516. Lindley, Fl. Med. 361.

The part employed, is the root; this is bitter and austere, especially in a fresh state. It is a powerful astringent, but is seldom used in this country. The dose of the powder is twenty or thirty grains, three or four times a-day.

#### INFUSION OF BISTORT.

R. Bistort one ounce  
Boiling water two pints  
Infuse for two hours, and strain. Recommended in passive hemorrhages. *Radius.*

#### ASTRINGENT CLYSTER.

R. Bistort one ounce  
Poppy heads two drachms  
Water one pint  
Infuse, and strain. As an injection, in diarrhoea and chronic dysentery.

*Guibourt.*

### BOLETUS LARICIS.

#### WHITE AGARIC.

A parasitic fungus growing on the trunk of the larch, in many parts of Europe. It is destitute of smell, but the taste, at first rapid and farinaceous, becomes bitter, acrid, and nauseous. Formerly much used as a purgative, in doses of half a drachm to a drachm. It has lately been thought to be very efficacious in arresting the colliquative sweats in phthisis.

#### POWDER OF AGARIC.

R. Powdered agaric twelve grains  
White sugar two drachms  
Mix well, and divide into six powders. One to be given at night, against colliquative sweats. *Radius.*

#### POWDER OF AGARIC AND OPIUM.

R. Powdered agaric eighteen grains  
Opium three grains  
Gum Arabic two drachms  
Mix, and divide into nine powders. One to be given at night, against colliquative sweats. *Radius.*

#### PILLS OF AGARIC AND OPIUM.

R. Powdered agaric fifteen grains  
Extract of opium two grains and a-half  
Mix, and form six pills. One or two at bedtime, in same cases as above noticed. *Rayer.*

### BROMINUM.

#### BROMINE.

This elementary substance has much analogy to iodine in its chemical characters, and in its action on the system. It is a dark red, volatile liquid, having a very caustic taste and a disagreeable odor. It has been employed in bronchocele, scrofula, chronic cutaneous diseases, and hypertrophy of the heart.

#### SOLUTION OF BROMINE.

R. Bromine one part  
Distilled water forty parts  
Mix. Dose, six drops several times a-day, gradually increasing the dose. *Pourché.*

#### ALCOHOLIC SOLUTION OF BROMINE.

R. Bromine ten drops  
Alcohol one ounce  
Mix. As an external application. The strength to be increased five drops to the ounce, daily. *Founet.*

#### LOTION OF BROMINE.

R. Bromine twenty to thirty drops  
Water one pint  
Mix. For scrofulous ulcers. *Glover.*  
R. Bromine four fl. drachms  
Water five fl. ounces  
Mix. *Pourché.*

### BRUCIA.

#### BRUCINE.

R. Powdered false *Angustura* bark, at will. Treat it three times with water acidulated with muriatic acid, mix the liquids, evaporate, add milk of lime, wash the precipitate, dry, and treat it with alcohol; eva-



porate this, combine the residue with sulphuric acid, dissolve the salt in water, treat with animal charcoal, crystallize, redissolve in water, and precipitate by means of ammonia. *Cottureau.*

A highly poisonous alkaloid, obtained from the bark of the *strychnos nux vomica*, or false Angustura bark. It is white, very bitter, and readily soluble in alcohol, but with difficulty in water. Acts on the system like strychnia, but with less energy, and has been given in same class of diseases.

#### PILLS OF BRUCIA.

R. Brucia twelve grains  
Conserve of roses half a drachm  
Mix, and divide into twenty-four pills. One to be given morning and evening, gradually increasing the dose. *Foy.*

#### SOLUTION OF BRUCIA.

R. Brucia six grains  
Distilled water four fl. ounces  
Sugar two drachms  
Mix. A tablespoonful, morning and evening. *Magendie.*

#### TINCTURE OF BRUCIA.

R. Brucia eighteen grains  
Alcohol (0.847) one ounce  
Dissolve. Six to twenty-four drops in some demulcent drink. *Soubeiran.*

### BRUCIÆ ACETAS. MURIAS ET SULPHAS.

#### ACETATE, MURIATE, AND SULPHATE OF BRUCINE.

These are all made by the same process, using for each the appropriate acids.

R. Brucine at will  
Acetic acid sufficient  
Put the brucine in a porcelain capsule on a water-bath; pour a small quantity of water on it, and then add the acid very gradually, constantly stirring till perfect saturation takes place; filter, evaporate to one-half, and then crystallize. *Cottureau.*

### BRYONIA.

#### BRYONY.

The roots of two species of bryony are used

in medicine, the *B. alba*, and *B. dioica*, the first in England, the latter in other parts of Europe, but are identical in their sensible properties and action on the system.

*Sex. Syst.* Monœc. syngen. *Nat. Syst.* Cucurbitaceæ.

*Linn. Sp. Pl.* 621. *Griffith, Med. Bot.* 311.

The part used is the root; this is acrid and purgative, causing copious, watery stools; in doses of ten grains to two scruples. It has also some reputation as a cataplasm.

#### CATAPLASM OF BRYONY.

R. Juice of bryony root  
Crumb of bread, each, sufficient  
to form a cataplasm. Used as an application to engorged glands of the neck. *Barthez.*

#### COMPOUND CATAPLASM OF BRYONY.

R. Rased bryony root three ounces  
Contused elder bark one ounce  
Gum ammoniac half an ounce  
Muriate of ammonia two drachms  
Oil of cicuta two ounces  
Vinegar sufficient  
Mix, and heat. An application to scrofulous tumors. *Plenck.*

#### WINE OF BRYONY.

R. Bryony root one ounce  
White wine one pint  
Boil gently, and filter. To be taken in wineglassfuls, in dropsies. *Hufeland.*

### BUXUS.

#### Box.

The box is a shrub or small tree, native of the south of Europe, but now generally cultivated in this country.

*Sex. Syst.* Monœc. triand. *Nat. Syst.* Euphorbiaceæ.

The parts used are the wood and leaves. The first is sudorific and possesses somewhat the properties of guaiacum; the latter are purgative, but neither are now used. The following preparation, however, has obtained some celebrity in the cure of gonorrhœa, and in epilepsy.

#### OIL OF BOX.

R. Rased box-wood sufficient  
Distil, separate the oil, and rectify it.  
Dose, four to five drops, three times a-day. *Wirttemberg Ph.*



## C.

## CACAO.

## COCOA. CHOCOLATE NUTS.

These are the seeds of *Theobroma Cacao*, a small tree, indigenous to South America, and are also produced from some other sources.

*Sex. Syst.* Polydelph. pentand. *Nat. Syst.* Byttneraceæ.

Linn. Sp. Pl. 1100. Lindley, Flor. Med. 138.

The nuts are of an almond shape, and consist of a white, sweetish, somewhat oleaginous substance, covered by a leathery like shell. They are principally used as an article of food, but also are employed in medicine.

## COMPOUND COCOA POWDERS.

## PALAMOND.

R. Roasted cocoa eight ounces  
Rice flour  
Potato starch, each, two pounds  
Red saunders, powdered, one ounce  
Mix. A drachm to an ounce, boiled in water, as a restorative diet. *Soubiran.*

## RACAHOUT.

R. Roasted cocoa two ounces  
Potato starch five ounces  
Powdered salep one ounce  
Sugar eight ounces  
Vanilla sufficient  
Mix. Two or three spoonfuls boiled with eight fl. ounces of water, form a nutritious diet for the convalescent. *Foy.*

## WIKANA.

R. Roasted cocoa two ounces  
Sugar five ounces and a-half  
Powdered cinnamon two drachms  
vanilla thirty-six grains  
Ambergris three grains  
Musk one grain and a-half  
Mix. Half a drachm, boiled with milk or water, more stimulating than the last. *Guibourt.*

## COMPOUND COCOA PASTE CHOCOLATE.

## VANILLA CHOCOLATE.

R. Cocoa paste six pounds  
Sugar ten pounds  
Vanilla eleven drachms  
Triturate thoroughly together, and form cakes. *Cottreau.*

## AROMATIC CHOCOLATE.

R. Cocoa  
Sugar, each, sixteen ounces  
Powdered cinnamon half an ounce

Cloves two drachms  
Cardamom  
Vanilla, each, one drachm  
Triturate together, and form cakes. *Weigleb.*

## SALEP CHOCOLATE.

R. Cocoa paste  
Sugar, each, ten pounds  
Powdered salep ten ounces  
Triturate well, and form cakes of half an ounce each. *Cadet.*

## WHITE CHOCOLATE.

R. Sugar six pounds  
Rice flour one pound twelve ounces  
Potato starch eight ounces  
Gum Arabic four ounces  
Tincture of vanilla half fl. ounce  
Butter of cocoa eight ounces  
Boiling water sufficient  
Triturate well, into a stiff paste. *Cottreau.*

These afford a good article of diet, for convalescents, and debilitated persons, when boiled with water or milk. The last is the least stimulating.

## CHOCOLATE CREAM.

R. Boiling milk sixteen parts  
Sugar  
Yolk of egg, each, two parts  
Ground cocoa one part  
Mix, and boil gently. *Beral.*

## BUTTER OF COCOA.

R. Cocoa, roasted at will  
Reduce to a paste in a warm iron mortar, triturate on a stone slab, add a fifth part boiling water, place it in a bag, and express between two heated metal plates; melt the product, and, on cooling, separate the water. *Guibourt.*

## BUTTER OF COCOA MIXTURE.

R. Butter of cocoa three ounces  
Oil of almonds one ounce and six drachms  
Syrup of red poppies one ounce  
Orange-flower water four drachms  
Mix. As a soothing demulcent in catarrh. In spoonful doses. *Cadet.*

## BUTTER OF COCOA OINTMENT.

R. Butter of cocoa  
Oil of almonds equal parts  
Melt over a water-bath. *Guibourt.*



CADMIUM.

CADMIUM.

A soft, crystallizable, ductile, volatilizable metal, not used in medicine in its metallic state, but affording the following remedial salt.

CADMII SULPHAS.

SULPHATE OF CADMIUM.

R. Oxide of cadmium                      one ounce  
Sulphuric acid                              sufficient  
Dissolve, evaporate, and crystallize.

*Cottureau.*

Used in solution and ointment, as an application in chronic ophthalmia.

SOLUTION OF SULPHATE OF CADMIUM.

R. Sulphate of cadmium                  two grains  
Laudanum                                      twenty drops  
Rose water                                      four fl. ounces  
Mix. As a wash in chronic ophthalmia

*Guibourt.*

OINTMENT OF SULPHATE OF CADMIUM.

R. Sulphate of cadmium                  one or two grains  
Lard    one drachm  
Mix. As an application in spots on the cornea.

*Radiis.*

CALAMUS.

CALAMUS.

This is the root of *Acorus Calamus*, an indigenous plant found in most parts of the U. S. in wet situations. It also grows in Europe and Asia.

*Ser. Syst. Hexand. monog. Nat. Syst. Orontiacæ.*

Linn. Sp. Pl. 462. Griffith, Med. Bot. 620.

The part employed is the root; this has a fragrant odor, and a warm, bitterish, aromatic taste. It is a stimulant tonic, and aromatic. The dose, in substance, is from a scruple to a drachm.

ELECTUARY OF CALAMUS.

R. Powdered calamus                      two drachms and a-half  
valerian    one drachm and a-half

Syrup of orange peel                      two ounces

Mix. A teaspoonful every two hours.

*Shubarth.*

INFUSION OF CALAMUS.

R. Calamus                                      one ounce  
Boiling water                                  one pint

Infuse for a quarter of an hour, and strain. Dose, a wineglassful or more.

COMPOUND INFUSION OF CALAMUS.

R. Calamus                                      ten drachms  
Boiling water                                  one pint

Infuse and strain, then add

Peppermint water                          two ounces

Muriatic ether                              one drachm and a-half

Syrup    one ounce

Mix. Dose, a spoonful, in disordered digestion.

*Augustin.*

TINCTURE OF CALAMUS.

R. Calamus bruised                          one part  
Alcohol (0.847)                              eight parts

Macerate for eight days, and filter.

*Guibourt.*

COMPOUND TINCTURE OF CALAMUS.

R. Contused calamus  
ginger  
coriander

Black pepper, each,                          one ounce

Alcohol    two pints and a-half

Macerate for four days, and filter. Dose, forty to fifty drops as a stomachic and carminative.

*Niemann.*

R. Contused calamus  
gentian, each,                                  three ounces  
geum    two ounces and a-half  
angelica    one ounce and a-half

ginger    half an ounce

fennel    two ounces

alcohol    twelve pints

Macerate for six days, and filter. A teaspoonful in wine.

*Tromsdorff.*

CALCIUM.

CALCIUM.

CALCII CHLORIDUM.

CHLORIDE OF CALCIUM.

R. Chalk    five ounces

Muriatic acid

Water, each,                                      ten fl. ounces

Mix the acid and water, and gradually add the chalk. When all action ceases, filter, and evaporate to dryness. Fuse, and pour out on a stone slab; when cool, break in fragments, and preserve in well-stopped bottles.

*Lond. Ph.*

CATAPLASM OF CHLORIDE OF CALCIUM.

R. Chloride of calcium  
soda, each,                                      half an ounce  
Water    half a pint

Linseed meal                                      sufficient

Make a cataplasm. In scrofulous and white swellings.

*Swediaur.*

SOLUTION OF CHLORIDE OF CALCIUM.

R. Chloride of calcium                      eight ounces  
Water (imp. meas.)                          twelve fl. ounces  
Dissolve, and filter.

*Edin. Ph.*



Dose, thirty drops two or three times a-day, gradually increased; to be given in milk, or some demulcent. Used in scrofula, goitre, tabes mesenterica, &c.

#### MIXTURE OF CHLORIDE OF CALCIUM.

R. Chloride of calcium           one drachm  
Extract of henbane           ten grains  
Syrup of liquorice           one ounce  
Water                           six ounces

Mix. A spoonful four times a-day in scrofula. *Phæbus.*

R. Chloride of calcium           one drachm  
Almond mixture           seven fl. ounces  
Syrup of gum Arabic       one fl. ounce

Mix. A teaspoonful every three hours. *Gräfe.*

#### PILLS OF CHLORIDE OF CALCIUM.

R. Chloride of calcium           one drachm  
Extract of opium           nine grains  
Mucilage of gum Arabic       sufficient

Mix, and make fifty-four pills. One every two or three hours, in gonorrhea, gradually increasing the dose, until eight, ten, or twelve are taken every hour. *Gräfe.*

### CALX CHLORINATA.

#### CHLORINATED LIME.

##### CHLORIDE OF LIME.

R. Hydrate of lime           a pound  
Chlorine                   sufficient  
Pass the chloride over the lime, spread in a proper vessel, until it is saturated.

*Lond. Ph.*

It is a dessicant and disinfectant, and is also used to ill-conditioned ulcers, burns, chilblains, to some cutaneous affections, &c.

#### PRESERVATIVE LIQUID.

R. Water                   sixteen parts  
Chlorinated lime           four parts  
Alum                       two parts  
Nitro                      one part

Mix. Said to be very efficient for the preservation of anatomical preparations.

*Reboulet.*

#### LOZENGES OF CHLORINATED LIME.

R. Chlorinated lime           two drachms  
Sugar                   eight ounces  
Starch                   one ounce  
Tragacanth           one drachm  
Cochineal           three grains

Rub well together, and make lozenges of three grains each. One to be taken three or four times a-day, in cases of bad breath. *Deschamps.*

#### COLLUTORY OF CHLORINATED LIME.

R. Chlorinated lime           fifteen to thirty grains

Mucilage of gum Arabic   one fl. ounce

Syrup of orange peel   half fl. ounce

Mix. To be applied by means of a piece of sponge or camel's hair brush, to ulcers in the mouth. *Angelot.*

R. Chlorinated lime           three drachms

Distilled water

Alcohol, each,           two fl. ounces

Oil of roses           four drops

Dissolve, and filter. A teaspoonsful in a glass of water, to correct fetid breath.

*Chevallier.*

#### CHLORINATED LIME DENTIFRICE.

R. Chloride of lime           four grains

Powdered red coral       two drachms

A toothbrush, slightly wetted, to be dipped in this powder, and rubbed on the teeth.

*Magendie.*

#### COMPOUND INJECTION OF CHLORINATED LIME.

R. Chloride of lime           two drachms

Decoction of rhatany thirteen fl. ounces

Dissolve, and filter. As an injection in ozena, three or four times a-day.

*Detmold.*

#### OINTMENT OF CHLORINATED LIME.

R. Chlorinated lime           one drachm

Lard                       one ounce

Rub together. In scrofulous swellings.

*Cima.*

R. Chlorinated lime           half a drachm.

Lard                       one ounce

Rub together. In goitre. *Werneck.*

R. Chlorinated lime

Borate of soda, each,   one drachm

Lard                       one ounce

Rub together thoroughly. In chilblains.

*Trusen.*

R. Chlorinated lime           one drachm

Powdered digitalis       two drachms

Vinegar                   two fl. scruples

Lard                       one ounce

Rub together. In indolent glandular tumors. *Phæbus.*

#### LINIMENT OF CHLORINATED LIME.

R. Chlorinated lime           half a drachm

Rub in a glass mortar, adding gradually,

Rose water           one fl. ounce

and when quite clear

Oil of almonds           one fl. ounce

As an application in tinea capitis.

*Trusen.*







ness of the stomach. Dose, one to two tablespoonfuls. *Ellis.*

#### ANTACID MIXTURE.

R. Lime water four ounces  
Solution of potassa  
Syrup of orange peel, each, one ounce  
Mix. Dose, a tablespoonful in a cup of water. Found useful in dyspepsia and heartburn. *Cooley.*

#### LIME OINTMENT.

R. Lime one ounce  
Linseed oil three ounces  
Mucilage of quince seeds two ounces  
Mix well. In burns and scalds. *Radiis.*

R. Slaked lime one drachm  
Carbonate of soda two drachms  
Extract of opium ten grains  
Lard two ounces

Rub well together. In obstinate cutaneous affections. *Bielt.*

R. Slaked lime  
Laudanum, each, half a drachm  
Cucumber ointment four drachms  
Rub well together. As an application to hemorrhoidal tumors. *Guibourt.*

R. Lime  
Sulphur, each, two ounces  
Muriate of soda one ounce  
Lard one pound  
Olive oil two pounds

Mix well. As a friction in itch. *Ferrara. Ph.*

### CALCIS CARBONAS.

#### CARBONATE OF LIME.

##### PRECIPITATED CARBONATE OF LIME.

R. Water of chloride of calcium five parts  
Carbonate of soda three parts  
Dissolved in four times its weight of distilled water. Mix. Wash the precipitate three times, and dry. *Dub. Ph.*

#### PREPARED CHALK.

R. Chalk at will  
Add a little water, and rub to fine powder. Throw into a vessel of water, and stir, pour off liquid, while yet turbid, into another vessel, and permit chalk to subside; pour off water, and dry the precipitate. *U. S. Ph.*

#### PREPARED OYSTER SHELL.

R. Oyster shell at will  
Free it from extraneous matter, wash with boiling water, reduce to powder, and proceed as with chalk. *U. S. Ph.*

As an antacid in bowel affections. In doses

of from ten to forty, or more grains, often repeated.

#### AROMATIC POWDER OF CHALK.

R. Prepared oyster shell two drachms  
Magnesia one drachm  
Powdered cinnamon one scruple  
Mix.

#### COMPOUND POWDER OF CHALK.

R. Prepared chalk half a pound  
Cinnamon four ounces  
Tormentil  
Gum Arabic, each, three ounces  
Long pepper half an ounce  
Beat separately into fine powder, and mix. *Lond. Ph.*

Stimulant, astringent, and antacid; useful in diarrhœa, with acidity, but without inflammation. Dose, ten to twenty grains.

#### COMPOUND POWDER OF CHALK WITH OPIUM.

R. Compound powder of chalk six ounces and a-half  
Powdered opium four scruples  
Mix well. *Lond. Ph.*

In diarrhœa of adults, in doses of ten or twenty grains, after each evacuation.

#### DENTIFRICE POWDER WITH CARBONATE OF LIME.

R. Powdered cuttlefish bone  
Peruvian bark  
myrrh equal parts  
Mix. *Pierquin.*

R. Powdered cuttlefish bone six ounces  
red coral two ounces  
dragon's blood  
red saunders  
orris root, each, one ounce and a-half  
cinnamon  
alum, each, half an ounce  
Mix. *Par. Ph.*

#### LOZENGES OF CHALK.

R. Prepared chalk four ounces  
Gum Arabic, powdered, one ounce  
Nutmeg " one drachm  
Sugar " six ounces  
Rub together till mixed; mix with sufficient water to form mass; divide into lozenges of ten grains each. *U. S. Ph.*

As a mild antacid astringent, in diarrhœa.

#### DENTIFRICE ELECTUARY.

R. Powdered red coral four ounces  
cuttlefish bone  
cinnamon, each, one ounce  
cochineal half an ounce  
alum half a drachm



oil of cloves six drops  
honey ten ounces  
Mix. *Taddei.*

CHALK MIXTURE.

R. Prepared chalk half an ounce  
Sugar  
Powdered gum Arabic, each, two drachms  
Cinnamon water  
Water, each, four fl. ounces  
Rub them together until thoroughly mixed  
*U. S. Ph.*

A tablespoonful, as may be required.

R. Prepared chalk one drachm and a-half  
White sugar  
Gum Arabic, each, sufficient  
Oil of cinnamon two drops  
Tincture of opium forty to fifty drops  
Distilled water four fl. ounces  
Mix. A tablespoonful, every two hours, in diarrhoea and dysentery.  
*Ellis.*

COMPOUND CHALK MIXTURE.

R. Chalk mixture five fl. ounces  
Aromatic confection one drachm  
Solution carbonate ammonia one fl. drachm  
Laudanum twenty minims  
Mix. A tablespoonful, occasionally, in diarrhoea.  
*Ainslie.*

CALCIS PHOSPHAS.

PHOSPHATE OF LIME.

R. Burnt bones, in powder one part  
Diluted muriatic acid  
Water, each, two parts  
Digest together for twelve hours, and filter. Add water of caustic ammonia, sufficient to precipitate phosphate of lime. Wash this, and dry.  
*Dub. Ph.*

BURNT HARTSHORN.

R. Burn pieces of hartshorn in an open vessel till quite white, and prepare them as directed for chalk.  
*Lond. Ph.*

Has been thought efficacious in mollities ossium and rachitis, but probably inert. Dose, twenty grains, or more.

DENTIFRICE POWDER OF PHOSPHATE OF LIME.

R. Phosphate of lime eighteen parts  
White bole  
Gum lac, each, twelve parts  
Cinnamon one part  
Myrrh two parts  
Powder well, and mix.  
*Van Mons.*

DENTIFRICE ELECTUARY OF PHOSPHATE OF LIME.

R. Powdered phosphate of lime two ounces and a-half  
gum lac one ounce  
cinnamon one drachm and a-half  
cloves one scruple  
Syrup of lemon sufficient  
Make a soft electuary.  
*Van Mons.*

CALENDULA.

MARYGOLD.

This is the officinal name of *Calendula officinalis*, a plant indigenous to Europe, and often cultivated in gardens in this country as a pot herb.

*Ser. Syst.* Syngen. necess. *Nat. Syst.* Asteraceæ. *Linn. Sp. Pl.* 1304. *Lindley, Flor. Med.* 466.

The parts used are the herb, flowers, and seed. Their smell is peculiar, and somewhat aromatic; the taste is feebly bitter, and a little acrid. It was formerly much used, but had fallen into disuse, when, of late years, it has again been brought forward as a remedy in cancerous affections.

EXTRACT OF MARYGOLD.

R. Marygold one part  
Tepid water four parts  
Macerate for twenty-four hours, boil for a quarter of an hour, express, boil residue with four parts of water, express, mix two decoctions, and evaporate to proper consistence.  
*Guibourt.*

PILLS OF THE EXTRACT OF MARYGOLD.

R. Extract of marygold  
cicuta  
Sub carbonate of iron, each, two drachms  
Powdered marygold one drachm and a-half  
Mix well, and divide into pills of two grains each. Five or six are to be taken three times a-day in schirrhus and cancerous affections.  
*Rust.*

LOTION OF EXTRACT OF MARYGOLD.

R. Extract of marygold  
cicuta, each, three drachms  
Cherry laurel water two ounces  
Tincture of opium half an ounce  
Mix. As a lotion to cancerous ulcerations.  
*Radiis.*

INFUSION OF MARYGOLD.

R. Marygold flowers three drachms  
Boiling water sufficient  
to obtain five ounces; infuse, and add to the strained liquid



Syrup of orange peel half an ounce  
Dose, two spoonfuls, three times a-day.

*Radiis.*

#### OINTMENT OF MARYGOLD.

R. Marygold flowers three drachms  
Water five fl. ounces  
Boil to three ounces, strain, and add  
Extract of marygold one drachm  
Tar two ounces  
Pyroligneous acid half an ounce  
Gum Arabic sufficient  
Mix. As an application to cancerous  
ulcers.

*Radiis.*

### CALOTROPIS.

#### MUDAR.

The *Calotropis gigantea* is a native of the East Indies, and is now naturalized in several of the West India Islands.

*Sex. Syst.* Pentand. monog. *Nat. Syst.* Asclepiadaceæ.

Brown, Tr. Wern. Soc. 1. 29. Griffith, Med. Bot. 456.

The part used is the bark of the root; this is of a whitish color, inodorous, but with a bitter, nauseous taste. It is purgative, alterative, and diaphoretic, and has been highly spoken of in the treatment of elephantiasis and lepra. Dose, three grains to thirty; in the latter dose it is emetic and purgative.

#### INFUSION OF MUDAR.

R. Mudar three drachms  
Boiling water eight ounces  
Infuse, and strain. As an alterative, one to two fl. drachms; emetic, two fl. ounces.  
*Cassanova.*

#### PILLS OF MUDAR.

R. Powdered mudar one drachm  
Honey sufficient  
Beat into mass, and divide into pills of four grains. Two or three to be taken a-day as alterative in lepra, &c. *Cassanova.*

#### OIL OF MUDAR.

R. Powdered mudar one drachm  
Olive oil seven ounces  
Infuse for half an hour over a water-bath, and strain. As an application to leprosy sores.  
*Cassanova.*

### CAMPHORA.

#### CAMPHOR.

A peculiar white, crystallizable, semi-transparent, volatile product of several plants, but that used in medicine is obtained from the *Camphora officinarum*, an evergreen tree, of considerable size, indigenous to China, and other Asiatic countries.

*Sex. Syst.* Enneand. monog. *Nat. Syst.* Lauraceæ.

Nees. Laurin. 88. Griffith, Med. Bot. 553.

It has narcotic, diaphoretic, sedative properties, and is employed in a variety of diseases, and used externally, as an anodyne. The dose is from three to ten grains. Camphor can be powdered by trituration, with a few drops of alcohol.

#### CAMPHOR POWDER.

R. Camphor nine grains  
Sugar one drachm  
Mix, and divide into nine powders.  
*Augustin.*

#### COMPOUND CAMPHOR POWDER

R. Powdered camphor  
Myrrh, each, two drachms  
Peruvian bark  
Chamomile, each, half an ounce  
Charcoal one ounce  
Mix. Dose two scruples, or more, in gangrene. *Rust.*

R. Powdered camphor  
Benzoic acid, each, six grains  
Sugar one drachm  
Mix, and divide into four powders. As an antispasmodic and sedative.

*Saunders.*

R. Powdered gum Arabic one ounce  
orange peel  
sugar, each, two drachms  
camphor ten grains  
opium five grains

Mix. A teaspoonful every hour.  
*Ammon.*

Said to have been useful in cholera.

#### CAMPHOR PILLS.

R. Powdered camphor  
Sugar  
Starch  
Crumb of bread, each, one scruple  
Mix, and make twenty pills. *Augustin.*

#### COMPOUND CAMPHOR PILLS.

R. Powdered camphor twenty-four grains  
musk eight grains  
opium two grains  
syrup sufficient  
Beat into a mass, and divide into six pills. In hospital gangrene. *Dupuytren.*

#### PILLS OF CAMPHOR AND LACTUCARIUM.

R. Camphor  
Lactucarium, each, fifty grains  
Mix, and make twenty pills. Four to six daily, as an aphrodisiac. *Ricord.*

#### PILLS OF CAMPHOR AND MUSK.

R. Camphor one scruple  
Musk ten grains



Gum ammoniac two scruples  
Opium four grains  
Mix, and divide into four grain pills.  
Dose, four or five in a day, in nervous disorders. *Richard.*

## CAMPHOR WATER.

R. Camphor two drachms  
Alcohol forty minims  
Carbonate of magnesia one drachm  
Distilled water two pints  
Rub the camphor with the alcohol, then with the magnesia, and afterwards, with the water gradually added, filter. *U. S. Ph.*

Dose, one or two tablespoonfuls, to be repeated as occasion may require.

## CAMPHOR WATER AND LAUDANUM.

R. Camphor water four fl. ounces  
Compound spirit of lavender two drachms

Tincture of opium forty to fifty drops  
Mix. A tablespoonful, every two hours, in diarrhoea and dysentery. *Ellis.*

## CAMPHOR WATER AND NITRIC ACID.

R. Camphor water four fl. ounces  
Nitric acid four drops  
Tincture of opium forty to fifty drops  
Mix. In the same dose, and in the same complaints, as the last. *Ellis.*

R. Camphor water ten fl. drachms  
Laudanum forty drops  
Spirit of sulphuric ether one fl. drachm

Syrup of poppies one fl. drachm  
Mix. To prevent a paroxysm of intermittent fever. To be given just before its accession. *Gregory.*

## CAMPHOR WATER AND HOFFMANN'S ANODYNE.

R. Camphor water four fl. ounces  
Hoffmann's anodyne two fl. drachms  
Mix. A desertspoonful, every hour or two, in some demulcent drink. In nervous affections and sleeplessness in fevers. *Ellis.*

## INFUSION OF CAMPHOR.

R. Camphor one ounce  
Boiling water eight fl. ounces  
Pour water on camphor, and let stand till cold, keep in well-stopped bottle. Dose, a tablespoonful, every hour or two, in nervous per vigiliam and nervous irritability. *Ellis.*

## CAMPHOR MIXTURE.

R. Powdered camphor one drachm  
gum Arabic

white sugar, each,  
one drachm and a-half

Tincture of opium forty drops  
Mint water four fl. ounces

Mix. A tablespoonful every two hours, in low conditions of the system. *Ellis.*

R. Powdered camphor one scruple  
gum Arabic one drachm  
sugar sufficient  
Cinnamon water one ounce

Mix. To relieve the pain in dysmenorrhœa. One-half to be given as soon as pain is felt, if not relieved in an hour or two, the remainder to be taken. *Dewees.*

R. Powdered camphor half a drachm  
gum Arabic two drachms  
sugar three drachms  
vinegar half an ounce  
water six ounces

Mix. As a stimulant. *Phœbus.*

## EMULSION OF CAMPHOR.

R. Camphor  
Olive oil, each, one drachm  
Powdered gum Arabic half an ounce  
Water four fl. ounces

Triturate the camphor and oil together, until uniformly mixed; add the gum, and then the water gradually, triturating till the emulsion is formed. *W. Procter.*

## CAMPHOR AND MILK.

R. Camphor one drachm  
Boiling milk four fl. ounces  
Dissolve. To be used as the last. *Ellis.*

## CAMPHOR WITH MYRRH.

R. Powdered camphor one drachm  
myrrh half a drachm  
sugar two drachms  
water six fl. ounces

Mix. A tablespoonful every two hours. *Ellis.*

## TINCTURE OF CAMPHOR.

R. Camphor four ounces  
Alcohol two pints  
Dissolve. *U. S. Ph.*

Chiefly used as an anodyne embrocation.

## TINCTURE OF CAMPHOR AND SAFFRON.

R. Camphor one ounce  
Saffron one scruple  
Alcohol one pint

Macerate for a day, and filter. *Guibourt.*

As a lotion, or fomentation.

## ETHEREAL TINCTURE OF CAMPHOR.

R. Camphor one ounce  
Sulphuric ether eight fl. ounces  
Dissolve. *Swedjaur.*



Dose, twenty to thirty drops, in wine, as a stimulant and antispasmodic.

#### MIXTURE OF TINCTURE OF CAMPHOR.

R. Tincture of camphor  
Goulard's extract                      equal parts  
Mix. As a lotion, several times a-day, to parts disposed to ulcerate by constant decubitus. *Tott.*

#### MIXTURE OF CAMPHOR AND CHLOROFORM.

R. Camphor                                      one drachm  
Chloroform                                  half fl. drachm  
Mix, and add  
Yolk of egg                                      one  
Triturate, and add gradually  
Water    six fl. ounces  
As a stimulant. Dose, two fl. drachms to fl. half ounce. *Keating.*

#### MIXTURE OF CAMPHOR AND ETHER.

R. Camphor water                              one ounce  
Sulphuric ether                              one scruple  
Syrup    one ounce  
Hot water                                      two pints  
Mix. In spoonful doses in gout in the stomach. *Pierquin.*

R. Camphor                                      one scruple  
Sulphuric ether                              two drachms  
Dissolve, and add  
Laudanum                                      twenty drops  
Cinnamon water                              six ounces  
Mix. A spoonful every three or four hours as a stimulant. *Augustin.*

#### CAMPHOR LINIMENT.

R. Camphor                                      half an ounce  
Olive oil    two fl. ounces  
Dissolve the camphor in the oil. *U. S. Ph.*  
As an anodyne embrocation.

#### COMPOUND CAMPHOR LINIMENT.

R. Camphor                                      two ounces and a-half  
Solution of ammonia,                      seven fl. ounces and a-half  
Spirit of lavender                              one pint  
Mix solution of ammonia with the spirit; then, from a glass retort, by a slow fire, distil a pint; dissolve camphor in the distilled fluid. *Lond. Ph.*

A rubefacient and anodyne embrocation.

R. Camphor                                      one drachm  
Oil of chamomile  
Wine of opium, each,                      two drachms  
Oil of hyoseyamus                              one ounce  
Mix. *Augustin.*

#### LINIMENT OF CAMPHOR AND VINEGAR.

R. Tincture of camphor                      three fl. ounces  
Acetic acid                                      one fl. drachm

Mix. As a lotion or embrocation. *Ellis.*

#### ETHEREAL CAMPHOR LINIMENT.

R. Soft soap  
Alcohol, each,                                  one ounce  
Dissolve, and add  
Camphor    half an ounce  
Dissolved in  
Sulphuric ether                              one fl. ounce  
As an embrocation. *Saunders.*

#### WARD'S ESSENCE FOR HEADACHE.

R. Camphor                                      two ounces  
Rectified spirit                              fourteen ounces  
Water of ammonia                              two ounces  
Oil of lavender                              half an ounce  
Mix. As an application to the forehead, &c., in headache. *Redwood.*

#### CAMPHOR WATER MIXTURE.

R. Camphor water  
Solution of acetate of ammonia, each,      half fl. ounce  
Antimonial wine  
Tincture of opium, each, twenty drops  
Mix. To be taken at bedtime, as a stimulating diaphoretic. *Ellis.*

#### CAMPHOR CLYSTER.

R. Camphor                                      one drachm  
Olive oil    two ounces  
Mix. An excellent enema in cases of ascarides, to be used for three or four successive nights. *Ellis.*  
R. Common clyster                              two pints  
Camphor    two drachms  
Yolk of egg                                      one  
Mix. A fourth part to be injected at a time, to be employed as occasion may require in low fevers. *Chaussier.*

#### CAMPHOR OINTMENT.

R. Lard  
Suet  
Oil of bayberries  
Wax, each,                                      half an ounce  
Melt together, and add  
Camphor    one drachm  
Said to be efficacious as an application to chilblains. *Radius.*  
R. Fresh butter                              three drachms  
Spermaceti                                      two drachms  
White wax    three ounces  
Melt together, and add  
Powdered camphor  
Oxide of zinc, each,                              three grains  
To be applied in chronic inflammation of eyelids, to the affected parts, at night. *Maternite.*  
R. Camphor                                      one drachm  
Basilicon ointment                              one ounce



Soap half an ounce  
Powdered mustard one scruple  
Mix. As an external application in lum-  
bago. *Ferriar.*

## CAMPHOR COLLYRIUM.

R. Camphor water six fl. ounces  
Solution acetate of  
ammonia two fl. ounces  
Mix. A mild astringent, stimulant in in-  
flammation of the eyes. *Ellis.*

LINIMENT OF CAMPHOR AND TURPEN-  
TINE.

R. Oil of turpentine fifteen parts  
Camphor one part  
Mix. As a stimulating embrocation.  
*Beral.*

## NITRATE OF CAMPHOR.

R. Nitric acid at will  
Camphor sufficient  
to saturate. Decant. Ten drops every  
two hours on sugar, or in syrup, in chronic  
pleurisy. Also used as a friction in rheu-  
matism and palsy. *Van Mons.*

## CAMPHORATED COUGH MIXTURE.

R. Camphor half a scruple  
Mucilage of gum Arabic three ounces  
Syrup of mallows one ounce and a-half  
Mix. A teaspoonful every two hours.  
*Saunders.*

## CANELLA.

## CANELLA.

This is the bark of the *Canella alba*, a large  
tree, native of Jamaica and other West India  
islands.

*Sex. Syst.* Dodecand. monog. *Nat. Syst.* Me-  
liaceæ.

Swartz. Trans. Linn. Soc. 1. 96. Griffith, Med.  
Bot. 181.

The part used is the bark, which is of a pale  
orange-yellow colour, with an aromatic odor, and  
warm, bitterish, pungent taste. It is principally  
used as an adjuvant to other remedies. The dose,  
as a stimulant, is from ten grains to a scruple.

## CANNABIS.

## HEMP.

The *Cannabis sativa* is an annual plant, a na-  
tive of Asia, but generally cultivated in Europe  
and the United States. The East Indian plant  
is more powerful in its action on the system than  
that growing in more temperate climates.

*Sex. Syst.* Diœc. pentand. *Nat. Syst.* Canna-  
binaceæ.

Linn. Sp. Pl. 1457. Griffith, Med. Bot. 572.

The parts used are the seeds, the leaves, and  
the stems; the two latter are covered with a resin-

ous exudation, which is narcotic and intoxicating.  
They have been recommended in a variety of  
diseases of a formidable character.

## EXTRACT OF HEMP.

R. Dried hemp tops at will  
Boil in alcohol until all the resinous mat-  
ter is dissolved; strain, and evaporate to  
proper consistence. *O' Shaughnessy.*

Has been given with success in tetanus, hydro-  
phobia, cholera, &c. Dose, ten grains. To be  
increased as occasion may require.

## TINCTURE OF HEMP.

R. Extract of hemp twenty-four grains  
Diluted alcohol one fl. ounce  
Dissolve. In tetanus, one drachm every  
half hour, till some effect is produced. In  
cholera, ten drops every half hour.  
*O' Shaughnessy.*

## EMULSION OF HEMP-SEED.

R. Hemp-seed bruised six drachms  
Gum Arabic one drachm  
Parsley water five fl. ounces  
Rub together so as to form emulsion, strain,  
and add

Syrup of balsam of Peru one ounce  
A spoonful every hour in gonorrhœa.

*Radius.*

## RESIN OF HEMP.

R. Dried hemp tops, bruised, at will  
Macerate several times in warm water, and  
then in a solution of carbonate of soda;  
afterwards, wash well with water, pressing  
after each operation. Dry, and digest in  
rectified spirit, to which milk of lime, con-  
taining an ounce of lime to each pound of  
hemp tops, has been added. Filter, add a  
slight excess of sulphuric acid; again filter,  
distil off most of the spirit, add to the resi-  
due three or four times its bulk of water,  
evaporate the rest of the spirit, decant the  
water, wash and dry. Dose, two-thirds of  
a grain. *Smith.*

## CANTHARIS.

## SPANISH FLY.

The *Cantharis vesicatoria* is an insect inhabiting  
those parts of Europe in which the vine grows  
naturally. It has a fetid odor and an acrid  
taste. The body is oblong, of a golden green  
color. Spanish flies are stimulant, diuretic,  
rubefacient, and vesicant. They are used inter-  
nally in dropsies, and in various disorders of the  
urinary organs; externally, to irritate or vesicate.  
Several species are natives of the United States,  
which are not inferior in their powers to the  
foreign insect. Dose, one to two grains.

## POWDER OF CANTHARIDES AND SAVINE.

R. Powdered cantharides two grains



savine one drachm  
Mix, and divide into four powders. One to be taken at night, as an emmenagogue, watching its effects closely. *Ellis.*

#### POWDER OF CANTHARIDES AND CAMPHOR.

R. Powdered cantharides four grains  
camphor eight grains  
Sugar of milk three drachms  
Mix, and divide into eight powders, one to be given twice a-day as a diuretic.

*Augustin.*

R. Powdered cantharides eighteen grains  
opium  
camphor, each,  
thirty-six grains  
Mix, and form into thirty-six pills. One to be taken at bedtime, in cases of impotency from general debility. *Ellis.*

#### EXTRACT OF CANTHARIDES.

R. Cantharides at will  
Digest, in four or five times their weight of alcohol; filter, repeat operation, unite the two tinctures, and evaporate. *Soubéiran.*

#### PILLS OF CANTHARIDES AND IRON.

R. Powdered cantharides one scruple  
Subcarbonate of iron two scruples  
Extract of liquorice one drachm  
Form a mass, and divide into pills of one grain each. Dose, three or four, three times a-day, gradually augmenting the dose. As a diuretic. *Meissner.*

#### PILLS OF CANTHARIDES AND CAPSICUM.

R. Powdered cantharides five grains  
capsicum eight grains  
camphor one scruple  
guaiaicum one drachm

Tincture of colocynth sufficient  
Form mass, and divide into eighty pills. Dose, one, gradually increasing to four or five, twice a-day, in paraplegia. *Sundelin.*

#### INFUSION OF CANTHARIDES.

R. Cantharides one scruple  
Boiling water sufficient  
to obtain three ounces of infusion, after digesting for half an hour, and straining. *Soubéiran.*

#### EMULSION OF CANTHARIDES.

R. Cantharides one scruple  
Sweet almonds six drachms  
Sugar one ounce  
Beat together, and gradually rub up with  
Lime water ten ounces  
Strain. A spoonful every two or three hours in neuralgia, and in catarrh of the bladder. *Berends.*

R. Cantharides half a drachm  
Sweet almonds one ounce  
Sugar half an ounce  
Beat together, and make an emulsion with  
Hot water ten ounces  
Strain. A spoonful, every two or three hours, in hooping-cough. *Hufeland.*

#### VINEGAR OF CANTHARIDES.

R. Powdered cantharides two ounces  
Acetic acid one pint  
Macerate for eight days, express, and filter. *Lond. Ph.*  
As an epispastic, or as a rubefacient lotion.

#### TINCTURE OF CANTHARIDES.

R. Cantharides, bruised, one ounce  
Diluted alcohol two pints  
Macerate for fourteen days, express, and filter. *U. S. Ph.*

A very convenient mode of administering cantharides. Dose, from twenty drops to a fluid drachm. Also useful as a rubefacient.

#### CAMPHORATED TINCTURE OF CANTHARIDES.

R. Tincture of cantharides half an ounce  
Camphor three drachms  
Alcohol two ounces  
Mix. As a rubefacient. *Pierquin.*

#### COMPOUND TINCTURE OF CANTHARIDES.

R. Cantharides one drachm  
Mustard  
Black pepper  
Camphor, each, half an ounce  
Garlic one head  
Vinegar six ounces  
Alcohol twelve ounces  
Macerate for some days, express, and filter. Said to be very effectual as a rubefacient lotion in rheumatism, &c. *Foy.*

R. Tincture of cantharides.  
Extract of rosemary, each,  
one drachm

Solution of carbonate  
of potassa one drachm and a-half  
Water four ounces  
Mix. As a lotion to promote the growth of hair, where it has been lost from an eruption, or from fever. *Sachse.*

#### TINCTURE OF CANTHARIDES AND GUAIAICUM.

R. Powdered cantharides one drachm  
guaiaicum forty-five grains  
Contused rhubarb one drachm and a-half  
shell lac half a drachm  
Alcohol twelve fl. ounces  
Digest, and strain. Dose, from thirty to



fifty drops, night and morning, in some demulcent. In gleet. *Ellis.*

#### ETHEREAL TINCTURE OF CANTHARIDES.

R. Powdered cantharides one part  
Acetic ether sixteen parts  
Digest for eight days, express, and filter. *Foy.*

As a rubefacient and vesicant lotion; it is very energetic.

#### ETHEREAL EXTRACT OF CANTHARIDES.

R. Powdered cantharides at will  
Sulphuric ether sufficient  
treat the powder with the ether in a displacement funnel, properly arranged, until exhausted, and distil off the ether. Flies yield 8 per cent. of this extract. It is employed for increasing the activity of blistering plasters, and for admixture with wax to make blistering tissue paper. *Soubeyran.*

#### CAMPHORATED ETHEREAL TINCTURE OF CANTHARIDES.

R. Powdered cantharides one part  
Nitrate of camphor three parts  
Alcohol fifty-six parts  
Digest and express, and filter at the end of a few days. Advised in dropsy, in the dose of ten to twenty drops, in white wine. *Van Mons.*

#### OIL OF CANTHARIDES.

R. Cantharides one part  
Olive oil eight parts  
Digest for six hours on a water-bath, express, and filter. *Guibourt.*

An active vesicant.

R. Cantharides one part  
Oil of rosemary eight parts  
Digest for fifteen days, express, and filter. *Beral.*

More active than the preceding.

#### LINIMENT OF CANTHARIDES.

R. Powdered cantharides one ounce  
Oil of turpentine half a pint  
Digest for three hours over a water-bath, and strain. *U. S. Ph.*

An active stimulant when applied to the skin; used with success in the low stages of typhus fevers. Requires caution in its use.

#### LINIMENT OF CANTHARIDES AND CAMPHOR.

R. Camphor three drachms  
Dissolve in  
Liniment of cantharides half fl. ounce  
Add,  
Strong mercurial ointment  
Simple ointment, each, one ounce

Mix, and form liniment. In low stages of fever, where the object is to arouse the system, and to affect it with mercury; also as an application to tumors. *Ellis.*

#### DECOCTION OF CANTHARIDES IN TURPENTINE.

R. Cantharides in powder four ounces  
Oil of turpentine twelve ounces  
Mix in a flask, place this in a salt-bath, and heat for four hours; then dissolve with oil of turpentine until twelve fl. ounces are obtained. *Dr. Hartshorne.*

#### LINIMENT OF CANTHARIDES AND SOAP.

R. Tincture of cantharides half an ounce  
Liquid ammonia an ounce and a half  
Turpentine soap two ounces  
Spirit of turpentine one ounce  
of camphor eight ounces

Mix. A powerful rubefacient. *Niemann.*

R. Cantharides, in fine powder, one pound  
Yellow wax  
Resin  
Lard, each, eight ounces  
Melt the wax, resin, and lard, together, and add the cantharides, stirring constantly till cold. *U. S. Ph.*

The common blistering plaster.

#### CANTHARIDES OINTMENT.

R. Cantharides, in powder, two ounces  
Distilled water half a pint  
Resin cerate eight ounces  
Boil down the cantharides in the water to one-half, strain, add cerate, and evaporate to proper consistence. *U. S. Ph.*

As a stimulating dressing to blisters.

R. Spanish flies, in fine powder,  
Olive oil, each, six ounces  
Oil of turpentine three ounces  
Yellow wax

Resin, each, four ounces

Mix the first three ingredients, place on a water-bath, and continue heat till the turpentine has nearly evaporated, stirring occasionally. Then add the wax and resin, previously melted together, and heat till well incorporated; remove from fire, and stir till cold. *W. Procter.*

R. Powdered Spanish flies four ounces  
Boiling water twenty fl. ounces  
Macerate for twelve hours, and evaporate by a gentle heat to the consistence of syrup; add  
Yellow wax four ounces  
Resin  
Olive oil







and rubefacient. Dose, five to ten grains in powder or pill.

#### CAYENNE PEPPER PILLS.

R. Powdered cayenne pepper one drachm  
Crumb of bread  
Distilled water, each, sufficient  
Rub into mass, and form twelve pills.  
One, three or four times a-day. In debility of the stomach, especially from intemperance. *Ellis.*

R. Powdered cayenne pepper one scruple  
Extract of gentian one drachm  
Powdered gentian sufficient  
Form mass, and divide into sixty pills.  
In chronic dyspepsia. *Radius.*

#### CAYENNE PEPPER LOZENGES.

R. Cayenne pepper three drachms  
Ginger  
Horseradish, each, two drachms  
Sugar one ounce  
Conserve of roses one drachm  
Mix, and make thirty lozenges. One to be chewed occasionally in catarrhal deafness. *Fosbroke.*

#### INFUSION OF CAYENNE PEPPER.

R. Cayenne pepper three drachms  
Boiling water half a pint  
Infuse, and strain. Dose, half a fl. ounce.

#### CAYENNE PEPPER GARGLE.

R. Tincture of cayenne pepper half fl. ounce  
Rose water eight fl. ounces  
Mix.

R. Powdered cayenne pepper two tablespoonfuls  
Common salt a teaspoonful  
Boiling vinegar water, each, half a pint  
Infuse for an hour, and strain. In bad cases of scarlatina. *Chapman.*

R. Powdered cayenne pepper half an ounce  
Magnesia one drachm  
Boiling water vinegar, each, eight ounces  
Infuse, and strain. Used in same cases, and in same manner as above. *Headley.*

#### TINCTURE OF CAYENNE PEPPER.

R. Cayenne pepper one ounce  
Diluted alcohol two pints  
Macerate for fourteen days, and filter. *U. S. Ph.*

As a stimulant in low states of fever with gastric insensibility. Dose, one to two fl. drachms. Also used in a diluted state as gargle.

#### TINCTURE OF CAYENNE PEPPER AND CANTHARIDES.

R. Cayenne pepper one drachm  
Cantharides ten drachms  
Alcohol one pint  
Digest for two days, and filter. Dose, ten drops. *Coxe.*

#### VINEGAR OF CAYENNE PEPPER.

R. Cayenne pepper one part  
Vinegar six parts  
Digest for some days, and filter. In doses of half a drachm to a drachm, in passive dropsies. *Beral.*

#### CAYENNE PEPPER CATAPLASM.

R. Powdered cayenne pepper one ounce  
Ground mustard  
Common soap, each, three ounces  
Alcohol sufficient  
Mix. An active rubefacient. *Ammon.*

#### LOTION OF CAYENNE PEPPER.

R. Tincture of cayenne pepper camphor, each, four ounces  
Solution of ammonia two ounces  
Mix. A very powerful rubefacient. *Ammon.*

### CARBO ANIMALIS.

#### ANIMAL CHARCOAL.

This article is prepared by charring animal substances, usually bones. It requires purification for internal use.

#### PURIFIED ANIMAL CHARCOAL.

R. Animal charcoal one pound  
Muriatic acid  
Water, each, twelve fl. ounces  
Mix the acid and water, and pour on charcoal; digest for two days, occasionally stirring. Decant, and wash the charcoal several times with water and dry it. *U. S. Ph.*

Weise advises it made thus,

R. Lean of beef or mutton two parts  
Broken bones one part  
Char by a moderate fire in a coffee-roaster, let cool and pulverize.

Has been given with advantage in scrofulous diseases, chronic glandular swellings, &c., in doses of ten to twenty grains.

#### POWDER OF ANIMAL CHARCOAL.

R. Animal charcoal four grains  
Sugar one drachm  
Mix. To be taken morning and evening, in scrofula. *Weise.*

R. Animal charcoal six grains  
Burnt sponge twelve grains



Powdered liquorice root half a drachm  
Mix, and form six powders, one to be taken  
morning and night, in scirrhus goitre.

*Riecke.*

R. Animal charcoal four grains

Powdered liquorice root four scruples  
Mix, and divide into eight powders, one to  
be taken morning and evening, dry; a little  
water drank afterwards. After the eight  
powders are taken, increase the dose, by  
half a grain at a time, to four grains. In  
scirrhus tumors, in mammae. A spare  
diet to be observed.

*Weise.*

R. Animal charcoal one scruple

Subcarbonate of iron

Sugar, each, two scruples

Mix, and divide into ten powders. One  
three or four times a-day in incipient hy-  
pertrophy of the womb or ovaries. *Clarus.*

#### OINTMENT OF ANIMAL CHARCOAL.

R. Animal charcoal one part  
Lard two parts

Rub together. *Giordano.*

R. Animal charcoal half an ounce

Ointment of mallows six drachms

Mix. As a friction to glandular engorge-  
ments. *Radius.*

### CARBO LIGNI.

#### CHARCOAL

Is employed in medicine as an antiseptic and  
absorbent, and has been found useful in dyspepsia,  
&c. It also enters into the composition of poulti-  
ces, &c.

#### CHARCOAL DENTIFRICE POWDER.

R. Powdered charcoal four parts

Peruvian bark two parts

myrrh one part

Mix. *Phœbus.*

R. Powdered charcoal three drachms

Peruvian bark two drachms

calamus one drachm

myrrh half a drachm

Oil of cloves three drops

Mix. *Sartori.*

R. Powdered charcoal

red saunders

orange peel

Peruvian bark

cremor tartar, each,

one drachm

Oil of bergamot

twenty drops

Mix. *Schubarth.*

#### POWDER OF CHARCOAL AND QUASSIA.

R. Powdered charcoal six drachms

quassia

Magnesia, each, one drachm

Mix. Dose, a teaspoonful, two or three  
times a-day, in pyrosis. *Heim.*

#### CHARCOAL LOZENGES.

R. Powdered charcoal

sugar, each, one ounce

chocolate three ounces

Mucilage of tragacanth sufficient

Beat together, and form lozenges of eigh-  
teen grains. For fetid breath.

*Chevallier.*

#### CHARCOAL ELECTUARY.

R. Powdered charcoal

Carbonate of soda, each, one drachm

Electuary of senna two ounces

Mix. Two or three teaspoonfuls a-day in  
obstinate constipation. *Radius.*

R. Powdered charcoal

orris root

Peruvian bark, each,

one scruple

Tincture of myrrh half a drachm

Clarified honey sufficient

Mix. Used as a dentifrice. *Radius.*

#### OINTMENT OF CHARCOAL.

R. Powdered charcoal

Storax ointment, each, one ounce

Camphor

Myrrh, each, two drachms

Spirit of turpentine sufficient

Rub well together. As a dressing to  
fetid ulcers. *Rust.*

#### CHARCOAL CATAPLASM.

R. Freshly prepared charcoal sufficient

Mix with

Simple cataplasm in a tepid state.

*Dub. Ph.*

As an application to foul and gangrenous  
ulcers.

R. Powdered charcoal half an ounce

Cataplasm of chamomile five pounds

Mix. As above. *Brera.*

#### CHARCOAL SUPPOSITORY.

R. Cork charcoal

Wax, each, one part

Fresh butter two parts

Mix. As a suppository in hemorrhoids.  
*Cadet.*

### CARBO MINERALIS.

#### MINERAL COAL.

#### ANTHRACOKALI.

R. Carbonate of potassa six ounces

Lime three and a-half ounces



Water four pints  
Proceed as in making liquor potassa, and concentrate the clear solution, by boiling in an iron pot, till an oily looking liquid remains, then stir in five ounces of finely powdered mineral coal; remove from the fire, and continue stirring till the whole is reduced to an uniform powder, which is to be immediately put in small, well-stopped bottles. *Polya.*

Much recommended in cutaneous diseases, scrofula, chronic rheumatism, &c. Dose, two grains, twice or thrice a-day.

## POWDER OF ANTHRACOKALI.

R. Anthracokali two grains  
Powdered liquorice five grains  
Mix. Two to four times a-day. *Polya.*

## COMPOUND POWDER OF ANTHRACOKALI.

R. Anthracokali two grains  
Washed sulphur three to five grains  
Powdered liquorice two to three grains  
Mix. In psora, &c. *Polya.*

R. Anthracokali two grains  
Calomel sixth of a grain  
Powdered liquorice three grains  
Mix. In syphilitic eruptions. *Polya.*

R. Anthracokali  
Golden sulphuret of antimony half a grain  
Powdered liquorice three grains  
Mix. *Polya.*

## SULPHURETTED ANTHRACOKALI.

R. Prepared like the simple anthracokali, adding four drachms of sulphur to the coal, and dissolving, &c., as above. The dose and uses are the same. *Polya.*

## CARBONIS IODIDUM.

## IODIDE OF CARBON.

R. Tincture of iodine two fl. ounces  
Solution of potassa sufficient  
Add the solution of potassa to the tincture, till the latter loses its color, then add  
Distilled water sufficient  
to precipitate the iodide, separate and dry by a gentle heat.

## OINTMENT OF IODIDE OF CARBON.

R. Iodide of carbon half a drachm  
Simple ointment six drachms  
Rub together. As an application to enlarged glands, and obstinate cutaneous affections. *Litchfield.*

## CARDAMINE.

## CUCKOO FLOWER.

This is the *Cardamine Pratensis*, a small herbaceous plant, a native of Europe, and also found in the more northern parts of America.

*Ser. Syst.* Tetrad. siliquos. *Nat. Syst.* Brassicaceæ.

Linn. Sp. Pl. 915. Baker, Med. Trans. 1. 442.

The part employed is the flowers, which are said to be diuretic and antispasmodic. It is not used in this country, and appears to be little deserving of notice.

## CARDAMOMUM.

## CARDAMOM.

Under the name of Cardamom a variety of aromatic capsules have been used in medicine, all possessing the same properties and somewhat the same appearance. That recognized by the U. S. Pharm. is the product of the *Elletaria Cardamomum*, a native of Malabar.

*Ser. Syst.* Monand. monog. *Nat. Syst.* Zingiberaceæ.

Maton. Trans. Linn. Soc. x. 254. Griffith, Med. Bot. 633.

The part used is the seeds; these are angular, somewhat rough, of a reddish or brownish colour, with an agreeable fragrant odour, and a warm, aromatic taste. They are seldom used alone, being chiefly employed in compound preparations.

## AROMATIC POWDER.

R. Ginger  
Cinnamon, each, two ounces  
Cardamom seeds  
Nutmeg, grated, each, one ounce  
Mix, and rub together into fine powder. *U. S. Ph.*

Stimulant and carminative. Dose, ten to thirty grains, in dyspepsia with flatulence.

## AROMATIC CONFECTION.

R. Aromatic powder five ounces and a-half  
Powdered saffron half an ounce  
Syrup of orange peel six ounces  
Clarified honey two ounces  
Rub together, and beat into a mass. *U. S. Pharm.*

Used as a stimulant to the stomach, in doses of ten to sixty grains.

## TINCTURE OF CARDAMOM.

R. Cardamom, bruised, four ounces  
Diluted alcohol two pints  
Macerate for fourteen days, express, and filter. *U. S. Ph.*

A pleasant aromatic stimulant. Dose, one to two fl. drachms.



## COMPOUND TINCTURE OF CARDAMOM.

R. Powdered cardamom  
                   caraway, each, two drachms  
   and a-half  
                   cochineal           one drachm  
                   cinnamon       five drachms  
           Raisins                   five ounces  
           Proof spirit           two pints  
 Macerate for fourteen days, express, and  
 filter. *Lond. Ph.*

An agreeable aromatic and carminative, in the  
 dose of one or two fl. drachms.

## SYRUP OF CARDAMOM.

R. Cardamom water           one part  
       Sugar                   two parts  
 Dissolve, and boil to proper consistence.  
*Taddei.*

## ETHEREAL EXTRACT OF CARDAMOM.

R. Cardamom in powder           at will  
       Sulphuric ether           sufficient  
 Treat, till twice the weight of the seed  
 is obtained. Evaporate spontaneously un-  
 til deprived of ether. This extract is fluid,  
 and consists of volatile and fixed oil. It  
 may be used for aromatizing powders, or  
 as an addition to pills. *W. Procter.*

## CAROTA.

## CARROT.

The seeds and root of the common carrot,  
*Daucus carota*, a native of Europe, but now ex-  
 tensively naturalized in the United States, and  
 generally cultivated as an esculent in both coun-  
 tries.

*Sex. Syst.* Pentand. digyn. *Nat. Syst.* Apiaceæ.  
*Linn. Sp. Pl.* 348. *Griffith, Med. Bot.* 337.

The parts used are the seeds and root; the  
 former are aromatic, and have a pungent, bitter-  
 ish taste; the root of the wild carrot has a strong  
 smell, and an acrid, disagreeable taste; both  
 these are similar in their remedial effects, being  
 aromatic and diuretic. The root of the cultivated  
 variety is milder, and is not used internally, but  
 has much reputation as an external application to  
 foul ulcers, &c.

## INFUSION OF CARROT SEEDS.

R. Carrot seed half an ounce—one ounce  
       Boiling water           one pint  
 Infuse. To be taken during the day as a  
 diuretic, in dropsy and nephritic complaints.

## CATAPLASM OF CARROT ROOT.

R. Root of garden carrot           at will  
 Scrape down to a pulp. As an application  
 to foul and cancerous ulcers. *Wood.*

R. Root of garden carrot           at will  
 Boil till soft, and mash. *Dub. Ph.*

This acts merely as an emollient, and is not  
 possessed of any virtues beyond those of the  
 common poultice.

## EXTRACT OF CARROT ROOT.

R. Clarified juice of carrot root   at will  
 Evaporate on a water-bath to the consist-  
 ence of honey. *Swediaur.*

As an application to ulcerated cancers.

## CARROT OINTMENT.

R. Carrot root, grated,           one pound  
       Lard                   twenty ounces  
       Yellow wax           two ounces  
 Heat them together till the water of vege-  
 tation is driven off, and the fat has acquired  
 a yellow color; then strain for use.

*W. Procter.*

## CARTHAMUS.

## SAFFLOWER.

This is the florets of *Carthamus tinctorius*, a  
 native of the Levant and Egypt, but cultivated  
 in other parts of Europe and in the United  
 States.

*Sex. Syst.* Syngen. æqual. *Nat. Syst.* Aster-  
 aceæ.

*Linn. Sp. Pl.* 1162. *Griffith, Med. Bot.* 410.

The seeds were formerly much used in medi-  
 cine, but are at present seldom resorted to. The  
 florets are somewhat stimulant, and, in warm  
 infusion, diaphoretic, and are used as a substitute  
 for saffron.

## INFUSION OF SAFFLOWER.

R. Safflower                   two drachms  
       Boiling water           one pint  
 Infuse, and strain.

Used in domestic practice in the exanthemata,  
 to bring out the eruption.

## CARUM.

## CARAWAY.

Caraway seeds are the product of *Carum carui*,  
 an umbelliferous plant, a native of Europe, and  
 cultivated both there and in this country.

*Sex. Syst.* Pentand. digyn. *Nat. Syst.* Apiaceæ.  
*Linn. Sp. Pl.* 378. *Griffith, Med. Bot.* 318.

The seeds, the only part used, are of a brown-  
 ish color, with a fragrant but peculiar odor, and  
 an aromatic, warm taste. They are a pleasant  
 aromatic and carminative. Dose, in powder,  
 from a scruple to a drachm.

## CARAWAY WATER.

R. Caraway seeds,  
       bruised,           one pound and a-half  
       Proof spirits           seven fl. ounces  
       Water           two gallons  
 Mix. Distil a gallon. *Lond. Ph.*

## INFUSION OF CARAWAY.

R. Caraway seeds           two drachms  
       Boiling water           one pint  
 Infuse, and strain. *Wood.*

In flatulent colic of infants.



## SPIRIT OF CARAWAY.

R. Bruised caraway twenty-two ounces  
 Proof spirit one gallon  
 Water two pints  
 Mix, and distil one gallon. *Lond. Ph.*

As a carminative. In doses of one to two fl. drachms.

## OIL OF CARAWAY.

R. Caraway seeds at will  
 Water sufficient  
 Macerate for some time, distil, and separate the oil. *Van. Mons.*

## EMBROCATION OF CARAWAY.

R. Oil of caraway  
 peppermint, each, one scruple  
 olives one ounce  
 Wine of opium half a fl. drachm  
 Mix. As an embrocation on the abdomen of infants in colic. *Radius.*

## CARYOPHYLLUS.

## CLOVES.

Cloves are the unexpanded flowers of *Caryophyllus aromaticus*, a small tree, a native of the Molucca islands, and also cultivated in many other parts of the East Indies.

*Sex. Syst.* Icosand. monog. *Nat. Syst.* Myrtaceæ.

*Linn. Sp. Pl.* 735. *Griffith, Med. Bot.* 298.

Cloves are somewhat nail shaped, of a brown color, with a strong fragrant smell, and a permanent, pungent, aromatic taste. They are highly stimulating, and are given as a carminative, and excitant of the gastric functions, in the dose of five to ten grains, but are more used as an adjunct to various official compounds.

## INFUSION OF CLOVES.

R. Bruised cloves two drachms  
 Boiling water one pint  
 Macerate for two hours in a covered vessel, and strain. *U. S. Ph.*

Dose, from one to two fl. ounces.

## CLOVE WATER.

R. Bruised cloves one part  
 Water eight parts  
 Digest for twelve hours, and distil twelve parts. As a stimulant and carminative. *Cottureau.*

## SPIRIT OF CLOVES.

R. Bruised cloves one part  
 Alcohol eight parts  
 Macerate for some days, and distil eight parts. *Guibourt.*

## TINCTURE OF CLOVES.

R. Bruised cloves one part  
 Alcohol eight parts

Treat with one-half the alcohol at a time, and filter. *Beral.*

## WINE OF CLOVES.

R. Bruised cloves  
 mace, each, one drachm  
 Red wine one pint  
 Boil, and strain. As a fomentation. *Saunders.*

## OIL OF CLOVES.

R. Bruised cloves at will  
 Water sufficient  
 Macerate for some time, distil, return the product, and redistil several times, separate the oil, which sinks to the bottom. *Van Mons.*

Dose, two to six drops.

## AROMATIC BALSAM OF CLOVES.

R. Oil of cloves  
 nutmeg, each, one scruple  
 Spirit of juniper berries two ounces  
 Mix. As a stimulating friction. *Bories.*

## ODONTALGIC MIXTURE.

R. Oil of cloves twelve drops  
 Tincture of pimpinella  
 Sulphuric ether, each, two drachms  
 Aromatic tincture one drachm  
 Mix. To be applied to the carious tooth. *Augustin.*

## CLOVE PLASTER.

R. Yellow wax twenty-two parts  
 Suet twenty-four parts  
 Melt, and add  
 Powdered olibanum sixteen parts  
 cloves eight parts  
 Oil of nutmegs six parts  
 peppermint one part  
 Mix. *Van Mons.*

As an application to the stomach, or over a pained part.

## CLOVE CATAPLASM.

R. Rye flour six or eight spoonfuls  
 Red wine sufficient  
 to make a cataplasm, add  
 Powdered cloves  
 Rasped nutmeg, each, a teaspoonful  
 Mix well. As an application to the stomach, to check nausea or pain. *Radius.*

## CLOVE BAG.

R. Powdered cloves two drachms  
 marjoram half an ounce  
 rosemary one ounce  
 Mix, and quilt between folds of linen, dip in Cologne water, and apply to the pit of the stomach, to check nausea. *Phæbus.*



## CASCARILLA.

## CASCARILLA.

This is the bark of *Croton eleutheria*, a shrub found in several of the West India islands, and also, perhaps, derived from other species.

*Ser. Syst.* Monœc. monadelph. *Nat. Syst.* Euphorbiaceæ.

Swartz. H. Ind. Oc. 11. 1183. Griffith, Med. Bot. 596.

In rolled fragments, whitish externally, and of a chocolate color within, of an aromatic odor, and a spicy, bitter taste. It is an aromatic tonic, and is useful in debilitated conditions of the stomach and bowels. Dose, in powder, from a scruple to half a drachm.

## COMPOUND CASCARILLA POWDER.

R. Powdered cascarilla ten grains  
opium  
ipecacuanha, each, one grain

Mix. To be taken every five hours in obstinate diarrhœa. *Radiis.*

R. Powdered cascarilla one drachm  
hartshorn  
cuttlefish bone, each, four drachms  
amber two drachms

Mix. One drachm to be taken in a glass of sugar and water, at bedtime, against nocturnal pollutions. *Pierquin.*

## INFUSION OF CASCARILLA.

R. Bruised cascarilla one ounce  
Boiling water one pint  
Macerate for two hours in a covered vessel, and strain. *U. S. Ph.*

Dose, two fl. ounces.

## ALKALINE INFUSION OF CASCARILLA.

R. Bruised cascarilla three ounces  
Carbonate of potassa two drachms  
Boiling water sixteen fl. ounces  
Macerate for two hours, and filter. Dose, a spoonful, two or three times a-day, as an antacid and tonic. *Palat. Ph.*

## EXTRACT OF CASCARILLA.

R. Bruised cascarilla one part  
Alcohol five parts  
Treat the cascarilla several times with the alcohol, distil, and evaporate the residue. Dose, from ten grains to a drachm.

*Guibourt.*

## MIXTURE OF EXTRACT OF CASCARILLA.

R. Extract of cascarilla one drachm  
White sugar six drachms  
Oil of chamomile twenty drops  
Tragacanth half a drachm  
Wine of opium twenty-five drops  
Cinnamon water two fl. ounces  
Peppermint water four fl. ounces

Mix. A spoonful every hour in dysentery. *Augustin.*

## TINCTURE OF CASCARILLA.

R. Powdered cascarilla five ounces  
Proof spirit two pints  
Macerate fourteen days, and filter.

*Lond. Ph.*

## COMPOUND WINE OF CASCARILLA.

R. Powdered cascarilla one ounce and a-half  
Contused orange peel one ounce  
cinnamon two drachms  
White wine twenty-seven fl. ounces  
Digest for twenty-four hours, and filter. As a stomachic and tonic, in doses of two ounces, four times a-day. *Phæbus.*

## CASSIA.

## PURGING CASSIA.

Cassia is the fruit or pod of the *Cassia fistula*, a large tree, a native of Egypt and many parts of Asia, and cultivated in most tropical regions.

*Ser. Syst.* Decand. monog. *Nat. Syst.* Fabaceæ. Linn. Sp. Pl. 540. Griffith, Med. Bot. 253.

The pods are about a foot in length, cylindrical, of a dark brown color, with two longitudinal furrows on one side, and one on the other. They are internally divided into numerous cells, each containing one seed enveloped in a soft pulp. This latter, which is the part used, has a faint nauseous smell, and a sweet, mucilaginous taste. It is laxative, in doses of one or two drachms, and purgative in those of one or two ounces.

## PULP OF PURGING CASSIA.

R. Purging cassia, bruised, sufficient  
Pour boiling water on it to soften the pulp, strain, and evaporate to proper consistence. *U. S. Pharm.*

## CONFECTION OF CASSIA.

R. Cassia pulp half a pound  
Manna two ounces  
Tamarind pulp one ounce  
Syrup of roses eight fl. ounces  
Mix well, and evaporate to proper consistence. *Lond. Ph.*

Slightly laxative.

## MIXTURE OF CASSIA PULP.

R. Pulp of cassia one ounce  
Infusion of rhubarb a drachm and a-half  
Syrup of roses  
Manna, each, one ounce  
Mix well. A mild laxative. *Pierquin.*

## CASSIA MARILANDICA.

## AMERICAN SENNA.

This species is an indigenous perennial plant,



with showy, bright yellow flowers, growing in moist places and on the banks of streams.

*Ser. Syst. Decand. monog. Nat. Syst. Fabaceæ.*

*Linn. Sp. Pl. 541. Griffith, Med. Bot. 260.*

The leaves, which are the officinal portions, are usually exposed for sale in small packages, like the other herbs put up by the Shakers. They have a faint, unpleasant odor, and a nauseous taste like senna, of which they possess all the qualities, but are not quite as active.

#### INFUSION OF AMERICAN SENNA.

R. American senna one ounce and a-half  
Coriander seed, bruised, one drachm  
Boiling water one pint  
Macerate in a covered vessel one hour, and strain. *Martin.*

### CASTOREUM.

#### CASTOR.

Castor is a peculiar substance obtained from membranous follicles in the *Castor fiber*, situated between the organs of generation and the anus. It is in the form of solid, unctuous masses, contained in small, wrinkled, brownish sacs. It has a strong, unpleasant smell, and a bitter, acrid taste. It is stimulant and antispasmodic, and has been much praised in the neuroses. In this country it is not much used. Dose, from ten to twenty grains.

#### COMPOUND POWDER OF CASTOR.

R. Castor twelve grains  
Powdered cascarilla one scruple  
Magnesia a scruple and a-half  
Mix, and divide into three powders. *Saunders.*

R. Castor  
Powdered valerian  
Sugar, each, ten grains  
Mix, and divide into three doses. *Phœbus.*  
As antispasmodics.

#### BOLUS OF CASTOREUM.

R. Assafetida  
Valerian  
Castor  
Amber, each, half an ounce  
Camphor one scruple  
Syrup sufficient  
Mix, and form boluses of ten grains each. Said to be efficacious in hysteria, neuralgia, and vertigo. *Foy.*

#### CASTOREUM PILLS.

R. Assafetida one drachm  
Galbanum  
Myrrh, each, half a drachm  
Castoreum fifteen grains  
Tincture of valerian sufficient  
Beat together, and divide into seventy-two pills. Dose, three to eight, three times a-day, in hysteria. *Phœbus.*

#### PILLS OF CASTOR AND SUCCINIC ACID.

R. Castor one drachm  
Succinic acid half a drachm  
Extract of gentian sufficient  
Beat together, and form twenty-four pills. Three, morning and night, in hysteria. *Ellis.*

#### TINCTURE OF CASTOR.

R. Bruised castor two ounces  
Alcohol two pints  
Digest for seven days, express, and filter. *U. S. Ph.*

Dose from thirty drops to two fl. drachms.

#### ETHEREAL TINCTURE OF CASTOR.

R. Bruised castor two ounces  
saffron one ounce  
Spirit of sulphuric ether twelve ounces  
Digest for twenty four hours, and filter. As an antispasmodic. In dose of thirty drops. *Spielmann.*

#### AMMONIATED TINCTURE OF CASTOR.

R. Bruised castor two ounces and a-half  
Contused assafetida two drachms  
Spirit of ammonia two pints  
Digest for seven days, in a closed vessel, express, strain, and filter. *Ed. Ph.*

A powerful stimulant, and antispasmodic, in cases of spasm of the stomach, hysteria, &c. Dose, thirty drops to two fl. drachms.

#### COMPOUND TINCTURE OF CASTOR.

R. Assafetida two drachms  
Opium half a drachm  
Oil of amber one drachm  
Castor half an ounce  
Alcohol four ounces  
Digest for four days, express, and filter. Dose, thirty to forty drops in wine, in hysteria. *Ferrara. Ph.*

#### COMPOUND PLASTER OF CASTOR.

R. Wax plaster sixteen ounces  
Soap four ounces  
Melt together, and add, on cooling,  
Camphor one ounce  
Castor half an ounce  
Incorporate well. Highly praised in headache, rheumatism, toothache, &c. *Giordano.*

#### COMPOUND SPIRIT OF CASTOR.

R. Castor half an ounce  
Assafetida two drachms  
Oil of amber one drachm  
savine  
rue, each, half a drachm  
Alcohol ten ounces  
Digest, distil, and add  
Empyreumat. carb. of ammonia two ounces



Camphor one drachm  
Distil again. Dose, twenty to forty drops.  
Cottureau.

## CATALPA.

## CATALPA.

The *Catalpa cordifolia* is a beautiful native tree, principally found in the southern and south-western States.

*Sex. Syst.* Diand. monog. *Nat. Syst.* Bignoniaceæ.

*Elliot. Bot. i.* 24. *Lindley, Flor. Med.* 499.

The bark is said to be vermifuge, but the part used in medicine is principally the pods. These have been found efficacious in asthma.

## DECOCTION OF CATALPA PODS.

R. Catalpa pods half an ounce  
Water sufficient

to obtain eight ounces of decoction; add  
Oxymel of squills half an ounce

R. Catalpa pods half an ounce  
Seneka two drachms  
Water sufficient

to obtain eight ounces of decoction; add  
Oxymel of squills one ounce

In spoonful doses, in chronic, nervous asthma.  
*Brera.*

R. Catalpa pods three or four  
Water twelve fl. ounces

Boil down to six ounces, to be given in two doses, morning and night.

*Antomachi.*

## CATARIA.

## CATNEP.

This is the leaves, or whole herb of *Nepeta Cataria*, a perennial herbaceous plant, a native of Europe, and naturalized in the United States.

*Sex. Syst.* Didynam. gymnos. *Nat. Syst.* Labiataceæ.

*Linn Sp. Pl.* 797. *Griffith, Med. Bot.* 512.

It has a peculiar, somewhat unpleasant odor, and a bitterish, aromatic taste. It is stimulant, carminative, &c., and is used in the flatulent colic of infants, and as an emmenagogue, &c.

## INFUSION OF CATNEP.

R. Catnep two drachms  
Boiling water. eight ounces

Infuse in a covered vessel. Dose, a teaspoonful occasionally, whilst hot, for an infant in flatulent colic; to be taken freely, as a diaphoretic, or emmenagogue.

## CATECHU.

## CATECHU.

Catechu is an extract procured from the wood of *Acacia catechu*, and also from other trees.

The *Acacia catechu* is a native of the East Indies, and is now cultivated in some of the West India islands.

*Sex. Syst.* Polygam. monœc. *Nat. Syst.* Fabaceæ.

*Linn. Sp. Pl.* 409. *Griffith, Med. Bot.* 268.

Catechu comes in masses of various forms, of a rusty brown color externally, but paler within. It is inodorous, but has a bitter and astringent taste, followed by a sensation of sweetness. It principally consists of tannin and extractive. It is tonic and astringent, and is used in cases where astringents are required. The dose is from ten grains to half a drachm, repeated as required.

## COMPOUND POWDER OF CATECHU.

R. Powdered catechu fifteen grains  
salts two grains

Mix, and make powder; to be taken after each liquid stool, in diarrhœa from a weakened condition of the bowels.

*A. T. Thomson.*

R. Powdered catechu  
cascarilla  
gum Arabic

Aromatic powder equal parts

Mix. Dose, a scruple every two hours, in chronic diarrhœa.  
*Augustin.*

## BOLUS OF CATECHU.

R. Powdered catechu twenty-four grains  
Extract of opium two grains  
Conserve of roses sufficient

Mix, and form two boluses, one to be taken morning and evening, in chronic diarrhœa.

*Foy.*

## PILLS OF ALUM AND CATECHU.

R. Alum six grains  
Extract of opium  
Catechu each one grain

Mix, and divide into six pills, one to be given every two to four hours. In passive hemorrhage and atonic mucous discharges.

*Ellis.*

## COMPOUND CATECHU BOLUS.

R. Powdered catechu one scruple  
Confection of opium twelve grains  
Aromatic confection sufficient

Make a bolus, to be taken twice a-day, in inordinate flow of the menses.

*Babington.*

R. Powdered catechu twelve parts  
alum six parts  
opium two parts

Syrup of red roses sufficient

Form a mass, and divide into pills of six grains each. One or two a-day, in the decline of gonorrhœa.  
*Foy.*

## PILLS OF CATECHU AND LIQUORICE.

R Catechu half a drachm



Gum Arabic

Extract of liquorice, each,

two drachms

Mastic

one drachm

Syrup of mallows

sufficient

Form mass, and divide into pills of three grains each. Four, three times a-day, in chronic catarrh. *Van Mons.*

## LOZENGES OF CATECHU.

R. Catechu one ounce  
 Sugar half an ounce  
 Cloves one scruple  
 Essence of citron half a scruple  
 Mucilage of tragacanth sufficient  
 Beat into mass, and divide into lozenges of ten grains, each. One, occasionally, to arrest fetor of breath. *Brunswick Ph.*

## ELECTUARY OF CATECHU.

R. Catechu  
 Kino, each, four ounces  
 Cinnamon  
 Nutmeg, each, one ounce  
 Opium diffused in a little sherry one drachm and a-half  
 Syrup of red roses, reduced to the consistence of honey, one pint and a-half  
 Pulverize the solids, mix the opium and syrup, add the powders, and beat into a mass. *Dub. Ph.*

In diarrhœa and chronic dysentery, in dose of half a drachm to a drachm.

R. Catechu  
 Balsam of tolu, each, one drachm  
 Peruvian bark one ounce  
 Syrup of red roses sufficient  
 Beat together. One drachm morning and evening, in leucorrhœa and gonorrhœa. *Brera.*

## INFUSION OF CATECHU.

R. Catechu one drachm  
 Boiling water one pint  
 Digest for six hours, and strain. *Guibourt.*

## COMPOUND INFUSION OF CATECHU.

R. Powdered catechu half an ounce  
 Bruised cinnamon one drachm  
 Boiling water one pint  
 Macerate for an hour in a covered vessel, and strain. *U. S. Ph.*

Dose, from one to three fl. ounces, three or four times a-day.

## TINCTURE OF CATECHU.

R. Catechu three ounces  
 Cinnamon, bruised, two ounces  
 Diluted alcohol two pints  
 Macerate for fourteen days, express, and filter. *U. S. Ph.*

A pleasant astringent. Dose, thirty drops to three fl. drachms.

## CATECHU COLLUTORY.

R. Catechu  
 Myrrh, each, eight parts  
 Balsam of Peru one part  
 Spirit of cochlearia  
 Alcohol, each, twenty-four parts  
 Mix, and digest for four days, and filter.  
 As a mouth-wash, in a spongy condition of the gums. *Saunders.*

## MIXTURE OF CATECHU AND LOGWOOD.

R. Extract of logwood three drachms  
 Tincture of catechu two fl. drachms  
 Water seven fl. ounces  
 Mix. Two spoonfuls, every three or four hours, in chronic stages of diarrhœa and dysentery. *Ellis.*

## ANTI EMETIC MIXTURE.

R. Catechu one drachm  
 Columbo thirty grains  
 Winter's bark twenty grains  
 Boiling water four ounces  
 Mix, and digest for eight hours, strain, and add  
 Syrup of red roses one ounce  
 In spoonful doses. *Pierquin.*

## INJECTION OF CATECHU.

R. Catechu  
 Myrrh, each, one drachm  
 Dissolve in  
 Lime water four ounces  
 and strain. As an injection, in chronic leucorrhœa and gonorrhœa.

## AROMATIC PASTILLES OF CATECHU.

R. Extract of liquorice  
 Water, each, three ounces and a-half  
 Dissolve on a water-bath, and add  
 Powdered catechu four hundred and sixty-two grains  
 gum Arabic two hundred thirty-one grains

Evaporate to the consistence of an extract, and incorporate,

Mastic  
 Cascarilla  
 Charcoal  
 Orris root, each, thirty grains  
 Reduce to proper consistence, remove from fire, and add

Oil of peppermint thirty drops  
 Tincture of ambergris  
 musk, each, ten drops  
 Mix, and form into one-grain pills. Used to aromatize the breath. *Redwood.*



## CEANOTHUS.

## NEW JERSEY TEA.

Several species of *Ceanothus* have medical properties, but the only one in use in this country, is the *C. Americanus*, a suffruticose plant, found in most parts of the United States.

*Sex. Syst.* Pentand. trigyn. *Nat. Syst.* Rhamnaceæ.

Linn. Sp. Pl. 284. Griffith, Med. Bot. 218.

The leaves and root are bitter and astringent, but the root is the most active. They have been used in a variety of cases, to which astringents are applicable.

## DECOCTION OF NEW JERSEY TEA.

R. New Jersey tea root      two drachms  
Water      one pint

Boil for a quarter of an hour, and strain. Said by Ferrein to be efficacious in gonorrhœa, and syphilis; and by Dr. Hubbard, in dysentery, and also as a gargle in aphthous sore mouth, and in ulcerations of the fauces.

## CENTAUREA BENEDICTA.

## BLESSED THISTLE.

An annual plant, native of the south of Europe, and become naturalized in some parts of the United States.

*Sex. Syst.* Syngen. frust. *Nat. Syst.* Asteraceæ.

Linn. Spl. Pl. 1294. Griffith, Med. Bot. (*Cnicus*.) 409.

The whole herb is used; it has a faint nauseous odor, and a very bitter taste. It is used as a tonic, diaphoretic, or emetic. Dose of powder, as tonic, a scruple to a drachm.

## INFUSION OF BLESSED THISTLE.

R. Blessed thistle      three drachms  
Boiling water      one quart  
Infuse for half an hour, and strain. As a diaphoretic. *Fr. Cod.*

R. Blessed thistle      half an ounce  
Cold water      one pint  
Infuse, and strain. As a tonic, in the dose of two fl. ounces. *Wood.*

## WINE OF BLESSED THISTLE.

R. Blessed thistle      one ounce and a-half  
Columbo      three drachms  
Cinnamon      two drachms  
Wine      two pints  
Macerate for twenty-four hours, and strain. Dose, one to four spoonfuls in the morning, as a stomachic. *Wurtemberg Ph.*

## CENTAURIUM.

## CENTAURY.

The herbs or flowering heads of *Erythræa centaurium*, a small annual plant, indigenous to many parts of Europe.

*Sex. Syst.* Pentand. monog. *Nat. Syst.* Gentianaceæ.

Linn. (*Chironia*.) Sp. Pl. 332. Griffith, Med. Bot. 459.

This plant is very bitter, and is analogous in its properties to gentian. In this country, its use has been superseded by the American centaury or sabbatia. Dose of the powder, from thirty grains to a drachm.

## PORTLAND POWDER.

R. Centaury  
Germander  
Gentian  
Round aristolochia  
Teucrium      equal parts  
Powder, and mix. Once highly celebrated in the cure of gout.

## EXTRACT OF CENTAURY.

R. Centaury      one part  
Hot water      eight parts  
Macerate in one-half the water, express, and strain; repeat the process with the remainder of the water, unite the liquids, and evaporate. *Guibourt.*

## PILLS OF EXTRACT OF CENTAURY.

R. Extract of centaury      one drachm  
Myrrh      two drachms  
Balsam of Peru      one scruple  
Mix, and form into pills of three grains. Four, three times a-day, as a tonic and stomachic. *St. Marie.*

## COMPOUND WINE OF CENTAURY.

R. Centaury  
Apocynum  
Madder, each,      one ounce  
White wine  
Water, each,      a pint and a-half  
Boil down to two-thirds, and add to the strained liquid  
Syrup of orange peel      two ounces  
A cupful, morning and evening, in jaundice. *Beral.*

## MIXTURE OF EXTRACT OF CENTAURY.

R. Extract of centaury      one drachm  
Bitter almonds      two drachms  
Chamomile water      two ounces  
Rub well together, and strain. As a febrifuge. *Foy.*

## CETACEUM.

## SPERMACEITI.

A peculiar concrete substance, obtained from the head of *Physeter macrocephalus*, or spermaceti whale. It is white, semi-transparent, friable, soft, somewhat oily to the touch, insipid, but with a faint odor. It has been used as a demulcent, and also enters into the composition of several ointments and cerates.



## COMPOUND POWDER OF SPERMACETI.

R. Spermaceti two drachms  
 Nitrate of potassa one drachm  
 Orris root  
 Sugar, each, one ounce  
 Powder well, and mix. A teaspoonful,  
 three or four times a-day, in catarrh.

*Radiis.*

## SPERMACETI MIXTURE.

R. Spermaceti two drachms  
 Sugar three drachms  
 Paregoric elixir half fl. ounce  
 Water eight fl. ounces  
 Rub spermaceti and sugar together with  
 the yolk of an egg; add the water and  
 paregoric gradually. A tablespoonful,  
 several times a-day. In catarrh. *Ellis.*

R. Spermaceti half an ounce  
 Gum arabic two drachms  
 Syrup half an ounce  
 Water of bitter almonds one drachm  
 Water two ounces and a-half  
 Rub together, and strain. A dessertspoon-  
 ful every two hours. In catarrh. *Phœbus.*

R. Spermaceti two drachms  
 Olive oil one drachm  
 Powdered gum Arabic half an ounce  
 Water four fl. ounces  
 Triturate the spermaceti with the oil, until  
 reduced to a paste, then add the gum, and  
 lastly the water, gradually. *W. Procter.*

R. Spermaceti two drachms  
 Yolk of egg one  
 Powdered gum Arabic two drachms  
 Paregoric elixir six drachms  
 Antimonial wine four drachms  
 White sugar three drachms  
 Water six ounces

Mix. Rub the spermaceti with the yolk  
 of egg, then add the other ingredients se-  
 parately; when well incorporated, introduce  
 the water gradually. A tablespoonful every  
 two or three hours to an adult. Useful in  
 catarrh, especially of measles. *Dewees.*

## SPERMACETI OINTMENT.

R. Spermaceti six drachms  
 White wax two drachms  
 Olive oil three fl. ounces  
 Mix over a gentle fire, and stir till cold.

*Lond. Ph.*

Mild dressing for open surfaces.

OINTMENT OF SPERMACETI AND ROSE  
WATER.

R. Rose water  
 Oil of almonds each two fl. ounces  
 Spermaceti half an ounce  
 White wax one drachm

Melt together, by a water-bath, the oil,  
 spermaceti and wax; add rose water, and  
 stir till cold. *U. S. Ph.*

A pleasant and cooling application to irritable  
 surfaces, well known as *cold cream*.

## COLD CREAM WITHOUT SPERMACETI.

R. White wax one ounce  
 Oil of almonds four fl. ounces  
 Rose water two fl. ounces  
 Borax half a drachm  
 Oil of roses five minims

Dissolve the wax in the oil of almonds, by  
 a gentle heat, also dissolve the borax in the  
 rose water, and add the solution to the  
 heated mixture, stirring till cold, then add  
 the oil of roses. *E. Turnbull.*

## SPERMACETI CERATE.

R. Spermaceti one ounce  
 White wax three ounces  
 Olive oil six fl. ounces  
 Melt wax and spermaceti together, then add  
 the oil previously heated, and stir till cold.  
*U. S. Ph.*

An excellent dressing for blisters, wounds, &c.

## SPERMACETI LIP SALVE.

R. Spermaceti  
 White wax, each, half an ounce  
 Melt, and add  
 Oil of almonds one ounce  
 lavender twenty drops  
 bergamot ten drops  
 cloves two drops  
 Rub well together. *Niemann.*

R. Spermaceti one ounce and a-half  
 White wax nine drachms  
 Oil of almonds twelve ounces  
 Alkanet root two ounces  
 Otto of roses one drachm  
 Digest the first four ingredients in the heat  
 of a water-bath for four hours, strain, and  
 add the otto of roses. *Gray.*

## SPERMACETI LINIMENT.

R. Spermaceti  
 White wax, each, half a drachm  
 Oil of almonds one ounce  
 White lead  
 Litharge, each, twenty grains  
 Gum lac sufficient

Mix. As a cosmetic, but the presence of  
 the lead requires it to be used with caution.  
*Pierquin.*

## SULTANA OINTMENT.

R. Spermaceti eight parts  
 Butter of cacao sixteen parts  
 Oil of almonds thirty-two parts  
 Balsam of Peru one part



Melt together, and add  
Orange-flower water one part  
Stir constantly till cold. *Niemann.*

R. Spermaceti  
White wax, each,  
one drachm and a-half  
Oil of almonds three ounces

Melt together, and add  
Rose water two ounces  
Tincture of benzoin thirty drops  
Mix well. *Taddei.*

## CETRARIA.

## ICELAND MOSS.

This is a lichen found in great abundance in the northern regions of both continents.

*Sex. Syst.* Cryptog. lichen. *Nat. Syst.* Lichenaceæ.

When dried it is inodorous, but has a bitter, mucilaginous taste. It is demulcent, tonic, and nutritious, and is useful in pulmonary and other complaints, in which the local disease is attended with a debility of the digestive organs, or of the general system. Dose of the powder, thirty grains to a drachm.

## DECOCTION OF ICELAND MOSS.

R. Iceland moss half an ounce  
Water one pint and a-half  
Boil down to a pint; express, and strain. *U. S. Ph.*

To be taken during the twenty-four hours.

R. Iceland moss one ounce and a-half  
Water one quart  
Macerate the moss in a weak solution of carbonate of potassa, before subjecting it to decoction.  
Boil, and strain. *Van Mons.*

In this preparation the bitter principle is extracted by the potassa, and the result is a mere mucilaginous demulcent.

## ICELAND MOSS JELLY.

R. Iceland moss two parts  
Water twenty parts  
Boil down to six parts and strain, then add  
Sugar four parts  
Set by to cool. *Augustin.*

## ICELAND MOSS MIXTURE.

R. Iceland moss two ounces  
Hartshorn shavings one ounce  
Water one quart  
Boil down to a pint, and strain, add  
Wine of opium fifteen drops  
In phthisis, to be taken during the day. *Brera.*

R. Iceland moss two drachms  
Powdered salep ten grains

cinnamon eight grains  
Water twelve fl. ounces  
Boil to six ounces, and add  
Syrup sufficient  
Said to be useful in phthisis, leucorrhœa, and hooping-cough, in doses of four ounces, three or four times a-day. *Augustin.*

R. Decoction of Iceland  
moss seven fl. ounces and a-half  
Diluted sulphuric acid one fl. drachm  
Syrup four fl. drachms  
Laudanum fifty drops  
Mix. A wineglassful, three times a-day, in phthisis. *A. T. Thomson.*

## ICELAND MOSS CHOCOLATE.

R. Fine chocolate four pounds  
Sugar two pounds  
Powdered Iceland  
moss, one pound and a-half  
Tragacanth  
Cinnamon, each, four ounces  
Water sufficient  
Rub into a uniform paste. *Cadet.*

## CETRARINE.

R. Iceland moss,  
in coarse powder, one pound  
Alcohol, .883 four pounds  
Boil for half an hour, permit to cool, till no vapors are given off, express, and add to fluid  
Muriatic acid three drachms  
and four times its bulk of distilled water; let rest for a night in a closed matrass; then decant, throw deposit on filter, and press; while still moist, wash with alcohol or ether; then treat with boiling alcohol, filter, and permit cetrarine to precipitate. *Herberger.*

## POWDER OF CETRARINE.

R. Cetrarine  
Gum Arabic, each, two grains  
White sugar half a scruple  
Useful in intermittent fevers, in doses of eight grains every two hours during the apyrexia. *Muller.*

## CHELIDONIUM.

## CELANDINE.

The herb of *Chelidonium majus*, a perennial herbaceous plant, native of Europe, but generally naturalized in this country.

*Sex. Syst.* Polyand. polyg. *Nat. Syst.* Papaveraceæ.

*Linn. Sp. Pl.* 723. *Griffith, Med. Bot.* 130.

Exudes, when broken, an orange-colored, fetid juice. Its taste is extremely bitter and acrid, leaving a burning sensation. The root is



the most powerful. It has been used as a drastic hydragogue, and also in scrofula, &c. The dose of the dried root is half a drachm to a drachm; of the fresh juice, thirty to forty drops.

## EXTRACT OF CELANDINE.

R. Juice of celandine at will  
Boil so as to coagulate; strain, evaporate, and at the close of evaporation add the coagulum. *Van Mons.*

Dose, five to fifteen grains, as a hydragogue purgative.

## PILLS OF CELANDINE.

R. Extract of celandine  
Gum ammoniac, each, one drachm and a-half  
Soap one drachm  
Mix, and form pills of four grains. Two or three a-day, in engorgements of the viscera, especially of the spleen. *Schubarth.*

## EXTRACT OF CELANDINE MIXTURE.

R. Extract of celandine two drachms and a-half  
henbane one scruple  
Sulphate of potassa one ounce  
Tartar emetic one grain  
Elder water six ounces  
Oxymel of squills one ounce  
Mix. A dessertspoonful every two hours, as a hydragogue. *Augustin.*

## CHENOPODIUM.

## WORMSEED.

This is the fruit of *Chenopodium anthelminticum*, a native plant, found in most parts of the country.

*Sex. Syst.* Pentand. digyn. *Nat. Syst.* Chenopodiaceæ.

The fruit is in small grains, of a greenish yellow color, with a bitterish, aromatic, pungent taste, and a nauseous peculiar smell. Besides the seeds, the expressed juice of the whole plant is used, and the essential oil. They are all efficient anthelmintics. The first are given in the form of an electuary; the second in tablespoonful doses.

## DECOCTION OF WORMSEED.

R. Fresh leaves one ounce  
New milk one pint  
Orange peel two drachms  
Boil, and strain. Dose, a wineglassful, twice a-day.

## OIL OF WORMSEED.

R. Wormseed at will  
Water sufficient  
to cover the seed; infuse for some hours, distil, and separate the oil. Dose, from four to eight drops for a child, night and

morning, for three or four days, to be followed by a cathartic. *Coxe.*

## WORMSEED OIL MIXTURE.

R. Oil of wormseed one drachm  
Sugar  
Gum Arabic, each, one drachm and a-half

Mix, and add

Mint water two and a-half fl. ounces  
A teaspoonful, four times a-day for two days, to a child, to be followed by a purge. *Ellis.*

## CHIMAPHILA.

## PIPSISSEWA.

The leaves of *Chimaphila umbellata*, an evergreen plant, found in the northern portions of both continents.

*Sex. Syst.* Decand. monog. *Nat. Syst.* Pyrolaceæ.  
Linn. (*Pyrola*) Sp. Pl. 568. Griffith, Med. Bot. 421.

Pipsissewa has a peculiar and rather pleasant odor, especially when fresh, and an astringent, bitter, and somewhat sweetish taste. It is tonic, diuretic, and astringent, and has been much used in nephritic complaints, dropsy, &c.

## DECOCTION OF PIPSISSEWA.

R. Pipsissewa one ounce  
Water one pint and a-half  
Boil down to a pint, and strain. *U. S. Ph.*

To be taken during the day, in dropsy and chronic affections of urinary organs.

## COMPOUND DECOCTION OF PIPSISSEWA.

R. Decoction of pipsissewa one pint  
Solution of carbonate of potassa two drachms  
Mix. Four tablespoonfuls, three times a-day, in same cases as above. *Foy.*

R. Pipsissewa half an ounce to one ounce  
Water twelve fl. ounces  
Boil to six ounces, and add  
Gin two ounces  
Let stand for two hours, and strain. Two spoonfuls, four times a-day, in dropsy. *Radius.*

R. Pipsissewa six drachms  
Water twelve ounces  
Boil to six ounces, and add at close  
Senna two drachms  
Strain. A tablespoonful every two hours as a diuretic. *Radius.*

## EXTRACT OF PIPSISSEWA.

R. Pipsissewa one pound  
Water one gallon  
Macerate for twenty-four hours, boil down to two pints, strain, and evaporate to proper consistence. Dose, ten to thirty grains.



## PIPSISSEWA PILLS.

R. Extract of pipsissewa  
 Resin of guaiacum,  
     each,           one drachm and a-half  
 Precipitated sulph. of  
     antimony                   twelve grains  
 Beat together, and make pills of two grains.  
 Dose, five, twice a-day, in arthritic affec-  
 tions.

## PIPSISSEWA SYRUP.

R. Powdered pipsissewa      four ounces  
 Water                   eight fl. ounces  
 Macerate for thirty-six hours, percolate so  
 as to obtain a pint of fluid, evaporate one-  
 half, and add  
 Sugar                   sixteen ounces  
 One to two tablespoonfuls at a time.  
                                   *W. Procter.*

## PIPSISSEWA BEER.

R. Pipsissewa                   half a pound  
 Water                   one gallon  
 Boil, strain, and add  
 Brown sugar                   one pound  
 Powdered ginger           half an ounce  
 Yeast                   sufficient  
 To produce fermentation. When this has  
 taken place, keep in well-stopped bottles.  
 Dose, a half-tumblerfull, three or four  
 times a-day. In strumous affections, espe-  
 cially of the joints.           *Dr. I. Parrish.*

## CHIRETTA.

## CHIRETTA.

This is the herb and root of *Agathotes chirayta*,  
 a native of the northern mountainous parts of  
 India.

*Ser. Syst.* Pentand. digyn. *Nat. Syst.* Gentia-  
 naceæ.

Don. Phil. Mag., 1836. Griffith, Med. Bot.  
 462.

The whole plant is bitter, but the root most so;  
 it is inodorous. It appears to have the same pro-  
 perties as gentian, and is applicable to the same  
 cases. Dose, of powder, is twenty grains.

## INFUSION OF CHIRETTA.

R. Chiretta                   four drachms  
 Boiling water                   one pint  
 Infuse for two hours, and strain. *Ed. Ph.*  
 Dose, one to two fl. ounces as a simple tonic  
 bitter.

## TINCTURE OF CHIRETTA.

R. Chiretta                   five ounces  
 Diluted alcohol                   two pints  
 Macerate for fourteen days, and filter.  
 Dose, a teaspoonful.           *Dunglison.*

## CHLORINUM.

## CHLORINE.

This is an elementary, gaseous fluid, of a  
 greenish yellow color, and a peculiar, penetrat-  
 ing smell. It has been used as an inhalation in  
 a diluted state, in diseases of the lungs, and also  
 as a fumigation in diseases of the liver, &c., and  
 also as a disinfectant.

## CHLORINE.

R. Peroxide of manganese      one part  
 Common salt                   three parts  
 Sulphuric acid  
 Water, each,                   two parts  
 Mix the acid and water, and pour on the  
 other substances in powder.   *Paris Cod.*

## CHLORINE WATER.

R. Dried muriate of  
     soda                   one hundred parts  
 Oxide of manganese           thirty parts  
 Sulphuric acid               eighty-seven parts  
 Water   one hundred and twenty parts  
 Add the acid gradually to the water; when  
 cold, pour on the other substances in pow-  
 der, and put into a retort; then with a  
 gradually increased heat, let the gas that  
 escapes be transmitted through two hun-  
 dred parts of distilled water; the process  
 is concluded, when effervescence in the  
 retort has subsided.           *Dub. Ph.*

To be kept in well-stopped bottles, and in  
 the dark. Chlorine water is stimulant, and an-  
 tiseptic, and has been used in a variety of dis-  
 eases. (See *Dunglison, New Rem.* 170 *et seq.*)  
 The dose is from one to four fl. drachms, pro-  
 perly diluted.

## CHLORINE COLLUTORY.

R. Chlorine water                   half an ounce  
 Syrup of mallows                   one ounce  
 Mix. In aphthous sore mouth. *Radius.*  
 R. Chlorine water                   half an ounce  
 Infusion of sage                   six ounces  
 Honey of roses                   one ounce  
 Mix. In same cases.           *Radius.*

## CHLORINE GARGLE.

R. Tragacanth                   twelve grains  
 Distilled water                   four fl. ounces  
 Chlorine water  
 Syrup, each,                   half an ounce  
 Mix. As a gargle in ulceration, and  
 chronic inflammation of the mouth, and  
 fauces.                   *Foy.*

## CHLORINE INJECTION.

R. Chlorine water                   half a drachm  
 Distilled water   one ounce and a-half  
 Extract of opium                   fifteen grains  
 Mix. As an injection in ulcers of the  
 uterus.                   *Phæbus.*



## CHLORINE CLYSTER.

- R. Chlorine water two drachms  
 Starch four drachms  
 Extract of opium half a grain  
 Water six ounces  
 Mix. As an enema in the diarrhoea of consumptive patients. *Cadet.*

## CHLORINE MIXTURE.

- R. Chlorine water two to four drachms  
 Decoction of mal-  
 lows two ounces and a-half  
 Mix. A teaspoonful every hour, in softening the stomach in children. *Radius.*  
 R. Chlorine water two ounces  
 Syrup six ounces  
 Mix. A spoonful every two hours, in typhoid fevers. *Swedjaur.*  
 R. Chlorine water two drachms  
 Strawberry water one ounce and a-half  
 Quince mucilage  
 Syrup of mallows, each, six drachms  
 Mix. A teaspoonful every two hours, to children in scarlatina. *Radius.*

## CHLORINE OIL.

- R. Olive oil at will  
 Pass a current of chlorine through it, at the end of two or three days, wash it with cold water. As an application to tinea, and lepra. *Deimann.*

## CHLORINE OINTMENT.

- R. Chlorine water one part  
 Lard eight parts  
 Triturate well together. Employed in itch. *Augustin.*

## CHLORINE LINIMENT.

- R. Chlorine water one fl. drachm  
 Olive oil one fl. ounce  
 Mix well. In itch, tinea, and herpes. *Deimann.*  
 R. White wax two drachms  
 Melt by a gentle heat, and add  
 Almond oil sufficient  
 to make a liniment; on cooling, add  
 Chlorine water one drachm and a-half  
 As an application to foul ulcers. *Ludwig.*

## CHLOROFORMUM.

## CHLOROFORM.

This has received various names, as *Chloride of Carbon*, *Chloride of Formyl*, *Chloric Ether*, *Trichloride of Formyl*, &c. It is a transparent, heavy fluid, of a peculiar, fragrant, ethereal, apple-like odor, and a slightly acid, intensely sweet taste. In large doses it is a narcotic poison; in medicine, a stimulant, sedative, antispasmodic anodyne, and anæsthetic; for which latter purpose it has

attained much celebrity. The usual dose, when inhaled, is a fl. drachm, to be increased in a few minutes, if no effect is produced. The best plan of inhaling is to twist a handkerchief into the form of a bird's nest, wet this with the chloroform, and apply it to the mouth and nose. The dose, internally, is from five to twenty minims.

- R. Powdered chloride  
 of lime four pounds  
 Water twelve pounds  
 Alcohol twelve fl. ounces  
 Mix, in a capacious still or retort, and distil as long as a dense liquid, which sinks in the water that comes over with it, is produced. Separate this, agitate it with sulphuric acid, and distil it from carbonate of barytæ. *Dumas.*

## EMULSION OF CHLOROFORM.

- R. Chloroform six drachms  
 Syrup half an ounce  
 Yolk of egg one  
 Distilled water four fl. ounces and a-half  
 Mix, by rubbing the yolk of the egg with the syrup, then adding the chloric ether and water so as to form an emulsion. *Moxon.*

## LINIMENT OF CHLOROFORM.

- R. Soap liniment two fl. ounces  
 Chloroform one fl. drachm  
 Mix. As an application in neuralgic pains, over the affected part. *Tuson.*

## CHONDRUS.

## IRISH MOSS. CARRAGEEN.

The fronds of *Chondrus crispus*, a sea-weed growing on rocks and stones in the seas of Northern Europe, especially on the southern and western coasts of Ireland.

*Sex. Syst. Cryptog. fuci. Nat. Syst. Ceramiceæ.*

Greville, Alg. Brit. 129. Dunglison, New Rem. 175.

It is of a yellowish, or purplish color, translucent, nearly tasteless, and scentless. It is nutritive and demulcent, and affords a useful article of diet to convalescents. It has also been recommended in scrofula, and bowel and pectoral affections. Before using it should be soaked in cold water to remove any salt or other impurities.

## DECOCTION OF CARRAGEEN.

- R. Carrageen, sliced, half an ounce  
 Water three pints  
 Boil for a quarter of an hour, express, and strain. *Beral.*  
 R. Carrageen, sliced, half a drachm  
 New milk, nine fl. ounces



Boil to five fl. ounces, strain, and add  
 Sugar half a drachm to a drachm  
 Bitter almond water one scruple  
 To be taken during the day. *Gräfe.*

R. Carrageen four scruples  
 New milk twenty-four fl. ounces  
 Boil for ten minutes, express, and add  
 Orange-flower water one ounce and a-half

## CARRAGEEN JELLY.

R. Carrageen two drachms  
 New milk one pint  
 Boil to consistence of jelly, strain, and add  
 Sugar one ounce  
 Bitter almonds two  
*Radius.*

R. Mucilage of carrageen five ounces  
 Sugar four ounces  
 Evaporate to eight ounces, by a gentle heat; strain, and add  
 Essence of citron fifty drops  
*Beral.*

## CARRAGEEN MIXTURE.

R. Carrageen half a drachm  
 Water sufficient  
 to obtain six ounces of mucilage; strain, and add  
 Phosphate of soda one drachm and a-half  
 Syrup of opium three drachms  
 A spoonful, every two hours, in hemoptysis.  
*Clarus.*

## CICHORIUM.

## SUCCORY.

The *Cichorium intybus* is a perennial herbaceous plant, indigenous to Europe, but naturalized in many places in this country.

*Sex. Syst.* Syngen. æqual. *Nat. Syst.* Asteraceæ.

The whole plant is bitter, the root the most so. It is tonic, aperient, and deobstruent. The dried and torrefied root is used in Europe to mix with, or as a substitute for coffee.

## EXTRACT OF SUCCORY.

R. Juice of succory at will  
 Evaporate to proper consistence. *Guibourt.*  
 Dose, from ten grains to half a drachm.

## INFUSION OF SUCCORY.

R. Dried succory half an ounce  
 Boiling water two pints  
 Infuse and strain, and add  
 Syrup of maidenhair one drachm  
*Cottureau.*

## COMPOUND SYRUP OF SUCCORY.

R. Juice of succory  
 dandelion

fumitory, each, six ounces  
 Sugar one pound  
 Boil to proper consistence, and strain.  
 Dose, one to two ounces. As a diuretic.  
*St. Marie.*

## CIMICIFUGA.

## BLACK SNAKEROOT.

This is the root of *Cimicifuga racemosa*, a native, found in many parts of the United States, in shady places.

*Sex. Syst.* Polyand. di pent. *Nat. Syst.* Ranunculaceæ.

Torrey and Gray, Fl. 1. 36. Griffith, Med. Bot. 92.

The root is thick, contorted, dark brown, with numerous radicles. The odor is peculiar and unpleasant, and the taste bitter and somewhat astringent. It is acro-narcotic, and is used in rheumatism, dropsy, pectoral affections, chorea, &c.. The dose of the powder, a drachm, several times a-day.

## DECOCTION OF BLACK SNAKEROOT.

R. Black snakeroot (contused) one ounce  
 Water one pint  
 Boil for a short time, and strain. Dose, one to two fl. ounces. *Wood.*

## TINCTURE OF BLACK SNAKEROOT.

R. Bruised black snakeroot four ounces  
 Alcohol one pint  
 Digest for fourteen days, and filter. Dose, twenty drops, three or four times a-day.  
*Carson.*

Used in rheumatic cases.

## CINCHONA.

## PERUVIAN BARK.

Cinchona is the bark of several species of *Cinchona*, all natives of South America. There are numerous varieties of this article used in medicine, but the state of our knowledge on the subject does not permit a reference of them to a particular species of the tree, except in a few cases. (See *U. S. Dispens.* 212, and Griffith, *Med. Bot.* 368.) Peruvian bark owes its properties to the presence of one or more alkaloids, called quinia, cinchonina, &c.

It is a bitter tonic and antiperiodic, and is used in a variety of diseases, and is given in a multitude of forms. The dose in powder as a febrifuge is a drachm, frequently repeated as a tonic from ten to thirty grains. Its use is now much superseded by that of quinia.

## COMPOUND POWDER OF PERUVIAN BARK.

R. Powdered Peruvian bark  
 Cremor tartar, each, one ounce  
 Powdered cloves one drachm  
 Mix. A drachm and a-half every two hours, in the apyrexia of fevers. *Ellis.*



R. Powdered Peruvian bark half an ounce  
 Snakeroot one drachm  
 Bicarbonate of soda two scruples  
 Mix, and divide into four powders, one to be given every two hours, in apyrexia of obstinate intermittents. *Ellis.*

R. Powdered Peruvian bark  
 uva ursi, each, one or two drachms  
 opium three grains  
 Mix, and divide into six powders, one to be taken three or four times a-day, followed by a draught of two ounces of lime water. Antilithic. *Ferriar.*

R. Powdered Peruvian bark one ounce  
 Tartar emetic two grains  
 Opium one grain  
 Mix, and divide into four powders; one to be taken every two hours. *Brera.*

R. Powdered Peruvian bark  
 Sulphate of magnesia, each, six drachms  
 Mix, and divide into four powders. One every two hours, where purging is required, and at same time a tonic impression to be made. *Ellis.*

R. Powdered Peruvian bark thirty grains  
 Aromatic powder ten grains  
 Make a powder, to be taken in a cup of milk every three hours. In convalescence from fevers.

*A. T. Thomson.*

#### POWDER OF PERUVIAN BARK AND ARNICA.

R. Powdered Peruvian bark  
 arnica  
 camphor equal parts  
 Mix. As an application to gangrenous ulcers. *Cadet.*

#### POWDER OF PERUVIAN BARK AND CASCARILLA.

R. Powdered Peruvian bark two drachms  
 cascarrilla twelve grains  
 Mix. In apyrexia of intermittents *Foy.*

#### POWDER OF PERUVIAN BARK AND CAMPHOR.

R. Powdered Peruvian bark half an ounce  
 camphor one scruple  
 gum Arabic two drachms  
 Mix. As an application to atonic ulcers. *Gen. Ph.*

#### PERUVIAN BARK DENTIFRICE POWDERS.

R. Powdered Peruvian bark one ounce  
 and a-half  
 red saunders half an ounce  
 Oil of bergamot

Oil of cloves, each, twelve drops  
 Mix. *Hufeland.*

R. Powdered Peruvian bark three ounces  
 cremor tartar  
 sage leaves  
 myrrh, each, half an ounce  
 catechu six drachms  
 Oil of cloves sixteen drops  
 Mix. *Hesse Ph.*

R. Powdered Peruvian bark  
 charcoal, each, one drachm  
 and a-half  
 cinnamon half a drachm  
 Mix. *Soubeyran.*

R. Powdered Peruvian bark one ounce  
 orris root  
 sage leaves  
 myrrh, each, half an ounce  
 Mix. *Fulda. Disp.*

R. Powdered Peruvian bark two ounces  
 muriate of ammonia half an ounce  
 orris root one ounce  
 catechu  
 myrrh, each, six drachms  
 Oil of cloves twelve drops  
 Mix. *Pideret.*

#### POWDER OF PERUVIAN BARK AND VALERIAN.

R. Powdered Peruvian bark two drachms  
 valerian one drachm  
 Mix. Divide into four powders. In nervous disorders. *Ratier.*

#### POWDER OF PERUVIAN BARK AND ISINGLASS.

R. Powdered Peruvian bark one ounce  
 isinglass one drachm  
 and a-half  
 Mix, and divide into sixteen powders. Dose, at first, from three to six a-day, in passive menorrhagia, leucorrhœa, chronic diarrhœa, &c. *Brera.*

#### PERUVIAN BARK MIXTURE.

R. Powdered Peruvian bark half an ounce  
 Confection of opium one drachm  
 Lemon juice two drachms  
 Port wine four fl. ounces  
 Mix. A wineglassful every two hours, in the apyrexia of intermittent fever. *Ellis.*

R. Powdered Peruvian bark two drachms  
 Wine of opium sixty drops  
 Nitrate of potassa five grains  
 Infusion of gentian six ounces  
 Mix. An ounce every two hours. *Pierquin.*



R. Powdered Peruvian bark two drachms  
Compound tincture of bark one fl. ounce  
Decoction of bark three fl. ounces  
Syrup half fl. ounce  
Mix. A tablespoonful every one or two hours in the apyrexia. *Ellis.*

R. Powdered Peruvian bark one ounce and a-half  
Magnesia six drachms  
Saffron one scruple  
Conserve of opium two drachms  
White wine  
Water, each, twelve ounces  
Mix. To be taken in divided doses, during the day, in apyrexia. *Giordano.*

R. Powdered Peruvian bark one ounce  
Alcohol eight ounces  
Sulphuric ether two drachms  
Mix. A spoonful every hour, or two hours. *Augustin.*

R. Powdered Peruvian bark one ounce  
ginger twenty grains  
Red wine eight fl. ounces  
Sugar one ounce  
Mix. To be taken during the apyrexia. *Radius.*

#### EXTRACT OF PERUVIAN BARK.

R. Peruvian bark, in coarse powder, one pound  
Alcohol four pints  
Water sufficient

Macerate the bark in the alcohol for four days, and obtain tincture by displacement. When the liquid measures four pints, set aside, and obtain six pints of infusion. Distil off alcohol from tincture, and evaporate infusion to the consistence of thin honey; mix, and evaporate to proper consistence. *U. S. Ph.*

Dose, ten to thirty grains.

#### VINOUS EXTRACT OF PERUVIAN BARK.

R. Powdered Peruvian bark one part  
White wine eight parts  
Digest for three days, express, strain, and evaporate to proper consistence. *Hesse Ph.*

#### COMPOUND BOLUS OF EXTRACT OF PERUVIAN BARK.

R. Extract of Peruvian bark twenty grains  
Nitrate of potassa thirty grains  
Conserve of orange peel sufficient  
Mix, and form five boluses. *Pierquin.*

#### COMPOUND PILLS OF EXTRACT OF PERUVIAN BARK.

R. Extract of Peruvian bark

Extract of gentian, each, one drachm  
Sulphate of iron half a drachm  
Powdered myrrh one drachm  
Oil of caraway ten drops  
Syrup of ginger sufficient  
Beat together into mass, and form forty pills. Three to be taken, three times a-day. *Ellis.*

R. Extract of Peruvian bark one drachm  
opium one grain  
Camphor twelve grains  
Powdered Peruvian bark sufficient  
Beat into a mass, and form twelve pills. One to be taken, three or four times a-day. *Ellis.*

R. Extract of Peruvian bark  
rhubarb  
gentian  
blessed thistle equal parts  
Beat into mass, and form pills of four grains. Dose, one or two, three times a-day. *St. Marie.*

R. Extract of Peruvian bark  
gentian, each, one drachm  
Sulphate of iron half a drachm  
Powdered myrrh one drachm  
Oil of caraway ten drops  
Syrup of ginger sufficient  
Beat well together, and divide into forty pills. Three to be taken three times a-day. In the apyrexia of intermittents. *Thomson.*

R. Extract of Peruvian bark two drachms  
Powdered alum one drachm  
Syrup sufficient  
Mix, and divide into thirty-six pills. Four to be taken every four or six hours. In passive hemorrhages. *A. T. Thomson.*

#### DECOCTION OF PERUVIAN BARK.

R. Peruvian bark, bruised, one ounce  
Water one pint  
Boil for ten minutes, and strain while hot. Dose, two fl. ounces. *U. S. Ph.*

R. Peruvian bark, bruised, one ounce  
Water one pint  
Boil for ten minutes, and while hot, add  
Snakeroot half an ounce  
Orange peel two drachms  
Infuse for half an hour, near the fire, in a covered vessel. Dose, a wineglassful every hour. *Ellis.*

R. Decoction of Peruvian bark six fl. ounces  
Compound tincture of bark one fl. ounce  
Aromatic sulphuric acid ten drops  
Mix. A tablespoonful every hour or two. *Ellis.*



## DECOCTION OF PERUVIAN BARK MIXTURE.

R. Decoction of Peruvian bark six ounces  
 Tincture of Peruvian bark one ounce  
 Sulphuric acid one drachm  
 Syrup of orange peel half an ounce  
 Mix. A tablespoonful every hour or two.  
*Foy.*

R. Decoction of Peruvian bark eight ounces  
 Infusion of arnica flowers six ounces  
 Camphor two scruples  
 Syrup of balsam of tolu one ounce  
 Emulsion of gum four ounces  
 Mix. To be taken in divided doses, during the day.  
*Brugnatelli.*

## DECOCTION OF PERUVIAN BARK AND CASCARILLA.

R. Peruvian bark one ounce  
 Cascarilla two drachms  
 Water twelve ounces  
 Boil to eight ounces, express, and strain, and add, when cold,  
 Sulphuric ether two drachms  
 Mix. *Copenhagen Ph.*

## DECOCTION OF PERUVIAN BARK AND RHATANY.

R. Peruvian bark  
 Rhatany, each, half an ounce  
 Water sufficient  
 to obtain nine ounces of decoction; strain, and add

Vinous extract of Peruvian bark four scruples  
 Compound tincture of Peruvian bark one fl. drachm  
 Citron water one ounce

Mix. In chronic diarrhoea. *Brera.*

## COMPOUND DECOCTION OF PERUVIAN BARK.

R. Peruvian bark one ounce  
 Virginia snakeroot half an ounce  
 Orange peel two drachms  
 Powdered cloves one drachm  
 Carbonate of potassa one drachm

Mix. Put the mixture into a proper vessel, with three half pints of water, boil down gently, to one pint; let settle, decant. Dose, a wineglassful every hour, for eight hours, in the apyrexia. *Dewees.*

## INFUSION OF PERUVIAN BARK.

R. Peruvian bark, bruised, one ounce  
 Boiling water one pint  
 Infuse for two hours, in a covered vessel, and strain.  
*U. S. Ph.*

Dose, two fl. ounces, as occasion may require.

## INFUSION OF PERUVIAN BARK WITH LIME WATER.

R. Bruised Peruvian bark two ounces  
 Lime water two pints  
 Infuse, and strain. A wineglassful to be given three or four times a-day. *Ellis.*

## INFUSION OF PERUVIAN BARK AND VALERIAN.

R. Bruised Peruvian bark one ounce and a-half  
 valerian half an ounce  
 Boiling water two pints  
 Infuse for twelve hours, and strain.  
*Fulda Disp.*

## INFUSION OF PERUVIAN BARK AND SERPENTARIA.

R. Bruised Peruvian bark ten drachms  
 Boiling water one pint  
 Infuse for four hours, then boil for half an hour, then add  
 Snakeroot two drachms  
 Infuse for two days, and strain.  
*Saunders.*

## INFUSION OF PERUVIAN BARK AND MAGNESIA.

R. Powdered Peruvian bark one ounce  
 Magnesia one drachm  
 Boiling water twelve ounces  
 Infuse for twelve hours, express, and strain. Said to be more astringent than other infusions.  
*Van Mons.*

## COMPOUND INFUSION OF PERUVIAN BARK.

R. Peruvian bark  
 Juniper berries, each, one drachm  
 Orange peel  
 Cinnamon, each, two drachms  
 Winter's bark half a drachm  
 Ipecacuanha fifteen grains  
 Boiling water seven ounces

Infuse, and strain, then add  
 Extract of juniper berries one drachm and a-half  
 In spoonful doses, in atonic dropsies.  
*Bories.*

R. Root of scorzonera one ounce  
 Contused lemon seeds two drachms  
 Water six pints

Boil to four pints; pour boiling liquid on  
 Bruised Peruvian bark two ounces  
 Contrayerva one ounce  
 Infuse for a quarter of an hour, strain, and add

Honey of elder three ounces  
 As a stimulating diuretic. *Span. Ph.*



## COMPOUND INFUSION OF PERUVIAN BARK.

R. Powdered Peruvian bark one ounce  
Aromatic sulphuric acid one fl. drachm  
Water one pint  
Macerate for twelve hours, and strain.  
*U. S. Ph.*

Dose, two fl. ounces.

## COMPOUND INFUSION OF PERUVIAN BARK AND IRON.

R. Peruvian bark, in coarse powder, one ounce  
Sliced Colombo root three drachms  
Bruised cloves two drachms  
Iron filings half an ounce  
Digest for three days in a close vessel, add  
Peppermint water sufficient  
to yield a mixture of twelve ounces, on filtering, then add

Compound tincture of cardamom three ounces

Tincture of orange peel three drachms

Dose, one to two fl. ounces. *Dub. Ph.*

## ELECTUARY OF PERUVIAN BARK.

R. Powdered Peruvian bark one ounce  
Syrup of orange peel sufficient  
Mix. A teaspoonful every hour. *Radius.*

## ASTRINGENT ELECTUARY OF PERUVIAN BARK.

R. Powdered Peruvian bark  
orange peel, each, six drachms  
crabs' eyes two drachms  
Conserve of red roses  
rose hips, each, six drachms  
Syrup of catechu sufficient  
Mix. Dose, one drachm, two or three times a-day, in chronic diarrhoea.  
*Saunders.*

## ELECTUARY OF PERUVIAN BARK AND CATECHU.

R. Powdered Peruvian bark one ounce  
catechu  
Balsam of tolu, each, one drachm  
Syrup of comfrey sufficient  
Divide into ten doses, one to be taken twice or three times a-day, in some appropriate vehicle. Much praised in hemoptysis.  
*Pierquin.*

## ELECTUARY OF PERUVIAN BARK AND TIN.

R. Powdered Peruvian bark one ounce  
tin  
valerian, each, half an ounce  
Syrup sufficient  
Mix. Dose, a drachm, morning and evening, in epilepsy. *Cadet.*

## ELECTUARY OF PERUVIAN BARK AND CLOVES.

R. Peruvian bark two ounces  
Cloves one drachm  
Simple syrup sufficient  
to form electuary. A piece, the size of a nutmeg, every two hours in the apyrexia of intermittent fever. *Devees.*

## ELECTUARY OF PERUVIAN BARK AND IRON.

R. Powdered Peruvian bark six drachms  
Pure oxide of iron  
Confection of opium, each, two drachms  
Syrup of cinnamon sufficient  
Mix. Dose, a drachm, morning and evening, in ascites, after the evacuation of the fluid. *Cadet.*

## ELECTUARY OF PERUVIAN BARK AND SULPHUR.

R. Powdered Peruvian bark one drachm  
sulphur  
crabs' eyes  
spermaceti, each, two drachms  
Extract of opium four grains  
Powdered mushrooms one drachm and a-half  
Conserve of roses four drachms  
Syrup of milfoil sufficient  
Mix. Dose, two scruples, three times a-day. Highly praised in phthisis. *Cadet.*

## TINCTURE OF PERUVIAN BARK.

R. Powdered Peruvian bark six ounces  
Diluted alcohol two pints  
Macerate for fourteen days, express, and filter. *U. S. Ph.*

Generally employed as an adjuvant to other preparations of bark, in the dose of one to four fluid drachms.

## COMPOUND TINCTURE OF PERUVIAN BARK.

R. Powdered Peruvian bark two ounces  
Bruised orange peel one ounce and a-half  
snakeroot three drachms  
Saffron, cut,  
Red saunders, rasped, each, one drachm  
Diluted alcohol twenty fl. ounces  
Macerate for fourteen days, express, and filter. *U. S. Ph.*

An excellent stomachic and useful adjunct to other preparations of bark.

## TINCTURE OF PERUVIAN BARK AND VALERIAN.

R. Tincture of Peruvian bark  
valerian, each, one fl. ounce  
cardamom two fl. drachms  
Mint water four fl. ounces



Mix. A tablespoonful every three hours, as a tonic in nervous temperaments. *Ellis*.

#### TINCTURE OF PERUVIAN BARK AND SNAKEROOT.

R. Powdered bark three ounces  
snakeroot two ounces  
contrayerva one ounce  
Alcohol sixteen fl. ounces  
Macerate for twenty-four hours, express, and filter. Dose, fifty to sixty drops.

*Van Mons.*

#### TINCTURE OF PERUVIAN BARK AND AMMONIA.

R. Powdered Peruvian bark one ounce  
Ammoniated alcohol eight fl. ounces  
Macerate for eight days in a covered vessel, express, and filter. *Swediaur.*

#### TINCTURE OF PERUVIAN BARK AND GENTIAN.

R. Bruised Peruvian bark eight parts  
gentian  
orange peel, each, three parts  
Diluted alcohol ninety-six parts  
Macerate for three days, express, and filter. *Foy.*

#### TINCTURE OF BARK AND CANTHARIDES.

R. Tincture of bark three ounces  
Paregoric elixir half an ounce  
Tincture of Spanish flies one drachm  
Mix. Used, it is stated, with great success, in whooping-cough. To be given in small doses, until a slight strangury is caused. *Sutcliffe.*

#### CATAPLASM OF PERUVIAN BARK.

R. Cataplasm of linseed meal one pound  
Powdered Peruvian bark four ounces  
Mix. *Foy.*

R. Powdered Peruvian bark  
charcoal, each, one ounce  
camphor one drachm and a-half  
Spirits of turpentine sufficient  
Mix. *Phæbus.*

These cataplasms are employed as applications to gangrenous ulcers, &c.

#### CERATE OF PERUVIAN BARK.

R. Simple cerate eight ounces  
Camphor one drachm and a-half  
Rub together, and gradually incorporate  
Decoction of Peruvian bark one ounce  
An application to ill-conditioned ulcers.

*Van Mons.*

#### PERUVIAN BARK POMATUM.

R. Extract of Peruvian bark half a drachm  
Oil of roses two drops  
bergamot eight drops

Beef marrow half an ounce  
Incorporate well. As an application to promote the growth of the hair. *Phæbus.*

#### PERUVIAN BARK GELATINE.

R. Bruised Peruvian bark  
Water, each, four ounces  
Boil, express, strain, and add  
Gelatine three drachms  
White sugar one ounce and a-half  
To be taken in spoonfuls, in the apyrexia of intermittents. *Pierquin.*

#### CLYSTER OF PERUVIAN BARK.

R. Decoction of Peruvian bark four ounces  
Yolk of egg one  
Powdered Peruvian bark three drachms  
Tincture of opium twelve drops  
Mix. *Radius.*

R. Extract of Peruvian bark half a drachm  
Warm water four fl. ounces  
Dissolve, and add  
Olive oil half a fl. ounce  
Tincture of opium ten drops  
Mix. To be administered every four hours, in the apyrexia of intermittents. *Ellis.*

#### GARGLE OF PERUVIAN BARK.

R. Tincture of Peruvian bark  
Syrup of mulberries, each, four fl. drachms  
Spirit of horseradish one drachm  
Infusion of sage six fl. ounces  
Mix. In obstinate sore throat, when active inflammation has subsided. *Cadet.*

#### LOTION OF PERUVIAN BARK.

R. Extract of Peruvian bark  
Sulphuric ether, each, one drachm  
Alcohol seven drachms  
Mix. As a friction, three times a-day, in weakness of the back. *Augustin.*

#### PERUVIAN BARK BEER.

R. Bruised Peruvian bark one and a-half ounces  
Cinnamon two drachms  
Rasped nutmeg seven drachms  
Yeast two ounces  
Sugar twenty-five ounces  
Water one hundred fl. ounces  
Mix, and expose to a warm temperature after fermentation, decant, and strain. Three or four glasses to be taken during the apyrexia of intermittents. *Ferrara Ph.*

COMPOUND SYRUP OF PERUVIAN BARK.  
R. Powdered Peruvian bark two drachms



Powdered seneka half a drachm  
 liquorice one drachm  
 Contused poppy heads three  
 ipecacuanha one scruple  
 Ground ivy  
 Hyssop  
 Pennyroyal, each, a handful  
 Sugar one pound  
 Water sufficient  
 Boil, and make a syrup. Three spoonfuls  
 a day, in hooping-cough. *Bories.*

## WINE OF PERUVIAN BARK.

R. Powdered Peruvian bark one part  
 White wine twelve parts  
 Digest for twenty-four hours, often stirring,  
 and filter. *Cottreau.*  
 R. Red wine twenty parts  
 Tincture of Peruvian bark one part  
 Mix. *Niemann.*

## WINE OF PERUVIAN BARK AND IRON.

R. Bruised Peruvian bark, one ounce and a-half  
 Oxide of iron half an ounce  
 Powdered cinnamon two drachms  
 Sugar one ounce  
 White wine two pints  
 Mix, and digest for two days, and filter.  
 Dose, an ounce, two or three times a-day,  
 as a tonic.

## WINE OF PERUVIAN BARK AND CALAMUS.

R. Bruised Peruvian bark  
 Calamus, each, one ounce and a-half  
 Quassia  
 Cinnamon  
 Elder flowers, each, six drachms  
 Alcohol three pints  
 Water eighteen pints  
 Macerate, express, strain, and add  
 Tincture of iron twelve ounces  
 Orange-flower water, one pint and a-half  
 Syrup six ounces  
 Mix. Dose, two drachms, morning and  
 evening, in leucorrhœa and gonorrhœa.  
*Cadet.*

## PERUVIAN BARK MIXTURE.

R. Peruvian bark six drachms  
 Confection of opium three drachms  
 Cremor tartar two drachms  
 Water six ounces  
 Mix. A tablespoonful every hour, in the  
 apyrexia. *Dewees.*

## SYRUP OF PERUVIAN BARK.

R. Extract of bark fourteen drachms  
 White wine two pounds  
 White sugar three pounds

Dissolve the extract in the wine, filter,  
 form a syrup with the sugar, with the aid  
 of a very moderate heat. *Monchon.*

## CINCHONIA.

## CINCHONIA.

An alkaloid existing in Peruvian bark; most  
 abundant in the pale kinds. It is white, trans-  
 lucent, and crystallizable, but little soluble in  
 water; very bitter, and has much the same pro-  
 perties as quinia.

## CINCHONIA.

R. Pale bark one thousand parts  
 Muriatic acid sixty-four parts  
 Water twelve thousand parts  
 Quicklime one hundred parts

Exhaust the bark by three successive boilings  
 with a third part of the acid and water, each  
 time, unite the decoctions, add the lime, diffused  
 in water, wash, and dry the precipitate; treat  
 it with alcohol, filter whilst hot, distil in a  
 water-bath, evaporate to dryness, digest in cold  
 alcohol, dissolve the residue in boiling alcohol,  
 with the addition of some animal charcoal, filter  
 whilst hot, and crystallize. *Par. Cod.*

## PILLS OF CINCHONIA.

R. Cinchonia one scruple  
 Conserve of roses sufficient  
 Mix, and form twenty-four pills. Dose,  
 two to eight. *Cadet.*

## MURIATE OF CINCHONIA.

R. Cinchonia at will  
 Muriatic acid sufficient  
 to saturate, filter, evaporate, and crystal-  
 lize. *Giordano.*

## KINATE OF CINCHONIA.

R. Alcoholic solution of sulphate  
 of cinchonia at will  
 Aqueous solution of kinate  
 of lime sufficient  
 to produce full precipitate; filter, evapo-  
 rate, redissolve, and crystallize. *Magendie.*

## SULPHATE OF CINCHONIA.

R. Cinchonia at will  
 Diluted sulphuric acid sufficient  
 to dissolve the cinchonia, evaporate, and  
 crystallize. *Guibourt.*

## PILLS OF SULPHATE OF CINCHONIA.

R. Sulphate of cinchonia half a drachm  
 Conserve of roses sufficient  
 Mix, and make thirty pills. Dose, one  
 to eight. *Cadet.*

## BOLUSES OF CINCHONIA.

R. Sulphate of cinchonia three grains  
 Crumb of bread  
 Honey  
 Liquorice, each, sufficient



Mix, and form boluses. One every two hours. *Brera.*

## SYRUP OF CINCHONIA.

R. Sulphate of cin-  
chonia forty-eight grains

Syrup one pint

Mix. A spoonful is a dose. *Cadet.*

R. Sulphate of cin-  
chonia twenty-four grains

Orange-flower water two drachms

Syrup twelve fl. ounces

Mix. As above. *Giordano.*

## WINE OF CINCHONIA.

R. Sulphate of cin-  
chonia twenty-four grains

Maderia wine two pints

Make solution, and filter. Dose, one to four ounces. *Cottureau.*

## TINCTURE OF CINCHONIA.

R. Sulphate of cinchonia twelve grains

Alcohol one ounce

Dissolve. *Foy.*

## MIXTURE OF SULPHATE OF CINCHONIA.

R. Sulphate of cinchonia three grains

Peppermint water three ounces

Syrup of capillaire one ounce

Mix. To be taken in four doses, during the day. *Cadet.*

## CINNAMOMUM.

## CINNAMON.

This is the bark of *Cinnamomum Zeylanicum*, and *C. aromaticum*, and also, perhaps, of other species; all natives of various parts of the tropical regions of Asia. That most commonly used in this country, is what is termed cassia, in commerce, and is imported from China.

*Ser. Syst.* Enneand. monog. *Nat. Syst.* Lauraceæ.

*Nees.* Laurineæ. 52. *Griffith, Med. Bot.* 555.

It is in tubular pieces, of a light brownish red color, of an aromatic, warm, agreeable taste, and of a fragrant, peculiar smell. It is a warm cordial stimulant. Dose, in powder, ten grains to a scruple.

## COMPOUND POWDER OF CINNAMON.

R. Powdered cinnamon  
ginger, each, two ounces

Cardamom seeds

Rasped nutmeg, each, one ounce

Rub together. *U. S. Ph.*

Stimulant, and carminative. Dose, ten to thirty grains, in debility of the stomach, with flatulence.

R. Powdered cinnamon one ounce

cloves

ginger

nutmeg, each, three drachms

Powdered mace two drachms

red saunders half an ounce

sugar thirty-two ounces

Mix well. Dose, from ten grains to a drachm. *Wirtemberg Ph.*

R. Powdered cinnamon twenty grains

cloves twelve grains

vanilla six grains

Rice flour six drachms

Sugar half an ounce

Mix. Dose, six grains to a scruple.

*Speilmann.*

## CINNAMON LOZENGES.

R. Powdered sugar

Cinnamon water, each, six ounces

Boil to syrup, and add

Powdered sugar one pound

Oil of cinnamon one drachm and a-half

Mix, make lozenges, and dry. *Taddei.*

## COMPOUND ELECTUARY OF CINNAMON.

R. Confection of orange peel, three ounces

nutmeg one ounce and a-half

ginger six drachms

Powdered cinnamon half an ounce

Syrup of orange peel sufficient

Mix. *Swediaur.*

## INFUSION OF CINNAMON.

R. Bruised cinnamon half an ounce

Boiling water two pints

Infuse. In dyspepsia and nervous colics. *Radius.*

## COMPOUND VINEGAR OF CINNAMON.

R. Cinnamon

Cloves

Nutmeg, each, one drachm

Peppermint half an ounce

Lavender

Rosemary, each, one ounce

Vinegar eight fl. ounces

Diluted alcohol twenty fl. ounces

Mix, and distil twenty-eight ounces. Used as an aromatic. *Van Mons.*

## COMPOUND WINE OF CINNAMON.

R. Powdered cinnamon

Cloves

Mace

Cardamom, each, half a drachm

White wine two pints

Boil, filter, and add

Sugar ten ounces

As a cordial stomachic and stimulant.

*Augustin.*

## TINCTURE OF CINNAMON.

R. Bruised cinnamon three ounces

Diluted alcohol two pints



Macerate for fourteen days, express, and filter. *U. S. Ph.*

Dose, one to four fl. drachms.

#### COMPOUND TINCTURE OF CINNAMON.

R. Bruised cinnamon one ounce  
cardamom seeds half an ounce  
ginger three drachms  
Diluted alcohol two pints  
Macerate for fourteen days, express, and filter. *U. S. Ph.*

A warm, aromatic tincture, useful in flatulence, gastrodynia, &c. Dose, one to two fl. drachms.

R. Bruised cinnamon one ounce  
cloves  
nutmeg  
Cut saffron, each, half an ounce  
Bruised calamus one drachm  
mace two drachms  
Fresh lemon peel no. two  
orange peel no. one  
Diluted alcohol one pint and a-half  
Digest, express, and filter. Dose fifty to eighty drops, as a stomachic. *Wirttemberg Ph.*

#### TINCTURE OF CINNAMON GARGLE.

R. Tincture of cinnamon one part  
Syrup of currants two parts  
Peppermint water sixteen parts  
Mix. In malignant sore throat. *Foy.*

#### ETHEREAL TINCTURE OF CINNAMON.

R. Bruised cinnamon three drachms  
cardamom half an ounce  
angelica one drachm and a-half  
long pepper one drachm  
Sulphuric ether five fl. ounces  
Diluted alcohol ten fl. ounces  
Macerate for eight days, express, and filter. Stomachic and carminative. Dose, thirty to fifty drops. *Niemann.*

#### WATER OF CINNAMON.

R. Oil of cinnamon half fl. drachm  
Carbonate of magnesia half a drachm  
Distilled water two pints  
Rub the oil with the magnesia, and both with the water, and filter. *U. S. Ph.*  
Mostly employed as a vehicle or adjuvant to other medicines.

#### COMPOUND CINNAMON WATER.

R. Sage  
Rosemary  
Peppermint  
Lavender, each, two parts  
Fennel  
Cinnamon, each, one part

Water one hundred and twenty parts  
Mix, and distil seventy-two parts. In spoonful doses. *Saxon Ph.*

#### OIL OF CINNAMON.

R. Bruised cinnamon at will  
Water sufficient  
Distil, and separate the oil. *Par. Cod.*

#### CINNAMON SPICE PLASTER.

R. Olibanum three ounces  
Yellow wax half an ounce  
Powdered cinnamon six drachms  
Oil of pimento  
lemons, each, two drachms  
Melt the wax and olibanum, and add the cinnamon, rubbed with the oils. *Coxe.*

### COCCULUS.

#### COCCULUS INDICUS.

This is the fruit of *Anamirta cocculus*, a woody vine, native of the East Indies.

*Ser. Syst.* Diœc. dodecand. *Nat. Syst.* Menispermaceæ.

Wight and Arnott, 1. 449. Griffith, Med. Bot. 105.

They are about the size of a pea, roundish, with a thin, dry, blackish, external coat, inclosing another of a lighter color, containing an oily, bitter, white kernel. They are inodorous, but of a permanent, bitter taste. They are acro narcotic, and owe their properties to the presence of a peculiar principle, called *picrotoxin*. They are seldom used in medicine.

#### OINTMENT OF COCCULUS INDICUS.

R. Cocculus indicus at will  
Separate the kernels, beat them in a mortar, first alone, then with a little lard, finally, add lard equal to five times the weight of kernels. *Ed. Ph.*

Used for the destruction of vermin, and in the treatment of ringworms and scabies of the scalp.

#### OINTMENT OF PICTROTOXIN.

R. Picrotoxin six grains  
Lard one ounce  
Mix well. In tinea. *Radius.*

### COCCUS.

#### COCHINEAL.

Cochineal is a small, hemipterous insect, inhabiting Mexico and some parts of South America, living principally on different species of cactus, and especially the *Opuntia cochinillefera*. It is in grains of reddish-black, often covered with a white powder. It has a feeble odor, and a bitterish, somewhat acidulous taste. The powder is of a reddish-purple color, tinging fluids of a deep red. It is said to be somewhat anodyne



and antispasmodic, but is principally used as a coloring agent.

#### TINCTURE OF COCHINEAL.

R. Powdered cochineal                    one part  
Diluted alcohol                    eight parts  
Macerate eight days, express, and filter.  
Advised in convulsive coughs.      *Beral.*

#### MIXTURE OF COCHINEAL AND CARB. POTASSA.

R. Carbonate of potassa            one scruple  
Powdered cochineal            half a scruple  
Sugar                    one drachm  
Distilled water            four fl. ounces  
Make mixture. Dose, for children, a teaspoonful, every two or three hours. A highly popular remedy in whooping-cough.      *Ellis.*

#### MIXTURE OF COCHINEAL AND COMMON SALT.

R. Common salt            three drachms  
Powdered cochineal            fifteen grains  
Divide into six doses. One to be taken every morning, the last followed by a purge. As a vermifuge.      *Rush.*

### COCHLEARIA.

#### SCURVY GRASS.

Scurvy grass is the *Cochlearia officinalis*, an herbaceous plant, native of the northern parts of Europe, and cultivated there and in the United States as a garden herb.

*Sex. Syst.* Tetradyn. silic. *Nat. Syst.* Brassicaceæ.

The whole herb is used; it has a penetrating and acrid smell when bruised, and a pungent, bitter taste. It is antiscorbutic, aperient, and diuretic.

#### ELECTUARY OF SCURVY GRASS.

R. Conserve of scurvy  
grass                    one ounce and a-half  
Calamus                    one drachm  
Syrup of orange peel  
Aromatic sulphuric acid, each, sufficient  
Mix. A teaspoonful every two hours, as an antiscorbutic.      *Augustin.*

#### GARGLE OF SCURVY GRASS.

R. Spirit of scurvy grass            one drachm  
Syrup of Peruvian bark            half an ounce  
Infusion of water cress            eight ounces  
Mix. As a gargle in scurvy.      *Foy.*

### CODEIA.

#### CODEIA.

An alkaloid obtained from opium, and bearing the same relation to morphia that cinchonia does

to quinia. It is precipitated by tannin, and is soluble in ether, but is not thrown down from a dilute solution of its salts by ammonia. It acts somewhat like morphia, but produces great itching of the skin.

#### CODEIA.

R. Opium                    at will  
Macerate in a sufficient quantity of water, decant, treat with chloride of calcium, filter, evaporate, and crystallize; decompose by ammonia, filter, concentrate fluid, crystallize, dissolve the crystals in water, add a slight excess of caustic potassa, dissolve the precipitate in alcohol and ether, and crystallize. Dose, one to two grains, in neuralgia, &c.      *Cottreau.*

R. Mother water of morphia (by Gregory's process)            at will  
Evaporate and crystallize; re-dissolve the crystals, and re-crystallize. Treat with solution of potassa, which dissolves the morphia and precipitates the codeia; wash the latter with a little water, dissolve in hot ether, and suffer the solution to evaporate spontaneously.

#### SYRUP OF CODEIA.

R. Codeia                    twenty-four grains  
Distilled water            four fl. ounces  
Sugar                    eight ounces  
Reduce codeia to fine powder, triturate with one-third of water, allow to settle and decant; treat residue with the remainder of water, at two operations. Treat in a matrass on water-bath till quite dissolved, add sugar, and dissolve.      *Cap.*

Dose, one drachm, night and morning; gradually increased in neuralgic irritation of the stomach, whooping-cough, &c.

#### MURIATE OF MORPHIA AND CODEIA.

R. Opium                    at will  
Treat with water, add a strong solution of chloride of calcium, filter with animal charcoal, evaporate, and crystallize. *Magendie.*  
Dose, half a grain to a grain.

### COFFEA.

#### COFFEE.

Coffee is the product principally of *Coffea Arabica*, a small tree, native of the eastern coast of Africa, but now cultivated in most tropical regions. The varieties are very numerous, and in some cases probably are derived from other species.

*Sex. Syst.* Pentand. monog. *Nat. Syst.* Cinchonaceæ.

Linn. Sp. Pl. 245. Griffith, Med. Bot. 361.

Raw coffee has febrifuge properties, and has been used with success in the treatment of inter-



mittents. In a roasted state it acts on the nervous system, producing wakefulness, and also somewhat excites the circulatory system. It is said to be useful in periodic asthma, and is an efficient anti-emetic.

#### DECOCTION OF RAW COFFEE.

R. Raw coffee one ounce  
Water eighteen ounces  
Boil to two-thirds, strain, and add  
Lemon juice two ounces  
To be taken warm, during the apyrexia.  
*Foy.*

#### VINEGAR OF COFFEE.

R. Roasted coffee, ground, three ounces  
Vinegar twelve ounces  
Boil, strain, and add  
Sugar one ounce and a-half  
Two spoonfuls occasionally, in poisoning by opium, after the evacuation of the poison.  
*Pierquin.*

#### SYRUP OF COFFEE.

R. Coffee, roasted, four ounces  
Water two pints  
Refined sugar three pounds  
Infuse the coffee in a pint and a-half of the water for six hours, boil in a water-bath, let settle, decant, and add remainder of the water. Let stand for some hours; decant, add sugar, form syrup, and strain. Dose, from one ounce to two.  
*Ferrari.*

### COLCHICUM.

#### COLCHICUM.

The *Colchicum autumnale*, or meadow saffron, is a bulbous plant, a native of most parts of Europe, flowering in the autumn, and putting forth its leaves, and perfecting its seed the succeeding spring.

*Sex. Syst.* Hexand. monog. *Nat. Syst.* Melanthaceæ.

Linn. Sp. Pl. 485. Griffith, Med. Bot. 644.

The parts used are the bulb and the seeds. The first is of a brown color externally, white within, rounded on one side, flattened on the other, of scarcely any smell, but an acrid, bitter taste; the seeds are small, rounded, of a brownish yellow color, and of the same taste as the bulb. They owe their properties to the presence of colchicia.

Colchicum is principally used in the various forms of gout and rheumatism. It is an active irritant, and in large doses, acro-narcotic. The dose of the powdered bulb is two to eight grains, every four or six hours, that of the seeds much the same.

#### EXTRACT OF COLCHICUM BULB.

R. Fresh colchicum bulb one pound  
Bruise in stone mortar, express juice, evaporate to proper consistence. *Lond. Ph.*

#### ACETIC EXTRACT OF COLCHICUM BULB.

R. Fresh colchicum bulb one pound  
Acetic acid three fl. ounces  
Bruise, gradually adding acid, express, and evaporate in porcelain vessel to proper consistence. *Lond. Ph.*

Dose, one to two grains, two or three times a-day.

#### COLCHICUM PILLS.

R. Powdered colchicum three grains  
Soap sufficient  
Make three pills, to be taken during the day, gradually increasing to five or six.  
*Ritton.*

#### COMPOUND COLCHICUM PILLS.

R. Acetic extract of colchicum three grains  
Dover's powder  
Compound extract of colocynth, each, one grain  
Mix, and make a pill. In gout, after having taken, for some days, thirty to forty drops of colchicum wine, morning and evening. *Halford.*

R. Extract of colchicum root two grains  
foxglove one grain  
col., comp., twenty grains  
Mix, and form five pills. One at night, in gout. *Lartiques.*

#### WINE OF COLCHICUM ROOT.

R. Colchicum root, bruised, one pound  
Sherry wine two pints  
Macerate for fourteen days, express, and filter. *U. S. Ph.*

#### WINE OF COLCHICUM SEEDS.

R. Colchicum seed, bruised, four ounces  
Sherry wine two pints  
Macerate for fourteen days, express, and filter. *U. S. Ph.*

The dose of the first, which is saturated, is from ten drops to a fl. drachm; for the second, one to two fl. drachms.

R. Colchicum three ounces  
Sherry wine, deprived of its spirit by evaporation two ounces  
Macerate for six days, and filter. *Taddei.*

This is said to be the formula of the *Eau medicinale of Husson.*

#### TINCTURE OF COLCHICUM SEEDS.

R. Colchicum seed, bruised, four ounces  
Diluted alcohol two pints  
Macerate for fourteen days, express, and filter. *U. S. Ph.*

Dose, from half a fl. drachm to two fl. drachms. Sometimes used as an embrocation in gout, rheumatism, and neuralgia.



## COMPOUND TINCTURE OF COLCHICUM SEEDS.

R. Colchicum seeds, bruised, five ounces  
Aromatic spirit of ammonia two pints  
Macerate for fourteen days, express, and filter. *Lond. Ph.*

More stimulating than the last. Dose, thirty drops to a fl. drachm.

## TINCTURE OF COLCHICUM FLOWERS.

R. Dried colchicum flowers one ounce  
Diluted alcohol one pint  
Macerate for eight days, express, and filter. Advised in acute rheumatism. *Bushell.*

## VINEGAR OF COLCHICUM ROOT.

R. Dried colchicum root two ounces  
Distilled vinegar two pints  
Alcohol one fl. ounce  
Macerate the root in the vinegar for seven days, express, and let dregs subside, pour off and add alcohol. *U. S. Ph.*

As a diuretic in dropsy, also given in gout, rheumatism, &c. Dose, thirty drops to two fl. drachms.

## OXYMEL OF COLCHICUM.

R. Fresh colchicum root, sliced, one ounce  
Distilled vinegar one pint  
Clarified honey two pounds  
Macerate the colchicum in the vinegar for two days, express, and strain; add honey, and boil to consistence of syrup. *Dub. Ph.*  
Dose, one fl. drachm twice a-day.

## VINEGAR OF COLCHICUM SEEDS.

R. Colchicum seeds one part  
Distilled vinegar four parts  
Macerate a month, and filter. *Beral.*

## SYRUP OF COLCHICUM.

R. Vinegar of colchicum root sixteen parts  
Sugar twenty-six parts  
Boil, by a gentle fire. *Swed. Pharm.*

## MIXTURE OF COLCHICUM AND MAGNESIA.

R. Magnesia one drachm  
Sugar  
Gum Arabic, each, sufficient  
Distilled water four fl. ounces  
Wine of colchicum forty drops  
Mix. A tablespoonful every two hours, till it operates. In gout and rheumatism. *Ellis.*

## MIXTURE OF COLCHICUM AND SULPHATE OF MAGNESIA.

R. Sulphate of magnesia one to two ounces  
Mint water ten fl. ounces  
Vinegar of colchicum  
Syrup of saffron, each, one fl. ounce  
Magnesia eight scruples

Mix. Dose, one to three tablespoonfuls, every two hours, till four to six evacuations are procured in the twenty-four hours. In gout. *Scudamore.*

## COMPOUND WINE OF COLCHICUM.

R. Tincture of colchicum seeds  
one scruple to half a fl. drachm  
Wine of colchicum seeds half a fl. ounce  
Mix. Fifteen to forty drops, three times a-day. *Weber.*

## TINCTURE OF COLCHICUM MIXTURE.

R. Tincture of colchicum seeds  
guaiaicum, each,  
three fl. drachms  
Mix. Thirty to forty drops, three times a-day, in chronic rheumatism. *Radius.*

## TINCTURE OF COLCHICUM AND DIGITALIS.

R. Tincture of colchicum seed  
digitalis, each,  
two fl. drachms  
Nitric ether one scruple

Mix. Twenty drops, morning and evening, on sugar, in hydrothorax. *Hildebrand.*

## COLCHICUM MIXTURE.

R. Wine of colchicum seed thirty drops  
Denarcotized laudanum twenty-five drops  
Sugar thirty grains  
Water one ounce  
Mix. Found useful in acute rheumatism. To be taken at night. *Dewees.*

## COLCHICUM MIXTURE.

R. Wine of colchicum seeds forty drops  
Acetic tincture of opium twenty drops  
Sugar thirty grains  
Water one ounce

Mix. In gout, after inflammation is allayed. To be taken at bedtime. *Dewees.*

R. Expressed juice of colchicum  
flowers two parts  
Brandy one part

Mix, and let rest for a few days; decant, and keep for use in well-stopped bottles. *Wilson.*

Said to be identical in every respect to the *Eau medicinale of Husson.*

## MIXTURE OF COLCHICUM AND ELATERIUM.

R. Elaterium one grain  
Spirit of nitric ether two fl. ounces  
Tincture of squills  
Oxymel of colchicum, each,  
half a fl. ounce  
Syrup one fl. ounce  
Mix. A teaspoonful, three or four times a-day, in hydrothorax. *Ferriar.*



## VINEGAR OF COLCHICUM MIXTURE.

R. Vinegar of colchicum  
Syrup, each, half a fl. ounce  
Carbonate of magnesia one drachm  
and a-half

Peppermint water four ounces  
Mix. In tablespoonful doses, as a sudorific in gout, &c. *Foy.*

## MIXTURE OF COLCHICUM AND SQUILLS.

R. Oxymel of colchicum  
squills  
Tincture of tobacco, each, half fl. ounce

Mix. A teaspoonful four times a-day. As a diuretic in dropsy. Requires caution.

*Ferriar.*

## MIXTURE OF WINE OF COLCHICUM.

R. Wine of colchicum half fl. drachm  
Carbonate of magnesia one drachm  
Cinnamon water

Distilled water, each, three fl. ounces  
Mix. A tablespoonful, three times a-day. As a diuretic. *Ellis.*

## MIXTURE OF COLCHICUM AND AMMONIA.

R. Oxymel of squills  
Solution acetate of ammonia, each, two fl. ounces  
Parsley water six fl. ounces

Mix. A spoonful, every two hours, as a diuretic. *Augustin.*

## LINIMENT OF COLCHICUM AND CAMPHOR.

R. Tincture of colchicum root  
camphor, each, three fl. ounces

Mix. As an embrocation, in gout, rheumatism, and neuralgia. *Leycock.*

## COLLINSONIA.

## HORSE BALM.

The *Collinsonia Canadensis* is a native plant, found in most parts of this country, in woods; it is from two to three feet in height.

*Sex. Syst.* Diand. monog. *Nat. Syst.* Lamiaceæ.  
Linn. Sp. Pl. 39. Griffith, Med. Bot. 513.

The whole plant has a strong, peculiar, unpleasant odor, and a warm, pungent taste: these are most developed in the root; this is knotty and hard. It is diuretic, diaphoretic, and tonic; and is popularly used in diseases of the bladder, leucorrhœa, and dropsy, and externally as a fomentation to wounds, ulcers, &c.

## DECOCTION OF HORSE BALM.

R. Horse balm root, bruised, two ounces  
Water two pints  
Boil for half an hour, and strain. Dose, a wineglassful, every two hours, in dropsy.

## TINCTURE OF HORSE BALM.

R. Bruised horse balm root two ounces

Diluted alcohol one pint  
Macerate for fourteen days, express, and filter. Dose, a teaspoonful.

## OIL OF HORSE BALM.

R. Horse balm at will  
Water sufficient  
Distil, return product, and redistil, collect product. Dose, five to ten drops.

## COLOCYNTHIS.

## COLOCYNTH.

Colocynth is the dried pulp of the fruit of *Citrullus colocynthis*, a cucurbitaceous vine found in many parts of Asia and Africa. The fruit is a round pepo, of the size and color of an orange, with a thin, but hard rind, and containing numerous seeds, enveloped in a white, spongy pulp.

*Sex. Syst.* Monœc. monadelph. *Nat. Syst.* Cucurbitaceæ.

Royle, Mat. Med. 396. Griffith, Med. Bot. 306.

It occurs in the shops in white, round, light balls, composed of a spongy, dried pulp, enclosing numerous seeds. This spongy substance has a faint odor, but an intensely bitter and nauseous taste. It is a powerful drastic and hydragogue purgative, and is seldom given alone.

## POWDER OF COLOCYNTH.

R. Colocynth one to three grains  
Gum Arabic

Liquorice

Sugar, each, five grains

Mix. As a hydragogue purgative.

*Augustin.*

## EXTRACT OF COLOCYNTH.

R. Sliced colocynth one pound  
Distilled water two gallons

Mix, and boil over a slow fire for six hours, occasionally adding more water. Strain while hot, and evaporate to proper consistence. *Lond. Ph.*

Dose, from five grains to half a drachm.

R. Colocynth pulp at will  
Diluted alcohol nine grains

Macerate the pulp in a sufficiency of menstruum to cover it for twenty-four hours, displace till exhausted, and evaporate to the proper consistence. This extract is more active than the aqueous preparation.

*W. Procter.*

## COMPOUND EXTRACT OF COLOCYNTH.

R. Colocynth sliced six ounces  
Powdered aloes twelve ounces

scammony four ounces

cardamom one ounce

Castile soap three ounces

Diluted alcohol one gallon

Macerate the colocynth in the alcohol, with



a mild heat for four days, express, and filter, then add the aloes, scammony, and soap, evaporate to proper consistence, and add the cardamom. *U. S. Ph.*

A valuable and safe cathartic. In small doses, laxative. Dose, five to thirty grains.

#### COMPOUND PILLS OF COLOCYNTH.

R. Compound extract of colocynth, powdered, half an ounce  
Extract of jalap, in powder,  
Calomel, each, three drachms  
Gamboge, in powder, two scruples  
Mix, and with water form mass, to be divided into one hundred and eighty pills. *U. S. Ph.*

A most excellent cathartic, when it is wished to act on the biliary organs. Dose, one to three or four pills.

R. Powdered aloes  
Scammony, each, eight parts  
Colocynth four parts  
Sulphate of potassa  
Oil of cloves, each, one part  
Rectified spirit sufficient  
Pulverize the aloes, scammony, and sulphate of potassa, together; mix the colocynth with them, add oil of cloves, and with the rectified spirit beat into a mass, to be divided into five-grain pills. *Ed. Ph.*

Dose, five to fifteen grains.

R. Compound extract of colocynth one scruple  
Alcoholic extract of jalap six grains  
Compound powder of scammony  
Calomel, each, ten grains  
Tartar emetic one grain  
Castile soap five grains  
Oil of cinnamon four drops

Beat well together, and form fifteen pills. Dose, one to three, at bedtime. *Meigs.*

R. Extract of colocynth two drachms  
Resin of jalap one drachm  
Soap one drachm and a-half  
Guaiaicum three drachms  
Tartar emetic eight grains  
Oil of juniper  
Rosemary, each, four drops  
Syrup of buckthorn sufficient  
Mix well, and divide into four-grain pills. Dose, one to three. *Barclay.*

#### PILLS OF COLOCYNTH AND HENBANE.

R. Extract of colocynth two parts  
henbane one part  
Beat together, with a few drops of rectified spirit, and divide into five-grain pills. *Ed. Ph.*

Dose, five to twenty grains.

#### COLOCYNTH CLYSTER.

R. Extract of colocynth one scruple  
Common salt three drachms  
Syrup of dogtooth grass one ounce and a-half  
Infusion of chamomile five ounces  
In cerebral affections. *Phœbus.*

#### COLOCYNTH MIXTURE.

R. Colocynth one drachm  
Boiling water six ounces  
Boil for ten minutes, strain, add, when cold,  
Hoffmann's anodyne one drachm  
Syrup of orange peel one ounce  
Mix. A spoonful, three times a-day, in ascites, or hydrothorax. *Augustin.*

#### TINCTURE OF COLOCYNTH.

R. Colocynth eight parts  
Star anise one part  
Alcohol ninety-six parts  
Macerate for three days, and filter. Dose, fifteen to twenty drops. *Van Mons.*

#### TINCTURE OF COLOCYNTH MIXTURE.

R. Tincture of colocynth two drachms  
Solution of antimoniated soap six drachms  
Mix. Dose, twenty to thirty drops a-day, in lepra and obstinate cutaneous affections. *Heim.*

#### COLOMBA.

##### COLUMBO.

Columbo is the root of *Cocculus palmatus*, a climbing plant, a native of Mozambique.

*Sex. Syst. Diœc. hexand. Nat. Syst. Menispermaceæ.*

Hooker, Bot. Mag. 2970-71. Griffith, Med. Bot. 103.

As found in the shops, the root is in round slices, externally of a brown, wrinkled appearance, internally yellow. It is somewhat aromatic, and has a very bitter taste. It is an excellent bitter tonic, with no astringency. It is useful in diseases where the pure bitters are required, and generally agrees with the stomach. Dose, in powder, is from ten to thirty grains.

#### POWDER OF COLUMBO AND IRON.

R. Powdered columbo  
Subcarbonate of iron  
Powdered rhubarb  
ginger, each, one drachm  
Mix, and make eleven powders; one to be taken every four hours. *A. T. Thomson.*

#### POWDER OF COLUMBO AND MAGNESIA.

R. Powdered columbo ten grains  
Magnesia two scruples  
Mix. For a dose. In cardialgia. *Brugnatelli.*



## POWDER OF COLUMBO AND GINGER.

R. Powdered columbo  
subcarbonate of iron  
rhubarb  
ginger, each, one drachm  
Mix, and divide into twelve powders. One  
to be taken three times a-day, in syrup. *Ellis.*

## POWDER OF COLUMBO AND TARTRATE OF IRON.

R. Tartrate of iron and  
potassa two scruples  
Powdered columbo half a drachm  
Mix, and divide into four powders. One  
every three or four hours, in syrup. *Ellis.*

## COMPOUND PILLS OF COLUMBO.

R. Powdered columbo four scruples  
Opium four grains  
Oil of peppermint ten drops  
Syrup of pinks sufficient  
Beat into mass, and form thirty pills. Two,  
three times a-day, in spasmodic vomiting.  
*St. Marie.*

R. Powdered columbo one drachm  
rhubarb two scruples  
Extract of chamomile two drachms  
Oil of caraway five drops  
Syrup of saffron sufficient  
Form mass, and divide into pills of four  
grains. Four to be taken a-day, in mania  
with amenorrhœa. *Augustin.*

## INFUSION OF COLUMBO.

R. Bruised columbo half an ounce  
Boiling water one pint  
Macerate two hours in a covered vessel,  
and strain. *U. S. Ph.*

Dose, two fl. ounces, three or four times a-day.

## INFUSION OF COLUMBO AND GINGER.

R. Bruised columbo one ounce  
ginger two drachms  
Boiling water one pint  
Infuse, and strain. A wineglass, cold,  
every two hours, in chronic diarrhœa.  
*Ellis.*

## INFUSION OF COLUMBO, RHUBARB, &amp;C.

R. Bruised caraway  
columbo  
rhubarb, each, one scruple  
Boiling water eight fl. ounces  
Digest for two hours, and strain.

R. To strained liquid three half fl. ounces  
Add  
Tincture of rhubarb one fl. drachm  
Syrup of ginger two fl. drachms  
Mix. Dose, a teaspoonful to a tablespoon-  
ful, in diarrhœa. *Ellis.*

## MIXTURE OF COLUMBO.

R. Columbo half a drachm  
Boil in  
Water three to five fl. ounces  
Strain, and add  
Carbonate of potassa ten grains  
Lemon juice three fl. drachms  
Tincture of opium twelve drops  
Mix. A tea to a table spoonful, every  
hour, as an anti-emetic. *Ellis.*

## COMPOUND DECOCTION OF COLUMBO.

R. Columbo  
Quassia, each, two drachms  
Orange peel one drachm  
Rhubarb one scruple  
Subcarbonate of potassa half a drachm  
Water twenty fl. ounces  
Boil down to a pint, strain, and add  
Tincture of lavender half fl. ounce  
*Coxe.*

## EXTRACT OF COLUMBO.

R. Bruised columbo two parts  
Alcohol three parts  
Water nine parts  
Macerate for twenty-four hours, distil off  
the alcohol, and evaporate to proper con-  
sistence. *Cottureau.*

## TINCTURE OF COLUMBO.

R. Columbo bruised four ounces  
Diluted alcohol two pints  
Macerate for fourteen days, and filter.  
*U. S. Ph.*

Dose, one to four fl. drachms.

## MIXTURE OF COLUMBO AND CASCARILLA.

R. Powdered columbo half an ounce  
Water ten fl. ounces  
Boil to six ounces, add, towards close,  
Cascarilla two drachms  
Strain, and add  
Tincture of orange peel two drachms  
Syrup of cinnamon one ounce  
A spoonful every hour, in chronic diarrhœa.  
*Berends.*

## MIXTURE OF COLUMBO AND SALEP.

R. Extract of columbo one drachm  
Decoction of salep three ounces  
Fennel sugar two scruples  
Mix. Two spoonfuls a-day, in the diarrhœa  
of children. *Wendt.*

## COMPTONIA.

## SWEET FERN.

A shrubby, indigenous plant, found in the  
northern and middle States.

*Sex. Syst. Monœc. triand. Nat. Syst. Myricaceæ.*  
*Aiton, Hort. Rev. iii. 334. Griffith. Med. Bot.*  
*584.*



It is aromatic, especially when bruised. It is a mild astringent, and is used in bowel complaints, in the form of decoction.

## CONIUM.

## HEMLOCK.

This is an umbelliferous, poisonous plant, the *Conium maculatum*, a native of Europe, but naturalized in many parts of this country, growing in waste places.

*Ser. Syst.* Pentand. digyn. *Nat. Syst.* Apiaceae.

*Linn. Sp. Pl.* 349. *Griffith, Med. Bot.* 339.

Both the seeds and the leaves are used. The latter have a strong, unpleasant, narcotic odor, and a somewhat bitterish taste; the seeds possess these qualities in a less degree. They are both narcotic, but neither stimulant nor sedative, and are given in a variety of complaints, to alleviate, especially in malignant tumors, in which they are thought to exert a curative influence. The dose, in powder, is from three to four grains of the leaves; and of the seeds, rather less; twice a-day, gradually increasing.

## POWDER OF HEMLOCK.

R. Powdered hemlock five grains  
liquorice six grains  
Make a powder, to be taken three times a-day. In scirrhus affections, scrofula, old ulcers, &c.

## FECULA OF HEMLOCK.

R. Hemlock juice at will  
Place on the fire, remove the coagulum, wash it well, dry, and pulverize. Dose, five to twenty grains. *Van Mons.*

## EXTRACT OF HEMLOCK.

R. Fresh hemlock leaves one pound  
Bruise in a stone mortar, sprinkling on them a little water; express the juice, heat to boiling point, strain, and evaporate to proper consistence. *U. S. Ph.*

Dose, one grain, morning and evening, gradually increased till it affects the system.

## ALCOHOLIC EXTRACT OF HEMLOCK.

R. Hemlock leaves, in coarse powder, one pound  
Diluted alcohol four pints  
Moisten powder with half a pint of the alcohol, and allow to macerate for twenty-four hours, place in percolating apparatus and add remainder of alcohol; when this has disappeared, add water so as to keep powder covered. Stop process when the liquid passing, causes precipitate in that previously obtained. Distil off alcohol, and evaporate to proper consistence.

*U. S. Ph.*

Dose, one to two grains, twice a-day, gradually increasing.

## PILLS OF HEMLOCK AND IPECACUANHA.

R. Extract of hemlock  
Dover's powder, each, ten grains  
Mix, and form five pills. Two at bed-time, in pulmonary irritation, with rheumatic and other local pains. *Ellis.*

R. Extract of hemlock five drachms  
Ipecacuanha one drachm  
Mucilage of gum Arabic sufficient  
Mix, and make four grains.

## PILLS OF HEMLOCK AND CALOMEL.

R. Extract of hemlock two scruples  
Calomel fifteen grains  
Mix, and form fifteen pills. One, three times a-day, in syphiloid affections. *Ellis.*

## PILLS OF HEMLOCK AND DANDELION.

R. Extract of hemlock  
dandelion  
Gum Arabic  
Soap, each, one drachm  
Honey sufficient  
Mix, and form three-grain pills. In glandular and other tumors. *Brera.*

## INFUSION OF HEMLOCK.

R. Hemlock leaves half an ounce  
Boiling water one pint  
Infuse and strain. As a wash to cancerous and malignant ulcers. *Ellis.*

## HEMLOCK COLLUTORY.

R. Hemlock leaves two drachms  
Poppy seeds one drachm  
Henbane seeds half a drachm  
Milk half a pint  
Boil slightly, and strain. In rheumatic toothache. *Phæbus.*

## TINCTURE OF HEMLOCK.

R. Hemlock leaves four ounces  
Diluted alcohol two pints  
Macerate for fourteen days, express, and filter. *U. S. Ph.*

Dose, thirty drops to a fl. drachm.

## ETHEREAL TINCTURE OF HEMLOCK.

R. Hemlock leaves one part  
Sulphuric ether four parts  
Macerate for two days, and filter. *Soubéiran.*

## OINTMENT OF HEMLOCK.

R. Fresh hemlock leaves  
Prepared lard, each, two pounds  
Boil till the leaves becomes crisp, then express through linen. *Dub. Ph.*

R. Juice of hemlock leaves one part  
Lard four parts  
Boil till moisture is expelled, and express



through a cloth. As a dressing to painful ulcers.

*Swediaur.*

#### OIL OF HEMLOCK.

R. Fecula of hemlock one part  
Olive oil four parts  
Mix, and strain. As a friction to the abdomen in ileus, and to the anus in hemorrhoids.

*Van Mons.*

#### HEMLOCK PLASTER.

R. Elemi two parts  
White wax one part  
Melt together, and add  
Alcoholic extract of hemlock nine parts  
As an application to painful ulcers and malignant tumors.

*Guibourt.*

R. Juice of hemlock twelve parts  
Powdered hemlock six parts  
Evaporate to a pasty consistence, and add to, a melted mixture of  
Yellow wax ten parts  
Turpentine four parts  
Olive oil six parts  
Used as above.

*Pideret.*

#### COMPOUND HEMLOCK PLASTER.

R. Yellow wax eight parts  
Resin sixteen parts  
Fecula of hemlock forty-eight parts  
Melt together, and add a melted mixture of  
Gum ammoniac four parts  
Turpentine one part  
Stir well till cold. Used as above.

*Van Mons.*

R. Hemlock plaster one ounce  
Burgundy pitch  
Diachylon, each, half an ounce

Mix, and spread, then sprinkle with from six to twelve grains of tartar, according to the age of the patient. Said to be very useful in whooping-cough.

R. Soap plaster one ounce  
Extract of hemlock half an ounce  
Powdered belladonna two drachms  
Mix.

*Hamburg Cod.*

#### CATAPLASM OF HEMLOCK.

R. Powdered hemlock  
Linseed meal, each, one ounce  
Fecula of hemlock  
Boiling water, each, sufficient

Mix. As an application to cancerous and other painful ulcers.

*Guibourt.*

R. Hemlock leaves a handful  
Henbane leaves half a handful  
Lard half an ounce  
Rose oil one ounce  
Decoction of mallows sufficient

Mix, and boil to the consistence of cataplasm.

*Pierquin.*

R. Powdered hemlock two ounces  
Carrot juice one pound  
Tincture of opium and saffron three drachms  
Linseed meal sufficient  
Make a cataplasm. In cancerous ulcers.

*Piderot.*

#### MIXTURE OF HEMLOCK AND PAREGORIC.

R. Extract of hemlock one drachm  
Paregoric elixir  
Syrup of tolu, each, half fl. ounce  
Rose water four fl. ounces  
Mix. From half to a teaspoonful once in four hours, to a child a year old, watching the effect, in pertussis.

*Pearson.*

#### SUPPOSITORY OF HEMLOCK.

R. Extract of hemlock  
White wax, each, one part  
Butter of cocoa four parts  
Mix, in painful hemorrhoids and spasms of the rectum.

*Beral.*

### CONTRAYERVA.

#### CONTRAYERVA.

Contrayerva is the root of *Dorstenia contrayerva*, and other species, all natives of South America, Mexico, and the West Indies. They are perennial, dwarf, herbaceous plants, usually growing in high, rocky places.

*Sex. Syst. Tetrand. monog. Nat. Syst. Monacæ.*

Linn. Mat. Med. No. 124. Griffith, Med. Bot. 577.

The root is oblong, hard, rough, and solid, of a reddish-brown color, with numerous, long, yellowish fibres. It has an aromatic odor, and a warm, pungent, somewhat bitterish taste. It is a stimulant, tonic, and diaphoretic, and has been found useful in low states of the system. The dose of the powder is about half a drachm.

#### COMPOUND POWDER OF CONTRAYERVA.

R. Powdered contrayerva six drachms  
Virginia snakeroot two drachms  
Prepared chalk one ounce  
Mix. Dose, a teaspoonful, two or three times a day, as a tonic.

*Spielmann.*

#### EXTRACT OF CONTRAYERVA.

R. Contrayerva, bruised, one part  
Boiling water six parts  
Infuse for three days, boil, strain, express, and evaporate to proper consistence. Dose, one to two scruples.

*Palat. Cod.*

#### DECOCTION OF CONTRAYERVA.

R. Contrayerva, bruised, one drachm



Water one pint  
Boil for ten minutes, and strain.

*Taddei.*

#### GARGLE OF CONTRAYERVA.

R. Contrayerva half an ounce  
Figs one ounce  
Water eighteen fl. ounces  
Mix, and boil down to two-thirds.

*Saunders.*

#### TINCTURE OF CONTRAYERVA.

R. Contrayerva one part  
Diluted alcohol eight parts  
Digest in one-half the alcohol for four days, strain and digest with the other half of the alcohol; strain, and filter the united liquids. Dose, thirty to forty drops.

*Guibourt.*

#### COMPOUND TINCTURE OF CONTRAYERVA.

R. Contrayerva  
Virginia snakeroot, each,  
one ounce and a-half  
Juniper berries one ounce  
Spirit of ammonia six fl. ounces  
Alcohol one pint  
Macerate for four days, express, and filter.  
Dose, forty to eighty drops, as a stimulating diaphoretic.

*Lip. Dispen.*

#### CONTRAYERVA MIXTURE.

R. Powdered contrayerva eight grains  
Cinnamon water one fl. ounce and a-half  
Nutmeg water two fl. drachms  
Syrup of cinnamon three drachms  
Mix. A spoonful as a stomachic.

*Swediaur.*

R. Gum tragacanth  
Comp. contrayerva powder, each,  
one scruple  
Distilled water one fl. ounce  
Cinnamon water three fl. drachms  
Syrup of saffron one fl. drachm  
Mix. As a quieting draught for a woman after delivery.

*Merriman.*

### CONVOLVULUS PANDURATUS.

#### WILD POTATO.

This is a native plant, with a large perennial root, found in sandy soils in most parts of the United States.

*Sex. Syst.* Pentand. monog. *Nat. Syst.* Convolvulaceæ.

Linn. Sp. Pl. 219. Griffith, Med. Bot. (Ipomæa) 476.

The root, as found in the shops, is in circular pieces, of a yellowish-brown color, scarcely any smell, and a bitterish, somewhat acrid taste. It is feebly cathartic, but is said to be a good diuretic

in calculous complaints. Dose, as purgative, forty grains. As a diuretic, it is given in decoction.

### COPAIBA.

#### COPAIBA.

This is the juice of *Copaifera officinalis* and other species, all large trees, natives of South America and principally of Brazil.

*Sex. Syst.* Decand. monog. *Nat. Syst.* Fabaceæ.

Linn. Sp. Pl. 557. Griffith, Med. Bot. 264.

This juice or balsam is clear and transparent, of a pale yellowish color; a peculiar and somewhat unpleasant odor, and a bitterish, pungent, nauseous taste. It is rather more consistent than olive oil when fresh, but, by age, becomes thicker, and may even assume the solidity of a resin. It is stimulant, diuretic, and cathartic, and is principally used in morbid discharges from mucous surfaces, especially those of the genito-urinary organs. The dose is from twenty drops to a fl. drachm, three times a-day.

#### BOLUSES OF COPAIBA.

R. Copaiba two drachms  
Gum Arabic one ounce  
Powdered liquorice sufficient  
Mix, and form mass. Dose, a scruple to two scruples.

*Radius.*

R. Copaiba one ounce  
Calcined magnesia one drachm  
Opium four grains  
Powdered cubebs three drachms  
Syrup of poppies sufficient  
Mix, and divide into boluses of half a drachm. Two to be taken at night, in gonorrhœa.

*Henschel.*

#### PILLS OF COPAIBA.

R. Copaiba two ounces  
Magnesia (freshly prepared) one drachm  
Mix, and set aside, till it concretes into a pilular mass; this to be divided into two hundred pills.

*U. S. Ph.*

Dose, two to six, two or three times a-day.

R. Copaiba  
Powdered cubebs, each, one drachm  
Mucilage of gum Arabic sufficient  
Mix, and divide into three-grain pills. Five to ten a-day.

*Hildenbrand.*

R. Copaiba one ounce  
Dragon's blood half an ounce  
Calomel one drachm  
Conserve of roses four ounces  
Mix, and make pills of six grains. Dose, from three to five a-day.

*Foy.*

#### PILLS OF COPAIBA AND CUBEBS.

R. White wax two drachms  
Melt, by a gentle heat, and add



Copaiba half an ounce  
Powdered cubebs one ounce  
Incorporate well, and divide into two hundred and forty pills. *Berens.*

PILLS OF COPAIBA, CUBEBS, AND TURPENTINE.

R. Oil of copaiba  
cubebs  
turpentine, each, one fl. drachm  
Magnesia two drachms  
Mix, and form sixty pills.

This is said to be an excellent preparation. It was communicated by Mr. Procter, who is unacquainted with the author of it.

COPAIBA CONFECTION.

R. Turpentine one ounce  
Copaiba half an ounce  
Mix well, in a warmed mortar, and add  
Mucilage of gum Arabic one ounce  
Conserve of roses four ounces  
Dose, two to three drachms, three times a-day, in obstinate gonorrhœa. *Swedjaur.*

R. Copaiba  
Powdered cubebs, each, four drachms and a-half  
Yolk of egg one  
Conserve of roses half an ounce  
Mix. A teaspoonful, three or four times a-day. *Voght.*

R. Copaiba  
Powdered cubebs, each, two ounces  
alum one ounce  
Opium five grains  
Incorporate well. One drachm to be taken in the pulp of a prune, night and morning, and rapidly increased to two drachms. Is very effectual in gonorrhœa. *Berton.*

EMULSION OF COPAIBA.

R. Copaiba  
Mucilage gum Arabic, each, two ounces  
Water, twelve fl. ounces  
Rub the copaiba gradually with the mucilage in a mortar, add the water by degrees, constantly rubbing. *Beral.*

INJECTION OF COPAIBA.

R. Copaiba two drachms  
Mucilage of gum Arabic half an ounce  
Lime water six fl. ounces  
Mix well. As an injection in ulcers of the rectum, vagina, or urethra. *Abernethy.*

R. Copaiba half an ounce  
Yolk of egg sufficient  
Lime water six ounces

Honey of roses three ounces  
Mix. As an injection in fistulous ulcers. *Plenck.*

COPAIBA MIXTURE.

R. Copaiba  
Sweet spirits of nitre, each, half a fl. ounce  
Powdered gum Arabic  
sugar, each, one drachm  
Compound spirits of  
lavender two fl. drachms  
Tincture of opium one fl. drachm  
Distilled water four fl. ounces

Mix. A tablespoonful three times a-day in gonorrhœa. *Chapman.*

R. Copaiba  
Powdered cubebs, each, half an ounce  
Gum Arabic two drachms  
Paregoric elixir half a fl. ounce  
Sugar two drachms  
Water eight fl. ounces

Mix. A tablespoonful every three hours. *Ellis.*

R. Copaiba half an ounce  
Mint water  
Alcohol  
Syrup, each, half a fl. ounce  
Sweet spirit of nitre half a fl. drachm

Mix. To be taken in four doses in a day, followed by demulcent drinks. In chronic gonorrhœa. *Ellis.*

R. Balsam copaiba  
Sweet spirit of nitre, each, half a fl. ounce  
Compound spirit of lavender  
Laudanum, each, one fl. drachm  
Powdered gum Arabic one drachm  
Water three fl. ounces

Mix. A tablespoonful three times a-day in gonorrhœa. *Chapman.*

R. Copaiba half an ounce  
Mucilage of gum Arabic two ounces  
Clarified honey six drachms  
Solution of potassa two drachms  
Essence of lemon half a drachm  
Rose water five fl. ounces

Mix. Two to three spoonfuls, three times a-day. *Stephenson.*

R. Copaiba one drachm and a-half  
Powdered kino one scruple  
Mucilage of gum Arabic two drachms  
Syrup of rhatany one ounce  
Decoction of chamomile four ounces

Mix. A spoonful every hour or two.

*Foy.*

R. Copaiba half a drachm  
Tincture of allspice



Syrup of mallows, each, two drachms  
Mucilage of gum Arabic sufficient  
Water one ounce  
Mix. To be taken morning and evening.  
*Saunders.*

R. Copaiba  
Syrup of lemon  
Mint water  
Orange-flower water, each, one ounce  
Sulphuric acid one drachm  
Tragacanth sufficient  
Mix. A spoonful, morning and evening.  
*Delpech.*

R. Copaiba  
Tolu  
Powdered gum Arabic, each,  
half an ounce  
Elixir of vitriol twenty drops  
Distilled water six fl. ounces  
Mix. A tablespoonful, in chronic hoop-  
ing-cough. *Ellis.*

R. Copaiba one drachm  
Carbonate of potassa thirty grains  
Resin of Peruvian bark twenty grains  
Gum Arabic twelve grains  
Aloes two grains  
Syrup of capillaire two ounces  
Cinnamon water  
Mint water, each, three fl. ounces  
Mix. Dose, a glass full, several times a-  
day, as an anthelmintic. *Pierquin.*

## TINCTURE OF COPAIBA.

R. Copaiba one part  
Alcohol eight parts  
Digest for several days, and filter. Dose,  
thirty to sixty drops. *Guibourt.*

## ALKALINE TINCTURE OF COPAIBA.

R. Copaiba one ounce  
Carbonate of potassa one drachm  
Alcohol four fl. ounces  
Digest, and filter. Dose, twenty-five to  
fifty drops. *Brunswick Ph.*

## COMPOUND TINCTURE OF COPAIBA.

R. Copaiba one ounce  
Guaiacum two drachms  
Oil of sassafras half a drachm  
Alcohol four and a-half fl. ounces  
Digest with a gentle heat, filter. *Foy.*  
Dose, a drachm, in some infusion.

R. Copaiba one ounce and a-half  
Balsam Peru half an ounce  
Carbonate of  
potassa one drachm and a-half  
Cut saffron two drachms  
Alcohol sixteen fl. ounces  
Mix, and digest for three days, and filter.

Two spoonfuls, two or three times a-day,  
in water or in wine. *Wirttemberg Ph.*

## OIL OF COPAIBA.

R. Copaiba one ounce  
Water one pint and a-half  
Distil, return product several times, and  
redistil; separate oil. *Ed. Ph.*  
Dose, ten to thirty drops, rubbed up with mu-  
cilage and water.

## COPAIBA CLYSTER.

R. Copaiba one to four drachms  
Extract of opium one grain  
Yolk of egg one  
Decoction of mal-  
lows four to six fl. ounces  
Make an emulsion, as a clyster in gonorr-  
hoea. *Guibourt.*

## COPAIBA PASTE.

R. Sweet almonds six drachms  
Mallow paste one drachm  
Catechu half a drachm  
Copaiba three drachms  
Rub together into a paste. Dose, a spoon-  
ful. *Phœbus.*

## COPTIS.

## GOLDTHREAD.

The root of *Coptis trifolia*, a small evergreen  
plant, found in the more northern parts of both  
continents, in wet and boggy situations.

*Sex. Syst.* Polyand. polyg. *Nat. Syst.* Ranun-  
culaceæ.

Salisbury. Linn. Trans. viii. 305. Griffith,  
Med. Bot. 87.

The roots, which are the part used, are long,  
slender, orange-yellow, with no smell, but a  
strong, and purely bitter taste. It is a simple  
tonic bitter, with no astringency. It is used as a  
stomachic, and as a local application to aphthous  
sore mouth. Dose, in powder, ten to thirty  
grains.

## INFUSION OF GOLDTHREAD.

R. Goldthread one ounce  
Boiling water one pint  
Infuse for an hour, and strain. Dose,  
half fl. ounce to two fl. ounces.

*Dunghison.*

## TINCTURE OF GOLDTHREAD.

R. Goldthread one ounce  
Diluted alcohol one pint  
Macerate for a week, and filter. Dose,  
one fl. drachm to three fl. drachms.

*Wood.*

## CORIANDRUM.

## CORIANDER.

This is the fruit of *Coriandrum sativum*, an  
annual plant, a native of the south of Europe,



but naturalized in many other parts of that quarter of the world, and also extensively cultivated.

*Sex. Syst.* Pentand. digyn. *Nat. Syst.* Apiaceæ.

Linn. Sp. Pl. 367. Griffith, Med. Bot. 341.

The fruit is somewhat globular, of a grayish, or brownish-yellow color, and often separated into two portions, or mericarps. It has a pleasant aromatic smell and taste. It is principally used as an adjuvant to other articles, to disguise their taste, or to modify their griping qualities. Dose, from a scruple to a drachm.

#### COMPOUND POWDER OF CORIANDER.

R. Powdered coriander one drachm and a-half  
Ivory filings  
Burnt hartshorn  
Prepared chalk, each, one scruple  
Powdered cinnamon half a scruple  
Sugar of roses sufficient  
Mix well. A teaspoonful, after eating, as a stomachic. *Pierquin.*

R. Powdered coriander  
rhubarb  
columbo, each, ten grains  
Mix. To be taken at bedtime, in dyspepsia with flatulence and costiveness. Stomachic. *Ainslie.*

#### TINCTURE OF CORIANDER.

R. Coriander one part  
Diluted alcohol eight parts  
Macerate for a week, and filter. *Beral.*

#### COMPOUND WATER OF CORIANDER.

R. Coriander eight ounces  
Citron water one fl. ounce  
Nutmeg  
Storax  
Benzoin, each, half an ounce  
Vanilla three drachms  
Alcohol forty-eight fl. ounces  
Macerate for two days, and distil. Dose, half fl. ounce, as a stomachic and carminative. *Spielmann.*

### CORNU.

#### HARTSHORN.

This is officially the horn of the *Cervus elaphas*, or European stag; but that of our native species has the same properties. It is found in the shops, in the form of shavings, which are of a yellowish white color, friable, and without smell or taste. They consist principally of gelatine and phosphate of lime, and are employed to make a jelly, which is not superior to any other animal jelly.

#### HARTSHORN JELLY.

R. Hartshorn four ounces  
Water eight ounces  
Muriatic acid one drachm

Beat together for ten minutes, wash in two or three waters, boil with fresh water for half an hour, express, strain, and add

Sugar four ounces

Boil down sufficiently for the mixture to jelly, on cooling. *Ferrez.*

#### COMPOUND JELLY OF HARTSHORN.

R. Hartshorn jelly eight ounces  
Paste of sweet almonds one ounce  
Sugar half an ounce  
Orange-flower water one drachm  
Essence of citron twelve drops  
Mix, and heat; then permit to cool.

*Guibourt.*

#### BURNT HARTSHORN.

R. Burn pieces of hartshorn in an open vessel till quite white; powder, and prepare like prepared chalk. *Lond. Ph.*

Dose, half a drachm to two drachms.

#### DENTIFRICE OF BURNT HARTSHORN.

R. Powdered burnt hartshorn one ounce  
Orris root two drachms  
Dragon's blood one drachm  
Oil of roses two drops  
Mix. *Steph. & Church.*

### CORNUS.

#### DOGWOOD.

### CORNUS CIRCINATA.

#### ROUND-LEAVED DOGWOOD.

A large shrub, native of the United States, with a bitter, astringent, aromatic bark.

### CORNUS FLORIDA.

#### TREE DOGWOOD.

A small tree found in most parts of the United States, flowering early in the spring. The bark, especially of the root, has a bitter, astringent, somewhat aromatic taste, and a faint odor.

### CORNUS SERICEA.

#### SWAMP DOGWOOD.

A large shrub, like the others a native of this country, with a bark of the same qualities as them.

*Sex. Syst.* Tetrand. monog. *Nat. Syst.* Cornaceæ.

The barks of these shrubs and trees are tonic and astringent, and have much the same properties as Peruvian bark, and may be used as a substitute for it. That of *C. Florida* is said to contain a peculiar bitter principle, which has been



called *cornine*. The dose of the powder is from a scruple to a drachm.

#### DECOCTION OF DOGWOOD.

R. Dogwood bark, bruised, one ounce  
Water one pint  
Boil for half an hour, and strain. A wine-glassful every hour, in the apyrexia of intermittents. *U. S. Ph.*

R. Dogwood bark  
Sassafras, each, six ounces  
Water eight pints  
Boil, and strain. Given as above. *Niemann.*

#### PILLS OF ROUND-LEAVED DOGWOOD.

R. Extract of round-leaved dogwood one drachm and a-half  
Powdered ginger ten grains  
Dried carbonate of soda ten grains  
Mix, and form twenty-four pills. *Reece.*

#### WINE OF ROUND-LEAVED DOGWOOD.

R. Extract of round-leaved dogwood three drachms  
White wine one pint and a-half  
Digest for fourteen days, and filter. *Reece.*

### COTULA.

#### MAYWEED.

A small annual syngenesious plant, a native of Europe, and extensively naturalized in the United States. It is the *Maruta cotula* of botanists.

*Sex. Syst. Syng. frust. Nat. Syst. Asteraceæ.*

*Decandolle, Prod. vi. 13. Griffith, Med. Bot. 399.*

The whole herb has a strong, unpleasant smell, and a bitter, warm taste. It has much the same properties as chamomile as an internal remedy, and may be used as a substitute for it; but its disagreeable odor renders it nauseous to most persons. It is also employed in amenorrhœa, and as an antispasmodic. It is usually given in infusion.

#### INFUSION OF MAYWEED.

R. Mayweed half an ounce  
Boiling water one pint  
Macerate for a quarter of an hour, in a covered vessel, and strain. Used in same cases, and in same manner as infusion of chamomile.

#### CATAPLASM OF MAYWEED.

R. Mayweed, fresh, at will  
Bruise, or cut very fine. When applied as a cataplasm, it produces redness and vesication as rapidly as Spanish flies. *Ashley.*

### CREASOTUM.

#### CREASOTE.

A peculiar substance, analogous to the volatile oils, existing in the products of the distillation of wood. It is a colorless, oleaginous fluid, of a burning, caustic taste, and of an odor resembling that of smoked meat. It is corrosive, but, in a diluted state, is an effectual preserver of animal substances. It is irritant, narcotic, styptic, &c., and has been employed in a variety of diseases, both internally and as a local application.

#### CREASOTE.

R. Tar at will  
Distil, changing the recipients several times, till the residue has the consistence of pitch; agitate the heavy oil that passes with a little concentrated sulphuric acid; then mix it with its own volume of water, and rectify in small retorts; dissolve the product which sinks to the bottom in a hot solution of potassa, aided by a gentle heat; when it cools, add a slight excess of sulphuric acid, separate the oil, and distil it in small quantities at a time; again rectify it two or three times. Dose, one to two drops, mixed with water.

#### CREASOTE MIXTURE.

R. Creasote six drops  
Powdered mallows  
Sugar, each, one drachm  
Mix well, and divide into three powders, to be taken during the day. In hemoptysis and phthisis. *Radius.*

#### CREASOTE PILLS.

R. Creasote one drachm  
Powdered mallow root sufficient  
to make one hundred and ten pills. Said to be beneficial in acute rheumatism. Four or five to be taken, morning and evening. *Riech.*

R. Creasote one drachm  
Extract of liquorice, each, one drachm and a-half  
Water sufficient  
Beat into mass, and make pills of two grains. Two, morning and evening, in phthisis and laryngitis. *Radius.*

R. Creasote ten drops  
Powdered liquorice  
Mucilage of gum Arabic, each, sufficient  
Mix, and form twenty pills. Dose, one, three times a-day. In bronchitis, neuralgia, &c. *Ellis.*

R. Creasote one drachm  
Extract of liquorice



Galbanum, each, half a drachm  
Powdered mallows two drachms  
Form mass, and divide into one hundred  
and twenty pills. Six, four times a-day, in  
phthisis. *Riech.*

## ALCOHOLIC SOLUTION OF CREASOTE.

R. Creasote one part  
Alcohol sixteen parts  
Mix. A drop or two placed in the cavity  
of an aching tooth, will afford relief.

*Marinus.*

R. Creasote one fl. drachm  
Alcohol two fl. ounces  
Mix. Used with water as a mouth wash,  
to correct fetor of the mouth, and to check  
caries of the teeth. *Buchner.*

R. Creasote one fl. drachm  
Alcohol one fl. ounce and a-half  
Tincture of cochineal six fl. drachms  
Oil of mint thirty-six drops  
Mix. Used as above. *Ringhini.*

## CREASOTE LOTION.

R. Creasote half a drachm  
Water five ounces  
Mix. As a lotion in psora, &c. *Radius.*

R. Creasote ten drops  
Vinegar two fl. drachms  
Water two fl. ounces  
Mix. As an application to phagedenic  
ulcerations and chancres. *Short.*

## CREASOTE COLLUTORY.

R. Creasote three drops  
Compound spirit of  
lavender twenty drops  
Distilled water half fl. ounce  
Mix. *Black.*

R. Creasote four drops  
Distilled water two fl. ounces  
Mix. In irritation of the gums.

*Fremanger.*

R. Creasote half a fl. drachm  
Powdered gum  
Arabic one ounce and a-half  
Camphor  
water two fl. ounces and a-half  
Mix. To be used every two hours in  
aphthous ulceration of the mouth.

*Magendie.*

## CREASOTE MIXTURE.

R. Creasote  
Acetic acid, each, sixteen drops  
Compound spirit of juniper  
Syrup each one fl. ounce  
Water fourteen fl. ounces  
Mix the creasote with the acid, gradually

add the water, and then the spirit and  
syrup. *Ed. Ph.*

Dose, one fl. ounce.

R. Creasote two to four drops  
Mucilage of gum Arabic one ounce  
Infusion of salep five ounces  
Sugar one drachm  
Mix. A spoonful every two hours in  
the colliquative diarrhoea of phthisis.

*Radius.*

R. Creasote five drops  
Mucilage of gum Arabic three fl. ounces  
Syrup of mallows one fl. drachm  
Mix. A spoonful every three hours in  
hemoptysis. *Santoni.*

R. Creasote one drop  
Camphor water  
Compound infusion of  
gentian, each, six fl. drachms.  
Mix. To check vomiting. *Santoni.*

## CREASOTE OINTMENT.

R. Creasote half a fl. drachm  
Lard one ounce  
Melt the lard, add the creasote, and stir  
till cold. *U. S. Ph.*

As an application in scaly, cutaneous eruptions, ulcers, &c.

R. Simple cerate  
Oil of almonds, each, one ounce.  
Creasote thirty drops  
Mix. As an application in scrofulous  
caries. *Fremanger.*

## COMPOUND OINTMENT OF CREASOTE.

R. Creasote  
Solution of subacetate of  
lead, each, ten drops  
Extract of opium one grain and a-half  
Lard one ounce  
Mix. As an application to chilblains.

*Devergie.*

R. Creasote half a drachm  
Purified animal charcoal one drachm  
Alcohol one fl. drachm and a-half  
Spermaceti  
ointment, one ounce and a-half  
Mix. Used as an application to burns,  
to be diluted, if necessary. *Sutro.*

## CROCUS.

## SAFFRON.

Saffron is the stigmas of *Crocus sativus*, a bulbous rooted plant, native of Greece, Asia Minor, &c., and extensively cultivated in many parts of Europe.

*Sax. Syst. Triand. monog. Nat. Syst. Iridaceæ.*



Saffron has a peculiar, sweetish, fragrant odor, a warm, bitter taste, and is of an orange red color. It was at one time much used as an excitant, narcotic aromatic and emmenagogue, but is seldom employed in this country, except as a coloring ingredient in compound preparations. Dose, from ten to thirty grains.

## PILLS OF SAFFRON.

R. Saffron  
Myrrh  
Sulphur, each, one drachm  
Inspissated bile sufficient  
Mix, and form one hundred and fifty pills.  
Dose, twelve a-day, as an emmenagogue.  
*Phæbus.*

## TINCTURE OF SAFFRON.

R. Saffron, cut fine two ounces  
Proof spirit two pints  
Digest for fourteen days, express, and filter.  
*Ed. Ph.*  
Dose, one to two fl. drachms.

## COMPOUND TINCTURE OF SAFFRON.

R. Saffron one ounce  
Myrrh two ounces  
Socotrine aloes ten ounces  
Cinnamon  
Mace  
Nutmeg, each, half an ounce  
Orange-flower water one pint  
Alcohol sixteen pints  
Digest for two days, distil off eight pints.  
*Cottureau.*

## ELIXIR DE GARUS.

R. Compound tincture of saffron eight pints  
Syrup of maidenhair ten pints  
Mix, color with caramel, dissolved in Orange-flower water eight fl. ounces  
Used as a stomachic and carminative.  
*Foy.*

## INFUSION OF SAFFRON.

R. Saffron two drachms  
Boiling water two pints  
Infuse, and strain. Advised in debility of the digestive organs.  
*Beral.*

## SYRUP OF SAFFRON.

R. Saffron ten drachms  
Boiling water one pint  
White sugar three pounds  
Macerate the saffron in the water for twelve hours, strain, and add the sugar.  
*Lond. Ph.*

Slightly stimulant.

## SAFFRON COLLYRIUM.

R. Saffron one part  
Boiling water two hundred parts  
Infuse, strain, and add

Tincture of opium one part  
*Phæbus.*

## SAFFRON OINTMENT.

R. Cut saffron one scruple  
Camphor two scruples  
Oil of henbane two drachms  
Goulard's cerate six drachms  
Mix. As an application to painful hemorrhoids.  
*Speilmann.*

## CUBEBA.

## CUBEBS.

Cubebs are the berries or fruit of *Piper cubeba*, a climbing perennial vine, native of many parts of the East Indies.

*Ser. Syst.* Diand. trigyn. *Nat. Syst.* Piperaceæ.

*Linn. Sp. Pl.* 90. *Griffith, Med. Bot.* 567.

Cubebs somewhat resemble black pepper, but are of a lighter color, and are furnished with a short stalk. The stalk is thin and hard, and the seed round, white, and oily. The taste is acrid, peppery, and camphoraceous; the odor aromatic and pleasant. Cubebs are stimulant, with a special action on the genito-urinary organs, and have been much used in gonorrhœa, leucorrhœa, &c., and have also been found useful in piles, bronchitis, &c. The dose of the powder is from ten grains to three drachms, according to circumstances.

## POWDER OF CUBEBS.

R. Powdered cubebs half a drachm to one drachm  
ipecacuanha eighth of a grain  
Oil of peppermint half a drop  
Mix. To be taken four times a-day, in gonorrhœa.  
*Schmidt.*

## POWDER OF CUBEBS AND ERGOT.

R. Powdered ergot two scruples  
cubebs one ounce  
cinnamon half a drachm  
sugar one drachm  
Mix, and divide into eight powders. One to be given three or four times a-day. In gonorrhœa, gleet, leucorrhœa, &c. *Ryan.*

## POWDER OF CUBEBS AND HEMLOCK.

R. Powdered cubebs half a drachm to a drachm  
Extract of hemlock half a grain  
Prepared oyster shell five grains  
Mix. To be taken four times a-day, in gonorrhœa.  
*Phæbus.*

## POWDER OF CUBEBS AND ALUM.

R. Powdered cubebs two ounces  
alum half an ounce  
Mix, and divide into nine powders. One to be taken three times a-day. *Ricord.*



## CUBEB LOZENGES.

R. Powdered cubebs two drachms  
Balsam of tolu six grains

Mix, and add

Extract of liquorice one ounce  
Syrup of balsam of Peru one drachm  
Gum Arabic sufficient

Rub well together, and form lozenges of ten grains. One of these, permitted gradually to melt in the mouth, alleviates the obstruction in the nose, in coryza. *Spitta.*

## ELECTUARY OF CUBEBS.

R. Powdered cubebs half an ounce  
Clarified honey sufficient

Mix. Three or four teaspoonfuls a-day, in gonorrhœa, mucous discharges from the bladder, &c. *Radius.*

R. Powdered cubebs  
Balsam copaiba, each, two ounces  
Powdered alum one ounce  
Extract of opium five grains

Rub together. Dose, one drachm, night and morning, in the pulp of a prune, rapidly increasing the dose to two drachms. In gonorrhœa. *Beral.*

## INJECTION OF CUBEBS.

R. Powdered cubebs one ounce  
Water one pint

Boil, and strain. Add

Extract of belladonna one scruple

In gonorrhœa and leucorrhœa. *Soubéiran.*

## CLYSTER OF CUBEBS.

R. Powdered cubebs one to four drachms  
Decoction of mallows six ounces

Mix. To be administered, night and morning, in gonorrhœa. *Foy.*

## ETHEREAL EXTRACT OF CUBEBS.

R. Powdered cubebs one pound  
Sulphuric ether sufficient

Introduce the powder into a displacer, insert the lower end into a closely fitting bottle, add the ether carefully, cover the top of the filter with wet bladder pierced with several pin-holes. The flow should be very gradual. Introduce the tincture into a large retort, and distil, by means of a water-bath, into a well-cooled receiver, till five-sixths have passed over. Continue the evaporation at a heat below 120° F., till the extract is of proper consistence. *W. Procter.*

## EMULSION OF ETHEREAL EXTRACT OF CUBEBS.

R. Ethereal extract of cubebs two drachms  
Powdered gum Arabic half an ounce  
sugar one drachm

Water three ounces and a-half  
Mix. A tablespoonful is equal to two drachms of cubebs. *W. Procter.*

## ALCOHOLIC ETHEREAL EXTRACT OF CUBEBS.

R. Powdered cubebs at will  
Exhaust by ether, by means of a displacement apparatus, act on residue with diluted alcohol, distil the two tinctures separately. Evaporate the residue of the alcoholic tincture by means of a water-bath; add both together, and let the remaining ether evaporate spontaneously. *Labelonye.*

## LOZENGES OF ALCOHOLIC ETHEREAL EXTRACT OF CUBEBS.

R. Extract of cubebs eight ounces  
Alcohol two pints

Dissolve, and add

Powdered sugar one pound  
Oil of peppermint eighteen drops

Mix, and allow alcohol to evaporate by a gentle heat; reduce to powder, and add

Mucilage tragacanth sufficient

Mix, and divide into lozenges of six to eighteen grains. *Labelonye.*

## CUBEB MIXTURE.

R. Powdered cubebs two drachms  
Carbonate of soda half a drachm

Mucilage of gum Arabic six fl. drachms

Mint water six fl. ounces

Mix. A tablespoonful every hour.

*Fosbroke.*

R. Powdered cubebs two drachms  
Subnitrate of Bismuth half a drachm

Mucilage of gum Arabic half fl. ounce

Syrup six fl. drachms

Water six fl. ounces

Mix. A tablespoonful four times a-day.

*Fosbroke.*

Both these are recommended in the various affections of mucous membranes.

R. Powdered cubebs two drachms  
Wine two ounces

Essence of bergamot one drop

Mix. To be taken every hour or two.

*Pierquin.*

## TINCTURE OF CUBEBS.

R. Bruised cubebs four ounces  
Diluted alcohol two pints

Macerate for fourteen days, express, and filter. *U. S. Ph.*

Dose, one to two drachms, as a stomachic and carminative, or in advanced stage of gonorrhœa.

## SYRUP OF ALCOHOLIC ETHEREAL EXTRACT OF CUBEBS.

R. Extract of cubebs three ounces



Mucilage of gum Arabic      sufficient  
Peppermint water      one pint  
White sugar      two pounds  
Mix. A teaspoonful, several times a-day,  
in a glass of water. *Labelonye.*

## OIL OF CUBEBS.

R. Powdered cubebs      one part  
Water      two parts  
Distil, and collect the oil. *Giordano.*  
Dose, ten to twelve drops, gradually increased.  
Given in emulsion, or in gelatin capsules.

MIXTURE OF OIL OF CUBEBS AND  
COPAIVA.

R. Oil of cubebs      one scruple  
Copaiba      two ounces  
Mix. A teaspoonful, four times a-day.  
*Beral.*

## CUNILA.

## DITTANY.

The whole herb of *Cunila mariana*. A native plant, found in most places in the United States.  
*Sax. Syst. Diand. monog. Nat. Syst. Lamiaceæ*  
*Linn. Sp. Pl. 30. Griffith, Med. Bot. 509.*

It has a powerful, aromatic odor, and a warm, pungent taste. It is stimulant, carminative, sudorific, and emmenagogue.

## INFUSION OF DITTANY.

R. Dittany      half an ounce  
Boiling water      one pint  
Infuse and strain. Used warm, as a diaphoretic and emmenagogue.

## OIL OF DITTANY.

R. Dittany      at will  
Water      sufficient  
Distil, and separate oil. As a carminative, in doses of two or three drops, with sugar.

## CUPRUM.

## COPPER.

A brilliant ductile, malleable metal, of a reddish color, having an unpleasant taste and smell. It is not used in its pure state in medicine, but affords numerous remedial preparations.

## CUPRI ACETAS.

## ACETATE OF COPPER.

R. Pulverized verdigris      at will  
Acetic acid      sufficient  
Dissolve, filter, evaporate, and crystallize.  
*Van Mons.*

## CUPRI SUBACETAS.

## VERDIGRIS.

## PREPARED SUBACETATE OF COPPER.

R. Verdigris, in powder,      at will  
Prepared as directed for chalk. *Dub. Ph.*

## POWDER OF VERDIGRIS AND SAVINE.

R. Verdigris  
Savine      equal parts  
Mix. To be dusted on sores with fungous excrescences. *Ellis.*

## PILLS OF ACETATE OF COPPER.

R. Acetate of copper      fifteen grains  
Boiling water      sufficient  
to dissolve, add  
Opium      five grains  
Extract of liquorice      one drachm  
Powdered liquorice      sufficient  
Mix, and make one hundred and eighty pills. Dose, three to ten, three times a-day. *Phœbus.*

## VERDIGRIS AND ALUM.

R. Verdigris  
Nitre  
Alum, each,      eight parts  
Melt together, and add  
Camphor      one to two parts  
Much used in Europe as a basis for ophthalmic solutions, under the name of Beer's divine stone. *Radius.*

## OPHTHALMIC WASHES.

R. Beer's divine  
stone      eight to twelve grains  
Water      four fl. ounces and a-half  
Wine of opium      a scruple to one fl. drachm  
Mix. In chronic ophthalmia. *Benedict.*

R. Beer's divine stone three to four grains  
Elder water      one fl. ounce  
Tincture of opium      twelve drops  
Solution of acetate of lead      five drops  
Mix. Used like the last. *Rust.*

## COMPOUND VERDIGRIS LOTION.

R. Verdigris  
Burnt alum, each,      two drachms  
Honey      half an ounce  
White wine      one pint  
Mix. As a wash for indolent venereal or scorbutic ulcers. *Scherf.*

## METZ'S BALSAM.

R. Linseed oil  
Olive oil, each,      six ounces  
Oil of laurel berries      one ounce



Turpentine two ounces  
Melt by a gentle heat, and add  
Powdered aloes two drachms  
verdigris three drachms  
white vitriol one drachm and a-half  
Pour into a bottle, and add  
Oil of juniper half an ounce  
cloves one drachm  
Mix. As a dressing to wounds and ulcers.  
*Guibourt.*

## VERDIGRIS OINTMENT.

R. Verdigris, in fine powder, one drachm  
Simple ointment fifteen drachms  
Melt the ointment, and add the verdigris,  
and stir till cold. *U. S. Ph.*

Used as a mild escharotic and stimulant to fungous ulcers, to warts, corns, &c., and obstinate cutaneous affections.

R. Verdigris two drachms  
Oxide of zinc  
Camphor, dissolved in  
alcohol, each, six drachms  
Triturate well, and incorporate with a  
melted mixture of  
Lard  
Suet, each, two ounces  
and stir till cold  
Recommended in scrofulous ophthalmia.  
*Swedjaur.*

## VERDIGRIS PLASTER.

R. Galbanum one ounce  
Pitch half an ounce  
Litharge plaster two drachms  
Melt, and add  
Verdigris  
Muriate of ammonia, each, one ounce  
This is a good application to corns and  
warts. *Saxon Ph.*

## EGYPTIAN OINTMENT.

R. Verdigris five parts  
Purified honey sixteen parts  
Strong vinegar seven parts  
Burnt alum half a part  
Mix, and melt by a gentle heat, stirring  
constantly. This is thought to form an  
excellent detergent application to ulcers.  
*Giordano.*

## LINIMENT OF VERDIGRIS.

R. Powdered verdigris one ounce  
Vinegar seven fl. ounces  
Honey fourteen ounces  
Dissolve the verdigris in the vinegar,  
strain, gradually add the honey, and boil  
down to proper consistence. *Lond Ph.*  
Used like the last, and also much diluted as a  
gargle in venereal sore throat.

## CUPRI MURIAS.

## MURIATE OF COPPER.

R. Chloride of potassium seven parts  
Sulphate of  
copper eleven and a-half parts  
Pulverize, mix, and gradually add twelve  
parts boiling water; on cooling, filter,  
separate the sulphate of potassa that  
forms, then permit the crystals of the  
muriate to form, separate, and dry.  
*Van Mons.*

## CUPREOUS ETHER.

R. Chloride of barium ten parts  
Sulphate of copper twelve parts  
Sulphuric ether six parts  
Triturate the salts together, add the ether,  
and decant when all the muriate of cop-  
per is dissolved. Has been recommended  
in small doses in epilepsy. *Van Mons.*

## MURIATE OF COPPER AND AMMONIA.

R. Muriate of copper  
ammonia equal parts  
Dissolve in water, add liquid ammonia,  
drop by drop, as long as any precipitate  
takes place, and is again dissolved by the  
addition of more ammoniac, filter, and  
evaporate. *Augustin.*

Used in epilepsy, in doses of two to ten grains.

SOLUTION OF AMMONIACAL MURIATE OF  
COPPER AND MERCURY.

R. Copper filings two drachms  
Calomel two ounces  
Dissolve the copper in two fl. ounces liquid  
ammonia, and the calomel in two fl. ounces  
muriatic acid, with fifteen drops of ni-  
tric acid; mix the solutions in such pro-  
portions that the precipitate that first  
forms is redissolved. *Radius.*

## KOECHLIN'S DROPS.

R. Solution of ammoniacal  
muriate of copper and  
mercury two fl. drachms  
Distilled water twenty fl. ounces  
Mix. Much praised in obstinate venereal  
affections, scrofula, &c. Dose, a teaspoon-  
ful after each meal. *Augustin.*

## CUPRUM AMMONIATUM.

## AMMONIATED COPPER.

R. Sulphate of copper half an ounce  
Carbonate of ammonia six drachms  
Rub together in a glass mortar till efferves-  
cence ceases, wrap in bibulous paper, and  
dry with a gentle heat. *U. S. Ph.*  
Has been used in epilepsy, chorea, hysteria,



&c. Dose, a quarter to half a grain twice a-day, and gradually increased.

#### SOLUTION OF AMMONIATED COPPER.

R. Ammoniated copper           one drachm  
Distilled water                   one pint  
Dissolve, and filter.           *Lond. Ph.*

As a lotion to foul and indolent ulcers, or internally, in same cases as the powder.

#### INJECTION OF AMMONIATED COPPER.

R. Ammoniated copper           five grains  
Rose water                       eight fl. ounces  
Mix. In gonorrhœa.           *Ellis.*

#### POWDER OF AMMONIATED COPPER AND BELLADONNA.

R. Ammoniated copper two to four grains  
Powdered belladonna  
root                               one to four grains  
Sugar                             one drachm  
Mix, and form six powders, one to be taken every two hours, in epilepsy. *Hildebrand.*

#### PILLS OF AMMONIATED COPPER.

R. Ammoniated copper           two grains  
Crumb of bread               sufficient  
Mix, and form four pills. One to be taken twice a-day, in epilepsy, gradually increasing the dose. *A. T. Thomson.*

R. Ammoniated copper           fifteen grains  
Crumb of bread               two scruples  
Sugar                            one scruple  
Liquid ammonia               sufficient  
Mix, and form thirty pills. Dose, one, three times a-day, gradually increased. In epilepsy. *Van Mons.*

#### COMPOUND PILLS OF AMMONIATED COPPER.

R. Ammoniated copper  
Opium, each,                   ten grains  
Extract of dandelion  
Powdered mallows, each, two scruples  
Mix, and form fifty pills. Dose, five, twice a-day, in diabetes. *Radius.*

#### GARGLE OF AMMONIATED COPPER.

R. Ammoniated copper           eight grains  
Savine water                   six fl. ounces  
Mix. In chronic sore throat. *Kopp.*

#### ointment of AMMONIATED COPPER.

R. Solution of ammoniated  
copper                           one fl. drachm  
Simple cerate, melted,       one ounce  
Mix well. As a stimulant to indolent ulcers. *Swediaur.*

### CUPRI SULPHAS.

#### SULPHATE OF COPPER.

##### POWDER OF SULPHATE OF COPPER.

R. Sulphate of copper           twelve grains  
Sugar of milk                   four scruples  
Mix, and divide into four powders. Much praised in croup—given as follows: one is to be administered at once, another dissolved in three spoonfuls of water, one of which is to be taken every hour. *Radius.*

R. Sulphate of copper           four grains  
Sugar of milk                   eight scruples  
Mix, and divide into eight powders. Two a-day in obstinate diarrhœa. *Elliotson.*

#### COMPOUND POWDER OF SULPHATE OF COPPER.

R. Sulphate of copper  
zinc  
Alum  
Carbonate of lead  
Armenian bole                   equal parts  
Melt the three first substances together, pulverize, add the two other articles, and sift.

This is a powerful astringent, and has proved useful as a styptic. *Cottureau.*

#### PILLS OF SULPHATE OF COPPER.

R. Sulphate of copper           four grains  
Extract of cinchona           thirty-two grains  
Mix, make mass, and divide into sixteen or twenty pills. One to be taken three times a-day, in obstinate intermittents. *Ellis.*

R. Sulphate of copper  
Ipecacuanha, each,           one drachm  
Syrup                           sufficient  
Form mass, and divide into pills of five grains each. Dose, two to four, every two or three days, in the morning, before eating, in phthisis. *Foy.*

R. Sulphate of copper           six grains  
Powdered calamus           two drachms  
Extract of liquorice  
Water, each,                   sufficient  
Form mass, and divide into ninety-six pills. Four to seven, two or three times, in the apyrexia of obstinate intermittents. *Adair.*

#### PILLS OF SULPHATE OF COPPER AND OPIUM.

R. Sulphate of copper           two grains  
Opium                           four grains  
Conserve of roses           sufficient



Mix, and make sixteen pills. One, three times a-day, in obstinate intermittents.

*Chapman.*

#### COMPOUND PILLS OF SULPHATE OF COPPER.

R. Sulphate of copper two grains  
Alcohol. ext. cascarilla seventeen grains  
Opium two grains  
Syrup of ginger sufficient  
Mix, and make eight pills. One, three times a-day, in epilepsy.

*Ainslie.*

#### ELECTUARY WITH SULPHATE OF COPPER.

R. Sulphate of copper one scruple  
Opium one grain  
Armenian bole  
Catechu, each, one drachm and a-half  
Syrup sufficient  
Mix. In obstinate diarrhoea.

*Saunders.*

#### GARGLE WITH SULPHATE OF COPPER.

R. Sulphate of copper twenty grains  
Infusion of sage six fl. ounces  
Tincture of myrrh  
catechu  
kino, each, one drachm  
pimpinella five drachms

Honey six drachms

Mix well. In obstinate salivation.

*Kopp.*

#### COLLYRIUM OF SULPHATE OF COPPER.

R. Sulphate of copper  
Armenian bole, each, eight grains  
Camphor two grains  
Boiling water eight fl. ounces  
Rub well together, permit to settle, and strain. As an application in purulent ophthalmia of infants.

*Ware.*

R. Sulphate of copper six grains  
Camphor one drachm  
Boiling water eight fl. ounces

Rub the camphor with the water, strain, and add sulphate of copper. As a substitute for the above.

*Ellis.*

#### SULPHATE OF COPPER LOTION.

R. Sulphate of copper one ounce and a-half  
Alum five drachms and a-half  
Sage leaves five ounces  
Vinegar one pint  
Solution of muriate of ammonia two pints

Boil together for half an hour. Used as an application, in a tepid state, to swellings and contusions.

*Purmann.*

#### INJECTION OF SULPHATE OF COPPER.

R. Sulphate of copper six grains  
Distilled water six fl. ounces

Tincture of opium one fl. drachm  
Dissolve. As an injection in chronic gonorrhoea.

*Ellis.*

R. Sulphate of copper six grains  
Distilled water four fl. ounces

Dissolve, and add  
Solution subacetate of lead twenty drops

As a wash and injection in phymosis.

*Swedjaur.*

#### OINTMENT OF SULPHATE OF COPPER.

R. Sulphate of copper  
Calamine, each, five grains  
Camphor two grains  
Fresh butter two drachms

Triturate well together. A small portion to be applied to the edges of the eyelids, in the evening, in psorophthalmia.

*Rust.*

R. Sulphate of copper  
Verdigris, each, two drachms  
Alum half an ounce  
Corrosive sublimate two scruples  
Lard one ounce  
Burgundy pitch one pound

Melt the pitch and lard by a gentle heat, and add the other ingredients, stirring till cold. As a dressing to venereal and fungous ulcers.

*Augustin.*

R. Powdered sulphate of copper.  
catechu, each, four drachms  
alum nine drachms  
resin four ounces

Olive oil sufficient

Make ointment. As an application to indolent and ill-conditioned ulcers.

*Kerr.*

#### SULPHATE OF COPPER STYPTIC.

R. Sulphate of copper three grains  
Sulphuric acid twenty drops  
Water two ounces

Dissolve. In epistaxis. Twenty to forty drops to be taken in water, every hour.

*Thatcher.*

## CYDONIA.

### QUINCE SEEDS.

These seeds are the product of *Cydonia vulgaris*, a small tree, native of some parts of Europe, but extensively cultivated in this country. The fruit is much used for the purpose of making preserves.

*Sex. Syst.* Icosand. pentag. *Nat. Syst.* Pomaceæ.

Persoon, *Enchirid.* ii. 40. Griffith, *Med. Bot.* 291.

The seeds are inodorous, insipid, and abounding with mucilage, one drachm rendering six ounces of water, viscid. Used like the other bland mucilages.



## DECOCTION OF QUINCE SEEDS.

R. Quince seeds two drachms  
 Distilled water one pint  
 Boil over a slow fire, for ten minutes, and strain. *Lond. Ph.*

R. Quince seeds one ounce  
 Boiling water six fl. ounces  
 Digest on hot coals, for two hours; strain.  
 Both these are used in ophthalmia, &c.

## QUINCE SEED MIXTURE.

R. Mucilage of quince seeds one ounce  
 Yolk of eggs two  
 Honey of roses three ounces  
 Mix. A teaspoonful occasionally, in cough and hoarseness. *Sainte Marie.*

## BANDOLINE FOR THE HAIR.

R. Mucilage of quince seeds eight fl. ounces  
 Cologne water, or Brandy eight fl. ounces  
 Mix.

Used as an application to the hair, to give gloss and smoothness.

## CYMINUM.

## CUMIN.

Cumin seeds are produced by *Cuminum cyminum*, an umbelliferous plant, a native of Egypt, but cultivated in southern Europe.

*Sex. Syst.* Pentand. digyn. *Nat. Syst.* Apiaceæ.

*Linn. Sp. Pl.* 365. *Lindley, Fl. Med.* 51.

These seeds have a peculiar, strong, heavy odor, and a warm, bitterish, aromatic taste. They are carminative and stimulant, are seldom used internally, but enter into the composition of some warm plasters.

## CUMIN PLASTER.

R. Cumin  
 Caraway  
 Laural berries, each, three ounces  
 Burgundy pitch three pounds  
 Yellow wax three ounces  
 Melt the pitch and wax together, and add the other ingredients, stirring well.  
*Lond. Ph.*

## D

## DELPHINIUM.

## LARKSPUR.

The *Delphinium consolida* is a native of Europe, but has become partially naturalized in this country, and is also much cultivated in gardens as an ornamental flower.

*Sex. Syst.* Polyand. trigyn. *Nat. Syst.* Ranunculaceæ.

*Linn. Sp. Pl.* 748. *Griffith, Med. Bot.* 88.

Several parts of this plant have been employed medicinally, as the flowers, seeds, and roots. They owe their properties to the presence of an alkaloid, called delphinia. The flowers are said to be diuretic, vermifuge, and emmenagogue, and the seeds and root to be diuretic, &c.

## TINCTURE OF LARKSPUR SEEDS.

R. Larkspur seeds one ounce  
 Diluted alcohol one pint  
 Macerate for some days, and filter. Dose, ten to twenty drops, three times a-day, in spasmodic asthma. *Augustin.*

## DELPHINIA.

## DELPHINIA.

This alkaloid is prepared from the seeds of different species of *Delphinium*, but generally from those of the *D. staphisagria*.

R. Larkspur seeds at will  
 Water sufficient  
 Boil, repeat the decoction with another portion of water, till all soluble portions of the seeds are dissolved. Concentrate the united decoctions, add calcined magnesia, and filter after a short ebullition. Wash the precipitate with cold water, and dry it, digest it in alcohol on a water-bath, and permit to crystallize. *Magendie.*

This is used in the same cases as veratria, in doses of a quarter to half a grain, to the extent of two to three grains a-day. It is also employed in ointment or in solution in alcohol.

## SOLUTION OF DELPHINIA.

R. Delphinia one scruple  
 Rectified spirit two fl. ounces  
 Dissolve. For outward use. *Turnbull.*

## OINTMENT OF DELPHINIA.

R. Delphinia ten to thirty grains  
 Olive oil one drachm  
 Rub together, and add  
 Lard one ounce  
 Mix well. *Turnbull.*



## DIANTHUS.

## PINK.

The only species that is officinal is the *D. caryophyllus* or clove pink, a native of the south of Europe, but generally cultivated in gardens, for the beauty and fragrance of its flowers.

*Sex. Syst.* Decand. digyn. *Nat. Syst.* Caryophyllaceæ.

Linn. Sp. Pl. 587. Woodville, Med. Bot. t. 205.

The parts used are the flowers, these should be of a dark red color, and very aromatic. They are principally employed to form a syrup, used as a vehicle for other medicines.

## SYRUP OF CLOVE PINK.

R. Petals of clove pink           one pound  
Boiling water                   two pints  
Infuse for twelve hours, strain, and add  
twice the weight of sugar.      *Guibourt.*

## DIGITALIS.

## FOXGLOVE.

The foxglove is a native of Europe, but is cultivated in this country, both as an ornamental plant and for medicinal purposes.

*Sex. Syst.* Didynam. angios. *Nat. Syst.* Scrophulariaceæ.

Linn. Sp. Pl. 868. Griffith, Med. Bot. 520.

The parts used are the leaves; these should be dried in the dark, and always kept from the light. When good, they are of a dull green color, a feeble, narcotic odour, and a bitter, unpleasant taste. Digitalis is narcotic, sedative, and diuretic, and, in large doses, poisonous. It is given to fulfil many indications; as a diuretic, as a sedative, and antispasmodic, to reduce the force of the circulation, &c. It is given in substance in the dose of one grain, two or three times a-day, gradually increasing the quantity until some effect is produced on the system.

## POWDERS OF FOXGLOVE.

R. Powdered foxglove           one grain  
cinnamon                   four grains  
sugar                       ten grains  
Mix. For a single dose, to be repeated  
two or three times a-day.      *Radius.*

R. Foxglove           a quarter to one grain  
Sulphate of  
quinine           a-half to two grains  
Fennel seed           six grains  
Sugar of milk           ten grains  
Mix. To be taken three or four times a-  
day, in phthisis.      *Gunther.*

R. Powdered foxglove           ten grains  
calomel                   six grains  
valerian               two scruples  
assafetida  
castor, each,   half a drachm  
sugar               one drachm  
Mix, and divide into twenty-four powders.

One, morning and evening, in the convulsions attendant on hydrocephalus.

*Pierquin.*

R. Powdered foxglove           fifteen grains  
nitre                   three drachms  
cremor tartar   half an ounce

Mix, and divide into six powders. One to be taken every two hours. As a diuretic in dropsies.      *Brera.*

R. Powdered foxglove           six grains  
Golden sulphuret of antimony  
Calomel, each,           three grains  
Sugar                   one drachm

Mix, and divide into six powders. One, every three hours in peripneumonia.

*Radius.*

R. Powdered foxglove           eight grains  
Extract of opium           four grains  
Sugar                   one ounce  
Oil of peppermint           four drops

Mix, and divide into eight powders. One in the evening in palpitation of the heart.

*Sainte Marie.*

## EXTRACT OF FOXGLOVE.

R. Fresh juice of foxglove           at will  
Evaporate to proper consistence, without removing fecula.      *Lond. Ph.*

Dose, from half a grain to two grains.

## BOLUS OF FOXGLOVE.

R. Powdered foxglove           fifteen grains  
Calomel                   eight grains  
Root of juniper           one drachm

Mix, and form four boluses. One, every four hours in encephalitis.      *Brera.*

R. Powdered foxglove           twelve grains  
Calomel                   six grains  
Tartar emetic           two grains

Purified honey  
Powdered liquorice root, each, sufficient  
Mix, and form four boluses. One, every four hours, in hemoptysis and engorgement of the abdominal viscera.      *Brera.*

## PILLS OF FOXGLOVE.

R. Powdered foxglove           twelve grains  
Calomel                   six grains  
Opium                   two grains  
Conserve of roses           sufficient

Mix, and form twelve pills. One to be taken every eight hours, in hydrothorax and ascites.      *Ellis.*

R. Powdered foxglove           four grains  
camphor               twelve grains  
Extract of henbane           eighteen grains  
Mix, and make six pills. Two to be taken at bedtime. Used in maniacal and spasmodic affections.      *A. T. Thomson.*



R. Powdered squills four grains  
myrrh one scruple  
foxglove ten grains  
calomel six grains

Triturate together, and add

Assafetida half a drachm  
Extract of gentian sufficient

Beat into mass, and divide into fifteen pills.  
One night and morning, as a diuretic in dropsies. *Paris.*

R. Powdered foxglove  
opium, each, six grains  
Conserve of roses sufficient  
Mix, and make twelve pills. One to be taken every four hours. In asthma, &c. *Ellis.*

R. Powdered foxglove one drachm  
Extract of hemlock  
rhubarb  
Aloes, each, half a drachm  
Gum ammoniac four scruples  
Oxymel of squills  
Powdered liquorice root, each, sufficient  
Mix, and make pills of six grains. Three to be taken morning and evening. In dropsy, &c. *Brera.*

#### INFUSION OF FOXGLOVE.

R. Foxglove leaves one drachm  
Boiling water half a pint  
Tincture of cinnamon one fl. ounce  
Macerate the foxglove with the water, in a covered vessel, for four hours, strain, and add the tincture. *U. S. Ph.*

The dose is half a fl. ounce, twice a-day, till the system is affected.

#### MIXTURE OF FOXGLOVE AND ACETATE OF POTASSA.

R. Infusion of foxglove four fl. ounces  
Tincture of foxglove one fl. drachm  
Acetate of potassa one drachm  
Tincture of opium ten drops  
Mix. A dessertspoonful three or four times a-day; as a diuretic. *Ellis.*

#### MIXTURE OF FOXGLOVE AND ACETATE OF LEAD.

R. Powdered foxglove three to five grains  
Solution of subacetate of lead five drops  
Syrup of orange flowers one ounce  
Infusion of poppies six ounces  
Mix. In spoonful doses, in hypertrophy of the heart. *Foy.*

#### MIXTURE OF FOXGLOVE AND TARTARIC ACID.

R. Infusion of foxglove half an ounce  
Tartaric acid one scruple  
Carbonate of soda twenty-four grains

Sweet spirit of nitre one drachm  
Tincture of squills four drops  
Mint water two ounces  
Mix. As a diuretic, in ascites. To be taken twice or thrice a-day. *Dewees.*

#### EXPECTORANT MIXTURE OF FOXGLOVE.

R. Foxglove leaves thirty grains  
Boiling water sufficient  
to obtain four ounces of strained infusion, add

Syrup of gum Arabic three ounces  
Kermes mineral six grains  
Syrup of mallows one ounce  
Mix. To be taken, in teaspoonful doses, in pneumonia and pleurisy. *Brera.*

#### TINCTURE OF FOXGLOVE.

R. Foxglove leaves four ounces  
Diluted alcohol two pints  
Macerate for fourteen days, and filter. *U. S. Ph.*

Dose, from ten to twenty drops, two or three times a-day.

#### MIXTURE OF TINCTURE OF FOXGLOVE.

R. Tincture of foxglove one fl. drachm  
opium sixty drops  
Distilled water two fl. ounces  
Mix. A teaspoonful, two or three times a-day, in hemoptysis and incipient phthisis. *Ellis.*

#### ETHEREAL TINCTURE OF FOXGLOVE.

R. Foxglove leaves one part  
Sulphuric ether four parts  
Macerate for two days, and decant. Dose, from ten to twenty drops. *Soubeiran.*

#### VINEGAR OF FOXGLOVE.

R. Foxglove leaves half an ounce  
Vinegar four fl. ounces  
Infuse for twenty-four hours, strain, and add  
Sugar five ounces  
A teaspoonful, several times a-day. Highly spoken of in incipient phthisis. *Martius.*

#### OINTMENT OF FOXGLOVE.

R. Bruised foxglove leaves one part  
Lard two parts  
Melt over a slow fire, until all moisture is driven off, and strain. As an application to chronic ulcers. *Soubeiran.*

#### PLASTER OF FOXGLOVE.

R. Foxglove leaves two drachms  
Vinegar one fl. ounce  
Macerate, and evaporate to consistence of an extract, and add  
Calomel ten grains  
Elemi ointment eight scruples  
In chronic glandular swellings. *Sundelin.*



## SYRUP OF FOXGLOVE.

R. Foxglove leaves (European) four ounces  
 Water sufficient  
 Sugar one pound  
 Bruise the foxglove, add sufficient water to render quite moist, exhaust, by process of displacement, evaporate in a water-bath to ten ounces, add sugar, and form syrup. One teaspoonful is equivalent to fifteen grains of digitalis, or about two ounces of infusion. *Duhamel.*

## DIGITALINA.

## DIGITALIN.

R. Powdered foxglove at will  
 Macerate for twenty-four hours in a sufficiency of sulphuric ether, decant, repeat the operation several times, the last on a water-bath. Unite, and filter the tinctures, and distil off the most of the ether, over a water-bath. Treat the residue with distilled water. Add, very gradually, finely powdered litharge, till there is no acid reaction. Evaporate to dryness, treat with sulphuric ether, and evaporate to crystallizing point. *Magendie.*

Dose, from one-sixteenth to one-eighth of a grain.

## PILLS OF DIGITALIN.

R. Digitalin three quarters of a grain  
 Powdered gum Arabic  
 Mucilage of gum Arabic, each, sufficient  
 Mix, and make twenty pills. Dose, one to four daily, in hypertrophy of the heart. *Bouchardat.*

## GRANULES OF DIGITALIN.

R. Digitalin fifteen and a-half grains  
 Sugar one ounce and a-half  
 Water sufficient  
 Mix, and form one thousand granules. Dose, from four to six a-day. *Homolle.*

## DIOSMA.

## BUCHU.

Buchu consists of the leaves of several species of *Diosma*, or more properly *Barosma*, especially of *B. crenata* and *B. seratifolia*. They are small shrubs, natives of the Cape of Good Hope.

*Ser. Syst.* Pentand. monog. *Nat. Syst.* Rutaceæ.

*Linn. Sp. Pl.* (*Diosma*) 287. *Griffith, Med. Bot.* 191.

These leaves have a strong, somewhat aromatic odor, and a bitterish, aromatic taste. They are gently stimulant and diuretic, and are much used in diseases of the urinary organs, and also in some other complaints. The dose, in substance, is from twenty to thirty grains.

## INFUSION OF BUCHU.

R. Buchu leaves one ounce  
 Boiling water one pint  
 Macerate for four hours, in a covered vessel, and strain. *U. S. Ph.*

The dose is from one to two fluid ounces.

## COMPOUND INFUSION OF BUCHU.

R. Buchu leaves  
 Urva ursi, each, half an ounce  
 Boiling water eight ounces  
 Digest for half an hour, strain, and add  
 Syrup of senega half a fl. ounce  
 One or two spoonfuls every two hours, in atony of the bladder, and mucous discharges. *Radius.*

## TINCTURE OF BUCHU.

R. Buchu leaves two ounces  
 Proof spirit one pint  
 Macerate for seven days, and filter. *Dub. Ph.*

Dose, from one to four fl. drachms.

## EXTRACT OF BUCHU.

R. Buchu, in coarse powder, a pound  
 Ether four fl. ounces  
 Alcohol twelve fl. ounces  
 Displace without maceration, add diluted alcohol until a pint of ethero-alcoholic tincture is obtained; suffer this to evaporate spontaneously; treat the residue in the displacer, with diluted alcohol, till two pints are obtained; evaporate to a syrup; add the product of the first tincture, and with a gentle heat concentrate to the proper consistence. *W. Procter.*

## FLUID EXTRACT OF BUCHU.

R. Buchu leaves eight ounces  
 Alcohol sixteen fl. ounces  
 Water sufficient  
 Reduce the leaves to a coarse powder, moisten them in a covered vessel, with twelve fl. ounces of the alcohol, macerate for six hours, and introduce into a displacer. When the clear fluid has ceased to pass, add the remaining alcohol, mixed with four fl. ounces of water, until the displaced liquid amounts to twelve fl. ounces, which is to be set aside until reduced to six fl. ounces by spontaneous evaporation. The residue in the displacer is then to be treated with a pint of cold water by maceration for twelve hours, and subjected to pressure till a pint is obtained. Evaporate this to ten fl. ounces, mix with six fl. ounces of the tincture, agitate for several days, filter, or strain. Dose, one to two teaspoonfuls. *W. Procter.*



## DIOSPYROS.

## PERSIMMON.

The Persimmon, or *Diospyros Virginiana*, is a native tree, found in the middle and southern States, most frequent in the latter.

*Sex. Syst.* Diœc. octand. *Nat. Syst.* Ebenaceæ.  
Linn. Sp. Pl. 1510. Griffith, Med. Bot. 435.

Several parts of the Persimmon have been used in medicine, as the bark and the unripe fruit. They are both powerful astringents, and have been employed with success in bowel complaints and hemorrhages. The bark may be given in infusion, the fruit, in syrup or vinous tincture.

## INFUSION OF PERSIMMON BARK.

R. Persimmon bark,  
bruised, half an ounce  
Boiling water one pint  
Macerate for two hours, and strain.

Used in doses of a fl. ounce, in intermittents, and as a gargle in ulcerated sore throat.

## WINE OF PERSIMMONS.

R. Crushed green persimmons one pound  
Port wine one pint and a-half  
Macerate for fourteen days, strain, and filter. Dose, two to four fl. drachms a day, as an astringent. *Beasley.*

## DIRCA.

## LEATHERWOOD.

The *Dirca palustris* is an indigenous shrub, found in boggy situations, with long, tough roots.

*Sex. Syst.* Octand. monog. *Nat. Syst.* Thymelaceæ.

Linn. Amœn. iii. 12. Griffith, Med. Bot. 561.

The part used is the bark, which is very fibrous; it causes redness and vesication, when applied to the skin; when chewed, it induces salivation. It has been given in infusion, or decoction, as a sudorific, and expectorant, and also in the same class of diseases as mezereon.

## DRACONTIUM.

## SKUNK CABBAGE.

A perennial plant, with very large leaves, indigenous to the United States, growing in wet situations: it is the *Symplocarpus fatidus*.

*Sex. Syst.* Tetrand. monog. *Nat. Syst.* Ononiticeæ.

Linn. Sp. Pl. 1372 (*Dracontium*). Griffith, Med. Bot. 619.

The root is officinal, but the seeds are equally powerful. They both have a very acrid taste when fresh. The leaves, when bruised, have a nauseous smell, and are a good stimulating application to blisters. The root and seeds are stimulant, antispasmodic, and narcotic; and have been employed with success in asthma, hysteria, &c. The dose in powder is from two to twenty grains, several times a-day.

## INFUSION OF SKUNK CABBAGE.

R. Skunk cabbage root one ounce  
Boiling water one pint

Infuse in a covered vessel for an hour, and strain. Dose, an ounce to two ounces.

## TINCTURE OF SKUNK CABBAGE ROOT.

R. Skunk cabbage root, sliced, one ounce  
Alcohol six fl. ounces

Macerate for fourteen days, and strain.

*Turner.*

## TINCTURE OF SKUNK CABBAGE SEED.

R. Skunk cabbage seed two drachms  
Alcohol four fl. ounces

Macerate for fourteen days, and strain.

*Turner.*

## DULCAMARA.

## BITTERSWEET.

The Bittersweet, or *Solanum dulcamara*, is a climbing shrub, indigenous to Europe, and naturalized extensively in this country.

*Sex. Syst.* Pentand. monog. *Nat. Syst.* solanaceæ.

Linn. Sp. Pl. 264. Griffith, Med. Bot. 480.

The officinal portions are the small stems. These have a heavy, unpleasant odor when fresh, but are scentless when dry. They have a somewhat bitter taste, followed by a peculiar sweetness. The properties of Bittersweet are those of a narcotic, diuretic, and diaphoretic. It has been used in chronic rheumatism, asthma, &c., and especially in chronic cutaneous disorders. The dose, in substance, is from thirty grains to a drachm, but it is seldom given in this form.

## DECOCTION OF BITTERSWEET.

R. Bittersweet, bruised, one ounce  
Water one pint and a-half  
Boil down to a pint, and strain.

*U. S. Ph.*

Dose, from one to two fl. ounces, three or four times a-day, gradually increased, till a pint is taken in the twenty-four hours. In chronic eruptions, &c.

## COMPOUND DECOCTION OF BITTERSWEET.

R. Bittersweet half an ounce  
Liquorice root  
Burdock  
Sassafras bark  
Guaicum, each, two drachms  
Water two pints

Boil down to sixteen ounces, and strain.

Dose, one to two fl. ounces, several times a-day, in chronic rheumatism, and venereal affections.

*Augustin.*

## EXTRACT OF BITTERSWEET.

R. Bittersweet, in coarse  
powder, one pound  
Water sufficient  
Mix the powder with a pint of the water,



let it stand for twenty-four hours, put it in a displacement apparatus, and add water gradually, till the liquor that passes is but slightly impregnated with the properties of the bittersweet. Heat the filtered fluid, strain, and evaporate to proper consistence. *U. S. Ph.*

Dose, from five to ten grains.

R. Freshly dried bittersweet, contused, eight ounces  
Diluted alcohol four pints  
Digest by a gentle heat, for fourteen days, express, filter, and evaporate on water-bath, to a proper consistence. Dose, five to eight grains. *J. W. Simes.*

#### SYRUP OF BITTERSWEET.

R. Coarsely powdered bitter-sweet four ounces  
Water twelve fl. ounces  
Alcohol four fl. ounces  
Mix the fluids, pour on the powder in a displacer, until one pound of tincture is obtained, adding water to displace the diluted alcohol. Evaporate to half a pint, add fifteen ounces of sugar, and make a syrup. *W. Procter.*

#### PILLS OF EXTRACT OF BITTERSWEET.

R. Extract of bittersweet one ounce  
Crude antimony  
Powder of bittersweet, each, half an ounce  
Mix, and form pills of two grains each. Dose, from fifteen to twenty, twice a-day, in obstinate cutaneous affections. *Radiis.*

#### MIXTURE OF BITTERSWEET.

R. Extract of bittersweet three drachms  
seneka two drachms  
hemlock one drachm  
Antimonial wine  
Cinnamon water, each, one ounce  
Mix. Forty to eighty drops, four times a-day, in serofulous swelling, spasmodic cough, &c. *Augustin.*  
R. Extract of bittersweet half a scruple  
Infusion of liquorice  
Decoction of mallows, each, one ounce and a-half  
Wine of opium two drops  
Oxymel three drachms  
Mix. A teaspoonful, in chronic catarrh in children. *Radiis.*

## E

### ELATERIUM.

#### ELATERIUM

Is a peculiar substance deposited by the juice of the fruit of *Ecbalium elaterium*, a native of the south of Europe, growing in waste places.

*Sex. Syst.* Monœc. monad. *Nat. Syst.* Cucurbitaceæ.

Richard. Griffith, Med. Bot. 305.

Elaterium is in light, friable, thin, somewhat curled flakes or fragments, of a pale, grayish-green color. The taste is acrid, and bitterish; but the odor is very slight. The dose is from one to two grains, of the common commercial kind, of the pure and genuine, not more than one-eighth to a quarter of a grain.

#### ELATERIUM PILLS.

R. Elaterium ten grains  
Extract of gentian three drachms  
Mix, and form pills of four grains each. One or two, twice a-day, as a hydragogue purgative. *Saunders.*

R. Elaterium six grains  
Extract of gentian  
Soap, each, nine grains

Mix, and form twelve pills. One to four in obstinate constipation. *Radiis.*

R. Elaterium half a drachm  
Aloes  
Gamboge, each, two drachms  
Gum ammoniac two ounces  
Tincture of wormwood sufficient  
Mix, and form pills of four grains each. *Swed. Ph.*

R. Elaterium fifteen grains  
Sulphate of potassa one scruple  
Soap one drachm  
Powdered ginger one drachm and a-half  
Rub the elaterium and sulphate of potassa together, and then with the ginger and soap, adding sufficient water to form mass, and form sixty pills. One to two every hour, until full evacuations are procured. *Sprague.*

#### ELATERIUM MIXTURE.

R. Elaterium one grain  
Spirit of nitric ether two fl. ounces



Tincture of squills

Oxymel of colchicum, each, half a fl. ounce

Syrup one fl. ounce

Mix. A teaspoonful, three or four times a-day, in a little water. In ascites, or hydrothorax. *Ferriar.*

R. Elaterium one grain

Parsley water six fl. ounces

Oxymel of squills

Syrup of buckthorn, each, half fl. ounce

Root of juniper one ounce

Aromatic tincture one drachm

Spirit of nitric ether two fl. drachms

Mix. Used as above, and in the same cases. *Hufeland.*

## OIL OF ELATERIUM.

R. Fruit of elaterium one part

Oil of olives two parts

Digest in a water-bath, for some days, then evaporate all moisture, and strain. As an application for sore nipples, frosted limbs, and painful hemorrhoids.

*Lond. Ph.*

## ELATERINA.

## ELATERIN.

R. Juice of ecballium at will  
Treat with water, dissolve the insoluble residue in alcohol, evaporate to consistence of syrup, purify the crystals that form, by washing with ether.

## TINCTURE OF ELATERIN.

R. Elaterin one grain

Alcohol one ounce

Nitric acid four drops

Mix. Dose, twenty to forty drops, in cinnamon water. *Morris.*

## ELEMI.

## ELEMI.

Elemi is a resinous exudation of various species of plants, principally belonging to the natural order of Amyridaceæ. It occurs in masses of various consistence and color, but usually diaphanous; it has a terebinthinate and somewhat aromatic odor, and a warm, unpleasant taste. It has the usual properties of the turpentine, but is only employed as an external application. It is seldom used in this country, but is extensively employed in Europe.

## OINTMENT OF ELEMI.

R. Elemi one pound

Common turpentine ten ounces

Suet two pounds

Olive oil two fl. ounces

Melt the elemi with the suet; remove from fire, and add turpentine and oil, and strain.

*Lond. Ph.*

As a gentle stimulant to flabby ulcers, &amp;c.

R. Elemi

Venice turpentine, each,

one ounce and a-half

Suet

two ounces

Balsam tolu

one ounce

Melt together, and strain. As a dressing to indolent and ill-conditioned ulcers.

*Neimann.*

## ELEMI CAUTERY PLASTER.

R. Elemi

Spermaceti, each, five parts

Turpentine six parts

White wax ten parts

Melt together, and spread on paper. To keep up discharge from issues. *Soubeiran.*

## EMETINA.

## EMETINE.

This is a peculiar alkaloid, found in the various kinds of ipecacuanha, and to which they owe their active properties. There are two varieties, the colored or impure, and the white or pure.

## IMPURE EMETINE.

R. Ipecacuanha one part

Alcohol (.835) four parts

Macerate for some days, express, and filter; treat the residue with three parts of alcohol as above, unite the tinctures, distil, dissolve the residue in four parts of cold water, filter, evaporate to the consistence of honey, and finish the drying by a gentle heat.

*Cottureau.*

It is reddish brown, inodorous, bitter, and deliquescent.

## PURE EMETINE.

R. Impure emetine at will  
Dissolve in water, treat with magnesia, wash with cold water, treat several times with alcohol, on a water-bath, evaporate the solution to dryness, dissolve the residues in very dilute sulphuric or acetic acid, boil with a little animal charcoal, add an alkaline solution to saturate the acid, dry the precipitate, dissolve it in alcohol, and evaporate to dryness.

*Cottureau.*

It is yellowish or white, pulverulent, not deliquescent. Dose, 1-16th to 1-4th of a grain.

## EMETINE MIXTURE.

R. Impure emetine four grains

Infusion of orange leaves two fl. ounces

Syrup of orange flowers half a fl. ounce



Mix. A dessertspoonful every half hour, to produce vomiting. *Magendie.*

R. Pure emetine, dissolved in  
Nitric acid one grain  
Infusion of linden three fl. ounces  
Syrup of mallows one fl. ounce

Mix. Given as the last. *Foy.*

#### EMETINE LOZENGES.

R. Impure emetine thirty-two grains  
Sugar two ounces  
Mucilage of tragacanth sufficient  
Rub together, and form lozenges of eighteen grains. One will usually vomit a child, and three or four an adult. *Magendie.*

R. Impure emetine thirty-two grains  
Sugar four ounces  
Mucilage tragacanth sufficient  
Mix, and form lozenges of nine grains each. One occasionally as an expectorant. *Guibourt.*

#### SYRUP OF EMETINE.

R. Impure emetine sixteen grains  
Dissolve in a little water, filter, and add to  
Syrup one pound

R. Pure emetine four grains  
Syrup one pound

Mix. The dose of these syrups is a teaspoonful. *Soubéiran.*

### ERGOTA.

#### ERGOT.

Ergot is the diseased seeds of the common rye, but the cause of this morbid growth has given rise to much controversy. The most general opinion, now is, that it is owing to a fungus, but authors are not agreed as to the character of this. The most recent investigator, Mr. Queckett, calls it *Ergotetia arborifaciens*. It is also a question whether the peculiar properties of ergot depend on the fungus, or on the morbid grain. Ergot is one of the few articles that act specifically on the uterus, increasing its contractile energy; and its principal use is as a stimulant to uterine action in labor, and also to check hemorrhage from that organ, but it has been employed in a variety of other diseases, and especially externally as a styptic. The dose, in substance, to aid labor, is from fifteen to twenty grains, to be repeated every twenty minutes, till the desired effect is produced, or till a drachm is taken.

#### INFUSION OF ERGOT.

R. Ergot one drachm  
Boiling water four fl. ounces  
Infuse. One-third as a dose, to be repeated every twenty minutes. *Wood.*

R. Ergot half a drachm  
Boiling water one fl. ounce and a-half  
Infuse for half an hour, and strain. To be repeated every four hours. *Ramsbotham.*

#### DECOCTION OF ERGOT.

R. Ergot one ounce  
Water two pints and a-half  
Boil to two pints, and strain. A table-spoonful every quarter of an hour.

#### WINE OF ERGOT.

R. Bruised ergot two ounces  
Sherry wine one pint  
Macerate for fourteen days, express, and filter. *U. S. Ph.*

Dose, in labor, two to three fl. drachms; for other purposes, one to two fl. drachms.

#### SYRUP OF ERGOT.

R. Ergot one ounce and a-half  
White wine two ounces  
Macerate for eight days, express, and filter; treat the residue three times with water, mix together all the decoctions, and add  
Sugar one pound

Evaporate to consistence of syrup, and add the vinous tincture. Dose, one to two fl. ounces. *Foy.*

R. Ergot twenty grains  
Extract of opium three-quarters of a grain

Syrup eight fl. ounces  
Mix. Dose, one fl. ounce occasionally in engorgements of the uterus. *Lisfranc.*

#### TINCTURE OF ERGOT.

R. Ergot two ounces and a-half  
Diluted alcohol one pint  
Macerate for fourteen days, and strain. Dose, twenty drops to a fl. drachm. As a sedative. *Guy's Hospital.*

#### COMPOUND POWDER OF ERGOT.

R. Powdered ergot two scruples  
cubeb one ounce  
cinnamon half a drachm  
sugar one drachm

Mix, and divide into eight powders; one to be taken, three or four times a-day, in leucorrhœa or gleet. *Ryan.*

R. Powdered ergot half a drachm  
Aromatic powder half a scruple  
Sugar half a drachm

Mix, and divide into ten powders. One to be given every hour to three hours, in hemorrhages. *Ryan.*

#### ERGOT MIXTURE.

R. Powdered ergot half a drachm  
Syrup half a fl. ounce  
Mint water one fl. ounce

Mix. One-third to be taken at a dose, and repeated every twenty minutes; to aid labor. *Soubéiran.*



R. Powdered ergot                    one drachm  
 Syrup                                    three fl. ounces  
 Tincture of opium                    twenty drops  
 Essence of bergamot                    six drops  
 Mix. A spoonful, to be repeated as may  
 be required. *Pierquin.*

## ERGOT CLYSTER.

R. Powdered ergot                    one to three drachms  
 Water                                    twelve ounces  
 Boil for ten minutes, and strain. As an  
 enema, to aid labor. *Foy.*

## COMPOUND ERGOT PILLS.

R. Powdered ergot                    half a drachm  
 Extract of gentian                    one drachm  
 Rub together, and make fifteen pills. In  
 dysmenorrhœa. One pill to be taken three  
 times a-day. *Dewees.*

## ERGOT INJECTION.

R. Powdered ergot                    half an ounce  
 Boiling water                        half a pint  
 Infuse, and strain. As an injection in  
 erythema of the vagina and urethritis.  
*Desruelles.*

## EXTRACT OF ERGOT.

R. Powdered ergot                    at will  
 Water                                    sufficient  
 Exhaust the ergot by the process of displace-  
 ment; heat the result on water-bath; filter,  
 and evaporate to consistence of syrup, add  
 alcohol to precipitate gummy principles,  
 permit to rest, decant, and evaporate to  
 consistence of soft extract. *Bonjean.*

Said to be very efficacious in hemorrhages,  
 in doses of two grains every two hours; also ap-  
 plied as a styptic to bleeding vessels.

## EXTRACT OF ERGOT MIXTURE.

R. Extract of ergot                    fifteen grains  
 Syrup of orange flowers                    one fl. ounce  
 Water                                    three fl. ounces  
 Mix. A tablespoonful, every quarter of an  
 hour, in uterine hemorrhage. Dose to be  
 increased, if requisite. *Bonjean.*

## PILLS OF EXTRACT OF ERGOT.

R. Extract of ergot                    one drachm  
 Powdered liquorice root                    sufficient  
 Mix, and make fifty pills. Dose, six to  
 ten, a-day.

## OIL OF ERGOT.

R. Powdered ergot                    at will  
 Sulphuric ether                        sufficient  
 Exhaust ergot by process of displacement;  
 evaporate result by a gentle heat. *Wright.*

Dose, from twenty to fifty drops, in some ap-  
 propriate vehicle, in parturition, &c. In doses of  
 ten drops, every three hours, in diarrhœa, gastric  
 irritability, &c.

## ERIGERON.

## 1. ERIGERON CANADENSE.

## CANADA FLEABANE.

A common native plant, with numerous white  
 flowers, found in waste places, in the northern  
 and middle States.

*Sex. Syst.* Syngen. super. *Nat. Syst.* Asteraceæ.  
*Willd. Sp. Pl.* iii. 1954. *Torrey & Gray;*  
*Fl.* ii. 175.

The whole plant is used; it has a rather plea-  
 sant smell, and a bitterish, acrid, somewhat  
 astringent taste. It is said to be diuretic, tonic,  
 and astringent, and to be useful in dropsies and  
 diarrhœa. Dose, in substance, thirty grains to a  
 drachm.

## INFUSION OF CANADA FLEABANE.

R. Canada fleabane                    one ounce  
 Boiling water                        one pint  
 Infuse. Dose, two to four fl. ounces.  
*Dupuy.*

## EXTRACT OF CANADA FLEABANE.

R. Canada fleabane                    one pound  
 Water                                    one gallon  
 Boil down to four pints, filter, and evapo-  
 rate to proper consistence. Dose, five to  
 ten grains.

## 2. ERIGERON ANNUUM.

## FLEABANE.

This species, which has been also called, *E.*  
*heterophyllum*, is very closely allied to, and iden-  
 tical in properties with, *E. strigosum* (*E. Philadel-*  
*phicum*). It is also a common plant in the  
 United States.

*Persoon, Synop.* ii. 431. *Griffith, Med. Bot.*  
 394.

The whole plant is used. It has a peculiar,  
 but not unpleasant odor, when bruised, and as-  
 tringent, bitterish taste. It is much celebrated as  
 a diuretic, especially in dysuria of children.

## INFUSION OF FLEABANE.

R. Fleabane                            one ounce  
 Boiling water                        one pint  
 Infuse. Dose, two to four fl. ounces,  
 every three or four hours. *Dewees.*

## ERYNGIUM.

## 1. ERYNGIUM MARITIMUM.

## SEA HOLLY.

This plant is a native of Europe, growing on  
 sandy beaches.

*Sex. Syst.* Pentand. digyn. *Nat. Syst.* Api-  
 acææ.

*Linn. Sp. Pl.* 337. *Griffith, Med. Bot.* 315.

It is thought to be diuretic, and was also es-  
 teemed aphrodisiac. The part used is the root,  
 which has a sweet, agreeable taste, and an aro-  
 matic odor.



## 2. ERYNGIUM AQUATICUM.

## BUTTON SNAKEROOT.

A native species, growing in wet places, from Virginia to Florida.

Linn. Sp. Pl. ed. 2. p. 336. Torrey & Gray, Fl. 1. 604.

The root is bitter, aromatic, and pungent. It is diaphoretic and expectorant, and is useful in the same cases to which senega is appropriate. Used to some extent in South Carolina.

## EUPATORIUM.

## BONESET.

This, the *E. perfoliatum*, is a native plant, found in abundance in most parts of the United States, in moist situations. Several other native species are almost identical in their properties, as *E. tenuifolium*, *E. purpureum*, &c.

Sex. Syst. Syngen. æqual. Nat. Syst. Asteraceæ.

The parts used are the leaves and flowers. These have a faint odor, and a peculiar, bitter taste. It is tonic and diaphoretic, and, in large doses, emetic and laxative. Dose, in substance, as a tonic, is twenty to thirty grains.

## INFUSION OF BONESET.

R. Boneset one ounce  
Boiling water one pint  
Infuse for two hours, in a covered vessel, and strain. *U. S. Ph.*

As a diaphoretic, to be taken freely, warm; as a tonic, a fl. ounce, cold.

## DECOCTION OF BONESET.

R. Boneset one ounce  
Water one pint and a-half  
Boil down to a pint.

Dose, a wineglassful, or more, as an emetic and cathartic.

## COMPOUND INFUSION OF BONESET.

R. Boneset  
Sage, each, half an ounce  
Cascarilla one drachm  
Boiling water one pint and a-half  
Infuse till cold, and strain. A wineglassful, every three or four hours. In hectic fever. *Ellis.*

## EUPHORBIA.

## SPURGE.

## 1. EUPHORBIA COROLLATA.

## BOWMAN'S ROOT.

A native plant, with numerous white flowers, found in many parts of the United States, in dry, sandy soil.

Sex. Syst. Dodecand. trigyn. Nat. Syst. Euphorbiaceæ.

Linn. Sp. Pl. 258. Griffith, Med. Bot. 592.

The root, which is the part used, is large, white, inodorous, and almost tasteless. It is a safe and certain emetic, in doses of fifteen or twenty grains, and diaphoretic in that of three or four.

## 2. EUPHORBIA IPECACUANHA.

## WILD IPECACUANHA.

Also a native plant, with numerous procumbent stems, and variously shaped leaves.

Linn. Sp. Pl. 653. Griffith, Med. Bot. 592.

The root, the part used, is large, white, almost inodorous, and a somewhat sweet taste. It is an energetic emetic, but, like the last species, apt to act on the bowels. Dose, ten to fifteen grains.

## COMPOUND POWDER OF WILD IPECACUANHA.

R. Powdered wild ipecacuanha  
opium, each, six grains  
Sulphate of potassa forty-eight grains  
Triturate well, and divide into six powders. As a substitute for Dover's powder, to which it is equal, if not superior. *W. P. C. Barton.*

## 3. EUPHORBIA HYPERICIFOLIA.

## BLACK PURSLANE.

A small, procumbent, native plant, with leaves often marked with a purple spot.

Linn. Sp. Pl. 652. Griffith, Med. Bot. 593.

The whole plant is used, and is said to be astringent and narcotic. It is inodorous, but has a sweetish, somewhat austere taste. It has been recommended in diarrhoea, fluor albus, &c.

## INFUSION OF BLACK PURSLANE.

R. Dried black purslane half an ounce  
Boiling water one pint  
Infuse for half an hour, and strain. Dose, in dysentery and diarrhoea, a tablespoonful, every hour, till the morbid symptoms yield, then less frequently; in menorrhagia and leucorrhoea, a wineglassful, three times a-day. *Zollickoffer.*

## 4. EUPHORBIA LATHYRIS.

## MOLE PLANT.

A tall species, native of Europe, but often cultivated in this country.

Linn. Sp. Pl. 655. Lindley, Flor. Med. 194.

The seeds, which are the officinal portion, afford a purgative oil, which is tasteless, and almost inodorous. It acts on the bowels, in doses of from four to twelve drops, but is apt to cause nausea and vomiting.



## MIXTURE OF OIL OF MOLE PLANT.

- R. Oil of mole plant            eight drops  
 Powdered gum Arabic        one drachm  
 Sugar                            two ounces  
 Distilled water               three fl. ounces

Mix. As a purgative draught.

*Pichonnier.*

## EUPHORBIIUM.

## EUPHORBIIUM

Is the concrete resinous juice obtained from several succulent species of euphorbia, principally natives of Africa. It is in the form of tears, or small, irregular masses, of a yellowish, or reddish color. The taste is at first slight, but soon becomes acrid and burning; the odor is very faint. It is a violent emetic and purgative, and is not used internally; as an external application it is employed in Europe, and also as a sternutatory.

## STERNUTATORY OF EUPHORBIIUM.

- R. Powdered euphorbium    one drachm  
       white hellebore four scruples
- Mix. Very violent in its action. *Pierquin.*

## OIL OF EUPHORBIIUM.

- R. Euphorbium                    one part  
 Olive oil                        ten parts
- Digest for ten days, and filter. As a friction in paralysis. *Van Mons.*

## PLASTER OF EUPHORBIIUM.

- R. Pitch  
 Yellow wax, each,            three ounces  
 Turpentine                    one ounce
- Melt together, and add  
 Gum ammoniac  
 Olibanum  
 Mastich  
 Calamine, each,              one ounce  
 Euphorbium  
 Pyrethrum  
 Common salt, each,        two ounces

Stir, till cold. A stimulating and rubefacient application, known as *capuchin* ointment. *Wirtemberg Ph.*

## EUPHRASIA.

## EYEBRIGHT.

The *Euphrasia officinalis* is a small, annual plant, a native of Europe, and the northern parts of Asia and America.

*Ser. Syst.* Didynam. gymnos. *Nat. Syst.* Scrophulariaceæ.

*Linn. Sp. Pl.* 841. *Lindley, Flor. Med.* 506.

The whole plant is used. It is slightly bitter and aromatic, and, at one time, was much used in diseases of the eyes. Dr. Kranichfeld says it is peculiarly beneficial in catarrhal ophthalmia, and also in cough and other catarrhal affections.

## F

## FERRUM.

## IRON.

A solid, hard, malleable, ductile metal, of a silvery white color, and fibrous texture, attracted by the magnet, of a slight styptic taste, and a faint, but peculiar odor, when rubbed. Oxidizable in the air, and when heated to whiteness, burning with great brilliancy. It is employed medicinally in a great number of forms, both in the metallic form, in those of oxides and salts. All these are powerfully tonic and astringent.

## FERRI FILUM.

## IRON WIRE.

## FERRI RAMENTA.

## IRON FILINGS.

These should always be perfectly pure, when

used internally, and in a state of minute division.

## PREPARED METALLIC IRON.

- R. Iron filings                    at will
- Pound in an iron mortar, with great force, sift through a fine sieve, and porphyryze, until all metallic brilliancy has disappeared, and again sift. Keep in a perfectly dry place. *Paris Cod.*

Dose, from five to twenty grains.

- R. Spread oxide of iron in a tube, heat the tube, and cause a stream of hydrogen gas to pass through it, till the iron is reduced. *Soubéiran.*

- R. Place precipitated carbonate of iron on layers of iron gauze, in a tube of wrought iron, pass a stream of hydrogen through it, heat to a dull red, and maintain this for some hours, keeping up a small current of gas, till cool. *W. Procter.*



**POWDER OF METALLIC IRON.**  
 R. Prepared metallic iron ten grains  
 Sugar one drachm  
 Mix. To be given, four times a-day, in chlorosis and amenorrhœa. *Brera.*

**COMPOUND POWDER OF METALLIC IRON.**  
 R. Prepared metallic iron  
 Powdered guaiacum, each, half a drachm  
 Prepared oyster shell half an ounce  
 Sugar three drachms  
 Mix. Divide into ten powders; one, twice a-day, in scrofula. *Gœlis.*

R. Prepared metallic iron  
 Powdered cinnamon, each, half a drachm  
 Sugar, three drachms  
 Mix, and divide into twelve powders.  
 One every three hours, as a tonic. *Guibourt.*

R. Prepared metallic iron one grain  
 Carbonate of magnesia five grains  
 Powdered cinnamon two grains  
 sugar one scruple  
 Mix. One-half in the morning, the other in the evening, in rachitis. *Hufeland.*

R. Prepared metallic iron  
 Powdered anise  
 Castor, each, two drachms  
 Powdered cinnamon  
 nutmeg, each, one drachm  
 Mix, and divide into twenty-four powders.  
 One, morning and evening, in chlorosis. *St. Marie.*

**BOLUS OF PREPARED IRON.**  
 R. Prepared metallic iron  
 Powdered myrrh, each, ten grains  
 Aromatic powder two grains  
 Powder of rosemary half a drachm  
 Syrup of orange peel sufficient  
 Make a bolus. To be taken morning and evening, in amenorrhœa. *Augustin.*

**PILLS OF PREPARED IRON.**  
 R. Prepared metallic iron two ounces  
 Extract of wormwood, sufficient  
 Mix, and make six-grain pills. Three or four, morning and evening, in chlorosis, dyspepsia, &c. *Soubeiran.*

R. Prepared metallic iron one drachm  
 Powdered black hellebore one scruple  
 Extract of gentian two drachms  
 Syrup of saffron sufficient  
 Mix, and form sixty pills. Two, three times a-day, as an emmenagogue. *St. Marie.*

R. Prepared metallic iron half an ounce

Powdered columbo four scruples  
 rhubarb  
 cinnamon, each, two scruples  
 Extract of quassia sufficient  
 Mix, and form pills of three grains each.  
 Three to four a-day, as a tonic in debility. *Jahn.*

**AROMATIC MIXTURE OF IRON.**  
 R. Powdered Peruvian bark one ounce  
 Sliced columbo three drachms  
 Bruised cloves two drachms  
 Iron filings half an ounce  
 Digest for three days in a covered vessel, and add as much peppermint water as will yield a mixture of twelve ounces, after filtration, then add compound tincture of cardamom, three ounces, and tincture of orange peel, three drachms. *Dub. Ph.*

As a tonic, in doses of one or two fl. ounces.

## FERRI ACETAS.

**ACETATE OF IRON.**  
 R. Subcarbonate of iron one part  
 Acetic acid six parts  
 Digest for three days, and filter. *Dub. Ph.*  
 Dose, from ten to twenty-five drops, in water.

**TINCTURE OF ACETATE OF IRON.**  
 R. Acetate of potassa two parts  
 Sulphate of iron one part  
 Rectified spirit twenty-six parts  
 Rub the acetate and sulphate together, dry by a moderate heat, and triturate with spirit. Digest in a closed bottle for seven days, and decant. *Dub. Ph.*  
 Dose, from thirty drops to a teaspoonful, in some convenient vehicle.

**ALCOHOLIC TINCTURE OF ACETATE OF IRON.**  
 R. Liquid acetate of iron one part  
 Diluted alcohol seven parts  
 Digest for two days, and filter. Dose, ten to fifty drops. *Guibourt.*

**ETHEREAL TINCTURE OF ACETATE OF IRON.**  
 R. Liquid acetate of iron nine ounces  
 Acetic ether one ounce  
 Alcohol two ounces  
 Mix. Dose, twenty to thirty drops. *Spielmann.*

**ACETATE OF IRON AND ALUM.**  
 R. Sulphate of iron two parts  
 Alum one part



Boiling water ten parts  
Filter the solution, and gradually add solution of subacetate of lead till precipitation ceases; let rest, and filter; evaporate to consistence of a jelly. As a marking ink; it resists alkalies, but not acids.  
*Guibourt.*

## FERRUM AMMONIATUM.

## AMMONIATED IRON.

R. Subcarbonate of iron three ounces  
Muriatic acid ten fl. ounces  
Muriate of ammonia two pounds and a-half  
Distilled water four pints  
Mix the subcarbonate with the acid, in a glass vessel, and digest for two hours, then add the muriate of ammonia, previously dissolved in the water, filter, and evaporate to dryness. *U. S. Ph.*

Said to be useful in amenorrhœa, scrofula, epilepsy, &c. Dose, four to twelve grains, several times a-day.

## COMPOUND POWDER OF AMMONIATED IRON.

R. Ammoniated iron  
Powdered quassia, each, two drachms  
Powdered chamomile half an ounce  
Mix, and divide into twenty-four powders. One, four times a-day, as a febrifuge. *Wurtzburg Ph.*

R. Ammoniated iron five grains  
Gum ammoniac three grains  
Powdered senega five grains  
liquorice one drachm  
Mix, and divide into twelve powders. One, every three hours, in gonorrhœa. *Foy.*

## BOLUS OF AMMONIATED IRON.

R. Ammoniated iron five grains  
Aromatic powder twelve grains  
Mucilage sufficient  
Mix, and make bolus. One to be taken twice a-day, in chlorosis and scrofula. *Ainslie.*

## PILLS OF AMMONIATED IRON.

R. Ammoniated iron  
Galbanum, each, one drachm  
Assafetida two drachms  
Castor one scruple  
Tincture of valerian sufficient  
Beat into mass, and form pills of three grains. Two, morning and evening, in atonic nervous disorders. *Radius.*

R. Ammoniated iron  
Aromatic powder

Extract of Peruvian bark  
Alum, each, half a drachm  
Oil of cinnamon twelve drops  
Beat into mass, and form pills of two grains. Five to ten, morning and evening, in asthenic menorrhagia. *Formey.*

## MIXTURE OF AMMONIATED IRON.

R. Powdered galanga  
rhubarb, each, half a drachm  
Boiling water sufficient  
to obtain three ounces of strained solution, add  
Ammoniated iron six grains  
Peppermint sugar four grains  
Mix. A dessertspoonful, every three hours, in indigestion and chronic diarrhœa of children. *Wendt.*

## TINCTURE OF AMMONIATED IRON.

R. Ammoniated iron four ounces  
Proof spirit one pint  
Dissolve, and filter. Dose, forty to sixty drops. *Lond. Ph.*

## FERRI ARSENIAS.

## ARSENIATE OF IRON.

R. Solution of sulphate of iron at will  
Add gradually a solution of arseniate of potassa, collect, and wash the precipitate.

## PILLS OF ARSENIATE OF IRON.

R. Arseniate of iron three grains  
Extract of hops two drachms  
Powdered mallows half a drachm  
Syrup sufficient  
Beat together, and form forty-eight pills. One or two to be given, daily, in scrofulous, herpetic, and cancerous affections. *Bielt.*

## OINTMENT OF ARSENIATE OF IRON.

R. Arseniate of iron half a drachm  
Phosphate of iron two drachms  
Spermaceti cerate six drachms  
Triturate together. As an application to cancerous ulcers. To be used with great caution. *Carmichael.*

## FERRI BROMIDUM.

## BROMIDE OF IRON.

R. Bromine one part  
Distilled water twenty parts  
Iron filings sufficient  
Boil on a sand-bath; when the solution becomes greenish, filter, and evaporate to dryness. *Magendie.*

## PILLS OF BROMIDE OF IRON.

R. Bromide of iron twelve grains



Conserve of roses      eighteen grains  
Gum Arabic      twelve grains  
Mix, and form twenty pills. Two, in the morning and evening, as a tonic and alterative. *Magendie.*

R. Bromide of iron      one drachm  
Extract of liquorice      sufficient  
Mix, and make sixty pills. One or two, in the morning and evening, in scrofula, &c. *Radiis.*

#### OINTMENT OF BROMIDE OF IRON.

R. Bromide of iron      one drachm  
Bromine      twelve grains  
Lard      one ounce  
Mix. In frictions on the inside of the thighs, in amenorrhœa. *Cadet.*

### FERRI CARBONAS.

#### CARBONATE OF IRON.

From the avidity with which carbonate of iron absorbs oxygen from the air and loses its carbonic acid, it is a very unstable preparation, and therefore what is usually prescribed under this name is merely a subcarbonate, or sesquioxide; by combination with saccharine matter; it, however, is protected from the oxidizing process, to a very great extent.

#### SACCHARINE CARBONATE OF IRON.

R. Sulphate of iron      four ounces  
Carbonate of soda      five ounces  
Pure sugar      two ounces  
Water      four pints  
Dissolve the sulphate and carbonate, each in half the water, add the solutions together, collect the precipitate, divest it of as much water as possible, and immediately triturate it with the sugar. Dry it at a low temperature. *Ed. Ph.*

Dose, from five to thirty grains.

#### PILLS OF CARBONATE OF IRON.

R. Sulphate of iron      four ounces  
Carbonate of soda      five ounces  
Clarified honey      two ounces and a-half  
Syrup  
Boiling water, each,      sufficient  
Dissolve the sulphate and carbonate, each in a pint of water, adding to each solution one fl. ounce of syrup; mix the two solutions in a bottle just large enough to hold the mixture, close accurately with a stopper, and set by to let the carbonate of iron subside; decant, wash, precipitate with water sweetened with syrup, until the washings cease to be saline, express, in flannel, as much of the water as possible, and mix with the honey. Heat over a

water-bath, to proper consistence. This is known as Vallet's carbonate of iron. Dose, from ten to thirty grains, in the course of the day, where iron is indicated. *U. S. Ph.*

#### SUBCARBONATE OF IRON.

R. Sulphate of iron      eight ounces  
Carbonate of soda      nine ounces  
Boiling water      one gallon  
Dissolve the sulphate and carbonate, each in four pints of the water, mix the solutions, stir, set by for powder to subside, decant, wash the precipitate with hot water, wrap in bibulous paper, and dry with gentle heat. *U. S. Ph.*

Dose, five to thirty grains, or even more, several times a-day.

#### POWDER OF SUBCARBONATE OF IRON.

R. Subcarbonate of iron      one drachm  
Divide into six powders. One every six hours, in syrup. In chorea, neuralgia, &c. *Ellis.*

#### MIXTURE OF SUBCARBONATE OF IRON.

R. Powdered myrrh      one drachm  
Sulphate of iron      one scruple  
Carbonate of potassa      one drachm  
Sugar      two drachms  
Water      six fl. ounces  
Mix. Dose, a tablespoonful, according to circumstances. *Ellis.*

R. Myrrh      one drachm  
Carbonate of potassa      twenty-five grains  
Rose-water      seven fl. ounces and a-half  
Powdered sulphate of iron      one scruple  
Spirit of lavender      half a fl. ounce  
White sugar      one drachm  
Rub the myrrh with the rose water, gradually added, mix with the spirit of lavender sugar and carbonate of potassa, and, lastly, the iron; pour into a well-stopped bottle, and keep closed. *U. S. Ph.*

These are well known under the name of Griffith's antihectic mixture. Dose, one or two fl. ounces, two or three times a-day, in chlorosis, debility of the gastric organs, &c.

R. Subcarbonate of iron  
Carbonate of potassa, each,      four ounces  
Black hellebore      two ounces  
Water      sufficient  
Boil, after macerating for some days, stirring from time to time; then let settle, decant, and evaporate to consistence of an extract, and add to this  
Spirit of scurvy  
grass      three times its weight  
And filter. Forty to sixty drops, in hypochondria, melancholia, &c. *Palat. Ph.*



## ARTIFICIAL CHALYBEATE WATER.

- R. Water fifty pints  
 Carbonate of lime five drachms  
                   magnesia ten drachms  
 Black oxide of iron two drachms  
 Sulphate of magnesia six drachms  
 Common salt one drachm

Mix, and condense a thousand cubic inches of carbonic acid in the solution. *Swediasur.*

## COMPOUND PILLS OF SUBCARBONATE OF IRON.

- R. Powdered foxglove half a drachm  
                   yew leaves  
 Carbonate of iron, each, one drachm  
 Ergot two drachms  
 Syrup sufficient

Mix, and make two hundred and fifty pills. Two to eight a-day, with an infusion of savine, in chlorosis and amenorrhœa.

*Neuman.*

- R. Powdered myrrh two drachms  
 Carbonate of soda  
 Sulphate of iron, each, one drachm  
 Syrup sufficient

Rub the myrrh with the carbonate of soda, then add the sulphate of iron, and rub well; beat with syrup into a mass, and divide into eighty pills. *U. S. Ph.*

Dose, two to six pills, three times a-day.

- R. Sulphate of iron  
 Bicarbonate of potassa, each, two drachms  
 Powdered mallowroot, thirty-six grains  
 Gum Arabic eighteen grains

Beat with water into a mass, and form forty-eight pills.

- R. Sulphate of iron  
 Myrrh, each, two scruples  
 Carbonate of potassa  
 Soap, each, half a drachm

Beat together, and form forty pills. Dose, two, three times a-day, as an emmenagogue. *Ellis.*

- R. Subcarbonate of iron  
 Extract of conium, each, one drachm  
 Mix, and divide into twenty-four pills. Two to be taken twice a-day. Given in fluor albus and scrofula. *A. T. Thomson.*

## IRON WINE.

- R. Carbonate of iron one ounce and a-half  
 Contused orange peel  
                   gentian, each, half an ounce  
 Port wine two pints  
 Mix, and digest at moderate heat for three days, repeatedly agitating. *Chapman.*

## IRON PLASTER.

- R. Subcarbonate of iron three ounces  
 Lead plaster two pounds  
 Burgundy pitch half a pound  
 Add the subcarbonate to the other ingredients previously melted together, and stir till cold. *U. S. Ph.*

As a strengthening plaster, in pains, and want of power in the muscles.

## FERRI CARBURETUM.

## CARBURET OF IRON.

This article, well known as Plumbago, or Black Lead, has been highly recommended, of late years, in cutaneous affections, and is recognized as officinal by many of the pharmacopœias of continental Europe.

## PREPARED CARBURET OF IRON.

- R. Pulverized carburet of iron one pound  
 Boil in water for an hour, decant, and digest in eight ounces of water, mixed with two ounces of nitric and muriatic acids; digest for twenty-four hours, decant, and wash well with water, and dry. Dose, from five to fifteen grains, two or three times a-day. *Pruss. Ph.*

## POWDER OF CARBURET OF IRON.

- R. Powdered carburet of iron  
 Sugar, each, half a drachm  
 Mix, and divide into six powders. One to be taken every hour, in lepra and lichen. *Radiis.*

## PILLS OF CARBURET OF IRON.

- R. Powdered carburet of iron  
 Extract of bittersweet, each, one drachm  
 Calomel twelve grains  
 Golden sulphuret of antimony six grains  
 Extract of liquorice sufficient  
 Mix, and form sixty pills. Dose, five, three times a-day. *Niemann.*

## ELECTUARY OF CARBURET OF IRON.

- R. Powdered carburet of iron  
 Flowers of sulphur  
 Crude antimony, each, half an ounce  
 Black oxide of mercury one scruple  
 Wild violet  
 Extract of pulsatilla  
                   bittersweet, each, two drachms  
 Camphor one drachm  
 Syrup of orange peel sufficient  
 Mix. To be used in eight days, in teaspoonful doses. *Phæbus.*



## MIXTURE OF CARBURET OF IRON.

R. Powdered carburet of iron  
Flowers in sulphur, each, one drachm  
Triturate together, and add  
Barley water one pint  
To be taken in four doses. In psora, &c.  
*Brera.*

## OINTMENT OF CARBURET OF IRON.

R. Powdered carburet  
of iron, half an ounce  
Flowers of zinc one drachm  
Lard one ounce  
Triturate together. As an application in  
herpetic affections. *Brera.*

R. Powdered carburet of iron, one ounce  
Flowers of sulphur four scruples  
Rose ointment six ounces  
Mix. *Taddei.*

## PLASTER OF CARBURET OF IRON.

R. Powdered carburet of iron one part  
Soap plaster four parts  
Triturate together. *Giordano.*

## FERRI CHLORIDUM.

## CHLORIDE OF IRON.

R. Subcarbonate of iron at will  
Muriatic acid sufficient  
to dissolve the carbonate, evaporate to dry-  
ness, by a water-bath. *Cottureau.*

R. Peroxide of iron five ounces  
Muriatic acid nineteen fl. ounces  
Boil for ten minutes, then evaporate to  
fifteen ounces, filter. Place under a bell-  
glass, with a vessel containing caustic po-  
tassa; in a few days mammillary granules  
will form; separate these, and preserve in  
a glass-stoppered bottle. *Beral.*

CHLORIDE OF IRON AND ACETATE OF  
LEAD.

R. Liquid chloride of iron four ounces  
Evaporate to one-half, and add  
Acetate of lead four ounces  
Dry, and pulverize. Four to twelve grains  
in hemorrhages. *Cadet de Gassicourt.*

## TINCTURE OF CHLORIDE OF IRON.

R. Subcarbonate of iron half a pound  
Muriatic acid one pint  
Alcohol three pints  
Pour the acid on the subcarbonate; digest  
for three days, shaking occasionally, set by  
to settle, decant, and add the alcohol to  
the clear solution. *U. S. Ph.*

An active and energetic preparation; tonic,  
diuretic, styptic, &c. Dose, ten to thirty drops,  
gradually increased, two or three times a-day.

## SYRUP OF CHLORIDE OF IRON.

R. Chloride of iron one part  
Syrup twenty-three parts  
Dissolve. Dose, one to four drachms.  
*Beral.*

## CHLORIDE OF IRON MIXTURE.

R. Saffron half a drachm  
Water four fl. ounces  
Infuse for an hour, strain, and add  
Chloride of iron half a drachm  
Syrup of valerian two ounces  
Mix. Dose, a tablespoonful several times  
a-day, in hemorrhages or chlorosis.  
*Cadet de Gassicourt.*

R. Mallow root two drachms  
Water sufficient  
to obtain two ounces of decoction, strain,  
and add

Gum Arabic two drachms  
Chloride of iron half a scruple  
Syrup of mallows six drachms  
Mix. Two teaspoonfuls, every hour or two,  
in softening of the stomach. *Radius.*

R. Tincture of chloride of iron  
Compound tincture of  
aloes, each, half a fl. ounce  
Tincture of castor two fl. drachms  
Mix. Ten to thirty drops, three times  
a-day, in infusion of chamomile, as an  
emmenagogue. *Ellis.*

ETHEREAL, OR BESTERCHEFFE'S TINC-  
TURE OF CHLORIDE OF IRON.

R. Chloride of iron one part  
Sulphuric ether four parts  
Mix, and add  
Alcohol eight parts  
*Van Mons.*

R. Tincture of chloride of iron  
Sulphuric ether equal parts  
*Guibourt.*

R. Chloride of iron one part  
Hoffmann's anodyne seven parts  
*Beral.*

These are active preparations. Dose, twenty  
to thirty drops.

## TINCTURE OF CHLORIDE OF IRON.

R. Muriatic acid two ounces (troy)  
Protocarbonate of iron q. s. to saturate  
Honey two ounces and a-half  
Alcohol sufficient  
Saturate the acid with the carbonate, add  
the honey and sufficient alcohol to make  
nineteen fl. ounces; after standing six hours,  
filter. *W. Procter.*



## FERRI CITRAS.

## CITRATE OF IRON.

R. Crystals of citric acid      three parts  
Hydrated oxide of iron (dry) two parts  
Distilled water at 180°      twelve parts  
Dissolve, filter, and evaporate to consist-  
ence of syrup, spread in thin layers, and  
dry. *Duhamel.*

## LIQUID CITRATE OF IRON.

R. Citrate of iron      four ounces  
Water      twelve fl. ounces  
Dissolve, with the aid of heat, and evapor-  
ate until the solution measures half a pint.  
*W. Procter.*

## SYRUP OF CITRATE OF IRON.

R. Liquid citrate of iron      one ounce  
Syrup      fifteen ounces  
Spirit of citron      two drachms  
Mix. Dose, two drachms to an ounce.  
*Beral.*

## WINE OF CITRATE OF IRON.

R. Liquid citrate of iron      eight scruples  
Rhine wine      sixteen ounces  
Digest for two or three days, and filter.  
*Beral.*

## AROMATIC WINE OF CITRATE OF IRON.

R. Iron filings      one ounce  
Lemon juice      three fl. ounces  
Contused gentian      half an ounce  
                                 cinnamon      two drachms  
Rhine wine      sixteen ounces  
Digest for twenty-four hours, and decant.  
The dose is a drachm to a fl. ounce, two or  
three times a-day. *Hamburgh Cod.*  
R. Iron filings      four ounces  
Bitter oranges      four  
Beat them together, and at the end of two  
days, add to the mixture  
Madeira wine      ten fl. ounces  
Spirit of orange peel      two fl. ounces  
Digest, and then express and filter. Dose,  
half a drachm to two drachms.

*Batav. Ph.*

Said to be one of the best compounds of iron.

## TINCTURE OF CITRATE OF IRON.

R. Liquid citrate of iron      two ounces  
Diluted alcohol      thirteen fl. ounces  
Spirit of citron      one fl. ounce  
Mix. Dose, ten to fifty drops. *Beral.*

## SYRUP OF CITRATE OF IRON.

R. Moist protocarbonate of iron,  
(prepared as directed for  
Vallet's mass, and washed  
with sweetened water)      at will  
Dissolve in a slight excess of citric acid,

dissolved in water, and evaporate to dry-  
ness to make the syrup.

R. Citrate of iron      one ounce  
Syrup      five fl. ounces  
Dissolve. Dose, thirty drops to a tea-  
spoonful. *E. Parrish.*

## AMMONIO CITRATE OF IRON.

R. Liquid citrate of iron      one ounce  
Water of ammonia      sufficient  
to saturate the excess of acid. More solu-  
ble than the citrate. *Beral.*

## CITRATE OF IRON AND QUINIA.

R. Liquid citrate of iron      four parts  
Solution of citrate of quinia      one part  
Mix, and evaporate. As a tonic, in doses  
of five grains, three times a-day. *Beral.*

FERRI FERROCYANU-  
RETUM.

## PRUSSIAN BLUE.

R. Sulphate of iron      four ounces  
Sulphuric acid      three fl. drachms and a-half  
Nitric acid      six fl. drachms  
Ferrocyanuret of  
                                 potassium      four ounces and a-half  
Water      two pints  
Dissolve the sulphate in a pint of the wa-  
ter, add the sulphuric acid, and boil.  
Add the nitric acid in small portions, boil-  
ing for a few moments after each addition,  
till a dark color is no longer produced,  
and allow to cool. Dissolve the ferrocy-  
anuret in the rest of the water, and add  
the solution by degrees, to the first liquid,  
stirring each time. Filter, wash the pre-  
cipitate with boiling water, until the wash-  
ings are tasteless. Dry, and powder.

*U. S. Ph.*

Tonic, febrifuge, and alterative. Dose, three  
to five grains, several times a-day, gradually in-  
creased till some effect is produced.

## POWDER OF PRUSSIAN BLUE.

R. Prussian blue      twelve grains  
Sugar      one drachm  
Mix, and divide into twelve powders.  
One, every two hours. In epilepsy.  
*Radius.*

## COMPOUND POWDER OF PRUSSIAN BLUE.

R. Prussian blue      twelve grains  
White pepper  
Mustard, each,      half an ounce  
Mix, and divide into twelve powders.  
One, every quarter of an hour, in the  
apoplexia of intermittents. *Foy.*



R. Prussian blue  
Powdered guaiacum, each, half to one drachm  
Mix, and divide into twelve powders.  
One, three times a-day, in intermittents.  
*Ellis.*

#### PILLS OF PRUSSIAN BLUE.

R. Prussian blue  
eighteen to thirty-six grains  
Gum ammoniac  
Extract of dandelion  
Rhubarb, each, one drachm  
Mix. Four to six, twice a-day. In disease of the ganglionic system. *Radius.*

#### MIXTURE OF PRUSSIAN BLUE.

R. Prussian blue one ounce  
Parsley water six ounces  
Cherry laurel water two drachms  
Mix. A spoonful, three times a-day. In stone and gravel. *Radius.*

#### OINTMENT OF PRUSSIAN BLUE.

R. Codliver oil one ounce  
Prussian blue twenty-four grains  
Cyanide of mercury eight grains  
Oil of cherry laurel four drops  
Mix. As an application in strumous ophthalmia. *Canon de Villards.*

#### BLUE INK.

R. Pure Prussian blue six parts  
Oxalic acid one part  
Triturate with a little water, to a perfectly smooth paste, then dilute with water, to a proper consistence. A little gum may be added to prevent the fluid from spreading.  
*Mohr.*

### FERRI GALLAS.

#### GALLATE OF IRON.

This is not used in medicine, but, with the tannate of iron, forms the basis of writing ink.

#### WRITING INK.

R. Aleppo galls, bruised, twelve pounds  
Water six gallons  
Boil in a copper vessel for an hour, adding water, to supply loss by evaporation, strain; boil residue with four gallons of water, for half an hour, and again boil with two gallons of water; mix decoctions, and add

Sulphate of iron four pounds  
Contused gum  
Arabic three pounds and a-half  
Agitate, permit to settle, strain through fine hair sieve, and keep in close vessels.  
*Lewis.*

R. Powdered galls eight pounds  
Rasped logwood four pounds  
Gum Arabic three pounds  
Sulphate of iron four pounds

copper  
Sugar candy, each, one pound  
Gum ammoniac two ounces  
Water sixteen pints  
Alcohol half a pint

Macerate for forty-eight hours, with a gentle heat, let rest for ten days, and decant into bottles. *Hunt.*

R. Powdered galls fifty parts  
Water eight hundred parts  
Digest for twenty-four hours, strain, and add

Sulphate of iron  
Gum Arabic, each, twenty-five parts  
When clear, add a solution of  
Muriate of ammonia eight parts  
Gum Arabic two parts  
Boiling water sixteen parts  
Oil of lavender one part  
Said to be indelible. *Guibourt.*

R. Bruised galls forty ounces  
Gum Arabic ten ounces  
Sulphate of iron nine ounces  
Soft water forty-five fl. ounces  
Macerate for three weeks, often stirring.  
A very permanent ink.

*English Exchequer.*

### FERRI IODIDUM.

#### IODIDE OF IRON.

R. Iodine two ounces  
Iron filings one ounce  
Distilled water one pint and a-half

Mix the iodine with a pint of the water, in a glass vessel, gradually add the iron filings, constantly stirring. Heat till liquid becomes of a light greenish color, filter and wash residuum with remainder of water, boiling hot. Evaporate liquid to dryness, in an iron vessel, at a heat not above 212°. Keep in well-closed bottle.

*U. S. Ph.*

Tonic, alterative, diuretic, and emmenagogue.  
Dose, three grains, gradually increased to eight, or more.

#### LOZENGES OF IODIDE OF IRON.

R. Iodide of iron thirty grains  
Saffron half a drachm  
Mucilage of tragacanth  
Infusion of cinnamon, each, sufficient  
Mix, and make two hundred and forty lozenges. Fifteen to twenty a-day, gra-



dually augmenting, in amenorrhœa and leucorrhœa.

*Pierquin.*

#### PILLS OF IODIDE OF IRON.

R. Iodide of iron                   one drachm  
Bromide of sodium           half a drachm  
Extract of liquorice           sufficient  
Mix, and form pills of two grains. One to three, morning and evening, in scrofula, &c.

*Wernick.*

#### PILLS OF PROTO-IODIDE OF IRON.

R. Crystallized sulphate  
of iron                   24.7 grains  
Iodide of potassium       32.4 grains  
Gum tragacanth           4.6 grains  
Sugar                   15.4 grains  
Syrup of marsh mallows  
Powder of marsh mallows,  
each,                   sufficient

Mix, and make thirty-six pills. Reduce the sulphate to a fine powder, then the iodide of potassium; triturate the mixture, then add the gum, sugar, and syrup, and, if necessary, the powder of marshmallow. Each contains .8 of a grain of dry iodide, or 1.09 of the hydrated iodide of iron. Keep in closed bottles, in a dry place.

*Callond.*

#### SOLUTION OF IODIDE OF IRON.

R. Iodide of iron                   one drachm  
Distilled water                   one fl. ounce  
Mix. Six, to ten drops, three times a-day, in some cold water.

*Ellis.*

R. Iodide of iron                   twenty-four grains  
Distilled water                   one fl. ounce  
Dissolve. A teaspoonful is a dose; this contains three grains.

*Ellis.*

#### OFFICIAL SOLUTION OF IODIDE OF IRON.

R. Iodine                   two ounces  
Iron filings                   one ounce  
Clarified honey               five fl. ounces  
Distilled water               sufficient

Mix the iodine with ten fl. ounces of the water, in a porcelain or glass vessel, gradually add the iron filings, constantly stirring. Heat gently till the mixture becomes of a light greenish color, add honey, continue heat for a short time, and filter. Wash the residuum on filter, and allow to pass till the whole of the filtered liquid amounts to twenty fl. ounces.

*U. S. Ph.*

Dose, from thirty to seventy-five drops.

#### SYRUP OF IODIDE OF IRON.

R. Dry iodine                   two hundred grains  
Fine iron wire, well  
cleaned,                   one hundred grains

White sugar, in

powder,                   four ounces and a-half

Distilled water                   six fl. ounces

Boil the iodine, iron, and water, together, in a glass matrass; at first gently, to avoid the expulsion of iodine vapor, afterwards briskly, till about two fl. ounces remain. Filter this quickly, while hot, into a matrass containing the sugar, dissolve with a gentle heat, and add distilled water to make up six fl. ounces. Twelve minims contain one grain of iodide of iron.

*Ed. Ph.*

#### SYRUP OF PROTO-IODIDE OF IRON.

R. Sulphate of iron           one drachm and a-half  
Iodide of potassium           two drachms  
Water                   half an ounce  
Syrup                   one ounce and a-half  
Rub the salts together, dissolve them in the water, add the syrup, and bottle at once.

*W. Procter.*

#### WINE OF IODIDE OF IRON.

R. Sulphate of iron                   eight parts  
Iodide of potassium,               ten and a-half parts  
White wine

three hundred and twenty parts

Pulverize the two salts, adding a few drops of wine, triturate for a few moments, add the rest of the wine, and filter. To be kept in glass-stopped bottles. Thirty-two parts contain one of the iodide.

*Callond.*

R. Iodide of iron                   half an ounce  
Claret wine                   one pint  
Dissolve. A dessertspoonful, morning and evening.

*Soubeiran.*

#### TINCTURE OF IODIDE OF IRON.

R. Iodide of iron                   one drachm  
Alcohol  
Distilled water, each,           one fl. ounce  
Mix. Thirty drops to a fl. drachm, twice a-day.

*Foy.*

#### TINCTURE OF PROTO-IODIDE OF IRON.

R. Sulphate of iron                   eight parts  
Iodide of potassium               ten and a-half parts  
Alcohol (.842)                   one hundred and sixty parts  
Triturate the salts, separately, and then together; add alcohol, and filter. To be kept in glass-stopped bottles. Sixteen parts contain one of iodide of iron.

*Callond.*

#### MIXTURE OF IODIDE OF IRON.

R. Iodide of iron                   sixteen grains



Compound tincture  
of gentian one fl. ounce  
Distilled water seven fl. ounces  
Mix. Two tablespoonfuls, two or three  
times a-day. *Ashwell.*

INJECTION OF IODIDE OF IRON.  
R. Iodide of iron half an ounce  
Distilled water one pint  
Dissolve. As an injection, or lotion, in  
amenorrhœa, leucorrhœa, &c. *Soubeiran.*  
R. Iodide of iron three grains  
Distilled water six fl. ounces  
Dissolve. In gonorrhœa, strength to be  
increased if required. *Ricord.*

SYRUP OF IODIDE AND CHLORIDE OF  
IRON.  
R. Iodine three hundred and  
eighty-four grains  
Muriatic acid  
(1.16) four ounces and a-half  
Powdered sugar twelve ounces  
Iron filings  
Distilled water, each, sufficient  
Mix the acid with an ounce and a-half of  
clear iron filings; allow to stand; agitate  
occasionally, when neutralized, filter. Mix  
the iodine with three ounces of water in a  
flask, add half the weight of iron filings,  
agitate till all the iodine is combined, and  
filter. Mix the solutions, add the sugar,  
and as much water as will make a pint.  
*W. Procter.*

SYRUP OF IODIDE AND CHLORIDE OF IRON  
MIXTURE.  
R. Syrup of iodide and  
chloride of iron two drachms  
Syrup of orange peel four drachms  
Infusion of cascarrilla four fl. ounces  
Mix. One-fourth to be taken twice a-day.  
*Battley.*

BATH OF IODIDE OF IRON.  
R. Iodide of iron two ounces  
Water two hundred pints  
In leucorrhœa, amenorrhœa, chlorosis, &c.  
The quantity of iodide may be increased  
gradually to double, for adults. *Pierquin.*

OINTMENT OF IODIDE OF IRON.  
R. Iodide of iron one drachm and a-half  
Lard one ounce  
Triturate together. A piece as large as a  
hazelnut to be rubbed on the inside of the  
thigh, morning and evening, in leucorrhœa,  
and amenorrhœa, also beneficial in glandu-  
lar swellings. *Pierquin.*

## FERRI LACTAS.

## LACTATE OF IRON.

R. Lactate of lime one hundred parts  
Boiling water five hundred parts  
Dissolve.  
Pure sulphate of iron sixty-eight parts  
Distilled water five hundred parts  
Dissolve  
Mix the two solutions; filter, add an ex-  
cess of lactic acid, heat in a water-bath,  
stirring constantly; filter, to separate the  
sulphate of lime, evaporate rapidly either  
in an iron vessel or porcelain one, with the  
addition of some iron filings; filter, and  
crystallize, wash the crystals with alcohol,  
and dry. *Lepage.*

Used in same cases as other preparations of  
iron. Dose, one or two grains, frequently re-  
peated.

LOZENGES OF LACTATE OF IRON.  
R. Lactate of iron one drachm  
Sugar one ounce and a-half  
Mucilage of gum Arabic sufficient  
Mix, and make lozenges of twelve grains.  
*Cap.*

PILLS OF LACTATE OF IRON.  
R. Lactate of iron  
Powdered mallows equal parts  
Clarified honey sufficient  
Make pills of ten grains each. *Cap.*

SYRUP OF LACTATE OF IRON.  
R. Lactate of iron one drachm  
White sugar twelve ounces and a-half  
Boiling distilled  
water six fl. ounces and a-half  
Triturate the lactate with half an ounce of  
the sugar, dissolve quickly in the boiling  
water, pour the solution into a matrass,  
place on a sand-bath, add the rest of the  
sugar gradually; when dissolved, filter,  
and when cold, pour in well-stopped bot-  
tles.

Dose, from two to four fl. drachms.

*Cap.*

## FERRI MALAS.

## MALATE OF IRON.

R. Black oxide of iron at will  
Juice of sour apples sufficient  
Heat the juice to the boiling point, add the  
oxide, as long as any is dissolved, cool,  
filter, and evaporate to dryness. Dose,  
eight grains to a scruple. *Van Mons.*

SOLUTION OF MALATE OF IRON.  
R. Malate of iron one ounce



Water twelve fl. ounces  
Dissolve, and filter. As an astringent, in doses of a fl. drachm. *Spieglmann.*

TINCTURE OF MALATE OF IRON.

R. Malate of iron one part  
Cinnamon water  
Alcohol, each, three parts  
Macerate for three days, and filter. Dose, forty to sixty drops. *Austrian Ph.*

MIXTURE OF MALATE OF IRON.

R. Extract of Peruvian bark  
gentian, each, one drachm  
Peppermint water  
Chamomile water, each, one fl. ounce  
Malate of iron half a drachm  
To be taken in divided doses, in dyspepsia, &c. *Augustin.*

BITTER INFUSION OF IRON.

R. Iron filings three ounces  
Bruised ginger  
gentian, each, one ounce  
orange peel half an ounce  
Infuse in a pint of strong old cider for a month, frequently agitating, and filter.

Dose, half a drachm to one drachm, thrice daily, as a tonic.

FERRI NITRAS.

NITRATE OF IRON.

R. Red oxide of iron four drachms  
Nitric acid six fl. drachms  
Dissolve, and add  
Water one fl. ounce  
Filter. Dose, six to twelve drops, in a convenient vehicle. In dyspepsia, and calculous complaints. *Swedjaur.*

SOLUTION OF PERSESQUINITRATE OF IRON.

R. Iron wire (cut small) one ounce and a-half  
Nitric acid three fl. ounces  
Water fifteen fl. ounces  
Dissolve, decant, and filter, add  
Muriatic acid one fl. drachm  
Diluted with sufficient water to make the whole twenty fl. ounces. Dose, ten drops several times a-day, gradually increasing. Has been of great benefit in chronic diarrhoea, &c. *Kerr.*

FERRI OXIDUM HYDRATUM.

HYDRATED OXIDE OF IRON.

R. Sulphate of iron four ounces  
Sulphuric acid three fl. drachms and a-half

Nitric acid six fl. drachms  
Solution of ammonia sufficient  
Water two pints

Dissolve the sulphate in the water, gradually add the sulphuric acid, boil, add nitric acid very gradually, boiling after each addition, till no dark color is produced. Filter, allow to cool, add solution of ammonia, stirring well. Wash precipitate with water, as long as washings give a precipitate with chloride of barium. Keep in close bottles, under water. *U. S. Ph.*

Used, in large doses, as an antidote to arsenic.

FERRI OXIDUM NIGRUM.

BLACK OXIDE OF IRON.

R. Sulphate of iron six ounces  
Sulphuric acid two fl. drachms, two fl. scruples  
Pure nitric acid four fl. drachms and a-half  
Stronger water of ammonia four fl. ounces and a-half  
Boiling water three pints  
Dissolve half the sulphate in half the water, add the sulphuric acid, and boil; add the nitric acid, gradually, boiling briskly after each addition. Dissolve the rest of the sulphate in the remainder of the water, mix the two solutions, and immediately add the ammonia, stirring briskly. Collect the precipitate on a muslin filter, wash in water till the washings scarcely give a precipitate with nitrate of baryta; dry, at a low temperature. *Ed. Ph.*

R. Wash blacksmiths' scales with water, dry, separate metallic iron by means of magnet, reduce to powder, and proceed as for prepared chalk. *Dub. Ph.*

R. Iron filings, any quantity; place in an earthenware pan, wet thoroughly with water; in an hour, heap up on an iron plate; twelve hours afterwards, moisten with hot water, and repeat this in twelve hours, and, a day afterwards, triturate with cold water, decant the turbid water, let it settle, collect the deposit on a filter, and when perfectly drained dry by a mild heat. *Par. Cod.*

These do not give identical products, but they are sufficiently alike to be used for the same purposes.

POWDER OF BLACK OXIDE OF IRON.

R. Black oxide of iron three parts  
Aromatic powder six parts  
Sugar eight parts



Mix. Dose, thirty to forty grains, twice a-day, in leucorrhœa and rachitis.

*Swedjaur.*

R. Black oxide of iron  
Columbo, each, one drachm  
Rind of bitter orange half an ounce  
Cinnamon one scruple

Mix, and divide into nine powders. Three to be given a-day, in chlorosis and amenorrhœa.

*Brera.*

#### PILLS OF BLACK OXIDE OF IRON.

R. Black oxide of iron eight grains  
Powdered saffron

valerian, each, sixteen grains

Syrup of wormwood sufficient

Mix, and form eight pills; four to eight a-day, in amenorrhœa.

*Foy.*

R. Black oxide of iron  
Powdered winter's

bark, each, one drachm

Extract of ox bile

centaury, each, three drachms

Mix, and form pills of four grains. Four to six a-day, in engorgements of the abdominal viscera.

*Pierquin.*

#### ELECTUARY OF BLACK OXIDE OF IRON.

R. Black oxide of iron half an ounce  
Carbonate of potassa two scruples

Prepared chalk

Ginger, each, two drachms

Syrup of orange-peel sufficient

Mix well. Dose, half a drachm, two or three times a-day, in chlorosis and leucorrhœa.

*Swedjaur.*

R. Black oxide of iron  
Soap, each, one drachm

Extract of hemlock twelve grains

squill one scruple

Oxymel sufficient

Mix well. Dose, half a drachm to a drachm, in dropsy, with engorgement of the abdominal viscera.

*Brera.*

### FERRI OXIDUM RUBRUM.

#### RED OXIDE OF IRON.

R. Sulphate of iron, any quantity; heat till water of crystallization is expelled, then roast by a powerful heat, till all acid is driven off, wash, till all traces of acid disappear, dry.

*Dub. Ph.*

Astringent and tonic, and has been advised in neuralgia, &c., in doses of five to fifteen grains, but is inferior to the subcarbonate.

#### PILLS OF RED OXIDE OF IRON.

R. Red oxide of iron

Powder of pot marygold

Extract of pot marygold, each, one drachm

Mucilage of gum Arabic sufficient

Mix, and form ninety pills. Highly spoken of in cancerous affections; five to eight to be taken, three times a-day.

*Rust.*

#### RED OXIDE OF IRON PLASTER.

R. Litharge plaster two pounds

Frankincense half a pound

Red oxide of iron three ounces

Melt the plaster and frankincense together, add the oxide, and stir well.

*Dub. Ph.*

As a strengthening plaster, in muscular relaxation and weakness.

R. Lead plaster twenty-four parts

Resin six parts

Olive oil

Yellow wax, each, three parts

Red oxide of iron eight parts

Triturate the oxide with the oil, and add to a mixture of the other substances, melted together.

*Guibourt.*

### FERRI PHOSPHAS.

#### PHOSPHATE OF IRON.

R. Sulphate of iron five ounces

Phosphate of soda six ounces

Water one gallon

Dissolve the sulphate and phosphate, each, in four pints of the water, mix the solutions, and let settle; decant, wash the precipitate with hot water, and dry with a gentle heat.

*U. S. Ph.*

Used with advantage in amenorrhœa, and certain cases of dyspepsia. Dose, five to ten grains.

### FERRI SULPHAS.

#### SULPHATE OF IRON.

R. Iron wire, in small pieces, twelve ounces

Sulphuric acid eighteen ounces

Water one gallon

Mix the acid and water, and add the iron; heat till effervescence ceases. Pour off the solution, add half a drachm of sulphuric acid, filter, the end of the funnel touching the bottom of the receiving vessel. Evaporate in a matrass, set aside to crystallize in a covered vessel, drain crystals in a funnel, dry, and keep in well-closed bottles.

*U. S. Ph.*

Astringent and tonic. Dose, from one to five grains.



## DRIED SULPHATE OF IRON.

R. Sulphate of iron at will  
Heat over a moderate fire, in a porcelain vessel, till converted into grayish white mass, cool, and powder. *Ed. Ph.*

Three grains are equal to five of the undried.

## POWDER OF SULPHATE OF IRON.

R. Powdered sulphate of iron four grains  
ginger ten to sixteen grains  
Mix. To be taken twice a-day, in amenorrhœa, chlorosis, &c. *Saunders.*

R. Powdered sulphate of iron half an ounce  
tansy three ounces  
Mix. Dose, a drachm. As a vermifuge. *Dan. Ph.*

## PILLS OF SULPHATE OF IRON.

R. Dried sulphate of iron two parts  
Extract of dandelion five parts  
Conserve of red roses two parts  
Powdered liquorice root three parts  
Beat together, and form into five-grain pills. *Ed. Ph.*

R. Sulphate of iron two drachms  
Extract of wormwood half an ounce  
Syrup of saffron sufficient  
Beat into mass, and divide into one hundred and fifty pills. Four to five, three times a-day. In chlorosis, leucorrhœa, &c. *Werthof.*

R. Sulphate of iron  
Kino  
Gentian  
Extract of gentian, each, one drachm and a-half  
Turpentine three drachms  
Powdered mallow root sufficient  
Beat into mass, and make two hundred and seventy pills. Ten, four times a-day, in secondary gonorrhœa. *Walch.*

## BLAND'S ANTI-CHLOROTIC PILLS.

R. Sulphate of iron, in fine powder one ounce  
Carbonate of potassa, in dry powder one ounce  
Mucilage of tragacanth  
Powdered liquorice root, each, sufficient  
To make ninety-nine pills. Triturate well, and divide into pills. To be taken, one in the morning and evening, for the first three days, a third pill at noon for the next three days, and so increased to nine a-day. *Bland.*

## PILLS OF IRON AND RHUBARB.

R. Dried sulphate of iron four parts  
Extract of rhubarb ten parts

Conserve of red roses five parts  
Beat into mass, and form five-grain pills. *Ed. Ph.*

Tonic and laxative. In dose of two or three.

## COMPOUND SULPHATE OF IRON PILLS.

R. Sulphate of iron one scruple  
Powdered senna  
jalap  
cremor tar-  
tar, each, half a scruple  
ginger twelve grains  
Syrup sufficient  
Form mass, and divide into twenty-five pills. *Ellis.*

These are known as *Hooper's pills*, as are also the following:—

R. Barbadoes aloes eight ounces  
Sulphate of iron four ounces  
Extract of hellebore  
Myrrh  
Soap, each, two ounces  
Powdered cinnamon  
ginger, each, one ounce  
Water sufficient  
Beat into mass, and divide into two and a-half grain pills. *Phil. Coll. Pharm.*  
As laxative and emmenagogue.

R. Dried sulphate of iron one scruple  
Powdered aloes two scruples  
cloves five grains  
Venice turpentine sufficient  
Form mass, and divide into twenty pills. One, three times a-day, as an emmenagogue. *Ellis.*

## SOLUTION OF SULPHATE OF IRON.

R. Sulphate of iron  
Alum, each, six ounces  
Water forty-eight ounces  
Dissolve, filter, and add  
Sulphuric acid four ounces  
In hemorrhages, in doses of ten or twelve drops in water. *Spielmann.*

## SULPHATE OF IRON INJECTION.

R. Sulphate of iron half to one drachm  
Sage water four ounces  
Mucilage gum Arabic half a drachm  
Mix. As an injection, in nasal and uterine hemorrhages. *Berends.*

## COMPOUND IRON MIXTURE.

R. Myrrh one drachm and a-half  
Oil of partridge berry six drops  
nutmeg two drops  
Carbonate of potassa two scruples  
White sugar one ounce  
Sulphate of iron half a drachm  
Distilled water seven fl. ounces



Rub down the myrrh and the oils with some of the water added very gradually, then add the carbonate of potassa and sugar, and lastly the sulphate of iron, dissolved in the rest of the water, and immediately bottle.

*Bond.*

#### MIXTURE OF SULPHATE OF IRON.

R. Sulphate of iron two grains  
Sulphuric acid four to six drops  
White sugar one drachm  
Water four fl. ounces

Mix. A teaspoonful, with water, every two to four hours, as a tonic. *Ellis.*

R. Sulphate of iron four grains  
Elixir of vitriol twenty drops  
Distilled water one fl. ounce

Mix. A teaspoonful, in a wineglass of water, three times a-day. *Ellis.*

R. Powdered myrrh one drachm  
gum Arabic two drachms  
Syrup one fl. ounce  
Infusion of chamomile six fl. ounces

Mix well, and add  
Powdered sulphate of iron fifteen grains  
Cinnamon water one fl. ounce

Mix. Two spoonfuls, every three hours, as an emmenagogue. *Frank.*

#### SYRUP OF SULPHATE OF IRON.

R. Sulphate of iron two drachms  
Water one fl. ounce

Dissolve, filter, and add  
Syrup of gum Arabic seventeen fl. ounces

Mix. A tablespoonful, twice a-day, in chlorosis, leucorrhœa, &c. *Soubéiran.*

### FERRI SULPHURETUM.

#### SULPHURET OF IRON.

Heat an iron rod to a white heat, apply to it a roll of sulphur, receiving the sulphuret of iron in water; separate it from the sulphur, dry, and keep in a well-stopped bottle. *Dub. Ph.*

Used to make hydrosulphuric acid, by the addition of diluted sulphuric acid.

### FERRI ET POTASSÆ TARTRAS.

#### TARTRATE OF IRON AND POTASSA.

R. Subcarbonate of iron three ounces  
Muriatic acid ten fl. ounces  
Solution of potassa five pints and a-half  
Bitartrate of potassa seven ounces and a-half

Distilled water one gallon and a-half  
Mix the subcarbonate with the acid, digest for two hours, pour into a gallon of the water, let rest for an hour, decant, add solution of potassa, wash precipitate, and whilst moist, mix with the bitartrate and half a gallon of the water, keep at a temperature of 140° for thirty hours, often stirring; filter, and evaporate on a water-bath, at same temperature, to dryness.

*U. S. Ph.*

Dose, from ten grains to half a drachm, in solution, or with an aromatic or bitter, in bolus.

#### POWDER OF TARTRATE OF IRON AND COLUMBO.

R. Tartrate of iron and potassa two scruples  
Powdered columbo half a drachm  
Mix, and divide into four powders. One, every three or four hours, in syrup. As a tonic. *Ellis.*

#### EFFERVESCENT POWDER OF TARTRATE OF IRON.

R. Subcarbonate of magnesia two drachms  
Carbonate of potassa half an ounce  
Tartrate of iron two scruples  
Two or three teaspoonfuls in water, during the day; to be taken whilst effervescing. *Radiis.*

#### SOLUTION OF TARTRATE OF IRON AND POTASSA.

R. Tartrate of iron and potassa one ounce  
Cinnamon water one pint  
Dissolve; as a lotion or fomentation for bruises. Internally in leucorrhœa, chlorosis, &c. *Swediaur.*

#### COMPOUND WINE OF TARTRATE OF IRON.

R. Subcarbonate of iron one ounce and a-half  
Powdered gentian  
orange peel, each, half an ounce  
Red wine two pints  
Macerate for three days, and give a wine-glassful two or three times a-day. In dyspepsia, &c. *Ellis.*

R. Subcarbonate of iron two ounces  
Powdered cinnamon one ounce  
Rhine wine two pints  
Macerate for some days, and filter. *Van Mons.*

#### WINE OF TARTRATE OF IRON.

R. Subcarbonate of iron one part  
Rhine wine twenty-four parts  
Macerate for some days, and filter. Dose, one or two ounces. *Soubéiran.*



## COMPOUND TINCTURE OF TARTRATE OF IRON.

R. Tartrate of iron wine  
Tincture of calamus  
Hoffmann's balsamic  
elixir, each, four fl. drachms  
Mix. Dose, seventy drops, morning and evening, in wine. Said to be an excellent remedy in ascarides. *Dorfmeuller.*

## FERRI AMMONIO-TARTRAS.

## AMMONIO-TARTRATE OF IRON.

R. Tartaric acid one part  
Boiling water sufficient  
to dissolve, add  
Iron filings three parts  
Digest in a warm place, for two or three days. Add caustic ammonia in slight excess, stir well, pour off the solution, set it aside, decant the clear liquid, and evaporate to dryness; redissolve in distilled water, add a little more ammonia, filter, and evaporate, in shallow, porcelain dishes, by a gentle heat, till it becomes brittle, then chip it off with a blunt-pointed knife.

*Aikin.*

R. Tartaric acid fifty drachms  
Water one gallon

Saturate with

Carbonate of ammonia

And add fifty drachms more of acid. Heat the solution in a water-bath, and add moist hydrated oxide of iron (derived from fifty-three and a-half drachms of subcarbonate of iron dissolved in muriatic acid, and precipitated by ammonia). Digest till the oxide is dissolved, filter, and evaporate to dryness by means of a water-bath.

*W. Procter.*

Has the same properties as the other preparations of iron, but has the advantage of being readily soluble, of being more palatable, and very permanent. The dose is five grains or more, in pills or solution.

## SOLUTION OF AMMONIO-TARTRATE OF IRON.

R. Ammonio-tartrate of iron half a drachm  
Distilled water seven fl. drachms  
Rectified spirit one fl. drachm  
Mix, and dissolve. *Aikin.*

## FERRI TANNAS.

## TANNATE OF IRON.

R. Tannic acid ninety parts

Boiling water sufficient  
to dissolve, add gradually

Subcarbonate of iron four hundred and forty parts  
Agitate till effervescence ceases. Evaporate at 176°, in a porcelain vessel, until it becomes thick, then dry on a plate, in a stove, at 95°. *Benedetti.*

Found useful in chlorosis, in doses of two or three grains, thrice a-day, increasing the dose as required.

## FERRI VALERIANAS.

## VALERIANATE OF IRON.

R. Clean iron filings at will  
Put in a wedgewood mortar, add gradually an equal weight of valerianic acid, stirring continually. In an hour, add distilled water, pour the whole into a flask, warm gently, and filter. The surface, exposed to the air, becomes covered with a crystalline layer of valerianate of iron. Collect this on a filter, repeat this as long as crystals are formed.

Dose, one to two grains. *Ruspini.*

## FICUS.

## FIGS.

Figs are the fruit of *Ficus carica*, a small tree, a native of Asia, and now cultivated in all warm climates. There are numerous varieties. *Sex. Syst.* Polygam. diœc. *Nat. Syst.* Monacœ.

Linn. Sp. Pl. 1513. Griffith, Med. Bot. 576.

They come to this country in a dry state, from the ports of the Levant. They are nutritious, laxative, and demulcent, and are used in medicine in demulcent decoctions, and sometimes externally, as a suppurative poultice.

## DECOCTION OF FIGS.

R. Figs one ounce  
Water one pint  
Boil, and strain. Demulcent and pectoral. *Cadet.*

## GARGLE OF FIGS.

R. Figs  
Mallow root, each, one ounce  
Milk two pints  
Boil away one-fourth. As an emollient gargle in sore throat. *Radiis.*

## COMPOUND DECOCTION OF FIGS.

R. Figs  
Raisins, each, two ounces  
Barley water two pints  
Boil for a quarter of an hour, and add  
Liquorice root half an ounce  
Infuse, and strain. *Foy.*



This is identical with the compound decoction of barley, of the Lond. Ph., and like it, is nutritive, demulcent, and laxative.

#### COFFEE OF FIGS.

R. Figs, roasted like coffee, two drachms  
Water one pint  
Boil gently, and strain. *Radius.*

This is said to be an excellent substitute for coffee, and to form an excellent diet in diseases of the chest.

#### PASTE OF FIGS.

R. Figs at will  
Reduce to a pulp, express through a hair sieve, mix with four times the weight of sugar, concentrate by a gentle heat, and cut into lozenges. *Soubeiran.*

### FILIX MAS.

#### MALE FERN.

The Male Fern is the rhizome of *Nephrodium filix mas*, a native of the temperate regions of the whole world, in shady, rather damp situations.

*Sex. Syst. Crypt. fil. Nat. Syst. Polypodiaceae.*  
*Linn. Sp. Pl. 1551. (Polypodium). Griffiths, Med. Bot. 670.*

The rhizome is of a brown color, externally, with a feeble, but peculiar odor, and a sweet, bitter, astringent taste. It should always be used as fresh as possible. It is tonic and anthelmintic, but is not much used in this country. The dose, in substance, is from one to three drachms, twice a-day.

#### POWDER OF MALE FERN.

R. Male fern one ounce  
Gamboge  
Calomel, each, fourteen grains  
Scammony eighteen grains  
Mix, and divide into three powders. In tapeworm. *Cadet.*

#### BOLUS OF MALE FERN.

R. Male fern half an ounce  
Rhubarb two drachms  
Semen contra half an ounce  
Calomel one drachm  
Syrup of wormwood sufficient  
Mix, and form twelve boluses. *Spielmann.*

#### ETHEREAL EXTRACT OF MALE FERN.

R. Powdered male fern at will  
Exhaust it in a displacement apparatus, with ether, and distil off the ether. Dose, from fifteen to twenty grains, twice a-day.

#### PILLS OF EXTRACT OF MALE FERN.

R. Ethereal extract of male fern one scruple  
Powder of male fern twelve grains  
Conserve of roses sufficient  
Mix, and make twelve to sixteen pills.

To be taken in the evening at short intervals. *Peschier.*

#### OIL OF MALE FERN.

R. Powdered male fern four parts  
Alcohol eight parts  
Digest for twenty-four hours, decant, repeat the operation with five parts of alcohol, unite the liquid, distil off the alcohol, mix the residue with three parts of water, distil, and permit to rest, and collect the oil which sinks to the bottom, and wash it with pure water. *Magendie.*

#### PILLS OF OIL OF MALE FERN.

R. Oil of male fern twenty-four drops  
Powdered mallow root three scruples  
Fresh lard two scruples  
Mix, and divide into twenty-four pills. One-half to be taken in the evening, and the remainder next morning, and, an hour afterwards, a dose of castor oil. *Jourdan.*

#### ELECTUARY OF MALE FERN.

R. Ethereal extract of male fern half a drachm  
Honey of roses one ounce  
Mix. Half in the evening, the remainder the next morning. *Radius.*

#### JELLY WITH MALE FERN.

R. Male fern  
Corsica moss, each, six drachms  
Isinglass one drachm  
Water eight ounces  
Boil to three ounces, and strain; add  
Ethereal extract of male fern one scruple  
Syrup of mulberries one ounce  
To be taken in divided doses, during the day. *Phacbus.*

#### MIXTURE OF MALE FERN.

R. Male fern half an ounce  
Elecampane two drachms  
Wormwood a handful  
Semen contra one drachm  
Water sufficient  
to obtain a quart of decoction; add  
Vermifuge syrup one ounce  
Mix. *Bories.*  
R. Powdered male fern four drachms  
Balm water three ounces  
Syrup of gum one ounce  
Mix. To be taken in the evening, and, next morning, an ounce of castor oil. *Radius.*

#### TINCTURE OF MALE FERN BUDS.

R. Buds of male fern one part  
Sulphuric ether eight parts



Macerate for five days, and decant. Dose,  
from eight to thirty drops. Foy.

## FENICULUM.

## FENNEL.

There are several varieties, or species of Fennel, closely resembling each other, and having the same properties; that recognized as official by the U. S. Pharm., is the *F. vulgare*, a native of Europe, and often cultivated in the gardens in this country.

*Sex. Syst.* Pentand. digyn. *Nat. Syst.* Apiaceæ.

Linn. Sp. Pl. 377. Griffith, Med. Bot. 321.

The part used is the seeds. These are fragrant, with a sweetish, warm, pleasant, aromatic taste. They are employed as a carminative, and as corrective of other, and more unpleasant remedies. The dose of the powder, is from a scruple to half a drachm.

## COMPOUND POWDER OF FENNEL.

R. Powdered fennel seed  
    anise  
    lettuce  
    white poppy  
    benne, each,           one ounce  
                                    and a-half  
    sugar                   six ounces

Mix. Dose, a drachm, two or three times a-day. Said to increase the secretion of milk in nursing women.

Neimann.

## INFUSION OF FENNEL.

R. Fennel seed                   one drachm  
    Boiling water               one pint  
Infuse for two hours, and strain. Given in teaspoonful doses to infants, in windy colic, or used as an enema, for the same purpose. Taddei.

## FENNEL WATER.

R. Oil of fennel               half a fl. drachm  
    Carbonate of magnesia   half a drachm  
    Distilled water           two pints  
Rub the oil with the magnesia, gradually add the water, constantly rubbing, and filter. U. S. Ph.

## OIL OF FENNEL.

R. Bruised fennel seed       at will  
    Water                   sufficient  
Mix, and distil. Guibourt.

Dose, five to fifteen drops.

## FENNEL OINTMENT.

R. Oil of fennel               thirty drops  
    Lard                   four ounces  
Triturate together. Said to destroy lice in the hair. Neimann.

## FRASERA.

## AMERICAN COLUMBO.

The *Frasera carolinensis* is one of the tallest and most elegant of our native herbaceous plants, sometimes attaining a height of ten feet, with a pyramid of yellowish-white flowers, of three or four feet in length.

*Sex. Syst.* Tetrand. monog. *Nat. Syst.* Gentianaceæ.

Walter. Fl. Carol. 87. Griffith, Med. Bot. 462.

The root, which is the official portion, is large and succulent, and is usually dried in circular sections, like columbo. It has an odor in a dried state, but possesses a sweetish, bitter taste. It is a mild, bitter tonic, suited to all cases to which the simple bitters are applicable. The dose, in powder, is from thirty grains to a drachm.

## INFUSION OF AMERICAN COLUMBO.

R. American columbo, bruised, one ounce  
    Boiling water               one pint  
Infuse for an hour, and strain. Wood.

Dose, one to two fl. ounces, several times a-day.

## TINCTURE OF AMERICAN COLUMBO.

R. Contused American columbo one ounce  
    Diluted alcohol           one pint  
Macerate for fourteen days, and filter.  
Dose, one to three drachms. Dunglison.

## FULIGO.

## SOOT.

The soot of wood is of a black color, has a peculiar smell, and a bitter, disagreeable, empyreumatic taste. It owes its properties to the presence of pyrogenous oil or resin and creasote, but also contains various salts, united to acetic acid. It was formerly official, but gradually was relinquished in regular practice; but, of late years, has again attracted attention, and has been employed with success in a variety of diseases.

## PILLS OF SOOT.

R. Extract of soot               half a drachm  
    Comp. galbanum pill       one scruple  
    Oil of valerian           ten minims  
Mix well, and form twelve pills, two, thrice a-day, in hysteria. Neligan.  
R. Soot                   one drachm and a-half  
    Carbonate of ammonia   half a drachm  
    Tar                   sufficient  
Mix, and make five-grain pills. Ferriar.

## DECOCTION OF SOOT.

R. Soot                   two handfuls  
    Water               two pints  
Boil, for half an hour, and strain. As a lotion, in herpetic, psoric, and venereal ulcers. Soubeiran.



## TINCTURE OF SOOT.

R. Soot one part  
Diluted alcohol eight parts  
Macerate for eight days, and filter. Dose,  
from thirty to sixty drops, as an antispas-  
modic, in hysteria, &c. *Guibourt.*

## MIXTURE OF SOOT.

R. Soot half an ounce  
Carbonate of potassa one ounce and a-half  
Carbonate of ammonia two drachms  
Elder-flower water nine fl. ounces  
Digest for some time by a gentle heat, and  
filter. Dose, thirty to sixty drops, several  
times a-day. In dyspepsia, suppressed  
cutaneous affections, &c. *Dunglison.*

R. Hickory ashes one quart  
Soot six ounces  
Boiling water one gallon  
Mix, and digest for twenty-four hours,  
frequently stirring, and decant. A tea-  
cupful, three times a-day, used in dyspep-  
sia. *Ellis.*

## SOOT MOUTH WASH.

R. Extract of soot one part  
Vinegar twelve parts  
Mix. A few drops, in a glass of tepid  
water, in aphthous sore mouth. *Soubeiran.*

## OIL OF SOOT.

R. Soot four ounces  
Olive oil twelve ounces  
Boil, and strain. As an application to  
ulcers in the mouth, or on the genitals.  
*Radius.*

## SOOT OINTMENT.

R. Soot one part  
Lard four parts  
Triturate together. As an application in  
tinea, scrofulous sores, &c. *Soubeiran.*

## COMPOUND SOOT OINTMENT.

R. Soot  
Lard, each, half an ounce  
Extract of belladonna one drachm  
Triturate together. As an application to  
cancerous sores. *Radius.*

R. Soot an ounce and a-half  
Lard four ounces  
Sulphate of zinc six drachms  
Triturate together. In cases of tinea.  
*Bories.*

R. Soot  
Powdered Peruvian bark, each, half an ounce  
Powdered charcoal  
sulphur, each, one ounce  
Lard sufficient  
to make an ointment. In tinea.

*Radius.*

## CATAPLASM OF SOOT.

R. Soot two ounces  
White of eggs six  
Mix. As an application to tinea, herpes,  
and scrofulous ulcers. *St. Marie.*

## INJECTION OF SOOT.

R. Decoction of soot one pint  
Alum half an ounce  
Water six ounces  
Mix. Advised as an injection, in leucor-  
rhœa. *Soubeiran.*

## FULIGOKALI.

## FULIGOKALI.

R. Potassa twenty parts  
Shining soot one hundred parts  
Water sufficient  
Boil for an hour, cool, dilute with water,  
filter, evaporate to dryness, keep in well-  
closed bottles. *Deschamps.*

Used, both internally and externally, in chro-  
nic cutaneous disorders. Dose, two to three  
grains, several times a-day.

## SULPHURETTED FULIGOKALI.

R. Fuligokali sixty parts  
Potassa fourteen parts  
Sulphur five parts  
Dissolve the sulphur and potassa, add the  
fuligokali, evaporate to dryness.

## PILLS OF SULPHURETTED FULIGOKALI.

R. Sulphuretted fuligokali five drachms  
Starch two drachms and a-half  
Tragacanth eight grains  
Syrup sufficient  
Mix, and make one hundred pills, and  
cover them with two or three coats of tra-  
gacanth. *Deschamps.*

## OINTMENT OF FULIGOKALI.

R. Fuligokali sixteen to thirty grains  
Lard one ounce  
Triturate together. Used as a deterative,  
resolvent, and somewhat stimulant appli-  
cation, in obstinate cutaneous affections.  
*Gibert.*



## G

## GALBANUM.

## GALBANUM.

This is the concrete juice of an unknown plant, but generally admitted to belong to the Apiaceæ. According to Don, and others, the Asiatic galbanum is the product of *Galbanum officinale*, a native of Persia.

Don, Trans. Linn. Soc. xvi. 603. Griffith, Med. Bot. 333.

Galbanum is in whitish or yellowish tears, or in brownish-yellow, or brownish masses, formed of these tears agglutinated, and containing various impurities. The odor is peculiar, and somewhat balsamic; the taste is hot, bitter, and acrid. It is stimulant, expectorant, and antispasmodic, in doses of ten to twenty grains, and also used externally to indolent swellings, &c.

## PURIFIED GALBANUM.

R. Galbanum one part  
Alcohol four parts

Dissolve by means of a water-bath, strain, treat the residue with a fresh quantity of alcohol, unite the two solutions, distil off the alcohol, and evaporate the residue.

*Guibourt.*

## COMPOUND GALBANUM PILLS.

R. Galbanum  
Myrrh, each, one ounce and a-half  
Assafetida half an ounce  
Syrup sufficient

Beat into mass, and divide into four hundred and eighty pills. *U. S. Ph.*

Dose, ten to twenty grains, in chlorosis and hysteria.

R. Galbanum two drachms  
Oil of savine four to twelve drops  
Malate of iron one drachm and a-half  
Powdered valerian sufficient

Beat into mass, and divide into one hundred and twenty pills. Two, three times a-day, in amenorrhœa, chlorosis, &c.

*Phœbus.*

## GALBANUM MIXTURE.

R. Galbanum two drachms  
Vinegar of squills  
Fennel water, each, two fl. ounces  
Spirit of mindererus two fl. drachms  
Nitric ether one fl. drachm  
Syrup of mallows half fl. ounce

Mix. Three or four spoonfuls a-day, in humid asthma. *Jahn.*

## TINCTURE OF GALBANUM.

R. Galbanum, in small pieces, two ounces

Proof spirit two pints  
Digest for seven days, and filter.

*Dub. Ph.*

Antispasmodic, &c. In doses of one to three fl. drachms.

## COMPOUND TINCTURE OF GALBANUM.

R. Tincture of galbanum  
wormwood

Spirit of rosemary equal parts  
Mix. Dose, forty to fifty drops. *Augustin.*

## GALBANUM PLASTER.

R. Lead plaster two pounds  
Galbanum half a pound  
Yellow wax four ounces

Melt the galbanum, add the plaster and wax, and melt the whole, and strain.

*Dub. Ph.*

## COMPOUND GALBANUM PLASTER.

R. Galbanum eight ounces  
Lead plaster three ounces  
Turpentine ten drachms  
Burgundy pitch three ounces

Melt the galbanum and turpentine together, strain, add the pitch, and then the lead plaster, melted; mix well. *U. S. Ph.*

A good application to chronic, scrofulous enlargements of the glands and joints.

R. Lead plaster ninety-six parts  
Ammoniac  
Galbanum

Vinegar, each, eight parts  
Powder of saffron three parts  
Melt together, and strain. *Lond. Ph.*

R. Galbanum and saffron  
plaster eight parts  
Camphor  
Petroleum  
Subcarbonate of ammonia, each,

one part

Melt together. In chronic rheumatism. *Augustin.*

R. Galbanum  
Turpentine, each, one part  
Melt, strain, and add a melted mixture of  
White wax four parts  
Spermaceti two parts  
Oil of almonds one part

Mix well. A milder application than any of the above. *Spielmann.*



## GALIUM VERUM.

## YELLOW LADIES BEDSTRAW.

A perennial, herbaceous plant, native of Europe, which is stated to have the power of coagulating milk, and also used to dye yellow.

*Sex. Syst.* Tetrand. monog. *Nat. Syst.* Rubiaceæ.

Linn. Sp. Pl. 155. Stokes, Bot. Mat. Med. i. 202.

The whole plant is used; it has an agreeable odor, and an astringent, acidulous, somewhat acrid taste. It was, at one time, much used in epilepsy and hysteria, and also as an application in cutaneous affections.

## INFUSION OF YELLOW LADIES BEDSTRAW.

R. Flowers of yellow ladies bedstraw two drachms  
Boiling water two pints  
Digest for an hour, and strain. This to be taken in a tepid state, in wineglassful doses, as a sedative and antispasmodic, in cerebral affections. *Chaussier.*

## WINE OF YELLOW LADIES BEDSTRAW.

R. Expressed juice of yellow ladies bedstraw six fl. ounces  
White wine one to two fl. ounces  
Mix. To be taken in epilepsy, as soon as indications of an attack manifest themselves. *Bories.*

## GALLA.

## GALLS.

Galls are morbid excrescences on *Quercus infectoria*, caused by the puncture of a hymenopterous insect, the *Diplolepis gallæ tinctoriæ*. They come from the Levant ports, and present several varieties, the best of which are dark colored. They are round, hard, solid, brittle, inodorous, and have a bitter, very astringent taste. They contain tannic and gallic acids. They are powerfully astringent. They are more used as external applications, than as internal remedies. Dose, ten to thirty grains, several times a-day.

## COMPOUND POWDER OF GALLS.

R. Powdered galls  
kino, each, half an ounce  
alum two drachms  
Mix. As a styptic. *Radius.*

## INFUSION OF GALLS.

R. Powdered galls one to two drachms  
Boiling water one pint  
Digest for half an hour, and strain. Used for injections, fomentations, gargles, &c. *Taddei.*

## COMPOUND INFUSION OF GALLS.

R. Infusion of galls four fl. ounces  
Prepared chalk half an ounce  
Tincture of opium half fl. drachm

Powdered gum Arabic one drachm  
Mix. Dose, a tablespoonful every two hours. In diarrhœa. *Ellis.*

## GARGLE OF GALLS.

R. Infusion of galls seven fl. ounces  
Alcohol one fl. ounce  
Mix. As a gargle in relaxation of uvula and soft palate. *Ellis.*

## LOTION OF GALLS.

R. Claret wine one pint  
Common salt one drachm  
Sulphate of iron two drachms  
Boil for a few minutes, and add  
Oxide of copper one drachm  
Boil for two minutes. Add  
Powdered galls two drachms  
This is used to color the hair; it is to be washed with this fluid, dried with a warm cloth, and then washed with common water. *Laforest.*

## SYRUP OF GALLS.

R. Bruised galls two drachms  
Brandy eight fl. ounces  
Mix. Burn out the spirit, and add  
Powdered cinnamon mace, each, two drachms  
While hot, add sugar by burning it over the inflamed brandy, till a syrup is formed. Decant. Dose, a tablespoonful every two hours. In obstinate chronic diarrhœa. *Parrish.*

## TINCTURE OF GALLS.

R. Bruised galls four ounces  
Diluted alcohol two pints  
Macerate for fourteen days, express, and filter. *U. S. Ph.*

A powerful astringent, in doses of one to three fl. drachms.

R. Bruised galls one pound  
Water two pints  
Macerate twenty-four hours, and add  
Alcohol two pints  
Eau de Cologne six fl. ounces  
Filter. As an injection, diluted with water, in leucorrhœa and gonorrhœa. *Gibert.*

## EXTRACT OF GALLS.

R. Galls, in coarse powder at will  
Macerate in water sufficient to cover them for twenty-four hours; then introduce the moist mass into a displacer, and act on it with water, until the galls are nearly exhausted; evaporate the infusion to dryness, on a water-bath.

## OINTMENT OF EXTRACT OF GALLS.

R. Extract of galls one drachm  
Lard one ounce



Triturate the extract with a little water, until it is reduced to a uniform syrupy consistence, then add the lard, and mix thoroughly. *D. S. Jones.*

## OINTMENT OF GALLS.

R. Galls, in very fine powder, one ounce  
Lard seven ounces  
Mix. As an application to piles, pro-  
lapsus ani, and also to indolent ulcers.  
*U. S. Ph.*

## COMPOUND OINTMENT OF GALLS.

R. Galls, in very fine powder, two drachms  
Powdered opium half a drachm  
Lard two ounces  
Mix well. *Lond. Ph.*  
As an application to irritable piles, &c.

R. Powdered galls  
Tincture of opium, each, two parts  
Camphor one part  
Lard eight parts  
Triturate well together. As an application  
to painful piles. *Foy.*

R. Finely powdered galls one scruple  
Simple cerate one drachm  
Essence of lemon twenty drops  
Lead water  
Laudanum, each, forty drops  
Rub well together. As an application to  
hemorrhoids, after subsidence of inflamma-  
tion. *Dewees.*

## GAMBOGIA.

## GAMBOGE.

This is the concrete juice of several species of East Indian trees, probably belonging to the genus *hebradendron*.

*Sex. Syst.* Polyand. monog. *Nat. Syst.* Clusiaceæ.

Graham, *Comp. Bot. Mag.* ii. 199. Griffith, *Med. Bot.* 152.

Gamboge is a gum resin, of a yellow-orange color, opaque, brittle, almost inodorous, of an insipid taste at first, but soon becoming acrid. It is a powerful hydragogue and drastic purgative, but apt to produce nausea and vomiting. Dose, from one to six grains, in pill, emulsion, or alkaline solution.

## COMPOUND POWDER OF GAMBOGE.

R. Bitartrate of potassa two drachms  
Powdered jalap one drachm  
gamboge six grains  
Mix, and divide into six powders. One to  
be given every two or three hours, in syrup,  
as a hydragogue purgative. *Ellis.*

R. Powdered gamboge two grains  
sulphate of iron six grains  
sugar one scruple

Oil of peppermint three drops  
Mix. To be taken twice a-day, against  
tape-worm. *Vogt.*

## COMPOUND PILLS OF GAMBOGE.

R. Powdered gamboge one drachm  
aloes one drachm and a-half  
ginger half a drachm  
Soap two drachms  
Beat into mass. *Lond. Ph.*  
Dose, ten to twelve grains, as a purgative.

R. Powdered gamboge ten grains  
guaiaicum  
Blanched almonds, each, one drachm  
Syrup sufficient  
Beat together, and form twenty-eight pills.  
Two to four, as a gentle purgative.  
*Saunders.*

## ALKALINE SOLUTION OF GAMBOGE.

R. Powdered gamboge one drachm  
Carbonate of potassa three drachms  
Water six drachms  
Dissolve. Dose, fifteen drops, every three  
hours, as a hydragogue in dropsies.  
*Van Mons.*

## GAMBOGE MIXTURE.

R. Powdered gamboge four grains  
Spirit of nitric ether one fl. drachm  
Tincture of senna two fl. drachms  
Mint water  
Syrup of buckthorn, each, half fl. ounce  
Mix. As a hydragogue purgative, in  
dropsy, especially in hydrothorax.  
*Ferriar.*

R. Gamboge two scruples  
Tartrate of potassa one ounce  
White sugar two drachms  
Water six ounces  
Make a solution. As a purgative, in as-  
cites. A tablespoonful, every two or  
three hours, until it operates. *Dewees.*

## GAMBOGE AND ELATINE MIXTURE.

R. Gamboge four grains  
Elatine half a grain  
Sweet spirit of nitre one ounce  
Water four ounces  
Mix. As a hydragogue purgative, in as-  
cites. A tablespoonful, to be taken every  
two or three hours, until free purgation is  
induced. *Dewees.*

## ALKALINE TINCTURE OF GAMBOGE.

R. Gamboge one part  
Carbonate of potassa two parts  
Diluted alcohol twenty-four parts  
Digest for some days, and filter. *Scherf.*



AMMONIACAL TINCTURE OF GAMBOGE.  
 R. Gamboge thirty-six grains  
 Spirit of ammonia four fl. ounces  
 Digest for eight days, and filter. Dose,  
 one to two fl. drachms. *Swediaur.*

## GAULTHERIA.

## PARTRIDGE BERRY.

The leaves of *Gaultheria procumbens*; a small, indigenous, evergreen plant, found in most parts of the United States, in dry, sandy situations.

*Sex. Syst.* Decand. monog. *Nat. Syst.* Ericaceae.

Kalm. Amoen. Acad. iii. 14. Griffith, Med. Bot. 424.

These leaves have a peculiar, aromatic taste and odor, with a little astringency. They are stimulant, and somewhat tonic and astringent, and are much used in the form of tea, in bowel diseases, and as an emmenagogue.

## OIL OF PARTRIDGE BERRY.

R. Leaves of partridge berry at will  
 Water sufficient  
 Distil, and collect the oil that passes. *Coxe.*

Principally used to flavor mixtures, &c. In large doses, is poisonous. Dissolved in diluted alcohol, is in common use as a carminative and stimulant.

## PARTRIDGE BERRY WATER.

R. Oil of partridge berry sixteen drops  
 Carbonate of magnesia one drachm  
 Water one pint  
 Triturate the oil, magnesia, and a little of the water together, then add the rest of the water, and filter.

## GENTIANA.

## GENTIAN.

The roots of several species of gentian are used in medicine; but that which is most generally recognized as officinal, is *Gentiana lutea*, a tall, perennial plant, native of the mountains of Europe.

*Sex. Syst.* Pentand. digyn. *Nat. Syst.* Gentianaceae.

Linn. Sp. Pl. 329. Griffith, Med. Bot. 460.

The root is of a grayish brown externally, and yellowish within; it has a slightly sweetish, but extremely bitter taste, and a faint, but peculiar odor. It is a pure and simple bitter tonic, with no astringency. The dose of the powder is from ten to forty grains, but is seldom given in this form.

## COMPOUND POWDER OF GENTIAN.

R. Powdered gentian  
 cascarilla  
 orange peel, each,  
 one drachm

Peppermint sugar three drachms  
 Mix. Dose, thirty grains, several times a-day, as a stomachic. *Augustin.*

## EXTRACT OF GENTIAN.

R. Gentian, in coarse powder, one pound  
 Water sufficient  
 Exhaust the powder by means of displacement. Heat the filtered liquid, strain, and evaporate to proper consistence. *U. S. Ph.*

Dose, ten to thirty grains, as a tonic; also as a basis for various metallic pills.

## COMPOUND GENTIAN PILLS.

R. Powdered aloes one drachm  
 Extract of gentian half a drachm  
 Oil of caraway ten drops  
 Syrup sufficient  
 Beat into mass, and form pills of four grains. Two or three for a dose, as a purgative. *Ellis.*

R. Extract of gentian  
 Powdered rhubarb  
 Soap equal parts  
 Water sufficient  
 Beat into mass, and form four-grain pills. Tonic and laxative. *Fulda Dispen.*

## PILLS OF GENTIAN AND AMMONIA.

R. Extract of gentian  
 Carbonate of ammonia, each,  
 Mix, and make twenty-four pills. Two to be taken twice or thrice a-day. For heart-burn in gouty habits. *Steph. and Church.*

## INFUSION OF GENTIAN.

R. Contused gentian two drachms  
 Boiling water one pint  
 Infuse for twelve hours, and strain. *Beral.*

## COMPOUND INFUSION OF GENTIAN.

R. Bruised gentian half an ounce  
 orange peel  
 coriander, each, one drachm  
 Diluted alcohol four fl. ounces  
 Water twelve fl. ounces  
 Pour on the alcohol, and, in three hours, add the water, macerate for twelve hours, and strain. *U. S. Ph.*

Dose, a fl. ounce, as a tonic.

R. Bruised gentian  
 calamus, each, one ounce  
 Centaury six drachms  
 Rosemary two drachms  
 Boiling water four pints  
 Infuse for twelve hours, and strain. Used as an emmenagogue, in doses of three ounces, twice a-day. *Spielmann.*



## INFUSION OF GENTIAN WITH RHUBARB.

R. Bruised gentian one drachm  
 rhubarb two drachms  
 Boiling water twelve fl. ounces  
 Macerate for an hour, strain, and add  
 Carbonate of ammonia two scruples  
 Dose, one to two fl. ounces, in dyspepsia,  
 chronic rheumatism, &c.

*Steph. and Church.*

## MIXTURE OF GENTIAN.

R. Carbonate of magnesia one drachm  
 Infusion of gentian six fl. ounces  
 Mix. A wineglassful, three times a-day,  
 as an antilithic, where uric acid abounds.

*Brande.*

R. Extract of gentian two drachms  
 Carbonate of potassa two scruples  
 Mint water five fl. ounces  
 Tincture of orange peel five fl. drachms  
 Sugar one ounce

Mix. A spoonful, three times a-day, as a  
 stomachic and carminative.

*Berends.*

## WINE OF GENTIAN.

R. Gentian, coarsely  
 powdered, half an ounce  
 Yellow bark, coarsely  
 powdered, one ounce  
 Orange peel two drachms  
 Canella, in coarse powder, one drachm  
 Proof spirit four fl. ounces and a-half  
 Sherry wine forty-six fl. ounces  
 Digest the root and barks in the spirit,  
 for twenty-four hours, add wine, digest for  
 seven days, express, strain, and filter.

*Ed. Ph.*

A stomachic bitter, in doses of four to eight fl.  
 drachms.

## SYRUP OF GENTIAN.

R. Gentian, in coarse  
 powder, forty-eight parts  
 Cold water sufficient  
 Syrup fifteen hundred parts  
 Put the gentian in a displacement appa-  
 ratus, and make a strong infusion, filter  
 this, and add to boiling syrup, of such a  
 consistence, as, after the addition, it will  
 make 30°.

*Tauvel.*

## SYRUP OF EXTRACT OF GENTIAN.

R. Aqueous extract of gen-  
 tian two drachms  
 Water eight fl. ounces  
 Sugar sixteen ounces (av.)  
 Dissolve the extract in the water, add the  
 sugar, and form a syrup with a gentle heat,  
 and strain.

*W. Procter.*

## TINCTURE OF GENTIAN.

R. Gentian one part

Diluted alcohol four parts  
 Macerate for six days, and filter. Dose,  
 one to two fl. drachms.

*Cottureau.*

## ACIDULATED TINCTURE OF GENTIAN.

R. Compound tincture of  
 gentian four fl. ounces  
 Elixir of vitriol half fl. drachm  
 Mix. A teaspoonful, in sugar and water,  
 three or four times a-day.

*Ellis.*

## COMPOUND TINCTURE OF GENTIAN.

R. Bruised gentian two ounces  
 Orange peel one ounce  
 Bruised cardamoms half an ounce  
 Diluted alcohol two pints  
 Macerate for fourteen days, express, and  
 filter.

*U. S. Ph.*

An excellent bitter tonic, in doses of one fl.  
 drachm.

## AMMONIACAL TINCTURE OF GENTIAN.

R. Bruised gentian one ounce  
 Carbonate of ammonia two drachms  
 Diluted alcohol two pints  
 Macerate the root in the alcohol, for eight  
 days, filter, and add the ammonia. Dose,  
 one to four fl. drachms, in scrofula.

*Taddei.*

## ALKALINE TINCTURE OF GENTIAN.

R. Bruised gentian one ounce  
 Carbonate of soda three drachms  
 Diluted alcohol two pints  
 Macerate for eight days, and filter. Used  
 in same cases, and same doses, as the for-  
 mer.

*Guibourt.*

## MIXTURE OF TINCTURE OF GENTIAN.

R. Tincture of gen-  
 tian one fl. ounce and a-half  
 Tincture of rhubarb

cinnamon half fl. ounce

opium ten drops

Mix. A dessertspoonful, every hour, in  
 diarrhoea.

*Foy.*

R. Extract of gentian  
 cascarrilla, each,  
 two drachms

Peppermint water four fl. ounces

Mix. In spoonful doses, as a tonic and  
 stomachic.

*St. Marie.*

GENTIAN AND SULPHURIC ACID MIX-  
TURE.

R. Compound infusion of  
 gentian five fl. ounces  
 Compound tincture of  
 gentian one fl. ounce  
 Diluted sulphuric acid one fl. drachm  
 Mix. A tablespoonful, three times a-day,  
 in dyspepsia.

*Brande.*



## GENTIANINA.

## GENTIANIN.

R. Powdered gentian at will  
 Macerate in cold ether, and repeat till exhausted; unite the tinctures, and distil; treat the residue several times with cold alcohol, distil the tinctures, evaporate the residue to dryness; mix it with an excess of magnesia, treat with sulphuric ether, then with oxalic acid, then again with the ether, and distil. *Magendie.*

Dose, one to two grains.

## PILLS OF GENTIANIN.

R. Gentianin five grains  
 Conserve of roses  
 Powdered liquorice, each, sufficient  
 Mix, and make six pills.

## SYRUP OF GENTIANIN.

R. Gentianin sixteen grains  
 Syrup one pound  
 Dissolve, and boil for a short time. Dose, half an ounce to an ounce. *Bories.*

## TINCTURE OF GENTIANIN.

R. Gentianin five grains  
 Alcohol one ounce  
 Mix. Dose, from one to four drachms. *Cadet.*

## GERANIUM.

## CRANESBILL.

X This is the root of *Geranium maculatum*, a native, perennial plant, found, in most parts of the country, in woods and shady places.

*Sex. Syst.* Monadelph. decand. *Nat. Syst.* Geraniaceæ.

Linn. Sp. Pl. 955. Griffith, Med. Bot. 209.

The root is in rough pieces, of a dark brown color externally, and flesh-colored within. It is inodorous; the taste is astringent, but not bitter. The dose is from ten to thirty grains, as an astringent, in all cases in which kino or rhatany are found useful.

## DECOCTION OF GERANIUM.

R. Geranium one ounce  
 Water one pint and a-half  
 Boil down to a pint. Dose, from one to two fl. ounces. *Wood.*

## EXTRACT OF GERANIUM.

R. Bruised geranium root one pound  
 Water one gallon  
 Boil down to one half, strain, and evaporate to due consistence. This extract is very similar to Rhatany, and may be given in the same cases, and in the same doses.

## GEOFFROYA INERMIS.

## CABBAGE TREE BARK.

The Cabbage Tree is a native of the West India Islands; it is a large tree, having a smooth, gray bark.

*Sex. Syst.* Diadelph. decand. *Nat. Syst.* Fabaceæ.

Linn. Sp. Pl. 1043. Griffith, Med. Bot. 247. (Andira.)

The bark is in long, fibrous pieces, of a brownish ash color externally, and yellowish within; it has a resinous fracture, an unpleasant odor, and a sweetish, mucilaginous, bitterish taste. It is a powerful vermifuge, in doses of a scruple to half a drachm. The best form of administration is the syrup.

## DECOCTION OF CABBAGE TREE BARK.

R. Cabbage tree bark one ounce  
 Water two pints  
 Boil down to one pint, and strain. Dose, one to two fl. ounces. *Ellis.*

## EXTRACT OF CABBAGE TREE BARK.

R. Cabbage tree bark one part  
 Water eight parts  
 Boil for a quarter of an hour, strain, add four parts of water, boil, and strain; evaporate the united decoctions to a proper consistence. Dose, three grains. *Belg. Ph.*

## MIXTURE OF CABBAGE TREE BARK.

R. Cabbage tree bark half an ounce  
 Water six ounces  
 Boil for half an hour, strain, and add  
 Tincture of cabbage tree bark one ounce  
 Syrup of orange peel half an ounce  
 A tablespoonful, morning and evening. *Niemann.*

## GEUM.

## AVENS.

## 1. GEUM RIVALE.

## WATER AVENS.

This plant is a native of Europe and the United States, in wet, boggy situations.

*Sex. Syst.* Icosand. polyg. *Nat. Syst.* Rosaceæ.

Linn. Sp. Pl. 717. English Bot. 1. 106.

Root reddish or purplish, hard and brittle, inodorous, of an astringent, bitterish taste. It is tonic and astringent, and has been used advantageously in passive hemorrhages, leucorrhœa, &c. The dose, in powder, is from a scruple to a drachm, three times a-day.

## DECOCTION OF WATER AVENS.

R. Water avens root one ounce



Water one pint and a-half  
Boil down to a pint, and strain, to be given  
in doses of one or two fl. ounces.

## 2. GEUM VIRGINIANUM. WHITE AVENS.

Also a native of the United States, growing  
in woods and along streams. Flower small and  
white.

Linn. Sp. Pl. 719. Griffith, Med. Bot. 279.

The root is of a brown color, somewhat aromatic, and of a bitterish, astringent taste. It has the same properties as the *G. rivale*, and is given in the same cases and doses.

## 3. GEUM URBANUM. AVENS.

A native of Europe, growing in woods and waste places. They are small, and of a yellow color.

Linn. Sp. Pl. 716. Eng. Bot. t. 1400.

The root consists of a short caudex with many long fibres. When fresh, it has somewhat the odor of cloves, but when dry, is almost inodorous. The taste is bitter and astringent. Like the other species, it is tonic and astringent.

### COMPOUND POWDER OF AVENS.

R. Powdered avens root  
gum Arabic, each, three  
drachms  
kino  
cinnamon, each, one drachm  
sugar half an ounce

Mix. A teaspoonful, every two or three hours, in diarrhoea. *Vogel.*

### MIXTURE OF AVENS.

R. Bruised avens one ounce  
Water twelve fl. ounces  
Boil to eight ounces, and to the strained cold decoction, add  
Extract of avens two drachms  
Sulphuric ether one drachm  
Mix. Two spoonfuls, every two hours.  
As a febrifuge. *Radius.*

## GILLENIA. INDIAN PHYSIC.

This is the root of *Gillenia trifoliata*, a native herbaceous plant, found in most parts of the country, in shady, and rather marsh situations.

*Sex. Syst.* Icosand. pentag. *Nat. Syst.* Rosaceæ.

Mænoch. Sp. Pl. 286. Griffith, Med. Bot. 282.

The roots are about as thick as a quill, wrinkled longitudinally, and irregularly undulated; a light brown color, and consisting of a thick cortical portion, and a slender ligneous centre. The

bark is of a disagreeable, bitter taste. It is a mild and certain emetic, and forms a good substitute for ipecacuanha. The dose, from twenty to thirty grains, to be repeated in half an hour, if it does not produce the desired effect.

## GLYCERINUM.

### GLYCERIN

Is a fatty body, which may be obtained by the saponification of olive oil, in making lead plaster. The water in which the plaster is malaxated, dissolves the glycerin, having a little lead in solution. This water is to be evaporated till of a syrupy consistence, sulphuretted hydrogen passed through to separate the lead, heated, to expel any excess of gas, digested with animal charcoal, filtered, and evaporated in vacuo, at a temperature of 150° F. In a pure state it is a nearly colorless, viscid fluid, and is uncrystallizable.

It has been used with success in lotions, poultices, baths, &c., to render them emollient and soothing; in diseases of the skin, as pityriasis, impetigo, chapped hands, nipples, face, &c.

### OINTMENT OF GLYCERIN.

R. Glycerin one fl. drachm  
Borax ten grains  
Rose water seven fl. drachms  
Mix. Used as a lotion in lichen, three times a-day; also in chapped hands.

*Clymer.*

## GLYCYRRHIZA.

### LIQUORICE ROOT.

Several species of this genus have sweet and demulcent roots, but the only officinal species is the *G. glabra*, a native of the south of Europe, and of some parts of Asia and Africa, and also successfully cultivated in more northern regions.

*Sex. Syst.* Diadelph. decand. *Nat. Syst.* Fabaceæ.

Linn. Sp. Pl. 1046. Griffith, Med. Bot. 236.

The root is in long pieces of various thicknesses, of a grayish-brown color externally, and yellowish within; it is inodorous, and of a sweet, mucilaginous taste, with a slight degree of acidity. It is an excellent demulcent, and is much used in affections of the mucous membranes, and as an addition to the more irritating or nauseous remedies, to abate their acrimony or conceal their taste.

### EXTRACT OF LIQUORICE.

R. Liquorice root,  
sliced, two pounds and a-half  
Boiling distilled water two gallons  
Macerate for twenty-four hours, boil down to a gallon, strain, evaporate to proper consistence. *Lond. Ph.*

### REFINED LIQUORICE.

R. Crude liquorice at will  
Dissolve in water, strain, and evaporate.



These two articles are useful demulcents, and additions to cough mixtures.

*U. S. Disp.*

#### LIQUORICE PASTE.

R. Extract of liquorice one pound  
Cold water four pints  
Dissolve, strain, and add  
Gum Arabic two pounds  
Sugar one pound  
Evaporate to proper consistence. *Soubeiran.*

R. Gum Arabic two pounds  
Extract of liquorice nine ounces  
Sugar one pound  
Water sufficient  
Evaporate to consistence of syrup, and add

Powdered orris root  
elecampane, each, half an ounce

Continue to evaporate, then add  
Oil of aniseed two scruples  
*Spielmann.*

#### LOZENGES OF LIQUORICE.

R. Extract of liquorice  
Gum Arabic, each, six ounces  
Sugar one pound  
Dissolve in sufficiency of boiling water, evaporate on water-bath, to proper consistence for lozenges. *Ed. Ph.*

#### LOZENGES OF LIQUORICE AND OPIUM.

R. Powdered opium half an ounce  
liquorice  
gum Arabic  
sugar, each, ten ounces  
Oil of anise two fl. drachms  
Mix the powder well, add the oil of anise, form mass with water, and divide into lozenges of six grains. *U. S. Ph.*

Each lozenge contains one-tenth of a grain of opium.

R. Powdered liquorice  
gum Arabic  
sugar, each, two drachms  
opium six grains  
Oil of anise, four drops  
Mix, and with sufficient water form mass, and divide into sixty lozenges. These are known as Dr. Wistar's cough lozenges. *Ellis.*

Both the above are demulcent and anodyne, and are well suited to allay cough, where opium is permissible.

#### MIXTURE OF LIQUORICE.

R. Powdered liquorice one drachm  
Chamomile water one fl. ounce

Syrup of mallows four fl. ounces  
Mix. In spoonful doses, to allay cough. *Foy.*

R. Powdered liquorice  
gum Arabic, each, two drachms  
Boiling water four fl. ounces  
Mix, and dissolve, then add  
Spirit of nitric ether  
Antimonial wine, each, one fl. drachm  
Tincture of opium forty to fifty drops  
A tablespoonful occasionally, in catarrh. *Ellis.*

R. Powdered liquorice  
gum Arabic, each, two drachms  
Boiling water four fl. ounces  
Dissolve, and add  
Antimonial wine two fl. drachms  
Tincture of opium twenty minims  
Mix. A tablespoonful, occasionally. *U. S. Disp.*

These two mixtures are well known under the name of *Brown mixture*.

#### SYRUP OF LIQUORICE.

R. Liquorice root two ounces  
Maidenhair one ounce  
Hyssop half an ounce  
Water two pints  
Infuse for twenty-four hours, boil to one-half, and add to strained decoction  
Honey eight ounces  
Sugar sixteen ounces  
Rose water four fl. ounces  
Make a syrup. As a cough mixture, in spoonful doses. *Wirtem. Ph.*

## GOSSYPIMUM.

### COTTON

Is the down or hair attached to the seeds of *Gossypium herbaceum*, and other species of the genus. It consists of fine, white filaments, having neither odor nor taste, and is insoluble in water, alcohol, ether, the oils, &c., but is soluble in strong alkaline solutions.

*Ser. Syst.* Monadelph. polyand. *Nat. Syst.* Malvaceæ.

*Linn. Sp. Pl.* 845. *Griffith, Med. Bot.* 162.

It is principally used for the fabrication of cloth, but is also employed in medicine, as an application to recent burns and scalds, to erysipelas, and to parts affected with rheumatism. It should be applied in the form of thin sheets. The seeds furnish much oil, and the root is said to be an active emmenagogue.

#### DECOCTION OF COTTON ROOT.

R. Inner part of cotton root four ounces  
Water one quart



Boil down to a pint. Said to be as effectual as ergot in promoting uterine contraction. Dose, a wineglassful, every twenty or thirty minutes. *Bouchelle.*

ETHEREAL SOLUTION OF COTTON.  
(COLLODION.)

This preparation, which is a solution of gun-cotton in ether, is used as a dressing to wounds; by its evaporation, it unites them closely, preserves them from the action of the air, is impervious to moisture; being transparent, it permits the progress of the wound to be examined, &c.

R. Nitric acid (1.45)  
Sulphuric acid (commercial), each, one fl. ounce  
Cleaned and bleached cotton two drachms  
Saturate the cotton thoroughly, with the acids, and macerate for twelve hours. Wash the cotton; dry rapidly, by artificial heat, in the shade, and dissolve it in Sulphuric ether (official) one pint and a-half  
*E. Parrish. W. Livermore.*

R. Finely powdered nitrate of potash 40 parts, by weight  
Concentrated sulphuric acid 60 " " "  
Carded cotton 2 " " "

Mix the nitre with the sulphuric acid, in a porcelain vessel, then add the cotton, and agitate the mass for three minutes, by the aid of two glass rods. Wash the cotton, without first pressing it, in a large quantity of water, and when all acidity is removed (indicated by litmus paper), press firmly in a cloth. Pull it out into a loose mass, and dry it in a stove at a moderate heat

Prepared cotton 8 parts, by weight  
Rectified sulphuric ether 125 " " "  
Rectified alcohol 8 " " "

Put the cotton, with the ether, into a well-stoppered bottle, and shake the mixture for some minutes. Then add the alcohol by degrees, and continue to shake, until the whole of the liquid acquires a syrupy consistency. It may then be passed through a cloth, the residue strongly pressed, and the liquid kept in a well-secured bottle. *Miahle.*

GRANATUM.

POMEGRANATE.

The *Punica granatum*, a small tree, a native of, and cultivated in warm climates, and much esteemed for its subacid fruit, affords two arti-

cles employed medicinally, the rind of the fruit, and the bark of the root.

*Ser. Syst.* Icosand. monog. *Nat. Syst.* Myrtacæ.

Linn. Sp. Pl. 676. Griffith, Med. Bot. 294.

The rind, as found in the shops, is in fragments of a dry, hard, brittle consistence; of a yellowish, or reddish-brown color, inodorous, and of an astringent, somewhat bitter taste. The bark of the root is in small pieces, of an ash-gray or yellowish color, externally, yellow within; brittle, with scarcely any odor, and astringent, but not bitter taste. The rind is a good astringent, in doses of ten to thirty grains; and the bark of the root, an efficient vermifuge, especially in a fresh, or recently dried state.

DECOCTION OF POMEGRANATE RIND.

R. Pomegranate rind two ounces  
Distilled water a pint and a-half  
Boil down to a pint, and strain. *Lond. Ph.*  
Dose, one fl. ounce. As an astringent.

GARGLE OF POMEGRANATE RIND.

R. Pomegranate rind half an ounce  
Red roses two drachms  
Boiling water six fl. ounces  
Infuse, strain, and add Clarified honey one fl. ounce  
Alum two drachms  
Used as gargle, in inflammation of fauces and throat. *Ellis.*

INJECTION OF POMEGRANATE RIND.

R. Pomegranate rind one part  
Water thirty-two parts  
Boil away one-half. *Beral.*

DECOCTION OF POMEGRANATE ROOT.

R. Bark of pomegranate root two ounces  
Water one pint and a-half  
Macerate for two hours, and then boil to a pint. To be taken in three doses. As a vermifuge, in cases of tænia. *Foy.*

EXTRACT OF POMEGRANATE ROOT.

R. Bark of pomegranate root at will  
Diluted alcohol sufficient  
Macerate for a week, distil off the alcohol, and evaporate the residue to proper consistence. *Soubeiran.*

ELECTUARY OF EXTRACT OF POMEGRANATE ROOT.

R. Extract of pomegranate root six drachms  
Gum tragacanth sufficient  
Lemon juice two fl. ounces  
Linden water three fl. ounces  
Mix. One-half to be taken, and half an hour afterwards the remainder. *Radius.*

MIXTURE OF EXTRACT OF POMEGRANATE ROOT.

R. Extract of pomegranate root six drachms



Lemon juice  
Mint water  
Linden water, each, two fl. ounces  
Mix. One-fourth to be taken, every  
quarter of an hour. *Radius.*

#### BOLUS OF POMEGRANATE ROOT.

R. Powdered bark of pome-  
granate root one drachm  
Powdered red Peruvian  
bark two drachms  
Balsam copaiba sufficient  
Mix, and form twelve boluses. One,  
morning and evening, in chronic mucous  
discharges. *Radius.*

R. Powdered bark of pome-  
granate root one drachm  
Assafetida half a drachm  
Oil of croton four drops  
Syrup of ether sufficient  
Mix, and make fifteen boluses. Five  
a-day, in the treatment of tænia. *Foy.*

### GRATIOLA.

#### HEDGE HYSSOP.

Most of the species of this genus have active  
properties, but only one of them has been em-  
ployed in medicine; the *G. officinalis* of Europe;  
a native species, the *G. aurea*, is possessed of  
identical properties, which, in both cases, depend  
on the presence of *veratrina*.

*Sex. Syst.* Diand. monog. *Nat. Syst.* Scrophu-  
liaceæ.

Linn. Sp. Pl. 24. Griffith, Med. Bot. 518.

The whole plant is used, it is almost inodor-  
ous, but has a bitter, nauseous taste. It is a dras-  
tic cathartic and emetic, with some diuretic pro-  
perties. It has been stated that it forms the  
basis of the *Eau medicinale of Husson*. The  
dose of the powder is from ten to twenty  
grains.

#### COMPOUND POWDER OF HEDGE HYSSOP.

R. Powdered hedge hyssop two scruples  
Calomel five grains  
Assafetida half a drachm  
Oil of peppermint three drops  
Mix, and divide into two powders. One  
to be taken every three hours, in tænia.  
*Augustin.*

#### WINE OF HEDGE HYSSOP.

R. Dried herb of hedge hyssop one ounce  
Contused root of hedge  
hyssop half an ounce  
White wine sixteen fl. ounces  
Macerate for eight days, with a gentle  
heat, and strain. A teaspoonful, at bed-  
time, drinking after it half a pint of beef  
tea. If necessary, repeat next evening.  
*Perkins.*

Said to be the original recipe for the *Eau me-  
dicinale*.

### GUAIAECUM.

#### GUAIAECUM.

The *Guaiacum officinale*, a large tree, a native  
of the West Indies and some parts of South  
America, affords two medicinal products, the  
wood and resin, or concrete juice.

*Sex. Syst.* Décand. monog. *Nat. Syst.* Zygo-  
phyllaceæ.

Linn. Sp. Pl. 546. Griffith, Med. Bot. 203.

The wood is very hard, of a greenish brown  
color, of a bitterish, somewhat acrid taste, and of  
a faint smell, except when heated. The resin  
is of a greenish or dark olive color, of a feeble  
odor, and of an acrid taste. They both are  
stimulant and alterative, and have been much  
used in chronic rheumatism, gout, obstinate cuta-  
neous affections, amenorrhœa, &c. The dose of  
the resin is from ten to thirty grains.

#### DECOCTION OF GUAIAECUM WOOD.

R. Rasped guaiacum wood one ounce  
Water one pint and a-half  
Boil down to a pint. To be taken in  
divided doses, during the day.

*U. S. Dispens.*

#### COMPOUND DECOCTION OF GUAIAECUM WOOD.

R. Guaiacum shavings three ounces  
Raisins two ounces  
Sassafras root, rasped,  
Liquorice root, bruised, each, one ounce  
Water eight pints

Boil the guaiacum and raisins in the water,  
down to five pints, adding the other ingre-  
dients, towards close of operation. Strain.

*Ed. Ph.*

From one to two pints may be taken, during  
the day, in divided doses, as an alterative in  
chronic rheumatism and cutaneous diseases.

R. Rasped guaiacum wood one ounce  
sassafras half an ounce  
Contused liquorice root one drachm  
Water five pints

Boil for two hours, strain, and add  
Cinnamon water three ounces  
A wineglassful, several times a-day, as  
alterative, &c. *Augustin.*

R. Rasped guaiacum wood two drachms  
Water sufficient  
to obtain one pint of strained decoction.  
After an hour's boiling, add

Wine of colchicum seed two drachms  
A wineglassful every two hours, in chronic  
pains of the joints. *Radius.*

#### EXTRACT OF GUAIAECUM WOOD.

R. Powdered Guaiacum wood at will



Exhaust with boiling water, by means of a displacement apparatus, and evaporate.

*Van. Mons.*

#### OIL OF GUAIACUM WOOD.

R. Guaiacum wood at will  
Distil on a sand-bath, separate the oil, and rectify it. Dose, four or five drops, in gonorrhœa, also as a friction in rheumatism.  
*Taddei.*

#### COMPOUND OIL OF GUAIACUM.

R. Oil of guaiacum three ounces  
Balsam of Peru one ounce  
Ammoniated alcohol half an ounce  
Mix. In caries of the teeth, and as a dressing to malignant ulcers. *Swediaur.*

#### COMPOUND POWDER OF GUAIACUM.

R. Powdered resin of guaiacum two drachms  
Cremor tartar half an ounce  
Sugar six drachms  
Mix. Three teaspoonfuls a-day, in gout.  
*Phœbus.*

R. Powdered resin of guaiacum  
nitrate of potassa, each, one drachm  
ipecacuanha three grains  
opium two grains  
Mix, and divide into six powders. One, every three hours; as a stimulating diaphoretic in acute rheumatism, after reduction of inflammation.  
*Ellis.*

#### PILLS OF GUAIACUM AND TURPENTINE.

R. Powdered resin of guaiacum one drachm  
Venice turpentine sufficient  
Mix, and make fifteen pills. One, three times a-day, in gleet and leucorrhœa.  
*Ellis.*

#### PILLS OF GUAIACUM, ALOES, &C.

R. Powdered guaiacum resin one drachm  
aloes thirty-six grains  
rhubarb two drachms  
Canada balsam sufficient  
Form mass, and divide into forty-eight pills. One, every three hours, as a stimulating cathartic.  
*Ellis.*

#### PILLS OF GUAIACUM, ANTIMONY, &C.

R. Powdered resin of guaiacum  
Golden sulphuret of antimony, each, half a drachm  
Calomel ten grains  
Extract of dandelion sufficient  
Form mass, and divide into three-grain pills. Three to four, three times a-day, in painful affections of the joints.  
*Hildebrand.*

#### PILLS OF GUAIACUM, SULPHUR, &C.

R. Powdered resin of guaiacum half an ounce  
Washed sulphur one ounce  
Crude antimony one drachm and a-half  
Extract of blessed thistle sufficient  
Form mass, and make pills of two grains. Ten, three times a-day, in gout, rheumatism, and obstinate cutaneous affections.  
*Quarin.*

#### MIXTURE WITH GUAIACUM, &C.

R. Powdered resin of guaiacum two drachms  
Powdered nitrate of potassa one drachm and a-half  
Powdered gum Arabic one drachm  
tartar emetic one grain  
liquorice one scruple  
Distilled or mint water eight fl. ounces  
Mix. A tablespoonful, three or four times a-day, in rheumatism, &c. *Ellis.*

#### GUAIACUM MIXTURE.

R. Resin of guaiacum three drachms  
Sugar half an ounce  
Gum Arabic mixture half a fl. ounce  
Cinnamon water nineteen fl. ounces  
Rub the resin with the sugar, then with the mixture of gum Arabic, and gradually add the cinnamon water while rubbing.  
*Lond. Ph.*

One to three tablespoonfuls, two or three times a-day.

#### MIXTURE OF GUAIACUM AND BITTER-SWEET.

R. Powdered resin of guaiacum two drachms  
Powdered gum Arabic three drachms  
Extract of bitter sweet three drachms  
Syrup of mallows one fl. ounce  
Distilled water seven fl. ounces  
Mix. A tablespoonful every three hours, in chronic rheumatism. *Berends.*

#### TINCTURE OF GUAIACUM.

R. Powdered resin of guaiacum half a pound  
Alcohol two pints  
Macerate for fourteen days, and filter.  
*U. S. Ph.*

Dose, one to three fl. drachms, three or four times a-day, in gout and chronic rheumatism; to be given in milk or mucilage.

R. Powdered resin of guaiacum four ounces  
Carbonate of soda or potassa one drachm and a-half  
Powdered pimento one ounce  
Diluted alcohol one pint



Digest for a few days. Add volatile spirit of ammonia, if required, in proportion of one or two drachms to each four ounces of tincture. Highly praised in idiopathic amenorrhœa, in doses of a teaspoonful, three times a-day, in milk or wine.

*Dewees.*

#### AMMONIATED TINCTURE OF GUAIAIACUM.

R. Powdered resin of  
guaiacum four ounces  
Aromatic spirit of  
ammonia one pint and a-half  
Macerate for fourteen days, and filter.

*U. S. Ph.*

Dose, one to two fl. drachms, two or three times a-day, in chronic rheumatism.

#### AMMONIATED TINCTURE OF GUAIAIACUM AND COPAIBA.

R. Ammoniated tincture of  
guaiacum one fl. ounce  
Copaiba half fl. ounce  
Mix. A teaspoonful, two or three times a-day, in amenorrhœa with general debility.

*Ellis.*

#### AMMONIATED TINCTURE OF GUAIAIACUM AND CAMPHOR.

R. Ammoniated tincture of guaiacum  
Camphorated tincture of  
opium, each, one fl. ounce  
Mix. Two teaspoonfuls every two hours, in misplaced and retrocedent gout. *Ellis.*

#### ODONTALGIC MIXTURE OF GUAIAIACUM.

R. Tincture of guaiacum six fl. drachms  
Oil of valerian one fl. ounce  
Syrup of scurvy grass  
Compound tincture of  
benzoin, each, two fl. drachms  
Tincture of opium one drachm  
Mix. A teaspoonful mixed with hot water, to be held in the mouth, in toothache.

*Carus.*

#### MIXTURE OF TINCTURE OF GUAIAIACUM AND HENBANE.

R. Tincture of guaiacum two fl. drachms  
henbane one fl. drachm  
Mix. Twenty to thirty drops, morning and evening, in spasm of the stomach and neuralgia.

*Radius.*

## H

### HÆMATOXYLON.

#### LOGWOOD.

This is the heart wood of *Hæmatoxylin campechianum*, a middle-sized tree, a native of Campechy, and naturalized in several of the West Indian islands. It is principally used for tinctorial purposes, but is also employed in medicine.

*Ser. Syst.* Decand. monog. *Nat. Syst.* Fabaceæ.

Linn. Sp. Pl. 549. Griffith, Med. Bot. 252.

It occurs in logs of a dark color externally, and deep red internally; for medical use, it is chipped or rasped. It has a feeble but peculiar odor, and a sweetish, somewhat astringent taste. It is used as a mild astringent in chronic diseases, and relaxed condition of the bowels.

#### INFUSION OF LOGWOOD.

R. Rasped logwood half an ounce  
Boiling water one pint  
Infuse for two hours, and strain. A table-spoonful every two or three hours, in cholera and diarrhœa of children. *Ellis.*

#### DECOCTION OF LOGWOOD.

R. Rasped logwood one ounce  
Water two pints  
Boil down to a pint, and strain. *U. S. Ph.*

A good astringent in diarrhœa, especially in

children. Dose for an adult, two fl. ounces; for a child about two years of age, two or three fl. drachms, several times a-day.

#### EXTRACT OF LOGWOOD.

R. Rasped logwood one pound  
Water one gallon  
Boil to four pints, strain while hot, then evaporate to proper consistence. *U. S. Ph.*  
Dose, from ten to thirty grains.

#### ELECTUARY OF EXTRACT OF LOGWOOD.

R. Extract of logwood  
liquorice, each, two drachms  
Peruvian bark three drachms  
cascarilla one drachm  
Mucilage of quince sufficient  
Mix. A teaspoonful, three times a-day, in chronic diarrhœa, &c. *Radius.*

#### MIXTURE OF EXTRACT OF LOGWOOD.

R. Extract of logwood three drachms  
Boiling water seven fl. ounces  
Dissolve, strain, and add  
Tincture of cinnamon six fl. drachms  
catechu two fl. drachms  
Mix. Dose, one fl. ounce, every six hours. *Beasley.*



- R. Extract of logwood three drachms  
 Tincture of catechu two fl. drachms  
 Water seven fl. ounces  
 Mix. Two spoonfuls, every three or four hours. In diarrhoea and dysentery. *Ellis*.  
 R. Extract of logwood three drachms  
 Spirit of cinnamon one fl. ounce and a-half  
 Water seven fl. ounces  
 Tincture of kino two fl. drachms  
 Mix. In the same doses, and in the same diseases, as the last. *Swediaur*.

## HAMAMELIS.

### WITCH HAZEL.

The *Hamamelis Virginica* is a large native shrub, found along streams in most parts of the country, presenting several varieties.

*Sex. Syst.* Tetrand. digyn. *Nat. Syst.* Hamamelidaceæ.

*Pursh.* Fl. Am. i. 116. *Griffith, Med. Bot.* 350.

The parts used are the bark and leaves. These are bitter and astringent, leaving a sensation of sweetness. They are considered sedative, astringent and tonic, and useful in bowel affections and hemorrhages, and externally as an application to tumours, painful hemorrhoids, &c. They are used in infusion, decoction, or poultice.

## HEDEOMA.

### PENNYROYAL.

*Hedeoma pulegioides* is a small, annual, aromatic plant, abundant in most parts of the United States, growing in dry, sterile situations.

*Sex. Syst.* Diand. monog. *Sex. Syst.* Lauriaceæ.  
*Persoon. Synop.* ii. 131. *Griffith, Med. Bot.* 508.

The whole plant is used. It has a warm, pungent taste, and a powerful aromatic odor. It is a stimulating aromatic, used to obviate nausea and relieve flatulence, and also, in domestic practice, as an emmenagogue.

### INFUSION OF PENNYROYAL.

- R. Pennyroyal two drachms  
 Boiling water sufficient  
 to afford six ounces of strained infusion. To be taken freely, in a warm state, at bedtime, the feet previously bathed in hot water, in amenorrhœa.

### OIL OF PENNYROYAL.

- R. Pennyroyal at will  
 Water sufficient  
 Distil, and collect the oil. Dose, two to ten drops, in flatulent colic and nausea.

## HELENIIUM.

### SNEEZEWORT.

Sneezewort, or *Helenium autumnale*, is an indigenous plant, found in wet situations, in most parts of the United States, flowering in the autumn.

*Sex. Syst.* Syngen. super. *Nat. Syst.* Asteraceæ.  
*Torrey & Gray, Fl. ii.* 384. *Griffith, Med. Bot.* 398.

This plant is almost inodorous, but has a bitter, somewhat pungent, acrid taste. It is tonic, diaphoretic, and powerfully errhine; this latter property is most developed in the flowers, and especially the central florets; these, when powdered, may be used in those cases to which errhines are applicable.

## HELIANTHEMUM.

### FROSTWEED.

The *Helianthemum Canadense* is a native perennial plant, growing in many parts of the United States, in dry, sandy soils.

*Sex. Syst.* Polyand. monog. *Nat. Syst.* Cistaceæ.  
*Mich. Fl. i.* 308. *Darlington, Fl. Cest.* 313.

It has little or no odor, but an astringent, somewhat aromatic, bitterish taste. It is tonic and astringent, and has proved beneficial in scrofula.

## HELLEBORUS.

### BLACK HELLEBORE.

Several species of *Helleborus* are possessed of almost identical medical properties, but the only one used in this country, is *H. nigra*, so called on account of the color of its roots. It is a native of mountain woods, in many parts of Europe, and is cultivated as an ornamental flowering plant.

*Sex. Syst.* Polyand. polyg. *Nat. Syst.* Ranunculaceæ.

*Linn. Sp. Pl.* 783. *Griffith, Med. Bot.* 85.

The parts used are the small fibres, or roots, which are about as thick as straw, and black; they proceed from a rhizome or caudex, and have a somewhat nauseous odor, and a bitter, unpleasant, acrid taste. Black hellebore is a drastic hydragogue purgative; having, also, considerable emmenagogue powers. The dose in powder is from ten to twenty grains, as a purge; two or three grains, as an alterative.

### COMPOUND BLACK HELLEBORE PILLS.

- R. Powdered black hellebore fifteen grains  
 Calomel five grains  
 Powdered ipecacuanha three grains  
 Syrup of ginger sufficient  
 Mix, and make four pills. Two to be taken every four hours, in dropsy, till full purgation is caused. *Ainslie*.



## EXTRACT OF BLACK HELLEBORE.

R. Powdered black hellebore at will  
Exhaust by means of the displacement  
process, with cold diluted alcohol. Distil  
off the alcohol, and evaporate to proper  
consistence. *U. S. Ph.*

Dose, ten to fifteen grains, as a drastic purge.

VINO-ALCOHOLIC EXTRACT OF BLACK  
HELLEBORE.

R. Powdered black hellebore two pounds  
Carbonate of potassa half a pound  
Diluted alcohol eight pints  
Macerate for twelve hours, express, and  
pour on the residuum  
White wine eight pints  
Digest for twenty-four hours, express, mix  
the two tinctures and evaporate.

*Cottureau.*

Dose, ten to fifteen grains.

COMPOUND PILLS OF EXTRACT OF BLACK  
HELLEBORE.

R. VINO-alcoholic extract of  
black hellebore  
Extract of myrrh, each, two ounces  
Powder of blessed thistle one ounce  
Beat together, and form pills of one grain.  
Ten to twenty a-day, in amenorrhœa,  
dropsy, &c. *Radius.*

These pills are much used in Europe, under  
the name of Bacher's pills.

R. Extract of black hellebore  
Assafetida  
Gum ammoniac  
Soap, each, two drachms  
Rhubarb sufficient  
Beat into mass, and form pills of two  
grains. Dose, ten to twelve, morning and  
evening, as a purgative and emmena-  
gogue. *Augustin.*

## TINCTURE OF BLACK HELLEBORE.

R. Bruised black hellebore four ounces  
Diluted alcohol two pints  
Macerate for fourteen days, express, and  
filter. *U. S. Ph.*

Used as an emmenagogue, in doses of thirty  
drops to a fl. drachm, night and morning, watch-  
ing its action.

MIXTURE WITH TINCTURE OF BLACK  
HELLEBORE.

R. Tincture of black  
hellebore half a fl. ounce  
Tincture of myrrh one fl. ounce  
Spanish flies two fl. drachms  
Mix.

Thirty drops, three times a day, in sugar and  
water, as an emmenagogue.

COMPOUND WINE OF BLACK HELLE-  
BORE.

R. Bruised black hellebore one ounce  
wormwood a handful  
White wine four pints  
Macerate for three days, express, and fil-  
ter. Two to three spoonfuls, in the  
morning, fasting. In dropsy. *Brunner.*

## OINTMENT OF BLACK HELLEBORE.

R. Powdered black  
hellebore one to two drachms  
Lard one ounce  
Mix. As an application to obstinate her-  
petic eruptions. *Souberian.*

## HEMIDESMUS.

## INDIAN SARSAPARILLA.

The *H. indicus* is a native of many parts of  
India. It is a climbing plant, and has been  
long used in the East as an efficient medicinal  
agent, but was almost unknown in Europe, or  
this country, until about 1819.

*Sex. Syst.* Pentand. digyn. *Nat. Syst.* Ascle-  
piadaceæ.

Brown, Hort. Kew. ii. 75. Griffith, Med.  
Bot. 453.

The root, which is the part used, is long, tor-  
tuuous, rugose, with longitudinal furrows; it is  
brownish, externally, and has a peculiar and  
somewhat aromatic odor, and a bitterish taste.  
It has the properties of sarsaparilla; and like it,  
is given in infusion, decoction, &c. These are  
made in the same manner as those of sarsapa-  
rilla, and given in the same doses, and in simi-  
lar diseases.

## SYRUP OF INDIAN SARSAPARILLA.

R. Indian sarsaparilla half a pound  
Boiling water one pint  
Digest in a covered vessel, with a gentle  
heat, for three or four hours, strain, add  
twice the weight of sugar, and make  
syrup. *Beasley.*

## MIXTURE OF INDIAN SARSAPARILLA.

R. Indian sarsaparilla ten drachms  
Extract of liquorice ten grains  
Distilled water ten fl. ounces  
Digest for twelve hours, strain, heat to  
180°, and again strain. One-third, three  
times a-day. *Beasley.*

R. Syrup of Indian sarsaparilla five ounces  
Solution of potassa half to one fl.  
drachm  
Orange-flower water one fl. ounce  
One fl. ounce, thrice a-day, in barley  
water. For gonorrhœa. *Bellinarye.*

## INFUSION OF INDIAN SARSAPARILLA.

R. Indian sarsaparilla two ounces  
Lime water one pint



Infuse in a close vessel for twelve hours.  
Dose, a wineglassful. *Ashburner.*

#### DECOCTION OF INDIAN SARSAPARILLA.

R. Indian sarsaparilla two ounces  
Water one pint and a-half  
Boil down to one pint. To be taken during the day, in wineglassful doses. *Pereira.*

### HEPATICA.

#### LIVERWORT.

The *H. triloba* is a small native plant, found in most parts of the United States, and also in the northern regions of Europe and Asia, having three lobed leaves, and presenting two marked varieties: one with the lobes of the leaves rounded, the other with them acute.

*Sex. Syst.* Polyand. polyg. *Nat. Syst.* Ranunculaceæ.

Torrey & Gray, Fl. i. 14. Griffith, Med. Bot. 81.

The whole herb is used. It is inodorous, and has a mucilaginous, slightly astringent and bitterish taste. It is a very mild, demulcent tonic and astringent, and was much employed at one time as a deobstruent in herpetic affections, and also as a remedy in diseases of the lungs. Its powers, however, are very slight. It is given in infusion; to be taken freely.

### HERACLEUM.

#### MASTERWORT.

This is the root of *Heracleum lanatum*, a very large, perennial plant, found in many parts of the United States, in waste places.

*Sex. Syst.* Pentand. digyn. *Nat. Syst.* Apiaceæ.

Torrey & Gray, Fl. i. 632. Griffith, Med. Bot. 335.

The part used is the root; this resembles the parsnip in appearance, but has a rank, unpleasant odor, and a pungent, acrid taste. It is said to be diuretic, expectorant, and antispasmodic, and has proved useful in epilepsy, attended with a disordered condition of the digestive organs. In doses of two or three drachms daily. It is also used in strong decoction, in dyspepsia with flatulence and cardialgia.

### HEUCHERA.

#### ALUM ROOT.

Most of the species of *Heuchera* are possessed of identical properties, but the only one recognized by the U. S. Pharmacopœia is *H. Americana*. This is an indigenous plant, growing in shady, rocky situations, in most parts of the country.

*Sex. Syst.* Pentand. digyn. *Nat. Syst.* Saxifragaceæ.

Torrey & Gray, Fl. i. 578. Griffith, Med. Bot. 313.

The root is rugose, irregular, yellowish, almost inodorous, and of a strong, styptic taste. It is a

powerful astringent, and may be employed in such cases as require medicines of this class.

### HORDEUM.

#### BARLEY.

There are several species of *Hordeum*, but those usually cultivated in this country are *H. vulgare* and *H. distichon*, the latter of which is recognized as officinal. The native country of these is unknown, but they have been cultivated from the earliest ages.

*Sex. Syst.* Triand. digyn. *Nat. Syst.* Graminaceæ.

Linn. Sp. Pl. 125. Griffith, Med. Bot. 664.

The seeds are used in various forms; in that of meal, malt, pearl barley, &c. They are one of the mildest and least irritating of the cerealia, and are much used in decoction, &c., as a nutritive and demulcent drink.

#### PREPARED BARLEY MEAL.

R. Barley meal at will  
Tie it in a linen or cotton cloth, and boil it for twelve hours, then let it cool, remove the outer crust, and pulverize the centre. A useful diet, boiled with milk, in bowel diseases. *Hanover Ph.*

R. Barley meal twelve ounces  
Sugar four ounces  
Powdered cinnamon half a drachm  
Mix, and place in a proper vessel, covering with wheat dough, put in an oven and bake, remove, cool, and pulverize. Half an ounce to two ounces, cooked with water or milk, forms an excellent diet, in debilitated conditions of the system. *Taddei.*

#### BARLEY SUGAR.

R. Decoction of barley a pint and a-half  
Sugar two pounds  
Boil to proper consistence, and form lozenges or rolls. As a demulcent in catarrh. *Giordano.*

#### DECOCTION OF BARLEY.

R. Pearl barley two ounces  
Water four pints and a-half  
Wash the barley in cold water, drain, pour on it half a pint of the water, boil for a short time, drain off this water, add the remainder in a boiling state, and boil down to one-half, and strain. *U. S. Ph.*

As a nutritive and demulcent drink in febrile and inflammatory diseases.

#### COMPOUND DECOCTION OF BARLEY.

R. Decoction of barley two pints  
Sliced figs two ounces and a-half  
liquorice root five drachms  
Stoned raisins two ounces and a-half



Water one pint  
Mix, and boil down to two pints, and strain.  
*Lond. Ph.*

A demulcent, nutritive, and somewhat laxative drink.

#### BARLEY WATER WITH NITRATE OF POTASSA.

R. Decoction of barley one pint  
Nitrate of potassa two drachms  
Lemon juice one fl. ounce  
Mix. To be used warm as a diaphoretic drink.  
*Ellis.*

### HUMULUS.

#### Hops.

By this is meant the strobiles of the *Humulus lupulus*, a climbing vine, a native of Europe, and probably of this country. It is extensively cultivated for its aments or strobiles, which are largely employed in the preparation of malt liquors, and also in medicine.

*Ser. Syst.* Dicc. pentand. *Nat. Syst.* Cannabaceæ.

*Linn. Sp. Pl.* 1457. *Griffith, Med. Bot.* 574.

Hops consist of thin, somewhat translucent, leaflike scales, of a greenish yellow color, having, near their base, two small, round, dark seeds. Their odor is strong and peculiar, somewhat narcotic, and fragrant; their taste is bitter, aromatic, and somewhat astringent. These properties depend on a peculiar secretion, called Lupuline (q. v.). Hops are tonic, and slightly narcotic, and are used in various conditions of the system. The dose, in substance, is from half a drachm to a drachm, but it is seldom administered in this form.

#### INFUSION OF HOPS.

R. Hops half an ounce  
Boiling water one pint  
Macerate for two hours in a covered vessel, and strain.  
*U. S. Ph.*

Dose, one to two fl. ounces, in dyspepsia, nervous tremors, &c.

#### EXTRACT OF HOPS.

R. Hops at will  
Diluted alcohol sufficient  
Treat in a displacement apparatus, distil off the alcohol, and evaporate the residue. Dose, from ten to thirty grains. *Cottrean.*

#### HOP MIXTURE.

R. Extract of hops two drachms  
Water of hops seven fl. ounces  
Tincture of hops half fl. ounce  
Syrup of orange peel one fl. ounce  
Mix. A tablespoonful every hour, as a tonic and stomachic.  
*Niemann.*

#### TINCTURE OF HOPS.

R. Hops five ounces  
Diluted alcohol two pints

Macerate for fourteen days, express, and filter.  
*U. S. Ph.*

Dose, from one to three fl. drachms, as a tonic, and narcotic, especially in the wakefulness and tremors of drunkards.

#### ALKALINE TINCTURE OF HOPS.

R. Hops  
Centaury, each, one ounce  
Peel of bitter orange two drachms  
Carbonate of potassa one scruple  
Diluted alcohol eighteen fl. ounces  
Macerate for eight days, express, and filter. Dose, half an ounce to an ounce.

#### CATAPLASM OF HOPS.

R. Hops one pound  
Flaxseed meal two ounces  
Beer sufficient  
Mix. As an application to bruises and indolent tumors.  
*Radiis.*

#### OINTMENT OF HOPS.

R. Hops two ounces  
Lard ten ounces  
Digest for some hours by a gentle heat, express, and strain; recommended to relieve the pain of cancerous sores.  
*Swedjaur.*

### HYDRARGYRUM.

#### MERCURY.

Mercury is a brilliant, silver-white, fluid metal, having neither taste nor smell. It becomes solid at 39°, and boils at 656°. Mercury, in masses, does not appear to act on the system, but when in a state of great division, it produces marked and peculiar effects. It acts as an alterative, deobstruent, sialagogue, &c. It has been employed in almost all diseases, in some of its preparations, each of which has some peculiarity of action.

#### PURIFIED MERCURY.

R. Mercury six parts  
Distil off four parts, very slowly.  
*Dub. Ph.*

#### MERCURY WITH ANTIMONY.

R. Mercury four parts  
Crude antimony three parts  
Sulphur two parts  
Triturate together till the mercury disappears. As a diaphoretic and alterative. Dose, one to four grains. *Giordano.*

This preparation is much used in France and Germany, under the name of antimonial ethiops.

#### COMPOUND MERCURIAL POWDER.

R. Powdered resin of  
guaiaicum one drachm  
Mercury with antimony  
Magnesia, each, one scruple



Mix. To be taken in two days, in divided doses, in chronic exanthematous affections. *Hufeland.*

R. Mercury with anti-  
mony twenty-four grains  
Subcarbonate of soda eighteen grains  
Powdered sassafras  
Sugar, each, one drachm  
Mix, and divide into six powders. One to be taken daily, in chronic diseases of the skin. *Sundelin.*

#### PILLS OF MERCURY AND ANTIMONY.

R. Mercury one drachm  
Crude antimony four scruples  
Flowers of sulphur two scruples  
Extract of opium one scruple  
Triturate till mercury disappears, with  
Syrup of mallows sufficient  
Divide into one hundred and fifty pills.  
Dose, five or six a-day. Highly spoken of by Huxham, in venereal pains, scrofula, &c. *Baldinger.*

R. Mercury with antimony one drachm  
Extract of hemlock two drachms  
Soap half an ounce  
Galbanum  
Extract of ox gall, each, half a drachm  
Beat into mass, and form pills of two grains. Dose, eight to ten, three times a-day, in scirrhus of the pylorus. *Radius.*

#### MERCURY WITH CHALK.

R. Mercury three ounces  
Prepared chalk five ounces  
Rub together till all the globules disappear. *U. S. Ph.*

A mild preparation, well suited as an alterative in complaints of children. Dose, five grains to half a drachm, twice a-day, for adults, two or three grains for a child.

#### MERCURY AND CHALK.

R. Mercury three ounces  
Resin six drachms  
Prepared chalk five ounces  
Alcohol sufficient  
Make a paste with the resin, and a small quantity of the alcohol; then add the mercury, which may be extinguished in a short time; add the chalk and alcohol gradually, so as to keep up the pasty consistence; then add sufficient alcohol to dissolve out the resin, and wash the powder on a filter, and dry.

*Dr. Stewart, modified by P. Lehman.*

#### MERCURY AND CHALK WITH IPECACUANHA.

R. Mercury with chalk one scruple

Powdered ipecacuanha ten grains  
Mix, and divide into six powders. One, night and morning, in syrup. In dyspepsia with biliary derangement. *Paris.*

#### MERCURY AND CHALK LINIMENT.

R. Mercury  
Prepared chalk, each, half an ounce  
Triturate till globules disappear, and add  
Honey of roses two ounces  
Mix well. As a dressing to venereal ulcers. *Bories.*

#### MERCURY AND GUM.

R. Mercury one part  
Gum Arabic two parts  
Make a mucilage with a quarter of the gum, rub the mercury with it till extinguished, add the remainder of the gum, with as much water as is required, mix well, dry by a gentle heat, and pulverize. *Paris Cod.*

#### PILLS OF MERCURY AND GUM WITH HEMLOCK.

R. Mercury one part  
Gum Arabic two parts  
Syrup of violets four parts  
Triturate till mercury is extinguished, and add  
Extract of hemlock one part  
Powdered liquorice sufficient  
Mix well, and form pills of two grains. Two to four, twice a-day, as alterative. *Plenck.*

#### MERCURIAL MUCILAGE OF GUM ARABIC.

R. Mercury one part  
Gum Arabic two parts  
Water sufficient  
Triturate till the mercury is extinguished. This is a good preparation of mercury; it is given in the dose of two teaspoonfuls, in syrup or mucilage, morning and night, as an alterative or anthelmintic. *Beral.*

#### MERCURIAL GARGLE.

R. Mercury half a drachm  
Gum Arabic three drachms  
Syrup of poppies half an ounce  
Calomel six grains  
Triturate till mercury is extinguished, and add  
Decoction of clematis twenty-six fl. ounces  
Honey of roses one ounce  
Essence of myrrh one drachm  
Mix. As a gargle in syphilitic angina, and ozena. *Plenck.*

#### MERCURIAL LOTION.

R. Mercury one drachm



Gum Arabic four drachms  
 Syrup of poppies sufficient  
 Rub together till globules disappear, and  
 add gradually, constantly rubbing,  
 Boiling milk eight fl. ounces  
 As a lotion in gonorrhœal ophthalmia,  
 ulcers on the penis, and also as a gargle  
 in venereal sore throat. *Plenck.*

#### MERCURY WITH MAGNESIA.

R. Purified mercury  
 Manna, each, two parts  
 Carbonate of magnesia one part  
 Rub the mercury with the manna, adding  
 a little water, till globules disappear; add  
 one-eighth part of the magnesia, still rub-  
 bing, when mixed; then sixteen parts of  
 hot water, and agitate; let sediment sub-  
 side; then decant, and repeat washing, till  
 all the manna is removed; mix the residue  
 with remainder of magnesia, and dry.

*Dub. Ph.*

A mild mercurial, well suited as an alterative  
 in children, especially when there is constipa-  
 tion. Dose, the same as mercury with chalk.

#### MERCURIAL BLUE PILLS.

R. Mercury one ounce  
 Confection of roses one ounce and a-half  
 Powdered liquorice root half an ounce  
 Rub the mercury with the confection, till  
 all the globules disappear, add the liquorice  
 root, and beat into mass. Divide into four  
 hundred and eighty pills. *U. S. Ph.*

One of the mildest and best of the mercurial  
 preparations. From five to fifteen grains as a  
 purgative; one pill every night, or every other  
 night, as alterative; one pill, two or three times  
 a-day, as an alterative. Each pill contains one  
 grain of mercury.

#### BLUE PILL WITH JALAP.

R. Blue pill  
 Powdered jalap  
 aloes, each, fifteen grains  
 Mix, and form twelve pills; three at night,  
 as a purgative. *Ellis.*

#### BLUE PILL WITH RHUBARB.

R. Blue pill nine grains  
 Powdered rhubarb  
 Bicarbonate of soda, each, twelve grains  
 Aromatic syrup of rhubarb sufficient  
 Beat into mass, and form twelve pills.  
 One, twice or thrice a-day, as an alterative,  
 in hepatic derangement. *Hartshorne.*

#### BLUE PILL WITH COLOCYNTH.

R. Blue pill  
 Extract of colocynth, each, five grains  
 Oil of caraway two drops  
 Mix, and make two pills. A very active  
 purgative. *Ellis.*

#### BLUE PILL AND QUINIA.

R. Blue pill  
 Sulphate of quinia  
 Powdered aloes, each, twelve grains  
 Aromatic syrup of rhubarb sufficient  
 Beat into mass, and form twelve pills.  
 One, twice to four times a-day, as a tonic  
 alterative in deranged condition of liver,  
 consequent to fevers. *Ellis.*

#### COMPOUND MERCURIAL PILLS.

R. Blue pill five grains  
 Powdered ipecacuanha two grains  
 Camphor one grain and a-half  
 Syrup of ginger sufficient  
 Mix, and make two pills. One, morning  
 and evening, in hepatitis, till mouth is  
 affected. *Ainslie.*

R. Blue pill four grains  
 Compound powder of squill six grains  
 Powdered ipecacuanha two grains  
 Syrup of ginger sufficient  
 Mix, and make three pills. To be taken  
 in a day; in hepatic obstruction, threaten-  
 ing dropsy. *Ainslie.*

R. Blue pill  
 Antimonial pow-  
 der, each, two and a-half grains  
 Opium half a grain  
 Syrup of ginger sufficient  
 Make a pill. To be taken at bedtime; in  
 venereal blotches, attended with diarrhœa.  
*Ainslie.*

#### ABERNETHY'S PILL.

R. Blue pill ten grains  
 Powdered jalap twenty grains  
 Syrup of buckthorn sufficient  
 Mix well, and divide into six pills. Two  
 at night, with a wineglassful of infusion  
 of senna, in the morning. *Cooley.*

#### MERCURIAL MIXTURE.

R. Mercury six drachms  
 Syrup of poppies one ounce  
 Triturate till globules disappear, and add  
 Orange-flower water two fl. ounces  
 Rose water one fl. ounce  
 Mix well. Dose, a spoonful, morning and  
 night, in gastrodynia and spasmodic vomit-  
 ing. *Cadet de Gassicourt.*

#### MERCURIAL OINTMENT.

R. Mercury two pounds  
 Lard twenty-three ounces  
 Suet one ounce  
 Rub the mercury with the suet and a little  
 of the lard, till the globules disappear;  
 add the remainder of the lard, mixing  
 well. *U. S. Ph.*



This is the strong mercurial ointment. It can be weakened by the addition of lard, as may be wished. Used as an inunction to mercurialize the system, by rubbing about a drachm on inside of thighs, twice a-day.

- R. Mercury two pounds  
Lard, rendered rancid by exposure in a damp, divided state four ounces  
Lard nineteen ounces  
Suet one ounce

Triturate the mercury and rancid lard, until the globules disappear, then add the lard and suet, and triturate until the texture of the mass is uniform. *W. Procter.*

#### CAMPHORATED MERCURIAL OINTMENT.

- R. Mercurial ointment one ounce  
Camphor one drachm  
Mix. Used like the former; also to disperse indolent swellings.

#### MERCURIAL CATAPLASM.

- R. Mercurial ointment half an ounce  
Camphor two drachms  
Boiling milk three ounces  
Crumb of bread sufficient  
Mix, and form cataplasm. *Radius.*

#### MERCURIAL CERATE.

- R. Mercurial ointment  
Simple cerate equal parts  
Mix well. As a dressing to venereal ulcers. *Guibourt.*

#### COMPOUND MERCURIAL CERATE.

- R. Mercurial ointment  
Soap cerate, each, four ounces  
Camphor one ounce  
Rub well together. *Lond. Ph.*

As an application to disperse indolent swellings.

#### COMPOUND MERCURIAL LINIMENT.

- R. Mercurial ointment  
Lard, each, four ounces  
Camphor one ounce  
Rectified spirit one fl. drachm  
Solution of ammonia four fl. ounces  
Rub the camphor with the spirit, then with the lard and mercurial ointment; lastly, add gradually the solution of ammonia, and mix well. *Lond. Ph.*

As a stimulating liniment, in chronic swellings, glandular tumors, &c.

#### OPIATED MERCURIAL LINIMENT.

- R. Mercurial ointment four parts  
Oil of sweet almonds forty-eight parts  
Tincture of opium three parts  
Mix well. Advised in inflamed ulcers of the glans penis, to be applied two or three times a-day. *St. Marie.*

#### COMPOUND MERCURIAL OINTMENT.

- R. Mercurial ointment two ounces  
Lard twelve ounces  
Powdered stavesacre three ounces  
Melt the lard, and add the other ingredients, mixing well. As an application to destroy lice. *Giordano.*

- R. Mild mercurial ointment eight parts  
Soft soap two parts  
Camphor one part  
Rub well together. Recommended in periostitis, and engorgements of the testicles. *Swedjaur.*

#### MERCURIAL AND BELLADONNA PLASTER.

- R. Mercurial ointment one ounce  
Ammoniac six drachms  
Extract of belladonna four drachms  
Hydrocyanic acid thirty drops  
Make mass with ammoniac, and extract with a little water, mix mercurial ointment and acid, and rub the whole together. Useful as an application to scirrhus and scrofulous tumors. *Med.-Chirurg. Pharm.*

#### OINTMENT OF MERCURY AND BELLADONNA.

- R. Strong mercurial ointment thirty parts  
Extract of belladonna four parts  
opium one part  
Balsam of Peru sufficient  
Mix well, as an application to painful hemorrhoidal tumors. *Mignot.*

#### MERCURIAL PLASTER.

- R. Mercury six ounces  
Olive oil  
Resin, each, two ounces  
Lead plaster one pound

Melt the oil and resin together, and, when cool, rub the mercury with them, till the globules disappear; gradually add the lead plaster, previously melted, and mix well.

*U. S. Ph.*

As an application to buboes, venereal nodes, &c.

- R. Mercurial ointment one ounce  
Oil of turpentine  
Camphor, each, two drachms  
Simple cerate one ounce

Mix well. Used as a rubefacient application over the region of the liver, when blisters cannot be used. *Ellis.*

#### HYDRARGYRI ACETAS.

##### ACETATE OF MERCURY.

##### ACETATE OF PROTOXIDE OF MERCURY.

- R. Protonitrate of mercury one part



Distilled water six parts  
Dissolve the salt in water, acidified with a little nitric acid, add a solution of acetate of soda or potassa, filter, wash, and dry the precipitate. *Beral.*

#### PILLS OF ACETATE OF MERCURY.

R. Red oxide of mercury one pound  
Distilled vinegar eight pints  
Dissolve, and triturate a pint of this solution with  
Flake manna two pounds  
Rubbing the mixture for a long time, dry before the fire, often stirring; when of a proper consistence, form pills of a grain and a-half. *Guibourt.*

These pills were highly esteemed under the name of *Keyser's pills*. They contain, at first, the acetate of the deutoxide, but this gradually changes to the acetate of the protoxide, then to an oxide; therefore, the following have been substituted.

R. Acetate of mercury  
Flake manna  
Powdered gum Arabic, each, one scruple  
Rose water sufficient  
Beat into mass, and form twenty pills. As a sialagogue, three to be taken at night, or one three times a-day. *Ellis.*

#### PILLS OF ACETATE OF MERCURY AND OPIUM.

R. Acetate of mercury  
Opium  
Camphor, each, thirty grains  
Syrup of poppies sufficient  
Mix, and make thirty pills. *Carmichael.*

#### SOLUTION OF ACETATE OF MERCURY.

R. Acetate of mercury ten or twelve grains  
Rose water five fl. ounces  
Dissolve. As a lotion in obstinate cutaneous affections. *Niemann.*

#### LINIMENT OF ACETATE OF MERCURY.

R. Acetate of mercury one part  
Olive oil two parts  
Lard six parts  
Rub the salt with a little of the lard, and then add gradually the remainder, and then the oil, constantly rubbing. Used as an application in herpes. *Van Mons.*

### HYDRARGYRUM AMMONIATUM.

#### WHITE PRECIPITATE.

R. Corrosive sublimate six ounces  
Distilled water one gallon  
Solution of ammonia eight fl. ounces

Dissolve the corrosive sublimate in the water, with the aid of heat, and when cold add the solution of ammonia, frequently stirring. Wash the precipitate thoroughly, and dry it. *U. S. Ph.*

This is seldom, if ever, used, except as an external remedy. *U. S. Ph.*

#### OINTMENT OF WHITE PRECIPITATE.

R. White precipitate one drachm  
Simple ointment one ounce and a-half  
Soften the ointment over a gentle fire, and mix the white precipitate. *U. S. Ph.*

As an application to cutaneous eruptions.

### HYDRARGYRI BORAS.

#### BORATE OF MERCURY.

R. Calomel twenty-two parts  
Borate of soda twenty-six parts  
Triturate together; in a quarter of an hour add a little water, then gradually, more constantly rubbing, permit to settle, decant, wash the precipitate till the washings are insipid, and dry. *Van Mons.*

The borate of mercury is said to resemble calomel in its action. Dose, two grains a-day, gradually augmenting.

### HYDRARGYRI BROMIDUM.

#### BROMIDE OF MERCURY.

There are two bromides of mercury; one the protobromide, answering to calomel; the other the bibromide, like corrosive sublimate, an irritant poison.

#### PROTOBROMIDE OF MERCURY.

R. Solution of bromide of potassium at will  
Add a weak solution of protonitrate of mercury, as long as it causes a precipitate; wash this, and dry by a gentle heat. Dose, one grain a-day, gradually increased, as an alterative; in four or five grains as a purgative. *Magendie.*

#### BIBROMIDE OF MERCURY.

R. Bromine  
Mercury, each, equal parts  
Mix, and sublime. Dose, one-twentieth of a grain, gradually increased to a fourth. In syphilis. *Magendie.*

#### ETHEREAL SOLUTION OF BROMIDE OF MERCURY.

R. Bibromide of mercury one grain  
Sulphuric ether one fl. drachm  
Dissolve. Dose, ten to twenty drops, in barley water. In syphilis and lepra. *Werneck.*



HYDRARGYRI CHLORIDUM  
CORROSIVUM.

## CORROSIVE SUBLIMATE.

Corrosive sublimate is a very energetic and poisonous preparation, requiring much care in its exhibition. It will produce the usual effects of the other mercurials, but is less apt to salivate. It is much used in secondary syphilis, and as an alterative, and also as an external application, a stimulant, and escharotic. The best antidote, when poisonous doses have been taken, is albumen, followed by an emetic.

## SOLUTION OF CORROSIVE SUBLIMATE.

R. Corrosive sublimate  
Muriate of ammonia, each, ten grains  
Distilled water (imp. meas.) one pint  
Dissolve. *Lond. Ph.*

A fl. ounce contains half a grain of the mercurial salt. The dose is from one to four fl. drachms, in some mucilaginous drink.

ALCOHOLIC SOLUTION OF CORROSIVE  
SUBLIMATE.

R. Corrosive sublimate one to two grains  
Diluted alcohol four fl. ounces  
Dissolve. A tablespoonful, night and morning, in decoction of sarsaparilla. *Ellis.*

POWDER OF CORROSIVE SUBLIMATE AND  
COPPER.

R. Corrosive sublimate  
Sulphate of copper, each, ten grains  
Nitrate of silver six grains  
Mix. Applied to venereal excrescences, previously moistened with water. *Kruzer.*

POWDER OF CORROSIVE SUBLIMATE AND  
ZINC.

R. Corrosive sublimate  
Sulphate of zinc equal parts  
Mix. In onychia maligna, sprinkled on the sore, and covered with lint soaked in tincture of myrrh. *Perkins.*

## PILLS OF CORROSIVE SUBLIMATE.

R. Corrosive sublimate five grains  
Distilled water thirty to forty drops  
Confection of roses one scruple  
Powdered liquorice sufficient  
Dissolve the corrosive sublimate in the water, and add the other articles, and rub well together. Make forty pills. One, from three to four times a-day, in syphilis. *Ellis.*

COMPOUND PILLS OF CORROSIVE  
SUBLIMATE.

R. Corrosive sublimate half a grain  
Extract of Peruvian bark ten grains  
opium half a grain  
Powdered Peruvian bark sufficient  
Beat together, and make two pills. One,

morning and evening. Each contains a quarter of a grain of the mercurial salt. They require much caution in their use. In syphilis. *Dupuytren.*

R. Corrosive sublimate six grains  
Dissolve in  
Distilled water sufficient  
and add

Extract of hemlock one drachm  
Powdered hemlock sufficient  
Beat into mass, and make forty-eight pills, to be given like the above. Each pill contains an eighth of a grain of corrosive sublimate. *Ellis.*

## CORROSIVE SUBLIMATE MIXTURE.

R. Corrosive sublimate four grains  
Muriate of ammonia eight grains  
Diluted alcohol two fl. ounces  
Mix. A teaspoonful twice a-day, in barley water, in syphilis. *Ainslie.*

R. Corrosive sublimate eight grains  
Muriatic acid twenty-four drops  
Comp. tincture cardamom eight fl. ounces

Mix. Dose, two drachms, morning and evening, in a wineglassful of sugar and water. In syphilis. *Guibourt.*

R. Corrosive sublimate two grains  
Distilled water six fl. ounces  
Spirit of cinnamon  
Syrup, each, one fl. ounce  
Mix. One or two spoonfuls, twice or thrice a-day, in venereal cases. *Ellis.*

## LOTION OF CORROSIVE SUBLIMATE.

R. Corrosive sublimate three to six grains  
Distilled water sufficient  
Dissolve, and add  
Extract of hemlock  
chamomile, each,

two drachms  
Tincture of opium one fl. drachm  
Honey of roses one ounce  
Mix. As an application to venereal ulcers, in the throat, and on the labia pudendi. *Rust.*

R. Corrosive sublimate one drachm  
Distilled water six fl. ounces  
Dissolve, and add  
Muriate of ammonia two drachms  
Nitrate of potassa half an ounce  
Dissolve. As a wash in itch. *Good.*

COSMETIC LOTION WITH CORROSIVE SUB-  
LIMATE.

R. Blanched sweet almonds one ounce  
bitter almonds four drachms  
Cherry water ten fl. ounces



Make an emulsion, and add

Corrosive sublimate            six grains  
Tincture of benzoin       six fl. drachms  
Lemon juice               four fl. drachms

Mix. As a wash for eruptions on the face, to be used morning and evening, previously to be shaken. It must be used with caution. *Cadet de Gassicourt.*

#### LOTION OF CORROSIVE SUBLIMATE AND COPPER.

R. Corrosive sublimate       two grains  
Acetate of copper           six grains  
Distilled water           two pints

Dissolve. As a wash in obstinate porrigo. *Augustin.*

#### LOTION OF CORROSIVE SUBLIMATE AND CAMPHOR.

R. Corrosive sublimate       half a drachm  
Camphor                   one drachm  
Diluted alcohol           one ounce

Dissolve. As an application to destroy condylomata. *Phæbus.*

#### BATEMAN'S MERCURIAL LOTION.

R. Corrosive sublimate       two grains  
Compound spirit of  
lavender                   one fl. ounce  
Distilled water           four fl. ounces

Dissolve. As a lotion in obstinate cutaneous eruptions. *Bateman.*

#### ANTACRID TINCTURE.

R. Powdered resin of guaiacum  
Canada balsam, each,       one ounce  
Corrosive sublimate       one scruple  
Oil of sassafras           two fl. drachms  
Alcohol                   eight fl. ounces

Dissolve the mercurial salt in one-half the alcohol, and then add the remainder, and the other ingredients; after a due digestion, filter. Ten to twenty drops, morning and evening, in wine or water, in syphilis. *Ellis.*

#### COLLYRIUM OF CORROSIVE SUBLIMATE.

R. Corrosive sublimate       two grains  
Opium                   ten grains  
Rose water               four fl. ounces

Dissolve, and add

Mucilage of quince seeds half an ounce  
Mix. *Van Mons.*

#### INJECTION OF CORROSIVE SUBLIMATE.

R. Corrosive sublimate       three grains  
Water of rosemary  
Distilled water, each,       three fl. ounces

Dissolve. As an injection in fistula lachrymalis. *Beer.*

R. Corrosive sublimate       one drachm  
Alcohol                   one fl. ounce

Dissolve, add from five to twenty drops to a solution of

Sulphate of zinc       five to ten grains  
In

Water                   four fl. ounces  
A drachm to be injected into the urethra, three times a-day, in gonorrhœa. *Whately.*

#### YELLOW WASH.

R. Corrosive sublimate one to three grains  
Lime water               one fl. ounce  
Dissolve. *Ellis.*

R. Corrosive sublimate       one scruple  
Carbonate of potassa       one drachm  
Rub together, and add gradually

Distilled water           one pint  
Mix. *Span. Ph.*

These preparations are employed as lotions to venereal and phagedenic ulcers, and should be shaken up when used.

#### COSMETIC WASH.

R. Blanched bitter almonds   six ounces  
Beef tea                   sixty-four ounces

Make an emulsion, and add

Corrosive sublimate       half an ounce  
Lemon juice               ten ounces  
Solution of carbonate of  
potassa                   half an ounce

Incorporate gradually

White of egg               six ounces

Strain, and add

Camphor, rubbed with  
mucilage gum Arabic half a drachm

Mix well. It is said to be an effectual wash for eruptions on the face, but must be used with extreme caution, and very seldom. *Van Mons.*

## HYDRARGYRI CHLORIDUM MITE.

### CALOMEL.

Calomel is prepared by several modes, on the large scale, and is seldom or never made by the apothecary. The several processes will be found, with judicious observations on each, in the U. S. Dispensatory. It is used in almost every disease, either as purgative, alterative, anthelmintic, &c. As a purgative it is peculiar, in not producing effects in proportion to the dose.

#### POWDER OF CALOMEL AND ANTIMONY.

R. Calomel  
Golden sulphuret of  
antimony                   equal parts  
Triturate together. This powder has been much celebrated under the name of *Plummer's Alterative*, as a deobstruent and alterative. The dose is from five to ten



grains a-day, in divided doses. It should be used recently prepared, as in a short time it changes to sulphuret of mercury, and oxide of antimony. *Guibourt.*

## POWDER OF CALOMEL AND JALAP.

R. Calomel five grains  
Powdered jalap ten grains  
Mix. As a purgative, to be mixed with syrup or molasses. *Ellis.*

R. Calomel three grains  
Powdered jalap  
Sugar, each, ten grains  
Mix. Make a powder to be taken at night, or early in the morning, in bilious fevers, and obstructed bowels. *A. T. Thomson.*

## POWDER OF CALOMEL, ANTIMONY, AND HENBANE.

R. Calomel one grain  
Golden sulphuret of antimony three grains  
Powdered extract of henbane one grain  
Powdered sugar half a drachm  
Mix. To be taken night and morning, in spasmodic diseases; is said to have proved very useful in nyctalopia. *Phæbus.*

## POWDER OF CALOMEL, JALAP, AND RHUBARB.

R. Calomel  
Powdered jalap  
rhubarb, each, five grains  
Oil of cinnamon one drop  
Mix. As a purgative, to be given in syrup or molasses. *Ellis.*

## POWDER OF CALOMEL AND FOXGLOVE.

R. Calomel three grains  
Powdered foxglove four grains  
Sugar one drachm  
Mix, and divide into twelve powders. Two a-day, in chronic hydrocephalus. *Berends.*

## POWDER OF CALOMEL, NITRATE OF POTASSA, &amp;c.

R. Calomel six grains  
Nitrate of potassa one drachm  
Tartar emetic half a grain  
Mix, and divide into six powders. One, every two hours, as a diaphoretic in febrile affections. *Ellis.*

## POWDER OF CALOMEL AND OPIUM.

R. Calomel sixteen grains  
Powdered opium four grains  
ipecacuanha eight grains  
Mix, and divide into eight powders. One,

to be taken every hour or two. In dysentery. *Chapman.*

## POWDER OF CALOMEL AND GAMBOGE.

R. Calomel five grains  
Powdered gamboge six to ten grains  
Mix. As an anthelmintic. *Ellis.*

R. Calomel  
Powdered gamboge  
jalap  
rhubarb  
cinnamon, each, two drachms  
Mix. Dose, five to twenty grains. *A. T. Thomson.*

## ANTHELMINTIC PURGATIVE.

R. Calomel three grains  
Compound powder of scammony twelve grains  
Mix. To be taken at once, in cases of lumbrici. *A. T. Thomson.*

## POWDER OF CALOMEL AND PINK ROOT.

R. Calomel four grains  
Powdered pink root ten grains  
Mix. To be taken two mornings in succession, also on afternoon of second day, followed by a mild purgative, as an anthelmintic for children over four years of age. *Ellis.*

## DRY COLLYRIUM OF CALOMEL.

R. Calomel  
Powdered sugar, each, half a drachm  
opium ten grains  
Mix. To be blown into the eye, in ulcers of the cornea. *Radius.*

## CALOMEL PILLS.

R. Calomel half an ounce  
Powdered gum Arabic one drachm  
Syrup sufficient  
Mix the calomel and gum, then beat with syrup into mass, and divide into two hundred and forty pills. *U. S. Ph.*

A very convenient form to give calomel, whether as purgative, alterative, &c.; each pill contains one grain of calomel.

## PILLS OF CALOMEL, QUINIA, &amp;c.

R. Calomel six grains  
Powdered opium three grains  
Sulphate of quinia twelve grains  
Syrup sufficient  
Beat into mass, and form twelve pills. One, night and morning, as an alterative in conditions following bilious fever. *Ellis.*

## PILLS OF CALOMEL AND ACETATE OF LEAD.

R. Acetate of lead half a drachm  
Calomel five grains  
Confection of roses sufficient



Form mass, and divide into ten pills. One to be given every two to four hours, in hematemesis, &c. *Ellis.*

#### PILLS OF CALOMEL AND DANDELION.

R. Calomel four grains  
Extract of dandelion eighteen grains  
Mix. To be taken in a day, in divided doses, in abdominal obstructions. *Brera.*

#### PILLS OF CALOMEL, SQUILL, &c.

R. Plummer's powder twelve grains  
Gum ammoniac two grains  
Extract of dandelion three grains  
Powdered squill half a drachm  
Beat together, and form pills of three grains. Dose, five, three or four times a-day, in engorgements of the abdominal viscera, with anasarca supervening on intermittent fever. *St. Marie.*

#### PILLS OF CALOMEL AND IRON.

R. Calomel  
Golden sulphuret of antimony  
Sulphate of iron  
Myrrh, each, one drachm  
Syrup sufficient  
Beat into mass, and form pills of three grains. Dose, four, morning and evening, in the same cases as above. *Swediaur.*

#### PILLS OF CALOMEL AND CATECHU.

R. Powdered catechu  
Balsam copaiba, each, three drachms  
Calomel one scruple  
Syrup of comfrey sufficient  
Beat into mass, and make one hundred and fifty pills. Four, thrice a-day, in leucorrhœa or gonorrhœa. *St. Marie.*

#### COMPOUND CALOMEL PILLS.

R. Calomel  
Oxysulphuret of antimony, each, two drachms  
Powdered guaiacum resin half an ounce  
Molasses two drachms  
Rub the calomel with the antimony, and then with the other ingredients, till well incorporated. *Lond. Ph.*

These pills are known as *Plummer's pills*, and have been much employed in chronic rheumatism, and obstinate cutaneous affections, especially when there is a syphilitic taint. The dose is from three to six grains, twice a-day.

R. Calomel two scruples  
Precipitated sulphuret of antimony one drachm  
Guaiacum two drachms  
Balsam copaiba sufficient  
Mix, and make sixty pills. Three at

night, in venereal herpes, till mouth is affected; also using frequent tepid baths. *Ainslie.*

#### COMPOUND CATHARTIC PILLS.

R. Powdered compound extract of colocynth half an ounce  
Extract of jalap  
Calomel, each, three drachms  
Gamboge two scruples  
Mix, and with water beat into a mass. To be divided into one hundred and eighty pills. *U. S. Ph.*

As a cathartic or laxative, in cases where there is derangement of the liver. Dose, as a purgative, about ten grains; as a laxative, one of the pills.

#### PILLS OF CALOMEL AND COLOCYNTH.

R. Compound extract of colocynth forty-eight grains  
Calomel one scruple  
Mix, and divide into twenty pills. Two or three will act as a cathartic. *Ellis.*

#### PILLS OF CALOMEL AND ANTIMONY.

R. Calomel ten grains  
Golden sulphuret of antimony one scruple  
Extract of pot marigold  
hemlock, each, two drachms  
Mix, and beat into mass, and form pills of two grains. Dose, five, thrice a-day, in chronic indurations. *Rust.*

R. Calomel six grains  
Kermes mineral twelve grains  
Syrup of elder sufficient  
to make six pills. One every two hours, on the decline of peripneumonia, and in the cough of children arising from worms. *Brera.*

#### PILLS OF CALOMEL AND GUAIACUM.

R. Calomel  
Resin of guaiacum, each, two drachms  
Powdered mallows four ounces  
Syrup sufficient  
Form mass, and make pills of four grains. Dose, four or five a-day, in syphilis. *Alibert.*

#### PILLS OF CALOMEL, JALAP, &c.

R. Calomel  
Resin of jalap  
Compound extract of rhubarb  
Soap equal parts  
Form mass, and make pills of one grain. Dose, ten to twelve grains, as purgative and anthelmintic. *Par. Cod.*

#### PILLS OF CALOMEL AND OPIUM.

R. Calomel two grains  
Opium half a grain



Mix. Thrice a-day, in neuralgia of the face. *Leslie.*

ELECTUARY WITH CALOMEL, &c.

R. Calomel ten grains  
Powdered rhubarb  
    semen contra  
    valerian, each, two drachms  
Conserve of worm-  
    wood one ounce and a-half  
Oxymel of squill sufficient  
Mix. A drachm to half an ounce, in epilepsy, complicated with worms.

*Swediaur.*

CALOMEL OINTMENT.

R. Calomel one drachm  
Rose ointment three drachms  
Mix. As an application in herpes.

*Dupuytren.*

R. Calomel one drachm  
Sublimed sulphur two drachms  
Lard one ounce

Mix well. In obstinate cutaneous affections.

*Fouquier.*

OINTMENT OF CALOMEL AND SQUILL.

R. Calomel  
Powdered squill, each, half a drachm  
Lard two drachms  
Oil of roses four drops

Mix. In chronic swellings of the joints.

*Dupuytren.*

OINTMENT OF CALOMEL AND ACETATE OF COPPER.

R. Calomel  
Acetate of copper, each, one scruple  
Lard eleven drachms

Mix well. As an application in porrigo.

*Cadet de Gassicourt.*

OINTMENT OF CALOMEL, ALUM, &c.

R. Calomel two drachms  
Burnt alum  
White oxide of  
    lead, each, half an ounce  
Oil of turpentine two drachms  
Simple cerate one ounce and a-half

Mix well. As an application to tinea capitis, to be applied at night, and washed off in the morning.

*Ellis.*

CALOMEL AND CAMPHOR OINTMENT.

R. Calomel half a drachm  
Camphor twelve grains  
Lard one ounce

Mix. As an application in lichen. *Bielt.*

16

HYDRARGYRI ET QUININÆ CHLORIDUM.

CHLORIDE OF MERCURY AND QUININE.

R. Bichloride of mercury one part  
Muriate of quinine three parts  
Dissolve separately in the smallest quantity of water, and mix the solutions. Collect the precipitate, and dry by a gentle heat.

*McDermott.*

PILLS OF CHLORIDE OF MERCURY AND QUININE.

R. Chloride of mercury  
    and quinine fifteen grains  
Opium six grains  
Crumb of bread sufficient  
Mix well, and make thirty pills. One, thrice a-day, to produce salivation.

*Hamilton.*

HYDRARGYRI CYANURETUM.

CYANURET OF MERCURY.

R. Ferrocyanuret of iron four ounces  
Red oxide of mercury three ounces  
Distilled water three pounds

Put the ferrocyanuret of iron and the oxide of mercury, well powdered and mixed, into a glass vessel, and pour on them two pints of the water. Boil, and stir, and if in half an hour a blue color remains, add a small portion of oxide of mercury, and continue boiling until the mixture becomes of a yellowish color; then filter, wash the residue in a pint of distilled water, and again filter; mix the solutions, and evaporate, and crystallize. Purify these by dissolving in distilled water, filtering and evaporating.

*U. S. Ph.*

It is given in doses of a sixteenth to an eighth of a grain, in syphilis, chronic inflammation of the thoracic and abdominal organs, &c., and also is employed externally in porrigo, and other cutaneous affections.

SOLUTION OF CYANURET OF MERCURY.

R. Cyanuret of mercury four grains  
Distilled water eight fl. ounces  
Dose, half a drachm to a drachm.

*Chaussier.*

COMPOUND PILLS OF CYANURET OF MERCURY.

R. Cyanuret of mercury six grains  
Opium twelve grains  
Crumb of bread one drachm  
Honey sufficient



Mix, and make ninety-six pills. One, morning and evening. *Guibourt.*

R. Cyanuret of mercury eighteen grains  
Muriate of ammonia  
Extract of aconite, each, three drachms  
box one ounce and a-half  
Oil of anise one scruple  
Beat together, and make four hundred pills. Two, morning and evening. *Cadet.*

#### COMPOUND TINCTURE OF CYANURET OF MERCURY.

R. Cyanuret of mercury eighteen grains  
Water fourteen fl. ounces  
Alcohol ten fl. ounces  
Muriate of ammonia  
Extract of aconite, each, three drachms  
box one ounce and a-half  
Oil of sassafras twenty-four drops  
Dissolve the cyanuret in the water, add the ammonia, extract, and alcohol, let stand for some hours, filter, and add essential oil.

#### GARGLE OF CYANURET OF MERCURY.

R. Cyanuret of mercury ten grains  
Barley water one pint  
Honey of roses one ounce  
Mix. As a gargle. *Brera.*

R. Sarsaparilla half an ounce  
Water sixteen fl. ounces  
Boil, and add towards close of operation  
Fresh hemlock two drachms  
Strain, and add to eight ounces of filtered liquid

Cyanuret of mercury two grains  
As a gargle. *Muller.*

#### OINTMENT OF CYANURET OF MERCURY.

R. Cyanuret of mercury twelve grains  
Lard one ounce  
Rub well together. As an application to venereal ulcers. *Foy.*

R. Cyanuret of mercury sixteen grains  
Lard one ounce  
Oil of lemon fifteen drops  
Rub together. As an application to moist tetter. *Bielt.*

### HYDRARGYRI IODIDUM.

#### IODIDE OF MERCURY.

R. Mercury one ounce  
Iodine five drachms  
Alcohol sufficient  
Rub the mercury and iodine together, adding sufficient alcohol to form a soft paste, and continue rubbing till globules disappear. Dry the iodide in the dark,

with a gentle heat, and keep from light, in a well-stopped bottle. *U. S. Ph.*

The dose is about a grain a-day, gradually increased to three or four. Used in scrofula, and scrofulous syphilis.

#### POWDER OF IODIDE OF MERCURY.

R. Iodide of mercury one to eight grains  
Magnesia one drachm  
Mix. Divide into twelve powders. One, three times a-day. *Radius.*

#### PILLS OF IODIDE OF MERCURY.

R. Iodide of mercury five grains  
Confection of roses sufficient  
Mix, and make thirty pills. One, three times a-day. *Ellis.*

R. Iodide of mercury one drachm  
Confection of dog rose three drachms  
Powdered ginger one drachm  
Beat together. Dose, five to ten grains. *Lond. Ph.*

R. Iodide of mercury one grain  
Extract of juniper twelve grains  
Powdered liquorice sufficient  
Mix, and make eight pills. Two, morning and evening, gradually increasing to double the number. *Magendie.*

#### COMPOUND PILLS OF IODIDE OF MERCURY.

R. Iodide of mercury six grains  
Extract of opium four grains  
Lactucarium twenty-four grains  
Extract of guaiacum forty-eight grains  
Beat together, and form forty-eight pills. In syphilis of children; one pill to a child six months old, and two to four at more advanced ages. *Ricord.*

R. Iodide of mercury half a drachm  
Extract of guaiacum one drachm  
lettuce two scruples  
sarsaparilla sufficient  
Mix, and make seventy-two pills. One, and then two daily. *Bielt.*

R. Iodide of mercury  
Extract of lettuce, each, half a drachm  
hemlock one drachm  
Mix, and make thirty pills. *Ricord.*

R. Iodide of mercury two scruples  
Aloes  
Sulphate of iron  
Myrrh, each, half a drachm  
Oil of savine twenty drops  
Mix, and make twenty-four pills. One, thrice a-day, in amenorrhœa. *Barbour.*

#### OINTMENT OF IODIDE OF MERCURY.

R. Iodide of mercury one scruple  
Lard one ounce and a-half



Rub well together. In scrofulous swellings, indolent granular tumors, &c.

*Soubeiran.*

R. Iodide of mercury           one ounce  
White wax                   two ounces  
Lard                           six ounces

Melt the wax and lard, and stir in the iodide. *Lond. Ph.*

As a dressing to scrofulous ulcers, &c.

R. Iodide of mercury           six grains  
Acetate of morphia       eight grains  
Lard                           one ounce

Rub well together. As an application to obstinate glandular swellings. *Pelletan.*

R. Iodide of mercury           one drachm  
Soap                          half a drachm  
Rose water                  two drachms  
Rose ointment              six drachms  
Mix. *Riecke.*

## HYDRARGYRI IODIDUM RUBRUM.

### RED IODIDE OF MERCURY.

R. Corrosive sublimate       one ounce  
Iodide of potassium       ten drachms  
Distilled water           two pints

Dissolve the corrosive sublimate in a pint and a-half, and the iodide in half a pint, of the water, and mix the solutions. Filter, and wash the collected precipitate with distilled water. Dry by a gentle heat, and keep in a well-stopped bottle. *U. S. Ph.*

Much more active than the iodide. Used in the same cases, in doses of a sixteenth, gradually increased to the fourth of a grain.

### PILLS OF RED IODIDE OF MERCURY.

R. Red iodide of mercury    one grain  
Extract of juniper       twelve grains  
Powdered liquorice       sufficient  
Mix, and make eight pills. Two, morning and evening. *Guibourt.*

R. Red iodide of mercury    five grains  
Syrup                       sufficient

Rub well together, and then with Crumb of bread Sugar, each,           sufficient to make sixty pills. Two, morning and evening, gradually increasing. *Radius.*

### WASH OF RED IODIDE OF MERCURY.

R. Red iodide of mercury    twelve grains  
Distilled water           six fl. ounces  
Mix. As a lotion to scrofulous ulcers. *Radius.*

### TINCTURE OF RED IODIDE OF MERCURY.

R. Red iodide of mercury    twenty grains

Alcohol (.837) one fl. ounce and a-half  
Dissolve. Dose, ten to twenty drops, in distilled water. Twenty-six drops contain about an eighth of a grain. *Foy.*

### ETHEREAL TINCTURE OF RED IODIDE OF MERCURY.

R. Red iodide of mercury    twenty grains  
Sulphuric ether       one ounce and a-half  
Dissolve. Rather more powerful than the last, and therefore to be given in smaller doses. *Magendie.*

### OINTMENT OF RED IODIDE OF MERCURY.

R. Prepared like the ointment of the iodide. *Lond. Ph.*

It is more active than the ointment of the iodide.

R. Red iodide of mercury    one scruple  
Lard                          one ounce and a-half  
Rub well together. As a dressing to obstinate venereal ulcers. *Soubeiran.*

R. Red iodide of mercury    fifteen grains  
Lard                          one ounce  
Essence of bergamot    twenty drops  
Mix well. As an application to chronic cutaneous affections. *Foy.*

## HYDRARGYRI NITRAS.

### NITRATE OF MERCURY.

### NITRATE OF THE PROTOXIDE OF MERCURY.

R. Mercury                   equal parts  
Nitric acid                   equal parts  
Dissolve, with aid of a gentle heat, then boil until a yellow sediment is formed, decant, and permit to crystallize. *Guibourt.*

### PILLS OF PROTONITRATE OF MERCURY.

R. Protonitrate of mercury   ten grains  
Extract of liquorice       forty grains  
Beat into mass with a little water, and form sixty pills. *Foy.*

### OINTMENT OF NITRATE OF MERCURY.

R. Mercury                   one ounce  
Nitric acid                   eleven fl. drachms  
Fresh neat's foot oil   nine fl. ounces  
Lard                          three ounces  
Dissolve the mercury in the acid, melt the oil and lard together, and when they become thick, on cooling, add the solution, and mix well. *U. S. Ph.*

A stimulant and alterative application, used in most cases diluted with lard. Employed in various skin diseases, &c. It is known under the name of *Citrine ointment*.







Simple ointment eight ounces  
 Soften the ointment by a gentle heat, and  
 add oxide of mercury and mix well.

*U. S. Ph.*

An efficacious, stimulating application to foul  
 and indolent ulcers, psorophthalmia, &c. Where  
 it is too stimulating, dilute with lard.

R. Red oxide of mercury one drachm  
 Venice turpentine one ounce  
 Mix. As an application to indolent ulcers,  
 and also used in itch. *Ellis.*

#### OINTMENT OF RED OXIDE OF MERCURY AND SULPHUR.

R. Red oxide of mercury  
 Mercury, each, one part  
 Precipitated sulphur sixteen parts  
 Triturate till globules disappear, and add  
 Lard thirty-two parts  
 Rub well together. As a friction in the  
 cure of itch. *Swediaur.*

#### BROWN OINTMENT OF RED OXIDE OF MERCURY.

R. Red oxide of mercury one part  
 Basilicon ointment six parts  
 Rub well together. Used in the treatment  
 of indolent venereal ulcers. *Saunders.*

#### OINTMENT OF RED OXIDE OF MERCURY AND TIN.

R. Red oxide of mercury two drachms  
 Amalgam of equal parts of  
 mercury and tin four drachms  
 Rose ointment one ounce  
 Rub well together, and add  
 Oil of peppermint twenty drops  
 Highly spoken of in hemorrhoidal tumors.  
*Cadet de Gassicourt.*

#### OINTMENT OF RED OXIDE OF MERCURY AND CINNABAR.

R. Red oxide of mercury one ounce  
 Cinnabar one drachm  
 Lard one pound  
 Rub well together. Employed in chronic  
 inflammations of the eyelids. *Giordano.*

#### OINTMENT OF RED OXIDE OF MERCURY AND ZINC.

R. Red oxide of mercury one part  
 Sulphate of zinc two parts  
 Lard ninety-six parts  
 Rub well together. In scrofulous ophthal-  
 mia, as an application to the edges of the  
 eyelids. *Dupuytren.*

#### OINTMENT OF RED OXIDE OF MERCURY AND LEAD.

R. Red Oxide of mercury  
 Prepared calamine

Burnt alum  
 Acetate of lead, each, one drachm  
 Corrosive sublimate twelve grains  
 Rose ointment one ounce  
 Rub well together. Celebrated in the  
 treatment of herpetic affections. *Guibourt.*

#### HYDRARGYRI PHOSPHAS.

##### PHOSPHATE OF MERCURY.

R. Solution of nitrate of mercury at will  
 Add to it a solution of phosphate of soda,  
 as long as any precipitate is formed; de-  
 cant, wash the precipitate well, and dry it.  
*Van Mons.*

This has been highly praised in secondary  
 syphilis. The dose is half a grain to a grain,  
 combined with sugar or some aromatic powder.

##### PILLS OF PHOSPHATE OF MERCURY.

R. Phosphate of mercury nine grains  
 Tartar emetic one grain  
 Opium six grains  
 Conserve of roses sufficient  
 Mix, and make nine pills. One in the  
 evening, at bedtime. *Radiis.*

#### HYDRARGYRI SULPHAS.

##### SULPHATE OF MERCURY. PERSULPHATE OF MERCURY.

R. Purified mercury  
 Sulphuric acid, each, six parts  
 Nitric acid one part  
 Expose them to heat in a glass vessel, and  
 increase the heat, until the mass becomes  
 white and perfectly dry. *Dub. Ph.*

Not used as a remedy, but serves to form cor-  
 rosive sublimate, calomel, and turpeth mineral.

##### YELLOW SULPHATE OF MERCURY.

R. Persulphate of mercury one part  
 Warm water twenty parts  
 Rub together in an earthenware mor-  
 tar, and pour off the supernatant liquid,  
 wash the yellow powder with warm dis-  
 tilled water, as long as the washings are  
 precipitated by caustic potassa. Dry.  
*Dub. Ph.*

This is alterative, errhine, and emetic. The  
 dose, as an alterative, is from a quarter to half  
 a grain; as an errhine, one grain, mixed with  
 five grains of some bland powder; as an emetic,  
 two to five grains.

##### COMPOUND POWDER OF SULPHATE OF MERCURY.

R. Sulphate of mercury one grain  
 Powdered asara-  
 bacca one drachm and a-half



Mix, and divide into eight powders. One to be used as an errhine, twice a-day; in gutta serena, ozaena, &c. *Ellis.*

#### OINTMENT OF SULPHATE OF MERCURY.

R. Sulphate of mercury one part  
Lard eight parts  
Rub well together. As an application to herpes, porrigo, &c. *Soubeiran.*

R. Sulphate of mercury two drachms  
Chloride of lime three drachms  
Oil of almonds six drachms  
Lard two ounces  
Melt together the oil and lard, and mix the two powders. More stimulating than the last. Used in obstinate cutaneous affections. *Chevallier.*

#### HYDRARGYRI SULPHURETUM NIGRUM.

##### BLACK SULPHURET OF MERCURY.

R. Mercury  
Sulphur, each, one pound  
Rub together till all globules disappear. *U. S. Ph.*

Employed as an alterative in cutaneous affections and glandular swellings. Dose, from five to thirty grains, or even more, several times a-day.

##### POWDER OF BLACK SULPHURET OF MERCURY.

R. Black sulphuret of mercury four drachms  
Prepared chalk two ounces  
Powdered amber one ounce  
Mix well, and divide into sixty-four powders. One, and afterwards two a-day, in a glass of sugar and water. In scrofulous affections. *St. Marie.*

R. Black sulphuret of mercury  
Prepared oyster shell, each, one drachm  
Powdered orange peel half a drachm  
Mix, and make ten powders. Three a-day, as an anthelmintic. *Phœbus.*

##### PILLS OF BLACK SULPHURET OF MERCURY.

R. Black sulphuret of mercury  
Extract of bitter-sweet, each, two drachms  
Powdered mallows sufficient  
Mix, and form one hundred and twenty pills. Four to five, three times a-day. In obstinate cutaneous affections. *Radius.*

R. Black sulphuret of mercury one scruple  
Extract of dandelion three drachms

Gum ammoniac two drachms  
Powdered squill half a drachm  
Syrup sufficient

Beat together, and make pills of three grains. Dose, five, thrice a-day. In ascites with engorgement of the abdominal viscera. *St. Marie.*

R. Black sulphuret of mercury half an ounce  
Guaiaicum  
Soap, each, two drachms  
Golden sulphuret of antimony two scruples  
Extract of horehound sufficient  
Beat together, and form pills of two grains. Ten, morning and evening. In chronic rheumatism. *Baldinger.*

#### HYDRARGYRI SULPHURETUM RUBRUM.

##### RED SULPHURET OF MERCURY.

R. Mercury forty ounces  
Sulphur eight ounces  
Melt the sulphur, and mix the mercury with it over the fire. When mass begins to swell, remove from fire, and cover the vessel, to prevent combustion; when cool, powder, and sublime. *U. S. Ph.*

At one time, much used as alterative and anthelmintic, but now seldom employed except for fumigation.

##### POWDER OF RED SULPHURET OF MERCURY.

R. Red sulphuret of mercury half an ounce  
Powdered sugar candy three ounces  
Oil of cinnamon half a drachm  
Rub well together. At one time celebrated as cordial, stomachic, and analeptic, in doses of a scruple. *Swediaur.*

##### PILLS OF RED SULPHURET OF MERCURY.

R. Red sulphuret of mercury  
Powdered Virginia snake-root, each, half a drachm  
Syrup sufficient  
Mix, and form twelve pills. Two to be taken thrice a-day; alterative and diaphoretic; useful in obstinate cutaneous affections. *Ellis.*

R. Red sulphuret of mercury  
Extract of wormwood  
hemlock, each, one drachm  
Beat together, and form pills of two grains. Dose, seven, morning and evening, in venereal glandular swellings. *Augustin.*



## FUMIGATING POWDER.

R. Red sulphuret of  
mercury four drachms  
Olibanum two drachms  
Mix. To be thrown on a red hot iron, and  
the diseased parts exposed to the fumes.  
In herpes, and venereal pustules. *Foy.*

## CERATE OF RED SULPHURET OF MERCURY.

R. Powdered red sulphuret  
of mercury fifteen grains  
Yellow resin half an ounce  
Yellow wax  
Lard, each, half a pound  
Melt the resin, wax, and lard together, and  
add the sulphuret. As a dressing to ill-  
conditioned ulcers. *Swedjaur.*

## OINTMENT OF RED SULPHURET OF MERCURY.

R. Red sulphuret of  
mercury one drachm and a-half  
Muriate of ammonia half a drachm  
Lard two ounces  
Rose water one fl. drachm  
Rub well together. Advised in phthiri-  
asis, to diminish the itching, and to destroy  
the insects. *Radius.*  
R. Red sulphuret of mercury one drachm  
Camphor one scruple  
Simple cerate one ounce  
Rub well together. In the treatment of  
herpes; to be applied at least twice a-day.  
*Alibert.*

## HYDRARGYRI TARTRAS.

## TARTRATE OF MERCURY.

R. Black oxide of mercury at will  
Tartaric acid sufficient  
Boil together, in a due proportion of water,  
until the oxide is dissolved; filter, and  
evaporate to the point of crystallization;  
wash the crystals. *Swedjaur.*

Dose, one to two grains, twice a-day.

## TARTRATE OF MERCURY AND POTASSA.

R. Boiling, saturated solution of  
Nitrate of mercury at will  
Add, drop by drop, a boiling saturated solu-  
tion of bitartrate of potassa, as long as a  
precipitate is formed, decant, wash the pre-  
cipitate well with cold water, dissolve in  
boiling water, and set aside to crystallize.  
*Van Mons.*

Dose, one to three grains. Once very cele-  
brated as Pressavin's vegeto-mercurial liquor.

## HIPPOCASTANUM.

## HORSE CHESTNUT.

The horse chestnut, or *Æsculus hippocastanum*,  
is a beautiful and lofty tree, a native of the cen-  
tral parts of Asia, but extensively cultivated in  
Europe, and the United States.

*Ser. Syst.* Heptand. monog. *Nat. Syst.* Sapin-  
daceæ.

*Linn. Sp. Pl.* 488. *Griffith, Med. Bot.* 213.

The part used is the bark; this is light, brittle,  
of a brownish red externally, of a yellow brown  
within. It has a somewhat aromatic odor, and  
an astringent, bitter taste. It is a somewhat ac-  
tive astringent, and has been much praised in  
intermittent fevers. The dose of the powder is  
two to four scruples, every three hours, till an  
ounce and a-half is taken.

COMPOUND POWDER OF HORSE CHEST-  
NUT.

R. Powdered horse chestnut bark  
willow bark, each, half an  
ounce  
gentian  
calamus  
cloves, each, two drachms  
Mix. *Hufeland.*

## DECOCTION OF HORSE CHESTNUT BARK.

R. Horse chestnut  
bark one ounce and a-half  
Water thirty fl. ounces  
Boil down to ten ounces, adding towards  
close of operation,  
Liquorice root one drachm  
Strain. A cupful every two hours.  
*Niemann.*

COMPOUND DECOCTION OF HORSE CHEST-  
NUT.

R. Horse chestnut bark  
Willow bark, each, half an ounce  
Calamus  
Root of water avens, each, two drachms  
Water sixteen fl. ounces  
Boil down to one-half. *Spielmann.*  
R. Horse chestnut  
bark one ounce and a-half  
Water eighteen fl. ounces  
Boil down to one-half, strain, and add  
Sulphuric ether one to two drachms  
Syrup of orange peel one ounce  
Mix. To be used during the apyrexia.  
*Phœbus.*

## EXTRACT OF HORSE CHESTNUT.

R. Horse chestnut bark at will  
Water sufficient  
Exhaust the bark in a displacement appa-  
ratus, and evaporate to proper consistence.  
Dose, five to fifteen grains. *Van Mons.*



## HYDRASTIS.

## YELLOW ROOT.

This is the root of *Hydrastis Canadensis*, a native plant, more common in the western than in the eastern States. The flowers are yellow and fugacious, and are succeeded by red fruit, resembling raspberries, but not eatable.

*Sex. Syst.* Polyand. polygyn. *Nat. Syst.* Ranunculaceæ.

Linn. Sp. Pl. 784. Griffith, Med. Bot. 82.

The root is contorted, rugose, and of a bright yellow color. It has a strong, somewhat narcotic odor, and a very bitter taste. It is a powerful tonic bitter, and is also used in decoction as a wash in chronic ophthalmia. It is given in powder or decoction.

## HYOSCYAMUS.

## HENBANE.

Several species of *Hyoscyamus* are medicinal, but the only one that is officinal, is the *H. niger*, a native of Europe, and now naturalized in some parts of the United States. It has a lurid appearance, and a rank, unpleasant odor.

*Sex. Syst.* Pentand. monog. *Nat. Syst.* Solanaceæ.

Linn. Sp. Pl. 257. Griffith, Med. Bot. 484.

All parts of the plant are active, but the leaves are generally employed, though the seeds also are officinal. The dried leaves have but little taste or smell. The seeds are small, roundish, somewhat uniform, of a grayish color; of an unpleasant smell, and an oily, somewhat bitter taste. They both owe their properties to an alkaloid, called *hyoscyamia*. They are anodyne and hypnotic, and are used to relieve pain, to procure sleep, and to quiet an excited nervous system. The dose of the powdered leaves is from five to ten grains, of the seeds rather less.

## AQUEOUS EXTRACT OF HENBANE.

R. Henbane leaves one pound  
Bruise them in a stone mortar, adding a little water; express, heat the juice to boiling point, strain, and evaporate.

*U. S. Ph.*

Dose, a grain, twice a-day, gradually increased till the system is affected.

## ALCOHOLIC EXTRACT OF HENBANE.

R. Henbane leaves, in coarse powder one pound  
Diluted alcohol four pints  
Moisten the henbane with half a pint of the alcohol, and let stand for twenty-four hours; then put in a displacement apparatus, and exhaust, with the remainder of the alcohol, adding sufficient water at the close, to drive off all the spirit. Distil off the alcohol, and evaporate the residue to proper consistence.

*U. S. Ph.*

Dose, as for the last, and in the same manner.

## COMPOUND POWDER OF HENBANE.

R. Powdered extract of henbane  
Oxide of zinc, each, ten grains  
Magnesia half a drachm  
Sugar of milk two drachms and a-half  
Mix, and divide into ten powders. One every three hours, as a sedative, in nervous attacks.

*Vogler.*

## COMPOUND PILLS OF HENBANE.

R. Extract of henbane  
opium  
belladonna  
hemlock, each, one ounce  
Beat together, and form pills of one grain. Said to be useful in neuralgia of the face, in doses of one to six a-day.

*Pierquin.*

R. Extract of henbane half a drachm  
opium two grains

Powdered foxglove

ipécacuanha, each, five grains

Beat together, and make thirty pills. One every three hours, in whooping-cough.

*Phabus.*

R. Extract of henbane  
valerian

Oxide of zinc equal parts

Beat together, and form pills of three grains. In neuralgia, in doses of one to ten.

*Foy.*

R. Extract of henbane two scruples  
Powdered myrrh one drachm and a-half  
squill half a drachm

Distilled water sufficient

Beat together, and make thirty pills. Two, night and morning, in catarrh and phthisis, with debility of the pulmonary organs.

*Paris.*

## PILLS OF HENBANE AND OPIUM.

R. Extract of henbane  
hemlock, each, fifteen grains  
Powdered opium four grains  
Beat into mass, and make ten pills. One at night, as an anodyne.

*Ellis.*

## PILLS OF HENBANE AND IPECACUANHA.

R. Extract of henbane ten grains  
Powdered ipécacuanha five grains  
Mix, and make ten pills. One, every hour or two, in irritability of the bowels, with flatulence.

*Ellis.*

## INFUSION OF HENBANE.

R. Leaves of henbane half an ounce  
Boiling water one pint  
Infuse. As a lotion to painful ulcers.

*Ellis.*

## COMPOUND INFUSION OF HENBANE.

R. Henbane leaves



Poppy heads  
Mallows, each, one ounce  
Boiling water four pints  
Infuse for an hour, and strain. As a fo-  
mentation to painful tumors, &c. *Radiis.*

MIXTURE OF HENBANE.

R. Oil of almonds half an ounce  
Powdered gum Arabic two drachms  
Oil of fennel one drop  
Extract of henbane six grains  
Sugar half an ounce  
Water two fl. ounces  
Rub well together. A spoonful every two  
hours, in catarrh. *Foy.*

MIXTURE OF HENBANE AND ANTIMONY.

R. Extract of henbane ten grains  
Antimonial wine two fl. drachms  
Dissolve. In hooping-cough, in doses of  
five to ten drops. *Augustin.*

TINCTURE OF HENBANE.

R. Henbane leaves four ounces  
Diluted alcohol two pints  
Macerate for fourteen days, express, and  
filter. *U. S. Ph.*

Dose, a fl. drachm.

ETHEREAL TINCTURE OF HENBANE.

R. Henbane leaves one part  
Sulphuric ether six parts  
Macerate for eight days, express, and fil-  
ter. Dose, ten to twelve drops, three or  
four times a-day. *Hesse Ph.*

MIXTURE OF HYOSCYAMUS AND SQUILL.

R. Tincture of squill ten drops  
Nitric acid five drops  
Extract of henbane three grains

Water one fl. ounce and a-half  
Mix. In spasmodic asthma; to be re-  
peated every three or four hours. *Bree.*

LINIMENT OF HENBANE.

R. Extract of henbane half a drachm  
White soap two drachms  
Linseed oil six ounces  
Mix. As an embrocation to glandular  
swellings. *Dupuytren.*

OINTMENT OF HENBANE.

R. Extract of henbane one drachm  
Lard one ounce  
Rub together. As an application to pain-  
ful hemorrhoids. *Taddei.*

COMPOUND OINTMENT OF HENBANE.

R. Leaves of henbane  
stramonium  
bittersweet  
elder, each, a handful  
Lard two pounds  
Heat together till all moisture is evapora-  
ted, and strain. As an application to  
frost-bites, chilblains, &c. *Spielmann.*  
R. Fresh poplar buds four ounces  
Lard twelve ounces  
Leaves of henbane

poppy  
belladonna  
black nightshade, each, one  
ounce

Heat together till all moisture is evapora-  
ted. As a soothing and anodyne oint-  
ment to painful local affections.

*Cottureau.*

This is much used in Europe, under the name  
of *Unguentum populeum*.

ICHTHYOCOLLA.

ISINGLASS.

This is the swimming bladder of several kinds  
of fish. The best, however, is procured from  
various species of sturgeon, inhabiting the more  
southern parts of Russia. It is a pure and nu-  
tritious gelatine, and is rather employed as food,  
and in the arts, than in medicine, in which it  
is only used as the basis of court plaster.

COURT PLASTER.

R. Isinglass one ounce  
Water eight fl. ounces

Macerate for twenty-four hours, and add  
Alcohol eight fl. ounces  
When perfectly dissolved, brush over  
strained silk, or sarsnet, several times, al-  
ternating with a layer of tincture of Peru  
or benzoin. *Cottureau.*

R. Isinglass nine drachms  
Dissolve in water, and add hot filtered mix-  
ture of

Alcohol (0.921) twelve fl. ounces  
Tincture of benzoin two fl. ounces  
Apply this to strained silk, till six coats



are given, permitting each to dry before applying another, then brush over twice with a mixture of

Tincture of benzoin      six fl. ounces  
Liquid turpentine      four ounces  
*Virey.*

R. Isinglass      one ounce  
Water      two fl. ounces  
Mix, and allow to stand till the isinglass is quite soft, then add

Rectified spirit three fl. ounces and a-half  
Mixed with

Water      one fl. ounce and a-half  
Expose to heat of water-bath, till dissolved, and apply to oiled silk, nailed to a board, by means of a soft brush; apply four coats.  
*Liston.*

R. Fasten a piece of fine muslin, linen, or silk, to a flat board, and give it a thin coat of flour paste; over this, when dry, two coats of colorless gelatine are to be applied. Said to be superior to the usual court plaster.  
*Deschamps.*

#### DIAMOND CEMENT.

R. Isinglass, softened in water, and dissolved in the smallest quantity of diluted alcohol      two ounces  
Ammoniac      ten grains  
And whilst still liquid, add  
Solution of mastic, made with half a drachm in three drachms of alcohol.  
Stir well together. *Redwood.*

### INDIGUM.

#### INDIGO.

A tinctorial material, obtained from several species of *Indigofera*. It is found in commerce, in the form of small, solid, brittle masses, having scarcely any taste or odor, of a deep blue color, but assuming a coppery lustre when rubbed. It is principally used in the arts, but has also been employed as a remedial agent, in various spasmodic diseases. The dose is about a scruple, two or three times a-day, rapidly increased to a drachm or more.

#### POWDER OF INDIGO.

R. Powdered indigo      half an ounce  
Aromatic powder      half a drachm  
Mix. A teaspoonful, three times a-day, in epilepsy. *Radius.*

#### ELECTUARY OF INDIGO.

R. Powdered indigo      half an ounce  
Aromatic powder      half a drachm  
Syrup      one fl. ounce  
Mix. To be taken in divided doses, during the day. *Phæbus.*

#### PILLS OF INDIGO.

R. Powdered indigo      seventy-five grains  
Assafetida      fifteen grains  
Castor      seven grains  
Mix, and form twenty pills. One, every hour. *Podreca.*

#### SULPHATE OF INDIGO.

R. Indigo (Bengal) in powder      four ounces  
Sulphuric acid      twenty-four ounces  
Dissolve, by maceration and agitation, and add water, until the solution measures four pints.

### INULA.

#### ELECAMPANE.

The root of *Inula helenium*, a large herbaceous plant, a native of Europe, and cultivated in gardens in this country, where it has also become naturalized in some places.

*Sex. Syst.* Syngen. super. *Nat. Syst.* Astera-ceæ.

Linn. Sp. Pl. 1236. Griffith, Med. Bot. 397.

The root is thick, fleshy, and branched. As dried and found in the shops, it is in slices of a yellowish gray color, with a peculiar and aromatic odor, and an unpleasant, bitterish taste, with an after sensation of pungency. It is tonic and expectorant, and also acts in some cases as a diuretic and diaphoretic. The dose of the powder is from a scruple to a drachm.

#### DECOCTION OF ELECAMPANE.

R. Elecampane root      half an ounce  
Water      one pint  
Boil. Dose, a fl. ounce, every hour or two.  
*U. S. Disp.*

#### COMPOUND PILLS OF ELECAMPANE.

R. Extract of elecampane      one drachm  
Powdered foxglove      ipecacuanha, each, ten grains  
Opium      five grains  
Beat into mass, and form pills of two grains. One every hour, in chronic catarrh. *Heim.*

#### EXTRACT OF ELECAMPANE.

R. Elecampane root      at will  
Water      sufficient  
Make a concentrated infusion, strain, and evaporate to the proper consistence. Dose, from a scruple to a drachm. *Cottureau.*

#### MIXTURE OF ELECAMPANE.

R. Extract of elecampane      two drachms  
Antimonial wine      one fl. ounce  
Syrup of orange flowers      half a fl. ounce  
Mix. Dose, twenty to thirty drops, every three hours, in asthma and chronic catarrh. *Jahn.*

#### COMPOUND OXYMEL OF ELECAMPANE.

R. Elecampane root      one ounce



Orris root half an ounce  
 Water thirty-six fl. ounces  
 Boil down to two-thirds, strain, and add  
 Honey sixteen ounces  
 Gum ammoniac one ounce  
 Vinegar eight fl. ounces  
 Evaporate to the consistence of honey. A spoonful occasionally, in catarrh and humid asthma. *Spielmann.*

## IODINUM.

## IODINE.

Iodine is an elementary non-metallic body, having many of the properties of chlorine. It exists in many marine vegetables and animals, and also in some mineral bodies and salt springs, but is principally obtained from kelp or the ashes of seaweeds. It is soft, friable, and opaque, of a bluish black color and metallic lustre, and has a strong and penetrating odor and an acrid taste. It is employed as a stimulant or alterant in morbid actions of the absorbent and glandular systems, both externally and internally, and, in over doses, acts as an irritant poison.

The dose, in substance, is from the eighth of a grain to a grain.

## PILLS OF IODINE.

R. Iodine six grains  
 Extract of gentian one drachm  
 Mucilage of gum Arabic sufficient  
 Mix, and form twenty-four pills. These are said to be useful in mercurial or scorbutic salivation. *Radius.*

## TINCTURE OF IODINE.

R. Iodine one ounce  
 Alcohol one pint  
 Dissolve. *U. S. Ph.*

Sixteen minims, or thirty-five drops, contain one grain of iodine. Dose, from ten to twenty drops, three times a-day; to be increased.

## SATURATED TINCTURE OF IODINE.

R. Iodine two scruples  
 Alcohol one fl. ounce  
 Compound spirit of lavender two fl. drachms  
 Mix. Five to ten drops, twice a-day, gradually increasing. In amenorrhœa, leucorrhœa, gonorrhœa, &c. *Durand.*

## COMPOUND TINCTURE OF IODINE.

R. Iodine half an ounce  
 Iodide of potassium one ounce  
 Alcohol one pint  
 Mix, and form a tincture. Dose, five to fifteen drops, as may be required. *Ellis.*

## ETHEREAL TINCTURE OF IODINE.

R. Iodine two scruples  
 Sulphuric ether one fl. ounce and a-half  
 Dissolve, and filter. *Magendie.*

R. Iodine six grains  
 Sulphuric ether two fl. drachms  
 Dissolve. Dose, ten drops, two or three times a-day, much diluted. In diseases of the skin or glands. *Ellis.*

## MIXTURE OF IODINE AND IODIDE OF POTASSIUM.

R. Iodine three grains  
 Iodide of potassium six grains  
 Distilled water one fl. ounce  
 Dissolve. Dose, six to twenty drops, thrice a-day, in a draught of cold water. In glandular disease and affections of the mucous membranes. *Morton.*

## BATH WITH IODINE.

For adults. 1 2 3  
 R. Iodide of potassium 231 grs. 308 grs. 370 grs.  
 Iodine 123 " 154 " 184 "  
 Water twenty fl. ounces  
 Dissolve, and add to bath of sixty gallons.

## For children.

R. Iodide of potassium 77 grs. 92 grs. 107 grs.  
 Iodine 38 " 46 " 61 "  
 Water ten fl. ounces  
 Dissolve, and add to bath of sixty gallons. Used in scrofulous affections. *Lugol.*

## SOLUTION OF IODINE.

For internal use. 1 2 3  
 R. Iodine  $\frac{1}{2}$  gr.  $\frac{2}{3}$  gr. 1 gr.  
 Common salt 12 grs. 12 grs. 12 grs.  
 Distilled water one pint

## For external use.

R. Iodine 2 grs. 3 grs. 4 grs.  
 Distilled water one pint  
 As a lotion. *Lugol.*

## LOTION OF IODINE.

R. Tincture of iodine half fl. ounce  
 Iodide of iron twelve grains  
 Chloride of antimony half an ounce  
 Mix. As an application to corns, by means of a camel's hair pencil, after the corn has been pared. *Henderson.*

R. Tincture of iodine one fl. drachm  
 Black drop two fl. drachms  
 Oil of almonds  
 Lime water, each, two fl. ounces  
 Mix carefully. As a lotion for chilblains. *Cadet.*

R. Iodine one to two grains  
 Alcohol one fl. drachm  
 Water one pint  
 Mix. To wash scrofulous ulcers. *Foy.*

## MIXTURE OF IODINE.

R. Tincture of iodine ninety drops



Decoction of Peruvian bark ten fl. ounces  
 Mix. Two spoonfuls, three times a-day.

*Radiis.*

R. Tincture of iodine ten drops  
 Water of soapwort four fl. ounces  
 Syrup of orange-flowers one fl. ounce

Mix. In spoonful doses. In the treatment of scrofulous ulcers.

*Foy.*

#### CLYSTER OF IODINE.

R. Gum Arabic four drachms  
 Water six fl. ounces

Dissolve, and add

Tincture of iodine five drops

In amenorrhœa, dysmenorrhœa, &c.

*Cadet.*

#### GARGLE OF IODINE.

R. Tincture of iodine one to two fl. drachms

Tincture of opium one fl. drachm

Water six fl. ounces

Mix. As a gargle, in ulceration of the tonsils.

*Ross.*

#### OINTMENT OF IODINE.

R. Iodine one scruple

Alcohol twenty minims

Lard one ounce

Rub the iodine with the alcohol, and then with the lard, till thoroughly mixed.

*U. S. Ph.*

In goitre, scrofulous swellings, &c.

R. Iodine one scruple

Rose ointment one ounce

Mix. To be applied to the tonsils, night and morning, by means of a brush. Not to be used till all inflammation is subdued.

*Cerchiari.*

R. Iodine one part

Lard sixteen parts

Rub together carefully.

*Guibourt.*

#### COMPOUND OINTMENT OF IODINE.

R. Iodine half a drachm

Iodide of potassium one drachm

Alcohol one fl. drachm

Lard two ounces

Rub the iodine and iodide of potassium with the alcohol, and then with the lard, till thoroughly mixed.

*U. S. Ph.*

Used in same cases as the iodine ointment, but is more active.

#### IODINE OINTMENT WITH OIL OF TOBACCO.

R. Iodine twelve grains

Iodide of potassium four scruples

Oil of tobacco fifty drops

Lard two drachms

Mix. To relax rigid muscles. *Dover.*

#### IODINE OINTMENT AND MERCURY.

R. Mercurial ointment sixteen parts

Iodine three parts

Mix. In frictions, in ovarian dropsy.

*Radiis.*

#### LINIMENT OF IODINE.

R. Liniment with opium one fl. ounce

Tincture of iodine one fl. drachm

Mix. *Manson.*

#### CATAPLASM OF IODINE.

R. Tincture of iodine half fl. ounce

Flaxseed meal one ounce

Oatmeal three ounces

Distilled water sufficient

Make a cataplasm. Used as an application to scrofulous tumors and goitre.

*Dunglison. N. Rem.*

#### IODINE PLASTER.

R. Iodine one drachm

Iodide of potassium one scruple

Water half a fl. drachm

Triturate in a mortar, and add to

Burgundy pitch three ounces

Wax

Suet, each, half an ounce

Melted together, and triturate till well mixed.

## IPECACUANHA.

### IPECACUANHA.

The root of *Cephalis ipecacuanha*, a small herbaceous plant, a native of Brazil.

*Sex. Syst.* Pentand. monog. *Nat. Syst.* Cinchonaceæ.

Richard, Bull. Med. iv. 92. Griffith, Med. Bot. 357.

This root is about the size of a large quill, of an irregular, twisted shape, apparently formed of small, unequal rings, about a line in height, separated by narrow interstices; when broken, it presents two distinct parts, a thin ligneous centre, and a thick cortical layer of a brownish color, and somewhat resinous appearance, having a herbaceous, sub-acrid, somewhat bitter taste, and nauseous odor. Externally it presents some varieties of color, but is usually brown. It is emetic, and sometimes purgative in full doses, and diaphoretic and expectorant in small, but repeated quantities. As an emetic, the dose is fifteen to twenty grains; as a nauseant, two to four grains; as an expectorant, one to two grains.

#### POWDER OF IPECACUANHA AND OPIUM, DOVER'S POWDER.

R. Ipecacuanha

Opium, each, in powder, one drachm

Sulphate of potassa one ounce

Triturate thoroughly. *U. S. Ph.*



R. Ipecacuanha  
Liquorice root  
Extract of opium, each,  
in powder, one ounce  
Sulphate of potassa  
Nitrate of potassa, each, four ounces  
Triturate well. *Par. Cod.*

This latter formula approaches nearest to the original formula of Dover, but is not superior to the first, now generally used. Dose, five to ten grains, as a diaphoretic.

#### POWDER OF IPECACUANHA AND TARTAR EMETIC.

R. Powdered ipecacuanha one scruple  
Tartar emetic one grain  
Mix. As an emetic, or in divided doses, as a nauseant and sudorific. *Ellis.*

#### POWDER OF IPECACUANHA AND RHUBARB.

R. Powdered ipecacuanha  
rhubarb, each, one scruple  
Mix. As an emeto-cathartic in full doses; or tonic alterative, in doses of three or four grains. *Beasley.*

#### POWDER OF IPECACUANHA AND MURIATE OF AMMONIA.

R. Powdered ipecacuanha a quarter grain  
Muriate of ammonia twelve grains  
Extract of liquorice  
Sugar of milk, each, six grains  
Mix. To be taken every two hours, as an expectorant, on the decline of pneumonia. *Kopp.*

#### POWDER OF IPECACUANHA AND ANTIMONY.

R. Ipecacuanha  
Golden sulphuret of antimony, each, one grain  
Camphor three grains  
Gum Arabic one scruple  
Triturate well. To be taken every two hours, in deficient expectoration in pneumonia. *Phœbus.*

#### POWDER OF IPECACUANHA AND CARBONATE OF SODA.

R. Powdered ipecacuanha  
opium, each, one grain  
Carbonate of soda twelve grains  
Mix, and rub well together. To be taken every eight hours, in spasmodic asthma, whooping-cough, &c. *A. T. Thomson.*

#### POWDER OF IPECACUANHA AND NITRE.

R. Dover's powder one drachm  
Nitrate one ounce  
Mix. To be taken in divided doses. *Beasley.*

R. Powdered ipecacuanha six grains  
nitrate of potassa  
one scruple and a-half  
myrrh twelve grains  
Mix, and divide into four powders; one to be taken every four hours. Found useful in asthma, and the earlier stages of phthisis. *A. T. Thomson.*

#### POWDER OF IPECACUANHA AND CALOMEL.

R. Powdered ipecacuanha six grains  
opium three grains  
Calomel one grain and a-half  
Powdered nitre half a drachm  
Mix, and make six powders. One, every three or four hours, as a diaphoretic. *Ellis.*

#### POWDER OF IPECACUANHA AND MYRRH.

R. Powdered ipecacuanha six grains  
myrrh twelve grains  
nitre half a drachm  
Mix, and make four powders. One, every four hours, as a stimulating expectorant. *Paris.*

#### POWDER OF IPECACUANHA AND TRAGACANTH.

R. Powdered ipecacuanha five grains  
Compound tragacanth powder ten grains  
Mix, make a powder, to be taken every four or six hours. Useful at the commencement of febrile diseases, after clearing the stomach and bowels. *A. T. Thomson.*

#### EXTRACT OF IPECACUANHA.

R. Coarsely powdered ipecacuanha two parts  
Proof spirit seven parts  
Treat by displacement; distil off the spirit and evaporate, on water-bath, to proper consistence. *Paris Cod.*

#### BOLUS OF IPECACUANHA.

R. Powdered ipecacuanha two grains  
Conserve of opium one scruple  
Prepared chalk sufficient  
Mix, and make eight boluses. Two, night and morning, in dysentery. *St. Marie.*  
R. Powdered ipecacuanha two grains  
Aromatic powder ten grains  
Syrup of cinnamon sufficient  
Mix, and make two boluses. One, in the evening, in dyspepsia. *Cadet.*

#### PILLS OF IPECACUANHA AND OPIUM.

R. Dover's powder three parts  
Confection of roses one part  
Mix, and make pills of four grains each. *Ed. Ph.*



## COMPOUND PILLS OF IPECACUANHA.

R. Dover's powder three drachms  
Powdered squill  
Gum ammoniac, each, one drachm  
Mucilage sufficient  
Mix, and form mass. Dose, fifteen to twenty grains, as a diaphoretic.

*Lond. Ph.*

## PILLS OF IPECACUANHA AND FOXGLOVE.

R. Powdered Ipecacuanha  
foxglove, each, half a drachm  
opium five grains  
Extract of elecampane sufficient  
Mix, and make sixty pills. One, every three hours, in spasmodic asthma. *Heim.*

## PILLS OF IPECACUANHA AND CENTAURY.

R. Powdered ipecacuanha  
Extract of centaury, each, fifteen grains  
Powdered rhubarb half a drachm  
Mix, and make thirty pills. One, night and morning, in dyspepsia. *Radius.*

## LOZENGES OF IPECACUANHA.

R. Powdered ipecacuanha half an ounce  
sugar fourteen ounces  
Arrow root four ounces  
Mucilage of tragacanth sufficient  
Mix the powders thoroughly, and form into a mass with the mucilage, and divide into lozenges of ten grains each. *U. S. Ph.*

A good expectorant in catarrh. Each contains a quarter of a grain of ipecacuanha.

## LOZENGES OF IPECACUANHA AND CAMPHOR.

R. Powdered ipecacuanha fifteen grains  
camphor  
sugar, each, one drachm  
Mucilage of tragacanth sufficient  
Mix, and make sixty lozenges. *Beasley.*

R. Powdered sugar five hundred parts  
Flake manna one hundred and twenty-five parts  
Powdered ipecacuanha eighteen parts  
Lactucarium eight parts  
Powdered squill four parts  
Mucilage of tragacanth sufficient  
Make into a uniform paste, and divide into lozenges of ten grains. Five or six a-day, in catarrh, &c. *Grunn.*

## CONFECTION OF IPECACUANHA.

R. Powdered ipecacuanha twelve grains  
Flowers of sulphur one scruple  
Powdered orris one drachm  
Syrup of mallows  
Manna, each, two ounces  
Mix. A teaspoonful, two or three times a-day, in whooping-cough. *Bories.*

## DECOCTION OF IPECACUANHA.

R. Bruised ipecacuanha two drachms  
Water twelve ounces  
One-third of the water is to be used at a time, so as to make three decoctions; these are to be united, and should amount to six ounces. Celebrated in dysentery; one-third to be taken at a dose. *Soubeiran.*

## MIXTURE OF IPECACUANHA.

R. Powdered ipecacuanha half a drachm  
Tartar emetic one grain  
Tincture of squill one fl. ounce  
Distilled water seven fl. ounces and a-half  
Mix. Four tablespoonfuls; then two every ten minutes, till vomiting is produced.

*Ellis.*

R. Powdered ipecacuanha five grains  
Syrup of poppies one drachm  
Spirit of cinnamon two drachms  
Chalk mixture thirteen drachms  
Mix. As a draught in dysentery. *Hooper.*

R. Powdered ipecacuanha two grains  
Carbonate of soda ten grains  
Syrup of poppies one ounce  
Mint water six ounces  
Mix. In spoonfuls, to check spasmodic vomiting. *St. Marie.*

R. Ipecacuanha one drachm  
Senna two drachms  
Boiling water six fl. ounces  
Infuse for twelve hours, strain, and add  
Oxymel of squills  
Syrup of hyssop, each, one ounce  
Mix. In spoonfuls, in whooping-cough. *Guibourt.*

## SYRUP OF IPECACUANHA.

R. Ipecacuanha, in coarse powder, one ounce  
Diluted alcohol one pint  
Syrup two pints  
Macerate the ipecacuanha in the alcohol for fourteen days, and filter. Evaporate to two fl. ounces, and filter; mix with the syrup, and evaporate on a water-bath to proper consistence. *U. S. Ph.*

As emetic for children, one to two fl. drachms; as expectorant, five to twenty minims.

## COMPOUND SYRUP OF IPECACUANHA.

R. Ipecacuanha one drachm  
Orris root two drachms  
Red Peruvian bark half an ounce  
Seneca  
Iceland moss, each, two ounces  
Sugar two pounds  
Water sufficient  
Mix, and make a syrup. In spoonful doses, as an expectorant. *Courtois.*



R. Syrup of ipecacuanha  
 poppies, each, two ounces  
 orange flowers one ounce  
 Oxy-mel of squill one ounce and a-half  
 Mix. Two spoonfuls every hour, in hoop-  
 ing cough. *Cadet.*

## WINE OF IPECACUANHA.

R. Bruised ipecacuanha two ounces  
 Sherry wine two pints  
 Macerate for fourteen days, express, and  
 filter. *U. S. Ph.*

As emetic and diaphoretic. Dose, for an  
 adult, as emetic, one fl. ounce; for a child, one  
 fl. drachm.

## COMPOUND WINE OF IPECACUANHA.

R. Ipecacuanha four parts  
 Sugar two parts  
 Diluted alcohol twenty-four parts  
 Digest for eight days, and add  
 Sherry wine ninety-six parts  
 Star anise one part  
 Sugar four parts  
 Filter at the end of six days. *Niemann.*

WINE OF IPECACUANHA AND TARTAR  
EMETIC.

R. Wine of ipecacuanha one fl. ounce  
 Tartar emetic one grain  
 Mix. A teaspoonful, every ten or fifteen  
 minutes, till vomiting is induced. *Ellis.*

## ALKALINE WINE OF IPECACUANHA.

R. Wine of ipecacuanha forty minims  
 Tincture of opium eight drops  
 Carbonate of soda sixteen grains  
 Syrup two fl. drachms  
 Water fourteen fl. drachms  
 Mix. A teaspoonful, to children, in hoop-  
 ing cough. *Pearson.*

## ELIXIR OF IPECACUANHA.

R. Powdered ipecacuanha  
 Balsam of tolu, each, half an ounce  
 Flowers of benzoin  
 Opium  
 Saffron, each, two drachms  
 Oil of anise one drachm  
 Camphor two scruples  
 Alcohol two pints  
 Digest, express, and filter. Dose, one to  
 two drachms, as a stimulating sudorific.  
*Cadet.*

## IRIS FLORENTINA.

## FLORENTINE IRIS. ORRIS.

The Florentine Iris is a native of the southern  
 parts of Europe, and affords a root, or rhizome,  
 known as orris root.

*Sex. Syst.* Triand. monog. *Nat. Syst.* Irida-  
 ceæ.

*Linn. Sp. Pl.* 55. *Griffith, Med. Bot.* 625.

The rhizome is horizontal, knotty, and frag-  
 rant; in a recent state it is acrid, and somewhat  
 bitter; this acrimony disappears in a great mea-  
 sure on dessication. It is seldom used in medi-  
 cine, but forms an ingredient in numerous tooth-  
 powders, &c.

## DENTIFRICE WITH ORRIS ROOT.

R. Cream of tartar two ounces  
 Powdered orris root one ounce  
 myrrh  
 kino, each, half an ounce  
 Mix well. *Redwood.*

R. Powdered Castile soap  
 orris root, each, two ounces  
 cuttlefish bone  
 prepared chalk, each,  
 three ounces

Oil of cloves  
 Essence of lemons, each, twenty drops  
 Mix. *Redwood.*

## J

## JALAPA.

## JALAP.

Jalap is the root of the *Ipomœa jalapa*, a native  
 of Mexico.

*Sex. Syst.* Pentand. monog. *Nat. Syst.* Con-  
 volvulacæ.

*Coxe, Am. Journ. Med. Sci.* 1830. *Griffith,*  
*Med. Bot.* 474.

The root, when dried, is in pieces of various  
 forms and size; usually entire, but sometimes

in slices. The epidermis is thin and brown;  
 within it is grayish, with brown, concentric rings.  
 The powder is of a brownish yellow, with a pe-  
 culiar, unpleasant odor, and a sweetish, sub-acrid,  
 nauseous taste. It is an active purgative, in doses  
 of ten to thirty grains.

## POWDER OF JALAP AND CALOMEL.

R. Powdered jalap ten grains  
 Calomel five grains



Mix. To be given in syrup or molasses.  
As a purgative. *Ellis.*

R. Powdered jalap twelve grains  
Calomel three grains  
Sulphate of potassa seven grains  
Mix; make a powder, to be taken at night;  
a useful purge in diminishing the action of  
the liver. *A. T. Thomson.*

#### POWDER OF JALAP AND CREMOR TAR- TAR.

R. Powdered jalap one drachm  
Cremor tartar two drachms  
Mix, and divide into six powders. One  
every two hours, as a hydragogue purga-  
tive. *Ellis.*

R. Jalap two ounces  
Cremor tartar one ounce  
Sugar thirteen ounces  
Oil of orange peel two drachms

Mix. Dose, one to two drachms. *Cadet.*

R. Powdered jalap one ounce  
Cremor tartar two ounces

Mix. *U. S. Ph.*

R. Powdered jalap three ounces  
Cremor tartar six ounces  
Ginger two drachms

Mix. *Lond. Ph.*

Dose, thirty grains to a drachm.

#### POWDER OF JALAP AND IPECACUANHA.

R. Jalap fifteen grains  
Ipecacuanha five grains  
Oil of cinnamon two drops

Mix. *Paris.*

#### POWDER OF JALAP AND SCAMMONY.

R. Jalap one drachm  
Scammony one scruple  
Ipecacuanha ten drachms

Mix, and divide into powders of eight  
grains. In mucous discharges. *Dumas.*

#### COMPOUND POWDER OF JALAP.

R. Powdered jalap three ounces  
Cremor tartar six ounces  
Powdered ginger two drachms

Mix. Half a drachm to one drachm, as  
a hydragogue purgative. *Lond. Ph.*

#### EXTRACT OF JALAP.

R. Jalap, in coarse powder, one pound  
Alcohol four pints  
Water sufficient

Macerate the jalap in the alcohol for four  
days, then exhaust by the process of dis-  
placement; first by alcohol, then by wa-  
ter. Distil off the alcohol from the tinc-  
ture, and evaporate the infusion, till both  
are of the consistence of thin honey, mix

them, and evaporate to proper consist-  
ence. *U. S. Ph.*

Dose, from ten to twenty grains.

#### RESIN OF JALAP.

R. Coarsely powdered jalap at will  
Moisten thoroughly with rectified spirit,  
put in a displacement apparatus and ex-  
haust, distil off the spirit, and evaporate  
to due consistence. *Ed. Ph.*

Dose, four to twelve grains.

#### ALKALINE EXTRACT OF JALAP.

R. Powdered jalap two pounds  
Alcohol (.921) one gallon  
Digest for some days, strain, and filter;  
evaporate slowly on the water-bath to a  
pilular consistence, adding gradually

Carbonate of potassa twelve drachms  
Dose, three to nine grains. *Durand.*

#### COMPOUND PILLS OF JALAP.

R. Jalap  
Rhubarb  
Aloes, each, one ounce  
Soap two ounces  
Syrup of ginger sufficient

Mix, and form mass. Dose, twenty to  
thirty grains. *Foy.*

#### PILLS OF JALAP AND CALOMEL.

R. Resin of jalap  
Soap, each, one drachm  
Oil of orange peel six drops  
Mix, and make sixty pills. *Alibert.*

#### ELECTUARY OF JALAP.

R. Powdered jalap  
Black sulphuret of  
mercury, each, two drachms  
Scammony one drachm  
Resin of jalap  
Squills, each, one drachm and a-half  
Syrup of buckthorn sufficient

Mix. Dose, half a drachm to a drachm,  
in dropsy or lead colic. *Cadet.*

R. Powdered jalap two drachms  
Cremor tartar one ounce and a-half  
Ginger one scruple  
Molasses two ounces and a-half

Mix. Dose, two drachms. *Beasley.*

R. Powdered jalap  
Cremor tartar  
Nitrate, each, half an ounce  
Confection of senna one ounce  
Simple syrup sufficient

Mix. A bolus the size of a hazelnut,  
three or four times a-day, in hemorrhoids.  
*Ellis.*

#### EMULSION OF JALAP.

R. Resin of jalap nine grains



Soap three grains  
Olive oil one ounce and a-half  
Triturate well. Said to be useful in dysentery. A tablespoonful, occasionally.  
*Vogel.*

## MIXTURE OF JALAP.

R. Infusion of senna six fl. ounces  
Extract of jalap half a drachm  
Tartrate of potassa four drachms  
Oil of ginger twenty minims  
Mix. *Beasley.*

## SOAP OF JALAP.

R. Resin of jalap  
Soap equal parts  
Dissolve in  
Alcohol sufficient  
Evaporate to pilular consistence. Dose, ten to fifteen grains. *Pruss. Ph.*

## TINCTURE OF JALAP.

R. Powdered jalap eight ounces  
Diluted alcohol two pints  
Macerate for fourteen days, express, and filter. *U. S. Ph.*

Seldom given alone, but added to cathartic mixtures, in portions of one to two fl. drachms.

## COMPOUND TINCTURE OF JALAP.

R. Jalap six drachms  
Black hellebore three drachms  
Juniper berries half an ounce  
Rasped guaiacum half an ounce  
Brandy twenty-four fl. ounces  
Digest for three days, express, and filter. *Ed. Ph. 1744.*

R. Jalap eight ounces  
Scammony two ounces  
Turberth one ounce  
Brandy twelve pints  
Digest for twelve hours, on a water-bath, filter, and add  
Syrup of senna four pounds  
Mix. *Giordano.*

## MIXTURE OF TINCTURE OF JALAP.

R. Tincture of jalap two fl. drachms  
Vinegar of squill one fl. drachm  
Mint water one fl. ounce  
Mix. To be taken three times a-day. *Ellis.*

## JUGLANS CINEREA.

## BUTTERNUT.

The Butternut, or White Walnut, is an indigenous tree, with numerous, nearly horizontal branches, found in many places in the United States.

*Sex. Syst. Monœc. polyand. Nat. Syst. Juglandaceæ.*

Linn. Sp. Pl. 1415. Griffith, Med. Bot. 589.

The part used is the inner bark, especially of the root; this is fibrous, has little odor, but a peculiar bitter, somewhat acrid taste. It is, at first, white, but gradually changes to a dark brown. It is a mild cathartic, and is well suited to cases of costiveness and bowel affections. It is usually given in extract.

## EXTRACT OF BUTTERNUT.

R. Butternut, in coarse powder, one pound  
Water sufficient  
Mix the butternut with a pint of the water, macerate for twenty-four hours, introduce into a displacement apparatus, and exhaust. Heat the fluid to a boiling point, strain, and evaporate to proper consistence. *U. S. Ph.*

Dose, from five to ten grains, as a laxative; from twenty to thirty, as a purgative.

## JUNIPERUS.

## JUNIPER.

The common Juniper, *J. communis*, is an erect, evergreen shrub, sometimes attaining a large size, found in Europe, and Northern Asia and America. The tops and the berries are officinal. *Sex. Syst. Diœc. monadelph. Nat. Syst. Pinales.*

Linn. Sp. Pl. 1470. Griffith, Med. Bot. 608.

The berries are oblong globular, marked with three radiating furrows at top, and contain three seeds. They are of a purplish black color, have a sweetish, somewhat terebinthinate taste, and a peculiar, aromatic odor. They are used as stimulating diuretics, especially as adjuvants to more active remedies. The tops have a balsamic odor, and a resinous, bitterish taste, with the same medicinal properties as the berries.

## EXTRACT OF JUNIPER.

R. Bruised juniper berries one pound  
Distilled water three pints  
Macerate at 77° to 86° for twenty-four hours; strain with gentle pressure, add more water to the berries, and let stand for twelve hours. Filter, and evaporate to proper consistence. Dose, one to three drachms. *Par. Cod.*

## INFUSION OF JUNIPER.

R. Juniper berries, bruised, one ounce  
Boiling water one pint  
Infuse, and strain. To be taken during the twenty-four hours. *Radius.*

## COMPOUND INFUSION OF JUNIPER.

R. Juniper berries, bruised,  
Mustard seed  
Ginger, each, half an ounce  
Horseradish, bruised,  
Parsley root, bruised, each, one ounce  
Cider two pints



Infuse, and strain. A wineglassful, two or three times a-day, in general dropsy.

*Parrish.*

R. Juniper berries, bruised, one ounce and a-half  
Cremor tartar one ounce  
Boiling water one pint

Infuse. A teacupful, several times a-day, with forty drops of sweet spirit of nitre. In dropsy.

*Ellis.*

R. Juniper berries, bruised, two ounces and a-half  
Boiling water one pint

Infuse for two hours, and strain; add

Compound spirit of juniper ten fl. drachms

Dose, two fl. ounces, three times a-day.

*Beasley.*

#### ELECTUARY OF JUNIPER.

R. Extract of juniper  
Clarified honey, each, one pound  
Juniper berries  
Cinnamon

Orange peel, each, half an ounce

Mix. Stomachic and diuretic. Dose, one to four drachms.

*Saxon Ph.*

#### MIXTURE WITH OIL OF JUNIPER.

R. Oil of juniper half a drachm  
Nitric ether  
Tincture of fox-glove, each, three drachms

Mix. Twenty to thirty drops, every three hours.

*Hufeland.*

R. Oil of juniper eight drops  
Sugar

Gum Arabic, each, a drachm and a-half

Distilled water four fl. ounces

Mix. A tablespoonful, every hour or two, as a diuretic.

*Ellis.*

#### COMPOUND SPIRIT OF JUNIPER.

R. Juniper berries, bruised, one pound  
Carraway, bruised,  
Fennel, bruised, each,

one ounce and a-half

Diluted alcohol one gallon

Water two pints

Macerate in the alcohol, for twenty-four hours, add the water, and with a slow fire, distil one gallon.

*U. S. Ph.*

Dose, two to four fl. drachms, as an addition to diuretic mixtures, &c., in debilitated cases of dropsy.

#### LINIMENT OF JUNIPER.

R. Oil of juniper one ounce and a-half  
Oil of anise six drops  
Lard two ounces

Mix well; as an application in tinea capitis.

*Sully.*

R. Spirit of juniper two ounces

Oil of cloves

Butter of nutmeg, each, half a drachm

Mix. As a friction in ecchymosis, and indolent tumors.

*Augustin.*

#### JUNIPERUS VIRGINIANA.

##### RED CEDAR.

The tops of the red cedar are officinal. This is an evergreen tree, sometimes attaining a great size, with small, fleshy, closely imbricated leaves. It is found in all parts of the United States, but is most abundant and vigorous in the south.

*Sax. Syst. Dicc. monadelph. Nat. Syst. Pinaceæ.*

*Linn. Sp. Pl. 1471. Griffith, Med. Bot. 609.*

The leaves or tops have a peculiar and somewhat aromatic odor, a bitterish and somewhat acrid taste. They have much the same properties as those of savine, but are not as active, though often used as a substitute for them.

#### K

#### KINO.

##### KINO.

This is an extract obtained from several different plants, but that most used and esteemed is the product of *Pterocarpus marsupium*, a large tree, indigenous to some parts of India.

It is usually in small, angular fragments, of a dark, brownish-black color, opaque and shining, of a brittle and friable consistence, an astringent and slightly bitter taste; adhering to the teeth

when chewed, and tinging the saliva of a red color. It is a powerful astringent, and much used in the treatment of mucous discharges, &c., and as a local application in gonorrhœa, to indolent ulcers, relaxed uvula, &c. The dose in powder is from ten to thirty grains.

#### COMPOUND POWDER OF KINO.

R. Powdered kino fifteen drachms  
opium one drachm  
cinnamon four drachms



Rub well together. Dose, five to twenty grains. *Lond. Ph.*

As an anodyne astringent, in diarrhœa and hemorrhage of the bowels.

R. Powdered kino  
Sugar of milk, each, one scruple  
Mix. To be taken every three or four hours, in cardialgia. *Radius.*

COMPOUND PILLS OF KINO.

R. Powdered kino one scruple  
opium two grains  
Mucilage of gum Arabic sufficient  
Mix, and make four pills. One, every four hours, in diarrhœa, pyrosis, &c. *Ellis.*

R. Powdered kino two scruples  
camphor thirty grains  
Aromatic powder one scruple  
Oxide of zinc ten grains  
Mucilage of gum Arabic sufficient  
Mix, and make twenty pills. Two, morning and evening, in diabetes. *Augustin.*

ELECTUARY OF KINO.

R. Powdered kino six drachms  
alum  
cinnamon, each, two drachms  
Syrup sufficient  
Mix into an electuary. One drachm, two or three times a-day, in diarrhœa, chronic dysentery, &c. *Swediaur.*

INFUSION OF KINO.

R. Kino two drachms  
Boiling water eight. fl. ounces  
Infuse, and strain when cool. Dose, a fl. ounce. *Wood.*

INJECTION OF KINO.

R. Kino one to two drachms  
Alum twenty to thirty grains  
Boiling water two pints  
Infuse for an hour, and strain. As an injection in chronic urethritis. *Swediaur.*

SYRUP OF KINO.

R. Kino two ounces  
Tepid water three pints  
Dissolve, and add  
Sugar four pounds  
Clarify, and evaporate to consistence of syrup. *Taddei.*

TINCTURE OF KINO. *X*

R. Powdered kino three ounces and a-half  
Rectified spirit two pints  
Macerate for fourteen days, and filter. *Lond. Ph.*

Dose, one to two fl. drachms, generally as an adjuvant to astringent mixtures.

KRAMERIA.

RHATANY.

The roots of several species of *Krameria* are possessed of analogous properties, but the only one that is officinal, is that of *K. triandra*, a small shrub found in Peru, in dry, sandy places. The stem is procumbent, and much branched, with sessile, hairy leaves, and dark red flowers. The root is long and branching.

*Sex. Syst.* Tetrand. monog. *Nat. Syst.* Polygalaceæ.

Ruiz & Pavon, Fl. Per. i. 93. Griffith, Med. Bot. 227.

The dried root is of a dark, reddish-brown color, with no odor, but a bitter and astringent taste. The small roots are the most powerful. It is somewhat tonic, and powerfully astringent. It is used for most purposes to which the vegetable astringents are applied. The dose in powder is from twenty to thirty grains.

COMPOUND POWDER OF RHATANY.

R. Extract of rhatany  
Catechu, each, ten grains  
Alum four grains  
Powdered liquorice ten grains  
Mix. To be taken three or four times a-day, in diarrhœa, &c. *Vogt.*

DENTRIFICE OF RHATANY.

R. Extract of rhatany one to two drachms  
myrrh one drachm  
Powdered orris six drachms  
Oil of cinnamon two drops  
myrrh ten drops  
Balsam Peru ten grains  
Mix well. *Phæbus.*

EXTRACT OF RHATANY.

R. Prepared like extract of gentian. *U. S. Ph.*

Dose, ten to twenty grains.

PILLS OF RHATANY.

R. Extract of rhatany one scruple  
Powdered kino sufficient  
Mix, and make eight pills. One, every hour or two, to check mucous discharges. *Ellis.*

R. Extract of rhatany one drachm  
rhubarb two scruples  
Ginger one scruple  
Mix, and make twenty-four pills. *Reece.*

ELECTUARY OF RHATANY.

R. Powdered rhatany one ounce  
ginger one drachm  
Syrup of orange peel sufficient  
Mix. A teaspoonful every two or three hours. *Ferrara Ph.*

INFUSION OF RHATANY.

R. Bruised rhatany one ounce



Boiling water one pint  
Infuse for four hours, and strain.

*U. S. Ph.*

Dose, one to two fl. ounces.

#### COLLUTORY OF RHATANY.

R. Extract of rhatany half a drachm  
Mucilage of quince

seeds two drachms and a-half

Sage water one ounce and a-half

Mix. As a mouth-wash in aphthæ, &c.

*Radius.*

#### CLYSTER OF RHATANY.

R. Extract of rhatany one scruple

Mucilage of quince

seeds three drachms

Infusion of sage one fl. ounce and a-half

Mix. To check diarrhœa, &c. *Ammon.*

#### INJECTION OF RHATANY.

R. Extract of rhatany half an ounce

Tincture of catechu

kino, each,

one drachm and a-half

To be added to a decoction of

Rhatany ten drachms

Water three pints

Reduced to two-thirds, and strained. As  
an injection in leucorrhœa, and prolapsus  
of the uterus. *Cadet.*

#### MIXTURE OF RHATANY.

R. Extract of rhatany one drachm

Syrup of poppies

Rose water, each, two fl. ounces

Mix. A teaspoonful occasionally, in pas-  
sive hemorrhage and chronic dysentery.

*Fouquier.*

R. Extract of rhatany one drachm

Chamomile water six fl. ounces

Elixir of vitriol half a drachm

Mix. As the last. *Hildebrand.*

#### SYRUP OF RHATANY.

R. Extract of rhatany two ounces

Water one pint

Sugar two pounds and a-half

Dissolve the extract in the water, add the  
sugar, and form syrup. *U. S. Ph.*

Dose, half fl. ounce.

#### TINCTURE OF RHATANY.

R. Powdered rhatany six ounces

Diluted alcohol two pints

Macerate for fourteen days, express, and  
filter. *U. S. Ph.*

Dose, one to two fl. drachms.

#### COMPOUND OINTMENT OF RHATANY.

R. Burgundy pitch eight parts

Venice turpentine two parts

White wax one part

Extract of rhatany two parts

Alum one part

Melt the first three ingredients, over a  
gentle fire, and add the rhatany and alum,  
in fine powder, and incorporate well.

Useful where an astringent application is in-  
dicated.

## L

### LAC.

#### MILK.

The milk of several animals has been used  
medicinally, but that generally employed is from  
the cow. It is mostly employed in the form of  
whey, and of sugar of milk.

#### SIMPLE WHEY.

R. Milk three pints

Prepared rennet sufficient

Cover up the mixture, and place before the  
fire, till a uniform curd is formed; divide  
this with a spoon, and strain. A good  
diluent in febrile affections; when boiled  
down to one half, it is nutritive.

*A. T. Thomson.*

#### VINEGAR WHEY.

R. Milk one pint

Vinegar half fl. ounce

Boil a few minutes, and separate the curd.

*Ellis.*

#### TAMARIND WHEY.

R. Boiling milk one pint

Tamarinds two ounces

Boil till coagulation is effected, and strain.

*Plenck.*

#### CREMOR TARTAR WHEY.

R. Boiling milk one quart

Cremor tartar half an ounce

Boil until the curd separates, and strain.

*Plenck.*



## ALUM WHEY.

R. Clear whey one pint  
Alum two drachms  
Dissolve. A glassful occasionally, in passive hemorrhages. *Foy.*

## AROMATIC ALUM WHEY.

R. Powdered alum one drachm  
Sugar of milk half a drachm  
Powdered cinnamon fifteen grains  
Milk two pints  
Boil, and strain when coagulated. In passive uterine hemorrhages. To be taken during the day. *Niemann.*

## MUSTARD WHEY.

R. Bruised mustard seed half an ounce  
Milk one pint  
Boil till the milk is coagulated, and strain. A cupful, as a diuretic, in dropsical affections. *A. T. Thomson.*

## WINE WHEY.

R. Boiling milk half a pint  
White wine one to two fl. ounces  
Sugar sufficient  
Boil for ten minutes, constantly stirring, then strain. A good mode of giving wine as a stimulant. Dose must depend on circumstances. *Ellis.*

## POWDER OF MILK.

R. Milk two pints  
Powdered carbonate of soda half a drachm  
Water one fl. ounce  
Sugar one pound  
Dissolve the soda in the water, add the solution to the milk, gently heat, and stir. When evaporated to one-third, add the sugar gradually, and stir. Remove from fire, pour into plates, and dry in an oven. When dry, powder, and keep in well stoppered bottles. One to two ounces, mixed with a quart of water, forms an agreeable drink, and is a good substitute for milk. *Legrip.*

## ARTIFICIAL GOATS' MILK.

R. Fresh suet one ounce  
Cut in small pieces, and tie in a muslin bag, and boil in  
Milk two pints  
Sugar candy two drachms  
A good diet in scrofulous emaciation, and in latter stages of phthisis. *A. T. Thomson.*

## MIXTURE OF MILK AND SUET.

R. Sheeps' suet two ounces  
Milk one pint  
Starch half an ounce

Boil slowly for half an hour. As a drink, in dysentery and cholera infantum. *Ellis.*

## MILK AND SODA WATER.

R. Milk a teacupful  
Heat nearly to boiling, and add  
Sugar a teaspoonful  
Put in a large tumbler, and add  
Soda water sufficient  
A good mode of taking milk when there is much acidity of stomach. *A. T. Thomson.*

## SUGAR OF MILK.

R. Plain whey at will  
Clarify with white of egg, and evaporate on water-bath, to crystallization, and set aside in a cool place. *Redwood.*

Cooling and laxative, in doses of two drachms to one ounce.

## SYRUP OF MILK.

R. Skimmed milk twelve pounds  
Evaporate to one-half, and add  
Sugar nine pounds  
Cherry laurel water three ounces  
*Guibourt.*

## MIXTURE OF SUGAR OF MILK AND ICELAND MOSS.

R. Iceland moss four drachms  
Water one pint and a-half  
Reduce to one-half by boiling, strain, and add  
Sugar of milk two drachms  
Milk one pint  
Syrup of asparagus three ounces  
A cupful, occasionally, in chronic catarrh. *Cadet.*

## MIXTURE OF SUGAR OF MILK AND GUM ARABIC.

R. Sugar of milk  
Sugar, each, two pounds  
Gum Arabic three pounds  
Extract of dog grass, seven ounces  
Mix. Half an ounce dissolved in a quart of water, forms a refreshing and demulcent drink. *Cadet.*

## SUGAR OF MILK DRAUGHT.

R. Sugar of milk half an ounce  
Boiling water two pounds  
Lemon juice sufficient  
Mix. Recommended in cholera, to appease the thirst. *Ammon.*

## PAROLIC CEMENT.

R. Curd of milk, dried and powdered ten ounces  
Quicklime, powdered, one ounce  
Camphor two drachms



Mix, and keep in closely-stopped bottles. When used, a portion is to be mixed with a little water into a paste, to be applied quickly. Used to cement broken glass, china, &c. *Cooley.*

## LACCA.

### LAC

Is a resinous substance obtained from several kinds of East Indian plants, but particularly from the *Croton lacciferum*. Many varieties are known in commerce, but the principal are seed, shell, and stick lac, of which the purest is the shell. This is in thin fragments of various shades of yellow, or reddish color; brittle, inodorous, and insipid. It is somewhat astringent, and was formerly used in medicine, but is now mainly employed in the arts, for the manufacture of varnishes, sealing wax, &c.

### WHITE OR BLEACHED LAC.

R. Shell lac at will  
Boil in solution of potassa, pass chlorine gas through the solution, and malaxate in hot water. *Redwood.*

### AQUEOUS SOLUTION OF LAC.

R. Shell lac five ounces  
Borax one ounce  
Water one pint  
Boil together, and strain. Used as a varnish, or a vehicle for colors. Mixed with lampblack, it forms an ink that will resist the action of acids. *Cooley.*

### CEMENT OF SHELL LAC.

R. Shell lac at will  
Alcohol sufficient  
to make a solution of the consistence of molasses. Exceedingly tenacious. *Cooley.*

### SEALING WAX, RED.

R. Shell lac two pounds  
Venice turpentine one pound  
Vermilion one pound and a-half  
Melt the lac and turpentine together, and stir in the vermilion. *Gray.*

The other colored waxes are made in the same manner, using the appropriate pigment.

## LACTUCA ELONGATA.

### WILD LETTUCE.

The Wild Lettuce is an indigenous, herbaceous perennial, from three to six feet high, with pale yellow flowers, growing in woods and fertile soils.

*Sax. Syst.* Syngen. æqual. *Nat. Syst.* Astera-ceæ.

Willd. Sp. Pl. iii. 1525. Big. Seq. 241.

Considered to resemble the *Lactuca virosa* in

medical properties. An extract made from it may be given in doses of five to fifteen grains.

## LACTUCA.

### LETTUCE.

This well known plant is commonly cultivated for the table as a salad. It is erect and leafy, the leaves being ovate, semi-amplexicaunt, and entire or toothed. The whole abounding in a bitter, milky juice.

It is anodyne and sedative, but is seldom used in this country except in the form of the extract, or lactucarium, but is employed to fulfil many indications in Europe.

### MIXTURE OF LETTUCE AND DANDELION.

R. Fresh leaves of lettuce  
and root of dandelion  
of chicory, each, three  
ounces

Beat the whole into a pulp, adding, gradually,

Peppermint water three ounces  
Express, and filter. To be taken, during the day, in hypochondriasis, with insomnia. *Voght.*

### EXTRACT OF LETTUCE. THRIDACE.

R. Juice of the bark of lettuce at will  
Evaporate in the stove by a gentle heat, on plates. *Par. Cod.*

### SYRUP OF THRIDACE.

R. Thridace two drachms  
Water two fl. ounces

Add to the solution

Boiling syrup one pound  
Evaporate, and strain. *Par. Cod.*

### EXTRACT OF LETTUCE. LACTUCARIUM.

R. Collect the milky juice that flows from incisions made in the stalks of lettuce.

*U. S. Ph.*

### PILLS OF LACTUCARIUM.

R. Lactucarium twelve grains  
Make six pills; one every two hours, till sleep is procured. *Ellis.*

R. Lactucarium twelve grains

Conserve of elder berries

Extract of liquorice, each, sufficient

Mix, and make four pills. One, every three hours, in obstinate coughs, without expectoration. *Brera.*

### MIXTURE OF LACTUCARIUM.

R. Lactucarium thirty grains

Decoction of Iceland moss two ounces

Mucilage half an ounce

Syrup one ounce

Mix. Two spoonfuls every two hours, in spasmodic cough, insomnia, hysteria, &c.

*Brera.*



### WATER OF LETTUCE.

R. Fresh lettuce	ten parts
Water	twenty parts
Distil off ten parts.	<i>Par. Cod.</i>
Dose, half a fl. ounce to two fl. ounces.	

SYRUP OF LETTUCE.

R. Water of lettuce	one pint
Sugar	two pounds
Make syrup.	<i>Par. Cod.</i>

MIXTURE OF WATER OF LETTUCE.

R. Water of lettuce  
linden, each, two fl. ounces  
orange flowers two drachms  
Syrup of poppies one ounce  
Mix. A spoonful every two hours. *Foy.*

LAURUS.

LAUREL. BAY TREE.

The berries (*Lauri baccæ*) and leaves (*Lauri folia*) of the *Laurus nobilis*, or bay tree, are used in medicine. This laurel is a small, evergreen tree, indigenous to the countries bordering on the Mediterranean.

*Sex. Syst.* Enneand. monog. *Nat. Syst.* Lauracere.

Linn. Sp. Pl. 529. Griffith, Med. Bot. 550.

The leaves are fragrant, and have an aromatic, bitterish taste. The berries are small, oval, of a deep blue, when fresh, of a brownish-black, when dried, aromatic, and with an acrid, bitterish taste; they contain a concrete, greenish oil. Neither the leaves nor berries, nor their products, are now much used in medicine; they are stimulant and narcotic.

### OIL OF LAUREL.

R. Laurel berries at will  
Crush, warm the pulp for a few moments  
by a gentle heat, and subject to pressure.  
*Guibourt.*

Used as a friction, in swelling and pain in the joints, colic, &c.

LAUREL OINTMENT.

R. Suet	eight ounces
Melt, and add	
Laurel oil	one pound
Essence of turpen-	
tine	one ounce and a-half
	<i>Port Ph.</i>

There are a variety of formulas for this ointment, which is much used under the name of Nervine ointment, or balsam.

R. Fresh laurel leaves  
berries, each, one pound  
Lard two pounds  
Digest the bruised leaves and berries with  
the lard, till all moisture is driven off, and  
express. *Par. Cod.*

Both these are used as stimulating frictions.



## LAURO CERASUS.

## CHERRY LAUREL.

This is the officinal appellation of the leaves of the *Prunus lauro cerasus*, a small evergreen tree, a native of Asia Minor, but cultivated in the temperate parts of Europe, both for ornament and for medicinal purposes.

*Sex. Syst.* Icosand. monog. *Nat. Syst.* Drupaceae.

Linn. Sp. Pl. 678. Griffith, Med. Bot. 289.

The leaves are ovate-oblong, smooth, and shining, with scarcely any smell when entire, but when bruised, giving out a strong bitter almond odor; they have an astringent, bitter taste, with a flavor like the peach kernel. They become scentless by desiccation. They possess properties analogous to those of hydrocyanic acid, and some of the preparations from them are used for similar purposes. From the plant not being cultivated in this country, these preparations are not in use.

## CHERRY LAUREL WATER.

R. Fresh leaves of cherry laurel one pound  
Water two pints and a-half  
Compound spirits of lavender one ounce  
Bruise the leaves, mix them with the water, and distil off one pint; agitate this well; if any milkiness remain, filter, then add the spirit of lavender. *Ed. Ph.*

A sedative narcotic, similar to diluted hydrocyanic acid, but of uncertain strength. Dose, thirty minims to a fl. drachm.

## INFUSION OF CHERRY LAUREL.

R. Fresh leaves of cherry laurel four ounces  
Boiling water two pints  
Infuse, strain, and add  
Clarified honey four ounces  
Used as a lotion in cancer of the lips, and in malignant ulcers. *Cadet.*

## OINTMENT OF CHERRY LAUREL.

R. Essential oil of cherry laurel one drachm  
Lard one ounce  
Mix. As an application to alleviate pain of cancers. *Soubeyran.*

## CHERRY LAUREL WATER LOTION.

R. Distilled cherry laurel water four ounces  
Sulphuric ether one ounce  
Extract of belladonna two drachms  
Mix. Said to be very beneficial as a lotion in cases of neuralgia. *Roux.*

## MIXTURE OF CHERRY LAUREL WATER.

R. Cherry laurel water sixty drops  
Peppermint water two fl. ounces  
Infusion of quassia four fl. ounces

Mix. In dysentery, &c., in doses of a tablespoonful, three times a-day. *Augustin.*

## CERATE OF CHERRY LAUREL.

R. Cherry laurel water twelve parts  
Oil of almonds sixteen parts  
White wax four parts  
Melt the oil and wax together, and mix the cherry laurel water, stirring well. As an application to burns. *Roux.*

## LAVANDULA.

## LAVENDER.

Several species of Lavender are used in medicine, but the only one recognized by the U. S. Ph. is the *Lavandula vera*. This is a small shrub, a native of the south of Europe, and commonly cultivated in our gardens; it has opposite, sessile, narrow leaves, and spikes of small, blue flowers.

*Sex. Syst.* Didym. gymnos. *Nat. Syst.* Lamiaceae.

Linn. Sp. Pl. 800. Griffith, Med. Bot. 501.

The part used is the flowers; these have a strong, fragrant odor, and an aromatic, warm, bitterish taste. Lavender is an aromatic stimulant, and much used in nervous debility, but is mostly employed as a perfume.

## COMPOUND POWDER OF LAVENDER.

R. Lavender  
Benzoin  
Cloves  
Sal ammoniac, each, one ounce  
Petals of red roses four ounces  
Pomegranate flowers  
Chamomile flowers  
Pot marygold flowers, each, two ounces  
Peppermint flowers six drachms  
Myrrh four drachms  
Cologne water two drachms  
Solution of ammonia eighteen drops  
Oil of cinnamon six drops  
Mix well. To perfume rooms and drawers. *Guibourt.*

## SPIRIT OF LAVENDER.

R. Fresh lavender two pounds  
Alcohol one gallon  
Water two pints  
Mix, and distil a gallon by a slow fire. *U. S. Ph.*

## LAVENDER WATER.

R. Oil of lavender one fl. ounce  
Diluted alcohol fifteen fl. ounces  
Mix. *Gray.*  
R. Alcohol five gallons  
Oil of lavender twenty ounces  
bergamot five ounces  
Essence of ambergris half an ounce  
Mix.



R. Oil of lavender	four ounces
Essence of musk	two ounces
Oil of berga-	
mot	ten fl. drachms and a-half
Oil of cloves	five drachms
roses	one drachm
origanum	half a drachm
rosemary	half a drachm
Alcohol	ten pints
Water	two pints

Mix.

These are used for perfumes.

#### COMPOUND SPIRIT OF LAVENDER.

R. Spirit of lavender	three pints
Spirit of rosemary	one pint
Bruised cinnamon	one ounce
cloves	two drachms
nutmeg	half an ounce
Rasped red saunders	three drachms

Macerate for fourteen days, express, and filter. *U. S. Ph.*

As a stimulant and carminative, in doses of thirty drops to a fl. drachm.

#### OINTMENT OF LAVENDER.

R. Oil of lavender	
nutmeg	
Butter of cocoa	equal parts

Mix. To increase the growth of hair.

#### ETHEREAL TINCTURE OF LAVENDER.

R. Oil of lavender	one part
Sulphuric ether	seven parts

Mix. Dose, five to ten drops. *Beral.*

#### COMPOUND ESSENCE OF LAVENDER.

R. Oil of lavender	forty-eight drops
cloves	thirty-two drops
orange peel	sixteen drops
bergamot	eight drops
Sweet spirit of nitre	eight drops
Oil of yellow sandal	
neroli	
roses, each,	two drops
cinnamon	one drop
Diluted alcohol	one ounce
Dissolve, and add	
Honey water	eight ounces
Essence of ambergris and	
musk	one ounce

Mix. As a perfume. *Redwood.*

### LIMON.

#### LEMON.

The Lemon is the fruit of the *Citrus limonum*, a native of Asia, but now generally cultivated in all warm climates.

*Sex. Syst.* Polydelph. icosand. *Nat. Syst.* Aurantiaceæ.

Risso, Ann. Mus. xx. 207. Griffith, Med. Bot. 168.

The parts used in medicine are, the juice and the outer rind; the latter has a fragrant odor, and an aromatic, bitter taste; the juice, as is well known, is very acid, with a peculiar, grateful, and refreshing taste.

#### ESSENCE OF LEMON.

R. Pure oil of lemon	one fl. ounce
Deodorized alcohol	eight fl. ounces
Exterior rind of	
lemon (fresh)	half an ounce

Mix. Macerate for forty-eight hours, and filter; used for flavoring mixtures, and for pastry. *W. Procter.*

#### LOZENGES OF OIL OF LEMON.

R. Oil of lemon	one drachm
Sugar	twelve ounces
Lemon water	sufficient

Boil one-third of the sugar, with the lemon water, to the consistence of honey, by a gentle heat, add the rest of the sugar and the oil, and form lozenges. *Cottureau.*

#### LOZENGES OF LEMON JUICE.

R. Lemon juice	two ounces and a-half
Sugar	sixteen ounces
Essence of lemon	one scruple

Mix, and form lozenges. *Radius.*

#### COLLUTORY OF LEMON JUICE.

R. Lemon juice	
Sugar, each,	one part
Claret wine	sixteen parts

As a mouth-wash in scurvy. *Brera.*

#### LEMONADE.

R. Lemon juice	four ounces
Fresh lemon peel	half an ounce
Sugar	four ounces
Boiling water	three pints

Mix, and let cool. *Ellis.*

R. Lemons	three
Boiling water	two pints
Sugar	two ounces

Cut the lemons in slices, pour the water on them, and add the sugar at the end of an hour, and strain. *Ratier.*

#### COMPOUND LEMONADE.

R. Lemons	two
Peel them, remove the seeds, and beat them with	
Toasted bread	two ounces
Adding, gradually,	
Infusion of malt	one pint and a-half
Strain, and add	
Syrup of mulberries	one ounce
Wine	six ounces

*Cadet.*



## LEMON PEEL WATER.

R. Pare the rind of one lemon, previously rubbed with half an ounce of loaf sugar; put both into a jar, and add a quart of boiling water; when cold decant, and add one tablespoonful of lemon juice.

*A. T. Thomson.*

## LEMON SYRUP.

R. Strained lemon juice one pint

Sugar two pounds

Mix, and form syrup. *U. S. Ph.*

## SYRUP OF LEMON PEEL.

R. Fresh lemon peel three ounces

Boiling water one pint

Infuse for three days, and add

Syrup three pounds

*Guibourt.*

## VINOUS SYRUP OF LEMON PEEL.

R. Fresh lemon peel half a pound

Wine a pint and a-half

Macerate for twenty-four hours, express, and add

Syrup one pound and a-half

Boil, strain, and add

Oil of lemon, rubbed

with sugar fifteen drops

*Sard. Ph.*

## SPIRIT OF LEMON.

R. Fresh lemon peel one part

Alcohol four parts

Water two parts

Macerate for two days, and then distil four parts. *Taddei.*

## ODORIFEROUS SPIRIT OF LEMON.

R. Oil of lemon

bergamot

lavender, each, one part

Acetic ether four parts

Mix. Used as a perfume. *Sax. Ph.*

## COLOGNE WATER.

R. Oil of bergamot

lemon

cedrat, each, three ounces

rosemary

neroli

lavender, each,

one ounce and a-half

cinnamon six drachms

Alcohol twenty-four pounds

Eau de melisse three pounds

Spirit of rosemary two pounds

Dissolve the oils in the alcohol, add the other articles, and distil four-fifths.

*Cottureau.*

R. Balm two pounds

Rosemary eight ounces

Lemon peel

six ounces

Nutmeg

Cloves

Cinnamon

Coriander, each, two ounces

Oil of bergamot one ounce

Alcohol six pounds

Water eight pounds

Mix, and distil by a gentle fire.

*Niemann.*

R. Oil of neroli

lemon

bergamot

orange

rosemary, each, twelve drops

Cardamom one drachm

Alcohol one pint

Mix, and distil two-thirds. Said by Troms-

*Niemann.*

R. Oil of bergamot

lemon

cedrat

neroli, each, four drachms

cinnamon

rosemary, each, one drachm

Alcohol three pounds

Spirit of rosemary three ounces

Digest for a few days, and distil almost to dryness, add to the product

Balm water nine ounces

Spirit of jasmine

orris, each, four drachms

*Giordano.*

R. Oil of bergamot

three ounces

lemon

two ounces

lavender three drachms and a-half

neroli two drachms and a-half

origanum two drachms

rosemary one drachm

Essence of vanilla two drachms

Musk ten grains

Alcohol thirteen pints

Rose water two pints

Orange-flower water one pint

Mix, and after fourteen days, filter. *Gray.*

R. Essence of bergamot

two ounces

rosemary

mint, each, one drachm

lemon two drachms

neroli ten grains

Balm water two ounces and a-half

Mix. *Ferrera Ph.*

R. Oil of bergamot

lemons, each,

one drachm

neroli

half a drachm

cedrat

twenty drops



Honey water one ounce  
 Alcohol one pint  
 Mix. *Gray.*  
 R. Oil of neroli  
     orange  
     lemon  
     marjoram  
     rosemary,  
         each, one drachm and a-half  
     cinnamon four drops  
 Civet three grains  
 Ambergris two grains  
 Alcohol four pounds and a-half  
 Macerate for a month, and filter.

*Swedisch.*

R. Oil of bergamot three drachms  
     lemon two drachms  
     lavender twenty-five drops  
     neroli fifteen drops  
     origanum ten drops  
 Alcohol two pints  
 Mix. *Gray.*

## UNPARALLELED WATER.

R. Oil of lemon half an ounce  
     bergamot two drachms and a-half  
     cedrat two drachms  
 Alcohol seventy-two ounces  
 Hungary water eight ounces  
 Mix, and distil all but nine ounces.  
*Guibourt.*

## LINUM.

## FLAXSEED.

Flaxseed is the product of *Linum usitatissimum*, or common flax, now cultivated in most parts of the world, but whose native country is unknown.

*Sex. Syst.* Pentand. pentag. *Nat. Syst.* Linaceæ.  
*Linn. Sp. Pl.* 397. *Griffith, Med. Bot.* 206.

Both the seeds and the oil extracted from them are officinal. The first are small, oval, shining, and of a brown color. They are devoid of smell, and have a mucilaginous taste when unbroken, but an oily one when chewed. When steeped in water, they afford a viscid, inodorous and almost tasteless mucilage. The oil, which is obtained by expression, is laxative and emollient, but is principally used in the arts.

## COMPOUND MEAL OF FLAXSEED.

R. Flaxseed meal  
     Barley meal  
     Rye meal equal parts  
 Mix. To make cataplasm. *Cottureau.*

## EMOLLIENT CATAPLASM.

R. Compound meal of flaxseed four ounces  
 Water sufficient.  
 Boil to proper consistence. *Par. Cod.*

## CATAPLASM OF FLAXSEED.

R. Boiling water one pint  
     Flaxseed meal sufficient  
 Mix, and make cataplasm. *Lond. Ph.*

## INFUSION OF FLAXSEED.

R. Flaxseed half an ounce  
     Liquorice, bruised, two drachms  
     Boiling water one pint  
 Macerate for four hours, in a covered vessel, and strain. *U. S. Ph.*

A pleasant demulcent drink, in inflammatory affections of the lungs, bladder, &c.

## FLAXSEED MIXTURE.

R. Flaxseed one drachm  
     Boiling water six ounces  
 Infuse, strain, and add  
     Syrup of poppies half an ounce  
     Orange-flower water two drachms  
 Mix. *St Marie.*

## CLYSTER OF FLAXSEED OIL.

R. Flaxseed oil two ounces  
     infusion eight ounces  
 Mix. *Swedisch.*

## FLAXSEED OIL LINIMENT.

R. Flaxseed oil twelve fl. ounces  
     Lime water seven fl. ounces  
 Mix. As an application to burns and scalds. *Ainslie.*

## LIRIODENDRON.

## TULIP TREE BARK.

The tulip tree, or *Liriodendron tulipifera*, is one of the most magnificent of our native trees. It is found in most parts of the United States, and is popularly known as the poplar.

*Sex. Syst.* Polyand. polyg. *Nat. Syst.* Magnoliaceæ.

*Linn. Sp. Pl.* 755. *Griffith, Med. Bot.* 98.

The officinal portion is the bark; this, when deprived of its epidermis, is of a yellowish white color, having a faint odor, but a bitter, pungent, aromatic taste. It is a stimulating tonic, and has been used as a febrifuge, as well as in dyspepsia, chronic rheumatism, &c. The dose in powder is from half a drachm to two drachms.

## INFUSION OF TULIP TREE BARK.

R. Tulip tree bark one ounce  
     Boiling water one pint  
 Infuse, and strain. Dose, one to two fl. ounces, as a stimulating tonic and diaphoretic. *Wood.*

## TINCTURE OF TULIP TREE BARK.

R. Tulip tree bark, bruised, four ounces  
     Diluted alcohol one pint  
 Macerate for a week, express, and filter.  
 Dose, about a fl. drachm. *Beasley.*



## LOBELIA.

## LOBELIA.

Several species of *Lobelia* are medicinal, but the only one recognized as officinal, is the *L. inflata*, a small annual plant, found in most parts of the United States, having numerous small blue flowers, on leafy racemes. The leaves are ovate, sessile, serrate, and hairy.

*Sex. Syst.* Pentand. monog. *Nat. Syst.* Lobeliaceæ.

Linn. Sp. Pl. 1006. Griffith, Med. Bot. 418.

The whole herb is officinal, but the root and seed vessels are the most active. When dried, it has a somewhat unpleasant odor, and an acrid, burning, nauseous taste, resembling that of tobacco. It is emetic, and, in small doses, expectorant and sudorific. It also possesses narcotic properties. It is principally used in asthma, it has also been used in other diseases of the respiratory organs, &c. Dose in substance, as an emetic, from five to twenty grains, as an expectorant, one to three grains.

## INFUSION OF LOBELIA.

R. *Lobelia* one ounce  
Boiling water one pint  
Infuse. A fl. ounce every half hour, till vomiting ensues. *Ellis.*

## TINCTURE OF LOBELIA.

R. *Lobelia* four ounces  
Diluted alcohol two pints  
Macerate for fourteen days, express, and filter. *U. S. Ph.*

Dose, as emetic, half fl. ounce; as antispasmodic and expectorant in asthma, one to two fl. drachms, every two or three hours.

## ETHEREAL TINCTURE OF LOBELIA.

R. *Lobelia*, in coarse powder, five ounces  
Spirit of sulphuric ether two pints  
Form tincture by displacement. *Ed. Ph.*  
Dose, the same as alcoholic tincture.

R. *Lobelia* one pound  
Alcohol four pints  
Spirit of nitric ether four pints  
sulphuric ether four ounces  
Macerate for fourteen days in a dark place, and filter. Dose, five to twenty minims. *Whitelaw.*

## MIXTURE OF LOBELIA.

R. Tincture of *lobelia* one drachm  
Decoction of mallows six ounces  
A spoonful, every two or three hours, as an expectorant. *Radiis.*

## ACETIC EXTRACT OF LOBELIA.

R. *Lobelia* seed, bruised, eight ounces  
Diluted alcohol four pints  
Acetic acid one fl. ounce  
Macerate the seed in the diluted alcohol, to which the acid has been added, for forty-eight hours; then displace until four pints

of tincture are obtained, using diluted alcohol to expel the last portion, and then evaporate to the consistence of an extract.

*W. Procter.*

## VINEGAR OF LOBELIA.

R. *Lobelia* leaves, in powder, four ounces  
Diluted acetic acid one and a-half pints  
Macerate for twelve hours, and displace until twenty-four fluid ounces are obtained—using diluted acetic acid to expel the last portion. *W. Procter.*

## SYRUP OF LOBELIA.

R. Vinegar of *lobelia* six fl. ounces  
Sugar twelve ounces (av.)  
Dissolve the sugar in the vinegar, by aid of heat, remove the scum which rises, and strain. *W. Procter.*

## LUPULINA.

## LUPULIN

Is the powder attached to the strobiles of *Humulus lupulus*, and has all the properties of hops. It is a yellowish powder, mixed with portions of the scales of the hop strobiles. It is tonic, and somewhat narcotic. It is obtained by rubbing, or threshing and sifting the strobiles, or by washing these several times in different portions of water, mixing the washings, and letting the lupulin deposit, when it is to be dried. The dose is from six to ten grains.

## POWDER OF LUPULIN.

R. Lupulin one part  
Sugar two parts  
Mix. Dose, ten to twenty grains. *Foy.*

## PILLS OF LUPULIN.

R. Lupulin at will  
Rub into mass in a warm mortar, and divide into two-grain pills. *Magendie.*

R. Lupulin two drachms  
Gum tragacanth  
Water, each, sufficient  
Make mass, and make pills of two grains.

## TINCTURE OF LUPULIN.

R. Lupulin four ounces  
Alcohol two pints  
Macerate for fourteen days, and filter. *U. S. Ph.*

Dose, one to two fl. drachms, in sweetened water.

## SYRUP OF LUPULIN.

R. Tincture of lupulin one part  
Syrup seven parts  
Mix. Dose, half an ounce to an ounce. *Foy.*

## EMULSION WITH LUPULIN.

R. Tincture of lupulin four scruples



Syrup of red poppies  
of orange flowers,  
each, one ounce  
Emulsion of almonds four ounces  
Water ten ounces  
Mix. *Beral.*

## OINTMENT OF LUPULIN.

R. Lupulin one part  
Lard three parts  
Digest, on a water-bath, for five or six  
hours, and strain. To appease the pain of  
cancerous ulcers. *Soubéiran.*

## M

## MACIS.

## MACE.

A laciniated and reticulated, smooth, thin, flexible membrane, of a saffron yellow color, which is the arillus investing the shell of the nutmeg, which, in taste and odor, it much resembles. It is an active, aromatic stimulant, but is more used for culinary purposes than in medicine. The dose is from ten grains to a scruple.

## TINCTURE OF MACE.

R. Mace one part  
Alcohol (.851) eight parts  
Macerate for eight days, express, and filter. *Beral.*

Dose, from thirty to forty drops, as a carminative and stomachic.

## CARMINATIVE DROPS.

R. Oil of mace half a drachm  
Nitric ether a drachm and a-half  
Mix. In flatulent colic, ten or twelve drops on sugar. *Radius.*

## NERVINE BALSAM.

R. Oil of mace four ounces  
Beef marrow four ounces  
Melt together, and add  
Oil of rosemary two drachms  
cloves one drachm  
Camphor one drachm  
Balsam tolu two drachms  
Dissolved in  
Rectified spirit four drachms  
As a liniment in rheumatism, &c. *Redwood.*

## MAGNESIUM.

This metal is not used, but many of its compounds are largely employed.

## MAGNESIA.

## MAGNESIA.

R. Carbonate of magnesia, any quantity.

Put in an earthen vessel, and expose it to a red heat for two hours, or till the carbonic acid is wholly expelled. *U. S. Ph.*

## HEAVY MAGNESIA.

R. Mix solutions of one hundred and twenty-three parts of crystallized sulphate of magnesia, and one hundred and forty-four parts of crystallized carbonate of soda; evaporate to dryness, and calcine till the carbonic acid is expelled. Dissolve out the sulphate of soda, wash, and dry.

*R. Phillips.*

## POWDER OF MAGNESIA AND RHUBARB.

R. Powdered rhubarb one scruple  
Magnesia ten grains  
Oil of cinnamon one drop  
Mix. As a purgative, to be given in sugar and water. *Ellis.*

## POWDER OF MAGNESIA AND SULPHUR.

R. Precipitated sulphur  
Magnesia, each, half an ounce  
Mix. A teaspoonful, three or four times a-day, as an aperient. *Ellis.*

## POWDER OF MAGNESIA AND ORANGE PEEL.

R. Magnesia four drachms  
Powdered orange peel  
fennelseed, each, one drachm  
Sugar two drachms  
Mix. Ten or twelve grains, three times a-day, as a stimulant to the digestive organs. *Foy.*

## MAGNESIA MIXTURE.

R. Magnesia one drachm  
Water of ammonia one fl. drachm  
Spirit of cinnamon three fl. drachms  
Water five and a-half fl. ounces  
Mix. Recommended by Dr. James, in the cardialgia of pregnant women. Two or three teaspoonfuls, to be taken as occasion may require. *Sims.*

R. Magnesia thirty grains



Syrup of ginger two drachms  
 Peppermint water two fl. ounces and a-half  
 Compound spirit of lavender half fl. drachm  
 Spirit of caraway half fl. ounce  
 Mix. A spoonful, every hour as an antacid. *Foy.*

## MAGNESIA AND GENTIAN.

R. Magnesia one drachm  
 Infusion of gentian six fl. ounces  
 Mix. A wineglassful, three times a-day, in cases of uric acid diathesis. *Brande.*

## MAGNESIA AND RHUBARB MIXTURE.

R. Magnesia half a-drachm  
 Powdered rhubarb two grains  
 sugar one drachm  
 Essence of peppermint six drops  
 Distilled water one fl. ounce and a-half  
 Mix. In bowel complaints of children. A teaspoonful, every two hours, till it operates. *Ellis.*

## MAGNESIÆ ACETAS.

## ACETATE OF MAGNESIA.

R. Carbonate of magnesia one hundred and twenty parts  
 Acetic acid sufficient  
 to saturate. Evaporate till the mixture weighs three hundred parts. It forms a syrupy fluid. One ounce of this solution mixed with three ounces of syrup of oranges, constitutes the weaker solution, and one ounce and a-half to three ounces of syrup, forms the stronger solution. *Renault.*

## MAGNESIÆ CARBONAS.

## CARBONATE OF MAGNESIA.

## SOLUTION OF MAGNESIA.

R. Sulphate of magnesia seven drachms  
 Bi-carbonate of soda nine drachms  
 Water twenty fl. ounces  
 Carbonic acid gas six volumes  
 Dissolve the salts in the water, and pass the gas through the mixture. The result will be two drachms of bi-carbonate of magnesia, in solution. *Phæbus.*

## DEWEES' CARMINATIVE.

R. Carbonate of magnesia half a drachm  
 Tincture of assafetida forty drops  
 opium twenty drops  
 Sugar one drachm  
 Distilled water one fl. ounce  
 Mix. In flatulent colic, diarrhœa, &c., of

children. Dose, twenty drops and upwards, according to age.

*Deweës.*

## DALBY'S CARMINATIVE.

R. Carbonate of magnesia two scruples  
 Oil of peppermint one drop  
 nutmeg two drops  
 aniseed three drops  
 Tincture of castor thirty drops  
 assafetida fifteen drops  
 Spirit of pennyroyal fifteen drops  
 Compound tincture of cardamom thirty drops  
 Peppermint water two ounces

*Paris.*

R. Carbonate of potassa half an ounce  
 magnesia twelve ounces  
 Laudanum six fl. ounces  
 Oil of peppermint  
 caraway  
 fennel, each, two fl. scruples  
 Sugar thirty-two ounces  
 Water ten pints  
 Triturate the oils with the sugar and magnesia, then add the remainder.

*Phil. Coll. Pharm.*

## CARBONATE OF MAGNESIA MIXTURE.

R. Carbonate of magnesia half a drachm  
 Sulphate of magnesia three drachms  
 Aromatic spirit of ammonia one fl. drachm  
 Tincture of rhubarb half fl. ounce  
 henbane half fl. drachm  
 Mint water four fl. ounces  
 Mix. As a carminative cathartic. A tablespoonful, two or three times a-day.

*Meigs.*

## MAGNESIA AND COLCHICUM.

R. Carbonate of magnesia one drachm  
 Sugar  
 Gum Arabic, each, sufficient  
 Wine of colchicum forty drops  
 Distilled water four fl. ounces

Mix. In gout and rheumatism. A tablespoonful, every hour, till it operates.

*Ellis.*

## MIXTURE OF MAGNESIA AND CAMPHOR.

R. Magnesia one drachm  
 Camphor half a drachm  
 Sugar  
 Gum Arabic, each, sufficient  
 Sulphuric ether half fl. drachm  
 Distilled water four fl. ounces

Mix. In flatulency and irritable stomach. A tablespoonful, four or five times a-day.

*Ellis.*



## MAGNESIÆ CITRAS.

## CITRATE OF MAGNESIA.

R. Citric acid at will  
Dissolve in water, and add  
Carbonate of magnesia sufficient  
to saturate; wash the powder, and dry by  
a gentle heat. Laxative, but not as ac-  
tive as the sulphate. *Beasley.*

## SOLUTION OF CITRATE OF MAGNESIA.

R. Carbonate of magnesia  
two hundred and ninety-two grains  
Crystallized citric acid  
four hundred and forty-six grains  
Water sixteen fl. ounces  
Lemon syrup two fl. ounces  
Dissolve one hundred and thirty-eight  
grains of the carbonate in two fl. ounces  
of water; holding in solution one hundred  
and seventy grains of citric acid, and pour  
it into a twelve ounce mineral water bot-  
tle. The remainder of the magnesia is  
then triturated with the rest of the water,  
and poured into the bottle, and one hun-  
dred and eighty-five grains of the acid  
added, and the bottle immediately corked  
and tied over. As soon as, with occasional  
agitation, the opaque fluid becomes slight-  
ly milky, remove the cork, filter the solu-  
tion, and pour it back into the bottle, with  
two ounces of lemon syrup, and ninety-  
one grains of citric acid; cork and wire.  
Each ounce contains one drachm of the ci-  
trate. Dose, from a half to a whole bot-  
tle. *Rabourdin.*

## CITRATE OF MAGNESIA WATER.

R. Subcarbonate of mag-  
nesia five drachms and a-quarter  
Citric acid five drachms and a-half  
Lemon syrup two drachms and a half  
Water sufficient  
to fill a Seidlitz water-bottle.  
Dissolve the acid in one sixth of the wa-  
ter, triturate the carbonate with the re-  
mainder, and add to it half the acid solu-  
tion; and, on the ceasing of the efferve-  
scence, pour it into the bottle with the  
syrup; add the acid solution, and cork  
instantly. As a purgative draught.  
*Bardet.*

## MAGNESIÆ SULPHAS.

## SULPHATE OF MAGNESIA.

## CHELTENHAM SALTS.

R. Sulphate of magnesia  
Muriate of soda, each, four parts  
Sulphate of soda three parts

To be well dried before being powdered  
and mixed. Dose, half an ounce, or more,  
in solution. *Cooley.*

## SEIDLITZ WATER.

R. Sulphate of magnesia two drachms  
Chloride of magnesia eighteen grains  
Soda water twenty fl. ounces  
Dissolve. As a purgative. *Foy.*

## PURGATIVE EMULSION.

R. Sulphate of magnesia two drachms  
Flake manna one ounce  
Simple emulsion four fl. ounces  
Dissolve. A tablespoonful, every two  
hours. *Wendt.*

SULPHATE OF MAGNESIA AND TARTAR  
EMETIC.

R. Sulphate of magnesia one ounce  
Tartar emetic one grain  
Flake manna one ounce  
Lemon juice half an ounce  
Water eight ounces  
Make a solution, and strain. A table-  
spoonful, every hour, till it operates.  
*Dewees.*

## SULPHATE OF MAGNESIA AND COFFEE.

R. Sulphate of magnesia one ounce  
Powdered roasted  
coffee two drachms and a-half  
Water sixteen ounces  
Boil well in an earthen vessel, for two  
minutes; remove from fire, and let infuse  
for some minutes; then filter, or strain.  
This will destroy the bitter taste of the  
salt. *Combes.*

## CLYSTER WITH SULPHATE OF MAGNESIA.

R. Sulphate of magnesia two ounces  
Olive oil one fl. ounce  
Tepid water one pint  
Mix. One-half to be injected, and if it  
does not produce the desired effect, the  
remainder to be administered. *Ellis.*

SULPHATE OF MAGNESIA AND SULPHURIC  
ACID.

R. Sulphate of magnesia sufficient  
to saturate  
Water seven fl. ounces  
Add to solution  
Diluted sulphuric acid one fl. ounce  
Dose, a tablespoonful, in a wineglassful of  
water, every hour, till it operates. *Henry.*

SULPHATE OF MAGNESIA AND ROCHELLE  
SALT.

R. Sulphate of magnesia  
Tartrate of potassa and  
soda equal parts



Dose, two to three drachms, in six ounces of water. *Christison.*

SULPHATE OF MAGNESIA AND NITRIC ACID.

R. Sulphate of magnesia half an ounce  
Tincture of jalap one fl. drachm  
Nitric acid two drops  
Mint water two fl. ounces  
Mix. To be taken for a draught. *Ellis.*

MAGNESII SULPHURETUM.

SULPHURET OF MAGNESIA.

SYRUP OF SULPHURET OF MAGNESIA.

R. Sulphuret of magnesia half an ounce  
Fennel water six ounces  
Dissolve, strain, and add  
Sugar fifteen ounces  
Said to be useful in chronic exanthemata, whooping-cough, &c. Dose, a spoonful, occasionally. *Radiis.*

MAGNESIÆ PHOSPHAS.

PHOSPHATE OF MAGNESIA.

R. Magnesia at will  
Phosphoric acid sufficient  
to saturate; filter, and evaporate to dryness. *Neimann.*

Stated to be useful in rachitis, in doses of ten to twenty grains.

MAGNOLIA.

The barks of several species of Magnolia are employed as stimulating tonics in the United States, but more especially that of *M. glauca*. It is a stimulating, bitter tonic, with some diaphoretic powers, and the decoction has been used with some success in intermittent fevers and rheumatism, as has also a tincture of the cones. Dose, in powder, half a drachm to a drachm.

TINCTURE OF MAGNOLIA.

R. Recently dried bark or  
cones of magnolia four ounces  
Diluted alcohol one pint  
Macerate for a week, express, and filter.  
In chronic rheumatism. *Beasley.*

MALTUM.

MALT.

Malt is barley germinated by warmth and moisture, and then subjected to such a degree of heat as to destroy the vital principle. It contains sugar, gum, and hordein, and is principally used in the manufacture of malt liquors.

INFUSION OF MALT.

R. Ground malt one pint  
Scalding water three pints  
Infuse for two hours, strain, and add sugar or lemon juice, if required. *Ellis.*

Much prescribed by the late Dr. Parrish, as a drink in inflammatory fevers.

EMOLLIENT CLYSTER.

R. Ground malt  
Mallow root  
Pearl barley, each, two drachms  
Water one quart  
Boil down to a pint, and strain. *Phæbus.*

MALVA.

COMMON MALLOW.

Several species of Mallow are recognized as officinal in the European Pharmacopœias, and, although they are very similar in their properties, that most generally employed is the *M. sylvestris*, the leaves and flowers being the officinal parts. They are emollient and demulcent, but are seldom or never employed in this country.

MANGANESIUM.

MANGANESE.

This metal is never employed in medicine, but several of its oxides and salts have obtained some celebrity.

MANGANESII CARBONAS.

CARBONATE OF MANGANESE.

R. Oxide of manganese at will  
Wash in very dilute muriatic acid, dissolve in strong muriatic acid, and evaporate to dryness. Dissolve part in water, and precipitate with carbonate of soda; wash precipitate, and digest it with a solution of the rest of the salt. Filter, and precipitate by carbonate of soda; wash, and dry. *Beasley.*

MANGANESII MURIAS.

MURIATE OF MANGANESE.

R. Muriatic acid at will  
Carbonate of manganese sufficient  
to saturate; evaporate, and crystallize. Preserve in closely-stopped bottles. A solution in water has been praised as a gargle in aphthous sore throat. *Jourdain.*

PILLS OF MURIATE OF MANGANESE.

R. Muriate of manganese two scruples  
Gum Arabic  
Extract of liquorice, each, one scruple



Mix, and form pills of two grains. Advised in obstinate cutaneous affections, in doses of from five to twenty. *Augustin.*

## MANGANESII OXIDUM.

## OXIDE OF MANGANESE.

## POWDER OF OXIDE OF MANGANESE.

R. Oxide of manganese two grains  
Magnesia one scruple  
Mix. To be taken every three hours, in chlorosis. *Brera.*

## BOLUS OF OXIDE OF MANGANESE.

R. Oxide of manganese fourteen grains  
Extract of savine  
Aloes, each, six grains  
Mix, and divide into six boluses; to be taken during the day, as an emmenagogue. *Niemann.*

## OINTMENT OF OXIDE OF MANGANESE.

R. Oxide of manganese one part  
Lard two parts  
*Giordano.*

## OINTMENT OF OXIDE OF MANGANESE AND SULPHUR.

R. Oxide of manganese  
Sulphur, each, one ounce  
Soap one ounce  
Lard three drachms  
Mix. *Beasley.*

Both these ointments have been recommended in porrigo, and other skin diseases.

## MANGANESII SULPHAS.

## SULPHATE OF MANGANESE.

R. Carbonate of manganese at will  
Diluted sulphuric acid sufficient to saturate; evaporate, and crystallize. Dose, one to two drachms, in half a pint of water; to be taken before breakfast, as a cholagogue cathartic. *Beasley.*

## MANNA.

## MANNA.

This is the concrete juice of several species of *Frazinus*, but principally from the *F. ornus*, or *Ornus Europæa*; a small tree found in the south of Europe.

*Sex. Syst. Diand. monog. Nat. Syst. Oleraceæ. Linn. Sp. Pl. 1510. Griffith, Med. Bot. 445.*

There are several varieties of manna—*flake*, common or in sorts, and *fat*. The first is the best and purest. It consists of sugar, a peculiar principle called mannite, and a viscous, nauseous matter. Manna is a gentle laxative, but some-

times causes flatulence and pain. Dose, for an adult, one to two ounces; for children, one to four drachms.

## LOZENGES OF MANNA.

R. Mallow root  
Water, each, four ounces  
Boil a short time, and add  
Manna twelve ounces  
When dissolved, strain, and add  
Sugar six pounds  
Opium, dissolved in water, twelve grains  
Evaporate to the consistence of an electuary, and add  
Orange-flower water three ounces  
Essence of bergamot citron, each, four drops  
Evaporate to proper consistence, pour on a marble slab, and divide into lozenges. *Guibourt.*

## SYRUP OF MANNA.

R. Manna six ounces  
Water twelve fl. ounces  
Dissolve, and strain, then add  
Sugar one pound  
Evaporate to syrup. *Pidereau.*

## EMULSION OF MANNA.

R. Sweet almonds half an ounce  
Manna two ounces  
Syrup of peach flowers one fl. ounce  
Infusion of liquorice four fl. ounces  
Orange-flower water half fl. ounce  
Make an emulsion. *Radius.*

R. Manna  
Oil of almonds, each, one ounce  
Carbonate of potassa twelve grains  
Cinnamon water  
Water, each, three fl. ounces  
Rub the first three ingredients together, and gradually add the waters. Two fl. ounces to be taken daily, in divided doses, in inflammation of the kidneys. *Babington.*

## MANNA MIXTURE.

R. Sulphate of magnesia one ounce  
Manna six drachms  
Tincture of senna one fl. drachm and a-half  
Boiling water five fl. ounces  
Mix. As a purge. *Ainslie.*  
R. Manna one ounce  
Sulphate of soda one ounce and a-half  
Water six fl. ounces  
*Augustin.*  
R. Manna four ounces  
Senna one ounce  
Cremor tartar half an ounce



Anise half an ounce  
Boiling water four pints  
Digest for some time, and strain.  
*Span. Pharm.*

## MANNITE.

## MANNITE.

R. Common manna six pounds  
Distilled water three pints  
White of egg one  
Mix, and boil for a few minutes; strain when cold; press impure mannite in a cloth; mix it with its weight of water, and again press. Dissolve in boiling water, with animal charcoal, filter, evaporate to a pellicle, and set aside to crystallize.

*Ruspini.*

Dose, from two drachms to two ounces.

## MARANTA.

## ARROWROOT.

Arrowroot is the fecula of the roots, or rhizomes of many plants, but that in common use is derived from the *Maranta arundinacea*, a native of the West Indies.

*Sex. Syst.* Monand. monog. *Nat. Syst.* Marantaceæ.

*Linn. Sp. Pl.* 2. *Griffith, Med. Bot.* 637.

West Indian arrowroot is a light white powder, tasteless and inodorous. It has firm feel, and crackles when pressed. It is nutritious and demulcent, and is an article of diet well suited to the sick and convalescent, especially in bowel complaints. It is prepared for use by adding to it a certain proportion of boiling water, usually about a pint to a tablespoonful of the fecula.

## WATER ARROWROOT.

R. Rub the arrowroot with a little cold water, till well mixed; then pour boiling water over it, stirring constantly; afterwards boil for five minutes. Sweeten with white sugar, and flavor with lemon juice, or some aromatic; if wine be added, none of the astringent kinds should be used.

*Thomson.*

## MILK ARROWROOT.

R. Arrowroot a tablespoonful  
Sweet milk  
Boiling water, each, half a pint  
Proceed as with water arrowroot. *Ellis.*  
R. Milk fifteen fl. ounces  
Sugar two ounces  
Arrowroot one ounce  
Proceed as above. *Beral.*

## ARROWROOT PUDDING.

Rub a tablespoonful of arrowroot with a

little cold water, and add to it, stirring constantly, a pint of boiling milk; with this mix one egg and three teaspoonfuls of sugar, previously beaten together. This may be boiled or baked, and forms a good diet in convalescence. *Thomson.*

## BEEF TEA ARROWROOT.

This is made in the same manner as the water arrowroot, except that beef tea is used in the boiling state instead of water, and that the mixture is to be boiled twenty minutes instead of five. *Thomson.*

## MARRUBIUM.

## HOREHOUND.

Horehound, *M. vulgare*, is a small, herbaceous plant, a native of Europe, and naturalized in some parts of this country.

*Sex. Syst.* Didynam. gymnos. *Nat. Syst.* Lamiales.

*Linn. Sp. Pl.* 816. *Griffith, Med. Bot.* 512.

The whole herb is officinal. It has a strong and peculiar odor in the fresh state, which is almost lost on drying. Its taste is bitter, and somewhat acrid. It is tonic, emmenagogue, antispasmodic, and, in large doses, laxative. The dose of the powder is from thirty grains to a drachm.

## EXTRACT OF HOREHOUND.

R. Powdered horehound sufficient  
Exhaust by the process of displacement, and evaporate to proper consistence. The dose is from a scruple to half a drachm.

## COMPOUND DECOCTION OF HOREHOUND.

R. Dried horehound one ounce  
Bruised liquorice root  
Flaxseed, each, half an ounce  
Boiling water one pint and a-half  
Macerate for four hours, and strain. Dose, from a fl. ounce to two fl. ounces, in catarrh. *Steph. and Church.*

## PECTORAL MIXTURE.

R. Extract of horehound  
cough grass, each, two drachms  
Decoction of dandelion ten fl. ounces  
Oxymel of squills  
Syrup of fennel, each, two fl. ounces  
Mix. A tablespoonful occasionally. *Augustin.*

## SYRUP OF HOREHOUND.

R. Dried horehound one ounce  
Horehound water two pints  
Digest for two hours, express, and add  
Sugar four pounds  
Form a syrup. *Cottèreau.*



## HOREHOUND CANDY.

R. Expressed juice of horehound one pint  
 Sugar ten pounds  
 Boil to proper consistence, pour into  
 moulds, or on marble slab, and divide.

*Cooley.*

Much used to allay irritation of throat, in catarrh.

## MASTICHE.

## MASTIC.

Mastic is a concrete, resinous exudation from the *Pistacia lentiscus*; a small tree, a native of most of the countries bordering on the Mediterranean.

*Sex. Syst.* Diœc. pentand. *Nat. Syst.* Anacardiaceæ.

Linn. Sp. Pl. 1454. Griffith, Med. Bot. 186.

Mastic occurs in yellowish, semi-transparent, brittle grains or tears, of mild, agreeable smell, and a resinous but not unpleasant taste. It is much used by the Turks as a masticatory to sweeten the breath and to strengthen the gums. It is more employed in the arts as the basis of varnishes, than in medicine.

## MASTIC DENTIFRICE.

R. Powdered mastic  
 Prepared chalk, each, two drachms  
 Powdered orris root half an ounce  
 Oil of roses three drops  
 Mix. *Phæbus.*

## AROMATIC MASTIC COLLUTORY.

R. Mastic two drachms  
 Balsam of Peru half a drachm  
 Gum Arabic two drachms  
 Orange-flower water six fl. ounces  
 Make an emulsion, and add  
 Tincture of myrrh two fl. drachms  
*Phæbus.*

## ANTI-ODONTALGIC PASTE.

R. Mastic  
 Sandarach, each, two drachms  
 Kino ten grains  
 Opium two grains  
 Oil of rosemary eight drops  
 Spirit of horseradish sufficient  
 Mix, and form a paste. Used to fill decayed teeth. *Gassicourt.*

## ETHEREAL TINCTURE OF MASTIC.

R. Mastic four parts  
 Sulphuric ether one part  
 Dissolve. To fill the cavity of carious teeth, for which purpose a small dossil of cotton is saturated with it, and introduced into the cavity. *Soubeiran.*

## PICTURE VARNISH.

R. Mastic three hundred and sixty parts  
 Venice turpentine forty-five parts

Camphor fifteen parts  
 Spirit of turpentine one thousand parts  
 Dissolve with heat. *Redwood.*

## CRYSTAL VARNISH.

R. Mastic three ounces  
 Alcohol one pint  
 Dissolve. Used to fix pencil drawings. *Cooley.*

## MATICO.

## MATICO.

This is the leaves of a Peruvian plant, or plants; in most cases of the *Piper angustifolium* (*Artanthe elongata*).

*Sex. Syst.* Diand. trigyn. *Nat. Syst.* Piperaceæ.

The leaves, as found in commerce, are strongly veined, reticulated; of a greenish ash color; mixed with fragments of the stem and flower spikes; of a somewhat aromatic odor, and a warm, spicy taste. They have attained much celebrity as a hemostatic, and have also been given in mucous and other discharges. As a styptic, they are applied in substance to the bleeding part.

## INFUSION OF MATICO.

R. Matico one ounce  
 Boiling water one pint  
 Infuse for two hours. Dose, two table-  
 spoonfuls, two or three times a-day, or of-  
 tener. In hemorrhagic and other dis-  
 charges. *Jeffreys.*

## INFUSION OF MATICO AND SENNA.

R. Matico  
 Senna, each, two drachms  
 Boiling water one pint  
 Infuse, and strain. Dose, one fl. ounce  
 and a-half, repeatedly. *Watmough.*

## DECOCTION OF MATICO.

R. Matico one ounce  
 Water one pint  
 Boil for ten or fifteen minutes, and strain.  
 Given as above. *Jeffreys.*

## TINCTURE OF MATICO.

R. Matico two and a-half ounces  
 Diluted alcohol one pint  
 Macerate for fourteen days, and strain. *Lane.*

Dr. Jeffreys directs three ounces of matico. Dose, from thirty to sixty drops, in water. In hemorrhagic discharges, &c.

## SYRUP OF MATICO.

R. Matico four ounces  
 Diluted alcohol one pint  
 Make tincture by displacement, evaporate  
 to one half; add  
 Sugar sufficient  
 and form syrup. *Ruschenberger.*



Used in same cases as above. Dose, one to two fl. drachms.

#### OINTMENT OF MATICO.

R. Powdered matico three drachms  
 opium three grains  
 Lard one ounce  
 Mix. As an application to hemorrhoids.  
 Young.

#### MATRICARIA.

##### GERMAN CHAMOMILE.

This is the flowers of *M. chamomilla*, a small, annual, herbaceous plant, native of many parts of Europe.

*Sex. Syst.* Syngen. super. *Nat. Syst.* Astera-ceæ.

Linn. Sp. Pl. 1256. Stokes, Bot. Mat. Med. iv. 238.

The flowers, which are the officinal part, are smaller than those of the true chamomile. They have a strong, penetrating, unpleasant odor, which is much diminished by drying. Their taste is bitter and somewhat nauseous. The medical properties are much the same as those of chamomile, and it is given in the same manner.

#### MEL.

##### HONEY.

This is a peculiar fluid, prepared from flowers by the *Apis mellifica*, or honey-bee.

In a recent state it is fluid, but by age concretes into a soft, granular mass. It has a peculiar, somewhat aromatic odor, and a sweet taste, followed by faint acidity. It resembles sugar in its properties, but is more laxative. It is principally used in medicine as a vehicle.

##### CLARIFIED HONEY.

R. Honey at will  
 Melt by means of a water-bath, and remove the scum.  
*U. S. Pharm.*

##### PREPARED HONEY.

R. Clarified honey half a pint  
 Diluted alcohol one pint  
 Prepared chalk half an ounce  
 Mix the honey and alcohol, add the chalk, let the mixture stand for two hours, occasionally stirring. Heat to boiling, filter, and evaporate on water-bath, so that when cold the sp. gr. will be 1.32.  
*U. S. Pharm.*

##### OXYMEL.

R. Honey two pounds  
 Distilled vinegar one pint  
 Boil to consistence of syrup, removing scum.  
*Dub. Pharm.*

##### HYDROMEL.

R. Honey two parts

Water thirty-two parts  
 Dissolve, and strain. A refreshing and slightly laxative drink.  
*Foy.*

##### PECTORAL HYDROMEL.

R. Washed Iceland moss two ounces  
 Hyssop leaves half an ounce  
 Hydromel two pounds  
 Macerate for five days, strain, and add  
 Sugar three ounces  
 In catarrh, &c., one or two spoonfuls occasionally.  
*Foy.*

##### EXPECTORANT MIXTURE.

R. Honey  
 Fresh butter, each, two ounces  
 Mix, and melt over a gentle fire. A spoonful occasionally.  
*St. Marie.*

##### CEROMEL.

R. Yellow wax one ounce  
 Honey four ounces  
 Melt the wax, and add the honey, stirring well. As an application to indolent ulcers.  
*Van Mons.*

##### HONEY WATER.

R. Honey two parts  
 Washed sand three parts  
 Introduce into a retort, and distil on a sand-bath, and remove the oil that floats on the product. Aperient and diuretic; in doses of a scruple to half a drachm.  
*Spielmann.*

R. Rectified spirit eight pints  
 Rose water two pints  
 Orange-flower water two pints  
 Oil of cloves half an ounce  
 lavender half an ounce  
 bergamot two drachms  
 sandal wood one drachm  
 Honey one ounce  
 Tincture of saffron one ounce  
 Musk ten grains  
 Macerate for a week, and filter. Used as a perfume.  
*Gray.*

R. Honey  
 Coriander, each, eight ounces  
 Fresh lemon peel one ounce  
 Cloves six drachms  
 Nutmeg  
 Benzoin  
 Storax, each, half an ounce  
 Vanilla three drachms  
 Rose water  
 Orange-flower water, each, five ounces  
 Alcohol forty-eight ounces  
 Distil by means of a water-bath. *Guibourt.*



## MELISSA.

## BALM.

Several species have been used in medicine, but the only one that is officinal in this country is *M. officinalis*, a small herbaceous plant, which is a native of the South of Europe, and is very generally cultivated in our gardens.

*Sex. Syst.* Didynam. gymnos. *Nat. Syst.* Lamiales.

*Linn. Sp. Pl.* 827. *Stokes, Bot. Mat. Med.* iii. 365.

The whole herb is used; it has an agreeable odor, somewhat like that of lemons, and an aromatic, slightly bitter taste. Its medicinal properties are very slight, but it communicates an agreeable flavor to infusions.

## COMPOUND SPIRIT OF BALM.

R. Fresh balm one pound and a-half  
Lemon peel four ounces  
Cinnamon  
Cloves  
Nutmegs, each, two ounces  
Angelica, dried,  
Coriander, each, one ounce  
Alcohol, .869, eight pounds  
Macerate for two days, and distil.

Used as a perfume.

R. Fresh mint thirteen ounces  
Sage three ounces and a-half  
Angelica eighteen drachms  
Hyssop  
Marjoram, each, fourteen drachms  
Thyme thirteen drachms  
Cinnamon  
Coriander, each, twelve drachms  
Rosemary  
Cloves  
Nutmeg, each, ten drachms  
Anise half an ounce  
Lemon peel one drachm  
Alcohol, (.927)  
one hundred and forty ounces

Macerate for two days, distil, and rectify.  
*Guibourt.*

This form of the compound spirit of balm is known as the *Eau des Carnus*. It is used as a perfume, stomachic, and stimulant.

R. Compound spirit of balm sixteen parts  
Spirit of mint  
rosemary, each, twelve parts  
sage nine parts  
thyme eight parts

Mix. This is known as *Eau de Dardel*, and is used as above. *Guibourt.*

## ANTI-HYSTERIC WATER.

R. Fresh balm one pound  
Laurel berries  
Cinnamon, each, one ounce  
Myrrh half an ounce

Castor two drachms  
White wine twelve pounds

Digest for some time, and distil off one-half. As a stimulant, and antispasmodic in hysteria. *Wirttemberg Ph.*

## MENTHA PIPERITA.

## PEPPERMINT.

Many species of *Mentha* are used in medicine, but two only are recognized in our pharmacopœia, the *M. piperita* and *M. viridis*, both natives of Europe, and naturalized in this country.

*Sex. Syst.* Didynam. gymnos. *Nat. Syst.* Lamiales.

*Smith, Eng. Bot.* 687. *Griffith, Med. Bot.* 502.

The whole plant is used; it has a peculiar, aromatic odor, and a balsamic, pungent, camphorated taste, followed by a sensation of coolness. It is aromatic, carminative, and stimulant, and is much employed to expel flatus, obviate nausea, &c.

## OIL OF PEPPERMINT.

R. Peppermint at will  
Water sufficient  
Distil, and collect the oil that floats on the product. *Par. Cod.*

## ESSENCE OF PEPPERMINT.

R. Oil of peppermint two fl. ounces  
Alcohol one pint  
Mix. *U. S. Ph.*

## PEPPERMINT LOZENGES.

R. Oil of peppermint one drachm  
Water of peppermint sufficient  
Sugar twelve ounces  
Mix, and form pastilles. *Cottureau.*

## CARMINATIVE MIXTURE.

R. Essence of peppermint one fl. ounce  
Peppermint water four fl. ounces  
Syrup of mallows three fl. ounces  
Oil of cinnamon two drops  
peppermint three drops  
Mix. A spoonful every two hours, in convulsive hiccup. *Alibert.*

R. Magnesia half a drachm  
Peppermint  
water two and a-half fl. drachms  
Compound spirit of  
lavender half fl. drachm  
Spirit of caraway four fl. drachms  
Syrup of ginger two fl. drachms  
Mix. A spoonful occasionally, as an antacid and carminative. *Paris.*

## MENTHA VIRIDIS.

## SPEARMINT.

This species, like the last, although a native of



Europe, has become extensively naturalized in many parts of the United States, principally in low, wet situations. It is less powerfully aromatic, and pungent than peppermint, but is more agreeable in odor and taste to most persons. It has the same properties, and its preparations are the same as of that article.

#### INFUSION OF MINT.

R. Dried spearmint two drachms  
Boiling water sufficient  
to afford six fl. ounces of strained infusion.  
*Dub. Ph.*

#### COMPOUND INFUSION OF MINT.

R. Dried spearmint two drachms  
Boiling water sufficient  
to afford six fl. ounces of strained infusion.

Add  
Sugar two drachms  
Oil of spearmint three drops

Dissolved in  
Compound tincture of  
cardamom half fl. ounce  
To allay nausea, &c. Dose, one to two fl. ounces.  
*Dub. Ph.*

#### ARQUEBUSADE WATER.

R. Dried mint  
angelica tops, each, one pound  
wormwood half a pound  
Angelica seeds five ounces  
Oil of juniper half a drachm  
Spirit of rose-  
mary three pints and a-half  
Rectified spirit five gallons  
Water four gallons  
Mix, and distil six gallons. Much celebrated as a vulnerary, for contusions, and for cleansing and healing ulcers and wounds, especially those caused by fire-arms.  
*Redwood.*

### MENYANTHES.

#### BUCKBEAN.

The *M. trifoliata*, or buckbean, is an aquatic, herbaceous plant, with ternate leaves, a native of both Europe and North America.

*Sex. Syst.* Pentand. monog. *Nat. Syst.* Gentianaceæ.

*Linn. Sp. Pl.* 207. *Griffith, Med. Bot.* 464.

All parts of the plant are medicinal. They are very bitter, but have very little odor. In small doses, the buckbean is tonic and astringent, in large ones cathartic, and even emetic. As a tonic, the dose of the powdered leaves, or root, is from twenty to thirty grains.

#### INFUSION OF BUCKBEAN.

R. Buckbean one ounce  
Boiling water one pint  
Macerate for two hours, and strain. Dose, one to two fl. ounces.  
*Saunders.*

#### EXTRACT OF BUCKBEAN.

R. Clarified juice of buckbean at will  
Evaporate to the proper consistence.  
*Guibourt.*

Dose, ten to fifteen grains.

#### COMPOUND PILLS OF BUCKBEAN.

R. Extract of buckbean  
valerian, each, two drachms  
Soap half an ounce  
Rhubarb one drachm and a-half  
Syrup sufficient  
Mix, and make pills of two grains. Ten, three times a-day, in a costive condition of the bowels.  
*Vogel.*

#### MIXTURE OF BUCKBEAN, FUMITORY, &c.

R. Extract of buckbean  
fumitory  
couch grass, each,  
half an ounce  
Chamomile water ten fl. ounces  
Compound infusion of  
horseradish four fl. ounces  
Sulphuric acid half a drachm  
Mix. Two tablespoonfuls a-day, as an antiscorbutic.  
*Selle.*

#### DIURETIC MIXTURE.

R. Extract of buckbean half an ounce  
Vinegar of squills two fl. ounces  
Tincture of assafetida half fl. ounce  
Mix. Dose, thirty drops, three times a-day, as a diuretic.  
*Augustin.*

### MEZEREUM.

#### MEZEREON.

This is the bark of several species of *Daphne*; two species of which are officinal in the U. S. Pharm., the *D. mezereum* and *D. gnidium*, both shrubs indigenous to Europe. The first being the most generally used.

*Sex. Syst.* Octand. monog. *Nat. Syst.* Thymelacææ.

*Linn. Sp. Pl.* 509. *Griffith, Med. Bot.* 560.

The officinal portion is the bark. This, as found in the shops, is in long, narrow strips, of a grayish color and fibrous texture. It is almost inodorous, with a sweetish taste at first, soon becoming extremely acrid. It is used as an external irritant and stimulant, and internally as an alterative diaphoretic.

#### DECOCTION OF MEZEREON.

R. Mezereon two drachms  
Water three pints  
Boil till reduced to a quart. Add  
Liquorice root half an ounce  
Strain. Four to eight fl. ounces a-day.  
*Thomson.*

In syphilitic affections, especially when there are night-pains and eruptions.



## COMPOUND DECOCTION OF MEZEREON.

R. Mezereon	two drachms
Bittersweet	half an ounce
Burdock	two ounces
Water	four pints

Boil down to three pints, and add

Liquorice root	two drachms
----------------	-------------

Strain. In the same cases as above, and in obstinate diseases of the skin.

*Van Mons.*

## EXTRACT OF MEZEREON.

R. Mezereon	three parts
Alcohol	five parts

Digest, and repeat the operation twice, with the same proportion of alcohol, each time; unite the tinctures; filter, and distil off three-fourths of the spirit; filter the residue, retaining what is left on the filter.

*Ham. Pharm.*

## GREEN OIL OF MEZEREON.

R. Extract of mezereon	at will
Sulphuric ether	sufficient

Macerate, express, and permit the ether to evaporate.

*Guibourt.*

## MEZEREON OINTMENT.

R. Extract of mezereon	one part
Alcohol	
White wax, each,	eight parts
Lard	seventy-two parts

Dissolve the extract in the alcohol; add the wax and lard; heat to expel the alcohol, and strain.

*Guibourt.*

R. Green oil of mezereon	one part
White wax	sixteen parts
Lard	one hundred and sixty parts

Mix, and melt by means of a water-bath.

Used as a stimulating application to blistered surfaces, and to indolent ulcers.

## MONARDA.

## HORSEMINT.

Almost all the species of *Monarda* are possessed of medicinal properties, but one only is recognized as officinal; the *M. punctata*. This is a native, perennial, herbaceous plant, with yellow flowers, spotted with brown; usually growing in sandy soil.

*Sex. Syst.* Diand. monog. *Nut. Syst.* Lamiaceae.

*Linn. Sp. Pl.* 126. *Griffith, Med. Bot.* 510.

The whole plant is aromatic, and abounds in a pungent, volatile oil. It is used in infusion, for flatulent colic, and as an emmenagogue.

## OIL OF HORSEMINT.

R. Fresh horsemint	at will
Water	sufficient

Distil, and collect the oil that floats on the product.

Internally, as a carminative; in doses of two or three drops, with sugar and water. Externally, as a rubefacient, in low states of fever, rheumatism, &c. In most cases it must be diluted.

## OIL OF HORSEMINT LINIMENT.

R. Oil of horsemint	half an ounce
Tincture of camphor	two ounces
Laudanum	two drachms

Mix. As a rubefacient application.

*Atlee.*

## MONESIA.

## MONESIA.

This is the extract of an unknown tree, perhaps a *Chrysophyllum*, coming from South America. It is in the form of hard, thick cakes, having at first a sweet taste, which soon becomes astringent and acrid. It is of a dark brown color, friable, and soluble in water. It has been used with success in various discharges, especially of a chronic character; in chronic bronchitis, &c., in doses of two to ten grains, frequently repeated; also as an application to atonic ulcers.

## PURIFIED MONESIA.

R. Monesia, bruised,	one pound
Boiling water	six pints

Infuse for twenty-four hours, stirring occasionally; decant, and evaporate by a water-bath.

*Beasley.*

## MONESIA MIXTURE.

R. Monesia	two scruples
Water	seven and a-half fl. ounces
Compound tincture of cardamoms	half a fl. ounce

Mix, and dissolve. Dose, a tablespoonful.

*Neligan.*

## SYRUP OF MONESIA.

R. Monesia	one drachm
Water	one fl. drachm
Boiling syrup	twelve fl. ounces

Mix. Dose, half a fl. ounce.

*Derosne.*

## COMPOUND SYRUP OF MONESIA.

R. Hot syrup of monesia	sixteen fl. ounces
Orange-flower water	half a fl. ounce
Extract of poppies	sixteen grains

Mix. As above.

*Derosne.*

## TINCTURE OF MONESIA.

R. Monesia	one ounce
Diluted alcohol	nine and a half fl. ounces
Water	two fl. ounces

Macerate, and decant.

*Donovan.*

R. Monesia	half an ounce
Alcohol	two fl. ounces
Water	seven and a-half fl. ounces

*St. Ange.*



As above. Used in injections; half a drachm to a drachm, to six fl. ounces of water.

#### MONESIA OINTMENT.

R. Monesia one drachm  
Lard one ounce  
Mix. *St. Ange.*

R. Monesia  
Water, each, one part  
White wax two parts  
Oil of almonds four parts  
Mix. *Derosne.*

As an application to indolent ulcers.

### MORA.

#### MULBERRIES.

Two species of *Morus* produce fruit which appears to have identical properties: *M. nigra*, a native of Europe, which is officinal in the foreign Pharmacopœias; and *M. rubra*, a native of the United States.

*Sex. Syst.* Monœc. tetrand. *Nat. Syst.* Moraceæ.

Mulberries are refreshing and laxative, and form the basis of a grateful drink in febrile cases.

#### SYRUP OF MULBERRIES.

R. Strained juice of mulberries one pint  
Sugar two and a-half pounds  
Dissolve with a gentle heat, set aside for twenty-four hours, remove the scum, and decant. *Lond. Ph.*

As an addition to cooling drinks in fever.

#### ROB OF MULBERRIES.

R. Strained juice of mulberries at will  
Evaporate to the consistence of honey. *Austr. Ph.*

Used as a detersive application to ulcers, and as an addition to gargles.

### MORPHIA.

#### MORPHIA.

R. Opium, sliced, one pound  
Distilled water  
Alcohol, each, sufficient  
Solution of ammonia six fl. ounces  
Macerate the opium with four pints of the water, for twenty-four hours; and having worked it with the hand, digest for twenty hours, and strain. Operate on the residue twice, in the same manner. Mix the infusions, evaporate to six pints, and filter; then add five pints of alcohol, and afterwards three fl. ounces of the solution of ammonia mixed with a half pint of alcohol. Let rest for twenty-four hours, add the rest of the ammonia, mixed as be-

fore, and set aside for twenty-four hours to crystallize. Purify the crystals by boiling them with two pints of alcohol, till dissolved, and filtering while hot through animal charcoal, and recrystallizing.

*U. S. Ph.*

R. Mix a concentrated infusion of opium with milk of lime (in which the lime is one-fourth the weight of the opium used); heat the mixture to boiling; filter while boiling hot, through linen, and add an excess of muriate of ammonia. As it cools, the morphia is precipitated. *Mohr.*

R. Muriate of morphia one ounce  
Distilled water one pint  
Dissolve, and add

Solution of ammonia five fl. drachms mixed with one fl. ounce of distilled water; shake together; wash the precipitate with distilled water, and dry by a gentle heat. *Lond. Ph.*

One-sixth of a grain is about equal in power to a grain of opium.

#### PILLS OF MORPHIA.

R. Morphia one grain  
Conserve of roses sufficient  
Mix, and make six pills. *Magendie.*

#### MORPHIA DRAUGHT.

R. Morphia a quarter of a grain  
Syrup of poppies one drachm  
Distilled water one fl. ounce  
Mix. At bedtime. *Brera.*

#### INJECTION OF MORPHIA.

R. Morphia two grains  
Almond oil  
Oil of lilies, each, half an ounce  
Mix.

R. Morphia two grains  
Yolk of egg one  
Oil of chamomile  
poppies, each, one ounce  
Mix. To ease pain in earache, in acute gonorrhœa and hemorrhoids. *Brera.*

### MORPHIÆ ACETAS.

#### ACETATE OF MORPHIA.

R. Powdered morphia one ounce  
Distilled water half a pint  
Acetic acid sufficient  
Mix the morphia with the water, and drop in the acid, constantly stirring, till the morphia is saturated and dissolved. Evaporate to consistence of syrup on a water-bath. Dry by a gentle heat, and powder. *U. S. Ph.*

Dose, one-eighth to a quarter of a grain.



## COMPOUND POWDER OF ACETATE OF MORPHIA.

R. Acetate of morphia                    one grain  
 Powdered foxglove                    six grains  
 Extract of pulsatilla                twelve grains  
 Sugar                                        one drachm  
 Mix, and form thirty-six powders. Four  
 a-day, in the cough and sleeplessness of  
 consumptive patients. *Cadet.*

## BOLUS OF ACETATE OF MORPHIA.

R. Acetate of morphia                    one grain  
 Olive oil                                    ten drops  
 Crumb of bread  
 Honey, each,                            sufficient  
 Mix, and make six boluses. *Brera.*

## PILLS OF ACETATE OF MORPHIA.

R. Acetate of morphia                    one grain  
 Conserve of roses                    sixteen grains  
 Mix, and make eight pills. *Mialhe.*

R. Acetate of morphia                    one grain  
 Golden sulphuret of  
 antimony  
 Extract of aconite, each,            two grains  
 Powdered liquorice  
 Honey, each,                            sufficient  
 Mix, and make eight pills. *Brera.*

R. Acetate of morphia                    one grain  
 Powdered digitalis                    six grains  
                   camphor                    ten grains  
                   gum Arabic                eight grains  
 Syrup of tolu                            sufficient  
 Beat into mass, and divide into six pills.  
 One may be taken every three hours.

*A. T. Thomson.*

R. Acetate of morphia                    fifteen grains  
 Ammon. sulphate of  
 copper                                    half a drachm  
 Inspissated bile  
 Powdered quassia, each, four scruples  
 Mix, and make one hundred pills. Five,  
 morning and evening, in diabetes mellitus.  
*Brendt.*

## SOLUTION OF ACETATE OF MORPHIA.

R. Acetate of morphia                    sixteen grains  
 Acetic acid                                two fl. drachms  
 Distilled water                        six fl. drachms  
 Mix. Dose, six to twenty-four drops.  
*Dunglison.*

## ALCOHOLIC SOLUTION OF MORPHIA.

R. Acetate of morphia                    sixteen grains  
 Alcohol                                    one fl. ounce  
 Dissolve. *Cottureau.*

## SYRUP OF ACETATE OF MORPHIA.

R. Acetate of morphia                    four grains

Dissolve in a small portion of water, and a  
 few drops of acetic acid, add to

Syrup                                        sixteen ounces  
*Par. Cod.*

## MIXTURE WITH ACETATE OF MORPHIA.

R. Solution of acetate of  
 morphia                                    twenty drops  
 Lactucarium                            ten grains  
 Infusion of chamomile                five ounces  
 Syrup of marsh mallow                half an ounce  
 Mix. Dose, a spoonful. *Beral.*

## CLYSTER WITH ACETATE OF MORPHIA.

R. Starch                                    one drachm  
 Hot water                                one pint  
 Acetate of morphia                    one grain  
 Mix. In chronic diarrhœa. *Cadet.*

## OINTMENT OF ACETATE OF MORPHIA.

R. Acetate of morphia six to eight grains  
 Lard                                        one to two drachms  
 Mix. As a friction in violent rheumatic  
 pains. *Foy.*

R. Acetate of morphia                    four grains  
 Mercurial ointment  
 Simple ointment, each,                two drachms  
 Mix. In frictions on the labia; twice a-  
 day, in cancer of the uterus. *Hildenbrand.*

## MORPHIÆ BIMECONAS.

## BIMECONATE OF MORPHIA.

R. Meconic acid                        two hundred grains  
 Boiling water                            sufficient  
 Dissolve, and add  
 Morphia                                    sufficient  
 to saturate. Evaporate to dryness.  
*Squire.*

Dose, one-fourth of a grain.

## SOLUTION OF BIMECONATE OF MORPHIA.

R. Bimeconate of morphia                ten grains  
 Rectified spirit                        one fl. drachm  
 Distilled water                        thirteen fl. drachms  
 Mix. About the strength of laudanum.  
*Beasley.*

## MORPHIÆ CITRAS.

## CITRATE OF MORPHIA.

R. Morphia                                sixteen grains  
 Crystallized citric acid                eight grains  
 Distilled water                        one ounce  
 Color with  
 Tincture of cochineal                two drachms  
 Mix. Dose, six to thirty drops, in the  
 twenty-four hours. *Magendie.*



## MORPHIÆ HYDRIODAS.

## HYDRIODATE OF MORPHIA.

R. Muriate of morphia two parts  
Iodide of potassium one part  
Make a strong solution of each, and mix, wash the precipitate in a little cold water, press in bibulous paper, redissolve in hot water, and let crystallize. *A. T. Thomson.*

## MORPHIÆ MURIAS.

## MURIATE OF MORPHIA.

R. Morphia in powder one ounce  
Distilled water half a pint  
Muriatic acid sufficient  
Mix the morphia with the water, drop in the acid, constantly stirring, till the morphia is saturated and dissolved. Evaporate by means of a water-bath to crystallization. Dry upon bibulous paper.

*U. S. Ph.*

## SOLUTION OF MURIATE OF MORPHIA.

R. Muriate of morphia one drachm and a-half  
Rectified spirit five fl. ounces  
Distilled water fifteen fl. ounces  
Mix the spirit and water, and dissolve the muriate in the mixture with the aid of heat.

*Ed. Ph.*

Eighteen minims contain one-sixth of a grain of the muriate, equal to a grain of opium.

## SYRUP OF MURIATE OF MORPHIA.

R. Muriate of morphia one grain  
Syrup four ounces  
Mix. Dose, a teaspoonful.

*Cadet.*

## MIXTURE OF MURIATE OF MORPHIA.

R. Muriate of morphia sixteen grains  
Acetic acid four drops  
Distilled water one fl. ounce  
Alcohol one fl. drachm  
Mix. Dose, six to twenty drops.

*Cadet.*

## COMPOUND SYRUP OF MURIATE OF MORPHIA.

R. Muriate of morphia two grains  
Syrup of pinks ten ounces  
balm four ounces  
orange flowers two ounces  
Mix. Dose, half an ounce to an ounce.

*Cadet.*

## LOZENGES OF MURIATE OF MORPHIA.

R. Muriate of morphia one scruple  
Tincture of tolu half a fl. ounce  
Sugar twenty-five ounces  
Dissolve the muriate in a little hot water, mix it and the tincture with the sugar,

beat into a mass with mucilage, and divide into lozenges of fifteen grains. Each lozenge contains a fortieth of a grain of the muriate.

*Ed. Ph.*

## LOZENGES OF MURIATE OF MORPHIA AND IPECACUANHA.

R. Made as the last, with the addition of one drachm of ipecacuanha.

*Ed. Ph.*

## MORPHIA NITRAS.

## NITRATE OF MORPHIA.

R. Morphia at will  
Dilute nitric acid sufficient  
Saturate, dissolve, evaporate, and let crystallize.

*Giordano.*

## MORPHIÆ PHOSPHAS.

## PHOSPHATE OF MORPHIA.

Make like the last, using dilute phosphoric acid instead of nitric.

## MORPHIÆ SULPHAS.

## SULPHATE OF MORPHIA.

R. Morphia, in powder, one ounce  
Distilled water half a pint  
Diluted sulphuric acid sufficient  
Mix the morphia with the water, drop in the acid, constantly stirring till the morphia is saturated and dissolved. Evaporate on water-bath and let crystallize. Dry crystals on bibulous paper.

*U. S. Ph.*

Dose, one-eighth to a quarter of a grain.

## BOLUS OF SULPHATE OF MORPHIA.

R. Sulphate of morphia two grains  
Oil of almonds sufficient  
Sulphate of iron four grains  
Crumb of bread  
Honey, each, sufficient  
Mix, and make eight boluses.

R. Sulphate of morphia one grain  
Ipecacuanha three grains  
Oil of almonds twelve drops  
Extract of aconite two grains  
Liquorice powder  
Honey, each, sufficient  
Mix, and make six boluses. One, every three or four hours.

*Brera.*

## PILLS OF SULPHATE OF MORPHIA.

R. Sulphate of morphia two grains  
Cyanide of potassium four grains  
Mucilage sufficient  
Mix, and make twenty-four pills. One every six hours, in neuralgia.

*Rougier.*



R. Sulphate of morphia      one grain  
 Olive oil      sufficient  
 Ipecacuanha      three grains  
 Nux vomica      two grains  
 Crumb of bread  
 Honey, each,      sufficient  
 Mix, and make six pills.      One, every two hours. *Brera.*

## SOLUTION OF SULPHATE OF MORPHIA.

R. Sulphate of morphia      eight grains  
 Distilled water      half a pint  
 Dissolve. *U. S. Ph.*

One fl. drachm contains an eighth of a grain.

R. Sulphate of morphia      four grains  
 Distilled water      one fl. ounce  
 Dissolve. One fl. drachm contains half a grain. *Copland.*

## SYRUP OF SULPHATE OF MORPHIA.

R. Sulphate of morphia      one grain  
 Dissolve in a little water, and add to  
 Syrup      four ounces  
 Each ounce contains one quarter of a grain of sulphate. *Cadet.*

## MIXTURE OF SULPHATE OF MORPHIA.

R. Sulphate of morphia      sixteen grains  
 Acetic acid      four drops  
 Distilled water      one ounce  
 Alcohol      one drachm  
 Mix. Dose, six to twenty drops. *Magendie.*

## LOTION OF SULPHATE OF MORPHIA AND BORAX.

R. Sulphate of morphia      six grains  
 Borax      half an ounce  
 Rose water      eight fl. ounces  
 Mix. As an application in pruritus vaginæ, washing first with tepid soap and water. *Meigs.*

## MORPHIÆ TARTRAS.

## TARTRATE OF MORPHIA.

R. Morphia      at will  
 Saturate with  
 Solution of tartaric acid      sufficient  
 Evaporate and crystallize. *A. T. Thomson.*

## MOSCHUS.

## MUSK.

This is a peculiar concrete substance obtained from the *Moschus moschiferus*, a small animal of the deer kind, inhabiting the mountainous regions of central Asia. The musk is secreted in the male, in an oval sac, situated near the generative organs. It is found in commerce in these sacs; it is concreted or granular; of a brownish color;

soft and greasy to the touch; of a powerful, penetrating odor, and of a bitter, unpleasant, somewhat acrid taste. From its high price it is very liable to adulteration. It is antispasmodic and stimulant, and has been much used in spasmodic diseases of all kinds, as well as a stimulant in low states of the system. The dose is from five to ten grains.

## POWDER OF MUSK.

R. Musk      three grains  
 Opium      half a grain  
 Gum Arabic      one scruple  
 Sugar      two drachms

Triturate together, and divide into six powders. One, every three hours, in whooping-cough. *Augustin.*

R. Musk      sixteen grains  
 Valerian      twenty-four grains  
 Camphor      eight grains

Mix. As an antispasmodic, in hysteria, &c.; in doses of three to twelve grains. *Guibourt.*

This is called by Jourdain, Tonquin powder, but this name belongs rather to the following.

R. Musk      sixteen grains  
 Cinnabar      twelve grains

Mix. For a single dose. *Spielmann.*

Once celebrated as a remedy in hydrophobia.

## MUSK BOLUS.

R. Musk      five to ten grains  
 Camphor      five grains  
 Syrup      sufficient

Make a bolus. Antispasmodic. *Ellis.*

R. Musk  
 Carbonate of ammonia, each,      ten grains  
 Conserve of roses      sufficient

Make a bolus. One, every three hours. In mortification, attended with spasmodic action. *Ellis.*

## MUSK PILLS.

R. Musk      eight grains  
 Opium      two grains  
 Camphor      twenty-four grains  
 Syrup      sufficient

Mix, and form six pills. To be taken during the day, in hospital gangrene. *Dupuytren.*

R. Musk      ten grains  
 Camphor      one scruple  
 Gum ammoniac      two scruples  
 Opium      four grains

Mix, and make pills of four grains. Four or five in the twenty-four hours, in nervous disorders. *Richard.*

R. Musk      twelve grains  
 Castor      twenty-four grains  
 Assafetida      thirty grains



Tincture of valerian sufficient  
Make twenty-four pills. Four, three  
times a-day, in nervous complaints.

*Radius.*

#### MUSK MIXTURE.

R. Musk two drachms  
Sugar  
Gum Arabic, each, one drachm  
Distilled water six fl. ounces  
Mix. A tablespoonful, every hour or  
two, in low fevers. *Ellis.*

R. Musk mixture six fl. ounces  
Paregoric elixir half fl. ounce  
Ammon. tinc. of  
valerian one fl. drachm  
Mix. A teaspoonful, three or four times  
a-day, in pertussis in children. *Ellis.*

R. Musk two grains  
Fennel water six fl. ounces  
Laudanum ten drops  
Syrup of poppies two drachms  
Mix. A spoonful, every hour, in tris-  
mus. *Cheselden.*

R. Musk mixture  
Camphor water, each, three fl. ounces  
and a-half  
Syrup of ginger  
Spirit of sulphuric  
ether, each, two fl. drachms  
Mix. A tablespoonful, every three or  
four hours, in low fevers. *Ainslie.*

#### MUSK CLYSTER.

R. Musk ten grains  
Valerian half an ounce  
Starch half a drachm  
Boiling water eight ounces  
Mix. As an antispasmodic and excitant.  
*Radius.*

R. Musk twelve grains  
Sugar two scruples  
Spirit of ammonia thirty drops  
Infusion of flaxseed four fl. ounces  
Mix. For children with convulsions.  
*Ellis.*

#### TINCTURE OF MUSK.

R. Musk one part  
Alcohol twelve parts  
Digest for twelve days, and filter. Dose,  
thirty to sixty drops. *Guibourt.*

### MUCUNA.

#### COWHAGE.

This is the bristles of the pods of *Mucuna pruriens*, a perennial, twining plant, native of the warmer parts of America, bearing somewhat

curved brown pods, covered with short bristles, which, when dry, readily separate.

*Sex. Syst.* Diadelph. decand. *Nat. Syst.* Fagaceæ.

Decandolle, Prod. ii. 405. Griffith, Med. Bot. 242.

The pod is shaped somewhat like the italic S, and is covered with brown, bristly hairs, which adhere to any substance coming in contact with them. These hairs or spicula are possessed of anthelmintic powers, but whether they act mechanically or not, has not been fully ascertained, though the probability is that they do.

#### ELECTUARY OF COWHAGE.

R. Cowhage two scruples  
Syrup half an ounce  
Mix. A teaspoonful every morning, fast-  
ing, for three days, to be followed by a  
dose of castor oil; in cases of lumbrici.

*Correa.*

R. Cowhage one drachm  
Honey sufficient  
to make electuary. To be given as above.  
*Ellis.*

### MYRISTICA.

#### NUTMEG.

The nutmeg is the kernel of the fruit of *Myristica moschata*, a small tree, a native of the Molucca Islands, and now cultivated in various tropical regions.

*Sex. Syst.* Diœc. monadelph. *Nat. Syst.* Myristicaceæ.

Thunberg. Act. Holm. 1782. Griffith, Med. Bot. 109.

The nutmeg is an oval, oblong almond, of a lightish brown color externally, and of a reddish gray with red veins, internally, of an agreeable, fragrant odor, warm, aromatic taste, and unctuous feel. It is principally used for culinary purposes, but is also employed in medicine as a stimulant, and to disguise the taste of nauseous remedies.

#### AROMATIC POWDER.

R. Cinnamon  
Ginger, each, two ounces  
Cardamom seeds  
Nutmeg, grated, each, one ounce  
Rub together into very fine powder.  
*U. S. Ph.*

Stimulant and carminative; dose, ten to thirty grains.

#### GOELIS' ANTIHECTIC POWDER.

R. Burnt hartshorn  
Powdered nutmeg  
Roasted laurel berries, each, one part  
Liquorice three parts  
Mix, and make a powder. Advised in the hectic attendant on scrofulous affections.

*Augustin.*

#### STIMULATING CLYSTER.

R. Powdered nutmeg one drachm



Powdered columbo one drachm  
 salep one scruple  
 Infusion of flaxseed four fl. ounces  
 Mix. As a stimulating enema. *Ammon.*

## SPIRIT OF NUTMEG.

R. Nutmeg, bruised, two ounces  
 Diluted alcohol one gallon  
 Water a pint  
 Mix, and distil one gallon, with a slow fire.  
*U. S. Ph.*

Dose, one to two fl. drachms; principally used to flavor other medicines.

## COMPOUND SPIRIT OF NUTMEG.

R. Nutmeg two ounces  
 Lemon peel  
 Orange peel  
 Spearmint  
 Balm, each, one ounce  
 Diluted alcohol four pints  
 Water twelve pints  
 Distil three pints. *Van Mons.*

As a stomachic, and an external application to contusions.

## MYROXYLON.

## BALSAM OF PERU.

This balsam is the product of *Myroxylon Peruvianum*, a large tree, indigenous to South America, where it is called *quinquina*.

*Sex. Syst.* Decand. monog. *Nat. Syst.* Fabaceae.

*Linn. Sp. Pl.* 233. *Griffith, Med. Bot.* 248.

The balsam is a thick, viscid fluid, of a dark, reddish-brown color, an agreeable, fragrant odor, and a warm, bitterish taste, occasioning a pungent sensation in the throat. It is stimulating, tonic, and expectorant, and has been much used in pectoral complaints, &c., and also externally as an application to indolent ulcers. The dose is about half fl. drachm.

## PILLS OF BALSAM OF PERU.

R. Balsam of Peru one drachm  
 Extract of bitter polygala two drachms  
 Marsh mallow sufficient  
 to make one hundred and twenty pills.  
 Ten, four times a-day, in chronic mucous discharges. *Schubert.*

## ACOUSTIC BALSAM.

R. Balsam of Peru half a drachm  
 Narcotic oil  
 Onion juice, each, one ounce  
 Mix. A dossil of cotton saturated with this oil, is to be introduced into the deaf ear, provided there is no inflammation or violent pain.

R. Beef gall three fl. drachms  
 Balsam of Peru one fl. drachm

Mix. To be occasionally dropped into the ear to correct a fetid discharge, syringing it also, daily, with a weak solution of soap and water. *Hugh Smith.*

## LINIMENT FOR CHILBLAINS.

R. Balsam of Peru half a drachm  
 Muriatic ether  
 Laudanum, each, two drachms  
 Mix. As a friction. *Henschel.*

## MAMMILLARY LOTION.

R. Balsam of Peru one drachm  
 Yolk of egg one  
 Spirit of wild thyme three ounces  
 Mix. The sore nipple is to be bathed with this, and then sprinkled with a powder, composed of one drachm of Peruvian bark, and two drachms of gum Arabic. *Iverg.*

## BALSAM OF PERU MIXTURE.

R. Balsam of Peru two drachms  
 Yolk of egg one  
 Extract of cinchona six drachms  
 Honey of roses three ounces  
 Mix. Two dessertspoonfuls, four times a-day, in chronic mucous discharges.

*St. Marie.*

R. Balsam of Peru half a drachm  
 Mucilage of gum Arabic sufficient  
 Cinnamon water  
 Water, each, half fl. ounce

Mix. To be taken three or four times a-day, as an expectorant in chronic catarrh. *Ellis.*

## SYRUP OF BALSAM OF PERU.

R. Tincture of balsam of Peru one fl. ounce  
 Tepid simple syrup two pounds  
 Mix, and agitate briskly. Dose, from one to two drachms. *Lisbon Ph.*

## TINCTURE OF BALSAM OF PERU.

R. Balsam of Peru one part  
 Alcohol eight parts  
 Macerate for some days, and filter. *Guibourt.*

Principally used as an external application to ulcers, &c.

## BALSAM OF PERU COLLUTORY.

R. Tincture of balsam of Peru one part  
 guaiacum four parts  
 Mix. A teaspoonful to a glass of water, to rinse the mouth. *Taddei.*

## LOCATELLI'S BALSAM.

R. Olive oil six ounces  
 Yellow wax four ounces  
 Wine five fl. ounces



Melt together by a gentle heat, till all moisture is evaporated, and add

Venice turpentine six ounces  
Balsam of Peru two drachms  
Red saunders half an ounce

Mix. Formerly used in phthisis, but now employed only as an external application. This is the original formula. *Spielmann.*

#### OINTMENT OF BALSAM OF PERU.

R. Balsam of Peru  
Spermaceti ointment equal parts  
Mix. As a dressing to painful ulcers. *Radiis.*

#### COMPOUND OINTMENT OF BALSAM OF PERU.

R. Lard two ounces  
White wax four drachms  
Melt in a water-bath, and add  
Balsam of Peru two drachms  
Oil of lavender twelve minims  
As an application to promote the growth of the hair. *Copland.*

#### PLASTER OF BALSAM OF PERU.

R. Powdered carbonate of lead sixteen parts  
litharge eight parts  
Rose oil forty-eight parts  
White wax thirty-two parts  
Melt together, and, at close of operation, add  
Balsam of Peru two parts  
As an application to indolent ulcers. *Foy.*

### MYRRHA.

#### MYRRH.

Myrrh is the concrete juice of *Balsamodendron myrrha*, a small shrubby tree, with spiny branches, a native of Arabia, &c.

*Sex. Syst.* Octand. monog. *Nat. Syst.* Amyridaceæ.

*Nees. Offic. Planz.* i. 17. *Griffith, Med. Bot.* 171.

Myrrh occurs in tears, or in agglutinated masses, of various shades of color; the best is of a reddish yellow color, somewhat translucent, having a peculiar, aromatic odor and a warm, bitter taste. It is a stimulating tonic. It is given in a variety of diseases, and is used externally as an application to foul ulcers, aphthæ, &c. The dose is from ten to thirty grains, usually in combination.

#### POWDER OF MYRRH AND IPECACUANHA.

R. Powdered myrrh twelve grains  
ipecacuanha six grains  
nitre half a drachm  
Mix, and divide into four powders. One, every fourth hour. Stimulating expectorant. *Paris.*

#### EMMENAGOGUE POWDER.

R. Myrrh twelve grains  
Saffron three grains  
Oil of cloves one drop  
Rub into powder. *Augustin.*

#### PILLS OF MYRRH AND ZINC.

R. Sulphate of zinc ten grains  
Powdered myrrh one drachm and a-half  
Conserve of roses sufficient  
Mix, and form twenty pills. Two, twice a-day. In pertussis. *Paris.*

#### PILLS OF MYRRH AND SULPHATE OF IRON.

R. Myrrh  
Sulphate of iron, each, two scruples  
Carbonate of potassa  
Soap, each, half a drachm  
Rub into mass, and form forty pills. Two, thrice a-day. In amenorrhœa. *Ellis.*

#### PILLS OF MYRRH AND SQUILL.

R. Myrrh one drachm and a-half  
Dried squill half a drachm  
Extract of henbane two scruples  
Distilled water sufficient  
Rub into mass, and form thirty pills. Two, night and morning. In catarrh and phthisis. *Paris.*

#### PILLS OF MYRRH AND CANADA BALSAM.

R. Myrrh three drachms  
Canada balsam one drachm and a-half  
Opium half a drachm  
Mix, and make pills of two grains. Two to four, every hour, in ulcerated phthisis. *Augustin.*

#### ALKALINE SOLUTION OF MYRRH.

R. Myrrh two ounces  
Sub. carbonate of soda one drachm  
Boiling water eight fl. ounces  
Macerate for two days, and filter. *Swediaur.*

#### COMPOUND EXTRACT OF MYRRH.

R. Myrrh two ounces  
Gum Arabic two drachms  
Triturate together; add sufficient water to make a thick emulsion, and mix well with the  
Extract of couchgrass four ounces  
Advised in phthisis, and ulcers of the uterus. Dose, one to three drachms, mixed with water, two or three times a-day. *Swediaur.*

#### MYRRH COLLUTORY.

R. Lime water one fl. ounce and a-half  
Tincture of myrrh two fl. drachms  
Honey of roses two drachms  
Mix. *Pharm. Chirurg.*



R. Tincture of myrrh two fl. drachms  
 Infusion of sage six fl. ounces  
 Honey of roses two ounces  
 Mix. *Augustin.*

## MYRRH GARGLE.

R. Tincture of myrrh  
 Vinegar, each, two fl. ounces  
 Honey one ounce  
 Infusion of contra-

yerva one pint and a-half  
 Mix. In putrid sore throat, scarlatina,  
 &c. *Fothergill.*

R. Honey of roses one ounce and a-half  
 Barley water twelve fl. ounces  
 Tincture of myrrh six fl. drachms  
 Vinegar one fl. ounce

Mix. As a gargle in putrid sore throat.  
*Ainslie.*

## STIMULATING INJECTION.

R. Myrrh one ounce  
 Quicklime two ounces  
 Water two pints

Infuse for a few days, and decant. As an  
 injection in fistulous ulcers. *St Marie.*

## TINCTURE OF MYRRH.

R. Myrrh, bruised, four ounces  
 Alcohol three pints

Macerate for fourteen days, and filter.  
*U. S. Ph.*

## TINCTURE OF MYRRH AND HELLEBORE.

R. Tincture of myrrh one fl. ounce  
 black helle-  
 bore half a fl. ounce  
 Spanish flies two fl. drachms

Mix. As an emmenagogue, in doses of  
 thirty drops, three times a-day, in a little  
 sugar and water. *Ellis.*

## GRIFFITH'S MYRRH MIXTURE.

R. Myrrh one drachm  
 Sulphate of iron one scruple  
 Carbonate of potassa one drachm  
 Sugar two drachms  
 Water six fl. ounces

Make mixture. As a tonic in phthisis, in  
 tablespoonful doses, according to circum-  
 stances. *Ellis.*

## COMPOUND MYRRH MIXTURE.

R. Powdered myrrh one drachm  
 Carbonate of potassa half a drachm  
 Sulphate of iron twelve grains  
 Mucilage gum Arabic two fl. drachms  
 Decoction of liquo-  
 rice six and a-half fl. ounces  
 Spirit of allspice one fl. ounce

Rub the myrrh, potassa, sulphate of iron,  
 and mucilage, well together; add gradually  
 the other ingredients. Dose, a tablespoon-  
 ful, twice or thrice a-day. *Babington.*

Nearly the same as Griffith's mixture, and  
 used in the same cases.

## OIL OF MYRRH.

R. Myrrh two parts  
 Washed sand three parts  
 Distil, separate the oil that passes, and  
 rectify it. *Wirt. Ph.*

## MYRRH PLASTER.

R. Balsam of Peru  
 Camphor  
 Powdered myrrh, each, one ounce and  
 a-half

Lead plaster thirty-two ounces  
 Triturate the three first ingredients to-  
 gether, and when intimately mixed, add  
 the melted plaster, and stir until chilled;  
 then form into rolls of half a pound each.

## N

## NAPHTHALINA.

## NAPHTHALINE

Is a product of the distillation of coal tar, and  
 is deposited from the rectified oil of coal tar, in  
 white, shining, concrete crystals, and may be  
 purified by dissolving in alcohol and recrystalliz-  
 ing. It is an active expectorant, in doses of eight  
 to thirty grains.

## SYRUP OF NAPHTHALINE.

R. Naphthaline sixteen grains

Dissolve in smallest quantity of hot alco-  
 hol, and triturate with

Syrup four ounces  
*Dupasquier.*

## OINTMENT OF NAPHTHALINE.

R. Naphthaline one scruple  
 Lard five drachms

Mix. As an application in dry tetter,  
 lepra, psoriasis, &c. *Emery.*



## LOZENGES OF NAPHTHALINE.

R. Naphthaline five scruples  
 Sugar twenty ounces  
 Oil of aniseed  
 Mucilage of tragacanth, each, sufficient  
 Mix, and form lozenges of fifteen grains.  
 Expectorant. One occasionally, to the  
 amount of twenty a-day. *Dupasquier.*

## NARCOTINA.

## NARCOTINE

Is a white, tasteless, inodorous principle, existing in opium. It forms bitter salts with the acids. It has been used with success as an antiperiodic, in doses of three grains, three times a-day.

R. Residue of opium, exhausted by  
 Water at will  
 Dry it, powder it coarsely, add cold acetic acid, express, and filter; add an excess of ammonia, wash the precipitate with cold water, dissolve in boiling alcohol, decolorize by means of animal charcoal, let cool, and crystallize.

## BOLUS OF NARCOTINE.

R. Narcotine one scruple  
 Oil of almonds six drops  
 Crumb of bread  
 Honey, each, sufficient  
 Mix, and make eight boluses. One, three times a-day, in the apyrexia of intermittents. *Brera.*

## MIXTURE OF NARCOTINE.

R. Narcotine one drachm  
 Lemon water three fl. ounces  
 Mix. Dose, a spoonful. *Brera.*

## NARCOTINÆ MURIAS.

## MURIATE OF NARCOTINE.

R. Opium two pounds  
 Alcohol twenty pounds  
 Rub together, gradually adding the spirit, till the opium is exhausted, decant, and press residue; to solution add ammonia, till turbid. Distil off one-fourth of alcohol, and let the product cool; wash the deposit with water, and then with a drachm of muriatic acid mixed with a quart of water. Filter the solution, and evaporate to dryness. Dose, about the same as of narcotine. *O'Shaughnessy.*

## NUX VOMICA.

## NUX VOMICA

Is the seed of the *Strychnos nux vomica*, a moderate sized tree, a native of many parts of the East Indies, having a very bitter wood.

*Sex. Syst.* Pentand. monog. *Nat. Syst.* Loganiaceæ.

*Linn. Sp. Pl.* 271. *Griffith, Med. Bot.* 469.

The seeds are flat, peltate, with a narrow annular striæ, somewhat downy on the surface; they are of a horny consistence, inodorous, and of a bitter, acrid, somewhat nauseous taste. *Nux vomica* is a violent excitant of the cerebro-spinal system, and in large doses an active poison. In small doses, frequently repeated, it is tonic, diuretic, and even laxative. It owes its energetic properties to the presence of strychnia and brucia. It is principally used in paralysis, in doses of about five grains, three or four times a-day.

## POWDER OF NUX VOMICA.

R. Powdered nux vomica three grains  
 Gum Arabic  
 Sugar, each, twelve grains  
 Mix. Advised in chronic dysentery. One, every day. *Soubeiran.*  
 R. Powdered nux vomica eighteen grains  
 ipecacuanha

twenty-four grains  
 Powdered rhubarb one drachm  
 Prepared chalk two scruples  
 Oil of peppermint two drops  
 Mix. To be divided into twelve powders. One every three or four hours, in cardialgia, &c. *Vogt.*

## EXTRACT OF NUX VOMICA.

R. Nux vomica one pound  
 Alcohol sufficient  
 Expose the nux vomica to steam, till softened, slice, dry, and reduce it to powder. Place this in a displacement apparatus, add alcohol till it comes away without bitterness. Distil off the alcohol from the tincture, and evaporate to a proper consistence. *U. S. Ph.*

Dose, half a grain to two grains.

## TINCTURE OF NUX VOMICA.

R. Powdered nux vomica one part  
 Alcohol four parts  
 Macerate for fourteen days, and filter. *Cottreau.*

Dose, five to twenty drops, in paralysis.

## COMPOUND TINCTURE OF NUX VOMICA.

R. Extract of nux vomica twenty-four grains  
 Camphor one drachm  
 Tincture of pellitory one fl. ounce



Mix. Dose, twenty drops, four times a-day, in paralysis. *Radius.*

R. Tincture of nux vomica  
Spanish flies, each,  
one fl. drachm

Phosphoric ether one fl. scruple  
Mix. Thirty drops, three or four times a-day, in paralysis. *Radius.*

#### PILLS OF NUX VOMICA.

R. Powdered nux vomica thirty grains  
Conserve of roses sufficient  
Mix, and make ten pills; one, twice or thrice a-day, in paralysis, closely watching the effects. *Ellis.*

#### PILLS OF EXTRACT OF NUX VOMICA.

R. Extract of nux vomica one scruple  
liquorice seven scruples  
Mix, and make eighty pills. Two to four, two or three times a-day, in paralysis. *Radius.*

#### COMPOUND PILLS OF NUX VOMICA.

R. Extract of nux vomica six grains

Black oxide of iron + one drachm  
Mix, and make twenty-four pills. Three a-day, in atonic incontinence of urine. *Mondiere.*

#### PILLS OF NUX VOMICA AND ALOES.

R. Extract of nux vomica ten grains  
Pills of aloes and myrrh four scruples  
Mix well, and form thirty-six pills. One or two, night and morning. *Copland.*

#### MIXTURE OF NUX VOMICA.

R. Extract of nux vomica ten grains  
Elder water six fl. ounces  
Mucilage of gum Arabic  
Spirit of mallows, each, one fl. ounce  
Mix. A spoonful every two hours, in chronic dysentery. *Ammon.*

#### LOTION OF NUX VOMICA.

R. Extract of nux vomica eight grains  
Alcohol two fl. ounces  
Strong water of ammonia half fl. ounce  
Mix. As a stimulating lotion to paralyzed limbs. *Radius.*

### O

#### OLEUM ANIMALE EMPYREUMATICUM.

##### DIPPEL'S ANIMAL OIL.

R. Oil of hartshorn at will  
Distil off one-fourth, by a slow heat, on a sand-bath, keeping it protected from the light. *Van Mons.*

Antispasmodic, diaphoretic, and anodyne, in doses of five or ten drops; poisonous in large doses.

##### LINIMENT OF DIPPEL'S OIL.

R. Dippel's oil one drachm  
Oil of almonds six drachms  
Turpentine soap half an ounce  
Mix. As an exciting liniment, also as an application to the abdomen, in cases of worms. *Sundelin.*

##### TINCTURE OF DIPPEL'S OIL.

R. Dippel's oil one part  
Sulphuric ether fifteen parts  
Dissolve. Fifteen to thirty drops, as a stimulant and antispasmodic. *Beral.*

##### MIXTURE OF DIPPEL'S OIL.

R. Dippel's oil one drachm  
Hoffmann's anodyne three drachms  
19

Dissolve. Twenty to thirty drops, four times a-day, in chorea and tetanus. *Radius.*

#### OLEUM CAJUPUTI.

##### OIL OF CAJEPUT.

This is a fluid, transparent, green oil, of an aromatic and pleasant odor, and a warm, pungent taste; obtained from the *Melaleuca cajuputi*, a small tree, a native of the Molucca islands.

*Ser. Syst.* Polydelph. icosand. *Nat. Syst.* Myrtaceæ.

Maton. Lond. Pharm. 1800. Griffith, Med. Bot. 296.

It is also furnished by other species. It is an active stimulant, when given internally, in doses of a few drops, and acts as a rubefacient, when applied to the skin.

##### MIXTURE OF OIL OF CAJEPUT.

R. Oil of cajeput half a drachm  
Dippel's animal oil one drachm  
Mix. As a stimulant. Dose, five to fifteen drops. *Augustin.*

##### CAJEPUT LINIMENT.

R. Cajeput oil  
Camphor, each, three drachms



Soft soap	two ounces
Alcohol	one pint
Water of ammonia	two fl. ounces
Mix. As an embrocation.	<i>Chapman.</i>

## OLEUM MORRHUÆ.

## COD LIVER OIL.

This, which is also called *Oleum jecoris aselli*, is procured from the livers of several kinds of fish, but principally from the Cod, though that from the Ray is said to be preferable. There are several varieties; one, clear and limpid, the other dark colored, and somewhat acrid, which is said by many writers to be the most powerful. The pale oil has a faint smell, and a bland taste. It has obtained much celebrity as an alterative, in strumous affections; in chronic gout, rheumatism, and cutaneous diseases, &c. The dose for an adult, is from half a spoonful to three spoonfuls, two or three times a-day.

## MIXTURE OF COD LIVER OIL.

R. Cod liver oil	four fl. ounces
Gum Arabic	
Sugar, each,	two drachms
Cinnamon, or mint	
water	four fl. ounces

Mix. One or two tablespoonfuls, morning and evening. *Ellis.*

R. Cod liver oil	four fl. ounces
Solution of carbonate of potassa	half fl. ounce
Syrup of orange peel	half fl. ounce
Peppermint water	seven fl. ounces

Mix. Dose, one and a-half fl. ounces to three fl. ounces. *Beasley.*

R. Cod liver oil	one fl. ounce
Solution carbonate of potassa	two fl. drachms
Syrup of orange peel	one fl. ounce
Oil of calamus	three drops

Mix. Dose, one to two fl. drachms, night and morning, for rickets in children. *Fehr.*

R. Cod liver oil	
Syrup of orange peel	
Aniseed water, each,	one fl. ounce
Oil of calamus	three drops

Mix. Three spoonfuls a-day, in rachitis, and gouty swellings. *Phæbus.*

## SYRUP OF COD LIVER OIL.

R. Cod liver oil	eight parts
Powdered gum Arabic	five parts
Simple syrup	four parts
Make an emulsion, and add	
Sugar	twenty-four parts
Dissolve by gentle heat, and add	
Orange-flower water	two parts

*Duclou.*

## LINIMENT OF COD LIVER OIL.

R. Cod liver oil	one fl. ounce
Water of ammonia	half fl. ounce
Mix.	<i>Brach.</i>

R. Cod liver oil	four fl. drachms
Lead water	two fl. drachms
Yolk of egg	three drachms

Mix. As an application to ulcers, &c. *Brefeld.*

## OINTMENT OF COD LIVER OIL.

R. Cod liver oil	three fl. ounces
Spermaceti	six drachms
White wax	two drachms
Melt together, and stir till cold.	<i>Beasley.</i>

R. Caustic soda	one ounce
Water	two fl. ounces and a-half

Dissolve, and add  
Cod liver oil eight fl. ounces  
Agitate briskly. *Deschamps.*

R. Cod liver oil	ten parts
Lead water	five parts
Lard	ten parts

Mix.

## COMPOUND OINTMENT OF COD LIVER OIL.

R. Cod liver oil	one fl. drachm
Red oxide of mercury	four grains
Simple cerate	two scruples
Mix.	<i>Cunier.</i>

R. Cod liver oil	
Extract of wood	
soot, each,	two drachms
Citrine ointment	one drachm
Beef marrow	six ounces
Mix.	<i>Carron.</i>

In opacities of the cornea, and scrofulous ophthalmia.

## COMPOUND OIL OF COD LIVER.

R. Cod liver oil	one fl. drachm
Walnut oil	two fl. drachms
Mix. A drop or two to be introduced between the eyelids, in opacities of the cornea.	<i>Radius.</i>

## OLEUM OLIVÆ.

## OLIVE OIL.

The oil of the fruit of *Olea Europæa*, a small tree, originally from Syria, but now extensively cultivated in the countries bordering on the Mediterranean.

*Sax. Syst. Diand. monog. Nat. Syst. Oleaceæ. Linn. Sp. Pl. 11. Griffith, Med. Bot. 442.*

Good olive oil is an unctuous fluid, of a pale yellow or greenish color; almost inodorous, and of a bland taste. It is principally used as an article of food, but is also employed in medicine as a demulcent, emollient, and laxative, and in



the composition of liniments, ointments, &c. The dose, as a laxative, is about a fl. ounce.

## OLIVE OIL MIXTURE.

- R. Syrup of gum four fl. ounces  
Olive oil half fl. ounce  
Mix. As a laxative. *Radius.*
- R. Olive oil eight fl. ounces  
Aromatic spirit of ammonia two fl. drachms  
Mix. Three spoonfuls, night and morning, as an anthelmintic. *Ellis.*
- R. Olive oil one fl. ounce  
Solution carbonate of potassa half fl. drachm  
Mint water seven fl. ounces  
Mix. *Guy's Hospit.*
- R. Olive oil one fl. ounce  
Water of carbonate of ammonia one fl. drachm  
Mint water seven fl. ounces  
Mix. *Guy's Hospit.*
- R. Olive oil mixture eight fl. ounces  
Manna one ounce and a-half  
Mix. Dose, three spoonfuls as a laxative. *St. Bart's Hospit.*

## OLIVE OIL CLYSTER.

- R. Common salt one tablespoonful  
Olive oil  
Molasses, each, two tablespoonfuls  
Warm water one pint  
Mix. *U. S. Dispens.*
- R. Manna one ounce  
Compound decoction of chamomile ten fl. ounces  
Dissolve, and add  
Olive oil one fl. ounce  
Sulphate of magnesia half an ounce  
*Dub. Ph.*

## OLEUM RICINI.

## CASTOR OIL.

The oil of the seeds of *Ricinus communis*, a perennial tree in tropical countries, but an annual herbaceous plant, in temperate latitudes. The seeds are ovate, compressed, bean-like, of a grayish ash color, marbled with reddish brown, not unlike the dog tick in appearance.

*Ser. Syst.* Monœc. monad. *Nat. Syst.* Euphorbiacæ.

Linn. Sp. Pl. 1430. Griffith, Med. Bot. 599.

The oil, which is generally obtained by expression, is a thick, viscous, colorless fluid, with a faint, but unpleasant odor, and a mild, but nauseous taste, followed by a slight sensation of acidity. It is a mild, but prompt cathartic, acting rather as an evacuant than as an excitant of the alvine secretions. The dose is about a fl. ounce; for infants, from one to four fl. drachms. Its dis-

gusting taste is best disguised by mixing it with froth of porter.

## OLEAGINOUS MIXTURE.

- R. Castor oil one ounce  
Powdered gum Arabic two drachms  
Sugar one drachm  
Rub together, and add gradually  
Mint water four fl. ounces  
Triturate well. A tablespoonful, every hour or two hours, till it acts. *Ellis.*
- In some cases, as in dysentery, it is of benefit, to add thirty or forty drops of laudanum to the mixture.
- R. Castor oil  
Mucilage of gum Arabic  
Syrup of orgeat, each, two ounces  
Water ten fl. ounces  
Make an emulsion. Dose, as last. *Beral.*
- R. Castor oil one ounce  
Yolk of egg one  
Water of orange peel two fl. drachms  
Water two fl. ounces  
Make an emulsion. *Radius.*
- R. Castor oil one ounce  
Yolk of egg one  
Orange-flower water  
Simple syrup, each, half an ounce  
Water two ounces  
Make an emulsion. *Cotteneau.*
- R. Castor oil eleven drachms  
Powdered tragacanth half a drachm  
White sugar seventy-five grains  
Water two and a-half fl. ounces  
Syrup of orange-flowers six fl. drachms  
Triturate the tragacanth with the sugar; add the syrup, and rub well in a mortar, until the mucilage begins to thicken, then add the oil, and continue rubbing till it is homogeneous, adding the water gradually, during the process. *Mannè.*

## EMULSION OF CASTOR OIL SEEDS.

- R. Castor oil seeds half an ounce  
Anise water four ounces  
Sugar two drachms  
Deprive the seeds of their exterior, triturate them to a uniform pulp with a little water, and the sugar, and finally add the remainder of the water gradually, and strain through a coarse cloth. *W. Procter.*

## ANTHELMINTIC EMULSION.

- R. Castor oil  
Mucilage of gum Arabic, each, one ounce  
Syrup of Corsica moss  
Water of semen contra chamomile, each, two fl. ounces



Emulsion of sweet  
almonds eight fl. ounces  
Mix, and make an emulsion. As an an-  
thelmintic. *Beral.*

## CASTOR OIL CLYSTER.

R. Castor oil one ounce and a-half  
Yolk of egg two  
Infusion of chamomile six fl. ounces  
Mix. *Radius.*

## MIXTURE OF CASTOR OIL AND ETHER.

R. Castor oil one ounce  
Sulphuric ether two drachms  
Mix. A spoonful every two hours, ad-  
vised as an anthelmintic, in cases of tape-  
worm. *Radius.*

## FIXATURE.

R. Castor oil two ounces  
Spermaceæ one drachm  
Arnotto half an ounce  
Oil of bergamot one drachm  
Otto of roses five drops  
Mix; melt by a moderate heat, and strain.  
To stiffen, and keep hair in form. *Redwood.*

## OLEUM TEREBINTHINÆ.

## OIL OF TURPENTINE.

This is usually known as *Spirits of turpentine*, and is the volatile oil obtained from the turpentine afforded by several species of *Pinus*. It is limpid, colorless, of a strong, penetrating, peculiar odor, and of a warm, pungent, somewhat bitterish taste. It is stimulant, diuretic, anthelmintic, and cathartic, and externally rubefacient.

## PURIFIED OIL OF TURPENTINE.

R. Oil of turpentine eight parts  
Alcohol one part  
Agitate together, and pour off the spirit,  
and repeat the process several times. *Nimmo.*

## LOTION FOR CHILBLAINS.

R. Oil of turpentine four parts  
Sulphuric acid one part  
Olive oil ten parts  
Mix. To be applied to the affected part,  
night and morning. *Gassicourt.*

## OIL OF TURPENTINE AND HONEY.

R. Oil of turpentine two fl. drachms  
Honey one fl. ounce  
Mix. A teaspoonful night and morning,  
in warm tea. In sciatica. *Ellis.*

## LOTION FOR TOOTHACHE.

X R. Oil of turpen-  
tine one fl. drachm and a-half  
Oil of cloves  
Oil of cajeput, each, half a fl. drachm

Balsam of Peru two drachms  
Opium two scruples  
To be well rubbed together. As an ap-  
plication to the face, in cases of toothache. *Beasley.*

## TURPENTINE MIXTURE.

R. Oil of turpen-  
tine one hundred and twenty drops  
Powdered gum Arabic  
sugar, each, two drachms  
Laudanum sixty drops  
Compound spirits of  
lavender two fl. drachms  
Mint water five fl. ounces  
Mix. A tablespoonful every two hours,  
in low forms of fever, &c. *Ellis.*

## VERMIFUGE EMULSION.

R. Oil of turpentine six fl. drachms  
Gum Arabic two drachms  
Chamomile water six fl. ounces  
Sulphuric ether two fl. drachms  
Mix. Two spoonfuls, night and morning,  
in cases of tapeworm. *Radius.*

## TURPENTINE MIXTURE.

R. Oil of turpentine one fl. ounce  
Powdered gum Arabic  
sugar, each, two drachms  
Mint water four fl. ounces  
Mix. A tablespoonful every two hours  
till it operates. An active purgative. *Frank.*

R. Oil of turpentine three fl. drachms  
Yolk of egg two  
Syrup of mint two fl. ounces  
orange flowers  
ether, each, one fl. ounce  
Tincture of cinnamon half a fl. drachm  
Mix. A spoonful, three times a-day, in  
neuralgia and rheumatism. *Foy.*

R. Oil of turpentine one fl. ounce  
Yolk of egg one  
Triturate together, and add, gradually,  
Emulsion of almonds four fl. ounces  
Syrup of orange two fl. ounces  
Compound spirit of  
lavender four fl. drachms  
Oil of cinnamon four drops  
One fl. ounce, three times a-day. Advised  
as a purge in Iritis. *Carmichael.*

## SPIRIT OF TURPENTINE MIXTURE.

R. Honey  
Spirit of turpentine  
Ammoniated tincture of  
guaiacum, each, two drachms  
Oil of cloves  
lemon, each, three drops



Mix. A teaspoonful, twice or three times a-day, in sciatica and lumbago. *Copland.*

## TURPENTINE CLYSTER.

R. Oil of turpentine one fl. ounce and a-half  
Yolk of egg one  
Tepid infusion of flaxseed one pint  
Mix. *Ellis.*

R. Oil of turpentine one fl. ounce  
Yolk of egg sufficient  
Decoction of barley nineteen fl. ounces  
*Lond. Ph.*

## TURPENTINE AND ETHER.

R. Oil of turpentine  
Sulphuric ether equal parts  
Mix. Dose, twenty to forty drops, in honey or syrup, in biliary calculi, and as an external application in rheumatism.  
*Cottureau.*

## TURPENTINE LINIMENT.

R. Oil of turpentine  
olives, each, two fl. ounces  
Tincture of camphor one fl. ounce  
Water of ammonia one fl. drachm  
Mix. As an external rubefacient.  
*Ellis.*

R. Soft soap two ounces  
Camphor one ounce  
Oil of turpentine sixteen fl. ounces  
Mix. A powerful rubefacient. *Lond. Ph.*

R. Oil of turpentine half a pint  
Resin cerate one pound  
Melt the cerate, and add the turpentine.  
*U. S. Ph.*

This is known as *Kentish's ointment*, and is much used as an application to burns and scalds.

## ACETIC TURPENTINE LINIMENT.

R. Oil of turpentine three fl. ounces  
Acetic acid five fl. drachms  
Rose water two and a-half fl. ounces  
Essence of lemon four fl. scruples  
Yolk of egg one  
Mix. As an external embrocation and liniment in phthisis.  
*Stokes.*

## SULPHURIC TURPENTINE LINIMENT.

R. Oil of turpentine three fl. ounces  
Sulphuric acid one fl. drachm  
Olive oil three fl. ounces  
Mix. *Brodie.*

## OPIATED TURPENTINE LINIMENT.

R. Oil of turpentine one fl. ounce  
chamomile two fl. ounces  
Laudanum one fl. drachm  
Mix. As a lotion in neuralgia.  
*Recamier.*

## STARKEY'S SOAP.

R. Dry carbonate of potassa  
Oil of turpentine  
Turpentine equal parts  
Mix the potassa with the oil, then add the resin, and triturate till the mixture is of the consistence of honey. Used in dropsy, and in gonorrhœa. Dose, eight to ten grains.  
*Guibourt.*

## DIURETIC WINE.

R. Oil of turpentine two fl. drachms  
Lemon juice one fl. ounce  
Wine four fl. ounces  
Mix. For a single dose. *Pierquin.*

## OLEUM TIGLII.

## CROTON OIL.

This oil is procured, for the most part, from the seeds of the *Croton tiglium*, but also from those of two or three other species. They are all natives of India, and the adjoining parts of Asia. The *C. tiglium* is a moderate sized shrub, bearing a somewhat triangular nut, containing three ovoid seeds, of a reddish brown color, having an oleaginous kernel, which affords, on pressure, the oil in question.

*Sex. Syst.* Monœc. monadelph. *Nat. Syst.* Euphorbiaceæ.

*Linn. Sp. Pl.* 1426. *Griffith, Med. Bot.* 597.

The oil, when pure and fresh, is nearly colorless, or yellowish; but when kept for some time, becomes of a reddish-brown or orange color. It has a faint, but peculiar smell, and an acrid and hot taste. It is a powerful hydragogue purgative, and has been much used in dropsy, apoplexy, and visceral obstructions; when applied externally, it causes irritation and inflammation of the skin, followed by a pustular eruption; and has been successfully employed in rheumatism, neuralgia, and bronchial and pulmonary affections. Dose for an adult, one to two drops.

## BOLUS OF CROTON OIL.

R. Croton oil one drop  
Powdered gum Arabic half a drachm  
Syrup of orange flowers sufficient  
Mix, and make four boluses. Two to four to be taken in the morning. *Foy.*

## PILLS OF CROTON OIL.

R. Croton oil six drops  
Soap half a drachm  
Oil of caraway eight drops  
Powdered liquorice root sufficient  
Mix, and make twelve pills. Dose, one or more. *Reece.*

R. Croton oil six drops  
Pill of aloes and myrrh one drachm and a-half  
Soap one scruple  
Powdered liquorice root sufficient



Mix, and make thirty pills. Dose, two to three, or more. *Copland.*

R. Croton oil one drop  
Crumb of bread sufficient

Mix, and make four pills. One, every hour, until they operate. *Ellis.*

R. Croton oil two drops  
Soap two grains  
Gum Arabic sufficient

Mix, and make four pills. *Foy.*

#### COMPOUND CROTON OIL PILLS.

R. Powdered scammony  
Powdered aloes, each, sixty-four parts  
Croton oil three parts  
Alcohol four parts

Dissolve the oil in the alcohol; add the solution, gradually, to a mixture of the powders, and beat into a mass. Divide into pills of five grains. One to three, for children of eight to fourteen years of age; three to five, for adults. *Beral.*

#### PILLS OF CROTON OIL AND QUININE.

R. Croton oil soap three grains  
Sulphate of quinine one drachm  
Extract of dandelion sufficient

Mix, and make twenty pills. One for a dose. *Caventou.*

#### PILLS OF CROTON OIL AND BLUE MASS.

R. Croton oil soap three grains  
Extract of henbane  
Blue mass, each, twenty-four grains  
Oil of pimento twelve minims

Mix, and make twelve pills. Two at bedtime. *Neligan.*

#### LOZENGES OF CROTON OIL.

R. Vanilla chocolate two drachms  
Sugar one drachm  
Starch one scruple  
Croton oil five drops

Mix, and make thirty lozenges. *Soubéiran.*

#### SOAP OF CROTON OIL.

R. Croton oil two parts  
Solution of caustic soda one part  
Mix; put into paper moulds; in a few days, slice, and keep in well-stopped bottles. Dose, one to three grains, in pills. *Foy.*

#### TINCTURE OF CROTON OIL.

R. Croton oil sixteen drops  
Alcohol one ounce  
Macerate for six or eight days, and filter. Dose, fifteen to twenty-five drops. *Soubéiran.*

R. Croton oil four drops

Tincture of myrrh one fl. ounce  
Mix, digest, and filter. Dose, one to two fl. drachms. *Bateman.*

R. Croton oil eight drops  
Alcohol one fl. ounce  
Mix, digest, and filter. Dose, half to one fl. drachm. *Nimmo.*

#### EMULSION OF CROTON OIL.

R. Croton oil three drops  
Almond oil half a fl. ounce  
Powdered gum Arabic two drachms  
Triturate well, and gradually add

Syrup of orange flowers one fl. ounce  
Chamomile water five fl. ounces  
A tablespoonful every two hours, till it operates. *Phœbus.*

R. Croton oil one drop  
Yolk of egg two drachms  
Orange-flower water  
Mint water, each, one ounce  
Make an emulsion. *Foy.*

#### MIXTURE OF CROTON OIL.

R. Croton oil one to two drops  
Mucilage of gum Arabic  
Distilled water, each, one fl. ounce  
Mix. A teaspoonful every two hours, until it operates. *Ellis.*

R. Croton oil two drops  
White sugar two drachms  
Gum Arabic half a drachm  
Tincture of cardamom half fl. drachm  
Distilled water one fl. ounce

Mix. Dose, two dessertspoonfuls every three or four hours. As it is agreeable to the taste, it is suited for children, but in smaller doses. *Beral.*

R. Tincture of croton oil twenty-five drops  
Mucilage of gum Arabic one drachm  
Water one ounce  
Mix. In the morning, fasting. *Foy.*

#### SAPONACEOUS SOLUTION OF CROTON OIL.

R. Croton oil eight drops  
Potassa six grains  
Distilled water two fl. drachms  
Mix. From three to six drops may be given for a dose. *Ellis.*

#### LINIMENT OF CROTON OIL.

R. Croton oil one part  
Olive oil five parts  
Mix. *Pereira.*

R. Croton oil one fl. drachm  
Oil of turpentine one fl. ounce  
Mix. *Corrigan.*

R. Croton oil four drops  
Carbonate of soda ten grains



Spirit of mint half an ounce  
In friction, in rheumatism. *Foy.*

R. Croton oil  
Solution of potassa, each, half fl. ounce  
Mix, and agitate.

R. The above solution thirty minims  
Rose water one fl. ounce

Mix. To be used twice a-day, till pustules  
appear. *J. Allen.*

#### EMBROCACTION OF CROTON OIL.

R. Croton oil twenty minims  
Tartar emetic one scruple  
Solution of potassa one fl. drachm  
Water eight fl. drachms

Mix. To keep up a mild eruption on the  
skin. *Morris.*

#### CERATE OF CROTON OIL.

R. Lard two and a-half parts  
Wax half a part

Melt together, and when nearly cold, mix  
with them

Croton oil one part  
*Caventou.*

R. Soap cerate four parts

Melt, and when semifluid, add  
Croton oil one part  
*Beasley.*

#### PLASTER OF CROTON OIL.

R. Lead plaster four parts  
Melt, and when nearly cold, add

Croton oil one part  
Spread on linen for an adhesive and irri-  
tating plaster *Bouchardat.*

R. Lead plaster eight parts  
Melt, and when nearly cold, add

Croton oil twenty parts  
To be spread as above. A very active  
counter irritant. *Bouchardat.*

#### OINTMENT OF CROTON OIL.

R. Croton oil ten minims  
Lard half an ounce

Mix. *Ainslie.*

### OPIUM.

#### OPIUM.

Opium is the inspissated juice of the unripe capsules of the *Papaver somniferum*, and presents many varieties, as the Turkey or Smyrna, the East Indian or Bengal, the Egyptian, &c.; of which the first is the best, and affords the largest proportion of morphia.

Opium contains various peculiar principles, several of which are officinal, and are treated of under their respective titles. Turkey opium is in flattish, rounded masses, of half a pound to two pounds in weight, covered externally with

the seed vessels of some species of dock. The texture is soft; the color is pale brown; the odor is strong and narcotic, and the taste bitter and acrid.

Opium is stimulant, in small and repeated doses, narcotic in large; and also antispasmodic, diaphoretic, sedative, and anodyne. It is used to fulfil a variety of indications; to procure sleep, to lull pain, to check morbid discharges, to alleviate cough, &c. The medium dose is one grain, but in spasm, &c., it is given in much larger doses.

#### EXTRACT OF OPIUM. (AQUEOUS.)

R. Opium one pound  
Water five pints

Cut the opium into small fragments, macerate it for twenty-four hours in a pint of water, break the pieces down with the hand, express; add another pint of water to the residuum, macerate for twenty-four hours, and again express; repeat this process till all the water has been used. Filter the several infusions, unite them, and evaporate on a water-bath to due consistence.

*Edin. Ph.*

#### EXTRACT OF OPIUM. (AQUEO-ALCOHOLIC.)

R. Opium one part  
Alcohol four parts

Cut the opium in small pieces, and digest it in the alcohol for two days, at a gentle heat, express, and treat the residue with four parts of warm water; express, unite the solutions, and evaporate to due consistence. *Taddei.*

#### EXTRACT OF OPIUM. (ACETOUS.)

R. Opium one ounce  
Distilled vinegar two pints

Cut the opium into small pieces, digest in the vinegar for two days, on a sand-bath, stirring from time to time, decant, filter, and evaporate to due consistence.

*Soubeyran.*

#### EXTRACT OF OPIUM. (ALCOHOLIC.)

R. Opium at will  
Alcohol sufficient

Reduce the opium to small pieces, digest in the alcohol in a closed vessel, by a gentle heat, often stirring, filter, and distil off the alcohol, till the residue is of due consistence. *Antwerp Ph.*

#### EXTRACT OF OPIUM. (VINOUS.)

R. Opium one part  
White wine four parts

Reduce the opium to small pieces, macerate it in the wine for twenty-four hours, occasionally stirring; express. Macerate the residue in two more parts of wine, and



express; unite the solutions, and evaporate to proper consistence. *Par. Cod.*

#### EXTRACT OF OPIUM. (DENARCOTIZED.)

R. Aqueous extract of opium at will  
Rub it with a little water, put it in a flask, add sulphuric ether, agitate, and decant; repeat the process with other portions of ether, as long as anything is taken up, and evaporate the residuum to a pilular consistence. *Robiquet.*

R. Aqueous extract of opium four parts  
Resin one part

Beat together, and add

Boiling water sixteen parts  
Boil till reduced one half, add as much cold water as has been boiled away, filter, and evaporate. *Limousin—Lamothe.*

#### EXTRACT OF OPIUM. (BY FERMENTATION.)

R. Opium one part  
Water eight parts

Mix, and add

Yeast sufficient

Let ferment for a week, at a temperature of 68° to 70°; then dilute with water, filter, boil till all vinous odor is dissipated, and evaporate to proper consistence. *Deyeux.*

R. Opium two ounces

Dissolve in a mixture of

Quince juice  
Water, each, sixty ounces

Digest for some days, with a gentle heat, strain, and add

White sugar four ounces

Let ferment for a month, stirring from time to time, then strain, and evaporate. *Langolst.*

#### EXTRACT OF OPIUM. (ROASTED.)

R. Powdered opium one part  
Heat it on a flat dish, over a moderate fire, constantly stirring, as long as fumes are given off. Treat it twice with six times its weight of cold water, filter, and evaporate. *Guibourt.*

The last three forms of Extract, whilst containing the full proportion of morphine, are deprived of the irritating and virose principles of opium.

#### COMPOUND POWDER OF OPIUM AND CHALK.

R. Powdered opium six grains  
cinnamon one drachm  
long pepper eight grains

Prepared chalk one drachm and a-half

Mix, and divide into twelve powders. One, three or four times a-day. In diarrhoea. *Ellis.*

#### POWDER OF ROASTED OPIUM.

R. Powdered opium at will  
Put it in a flat dish; moisten it with white wine, expose to a moderate heat, constantly stirring till it is perfectly dry, repeat the operation, till the opium is one-half reduced, moisten with vinegar, and rub into paste, dry, and pulverize. As an astringent in hemorrhages, and mucous discharges, in doses of three or four grains, with some bitter extract. *Giordano.*

#### POWDER OF OPIUM, CAMPHOR, &C.

R. Powdered opium ten grains  
camphor two scruples  
Carbonate of ammonia four scruples  
Starch fifteen grains  
Mix, and make eight powders. One, every hour or two, as antispasmodic. *Swedjaur.*

#### POWDER OF OPIUM AND NITRE.

R. Powdered opium four grains  
nitre

Sugar of

milk, each, one drachm and a-half  
Mix, and make six powders. As an antispasmodic. *Phabus.*

#### POWDER OF OPIUM AND SULPHUR.

R. Powdered opium  
camphor, each, two grains  
sulphur  
sugar, each, half a drachm  
Mix, and make four powders. One, every three hours, in lead colic. *Hildebrand.*

#### POWDER OF OPIUM AND MUSK.

R. Powdered opium two grains  
Musk five grains  
Magnesia four grains  
Sugar of milk ten grains  
Mix. To be taken every two to four hours, in delirium tremens. *Vogt.*

#### POWDER OF OPIUM AND ANTIMONY.

R. Dover's powder  
James' powder, each, four grains  
Mix. To be taken every four hours, in obstinate rheumatic pains. *Brera.*

#### PILLS OF OPIUM.

R. Powdered opium one drachm  
Soap twelve grains  
Beat into a mass, and divide into sixty pills. *U. S. Ph.*

#### PILLS OF OPIUM, HENBANE, AND HEMLOCK.

R. Powdered opium four grains  
Extract of henbane  
hemlock, each, fifteen grains



Mix, and divide into ten pills. One at night, where an anodyne is required.

*Ellis.*

PILLS OF OPIUM AND SULPHURET OF ANTIMONY.

R. Extract of opium ten grains  
Precip. sulph. antimony twelve grains  
Nitrate of potassa twenty-four grains  
Syrup sufficient

Mix, and make six pills. One at bedtime, to cause perspiration, and to ease pain in rheumatism.

*Recamier.*

PILLS OF OPIUM AND FOXGLOVE.

R. Powdered opium  
foxglove, each, six grains  
Conserve of roses sufficient

Mix, and make twelve pills. One, every four hours, in asthma, &c.

*Ellis.*

AROMATIC PILLS OF OPIUM.

R. Extract of opium  
Saffron  
Powdered cinnamon  
nutmeg  
cardamom, each, one drachm  
Syrup of orange flowers sufficient

Mix, and make pills of three grains. Known as Oriental pills, and considered to be aphrodisiac. One to three, at bedtime.

*Cadet.*

PILLS OF OPIUM AND CAMPHOR.

R. Extract of opium three grains  
Camphor six grains  
Syrup sufficient

Mix, and make six pills. One to three a-day, as an anodyne, and antispasmodic.

*Foy.*

PILLS OF OPIUM AND BUTTER OF CACAO.

R. Butter of cacao  
Powdered gum  
Arabic, each, forty-eight grains  
Extract of opium twelve grains  
Syrup of ipecacuanha sufficient

Mix, and make pills of five grains. One in the evening, as an anodyne and expectorant.

*Petit.*

PILLS OF OPIUM AND MUSK.

R. Extract of opium twelve grains  
valerian

Musk, each, twenty-four grains  
Mix, and make sixteen pills. One, then two, then three a-day, in hysteria.

*Foy.*

PILLS OF OPIUM AND SULPHATE OF ZINC.

R. Extract of opium one grain  
Sulphate of zinc four grains  
Syrup of gum sufficient

Mix, and make four pills. Two a-day, in

painful mucous discharges from the urethra or vagina.

*Foy.*

PILLS OF OPIUM, HEMLOCK, CALOMEL.

R. Extract of opium eighteen grains  
hemlock one drachm  
Calomel thirty-six grains  
Syrup of mallows sufficient

Mix, and make thirty-six pills. Two to six a-day, as alterative and sedative, in organic affections.

*Foy.*

R. Opium four grains  
Calomel six grains  
Tartar emetic one grain  
Extract of hemlock one scruple

Mix, and make eight pills; two to be taken at bedtime, in rheumatic pains.

*Brande.*

PILLS OF OPIUM AND ACETATE OF LEAD.

R. Extract of opium one grain  
Acetate of lead four grains  
Powdered henbane eight grains

Mix, and make eight pills. One, morning and evening, in epilepsy.

*Recamier.*

R. Acetate of lead twenty-four grains  
Powdered opium three grains  
Syrup sufficient

Mix, and make twelve pills. One every three hours, in hemorrhages, dysentery, and cholera.

*Chapman.*

PILLS OF OPIUM AND ACETATE OF MERCURY.

R. Extract of opium  
Acetate of mercury  
Camphor, each, twelve grains  
Syrup of poppies sufficient

Mix, and make thirty pills. One, morning and evening, in syphilis.

*Carmichael.*

PILLS OF OPIUM, NITRATE OF SILVER, &c.

R. Extract of opium seventy-two grains  
Nitrate of silver six grains  
Musk forty-eight grains  
Camphor ninety-six grains

Mix, and make ninety-six pills. One, morning and evening, gradually increasing the dose, in epilepsy, paralysis, &c.

*Foy.*

PILLS OF OPIUM, CASTOR, &c.

R. Opium half a grain  
Castor six and a-half grains  
Powdered digitalis one grain  
Syrup sufficient

Make two pills. One to be taken twice or thrice a-day. In spasmodic asthma and dyspnoea.

*A. T. Thomson.*

PILLS OF OPIUM AND LIQUORICE.

R. Powdered opium ten grains



Extract of liquorice one drachm  
Mix, and make eighty pills. One, occasionally, as an expectorant. *Wurt. Ph.*

## CONFECTION OF OPIUM.

R. Powdered opium four drachms and a-half  
Aromatic powder six ounces  
Clarified honey fourteen ounces  
Rub the opium with the aromatic powder, add the honey, and beat together till thoroughly mixed. *U. S. Ph.*

As a stimulant narcotic, in atonic gout, flatulent colic, &c. It contains one grain of opium in thirty-six.

R. Powdered catechu four ounces  
kino three ounces  
nutmeg  
cinnamon, each, one ounce  
Opium, dissolved in wine, one drachm and a-half  
Syrup of red roses twenty-seven ounces  
Mix. Each drachm contains rather less than half a grain of opium. *Soubairan.*

## ANTI-ODONTALGIC MASS.

R. Opium two grains  
White wax two drachms  
Mastic one drachm  
Oil of almonds three drachms  
cloves twelve drops  
cochineal eight grains  
Mix well into a uniform mass. To fill carious teeth. *Clarus.*

R. Opium five grains  
Oil of cloves three drops  
Extract of henbane five grains  
belladonna ten grains  
Powdered pellitory sufficient  
Mix, and make a consistent mass. Used as above. *Rust.*

## BALSAM FOR THE TOOTHACHE.

R. Opium one scruple  
Oil of turpentine one drachm and a-half  
Oil of cloves  
cajuput, each, half a drachm  
Balsam of Peru two drachms  
Mix. *Beasley.*

R. Opium  
Camphor, each, two scruples  
Dissolve in a little alcohol, and add  
Oil of cloves one drachm  
guaiacum two drachms  
nutmeg six drachms  
Mix. *Van Mons.*

## TOOTHACHE DROPS.

R. Opium  
Camphor, each, ten grains

Alcohol sufficient  
Oil of cloves  
cajuput, each, one drachm  
Mix. *Copland.*

## ANTIDYSENTERIC OPIATE.

R. Purified opium four grains  
Ipecacuanha half a drachm  
Tormentilla one drachm  
Syrup of whortleberries  
Conserve of red roses, each, six drachms  
Mix. Dose, one drachm, every hour. *Quarin.*

## PLASTER OF OPIUM.

R. Powdered opium two ounces  
Burgundy pitch three ounces  
Lead plaster one pound  
Boiling water four fl. ounces  
Melt together the pitch and plaster; then add the opium mixed with the water, and boil to the proper consistence. *U. S. Ph.*  
As an application in rheumatic and other pains.

## PLASTER OF OPIUM AND CAMPHOR.

R. Powdered opium  
camphor  
soap, each, one drachm  
Laudanum sufficient  
to make a plaster *Ellis.*

R. Opium  
Camphor, each, half a drachm  
Lead plaster sufficient  
Mix. For local pains. *Paris.*

R. Opium one drachm  
Simple plaster two ounces  
Galbanum one ounce  
Oil of caraway two drachms  
Melt the last three ingredients, and add the opium. As an application to the abdomen, in flatulent colic, diarrhoea, and dysentery. *Swediaur.*

## LINIMENT OF OPIUM.

R. Soap six ounces  
Opium one ounce and a-half  
Camphor three ounces  
Oil of rosemary six fl. drachms  
Rectified spirit two pints  
Macerate the soap and opium in the spirit for three days; filter; add the oil and camphor, and agitate briskly. *Ed. Ph.*

As an embrocation in rheumatic pains, sprains, &c.

## LINIMENT OF OPIUM AND OIL OF CHAMOMILE.

R. Opium  
Oil of chamomile, each, half a drachm  
almonds two drachms



Mix. As a friction around the eyes, in spasm of the eyelids. *Weller.*

ANTI-OTITIC MIXTURE.

R. Opium four grains  
Saffron ten grains  
Myrrh half a drachm  
Juice of mallows half an ounce  
Oil of almonds two ounces  
Triturate well together, and strain. As an injection into the ear, in pain in that organ. *Pierquin.*

ANODYNE OINTMENT.

R. Opium  
Saffron, each, one scruple  
Yolk of egg one  
Poplar ointment  
Acetate of lead ointment, each, one ounce

Mix. As an application to painful hemorrhoids. *Fulda Ph.*

R. Syrup of opium one ounce  
Lard three ounces  
Essence of roses four drops

Mix. For chapped lips. *Pierquin.*

R. Opium half a drachm  
Extract of hemlock one drachm  
Basilicon ointment half an ounce

Mix. As an application to gangrenous ulcers. *Carus.*

R. Opium ten grains  
Alum fifteen grains  
Lard half an ounce

Mix. In the treatment of ulcerated, atonic buboes. *Simon.*

OINTMENT OF OPIUM AND TAR.

R. Powdered opium two drachms  
Tar ointment one ounce

Mix. As an application to hemorrhoids. *Ellis.*

CERATE OF OPIUM.

R. Opium half a drachm  
Yolk of egg one

Mix well, and add Simple cerate one ounce

Triturate well together. *Lagneau.*

SYRUP OF OPIUM.

R. Extract of opium one part  
Water sixteen parts  
Syrup five hundred parts  
Dissolve the opium in the water, filter, add it to the syrup, heated to boiling, strain. *Par. Cod.*

SUCCINATED SYRUP OF OPIUM.

R. Syrup of opium one ounce  
Spirit of amber two grains  
Mix. *Soubeiran.*

ANODYNE COLLYRIUM.

R. Extract of opium ten grains  
Camphor six grains  
Boiling water twelve fl. ounces  
Rub the opium and camphor together, and add the water, and strain. *Ellis.*

R. Extract of opium four grains  
Rose water four ounces  
Dissolve, and strain. *Par. Cod.*

OPIUM FOMENTATION.

R. Extract of opium two drachms  
Boiling water one pint  
Dissolve, and strain. As a fomentation in pruriginous affections. *Radius.*

INJECTION OF OPIUM.

R. Opium twelve grains  
Solution of subacetate of lead twelve drops  
Water nine ounces

Mix. As an injection in gonorrhœa. *Girtanner.*

R. Extract of opium six grains  
belladonna

one drachm and a-half  
Decoction of wild lettuce one pound  
Dissolve. As an injection in neuralgia, and hemorrhage of the urethra and vagina. *Foy.*

R. Extract of opium one and a-half grain  
Distilled water five fl. drachms  
Dissolve. As an injection in coryza, and other inflammations of the nasal, mucous membrane. *Deschamps.*

He directs one nostril to be closed by the finger, and the liquid to be drawn up into the other, &c.

CLYSTER OF OPIUM.

R. Powdered opium two grains  
Mucilage of gum Arabic half fl. ounce  
Tepid milk two fl. ounces

Mix. *Ellis.*

R. Flaxseed one ounce  
Boiling water six ounces  
Infuse for an hour, strain, and add  
Extract of opium two grains  
Dissolve. *Spielman.*

SUPPOSITORY OF OPIUM.

R. Powdered opium two grains  
Soap four grains  
Mix. *Ellis.*

LINCTUS WITH OPIUM.

R. Extract of opium one grain  
Peruvian bark four grains  
Camphor six grains  
Sugar one drachm  
White linctus four ounces



Mix. In bronchitis, when the cough is violent. *Foy.*

#### MIXTURE OF OPIUM AND SYRUP OF POPPIES.

R. Opium two to three grains  
Syrup of poppies one ounce  
Chamomile water six ounces

Mix. A spoonful every half hour, in dysentery. *Swediaur.*

#### MIXTURE OF OPIUM AND LIME WATER.

R. Extract of opium one grain  
Lime water

Oil of almonds, each, three fl. drachms

Mix. For the treatment of sore nipples, to be applied on dossils of lint.

*Sibergundi.*

#### MIXTURE OF OPIUM AND CINNAMON WATER.

R. Powdered opium ten grains  
Sugar one drachm  
Cinnamon water six fl. ounces

Mix. A tablespoonful every two hours, in tetanus and colica pictonum. *Ellis.*

#### WATER OF OPIUM.

R. Opium, in small pieces one pound  
Water six pounds

Macerate for forty-eight hours, and distil one pound. It is given in the dose of one to four drachms, in syrup of sugar. *Foy.*

#### MURIATE OF OPIUM.

R. Powdered opium one ounce  
Muriatic acid one ounce  
Distilled water twenty ounces

Mix, and shake the mixture frequently, for fourteen days, strain, and filter. Dose, from twenty to forty drops. Said not to cause headache. *Nichol.*

#### VINEGAR OF OPIUM. BLACK DROP.

R. Powdered opium eight ounces  
Coarsely powdered  
nutmeg one ounce and a-half  
Saffron half an ounce  
Sugar twelve ounces  
Distilled vinegar sufficient

Digest the opium, nutmeg, and saffron, with a pint and a-half of distilled vinegar, on a sand-bath, with a gentle heat, for forty-eight hours, and strain. Digest the residue with an equal quantity of distilled vinegar, in the same manner, for twenty-four hours. Put the whole into a displacement apparatus, and return the filtered liquor as it passes, until it comes away clear. When the filtration has ceased, pour distilled vinegar gradually on the residue in the filter, until the whole quan-

tity of filtered liquor amounts to three pints. Lastly, add the sugar, and, by means of a water-bath, evaporate to three pints and four fl. ounces. Diluted acetic acid may be substituted for the distilled vinegar. *U. S. Ph.*

Dose, about six minims, which are nearly equivalent to one grain of opium.

#### LANCASTER BLACK DROP.

R. Opium half a pound  
Verjuice three pints  
Bruised nutmeg one ounce and a-half  
Saffron half an ounce

Boil to a proper consistence; add two spoonfuls of yeast, and let stand in a warm place for six or eight weeks, and then in the open air till of the consistence of syrup, then decant, filter, and bottle; adding a little sugar to each bottle. Dose, six to ten drops. *Armstrong.*

#### HOULTON'S BLACK DROP.

R. Opium two ounces and a-half  
Diluted acetic acid thirty-two ounces  
Digest for six days with a gentle heat, and filter, evaporate to an extract; mace-

rate in  
Rectified spirit five fl. ounces  
Distilled water thirty-five ounces  
for eight days, and filter. *Beasley.*

About the strength of laudanum.

#### GUY'S HOSPITAL BLACK DROP.

R. Powdered opium eight ounces  
Juice of crab apples two pints  
boil gently for half an hour, decant, and boil residue with one pint more of the juice, for a quarter of an hour; express, and strain; mix the two liquors, and add  
Bruised nutmeg one ounce  
Saffron half an ounce  
Yeast half fl. ounce.

Ferment for some days; macerate for fourteen days; filter, and evaporate by a water-bath to consistence of thin syrup. Dose, two to ten minims. *Beasley.*

#### ROUSSEAU'S BLACK DROP.

R. Opium four ounces  
Honey twelve ounces  
Hot water five pounds  
Yeast two drachms

Dissolve the opium and honey separately in the hot water, mix, and add the yeast; keep at about 80° F. for a month; express; filter; distil off sixteen ounces, and evaporate residue to ten ounces; add to it one and a half ounce of strong spirit; mix, and filter. Four drops are equivalent to one grain of opium. *Beasley.*



PORTER'S BLACK DROP.

R. Opium four ounces  
Citric acid two ounces  
Beat together in a mortar, and add  
Boiling distilled water one pint  
Triturate well together, let stand for  
twenty-four hours, and filter. Dose, from  
six to twenty-four drops. *Redwood.*

ACETATED TINCTURE OF OPIUM.

R. Opium two ounces  
Vinegar twelve fl. ounces  
Alcohol half a pint  
Rub the opium with the vinegar, then  
add the alcohol, macerate for fourteen  
days, express, and filter through paper.  
*U. S. Ph.*

Dose, ten minims or twenty drops, which are  
equivalent to a grain of opium.

MIXTURE WITH BLACK DROP.

R. Houlton's black drop ten drops  
Spirit of nitric ether half a drachm  
Distilled water one ounce  
Mix. To be taken at once. *Foy.*

SEDATIVE MIXTURE.

R. Opium two grains  
Distilled vinegar half an ounce  
Plantain water six ounces  
Syrup of white poppy one ounce  
Mix, and filter. In spoonful doses, in  
hæmoptysis with spasms. *Pierquin.*

WINE OF OPIUM.

R. Powdered opium two ounces  
Bruised cinnamon  
cloves, each, one drachm  
Wine one pint  
Macerate for fourteen days, agitating occa-  
sionally, express, and filter. *U. S. Ph.*

Dose, about thirteen minims or twenty-five  
drops, equivalent to a grain of opium.

R. Extract of opium two ounces  
Cinnamon water ten ounces  
Alcohol two ounces  
White wine four ounces  
Mix, and macerate for four days, and filter.  
*Brugnatelli.*

About the same strength as the last.

R. Acetous extract of opium six drachms  
Sherry wine ten ounces  
Brandy two ounces  
Mix; macerate for four days, and filter;  
each drachm contains four grains of opium.  
*Lalouette.*

LAUDANUM OF SYDENHAM.

R. Opium two ounces  
Saffron one ounce

Bruised cinnamon

cloves, each, one drachm  
Sherry wine one pint  
Mix, and macerate for fifteen days, and  
filter. *Par. Cod.*  
Twenty drops are equivalent to a grain of  
opium.

COLLYRIUM OF WINE OF OPIUM.

R. Decoction of flaxseed four ounces  
Saffron one drachm  
Wine of opium one drachm  
Macerate the saffron in the flaxseed decoc-  
tion, strain, and add the wine of opium.  
*Foy.*

R. Acetate of copper three grains  
Dissolve in  
Rose water eight ounces  
add,  
Wine of opium one drachm  
In chronic ophthalmia. *Foy.*

FOMENTATION OF WINE OF OPIUM.

R. Opium one ounce  
Wine two pints  
Boil down to one pint. As an anodyne  
application, in gouty and rheumatic pains,  
&c. *Pierquin.*

ODONTALGIC DROPS.

R. Wine of opium  
Hoffmann's anodyne  
Oil of peppermint equal parts  
Mix. In frictions on the cheek, and ap-  
plied to carious teeth, on cotton.  
*Dobberan.*

MIXTURE OF WINE OF OPIUM.

R. Wine of opium ten drops  
Cinnamon water one ounce  
Balm water two ounces  
Tincture of castor twenty drops  
Syrup of opium half an ounce  
Mix. In spoonful doses, in uterine colic.  
*Augustin.*

R. Cascarella  
Colombo, each, two drachms  
Boiling water sufficient  
to obtain seven ounces of strained infusion.  
Add to this

Wine of opium  
Sulphuric ether, each, twenty drops  
To be taken by degrees, in chronic diar-  
rhœa. *Brera.*

TINCTURE OF OPIUM. LAUDANUM.

R. Opium, powdered, two ounces and  
a-half  
Diluted alcohol two pints  
Macerate for fourteen days, and filter.  
*U. S. Ph.*



Dose, thirteen minims or twenty-five drops, equivalent to a grain of opium.

#### BATTLE'S SEDATIVE DROPS.

R. Hard extract of opium three ounces  
Boiling distilled water thirty ounces  
Dissolve, when cold filter, and add  
Rectified spirit six ounces  
and water sufficient to make up two pints.  
*Cooley.*

#### SMITH'S CONCENTRATED LAUDANUM.

R. Denarcotized opium four ounces  
Dissolve in alcohol, filter, evaporate to consistence of an extract, redissolve in water, and evaporate the filtered solution to twelve ounces; add  
Rectified spirit twenty-two drachms  
Distilled water sufficient  
to make up sixteen ounces. Dose, three to five drops.  
*Beasley.*

#### COMPOUND TINCTURE OF OPIUM.

R. Extract of liquorice  
Opium, each, half an ounce  
Carbonate of potassa one drachm  
Water three pints  
Boil down to one pint, filter, and evaporate to twelve ounces, then add  
Spirit of pimento five fl. ounces  
Powdered cochineal half a drachm  
Let rest for some time, and filter.  
*Med.-Chirurg. Rev.*

#### AMMONIATED TINCTURE OF OPIUM.

R. Benzoic acid  
Chopped saffron, each, six drachms  
Sliced opium half an ounce  
Oil of anise one drachm  
Spirit of ammonia two pints  
Digest for seven days, and filter. *Ed. Ph.*

Also called *Scotch Paregoric*. Eighty minims should contain one grain of opium.

#### CAMPHORATED TINCTURE OF OPIUM.

##### PAREGORIC.

R. Powdered opium  
Benzoic acid, each, one drachm  
Oil of anise one fl. drachm  
Clarified honey two ounces  
Camphor two scruples  
Diluted alcohol two pints  
Macerate for fourteen days, and filter.  
*U. S. Ph.*

Half a fl. ounce contains rather less than a grain of opium. Dose, for an infant, five to twenty drops; for an adult, one to two fl. drachms.

#### BATEMAN'S PECTORAL DROPS.

R. Diluted alcohol four gallons  
Rasped red saunders two ounces

Digest for twenty-four hours, filter, and add

Powdered opium  
catechu  
camphor, each, two ounces

Oil of anise four fl. drachms  
Digest for ten days. About as strong as camphorated tincture of opium, or two grains of opium to the fl. ounce.

*Phil. Coll. Pharm.*

#### GODFREY'S CORDIAL.

R. Tincture of opium one pint and a-half  
Sugar-house molasses sixteen pints  
Alcohol two pints  
Water twenty-six pints

Carbonate of  
potassa two ounces and a-half  
Oil of sassafras four fl. drachms

Dissolve the carbonate of potassa in the water; add the molasses; heat over a gentle fire, till they simmer; remove the scum; add the laudanum and oil of sassafras, previously mixed together.

*Phil. Coll. Pharm.*

Contains rather more than one grain of opium to the fl. ounce.

#### TINCTURE OF OPIUM AND SOAP.

R. Opium half an ounce  
Soap two ounces  
Alcohol sixteen ounces  
Digest for three days on a water-bath, filter, and dissolve in the liquor

Camphor six drachms  
Oil of rosemary one drachm

Principally used in frictions, in pains in the limbs, &c., but also given internally, in doses of thirty to fifty drops, in wine.

*Turin Ph.*

#### SUCCINATED TINCTURE OF OPIUM.

R. Opium forty grains  
Camphor  
Amber, each, half a drachm  
Alcohol six ounces

Digest for four days on a water-bath, and filter. As a friction in spasmodic attacks.  
*Dumas.*

#### SWEDIAUR'S TINCTURE OF OPIUM.

R. Extract of opium one drachm  
Distilled water four drachms  
Alcohol half a drachm  
Dissolve, and filter. Five drops are equivalent to a grain of opium. *Swediaur.*

#### TINCTURE OF OPIUM AND ASARABACCA.

R. Asarabacca root  
Sassafras, each, one ounce  
Aloes wood half an ounce



Opium three drachms  
 Carbonate of ammonia one drachm  
 Alcohol sixteen ounces  
 Digest for twenty days, and filter. Dose,  
 six to thirty drops. In hysteria and epi-  
 lepsy. *Cottureau.*

#### WARNER'S TINCTURE OF OPIUM.

R. Opium  
 Soap, each, six drachms  
 Nutmeg one drachm  
 Camphor four drachms  
 Saffron forty grains  
 Spirit of ammonia nine ounces  
 Digest for ten days, agitating occasionally,  
 then filter. *Van Mons.*

#### LETTSON'S ELIXIR.

R. Opium  
 Benzoic acid  
 Saffron, each, two drachms  
 Camphor four scruples  
 Oil of anise one drachm  
 Ipecacuanha  
 Balsam of tolu, each, half an ounce  
 Alcohol two pounds  
 Macerate for ten days, and filter. Dose,  
 five to twenty drops, in hooping-cough.  
*Augustin.*

#### CLYSTER OF LAUDANUM.

R. Tincture of opium half a drachm  
 Infusion of flax-  
 seed two to four fl. ounces  
 Mix. *Ellis.*  
 R. Decoction of starch four fl. ounces  
 Tincture of opium thirty minims  
 Mix. In obstinate vomiting, strangury  
 from blisters, painful affections of the  
 genito-urinary apparatus, tenesmus, &c.  
*Lond. Ph.*

#### CLYSTER OF LAUDANUM AND VALERIAN.

R. Tincture of opium five to six drops  
 Infusion of valerian three ounces  
 Mix, in spasms, in children. *Swedaur.*

#### OPIATE LINIMENT.

R. Olive oil two fl. ounces  
 Tincture of opium two fl. drachms  
 Solution of acetate of  
 lead half a fl. ounce  
 Mix. *Ellis.*  
 R. Sulphuric  
 ether two fl. drachms and a-half  
 Spirit of cam-  
 phor two fl. ounces and a-half  
 Laudanum half a fl. ounce  
 Mix; as an embrocation in flatulent colic.  
*Ainslie.*

R. Tincture of opium  
 Spirit of sulphuric ether  
 camphor, each, two fl. ounces  
 Mix. As an embrocation in rheumatism,  
 tooth, or earache, &c. *Ellis.*

#### LINIMENT OF LAUDANUM AND LIME WATER.

R. Tincture of  
 opium two drachms and a-half  
 Lime water one ounce  
 Oil of almonds two ounces  
 Mix. To be applied on lint; four times  
 a-day, on painful syphilitic pustules.  
*Phabus.*

#### LAUDANUM OINTMENT.

R. Tincture of opium one drachm  
 Spermaceti ointment two ounces  
 Mix. As an application, morning and  
 evening, to painful hemorrhoids. *Brera.*

#### BROWN COUGH MIXTURE.

R. Paragoric elixir one ounce  
 Antimonial wine half an ounce  
 Powdered gum Arabic  
 Extract of liquor-  
 ice, each, three drachms  
 Water six ounces  
 Mix well. A tablespoonful, every three  
 or four hours, in catarrh. *Dewees.*

R. Extract of liquorice  
 Powdered gum  
 Arabic, each, two drachms  
 Boiling water four fl. ounces  
 Dissolve, and add  
 Antimonial wine two fl. drachms  
 Tincture of opium twenty minims  
 Mix. A tablespoonful occasionally.  
*Wood.*

#### OPIATE LINCTUS.

R. Tincture of opium two fl. drachms  
 Diluted sulphuric  
 acid two fl. drachms and a-half  
 Molasses eight fl. ounces  
 Water three fl. ounces  
 Mix. A teaspoonful occasionally, to  
 quiet cough. *Beasley.*

#### ANODYNE DRAUGHT.

R. Camphor mixture nine drachms  
 Nitrate of potassa six grains  
 Comp. spirit of ether one drachm  
 Tincture of opium ten to twelve minims  
 Syrup of poppies two drachms  
 Mix. To be taken at bedtime. *Copland.*  
 R. Tincture of opium twelve minims  
 Water one fl. ounce  
 Pimento water three drachms



Syrup of poppies one drachm  
 Mix. *Beasley.*  
 R. Tincture of  
   opium fifteen to twenty-five drops  
   Syrup of poppies two fl. drachms  
   Spirit of cinnamon one fl. drachm  
   Distilled water one fl. ounce and a-half  
 Mix. *Ellis.*

#### MIXTURE OF LAUDANUM AND TARTAR EMETIC.

R. Laudanum one fl. drachm  
 Tartar emetic four grains  
 Camphor mixture eight fl. ounces  
 Mix. In delirium tremens, and the ad-  
 vanced stage of low fevers. Dose, half a  
 fl. ounce to one fl. ounce. *Graves.*

#### WISTAR'S COUGH LOZENGES.

R. Powdered gum Arabic  
   extract of liquorice  
   sugar, each, two drachms  
   opium six grains  
 Oil of anise four drops  
 Mix, and add  
   Distilled water sufficient  
 Make into mass, and divide into sixty  
 lozenges. One, three or four times a-day.  
*Ellis.*

### OPOPONAX.

#### OPOPONAX.

A gum resin obtained from the *Opoponax chi-  
 ronium*, a tall, parsnep-like plant, a native of the  
 warm countries of Europe and the Levant.

*Ser. Syst.* Pentand. digyn. *Nat. Syst.* Apiaceæ.  
*Decandolle.* iv. 170. *Griffith, Med. Bot.* 323.

It occurs in tears and irregular lumps, or frag-  
 ments, of a reddish-yellow color. Its odor is  
 strong, peculiar, and unpleasant, and its taste bit-  
 ter and acrid. It is seldom used in this country,  
 but was formerly much esteemed in a variety of  
 diseases. The dose is from ten to thirty grains.

#### TINCTURE OF OPOPONAX.

R. Opoponax one part  
 Alcohol five parts  
 Macerate for some days, and filter. *Beral.*

#### COMPOUND TINCTURE OF OPOPONAX.

R. Round birthwort  
 Long birthwort  
 Orris root, each, half an ounce  
 Opoponax  
 Sagapenum, each, two drachms  
 Guaiacum four scruples  
 Cloves two drachms  
 Camphor three drachms  
 Alcohol ten ounces  
 Macerate for twenty-four hours, and filter.  
 As an application to foul venereal ulcers.  
*Brera.*

#### EMULSION OF OPOPONAX.

R. Opoponax  
 Soap, each, one drachm  
 Yolk of egg two drachms  
 Syrup of worm-  
   wood one ounce and a-half  
 Fennel water three ounces  
 Make an emulsion. A teaspoonful every  
 hour, as a vermifuge. *Bories.*

### ORIGANUM.

#### ORIGANUM.

The article so designated in the U. S. Pharm.  
 is the herb of the *Origanum vulgare*, a native of  
 Europe, but extensively naturalized in the United  
 States.

*Ser. Syst.* Didynam. gymnos. *Nat. Syst.* La-  
 miaceæ.

*Linn. Sp. Pl.* 834. *Griffith, Med. Bot.* 511.

The dried herb has an aromatic, agreeable  
 odor, and a hot, pungent taste, depending on the  
 presence of a volatile oil. It is somewhat tonic  
 and stimulating, and has been used in the form  
 of infusion as a diaphoretic and emmenagogue.  
 The oil is employed in stimulating liniments.

### ORYZA.

#### RICE.

The seed or grain of *Oryza sativa*, an annual  
 plant, said to be a native of Ethiopia, but now  
 extensively cultivated in most warm countries.

*Ser. Syst.* Hexand. digyn. *Nat. Syst.* Grami-  
 naceæ.

*Linn. Sp. Pl.* 475. *Griffith, Med. Bot.* 660.

This grain as found in commerce is deprived  
 of its cuticle, is white, dry, hard, inodorous, and  
 of a farinaceous taste. It is very nutritive, and,  
 in the form of a decoction, emollient.

#### RICE WATER.

R. Rice, well washed, two ounces  
 Water two quarts  
 Boil for an hour and a half, then add sugar  
 and flavoring, as may be required. *Ellis.*

R. Rice five drachms  
 Water sufficient  
 to obtain a quart of decoction; add  
 Liquorice root three drachms  
 Let stand for some time, and strain, add-  
 ing sugar, &c., at will. *Cottureau.*

#### RICE GRUEL.

R. Ground rice one ounce  
 Cinnamon one drachm  
 Water one quart  
 Boil for forty minutes, adding the aro-  
 matic near the close. Strain, and sweeten.  
*Ellis.*

#### MUCILAGE OF RICE.

R. Rice one ounce



Macerate it for three hours in

Tepid water one quart  
Then boil slowly for an hour, and strain.

*A. T. Thomson.*

#### RICE JELLY.

R. Rice sufficient  
Macerate in as much water as will cover it, boil slowly, adding water as it evaporates, until the rice is reduced to a pap, sweeten and flavor, and pass through a fine sieve. On cooling, it becomes a moderately consistent jelly. A good diet in dyspepsia, &c.

### OVUM.

#### EGGS.

#### OIL OF EGGS.

R. Yolk of eggs at will  
Heat gently till the moisture is dissipated, introduce into a displacer, and exhaust by ether, distil the product in a water-bath, heat the residue, till the albuminous matter coagulates, then strain. *Par. Cod.*

Was, at one time, much esteemed as an application to hemorrhoids, chaps, and excoriations, &c.

#### EMULSION OF EGGS.

R. Yolk of eggs two  
Powdered sugar one ounce  
Boiling water fourteen ounces  
Triturate the eggs and sugar in a marble mortar, gradually adding the water.

*Beral.*

#### EMOLLIENT CLYSTER OF EGGS.

R. Yolk of eggs two  
Decoction of bran one pound  
Mix. *Pierquin.*

#### RESTORATIVE CLYSTER OF EGGS.

R. Yolk of egg one  
White wine two ounces  
Beef tea, without salt, eight ounces  
Mix. *Radius.*

#### MIXTURE WITH EGGS.

R. Yolk of egg one  
Cream six ounces

Cinnamon barley water six drachms  
Sugar one ounce

Mix. To be taken in teaspoonful doses, in the convalescence of children. *Foy.*

R. Yolk of egg one  
Water two pints

Make an emulsion, and add  
Common salt half a drachm

Mix. In mesenteric atrophy of children. *Hufeland.*

#### EGG AND BRANDY MIXTURE.

R. Brandy  
Cinnamon water, each, four fl. ounces  
Yolk of eggs two  
Sugar half an ounce  
Oil of cinnamon two minims  
Mix. *Lond. Ph.*

As a stimulant in the sinking stage of fevers.

#### EGG AND WINE MIXTURE.

R. Yolk of eggs two  
Oil of cinnamon twenty drops  
Mix, and add  
Madeira wine  
Cinnamon water, each, three fl. ounces  
Distilled water two fl. ounces  
Sugar two drachms  
Mix. Three or four spoonfuls for a dose. In convalescence from low fevers. *Ellis.*

#### LINIMENT OF EGGS.

R. Yolk of egg one  
Flaxseed oil two ounces  
Mix well. As an application to burns. *Radius.*

R. White of egg one  
Flaxseed oil three ounces  
Mix well. As the last. *Mysicht.*

#### CERATE OF EGGS.

R. Yolk of egg one part  
Simple cerate two parts  
Mix well. *Foy.*

R. Yolk of a hard-boiled egg one  
Yellow wax half an ounce  
Almond oil one ounce and a-half  
Melt the wax and oil together, and add the egg, triturating them thoroughly together. As an application to burns. *Soubairan.*



## P

## PAPAYER.

## POPPY HEADS.

The ripe capsules of *Papaver somniferum*. These capsules are of a more or less globular form, crowned by a radiated, persistent stigma; of a light brown color, a papery and brittle texture, inodorous, but of a slightly bitter taste. They are analogous in properties, but in a slight degree, to opium. They contain numerous, small, oleaginous seed, which are used in many preparations.

## DECOCTION OF POPPY HEADS.

R. Poppy heads, sliced, four ounces  
Water four pints  
Boil, and strain. *Lond. Ph.*

As a soothing and anodyne fomentation, in painful tumors and inflammations.

## SYRUP OF POPPY HEADS.

R. Poppy heads, bruised, fourteen ounces  
Water two gallons and a-half  
Macerate for twenty-four hours, boil down to a gallon, express, evaporate to two pints, and add

Sugar two pounds  
Dose, two fl. drachms to a fl. ounce, for an infant half a fl. drachm. *Dub. Ph.*

As a sedative and hypnotic.

R. Poppy heads sixteen ounces  
Diluted alcohol four pints  
Sugar thirty ounces

Deprive the heads of their seeds; bruise them thoroughly, macerate them in twice their weight of diluted alcohol, for two days, express powerfully, add the remainder of the alcohol, and after twenty-four hours, again express. Evaporate the liquids to one pint, strain, and add the sugar, and dissolve by the aid of a gentle heat.

*W. Procter.*

## EXTRACT OF POPPY HEADS.

R. Poppy heads, bruised, fifteen ounces  
Boiling water one gallon  
Macerate for twenty-four hours, boil down to four pints, strain, and evaporate over a vapor-bath, to proper consistence.

*Ed. Ph.*

The dose is from five to ten grains.

## PECTORAL SYRUP.

R. Dates two pounds  
Jujubes one pound  
Liquorice root half a pound

Mallow root

Maidenhair

Poppy heads, each, four ounces  
Water sixteen pints

Boil, strain, and add

Sugar eight pounds

Evaporate to the consistence of syrup.

Dose, one to two ounces. *Gassicourt.*

## SEDATIVE INJECTION.

R. Simple emulsion five fl. ounces  
Decoction of poppy heads one pint  
White of egg one drachm  
Mix. As an injection in acute gonorrhœa. *Gassicourt.*

## SEDATIVE MIXTURE.

R. Flaxseed two drachms  
Poppy head one  
Water sufficient  
to obtain eight ounces of infusion, add  
Yolk of egg one  
Mix well. In painful diarrhœa. *St. Marie.*

## PAREIRA.

## PAREIRA BRAVA.

This is the root of *Cissampelos pareira*, a climbing shrub, with large, cordate, villous leaves, a native of the West Indies and South America.

*Sex. Syst.* Monœc. monad. *Nat. Syst.* Menispermaceæ.

*Linn. Sp. Pl.* 1473. *Griffith, Med. Bot.* 106.

The root, which is the officinal portion, as found in the shops, is usually in large, crooked pieces, of a dark color, externally, and yellowish within; it has at first a sweetish, and somewhat aromatic taste, but leaves a bitterish, unpleasant impression; the odor is very slight. It is tonic and alterative, and acts specifically on the bladder, diminishing irritability, and diminishing mucous secretion. The dose, in substance, is from thirty grains to a drachm.

## INFUSION OF PAREIRA BRAVA.

R. Pareira brava six drachms  
Boiling water a pint  
Macerate for two hours, and strain. *Edin. Ph.*

Dose, one to two fl. ounces, in irritation and chronic inflammation of urinary passages.

## DECOCTION OF PAREIRA BRAVA.

R. Pareira brava four drachms  
Water one pint and a-half



Boil to a pint, and strain. Dose, from eight to twelve fl. ounces, during the day.

*Brodie.*

#### EXTRACT OF PAREIRA BRAVA.

R. Powdered pareira brava      one pound  
Water      sufficient

Mix the powder with half a pint of the water; introduce into a displacement apparatus, and pour on water, till the powder is exhausted. Heat the fluid to the boiling point; strain, and evaporate to proper consistence.

*Edin. Ph.*

Dose, ten grains to half a drachm.

#### TINCTURE OF PAREIRA BRAVA.

R. Pareira brava      two ounces  
Diluted alcohol      one pint

Digest for seven days, and filter. *Brodie.*

Dose, fifty to sixty drops.

### PAULLINIA.

#### PAULLINIA

Is a preparation from the seeds of *Paullinia sorbilis*, a climbing plant, indigenous to Brazil. The seeds are dried, pounded, mixed with cocoa and cassava, and with water; formed into a paste, which is dried. This preparation is of a brown color, hard, light, inodorous, and of a somewhat astringent taste. It is highly esteemed in Brazil, in diseases of the bowels and bladder, and has been used in Europe, as a tonic in these, and in chlorosis, &c., with much success.

*Ser. Syst.* Octan. trigyn. *Nat. Syst.* Sapindaceæ.

*Martins. Mat. Med. Bras.* 59.

#### LOZENGES OF PAULLINIA.

R. Paullinia      five drachms and a-half  
Vanilla sugar      one pound and a-half  
Water      sufficient

Mix, and form lozenges of ten grains each.

*Gavrelle.*

#### SYRUP OF PAULLINIA.

R. Paullinia      two drachms and a-half  
Simple syrup      two pints

Mix. Dose, half a fl. ounce.

*Gavrelle.*

#### PILLS OF PAULLINIA.

R. Paullinia      sufficient  
Make into pills of a grain and a-half.  
Five to ten, as occasion may require.

*Gavrelle.*

#### ALCOHOLIC EXTRACT OF PAULLINIA.

R. Powdered paullinia      at will  
Alcohol      sufficient

Introduce into a displacement apparatus, and pass the alcohol through till the powder is exhausted; distil off the spirit, and

evaporate to proper consistence. Eight to ten grains, during the day.

*Dechastelus.*

#### CHOCOLATE WITH PAULLINIA.

R. Paullinia      one ounce  
Chocolate      sixteen ounces

Mix, and form a paste. As a restorative, in diseases of debility, chlorosis, &c.

*Beasley.*

### PETROLEUM.

#### PETROLEUM.

A liquid bitumen, of a fluid consistence, of a brownish black, or reddish brown color, having a bituminous odor, and an acrid, strong taste. It is found in various parts of the world, issuing from the earth in the form of springs. Many of these exist in the United States, as on the shores of Seneca Lake, on the Kenhawa, &c. These products are light colored and more liquid than the Barbadoes and other foreign varieties, and are known as Seneca oil.

Petroleum is a stimulating antispasmodic and sudorific, with some anthelmintic properties. The dose is from half a drachm to a drachm.

#### BRITISH OIL.

R. Oil of turpentine  
flaxseed, each,      eight fl. ounces  
amber  
juniper, each,      four fl. ounces  
Barbadoes petroleum      three fl. ounces  
Seneca oil      one fl. ounce

Mix. As a rubefacient liniment.

*Phil. Coll. Pharm.*

R. Oil of turpentine      eight fl. ounces  
Barbadoes petroleum      four fl. ounces  
Oil of rosemary      four fl. drachms

Mix. *Gray.*

#### EMBROCATION OF PETROLEUM.

R. Petroleum      half an ounce  
Oil of turpentine      one drachm  
Spirit of juniper      four ounces

Mix. As a friction to the loins, in atony of the urinary passages and dropsy.

*Radiis.*

R. Petroleum      half an ounce  
Spirit of lavender      one ounce  
Laudanum      two drachms

Mix. As a lotion to frozen limbs.

*Phœbus.*

R. Petroleum      one ounce  
Alcohol, each,      one ounce  
Mix. *Saunders.*

R. Petroleum      one drachm and a-half  
Spirit of ammonia      two ounces

Mix. As a remedy for chilblain. *Radiis.*

#### OINTMENT OF PETROLEUM.

R. Petroleum      three drachms



Camphor one scruple  
Simple ointment one ounce and a-half  
Mix. *Radiis.*

## PLASTER OF PETROLEUM.

R. Resin half an ounce  
Camphor one drachm  
Opium half a drachm  
Petroleum sufficient  
Mix. As an application in chronic rheumatism. *Radiis.*

## ANTHELMINTIC MIXTURE OF PETROLEUM.

R. Petroleum half an ounce  
Tincture of assafetida six drachms  
Mix. Forty drops, three times a-day, in cases of tape-worm. *Schwartz.*

## DIURETIC MIXTURE OF PETROLEUM.

R. Petroleum two drachms  
Tincture of squill one drachm  
Spirit of juniper six drachms  
Mix. Forty drops, three times a-day, in dropsy. *Vogt.*

## PETROSELINUM.

## PARSLEY.

Parsley, or *Petroselinum sativum*, is a biennial, herbaceous plant, with a fusiform root; a native of southern Europe, but generally cultivated in our gardens, for the sake of its leaves, which are used for culinary purposes.

*Sex. Syst.* Pentand. digyn. *Nat. Syst.* Apiaceæ.  
Hoffmann, Umb. 1, t. 1. Griffith, Med. Bot. 318.

The root, which is the officinal portion, is fusiform, white, fleshy, of a pleasant odor and a somewhat sweetish, aromatic taste, becoming inert by drying; it should therefore be used in recent state. It is diuretic and slightly aperient.

## INFUSION OF PARSLEY ROOT.

R. Parsley root one ounce  
Boiling water one pint  
Infuse and strain. A teacupful, with a drachm of sweet spirit of nitre, every three or four hours. *Ellis.*

In strangury from blisters; and highly spoken of by Dr. Chapman, in dropsies.

## OIL OF PARSLEY.

R. Parsley seeds one part  
Water four parts  
Distil, and separate the oil. *Brunswick Ph.*

Carminative, and also said to be very efficacious in gonorrhœa, in doses of three or four drops, three times a-day.

## PHLORIDZINA.

## PHLORIDZINE.

A bitter principle obtained from the bark of the apple, pear, cherry, and plum, especially from that of the root. It is white, very bitter, and is said to be a powerful febrifuge. It is obtained by boiling the fresh bark of the root for two hours in sufficient water to cover it, decanting the decoction, and renewing the process. The two decoctions are united, permitted to stand for twenty-four hours, when a deposit of phloridzine will take place, and, by evaporation, an additional portion will be obtained. This impure product is to be treated with distilled water and animal charcoal, to purify it.

## POWDER OF PHLORIDZINE.

R. Phloridzine ten to fourteen grains  
Sugar one drachm  
Mix. In the apyrexia of intermittent fever. *De Koninck.*

## PHOSPHORUS.

## PHOSPHORUS.

An elementary substance, usually in the form of thin sticks, which are translucent, of a pale yellowish color, and flexible consistence. It has no perceptible taste, but an alliaceous odor. It is principally made from bone ashes, which consist mainly of phosphate of lime. It is a highly powerful, diffusible stimulant, and ought never to be given in substance, and its effects should be always closely watched. It has been given in cases of extreme prostration of the vital powers.

## PHOSPHORATED OIL.

R. Phosphorus one part  
Almond oil sixteen parts  
dissolve, and add  
Oil of bergamot four drops  
Mix. Dose, five to ten drops in some mucilaginous liquid. *Magendie.*

Has been advised in gout, chronic rheumatism, &c., and also externally, as a friction.

## PHOSPHORATED ETHER.

R. Phosphorus one part  
Sulphuric ether sixty parts  
Macerate for a month in a bottle covered with black paper; decant into small bottles similarly prepared. Dose, five to ten drops, in emollient liquid, every four hours. *Van Mons.*  
R. Phosphorus two grains  
Oil of peppermint half a drachm  
add to the solution  
Sulphuric ether half fl. ounce



Mix well. Two to six drops, every four hours, on sugar. In epilepsy, paralysis, &c. *Augustin.*

#### PHOSPHORATED CERATE.

R. Phosphorated ether five parts  
Simple cerate twenty-four parts  
Mix. As a friction in obstinate cutaneous affections. *Foy.*

#### PHOSPHORATED OIL OF TURPENTINE.

R. Phosphorus two grains  
Oil of turpentine four fl. drachms  
Mix, and dissolve. Dose, ten to twelve drops, in obstinate intermittents; to be given in a cupful of oatmeal gruel. *Hufeland.*

#### PHOSPHORATED LINIMENT.

R. Phosphorus six grains  
Oil of almonds one ounce  
Caustic ammonia  
Camphor, each, two grains  
Mix. *Augustin.*

#### PHOSPHORATED OINTMENT.

R. Phosphorated ether one drachm  
Lard one ounce  
Rub together. *Guibourt.*

#### PHOSPHORUS PASTE.

R. Phosphorus eight parts  
Lukewarm water one hundred and eighty parts  
Rye meal one hundred and eighty parts  
Melted butter one hundred and eighty parts  
Sugar one hundred and twenty parts  
Liquefy the phosphorus in the water; pour into a mortar, and add the meal; when cold, add the butter and sugar. An active poison for rats and mice. *Simon.*

### PHYTOLACCA.

#### POKE.

Both the berries and the root of the Poke, or *Phytolacca decandra*, are officinal. It is a large, herbaceous plant, with a perennial root, and bearing numerous clusters of dark purple berries.

*Sex. Syst.* Decand. decagyn. *Nat. Syst.* Phytolaccaceæ.

*Linn. Sp. Pl.* 631. *Griffith, Med. Bot.* 535.

The root, as found in the shops, is in transverse slices, of a whitish brown externally; and exhibiting on the cut surfaces numerous concentric rings. The taste is somewhat saccharine, followed by a sensation of acrimony; the odor of the fresh root is somewhat like that of ginseng, but this disappears on drying. It is emetocathartic, with some narcotic properties. It has been used as an alterative, in syphilis, rheumatism, and chronic eruptions. The berries have

also been praised in the same complaints. The dose of the powdered root, is from one to five grains, as an alterative; from ten to thirty as an emetic.

#### OINTMENT OF POKE.

R. Powdered root, or leaves of poke one drachm  
Lard one ounce  
Rub together. As an application in tinea capitis, psora, &c. *G. B. Wood.*

### PIMENTA.

#### ALLSPICE.

This, which is also called *Jamaica pepper*, is the unripe berries of *Eugenia pimenta*, a beautiful evergreen tree, which is indigenous to the West Indies and South America.

*Sex. Syst.* Icosand. monog. *Nat. Syst.* Myrtaceæ.

*Decandolle, Prod.* iii. 285. *Griffith, Med. Bot.* 300.

The berries, which are the officinal part, are collected before they are ripe, and dried in the sun. In this state they are round, brown, somewhat rough, and a little larger than a peppercorn. They have an aromatic, agreeable odor, and a powerful, clove-like taste. Pimento is a warm, aromatic stimulant, but is more employed as a condiment than in medicine. The dose, in powder, is from ten to forty grains.

#### BOLUS OF ALLSPICE.

R. Powdered allspice  
cinnamon  
saffron, each, four grains  
opium one grain  
conservé of roses sufficient  
Mix, and make two boluses. In chronic diarrhoea. *Foy.*

#### WATER OF ALLSPICE.

R. Allspice one pound  
Diluted alcohol seven fl. ounces  
Water sixteen pints  
Distil eight pints. *Lond. Ph.*  
Used as a carminative. Dose, one to two fl. ounces.

#### SPIRIT OF ALLSPICE.

R. Allspice, bruised, three ounces  
Diluted alcohol one gallon  
Water sufficient  
To prevent empyreuma. Macerate for twenty-four hours; distil off a gallon. Used in same cases as other aromatic spirits, in doses of one to two fl. drachms. *Dub. Ph.*

#### TINCTURE OF ALLSPICE.

R. Allspice one part  
Alcohol five parts  
Macerate for fifteen days, and filter. Dose, from ten to twelve drops. *Swed. Jour.*



## PIPER.

## BLACK PEPPER.

This article is the berries of *Piper nigrum*, a perennial, shrubby vine, from eight to twelve feet high, a native of India, and extensively cultivated in many parts of tropical Asia.

*Sex. Syst.* Decand. trigyn. *Nat. Syst.* Piperaceæ.  
Linn. Sp. Pl. 40. Griffith, Med. Bot. 564.

The fruit, which is the officinal part, is gathered before it is quite ripe, and by the drying of the pulp over the grayish white seed, become wrinkled and black; in this state they are known as *black pepper*; if permitted to ripen, and deprived of their skin by maceration, and dried, they form what is called *white pepper*, which is less active than the black.

Black pepper is a warm, carminative stimulant, but is principally used as a condiment. It has, however, been used in gonorrhœa, &c., and in the treatment of intermittent fevers, and also as an external application, in the form of ointment, to tinea capitis. The dose is from ten to twenty grains.

## CONFECTION OF BLACK PEPPER.

R. Powdered black pepper  
elecampane, each, one pound  
fennel seeds three pounds

Honey

Sugar, each, two pounds

Rub the first three ingredients and the sugar together, and keep in a covered vessel. When to be used, add the honey, and incorporate thoroughly (or seven ounces of the powder with two ounces of the honey). *Lond. Ph.*

This, which is the same with *Ward's paste*, so celebrated in the cure of piles, must be used for a length of time, to be of service. Dose, from one to two drachms, two or three times a-day.

## WARD'S PASTE.

R. Powdered black pepper  
elecampane, each, eight ounces  
fennel seed, twenty-two ounces

Honey

Sugar, each, one pound

Triturate well together, and form an electuary. As a remedy in hemorrhoids. A piece, the size of a nutmeg, to be taken three times a-day. *Gray.*

## ELECTUARY OF BLACK PEPPER.

R. Powdered black pepper  
Conserve of orange, each, one ounce  
Syrup of orange peel sufficient  
Make an electuary. As a stomachic and carminative. A drachm, twice a-day. *Saunders.*

## RESINOUS OIL OF BLACK PEPPER.

R. Black pepper at will

Grind it, and extract by means of alcohol (8.35); distil off the alcohol. Treat the resinous residue with solution of potassa, which removes all but the piperine. The alcoholic liquid is then to be saturated with diluted muriatic acid, and the fluid oleo-resin, that precipitates, is, after being washed with water, ready for use.

*W. Procter.*

## VOLATILE OIL OF BLACK PEPPER.

R. Powdered black pepper at will  
Water sufficient  
Distil, and separate the oil from the product.

## OINTMENT OF BLACK PEPPER.

R. Powdered black pepper four ounces  
Lard one pound  
Mix. As an application in tinea capitis. *Dub. Ph.*

## PLASTER OF BLACK PEPPER.

R. Powdered black pepper  
Galbanum, each, three drachms  
Pitch two ounces  
Oil of laurel berries sufficient  
Mix. As a stimulating plaster, in rheumatism, &c. *Augustin.*

## CATAPLASM OF BLACK PEPPER.

R. Flour of mustard half a pound  
Black pepper  
Ginger, each, one drachm  
Syrup sufficient  
to make a soft paste. To be applied to the pit of the stomach in colic, &c.; or to the feet, as a revulsive. *Ellis.*

PIPER LONGUM,  
LONG PEPPER,

Is the dried spikes of the *Piper longum*, or *Chavica officinarum*, &c., a climbing shrub, with diœcious flowers, a native of some of the East India Islands, and cultivated in Sumatra and Java.

Linn. Sp. Pl. 41. Griffith, Med. Bot. 566.

The officinal portion is the fruit catkins, dried in the sun. These are aromatic; of a pungent, fiery taste; of a grayish brown color; cylindrical. This pepper has the same properties as the black, but it is little used in this country.

## CATAPLASM OF LONG PEPPER.

R. Powdered long pepper  
ginger, each, half an ounce  
White of egg sufficient  
Mix well into a paste. As a rubefacient application in pleurisy. *Foy.*



PIPERINUM.

PIPERINE.

A peculiar principle obtained from *Piper nigrum*. It is white, and crystallizes in four-sided prisms. It was thought to be an alkali, but the experiments of Pelletier show that it is more analogous to the resins. It has been much praised in Italy, as a febrifuge; and is also spoken of with favor by physicians of other countries. It has been supposed that its acrid taste and remedial properties were dependent on an admixture of acrid oil, but Dr. Christison has shown that such is not the case.

R. Alcoholic extract of black pepper at will  
Treat it with water, containing a hundredth of caustic potassa; wash the residue in cold water; dissolve in alcohol; filter, and permit to evaporate and crystallize. *Guibourt.*

Dose, from two to ten grains.

PILLS OF PIPERINE.

R. Piperine twenty-four grains  
Crumb of bread sufficient  
Mix, and make twelve pills. One, every two hours. *Meli.*  
R. Piperine twelve grains  
Extract of gentian sufficient  
Mix, and make twelve pills. One, every hour, during the apyrexia. *Ellis.*

PILLS OF PIPERINE AND MERCURY.

R. Blue pill one grain  
Piperine  
Sulphate of quinine, each, two grains  
Syrup sufficient  
Mix, and make a pill. *Hartle.*

One, three times a day, for a few days, after the paroxysm has been checked by piperine.

TINCTURE OF PIPERINE.

R. Piperine one part  
Alcohol seven parts  
Dissolve, and filter. *Beral.*

PISCIDIA EYTHRINA.

JAMAICA DOGWOOD.

This is a small West Indian tree, with a hard, heavy wood, and a narcotic bark, employed to intoxicate fish.

*Ser. Syst. Diadelph. decand. Nat. Syst. F. baceæ.*

*Linn. Sp. Pl. 993. Griffith, Med. Bot. 246.*

The part used is the bark of the root, which appears to be a stimulating narcotic; causing excitement of the system, copious perspiration, and profound sleep. Applied topically, to allay pain.

TINCTURE OF JAMAICA DOGWOOD.

R. Bruised Jamaica dogwood one ounce

Diluted alcohol four fl. ounces  
Digest for seven days, and filter. Full dose, as a narcotic, one fl. drachm.

*Hamilton.*

PIX ABIETIS.

BURGUNDY PITCH.

This is the prepared concrete juice of *Abies excelsa*, a lofty tree, a native of Europe and northern Asia.

*Ser. Syst. Monœc. monadelph. Nat. Syst. Pinacæ.*

*Decandolle, Fl. Fr. iii. 375. Griffith, Med. Bot. 606.*

It is also obtained from *A. picea*.

When pure, it is hard, brittle, opaque, of a yellowish, or brownish yellow color, and of a weak, turpentine-like taste and odor. The *Abietis resina* of the London College, is the concrete juice of several of the pine tribe, as it exudes. It is in solid, brittle tears, of a brownish-yellow color, and emits an agreeable odor when burned.

Both are used as ingredients in plaster.

PITCH PLASTER.

R. Burgundy pitch two pounds  
Resin of spruce fir one pound  
Resin  
Wax, each, four ounces  
Expressed oil of nutmeg one ounce  
Olive oil  
Water, each, two fl. ounces  
Melt the pitch, resin, and wax together, and add first the resin of the spruce fir, then the olive oil, oil of nutmegs, and water; mix, and boil to a proper consistence. *Lond. Ph.*

As a rubefacient plaster in pectoral, rheumatic, &c., affections. It often causes a serous discharge, and much irritation.

R. Burgundy pitch six ounces  
Wax half an ounce  
Turpentine one drachm  
Melt, and mix. *Guy's Hosp.*

AROMATIC PLASTER.

R. Burgundy pitch three ounces  
Yellow wax half an ounce  
Powdered cinnamon six drachms  
Oil of pimento  
lemons, each, two drachms  
Melt the resin and wax together, and strain. When they begin to thicken, on cooling, mix in the cinnamon, previously rubbed with the oils, and make a plaster. *Dub. Ph.*

A good local stimulant, when applied to the region of the stomach, allaying nausea and vomiting, and relieving gastric uneasiness.

COMPOUND PITCH PLASTER.

R. Burgundy pitch two parts



White wax four parts  
 Turpentine  
 Balsam of tolu, each, one part  
 Mix, and melt. *Beral.*  
 R. Burgundy pitch twenty-four parts  
 Resin  
 Yellow wax, each, four parts  
 Resin of spruce fir twelve parts  
 Oil of nutmegs one part  
 Melt together. *Lisbon Ph.*

## PITCH CERATE.

R. Burgundy pitch six parts  
 Suet four parts  
 White wax three parts  
 Melt together. *Beral.*

## PILLS OF BURGUNDY PITCH.

R. Burgundy pitch five drachms  
 Mucilage sufficient  
 Mix, and make one hundred pills. Six to eight, three times a-day, in cutaneous affections. *Ulrich.*

## PIX CANADENSIS.

## CANADA, OR HEMLOCK PITCH.

The prepared concrete juice of the *Abies Canadensis*, a lofty tree, found in the more northern parts of the United States, in Canada, &c., and also in the mountainous portions of the Middle States.

*Ser. Syst.* Monœc. monadelph. *Nat. Syst.* Pinaceæ.

Mich. N. A. Sylv. iii. 185. Griffith, Med. Bot. 606.

The resin exudes spontaneously, and hardens on the bark, from which it is separated by boiling in water. It, however, requires a purification, by melting and straining, before it is fit for use. In this state, it is hard, brittle, of a dark yellowish brown color, which becomes darker by exposure to the air, of a peculiar, but faint odor, and a scarcely perceptible taste.

It is a mild rubefacient, much resembling Burgundy pitch in its action and powers, and is employed for the same purposes.

## PIX LIQUIDA.

## TAR.

This is an impure turpentine, obtained by the slow combustion of various species of the pine tribe. It is made in large quantities in North Carolina, and in various other places in America and Europe.

It is of a brownish black color, of a tenacious consistence, has a peculiar empyreumatic odor, and a somewhat bitter, resinous, subacid taste.

Its properties are analogous to those of the turpentines. It is used both internally and externally. The dose is from half a drachm to one drachm, several times a-day.

## TAR PILLS.

R. Tar one drachm  
 Powdered elecampane sufficient  
 Mix, and form a mass. *Beasley.*  
 R. Tar one drachm  
 Wheat flour sufficient  
 Mix, and form a mass. *G. B. Wood.*  
 R. Tar two scruples  
 Liquorice powder one scruple  
 Mix, and make sixteen pills. *Seymour.*

## TAR WATER.

R. Tar two pints  
 Water one gallon  
 Mix, stirring with a wooden rod for fifteen minutes; when the tar has subsided, strain, and keep in well-stopped bottles. *Dub. Ph.*

It is stimulant and diuretic, and was once much used in a variety of diseases, especially in those of the lungs. The dose is from one to two pints, in divided doses, in the course of the day. It is also used as a lotion in cutaneous diseases.

## TAR OINTMENT.

R. Tar  
 Suet, each, one pound  
 Melt the suet with a moderate heat, add the tar, and stir till cold. *U. S. Ph.*

A stimulant application to various cutaneous eruptions, as psoriasis and tinea capitis.

## COMPOUND TAR OINTMENT.

R. Tar ointment  
 Cerate of acetate of lead, each, half a pound  
 Mix. *St. Bart's Hosp.*  
 R. Tar ointment  
 Sulphur ointment equal parts  
 Mix. *Guy's Hosp.*

## PIX NIGRA.

## BLACK PITCH.

This is the solid black mass that remains after the evaporation of tar. It has a shining fracture. It is gently stimulant and tonic, and has been used internally in some cutaneous diseases, and in piles. The dose is from ten grains to a drachm, in piles. It is also employed externally.

## OINTMENT OF BLACK PITCH.

R. Black pitch  
 Wax  
 Resin, each, nine ounces  
 Olive oil sixteen fl. ounces  
 Melt together, and strain. *Lond. Ph.*  
 Used for the same purposes as tar ointment.

R. Black pitch  
 Resin  
 Yellow wax, each, one part  
 Olive oil four parts



Melt the two resins together, add the wax, and then the oil. *Guibourt.*

R. Black pitch sixteen parts  
Yellow wax twelve parts  
Lard ten parts  
Olive oil two parts  
Melt together. *Pipon.*

#### PLASTER OF BLACK PITCH.

R. Black pitch  
Resin  
Suet  
Turpentine, each, two parts  
Yellow wax three parts  
Melt together. *Niemann.*

R. Black pitch  
Resin  
Wax, each, eight parts  
Suet one part  
Melt together. *Wirtem. Ph.*

### PLATINUM. PLATINA.

#### PLATINI BICHLORIDUM.

##### BICHLORIDE OF PLATINA.

Made by dissolving platinum in nitromuriatic acid, and evaporating the solution to dryness by a gentle heat. *Beasley.*

It, as well as a concentrated solution, is of a deep red color. It is very deliquescent and soluble in water. It is poisonous in the dose of fifteen grains, but has been used as an alterative in syphilis.

##### MIXTURE OF BICHLORIDE OF PLATINUM.

R. Bichloride of platinum four grains  
Sugar two drachms  
Distilled water three fl. ounces  
Dissolve. Dose, four fl. drachms, three times a-day. *Hoefler.*

##### PILLS OF BICHLORIDE OF PLATINUM.

R. Bichloride of platinum seven and a-half grains  
Guaiacum one drachm  
Powdered liquorice root sufficient  
Mix, and form twenty pills. One pill three times a-day. *Hoefler.*

##### OINTMENT OF BICHLORIDE OF PLATINUM.

R. Bichloride of platinum one drachm  
Extract of belladonna two drachms  
Lard four ounces  
Rub well together. As an application to indolent ulcers. *Hoefler.*

### SODII CHLOROPLATINAS.

#### CHLOROPLATINATE OF SODIUM.

Mix a solution of six parts of chloride of sodium with a solution of seventeen parts of bichloride of platina; evaporate, and crystallize. The crystals are of a deep yellow color, soluble in water and alcohol. They are analogous in their medical properties to the analogous salt of gold. Dose, one to three grains. *Beasley.*

#### INJECTION OF CHLOROPLATINATE OF SODIUM.

R. Chloroplatinate of sodium half a drachm  
Decoction of poppies eight fl. ounces  
Dissolve. *Hoefler.*

### PLUMBUM.

#### LEAD.

A softish, blue gray, malleable metal, with a perceptible taste, and a peculiar smell when rubbed. It is not officinal in its metallic state, but its preparations are much employed, as sedatives and astringents.

### PLUMBI ACETAS.

#### ACETATE OF LEAD.

##### POWDER OF ACETATE OF LEAD.

R. Acetate of lead  
Powdered opium, each, six grains  
Sugar twenty-four grains  
Mix. Three grains, morning and evening, to check night sweats and diarrhoea, in phthisis. *Foy.*

R. Calomel one to two grains  
Acetate of lead half to one grain  
Mix, and divide into four powders. One, every three hours, in cholera infantum of infants. *Ellis.*

R. Acetate of lead two to three grains  
Opium quarter of a grain  
Acetate of soda three grains  
Sugar of milk five grains  
Mix. To be taken every hour, in hemorrhages. *Phœbus.*

##### PILLS OF ACETATE OF LEAD.

R. Acetate of lead  
Powdered mallows, each, one drachm  
Simple syrup sufficient  
Mix, and make thirty-six pills. Four to five a-day, to check sweats in phthisis. *Radius.*

R. Acetate of lead half a drachm  
Calomel five grains  
Conserve of roses sufficient



Mix, and make ten pills; one, every two to four hours, in hematemesis. *Ellis.*

R. Acetate of lead thirty grains  
Powdered colchicum twenty grains  
opium three grains  
Mucilage of gum Arabic sufficient  
Mix thoroughly, and form into ten pills.  
One to be taken every six hours. Used  
in active hemorrhages, washed down with  
a draught of one drachm of distilled vinegar  
to one fl. ounce of water. Also, given in  
phthisis, after bleeding, one twice a-day.

*A. T. Thomson.*

R. Acetate of lead one scruple  
Opium one grain  
Conserve of roses sufficient  
Mix, and make twelve pills. One every  
hour at first, then every two hours, in  
cholera. *Graves.*

R. Acetate of lead twelve grains  
Opium six grains  
Conserve of roses sufficient  
Mix, and make six pills. One to be  
repeated according to circumstances, in  
hemoptysis, &c. *Ellis.*

R. Acetate of lead six grains  
Powdered mallows  
Extract of seneka, each, one drachm  
Mix, and make sixty pills. Two to five,  
several times a-day, in hemoptysis. *Kopp.*

#### RUBEFACIENT PASTE.

R. Acetate of lead one ounce  
Acid sulphate of potassa three ounces  
Water sufficient  
Rub into a paste. It acts powerfully on  
the skin. *Clarus.*

#### COLLYRIUM OF ACETATE OF LEAD.

R. Solution of acetate of  
lead twelve drops  
Wine of opium forty drops  
Rose water four fl. ounces  
Mix. *Ellis.*

R. Acetate of lead one scruple  
Lime water twelve fl. ounces  
Catechu three drachms  
Honey of roses two ounces  
Dissolve, and filter. This is much more  
astringent than the last. *Augustin.*

#### SOLUTION OF SUBACETATE OF LEAD.

R. Acetate of lead sixteen ounces  
Litharge nine ounces and a-half  
Distilled water four pints  
Boil together, for half an hour, adding  
distilled water, so as to preserve the mea-  
sure, filter, and keep in close-stopped bot-  
tles. *U. S. Ph.*

The *diluted solution* is made by mixing two fl. drachms of the above solution to a pint of dis-  
tilled water.

#### ACETATE OF LEAD CERATE.

Acetate of lead four parts  
Soap one part and a-half  
Mix thoroughly in a heated mortar, and  
add  
White wax three parts  
Previously melted with  
Olive oil three parts  
Mix well. *Van Mons.*

#### PLASTER OF ACETATE OF LEAD.

R. Acetate of lead one drachm  
Flowers of sulphur half an ounce  
Resin two ounces  
Melt together. As an application to vene-  
real buboes. *Bories.*

#### PILE OINTMENT.

R. Acetate of lead fifteen grains  
Burnt cork half an ounce  
Fresh butter two ounces  
Triturate well together. *Bories.*

#### ACETATE OF LEAD OINTMENT.

R. Acetate of lead  
Extract of belladonna, each, one part  
Lard six parts  
Mix. As an application to fissures of the  
anus. *Foy.*

#### SUBACETATE OF LEAD CATAPLASM.

R. Common cataplasm four ounces  
Solution of subacetate  
of lead one fl. ounce  
Muriate of ammonia half a drachm  
Mix. As an application to indolent tu-  
mors. *Radius.*

#### SUBACETATE OF LEAD COLLYRIUM.

R. Solution of subacetate of lead five drops  
Elder water half a fl. ounce  
Laudanum ten drops  
Mix. *Rust.*

#### SUBACETATE OF LEAD GARGLE.

R. Solution of subacetate  
of lead half a fl. drachm  
Barley water one pint  
Simple syrup one ounce  
Mix. *Radius.*

#### SUBACETATE OF LEAD INJECTION.

R. Solution of subacetate  
of lead two fl. drachms  
Distilled vinegar eight fl. ounces  
Rose water twenty-four fl. ounces  
Mix. In leucorrhœa. *Young.*

#### LINIMENT FOR MILK ABSCESS.

R. Olive oil two ounces



Lead water                    one drachm  
Sulphuric ether            two drachms  
Laudanum                   one drachm  
Mix. A rag moistened with this liniment  
to be frequently applied to the inflamed  
part. *Dewees.*

R. Acetate of lead            one drachm  
Distilled vinegar           two ounces  
Dissolve, and add  
Alcohol                    one ounce  
Distilled water            five ounces  
Mix. To be applied as above. *Clarke.*

## ACETATE OF LEAD LOTION.

R. Diluted solution of sub-  
acetate of lead            one fl. ounce  
Spirit of camphor        three fl. ounces  
Water                    one pint  
Mix. As an application to old ulcers.  
*Ellis.*

## ACETATE OF LEAD CATAPLASM.

R. Crumb of bread           one pound  
Milk                    four pounds  
Boil together, and add  
Solution of subacetate  
of lead                    one fl. ounce  
*Spielmann.*

## VEGETO MINERAL WATER OF GOULARD.

R. Solution of subacetate of lead one part  
Alcohol (.921.)            four parts  
Distilled water            forty-eight parts  
Mix. *Taddei.*

Used in fomentations, lotions, and cataplasms,  
and may always be replaced by the diluted so-  
lution of the subacetate.

## ACETATE OF LEAD GARGLE.

R. Acetate of lead            sixty grains  
Extract of opium           six grains  
Rose water                eight fl. ounces  
Syrup of mulberries       one ounce  
Mix. *Pierquin.*

## ACETATE OF LEAD MIXTURE.

R. Acetate of lead            half an ounce  
Sulphate of iron           three drachms  
Vinegar  
Alcohol, each,            two fl. ounces  
Rose water                six fl. drachms  
Dissolve the acetate in the vinegar with a  
gentle heat; add the sulphate, and then  
the alcohol, mixed with the rose water.  
Highly praised in the sweats and colliqua-  
tive diarrhoea of phthisis; in gonorrhoea  
and nocturnal emissions. *Gormann.*

## GOULARD'S BALSAM.

R. Oil of turpentine           at will

Heat, and gradually add  
Acetate of lead            sufficient  
Constantly stirring, till no more will dis-  
solve; let rest, and decant while hot.  
Used as an application to eroding and  
painful ulcers. *Van Mons.*

## SUBACETATE OF LEAD CERATE.

R. Solution of subacetate  
of lead            two fl. ounces and a-half  
White wax                four ounces  
Olive oil                nine fl. ounces  
Camphor                half a drachm  
Melt the wax, and mix with eight fl.  
ounces of the oil; remove from fire, and  
when it begins to thicken, gradually add  
solution of subacetate of lead; stir till  
cool, then add camphor, dissolved in rest  
of oil, and mix. *U. S. Ph.*

INJECTION OF SUBACETATE OF LEAD  
AND LIME WATER.

R. Solution of subacetate  
of lead            thirty-four drops  
Lime water            four fl. ounces  
Olive oil                two fl. drachms  
Mix, and shake whenever used. In in-  
flammation of the prostate and urethra.  
*Foy.*

## LINIMENT OF SUBACETATE OF LEAD.

R. Solution of subacetate of lead  
Laudanum  
Honey of roses, each,    two fl. drachms  
Conserve of roses        one ounce  
Mix. *Guy's Hos.*

## SUBACETATE OF LEAD OINTMENT.

R. Cerate of subacetate of lead  
Poplar ointment, each,    one ounce  
Powdered camphor        two drachms  
Mix. In inflamed acne and other cutane-  
ous affections. *St. Marie.*

R. Solution of subacetate of lead  
White wax, each,        one drachm  
Olive oil                one ounce  
Mix. As an application in ophthalmia.  
*Armstrong.*

## SUBACETATE OF LEAD PLASTER.

R. Olive oil                fourteen fl. ounces  
Yellow wax                twenty-four ounces  
Melt together, and add  
Solution of subacetate  
of lead                one fl. ounce  
Camphor, rubbed in a  
little oil                half a drachm  
Mix well, and spread on linen or skin.  
It has been much praised in rheumatic  
pains, and indolent ulcers. *Van Mons.*



## PLUMBI CARBONAS.

## CARBONATE OF LEAD.

## CAMPHORATED POWDER OF CARBONATE OF LEAD.

R. Carbonate of lead           ten drachms  
 Starch                           two drachms  
 Sarcocolla  
 Gum Arabic  
     tragacanth, each,       one drachm  
 Camphor                   half a drachm  
 Triturate together. As an application to  
 ulcers. *Giannini.*

## OINTMENT OF CARBONATE OF LEAD.

R. Carbonate of lead           two ounces  
 Simple ointment           one pound  
 Powder the carbonate very fine, add it to  
 the ointment previously melted, and mix.  
*U. S. Ph.*

As an application to excoriated or ulcerated  
 surfaces.

R. Carbonate of lead           six ounces  
 Camphor                   one ounce  
 Olibanum                  two ounces  
 Rose oil                   three ounces  
 White of eggs               seven  
 Triturate together. Same doses as above.  
*Harrer.*

R. Carbonate of lead           one ounce  
 Litharge                  two drachms  
 Bole armenian           one ounce  
 Honey of roses           half an ounce  
 Lard                       sufficient  
 Triturate together. As a dressing to pha-  
 gedenic and cancerous ulcers. *Swediaur.*

R. Carbonate of lead           one ounce  
 Opium                   one drachm  
 Lard                   half an ounce  
 Anodyne balsam           sufficient  
 Mix. As an application in facial neural-  
 gia. *Fouquier.*

## CAMPHORATED CARBONATE OF LEAD

## OINTMENT.

R. Carbonate of lead           one ounce  
 Simple ointment           five ounces  
 Camphor                   two scruples  
 Mix. *Beasley.*

## PLASTER OF CARBONATE OF LEAD.

R. Carbonate of lead           one pound  
 Olive oil                  two pounds  
 White wax               three ounces  
 Water                   two pounds  
 Melt the wax and oil together, add the  
 carbonate of lead, and stir till mixed, then  
 add the water and boil; malaxate, and  
 melt again without the wax. *Par. Cod.*

R. Carbonate of lead           one pound  
 Olive oil                  two pints  
 Yellow wax               four ounces  
 Lead plaster           one pound and a-half  
 Powdered orris root       nine ounces

Boil together the oil and carbonate of lead,  
 adding a little water, and constantly stir-  
 ring, till they are perfectly incorporated,  
 then add the wax and plaster; when these  
 are melted, mix in the orris, and stir well.

*U. S. Ph., 1830.*

As a substitute for Mahy's plaster, and a good  
 application to inflamed and excoriated surfaces,  
 as bed sores, &c.

## PLUMBI CHLORIDUM.

## CHLORIDE OF LEAD.

R. Acetate of lead           nineteen ounces  
 Chloride of sodium       six ounces  
 Dissolve the two salts separately, the for-  
 mer in three pints of boiling, distilled  
 water, the latter in one pint of distilled  
 water; mix the solutions, and wash the  
 precipitate with distilled water.

*Lond. Ph.*

## OINTMENT OF CHLORIDE OF LEAD.

R. Chloride of lead           one drachm  
 Simple cerate           one ounce  
 Mix. As an application in cancerous ul-  
 cerations. *Tuson.*

## LOTION OF CHLORIDE OF LEAD.

R. Chloride of lead           one drachm  
 Water                   one pint  
 Dissolve; as a wash to cancerous ulcera-  
 tions, and in painful neuralgic tumors.  
*Tuson.*

## PLUMBI IODIDUM.

## IODIDE OF LEAD.

R. Solution of iodide of potassium  
     acetate of lead, each,  
    sufficient

Add one solution gradually to the other,  
 till there is no longer a precipitate, wash  
 this in cold water, and dry it. *Cottureau.*

R. Iodide of potassium  
     Nitrate of lead, each,   half an ounce  
     Water                   a pint and a-half

Dissolve the salts separately, each in one-  
 half of the water; mix the solutions, col-  
 lect the precipitate on a filter of linen or  
 muslin, and wash it with water. Boil the  
 powder in three gallons of water, acidulated  
 with three ounces of pyroligneous acid.  
 Let any undissolved matter subside, main-  
 taining the temperature near the boiling



point, and pour off the clear liquor, from which the iodide of lead will crystallize on cooling. *Ed. Ph.*

This is a preferable formula to that made with the acetate of lead, as above, or that of the *Lond. Ph.*

#### PILLS OF IODIDE OF LEAD.

R. Iodide of lead four grains  
Conserve of roses one scruple  
Mix, and form twelve pills. One, night and morning, gradually increased, in scrofulous affections. *Foy.*

R. Iodide of lead half a drachm  
Conserve of roses sufficient  
Mix, and form one hundred and twenty pills. Administered as above. *Gassicourt.*

#### POWDER OF IODIDE OF LEAD.

R. Iodide of lead  
Powdered foxglove, each, six grains  
Alcoholic extract of stramonium two grains  
Sugar one drachm  
Mix, and divide into twenty-four powders. One, four times a-day. *Gassicourt.*

#### OINTMENT OF IODIDE OF LEAD.

R. Iodide of lead one ounce  
Lard eight ounces  
Mix. As an application to indolent and scrofulous swellings. *Lond. Ph.*

### PLUMBI OXIDUM SEMIVITREUM.

#### LITHARGE.

##### LEAD PLASTER.

R. Litharge, in fine powder, five pounds  
Olive oil one gallon  
Water two pints  
Boil together over a gentle fire, stirring constantly, till a plaster is formed. *U. S. Ph.*

##### LEAD CERATE.

R. Lead plaster six parts  
White wax half a part  
Melt, mix, and add  
Olive oil two parts and a-half  
*Van Mons.*

##### LEAD OINTMENT.

R. Olive oil eight parts  
Lard  
Butter  
Mutton fat  
Yellow wax  
Litharge, each, two parts  
Heat the oil, fat, and wax together, till they emit fumes; gradually add the li-

tharge; constantly stirring, till the mixture assumes a dark brown color, and add

Pitch one part  
Strain. *Foy.*

This preparation is much used in France, under the name of *Onguent de la Mere*, as a maturant to buboes, abscesses, &c.

#### COMPOUND LEAD OINTMENT.

R. Lead plaster three pounds  
Olive oil one pint  
Melt together, and add a mixture of  
Prepared chalk eight ounces  
Distilled vinegar six fl. ounces  
whilst warm, to the warm plaster and oil, and stir till cold. *Lond. Ph.*

R. Fresh root of figwort three ounces  
Leaves of henbane three handfuls  
Fresh butter one pound and a-half  
Litharge two ounces  
Yolk of eggs two  
Boil the root and leaves in the butter, till all moisture is driven off; express, and strain; add the litharge; stir till cold, and add the yolk of eggs, and incorporate well. *Saxon Ph.*

Celebrated as an application to painful hemorrhoids.

#### DIAPALMA PLASTER.

R. Lead plaster nine pounds  
White wax nine ounces  
Melt by a gentle heat, and add  
Sulphate of zinc four ounces  
Dissolve in a little water; continue the heat, constantly stirring till all moisture is evaporated. *Guibourt.*

#### GAULTHER'S PLASTER.

R. Diapalma plaster twelve parts  
Olive oil  
White wax, each, one part  
Turpentine two parts  
Melt the first three articles together, with a gentle heat; add the turpentine. *Guibourt.*

#### BAYNTON'S ADHESIVE PLASTER.

R. Lead plaster one pound  
Resin six drachms  
Melt together. *Beasley.*

#### LEAD CATAPLASM.

R. Flaxseed meal  
Arrowroot, each, two ounces  
Water eleven fl. ounces  
Lead ointment one ounce  
Mix, and heat to a proper consistence. As a maturing application. *Beral.*

#### BANYER'S OINTMENT.

R. Litharge two ounces



Burned alum  
Calomel, each, one ounce and a-half  
Lard two pounds  
Venice turpentine half a pound  
Mix. As an application in porrigo.  
*Cazenave & Schledel.*

## PLUMBI OXIDUM RUBRUM.

## RED LEAD.

## NUREMBERG PLASTER.

R. Red lead eight ounces  
Olive oil one pound  
Mix, and heat till it assumes a brown color, and add  
Resin half an ounce  
Yellow wax one ounce and a-half  
Camphor two drachms  
Mix, and stir well till incorporated.  
*Hessian Ph.*

## PLUMBI SACCHARAS.

## SACCHARATE OF LEAD.

R. Mix one part of sugar with two parts of nitric acid, diluted with ten of water, and apply heat as long as any reaction takes place; neutralize with chalk, filter, and add to the filtered solution acetate of lead as long as any precipitate is found; wash this, and dry it.  
*Beasley.*

## NITRO-SACCHARATE OF LEAD.

R. Saccharate of lead at will  
Dissolve it in nitric acid, diluted with nineteen parts of water, filter, evaporate, and set aside to crystallize.  
*Hoskins.*

## SOLUTION OF NITRO-SACCHARATE OF LEAD.

R. Nitro-saccharate of lead five grains  
Saccharic acid twenty-five drops  
Water five ounces  
Dissolve. Proposed as a solvent for phosphatic calculi.  
*Hoskins.*

## PLUMBI TANNAS.

## TANNATE OF LEAD.

R. Strong infusion of galls at will  
Solution of acetate of lead sufficient  
Add the solution of the acetate, drop by drop, till it no longer causes a precipitate; wash this, and dry it.

## LINIMENT OF TANNATE OF LEAD.

R. Decoction of oak bark eight fl. ounces  
Solution of acetate of lead sufficient

Add the solution of lead to the decoction by degrees, till it causes no precipitate; wash the precipitate, and add  
Alcohol, two drachms  
Recommended in bed sores. *Sundelin.*

## OINTMENT OF TANNATE OF LEAD.

R. Recent tannate of lead twelve parts  
Lard thirty parts  
Mix. As an application to bed sores.  
*Tott.*

R. Decoction of oak bark six ounces  
Solution of acetate of lead one ounce and a-half  
Mix. Collect the precipitate, and mix whilst moist,  
Lard one ounce  
Camphor ten grains  
As an application to excoriation. *Sundelin.*

## PODOPHYLLUM.

## MAY APPLE.

This is the root or rhizome of the *Podophyllum peltatum*, a native, herbaceous plant, growing in damp situations, having two large peltate, palmate leaves.

*Sex. Syst.* Polyand. monog. *Nat. Syst.* Berberidaceæ.

The dried root is about as thick as a quill, of a blackish brown color externally, and somewhat corrugated; within, it is of a dirty white; it has a faint but unpleasant odor, and a bitterish, somewhat sweetish taste. It is a purgative of the same character as jalap, and may be given in the same combinations. Dose, ten grains to a scruple.

## EXTRACT OF MAY APPLE.

R. May apple root, in coarse powder one pound  
Alcohol four pints  
Water sufficient  
Macerate the May apple root for four days in the alcohol, and then obtain tincture by displacement with the water, then by same means obtain six pints of infusion; distil off the alcohol from tincture, and evaporate the infusion, till each are of the consistence of thin honey. Mix, and evaporate to a due consistence. Dose, five to fifteen grains. *U. S. Ph.*

## PODOPHYLLIN.

R. May apple root  
Alcohol a sufficient quantity  
Animal charcoal, in coarse powder, six ounces  
Reduce the may apple root to powder, moisten it with half its weight of alcohol, introduce into a displacer, and pour on al-



cohol until four pints of tincture are obtained. Distil off the alcohol, treat the residual extract with ether to remove fixed oil, boil the residue with water, and dissolve what remains in alcohol, 42° B; treat it with purified animal charcoal, and allow the decolorized solution to evaporate spontaneously. The dry, amorphous mass is podophyllin, and, in cathartic power, six grains are equal to eight of jalap resin.

*J. R. Lewis.*

## POLYGONUM.

### POLYGONUM BISTORTA.

#### BISTORT.

A sub-aquatic plant, with a perennial root and annual, herbaceous stem, growing in moist places in Europe and the northern parts of Asia and America.

*Sex. Syst.* Octand. trigyn. *Nat. Syst.* Polygonaceæ.

Linn. Sp. Pl. 516. Steph. and Church. 1 t. 47.

The root is twisted or bent, of a deep brown color externally, reddish within, inodorous, but of a rough, astringent taste. It is a powerful astringent, but is seldom used in this country. Dose, from twenty to thirty grains.

#### EXTRACT OF BISTORT.

*R.* Powdered bistort one pound  
Treat it by the process of displacement, and evaporate the filtered fluid.

*Pharm. Hosp. Mil.*

#### MIXTURE OF BISTORT.

*R.* Powdered bistort two drachms  
Syrup of quince one ounce  
Rub together, and add  
Tincture of catechu two fl. drachms  
Water four fl. ounces

A spoonful every hour, shaking the bottle each time, in passive hemorrhages, atonic mucous discharges.

*Foy.*

## POTASSIUM.

### POTASSIUM

Is a soft, silver-white metal, readily oxidizable by the action of the air; when thrown on water it takes fire and burns with a rose-colored flame, combining with oxygen, and forming potassa, which is dissolved in the water. It must be kept in liquids, into whose composition no oxygen enters, as naphtha, &c. It is not used in medicine, but its numerous compounds are highly important.

*R.* Fused carbonate of potassa one pound  
Iron filings three quarters of a pound  
Powdered charcoal one pound

Mix, and introduce into a gun barrel or iron bottle, placed in a furnace so made that the flame of a very strong fire may surround it, and maintain every part at a uniform heat approaching to whiteness; the gun barrel or bottle is to be connected by an iron tube with a copper receiver partly filled with naphtha and surrounded by ice; the heat is to be gradually raised until the requisite temperature is reached, when the potassium will distil over.

*Curaudau.*

## POTASSI BROMIDUM.

### BROMIDE OF POTASSIUM.

*R.* Bromine two ounces  
Iron filings one ounce  
Distilled water three pints

To a pint and a half of the water, first add the iron and then the bromine. Set them aside for half an hour, occasionally stirring. Apply a gentle heat, and when the liquid becomes greenish, add seventeen drachms of carbonate of potassa dissolved in the rest of the water. Strain, and wash what remains in two pints of boiling distilled water, and again strain. Evaporate the mixed liquors to crystallization. *Lond. Ph.*

Alterative and resolvent; used both externally and internally in bronchocele and scrofula, in doses of two to ten grains, three times a-day.

### SOLUTION OF BROMIDE OF POTASSIUM.

*R.* Bromide of potassium twelve grains  
Lettuce water three ounces  
Syrup of marsh mallows one ounce  
Mix. To be given in spoonful doses, in the twenty-four hours, in scrofula. *Foy.*

### OINTMENT OF BROMIDE OF POTASSIUM.

*R.* Bromide of potassium thirty-four grains  
Lard one ounce  
Mix. In frictions on scrofulous swellings, and to tinea and tetter. *Magendie.*

### COMPOUND OINTMENT OF BROMIDE OF POTASSIUM.

*R.* Bromide of potassium twenty-four grains  
Liquid bromine six to twelve drops  
Lard an ounce  
Mix. *Magendie.*

## POTASSII CHLORIDUM.

### CHLORIDE OF POTASSIUM.

*R.* Carbonate of potassa at will  
Dissolve in a sufficient quantity of water, filter, and add muriatic acid to saturation,



concentrate the solution by evaporation, and cool slowly, that crystals may form.

*Guibourt.*

Formerly known as the Salt of Sylvius, and used in fever to excite perspiration, and as a diuretic, in doses of from one to two scruples.

#### SOLUTION OF CHLORINATED POTASSA.

R. Carbonate of potassa                      one part  
Water    ten parts  
Dissolve, and pass a current of chlorine through the solution, till it is saturated, and preserve in well-stopped bottles. This is known as *Eau de Javelle*, and is principally used for bleaching, but may be employed as a disinfectant, like Labarraque's liquid.

*Guibourt.*

### POTASSII CYANURETUM.

#### CYANURET OF POTASSIUM.

R. Ferrocyanuret of potassium                      eight ounces  
Distilled water                      six fl. ounces  
Heat the ferrocyanuret moderately, until it becomes nearly white, and is deprived of its water of crystallization. Put the residue into an earthen retort, with the beak loosely stopped, and expose it to a red heat for two hours, or till gas ceases to be evolved. Withdraw it from the fire, close the orifice, and let remain till cold. Break the retort, remove the contents, reduce them to a coarse powder, introduce them in a bottle of the capacity of twelve fl. ounces, and add the distilled water; agitate occasionally for half an hour, throw it on a filter, evaporate the filtered solution rapidly to dryness, and keep in a closely stopped bottle.

*U. S. Ph.*

R. Ferrocyanuret of potassium, dried,                      eight ounces  
Pure carbonate of potassa                      three ounces  
Charcoal, in fine powder,                      one ounce  
Mix well, fuse in an iron vessel until effervescence ceases, and the fusion is tranquil, pour out the fused mass on an iron plate, and cool. Powder when cool, dissolve in alcohol (9.00), and separate and dry the crystals which form on cooling, and preserve them in well-closed bottles. *Wöhler.*

R. Ferrocyanuret of potassium, dried,                      eight parts  
Dried carbonate of potassa                      three parts  
Fuse together, when the mass is fluid, stir with a glass rod, until it becomes quite colorless, allow to settle, and pour out on a marble slab.

*Liebig.*

This contains one-eighth of cyanate of potassa.

#### PILLS OF CYANURET OF POTASSIUM.

R. Cyanuret of potassium  
Starch mixed with syrup of gum, each,                      half a grain  
Mix, and make a pill. Dose, one morning and evening. In spasmodic affections of the chest.

*Foy.*

#### SOLUTION OF CYANURET OF POTASSIUM.

R. Cyanuret of potassium                      one drachm  
Distilled water                      one ounce  
Dissolve.

*Magendie.*

R. Cyanuret of potassium                      twenty-two grains  
Diluted alcohol                      nine fl. drachms  
Dissolve. Equal to hydrocyanic acid containing one grain to a fl. drachm.

*Laming.*

#### MIXTURE OF CYANURET OF POTASSIUM.

R. Solution of cyanuret of potassium                      one fl. drachm  
Distilled water                      one pint  
Sugar                      one ounce and a-half  
Mix. A tablespoonful, night and morning, in pectoral affections.

*Magendie.*

R. Cyanuret of potassium                      four grains  
Distilled water                      two ounces  
Syrup of sugar                      three drachms  
Mix. A teaspoonful, every three or four hours, in asthma and chronic catarrh.

*Cadet.*

R. Tincture of castor  
Musk  
Nitrate of potassa, each,                      four grains  
Cyanuret of potassium                      two grains  
Linden-flower water                      eight fl. ounces  
Mix. To be taken in divided doses, in twenty-four hours, in chorea.

*Fouquier.*

R. Lettuce water                      two ounces  
Cyanuret of potassium                      half a grain to a grain and a-half  
Syrup of marsh mallows                      one ounce  
Mix. A tablespoonful, every two hours.

*Magendie.*

#### SYRUP OF CYANURET OF POTASSIUM.

R. Solution of cyanuret of potassium                      one drachm  
Simple syrup                      one pound  
Mix.

*Magendie.*

#### LOTION OF CYANURET OF POTASSIUM.

R. Cyanuret of potassium                      one to four grains  
Distilled water                      one ounce  
Mix.

*Foy.*

R. Cyanuret of potassium                      eight grains  
Distilled water                      one ounce  
Mix. To be applied by means of com-



presses, in cases of neuralgia, articular rheumatism, &c. *Foy.*

R. Cyanuret of potassium ten grains  
Emulsion of bitter almonds six ounces

Mix. As an application to chronic eruptions, attended with much itching.

*Cazenave.*

OINTMENT OF CYANURET OF POTASSIUM.

R. Cyanuret of potassium one to four grains

Lard one ounce

Mix. *Lombard.*

R. Cyanuret of potassium twelve grains

Oil of almonds two drachms

Cold cream two ounces

Mix. To be rubbed on the skin to relieve neuralgic pains. *Cazenave.*

POTASSII IODIDUM.

IODIDE OF POTASSIUM.

R. Iodine six ounces

Iron filings three ounces

Carbonate of potassa sufficient

Distilled water four pints

Mix the iodine with three pints of the water, and add the iron filings, stirring frequently for half an hour. Apply a gentle heat, and when the mixture assumes a greenish color, add gradually the carbonate of potassa, dissolved in half a pint of the water, until it ceases to produce a precipitate. Continue the heat for half an hour, and filter. Wash the residuum with half a pint of the distilled water, boiling hot and filter. Mix the filtered liquors, and evaporate, so that crystals may form. Pour off the liquid, and dry the crystals on bibulous paper. *U. S. Ph.*

R. Iodine sixteen ounces

Distilled water eight pints

Sulphuret of barium sufficient

Sulphate of potassa twelve ounces

Mix the iodine with the water, and gradually add the powdered sulphuret, until the solution becomes colorless, constantly stirring. Heat to the boiling point, add the sulphate of potassa, boil for a quarter of an hour, and filter. Evaporate to crystallization. *Mohr.*

R. Caustic potash four ounces

Iodine sufficient

Charcoal, in powder, one ounce

Water sufficient

Dissolve the potash in a pint of water, add the iodine in powder gradually, as long as the solution becomes colorless by stirring,

then add the charcoal, and evaporate to dryness in an iron pot. Cover this and heat to dull redness for half an hour, remove the mass from the pot, treat it with the smallest quantity of water necessary to dissolve the salt, filter, evaporate, and crystallize. The mother liquid yields more crystals by evaporation.

*Turner—modified.*

R. Iodine one hundred grains

Carbonate of potassa seventy-five grains

Water two drachms

Iron filings thirty grains

Mix, and heat slightly, then to redness; the resulting red powder is to be treated with water; filtered, and evaporated to dryness. *Pypues.*

Dose, two to fifteen grains, but has been given in still larger quantities.

PILLS OF IODIDE OF POTASSIUM.

R. Iodide of potassium five drachms

Distilled water six drachms

Crumb of bread sufficient

Mix, and make three hundred pills. In goitre, leucorrhœa, &c. *Pierquin.*

R. Iodide of potassium fifteen grains

Burnt sponge

Extract of bittersweet, each, five drachms

Distilled water sufficient

Mix, and make one hundred and eighty pills. Six, twice a-day, in scrofula and goitre. *Voght.*

LOZENGES OF IODIDE OF POTASSIUM.

R. Iodide of potassium one drachm

Sugar three ounces

Mucilage of tragacanth sufficient

Mix, and make lozenges of twelve grains. One to six, or more, a-day. *Giordano.*

R. Iodide of potassium seventy-five grains

Powdered coffee half a drachm

sugar four ounces

Mucilage of gum, made with

a strong infusion of coffee sufficient

Mix, and make three hundred lozenges. Each contains a quarter of a grain of iodide. In goitre, scrofula, &c. *Pierquin.*

SOLUTION OF IODIDE OF POTASSIUM.

R. Iodide of potassium thirty-six grains

Distilled water one ounce

Dissolve. Dose, ten minims. *Magendie.*

R. Iodide of potassium one drachm

Distilled water one ounce

Dissolve. Fifteen drops, gradually increased; three times a-day, in ovarian dropsy. *Elliotson.*



R. Iodide of potassium four ounces  
 Distilled water eight fl. ounces  
 Dissolve. Ten minims contain five grains  
 of iodide. *Guy's Hosp.*

#### INJECTION OF IODIDE OF POTASSIUM.

R. Iodide of potassium one to three grains  
 Distilled water one pint  
 Dissolve. To stimulate fistulous sinuses  
 in scrofulous patients. *Foy.*

#### MIXTURE OF IODIDE OF POTASSIUM.

R. Iodide of potassium one drachm  
 Lettuce water eight ounces  
 Mint water two drachms  
 Syrup of marsh mallow one ounce  
 Mix. *Foy.*

R. Iodide of potassium four drachms  
 Lettuce water eight ounces  
 Orange-flower water two drachms  
 Syrup of marsh mallow one ounce  
 Mix. *Foy.*

R. Iodide of potassium four drachms  
 Lettuce water eight ounces  
 Orange-flower water one drachm  
 Tincture of fox-  
 glove one to two drachms  
 Syrup of marsh mal-  
 low one ounce and a-half  
 Mix. *Magendie.*

In hypertrophy of the heart. A teaspoonful,  
 morning and evening.

R. Iodide of potassium two drachms  
 Distilled water sixteen ounces  
 Syrup two ounces  
 Mix. Two or three spoonfuls a-day.  
*Cazenave.*

R. Quassia  
 Gentian, each, one drachm  
 Boiling water sixteen fl. ounces  
 Macerate for an hour, strain, and add  
 Iodide of potassium thirty-six grains  
 Carbonate of potassa two drachms  
 Mix. A tablespoonful, three times a-day.  
*Cumming.*

R. Solution of iodide of  
 potassium fifteen drops  
 Medicinal hydrocya-  
 nic acid ten to twelve drops  
 Lettuce water four ounces  
 Syrup of marsh mallow one ounce  
 Mix. A teaspoonful every hour, in phthisis.  
*Magendie.*

R. Iodide of potassium two grains  
 Sulphate of magnesia half an ounce  
 Tartar emetic half a grain  
 Distilled water six ounces  
 Mix. A teaspoonful, three or four times  
 a-day, in scrofula. *Radius.*

#### SYRUP OF IODIDE OF POTASSIUM.

R. Iodide of potassium one drachm  
 Simple syrup one pound  
 Dissolve. *Cadet.*

R. Iodide of potassium twenty grains  
 Peppermint water two drachms  
 Simple syrup ten ounces  
 Mix. *Giordano.*

R. Iodide of potassium two drachms  
 Simple syrup six ounces  
 Mix. *Cazenave.*

#### IODIDE OF POTASSIUM AND SARSAPA- RILLA.

R. Iodide of potassium one drachm  
 Decoction of sarsaparilla two pints  
 Syrup of orange peel two ounces  
 Mix. To be taken in glassfuls, during the  
 day. *Magendie.*

#### OINTMENT OF IODIDE OF POTASSIUM.

R. Iodide of potassium half a drachm  
 Lard one ounce and a-half  
 Mix. *Magendie.*

The strength of this ointment may vary much,  
 from containing a twenty-fourth of the salt, to an  
 eighth or more, according as it may be required.  
 In goitre, scrofulous swellings, &c.

R. Iodide of potassium half a drachm  
 Solution of potassa two drops  
 Lard one ounce and a-half  
 Wax three drachms  
 Mix. *Coindet.*

R. Spermaceti half an ounce  
 Olive oil six drachms  
 White wax two drachms  
 Iodide of potassium four scruples  
 Oil of lemons  
 roses, each, three drops  
 Mix. This ointment does not change  
 color by keeping. *Kallhofert.*

R. Iodide of potassium one drachm  
 Magnesia six grains  
 Rub together with a few drops of water,  
 and add  
 Rose water ointment one ounce  
 Mix well. *Hamb. Ph.*

R. Iodide of potassium  
 Carbonate of soda, each,  
 half a drachm  
 Rose water ointment six drachms  
 Mix well. In chronic enlargement of the  
 testicle. *Walthen.*

#### OINTMENT OF IODIDE OF POTASSIUM AND OPIUM.

R. Iodide of potas-  
 sium one drachm and a-half  
 Lard one ounce and a-half



Laudanum one fl. drachm  
 Mix. *A. T. Thomson.*  
 R. Iodide of potassium half a drachm  
 Extract of opium ten grains  
 Simple cerate one ounce  
 Mix. As a dressing to cancerous ulcers.

OINTMENT OF IODIDE OF POTASSIUM  
 AND MERCURY.

R. Iodide of potassium twelve to twenty grains  
 Mercurial ointment half an ounce  
 Mix. In various diseases of the skin.  
*Blarius.*

R. Ointment of iodide of potassium  
 Mercurial ointment  
 Oil of henbane, each, two ounces  
 juniper one drachm  
 Mix. *Schonlein.*

When iodide of potassium and mercurial ointment are mixed, the globules soon appear, but if the iodide be artificially dried, previously, a uniform ointment is produced.

R. Iodide of potassium three parts  
 Iodide of mercury two parts  
 Camphor two parts  
 Galen's cerate thirty-two parts  
 Mix. Used as friction in acute hydrocephalus, in doses of a half a drachm to a drachm.  
*Golfin.*

OINTMENT OF IODIDE OF POTASSIUM  
 AND MORPHIA.

R. Iodide of potassium fifteen grains  
 Muriate of morphia half a drachm  
 Lard ten ounces  
 Mix. In painful tumors of the breasts, to be applied twice a-day.  
*Chomel.*

LINIMENT OF IODIDE OF POTASSIUM.

R. Iodide of potassium one ounce  
 Alcohol four ounces  
 Dissolve.  
 R. Animal soap one ounce and a-half  
 Alcohol four fl. ounces  
 Dissolve. Add the two solutions together, aromatize with a few drops of oil of lavender, and pour into wide-mouthed bottles.  
*Boudet.*

R. White soap seven drachms  
 Diluted alcohol two fl. ounces  
 Dissolve by a gentle heat, and whilst warm, add  
 Iodide of potassium four drachms  
 dissolved in  
 Diluted alcohol four fl. drachms  
*Beasley.*

R. Iodide of potassium four drachms  
 Diluted alcohol two ounces  
 Dissolve.

R. Curd soap six drachms  
 Diluted alcohol two ounces  
 Dissolve by a gentle heat, and add the solution of iodide of potassium, and aromatize.  
*Foy.*

These are used as liniments in goitre.

R. White soap ten drachms  
 Oil of almonds ten drachms  
 Iodide of potassium one drachm  
 Water one drachm  
 Dissolve the iodide in the water, and add it to the soap and oil, melted together.  
*Guibourt.*

SOAP OF IODIDE OF POTASSIUM.

R. Camphor one drachm  
 Tincture of benzoin three drachms  
 Add to the solution, triturating well,  
 Iodide of potassium two drachms  
 Goulard's extract four drachms  
 Add to the mixture  
 Oil of almonds thirty-four drachms  
 Solution of potassa two ounces  
 Essence of lavender half a drachm  
 Mix well. As an application to chilblains.  
*Cadet.*

POTASSII BINIODIDUM.

BINIODIDE OF POTASSIUM.

R. Iodide of potassium twenty parts  
 Iodine six parts  
 Mix, and triturate together till the mass is homogeneous, and of a dark brown color.  
*Giordano.*

To have a perfectly saturated salt, iodine in excess is to be added to a solution of the iodide until no more is dissolved. In the following preparations, there is a mixture of iodide and biniodide.

IODURETTED WATER.

R. Iodide of potassium six grains  
 Iodine one grain  
 Water two pints  
 Dissolve. To be used instead of common water at meals.  
*Magendie.*

IODURETTED WATER, (LUGOL'S.)

R. Iodide of potassium one grain and a-half  
 Iodine three-quarters of a grain  
 Distilled water eight ounces  
 Dissolve.  
 R. Iodide of potassium two grains  
 Iodine one grain  
 Distilled water eight ounces  
 Dissolve.  
 R. Iodide of potassium two grains and a-half  
 Iodine one grain and a quarter



Distilled water eight ounces  
Dissolve. The first is for young children, to be taken in divided doses, in three days; the second for those of more advanced age, in two days; the third for adults, in one day. *Guibourt.*

R. Iodide of potassium six grains  
Iodine three grains  
Water sixteen ounces

Dissolve. In poisoning by the vegetable alkaloids, in wineglassfuls, after the stomach has been emptied. *Bouchardat.*

R. Iodide of potassium ten grains  
Iodine five grains  
Distilled water twenty fl. ounces

Dissolve. Dose, two to six fl. drachms. *Lond. Ph.*

#### COMPOUND SOLUTION OF IODIDE OF POTASSIUM.

R. Iodide of potassium one ounce and a-half  
Iodine six drachms  
Distilled water one pint

Dissolve. Dose, six drops, three times a-day, in sweetened water. *U. S. Ph.*

#### IODURETTED BATH.

R. Iodide of potassium four scruples  
Iodine two scruples  
Water ten ounces

Dissolve. To be added to a sufficient quantity of water. For children.

R. Iodide of potassium six drachms  
Iodine three drachms  
Water twenty ounces

Dissolve. To be added to a sufficient quantity of water. For adults. *Foy.*

#### COLLYRIUM OF BINIODIDE OF POTASSIUM.

R. Iodide of potassium twenty-four grains  
Iodine one to two grains  
Rose water six ounces

Dissolve. To be used four times a-day, in scrofulous ophthalmia. *Magendie.*

#### IODURETTED INJECTION.

R. Iodide of potassium four grains  
Iodine two grains  
Distilled water sixteen ounces

Dissolve.

R. Iodide of potassium eight grains  
Iodine four grains  
Distilled water sixteen ounces

Dissolve. To stimulate fistulous sinuses. *Guibourt.*

#### SYRUP OF BINIODIDE OF POTASSIUM.

R. Biniodide of potassium twenty grains  
Peppermint water two drachms

Simple syrup two ounces  
Dissolve the biniodide in the peppermint water, and add the solution to the syrup. *Giordano.*

#### COMPOUND LOTION OF IODIDE OF POTASSIUM.

R. Iodide of potassium two to four grains  
Iodine one to two grains  
Distilled water eight fl. ounces

Dissolve. As an application in scrofulous ophthalmia, fistulas, &c. *Laugel.*

R. Iodide of potassium two ounces  
Iodine one ounce  
Distilled water three ounces

Dissolve. *Guibourt.*

R. Iodide of potassium one ounce  
Iodine half an ounce  
Distilled water six ounces

Dissolve. *Soubéiran.*

These two preparations are used to touch the eyelids in scrofulous ophthalmia.

#### CAUSTIC SOLUTION OF BINIODIDE OF POTASSIUM.

R. Iodide of potassium  
Iodine, each, one ounce  
Distilled water two ounces

Mix. *Guibourt.*

R. Iodide of potassium  
Iodine  
Distilled water, each, one ounce

Mix. *Cadet.*

These are used to touch the surfaces of scrofulous ulcers.

#### COMPOUND TINCTURE OF IODIDE OF POTASSIUM.

R. Iodide of potassium one ounce  
Iodine half an ounce  
Alcohol one pint

Dissolve. *U. S. Ph.*

R. Iodide of potassium two ounces  
Iodine one ounce  
Alcohol two pints (imp.)

Dissolve. *Lond. Ph.*

Dose, from fifteen to thirty drops.

#### MIXTURE OF BINIODIDE OF POTASSIUM.

R. Iodide of potassium four drachms  
Iodine two grains  
Orange-flower water

Mint water, each, three ounces

Mix. A dessertspoonful, three times a-day, in epilepsy. *Magendie.*

R. Iodide of potassium half a drachm  
Iodine half a grain  
Syrup of poppies half a fl. ounce  
Distilled water half a pint



Mix. Two tablespoonfuls, three times a-day, in syphilis combined with scrofula.  
*Tyrrell.*

OINTMENT OF BINIODIDE OF POTASSIUM.

R. Iodide of potassium one drachm  
Iodine half a drachm  
Alcohol one fl. drachm  
Lard two ounces

Rub the iodide and iodine with the alcohol, and then with the lard, until they are thoroughly mixed.  
*U. S. Ph.*

R. Iodide of potassium twenty-four grains  
Iodine twelve grains  
Lard two ounces

Mix. *Lugol.*

R. Iodide of potassium two drachms  
Iodine eighteen grains  
Lard two ounces

Mix. *Lugol.*

R. Iodide of potassium two drachms and a-half  
Iodine twenty-one grains  
Lard two ounces

Mix. *Lugol.*

R. Iodide of potassium two drachms and a-half  
Iodine twenty-four grains  
Lard two ounces

Mix. In scrofulous ulcers, &c. *Lugol.*

OINTMENT OF BINIODIDE OF POTASSIUM AND OPIUM.

R. Iodide of potassium one drachm  
Iodine fifteen grains  
Laudanum of Rousseau two drachms  
Lard two ounces

Mix. As an application to scrofulous ulcers.  
*Foy.*

PLASTER OF BINIODIDE OF POTASSIUM.

R. Iodide of potassium  
Iodine, each, ten grains to one scruple  
Mercurial or soap plaster two ounces

Mix. As an application to syphilitic and gouty swellings.  
*Ebers.*

POTASSII HYDRARGYRO-  
IODIDUM.

HYDRARGYRO-IODIDE OF POTASSIUM.

R. Iodide of potassium eight grains  
Dissolved in ten or fifteen minims of water

Red iodide of mercury eleven grains  
Mix. This contains twenty-grains of the hydrargyro-iodide, and is not decomposed by water.  
*Channing.*

R. Bichloride of mercury one equivalent  
Dissolve in a solution of

Iodide of potassium four equivalents  
Evaporate to dryness, treat with alcohol, and evaporate to crystallization.

*Amb. Smith.*

Dose, one-twelfth of a grain, three times a-day; in many cases, a much smaller quantity is required to be given.

SOLUTION OF HYDRARGYRO-IODIDE OF  
POTASSIUM.

R. Iodide of potassium three grains and a-half  
Red iodide of mercury four grains and a-half  
Distilled water one fl. ounce

Dissolve first, the iodide of potassium, and then the mercurial salt in the water. The compound salt amounts to about eight grains. The dose is from two to five drops, three times a-day, much diluted, in dyspepsia, enlargement of the spleen, dropsy, &c.  
*Channing.*

R. Iodide of potassium  
Red iodide of mercury, each, eight grains  
Distilled water eight ounces

Mix. Dose, two fl. drachms and upwards, in the twenty-four hours.  
*Puche.*

R. Hydrargyro-iodide of potassium twelve grains  
Water sixteen fl. ounces  
*Limonium Lamothe.*

TINCTURE OF HYDRARGYRO-IODIDE OF  
POTASSIUM.

R. Hydrargyro-iodide of potassium one grain  
Diluted alcohol one fl. ounce  
Dissolve. Ten drops, three times a-day.  
*Channing.*

PILLS OF HYDRARGYRO-IODIDE OF  
POTASSIUM.

R. Red iodide of mercury  
Iodide of potassium, each, eight grains  
Sugar of milk sixty-four grains  
Mucilage of gum Arabic sufficient  
Mix, and make thirty-two pills.

R. Iodide of potassium  
Red iodide of mercury, each, six grains  
Opium twelve grains  
Mix, and make twenty-four pills. *Mialhe.*

OINTMENT OF HYDRARGYRO-IODIDE OF  
POTASSIUM.

R. Red iodide of mercury seven grains



Iodide of potassium two scruples  
Lard one ounce  
Mix. To be applied to tumors, two or three times a-day. *Hildreth.*

R. Hydrargyro-iodide of potassium one scruple  
Lard one ounce  
Mix. *Lamothé.*

### POTASSII SULPHO-CYANURETUM.

SULPHO-CYANURET OF POTASSIUM.  
R. Ferro-cyanuret of potassium three parts  
Sulphur one part  
Pack in a crucible, heat to redness, for an hour, treat with alcohol, and evaporate to crystallization. *Van Mons.*

R. Prussian blue three parts  
Sulphuret of potassa one part  
Put the mixture into a covered crucible, and heat to a dull red for half an hour, treat with alcohol, filter the solution, evaporate, and crystallize. *Van Mons.*

R. Digest an aqueous solution of cyanuret of potassium with sulphur, of which it will take up one-third. Filter, and evaporate. *Beasley.*

### POTASSII SULPHURETUM.

SULPHURET OF POTASSIUM.  
R. Sulphur one ounce  
Carbonate of potassa two ounces  
Rub the carbonate of potassa, previously dried with the sulphur, melt the mixture in a covered crucible over the fire; then pour it out, and, when cold, put it in a bottle, which is to be well stopped. *U. S. Ph.*

A stimulant, expectorant, and diaphoretic, in small doses, poisonous in large; used externally in many cutaneous diseases. Dose, from two to ten grains, several times a-day.

### HYPOSULPHITED SULPHURET OF POTASSIUM.

R. Sublimed sulphur one part  
Solution of potassa three parts  
Mix, and heat on a sand-bath, until it marks 39° B. *Guibourt.*

It consists of three parts of sulphuret and one of hyposulphite.

FERRO-SULPHURET OF POTASSIUM.  
R. Carbonate of potassa  
Sulphur, each, one ounce  
Black oxide of iron two drachms  
Mix, and melt in a crucible, pour out on

a slab of marble, break in pieces, and keep in a well-closed bottle. Dose, three to four grains. *Ferrara Ph.*

### SULPHURET OF POTASSIUM AND CREMOR TARTAR.

R. Sulphuret of potassium four scruples  
Cremor tartar four drachms  
Mix, and divide into twenty-four powders. Dose, one every four hours, in a glass of sweetened water. Said to be efficacious in mercurial salivation. *Bories.*

### BOLUS OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium three grains  
Conserve of elder berries sufficient  
Mix. Make six boluses; one every three hours. In mercurial salivation. *Brera.*

R. Sulphuret of potassium six grains  
Black oxide of iron three grains  
Extract of quassia ten grains  
Burnt sponge sufficient  
Mix, and make a bolus. One, morning and evening, in goitre and glandular affections. *Phæbus.*

### PILLS OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium one drachm  
Extract of liquorice sufficient  
Mix, and make thirty pills. Two to five, several times a-day.

### COMPOUND PILLS OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium  
Gum ammoniac  
Extract of dandelion, each, one drachm  
Soap  
Rhubarb, each, half a drachm  
Opium three grains  
Mix, and make pills of two grains. *Radius.*

### PILLS OF SULPHURET OF POTASSIUM AND CHARCOAL.

R. Sulphuret of potassium one drachm  
Extract of blessed thistle  
Powdered charcoal, each, two grains  
Mix, and make pills of two grains. Fifteen to twenty, four times a-day, in phthisis. *Augustin.*

### ELECTUARY OF SULPHURET OF POTASSIUM.

R. Butter of cocoa two drachms  
Oil of almonds half an ounce  
Melt together, and add, triturating well  
Sulphuret of potassium ten grains  
Sugar three drachms  
As an alterative for children, in teaspoonful doses. *Phæbus.*



SYRUP OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium eight grains  
Distilled water sixteen grains  
Dissolve, add to it  
Simple syrup one ounce  
*Par. Cod.*

A teaspoonful, for croup in children.

COMPOUND SYRUP OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium one ounce  
Fennel water sixteen ounces  
Simple syrup thirty-eight ounces  
Mix. Once much celebrated as Willis' bechic syrup, in croup, &c. *Giordano.*

SYRUP OF HYPOSULPHITED SULPHURET OF POTASSIUM.

R. Hyposulphited sulphuret of potassium sixteen grains  
Simple syrup one ounce  
Mix. *Guibourt.*

SOLUTION OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium one part  
Water sufficient  
to make a solution of 30° B. *Soubéiran.*

WATER OF SULPHURET OF POTASSIUM.

R. Washed sulphur one part  
Water of caustic potassa eleven parts  
Boil for ten minutes, and filter. Keep in well-stopped bottles. *Dub. Ph.*

This is not analogous to a solution of sulphuret of potassium, as it contains much hyposulphite of potassa. Dose, ten minims to one fl. drachm, three times a-day.

TINCTURE OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium four ounces  
Diluted alcohol sixteen ounces  
Digest for twenty-four hours, and strain. *Quincy.*

R. Carbonate of potassa one ounce  
Sulphur two ounces  
Melt together, and digest with  
Alcohol one pound  
Filter. *Saxon. Ph.*

Recommended in itch, in doses of sixty drops.

LOTION OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium one ounce  
Water half a pint  
Dissolve. As a wash in herpetic and other cutaneous eruptions. *Ellis.*

R. Sulphuret of potassium one to two parts  
Water sixteen parts  
Dissolve. *Foy.*

COMPOUND LOTION OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium half an ounce

Soap one ounce  
Alcohol four fl. ounces  
Tincture of myrrh half a fl. ounce  
Lime water one pint  
Mix. As an application in tinea capitis. *Ellis.*

R. Sulphuret of potassium one to two ounces  
Water one pint  
Dissolve.

R. Muriatic acid one fl. ounce to two fl. ounces  
Distilled water two pints  
Mix an ounce of each solution with four ounces of warm water. As an application in psora. *Alibert.*

R. Sulphuret of potassium three drachms  
Soap one drachm and a-half  
Lime water seven and a-half fl. ounces  
Diluted alcohol two fl. ounces  
Mix. *Burns.*

Known as Barlow's lotion, and used in various cutaneous diseases.

R. Sulphuret of potassium two drachms  
Soap two drachms and a-half  
Lime water seven fl. ounces  
Alcohol one fl. drachm  
Mix. *Bielt.*

As a lotion in porrigo.

R. Sulphuret of potassium four ounces  
Water one pint and a-half  
Sulphuric acid half an ounce  
Mix. As a lotion in itch, to be used morning and evening. *Dupuytren.*

SULPHURET OF POTASSIUM BATH.

R. Sulphuret of potassium four ounces  
Water twelve ounces  
Dissolve, and mix with the water of a bath. *Beral.*

R. Sulphuret of potassium one ounce  
Common salt two ounces  
Carbonate of soda four drachms  
Leaves of sage one to two handfuls  
Water six quarts

Boil for twenty-four hours. In fomentations, douches, &c., in spina ventosa and scrofula. *St. Marie.*

R. Sulphuret of potassium four ounces  
Water two hundred pints  
Mix, and add  
Glue two pounds  
Dissolved in ten pints of boiling water. *Foy.*

INJECTION OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium one drachm



Distilled water eight to twelve ounces  
Dissolve. In gonorrhœa. *Wedekind.*

#### MIXTURE OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium one drachm and a-half  
Bicarbonate of potassa ten grains  
Oil of peppermint one to two drops  
Syrup of orange peel two ounces  
Mucilage one ounce

Mix. A teaspoonful, every two hours, as an alterative in scrofula. *Lockstaedt.*

R. Sulphuret of potassium one scruple  
Carbonate of potassa ten grains  
Peppermint water two ounces  
Syrup of saffron one ounce

Mix. A spoonful, every two hours, in cutaneous diseases. *Phœbus.*

R. Sulphuret of potassium one scruple  
Water one ounce  
Syrup of cinnamon two drachms

Mix. To be taken in three doses, in mercurial disease. *Dzondi.*

R. Sulphuret of potassium one scruple  
Kermes mineral ten grains  
Syrup of senega two ounces  
Anisated ammonia one scruple

Mix. A spoonful, every hour in croup. *Hagen.*

R. Sulphuret of potassium  
Sugar, each, one drachm  
Water one pint

Mix. Has been advised as an antidote in poisoning by arsenic. *Augustin.*

#### OINTMENT OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium  
Carbonate of soda, each, three drachms  
Lard three ounces

Mix. As an application in tinea capitis. *Alibert.*

R. Soap one pound  
Water one ounce

Soften by means of a water-bath, and add  
Sulphuret of potassium three ounces  
Oil of pinks two pounds

Mix well. *Jadelot.*

R. Sulphuret of potassium six ounces  
Water two ounces

Dissolve, and add  
Soap two pounds  
Oil of pinks four pounds  
thyme two drachms

Mix. *Giordano.*

R. Powdered soap two ounces  
Sulphuret of potassium  
Water, each, one ounce

Mix well, and add, gradually,  
Olive oil five ounces  
This liniment alters rapidly when exposed to the air. *Beral.*

#### COMPOUND PLASTER OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium  
Powdered hemlock, each, two drachms  
Camphor  
Turpentine, each, four drachms  
Soap half a drachm  
Yellow wax one ounce  
Simple plaster four ounces

Mix well. As a dressing to tumefied lymphatic glands. *Bories.*

#### SOAP OF SULPHURET OF POTASSIUM.

R. Sulphuret of potassium one part  
Boiling water two parts

Dissolve, and add  
Yellow wax one part

Evaporate, stirring continually. *Bavar. Ph.*

#### POTASSA.

##### CAUSTIC POTASSA.

R. Solution of potassa one gallon  
Evaporate quickly, in a clean iron vessel, over the fire, till ebullition ceases and the potassa melts. Pour into moulds, and keep in well-stopped bottles. *U. S. Ph.*

A powerful escharotic, used to form issues and in opening abscesses.

##### SOLUTION OF CAUSTIC POTASSA.

R. Caustic potassa one drachm and a-half  
Distilled water two fl. ounces  
Dissolve. As a rubefacient in tetanus, to be applied to the spine. *Hartshorne.*

##### POTASSA WITH LIME.

R. Caustic potassa  
Lime, each, one ounce  
Rub them together, and keep in a well-stopped bottle. *Lond. Ph.*

Used as the above, but is slower in producing an effect. It is to be made into a paste with a little alcohol.

##### SOLUTION OF POTASSA.

R. Carbonate of potassa one pound  
Lime half a pound  
Boiling distilled water one gallon  
Dissolve the carbonate of potassa in half a gallon of the water, pour a little of the water on the lime, and, when slaked, add the remainder. Mix the hot liquors, and boil for ten minutes, stirring continually, set the mixture aside in a covered vessel,



till clear. Decant, and keep in well-stopped bottles of green glass. *U. S. Ph.*

R. Caustic potassa one scruple  
Water one fl. ounce  
Dissolve. This solution very nearly represents the above. *W. Procter.*

LITHONTRIPTIC SOLUTION OF CAUSTIC POTASSA.

R. Calcined carbonate of potassa half a pound  
Burnt oyster shells one pound and a-half  
Boiling water eight pints and a-half  
Mix, and let rest for twenty-four hours, and then filter. One fl. drachm three times a-day, in flaxseed tea, adding fifteen grains of magnesia to each dose.

*Saunders.*

SAVIARD'S STIMULANT LOTION.

R. Caustic potassa two drachms  
Camphor two scruples  
Sugar two ounces  
Water two pints  
Mix. As an application to indolent ulcers. *Foy.*

COLLYRIUM OF CAUSTIC POTASSA.

R. Caustic potassa one grain  
Distilled water one ounce  
Dissolve. A drop or two to be introduced into the eye, to remove specks on the cornea. *Gimbernati.*  
R. Caustic potassa one grain  
Extract of opium four grains  
Distilled water four ounces  
Dissolve, and filter. In chronic ophthalmia. *Hamb. Ph.*

MIXTURE OF CAUSTIC POTASSA.

R. Tincture of caustic potassa half an ounce  
Volatile tincture of guaiacum two drachms  
Laudanum half a drachm  
Mix. Twenty drops, three times a-day, in gout. *Græfe.*

INJECTION OF CAUSTIC POTASSA.

R. Caustic potassa half a grain  
Chamomile water one ounce and a-half  
Laudanum five drops  
Mix. As an injection into the ear, in deafness. *Rust.*  
R. Caustic potassa two grains  
Distilled water one ounce  
Dissolve. As an injection, at the commencement of gonorrhœa. *Girtanner.*

POTASSÆ ACETAS.

ACETATE OF POTASSA.

R. Acetic acid one pint  
Carbonate of potassa sufficient  
to saturate; to be added gradually; filter; evaporate by means of a sand-bath to dryness; keep in closely-stopped bottles. *U. S. Ph.*

A diuretic, in doses of a scruple to a drachm; as a laxative, in doses of two or three drachms.

SCILLITIC ACETATE OF POTASSA.

R. Carbonate of potassa half an ounce  
Vinegar of squills twelve fl. ounces  
Evaporate to the consistence of honey, and add  
Alcohol (.842) six fl. ounces  
Decant, after digesting for a few days. Dose, forty to sixty drops, in some demulcent fluid, in dropsy and asthma. *Keup.*

BOLUS OF ACETATE OF POTASSA.

R. Acetate of potassa one scruple  
Conserve of pot marigold sufficient  
Mix. To be taken in the morning, for some days, to retard the secretion of milk. *St. Marie.*

COMPOUND PILLS OF ACETATE OF POTASSA.

R. Acetate of potassa  
Gum ammoniac  
Pills of aloes and myrrh, each, one part  
Soap two parts  
Simple syrup sufficient  
Mix, and make pills of five grains. *Van Mons.*

CONSERVE WITH ACETATE OF POTASSA.

R. Acetate of potassa half an ounce  
Sulphate of soda one drachm  
Juice of scurvy grass  
fumitory  
dandelion, each, two ounces  
Sugar sufficient  
Mix, and form conserve. A teaspoonful, two or three times a-day, in obstructions of the bowels. *Bories.*

R. Acetate of potassa  
Powdered burnt sponge, each, two drachms  
Calomel twelve grains  
Sulphuret of antimony  
one drachm and a-half  
Jalap sixteen grains  
Sulphur one grain



Simple syrup sufficient  
 Mix. Dose, twelve to twenty-four grains,  
 twice a-day, in scrofula. *Baumes.*

#### LIQUID ACETATE OF POTASSA.

R. Acetate of potassa one part  
 Distilled water two parts  
 Dissolve. *Amster. Ph.*

R. Acetate of potassa at will  
 Expose to the action of the air, till it de-  
 liquescs, and filter. *Pideret.*

Dose, forty to eighty drops, in an appropriate  
 vehicle, as a diuretic.

#### MIXTURE WITH LIQUID ACETATE OF POTASSA.

R. Liquid acetate of potassa two ounces  
 Extract of Peruvian bark two drachms  
 cascarilla one drachm

Elixir proprietatis one ounce  
 Mix. In intermittent fevers, complicated  
 with obstruction and dropsy; in doses of  
 thirty drops, every two hours, during the  
 apyrexia. *Bories.*

R. Liquid acetate of potassa four ounces  
 Extract of centaury two drachms  
 Sulphuric ether twenty drops  
 Syrup of kermes two ounces  
 Mix. In dropsy and obstructions. In  
 doses of thirty drops. *Saunders.*

#### TINCTURE OF ACETATE OF POTASSA.

R. Acetate of potassa two ounces  
 Alcohol four ounces  
 Digest for some days, and filter. Dose,  
 thirty to sixty drops. *Bruns. Ph.*

#### MIXTURE OF ACETATE OF POTASSA.

R. Acetate of potassa  
 Extract of dandelion  
 hemlock, each, half an ounce

Fennel water six ounces  
 Syrup of marsh mallow one ounce  
 Mix. Two teaspoonfuls, every four hours,  
 in jaundice. *Quarin.*

R. Carbonate of potassa fifteen grains  
 Distilled vinegar sufficient  
 to dissolve; add to the solution

Water half an ounce  
 Mint water one ounce  
 Pimento water  
 Vinegar of squill, each, one drachm  
 Simple syrup two drachms  
 Mix. Dose, half a drachm to a drachm,  
 every six hours, in dropsy. *Milman.*

R. Arnica root one drachm  
 Boiling water sufficient  
 to obtain four ounces of infusion; add  
 Carbonate of potassa, saturated  
 with vinegar of squill, half a drachm

Syrup of orange flowers half an ounce  
 Mix. A spoonful every two hours, in  
 dropsy. *Wendt.*

R. Acetate of potassa  
 Oxymer of squill, each, one drachm  
 Linden water four drachms  
 Wine of opium fifteen drops  
 Syrup of marsh mallow one ounce  
 Mix. In spoonful doses, as a diuretic  
 and sedative. *Pierquin.*

#### POTASSÆ ARSENITIS LIQUOR.

##### ARSENICAL, OR FOWLER'S SOLUTION.

R. Arsenious acid in small fragments  
 Pure carbonate of  
 potassa, each, sixty-four grains  
 Distilled water sufficient  
 Compound spirit of  
 lavender half a fl. ounce

Boil the arsenious acid and carbonate of  
 potassa with twelve fl. ounces of the water,  
 in a glass vessel, till the acid is entirely  
 dissolved. To the solution, when cold,  
 add the spirit of lavender, and afterwards,  
 sufficient distilled water to make it fill  
 exactly the measure of a pint. *U. S. Ph.*

Used in the same cases as arsenious acid,  
 especially in intermittent fever. Each fl. drachm  
 contains half a grain of arsenious acid. Dose,  
 about ten drops, two or three times a-day.

#### POTASSÆ ARSENIAS. ARSENIATE OF POTASSA.

R. White arsenic  
 Nitre, each, one ounce  
 Pulverize separately, mix, and introduce  
 into an earthenware retort; heat to red-  
 ness, as long as nitrous fumes are evolved,  
 let cool, dissolve the residuum in four pints  
 of boiling distilled water, evaporate, and  
 crystallize. *Guibourt.*

Dose, one-sixteenth to one-eighth of a grain.

##### MIXTURE OF ARSENIATE OF POTASSA.

R. Arseniate of potassa one fifth of a grain  
 Mint water three ounces  
 Simple syrup half an ounce  
 Mix. In spoonful doses, in the apyrexia  
 of intermittents. *Foy.*

#### POTASSÆ BORAS. BORATE OF POTASSA.

R. Boracic acid six parts



Bicarbonate of potassa five parts  
Mix, and heat to redness in a crucible,  
dissolve the residuum in water, filter, and  
evaporate to dryness. *Beasley.*

# POTASSÆ BOROTARTRAS.

## BOROTARTRATE OF POTASSA.

R. Powdered bitartrate of  
potassa four ounces  
Boracic acid one ounce  
Water three pints

Mix, and put in a silver basin, and boil  
till most of the water is evaporated, and  
continue the evaporation by a gentle heat,  
constantly stirring. When the matter has  
become thick, take it up in portions, flat-  
ten them, and dry in a stove, reduce them  
to powder, and keep in well-closed bottles.

*Par. Cod.*

R. Borax two hundred and fifty parts  
Bitartrate of potassa seventy-five parts  
Dissolve in boiling water, evaporate until  
a portion dropped on a cold slab solidifies.  
Powder in a warm mortar, and keep in  
well-stopped bottles. *Kaepeler.*

R. Bitartrate of potassa twelve ounces  
Borate of soda six ounces  
Tartaric acid three drachms  
Dissolve in water, clarify with white of  
egg, and proceed as directed by the Paris  
Codex. *Cambranae.*

This is known as soluble cream of tartar. It  
is purgative in about the same dose as cream of  
tartar.

## POWDER OF BOROTARTRATE OF POTASSA.

R. Cream of tartar one ounce  
Borax three drachms  
Sugar two ounces

Mix, and divide into three powders; one  
to be taken, in a glass of water, every half  
hour. *Pierquin.*

## COMPOUND PILLS OF BOROTARTRATE OF POTASSA.

R. Borotartrate of  
potassa one drachm and a-half  
Extract of buckbean two ounces  
Seneka snakeroot two drachms  
Colchicum one drachm and a-half  
Mix, and make pills of two grains. Dose,  
six to twelve every two hours, as a hydra-  
gogue purgative. *Augustin.*

## SOLUTION OF BOROTARTRATE OF POTASSA.

R. Cream of tartar  
Borax, each, one drachm

Boiling water one pint  
Dissolve. *Bories.*

## MIXTURE OF BOROTARTRATE OF POTASSA.

R. Borotartrate of  
potassa four to six drachms

Juniper water  
Parsley water, each, three ounces

Dissolve, and add  
Nitric ether two drachms

Syrup of poppies one ounce

Mix. Two spoonfuls every two hours, as  
a diuretic. *Niemann.*

R. Borotartrate of potassa one ounce

Mint water eight ounces

Nitric ether two drachms

Oxymel of squill one ounce

Mix. Two spoonfuls, every two hours,  
as a diuretic. *Selig.*

R. Borotartrate of potassa half an ounce

Tartar emetic one grain

Water four ounces

Mix. In spoonful doses, in congestion of  
the head. *Radius.*

## BOROTARTRATE OF POTASSA AND MAG- NESIA.

R. Borotartrate of potassa one part

Carbonate of mag-

nesia one-fourth part

Water six parts

Dissolve the borotartrate in the water, in a  
silver basin, saturate with the magnesia,  
evaporate carefully, till reduced to a tena-  
cious paste, which divide into small masses,  
and dry in a stove. *Mailliere Renault.*

## MIXTURE OF BOROTARTRATE OF POTASSA AND MAGNESIA.

R. Borotartrate of potassa  
and magnesia thirty parts

Citric acid two parts

Syrup of lemon sixty parts

Warm water three hundred parts

Dissolve. *Mailliere Renault.*

# POTASSÆ CARBONAS.

## CARBONATE OF POTASSA.

R. Impure carbonate  
of potassa three pounds

Water two pints and a-half

Dissolve the impure carbonate in the water,  
filter, pour into a clean iron vessel, and  
evaporate over a gentle fire, till the solu-  
tion thickens, then remove from the fire,  
and stir constantly, with an iron spatula,  
till the salt granulates. *U. S. Ph.*



Used as an antacid, and diuretic, &c. Dose, from five to twenty grains.

#### PURE CARBONATE OF POTASSA.

R. Bitartrate of potassa two pounds  
Nitrate of potassa one pound  
Rub them separately into powder, mix, and throw the mixture into a brass vessel, heated nearly to redness, from the residue prepare as directed for the carbonate.

*U. S. Ph.*

R. Bicarbonate potassa at will  
Put it in an iron crucible, heat gradually to redness for half an hour; when cold, remove the carbonate, and if discolored by the iron vessel, dissolve in water, filter, and evaporate to dryness in a porcelain crucible.

*W. Procter.*

#### SOLUTION OF CARBONATE OF POTASSA.

R. Carbonate of potassa one pound  
Distilled water twelve fl. ounces  
Dissolve, and filter.

*U. S. Ph.*

Dose, ten minims to a fl. drachm, properly diluted.

#### POWDER OF CARBONATE OF POTASSA.

R. Carbonate of potassa ten grains  
Powdered chamomile  
myrrh, each,  
twenty-four grains

Mix. To be taken every hour, in the dyspepsia of intermittents.

*Foy.*

R. Carbonate of potassa one part  
Powdered gum Arabic eight parts  
Triturate together for a long time. Dose, half a drachm to a drachm, dissolved in water, in engorgement of the bowels.

*Guibourt.*

#### COLLYRIUM OF CARBONATE OF POTASSA.

R. Carbonate of potassa two to six grains  
Water one ounce

Mix. To be dropped in the eye every two to four hours, in specks on the cornea.

*Himley.*

R. Carbonate of potassa twenty-five grains  
Veal broth, filtered, sufficient  
Celandine water two ounces  
Digest for twenty-four hours on a sand-bath, filter, and add

Tincture of aloes twenty-four drops  
In specks on the cornea.

*Borès.*

#### TINCTURE OF CARBONATE OF POTASSA.

R. Carbonate of potassa one part  
Calcine it with a strong heat in a crucible for two hours, pour it in a heated mortar, pulverize it rapidly, and pour on the hot powder

Alcohol four parts

Digest for fifteen days in a stove, often agitating, and filter.

*Guibourt.*

This is a weak alcoholic solution of caustic potassa.

#### COLLUTORY OF CARBONATE OF POTASSA.

R. Alcohol four ounces  
Essence of mint twenty drops  
roses eight drops

Cochineal

Carbonate of potassa, each, ten grains  
Macerate for twenty-four hours, and filter.

Much esteemed as a mouth-wash and gargle, under the name of "Oriental Water."

A teaspoonful is to be used, mixed with a glass of water.

*Delabarre.*

#### EMULSION WITH CARBONATE OF POTASSA.

R. Carbonate of potassa one scruple  
Yolk of egg three drachms  
Mucilage one drachm

Oil of almonds

Cherry water, each, one ounce

Linden water two ounces

Mix. A teaspoonful, every hour, in the colic of children.

*Rosenstein.*

#### FOMENTATION WITH CARBONATE OF POTASSA.

R. Carbonate of potassa  
Soap, each, one ounce  
Elder water two pounds

Dissolve, filter, and add

Sal ammoniac two drachms

As an application to bruises, &c.

*Cadet.*

#### LINIMENT WITH CARBONATE OF POTASSA.

R. Solution of carbonate of potassa two ounces  
Olive oil four ounces  
Yolk of eggs two

Mix. As an application to chaps.

*Plenck.*

#### LOTION OF CARBONATE OF POTASSA.

R. Carbonate of potassa three drachms  
Water four ounces

In pruritus vaginæ.

*Trousseau.*

R. Carbonate of potassa one ounce  
Rose water one pound

Dissolve, and filter. As an application to chilblains.

*Brugnatelli.*

R. Carbonate of potassa three drachms  
Common salt two drachms

Rose water eight ounces

Orange flower-water two ounces

Mix. As a lotion in sunburn and tan.

*Sundelin.*

#### MIXTURE OF CARBONATE OF POTASSA.

R. Carbonate of potassa two drachms  
Antimonial wine one fl. drachm



Laudanum forty drops  
Compound spirit of lavender two fl. drachms  
Distilled water four fl. ounces  
Mix. A tablespoonful, every hour or two, as an expectorant. *Ellis.*  
R. Carbonate of potassa one scruple  
Lemon juice four fl. drachms  
Cinnamon water seven fl. drachms  
Wine of ipecacuanha fifteen to twenty drops  
Mix. To be taken every three or four hours, as a diaphoretic. *Burke.*  
R. Carbonate of potassa one scruple  
Powdered gum Arabic half a drachm  
Oil of mint two drops  
Laudanum ten drops  
Water three drachms  
Mix. A tablespoonful for a dose, as may be required, in sickness of the stomach. *Ellis.*  
R. Carbonate of potassa one drachm  
Peppermint water four ounces  
Hoffmann's anodyne two scruples  
Simple syrup half an ounce  
Mix. Dose, a tablespoonful, in asthma, with acidity of the stomach. *Augustin.*  
R. Carbonate of potassa  
White sugar, each, one drachm  
Compound spirit of lavender two fl. drachms  
Laudanum forty drops  
Mint water four fl. ounces  
Mix. A tablespoonful, every hour or two, in sickness of the stomach, with acidity. *Ellis.*  
R. Aromatic spirit of ammonia one fl. drachm  
Carbonate of potassa two drachms  
Cinnamon water four fl. ounces  
Mix. A teaspoonful occasionally, in pyrosis. *Ellis.*  
R. Carbonate of potassa  
Ipecacuanha  
Calomel, each, one drachm and a-half  
Water two pints  
Boil down to one-half in an earthen vessel. Two spoonfuls a-day, in a quart of infusion of sarsaparilla. *St. Marie.*  
R. Carbonate of potassa one drachm  
Elder-flower water six ounces  
Syrup of marsh mallow one ounce  
Mix. A spoonful every hour, in angina, accompanied with aphtha, in catarrhal fevers. *Radiis.*  
R. Carbonate of potassa one scruple  
Cochineal half a drachm

Hyssop water four ounces  
Syrup of poppies  
tolu, each, half an ounce  
Orange flower-water  
Syrup of ipecacuanha, each, one ounce  
Mix. A spoonful, every two hours, in hooping-cough. *Bories.*

COMPOUND TINCTURE OF CARBONATE OF POTASSA.

R. Carbonate of potassa  
Cinnamon water, each, one pound  
Opium two ounces  
Vinous cinnamon water four ounces  
Digest in a water-bath for three weeks, often agitating; and add to the filtered solution.  
Gum Arabic two ounces  
Carbonate of ammonia one ounce  
Cinnamon water six ounces  
In syphilis. *Niemann.*

OINTMENT OF CARBONATE OF POTASSA.

R. Carbonate of potassa one ounce  
Flowers of sulphur two ounces  
Lard four ounces  
Mix. As a friction in itch. *Brera.*

POTASSÆ BICARBONAS.

BICARBONATE OF POTASSA.

R. Carbonate of potassa four pounds  
Distilled water ten pints  
Dissolve the carbonate in the water, and pass carbonic acid through the solution, till it is fully saturated. Then filter and evaporate, that crystals may form, taking care that the heat does not exceed 160°. Pour off the supernatant liquor, and dry the crystals on bibulous paper. Carbonic acid is obtained from marble, by the addition of dilute sulphuric acid. *U. S. Ph.*

Properties are the same as those of the carbonate, but it generally agrees better with the stomach. Dose, twenty grains to a drachm.

EFFERVESCING POWDERS OF BICARB. POTASSA.

R. Bicarbonate of potassa six hundred and forty grains.  
Divide into sixteen powders.  
R. Tartaric acid one ounce  
Divide into sixteen powders.  
Keep the acid and alkaline powders in paper of different colors. *Ed. Ph.*  
Usually made with the bicarbonate of soda.

EFFERVESCING SOLUTION OF POTASSA.

R. Bicarbonate of potassa one drachm  
Distilled water one pint (Imp.)



Dissolve, and pass into it carbonic acid, compressed by force, more than is sufficient for saturation; keep in a well-stopped vessel.  
*Lond. Ph.*

#### MIXTURE OF BICARBONATE OF POTASSA.

R. Bicarbonate of potassa ten to fifteen grains  
Seltzer water six fl. ounces  
Dissolve. To be taken three or four times a-day, as an antilithic.  
*Ellis.*

R. Bicarbonate of potassa one drachm  
Mint water six ounces  
Dissolve. A spoonful every hour—has been recommended in cholera.  
*Ammon.*

### POTASSÆ ET AMMONIÆ CARBONAS.

#### CARBONATE OF POTASSA AND AMMONIA.

R. Carbonate of potassa four parts  
ammonia one part  
Water sufficient

to dissolve the two salts; pass a current of carbonic acid through the solution to saturation, heat gently, and let crystallize.

In doses of half a drachm, in half an ounce of mint water, four times a-day, in diabetes, dyspepsia, gravel, &c.

*Swedjaur.*

### POTASSÆ CHLORAS.

#### CHLORATE OF POTASSA.

R. Carbonate of potassa two parts  
Quicklime one part  
and expose to a current of chlorine. When saturated, heat the mixture gently, digest it in water, and separate the chlorate from the filtered liquid, by crystallization.

*Graham.*

Has been used in scurvy, chronic hepatitis, syphilis, &c., in doses of fifteen to twenty grains.

#### POWDER OF CHLORATE OF POTASSA.

R. Chlorate of potassa six to eight grains  
Sugar one scruple  
Mix. To be given two to four times a-day, in phthisis, to diminish the febrile state.  
*Radius.*

#### COMPOUND POWDER OF CHLORATE OF POTASSA.

R. Chlorate of potassa seven grains  
Muriate of soda one scruple  
Carbonate of soda half a drachm  
Mix. For a dose.  
*Stevens.*

Recommended in cholera.

#### SOLUTION OF CHLORATE OF POTASSA.

R. Chlorate of potassa one drachm  
Distilled water twelve fl. ounces  
Dissolve.  
*Copland.*

As a lotion to indolent ulcers.

R. Chlorate of potassa half a drachm  
Syrup two and a-half drachms  
Water twelve and a-half drachms  
Mix. To be given in the course of the day, in tablespoonful doses, in cancrum oris.  
*H. Hunt.*

R. Chlorate of potassa one drachm and a-half  
Distilled water five ounces  
Dissolve. A tablespoonful, every two hours, in obstinate rheumatism, tic-douloureux, &c.  
*Knod.*

### POTASSÆ CITRAS.

#### CITRATE OF POTASSA.

R. Citric acid ten ounces  
Bicarbonate of potassa fourteen ounces  
Water two pints

Dissolve the acid in the water, and gradually add the bicarbonate, when effervescence ceases; filter the solution, if necessary, through paper, free from iron, and evaporate to dryness, in a porcelain capsule, observing to stir constantly, as soon as the salt begins to granulate.  
*W. Procter.*

R. Carbonate of potassa three drachms  
Recent lemon juice sufficient  
to saturate. Let stand for twenty-four hours, filter, and evaporate to dryness.  
*Van Mons.*

#### SOLUTION OF CITRATE OF POTASSA.

R. Citric acid half an ounce  
Oil of lemons two minims  
Water half a pint  
Carbonate of potassa sufficient  
Rub the citric acid with the oil of lemons, and then with the water, till dissolved; lastly, add the carbonate of potassa gradually, till the acid is perfectly saturated, then filter.  
*U. S. Ph.*

This neutral mixture is used where fresh lemon juice cannot be procured; it is as efficacious, but is not as agreeable.

R. Citrate of potassa two drachms  
Distilled water four to six fl. ounces  
Oil of lemons two drops  
White sugar two drachms

Mix. A tablespoonful to be taken every two hours. In same cases as the neutral mixture.  
*Ellis.*



COMPOUND NEUTRAL MIXTURE.

R. Lemon juice one ounce  
Carbonate of potassa sufficient  
to saturate, add  
Mint water seven ounces  
Tartar emetic one grain  
Syrup half an ounce  
Mix. *Lisbon Ph.*

MIXTURE OF CITRATE OF POTASSA, AND PERUVIAN BARK.

R. Lemon juice one fl. ounce and a-half  
Carbonate of potassa one drachm  
Tincture of Peruvian bark one fl. ounce  
Cinnamon water three fl. ounces  
Mix. A tablespoonful every two hours.  
To promote insensible perspiration while taking bark. *Ellis.*

POTASSÆ IODAS.

IODATE OF POTASSA.

R. Iodine at will  
Caustic potassa sufficient  
Dissolve the iodine in the alkali, till the solution begins to become colored. Evaporate to dryness; treat the residue with alcohol, and preserve the residue. The alcohol takes up the iodide of potassium. *Ferrara Ph.*

SYRUP OF IODATE OF POTASSA.

R. Iodate of potassa twenty grains  
Peppermint water two drachms  
Simple syrup ten ounces  
Mix. *Giordano.*

POTASSÆ NITRAS.

NITRATE OF POTASSA.

POTASSÆ NITRAS PURIFICATUM.

PURIFIED NITRATE OF POTASSA.

R. Nitrate of potassa one part  
Boiling water two parts  
Dissolve, filter, and set aside, so that, on cooling, crystals may form. *Dub. Ph.*

FUSED NITRATE OF POTASSA. SAL PRUNELLE.

R. Nitrate of potassa at will  
Melt in a crucible, and cast it into moulds. *Guibourt.*

R. Nitrate of potassa two pounds and a-half  
Melt in a crucible, adding gradually,  
Flowers of sulphur half an ounce  
After the deflagration, and the mixture is

somewhat cooled, pour it into a basin, and turn this in all directions, so that the mass may cool in a third and uniform layer. When cool, break this in pieces.

*Span. Ph.*

This preparation will contain some sulphate of potassa.

POWDER OF NITRATE OF POTASSA AND ORRIS ROOT.

R. Nitrate of potassa one drachm  
Spermaceæ two drachms  
Sugar  
Orris root, each, one ounce  
Mix. A teaspoonful, in catarrhal affections. *Augustin.*

COMPOUND NITROUS POWDERS.

R. Powdered nitre one drachm and a-half  
Tartar emetic one grain  
Calomel four grains  
Mix, and divide into eight powders. One every two hours. As a diaphoretic, &c. in fever. *Dewees.*

R. Nitrate of potassa two ounces  
Bitartrate of potassa four drachms  
Tartar emetic four grains  
Mix. *Beasley.*

POWDER OF NITRATE OF POTASSA AND SQUILL.

R. Nitrate of potassa fifteen grains  
Powdered squill  
pimento, each, ten grains  
Mix. Dose, ten to twenty grains, two or three times a-day, as a diuretic. *Swedjaur.*

POWDER OF NITRATE OF POTASSA AND CAMPHOR.

R. Nitrate of potassa ten grains  
Camphor four to eight grains  
Gum Arabic twenty-four grains  
Mix, and triturate well; one-third to one-half at a dose. *Foy.*

PILLS OF NITRATE OF POTASSA.

R. Nitrate of potassa six drachms  
Gum Arabic three drachms  
Liquorice root  
Marsh mallow, each, three ounces  
Simple syrup sufficient  
Mix, and make pills of five grains. In inflammation of the urethra and dysuria; five or six, three times a-day. *Foy.*

PILLS OF NITRATE OF POTASSA AND CAMPHOR.

R. Nitrate of potassa four parts  
Camphor  
Conserve of roses, each, two parts  
Mix, and make pills of four grains. Two to ten a-day, in gonorrhœa. *Guibourt.*



## NITRATED EMULSION.

R. Nitrate of potassa two drachms  
 Sugar of milk one ounce  
 Extract of henbane half a scruple  
 Emulsion of almonds one pound  
 Mix, and dissolve. A spoonful every hour, in gonorrhœa. *Phœbus.*

## COOLING LOTION OF NITRATE OF POTASSA.

R. Nitrate of potassa  
 Sal ammonia, each, one part  
 Water forty-eight parts  
 Dissolve, and add  
 Vinegar four parts  
 As a lotion and application, by means of compresses, to contusions and ecchymoses. *Hep. Ph.*

## NITRATE OF POTASSA GARGLE.

R. Nitrate of potassa seven drachms  
 Barley water fourteen fl. ounces  
 Oxymel one fl. ounce and a-half  
 Mix. As gargle, in inflammatory sore throat. *Ainslie.*

## MIXTURE OF NITRATE OF POTASSA.

R. Nitrate of potash eight grains  
 Tincture of digitalis fifteen to twenty drops  
 Water ten fl. drachms  
 Sweet spirit of nitre  
 Syrup of roses, each, half a fl. drachm  
 Mix. To be taken twice a-day, as a diuretic. *Burke.*

R. Nitrate of potassa one drachm and a-half  
 Mucilage of gum Arabic two fl. ounces  
 Antimonial wine forty minims  
 Syrup of orange peel half a fl. ounce  
 Water four fl. ounces  
 Mix. One fl. ounce, three times a-day, in remittent fever, with hot skin. *Ainslie.*

R. Nitrate of potash eight to ten grains  
 Water eleven fl. drachms  
 Solution of tartar emetic fifteen drops  
 Simple syrup  
 Sweet spirit of nitre, each, half a fl. drachm  
 Mix. To be taken twice a-day, as a diaphoretic. *Burke.*

R. Extract of chamomile one drachm  
 Nitrate of potassa one ounce  
 Chamomile water four ounces

Mix. A spoonful every two or three hours, as an antispasmodic. *Radius.*

R. Nitrate of potassa half an ounce  
 Barley water one pound  
 Syrup of marsh mallows six ounces  
 Oil of almonds four ounces

Mix. A glassful every four hours, in dysuria and strangury. *Cadet.*

R. Nitrate of potassa two drachms  
 Decoction of asparagus two pounds  
 Oxymel of squill half an ounce

Mix. To be taken in divided doses, as a diuretic. *Brera.*

R. Juniper berries, bruised, two ounces  
 Boiling water one pint

When cold, strain, and add  
 Nitrate of potassa two drachms  
 Syrup of ginger one fl. ounce  
 One ounce to an ounce and a-half, every three or four hours, in dropsy. *Hartman.*

## POTASSÆ OXALAS.

## OXALATE OF POTASSA.

The article usually sold for oxalate of potassa, salt of sorrel, &c., is the binoxalate of potassa. It is not much used in medicine, but is employed in the arts, to remove ink and iron stains from linen and cotton; to bleach the straw for bonnets, &c.

R. Oxalic acid one part  
 Carbonate of potassa sufficient  
 to saturate; add to the solution three parts more of the acid; evaporate, and crystallize. *Cooley.*

## POWDER OF OXALATE OF POTASSA.

R. Oxalate of potassa twenty grains  
 Tartrate of potassa  
 Sulphate of potassa, each, one drachm  
 Scammony fifteen grains  
 Red saunders ten grains  
 Mix. To be taken in the morning, in intermittent fevers. Two doses are stated to be sufficient. *Giordano.*

## LOZENGES OF OXALATE OF POTASSA.

R. Oxalate of potassa one drachm and a-half  
 White sugar eight ounces  
 Gum tragacanth two scruples  
 Lemon water five drachms  
 Oil of lemons eight drops  
 Mix, and make lozenges of twelve grains. *Guibourt.*

## POTASSÆ SILICAS.

## SILICATE OF POTASSA.

R. Powdered quartz or flint one part  
 Subcarbonate of potassa two parts  
 Mix, and fuse in a crucible; when cool, dissolve in water; filter, and evaporate to dryness.



SOLUTION OF SILICATE OF POTASSA.

R. Silicate of potassa ten to fifteen grains  
Distilled water six to eight fl. ounces  
Dissolve. To be taken twice a-day, to  
remove gouty concretions. *Ure.*

SOLUBLE GLASS.

R. Subcarbonate of potassa seventy parts  
Subcarbonate of soda fifty-four parts  
Silex one hundred and ninety-two parts  
Melt together. The resulting glass is  
soluble in boiling water. The solution  
forms a fine, transparent, elastic varnish.  
*Dobereiner.*

POTASSÆ SULPHAS.

SULPHATE OF POTASSA.

R. Residuum of the prepara-  
tion of nitric acid two pounds  
Boiling water two gallons  
Expel the excess of acid by heating the  
salt in a crucible; boil the remainder in  
the water, till a pellicle forms; filter the  
solution; set aside to crystallize; pour  
off the water, and dry the crystals.  
*Lond. Ph.*

A mild cathartic, in doses of four or five  
drachms, but in large doses, acts as an irritant.

COMPOUND SALINE POWDER.

R. Sulphate of potassa three ounces  
Muriate of soda  
Sulphate of magnesia, each, four ounces  
Dry the salts separately, with a gentle  
heat; then triturate them well together,  
and preserve in glass vessels. *Ed. Ph.*

Aperient, in doses of two or three drachms,  
dissolved in half a pint of carbonic acid water,  
before breakfast.

POWDER OF SULPHATE OF POTASSA AND  
RHUBARB.

R. Sulphate of potassa one drachm  
Powdered rhubarb  
one scruple and a-half  
chamomile one drachm  
Mix, and divide into six powders; one,  
twice a-day, in sugar and water, in dys-  
pepsia and torpor of the bowels. *Ellis.*

R. Sulphate of potassa two ounces  
Powdered rhubarb one ounce  
Sal ammoniac half an ounce  
Mix. Half a drachm, in same cases as  
the last. *Pfaff.*

PILLS OF SULPHATE OF POTASSA.

R. Sulphate of potassa two drachms  
Powdered rhubarb two scruples  
Oil of fennel six drops  
Extract of blessed thistle sufficient

Mix, and make sixty pills. Dose, five or  
six a-day, as a purgative. *Phœbus.*

MIXTURE WITH SULPHATE OF POTASSA.

R. Centaury  
Chamomile, each, one ounce  
Water sufficient  
to obtain three pints of infusion; add  
Sulphate of potassa  
Honey, each, two ounces  
Three glassfuls a-day, in fevers.  
*Swediaur.*

POTASSÆ BISULPHAS.

BISULPHATE OF POTASSA.

R. Salt remaining after dis-  
tillation of nitric acid two pounds  
Sulphuric acid one pound  
Boiling water six pints (Imp.)  
Dissolve the salt in the water, add the acid,  
and mix. Boil down the solution, and set  
aside, that crystals may form. *Lond. Ph.*  
Aperient and tonic, in doses of one to two  
drachms.

EFFERVESCING DRAUGHT OF SULPHATE  
OF POTASSA.

R. Sulphate of potassa  
Carbonate of soda, each, one drachm  
Dissolve separately, in two fl. ounces of  
water, each; mix, and take whilst effervesc-  
ing. *Barker.*

DISINFECTING POWDER.

R. Acid of sulphate of  
potassa four hundred and ten parts  
Subacetate of lead seventy parts  
Manganese thirty parts  
Reduce these, separately, to fine powder,  
and, when wanted, mix in a proper vessel.  
*Keist.*

POTASSÆ ET AMMONIÆ  
SULPHAS.

SULPHATE OF POTASSA AND AMMONIA.

R. Bisulphate of potassa one part  
Boiling water two parts  
Dissolve, and add  
Solution of ammonia sufficient  
to saturate to excess, and crystallize.  
*Van Mons.*

POTASSÆ ET MAGNESIÆ  
SULPHAS.

SULPHATE OF POTASSA AND MAGNESIA.

R. Sulphate of  
potassa three and a-half parts



Sulphate of magnesia      ten parts  
Boiling water              sufficient  
Filter, and crystallize.

### POTASSÆ SULPHAS CUM SULPHURE.

SULPHATE OF POTASSA WITH SULPHUR.

R. Nitrate of potassa  
Sulphur                      equal parts  
Mix, and throw in small successive portions into a red-hot crucible; when cool, reduce to powder, and keep in well-closed bottles. *Ed. Ph.*

Has much the same properties as the sulphate of potassa. Dose, half a drachm to a drachm.

### POTASSÆ TARTRAS.

TARTRATE OF POTASSA.

R. Carbonate of potassa      sixteen ounces  
Bitartrate of  
potassa                      three pounds, or q. s.  
Boiling water              one gallon  
Dissolve the carbonate of potassa in the water, then gradually add the bitartrate in fine powder to the solution, till it is perfectly saturated, and boil. Filter the liquor, evaporate till a pellicle forms, and set aside to crystallize. Pour off the liquid, and having dried the crystals on bibulous paper, keep them in closely stopped bottles. *U. S. Ph.*

A mild, cooling purgative, in doses from a drachm to an ounce.

### POWDER OF TARTRATE OF POTASSA AND RHUBARB.

R. Tartrate of potassa      two drachms  
Powdered rhubarb  
orange peel, each, one drachm  
Oil of cajeput              one scruple  
Mix. A teaspoonful, three times a-day, in obstructions of the portal system. *St. Marie.*

### MIXTURE OF TARTRATE OF POTASSA.

R. Tartrate of potassa six to eight drachms  
Infusion of chicory      twenty ounces  
Manna                      two ounces  
Mix. To be taken in the morning, in four portions, every day or every second day, for a fortnight, in chronic affections of the liver. *Grant.*  
R. Tartrate of potassa      one ounce  
Extract of soapwort      half an ounce  
Balm water                  six ounces

Mix. Two spoonfuls, night and morning, in all forms of hemorrhoidal disease.

*Radiis.*

R. Tartrate of potassa      half an ounce  
Extract of centaury      two drachms  
Water                      eight ounces

Mix. Two spoonfuls every hour or two, in obstructions of the liver. *Swed. iaur.*

R. Tartrate of potassa      one drachm  
Nitrate of potassa      half a drachm  
Manna                      one ounce  
Decoction of dandelion      six ounces

Mix. Two spoonfuls, every two hours, in dropsy consecutive to scarlet fever. *Phæbus.*

### POTASSÆ BITARTRAS.

BITARTRATE OF POTASSA.

#### CREAM OF TARTAR.

This is cathartic, diuretic, and cooling. The dose is from one to two drachms, as an aperient; and from half an ounce to one ounce, as a hydragogue purgative.

### DENTIFRICE OF BITARTRATE OF POTASSA.

R. Bitartrate of potassa      six parts  
Powdered shell lac  
cuttlefish bone  
orris root, each, eight parts  
cloves  
myrrh  
mastic, each, two parts  
Triturate well together. *Guibourt.*

R. Bitartrate of potassa  
Rhatany                      equal parts  
Orris root                      sufficient  
to give an agreeable odor. Triturate well together. *Ferrara Ph.*

### POWDER OF BITARTRATE OF POTASSA.

R. Bitartrate of potassa      three ounces  
Nitrate of potassa      three drachms  
Sugar                      four ounces  
Mix. Dose, one to two drachms, in whey, as a laxative. *St. Marie.*

R. Bitartrate of  
potassa                      a drachm and a-half  
Powdered squill              two grains  
foxglove                      one grain  
ginger                      five grains  
Mix. Make a powder, to be taken every eight hours. In ascites and anasarca. *A. T. Thomson.*

R. Bitartrate of potassa      thirty grains  
Powdered squill              two grains  
ginger                      four grains  
Mix. Make a powder, to be taken every eight hours. Useful in ascites. *A. T. Thomson.*



ELECTUARY OF BITARTRATE OF POTASSA.

R. Bitartrate of potassa one ounce  
Powdered ginger  
Conserve of roses, each, one drachm  
Syrup of orange peel sufficient  
Mix. To be taken in spoonful doses, as a hydragogue purgative. *Monro.*

OXYMEL OF BITARTRATE OF POTASSA.

R. Bitartrate of potassa one part  
Clarified honey two parts  
Mix. As a purgative in bilious fever and dropsies. *Swediaur.*

SOLUTION OF BITARTRATE OF POTASSA.

R. Bitartrate of potassa one ounce  
Water two pints  
Dissolve. To be taken freely during the day, as a diuretic. *Ellis.*

COMPOUND SOLUTION OF BITARTRATE OF POTASSA.

R. Bitartrate of potassa two drachms  
Manna two ounces  
Water eight ounces  
Lemon juice half an ounce  
Mix, and clarify with the white of an egg, infuse a small quantity of orange peel in the fluid, and strain on cooling. A pleasant laxative. *Taddei.*

MIXTURE OF BITARTRATE OF POTASSA.

R. Bitartrate of potassa two scruples  
Antimonial wine  
Vinegar of squill, each, half a drachm  
Parsely water one ounce and a-half  
Syrup of seneka six drachms  
Mix. A teaspoonful every two hours, in dropsy supervening on scarlatina. *Vogt.*  
R. Bitartrate of potassa half an ounce  
Barley water two pounds  
Dissolve, and add  
Brandy one to two ounces  
Much praised as a diuretic in dropsies. *Milmann.*

R. Bitartrate of potassa one ounce  
Borax two drachms  
Boiling water sufficient  
to dissolve. To ten ounces of the cooled solution add  
Nitrate of potassa two drachms  
Oxymel two ounces  
Mix. *Beasley.*

R. Bitartrate of potassa  
Extract of horehound, each, two drachms  
Balm water three ounces  
Mix. One-half to be taken morning and evening, in hemorrhoids, with constipation. *Augustin.*

IMPERIAL.

R. Bitartrate of potassa half an ounce  
Lemon, cut into slices, one  
White sugar half a pound  
Water three pints  
Mix. Let stand for half an hour, and strain. A pleasant, cooling drink. *Gray.*

POTASSÆ ET AMMONIÆ TARTRAS.

TARTRATE OF POTASSA AND AMMONIA.

R. Bitartrate of potassa at will  
Boiling water sufficient  
to dissolve; add, gradually,  
Solution of ammonia  
to saturate the excess of the acid; filter, and evaporate, so that crystals may form. *Hamb. Ph.*

SOLUTION OF TARTRATE OF POTASSA AND AMMONIA.

R. Solution of carbonate of ammonia six ounces  
Add, gradually,  
Bitartrate of potassa sufficient  
to saturate, let stand for some hours, and filter. Diuretic, in doses of a drachm. *Liepsic Ph.*

MIXTURE OF TARTRATE OF POTASSA AND AMMONIA.

R. Tartrate of potassa and ammonia half an ounce  
Fennel water six ounces  
Extract of dandelion  
Clarified honey, each, one ounce  
Mix. A spoonful every hour, as a diuretic. *Phæbus.*

PRINOS.

BLACK ALDER.

This, which is also known as *Winter berry*, is an indigenous shrub, found in most parts of the country, principally in low, moist situations; flowering in June, and bearing numerous scarlet berries, which remain on the bush after the fall of the leaves.

*Sax. Syst.* Hexand. monog. *Nat. Syst.* Aquifoliaceæ.

*Linn. Sp. Pl.* 471. *Griffith, Med. Bot.* 434.

The officinal portion is the bark, which, when dried for use, is in slender pieces, of a greenish white color internally, and of an ash gray, mixed with brown externally; brittle, inodorous, and of a bitter, astringent taste. It is tonic and astringent, and has been used with success in diarrhœa, intermittent fevers, &c. The dose is from thirty grains to a drachm, three or four times a-day.

DECOCTION OF BLACK ALDER.

R. Black alder two ounces



Boiling water three pints  
Boil down to a quart, and strain. One  
gill, every two hours. *W. P. C. Barton.*

## PRUNUM.

## PRUNES.

Prunes are the dried fruit of various varieties of *Prunus domestica*, or cultivated Plum tree. They are principally derived from the south of France, and are much used as an article of dessert, but are also employed in medicine, as a laxative, either alone or in combination, especially with senna, as in the confection of senna, &c., in the form of pulp, which is prepared by softening them in the vapor of boiling water; separating the stones, and pressing through a hair sieve.

## PRUNUS VIRGINIANA.

## WILD CHERRY BARK.

This is the bark of *Cerasus serotina*, though the U. S. Pharm. still retains the old, but incorrect designation. The *Cerasus serotina* is found in most parts of the United States, and in some situations attains a very large size.

*Sex. Syst.* Icosand. monog. *Nat. Syst.* Drupaceae.

Decandolle. Prod. ii. 540. Griffith, Med. Bot. 288.

The bark of both the roots and branches, is used, but the former is to be preferred. As dried for use, it is in pieces of various sizes; deprived of epidermis; of a reddish brown color, and readily pulverized; when fresh it has an odor of bitter almonds, which is much diminished on drying, but reappears on maceration; the taste is bitter and aromatic. It is tonic, but also exercises a sedative action on the circulatory and nervous systems, and is much used in a variety of diseases. The dose in powder is half a drachm to two drachms.

## INFUSION OF WILD CHERRY BARK.

*R.* Wild cherry bark,  
bruised, half an ounce  
Cold water one pint  
Macerate for twenty-four hours, and  
strain. *U. S. Ph.*

Dose, two or three fl. ounces, three or four times a-day.

## SYRUP OF WILD CHERRY BARK.

*R.* Powdered wild cherry  
bark four ounces  
Water twelve fl. ounces  
Macerate for two days; put in a displacement apparatus; add water till twelve fl. ounces are obtained, returning the first portions till it comes away clear, then add

Sugar twenty-four ounces  
And form syrup. *Procter.*

Dose, from a fl. drachm to a fl. ounce.

*R.* Same as the above, but with half the quantity of sugar. *D. S. Jones.*

## PYRETHRUM.

## PELLITORY.

This is the root *Anacyclus pyrethrum*, a small plant with perennial roots and annual stems, a native of the countries bordering on the Mediterranean.

*Sex. Syst.* Syngen. super. *Nat. Syst.* Astera-ceae.

Decandolle. prod. vi. 15. Griffith, Med. Bot. 402.

The dried root is about the size of the little finger, with a thick; brown bark, marked with black, shining points; it is inodorous; its taste is at first slight, but afterwards extremely acrid, leaving a burning, pricking sensation. It is a powerful local irritant, and is principally used as a masticatory and sialagogue, but has also been employed in intermittents, palsies, &c.

## LOZENGES OF PELLITORY.

*R.* Powdered pellitory  
mastich, each, one drachm  
Mucilage of tragacanth sufficient  
Mix, and make lozenges of twelve grains  
each. As a masticatory in toothache.

*Foy.*

## TINCTURE OF PELLITORY.

*R.* Pellitory one part  
Alcohol four parts  
Macerate for eight days, and filter.

*Par. Cod.*

## COMPOUND TINCTURE OF PELLITORY.

*R.* Pellitory four drachms  
Camphor three drachms  
Opium one drachm  
Oil of cloves two drachms  
Alcohol six fl. ounces

Macerate for eight days, and filter. As a sialagogue in toothache. *Brande.*

*R.* Pellitory one ounce  
Para cress four ounces  
Italian elecampane one ounce  
Alcohol eight fl. ounces

Macerate for fifteen days, express, and filter. Much celebrated as an odontalgic, under the name of *Paraguay Roux*.

## LINIMENT OF PELLITORY.

*R.* Tincture of pellitory six fl. drachms  
Camphorated oil half a fl. ounce  
Water of ammonia half a fl. drachm  
Mix. As a lotion for chilblains, and in rheumatic pains. *Radiis.*

## PLASTER OF PELLITORY.

*R.* Yellow wax three ounces  
Turpentine two ounces and a-half  
Melt together, and add



Ammoniac  
Sagapenum  
Galbanum

Powdered pellitory

mustard, each, one ounce

Mix well. As a rubefacient application  
in rheumatism, &c. *Cadet.*

#### EXTRACT OF PELLITORY.

R. Pellitory, ground, one pound  
Alcohol two pints  
Ether half a pint

Mix the ether with a pint of the alcohol, pour it gradually on the powder, put it in a displacer, when it ceases to drop, add the remainder of the alcohol, and finally sufficient diluted alcohol, to displace two pints and a half in all. Allow the ethero-alcoholic tincture to evaporate spontaneously, or with a moderate heat, until a soft extract is obtained.

Employed to destroy the sensibility of the nerves of teeth, previous to plugging, or for toothache. *W. Procter.*

## Q

### QUASSIA.

#### QUASSIA.

Under the name of Quassia, the wood of two different plants is used,—that of *Q. amara*, and of *Q. (Picræna) excelsa*, the latter being alone recognized in the U. S. Pharm. This is furnished by a large tree found in many of the West India islands, and known as the *bitter ash*.

*Sex. Syst.* Pentand. monog. *Nat. Syst.* Simarubaceæ.

Lind. Flor. Med. 208. Griffith, Med. Bot. 200.

It is found in commerce, in billets of various sizes, having a smooth, brittle bark. The wood is white, but becomes yellowish on exposure to the air; it has scarcely any smell, but possesses an intense, permanent bitter taste. It is a pure and powerful tonic, and is much used for that purpose. It is seldom given in substance.

#### INFUSION OF QUASSIA.

R. Rasped quassia two drachms  
Cold water one pint

Macerate for twelve hours, and strain.

*U. S. Ph.*

Dose, two fl. ounces, three or four times a-day.

#### COMPOUND INFUSION OF QUASSIA.

R. Quassia  
Serpentaria  
Orange peel, each, half an ounce  
Boiling water two pints

Infuse and strain. A teacupful, cold, three times a-day. *Ellis.*

#### EXTRACT OF QUASSIA.

R. Prepared like extract of gentian.  
*U. S. Ph.*

Dose, about five grains. A very efficient, bitter tonic.

#### TINCTURE OF QUASSIA.

R. Rasped quassia two ounces

Diluted alcohol two pints

Macerate for fourteen days, express, and filter; or it may be made by displacement.

*U. S. Ph.*

Dose, two fl. drachms.

#### COMPOUND TINCTURE OF QUASSIA.

R. Bruised cardamoms  
cochineal, each, half an ounce

Powdered cinnamon

Chipped quassia, each, six drachms

Raisins seven ounces

Diluted alcohol two pints (impl.)

Digest for seven days, strain, express residue, and filter. Dose, one or two fl. drachms. *Ed. Ph.*

#### WINE OF QUASSIA.

R. Rasped quassia half an ounce  
Orange peel two drachms  
Wine one pint and a-half

Macerate for twenty-four hours, express, and filter. Dose, three fl. drachms to a fl. half ounce, twice a-day. *Radius.*

### QUERCUS.

#### OAK BARK.

Several species of oak are recognized as officinal in the Pharmacopœias, as the *Q. alba*, or *White oak*, and *Q. tinctoria*, or *Black oak*, in the U. S.; *Q. pedunculata*, European *White oak*, in the Lond. and Edin.; *Q. robur*, European oak, in the Dublin.

*Sex. Syst.* Monœc. polyand. *Nat. Syst.* Corylaceæ.

The part used is the bark; this, in all the species, is astringent and tonic, and has been used in a variety of diseases, but more generally as an external application than as an internal



remedy. The dose is from thirty grains to a drachm.

#### COMPOUND POWDER OF OAK BARK.

R. Powdered oak bark                   one scruple  
Calamus  
Gentian, each,                           five grains  
Mix. To be taken every three hours, in  
apyrexia of intermittents.           *Augustin.*

#### EXTRACT OF OAK BARK.

R. To be prepared like extract of gentian.  
*Dub. Ph.*

The dose is from ten to forty grains.

#### DECOCTION OF OAK BARK.

R. Oak bark                               eight drachms  
Water                                       two pints  
Boil down to one pint, and strain.  
*Dub. Ph.*

The *U.S. Ph.*, for decoction of white oak bark, orders one ounce of the bark, and water a pint and a-half. The dose is a wineglassful, as an astringent in chronic diarrhoea, &c.

#### GARGLE OF OAK BARK.

R. Decoction of oak bark               one pint  
Add  
Alum                                       half a drachm  
Brandy                                   two fl. ounces  
As a gargle in chronic sore throat, with relaxed uvula, and as an injection in leucorrhœa.                                       *Ellis.*

#### CATAPLASM OF OAK BARK.

R. Powdered oak bark  
catechu  
Barley meal, each,                   one ounce  
Water                                   sufficient  
Boil to the proper consistence.       *Foy.*

As an application in gangrene and mortification.

#### CONFECTION OF ACORNS.

R. Powdered acorns                   three ounces  
red coral  
catechu, each,                       one ounce and a-half  
Confection of dog rose               ten ounces  
Syrup of red roses                   sufficient  
Mix. One drachm every four hours, in chronic diarrhoea.                       *Bories.*

#### ACORN COFFEE.

R. Acorns, roasted,                   six drachms  
Coffee, roasted,                       two drachms  
Water                                   sufficient  
Boil for a short time, and strain. Two or three cupfuls a-day, in scrofula and rachitis.  
*Augustin.*

### QUINIA.

#### QUININE.

R. Sulphate of quinine               one part  
Boiling water                       thirty parts  
Dissolve. Add water of ammonia sufficient to precipitate the quinine, wash the precipitate in warm distilled water, and dry it.

#### TINCTURE OF QUININE.

R. Quinine                               one part  
Alcohol                               seven parts  
Dissolve. Dose, twenty to forty drops.  
*Beral.*

#### IMPURE QUININE.

R. Yellow bark                       one hundred parts  
Muriatic acid                       five parts  
Water                               five hundred parts  
Boil, decant, and repeat process three times with the same proportions of acid and water; unite decoctions, precipitate with milk of lime, wash and dry the precipitate, treat it several times with boiling alcohol, mix, and filter the solutions, and distil off the spirit.                                       *Trousseau.*

This is said to be as active as the sulphate, and to be less unpleasant to the taste. It is given in the same doses.

#### TINCTURE OF IMPURE QUININE.

R. Impure quinine                   one ounce  
Alcohol  
Distilled water, each, twelve fl. ounces  
Mix.                                       *Piorry.*

#### AMORPHOUS QUININE, OR CHINOÏDINE.

R. Mother waters of sulphate of quinine                               at will  
Solution of carbonate of potassa                               sufficient to precipitate. Wash and dry this precipitate, dissolve in sulphuric ether, decant, and evaporate by a gentle heat. *Neligan.*

### QUININÆ ACETAS.

#### ACETATE OF QUININE.

R. Quinine                               two parts  
Distilled water                       three parts  
Heat, and add as much acetic acid as will dissolve the quinine, and render the solution somewhat acid. Filter whilst boiling, and set aside to crystallize.       *Cottureau.*

Acts like the other salts of quinine, and is in no way superior to them.



## QUININÆ ARSENIAS.

### ARSENATE OF QUININE.

R. Arsenic acid      one drachm and a-half  
Distilled water      six fl. ounces  
Quinine              five drachms  
Boil till solution takes place. Filter, let  
crystallize, and purify by recrystallization.  
In intermittents; dose, one-fifth of a grain.  
*Bourrières.*

## QUININÆ CITRAS.

### CITRATE OF QUININE.

R. Quinine              two parts  
Distilled water      three parts  
Heat, and add sufficient citric acid to  
acidulate the mixture; when a perfect so-  
lution is effected, filter, and set aside to  
crystallize. *Magendie.*

The dose is the same as that of the sulphate:  
in the same cases.

### SYRUP OF CITRATE OF QUININE.

R. Acid citrate of  
quinine,              thirty-six grains  
Simple syrup              one pint  
Mix. Half fl. ounce to fl. ounce, in the  
twenty-four hours. *Magendie.*

## QUININÆ ET FERRI CITRAS.

### CITRATE OF QUININE AND IRON.

See *Ferri et Quinix Citras.*

## QUININÆ FERROCYANAS.

### FERROCYANATE OF QUININE.

R. Sulphate of quinine one hundred parts  
Ferrocyanide of  
potassa              thirty-one parts  
Distilled water

two thousand five hundred parts  
Boil for a few minutes, let the solution  
cool; separate the oily compound, and  
wash it with a little water. Dissolve it  
in boiling alcohol, and let crystallize.  
*Par. Cod.*

Said to be more efficacious than any other salt  
of quinia. Dose, about the same as the sul-  
phate.

### MIXTURE OF FERROCYANATE OF QUI- NINE.

R. Ferrocyanate of quinine      four grains  
Alcohol                              one fl. drachm  
Dissolve, and add  
Camphor water              seven fl. drachms  
To be taken as required, shaking the vial.  
*Donovan.*

## PILLS OF FERROCYANATE OF QUININE.

R. Ferrocyanate of  
quinine              twenty-four grains  
Mucilage of gum Arabic      sufficient  
Mix, and make twelve pills. Two for a  
dose. *Donovan.*

## QUININÆ HYDRIODAS.

### HYDRIODATE OF QUININE.

R. Add, by drops, a solution of twenty-  
four parts of iodide of potassium, in eight  
parts of water, to a strong solution of  
twenty parts of sulphate of quinine; wash  
the precipitate quickly, and dry in the  
shade. *Righini.*

In obstinate intermittents and scrofulous af-  
fections.

## QUININÆ ET FERRI IODI- DUM.

### IODIDE OF QUININE AND IRON.

R. Pour a strong solution of acid sul-  
phate of quinine into a fresh solution of  
iodide of iron; collect the precipitate;  
dry by pressing between blotting paper,  
and keep it from the air. *Bouchardat.*

## QUININÆ HYDRIODAS IODU- RETA.

### IODURETTED HYDRIODATE OF QUININE.

R. Add a solution of iodide of iron, con-  
taining a slight excess of iodine, to an  
acid solution of quinine. Treat the pre-  
cipitate with boiling alcohol, and filter  
whilst hot, and set aside to crystallize.  
*Bouchardat.*

These preparations are useful where an al-  
terative and tonic are required.

## QUININÆ ET HYDRARGYRI- CHLORIDUM.

### DOUBLE CHLORIDE OF QUININE AND MERCURY.

R. Corrosive sublimate      one part  
Hydrochlorate of quinine      three parts  
Dissolve separately, in the smallest quan-  
tity of water; mix the solutions; collect  
the precipitate, and dry by a gentle heat.  
*McDermott.*

### PILLS OF DOUBLE CHLORIDE OF QUI- NINE AND MERCURY.

R. Double chloride of quinine  
and mercury              fifteen grains  
Opium                              six grains



Crumb of bread sufficient  
 Mix, and form thirty pills. One, three  
 times a-day, to produce salivation.  
*Hamilton.*

## QUINIÆ KINAS.

## KINATE OF QUININE.

R. Alcoholic solution of sul-  
 phate of quinine at will  
 Aqueous solution of kinate  
 of lime sufficient  
 to precipitate; filter, evaporate; redissolve,  
 and crystallize. *Magendie.*

## PILLS OF KINATE OF QUININE.

R. Kinate of quinine  
 Black pepper  
 Extract of worm-  
 wood, each, one drachm  
 Mix, and make sixty pills. Three, every  
 two or three hours, in obstinate intermit-  
 tents. *Ronander.*

## QUINIÆ LACTAS.

## LACTATE OF QUININE.

R. Lactic acid at will  
 Quinine sufficient  
 to saturate; leave the solution to evapo-  
 rate spontaneously, in a shallow vessel, in  
 a warm room, till crystals are formed.  
*Bouchardat.*

## PILLS OF LACTATE OF QUININE.

R. Lactate of quinine half a drachm  
 Extract of juniper sufficient  
 Mix, and make twenty pills. Two to six  
 a-day, in intermittents. *Bouchardat.*

## MIXTURE OF LACTATE OF QUININE.

R. Lactate of quinine seven grains  
 Mint water five fl. drachms  
 Syrup of cloves one fl. drachm  
 Water three and a-half fl. ounces  
 Mix. To be taken in divided doses dur-  
 ing the apyrexia of intermittents.  
*Bouchardat.*

## SYRUP OF LACTATE OF QUININE.

R. Lactate of quinine fifteen grains  
 Dissolve in  
 Water one fl. ounce  
 Add  
 Sugar two ounces  
 Form syrup. A teaspoonful, in the inter-  
 mittents of children. *Bouchardat.*

## QUINIÆ MURIAS.

## MURIATE OF QUININE.

R. Sulphate of  
 quinine one ounce and a-half  
 Chloride of barium half an ounce  
 Dissolve separately, in boiling distilled  
 water; mix the solutions, filter, and eva-  
 porate to crystallization. *Guibourt.*  
 R. Diluted muriatic acid at will  
 Quinine sufficient  
 to saturate; evaporate, and crystallize.  
 Dose, from half a grain to a grain.  
*Cottureau.*

## MIXTURE OF MURIATE OF QUININE.

R. Muriate of quinine twelve grains  
 Diluted muriatic acid five minims  
 Distilled water seven fl. ounces  
 Syrup of orange flowers one fl. ounce  
 Mix. Dose, one fl. ounce. *Neligan.*  
 R. Muriate of quinine eight grains  
 Fennel water five fl. ounces  
 Muriatic ether one fl. drachm  
 Sugar half an ounce  
 Mix. A spoonful every two hours.  
*Radius.*

R. Muriate of quinine one scruple  
 Peppermint water one half fl. ounce  
 Mix. Twenty to sixty drops, every two  
 hours, in intermittents of children.  
*Radius.*

## QUINIÆ NITRAS.

## NITRATE OF QUININE.

R. Diluted nitric acid at will  
 Quinine sufficient  
 to saturate; boil with animal charcoal,  
 filter, evaporate, and let crystallize.  
*Taddei.*

## QUINIÆ PHOSPHAS.

## PHOSPHATE OF QUININE.

R. Quinine two parts  
 Water three parts  
 Boil, and add  
 Phosphoric acid sufficient  
 to saturate; filter while hot, and let crys-  
 tallize. *Turin Ph.*

## QUINIÆ SULPHAS.

## SULPHATE OF QUININE.

R. Yellow bark, in coarse  
 powder, four pounds







Extract of chamomile fifteen grains  
Mix, and make six pills. To be taken  
during the apyrexia of intermittent fevers.  
*Elliotson.*

#### PILLS OF SULPHATE OF QUININE AND GENTIAN.

R. Sulphate of quinine one scruple  
Extract of gentian two scruples  
Mix, and make twenty pills.

#### COMPOUND PILLS OF SULPHATE OF QUININE.

R. Calomel six grains  
Powdered opium three grains  
Sulphate of quinine twelve grains  
Syrup sufficient  
Beat into mass, and divide into twelve pills. One, night and morning, as alternative.  
*Ellis.*

R. Blue pill mass  
Sulphate of quinine  
Powdered aloes, each, twelve grains  
Aromatic syrup of rhubarb sufficient  
Beat into mass, and divide into twelve pills. One, three or four times a-day. This, or the last combination is suited to the condition following bilious remittent or intermittent fevers.  
*Ellis.*

R. Sulphate of quinine twelve grains  
Extract of gentian one scruple  
Compound rhubarb pill two scruples  
Blue pill mass six grains  
Mix, and make twelve pills. One, three times a-day.  
*Ryan.*

R. Sulphate of quinine ten to fifteen grains  
Dry phosphoric acid two scruples  
Powdered mallow root four scruples  
Extract of centaury sufficient  
Mix, and make sixty pills. Three or four pills, two or three times a-day, in nervous debility with tendency to abortion.  
*Radius.*

R. Powdered camphor one scruple  
Sulphate of quinine two scruples  
Pill aloes and myrrh one drachm and a-half  
Syrup of ginger sufficient  
Mix, and make forty pills. One, twice a-day.  
*Copland.*

#### SYRUP OF QUININE.

R. Sulphate of quinine thirty-two grains  
Distilled water two fl. drachms  
Alcoholized sulphuric acid six drops  
Dissolve, and add to  
Syrup sixteen fl. ounces  
*Par Cod.*

#### SYRUP OF QUININE AND COFFEE.

R. Ground roasted coffee four ounces  
Boiling water two pints four fl. ounces  
Make an infusion, let cool, and add  
Sugar four pounds  
Dissolve by means of a water-bath, and add  
Sulphate of quinine one drachm  
Dissolve in a little water acidulated with sulphuric acid.  
*Bories.*

#### MIXTURE OF SULPHATE OF QUININE.

R. Sulphate of quinine twenty grains  
Sulphuric acid one drop  
White sugar one drachm  
Cinnamon water two fl. ounces and a-half  
Mix. A teaspoonful every hour, in the apyrexia of intermittents.  
*Ellis.*

#### MIXTURE OF SULPHATE OF QUININE AND COFFEE.

R. Strong infusion of coffee five fl. ounces  
Sulphate of quinine twenty-four grains  
Sugar four drachms  
Mix. Dose, a tablespoonful. The coffee conceals the bitterness of the quinine.  
*Beasley.*

#### TINCTURE OF QUININE.

R. Sulphate of quinine one scruple  
Alcohol half a fl. ounce  
Dissolve. Dose, ten to twenty drops.  
*Ellis.*

#### COMPOUND TINCTURE OF QUININE.

R. Sulphate of quinine forty-eight grains  
Compound tincture of orange peel five and a-half fl. ounces  
Elixir of vitriol forty-five drops  
Mix. Dose, half a fl. drachm to two fl. drachms.  
*Copland.*

#### WINE OF QUININE.

R. Sulphate of quinine twelve grains  
Madeira wine two pints  
Dissolve. Dose, one or two fl. ounces.  
*Magendie.*

#### AROMATIC WINE OF QUININE.

R. Sulphate of quinine eighteen grains  
Citric acid fifteen grains  
Orange wine twenty-four fl. ounces  
Mix.  
*Collier.*

#### LINIMENT OF SULPHATE OF QUININE.

R. Sulphate of quinine half a drachm  
Tartar emetic six grains  
Extract of opium twelve grains  
Spirit of camphor eighteen fl. drachms  
Mix. Three fl. drachms to be rubbed on the epigastrium, three times a-day, in intermittents.  
*Schuster.*



## OINTMENT OF SULPHATE OF QUININE.

R. Sulphate of quinine	one ounce
Alcohol (.83)	
Sulphuric acid, each,	sufficient
Lard	four ounces

Rub together. Half an ounce to be rubbed into the groins, in malignant intermittents. *Antonini.*

## PLASTER OF SULPHATE OF QUININE.

R. Simple plaster	six drachms
Resin	two drachms

Melt together, and add  
Sulphate of quinine one drachm  
Oil of cajeput  
Camphor, each, one scruple  
Mix. To be applied to the epigastrium as a prophylactic in cholera. *Ammon.*

## ENEMA OF SULPHATE OF QUININE.

R. Sulphate of quinine	twelve to twenty grains
Flaxseed tea	four fl. ounces

Mix. To be used every four to six hours. *Ellis.*

## DENTIFRICE OF SULPHATE OF QUININE.

R. Sulphate of quinine	four grains
Prepared coral	one ounce
Carmin lake	eight grains
Essence of myrrh	two drops

Mix. *Pelletier.*

## GARGLE OF SULPHATE OF QUININE.

R. Sulphate of quinine	twelve grains
copper	sixteen grains
Elixir of vitriol	one fl. drachm
Water	eight fl. ounces

Mix. To be used three or four times a-day, in obstinate sore throat.

*Hartshorne.*

## QUININÆ SULPHO-TARTRAS.

## SULPHO-TARTRATE OF QUININE.

R. Sulphate of quinine	four drachms
------------------------	--------------

Tartaric acid four drachms and a-half  
Distilled water two fl. ounces  
Dissolve. Half a fl. drachm to a drachm, in the course of the day. *Righini.*

## QUININÆ TANNAS.

## TANNATE OF QUININE.

R. Sulphate of quinine	one part
Water	twenty parts
Diluted sulphuric acid	sufficient

Dissolve, filter, and add  
Infusion of galls sufficient  
to precipitate; collect precipitate, wash and dry it. *Greek Ph.*

Found useful in intermittent neuralgia.

## QUININÆ TARTRAS.

## TARTRATE OF QUININE.

R. Quinine	two parts
Water	three parts

Mix, boil, and add  
Tartaric acid sufficient  
to dissolve the quinine; filter whilst hot, and let crystallize. *Par. Cod.*

## QUININÆ VALERIANAS.

## VALERIANATE OF QUININE.

R. Fresh precipitated quinine	three parts
Valerianic acid	one part
Water	sixty parts

Mix; boil; filter while hot, and let crystallize; dry the crystals under 122° F. *Wettstein.*

Said to be more powerful than the sulphate, and not to cause nervous symptoms. It is given in solution, pill, enema, &c.

## R

## RESINA.

## RESIN.

*Resin*, or as it is vulgarly called *Rosin*, is the residuum after the distillation of the volatile oil from various species of the Pine tribe. In this state it is *Yellow resin*, or *Colophone*, which, when

melted and agitated with water, becomes of a whitish color, and forms *White resin*. Resin, when pure, is yellowish brown, inclining to olive or brown; it is solid, brittle, of a smooth, and shining fracture; of faint odor, and a somewhat acrid taste. It is principally used in medicine as an ingredient in plasters and ointments.



## HÆMOSTATIC POWDER.

R. Powdered resin four parts  
     gum Arabic  
     charcoal, each, one part

Mix. As an application to check external bleeding. *Guibourt.*

R. Resin two parts  
     Gum Arabic one part  
     Charcoal half a part

Pulverize well, and mix. Has proved useful in checking bleeding from wounds, &c. *Bonnefoux.*

## RESIN CERATE, OR BASILICON OINTMENT.

R. Resin five ounces  
     Lard eight ounces  
     Yellow wax two ounces

Melt together; strain through linen; and stir till cool. *U. S. Ph.*

As a stimulant application to blistered surfaces, burns, ulcers, &c.

## COMPOUND RESIN CERATE.

R. Resin  
     Suet  
     Yellow wax, each, one pound  
     Turpentine half a pound  
     Flaxseed oil half a pint

Melt together; strain through linen, and stir till cold. *U. S. Ph.*

Known as *Deshler's Salve*; is rather more stimulating than the last.

## RESIN, OR ADHESIVE PLASTER.

R. Powdered resin half a pound  
     Lead plaster three pounds  
 Melt the plaster by a gentle heat, and add the resin, mixing well. *U. S. Ph.*

## PLASTER OF ST. ANDREW.

R. White resin eight ounces  
     Elemi two ounces  
     Venice turpentine  
     Oil of bay laurel, each, one ounce  
 Melt together, and strain. An adhesive plaster. *Foy.*

## RHEUM.

## RHUBARB.

Rhubarb is the product of one or more species of *Rheum*, but of which, is not known with any certainty. All that is clearly ascertained is, that they grow in the deserts of Tartary.

*Sex. Syst.* Enneand. trigyn. *Nat. Syst.* Polygonacea.

*U. S. Dispens.* 587. *Griffith, Med. Bot.* 539.

Two kinds of Rhubarb are usually found in commerce, the Russian or Turkey, and the Chinese, the first of which is the most valuable, but the latter is the most generally used, and when

good, fulfils all that is required. They both are purgatives, with some tonic and astringent powers, and are much used in various forms of disease. The dose, in substance, to produce a full effect, is from twenty to thirty grains. They are given in a vast variety of forms.

## POWDER OF RHUBARB AND MAGNESIA.

R. Powdered rhubarb one scruple  
     Magnesia ten grains

Mix. To be given in syrup, or sugar and water. *Ellis.*

## POWDER OF RHUBARB AND CHALK.

R. Powdered rhubarb fifteen grains  
     Compound chalk

powder twenty-four grains  
 Mix. In the evening, in cardialgia. *Foy.*

## POWDER OF RHUBARB AND SULPHATE OF POTASSA.

R. Powdered rhubarb one drachm  
     sulphate of  
     potassa two drachms

Mix. Ten grains to a drachm, every morning. *Fordyce.*

## COMPOUND POWDERS OF RHUBARB.

R. Magnesia  
     Cremor tartar  
     Powdered rhubarb  
     chamomile  
     Oleo-sacch. of  
     fennel, each, half an ounce

Mix. A teaspoonful, twice or thrice a-day, in obstructions of the abdominal viscera. *Selle.*

R. Powdered rhubarb thirty grains  
     sulphate of potassa  
     chamomile, each, one drachm

Mix, and divide into six powders. One, twice a-day, in dyspepsia with torpor of the bowels. *Ellis.*

R. Powdered rhubarb four ounces  
     Magnesia one pound  
     Powdered ginger two ounces

Mix well, and keep in well-closed bottles. *Ed. Ph.*

A good antacid laxative. Dose, five grains to a drachm, according to age.

## ROASTED RHUBARB.

R. Coarsely powdered rhubarb at will  
 Place it in a shallow iron dish, heat regularly, so as to scorch the powder, which should be so constantly stirred, that the influence of the heat should be uniform throughout; when the color has changed to brown, the process is to be stopped, and the rhubarb reduced to fine powder, for use. *W. Procter.*

R. Powdered rhubarb at will



Treat in an iron vessel, constantly stirring, till it becomes almost black, then smother it in a covered jar. Dose, five to ten grains, as an astringent in diarrhoea.

*Hoblyn.*

#### PILLS OF RHUBARB.

R. Powdered rhubarb six drachms  
Soap two drachms  
Beat them with water, so as to form a mass, and divide into one hundred and twenty pills. *U. S. Ph.*

Each pill contains three grains.

#### COMPOUND RHUBARB PILLS.

R. Powdered rhubarb one ounce  
aloes six drachms  
myrrh half an ounce  
Oil of peppermint half a fl. drachm  
Syrup of orange peel sufficient  
Beat together into mass, and divide into two hundred and forty pills. *U. S. Ph.*

A warm, tonic laxative, useful in costiveness with debility of the stomach. Dose, two to four a-day.

R. Powdered rhubarb one drachm and a-half  
Sulphate of iron half a drachm  
Soap two scruples  
Distilled water sufficient  
Beat into mass, and divide into forty pills. In similar cases as last, three or four to be taken at bedtime. *Griffitts.*

#### PILLS OF RHUBARB AND IPECACUANHA.

R. Powdered rhubarb one scruple  
ipecacuanha ten grains  
Opium three grains  
Oil of cinnamon five drops  
Gum Arabic sufficient  
Triturate together, and divide into ten pills. One to be given every two or three hours, in dysentery, to relieve tormina and tenesmus. *Chapman.*

#### PILLS OF RHUBARB AND CARAWAY.

R. Powdered rhubarb two drachms  
Syrup one fl. drachm  
Oil of caraway ten minims  
Mix, and divide into forty pills. *Kitchener.*

#### PILLS OF RHUBARB AND CHAMOMILE.

R. Powdered rhubarb  
aloes  
myrrh  
Extract of chamomile, each, one drachm  
Oil of chamomile twelve drops  
Mix, and divide into sixty pills.

These are known as *Speediman's pills.* *Beasley.*

R. Powdered rhubarb  
ginger, each, half a drachm

Extract of chamomile one drachm  
Beat together, and divide into thirty pills; three to be taken before each meal. Have been recommended in dyspepsia and chlorosis. *A. T. Thomson.*

#### PILLS OF RHUBARB AND SODA.

R. Powdered rhubarb  
Carbonate of soda  
Extract of gentian, each, one drachm and a-half  
Mix, and make sixty pills. *Guy's Hosp.*  
R. Powdered rhubarb  
Dried carbonate of soda  
Extract of gentian, each, one scruple  
Calomel three grains  
Mix, and make twenty pills. Two, occasionally, in dyspepsia. *Ellis.*

#### PILLS OF RHUBARB AND OX GALL.

R. Inspissated ox gall  
Gum ammoniac  
Powdered rhubarb equal parts  
Mix, and form pills of two grains each. *Leipsic Ph.*

In constipation, deficiency of bile, &c.

#### LOZENGES OF RHUBARB.

R. Powdered rhubarb  
Cremor tartar, each, two drachms  
Fresh orange peel half a drachm  
Sugar, dissolved in  
Orange-flower water four ounces  
Mix, and make lozenges of eighteen grains. *Beral.*

#### SUPPOSITORY OF RHUBARB.

R. Extract of rhubarb half a drachm  
Soap three drachms  
Powdered rhubarb sufficient  
Mix, and make three suppositories. *Radiis.*

#### ELECTUARY OF RHUBARB.

R. Powdered rhubarb one drachm and a-half  
Sulphate of potassa one drachm  
Cremor tartar half an ounce  
Pulp of tamarinds two ounces  
Mix. A teaspoonful. *Saunders.*

#### INFUSION OF RHUBARB.

R. Bruised rhubarb one drachm  
Boiling water eight fl. ounces  
Infuse for two hours, and strain. *Guibourt.*

One or two fl. ounces, every three or four hours, till it operates.

R. Bruised rhubarb one ounce  
Boiling water eighteen fl. ounces  
Infuse for twelve hours, in a covered vessel; add



Spirit of cinnamon two fl. ounces  
and strain. Dose, as above. *Ed. Ph.*

#### ALKALINE INFUSION OF RHUBARB.

R. Bruised rhubarb two drachms  
Carbonate of potassa one drachm  
Boiling water half a pint  
Infuse for four hours, strain, and add  
Tincture of cinnamon half a fl. ounce  
*Copland.*

#### EXTRACT OF RHUBARB.

R. Powdered rhubarb fifteen ounces  
Proof spirit one pint  
Distilled water seven pints  
Macerate for four days by a gentle heat;  
strain; set aside to settle; decant, and  
evaporate to proper consistence. *Dub. Ph.*  
Dose, ten to thirty grains.

#### FLUID EXTRACT OF RHUBARB.

R. Rhubarb eight ounces  
Sugar five ounces  
Diluted alcohol two pints  
Powder the rhubarb coarsely, and mix it  
with its bulk of coarse sand; moisten the  
mixture with sufficient of the diluted alco-  
hol to form a pasty mass; let stand until  
the particles have absorbed, and are swol-  
len by the fluid. Then introduce into a  
displacer, and shake till uniformly settled,  
and cover with cloth or paper; pour on  
the rest of the alcohol, until the product  
has but little odor or taste of the root.  
The tincture thus obtained, is to be eva-  
porated in a water-bath to five and a-half  
fl. ounces; the sugar added; when the  
whole should measure eight fl. ounces.  
Dose, fifteen drops to half a drachm.

*W. Procter.*

#### SYRUP OF RHUBARB.

R. Rhubarb, bruised, two ounces  
Boiling water one pint  
Macerate for twenty-four hours, and  
strain, then add  
Sugar two pounds  
and make syrup. *U. S. Ph.*

Mild cathartic. Dose, for young children, one  
to two fl. drachms.

#### AROMATIC SYRUP OF RHUBARB.

R. Rhubarb, bruised, two ounces and a-half  
Cloves, bruised,  
Cinnamon, bruised, each, half an ounce  
Nutmeg, bruised, two drachms  
Diluted alcohol two pints  
Macerate for fourteen days, and strain;  
evaporate on water-bath to one pint, and  
add

Syrup six pints  
previously heated. *U. S. Ph.*

Warm laxative in cases of children. Dose,  
one fl. drachm, in bowel complaints, to be re-  
peated every two hours.

#### SYRUP OF RHUBARB AND SENNA.

R. Bruised rhubarb one ounce  
Senna two ounces  
Fennel seed  
Bruised cinnamon, each, two drachms  
Boiling water two pints and a-half  
Macerate for twelve hours, strain, and  
add  
Sugar three pounds  
Make syrup. *Ed. Ph.*

#### TINCTURE OF RHUBARB.

R. Rhubarb, bruised, three ounces  
Cardamoms, bruised, half an ounce  
Diluted alcohol two pints  
Macerate for fourteen days, express, and  
filter. *U. S. Ph.*

Dose, as purgative, half to one fl. ounce; as  
stomachic and tonic, one to two fl. drachms.

#### TINCTURE OF RHUBARB AND GENTIAN.

R. Rhubarb, bruised, two ounces  
Gentian, bruised half an ounce  
Diluted alcohol two pints  
*U. S. Ph.*

Macerate for fourteen days, express, and filter.  
In flatulent colic, &c. Dose, as above.

R. Rhubarb, bruised two ounces  
Gentian, bruised half an ounce  
Virginia snake-  
root one drachm and a-half  
Diluted alcohol two pints  
Macerate for three days, express, and filter.  
*Van Mons.*

#### TINCTURE OF RHUBARB AND ALOES.

R. Rhubarb, bruised, ten drachms  
Aloes, powdered, six drachms  
Cardamoms, bruised, half an ounce  
Diluted alcohol two pints  
Macerate for fourteen days, express, and  
filter. *U. S. Ph.*

This was formerly known as *Elixir sacrum*,  
and was much used.

#### TINCTURE OF RHUBARB AND SENNA.

R. Rhubarb, bruised, one ounce  
Senna two drachms  
Coriander, bruised,  
Fennel, bruised, each, one drachm  
Red saunders, rasped, two drachms  
Saffron  
Extract of liquorice, each, half a drachm  
Raisins, stoned, half a pound  
Diluted alcohol three pints



Macerate for fourteen days, express, and filter. *U. S. Ph.*

Well known and much used under the name of *Warner's gout cordial*, as a stomachic and slight purgative. Dose, from half to two fl. ounces.

#### WARNER'S CORDIAL.

R. Contused rhubarb one ounce  
Senna half an ounce  
Saffron one drachm  
Fennel seed  
Coriander, each, two drachms  
Extract of liquorice four drachms  
Raisins, pounded, one pound  
Brandy three pints

Mix, and macerate for a week, and strain.  
Dose, half a wineglassful, in atonic gout, with flatulence. *Chapman.*

#### TINCTURE OF RHUBARB AND ANISEED.

R. Rhubarb, bruised,  
Liquorice root, bruised, each, two ounces  
Aniseed, bruised,  
Sugar, each, one ounce  
Diluted alcohol two pints

Macerate for fourteen days, express, and filter. *Copland.*

Used as the tincture.

#### ALKALINE TINCTURE OF RHUBARB.

R. Rhubarb, bruised, one ounce and a-half  
Carbonate of potassa three drachms  
Boiling water twelve fl. ounces  
Macerate for twelve hours, strain, and add  
Tincture of cinnamon two fl. ounces  
*Prus. Ph.*

As a mild purgative, where there is acidity of stomach, in doses of half to two fl. ounces.

#### WINE OF RHUBARB.

R. Rhubarb, bruised, two ounces  
Canella, bruised, one drachm  
Wine one pint  
Macerate for fourteen days, occasionally agitating, express, and filter. *U. S. Ph.*

A warm, cordial laxative, in doses of one to four fl. drachms.

R. Rhubarb, bruised, two ounces and a-half  
Cardamoms, bruised, half an ounce  
Saffron two drachms  
Wine two pints  
Alcohol eight fl. ounces  
Macerate for ten days, express, and filter.  
*Par. Cod.*

Used as the last.

#### WINE OF RHUBARB AND GENTIAN.

R. Rhubarb, bruised, one ounce  
Gentian, bruised, two drachms  
Canella one drachm

Wine one pint  
Macerate for three days, express, and filter.  
*Swed. Ph.*

#### COMPOUND WINE OF RHUBARB.

R. Rhubarb, bruised,  
Orange peel, bruised, each, half an ounce  
Wine two pints  
Macerate for twenty hours, strain, and add  
Oleo-sacch. of mace one ounce  
Hoffmann's elixir one drachm  
As a stomachic, a tablespoonful three times a-day. *Phæbus.*

#### RHUBARB MIXTURE.

R. Bruised rhubarb one drachm  
fennel one scruple  
Sulphate of magnesia half an ounce  
Manna one ounce  
Boiling water six fl. ounces  
Infuse and strain, when cold. As a mild purgative; one-half to be taken at a dose. *Radiis.*

R. Infusion of rhubarb one fl. drachm and a-half  
Spirit of hartshorn two fl. scruples  
Cinnamon water one fl. ounce and a-half  
Syrup of opium six fl. drachms  
Mix. A teaspoonful every two hours, in the chronic diarrhœa of young children. *Vogt.*

R. Powdered rhubarb one drachm  
Carbonate of soda two drachms  
Tincture of orange peel one fl. ounce and a-half  
Decoction of liquorice ten fl. ounces and a-half  
Mix. Dose, half a fl. ounce to one fl. ounce, three times a-day. *Guy's Hosp.*

R. Powdered rhubarb one drachm  
Magnesia one drachm and a-half  
Powdered ginger one scruple  
Peppermint water one pint  
Mix. Dose, half a fl. ounce. *Gregory.*

R. Powdered rhubarb two scruples  
Tartrate of potassa one ounce  
Peppermint water six fl. ounces  
Tincture of senna  
Syrup of ginger, each, half a fl. ounce  
Mix. Dose, one fl. ounce. *Brande.*

#### RHŒAS.

##### RED POPPY.

The *Papaver rhœas*, or Red Poppy, is a native of Europe, where it is found in great abundance in cultivated grounds, and has become naturalized in many places in this country.



*Ser. Syst.* Polyand. monog. *Nat. Syst.* Papaveraceæ.

Linn. Sp. Pl. 726. Woodville, Med. Bot. 387.

The parts used are the petals; these have a mucilaginous, somewhat bitter taste, and a narcotic smell, which latter is lost on drying. Their action on the system is slight, but they are used in Europe in the preparation of a syrup, prescribed as an anodyne in the catarrhal affections of children.

R. Red poppy petals                      one pound  
Water    one pint

Heat the water on a water-bath, gradually add the petals, remove from the fire, macerate for twelve hours, express, and strain, and add

Sugar                      two pounds and a-half  
And make syrup.                      *Lond. Ph.*

#### INFUSION OF RED POPPY.

R. Red poppy petals                      one drachm  
Boiling water                                      one pint  
Infuse and strain. To be taken freely, in catarrhal affections.                      *Cotteneau.*

#### COMPOUND INFUSION OF RED POPPY.

R. Red poppy petals                      two ounces  
Diluted sulphuric acid                      fifteen drops  
Sugar    two ounces  
Decoction of barley                      one pint  
Infuse, and strain. Used as above.                      *St. Marie.*

### ROSA CANINA.

#### DOG ROSE.

This shrub is a native of Europe, growing in hedges and thickets.

*Ser. Syst.* Icosand. polyg. *Nat. Syst.* Rosaceæ.

Linn. Sp. Pl. 704. Woodville, Med. Bot. 493.

The fruit, which is the officinal portion, is smooth, oval, red, and of a pleasant acidulous taste. It is principally used in confection.

#### CONFECTION OF DOG ROSE.

R. Pulp of fruit of dog rose                      one pound  
Sugar    twenty ounces  
Heat the pulp gently and gradually, add the sugar, and rub together till well mixed.                      *Lond. Ph.*

Has been advised in diarrhœa and dysentery, as an astringent, but is principally used as a pill basis.

### ROSA CENTIFOLIA.

#### HUNDRED-LEAVED ROSE.

This species, of which there are innumerable varieties, is cultivated in every garden, but its native country is unknown, though probably it is from Persia.

The petals are the only part used; these are fragrant, and have an acidulous, somewhat bit-

terish taste. They are slightly laxative, but are principally used for the making of rose water.

#### ROSE WATER.

R. Fresh hundred-leaved  
rose petals                      eight pounds  
Water    two gallons  
Mix, and distil one gallon.                      *U. S. Ph.*

#### OINTMENT OF ROSE WATER.

R. Rose water  
Oil of almonds, each,                      two fl. ounces  
Spermaceti                                      half an ounce  
White wax                                      one drachm  
Melt together, on a water-bath, the last three ingredients, and stir in the rose water till cold.                      *U. S. Ph.*

Much used, under the name of *Cold cream*, as a cooling application to irritated surfaces.

#### ROSE LOZENGES.

R. Powdered sugar  
Rose water, each,                      six ounces  
Evaporate gently, to a thick syrup, and add  
Coarsely powdered sugar                      one pound  
Oil of roses                      one drachm and a-half  
When dissolved, pour in drops, on a cold, oiled, marble slab, and dry.                      *Cotteneau.*

#### COLLUTORY OF ROSE WATER.

R. Rose water                      three fl. ounces  
Cream  
White of egg  
Syrup of violets, each,                      one fl. ounce  
Mix.    *St. Marie.*

#### OIL OF ROSES.

R. Fresh rose leaves                      at will  
Water    sufficient  
to moisten the leaves; distil, and collect the oil that floats on the product.                      *Guibourt.*

#### ESSENCE OF ROSES.

R. Pounded rose leaves  
Alcohol (.874), each,                      ten parts  
Infuse for twenty-four hours, by a gentle heat, and distil.                      *Taddei.*

#### ROSE OIL.

R. Pounded rose leaves                      one part  
Olive oil    four parts  
Digest with gentle heat for four days, express, and strain; repeat process with fresh leaves, a second and third time; separate the water, and filter.                      *Soubeiran.*

As an application to chilblains and irritated surfaces.

### ROSA GALLICA.

#### RED ROSE.

This species is a native of the south of Eu-



rope, and is generally cultivated in our gardens. The varieties are very numerous.

Linn. Sp. Pl. 704. Griffith, Med. Bot. 273.

The parts used in medicine are the petals of the unexpanded flowers, deprived of their white claws, and dried. In this state they are of a purplish red color; of an agreeable odor, and bitterish, astringent taste. They have mild astringent properties, but are more used as a vehicle, than for their own powers.

#### CONFECTION OF ROSES.

R. Red roses, in powder four ounces  
Powdered sugar thirty ounces  
Clarified honey six ounces  
Rose water eight fl. ounces  
Rub the roses with the rose water, at a boiling heat; add gradually the sugar and honey, and beat together. *U. S. Ph.*

Principally used as a pill basis.

#### ACID INFUSION OF ROSES.

R. Red roses half an ounce  
Boiling water forty-eight fl. ounces  
Diluted sulphuric acid three fl. drachms  
Digest in a glass vessel for half an hour; express and strain, and add  
Sugar one ounce and a-half  
As a cooling drink, mixed with water. *Dub. Ph.*

#### HONEY OF ROSES.

R. Red roses two ounces  
Clarified honey two pints  
Boiling water a pint and a-half  
Macerate the roses in the water for two hours, and strain; add the honey, and evaporate to proper consistence. *U. S. Ph.*

As an addition to gargles.

#### ELECTUARY OF ROSES.

R. Confection of roses sixteen parts  
Syrup of tolu four parts  
poppy one part  
Mix. Used as an astringent in bowel affections, in doses of a teaspoonful. *Foy.*

#### TINCTURE OF ROSES.

R. Red roses five ounces  
Alcohol two fl. ounces  
Rose water eight fl. ounces  
Digest for four days, express, and filter; digest residuum with half a pint of alcohol for three days; express, filter, and mix the liquors. *Squire.*

Used principally as a perfume.

#### WINE OF ROSES.

R. Red roses one part  
Red wine sixteen parts  
Infuse for half an hour, and strain. As an application to indolent ulcers. *Beral.*

#### COMPOUND VINEGAR OF ROSES.

R. Red roses one ounce and a half  
Sumach leaves one ounce  
Wine of quinces four fl. ounces  
Vinegar of red wine ten fl. ounces  
Macerate for ten days, express, and filter. As an application to contusions, and as a stypic. *Van Mons.*

#### SYRUP OF RED ROSES.

R. Red roses two ounces  
Boiling water one pint  
Sugar twenty ounces  
Macerate the rose leaves in the water for twelve hours, strain, evaporate, add sugar, and form syrup. *Ed. Ph.*

### ROSMARINUS.

#### ROSEMARY.

This is the *Rosmarinus officinalis*, a small evergreen shrub, a native of the south of Europe, and generally cultivated in gardens elsewhere.

*Sax. Syst.* Diand. monog. *Nat. Syst.* Lamiaceæ.

Linn. Sp. Pl. 33. Griffith, Med. Bot. 507.

The officinal portions are the flowering extremities. These have a fragrant odor, and a bitter, pungent taste. Rosemary is stimulant and carminative, but is principally employed as an aromatic addition to lotions and liniments, as an ingredient in articles of perfumery.

#### AROMATIC BATH.

R. Rosemary  
Thyme  
Sage  
Origanum  
Mint, each, half a pound  
Boiling water six pints  
Macerate for twelve hours, strain, and add  
Essence of soap four ounces  
Muriate of ammonia two ounces  
Pour the whole into water sufficient for a bath. *Foy.*

In chronic rheumatism, cutaneous affections, dyspepsia, &c.

#### AROMATIC FOMENTATION.

R. Rosemary half an ounce  
Red wine  
Water, each, three fl. ounces  
Infuse, and express. As a fomentation in contusions. *Augustin.*

#### AROMATIC VINEGAR.

R. Rosemary  
Sage, each, one ounce  
Lavender one ounce and a-half  
Cloves half a drachm  
Vinegar two pints



Infuse for eight days, and strain. As a lotion in contusions, strains, &c.

*Spielman.*

#### TINCTURE OF ROSEMARY.

R. Rosemary one part  
Spirit of rosemary four parts  
Macerate, express, and filter.

*Brunsw. Ph.*

#### OIL OF ROSEMARY.

R. Rosemary at will  
Water sufficient  
Distil, and collect the oil that floats on the product. Principally used in stimulating liniments, but sometimes given as carminative in doses of two or three drops.

*U. S. Ph.*

#### ROSEMARY OIL.

R. Rosemary one part  
Olive oil three parts  
Heat till the water has evaporated, express, and strain. As a stimulating application.

*Span. Ph.*

#### SPIRIT OF ROSEMARY.

R. Oil of rosemary two drachms  
Alcohol one gallon  
Water one pint  
Mix, and distil one gallon, by a slow fire.

*U. S. Ph.*

As an ingredient in lotions, and liniments, &c.

#### HUNGARY WATER.

R. Rosemary three parts  
Lavender one part  
Diluted alcohol six parts  
Water twelve parts  
Mix, and distil three-fourths. Used as a perfume.

*Van Mons.*

### RUBIA.

#### MADDER.

The *Rubia tinctorum* or Madder is a perennial plant, native of the south of Europe, and extensively cultivated in many countries for the sake of its roots, which are used in dying reds.

*Sex. Syst.* Tetrand. monog. *Nat. Syst.* Rubiaceæ.

Linn. Sp. Pl. 158. Griffith, Med. Bot. 382.

The root is the officinal portion, and as found in commerce, is in long, cylindrical pieces, about as thick as a quill, of a deep red or brown color. It has a peculiar and unpleasant odor, and a bitterish, astringent taste. It was much used formerly as a diuretic, but is at present but seldom prescribed, except as an emmenagogue, for which purpose it is highly spoken of by many practitioners.

#### COMPOUND POWDER OF MADDER.

R. Powdered madder  
sugar, each, two drachms

Powdered sulphate of

potassa half an ounce

Mix well. Dose, ten grains to a scruple, in rachitis.

*Radius.*

#### DECOCTION OF MADDER.

R. Powdered madder one ounce  
Boiling water one pint  
Simmer for fifteen minutes, and add  
Bruised cloves one drachm  
When cool, strain. A wineglassful every three hours, a short time previous to the expected time of the menstrual discharge.

*Dewees.*

R. Powdered madder one ounce  
mace two drachms  
Water one pint and a-half  
Boil down to one pint, strain, and add  
Aromatic tincture two fl. drachms  
Syrup of citric acid two fl. ounces  
Dose, three fl. ounces, three or four times a-day, in amenorrhœa, chlorosis, &c.

*Swedjaur.*

R. Powdered madder half an ounce  
hops one drachm  
English walnut leaves three  
Water two pints  
Boil down to a pint and a-half, strain, and add

Tincture of tartrate

of iron a teaspoonful

Advised in scrofula, in doses of two ounces, night and morning.

*St. Marie.*

### RUBUS TRIVIALIS.

#### DEWBERRY.

### RUBUS VILLOSUS.

#### BLACKBERRY.

These two species of shrubby plants are very common in the United States, and are well known for their pleasant fruit.

*Sex. Syst.* Icosand. polyg. *Nat. Syst.* Rosaceæ.

Torrey and Gray, i. 454, 455. Griffith, Med. Bot. 275, 276.

The officinal portions are the roots; these are identical in their effects; they have a very faint odor, and a bitter, astringent taste. The smaller are to be preferred. They are efficient, but mild astringents, and have been found very useful in a relaxed condition of the bowels.

#### DECOCTION OF BLACKBERRY ROOT.

R. Small roots of blackberry one ounce  
Water one pint and a-half  
Boil down to one pint, and strain. Dose, one or two fl. ounces, three or four times a-day, in diarrhœa, &c.



SYRUP OF BLACKBERRY.

R. Juice of blackberries twenty ounces  
 Sugar three pounds  
 Boil, and strain. Said to be very useful  
 in dysentery.

RUTA.

RUE.

The only species recognized in medicine is the *Ruta graveolens*, a small shrub with glaucous leaves and yellow flowers, a native of the south of Europe, but generally cultivated in gardens.

*Ser. Syst.* Decand. monog. *Nat. Syst.* Rutaceæ.

Linn. Sp. Pl. 523. Griffith, Med. Bot. 189.

The leaves are the part usually employed; they have a nauseous odor, and a bitter, acrid, disagreeable taste. They are stimulant and antispasmodic, and have been much used as an emmenagogue and vermifuge. The dose of the powder is from ten to twenty grains, two or three times a-day.

CONFECTION OF RUE.

R. Dried rue  
 Caraway  
 Laurel berries, each, one ounce and  
 a-half  
 Sagapenum half an ounce  
 Black pepper two drachms  
 Clarified honey sixteen ounces  
 Mix. *Dub. Ph.*

It is used as antispasmodic, in the form of an

enema made with a scruple to a drachm, mixed with half a pint of warm, mucilaginous fluid.

EXTRACT OF RUE.

R. Dried rue two pounds  
 Alcohol (.927) seven pints  
 Exhaust the rue by the process of displacement, distil the tincture, and evaporate the residue. *Wert. Ph.*

MIXTURE OF RUE AND SQUILL.

R. Juice of rue leaves one ounce  
 Oxy-mel of squill half an ounce  
 Mix. A teaspoonful occasionally, in hysteric affections. *Pierquin.*

SYRUP OF RUE.

R. Coarsely powdered sugar fifteen ounces  
 Tincture of rue two fl. ounces  
 Distilled water seven fl. ounces  
 Dissolve, and strain. *Beral.*

TINCTURE OF RUE.

R. Expressed juice of rue  
 Alcohol (.847), each, four fl. ounces  
 Mix, and filter at the end of twenty-four hours. *Beral.*

MIXTURE OF RUE.

R. Rue  
 Savin  
 Wormwood, each, three drachms  
 Boiling water one pint  
 Infuse, strain, and add  
 Castor oil half an ounce  
 As an injection, in cases of ascarides. *Radius.*

S

SABADILLA.

CEVADILLA.

This article is the seed of one or more plants belonging to the natural order of Melanthaceæ; the U. S. Ph. attributes it to *Veratrum sabadilla*, and the Lond. Ph. to *Helonias (Asagrea) officinale*.

The Cevadilla of the shops consists of follicles, seeds, &c. The first are ovate, oblong, acuminate, of a pale, yellowish-brown color, and a thin, papery consistence. The seeds are small, curved, acuminate, shining, wrinkled, and of a blackish-brown color. They have very little odor, but their taste is bitter and acrid. These seeds are a drastic emeto-cathartic, and have been used as an anthelmintic, and in nervous diseases, &c.; but are principally used in the preparation of veratria. The dose is from five to ten grains, in powder; but it should always be given with much caution.

COMPOUND POWDER OF CEVADILLA.

R. Powdered cevadilla two grains  
 Sulphate of iron one grain  
 Powdered semen contra  
 Magnesia, each, half a scruple  
 Mix. To be given four times a-day, in cases of tenia. *Radius.*

TINCTURE OF CEVADILLA.

R. Seeds of cevadilla, bruised, at will  
 Alcohol sufficient  
 to cover them; digest for ten days, and filter. As a lotion in rheumatism. *Turnbull.*

EXTRACT OF CEVADILLA.

R. Tincture of cevadilla at will  
 Evaporate to proper consistence. Dose,







## OIL OF SAVINE.

R. Savine at will  
Water sufficient  
Distil, and collect the oil that floats on the product. *Guibourt.*

A powerful stimulant, emmenagogue, and rubefacient. Dose, two to five drops.

## INFUSION OF SAVINE.

R. Savine one drachm  
Boiling water eight fl. ounces  
Infuse for half an hour, and strain. Dose, half a fl. ounce. *Pereira.*

R. Savine one drachm  
Camphor six grains  
Boiling water five fl. ounces  
Infuse, and strain. *Horn.*

## TINCTURE OF SAVINE.

R. Savine one part  
Alcohol four parts  
Macerate for ten days, and filter. Dose, ten to fifteen drops. *Soubeiran.*

## COMPOUND TINCTURE OF SAVINE.

R. Extract of savine one ounce  
Tincture of castor sixteen fl. ounces  
Tincture of myrrh eight fl. ounces  
Digest till dissolved. As an emmenagogue. *Lond. Ph., 1788.*

R. Compound tincture of savine one fl. ounce  
Tincture of black hellebore half a fl. ounce  
castor two fl. drachms  
Mix. Forty drops, three times a-day. *Ellis.*

## SAVINE CERATE.

R. Powdered savine two ounces  
Resin cerate one pound  
Melt the cerate, and mix in the savine. *U. S. Ph.*

As a dressing to keep up the discharge of blisters, &c.

## SAVINE OINTMENT.

R. Powdered savine  
Lard equal parts  
Mix. As an application to venereal warts. *Radiis.*

## SACCHARUM.

## SUGAR.

Sugar is procured from the *Saccharum officinarum* and other species, and is used in several forms, as molasses, brown, and refined sugar, the latter of which only is recognized in the U. S. Pharm.

## SYRUP.

R. Sugar two pounds and a-half

## Water

one pint

Dissolve the sugar in the water, by the aid of heat, remove any scum that may form, and strain while hot. *U. S. Ph.*

## CAMEL OR BURNT SUGAR.

R. Sugar at will  
Place in a shallow vessel, and heat over a gentle fire, till it assumes a dark brown color. Used as a coloring for spirits, &c. *Gray.*

## SAGAPENUM.

## SAGAPENUM.

This is a gum resin, derived most probably from some plant belonging to the *Apiaceæ*. It is in irregular masses composed of cohering fragments, of a yellowish brown, olive, or reddish yellow color, of a waxy consistence; of a somewhat alliaceous odor, and a pungent, bitterish, unpleasant taste. It is a mild stimulant, somewhat resembling assafetida in its properties, but inferior to it. It has been used as an emmenagogue and antispasmodic, but it is now seldom prescribed internally, but is still employed as an external application as a descutient.

## COMPOUND SAGAPENUM PILLS.

R. Sagapenum one ounce  
Aloes half a drachm  
Syrup of ginger sufficient  
Beat together. Dose, ten to thirty grains. *Lond. Ph.*

As a stimulant, antispasmodic laxative, in flatulent colic, &c.

## SAGO.

## SAGO.

This is the prepared fecula of the pith of various species of *Palmaceæ* and *Cycadaceæ*. It is prepared in two forms, meal and pearl sago, the latter of which is most generally employed. This is in small, hard, whitish or brownish grains, inodorous, and of a mucilaginous taste. It is insoluble in cold water, but by long boiling forms a glutinous solution. It is nutritive, easily digestible, and forms an excellent article of diet for the sick and convalescents.

## SAGO MUCILAGE.

R. Sago one ounce  
Water one pint  
Macerate by a gentle heat, for two hours, then boil for fifteen minutes, stirring continually, till the grains are perfectly dissolved, add sugar or flavoring, according to circumstances. *A. T. Thomson.*

## SAGO POSSET.

R. Sago two ounces  
Water two pints



Macerate for two hours, by a gentle heat, then boil, till a mucilage is formed; then rub half an ounce of sugar on the rind of a lemon, add this and one fl. drachm of tincture of ginger to half a pint of sherry wine, pour the whole into the sago mucilage, and boil for five minutes. Useful in convalescence to restore strength. A wineglassful to be taken every four or five hours.

A. T. Thomson.

#### SAGO MILK.

R. Sago one ounce  
Water one pint  
Macerate for half an hour, then add  
Milk one pint and a-half  
Boil slowly, till the sago is perfectly dissolved.

A. T. Thomson.

#### SALICINA.

##### SALICIN.

R. Decoction of willow bark at will  
Treat with slaked lime, filter, and evaporate to consistence of syrup; add alcohol, again filter, evaporate, and let crystallize; wash the crystals with cold water.

Cottureau.

Dose, four to six grains every three hours, in apyrexia of intermittents.

##### PILLS OF SALICIN.

R. Salicin  
Extract of gentian, each, twelve grains  
Liquorice powder sufficient  
Mix, and make six pills.

Foy.

##### COMPOUND SALICIN PILLS.

R. Salicin one scruple  
Compound rhubarb pill two scruples  
Mix, and make twelve pills.

Beasley.

##### COMPOUND POWDER OF SALICIN.

R. Salicin two scruples  
Aromatic powder one drachm  
Mix, and divide into twelve powders.

Neligan.

R. Salicin fifteen grains  
Tartar emetic one grain  
Powdered sugar fifty grains  
Mix, and divide into ten powders. One, thrice a-day.

Krombolz.

#### SALIX.

##### WILLOW.

Many species of willow have been used in medicine, and are recognized in the different Pharmacopœias; that admitted by the U. S. Ph. is the *Salix alba*, a small tree, a native of Eu-

rope, but extensively cultivated in the United States.

Sex. Syst. Diœcia diand. Nat. Syst. Sabicacœe.

Linn. Sp. Pl. 1449. Lindley, Flor. Med. 318.

The officinal portion is the bark; this is thin, flexible, fibrous pieces, of a brown color. It has a peculiar, aromatic odor, and an astringent, bitter taste. It is tonic and astringent, and has been used as a substitute for Peruvian bark.

##### COMPOUND POWDER OF WILLOW BARK.

R. Powdered willow bark  
horsechestnut bark  
gentian  
calamus  
avens root equal parts  
In intermittent fevers, in drachm doses.

Hufeland.

##### WILLOW BARK DENTIFRICE.

R. Powdered willow bark  
charcoal, each, half an ounce  
Extract of myrrh two drachms  
Balsam of Peru half a scruple  
Oil of cinnamon three drops  
Triturate well together.

Phœbus.

##### DECOCTION OF WILLOW BARK.

R. Contused willow  
bark one ounce and a-half  
Water fifteen fl. ounces  
Boil down to one-half.

Nieman.

##### OINTMENT OF WILLOW LEAVES.

R. Expressed juice of willow  
leaves two fl. ounces  
Lard sufficient  
Mix, and heat till moisture is driven off.  
Useful as a dressing to foul ulcers.

Hufeland.

#### SALVIA.

##### SAGE.

Though many species of *Salvia* possess analogous remedial properties, one only is recognized by the U. S. Ph.—the *Salvia officinalis*, a perennial plant, native of the south of Europe, but generally cultivated in our gardens, flowering in June, at which time it should be collected.

Sex. Syst. Diand. monog. Nat. Syst. Lamiacœe.

Linn. S. Pl. 34. Griffith, Med. Bot. 505.

The leaves are the officinal portion. These have a strong, fragrant odor, and a warm, bitterish, aromatic, somewhat astringent taste. Sage is stimulant, with some astringent and tonic powers. It is principally used in the composition of gargles, and is also said to abate the excessive sweats in hectic fever. The dose, in powder, is from twenty to thirty grains.

##### INFUSION OF SAGE.

R. Sage half an ounce  
Boiling water two pints  
Infuse for half an hour, and strain. Sugar



and lemon juice may be added, according to circumstances. As a drink in fevers.  
*Ellis.*

## COMPOUND INFUSION OF SAGE.

R. Sage  
Boneset, each, half an ounce  
Cascarilla one drachm  
Water one pint and a-half  
Infuse till cold, and strain. Dose, a wineglassful every three or four hours.  
In hectic fever. *Ellis.*

## GARGLE OF SAGE.

R. Infusion of sage one pint  
Diluted sulphuric acid two drachms  
Honey of roses one ounce  
Mix. In relaxation of the uvula, &c.  
*Radiis.*

R. Infusion of sage two pints  
Tincture of Peruvian bark  
Syrup of mulberries, each, half a fl. ounce  
Spirit of horseradish one drachm  
Mix. More active than the last. *Cadet.*

## VINEGAR OF SAGE.

R. Sage one part  
Vinegar six parts  
Mix, and distil four parts. *Beral.*  
As a gargle, mixed with water.

SAMBUCUS.  
ELDER.

The species recognized by the U. S. Ph., is the *Sambucus Canadensis*, which appears to be perfectly identical in properties with the European kind, the *S. nigra*.

*Sex. Syst.* Pentand. trigyn. *Nat. Syst.* Caprifoliaceæ.

*Willd. Sp. Pl.* 1494.

The officinal portion is the flowers, but the berries, bark, and leaves are also used. The flowers have a faint but peculiar odor when dried, and a bitterish taste. They are slightly stimulant and diaphoretic. The berries are diaphoretic and laxative; the bark acts as a hydragogue purgative, as do also the leaves.

## ELDER FLOWER WATER.

R. Fresh elder flowers ten pounds  
Proof spirit seven fl. ounces  
Water two gallons  
Mix, and distil a gallon. *Lond. Ph.*

Principally used as a flavoring ingredient in mixtures and emulsions.

## VINEGAR OF ELDER FLOWERS.

R. Elder flowers one part  
Vinegar of red wine twelve parts  
Macerate for eight days, express, and filter. As a gargle. *Soubiran.*

## GARGLE OF ELDER FLOWERS.

R. Elder flowers one ounce  
Boiling water one pint  
Infuse, strain, and add  
Nitrate of potassa half an ounce  
Tincture of burnet three fl. drachms  
Oxymel two fl. ounces  
Mix. *Saunders.*

## EXTRACT OF ELDER BERRIES.

R. Elder berries one pound  
Bruise them in a mortar; express the juice, and evaporate to due consistence.  
*Dub. Ph.*

Has been praised in rheumatic, gouty, and eruptive affections; in doses of one to four drachms.

## MIXTURE OF EXTRACT OF ELDER BERRIES.

R. Extract of elder berries  
Pulp of prunes  
Syrup of red poppies, each, two ounces  
Nitrate of potassa one drachm  
Mix. Two or three spoonfuls a-day, in asthma. *St. Marie.*

## DECOCTION OF ELDER BARK.

R. Elder bark three handfuls  
Water two pints  
Boil to one-half. Half a pint, morning and evening, in dropsy. *Sydenham.*

## OINTMENT OF ELDER LEAVES.

R. Fresh elder leaves three pounds  
Lard four pounds  
Suet two pounds  
Boil the leaves in the lard till crisp; express; strain; add the suet, and melt.  
*Dub. Ph.*

As a cooling application.

## OINTMENT OF ELDER FLOWERS.

R. Elder flowers  
Lard equal parts  
Melt, and continue heat, till all moisture is driven off, and express. *Beral.*  
Used as the last.

## SANGUINARIA.

## BLOODROOT.

This is the rhizome of the *Sanguinaria Canadensis*, popularly known as bloodroot or puccoon. It is a small, herbaceous perennial, flowering at the very commencement of the spring, at which time the leaf is small, but attains a large size during the summer.

*Sex. Syst.* Polyand. monog. *Nat. Syst.* Papaveraceæ.

*Linn. Sp. Pl.* 723. *Griffith, Med. Bot.* 127.

The root is horizontal, abrupt, often contorted, of a reddish brown color, externally, and of a



bright orange red within. When dried it is somewhat flattened, much wrinkled and twisted, of a reddish brown color externally, and of a bright orange within, becoming brown by exposure. The powder is brownish red. It has a faint narcotic odor, and a bitterish, acrid taste, which is very persistent. It is an acrid emetic, with narcotic and stimulant properties. Dose, as an emetic, from ten to twenty grains; for other purposes, from one to five grains.

#### PILLS OF BLOODROOT.

R. Powdered bloodroot            one drachm  
 Conserve of roses                sufficient  
 Mix, and make thirty pills. One to two as alterative, &c., five to ten as an emetic.

#### INFUSION OF BLOODROOT.

R. Bloodroot                        half an ounce  
 Boiling water                      one pint  
 Infuse for two hours. Dose, half to one fl. ounce. *Beasley.*

#### TINCTURE OF BLOODROOT.

R. Bloodroot, bruised,            four ounces  
 Diluted alcohol                   two pints  
 Macerate for fourteen days, express, and filter. *U. S. Ph.*

As emetic, three or four fl. drachms; as expectorant, alterative, &c., thirty to sixty drops.

### SAPO. SOAP.

Soaps are all combinations of animal or vegetable oils, with one or more of the alkalies; they are principally used for washing, but some of them are official.

### SAPO DURUS. HARD SOAP.

This is official under the name of *Sapo*, in the U. S. and Lond. Phs., and is a combination of olive oil and soda.

#### SPANISH, OR CASTILE SOAP.

This is the official soap of the U. S. Ph., and presents two varieties, the white and the marbled, the first of which only should be used.

### SAPO MOLLIS. SOFT SOAP.

This soap is a combination of olive or other vegetable oils and potassa, or animal oils with the same alkali.

### SAPO VULGARIS, COMMON SOAP,

Is a hard soap, made from tallow and caustic soda.

Besides these official soaps, there are many others, which are used medicinally, or for cleansing purposes.

#### ALMOND SOAP.

R. Caustic solution  
       of soda                        one thousand parts  
 Oil of almonds                   two thousand one hundred parts  
 Mix, and keep the mixture for some days, at a temperature of 68° F., stirring from time to time, until it acquires the consistence of a soft paste; place in moulds until dry. It should not be used for medicinal purposes, until it has been exposed to the action of the air for a month or two.

*Par. Cod.*

#### BEEF MARROW SOAP.

R. Purified beef  
       marrow                        five hundred parts  
 Caustic solution  
       of soda                      two hundred and fifty parts  
 Common salt                      one hundred parts  
 Water                              one thousand parts  
 Put the marrow, with hot water, into a porcelain vessel, and heat, until it is melted; then add the solution of caustic soda, by degrees, constantly stirring, till saponification is complete; then add the salt, slightly stirring, collect the soap which rises, drain it, melt it by a gentle heat, pour into moulds, and allow it to solidify.

*Par. Cod.*

#### SOAP OF TURPENTINE. STARKEY'S SOAP.

R. Dry carbonate of potassa  
       Oil of turpentine  
       Venice turpentine            equal parts  
 Triturate the potassa with the oil, and then with the turpentine, when the mass has attained a proper consistence. Preserve in an earthenware vessel.

*Ratier.*

Was supposed to be a corrector of the injurious effects of opium, hellebore, &c. It is now principally given in gonorrhoea and dropsy. Dose, eight to ten grains.

#### AROMATIC SOAP.

R. Concentrated solution of  
       soap                            six ounces  
 Oil of bergamot  
       lavender  
       rosemary, each,            two drachms  
 Mix. Employed in baths, as tonic and antispasmodic.

*Niemann.*

#### CAMPHORATED SOAP.

R. White soap                    sixteen ounces  
 Boiling water                    eight ounces  
 Olive oil                           six ounces  
 Powdered camphor              one drachm  
 Dissolve the soap in the water, evaporate



gently to the consistence of a soft paste, add the camphor incorporated in the oil, mix well, and pour into moulds.

Said to be useful in chaps and excoriations. *Wetzler.*

#### COSMETIC SOAP POWDER.

R. White soap twelve ounces  
Carbonate of potassa two ounces  
Powdered orris root three ounces  
horse-chestnuts two pounds  
Oil of lavender  
bergamot, each, forty drops  
lemon thirty drops  
cloves ten drops  
Sugar half an ounce  
Mix, and form a powder. *Niemann.*

#### ARSENICAL SOAP.

R. Arsenious acid two pounds  
Carbonate of potassa twelve ounces  
Camphor five ounces  
White soap two pounds  
Powdered lime eight ounces  
Reduce each to powder, and mix. Used as a preservative for specimens of natural history against the attacks of insects. Known as Beconi's arsenical soap. *Gannal.*

#### SAPONINE. (For cleaning gloves.)

R. Powdered soap two hundred and fifty parts  
Solution of chloride of potassa one hundred and sixty-five parts  
Solution of ammonia ten parts  
Water one hundred and fifty-five parts  
Mix, and form a paste: a small portion rubbed on a glove with a piece of flannel will cleanse it. *Duvignan.*

#### GREASE BALLS.

R. Fuller's earth two pounds four ounces  
Carbonate of soda  
Soap, each, eight ounces  
Add, Yolks of eggs eight  
Well beaten with Ox gall eight ounces  
Levigate thoroughly, form into cakes or balls, and dry. *Lenormand.*

#### BOLUS OF SOAP.

R. White soap two scruples  
Oil of caraway two or three drops  
Syrup sufficient  
Mix, and make two boluses. They are purgative. *Swedjaur.*

#### PILLS OF SOAP.

R. White soap one hundred and twenty-five parts

Marsh mallow root, powdered, sixteen parts  
Nitrate of potassa four parts  
Beat together till well incorporated, and divide into four-grain pills. *Par. Cod.*

#### PILLS OF SOAP AND OX GALL.

R. Powdered white soap two drachms  
Extract of ox gall one drachm  
Mix, and incorporate Powdered guaiacum  
Calomel, each, half a drachm  
Powdered guaiacum wood sufficient  
Mix, and make four-grain pills. In gout, one to two, morning and evening. *Vicq. D'Azir.*

#### COMPOUND SOAP PILLS.

R. White soap three drachms  
Ammoniac  
Rhubarb, each, one drachm  
Aloes ten grains  
Assafetida  
Saffron, each, thirty-six grains  
Mix, and make three-grain pills. Purgative and alterative; four to six a-day. *Recamier.*

#### ELECTUARY OF SOAP, &c.

R. White soap one drachm and a-half  
Venice turpentine one drachm  
Seneka  
Marsh mallow, each, two drachms  
Rob of juniper sufficient  
Mix. Four teaspoonfuls a-day. *Vogt.*

#### ESSENCE OF SOAP.

R. White soap twenty-four parts  
Distilled water thirty-two parts  
Alcohol (.923) sixty-four parts  
Carbonate of potassa one part  
Essence of lemons sufficient  
Dissolve the soap in the water and alcohol; add the potassa and essence, and filter. Used for the toilette. *Soubeyran.*

#### CAMPHORATED ESSENCE OF SOAP.

R. White soap three parts  
Camphor one part  
Spirit of rosemary sixteen parts  
Dissolve the camphor, and then the soap, in the spirit. As an embrocation in rheumatic pains, &c. *Guibourt.*

#### ETHEREAL SOLUTION OF SOAP.

R. Beef marrow soap one part  
Acetic ether five parts  
Dissolve by aid of a water-bath. Used as an embrocation in rheumatic pains, &c. *Foy.*

#### CAMPHORATED ACETIC BALSAM OF SOAP.

R. Common soap  
Camphor, each, one drachm



Acetic ether                      one ounce  
Oil of thyme                      ten drops  
Dissolve the soap in the ether with the aid of heat; add the camphor, then the oil, and filter. Used as above.

*Cottureau.*

#### CATAPLASM OF SOAP.

R. Common soap                      four drachms  
Roasted onion  
Mustard, each,                      two ounces  
Water                      sufficient

Heat together, and mix into a cataplasm. As a maturing application to boils, abscesses, &c. *Foy.*

#### SOAP SUPPOSITORY.

R. Soap                      two ounces  
Common salt                      one ounce  
Honey                      sufficient

Mix, and form into conical suppositories, and oil them on the surface. *Spielmann.*

#### CLYSTER OF SOAP.

R. Barley water                      six ounces  
White soap                      one drachm  
Honey                      six drachms  
Mix, and dissolve. *Brera.*

#### SOAP OF COD LIVER OIL.

R. Cod liver oil                      two ounces  
Caustic soda                      two drachms  
Water                      five drachms

Dissolve the soda in the water, and mix the solution with the oil. *Deschamps.*

#### IODURETTED SOAP OF COD LIVER OIL.

R. Soap of cod liver oil                      one ounce  
Iodide of potassium                      one drachm  
Water                      one drachm

Dissolve the iodide in the water, and add it to the soap. *Deschamps.*

#### SOAP, WITH SULPHUR.

R. Soap  
Sulphur, each,                      four ounces  
Oil of bergamot                      half a drachm  
Water                      sufficient

Beat together. As an application in itch. *Frank.*

#### ITCH OINTMENT.

R. Brown soap                      one ounce  
Common salt  
Sulphur, each,                      half an ounce  
Alcohol                      one fl. drachm  
Vinegar                      two fl. drachms  
Chloride of lime                      half a drachm

Rub well together. One-fourth to be used night and morning, as a friction. It is effectual, cheap, and inoffensive.

*Emory.*

## SARSAPARILLA.

### SARSAPARILLA.

This is the official name for the roots of several species of *Smilax* found in Mexico and various parts of South America. These roots are known in commerce by names of the places from whence they are shipped, as Honduras, Jamaica, Vera Cruz, &c.

Sarsaparilla, as found in commerce, is in packages, composed of dried roots, several feet in length, about the thickness of a quill, more or less wrinkled, of an ash gray to a dark brown color externally, and white to brownish within. The odor is slight but peculiar, the taste is at first mucilaginous, but finally somewhat acrid. It is considered alterative and tonic, but its real action is not well understood. The dose, in powder, is half a drachm to a drachm, three or four times a-day.

### POWDER OF SARSAPARILLA AND PERUVIAN BARK.

R. Powdered sarsaparilla                      one ounce  
Peruvian bark three drachms  
Carbonate of soda                      two drachms  
Mix, and divide into sixteen powders. One, thrice a-day, as an alterative. *Cline.*

### INFUSION OF SARSAPARILLA.

R. Sarsaparilla, bruised,                      one ounce  
Boiling water                      one pint  
Digest for two hours, in a covered vessel, and strain. *U. S. Ph.*

One to four fl. ounces, three times a-day.

### ALKALINE INFUSION OF SARSAPARILLA.

R. Sarsaparilla, bruised,                      twelve ounces  
Liquorice root,                      one ounce and a-half  
bruised,  
Solution of                      one fl. ounce and a-half  
potassa                      five pints and a-half  
Boiling water  
Macerate for twenty-four hours, and strain. Dose, from eight fl. ounces to one pint daily. As alterative. *St. Geo. Hosp.*

### DECOCTION OF SARSAPARILLA.

R. Sliced sarsaparilla                      five ounces  
Boiling distilled water                      four pints  
Macerate for four hours in a covered vessel, remove sarsaparilla, bruise it, return it to liquor, and macerate for two hours, and boil down to two pints, then strain. *Lond. Ph.*

Dose, four to six fl. ounces a-day.

### COMPOUND DECOCTION OF SARSAPARILLA.

R. Sarsaparilla, sliced,                      six ounces  
Bark of sassafras, sliced,  
Guaiacum wood, rasped  
Liquorice root, bruised,  
each,                      one ounce  
Mezereon, sliced,                      three drachms  
Water                      four pints



Boil for a quarter of an hour, and strain.

*U. S. Ph.*

As an alterative and diaphoretic, in secondary syphilis, cutaneous affections, &c. Dose, four to six fl. ounces, a-day.

**FELTZ'S DECOCTION OF SARSAPARILLA.**

R. Sarsaparilla, bruised      three ounces  
Isinglass      half an ounce  
Crude antimony (tied  
in a rag)      three drachms  
Water      five pints

Boil to two pints and a-half, and strain.

*Beasley.*

**VINACHE'S DECOCTION OF SARSAPARILLA.**

R. Sarsaparilla, bruised,  
China root, bruised,  
Guaiacum wood, rasped,  
each,      one ounce and a-half

Crude antimony (in a rag) two ounces

Water      six pints

Macerate for twelve hours, boil to three pints, and add

Sassafras bark, sliced,

Senna, each,      half an ounce

Infuse for an hour, strain, let settle, and decant.

*Foy.*

**LISBON DIET DRINK.**

R. Guaiacum wood, rasped,      one ounce  
Sarsaparilla, bruised,      three ounces  
Mezereon, sliced,      half an ounce  
Crude antimony (in a rag) two ounces  
Water      twelve pints

Boil down to eight pints, and add

Red saunders, rasped,  
White sandal, rasped,  
each,      three ounces

Rose wood, rasped,

Sassafras bark, sliced, each, one ounce

Liquorice root, sliced,      half an ounce

Infuse for four hours, strain, and add syrup according to taste.

*Foy.*

Dose, a pint to two pints a-day.

R. Sarsaparilla, bruised,      four ounces  
Dried walnut peel      four ounces  
Guaiacum, rasped, one ounce and a-half  
Crude antimony (in a rag) half an ounce  
Water      four pints

Boil down to three pints.

*Pearson.*

**ZITTMAN'S DECOCTION.**

R. Sarsaparilla, bruised,      twelve ounces  
Spring water      fifty-six pints

Macerate for twenty-four hours, and introduce, tied up in a rag.

Styptic powder (four parts alum

and one of kino) one ounce and a-half

Calomel      half an ounce

Cinnabar      one drachm

Boil down to eighteen pints, and add

Aniseed

Fennel seed, each,      half an ounce

Senna      three ounces

Liquorice root      one ounce and a-half

Strain, and put aside as *strong decoction*.

Add to the residue

Sarsaparilla, bruised,      six ounces

Spring water      fifty-six pints

Boil down to eighteen pints, and add

Lemon peel

Cinnamon

Cardamoms

Liquorice root, each,      three drachms

Strain, and label, *Weak decoction*. To be

given freely in the treatment of syphilitic and cutaneous affections, in chronic rheumatism, &c.

*Radius.*

**JAUPERAND'S DECOCTION.**

R. Bruised sarsaparilla

China root

Crude antimony, each,      two ounces

Carbonate of potassa      one drachm

Senna

Sassafras, sliced, each,      one ounce

Rhubarb

Peruvian bark, each,      half an ounce

Water      sufficient

Boil by a gentle fire for eight hours, till twelve pints of decoction are obtained.

Let stand for some time, and strain. Advised in scrofula, two fl. ounces, three times a-day.

*Bories.*

**EXTRACT OF SARSAPARILLA.**

R. Sarsaparilla, in coarse

powder,      one pound

Diluted alcohol      four pints

Proceed as directed for alcoholic extract of aconite. *U. S. Ph. and Par. Cod.*

Dose, from ten to twenty grains, three or four times a-day.

**FLUID EXTRACT OF SARSAPARILLA.**

R. Sarsaparilla, bruised,      sixteen ounces

Liquorice root, bruised,

Guaiacum wood, rasped,

Sassafras bark, sliced, each, two ounces

Mezereon, sliced,      six drachms

Diluted alcohol,      eight pints

Digest for fourteen days, strain, express, and filter. Evaporate in water-bath to

twelve fl. ounces, add eight ounces of sugar, and remove from fire when this is dissolved. Dose, one fl. drachm, three or

four times a-day. *W. Hodgson.*

**ESSENCE OF SARSAPARILLA.**

R. Sarsaparilla, bruised,      ten ounces



Distilled water six pints  
Macerate at a temperature of 120° for six hours, and strain; repeat with same quantity of water. Mix the liquors, evaporate in china vessels, at 160°.

If reduced to ten fl. ounces (or to nine fl. ounces, with one fl. ounce of spirit added), one fl. drachm mixed with seven fl. drachms, is equal to the decoction of usual strength. If reduced to five fl. ounces, one fl. drachm is equal to two fl. ounces of decoction.

*Hening.*

R. Alcoholic extract of sarsaparilla one pound  
Sherry wine three pints  
Dissolve and filter. Half a fl. ounce to one fl. ounce, mixed with a quart of water, is equal to the decoction.

*Beral.*

#### COMPOUND ESSENCE OF SARSAPARILLA.

R. Sarsaparilla, bruised, eight ounces  
Boiling water sufficient  
to exhaust the root, by successive macerations; unite the liquors, and evaporate to ten fl. ounces; strain, and add, when cool, Alcohol (.842)

Tincture of guaiacum, each, four fl. drachms  
White wine one fl. ounce  
Essence of sassafras twelve drops  
Extract of liquorice two drachms

Filter. Dose, a spoonful, morning and evening, in some convenient vehicle.

*Cadet.*

#### SYRUP OF SARSAPARILLA.

R. Sarsaparilla, sliced, fifteen ounces  
Boiling water one gallon  
Macerate for twenty-four hours, boil down to four pints, strain, and add

Sugar fifteen ounces  
And evaporate to proper consistence.

*Lond. Ph.*

R. Alcoholic extract of sarsaparilla six ounces  
Water four pints

Dissolve by means of a water-bath, strain, and add

Sugar eight pounds  
Dissolve whilst hot.

*Beral.*

Each fl. ounce is equal to fifteen grains of the extract.

#### COMPOUND SYRUP OF SARSAPARILLA.

R. Sarsaparilla, bruised, two pounds  
Guaiacum wood, rasped, three ounces  
Hundred-leaved roses

Senna  
Liquorice root, bruised, each, two ounces

Oil of sassafras  
Oil of anise, each, five minims  
Oil of partridgeberry three minims  
Diluted alcohol ten pints  
Sugar eight pounds

Macerate the first five ingredients in the alcohol for fourteen days, express, and filter; evaporate the tincture to four pints, filter, add the sugar, and form a syrup; then, having rubbed the oils with a little of the syrup, mix well with the remainder.

*U. S. Ph.*

R. Sarsaparilla two pounds  
Bittersweet  
Pipsissewa, each, half a pound  
Guaiacum  
Liquorice root, each, four ounces  
Sassafras  
Partridgeberry leaves, each, two ounces  
Sugar twelve pounds

Reduce the ingredients to a coarse powder, macerate them in diluted alcohol, 18° B. for two days, put in a displacement apparatus, and displace slowly until two gallons of liquid have passed. Evaporate on a water-bath till reduced to six pints, then add the sugar, and form a syrup.

*Staples.*

Dose, half a fl. ounce, three or four times a-day.

#### SYRUP OF LAFFECTEUR.

R. Sarsaparilla, bruised,  
Marsh reed grass, each, thirty ounces  
Borage flowers eight ounces  
Senna  
White roses, each, two ounces  
Sugar  
Honey, each, six pounds  
Water eighteen pints

Boil the first two ingredients in half the water for an hour, strain off the decoction, and repeat the process with the remainder of the water, and boil two hours; towards close, add the senna, rose leaves, and borage, strain, mix the decoctions, add the sugar and honey, and boil to the consistence of syrup. Dose, six tablespoonfuls, early in the morning.

*Ellis.*

R. Sarsaparilla, bruised, nine pounds  
Guaiacum wood, rasped,  
China root, bruised, each, six pounds  
Yellow Peruvian bark, bruised, three pounds

Macerate for twenty-four hours in

Water one hundred and forty pints  
Boil down to one-third, express, and strain. Repeat twice, with same proportions of water; strain, unite the three decoctions, add

Purified molasses thirty pounds



Evaporate to the consistence of syrup, and pour, boiling, on  
 Sassafras, rasped, five pounds  
 Aniseed four ounces  
 Borage flowers one pound and a-half  
 Finally, strain. *Cadet.*

One to three ounces a-day, in syphilitic and cutaneous disorders.

#### SYRUP OF GESNOUIN.

R. Syrup of sarsaparilla four pounds  
 senna three ounces  
 Extract of borage five ounces  
 Conserve of elder berries one ounce  
 Mix, with heat. Four to six spoonfuls, early in the morning, using the infusion of sarsaparilla for drink. *Foy.*

#### SARSAPARILLA BEER.

R. Sarsaparilla, bruised, two pounds  
 Bark of guaiacum, powdered, eight ounces  
 Guaiacum wood, rasped, Anise seeds  
 Liquorice root, each, four ounces  
 Mezereon, bark of root, two ounces  
 Molasses two pounds  
 Cloves, bruised, twelve  
 Boiling water four gallons  
 Shake, thrice a-day. Let ferment; when fully fermented, to be taken in doses of a small tumblerful, two or three times a-day. *Hancock.*

### SASSAFRAS.

#### SASSAFRAS.

Two parts of the Sassafras, *Laurus sassafras* (*Sassafras officinale*), are officinal; the pith of the young branches and the bark of the root.

Sassafras is a medium-sized tree, found in most parts of the United States, growing in woods and open places, flowering before the appearance of the leaves.

*Sex. Syst.* Enneand. monog. *Nat. Syst.* Lauracæ.

Nees. Pl. Med. i. 131. Griffith, Med. Bot. 551.

The pith is in slender, cylindrical pieces, very spongy and light, with a mucilaginous and somewhat aromatic taste, affording a clear mucilage on the addition of water. The bark is in small fragments, of a reddish brown color, brittle, of a fragrant odor, and a pleasant aromatic taste. It is stimulant and diaphoretic.

#### INFUSION OF SASSAFRAS BARK.

R. Sassafras bark half an ounce  
 Boiling water one pint  
 Macerate for six hours, and strain. As a stimulating diaphoretic. *Niemann.*

#### COMPOUND INFUSION OF SASSAFRAS BARK.

R. Sassafras bark half an ounce

Liquorice root two drachms  
 Boiling water one pint  
 Infuse. One-half, morning and evening.  
 In scrofulous complaints. *Hufeland.*

#### MUCILAGE OF SASSAFRAS PITH.

R. Sassafras pith one drachm  
 Boiling water one pint  
 Infuse, and strain. As a soothing application in ophthalmia, and as a demulcent drink in diseases of the bowels, lungs, and bladder. *Wood.*

#### TINCTURE OF SASSAFRAS.

R. Sassafras, sliced, one part  
 Alcohol six parts  
 Macerate for three days, express, and filter.  
 Dose, one fl. drachm. *Beral.*

#### COMPOUND TINCTURE OF SASSAFRAS.

R. Sassafras, bruised, four ounces  
 Sage two ounces  
 Nutmeg, bruised, one ounce  
 Cinnamon, bruised, half an ounce  
 Star anise three drachms  
 Alcohol two pints  
 Macerate for three days, express, and filter. *Wurt. Ph.*

As a stomachic and carminative.

#### OIL OF SASSAFRAS.

R. Bruised sassafras at will  
 Water sufficient  
 Distil, and collect the oil which sinks to the bottom of the product. *Guibourt.*  
 Dose, two to ten drops, as carminative and stimulant.

### SCAMMONIUM.

#### SCAMMONY.

This is the concrete juice of the root of *Convolvulus scammonia*, a twining perennial plant, a native of Syria. It has a large, succulent root, affording a milky juice, which concretes by exposure to the air.

*Sex. Syst.* Pentand. monog. *Nat. Syst.* Convolvulacæ.

Linn. Sp. Pl. 218. Griffith, Med. Bot. 477.

The best scammony is light, resinous when broken, friable, not effervescing with an acid, of various shades of color, from dark ash to dark olive; forming an emulsion with water. The odor is peculiar and unpleasant, the taste is bitterish and acrid. It is an active purgative, causing pain, and usually operating with harshness when given alone, but much milder in combination. The dose is from five to fifteen grains.

#### POWDER OF SCAMMONY AND CREMOR TARTAR.

R. Powdered scammony  
 Cremor tartar equal parts  
 Mix. *Ed. Ph.*  
 Dose, fifteen to thirty grains.



## POWDER OF SCAMMONY AND SOOT.

R. Powdered scammony one drachm  
Soot one drachm and a-half  
Powdered resin two drachms  
Mix. Dose, twenty to thirty grains.

*Beasley.*

## COMPOUND POWDER OF SCAMMONY.

R. Scammony  
Hard extract of jalap, each, two ounces  
Ginger half an ounce  
Rub separately into fine powder, and mix.  
Dose, ten to twenty grains. *Lond. Ph.*

## PILLS OF SCAMMONY.

R. Powdered scammony fifteen grains  
Sugar ten grains  
Rub together, and add  
Oil of caraway four minims

## PILLS OF SCAMMONY AND OX GALL.

R. Powdered scammony two drachms  
Ox gall three drachms  
Extract of gentian half an ounce  
Mix, and make one hundred and sixty-two pills. Dose, four to six a-day. *Cadet.*

## COMPOUND PILLS OF SCAMMONY.

R. Powdered scammony  
Extract of henbane  
Powdered gamboge  
Compound extract of colocynth  
Soap, each, twelve grains  
Water sufficient  
Beat into mass, and form twelve pills.  
Dose, two to three. *Guy's Hosp.*

## CONFECTION OF SCAMMONY.

R. Powdered scammony one ounce and a-half  
Cloves, bruised,  
Powdered ginger, each, six drachms  
Oil of caraway half fl. drachm  
Syrup of roses sufficient  
Rub the dry ingredients into fine powder, add syrup, and rub well; then the oil of caraway, and mix. Dose, twenty to forty grains. *Dub. Ph.*

R. Powdered scammony  
jalap, each, two drachms  
squill one drachm and a-half  
resin of jalap half a drachm  
Syrup of buckthorn sufficient  
Mix, and form confection. A hydragogue, in dose of ten grains to a scruple. *Foy.*

## EXTRACT OR RESIN OF SCAMMONY.

R. Scammony, in fine powder, at will  
Boil in successive portions of proof spirit, till the spirit ceases to dissolve anything, filter, distil off the spirit, pour off the watery solution from the resin, wash this

well with boiling water, dry in a heat not above 240°. *Ed. Ph.*

Dose, five to ten grains.

## EMULSION OF SCAMMONY.

R. Resin of scammony seven grains  
Milk three fl. ounces  
Triturate the resin with a little of the milk, and gradually add the rest. *Ed. Ph.*  
The whole at a dose.

## COMPOUND EMULSION OF SCAMMONY.

R. Scammony seven grains  
Sugar two drachms  
Triturate, and add gradually  
New milk three fl. ounces  
Cherry laurel water three or four drops for a dose. *Blanche.*

## SCAMMONY MIXTURE.

R. Resin of scammony twelve grains  
Spirit of rosemary one fl. drachm  
Orange-flower water  
Syrup of peach flowers, each, one fl. ounce  
Mix. *Cadet.*

## SCAMMONY OIL.

R. Powdered scammony one scruple  
Oil of almonds one ounce  
Rub well together. Dose, half an ounce. *Van Mons.*

## ELIXIR OF SCAMMONY.

R. Scammony two drachms  
Proof spirit eight fl. ounces  
Heat, and then set fire to the spirit, and add  
Sugar four ounces  
When it is dissolved, extinguish the flame, and add  
Syrup of violets two fl. ounces  
Filter. It will give twelve ounces, containing twelve grains of scammony to the ounce. *Guibourt.*

## SOAP OF SCAMMONY.

R. Scammony one drachm  
Soap two drachms  
Alcohol (.874) two fl. ounces  
Dissolve by means of a water-bath. *Guibourt.*

## PURGATIVE PASTILLES.

R. Scammony six grains  
Tincture of senna forty drops  
Carbonate of magnesia five scruples  
White sugar eight scruples  
Powdered liquorice eight grains  
Gum tragacanth five grains  
Oil of aniseed one drop  
Syrup of violets sufficient  
to make eight pastilles. These are known



as *Pastilles de Santé*, and are useful in cases of acid eructation and flatulence. One or two, taken early in the morning, act as a laxative; by repeating the dose every quarter of an hour, full purgation ensues. *Delvincourt.*

#### PURGATIVE CHOCOLATE.

R. Chocolate five drachms  
Scammony resin six grains  
Calomel (hydrosulphate) one and a-half grains  
Sugar half a drachm

Triturate together in a mortar the scammony and sugar, then add the calomel, and, when thoroughly mixed, add the chocolate, previously softened by the heat of a water-bath. Make into forms. To be taken dry, and followed by some bland liquid. *Mialhe.*

#### SCILLA, SQUILL,

Is the bulb of *Scilla maritima*, a perennial plant, growing on the seacoast of the countries bordering on the Mediterranean. The bulb is large, pyriform, and tunicated; there are two varieties, the red and the white, but identical in their medical qualities.

*Sex. Syst.* Hexand. monog. *Nat. Syst.* Liliaceæ.

Linn. Sp. Pl. 442. Griffith, Med. Bot. 652.

The bulb is sliced and dried for use, in this state: it is in white, or yellowish-white, semi-transparent pieces, which are flexible when moist, but brittle when dry; of a mucilaginous, bitter taste, but scarcely any odor.

Squill is expectorant, diuretic, and in large doses emetic, and even purgative. Dose, as diuretic or expectorant, one grain two or three times a-day; as emetic, six to twelve grains.

#### POWDER OF SQUILL AND NITRE.

R. Powdered squill six grains  
nitre one drachm  
Mix, and divide into six powders. One, three times a-day, in sugar and water. As a diuretic. *Ellis.*

#### POWDER OF SQUILL AND IPECACUANHA.

R. Powdered squill twelve grains  
ipecacuanha twenty-four grains  
Mix, and divide into six powders. One, every two hours in chronic catarrh. *Foy.*

#### POWDER OF SQUILL AND CREMOR TARTAR.

R. Powdered squill one drachm  
Cremor tartar nine drachms  
Mix. Twenty to thirty grains, two or three times a-day, as a diuretic. *Guy's Hospt.*

#### POWDER OF SQUILL AND SULPHUR.

R. Powdered squill one part  
Flowers of sulphur two parts  
Powdered sugar three parts  
Mix. Dose, from five to twenty grains, according to age, in catarrhs. *Guibourt.*

#### POWDER OF SQUILL AND CALOMEL.

R. Powdered squill three grains  
Calomel  
Powdered gamboge, each, half a grain  
Mix. To be taken in the evening. As a diuretic. *A. Cooper.*

#### COMPOUND POWDER OF SQUILL.

R. Powdered squill  
foxglove, each, one grain  
Oil of juniper two drops  
Borate of soda  
Powdered liquorice, each, one scruple  
Powdered cinnamon two grains  
Mix. To be repeated, two or three times a-day, as a diuretic. *Radius.*

R. Powdered squill  
foxglove, each, five grains  
Resin of jalap six grains  
Acetate of potassa thirty grains  
Sulphate of soda two drachms  
Mix. To be taken early in the morning, as a purgative and diuretic. *Bories.*

#### PILLS OF SQUILL AND CALOMEL.

R. Calomel  
Powdered squill, each, twelve grains  
Conserve of roses sufficient  
Mix, and form twelve pills. One, night and morning. In dropsy, arising from visceral derangement. *Ellis.*

#### PILLS OF SQUILL AND AMMONIAC.

R. Powdered squill thirty grains  
Ammoniac one drachm and a-half  
Extract of hemlock thirty grains  
Mix, and form thirty pills; two, twice a-day. In asthma and chronic catarrh. *Ellis.*

#### PILLS OF SQUILL AND CROTON OIL.

R. Compound squill pill two scruples  
extract of colocynth two scruples  
Croton oil six minims  
Mix, and form eighteen pills. Three, twice a-week, in dropsy. *Schwyn.*

#### PILLS OF SQUILL AND IPECACUANHA.

R. Powdered squill half an ounce  
ipecacuanha two drachms  
Extract of opium fifteen grains  
Butter of cocoa one ounce  
Syrup of gum sufficient  
Beat together, and make pills of four



grains. One, three times a-day, as an expectorant. *Cadet.*

R. Powdered squill twelve grains  
       ipecacuanha twelve grains  
 Extract of dandelion three scruples  
 Blue pill ten grains  
 Simple syrup sufficient  
 Mix, and make twenty-four pills. Two, to be taken morning and evening, in dropsy. *Ainslie.*

#### PILLS OF SQUILL AND HENBANE.

R. Powdered squill half a drachm  
 Extract of henbane two scruples  
 Myrrh one drachm and a-half  
 Water sufficient  
 Beat together, and make thirty pills. Two, morning and evening, as expectorant. *Paris.*

#### COMPOUND PILLS OF SQUILL.

R. Calomel three grains  
 Recent squill  
 Ammoniac, each, one scruple  
 Dover's powder half a drachm  
 Conserve of roses sufficient  
 Make mass, and divide into thirty pills. One, three or four times a-day. In troublesome catarrh. *Latham.*

R. Powdered squill half a drachm  
 Calomel twenty-four grains  
 Tartar emetic six grains  
 Powdered opium eighteen grains  
 Conserve of roses sufficient  
 Beat together, and make twenty-four pills. One at night, in catarrh. *Ellis.*

R. Powdered squill one drachm  
       ginger  
       ammoniac, each, two drachms  
 Soap three drachms  
 Syrup sufficient  
 Beat into mass, and form one hundred and twenty pills. *U. S. Ph.*

Five to ten grains, three or four times a-day, as a stimulant expectorant.

R. Powdered squill fifteen grains  
       ammoniac one drachm and a-half  
       seneka two drachms  
 Golden sulph. of anti-  
       mony half a drachm  
 Extract of dandelion sufficient  
 Mix, and make pills of two grains. Diuretic. *Berends.*

R. Powdered squill twenty-four grains  
 Calomel sixteen grains  
 Soap one drachm and a half  
 Galbanum two drachms

Extract of dandelion three drachms and a-half  
 Essence of orange sufficient  
 Mix, and make pills of two grains. Three or four a-day, as purgative and diuretic. *Pideret.*

#### COMPOUND SYRUP OF SQUILL.

R. Bruised squill  
       seneka, each, four ounces  
 Tartar emetic forty-eight grains  
 Water four pints  
 Sugar three pounds and a-half  
 Pour the water on the squill and seneka, boil to one-half, strain, and add the sugar; evaporate to three pints, and, whilst hot, add the tartar emetic. *U. S. Ph.*

R. Squill  
       Seneka, in coarse  
       powder, each, four ounces  
 Tartar emetic forty-eight grains  
 Alcohol half a pint  
 Water sufficient  
 Sugar three pounds and a-half  
 Mix the alcohol with two pints and a-half of the water, and macerate the squill and seneka in the mixture for twenty-four hours. Put the whole into a displacement apparatus, and add as much water as may be necessary to obtain three pints of filtered liquor. Boil this for a few minutes, evaporate to one-half, and strain, then add the sugar, and evaporate to three pints; lastly, dissolve the tartar emetic in the syrup, whilst hot. *U. S. Ph.*

Known as *Hive syrup*. Dose, as expectorant, twenty to thirty drops, for adult; for children, five to ten drops; in croup, ten drops to a fl. drachm, repeated till it vomits.

R. Extract of seneka and squill one pound  
 Clarified honey, at 160°, six pounds  
 Mix, and add  
       Tartar emetic sixteen grains  
 to each sixteen fl. ounces of the syrup. *Ecky.*

R. Oxy-mel of  
       squill one fl. ounce and a-half  
 Syrup of ipecacuanha  
       poppies, each, two fl. ounces  
       orange-flowers half an ounce  
 Mix. Dose, half to one fl. ounce in whooping-cough. *Cadet.*

#### WINE OF SQUILL.

R. Squill one ounce  
 White wine sixteen fl. ounces  
 Macerate for twelve days, express, and filter. *Par. Cod.*



## COMPOUND WINE OF SQUILL.

R. Bruised squill one ounce  
 orange peel  
 calamus, each, three drachms  
 juniper berries two drachms  
 White wine four pints  
 Macerate for three days, express, filter,  
 and add  
 Oxymel of squills two fl. ounces  
 R. Dried squill, bruised,  
 Rhubarb  
 Juniper berries, each, one drachm  
 Cinnamon three drachms  
 Zedoary two drachms  
 Carbonate of  
 potassa one drachm and a-half  
 White wine two pints  
 Macerate, express, and filter. Dose, two  
 to four small wineglassfuls a-day, as a  
 diuretic. *Taddei.*

## BITTER WINE OF SQUILL.

R. Squill  
 Root of swallow-  
 wort, each, half an ounce  
 Peruvian bark  
 Winter's bark  
 Lemon peel, each, two ounces  
 Angelica half an ounce  
 Wormwood  
 Balm, each, one ounce  
 Juniper berries  
 Mace, each, half an ounce  
 White wine eight pints  
 Digest by a gentle heat, for four days, ex-  
 press, and filter. As a diuretic, in doses  
 of one to three fl. ounces. *Cottureau.*

## TINCTURE OF SQUILL.

R. Squill four ounces  
 Diluted alcohol two pints  
 Macerate for fourteen days, express, and  
 filter. *U. S. Ph.*  
 Dose, as diuretic or expectorant, twenty to  
 forty drops.

## ALKALINE TINCTURE OF SQUILL.

R. Squill two ounces  
 Solution of potassa two fl. drachms  
 Diluted alcohol twelve fl. ounces  
 Macerate for three days, express, and filter.  
*Van Mons.*

## TINCTURE OF SQUILL AND ELATERIUM.

R. Tincture of squill  
 Vinegar of colchi-  
 cum, each, two fl. drachms  
 Spirit of nitric ether one fl. ounce  
 Extract of elaterium one grain  
 Mix. Dose, fifteen minims to one fl.  
 drachm. *St. Bart. Hosp.*

COMPOUND TINCTURE OF SQUILL AND  
BENZOIN.

R. Squill  
 Orris root  
 Elecampane, each, twenty-five drachms  
 Benzoin two drachms  
 Licorice root  
 Aniseed  
 Myrrh, each, four scruples  
 Gum ammoniac two scruples  
 Saffron eighteen grains  
 Diluted alcohol one pint, six fl. ounces  
 Macerate for fifteen days, express, and fil-  
 ter. Celebrated as *Wedel's elixir*. Dose,  
 forty to sixty drops, in catarrh, asthma, &c.  
*Cadet.*

## ETHEREAL TINCTURE OF SQUILL.

R. Squill one ounce and a-half  
 Muriatic ether  
 Spirit of juniper, each, half a pound  
 Water sufficient  
 Distil one pound. Dose, ten to twenty  
 drops in water, as a diuretic. *Augustin.*

## MIXTURE OF SYRUP OF SQUILLS, &amp;c.

R. Syrup of squill half fl. ounce  
 Paregoric elixir two fl. drachms  
 Antimonial wine one fl. drachm  
 Powdered gum Arabic half a drachm  
 Distilled water three fl. ounces  
 Mix. A teaspoonful, every two or three  
 hours, as expectorant; suited to children.  
*Ellis.*

R. Syrup of squill  
 Peppermint water, each, two fl. ounces  
 Paregoric elixir  
 Compound spirit of  
 lavender, each, half fl. ounce  
 Syrup one fl. ounce  
 Mix. A spoonful, three or four times a-  
 day. A valuable expectorant. *Christison.*

MIXTURE OF OXYMEL OF SQUILL AND  
VALERIAN.

R. Powdered valerian two drachms  
 Oxymel of squill one fl. ounce  
 Laudanum twenty drops  
 Water one fl. ounce  
 Mix. A teaspoonful, every hour, after an  
 emetic in croup. *Kimbel.*

MIXTURE OF EXTRACT OF SQUILL AND  
RHUBARB.

R. Extract of squill sixteen grains  
 Liquid acetate of potassa half an ounce  
 Parsley water four fl. ounces  
 Syrup of rhubarb one fl. ounce  
 Mix. As a diuretic. *Radius.*



## MIXTURE OF OXYMEL OF SQUILL AND AMMONIAC.

R. Oxymel of squill six fl. drachms  
 Gum ammoniac one drachm  
 Extract of elecampane twenty-four grains  
 Syrup of hyssop six fl. drachms  
 Hyssop water four fl. ounces  
 Mix. A spoonful every hour, as an expectorant. *Foy.*

## MIXTURE OF VINEGAR OF SQUILL.

R. Vinegar of squill one fl. drachm  
 Barley water five fl. ounces  
 Syrup of saffron one fl. ounce  
 Mix. A tablespoonful, three or four times a-day, in troublesome cough. *Ainslie.*

## MIXTURE OF VINEGAR OF SQUILL AND AMMONIA.

R. Vinegar of squill two fl. drachms  
 Solution of acetate of ammonia  
 Compound spirit of horse-radish, each, one fl. ounce and a-half  
 Diluted pimento water four fl. ounces and a-half  
 Mix. Dose, two fl. ounces, three times a-day. As a diuretic. *St. Bart. Hosp.*

## MIXTURE OF OXYMEL OF SQUILL AND IPECACUANHA.

R. Ipecacuanha one drachm and a-half  
 Lemon peel two drachms  
 Cremor tartar half an ounce  
 Water four fl. ounces  
 Boil, strain, and add  
 Oxymel of squill half a fl. ounce  
 Mix well, as an expectorant. *Cadet.*

## MIXTURE OF OXYMEL OF SQUILL AND MARSH MALLOW.

R. Oxymel of squill  
 Syrup of marsh mallow  
 Mucilage of gum Arabic equal parts  
 Mix. As expectorant, in spoonful doses. *Foy.*

## EMULSION WITH SQUILL.

R. Oil of almonds  
 Mucilage of gum Arabic, each, one fl. ounce  
 Wine of squill eight fl. scruples  
 ipecacuanha four fl. scruples  
 Syrup of tolu two fl. ounces  
 Water of hyssop four fl. ounces  
 Water eight fl. ounces  
 Make an emulsion. *Beral.*

## OINTMENT OF SQUILL.

R. Squill three ounces  
 Solution of potassa sufficient  
 Reduce to mucilage by boiling, and add

Lard sufficient  
 to form an ointment. As friction to indolent tumors. *Hufeland.*

## COMPOUND OINTMENT OF SQUILL.

R. Fresh squill  
 Root of blue iris  
 male fern, each, six ounces  
 Juice of elaterium two pounds  
 bryony three pounds  
 Macerate for twenty-four hours, boil, express, and evaporate to soft consistence, and add  
 White wax fifteen ounces  
 Oil of mucilage four pounds  
 Mix. As a resolvent application. *Saxon Ph.*

## EXTRACT OF SQUILL.

R. Squill one pound  
 Proof spirit four pints  
 Macerate for some days, express, strain and filter; digest the residue with two pints of spirit, express, strain, and filter. Distil off the spirit, and evaporate to proper consistence. Dose, one to three grains. *Par. Cod.*

## ELECTUARY OF SQUILL.

R. Oxymel of squill two fl. ounces  
 Cremor tartar three ounces  
 Mix. Dose, two drachms. *Beasley.*  
 R. Powder of squill and nitre  
 Gum ammoniac  
 Tartrate of potassa, each, two drachms  
 Muriate of ammonia one scruple  
 Syrup of cinnamon sufficient  
 Mix. Dose, two drachms, three times a-day, in dropsy. *Radius.*

## VINEGAR OF SQUILL.

R. Bruised squill four ounces  
 Distilled vinegar two pints  
 Alcohol one fl. ounce  
 Macerate the squill with the vinegar in a close glass vessel for seven days, express, and let settle, pour off the clear liquid, and add alcohol. *U. S. Ph.*

Diuretic and expectorant. Dose, thirty minims to a fl. drachm.

## OXYMEL OF SQUILL.

R. Vinegar of squill two pints  
 Clarified honey three pounds  
 Mix, and evaporate by water-bath to the proper consistence. *U. S. Ph.*  
 Dose, one to two fl. drachms.

## SYRUP OF SQUILL.

R. Vinegar of squill one pint  
 Sugar two pounds



Mix, and form syrup. *U. S. Ph.*

As expectorant. Dose, about a fl. drachm.

#### AROMATIC SYRUP OF SQUILL.

R. Vinegar of squill half a pound

Syrup of cinnamon

ginger, each, one pound

Mix. As a stimulating expectorant.

*Swed. iaur.*

### SCOPARIUS.

#### BROOM.

This is the fresh tops of the *Cytisus scoparius*, a shrub, native of many parts of Europe, and cultivated in this country, with numerous, bright yellow flowers.

*Sex. Syst.* Diadel. decand. *Nat. Syst.* Fabaceæ.

Link. Enum. ii. 241. Griffith. Med. Bot. 233.

The officinal portion is the tops of the branches, but the seeds are also used. The branches have a bitter, nauseous taste, and a strong, peculiar odor. It is diuretic and cathartic, and even emetic, and has been advantageously used in dropsy.

#### INFUSION OF BROOM.

R. Broom tops one ounce

Boiling water one pint

Digest for four hours in a covered vessel, and strain. *Lond. Ph.*

Dose, two to four fl. ounces, three times a-day.

#### DECOCTION OF BROOM.

R. Broom tops

Juniper tops, each, half an ounce

Cremor tartar three drachms

Water one pint and a-half

Boil down to a pint, and strain. *Ed. Ph.*

#### COMPOUND DECOCTION OF BROOM.

R. Broom tops

Juniper berries

Dandelion root, each, half an ounce

Water one pint and a-half

Boil down to a pint, and strain.

*Lond. Ph.*

Dose, same as infusion.

#### EXTRACT OF BROOM.

R. Tops of broom at will

Boil, in eight times their weight in water, down to one-half; express, let settle, and strain. Evaporate by vapor-bath to proper consistence. *Dub. Ph.*

Dose, a scruple to a drachm.

#### CONSERVE OF BROOM.

R. Broom flowers one part

Sugar two parts

Rub together. Dose, a drachm to two drachms, in gout, rheumatism, &c.

*Van Mons.*

### SCROPHULARIA NODOSA.

#### FIGWORT.

A semi-aquatic plant, native of many parts of Europe, flowering in July.

*Sex. Syst.* Didyn. angios. *Nat. Syst.* Scrophulariaceæ.

Linn. Sp. Pl. 863. Griffith. Med. Bot. 518.

The leaves are the officinal part; they have a rank, unpleasant odor, and a bitter, disagreeable taste. They have been used as tonic, diuretic, and anthelmintic, but are principally used as an external discutient application.

#### OINTMENT OF FIGWORT.

R. Fresh leaves of figwort

Prepared lard, each, two drachms

suet one pound

Boil till the leaves are crisp, express, and strain. *Dub. Ph.*

As an application to piles, painful swellings, and cutaneous eruptions.

#### OIL OF FIGWORT.

R. Leaves of figwort three parts

Olive oil one part

Infuse, express, and strain. Used as above.

*Van Mons.*

### SENEGA.

#### SENEKA.

The officinal name for the root of *Polygala senega*, a native, perennial plant, found in most parts of the United States, on hill sides and dry woods, flowering from June to August.

*Sex. Syst.* Diadel. octand. *Nat. Syst.* Polygalaceæ.

Linn. Sp. Pl. 990. Griffith. Med. Bot. 225.

The root is contorted, knotted, tapering, branched, with a ridge or carina. It has a thick, resinous, grayish yellow, cortical portion, which only is active. It has a peculiar, but unpleasant odor; and a mucilaginous, sweetish taste, which becomes acrid and irritating. Its action is very various, being emetic, sudorific, diuretic, and expectorant, and even emmenagogue and purgative. It is principally used as a stimulating expectorant and diuretic. The dose, in powder, is from ten to twenty grains, but it is seldom given in this form.

#### COMPOUND POWDER OF SENEKA.

R. Powdered seneka half an ounce

Cremor tartar six drachms

Magnesia two drachms

Mix. A teaspoonful, twice a-day, in pannus of the eye. *Radius.*

#### PILLS OF SENEKA.

R. Powdered seneka six drachms

Extract of dandelion sufficient

Mix, and make pills of two or three grains. Six to twelve, three times a-day, in hypopyon and pannus. *Radius.*



## INFUSION OF SENEKA.

R. Seneka ten drachms  
Boiling water one pint  
Infuse for four hours in a covered vessel,  
and strain. *Ed. Ph.*

Dose, one to three fl. ounces.

## COMPOUND INFUSION OF SENEKA.

R. Seneka two drachms  
Boiling water six fl. ounces  
Infuse, strain, and add  
Syrup of tolu one fl. ounce  
Ammoniac half a drachm  
Mix well. A tablespoonful, every two  
hours. In catarrh. *Ellis.*

## DECOCTION OF SENEKA.

R. Seneka, bruised, one ounce  
Water one pint and a-half  
Boil down to one pint, and strain.  
*U. S. Ph.*

Dose, a tablespoonful, every three hours.

R. Seneka, bruised, one ounce  
Liquorice root half an ounce  
Distilled water one pint and a-half  
Boil down to a pint, and strain. Dose, a  
tablespoonful, in pectoral diseases. *Ellis.*

R. Bruised seneka one ounce  
Orange peel sufficient  
Water one pint  
Simmer by a slow fire, till reduced one-  
third; add orange peel; cover till cold,  
then strain. Four fl. ounces a-day, at  
first, then increase, as an emmenagogue.  
*Chapman.*

## SYRUP OF SENEKA.

R. Bruised seneka four ounces  
Water one pint  
Sugar one pound  
Boil the seneka with the water, to one-  
half; strain, add the sugar, and form  
syrup. *U. S. Ph.*

Dose, one to two fl. drachms, in pectoral com-  
plaints.

## EMULSION WITH SENEKA.

R. Bruised seneka half an ounce  
Water nine fl. ounces  
Boil down to one-third, strain, and add  
Camphor one scruple  
Gum Arabic two drachms  
Make an emulsion, and add  
Compound tincture of  
ammonia two fl. drachms  
A spoonful, every two hours, as a stimu-  
lating expectorant. *Phœbus.*

## PECTORAL JELLY WITH SENEKA.

R. Bruised seneka two drachms  
Water twelve fl. ounces

Boil down to nine fl. ounces, strain, and  
add

Isinglass sufficient  
Syrup of orange flowers one ounce  
Form a jelly. A spoonful, occasionally.  
*Radius.*

## MIXTURE WITH SENEKA.

R. Seneka, bruised, two drachms  
Water sufficient  
to obtain eight fl. ounces of strained de-  
coction; add

Camphor one drachm  
Mix. A spoonful, occasionally. *Brera.*

R. Infusion of seneka four fl. ounces  
Syrup of ipecacuanha one fl. ounce  
Oxymel of squill three fl. drachms  
Tartar emetic one grain and a-half  
Mix. A spoonful every quarter of an  
hour, till vomiting is produced, in croup.  
*Jadelot.*

R. Bruised seneka one drachm and a-half  
Boiling water sufficient  
to obtain four fl. ounces of the strained  
decoction; add

Antimonial wine two fl. scruples  
Syrup of marsh mallow one fl. ounce  
A small spoonful, occasionally, in the  
second stage of croup, as an expectorant.  
*Wendt.*

## EXTRACT OF SENEKA AND SQUILL.

R. Seneka  
Squill, each, in coarse  
powder, two pounds  
Proof spirit sufficient  
to cover; transfer to displacement appa-  
ratus; pass proof spirit through till near-  
ly tasteless; strain; distil off spirit, and  
evaporate to proper consistence. *Ecky.*

## POLYGALIC ACID.

R. Seneka, in powder, sixteen ounces  
Alcohol (35° B.) six pints  
Ether sufficient  
Purified animal charcoal two ounces  
Boil the seneka in four pints of the alcohol  
for fifteen minutes; suffer it to cool, then  
throw the whole on a displacement filter,  
and when the liquid ceases to pass add  
more alcohol, until six pints of tincture  
are obtained. Distil off the alcohol till of a  
syrupy consistence, wash this with ether,  
to remove fatty matter, and throw the  
residue into several times its bulk of con-  
centrated alcohol. After standing for  
twenty-four hours, the polygalic acid will  
be precipitated as a light brown powder,  
which is collected on a filter, washed with  
cold, strong alcohol, and dried. *W. Procter.*



SYRUP OF POLYGALIC ACID.

R. Polygalic acid                      eight grains  
Boiling water                      half a fl. ounce  
Syrup of gum  
Arabic                      one fl. ounce and a-half  
Mix, and make.  
Dose, a teaspoonful.

SENNA.

SENNA.

Under this name is included the leaves of several species of *Cassia*, as *C. acutifolia*, *C. elongata*, *C. obovata*, &c., forming what are termed Alexandrian, Tripoli, India, and Mecca sennas. They are all perennial plants, growing in warm climates.

*Ser. Syst.* Decand. monog. *Nat. Syst.* Fabaceæ.

Christison, Dispen. 858. Griffith, Med. Bot. 254.

The leaves only are official in the U. S. and British Pharms., but the pods are recognized by several of the European authorities. The leaves have a faint, but unpleasant odor, and a somewhat bitter, very nauseous taste. Senna is an active purgative, usually causing nausea and colic when given alone. It is seldom administered in substance, the dose is from half a drachm to two drachms.

COMPOUND POWDER OF SENNA.

R. Powdered senna  
Cremor tartar, each,                      two ounces  
Powdered scammony                      half an ounce  
ginger                      two drachms  
Mix. Dose, a scruple to a drachm.  
*Lond. Ph.*, 1824.

POWDER OF SENNA AND GUAIAIACUM.

R. Powdered senna  
guaiacum                      equal parts  
Dose, two teaspoonfuls, twice or thrice a-day. In jaundice. *Augustin.*

ALCOHOLIC EXTRACT OF SENNA.

R. Senna                      one part  
Alcohol                      four or five parts  
Heat slowly on a water-bath, to boiling point, let cool; in twenty-four hours express, and strain; repeat process with same quantity of alcohol, unite the tinctures, distil off the alcohol, and evaporate.  
*Guibourt.*

PILLS OF SENNA.

R. Powdered senna                      one drachm  
Extract of dandelion                      sufficient  
Mix, and make thirty pills. Dose, five to eight. *Hufeland.*

CONFECTION OF SENNA.

R. Senna                      eight ounces  
Coriander                      four ounces

Liquorice, bruised,                      three ounces  
Figs                      one pound  
Pulp of prunes  
tamarinds  
purging cassia, each,  
half a pound

Sugar                      two pounds and a-half  
Water                      four parts

Rub the senna and coriander together, and separate ten ounces with a sieve. Boil the residue with the figs and liquorice root in the water, to one-half, express, and strain. Evaporate by a water-bath to a pint and a-half, add the sugar and form a syrup, then rub the pulps with the syrup, add the sifted powder, and mix well. *U. S. Ph.*

A good laxative. Dose, about two drachms, taken at bedtime.

COMPOUND CONFECTION OF SENNA.

R. Confection of senna                      two ounces  
Powdered jalap                      one drachm  
Cremor tartar                      two drachms  
Powdered ginger one drachm and a-half  
Syrup                      sufficient  
Mix. Dose, one drachm. *Beasley.*

ELECTUARY OF SENNA AND RHUBARB.

R. Powdered rhubarb  
senna, each,                      two drachms  
fennel                      three drachms  
liquorice                      half an ounce  
Pulp of purging cassia  
tamarinds, each,                      two ounces  
Syrup of male fern                      sufficient  
Mix. Dose, two drachms to half an ounce. *Van Mons.*

ELECTUARY OF SENNA AND CREMOR TARTAR.

R. Powdered senna                      three drachms  
Cremor tartar                      one drachm  
Manna                      three ounces and a-half  
Pulp of purging cassia  
tamarinds,  
each, one ounce and a-half  
Syrup of violets                      sufficient  
Mix. Dose, about two drachms. *Van Mons.*

ELECTUARY OF SENNA AND FIGS.

R. Powdered senna                      half an ounce  
Pulp of figs                      sufficient  
Make an electuary. A piece the size of a hazelnut, night and morning, to obviate costiveness. *Ellis.*

ELECTUARY OF SENNA AND SULPHUR.

R. Powdered senna  
Flowers of sulphur  
Cremor tartar, each,                      one ounce



Manna six ounces  
 Pulp of tamarinds eight ounces  
 Syrup of senna sufficient  
 Mix. Said to be useful in hemorrhoids.  
 Dose, two to three drachms. *Swed. aur.*

## COMPOUND WINE OF SENNA.

R. Senna one ounce  
 Rhubarb six drachms  
 Cloves  
 Saffron, each, one drachm  
 Sherry wine two pints  
 Macerate for five days, often agitating, and decant. As a tonic purgative. Dose, about two fl. ounces. *Pierquin.*

R. Senna four ounces  
 Coriander  
 Fennel, each, two drachms  
 Sherry wine two pints and a-half  
 Digest for three days; add  
 Stoned raisins three ounces  
 Macerate for twenty-four hours, express, and strain. *Swed. Ph.*

## COMPOUND TINCTURE OF SENNA.

R. Senna three ounces and a-half  
 Caraway, bruised, three drachms and a-half  
 Cardamom, bruised, one drachm  
 Raisins five ounces  
 Proof spirit two pints  
 Macerate for fourteen days, express, and filter. *Lond. Ph.*

A warm, stomachic purgative, in atonic gout, &c., in doses of two fl. drachms to a fl. ounce.

## TINCTURE OF SENNA AND JALAP.

R. Senna three ounces  
 Jalap, in powder, one ounce  
 Coriander, bruised,  
 Caraway, bruised, each, half an ounce  
 Cardamom, bruised, two drachms  
 Sugar four ounces  
 Diluted alcohol three pints  
 Macerate for fourteen days, express, and filter. *U. S. Ph.*

Similar to the last, used in same cases, and in same doses.

## TINCTURE OF SENNA AND GENTIAN.

R. Senna four ounces  
 Gentian, bruised,  
 Orange peel, bruised, each, one ounce  
 Proof spirit sixty-four ounces  
 Infuse for some days, often stirring, express, and filter. *Swed. aur.*

## INFUSION OF SENNA.

R. Senna one ounce  
 Coriander, bruised, one drachm  
 Boiling water one pint

Macerate for an hour, in a covered vessel, and strain. *U. S. Ph.*

Dose, about four fl. ounces.

## COMPOUND INFUSION OF SENNA.

R. Senna half an ounce  
 Sulphate of magnesia  
 Manna, each, one ounce  
 Fennel seed one drachm  
 Boiling water half a pint  
 Macerate in a covered vessel till cool, and strain. Dose, one third, every four or five hours, till it operates. *Wood.*

R. Senna  
 Mint, each, ten drachms  
 Boiling water two pints  
 Macerate for an hour, strain, and add  
 Sulphate of magnesia eight ounces  
 Dose, two to four fl. ounces. *Guy's Hosp.*

## INFUSION OF SENNA AND TAMARINDS.

R. Senna one to three drachms  
 Tamarinds one ounce  
 Coriander, bruised, one drachm  
 Brown sugar half an ounce  
 Boiling water eight fl. ounces  
 Infuse for four hours in a covered vessel, and strain. Dose, two to four fl. ounces. *Ed. Ph.*

## INFUSION OF SENNA AND COFFEE.

R. Senna two drachms  
 Roasted coffee one drachm  
 Boiling water  
 Hot milk, each, three fl. ounces  
 Infuse for twelve hours, and strain. To be taken at once, in the morning. *Foy.*

R. Senna ten grains  
 Hot infusion of coffee  
 milk at will  
 Mix, and, when cool, strain, and add  
 Sugar sufficient  
 An agreeable purge for children. The amount of senna to be increased according to the age. *Guersant and Blake.*

R. Senna  
 Sulphate of magnesia  
 Coffee, roasted and ground, each, half an ounce  
 Boiling water four fl. ounces  
 Digest for half an hour, strain, and add  
 Sugar one and a-half ounces  
 As efficient as the "black draught," but much more agreeable. *Combes.*

R. Coffee, roasted and ground,  
 Senna, each, one ounce  
 Boiling water sufficient  
 Make an infusion, and evaporate to one ounce, and add



Simple syrup three fl. ounces

*Combes.*

INFUSION OF SENNA AND BUCKTHORN.

R. Senna two drachms  
Sulphate of soda half an ounce  
Boiling water four fl. ounces  
Syrup of buckthorn one fl. ounce  
Infuse, cool, and strain. *Ellis.*

INFUSION WITH TINCTURE OF SENNA.

R. Infusion of senna six fl. drachms  
Tincture of senna two fl. drachms  
Manna two drachms  
Mint water  
Distilled water, each, one fl. ounce and a-half

Mix. One-half at a dose; repeated, if it does not operate. *Abernethy.*

ANTHELMINTIC EMULSION.

R. Infusion of senna ten fl. drachms  
Syrup of buckthorn one fl. drachm  
Confection of scammony two scruples  
Copaiva thirty minims  
Spirit of turpentine six fl. drachms  
Mix, and make emulsion. Very efficient as an anthelmintic; to be taken early in the morning. *Med. Chirur. Rev.*

SENNÆ MIXTURE.

R. Infusion of senna one fl. ounce  
Tincture of jalap one fl. drachm  
Sulphate of magnesia four drachms  
Syrup of ginger one fl. drachm  
Mix. As a purgative draught. *Ainslie.*

INFUSION OF SENNA AND RHUBARB.

R. Senna six drachms  
Manna one ounce  
Bruised rhubarb  
cardamom, each, two drachms  
Boiling water one pint  
Infuse for one hour, and strain. Half a teacupful, every hour till it operates. *Ellis.*

INFUSION OF SENNA AND LEMON JUICE.

R. Senna one ounce and a-half  
Fresh lemon peel one ounce  
Lemon juice one fl. ounce  
Boiling water sixteen fl. ounces  
Infuse for two hours, and strain. *Beasley.*

SYRUP OF SENNA.

R. Senna two ounces  
Fennel seed, bruised, one ounce  
Boiling water one pint  
Sugar fifteen ounces  
Digest the senna and fennel seed in the

water, with a gentle heat, for an hour, strain, add the sugar, and form syrup. *U. S. Ph.*

Dose, one or two fl. drachms, to children.

FLUID EXTRACT OF SENNA.

R. Senna, coarsely powdered, thirty-two ounces  
Diluted alcohol four pints  
Water one gallon  
Sugar twenty ounces  
Oil of fennel sixty-four drops  
Hoffmann's anodyne sufficient  
Mix the senna and alcohol, and let rest for a night; introduce into a displacement apparatus, and, by means of the water, obtain one and a-half gallons of fluid; evaporate this to twenty fl. ounces, and, whilst hot, add the sugar; strain, and, when cold, add the oil of fennel, dissolved in the Hoffmann anodyne. Dose, a tablespoonful, as a full purgative. *Duhamel.*

R. Senna, well bruised, sixteen ounces (Troy)  
Diluted alcohol sufficient  
Oil of caraway  
Oil of anise, each, sixteen drops  
Sugar twelve ounces

Macerate the senna in two pints of the diluted alcohol for twenty-four hours; displace slowly with same menstruum until four pints pass. Evaporate this in a water-bath to ten ounces, and add the sugar; when this is dissolved, remove from the fire, and add the oils, dissolved in diluted alcohol. *Duhamel, modified by Procter.*

R. Senna fifteen pounds  
Boiling water sufficient  
Exhaust the senna by process of displacement, evaporate the fluid to ten pounds, add molasses, ten pounds, previously boiled to a candy; add further, twenty-four fl. ounces of rectified spirit, and water sufficient to make fifteen pints. Dose, two drachms. *Duncan.*

SYRUP OF SENNA AND RHUBARB.

R. Syrup of senna four fl. ounces  
rhubarb two fl. ounces  
cinnamon one fl. ounce  
Mix. *Van Mons.*

SYRUP OF SENNA AND APPLES.

R. Fermented juice of apples four pounds  
Senna four ounces and a-half  
Digest for a night, express, and strain, then add  
Sugar three pounds  
Clarify, and form syrup. *Sard. Ph.*



## SERPENTARIA.

## VIRGINIA SNAKEROOT.

Virginia snakeroot is furnished by several species of *Aristolochia*, as the *A. serpentaria*, *A. hirsuta*, and *A. reticulata*, all natives of the United States, especially the western portions.

*Sex. Syst.* Gynand. hexand. *Nat. Syst.* Aristolochiaceæ.

U. S. Dispens. 657. Griffith, Med. Bot. 529.

The root, which is the officinal portion, is very similar in the above-mentioned species. It is in tufts of slender, long, matted fibres, attached to a knotty, rugged head. These fibres are brittle, of a yellowish or brownish color, with an aromatic odor, and a pungent, bitter, camphorated taste. It is a stimulating tonic, and may also act as a diaphoretic, and diuretic. Dose, of the powder, is from ten to thirty grains.

## COMPOUND PILLS OF VIRGINIA SNAKEROOT.

R. Powdered Virginia  
snakeroot twenty-four grains  
Camphor forty-eight grains  
Conserve of roses sufficient  
Mix, and make eighteen pills; to be taken during the apyrexia, in malignant intermittents. *Foy.*

## ELECTUARY OF VIRGINIA SNAKEROOT.

R. Powdered Virginia snakeroot  
contrayerva  
Aromatic confection, each, one drachm  
Preserved ginger six drachms  
Syrup of parsley sufficient  
Make electuary. Half a drachm, every four hours, as a febrifuge. *Cadet.*

## INFUSION OF VIRGINIA SNAKEROOT.

R. Virginia snakeroot half an ounce  
Boiling water one pint  
Macerate for two hours in a covered vessel, and strain. *U. S. Ph.*

Dose, one to two fl. ounces, every two hours, in low forms of fever.

## COMPOUND INFUSION OF VIRGINIA SNAKEROOT.

R. Virginia snakeroot  
Contrayerva, each, five drachms  
Boiling water one pint  
Macerate for two hours, strain, and add  
Tincture of Virginia  
snakeroot two fl. ounces  
*Beasley.*

## TINCTURE OF VIRGINIA SNAKEROOT.

R. Virginia snakeroot,  
bruised, three ounces  
Diluted alcohol two pints  
Macerate for fourteen days, express, and filter. Dose, one to two fl. drachms.  
*U. S. Ph.*

## MIXTURE OF VIRGINIA SNAKEROOT AND ALLSPICE.

R. Comp. infus. of Virginia  
snakeroot twelve fl. ounces  
Tincture of allspice four fl. ounces  
Mix. Four spoonfuls, every six hours, in retrocedent and atonic exanthemata.

*Saunders.*

## ACETATED MIXTURE OF VIRGINIA SNAKEROOT AND ALLSPICE.

R. Add vinegar, two fl. ounces, to sixteen fl. ounces of last-mentioned mixture. Dose, as above, in petechial typhus. *Swedliaur.*

## INFUSION OF VIRGINIA SNAKEROOT AND ETHER.

R. Virginia snakeroot six drachms  
Boiling water eight fl. ounces  
Infuse, and strain; when cold, add  
Sulphuric ether two fl. drachms  
A tablespoonful, every hour. *Richter.*

## WINE OF VIRGINIA SNAKEROOT AND VANILLA.

R. Virginia snakeroot six drachms  
Vanilla two drachms  
Sherry wine sufficient  
Macerate for four hours, and add to each pint of the strained liquid,  
Camphor half a drachm  
Acetic ether one fl. drachm  
Syrup of cinnamon half a fl. ounce  
Dose, two spoonfuls an hour, in low states of the system. *Vogt.*

## TINCTURE OF VIRGINIA SNAKEROOT AND BALSAM OF PERU.

R. Virginia snakeroot six drachms  
Proof spirit five fl. ounces  
Water sufficient  
to obtain six fl. ounces of strained fluid; after one hour of infusion, add  
Camphor half a drachm  
Balsam of Peru one drachm  
Gum Arabic two drachms  
Make an emulsion, and add  
Acetic ether one fl. drachm  
A spoonful, every hour, in cases requiring stimulation. *Phœbus.*

## SESAMUM.

## BENNE.

Two species of *Sesamum*, the *S. orientale*, and the *S. indicum*, afford leaves and seeds that are recognized as officinal. They are both natives of Asia, and are cultivated in the West Indies and in the southern parts of the United States.

*Sex. Syst.* Didyn. angios. *Nat. Syst.* Pedaliaceæ.



Linn. Sp. Pl. 883, 884. Griffith, Med. Bot. 515.

The seeds are small, inodorous, and afford a bland oil, which may be used for all the purposes of olive oil. The leaves abound in a gummy substance which forms, with water, a rich, bland mucilage, useful in all cases requiring the use of demulcents.

#### INFUSION OF BENNE.

R. Fresh leaves of benne No. 2  
Cool water half a pint  
Infuse till a mucilage is formed. When dried leaves are used, the water should be hot. Wood.

### SIMARUBA.

#### SIMARUBA.

This is the bark of the root of *Simaruba officinalis* (amara), a large tree, native of Jamaica, and several parts of South America.

Sex. Syst. Decand. monog. Nat. Syst. Simarubaceæ.

Decand. i. 733. Griffith, Med. Bot. 198.

The bark is in long pieces, much rolled or quilled. It is fibrous, tough; of a grayish color externally, and yellowish within. It is inodorous, but extremely bitter. It has the properties of the pure bitters, but is apt to act on the stomach and bowels.

#### INFUSION OF SIMARUBA.

R. Simaruba bark, bruised, three drachms  
Boiling water one pint  
Macerate for two hours, and strain. Ed. Ph.

Dose, two fl. ounces.

#### COMPOUND INFUSION OF SIMARUBA.

R. Simaruba  
Wormwood, each, two drachms  
Boiling water one pint  
Infuse for fifteen minutes; strain, and add

Syrup of gentian one fl. ounce  
To be taken in wineglassful doses. Foy.

### SINAPIS.

#### MUSTARD.

The seeds of two species of Sinapis are used in medicine, those of *S. nigra* and *S. alba*, both annual plants indigenous to many parts of Europe, and cultivated in our gardens.

Sex. Syst. Tetradyn. siliq. Nat. Syst. Brassicaceæ.

Linn. Sp. Pl. 933. Griffith, Med. Bot. 133.

Black mustard seeds are small, globular, of a deep brown color, when bruised having a peculiar smell, which becomes pungent, on the flour being mixed with water; their taste is bitterish and pungent. White mustard seeds are larger, of a yellowish color, and less pungent. These seeds act as a laxative; when in the

state of flour, they operate as an emetic, and when applied to the skin as a rubefacient, or even vesicant.

#### CATAPLASM OF MUSTARD.

R. Flour of mustard  
Flaxseed meal, each, half a pound  
Boiling vinegar sufficient  
Form a cataplasm. Lond. Ph.

R. Flour of mustard two ounces  
Crumb of bread four ounces  
Common salt half an ounce  
Vinegar sufficient  
Mix. Ammon.

Used as rubefacients and revulsives.

#### COMPOUND CATAPLASM OF MUSTARD.

R. Flour of mustard eight ounces  
Powdered white pepper  
ginger, each, one drachm  
Oxymel sufficient  
Mix. Foy.

R. Flour of mustard four ounces  
Yeast one ounce and a-half  
Muriate of ammonia one drachm  
Rue half an ounce  
Vinegar sufficient  
Mix. Span. Ph.

Used as above.

#### MUSTARD FOMENTATION.

R. Flour of mustard four ounces  
Hot water one pint  
Mix. As a rubefacient. Ellis.

#### MUSTARD GARGLE.

R. Flour of mustard one drachm  
Water of angelica  
hyssop, each, six fl. ounces  
Spirit of horseradish half a fl. ounce  
muriate of ammonia four fl. scruples  
Honey of roses four ounces  
Mix. Spielman.

#### MUSTARD FOOT BATH.

R. Flour of mustard four ounces  
Hot water sufficient  
Mix. Beral.

#### MUSTARD WHEY.

R. Milk one pint  
Water two pounds  
Bruised mustard-seed one ounce and a-half  
Boil till curdled, and strain. Beasley.

R. Milk one pint  
Bruised mustard seed one ounce  
Boil till curdled, and strain, then add  
Sherry wine six fl. ounces  
A draught every hour or two, in low states of the system. Ellis.



### MUSTARD OINTMENT.

R. Flour of mustard	three ounces
Oil of almonds	half a fl. ounce
Lemon juice	sufficient
Mix. In sun-burn, &c.	<i>Frank.</i>

COMPOUND MUSTARD OINTMENT.

R. Resin ointment	one ounce
Soap	half an ounce
Camphor	two scruples
Flour of mustard	one scruple
Mix.	<i>Ferriar.</i>

### LOTION OF MUSTARD.

R. Mustard flour                      one ounce  
Express the oil, and add to the mass  
Oil of turpentine                    eight ounces  
Macerate for three days, express, and  
strain. As a friction in paralysis.  
*Niemann.*

VOLATILE OIL OF MUSTARD.

R. Bruised mustard seed    twenty pounds  
Cold water    sufficient  
Distil, and separate the oil.    *Magendie.*  
A very active rubefacient.

TINCTURE OF OIL OF MUSTARD.

R. Volatile oil of mustard    twelve parts  
 Alcohol    two hundred and fifty parts  
 Dissolve.    A good rubefacient.  
*Soubeiran.*

### LINIMENT OF MUSTARD.

R. Lard  
Oil of almonds, each, twelve ounces  
Yellow wax four ounces  
Melt by a gentle heat, pour into a heated mortar, and when cooling add  
Oil of lavender three ounces  
Carbonate of ammonia half an ounce  
Tincture of mustard two fl. ounces  
Camphor one ounce  
Triturate the ammonia with the oil; dissolve the camphor in the tincture, and add both to the melted mass, incorporating well. As an application to chilblains.

*Beral.*

## SODIUM.

## SODIUM.

SODII BROMIDUM.

BROMIDE OF SODA.

R. Solution of bromide of iron            at will  
   carbonate of soda sufficient  
to precipitate; filter, and evaporate.  
*Magendie.*

OINTMENT OF BROMIDE OF SODA.

R. Bromide of sodium thirty-four grains  
Lard one ounce  
Mix. As an application to obstinate  
cutaneous eruptions.

SODII CHLORIDUM.

### COMMON SALT.

POWDER OF COMMON SALT AND COCHINEAL.

P. Powdered common salt three drachms  
 cochineal fifteen grains  
 Mix, and divide into six powders. Anthelmintic. One every morning, and finally a dose of some purgative. *Rush.*

COMPOUND POWDER OF COMMON SALT.

R. Common salt  
Sulphate of magnesia, each, four ounces  
Sulphate of potassa three ounces  
Dry the salts separately, and pulverize; rub  
them well together, and keep in well-closed  
vessels. *Ed. Ph.*

### BATH OF COMMON SALT AND GELATINE.

R. Common salt	one pound
Water	four pints
Pour the solution in a bath, and add	
Flanders glue	two pounds
Dissolved in	
Water	six pints
Recommended in scrofulous affections.	
	<i>Foy.</i>

### FOMENTATION OF COMMON SALT.

R. Common salt                                two ounces  
Water    six fl. ounces  
Vinegar  
Brandy, each,                               three fl. ounces  
Mix. As a fomentation to bruises. *Vogt.*

CLYSTER WITH COMMON SALT.

R. Common salt	one ounce
Barley water	half a pint
Add to the solution	
Olive oil	one fl. ounce
	<i>Mid. Hosp.</i>

R. Common salt                      one to two ounces  
Tepid flaxseed infusion                      one pint  
Dissolve, and add  
Castor or olive oil  
Molasses, each,                      two fl. ounces  
One-half to be used at a time, the remain-  
der in half an hour afterwards, if needed.  
*Ellis.*



CLYSTER OF COMMON SALT AND ARNICA.  
 R. Common salt half to one ounce  
 Infusion of arnica twelve fl. ounces  
 Said to be useful in apoplexy and paralysis.

*Foy.*

MIXTURE OF COMMON SALT AND LEMON JUICE.

R. Lemon juice at will  
 Common salt sufficient  
 to saturate, filter. A tablespoonful every hour, in apyrexia of intermittents. *Bories.*

OINTMENT OF COMMON SALT.

R. Common salt one to four drachms  
 Lard one ounce  
 Rub together till smooth. The strength to be gradually increased from one drachm of the salt to four. To inflamed eyelids.

*Tavignot.*

COMPOUND OINTMENT OF COMMON SALT.

R. Common salt one ounce and a-half  
 Simple cerate three ounces  
 Rose water ointment one ounce

Rub together. In tinea capitis, as a friction, morning and evening.

*St. Marie.*

R. Common salt two drachms  
 Nut oil one drachm  
 Ox gall one ounce and a-half

Digest together for thirty-six hours, and triturate well. As a friction to scrofulous tumors.

*Roncalli.*

## SODII SULPHURETUM.

SULPHURET OF SODA.

R. Flowers of sulphur two hundred parts  
 Carbonate of potassa, dried, two hundred and sixty-seven parts  
 Mix; heat in a matrass till melted; on cooling, detach, and keep in a well-closed vessel.

*Cottureau.*

BATH WITH SULPHURET OF SODIUM.

R. Sulphuret of sodium ten ounces  
 Solution of common salt and gelatin four fl. ounces  
 Mix with the water of the bath, at the moment the patient enters it. In chronic cutaneous affections.

*Cadet.*

MIXTURE OF SULPHURET OF SODIUM AND SAL AMMONIAC.

R. Sulphuret of sodium one drachm  
 Muriate of ammonia fifty grains  
 Dissolve each in six ounces of water, mix the solutions, after having filtered them. As a lotion in itch, &c.

*Van Mons.*

LOTION WITH SULPHURET OF SODIUM.

R. Sulphuret of sodium three ounces

Soap one ounce and a-half  
 Dissolve in  
 Alcohol one pound  
 Add to the solution  
 Lime water eight pounds  
 In tinea capitis; the head to be washed with it every day, without removing the hair.

LINIMENT OF SULPHURET OF SODIUM.

R. Sulphuret of sodium four parts and a-half  
 Flowers of sulphur  
 Oil of almonds, each, one part and a-half  
 Lard six parts and a-half  
 Rub the powders with the lard, and gradually add the oil. As a lotion in chronic cutaneous diseases.

*Van Mons.*

OINTMENT OF SULPHURET OF SODIUM.

R. Sulphuret of sodium three drachms  
 Lard one ounce and a-half  
 Rub well together. As an application in itch.

*Swediaur.*

## SODA.

CAUSTIC SODA.

R. Solution of soda at will  
 Evaporate in a silver capsule, till it will solidify, on cooling; pour into moulds.

*Guibourt.*

SOLUTION OF SODA.

R. Crystallized carbonate of soda twenty-five parts  
 Quicklime six parts  
 Boil with a sufficient quantity of water, strain, and evaporate, till it marks 36° of the areometer.

*Guibourt.*

## SODÆ ACETAS.

ACETATE OF SODA.

R. Carbonate of soda at will  
 Distilled vinegar sufficient  
 to saturate; evaporate to sp. gr. 1.276, and set aside to crystallize.  
 Diuretic or cathartic, according to dose, from one scruple to four drachms.

*Dub. Ph.*

COMPOUND PILLS OF ACETATE OF SODA.

R. Acetate of soda five drachms  
 Camphor  
 Nitre, each, two drachms  
 Conserve of elder berries sufficient  
 Mix, and make one hundred and twenty pills. Dose, two, morning and evening, to check the secretion of milk.

*Bories.*



R. Acetate of soda  
Powdered rhubarb  
Inspissated ox gall, each, one drachm  
Powdered gum Arabic half a drachm  
Mix, and make pills of two grains. Dose,  
ten to fifteen, twice a-day, as a laxative.  
*Augustin.*

## SODÆ ARSENIAS.

## ARSENIATE OF SODA.

R. White arsenic one hundred parts  
Nitrate of soda one hundred and sixteen parts  
Pulverize, and introduce into a luted retort;  
heat in a reverberatory furnace, break the  
retort, dissolve the salt in water, add car-  
bonate of soda, till the solution is neutral,  
filter, evaporate, and crystallize. In inter-  
mittents, and some diseases of the skin.  
Dose, one-sixteenth to one-eighth of a  
grain. *Cottureau.*

## SOLUTION OF ARSENIATE OF SODA.

R. Arseniate of soda six grains  
Peppermint water two fl. ounces and a-half  
Add to the solution  
Wine of cinnamon half a fl. ounce  
Laudanum one fl. drachm  
Dose, forty to fifty drops, four times a-day.  
*Heineke.*

R. Crystallized arseniate of soda one grain  
Distilled water one fl. ounce  
Dissolve. Dose, from a scruple to half a  
drachm. *Foy.*

## SODÆ BORAS.

## BORAX.

## COMPOUND POWDER OF BORAX.

R. Powdered borax one drachm  
saffron ten grains  
Oil of cinnamon two drops  
Mix, for two doses. As an emmena-  
gogue. *Wedel.*  
R. Powdered borax half an ounce  
castor  
saffron, each,  
one drachm and a-half  
Oil of cinnamon eight drops  
amber six drops  
Mix, and form powder. Dose, from a  
scruple to half a drachm, to facilitate par-  
turation, expel placenta, &c. *Wert. Ph.*

## LOZENGES OF BORAX.

R. Powdered borax two drachms  
sugar half an ounce

Mucilage sufficient  
Mix, and make thirty lozenges. One,  
occasionally, in aphthous sore mouth.  
*Phœbus.*

## COLLUTORY OF BORAX.

R. Borax two drachms  
Water  
Tincture of myrrh, each, one fl. ounce  
Honey of roses two ounces  
Mix. *Svediaur.*

R. Borax three drachms  
Mucilage of quince  
seeds eight fl. ounces  
Honey of roses two ounces  
Mix. *Bahi.*

R. Borax three drachms  
Infusion of roses one fl. ounce and a-half  
Honey of roses two ounces  
Mix. *Pringle.*

## COLLYRIUM OF BORAX.

R. Borax one drachm  
Laudanum half fl. drachm  
Rose water four fl. ounces  
Mix. *Fricke.*

R. Borax half a drachm  
Sugar one drachm  
Rose water two fl. ounces  
Mix. *Richard.*

## GARGLE OF BORAX.

R. Borax two drachms  
Oxymel half fl. ounce  
Water eleven fl. ounces  
Mix. *Beasley.*

R. Borax one drachm  
Tincture of myrrh half fl. ounce  
Clarified honey one fl. ounce  
Rose water four fl. ounces.  
Mix. *Ellis.*

In aphthous sore mouth, ptyalism, &c.

## LOTION OF BORAX.

R. Borax half a drachm  
Orange-flower water  
Rose water, each, half fl. ounce  
Dissolve. In pruritus. *Cadet.*

## HONEY OF BORAX.

R. Powdered borax one drachm  
Clarified honey one ounce  
Mix. *Lond. Ph.*  
In aphtha.

## VINEGAR OF BORAX.

R. Distilled vinegar two fl. ounces  
Borax one drachm  
Dissolve. As lotion to ringworm of the  
scalp. *Christison.*



## MIXTURE OF BORAX.

- R. Borax two drachms  
 Chamomile water four fl. ounces  
 Syrup of orange peel half fl. ounce  
 Liquid succinate of ammonia two fl. drachms  
 Wine of opium half fl. drachm  
 Mix. As an emmenagogue, and to facilitate parturition. Dose, two spoonfuls, every half hour. *Swedjaur.*

## OINTMENT OF BORAX.

- R. Powdered borax fifteen grains  
 Fresh butter two drachms  
 Rub together. To apply to scaly eruptions. *Radiis.*  
 R. Powdered borax two drachms  
 Rose ointment one ounce  
 Mix. For chilblains. *Cadet.*

## SODÆ CARBONAS.

## CARBONATE OF SODA.

## DRIED CARBONATE OF SODA.

- R. Carbonate of soda at will  
 Heat in a clean iron vessel, till perfectly dried, stirring constantly; rub into powder. *U. S. Ph.*

## SOLUTION OF CARBONATE OF SODA.

- R. Carbonate of soda one ounce  
 Distilled water one pint  
 Dissolve. *Dub. Ph.*

Dose, one to two fl. ounces diluted, two or three times a-day, as antacid, &c.

## BICARBONATE OF SODA.

- R. Crystallized carbonate of soda at will  
 Saturate in proper apparatus, with carbonic acid.

- R. Carbonate of soda one part  
 Water two parts  
 Dissolve, and pass a current of carbonic acid through the solution; let crystallize. *Cottureau.*

## EFFERVESCENT SOLUTION OF SODA.

- R. Bicarbonate of soda one drachm  
 Distilled water one pint  
 Dissolve, and force into it more carbonic acid. Keep in a well-closed vessel. *Lond. Ph.*

## EFFERVESCENT POWDERS OF SODA.

- R. Bicarbonate of soda one ounce, fifty-four grains  
 Tartaric acid one ounce  
 Powder each, separately, and divide each into sixteen powders, and preserve in papers of different colors. *Ed. Ph.*

Known under the name of *Soda powders.*

## PILLS OF SODA AND RHUBARB.

- R. Powdered rhubarb  
 Dried carbonate of soda  
 Extract of gentian, each, one scruple  
 Calomel three grains  
 Mix, and make twenty pills. Two occasionally, in dyspepsia. *Ellis.*

## PILLS OF SODA AND IPECACUANHA.

- R. Carbonate of soda one drachm  
 Bitter almonds half a drachm  
 Ipecacuanha three grains  
 Extract of madder twenty-four grains  
 Water sufficient  
 Beat into mass, and form sixty pills. Three morning and evening, in obstinate cutaneous diseases. *Foy.*

## CARBONATE OF SODA PILLS.

- R. Extract of chamomile one drachm  
 Powdered rhubarb one scruple  
 Carbonate of soda half a drachm  
 Oil of caraway ten minims  
 Syrup of ginger sufficient  
 Mix, and make twenty-four pills, two to be taken thrice a-day, in dyspepsia. *Ainslie.*

## PILLS OF CARBONATE OF SODA.

- R. Dried carbonate of soda one drachm  
 Soap two scruples  
 Water sufficient  
 Beat together, and make thirty pills. Three, thrice a-day. In calculous affections. *Ellis.*

## POWDER OF CARBONATE OF SODA AND MAGNESIA.

- R. Bicarbonate of soda  
 Magnesia  
 Powdered ginger, each, one scruple  
 Mix. A small teaspoonful, occasionally. *Ellis.*

## POWDER OF CARBONATE OF SODA AND RHUBARB.

- R. Dried carbonate of soda one ounce  
 Rhubarb four drachms  
 Ginger one scruple  
 Mix. In cardialgia, and dyspepsia. *Beasley.*

## POWDER OF CARBONATE OF SODA AND MERCURY.

- R. Dried carbonate of soda five drachms  
 Calomel one drachm  
 Compound chalk powder ten drachms  
 Mix. Dose, eight to sixteen grains. *Guy's Hosp.*

## LOZENGES OF BICARBONATE OF SODA.

- R. Bicarbonate of soda one ounce  
 Powdered gum Arabic half an ounce  
 sugar three ounces



Mucilage sufficient  
Beat together, and form lozenges.

*Ed. Ph.*

Antacid and antilithic.

R. Bicarbonate of soda two ounces  
Sugar thirty-four ounces  
Oil of peppermint eighteen drops  
Mucilage thirty-six drachms  
Make lozenges of eighteen grains.

*Magendie.*

R. Bicarbonate of soda four parts  
Sugar sixty parts  
Chocolate eight parts  
Mucilage nine parts  
Make lozenges of eighteen grains.

*Beral.*

R. Bicarbonate of soda four drachms  
Sugar ten ounces  
Balsam of tolu one drachm  
Mucilage of tragacanth sufficient  
Make lozenges of twenty grains.

*Cadet.*

#### LOTION OF CARBONATE OF SODA.

R. Saturated solution of  
carbonate of soda one ounce  
Warm water two pints  
Mix. As a lotion in pruritus.

*Radius.*

#### MIXTURE OF CARBONATE OF SODA AND QUASSIA.

R. Carbonate of soda two drachms and a-half  
Rasped quassia one drachm and a-half  
Boiling water one pint  
Infuse for one hour, and strain. Two to four spoonfuls a-day, in dyspepsia.

*Augustin.*

#### MIXTURE OF CARBONATE OF SODA AND GENTIAN.

R. Carbonate of soda four scruples  
Compound infusion of gentian  
Cinnamon water, each, three fl. ounces  
Tincture of cardamom half a fl. ounce  
Mix. A tablespoonful, every two or three hours, in acidity and flatulence.

*Ellis.*

#### MIXTURE OF CARBONATE OF SODA AND COPAIBA.

R. Bicarbonate of soda one drachm  
Almond mixture four fl. ounces  
Copaiba two fl. drachms  
Laudanum sixty drops  
Mix. A tablespoonful occasionally, in acute pain in the bladder.

*Ellis.*

#### MIXTURE OF CARBONATE OF SODA AND CHAMOMILE.

R. Carbonate of soda one drachm  
Chamomile water three fl. ounces  
Syrup of gentian one fl. ounce

#### Tincture of Peruvian

bark one fl. drachm  
Mix. A spoonful, occasionally, in serofula.

*Foy.*

#### MIXTURE OF CARBONATE OF SODA AND IPECACUANHA.

R. Carbonate of soda twelve grains  
Wine of ipecacuanha twenty drops  
Laudanum four drops  
Distilled water one fl. ounce

Mix. A teaspoonful, every two or three hours, in pertussis of children.

*Ellis.*

#### DIURETIC SOLUTION WITH CARBONATE OF SODA.

R. Carbonate of soda three drachms  
Nitrate one drachm and a-half  
Honey one ounce and a-half  
Decoction of couchgrass three pints  
Mix. A glassful every hour, in dropsy.

*Cadet.*

#### ANTI-EMETIC MIXTURE.

R. Bicarbonate of soda one drachm and a-half  
Powdered gum Arabic two drachms  
Oil of mint four drops  
White sugar two drachms  
Carbonic acid water four ounces  
Mix. A tablespoonful occasionally, to calm stomach, or prevent vomiting.

*Dewees.*

#### INJECTION OF CARBONATE OF SODA.

R. Carbonate of soda one drachm  
Soap two ounces  
Water twelve fl. ounces  
Advised as an injection, in cases of uric calculi.

*Bories.*

#### EMULSION WITH CARBONATE OF SODA.

R. Bicarbonate of soda sixteen grains  
Nitrate four grains  
Syrup of marsh mallow  
violets, each, one fl. ounce  
Lettuce water two fl. ounces  
Emulsion of almonds twelve fl. ounces  
Mix. In nephritic complaints.

*Beral.*

#### OINTMENT OF CARBONATE OF SODA.

R. Carbonate of soda two drachms  
Wine of opium one fl. drachm  
Lard one fl. ounce  
Mix. In cutaneous affections, where there is pain.

*Soubeiran.*

R. Carbonate of soda  
Sulphuret of sodium, each, three drachms  
Lard one ounce  
Mix. In tinea capitis, having removed the crusts.

*Alibert.*



- R. Impure carbonate of soda three drachms  
 Slaked lime two drachms  
 Lard two ounces  
 Mix. As the last. *Cadet.*  
 R. Carbonate of soda two drachms  
 Slaked lime one drachm  
 Opium two grains  
 Lard two ounces  
 Mix. In prurigo. *Bielt.*

## SODÆ CHLORAS.

## CHLORATE OF SODA.

- R. Tartaric acid five ounces  
 Boiling water two pints  
 Dissolve, and add  
 Carbonate of  
 soda forty-five drachms and a-half  
 Also, dissolve  
 Chlorate of  
 potassa four ounces, fifteen grains  
 Boiling water sixteen fl. ounces  
 Mix the solutions whilst boiling, filter,  
 evaporate, and crystallize. *Hamb. Ph.*

## LOTION OF CHLORATE OF SODA.

- R. Chlorate of soda two drachms  
 Water four fl. ounces  
 In pruritus. *Darling.*

## GARGLE OF CHLORATE OF SODA.

- R. Chlorate of soda one to three scruples  
 Barley water three ounces  
 Honey of roses one ounce  
 Mix. *Radius.*

SODÆ CHLORINATÆ LI-  
QUOR.

## SOLUTION OF CHLORIDE OF SODA.

- R. Chlorinated lime one pound  
 Carbonate of soda two pounds  
 Water one gallon and a-half  
 Dissolve the carbonate of soda in three pints  
 of the water, with the aid of heat. Add  
 to the remainder of the water the chlorin-  
 ated lime, in powder, in small portions, at  
 a time. Set aside for some hours, to set-  
 tle; decant clear solution, and add that  
 of the carbonate of soda; decant, and  
 strain through linen, and keep in opaque  
 bottles. *U. S. Ph.*  
 R. Carbonate of soda one part  
 Water ten parts  
 Dissolve, and pass a current of chlorine  
 through the solution. *Guibourt.*

This is known as *Labarraque's Disinfecting So-  
 lution*; it is used as a disinfectant, but is also

used internally, in doses of thirty drops to one  
 fl. drachm, in low fevers, &c.

DILUTED SOLUTION OF CHLORIDE OF  
SODA.

- R. Solution of chloride  
 of soda half a fl. drachm  
 Distilled water four fl. ounces  
 To be taken in spoonful doses, in the apy-  
 rexia of intermittents. *Gouzee.*  
 R. Solution of chloride soda one part  
 Water twelve to sixteen parts  
 As a lotion to foul and cancerous ulcers.  
*Foy.*

## GARGLE OF CHLORIDE OF SODA.

- R. Chloride of soda sixteen drops  
 Hydromel one fl. ounce  
 Distilled water seven fl. ounces  
 Mix. *Beral.*

## SODÆ HYDROSULPHAS.

## HYDROSULPHATE OF SODA.

- R. Carbonate of soda ten ounces  
 Distilled water one pint  
 Dissolve, and mix with the solution  
 Flowers of sulphur ten drachms  
 Then pass an excess of sulphurous acid  
 gas into the liquid; boil for a few minutes  
 in a matrass; filter, and evaporate to one  
 third, and set aside to crystallize.  
*Par. Cod.*

Used in diseases of the skin, in doses of ten  
 grains to one drachm.

- R. Subcarbonate of soda  
 seven hundred and thirty parts  
 Sulphur forty-five parts  
 Water fifteen hundred parts  
 Mix the sulphur with a small portion of  
 the water, and add the soda, dissolved in  
 the rest. Introduce into two two-necked  
 bottles, so as to fill them about two-thirds.  
 Clean iron filings fifteen hundred parts  
 Sulphuric acid  
 (1.84) three thousand parts  
 Put into a flask, let cool, place on a sand-  
 bath, and convey the gas by rather large  
 tubes, first into a washing bottle, and then  
 through the solution in the two-necked  
 bottles. Continue the process for ten or  
 twelve hours, mix the contents of the two  
 bottles, filter, and evaporate to crystalli-  
 zation. *Legrip.*

Principally used in daguerreotyping, and as  
 above.

## SYRUP OF HYPOSULPHITE OF SODA.

- R. Hyposulphite of soda one ounce  
 Water twelve fl. ounces



Sugar twenty-three ounces  
Dissolve the salt in the water, add the sugar, and form syrup. Dose, from one to two ounces, twice a-day. *Mouchon.*

## SODÆ PHOSPHAS.

## PHOSPHATE OF SODA.

R. Powdered burnt bones ten pounds  
Sulphuric acid six pounds  
Carbonate of soda sufficient  
Mix the powdered bone with the acid in an earthen vessel, add a gallon of water, and stir. Digest for three days, occasionally adding a little water, then pour in a gallon of boiling water, strain; adding boiling water till the liquid passes almost tasteless. Let settle, decant, and boil down to a gallon. After settling, pour it into an iron vessel, heat, and gradually add the carbonate of soda dissolved in hot water, as long as there is effervescence, then filter, and let crystallize. More crystals will be obtained by adding carbonate of soda in excess, and again evaporating to crystallization. Keep in a well-stopped bottle. *U. S. Ph.*

A mild purgative, in doses of four drachms to one ounce.

R. Acid of phosphate of lime at will  
Add carbonate of soda to saturation, let settle, decant, evaporate, and crystallize. *Guibourt.*

## COMPOUND POWDER OF PHOSPHATE OF SODA.

R. Phosphate of soda two drachms  
Carbonate of soda one drachm  
Fennel sugar half an ounce  
Mix. Two to three teaspoonfuls a-day, to aid digestion. *Berands.*

## MIXTURE OF PHOSPHATE OF SODA.

R. Phosphate of soda half an ounce  
Decoction of carrageen six fl. ounces  
Syrup of orgeat half an ounce  
Mix. As a purgative. *Radius.*

R. Phosphate of soda  
Syrup of marsh mallow, each, half an ounce  
Water four fl. ounces  
Mix. As a purgative. *Augustin.*

## SODÆ SULPHAS.

## SULPHATE OF SODA.

## POWDER OF SULPHATE OF SODA AND TARTAR EMETIC.

R. Sulphate of soda two ounces

Tartar emetic  
Opium, each, one grain  
Gum Arabic  
Powdered liquorice, each, one drachm  
Mix, to be taken in water, in divided doses during the day, as a purgative and diuretic. *Swedjaur.*

## POWDER OF SULPHATE OF SODA AND NITRE.

R. Dried sulphate of soda eighteen drachms  
Nitro half a drachm  
Tartar emetic one grain  
Mix. One third in water or broth. *Beasley.*

## POWDER OF SULPHATE OF SODA AND OPIUM.

R. Sulphate of soda four scruples  
Powdered opium two grains  
Mix. In hemorrhagies and inflammations, after bleeding. *Radius.*

## COMPOUND PILLS OF SULPHATE OF SODA.

R. Sulphate of soda half a drachm  
Bitter almonds one drachm  
Ipecacuanha two grains  
Extract of madder sufficient  
Beat into mass, and make sixty pills. Three, morning and evening, in obstinate cutaneous affections. *Foy.*

## ELECTUARY WITH SULPHATE OF SODA.

R. Dried sulphate of soda half an ounce  
Pulp of tamarinds one ounce  
Syrup of lemon juice sufficient  
Mix. Two spoonfuls every hour, till it operates. *Phæbus.*

## COMPOUND SOLUTION OF SULPHATE OF SODA.

R. Sulphate of soda fourteen drachms  
Acetate of potassa twenty-four grains  
Nitro eighteen grains  
Tartar emetic quarter of a grain  
Water two pints  
Dissolve, and filter; in wineglassfuls, in habitual constipation. *Soubéiran.*

## EMULSION WITH SULPHATE OF SODA.

R. Sulphate of soda one ounce  
Oil of almonds one fl. ounce  
Yolk of eggs two  
Syrup of manna two fl. ounces  
Fennel water six fl. ounces  
Mix. Dose, two spoonfuls. *Phæbus.*

## LEMONADE WITH SULPHATE OF SODA.

R. Sulphate of soda one ounce and a-half  
Lemon juice half an ounce  
Sugar three ounces



Water sixteen fl. ounces  
Mix. *Radius.*

## CLYSTER OF SULPHATE OF SODA.

R. Sulphate of soda one ounce  
Barley water six fl. ounces  
Dissolve, and add  
Oil of flaxseed two ounces  
*Swedjaur.*

## LOTION OF SULPHATE OF SODA.

R. Sulphate of soda six ounces  
Soap three drachms  
Lime water eighteen fl. ounces  
Rectified spirit six fl. drachms  
Mix. In chronic cutaneous affections.  
*Van Mons.*

## SUPPOSITORY OF SULPHATE OF SODA.

R. Sulphate of soda  
Soap, each, half an ounce  
Honey sufficient  
Mix, and make four suppositories.  
*Phœbus.*

## BISULPHATE OF SODA.

R. Dried sulphate of soda ten parts  
Sulphuric acid seven parts  
Heat gently in a crucible. Purgative in  
doses from two drachms to six. *Beasley.*

## BISULPHATE OF SODA AND MAGNESIA.

R. Bisulphate of soda at will  
Hot water sufficient  
Dissolve, add magnesia to saturation; filter, and crystallize. Laxative and diuretic, in doses from half a drachm to two drachms, several times a-day. *Van Mons.*

SODÆ ET POTASSÆ  
TARTRAS.

## TARTARIZED SODA OR ROCHELLE SALT.

R. Carbonate of soda one pound  
Cremor tartar sixteen ounces  
Boiling water five pints  
Dissolve the carbonate of soda in the water, and add gradually the cremor tartar; filter, and evaporate to a pellicle, and set aside to crystallize. Decant, and dry crystals.  
*U. S. Ph.*

Mild purgative, in doses of half an ounce to an ounce.

POWDER OF TARTARIZED SODA AND  
RHUBARB.

R. Tartarized soda ten grains  
Powdered rhubarb three to seven grains  
Mix. To be taken every morning for two weeks, in enlargements of the mesentery, in children.  
*Fordyce.*

## SEIDLITZ POWDERS.

R. Tartarized soda two drachms  
Bicarbonate of soda two scruples  
Mix, and dissolve in half a tumbler of water,  
Powdered tartaric acid twenty-five grains  
Dissolve in an equal quantity of water, as above, and mix; to be taken during effervescence. *Ellis.*

## MIXTURE OF TARTARIZED SODA.

R. Tartarized soda half an ounce  
Sulphate of potassa two drachms  
Infusion of succory one pint  
Oxymel of squill four fl. ounces  
Mix. Four spoonfuls, in the morning, with an hour's interval. In visceral obstructions. *St. Marie.*  
R. Tartarized soda one ounce  
Extract of henbane six grains  
Syrup of chamomile one fl. ounce  
Water six fl. ounces  
Mix. A spoonful every hour, in constipation. *Phœbus.*

## WHEY WITH TARTARIZED SODA.

R. Root and leaves of dandelion  
Fumitory  
Water-cress  
Chervil, each, a handful  
Clarified whey four pints  
Boil for five minutes, express, strain, and add  
Tartarized soda two drachms  
Honey six drachms  
To be taken every morning, early, in four doses, with the interval of an hour. Much praised in visceral obstructions. *St. Marie.*

## CLYSTER WITH TARTARIZED SODA.

R. Tartarized soda  
Extract of dandelion, each, five drachms  
Infusion of elder flowers five fl. ounces  
Dissolve. *Radius.*

## SPIGELIA.

## PINKROOT.

This is the root of *Spigelia marilandica*, a beautiful, herbaceous, perennial plant, a native of the southern and south-western parts of the United States.

*Sex. Syst.* Pentand. monog. *Nat. Syst.* Loganiaceæ.

*Linn. Sp. Pl.* 249. *Griffith, Med. Bot.* 466.

The root consists of numerous small, blackish fibres, forming a dense bunch, and arising from a short rhizome. The odor is faint, and the taste sweetish, bitter, and unpleasant. It is a powerful and certain anthelmintic, with some purgative



and narcotic properties. It is given in powder, in the dose of one to two drachms for an adult, ten to twenty grains to a child; to be repeated morning and evening, for a few days, and followed by an active purgative.

#### PINKROOT WITH CALOMEL.

R. Powdered pinkroot            ten grains  
Calomel                            four grains  
Mix. To be taken two mornings in succession, and on the afternoon of the second day, followed by a purgative. For children above four years of age. *Ellis.*

#### PINKROOT WITH SAVINE.

R. Powdered pinkroot  
senna, each,            two scruples  
savine                    twelve grains  
Mix. One, every morning, for three days, followed by a purgative. *Ellis.*

#### INFUSION OF PINKROOT.

R. Pinkroot                    half an ounce  
Boiling water                one pint  
Macerate for two hours, in a covered vessel, and strain. *U. S. Ph.*

Four fl. drachms to a fl. ounce, to children; four to eight fl. ounces to adults; morning and evening, followed by purgative.

#### COMPOUND INFUSION OF PINKROOT.

R. Pinkroot                    half an ounce  
Senna                          two drachms  
Manna                          one ounce  
Fennel-seed            two drachms to half an ounce  
Boiling water                one pint  
Infuse. Half a wineglassful to a child, two years old, three times a-day. *Ellis.*

R. Pinkroot                    one ounce  
Bruised rhubarb            one drachm  
Senna                          two drachms  
Semen contra                one drachm  
Manna                          two drachms  
Coriander                    half a drachm  
Boiling water                one quart  
Infuse. A small teacupful, three times a-day. *Hays.*

#### SIMPLE FLUID EXTRACT OF PINKROOT.

R. Pinkroot                    sixteen ounces  
Diluted alcohol            sufficient  
Sugar                          twelve ounces  
Reduce the root to powder, macerate it in twice its weight of the diluted alcohol for twenty-four hours, then displace until four pints of tincture are obtained. Evaporate to ten fl. ounces, add the sugar, heat till dissolved: the whole to measure one pint. *W. Procter.*

#### COMPOUND FLUID EXTRACT OF SPIGELIA.

R. Bruised spigelia            four ounces  
Senna                          three ounces  
Savine                          one drachm  
Manna                          one ounce  
Sugar                          eight ounces  
Alcohol                        half a pint  
Boiling water                two pints

On the pinkroot, senna, and savine pour the boiling water, and cover till cool. Then add the alcohol, and let macerate for twenty-four hours; transfer to a displacement apparatus, and, having displaced, evaporate the product, at a low heat, to twelve fl. ounces, in which dissolve the manna and sugar, and evaporate to one pint. One fl. ounce contains the strength of two drachms of pinkroot and a drachm and a-half of senna. Dose, for a child, a teaspoonful. *T. Estlack.*

R. Pinkroot                    sixteen ounces (Av.)  
Senna                          eight ounces  
Powdered sugar            twenty-four ounces  
Carbonate of potassa        one ounce  
Oil of caraway  
Oil of anise, each,            half a drachm  
Diluted alcohol                sufficient

Reduce the pinkroot and senna to a coarse powder, mix them with two pints of diluted alcohol, and macerate for two days. Put in a displacer, gradually add more diluted alcohol, until five pints have passed. Evaporate this in a water-bath to twenty fl. ounces, add the carbonate of potassa. Triturate the oils with a portion of the sugar, then with the whole, mix this with the evaporated fluid, and apply a gentle heat. The whole should measure two pints.

Dose, from half a teaspoonful, to a child of one to two years of age, to a tablespoonful for an adult. *W. Procter.*

#### SPIRÆA.

##### HARDHACK.

The U. S. Pharm. indicates the root of *Spiræa tomentosa* as officinal; but this portion is less active than the leaves and flowers. It is a beautiful shrub, with spikes of rose-colored flowers, and leaves of a dark green above, and white beneath.

*Sex. Syst.* Icosand. pent. *Nat. Syst.* Rosaceæ.  
*Linn. Sp. Pl.* 701. *Griffith, Med. Bot.* 280.

The leaves and flowers are usually sold in packages; they have an odor not unlike that of black tea, and a bitter, very astringent taste.



The Hardhack is tonic and astringent, of considerable powers, and has the advantage of seldom being offensive to the stomach.

## DECOCTION OF HARDHACK.

R. Hardhack one ounce  
Water one pint and a-half  
Boil down to one pint. Dose, from one to two fl. ounces.

## EXTRACT OF HARDHACK.

R. Hardhack at will  
Water sufficient  
Exhaust by the process of displacement, evaporate the product by means of a water-bath to proper consistence. Dose, five grains to a scruple.

## SPONGIA.

## SPONGE.

Sponge is the horny skeletons of small polymorphous marine animals, found attached to submerged rocks in the seas of most warm climates. As found in the shops, it is in the form of a light, porous mass, of a yellowish brown color, and very elastic, and readily imbibing fluids.

It is used for the purpose of cleansing the surfaces of wounds and ulcers, and, in a compressed state, to form tents to dilate sinuses, &c. When burnt, or reduced to charcoal, it has been given successfully in some diseases; probably owing its efficacy to the presence of iodine.

## BURNT SPONGE.

R. Sponge at will  
Cut into pieces, separate any extraneous matters by beating, burn in a closed iron vessel till it becomes black and friable, then rub into fine powder. *U. S. Ph.*

Used in goitre, glandular swellings of a scrofulous nature, &c.; in doses of one to three drachms.

## POWDER OF BURNT SPONGE.

R. Burnt sponge ten grains  
Powdered rhubarb four grains  
Mix. To be taken morning and evening. In scrofula. *Hulse.*

## COMPOUND POWDERS OF BURNT SPONGE.

R. Burnt sponge six drachms  
Powdered cinnamon  
ginger, each, one drachm  
Sulphate of potassa two drachms  
Sugar five drachms  
Mix. *Greek Ph.*

Dose, a teaspoonful, three times a-day, in scrofula.

R. Burnt sponge four drachms  
Carbonate of magnesia  
Nitric

White sugar, each, two drachms  
Make powder. Dose, as above. *Clarus.*

## BOLUS OF BURNT SPONGE.

R. Burnt sponge one scruple  
Sulphate of potassa fifteen grains  
Balsam of sulphur ten drops  
Syrup sufficient  
Mix, and form two boluses. One, morning and evening, in scrofula. *Cadet.*

## ELECTUARY OF BURNT SPONGE.

R. Burnt sponge half an ounce  
Syrup of orange  
peel one ounce and a-half  
Mix. Two to four spoonfuls a-day, in goitre, &c. *Radius.*

## STANNUM.

## TIN.

## POWDER OF TIN.

R. Tin at will  
Melt in an iron vessel; while cooling, stir till reduced to powder, which is then to be sifted. *U. S. Ph.*

As an anthelmintic, in doses of half an ounce, mixed with syrup or molasses, for several mornings, followed by an active purgative.

## ELECTUARY OF TIN.

R. Powder of tin one ounce  
Extract of wormwood  
Powdered jalap, each, one drachm  
Compound syrup of chicory sufficient  
Mix. To be taken in twelve doses. *Foy.*

## OINTMENT OF TIN.

R. Amalgam of tin half an ounce  
Rose ointment one ounce  
Red precipitate two drachms  
Oil of peppermint twenty drops  
Rub well together. Said to be useful in hemorrhoids. *Brera.*

## STANNI CHLORIDUM.

## CHLORIDE OF TIN.

R. Tin at will  
Muriatic acid sufficient  
Dissolve, and crystallize. *Van Mons.*  
A violent purgative, in doses of two grains.

## STANNI SULPHURETUM.

## SULPHURET OF TIN.

R. Tin three parts  
Sulphur one part  
Heat in a crucible, as long as a flame



arises, cool, and pulverize. As a vermifuge, in doses of ten to twenty grains.

*Van Mons.*

#### COMPOUND POWDER OF SULPHURET OF TIN.

R. Sulphuret of tin                      four parts  
Powdered jalap                      two parts  
Assafetida                      one part  
Mix. Twelve to twenty grains, three times a-day, as a vermifuge. *Port. Ph.*

#### STANNI OXIDUM.

##### OXIDE OF TIN.

R. Tin    at will  
Keep it melted in an open vessel, constantly stirring till it is reduced into a gray powder, and sift. *Swed. Pharm.*

Has been recommended in tape-worm, in doses of five or six grains, several times a-day.

#### STAPHISAGRIA.

##### STAVESACRE.

The seeds of *Delphinium staphisagria*, a beautiful annual plant, with terminal racemes of blue flowers; a native of the south of Europe.

*Sex. Syst.* Polyand. trigyn. *Nat. Syst.* Ranunculaceæ.

*Linn. Sp. Pl.* 750. *Lindley, Med. Flor.* 9.

The seeds are large, somewhat triangular, of a brown color, a slight but unpleasant odor, and an acrid, bitter, nauseous taste. They are principally used to destroy lice in the hair, but also for other purposes.

##### DECOCTION OF STAVESACRE.

R. Stavesacre                                      one ounce  
Water                                      one pint and a-half  
Boil down to two-thirds, and strain. As a lotion in itch. *Swed. Pharm.*

##### VINEGAR OF STAVESACRE.

R. Stavesacre                                      one part  
Vinegar                                      sixteen parts  
Macerate and express. As a wash to destroy vermin in the hair. *Berol.*

##### OINTMENT OF STAVESACRE.

R. Stavesacre                                      two parts  
Lard                                      three parts  
Suet, each,                                      three parts  
Melt for some time, and strain. To destroy lice. *Bruns. Ph.*

#### STRAMONIUM.

##### STRAMONIUM.

Several parts of the *Datura stramonium* are used in medicine, as the root, leaves, and seeds. It is an annual herbaceous plant, a native of the

United States, having a rank, fetid odor, and bearing large, funnel-shaped, white flowers.

*Sex. Syst.* Pentand. monog. *Nat. Syst.* Solanaceæ.

*Linn. Sp. Pl.* 255. *Griffith, Med. Bot.* 490.

The leaves have a fetid, narcotic odor, and a bitter, nauseous taste; the odor is lost on drying, but the taste remains. The seeds are small, reniform, of a brownish-black color, inodorous, of a bitter, somewhat acrid taste. Stramonium is a powerful narcotic, and is much used in a variety of affections, both internally and externally. The dose of the powdered leaves is from two to three grains; of the seeds, about a grain.

##### PILLS OF STRAMONIUM SEEDS.

R. Powdered stramonium seeds ten grains  
camphor                      one drachm  
savine                      five scruples  
Extract of seneka                      four scruples  
Mix, and make pills of two grains. Dose, six, three times a-day. In rheumatism. *Vogt.*

##### COMPOUND STRAMONIUM PILLS.

R. Extract of stramonium                      one drachm  
Soap                      two drachms  
Powdered gum Arabic                      one scruple  
liquorice                      two scruples  
Mucilage of tragacanth                      sufficient  
Make mass, and divide into sixty pills. One, night and morning, in asthma. *Halford.*

##### EXTRACT OF STRAMONIUM SEEDS.

R. Powdered stramonium seeds one pound  
Diluted alcohol                      sufficient  
Rub the powder with half a pint of the alcohol, and introduce into a displacement apparatus, exhaust by means of alcohol. Distil the filtered liquor, and evaporate the residue to a proper consistence. *U. S. Ph.*

Dose, a quarter to half a grain twice a-day, gradually increasing.

##### EXTRACT OF STRAMONIUM LEAVES.

R. Stramonium leaves                      one pound  
Bruise in a stone mortar, with a little water. Express, and heat the juice to boiling; strain, and evaporate to proper consistence. *U. S. Ph.*

Dose, one grain, twice a-day, gradually increasing till it produces its effects.

##### MIXTURE OF EXTRACT OF SEEDS OF STRAMONIUM.

R. Extract of stramonium  
seeds                      twelve grains  
Antimonial wine                      half a fl. ounce  
Dissolve. Ten drops, every three hours, in an infusion of balm, as an antispasmodic. *Hufeland.*

##### ANTIRHEUMATIC LOTION.

R. Seeds of stramonium                      one ounce



Alcohol one pint  
 Infuse, strain, and add  
 Opium one ounce  
 Camphorated alcohol two fl. ounces  
 Mix. In frictions to the diseased part.  
*Pierquin.*

SYRUP OF STRAMONIUM.

R. Sugar fifteen ounces  
 Tincture of stramonium two ounces  
 Distilled water seven fl. ounces  
 Mix, and form syrup without heat, and strain.  
*Beral.*

R. Stramonium seeds one ounce  
 Vinegar one pound  
 Infuse for two days, strain, and add  
 Sugar two pounds  
 Dissolve.  
*Wurt. Ph.*

TINCTURE OF STRAMONIUM SEEDS.

R. Stramonium seeds, bruised, four ounces  
 Diluted alcohol two pints  
 Macerate for fourteen days, express, and filter.  
*U. S. Ph.*

Dose, from twenty to forty drops, two or three times a-day, increasing till it affects the system.

TINCTURE OF STRAMONIUM LEAVES.

R. Filtered juice of stramonium  
 Alcohol (.847) equal parts  
 Mix, and filter at the end of twenty-four hours.  
*Beral.*

ETHEREAL TINCTURE OF STRAMONIUM.

R. Stramonium seeds one part  
 Sulphuric ether four parts  
 Macerate for a week, and filter. Dose, two to three drops.  
*Soubeiran.*

WINE OF STRAMONIUM.

R. Stramonium seeds two ounces  
 Wine eight fl. ounces  
 Alcohol one ounce  
 Macerate for some days, and filter. Dose, six drops to a fl. scruple.  
*Van Mons.*

OINTMENT OF STRAMONIUM.

R. Fresh stramonium leaves one pound  
 Lard three pounds  
 Wax half a pound  
 Boil the leaves in the lard, till they become crisp, then strain through linen, afterwards add the wax, previously melted, and stir till cold.  
*U. S. Ph.*

A valuable anodyne application to painful hemorrhoids and tumors.

R. Extract of stramonium one drachm  
 Lard one ounce  
 Rub together.  
*Giordano.*

OIL OF STRAMONIUM.

R. Stramonium leaves one part

Olive oil two parts  
 Heat till moisture is driven off, express, and strain.  
*Cottureau.*

STYRAX,

STORAX,

Is the concrete juice of *Styrax officinale*, a small tree, a native of the warm parts of Europe, Syria, &c. It is procured by making incisions in the bark.

*Sex. Syst.* Decand. monog. *Nat. Syst.* *Styracaceæ.*

*Linn. Sp. Pl.* 635. *Griffith, Med. Bot.* 437.

Several kinds of Storax are employed, that in grains, in mass, in calamita, and the liquid. It has a fragrant odor, and warm, aromatic taste. It is a stimulating expectorant. Dose, ten to twenty grains, twice a-day.

PURIFIED STORAX.

R. Storax at will  
 Alcohol sufficient  
 Dissolve, and strain; distil off alcohol by gentle heat, till the storax is of proper consistence.  
*U. S. Ph.*

COMPOUND PILLS OF STORAX.

R. Purified storax three drachms  
 Powdered opium  
 Saffron, each, one drachm  
 Beat together. Five grains contain one of opium.  
*Lond. Ph.*

SYRUP OF STORAX.

R. Purified storax three ounces  
 Alcohol one fl. ounce  
 Water one pint  
 Digest for twelve hours on a water-bath, and add  
 Sugar two pounds  
 Orange-flower water one fl. ounce  
 Mix.  
*Giordano.*

STRYCHNIA.

STRYCHNINE.

R. Nux vomica, rasped, four pounds  
 Powdered lime six ounces  
 Muriatic acid three fl. ounces  
 Alcohol  
 Diluted sulphuric acid  
 Solution of ammonia  
 Purified animal charcoal  
 Water, each, sufficient

Digest the nux vomica in two gallons of water, acidulated with a fl. ounce of muriatic acid, for twenty-four hours; boil for two hours, express, and strain through linen. Boil residue twice successively, in the same quantity of acidulated water, and proceed as before. Mix the decoctions,



and evaporate to consistence of thin syrup; add lime, previously mixed with a pint of water; boil for ten minutes, often stirring. Pour mixture into a double linen bag, and wash well with water; press, dry, and powder the precipitate. Treat repeatedly with boiling alcohol, till it loses its bitterness; mix the solutions, and distil off the alcohol in a water-bath; mix the residue with water; apply heat; add sufficient diluted sulphuric acid to dissolve the strychnia; treat with animal charcoal at boiling temperature; filter, evaporate, and crystallize. Dissolve the crystals in water; add solution of ammonia to precipitate the strychnia. Dry on bibulous paper.

*U. S. Ph.*

R. Powdered nux vomica    nine pounds  
Water  
Sulphuric acid, each,    sufficient  
Powdered quicklime    ten ounces  
Alcohol    fifteen pints

Mix the nux vomica with sufficient water to form a thin paste, and keep at the temperature of 70 or 80°, until gas ceases to form; express, and boil the residue in several successive portions of water, and express. Set liquid aside to deposit; decant, and evaporate to three gallons. Add nine ounces of quicklime, and after six hours contact, express strongly; heat the liquid to the boiling point, and add a slight excess of sulphuric acid; separate the liquid by decantation, and evaporate to four pints; add one ounce of quicklime, and act as before, adding this precipitate to the former; dry, and powder; digest in five pints of alcohol diluted with five pints of water, at a gentle heat; separate precipitate, and boil in five pints of alcohol twice; mix, and filter solutions; distil off four-fifths, and set aside to crystallize.

*Molyn.*

Very active and dangerous. Dose, one-sixteenth to one-tenth of a grain, at first carefully watching the effects, and slowly increasing.

#### COMPOUND POWDER OF STRYCHNINE.

R. Strychnine    one grain  
Black oxide of iron  
Sugar  
Gum Arabic, each,    one drachm  
Rub well together, and divide into twelve or sixteen powders.

*Brera.*

#### PILLS OF STRYCHNINE.

R. Strychnine    two grains  
Conserve of roses    thirty-six grains  
Mix, and make twenty-four pills. One

to two, morning and evening, in paralysis.

*Magendie.*

R. Strychnine    two grains  
Extract of valerian    sufficient  
Mix well, and make thirty-two pills. One early in the morning, for five days, then one, morning and evening, in amaurosis.

*Furnari.*

#### STIMULANT PILLS.

R. Strychnia    one grain  
Acetic acid    one minim  
Crumb of bread    one scruple  
Mix thoroughly, and make ten pills. One to be taken every six hours, in the paralysis arising from lead.

*A. T. Thomson.*

#### TINCTURE OF STRYCHNINE.

R. Strychnine    three grains  
Alcohol (.842)    one fl. ounce  
Dissolve. Dose, from six to twenty-four drops, twice a-day.

*Magendie.*

#### MIXTURE OF STRYCHNINE.

R. Strychnine    one grain  
Distilled water    two fl. ounces  
White sugar    two drachms  
Acetic acid    three drops  
Mix. One fl. drachm contains one-sixteenth of a grain.

*Beasley.*

#### OINTMENT OF STRYCHNINE.

R. Strychnine    sixteen grains  
Lard    one ounce  
Rub well together. As a friction on paralyzed parts.

*Bouchardat.*

#### STRYCHNIA COLLYRIUM.

R. Strychnia    two grains  
Diluted acetic acid    one drachm  
Distilled water    one ounce  
Mix. A few drops of this applied to the eye, several times a-day, is stated to be beneficial in amaurosis.

*Henderson.*

#### LINIMENT OF STRYCHNINE.

R. Strychnine    thirty grains  
Olive oil    one ounce and a-half  
Rub well together. Ten drops to be rubbed on the temples, in amaurosis.

*Neligan.*

#### STRYCHNIAE ACETAS.

##### ACETATE OF STRYCHNINE.

R. Strychnine    at will  
Acetic acid    sufficient  
to dissolve; evaporate, and crystallize.

##### SOLUTION OF ACETATE OF STRYCHNINE.

R. Acetate of strychnine    three grains  
Alcohol    one fl. drachm



Cinnamon water seven fl. drachms  
Dissolve. Five drops twice a-day.

*Radiis.*

R. Strychnine one grain  
Distilled vinegar one fl. drachm  
water nine fl. drachms  
Mix. Dose, one fl. drachm containing one-tenth of a grain. *A. T. Thomson.*

#### TINCTURE OF ACETATE OF STRYCHNINE.

R. Acetate of strychnine one grain and a-half  
Alcohol half fl. ounce  
Dissolve. Twenty to thirty drops, twice a-day, in syphilitic pains in the bones.

*Radiis.*

#### STRYCHNINÆ MURIAS.

##### MURIATE OF STRYCHNINE.

R. Strychnine at will  
Muriatic acid sufficient  
to saturate; evaporate, and crystallize. *Cottureau.*

#### STRYCHNINÆ IODAS.

##### IODATE OF STRYCHNINE.

R. Strychnine at will  
Iodic acid sufficient  
to saturate; dissolve in boiling alcohol, filter, and crystallize. *Magendie.*

R. Solution of muriate of strychnine at will  
Solution of iodate of soda sufficient  
to precipitate; treat as the last. *Jourdan.*

Dose, one-eighth of a grain, in pill. One, morning, and evening, gradually increasing. In paraplegia.

#### STRYCHNINÆ NITRAS.

##### NITRATE OF STRYCHNINE.

R. Strychnine at will  
Nitric acid sufficient  
to saturate, with the aid of heat; filter while hot, evaporate, and crystallize. More active than strychnine. *Giordano.*

#### OINTMENT OF NITRATE OF STRYCHNINE.

R. Nitrate of strychnine one grain and a-half  
Lard two drachms  
Rub well together. *Wendt.*

#### STRYCHNINÆ SULPHAS.

##### SULPHATE OF STRYCHNINE.

R. Strychnine at will

Sulphuric acid sufficient  
to saturate at a gentle heat; filter, whilst hot, evaporate, and crystallize. *Cottureau.*

#### SUCCINUM,

##### AMBER,

Is a solid body, of various shades of yellow, translucent, or opaque, of a vitreous fracture, brittle, tasteless, inodorous, except when heated, then exhaling a peculiar, penetrating, but rather agreeable smell. It is a resin found in a fossil state, in various parts of the world. Not much used in medicine, but extensively employed in the arts.

##### FUMIGATING POWDER.

R. Amber  
Mastic  
Juniper berries, each, four parts  
Cascarilla one part  
Powder, and mix. *Scherf.*

##### OIL OF AMBER.

R. Powdered amber at will  
Mix, with equal weight of sand in a glass retort, to fill not more than one-half of it. Distil on a sand-bath, gradually increasing the heat. Separate the oil from the product, and keep in well-closed bottles. *U. S. Ph.*

##### PURIFIED OIL OF AMBER.

R. Oil of amber one pint  
Water six pints  
Mix in a glass retort; distil till four pints of water, with the oil, have passed over; separate the oil. *U. S. Ph.*

Stimulant and antispasmodic; in doses of five to fifteen drops; also used externally.

##### ARTIFICIAL MUSK.

R. Oil of amber one part  
Add gradually  
Nitric acid two parts  
Let react for twenty-four hours, and wash in cold water. *Van Mons.*

R. Strong nitric acid three drachms and a-half  
Add very gradually to  
Rectified oil of amber one drachm  
In a large, glass vessel. When action has ceased, permit to rest for twenty-four hours, remove the upper or resinous portion, and wash it well in cold and then in hot water, till all acid is removed. *Williams.*

Useful in those cases in which musk is applicable.

##### TINCTURE OF ARTIFICIAL MUSK.

R. Artificial musk one part  
Alcohol eight parts



Dissolve with a gentle heat. Forty drops as an antispasmodic. Useful in doses of five or six drops, in pertussis in children.

*Van Mons.*

R. Artificial musk two drachms  
Alcohol eight ounces

Dissolve, and filter. Dose, from twenty-five drops to a drachm, according to age.

*Williams.*

#### EMULSION OF ARTIFICIAL MUSK.

R. Artificial musk twelve grains  
Blanched almonds No. four  
Triturate well together, and gradually add  
Water six ounces

Dose, for a child of two years, two teaspoonfuls. Valuable in whooping-cough.

*Hufeland.*

#### LINIMENT OF OIL OF AMBER.

R. Oil of amber two fl. drachms  
olives half fl. ounce  
Laudanum two fl. drachms  
Brandy three fl. ounces

Mix. To be rubbed between the shoulders, in pertussis and infantile convulsions.

*Parrish.*

R. Oil of amber  
cloves, each, half an ounce  
olives one ounce

Mix. *Beasley.*

R. Oil of amber one drachm  
nutmeg two drachms and a-half

Mix. As a friction in hysteria, and cephalalgia. *Wurt. Ph.*

#### TINCTURE OF AMBER.

R. Powdered amber one ounce  
Diluted alcohol sixteen ounces

Digest for six days, and filter. *Par. Cod.*

Dose, forty to sixty drops.

#### ETHEREAL TINCTURE OF AMBER.

R. Oil of amber one part  
Sulphuric ether fifteen parts

Mix. Dose, fifteen to thirty drops. *Beral.*

#### ALKALINE TINCTURE OF AMBER.

R. Powdered amber two ounces  
Solution of carbonate of potassa half an ounce

Diluted alcohol eight fl. ounces  
Digest for some days, and filter. Dose, twenty to forty drops. *Spielman.*

#### BALSAM OF AMBER.

R. Essence of amber two fl. ounces  
myrrh one fl. ounce  
turpentine half fl. ounce

Mix. Digest at a gentle heat. As an antispasmodic friction. *Radius.*

R. Powdered amber one ounce  
Oil of turpentine two parts  
Digest at gentle heat. *Bate.*

#### EAU DE LUCE.

R. Oil of amber two drachms  
White soap

Balsam of Mecca, each, fifteen grains

Rectified spirit six ounces

Macerate for eight days, and filter. To one fl. drachm of this add two fl. ounces of water of ammonia. Used in the treatment of snake bites, as an antispasmodic, &c. *Redwood.*

#### MIXTURE OF OIL OF AMBER.

R. Oil of amber eighty drops  
Gum Arabic  
Sugar, each, half an ounce  
Tincture of tolu two fl. drachms  
Distilled water four fl. ounces

Mix. A tablespoonful, every two or three hours, in spasmodic cough. *Ellis.*

## SULPHUR.

### SULPHUR.

Three official forms of Sulphur are admitted in the U. S. Ph. Sulphur, or *Sublimed Sulphur*, *Washed Sulphur*, and *Precipitated Sulphur*.

### SUBLIMED SULPHUR.

R. Sulphur at will  
Heat at a temperature of 500° to 600°, in an earthen vessel, and condense the fumes in a large receiver.

### WASHED SULPHUR.

R. Sublimed sulphur at will  
Wash thoroughly with water, as long as any acidity is present.

### PRECIPITATED SULPHUR.

R. Sublimed sulphur one pound  
Lime one pound and a-half  
Water two gallons  
Muriatic acid sufficient

Slake the lime with a little water, mix it with the sulphur, add the rest of the water, boil for two or three hours, occasionally adding water, and filter. Dilute the liquid with an equal bulk of water, add sufficient muriatic acid to precipitate the sulphur. Wash the precipitate, till the washings are tasteless, and dry it.

### POWDER OF SULPHUR AND CAMPHOR.

R. Washed sulphur  
Sugar, each, half a drachm  
Camphor six grains  
Mix, and divide into six powders. One, every hour in mercurial cachexy. *Radius.*



## POWDER OF SULPHUR AND LIQUORICE.

R. Flowers of sulphur           one drachm  
       Powdered fennel seed       two drachms  
               liquorice           four drachms  
 Mix. A teaspoonful occasionally, in catarrh. *Phæbus.*

## POWDER OF SULPHUR AND ORRIS ROOT.

R. Washed sulphur           eight to eighteen grains  
       Sugar of milk           one scruple  
       Powdered orris root       one to three drachms  
 Mix, and divide into eight powders. One, every two hours, in milk, in the catarrh of children. *Kopp.*

## COMPOUND POWDER OF SULPHUR.

R. Flowers of sulphur           one ounce  
       Powdered fennel seed       one drachm  
               gum Arabic  
               sugar, each,       two ounces  
 Mix. Dose, half a drachm to a drachm, in dysentery. *Van Mons.*

R. Flowers of sulphur       half an ounce  
       Powdered liquorice       one ounce  
               orris root       two drachms  
               benzoin       one scruple  
               sugar       two ounces  
       Oil of fennel  
               anise, each,       ten drops

Mix. Dose, from a scruple to half a drachm. *Wert. Ph.*

R. Flowers of sulphur       two drachms  
       Cremor tartar       six drachms  
       Powdered fennel seed   one drachm  
               sugar       half an ounce

Mix. A spoonful two or three times a-day, as a laxative, in constipation. *Radius.*

R. Sublimed sulphur       one drachm and a-half  
       Brown sugar  
       Cremor tartar, each,   two drachms  
 Mix, and make powder. To be taken twice a-day, in itch, impetigo, or acne. *Ainslie.*

## POWDER OF SULPHUR AND CREMOR TARTAR.

R. Sublimed sulphur       half an ounce  
       Cremor tartar       one ounce  
 Mix with molasses. A teaspoonful, four or five times a-day, to children, in cutaneous affections. *Ellis.*

## POWDER OF SULPHUR AND MAGNESIA.

R. Precipitated sulphur  
       Magnesia, each,       half an ounce

Mix. A teaspoonful, four or five times a-day, as an aperient. *Ellis.*

## POWDER OF SULPHUR AND ANTIMONY.

R. Washed sulphur       two drachms  
       Golden sulph. of antimony  
       Camphor, each,       eight grains  
       Sugar       two scruples

Mix, and make a powder. *Hufeland.*  
 As a diaphoretic.

## SULPHUR ELECTUARY.

R. Washed sulphur  
       Cremor tartar, each,   half an ounce  
       Simple or lemon syrup   sufficient  
 Mix, and make electuary. To keep the bowels open in hemorrhoids of pregnant women. A teaspoonful, at night. *Dewees.*

R. Precipitated sulphur   one drachm and a-half  
       Orange marmalade       half an ounce  
       Pulp of tamarinds       two ounces  
       Sugar       one ounce

Mix. Three spoonfuls a-day, in hemorrhoids. *Radius.*

R. Washed sulphur one ounce and a-half  
       Confection of senna       two ounces  
       Nitre       one drachm  
       Syrup of orange peel   sufficient

Mix. One, or two drachms, once or twice a-day, in hemorrhoidal affections. *Ellis.*

R. Flowers of sulphur  
       Burnt sponge, each,   two drachms  
       Prepared oyster shell   one drachm  
       Conserve of roses       two ounces and a-half

Mix. A teaspoonful, three or four times a-day, as an alterative, in scrofulous affections. *Vogt.*

## COMPOUND ELECTUARY OF SULPHUR.

R. Sublimed sulphur       half an ounce  
       Cremor tartar       one drachm and a-half  
       Confection of senna       one ounce  
       Simple syrup       sufficient  
 Mix, and make electuary. A teaspoonful, at bedtime, for hemorrhoids. *Ainslie.*

## LINCTUS WITH SULPHUR.

R. Washed sulphur       half a drachm  
       Flowers of benzoin       fifteen grains  
       Syrup of violets       one ounce and a-half  
       Oxymel       one ounce

Mix. A spoonful, occasionally, as an expectorant. *Bories.*

## ELECTUARY OF SULPHUR.

R. Washed sulphur       half an ounce  
       Confection of senna   one ounce and a-half



Syrup sufficient  
 Mix. A teaspoonful, morning and evening, in hemorrhoids. *Lewis Disp.*

## SULPHUR MIXTURE.

R. Precipitated sulphur two drachms  
 Laudanum twenty drops  
 Water four fl. ounces

Mix. Three spoonfuls, a-day, in mercurial disease. *Radius.*

R. Precipitated sulphur two drachms  
 Fennel water three fl. ounces  
 Cinnamon water one fl. ounce  
 Syrup of opium half a fl. ounce

Mix. A spoonful every two hours, in diabetes mellitus. *Augustin.*

R. Washed sulphur one scruple  
 Mucilage seven drachms  
 Sugar half an ounce  
 Rose water one fl. drachm

A teaspoonful, every hour, in pectoral affections of young children. *Phæbus.*

## BALSAM OF SULPHUR.

R. Washed sulphur two ounces  
 Olive oil sixteen fl. ounces

Heat the oil, gradually stir in the sulphur till it combines. *Lond. Ph., 1824.*

Used as an external application to foul ulcers.

## ETHEREAL BALSAM OF SULPHUR.

R. Terebinthinated balsam of sulphur  
 Sulphuric ether, each, half a drachm  
 Dippel's animal oil six drachms

Mix. In flatulent colic. *Augustin.*

## TEREBINTHINATED BALSAM OF SULPHUR.

R. Flowers of sulphur one part  
 Oil of turpentine four parts  
 Dissolve on a sand-bath. As an external application to ulcers, &c. *Van Mons.*

## OINTMENT OF SULPHUR.

R. Flowers of sulphur one pound  
 Lard two pounds  
 Mix. *U. S. Ph.*

As an application in itch.

## COMPOUND SULPHUR OINTMENT.

R. Flowers of sulphur one ounce  
 Ammoniated mercury  
 Benzoic acid, each, one drachm  
 Oil of bergamot  
 Sulphuric acid, each, one fl. drachm  
 Nitrate of potassa two drachms  
 Lard half a pound

Melt the lard, add the other ingredients, and stir till cold. *U. S. Ph.*

As an application in itch, tinea capitis, crusta lactea, &c.

R. Sulphur half a pound

Powdered white hellebore two ounces  
 Nitrate of potassa one drachm  
 Soft soap half a pound  
 Lard one pound and a-half  
 Oil of bergamot thirty minims

Mix. *Lond. Ph.*

Used as the last, but more irritating.

R. Flowers of sulphur two parts  
 Carbonate of potassa one part  
 Lard eight parts

Mix. In itch. *Foy.*

R. Flowers of sulphur two ounces  
 Powdered muriate of ammonia two drachms  
 Oil of mint one drachm

Lard four ounces

Mix. In cases of inveterate itch. *Dewees.*

R. Washed sulphur ten ounces  
 Carbonate of potassa half an ounce  
 Cinnabar one ounce

Oil of bergamot half a fl. ounce  
 Lard ten ounces

Mix. In itch. *Bateman.*

## OINTMENT OF SULPHUR AND CAMPHOR.

R. Washed sulphur half a drachm  
 Camphor one scruple  
 Rose ointment one ounce  
 Oil of roses three drops

Mix. As an application in psorophthalmia. *Allen.*

## OINTMENT OF SULPHUR AND SOAP.

R. Flowers of sulphur one part  
 Soft soap two parts

Mix. In itch. *Radius.*

## OINTMENT OF SULPHUR AND ZINC.

R. Flowers of sulphur two ounces  
 Sulphate of zinc  
 Oil of bayberries, each, one ounce  
 Lard six ounces

Mix. In tinea capitis. *Sax. Ph.*

R. Flowers of sulphur four parts  
 Oxide of zinc three parts  
 Oil of bayberries six parts  
 Lard twelve parts

Mix. As last. *Van Mons.*

## LINIMENT OF SULPHUR AND SOAP.

R. Soap three ounces  
 Water six ounces

Dissolve with a gentle heat, and add  
 Flowers of sulphur three ounces

Used as a lotion in itch. *Lugol.*

## CERATE OF SULPHUR.

R. Flowers of sulphur two parts  
 Cold cream seven parts

Oil of almonds one part

Mix. *Par. Cod.*



SULPHURIS CARBURETUM,  
OR,  
CARBONIS SULPHURETUM.

## SULPHURET OF CARBON.

R. Sulphuret of iron eleven parts  
Charcoal three parts

Mix, and introduce into a stone retort, furnished with a glass tube, dipping into water. Separate the sulphuret which collects at the bottom of the water in the recipient, and redistil it from chloride of lime. *Van Mons.*

R. Sulphur at will  
Pass the vapor slowly through charcoal, heated to redness, in a porcelain tube, collect the product in a receiver, and purify by another distillation. *Cottureau.*

As a sudorific in rheumatism; two to three drops. Also used externally.

## DROPS OF SULPHURET OF CARBON.

R. Sulphuret of carbon one fl. drachm  
Alcohol half fl. ounce

Mix. Four to six drops, every two hours, in rheumatism. *Wutzer.*

## MIXTURE OF SULPHURET OF CARBON.

R. Sulphuret of carbon one scruple  
Cow's milk six fl. ounces  
Sugar two drachms

Mix. A tablespoonful four times a-day, in hypertrophy of the stomach and contraction of the œsophagus. *Clarus.*

## LINIMENT OF SULPHURET OF CARBON.

R. Sulphuret of carbon half an ounce  
Oil of almonds one ounce

Mix. As an embrocation in gouty nodes. *Mansfield.*

R. Sulphuret of carbon one drachm  
Camphorated oil one ounce  
*Beasley.*

R. Camphor two drachms  
Dissolve in  
Sulphuret of carbon half fl. ounce

And add

Alcohol one fl. ounce  
As an embrocation in rheumatism.

*Lampadius.*

R. Sulphuret of carbon two fl. drachms  
Camphorated ammon.

liniment two fl. ounces

Mix. As an embrocation, in rheumatism. *Wutzer.*

## SULPHURIS IODIDUM.

## IODIDE OF SULPHUR.

R. Iodine four ounces  
Sulphur one ounce

Rub the iodine and sulphur together, in a porcelain or glass mortar. Put the mixture into a matrass, close the orifice slightly, and apply a gentle heat, so as to darken the mass, but not melt it. When uniformly dark, increase the fire so as to melt the iodine, then incline the vessel in different directions, to return to the mass any portions that have been condensed on the surface of the vessel; lastly, allow to cool, break the matrass, and put the iodide into well-stopped bottles. *U. S. Ph.*

Used as an external application, in cutaneous affections.

## POWDER OF IODIDE OF SULPHUR.

R. Powdered iodide of sulphur ten grains  
Powdered gum Arabic sufficient

Mix, and divide into six powders. One, morning and evening, for an adult. Useful in porrigo, prurigo, impetigo, and tinea, also in nocturnal incontinence of urine. *Escobar.*

## OINTMENT OF IODIDE OF SULPHUR.

R. Iodide of sulphur five parts  
Lard ninety-six parts

Mix. *Foy.*

The strength is to be varied according to circumstances.

## T

## TABACUM.

## TOBACCO.

Tobacco is the leaves of *Nicotiana tabacum*, an annual, herbaceous plant, a native of the warmer parts of America, and extensively cultivated in the United States, and elsewhere. There are

numerous varieties, but, in all of them, the leaves are the part that is used.

*Ser. Syst.* Pentand. monog. *Nat. Syst.* Solanaceæ.

*Linn. Sp. Pl.* 258. *Griffith, Med. Bot.* 493.

Tobacco, as found in commerce, is in bundles of yellowish brown, dried leaves, having a narcotic,



penetrating odor, and a bitterish, nauseous, acrid taste. It has the powers of a sedative narcotic, with those of an emetic and diuretic; it also acts as an errhine and sialogogue. In large doses it acts as a poison. It is employed to produce relaxation in spasmodic affections, as a diuretic, &c.

#### COMPOUND POWDER OF TOBACCO.

R. Powdered tobacco  
valerian, each, two drachms  
Oil of lavender  
marjoram, each, three drops  
Mix. Used as a cephalic snuff. *Boeli.*  
R. Powdered tobacco two grains  
Tartar emetic one grain  
Powdered sugar two drachms  
gum Arabic half a drachm  
Mix, and form twenty powders. One, every two hours, in whooping-cough. *Petschaft.*

#### EXTRACT OF TOBACCO.

R. Cut tobacco four ounces  
Water two pints  
Boil, and let simmer for two or three hours, strain, and evaporate to consistence of extract. For external use in neuralgia. *Chippendale.*

#### PILLS OF TOBACCO.

R. Extract of tobacco one drachm  
liquorice one ounce  
Vinegar of squills sufficient  
Mix, and make one hundred and eighty pills. Dose, one to three in dropsy. *Van Mons.*

#### WINE OF TOBACCO.

R. Tobacco, cut, one ounce  
Wine one pint  
Macerate for fourteen days, express, and filter. *U. S. Ph.*

From ten to twenty minims, as a diuretic.

#### TINCTURE OF TOBACCO.

R. Cut tobacco one ounce  
Diluted alcohol one pint  
Digest for three days, express, and filter. Dose, as the wine. *Augustin.*

#### INFUSION OF TOBACCO.

R. Tobacco one drachm  
Boiling water one pint  
Macerate for an hour, in a covered vessel, and strain. *U. S. Ph.*

Used as an enema; one-half only should be used at a time; employed in strangulated hernia, obstinate colic, &c. Great caution must be used, as dangerous effects have followed its administration.

#### MIXTURE OF TOBACCO.

R. Tobacco one drachm  
Boiling water two ounces  
Infuse for twenty minutes, filter, and add

Alcohol two fl. drachms  
Thirty to fifty drops, twice or thrice a-day, in hydrothorax and dysuria. *Fowler.*

#### LOTION OF TOBACCO.

R. Tobacco two drachms to half an ounce  
Boiling water one pint  
Infuse. As a lotion in psora, but must be used with great caution. *Ellis.*

#### CATAPLASM OF TOBACCO.

R. Tobacco one ounce  
Beat up with water, and form cataplasm. To be applied to throat in croup, and spasm of glottis, &c. *Ellis.*

#### OINTMENT OF TOBACCO.

R. Fresh tobacco, cut, one ounce  
Lard one pound  
Boil over a gentle fire, till the leaves become friable, strain through linen. *U. S. Ph.*

In irritable ulcers, tinea capitis, &c. To be used with caution.

R. Powdered tobacco  
Flowers of sulphur, each, four ounces  
Powdered white hellebore two ounces  
Oleander leaves one ounce  
Common salt ten drachms  
Rose ointment two pounds  
Mix. As an application in psora, &c. *Taddei.*

R. Extract of tobacco one drachm  
Simple cerate one ounce  
Mix. As a friction in neuralgia. *Chippendale.*

#### OIL OF TOBACCO.

R. Kentucky tobacco at will  
Introduce it into an earthen or iron retort, connected with a receiver, which should be furnished with a tube, to conduct the incondensable gases to a chimney; heat the retort to the temperature of 600° Fahr., or thereabout, until the empyreumatic oil ceases to distil; separate the black tarry from the acid liquor in the receiver, and preserve it for use.

#### NICOTINA.

R. Kentucky tobacco at will  
Cut in pieces, and extract it with water; evaporate to an extract, exhaust this with alcohol; distil off the alcohol, treat the soft extract with an excess of potash in water; agitate this with ether, in separate portions, till the impure nicotine is removed; unite the ethereal liquid, add an excess of pulverized oxalic acid, wash the oxalate of nicotine which precipitates with ether, then treat it with an aqueous solution of potash,



and again agitate the liquid with ether to remove the alkaloid, which is obtained by distilling off the ether in the form of a light brown, syrupy liquid, and consists of nicotine, ether, water, and some ammonia. The last three bodies may be removed by heating the nicotine in a retort to the temperature of 284° F., in a current of hydrogen, for twelve hours; then, by raising the heat to 356° F., the nicotine distils over pure. *Schloessing.*

## TAMARINDUS.

## TAMARINDS.

Tamarinds are the preserved fruit of the *Tamarindus Indica*, a large tree, a native of the East Indies, and extensively cultivated in the tropical parts of America. Those brought here, are principally derived from the West Indies.

*Sex. Syst.* Diadelph. triand. *Nat. Syst.* Fabaceæ.

*Linn. Sp. Pl.* 48. *Griffith, Med. Bot.* 262.

The preserved pods, as they come to us, are in a dark-colored adhesive mass, formed of pulp, fragments of the pods, seeds, and syrup, of a sweet acidulous taste. They are cooling and laxative; and, when mixed with water, form a grateful drink in febrile diseases.

## PULP OF TAMARINDS.

*R.* Tamarinds at will  
Digest in a small quantity of water, then pass through a sieve. *U. S. Ph.*

## ELECTUARY OF TAMARINDS.

*R.* Pulp of tamarinds one ounce and a-half  
Cremor tartar half an ounce  
Syrup of raspberries sufficient  
Mix. Two teaspoonfuls morning and evening, as a laxative. *Radius.*

## INFUSION OF TAMARINDS.

*R.* Pulp of tamarinds one ounce  
Boiling water two pints  
Infuse for one hour, and strain. As a cooling drink in fevers. *Cotteneau.*

## TAMARIND WHEY.

*R.* Milk three pints  
Tamarinds one ounce  
Boil, and strain. As a cooling drink. *Foy.*

## TANACETUM.

## TANSY.

Tansy or *Tanacetum vulgare*, is a perennial herbaceous plant, with numerous yellow flowers; a native of Europe, and generally cultivated in our gardens, and has also become naturalized in some places.

*Sex. Syst.* Syngen. super. *Nat. Syst.* Asteraceæ.  
*Linn. Sp. Pl.* 1184. *Griffith, Med. Bot.* 406.

The whole plant is officinal; it has a peculiar, and strong odor, diminished by drying; and a warm, bitter, aromatic taste. It is an aromatic bitter, and has been praised in intermittents, hysteria, amenorrhœa, as an anthelmintic, &c. As a vermifuge, the seeds are to be preferred.

## INFUSION OF TANSY.

*R.* Fresh tansy one ounce  
Boiling water one pint  
Infuse, and strain. Internally, and as an injection against ascarides. *Niemann.*

## EXTRACT OF TANSY.

*R.* Tansy  
Alcohol, each, one pound  
Water eight pints  
Digest for three days, express, distil off the alcohol, and evaporate to proper consistence. Dose, six to twenty grains. *Giordano.*

## OIL OF TANSY.

*R.* Tansy at will  
Water sufficient  
Distil, and separate the oil. *Guibourt.*

## PILLS OF TANSY.

*R.* Oil of tansy one scruple  
Extract of English walnut two drachms  
Powdered marsh mallow sufficient  
Mix, and make sixty pills. Three to five every two hours, as a vermifuge. *Radius.*

## COMPOUND TINCTURE OF TANSY.

*R.* Tansy one ounce  
Wormwood  
Rhubarb, each, three ounces  
Sherry wine two fl. ounces  
Diluted alcohol twenty fl. ounces  
Digest for eight days, and filter. Dose, one to two fl. drachms, two or three times a-day, as a vermifuge. *Van Mons.*

## TAPIOCA.

## TAPIOCA

Is a fecula obtained from the root of *Janipha manihot*, a shrub-like, herbaceous plant, a native of the tropical parts of America, where it is largely cultivated under the name of Cassava, and forms an important article of food.

*Sex. Syst.* Monœc. monadelph. *Nat. Syst.* Euphorbiaceæ.

*Kunth. ii.* 85. *Griffith, Med. Bot.* 601.

Tapioca is found in the shops, in the form of irregular, rough, white grains, having little odor or taste, soluble in hot water, and affording a bland and nutritious diet for the sick and convalescent.

## TAPIOCA JELLY.

*R.* Tapioca two spoonfuls  
Water one pint



Boil gently for an hour, or till it becomes gelatinous; flavor with sugar, wine, &c., according to circumstances. *Ellis.*

#### TAPIOCA PUDDING.

R. Yolks of eggs two  
Sugar half an ounce  
Beat together, and stir the mixture with  
Tapioca mucilage one pint  
Bake in a slow oven. The mucilage should be made with milk, not water.

*A. T. Thomson.*

### TARAXACUM.

#### DANDELION.

This officinal article is the root of *Taraxacum dens lionis*, a small, herbaceous plant, with a perennial, fusiform root. It is a native of Europe, but has become abundantly naturalized in this country.

*Sex. Syst.* Syngen. equal. *Nat. Syst.* Asteraceæ. *Haller*, i. 23. *Griffith*, *Med. Bot.* 414.

The root is fusiform, of a light brown color externally, succulent, inodorous, and of a mucilaginous, bitterish taste. It is aperient, diuretic, and somewhat tonic, and thought to be resolvent in engorgements of the liver.

#### DECOCTION OF DANDELION.

R. Bruised dandelion root two ounces  
Water two pints  
Boil down to a pint, and strain. *U. S. Ph.*

A wineglassful, two or three times a-day, as a diuretic, &c.

#### COMPOUND INFUSION OF DANDELION.

R. Infusion of dandelion four fl. ounces  
Extract of dandelion two drachms  
Carbonate of soda half a drachm  
Tartrate of potassa three drachms  
Tincture of rhubarb three fl. drachms  
henbane twenty drops

Mix. One-third part, three times a-day, in dropsical and visceral affections.

*Meigs.*

#### FLUID EXTRACT OF DANDELION.

R. Dandelion root, fresh, thirty-two ounces  
Slice it, and reduce to a pulp. Mix this with one-sixth its bulk of alcohol, macerate for twenty-four hours, and express strongly. Add a pint of water containing a little alcohol, and again express. Evaporate the mixed products to twelve fl. ounces, add four fl. ounces of alcohol, and filter.

A teaspoonful is equal to half a drachm of the extract obtained from the expressed juice. It may be made from sixteen ounces of the dried root, reduced to a coarse powder, macerated in two pints of

water mixed with half a pint of alcohol, for two days, expressed, &c., as above.

*W. Procter.*

#### EXTRACT OF DANDELION.

R. Root of dandelion, bruised, one pound  
Water one gallon  
Boil down to four pints, express, and strain, then evaporate to proper consistence. *U. S. Ph.*

Dose, a scruple to a drachm, three times a-day, alone, or dissolved in cinnamon or mint water.

R. Bruise the recent root collected in September; add one-fourth its bulk of alcohol; allow the whole to macerate twelve hours, express powerfully, add a little dilute alcohol, and again express. Evaporate the mixed liquids in a water-bath to the proper consistence. *W. Procter.*

#### PILLS OF EXTRACT OF DANDELION.

R. Extract of dandelion half a drachm  
Powdered liquorice sufficient  
Mix, and make eight pills. One, three times a-day. In dropsy, and diseases of the urinary apparatus. *Ellis.*

R. Extract of dandelion  
Soap, each, one ounce  
Liquid acetate of potassa sufficient  
Mix, and make pills of four grains. Four to six in a day, as diuretic, &c. *St. Marie.*

#### PILLS OF DANDELION AND BLUE MASS.

R. Extract of dandelion half a drachm  
Blue pill five to ten grains  
Powdered uva ursi sufficient  
Mix, and make ten pills. One, thrice a-day. In dropsy connected with liver disease. *Ellis.*

#### MIXTURE OF DANDELION.

R. Extract of dandelion two ounces  
Peppermint water six fl. ounces  
Clarified honey one ounce  
Mix. Two spoonfuls, every three hours. *Richter.*

R. Extract of dandelion half a drachm  
Decoction of dandelion nine fl. drachms  
Spirit of nitric ether one fl. drachm  
Syrup of ginger two fl. drachms  
Mix. To be taken three times a-day, as a diuretic in anasarca and ascites. *Sprague.*

R. Cremor tartar two drachms  
Bruised cloves two scruples  
Sugar one ounce  
Decoction of dandelion one pound  
Macerate for two hours, and strain. One-fourth to be given every six hours, in dropsy. *Sprague.*



## CLYSTER OF DANDELION.

R. Dandelion three ounces  
 Bran one ounce  
 Water sixteen fl. ounces  
 Boil down to eight fl. ounces; add, at close,  
 Chamomile  
 Valerian, each, two drachms  
 Strain, for two injections. Much praised  
 in obstructions of the bowels. *Berands.*

## TEREBINTHINA.

## TURPENTINE.

Turpentine is the juice of several species of the Pine tribe, and is also procured from trees of other orders. That used in the United States is from several species of *Pinus* and *Abies*. All the turpentines are possessed of much the same properties; they have an aromatic odor, and a somewhat pungent and bitterish taste. They are stimulant, diuretic, anthelmintic, and, in large doses, purgative. They are also used externally as rubefacients.

## TURPENTINE PILLS.

R. Turpentine one drachm  
 Divide into fifteen pills. One, four or five times a-day, in affections of the urinary organs. *Ellis.*

## PILLS OF TURPENTINE AND GUAIAIACUM.

R. Powdered guaiacum one drachm  
 Turpentine sufficient  
 Mix, and make fifteen pills. One, thrice a-day; in gleet and leucorrhœa. They sometimes cause strangury. *Ellis.*

## PILLS OF TURPENTINE AND MAGNESIA.

R. Turpentine fourteen drachms  
 Calcined magnesia thirty-six grains  
 Mix, and make pills of six grains, at the end of twelve hours. In diseases of the urinary organs. *Monchon.*

## PILLS OF TURPENTINE AND RHUBARB.

R. Turpentine two ounces  
 Extract of rhubarb three drachms  
 Camphor two drachms  
 Mix, and make pills of four grains. Dose, three, thrice a-day, in leucorrhœa and gonorrhœa. *St. Marie.*

## PILLS OF TURPENTINE AND MYRRH.

R. Turpentine  
 Spermaceti half an ounce  
 Powdered myrrh two drachms  
 olibanum sufficient  
 Mix, and make pills of three grains. One every three hours, in chronic catarrh. *Cadet.*

## PILLS OF TURPENTINE AND JALAP.

R. Turpentine one drachm

Soap of jalap half a drachm  
 Extract of henbane four grains  
 Calomel eight grains  
 Make pills of twelve grains. Dose, four every three hours, for two days. As an anthelmintic. *Augustin.*

## ELECTUARY OF TURPENTINE.

R. Turpentine  
 Soap  
 Powdered rhubarb, each, one drachm  
 Syrup sufficient  
 Mix. Three teaspoonfuls a-day, in dropsy. *Radius.*

## TURPENTINE CLYSTER.

R. Venice turpentine half a fl. ounce  
 Yolk of egg one  
 Infusion of flaxseed ten fl. ounces  
 Rapeseed oil one fl. ounce  
 Mix. In colic, and obstinate constipation. *Ph. Noscom. Ed.*

## LINIMENT OF TURPENTINE.

R. Yellow wax one part  
 Melt, and add  
 Turpentine  
 Oil of turpentine, each, one part  
 Mix. Much praised in chronic ulcers on the legs. *Van Mons.*

R. Yellow wax half a pound  
 Olive oil one pound  
 Red saunders two ounces  
 Turpentine one pound  
 Melt together, strain, and add  
 Camphor two drachms

As an application to contusions and ulcerations. *Van Mons.*

R. Spirit of turpentine  
 Olive oil, each, one ounce and a-half  
 Dilute sulphuric acid three drachms  
 Mix. As an embrocation in rheumatism. *Chapman.*

## OINTMENT OF TURPENTINE.

R. Turpentine two pounds  
 Simple ointment nine ounces  
 Melt with a gentle heat, constantly stirring. *Austrian Ph.*

## COMPOUND OINTMENT OF TURPENTINE.

R. Mucilage one pound  
 Fresh butter two pounds  
 Olive oil three pounds and a-half  
 Melt together, and add  
 Turpentine three ounces  
 Mix well. As an application in engorgement of the breasts, in nephritic pains, &c. *Swediaur.*

## TURPENTINE PLASTER.

R. Turpentine one ounce



White of egg  
Wheat flour  
Sugar  
Honey  
Brandy

Olive oil, each, one ounce

Mix the turpentine and honey, gradually adding the other ingredients, rubbing well. As an application to painful ulcers.

*St. Marie.*

#### COMPOUND TURPENTINE PLASTER.

R. Turpentine eight parts  
Suet twenty-four parts  
Yellow wax thirty-two parts  
Essence of mint  
cloves, each, one part  
Oil of mace six parts  
Powder benzoin eight parts  
olibanum sixteen parts

Mix, with the assistance of heat. Useful in diarrhœa, dyspepsia, flatulence, colic.

*Ph. Sued.*

#### TURPENTINE MIXTURE.

R. Turpentine two drachms  
Yolk of egg one

Rub together, and add  
Extract of rhubarb

Sal prunelle, each, one drachm

Add gradually

Infusion of liquorice ten fl. ounces

Syrup of marsh mallows one fl. ounce

Mix. A spoonful every hour, in gonorrhœa.

*Bories.*

#### TURPENTINE MIXTURE.

R. Turpentine one ounce  
Mint water four fl. ounces  
Gum Arabic sufficient  
Simple syrup one ounce  
Extract of belladonna one grain

Mix, and make emulsion. Used in gonorrhœa, after the reduction of the inflammatory stage; when it acts too powerfully on the bowels, it must be intermitted for a few days.

*Ebriart.*

R. Spirit of turpentine one fl. ounce  
Oil of mint one fl. drachm

Mix. In flatulent colic, and gout in the stomach. Dose, a teaspoonful.

*Dewees.*

### TILIA EUROPÆA.

#### LINDEN.

The linden is a handsome tree, a native of the north of Europe, and much cultivated in the United States for ornament and shade.

*Sex. Syst.* Polyand. monog. *Nat. Syst.* Tiliaceæ.

Linn. Sp. Pl. 733. Lindley, Flor. Med. 147.

The part used is the flowers, which, when dried, have a faint but agreeable odor, and a sweetish, mucilaginous taste. They are antispasmodic, and diaphoretic, and are much used in France. It is probable that the flowers of our native species have the same properties.

#### INFUSION OF LINDEN FLOWERS.

R. Linden flowers two drachms  
Boiling water two pints  
Infuse for half an hour, and strain. A mild antispasmodic.

*Par. Cod.*

#### COMPOUND INFUSION OF LINDEN FLOWERS.

R. Linden flowers  
Chamomile  
Orange leaves, each, two drachms  
Boiling water two pints  
Infuse for half an hour, strain, and add  
Syrup two fl. ounces  
A small cupful, occasionally.

*Foy.*

### TOLUTANUM.

#### BALSAM OF TOLU.

This is the concrete balsamic juice of *Myrospermum toluiferum*, a large tree, of which little is known, growing in several parts of South America.

*Sex. Syst.* Diadelph. decand. *Nat. Syst.* Fabaceæ.

Decandolle, Prod. ii. 95. Griffith, Med. Bot. 250.

At first, the balsam is soft and tenacious, but gradually becomes hard and brittle, when it is transparent, shining, of a yellowish, or reddish brown color, of a fragrant odor, and a sweetish, warm taste. It is a stimulating expectorant. The dose is from ten to thirty grains.

#### LOZENGES OF BALSAM OF TOLU.

R. Balsam of Tolu one ounce  
Alcohol one fl. ounce

Dissolve, and add

Water two fl. ounces

Heat in a water-bath, and filter; add,

Tragacanth four scruples

Sugar sixteen ounces

Beat into paste, and make lozenges of twelve grains.

*Par. Cod.*

#### TINCTURE OF BALSAM OF TOLU.

R. Balsam of Tolu three ounces  
Alcohol two pints

Macerate till dissolved, then filter.

*U. S. Ph.*

A highly stimulating expectorant. Dose, one to two fl. drachms.

#### COMPOUND TINCTURE OF BALSAM OF TOLU.

R. Balsam of Tolu two ounces



Balsam of Peru one ounce  
 Flowers of benzoin  
 Saffron, each, half an ounce  
 Alcohol twenty-four fl. ounces  
 Digest for three days, and filter.

*Spielman.*

#### TINCTURE OF TOLU WITH FOXGLOVE.

R. Tincture of balsam  
 of Tolu one fl. ounce and a-half  
 Elixir of vitriol half fl. drachm  
 Tincture of foxglove one fl. drachm  
 Antimonial wine two fl. drachms  
 Clarified honey one fl. ounce and a-half  
 Powdered liquorice half a drachm  
 Distilled water six fl. ounces  
 Mix. A tablespoonful, according to circumstances.

*Ellis.*

#### EMULSION OF BALSAM OF TOLU.

R. Balsam Tolu three drachms  
 Mecca eight drops  
 Sweet almonds half an ounce  
 Barley water eighteen fl. ounces  
 Sugar six drachms  
 Make an emulsion. A spoonful occasionally, in chronic catarrh, &c.

*Spielman.*

#### SYRUP OF BALSAM OF TOLU.

R. Tincture of Tolu one fl. ounce  
 Syrup one pint and a-half  
 Mix, and evaporate on water-bath to the proper consistence.

*U. S. Ph.*

R. Balsam of Tolu ten drachms  
 Boiling water one pint  
 Sugar two pounds and a-half  
 Boil the balsam in the water for half an hour, occasionally stirring; strain when cold, add the sugar, and form syrup.

*Lond. Ph.*

#### MIXTURE OF TOLU AND MORPHIA.

R. Oxymel of squill one ounce and a-half  
 Syrup of Tolu half an ounce  
 Acetate of morphia half to one grain  
 Mix. A teaspoonful, as occasion may require, in catarrh.

*Ellis.*

#### MIXTURE OF TOLU AND BELLADONNA.

R. Syrup of Tolu three fl. ounces and a-half  
 seneka half a fl. ounce  
 Extract of belladonna eight to twelve grains  
 Wine of ipecacuanha one fl. drachm  
 Mix. A teaspoonful, three or four times a-day, in cough.

*Ellis.*

#### MIXTURE OF TOLU AND ALMOND

##### EMULSION.

R. Almond emulsion fifteen parts  
 Syrup of Tolu one part  
 Mix. In cough.

*Beral.*

#### MIXTURE OF TOLU AND COPAIBA.

R. Balsam of Tolu  
 copaiba  
 Powdered gum  
 Arabic, each, half an ounce  
 Elixir of vitriol twenty drops  
 Distilled water six fl. ounces  
 Mix. A tablespoonful occasionally, in chronic hooping-cough.

*Ellis.*

#### TOLU MIXTURE.

R. Tincture of Tolu one ounce  
 Laudanum two drachms  
 Tincture of foxglove one drachm  
 Mix. In the chronic cough of hæmoptysis. Forty to fifty drops, every three or four hours.

*Dewees.*

#### MIXTURE OF TOLU AND OPIUM.

R. Balsam of Tolu one drachm  
 Powdered gum  
 Arabic one drachm and a-half  
 Water four fl. ounces  
 Syrup of opium half a fl. ounce  
 Make an emulsion.

*Niemann.*

#### BALSAM OF TOLU INHALATION.

R. Balsam of Tolu one ounce  
 Boiling water one pint  
 Mix. The vapor to be inhaled.

*Ellis.*

## TORMENTILLA.

### TORMENTIL.

The root of *Potentilla tormentilla*, a small, perennial plant, with a large, woody, dark brown root, a native of many parts of Europe, growing in barren spots.

*Sex. Syst.* Icosand. polyg. *Nat. Syst.* Rosaceæ.

Sibthorp, Fl. Ox. 162. Griffith, Med. Bot. 278.

The root, when dried, is in irregular pieces, which are knotty, tuberculated, and of a dark reddish-brown color. Its odor is slightly aromatic, and its taste very astringent. It is a simple and somewhat active astringent. Dose, thirty grains to a drachm.

#### COMPOUND POWDER OF TORMENTIL.

R. Powdered tormentil  
 gum Arabic,  
 each, three ounces  
 Armenian bole six ounces  
 Powdered cinnamon four ounces  
 long pepper half an ounce  
 Mix.

*Geneva Ph.*

#### EXTRACT OF TORMENTIL.

R. Tormentil one part  
 Water eight parts  
 Boil, and strain; repeat with same quantity of water, mix the two decoctions, strain, and evaporate to the proper con-



sistence. Dose, from twenty to thirty grains. *Amst. Ph.*

#### DECOCTION OF TORMENTIL.

R. Tormentil, bruised, two ounces  
Distilled water one pint and a-half  
Boil down to a pint, and strain.

*Lond. Ph.*

Dose, one to two fl. ounces, three or four times a day.

#### GARGLE OF TORMENTIL.

R. Tormentil one ounce  
Water twelve fl. ounces  
Boil down one-third, strain, and add  
Alum one drachm  
Honey one ounce  
Mix. *Radius.*

### TOXICODENDRON.

#### POISON OAK.

The leaves of *Rhus toxicodendron*, a shrubby plant, native of the United States; sometimes becoming a vine, when it has received the name of *Rhus radicans*, or Poison vine. They both grow in woods, along fence rows, &c., and are extremely poisonous to some persons, whilst other individuals are not affected by them.

*Sex. Syst.* Pentand. trigyn. *Nat. Syst.* Anacardiaceæ.

*Linn. Sp. Pl.* 381. *Griffith, Med. Bot.* 184.

The officinal portion is the leaves, which are inodorous, and of a disagreeable, acrid taste. They are narcotic and stimulant, acting like the acro narcotic poisons in large doses. They have been used with success in obstinate cutaneous affections, chronic rheumatism, &c. The dose is half a grain, gradually increased.

#### EXTRACT OF POISON OAK.

R. Leaves of poison oak one part  
Alcohol four parts  
Infuse, strain, distil off the spirit, and evaporate to proper consistence.

*Par. Cod.*

Dose, one grain, gradually increased till some effect is induced.

#### PILLS OF POISON OAK.

R. Extract of poison oak  
Camphor, each, fifteen grains  
Extract of arnica  
valerian, each, thirty grains  
Powdered calamus sufficient  
Beat into mass, and make thirty pills.  
One, three times a-day, in hemiplegia.

*Brera.*

#### TINCTURE OF POISON OAK.

R. Clarified juice of poison oak  
Alcohol equal parts  
Mix, and filter at end of twenty-four hours.  
Dose, five drops, gradually increased.

### TRAGACANTHA.

#### TRAGACANTH.

A gummy exudation from *Astragalus verus*, and other species of *Astragalus*. All of them are small shrubs, with thorny petioles, found in the countries bordering on the Levant, in Persia, &c.

*Sex. Syst.* Diadelph. decand. *Nat. Syst.* Fabaceæ.

*Lindley, Flor. Med.* 247. *Griffith, Med. Bot.* 239.

Tragacanth is in contorted, vermicular pieces, rounded or flattened, of a whitish, or whitish yellow color; semi-translucent and corneous; tough, except when quite dry. It is inodorous, and of bland, mucilaginous taste. It forms a paste or mixture with water, but does not dissolve. It is demulcent and nutritive.

#### COMPOUND TRAGACANTH POWDER.

R. Powdered tragacanth  
gum Arabic  
starch, each, one ounce and  
a-half

Sugar three ounces  
Mix well. *Lond. Ph.*

Demulcent. Dose, from thirty grains to a drachm.

#### MUCILAGE OF TRAGACANTH.

R. Tragacanth one ounce  
Boiling water one pint  
Macerate for twenty-four hours, occasionally stirring, triturate till uniform, and strain forcibly through linen. *U. S. Ph.*

Principally used as a basis for more active medicines.

#### PASTE OF TRAGACANTH.

R. Tragacanth thirty-two parts  
Macerate in  
Water one thousand parts  
Also dissolve  
Isinglass forty-eight parts  
In

Water eight hundred parts  
Strain through a linen cloth, and mix with the macerated gum, and add

Syrup one thousand parts  
Evaporate by a gentle fire, to the consistence of soft paste, and gradually add

Orange-flower water sixty-four parts  
Evaporate to proper consistence on a water-bath. As expectorant. *Foy.*

R. White tragacanth sixty-four parts  
Isinglass ninety-six parts  
Water three thousand parts  
Simple syrup two thousand parts  
Orange-flower water  
one hundred and twenty-eight parts  
Bruise the tragacanth, and place it in a vessel for two days with five-sixths of the



water, and dissolve the isinglass in the rest of the water; strain both solutions; heat the syrup to boiling, and mix the whole, and continue heat, till the mixture is of the consistence of soft paste, stirring well. Remove from fire; add orange-flower water, and evaporate on water-bath, till sufficiently firm; pour on marble slab, and divide into lozenges. *Mouchon.*

## TUSSILAGO.

## COLTSFOOT.

Coltsfoot is a small, perennial plant, with large, radical, cordate leaves, and flowers on simple, leafless scapes. It is the *Tussilago farfara*, and is a native both of Europe and North America.

*Sex. Syst.* Syngen. super. *Nat. Syst.* Asteraceæ.

*Linn. Sp. Pl.* 1214. *Griffith, Med. Bot.* 393.

The whole herb is used, but the leaves and flowers should be preferred. The latter have an agreeable smell, the former are inodorous, but have a rough, bitterish, mucilaginous taste. They

both are demulcent, and have been much used in pectoral affections.

## DECOCTION OF COLTSFOOT.

R. Flowers of coltsfoot           one ounce  
Water                               two pints  
Boil down to a pint, and strain. *Beasley.*

## SYRUP OF COLTSFOOT.

R. Flowers of coltsfoot           one pound  
Boiling water                   two pints  
Macerate for twelve hours, express, strain, and add  
Sugar                               four pounds  
Make syrup. *Par. Cod.*

## COMPOUND DECOCTION OF COLTSFOOT.

R. Flowers of coltsfoot           six ounces  
Figs  
Raisins  
Jujubes, each,                   two ounces  
Water                               twelve pints  
Boil down one-third, adding  
Liquorice root                   two ounces  
Strain. As a demulcent drink. *Taddei.*

## U

## ULMUS CAMPESTRIS.

## ELM BARK.

This is a large tree, a native of Europe, having strong, spreading branches, and a rough, cracked bark.

*Sex. Syst.* Pentand. digyn. *Nat. Syst.* Ulmaceæ.

*Linn. Sp. Pl.* 327. *Lindley Flor. Med.* 307.

The part used is the inner bark; this is thin, tough, of a brownish yellow color, inodorous, of a mucilaginous, slightly astringent taste. It is demulcent, and somewhat astringent, and also acts as an alterative.

## DECOCTION OF ELM BARK.

R. Fresh elm bark,  
bruised,                   two ounces and a-half  
Distilled water                   two pints  
Boil down to one pint, and strain.  
*Lond. Ph.*

From four to six fl. ounces, two or three times a-day in cutaneous affections.

## COMPOUND INFUSION OF ELM BARK.

R. Elm bark  
Burdock  
Dock root  
Bittersweet  
Fumitory, each,                   two drachms

Boiling water                   one pint  
Infuse for four hours, strain, and add  
Syrup of sarsaparilla           one ounce  
To be taken in twenty-four hours, in divided doses, in chronic exanthemata.  
*Cadet.*

## ULMUS FULVA.

## SLIPPERY ELM.

A very lofty tree, found in the United States, especially west of the mountains, growing in dry, elevated situations.

*Sex. Syst.* Pentand. digyn. *Nat. Syst.* Ulmaceæ.

*Mich. Fl. Am.* 172. *U. S. Dispen.* 726.

The inner bark is the part used. This is in long, flat strips, of a tawny color on the outer surface, and reddish on the inner, of a peculiar, though not strong odor, and a mucilaginous taste. It is a good demulcent, and is much used in diseases of the mucous membranes. It is also very nutritious.

## INFUSION OF SLIPPERY ELM BARK.

R. Slippery elm bark, bruised, one ounce  
Boiling water                   one pint  
Macerate for two hours, and strain.  
*U. S. Ph.*

To be used freely, as a demulcent.



CATAPLASM OF SLIPPERY ELM BARK.  
 R. Ground slippery elm bark at will  
 Hot water sufficient  
 Make cataplasm. A soothing application  
 to irritable ulcers, &c.

## UVA URSI.

## UVA URSI.

*Uva ursi* is the leaves of *Arctostaphylos uva-ursi*, a small, evergreen shrub, inhabiting the northern parts of both continents, and high mountains in more southern latitudes.

*Sex. Syst.* Decand. monog. *Nat. Syst.* Ericaceæ.

Sprengel, ii. 287. Griffith, Med. Bot. 425.

The leaves, when dried, have a somewhat hay-like smell; their taste is astringent and bitterish, and finally sweetish. They are astringent and tonic, with some special action on the urinary organs. They are much used in diseases of the genito-urinary organs. Dose, in powder, from a scruple to a drachm, three or four times a-day.

## COMPOUND POWDER OF UVA URSI.

R. Powdered *uva ursi*  
 Peruvian  
 bark, each, two drachms  
 opium three grains  
 Mix, and divide into six powders. One,  
 twice a-day, with lime water, in nephritic  
 complaints, and as an antilithic. *Ferriar.*

R. Powdered  
*uva ursi* one drachm and a-half  
 Bicarbonate of soda one drachm  
 Mix, and divide into twelve powders. One  
 thrice a-day, in diseases of the kidney and  
 bladder. *Ellis.*

R. Powdered *uva ursi* half an ounce  
 gum Arabic two drachms  
 jalap one drachm

Powdered sugar half an ounce  
 Oil of orange peel six drops  
 Mix. Dose, one drachm, in nephritic  
 colic. *Quarin.*

## DECOCTION OF UVA URSI.

R. *Uva ursi* one ounce  
 Water twenty fl. ounces  
 Boil down to a pint, and strain.

*U. S. Ph.*

Dose, from one to two fl. ounces, three or four  
 times a-day.

## EXTRACT OF UVA URSI.

R. *Uva ursi* two pounds and a-half  
 Boiling water two gallons  
 Macerate for twenty-four hours, boil down  
 to a gallon, strain while hot, evaporate to  
 a proper consistence. *Lond. Ph.*

Dose, five to thirty grains.

## MIXTURE OF UVA URSI.

R. *Uva ursi* one ounce and a-half  
 Milfoil one ounce  
 Water three pints  
 Boil down to two pints, and add, at close,  
 Liquorice one ounce  
 Strain, and add  
 Syrup of cinnamon two ounces  
 To be taken in divided doses, in passive  
 hemorrhages of lungs and bladder.

*Radius.*

## SYRUP OF UVA URSI.

R. *Uva ursi* eight ounces  
 Boiling water two pints  
 Infuse the finely bruised leaves in the  
 water for three hours, put in a displace-  
 ment filter, and add water sufficient to  
 obtain two pints of infusion. Evaporate  
 to one pint, and add two pounds (av.) of  
 sugar, and make syrup. *W. Procter.*

## V

## VALERIANA.

## VALERIAN.

Several species of Valerian are used in medi-  
 cine, but one only is officinal, *Valeriana officinalis*, a handsome, perennial, herbaceous plant,  
 with white or rose-colored flowers, in terminal  
 corymbs.

*Sex. Syst.* Triand. monog. *Nat. Syst.* Valeriana-  
 ceæ.

Linn. Sp. Pl. 45. Griffith, Med. Bot. 384.

The part used is the root; this consists of

many long, slender fibres, arising from a tuber-  
 culated rhizome; the color is yellowish, or  
 brownish; the odor is strong and peculiar, and  
 the taste is bitter and aromatic. It is an active  
 antispasmodic, and is much used in nervous dis-  
 eases. The dose is from half a drachm to a  
 drachm, two or three times a-day.

## COMPOUND POWDER OF VALERIAN.

R. Powdered valerian one drachm  
 orange leaves one scruple  
 Muriate of ammonia two grains



Oil of cajeput four drops  
 Mix. A teaspoonful, four times a-day, in  
 hot tea. In epilepsy. *Niemann.*

R. Powdered valerian one ounce  
 Flowers of zinc one scruple  
 Musk ten grains

Mix. As an antispasmodic, in teaspoonful  
 doses. *Brera.*

R. Powdered valerian one scruple  
 Aromatic powder ten grains

Mix. To be taken three or four times  
 a-day. In hysteria, hemicrania, chlorosis,  
 &c. *A. T. Thomson.*

#### BOLUS OF VALERIAN AND IRON.

R. Powdered valerian one drachm  
 Carbonate of iron ten grains  
 Mucilage of gum Arabic sufficient

Mix, and make bolus. One, three times  
 a-day, as antispasmodic. *Ellis.*

#### BOLUS OF VALERIAN AND SULPHATE OF POTASSA.

R. Powdered vale-  
 rian two drachms and a-half  
 Sulphate of potassa eighteen grains  
 Syrup of orange peel sufficient

Mix, and make eighteen boluses. *Cadet.*

#### COMPOUND PILLS OF VALERIAN.

R. Powdered valerian half a drachm  
 Castor

Oxide of zinc, each, one scruple  
 Mix, and make eighteen pills. Three,  
 thrice a-day. *Dupuytren.*

#### ELECTUARY OF VALERIAN.

R. Powdered valerian  
 orange leaves, each, six  
 drachms

Syrup of wormwood sufficient  
 Mix. Two or three teaspoonfuls a-day.  
*Swediaur.*

R. Powdered valerian two ounces  
 Peruvian bark half an ounce  
 Carbonate of ammonia two drachms  
 Ginger syrup sufficient

Mix. Dose, a drachm every hour or two,  
 in periodic hemicrania. *Donald Monro.*

#### INFUSION OF VALERIAN.

R. Valerian half an ounce  
 Boiling water one pint

Macerate for an hour in a covered vessel,  
 and strain. *U. S. Ph.*

Dose, two fl. ounces, three or four times a-day.

#### COMPOUND INFUSION OF VALERIAN.

R. Valerian one ounce and a-half  
 Decoction of Peruvian bark two pints

Make an infusion, strain, and add

Syrup of chamo-  
 mile one fl. ounce and a-half

Camphor twenty grains

Mix. One to two fl. ounces, three times  
 a-day. *Saunders.*

#### TINCTURE OF VALERIAN.

R. Valerian, bruised, four ounces  
 Diluted alcohol two pints

Macerate for fourteen days, express, and  
 filter. *U. S. Ph.*

Dose, from one to four fl. drachms.

#### AMMONIATED TINCTURE OF VALERIAN.

R. Valerian, bruised, four ounces  
 Aromatic spirit of ammonia two pints

Macerate for fourteen days, express, and  
 filter. *U. S. Ph.*

Dose, one to two fl. drachms, in milk, or some  
 mucilaginous fluid.

#### COMPOUND AMMONIATED TINCTURE OF VALERIAN.

R. Valerian one ounce  
 Cloves

Mace, each, one drachm  
 Lemon peel one drachm and a-half

Aromatic spirit of  
 ammonia twenty-four fl. ounces

Macerate for a week, express, and filter.

Dose, a half to one fl. drachm. *Van Mons.*

#### COMPOUND TINCTURE OF VALERIAN.

R. Valerian  
 Castor, each, two ounces

Saffron one ounce

Peppermint water  
 Alcohol, each, twenty fl. ounces

Macerate for a week, express, and filter.

Dose, a half to one fl. drachm. *Wertz. Ph.*

#### WINE OF VALERIAN.

R. Valerian one ounce  
 Sherry wine one pint

Macerate for twenty-four hours, express,  
 and filter. A spoonful, as may be  
 required. *Niemann.*

#### SYRUP OF VALERIAN.

R. Bruised valerian one pound  
 Water eight pints

Macerate for twelve hours, then distil off  
 one pint and a-half, strain, and filter the  
 remainder. Mix with eight pounds of  
 syrup; evaporate to six and a-half pounds,  
 and add the distilled fluid. *Par. Cod.*

Mix. Dose, a drachm every hour or two,  
 in periodic hemicrania. *Donald Monro.*

Macerate for an hour in a covered vessel,  
 and strain. *U. S. Ph.*

Dose, two fl. ounces, three or four times a-day.

#### COMPOUND INFUSION OF VALERIAN.

R. Valerian one ounce and a-half  
 Decoction of Peruvian bark two pints

Make an infusion, strain, and add

R. Valerian one scruple  
 Carbonate of ammonia ten grains

Cinnamon water two fl. ounces

Mix. As a draught, every fourth hour,  
 in nervous headache, &c. *Ellis.*



## MIXTURE OF VALERIAN AND HOFFMANN'S ANODYNE.

- R. Valerian six drachms  
Boiling water eight fl. ounces  
Make an infusion, strain, and add  
Cinnamon water two fl. ounces  
Hoffmann's anodyne two fl. drachms  
Syrup one fl. ounce  
Mix. A spoonful, as required. *Foy.*

## TINCTURE OF VALERIAN AND HOFFMANN'S ANODYNE.

- R. Tincture of valerian  
Hoffmann's anodyne, each, one fl. ounce  
Mix. A teaspoonful, as may be required. *Ellis.*

## OIL OF VALERIAN.

- R. Bruised valerian at will  
Water sufficient  
Distil, and separate the oil from the product.

## MIXTURE OF OIL OF VALERIAN AND AMMONIA.

- R. Oil of valerian eight drops  
Aromatic spirit of ammonia one fl. drachm  
Water four fl. ounces  
Sugar two drachms  
Mix. A tablespoonful, every two or three hours. *Ellis.*

## EXTRACT OF VALERIAN.

- R. Coarsely powdered valerian at will  
Moisten with twice its weight of alcohol (.834), let macerate for three days, put into a displacement apparatus, and displace with alcohol (.915). The first product, on being distilled over a water-bath, affords a dark green resin, containing most of the volatile oil, and valerianic acid. The product of the distillation is to be diluted with distilled water, to reduce it to (.935), and the valerian exhausted by additional spirit. Mix the tinctures, filter and distil. The residue is to be evaporated and mixed with the first product. One grain is equal to five of valerian. *Burin.*

## FLUID EXTRACT OF VALERIAN.

- R. Valerian eight ounces  
Ether four fl. ounces  
Alcohol twelve fl. ounces  
Diluted alcohol sufficient  
Powder the valerian coarsely, and introduce it uniformly, but without packing, into a displacer. Add a mixture of the ether and alcohol, and let it percolate slowly through the powder, and pour on the diluted alcohol until the ethereal men-

struum has been displaced, when the product will amount to nearly a pint; add more diluted alcohol, until ten fl. ounces pass; place the ethereal tincture in a shallow vessel, and let it evaporate spontaneously, till it is reduced to six fl. ounces. Pour off the clear liquid, and mix it with the ten ounces of tincture, and the oleoresin adhering to the sides of the vessel, dissolved in a little alcohol and added to the rest; let stand, with occasional agitation, for three or four hours, and then filter. One fl. ounce is equal to half an ounce of the root. *W. Procter.*

- R. Valerian eight ounces  
Alcohol eight fl. ounces  
Sulphuric ether four fl. ounces  
Water sufficient

Bruise the valerian finely, put it in a displacement apparatus, and add the alcohol and ether, mixed; remove the last portions of the tincture by water, and set the solution aside in a warm room to evaporate, till all the ether and half the alcohol have escaped. The residue will amount to four fl. ounces, of a greenish color. Proceed to displace with the water, setting aside the first eight fl. ounces of the product, and proceed with the process till the root is exhausted. Evaporate this to four fl. ounces; mix all the solutions. The dose is about a teaspoonful. *E. T. Ellis.*

## VALERIANIC ACID.

- R. Valerian root at will  
Distil with water until the distillate has but a slight acid reaction; add carbonate of soda to the distilled water gradually, until it is rendered slightly alkaline, observing the weight of the carbonate used. Evaporate the solution of valerianate of soda to a small quantity; add sufficient sulphuric acid to saturate the soda used, and distil in a glass retort. The valerianic acid will be found in the receiver floating on the water, and should be removed, and preserved in glass vials.

## VANILLA.

## VANILLA.

Under this name are included the seed pods of several species of *Vanilla*, but that most generally known is the *V. aromatica*, a sarmentose plant, a native of Mexico and South America.

*Sex. Syst.* Gynand. monand. *Nat. Syst.* Orchidaceæ.

Swartz, *Occident*, iii. 1518. *Flore Medicale*, vi. 345.



The part used is the pod; this is long, slender, cylindrical, compressed, of a blackish-brown color, formed of two valves, and filled with a brown pulp, containing a great quantity of small, blackish, shining seeds. The odor is peculiar but pleasant, the taste aromatic, hot, and pungent. It is principally used in this country for flavoring ice creams, &c., but is esteemed by European writers as an aromatic stimulant, increasing the power of the muscular system, and exciting the sexual feelings. It has been given in hysteria, rheumatism, impotence, &c. The dose is from eight to twelve grains.

VANILLA SUGAR.

R. Tincture of vanilla one ounce  
Sugar eight ounces  
Mix well, dry in a stove, and pulverize.  
*Beral.*

VANILLA LOZENGES.

R. Vanilla one ounce  
Sugar seven ounces  
Gum tragacanth half a drachm  
Water half an ounce  
Mix, and make lozenges of twelve grains.  
*Beral.*

TINCTURE OF VANILLA.

R. Vanilla one part  
Alcohol four parts  
Macerate for fourteen days, express, and filter. Dose, forty drops.  
*Cottureau.*

SPIRIT OF VANILLA.

R. Vanilla one part  
Alcohol  
Water, each, twelve parts  
Distil twelve parts.  
*Soubeiran.*

VANILLA ARROWROOT.

R. Milk fifteen ounces  
Sugar  
Vanilla sugar  
Arrowroot, each, one ounce  
Mix, and boil, constantly stirring.  
*Beral.*

VANILLA MILK.

R. Milk sixteen ounces  
Vanilla sugar half an ounce  
Dissolve, and strain.  
*Beral.*

ESSENCE OF VANILLA.

R. Vanilla nine ounces  
Spirit of abelmoschus two pints  
Cloves thirty grains  
Musk seven grains  
Used as a perfume, and for flavoring.  
*Cooley.*

FLUID EXTRACT OF VANILLA.

R. Vanilla one ounce  
Sugar three ounces  
Syrup  
Water, each, half a pint  
Deodorized alcohol one fl. ounce

Cut the vanilla in thin, transverse slices, triturate it with the sugar, till reduced to powder moderately fine, add the syrup with two ounces of the water, put the mixture into a strong pint bottle, cork, and tie it over; place it in a vessel of water, which is then to be heated to the boiling point, and kept there for half an hour; remove the cork, and strain. The residue is then to be mixed with the remainder of the water and the alcohol, the mixture returned to the bottle, and heated as before, strained, and mixed with the first product.  
*W. Procter.*

VERATRIA.

VERATRIN.

R. Cevadilla, bruised, two pounds  
Alcohol three gallons  
Diluted sulphuric acid  
Solution of ammonia  
Purified animal charcoal  
Magnesia, each, sufficient

Boil the cevadilla in a gallon of the alcohol, in a retort with a receiver attached, for an hour, and pour off the liquor. To the residue, add another gallon of alcohol, together with the distilled portion, boil for an hour, and pour off the liquor; repeat a third time. Express, mix, and strain the liquors, and distil off the alcohol, on a water-bath. Boil the residue three or four times in water, acidulated with sulphuric acid, mix, and strain the liquors, and evaporate to the consistence of syrup. Add magnesia in slight excess, shake frequently, express, and wash residue. Repeat expression and washing several times. Dry the residue, digest it with gentle heat, several times in alcohol, straining after each digestion. Distil off alcohol from the mixed liquors, boil the residue for fifteen minutes in water, with a little sulphuric acid and animal charcoal, and strain. Thoroughly wash the residue, mix the washings with the strained liquor, evaporate gently to consistence of syrup, add as much ammonia as will precipitate the veratria, separate this and dry it. *U. S. Ph.*

Veratria is a violent stimulant, and active poison. It is given internally in minute doses, but is more generally used externally, in gout, rheumatism, nervous affections, &c. The dose is from one-twelfth to one-sixth of a grain.

PILLS OF VERATRIN.

R. Veratrin half a grain  
Syrup of gum  
Powdered gum Arabic, each, sufficient



Mix, and make six pills. Dose, one, to be increased to three. *Magendie.*

#### PILLS OF VERATRIN AND HENBANE.

R. Veratrin one grain  
Extract of henbane  
Powdered liquorice, each, ten grains  
Mix, and make twelve pills. One, thrice a-day. *Turnbull.*

#### TINCTURE OF VERATRIN.

R. Veratrin four grains  
Alcohol one fl. ounce  
Dissolve. Dose, from ten to twenty drops in a wineglassful of water, in dropsy, gout, rheumatism, &c. *Magendie.*

#### LOTION OF VERATRIN.

R. Veratrin one scruple to one drachm  
Alcohol two fl. ounces  
Mix. As embrocation, to be used with caution. *Turnbull.*

#### LINIMENT OF VERATRIN.

R. Veratrin eight grains  
Alcohol  
Soap liniment, each, four fl. drachms  
Mix well. *Brande.*

#### OINTMENT OF VERATRIN.

R. Veratrin five to twenty grains  
Lard one ounce  
Mix. A piece about the size of a hazelnut to be rubbed for five or fifteen minutes, over the seat of the disease, in rheumatism, &c. *Turnbull.*

R. Veratrin one scruple  
Rub with  
Olive oil one drachm  
Add  
Spermaceti ointment seven drachms  
Mix. As the last. *Turnbull.*

#### OINTMENT OF VERATRIN AND OPIUM.

R. Veratrin half a drachm  
Powdered opium one drachm  
Lard one ounce and a-half  
Mix. As a friction, in rheumatism. *Dunghison.*

#### OINTMENT OF VERATRIN AND IODINE.

R. Veratrin one scruple  
Iodide of potassium half a drachm  
Lard one ounce  
Mix. *Jourdan.*

### VERATRIÆ MURIAS.

#### MURIATE OF VERATRIN.

R. Veratria at will  
Muriatic acid sufficient  
to saturate; filter, evaporate, and crystallize. *Giordano.*

### VERATRIÆ NITRAS.

#### NITRATE OF VERATRIN.

### VERATRIÆ SULPHAS.

#### SULPHATE OF VERATRIN.

Are made in same manner, using the appropriate acids.

#### SOLUTION OF SULPHATE OF VERATRIN.

R. Sulphate of veratria one grain  
Distilled water two fl. ounces  
Dissolve. Dose, one to two fl. drachms, in a mixture, as a substitute for the *Eau Medicinale.* *Cadet.*

### VERATRUM ALBUM.

#### WHITE HELLEBORE.

A perennial, herbaceous plant, with a fleshy, fusiform rhizome, having numerous fibres. It is a native of the mountainous parts of Europe. *Sex. Syst.* Hexand. trigyn. *Nat. Syst.* Melanthaceæ.

*Linn. Sp. Pl.* 1479. *Griffith, Med. Bot.* 643.

The root, which is the officinal part, is in cylindrical, somewhat conical pieces, with numerous radicles; these are corrugated, and of a blackish brown color. Their odor is slight, and their taste at first sweetish, and then bitter and acrid. White hellebore is an active irritant, causing violent vomiting and purging. Externally, it acts like the other powerful acids. Internally, in small doses, it has been used in many diseases, but is now principally employed externally in the treatment of cutaneous affections, &c. Dose, from a grain to two grains, gradually increased.

#### POWDER OF WHITE HELLEBORE.

R. Powdered white hellebore three grains  
starch one scruple  
Mix. As an errhine in amaurosis, and lethargic affections. *Radius.*

#### DECOCTION OF WHITE HELLEBORE.

R. White hellebore ten drachms  
Distilled water two pints  
Boil down to a pint, and add  
Rectified spirit three fl. ounces  
and strain. *Lond. Ph.*

As a wash, in cutaneous diseases.

#### WINE OF WHITE HELLEBORE.

R. Bruised white hellebore four ounces  
Wine one pint  
Macerate for fourteen days, express, and filter. *U. S. Ph.*

Dose, ten minims, gradually increased, two or three times a-day, in gout and rheumatism.

#### TINCTURE OF WHITE HELLEBORE.

R. Bruised white hellebore four ounces  
Diluted alcohol one pint  
Macerate for eight days, express, and filter. *Ed. Ph.*

Dose, as of the wine, in same cases.



OINTMENT OF WHITE HELLEBORE.

R. Powdered white hellebore two ounces  
Lard eight ounces  
Oil of lemons twenty minims  
Mix. *U. S. Ph.*

As a friction in itch, &c. To be used with caution on children.

COMPOUND OINTMENT OF WHITE HELLEBORE.

R. Powdered white hellebore two ounces  
nitre one drachm  
Soft soap  
Flowers of sulphur, each, six ounces  
Lard one pound and a-half  
Mix. As a friction in itch. *Phæbus.*

VERATRUM VIRIDE.

AMERICAN HELLEBORE.

A tall, perennial plant, with a thick, fleshy rhizome, having a tunicated top, and numerous fibrous radicles. It is a native of the United States, growing in damp meadows, and wet situations.

*Sex. Syst.* Hexand. trigyn. *Nat. Syst.* Melanthaceæ.

Aiton, Hort. Kew. iii. 422. Griffith, Med. Bot. 643.

The root, in a fresh state, has an unpleasant odor, which disappears on drying. Its taste is bitter and acrid. Its properties are much the same as those of the white hellebore, but is stated not to purge, and to rapidly reduce the frequency and force of the pulse. Dose, one grain, gradually increasing.

TINCTURE OF AMERICAN HELLEBORE.

R. American hellebore,  
bruised, six ounces  
Diluted alcohol one pint  
Macerate for fourteen days, express, and filter. Dose, a half to one fl. drachm.  
*Osgood.*

WINE OF AMERICAN HELLEBORE.

R. American hellebore,  
bruised, six ounces  
White wine fourteen fl. ounces  
Diluted alcohol two ounces  
Macerate for fourteen days, express, and filter. Dose, as of tincture. *Osgood.*

The recent root should be used in these two preparations.

EXTRACT OF AMERICAN HELLEBORE.

R. Recent root of American hellebore at will  
Bruise well, and express the juice, and evaporate to proper consistence, by a gentle heat, or by exposure to the sun. Dose, from a fourth to half a grain. *Osgood.*

OINTMENT OF AMERICAN HELLEBORE.

R. Powdered extract of American hellebore one drachm

Simple cerate one ounce  
Oil of lemons three minims  
Mix. *Osgood.*

PILLS OF AMERICAN HELLEBORE.

R. Extract of American hellebore ten grains  
Opium five grains  
Soap fifteen grains  
Mucilage of gum Arabic sufficient  
Mix, and make thirty pills. One, every three or four hours, in rheumatism, &c.  
*Osgood.*

MIXTURE OF AMERICAN HELLEBORE.

R. Tincture of American hellebore bloodroot  
Camphorated tincture of opium equal parts  
Mix. Dose, fifty to sixty minims, in catarrh. *Osgood.*

VIOLA ODORATA.

SWEET VIOLET.

A well-known, small, perennial plant, with fragrant, blue flowers, a native of Europe, and generally cultivated in our gardens.

*Sex. Syst.* Pentand. monog. *Nat. Syst.* Violaceæ.

Linn. Sp. Pl. 1324. Griffith, Med. Bot. 140.

The flowers are the officinal portion. These are of a blue color, almost inodorous when dried, and of a slightly bitter taste. They are principally used as a demulcent, and laxative for infants.

CONSERVE OF VIOLETS.

R. Violet flowers, fresh, one part  
Sugar three parts  
Rub together. *Soubéiran.*

SYRUP OF VIOLETS.

R. Fresh violet flowers one pound  
Boiling water two pints and a-half  
Infuse for twenty-four hours, strain, and dissolve.  
Sugar seven pounds and a-half  
And form syrup. *Ed. Ph.*

A gentle laxative for infants, in doses of one to two fl. drachms.

HONEY OF VIOLETS.

R. Flowers of violets  
Boiling water, each, six pounds  
Infuse, strain, and add  
Honey six pounds  
Evaporate to a proper consistence. *Wurt. Ph.*

OIL OF VIOLETS.

R. Fresh violet flowers fifteen ounces  
Almond oil five pounds  
Infuse, with three equal quantities of the flowers, and strain. Used as an article of perfumery. *Taddei.*



## W

## WINTERA.

## WINTER'S BARK.

This is derived from an evergreen tree, found in the southern parts of South America, the *Drimys winteri*, and also from other species, growing in Brazil, &c.

*Sex. Syst.* Polyand. trigyn. *Nat. Syst.* Magnoliaceæ.

Forster, Gen. 84. Griffith, Med. Bot. 100.

The bark is in quilled pieces of some length, of a pale, yellowish, or dull, reddish gray, with

darker spots externally, and of a dark cinnamon internally. It has an aromatic odor, and a warm, spicy taste. It is a stimulating, aromatic tonic. The dose, in powder, is about half a drachm.

## TINCTURE OF WINTER'S BARK.

R. Bruised winter's bark                    one part  
Diluted alcohol                                eight parts  
Digest with one-half of the alcohol, and repeat with the other. Unite the two tinctures, and filter.

## Z

## ZINCUM.

## ZINC.

## ZINCI OXIDUM.

## OXIDE OF ZINC.

R. Sulphate of zinc                                one pound  
Carbonate of ammonia                    six ounces and a-half  
Distilled water                                three gallons  
Dissolve the sulphate of zinc and carbonate of ammonia, separately, in twelve pints of the water, strain, and mix the solutions. Wash the precipitate well with water, and expose to a strong heat.

*U. S. Ph.*

Tonic and antispasmodic, and given in nervous and spasmodic affections, in doses of two to ten grains and upwards several times a-day. Also used externally.

## POWDER OF OXIDE OF ZINC.

R. Oxide of zinc                                three grains  
White sugar                                    five grains  
Mix. One, every four hours. Useful in gastric or spasmodic cough.

*A. T. Thomson.*

## COMPOUND POWDER OF OXIDE OF ZINC.

R. Oxide of zinc                                six grains  
Extract of henbane                    one grain  
Oil of valerian                                one drop  
Powdered valerian                    half a drachm

Mix, and divide into nine powders. One, thrice a-day, in nervous disorders.

*Radiis.*

R. Oxide of zinc                                one drachm  
Powdered opium                                ten grains  
Peruvian bark  
Fennel, each,                                one ounce

Mix, and divide into fifty powders. One, every two hours, in whooping-cough.

*Augustin.*

## POWDER OF OXIDE OF ZINC AND COLOMBA.

R. Oxide of zinc                                four grains  
Powdered colomba                                thirty grains  
Mix, and divide into four powders. One, every three hours, in dyspepsia. *Brera.*

## PILLS OF OXIDE OF ZINC.

R. Oxide of zinc                                two scruples  
Conserve of roses                                sufficient  
Mix, and make ten pills. One, three or four times a-day, in epilepsy, chorea, &c.

*Ellis.*

R. Oxide of zinc                                three grains  
White sugar                                    five grains  
Mix. Make a powder, to be taken every four hours. In gastric or spasmodic cough.

*A. T. Thomson.*

## COMPOUND PILLS OF OXIDE OF ZINC.

R. Oxide of zinc                                five grains  
Extract of cascarrilla                    six grains  
Syrup of ginger                                sufficient



Mix, and make three pills; one to be taken three times a-day, in epilepsy. *Ainslie.*

R. Oxide of zinc                   one scruple  
Powdered valerian           thirty grains  
Castor                           four grains  
Syrup                           sufficient

Mix, and make twelve pills. To be taken during the day, in epilepsy, &c. *Foy.*

#### LOTION WITH OXIDE OF ZINC.

R. Oxide of zinc                   one ounce  
Elder flower  
water                   one fl. ounce and a-half

Agitate together. In pustular erysipelas. *Augustin.*

#### OINTMENT OF OXIDE OF ZINC.

R. Oxide of zinc                   one ounce  
Lard                           six ounces  
Mix. *U. S. Ph.*

A mild, astringent application, in chronic ophthalmia, and in excoriation and ulceration.

#### OINTMENT OF OXIDE OF ZINC AND OPIUM.

R. Oxide of zinc                   half a drachm  
Fresh butter                   one drachm  
Powdered opium               one grain  
Mix. In chronic impetigo. *Henke.*

#### OINTMENT OF OXIDE OF ZINC AND CALOMEL.

R. Oxide of zinc                   half a drachm  
Calomel  
Powdered aloes, each,       one grain  
Fresh butter                   two drachms  
Mix. In scrofulous ophthalmia. *Scarpa.*

#### CERATE OF OXIDE OF ZINC AND LYCOPodium.

R. Oxide of zinc  
Lycopodium, each,       fifteen grains  
Simple cerate               half an ounce  
Mix. In ulceration of the breasts, tetter, &c. *Hufeland.*

### ZINCI CHLORIDUM.

#### CHLORIDE OF ZINC.

R. Zinc, in small  
pieces,                   two ounces and a-half  
Nitric acid  
Prepared chalk, each,   one drachm  
Muriatic acid               sufficient

Add the zinc and muriatic acid together in a glass vessel, and dissolve; strain, add the nitric acid, evaporate to dryness. Dissolve in water, add the chalk, let rest for twenty-four hours, filter, and again evaporate. *U. S. Ph.*

R. Solution of chloride of barium at will

Solution of sulphate of zinc sufficient to precipitate sulphate of baryta. Filter and evaporate the fluid to obtain crystals of chloride of zinc. *Righini.*

Principally used externally, as a caustic and stimulant; also given internally, in very small doses, in the neuroses, &c.

#### CHLORIDE OF ZINC. (*Disinfectant.*)

R. Granulated zinc       four pounds (av.)  
Muriatic acid       four pounds, or sufficient  
Water               two gallons two pints

Add the acid gradually to the zinc, in a glass or porcelain vessel, till it is dissolved, taking care there is no excess of acid; strain through a coarse cloth, and add the water. This contains one part of the chloride in twelve, and is of the strength for a disinfectant. *E. Parrish.*

#### ETHEREAL TINCTURE OF CHLORIDE OF ZINC.

R. Chloride of zinc       four drachms  
Alcohol               one fl. ounce  
Sulphuric ether       two fl. ounces

Mix. Five drops every four hours, in a little sugar and water, gradually increasing the dose, in chorea, epilepsy, &c.

*Hufeland.*

#### LOTION OF CHLORIDE OF ZINC.

R. Chloride of zinc       eight grains  
Extract of aloes       two scruples  
Distilled water       four fl. ounces

Mix. As an application to atonic and foul ulcers. *Vogt.*

#### INJECTION OF CHLORIDE OF ZINC.

R. Liquid chloride  
of zinc               twenty-four drops  
Water               four fl. ounces

Mix, and filter. A small quantity to be injected a short distance into the urethra, two or three times a-day, in gleet.

*Gaudriot.*

#### VAGINAL SUPPOSITORY OF CHLORIDE OF ZINC.

R. Liquid chloride of zinc   five drops  
Sulphate of morphia       half a grain  
Mix with the following paste, three drachms.

Mucilage of tragacanth   six parts  
White sugar               three parts  
Starch                   nine parts

Make a suppository. To be used every day, or every other day. *Gaudriot.*

#### PASTE OF CHLORIDE OF ZINC.

R. Chloride of zinc       one part  
Wheat flour, or sulphate  
of lime,               four parts  
Water               sufficient



- R. Chloride of zinc                    one part  
 Wheat flour                    three parts  
 Water                    sufficient
- R. Chloride of zinc                    one part  
 Wheat flour                    two parts  
 Water                    sufficient

The chloride and flour are to be carefully and quickly mixed; add the water to half the quantity, so as to form a soft paste, mix with this the rest of the powder. Form into cakes of half a line to four lines in thickness. To remain on twenty-four hours, or more (six to ten are often sufficient). To produce an eschar in cancer, lupus, &c. *Canquoin.*

- R. Chloride of zinc                    one part  
 Chloride of antimony            half a part  
 Flour                    two and a-half parts  
 Water                    sufficient

To be mixed as above, so as to be moulded into any desired form. As a caustic, in nodulated cancerous tumors.

*Canquoin.*

Powdered opium may be added to any of the above, to mitigate the pain.

## ZINCI CYANURETUM.

### CYANIDE OF ZINC.

- R. Sulphate of zinc                    one part  
 Distilled water                    ten parts
- Dissolve; add  
 Solution of cyanide of  
 potassium                    sufficient  
 to form precipitate, being added by drops.  
 Filter, wash the precipitate, and dry.

*Guibourt.*

Used in spasmodic affections, &c., in doses of a twelfth to a quarter of a grain.

### POWDER OF CYANIDE OF ZINC.

- R. Cyanide of zinc                    three grains  
 Magnesia                    twenty-four grains  
 Powdered cinnamon            twelve grains
- Mix, and divide into twelve powders. One, every four hours, in nervous affections, &c. *Guibourt.*

### PILLS OF CYANIDE OF ZINC.

- R. Cyanide of zinc                    fifteen grains  
 Extract of liquorice            two drachms
- Mix, and make sixty pills. One, three times a-day. *Kopp.*

## ZINCI FERROCYANURETUM.

### FERROCYANIDE OF ZINC.

- R. Ferrocyanide of potassium    sixty parts

Sulphate of zinc                    eighty parts  
 Dissolve separately in boiling water, add the solutions together; collect the precipitate of ferrocyanide of zinc, wash it well, and dry. *Schindler.*

Given in doses of one grain, twice or thrice a-day, gradually increasing the quantity, in nervous affections, nocturnal pains, &c.

### POWDER OF FERROCYANIDE OF ZINC.

- R. Ferrocyanide of zinc            six grains  
 Magnesia                    four grains  
 Powdered cinnamon            three grains
- Mix. Every four hours, in nervous affections, cramp of the stomach, &c. *Radius.*

The quantity of the salt of zinc is too large, but it is thus ordered in various authors.

### COMPOUND PILLS OF FERROCYANIDE OF ZINC.

- R. Ferrocyanide of zinc            fifteen grains  
 Powdered valerian            half a drachm  
 Extract of valerian            sufficient
- Mix, and make sixty pills. Two, morning and evening, in chorea. *Rosensteil.*

- R. Ferrocyanide of zinc            ten grains  
 Magnesia                    two scruples  
 Oil of valerian                    twenty drops  
 Syrup of chamomile            five scruples  
 Powdered valerian            ten scruples

Mix, and make into three-grain pills. Six, three times a-day, gradually increasing. *Vogt.*

### SOLUTION OF FERROCYANIDE OF ZINC.

- R. Ferrocyanide of zinc            four grains  
 Distilled water                    two fl. ounces
- One-fourth part, four times a-day, in chorea. *Radius.*

### MIXTURE OF FERROCYANIDE OF ZINC.

- R. Ferrocyanide of  
 zinc                    eight to twenty grains  
 Powdered gum Arabic            two drachms  
 Cherry water                    three fl. ounces  
 Wine of opium                    one fl. scruple  
 Cherry laurel water            half a fl. ounce
- Mix. As a collyrium, in scrofulous and rheumatic ophthalmia. *Ammon.*

## ZINCI IODIDUM.

### IODIDE OF ZINC.

- R. Iodine                    one hundred and seventy parts  
 Powdered zinc                    twenty parts
- Heat in matrass till they sublime. *Magendie.*

- R. Iodine                    two parts  
 Powdered zinc                    one part



Digest in a little water, and agitate till the solution is colorless. Evaporate with a gentle heat.

*Beasley.*

Used externally.

#### OINTMENT OF IODIDE OF ZINC.

R. Iodide of zinc                   one drachm  
Lard                                   one ounce

As a friction to tumors, &c.                   *Ure.*

#### COLLYRIUM OF IODIDE OF ZINC.

R. Iodide of zinc                   four grains  
Distilled water                   six fl. ounces

Dissolve.

#### SOLUTION OF IODIDE OF ZINC.

R. Iodide of zinc           ten to thirty grains  
Distilled water           one fl. ounce

Dissolve. To be applied, by means of a sponge tied to a quill, to enlarged tonsils; to be followed, after some days' perseverance, by a saturated solution applied by a camel's hair brush.

*T. Ross.*

#### SYRUP OF IODIDE OF ZINC.

R. Iodine                   four drachms  
Powdered zinc           two drachms  
Water                   four fl. ounces

Agitate till colorless; filter into

Syrup                   twelve fl. ounces.  
*A. T. Thomson.*

### ZINCI ACETAS.

#### ACETATE OF ZINC.

R. Acetate of lead                   one pound  
Granulated zinc                   nine ounces  
Distilled water                   three pints

Dissolve the acetate in the water, and filter. Add the zinc, and agitate occasionally, in a stopped bottle, till the liquid gives no precipitate, with a solution of iodide of potassium. Filter, evaporate by a gentle heat to one-fifth, and set aside to crystallize. Decant, and dry the crystals on bibulous paper. If they are colored, dissolve in distilled water, heat, and drop into the hot solution a filtered solution of chlorinated lime, as long as any precipitate of iron takes place; filter, acidulate with a few drops of acetic acid, evaporate, and crystallize.

*U. S. Ph.*

Used as an external application.

#### COLLYRIUM OF ACETATE OF ZINC.

R. Sulphate of zinc  
Acetate of lead, each,           six grains  
Rose-water                   four fl. ounces

Mix, and filter.                   *Ellis.*

R. Acetate of zinc   fifteen to thirty grains

Distilled water           twelve fl. ounces  
Dissolve.                   *Ware.*

R. Sulphate of zinc           one drachm  
Acetate of lead           half a drachm  
Camphor                   one scruple  
Rose water                   twelve ounces  
Filter the solution.                   *Spielman.*

#### INJECTION OF ACETATE OF ZINC.

R. Acetate of zinc           eight grains  
Rose water                   four fl. ounces  
Dissolve. In gonorrhœa.                   *Ellis.*

#### LOTION OF ACETATE OF ZINC.

R. Acetate of zinc           four scruples  
Distilled vinegar           sixteen fl. ounces  
Dissolve.                   *Beral.*

#### TINCTURE OF ACETATE OF ZINC.

R. Sulphate of zinc  
Acetate of potassa, each,   one ounce  
Rub together, and add  
Diluted alcohol                   one pint  
Macerate for a week, stirring occasionally,  
and filter.                   *Dub. Ph.*

As collyrium and injection, properly diluted.

### ZINCI CARBONAS.

#### CARBONATE OF ZINC, OR CALAMINE.

##### PREPARED CARBONATE OF ZINC.

R. Carbonate of zinc                   at will  
Heat it to redness, and pulverize; then reduce to a very fine powder, as directed for prepared chalk.                   *U. S. Ph.*

Used externally, as a mild astringent and absorbent to excoriations and ulcerations, and also as an ingredient of cerates, &c.

##### PRECIPITATED CARBONATE OF ZINC.

R. Sulphate of zinc  
Carbonate of soda, each,   two pounds  
Water                   sufficient  
Dissolve the salts separately in a gallon of water, mix the solutions; wash the precipitated carbonate, first by decantation, and afterwards on a filter; then press, and dry it.

##### CERATE OF CARBONATE OF ZINC.

R. Prepared carbonate of zinc  
Yellow wax, each,           half a pound  
Lard                   two pounds  
Melt the lard and wax together, and when, on cooling, they begin to thicken, add the carbonate of zinc, and stir till cold.                   *U. S. Ph.*

Much used as a dressing to excoriations, shallow ulcerations, &c.



## PLASTER OF CARBONATE OF ZINC.

- R. Simple plaster  
Powdered carbonate of zinc, each, two pounds  
Yellow wax five ounces  
Suet  
Turpentine, each, one ounce and a-half  
Melt, and mix thoroughly. *Van Mons.*

- R. Prepared carbonate of zinc  
White lead, each, one ounce and a-half  
Yellow wax four ounces  
Olive oil six ounces  
Melt the wax with the oil, and add the two powders, stirring well. To diminish a too abundant suppuration in ulcers.

*Cadet.*

## ZINCI SULPHAS.

## SULPHATE OF ZINC.

- R. Zinc, in small pieces, four ounces  
Sulphuric acid six ounces  
Distilled water four pints  
Introduce the zinc and water into a glass vessel, and add the sulphuric acid by degrees; when all effervescence has ceased, filter, boil to a pellicle, and set aside to crystallize. *U. S. Ph.*

It is astringent, tonic, and, in large doses, a prompt and effective emetic. Dose, as a tonic, one to two grains; as emetic, ten to thirty.

## PILLS OF SULPHATE OF ZINC.

- R. Sulphate of zinc ten grains  
Conserve of roses sufficient  
Mix, and make twenty pills. One, three times a-day. In pertussis, &c. *Ellis.*

## PILLS OF SULPHATE OF ZINC AND TURPENTINE.

- R. Sulphate of zinc two drachms  
Venice turpentine sufficient  
Mix, and make sixty pills. As a tonic. *Beasley.*

## PILLS OF SULPHATE OF ZINC AND MYRRH.

- R. Sulphate of zinc ten grains  
Powdered myrrh one drachm and a-half  
Conserve of roses sufficient  
Mix, and make twenty pills. From two to four pills a-day, in hooping-cough. *Paris.*

## COMPOUND PILLS OF SULPHATE OF ZINC.

- R. Sulphate of zinc one grain  
Extract of gentian eight grains  
Syrup of ginger sufficient  
Mix, and make two pills, to be taken dur-

ing the day, with two fl. drachms of infusion of chamomile, in epilepsy. *Ainslie.*

## SOLUTION OF SULPHATE OF ZINC.

- R. Sulphate of zinc two grains  
Chamomile water three fl. ounces  
Dissolve. A spoonful, two or three times a-day, in chronic dysentery. *Radius.*

## COLLYRIUM OF SULPHATE OF ZINC.

- R. Sulphate of zinc six grains  
Acetate of lead ten grains  
Laudanum one drachm  
Water five fl. ounces  
Mix. As an application to the eyes in ophthalmia, after reduction of the inflammation. *Gregory.*

- R. Sulphate of zinc six grains  
Mucilage of quince seed one fl. ounce  
Distilled water six fl. ounces  
Spirit of camphor four or five drops  
Mix, and strain. *Scarpa.*

To be used as the last.

## COLLYRIUM OF SULPHATE OF ZINC AND CAMPHOR.

- R. Sulphate of zinc one scruple  
Camphor water  
Rose water, each, eight fl. ounces  
Mix. *Ainslie.*

## FOMENTATION WITH SULPHATE OF ZINC.

- R. Sulphate of zinc one drachm and a-half  
Decoction of pomegranate four pints  
Peruvian bark two pints  
Laudanum one fl. ounce  
Mix. As an application to external piles, and in vaginal gonorrhœa. *Brera.*

## GARGLE OF SULPHATE OF ZINC.

- R. Sulphate of zinc one drachm  
Honey half a fl. ounce  
Tincture of myrrh  
Brandy, each, one fl. ounce  
Rose water six fl. ounces  
Mix. In ulcerations of gums, &c., occasioned by excessive salivation. *Ellis.*

## INJECTION OF SULPHATE OF ZINC.

- R. Sulphate of zinc one drachm  
Camphor water two fl. ounces  
Water two pints  
Mix. *Swediaur.*

- R. Sulphate of zinc ten grains  
Powdered gum Arabic two drachms  
Laudanum one fl. drachm  
Distilled water eight fl. ounces  
Mix. In gonorrhœa. *Ellis.*

## LOTION OF SULPHATE OF ZINC.

- R. Sulphate of zinc sixteen grains



Rose water	sixteen fl. ounces
Mix.	<i>Beral.</i>
R. Sulphate of zinc	half a drachm
Water	half a pint
Lead water	half a pint
Mix.	<i>Beasley.</i>
R. Sulphate of zinc	four scruples
Distilled vinegar	one pint
Mix.	<i>Beral.</i>

## SULPHATE OF ZINC MIXTURE.

R. Sulphate of zinc	one grain
Decoction of cascarilla	twelve fl. drachms
Simple syrup	four fl. drachms
Mix, to be divided into four doses, one to be taken three times a-day, in hooping-cough.	<i>Ainslie.</i>

## ZINCI VALERIANAS.

## VALERIANATE OF ZINC.

R. Valerian root,	one hundred parts
recent,	
Water	five hundred parts
Sulphuric acid	ten parts
Bichromate of potassa	six parts
Macerate the coarsely powdered valerian in the water, to which had been added the sulphuric acid and the bichromate, for twenty-four hours, introduce the mixture into a still, and apply heat; returning the first fourth of the product to the still; the process is then to be continued, until the distilled water ceases to redden litmus paper. The product is then to be placed in a large matrass, and an excess of pure hydrated carbonate of zinc, is to be added, and allowed to digest at 190° F., for three or four hours, on a sand-bath, until saturated. The solution is then to be filtered and evaporated over a naked fire, till reduced to fifty parts, then placed on plates, and suffered to evaporate in a drying room, or by a moderate temperature.	
	<i>Lefort.</i>

R. Contused valerian	thirty-two ounces
Water	eight pounds
Sulphuric acid	three ounces
Macerate for two days, and distil as long as the product reddens litmus paper. Expose to the air for a month, then put it in a matrass with two hundred and twenty-five grains of recently prepared oxide of zinc, and digest on a sand-bath for eight or ten hours, at 175° F. agitating occasionally; filter whilst hot, evaporate to three-fourths, and dry the residue on earthen plates.	
	<i>Buisson.</i>

Has been recommended as a powerful tonic and antispasmodic, in neuralgia and hysteria, but has not been sufficiently tried to ascertain its true powers. Dose, one to two grains.

## PILLS OF VALERIANATE OF ZINC.

R. Valerianate of zinc	nine grains
Tragacanth	thirty grains
Mix, and make twelve pills. One, morning and evening, in nervous headache, &c.	
	<i>Devay.</i>

## MIXTURE OF VALERIANATE OF ZINC.

R. Valerianate of zinc	one grain and a-half
Distilled water	five drachms, twenty drops
Syrup	one ounce
Mix. A spoonful, every half hour.	
	<i>Devay.</i>

## ZINGIBER.

## GINGER.

The rhizome of *Zingiber officinale*, a small, herbaceous, perennial plant, a native of the East Indies.

*Sex. Syst.* Monand. monog. *Nat. Syst.* Zingiberaceæ.

Roscoe, Trans. Linn. Soc. viii. 348. Griffith, Med. Bot. 629.

The rhizome, or root, is creeping, knotty, lobated or palmated, and fleshy, with an agreeable, pungent, camphoraceous odor, and an aromatic, acrid, burning taste. Two kinds are met with in commerce, the black and the white; the first consists of the roots, scalded in hot water and dried; the second is composed of these roots, deprived of their epidermis previous to desiccation. Ginger contains a volatile oil, a soft, acrid resin, some gum, various salts, and the usual vegetable constituents. It is stimulating; somewhat acrid, aromatic, and stomachic. It is given in doses of ten grains to a scruple, or more.

## GINGER LOZENGES.

R. Powdered ginger	two ounces
sugar	fourteen ounces
Gum tragacanth	
Water, each,	twelve drachms
Mix well, and form lozenges of sixteen grains, each.	
	<i>Soubeiran.</i>

## SYRUP OF GINGER.

R. Ginger, sliced,	three ounces
Boiling water	three pints
Infuse for four hours, strain, and add double its weight of sugar, and form syrup.	
	<i>Guibourt.</i>

## GINGER BEER.

R. Sugar	three pounds
Bruised ginger	two ounces
Cream of tartar	one ounce
Lemons, sliced,	four



Yeast eight fl. ounces  
Boiling water four gallons  
Pour the water on the four first-named ingredients, infuse for two hours, strain, add the yeast, and, when fermentation has commenced, pour into strong bottles, and wire down the corks. *Redwood.*

R. White sugar twenty pounds  
Lemon juice eighteen fl. ounces  
Honey one pound  
Bruised ginger seventeen ounces  
Water eighteen gallons

Boil the ginger in three gallons of the water for half an hour; add the sugar, lemon juice, and honey, with the remainder of the water, and strain; when cold, add the white of an egg, and half a fl. ounce of essence of lemon; let rest for four days, and bottle. *Pereira.*

## OIL OF GINGER.

R. Ginger one part  
Water six parts  
Digest for five or six days; distil, and collect the oil.

## ETHEREAL EXTRACT OF GINGER.

R. Ginger four ounces  
Sulphuric ether six ounces  
Treat the powdered ginger with the ether, in a displacement apparatus, and evaporate the product by means of a water-bath; one part corresponds to sixteen of ginger. *Beral.*

## TINCTURE OF ETHEREAL EXTRACT OF GINGER.

R. Ethereal extract of ginger one part  
Alcohol (.825) fifteen parts  
Dissolve. *Beral.*

## SYRUP OF ETHEREAL EXTRACT OF GINGER.

R. Ethereal extract of ginger one scruple  
Syrup of sugar one ounce  
Dissolve, and strain, after twenty-four hours' digestion. *Beral.*

## ESSENCE OF GINGER.

R. Jamaica ginger four ounces  
French brandy one pint  
Powder the ginger coarsely, moisten it with the brandy, put it in a displacer, and pour the remainder of the brandy on it; when it ceases to pass, add diluted alcohol, till one pint has passed in all.

## SPICE PLASTER.

R. Powdered ginger two ounces  
cloves  
cinnamon, each, one ounce  
red pepper two drachms  
Tincture of ginger half an ounce  
Honey sufficient  
Mix the powders; add the tincture, and sufficient honey to make of proper consistence for a stiff cataplasm. *W. Procter.*



## DIETETIC PREPARATIONS.

NOT INCLUDED AMONG THE PREVIOUS PRESCRIPTIONS.

### TOUS LES MOIS.

This is a variety of arrowroot, prepared from the rhizomes of a species of *Canna*, and is to be employed in the same manner, and in same cases, as arrowroot.

### MUCILAGE OF SALEP.

Salep is the prepared bulbs of several species of orchidaceous plants. It is more nutritive than arrowroot or sago, and hence is adapted for the convalescent, rather than for the sick. These roots are powdered with much difficulty. The mucilage is prepared by dissolving the powder in boiling water, constantly stirring, and adding to the solution sugar and milk. *Soubciran* orders powdered salep, half an ounce, to be boiled with a sufficient quantity of water till dissolved, four ounces of sugar added, and to be aromatized according to taste.

### BISCUIT JELLY.

White biscuit, four ounces, to be boiled down in four pints of water, to one-half, strained, and evaporated to one pint; one pound of white sugar, four ounces of port wine, and one drachm of cinnamon water added, and the whole well mixed. It has been found useful in debility of the digestive organs.

*A. T. Thomson.*

### BREAD JELLY.

Cut a French roll into slices, toast these on both sides, and boil in a quart of water, until the whole forms a jelly, adding more water if required; strain, and flavor. This is nutritious, and may be made more so by using broth, wholly deprived of fat, instead of water.

*Cooley.*

### PANADA.

Stale wheat bread, one ounce; cinnamon, one drachm; water, one pint; cover up and let stand for an hour, beat up and boil for ten minutes, adding a little grated nutmeg

and sugar. Wine may be added, if required.

*Ellis.*

### COMPOUND SALEP POWDERS.

Powdered salep, tragacanth, and sago, each, four ounces; cochineal, half a drachm; prepared oyster shells, one ounce. These are to be carefully mixed and divided into powders, of one drachm each. Stir one of these into a pint of milk, and boil for ten or fifteen minutes. To be taken freely in diarrhoea and dysentery.

*Ellis.*

These are known as *Castanello's powders*, for which the following is used as a substitute. Powdered gum Arabic, tragacanth, arrowroot, sago, and tapioca, each, two drachms. Mix them well together, boil in a pint of milk, flavored with nutmeg or cinnamon. Used as a diet in dysentery, diarrhoea, &c.

### VEGETABLE BROTH.

Take two potatoes, one carrot, one turnip, and one onion. Slice them, and boil in a quart of water for an hour, adding more water from time to time, so as to keep up the original quantity; flavor with salt and a small portion of potherbs; strain. When advisable, a small quantity of mushroom catsup added to this broth greatly improves its taste.

A good substitute for animal food, when this last is inadmissible.

*Ellis.*

### GLOUCESTER JELLY.

Rice, sago, pearl barley, hartshorn shavings, eryngium root, each, one ounce; boil in three pints of water to one pint, and strain. This is very nutritive, dissolved in broth, milk, or wine.

*A. T. Thomson.*

### ALMOND JELLY.

Blanched sweet almonds and white sugar, each, one ounce; water, four ounces; make an emulsion; strain, and add melted hartshorn jelly, half a pound; orange-flower



water one drachm; and essence of lemon two or three drops.

*Cooley.*

#### BOILED FLOUR.

Take of fine flour a pound, tie it up in a linen cloth as tight as possible, and after frequently dipping it into cold water, dredge the outside with flour, till a crust is formed round it, which will prevent the water soaking into it, whilst boiling. Boil for a long time, and permit to cool, when it will become a hard, dry mass.

This to be grated and prepared like arrowroot. A good diet for children, in diarrhoea, &c.

*Ellis.*

#### BEEF TEA.

Take of lean beef, cut into shreds, one pound; water, one quart. Boil for twenty minutes, removing any scum that arises. When it has become cold, strain.

*Ellis.*

Take half a pound of good rump stake, cut it into thin slices, and spread these in a hollow dish; sprinkle a little salt over them, and pour upon the whole a pint of boiling water. Cover the dish and place it near the fire for half an hour; then remove to a pan and boil for fifteen minutes; strain through a fine sieve.

The quantity of water is too small for the strength of the tea for invalids, but is sufficient to extract all the soluble parts of the beef, and the tea can be reduced to the proper strength by the addition of boiling water.

*A. T. Thomson.*

#### ESSENCE OF BEEF.

Take of lean beef, sliced, a sufficient quantity to fill the body of a porter bottle, cork up loosely, and place it in a pot of cold water, attaching the neck, by means of a string, to the handle of the pot. Boil for an hour and a-half to two hours, then decant the liquid and skim it.

To this preparation may be added spices, salt, wine, brandy, &c., according to the taste of the patient and nature of the disease.

*Ellis.*

#### MUTTON TEA.

Take one pound of good mutton, freed from the fat, and cut into thin slices; pour over it a pint and a-half of boiling water, in the same manner as directed for beef tea; but it requires to be boiled for half an hour, previous to straining. If the invalid desires the addition of barley, an ounce of clean pearl barley, washed, and macerated in boiling water for an hour, may be boiled

with the mutton tea, and the undissolved barley, separated by straining.

*A. T. Thomson.*

#### VEAL TEA.

This is to be made in the same manner as beef tea, using a pound of the fillet of veal, free from fat and sliced, and a pint and a-half of boiling water; boiling for half an hour. It may also be made with the same quantity of the fleshy part of a knuckle of veal. By boiling this latter down to one-half, and straining, the decoction will gelatinize, and, if poured into small cups, will keep good for several days. By the addition of an equal quantity of boiling water to this jelly, it is fit for use in a few minutes.

*A. T. Thomson.*

#### CHICKEN WATER.

Take half a chicken, divested of all fat, and break the bones; add to this half a gallon of water, boil for half an hour, strain, and season with salt.

*Ellis.*

Take a small chicken, free it from the skin, and from all the fat between the muscles; divide it longitudinally into halves; remove the lungs, liver, &c., then cut it, bones and muscles, into thin slices, put these into a pan with a sufficient quantity of salt; add a quart of boiling water, cover the pan, and simmer slowly for two hours; strain through a fine sieve.

*A. T. Thomson.*

#### CALVES' FOOT JELLY.

Take two calves' feet, and add to them one gallon of water, and boil down to one quart; strain, and when cold, remove all fat; then add the white of six or eight eggs, well beaten; a pint of wine, half a pound of loaf sugar, and the juice of four lemons, and mix well. Boil for a few minutes, constantly stirring; then strain through a flannel bag. The wine may be omitted if necessary.

*Ellis.*

#### TOAST WATER.

Toast thoroughly a slice of stale bread, put it in a jug and pour over it a quart of water, which has been boiled and cooled, and in two hours decant; a small piece of orange or lemon peel put into the jug with the bread, improves the flavor greatly.

This forms a good drink in febrile affections.

*A. T. Thomson.*

#### APPLE WATER.

Slice two large apples, and pour over the



slices a pint of boiling water. Let stand for an hour, and decant; if necessary, sweeten with a little white sugar.

*A. T. Thomson.*

It is also prepared by boiling for an hour ten ounces of sliced apple in two pints of water, and straining.

*Soubeyran.*

#### LEMON-PEEL WATER.

Pare the rind of one lemon, which has been previously rubbed with half an ounce of refined loaf sugar, put the peeling and sugar into a jar, and pour over them a quart of boiling water. When cold, pour off the fluid, and add a tablespoonful of lemon juice. If wine be not improper, a glass of sherry may be added, instead of the lemon juice.

*A. T. Thomson.*

#### BRAN TEA.

Fresh wheat bran one pound, water three quarts. Boil down to one quart, strain, and add sugar, honey, or molasses, according to the taste of the patient.

#### CHICKEN PANADA.

Take the white meat of a boiled or roasted chicken, free it from the skin, and cut it into small morsels; pound these in a mortar with an equal quantity of stale bread, and a sufficiency of salt, adding gradually some of the water in which the chicken has been boiled, or some beef tea, until the whole forms a thin, fluid paste; put this into a pan, and boil for ten minutes, continually stirring.

*A. T. Thomson.*

#### MUTTON BROTH.

Take a pound of mutton, freed from fat, put it into a pan with three pints of water, and simmer for two hours. Take three carrots, the same number of turnips, peel, and cut them in slices, boil them for half an hour in a quart of water, throw them on a colander to drain off the water, and having boiled two onions sliced in a pint of water, and poured off the water, add the whole of these vegetables to the mutton liquor; after removing the meat, season with salt and a little celery seed. Simmer slowly for four hours, put in the meat again, and continue to simmer for another hour. The meat may be served with the broth. A palatable and very nutritive dish for convalescents.

*A. T. Thomson.*

#### FOWL WITH RICE.

Free a young fowl from the skin and the fat on the exterior of the body, and sim-

mer it in good beef tea, till it is very tender; season with salt, and, having boiled some rice, add it to the liquor before the fowl is dished.

*A. T. Thomson.*

#### WATER SOUCHY.

Take two small, fresh flounders, boil them in a quart of water to one-third, or long enough to reduce the fish to a pulp; strain the liquor through a sieve, and, having cut off the fins of four other small flounders, put the latter into the liquor, with a sufficiency of salt, a few grains of cayenne pepper, and a small quantity of chopped parsley, and boil till the fish is perfectly done. The fish and liquor are to be eaten together.

Few dishes are more relished by convalescents from fever.

*A. T. Thomson.*

Scale, gut, and wash two perch; put salt in the water; when it boils, put in the fish, with an onion cut into slices and separated into rings, a handful of parsley, picked and washed clean; when the fish are done, put them in a soup-dish, and pour the liquor over them.

*Hofland.*

#### MULLED WINE.

Take a quarter of an ounce of bruised cinnamon, half a nutmeg, grated, and ten bruised cloves; infuse them in half a pint of boiling water for an hour, strain, and add half an ounce of white sugar; pour the whole into a pint of hot port or sherry wine.

A good cordial and restorative in the low stages of fever, or in the debility of convalescence from fevers.

*A. T. Thomson.*

#### COLD CUSTARD.

Take the yolk and white of an egg, and a tablespoonful of sugar; beat together, till the tenacity of the white of the egg is entirely destroyed; add gradually, constantly stirring, a half a pint of cold water, two teaspoonfuls of rose water, and a little grated nutmeg. A wineglassful to be taken every two or three hours.

*Devees.*

#### MOLASSES POSSET.

Put in a saucepan a pint of best molasses, a teaspoonful of powdered white ginger, and a quarter of a pound of fresh butter. Simmer for half an hour, stirring frequently. Then stir in the juice of two lemons, or two tablespoonfuls of vinegar, cover the pan, and let it stand by the fire five minutes.

*A. T. Thomson.*

#### CHICKEN JELLY.

Cut a chicken into small pieces, bruise the bones, and put the whole into a stone



jar, with a cover that fits water tight. Set the jar in a large kettle of boiling water, and keep it boiling for three hours. Then strain, and season with salt, pepper, and mace, or with loaf sugar and lemon juice, according to circumstances and taste.

*A. T. Thomson.*

#### CHOCOLATE.

Put milk and water on to boil; then scrape the chocolate fine, one or two squares to a pint, as will best suit the stomach; when the milk and water boils, take it off the fire, throw the chocolate into it, mill it well, and serve it up with the froth. The sugar may be mixed with the scraped chocolate, or added afterwards. It should never be made before it is wanted, as heating it again injures the flavor, and causes a separation of the oil. *Cooley.*

#### CHOCOLATE MILK.

Dissolve an ounce of scraped chocolate in a pint of boiling new milk. *Cooley.*

#### COFFEE MILK.

Boil a dessertspoonful of ground coffee in a pint of milk, for a quarter of an hour; then clear it with white of egg or isinglass, let it boil for a few minutes, and set it by the side of the fire to fine. Sweeten according to taste.

This is a suitable breakfast for those of spare habit, and disposed to affections of the lungs. *Cooley.*

#### RICE CUSTARD.

Boil half a cupful of the best ground rice in a pint of milk, until dissolved; then mix it with a quart of cream; flavor with nutmeg, mace, and a little brandy.

*Cooley.*

#### FRUMENTY.

Bruised wheat, boiled in water until quite soft; drain, thin with milk, sweeten, and flavor according to taste.

The bruised wheat boiled to a paste, and, when cold, eaten with milk, in the evening, for some time, will often relieve costiveness.

#### GINGER BEER.

Take of white sugar, three pounds; bruised ginger, two ounces; cream of tartar, one ounce; lemons, sliced, four; boiling water, four gallons; yeast, eight ounces. Pour the water on the four first-named ingredients, and infuse for two hours; then strain, add the yeast, and, when fermenta-

tion has continued for some hours, put into stone bottles and tie down the corks.

*Gray.*

#### SPRUCE BEER.

Take of sugar, six pounds; essence of spruce, four ounces; boiling water, ten gallons; yeast, eight ounces. Add the water to the sugar and essence, ferment with the yeast, and bottle in the same way as ginger beer.

*Gray.*

#### MOLASSES BEER.

Take of molasses, fourteen pounds; hops, a pound and a-half; water, thirty-six gallons; yeast, a pound. Boil the hops in the water, add the molasses, and ferment.

*Gray.*

#### CAUDLE.

Into a pint of thin gruel, put, while it is boiling hot, the yolk of an egg, beaten with sugar, and mixed with a tablespoonful of cold water, a glass of wine, and some nutmeg; mix well together.

A nourishing, restorative mixture, given during convalescence. *Gray.*

#### BARLEY MIXTURE.

Take of pearl barley, sliced figs, stoned raisins, of each, two ounces and a-half; liquorice root, sliced and bruised, five drachms; water, five pints and a-half. Clean the barley by washing, boil it in four and a-half pints of the water, down to two pints; add the figs, raisins, and liquorice root, with the remaining pint of water; boil down to two pints, and strain.

This is nutritive and demulcent. *Gray.*

#### ARTIFICIAL ASSES' MILK.

Boil together a pint of water and an ounce of hartshorn shavings, until reduced to a jelly; add two ounces of white sugar; when cold, mix with a pint of new milk, and a teaspoonful of syrup of Tolu. To be taken freely, as a nutritive beverage.

*Cooley.*

#### ISINGLASS BLANCMANGE.

Isinglass, a quarter of a pound; rose water, half a pint; milk, two quarts; milk of almonds, half a pint. Boil to a proper consistence, and permit to cool.

*Hoffmann.*

#### RICE BLANCMANGE.

Ground rice, two ounces; milk, one pint; lump sugar, three ounces; a little lemon peel and cinnamon; dissolve the rice in



the milk, by boiling, reduce it to a proper consistence, then add the spice and sugar; boil for a few minutes, strain, and let cool. The rice should be rubbed up with a little water, before adding it to the milk, to prevent it from being in lumps.

*Cooley.*

#### SODA CAKES.

Flour, one pound; bicarbonate of soda,

a quarter of an ounce; sugar and butter, each, half a pound; make a paste with milk; add candied orange, lemon, or citron, or the fresh peel, grated, according to taste. They may be made with the same quantity of carbonate of magnesia, instead of the soda, and the candied peel omitted. In dyspepsia, with acidity.

*Cooley.*



## LIST OF INCOMPATIBLES.

---

- |   |   |
|---|---|
| <p><b>ABSINTHIUM</b>, with sulphates of iron and zinc, acetates of lead, nitrate of silver, tartar emetic.</p> <p><b>ACACIA</b>, with Goulard's extract, alcohol, nitric acid, muriated tincture of iron.</p> <p><b>ACIDUM ACETICUM</b>, with alkalies, alkaline and earthy carbonates.</p> <p><b>ACIDUM CITRICUM</b>, with mineral acids, acetates of lead, nitrate and acetate of mercury, alkalies, alkaline sulphurets.</p> <p><b>ACIDUM ARSENIOSUM</b>, with lime water, hydrosulphate of potassa, astringent vegetable infusions and decoctions.</p> <p><b>ACIDUM GALLICUM</b>, with lime water, carbonate of potassa, acetate of lead, sulphate of copper, nitrate of silver, iodide of iron, sulphate of iron, tartar emetic, nitrate of silver, solution of isinglass, solution of opium, &amp;c.</p> <p><b>ACIDUM HYDROCYANICUM</b>, with mineral acids, metallic oxides, chlorine, &amp;c.</p> <p><b>ACIDUM MURIATICUM</b>, with alkalies and their carbonates, alkaline earths, metallic oxides, sulphuret of potassium, tartrate of potassa, and most metallic salts, especially those of silver.</p> <p><b>ACIDUM NITRICUM</b>, with the metallic oxides, the salifiable bases, the essential oils, &amp;c.</p> <p><b>ACIDUM NITRO MURIATICUM</b>, with oxides, earths and alkalies, the sulphurets, &amp;c.</p> <p><b>ACIDUM PHOSPHORICUM</b>, with the soluble salts of lime, baryta, and lead.</p> <p><b>ACIDUM SULPHURICUM</b>, with the earths, alkalies, and their carbonates, the sulphurets, &amp;c.</p> <p><b>ACIDUM TANNICUM</b>, per salts of iron, albumen, gelatin, &amp;c.</p> <p><b>ACIDUM TARTARICUM</b>, with alkalies and their carbonates, and the salts of potassa.</p> <p><b>ÆTHER HYPONITROSUS</b>, with alcoholic solution of caustic potassa.</p> | <p><b>ÆTHER MURIATICUS</b>, with solution of caustic potassa.</p> <p><b>ALUMINÆ ET POTASSÆ SULPHAS</b>, with the alkalies and alkaline carbonates, lime, magnesia, acetate of lead, infusion of galls, &amp;c.</p> <p><b>AMMONIA</b>, with acids, mineral salts, alum, &amp;c.</p> <p><b>AMMONIÆ ACETAS</b>, with alkalies, strong acids, corrosive sublimate, nitrate of silver, &amp;c.</p> <p><b>AMMONIÆ CARBONAS</b>, with acids, caustic potassa, magnesia, alum, chloride of calcium, bitartrate and bisulphate of potassa, the salts of iron, bichloride of mercury, salts of lead, sulphate of zinc, &amp;c.</p> <p><b>AMMONIÆ MURIAS</b>, with sulphuric and nitric acids, salts of lead and silver, potassa, the carbonates of potassa and soda, lime, &amp;c.</p> <p><b>ANGUSTURA</b>, with sulphates of iron and copper, nitrate of silver, tartar emetic, acetates of lead, bichloride of mercury, potassa, infusion of galls, &amp;c.</p> <p><b>ANTHEMIS</b>, with solution of gelatin, infusions containing gallic acid, salts of iron, nitrate of silver, salts of lead, bichloride of mercury, &amp;c.</p> <p><b>ANTIMONII SULPHURETUM</b>, with nitric and nitro-muriatic acids.</p> <p><b>ANTIMONII ET POTASSÆ TARTRAS</b>, with alkalies and earths, and their carbonates, strong acids, hydrosulphurets, lime water, chloride of calcium, salts of lead, soaps, infusions containing gallic acid, rhubarb, &amp;c.</p> <p><b>ARMORACIA</b>, with carbonates of the alkalies, bichloride of mercury, nitrate of silver, vegetable bitters and astringents, &amp;c.</p> <p><b>ARGENTI NITRAS</b>, with the fixed alkalies, sulphuric, muriatic, and arsenious acids, and their salts, lime, the chlorides and sulphurets, astringent vege-</p> |
|---|---|



- table infusions, solutions of the salts of mercury and copper, &c.
- ARNICA, with sulphates of iron and zinc, acetate of lead, mineral acids, &c.
- AURANTII CORTEX, with infusion of bark, sulphate of iron, lime water, &c.
- BARYTÆ MURIAS, with alum, nitrate of silver, &c.
- BENZOINUM, with the acids and alkalies.
- BISTORTA, with salts of iron, gelatin, &c.
- BELLADONNA, with caustic alkaline solutions.
- BISMUTHI NITRAS, with the alkalies, vegetable astringents.
- CALAMUS, with acetate of lead.
- CALCH CHLORIDUM LIQUOR, with carbonates of soda and potassa, and carbonate of magnesia, &c.
- CALCIS LIQUOR, with the mineral and acetic, phosphoric, tartaric, and citric acids, muriate of ammonia, the alkaline carbonates, soap, the vegetable astringents, alum, sulphates of iron and zinc, sulphate of magnesia, chlorides of mercury, nitrate of silver.
- CALCIS CARBONAS, with the acids and acidulous salts, alum, muriate of ammonia.
- CAPSICUM, with corrosive sublimate, acetate of lead, nitrate of silver, sulphates of iron, zinc, or copper, carbonates of the alkalies.
- CARDAMOMUM, with the acids, sulphate of iron, bichloride of mercury, &c.
- CARYOPHYLLUS, with tartar emetic, sulphates of iron and zinc, &c.
- CASCARILLA, with lime water, sulphates of iron and zinc, infusions containing gallic acid.
- CASSIA FISTULÆ, with alcohol.
- CATECHU, with alkalies, the salts of iron, gelatin, &c.
- CINCHONA, with strong acids, alkalies, sulphates of iron and zinc, nitrate of silver, tartar emetic, lime, magnesia, &c.
- COCCUS, with acetate of lead, sulphate of zinc and iron.
- COLCHICUM, with acids which render the vinous tincture drastic; alkalies, on the contrary, render it milder in its operation.
- COLOCYNTHIS, with fixed alkalies, sulphate of iron, nitrate of silver, acetate of lead, &c.
- COLUMBA, with ammonia, lime water, mineral acids, muriate of iron, nitrate of silver, acetate of lead, isinglass.
- CONIUM, with the acids.
- CONTRAYERVA (tincture), with water.
- COPAIBA, with the mineral acids.
- COPTIS TRIFOLIATA, with nitrate of silver, acetate of lead.
- CRETA PRÆPARATA, with acids and acidulous salts, alum, muriate of ammonia.
- CUPRUM AMMONIATUM, with acids, potassa and soda, lime water.
- CUPRI SULPHAS, with fixed alkalies, ammonia, and its carbonate, bichloride of mercury, arsenite of potassa, nitrate of silver, acetate of lead, vegetable infusion, &c.
- CYDONIA, with acids, most metallic salts, alcohol.
- DIGITALIS, with sulphate of iron, infusion of Peruvian bark, acetate of lead.
- FERRUM AMMONIATUM, with acids, the fixed alkalies, lime water, astringent infusions.
- FERRI IODIDUM, with fixed alkalies, lime water, vegetable astringents, &c.
- FERRI ET POTASSÆ TARTRAS, with the mineral acids, alkalies, vegetable astringents.
- FERRI SUBCARBONAS, with the mineral acids, acidulous salt, &c.
- FERRI MURIATIS TINCTURA, with alkalies and their carbonates, lime water, carbonate of lime, magnesia, and its carbonate, solution of gum, vegetable astringents.
- FERRI SULPHAS, with nitric acid, fixed alkalies, and their carbonates, lime water, nitrate of potassa, borate of soda, acetate of lead, iodide of potassium, vegetable astringents, &c.
- GALLA, with the carbonates of the alkalies, lime water, sulphates of iron and zinc, acetate of lead, bichloride of mercury, gelatin, &c.
- GUAIACI TINCTURE, with water, the mineral acids, spirit of nitric ether, earthy and metallic salts, &c.



- GRANATUM (*Cortex*), with sulphate of iron, &c.
- HÆMATOXYLON, with mineral acids, solution of alum, sulphate of iron and copper, tartar emetic, acetate of lead.
- HYDRARGYRUM AMMONIATUM, with muriatic acid, the fixed alkalies.
- HYDRARGYRI CHLORIDUM MITE, with the alkalies, lime, muriate of ammonia, chlorides of potassium and sodium, carbonates of the alkalies, nitric acid, salts of iron, lead, and copper, iodide of potassium, soaps, &c.
- HYDRARGYRI CHLORIDUM CORROSIVUM, with alkalies and their carbonates, carbonates of lime, tartar emetic, sulphuret of potassium, soaps, iron, copper, lead, and their salts, nitrate of silver, albumen, gelatin, vegetable astringents, fixed oils, &c.
- HYDRARGYRI OXIDUM RUBRUM, with the mineral acids.
- HYDRARGYRI OXIDUM NIGRUM, with the mineral and acetic acids.
- HYDRARGYRI IODIDUM, with the mineral acids, iodide of potassium, chloride of sodium.
- HYDRARGYRI IODIDUM RUBRUM, with the mineral acids, iodide of potassium, chloride of sodium.
- HYDRARGYRUM CUM CRETA, with the mineral and acetic acids, alum, &c.
- HYOSCYAMUS, with acetate of lead, nitrate of silver, sulphate of iron.
- IODINE.
- IPECACUANHA, with the vegetable acids, the vegetable astringents, acetate of lead, &c.
- KINO, with the salts of iron, the mineral acids, tartar emetic, &c.
- KRAMERIA, with salts of iron, gelatin, mineral acids, &c.
- LUPULUS, with the salts of iron.
- MAGNESIA, with acids and acidulous salts, muriate of ammonia, metallic salts.
- MAGNESIÆ CARBONAS, with acids and acidulous salts, metallic salts, lime water, muriate of ammonia, bitartrate of potassa.
- MAGNESIÆ SULPHAS, with ammonia, acetate of lead, chloride of calcium, lime water, potassa and soda, and their carbonates, &c.
- MENTHA, with sulphate of iron, nitrate of silver, acetate of lead, &c.
- MORPHIÆ ACETAS, with alkaline carbonates, ammonia, all articles incompatible with infusion of opium, except acetate of lead.
- MORPHIÆ MURIAS, with acetate of lead, ammonia, alkaline carbonates, &c.
- MOSCHUS, with mineral acids, bichloride of mercury, sulphate of iron, nitrate of silver, infusion of bark, &c.
- OPIUM, with ammonia, carbonates of potassa and soda, nitrate of silver, acetate of lead, salts of copper, iron, and zinc, infusion of galls, &c.
- POTASSA, with acids and acidulous salts, earthy and metallic salts, ammonia and its salts, &c.
- POTASSÆ ACETAS, with mineral acids, sulphates of soda and magnesia, tartaric acid, &c.
- POTASSÆ ARSENITIS LIQUOR, with lime water, acids, chloride of calcium, sulphate of magnesia, alum, sulphate of iron, iodide of iron, nitrate of silver, vegetable astringents.
- POTASSÆ BITARTRAS, with lime water, ammonia, carbonates of potassa and soda, magnesia, sulphate of magnesia, &c.
- POTASSÆ CARBONAS, with acids, metallic salts, lime water, sulphate of magnesia, muriate of ammonia, &c.
- POTASSIÆ CITRAS, with salts of lime, lead, and silver, sulphuric acid.
- POTASSÆ NITRAS, with alum, sulphate of magnesia, metallic sulphates, sulphuric acid, muriatic acid (when heated).
- POTASSÆ SULPHAS, with tartaric acid, acetate of lead, nitric and muriatic acids, chloride of calcium, bichloride of mercury, nitrate of silver, sulphate of magnesia.
- POTASSÆ SULPHURETUM, with acids, most metallic salts.
- POTASSÆ TARTRAS, with all acids, lime water, acetate of lead, nitrate of silver.



- POTASSII IODIDUM, with acetate of lead, bichloride of mercury, tartaric acid, metallic salts, all acids and acidulous salts, except bitartrate of potassa.
- PIMENTA, with alum, ammonia, alkaline carbonates, salts of iron, copper, zinc, and silver, vegetable astringents.
- PIPER NIGRUM, with vegetable astringents.
- PLUMBI ACETAS, with the alkalies and their carbonates, tartaric, citric, and sulphuric acids, tartrate and bitartrate of potassa, alkaline and metallic sulphates, alum, borax, lime water, vegetable astringents, infusion of opium, the soaps, milk.
- PLUMBI IODIDUM, with solution of potassa, sulphuric acid.
- QUASSIA, with nitrate of silver, acetate of lead, &c.
- QUINÆ SULPHAS, with tartaric acid, tartrate of potassa, alkalies and their carbonates, lime water, infusion of galls.
- RHEUM, with strong acids, lime water, sulphates of iron and zinc, tartar emetic, bichloride of mercury, vegetable astringents.
- ROSA GALLICA, with sulphates of iron and zinc, gelatin, lime water.
- SALIX, with lime water, sulphate of iron, alkaline carbonates, solution of isinglass.
- SALVIA, with salts of iron.
- SAPO (*and liniments, &c., containing it*), with sulphates of lime and magnesia, chloride of calcium, alum, metallic salts, lime water.
- SARSAPARILLA, with infusion of galls, lime water, acetate of lead.
- SCILLA, with lime water, alkaline carbonates, nitrate of silver, acetate of lead, &c.
- SENNA, with strong acids, carbonates of the alkalies, lime water, tartar emetic.
- SERPENTARIA, with acetate of lead.
- SODÆ CARBONAS, with acids, bitartrate of potassa, metallic and earthy salts, lime water.
- SODÆ PHOSPHAS, with mineral acids, lime, magnesia, &c.
- SODÆ SULPHAS, with salts precipitated by sulphuric acid, carbonates of potassa, acetate of potassa.
- SODÆ ET POTASSÆ TARTRAS, with most acids and acidulous salts, acetate of lead.
- SPIRITUS ÆTHERIS NITRICI, with sulphate of iron, tincture of guaiacum, alkaline and earthy carbonates, especially those with an excess of acid.
- TAMARINDUS, with the salts of potassa, the alkaline carbonates, lime water, tartar emetic.
- TARAXACUM, with corrosive sublimate, sulphate of iron, nitrate of silver, acetate of lead, infusion of galls, &c.
- THEA, with salts of iron, gelatin, lime water, &c.
- TORMENTILLA, with solution of isinglass, salts of iron, alkalies, &c.
- TRAGACANTH, with sulphates of iron and copper, acetate of lead, &c.
- ULMUS, with alcohol, tinctures, if added in quantity.
- UVA URSI, with salts of iron.
- VALERIANA, with salts of iron.
- ZINCI OXIDUM, with potassa, soda, and ammonia.
- ZINCI SULPHAS, with potassa, soda, and ammonia, and their carbonates, the hydrosulphurets, milk, mucilages, &c.



# POSOLOGICAL TABLE

OF THE

## MOST IMPORTANT MEDICINES.

Absinthium, Infusion, f̄j to ij.	Ammonia, Alcohol, gtt. v to xx.
Extract, grs. x to xx.	aromatic, gtt. x to xx.
Tincture, gtt. xx to f̄ij.	Citrate, f̄ss to j.
Oil, gtt. ij to iv.	Hydrosulphate, gtt. v to vj.
Acetosella, Extract, ̄j to 3ss.	Muriate, gr. j to v.
Acidum Arseniosum, gr. $\frac{1}{16}$ to $\frac{1}{8}$ .	Nitrate, ̄j to ij.
Benzoicum, grs. x to xx.	Phosphate, f̄ss.
Boracicum, grs. v to xx.	Succinate, Spirit, gtt. xx to l.
Gallicum, grs. ij to x.	Sulphate, ̄j. to 3ss.
Hydrocyanicum (medicinal), gtt. j to ii.	Ammoniacum, grs. v to xv.
Hydriodicum, gtt. v to lx.	mixture, f̄ss to j.
Muriaticum, gtt. v to x.	and Nit. acid, f̄j to iv.
diluted, gtt. xx to xxx.	Amylum, Iodide, 3ss to ij.
Nitricum, gtt. v to x.	Angelica, Tincture, f̄j to ij.
diluted, gtt. xx to xxx.	Angustura, Powder, grs. x to ̄j.
Phosphoricum, diluted, gtt. x to f̄j.	Infusion, f̄ij.
Sulphuricum, gtt. ij to v.	Tincture, f̄j to ij.
diluted, gtt. x to xxx.	Anthemis, Infusion, f̄j to ij.
aromatic, gtt. v to x.	Extract, grs. x to xx.
Tannicum, gr. j to iij.	Syrup, f̄ss.
Tartaricum, 3j to ij.	Anthracokali, grs. ij to iv.
Aconitum, Powder, gr. j to ij.	Antimonium, grs. x to xx.
Extract, gr. j to ij.	Sulphuret, grs. v to x.
alcoholic, gr. $\frac{1}{8}$ to $\frac{1}{4}$ .	precipitated, grs. j to iij.
Tincture, gtt. v to vj.	Kermes mineral, gr. $\frac{1}{2}$ to x.
Æther, Acetic, gtt. x to f̄j.	and Lime, grs. j to v.
Hyponitrosus, gtt. x to lx.	Tartrate and Potassa, gr. $\frac{1}{2}$ to iij.
spirit of, f̄ss to ij.	Wine, gtt. xxx to 3j.
Hydrocyanic, gtt. ij to vj.	Antimonial Powder, grs. iij to x.
Muriatic, gtt. xxx to lx.	Apocynum, Decoction, f̄j to ij.
Sulphuric, f̄ss to j.	Extract, grs. iij to v.
spirit, f̄j to iij.	Argentum, Chloride, gr. $\frac{1}{10}$ to gr. ij.
compound, f̄ss to ij.	and Ammonia, gr. $\frac{1}{4}$ to $\frac{1}{10}$ .
Allium, Syrup, f̄j.	Cyanide, gr. $\frac{1}{12}$ to $\frac{1}{8}$ .
Aloe, Powder, gr. j to v.	Iodide, $\frac{1}{2}$ to j.
and Canella, grs. v to xx.	Nitrate, gr. $\frac{1}{4}$ to ij.
Wine, f̄j. to 3j.	Oxide, gr. $\frac{1}{4}$ to ij.
Tincture, f̄j. to 3j.	Armoracia, Infusion, f̄j to ij.
and Myrrh, f̄ss to jj.	Spirit. comp. f̄j to iv.
Aluminæ et Potassæ Sulphas, grs. v to x.	Arnica, Powder, grs. v to x.
Ammonia, Water, gtt. v to xx.	Extract, grs. v to ̄j.
Acetate, f̄ss to j.	Infusion, f̄ss to j.
Arseniate, gtt. x to xx.	Tincture, gtt. xxx to l.
Carbonate, grs. v to x.	
tincture comp., gtt. x to xl.	



Arsenicum, Iodide, gr. $\frac{1}{10}$ to $\frac{1}{5}$ .	Cannabis, Resin, gr. $\frac{1}{4}$ to gr. j.
Donovan's Solution, gtt. xv to f3ss.	Cantharis, gr. ss to j.
Arum, Powder, grs. x.	Tincture, gtt. xx to f3j.
Asclepias Tuberosa, Infusion, f3iij to iv.	Capsicum, grs. v to x.
Asparagus, Extract, ʒj to 3j.	Infusion, f3ss.
Syrup, 3j to ij.	Tincture, f3j to ij.
Assafetida, grs. ij to x.	Carbo Animalis, grs. x to xx.
Mixture, f3j to ij.	Ligni, 3ss to j.
Tincture, f3j.	Mineralis, Anthracokali, grs. ij to iv.
ammoniated, gtt. v to f3j.	Cardamomum, Tincture, f3j to ij.
Aurum, gr. $\frac{1}{4}$ to j.	Carota, Infusion, seeds, f3ij to iv.
Chloride, gr. $\frac{1}{20}$ to $\frac{1}{10}$ .	Carum, Spirit, f3j to ij.
and Sodium, gr. $\frac{1}{20}$ to $\frac{1}{10}$ .	Caryophyllus, Powder, grs. v to x.
Cyanide, gr. $\frac{1}{18}$ to $\frac{1}{10}$ .	Infusion, f3j to ij.
Iodide, gr. $\frac{1}{20}$ to $\frac{1}{10}$ .	Tincture, f3j to ij.
Oxide, gr. $\frac{1}{10}$ to $\frac{3}{4}$ .	Oil, gtt. ij to v.
Azederach, Decoction, f3ss to j.	Cascarilla, Powder, ʒj to 3ss.
Ballotta Lanata, Decoction, f3iij to iv.	Infusion, f3j to ij.
Baptisia Tinctoria, Decoction, f3ss.	Extract, grs. x to 3j.
Barium, Chloride, grs. ij to ij.	Tincture, f3j.
Solution, gtt. v.	Cassia Fistula, 3ij to 3ss.
Iodide, gr. $\frac{1}{8}$ .	Costoreum, grs. v to xx.
Baryta, Solution, gtt. iv to vj.	Tincture, gtt. xxx to f3ij.
Acetate, gtt. iv to vj.	Catalpa, Decoction, f3ij to iij.
Bebeerina, Sulphate, grs. v to ʒj.	Catechu, grs. x to ʒj.
Belladonna, gr. j to ij.	Infusion, comp. f3j to iij.
Extract, gr. $\frac{1}{2}$ to j.	Tincture, gtt. xxx to f3iij.
alcoholic, gr. $\frac{1}{2}$ to j.	Centaurea Benedicta, ʒj to 3j.
Syrup, 3j to ij.	Infusion, f3j to ij.
Tincture, gtt. xv to xxx.	Cetraria, 3ss to j.
Benzoin, Tincture, gtt. x to xx.	Cetrarine, grs. ij to v.
Bismuthum, Subnitrate, grs. ij to x.	Chelidonium, Extract, grs. v to xv.
Bistort, Infusion, f3j to ij.	Chenopodium, Juice, f3ss.
Boletus Laricis, gr. j to iij.	Oil, gtt. iv to x.
Brominum, Solution, gtt. v to x.	Chimaphila, Decoction, f3ij to iv.
Brucia, gr. $\frac{1}{8}$ to $\frac{1}{2}$ .	Extract, grs. x to xxx.
Tincture, gtt. v to xx.	Chiretta, ʒj.
Calamus, Infusion, f3j to ij.	Chlorinum, Water, f3j to iv.
Calcium, Chloride, Solution, gtt. xx to xxx.	Chloroformum, gtt. v to xx.
Iodide, grs. ij to iij.	Cimicifuga, Decoction, f3j to ij.
Sulphuret, grs. v to x.	Tincture, gtt. xx.
Calx. Lime water, f3ss to ij.	Cinchona, Powder, 3ss to j.
Calcis Carbonas, Powder, grs. x to xx.	Extract, grs. x to xxx.
comp. grs. x to xx.	Decoction, f3j.
Phosphate, ʒj to ij.	Infusion, f3ij.
Calendula, Extract, grs. ij to v.	Tincture, f3j to iv.
Calomel, grs. ij to x.	Cinchonia, grs. ij to iv.
Calotropis, Infusion, 3j to 3j.	Sulphate, gr. j to ij.
Powder, grs. iij to xxx.	Cinnamomum, Powder, grs. x to ʒj.
Camphora, grs. iij to x.	comp. grs. x to xxx.
Water, f3ss to j.	Tincture, f3j to iv.
Tincture, gtt. v to xx.	Codeia, Syrup, f3j.
Canella, grs. x to ʒj.	Muriate and Morphia, gr. $\frac{1}{2}$ to j.
Cannabis, Extract, grs. x.	Colchicum, Powder, grs. ij to viij.
Tincture, gtt. x to 3j.	Extract, bulb, gr. j to ij.
	Wine, root, gtt. x to xx.
	seeds, f3j.
	Tincture, seeds, f3ss to ij.



- Colocyntsis, Powder, gr. j to ij.  
Extract, grs. v to x.  
comp. grs. v to xxx.
- Colomba, Powder, grs. x to xxx.  
Infusion, f3ij.  
Tincture, f3j to iv.  
Extract, grs. j to v.
- Conium, Powder, grs. iij to v.  
Extract, gr. j to ij.  
alcoholic, gr. j to ij.  
Tincture, gtt. xx to f3j.
- Contrayerva, Powder, 3ss.  
Extract, 3j to ij.  
Tincture, gtt. xxx to xl.
- Copaiba, grs. xx to f3j.  
Tincture, gtt. xxx to f3j.  
Oil, gtt. x to xxx.
- Coptis, Powder, grs. x to xxx.  
Infusion, f3ss to ij.  
Tincture, f3j to iij.
- Cornus, Powder, 3j to 3j.
- Creasotum, gtt. j to ij.
- Crocus, grs. x to xxx.
- Cubeba, Powder, grs. x to 3iij.  
Tincture, f3j to ij.  
Oil, gtt. v to x.
- Cupri Acetas, gr. j.  
Sulphas (emetic), gr. j to iij.
- Cuprum Ammoniatum, gr.  $\frac{1}{4}$  to  $\frac{1}{2}$ .
- Delphinium, Tincture, gtt. x to xx.
- Delphniia, gr.  $\frac{1}{4}$  to  $\frac{1}{2}$ .
- Digitalis, Powder, gr.  $\frac{1}{2}$  to j.  
Infusion, f3ss.  
Tincture, gtt. x to xx.
- Digitalina, gr.  $\frac{1}{16}$  to  $\frac{1}{8}$ .
- Diosma, Powder, grs. xx to xxx.  
Infusion, f3j to ij.  
Tincture, f3j to iv.  
Extract, fluid, f3j to ij.
- Diospyros, Infusion, f3j.
- Dulcamara, Powder, grs. xxx to 3j.  
Decoction, f3j to ij.  
Extract, grs. v to x.
- Elaterium (common), gr. j to ij.  
(genuine), gr.  $\frac{1}{8}$  to  $\frac{1}{4}$ .
- Elaterina, Tincture, gtt. xx to xl.
- Emetina, gr.  $\frac{1}{16}$  to  $\frac{1}{4}$ .  
Syrup, f3j.
- Ergota, Powder, grs. x to xxx.  
Infusion, f3j.  
Wine, f3ij to iij.  
Tincture, gtt. xx to f3j.  
Extract, gr. ij.  
Oil, gtt. xx to l.
- Erigeron Annuum, Infusion, f3ij to iv.  
Canadense, Infusion, f3ij to iv.  
Extract, grs. v to x.
- Eupatorium, Powder, grs. xx to xxx.  
Infusion, f3j.
- Euphorbia Corollata, Powder, grs. xv to xx.  
Ipecacuanha, Powder, grs. x.  
to xv.  
Hypericifolia, Infusion, f3ss  
to ij.  
Lathyrus, Oil, gtt. iv to xii.
- Ferrum (prepared), grs. v to xx.  
Ammoniated, grs. iv to xii.  
Tincture, gtt. xl  
to f3j.
- Ferri, Acetate, gtt. x to xxv.  
Tincture, gtt. xxx to f3j.
- Arseniate, gr.  $\frac{1}{16}$  to  $\frac{1}{8}$ .
- Bromidum, grs. ij to v.
- Carbonate, grs. x to 3ij.  
Saccharine, grs. x to 3ss.
- Carburet, grs. v to xv.
- Chloride, gr. j to ij.  
Tincture, gtt. x to xxx.
- Citrate, grs. iv to viij.  
and Quinia, grs. v. to x.
- Ferrocyanide, grs. iij to v.
- Iodide, grs. ij to iij.  
Solution, gtt. xxx to f3j.  
Syrup, f3j.  
Tincture, gtt. xxx to f3j.
- Lactate, grs. ij to iv.
- Malate, grs. v to 3j.
- Nitrate, gtt. vj to xii.
- Persesquinitrate, gtt. x to xii.
- Oxide, Black, grs. v to xx.  
Red, grs. v to xv.
- Phosphate, grs. v to x.
- Sulphate, grs. j to v.  
dried, gr. j to iij.
- Tartrate, grs. v to x.  
Wine, f3j.  
Ammonio, grs. v.
- Tannate, gr. ij to iij.
- Valerianate, grs. j to ij.
- Filix Mas, 3j to ij.  
Extract, grs. xv to xx.
- Frasera, Powder, grs. xxx to 3j.  
Infusion, f3j to ij.
- Fuligo, Tincture, gtt. xxx to f3j.
- Fuligokali, gr. ij to iij.
- Galbanum, grs. x to xx.
- Galla, Powder, grs. v to xxx.  
Infusion, f3j. to ij.  
Tincture, f3j to iij.  
Syrup, f3ss.
- Gambogia, gr. j to vj.  
Solution, alkaline, gtt. xv.
- Gaultheria, Oil, gtt. ij to x.
- Gentiana, Powder, grs. x to xl.



Gentiana, Extract, grs. x to ʒj.  
 Infusion, fʒj.  
 Tincture, comp., fʒj to ij.  
 Geranium, grs. x to xxx.  
 Extract, grs. x to xx.  
 Geoffroya, ʒj to ʒss.  
 Decoction, fʒj to ij.  
 Gillenia, grs. xx to xxx.  
 Granatum, Decoction (rind), fʒj.  
 (bark), fʒij to iv.  
 Gratiola, Wine, fʒj.  
 Guaiacum, Resin, grs. x to ʒj.  
 (Wood, decoction,) fʒj to ij.  
 Oil, gtt. iv to v.  
 Tincture, fʒj to iij.  
 Ammoniated, fʒj to ij.  
 Hæmatoxylon, Infusion, fʒss.  
 Decoction, fʒij to fʒij.  
 Extract, grs. x to xxx.  
 Helleborus, Powder, grs. x to ʒj.  
 Extract, grs. x to xv.  
 Tincture, gtt. xxx to fʒj.  
 Humulus, Infusion, fʒj to ij.  
 Tincture, fʒj to iij.  
 Hydrargyrum,  
 with Antimony, gr. j to iv.  
 Chalk, grs. v to ʒj.  
 Magnesia, grs. v to ʒj.  
 Blue pill, grs. v to x.  
 Acetate, gr. j.  
 Borate, gr. ij.  
 Bromide, gr. j to v.  
 Bibromide, gr.  $\frac{1}{20}$  to  $\frac{1}{10}$ .  
 Chloride, Corrosive, gr.  $\frac{1}{16}$   
 to  $\frac{1}{4}$ .  
 Mild, gr. ij to x.  
 Cyanuret, gr.  $\frac{1}{16}$  to  $\frac{1}{8}$ .  
 Iodide, gr.  $\frac{1}{4}$  to  $\frac{1}{2}$ .  
 Red, gr.  $\frac{1}{16}$  to  $\frac{1}{8}$ .  
 Oxide, Black, gr. j to ij.  
 Red, gr.  $\frac{1}{2}$  to j.  
 Phosphate, gr.  $\frac{1}{2}$  to j.  
 Sulphate, gr.  $\frac{1}{4}$  to ij.  
 Sulphuret, Black, grs. v to xx.  
 Red, grs. x to xv.  
 Tartrate, gr. j to ij.  
 Hippocastanum, Powder, ʒij to iv.  
 Hyoseyamus, Powder, grs. ij to v.  
 Extract, gr. j.  
 Tincture, gtt. xx  
 to fʒss.  
 Indigum, Powder, ʒj to ij.  
 Iodinum, gr.  $\frac{1}{8}$  to j.  
 Tincture, gtt. x to xx.  
 Ipecacuanha, Powder, grs. xv to xxx.  
 comp. grs. v to x.  
 Syrup, fʒj to ij.

Ipecacuanha, Wine, fʒj to fʒj.  
 Jalapa, Powder, grs. x to xxx.  
 comp., ʒss to j.  
 Extract, grs. v to x.  
 alkaline grs. iij to ix.  
 Soap, grs. x to xv.  
 Tincture, fʒj to ij.  
 Juglans Cinerea, Extract, grs. v to xxx.  
 Juniperus, Extract, ʒj to iij.  
 Infusion, fʒij to iv.  
 Oil, gtt. iij to vj.  
 Kino, Powder, grs. x to xx.  
 Tincture, fʒss to ij.  
 Krameria, Powder, grs. xx to xxx.  
 Extract, grs. x to xx.  
 Infusion, fʒi to ij.  
 Syrup, fʒss.  
 Tincture, fʒj to ij.  
 Lactucarium, grs. iij.  
 Tincture, fʒss to ij.  
 Lauro Cerasus, Water, fʒss to j.  
 Liriodendron, Powder, ʒss to ij.  
 Infusion, fʒj to ij.  
 Tincture, fʒj to ij.  
 Lobelia, Powder, grs. v to x.  
 Tincture, fʒss to iv.  
 Lupulina, grs. v to x.  
 Tincture, fʒi to ij.  
 Macis, grs. x to ʒj.  
 Tincture, gtt. xxx to xl.  
 Magnesia, ʒj.  
 Carbonate, ʒj to ʒj.  
 Citrate, Solution, ʒj.  
 Sulphate, ʒj.  
 Sulphuret, Syrup, fʒss.  
 Phosphate, grs. x to xx.  
 Manolia, Powder, ʒj to ij.  
 Mangesii, Carbonate, ʒi to ʒi.  
 Muriate, grs. iv to x.  
 Oxide, grs. ij to iv.  
 Sulphate, ʒj to ij.  
 Manna, ʒj to ʒij.  
 Mannite, ʒij to ʒij.  
 Marrubium, Extract, ʒi to ʒss.  
 Matico, Infusion, fʒj.  
 Tincture, gtt. xxx to fʒj.  
 Syrup, fʒj to ij.  
 Mentha Piperita, Oil, gtt. i to iij.  
 Essence, gtt. x to xx.  
 Mentha Viridis, Infusion, fʒj to ij.  
 Menyanthes, Powder, grs. xx to xxx.  
 Infusion, fʒj to ij.  
 Extract, grs. x to xv.  
 Mezereum, Decoction, fʒj to iv.  
 Monarda, Oil, gtt. j to iij.



- Monesia, grs. ij to x.  
     Syrup, f3ss.  
 Morphia, gr.  $\frac{1}{8}$  to  $\frac{1}{4}$ .  
     Acetate, gr.  $\frac{1}{8}$  to  $\frac{1}{4}$ .  
     Bimeconate, gr.  $\frac{1}{4}$ .  
     Citrate, gtt. v to x.  
     Muriate, gr.  $\frac{1}{8}$  to  $\frac{1}{4}$ .  
     Sulphate, gr.  $\frac{1}{8}$  to  $\frac{1}{4}$ .  
 Moschus, grs. v to x.  
     Tincture, gtt. xxx to f3j.  
 Mucuna, Electuary, 3i to ij.  
 Myroxylon, f3ss.  
     Syrup, f3j to ij.  
 Myrrha, grs. x to xxx.  
  
 Narcotina, grs. iij.  
     Muriate, grs. ij to iv.  
 Nux vomica, Powder, grs. iij to v.  
     Extract, gr.  $\frac{1}{2}$  to ij.  
     Tincture, gtt. v to xx.  
  
 Oleum Animale Empyreumatica, gtt. v to x.  
     Cajuputi, gtt. ij to v.  
     Morrhuae, f3ss to f3ij.  
     Olivæ, f3j.  
     Ricini, f3ss to j.  
     Terebinthinæ, gtt. v to f3i.  
     Tiglii, gtt. i to ij.  
 Opium, gr. j.  
     Vinegar, Black drop, gtt. vii to x.  
     Lancaster " " gtt. vj to x.  
     Houlton's " " gtt. xx.  
     Rousseau's " " gtt. iv.  
     Porter's " " gtt. v to xx.  
     Tincture, gtt. xxv.  
         acetated, gtt. xx.  
     Wine, gtt. xxv.  
     Tincture, Smith's, gtt. iij to v.  
         ammoniated, f3i to ij.  
         Camphorated, f3j to iij.  
         Swediaur's, gtt. v.  
     Elixir, gtt. v to xx.  
 Opoponax, grs. x to xx.  
  
 Papaver, Syrup, f3ij to f3j.  
 Pareira, Powder, grs. xxx to 3j.  
     Infusion, f3j to ij.  
     Decoction, f3j to ij.  
     Extract, grs. x to 3ss.  
     Tincture, gtt. l to lx.  
 Paullinia, grs. x to xx.  
 Petroleum, f3ss to j.  
 Petroselinum, Infusion, Oss.  
 Phloridzina, grs. v to xv.  
 Phosphorus, gr.  $\frac{1}{16}$ .  
 Phytolacca, Powder, grs. x to xxx.  
 Piper Nigrum, Confection, 3j to ij.  
 Piperinum, gr. j to ij.  
 Piscidia Erythrina, Tincture, f3j.  
  
 Pix liquida, 3ss to j.  
 Platini, Bichloridum, gr. j to ij.  
 Plumbi, Acetate, gr. j to iv.  
     Iodide, gr.  $\frac{1}{2}$  to j.  
 Podophyllum, Powder, grs. x to xx.  
     Extract, grs. v to xv.  
 Podophyllin, grs. v to vj.  
 Potassi, Bromide, grs. ij to x.  
     Chloride, 3j to ij.  
     Cyanuret, grs.  $\frac{1}{8}$  to  $\frac{1}{4}$ .  
     Iodide, gr. j to x.  
     Iodo-hydrargyrate, gtt. ij to v.  
     Silica, Solution, f3iij.  
     Sulphate, 3iv to v.  
     Bisulphate, 3j to ij.  
         with Sulphur, 3ss to j.  
     Tartrate, 3j to 3j.  
     Bitartrate, 3j to 3ss.  
         and Ammonia, 3j.  
 Prinos, Powder, grs. xxx to 3j.  
     Decoction, f3ij.  
 Prunus Virginiana, Powder, grs. xxx to 3ij.  
     Infusion, f3ij to iij.  
     Syrup, f3j to f3j.  
  
 Quassia, Infusion, f3j.  
     Extract, grs. ij to v.  
     Tincture, f3j to ij.  
 Quereus, Powder, grs. xxx to 3j.  
     Extract, grs. x to xl.  
     Decoction, f3ij to iij.  
 Quinia, gr. j to ij.  
     Amorphous, gr. j to iv.  
     Acetate, gr. j to ij.  
     Arseniate, gr.  $\frac{1}{8}$ .  
     Citrate, gr. j to ij.  
     Ferrocyanate, gr. j to ij.  
     Chloride and Mercury, gr.  $\frac{1}{2}$ .  
     Kinate, gr. j to ij.  
     Lactate, gr. j to ij.  
     Muriate, gr.  $\frac{1}{2}$  to j.  
     Sulphate, gr. j to ij.  
     Sulpho-tartrate, f3ss to j.  
     Valerianate, gr.  $\frac{1}{2}$  to j.  
  
 Rheum, Powder, grs. x to xxx.  
     Roasted, grs. v to x.  
     Infusion, f3j to ij.  
     Extract, grs. x to xx.  
         Fluid, gtt. xv to f3ss.  
     Syrup, f3j to ij.  
         Aromatic, f3j to ij.  
     Tincture, f3j to ij.  
         and Senna, f3ss to ij.  
     Wine, f3j to iv.  
 Rubia, Powder, 3ss.  
     Decoction, f3iij.  
 Rubus, Decoction, f3j to ij.  
 Ruta, Powder, grs. x to xx.



- Sabadilla, Powder, grs. ij to v.  
Extract, gr.  $\frac{1}{2}$ .
- Sabbatia, Infusion, f $\overline{3}$ ij to iv.
- Sabina, Powder, grs. v to x.  
Oil, gtt. ij to v.
- Salicina, grs. iv to vj.
- Salvia, Infusion, f $\overline{3}$ ij to iv.
- Sanguinaria, Powder, grs. x to xx.  
Tincture, f $\overline{3}$ j to iv.
- Sarsaparilla, Powder,  $\overline{3}$ ss to j.  
Decoction, f $\overline{3}$ iv to vj.  
Extract, grs. x to xx.  
Fluid, f $\overline{3}$ j.  
Syrup, f $\overline{3}$ ss to j.  
Compound, f $\overline{3}$ ss to j.
- Sassafras, Infusion, f $\overline{3}$ j to ij.  
Oil, gtt. ij to v.
- Scammonium, Powder, grs. v to x.  
Confection, grs. xx to xxx.  
Resin, grs. v to x.
- Scilla, Powder, grs. v to x.  
Syrup, f $\overline{3}$ j.  
Compound, gtt. x to f $\overline{3}$ j.  
Tincture, gtt. xx to xl.  
Extract, gr. j to iij.  
Vinegar, f $\overline{3}$ ss to j.  
Oxymel, f $\overline{3}$ j to ij.
- Scoparius, Infusion, f $\overline{3}$ ij to iv.  
Extract,  $\overline{3}$ j to  $\overline{3}$ j.
- Senega, Powder, grs. x to xx.  
Infusion, f $\overline{3}$ j to ij.  
Decoction, f $\overline{3}$ ss.  
Syrup, f $\overline{3}$ j to ij.
- Senna, Powder,  $\overline{3}$ ss to ij.  
Confection,  $\overline{3}$ ij.  
Infusion, f $\overline{3}$ iv.  
Syrup, f $\overline{3}$ j to ij.  
Extract, fluid, f $\overline{3}$ ss.
- Serpentaria, Powder, grs. x to xx.  
Infusion, f $\overline{3}$ j to ij.  
Tincture, f $\overline{3}$ j to ij.
- Simaruba, Infusion, f $\overline{3}$ ij.
- Sodii Chloridum, Powder, grs. x to  $\overline{3}$ ss.  
Comp.  $\overline{3}$ ij to iij.
- Soda, Acetate,  $\overline{3}$ j to  $\overline{3}$ iv.  
Arsenate, gr.  $\frac{1}{16}$  to  $\frac{1}{8}$ .  
Borate, grs. xx to xxx.  
Carbonate, grs. x to xx.  
Solution, f $\overline{3}$ j to ij.  
Hydrosulphate, grs. x to  $\overline{3}$ j.  
Phosphate,  $\overline{3}$ iv to  $\overline{3}$ j.  
Sulphate,  $\overline{3}$ j.  
Tartrate,  $\overline{3}$ ss to j.
- Spigelia, Powder,  $\overline{3}$ j to ij.  
Infusion,  $\overline{3}$ iv to  $\overline{3}$ j.  
Comp.  $\overline{3}$ j to ij.  
Extract, Fluid,  $\overline{3}$ j to  $\overline{3}$ ss.
- Spiraea, Decoction, f $\overline{3}$ j to ij.
- Spiraea, Extract, grs. v to  $\overline{3}$ j.
- Spongia, Burnt,  $\overline{3}$ j to ij.
- Stannum, Powder,  $\overline{3}$ ss.  
Chloride, gr. ij.  
Sulphuret, grs. x to xx.  
Oxide, grs. v to vj.
- Stramonium, Powder, gr. ij to iij.  
Seeds, gr. j.  
Extract (seeds), gr.  $\frac{1}{4}$  to  $\frac{1}{2}$ .  
(leaves), gr. j.  
Tincture (seeds), gtt. xx to xxx.
- Styrax, grs. x to xx.
- Strychnia, gr.  $\frac{1}{16}$  to  $\frac{1}{10}$ .  
Acetate, Solution, gtt. v.  
Iodate, gr.  $\frac{1}{2}$ .
- Succinum, Oil, gtt. v to xv.  
Tincture, gtt. xl to lx.
- Sulphur, Prepared,  $\overline{3}$ ss to ij.  
Electuary,  $\overline{3}$ j to ij.
- Sulphuris Carburetum, gtt. ij to iij.
- Tabacum, Wine, m $\overline{x}$  to xx.  
Tincture, m $\overline{x}$  to xx.
- Tanacetum, Extract, grs. v to  $\overline{3}$ j.
- Taraxacum, Decoction,  $\overline{3}$ j to iij.  
Extract,  $\overline{3}$ j to  $\overline{3}$ j.  
Fluid, f $\overline{3}$ j.
- Terebinthina, grs. ij to v.
- Tolutanum, grs. x to xxx.  
Tincture, f $\overline{3}$ j to ij.  
Syrup, f $\overline{3}$ j to f $\overline{3}$ ss.
- Tormentilla, Powder, grs. xxx to  $\overline{3}$ j.  
Decoction, f $\overline{3}$ j to ij.
- Toxicodendron, Powder, gr.  $\frac{1}{2}$  to j.  
Extract, gr. j.  
Tincture, gtt. v.
- Tussilago, Decoction, f $\overline{3}$ ij to iv.  
Syrup, f $\overline{3}$ j to f $\overline{3}$ ss.
- Ulmus Campestris, Decoction, f $\overline{3}$ iv to vj.
- Ulmus Fulva, Decoction, f $\overline{3}$ iv to vj.
- Uva Ursi, Powder,  $\overline{3}$ j to  $\overline{3}$ j.  
Decoction, f $\overline{3}$ j to ij.  
Extract, grs. v to xxx.
- Valeriana, Powder,  $\overline{3}$ ss to j.  
Electuary,  $\overline{3}$ j to  $\overline{3}$ ij.  
Infusion, f $\overline{3}$ ij.  
Tincture, f $\overline{3}$ j to iv.  
Ammon. f $\overline{3}$ j to ij.  
Wine, f $\overline{3}$ j to iv.  
Oil, gtt. ij to v.  
Extract, grs. ij to x.  
Fluid, f $\overline{3}$ j to ij.
- Vanilla, Powder, grs. v to x.  
Tincture, gtt. xxx to xl.
- Veratria, grs.  $\frac{1}{12}$  to  $\frac{1}{6}$ .  
Tincture, gtt. v to x.



Veratrum Album, Powder, gr. j to ij.  
 Wine, ℥x.  
 Tincture, ℥x.  
 Veratrum Viride, Powder, gr. j.  
 Tincture, fʒss to j.  
 Wine, fʒss to j.  
 Extract, grs.  $\frac{1}{4}$  to  $\frac{1}{2}$ .  
 Viola Odorata, Syrup, fʒj to ij.  
 Wintera, Powder, ʒss to j.

Zinci Oxidum, grs. ij to x.  
 Chloridum, Solution, gtt. v.  
 Tincture, gtt. v.  
 Cyanuretum, gr.  $\frac{1}{12}$  to  $\frac{1}{4}$ .  
 Ferrocyanuretum, gr. j.  
 Solution, fʒss.  
 Sulphas, gr. j to xxx.  
 Solution, fʒss.  
 Valerianas, gr. j to ij.  
 Zingiber, Powder, gr. x to ʒj.  
 Tincture, fʒj to ij.



# TABLE OF PHARMACEUTICAL NAMES,

WHICH

DIFFER IN THE UNITED STATES, AND THE LONDON, EDINBURGH,  
AND DUBLIN PHARMACOPŒIAS.

UNITED STATES.	LONDON.	EDINBURGH.	DUBLIN.
Absinthium,	idem,	idem,	Artemisia absinthium.
Acacia,	idem,	Gummi acaciæ,	A. Arabica et A. vera.
Acetum,	idem,	Acetum gallicum,	Acetum vini.
Acetum destillatum,	idem,	idem,	Acetum distillatum.
Acidum arseniosum,	idem,	Arsenicum album,	Arsenici oxydum album.
Acidum hydrocyanicum,	A. hydrocyanicum dilutum,	A. hydrocyanicum,	Acidum Prussicum.
Acidum muriaticum,	A. hydrochloricum,	A. muriatum et m. purum,	A. muriaticum.
Acidum dilutum,	A. hydrochloricum dilutum,	A. muriaticum dilutum,	idem.
Aconitum,	Aconiti folia et radix,	Aconitum,	Aconitum paniculatum. Folia.
Adeps,	idem,	Axungia,	Adeps suillus præparatus.
Alcohol,	Spiritus rectificatus,	idem,	idem.
Alcohol dilutum,	Spiritus tenuior,	idem,	idem.
Allium,	idem,	idem,	Allium sativum.
Aloe,	idem,	A. Barbadosis, A. Indica, A. socotrina,	A. hepatica, A. socotorina.
Althæa,	Althææ radix et folia,	idem,	Althæa officinalis.
Alumen exsiccatum,	idem,	idem,	Alumen siccatum.
Ammoniæ murias,	Ammoniæ hydrochloras,	Ammoniæ murias,	idem.
Ammoniæ carbonas,	Ammoniæ sesquicarbonas,	Ammoniæ carbonas,	idem.
Ammoniacum,	idem,	idem,	Ammoniacum gummi.
Amygdala amara,	idem,	idem,	Amygdalæ amaræ.
Amygdala dulcis,	idem,	idem,	Amygdalæ dulces.
Angustura,	Cusparia,	idem,	Angustura.
Anthemis,	idem,	idem,	Chamæmelum.



UNITED STATES.	LONDON.	EDINBURGH.	DUBLIN.
Antimonii et potassæ tartras,	Antimonii potassio-tartras,	Antimonium tartarizatum,	Antimonii et potassæ tartras <i>sive</i> tartarum emeticum.
Antimonii sulphuretum præcipitatum,	Antimonii oxysulphuretum,	Antimonii sulph. aureum,	Sulph. antimoniatum fuscum.
Antimonii sulphuretum,	Antimonii sesquisulphuretum,	Antimonii sulphuretum,	idem.
Aqua destillata,	idem,	idem,	Aqua distillata.
Aqua camphoræ,	Mistura camphoræ,	idem,	idem.
	Aqua menthæ pulegii,	Aqua pulegii,	idem.
Argenticyanuretum,	Argenticyanidum,		
Argentinitras,	idem,	idem,	Argentinitras fusum.
Arnica,			Arnica montana.
	Asarum,		Asarum Europæum.
Barii chloridum,	idem,	Barytæ murias,	idem.
Belladonna,	idem,	idem,	Atropa belladonna.
Benzoinum,	idem,	idem,	Styrax benzoin. Resina.
Bismuthi subnitrates,	Bismuthi trisnitrates,	Bismuthum album,	Bismuthi subnitrates.
Calamus,	Acorus,	Calamus aromaticus,	
Calcii chloridum,	idem,	Calcis murias,	idem.
Canella,	idem,	idem,	Canella alba. Cortex.
Cantharis,	idem,	idem,	Cantharis vesicatoria.
Capsicum,	idem,	idem,	Capsicum annuum.
Cardamomum,	idem,	idem,	Amomum cardamomum.
Carota,	Dauci fructus,		Daucus carota var. sylvestris.
Carum,	Carui,	idem,	Carum carui.
Cassia fistula,	Cassia,	Cassiae pulpa,	Cassia fistula.
Cassiae fistulæ pulpa,	Cassia,	Cassiae pulpa,	Cassia fistula.
	Cataplasma fermenti,		Cataplasma fermenti cerevisiæ.
Ceratum cantharidis,	Emplastrum cantharidis,	idem,	idem.
Ceratum cetacei,	idem,	Ceratum simplex,	Unguentum cetacei.
Ceratum plumbi subacetatis,	Ceratum plumbi compositum,		
Ceratum resinæ,	idem,	Unguentum resinosum,	Ung. resinæ albæ.
Ceratum sabinæ,	idem,	idem,	Unguentum sabinæ.
Ceratum simplex,	Ceratum,		
Ceratum zinci carbonatis,	Ceratum calaminæ,	idem,	Unguentum calaminæ.
Cetraria,	idem,	idem,	Lichen Islandicus.
Chimaphila,	idem,	Pyrola,	Pyrola umbellata.
Cinchona flava,	Cinchona cordifolia,	Cinchona flava,	Cinchona cordifolia.
Cinchona pallida,	Cinchona lancifolia,	Cinchona cinerea,	Cinchona lancifolia.
		Cinchona coronæ,	
Cinchona rubra,	Cinchona oblongifolia,	Cinchona rubra,	Cinchona oblongifolia.
Cinnamomum,	idem,	idem et Cassiæ cortex,	idem et Cassia.
Coccus,	Cocci,	idem,	Coccus cacti.
Colchici radix,	Colchici cormus,	idem,	Colchicum autumnale. Bulbus.



UNITED STATES.	LONDON.	EDINBURGH.	DUBLIN.
Colchici semen,	Colchici semina,	idem,	Colchicum autumnale. Semina.
Colocynthis,	idem,	idem,	Cucumis colocynthis.
Colomba,	Calumba,	idem,	Colomba.
	Confectio amygdalæ,	Conserva amygdalarum,	Confectio amygdalarum.
Confectio aromatica,	idem,	Electuarium aromaticum,	Confectio aromatica.
Confectio aurantii corticis,	Confectio aurantii,	Conserva aurantii,	
	Confectio cassiæ,		Electuarium cassiæ.
Confectio opii,	idem,	Electuarium opii,	Confectio opii.
	Confectio piperis nigri,	Electuarium piperis,	Confectio piperis nigri.
Confectio rosæ,	Confectio rosæ gallicæ,	Conserva rosæ,	idem.
	Confectio rosæ cainæ,	Conserva rosæ fructus,	
	Confectio rutæ,		Conserva rutæ.
	Confectio scammonii,		Electuarium scammonii.
Confectio sennæ,	idem,	Electuarium sennæ,	idem.
Conii folia,	idem,	Conium,	Conium maculatum.
Conii semen,	Conii fructus,		
Contrayerva,	Contrajerva,		
Copaiba,	idem,	idem,	Copaifera officinalis.
Coriandrum,	idem,	idem,	Coriandrum sativum.
	Cornu,	idem,	Cornua cervina.
Creasotum,	Creasoton,	Creasotum,	
Creta,	idem,	idem,	Calcis carbonas. Creta alba.
Crocus,	idem,	idem,	Crocus sativus.
Cubeba,	Piper cubeba,	Cubebæ,	Cubeba.
Cuprum ammoniatum,	Cupri ammonio sulphas,	Cuprum ammoniatum,	idem.
Cupri subacetas,	Ærugo,	idem,	Cupri subacetas.
Curcuma,	idem,	idem,	Curcuma longa.
	Cuminum,	Cuminum,	
	Decoctum aloës compositum,	Decoctum aloës,	Decoctum aloës compositum.
Decoctum cetrariæ,	idem,		Decoctum lichenis islandici.
Decoctum chimaphilæ,	idem,		Decoctum pyrolæ.
Decoctum cinchonæ,	Decoctum cinchonæ cordifoliæ,	Decoctum cinchonæ,	idem.
	Decoctum C. lancifoliæ,		
	Decoctum C. oblongifoliæ,		
	Decoctum hordei compositum,	Mistura hordei,	Decoctum hordei compositum.
Decoctum quercûs albæ,	Decoctum quercûs,	idem,	idem.
	Decoctum sarzæ,	idem,	Decoctum sarsaparillæ.



UNITED STATES.	LONDON.	EDINBURGH.	DUBLIN.
Decoctum sarsaparillæ compositum,	Decoctum sarzæ compositum, Decoctum scoparii compositum,	idem, Decoctum scoparii,	Decoctum sarsaparillæ compositum.
Digitalis,	Digitalis folia, D. semina,	Digitalis,	Digitalis purpurea.
Diosma,	idem,	Bucku,	Diosma crenata.
Elaterium,	Extractum elaterii, Emplastrum ammoniaci cum hydrargyro, Emplastrum ceræ,	Elaterium, Emplastrum ammoniaci et hydrargyri, Emplastrum simplex, Emplastrum ferri,	Extractum elaterii. Emplastrum ammoniaci cum hydrargyro. Emplastrum thuris.
Emplastrum ferri, Emplastrum galbani compositum, Emplastrum picis cum cantharide, Emplastrum plumbi,	Emplastrum galbani, idem,	Emplastrum lithargyri, Emplastrum resinosum,	Emplastrum calefaciens. idem.
Emplastrum resinæ,	idem, Euphorbium,	idem,	Emplastrum lithargyri cum resinæ. Euphorbia canariensis.
Extractum aconiti,	idem, Extractum aloës purificatum,	idem, Extractum anthemidis,	Succus spissatus aconiti. Extractum aloës hepaticæ. Extractum chamæmeli.
Extractum belladonnæ, Extractum cinchonæ,	idem, Extractum cinchonæ cordifoliæ, Extractum C. lancifoliæ, Extractum C. oblongifoliæ, Extractum colocynthidis,	idem, Extractum cinchonæ, idem,	Succus spissatus belladonnæ. idem.
Extractum conii,	idem,	idem,	Extractum colocynthidis simplex. Succus spissatus conii.
Extractum hæmatoxyli, Extractum hyoscyami,	idem, idem, Extractum lupuli,	idem, idem, idem,	Extractum hæmatoxyli campechiani. Succus spissatus hyoscyami. Extractum humuli lupuli.
Extractum sarsaparillæ,	Extractum opii purificatum, Extractum sarzæ,	Extractum opii, Extractum sarzæ fluidum,	Extractum opii aquosum. Extractum sarsaparillæ. Extractum sarsaparillæ fluidum.
Ferri filum,		Ferri filum,	Ferrum fila.



UNITED STATES.	LONDON.	EDINBURGH.	DUBLIN.
Ferri ramenta,	Ferrum. Ramenta,	Ferri limatura,	Ferrum. Scobs.
Ferri et potassæ tar- tras,	Ferri potassio-tar- tras,	Ferrum tartarizat- um,	Ferri tartarum.
Ferri ferrocyanuret- um,	Ferri percyanidum,		Ferri cyanuretum.
Ferri oxidum hy- dratum,		Ferrugo,	
		Ferri oxidum ni- grum,	Ferri oxydum ni- grum.
Ferri subcarbonas,	Ferri sesquioxylum,	Ferri oxidum ru- brum,	Ferri carbonas.
Ferrum ammoniat- um,	Ferri ammonio-chlo- ridum,		
Ficus,	Fici,	idem,	Ficus carica.
Filix mas,	Aspidium,	Filix,	Filix mas.
		idem,	
Galla,	Gallæ,		Gallæ.
Gambogia,	Cambogia,	Cambogia,	Gambogia.
Gentiana,	idem,	idem,	Gentiana lutea.
Glycyrrhiza,	idem,	Glycyrrhizæ radix,	Glycyrrhiza glabra.
Granati fructûs cor- tex,	Granatum,		Punica granatum.
Granati radicis cor- tex,		Granati radix,	Punica granatum.
Guaiaçi lignum,	idem,	idem,	Guaiaicum officinale.
Guaiaçi resina,	idem,	Guaiaicum,	Guaiaicum officinale.
Hæmatoxylon,	Hæmatoxylum,	Hæmatoxylon,	Hæmatoxylum cam- pechianum.
Helleborus,	idem,	idem,	Helleborus niger.
Hordeum,	idem,	idem,	Hordeum distichon.
Humulus,	Lupulus,	idem,	Humulus lupulus.
Hydrargyri chlori- dum corrosivum,	Hydrargyri bichlori- dum,	Sublimatus corro- sivus,	Hydrargyri murias corrosivum.
Hydrargyri chlori- dum mite,	Hydrargyri chlori- dum,	Calomelas,	Calomelas sublimat- um.
Hydrargyri cyanu- retum,	Hydrargyri bician- idum,		Hydrargyri cyanure- tum.
Hydrargyri iodidum rubrum,	Hydrargyri biniodi- dum,	idem,	
Hydrargyri oxidum nigrum,	Hydrargyri oxy- dum,		Hydrargyri oxydum nigrum.
Hydrargyri oxidum rubrum,	Hydrargyri nitrico- oxydum,	Hydrargyri oxidum rubrum,	Hydrargyri oxydum nitricum.
Hydrargyri sulphas flavus,			Hydrargyri oxydum sulphuricum.
Hydrargyri sulphu- retum nigrum,	Hydrargyri sulphu- retum cum sul- phure,		Hydrargyri sulphu- retum nigrum.
Hydrargyri sulphu- retum rubrum,	Hydrargyri bisul- phuretum,	Cinnabaris,	Hydrargyri sulphu- retum rubrum.
Hydrargyrum am- moniatum,	Hydrargyri ammo- nio-chloridum,	Hydrargyri precipi- tatum album,	Hydrargyri submu- rias ammoniatum.
Hyoseyami folia,	idem,	Hyoseyamus,	Hyoseyamus niger.
Hyoseyami semen,	Hyoseyami semina,		
Infusum angusturæ,	Infusum cuspariæ,	idem,	Infusum angusturæ.



UNITED STATES.	LONDON.	EDINBURGH.	DUBLIN.
Infusum anthemidis,	idem,	idem,	Infusum chamaemeli.
Infusum armoraciæ,	Infusum armoraciæ compositum,		Infusum armoraciæ compositum.
	Infusum aurantii compositum,	Infusum aurantii,	Infusum aurantii com- positum.
Infusum caryophyl- li,	idem,	idem,	Infusum caryophyllo- rum.
Infusum catechu compositum,	idem,	Infusum catechu,	Infusum catechu com- positum.
Infusum colombæ,	Infusum calumbæ,	idem,	Infusum colombæ.
Infusum diosmæ,	idem,	Infusum bucku,	Infusum buchu.
Infusum gentianæ compositum,	idem,	Infusum gentianæ,	Infusum gentianæ compositum.
Infusum humuli,	Infusum lupuli,		
Infusum lini,	Infusum lini com- positum,	Infusum lini,	Infusum lini composi- tum.
Infusum rosæ com- positum,	idem,	Infusum rosæ,	Infusum rosæ acid- um.
Infusum sarsaparil- læ,			Infusum sarsaparillæ compositum.
Infusum sennæ,	Infusum sennæ com- positum,	Infusum sennæ,	Infusum sennæ com- positum.
		Infusum sennæ com- positum,	Infusum sennæ cum tamarindis.
Infusum tabaci,	Enema tabaci,	idem,	Infusum tabaci.
Inula,	idem,		Inula helenium.
Iodinium,	Iodinium,	Iodineum,	Iodinium.
Ipecacuanha,	idem,	idem,	Cephaelis ipecacuan- ha.
Juniperus,	Juniperi cacumina,	idem,	Juniperus communis. Cacumina.
	Juniperi fructus,	idem,	Juniperus communis. Baccæ.
Krameria,	idem,	idem,	Rhatania.
	Lacmus,	idem,	Litmus.
	Lactuca,		Lactuca sativa.
	Lauri baccæ, Lauri folia,		Laurus nobilis.
		Lauro cerasus,	Prunus lauro-cerasus.
Lavandula,	idem,	idem,	Lavandula spica.
Limon,	Limones,	idem,	idem.
Limonis cortex,	Limonum cortex,	idem,	Citrus medica.
Linimentum cam- phoræ,	idem,	idem,	Oleum camphoratum.
	Linimentum opii,	idem,	Linimentum saponis cum opio <i>vel</i> lini- mentum anodyn- um.
Linimentum terebin- thinæ,	idem,	Linimentum terebin- thinatum,	Linimentum terebin- thinæ.
Linum,	Lini semina,	Lini farina,	Linum usitatissimum.
Liquor ammoniæ,	idem,	Ammoniæ aqua,	Ammoniæ causticæ aqua.



UNITED STATES.	LONDON.	EDINBURGH.	DUBLIN.
Liquor ammoniæ acetatis,	idem,	Ammoniæ acetatis aqua,	idem.
Liquor ammoniæ fortior,	Ammoniæ liquor fortior,	Ammoniæ aqua fortior,	
	Liquor ammoniæ sesquicarbonatis,	Ammoniæ carbonatis aqua,	idem.
Liquor barii chloridi,	idem,	Solutio barytæ muriatis,	Barytæ muriatis aqua.
Liquor calcii chloridi,	idem,	Calcis muriatis solutio,	Calcis muriatis aqua.
Liquor calcis,	idem,	Aqua calcis,	idem.
	Liquor cupri ammonio-sulphatis,	Cupri ammoniati solutio,	Cupri ammoniati aqua.
Liquor ferri iodidi,		Ferri iodidi syrupus,	
Liquor iodini compositus,	Liquor potassii iodidi compositus,	Iodinei liquor compositus,	
Liquor plumbi subacetatis,	Liquor plumbi diacetatis,	Plumbi diacetatis solutio,	Plumbi subacetatis liquor.
Liquor plumbi subacetatis dilutus,	Liquor plumbi diacetatis dilutus,		Plumbi subacetatis liquor compositus.
Liquor potassæ,	idem,	Potassæ aqua,	Potassæ causticæ aqua.
Liquor potassæ arsenitis,	idem,	Liquor arsenicalis,	idem.
Liquor potassæ carbonatis,	idem,		Potassæ carbonatis aqua.
	Liquor potassæ effervescens,	Potassæ aqua effervescens,	
	Liquor sodæ effervescens,	Sodæ aqua effervescens,	
Liquor sodæ chlorinatae,	idem,		
	Manganesii binoxidum,	Manganesii oxidum,	Manganesii oxydum.
Marmor,	idem,	idem,	Calcis carbonas, Marmor album.
Marrubium,	idem,		Marrubium vulgare.
Melissa,		Melissa,	Melissa officinalis.
	Mentha pulegium,	Pulegium,	Mentha pulegium.
	Menyanthes,	idem,	Menyanthes trifoliata.
Mezereum,	idem,	Mezereon,	idem.
		Mistura acaciæ,	Emulsio Arabica.
Mistura amygdalæ,	idem,	Mistura amygdalarum,	idem.
Mistura assafoetidæ,	idem,		Mistura assæfoetidæ.
	Mora,		Morus nigra.
Morphiæ murias,	Morphiæ hydrochloras,	Morphiæ murias,	
Mucilago acaciæ,	Mistura acaciæ,	Mucilago,	Mucilago gummi Arabici.
	Decoctum amyli,	Mucilago amyli,	idem.
Mucilago tragacanthæ,		idem,	Mucilago gummi tragacanthæ.
Mucuna,	Mucuna pruriens,	idem,	Dolichos pruriens.
Myristica,	idem,	idem,	Nux moschata.



UNITED STATES.	LONDON.	EDINBURGH.	DUBLIN.
Myroxylon,	Balsamum Peruvianum,	idem,	Myroxylum Peruvianum.
Oleum amygdalæ,	Amygdalæ oleum,		Oleum amygdalarum.
Oleum bergamii,	Bergamii oleum,	Bergamottæ oleum,	
Oleum cajuputi,	Cajuputi,	Cajuputi oleum,	Melaleuca leucadendron.
Oleum caryophylli,	Caryophylli oleum,	idem,	Eugenia caryophyllata.
Oleum cinnamomi,	Cinnamomi oleum,	idem et Cassiæ oleum,	Laurus cinnamomum.
Oleum limonis,	Limonum oleum,	idem,	Citrus medica.
Oleum lini,	Lini oleum,	idem,	Oleum lini.
Oleum myristicæ,	Myristicæ oleum,	idem,	Myristica moschata.
Oleum olivæ,	Olivæ oleum,	idem,	Olea Europæum.
Oleum ricini,	Ricini oleum,	idem,	Ricinus communis.
Oleum rosæ,		Rosæ oleum,	
Oleum terebinthinæ,	Terebinthinæ oleum,	idem,	Oleum terebinthinæ.
Oleum tiglii,	Tiglii oleum,	Crotonis oleum,	Croton tiglium.
Oleum cari,	Oleum carui,	idem,	idem.
Oleum fœniculi,		idem,	Oleum fœniculi dulcis.
Oleum menthæ piperitæ,	idem,	idem,	Oleum menthæ piperitidis.
	Oleum menthæ pulegii,	Oleum pulegii,	idem.
Oleum rosmarini,	idem,	idem,	Oleum rosmarini.
Oleum succini rectificatum,	Oleum succini,		Oleum succini rectificatum.
	Oleum terebinthinæ purificatum,	idem,	Oleum terebinthinæ rectificatum.
Origanum,	idem,	idem,	Origanum vulgare.
Os,			Ossa.
Papaver,	idem,	idem,	Papaver somniferum.
	Petroleum,	idem,	idem et Bitumen petroleum.
Pilulæ aloës et myrrhæ,	Pilulæ aloës cum myrrhâ,	Pilulæ aloës et myrrhæ,	Pilulæ aloës cum myrrhâ.
	Pilulæ hydrargyri chloridi compositæ,	Pilulæ calomelanos compositæ,	idem.
		Pilulæ colocynthidis,	Pilulæ colocynthidis compositæ.
Pilulæ galbani compositæ,	idem,	Pilulæ assafœtidæ,	Pilulæ galbani compositæ.
	Pilulæ cambogiæ compositæ,	Pilulæ cambogiæ,	Pilulæ gambogiæ compositæ.
Pilulæ opii,		Pilulæ opii sive thebaicæ,	
Pilulæ saponis compositæ,	idem,		Pilulæ saponis cum opio.
Pilulæ scillæ compositæ,	idem,	Pilulæ scillæ,	Pilulæ scillæ compositæ.
	Pilulæ styracis compositæ,	Pilulæ styracis,	Pilulæ è styrace.
Piper,	Piper nigrum,	idem,	idem.
Pix abietis,	Pix abietina,	Pix Burgundica,	idem.



UNITED STATES.	LONDON.	EDINBURGH.	DUBLIN.
	Pix nigra, idem,	Pix arida, idem,	idem et Cerussa.
Plumbi carbonas, Plumbi oxidum semivitreum,	Plumbi oxydum,	Lithargyrum,	Plumbi oxydum semi- vitreum <i>sive</i> lithar- gyrum.
Potassæ carbonas impurus,	Potassæ carbonas impura,		Lixivus cinis.
Potassæ carbonas,	idem,	idem,	Potassæ carbonas è lixivo cinere.
Potassæ carbonas purus,		Potassæ carbonas purum,	Potassæ carbonas è tartari crystallis.
Potassii ferrocyanu- retum,	Potassii ferrocyani- dum,	idem,	
Potassa,	Potassæ hydras, Potassa cum calce,	Potassa, idem,	Potassa caustica. Potassa caustica cum calce.
Potassii iodidum,	idem,	idem,	Potassæ hydriodas.
Potassii sulphuret- um,	idem,	idem,	Potassæ sulphuretum.
Prunum,	Pruna,		Prunus domestica.
Pulvis aloës et canel- læ,			Pulvis aloës cum ca- nella.
Pulvis aromaticus,	Pulvis cinnamomi compositus,	Pulvis aromaticus,	idem.
	Pulvis cretæ compos. cum opio,	Pulvis cretæ opiatus,	Pulvis cretæ compos. cum opio.
Pulvis ipecacuanhæ et opii,	Pulvis ipecacuanhæ compositus,	idem,	idem.
Pulvis stanni,		Stanni pulvis,	idem.
Quassia,	idem,	idem,	Quassia excelsa.
Quercus alba et Q. tinctoria,	Quercus,	Quercus cortex,	Quercus robur.
Quinæ sulphas,	Quinæ disulphas,	Quinæ sulphas,	Quinæ sulphas.
Resina,	idem,	idem,	Pinus sylvestris, Re- sina.
	Rhamnus,	Rhamni baccæ,	Rhamnus catharticus, Baccæ.
Rheum,	idem,	idem,	Rheum palmatum et R. undulatum. Ra- dix.
	Rhœas,	idem,	Papaver rhœas. Peta- la.
	Rosa canina, idem,	Rosæ fructus, idem,	Rosa canina. Fructus. Rosmarinus officinalis. Cacumina.
Rosmarinus,			Rubia tinctorum.
Rubia,	Rumex,		Rumex acetosa.
Ruta,	idem,	idem,	Ruta graveolens.
Sabina,	idem,	idem,	Juniperus sabina.
Saccharum,	idem,	Saccharum purum,	Saccharum officina- rum purificatum.
		Saccharum com- mune,	Saccharum non-puri- ficatum.



UNITED STATES.	LONDON.	EDINBURGH.	DUBLIN.
	Sacchari faex,	idem,	Syrupus empyreumaticus,
Salix,		Salicis cortex,	Salix alba, S. fragilis, S. caprea.
Sambucus,	idem,	idem,	Sambucus nigra.
Santalum,	Pterocarpus,	idem,	Santalum rubrum. Pterocarpus santalinus.
Sapo,	idem,	Sapo durus,	idem.
	Sapo mollis,	idem,	idem.
Sarsaparilla,	Sarza,	idem,	Sarsaparilla.
Sassafras radicis cortex,	Sassafras,	idem,	idem, Laurus sassafras.
Scilla,	idem,	idem,	Scilla maritima.
Scoparius,	idem,	Scoparium,	Spartium scoparium.
Senega,	idem,	idem,	Polygala senega.
Senna,	idem,	Senna Alexandrina et Senna Indica,	Senna.
Serpentaria,	idem,	idem,	Aristolochia serpentaria.
Sevum,	idem,	idem,	Adeps ovillus præparatus.
Simaruba,	idem,	idem,	Quassia simaruba.
Sinapis,	idem,	Sinapi,	Sinapis alba, S. nigra.
Sodæ boras,	Borax, Sodæ carbonas impura,	idem,	Sodæ boras, Borax. Sodæ carbonas venale. Barilla.
Sodæ carbonas exsiccatus,	Sodæ carbonas exsiccata,	Sodæ carbonas siccatum,	idem.
Sodii chloridum,	idem,	Sodæ murias,	idem.
Sodæ bicarbonas,	Sodæ sesquicarbonas,	Sodæ bicarbonas,	idem.
Sodæ et potassæ tartaras,	Sodæ potassio tartaras,	Potassæ et sodæ tartaras,	Sodæ et potassæ tartaras.
Spigelia,	idem,	idem,	Spigelia marilandica.
Spiritus lavandulæ compositus,	Tinctura lavandulæ compositæ, Spiritus menthæ piperitæ,	Spiritus lavandulæ compositus, Spiritus menthæ,	idem. Spiritus menthæ piperitæ.
Spiritus myristicæ,	idem,	idem,	Spiritus nucis moschatae.
Spiritus rosmarini,	idem,	idem,	Spiritus rosmarini.
Spongia,		idem,	Spongia officinalis.
Spongia usta,	Staphisagria,	idem,	Pulvis spongiæ ustæ. Delphinium staphisagria.
Stramonii folia,	idem,	Stramonium,	idem, Datura stramonium.
Stramonii semen,	Stramonii semina,		idem.
Styrax,	idem,	idem,	Styrax officinale.
Styrax purificata,	Styrax colatus,	Extractum styracis,	
Sulphur,	idem,	idem,	Sulphur sublimatum.
Sulphur lotum,		Sulphur sublimatum,	Sulphur lotum.
Syrupus,	idem,	Syrupus simplex,	idem.



UNITED STATES.	LONDON.	EDINBURGH.	DUBLIN.
Syrupus aurantii	Syrupus aurantii,	idem,	idem.
Syrupus corticis,			
Syrupus limonis,	Syrupus limonum,	idem,	Syrupus limonis.
	Syrupus papaveris,	idem,	Syrupus papaveris somniaferi.
	Syrupus rhœados,	idem,	Syrupus papaveris rhœadis.
	Syrupus rosæ,	Syrupus rosæ centi- foliæ,	Syrupus rosæ.
	Syrupus sarzæ,	idem,	Syrupus sarsaparillæ.
Syrupus tolutani,	Syrupus tolutani,	idem,	Syrupus balsami tolu- tani.
Tabacum,	idem,	idem,	Nicotiana tabacum.
Tamarindi pulpa,	Tamarindus,	idem,	Tamarindus Indica.
Tamarindus,	idem,	idem,	Tamarindus Indica.
Tanacetum,			Tanacetum vulgare.
Taraxacum,	idem,	idem,	Leontodon taraxa- cum.
Terebinthina Cana- densis,	idem,	Balsamum Cana- dense,	idem.
Testa,	Testæ,		
Tinctura aloes et myrrhæ,	Tinctura aloes com- posita,	Tinctura aloes et myrrhæ,	Tinctura aloes com- posita.
Tinctura assafoeti- dæ,	idem,	idem,	Tinctura assafoetidæ.
Tinctura benzoini composita,	idem,	idem,	Tinctura benzois com- posita.
Tinctura camphoræ,	idem,	Tinctura bucku, idem,	Tinctura buchu. idem et Spiritus cam- phoratus.
Tinctura castorei,	idem,	idem,	Tinctura castorei ros- sici.
Tinctura colchici se- minis,	Tinctura colchici,	idem,	Tinctura seminum colchici.
Tinctura cubebæ,	idem,		Tinctura piperis cu- bebæ.
Tinctura gallæ,	idem,	Tinctura gallarum,	idem.
Tinctura guaiaci am- moniata,	Tinctura guaiaci composita,	Tinctura guaiaci am- moniata,	idem.
Tinctura hellebori,	idem,		Tinctura hellebori ni- gri.
Tinctura humuli,	Tinctura lupuli,		Tinctura humuli.
Tinctura iodini,		Tinctura iodinei,	Iodini tinctura.
Tinctura iodini com- positæ,	Tinctura iodinii com- posita,		
Tinctura lupulinæ,		Tinctura lupuli,	
Tinctura opii cam- phorata,	Tinctura camphoræ composita,	Tinctura opii cam- phorata,	idem.
Tinctura saponis camphorata,	Linimentum sapo- nis,	idem,	idem.
Tinctura sennæ et jalapæ,		Tinctura sennæ com- posita,	
Tinctura tolutani,		Tinctura tolutani,	Tinctura balsami tolu- tani.
Tinctura valerianæ ammoniata,	Tinctura valerianæ composita,	Tinctura valerianæ ammoniata,	idem.



UNITED STATES.	LONDON.	EDINBURGH.	DUBLIN.
Tolutanum,	Balsamum tolutan- um,	idem,	Toluifera balsamum.
Tormentilla,	idem,	idem,	Tormentilla officina- lis.
Toxicodendron,	idem,	idem,	Rhus toxicodendron.
Tragacantha,	idem,	idem,	Tragacantha gummi.
Trochisci glycyrrhi- zæ et opii,		Trochisci opii,	
	Tussilago,		Tussilago farfara.
Unguentum antimo- nii,	Unguentum antimo- nii potassio tar- tratis,	Unguentum antimo- niale,	Unguentum tartari emetici.
Unguentum cantha- ridis,	idem,	Unguentum infusi cantharidis,	Unguentum cantha- ridis.
	Ceratum cantharidis,	Unguentum cantha- ridis,	
Unguentum cupri subacetatis,		Unguentum ærugi- nis,	Unguentum cupri subacetatis.
Unguentum gallæ,			Unguentum galla- rum.
	Unguentum gallæ compositum,	Unguentum gallæ et opii,	
Unguentum hydrar- gyri,	Unguentum hydrar- gyri fortius,	Unguentum hydrar- gyri,	idem.
Unguentum hydrar- gyri ammoniati,	Unguentum hydrar- gyri ammonio chloridi,	Unguentum præcipi- tati albi,	Unguentum hydrar- gyri submuriatis ammoniati.
Unguentum hydrar- gyri nitratis,	idem,	Unguentum citri- num,	Unguentum hydrar- gyri nitratis <i>vel</i> Un- guentum citrinum.
Unguentum hydrar- gyri oxidi rubri,	Unguentum hydrar- gyri nitrico-oxydi,	Unguentum oxidi hydrargyri,	Unguentum hydrar- gyri oxydi nitrici.
Unguentum iodini,			Unguentum iodinii.
Unguentum iodini compositum,	Unguentum iodinii compositum,	Unguentum iodinei,	
	Ceratum plumbi ace- tatis,	Unguentum plumbi acetatis,	idem.
Unguentum simplex,		idem,	Unguentum ceræ al- bæ.
			Unguentum ceræ fla- væ.
			idem.
Unguentum veratri albi,	Unguentum veratri,		
Unguentum zinci oxidi,	Unguentum zinci,	idem,	Unguentum zinci ox- ydi.
Uva passa,	Uva,	Uvæ passæ,	Vitis vinifera, Fruc- tus siccatus.
Valeriana,	idem,	idem,	Valeriana officinalis.
Veratrum album,	Veratrum,	idem,	Veratrum album.
Vinum,	Vinum xericum,	Vinum album,	Vinum album His- panum.
Vinum antimonii,	Vinum antimonii po- tassio-tartratis,	Vinum antimoniale,	Liquor tartari emet- ici.
Vinum colchici ra- dicis,	Vinum colchici,	idem,	



UNITED STATES.	LONDON.	EDINBURGH.	DUBLIN.
Vinum veratri albi,	Vinum veratri,		
Wintera,		Viola,	Viola odorata.
			Wintera aromatica.
Zinci carbonas,	Calamina,		Zinci carbonas impu- rum, calamina.
Zinci carbonas præ- paratus,	Calamina præparata,	idem,	Zinci carbonas impu- rum præparatum.
Zinci oxidum,	Zinci oxydum,	Zinci oxidum,	Zinci oxydum.



## OFFICINAL PREPARATIONS AND DIRECTIONS.

### INTERNAL REMEDIES.

**POWDERS.**—These are of two kinds: *simple* and *compound*. The first are prepared by pulverization; and the second by the mixture of two or more simple powders, except where one of the ingredients is employed to facilitate the more minute division of the others, as in the case of the powder of ipecacuanha and opium. Many of the most important articles used in this form are powdered by persons who make it a special business, by grinding and stamping. When this operation is performed by the apothecary, or medical practitioner, it is most frequently accomplished by means of the pestle and mortar, and the sieve: but, in some cases, a stone slab and muller are required; whilst in others the article is merely rubbed through a sieve. Whenever a substance cannot be dried completely, without an alteration of its properties, recourse must be had to an intermedium, by which the moisture may be absorbed, or its state of aggregation modified. Thus, sugar is the best intermedium in pulverizing vanilla or nutmeg. When camphor is to be pulverized, the addition of a small quantity of alcohol will much facilitate the operation. In other cases, the intermedium should be of so hard a consistence as to assist in breaking down the substance to be pulverized; thus, gold-leaf is best reduced to powder by rubbing it with sulphate of potassa, and afterwards removing this latter by means of water.

The rules to be observed in the preparation of powders are:—

1. Operate, if possible, on perfectly dry articles, and in dry weather.
2. Adapt the nature of the mortar, and the mode of operating, to the nature of the substance: thus, woods and barks are to be pulverized in an iron mortar; sugar, alum, and nitre, in one of marble; corrosive sublimate in one of glass.
3. The mortar should be provided with a cover to prevent particles from being forced out by the action of the pestle, and also to arrest the escape of the finer particles, which would otherwise be diffused through the atmosphere, causing a loss of the product, and an annoyance to the operator.
4. Separate, from time to time, by aid of a sieve, the pulverized portions, returning the coarser particles to the mortar; and repeat this alternate pulverization and sifting until the process is completed.

*Compound Powders.*—1. Each substance is to be pulverized separately, and the quantity ordered in the prescription then weighed off; otherwise, the due proportions will not be maintained.

2. Pulverize soft substances with those which are the reverse, which will thus serve as an intermedium.

3. Pass all the substances through the same sieve, as this will render them more homogeneous, and will also prevent unnecessary loss.

Some substances, however well dried, are reduced to powder with great difficulty by the usual modes of pulverization; as, for instance, nux vomica and jalap, which require to be exposed to the steam of boiling water for some time, and then rapidly dried.

Few articles, except those containing volatile constituents, are deteriorated in their properties by being pulverized. Some, on the contrary, become more active by this process, from their less active portions being separated; for instance, the powder



of ipecacuanha, if properly prepared, and the ligneous part rejected, is far stronger than the root—all the efficient properties existing in the cortical portion.

Some other processes are employed to obtain certain powders.

*Precipitation.*—Tartar emetic is obtained in a perfectly fine powder, by precipitation from a concentrated aqueous solution by means of alcohol; so, also, the precipitated carbonate of lime is prepared by the action of a solution of carbonate of potassa on water of muriate of lime.

*Elutriation.*—This is a method by which the finer particles of a powder are separated from the coarser. It is performed by diffusing the powder through water, permitting the heavier portions to subside, then withdrawing the fluid, holding the finer particles in suspension, into another vessel, and allowing them to subside. This process is employed in making prepared chalk, &c.

*Granulation.*—This is used to obtain certain metals in a finely divided state; as zinc, tin, iron, &c.

Zinc is granulated either coarsely or finely: the first form is procured by melting the metal in an iron ladle, and pouring it slowly, in a fine stream, into cold water; the fine powder is obtained by rubbing the melted metal in an iron mortar. This metal loses its ductile properties at about  $400^{\circ}$ , and becomes so brittle as to be pulverized without difficulty. It does not melt, except at a temperature of  $773^{\circ}$ , so that the melting is many degrees above its brittle point; hence, by beginning the trituration at the melting point, it gradually reaches that at which it is most readily pulverized. The mortar should be thick and well heated before the fluid metal is poured into it. After being granulated, it should be properly sifted.

The granulation of tin may be effected in the same manner; but a more common mode is to pour the melted metal into a strong wooden box, provided with a closely fitting lid, and agitating till the tin is cold; by this means, it is rapidly converted into powder and small grains, which can be separated from each other by elutriation or sifting.

Iron is granulated by filing, or by means of a pestle and mortar (see page 199); but the best means to procure metallic iron in a finely divided state is by reducing it from the state of the carbonate, by means of hydrogen gas (see page 199).

**PILLS AND BOLUSES.**—Pills are small globular masses, of a semi-solid consistence, of the weight of half a grain to six grains, having as a base a powdery extract, &c., and rendered of the proper consistence by a syrup, pulp, &c. A bolus differs from a pill merely in being much larger, and generally less solid, weighing from six to twelve grains, and even more.

No form of medicinal agents is more frequently employed than that of the pill; not only because of the facility with which it is administered, and its comparatively little taste, but because this form answers so excellent a purpose in the preservation of certain compounds.

The article, or articles, to be administered in this mode, should first be formed into a consistent, moderately firm mass—sufficiently plastic to admit of its being moulded into shape, without adhesion to the moulding instrument, and yet of so firm a consistence as to retain the form given to it. To accomplish this, the following rules must be attended to:—

1. All the constituents which can be pulverized must be reduced to the state of a fine powder, and then thoroughly mixed together.

2. If soft ingredients enter into the composition, they must be triturated with the harder articles, which thus serve as an intermedium.

3. No deliquescent salt should enter into the composition of officinal pills; otherwise they will become altered and decomposed.

4. The nature of the excipient should be suited to that of the constituents: thus, syrups are to be used for most vegetable powders; soap for fatty matters; calcined magnesia for copaiba and turpentine. In some cases, no excipient is required, as for most of the gum resins. Extracts, also, when of proper consistence, can be made into pills without any addition. Mucilage, which is often ordered in the formation of various pills, is only suited to those which are to be used in a short time after



they are made; but is not suited to such as are to be kept, because pills into which it enters as an excipient become so hard as to pass through the bowels almost unaltered.

5. The excipient should be gradually added, and the mass beaten and triturated till it is perfectly homogeneous.

6. When the pilular mass is properly made, the next operation consists in dividing it into pills of equal size, which is effected by dividing it with a spatula into morsels of the requisite weight, and rolling them between the fingers; or, in a more effectual, rapid, and cleanly manner, by means of a pill-machine.

7. To prevent any adhesion of the pills after they are made, and to prevent any disagreeable taste, they are, in most cases, covered with an inert powder: as, powdered liquorice-root, starch, orris-root, lycopodium, and frequently with magnesia; but this latter is wholly unsuited to some preparations—for instance, to pills of calomel—as decomposition will be caused. In Europe, they are often coated with gold or silver leaf; but this plan is seldom adopted in this country. Where, from their nauseous taste, or other circumstances, it becomes expedient to cover them, this is best done by means of gelatine. Each pill, being stuck on the point of a thin wire four or five inches in length, is dipped into a solution of gelatine, so as to coat it completely, and the wire is then inserted into a pin-cushion, or a vessel containing fine sand, and left till the gelatine is firm, which occurs in about a quarter of an hour, when the pill can be removed. (See *Procter's Mohr and Redwood's Pharmacy*, 509.)

EXTRACTS.—This name is given to all products resulting from the evaporation of a solution, maceration, digestion, infusion, or of an expressed juice. They are generally of a soft consistence, of a dark color, and of an odor and taste analogous to those of the substance from which they are derived. They have been classed, according to the fluid employed in obtaining them; as watery, alcoholic, ethereal, vinous, acetic, &c. Some, however, are made from the expressed juices of plants, without any intermedium. The great object, in selecting one or more of these menstrua, is to employ that which takes up the largest proportion of the active ingredients of the root, bark, &c.; for the most advantageous solvent for one constituent may be wholly inefficient for others: thus, alcohol is the proper menstruum for resinous substances, water for bitter extractive, vinegar for articles containing the alkaloids, &c.; but, in many cases, an extract, containing all the active properties of the drug, can be obtained only by a combination or a succession of different solvents.

Extracts made by inspissating the expressed juices of plants have been considered to contain the active principles of the vegetable in a less altered form than by any other mode. But it has been shown that such is far from being the case, as, in some instances, the expressed juice does not contain the active constituents. Thus, in aconitum, much of the efficient principle is left in the residuum; added to which, in some extracts made in this manner, the medicinal ingredients are destroyed, or much deteriorated, in the evaporation. This is obviated, in a great measure, by allowing the expressed juice to evaporate spontaneously at ordinary temperatures; or, by carrying on this process in vacuo. It has been found that extracts made by the latter method have more of the aroma and taste of the original article, keep better, and are more efficient. When inspissation of the natural juices of the plant is not practicable, a solution of the active ingredients is to be obtained by the use of certain menstrua—as water and alcohol—according to the substance to be acted upon. When the active principles are soluble in water, that fluid is to be used as the vehicle of extraction; when resinous, alcohol is to be employed; when oleo-resinous, ether will be found the best menstruum. When it is desired that all the principles soluble in the various medicines should be re-united in the extract, the vegetable substance is first to be subjected to the action of one, and then to another, of the solvents; and the solutions thus obtained are to be mixed together, and properly inspissated. These solutions are obtained by maceration, digestion, infusion, and displacement. Decoction is seldom to be employed, as it affords extracts containing much inert matter, and, of course, of inferior quality. Of late years, the process of displacement has been much used in the formation of extracts, whatever the menstruum selected, as it affords a solution of



the soluble principles in the smaller quantity of fluid; and hence, prevents the injurious effects of too long an exposure to heat in the subsequent evaporation. (See Tinctures.)

With regard to the mode of evaporating extracts, much caution is requisite, as the various constituents of these bodies are very liable to undergo decomposition when exposed to the action of heat and atmospheric air. Sometimes the evaporation, especially of aqueous solutions, is performed over the naked fire; but this plan is liable to many objections, more particularly towards the close of the operation, when a small excess of heat may cause such a change in the character of the ingredients as to deprive them of all useful properties; and it is from this cause that so many of the extracts found in the shops are wholly worthless. Other and far better methods consist in the use of water and steam baths, which prevent all danger of burning the extract. In the inspissation, when conducted in open vessels, it is of importance that the evaporation should be carried on rapidly, that the solution may be exposed to the action of the air for as short a time as possible. Evaporation in vacuo is always to be preferred when practicable, as the process can not only be carried on at a lower temperature, but the deteriorating influence of the air is avoided; the great objection is the additional expense it entails on the operation. Spontaneous evaporation has also been successfully employed; though necessarily slow, it affords excellent products. It is always advantageous to aid this process by employing artificial heat to a certain degree, by conducting it in a drying room or warm closet, and by causing a current of dry air, heated to about 100°, to pass over the surface of the evaporating fluid. Extracts made in this manner keep well, not undergoing the spontaneous decomposition so detrimental to extracts in general. When alcoholic solutions are to be concentrated, distillation should always be used, as not only can the alcohol be saved, but, from the process being carried on out of the atmospheric air, one great source of injury to the product is avoided.

The consistence of extracts varies according to their ingredients, and the purposes to which they are to be applied. Two different degrees are recognized in the Pharmacopœias: one so soft as to be readily made into pills; the other so hard as to be pulverizable. These different forms of extracts are useful because of the readiness with which they can be formed into pills or powders prescribed; but, in many cases, extracts undergo a change, some becoming soft and others hard; and one of the difficult parts of an apothecary's business is to keep these preparations of a uniform consistence. They should be preserved in glazed earthenware pots, protected as much as possible from the action of the air. Notwithstanding every precaution that can be taken, a diminution of activity will, sooner or later, take place; hence, these preparations should be examined from time to time, and, if necessary, renewed. The safest plan is to renew those made from recent plants once a year.

**CONFECTIONS, CONSERVES, ELECTUARIES.**—As there is no essential difference in these preparations, they may be advantageously treated under one head. They are simple and compound: the first being merely a mixture of a recent vegetable substance and sugar, beaten into a uniform mass; the second, of the same substances, but in greater numbers. In consequence of the greater simplicity of prescriptions, of late years, the number of preparations of this kind have greatly diminished, and they are employed more as vehicles for other medicines than for their own remedial powers. The following rules are to be observed in their preparation and preservation:—

1. The powders to be incorporated are to be in a state of fine division, and the pulps used, perfectly homogeneous, and not too fluid.
2. The mixture should be intimate, and the consistence of the resulting mass should be solid enough to prevent a separation of the ingredients, and yet sufficiently soft to allow of its being swallowed without being masticated.
3. If the mixture swells up, and gives off carbonic acid, it is to be well rubbed in a mortar, to break down any sugar that may have crystallized, and any hard lumps that may have formed, so as again to form a uniform mass.
4. They should be preserved in earthenware or porcelain pots, well covered, and kept in a moderately moist situation.



**PULPS** are simple medicinal preparations, of a soft consistence, formed of the tissues of vegetable substances, or of some of its thin parts; they differ from extracts in containing insoluble as well as soluble principles. They are to be prepared by rubbing the fruit, or other articles from which they are made, through a metal sieve, so as to separate the vegetable fibre from the soft parts by which it is surrounded. When the fruit is hard or dry, it is to be softened by means of boiling water, previous to expression through the sieve. The expressed pulp, if not sufficiently consistent, is to be properly evaporated. Pulps are to be preserved in the same manner as confections. Very few of these preparations are now used in medicine.

**SYRUPS.**—These are liquid, viscous medicines, consisting of a concentrated solution of sugar in aqueous fluids.

All fluids susceptible of dissolving more than their weight of sugar can be formed into syrups. These syrups are of two kinds: simple or compound. Simple syrup is prepared by dissolving sugar in pure water; and compound syrups are obtained by dissolving the sugar in solutions of various substances, formed by infusion, decoction, expression, &c. The former is usually made in this country from refined sugar, and not, as in Europe, from the impure and unrefined qualities of this article. All syrups require to be perfectly filtered, so as to be limpid, and should have a certain viscosity of consistence, and be capable of being preserved without entering into fermentation, or crystallizing. These latter properties depend on their containing a proper proportion of sugar—an excess being deposited in a crystalline form, and a deficiency causing the solution to run into fermentation. The best mode of ascertaining the proper point of concentration is by means of its specific gravity at different temperatures. The specific gravity of well-prepared simple syrup is, when boiling, about 1.261, and when cold 1.319; but the proper degree of concentration is more readily obtained by means of Baumé's hydrometer. This should stand at about 30° in boiling syrup, and at 35° when it is cold. Other modes are also employed, which, although sufficiently accurate in the hands of an experienced operator, are not to be generally depended upon. They are derived from the degree of viscosity acquired by the syrup, as shown by the time required for the parts of a drop to re-unite, and by the length of the thread which a drop will produce before detaching itself, when poured from a spoon or ladle. When the syrup, on cooling, presents a crystalline pellicle, it is a proof that the evaporation has been carried too far; but, when the sugar has been mixed with an acid, or when the process has been too much prolonged, the sugar loses its power of crystallization, however much the syrup is concentrated, and, therefore, does not form a pellicle.

The compound syrups, when kept any time, are liable to various alterations, depending on their nature, and the degree of care used in their preparation. Thus, the acid syrups, as the syrup of lemons, when too concentrated, deposit a copious white precipitate; and, in some cases, solidify entirely. By heating them, they again become liquid; but again let fall a precipitate on cooling. This deposit is analogous to grape sugar, and is caused by the action of the acid on the sugar. When the sugar bears too small a proportion to the liquid, syrups are apt to run into fermentation. Even when the sugar is in proper proportion, this change often takes place, if the solution contains much amylaceous or extractive vegetable matter. Even when too much concentrated, they may also undergo this change, from part of the sugar being deposited in a crystalline state; and the crystal, attracting the sugar necessary to the preservation of the syrup, reduces its strength, and renders it liable to the same change as though it was originally too weak.

Syrups, especially those containing the juice of fruit, should be bottled whilst hot, and, when cold, well stopped and sealed; and these, as well as all other kinds, should be kept in a temperature not exceeding 60° F. Various plans have been devised to preserve syrups; but the best is to prepare them only in such quantities as will be used within a short time. The addition of chlorate of potassa, as advised by Macculloch, and of sugar of milk, as advised by Chereau, has proved useful; but the best mode is that of Mr. Durand, viz., adding about one drachm of Hoffmann's anodyne to



each pint of syrup; this appears to have the property of arresting or preventing any tendency to fermentation.

MELLITES, OR HONEYS, are liquid, viscid medicines, somewhat analogous to syrups, but in which the sugar is replaced by honey. Like syrups, they may be divided into simple and compound, or medicated. Their preparation, medical properties, modes of administration, and preservation are very similar to those of the syrups, and do not, therefore, require further notice. Though formerly much used, they are now seldom prescribed, as they are often found to disagree with the stomach, especially if made with the honey of commerce in an unpurified state.

INFUSIONS are aqueous solutions, made by treating vegetable products with cold or hot water, but never carried to ebullition. They are seldom made by the apothecary, who merely furnishes the medicinal ingredients; whilst the preparation is usually confided to the nurse or other attendant on the patient. This plan, although more economical than when the preparation is compounded by the apothecary, often renders the prescription of the physician of little avail, from the infusion being erroneously made. They are generally prepared by pouring boiling water on the ingredients, and macerating in a tightly-covered vessel until the liquid cools; or the vessel is kept for some time at a low heat before a fire. In most cases, an infusion should be strained or filtered before being used; but, in general, nurses are content merely to decant them for use. The vessels in which infusions are made are usually of stone or earthenware. When it is an object to retain the heat as long as possible, polished metallic vessels, provided they are not acted upon by the infusion, are to be preferred, as, from their inferior radiating powers, they retain heat better than earthenware. Infusions are mostly made for extemporaneous use, as they are exceedingly liable to decomposition, and consequently cannot be kept for any time without spoiling. They may, however, be preserved for some time by pouring them, whilst hot, into bottles, so as to fill them, and forcing in stout corks. If the operation be properly performed, so as to perfectly exclude the air, the infusion will keep for some months. An infusion made with cold water, by the process of displacement, has less tendency to spoil than one made in the usual manner with hot water.

DECOCTIONS differ from infusions in the circumstance that the substance to be acted upon is subjected to the solvent action of the menstruum at its boiling point, and continued for a longer or shorter period, according to the solubility of the substance, or its constituents. By decoction, all the principles soluble in water can be obtained; and even many substances not properly soluble in that fluid are diffused through it, and held in suspension. On the other hand, many substances are injured or destroyed by decoction, especially when their active principles are volatile, and when, during ebullition, chemical changes take place, by which the active constituents are rendered insoluble, or are decomposed. In fact, protracted ebullition is seldom required since the process of displacement has been known, as it has been found that water at a common temperature, by this method, will exhaust most vegetable substances more effectually than the same fluid at  $212^{\circ}$  by decoction. There are some exceptions; as all emulsive preparations, and certain gummy solutions, require ebullition. As in the case of infusions, decoctions are only made extemporaneously, for they readily decompose by keeping. They should be prepared in closed vessels, and generally in those of earthenware. Copper vessels should be used with great care, as many vegetable decoctions corrode them, especially if permitted to cool in them. Iron vessels are not to be employed when the decoction contains tannin or gallic acid. Zinc is very rapidly acted upon by many vegetable substances, and should, therefore, be avoided. In all cases, the vessels employed should be deep and narrow, rather than broad and shallow, to obviate as much as possible the influence of the air on the product.

TINCTURES are solutions of vegetable, animal, and, in some cases, of mineral substances in spirituous fluids. The spirit employed is alcohol, either diluted or undiluted, either pure or medicated; and, in some cases, ether. The form of tincture is one much



used in pharmacy; it presents the active principles of drugs in a small volume, it can be preserved in an unaltered state for a long time, and is, in most cases, well adapted to unite with other substances in extemporaneous prescriptions. Tinctures are made by maceration, or by displacement.

*Maceration* is an operation in which, by the action of a fluid at common temperatures, continued for a certain length of time, a solution of the principles of an organic substance in that fluid is obtained. As before stated, in making tinctures, the strength of the spirituous menstruum employed is varied according to circumstances. If the substance to be acted upon is resinous, alcohol is to be employed; if it contains also matters soluble in water, and insoluble in pure alcohol, this article, in a diluted state, is to be used. The period of maceration varies from some hours to several weeks; and, during the process, the vessel in which it is performed should be kept closed, to prevent evaporation. When the spirit has remained upon the substance for the directed period, it should be decanted, instead of being left, as is too often the case, standing on the dregs until it is used. This practice is erroneous, and may be attended with evil consequences; for the characters and properties of a tincture which has been suffered to remain too long in contact with the solid ingredients, will often be found to differ considerably from what they would have been if the process had been terminated at the indicated time. When a tincture is made by maceration, it requires to be filtered, after being separated from the dregs; otherwise, it will be turbid, and deposit much insoluble matter on standing.

*Displacement.*—This process is of comparatively late introduction in the making of tinctures, &c. For a great proportion of these preparations, it is decidedly to be preferred to maceration; but for others it has not proved so satisfactory. Mohr and Redwood observe, on this point:—

“When tinctures are made in large quantities, displacement is never likely to supersede maceration, on account of any practical advantages it may possess. If the prescribed directions be duly attended to, the process of maceration is unexceptionable. The process is more simple than the other; the mode of operating is more uniform, it is, in fact, always the same; it requires less of skill and dexterity in conducting it; it requires less constant attention during its progress, which, in operating on large quantities, is a consideration; and, finally, the apparatus required is less complicated.

“When only small quantities of tincture are made at a time, and kept in stock, the adoption of the process of displacement will often be found convenient and advantageous. It offers the means of making a tincture in two or three hours, which, by the other process, would require as many weeks. The process being thus completed in so short a time, for the quantity contemplated might be made at one operation, it would not be so likely as the other to experience neglect during the performance of it, or a deviation from the prescribed instructions; the product would, therefore, be more uniform. Finally, in many cases, the tincture and spirit may be removed from the dregs more completely, in operating on small quantities, by this process, than by the other.”

Several forms of displacement-apparatus are used in this country, but the one most frequently employed is that of Boullay. It is a vessel nearly cylindrical, made of tin, with its lower end funnel-shaped. It is made of various sizes: with the cylinder 12 inches in length by 2½ in diameter, or 14 by 4, or 17 by 6. A movable metallic plate, pierced with numerous holes, and provided with a knob or handle in the centre, and fitted accurately to the lower part of the cylinder, is to be placed in the projecting supports. On this, a thin stratum of carded cotton, covered with the powder to be operated upon, is to be laid, and the process carried on as directed above. This apparatus should be furnished with a stopcock at the lower end. Soubeiran has made an improvement on this filter by adapting to it a tin receiver, provided with a cock at the lower part, so that the filtered liquor can be drawn off at will. The substance to be exhausted must be reduced to various degrees of division in different instances; but, for the most part, the form of coarse powder is the most suitable, because, when the powder is very fine, the fluid is apt to be arrested in its passage, after becoming saturated with the soluble principles. In general, it is proper to soak the powder for a few hours with a sufficiency of the spirit, to make it into a stiff pulp, before putting it into



the cylinder; otherwise, the spirit may make narrow passages for itself, and leave a part of the material unacted on: but, in some instances, no such preliminary step is necessary, and here, as in the case of tincture of myrrh, the best procedure is to introduce the powder in the dry state, and to pour the spirit over it. Before the spirit is poured over the pulp or powder, it must be packed in the cylinder. The degree of pressure to be used, which varies in different instances, is to be learned only from experience in each case; and it may be observed, that this is the operation on the correct performance of which the success of the process chiefly depends, and the only one which requires some skill and nicety of management. If the material be loosely packed, the spirit will pass through too quickly, and not sufficiently saturated; if too firm pressure be employed, percolation will take place too slowly, or not at all. On the whole, the firmness of the packing should be inversely as the solvent power of the spirit upon the solid materials; but to this rule there are many exceptions. Since the fluid, for an obvious reason, passes after a time with increasing speed, it is occasionally necessary to press down the matter in the cylinder as the process advances; but neither does this rule always apply. In some operations, especially those on the large scale, it is found useful to promote the displacement by pressure. This may be produced simply by supplying the solvent through a tube several feet high, adapted to the upper end of the cylinder, and terminating at the top in a funnel. A still better method, as involving the use of less spirit, consists in the use of compressed air. By means of a condensing syringe, or a column of water or mercury in a tube, air is condensed in the bottle below; and when the material and spirit have been introduced into the percolator, the top is tightly secured by a screw, and the compressed air admitted by a stopcock. This method answers excellently on the small scale with a column of mercury.

The solution which first passes through in this process is always in a state of high concentration. In general, it is a simple solution of the soluble ingredients of the crude drug in the fluid employed. But sometimes the solvent, if compound, is resolved into its component parts; and the fluid which passes through is only one of these, holding the soluble parts of the drug in solution. Thus, if diluted alcohol be poured over powder of myrrh, in the cylinder, the fluid which first drops into the receiver is a solution of oily consistence, composed of resin and volatile oil dissolved in alcohol. And, if powder of galls be acted on in like manner, by aqueous sulphuric ether, two layers of fluid are obtained, a highly concentrated solution of tannin in the water of the ether, and a weak solution of the same principle in pure ether.

Displacement is accomplished, in the manner here described, with little or no intermixture of the liquid above, with that which is below, in the pulpy mass. If, after the pulp has been allowed to drain till drops cease to fall, the quantity of spirit retained by it be ascertained—that is, by subtracting what has dropped from what has been used to make the pulp,—and the same quantity be poured gently over the mass, repeatedly, as soon as the dropping caused by each successive addition ceases, a series of liquids will be obtained by the simple displacement of one another in the pulp; and the last portion of spirit used may be obtained by pouring over the pulp the same quantity of water. Hence, one of the many advantages possessed by this method of making tinctures, over the ordinary mode by maceration and expression, is that no part of the tincture is lost by being left behind in the residuum.

Even when a strict attention is paid to the prescribed formula, the resulting tinctures may be found to vary in medicinal strength; for any variation in the activity of the drug will influence the tincture. These preparations, although retaining their properties better than other vegetable solutions, will gradually undergo changes, from age and exposure, either losing their color, depositing insoluble matter, or, in some cases, as in that of the tincture of kino, becoming less astringent, and gradually gelatinizing.

WINES are tinctures in which the menstruum is wine. The wine ordered to be employed by the U. S. and British Pharmacopœias is Sherry. The solvent power of wines on vegetable products depends on the water and alcohol they contain; the other principles found in them are injurious to it as an excipient, the mucilage disposing



to fermentation, and the coloring matters and tannin often decomposing the active principles of the vegetable substances dissolved in it. On this account, these preparations are now going out of use, and mixtures of alcohol and water, of the proper strength, substituted as solvents. These can always be made of definite strength, which can seldom be attained when wine is used.

Medicated wines are made by maceration or displacement; but the latter process is seldom used. In whatever way they are prepared, they are very liable to undergo a change, and hence should be made in small quantities only, and kept in well-closed bottles, and in a cool and dark place.

VINEGARS are solutions of certain parts of vegetables in distilled vinegar or diluted acetic acid. Few of these preparations are officinal; but, in some cases, the state of solution in distilled vinegar is the best that can be desired, at least as far as energy is concerned. This is particularly the case when the activity of the medicinal substance depends on the presence of an alkaloid, which, by uniting with the acetic acid, forms a salt that is readily soluble; and, therefore, may be more perfectly extracted than by other menstrua.

As even distilled vinegar contains vegetable matter, rendering it liable to decomposition, it has been found advantageous to substitute acetic acid, properly diluted. In consequence of this liability to become decomposed, medicated vinegars should be made in small quantities, and renewed at short intervals. In many cases, the acetates of the vegetable alkaloids, dissolved in water or diluted alcohol, may be substituted for medicated vinegars, producing all their good effects, and attended with but few of their disadvantages.

MIXTURES are liquid medicines containing two or more ingredients, generally of extemporaneous preparation, though a few are recognized as officinal. They are in very constant use in the administration of remedies. The ingredients which usually enter into their composition are salts, and other solid bodies which are soluble, or readily miscible with aqueous fluids; also tinctures, spirits, syrups, decoctions, &c.

In making a mixture, the objects to be attained are the perfect solution of the soluble ingredients, and an equal diffusion of those which are not so, through the excipient, as well as the thorough admixture of all the constituents, so that each dose shall be similarly composed. The combination of articles in a mixture is effected either by agitation, or by rubbing the ingredients together in a mortar. When volatile substances enter into a mixture, they should be added last. Insoluble powders, which are easily diffused through a fluid, may be mixed with the liquid ingredients, by merely shaking them together in the bottle. When, however, powders do not mix readily with liquids, they must be triturated in a mortar with a small quantity of the fluid, gradually adding more, until they are of the consistence of thin paste. In the same manner, extracts and electuaries are to be rubbed down with a portion of the excipient previous to putting them in the bottle.

*Draughts* differ from mixtures only in being designed to be taken at a single dose. This is not a common mode of prescribing medicines, but in some cases it is advantageous, as, when it is wished to apportion the doses of a medicine accurately, or when the medicine is liable to undergo a change from the action of air.

*Emulsions* are mixtures containing substances of an oleaginous or resinous nature suspended in aqueous fluids by the aid of gum, syrup, yolk of eggs, or any viscid matter. It is a mechanical compound, and the union of its ingredients is promoted and rendered more or less permanent by the addition of an alkali. Emulsions are made by triturating certain oleaginous seeds with water, or by mixing the oil procured from them with the necessary ingredients; and also from gum resins in the same manner. Sometimes they are made with oils and an alkali. In making an emulsion of the oils or oleo-resins with mucilage, the former should be gradually added to the latter; by adding the mucilage to the oil, a good emulsion cannot be formed. Mucilage is preferable to an alkali in making many emulsions, as those with castor oil or copaiba; an alkali is preferable in forming an emulsion of almond or olive oil; but an emulsion formed with either of these agents alone will often separate, if the other be added. The presence of soluble salts in an emulsion is also apt to



cause a separation of the oil. Much spirit will produce the same effect in emulsions made with mucilage, and an acid in those made with an alkali. Some substances cannot be formed into good emulsions either with an alkali or with mucilage. This is the case with spermaceti and oil of turpentine. In such cases, the yolk of an egg is the best agent for effecting the admixture, by rubbing down the ingredients in a mortar.

Volatile oils are most readily made into an emulsion by being previously mixed with one of the terebinthinate fixed oils. Scammony is formed into an emulsion by means of milk; but resin of jalap will not unite with this fluid, and is best emulsified by triturating it with almonds and water. Emulsions should be made with cold water, and strained.

Mr. W. Procter gives the following valuable directions for making emulsions (*Phil. Journ. Pharm.* xv. 11): "In making an emulsion, a good deal depends even on so slight a circumstance as the form and material of the mortar and pestle. An emulsion may, indeed, be made in a smooth porcelain mortar, but the process is unquestionably more successful and easy in a Wedgewood one, and still more so in the old-fashioned marble mortar, which is generally used to this day by the druggists and apothecaries in England. But whatever be the composition of the mortar, it is essential that it should be perfectly *round* at the bottom; not flattish, as is sometimes the case; and the pestle should be so formed as in its motion to leave no hollows between its base and the concave surface of the mortar. Otherwise, the emulsion will most probably be imperfect. The mucilage or other viscid substance should always be put into the mortar before anything else: the oil (or balsam) may then be very gradually rubbed in, taking care not to add it more quickly than it can be subdued by the pestle; and if, during this part of the manipulation, the mixture should begin to assume a breaking or curdling appearance at the edges, a few drops of water must be immediately incorporated with it, before adding the remainder of the oil. For want of this precaution, I have often known an emulsion suddenly to lose its tenacious consistence in the mortar, and it is then in vain to endeavor to restore it. After the oil is thoroughly incorporated, some care is requisite to avoid separating it again by too hasty an effusion of the water or other fluid of the mixture; and, if any alcoholic or acid liquid is to be added, it must be at the very end of the process. Indeed, an acid liquid, even a slightly acescent syrup, will often entirely destroy the emulsion. Mixtures of copaiba are frequently spoiled by the addition of spir. eth. nit.; which might be avoided by first diluting it with one or two portions of water."

**MEDICATED WATERS.**—This term is used by the U. S. Pharmacopœia to designate all preparations of water impregnated with medicinal substances, not appertaining to other classes of preparations. It includes the **DISTILLED WATERS** of the British Colleges, and is a more appropriate designation, because many of the latter preparations are not distilled waters, but are made with an essential oil united to the water by trituration or agitation, no distillation being employed.

*By Distillation.*—There are two modes of obtaining medicated waters by this process: 1st, by distilling the water directly off the substances; 2d, by employing the essential oil already separated from the vegetable, and distilling it over with the water. In most cases, where the fresh plant can be procured, it is to be preferred. Many of these substances lose their volatile oil and fragrant properties when dried; but, in some instances, the oil is retained, notwithstanding desiccation. Many which lose the oil by the process of drying retain it fully for a length of time, when preserved by being beat into a pulp with common salt; and, when used in this state, afford very good distilled waters.

The material which supplies the volatile oil is, in general, simply mixed with the water in a state of fine division; and this is probably the best mode, where heat is used in such a way as to exclude the risk of empyreuma. If it does not require to be finely divided, as in the case of fresh leaves and flowers, it may be put conveniently into a net-bag, which, suspended in the middle of the still, may be withdrawn with facility, when its contents are exhausted. Some manufacturers use, not water, but steam, for obtaining distilled waters: that is, the material to be distilled is spread



over a fine gauze partition, or a plate perforated with numerous small holes, and steam is driven through the mass.

When the vegetable substance to be exhausted is a bark, wood, or other solid matter, it must be reduced to a state of moderately fine division. But this is not generally necessary in the case of leaves or flowers, because boiling water breaks down the cells in which the volatile oil is contained. When leaves, however, are thick and leathery, as in the instance of the cherry-laurel, the process is facilitated by chopping them down; and, in most cases, where leaves are large, it is difficult to get a sufficient quantity into the still without cutting them into pieces.—In preparing the finer kinds of distilled waters, it is necessary to clean the materials carefully, to remove all decayed leaves or flowers, or those infested by insects, and sometimes also to separate the leaf-stalks, or the green claw of the petals.

Heat may often be applied directly to the vessel; but in this way empyreuma is apt to be occasioned, especially in large operations, in consequence of the solid matters remaining fixed at the bottom. To avoid this, it is usual to apply the heat, in limited operations, through the medium of a solution of hydrochlorate of lime, which raises a temperature between  $212^{\circ}$  and  $270^{\circ}$ , according to its strength—or by means of an oil-bath, with a thermometer to regulate the temperature; and, on the great scale, it is best applied by means of steam admitted under pressure into a space surrounding the still.—Another cause of the empyreumatic taint of some distilled waters is the formation of a species of mucilaginous substance, at the expense of the volatile oil. This substance, which forms chiefly when the distillation is pushed too fast, or too far, and is seen encrusting globules of volatile oil, undissolved in the water, is apt to deposit itself on the side of the still, above the boiling materials, where it is afterwards decomposed by the heat. This fact explains the well-known observation, that the finest distilled waters are obtained by gentle distillation, and by abstaining from complete exhaustion of the materials. A still greater improvement is to prepare them with the vacuum-still, in the same way as is now often practiced in making extracts.

The other mode of obtaining medicated waters by distillation is by employing the volatile oil, previously separated from the plant, for the vegetable itself. This mode of operating affords a product less liable to change than the above; but it is not so aromatic as when obtained from the fresh plant.

Distilled waters, however carefully they may be kept, are apt, sooner or later, to lose their aroma; and some of them even become mouldy, and acquire thereby an unpleasant odor. They have been thought to keep better with the addition of about a fortieth part of rectified spirit; which may be either put into the still with the water, or added afterwards to the distilled fluid. But the advantages of this addition, although sanctioned by the authority of the Dublin Pharmacopœia, are doubted by practical men. It is believed that the most effectual precaution for preserving them is to prepare them with extremely pure natural waters, such as snow, rain, or very fine spring water (Müller), free especially of any unusual proportion of carbonic acid; and to keep them in black, orange, or red bottles, not in bottles of clear glass (Hanle). A better mode of procedure is to re-distil the water as soon as any change is perceived in it; this restores its original odor, and renders it less subject to alteration.

*By Admixture.*—Another method of making medicated waters is by impregnating the water with the volatile oil, by triturating them together with the addition of carbonate of magnesia, and filtering to remove the latter. This affords an excellent product, as it is pure and permanent. The addition of the carbonate of magnesia is merely to enable the operator to produce such a minute division of the oil as will enable water to act on it more efficiently. Other powders act in the same way, and are preferred by some manufacturers.

**DISTILLED, ESSENTIAL, OR VOLATILE OILS.**—Volatile oils are contained in vegetable cells, generally peculiar, and often so large as to be distinct to the naked eye. Sometimes they exist in such abundance that the oil may be obtained by mere expression. The oils of lemon, orange, bergamot, and citron are prepared in this way by the manufacturer; and from many other substances, such as the unripe germen of



rue, and the undeveloped corolla of the clove-tree, oils may be squeezed out by pressure with the nail. In some rare instances, as in that of the Liquid Borneo-Camphor from the *Dryobalanops Camphora*, and that of the Laurel-oil of Guiana from a species of *Ocotea*, volatile oil is obtained largely, and of considerable purity, by exudation from incisions. Much more frequently, however, it is exuded spontaneously, or from incisions, as a turpentine in combination with resin, or as a gum-resin in union with both gum and resin. Most frequently of all, it cannot be obtained by any of these modes, but adheres with more or less force to the flowers, leaves, fruit, bark, or wood, which contain it. In that case, it is sometimes destroyed or dispersed when the plant is dried, more especially if the organ which contains it is the flower; but often, when contained in the leaf, and very generally, if contained in the seed, bark, or wood, it is retained in part, or altogether, under desiccation, and even under long keeping. In some instances, it appears that the volatile oil obtained from plants does not exist ready formed, but is produced, on bruising or distilling them with water, through the reaction of other principles on one another. Of this mode of production, two remarkable examples exist—the volatile oils of the bitter almond and black mustard seed; and it is probable that the oils of cherry-laurel leaves, peach leaves, and of the leaves of other amygdalaceous plants, are similarly circumstanced.

A few volatile oils used in medicine are obtained by expression, such as the oils of orange, lemon, and bergamot, already mentioned. When oils exude along with resin in the form of turpentine, they may be separated from the resin by heat alone, as they are volatilizable at about the temperature of  $400^{\circ}$ . But thus obtained, they cannot be of fine quality, because the heat required is too near that at which resins, as well as the oils themselves, undergo decomposition. In general, therefore, this plan is not followed, for pharmaceutic purposes; and volatile oils are obtained from turpentines by distilling them with water. For, although they do not enter into ebullition at the temperature of boiling water, their vapors pass over in large quantity with steam; and, condensing along with it, form distilled waters—with volatile oils either floating on the water, or, more rarely, sinking to the bottom. The volatile oils of many vegetable substances are to be separated in this way alone, because the other principles, contained along with them in the crude substances, are empyreumatized by the higher temperature which is required to disengage the oil without the co-operation of watery vapor. In some instances, it has been thought advantageous to substitute for water in the still a strong solution of common salt, because a somewhat higher temperature is required to boil it. There seems an advantage, in this change, for the rectification of oils previously obtained from the raw materials in which they reside; but the advantage in the case of distillation from crude vegetable substances is doubtful.

The method of distilling volatile oils differs little from that described above for preparing distilled waters. The same precautions must be observed in applying heat, and for the same reasons. The formation of mucilaginous matter at the expense of the oil, towards the close of the distillation, is shown by the globules of oil coming over enveloped in a fine pellicle of it. The quantity of water must be proportionally less, or, rather, the same portion of water is to be used with successive portions of the material which yields the oil; otherwise, a material loss is sustained by solution of the oil in the water. In some instances, where the oil exists in low proportion, and is of great value, the distilled fluid should be left at rest for some time, and exposed to as low a temperature as can be commanded.

The mixed vapors which pass over condense into a milky-looking fluid, which, after standing some time in the receiver, separates into two portions, one a solution of a part of oil in water, and the other of the oil itself, which occupies the upper or lower part, according as it is lighter or heavier than the water.

Chevallier gives the following rules for the distillation of volatile oils:—

1. To operate upon large quantities, in order to obtain a greater product, and of better quality.
2. To conduct the distillation rapidly.
3. To divide the substances minutely, in order to facilitate the extrication of the oil.
4. To employ only sufficient water to prevent the plant from burning.



5. For substances whose oil is heavier than water, to saturate the water in the still with common salt, to raise the boiling point, and thus to enable the vapor to carry over more of the oil.

6. To employ, when possible, water which had already been distilled from off the same substances, and had thus become saturated with oil.

7. For oils naturally fluid, to cool the liquid in the refrigeratory frequently; but to retain it at 80° to 90° for those oils which easily become solid.

After the distillation, the oil should be separated from the water; for this purpose, several ingenious instruments have been devised, an account of which will be found in Procter's edition of *Mohr and Redwood's Pharmacy*, pp. 354, 355.

The following table, from *Christison's Dispensatory*, showing the proportion of volatile oil obtained from the chief medicinal vegetable substances, according to the most recent experiments, will be found useful. The data are chiefly extracted or calculated from experiments by M. Raybaud in the *Journal de Pharmacie*, xx.—by Dr. Martius in *Repertorium für die Pharmacie*, xxxix.—by Dr. Bley in the same work, xlviii.—by M. Dann and by M. Voelter in the same work, lv.;—and a few have been added from experiments of my own. The numbers represent the number of ounces obtained from 100 pounds avoirdupois. The letters before the figures refer to the authority for each.

Amygdalus communis.	Bitter almond	-	-	-	-	-	Ra.	0.38
Amygdalus communis.	Bitter almond	-	-	-	-	-	Vo.	7.70
Amygdalus communis.	Bitter almond (Duflos)	-	-	-	-	-	0.8 to	4.80
Angelica Archangelica—	dried root	-	-	-	-	-	Ra.	4.50
Anthemis nobilis—	fresh flowers raised at Grasse	-	-	-	-	-	Ra.	0.75
Anthemis nobilis—	dried flowers do.	-	-	-	-	-	Ra.	1.38
Anthemis nobilis—	dried flowers, long kept, Germany	-	-	-	-	-	Bl.	4.50
Anthemis nobilis—	flowers freshly dried	-	-	-	-	-	Steer.	5.33
Anthemis nobilis—	flowers 12 months dried	-	-	-	-	-	Steer.	3.0
Apium graveolens—	dried fruit	-	-	-	-	-	Ra.	9.00
Apium Petroselinum—	fresh herb, after flowering	-	-	-	-	-	Ra.	3.38
Apium Petroselinum—	dry fruit, France	-	-	-	-	-	Ra.	12.0
Apium Petroselinum—	dry fruit, Germany	-	-	-	-	-	Da.	30.0
Artemisia Absinthium—	fresh herb, Paris	-	-	-	-	-	Ra.	2.0
Artemisia Absinthium—	dried herb, recent, Germany	-	-	-	-	-	Ma.	16.0
Artemisia Absinthium—	dried herb, a year old, Germany	-	-	-	-	-	Bl.	8.75
Artemisia Absinthium—	dried herb, 3 years old, Germany	-	-	-	-	-	Ma.	5.0
Artemisia —?	Wormseed of commerce	-	-	-	-	-	Ra.	3.0
Artemisia —?	Levant wormseed	-	-	-	-	-	Vo.	10.8
Calamus aromaticus—	fresh root, Germany	-	-	-	-	-	Ma.	16.0
Calamus aromaticus—	recently dried, Germany	-	-	-	-	-	Bl.	17.5
Calamus aromaticus—	long dried, Germany	-	-	-	-	-	Da.	14.3
Carum Carui—	dried fruit of French commerce	-	-	-	-	-	Ra.	50.12
Carum Carui—	dried fruit of German commerce	-	-	-	-	-	Ma.	66.5
Carum Carui,	do. do.	-	-	-	-	-	Da.	46.6
Carum Carui,	do. do.	-	-	-	-	-	Vo.	70.0
Caryophyllus aromaticus.	Cloves, Bourbon	-	-	-	-	-	Ra.	144.0
Caryophyllus aromaticus.	Cloves, Cayenne	-	-	-	-	-	Ra.	152.0
Caryophyllus aromaticus.	Cloves, do.	-	-	-	-	-	Bl.	125.0
Caryophyllus aromaticus.	Cloves, Molucca: French commerce	-	-	-	-	-	Ra.	148.0
Caryophyllus aromaticus.	Cloves, do. English commerce	-	-	-	-	-	Ra.	112.5
Caryophyllus aromaticus.	Cloves, average, German commerce	-	-	-	-	-	Vo.	226.0
Caryophyllus aromaticus.	Cloves, finest, do.	-	-	-	-	-	Da.	250.0
Caryophyllus aromaticus.	Cloves; German commerce	-	-	-	-	-	Steer.	272.0
Cinnamomum zeylanicum—	cinnamon of commerce	-	-	-	-	-	Ra.	1.56
Cinnamomum Cassia—	cassia bark of commerce	-	-	-	-	-	Ra.	12.0
Citrus Aurantium—	Sweet orange flowers, 1 May, Nice	-	-	-	-	-	Ra.	5.0
Citrus vulgaris—	Bitter orange flowers, 7 May, Nice	-	-	-	-	-	Ra.	5.9
Citrus vulgaris,	do. do. 12 May, Carmet	-	-	-	-	-	Ra.	4.12
Citrus vulgaris,	do. do. 16 July, Paris	-	-	-	-	-	Ra.	0.9
Citrus vulgaris,	do. do. 14 Dec., Paris	-	-	-	-	-	Ra.	6.5
Citrus Aurantium—	rind of 100 oranges, by expression	-	-	-	-	-	Ra.	2.5
Citrus Aurantium,	do. by distillation	-	-	-	-	-	Ra.	2.75
Citrus vulgaris—	rind of 100 oranges, by expression	-	-	-	-	-	Ra.	4.0
Citrus vulgaris,	do. by distillation	-	-	-	-	-	Ra.	4.25
Citrus Limetta—	rind of 100 limes, by distillation	-	-	-	-	-	Ra.	2.12



Citrus Bergamium—rind of 100 bergamots, by distillation	-	-	-	Ra.	2.9
Citrus Limonum—rind of 100 lemons, by expression	-	-	-	Ra.	1.9
Citrus Limonum, do. by distillation	-	-	-	Ra.	1.4
Cochlearia Armoracia—fresh seeds	-	-	-	Ra.	0.9
Coriandrum sativum—dry fruit of French commerce	-	-	-	Ra.	2.3
Coriandrum sativum—dry fruit of German commerce	-	-	-	Da.	9.0
Croton Eleutheria—cascarilla bark	-	-	-	Bl.	5.62
Cuminum Cyminum—dry fruit of French commerce	-	-	-	Ra.	44.0
Cuminum Cyminum—dry fruit of German commerce	-	-	-	Bl.	32.5
Daucus Carota—dry fruit	-	-	-	Ra.	0.66
Daucus Carota—fresh root	-	-	-	Ra.	0.14
Dracocephalum moldavicum—flowering herb	-	-	-	Ra.	2.10
Drimys Winteri—Winter's bark (probably, however, only <i>Canella alba</i> )	-	-	-	Ra.	0.50
Eugenia Pimenta—pimenta berries, <i>Jamaica</i>	-	-	-	Ra.	12.38
Fœniculum officinale—dry fruit of French commerce	-	-	-	Ra.	33.0
Fœniculum officinale—dry fruit of German commerce	-	-	-	Ma.	56.6
Fœniculum officinale, do. do.	-	-	-	Bl.	83.0
Fœniculum officinale, do. do.	-	-	-	Da.	60.4
Fœniculum officinale—flowering herb, <i>Grasse</i>	-	-	-	Ra.	4.9
Fœniculum officinale—herb after flowering, <i>Grasse</i>	-	-	-	Ra.	6.0
Galipea officinalis—Cusparia-bark of commerce	-	-	-	Ra.	1.5
Genista canariensis—Rhodium-wood	-	-	-	Ra.	0.47
Geum urbanum—dry roots	-	-	-	Ra.	0.53
Hyssopus officinalis—flowering herb, <i>Grasse</i>	-	-	-	Ra.	5.30
Illicium anisatum—star-anise-fruit	-	-	-	Ra.	34.21
Illicium anisatum—star-anise-fruit	-	-	-	Da.	25.5
Juniperus communis—green berries, 12 Sept.	-	-	-	Ra.	3.9
Juniperus communis—ripe berries, 1 Dec., <i>France</i>	-	-	-	Ra.	7.75
Juniperus communis, do. fresh, <i>Germany</i>	-	-	-	Do.	15.5
Juniperus communis, do. a year old, <i>Germany</i>	-	-	-	Ma.	10.8
Juniperus communis, do. a year old, <i>Germany</i>	-	-	-	Bl.	16.25
Juniperus Sabina—fresh twigs, 5 March, <i>Grasse</i>	-	-	-	Ra.	19.05
Juniperus Sabina—fresh twigs, 2 Oct., <i>Paris</i>	-	-	-	Ra.	14.25
Juniperus Sabina—dried twigs, recent, <i>Germany</i>	-	-	-	Ma.	40.0
Juniperus Sabina—dried twigs, a year old, <i>Germany</i>	-	-	-	Ma.	25.0
Larix Cedrus—fresh cedar wood, <i>Paris</i>	-	-	-	Ra.	0.3
Larix Cedrus—cedar wood of commerce	-	-	-	Ra.	4.25
Laurus nobilis—fresh leaves, 26 Jan., <i>Paris</i>	-	-	-	Ra.	5.25
Laurus nobilis—leaves some years dried, <i>Germany</i>	-	-	-	Bl.	4.10
Laurus nobilis, { fresh leaves, } poor soil, low site	-	-	-	Chr.	7.33
Laurus nobilis, { early in Oct. } poor soil, high site	-	-	-	Chr.	6.9
Laurus nobilis, { near <i>Edinb.</i> } very fine soil, low site	-	-	-	Chr.	17.12
Lavandula vera—flowering herbs, 2 Aug., <i>Grasse</i>	-	-	-	Ra.	11.5
Lavandula vera, do. 2 Aug., <i>Grasse</i> , north exposure	-	-	-	Ra.	9.12
Lavandula vera—flowering herb, 26 July, <i>Soureillas</i>	-	-	-	Ra.	9.0
Lavandula vera—herb after flowering, 26 Sept., <i>Soureillas</i>	-	-	-	Ra.	15.0
Lavandula Spica—fresh herb, 24 July, <i>Paris</i>	-	-	-	Ra.	7.62
Lavandula Spica—fresh herb, 4 Aug., <i>Grasse</i>	-	-	-	Ra.	12.5
Lavandula Stoechas—dried spikes	-	-	-	Ra.	6.43
Ligusticum Levisticum—fresh herb, <i>Paris</i>	-	-	-	Ra.	1.12
Melissa officinalis—fresh flowering herb	-	-	-	Ra.	0.25
Mentha piperita—fresh tops in flower, <i>Grasse</i>	-	-	-	Ra.	6.25
Mentha piperita—fresh tops in flower, <i>Paris</i>	-	-	-	Ra.	3.40
Mentha piperita—dried tops in flower, <i>Germany</i>	-	-	-	Bl.	15.62
Mentha piperita—dried tops in flower, <i>Germany</i>	-	-	-	Ma.	21.0
Mentha Pulegium—fresh flowering herb	-	-	-	Ra.	1.0
Myristica moschata—mace of commerce, finest	-	-	-	Vo.	154.0
Myristica moschata, do. fine	-	-	-	Bl.	125.0
Myristica moschata, do. worm-eaten	-	-	-	Bl.	65.6
Myristica moschata—nutmegs of commerce, fine	-	-	-	Bl.	108.25
Myristica moschata, do. worm-eaten	-	-	-	Bl.	64.1
Myrtus communis—fresh leaves, September 20, <i>Grasse</i>	-	-	-	Ra.	4.5
Myrtus communis—fresh leaves, September 6, <i>Paris</i>	-	-	-	Ra.	2.5
Origanum Majorana—fresh flowering herb, August 3, <i>Grasse</i>	-	-	-	Ra.	8.5
Origanum Majorana do. August 3, <i>Paris</i>	-	-	-	Ra.	4.4
Origanum vulgare do. September 15, <i>Paris</i>	-	-	-	Ra.	0.4
Pimpinella Anisum—dry fruit of French commerce	-	-	-	Ra.	35.12
Pimpinella Anisum—dry fruit, new, German commerce	-	-	-	Ma.	37.5
Pimpinella Anisum—dry fruit, old, do.	-	-	-	Ma.	27.0
Pimpinella Anisum—dry fruit of do.	-	-	-	Vo.	25.0







the seeds too much; this is peculiarly the case with linseed oil, which is seldom seen of a light color.

*By Decoction.*—This is effected by boiling the bruised seeds in water, and skimming off the oil as it rises to the surface. This plan is also employed in making cod-liver oil, as well as in the manufacture of castor oil in the West Indies.

*By Solution.*—This affords a very pure oil, but is seldom employed on account of expense. It is, however, useful in facilitating the extraction of some of the thick and viscid oils. Thus, croton oil is more readily obtained by mixing the ground seeds with half their weight of alcohol, and, after letting the mixture stand for some time, submitting it to pressure, and distilling off the spirit from the product.

Solution in ether is also useful in some cases, when expense is not an object; oil of ergot is best obtained in this manner.

The solid oils, as oil of cocoa, &c., require the aid of heat in their expression, and the ground material containing them is therefore placed between heated plates in the press.

*Animal Fats.*—Formerly, the fats of many animals were employed in pharmacy; but, at present, those principally used are lard, suet, and beef's marrow. To render these fit for pharmaceutic purposes, the crude material is to be cut in small pieces, freed as much as possible from all extraneous substances, and placed in a boiler with water, and heated until it is fused; the heat is continued till the water has evaporated, and the fluid fat is then strained. These fats are best preserved by being run into glazed jars, and kept from the action of the air.

**ALKALOIDS.**—The vegetable alkaloids may be obtained by a variety of processes; but these invariably comprehend decomposition of the alkaloidal salt in the crude drug, either by the superior affinity of an alkali, earth, or alkaline carbonate, or by double decomposition with some compound salt whose base forms an insoluble salt with the acid in the drug. Active neutral principles—such as narcotin from opium, piperin from white pepper, picrotoxin from *cocculus indicus*, and elaterin from *elaterium*—may be obtained through the agency of such simple solvents as water, rectified spirit, and sulphuric ether, used singly or successively. But no alkaloid can be thus detached.

The solutions from which vegetable alkaloids are prepared are, in general, obtained best by the method of percolation, or displacement. Some experience is required to apply this process in all cases with success. But, when well performed, it is greatly superior in general to any other mode of extracting the active matters of vegetable drugs, in cases in which the liquid used is spirituous or ethereal; and it is often not less advantageous in the instance of water, as well as acidulous fluids. The precautions for applying it successfully has been considered under the head of tinctures.

**SPIRITS** are alcoholic solutions of volatile principles obtained by distillation.

When spirit is distilled with aromatic vegetables which contain volatile oil, the oil, for the most part, rises with the spirituous vapor, and condenses along with it in a state of solution. In some cases, the volatile oil rises with the vapor of strong spirit, so that alcohol may be employed for the purpose; and this is occasionally necessary for keeping the oil of the distilled spirit in solution. In other instances, the oil does not begin to pass over until watery vapor also passes in considerable proportion with the spirituous vapor; so that diluted alcohol is required for the process.

For most medicinal purposes, but especially the present, the menstruum should be made by diluting alcohol with water.

The best apparatus for preparing distilled spirits is that represented at page 118, fig. 115, Procter's edition of *Mohr and Redwood's Practical Pharmacy*. It has been proposed to prepare them by distillation in a vacuum-still; but many volatile oils will not rise with spirit-vapor at the low temperature at which spirit boils in a vacuum.

These preparations, like distilled waters, may be obtained either from crude vegetable substances, or from their volatile oils. They are seldom, however, of such fine aroma when prepared in the latter as in the former way, unless care be taken to employ those volatile oils only which have been recently as well as carefully distilled.



*Essences* differ from spirits in being volatile substances dissolved in alcohol; but this name is generally applied to solutions of the volatile oils.

TROCHES, OR LOZENGES, are small, dry, solid masses, consisting of powders incorporated with sugar and mucilage. Some writers include under this name *drops* and *pastes*, making the following distinction between them:—

*Lozenges*, when the principal basis is sugar, and when the ingredients are combined without the aid of heat.

*Drops*, when the principal basis is sugar, but when the ingredients are combined with the aid of heat.

*Pastes*, when the principal basis is principally a vegetable juice or pulp, and when the mixture is of a soft consistence.

Lozenges are much more employed in Europe than in this country as a mode of administering medicines. In making them, the sugar is employed in a powdered state; the more active ingredients added in powder, or in a liquid state; and the whole mixed into a paste by the addition of mucilage. The mucilage generally used is that of tragacanth, as being more tenacious than that of gum Arabic. After the ingredients are properly incorporated, the paste is rolled out into a uniform sheet on a marble slab, previously sprinkled with some powdered starch to prevent adhesion, and cut into small cakes by means of a punch. These cakes are then to be placed on sieves, and kept in a drying-room until they have become perfectly dry and hard, when they are to be sifted, and kept in well-closed bottles.

Drops are formed from coarsely-powdered sugar, to which the flavoring or medicinal ingredients are added in a liquid state, so as to moisten the sugar. The mixture is then melted in a proper vessel, over a clear fire, after which the melted mass should be allowed to drop in small portions on a marble slab or greased metallic plate, and, when cold, removed, and kept like lozenges.

Pastes are usually formed of inspissated vegetable juices, or decoctions, with the addition of sugar, gum, &c. These are to be brought to a proper consistence, and the mixture poured into flat, shallow moulds, or rolled out, as in the case of lozenges, and divided into pieces of the desired size.

INHALATIONS.—These have, at different times, been much in vogue as a means of combating disease, especially complaints of the respiratory organs. Within the last few years, much interest has been excited relative to this mode of administering medicinal agents, from the remarkable anæsthetic effects caused by the inhalation of the vapor of ether and chloroform. The simplest form of using either of these is by pouring the requisite quantity on a hollow sponge, and applying this over the mouth of the patient. Various forms of inhaling apparatus have been invented, which will be found described in *Mohr and Redwood's Pharmacy*, 535–9, with directions for their use.

## EXTERNAL REMEDIES.

BATHS.—By the term bath is meant the complete or partial immersion of the body in a fluid or gaseous medium, differing, in some circumstances, from that to which it has been accustomed. In order to obtain the full efficacy of a bath, without injurious consequences, its temperature should be regulated, and particularly prescribed, by the physician ordering it. The following temperatures are those given by Dr. Forbes in the *Cyclopædia of Practical Medicine*:—

Cold Bath	-	-	-	-	-	between	33° and 60° F.
Cool Bath	-	-	-	-	-	"	64° and 75° F.
Temperate Bath	-	-	-	-	-	"	75° and 85° F.
Tepid Bath	-	-	-	-	-	"	85° and 92° F.
Warm Bath	-	-	-	-	-	"	92° and 98° F.
Hot Bath	-	-	-	-	-	"	98° and 112° F.



The effects of these different baths on the system are very dissimilar, according to their temperature, and the time during which the patient is subjected to their influence.

**COLD BATH.**—When a person plunges into a cold bath, he is first sensible of a sudden sensation of cold upon the surface, accompanied by an oppression of breathing, causing this function to be performed in convulsive gasps. This is called the *shock*, and is caused by a rapid contraction of the cutaneous capillaries, and a retrocession of the blood to the lungs and other internal organs. In a short time, the difficulty of breathing disappears, the temperature becomes agreeable, and if the person now leaves the water, a warmth of the surface comes on, termed the *glow*, succeeded by a sense of invigoration of the whole system. Should the person remain in the water for too long a time, another train of symptoms manifest themselves: the sensation of cold soon attains to an unpleasant degree of chilliness, followed by rigors; a bluish tint is perceptible on the surface of the body; the blood accumulates in the internal organs; and, on leaving the water, there is no reaction, or a very feeble one, the surface remaining cold, the extremities benumbed; and headache, difficult respiration, often pain in the chest, ensue, with a sense of depression and lassitude. The use of proper means will often remove these symptoms; but they may lead to a variety of diseases of the internal organs.

The objects, in prescribing a cold bath, are the production of a sudden and powerful impression on the nervous system, and the tonic influence it exercises when followed by due reaction. In the first of these, it has been found useful in certain affections where there is a derangement of the functions of sensation, of motion or sensation unattended with a congestive or inflammatory condition of the internal organs; but it is more frequently ordered to fulfil the second indication.

It is always contra-indicated when, from debility, the system does not react so as to produce a glow; when there is a tendency to congestion of the cerebral vessels, or any serious organic affection of the heart, lungs, or kidneys. In all cases, it is advantageous, before taking the cold bath, to take such exercise as will raise the circulation, without occasioning fatigue or perspiration, for reaction is almost certain to follow the immersion, except when the person has remained in the water too long a time. The period of immersion should not exceed five minutes.

**COOL BATH.**—The action and uses of this are similar to the last, but are less powerful. It is, therefore, better calculated for those who are much debilitated.

**TEMPERATE BATH.**—As the temperate bath is of a temperature closely approaching that of the body, the shock and subsequent reaction are almost wanting. It is, therefore, much more employed for purposes of comfort and cleanliness than as a remedial agent. In delicate persons, it should always be used instead of the cold or cool bath, and is always better suited to very young children than lower temperatures.

**TEPID BATH.**—This is intermediate in operation between the temperate and the warm bath, and varies in effects and uses according to the temperature. In perfect health, it should not be used as an habitual indulgence; but, for the purposes of cleanliness, an occasional recurrence to it allows of a more perfect ablution than can be effected by cooler baths. It is better to use it about noon, when the first process of digestion of the morning meal is over, and immediately afterwards to take brisk exercise in the open air. In cases of fatigue and febrile irritation, from over-exertion or a long journey, the tepid bath is generally found very beneficial. It is also serviceable to persons of sedentary habits, &c. In all such cases, however, it is not to be employed immediately after a meal, or when the individual is unduly excited, either mentally or corporeally. In one class of complaints, those dependent on gastric irritation, the tepid, or even the warm bath proves of much service.

**WARM BATH.**—The first effect of a warm bath is to produce a sensation of heat upon the surface, and to increase the pulse in quickness and fulness, though in most cases to diminish its tenseness. The cutaneous circulation more especially becomes affected,



and the body is increased in bulk, as shown by the increased pressure of ligatures, or of rings upon the fingers. The secondary effects, when the immersion is continued for some time, are muscular relaxation, sometimes to a considerable degree; even after leaving the bath, a disposition to lassitude continues for some time, with a tendency to perspiration.

The remedial effects of a warm bath depend on its temperature, the time a patient remains in it, and the subsequent treatment. The medium time for remaining in the bath is from twenty to twenty-five minutes; but this must be regulated by the effect produced. It is beneficial in incipient catarrh, in some congestions of the internal organs, chronic rheumatism, and in spasmodic affections, especially those of children; but is contra-indicated in active fever, or when there is congestion, or a determination of blood to the head.

In the convulsions of children, its effects are remarkably beneficial, as it not only relaxes spasm, and relieves for the moment, but soothes nervous irritation. In cases where the convulsions are severe, it will be found advantageous to apply cold water to the head. When a warm bath is administered to a child, care must be taken not to expose it to the cold air for the purpose of drying its body; the best plan is to envelop it in a warm blanket, and to place it in bed at once. By this plan, it is not liable to take cold, which is a common objection to the use of the warm bath for children.

**HOT BATH.**—From its temperature being above that of the body, the hot bath is far more stimulating than the preceding, as evinced by the excitement of the pulse, the sensations of fulness in the head and throbbing of the cerebral vessels. Its use is principally confined to cases where it is wished to arouse nervous energy and vital action, as in Asiatic cholera, &c.; or where there is a sudden retrocession of cutaneous diseases. It has also proved useful in certain forms of rheumatism, and paralysis. As the intention is mainly to induce excitement, the patient is not to be exposed to its action long enough to cause exhaustion.

Whatever description of bath is ordered to be used, the original temperature is to be maintained during the whole time the patient remains in the water. At the end of some minutes, therefore, the heat should be tested by a thermometer, and, if requisite, hot water added. The sensations of the bather are always a fallacious criterion.

**SHOWER BATH.**—This is a modification of the cold affusion, being attended with the same effects, but in a less degree. The short duration of it renders it less refrigerant than the cold bath, and causes the primary shock to be the most important part of the influence it exercises, which may be modified by increasing or diminishing the temperature of the water, or that of the height from which the fluid falls. It is used either cold or tepid. The former is most advantageous when the powers of the system are sufficient to cause a reaction, that results in a glow on the surface, soon after using the bath; but, where this is not the case, as in debilitated individuals, the temperature is to be increased. It is adapted for those cases where a powerful impression is wished to be made upon the nervous system, as in chorea, hysteria, &c., and is also of much benefit to persons subject to determination to the head. In such cases, it is found advantageous to make the patient stand in hot water at the time of taking the bath. The reaction following its use is much promoted by using friction to the surface. The improved and portable shower baths, now to be procured, obviate many of the objections to its use which formerly existed. For children, the best and most convenient apparatus that can be employed is one described by Dr. A. T. Thomson. (*Domestic Management of the Sick Room.*)

“It consists of a hollow vessel made of tin, with a perforated bottom. The body of the vessel is of a bell-shape, with a hollow tube rising from the top, and terminating in a broad perforated rim. When the bath is to be used, it must be sunk in a bucket of water, until it is completely submerged; the air is thus driven out of the bath, through the tube, and the bath filled with water. The thumb of an attendant is then to be placed on the opening in the centre of the rim, and the bath raised from the bucket of water. The pressure of air upon the holes in the bottom retains the



water in the bath; and, on raising the thumb from the upper orifice, the whole is rapidly discharged. In using it, the child must be placed in an empty tub, and the bath, being held over his head, is then to be discharged; and the child immediately afterwards dried, with friction. When salt water is used for this bath, the hair should be kept dry, by means of an oil skin cap."

**LOCAL BATHS.**—These are most frequently used warm. The most common are the *hip-bath* and *foot-bath*. They are principally employed as revulsives. The first has been found a valuable remedy in diseases of the womb, and in irritations of the pelvic organs. Where it is employed merely to soothe pain, the temperature should not exceed 80° to 90°, and the patient is to remain in it for some time; but when it is intended to excite the uterus to greater action, it should be as hot as can be borne by the patient, though the continuance in it should not exceed ten to fifteen minutes. The hot foot-bath should be of as high a temperature as can be borne, so as to redden the skin of the immersed parts effectually. The vessel used should be sufficiently deep to allow the legs to be immersed nearly to the knees. It is a valuable remedy in the early stages of catarrh, and local congestions of the head, chest, or abdomen; and in the dyspnœa accompanying hypertrophy of the heart, it has often proved very beneficial. It may be rendered more stimulating by the addition of common salt, carbonate of potassa, or flour of mustard. In torpid states of the liver, a mixture of nitric and muriatic acids, in the proportion of about an ounce of each to every gallon of water, often proves highly useful.

**VAPOR BATH.**—This consists in either wholly enveloping the patient in the steam of hot water, or merely his body, or some of its parts. A much more elevated temperature can thus be borne than where the body is immersed in water, and its action is more confined to the skin; hence, although less stimulating, it is more diaphoretic than the hot bath. Where the vapor is inhaled, the heated surface being extended, it is more powerful in its effects. It has proved of great benefit where an active revulsion to the surface is indicated, and also in the treatment of cutaneous diseases. It can be applied with little trouble. All that is required is that the patient be seated on a chair with a vessel of hot water placed beside him, and the whole enveloped with a blanket, to be thrown over his head if the vapor is to be breathed, or pinned around his neck where this is not the case. The steam soon surrounds his body, and causes a copious perspiration; and, should it cease too soon to be evolved, its generation may be restored by dropping a heated brick or stone into the water. Where the vapor is not respired, it may be used of a higher temperature.

**WARM AIR BATH** is more stimulating than the vapor bath, producing—especially where the warm air is also breathed—general uneasiness, heat of skin, excitement of the pulse, until a general perspiration ensues. It is readily administered even to persons in bed. The bedclothes are to be elevated by a light frame, and the heated air admitted by means of a tube. It has been found beneficial in chronic rheumatism, and was at one time much praised in the collapsed stage of cholera.

**DOUCHES** consist in the forcible application of a stream of hot or cold water to a part of the body, in such a manner that the force of the stream of the fluid shall aid the stimulant effect. The hot douche may be formed by pouring from a height hot water from a tea-kettle, or through a tube about an inch and a half in diameter, so as to cause the fluid to strike forcibly upon the affected part of the body, whilst at the same time percussion is employed; or, in other words, the part is beaten by means of an Indian rubber bottle, stuffed with wool, and affixed to a cane handle.

A temperature of 160° is readily borne in douching. When the disease is chronic rheumatism, or when deep-seated pains are to be removed, the douching should be continued for either half or three-quarters of an hour at a time; and the patient should be afterwards placed in bed between blankets, in order to encourage sweat-



ing. The most decisive advantage, in enlarged, stiff, and painful joints, results from this mode of douching.

In cases of complicated fevers, and in some other diseases, in which inflammation of the brain occurs, the *cold douche* is frequently ordered. The mode of applying it consists in pouring a stream of cold water on the shaved head. The patient should be raised in bed, and the head held over a basin, or other vessel, whilst a stream of cold water is directed on the crown, and the fall of the water gradually increased. After this, the head should be merely dried, but not rubbed. It not only soothes the patient, but often induces sleep, although it usually causes a most uncomfortable feeling at first; but relief is so quickly experienced, that its repetition is generally requested by the invalid.

A *vapor douche* is very stimulating, and when of a high temperature, and continued for any time, will cause vesication. It may be used as a counter-irritant instead of moxa.

MEDICATED BATHS are such as contain medicinal agents in solution in the water, and are used both topically and generally. Saline, acid, and alkaline substances, with iodine and sulphur, are the materials most commonly employed. Medicated baths are sometimes natural, as those of sea water, and certain mineral springs; others consist of solutions of various articles, artificially prepared. Of the medicated baths, sea water is the most generally used, more, however, for pleasure, than for remedial purposes. It is not only stimulating, on account of its saline constituents, and its use followed by a perfect glow, but these constituents also exert a medicinal action on the system, in certain diseases of debility. The nitromuriatic bath has been found useful in diseases of the liver; and alkaline, ioduretted, and alkaline sulphuretted baths, are advantageously employed in many forms of cutaneous disease.

Medicated *vapor* baths are prepared by impregnating aqueous vapor with the volatile principles of medicinal plants, though it is very doubtful whether they produce any effects that cannot be obtained from the simple vapor bath. But the fumes of sulphur, chlorine, camphor, mercury, &c., in combination with aqueous vapor, have been found to exert powerful effects on the system, and to be very effectual in the cure of certain diseases.

AFFUSION.—This consists in pouring or dashing a quantity of water on the body, or a portion of the body, from an inconsiderable height, most generally for the purpose of reducing its temperature; but it is also employed as a revulsive agent, and to give an impulse to the nervous system. Affusions are made with cold or tepid water. Cold affusion is one of the most powerful general means of abstracting caloric from the body, we possess; it not only lessens the temperature of the surface, but it diminishes the action of the heart. It has been successfully employed in the treatment of fevers, but its use requires much judgment, as respects the time of its application, which, according to Currie and others, is, "when there is no sense of chilliness present, when the heat of the skin is steadily above what is natural, and when there is no general or profuse perspiration." It should not be employed either in the cold or sweating stage of fever, or in the hot stage where the heat does not much exceed the natural standard. From inattention to these circumstances, and from patients being injured by its effects, affusion is not at present often resorted to in the treatment of fevers. In scarlatina and some other of the exanthemata, it has been employed with benefit, to reduce the morbid heat of skin. It is, however, more relied upon in inflammatory diseases of the brain; in these cases, the water is to be poured upon the head, inclined over a basin or tub, by means of a pitcher. In children, it is sufficient to squeeze a large sponge, previously saturated with cold water, at some height above the head. Where a general affusion is deemed necessary, the patient should be taken out of bed, his head having been previously shaved; and, being stripped naked, and placed on a stool in an empty tub, from *three to five* gallons of water, at a temperature not under 40° Fahr., should be thrown over him. This affusion should be repeated until he feels cold, or rather until a rigor or shivering, or even chattering of the teeth, is



experienced. He should then be dried, placed in bed, and a little warm wine and water administered to him, to aid the reaction and the consequent perspiration.

But in cases where the invalid faints on raising him into the erect position; where he feels chilly, although the thermometer indicates a high febrile temperature; where the skin is bedewed with perspiration; where the patient is a woman, and the monthly change is present; or where the operation is dreaded, it ought not to be employed. The evening is the best time for using the cold affusion.

*Cool* affusion is sometimes used instead of the cold, and is preferable for weak, irritable individuals. *Tepid* affusion is also beneficial in certain cases; especially where there is a fear that perfect reaction will not take place after the application of cold water; or where disease of the pulmonary organs exists. It has been found very efficacious in scarlatina, as also in hectic fever. Warm affusion has likewise been used; but its effects are far more temporary than those of the warm bath.

**SPONGING.**—The object of sponging the body, either with cold or with tepid water, is to reduce the heat of the surface by means of evaporation, with the view of softening the pulse and diminishing its frequency; of rendering the skin perspirable; of relieving headache, checking delirium, and promoting sleep. In order to perform it well, the patient should be taken out of bed; and, being undressed and placed in a chair, two or three persons should, at the same time, pass large sponges, wet with simple cold or tepid water, or vinegar and water, rapidly over different portions of the naked body, until the whole of it shall have been successively sponged, and a chill has come over the patient, who is then to be dried and placed in bed.

This appears to be a simple process; but it requires both caution and judgment to determine on its use, and to render it not only salutary, but free from danger. For example, if, notwithstanding the continuance of great heat and dryness of the skin, a sensation of chilliness comes over the patient when the sponging is about to be done, the operation should be delayed; neither should it be done if the smallest tendency to perspiration, indicated by moisture in the axillæ, or on the palms of the hands, displays itself. But, if it has been done, and if partial relief has been obtained, if the heat of the skin returns, and the pulse rises, it may again be resorted to and repeated.

But, besides the advantages to be obtained from *sponging*, during the presence of disease, it is also productive of great benefit in warding off diseases, or in a prophylactic point of view. The sponging, however, under such circumstances, is to be performed in a different manner from that which has been described.

In those predisposed to asthma and to bronchitis, known in ordinary language by the term *cold*, or *cough*, and in those, also, who are predisposed to *croup*, nothing is so serviceable in warding off the paroxysm of the former disease, or in preventing the attack of the latter malady, as cold and tepid sponging of the chest and the trunk of the body. It is also equally serviceable during dentition in infants, and as a general tonic in delicate conditions of the habit. In these cases, it should be followed by gentle friction over the surface of the body.

Sponging the body, as a prophylactic, or as a tonic, is always most efficacious when it is performed whilst the patient is still in bed. Let him sit up; or, if unable to do so, let him be raised into the sitting position, keeping the lower limbs under the bed-clothes. The night-shirt or night-dress is then to be stripped down, and a large towel or sheet put round the waist, to prevent the bed-clothes from being wetted. The naked trunk of the body, and the upper part of the arms, are then to be moistened with a sponge, or a piece of flannel, dipped either in cold or tepid water, or vinegar and water, or salt water, as the case may demand; after which, in drying the skin, if water only has been used, brisk friction should be employed, so as to cause a general glow upon the surface. In some cases, salt and water, or vinegar and water, are preferable to the simple water. The best proportions are two ounces of *salt*, or the same quantity of common *vinegar*, to a pint of water. When these adjuncts are used, friction is not required; indeed, when salt water is employed, the skin should be dried with a soft, old towel, fitted to soak up the water only, and to leave the saline particles upon the surface.

The best time for using such spongings is just before getting out of bed in the



morning. They may be used in winter as well as in summer. In efficacy, they are certainly nearly equal to the shower-bath, which may be resorted to in vigorous habits; but, when the habit is delicate, and not equal to the reaction, on which the beneficial influence of the shower-bath rests, sponging is preferable. It should not be used when perspiration is present.

Sponging is as beneficial as the shower-bath, in rendering the habit less susceptible of cold; and, when conjoined with exercise in the open air, and proper regimen, not only asthma, coughs, and catarrhs, may be warded off by its employment, but the predisposition to tubercular consumption may be lessened, in a very remarkable degree, by its daily employment.

**FOMENTATIONS.**—These may be regarded as a species of local bathing. Although the decoctions usually employed are useful in aiding warmth, by their soothing or sedative influence, yet they are secondary objects in the application of fomentations, the intention being to convey heat, combined with moisture, to the part fomented. Flannel cloths, wrung out of boiling water, by means of two sticks turned in opposite directions, form the best fomentations. If they be shaken up, and laid lightly over the part, they involve a considerable quantity of air, which, being a bad conductor, retains the heat in them for a considerable time. In every process of fomenting, there should be two flannels, each three yards long, with the ends sewed together, to admit of the boiling water being wrung out of them; and the one flannel should be got ready whilst the other is applied. The fineness or the coarseness of the flannel is not a matter of indifference: the coarser it is, the less readily does it conduct heat, and the longer it retains its warmth; therefore, it is more efficient for fomenting. White flannel also retains the heat longer than colored flannel.

*Stuping* is a variety of fomentation useful in many cases, but especially in affections of the eyes. The patient should be his own operator. He should sit up in bed, and should place, in the hollow of his hand, a small piece of flannel, wrung out of boiling water, and hold the hand at such a distance under the eyes that the vapors may rise to them—changing the flannel as often as it ceases to give out warm vapor. If narcotic or sedative additions are directed, they should be poured hot upon the flannel, each time it is changed.

**CATAPLASMS, or POULTICES,** are modifications of fomentations; they generally consist of pulpy substances capable of absorbing much moisture, and of such consistence as to be applied accurately to any surface, however irregular. Their action, in most cases, depends upon the liquids with which they are moistened and the heat retained by the mass. Cataplasms may be emollient, medicated, or revulsive.

*Emollient Cataplasms* are usually made of bread and water, linseed meal, Indian meal, slippery elm, &c., and water; or of mashed vegetable substances, such as carrots, and marshmallows.

As warmth and moisture are the principal effects of emollient poultices, an excellent substitute for them is lint, dipped in warm water, and laid over the diseased part; the lint should be covered with a towel, in order to prevent evaporation. This mode is much neater and more cleanly than any poultice, and where merely warmth and moisture are required, it will answer all the purposes intended. A peculiar fabric, called *spongio-piline*, consisting principally of sponge, has lately been invented in England, which is said to answer very well, when properly imbued with hot water, as a substitute for the simple emollient poultice.

The addition of a little lard is useful, if the poultice does not require to be frequently changed. When the object, however, is to promote suppuration, the poultice should be frequently renewed, in which case the addition of the lard is unnecessary.

Poultices should never be heavy, nor very bulky, but they should be frequently repeated. They are useful, in all cases of inflammation which cannot be backed, to assist the suppurative process and the advancement of the matter to the surface. After an abscess is opened, the poultices should still be continued for a few days. Poultices may be used, also, as fomentations in colic, and in inflammation of the



bowels. In such cases, however, as the chief object is to allay pain and to soothe, they should be made with a decoction of poppy-heads, instead of water.

*Medicated Cataplasms* are those in which to the simple emollient poultice are added other ingredients, intended to exercise an influence on the part independent of mere warmth and moisture; as, for instance, the fermented poultice, in which yeast forms a constituent; the charcoal poultice, and various others enumerated in the Formulary, all intended to fulfil certain indications.

*Revulsive Cataplasms* are those which, by inducing a local irritation or inflammation of the parts to which they are applied, act as derivatives. To this class belong mustard poultices, or *Sinapisms*, Cayenne pepper, garlic poultices, &c.

Mustard poultices are made by mixing the flour of mustard, of the shops, with water, into the consistence of a paste. When it is wished to have a weaker preparation, wheaten, rye, or linseed flour is to be added to the mustard in such proportions as are deemed sufficient. The water used should be tepid, or merely warm; a boiling temperature is injurious, as it tends to coagulate the albumen of the mustard, and thus to prevent the extrication of the volatile oil, which is the active principle. Vinegar is often used, but never increases the energy of the preparation, and, in the case of the black mustard, diminishes its power. This poultice should be spread thickly on a rag, as soon as it is prepared, and covered with gauze or tissue paper, in order to prevent adhesion to the skin. In a few minutes after it is applied, the sinapism causes a sensation of heat, which increases to almost intolerable burning. After some time, the pain subsides, and is followed by a sense of weight and throbbing of the arteries. If the sinapism be left on, however, the burning pain again returns, and becomes so great, that few persons can bear the action of a mustard cataplasm longer than three quarters of an hour.

In delicate, sensitive people, the sinapism need not remain applied above six or eight minutes, as the effect continues some time after its removal. If it remain too long on, gangrene may supervene; this cannot happen unless the invalid be insensible, as too much pain is felt to permit the lengthened application of a sinapism; but, in cases of stupor or insensibility, or in low fevers, a sinapism, if left on longer than an hour, may cause both vesication and gangrene. If the redness only is required, the pain may be moderated by mixing with the paste a drachm of tincture of opium, or two drachms of tincture of henbane, neither of which, although they allay the pain, diminishes the rubefacient activity of the sinapism.

As a substitute for the mustard poultice, where mere external irritation is required, the capsicum or red pepper will answer admirably, and it has the advantage of not vesicating: it may be used as a poultice, by mixing the powder with bread and milk, or Indian meal, or as a lotion mixed with warm spirits.

**LOTIONS, LINIMENTS, EMBROCATIONS**, are medicines of a more or less fluid character, which are employed as external applications. They may be used as mere fomentations to soothe pain and remove inflammation, as a mode of introducing medicinal substances into the system, or as revulsive agents, according to their nature and composition. Liniments, which generally contain oily constituents, are usually aided by friction of the part to which they are applied, either with the hand alone, or with some article capable of exciting some irritation of the skin, as a piece of flannel or a soft flesh brush. In many cases, in which liniments are found beneficial, the advantage obtained from them is attributable rather to the friction than to any medicinal power in the lotion itself.

*Frictions*, whether simple or conjoined with liniments, are frequently important aids to medical treatment. Simple friction is performed by the hand alone, or with a piece of flannel, a hair glove, or a flesh brush. If it be properly performed—namely, by short, brisk strokes with the tip of the fingers, and with great celerity, when the naked hand is the agent; and if it be continued for an hour or upwards, and repeated several times a-day—its influence in reducing swelled glands and swellings of the joints, as well as in alleviating rheumatic pains, is very great; but, besides being well performed, the friction should be continued for half an hour, in order to render it useful.



Friction, when performed in a gentle, slow, and equable manner, by producing a continued repetition of an agreeable impression on the nervous system, acts as a soother of pain, independent of any aid from opiate liniments, and also induces sleep. The beneficial influence depends, in a great measure, on its transferring the attention of the sufferer from the seat of pain, and confining it to the mild and agreeable sensation impressed by the friction. In performing it, the operator should sit by the side of the bed, and, introducing the hand under the bed-clothes, rub the legs or the arms gently, with equally lengthened but slow movements. When the invalid is a child, its influence is more powerful when aided by a monotonous, but a soft tune, which, although it operates upon a distinct sense, yet, by combination, renders the friction more soporific.

When friction is intended to stimulate the surface, it is necessary to aid its influence by liniments containing ammonia, or camphor, or some other excitant or acrid substance: but these should be ordered by the medical attendant. In applying them, the same method should be adopted as when simple friction is employed; but the hand of the operator must be guarded by a glove from the influence of the acrimony of the liniment.

Friction is also used to introduce various substances—namely, mercurials, iodine, opium, &c.—into the body through the medium of the skin. The rubbing, in such cases, ought to be brisk, and applied with sufficient force to abrade slightly the cuticle; and continued until the substance, which is usually in the form of an ointment, disappears, owing to the absorption by the skin induced by the friction. In using such frictions, the hand of the operator should be guarded by a glove; otherwise, he is likely to suffer salivation, or some other unpleasant result, from the introduction of the medicinal agent into his own system.

VESICATORIES, or BLISTERS, have already been noticed in the introduction (p. 58), and do not require to be further commented upon.

ISSUES are small ulcers artificially established for the purpose of procuring and keeping up a discharge of pus. They form an important derivative remedy in many diseases, and are to be formed by the following methods: The simplest is by the application of a small blister, which is to be dressed with some epispastic or irritating ointment. It is, however, difficult to prevent the sore from healing, and the discharge is seldom sufficiently copious. They are also formed by making incisions through the skin, and inserting issue peas into the wound. The best and most effectual plan is by the use of caustic; this is done by means of a piece of leather, spread with adhesive plaster, through which a hole is made of the size of the intended issue; this is to be warmed and applied to the skin, so as to apply closely. The aperture is then to be filled with the caustic, in the form of a paste, and covered with another piece of leather, spread as above. These are to be left on until the skin is destroyed, and then to be removed and a poultice applied until the eschar separates. The late Dr. Wistar was in the habit of recommending the application of a small blister, and of rubbing the vesicated surface with caustic potassa for a few minutes. This is an expeditious and excellent method, as it is effectual and gives but little pain. The issue is kept open either by the introduction of peas, or a piece of lint smeared with the ointment of savine, into the wound. The peas operate by mechanical irritation; and, as they swell from the action of the heat and the moisture of the ulcer, they tend also to enlarge the wound. The peas should be changed at least once in twenty-four hours. Instead of the peas, it has been advised to use a small block of hard wood, with the lower surface cut into lozenges; these, pressing on the surface of the sore, occasion a copious discharge, and it will be found more convenient of application than peas. The very irritation, however, by which the discharge is maintained, is apt to cause fungous growths from the bottom of the wound, which should be kept down; this is to be done by dressing the issue occasionally with some escharotic, as burnt alum or red precipitate. Issues should be washed twice a-day with warm water; otherwise, they are apt to become very offensive.



SETONS are wounds kept in a suppurating state by means of some foreign body, which prevents their healing. They are made by passing a seton needle, armed with a skein of silk, thread, or a slip of gum elastic, through a folded portion of the skin, withdrawing the needle and leaving the silk in the wound. If a seton needle cannot be procured, the puncture may be made with a thumb-lancet or bistoury, and the silk passed through the wound by means of an eyed probe. The edges of the wound are apt, however, after it has been discharging for some length of time, to become callous, and the discharge to cease. When this occurs, the irritating substance should be smeared with some stimulating ointment. It should be recollected that much mischief may result from suddenly suppressing or removing either an issue or a seton. If this takes place, purgatives should be administered, and the discharge of the issue or the seton checked as gradually as possible.

OINTMENTS are simple or compound fatty bodies of a soft consistence, which are applied to the skin by inunction. They are more consistent than liniments, but softer than cerates. Simple ointment consists of one part of wax and four of lard, melted together and thoroughly mixed. This preparation is the basis of many compound ointments. Compound ointments are of three kinds: one consisting of simple ointment, combined mechanically with various substances; the second, formed by dissolving the active ingredients in melted ointment or lard; the third, made by uniting substances to the fatty matters that induce a chemical action between them.

Compound ointments of the first class are mixtures of the fatty basis with various solid and liquid substances; these are incorporated by trituration in a mortar, or on a stone slab. When hard substances enter into the combination, they require to be previously reduced to a fine powder, or, if they are soluble, to be rubbed into a paste with an appropriate menstruum, and then admixed. When extracts form constituents of an ointment, if they are not of a soft consistence, they should be rendered so by trituration with water before they are mixed with the fatty body. The mixture of two or more ointments may be effected by rubbing them together on a stone slab by means of a spatula.

Compound ointments of the second kind are made by boiling recent vegetable substances in the fatty basis, until all the water of vegetation is driven off; the heat should then be reduced to about that of boiling water, when the ointment is to be strained. Those of the third kind, as citrine ointment, are prepared by adding the constituents gradually together, aided by heat and constantly stirring.

Ointments are preserved by keeping them in glazed jars, with the surface covered with tin foil. According to M. Deschamps, the admixture of a twenty-fifth part of benzoin with the fatty matter used to make the ointment, will prevent or greatly retard the process of decomposition (see p. 128). In the case of simple ointment, and those for highly irritable surfaces, this plan answers well; where it is inexpedient, the ointments should be renewed at short intervals.

CERATES consist of a basis of wax and a fatty matter, with which other substances are incorporated. They are of such a consistence as to be softened, but not melted, by the warmth of the body. The general directions for making them are similar to those for ointments.

Many preparations of the present class are made by simply mixing or triturating their component parts together. But when resins, wax, spermaceti, or concrete oils are to be mixed either together or with fluid oils, it is better to unite them with the aid of heat, which not only liquefies the solids, but likewise renders them much more soluble in the liquids. Brisk agitation is required while the mixture concretes on cooling; otherwise, the solid ingredients will separate either mechanically or by crystallization. Even in purifying lard or suet by fusion and filtration, it is right to stir the mass as it cools; otherwise, the stearin or solid oil is apt to separate in part from the elain or liquid oil, so that an irregular mixture of the two principles is obtained. A gentle heat is sufficient for making ointments, cerates, and the like, because most of the ingredients are easily fusible; and a strong heat must be avoided, for in that case acrid acids are engendered, which may alter the properties of the preparation. Hence



the vapor-bath is generally used for applying heat; and it ought always to be employed in operations on a large scale.

When the cerate is intended to irritate and not to soothe, a high temperature may be employed, which, in some cases, is advantageous, as in the cerate of cantharides, by more effectually dissolving out the active principle.

PLASTERS are more consistent than cerates, adhesive at the temperature of the body, and requiring the aid of heat to soften them sufficiently to be spread. They are of two kinds: one consisting of a compound of olive oil and litharge, or an oleo-margarate of lead, having the common lead plaster as a basis, united to resinous or other substances; the other, composed wholly of resinous matters, or these mixed with fatty substances or wax.

In making plasters, in most cases, it is necessary to operate at a temperature that will not volatilize or carbonize the ingredients; hence, a water-bath or steam heat is preferred. In the case of the burnt plaster (*onguent de la mere*) of the French Codex, sufficient heat is employed to cause the fats to fume. Plasters should be made in metallic vessels, much larger than will contain the substances to be operated upon, as the mass increases in bulk by the action of heat causing an extrication of vapors. The water ordered in making lead plaster should be added at the commencement of the process; otherwise, when poured on the melted mass, it may cause a projection of portions of the heated materials on the operator. The ingredients should be thoroughly stirred together during the process of melting. When this operation is concluded, the plaster is to be removed, in small portions at a time, into a vessel constantly supplied with cold water; and, when sufficiently consistent, rolled into cylinders on a wet marble, and again placed in cold water to harden. To render plasters of a lighter color, they are kneaded and pulled under water; but this process should not be practiced on compound plasters containing substances soluble in water; nor should these plasters be cooled in water, but suffered to chill on a marble slab until fit to be made into rolls. Some of the officinal plasters belong to the second class, and are made by melting resinous matters with the addition of certain oleaginous substances.

Plasters are preserved by enveloping the rolls with paper, to exclude the air as much as possible. When kept for any length of time, they are apt to become hard and brittle, and to lose their color. When this is the case, they should be re-melted by a gentle heat, and sufficient oil added to restore their proper consistence.

Spreading plasters for use, requires skill and address on the part of the operator. They are spread on various textures, but most generally on linen or cotton cloth, or leather. The shape and size must be regulated by the part to which they are to be applied. The plaster is to be spread on the leather or cloth by means of a heated spatula of a peculiar form; this is brought in contact with the end of a roll, which, becoming liquefied, can then be evenly diffused over the surface. Where the plaster is sufficiently adhesive to maintain its place when applied to the body, it is to be spread so as to leave about half an inch of margin uncovered; but where it does not possess this quality in sufficient degree, it should be surrounded with an adhesive margin, which is to be prepared before spreading the body of the plaster. Full and particular directions for spreading plasters will be found in Procter's edition of *Mohr and Redwood*, p. 516, *et seq.*, and to which we would refer those who wish for clear and accurate directions on the subject.

FUMIGATIONS are extrications of vapors or gases, designed to modify the air of a sick chamber, and to mask any unpleasant odors that may be present, or to produce a medicinal effect on those parts of the body with which they are brought in contact.

Fumigations, for the purpose of obviating or masking unpleasant odors in a sick room, should never be employed to supersede ventilation and cleanliness; for most of them, instead of purifying the air, only render it more unfit for respiration. They are generally made by burning pastilles, sugar, juniper berries, benzoin, &c., so as to create an odoriferous smoke. As disinfecting agents, they are utterly useless; and are relics of an ancient custom of burning frankincense, and other odorous substances in vitiated air, to overcome the feter which is more or less present. They disguise unpleasant



odors; but they accomplish nothing more. The infection remains not only unaltered by the diffusion of the most powerful aromatic vapors, but its deleterious properties are sometimes augmented by them.

The fumes of burning camphor, and the vapors arising from its tincture, have more pretensions than either of the above to the name of a useful article of fumigation. It is much employed and confided in on the Continent; but the experience both of American and British physicians does not warrant the opinion that it possesses any power of destroying infection or contagion. Nothing is more ridiculous than the custom, which was at one time very general, and which is still continued to a certain extent—namely, that of carrying a camphor-bag about the person, as a protection against infectious diseases.

Vinegar is, not without reason, regarded as possessing some chemical influence in decomposing infectious and contagious matters; and, consequently, it is almost invariably sprinkled over the floor of the rooms of those suffering under infectious diseases; or the vapor of hot vinegar is diffused through their apartments. It is thought to be still more salubrious, and a more powerful disinfectant, when it holds camphor or aromatic oils in solution; hence the great popularity of the preparations called *Aromatic Vinegar* and *Thieves' Vinegar* (see p. 68). The repute of the latter is founded upon a story, that four thieves, who plundered the dead bodies during the plague at Marseilles, with perfect security, on being questioned respecting the cause of this impunity, confessed, on the condition of their lives being spared, that they attributed it solely to the use of *Aromatic Vinegar*.

Vinegar, in this state of combination, is extremely agreeable and refreshing, both to the invalid and the attendants of the sick room. The benefit which it produces depends upon a certain degree of stimulus imparted to the sensitive nerves, which are generally in a low condition in an infectious atmosphere; but, as a chemical agent, its powers are too feeble to be followed by much benefit. It is, however, always refreshing, and is much better adapted for overpowering the unpleasant odors of the sick room than any of the former substances.

The most efficacious fumigations that can be employed, are those of chlorine. This disinfecting gas is extricated from a mixture of common salt, black oxide of manganese, and sulphuric acid (see p. 158). Instead of chlorine disengaged in this mode, it is more common to use the chloride of lime, the chloride of soda, or the chloride of zinc; solutions of either of these, placed in shallow vessels in a room, will give off chlorine in a gradual manner, by a decomposition of their constituents. Some difference of opinion exists as to the mode in which this gas acts. Dr. A. T. Thomson is of opinion that it operates by decomposing the infected air:—

“With regard to the mode in which chlorine operates, the most probable opinion is, that it decomposes the infectious matter, which is a compound. One principle of this compound seems to be hydrogen; another, ammonia; and a third, a fetid volatile oil; besides the unknown matter of infection. The chlorine unites with the hydrogen, and forms hydrochloric acid, which combines with the second component of the infectious medium—namely, ammonia—and forms sal ammoniac; the withdrawing of both of which from the air causes the precipitation of the volatile oil; whilst the chlorine also neutralizes the real matter of infection, whatever it may be; and thus the vitiated atmosphere is purified.”

Although chlorine is capable of destroying fetid and unpleasant odors in the sick room, it is liable to many objections. However extricated, it is apt to create irritation of the respiratory organs and coughing when respired; it corrodes all metallic substances, and must, therefore, be used with caution; from the chlorides evolving the gas more slowly, they are, in most cases, preferable to the immediate extrication of chlorine from the salt and manganese.

Other acid fumigations have also been found useful as disinfectants; as those of nitrous acid, so highly recommended by Smith and Carmichael, and considered by Christison even as superior to chlorine. Sulphurous acid fumes, produced by burning sulphur, have likewise been recommended. Both are capable of destroying unpleasant odors and emanations; but the same objections apply against their use in a sick



chamber—that of causing much irritation of the organs of respiration; hence they are unfitted to cases where there is any disease or weakness of these parts.

Fumigations, used as medicinal agents, are much more prescribed on the Continent of Europe than in Great Britain or the United States. They are sometimes employed as inhalations—as emollient vapors in the treatment of bronchial affections; balsamic vapors in affections of the chest. They are also used externally, as the fumes of benzoin in rheumatic and gouty complaints; those of aloes in deficient menstruation; those of sulphur and mercury in various diseases of the skin.

The effects of fumigations vary according to the temperature at which the vapors are used; when it is high, and the whole body of the patient is exposed to it, it produces much the same train of symptoms as those caused by a vapor-bath. When the fumigation is made with substances which give off no watery vapors, the phenomena that result are analogous to those of a heated air-bath, but more marked, especially those which occur in the skin and subcutaneous tissue.

### BLOOD-LETTING.

Various methods are practiced for this purpose. General bleeding is performed by opening a vein, called venesection; or by puncturing an artery, which has received the name of arteriotomy. Topical blood-letting is practiced by opening the small vessels of a part by means of slight incisions, or what is called scarification, by the application of leeches, and by cupping.

GENERAL BLOOD-LETTING.—*Venesection* is one of the most frequently employed of the different modes of detracting blood. It is usually practiced on one of the veins of the arm. A band or ligature is to be applied around the arm above the point at which the vein is to be opened, so as to obstruct the passage of the blood to the heart; this causes an enlargement and turgidity of the vessel below the ligature. This bandage should never be applied so tightly as to prevent the passage of blood in the arteries of the part. The vein is to be opened by means of a common or thumb lancet, or a spring lancet. In using the first, the vein is to be kept from rolling by means of the thumb of the left hand placed a small distance below where the puncture is to be made; but care should be taken not to alter the relative position of the skin and vein, because in resuming their position after the lancet is inserted, the aperture through the skin will no longer correspond with that in the vein, in consequence of which a thrombus or ecchymosis will form—a very common accident in venesection. The lancet is to be pushed into the vein, and when its point is within the cavity of the vessel, it is to be carried forward a little, so as to enlarge sufficiently the opening. The arm is to be kept extended after the operation until the necessary quantity of blood is taken. If the blood should not flow as freely as is desired, the patient should exert the muscles of the arm by grasping a stick, or moving the fingers.

The spring-lancet is much used in some parts of the United States, and is almost universally employed in Philadelphia. This has arisen, in great measure, from its being preferred to the common lancet by Dr. Physick as well as by Dr. Dorsey, whose reasons for this preference are thus given in his “*Elements of Surgery*.”

“In a country, situated like the United States, where every surgeon, except those residing in the large cities, is compelled to be his own cutler, at least so far as to keep his instruments in order, the spring-lancet has a decided preference over the lancet; the blade of this can with great ease be sharpened by any man of common dexterity, and if not very keen, it does no mischief, whereas a dull lancet is a most dangerous instrument; and no one can calculate with certainty the depth to which it will enter: to sharpen a lancet, is regarded by the cutler as one of his nicest and most difficult jobs; it is one to which few surgeons are competent.

“The *safety* of using the fleam is demonstrated by daily experience; there is no country in which venesection is more frequently performed than in the United States, and, *perhaps, none where fewer accidents from the operation* have occurred; of these few, I beg leave to state that all the aneurisms produced by bleeding, which I have



seen, have been in cases where the lancet was used. I have since, however, met with an exception to this statement. I have seen the brachial artery opened by a spring-lancet, but it was by an old barber, half blind, and very clumsy."

"The manner of using the spring-lancet differs in nothing from the operation with the common lancet, excepting that the surgeon must place the instrument in such a situation over the vein that, when the spring is touched, the orifice into the vein will have a proper size and direction. Dexterity in this is very readily and speedily acquired. In point of *facility* in its use, it has a great advantage over the lancet.

"Among the advantages of the spring-lancet, *economy* is not the least. A country practitioner who is constantly employing the English lancets, and who is particular in using none but the best, must necessarily consume half the emolument derived from the operation in the purchase of his instruments. One spring-lancet, with an occasional new blade, will serve him all his life."

After a sufficient quantity of blood has been drawn, the flow is to be stopped by removing the ligature and placing a finger on the orifice, when any blood that has soiled the arm is to be wiped off, and the edges of the orifice carefully brought together, that they may unite by the first intention. To accomplish this, let the bleeder with the thumb of that hand which holds the arm push the skin towards the orifice, while he draws it on the other side to the same point with the compress; thus the skin will be thrown into folds at the wound, over which he is immediately to apply the compress, which should be broad to keep the skin better together, and thick to make the compression more certain.

When blood-letting is ordered, everything required for the operation should be at hand. The ligature for tying the arm, so as to obstruct the flow of blood in the veins below it, should be a soft old ribbon; and the same may be employed for securing the pledget over the orifice, when the flow of the blood is to be stopped. The pledget, or compress, should be a piece of linen, folded into a square form, about an inch in diameter, and comprehending three or four thicknesses of the material. It is always proper to use a graduated basin for receiving the blood, that the exact quantity drawn may be ascertained. If the invalid be bled in bed, he should sit erect; if he be up, he should be placed erect in a chair. When the operator is dexterous, there is little or no necessity for guarding either the bed or the clothes of the patient from the blood; as the pressure of the thumb of the operator, placed below the point where the vein is to be punctured, should not be relaxed until the utensil for receiving the blood be conveniently placed for that purpose. A basin with a little tepid water, and a clean, soft towel, should be ready for washing and drying the arm, before the compresses and the bandage for preventing the further escape of the blood be applied.

When a nurse or an attendant is not fully instructed in her duty, there is always considerable anxiety displayed to provide smelling salts and other means to obviate fainting. But, in many cases, much of the benefit of blood-letting depends on the fainting which it causes; and the propriety of checking that effect should be left entirely to the physician. No person should be present at this operation, nor should any one attempt to hold the basin for receiving the blood who is liable to become sick or faint at the sight of blood.

The arm should not be used for some hours after a vein has been opened in it; but, if everything has gone on well, the bandage may be removed at the end of two days. When this has been applied too tightly, the forearm and hand are apt to swell and become painful, in which case the bandage is to be loosened. If, on removing the bandage, it is found that the orifice, instead of being closed, presents an inflamed appearance, with its lips swelled and red, the arm is to be kept perfectly quiet, the edges of the wound brought in contact, and some cool application made to the part, to relieve the inflammation. In those cases where the inflammation extends beyond the orifice, and assumes an erysipelatous appearance, with a hard and painful tumor around the wound, a compress is to be applied upon the vein at the inflamed part, to make the two sides adhere together; if they do not adhere, simple contact will still be sufficient to prevent suppuration in this part; or, if inflammation has gone so far as to make the surgeon suspect that suppuration has taken place,



then the compress must be put upon that part of the vein just above the suppuration.

The remedy for inflammation of the veins consequent on venesection, which answers best, is the application of a blister over the inflamed part. As soon as the inflammation commences, a small plaster of simple cerate, spread on linen, should be applied on the orifice, and over this, a blister large enough to cover the whole inflamed part, extending three or four inches from the orifice in every direction. This remedy was first proposed, and made use of, by Dr. Physick, and has been employed by many physicians with great success, even after extensive inflammation of the vein has taken place.

Sometimes it becomes expedient to open the external jugular vein. To do this, the head should be laid on one side, and the vein compressed by the operator's thumb near the clavicle, and the opening made in that part of the vein which lies over the sternocleido-mastoid muscle. The blood soon ceases to escape after the pressure is removed, but a small strip of adhesive plaster is sometimes necessary.

In children, especially where they are very fat, a vein in the leg or foot can often be found, especially about or above the malleolus, when none are perceptible in the arm. To promote their distension, and to increase the bleeding when they are opened, it is found useful to immerse the limb in warm water.

In bleeding young children from the arm, when they are restless and fretful, it is a good plan to attach the child's arm to the left forearm of the operator, by means of ligatures at the wrist and centre of the arm; by this means, complete control is obtained over the motion of the child's arm, and the venesection can be accomplished with comparative ease.

**ARTERIOTOMY** is the opening of an artery to detract blood. It is much more limited than venesection in its application, being practiced only upon the temporal and posterior auricular arteries; and, in most cases, is restricted to the former only. In this, the anterior branch is preferred, where it passes above the exterior angle of the eyebrow. In performing the operation, the vessel is to be partially divided transversely, and not longitudinally. When as much blood has been drawn as is deemed necessary, the incision is to be deepened so as to divide the artery entirely. On the consequent retraction of the cut extremities, the bleeding will generally cease; if it should not, a graduated compress, maintained in place by a bandage around the head, is to be applied.

**TOPICAL BLOOD-LETTING.**—This consists of the abstraction of blood by cupping, leeching, or scarification.

*Cupping* is performed in the following manner: The skin being softened by means of a sponge and warm water, a small bell-like glass, known as a cupping-glass, having the air contained in it rarefied by being passed over the flame of a lamp, or by other means, is immediately applied to the part, all hair and other extraneous substances being previously removed; from the formation of a vacuum beneath the cup, the pressure of the air on the surrounding surface causes that portion included in the cup to swell, and the vessels to become turgid. When this has taken place, the cup is removed and several incisions made by means of a scarificator, an instrument containing numerous lancets, which, by means of a spring, can make a corresponding number of incisions at the same moment; the depth of these incisions can be regulated by means of a screw which protrudes or withdraws the lancet, according to the vascularity of the part, the quantity of blood to be drawn, &c. When a sufficient quantity of blood has collected in the cup, it is to be removed by introducing the nail of one of the fingers under the edge, by which means, air being allowed to enter, the cup becomes detached. The part being washed with warm water to remove any clots of blood, the cup is again to be applied as above, and the operation continued till a sufficient quantity of blood is obtained. Sometimes, especially when applied to the scalp, the cups fill so rapidly with blood, as to become detached almost immediately on being applied.

For the operation of cupping, a basin of hot water, sponges, and clean soft towels



are to be provided. It is generally considered as a severe and painful operation; but this is not the case, if the operator understand his business. This is readily ascertained by observing the manner in which the cups are applied, and the rapidity with which they are filled. A good cupper does not exhaust much of the air in the cup before applying it, but simply passes its mouth rapidly over the flame of a lamp; for, when it is held over the flame even for a few seconds, the compression of the edge of the cup upon the skin is so great, that it operates as a ligature, and checks the flow of the blood to the scarified part; thence very little blood is procured. A good cupper, also, removes the cups without spilling the blood which they contain; and the whole operation is completed in a short time.

In our large towns, it is but seldom that physicians, nurses, or those attending in a sick room, are called upon to perform the operation of cupping, as professional cuppers are readily to be obtained; but in small towns or villages, this is not the case, and the duty devolves either upon the physician or on those acting as nurses; and hence some directions are necessary to those unaccustomed to this task. Few are found sufficiently expert to exhaust the air in the cup by means of the lamp; and even among the professional cuppers in this city this mode is very rare, though, when properly performed, it is by far the best. The usual plan is to rarify the air in the cup by means of a small cone of paper, dipped in spirits of wine, or strong brandy; this is ignited and thrown in the cup, which is instantly to be applied to the intended spot. Where the proper cupping-glasses and scarificator are not to be had, wine glasses or very small tumblers may be substituted for the first, and small incisions by means of a thumb lancet will answer the purpose of the latter. A very convenient apparatus may be obtained at most of the surgical-instrument makers, consisting of cups and an exhausting syringe fitted to them, by which even the most inexperienced nurse can perform the operation of cupping.

The cicatrices of the scarifications leave a permanent mark; on which account, females should not be cupped upon the nape of the neck; but when blood is to be drawn from the head, the glasses should be applied behind the ears, and a portion of hair removed in such a manner that it may be covered by what remains.

When cupping-glasses are applied without any scarification being practiced, the operation is called *dry cupping*, and is much used to cause a speedy irritation of the skin for the relief of oppression of breathing, local pains, &c. To obtain the full benefit from this operation, the cups should be suffered to remain upon the part until they cause an exudation of a small quantity of serum, or great irritation of the part. The application of dry cups has been found extremely beneficial in poisoned wounds; they act not only by abstracting the poison, but also, by the pressure they exercise on and around the part, in preventing the absorption of it. The experiments of Dr. Barry in England, and of Drs. Pennock and Rodrigue in this country, show unequivocally the value of this plan of treatment.

*Leeching* consists in the application of leeches to any vascular part of the body. These are annulated animals, having an extensile, soft body, terminated at each extremity with a muscular disk, the anterior of which is furnished below with a mouth having three crescentic jaws, armed with numerous sharp teeth. To draw blood, the leech first renders the part tense by means of the anterior disk, which acts like a cupping-glass, and then makes a triangular wound by a saw-like motion of its tri-radiate jaws; it then draws the blood by suction, until it is completely gorged, when it detaches itself and falls off.

Leeches are seldom properly applied or managed. The part to which they are intended to be applied should be washed with a little soap and warm water, then with simple cold water, and, lastly, it should be well dried. If the part be hot and inflamed, the leeches should be put, for a few minutes, into tepid water; and this should be done, also, when they are to be applied in the mouth, or to any part of the body warmer than the general surface; but, at all times, before they are applied, they should be dried between the folds of a clean soft towel. The easiest and best mode of applying them is, first, to place the number to be used in a hollow made with the points of the fingers in a towel folded like a napkin; then, so to turn the towel and the leeches upon the part where it is intended they should fix, that the towel will cover them. The hand must



be kept over the towel, to prevent their escape, until they all bite, which usually happens in a few minutes: after which, the towel may be removed. By this method, twenty or thirty leeches can be applied more rapidly and with less trouble than two, when each leech is separately applied. Another plan is to put the leeches into a pill-box or small glass, and apply this over the part to which it is designed that they should attach themselves. If this plan, however, cannot be pursued, owing to the nature of the part to which they are to be applied—as, for example, the inner or outer angle of the eye—then the simplest method is to scratch the skin with the point of a needle, and to apply the leech to the spot moistened with blood. When they are to be applied within the mouth, or any open cavity, each leech should be put into a large quill, with its head towards the open end of the quill, which should be applied to the part, and retained upon it until the leech is fixed, when the quill may be gently withdrawn; but a thread should be tied round the tail of the leech when it is to be applied within the mouth, to prevent it from being swallowed—an accident which has occasionally happened, and has been productive of serious consequences.

Leeches should never be forcibly detached, as their teeth are apt to separate, and, being left in the wound, to cause an erysipelatous inflammation on the part. They should be permitted to drop spontaneously, which being the result of a temporary suffocation (*asphyxia*), all muscular energy ceases in the animal, and, the teeth shrinking, it drops off entire. A bread-and-water poultice, not too hot, should then be laid over the bites, to encourage the bleeding. The invalid should be kept warm in bed, when it is necessary to abstract a large quantity of blood. In general, the bites soon cease to bleed; but, in some instances, a copious flow takes place; and therefore, to prevent exhaustion, the poultice should be frequently examined. This exhaustion is more likely to occur in children than in adults; and, for the same reason, leeches should not be applied upon children late in the evening, unless they are very urgently required. It is also proper, in young patients, to select for their application a part which admits of pressure; for example, the thorax and temples, where the bones are covered with a thin layer of soft parts. Sometimes the hemorrhage continues so as to become alarming; in such cases, where pressure is unavailing or cannot be practiced, creasote or some of the styptics applied to the bites will be found useful. Powdering the spot with rye flour has also proved effectual. When these means fail, cauterizing the bites by means of a fine point of nitrate of silver, or with a red-hot probe or knitting needle generally proves successful.

*Scarifications* are slight incisions made in a vascular part, in order to relieve the engorged capillaries; they are practiced in inflammations of the conjunctiva, and in those of the tonsils, with much benefit in many cases. The operation is very simple, consisting merely in dividing the surface of the inflamed parts to a slight depth with the point of a lancet or scalpel, and encouraging the bleeding by means of warm water.



## POISONS.

---

POISONS are usually defined to be substances of an animal, vegetable, or mineral nature, which, when administered in a small quantity, are capable of producing deleterious effects on the animal economy, and, in some instances, of causing the same consequences when applied to the surface of the body. But this definition is not strictly accurate, and is not applicable to all cases of poisoning; nor is it, perhaps, possible to give in a few words such a one as would include all poisonous agents. Poisons are usually arranged according to their action on the system; but, as the alphabetical plan has been pursued in the main body of this work, it will be continued in the present instance. For full information on the subject, the reader is referred to the work of A. S. Taylor "On Poisons in relation to Medical Jurisprudence and Medicine."

### ACID, ACETIC.

*Symptoms.*—Great heat and burning pain in the stomach, convulsions, death.

*Morbid Appearances.*—Mouth and fauces brownish; lingual papillæ enlarged; œsophagus lined with a brownish adventitious membrane; stomach livid and even blackened; vessels much injected.

*Antidotes.*—Magnesia or its carbonates mixed into a cream with water.

### ACID ARSENIOS, or ARSENIC. A metallic, corrosive poison.

*Symptoms.*—A metallic, austere taste; a great flow of saliva; nausea and vomiting; fainting, great thirst; a sensation of heat in the stomach, which, in many cases, rejects the mildest fluids; much griping, tenesmus, and purging; the stools being dark-colored and very offensive; the urine scanty and high-colored; the pulse small, frequent, and often intermittent; distressing palpitation of the heart, with labored respiration and cold sweats; prostration of strength; sometimes paralysis of the extremities, delirium, convulsions, &c.

*Morbid Appearances* generally confined to the stomach and intestines; in the former, in the form of intense inflammation, but not of erosion or abrasion; the inflammation also evident in the upper intestines, but slight in the colon, though often violent in the rectum. The morbid appearances of other organs various. Sometimes no morbid phenomena to be found.

*Tests.*—In the solid state: entirely sublimed by heat. If previously mixed with charcoal (and placed in a suitable test-tube), deoxidated arsenic will be obtained in form of a metallic coating inside the tube; and this may be re-converted into arsenious acid by urging it in various directions along the tube with the aid of a minute (spirit-lamp) flame; the facets of the crystals thus formed (on the cooler situations) will be seen in some places with the unassisted eye, but more distinctly by means of a 4-power lens.

In solution: ammoniacal nitrate of silver produces a lemon-colored (*arsenite of silver*) precipitate. Ammoniacal nitrate of copper throws down a fresh, grass-green (*arsenite of copper*) precipitate. Transmission of sulphuretted hydrogen produces a bright yellow (*sesquisulphuret of arsenic*) precipitate. Lime-water precipitates a white (*arsenite of lime*) powder; but this test is not one to be relied upon.

Placed with zinc, and diluted sulphuric acid, in Marsh's apparatus, the arseniuretted hydrogen thence arising, when lighted, will deposit metallic arsenic on a piece of glass held within the flame.



The precipitates referred to, if washed and dried, may be treated as directed above for arsenic in the dry or solid condition.

[NOTE.—In testing suspected matters obtained from the stomach, these (and, in cases of death, the stomach itself) must be cut or broken down, and boiled during at least three-quarters of an hour: if not sufficiently fluid, add distilled water. Strain, and with addition of a small quantity of potash, again boil during a quarter of an hour, and filter. If this liquor manifests either alkaline or acid reaction, neutralize with potash, or with acetic acid, as may be required; then acidulate it faintly with hydrochloric acid. Solution of nitrate of silver will remove the acid; and solution of muriate of soda will precipitate any excess of silver. The liquor may now be experimented on with the above tests.]

*Treatment.*—Abundant draughts of sweet milk, gruel, decoctions of starch, or oily mixtures; tickling the fauces, &c., to induce vomiting; the stomach-pump; emetics of sulphate of zinc. Hydrated sesquioxide of iron (newly prepared, *v. process*, page 209) in large doses is the only antidote of reliance.

Afterwards combat any inflammatory symptoms by the usual means, and let the patient subsist, for a long time, wholly on the blandest diet.

All arsenical poisons have much the same action, and similar means are to be used for their detection and for counteracting their effects.

**ACID, BORACIC.**—This is said to be very virulent; but no instances have been recorded of its fatal effects on man.

*Tests.*—Not very soluble in water; soluble in alcohol, which, when inflamed, burns with a rich green color; soluble in the caustic alkalies.

**ACID, CARBONIC.**—This gas is freely liberated in respiration, combustion, and fermentation, also in the calcination of lime, and is found in coal mines, wells, and cellars.

*Symptoms.*—Great drowsiness, giddiness, difficulty of respiration, loss of muscular power and sensibility, and coma. The whole body, but especially the face, appears swollen and livid; eyes are usually prominent, and retain their brilliancy for some time.

*Tests.*—It extinguishes a taper if the proportion be above twelve or fifteen per cent.; lime-water, or a solution of subacetate of lead, is precipitated white by it.

*Treatment.*—Remove the person into the open air, and elevate the head, dash cold water over him, apply stimulating frictions to the thorax and extremities, use artificial inflation of the lungs; as soon as the patient can swallow, stimulants may be cautiously administered; but, if there are signs of oppression of the brain, venesection is to be performed.

**ACID, CITRIC.**—In large doses, acts as an irritant poison; but no fatal case in the human subject has been recorded.

**ACID, HYDROCYANIC or PRUSSIC.**—An extremely active, sedative poison.

*Symptoms.*—When the dose is large, almost immediate death ensues; in smaller quantities, it causes pain in the head and stupor, nausea, faintness, vertigo, and loss of sight, followed by difficulty of respiration, dilated pupils, a small vibrating pulse, and syncope, which will end in death, if curative means are not employed.

*Morbid Appearances.*—None; but a strong odor of the acid is exhaled from the stomach.

*Tests.*—The (bitter almond) peculiarity of its odor. When a little potash is mixed with liquids containing this poison, and solution of sulphate, or sesquichloride of iron added, a grayish-green precipitate is thrown down—which deepens to a Prussian blue tint, on addition of a few drops of sulphuric acid. The nitrate of silver produces a white (*cyanide of silver*) precipitate; which, after being washed and dried, and then held on a watch-glass over a flame, burns with a fresh rose-color, cyanogen being at the same time evolved.

When a mixture is to be examined, containing matters from the stomach, &c., if alkaline, it must first be neutralized by addition of sulphuric acid, then one-eighth



part cautiously distilled therefrom into a receiver immersed in some frigorific mixture; and the product may then be tested by nitrate of silver, as above.

*Treatment.*—The internal remedy best calculated to act chemically on this poison, is carbonate of potash in solution, quickly succeeded by watery solutions of sulphate of iron (with the intention of decomposing the acid, and forming the less injurious substance named *Prussian blue*); but the instances of successful treatment by these, or, indeed, by any other kind of treatment, are extremely rare. Inhalation of chlorine gas, ammonia, &c.; artificial respiration; exhibition of energetic stimulants, such as brandy, liquid ammonia; chlorine-water, chloride of soda and of lime, have all been extolled, and may be tried. Mustard poultices to the stomach, and cow-hage applied over the chest; venesection at the neck, temples, &c.; and affusion of cold water to the head and spinal district, are more likely to prove timely excitant and effective remedies. Employment of the stomach-pump, emetics, &c., seems a most improbable means of relief—or, rather, a waste of time.

**ACID, MURIATIC or HYDROCHLORIC.**—A corrosive mineral poison.

*Symptoms.*—Sensation of burning in the throat and stomach, styptic taste in the mouth, much thirst; the eyes red and sparkling; the pulse frequent and tense; the skin hot and dry; the tongue red and glazed; the lips black; vomiting of blood and yellow matter, having the peculiar smell of the acid; cold sweats, delirium, &c. Orfila also says, a thick white fume, having the penetrating odor of the acid, issues from the mouth.

*Morbid Appearances.*—Mouth, fauces, throat, and stomach highly inflamed, of a deep-red color; mucous membrane sometimes detached or destroyed, sometimes perforated; contents of the stomach yellow or dark green.

*Tests.*—Characteristic acid taste. The presence of muriatic acid (in small proportion) is at all times demonstrable in the juices of the stomach; therefore proof of its presence in *excess* is required in cases of poisoning. When a piece of glass, wetted with this acid, is held close to *liquor ammoniæ*, so as to let the vapors of both meet, white (*muriate of ammonia*) fumes are formed. Solution of nitrate of silver produces a white (*chloride of silver*), curdy-looking precipitate.

*Treatment.*—The immediate administration of magnesia, soda, or potassa, mixed in bland demulcent drinks, to be followed by the free use of emollients and mucilages. If patient survives the first effects of the poison, employ antiphlogistic means to overcome the supervening inflammation.

**ACID, NITRIC, or AQUAFORTIS.**—A corrosive mineral poison.

*Symptoms.*—Much the same as those produced by the last-mentioned poison.

*Morbid Appearances.*—In cases where death has occurred soon after the ingestion of the poison, the most striking appearance is a layer of yellow matter on all parts reached by the acid. The lips, chin, and hands of the person are also often stained with orange-colored spots. Perforations of the stomach are found in most cases.

*Tests.*—The orange-colored fumes that are given off, when it is boiled with copper filings, are characteristic. It reddens morphia, and blackens a solution of protosulphate of iron. A fluid containing it, on the addition of carbonate of potassa forms nitre, which may be known by its deflagration, or by powdering a small portion, and placing it in a glass tube with some fine copper filings, moistening with water, and adding a few drops of sulphuric acid—when, if it be nitre, there will be an evolution of the orange-colored fumes of nitrous acid.

*Treatment.*—The administration of magnesia, or chalk, in some bland fluid, then the free use of demulcents, and subsequently the employment of antiphlogistic means to subdue inflammation.

**ACID, OXALIC.**—A corrosive vegetable poison.

*Symptoms.*—Burning heat of the stomach, nausea; sometimes vomiting, but at others, ineffectual efforts to discharge the contents of the stomach; great prostration of strength, violent pain, spasmodic respiration, convulsions, &c. When the patient



survives the first effects of the poison, all the symptoms of violent inflammation of the alimentary canal are developed. In a diluted form, it appears to cause paralysis of the heart or symptoms of cerebro-spinal disease.

*Morbid Appearances.*—Tongue and fauces usually covered with a viscid white coat; the stomach containing a dark-brown mucous fluid, and its substance of almost a gelatinous consistency. In some cases, no traces of the action of the poison are perceptible; but, when death does not occur for some time, the alimentary canal is found inflamed.

*Tests.*—In solid state: Has the characteristic sour taste of most acids; and produces the same changes as they do on vegetable blues. When placed in water, a crackling sound accompanies the solution of its crystals.

In solution: Solution of chloride of lime produces a white (*oxalate of lime*) precipitate (which is insoluble in hydrochloric acid, but soluble in nitric acid). Solution of nitrate of silver also throws down a white precipitate (which, on being carefully dried by aid of heat, on bibulous paper, acquires a brownish outline, and towards the end shows some slight explosions).

*Treatment.*—Mixtures of magnesia, whiting (or plaster scraped from off the inside wall of an apartment) mixed with water (the amount of the latter being as small as convenient for swallowing); abundant drinks *subsequently*. No alkalies should be employed.

ACID, SULPHURIC.—A corrosive mineral poison.

*Symptoms.*—An austere styptic taste in the mouth; a sensation of burning heat in the throat and stomach, followed by nausea, vomiting, and much fetor of the breath. The matters vomited contain both venous and arterial blood. Signs of great inflammation of the abdominal viscera soon manifest themselves, with difficult respiration, a croupy cough, and a small, frequent, irregular pulse; great anxiety and restlessness, convulsive motions of the muscles of the face and lips; sometimes a papulous eruption preceding death.

*Morbid Appearances.*—These are not always to be found, except in the fauces and larynx, which, in most cases, present evidences of the highly corrosive action of the poison. The stomach sometimes is found to contain a quantity of dark grumous matter, and is much distended with fetid gas—its membranes ulcerated, dark-colored, and having numerous corroded spots, and even perforations.

*Tests.*—The acid may be in a concentrated or a diluted state. If in the former, any organic matter placed in contact with it is blackened and charred; when mixed with an equal bulk of water, much heat is evolved; when boiled with copper filings or mercury, sulphurous acid gas is evolved. When the acid is in a diluted state, the best test is nitrate of baryta, which causes a dense white precipitate of sulphate of baryta; this can be verified by calcining it for some minutes with an equal weight of charcoal, wrapped in platina foil, then introducing the residue into a glass-tube and adding a few drops of muriatic acid. This will cause an extrication of sulphuretted hydrogen, which can be recognized by its odor, and by its blackening carbonate or acetate of lead.

*Treatment.*—The same as for the other mineral acids.

ACID, TARTARIC.—A corrosive vegetable poison.

*Symptoms.*—Very analogous to those caused by oxalic acid.

*Morbid Appearances* are likewise similar to those produced by oxalic acid.

*Tests.*—When heated on platina foil, it burns with a pale reddish flame, and exhales a peculiar acrid vapor, leaving much carbonized matter. When a solution is treated with lime-water, it affords a white precipitate, soluble in an excess of the acid; when treated with caustic potassa, it affords a granular precipitate of the bitartrate.

*Treatment.*—The same as for oxalic acid.

ACETATE OF COPPER. *See* Copper.

ACETATE OF LEAD. *See* Lead.



ACETATE OF MORPHIA. *See* Opium.

ACONITE. *See* Vegetable Poisons.

ALCOHOL.—A narcotico-acrid poison.

*Symptoms.*—These vary according to the dose. In small quantities, there is mere excitement. In large doses, much excitement, with delirium, confusion of intellect, followed by somnolency; nausea and vomiting, and even coma and apoplexy. When an undue portion has been swallowed, it often proves instantly fatal; it may also prove fatal by occasioning or aggravating other diseases. The effects of an habitual use of it are diseases of the viscera, and various mental and nervous disorders. The symptoms of poisoning with alcohol may be mistaken for those of epilepsy or apoplexy.

*Morbid Appearances* are inflammation, softening, &c., of the mucous membrane of the stomach, congestion of the cerebral vessels, sanguineous or serous extravasation in the brain or lungs.

*Tests.*—Odor of the contents of the stomach, or of matters ejected from it. The chemical proof consists in removing the contents of the stomach, mixing them with distilled water, filtering and distilling in union with carbonate of potassa or soda; the product is mixed with fused chloride of sodium, and again distilled. Alcohol will be found in the receiver.

*Treatment.*—Withdraw the contents of the stomach, as speedily as possible, by a stomach-pump; afterwards an emetic of salt and water should be given in large doses, and repeated, at short intervals, till the stomach is well cleared. The cold affusion is useful in some cases. Ammonia may be employed as a stimulant, and general symptoms obviated by blood-letting; but this must be employed with great caution.

#### ALKALIES.

AMMONIA.—A corrosive mineral poison.

*Symptoms.*—Excoriation of the mouth and fauces; burning sensation in the throat and stomach, usually followed by vomiting and purging, the ejected matters being often bloody. When the quantity taken is large, an immediate feeling of strangulation ensues, attended with convulsions. If the result is fatal, it quickly follows the ingestion of the poison. The inhalation of ammonia by the nostrils, when it is too freely used in cases of fainting, has caused the same symptoms as when taken into the stomach.

*Morbid Appearances.*—Marks of high inflammation of the parts with which the poison has come in contact.

*Tests.*—The pungent odor; its alkaline reaction—but which is dissipated by heat. By causing a yellow precipitate with a mixture of arsenious acid and nitrate of silver; by producing a rich violet-blue solution with sulphate of copper; and forming white fumes with chlorine.

*Treatment.*—The immediate administration of one of the vegetable acids, and afterwards the copious use of demulcents. When ammoniacal vapor has been inhaled, the patient should inspire the vapor of vinegar.

POTASSA.—A corrosive mineral poison.

*Symptoms.*—An acrid, caustic, urinous taste in the mouth; a sensation of burning heat in the throat; nausea, and sometimes vomiting of bloody matters. The surface cold and clammy; the pulse quick and feeble; hypercatharsis, and violent colicky pains.

*Morbid Appearances.*—Strong marks of inflammation in the alimentary canal, softening, erosion of the mucous coat, and, in some cases, perforation of the stomach.

*Tests.*—Alkaline reaction; precipitation of nitrate of silver in the form of a dark-colored oxide. Carbonic acid water causes no precipitate. A concentrated solution affords a canary-yellow precipitate with bichloride of platina.



*Treatment.*—Vinegar and the diluted vegetable acids, to be followed by a free use of demulcents, or oleaginous mixtures.

**SODA.**—A corrosive mineral poison.

*Symptoms and Morbid Appearances.*—Analogous to those following the use of potassa.

*Tests.*—Caustic soda in solution is not precipitated by bichloride of platina or tartaric acid; its alkaline nature can be ascertained by the usual tests.

*Treatment.*—The same as for potassa.

**ANTIMONY.**—Metallic antimony is not regarded as a poison; but its vapor is said to cause unpleasant symptoms when respired.

**ANTIMONY, TARTARIZED**, generally known as **TARTAR EMETIC**, is an irritant metallic poison.

*Symptoms.*—Nausea and severe vomiting, hiccough, cardialgia, burning heat at the epigastrium, severe colic and purging, small and rapid pulse, cold skin, syncope, difficult respiration, vertigo, insensibility to external stimulants, painful cramps in the lower extremities, great prostration of strength.

*Morbid Appearances.*—The stomach and intestines often much inflated with gas; their mucous membrane red, and covered with a slightly adhering viscid layer; the peritoneum of a dark brick-red hue; the membranes of the brain displaying marks of inflammation.

*Tests.*—In a solid state, add charcoal, introduce into a test-tube and expose to heat; metallic antimony will be found of a grayish-black lustre. In the state of solution, diluted nitric acid causes a white precipitate. Sulphuretted hydrogen throws down a reddish-orange precipitate. Solutions containing tannin cause a copious, curdled, whitish-yellow precipitate. Should matters from the stomach be present, the solution must be well agitated along with a small portion of muriatic and tartaric acid; then filtered previous to being experimented on.

*Treatment.*—Encourage vomiting by free administration of warm water and other diluents, or employ the stomach-pump, if necessary. Infusions containing tannin are very useful by decomposing the poison.

**BARYTA, AND PREPARATIONS OF.**—The only two preparations of baryta that have caused death are the **CARBONATE** and **CHLORIDE**.

*Symptoms.*—Those of irritation, combined with an affection of the brain and nervous system, as vertigo, convulsions, and paralysis.

*Morbid Appearances.*—Evidences of inflammation of the mucous coat of the stomach.

*Tests.*—In solution: sulphuric acid throws down a profuse dense white (*sulphate of baryta*) precipitate. Solution of nitrate of silver also produces a white (*sulphate of silver*) precipitate. Sulphuretted hydrogen produces no change of appearance; which circumstance, being peculiar to salts of barium, is valuable as a *negative* test.

*Treatment.*—Sulphates of magnesia or soda in solution in water, if the patient is seen early; they are, however, of little use where the carbonate has been taken, for in this case emetics and the stomach-pump should be used.

**BELLADONNA.** See Vegetable Poisons.

**BISMUTH.**—The only preparation that has caused death is the **SUBNITRATE**.

*Symptoms.*—Burning pain in the throat, nausea, vomiting and diarrhoea, coldness of the extremities, and spasms.

*Morbid Appearances.*—Inflammation of the fauces and œsophagus, redness of the stomach and intestines.

*Tests.*—In substance: Dissolves in nitric acid, without change of color, and without effervescing (which distinguishes it from a *carbonate*). Sulphuric acid produces no



precipitate (disproving the possible idea of its being a preparation of *lead*). Dissolve it in nitric acid, and add solution of potash so long as any precipitate is thrown down; gather this in a filter, and, after washing and drying it, place it on charcoal (and treat it as directed for preparations of lead), when its reduction to the metallic condition will be effected.

*Treatment*.—Induce vomiting by tickling the fauces, &c., and give continuous full draughts of sweet milk.

**BROMINE**.—An irritant, corrosive mineral poison.

*Symptoms*.—Irritation and inflammation of the parts with which it comes in contact, whether in substance or in vapor.

*Morbid Appearances*.—Fauces, œsophagus, and stomach inflamed and corroded, mucous membrane gelatinized, duodenum of a yellow color, and thickened; the parts retaining a strong smell of the poison.

*Tests*.—Its color, odor, and volatility. All solutions containing it are rendered colorless by the addition of potassa. On being evaporated, the residue is to be incinerated at a low temperature, the ash dissolved in distilled water, filtered, and chlorine passed through the solution. On the addition of ether, the bromine is taken up by that fluid.

*Treatment*.—The immediate and free use of albumen, or starch.

**BRUCIA**. See Vegetable Poisons.

**CALOMEL**. See Mercury.

**CAMPHOR**. See Vegetable Poisons.

**CANTHARIDES**.—An acrid and corroding animal poison.

*Symptoms*.—A burning sensation in the throat; violent pain in the stomach and bowels; nausea, vomiting, and purging—the ejections being frequently bloody and purulent; great heat and irritation of the urinary organs, sometimes the most painful priapism; pulse quick and hard; convulsions, tetanus, delirium, syncope.

*Morbid Appearances*.—Inflammation and erosion of the stomach; if in substance, fragments of the flies, adhering to the mucous coat, or mixed with contents; marks of inflammation in the intestines and urinary organs: these are most evident when death does not soon follow the ingestion of the poison.

*Tests*.—The characteristic green, shining appearance of the fragments of the flies, and the character of the symptoms.

*Treatment*.—The promotion of vomiting by means of warm demulcents; copious dilution, bleeding, the warm bath, opiate frictions, enemata of mutton broth, laudanum, &c. Camphor, though not an antidote, alleviates some of the most distressing symptoms.

**CARBONIC ACID GAS**.

*Symptoms*.—Great drowsiness; difficulty of respiration; loss of sensibility; the countenance of a livid or deep leaden color.

*Morbid Appearances*.—Whole body, but especially the face, swollen, which is also usually livid, and the features distorted; eyes generally prominent, and retaining their brilliancy. Vessels of the brain and lungs in a state of congestion.

*Tests*.—Expose a vessel, filled with lime-water, to the deleterious atmosphere, when, if carbonic acid be present, it will assume a milky appearance, and let fall a white precipitate. A lighted candle will detect the presence of carbonic acid gas (if in excess) by being immediately extinguished in the tainted atmosphere; it must, for this purpose, be held near the floor. [The utmost caution should be observed in introducing lighted tapers into close situations supposed to contain this gas, lest an explosive hydrocarbon may, instead, be met with.]

*Treatment*.—Remove the patient from the situation or apartment in which the poison has been inhaled; or open the doors, windows, &c., for admission of pure air.



Continued application of cold water to the head and neck. Bleeding may be necessary; or cupping at back of the neck. Cautious inhalation of steam from warm water placed in a convenient vessel.

**CHLORINE.**—An acrid gaseous poison.

*Symptoms.*—Great and painful constriction of the glottis, cough, sensation of suffocation, alternating with asphyxia; if the case is protracted, inflammation of the larynx, and pneumonic inflammation; odor of the gas perceptible in the breath and in vomited matters.

*Morbid Appearances.*—Marks of irritation and inflammation in all the parts with which the poison has come in contact.

*Tests.*—A solution of this gas has a yellowish color, and a strong peculiar smell, which latter is evolved on boiling. It gives a white precipitate with nitrate of silver, which is insoluble in muriatic acid.

*Treatment.*—The best remedy is a free use of albumen. Magnesia with mucilaginous fluids may also be freely exhibited.

**CHROME.**—The only form of this metal that requires notice is the

**BICHROMATE OF POTASSA.**—An acrid mineral poison.

*Symptoms.*—Burning heat in the mouth, throat, and stomach; excessive and painful vomiting of bloody mucus; convulsions, palsy, &c.

*Morbid Appearances.*—Softening and abrasion of the mucous coat of the stomach and intestines.

*Tests.*—In substance, by its orange-red color. In solution, by affording a rich red precipitate with nitrate of silver, a bright yellow with acetate of lead, a dingy green with sulphuretted hydrogen.

*Treatment.*—The use of the carbonates of potassa or soda to neutralize the excess of chromic acid, followed by the administration of emetics.

**COCCULUS INDICUS.** *See Vegetable Poisons.*

**COLCHICUM AUTUMNALE.** *See Vegetable Poisons.*

**CONIUM MACULATUM.** *See Vegetable Poisons.*

**COPPER.**—This metal is not poisonous in a metallic state, but becomes highly so when oxidated, or in combination.

The most common preparations of copper are the **ACETATE** (*verdigris*), **SULPHATE** (*blue vitriol*), the **CARBONATE**, and **AMMONIURET**.

*Symptoms.*—Violent headache, vomiting, pain in the bowels, cramps in the lower extremities, a peculiar and permanent coppery taste, diarrhoea, convulsions, palsy, and insensibility.

*Morbid Appearances.*—Marks of inflammation in the stomach and intestines. When the case has been protracted, there is often a green tinge of the lining membrane, and a jaundiced appearance of the skin.

*Tests.*—The Sulphate in a solid state: bright blue color, astringent metallic impression on the tongue.

In solution: transmission of sulphuretted hydrogen affords a brownish-black (*sulphuret of copper*) precipitate. Solution of ammonia precipitates a pure green substance, which, on further addition of ammonia, becomes dissolved; but when only a small amount of the poison is present, no precipitation occurs, a clear violet-colored solution being then at once presented. Ferrocyanide of potassium throws down a reddish-brown (*ferrocyanide of copper*) precipitate. A piece of polished zinc or iron (as the blade of a table-knife) acquires a plating of copper, if placed in a liquid containing the above salt. The tests for sulphuric acid will indicate its presence here as the combining acid.

The Acetate: Same tests as preceding, excepting the last.



The Ammonio-sulphate: has a brilliant violet-blue color, and an ammoniacal odor. Solution of arsenious acid affords a fresh apple-green precipitate; this experiment is, conversely, that for arsenious acid.

*Treatment.*—Vomiting to be promoted by copious draughts of warm water, milk, or mucilaginous fluids; various antidotes have been highly praised. Sugar was formerly recommended, but has not been as successful as was supposed. Albumen and milk form an insoluble compound with copper, provided they are in large excess. The protosulphuret of iron and iron filings have also been employed with advantage; but their action is too slow. The hydrated oxide of iron has been successfully administered where the arsenite of copper had been taken.

CORROSIVE SUBLIMATE.—See Mercury.

DIGITALIS.—See Vegetable Poisons.

#### IODINE.

*Symptoms.*—In large doses, this substance causes heat and constriction of the fauces, nausea, offensive eructations, pain in the stomach, retching, colic, diarrhoea, quickness of the pulse, tremblings, great thirst, salivation, convulsions, &c.

*Morbid Appearances.*—Distension and inflammation of the stomach and bowels; sphacelation in some places: in protracted cases, increased size and paleness of the liver.

*Tests.*—In the solid form, it is in bluish-gray scales, having somewhat the odor of chlorine. When heated in a tube, it gives off violet fumes; when added to the mucilage of starch, it strikes a deep blue color. When it is mixed with other matters, pass a stream of sulphuretted hydrogen through the mixture, boil it, saturate with potassa, add mucilage of starch, filter, add nitric acid, and the color will indicate iodine.

*Treatment.*—Promote vomiting by the free administration of amylaceous and mucilaginous fluids, and treat the inflammatory symptoms as they occur.

#### IODIDE OF POTASSIUM.

*Symptoms.*—Uneasiness of the stomach, followed by nausea, and a burning sensation in the stomach, vomiting, cephalalgia, vertigo, and tremors.

*Morbid Appearances.*—Stomach contracted; mucous membrane with ecchymosed spots, and some abrasions; marks of inflammation in the intestines.

*Tests.*—Solid state. Heat with the protoxide of manganese; moisten the powder with a mixture of equal parts of sulphuric acid and water; heat in a tube, and the purple vapor of iodine makes its appearance.

In solution: add to the filtered fluid an equal quantity of mucilage of starch, and a few drops of nitric acid, and the characteristic blue color will be produced. Test the urine in the same manner. It may also be detected in the solids of the body, by drying them, incinerating and lixiviating, filtering, and using starch and nitric acid.

*Treatment.*—The same as for iodine.

IRON.—Not poisonous in a metallic state; but some of its preparations act as irritants, especially the sulphate and the muriate.

*Symptoms.*—Nausea, vomiting, and purging; sensation of heat in the stomach and bowels; dejections of a black color, &c.

*Morbid Appearances.*—Softening of the mucous coat of the stomach and intestines; the whole extent of the alimentary canal of a black color; marks of inflammation.

*Tests.*—For the sulphate: ferrocyanide of potassium will give a greenish-blue precipitate, becoming a deep blue by exposure to the air; hydrosulphuret of ammonia will give a black precipitate. The presence of sulphuric acid can be ascertained by means of nitrate of baryta.

*Treatment.*—In poisoning by either salt, magnesia or the alkaline carbonates should be freely given.



**LEAD.**—In a metallic state, does not appear to be poisonous; but most of its compounds are so, especially the acetate, chloride, carbonate, and oxide.

*Symptoms.*—Obstinate constipation, violent colic with retraction of the abdomen; vomiting, pulse small and hard, laborious breathing, and tremors, gums with a blue tinge, paralysis of the extremities, &c.

*Morbid Appearances.*—The appearances on dissection are very various, but are principally those of inflammation.

*Tests.*—In a solid state: Mixed with potash and charcoal, and exposed to a red heat (or placed in a suitably sized hollow in a piece of charcoal, and exposed to a blow-pipe flame), metallic lead is formed.

In solution: Sulphuric acid, and sulphates, throw down a white (*sulphate of lead*) precipitate. Solution of carb. of soda gives a white (*carbonate of lead*) deposit. Solution of chromate of potassa a rich yellow (*chromate of lead*) precipitate. Transmission of hydrosulphuric acid produces a blackish (*sulphuret of lead*) precipitate. Zinc precipitates lead from solution of any of its salts.

To render the *carbonate* amenable to the foregoing tests, it must first be dissolved in a sufficiency of nitric acid diluted with distilled water.

*Treatment.*—Administer solution of sulphate of zinc in full doses; induce and continue vomiting. If an insoluble salt of lead is the poison, almond or olive oil must at first be given in considerable quantity. The free exhibition of the sulphate of soda or magnesia, or of milk or albumen, to be followed by an emetic. In colic, the treatment is to consist of active purgatives, followed by full doses of opium or Dover's powder. Inflammatory symptoms are to be counteracted by antiphlogistic means.

**LOBELIA.**—See Vegetable Poisons.

**MERCURY.**—In a metallic state exercises but a slight influence on the body, but in a state of vapor is capable of causing violent symptoms. All the salts of mercury are poisonous; but the most important is corrosive sublimate.

*Symptoms.*—The symptoms caused by corrosive sublimate resemble those produced by arsenic, but, from the salt being more soluble, they are more immediate and violent; there is a more marked taste, and the evacuations are more frequently bloody. There are three varieties of poisoning with mercury. In the first, the leading symptoms are, violent irritation of the alimentary tube, namely, vomiting, purging, pain at the pit of the stomach, and irritation in the throat; metallic, styptic taste, corrosion of the mouth, tongue, and palate; constriction of the throat, and difficulty of swallowing. Blood evacuated both by vomiting and by stools, suppression of urine, countenance flushed, tumid, and bloated. In the second variety, salivation and sloughing of the mouth succeed to the irritation and inflammation. In the third, mercurial erethism comes on, and is not preceded by the symptoms of local irritation. The first variety arises from the more soluble salts of mercury, in large doses; the second, from the same preparations, but in smaller doses, and more diluted; the third, by the more insoluble and refractory compounds.

*Morbid Appearances.*—These are similar to those attendant on irritative or corrosive poisoning. There are frequently shrivelling of the tongue, and enlargement of the papillæ and root. In some cases, red and black spots in the cavities of the heart. Corrosion, ulceration, and disorganization of the mucous coat of the stomach and intestines. An inflamed condition of the urinary organs is also frequently observed.

*Tests.*—Corrosive sublimate in a solid state: Is sublimed when heated in a test-tube; and the acrimonious fumes speedily condense into a crystalline, semi-transparent mass. Placed in a test-tube, and potassa or soda added in solution, a yellow (*peroxide of mercury*) precipitate is thrown down (lime has the same effect). Triturated with an equal weight of well-dried carbonate of potassa, metallic globules of mercury will become visible.

In solution: Ammonia sends down a white (*ammonio-chloride of mercury*) precipitate. Solution of *proto*-chloride of tin affords a (*calomel*) precipitate, which at first is white, but acquires a leaden color on adding more of said test; when this precipitate (after being well agitated) is dried, minute globules of quicksilver are



formed. Transmission of sulphuretted hydrogen produces a (*bisulphuret of mercury*) precipitate, which at first is leaden-colored, then black. Solution of iodide of potassium affords an intensely brilliant scarlet (*proto-ioduret of mercury*) precipitate. Corrosive sublimate may be reduced to the metallic state through galvanic influence applied as follows: A drop of the suspected fluid being laid on a piece of polished gold, and both it and the gold touched at the same instant by a point of iron (as a thick needle, or the end of a penknife), a small silvery coating of mercury soon becomes apparent on the gold. Being most soluble in sulphuric ether, addition of this fluid is of much service when the poison is found mingled with organic or other matters.

**The Bicyanide:** When heated in a tube, cyanogen gas is evolved, which, when inflamed at the mouth, burns with a rose-red flame, with a blue halo.

**Calomel:** Is sublimated by heat. When heated along with potash, it at first assumes a black appearance, and then quicksilver globules are observed. Hydro-sulphuret of ammonia is equally valuable in treatment of calomel, as in the foregoing.

**The Iodide:** On cautious application of heat, it sublimes in red-colored crystals, which soon change to yellow, and subsequently to a dusky hue. On exposure to a sufficient degree of heat, iodine fumes are disengaged. When mixed with potash (in equal weights) and heated in a test-tube, decomposition occurs; metallic mercury being sublimed, and iodide of potassium deposited in the tube.

**Red Precipitate:** Exposed to heat in a test-tube, metallic mercury is sublimed; evolution of oxygen being at the same time demonstrable.

**The Persulphate:** Treated in same manner as foregoing, sublimation of the mercury takes place; and sulphurous acid gas is perceptibly evolved.

**Treatment.** If vomiting does not already exist, it must be produced by the exhibition of emetics. Various antidotes have been suggested for this poison. The best appears to be albumen of eggs in continuous large doses, and infusion of catechu afterwards; sweet milk; mixtures of flour with water (for sake of gluten) in successive cupfuls; iron filings (emetics of tartrate of antimony have been recommended, to combat ptyalism, when present); but to check excessive salivation, small doses of the chlorate of potassa appear to be the most successful.

**MORPHIA.** See Vegetable Poisons. Opium.

**NUX VOMICA.** See Vegetable Poisons.

**OPIUM.** See Vegetable Poisons.

**PHOSPHORUS.**—A corrosive poison.

**Symptoms** are protracted in their appearance. A hot alliaceous taste in the mouth; acrid burning sensation in the throat and stomach; nausea and vomiting. The vomited matters are of a dark color, and emit white fumes; pulse small and frequent; frequently violent convulsions.

**Morbid Appearances.**—A general inflammatory appearance of the stomach and intestines, with sphacelated spots in various parts.

**Tests.**—The peculiarity of its odor; its highly inflammable property, when only moderately heated.

**Treatment.**—Mixture of magnesia and cold water, in repeated draughts. Promote continuous vomiting by tickling application of a feather to the fauces. The exhibition of oil is injurious, as it tends to dissolve the phosphorus.

**STRYCHNIA.**—See Vegetable Poisons. Nux vomica.

**TIN.**—This is not poisonous in a metallic state; but some of its salts, as the chlorides, are violent irritants.

**Symptoms.**—An austere metallic taste; constriction of the throat, impeded respiration, violent vomiting, cramp of the stomach and violent colic, purging, pulse small and quick, convulsions, sometimes paralysis, asphyxia.



*Morbid Appearances.*—Inflammation and erosion of the stomach and intestines.

*Tests.*—Chloride of gold gives a deep purple-brown, almost black precipitate. Nitrate of silver affords a white precipitate, insoluble in nitric acid. Sulphuretted hydrogen throws down a deep chocolate-brown precipitate, even in diluted solutions.

*Treatment.*—Copious draughts of milk; then excite vomiting. Inflammatory symptoms to be combated by antiphlogistics, and nervous irritation by opiates and antispasmodics.

**VEGETABLE POISONS.**—These are very numerous, and possessed of the most different qualities; some being mere irritants, others narcotic irritants—whilst another class are pure narcotics.

**IRRITANTS.**—This includes a great number of vegetable substances, the principal of which are aloes, arum, capsicum, colocynth, creasote, croton seeds, elaterium, euphorbium, gamboge, jalap, mezereon, ricinus or castor seeds, savin, scammony, &c.

*Symptoms.*—Many of these articles act especially on the bowels, and in moderate doses are efficaciously used as purgatives. In large doses, they cause hypercatharsis, and much irritation of the stomach and bowels.

*Morbid Appearances.*—These are various degrees of inflammation of the stomach and bowels.

*Tests.*—None by means of reagents. The rich yellow color of *gamboge* may sometimes prove a means of its detection; and familiar acquaintance with others of the above substances may lead to their recognition by smell, taste, &c.

*Treatment.*—The expulsion of the poison by means of emetics, and, when this is effected, copious demulcent drinks in a warm state; succeeded by liquid cordial, stimulant, or opiate medicines suited to particular circumstances.

**NARCOTICS.**—This is a small class of the vegetable poisons, the most important of which are opium, hydrocyanic acid, and hyoscyamus.

#### OPIUM.

*Symptoms.*—Drowsiness, stupor, perfect insensibility, followed by delirium or profound coma; pallid countenance, deep and stertorous breathing, cold sweats, slow and full pulse, cold and livid skin, suspension of all the secretions except perspiration; sometimes convulsions, particularly in children.

*Morbid Appearances.*—Sometimes redness of the stomach and intestines, fluidity of the blood, engorgement of the lungs; but the only post-mortem appearance that appears to occur in all cases is turgescence of the vessels of the brain, sometimes with effusion of water on its surface or into the ventricles. But this condition of the brain does not furnish of itself any evidence of poisoning by opium, as it is frequently found to arise from a variety of other causes.

*Tests.*—Opium in substance: its peculiar taste, color, and odor (especially the latter quality when arising from opium in a softened or moist state, and more so if it be moderately heated; as when just obtained from the stomach).

In solution: Nitric acid throws down a red-colored (*nitrate of morphia*) precipitate. Tincture of the muriate of iron produces a (*meconate of iron*) deposit, which is also red.

When the suspected fluid has been procured from the stomach, it usually contains a mingling of organic substances; these must be broken down, and the whole rendered sufficiently fluid by addition of distilled water *when necessary*, then slightly acidulated with acetic acid; the liquor thus obtained, when well agitated, and filtered, must be evaporated to thickness of syrup. And, after being digested with alcohol, it must be boiled, and again concentrated to nearly the consistence of syrup. When this alcoholic extract (as it has been called) is dissolved in distilled water, solution of ammonia cautiously added produces a (*morphia*) precipitate, which acquires a yellow color when nitric acid is dropped into it. When the last named precipitate is removed by filtration, and acetate of lead added to the supernatant (*meconate of ammonia*) liquor, a (*meconate of lead*) deposit is afforded; which, when suspended in water, is decomposed



by transmission of sulphuretted hydrogen; a dark precipitate being thereby thrown down. The fluid remaining from this last process acquires a deep cherry-red hue on addition of the sesquichloride of iron.

**Morphia** in the solid state: Is soluble only in a slight degree in water. Nitric acid turns it first to an orange, and then to a deep orange-red. Chloride of iron gives an inky-blue color, which is destroyed by acids. Iodic acid is decomposed by morphia. To show this, the acid is to be mixed with starch; on the addition of morphia, the iodine combines with the starch, which becomes of a blue color, if the quantity is large, or of a reddish or purplish tint, if small.

*Treatment.*—The stomach to be evacuated by means of the stomach-pump, as speedily as possible. This is to be pursued until the water used no longer has the smell or taste of opium. When a stomach-pump cannot be procured, emetics of sulphate of zinc are to be administered, until the stomach is freed; strong decoctions of tea or coffee are found useful in the intervals. Cold affusions on the head, chest, and spine have been used with great success. Flagellation and other means of arousing the patient from a state of lethargy must also be employed. Bleeding should not be resorted to until all the poison has been eliminated, as the abstraction of blood tends to promote absorption. The administration of vinegar is also objectionable. The best liquid that can be given is a strong decoction of coffee. The various antidotes that have been recommended are useless.

#### HYOSCYAMUS.

*Symptoms.*—Sickness, stupor, dimness of sight and delirium, followed by coma and much dilatation of the pupils; pulse at first hard, but becoming weak and tremulous; petechia often making their appearance before death.

*Morbid Appearances.*—Inflammation of the stomach, bowels, and brain.

*Tests.*—There are no tests that can be relied upon, except a recognition of the plant and the nature of the symptoms.

*Treatment.*—The speedy evacuation of the poison, by means of emetics and purgatives, and the subsequent use of acidulous drinks.

**NARCOTICO-IRRITANTS.**—These are very closely allied in their effects to the last class, but have a more direct action on the spinal marrow and nerves, as shown by the more frequent occurrence of convulsions and paralysis. They differ much from each other in their action on the system; most of them, however, owing their properties to the presence of an alkaloidal principle. The most important are: digitalis, veratrum, conium, colchicum, lobelia, aconitum, belladonna, stramonium, tobacco, nux vomica, &c.

*Symptoms.*—These are various, but in general are vertigo, coma, delirium, paralysis, or convulsions, with disturbance and pain in the stomach and intestines. Those belonging to the nux vomica tribe have marked effects on the spinal marrow, causing tetanus and convulsions, but seldom coma or delirium; whilst, on the other hand, squill and foxglove produce symptoms of narcotism, preceded by vomiting, and other signs of irritant action on the stomach.

*Morbid Appearances.*—These, like the symptoms, are very various. In some cases, there is inflammation of the stomach and intestines; in others, this is wholly wanting. Where there have been symptoms of cerebral disturbance, traces of congestion of the brain are usually discernible, but are not found in every case.

*Tests.*—Most of these poisons, as before said, owe their deleterious powers to the presence of an alkaloidal principle, which is exceedingly difficult to detect by chemical tests; hence the strongest evidence is to be derived from an inspection of the fragments of the plant itself, if it has been administered in substance; but this mode of discrimination fails where the poison has been taken in the form of extract, infusion, or decoction. In such cases, the only reliance is on the symptoms and concomitant circumstances.

*Treatment.*—This consists in the prompt use of emetics, or the stomach-pump, followed by the administration of purgatives. No antidote can be relied upon; but, as



tannin decomposes these alkaloids, the free use of decoctions or infusions containing it may be resorted to with advantage.

**ZINC.**—This has no poisonous properties in a metallic state, but several of its salts possess active powers. Of these, the sulphate or white vitriol is the most common and most likely to occasion unpleasant symptoms.

*Symptoms.*—Pain in the abdomen, violent vomiting and diarrhoea, quick pulse, paleness and contraction of the features, coldness of the extremities. There is always a very austere taste in the mouth.

*Morbid Appearances.*—Marks of inflammation in the stomach and intestines.

*Tests.*—Ammonia gives a white precipitate, soluble in an excess of the alkali. Chromate of potassa affords an orange-red deposit, the chromate of zinc.

*Treatment.*—The free exhibition of warm water with milk. Infusions containing tannin. Where the poison has entered the bowels, emollient clysters are to be given.

### TABLE OF POISONS AND ANTIDOTES.

The following table, from Taylor "On Poisons," gives a succinct view of the most approved antidotes for the several poisons noticed in the foregoing pages:—

#### NON-METALLIC POISONS.

<i>Poisons.</i>	<i>Antidotes.</i>
Mineral Acids. { Sulphuric, Nitric, Muriatic, Nitro-Muriatic.	{ Magnesia mixed with water or milk; carbonate of lime; compound chalk powder; soda, potassa.
Vegetable Acids. { Oxalic, Tartaric.	{ Carbonate of lime.
Salts. - - - { Binoxalate of potassa.	Carbonate of lime. Sulphate of lime and water.
	Carbonate of soda in solution.
Alkalies. - - { Potassa, soda, ammonia, and their carbonates.	{ Vinegar, lemon juice, citric acid, oil.
	Sulphate of soda, potassa, magnesia or lime.
Salts - - - { Baryta and its soluble salts.	Mixture of sulphate of magnesia and vinegar.
	Carbonate of soda or ammonia.
	Alum.

#### METALLIC POISONS.

Arsenic and soluble arsenites.	{ Hydrated oxide of iron; hydrated magnesia.
Corrosive sublimate and salts of mercury.	{ Mixture of oil and lime water.
Soluble salts of lead.	{ Albumen, gluten or flour, diffused in water.
Carbonate of lead.	The alkaline sulphates.
Soluble salts of copper.	Sulphate of magnesia and vinegar.
	{ Albumen, gluten, flour diffused in water; milk.
Tartar emetic.	{ Decoctions and tinctures containing tannin.
	Magnesia.
Chloride of antimony.	Carbonate of soda; magnesia.
Salts of tin.	Milk; carbonate of soda; magnesia.
Sulphate or acetate of zinc.	Milk; carbonate of soda; magnesia.
Sulphate of iron.	Carbonate of soda or ammonia.
Nitrate of silver.	Chloride of sodium.



## NARCOTIC POISONS.

Opium; hyoseyamus.	{ Emetics; stomach pump; cold affusion. Strong decoction of coffee; electro- magnetism.
Prussic acid.	
	Ammonia; chlorine; cold affusion.

## MODES OF DISTINGUISHING SOME OF THE VEGETABLE ALKALOIDS, WHEN IN POWDER.

Treat the powder with nitric acid; this is colored red by *Brucia*, *Delphia*, *Morphia*, and by the *Strychnia* of commerce, but not the pure. If the reddened acid becomes violet on the addition of protochloride of tin, it is *Brucia*; if it becomes black and carbonaceous, it is *Delphia*. If the powder is fusible without decomposition, and decomposes iodic acid, it is *Morphia*; if it is not fusible without decomposition, and does not decompose iodic acid, it is *Strychnia*. If the powder strikes a green with nitric acid, it is *Solania*; if insoluble in ether, and does not redden nitric acid, it is *Emetia*; if soluble in ether, does not redden nitric acid, but melts and volatilizes with heat, it is *Atropia*; if thus affected by ether or nitric acid, but does not volatilize, it is *Veratria*.







# I N D E X

## OF

### DISEASES AND THEIR REMEDIES.

---

- ABSCCESS.** Lead oint., 317. Lead cataplasm, 317. Soap cataplasm, 362.
- ABSCCESS, MAMMARY.** Liniment acet. lead, 314. Oint. iod. potass. and morphia, 323.
- ACIDITY OF STOMACH.** Ammonia, 91. Lime water and milk, 135, 136. Prepared chalk, 136. Alkaline infusion cascarrilla, 150. Milk and soda water, 261. Magnesia mixt., 269. Carbonate potass., 331. Alkaline tincture, rhubarb, 351. Purgative pastilles, 366. Carb. soda, 381. Lozenges, soda, 382.
- ACNE.** Wash of benzoin, 129. Oint. subacetate lead, 315. Compound powder sulphur, 393.
- AMAUROSIS.** Ammoniac mixture, 98. Anemone, 102. Comp. powder sulphate mercury, 245. Pills strychnia, 390. Collyrium and liniment strychnia, 390. Powder white hellebore, 408.
- AMENORRHŒA.** Hiera picra, 82, and other preparations of aloes. Injection ammonia, 91. Muriate ammonia, 95. De Haen's pills, 97. Iodide calcium, 135. Cantharides and savine, 141. Infusion dittany, 185. Metallic iron, 200. Bolus metallic iron, 200. Ammoniated iron, 201. Ointment bromide iron, 202. Pills carbonate iron, 203. Mixture chloride iron, 204. Lozenges iodide iron, 206. Ointment iodide iron, 208. Black oxide iron, 211. Phosphate iron, 212. Sulphate iron, 211. Hooper's pills, 211. Compound galbanum pills, 217. Compound infusion gentian, 220. Compound tincture guaiacum, 227. Ammoniated tincture guaiacum, 228. Infusion pennyroyal, 229. Bacher's pills, 230. Tincture black hellebore, 230. Compound pills iodide mercury, 242. Saturated tincture iodine, 251. Clyster iodine, 252. Muriate manganese, 273. Horehound, 274. Myrrh and sulphate iron, 286. Tincture myrrh and hellebore, 287. Madder, 354. Decoction madder, 354. Savine and ginger, 356. Savine pills, 356. Oil savine, 357. Compound infusion senna, 372. Mixture of borax, 381.
- ANASARCA.** Calomel and squill, 240. Cremor tartar, 338. Mixture dandelion, 398.
- ANÆSTHETIC REMEDIES.** Sulphuric ether, 80. Chloroform, 159.
- ANGINA PECTORIS.** Assafetida and musk, 117.
- ANGINA SYPHILITIC.** Mercurial gargle, 233.
- APHONIA.** Decoction arnica, 113.
- APHTHÆ.** Decoction wild indigo, 124. Decoction barberry, 129. New Jersey tea, 154. Chlorine, 158. Infusion goldthread, 179. Collutory creasote, 182. Soot mouth wash, 216. Collutory rhatany, 260. Mixture carbonate potass., 333. Lozenges borax, 380. Collutory borax, 380. Honey borax, 380.
- ARSENIC, POISONING BY.** Hydrated oxide of iron, 209.
- ASCARIDES.** Suppository aloes, 84. Injection aloes, 85. Clyster Southern wood, 114. Assafetida and iron, 117. Clyster assafetida, 118. Clyster camphor, 140. Compound tincture tartrate of iron, 213. Mixture senna, 355. Infusion tansy, 397.



**ASCITES.** Electuary Per. bark and iron, 164. Pill foxglove, 196. Foxglove and tartaric acid, 191. Elaterium mixture, 194. Gamboge mixture, 219. Cremor tartar, 338. Mixture dandelion, 398.

**ASTHMA.** Ammoniac pill, 97. Ammoniac mixture, 98. Assafetida mixture, 117. Benzoin, 128. Catalpa, 152. Tincture larkspur, 189. Pill foxglove, 193. Galbanum mixture, 217. Henbane and squill, 249. Mixture elecampane, 250. Oxymel elecampane, 251. Ipecacuanha and carbonate potass., 253. Ipecacuanha and nitre, 253. Ipecacuanha and foxglove, 254. Tincture lobelia, 268. Opium and foxglove, 296. Opium and castor, 297. Solution cyanide potass., 320. Scillitic acetate potass., 329. Mixture extract elder berries, 359. Squill and ammoniac, 367. Compound tincture squill and benzoin, 369. Compound stramonium pills, 388.

**ATROPHY.** Mixture eggs, 305.

**BED SORES.** Alum liniment, 89. Mixture of tincture camphor, 140. Carbonate lead ointment, 316. Liniment and ointment tannate of lead, 318.

**BLADDER, DISEASES OF.** Emulsion cantharides, 142. Buchu, 192. Pareira brava, 306. Paullinia, 307. Mixture carbonate soda and copaiba, 382. Compound powder uva ursi, 404.

**BONES, PAINS OF.** Decoction mezereon, 278. Tincture acetate strychnia, 391.

**BRONCHITIS.** Anisated ammoniated alcohol, 94. Nitrate of silver, 111. Indian turnip, 115. Iodide calcium, 135. Creasote pills, 181. Monesia, 279. Linctus opium, 299.

**BUBO.** Mercurial plaster, 235. Anodyne ointment, 299. Plaster acetate lead, 314. Lead ointment, 317. Lead cataplasm, 317.

**BURNS, SCALDS.** Ointment wild indigo, 124. Lotion benzoin, 129. Solution chlorinated lime, 134. Lime water and oil, 135. Compound creasote ointment, 182. Carded cotton, 224. Cerate cherry laurel, 264. Liniment flaxseed, 267. Kentish's ointment, 293. Liniment eggs, 305. Basilicon ointment, 348.

**CALCULOUS AFFECTIONS.** Draught muriatic acid, 74. Phosphate of ammonia, 96. Lime water and carbonate potass., 135. Wild potato, 177. Mixture Prussian blue, 206. Nitrate of iron, 209. Magnesia and gentian, 270. Solution nitrosaccharate lead, 318. Solution caustic potass., 329. Mixture bicarbonate potass., 334. Pills carbonate soda, 381. Lozenges carbonate soda, 381. Injection carbonate soda, 382. Uva ursi, 404.

**CALCULI, BILIARY.** Turpentine and ether, 293.

**CANCER.** Arsenical powder, 69. Solution of arsenic, 69. Injection alum, 89. Acetate ammonia, 92. Iodide arsenic, 113. Chloride gold, 121. Chloride barium, 125. Compound pills belladonna, 127. Extract marigold, 137. Ointment marigold, 138. Carrot ointment, 148. Infusion hemlock, 175. Cataplasm hemlock, 178. Arseniate of iron, 201. Pills red oxide iron, 212. Compound soot ointment, 216. Hop ointment, 232. Infusion cherry laurel, 264. Ointment cherry laurel, 264. Ointment turpentine, 269. Ointment carbonate lead, 316. Ointment chloride lead, 316. Lotion chloride lead, 316. Ointment iodide potass. and opium, 323. Solution chloride soda, 383. Chloride zinc, 412.

**CANCERUM ORIS.** Pyroligneous acid, 69. Solution chloride potass., 334.

**CARDIALGIA.** Oil of wormwood, 64. Mixture boracic acid, 70. Acetic ether, 79. Ammonia, 91. Lime water and potass., 136. Pills gentian and ammonia, 220. Magnesia mixture, 269. Nux vomica, 288. Rhubarb and chalk, 348. Carbonate soda and rhubarb, 381.

**CARIES.** Phosphoric acid, 75. Creasote ointment, 182.

**CATARACT.** Anemone, 102.

**CATARRH.** Gum mixture, lozenges, &c., 66. Mixture benzoic acid, 70. Garlic, 81. Ammoniated anisated alcohol, 94. Ammoniac mixture, 98. Ammoniac and nitric



- acid, 98. Almond emulsion, 101. Starch lozenges, 102. Lozenges kermes mineral, 106. Syrup kermes mineral, 106. Catechu and liquorice, 152. Powder spermaceti, 155. Spermaceti mixture, 155. Mixture bittersweet, 194. Prepared liquorice, 224. Mixture henbane, 249. Compound pills elecampane, 250. Mixture elecampane, 250. Compound oxymel elecampane, 251. Ipecacuanha lozenges, 254. Lozenges ipecacuanha and camphor, 254. Lozenges lactucarium, 263. Horehound syrup and candy, 273. Pectoral hydromel, 276. Mixture balsam Peru, 285. Myrrh and squills, 286. Solution cyanide potass., 320. Nitrate potass. and orris root, 335. Syrup red poppies, 352. Squill and ipecacuanha, 367. Seneka and prep., 371-2. Sulphur and liquorice, 393. Sulphur and orris root, 393. Turpentine and myrrh, 399. Emulsion balsam tolu, 401. Mixture tolu and morphia, 401.
- CEPHALALGIA.** Asarabacca, 115. Ward's essence, 140. Liniment oil amber, 392.
- CEREBRAL AFFECTIONS.** Clyster tartar emetic, 108. Cataplasm horseradish, 112. Clyster colocynth, 173. Infusion galium, 218. Mixture borotartrate potass., 331. Clyster common salt and arnica, 379.
- CHANCER.** Syrup gold, 120. Ointment gold, 120. Creasote lotion, 182. Mercurial lotion, 233. Mercurial liniment, 235. Black wash, 244.
- CHAPS.** Glycerin, 233. Anodyne ointment, 299. Liniment carbonate potass., 332. Cold cream, 352. Ointment elder leaves, 359. Camphor soap, 360.
- CHILBLAINS.** Muriatic acid lotion, 74. Sulphuric acid liniment, 76. Alum ointment, 90. Solution chloride lime, 134. Camphor ointment, 140. Compound ointment creasote, 183. Ointment henbane, 249. Lotion iodine, 251. Liniment balsam Peru, 285. Turpentine lotion, 292. Embrocation petroleum, 307. Soap iodide potass., 323. Lotion carbonate potass., 332. Liniment pellitory, 340. Rose oil, 352. Mustard liniment, 378. Borax ointment, 381.
- CHLOROSIS.** Pills aloes and iron, 82. De Haen's pills, 97. Metallic iron, 200. Compound powder metallic iron, 200. Pills metallic iron, 200. Bolus ammoniated iron, 201. Pills carbonate iron, 203. Tincture chloride iron, 204. Bath iodide iron, 208. Black oxide iron, 210. Electuary black oxide iron, 210. Sulphate iron, 211. Bland's pills, 211. Syrup sulphate iron, 212. Tartrate of iron and potass., 212. Tannate iron, 213. Compound galbanum pills, 217. Oxide manganese, 273. Chocolate with paullinia, 307. Compound rhubarb pills, 349. Decoction madder, 354. Compound powder valerian, 405.
- CHOLERA.** Mixture nitric acid, 74. Liniment nitric acid, 74. Compound bismuth powder, 130. Camphor powder, 138. Extract hemp, 141. Infusion logwood, 228. Sugar milk draught, 261. Opium and acetate lead, 297. Compound pill acetate lead, 314. Mixture bicarbonate potass., 334. Compound powder chloride potass., 334. Plaster sulphate quinine, 347.
- CHOLERA INFANTUM.** Milk and suet, 261. Acetate lead, 313. Infusion Beane, 377.
- CHOREA.** Ammoniated copper, 186. Carbonate iron, 202. Mixture Dippel's animal oil, 289. Mixture cyanide potass., 320. Oxide zinc, 410. Ethereal tincture chloride zinc, 411.
- COLIC.** Alkaline tincture wormwood, 64. Aniseed, 103. Dewees' carminative, 118, 270. Clyster assafetida, 118. Infusion caraway, 148. Embrocation caraway, 149. Clove plaster, 149. Infusion catnep, 152. Infusion cinnamon, 167. Infusion fennel, 215. Oil pennyroyal, 229. Oil laurel, 267. Mace, 269. Dalby's carminative, 270. Plaster opium and camphor, 298. Opium liniment, 303. Cataplasm black pepper, 310. Emulsion carbonate potass., 332. Tinct. rhubarb and gentian, 350. Compound sagapenum pills, 357. Terebinthinate balsam sulphur, 394. Infusion tobacco, 396. Clyster turpentine, 399. Turpentine mixture, 400.
- COLICA PICTONUM.** Sulphuretted water, 73. Sulphuric acid lemonade, 76. Alum julep, 89. Alum mixture, 90. Liniment belladonna, 128. Opium and sulphur, 296. Opium and cinnamon water, 300.



- CONDYLOMATA.** Lotion corrosive sublimate and camphor, 238. Red oxide mercury, 244.
- CONSTIPATION.** Acetic acid clyster, 68. Preparations aloes, 82. Assafetida and aloes, 82. Pills elaterium, 194. Extract butternut, 257. Compound pills buckbean, 278. Compound rhubarb pills, 349. Rhubarb and ox gall, 349. Compound solution sulphate of soda, 384. Rochelle salts, 385. Compound powder sulphur, 393. Clyster turpentine, 399.
- CONTUSIONS.** Sulph. acid and alcohol, 76. Lotion sal ammoniac, 95. Liniment sal ammoniac, 95. Arnica and rue, 113. Lotion sulphate copper, 188. Cataplasm hops, 232. Arquebusade water, 298. Compound spirit nutmeg, 285. Fomentation carbonate potass., 332. Lotion nitrate potass., 336. Wine of roses, 353. Aromatic fomentation, 353. Aromatic vinegar, 353. Fomentation common salt, 378. Turpentine liniment, 399.
- CONVULSIONS.** Garlic, 81. Assafetida mixture, 117. Musk clyster, 284.
- CORNEA, AFFECTIONS OF.** Anemone, 102. Ointment cyanide silver, 110. Ointment iodide silver, 111. Ointment sulphate cadmium, 133. Powder calomel and sugar, 239. Compound oint. cod liver oil, 290. Collyrium caustic potass., 329. Collyrium carbonate potass., 332.
- CORNS.** Plaster ammonia, 91. Verdigris ointment, 186. Verdigris plaster, 186. Lotion iodine, 251.
- CORROSIVE SUBLIMATE, ANTIDOTE.** Powder gold and iron, 120.
- CORYZA.** Lozenges cubebs, 184. Injection opium, 297.
- COUGH.** Mixture gum Arabic, 65. Almond emulsion, 100. Green linctus, 100. White linctus, 101. Antimonial wine and ammoniac, 109. Assafetida and squill, 117. Camphorated cough mixture, 141. Tincture cochineal, 169. Mixture quince seeds, 189. Barley sugar, 231. Lactucarium, 262. Brown cough mixture, 303. Opiate linctus, 303. Pectoral syrup, 306. Mixture oil amber, 392. Mixture tolu and belladonna, 401. Oxide zinc, 410.
- CRAMP IN THE STOMACH.** Ammoniated tincture of castor, 151. Ferrocyanide zinc, 412.
- CROUP.** Sulphuretted syrup, 95. Sulphate copper, 187. Syrup sulphuret potass., 327. Compound syrup sulphuret potass., 327. Mixture sulphuret potass., 328. Compound syrup squill, 368. Oxymel squill and valerian, 369. Mixture seneka, 372. Cataplasm tobacco, 396.
- CUTANEOUS DISEASES.** Hydrosulphuretted bath, 72. Muriatic acid bath, 74. Lotion alum, 89. Arseniate ammonia, 92. Hydriodate ammonia, 94. Sulphuretted hydrosulphate ammonia, 95. Sulphuret antimony, 105. Lotion tartar emetic, 108. Iodide arsenic, 113. Donovan's solution, 114. Chloride barium, 123. Solution bromine, 130. Lime ointment, 136. Anthracokali, 146. Ointment iodide carbon, 147. Creasote ointment, 182. Decoction bittersweet, 193. Extract bittersweet, 194. Mixture soot, 216. Fuligokali, 216. Decoction guaiacum, 226. Guaiacum and sulphur, 227. Compound mercurial powder, 233. Solution acetate mercury, 236. Ointment white precipitate, 236. Bateman's lotion, 238. Plummer's pill, 240. Calomel ointment, 241. Ointment red iodide mercury, 243. Ointment nitrate mercury, 243. Ointment sulphate mercury, 246. Black sulphuret mercury, 246. Red sulphuret mercury, 246. Muriate manganese, 273. Compound decoction mezereon, 279. Cod liver oil, 290. Phosphorated cerate, 309. Pills Burgundy pitch, 312. Tar water, 312. Lotion cyanide potass., 321. Ointment iodide potassa and mercury, 323. Ioduretted water, 323. Sulphuret potassa, 326. Lotion sulphuret potass., 329. Barlow's lotion, 327. Mixture sulphuret potass., 328. Aromatic bath, 353. Extract elder berries, 359. Sarsaparilla, 362-5. Ointment bromide soda, 379. Sulphuret soda, 379. Arseniate soda, 380. Soda and ipecacuanha, 381. Ointment carbonate soda, 382. Hydrosulphate soda, 383. Compound pills sulphuret soda, 384. Lotion sulphuret soda, 385. Sulphur and cremor tartar, 393. Poison oak, 402. Decoction elm bark, 403. Decoction white hellebore, 408. Iodide zinc, 412.



- DEAFNESS.** Cayenne pepper lozenges, 145. Acoustic balsam, 285. Injection caustic potass., 329.
- DEBILITY.** Preparations cacao, 132. Camphor mixture, 139. Hartshorn jelly, 180. Preparations iron, 200. Barley meal, 231. Clyster eggs, 305. Mixture eggs, Chocolate and Paullinia, 306.
- DELIRIUM TREMENS.** Opium and musk, 296. Laudanum and tartar emetic, 304.
- DIABETES.** Phosphoric acid, 75. Hydrosulphate ammonia, 94. Compound pills ammoniated copper, 187. Compound pills kino, 259. Pills acetate morphia, 281. Carbonate potass. and ammonia, 334. Sulphur mixture, 394.
- DIARRHOEA.** Tannic acid, 76. Acetate alumine, 87. Alum, 87-8. Angustura, 103. Mixture chamomile, 104. Antimonial wine and laudanum, 109. Arnica, 112. Dewees' carminative, 118, 270. Clyster bistort., 130. Compound powder chalk, 136. Lozenges chalk, 136. Chalk mixture, 137. Camphor water and laudanum, 139. Camphor water and nitric acid, 139. Compound cascarilla powder, 150. Catechu, 152. Chlorine clyster, 159. Peruvian bark and rhatauy, 163. Electuary Peruvian bark, 164. Infusion columbo and ginger, 174. Columbo and cascarilla, 174. Creasote mixture, 182. Sulphate copper, 187. Electuary sulphate copper, 188. Wine persimmons, 193. Infusion black purslane, 195. Mixture ammoniated iron, 201. Persesquinitrate iron, 209. Compound infusion galls, 218. Syrup galls, 218. Geranium, 222. Decoction pomegranate rind, 225. Infusion logwood, 228. Electuary logwood, 228. Compound powder kino, 258. Electuary kino, 259. Compound powder rhatauy, 259. Monesia, 279. Injection acetate morphia, 281. Opium and chalk, 296. Roasted opium, 296. Plaster opium and camphor, 298. Mixture wine of opium, 301. Sedative mixture, 306. Bolus allspice, 309. Acetate lead, 313. Acetate lead mixture, 315. Decoction oak bark, 342. Confection acorns, 342. Roasted rhubarb, 348. Rhubarb mixture, 357. Confection dog rose, 352. Electuary roses, 353. Decoction blackberry root, 354. Infusion benne, 377. Hardhack, 386. Compound turpentine plaster, 400. Tormentil, 401. Infusion slippery elm, 403.
- DISINFECTION.** Prophylactic vinegar, 68. Gaseous muriatic acid, 73. Chloride of lime, 134. Chlorine, 158. Solution chloride potass., 320. Solution chloride soda, 383.
- DROPSY.** Indian hemp, 109. Infusion horseradish, 112. Compound spirit horseradish, 112. Decoction asparagus, 116. Extract asparagus, 116. Chloride gold and soda, 121. Hairy horehound, 124. Wine bryony, 131. Camphorated ethereal tincture cantharides, 143. Vinegar cayenne pepper, 145. Infusion carrot seeds, 148. Celandine, 157. Pipsissewa, 157. Black snakeroot, 160. Vinegar colchicum, 171. Colchicum and squill, 172. Horsebalm, 172. Colocynth, 172. Powdered foxglove, 190. Pills foxglove, 191. Infusion foxglove, 191. Elaterium, 194. Canada fleabane, 197. Gamboge, 219. Black hellebore, 229. Bacher's pills, 230. Compound wine black hellebore, 230. Jalap and cremor tartar, 256. Compound powder jalap, 256. Compound infusion juniper, 257. Preparations juniper, 257. Mustard whey, 261. Aromatic tincture lettuce, 263. Starkey's soap, 293, 360. Embrocation petroleum, 307. Mixture petroleum, 308. Infusion parsley root, 308. Solution hydriodate potass., 325. Scillitic acetate potass., 329. Liquid acetate potass., 330. Mixture acetate potass., 330. Borotartrate potass., 331. Nitrate potass. and squill, 335. Mixture nitrate potass., 336. Cremor tartar, 338. Tartrate potass. and ammonia, 339. Decoction elder bark, 359. Confection scammony, 366. Squill, 366-71. Squill and nitrate potass., 367. Squill and cremor tartar, 367. Infusion and decoction broom, 371. Acetate soda, 379. Solution carbonate soda, 382. Pills tobacco, 396. Wine tobacco, 396. Decoction dandelion, 398. Extract dandelion, 398. Electuary turpentine, 399. Veratria, 407.
- DROPSY, OVARIAN.** Ointment iodine and mercury, 252. Solution iodide potass., 321.
- DYSENTERY.** Mixture nitric acid, 74. Starch jelly, 102. Starch and suet, 102. Angustura, 103. Clyster bistort., 130. Camphor water and laudanum, 139.



- Camphor water and nitric acid, 139. Electuary catechu, 183. Infusion black purslane, 198. Mixture extract logwood, 225. Calomel and opium, 239. Decoction ipecacuanha, 254. Mixture ipecacuanha, 254. Emulsion jalap, 257. Electuary kino, 259. Mixture rhatany, 260. Milk and suet, 261. Mixture cherry laurel, 264. Injection morphia, 280. Nux vomica, 288. Mixture nux vomica and aloes, 289. Oleaginous mixture, 291. Opium and acetate lead, 297. Opiate, 298. Plaster opium and camphor, 298. Opium and syrup poppies, 300. Confection dogrose, 352. Syrup blackberries, 355. Compound powder sulphur, 393. Infusion slippery-elm, 403. Compound pills sulphate zinc, 414.
- DYSMENORRŒA.** Infusion Southern wood, 114. Compound pills ergot, 197. Clyster iodine, 252.
- DYSPEPSIA.** Wormwood, 64. Aloes pills, 82-84. Aniseed, 103. Orange peel and rhubarb, 119. Cayenne pepper pills, 145. Infusion cinnamon, 167. Powder coriander, 180. Pill prepared metallic iron, 200. Mixture malate iron, 209. Nitrate iron, 209. Mixture soot, 216. Infusion gentian and rhubarb, 221. Mixture gentian and sulphuric acid, 221. Masterwort, 231. Infusion hops, 232. Mercury, chalk, and ipecacuanha, 233. Bolus ipecacuanha, 253. Ipecacuanha and centaury, 254. Rice jelly, 305. Solution hydrargyro-iodide potass., 325. Sulphate potass. and rhubarb, 337. Rhubarb pills, 349. Soda and rhubarb, 381. Mixture carbonate soda and quassia, 382. Compound turpentine plaster, 400. Oxide zinc and columbo, 410.
- DYSPNŒA.** Opium and castor, 297.
- DYSURIA.** Benzoic acid and copaiba, 70. Tartar emetic and phosphate lime, 107. Infusion fleabane, 197. Pills nitrate potass., 335. Mixture tobacco, 396.
- EAR, AFFECTIONS OF.** Injection morphia, 280. Acoustic balsam, 285. Antitotic mixture, 299. Opium liniment, 303.
- ECCHYMOSES.** Juniper liniment, 258. Lotion nitrate potass., 336.
- ELEPHANTIASIS.** Mudar, 138.
- ENLARGEMENTS, GLANDULAR.** Ointment animal charcoal, 146.
- ENGORGEMENTS, VISCERAL.** Pills black oxide iron, 212. Electuary black oxide iron, 212. Calomel and dandelion, 240. Calomel and squill, 240. Sulphate mercury, 246. Compound powder rhubarb, 348. Tartrate soda and rhubarb, 385. Mixture tartrate soda, 385. Tartrate soda whey, 385. Compound infusion dandelion, 398.
- EPILEPSY.** Nitrate silver, 111. Belladonna, 126. Oil box, 131. Electuary Peruvian bark, 164. Muriate copper, 186. Ammoniated copper, 186. Ammoniated copper and belladonna, 187. Compound pills sulphate copper, 188. Ammoniated iron, 201. Prussian blue, 205. Wine galium, 218. Masterwort, 231. Calomel and opium, 241. Indigo, 250. Opium and nitrate silver, 297. Tincture opium and asarabacca, 302. Phosphuretted oil, 308. Mixture biniodide potass., 324. Compound powder valerian, 404. Oxide zinc, 410. Compound pills oxide zinc, 410. Ethereal tincture chloride zinc, 411. Compound pills sulphate zinc, 414.
- EPISTAXIS.** Alum, 88-9. Sulphate copper styptic, 188. Injection sulphate iron, 211.
- EXCORIATION.** Cold cream, 155, 352. Carbonate lead ointment, 316. Plaster carbonate lead, 316. Ointment tannate lead, 318. Rose oil, 352. Camphor soap, 360. Ointment oxide zinc, 411. Cerate carbonate zinc, 413.
- EXANTHEMATA.** Compound mercurial powder, 233. Sulphuret magnesia, 272. Mixture Virginia snakeroot and allspice, 376. Compound infusion elm bark, 403.
- EYE, AFFECTIONS OF.** Atropia, 118. Camphor ointment, 140. Camphor collyrium, 141.
- FACE, ERUPTIONS ON, BLOTCHES, &c.** Wash sal ammoniac, 96. Almond paste,



98. Almond powder, 99. Compound almond lotion, 99. Milk roses, 99, 100, 129. Goulard's lotion, 100. Cosmetic liniment, 101. Water anemone, 102. Wash of benzoin, 129. Emulsion benzoin, 129. Spermaceti liniment, 155. Sultana ointment, 155. Emulsion corrosive sublimate, 237. Cosmetic wash, 238. Lotion carbonate potass., 332. Beef marrow soap, 360. Aromatic soap, 360. Cosmetic soap powder, 361. Essence of soap, 361.

FAINTING. Aromatic vinegar, 68. Aromatic carbonate of ammonia, 92.

FEVER. Vinegar mixture, 67. Syrup vinegar, 67. Acetate ammonia, 91. Citrate ammonia, 94. Nitrate ammonia, 96. Tartar emetic, 107. Mixture extract centaury, 154. Boneset, 198. Compound powder ammoniated iron, 201. Mixture avens, 223. Calomel and jalap, 239. Calomel and nitrate potass., 239. Ipecacuanha and tragacanth, 283. Infusion malt, 272. Citrate potass., 324. Nitrate potass., 335. Nitrous powders, 335. Mixture nitrate potass., 336. Mixture sulphate potass., 337.

FEVER, HECTIC. Compound infusion boneset, 198. Gœlis' powder, 284. Compound powder sulphate quinine, 345. Compound infusion sage, 359.

FEVER, INTERMITTENT. Pills arsenic and opium, 70. Lockstadt's pills, 101. Golden sulphuret antimony, 106. Tartar emetic and quinine, 107. Electuary orange peel, 119. Bebeerine, 126. Camphor water and ether, 139. Cetrarine, 156. Preparations Peruvian bark, 160-66. Cinchonia, 166. Decoction coffee, 170. Dogwood, 180. Pills sulphate copper, 187. Sulphate copper and opium, 188. Persimmon, 193. Compound powder Prussian blue, 205. Horse chestnut, 247. Tulip tree bark, 267. Magnolia, 272. Narcotina, 288. Phloridzine, 308. Phosphorated oil turpentine, 309. Piperine, 311. Mixture acetate potass., 330. Arsenical solution, 330. Arseniate potass., 330. Mixture citrate potass. and bark, 335. Oxalate potass., 336. Quassia, 341. Oak bark, 341. Quinine and salts, 242-47. Salicine, 358. Willow bark, 358. Virginia snakeroot, 376. Common salt and lemon juice, 319. Arseniate soda, 380. Solution chloride soda, 383.

FEVER, TYPHUS. Carbonate ammonia, 93. Wild indigo, 124. Clyster camphor, 140. Liniment cantharides, 143. Tincture cayenne pepper, 145. Musk mixture, 283. Turpentine mixture, 292. Laudanum and tartar emetic, 303. Egg and brandy mixture, 305. Egg and wine mixture, 305. Mixture Virginia snakeroot and allspice, 376. Wine Virginia snakeroot and vanilla, 376. Tincture Virginia snakeroot and balsam Peru, 376. Mustard whey, 377.

FISSURES, ANUS. Ointment acetate lead, 314.

FISTULÆ. Injections copaiba, 178. Injection corrosive sublimate, 238. Injection myrrh, 287. Injection iodide potass., 321. Injection iodine, 324. Compound lotion iodide potass., 324.

FLATULENCE. Ammoniated alcohol, 94. Tincture angelica, 103. Aniseed, 103. Chamomile pills, 104. Tincture assafetida and soot, 118. Assafetida plaster, 118. Elixir orange peel, 119. Tincture benzoin, 129. Calamus, 133. Aromatic powder and confection, 147. Compound tincture cardamom, 148. Spirit caraway, 149. Preparations cloves, 149. Cinnamon, 167. Coriander, 180. Magnesia and camphor, 270. Peppermint, 277. Nutmeg, 284. Confection opium, 298. Water allspice, 309. Electuary black pepper, 310. Compound tincture sassafras, 361. Purgative pastilles, 366. Mixture carbonate soda and gentian, 382. Compound turpentine plaster, 400. Ginger, 415.

FÆTOR ORIS. Mixture nitric acid, 94. Alum, 88. Lozenges chloride of lime, 134. Charcoal lozenges, 146. Lozenges catechu, 153. Pastilles catechu, 153.

FROST, EFFECTS OF. Rust's ointment, 90. Oil elaterium, 195. Compound ointment henbane, 249. Embrocation petroleum, 307.

FUNGOUS FLESH. Burnt alum, 87. Chloride antimony, 105. Lotion orpiment, 114. Verdigris and savine, 185. Verdigris ointment, 186.

GANGLIONIC SYSTEM, DISEASES OF. Pills Prussian blue, 206.



- GANGRENE.** Compound camphor pills, 138. Peruvian bark and arnica, 161. Cataplasm Peruvian bark, 166. Anodyne ointment, 299. Cataplasm oak bark, 342.
- GASTRALGIA.** Mixture aconite, 78. Subnitrate bismuth, 129. Syrup codeia, 169.
- GASTRODYNIA.** Oxide silver, 111. Compound bismuth powder, 129. Compound tincture cinnamon, 168. Oil of ergot, 197. Mercurial mixture, 274.
- GLANDS ENLARGED.** Plaster ammoniac with mercury, 98. Cataplasm bryony, 131. Ointment chloride lime, 134. Animal charcoal, 145. Hemlock and dandelion, 175. Plaster foxglove, 191. Ointment iodide iron, 208. Compound galbanum plaster, 217. Compound mercurial liniment, 235. Ointment iodide mercury, 243. Red sulphuret mercury, 246. Liniment henbane, 249. Ethereal tincture iodine, 251. Mixture iodine and iodide potass., 231. Sulphuret potass., 326. Compound plaster sulphuret potass., 328. Burnt sponge, 387.
- GLEET.** Tannic acid, 76. Tincture cantharides and guaiacum, 142. Cubebs and ergot, 183. Compound powder ergot, 196. Pills guaiacum and turpentine, 227, 399. Injection chloride zinc, 411.
- GLOTTIS, SPASM OF.** Cataplasm tobacco, 396.
- GOITRE.** Solution chloride calcium, 133. Animal charcoal, 145. Iodine ointment, 252. Cataplasm iodine, 252. Iodide potass., 321. Ointment iodide potass., 322. Liniment iodide potass., 323. Sulphuret potass., 326. Burnt sponge, 387.
- GONORRHOEA.** Alumine, 87. Alum, 89. Nitrate silver, 112. Assafetida and opium, 117. Oil box, 131. Pills chloride calcium, 134. Emulsion hemp, 141. Compound bolus catechu, 152. Electuary catechu, 153. Wine Peruvian bark and calamus, 166. Copaiba and its preparations, 177-8. Cubebs, 183-4. Injection ammoniated copper, 187. Injection sulphate copper, 188. Compound powder ammoniated iron, 201. Injection iodide iron, 208. Pills sulphate iron, 211. Tincture galls, 218. Oil guaiacum, 227. Injection corrosive sublimate, 238. Calomel and catechu, 240. Saturated tincture iodine, 251. Injection opium, 299. Sedative injection, 306. Oil parsley, 308. Acetate lead mixture, 315. Injection sulphuret potass., 327. Injection caustic potass., 329. Pills nitrate potass. and camphor, 335. Nitrated emulsion, 336. Starkey's soap, 360. Turpentine and rhubarb, 399. Turpentine mixture, 400. Injection acetate zinc, 413. Fomentation sulphate zinc, 414. Injection sulphate zinc, 414.
- GOUT.** Aconite, 77. Ammoniacal liniment, 91. Carbonate ammonia, 93. Phosphate ammonia, 96. Ammoniacal succinic acid, 97. Golden sulphuret antimony, 106. Sulphuret calcium, 135. Camphorated ether, 140. Portland powder, 154. Pills pipsissewa, 158. Colchicum, 170, 171. Wine hedge hyssop, 226. Pills guaiacum and sulphur, 227. Ammoniacal tincture guaiacum, 228. Magnesia and colchicum, 270. Confection opium, 298. Phosphorated oil, 308. Mixture caustic potass., 329. Solution silicate potass., 337. Warner's cordial, 350-1. Extract elder berries, 359. Pills soap and ox gall, 361. Conserve broom, 369. Compound tincture senna, 374. Liniment sulphuret carbon, 395. Turpentine mixture, 400. Veratria, 407. Wine white hellebore, 418.
- GRAVEL.** Mixture Prussian blue, 206. Carbonate potass. and ammonia, 334.
- GUMS, AFFECTIONS OF.** Boracic acid, 70. Alumine, 87. Electuary alum, 88. Gargle alum, 89. Collutory catechu, 153. Collutory creasote, 182. Gargle sulphate zinc, 414.
- HEADACHE.** Mixture vinegar and cardamom, 67. Lotion ammonia, 91. Carbonate ammonia, 93. Vicat's mixture, 94. Ward's essence, 140. Castor plaster, 151. Sneezewort, 229. Mixture valerian and ammonia, 405. Valerianate zinc, 415.
- HEMATEMESIS.** Pills acetate lead, 313.
- HEMOPTYSIS.** Powder starch, 101. Carrageen mixture, 160. Electuary Peruvian bark and catechu, 164. Creasote mixture, 181. Bolus foxglove, 190. Mixture tincture foxglove, 191. Sedative mixture, 311. Compound pills acetate lead, 314. Tolu mixture, 401. Mixture uva ursi, 404.



- HEMORRHAGES.** Alum, 87, 88. Alum whey, 89. Bistort, 130. Alum and catechu, 152. Sulphate copper, 187. Compound powder ergot, 196. Extract ergot, 197. Chloride iron and acetate lead, 204. Tincture chloride iron, 204. Solution sulphate iron, 211. Compound powder galls, 218. Avena, 222. Compound powder kino, 259. Mixture rhatany, 260. Alum whey, 261. Matico, 275. Roasted opium, 296. Opium and acetate lead, 313. Compound pills acetate lead, 314. Acetate lead, 313. Mixture bistort, 319. Powder sulphate soda and opium, 384.
- HEMORRHOIDS.** Aloetic mixture, 85. Alum ointment, 90. Lime ointment, 136. Charcoal suppository, 146. Extract Peruvian bark, 166. Hemlock oil, 176. Suppository hemlock, 176. Saffron ointment, 183. Oil elaterium, 195. Ointment galls, 219. Compound gall ointment, 219. Witch hazel, 229. Ointment mercury and belladonna, 235. Ointment red oxide mercury and tin, 245. Ointment henbane, 249. Electuary jalap, 256. Ointment matico, 276. Injection morphia, 280. Anodyne ointment, 299. Ointment opium and tar, 299. Laudanum ointment, 303. Confection black pepper, 310. Ward's paste, 310. Pile ointment, 314. Lead ointment, 317. Mixture tartrate potass., 338. Mixture bitartrate potass., 339. Ointment figwort, 371. Electuary senna and sulphur, 373. Ointment tin, 387. Ointment stramonium, 389. Sulphur electuary, 393. Fomentation sulphate zinc, 414.
- HAIR, LOSS OF, &c.** Garlic, 81. Almond cream, 100. Macassar oil, 101. Fixature, 101, 292. Hair dye, 111. Compound tincture cantharides, 142. Pomatum cantharides, 144. Shampoo liquid, 144. Pomatum Peruvian bark, 165. Bandonline, 189. Lotion galls, 218. Compound ointment balsam Peru, 286.
- HAIR, TO REMOVE.** Orpiment and Powder orpiment, 114.
- HEART, AFFECTIONS OF.** Syrup asparagus, 116. Bromine, 130. Foxglove and acetate lead, 191. Digitalin, 192. Mixture iodide potass., 322.
- HEMICRANIA.** Compound powder valerian, 405. Electuary valerian, 405.
- HEMIPLEGIA.** Pills poison oak, 402.
- HEPATITIS.** Mixture nitric acid, 74. Bath nitromuriatic acid, 75. Compound mercurial pill, 234. Chlorate potass., 334. Dandelion, 398.
- HERNIA.** Clyster belladonna, 128. Infusion tobacco, 396.
- HERPES.** Chlorine liniment, 159. Arseniate iron, 201. Ointment carburet iron, 204. Decoction soot, 215. Ointment black hellebore, 230. Liniment acetate mercury, 236. Compound calomel pill, 240. Calomel ointment, 241. Ointment red oxide mercury and lead, 245. Ointment sulphate mercury, 246. Red sulphuret mercury, 247. Ointment red sulphuret mercury, 247. Lotion sulphuret potass., 327.
- HICCUGH.** Ammoniated alcohol, 94. Assafetida, 111.
- HOOPING-COUGH.** Succinated ammonia, 96. Extract anemone, 102. Kermes' mineral, 106. Antimonial wine and bittersweet, 109. Assafetida mixture, 117. Assafetida and tolu, 117. Assafetida plaster, 118. Belladonna, 126-7. Fumigating powder, 128. Emulsion cantharides, 142. Tincture bark and cantharides, 165. Syrup Peruvian bark, 166. Cochineal and carbonate potass., 169. Confection ipecacuanha, 254. Mixture ipecacuanha, 254. Compound syrup ipecacuanha, 255. Alkaline wine ipecacuanha, 255. Syrup sulphuret magnesia, 272. Myrrh and zinc, 304. Lettsom's elixir, 303. Mixture carbonate potass., 333. Compound syrup squill, 368. Carbonate soda and ipecacuanha, 382. Tincture artificial musk, 391. Liniment oil of amber, 392. Compound powder tobacco, 396. Mixture balsam tolu and copaiba, 401. Sulphate zinc, 414. Mixture sulphate zinc, 415.
- HYDROPHOBIA.** Extract hemp, 141. Tonquin powder, 282.
- HYDROCEPHALUS.** Powder foxglove, 190. Calomel and foxglove, 239. Ointment iodide potass. and mercury, 323.
- HYDROTHORAX.** Tincture colchicum and foxglove, 171. Mixture colocynth, 173.



- Pills foxglove, 190. Mixture elaterium, 194. Gamboge mixture, 219. Extract lettuce, 263. Mixture tobacco, 396.
- HYPOCHONDRIA.** Assafetida, 117. Mixture lettuce and dandelion, 262.
- HYPOPION.** Pills seneka, 371.
- HYSTERIA.** Mixture acetic ether, 79. Carbonate ammonia, 93. Succinate ammonia, 97. Mixture chamomile, 104. Infusion Southern wood, 114. Assafetida and iron, 117. Assafetida mixture, 117. Alkaline tincture assafetida, 117. Tincture assafetida and soot, 118. Assafetida and castor, 118. Bolus castor, 151. Castor pills, 151. Skunk cabbage, 193. Soot pills, 215. Compound galbanum pills, 217. Mixture lactucarium, 262. Musk, 283. Pills opium and musk, 297. Tincture opium and asarabacca, 302. Mixture rue and squills, 355. Liniment oil of amber, 392. Compound powder valerian, 405. Vanilla, 407. Valerianate zinc, 415.
- ILEUS.** Hemlock oil, 176.
- IMPETIGO.** Lotion hydrocyanic acid, 72. Glycerine, 223. Compound powder sulphur, 393. Powder iodide sulphur, 395. Ointment oxide zinc and opium, 411.
- INDURATIONS.** Pills calomel and antimony, 240.
- IRITIS.** Turpentine mixture, 292.
- ISCHURIA.** Assafetida and opium, 117.
- ITCH.** Sulphuric acid ointment, 76. Lime water and sulphur, 135. Lime ointment, 136. Chlorine ointment, 159. Lotion corrosive sublimate, 237. Ointment red oxide mercury and sulphur, 245. Tincture sulphuret potass., 327. Compound lotion sulphuret potass., 327. Ointment carbonate potass., 333. Soap and sulphur, 362. Lotion sulphuret iodine, 379. Decoction stavesacre, 388. Compound powder sulphur, 393. Sulphur ointment, 394. Compound sulphur ointment, 394. Ointment white hellebore, 409.
- JAUNDICE.** Barberry, 129. Compound wine centaury, 184.
- JOINTS, AFFECTIONS OF.** Pipsissewa beer, 158. Compound decoction guaiacum, 226. Pills guaiacum and antimony, 227. Ointment calomel and squills, 241. Oil laurel, 263.
- KIDNEYS, AFFECTIONS OF.** Emulsion manna, 273. Emulsion carbonate soda, 382. Ointment turpentine, 399.
- LABOR.** Contrayerva mixture, 177. Ergot and its preparations, 196-7. Decoction cotton root, 224. Compound powder borax, 380.
- LARYNGITIS.** Creasote pills, 181. Liniment croton oil, 294. Plaster croton oil, 295.
- LEUCORRHEA.** Injection gallic acid, 71. Tannic acid, 76. Whytt's pills, 84. Alum, 88. Electuary catechu, 153. Peruvian bark and isinglass, 161. Wine Peruvian bark and calamus, 166. Cubebs and ergot, 183. Compound powder ergot, 196. Infusion black purslane, 198. Compound pills carbonate iron, 203. Lozenges iodide iron, 206. Bath iodide iron, 208. Ointment iodide iron, 208. Black oxide iron, 210. Electuary black oxide iron, 211. Sulphate iron, 211. Syrup sulphate iron, 212. Solution tartrate iron and potass., 212. Injection soot, 216. Tincture galls, 218. Avens, 222. Compound powder guaiacum, 227. Calomel and catechu, 240. Saturated tincture iodine, 251. Injection rhatany, 260. Injection subacetate lead, 314. Iodide potass., 321. Decoction oak bark, 342. Pills turpentine and guaiacum, 399.
- LEPRA.** Pills arsenic and opium, 70. Iodide arsenic, 113. Donovan's solution, 114. Barytic liniment, 126. Mudar, 138. Chlorine oil, 159. Mixture tincture colocynth, 173. Carburet iron, 203. Ethereal solution bromide mercury, 276. Oint. naphthaline, 287.



- LICHEN.** Carburet iron, 203. Ointment glycerine, 223. Ointment calomel and camphor, 241. Ointment cyanide mercury, 242.
- LIPS, EXCORIATION OF.** Spermaceti lip salve, 155.
- LIVER, AFFECTIONS OF.** Muriatic acid pediluvium, 74. Terebinthinated ether, 81. Ointment tartar emetic, 108. Belladonna, 127. Blue pill and jalap, 234. Blue pill and quinine, 234. Compound mercurial pill, 234. Compound cathartic pills, 240. Calomel and jalap, 256. Tartrate of potass. and rhubarb, 338. Mixture tartrate potass., 338.
- LUMBAGO.** Camphor ointment, 140. Spirit turpentine mixture, 292.
- LUNGS, AFFECTIONS OF.** Syrup chamomile, 104. Tartar emetic, 107. Tartar emetic and opium, 107. Lotion tartar emetic, 108. Ointment tartar emetic, 108. Syrup antimonial wine, 109. Fig coffee, 214. Syrup asparagus, 116. Compound tincture benzoin, 129. Compound pills henbane, 248. Ipecacuanha and antimony, 253. Myrrh and ipecacuanha, 286. Naphthaline, 287. Tar water, 312. Solution cyanide potass., 320. Decoction and syrup seneka, 372. Storax, 389. Balsam tolu, 400. Colt's foot, 403.
- MAMMÆ, AFFECTIONS OF.** Ointment carbonate ammonia, 93. Fomentation sal-ammoniac, 95. Cataplasm sal-ammoniac, 95. Lime water and oil, 135. Ointment turpentine, 399. Ointment oxide zinc and lycopodium, 411.
- MANIA.** Compound pills columbo, 173. Pills foxglove, 193.
- MARASMUS.** Jelly orange leaves, 119.
- MEASLES.** Infusion safflower, 148. Spermaceti mixture, 155.
- MENORRHAGIA.** Pills gallic acid, 71. Pills tannic acid and opium, 76. Alum, 88. Compound bolus catechu, 152. Peruvian bark and isinglass, 161. Mixture extract ergot, 197. Infusion black purslane, 198. Pills ammoniated iron, 201. Injection sulphate iron, 211. Aromatic alum whey, 261.
- MERCURIAL DISEASE.** Mixture sulphuret potass., 328. Sulphur mixture, 394.
- MILK, TO INCREASE SECRETION OF.** Compound powder fennel, 215.
- MILK, TO RETARD SECRETION OF.** Bolus acetate potass., 329. Pills acetate soda, 379.
- MOUTH, AFFECTIONS OF.** Vinegar gargle, 67. Muriatic acid gargle, 74. Collutory nitric acid, 74. Alum, 89. Chlorine gargle, 158. Collutory balsam Peru, 285. Collutory carbonate potass., 332.
- MOLLITIES OSSIIUM.** Phosphate lime, 137.
- MUSCLES, RIGIDITY OF.** Ointment iodine and oil tobacco, 252.
- NAUSEA.** Clove cataplasm, 149. Clove bag, 149. Pennyroyal, 229. Compound infusion mint, 278. Aromatic plaster, 311. Mixture carbonate potass., 332.
- NEPHRITIS.** Compound decoction oatmeal, 123. Infusion carrot seed, 148.
- NEURALGIA.** Aconite plaster, 77, 78. Plaster belladonna, 128. Oil benzoin, 129. Emulsion cantharides, 142. Bolus castor, 151. Liniment chloroform, 159. Tincture colchicum, 170. Colchicum and camphor, 172. Creasote pills, 181. Carbonate iron, 202. Mixture tincture guaiacum, 228. Calomel and opium, 240. Compound pills henbane, 248. Lotion cherry laurel, 264. Sulphate morphia, 282. Turpentine mixture, 292. Opiated turpentine liniment, 293. Carbonate lead ointment, 316. Lotion chloride lead, 316. Lotion cyanide potass., 320. Ointment cyanide potass., 321. Solution chloride potass., 334. Extract cevadilla, 355. Extract tobacco, 396. Ointment tobacco, 396. Veratria, 407. Valerianate zinc, 411.
- NERVOUS DISORDERS.** Camphor and musk, 138. Camphor water and Hoffmann's anodyne, 139. Peruvian bark and valerian, 161. Pills ammoniated iron, 201. Infusion hops, 232. Compound powder henbane, 248. Musk, 283. Wild cherry bark, 340. Compound pills sulphate quinine, 346. Oxide zinc, 410. Cyanide zinc, 412. Ferrocyanide zinc, 412.



- NIPPLES, EXCORIATED.** Oil elaterium, 195. Glycerine, 223. Mammillary lotion, 285. Opium and lime water, 300.
- NODES, VENEREAL.** Mercurial plaster, 235.
- NUX VOMICA, POISONING BY.** Sulphuric ether and turpentine, 81.
- NYCTALOPIA.** Calomel, antimony, and henbane, 239.
- ŒDEMA.** Fomentation arnica, 113.
- ONYCHIA.** Corrosive sublimate and zinc, 237. Red oxide mercury, 244.
- OPHTHALMIA.** Alum collyrium, 88. Alum curd, 88. Acetate ammonia, 92. Antimonial wine, 108. Nitrate silver, 111. Ointment nitrate silver, 112. Chloride gold, 121. Chloride barium, 125. Sulphate cadmium, 133. Verdigris and alum, 185. Verdigris ointment, 186. Collyrium sulphate copper, 188. Decoction quince-seeds, 189. Eyebright, 199. Ointment Prussian blue, 206. Ointment red oxide and zinc, 245. Yellow root, 248. Wine opium, 301. Collyrium acetate lead, 314. Collyrium subacetate lead, 314. Ointment subacetate lead, 315. Collyrium biniodide potass., 324. Compound lotion iodide potass., 324. Collyrium caustic potass., 329. Ointment oxide zinc, 411. Ointment oxide of zinc and calomel, 411. Mixture ferrocyanide zinc, 412. Collyrium acetate zinc, 413. Collyrium sulphate zinc, 414.
- OPHTHALMIA, GONORRHOEAL.** Mercurial lotion, 234.
- OPIUM, POISONING BY.** Vinegar coffee, 70.
- ORCHITIS.** Mixture chloride barium, 125.
- OZÆNA.** Angelica, 103. Chloride lime, 134. Mercurial gargle, 233. Compound powder sulphate mercury, 245.
- PALPITATION.** Syrup asparagus, 116. Powder foxglove, 190. Wild cherry bark, 340.
- PANNUS.** Compound powder seneka, 371. Pills seneka, 371.
- PARALYSIS.** Infusion horseradish, 112. Extract arnica, 113. Assafetida, 116. Brucine, 131. Nitrate camphor, 141. Cantharides and Cayenne pepper, 142. Nux vomica, 288. Tincture nux vomica, 288-9. Pills opium and nitrate silver, 297. Phosphorated oil, 308. Lotion mustard, 378. Bath common salt and arnica, 379. Strychnine, 389. Iodate strychnine, 391.
- PERIPNEUMONIA.** Golden sulphuret antimony, 106. Powder foxglove, 190. Pills calomel and antimony, 240.
- PERIOSTITIS.** Compound mercurial ointment, 235.
- PHYMOSIS.** Solution sulphate copper, 188.
- PHTHISIS.** Mixture hydrocyanic acid, 72. Alum and benzoin, 88. Mixture chloride barium, 125. Belladonna, 127. Iodide calcium, 135. Iceland moss, 156. Electuary Peruvian bark and sulphur, 164. Pills creasote, 181. Pills sulphate copper, 187. Vinegar foxglove, 191. Ipecacuanha and nitrate potass., 253. Artificial goat's milk, 261. Locatelli's balsam, 285. Myrrh and Canada balsam, 286. Compound extract myrrh, 286. Griffith's mixture, 287. Acetic turpentine liniment, 293. Compound pills acetate lead, 314. Mixture iodide potass., 322. Sulphuret potass. and charcoal, 326. Chlorate potass., 334.
- PLEURITIS.** Mixture foxglove, 191. Cataplasm long pepper, 310.
- PNEUMONIA.** Benzoic acid and ipecacuanha, 70. Mixture foxglove, 191. Powder ipecacuanha and muriate ammonia, 253. Powder ipecacuanha and antimony, 253.
- PORRIGO.** Ointment nitric acid, 74. Lotion corrosive sublimate and copper, 238. Ointment calomel and acetate lead, 241. Ointment nitrate mercury and lead, 244. Ointment sulphate mercury, 246. Ointment oxide manganese, 173. Banyer's ointment, 317. Compound lotion sulphuret potass., 327. Powder iodide sulphur, 395.



- PRURIGO.** Fomentation opium, 299. Ointment carbonate soda, 382. Powder iodide sulphur, 395.
- PRURITUS.** Lotion borax, 380. Lotion carbonate soda, 382. Lotion chloride soda, 383.
- PSORIASIS.** Arsenic and pepper pills, 69. Ointment nitric acid, 74. Ointment subnitrate bismuth, 130. Compound powder anthracokali, 147. Decoction soot, 215. Naphthaline ointment, 287. Ointment poke, 309. Tar ointment, 312. Compound lotion sulphuret potass., 327. Lotion tobacco, 396. Ointment tobacco, 396.
- PSOROPHTHALMIA.** Ointment sulphate copper, 188. Ointment red oxide mercury, 204. Ointment red oxide mercury and cinnabar, 245. Ointment common salt, 379. Compound sulphur ointment, 394.
- PYROSIS.** Alkaline wine aloes, 85. Compound pills kino, 259. Mixture carbonate potass., 333.
- RACHITIS.** Phosphate lime, 137. Black oxide iron, 211. Phosphate magnesia, 272. Mixture cod liver oil, 290. Acorn coffee, 342. Compound powder madder, 354.
- RECTUM, DISEASES OF.** Suppository hemlock, 176. Injection copaiba, 178.
- RHEUMATISM.** Aconite, 77. Aletris, 81. Ammoniacal liniment, 91. Phosphate ammonia, 96. Golden sulphuret antimony, 106. Tartar emetic and opium, 107. Plaster tartar emetic, 108. Antimonial powder, 109. Burdock, 110. Extract arnica, 113. Ointment chloride gold, 121. Sulphuret lime, 135. Nitrate camphor, 141. Compound tincture cantharides, 142. Plaster castor, 151. Black snakeroot, 160. Colchicum, 170-1. Tincture colchicum flowers, 171. Creasote pills, 181. Compound decoction bittersweet, 193. Compound galbanum plaster, 217. Hedge hyssop, 226. Decoction guaiacum, 226. Oil guaiacum, 227. Compound powder guaiacum, 227. Guaiacum mixture, 227. Guaiacum and bitter-sweet, 227. Tincture guaiacum, 227. Plummer's pills, 240. Black sulphuret mercury, 246. Nervine balsam, 269. Magnesia and colchicum, 270. Tincture magnolia, 272. Oil of horsemint, 279. Turpentine mixture, 292. Opium and antimony, 296. Opium plaster, 298. Fomentation wine opium, 301. Opium liniment, 303. Plaster petroleum, 308. Phosphorated oil, 308. Plaster black pepper, 310. Pitch plaster, 311. Plaster subacetate lead, 315. Solution chlorate potass., 334. Plaster pellitory, 340. Tincture cevadilla, 355. Camphorated essence soap, 361. Zittman's decoction, 363. Conserve broom, 371. Pills stramonium seeds, 388. Lotion stramonium, 388. Sulphuret carbon, 395. Turpentine liniment, 399. Poison oak, 402. Veratria 407. Wine white hellebore, 408.
- RINGWORM.** Ointment cocculus indicus, 168. Vinegar borax, 380.
- SALIVATION.** Sulphuretted hydrogen, 72. Tannic acid, 76. Gargle sulphate copper, 188. Pills iodine, 251. Sulphuret potass. and cremor tartar, 326. Gargle borax, 380.
- SCARLATINA.** Mixture hyponitrous ether, 80. Acetate ammonia, 92. Plaster tartar emetic, 100. Belladonna, 127-8. Cayenne pepper gargle, 145. Chlorine mixture, 159. Myrrh gargle, 287. Mixture tartrate potass., 338. Mixture cremor tartar, 339.
- SCIATICA.** Liniment succinated ammonia, 97. Plaster sulphuret antimony, 107. Oil turpentine and honey, 292. Oil turpentine mixture, 292.
- SCHIRRHUS.** Bolus sal ammoniac, 95. Extract marigold, 137. Charcoal, 146. Hemlock, 175. Mercury and antimony, 239. Ointment mercury and belladonna, 235.
- SCROFULA.** Hydriodate ammonia, 94. Chloride barium, 125. Iodide barium, 125. Barytes, 125. Compound cataplasm bryony, 131. Solution chloride calcium, 133. Chloride lime, 134. Animal charcoal, 145. Anthracokali, 146. Hemlock, 175.



- Koechlin's drops, 186. Mixture bittersweet, 194. Compound powder metallic iron, 200. Ammoniated iron, 201. Bromide iron, 202. Compound pills carbonate iron, 203. Lozenges iodide iron, 207. Ammoniated tincture gentian, 231. Frost weed, 229. Ointment mercury and belladonna, 235. Mercury and antimony, 233. Iodide mercury, 242. Ointment iodide mercury, 243. Biniodide mercury, 243. Black sulphuret mercury, 246. Iodine bath, 251. Mixture iodine, 252. Cod liver oil, 290. Iodide lead, 317. Ointment iodide lead, 317. Solution bromide potass., 319. Ointment bromide potass., 319. Ointment iodide potass., 322. Mixture iodide potass., 322. Bath sulphuret potass., 327. Mixture sulphuret potass., 328. Conserve acetate potass., 329. Acorn coffee, 342. Hydriodate quinine, 343. Decoction madder, 354. Jauperand's decoction, 363. Compound infusion sassafras, 365. Compound ointment common salt, 379. Bath common salt and gelatine, 378. Carbonate soda and chamomile, 383. Sulphur electuary, 393.
- **SCURVY.** Wood sorrel, 67. Alumine, 87. Mixture horseradish, 112. Brook lime, 126. Scurvy grass, 169. Collutory lemon juice, 265. Mixture buckbean, 278. Chlorate potass., 334.
- SEA SICKNESS.** Plaster carbonate ammonia, 93.
- SLEEPLESSNESS.** Camphor water and Hoffmann's anodyne, 139. Tincture hops, 232. Mixture lactucarium, 262. Acetate morphia, 281. Preparations opium, 295. Syrup poppy heads, 306. Jamaica dogwood, 311.
- SNAKE BITES.** Ammonia and ether, 91. Eau de luce, 93, 392.
- SORE THROAT, VENEREAL.** Mercurial lotion, 233.
- SPASM EYELIDS.** Lotion opium and oil chamomile, 298.
- SPASMODIC DISEASES.** Sulphuric acid and ether, 76. Succinated ammonia, 96. Nitrate silver, 111. Electuary orange leaves, 119. Camphor, 138. Ethereal tincture camphor, 139. Castor, 151. Chloroform, 159. Mayweed, 181. Pills foxglove, 190. Tincture galbanum, 217. Calomel, antimony, and henbane, 239. Anti-hysterical water, 277. Musk, 283. Dippel's animal oil, 289. Opium and its preparations, 291. Succinated tincture opium, 302. Clyster opium and valerian, 303. Cyanide potass., 320. Mixture nitrate potass., 336. Confection rue, 365. Extract stramonium, 388. Purified oil amber, 391. Artificial musk, 391. Balsam amber, 392. Infusion linden, 400. Valerian, 404. Oxide zinc, 410. Cyanide zinc, 412.
- SPERMATORRHOEA.** Compound powder cascarilla, 150. Mixture acetate lead, 315.
- SPINA VENTOSA.** Bath sulphuret potass., 327.
- SPLEEN, DISEASES OF.** Solution hydrargyro-iodide potass., 325.
- SPRAINS.** Lotion sal ammoniac, 95. Liniment opium, 298. Aromatic vinegar, 353.
- STOMACH, AFFECTIONS OF.** Cayenne pepper pills, 145. Chlorine mixture, 159. Tincture chloride iron, 204. Mixture tincture guaiacum, 228. Aromatic plaster, 311. Compound powder sulphate quinine, 345. Mixture sulphuret carbon, 395.
- STRANGURY.** Clyster laudanum, 303. Infusion parsley, 308. Mixture nitrate potass., 336.
- SWEATING, PROFUSE.** Mixture nitric acid, 74. Agaric, 130. Acetate lead, 313. Mixture acetate lead, 315.
- SYPHILIS.** Mixture nitric acid, 74. Sulphuretted hydrosulphate ammonia, 95. Chloride silver, 110. Cyanide silver, 110. Gargle orpiment, 114. Preparations gold, 120-2. Compound powder anthracokali, 147. Hemlock and calomel, 175. Koechlin's drops, 186. Mercury and its preparations, 232. Opiated acetate mercury, 297. Bichloride platina, 313. Chloroplatinate soda, 313. Mixture biniodide potass., 324. Compound tincture carbonate potass., 333. Double chloride quinine and mercury, 343. Zittman's decoction, 363.
- TAPEWORM.** Chloride barium, 125. Male fern, 214. Compound powder gamboge,



219. Decoction pomegranate root, 225. Anthelmintic emulsion, 292. Mixture petroleum, 308. Compound powder cevadilla, 355. Oxide tin, 388.

**TEETH, DISEASES OF.** Alum and ether, 88. Vicat's mixture, 94. Chloride lime, 134. Cantharides plaster, 144. Oil cloves, 149. Plaster castor, 151. Chlorine oil, 159. Hemlock collyrium, 175. Solution creasote, 182. Compound oil guaiacum, 227. Mixture guaiacum, 228. Mastic paste, 275. Turpentine lotion, 292. Antiodontalgic mass, 298. Balsam and drops for toothache, 298. Odontalgic drops, 301. Pellitory and preparations, 340.

**TESTICLE, ENGORGED.** Compound mercurial ointment, 235. Ointment iodide potass., 322.

**TETANUS.** Extract hemp, 141. Mixture Dippel's animal oil, 289. Opium and cinnamon water, 300. Solution caustic potass., 128.

**THROAT, AFFECTIONS OF.** Nitrate silver, 111. Cayenne pepper gargle, 145. Gargle Peruvian bark, 165. Liniment verdigris, 186. Gargle ammoniated copper, 187. Gargle figs, 213. Gargle galls, 218. Gargle pomegranate rind, 225. Muriate manganese, 272. Gargle myrrh, 287. Gargle nitrate potass., 336. Gargle sulphate quinine, 347. Gargle sage, 359. Vinegar sage, 359. Vinegar elder flowers, 359.

**TINEA CAPITIS.** Lotion hydrocyanic acid, 72. Ammoniacal liniment, 91. Sulphuret antimony, 105. Ointment azedarach, 123. Liniment chloride lime, 134. Compound soot ointment, 216. Ointment calomel and alum, 241. Ointment poke, 309. Oil black pepper, 310. Tar ointment, 312. Ointment black pitch, 312. Ointment bromide potass., 319. Compound lotion sulphuret potass., 329. Ointment sulphuret potass., 328. Compound ointment common salt, 379. Lotion sulphuret soda, 379. Ointment carbonate soda, 382. Compound sulphur ointment, 394. Ointment sulphur and zinc, 394. Ointment sulphuret carbon, 395.

**TONSILS INFLAMED, &c.** Gargle alum, 88. Gargle iodine, 252. Ointment iodine, 252.

**TENESMUS.** Clyster, laudanum, 303. Pills rhubarb and ipecacuanha, 349.

**TRISMUS.** Musk mixture, 284.

**ULCERS.** Nitric acid, 74. Sulphate alumine, 87. Burnt alum, 87. Anglo Saxon ointment, 90. Ointment oxide silver, 111. Ointment nitrate silver, 112. Chloride lime, 134. Ointment charcoal, 146. Charcoal cataplasm, 146. Chlorine liniment, 159. Peruvian bark and camphor, 161. Cerate Peruvian bark, 166. Hemlock ointment, 175. Verdigris lotion, 185. Metz's balsam, 185. Egyptian ointment, 186. Sal ammoniated copper, 187. Ointment ammoniated copper, 187. Ointment sulphate copper, 188. Ointment foxglove, 191. Ointment oleine, 195. Ointment soot, 216. Ointment galls, 219. Compound oil guaiacum, 227. Ointment red oxide mercury, 245. Lotion henbane, 248. Infusion cherry laurel, 264. Ceromel, 276. Mezereon ointment, 279. Ointment monesia, 280. Root of mulberries, 280. Tincture balsam Peru, 285. Ointment balsam Peru, 286. Plaster balsam Peru, 286. Cod liver oil, 290. Ointment bichloride platina, 313. Lotion acetate lead, 315. Goulard's lotion, 315. Ointment subacetate lead, 315. Compound powder carbonate lead, 316. Ointment carbonate lead, 316. Caustic solution iodide potass., 324. Ointment biniodide potass., 325. Saviard's lotion, 329. Solution chloride potass., 334. Resin cerate, 348. Wine roses, 313. Ointment willow leaves, 358. Balsam sulphur, 394. Turpentine liniment, 399. Turpentine plaster, 399. Cataplasm slippery-elm, 404. Lotion chloride zinc, 411. Plaster carbonate zinc, 414.

**ULCERS VENEREAL.** Mercurial liniment, 235. Lotion corrosive sublimate, 237. Yellow wash, 238. Ointment cyanide mercury, 242. Ointment red iodide mercury, 243. Black wash, 244. Compound tincture opoponax, 304. Plaster biniodide potass. and opium, 321.

**URETHRITIS.** Injection kino, 259. Injection subacetate lead and lime water, 315. Pills nitrate potass., 325.



- URINARY ORGANS, AFFECTIONS OF. Pipsissewa, 157. Buchu, 192. Clyster laudanum, 303. Pareira brava, 306. Embrocation petroleum, 307. Extract dandelion, 398. Turpentine pills, 399. Pills turpentine and magnesia, 399. Compound uva ursi, 404.
- URINE, INCONTINENCE. Chloride gold and soda, 121. Compound pills nux vomica, 289. Powder iodide sulphur, 395.
- URINE, RETENTION. Sal ammoniac, 95.
- UTERUS, DISEASES OF. Animal charcoal, 146. Chlorine injection, 158. Syrup ergot, 196. Injection rhatany, 260. Ointment acetate morphia, 281. Mixture wine opium, 301.
- UVULA, RELAXATION OF. Gargle galls, 218. Gargle oak bark, 342. Gargle sage, 359.
- VAGINA, DISEASES OF. Injection copaiba, 178. Injection opium, 299. Suppository chloride zinc, 411.
- VERMIN, TO DESTROY. Ointment cocculus indicus, 168. Ointment fennel, 215. Ointment cevadilla, 356. Capuchin powder, 356. Vinegar stavesacre, 388. Ointment stavesacre, 388.
- VERTIGO. Bolus castor, 158.
- VOMITING, TO CHECK. Mixture catechu, 153. Compound pills colombo, 174. Mixture colombo, 174. Creasote mixture, 182. Clyster laudanum, 303. Aromatic plaster, 311. Antiemetic mixture, 382.
- VOMITING, SPASMODIC. Mercurial mixture, 234. Mixture ipecacuanha, 254.
- WARTS. Escharotic powder alum, 87. Ointment chloride antimony, 105. Verdigris ointment, 186. Verdigris plaster, 186. Corrosive sublimate and copper, 237. Savine and verdigris, 386. Savine ointment, 357.
- WEAKNESS OF BACK. Lotion Peruvian bark, 165. Iron plaster, 203. Plaster red oxide iron, 212.
- WHITE SWELLING. Ointment nitrate silver, 112. Chloride calcium, 133.
- WORMS. Tincture and wine wormwood, 64. Ointment aloes, 86. Semen contra, 115. Assafetida and iron, 117. Azedarach, 123. Wormseed, 157. Cochineal and common salt, 169. Mixture copaiba, 179. Male fern, 214. Cabbage tree bark, 222. Mercurial mucilage, 233. Calomel and scammony, 219. Calomel and jalap, 240. Black sulphuret mercury, 246. Cowhage, 284. Dippel's animal oil, 289. Mixture olive oil, 291. Anthelmintic emulsion, 291. Castor oil and ether, 292. Emulsion opoponax, 304. Savine and pinkroot, 356. Anthelmintic emulsion, 375. Pinkroot, 385. Powder tin, 387. Sulphuret tin, 387. Tansy, 397. Turpentine and jalap, 399.



# INDEX

OF

## PHARMACEUTICAL AND BOTANICAL NAMES.

- Abies canadensis, 312  
 excelsa, 311  
 picea, 311  
 Abietis resina, 311  
 Absinthium, 65  
 Acacia, 66  
 arabica, 66  
 catechu, 152  
 senegal, 66  
 vera, 66  
 Acetosella, 67  
 Acetum, 67  
 Acidum aceticum, 68  
 empyreumaticum, 69  
 arseniosum, 69  
 benzoicum, 70  
 boracicum, 70  
 citricum, 71  
 gallicum, 71  
 hydrocyanicum, 71  
 hydriodicum, 72  
 hydrosulphuricum, 72  
 lacticum, 73  
 muriaticum, 73  
 nitricum, 74  
 nitro-muriaticum, 75  
 oxalicum, 75  
 phosphoricum, 75  
 sulphuricum, 75  
 tannicum, 76  
 tartaricum, 77  
 valerianicum, 77, 406  
 Aconitina, 78  
 Aconitum, 77  
 napellus, 77  
 paniculatum, 77  
 Acorus calamus, 133  
 Adiantum, 79  
 capillus veneris, 79  
 pedatum, 79  
 Æsculus hippocastanum, 247  
 Æther aceticus, 79  
 hydrocyanicus, 80  
 hyponitrosus, 79  
 muriaticus, 80  
 sulphuricus, 80  
 terebinthinatus, 81  
 Agathotes chirayta, 158  
 Aletris, 81  
 farinosa, 81  
 Allium, 81  
 sativum, 81  
 Aloe, 82  
 socotrina, 82  
 spicata, 82  
 vulgaris, 82  
 Althæa, 86  
 officinalis, 86  
 Alumina, 86  
 Alumina acetas, 87  
 et potassæ sulphas, 87  
 sulphas, 87  
 Ammonia, 90  
 Ammonia acetas, 91  
 arsenias, 92  
 carbonas, 92  
 citras, 94  
 hydriodas, 94  
 hydrosulphas, 94  
 murias, 95  
 nitras, 96  
 phosphas, 96  
 succinas, 96  
 sulphas, 97  
 Ammoniacum, 97  
 Amygdala, 98  
 amara, 98  
 dulcis, 98  
 Amygdalus communis, 98  
 Amylum, 101  
 Amyli iodidum, 102  
 Anacylus pyrethrum, 340  
 Anamirta cocculus, 168  
 Anemone, 102  
 pratensis, 102  
 pulsatilla, 102  
 Angelica, 102  
 atropurpurea, 102  
 officinalis, 102  
 Angustura, 103  
 Anisum, 103  
 Anthemis, 104  
 nobilis, 104  
 Antimonialis pulvis, 109  
 Antimonium, 104  
 Antimonii chloridum, 105  
 et potassæ tartras, 107  
 sulphas, 107  
 sulphuretum, 105  
 Apis mellifica, 276  
 Apocynum, 109  
 androsæmifolium, 109  
 cannabinum, 109  
 Aralia nudicaulis, 110  
 spinosa, 110  
 Arctium, 110  
 lappa, 110  
 Arctostaphylos uva ursi, 404  
 Argentum, 110  
 Argenti chloridum, 110  
 cyanidum, 110  
 iodidum, 110



- Argenti oxidum, 111  
     nitras, 111  
 Aristolochia hirsuta,  
     reticulata,  
     serpentaria,  
 Armoracia, 112  
 Arnica, 112  
     montana, 112  
 Arsenicum, 113  
 Arsenici iodidum, 113  
     et hydrargyri iodidum, 114  
     sulphuretum, 114  
 Artanthe elongata, 275  
 Artemisia, 114  
     absinthium, 65  
     abrotanum, 114  
     contra, 115  
     judaica, 115  
     moxa, 114  
     santonica, 115  
 Arum, 115  
     triphyllum, 115  
 Asagraea officinalis, 355  
 Asarum, 115  
     canadense, 115  
     europaea, 115  
 Asclepias, 115  
     incarnata, 116  
     syriaca, 116  
     tuberosa, 115  
 Asparagus, 116  
     officinalis, 116  
 Assafetida, 116  
 Astragalus tragacantha, 402  
 Atropa belladonna, 126  
 Atropia, 118  
 Aurantium, 118  
 Aurantii cortex, 119  
     flores, 120  
     folia, 119  
 Aurum, 120  
     ammoniatum, 123  
     stanno paratum, 123  
 Auri chloridum, 120  
     cyanidum, 122  
     et sodii chloridum, 121  
     iodidum, 122  
     oxidum, 122  
 Avena sativa, 123  
 Azedarach, 124  
  
 Ballota lanata, 124  
 Balsamodendron myrrha, 286  
 Baptisia tinctoria, 124  
 Barium, 124  
 Barii chloridum, 124  
     iodidum, 125  
     sulphuretum, 125  
 Barosma crenata, 192  
     serratifolia, 192  
 Baryta, 125  
 Baryta acetat, 126  
 Bebeerina, 126  
 Beccabunga, 126  
 Belladonna, 126  
 Benzoinum, 128  
 Berberis, 129  
     vulgaris, 129  
 Bismuthum, 129  
 Bismuthi subnitrates, 129  
 Bistorta, 130  
 Boletus laricis, 130  
 Bromium, 130  
 Brucia, 130  
 Bruciae acetat, 131  
     murias, 131  
     sulphas, 131  
  
 Bryonia, 131  
     alba, 131  
     dioica, 131  
 Buxus, 131  
     sempervirens, 131  
  
 Cacao, 132  
 Cadmium, 133  
 Cadmii sulphas, 133  
 Calamus, 133  
 Calcis carbonas, 136  
     phosphas, 137  
 Calcii chloridum, 133  
     iodidum, 135  
     sulphuretum, 135  
 Calcium, 133  
 Calendula, 137  
     officinalis, 137  
 Calotropis, 138  
     gigantea, 138  
 Calx, 135  
     chlorinata, 134  
 Camphora, 138  
     officinarum, 138  
 Canella, 141  
     alba, 141  
 Cannabis, 141  
     sativa, 141  
 Cantharis, 141  
     vesicatoria, 141  
 Capsicum, 144  
     annuum, 144  
     baccatum, 144  
     frutescens, 144  
 Carbo animalis, 145  
     ligni, 146  
     mineralis, 146  
 Carbonis iodidum, 147  
     sulphuretum, 395  
 Cardamine, 147  
     pratensis, 147  
 Cardamomum, 147  
 Carota, 148  
 Carthamus, 148  
     tinctoria, 148  
 Carum, 148  
     carui, 148  
 Caryophyllus, 149  
     aromaticus, 149  
 Cascarilla, 150  
 Cassia acutifolia, 373  
     elongata, 373  
     fistula, 150  
     marilandica, 150  
     obovata, 373  
 Castor fiber, 151  
 Castoreum, 151  
 Catalpa, 152  
     cordifolia, 152  
 Cataria, 152  
 Catechu, 152  
 Ceanothus, 154  
     americanus, 154  
 Centaurea benedicta, 154  
 Centaurium, 154  
 Cephalis ipecacuanha, 252  
 Cerasus serotina, 340  
 Cetaceum, 154  
 Cetraria, 156  
 Chelidonium, 156  
     majus, 156  
 Chenopodium, 156  
     anthelminticum, 156  
 Chimaphila, 157  
     umbellata, 157  
 Chiretta, 158  
 Chlorinum, 158



Chloriformum, 159  
 Chondrus, 159  
     *crispus*, 159  
 Chrysophyllum, 279  
 Cichorium, 160  
     *intybus*, 160  
 Cimicifuga, 160  
     *racemosa*, 160  
 Cinchona, 160  
 Cinchonina, 166  
 Cinnamomum, 167  
     *aromaticum*, 167  
     *zeylanicum*, 167  
 Cissampelos pareira, 306  
 Citrullus colocynthis, 172  
 Citrus aurantium, 118  
     *limonum*, 264  
 Cocculus, 168  
     *palmatum*, 173  
 Coccus, 168  
 Cochlearia, 199  
     *armoracia*, 112  
     *officinalis*, 169  
 Codeia, 169  
 Coffea, 169  
     *arabica*, 169  
 Colchicum, 170  
     *autumnale*, 170  
 Collinsonia, 172  
     *canadensis*, 172  
 Colocynthis, 172  
 Colomba, 173  
 Comptonia, 174  
 Conium, 175  
     *maculatum*, 175  
 Contrayerva, 176  
 Convolvulus panduratus, 177  
     *scammonia*, 365  
 Copaiba, 177  
 Copaifera officinalis, 177  
 Coptis, 179  
     *trifolia*, 179  
 Coriandrum, 179  
     *sativum*, 179  
 Cornu, 178  
 Cornus, 180  
     *circinata*, 180  
     *florida*, 180  
     *sericea*, 180  
 Cotula, 181  
 Creasotum, 181  
 Crocus, 182  
     *sativus*, 182  
 Croton eleutheria, 150  
     *lacciferum*, 262  
     *tigium*, 293  
 Cubeba, 183  
 Cuminum cyminum, 189  
 Cunila, 185  
     *mariana*, 185  
 Cuprum, 185  
     *ammoniatum*, 186  
 Cupri acetas, 185  
     *urias*, 186  
     *sulphas*, 187  
 Cydonia, 188  
     *vulgaris*, 188  
 Cuminum, 189  
 Cytisus scoparius, 371  
 Daphne gnidium, 278  
     *mezereum*, 278  
 Datura stramonium, 388  
 Daucus carota, 148  
 Delphinium, 189  
     *consolida*, 189  
     *staphisagria*, 388

Delphinia, 189  
 Dianthus, 190  
     *caryophyllus*, 190  
 Digitalis, 190  
     *purpurea*, 190  
 Digitalina, 192  
 Diosma, 192  
 Diospyros, 193  
     *virginiana*, 193  
 Diplolepis gallæ tinctoriæ, 218  
 Dirca, 193  
     *palustris*, 193  
 Dorema ammoniacum, 97  
 Dorstenia contrayerva, 176  
 Dracontium, 193  
 Drimys winteri, 410  
 Dulcamara, 193  
 Ecbalium elaterium, 194  
 Elaterina, 195  
 Elaterium, 194  
 Elemi, 195  
 Elettaria cardamomum, 147  
 Emetina, 195  
 Ergota, 196  
 Ergotetia abortifaciens, 196  
 Erigeron, 197  
 Erigeron annuum, 197  
     *canadense*, 197  
     *heterophyllum*, 197  
     *philadelphicum*, 197  
     *strigosum*, 197  
 Eryngium, 197  
     *aquaticum*, 198  
     *maritimum*, 197  
 Erythraea centaurium, 154  
 Eugenia pimenta, 309  
 Eupatorium, 198  
     *perfoliatum*, 198  
     *purpureum*, 198  
     *teucrifolium*, 198  
 Euphorbia, 198  
     *corollata*, 198  
     *hypericifolia*, 198  
     *ipecacuanha*, 198  
     *lathyrus*, 198  
 Euphorbium, 199  
 Euphrasia, 199  
     *officinalis*, 199  
 Ferrum, 199  
     *ammoniatum*, 201  
     *sulphuretum*, 212  
 Ferri acetas, 200  
     *ammonio-tartras*, 212  
     *arsenias*, 201  
     *bromidum*, 201  
     *carbonas*, 202  
     *carburetum*, 203  
     *chloridum*, 204  
     *citras*, 205  
     *et potassæ tartras*, 212  
     *ferrocyanuretum*, 205  
     *filum*, 199  
     *gallas*, 206  
     *iodidum*, 206  
     *lactas*, 208  
     *malas*, 208  
     *nitras*, 209  
     *oxidum hydratum*, 209  
         *nigrum*, 209  
         *rubrum*, 210  
     *phosphas*, 210  
     *ramenta*, 199  
     *sulphas*, 210  
     *tannas*, 213  
     *valerianas*, 213



- Ferula assafoetida*, 116  
*tingitana*, 97  
*Ficus*, 213  
*carica*, 213  
*Filix mas*, 214  
*Fœniculum*, 215  
*vulgare*, 215  
*Frasera*, 215  
*carolinensis*, 215  
*Fraxinus ornus*, 273  
*Fuligo*, 215  
*Fuligokali*, 216  
  
*Galbanum*, 217  
*officinale*, 217  
*Galipea officinalis*, 103  
*Galium verum*, 218  
*Galla*, 218  
*Gambogia*, 219  
*Gaultheria*, 220  
*procumbens*, 220  
*Gentiana*, 220  
*lutea*, 220  
*Gentianina*, 222  
*Geranium*, 222  
*maculatum*, 222  
*Geoffroya inermis*, 222  
*Geum*, 222  
*rivale*, 222  
*urbanum*, 223  
*virginianum*, 223  
*Gillenia*, 223  
*trifoliata*, 223  
*Glycerinum*, 223  
*Glycyrrhiza*, 223  
*glabra*, 223  
*Gossypium*, 224  
*herbaceum*, 224  
*Granatum*, 225  
*Gratiola*, 226  
*aurea*, 226  
*officinalis*, 226  
*Guaiacum*, 226  
*officinale*, 226  
  
*Hæmatoxylon*, 228  
*campechianum*, 228  
*Hamamelis*, 229  
*virginiana*, 229  
*Hebradendron*, 219  
*Hedeoma*, 229  
*pulegioides*, 229  
*Helenium*, 229  
*autumnale*, 229  
*Helianthemum*, 229  
*canadense*, 229  
*Helleborus*, 229  
*niger*, 229  
*Helonias officinale*, 355  
*Hemidesmus*, 230  
*indicus*, 230  
*Hepatica*, 230  
*triloba*, 230  
*Heracleum*, 231  
*lanatum*, 231  
*Heuchera*, 231  
*americana*, 231  
*Hippocastanum*, 247  
*Hordeum*, 231  
*distichon*, 231  
*vulgare*, 231  
*Humulus*, 232  
*lupulus*, 232  
*Hydrargyrum*, 232  
*ammoniatum*, 236  
*Hydrargyri acetas*, 235  
*boras*, 236  
  
*Hydrargyri bromidum*, 236  
*chloridum corrosivum*, 237  
*mite*, 238  
*cyanuretum*, 241  
*et quiniæ chloridum*, 241  
*iodidum*, 242  
*rubrum*, 243  
*nitras*, 243  
*oxidum nigrum*, 244  
*rubrum*, 244  
*phosphas*, 245  
*sulphas*, 245  
*sulphuretum nigrum*, 246  
*rubrum*, 246  
*tartras*, 247  
*Hydrastis canadensis*, 248  
*Hyoscyamus*, 248  
*niger*, 248  
  
*Ichthyocolla*, 249  
*Indigofera*, 250  
*Indigum*, 250  
*Inula*, 250  
*helenium*, 250  
*Iodidum*, 251  
*sulphuris*, 395  
*Ipecacuanha*, 252  
*Ipomœa jalapa*, 255  
*Iris florentina*, 255  
  
*Jalapa*, 255  
*Janipha manihot*, 397  
*Juglans cinerea*, 257  
*Juniperus*, 257  
*communis*, 257  
*sabina*, 356  
*virginiana*, 258  
  
*Kino*, 258  
*Krameria*, 259  
*triandra*, 259  
  
*Lac*, 260  
*Lacca*, 262  
*Lactuca*, 262  
*elongata*, 262  
*Lactucarium*, 262  
*Lauri baccæ*, 263  
*folia*, 263  
*Lauro cerasus*, 264  
*Laurus*, 263  
*nobilis*, 263  
*sassafras*, 365  
*Lavandula*, 264  
*vera*, 264  
*Limon*, 265  
*Linum*, 267  
*usitatissimum*, 267  
*Liriodendron*, 267  
*tulipifera*, 267  
*Lobelia*, 268  
*inflata*, 268  
*Lupulina*, 268  
  
*Macis*, 269  
*Magnesia*, 269  
*Magnesiæ acetas*, 270  
*carbonas*, 270  
*citras*, 271  
*phosphas*, 272  
*sulphas*, 271  
*Magnesii sulphuretum*, 272  
*Magnesium*, 269  
*Magnolia*, 272  
*glaucæ*, 272  
*Maltum*, 272  
*Malva*, 272



- Malva sylvestris*, 272  
*Manganesii carbonas*, 272  
     *murias*, 272  
     *oxidum*, 273  
     *sulphas*, 273  
*Manganeseum*, 272  
*Manna*, 273  
*Mannita*, 274  
*Maranta*, 274  
     *arundinacea*, 274  
*Marrubium*, 274  
     *ulgare*, 274  
*Maruta cotula*, 181  
*Mastiche*, 275  
*Matico*, 275  
*Matricaria*, 276  
     *chamomilla*, 276  
*Mel*, 276  
*Melaleuca cajuputi*, 289  
*Melia azedarach*, 124  
*Melissa*, 277  
     *officinalis*, 277  
*Mentha piperita*, 277  
     *viridis*, 277  
*Menyanthes*, 278  
     *trifoliata*, 278  
*Mezereum*, 278  
*Monarda*, 279  
     *punctata*, 279  
*Monesia*, 279  
*Mora*, 280  
*Morphia*, 280  
*Morphiæ acetas*, 280  
     *bimeconas*, 281  
     *citras*, 281  
     *hydriodas*, 282  
     *murias*, 282  
     *nitras*, 282  
     *phosphas*, 282  
     *sulphas*, 282  
     *tartras*, 283  
*Morus nigra*, 280  
     *rubra*, 280  
*Moschus*, 283  
     *moschiferus*, 283  
*Mucuna*, 284  
     *pruriens*, 284  
*Myristica*, 284  
     *moschata*, 284  
*Myroxylon*, 285  
     *peruiferum*, 285  
*Myrospermum toluiferum*, 400  
*Myrrha*, 287  
  
*Naphthalina*, 287  
*Narcotina*, 288  
*Narcotinæ murias*, 288  
*Narthex*, 116  
*Nectandra radæi*, 126  
*Nepeta cataria*, 152  
*Nephrodium filix mas*, 214  
*Nicotiana tabacum*, 395  
*Nux vomica*, 288  
  
*Olea europæa*, 290  
*Oleum animale empyreumaticum*, 289  
     *cajuputi*, 289  
     *jecoris aselli*, 290  
     *morrhææ*, 290  
     *olivæ*, 290  
     *ricini*, 291  
     *terebinthinæ*, 292  
     *tiglii*, 293  
*Opium*, 295  
*Opoponax*, 304  
     *chironium*, 304  
*Origanum*, 304  
     *vulgare*, 304  
*Ornus europæa*, 273  
*Oryza*, 304  
     *sativa*, 304  
*Ovum*, 305  
*Oxalis acetosella*, 67  
  
*Papaver*, 306  
     *rhœas*, 351  
     *somniferum*, 295, 306  
*Pareira*, 306  
     *brava*, 306  
*Paullinia*, 307  
     *sorbilis*, 307  
*Petroleum*, 307  
*Petroselinum*, 308  
     *sativum*, 308  
*Phloridzina*, 308  
*Phosphorus*, 308  
*Phytolacca*, 309  
     *decandra*, 309  
*Physeter macrocephalus*, 154  
*Picræna excelsa*, 341  
*Pimenta*, 309  
*Pimpinella anisum*, 103  
*Piper*, 310  
     *angustifolium*, 275  
     *cubeba*, 183  
     *longum*, 311  
     *nigrum*, 310  
*Piperinum*, 311  
*Piscidia erythrina*, 311  
*Pistacia lentiscus*, 275  
*Pix abietis*, 311  
     *canadensis*, 312  
     *liquida*, 312  
     *nigra*, 312  
*Platina bichloridum*, 313  
*Platinum*, 313  
*Plumbi acetas*, 313  
     *carbonas*, 316  
     *chloridum*, 316  
     *iodidum*, 316  
     *oxidum semivitreum*, 317  
         *rubrum*, 318  
     *saccharas*, 318  
     *tannas*, 318  
*Plumbum*, 313  
*Podophyllin*, 318  
*Podophyllum*, 318  
     *peltatum*, 318  
*Polygala senega*, 371  
*Polygonum*, 319  
     *bistorta*, 130, 319  
*Potassa*, 328  
*Potassæ acetas*, 329  
     *arsenitis liquor*, 330  
     *arsenias*, 330  
     *bicarbonas*, 333  
     *bisulphas*, 337  
     *bitartras*, 338  
     *boras*, 330  
     *borotartras*, 331  
     *carbonas*, 331  
     *chloras*, 334  
     *citras*, 334  
     *cum sulphure*, 338  
     *et ammoniæ carbonas*, 334  
         *sulphas*, 337  
         *tartras*, 339  
     *et magnesiæ sulphas*, 337  
     *iodas*, 335  
     *nitras*, 335  
         *purificatum*, 335  
     *oxalas*, 336  
     *silicas*, 336  
     *sulphas*, 337



- Potassæ tartras, 338  
 Potassii biniodidum, 323  
     bromidum, 319  
     chloridum, 319  
     cyanuretum, 320  
     hydrargyro-iodidum, 325  
     iodidum, 321  
     sulpho-cyanuretum, 326  
     sulphuretum, 326  
 Potassium, 319  
 Potentilla tormentilla, 401  
 Prinos, 339  
     verticillatus, 339  
 Prunum, 340  
 Prunus domestica, 340  
     lauro-cerasus, 264  
     Virginiana, 340  
 Pterocarpus marsupium, 258  
 Punila granatum, 225  
 Pyrethrum, 340  
  
 Quassia, 341  
     amara, 341  
     excelsa, 341  
 Quercus, 341  
     alba, 341  
     infectoria, 218  
     pedunculata, 341  
     robur, 341  
     tinctoria, 341  
 Quinia, 342  
 Quiniæ acetat, 342  
     arsenias, 343  
     citras, 343  
     et ferri citras, 343  
         iodidum, 343  
     et hydrargyri chloridum, 343  
     ferrocyanas, 343  
     hydriodas, 343  
         iodureta, 343  
     kinas, 344  
     lactas, 344  
     murias, 344  
     nitras, 344  
     phosphas, 344  
     sulphas, 344  
     tannas, 347  
     tartras, 347  
     valerianas, 347  
  
 Resina, 347  
 Rheum, 348  
 Rhœas, 351  
 Rhus toxicodendron, 402  
 Ricinus communis, 291  
 Rosa canina, 352  
     centifolia, 352  
     gallica, 352  
 Rosmarinus, 353  
     officinalis, 353  
 Rubia, 354  
     tinctorum, 354  
 Rubus trivialis, 354  
     villosus, 354  
 Ruta, 355  
     graveolens, 355  
  
 Sabadilla, 355  
 Sabbatia, 356  
     angularis, 356  
 Sabina, 356  
 Saccharum, 357  
     officinarum, 357  
 Sagapenum, 357  
 Sago, 357  
 Salicina, 358  
  
 Salix, 358  
     alba, 358  
 Salvia, 358  
     officinalis, 358  
 Sambucus, 359  
     canadensis, 359  
     nigra, 359  
 Sanguinaria, 359  
     canadensis, 359  
 Sapo, 360  
     durus, 360  
     mollis, 360  
     vulgaris, 360  
 Sarsaparilla, 362  
 Sassafra, 365  
     officinale, 365  
 Scammonium, 365  
 Scilla, 367  
     maritima, 367  
 Scoparius, 371  
 Scrophularia nodosa, 371  
 Senega, 371  
 Senna, 373  
 Serpentaria, 376  
 Sesamum, 376  
     indicum, 376  
     orientale, 376  
 Simaruba, 377  
     amara, 377  
     officinalis, 377  
 Sinapis, 377  
     alba, 377  
     nigra, 377  
 Smilax, 362  
 Soda, 379  
 Sodæ acetat, 379  
     arsenias, 380  
     boras, 380  
     carbonas, 381  
     chloras, 383  
     chlorinatæ liquor, 383  
     et potassæ tartras, 385  
     hydrosulphas, 383  
     phosphas, 384  
     sulphas, 384  
 Sodium, 378  
 Sodii bromidum, 378  
     chloridum, 378  
     chloro-platinas, 313  
     sulphuretum, 379  
 Solanum dulcamara, 193  
 Spigelia, 385  
     marilandica, 385  
 Spiræa, 386  
     tomentosa, 386  
 Spongia, 387  
 Stanni chloridum, 387  
     oxidum, 388  
     sulphuretum, 387  
 Stannum, 387  
 Staphisagria, 388  
 Stramonium, 388  
 Strychnia, 389  
 Strychniæ acetat, 390  
     iodas, 391  
     murias, 391  
     nitras, 391  
     sulphas, 391  
 Strychnos nux vomica, 288  
 Styra, 389  
     benzoin, 128  
     officinale, 389  
 Succinum, 391  
 Sulphur, 392  
 Sulphuris carburetum, 395  
     iodidum, 395  
 Symplocarpus fœtidus, 197



Tabacum, 395  
Tamarindus, 397  
    indicus, 397  
Tanacetum, 397  
    vulgare, 397  
Tapioca, 397  
Taraxacum, 398  
    dens leonis, 398  
Terebinthina, 399  
Theobroma cacao, 132  
Tilia europæa, 400  
Tolutanum, 400  
Tormentilla, 401  
Toxicodendron, 402  
Tragacantha, 402  
Tussilago, 403  
    farfara, 403  
  
Ulmus campestris, 403  
    fulva, 403  
Uva ursi, 404  
  
Valeriana, 404  
    officinalis, 404  
Vanilla, 406  
    aromatica, 406

Veratria, 407  
Veratriæ murias, 408  
    nitræ, 408  
    sulphas, 408  
Veratrum album, 408  
    sabadilla, 355  
    viride, 409  
Veronica anagallis, 126  
    beccabunga, 126  
Viola odorata, 409  
  
Wintera, 410  
  
Zincum, 410  
Zinci acetæ, 413  
    carbonas, 413  
    chloridum, 411  
    cyanuretum, 412  
    ferro-cyanuretum, 412  
    iodidum, 412  
    oxidum, 410  
    sulphas, 414  
    valerianas, 415  
Zingiber, 415  
    officinale, 415







# GENERAL INDEX.

Abbreviations in pharmaceutical formulæ, 33

Acetate of alumina, 87

— ammonia, 91

— copper, 185

— iron, 200

— lead, 313

— mercury, 235

— morphia, 280

— potassa, 329

— quinine, 342

— soda, 279

— strychnine, 390

— zinc, 413

Acetic acid, 68

— cataplasm, 68

— clyster, 68

— diluted, 68

Acid, acetic, 68

— arsenious, 69

— benzoic, 70

— boracic, 70

— citric, 71

— gallic, 71

— hydriodic, 72

— hydrocyanic, 71

— hydrosulphuric, 72

— lactic, 73

— muriatic, 73

— nitric, 74

— nitro-muriatic, 75

— oxalic, 75

— phosphoric, 75

— polygalic, 372

— pyroligneous, 69

— sulphuric, 75

— tannic, 76

— tartaric, 77

— valerianic, 77, 406

Acids, 53

Aconite, 77

— extract, 77

— alcoholic, 77

— ammoniated, 78

— pills of, 77

— mixture, 78

— ointment, 78

— ammoniated, 78

— plaster, 77

— powder, compound, 77

— tincture, 78

— wine, compound, 78

Aconitine, 78

— lotion, 78

— ointment, 78

Adhesive plaster, 348

— Baynton's, 317

Administration of medicines, 47

Affusion, 466

Agaric, white, 130

— pills, with opium, 130

— powder, 130

Agaric, white, powder, with opium, 130

Air in convalescence, 63

Alcohol, ammoniated, 94

— aromatic, 94

— mixture, 94

— anodyne, 94

— anisated, ammoniated, 94

— lotion, 94

— mixture, 94

Alkaloids, 461

Allspice, 309

— bolus of, 309

— spirit of, 309

— tincture of, 309

— water of, 309

Almonds, 98

— bitter, 98

— butter of, 99

— clyster, 100

— cold cream, 100

— confection of, 99

— cream of, 100

— emulsion of, 99, 100

— aromatic, 100

— fixature, 101

— jelly, 417

— linctus, green, 100

— white, 101

— liniment, cosmetic, 101

— Lockstadt's pills, 101

— lotion, 99

— Goulard's, 100

— Macassar oil, 101

— milk anodyne, 99

— artificial, 99

— of roses, 99

— mixtures, 101

— oil, 100

— of bitter, 100

— paste, 99

— powder, 99

— soap, 360

— sweet, 98

— syrup of orgeat, 99

— water of bitter, 100

— diuretic, 100

Aloes, 82

— clyster, 85

— decoction, compound, 84

— electuary, 84

— elixir, Boerhaave's, 85

— Clauder's, 85

— Garus', 86

— Stoughton's, 86

— injection, 85

— mixture, 85

— ointment, 86

— pills, 82

— Anderson's, 83

— antichlorotic, 83

— aperient, 84



- Aloes, pills, Barthez's, 84  
 ————— Becker's, 84  
 ————— Chapman's, 82  
 ————— compound, 82  
 ————— Duchesne's, 83  
 ————— Frank's, 84  
 ————— Fuller's, 83  
 ————— Griffith's, 84  
 ————— Hooper's, 83  
 ————— James's, 83  
 ————— Lady Webster's, 83  
 ————— Mitchell's, 82  
 ————— Morrison's, 83  
 ————— Peter's, 83  
 ————— Pittscharf's, 84  
 ————— Speediman's, 83  
 ————— splenetic, 83  
 ————— Whytt's, 84  
 ————— with assafetida, 82  
 ————— blue mass, 84  
 ————— iron, 82  
 ————— myrrh, 82  
 ————— powder, 82  
 ————— and canella, 82  
 ————— compound, 82  
 ————— emmenagogue, 82  
 ————— suppository, 84  
 ————— tincture, 85  
 ————— and myrrh, 85  
 ————— ethereal, 85  
 ————— wine, 85  
 ————— alkaline, 85  
 ————— balsamic, 85
- Alum, 87  
 ————— bolus, 88  
 ————— burnt, 87  
 ————— collyrium, 88  
 ————— curd, 88  
 ————— draught, 90  
 ————— electuary, 88  
 ————— errhine, 88  
 ————— gargle, 88, 89  
 ————— injection, 89  
 ————— julep, 89  
 ————— liniment, 89  
 ————— lotion, 89  
 ————— ointment, 90  
 ————— Anglo-Saxon, 90  
 ————— chilblain, 90  
 ————— pile, 90  
 ————— Rust's, 90  
 ————— pills, 88  
 ————— with benzoin, 88  
 ————— powder, 87  
 ————— root, 231  
 ————— solution, 89  
 ————— odontalgic, 88  
 ————— whey, 89, 261  
 ————— aromatic, 89, 261
- Alumine, 86  
 ————— acetate, 87  
 ————— astringent, Rust's, 87  
 ————— electuary, 87  
 ————— sulphate, 87  
 ————— wash, detergent, 87
- Amber, 391  
 ————— artificial musk, 391  
 ————— emulsion, 391  
 ————— tincture, 391  
 ————— balsam, 392  
 ————— eau de luce, 392  
 ————— liniment, 392  
 ————— mixture of oil, 392  
 ————— oil, 391  
 ————— purified, 391  
 ————— powder, fumigating, 391  
 ————— tincture, 392
- Amber tincture, alkaline, 392  
 ————— ethereal, 392
- American centaury, 356  
 ————— extract, 356  
 ————— infusion, 356  
 ————— tincture, 356  
 ————— columbo, 215  
 ————— infusion, 215  
 ————— tincture, 215  
 ————— hellebore, 409  
 ————— extract, 409  
 ————— mixture, 409  
 ————— ointment, 409  
 ————— pills, 409  
 ————— tincture, 409  
 ————— wine, 409  
 ————— senna, 150  
 ————— infusion, 151
- Ammonia, 90  
 ————— acetate, 91  
 ————— cataplasm, 92  
 ————— collyrium, 92  
 ————— mixture, 92  
 ————— solution, 91  
 ————— arseniate, 92  
 ————— solution, 92  
 ————— carbonate, 92  
 ————— aromatic, 92  
 ————— draught, 93  
 ————— with ginger, 93  
 ————— drops, 93  
 ————— liniment, 93  
 ————— mixture, 93  
 ————— ointment, 93  
 ————— Gondret's, 93  
 ————— pills, 93  
 ————— plaster, 93  
 ————— potion, 92  
 ————— tincture, compound, 93  
 ————— citrate, 94  
 ————— gargle, 90  
 ————— hydriodate, 94  
 ————— ointment, 94  
 ————— hydrosulphate, 94  
 ————— liniment, 95  
 ————— sulphuretted, 95  
 ————— syrup, 95  
 ————— injection, 91  
 ————— liniment, 90  
 ————— compound, 90  
 ————— sulphuretted, 91  
 ————— terebinthinate, 91  
 ————— lotion, 91  
 ————— Granville's, 90  
 ————— mixture, 91  
 ————— muriate, 95  
 ————— bolus, 95  
 ————— cataplasm, 95  
 ————— collutory, 95  
 ————— draught, 95  
 ————— emulsion, 96  
 ————— fomentation, 95  
 ————— liniment, 95  
 ————— lotion, 95  
 ————— powder, 95  
 ————— wash, 96  
 ————— nitrate, 96  
 ————— mixture, 96  
 ————— phosphate, 96  
 ————— solution, 96  
 ————— plaster, 91  
 ————— sinapism, 91  
 ————— spirit, fetid, 91  
 ————— succinate, 96  
 ————— impure, 96  
 ————— liniment, 97  
 ————— mixture, 96, 97



- Ammonia, succinate, spirit, 96  
     — with ether, 97  
     — succinic acid, 96  
     — sulphate, 97  
     — water, 90  
 Ammoniac mixture, 97, 98  
     — compound, 98  
     — with nitric acid, 98  
     — pills, 97  
     — compound, 97  
     — De Haen's, 97  
     — Klein's, 97  
     — with rhubarb, 97  
     — plaster, 98  
     — gum, 98  
     — with hemlock, 98  
     — mercury, 98  
 Ammoniated alcohol, 94  
     — anisated, 94  
     — lotion, 94  
     — mixture, 94  
     — aromatic, 94  
     — mixture, 94  
     — Vicat's, 94  
 Ammonio-citrate of iron, 205  
     — tartrate of iron, 213  
     — solution, 213  
 Amorphous quinine, 342  
 Anemone, 102  
     — collyrium, 102  
     — extract, 102  
     — pills, 102  
     — water, 102  
 Angelica, 102  
     — conserve, 103  
     — essence, 103  
     — errhine mixture, 103  
     — tincture, 103  
     — compound, 102  
     — tree, 110  
     — water, 103  
 Angustura, 103  
     — electuary, 103  
     — infusion, 103  
     — mixture, 103  
     — tincture, 103  
 Animal charcoal, 145  
 Anise, 103  
     — lozenges, 103  
     — mixture, 104  
     — oil of, 104  
     — spirit, 103  
     — tincture, 104  
 Antacids, 53  
 Anthelmintic purgative, 239  
 Anthelmintics, 54  
 Anthracokali, 146  
     — powder, 147  
     — compound, 147  
     — sulphuretted, 147  
 Anti-dysenteric opiate, 298  
 Antilithics, 54  
 Antimonial powder, 109  
     — James's, 109  
     — pills, with calomel, 109  
     — wine, 108  
 Antimony, 104  
     — chloride, 105  
     — nitro-muriatic oxide, 105  
     — ointment, 105  
     — sulphuret, 105  
     — golden, 106  
     — powder, 106  
     — Kermes mineral, 105  
     — emulsion, 106  
     — lozenges, 106  
     — mixture, 106  
 Antimony, sulphuret, Kermesine powder, 106  
     — compound, 106  
     — lozenges, 105  
     — mixture, 106  
     — with lime, 106  
     — pills, 106  
     — plaster, 107  
     — ointment, 105  
     — pills, 105  
     — precipitated, 105  
     — sulphate, 107  
     — tartrate and potassa, 107  
 Anti-odontalgic mass, 298  
 Antispasmodics, 54  
 Apothecaries' measure, 16  
     — weight, 12  
 Apple water, 418  
 Approximative measures, 17  
 Arrowroot, 274  
     — beef tea, 274  
     — milk, 274  
     — pudding, 274  
     — vanilla, 407  
     — water, 274  
 Arseniate of ammonia, 92  
     — iron, 201  
     — potassa, 330  
     — quinine, 343  
     — soda, 380  
     — solution, 380  
 Arsenic, 113  
     — iodine, 113  
     — sesquiodide, 113  
     — ointment, 113  
     — pills, 113  
     — solution with mercury, 114  
     — draught, 114  
 Arsenical solution, 330  
 Arsenious acid, 69  
     — cerate, 69  
     — ointment, 69  
     — pills, 69  
     — with opium, 70  
     — pepper, 69  
     — powder, 69  
     — compound, 69  
     — solution, 69  
 Arsenite of potassa, solution, 330  
 Arteriotomy, 476  
 Artificial musk, 391  
     — sulphuretted water, 73  
 Asarabacca, 115  
     — powder, compound, 115  
 Asparagus, 116  
     — decoction of roots, 116  
     — extract of roots, 116  
     — of shoots, 116  
     — mixture, 116  
     — syrup of shoots, 116  
 Assafetida, 116  
     — Dewees' carminative, 118  
     — enema, 118  
     — mixture, 117  
     — compound, 118  
     — with oxymel squills, 117  
     — tolu, 117  
     — pills, 116  
     — with aloes, 118  
     — iron, 117  
     — lactucarium, 116  
     — musk, 117  
     — opium, 117  
     — plaster, 118  
     — tincture, 117  
     — alkaline, 117  
     — ammoniated, 118  
     — and castor, 118



- Assafetida tincture and soot, 118  
 Asses' milk, artificial, 420  
 Atropine, 118  
     — solution, 118  
 Avena, 222  
     — mixture, 223  
     — powder, compound, 223  
     — water, 222  
     — decoction, 222  
     — white, 223  
 Azedarach, 124  
     — decoction, 124  
     — ointment, 124  
  
 Balm, 277  
     — spirit, compound, 277  
     — water of, antihysterical, 277  
 Balsam acoustic, 285  
     — amber, 392  
     — cloves, aromatic, 149  
     — emulsion, 401  
     — Goulard's, 315  
     — inhalation, 401  
     — Locatelli's, 285  
     — lozenges, 400  
     — Metz's, 185  
     — mixture with almond emulsion, 401  
         — belladonna, 401  
         — copaiba, 401  
         — opium, 401  
         — morphia, 401  
     — nervine, 269  
     — of Peru, 285  
     — acoustic balsam, 285  
     — collutory, 285  
     — liniment, 285  
     — lotion, mammillary, 285  
     — mixture, 285  
     — ointment, 286  
     — compound, 286  
     — pills, 285  
     — plaster, 286  
     — syrup, 285  
     — tincture, 285  
     — of tolu, 400  
     — soap, camphorated acetic, 361  
     — sulphur, 395  
         — ethereal, 395  
         — terebinthinated, 395  
     — tincture, 400  
         — compound, 400  
         — with foxglove, 401  
     — toothache, 298  
     — Turlington's, 129  
     — syrup, 401  
 Bandoline, 189  
 Barberry, 129  
     — lemonade, 129  
 Barium, 124  
     — chloride, 124  
     — collyrium, 125  
     — mixture, 125  
     — pills, 125  
     — powders, 124  
     — solution, 125  
     — iodide, 125  
     — ointment, 125  
     — powder, 125  
     — sulphuret, 125  
 Barley, 231  
     — decoction, 231  
     — compound, 231  
     — with nitrate of potassa, 231  
     — meal, 231  
     — sugar, 231  
     — mixture, 420  
 Barytes, 125  
     — acetate, 126  
     — liniment, 126  
     — solution, 125  
 Basilicon ointment, 348  
 Baths, 462  
     — aromatic, 353  
     — cold, 463  
     — common salt and gelatine, 378  
     — cool, 463  
     — douche, 465  
     — hot, 463  
     — hydrosulphuretted, 72  
     — iodide of iron, 208  
     — iodine, 251  
     — ioduretted, 324  
     — local, 465  
     — medicated, 466  
     — muriatic acid, 74  
     — nitro-muriatic acid, 75  
     — shower, 464  
     — sulphuret of potassium, 327  
         — sodium, 379  
     — temperate, 463  
     — tepid, 463  
     — vapor, 465  
     — warm, 463  
         — air, 465  
 Baumé's hydrometer, 23  
     — table of agreement with  
         sp. gr., 25  
 Bay tree, 263  
 Bebeerine, 126  
     — pills of sulphate, 126  
     — solution of, 126  
 Beef essence, 418  
     — tea, 418  
     — arrowroot, 274  
 Beer, ginger, 415, 420  
     — molasses, 420  
     — Peruvian bark, 166  
     — pipsissewa, 158  
     — sarsaparilla, 365  
     — spruce, 420  
 Belladonna, 126  
     — clyster, 128  
     — extract, 127  
     — alcoholic, 127  
     — solution of, 127  
     — fumigation, 127  
     — liniment, 128  
     — mixture, 128  
     — ointment, 128  
     — opiated, 129  
     — pills, compound, 127  
     — plaster, 128  
     — powder, 126  
     — and rhubarb, 129  
     — syrup, 127  
     — Gillet's, 127  
     — tincture, 127  
     — ethereal, 128  
 Benne, 376  
     — infusion, 877  
 Benzoic acid, 70  
     — and copaiba mixture, 70  
     — and ipecacuanha powder, 70  
     — mixture, 70  
 Benzoin, 128  
     — balsam, Turlington's, 127  
     — emulsion, 129  
     — lard, benzoated, 128  
     — lotion, 129  
     — milk of roses, 129  
     — oil, 129  
     — pastiles, fumigating, 128  
     — powder, 128  
     — fumigating, 128



- Benzoin tincture, 129  
     — compound, 129  
     — wash, 129  
 Bibromide of mercury, 236  
 Bicarbonate of potassa, 333  
     — of soda, 381  
 Biniodide of potassium, 323  
 Binoxalate of potassa, 336  
 Biscuit jelly, 417  
 Bismuth, 129  
     — subnitrate, 129  
     — ointment, 130  
     — pills, 130  
     — powder, compound, 129  
 Bistort, 130, 319  
     — clyster, astringent, 130  
     — extract, 319  
     — infusion, 130  
     — mixture, 319  
 Bisulphate of potassa, 337  
     — soda, 385  
 Bitartrate of potassa, 338  
 Bitter almonds,  
 Bittersweet, 193  
     — decoction, 193  
     — compound, 193  
     — extract, 193  
     — pills, 194  
     — mixture, 194  
     — syrup, 194  
 Black alder, 339  
     — decoction, 339  
     — drop, 300  
     — Guy's Hospital, 300  
     — Houlton's, 300  
     — Lancaster, 300  
     — Porter's, 300  
     — Rousseau's, 300  
     — hellebore, 229  
     — extract, 230  
     — ointment, 230  
     — pills, compound, 229  
     — tincture, 230  
     — mixture, 230  
     — vino-alcoholic, 230  
     — pills, 230  
     — wine, compound, 230  
     — oak, 341  
     — oxide of iron, 209  
     — mercury, 244  
     — pepper, 310  
     — cataplasm, 310  
     — confection, 310  
     — electuary, 310  
     — oil, resinous, 310  
     — volatile, 310  
     — ointment, 310  
     — plaster, 310  
     — Ward's paste, 310  
     — pitch, 312  
     — ointment, 312  
     — plaster, 313  
     — purslane, 198  
     — infusion, 198  
     — snakeroot, 160  
     — decoction, 160  
     — tincture, 160  
     — sulphuret of mercury, 246  
 Blackberry, 354  
     — decoction, 354  
     — syrup, 355  
 Blancmange isinglass, 420  
     — rice, 420  
 Blessed thistle, 154  
     — infusion, 154  
     — wine, 154  
 Blistering court plaster, 144  
 Blisters, 38, 470  
 Blood-letting, 474  
     — general, 474  
     — topical, 474  
 Bloodroot, 359  
     — infusion, 360  
     — pills, 360  
     — tincture, 360  
 Blue pills, 284  
     — and colocynth, 284  
     — jalap, 284  
     — quinine, 284  
     — rhubarb, 284  
 Boiled flour, 418  
 Bolus, 447  
     — acetate of morphia, 282  
     — potassa, 329  
     — allspice, 309  
     — alum, 88  
     — ammoniated iron, 201  
     — burnt sponge, 387  
     — castor, 151  
     — catechu, 152  
     — chloride of gold, 121  
     — cinchonia, 166  
     — copaiba, 177  
     — croton oil, 293  
     — foxglove, 190  
     — ipecacuanha, 253  
     — iron, metallic, 200  
     — male fern, 214  
     — musk, 283  
     — narcotine, 288  
     — oxide of manganese, 273  
     — Peruvian bark, extract of, 162  
     — pomegranate root, 226  
     — sal ammoniac, 95  
     — soap, 361  
     — sulphate of morphia, 282  
     — sulphuret of potassium, 326  
     — tartar emetic, 107  
     — valerian and iron, 405  
     — sulphate of potassa, 405  
 Boneset, 198  
     — decoction, 198  
     — infusion, 198  
     — compound, 198  
 Boracic acid, 70  
     — collutory, 70  
     — mixture, 70  
 Borate of soda, 380  
 Borax, 380  
     — collutory, 380  
     — collyrium, 380  
     — gargle, 380  
     — honey, 380  
     — lotion, 380  
     — lozenges, 380  
     — mixture, 381  
     — ointment, 381  
     — powder, compound, 380  
     — vinegar, 380  
 Boullay's filter, 452  
 Bowman's root, 198  
 Box, 131  
     — oil, 131  
 Bran tea, 419  
 Bread jelly, 417  
 British oil, 307  
 Bromide of iron, 201  
     — mercury, 236  
     — potassium, 319  
     — sodium, 378  
 Bromine, 130  
     — solution, 130  
     — alcoholic, 130  
 Brooklime, 126



- Brooklime, syrup, 126  
     — water, 126  
 Broom, 371  
     — conserve, 371  
     — decoction, 371  
     — compound, 371  
     — extract, 371  
     — infusion, 371  
 Broth, mutton, 417  
     — vegetable, 417  
 Brown mixture, 224  
 Brucine, 130  
     — acetate, 131  
     — muriate, 131  
     — pills, 131  
     — solution, 131  
     — sulphate, 131  
     — tincture, 131  
 Bryony, 131  
     — cataplasm, 131  
     — compound, 131  
     — wine, 131  
 Buchu, 192  
     — extract, 192  
     — fluid, 192  
     — infusion, 192  
     — compound, 192  
     — tincture, 192  
 Buckbean, 278  
     — extract, 278  
     — infusion, 278  
     — mixture, 278  
     — diuretic, 278  
     — pills, compound, 278  
 Burdock, 110  
     — decoction, 110  
     — infusion, 110  
 Burgundy pitch, 311  
 Burnt alum, 87  
     — hartshorn, 137, 180  
     — sponge,  
     — sugar, 357  
 Butter of almonds, 99  
     — cocoa, 132  
     — mixture, 132  
     — ointment, 132  
 Butternut, 257  
     — extract, 257  
 Button snakeroot, 257  
  
 Cabbage tree bark, 222  
     — decoction, 222  
     — extract, 222  
     — mixture, 222  
 Cadmium, 133  
     — sulphate, 133  
     — ointment, 133  
     — solution, 133  
 Cajeput oil, 289  
     — liniment, 289  
     — mixture, 289  
 Cakes, soda, 421  
 Calamine, 413  
 Calamus, 133  
     — electuary, 133  
     — infusion, 133  
     — compound, 133  
     — tincture, 133  
     — compound, 133  
 Calcium, 133  
     — chloride, 133  
     — cataplasm, 133  
     — mixture, 134  
     — pills, 134  
     — solution, 133  
     — iodide, 135  
     — pills, 135  
 Calcium, sulphuret, 135  
     — liniment, 135  
     — pills, compound, 135  
 Calomel, 238  
     — collyrium, dry, 239  
     — electuary with, 241  
     — pills, 239  
     — and acetate of lead, 239  
     — antimony, 240  
     — catechu, 230  
     — colocynth, 240  
     — dandelion, 240  
     — guaiacum, 240  
     — jalap, 240  
     — iron, 240  
     — opium, 240  
     — quinine, 239  
     — squill, 240  
     — cathartic, compound, 240  
     — compound, 240  
     — ointment, 241  
     — with acetate of copper, 241  
     — alum, 241  
     — camphor, 241  
     — squill, 241  
     — powder with antimony, 238  
     — and henbane, 239  
     — anthelmintic, 239  
     — with foxglove, 239  
     — gamboge, 239  
     — jalap, 239  
     — and rhubarb, 239  
     — nitrate of potassa, 239  
     — opium, 239  
     — pink root, 239,  
  
 Calves' foot jelly, 418  
 Camphor, 138  
     — and chloroform, 140  
     — ether, 140  
     — Hoffmann's anodyne, 139  
     — laudanum, 139  
     — milk, 139  
     — nitric acid, 139  
     — clyster, 140  
     — collyrium, 141  
     — cough mixture, 141  
     — essence, Ward's, 140  
     — emulsion, 139  
     — infusion, 139  
     — liniment, 140  
     — and turpentine, 141  
     — vinegar, 140  
     — compound, 140  
     — ethereal, 140  
     — mixture, 139  
     — nitrate, 141  
     — ointment, 140  
     — pills, 138  
     — and lactucarium, 138  
     — musk, 138  
     — compound, 138  
     — powder, 138  
     — compound, 138  
     — tincture, 139  
     — and saffron, 139  
     — ethereal, 139  
     — mixture, 140  
     — water, 139  
 Canada extract, 197  
     — fleabane, 197  
     — infusion, 197  
     — pitch, 312  
 Canella, 141  
 Cantharides, 141  
     — court-plaster, blistering, 144  
     — decoction, 143



- Cantharides, emulsion, 142  
     — extract, 142  
     —      ethereal, 143  
     — infusion, 142  
     — liniment, 143  
     —      and camphor, 143  
     —      soap, 143  
     —      stimulating, 144  
     — oil, 143  
     — ointment, 143  
     —      issue, 144  
     — pills with capsicum, 142  
     —      iron, 142  
     — plaster, 144  
     —      camphorated, 144  
     —      compound, 144  
     —      odontalgic, 144  
     — pomatum, 144  
     — powder with camphor, 142  
     —      savine, 142  
     — shampoo liquid, 144  
     — tincture, 142  
     —      camphorated, 142  
     —      ethereal, 143  
     —      camphorated, 143  
     —      with guaiacum, 142  
     — vinegar, 142  
 Caramel, 357  
 Caraway, 148  
     — embrocation, 149  
     — infusion, 148  
     — oil, 149  
     — spirit, 149  
     — water, 148  
 Carbon, iodide, 147  
     — ointment, 147  
     — sulphuret of, 395  
     — drops, 395  
     — liniment, 395  
     — mixture, 395  
 Carbonate of ammonia, 92  
     — iron, 202  
     — lead, 316  
     — lime, 136  
     — magnesia, 270  
     — manganese, 272  
     — potassa, 331  
     — soda, 381  
     — zinc, 413  
 Carburet of iron, 203  
     — sulphur, 395  
 Cardamom, 147  
     — confection, 147  
     — extract, ethereal, 148  
     — powder, aromatic, 147  
     — syrup, 148  
     — tincture, 147  
     — compound, 148  
 Carminative, Dalby's, 270  
     — Dewees', 118, 270  
 Carrageen, 159  
     — decoction, 159  
     — jelly, 160  
     — mixture, 160  
 Carrot, 148  
     — cataplasm, 148  
     — extract, 148  
     — infusion of seeds, 148  
     — ointment, 148  
 Casearilla, 150  
     — extract, 150  
     —      mixture, 150  
     — infusion, 150  
     —      alkaline, 150  
     — powder, compound, 150  
     — tincture, 150  
     — wine, compound, 150  
 Cassia, purging, 150  
     — confection, 150  
     — mixture, 150  
     — pulp, 150  
 Castanillo's powders, 417  
 Castor, 151  
     — lotus, 151  
     — oil, 291  
     — clyster, 292  
     — emulsion of, 291  
     —      seeds, 291  
     — fixture, 292  
     — mixture, 291  
     —      with ether, 292  
     — pills, 151  
     — plaster, compound, 151  
     — powder, compound, 151  
     — spirit, compound, 151  
     — tincture, 151  
     — ammoniated, 151  
     — compound, 151  
     — ethereal, 151  
     — with succinic acid, 151  
 Catalpa, 152  
     — decoction, 152  
 Cataplasms, 468  
     — acetate of lead, 315  
     — acetic acid, 68  
     — black pepper, 310  
     — bryony, 131  
     —      compound, 131  
     — carrot-root, 148  
     — Cayenne pepper, 145  
     — charcoal, 146  
     — chloride of calcium, 133  
     — clove, 149  
     — emollient, 267  
     — flaxseed, 267  
     — foxglove, 191  
     — garlic, 81  
     — hemlock, 176  
     — hops, 232  
     — horseradish, 112  
     — iodine, 252  
     — lead, 317  
     — long pepper, 310  
     — mayweed, 181  
     — mercurial, 235  
     — mustard, 377  
     —      compound, 377  
     — oak bark, 342  
     — oatmeal, 124  
     — Peruvian bark, 165  
     — powder for, 124  
     — pyroligneous acid, 69  
     — sal ammoniac, 95  
     — slippery elm, 404  
     — soap, 362  
     — soot, 216  
     — subacetate of lead, 314  
     — tobacco, 396  
     — vinegar, 67  
 Catechu, 152  
     — bolus, 152  
     — compound, 152  
     — collutory, 153  
     — electuary, 153  
     — infusion, 153  
     —      compound, 153  
     — injection, 153  
     — lozenges, 153  
     — mixture, 153  
     —      anti-emetic, 153  
     —      with logwood, 153  
     — pastilles, aromatic, 153  
     — pills with alum, 152  
     — liquorice, 152



- Catechu powder, compound, 152  
     tincture, 153  
 Cathartics, 55  
 Catnep, 152  
     infusion, 152  
 Caudle, 420  
 Caustic chloride of gold, 120  
     potassa, 328  
     soda, 379  
 Caustics, 60  
 Cayenne pepper, 144  
     cataplasm, 145  
     gargle, 145  
     infusion, 145  
     lotion, 145  
     lozenges, 145  
     pills, 145  
     tincture, 145  
     with cantharides, 145  
     vinegar, 145  
 Celandine, 156  
     extract, 157  
     mixture, 157  
     pills, 157  
 Cement, diamond, 250  
     parolic, 261  
     shell lac, 262  
 Centaury, 154  
     American, 154  
     extract, 154  
     mixture, 154  
     pills, 154  
     Portland powder, 154  
     wine, compound, 154  
 Cerate, 471  
     acetate of lead, 314  
     arsenical, 69  
     carbonate of zinc, 413  
     cherry laurel, 264  
     croton oil, 295  
     eggs, 305  
     lead, 317  
     mercurial, 235  
     compound, 235  
     opium, 299  
     oxide of zinc and lycopodium, 411  
     Peruvian bark, 165  
     phosphorated, 309  
     pitch, 312  
     red sulphuret of mercury, 247  
     resin, 348  
     compound, 348  
     savine, 357  
     spermaceiti, 155  
     subacetate of lead, 315  
     sulphur, 394  
 Ceromel, 276  
 Cetrarine, 156  
     powder, 156  
 Cevadilla, 355  
     Capuchin, 356  
     extract, 355  
     ointment, 356  
     powder, compound, 355  
     tincture, 355  
 Chalk, 136  
     mixture, 136  
     prepared, 136  
 Chalybeate water, artificial, 203  
 Chamomile, 104  
     decoction, compound, 104  
     extract, 104  
     pills, compound, 104  
     German, 276  
     infusion, 104  
     mixture, 104  
 Chamomile syrup, 104  
 Charcoal, animal, 145  
     ointment, 146  
     powder, 145  
     purified, 145  
     cataplasm, 146  
     electuary, 146  
     lozenges, 146  
     mineral, 146  
     ointment, 146  
     powder, dentifrice, 146  
     and quassia, 146  
     suppository, 146  
     wood, 146  
 Cheltenham salts, 271  
 Cherry cerate, 264  
     infusion, 264  
     laurel, 264  
     ointment, 264  
     water, 264  
     lotion, 264  
     mixture, 264  
 Chicken jelly, 419  
     panada, 419  
     water, 418  
 Chinoidine, 342  
 Chiretta, 158  
     infusion, 158  
     tincture, 158  
 Chlorate of potassa, 334  
     soda, 383  
 Chloride of barium, 124  
     calcium, 133  
     gold, 120  
     and sodium, 121  
     iron, 204  
     lead, 316  
     lime, 134  
     potassium, 319  
     silver, 110  
     soda, 383  
     sodium, 378  
     zinc, 411  
 Chlorinated lime, 134  
     soda, 383  
 Chlorine, 158  
     clyster, 159  
     collutory, 158  
     gargle, 158  
     injection, 158  
     liniment, 159  
     mixture, 159  
     oil, 159  
     ointment, 159  
     water, 158  
 Chloroform, 159  
     emulsion, 159  
     liniment, 159  
 Chloroplatinate of sodium, 313  
     injection of, 313  
 Chocolate, 420  
     aromatic, 132  
     cream, 132  
     Iceland moss, 156  
     milk, 420  
     paullinia, 307  
     purgative, 367  
     salep, 132  
     vanilla, 132  
     white, 132  
 Cinchonia, 166  
     boluses, 167  
     kinate, 166  
     muriate, 166  
     pills, 166  
     sulphate, 166  
     pills, 167



Cinchonia, sulphate, wine, 167

— syrup, 167

— tincture, 167

— wine, 167

Cinnamon, 167

— electuary, compound, 167

— infusion, 167

— lozenges, 167

— oil, 168

— plaster, spiced, 168

— powder, compound, 167

— tincture, 167

— compound, 168

— ethereal, 168

— vinegar, compound, 167

— water, 168

— compound, 168

— wine, compound, 167

Citrate of ammonia, 95

— iron, 205

— and quinia, 205, 343

— magnesia, 271

— morphia, 281

— potassa, 334

— quinine, 343

Citric acid, 71

— lozenges, 71

— syrup, 71

Cleanliness in sick room, 46

Clovés, 149

— bag, 149

— balsam, 149

— cataplasm, 149

— infusion, 149

— mixture, odontalgic, 149

— oil, 149

— plaster, 149

— spirit, 149

— tincture, 149

— water, 149

— wine, 149

Clysters, 55

— acetate of morphia, 281

— acetic acid, 68

— anthelmintic, of aloes, 85

— assafetida, 118

— belladonna, 128

— bistort, 130

— camphor, 140

— castor oil, 292

— chlorine, 159

— colocynth, 173

— common salt, 378

— and arnica, 379

— copaiba, 179

— cubebs, 184

— dandelion, 399

— egg, 305

— emollient, oil of almonds, 100

— malt, 272

— ergot, 197

— flaxseed oil, 267

— iodine, 252

— laudanum, 303

— and valerian, 303

— musk, 284

— olive oil, 291

— opium, 299

— Peruvian bark, 165

— rhatany, 260

— soap, 362

— southernwood, 114

— stimulating, with nutmeg, 284

— sulphate of magnesia, 271

— quinine, 347

— soda, 385

— tartar emetic, 108

Clysters, tartarized soda, 385

— turpentine, 399

— oil, 293

— wormwood, 65

Cocculus indicus, 168

— ointment, 168

— picrotoxin, 168

Cochineal, 168

— tincture, 169

— with carbonate of potassa, 169

— common salt, 169

Cocoa, 132

— butter, 132

— ointment, 132

— mixture, 132

— chocolates, 132

— aromatic, 132

— cream, 132

— salep, 132

— vanilla, 132

— white, 132

— powders, compound, 132

— palamond, 132

— racahout, 132

— wikana, 132

Codeia, 169

— and muriate of morphia, 169

— syrup, 169

Cod liver oil, 290

— compound, 290

— liniment, 290

— mixture, 290

— ointment, 290

— compound, 290

— syrup, 290

Coffee, 169

— acorn, 342

— decoction, 170

— figs, 214

— milk, 420

— syrup, 170

— vinegar, 170

Colchicum, 170

— extract bulb, 170

— acetic, 170

— liniment, with camphor, 172

— mixture, 171

— with ammonia, 172

— elaterium, 171

— magnesia, 171

— squill, 172

— sulphate magnesia, 171

— oxymel, 171

— pills, 170

— compound, 170

— syrup, 171

— tincture,

— flowers, 171

— seeds, 170

— and foxglove, 171

— compound, 171

— mixture, 171

— vinegar, root, 171

— seeds, 171

— mixture, 172

— wine, compound, 171

— mixture, 172

— root, 170

— seeds, 170

Cold bath, 463

— cream, 100, 155

— without spermaceti, 155

— custard, 419

Collodion, 225

Collutory, balsam of Peru, 285

— boracic acid, 70

— borax, 380



- Collutory, carbonate of potassa, 332  
 ----- catechu, 153  
 ----- chlorinated lime, 134  
 ----- chlorine, 158  
 ----- creasote, 182  
 ----- hemlock, 175  
 ----- lemon juice, 265  
 ----- mastic, aromatic, 275  
 ----- myrrh, 286  
 ----- nitric acid, 74  
 ----- pyroligneous acid, 69  
 ----- rhatany, 260  
 ----- rose water, 352  
 ----- sal ammoniac, 95  
 ----- soot, 216  
 Collyrium, acetate of lead, 314  
 ----- zinc, 413  
 ----- alum, 88  
 ----- anemone, 102  
 ----- anodyne, 299  
 ----- antimonial wine, 108  
 ----- biniodide of potassium, 324  
 ----- borax, 380  
 ----- calomel, dry, 239  
 ----- camphor, 141  
 ----- carbonate of potassa, 332  
 ----- caustic potassa, 329  
 ----- chloride of barium, 125  
 ----- gold, 121  
 ----- corrosive sublimate, 238  
 ----- iodide of zinc, 413  
 ----- nitrate of silver, 111  
 ----- saffron, 183  
 ----- strychnine, 390  
 ----- subacetate of lead, 314  
 ----- sulphate of copper, 188  
 ----- zinc, 414  
 ----- and camphor, 414  
 ----- wine of opium, 301  
 Colocynth, 172  
 ----- clyster, 173  
 ----- extract, 172  
 ----- compound, 172  
 ----- mixture, 173  
 ----- pills, and henbane, 173  
 ----- compound, 173  
 ----- powder, 172  
 ----- tincture, 173  
 ----- mixture, 173  
 Columbo, 173  
 ----- American, 215  
 ----- decoction, compound, 174  
 ----- extract, 174  
 ----- infusion, 174  
 ----- and ginger, 174  
 ----- rhubarb, 174  
 ----- mixture, 174  
 ----- and cascarrilla, 174  
 ----- salep, 174  
 ----- pills, compound, 174  
 ----- powder and iron, 173  
 ----- ginger, 174  
 ----- magnesia, 173  
 ----- tartrate of iron, 174  
 ----- tincture, 174  
 Coltsfoot, 403  
 ----- decoction, 403  
 ----- compound, 403  
 ----- syrup, 493  
 Common salt, 378  
 ----- bath, with gelatine, 378  
 ----- clyster, 378  
 ----- with arnica, 379  
 ----- fomentation, 378  
 ----- mixture with lemon juice, 379  
 ----- ointment, 379  
 ----- compound, 379  
 Common salt powder, compound, 378  
 ----- with cochineal, 378  
 Confections, 449  
 Confection acorns, 342  
 ----- almonds, 99  
 ----- aromatic, 147  
 ----- black pepper, 310  
 ----- cassia, 150  
 ----- copaiba, 178  
 ----- dog rose, 352  
 ----- ipecacuanha, 254  
 ----- orange flowers, 120  
 ----- peel, 119  
 ----- roses, 353  
 ----- scammony, 366  
 ----- senna, 373  
 ----- compound, 373  
 Conserves, 449  
 Conserve of acetate of potassa, 329  
 ----- angelica, 103  
 ----- broom, 371  
 ----- violets, 409  
 Contrayerva, 176  
 ----- decoction, 176  
 ----- extract, 176  
 ----- gargle, 177  
 ----- mixture, 177  
 ----- powder, compound, 176  
 ----- tincture, 177  
 ----- compound, 177  
 Copaiba, 177  
 ----- boluses, 177  
 ----- clyster, 179  
 ----- confection, 178  
 ----- emulsion, 178  
 ----- injection, 178  
 ----- mixture, 178  
 ----- oil, 179  
 ----- paste, 179  
 ----- pills, 177  
 ----- and cubebs, 177  
 ----- cubebs and turpentine, 178  
 ----- tincture, 179  
 ----- alkaline, 179  
 ----- compound, 179  
 Copper, 185  
 ----- acetate, 185  
 ----- ammoniated, 186  
 ----- gargle, 187  
 ----- injection, 187  
 ----- ointment, 187  
 ----- pills, 187  
 ----- compound, 187  
 ----- powder, with belladonna, 187  
 ----- solution, 187  
 ----- muriate, 186  
 ----- and ammonia, 186  
 ----- cupreous ether, 186  
 ----- Koechlin's drops, 186  
 ----- solution, with mercury, 186  
 ----- subacetate, 185  
 ----- balsam, Metz's, 185  
 ----- liniment, 186  
 ----- lotion, compound, 185  
 ----- ointment, 186  
 ----- Egyptian, 186  
 ----- pills, 185  
 ----- with alum, 185  
 ----- plaster, 186  
 ----- powder, with savine, 185  
 ----- wash, ophthalmic, 185  
 ----- sulphate, 187  
 ----- collyrium, 188  
 ----- electuary, 188  
 ----- gargle, 188  
 ----- lotions, 188



- Copper, sulphate, ointment, 188  
     \_\_\_\_\_ pills, 189  
     \_\_\_\_\_ and opium, 187  
     \_\_\_\_\_ compound, 188  
     \_\_\_\_\_ powder, compound, 187  
     \_\_\_\_\_ styptic, 188  
 Cordial, Godfrey's, 302  
     \_\_\_\_\_ Warner's, 351  
 Coriander, 179  
     \_\_\_\_\_ powder, compound, 180  
     \_\_\_\_\_ tincture, 180  
     \_\_\_\_\_ water, compound, 180  
 Corrosive sublimate, 237  
     \_\_\_\_\_ collyrium, 238  
     \_\_\_\_\_ injection, 238  
     \_\_\_\_\_ lotion, 237  
     \_\_\_\_\_ Bateman's, 238  
     \_\_\_\_\_ cosmetic, 237  
     \_\_\_\_\_ with camphor, 238  
     \_\_\_\_\_ copper, 238  
     \_\_\_\_\_ mixture, 237  
     \_\_\_\_\_ pills, 237  
     \_\_\_\_\_ compound, 237  
     \_\_\_\_\_ powder with copper, 237  
     \_\_\_\_\_ zinc, 237  
     \_\_\_\_\_ solution, 237  
     \_\_\_\_\_ alcoholic, 237  
     \_\_\_\_\_ tincture, antacid, 238  
     \_\_\_\_\_ wash, cosmetic, 238  
     \_\_\_\_\_ yellow, 238  
 Cotton, 224  
     \_\_\_\_\_ decoction of root, 224  
     \_\_\_\_\_ solution, ethereal, 225  
 Court plaster, 249  
 Cowhage, 284  
     \_\_\_\_\_ electuary, 284  
 Cranesbill, 222  
     \_\_\_\_\_ decoction, 222  
     \_\_\_\_\_ extract, 222  
 Cream, chocolate, 132  
     \_\_\_\_\_ cold, 100, 155  
     \_\_\_\_\_ without spermaceti, 155  
     \_\_\_\_\_ of tartar, 338  
 Creasote, 181  
     \_\_\_\_\_ collutory, 182  
     \_\_\_\_\_ lotion, 182  
     \_\_\_\_\_ mixture, 181, 182  
     \_\_\_\_\_ ointment, 182  
     \_\_\_\_\_ compound, 182  
     \_\_\_\_\_ pills, 181  
     \_\_\_\_\_ solution, alcoholic, 182  
 Cremor tartar, 338  
 Croton oil, 293  
     \_\_\_\_\_ bolus, 293  
     \_\_\_\_\_ cerate, 295  
     \_\_\_\_\_ embrocation, 295  
     \_\_\_\_\_ emulsion, 294  
     \_\_\_\_\_ liniment, 294  
     \_\_\_\_\_ lozenges, 294  
     \_\_\_\_\_ mixture, 294  
     \_\_\_\_\_ ointment, 295  
     \_\_\_\_\_ pills, 293  
     \_\_\_\_\_ compound, 294  
     \_\_\_\_\_ with blue mass, 294  
     \_\_\_\_\_ quinine, 294  
     \_\_\_\_\_ plaster, 295  
     \_\_\_\_\_ soap, 294  
     \_\_\_\_\_ solution, saponaceous, 294  
     \_\_\_\_\_ tincture, 294  
 Cubebs, 183  
     \_\_\_\_\_ clyster, 184  
     \_\_\_\_\_ electuary, 184  
     \_\_\_\_\_ extract, alcoholic ethereal, 184  
     \_\_\_\_\_ emulsion, 184  
     \_\_\_\_\_ ethereal, 184  
     \_\_\_\_\_ lozenges, 184  
     \_\_\_\_\_ lozenges, alcoholic ethereal extract, 184  
 Cubebs, mixture, 184  
     \_\_\_\_\_ oil and copaiba, 185  
     \_\_\_\_\_ oil, 185  
     \_\_\_\_\_ powder, 183  
     \_\_\_\_\_ and alum, 183  
     \_\_\_\_\_ ergot, 183  
     \_\_\_\_\_ hemlock, 183  
     \_\_\_\_\_ syrup, alco-ether. extract, 184  
     \_\_\_\_\_ tincture, 184  
 Cuckoo flower, 147  
 Cumin, 189  
     \_\_\_\_\_ plaster, 189  
 Cyanide of gold, 122  
     \_\_\_\_\_ mercury, 241  
     \_\_\_\_\_ potassa, 320  
     \_\_\_\_\_ silver, 110  
     \_\_\_\_\_ zinc, 412  
 Cyanuret of gold, 122  
     \_\_\_\_\_ mercury, 241  
     \_\_\_\_\_ potassium, 320  
     \_\_\_\_\_ silver, 110  
     \_\_\_\_\_ zinc, 412  
 Cupping, 476  
 Custard, cold, 419  
     \_\_\_\_\_ rice, 420  
 Dalby's carminative, 270  
 Dandelion, 398  
     \_\_\_\_\_ clyster, 399  
     \_\_\_\_\_ decoction, 398  
     \_\_\_\_\_ extract, 398  
     \_\_\_\_\_ fluid, 398  
     \_\_\_\_\_ pills, 398  
     \_\_\_\_\_ with blue mass, 398  
     \_\_\_\_\_ infusion, compound, 398  
     \_\_\_\_\_ mixture, 398  
 Decoction, 451  
     \_\_\_\_\_ aloes, compound, 84  
     \_\_\_\_\_ arnica, 113  
     \_\_\_\_\_ asparagus, 116  
     \_\_\_\_\_ azedarach, 124  
     \_\_\_\_\_ barley, 231  
     \_\_\_\_\_ compound, 231  
     \_\_\_\_\_ bittersweet, 193  
     \_\_\_\_\_ compound, 193  
     \_\_\_\_\_ black alder, 339  
     \_\_\_\_\_ snakeroot, 160  
     \_\_\_\_\_ blackberry root, 355  
     \_\_\_\_\_ boneset, 198  
     \_\_\_\_\_ broom, 371  
     \_\_\_\_\_ compound, 371  
     \_\_\_\_\_ burdock, 110  
     \_\_\_\_\_ cabbage tree bark, 222  
     \_\_\_\_\_ cantharides in turpentine, 143  
     \_\_\_\_\_ carrageen, 159  
     \_\_\_\_\_ catalpa, 152  
     \_\_\_\_\_ chamomile, compound, 104  
     \_\_\_\_\_ coffee, raw, 170  
     \_\_\_\_\_ coltsfoot, 403  
     \_\_\_\_\_ compound, 403  
     \_\_\_\_\_ columbo, compound, 174  
     \_\_\_\_\_ contrayerva, 176  
     \_\_\_\_\_ cotton-root, 224  
     \_\_\_\_\_ dandelion, 398  
     \_\_\_\_\_ dogwood, 181  
     \_\_\_\_\_ elder bark, 359  
     \_\_\_\_\_ elecampane, 250  
     \_\_\_\_\_ elm bark, 403  
     \_\_\_\_\_ ergot, 196  
     \_\_\_\_\_ figs, 213  
     \_\_\_\_\_ compound, 213  
     \_\_\_\_\_ geranium, 222  
     \_\_\_\_\_ guaiacum wood, 226  
     \_\_\_\_\_ compound, 226  
     \_\_\_\_\_ hairy horehound, 124  
     \_\_\_\_\_ hardhack, 387  
     \_\_\_\_\_ horehound, compound, 274



- Decoction, horse balm, 172  
 ——— horsechestnut bark, 247  
 ——— compound, 247  
 ——— Iceland moss, 156  
 ——— Indian hemp, 109  
 ——— sarsaparilla, 231  
 ——— ipecacuanha, 254  
 ——— logwood, 228  
 ——— madder, 354  
 ——— marshmallow, 86  
 ——— matico, 275  
 ——— mezereon, 278  
 ——— compound, 279  
 ——— New Jersey tea, 154  
 ——— oak bark, 342  
 ——— oatmeal, compound, 123  
 ——— Pareira brava, 306  
 ——— Peruvian bark, 162  
 ——— and cascarilla, 163  
 ——— rhatany, 163  
 ——— compound, 163  
 ——— pipsissewa, 157  
 ——— compound, 157  
 ——— pomegranate rind, 225  
 ——— root, 225  
 ——— poppy heads, 306  
 ——— sarsaparilla, 362  
 ——— compound, 362  
 ——— Feltz's, 363  
 ——— Jauperand's, 363  
 ——— Vinache's, 363  
 ——— Zittman's, 363  
 ——— seneka, 372  
 ——— soot, 215  
 ——— stavesacre, 388  
 ——— tormentil, 402  
 ——— uva ursi, 404  
 ——— water avens, 222  
 ——— wild indigo, 124  
 ——— willow bark, 358  
 ——— wormseed, 157  
 Delphinia, 189  
 ——— ointment, 189  
 ——— solution, 189  
 Demulcents, 56  
 Dentifrice, bitartrate potassa, 338  
 ——— burnt hartshorn, 180  
 ——— charcoal, 146  
 ——— chlorinated lime, 134  
 ——— electuary, 136, 137  
 ——— mastic, 275  
 ——— orris root, 255  
 ——— Peruvian bark, 161  
 ——— phosphate of lime, 137  
 ——— rhatany, 259  
 ——— sulphate of quinine, 347  
 ——— willow bark, 358  
 ——— with carbonate of lime, 136  
 Dewberry, 354  
 Dewees' carminative, 118, 270  
 Diaphoretics, 57  
 Diet-drink, Lisbon, 363  
 Diet in convalescence, 63  
 Digitalin, 192  
 ——— pills, 192  
 ——— granules, 192  
 Diluents, 57  
 Dippel's animal oil, 289  
 ——— liniment, 289  
 ——— mixture, 289  
 ——— tincture, 289  
 Displacement, 452  
 Distillation, 455  
 Distilled oils, 456  
 ——— waters, 455  
 Dittany, 185  
 ——— infusion, 185  
 Dittany, oil, 185  
 Diuretics, 57  
 Dog rose, 352  
 ——— confection, 352  
 Dogwood, 180  
 ——— decoction, 181  
 ——— Jamaica, 311  
 ——— pills, 181  
 ——— round-leaved, 180  
 ——— swamp, 180  
 ——— tree, 180  
 ——— wine, 181  
 Dogsbane, 109  
 Donovan's solution, 114  
 Dose of medicines, 50  
 Drops, 17, 462  
 ——— Bateman's, 302  
 ——— Battley's, 302  
 ——— carminative, 269  
 ——— Koechlin's, 186  
 ——— odontalgic, 301  
 ——— table of, 17  
 ——— toothache, 298  
 Draughts, 454  
 Draught, anodyne, 303  
 ——— morphia, 280  
 ——— solution, iodide, arsenic and mercury,  
 114  
 ——— sugar of milk, 267  
 ——— sulphate potassa, effervescing, 337  
 Eau de Javelle, 320  
 ——— de luce, 97, 392  
 Effervescing draught, 333, 381  
 ——— powders, 333, 381  
 Eggs, 305  
 ——— cerate, 305  
 ——— clyster, emollient, 305  
 ——— restorative, 305  
 ——— emulsion, 305  
 ——— liniment, 305  
 ——— mixture, 305  
 ——— with brandy, 305  
 ——— wine, 305  
 ——— oil, 305  
 Elaterine, 195  
 ——— tincture, 195  
 Elaterium, 194  
 ——— mixture, 194  
 ——— oil, 195  
 ——— pills, 194  
 Elder, 359  
 ——— decoction of bark, 359  
 ——— extract of berries, 359  
 ——— mixture, 359  
 ——— gargle, 359  
 ——— ointment, 359  
 ——— leaves, 359  
 ——— vinegar, 359  
 ——— water of flowers, 359  
 Elecampane, 250  
 ——— decoction, 250  
 ——— extract, 250  
 ——— mixture, 250  
 ——— oxymel, compound, 250  
 ——— pills, compound, 250  
 Electuaries, 449  
 Electuary, anthelmintic, 84  
 ——— dentifrice, 136, 137  
 ——— laxative, 84  
 ——— for teeth, 87  
 ——— of alum, 88  
 ——— angustura, 103  
 ——— bitartrate of potassa, 339  
 ——— black oxide of iron, 210  
 ——— pepper, 310  
 ——— burnt sponge, 387



- Electuary of calamus, 133  
 — calomel, 241  
 — carburet of iron, 203  
 — catechu, 153  
 — charcoal, 146  
 — cinnamon, compound, 167  
 — cowhage, 284  
 — cubebs, 184  
 — indigo, 250  
 — jalap, 256  
 — juniper, 258  
 — kino, 259  
 — logwood, extract of, 228  
 — male fern, 214  
 — orange leaves, 119  
 — peel, 119  
 — Peruvian bark, 164  
     — astringent, 164  
     — with catechu, 164  
     — cloves, 164  
     — iron, 164  
     — sulphur, 164  
     — tin, 164  
 — pomegranate root, 225  
 — rhatany, 259  
 — rhubarb, 349  
 — roses, 353  
 — scurvy grass, 169  
 — semen contra, 115  
 — senna and cremor tartar, 373  
     — figs, 373  
     — rhubarb, 373  
     — sulphur, 373  
 — soap, 361  
 — squill, 370  
 — sulphate of copper, 188  
     — soda, 384  
 — sulphur, 393  
     — compound, 393  
 — sulphuret of potassium, 326  
 — tamarinds, 397  
 — tin, 387  
 — turpentine, 399  
 — valerian, 405  
 — Virginia snakeroot, 376  
 Elemi, 195  
 — cautery plaster, 195  
 — ointment, 195  
 Elixir, vitriol, 75  
 — Boerhaave's, 85  
 — Clauder's, 85  
 — Garus', 86, 183  
 — ipecacuanha, 255  
 — Lettsom's, 303  
 — orange peel, 119  
 — scammony, 366  
 — Stoughton's, 86  
 Elm bark, 403  
 — decoction, 403  
 — infusion, 403  
 — slippery, 403  
 Elutriation, 447  
 Embrocation, caraway, 149  
 — croton oil, 295  
 — petroleum, 307  
 Embrocations, 469  
 Emetic, tartar, 107  
 Emetics, 57  
 Emetin, 195  
 — impure, 195  
 — lozenges, 196  
 — mixture, 195  
 — pure, 195  
 — syrup, 196  
 Emmenagogues, 58  
 Emollients, 56  
 Emulsions, 454  
 — almond, 99, 100  
     — aromatic, 100  
 — anthelmintic, 291, 375  
 — artificial musk, 392  
 — benzoin, cosmetic, 129  
 — camphor, 139  
 — cantharides, 142  
 — carbonate of potassa, 332  
 — castor oil seeds, 291  
 — chloroform, 159  
 — copaiba, 178  
 — croton oil, 294  
 — cubebs, ethereal extract, 184  
 — egg, 305  
 — hemp seed, 141  
 — jalap, 256  
 — kermes mineral, 106  
 — lupulin, 268  
 — manna, 373  
 — nitrated, 336  
 — opoponax, 304  
 — purgative, 271  
 — sal ammoniac, 96  
 — scammony, 366  
     — compound, 366  
     — seneka, 372  
     — squill, 370  
     — sulphate soda, 384  
     — tartar emetic, 108  
 Enemata, 55  
 Epispastics, 58  
 Ergot, 196  
 — clyster, 197  
 — decoction, 196  
 — extract, 197  
 — infusion, 196  
 — injection, 197  
 — mixture, 196  
     — extract, 197  
 — oil, 197  
 — pills, compound, 197  
     — extract, 197  
 — powder, compound, 196  
 — syrup, 196  
 — tincture, 196  
 — wine, 196  
 Errhine, alum, 88  
     — euphorbium, 199  
 Errhines, 60  
 Escharotics, 60  
 Essence of angelica, 103  
     — ginger, 416  
     — lavender, compound, 265  
     — lemon, 265  
     — peppermint, 277  
     — roses, 352  
     — sarsaparilla, 368  
     — compound, 369  
     — soap, 361  
     — camphorated, 361  
     — vanilla, 407  
     — Ward's for headache, 140  
 Essential oils, 456  
 Ether, acetic, 79  
     — mixture, 79  
     — spirit, 79  
     — cupreous, 186  
     — hydrocyanic, 80  
     — muriatic, 80  
     — mixture, 80  
     — spirit of, 80  
     — nitrous, 79  
     — mixture, 80  
     — sweet spirit of, 79  
     — mixture of, 80  
     — phosphorated, 308



Ether, sulphuric, 80  
     — ethereal oil, 80  
     — Hoffmann's anodyne, 80  
     — lotion, 80  
     — mixture with camphor, 81  
     —                   turpentine, 81  
     — rectified, 80  
     — spirit, 80  
     — syrup, 81  
     — terebinthinated, 81  
 Ethereal oil, 80  
 Euphorbium, 199  
     — oil, 199  
     — plaster, 199  
     — sternutatory, 199  
 Examination of excretions, 47  
 Exercise in convalescence, 85  
 Expectorants, 60  
 Extracts, 448  
 Extract of aconite, 77  
     — alcoholic, 77  
     — ammoniated, 78  
     — American centaury, 356  
     —                   hellebore, 409  
     — anemone, 142  
     — arnica, 113  
     — asparagus roots, 116  
     —                   shoots, 116  
     — belladonna, 127  
     —                   alcoholic, 127  
     — bistort, 319  
     — bittersweet, 193  
     — black hellebore, 230  
     —                   alcoholic, 230  
     — broom, 371  
     — buchu, 192  
     —                   fluid, 192  
     — buckbean, 278  
     — butternut, 257  
     — cabbage tree bark, 222  
     — Canada fleabane, 197  
     — cantharides, 142  
     —                   ethereal, 143  
     — cardamom, 148  
     — carrot root, 148  
     — cascarilla, 150  
     — celandine, 157  
     — centaury, 154  
     — cevadilla, 355  
     — chamomile, 104  
     — colchicum bulb, 170  
     —                   acetic, 170  
     — colocynth, 172  
     —                   compound, 172  
     — columbo, 174  
     — contrayerva, 176  
     — cubebs, 184  
     —                   alcoholic ethereal, 184  
     — dandelion, 398  
     —                   fluid, 398  
     — elder berries, 359  
     — elecampane, 250  
     — ergot, 197  
     — foxglove, 190  
     — galls, 218  
     — gentian, 220  
     — ginger, ethereal, 416  
     — guaiacum wood, 227  
     — hardhack, 387  
     — hemlock, 175  
     —                   alcoholic, 175  
     — hemp, 141  
     — henbane, alcoholic, 248  
     —                   aqueous, 248  
     — hops, 232  
     — horehound, 274  
     — Indian hemp, 109

Extract of ipecacuanha, 253  
     — jalap, 256  
     —                   alkaline, 256  
     — juniper, 257  
     — lettuce, 262  
     —                   watery, 263  
     — liquorice, 223  
     — lobelia, acetic, 268  
     — logwood, 228  
     — male fern, 214  
     — marygold, 137  
     — may apple, 318  
     — mezereon, 279  
     — myrrh, 286  
     — nux vomica, 288  
     — oak bark, 342  
     — opium, acetous, 295  
     —                   alcoholic, 295  
     —                   aqueo-alcoholic, 295  
     —                   aqueous, 295  
     —                   by fermentation, 296  
     —                   denarcotized, 296  
     —                   roasted, 296  
     —                   vinous, 295  
     — pareira brava, 307  
     — paullinia, alcoholic, 307  
     — pellitory, 341  
     — Peruvian bark, 162  
     —                   vinous, 162  
     — pink-root, compound, 386  
     —                   fluid, 386  
     — pipsissewa, 157  
     — poison oak, 402  
     — pomegranate root, 225  
     — poppy heads, 306  
     — quassia, 341  
     — rhatany, 259  
     — rhubarb, 350  
     —                   fluid, 350  
     — rue, 355  
     — sarsaparilla, 363  
     —                   fluid, 363  
     — scammony, 366  
     — seneka, with squill, 372  
     — senna, alcoholic, 373  
     —                   fluid, 375  
     — squill, 370  
     — stramonium leaves, 388  
     —                   seeds, 388  
     — succory, 160  
     — tansy, 397  
     — tobacco, 396  
     — tormentil, 401  
     — uva ursi, 404  
     — valerian, 406  
     —                   fluid, 406  
     — vanilla, fluid, 407  
     — wood sorrel, 67  
     — wormwood, 65  
 Eyebright, 199  
 False sarsaparilla, 110  
 Fats, 460  
 Fecula of hemlock, 175  
 Fennel, 215  
     — infusion, 215  
     — oil, 215  
     — ointment, 215  
     — powder, compound, 215  
     — water, 215  
 Fern, male, 214  
     — bolus, 214  
     — electuary, 214  
     — extract, ethereal, 214  
     —                   pills, 214  
     — jelly, 214  
     — mixture, 214



- Fern, male, oil, 214  
     — pills, 214  
     — powder, 214  
     — tincture, 214  
 Ferrocyanate of quinine, 343  
     — zinc, 412  
 Ferrocyanuret of iron, 205  
 Figs, 213  
     — coffee, 214  
     — decoction, 213  
     — compound, 213  
     — gargle, 213  
     — paste, 214  
 Figwort, 371  
     — oil, 371  
     — ointment, 371  
 Filter, Boullay's, 452  
 Fixture, 101, 292  
 Fixed oils, 460  
 Flaxseed, 267  
     — cataplasm, 267  
     — clyster of oil, 267  
     — infusion, 267  
     — liniment of oil, 267  
     — meal, cataplasm, 267  
     — compound, 257  
     — mixture, 267  
 Fleabane, 197  
     — Canada, 197  
     — infusion, 197  
     — extract, 197  
     — infusion, 197  
 Flies, Spanish, 141  
 Florentine orris, 255  
 Flour, boiled, 418  
     — buchu, 192  
 Fluid extract of pinkroot, 386  
     — rhubarb, 350  
     — sarsaparilla, 363  
     — senna, 375  
     — spigelia, compound, 386  
     — valerian, 406  
     — vanilla, 407  
 Flummery of oatmeal, 123  
 Fomentations, 468  
 Fomentation, arnica, 113  
     — with rue, 113  
     — aromatic, 353  
     — carbonate of potassa, 332  
     — common salt, 378  
     — mustard, 377  
     — nitric acid, 74  
     — opium, 299  
     — phosphoric acid, 75  
     — sal ammoniac, 95  
     — sulphate of zinc, 414  
     — wine of opium, 301  
 Fowl, with rice, 419  
 Fowler's solution, 330  
 Foxglove, 190  
     — bolus, 190  
     — extract, 190  
     — infusion, 190  
     — mixture, expectorant, 191  
     — with acetate of lead, 191  
     — potassa, 191  
     — tartaric acid, 191  
     — ointment, 191  
     — pills, 190  
     — plaster, 191  
     — powder, 190  
     — syrup, 192  
     — tincture, 191  
     — ethereal, 191  
     — mixture, 191  
     — vinegar, 191  
 Frictions, 469  
 Frost-weed, 229  
 Frumenty, 420  
 Fuligokali, 216  
     — ointment, 216  
     — sulphuretted, 216  
     — pills, 216  
 Fumigations, 472  
 Fumigation with belladonna, 127  
 Furniture of sick room, 48  
 Galbanum, 217  
     — mixture, 217  
     — pills, compound, 217  
     — plaster, 217  
     — compound, 217  
     — purified, 217  
     — tincture, 217  
     — compound, 217  
 Gallic acid, 71  
     — injection, 71  
     — pills, 71  
 Galls, 218  
     — extract, 218  
     — ointment, 218  
     — gargle, 218  
     — infusion, 218  
     — compound, 218  
     — lotion, 218  
     — ointment, 219  
     — compound, 219  
     — powder, compound, 218  
     — syrup, 218  
     — tincture, 218  
 Gamboge, 219  
     — pills, compound, 219  
     — powder, compound, 219  
     — mixture, 219  
     — with elatine, 219  
     — solution, alkaline, 219  
     — tincture, alkaline, 219  
     — ammoniacal, 220  
 Gargle, acetate of ammonia, 92  
     — lead, 315  
     — alum, 88, 89  
     — ammoniated copper, 187  
     — borax, 380  
     — carbonate of soda, 383  
     — cayenne pepper, 145  
     — chloride of soda, 383  
     — chlorine, 158  
     — cinnamon, tincture, 168  
     — contrayerva, 117  
     — cyanuret of mercury, 242  
     — elder flowers, 359  
     — figs, 213  
     — galls, 218  
     — iodine, 252  
     — mercurial, 233  
     — muriatic acid, 74  
     — mustard, 377  
     — myrrh, 287  
     — nitrate of potassa, 336  
     — oak bark, 342  
     — orpiment, 114  
     — Peruvian bark, 165  
     — pomegranate rind, 225  
     — sage, 359  
     — scurvy grass, 169  
     — subacetate of lead, 314  
     — sulphate of copper, 188  
     — quinine, 347  
     — zinc, 414  
     — tannic acid, 76  
     — tormentil, 402  
     — vinegar, 67  
 Garlic, 81  
     — cataplasm, 81



- Garlic, lotion, capillary, 81  
 — syrup, 81  
 Gelatine and Peruvian bark, 165  
 Gentian, 220  
 — extract, 220  
 — syrup, 221  
 — infusion, 221  
 — compound, 221  
 — with rhubarb, 221  
 — mixture, 221  
 — pills, compound, 220  
 — with ammonia, 220  
 — powder, compound, 220  
 — syrup, 221  
 — tincture, 221  
 — acidulated, 221  
 — alkaline, 221  
 — ammoniacal, 221  
 — compound, 221  
 — mixture, 221  
 — with sulphuric acid, 221  
 — wine, 221  
 Gentianin, 222  
 — pills, 222  
 — syrup, 222  
 — tincture, 222  
 German chamomile, 276  
 Ginger, 415  
 — beer, 415, 420  
 — essence, 416  
 — extract, ethereal, 416  
 — syrup, 416  
 — tincture, 416  
 — lozenges, 415  
 — oil, 416  
 — spice plaster, 416  
 — syrup, 415  
 Glass, soluble, 337  
 Gloucester jelly, 417  
 Glycerin, 223  
 — ointment, 223  
 Godfrey's cordial, 302  
 Gold, 120  
 — ammoniated, 123  
 — pills, 123  
 — chloride, 120  
 — bolus, 121  
 — caustic, 121  
 — collyrium, 121  
 — ointment, 121  
 — pills, 121  
 — powder, 120  
 — tincture, 121  
 — with sodium, 121  
 — lozenges, 121  
 — ointment, 122  
 — pills, 121  
 — powder, 121  
 — solution, 121  
 — syrup, 121  
 — compound, 122  
 — cyanide, 122  
 — lozenges, 122  
 — pills, 122  
 — powder, 122  
 — solution, 122  
 — iodide, 122  
 — ointment, 120  
 — oxide, 122  
 — pills, 123  
 — powder, 122  
 — powder, 120  
 — with iron, 120  
 — purple of Cassius, 123  
 — syrup, 120  
 Golden sulphuret of antimony, 106  
 Goldthread, 179  
 — infusion, 179  
 — tincture, 179  
 Gondret's ointment, 93  
 Goulard's balsam, 315  
 Granulation, 447  
 Granules of digitalin, 192  
 Gravity, specific, 20  
 Grease balls, 361  
 Griffith's mixture, 202  
 Griffith's pills, 84  
 Groat gruel, 123  
 Gruel, oatmeal, 123  
 Guaiacum, 226  
 — decoction, 226  
 — compound, 226  
 — extract, 226  
 — mixture, 227  
 — with bittersweet, 227  
 — oil, 227  
 — compound, 227  
 — pills with aloes, 227  
 — antimony, 227  
 — sulphur, 227  
 — turpentine, 227  
 — powder, compound, 227  
 — tincture, 227  
 — ammoniated, 228  
 — and camphor, 228  
 — copaiba, 228  
 — mixture, odontalgic, 228  
 — with henbane, 228  
 Gum Arabic, 66  
 — lozenges of, 66, 67  
 — mixture of, 66, 67  
 — mucilage of, 66  
 — paste of, 66  
 — syrup of, 66  
 — compound, 66  
 — tincture of, 67  
 — plaster, 98  
 — tragacanth, 402  
 Hair dye, 111  
 Hairy horehound, 124  
 — decoction, 124  
 Hardhack, 386  
 — decoction, 387  
 — extract, 387  
 Hartshorn, 180  
 — burnt, 180  
 — dentifrice, 180  
 — jelly, 180  
 — compound, 180  
 Hedge hyssop, 226  
 — powder, compound, 226  
 — wine, 226  
 Hellebore, American, 409  
 — black, 229  
 — white, 408  
 Hemlock, 175  
 — cataplasm, 176  
 — collutory, 175  
 — extract, 175  
 — alcoholic, 175  
 — fecula, 175  
 — infusion, 175  
 — mixture, 176  
 — oil, 176  
 — ointment, 175  
 — pills with calomel, 175  
 — dandelion, 175  
 — ipecacuanha, 175  
 — pitch, 312  
 — plaster, 176  
 — powder, 175  
 — compound, 176  
 — suppository, 176



- Hemlock, tincture, 175  
     — etheral, 175  
 Hemp, 141  
     — emulsion, 141  
     — extract, 141  
     — resin, 141  
     — tincture, 141  
     — Indian, 109  
 Henbane, 248  
     — extract, alcoholic, 248  
     — aqueous, 248  
     — infusion, 248  
     — compound, 248  
     — liniment, 249  
     — mixture, 249  
     — with antimony, 249  
     — squill, 249  
     — ointment, 249  
     — compound, 249  
     — pills, compound, 248  
     — and ipecacuanha, 248  
     — opium, 248  
     — powder, compound, 248  
     — tincture, 249  
     — etheral, 249  
 Hive syrup, 368  
 Hoffmann's anodyne, 80  
 Holly, sea, 197  
 Honeys, 451  
 Honey, borax, 380  
     — clarified, 276  
     — ceromel, 276  
     — hydromel, 276  
     — mixture, expectorant, 276  
     — oxymel, 276  
     — pectoral, 276  
     — prepared, 276  
     — roses, 353  
     — violets, 409  
     — water, 276  
 Hooper's pills, 83  
 Hops, 232  
     — cataplasm, 232  
     — extract, 232  
     — infusion, 232  
     — mixture, 232  
     — ointment, 232  
     — tincture, 232  
     — alkaline, 232  
 Horehound, 274  
     — candy, 275  
     — decoction, compound, 274  
     — extract, 274  
     — hairy, 124  
     — mixture, pectoral, 274  
     — syrup, 274  
 Horse balm, 172  
     — decoction, 172  
     — oil, 172  
     — tincture, 172  
     — chestnut, 247  
     — decoction, 247  
     — compound, 247  
     — extract, 247  
     — powder, compound, 247  
 Horsemint, 279  
     — liniment, 279  
     — oil, 279  
 Horseradish, 112  
     — cataplasm, compound, 112  
     — infusion, 112  
     — mixture, 112  
     — spirit, compound, 112  
 Hot bath, 464  
 Hundred-leaved rose, 352  
 Hungary water, 354  
 Hydrargyro-iodide of potassium, 325  
 Hydriodate of ammonia, 94  
 Hydriodic acid, 72  
 Hydrocyanic acid, 71  
     — extemporaneous, 71  
     — injection, 72  
     — julep, 72  
     — lotion, 72  
     — mixture, 72  
     — syrup, 72  
     — ether, 80  
 Hydromel, 276  
     — pectoral, 276  
 Hydrometers, 20  
 Hydrometrical equivalents, 26  
 Hydrosulphuric acid, 72  
     — artificial sulphuretted water, 73  
     — bath, hydrosulphuretted, 72  
     — lotion, hydrosulphuretted, 73  
     — powder against salivation, 72  
 Hyposulphited sulphuret of potassium, 326  
 Iceland moss, 156  
     — cetrarine, 156  
     — powder, 156  
     — chocolate, 156  
     — decoction, 156  
     — jelly, 156  
     — mixture, 156  
 Imperial, 339  
     — measure, 16  
 Incompatibles, table of, 422  
 Indelible ink, 111  
 Indian hemp, 109  
     — decoction, 109  
     — extract, 109  
     — physic, 223  
     — sarsaparilla, 230  
     — decoction, 231  
     — infusion, 230  
     — mixture, 230  
     — syrup, 230  
     — turnip, 115  
 Indigo, 250  
     — electuary, 250  
     — pills, 250  
     — powder, 250  
     — sulphate, 250  
     — wild, 124  
     — decoction, 124  
     — ointment, 124  
 Infusions, 451  
 Infusion, American centaury, 356  
     — columbo, 215  
     — senna, 151  
     — ammoniated copper, 187  
     — angustura, 103  
     — arnica, 113  
     — compound, 113  
     — benne, 377  
     — bistort, 130  
     — black purslane, 198  
     — blessed thistle, 154  
     — bloodroot, 360  
     — boneset, 198  
     — compound, 198  
     — broom, 371  
     — buchu, 192  
     — compound, 192  
     — buckbean, 278  
     — burdock, 110  
     — calamus, 133  
     — camphor, 139  
     — Canada fleabane, 197  
     — cantharides, 142  
     — caraway, 148  
     — carrot seeds, 148



- Infusion, cascarilla, 150  
     alkaline, 150  
     catechu, 153  
     compound, 153  
     catnep, 152  
     Cayenne pepper, 145  
     chamomile, 104  
     cherry laurel, 264  
     chiretta, 158  
     cinnamon, 167  
     cloves, 149  
     columbo, 174  
     with ginger, 174  
     rhubarb, 174  
     dandelion, 398  
     dittany, 185  
     elm bark, 403  
     ergot, 196  
     fennel, 215  
     flaxseed, 267  
     fleabane, 197  
     foxglove, 191  
     galls, 218  
     compound, 218  
     gentian, 220  
     compound, 220  
     with rhubarb, 221  
     goldthread, 179  
     hemlock, 175  
     henbane, 248  
     compound, 248  
     hops, 232  
     horseradish, 112  
     compound, 112  
     Indian sarsaparilla, 230  
     iron, bitter, 209  
     juniper, 257  
     compound, 257  
     kino, 259  
     linden flowers, 400  
     compound, 400  
     lobelia, 268  
     logwood, 228  
     maidenhair, 79  
     malt, 272  
     marygold, 137  
     matico, 275  
     compound, 275  
     mayweed, 181  
     mint, 278  
     compound, 278  
     mudar, 138  
     orange peel, 119  
     compound, 119  
     pareira brava, 306  
     parsley root, 308  
     pennyroyal, 229  
     persimmon bark, 193  
     Peruvian bark, 163  
     compound, 163  
     with iron, 164  
     lime water, 163  
     magnesia, 163  
     serpentaria, 163  
     valerian, 163  
     pinkroot, 386  
     compound, 386  
     pleurisy root, 115  
     quassia, 341  
     compound, 341  
     red poppy, 352  
     rhatany, 259  
     rhubarb, 349  
     alkaline, 350  
     roses, acid, 353  
     safflower, 148  
     saffron, 183
- Infusion, sage, 358  
     compound, 359  
     sarsaparilla, 362  
     alkaline, 362  
     sassafras bark, 365  
     compound, 365  
     savine, 357  
     seneka, 372  
     compound, 372  
     senna, 374  
     compound, 374  
     with buckthorn, 375  
     coffee, 374  
     lemon juice, 375  
     rhubarb, 375  
     tamarinds, 374  
     tincture of senna, 375  
     simaruba, 377  
     compound, 377  
     skunk cabbage, 193  
     slippery elm, 403  
     southern wood, 114  
     succory, 160  
     tamarind, 397  
     tansy, 397  
     tobacco, 396  
     tulip tree bark, 267  
     valerian, 405  
     compound, 405  
     Virginia snakeroot, 376  
     compound, 376  
     with ether, 377  
     yellow ladies' bedstraw, 218  
     wild cherry bark, 340  
     ginger, 115  
     wormwood, 65
- Inhalations, 462  
 Inhalation, balsam of tolu, 401  
 Injection, acetate of zinc, 413  
     alum, 89  
     ammonia, 91  
     carbonate of soda, 382  
     catechu, 153  
     caustic potassa, 329  
     chloride of zinc, 411  
     chlorinated lime, compound, 134  
     chlorine, 158  
     chloroplatinate of sodium, 313  
     copaiba, 178  
     corrosive sublimate, 238  
     cubeb, 184  
     detersive, 85  
     ergot, 197  
     gallic acid, 71  
     hydrocyanic acid, 72  
     iodide of iron, 208  
     potassium, 322  
     ioduretted, 324  
     kino, 259  
     morphia, 280  
     opium, 299  
     pyroligneous acid, 69  
     rhatany, 260  
     sedative, 306  
     soot, 216  
     stimulating, 287  
     subacetate of lead, 314  
     and lime water, 315  
     sulphate of copper, 188  
     iron, 211  
     zinc, 414  
     sulphuret of potassium, 327  
     tannic acid, 76
- Ink, blue, 206  
     indelible, 111  
     without mordant, 111  
     writing, 206



- Inspissated juices, 448  
 Introduction, 9  
 Iodide of arsenic, 113  
     — and mercury, 114  
     — barium, 125  
     — gold, 122  
     — iron, 206  
     — lead, 316  
     — mercury, 242  
     — red, 243  
     — potassium, 321  
     — silver, 110  
     — starch, 102  
     — sulphur, 395  
     — zinc, 412  
 Iodine, 251  
     — bath, 251  
     — cataplasm, 252  
     — clyster, 252  
     — gargle, 252  
     — liniment, 252  
     — lotion, 251  
     — mixture, 251  
     — with iodide of potassium, 251  
     — ointment, 252  
     — compound, 252  
     — with mercury, 252  
     — oil of tobacco, 252  
     — pills, 251  
     — plaster, 252  
     — tincture, 251  
     — compound, 251  
     — ethereal, 251  
 Ipecacuanha, 252  
     — bolus, 253  
     — confection, 254  
     — decoction, 254  
     — elixir, 255  
     — extract, 253  
     — lozenges, 254  
     — mixture, 254  
     — pills, compound, 254  
     — with centaury, 254  
     — foxglove, 254  
     — opium, 253  
     — powder with antimony, 253  
     — calomel, 253  
     — carb. soda, 253  
     — muriate ammonia, 253  
     — myrrh, 253  
     — nitre, 253  
     — opium, 252  
     — rhubarb, 253  
     — tartar emetic, 253  
     — tragacanth, 253  
     — syrup, 254  
     — compound, 254  
     — wild, 198  
     — wine, 255  
     — alkaline, 255  
     — compound, 255  
     — with tartar emetic, 255  
 Iris, Florentine, 255  
 Iron, 199  
     — acetate, 200  
     — and alum, 200  
     — tincture, alcoholic, 200  
     — ethereal, 200  
     — ammoniated, 201  
     — bolus, 201  
     — mixture, 201  
     — pills, 201  
     — powder, compound, 201  
     — tincture, 201  
     — arseniate, 201  
     — ointment, 201  
     — pills, 201  
     — bromide, 201  
     — ointment, 202  
     — pills, 201  
     — carbonate, 202  
     — artificial chalybeate water, 203  
     — pills, 202  
     — plaster, 203  
     — saccharine, 202  
     — subcarbonate, 202  
     — mixture, 202  
     — pills, comp., 203  
     — powder, 202  
     — wine, 203  
     — carburet, 203  
     — electuary, 203  
     — mixture, 204  
     — ointment, 204  
     — pills, 203  
     — plaster, 204  
     — powder, 203  
     — prepared, 203  
     — chloride, 204  
     — and acetate of lead, 204  
     — mixture, 204  
     — syrup, 204  
     — tincture, 204  
     — ethereal, 204  
     — citrate, 205  
     — ammonio, 205  
     — and quinine, 205  
     — liquid, 205  
     — syrup, 205  
     — tincture, 205  
     — wine, 205  
     — aromatic, 205  
     — ferrocyanuret, blue ink, 206  
     — mixture, 206  
     — pills, 206  
     — powder, 205  
     — compound, 205  
     — Prussian blue, 205  
     — filings, 199  
     — gallate, 206  
     — writing ink, 206  
     — iodide, 206  
     — bath, 208  
     — injection, 208  
     — lozenges, 206  
     — mixture, 207  
     — pills, 207  
     — solution, 207  
     — officinal, 207  
     — syrup, 207  
     — and chloride of iron, 208  
     — tincture, 207  
     — wine, 207  
     — lactate, 208  
     — lozenges, 208  
     — pills, 208  
     — syrup, 208  
     — malate, 208  
     — bitter infusion, 209  
     — mixture, 209  
     — solution, 208  
     — tincture, 209  
     — metallic, prepared, 199  
     — bolus, 200  
     — mixture, 200  
     — pills, 200  
     — powder, 200  
     — compound, 200  
     — nitrate, 209  
     — solution, 209  
     — oxide, black, 209  
     — electuary, 210  
     — pills, 210  
     — powder, 209



- Iron, oxide, hydrated, 209  
     — red, 210  
     — pills, 210  
     — plaster, 210  
     — persesquintrate, 209  
     — phosphate, 210  
     — proto-iodide, 207  
     — pills, 207  
     — syrup, 207  
     — tincture, 207  
     — sulphate, 210  
     — dried, 211  
     — injection, 211  
     — mixture, 212  
     — compound, 211  
     — pills, 211  
     — Bland's, 211  
     — compound, 211  
     — with rhubarb, 211  
     — powder, 211  
     — solution, 211  
     — syrup, 212  
     — sulphuret, 212  
     — tannate, 213  
     — tartrate, and potassa, 212  
     — powder, and columbo 212,  
     — effervescent, 212  
     — solution, 212  
     — tincture, compound, 213  
     — wine, 212  
     — compound, 212  
     — ammonio, 213  
     — solution, 213  
     — valerianate, 213  
 Isinglass, 249  
     — blanc mange, 420  
     — court plaster, 249  
     — diamond cement, 250  
 Issues, 470  
 Jalap, 255  
     — electuary, 256  
     — emulsion, 256  
     — extract, 256  
     — alkaline, 256  
     — mixture, 257  
     — pills, compound, 256  
     — with calomel, 256  
     — powder, compound, 256  
     — with calomel, 255  
     — cremor tartar, 256  
     — ipecacuanha, 256  
     — scammony, 256  
     — resin, 256  
     — soap, 257  
     — tincture, 257  
     — compound, 257  
     — mixture, 257  
 Jamaica dogwood, 311  
     — pepper, 309  
     — tincture, 311  
 James's powder, 109  
 Jelly almond, 417  
     — biscuit, 417  
     — bread, 417  
     — calves' foot, 418  
     — carrageen, 160  
     — chicken, 419  
     — Gloucester, 417  
     — hartshorn, 180  
     — compound, 180  
     — Iceland moss, 156  
     — male fern, 214  
     — orange leaves, 119  
     — pectoral, 372  
     — rice, 305  
     — starch, 102  
     — Jelly, tapioca, 397  
     — Julep, alum, 89  
     — hydrocyanic, 72  
 Juniper, 257  
     — electuary, 258  
     — extract, 257  
     — infusion, 257  
     — compound, 257  
     — liniment, 258  
     — mixture of oil, 258  
     — spirit, compound, 258  
 Kermes mineral, 105  
 Kermesine powder, 106  
 Kinate of cinchonia, 166  
     — quinine, 344  
 Kino, 258  
     — electuary, 259  
     — infusion, 259  
     — injection, 259  
     — pills, compound, 259  
     — powder, compound, 258  
     — syrup, 259  
     — tincture, 259  
 Labarraque's solution, 383  
 Lac, 262  
     — bleached, 262  
     — cement, 262  
     — sealing wax, 262  
     — solution, aqueous, 262  
 Lactate of iron, 208  
     — quinine, 344  
 Lactic acid, 73  
     — lemonade, 73  
     — lozenges, 73  
 Lactucarium, 262  
     — mixture, 262  
     — pills, 262  
     — syrup, 262  
     — tincture, 262  
 Ladies' bed straw, yellow, 218  
     — infusion, 218  
     — wine, 218  
 Lady Webster's pills, 83  
 Lancaster black drop, 300  
 Larkspur, 189  
     — tincture, 189  
 Laudanum, 301  
     — Smith's, 302  
     — Swediaur's, 302  
     — Sydenham's, 301  
 Laurel, 263  
     — cherry, 264  
     — oil, 263  
     — ointment, 263  
 Lavender, 264  
     — essence, compound, 265  
     — ointment, 265  
     — powder, compound, 264  
     — spirit, 264  
     — compound, 265  
     — tincture, ethereal, 265  
     — water, 264  
 Lead, 313  
     — acetate, 313  
     — cataplasm, 315  
     — cerate, 314  
     — collyrium, 314  
     — gargle, 315  
     — Goulard's balsam, 315  
     — liniment, 314  
     — lotion, 315  
     — mixture, 315  
     — ointment, 314  
     — pile, 314  
     — paste, rubefacient, 314



- Lead, acetate, pills, 313  
     — plaster, 314  
     — powder, 313  
     — carbonate, 316  
         — ointment, 316  
         — camphorated, 316  
         — plaster, 316  
         — powder, camphorated, 316  
     — chloride, 316  
         — lotion, 316  
         — ointment, 316  
     — iodide, 316  
         — ointment, 317  
         — pills, 317  
         — powder, 317  
     — litharge, 317  
         — cataplasm, 317  
         — cerate, 317  
         — ointment, 317  
             — Banyer's, 317  
             — compound, 317  
         — plaster, 317  
             — Baynton's, 317  
             — diapalma, 317  
             — Gauthier's, 317  
     — red, 318  
         — plaster, Nuremberg, 318  
     — saccharate, 318  
         — nitro, 318  
         — solution, 318  
     — subacetate, 313  
         — cataplasm, 314  
         — cerate, 315  
         — collyrium, 314  
         — gargle, 314  
         — Garland's vegeto-mineral  
             water, 315  
         — injection, 314  
             — with lime, 315  
         — liniment, 315  
         — ointment, 315  
         — plaster, 315  
         — solution, 314  
 Leather wood, 193  
 Leeching, 477  
 Lemon, 265  
     — collutory of juice, 265  
     — Cologne water, 266  
     — essence, 265  
     — juice, artificial, 71  
     — lozenges of juice, 265  
         — oil, 265  
     — spirit, 266  
         — odoriferous, 266  
     — syrup, 266  
         — vinous of peel, 266  
     — water of peel, 266, 419  
     — unparalleled, 267  
 Lemonade, 265  
     — barberries, 129  
     — compound, 265  
     — dry, 71  
     — lactic acid, 73  
     — phosphoric acid, 75  
     — sulphuric acid, 76  
     — tartaric acid, 77  
     — with sulphate of soda, 384  
 Lettuce, wild, 262  
     — extract, 262  
         — lactucarium, 262  
             — lozenges, 263  
             — mixture, 262  
             — pills, 262  
             — syrup, 262  
             — tincture, 262  
     — thridace, 262  
     — syrup, 262  
 Lettuce, extract, watery, 263  
     — mixture, 263  
     — powder, 263  
     — mixture, with dandelion, 262  
     — syrup, 263  
     — tincture, 263  
     — water, 263  
         — mixture, 263  
 Lime, 135  
     — carbonate, 136  
         — dentifrice, 136  
         — electuary, 136  
         — lozenges, 136  
         — mixture, 137  
             — compound, 137  
             — powder, aromatic, 136  
             — compound, 136  
                 — with opium, 136  
         — precipitated, 136  
         — prepared chalk, 136  
             — oyster shells, 136  
     — chlorinated, 134  
         — collutory, 134  
         — dentifrice, 134  
         — injection, 134  
         — liniment, 134  
         — lozenges, 134  
         — ointment, 134  
         — preservative liquid, 134  
     — liniment, with sulphur, 135  
     — ointment, 136  
     — phosphate, 137  
         — electuary, dentifrice, 137  
         — hartshorn, burnt, 137  
         — powder, dentifrice, 137  
     — water, 135  
         — compound, 135  
         — and carbonate of potassa, 135  
             — milk, 135  
             — liniment, 135  
                 — with alcohol, 135  
                 — oil, 135  
                 — opium, 135  
                 — sulphur, 135  
         — mixture, antacid, 136  
 Linctus, gum Arabic, 67  
     — green, 100  
     — opiate, 303  
     — opium, 299  
     — sulphur, 393  
     — white, 101  
 Linden, 400  
     — infusion, 400  
     — compound, 400  
 Liniments, 469  
 Liniment, acetate of mercury, 236  
     — alum, 89  
     — amber, oil of, 392  
     — ammonia, 90  
         — camphorated, 93  
         — compound, 90  
         — sulphuretted, 91  
         — terebinthinate, 91  
     — belladonna, 128  
     — cajeput oil, 289  
     — camphor, 140  
         — compound, 140  
         — ethereal, 140  
         — turpentine, 141  
         — with vinegar, 140  
     — cantharides, 143  
         — aromatic, 144  
         — soap, 143  
         — stimulating, 144  
         — with camphor, 143  
     — carbonate of potassa, 332  
     — chlorinated lime, 134



- Liniment, chlorine, 159  
 — chloroform, 159  
 — cod liver oil, 172  
 — colchicum and camphor, 172  
 — cosmetic, 101  
 — croton oil, 294  
 — Dippel's oil, 289  
 — eggs, 305  
 — flaxseed oil, 267  
 — for chilblains, 285  
 — milk abscess, 314  
 — sore breasts, 136  
 — garlic, 81  
 — henbane, 249  
 — horsemint, 279  
 — iodide of potassium, 323  
 — iodine, 252  
 — juniper, 258  
 — laudanum and lime water, 303  
 — lime and sulphur, 136  
 — water and alcohol, 136  
 — oil, 136  
 — opium, 136  
 — mercurial, compound, 235  
 — opiated, 235  
 — mercury and chalk, 233  
 — muriatic acid, 74  
 — mustard, 378  
 — nitrate of mercury, 244  
 — nitric acid, 74  
 — opiate, 303  
 — opium, 298  
 — with oil of chamomile, 298  
 — pellitory, 340  
 — phosphorated, 309  
 — sal ammoniac, 95  
 — spermaceti, 155  
 — strychnine, 390  
 — subacetate of lead, 315  
 — succinate of ammonia, 97  
 — sulphate of quinine, 346  
 — sulphur and soap, 394  
 — sulphuret of calcium, 135  
 — carbon, 395  
 — sodium, 379  
 — sulphuric acid, 76  
 — tannate of lead, 318  
 — tartar emetic, ammoniacal, 108  
 — turpentine, 399  
 — acetic, 293  
 — oil, 293  
 — opiated, 293  
 — sulphuric, 293  
 — veratrine, 408  
 — verdigris, 186  
 — Wilkinson's, 95  
 Lip salve, 155  
 Liquorice root, 223  
 — extract, 223  
 — refined, 223  
 — lozenges, 224  
 — with opium, 224  
 — mixture, 224  
 — paste, 224  
 — syrup, 224  
 Lisbon diet drink, 363  
 Litharge, 317  
 Lithontriptics, 54  
 Liverwort, 231  
 Lobelia, 268  
 — extract, acetic, 268  
 — infusion, 268  
 — mixture, 268  
 — syrup, 268  
 — tincture, 268  
 — ethereal, 268  
 — vinegar, 268  
 Logwood, 228  
 — decoction, 228  
 — extract, 228  
 — electuary, 228  
 — mixture, 228  
 — infusion, 228  
 Long pepper, 310  
 — cataplasm, 310  
 Lotions, 469  
 Lotion, anti-rheumatic, 389  
 — Barlow's, 327  
 — Bateman's, 238  
 — capillary, 81  
 — for burns, 129  
 — chilblains, 292  
 — hiccough, 94  
 — toothache, 292  
 — Granville's, 90  
 — Goulard's, 100  
 — hydrosulphuretted, 73  
 — mammillary, 385  
 — mercurial, 237  
 — Saviard's, 329  
 — of acetate of lead, 315  
 — zinc, 413  
 — aconitine, 78  
 — almond, compound, 99  
 — alum, 89  
 — ammonia, 91  
 — borax, 380  
 — bromine, 130  
 — carbonate of potassa, 332  
 — soda, 382  
 — Cayenne pepper, 145  
 — cherry laurel water, 264  
 — chlorate of soda, 383  
 — chloride of lead, 316  
 — zinc, 411  
 — copper, 188  
 — corrosive sublimate, 237  
 — and camphor, 238  
 — copper, 238  
 — cosmetic, 237  
 — creasote, 182  
 — cyanuret of potassium, 320  
 — galls, 218  
 — hydrocyanic acid, 72  
 — iodide of potassium, compound, 324  
 — iodine, 251  
 — marygold, extract of, 137  
 — muriatic acid, 74  
 — mustard, 378  
 — nitrate of potassa, 336  
 — nitric acid, 74  
 — nux vomica, 289  
 — orpiment, 114  
 — oxide of zinc, 411  
 — Peruvian bark, 166  
 — sal ammoniac, 95  
 — sulphate of morphia, and borax, 283  
 — potassium, 327  
 — compound, 327  
 — soda, 385  
 — sodium, 379  
 — zinc, 414  
 — sulphuric ether, 80  
 — tartar emetic, 108  
 — and camphor, 108  
 — corrosive sublimate, 108  
 — tobacco, 396  
 — veratrin, 408  
 — verdigris, compound, 185  
 Lozenges, 462  
 — of anise, 103  
 — balsam of tolu, 400  
 — bicarbonate of soda, 381



- Lozenges of borax, 380  
 — catechu, 153  
 — Cayenne pepper, 145  
 — chalk, 136  
 — charcoal, 146  
 — chloride of gold and sodium, 121  
 — chlorinated lime, 134  
 — cinnamon, 167  
 — citric acid, 71  
 — croton oil, 294  
 — cubebs, 184  
 — — extract, 184  
 — emetine, 196  
 — ginger, 415  
 — gum, 66  
 — iodide of iron, 206  
 — — potassium, 321  
 — ipecacuanha, 254  
 — — and camphor, 254  
 — kermes mineral, 106  
 — lactate of iron, 208  
 — lactic acid, 73  
 — lactucarium, 263  
 — lemon juice, 265  
 — — oil, 265  
 — liquorice, 224  
 — — and opium, 224  
 — manna, 273  
 — marsh mallow, 86  
 — muriate of morphia, 282  
 — — and ipecacuanha, 282  
 — naphthaline, 258  
 — oxalate of potassa, 336  
 — oxalic acid, 75  
 — paullinia, 307  
 — pellitory, 340  
 — peppermint, 277  
 — rose, 352  
 — starch, 102  
 — tartaric acid, 77  
 — Tronchin's gum, 67  
 — vanilla, 407  
 — Wistar's, 224, 304  
 Lunar caustic, 111  
 Lupuline, 268  
 — emulsion, 268  
 — ointment, 269  
 — pills, 268  
 — powder, 268  
 — syrup, 268  
 — tincture, 268  
 Mace, 269  
 — balsam, neurine, 269  
 — drops, carminative, 269  
 — tincture, 269  
 Maceration, 452  
 Mackenzie's solution, 111  
 Madar, 138  
 Madder, 354  
 — decoction, 354  
 — powder, compound, 354  
 Magnesia, 269  
 — acetate, 270  
 — carbonate, 270  
 — — carminative, Dalby's, 270  
 — — Dewees', 270  
 — — mixture, 270  
 — — with camphor, 270  
 — — colchicum, 270  
 — — solution, 270  
 — citrate, 271  
 — — solution, 271  
 — — water, 271  
 — heavy, 269  
 — mixture, 269  
 Magnesia, mixture, and gentian, 270  
 — — rhubarb, 270  
 — — phosphate, 272  
 — — powder, with orange peel, 269  
 — — rhubarb, 269  
 — — sulphur, 269  
 — — sulphate, 271  
 — — Cheltenham salts, 271  
 — — clyster, 271  
 — — mixture, with coffee, 271  
 — — nitric acid, 272  
 — — Rochelle salt, 271  
 — — sulphuric acid, 271  
 — — tartar emetic, 271  
 — — sulphuret, 272  
 — — syrup, 272  
 Magnolia, 272  
 — tincture, 272  
 Maidenhair, 79  
 — infusion, 79  
 — syrup, 79  
 — — compound, 79  
 Male fern, 214  
 — bolus, 214  
 — electuary, 214  
 — extract, ethereal, 214  
 — — pills, 214  
 — — jelly, 214  
 — — mixture, 214  
 — — oil, 214  
 — — pills, 214  
 — — powder, 214  
 — — tincture of buds, 214  
 Mallow, common, 272  
 Malt, 272  
 — clyster, 272  
 — infusion, 272  
 Management of convalescence, 63  
 Manganese, 272  
 — carbonate, 272  
 — muriate, 272  
 — — pills, 272  
 — oxide, 273  
 — bolus, 273  
 — ointment, 273  
 — — compound, 273  
 — — powder, 273  
 — — sulphate, 273  
 Manna, 273  
 — emulsion, 273  
 — lozenges, 273  
 — mixture, 273  
 — syrup, 273  
 Mannite, 274  
 Marsh mallow, 86  
 — decoction, 86  
 — lozenges, 86  
 — ointment, 86  
 — paste, 86  
 — powder, 86  
 — syrup, 86  
 Marygold, 137  
 — extract, 137  
 — — lotion, 137  
 — — pills, 137  
 — — infusion, 137  
 — — ointment, 138  
 Masterwort, 231  
 Mastic, 275  
 — collutory, 275  
 — dentifrice, 275  
 — paste, anti-odontalgic, 275  
 — tincture, ethereal, 275  
 — varnish, crystal, 275



- Mastic, varnish, picture, 275  
 Matico, 275  
     — decoction, 275  
     — infusion, 275  
     — with senna, 275  
     — ointment, 276  
     — syrup, 275  
     — tincture, 275  
 May apple, 318  
     — extract, 318  
     — podophyllin, 318  
 Mayweed, 181  
     — cataplasm, 181  
     — infusion, 181  
 Meal, barley, 231  
     — flaxseed, compound, 267  
 Measures, 15  
     — approximative, 15  
     — drops, 17  
     — foreign, 19  
     — French, 19  
     — imperial, 15  
     — wine, 15  
 Medicated waters, 455  
 Mellites, 451  
 Mercury, 232  
     — acetate, 235  
         — liniment, 236  
         — pills, 236  
         — with opium, 236  
         — solution, 236  
     — ammoniated, 236  
         — ointment, 236  
     — and antimony, 232  
         — chalk, 233  
         — liniment, 233  
         — with ipecacuanha, 233  
     — gum, 233  
         — pills, with hemlock, 233  
         — magnesia, 234  
     — borate, 236  
     — bromide, 236  
         — bibromide, 236  
         — protobromide, 236  
         — solution, ethereal, 236  
     — calomel, 238  
     — cataplasm, 235  
     — cerate, 235  
         — compound, 235  
     — corrosive sublimate, 237  
     — cyanuret, 241  
         — gargle, 242  
         — ointment, 242  
         — pills, compound, 241  
         — solution, 241  
         — tincture, compound, 242  
     — gargle, 233  
     — iodide, 242  
         — ointment, 242  
         — pills, 242  
         — compound, 242  
         — powder, 242  
         — red, 243  
         — ointment, 243  
         — pills, 243  
         — tincture, 243  
         — ethereal, 243  
     — liniment, compound, 235  
         — opiated, 235  
     — lotion, 233  
     — mixture, 234  
     — mucilage, 232  
     — nitrate, 243  
         — liniment, 244  
         — ointment, 243  
         — with lead, 244  
         — pills, 243  
     — nitrate, protoxide, 243  
         — ointment, 234  
         — camphorated, 235  
         — compound, 235  
         — with belladonna, 235  
         — oxide, black, 244  
             — wash, 244  
             — ointment, 244  
             — pills, 244  
         — red, 244  
             — ointment, 244  
                 — brown, 245  
                 — with cinnabar, 245  
                 — lead, 245  
                 — sulphur, 245  
                 — tin, 245  
                 — zinc, 245  
         — pills, 244  
         — powder, 244  
         — phosphate, 245  
         — pills, 245  
         — pills, 234  
             — Abernethy's, 234  
             — and antimony, 233  
             — colocynth, 234  
             — jalap, 234  
             — quinine, 234  
                 — compound, 234  
             — rhubarb, 234  
                 — compound, 234  
         — plaster, 235  
             — with belladonna, 235  
         — powder, compound, 232  
         — purified, 232  
         — sulphate, 245  
             — ointment, 246  
             — powder, compound, 245  
             — yellow, 245  
         — sulphuret, black, 246  
             — pills, 246  
             — powder, 246  
             — red, 246  
                 — cerate, 247  
                 — ointment, 247  
                 — pills, 246  
                 — powder, 246  
                 — fumigating, 247  
         — tartrate, 247  
             — and potassa, 247  
 Method of displacement, 452  
 Mezereon, 278  
     — decoction, 278  
     — compound, 279  
     — extract, 279  
     — oil, 279  
     — ointment, 279  
 Milk, 260  
     — anodyne, 99  
     — arrowroot, 274  
     — artificial, 99  
         — asses', 420  
         — goats', 261  
     — cement, parolic, 261  
     — chocolate, 420  
     — coffee, 420  
     — mixture, with suet, 261  
         — soda water, 261  
     — powder, 261  
     — roses, 99, 129  
     — sago, 358  
     — sugar of, 261  
         — draught, 261  
         — with gum Arabic, 261  
         — Iceland moss, 261  
     — syrup, 261  
     — vanilla, 407  
     — whey alum, 260



- Milk whey, aromatic, 261  
     cremor tartar, 260  
     mustard, 260  
     simple, 260  
     tamarind, 260  
     vinegar, 260  
     wine, 261  
 Mineral, Kermes, 105  
 Mint, 277  
 Mixture, acetate of ammonia, 92  
     lead, 315  
     morphia, 281  
     potassa, 320  
     acetic ether, 79  
     aconite, 78  
     American hellebore, 409  
     ammonia, 91  
         and ether, 91  
     ammoniac, 97, 98  
         and nitric acid, 98  
         compound, 98  
     ammoniated alcohol, 94  
         iron, 201  
     angustura, 103  
     anisated ammoniated alcohol, 94  
     aniseed, 104  
     antacid, 136  
     anthelmintic, 308  
     anti-emetic, 153, 382  
     anti-epileptic, 111  
     anti-otitic, 299  
     antimonial wine and ammoniac, 109  
         bittersweet, 109  
         laudanum, 109  
     anodyne, Vicat's, 94  
     arseniate of potassa, 330  
     asparagus, diuretic, 116  
     assafetida, 117  
         and oxymel of squills, 117  
         tolu, 117  
         compound, 118  
     avens, 223  
     balsam of Peru, 285  
         tolu, 401  
         and almond emulsion, 401  
         belladonna, 401  
         copaiba, 401  
         morphia, 401  
         opium, 401  
     belladonna, 128  
     benzoic acid, 70  
     bicarbonate of potassa, 334  
     bichloride of platina, 313  
     biniodide of potassium, 324  
     bistort, 319  
     bitartrate of potassa, 339  
     bittersweet, 194  
     black drop, 301  
         hellebore, 230  
     boracic acid, 70  
     borax, 381  
     borotartrate of potassa, 331  
         and magnesia, 331  
     brown, 224, 303  
     buckbean, 278  
         diuretic, 278  
     butter of cocoa, 132  
     cabbage tree bark, 222  
     camphor, 139  
         and chloroform, 140  
         ether, 140  
         water, 140  
     carbonate of ammonia, 93  
         and ginger, 93  
         magnesia, 270  
     Mixture, carbonate of magnesia and camphor, 270  
         colchicum, 270  
         potassa, 332  
         soda and chamomile, 382  
         copaiba, 382  
         gentian, 382  
         ipecacuanha, 382  
         quassia, 382  
     carburet of iron, 204  
     carminative, 277  
     carrageen, 160  
     cascarilla, 150  
     cassia pulp, 150  
     castor oil and ether, 192  
     catechu and logwood, 153  
     caustic potassa, 329  
     celandine, 157  
     centaury, 154  
     chalk, 137  
         compound, 137  
     chamomile, 104  
     cherry laurel water, 264  
     chloride of barium, 125  
         calcium, 134  
         iron, 204  
     chlorine, 159  
     citrate of potassa and Peruvian bark, 335  
     cochineal and carbonate of potassa, 169  
         common salt, 169  
     cod liver oil, 290  
     colchicum, 171  
         and elaterium, 171  
         magnesia, 171  
         sulphate of magnesia, 171  
     tincture, 171  
         with foxglove, 171  
         vinegar, 172  
         with squills, 172  
         wine, 172  
     colocynth, 173  
         tincture, 173  
     columbo, 174  
         and cascarilla, 174  
         salep, 174  
     common salt and lemon juice, 379  
     contrayerva, 177  
     copaiba, 178  
     corrosive sublimate, 237  
     creasote, 181  
     croton oil, 294  
     cubebs, 184  
         oil and copaiba, 185  
     cyanuret of potassium, 320  
     dandelion, 398  
     Dippel's oil, 289  
     diuretic, 308  
     eggs, 305  
         and brandy, 305  
         wine, 305  
     elaterium, 194  
     elder, 359  
     elecampane, 250  
     emetine, 195  
     ergot, 196  
         extract, 197  
     errhine, 103  
     expectorant, 276  
         ferrocyanide of quinine, 343  
         zinc, 412  
     flaxseed, 267  
     foxglove and acetate potassa, 191  
         tartaric acid, 191  
         expectorant, 191  
         tincture, 191  
     galbanum, 217  
     gamboge, 219  
         and elatine, 219



- Mixture, gentian, 221  
     — and sulphuric acid, 221  
     — tincture, 221  
     — guaiacum, 227  
         — and bittersweet, 227  
         — camphor, 228  
         — copaiba, 228  
         — henbane, 228  
         — odontalgic, 228  
     — gum Arabic, 66  
         — Wendt's, 67  
     — hemlock and paregoric, 176  
     — henbane, 249  
         — and antimony, 249  
         — squill, 249  
     — hops, 232  
     — horseradish, 112  
     — hydrocyanic acid, 72  
     — hyponitrous ether, 80  
     — Iceland moss, 156  
     — Indian sarsaparilla, 230  
     — iodide of iron, 207  
         — and chloride, syrup of, 208  
         — potassium, 322  
         — and sarsaparilla, 322  
     — iodine and iodide of potassium, 251  
     — ipecacuanha, 254  
     — iron, aromatic, 200  
         — compound, 211  
     — jalap, 257  
         — tincture, 257  
     — Kermes mineral, 106  
     — lactate of quinine, 364  
     — lactucarium, 262  
         — extract, 263  
         — water, 263  
     — laudanum and asarabacca, 302  
         — soap, 302  
         — tartar emetic, 304  
     — lettuce and dandelion, 262  
     — liquorice, 224  
     — lobelia, 268  
     — logwood, extract, 228  
     — magnesia, 268  
         — and gentian, 270  
         — rhubarb, 270  
     — malate of iron, 209  
     — male fern, 214  
     — manna, 273  
     — mercurial, 234  
     — milk and suet, 261  
     — monesia, 279  
     — muriate of morphia, 282  
         — quinine, 244  
     — muriatic ether, 80  
     — myrrh, compound, 287  
         — Griffith's, 287  
     — narcotine, 288  
     — neutral, 334  
         — compound, 335  
     — nitrate of ammonia, 96  
         — potassa, 336  
     — nitric acid, 74  
     — nux vomica, 289  
     — odontalgic, 149  
     — oil of almonds, 101  
         — amber, 392  
         — anise, 104  
         — cajeput, 289  
         — juniper, 258  
         — mole plant, 199  
         — valerian, 406  
     — oleaginous, 291  
     — olive oil, 291  
     — opium and cinnamon water, 300  
         — lime water, 300  
     — syrup of poppies, 300
- Mixture, pectoral, 274  
     — Peruvian bark, 161, 166  
     — phosphate of soda, 384  
     — pomegranate root, extract, 225  
     — Prussian blue, 206  
     — quince seed, 189  
     — rhatany, 260  
     — rhubarb, 351  
     — rue, 355  
     — scammony, 366  
     — sedative, 301, 306  
     — semen contra, 115  
     — seneka, 372  
     — senna, 375  
     — soot, 216  
     — spermaceti, 155  
     — squill, syrup, 369  
         — extract, and rhubarb, 369  
         — oxymel and ammoniac, 370  
         — ipecacuanha, 370  
         — marsh mallow, 370  
         — valerian, 369  
     — vinegar, 370  
         — and ammoniac, 370  
     — starch and suet, 102  
     — stramonium, 388  
     — strychnine, 390  
     — subcarbonate of iron, 202  
     — succinate of ammonia, 96  
     — sugar of milk and gum Arabic, 261  
     — sulphate of iron, 212  
         — magnesia and coffee, 271  
         — nitric acid, 272  
         — Rochelle salt, 271  
         — sulphuric acid, 271  
         — tartar emetic, 271  
     — morphia, 283  
     — potassa, 337  
     — quinine, 346  
         — and coffee, 346  
     — zinc, 415  
     — sulphur, 394  
     — sulphuret of antimony, 106  
         — and lime, 106  
     — carbon, 395  
     — potassium, 328  
     — sodium, 379  
     — sulphuric ether and camphor, 81  
     — turpentine, 87  
     — sweet spirit of nitre, 80  
     — tannic acid, 76  
     — tartarized soda, 385  
     — tartrate of potassa, 338  
         — and ammonia, 339  
     — tobacco, 396  
     — turpentine, 403  
         — oil, 292  
         — and ether, 293  
     — uva ursi, 404  
     — valerian and ammonia, 405  
         — Hoffmann's anodyne, 406  
     — vinegar, 67  
         — and cardamoms, 67  
     — Virginia snakeroot, acetated, 376  
         — and allspice, 376  
     — wine of opium, 301  
     — wormseed oil, 157
- Mode of administration of medicine, 47  
 Molasses beer, 420  
     — posset, 419  
 Mole plant, 198  
     — mixture of oil, 199  
 Monesia, 279



- Monesia, mixture, 279  
 — ointment, 280  
 — purified, 279  
 — syrup, 279  
 — compound, 279  
 — tincture, 279  
 Monkshood, 17  
 Morphia, 280  
 — acetate, 280  
 — bolus, 281  
 — clyster, 281  
 — mixture, 281  
 — ointment, 281  
 — pills, 281  
 — powder, compound, 281  
 — solution, 281  
 — alcoholic, 281  
 — bimeconate, 281  
 — solution, 281  
 — citrate, 281  
 — draught, 280  
 — hydriodate, 282  
 — injection, 280  
 — muriate, 282  
 — lozenges, 282  
 — and ipecacuanha, 282  
 — mixture, 282  
 — solution, 282  
 — syrup, 282  
 — compound, 282  
 — nitrate, 282  
 — pills, 280  
 — phosphate, 282  
 — sulphate, 282  
 — bolus, 282  
 — lotion, 283  
 — mixture, 283  
 — pills, 282  
 — solution, 283  
 — syrup, 283  
 — tartrate, 283  
 Mucilage, gum Arabic, 66  
 — mercurial, 233  
 — rice, 304  
 — salep, 417  
 — sassafras pith, 365  
 — starch, 101  
 — tragacanth, 402  
 Mudar, 138  
 — infusion, 138  
 — oil, 138  
 — pills, 138  
 Mulberries, 280  
 — syrup, 280  
 — root, 280  
 Mulled wine, 419  
 Muriate of ammonia, 95  
 — cinchonine, 104  
 — manganese, 272  
 — morphia, 282  
 — narcotine, 288  
 — opium, 300  
 — quinine, 166  
 Muriatic acid, 73  
 — bath of, 74  
 — diluted, 73  
 — draught, 74  
 — gargle, 74  
 — gaseous, 73  
 — liniment, 74  
 — lotion, 74  
 — pediluvium, 74  
 Musk, 283  
 — artificial, 391  
 — emulsion, 392  
 — tincture, 391  
 — bolus, 283  
 Musk, clyster, 284  
 — mixture, 284  
 — pills, 283  
 — powder, 283  
 — tincture, 284  
 Mustard, 377  
 — cataplasm, 377  
 — compound, 377  
 — fomentation, 377  
 — foot-bath, 377  
 — gargle, 377  
 — liniment, 378  
 — lotion, 378  
 — oil, 378  
 — tincture, 378  
 — ointment, 378  
 — compound, 378  
 — whey, 377  
 Mutton tea, 418  
 — broth, 419  
 Myrrh, 286  
 — collutory, 286  
 — extract, compound, 286  
 — gargle, 287  
 — injection, 287  
 — mixture, compound, 287  
 — Griffith's, 287  
 — oil, 287  
 — pills and Canada balsam, 286  
 — squill, 286  
 — sulphate of iron, 286  
 — zinc, 286  
 — plaster, 287  
 — powder, emmenagogue, 286  
 — and ipecacuanha, 286  
 — solution, alkaline, 286  
 — tincture, 287  
 — and hellebore, 287  
 Naphthaline, 287  
 — lozenges, 288  
 — ointment, 287  
 — syrup, 287  
 Narcotics, 61  
 Narcotine, 288  
 — bolus, 288  
 — mixture, 288  
 — muriate, 288  
 Neutral mixture, 334  
 New Jersey tea, 154  
 — decoction, 154  
 Nicotina, 396  
 Nitrate of camphor, 141  
 — potassa, 335  
 — quinine, 344  
 — silver, 111  
 — veratrine, 408  
 Nitre, 335  
 — sweet spirit, 79  
 Nitric acid, 74  
 — collutory, 74  
 — diluted, 74  
 — fomentation, 74  
 — liniment, 74  
 — lotion, 74  
 — mixture, 74  
 — ointment, 74  
 Nitro-muriatic acid, 75  
 — bath, 75  
 Nutmeg, 284  
 — clyster, 284  
 — powder, anti-hectic, 284  
 — aromatic, 284  
 — spirit, 285  
 — compound, 285  
 Nux vomica, 288  
 — extract, 288



Nux vomica, extract, pills, 289

——— lotion, 289

——— mixture, 289

——— pills, 289

——— and aloes, 289

——— compound, 289

——— powder, 288

——— tincture, 289

——— compound, 289

Oak bark, 341

—— acorn coffee, 342

—— cataplasm, 342

—— confection, 342

—— decoction, 342

—— extract, 342

—— gargle, 342

—— powder, compound, 342

Oat, 123

—— cataplasm, 124

—— decoction, 123

—— flummery, 123

—— gruel, groat, 123

—— powder for cataplasm, 124

Oatmeal gruel, 123

Observations on management in the sick room, 45

Official preparations and directions, 446

Oil of almond, 100

—— bitter, 100

—— amber, 391

—— purified, 391

—— benzoin, 129

—— black pepper, resinous, 310

—— volatile, 310

—— box, 131

—— cajeput, 289

—— cantharides, 143

—— carraway, 149

—— castor, 291

—— chlorine, 159

—— cinnamon, 168

—— cloves, 149

—— cod liver, 290

—— copaiba, 179

—— croton, 293

—— cubebs, 185

—— dittany, 185

—— eggs, 305

—— elaterium, 195

—— ergot, 197

—— euphorbium, 199

—— fennel, 215

—— figwort, 371

—— ginger, 416

—— guaiacum, 227

—— compound, 227

—— hedge hyssop, 226

—— hemlock, 176

—— horsebean, 172

—— horsemint, 279

—— laurel, 263

—— macassar, 214

—— male fern, 214

—— mezereon, green, 279

—— mudar, 138

—— mustard, 378

—— myrrh, 287

—— olive, 290

—— orange flowers, 120

—— parsley, 308

—— partridge berry, 220

—— pennyroyal, 229

—— peppermint, 277

—— rose, 352

—— rosemary, 354

—— roses, 352

Oil of sassafras, 365

—— savine, 357

—— scammony, 366

—— soot, 216

—— stramonium, 389

—— tansy, 397

—— tobacco, 396

—— turpentine, 292

—— and ether, 293

—— honey, 292

—— clyster, 293

—— emulsion, 292

—— liniment, 293

—— acetic, 293

—— opiated, 293

—— sulphuric, 293

—— lotion, 292

—— for toothache, 292

—— mixture, 292

—— phosphorated, 309

—— purified, 292

—— soap, Starkey's, 293

—— wine, 293

—— valerian, 406

—— violets, 409

—— wormseed, 157

—— wormwood, 65

—— essential, 65

—— British, 307

—— Dippel's, 289

—— distilled, 456

—— essential, 556

—— fixed, 460

—— phosphorated, 308

—— volatile, 456

Ointments, 471

Ointment, aconitine, 78

—— ammoniacal, 93

—— Anglo-Saxon, 90

—— anodyne, 299

—— arsenical, 69

—— Banyer's, 317

—— basilicon, 348

—— camphor, 140

—— Egyptian, 186

—— for chilblains, 90

—— issue, 144

—— itch, 362

—— laudanum, 303

—— mercurial, 234

—— camphorated, 235

—— compound, 235

—— and belladonna, 235

—— naphthaline, 287

—— phosphorated, 309

—— pile, 90, 314

—— Rust's, 90

—— saffron, 183

—— savine, 357

—— soot, 216

—— compound, 216

—— spermaceti, 155

—— and rose water, 155

—— sulphuric acid, 76

—— sultana, 155

—— of acetate of lead, 314

—— morphia, 281

—— aconite, 78

—— ammoniated, 78

—— aloes, 86

—— American hellebore, 409

—— Gondret's, 93

—— ammoniated copper, 187

—— animal charcoal, 146

—— arseniate of iron, 201

—— azedarach, 124

—— balsam of Peru, 286



- Ointment of balsam of Peru, compound, 286  
     belladonna, 128  
     bichloride of platina, 313  
     biniodide of potassium, 325  
         and opium, 325  
     black hellebore, 230  
         oxide of mercury, 244  
         pepper, 310  
         pitch, 312  
     borax, 381  
     bromide of iron, 202  
         potassium, 319  
         compound, 319  
     calomel, 241  
         and acetate of copper, 241  
         alum, 241  
         camphor, 241  
         squill, 241  
     cantharides, 143  
     carbonate of ammonia, 93  
         lead, 316  
         camphorated, 316  
         potassa, 333  
         soda, 382  
     carburet of iron, 204  
     carrot, 148  
     cevadilla, 356  
         extract, 356  
     charcoal, 146  
     cherry laurel, 264  
     chloride of antimony, 145  
         gold, 121  
         and sodium, 122  
         lead, 316  
     chlorinated lime, 134  
     chlorine, 159  
     cocculus indicus, 168  
     cod liver oil, 290  
         compound, 290  
     common salt, 379  
         compound, 379  
     creasote, 182  
         compound, 182  
     croton oil, 295  
     cyanide of silver, 110  
     cyanuret of mercury, 242  
         potassium, 321  
     delphinia, 189  
     elder flowers, 359  
         leaves, 359  
     elemi, 195  
     fennel, 215  
     figwort, 371  
     foxglove, 191  
     fuligokali, 216  
     galls, 219  
         compound, 219  
         extract, 218  
     glycerin, 223  
     gold, 120  
     hemlock, 175  
     henbane, 249  
         compound, 249  
     hops, 232  
     hydrargyro-iodide of potassium, 325  
     hydriodate of ammonia, 94  
     iodide of arsenic, 113  
         barium, 125  
         carbon, 147  
         iron, 208  
         lead, 317  
         mercury, 242  
         potassium, 322  
         and opium, 322  
         mercury, 323  
         morphia, 323  
         silver, 111  
     Ointment of iodide of sulphur, 395  
         zinc, 413  
         iodine, 252  
         compound, 252  
         with mercury, 252  
         oil of tobacco, 252  
     laurel, 263  
     lavender, 265  
     lead, 317  
         compound, 317  
     lime, 136  
     lupulin, 269  
     marygold, 138  
     marsh mallow, 86  
     mezereon, 279  
     monesia, 280  
     mustard, 378  
         compound, 378  
     nitrate of bismuth, 130  
         mercury, 243  
         silver, 112  
         compound, 112  
         strychnine, 391  
     nitric acid, 74  
     opium and tar, 299  
     oxide of manganese, 273  
         and sulphur, 273  
         silver, 111  
     petroleum, 307  
     picrotoxin, 168  
     poke, 309  
     Prussian blue, 206  
     red iodide of mercury, 243  
         oxide of mercury, 244  
         and basilicon, 245  
         cinnabar, 245  
         lead, 245  
         sulphur, 245  
         tin, 245  
         zinc, 245  
     sulphuret of mercury, 247,  
     rhatany, compound, 260  
     rose water, 352  
     squill, 370  
         compound, 370  
     stavesacre, 388  
     stramonium, 389  
     strychnine, 390  
     subacetate of lead, 315  
     sulphate of cadmium, 133  
         copper, 188  
         mercury, 246  
         quinine, 347  
     sulphur, 394  
         and camphor, 394  
         soap, 394  
         zinc, 394  
         compound, 394  
     sulphuret of antimony, 105  
         potassium, 328  
         sodium, 379  
     tannate of lead, 318  
     tannic acid, 76  
     tar, 312  
         compound, 312  
     tartar emetic, 108  
         compound, 108  
     tin, 387  
     tobacco, 396  
     turpentine, 399  
         compound, 399  
     veratrin, 408  
         and iodine, 408  
         opium, 408  
     verdigris, 186  
     white hellebore, 409  
         compound, 409



- Ointment of white precipitate, 236  
     — wild indigo, 124  
     — willow leaves, 358  
     — zinc, oxide, 411  
     — and calomel, 411  
     — lycopodium, 411  
     — opium, 411  
 Olive oil, 290  
     — clyster, 291  
     — mixture, 291  
 Opiate, anti-dysenteric, 298  
 Opium, 295  
     — balsam, 298  
     — cerate, 299  
     — clyster, 299  
     — collyrium, 299  
     — confection, 298  
     — draught, anodyne, 303  
     — drops, 298  
     — elixir, Lettsom's, 303  
     — extract, acetous, 295  
         — alcoholic, 295  
         — aqueous, 295  
         — aqueo-alcoholic, 295  
         — denarcotized, 296  
         — fermented, 296  
         — roasted, 296  
         — vinous, 295  
     — Godfrey's cordial, 302  
     — fomentation, 299  
     — injection, 299  
     — linctus, 299, 303  
     — liniment, 298  
         — with oil of chamomile, 298  
     — lozenges, Wistar's, 304  
     — mass, anti-odontalgic, 298  
     — mixture, anti-otitic, 299  
         — brown cough, 303  
         — with cinnamon water, 300  
         — lime water, 300  
         — syrup of poppies, 300  
         — sedative, 301  
     — muriate, 300  
     — ointment, 299  
         — and tar, 299  
     — pills, 296  
         — aromatic, 297  
         — with acetate of lead, 297  
             — mercury, 297  
             — antimony, 296  
             — butter of cocoa, 297  
             — camphor, 297  
             — castor, 297  
             — foxglove, 297  
             — hemlock, 297  
             — and henbane, 296  
             — liquorice, 297  
             — musk, 297  
             — nitrate of silver, 297  
             — sulphate of zinc, 297  
             — sulphuret antimony, 297  
     — plaster, 298  
         — with camphor, 298  
     — powder, compound, with chalk, 296  
         — roasted, 296  
         — with antimony, 296  
             — camphor, 296  
             — musk, 296  
             — nitre, 296  
             — sulphur, 296  
     — suppository, 299  
     — syrup, 299  
         — succinated, 299  
     — tincture, 301  
         — acetated, 301  
         — ammoniated, 302  
         — and asarabacca, 302  
     — Opium tincture and soap, 302  
         — tartar emetic, 304  
         — Bateman's, 302  
         — Battley's, 302  
         — camphorated, 302  
         — clyster, 303  
         — compound, 302  
         — liniment, 303  
         — and lime water, 303  
         — ointment, 303  
         — Smith's, 302  
         — succinated, 302  
         — Swediaur's, 302  
         — Warner's, 303  
         — Guy's hospital, 300  
         — Houlton's, 300  
         — Lancaster, 300  
         — mixture, 301  
         — vinegar, or black drop, 300  
         — Porter's, 301  
         — Rousseau's, 300  
         — water, 300  
         — wine, 301  
         — collyrium, 301  
         — drops, odontalgic, 301  
         — fomentation, 301  
         — mixture, 301  
         — Sydenham's, 301  
 Opodeldoc, liquid, 91  
     — Steer's, 91  
 Opoponax, 304  
     — emulsion, 304  
     — tincture, 304  
         — compound, 304  
 Orange, 118  
     — flowers, 120  
         — confection, 120  
         — oil, 120  
         — pastilles, 120  
         — syrup, 120  
         — water, 120  
     — leaves, 119  
         — electuary, 119  
         — jelly, 119  
     — peel, 119  
         — confection, 119  
         — electuary, 119  
         — elixir, 119  
         — infusion, 119  
         — compound, 119  
         — powder, with rhubarb, 119  
         — spirit, 119  
         — syrup, 119  
         — tincture, 119  
 Orgeat, syrup of, 99  
 Origanum, 304  
 Orpiment, 114  
     — gargle, 114  
     — lotion, 114  
     — powder, 114  
 Orris root, 255  
     — dentifrice, 255  
 Oxalic acid, 75  
     — lozenges, 75  
 Oxide, antimony, nitro-muriatic, 105  
     — gold, 122  
     — manganese, 273  
     — silver, 111  
     — zinc, 410  
 Oxymel, 276  
     — bitartrate of potassa, 339  
     — colchicum, 171  
     — elecampane, compound, 250  
     — squill, 370  
 Oyster-shell, prepared, 136  
 Panada, 417



- Panada, chicken, 419  
 Paregoric elixir, 302  
 Pareira brava, 306  
     — decoction, 307  
     — extract, 307  
     — infusion, 306  
     — tincture, 307  
 Parsley, 308  
     — infusion of, 308  
     — oil of, 308  
 Partridge-berry, 220  
     — oil, 220  
     — water, 220  
 Pastes, 462  
 Paste, almond, 99  
     — chloride of zinc, 411  
     — cocoa, compound, 132  
     — copaiba, 179  
     — de guimauve, 66  
     — fig, 214  
     — liquorice, 224  
     — marshmallow, 86  
     — odontalgic, 275  
     — pectoral, of gum Arabic, 66  
     — phosphorus, 309  
     — tragacanth, 402  
     — rubefacient, 314  
     — Ward's, 310  
 Pastilles, catechu, 153  
     — fumigating, 128  
     — orange flowers, 120  
     — purgative, 366  
     — vinegar, 67  
 Paullinia, 307  
     — chocolate, 307  
     — extract, alcoholic, 307  
     — lozenges, 307  
     — pills, 307  
     — syrup, 307  
 Pediluvium, mustard, 377  
     — muriatic acid, 74  
 Pellitory, 340  
     — extract, 340  
     — liniment, 340  
     — lozenges, 340  
     — plaster, 340  
     — tincture, 340  
         — compound, 340  
 Pennyroyal, 229  
     — infusion, 229  
     — oil, 229  
 Pepper, black, 310  
     — cubebs, 183  
     — Jamaica, 310  
     — white, 310  
 Peppermint, 277  
     — essence, 277  
     — lozenges, 277  
     — mixture, 277  
     — oil, 277  
 Persimmon, 193  
     — infusion of bark, 193  
     — wine of fruit, 193  
 Persulphate of mercury, 245  
 Peruvian bark, 160  
     — beer, 165  
     — cataplasm, 165  
     — cerate, 165  
     — clyster, 165  
     — decoction, 162  
     — compound, 163  
     — mixture, 163  
     — with cascarilla, 163  
     — rhatany, 163  
     — electuary, 164  
     — astringent, 164  
     — with catechu, 164  
     — Peruvian bark, electuary, with cloves, 164  
         — iron, 164  
         — sulphur, 164  
         — tin, 164  
     — extract, 162  
         — bolus, 162  
         — pills, 162  
         — vinous, 162  
     — gargle, 165  
     — gelatine, 165  
     — infusion, 163  
     — compound, 164  
         — with iron, 164  
         — with limewater, 163  
         — magnesia, 163  
         — snakeroot, 163  
         — valerian, 163  
     — lotion, 165  
     — mixture, 161, 166  
     — pomatum, 165  
     — powder, compound, 160  
         — dentifrice, 161  
         — with arnica, 161  
         — camphor, 161  
         — cascarilla, 161  
         — isinglass, 161  
         — valerian, 161  
     — syrup, 166  
         — compound, 165  
         — tincture, 164  
         — compound, 164  
         — with ammonia, 165  
         — cantharides, 165  
         — gentian, 165  
         — snakeroot, 165  
         — valerian, 164  
     — wine, 166  
         — and calamus, 166  
         — iron, 166  
 Petroleum, 307  
     — anthelmintic mixture, 308  
     — British oil, 307  
     — diuretic mixture, 308  
     — embrocation, 307  
     — ointment, 307  
     — plaster, 308  
 Pharmaceutical names, table of, 433  
 Phloridzine, 308  
     — powder of, 308  
 Phosphate of ammonia, 96  
     — iron, 210  
     — lime, 137  
     — mercury, 245  
     — quinine, 344  
     — soda, 384  
 Phosphoric acid, 75  
     — diluted, 75  
     — fomentation, 75  
     — lemonade, 75  
     — pills, 75  
     — tincture of myrrh, 75  
 Phosphorated ether, 308  
     — liniment, 309  
     — oil, 308  
     — turpentine, 309  
     — ointment, 309  
 Phosphorus, 308  
     — cerate, 309  
     — paste, 309  
 Picrotoxin, 168  
 Pills, 447  
     — Abernethy's, 234  
     — acetate of copper, 185  
     — lead, 313  
     — mercury, 236  
     — and opium, 236  
     — morphia, 281



- Pills, acetate of potassa, compound, 329  
     — soda, compound, 379  
     — aconite, extract, 77  
     — agaric and opium, 134  
     — aloes, 82  
         — and assafetida, 82  
         — blue mass, 84  
         — iron, 82  
         — myrrh, 82  
         — rhubarb, 84  
     — Anderson's, 83  
     — antichlorotic, 83  
     — aperient, 84  
     — Barthez's, 84  
     — Bicker's, 84  
     — Chapman's, 82  
     — compound, 82  
     — Duchesne's, 83  
     — Frank's, 84  
     — Fuller's, 83  
     — Griffiths', 84  
     — James's analeptic, 83  
     — Hooper's, 83, 211  
     — Lady Webster's, 83  
     — Mitchell's, 82  
     — Morrison's, 83  
     — Peters', 83  
     — Pitschaft's eccoprotic, 84  
     — Rufus', 82  
     — Speediman's, 83, 349  
     — splenitic, 83  
     — Whytt's, 84  
     — alum, 88  
         — and benzoin, 88  
         — astringent, 88  
     — American hellebore, 409  
     — ammoniac, 97  
         — and rhubarb, 97  
         — compound, 97  
         — De Haen's, 97  
         — Klein's, 97  
     — ammoniated copper, 187  
         — compound, 187  
         — gold, 123  
         — iron, 201  
     — anemone, 102  
     — antimonial powder and calomel, 109  
     — arseniate of iron, 201  
     — arsenic and opium, 70  
         — pepper, 69  
     — arsenical, 69  
     — assafetida, 116  
         — and aloes, 48  
         — lactucarium, 116  
         — iron, 117  
         — musk, 117  
         — opium, 117  
     — Bacher's, 230  
     — balsam of Peru, 285  
     — belladonna, compound, 127  
     — bichloride of platina, 313  
     — bittersweet, extract, 194  
     — black hellebore, 229  
         — extract of, 230  
     — black oxide of mercury, 244  
     — Bland's antichlorotic, 211  
     — bloodroot, 360  
     — blue, or mercurial, 234  
         — and colocynth, 234  
         — jalap, 234  
         — quinine, 234  
         — rhubarb, 234  
         — compound, 234  
     — borotartarate of potassa, compound, 331  
     — bromide of iron, 201  
     — brucia, 131  
     — buckbean, compound, 278  
     — Pills, calomel, 239  
         — and acetate of lead, 239  
         — antimony, 240  
         — catechu, 240  
         — colocynth, 240  
         — dandelion, 240  
         — guaiacum, 240  
         — iron, 240  
         — jalap, 240  
         — opium, 240  
         — quinine, 239  
         — squill, 240  
         — compound, 240  
     — camphor, 138  
         — compound, 138  
         — and lactucarium, 138  
         — musk, 138  
     — cantharides and capsicum, 142  
         — iron, 142  
     — carbonate of ammonia, 93  
         — iron, 202  
         — compound, 203  
         — soda, 381  
     — carburet of iron, 203  
     — castor, 151  
         — and succinic acid, 151  
     — catechu and alum, 152  
         — liquorice, 152  
     — cathartic, compound, 240  
     — Cayenne pepper, 145  
     — celandine, 157  
     — centaury, extract, 154  
     — chamomile, compound, 144  
     — chloride of barium, 125  
         — calcium, 134  
         — gold, 121  
         — and sodium, 121  
         — mercury and quinine, 241, 343  
         — silver and ammonia, 110  
     — colchicum, 170  
         — compound, 170  
     — colocynth, compound, 173  
         — and henbane, 173  
     — columbo, compound, 174  
     — copaiba, 177  
         — and cubebs, 177  
         — and turpentine, 178  
     — corrosive sublimate, 237  
         — compound, 237  
     — creasote, 181  
     — croton oil, 293  
         — compound, 294  
         — and blue mass, 294  
         — quinine, 294  
     — cyanide of gold, 122  
         — zinc, 412  
     — cyanuret of mercury, compound, 241  
         — potassium, 320  
     — dandelion, 398  
         — and blue mass, 398  
     — digitalin, 192  
     — dogwood, round-leaved, 181  
     — elaterium, 194  
     — elecampane, compound, 250  
     — ergot, extract of, 197  
         — compound, 197  
     — ferrocyanate of quinine, 343  
         — zinc, compound, 412  
     — foxglove, 190  
     — fuligokali, sulphuretted, 216  
     — galbanum, compound, 217  
     — gallic acid, 71  
     — gamboge, compound, 219  
     — gentian, compound, 220  
         — and ammonia, 220  
     — gentianin, 222



- Pills, guaiacum and aloes, 227  
     — antimony, 227  
     — sulphur, 227  
     — turpentine, 227  
     — hemlock and calomel, 175  
     — dandelion, 175  
     — ipecacuanha, 175  
     — henbane, compound, 248  
     — and ipecacuanha, 248  
     — opium, 248  
     — hydrargyro-iodide of potassium, 325  
     — iodide of arsenic, 113  
     — calcium, 135  
     — iron, 207  
     — lead, 317  
     — mercury, 242  
     — compound, 242  
     — potassium, 321  
     — iodine, 251  
     — indigo, 250  
     — ipecacuanha and centaury, 254  
     — foxglove, 254  
     — opium, 253  
     — compound, 254  
     — iron, prepared, 200  
     — jalap and calomel, 256  
     — compound, 256  
     — Keyser's, 236  
     — kinate of quinine, 344  
     — kino, compound, 259  
     — lactate of iron, 208  
     — quinine, 344  
     — lactucarium, 262  
     — Lockstadt's, 101  
     — lupulin, 268  
     — male fern, extract, 214  
     — oil, 214  
     — marygold, extract, 137  
     — mercury and antimony, 233  
     — gum, with hemlock, 233  
     — morphia, 280  
     — mudar, 138  
     — muriate of manganese, 272  
     — musk, 283  
     — myrrh and Canada balsam, 286  
     — squill, 286  
     — sulphate of iron, 286  
     — zinc, 286  
     — nitrate of bismuth, 130  
     — potassa, 335  
     — and camphor, 335  
     — silver, 111  
     — nux vomica, 289  
     — and aloes, 289  
     — compound, 289  
     — extract, 289  
     — compound, 289  
     — opium, 296  
     — and acetate of lead, 297  
     — mercury, 297  
     — butter of cacao, 297  
     — camphor, 297  
     — castor, 297  
     — foxglove, 297  
     — hemlock, 297  
     — henbane, 296  
     — liquorice, 297  
     — musk, 297  
     — nitrate of silver, 297  
     — sulphate of zinc, 297  
     — sulphuret antimony, 297  
     — aromatic, 297  
     — oxide, black, of iron, 210  
     — gold, 123  
     — red, of iron, 210  
     — zinc, 410  
     — compound, 410
- Pills, paullinia, 307  
     — Peruvian bark, extract, 162  
     — phosphate of mercury, 245  
     — phosphoric acid, 75  
     — piperine, 311  
     — and mercury, 311  
     — pipsissewa, 158  
     — Plummer's, 240  
     — poison oak, 402  
     — proto-nitrate of iron, 207  
     — mercury, 243  
     — Prussian blue, 206  
     — red iodide of mercury, 243  
     — oxide of mercury, 244  
     — rhatany, 259  
     — rhubarb, 349  
     — compound, 349  
     — and caraway, 349  
     — chamomile, 349  
     — ipecacuanha, 349  
     — ox-gall, 349  
     — soda, 349  
     — saffron, 183  
     — sagapenum, compound, 357  
     — salicine, 358  
     — compound, 358  
     — savine, 356  
     — scammony, 366  
     — and ox-gall, 366  
     — compound, 366  
     — seneka, 371  
     — senna, 373  
     — soap, 361  
     — and ox-gall, 361  
     — compound, 361  
     — soda and ipecacuanha, 381  
     — rhubarb, 381  
     — soot, 215  
     — squill and ammoniac, 367  
     — calomel, 367  
     — croton oil, 367  
     — ipecacuanha, 367  
     — henbane, 368  
     — compound, 368  
     — stimulant, 390  
     — storax, 389  
     — stramonium, 388  
     — compound, 388  
     — strychnine, 390  
     — compound, 390  
     — sulphate of cinchonia, 166  
     — copper, 187  
     — and opium, 187  
     — compound, 188  
     — iron, 211  
     — and rhubarb, 211  
     — compound, 211  
     — morphia, 282  
     — potassa, 337  
     — quinine, 345  
     — compound, 346  
     — with gentian, 346  
     — soda, compound, 384  
     — zinc, 414  
     — compound, 414  
     — with myrrh, 414  
     — turpentine, 414  
     — sulphuret of antimony, 105  
     — and lime, 106  
     — calcium, compound, 135  
     — mercury, black, 246  
     — red, 246  
     — potassium, 326  
     — and charcoal, 326  
     — compound, 326  
     — tin, compound, 388  
     — tannic acid, 76



- Pills, tannic acid, and opium, 76  
     — tansy, 397  
     — tar, 392  
     — tartar emetic and camphor, 107  
     —                      guaiacum, 107  
     —                      opium, 107  
     — tobacco, 396  
     — turpentine, 399  
     —                      and guaiacum, 399  
     —                      jalap, 399  
     —                      magnesia, 399  
     —                      myrrh, 399  
     —                      rhubarb, 399  
     — valerian, compound, 405  
     — valerianate of zinc, 415  
     — veratrin, 407  
     —                      and henbane, 408  
     — Virginia snakeroot, compound, 376  
 Pink, 190  
     — syrup, 190  
 Pinkroot, 386  
     — and calomel, 386  
     —                      savine, 386  
     — decoction, 387  
     — extract, 387  
     —                      fluid, 386  
     —                      compound, 386  
     — infusion, compound, 386  
 Piperine, 311  
     — pills of, 311  
     —                      and mercury, 311  
     — tincture of, 311  
 Pipsissewa, 157  
     — beer, 158  
     — decoction, 157  
     —                      compound, 157  
     — extract, 157  
     — pills, 158  
     — syrup, 158  
 Pitch, black, 312  
     — ointment of, 312  
     — plaster of, 313  
     — Burgundy, 311  
     — Canada, 312  
     — cerate, 312  
     — hemlock, 312  
     — pills of, 312  
     — plaster, 311  
     —                      aromatic, 311  
     —                      compound, 311  
 Plasters, 472  
 Plaster, acetate of lead, 314  
     — adhesive, 348  
     —                      Baynton's, 317  
     — ammonia, 91  
     — ammoniac, 98  
     —                      with hemlock, 98  
     —                      mercury, 98  
     — assafetida, 118  
     — balsam of Peru, 286  
     — belladonna, 128  
     — biniodide of potassium, 325  
     — black pepper, 310  
     —                      pitch, 313  
     — cantharides, camphorated, 144  
     —                      compound, 144  
     —                      odontalgic, 144  
     — carbonate of ammonia, 93  
     —                      lead, 316  
     —                      zinc, 414  
     — carburet of iron, 204  
     — cinnamon, spice, 168  
     — clove, 149  
     — court, 249  
     —                      blistering, 144  
     — croton oil, 295  
     — cumin, 189  
 Plaster, diapalma, 317  
     — elemi, cautery, 195  
     — euphorbium, 199  
     — foxglove, 191  
     — galbanum, 217  
     —                      compound, 217  
     — Gauthier's, 317  
     — gum, 98  
     — hemlock, 176  
     —                      compound, 176  
     — iodine, 252  
     — iron, 203  
     — lead, 317  
     — mercurial, 235  
     —                      and belladonna, 235  
     — myrrh, 287  
     — Nuremberg, 318  
     — opium, 298  
     —                      and camphor, 298  
     — pellitory, 340  
     — petroleum, 308  
     — pitch, 311  
     —                      aromatic, 311  
     —                      compound, 311  
     — red oxide of iron, 210  
     — resin, 348  
     — spice, 416  
     — St. Andrew's, 348  
     — subacetate of lead, 315  
     — sulphate of quinine, 347  
     — sulphuret of antimony and lime, 107  
     —                      potassium, compound, 328  
     — tartar emetic, 108  
     — turpentine, 399  
     —                      compound, 400  
     — verdigris, 186  
 Platina, 313  
     — bichloride of, 313  
     — mixture of, 313  
     — ointment of, 313  
     — pills of, 313  
 Pleurisy root, 115  
     — infusion, 115  
 Plummer's pills, 240  
 Podophyllin, 318  
 Poids de Marc, 12  
 Poison, extract, 402  
     — oak, 402  
     — pills, 402  
     — tincture, 402  
 Poisons, 479  
 Poke, 309  
     — ointment of, 309  
 Polygalic acid, 372  
 Pomatum, cantharides, 144  
 Pomegranate, 225  
     — bolus of root, 226  
     — decoction of rind, 225  
     —                      root, 225  
     — extract of root, 225  
     —                      electuary, 225  
     —                      mixture, 225  
     — gargle of rind, 225  
     — injection of rind, 225  
 Poppy heads, 306  
     — decoction, 306  
     — injection, sedative, 306  
     — mixture, sedative, 306  
     — syrup, 306  
     —                      pectoral, 306  
     — red, 357  
 Portland powder, 154  
 Posological table, 426  
 Posset, molasses, 419  
 Potassa, 328  
     — acetate, 329  
     — scillitic, 329



Potassa, acetate, bolus, 329  
 — conserve, 329  
 — liquid, 330  
 — mixture, 330  
 — pills, compound, 329  
 — tincture, 330  
 — arseniate, 330  
 — mixture, 330  
 — solution, 330  
 — bicarbonate, 333  
 — effervescing powders, 333  
 — solution, 333  
 — mixture, 334  
 — bisulphate, 337  
 — disinfecting powder, 337  
 — effervescing draught, 337  
 — bitartrate, 338  
 — dentifrice, 338  
 — electuary, 339  
 — imperial, 339  
 — mixture, 339  
 — oxymel, 339  
 — powder, 338  
 — solution, 339  
 — compound, 339  
 — borate, 330  
 — borotartrate, 331  
 — and magnesia, 331  
 — mixture, 331  
 — pills, compound, 331  
 — powder, 331  
 — solution, 331  
 — carbonate, 331  
 — and ammonia, 334  
 — collutory, 332  
 — collyrium, 332  
 — emulsion, 332  
 — fomentation, 332  
 — liniment, 332  
 — lotion, 332  
 — mixture, 332  
 — ointment, 333  
 — powder, 332  
 — pure, 332  
 — solution, 332  
 — tincture, 332  
 — compound, 333  
 — caustic, 328  
 — collyrium, 329  
 — injection, 329  
 — lotion, stimulant, 329  
 — mixture, 329  
 — solution of, 328  
 — lithontriptic, 329  
 — chlorate, 334  
 — powder, 334  
 — compound, 334  
 — solution, 334  
 — citrate, 334  
 — mixture, 335  
 — solution, 334  
 — compound, 335  
 — iodate, 335  
 — syrup, 335  
 — nitrate, 335  
 — emulsion, 336  
 — fused, 335  
 — gargle, 336  
 — lotion, 336  
 — mixture, 336  
 — pills, 335  
 — with camphor, 335  
 — powder, compound, 335  
 — with camphor, 335  
 — orris root, 335  
 — squill, 335

Potassa, nitrate, purified, 335  
 — oxalate, 336  
 — lozenges, 336  
 — powder, 336  
 — silicate, 336  
 — soluble glass, 337  
 — solution, 337  
 — sulphate, 337  
 — and ammonia, 337  
 — magnesia, 337  
 — sulphur, 338  
 — mixture, 337  
 — pills, 337  
 — powder, compound, 337  
 — and rhubarb, 337  
 — tartrate, 338  
 — and ammonia, 339  
 — mixture, 339  
 — solution, 339  
 — mixture, 338  
 — powder, and rhubarb, 338  
 — with lime, 328  
 — solution of, 328  
 Potassium, 319  
 — biniodide of, 323  
 — collyrium, 324  
 — ioduretted bath, 324  
 — water, 323  
 — Lu-  
 gol's, 323  
 — lotion, compound, 324  
 — mixture, 324  
 — ointment, 325  
 — and opium, 325  
 — plaster, 325  
 — solution, compound, 324  
 — caustic, 324  
 — syrup, 324  
 — tincture, compound, 324  
 — bromide of, 319  
 — ointment of, 319  
 — compound, 319  
 — solution of, 319  
 — chloride of, 319  
 — solution of, 320  
 — cyanuret of, 320  
 — lotion of, 320  
 — mixture of, 320  
 — ointment of, 321  
 — pills of, 320  
 — solution of, 320  
 — syrup of, 320  
 — hydrargyro-iodide of, 325  
 — ointment, 325  
 — pills, 325  
 — solution, 325  
 — tincture, 325  
 — iodide of, 321  
 — and sarsaparilla, 322  
 — injection of, 322  
 — liniment of, 323  
 — lozenges of, 321  
 — mixture of, 322  
 — ointment of, 322  
 — and mercury, 323  
 — morphia, 323  
 — opium, 322  
 — pills of, 321  
 — soap of, 323  
 — solution of, 321  
 — syrup of, 322  
 — sulpho-cyanuret of, 326  
 — sulphuret of, 326  
 — and cremor tartar, 326  
 — bath, 327  
 — bolus, 326  
 — electuary, 326



- Potassium, sulphuret of, ferro, 326  
 ————— hyposulphited, 326  
 ————— syrup, 327  
 ————— injection, 327  
 ————— lotion, 327  
 ————— compound, 327  
 ————— mixture, 328  
 ————— ointment, 328  
 ————— pills, 326  
 ————— compound, 326  
 ————— and charcoal, 326  
 ————— plaster, compound, 328  
 ————— soap, 328  
 ————— solution, 327  
 ————— syrup, 327  
 ————— compound, 327  
 ————— tincture, 327  
 ————— water, 327  
 Potion of carbonate of ammonia, 92  
 Poultices, 468  
 Powders, 446  
 ————— compound, 446  
 ————— simple, 446  
 Powder, acetate of lead, 313  
 ————— morphia, compound, 281  
 ————— aconite, compound, 77  
 ————— agaric, 130  
 ————— and opium, 130  
 ————— almond, 99  
 ————— aloes, 82  
 ————— and canella, 82  
 ————— compound, 82  
 ————— emmenagogue, 82  
 ————— alum, 87  
 ————— escharotic, 87  
 ————— ammoniated copper and belladonna, 187  
 ————— iron, 201  
 ————— animal charcoal, 145  
 ————— anthracokali, 147  
 ————— compound, 147  
 ————— antihectic, 284  
 ————— arnica, compound, 112  
 ————— aromatic, 147, 284  
 ————— arsenical, 69  
 ————— arsenic, compound, 69  
 ————— asarabacca, 115  
 ————— avens, compound, 223  
 ————— belladonna, 126  
 ————— and rhubarb, 127  
 ————— benzoic acid and ipecacuanha, 70  
 ————— benzoin, 128  
 ————— bicarbonate of potassa, 333  
 ————— bismuth, compound, 129  
 ————— bitartrate of potassa, 338  
 ————— black oxide of iron, 209  
 ————— sulphuret of mercury, 246  
 ————— borax, compound, 380  
 ————— borotartrate of potassa, 331  
 ————— burnt sponge, 387  
 ————— compound, 387  
 ————— calomel and antimony, 238  
 ————— jalap, 239  
 ————— foxglove, 239  
 ————— nitrate of potassa, 239  
 ————— opium, 239  
 ————— gamboge, 239  
 ————— pinkroot, 239  
 ————— antimony and henbane, 239  
 ————— jalap and rhubarb, 239  
 ————— camphor, 138  
 ————— compound, 138  
 ————— cantharides and camphor, 142  
 ————— savine, 141  
 ————— capuchin, 356  
 ————— carbonate of lead, camphorated, 316  
 ————— potassa, 332  
 Powder, carbonate of soda, effervescing, 381  
 ————— and magnesia, 381  
 ————— mercury, 381  
 ————— rhubarb, 381  
 ————— zinc, 414  
 ————— carburet of iron, 203  
 ————— cascarrilla, compound, 150  
 ————— castor, compound, 151  
 ————— catechu, compound, 152  
 ————— cevadilla, compound, 355  
 ————— chalk, aromatic, 136  
 ————— compound, 136  
 ————— with opium, 136  
 ————— charcoal and quassia, 146  
 ————— chlorate of potassa, 334  
 ————— compound, 334  
 ————— chloride of barium, 124  
 ————— chloride of gold, 120  
 ————— and sodium, 121  
 ————— silver, 110  
 ————— cinnamon, compound, 167  
 ————— cocoa, compound, 132  
 ————— colocynth, compound, 172  
 ————— columbo and ginger, 174  
 ————— iron, 173  
 ————— magnesia, 173  
 ————— tartrate of iron, 174  
 ————— common salt and cochineal, 378  
 ————— compound, 378  
 ————— contrayerva, compound, 176  
 ————— coriander, compound, 180  
 ————— corrosive sublimate and copper, 237  
 ————— zinc, 237  
 ————— cubebs, 183  
 ————— and alum, 183  
 ————— ergot, 183  
 ————— hemlock, 183  
 ————— cyanide of gold, 122  
 ————— disinfecting, 337  
 ————— Dover's, 252  
 ————— emmenagogue, 286  
 ————— ergot, compound, 196  
 ————— fennel, compound, 215  
 ————— foxglove, 190  
 ————— fumigating, 128, 247  
 ————— galls, compound, 218  
 ————— gamboge, compound, 219  
 ————— gentian, compound, 220  
 ————— gold, 120  
 ————— golden sulphuret of antimony, 106  
 ————— guaiacum, compound, 227  
 ————— hæmostatic, 348  
 ————— hedge hyssop, 226  
 ————— hemlock, 175  
 ————— henbane, compound, 248  
 ————— horse chestnut, compound, 247  
 ————— indigo, 250  
 ————— iodide of barium, 125  
 ————— lead, 317  
 ————— mercury, 242  
 ————— sulphur, 395  
 ————— ipecacuanha and antimony, 253  
 ————— calomel, 253  
 ————— carbonate of soda, 253  
 ————— muriate of ammonia, 253  
 ————— myrrh, 253  
 ————— nitre, 253  
 ————— opium, 252  
 ————— rhubarb, 253  
 ————— tartar emetic, 253  
 ————— tragacanth, 253  
 ————— jalap and calomel, 255  
 ————— cremor tartar, 256  
 ————— ipecacuanha, 256  
 ————— scammony, 256  
 ————— compound, 256



- Powder, James's, 109  
 — kermesine, 106  
 — compound, 106  
 — kino, compound, 258  
 — lavender, compound, 264  
 — lupuline, 268  
 — madder, compound, 354  
 — magnesia and orange peel, 269  
 — rhubarb, 269  
 — sulphur, 269  
 — male fern, 214  
 — marsh mallow, 86  
 — mercurial, compound, 233  
 — metallic iron, 200  
 — compound, 200  
 — milk, 261  
 — musk, 283  
 — myrrh and ipecacuanha, 286  
 — nitrate of potassa and camphor, 335  
 — orris root, 335  
 — squill, 335  
 — compound, 335  
 — nux vomica, 288  
 — oak bark, compound, 342  
 — opium and antimony, 296  
 — camphor, 296  
 — chalk, compound, 296  
 — musk, 296  
 — nitre, 296  
 — sulphur, 296  
 — roasted, 296  
 — orange peel and rhubarb, 119  
 — orpiment, 114  
 — oxalate of potassa, 336  
 — oxide of gold, 122  
 — manganese, 273  
 — Peruvian bark, and arnica, 161  
 — camphor, 161  
 — cascarilla, 161  
 — isinglass, 161  
 — valerian, 161  
 — compound, 160  
 — phloridzine, 308  
 — phosphate of soda, compound, 384  
 — pinkroot and calomel, 386  
 — savine, 386  
 — Portland, 154  
 — Prussian blue, 205  
 — compound, 205  
 — red oxide of mercury, 244  
 — sulphuret of mercury, 246  
 — rhatany, compound, 259  
 — rhubarb and chalk, 348  
 — magnesia, 348  
 — sulphate of potassa, 348  
 — compound, 348  
 — sal ammoniac, 95  
 — salicin, compound, 358  
 — salivation, against, 72  
 — sarsaparilla and Peruvian bark, 362  
 — savine and ginger, 356  
 — pinkroot, 356  
 — Spanish flies, 356  
 — verdigris, 356  
 — scammony and cremor tartar, 365  
 — soot, 366  
 — compound, 366  
 — Seidlitz, 385  
 — semen contra, 115  
 — seneka, compound, 371  
 — senna, and guaiacum, 373  
 — compound, 373  
 — soap, cosmetic, 361  
 — spermaceti, compound, 155  
 — squill and calomel, 367  
 — cremor tartar, 367  
 — ipecacuanha, 367  
 Powder, squill and nitre, 367  
 — sulphur, 367  
 — compound, 367  
 — starch, compound, 101  
 — subcarbonate of iron, 202  
 — sulphate of copper, 187  
 — compound, 187  
 — iron, 211  
 — mercury, compound, 245  
 — potassa, and rhubarb, 337  
 — compound, 337  
 — quinine, 345  
 — and morphia, 345  
 — soda, 345  
 — tartar emetic, 345  
 — tartaric acid, 345  
 — compound, 345  
 — soda and opium, 384  
 — nitre, 384  
 — tartar emetic, 384  
 — sulphur and antimony, 393  
 — camphor, 392  
 — cremor tartar, 393  
 — liquorice, 393  
 — magnesia, 393  
 — orris root, 393  
 — compound, 393  
 — sulphuret of potassium and cremor tar-  
 tar, 326  
 — tartar emetic, 107  
 — and ipecacuanha, 107  
 — phosphate lime, 107  
 — quinine, 107  
 — compound, 107  
 — tartarized soda and rhubarb, 385  
 — tartrate of iron and columbo, 212  
 — effervescent, 212  
 — potassa and rhubarb, 338  
 — tragacanth, compound, 402  
 — uva ursi, compound, 404  
 — valerian, compound, 404  
 — verdigris and savine, 185  
 — wild ipecacuanha, 198  
 — willow bark, compound, 358  
 — white hellebore, 408  
 — zinc, oxide, 410  
 — and colombo, 410  
 — compound, 410  
 — cyanide, 412  
 — ferricyanide, 412  
 Precipitation, 447  
 Preservative liquid, 134  
 Proper use of utensils, 49  
 Prunes, 340  
 Prussian blue, 205  
 — ink, 206  
 — mixture, 206  
 — ointment, 206  
 — pills, 206  
 — powder, 205  
 — compound, 205  
 Pudding, arrowroot, 274  
 — tapioca, 398  
 Pulp of purging cassia, 144  
 — tamarinds, 397  
 Purging cassia, 150  
 Purified animal charcoal, 145  
 Pyroligneous acid, 69  
 — cataplasm, 69  
 — injection, 69  
 — mouth wash, 69  
 Quassia, 341  
 — extract, 341  
 — infusion, 341



- Quassia, infusion, compound, 341  
 ———— tincture, 341  
 ———— compound, 341  
 ———— wine, 341  
 Quiet in sick room, 47  
 Quince seeds, 188  
 ———— bandoline, 180  
 ———— decoction, 189  
 ———— mixture, 189  
 Quinine, 342  
 ———— acetate, 342  
 ———— amorphous, 342  
 ———— and chloride mercury, 343  
 ———— pills, 344  
 ———— arseniate, 343  
 ———— citrate, 343  
 ———— and iron, 343  
 ———— syrup, 343  
 ———— ferrocyanate, 343  
 ———— mixture, 343  
 ———— pills, 343  
 ———— hydriodate, 343  
 ———— impure, 342  
 ———— tincture, 342  
 ———— iodide, with iron, 343  
 ———— ioduretted hydriodate, 343  
 ———— kinate, 344  
 ———— pills, 344  
 ———— lactate, 344  
 ———— mixture, 344  
 ———— pills, 344  
 ———— syrup, 344  
 ———— muriate, 344  
 ———— mixture, 344  
 ———— nitrate, 344  
 ———— phosphate, 344  
 ———— sulphate, 344  
 ———— dentifrice, 347  
 ———— enema, 347  
 ———— gargle, 347  
 ———— liniment, 346  
 ———— mixture, 346  
 ———— with coffee, 346  
 ———— ointment, 347  
 ———— pills, 345  
 ———— and gentian, 346  
 ———— compound, 346  
 ———— plaster, 347  
 ———— powder, 345  
 ———— and morphia, 345  
 ———— soda, 345  
 ———— tartar emetic, 345  
 ———— tartaric acid, 345  
 ———— compound, 345  
 ———— syrup, 346  
 ———— and coffee, 346  
 ———— tincture, 346  
 ———— compound, 346  
 ———— wine, 346  
 ———— aromatic, 346  
 ———— sulpho-tannate, 347  
 ———— tannate, 347  
 ———— tartrate, 347  
 ———— valerianate, 347  
 Racahout, 132  
 Ratafia of wormwood, 66  
 Red cedar, 258  
 ———— iodide of mercury, 243  
 ———— lead, 318  
 ———— oxide of iron, 210  
 ———— mercury, 244  
 ———— poppy, 351  
 ———— infusion, 352  
 ———— compound, 352  
 Red poppy, syrup, 352  
 ———— roses, 352  
 ———— sulphuret of mercury, 246  
 Refrigerants, 61  
 Resin, 347  
 ———— cerate, 348  
 ———— compound, 348  
 ———— hemp, 141  
 ———— jalap, 256  
 ———— plaster, 348  
 ———— St. Andrew's, 348  
 ———— powder, hæmostatic, 348  
 ———— scammony, 366  
 Rhatany, 259  
 ———— clyster, 260  
 ———— collutory, 260  
 ———— dentifrice, 259  
 ———— electuary, 259  
 ———— extract, 259  
 ———— infusion, 259  
 ———— injection, 260  
 ———— mixture, 260  
 ———— ointment, compound, 260  
 ———— pills, 259  
 ———— powder, compound, 259  
 ———— syrup, 260  
 ———— tincture, 260  
 Rhubarb, 348  
 ———— electuary, 349  
 ———— extract, 350  
 ———— fluid, 350  
 ———— infusion, 349  
 ———— alkaline, 350  
 ———— lozenges, 349  
 ———— mixture, 351  
 ———— pills, 349  
 ———— compound, 349  
 ———— and caraway, 349  
 ———— chalk, 348  
 ———— chamomile, 349  
 ———— ipecacuanha, 349  
 ———— oxgall, 349  
 ———— soda, 349  
 ———— powder and magnesia, 348  
 ———— sulphate of potassa, 348  
 ———— compound, 348  
 ———— roasted, 348  
 ———— suppository, 349  
 ———— syrup, 350  
 ———— aromatic, 350  
 ———— and senna, 350  
 ———— tincture, 350  
 ———— alkaline, 351  
 ———— and aloes, 350  
 ———— aniseed, 351  
 ———— gentian, 350  
 ———— senna, 350  
 ———— Warner's cordial, 351  
 ———— wine, 351  
 ———— and gentian, 351  
 ———— compound, 351  
 Rice, 304  
 ———— blanchmange, 420  
 ———— custard, 420  
 ———— gruel, 304  
 ———— mucilage, 304  
 ———— water, 304  
 Rochelle salt, 385  
 Rose, dog, 352  
 ———— confection, 352  
 ———— hundred-leaved, 352  
 ———— collutory, 352  
 ———— essence, 352  
 ———— lozenges, 352  
 ———— oil, 352  
 ———— water, 352  
 ———— ointment, 352



- Rose, red, 352  
     confection, 353  
     electuary, 353  
     honey, 353  
     infusion, acid, 353  
     syrup, 353  
     tincture, 353  
     vinegar, compound, 353  
     wine, 353  
 Rosemary, 353  
     aromatic bath, 353  
     fomentation, 853  
     vinegar, 353  
     Hungary water, 354  
     oil, 354  
     spirit, 354  
     tincture, 354  
 Rosin, 347  
 Round-leaved dogwood, 180  
 Rue, 355  
     confection, 355  
     extract, 355  
     mixture, 355  
     and squill, 355  
     syrup, 355  
     tincture, 355  
 Rules for the administration of medicine, 53  
 Rust's astrigent, 87  
  
 Saccharate of lead, 318  
 Saccharine carbonate of iron, 202  
 Safflower, 148  
     infusion, 148  
 Saffron, 182  
     collyrium, 183  
     elixir, 183  
     infusion, 183  
     ointment, 183  
     pills, 183  
     syrup, 183  
     tincture, 183  
     compound, 183  
 Sagapenum, 357  
     pills, compound, 357  
 Sage, 358  
     gargle, 359  
     infusion, 358  
     compound, 359  
     vinegar, 359  
 Sago, 357  
     milk, 358  
     mucilage, 357  
     posset, 357  
 Sal ammoniac, 95  
 Salep, mucilage, 417  
     powder, compound, 417  
 Salicin, 358  
     pills, 358  
     compound, 358  
     powder, compound, 358  
 Salt, common, 378  
 Saltpetre, 335  
 Saponine, 361  
 Sarsaparilla, 362  
     beer, 365  
     decoction, 362  
     compound, 362  
     Feltz's, 363  
     Jauperand's, 363  
     Vinache's, 363  
     Zittman's, 363  
     essence, 363  
     compound, 364  
     extract, 363  
     fluid, 363  
     Indian, 230  
     decoction, 231  
     Sarsaparilla, Indian, infusion, 230  
     mixture, 230  
     syrup, 230  
     infusion, 362  
     alkaline, 362  
     Lisbon diet drink, 363  
     powder, with Peruvian bark, 362  
     syrup, 364  
     compound, 364  
     Gesnouin's, 365  
     Laffecteur's, 364  
 Sassafras, 364  
     bark, 364  
     infusion, 364  
     compound, 364  
     oil, 364  
     tincture, 364  
     compound, 364  
     pith, 364  
     mucilage, 364  
 Savine, 356  
     cerate, 357  
     extract, 356  
     infusion, 357  
     oil, 357  
     ointment, 357  
     pills, 356  
     powder with cantharides, 356  
     ginger, 356  
     pinkroot, 356  
     verdigris, 356  
     tincture, 357  
     compound, 357  
 Scammony, 365  
     chocolate, 367  
     confection, 366  
     elixir, 366  
     emulsion, 366  
     compound, 366  
     mixture, 366  
     oil, 366  
     pastilles, 366  
     de santé, 366  
     pills, 366  
     compound, 366  
     with ox-gall, 366  
     powder with cream tartar, 365  
     soot, 365  
     compound, 366  
 Scarifications, 477  
 Scurvy grass, 169  
     electuary, 169  
     gargle, 169  
 Sea holly, 197  
 Sealing-wax, red, 262  
 Sedatives, 61  
 Seidlitz powders, 385  
 Semen contra, 115  
     electuary, 115  
     mixture, 115  
     powder, 115  
 Semi-vitrified oxide of lead, 317  
 Seneka, 371  
     decoction, 372  
     emulsion, 372  
     extract with squill, 372  
     infusion, 372  
     compound, 372  
     jelly, 372  
     mixture, 372  
     pills, 371  
     polygalic acid, 372  
     syrup, 373  
     powder, compound, 371  
     syrup, 327  
 Senna, 373  
     American, 150



- Senna, American, infusion, 151  
 — confection, 373  
 — compound, 373  
 — electuary with cremor tartar, 373  
 — figs, 373  
 — rhubarb, 373  
 — sulphur, 373  
 — emulsion, 375  
 — extract, alcoholic, 373  
 — fluid, 375  
 — infusion, 374  
 — compound, 374  
 — with buckthorn, 375  
 — coffee, 374  
 — lemon juice, 375  
 — rhubarb, 375  
 — tamarinds, 374  
 — tincture, 375  
 — mixture, 375  
 — pills, 373  
 — powder, compound, 373  
 — and guaiacum, 373  
 — syrup, 375  
 — and apples, 375  
 — rhubarb, 375  
 — tincture, compound, 374  
 — with gentian, 374  
 — jalap, 374  
 — wine, compound, 374  
 Setons, 471  
 Shampoo liquid, 144  
 Sialagogues, 61  
 Silicate of potassa, 336  
 Silk weed, 116  
 Silver, 110  
 — chloride, 110  
 — and ammonia, 110  
 — pills, 110  
 — powder, 110  
 — cyanide, 110  
 — ointment, 110  
 — iodide, 110  
 — ointment, 111  
 — nitrate, 111  
 — collyrium, 111  
 — hair dye, 111  
 — indelible ink, 111  
 — mixture, 111  
 — ointment, 112  
 — compound, 112  
 — pills, 111  
 — solution, 111  
 — Mackensie's, 111  
 — oxide, 111  
 — ointment, 111  
 Simaruba, 377  
 — infusion, 377  
 — compound, 377  
 Sinapisms, 469  
 Skunk cabbage, 193  
 — infusion, 193  
 — tincture, 193  
 Slippery elm, 403  
 — cataplasm, 404  
 — infusion, 403  
 Snakeroot, black, 160  
 — decoction, 160  
 — tincture, 160  
 — Virginia, 376  
 — electuary, 376  
 — infusion, 376  
 — compound, 376  
 — mixture, with allspice, 376  
 — acetated, 376  
 — ether, 376  
 — pills, compound, 376  
 Snakeroot, Virginia, tincture, 376  
 — with balsam Peru, 376  
 — wine, with vanilla, 376  
 Sneezewort, 229  
 Soap, 360  
 — almond, 360  
 — aromatic, 360  
 — arsenical, 361  
 — balsam, camphorated acetic, 361  
 — beef marrow, 360  
 — bolus, 361  
 — camphorated, 360  
 — castile, 360  
 — cataplasm, 362  
 — clyster, 362  
 — cod liver oil, 362  
 — ioduretted, 362  
 — common, 360  
 — electuary, 361  
 — essence, 361  
 — camphorated, 361  
 — grease balls, 361  
 — hard, 360  
 — iodide of potassium, 323  
 — itch ointment, 362  
 — jalap, 257  
 — pills, 361  
 — and ox-gall, 361  
 — compound, 361  
 — powder, aromatic, 361  
 — saponine, 361  
 — soft, 360  
 — solution, ethereal, 361  
 — Spanish, 360  
 — Starkey's, 293, 360  
 — sulphuret of potassium, 328  
 — suppository, 362  
 — turpentine, 360  
 — with sulphur, 362  
 Soda, 379  
 — acetate, 379  
 — pills, compound, 379  
 — arseniate, 380  
 — solution, 380  
 — bicarbonate, 381  
 — lozenges, 381  
 — bisulphate, 385  
 — and magnesia, 385  
 — borate, 380  
 — cakes, 421  
 — carbonate, 381  
 — dried, 381  
 — emulsion, 382  
 — injection, 382  
 — lotion, 382  
 — mixture, anti-emetic, 382  
 — with chamomile, 382  
 — copaiba, 382  
 — gentian, 382  
 — ipecacuanha, 382  
 — quassia, 382  
 — ointment, 382  
 — pills, 381  
 — powder, with magnesia, 381  
 — mercury, 381  
 — rhubarb, 381  
 — solution, diuretic, 382  
 — caustic, 379  
 — chlorate of, 383  
 — gargle, 383  
 — lotion, 383  
 — chlorinated, 383  
 — gargle, 383  
 — solution, 383  
 — diluted, 383  
 — hydrosulphate of, 383



- Soda, hyposulphite of, syrup, 383  
     phosphate, 384  
         mixture, 384  
         powder, compound, 384  
     pills, with ipecacuanha, 381  
         rhubarb, 381  
     powder, effervescing, 381  
     solution, 379  
         effervescing, 381  
     sulphate, 384  
         clyster, 385  
         electuary, 384  
         emulsion, 384  
         lemonade, 384  
         lotion, 385  
         pills, compound, 384  
         powder, with nitre, 384  
             opium, 384  
             tartar emetic, 384  
         solution, compound, 384  
         suppository, 385  
     tartarized, 385  
         clyster, 385  
         mixture, 385  
         powder, with rhubarb, 385  
         Seidlitz's, 385  
         whey, 385
- Sodium, 378  
     bromide, 378  
         ointment, 378  
     chloride, 378  
         bath, with gelatine, 378  
         clyster, 378  
             and arnica, 379  
         fomentation, 378  
         mixture, with lemon juice, 379  
         ointment, 379  
             compound, 379  
         powder, compound, 378  
             and cochineal, 378  
     chloroplatinate, 313  
         injection, 313  
     sulphuret, 379  
         bath, 379  
         liniment, 379  
         lotion, 379  
         mixture with sal ammoniac, 379  
             ointment, 379
- Soluble glass, 337
- Solution, arsenical, 330  
     Donovan's, 114  
     Fowler's, 330  
     Labarraque's, 383  
     of acetate ammonia, 91  
         barytes, 126  
         mercury, 236  
         morphia, 281  
             alcoholic, 281  
     alum, 89  
         odontalgic, 88  
     arsenate of ammonia, 92  
         soda, 380  
     arsenic, 69  
     ammoniated copper, 187  
         muriate of copper and mercury, 186  
         ammonio-tartrate of iron, 213  
         atropine, 118  
         barytes, 125  
         belladonna, extract, 127  
         bimeconate of morphia, 281  
         bitartrate of potassa, 339  
             compound, 339  
         borotartrate of potassa, 331  
         bromide of mercury, ethereal, 236  
             potassium, 319
- Solution of bromine, 130  
     alcoholic, 130  
     brucia, 131  
     carbonate of potassa, 332  
         soda, 382  
             diuretic, 382  
             effervescing, 381  
     caustic potassa, 328  
     chlorate of barium, 125  
         potassa, 334  
     chloride of calcium, 133  
         gold and sodium, 121  
         soda, 383  
             diluted, 383  
     chlorinated potassa, 320  
     citrate of magnesia, 271  
         potassa, 234  
     corrosive sublimate, 237  
         alcoholic, 237  
     cotton, ethereal, 225  
     creasote, alcoholic, 183  
     croton oil, saponaceous, 294  
     cyanide of gold, 122  
     cyanuret of mercury, 241  
         potassium, 320  
     delphinia, 189  
     ferrocyanide of zinc, 412  
     gamboge, alkaline, 219  
     hydrargyro-iodide of potassium, 325  
     iodide of iron, 207  
         official, 207  
         potassium, 321  
             caustic, 324  
             compound, 324  
         zinc, 413  
     lac, 262  
     magnesia, 270  
     muriate of morphia, 282  
     myrrh, alkaline, 286  
     nitrate of silver, 111  
     nitro-saccharate of lead, 318  
     potassa, 328  
         effervescing, 333  
         lithontriptic, 329  
     persesquinitrate of iron, 209  
     phosphate of ammonia, 96  
     soap, ethereal, 361  
     soda, 379  
     subacetate of lead, 314  
     sulphate of bebeerine, 126  
         cadmium, 133  
         iron, 211  
         morphia, 283  
         soda, compound, 384  
         veratrine, 408  
         zinc, 414  
     sulphuret of potassium, 327  
     tartrate of iron and potassa, 212  
         potassa and ammonia, 339
- Soot, 215  
     cataplasm, 216  
     decoction, 215  
     injection, 216  
     mixture, 216  
     mouth wash, 216  
     oil, 216  
     ointment, 216  
         compound, 216  
     pills, 215  
     tincture, 216
- Southernwood, 114  
     clyster, 114  
     infusion, 114
- Spanish flies, 141
- Spearmint, 277  
     arquebusade water, 278  
     infusion, 278



- Spearmint, infusion, compound, 278  
 Specific gravity, 20  
 Spermaceti, 154  
     — cerate, 155  
     — liniment, 155  
     — lip salve, 155  
     — mixture, 155  
     — ointment, 155  
     — and rosewater, 155  
     — sultana, 155  
     — powder, compound, 155  
 Spirits, 461  
 Spirit of acetic ether, 79  
     — allspice, 309  
     — ammonia, and succinic acid, 86  
         — fetid, 91  
     — aniseed, 103  
     — balm, compound, 277  
     — caraway, 149  
     — castor, compound, 151  
     — cloves, 149  
     — horseradish, compound, 112  
     — juniper, compound, 258  
     — lavender, 264  
         — compound, 265  
     — lemon, 266  
         — odoriferous, 266  
     — muriatic ether, 80  
     — nutmeg, 285  
         — compound, 285  
     — orange peel, 119  
     — rosemary, 354  
     — succinated ammonia, 96  
     — sulphuric ether, 80  
     — sweet of nitre, 79  
     — vanilla, 407  
     — vinegar, aromatic, 68  
     — wormwood, compound, 65  
 Sponge, 387  
     — burnt, 387  
     — bolus, 387  
     — electuary, 387  
     — powder, 387  
         — compound, 387  
 Sponging, 467  
 Spruce beer, 420  
 Spurge, 198  
 Squill, 367  
     — electuary, 370  
     — emulsion, 370  
     — extract, 370  
         — and rhubarb, 369  
     — ointment, 370  
         — compound, 370  
     — oxymel, 370  
         — mixture, with ammonia, 370  
             — ipecacuanha, 370  
             — marsh mallow, 370  
     — pills and ammoniac, 367  
         — calomel, 367  
         — croton oil, 367  
         — henbane, 368  
         — ipecacuanha, 367  
         — compound, 368  
     — powder with calomel, 367  
         — cremor tartar, 367  
         — ipecacuanha, 367  
         — nitre, 367  
         — sulphur, 367  
         — compound, 367  
     — syrup, 370  
         — aromatic, 371  
         — compound, 368  
         — mixture, 369  
     — tincture, 369  
         — alkaline, 369  
         — and elaterium, 369  
     — Squill, tincture, and benzoin, compound, 369  
         — ethereal, 369  
         — vinegar, 370  
         — mixture, 370  
             — and ammonia, 370  
         — wine, 368  
         — compound, 369  
         — bitter, 369  
 Star grass, 81  
     — tincture, 81  
 Starch, 101  
     — iodide, 102  
     — jelly, 102  
     — lozenges, 102  
     — mixture, with suet, 102  
     — mucilage, 101  
         — powder, compound, 101  
 Starkey's soap, 273, 360  
 Stavesacre, 388  
     — decoction, 388  
     — ointment, 388  
     — vinegar, 388  
 Stimulants, 62  
 Storax, 389  
     — pills, compound, 389  
     — purified, 389  
     — syrup, 389  
 Stramonium, 388  
     — extract of leaves, 388  
     — seeds, 388  
     — lotion, 388  
     — mixture, 388  
     — ointment, 389  
     — pills, 388  
         — compound, 388  
     — syrup, 389  
     — tincture of leaves, 389  
         — seeds, 389  
         — ethereal, 389  
     — wine, 389  
 Strychnine, 389  
     — acetate, 390  
     — solution, 390  
     — tincture, 391  
     — collyrium, 390  
     — iodate, 391  
     — liniment, 390  
     — mixture, 390  
     — muriate, 390  
     — nitrate, 391  
         — ointment, 391  
     — ointment, 390  
     — pills, 390  
         — stimulant, 390  
     — powder, compound, 390  
     — sulphate, 391  
     — tincture, 390  
 Styptic of sulphate of copper, 188  
 Stuping, 468  
 Subacetate of copper, 185  
     — lead, 314  
 Subcarbonate of iron, 202  
 Subnitrate of bismuth, 129  
 Succinate of ammonia, 96  
 Succory, 160  
     — extract, 160  
     — infusion, 160  
     — syrup, compound, 160  
 Sugar, 357  
     — barley, 231  
     — burnt, 357  
     — milk, 261  
     — syrup, 357  
     — vanilla, 407  
 Sulphate of alumina, 87  
     — and potassa, 87  
     — ammonia, 97



- Sulphate of antimony, 107  
     — brucia, 131  
     — cinchonia, 166  
     — copper, 187  
     — indigo, 250  
     — iron, 210  
     — magnesia, 271  
     — manganese, 273  
     — morphia, 282  
     — potassa, 337  
     —      and ammonia, 337  
     —      magnesia, 337  
     —      sulphur, 338  
     — quinine, 344  
     — soda, 384  
     — zinc, 414  
 Sulpho-cyanuret of potassium, 326  
     — tartrate of quinia, 347  
 Sulphur, 392  
     — balsam, 394  
     —      etheral, 394  
     —      terebinthinated, 394  
     — cerate, 394  
     — electuary, 393  
     —      compound, 393  
     — iodide, 395  
     —      ointment, 395  
     —      powder, 395  
     — linctus, 393  
     — liniment, with soap, 394  
     — mixture, 394  
     — ointment, 394  
     —      compound, 394  
     —      with camphor, 394  
     —      soap, 394  
     —      zinc, 394  
     — powder, compound, 393  
     —      with antimony, 393  
     —      camphor, 392  
     —      cremor tartar, 393  
     —      liquorice, 393  
     —      magnesia, 393  
     —      orris root, 393  
     — precipitated, 392  
     — sublimed, 392  
     — washed, 392  
 Sulphuret of antimony, 105  
     — arsenic, 114  
     — barium, 125  
     — calcium, 135  
     — carbon, 395  
     — iron, 212  
     — magnesia, 272  
     — mercury, black, 246  
     —      red, 246  
     — potassium, 326  
     — soda, 379  
 Sulphuretted water, artificial, 73  
 Sulphuric acid, 75  
     — and alcohol, 76  
     —      nitric ether, 76  
     — dilute, 75  
     — lemonade, 76  
     — liniment, 76  
     — ointment, 76  
     — ether, 80  
     — etheral oil, 80  
     — Hoffmann's anodyne, 80  
     — lotion, 80  
     — mixture, with camphor, 81  
     —      turpentine, 81  
     — rectified, 80  
     — spirit, 80  
     — syrup, 81  
 Suppositories, 56  
 Suppository, anthelmintic, 84  
     — charcoal, 146  
 Suppository, hemlock, 176  
     — opium, 299  
     — rhubarb, 349  
     — soap, 362  
     — zinc, chloride of, 411  
 Swamp dogwood, 180  
 Sweet almonds, 98  
     — fern, 174  
     — spirit of nitre, 79  
     — violet, 409  
 Sydenham's laudanum, 301  
 Syrups, 450  
 Syrup, 357  
     — acetate of morphia, 281  
     — antimonial wine, 109  
     — asparagus shoots, 116  
     — balsam of Peru, 285  
     —      tolu, 401  
     — belladonna, 127  
     —      Gillet's, 127  
     — biniodide of potassium, 324  
     — bittersweet, 194  
     — blackberry, 355  
     — brooklime, 126  
     — cadmium, 148  
     — capillaire, 79  
     — chamomile, 104  
     — chloride of gold and sodium, 121  
     —      compound, 122  
     —      iron, 204  
     — cinchonia, 167  
     — citrate of iron, 205  
     — citric acid, 71  
     — clove pink, 190  
     — cod liver oil, 290  
     — codeia, 169  
     — coffee, 170  
     — colchicum, 171  
     — coltsfoot, 403  
     — cubebs, alcoholic etheral extract, 184  
     — cyanuret of potassium, 320  
     — emetine, 196  
     — ergot, 196  
     — galls, 218  
     — garlic, 81  
     — gentian, 221  
     —      extract, 221  
     — gentianin, 222  
     — ginger, 415  
     —      etheral extract, 416  
     — gold, 120  
     — gum, 66  
     —      compound, 66  
     — hive, 368  
     — horehound, 274  
     — hydrocyanic acid, 72  
     — hyposulphite of soda, 383  
     — hyposulphited sulphuret of potassium, 327  
     — Indian sarsaparilla, 230  
     — iodate of potassa, 335  
     — iodide and chloride of iron, 208  
     —      of iron, 207  
     —      potassium, 322  
     —      zinc, 413  
     — ipecacuanha, 254  
     —      compound, 254  
     — kino, 259  
     — lactate of iron, 208  
     —      quinine, 344  
     — lactucarium, 263  
     — lemon, 266  
     — peel, 266  
     —      vinous, 266  
     — lettuce, 263  
     — liquorice, 224  
     — lobelia, 268  
     — lupuline, 268



- Syrup, maidenhair, 79  
     — compound, 79  
     — manna, 273  
     — marsh mallow, 86  
     — matico, 275  
     — milk, 261  
     — monesia, 279  
     — compound, 279  
     — mulberries, 280  
     — muriate of morphia, 282  
     — compound, 282  
     — naphthaline, 287  
     — opium, 299  
     — succinated, 299  
     — orange flowers, 120  
     — peel, 119  
     — orgeat, 99  
     — paullinia, 307  
     — pectoral, 306  
     — Peruvian bark, 166  
     — pipsissewa, 158  
     — poppy heads, 306  
     — proto-iodide of iron, 207  
     — red roses, 353  
     — rhatany, 260  
     — rhubarb, 350  
     — and senna, 350  
     — aromatic, 350  
     — rue, 355  
     — saffron, 183  
     — sarsaparilla, 364  
     — compound, 364  
     — Gesnouin's, 365  
     — Laffecteur's, 364  
     — seneka, 372  
     — senna, 375  
     — and apples, 375  
     — rhubarb, 375  
     — squill, 370  
     — aromatic, 371  
     — compound, 368  
     — storax, 389  
     — stramonium, 389  
     — succory, compound, 160  
     — sulphate of iron, 212  
     — morphia, 283  
     — quinine, 346  
     — and coffee, 346  
     — sulphuret of magnesia, 272  
     — potassium, 327  
     — compound, 327  
     — sulphuretted, 95  
     — sulphuric ether, 81  
     — thridace, 262  
     — uva ursi, 404  
     — valerian, 405  
     — vinegar, 67  
     — violets, 409  
     — wild cherry bark, 340  
     — wormwood, 65
- Table of avoirdupois weights, 10  
     — distilled oils, 458  
     — drops and minims, 18  
     — foreign weights, 14  
     — hydrometrical equivalents, 28  
     — incompatibles, 422  
     — pharmaceutical names, 433  
     — posological, 426  
     — sp. grs. and Baumé's hydrometer, 25  
     — of preparations, 31
- Tamarinds, 397  
     — electuary, 397  
     — infusion, 397  
     — pulp, 397  
     — whey, 397
- Tannate of lead, 318  
     — of quinine, 347  
 Tannic acid, 76  
     — gargle, 76  
     — injection, 76  
     — mixture, 76  
     — ointment, 76  
     — pills, 76  
     — and opium, 76
- Tansy, 397  
     — extract, 397  
     — infusion, 297  
     — oil, 397  
     — pills, 397  
     — tincture, compound, 397
- Tapioca, 397  
     — jelly, 397  
     — pudding, 397
- Tar, 312  
     — ointment of, 312  
     — compound, 312  
     — pills, 312  
     — water, 312
- Tartaric acid, 77  
     — lemonade, 77  
     — lozenges, 77
- Tartar emetic, 107  
     — bolus, 107  
     — clyster, 108  
     — emulsion, 108  
     — liniment, with ammonia, 108  
     — lotion, 108  
     — and camphor, 108  
     — corrosive sublimate, 108  
     — ointment, 108  
     — compound, 108  
     — pills and camphor, 107  
     — guaiacum, 107  
     — opium, 107  
     — plaster, 108  
     — powder, 107  
     — and ipecacuanha, 107  
     — phosphate of lime, 107  
     — quinine, 107  
     — compound, 107  
     — solution, 108  
     — and squill, 108  
     — wine, 108  
     — collyrium, 108  
     — mixture, with ammonia, 109  
     — bitter sweet, 109  
     — laudanum, 109  
     — syrup, 109
- Tartarized soda, 385  
 Tartrate of antimony and potassa, 107  
     — iron and potassa, 212  
     — mercury and potassa, 247  
     — potassa, 338  
     — and ammonia, 339  
     — quinine, 347  
     — soda and potassa, 385
- Tea, beef, 418  
     — bran, 419  
     — mutton, 418  
     — veal, 418
- Temperatures in pharmaceutical operations, 28  
 Tepid bath, 463  
 Thermometrical scales, 32  
 Thridace, 262  
 Tin, 387  
     — chloride, 387  
     — electuary, 387  
     — ointment, 387  
     — oxide, 388  
     — powder, 387



- Tin, sulphuret, 387  
     powder, compound, 388  
 Tinctures, 451  
 Tincture, acetate of iron, 200  
     alcoholic, 200  
     ethereal, 200  
     morphia, 330  
     potassa, 391  
     zinc, 413  
     aconite, 78  
         root, 78  
     aloes, 85  
         and myrrh, 85  
     allspice, 309  
     amber, 392  
         alkaline, 392  
         ethereal, 392  
     American centaury, 356  
         columbo, 215  
         hellebore, 409  
     ammonia, compound, 93  
     ammoniated iron, 201  
     angelica, 103  
         compound, 102  
     angustura, 102  
     aniseed, 104  
     antacid, 238  
     arnica, 113  
     assafetida, 117  
         alkaline, 117  
         ammoniated, 118  
         and castor, 118  
         soot, 118  
     balsam of Peru, 285  
         tolu, 400  
         compound, 400  
         with foxglove, 401  
     belladonna, 127  
         ethereal, 128  
     benzoin, 129  
         compound, 129  
     black hellebore, 230  
         snakeroot, 160  
     bloodroot, 360  
     brucia, 131  
     buchu, 192  
     calamus, 133  
         compound, 133  
     camphor, 139  
         and saffron, 139  
         ethereal, 139  
     cantharides, 142  
         and guaiacum, 142  
         camphorated, 142  
         ethereal, 143  
         compound, 142  
         ethereal, 143  
     carbonate of potassa, 332  
         compound, 333  
     cardamom, 147  
         compound, 148  
     cascarilla, 150  
     castor, 151  
         ammoniated, 151  
         compound, 151  
         ethereal, 151  
     catechu, 153  
     Cayenne pepper, 145  
         and cantharides, 145  
     cevadilla, 355  
     chiretta, 158  
     chloride of gold, 121  
         iron, 204  
         ethereal, 204  
         zinc, ethereal, 411  
     cinchonia, 167  
     cinnamon, 167  
     Tincture, cinnamon, compound, 168  
         ethereal, 168  
         cochineal, 169  
         colchicum flowers, 171  
         seeds, 170  
         compound, 171  
         colocynth, 173  
         columbo, 174  
         contrayerva, 177  
         compound, 177  
         copaiba, 179  
         alkaline, 179  
         compound, 179  
         coriander, 180  
         Croton oil, 294  
         cubeb, 184  
         cyanuret mercury, compound, 242  
         Dippel's oil, 289  
         elaterin, 195  
         ergot, 196  
         foxglove, 191  
         ethereal, 191  
         galbanum, 217  
         compound, 217  
         galls, 218  
         gamboge, alkaline, 219  
         ammoniacal, 220  
         gentian, 221  
         acidulated, 221  
         alkaline, 221  
         ammoniacal, 221  
         compound, 221  
         gentianine, 222  
         ginger, ethereal extract, 416  
         goldthread, 179  
         guaiacum, 227  
         ammoniated, 228  
         hemlock, 175  
         ethereal, 175  
         hemp, 141  
         henbane, 249  
         ethereal, 249  
         hops, 232  
         alkaline, 232  
         horse balm, 172  
         hydrargyro-iodide of potassium, 325  
         iodide of iron, 207  
         potassium, compound, 324  
         iodine, 251  
         compound, 251  
         ethereal, 251  
         saturated, 251  
         jalap, 257  
         compound, 257  
         Jamaica dogwood, 311  
         kino, 259  
         lactucarium, 263  
         larkspur seeds, 189  
         lavender, ethereal, 265  
         lettuce, aromatic, 263  
         lobelia, 268  
         ethereal, 268  
         lupuline, 268  
         mace, 269  
         magnolia, 272  
         malate of iron, 209  
         male fern buds, 214  
         mastic, ethereal, 275  
         matco, 275  
         monesia, 279  
         musk, 284  
         artificial, 391  
         myrrh, 287  
         and hellebore, 287  
         nux vomica, 288  
         compound, 288  
         opium, 301



- Tincture, opium, acetated, 301  
 ————— ammoniated, 302  
 ————— and asarabacca, 302  
 ————— soap, 302  
 ————— Bateman's, 302  
 ————— camphorated, 302  
 ————— compound, 302  
 ————— Smith's, 302  
 ————— succinated, 302  
 ————— Swediaur's, 302  
 ————— Warner's, 303  
 ————— opoponax, 304  
 ————— compound, 304  
 ————— orange peel, 119  
 ————— pareira brava, 307  
 ————— pellitory, 340  
 ————— compound, 340  
 ————— Peruvian bark, 164  
 ————— and ammonia, 165  
 ————— cantharides, 165  
 ————— gentian, 165  
 ————— snakeroot, 165  
 ————— valerian, 164  
 ————— compound, 164  
 ————— phosphoric, of myrrh, 75  
 ————— piperine, 311  
 ————— poison oak, 402  
 ————— proto-iodide of iron, 207  
 ————— quassia, 341  
 ————— compound, 341  
 ————— quinine, 342  
 ————— impure, 342  
 ————— red iodide mercury, 243  
 ————— ethereal, 243  
 ————— rhatany, 260  
 ————— rhubarb, 350  
 ————— and aloes, 350  
 ————— aniseed, 351  
 ————— gentian, 350  
 ————— senna, 350  
 ————— alkaline, 351  
 ————— rosemary, 354  
 ————— roses, 353  
 ————— rue, 355  
 ————— saffron, 183  
 ————— compound, 183  
 ————— sassafras, 365  
 ————— compound, 365  
 ————— savine, 357  
 ————— compound, 357  
 ————— senna, and jalap, 374  
 ————— gentian, 374  
 ————— compound, 374  
 ————— skunk cabbage root, 193  
 ————— seed, 193  
 ————— snakeroot, black, 160  
 ————— Virginian, 376  
 ————— and balsam Peru, 376  
 ————— soot, 216  
 ————— squill, 369  
 ————— alkaline, 369  
 ————— and benzoin, 369  
 ————— elaterium, 369  
 ————— ethereal, 369  
 ————— stargrass, 81  
 ————— stramonium ethereal, 389  
 ————— leaves, 389  
 ————— seeds, 389  
 ————— strychnine, 390  
 ————— sulphate of quinine, 346  
 ————— compound, 346  
 ————— sulphuret of potassium, 327  
 ————— tansy, compound, 397  
 ————— tartrate of iron, compound, 213  
 ————— tobacco, 396  
 ————— tulip tree bark, 267  
 Tincture, valerian, 405  
 ————— ammoniated, 405  
 ————— compound, 405  
 ————— compound, 405  
 ————— vanilla, 407  
 ————— veratrine, 408  
 ————— white hellebore, 408  
 ————— winter's bark, 410  
 ————— wormwood, 65  
 ————— alkaline, 65  
 Toast water, 418  
 Tobacco, 395  
 ————— cataplasm, 396  
 ————— extract, 396  
 ————— infusion, 396  
 ————— lotion, 396  
 ————— mixture, 396  
 ————— nicotina, 396  
 ————— oil, 396  
 ————— ointment, 396  
 ————— pills, 396  
 ————— powder, compound, 396  
 ————— tincture, 396  
 ————— wine, 396  
 Tolu, balsam of, 400  
 Tonics, 62  
 Tormentil, 401  
 ————— decoction, 402  
 ————— extract, 401  
 ————— gargle, 402  
 ————— powder, compound, 401  
 Tous les mois, 417  
 Tragacanth, 402  
 ————— mucilage, 402  
 ————— paste, 402  
 ————— powder, compound, 402  
 Troches, 462  
 Tulip tree bark, 267  
 ————— infusion, 267  
 ————— tincture, 267  
 Turlington's balsam, 129  
 Turpentine, 399  
 ————— clyster, 399  
 ————— electuary, 399  
 ————— liniment, 399  
 ————— mixture, 400  
 ————— oil, 292  
 ————— ointment, 399  
 ————— compound, 399  
 ————— pills, 399  
 ————— with guaiacum, 399  
 ————— jalap, 399  
 ————— magnesia, 399  
 ————— myrrh, 399  
 ————— rhubarb, 399  
 ————— plaster, 399  
 ————— compound, 400  
 Uva ursi, 404  
 ————— decoction, 404  
 ————— extract, 404  
 ————— mixture, 404  
 ————— powder, compound, 404  
 ————— syrup, 404  
 Valerian, 404  
 ————— bolus, with iron, 405  
 ————— sulphate of potassa, 405  
 ————— electuary, 405  
 ————— extract, 406  
 ————— fluid, 406  
 ————— infusion, 405  
 ————— compound, 405  
 ————— mixture, with ammonia, 405  
 ————— Hoffmann's anodyne, 406  
 ————— oil, 406  
 ————— with ammonia, 406



- Valerian, pills, compound, 405  
 — powder, compound, 404  
 — syrup, 405  
 — tincture, 405  
 — ammoniated, 405  
 — compound, 405  
 — compound, 405  
 — with Hoffmann's anodyne, 406  
 — wine, 405  
 Valerianate of quinine, 347  
 — zinc, 415  
 Valerianic acid, 77, 406  
 Vanilla, 406  
 — arrowroot, 407  
 — essence, 407  
 — extract, fluid, 407  
 — lozenges, 407  
 — milk, 407  
 — spirit, 407  
 — sugar, 407  
 — tincture, 407  
 Vapor bath, 465  
 Varnish, crystal, 275  
 — picture, 275  
 Veal tea, 418  
 Vegetable broth, 417  
 Venesection, 474  
 Ventilation of sick room, 45  
 Veratrin, 407  
 — liniment, 408  
 — lotion, 408  
 — muriate, 408  
 — nitrate, 408  
 — ointment, 408  
 — with iodine, 408  
 — opium, 408  
 — pills, 407  
 — with henbane, 408  
 — sulphate, 408  
 — solution, 408  
 — tincture, 408  
 Verdigris, 185  
 Vesicatories, 470  
 Vinegar, 67  
 — aromatic, 68  
 — spirit, 68  
 — borax, 380  
 — cantharides, 142  
 — cataplasm, 67  
 — Cayenne pepper, 145  
 — cinnamon, compound, 167  
 — coffee, 176  
 — colchicum root, 171  
 — seeds, 171  
 — distilled, 68  
 — elder flowers, 359  
 — four thieves, 68  
 — gargle, 67  
 — lobelia, 268  
 — mixture, 67  
 — and cardamom, 67  
 — opium, 300  
 — pastilles, 67  
 — prophylactic, 68  
 — rosemary, aromatic, 353  
 — roses, compound, 353  
 — sage, 359  
 — squill, 370  
 — syrup, 67  
 — stavesacre, 388  
 Vinegars, 454  
 Violet, sweet, 409  
 — conserve, 409  
 — honey, 409  
 — oil, 409  
 Vocabulary of words used in prescriptions, 34  
 Volatile oils, 456  
 Walnut, white, 257  
 Warm bath, 463  
 Warner's cordial, 317  
 Wash, black, 244  
 — cosmetic, 129, 238  
 — detergent, 87  
 — ophthalmic, 185  
 — sal ammoniac, 96  
 — yellow, 238  
 Water, allspice, 309  
 — ammonia, 90  
 — anemone, 102  
 — angelica, 103  
 — anti-hysterical, 377  
 — apple, 418  
 — arquebusade, 278  
 — avens, 227  
 — barley, with nitrate potassa, 232  
 — bitter almonds, 100  
 — brooklime, 126  
 — camphor, 139  
 — and laudanum, 139  
 — nitric acid, 139  
 — Hoffmann's anodyne, 139  
 — caraway, 148  
 — chalybeate, artificial, 203  
 — cherry laurel, 264  
 — chlorine, 158  
 — cinnamon, 168  
 — compound, 168  
 — citrate of magnesia, 271  
 — clove, 149  
 — Cologne, 266  
 — coriander, compound, 180  
 — distilled, 455  
 — diuretic, 100  
 — elder flower, 359  
 — fennel, 215  
 — Goulard's vegeto-mineral, 315  
 — honey, 276  
 — Hungary, 354  
 — ioduretted, 323  
 — lavender, 264  
 — lemon peel, 266, 419  
 — lettuce, 263  
 — lime, 135  
 — and carbonate of potassa, 135  
 — milk, 135  
 — compound, 135  
 — medicated, 455  
 — opium, 300  
 — orange flower, 120  
 — partridge berry, 220  
 — rice, 304  
 — rose, 352  
 — Seidlitz, 271  
 — souchy, 419  
 — sulphuret of potassium, 327  
 — sulphuretted, artificial, 73  
 — tar, 312  
 — toast, 418  
 — unparalleled, 267  
 Weights and measures, 9  
 — apothecaries', 12  
 — avoirdupois, 9  
 — foreign, 12  
 — French, 12  
 — metrical, 12  
 — Troy, 9  
 Whey, 260  
 — alum, 89, 260  
 — aromatic, 89, 261  
 — cremor tartar, 260  
 — mustard, 261, 377  
 — tamarind, 260, 397  
 — tartarized soda, 385  
 — vinegar, 260



- Whey, wine, 261  
 White hellebore, 408  
     — decoction, 408  
     — ointment, 409  
     — compound, 409  
     — powder, 408  
     — tincture, 408  
     — wine, 408  
     — precipitate, 236  
     — walnut, 257  
 Wikana, 132  
 Wild cherry bark, 340  
     — infusion, 340  
     — syrup, 340  
     — ginger, 115  
     — infusion, 115  
     — ipecacuanha, 198  
     — powder, compound, 198  
     — potato, 177  
 Wilkinson's liniment, 95  
 Willow bark, 358  
     — decoction, 358  
     — dentifrice, 358  
     — ointment, 358  
     — powder, compound, 358  
 Wine of aconite, compound, 78  
     — aloes, 85  
     — alkaline, 85  
     — balsamic, 85  
     — American hellebore, 409  
     — black hellebore, 230  
     — blessed thistle, 154  
     — bryony, 131  
     — centaury, compound, 154  
     — cinchonia, 167  
     — cinnamon, compound, 167  
     — citrate of iron, 205  
     — aromatic, 205  
     — cloves, 149  
     — colchicum root, 170  
     — seeds, 170  
     — compound, 171  
     — dogwood, round-leaved, 181  
     — ergot, 196  
     — gentian, 221  
     — iodide of iron, 207  
     — ipecacuanha, 255  
     — alkaline, 255  
     — and tartar emetic, 255  
     — compound, 255  
     — iron, 203  
     — opium, 301  
     — Sydenham's, 301  
     — Peruvian bark, 166  
     — persimmons, 193  
     — and calamus, 166  
     — iron, 166  
     — quassia, 341  
     — rhubarb, 351  
     — and gentian, 351  
     — compound, 351  
     — roses, 353  
     — senna, compound, 374  
     — squill, 368  
     — bitter, 369  
     — compound, 369  
     — stramonium, 389  
     — sulphate of quinine, 346  
     — aromatic, 346  
     — tartrate of iron, 212  
     — compound, 212  
     — tobacco, 396  
     — valerian, 405  
     — Virginia snakeroot, 376  
     — white hellebore, 408  
     — wormwood, 65  
     — yellow ladies' bedstraw, 218  
 Wine antimonial, 108  
     — diuretic, 293  
     — mulled, 419  
     — whey, 261  
 Wines, 453  
 Winter's bark, 410  
     — tincture, 410  
 Wistar's cough lozenges, 304  
 Witch hazel, 209  
 Wolfsbane, 112  
     — decoction, 113  
     — extract, 113  
     — fomentation, 113  
     — infusion, 113  
     — compound, 113  
     — powder, compound, 112  
     — tincture, 113  
 Wood sorrel, 67  
     — extract of, 67  
 Wormseed, 157  
     — decoction, 157  
     — oil, 157  
     — mixture, 157  
 Wormwood, 65  
     — clyster, 65  
     — essential oil, 65  
     — infusion, 65  
     — oil, 65  
     — ratafia, 66  
     — syrup, 65  
     — tincture, 65  
     — alkaline, 65  
     — wine, 65  
 Yellow ladies' bedstraw, 218  
     — root, 248  
     — sulphate of mercury, 245  
 Zinc, 410  
     — acetate, 413  
     — collyrium, 413  
     — injection, 413  
     — lotion, 413  
     — tincture, 413  
     — carbonate, 413  
     — cerate, 413  
     — plaster, 414  
     — precipitated, 413  
     — prepared, 413  
     — chloride, 411  
     — disinfectant, 411  
     — injection, 411  
     — lotion, 411  
     — paste, 411  
     — suppository, 411  
     — tincture, ethereal, 411  
     — cyanide, 412  
     — pills, 412  
     — powder, 412  
     — ferrocyanide, 412  
     — mixture, 412  
     — pills, compound, 412  
     — powder, 412  
     — solution, 412  
     — iodide, 412  
     — collyrium, 413  
     — ointment, 413  
     — solution, 413  
     — syrup, 413  
     — oxide, 410  
     — cerate, with lycopodium, 411  
     — lotion, 411  
     — ointment, 411  
     — with calomel, 411  
     — opium, 411  
     — pills, 410  
     — compound, 410



Zinc, oxide, powder, 410  
——— compound, 410  
——— with colomba, 410  
—— sulphate, 413  
——— collyrium, 414  
——— with camphor, 414  
——— fomentation, 414  
——— gargle, 414  
——— injection, 414  
——— lotion, 414

Zinc, sulphate mixture, 415  
——— pills, 414  
——— compound, 414  
——— with turpentine, 414  
——— myrrh, 414  
——— solution, 414  
——— valerianate, 413, 415  
——— mixture, 415  
——— pills, 415

THE END.







NOVEMBER, 1851.

## CATALOGUE

OF

# MEDICAL AND SURGICAL WORKS,

PUBLISHED BY

BLANCHARD & LEA,

PHILADELPHIA,

AND

FOR SALE BY ALL BOOKSELLERS.



## TO THE MEDICAL PROFESSION.

The Subscribers subjoin a list of their publications in medical and other sciences, to which they would invite the attention of the Profession, with the full confidence that they will be found to correspond in every respect with the description. They are to be had of all the principal booksellers throughout the Union, from whom, or from the publishers particulars respecting price, &c., may be had on application.

Philadelphia, Nov., 1851.

BLANCHARD & LEA,  
(Late LEA & BLANCHARD.)

### DICTIONARIES, JOURNALS, &c.

American Journal of the Medical Sciences, quarterly, at \$5 a year.  
Cyclopædia of Practical Medicine, by Forbes, Tweedie, &c., edited by Dunglison, in 4 super royal volumes, 3154 double columned pages.  
Dunglison's Medical Dictionary, 8th ed., 1 vol. imp. 8vo., 928 large pages, (now ready.)  
Hoblyn's Dictionary of Medical Terms, by Hays, 1 vol. large 12mo., 402 pages, double columns.  
Neill and Smith's Compend of the Medical Sciences, 1 vol., large 12mo., 900 pp., 350 cuts.  
Transactions of the American Medical Association, Vols. I, II, and III, cloth or paper.  
Medical News and Library, monthly, at \$1 a year.

### ANATOMY.

Anatomical Atlas, by Smith and Horner, large imp. 8vo., 650 figures. New and cheaper ed.  
Horner's Special Anatomy and Histology, new edition, 2 vols. 8vo., many cuts, (now ready.)  
Horner's United States Dissector, 1 vol. large royal 12mo., many cuts, 444 pages.  
MacLise's Surgical Anatomy, now complete in 1 large imp. 4to. vol., strongly bound, with 68 magnificent colored plates.  
Sharpey and Quain's Anatomy, by Leidy, 2 vols. 8vo., 1300 pages, 511 wood-cuts.  
Wilson's Human Anatomy, by Goddard, 4th edition, 1 vol. 8vo., 252 wood-cuts, 580 pp.  
Wilson's Dissector, by Goddard. New edition, with cuts, 1 vol. 12mo., 458 pages.

### PHYSIOLOGY.

Carpenter's Principles of Human Physiology, 1 vol. 8vo., 752 pp., 300 cuts and 2 plates, new and improved edition, (nearly ready.)  
Carpenter's Elements, or Manual of Physiology, new and improved ed., 1 vol. 8vo., (now ready.)  
Carpenter's General and Comparative Physiology, 1 vol. 8vo., pp. 1122, 321 cuts, (now ready.)  
Dunglison's Human Physiology, 7th edition, 2 vols. 8vo., 1428 pages, and 472 wood-cuts.  
Harrison on the Nerves, 1 vol. 8vo., 292 pages.  
Kirkes and Paget's Physiology, 1 vol. 12mo., many cuts, 550 pages.  
Longet's Physiology. Translated by F. G. Smith. 2 vols. 8vo., many cuts, (preparing.)  
Matteucci on the Physical Phenomena of Living Beings, 1 vol. 12mo., 388 pp., cuts.  
Solly on the Brain, 1 vol. 8vo., 496 pp., 118 cuts.  
Todd and Bowman's Physiological Anatomy and Physiology of Man, with numerous wood-cuts. Parts I., II. and III., 1 vol. 8vo., 156 wood-cuts.

### PATHOLOGY.

Abercrombie on the Brain, 1 vol. 8vo., 324 pp.  
Blakiston on Diseases of the Chest, 1 vol., 384 pp.  
Blood and Urine Manuals, by Reese, Griffith, Markwick, Bird, and Frick, 2 vols. 12mo., many cuts and plates.  
Budd on the Liver, 1 vol. 8vo., 392 pages, plates and wood-cuts.  
Burrows on Cerebral Circulation, 1 vol. 8vo., 216 pages, with 6 colored plates.  
Billing's Principles, new and improved edition, 1 vol. 8vo., 250 pages, (just issued.)  
Bird on Urinary Deposits, 12mo., new and improved edition, (just issued.)  
Copland on Palsy and Apoplexy, 1 vol. 12mo., 236 pp.  
Frick on Renal Affections, 1 vol. 12mo., cuts.  
Hasse's Pathological Anatomy, 8vo., 379 pages.  
Hope on the Heart, new ed., pl's, 1 vol. 8vo., 572 p.  
Hughes on the Lungs, &c., 1 vol. 12mo., 270 pp.

Lallemand on Spermatorrhœa, 1 vol. 8vo., 320 pp.  
Mitchell on Fevers, 1 vol. 12mo., 138 pages.  
Philip on Protracted Indigestion, 8vo., 240 pp.  
Philips on Scrofula, 1 vol. 8vo., 350 pages.  
Ricord on Venereal, new ed., 1 vol. 8vo., 340 pp.  
Stanley on Diseases of the Bones, 1 vol. 8vo., 286 pages.  
Vögel's Pathological Anatomy of the Human Body, 1 vol. 8vo., 536 pages, col. plates.  
Wilson on the Skin, 1 vol. 8vo., new ed., 440 pp. Same work, with colored plates.  
Whitehead on Sterility and Abortion, 1 vol. 8vo., 368 pages.  
Williams' Principles of Medicine, by Clymer, 2d edition, 440 pages, 1 vol. 8vo.  
Williams on the Respiratory Organs, by Clymer, 1 vol. 8vo., 500 pages.

### PRACTICE OF MEDICINE.

Ashwell on Females, 2d ed., 1 vol. 8vo., 520 pp.  
Barlow's Practice of Medicine, (preparing.)  
Bennet on the Uterus, 2d and enlarged edition, 1 vol. 8vo., 356 pages.  
Bartlett on Fevers, 2d edition, 550 pages.  
Benedict's Compendium of Chapman's Lectures, 1 vol. 8vo., 258 pages.  
Chapman on Fevers, Gout, Dropsy, &c. &c., 1 vol. 8vo., 450 pages.  
Colombat de L'Isère on Females, by Meigs, 1 vol. 8vo., 720 pages, cuts. New edition.  
Condie on the Diseases of Children, 3d edition, 1 vol. 8vo.  
Churchill on the Diseases of Infancy and Childhood, 1 vol. 8vo.  
Churchill on the Diseases of Females, by Huston, 5th ed., rev'd by the author, 1 vol. 8vo., 632 pp.  
Churchill's Monographs of the Diseases of Females, 1 vol. 8vo., now ready, 450 pages.  
Clymer and others on Fevers, a complete work in 1 vol. 8vo., 600 pages.  
Day on Old Age, 1 vol. 8vo., 226 pages.  
Dewees on Children, 9th ed., 1 vol. 8vo., 548 pp.  
Dewees on Females, 9th ed., 1 vol. 8vo., 532 p. pls.  
Dunglison's Practice of Medicine, 3d edition, 2 vols. 8vo., 1500 pages.  
Esquirol on Insanity, by Hunt, 8vo., 496 pages.  
Meigs' Letters on Diseases of Females, 1 vol. 8vo., 690 pp., 2d ed., improved, (lately issued.)  
Meigs on Certain Diseases of Infancy, 1 vol. 8vo., 216 pp., (a new work.)  
Thomson on the Sick Room, &c., 1 vol. large 12mo., 360 pages, cuts.  
Watson's Principles and Practice of Physic, 3d edition by Condie, 1 vol. 8vo., 1060 large pages.  
West's Lectures on the Diseases of Infancy and Childhood. 1 vol. 8vo., 452 pp.  
Walshe on the Heart and Lungs. A new work, now ready, 1 vol. royal 12mo. ex. cloth.

### SURGERY.

Brodie on Urinary Organs, 1 vol. 8vo., 214 pages.  
Brodie on the Joints, 1 vol. 8vo., 216 pages.  
Brodie's Lectures on Surgery, 1 vol. 8vo., 350 pp.  
Brodie's Select Surgical Works, 780 pp. 1 vol. 8vo.  
Chelius' System of Surgery, by South and Norris, in 3 large 8vo. vols., near 2200 pages.  
Cooper on Dislocations and Fractures, 1 vol. 8vo., 500 pages, many cuts.  
Cooper on Hernia, 1 vol. imp. 8vo., many plates.  
Cooper on the Testis and Thymus Gland, 1 vol. imperial 8vo., many plates.  
Cooper on the Anatomy and Diseases of the Breast, Surgical Papers, &c. &c., 1 vol. imp. 8vo., pl'ts.  
Druitt's Principles and Practice of Modern Surgery, 1 vol. 8vo., 576 pages, 193 cuts, 4th ed.



Duften on Deafness and Diseases of the Ear, 1 vol. 12mo., 120 pages.  
 Durlacher on Corns, Bunions, &c., 12mo., 134 pp.  
 Ear, Diseases of, a new work, (preparing.)  
 Fergusson's Practical Surgery, 1 vol. 8vo., 3d edition, 630 pages, 274 cuts.  
 Guthrie on the Bladder, 8vo., 150 pages.  
 Gross on Injuries and Diseases of Urinary Organs, 1 lrg. vol. 8vo., 726 pp. many cuts, (now ready.)  
 Jones' Ophthalmic Medicine and Surgery, by Hays, 1 vol. 12mo., 529 pp., cuts and plates.  
 Liston's Lectures on Surgery, by Mütter, 1 vol. 8vo., 566 pages, many cuts.  
 Lawrence on the Eye, by Hays, new ed. much improved, 863 pp., many cuts and plates.  
 Lawrence on Ruptures, 1 vol. 8vo., 480 pages.  
 Miller's Principles of Surgery, 2d edition, 1 vol. 8vo., 538 pp., 1848.  
 Miller's Practice of Surgery, 1 vol. 8vo., 496 pp.  
 Malgaigne's Operative Surgery, by Brittan, with cuts. Now complete in one 8vo. vol., of about 600 pages.  
 Maury's Dental Surgery, 1 vol. 8vo., 286 pages, many plates and cuts.  
 Skey's Operative Surgery, 1 vol. large 8vo., many cuts, 662 pages, a new work, (just issued.)  
 Sargent's Minor Surgery, 1 vol. royal 12mo., 380 pages, 128 cuts.  
 Smith on Fractures, 1 vol. 8vo., 200 cuts, 314 pp.

#### MATERIA MEDICA AND THERAPEUTICS.

Bird's (Golding) Therapeutics, (preparing.)  
 Christison's and Griffith's Dispensatory, 1 large vol. 8vo., 216 cuts, over 1000 pages.  
 Carpenter on Alcoholic Liquors in Health and Disease, 1 vol. 12mo.  
 Carson's Synopsis of Lectures on Materia Medica and Pharmacy in the University of Pennsylvania, 1 vol. 8vo., 208 pages, (now ready.)  
 Dunglison's Materia Medica and Therapeutics, now ready, 4th ed., much improved, 182 cuts, 2 vols. 8vo., 1850.  
 Dunglison on New Remedies, 6th ed., much improved, 1 vol. 8vo., 750 pages.  
 De Jongh on Cod-Liver Oil, 12mo.  
 Ellis' Medical Formulary, 9th ed., much improved, 1 vol. 8vo., 268 pages.  
 Griffith's Universal Formulary, 1 large vol. 8vo., 560 pages.  
 Griffith's Medical Botany, a new work, 1 large vol. 8vo., 704 pp., with over 350 illustrations.  
 Mayne's Dispensatory, 1 vol. 12mo., 330 pages.  
 Mohr, Redwood, and Procter's Pharmacy, 1 vol. 8vo., 550 pages, 506 cuts.  
 Pereira's Materia Medica, by Carson, 3d ed., 2 vols. 8vo., much improved and enlarged, with 400 wood cuts, (nearly ready.)  
 Royle's Materia Medica and Therapeutics, by Carson, 1 vol. 8vo., 689 pages, many cuts.

#### OBSTETRICS.

Churchill's Theory and Practice of Midwifery, a new and improved ed., by Condie, 1 vol. 8vo., 510 pp., many cuts, (just issued.)  
 Dewees' Midwifery, 11th ed., 1 vol. 8vo., 660 pp., plates.  
 Lee's Clinical Midwifery, 12mo., 238 pages.  
 Meigs' Obstetrics; the Science and the Art; 1 vol. 8vo., 686 pages, 121 cuts.  
 Ramsbotham on Parturition, with many plates, 1 large vol., imperial 8vo., 520 pp. 6th edition, (now ready.)  
 Rigby's Midwifery, new edition, 1 vol. 8vo., (just issued,) 422 pages.  
 Smith (Tyler) on Parturition, 1 vol. 12mo., 400 pp.

#### CHEMISTRY AND HYGIENE.

Bowman's Practical Chemistry, 1 vol. 12mo., 97 cuts, 350 pages.  
 Brigham on Excitement, &c., 1 vol. 12mo., 204 pp.

Beale on Health of Mind and Body, 1 vol. 12mo., extra cloth, (now ready.)  
 Bowman's Medical Chemistry, 1 vol. 12mo., many cuts, just ready, 288 pages.  
 Dunglison on Human Health, 2d ed., 8vo., 464 pp.  
 Fowne's Elementary Chemistry, 3d ed., 1 vol. 12mo., much improved, many cuts, now ready.  
 Graham's Chemistry, by Bridges, new and improved edition. Part I, (in press.)  
 Gardner's Medical Chemistry, 1 vol. 12mo. 400 pp.  
 Griffith's Chemistry of the Four Seasons, 1 vol. royal 12mo., 451 pages, many cuts.  
 Knapp's Chemical Technology, by Johnson, 2 vols. 8vo., 936 pp., 460 large cuts.  
 Simon's Chemistry of Man, 8vo., 730 pp., plates.

#### MEDICAL JURISPRUDENCE, EDUCATION, &c.

Bartlett's Philosophy of Medicine, 1 vol. 8vo., 312 pages.  
 Bartlett on Certainty in Medicine, 1 vol. small 8vo., 84 pages.  
 Dunglison's Medical Student, 2d ed. 12mo., 312 pp.  
 Taylor's Medical Jurisprudence, by Griffith, 1 vol. 8vo., new edition, 1850, 670 pp.  
 Taylor on Poisons, by Griffith, 1 vol. 8vo., 688 pp.

#### NATURAL SCIENCE, &c.

Arnott's Physics, 1 vol. 8vo., 484 pp., many cuts.  
 Ansted's Ancient World, Popular Geology, in 1 12mo. volume, with numerous cuts, 382 pages.  
 Bird's Natural Philosophy, 1 vol. royal 12mo., 402 pages and 372 wood-cuts.  
 Brewster's Optics, 1 vol. 12mo. 423 pp. many cuts.  
 Broderip's Zoological Recreations, 1 vol. 12mo., 376 pp.  
 Coleridge's Idea of Life, 12mo., 94 pages.  
 Carpenter's General and Comparative Physiology, 1 large 8vo. vol., many wood-cuts, (now ready.)  
 Dana on Zoophytes, being vol. 8 of Ex. Expedition, royal 4to., extra cloth.  
 Atlas to "Dana on Zoophytes," im. fol., col. pl's.  
 Gregory on Animal Magnetism, 1 vol., royal 12mo., (now ready.)  
 De la Beche's Geological Observer, 1 large 8vo. vol., many wood-cuts, (just ready.)  
 Hale's Ethnography and Philology of the U. S. Exploring Expedition, in 1 large imp. 4to. vol.  
 Herschel's Treatise on Astronomy, 1 vol. 12mo., 417 pages, numerous plates and cuts.  
 Herschel's Outlines of Astronomy, 1 vol. small 8vo., plates and cuts. (A new work.) 620 pp.  
 Humboldt's Aspects of Nature, 1 vol. 12mo., new edition.  
 Johnston's Physical Atlas, 1 vol. imp. 4to., half bound, 25 colored maps.  
 Kirby and Spence's Entomology, 1 vol. 8vo., 600 large pages; plates plain or colored.  
 Knox on Races of Men, 1 vol. 12mo.  
 Lardner's Handbooks of Natural Philosophy, 2 vols. royal 12mo., with 800 cuts, (in press.)  
 Müller's Physics and Meteorology, 1 vol. 8vo., 636 pp., with 540 wood-cuts and 2 col'd plates.  
 Small Books on Great Subjects, 12 parts, done up in 3 handsome 12mo. volumes, extra cloth.  
 Somerville's Physical Geography, 1 vol. 12mo., cloth, 540 pages, enlarged edition, now ready.  
 Weisbach's Mechanics applied to Machinery and Engineering, Vol. I. 8vo., 486 p. 550 wood-cuts.  
 Vol. II., 8vo., 400 pp., 340 cuts.

#### VETERINARY MEDICINE.

Clater and Skinner's Farrier, 1 vol. 12mo., 220 pp.  
 Youatt's Great Work on the Horse, by Skinner, 1 vol. 8vo., 448 pages, many cuts.  
 Youatt and Clater's Cattle Doctor, 1 vol. 12mo., 282 pages, cuts.  
 Youatt on the Dog, by Lewis, 1 vol. demy 8vo., 403 pages, beautiful plates.  
 Youatt on the Pig, a new work, with beautiful illustrations of all the different varieties, 12mo.

Other new and important works are in preparation.



TWO MEDICAL PERIODICALS FOR FIVE DOLLARS.

# THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES,

EDITED BY ISAAC HAYS, M. D.,

IS PUBLISHED QUARTERLY, ON THE FIRST OF JANUARY, APRIL, JULY, AND OCTOBER,

*By BLANCHARD & LEA, Philadelphia.*

Each Number contains about Two Hundred and Eighty Large Octavo Pages,

Appropriately Illustrated with Engravings on Copper, Wood, Stone, &c.

## THE MEDICAL NEWS AND LIBRARY

Is Published Monthly, and consists of

THIRTY-TWO VERY LARGE OCTAVO PAGES,

Containing the Medical Information of the day, as well as a Treatise of high character on some prominent department of Medicine.

In this manner its subscribers have been supplied with

WATSON'S LECTURES ON THE PRACTICE OF MEDICINE,

BRODIE'S CLINICAL LECTURES ON SURGERY,

TODD & BOWMAN'S PHYSIOLOGY,

AND WEST ON THE DISEASES OF INFANCY AND CHILDHOOD.

And the work at present appearing in its columns is

**MALGAIGNE'S OPERATIVE SURGERY,**

TRANSLATED AND EDITED BY BRITTAN,

With Engravings on Wood.

Which will be completed in the present year, and be succeeded, in 1852, by a work of equal value.

### TERMS.

THE SUBSCRIPTION TO THE

## AMERICAN JOURNAL OF THE MEDICAL SCIENCES IS FIVE DOLLARS PER ANNUM.

When this amount is paid in advance, the subscriber thereby becomes entitled to the

## MEDICAL NEWS AND LIBRARY FOR ONE YEAR, GRATIS.

When ordered separately, the price of the "News" is ONE DOLLAR per annum, invariably in advance.

For the small sum, therefore, of FIVE DOLLARS, the subscriber can obtain a Quarterly and a Monthly Journal of the highest character, presenting about

**Fifteen hundred large octavo pages, with numerous Illustrations,**

Rendering these among

## The Cheapest of American Medical Periodicals.

Those who are desirous of subscribing are recommended to forward their names without loss of time, as the increase of the subscription list has almost exhausted the whole edition printed for the present year, and the publishers cannot pledge themselves to supply copies unless ordered early.

### REDUCTION OF POSTAGE.

Under the new postage law, subscribers to the American Journal of the Medical Sciences are entitled to the benefit of the half rates of postage for advance payment, which is thus defined in the Post Master General's circular of June 11th, 1851:—

"When a periodical is published only quarterly, the actual and bona fide subscriber to such periodical may pay in advance, and have the benefit of such advance payment, provided he pays to the postmaster at the office where he is to receive the periodical, before its delivery."

It will therefore be seen that the subscriber has only to pay for each number before taking it out of the office, in order to secure the benefit of the half postage, which is as follows:—

For any distance under 500 miles, - - - ½ cent per ounce,

" " between 500 and 1500 miles, - 1 " "

" " " 1500 and 2500 " - 1½ " "

When the postage is not thus paid in advance, it will be at double these rates.

The Medical News and Library pays postage as a newspaper. Each number weighs between one and two ounces. Subscribers will therefore, under the new postage law, pay in advance for each quarter as follows:—

For any distance under 50 miles, - - - 1½ cents per quarter.

" " between 50 and 300 miles, - 2½ " "

" " " 300 and 1000 miles, - 3½ " "

" " " 1000 and 2000 miles, 5 " "



NEILL & SMITH'S COMPENDIUM.  
NEW EDITION, Nearly Ready.

**AN ANALYTICAL COMPENDIUM**  
OF THE VARIOUS BRANCHES OF MEDICAL SCIENCE,  
FOR THE USE AND EXAMINATION OF STUDENTS.

BY JOHN NEILL, M. D.,

Demonstrator of Anatomy in the University of Pennsylvania, Lecturer on Anatomy in the Medical Institute of Philadelphia, &c.,

AND

FRANCIS GURNEY SMITH, M. D.,

Lecturer on Physiology in the Philadelphia Association for Medical Instruction, &c. &c.

*Second Edition, Revised and Improved.*

*In one very large and handsomely printed volume, royal 12mo., of over 900 large pages, with about 350 illustrations, strongly bound in leather, with raised bands.*

The rapid exhaustion of a large edition of this work is sufficient evidence that it has fulfilled the object for which it was designed by the authors, not only as a manual for the use and examination of students, enabling them to recall the facts presented in the lecture room or in the larger textbooks, but also as an aid to the practitioner who may be desirous of refreshing his knowledge by a volume furnishing a rapid, but exact, outline of all the various branches of medical science. In the preparation of the new edition every effort has been made to keep it on a level with the advance of investigation and discovery in all its different departments, and the experience which its use has afforded, has been employed to adapt it still further to the wants of the student. To accomplish this it has been thoroughly revised, and in many parts remodelled and rewritten, while additional illustrations have been introduced, where necessary. An improvement will also be found in its mechanical execution, while the very large size of the pages, and the abundance of illustrative engravings render it one of the cheapest works offered to the notice of the profession.

The arrangement adopted will be found at once concise and clear; while its mechanical execution, its copious pictorial illustrations in the branches of anatomy, physiology, surgery, obstetrics, materia medica and chemistry, together with its neat, cheap, and convenient form, will recommend it to all such students and practitioners who may desire to avail themselves of what cannot fail to prove, if kept within its proper sphere, a convenient and useful remembrance.—*American Journal of the Medical Sciences.*

We do not share in the opinion entertained by some, that compendiums of science are not desirable, or with the still smaller number, who esteem them useless. On the contrary, when well executed, they are of essential service to the student. Taking the work before us, we can certainly say that no one who has not occupied himself with the different scientific treatises and essays that have appeared recently, and has withal a rare memory, could pretend to possess the knowledge contained in it; and hence we can recommend it to such—as well as to students especially—for its general accuracy and adequacy for their purposes; and to the well-informed practitioner to aid him in recalling what may easily have passed from his remembrance. We repeat our favorable impression as to the value of this book, or series of books; and recommend it as decidedly useful to those especially who are commencing the study of their profession.—*The Medical Examiner.*

We have no hesitation in recommending it to students.—*Southern Medical and Surgical Journal.*

Books of this description are most erroneously denounced, from the supposition that they are intended to take the place of elaborate treatises; but their object is rather to assist the student in mastering the elements of medicine, and to aid the practitioner by refreshing his recollection of former studies. We have looked through this compendium, and we find that the authors have really succeeded in compressing a large amount of valuable information into a very small compass. We recommend this work especially to the notice of our junior readers. To those who are about to commence their studies in a medical school it will be found a serviceable guide.—*London Medical Gazette.*

PROFESSOR DICKSON'S ESSAYS—Nearly Ready.

**ESSAYS ON LIFE, SLEEP, PAIN, INTELLECTION, HYGIENE, AND DEATH.**

BY SAMUEL HENRY DICKSON, M. D.,

Professor of the Institutes and Practice of Medicine in the Charleston Medical College.

*In one very handsome volume, royal 12mo.*

**MALGAIGNE'S SURGERY.—Now Ready.**

**OPERATIVE SURGERY,**  
BASED ON NORMAL AND PATHOLOGICAL ANATOMY.

BY J. F. MALGAIGNE.

TRANSLATED FROM THE FRENCH,

BY FREDERICK BRITTAN, A. B., M. D., M. R. C. S. L.

WITH NUMEROUS ILLUSTRATIONS ON WOOD.

*In one handsome octavo volume of nearly 600 pages.*

This work has, during its passage through the columns of the "Medical News and Library" in 1850 and 1851, received the unanimous approbation of the profession, and in presenting it in a complete form the publishers confidently anticipate for it an extended circulation.

Certainly one of the best books published on operative surgery.—*Edinburgh Med. Journal.*

We can strongly recommend it both to practitioners and students, not only as a safe guide in the dissecting-room or operating-theatre, but also as a concise work of reference for all that relates to operative surgery.—*Forbes' Review.*

Dr. Brittan has performed his task of translator and editor with much judgment. The descriptions are perfectly clear and explicit; and the author's occasional omissions of important operations proposed by British surgeons are judiciously supplied in brief notes.—*Medical Gazette.*



**GROSS ON URINARY ORGANS—(Just Issued.)****A PRACTICAL TREATISE ON THE  
DISEASES AND INJURIES OF THE URINARY ORGANS.**

BY S. D. GROSS, M. D., &c.,  
Professor of Surgery in the New York University.

In one large and beautifully printed octavo volume, of over seven hundred pages.

With numerous Illustrations.

The author of this work has devoted several years to its preparation, and has endeavored to render it complete and thorough on all points connected with the important subject to which it is devoted. It contains a large number of original illustrations, presenting the natural and pathological anatomy of the parts under consideration, instruments, modes of operation, &c. &c., and in mechanical execution it is one of the handsomest volumes yet issued from the American press.

A very condensed summary of the contents is subjoined.

INTRODUCTION.—CHAPTER I. Anatomy of the Perinæum.—CHAPTER II. Anatomy of the Urinary Bladder.—CHAPTER III. Anatomy of the Prostate.—CHAPTER IV. Anatomy of the Urethra.—CHAPTER V. Urine.

PART I. DISEASES AND INJURIES OF THE BLADDER.

CHAPTER I. Malformations and Imperfections.—CHAPTER II. Injuries of the Bladder.—CHAPTER III. Inflammation of the Bladder.—CHAPTER IV. Chronic Lesions of the Bladder.—CHAPTER V. Nervous Affections of the Bladder.—CHAPTER VI. Heterologous Formations of the Bladder.—CHAPTER VII. Polypous, Fungous, Erectile, and other Morbid Growths of the Bladder.—CHAPTER VIII. Worms in the Bladder.—CHAPTER IX. Serous Cysts and Hydatids.—CHAPTER X. Fœtal Remains in the Bladder.—CHAPTER XI. Hair in the Bladder.—CHAPTER XII. Air in the Bladder.—CHAPTER XIII. Hemorrhage of the Bladder.—CHAPTER XIV. Retention of Urine.—CHAPTER XV. Incontinence of Urine.—CHAPTER XVI. Hernia of the Bladder.—CHAPTER XVII. Urinary Deposits.—CHAPTER XVIII. Stone in the Bladder.—CHAPTER XIX. Foreign Bodies in the Bladder.

PART II. DISEASES AND INJURIES OF THE PROSTATE GLAND.

CHAPTER I. Wounds of the Prostate.—CHAPTER II. Acute Prostatitis.—CHAPTER III. Hypertrophy of the Prostate.—CHAPTER IV. Atrophy of the Prostate.—CHAPTER V. Heterologous Formations of the Prostate.—CHAPTER VI. Cystic Disease of the Prostate.—CHAPTER VII. Fibrous Tumors of the Prostate.—CHAPTER VIII. Hemorrhage of the Prostate.—CHAPTER IX. Calculi of the Prostate.—CHAPTER X. Phlebitis of the Prostate.

PART III. DISEASES AND INJURIES OF THE URETHRA.

CHAPTER I. Malformations and Imperfections of the Urethra.—CHAPTER II. Laceration of the Urethra.—CHAPTER III. Stricture of the Urethra.—CHAPTER IV. Polypoid and Vascular Tumors of the Urethra.—CHAPTER V. Neuralgia of the Urethra.—CHAPTER VI. Hemorrhage of the Urethra.—CHAPTER VII. Foreign Bodies in the Urethra.—CHAPTER VIII. Infiltration of Urine.—CHAPTER IX. Urinary Abscess.—CHAPTER X. Fistula of the Urethra.—CHAPTER XI. False Passages.—CHAPTER XII. Lesions of the Gallinaginous Crest.—CHAPTER XIII. Inflammation and Abscess of Cowper's Glands.

**COOPER ON DISLOCATIONS.—New Edition—(Just Issued.)****A TREATISE ON  
DISLOCATIONS AND FRACTURES OF THE JOINTS.**

By SIR ASTLEY P. COOPER, BART., F. R. S., &c.

EDITED BY BRANSBY B. COOPER, F. R. S., &c.

WITH ADDITIONAL OBSERVATIONS BY PROF. J. C. WARREN.

A NEW AMERICAN EDITION,

In one handsome octavo volume, with numerous illustrations on wood.

After the fiat of the profession, it would be absurd in us to eulogize Sir Astley Cooper's work on Dislocations. It is a national one, and will probably subsist as long as English Surgery.—*Medico-Chirurg. Review.*

**WORKS BY THE SAME AUTHOR.**

COOPER (SIR ASTLEY) ON THE ANATOMY AND TREATMENT OF ABDOMINAL HERNIA. 1 large vol., imp. 8vo., with over 130 lithographic figures.

COOPER ON THE STRUCTURE AND DISEASES OF THE TESTIS, AND ON THE THYMUS GLAND. 1 vol., imp. 8vo., with 177 figures on 29 plates.

COOPER ON THE ANATOMY AND DISEASES OF THE BREAST, WITH TWENTY-FIVE MISCELLANEOUS AND SURGICAL PAPERS. 1 large vol., imp. 8vo., with 252 figures on 36 plates.

These three volumes complete the surgical writings of Sir Astley Cooper. They are very handsomely printed, with a large number of lithographic plates, executed in the best style, and are presented at exceedingly low prices.

**LISTON & MUTTER'S SURGERY.****LECTURES ON THE OPERATIONS OF SURGERY,  
AND ON DISEASES AND ACCIDENTS REQUIRING OPERATIONS.**

BY ROBERT LISTON, Esq., F. R. S., &c.

EDITED, WITH NUMEROUS ADDITIONS AND ALTERATIONS,

BY T. D. MUTTER, M. D.,

Professor of Surgery in the Jefferson Medical College of Philadelphia.

In one large and handsome octavo volume of 566 pages, with 216 wood-cuts.



## LIBRARY OF SURGICAL KNOWLEDGE.

**A SYSTEM OF SURGERY.**

BY J. M. CHELIUS.

TRANSLATED FROM THE GERMAN,  
AND ACCOMPANIED WITH ADDITIONAL NOTES AND REFERENCES,  
BY JOHN F. SOUTH.

Complete in three very large octavo volumes of nearly 2200 pages, strongly bound, with raised bands and double titles: or in seventeen numbers, at fifty cents each.

We do not hesitate to pronounce it the best and most comprehensive system of modern surgery with which we are acquainted.—*Medico-Chirurgical Review*.The fullest and ablest digest extant of all that relates to the present advanced state of Surgical Pathology.—*American Medical Journal*.If we were confined to a single work on Surgery, that work should be Chelius's.—*St. Louis Med. Journal*.As complete as any system of Surgery can well be.—*Southern Medical and Surgical Journal*.The most finished system of Surgery in the English language.—*Western Lancet*.The most learned and complete systematic treatise now extant.—*Edinburgh Medical Journal*.No work in the English language comprises so large an amount of information relative to operative medicine and surgical pathology.—*Medical Gazette*.A complete encyclopedia of surgical science—a very complete surgical library—by far the most complete and scientific system of surgery in the English language.—*N. Y. Journal of Medicine*.One of the most complete treatises on Surgery in the English language.—*Monthly Journal of Med. Science*.The most extensive and comprehensive account of the art and science of Surgery in our language.—*Lancet*.**A TREATISE ON THE DISEASES OF THE EYE.**

BY W. LAWRENCE, F. R. S.

*A new Edition. With many Modifications and Additions, and the introduction of nearly 200 Illustrations,*

BY ISAAC HAYS, M. D.

In one very large 8vo. vol. of 860 pages, with plates and wood-cuts through the text.

**JONES ON THE EYE.****THE PRINCIPLES AND PRACTICE  
OF OPHTHALMIC MEDICINE AND SURGERY.**

BY T. WHARTON JONES, F. R. S., &amp;c. &amp;c.

EDITED BY ISAAC HAYS, M. D., &amp;c.

In one very neat volume, large royal 12mo. of 529 pages, with four plates, plain or colored, and ninety-eight well executed wood-cuts.

## MILLER'S PRINCIPLES AND PRACTICE OF SURGERY.

**THE PRINCIPLES OF SURGERY.**

Second edition, one vol. 8vo.

**THE PRACTICE OF SURGERY.**

Second edition, one vol. 8vo.

BY JAMES MILLER, F. R. S. E.,

Professor of Surgery in the University of Edinburgh, &amp;c.

STANLEY ON THE BONES.—A Treatise on Diseases of the Bones. In one vol. 8vo., extra cloth. 286 pp.

BRODIE'S SURGICAL LECTURES.—Clinical Lectures on Surgery. 1 vol. 8vo., cloth. 350 pp.

BRODIE ON THE JOINTS.—Pathological and Surgical Observations on the Diseases of the Joints. 1 vol. 8vo., cloth. 216 pp.

BRODIE ON URINARY ORGANS.—Lectures on the Diseases of the Urinary Organs. 1 vol. 8vo., cloth. 214 pp.

\* \* \* These three works may be had neatly bound together, forming a large volume of "Brodie's Surgical Works." 780 pp.

RICORD ON VENEREAL.—A Practical Treatise on Venereal Diseases. With a Therapeutical Summary and Special Formulary. Translated by Sidney Doane, M. D. Fourth edition. 1 vol. 8vo. 340 pp.

DURLACHER ON CORNS, BUNIONS, &amp;c.—A Treatise on Corns, Bunions, the Diseases of Nails, and the General Management of the Feet. In one 12mo. volume, cloth. 134 pp.

GUTHRIE ON THE BLADDER, &amp;c.—The Anatomy of the Bladder and Urethra, and the Treatment of the Obstructions to which those Passages are liable. In one vol. 8vo. 150 pp.

LAWRENCE ON RUPTURES.—A Treatise on Ruptures, from the fifth London Edition. In one 8vo. vol. sheep. 480 pp.

MAURY'S DENTAL SURGERY.—A Treatise on the Dental Art, founded on Actual Experience. Illustrated by 241 lithographic figures and 54 wood-cuts. Translated by J. B. Savier. In 1 8vo. vol., sheep. 286 pp.

DUFTON ON THE EAR.—The Nature and Treatment of Deafness and Diseases of the Ear; and the Treatment of the Deaf and Dumb. One small 12mo. volume. 120 pp.

SMITH ON FRACTURES.—A Treatise on Fractures in the vicinity of Joints, and on Dislocations. One vol. 8vo., with 200 beautiful wood-cuts.



NEW AND IMPORTANT WORK ON PRACTICAL SURGERY.—(JUST ISSUED.)

**OPERATIVE SURGERY.**

BY FREDERICK C. SKEY, F. R. S., &amp;c.

In one very handsome octavo volume of over 650 pages, with about one hundred wood-cuts.

The object of the author, in the preparation of this work, has been not merely to furnish the student with a guide to the actual processes of operation, embracing the practical rules required to justify an appeal to the knife, but also to present a manual embodying such *principles* as might render it a permanent work of reference to the practitioner of operative surgery, who seeks to uphold the character of his profession as a science as well as an art. In its composition he has relied mainly on his own experience, acquired during many years' service at one of the largest of the London hospitals, and has rarely appealed to other authorities, except so far as personal intercourse and a general acquaintance with the most eminent members of the surgical profession have induced him to quote their opinions.

*From Professor C. B. Gibson, Richmond, Virginia.*

I have examined the work with some care, and am delighted with it. The style is admirable, the matter excellent, and much of it original and deeply interesting, whilst the illustrations are numerous and better executed than those of any similar work I possess.

In conclusion we must express our unqualified praise of the work as a whole. The high moral tone, the liberal views, and the sound information which pervades it throughout, reflect the highest credit upon the talented author. We know of no one who has succeeded, whilst supporting operative surgery in its proper rank, in promulgating at the same time sounder and more enlightened views upon that most important of all subjects, the principle that should guide us in having recourse to the knife.—*Medical Times.*

The treatise is, indeed, one on operative surgery, but it is one in which the author throughout shows that he is most anxious to place operative surgery in its just position. He has acted as a judicious, but not partial friend; and while he shows throughout that he is able and ready to perform any operation which the exigencies and casualties of the human frame may require, he is most cautious in specifying the circumstances which in each case indicate and contraindicate operation. It is indeed gratifying to perceive the sound and correct views which Mr. Skey entertains on the subject of operations in general, and the gentlemanly tone in which he impresses on readers the lessons which he is desirous to inculcate. His work is a perfect model for the operating surgeon, who will learn from it not only when and how to operate, but some more noble and exalted lessons which cannot fail to improve him as a moral and social agent.—*Edinburgh Medical and Surgical Journal.*

## THE STUDENT'S TEXT-BOOK.

**THE PRINCIPLES AND PRACTICE OF MODERN SURGERY.**

BY ROBERT DRUITT, Fellow of the Royal College of Surgeons.

**A New American, from the last and improved London Edition.**

EDITED BY F. W. SARGENT, M. D., Author of "Minor Surgery," &amp;c.

ILLUSTRATED WITH ONE HUNDRED AND NINETY-THREE WOOD ENGRAVINGS.

In one very handsomely printed octavo volume of 576 large pages.

*From Professor Brainard, of Chicago, Illinois.*

I think it the best work of its size, on that subject, in the language.

*From Professor Rivers, of Providence, Rhode Island.*

I have been acquainted with it since its first republication in this country, and the universal praise it has received I think well merited.

*From Professor May, of Washington, D. C.*

Permit me to express my satisfaction at the republication in so improved a form of this most valuable work. I believe it to be one of the very best text-books ever issued.

*From Professor McCook, of Baltimore.*

I cannot withhold my approval of its merits, or the expression that no work is better suited to the wants of the student. I shall commend it to my class, and make it my chief text-book.

FERGUSON'S OPERATIVE SURGERY. NEW EDITION.

**A SYSTEM OF PRACTICAL SURGERY.**

BY WILLIAM FERGUSSON, F. R. S. E.,

Professor of Surgery in King's College, London, &amp;c. &amp;c.

THIRD AMERICAN, FROM THE LAST ENGLISH EDITION.

With 274 Illustrations.

In one large and beautifully printed octavo volume of six hundred and thirty pages.

It is with unfeigned satisfaction that we call the attention of the profession in this country to this excellent work. It richly deserves the reputation conceded to it, of being the best practical Surgery extant, at least in the English language.—*Medical Examiner.*

**A NEW MINOR SURGERY.****ON BANDAGING AND OTHER POINTS OF MINOR SURGERY.**

BY F. W. SARGENT, M. D.

In one handsome royal 12mo. volume of nearly 400 pages, with 128 wood-cuts.

*From Professor Gilbert, Philadelphia.*

Embracing the smaller details of surgery, which are illustrated by very accurate engravings, the work becomes one of very great importance to the practitioner in the performance of his daily duties, since such information is rarely found in the general works on surgery now in use.



THE GREAT ATLAS OF SURGICAL ANATOMY.  
(NOW COMPLETE.)

**SURGICAL ANATOMY.**

BY JOSEPH MACLISE, SURGEON.

IN ONE VOLUME, IN VERY LARGE IMPERIAL QUARTO.

**With Sixty-eight large and splendid Plates, drawn in the best style, and beautifully colored,**

**Containing one hundred and ninety Figures, many of them the size of life.**

TOGETHER WITH COPIOUS EXPLANATORY LETTER-PRESS.

Strongly and handsomely bound, being one of the best executed and cheapest surgical works ever presented in this country.

This great work being now complete, the publishers confidently present it to the attention of the profession as worthy in every respect of their approbation and patronage. No complete work of the kind has yet been published in the English language, and it therefore will supply a want long felt in this country of an accurate and comprehensive Atlas of Surgical Anatomy to which the student and practitioner can at all times refer, to ascertain the exact relative position of the various portions of the human frame towards each other and to the surface, as well as their abnormal deviations. The importance of such a work to the student in the absence of anatomical material, and to the practitioner when about attempting an operation, is evident, while the price of the book, notwithstanding the large size, beauty, and finish of the very numerous illustrations is so low as to place it within the reach of every member of the profession. The publishers therefore confidently anticipate a very extended circulation for this magnificent work.

To present some idea of the scope of the volume, and of the manner in which its plan has been carried out, the publishers subjoin a very brief summary of the plates.

- Plates 1 and 2.—Form of the Thoracic Cavity and Position of the Lungs, Heart, and larger Blood-vessels.
- Plates 3 and 4.—Surgical Form of the Superficial Cervical and Facial Regions, and the Relative Positions of the principal Blood-vessels, Nerves, &c.
- Plates 5 and 6.—Surgical Form of the Deep Cervical and Facial Regions, and Relative Positions of the principal Blood-vessels, Nerves, &c.
- Plates 7 and 8.—Surgical Dissection of the Subclavian and Carotid Regions, and Relative Anatomy of their Contents.
- Plates 9 and 10.—Surgical Dissection of the Sterno-Clavicular or Tracheal Region, and Relative Position of its main Blood-vessels, Nerves, &c.
- Plates 11 and 12.—Surgical Dissection of the Axillary and Brachial Regions, displaying the Relative Order of their contained parts.
- Plates 13 and 14.—Surgical Form of the Male and Female Axillæ compared.
- Plates 15 and 16.—Surgical Dissection of the Bend of the Elbow and the Forearm, showing the Relative Position of the Arteries, Veins, Nerves, &c.
- Plates 17, 18 and 19.—Surgical Dissections of the Wrist and Hand.
- Plates 20 and 21.—Relative Position of the Cranial, Nasal, Oral, and Pharyngeal Cavities, &c.
- Plate 22.—Relative Position of the Superficial Organs of the Thorax and Abdomen.
- Plate 23.—Relative Position of the Deeper Organs of the Thorax and those of the Abdomen.
- Plate 24.—Relations of the Principal Blood-vessels to the Viscera of the Thoracic-Abdominal Cavity.
- Plate 25.—Relations of the Principal Blood-vessels of the Thorax and Abdomen to the Osseous Skeleton, &c.
- Plate 26.—Relation of the Internal Parts to the External Surface of the Body.
- Plate 27.—Surgical Dissection of the Principal Blood-vessels, &c., of the Inguino-Femoral Region.
- Plates 28 and 29.—Surgical Dissection of the First, Second, Third, and Fourth Layers of the Inguinal Region, in connection with those of the Thigh.
- Plates 30 and 31.—The Surgical Dissection of the Fifth, Sixth, Seventh and Eighth Layers of the Inguinal Region, and their connection with those of the Thigh.
- Plates 32, 33 and 34.—The Dissection of the Oblique or External and the Direct or Internal Inguinal Hernia.
- Plates 35, 36, 37 and 38.—The Distinctive Diagnosis between External and Internal Inguinal Hernia, the Taxis, the Seat of Stricture, and the Operation.
- Plates 39 and 40.—Demonstrations of the Nature of Congenital and Infantile Inguinal Hernia, and of Hydrocele.
- Plates 41 and 42.—Demonstrations of the Origin and Progress of Inguinal Hernia in general.
- Plates 43 and 44.—The Dissection of Femoral Hernia, and the Seat of Stricture.
- Plates 45 and 46.—Demonstrations of the Origin and Progress of Femoral Hernia, its Diagnosis, the Taxis, and the Operation.
- Plate 47.—The Surgical Dissection of the principal Blood-vessels and Nerves of the Iliac and Femoral Regions.
- Plates 48 and 49.—The Relative Anatomy of the Male Pelvic Organs.
- Plates 50 and 51.—The Surgical Dissection of the Superficial Structures of the Male Perineum.
- Plates 52 and 53.—The Surgical Dissection of the Deep Structures of the Male Perineum.—The Lateral Operation of Lithotomy.



## MACLISE'S SURGICAL ANATOMY—(Continued.)

- Plates 54, 55 and 56.—The Surgical Dissection of the Male Bladder and Urethra.—Lateral and Bilateral Lithotomy compared.
- Plates 57 and 58.—Congenital and Pathological Deformities of the Prepuce and Urethra.—Structure and Mechanical Obstructions of the Urethra.
- Plates 59 and 60.—The various forms and positions of Strictures and other Obstructions of the Urethra.—False Passages.—Enlargements and Deformities of the Prostate.
- Plates 61 and 62.—Deformities of the Prostate.—Deformities and Obstructions of the Prostatic Urethra.
- Plates 63 and 64.—Deformities of the Urinary Bladder.—The Operations of Sounding for Stone, of Catheterism, and of Puncturing the Bladder above the Pubes.
- Plates 65 and 66.—The Surgical Dissection of the Popliteal Space, and the Posterior Crural Region.
- Plates 67 and 68.—The Surgical Dissection of the Anterior Crural Region, the Ankles, and the Foot.

Notwithstanding the short time in which this work has been before the profession, it has received the unanimous approbation of all who have examined it. From among a very large number of commendatory notices with which they have been favored, the publishers select the following:—

*From Prof. Kimball, Pittsfield, Mass.*

I have examined these numbers with the greatest satisfaction, and feel bound to say that they are altogether, the most perfect and satisfactory plates of the kind that I have ever seen.

*From Prof. Brainard, Chicago, Ill.*

The work is extremely well adapted to the use both of students and practitioners, being sufficiently extensive for practical purposes, without being so expensive as to place it beyond their reach. Such a work was a desideratum in this country, and I shall not fail to recommend it to those within the sphere of my acquaintance.

*From Prof. P. F. Eve, Augusta, Ga.*

I consider this work a great acquisition to my library, and shall take pleasure in recommending it on all suitable occasions.

*From Prof. Peaslee, Brunswick, Me.*

The second part more than fulfils the promise held out by the first, so far as the beauty of the illustrations is concerned; and, perfecting my opinion of the value of the work, so far as it has advanced, I need add nothing to what I have previously expressed to you.

*From Prof. Gunn, Ann Arbor, Mich.*

The plates in your edition of MacLise answer, in an eminent degree, the purpose for which they are intended. I shall take pleasure in exhibiting it and recommending it to my class.

*From Prof. Rivers, Providence, R. I.*

The plates illustrative of Hernia are the most satisfactory I have ever met with.

*From Professor S. D. Gross, Louisville, Ky.*

The work, as far as it has progressed, is most admirable, and cannot fail, when completed, to form a most valuable contribution to the literature of our profession. It will afford me great pleasure to recommend it to the pupils of the University of Louisville.

*From Professor R. L. Howard, Columbus, Ohio.*

In all respects, the first number is the beginning of a most excellent work, filling completely what might be considered hitherto a vacuum in surgical literature. For myself, in behalf of the medical profession, I wish to express to you my thanks for this truly elegant and meritorious work. I am confident that it will meet with a ready and extensive sale. I have spoken of it in the highest terms to my class and my professional brethren.

*From Prof. C. B. Gibson, Richmond, Va.*

I consider MacLise very far superior, as to the drawings, to any work on Surgical Anatomy with which I am familiar, and I am particularly struck with the exceedingly low price at which it is sold. I cannot doubt that it will be extensively purchased by the profession.

*From Prof. Granville S. Pattison, New York.*

The profession, in my opinion, owe you many thanks for the publication of this beautiful work—a work which, in the correctness of its exhibitions of Surgical Anatomy, is not surpassed by any work with which I am acquainted; and the admirable manner in which the lithographic plates have been executed and colored is alike honorable to your house and to the arts in the United States.

*From Prof. J. F. May, Washington, D. C.*

Having examined the work, I am pleased to add my testimony to its correctness, and to its value as a work of reference by the surgeon.

*From Prof. Alden Marsh, Albany, N. Y.*

From what I have seen of it, I think the design and execution of the work admirable, and, at the proper time in my course of lectures, I shall exhibit it to the class, and give it a recommendation worthy of its great merit.

*From H. H. Smith, M. D., Philadelphia.*

Permit me to express my gratification at the execution of MacLise's Surgical Anatomy. The plates are, in my opinion, the best lithographs that I have seen of a medical character, and the coloring of this number cannot, I think, be improved. Estimating highly the contents of this work, I shall continue to recommend it to my class as I have heretofore done.

*From Prof. D. Gilbert, Philadelphia.*

Allow me to say, gentlemen, that the thanks of the profession at large, in this country, are due to you for the republication of this admirable work of MacLise. The precise relationship of the organs in the regions displayed is so perfect, that even those who have daily access to the dissecting-room may, by consulting this work, enliven and confirm their anatomical knowledge prior to an operation. But it is to the thousands of practitioners of our country who cannot enjoy these advantages that the perusal of those plates, with their concise and accurate descriptions, will prove of infinite value. These have supplied a desideratum, which will enable them to refresh their knowledge of the important structures involved in their surgical cases, thus establishing their self-confidence, and enabling them to undertake operative procedures with every assurance of success. And as all the practical departments in medicine rest upon the same basis, and are enriched from the same sources, I need hardly add that this work should be found in the library of every practitioner in the land.



## MACLISE'S SURGICAL ANATOMY—(Continued.)

From Professor J. M. Bush, Lexington, Ky.

I am delighted with both the plan and execution of the work, and shall take all occasions to recommend it to my private pupils and public classes.

The most accurately engraved and beautifully colored plates we have ever seen in an American book—one of the best and cheapest surgical works ever published.—*Buffalo Medical Journal*.

It is very rare that so elegantly printed, so well illustrated, and so useful a work, is offered at so moderate a price.—*Charleston Medical Journal*.

A work which cannot but please the most fastidious lover of surgical science, and we hesitate not to say that if the remaining three numbers of this work are in keeping with the present, it cannot fail to give universal satisfaction. In it, by a succession of plates, are brought to view the relative anatomy of the parts included in the important surgical divisions of the human body, with that fidelity and neatness of touch which is scarcely excelled by nature herself. The part before us differs in many respects from anything of the kind which we have ever seen before. While we believe that nothing but an extensive circulation can compensate the publishers for the outlay in the production of this edition of the work—furnished as it is at a very moderate price, within the reach of all—we desire to see it have that circulation which the zeal and peculiar skill of the author (he being his own draughtsman), the utility of the work, and the neat style with which it is executed, should demand for it in a liberal profession.—*N. Y. Journal of Medicine*.

This is an admirable reprint of a deservedly popular London publication. Its English prototype, although not yet completed, has already won its way, amongst our British brethren, to a remarkable success. Its plates can boast a superiority that places them almost beyond the reach of competition. And we feel too thankful to the Philadelphia publishers for their very handsome reproduction of the whole work, and at a rate within everybody's reach, not to urge all our medical friends to give it, for their own sakes, the cordial welcome it deserves. In a speedy and extensive circulation.—*The Medical Examiner*.

The plates are accompanied by references and explanations, and when the whole has been published it will be a complete and beautiful system of Surgical Anatomy, having an advantage which is important, and not possessed by colored plates generally, viz. its cheapness, which places it within the reach of every one who may feel disposed to possess the work. Every practitioner, we think, should have a work of this kind within reach, as there are many operations requiring immediate performance in which a book of reference will prove most valuable.—*Southern Medical and Surg. Journal*.

The work of MacLise on Surgical Anatomy is of the highest value. In some respects it is the best publication of its kind we have seen, and is worthy of a place in the library of any medical man, while the student could scarcely make a better investment than this.—*The Western Journal of Medicine and Surgery*.

No such lithographic illustrations of surgical regions have hitherto, we think, been given. While the operator is shown every vessel and nerve where an operation is contemplated, the exact anatomist is refreshed by those clear and distinct dissections which every one must appreciate who has a particle of enthusiasm. The English medical press has quite exhausted the words of praise in recommending this admirable treatise. Those who have any curiosity to gratify in reference to the perfectibility of the lithographic art in delineating the complex mechanism of the human body, are invited to examine our specimen copy. If anything will induce surgeons and students to patronize a book of such rare value and every-day importance to them, it will be a survey of the artistical skill exhibited in these fac-similes of nature.—*Boston Medical and Surg. Journal*.

The fidelity and accuracy of the plates reflect the highest credit upon the anatomical knowledge of Mr. MacLise. We strongly recommend the descriptive commentaries to the perusal of the student both of surgery and medicine. These plates will form a valuable acquisition to practitioners settled in the country, whether engaged in surgical, medical, or general practice.—*Edinburgh Medical and Surgical Journal*.

We are well assured that there are none of the cheaper, and but few of the more expensive works on anatomy, which will form so complete a guide to the student or practitioner as these plates. To practitioners, in particular, we recommend this work as far better, and not at all more expensive, than the heterogeneous compilations most commonly in use, and which, whatever their value to the student preparing for examination, are as likely to mislead as to guide the physician in physical examination, or the surgeon in the performance of an operation.—*Monthly Journal of Medical Sciences*.

The dissections from which these various illustrations are taken appear to have been made with remarkable success; and they are most beautifully represented. The surgical commentary is pointed and practical. We know of no work on surgical anatomy which can compete with it.—*Lancet*.

This is by far the ablest work on Surgical Anatomy that has come under our observation. We know of no other work that would justify a student, in any degree, for neglect of actual dissection. A careful study of these plates, and of the commentaries on them, would almost make an anatomist of a diligent student. And to one who has studied anatomy by dissection, this work is invaluable as a perpetual remembrancer, in matters of knowledge that may slip from the memory. The practitioner can scarcely consider himself equipped for the duties of his profession without such a work as this, and this has no rival, in his library. In those sudden emergencies that so often arise, and which require the instantaneous command of minute anatomical knowledge, a work of this kind keeps the details of the dissecting-room perpetually fresh in the memory. We appeal to our readers, whether any one can justifiably undertake the practice of medicine who is not prepared to give all needful assistance, in all matters demanding immediate relief.

We repeat that no medical library, however large, can be complete without MacLise's Surgical Anatomy. The American edition is well entitled to the confidence of the profession, and should command, among them, an extensive sale. The investment of the amount of the cost of this work will prove to be a very profitable one, and if practitioners would qualify themselves thoroughly with such important knowledge as is contained in works of this kind, there would be fewer of them sighing for employment. The medical profession should spring towards such an opportunity as is presented in this republication, to encourage frequent repetitions of American enterprise of this kind.—*The Western Journal of Medicine and Surgery*.

It is a wonderful triumph, showing what ingenuity, skill, and enterprise can effect if supported by a sufficient number of purchasers. No catchpenny sketches on flimsy material and with bad print, but substantial lithographs on fine paper and with a bold and legible type. The drawings are of the first class, and the light and shade so liberally provided for, that the most ample expression, with great clearness and sharpness of outline, is secured.—*Dublin Medical Press*.

Our hearty good wishes attend this work, which promises to supply, when complete, a far better series of delineations of surgical regions than has been yet given, and at a price as low as that of the most ordinary series of illustrations.—*The British and Foreign Medico-Chirurgical Review*.

The plates continue to be of the same excellent character that we have before ascribed to them, and their description all that might naturally be expected from so good an anatomist as Mr. MacLise. The work ought to be in the possession of every one, for it really forms a valuable addition to a surgical library.—*The Medical Times*.

It is, and it must be unique, for the practical knowledge of the surgeon, the patience and skill of the dissector, in combination with the genius of the artist, as here displayed, have never before been, and perhaps, never will be again associated to a similar extent in the same individual.—*Lancet*.

The plates are accurate and truthful; and there is but one word in the English language descriptive of the letter-press—faultless.

For the quality, it is the cheapest work that we have seen, and will constitute a valuable contribution to the surgeon's library.—*The N. W. Medical and Surgical Journal*.



**SHARPEY AND QUAIN'S ANATOMY.—Lately Issued.**

# HUMAN ANATOMY.

BY JONES QUAIN, M. D.

FROM THE FIFTH LONDON EDITION.

EDITED BY

RICHARD QUAIN, F. R. S.,

AND

WILLIAM SHARPEY, M. D., F. R. S.,

Professors of Anatomy and Physiology in University College, London.

**REVISED, WITH NOTES AND ADDITIONS,**

BY JOSEPH LEIDY, M. D.

Complete in Two large Octavo Volumes, of about Thirteen Hundred Pages.

BEAUTIFULLY ILLUSTRATED

With over Five Hundred Engravings on Wood.

We have here one of the best expositions of the present state of anatomical science extant. There is not probably a work to be found in the English language which contains so complete an account of the progress and present state of general and special anatomy as this. By the anatomist this work must be eagerly sought for, and no student's library can be complete without it.—*The N. Y. Journal of Medicine.*

We know of no work which we would sooner see in the hands of every student of this branch of medical science than Sharpey and Quain's Anatomy.—*The Western Journal of Medicine and Surgery.*

It may now be regarded as the most complete and best posted up work on anatomy in the language. It will be found particularly rich in general anatomy.—*The Charleston Medical Journal.*

We believe we express the opinion of all who have examined these volumes, that there is no work superior to them on the subject which they so ably describe.—*Southern Medical and Surgical Journal.*

It is one of the most comprehensive and best works upon anatomy in the English language. It is equally valuable to the teacher, practitioner, and student in medicine, and to the surgeon in particular.—*The Ohio Medical and Surgical Journal.*

To those who wish an extensive treatise on Anatomy, we recommend these handsome volumes as the best that have ever issued from the English or American Press.—*The N. W. Medical and Surgical Journal.*

We believe that any country might safely be challenged to produce a treatise on anatomy so readable, so clear, and so full upon all important topics.—*British and Foreign Medico-Chirurgical Review.*

It is indeed a work calculated to make an era in anatomical study, by placing before the student every department of his science, with a view to the relative importance of each; and so skillfully have the different parts been interwoven, that no one who makes this work the basis of his studies will hereafter have any excuse for neglecting or undervaluing any important particulars connected with the structure of the human frame; and whether the bias of his mind lead him in a more especial manner to surgery, physis, or physiology, he will find here a work at once so comprehensive and practical as to defend him from exclusiveness on the one hand, and pedantry on the other.—*Monthly Journal and Retrospect of the Medical Sciences.*

We have no hesitation in recommending this treatise on anatomy as the most complete on that subject in the English language; and the only one, perhaps, in any language, which brings the state of knowledge forward to the most recent discoveries.—*The Edinburgh Medical and Surgical Journal.*

Admirably calculated to fulfil the object for which it is intended.—*Provincial Medical Journal.*

The most complete Treatise on Anatomy in the English language.—*Edinburgh Medical Journal.*

There is no work in the English language to be preferred to Dr. Quain's Elements of Anatomy.—*London Journal of Medicine.*

THE STUDENT'S TEXT-BOOK OF ANATOMY.

NEW AND IMPROVED EDITION—JUST ISSUED.

## A SYSTEM OF HUMAN ANATOMY, GENERAL AND SPECIAL.

BY ERASMUS WILSON, M. D.

FOURTH AMERICAN FROM THE LAST ENGLISH EDITION.

EDITED BY PAUL B. GODDARD, A. M., M. D.

WITH TWO HUNDRED AND FIFTY ILLUSTRATIONS.

Beautifully printed, in one large octavo volume of nearly six hundred pages.

In many, if not all the Colleges of the Union, it has become a standard text-book. This, of itself, is sufficiently expressive of its value. A work very desirable to the student; one, the possession of which will greatly facilitate his progress in the study of Practical Anatomy.—*New York Journal of Medicine.*

Its author ranks with the highest on Anatomy.—*Southern Medical and Surgical Journal.*

It offers to the student all the assistance that can be expected from such a work.—*Medical Examiner.*

The most complete and convenient manual for the student we possess.—*American Journal of Med. Science.*

In every respect this work, as an anatomical guide for the student and practitioner, merits our warmest and most decided praise.—*London Medical Gazette.*



**HORNER'S ANATOMY.****MUCH IMPROVED AND ENLARGED EDITION.**—(*Just Issued.*)**SPECIAL ANATOMY AND HISTOLOGY.**

BY WILLIAM E. HORNER, M. D.,

Professor of Anatomy in the University of Pennsylvania, &amp;c.

EIGHTH EDITION.

EXTENSIVELY REVISED AND MODIFIED TO 1851.

In two large octavo volumes, handsomely printed, with several hundred illustrations.

This work has enjoyed a thorough and laborious revision on the part of the author, with the view of bringing it fully up to the existing state of knowledge on the subject of general and special anatomy. To adapt it more perfectly to the wants of the student, he has introduced a large number of additional wood engravings, illustrative of the objects described, while the publishers have endeavored to render the mechanical execution of the work worthy of the extended reputation which it has acquired. The demand which has carried it to an EIGHTH EDITION is a sufficient evidence of the value of the work, and of its adaptation to the wants of the student and professional reader.

NEW AND CHEAPER EDITION OF

**SMITH & HORNER'S ANATOMICAL ATLAS.****AN ANATOMICAL ATLAS,**

ILLUSTRATIVE OF THE STRUCTURE OF THE HUMAN BODY.

BY HENRY H. SMITH, M. D., &amp;c.

UNDER THE SUPERVISION OF

WILLIAM E. HORNER, M. D.,

Professor of Anatomy in the University of Pennsylvania.

In one volume, large imperial octavo, with about six hundred and fifty beautiful figures.

With the view of extending the sale of this beautifully executed and complete "Anatomical Atlas," the publishers have prepared a new edition, printed on both sides of the page, thus materially reducing its cost, and enabling them to present it at a price about forty per cent. lower than former editions, while, at the same time, the execution of each plate is in no respect deteriorated, and not a single figure is omitted.

These figures are well selected, and present a complete and accurate representation of that wonderful fabric, the human body. The plan of this Atlas, which renders it so peculiarly convenient for the student, and its superb artistical execution, have been already pointed out. We most congratulate the student upon the completion of this Atlas, as it is the most convenient work of the kind that has yet appeared; and we must add, the very beautiful manner in which it is "got up" is so creditable to the country as to be flattering to our national pride.—*American Medical Journal.*

**HORNER'S DISSECTOR.****THE UNITED STATES DISSECTOR;**

Being a new edition, with extensive modifications, and almost re-written, of

**"HORNER'S PRACTICAL ANATOMY."**

In one very neat volume, royal 12mo., of 440 pages, with many illustrations on wood.

**WILSON'S DISSECTOR, New Edition.**—(*Just Issued.*)**THE DISSECTOR;  
OR, PRACTICAL AND SURGICAL ANATOMY.**

BY ERASMUS WILSON.

MODIFIED AND RE-ARRANGED BY

PAUL BECK GODDARD, M. D.

A NEW EDITION, WITH REVISIONS AND ADDITIONS.

In one large and handsome volume, royal 12mo., with one hundred and fifteen illustrations.

In passing this work again through the press, the editor has made such additions and improvements as the advance of anatomical knowledge has rendered necessary to maintain the work in the high reputation which it has acquired in the schools of the United States as a complete and faithful guide to the student of practical anatomy. A number of new illustrations have been added, especially in the portion relating to the complicated anatomy of Hernia. In mechanical execution the work will be found superior to former editions.



WORKS BY W. B. CARPENTER, M. D.

COMPARATIVE PHYSIOLOGY—(Now Ready.)

# PRINCIPLES OF PHYSIOLOGY, GENERAL AND COMPARATIVE.

THIRD EDITION, GREATLY ENLARGED.

In one very handsome octavo volume, of over 1100 pages, with 321 beautiful wood-cuts.

This great work will supply a want long felt by the scientific public of this country, who have had no accessible treatise to refer to, presenting, in an intelligible form, a complete and thorough outline of this important subject. The high reputation of the author, on both sides of the Atlantic, is a sufficient guarantee for the completeness and accuracy of any work to which his name is prefixed; but this volume comes with the additional recommendation that it is the one on which the author has bestowed the greatest care, and on which he is desirous to rest his reputation. Two years have been devoted to the preparation of this edition, which has been thoroughly remoulded and rewritten, so as, in fact, to constitute a new work. The amount of alterations and additions may be understood from the fact that of the ten hundred and eighty pages of the text, but one hundred and fifty belong to the previous edition. Containing, as it does, the results of years devoted to study and observation, it may be regarded as a complete exposition of the most advanced state of knowledge in this rapidly progressive branch of science, and as a storehouse of facts and principles in all departments of Physiology, such as perhaps no man but its author could have accumulated and classified.

In every point of mechanical execution, and profuseness and beauty of illustration, the publishers risk nothing in saying that it will be found all that the most fastidious taste could desire.

A truly magnificent work. In itself a perfect physiological study.—*Ranking's Abstract*, July 21, 1851.

This work stands without its fellow. It is one few men in Europe could have undertaken; it is one no man, we believe, could have brought to so successful an issue as Dr. Carpenter. It required for its production a physiologist at once deeply read in the labours of others, capable of taking a general, critical, and unprejudiced view of those labours, and of combining the varied, heterogeneous materials at his disposal, so as to form an harmonious whole. We feel that this abstract can give the reader a very imperfect idea of the fulness of this work, and no idea of its unity, of the admirable manner in which material has been brought, from the most various sources, to conduce to its completeness, of the lucidity of the reasoning it contains, or of the clearness of language in which the whole is clothed. Not the profession only, but the scientific world at large, must feel deeply indebted to Dr. Carpenter for this great work. It must, indeed, add largely even to his high reputation.—*Medical Times*.

CARPENTER'S ELEMENTS OF PHYSIOLOGY, New and Improved Edition—(Now Ready.)

## ELEMENTS OF PHYSIOLOGY; INCLUDING PHYSIOLOGICAL ANATOMY.

WITH NEARLY TWO HUNDRED ILLUSTRATIONS.

*Second American, from the Second London Edition.*

In one handsome octavo volume, of about six hundred pages.

This work has been subjected to a thorough revision by the author, who, without materially increasing its size, has brought it up completely with the most recent researches and investigations. It may, therefore, be regarded as the latest exponent of the views of the author, with respect to all the modern doctrines of physiology. Numerous wood-cuts have been added, the work has been carefully read during its passage through the press, and in every point of mechanical execution it will be found equal, if not superior, to the former edition.

To say that it is the best manual of Physiology now before the public, would not do sufficient justice to the author.—*Buffalo Med. Journal*.

In his former works it would seem that he had exhausted the subject of Physiology. In the present, he gives the essence, as it were, of the whole.—*N. Y. Journal of Medicine*.

The best and most complete exposé of modern physiology, in one volume, extant in the English language.—*St. Louis Med. Journal*.

Those who have occasion for an elementary treatise on physiology, cannot do better than to possess themselves of the manual of Dr. Carpenter.—*Medical Examiner*.

CARPENTER'S HUMAN PHYSIOLOGY—(New Edition, Preparing.)

## PRINCIPLES OF HUMAN PHYSIOLOGY,

WITH THEIR CHIEF APPLICATIONS TO

PATHOLOGY, HYGIENE, AND FORENSIC MEDICINE.

A NEW EDITION, WITH EXTENSIVE ADDITIONS AND IMPROVEMENTS BY THE AUTHOR.

With Two Lithographic Plates, and over 300 Wood-cuts.

In one large and handsomely printed octavo volume of over seven hundred and fifty pages.

In preparing a new edition of this very popular text-book, it has been thoroughly revised by the author, who, without materially increasing its bulk, has embodied in it all the recent investigations and discoveries in physiological science, and has rendered it in every respect on a level with the improvements of the day. Although the number of the wood-engravings has been but little increased, a considerable change will be found. In passing the volume through the press in this country, the services of a professional gentleman have been secured, in order to insure the accuracy so necessary to a scientific work.

In recommending this work to their classes, Professors of Physiology can rely on their being always able to procure editions brought thoroughly up with the advance of science.



**DUNGLISON'S PHYSIOLOGY.**  
**New and much Improved Edition.—(Just Issued.)**

# HUMAN PHYSIOLOGY.

BY ROBLEY DUNGLISON, M. D.,  
 Professor of the Institutes of Medicine in the Jefferson Medical College, Philadelphia, etc. etc.  
 SEVENTH EDITION.

Thoroughly revised and extensively modified and enlarged,  
 With nearly Five Hundred Illustrations.

In two large and handsomely printed octavo volumes, containing nearly 1450 pages.

On no previous revision of this work has the author bestowed more care than on the present, it having been subjected to an entire scrutiny, not only as regards the important matters of which it treats, but also the language in which they are conveyed; and on no former occasion has he felt as satisfied with his endeavors to have the work on a level with the existing state of the science. Perhaps at no time in the history of physiology have observers been more numerous, energetic, and discriminating than within the last few years. Many modifications of fact and inference have consequently taken place, which it has been necessary for the author to record, and to express his views in relation thereto. On the whole subject of physiology proper, as it applies to the functions executed by the different organs, the present edition, the author flatters himself, will therefore be found to contain the views of the most distinguished physiologists of all periods.

The amount of additional matter contained in this edition may be estimated from the fact that the mere list of authors referred to in its preparation alone extends over nine large and closely printed pages. The number of illustrations has been largely increased, the present edition containing four hundred and seventy-four, while the last had but three hundred and sixty-eight; while, in addition to this, many new and superior wood-cuts have been substituted for those which were not deemed sufficiently accurate or satisfactory. The mechanical execution of the work has also been improved in every respect, and the whole is confidently presented as worthy the great and continued favor which it has so long received from the profession.

It has long since taken rank as one of the medical classics of our language. To say that it is by far the best text-book of physiology ever published in this country, is but echoing the general testimony of the profession.  
 —*N. Y. Journal of Medicine.*

The most full and complete system of Physiology in our language.—*Western Lancet.*

The most complete and satisfactory system of Physiology in the English language.—*Amer. Med. Journal.*

The best work of the kind in the English language.—*Silliman's Journal.*

We have, on two former occasions, brought this excellent work under the notice of our readers, and we have now only to say that, instead of falling behind in the rapid march of physiological science, each edition brings it nearer to the van.—*British and Foreign Medical Review.*

A review of such a well-known work would be out of place at the present time. We have looked over it, and find, what we knew would be the case, that Dr. Dunglison has kept pace with the science to which he has devoted so much study, and of which he is one of the living ornaments. We recommend the work to the medical student as a valuable text-book, and to all inquirers into Natural Science, as one which will well and delightfully repay perusal.—*The New Orleans Medical and Surgical Journal.*

**KIRKES AND PAGET'S PHYSIOLOGY.—(Lately Issued.)**

## A MANUAL OF PHYSIOLOGY, FOR THE USE OF STUDENTS.

BY WILLIAM SENHOUSE KIRKES, M. D.,

ASSISTED BY JAMES PAGET,

Lecturer on General Anatomy and Physiology in St. Bartholomew's Hospital.

In one handsome volume, royal 12mo., of 550 pages, with 118 wood-cuts.

An excellent work, and for students one of the best within reach.—*Boston Medical and Surgical Journal.*  
 One of the best little books on Physiology which we possess.—*Braithwaite's Retrospect.*

Particularly adapted to those who desire to possess a concise digest of the facts of Human Physiology.—*British and Foreign Med.-Chirurg. Review.*

One of the best treatises which can be put into the hands of the student.—*London Medical Gazette.*

We conscientiously recommend it as an admirable "Handbook of Physiology."—*London Jour. of Medicine.*

### SOLLY ON THE BRAIN.

## THE HUMAN BRAIN; ITS STRUCTURE, PHYSIOLOGY, AND DISEASES.

WITH A DESCRIPTION OF THE TYPICAL FORM OF THE BRAIN IN THE ANIMAL KINGDOM.

BY SAMUEL SOLLY, F. R. S., &c.,

Senior Assistant Surgeon to the St. Thomas' Hospital, &c.

From the Second and much Enlarged London Edition. In one octavo volume, with 120 Wood-cuts.

**HARRISON ON THE NERVES.**—An Essay towards a correct theory of the Nervous System. In one octavo volume, 292 pages.

**MATTEUCCI ON LIVING BEINGS.**—Lectures on the Physical Phenomena of Living Beings. Edited by Pereira. In one neat royal 12mo. volume, extra cloth, with cuts—388 pages.

**ROGET'S PHYSIOLOGY.**—A Treatise on Animal and Vegetable Physiology, with over 400 illustrations on wood. In two octavo volumes, cloth.

**ROGET'S OUTLINES.**—Outlines of Physiology and Phrenology. In one octavo volume, cloth—516 pages.  
**ON THE CONNECTION BETWEEN PHYSIOLOGY AND INTELLECTUAL SCIENCE.** In one 12mo. volume, paper, price 25 cents.

**TODD & BOWMAN'S PHYSIOLOGY.**—Physiological Anatomy and Physiology of Man. With numerous handsome wood-cuts. Parts I, II, and III, in one 8vo. volume, 552 pp. Part IV will complete the work.



*WILLIAMS' PRINCIPLES—New and Enlarged Edition.***PRINCIPLES OF MEDICINE;**

Comprising General Pathology and Therapeutics,

AND A BRIEF GENERAL VIEW OF

ETIOLOGY, NOSOLOGY, SEMEIOLOGY, DIAGNOSIS, PROGNOSIS, AND HYGIENICS.

BY CHARLES J. B. WILLIAMS, M. D., F. R. S.,

Fellow of the Royal College of Physicians, &amp;c.

EDITED, WITH ADDITIONS, BY MEREDITH CLYMER, M. D.,

Consulting Physician to the Philadelphia Hospital, &amp;c. &amp;c.

THIRD AMERICAN, FROM THE SECOND AND ENLARGED LONDON EDITION.

In one octavo volume, of 440 pages.

**BILLING'S PRINCIPLES, NEW EDITION—(Just Issued.)****THE PRINCIPLES OF MEDICINE.**

BY ARCHIBALD BILLING, M. D., &amp;c.

Second American from the Fifth and Improved London Edition.

In one handsome octavo volume, extra cloth, 250 pages.

We can strongly recommend Dr. Billing's "Principles" as a code of instruction which should be constantly present to the mind of every well-informed and philosophical practitioner of medicine.—*Lancet*.**MANUALS ON THE BLOOD AND URINE.**

In one handsome volume royal 12mo., extra cloth, of 460 large pages, with numerous illustrations.

CONTAINING

I. A Practical Manual on the Blood and Secretions of the Human Body. BY JOHN WILLIAM GRIFFITH, M. D., &amp;c.

II. On the Analysis of the Blood and Urine in health and disease, and on the treatment of Urinary diseases. BY G. OWEN REESE, M. D., F. R. S., &amp;c. &amp;c.

III. A Guide to the Examination of the Urine in health and disease. BY ALFRED MARKWICK.

The importance now attached to the Diagnosis of the Blood and Urine, and the rapid increase of our knowledge respecting the pathological conditions of the fluids of the human body, have induced the publishers to present these manuals in a cheap and convenient form, embracing the results of the most recent observers in a practical point of view. On the subject of the chemical and microscopical examinations of these fluids, they would also call the attention of the student to BOWMAN'S MEDICAL CHEMISTRY, and SIMON'S ANIMAL CHEMISTRY. See p. 30.

NEW EDITION—(JUST ISSUED.)

**URINARY DEPOSITS;  
THEIR DIAGNOSIS, PATHOLOGY, AND THERAPEUTICAL INDICATIONS.**

BY GOLDING BIRD, A. M., M. D., &amp;c.

A NEW AMERICAN, FROM THE THIRD AND IMPROVED LONDON EDITION.

In one very neat volume, royal 12mo., with over sixty illustrations.

Though the present edition of this well-known work is but little increased in size, it will be found essentially modified throughout, and fully up to the present state of knowledge on its subject. The unanimous testimony of the medical press warrants the publishers in presenting it as a complete and reliable manual for the student of this interesting and important branch of medical science.

**THE PATHOLOGICAL ANATOMY OF THE HUMAN BODY.**

BY JULIUS VOGEL, M. D., &amp;c.

Translated from the German, with Additions,

BY GEORGE E. DAY, M. D., &amp;c.

ILLUSTRATED BY UPWARDS OF ONE HUNDRED FIGURES, PLAIN AND COLORED.

In one neat octavo volume.

ABERCROMBIE ON THE BRAIN.—Pathological and Practical Researches on Diseases of the Brain and Spinal Cord. A new edition, in one small 8vo. volume, pp. 324.

BURROWS ON CEREBRAL CIRCULATION.—On Disorders of the Cerebral Circulation, and on the Connection between Affections of the Brain and Diseases of the Heart. In one 8vo. vol., with colored plates, pp. 216.

BLAKISTON ON THE CHEST.—Practical Observations on certain Diseases of the Chest, and on the Principles of Auscultation. In one volume, 8vo., pp. 384.

HASSE'S PATHOLOGICAL ANATOMY.—An Anatomical Description of the Diseases of Respiration and Circulation. Translated and Edited by Swaine. In one volume, 8vo., pp. 379.

HUGHES ON THE LUNGS AND HEART.—Clinical Introduction to the Practice of Auscultation, and other modes of Physical Diagnosis. In one 12mo. volume, with a plate, pp. 270.

FRICK ON THE URINE.—Renal Affections, their Diagnosis and Pathology. In one handsome volume, royal 12mo., with illustrations.

COPLAND ON PALSY.—Of the Causes, Nature, and Treatment of Palsy and Apoplexy. In one volume, royal 12mo. (Just Issued.)



**DUNGLISON'S PRACTICE OF MEDICINE.**  
 ENLARGED AND IMPROVED EDITION.  
**THE PRACTICE OF MEDICINE.**  
 A TREATISE ON  
**SPECIAL PATHOLOGY AND THERAPEUTICS.**  
 THIRD EDITION.

BY ROBLEY DUNGLISON, M. D.,

Professor of the Institutes of Medicine in the Jefferson Medical College; Lecturer on Clinical Medicine, &c.

In two large octavo volumes, of fifteen hundred pages.

The student of medicine will find, in these two elegant volumes, a mine of facts, a gathering of precepts and advice from the world of experience, that will nerve him with courage, and faithfully direct him in his efforts to relieve the physical sufferings of the race.—*Boston Medical and Surgical Journal*.

Upon every topic embraced in the work the latest information will be found carefully posted up. *Medical Examiner*.

It is certainly the most complete treatise of which we have any knowledge. There is scarcely a disease which the student will not find noticed.—*Western Journal of Medicine and Surgery*.

One of the most elaborate treatises of the kind we have.—*Southern Medical and Surg. Journal*.

A New Work. Now Ready.

**DISEASES OF THE HEART, LUNGS, AND APPENDAGES;  
 THEIR SYMPTOMS AND TREATMENT.**

BY W. H. WALSHE, M. D.,

Professor of the Principles and Practice of Medicine in University College, London, &c.

In one handsome volume, large royal 12mo.

The author's design in this work has been to include within the compass of a moderate volume, all really essential facts bearing upon the symptoms, physical signs, and treatment of pulmonary and cardiac diseases. To accomplish this, the first part of the work is devoted to the description of the various modes of physical diagnosis, auscultation, percussion, mensuration, &c., which are fully and clearly, but succinctly entered into, both as respects their theory and clinical phenomena. In the second part, the various diseases of the heart, lungs, and great vessels are considered in regard to symptoms, physical signs and treatment, with numerous references to cases. The eminence of the author is a guarantee to the practitioner and student that the work is one of practical utility in facilitating the diagnosis and treatment of a large, obscure and important class of diseases.

THE GREAT MEDICAL LIBRARY.

**THE CYCLOPEDIA OF PRACTICAL MEDICINE;**

COMPRISING

Treatises on the Nature and Treatment of Diseases, Materia Medica, and Therapeutics, Diseases of Women and Children, Medical Jurisprudence, &c. &c.

EDITED BY

JOHN FORBES, M. D., F. R. S., ALEXANDER TWEEDIE, M. D., F. R. S.

AND JOHN CONNOLLY, M. D.

Revised, with Additions,

BY ROBLEY DUNGLISON, M. D.

THIS WORK IS NOW COMPLETE, AND FORMS FOUR LARGE SUPER-ROYAL OCTAVO VOLUMES, Containing Thirty-two Hundred and Fifty-four unusually large Pages in Double Columns, Printed on Good Paper, with a new and clear type.

THE WHOLE WELL AND STRONGLY BOUND WITH RAISED BANDS AND DOUBLE TITLES.

This work contains no less than FOUR HUNDRED AND EIGHTEEN DISTINCT TREATISES,

By Sixty-eight distinguished Physicians.

The most complete work on Practical Medicine extant; or, at least, in our language.—*Buffalo Medical and Surgical Journal*.

For reference, it is above all price to every practitioner.—*Western Lancet*.

One of the most valuable medical publications of the day—as a work of reference it is invaluable.—*Western Journal of Medicine and Surgery*.

It has been to us, both as learner and teacher, a work for ready and frequent reference, one in which modern English medicine is exhibited in the most advantageous light.—*Medical Examiner*.

We rejoice that this work is to be placed within the reach of the profession in this country, it being unquestionably one of very great value to the practitioner. This estimate of it has not been formed from a hasty examination, but after an intimate acquaintance derived from frequent consultation of it during the past nine or ten years. The editors are practitioners of established reputation, and the list of contributors embraces many of the most eminent professors and teachers of London, Edinburgh, Dublin, and Glasgow. It is, indeed, the great merit of this work that the principal articles have been furnished by practitioners who have not only devoted especial attention to the diseases about which they have written, but have also enjoyed opportunities for an extensive practical acquaintance with them.—and whose reputation carries the assurance of their competency justly to appreciate the opinions of others, while it stamps their own doctrines with high and just authority.—*American Medical Journal*.



WATSON'S PRACTICE OF MEDICINE—New Edition.

LECTURES ON THE  
**PRINCIPLES AND PRACTICE OF PHYSIC.**

BY THOMAS WATSON, M. D., &amp;c. &amp;c.

Third American, from the last London Edition.

REVISED, WITH ADDITIONS, BY D. FRANCIS CONDIE, M. D.,  
Author of "A Treatise on the Diseases of Children," &c.

IN ONE OCTAVO VOLUME,

Of nearly ELEVEN HUNDRED LARGE PAGES, strongly bound with raised bands.

To say that it is the very best work on the subject now extant, is but to echo the sentiment of the medical press throughout the country.—*N. O. Medical Journal.*Of the text-books recently republished Watson is very justly the principal favorite.—*Holmes' Report to Nat. Med. Assoc.*By universal consent the work ranks among the very best text-books in our language.—*Ill. and Ind. Med. Journal.*Regarded on all hands as one of the very best, if not the very best, systematic treatise on practical medicine extant.—*St. Louis Med. Journal.*Confessedly one of the very best works on the principles and practice of physic in the English or any other language.—*Med. Examiner.*As a text-book it has no equal; as a compendium of pathology and practice no superior.—*N. Y. Annalist.*We know of no work better calculated for being placed in the hands of the student, and for a text-book, on every important point the author seems to have posted up his knowledge to the day.—*Amer. Med. Journal.*One of the most practically useful books that ever was presented to the student.—*N. Y. Med. Journal.*

WILSON ON THE SKIN.

**ON DISEASES OF THE SKIN.**

BY ERASMUS WILSON, F. R. S.,

Author of "Human Anatomy," &amp;c.

SECOND AMERICAN FROM THE SECOND LONDON EDITION.

In one neat octavo volume, extra cloth, 440 pages.

Also, to be had with eight beautifully colored steel plates.

Also, the plates sold separate, in boards.

Much Enlarged Edition of BARTLETT ON FEVERS.

THE HISTORY, DIAGNOSIS, AND TREATMENT OF THE  
**FEVERS OF THE UNITED STATES.**

BY ELISHA BARTLETT, M. D.,

In one octavo volume of 550 pages, beautifully printed and strongly bound.

CLYMER AND OTHERS ON FEVERS.

**FEVERS; THEIR DIAGNOSIS, PATHOLOGY, AND TREATMENT.**

PREPARED AND EDITED, WITH LARGE ADDITIONS,

FROM THE ESSAYS ON FEVER IN TWEEDIE'S LIBRARY OF PRACTICAL MEDICINE,

BY MEREDITH CLYMER, M. D.

In one octavo volume of six hundred pages.

BENEDICT'S CHAPMAN.—Compendium of Chapman's Lectures on the Practice of Medicine. One neat volume, 8vo., pp. 258.

BUDD ON THE LIVER.—On Diseases of the Liver. In one very neat 8vo. vol., with colored plates and wood-cuts, pp. 392.

CHAPMAN'S LECTURES.—Lectures on Fevers, Dropsy, Gout, Rheumatism, &amp;c. &amp;c. In one neat 8vo. volume, pp. 450.

ESQUIROL ON INSANITY.—Mental Maladies, considered in relation to Medicine, Hygiene, and Medical Jurisprudence. Translated by R. K. Hunt, M. D., &amp;c. In one 8vo. volume, pp. 496.

THOMSON ON THE SICK ROOM.—Domestic management of the sick Room, necessary in aid of Medical Treatment for the cure of Diseases. Edited by R. E. Griffith, M. D. In one large royal 12mo. volume, with wood-cuts, pp. 360.

HOPE ON THE HEART.—A Treatise on the Diseases of the Heart and Great Vessels. Edited by Pen-nock. In one volume, 8vo., with plates, pp. 572.

LALLEMAND ON SPERMATORRHOEA.—The Causes, Symptoms, and Treatment of Spermatorrhœa. Translated and Edited by Henry J. McDougal. In one volume, 8vo., pp. 320.

PHILIPS ON SCROFULA.—Scrofula: its Nature, its Prevalence, its Causes, and the Principles of its Treatment. In one volume, 8vo., with a plate, pp. 350.

WHITEHEAD ON ABORTION, &amp;c.—The Causes and Treatment of Abortion and Sterility; being the Result of an Extended Practical Inquiry into the Physiological and Morbid Conditions of the Uterus. In one volume, 8vo., pp. 368.

WILLIAMS ON RESPIRATORY ORGANS.—A Practical Treatise on Diseases of the Respiratory Organs; including Diseases of the Larynx, Trachea, Lungs, and Pleuræ. With numerous Additions and Notes by M. Clymer, M. D. With wood-cuts. In one octavo volume, pp. 508.

DAY ON OLD AGE.—A Practical Treatise on the Domestic Management and more important Diseases of Advanced Life. With an Appendix on a new and successful mode of treating Lumbago and other forms of Chronic Rheumatism. 1 vol. 8vo., pp. 226.



**MEIGS ON FEMALES, New and Improved Edition—(Lately Issued.)**  
**WOMAN; HER DISEASES AND THEIR REMEDIES;**  
**A SERIES OF LETTERS TO HIS CLASS.**

BY C. D. MEIGS, M. D.,

Professor of Midwifery and Diseases of Women and Children in the Jefferson Medical College of Philadelphia, &c. &c.

In one large and beautifully printed octavo volume, of nearly seven hundred large pages.

"I am happy to offer to my Class an enlarged and amended edition of my Letters on the Diseases of Women; and I avail myself of this occasion to return my heartfelt thanks to them, and to our brethren generally, for the flattering manner in which they have accepted this fruit of my labor."—PREFACE.

The value attached to this work by the profession is sufficiently proved by the rapid exhaustion of the first edition, and consequent demand for a second. In preparing this the author has availed himself of the opportunity thoroughly to revise and greatly to improve it. The work will therefore be found completely brought up to the day, and in every way worthy of the reputation which it has so immediately obtained.

Professor Meigs has enlarged and amended this great work, for such it unquestionably is, having passed the ordeal of criticism at home and abroad, but been improved thereby; for in this new edition the author has introduced real improvements, and increased the value and utility of the book immeasurably. It presents so many novel, bright and sparkling thoughts; such an exuberance of new ideas on almost every page, that we confess ourselves to have become enamored with the book and its author; and cannot withhold our congratulations from our Philadelphia conferees, that such a teacher is in their service. We regret that our limits will not allow of a more extended notice of this work, but must content ourselves with thus commending it as worthy of diligent perusal by physicians as well as students, who are seeking to be thoroughly instructed in the important practical subjects of which it treats.—*N. Y. Med. Gazette.*

It contains a vast amount of practical knowledge, by one who has accurately observed and retained the experience of many years, and who tells the result in a free, familiar, and pleasant manner.—*Dublin Quarterly Journal.*

There is an off-hand fervor, a glow and a warm-heartedness infecting the effort of Dr. Meigs, which is entirely captivating, and which absolutely hurries the reader through from beginning to end. Besides, the book teems with solid instruction, and it shows the very highest evidence of ability, viz., the clearness with which the information is presented. We know of no better test of one's understanding a subject than the evidence of the power of lucidly explaining it. The most elementary, as well as the obscurest subjects, under the pencil of Prof. Meigs, are isolated and made to stand out in such bold relief, as to produce distinct impressions upon the mind and memory of the reader.—*The Charleston Medical Journal.*

The merits of the first edition of this work were so generally appreciated, and with such a high degree of favor by the medical profession throughout the Union, that we are not surprised in seeing a second edition of it. It is a standard work on the diseases of females, and in many respects is one of the very best of its kind in the English language. Upon the appearance of the first edition, we gave the work a cordial reception, and spoke of it in the warmest terms of commendation. Time has not changed the favorable estimate we placed upon it, but has rather increased our convictions of its superlative merits. But we do not now deem it necessary to say more than to commend this work, on the diseases of women, and the remedies for them, to the attention of those practitioners who have not supplied themselves with it. The most select library would be imperfect without it.—*The Western Journal of Medicine and Surgery.*

He is a bold thinker, and possesses more originality of thought and style than almost any American writer on medical subjects. If he is not an elegant writer, there is at least a freshness—a raciness in his mode of expressing himself—that cannot fail to draw the reader after him, even to the close of his work: you cannot nod over his pages; he stimulates rather than narcotises your senses, and the reader cannot lay aside these letters when once he enters into their merits. This, the second edition, is much amended and enlarged, and affords abundant evidence of the author's talents and industry.—*N. O. Medical and Surgical Journal.*

The practical writings of Dr. Meigs are second to none.—*The N. Y. Journal of Medicine.*

The excellent practical directions contained in this volume give it great utility, which we trust will not be lost upon our older colleagues; with some condensation, indeed, we should think it well adapted for translation into German.—*Zeitschrift für die Gesamte Medicin.*

NEW AND IMPROVED EDITION—(Lately Issued.)

**A TREATISE ON THE DISEASES OF FEMALES,**  
**AND ON THE SPECIAL HYGIENE OF THEIR SEX.**  
**BY COLOMBAT DE L'ISERE, M. D.**

TRANSLATED, WITH MANY NOTES AND ADDITIONS, BY C. D. MEIGS, M. D.

SECOND EDITION, REVISED AND IMPROVED.

In one large volume, octavo, of seven hundred and twenty pages, with numerous wood-cuts.

We are satisfied it is destined to take the front rank in this department of medical science. It is in fact a complete exposition of the opinions and practical methods of all the celebrated practitioners of ancient and modern times.—*New York Journ. of Medicine.*

**ASHWELL ON THE DISEASES OF FEMALES.**  
**A PRACTICAL TREATISE ON THE DISEASES PECULIAR TO WOMEN.**

ILLUSTRATED BY CASES DERIVED FROM HOSPITAL AND PRIVATE PRACTICE.

BY SAMUEL ASHWELL, M. D. WITH ADDITIONS BY PAUL BECK GODDARD, M. D.

Second American edition. In one octavo volume, of 520 pages.

One of the very best works ever issued from the press on the Diseases of Females.—*Western Lancet.*

**ON THE CAUSES AND TREATMENT OF ABORTION AND STERILITY.** By James Whitehead, M. D., &c. In one volume octavo, of about three hundred and seventy-five pages.



NEW AND IMPROVED EDITION—(Lately Issued.)

## THE DISEASES OF FEMALES. INCLUDING THOSE OF PREGNANCY AND CHILD-BEED.

BY FLEETWOOD CHURCHILL, M. D., M. R. I. A.,

Author of "Theory and Practice of Midwifery," "Diseases of Females," &amp;c.

A New American Edition (The Fifth), Revised by the Author.

WITH THE NOTES OF ROBERT M. HUSTON, M. D.

In one large and handsome octavo volume of 632 pages, with wood-cuts.

To indulge in panegyric, when announcing the fifth edition of any acknowledged medical authority, were to attempt to "gild refined gold." The work announced above, has too long been honored with the term "classical" to leave any doubt as to its true worth, and we content ourselves with remarking, that the author has carefully retained the notes of Dr. Huston, who edited the former American edition, thus really enhancing the value of the work, and paying a well merited compliment. All who wish to be "posted up" on all that relates to the diseases peculiar to the wife, the mother, or the maid, will hasten to secure a copy of this most admirable treatise.—*The Ohio Medical and Surgical Journal.*

We know of no author who deserves that approbation, on "the diseases of females," to the same extent that Dr. Churchill does. His, indeed, is the only thorough treatise we know of on the subject, and it may be commended to practitioners and students as a masterpiece in its particular department. The former editions of this work have been commended strongly in this journal, and they have won their way to an extended, and a well deserved popularity. This fifth edition, before us, is well calculated to maintain Dr. Churchill's high reputation. It was revised and enlarged by the author, for his American publishers, and it seems to us, that there is scarcely any species of desirable information on its subjects, that may not be found in this work.—*The Western Journal of Medicine and Surgery.*

We are gratified to announce a new and revised edition of Dr. Churchill's valuable work on the diseases of females. We have ever regarded it as one of the very best works on the subjects embraced within its scope, in the English language; and the present edition, enlarged and revised by the author, renders it still more entitled to the confidence of the profession. The valuable notes of Prof. Huston have been retained, and contribute, in no small degree, to enhance the value of the work. It is a source of congratulation that the publishers have permitted the author to be, in this instance, his own editor, thus securing all the revision which an author alone is capable of making.—*The Western Lancet.*

As a comprehensive manual for students, or a work of reference for practitioners, we only speak with common justice when we say that it surpasses any other that has ever issued on the same subject from the British press.—*The Dublin Quarterly Journal.*

Churchill's Monographs on Females—(Lately Issued.)

## ESSAYS ON THE PUERPERAL FEVER, AND OTHER DISEASES PECULIAR TO WOMEN.

SELECTED FROM THE WRITINGS OF BRITISH AUTHORS PREVIOUS TO THE CLOSE OF  
THE EIGHTEENTH CENTURY.

Edited by FLEETWOOD CHURCHILL, M. D., M. R. I. A.,

Author of "Treatise on the Diseases of Females," &amp;c.

In one neat octavo volume, of about four hundred and fifty pages.

To these papers Dr. Churchill has appended notes, embodying whatever information has been laid before the profession since their authors' time. He has also prefixed to the essays on puerperal fever, which occupy the larger portion of the volume, an interesting historical sketch of the principal epidemics of that disease. The whole forms a very valuable collection of papers by professional writers of eminence, on some of the most important accidents to which the puerperal female is liable.—*American Journal of Medical Sciences.*

MUCH ENLARGED AND IMPROVED EDITION—(Lately Issued.)

## A PRACTICAL TREATISE ON INFLAMMATION OF THE UTERUS AND ITS APPENDAGES, AND ON ULCERATION AND INDURATION OF THE NECK OF THE UTERUS.

BY HENRY BENNETT, M. D.,

Obstetric Physician to the Western Dispensary.

Second Edition, much enlarged.

In one neat octavo volume of 350 pages, with wood-cuts.

This edition is so enlarged as to constitute a new work. It embraces the study of inflammation in all the uterine organs, and its influence in the production of displacements and of the reputed functional diseases of the uterus.

Few works issue from the medical press which are at once original and sound in doctrine; but such, we feel assured, is the admirable treatise now before us. The important practical precepts which the author inculcates are all rigidly deduced from facts. . . . Every page of the book is good, and eminently practical. So far as we know and believe, it is the best work on the subject on which it treats.—*Monthly Journal of Medical Science.*

## A TREATISE ON THE DISEASES OF FEMALES.

BY W. P. DEWEES, M. D.

NINTH EDITION.

In one volume, octavo. 532 pages, with plates.



## MEIGS ON CHILDREN—Just Issued.

OBSERVATIONS ON  
CERTAIN OF THE DISEASES OF YOUNG CHILDREN.

BY CHARLES D. MEIGS, M. D.,

Professor of Midwifery and of the Diseases of Women and Children in the Jefferson  
Medical College of Philadelphia, &c. &c.

In one handsome octavo volume of 214 pages.

While this work is not presented to the profession as a systematic and complete treatise on Infantile disorders, the importance of the subjects treated of, and the interest attaching to the views and opinions of the distinguished author must command for it the attention of all who are called upon to treat this interesting class of diseases.

It puts forth no claims as a systematic work, but contains an amount of valuable and useful matter, scarcely to be found in the same space in our home literature. It can not but prove an acceptable offering to the profession at large.—*N. Y. Journal of Medicine.*

The work before us is undoubtedly a valuable addition to the fund of information which has already been treasured up on the subjects in question. It is practical, and therefore eminently adapted to the general practitioner. Dr. Meigs' works have the same fascination which belongs to himself.—*Medical Examiner.*

This is a most excellent work on the obscure diseases of childhood, and will afford the practitioner and student of medicine much aid in their diagnosis and treatment.—*The Boston Medical and Surgical Journal.*

We take much pleasure in recommending this excellent little work to the attention of medical practitioners. It deserves their attention, and after they commence its perusal, they will not willingly abandon it, until they have mastered its contents. We read the work while suffering from a carbuncle, and its fascinating pages often beguiled us into forgetfulness of agonizing pain. May it teach others to relieve the afflictions of the young.—*The Western Journal of Medicine and Surgery.*

All of which topics are treated with Dr. Meigs' acknowledged ability and original diction. The work is neither a systematic nor a complete treatise upon the diseases of children, but a fragment which may be consulted with much advantage.—*Southern Medical and Surgical Journal.*

## NEW WORK BY DR. CHURCHILL.

ON THE  
DISEASES OF INFANTS AND CHILDREN.

BY FLEETWOOD CHURCHILL, M. D., M. R. I. A.,

Author of "Theory and Practice of Midwifery," "Diseases of Females," &amp;c.

In one large and handsome octavo volume of over 600 pages.

From Dr. Churchill's known ability and industry, we were led to form high expectations of this work; nor were we deceived. Its learned author seems to have set no bounds to his researches in collecting information which, with his usual systematic address, he has disposed of in the most clear and concise manner, so as to lay before the reader every opinion of importance bearing upon the subject under consideration.

We regard this volume as possessing more claims to completeness than any other of the kind with which we are acquainted. Most cordially and earnestly, therefore, do we commend it to our professional brethren, and we feel assured that the stamp of their approbation will in due time be impressed upon it.

After an attentive perusal of its contents, we hesitate not to say, that it is one of the most comprehensive ever written upon the diseases of children, and that, for copiousness of reference, extent of research, and perspicuity of detail, it is scarcely to be equalled, and not to be excelled in any language.—*Dublin Quarterly Journal.*

The present volume will sustain the reputation acquired by the author from his previous works. The reader will find in it full and judicious directions for the management of infants at birth, and a compendious, but clear, account of the diseases to which children are liable, and the most successful mode of treating them. We must not close this notice without calling attention to the author's style, which is perspicuous and polished to a degree, we regret to say, not generally characteristic of medical works. We recommend the work of Dr. Churchill most cordially, both to students and practitioners, as a valuable and reliable guide in the treatment of the diseases of children.—*Am. Journ. of the Med. Sciences.*

After this meagre, and we know, very imperfect notice, of Dr. Churchill's work, we shall conclude by saying, that it is one that cannot fail from its copiousness, extensive research, and general accuracy, to exalt still higher the reputation of the author in this country. The American reader will be particularly pleased to find that Dr. Churchill has done full justice throughout his work, to the various American authors on this subject. The names of Dewees, Eberle, Condie, and Stewart, occur on nearly every page, and these authors are constantly referred to by the author in terms of the highest praise, and with the most liberal courtesy.—*The Medical Examiner.*

We know of no work on this department of Practical Medicine which presents so candid and unprejudiced a statement or posting up of our actual knowledge as this.—*N. Y. Journal of Medicine.*

Its claims to merit, both as a scientific and practical work, are of the highest order. Whilst we would not elevate it above every other treatise on the same subject, we certainly believe that very few are equal to it, and none superior.—*Southern Med. and Surg. Journal.*



New and Improved Edition.

# A PRACTICAL TREATISE ON THE DISEASES OF CHILDREN.

BY D. FRANCIS CONDIE, M. D.,

Fellow of the College of Physicians, &amp;c. &amp;c.

Third edition, revised and augmented. In one large volume, 8vo., of over 700 pages.

In the preparation of a third edition of the present treatise, every portion of it has been subjected to a careful revision. A new chapter has been added on Epidemic Meningitis, a disease which, although not confined to children, occurs far more frequently in them, than in adults. In the other chapters of the work, all the more important facts that have been developed since the appearance of the last edition, in reference to the nature, diagnosis, and treatment of the several diseases of which they treat, have been incorporated. The great object of the author has been to present, in each succeeding edition, as full and connected a view as possible of the actual state of the pathology and therapeutics of those affections which most usually occur between birth and puberty.

To the present edition there is appended a list of the several works and essays quoted or referred to in the body of the work, or which have been consulted in its preparation or revision.

Every important fact that has been verified or developed since the publication of the previous edition, either in relation to the nature, diagnosis, or treatment of the diseases of children, have been arranged and incorporated into the body of the work: thus putting up to date, to use a counting-house phrase, all the valuable facts and useful information on the subject. To the American practitioner, Dr. Condie's remarks on the diseases of children will be invaluable, and we accordingly advise those who have failed to read this work to procure a copy, and make themselves familiar with its sound principles.—*The New Orleans Medical and Surgical Journal*.

We feel persuaded that the American Medical profession will soon regard it, not only as a very good, but as the very best "Practical Treatise on the Diseases of Children."—*American Medical Journal*.

We pronounced the first edition to be the best work on the Diseases of Children in the English language, and, notwithstanding all that has been published, we still regard it in that light.—*Medical Examiner*.

*From Professor Wm. P. Johnston, Washington, D. C.*

I make use of it as a text-book, and place it invariably in the hands of my private pupils.

*From Professor D. Humphreys Storer, of Boston.*

I consider it to be the best work on the Diseases of Children we have access to, and as such recommend it to all who ever refer to the subject.

*From Professor M. M. Pallen, of St. Louis.*

I consider it the best treatise on the Diseases of Children that we possess, and as such have been in the habit of recommending it to my classes.

Dr. Condie's scholarship, acumen, industry, and practical sense are manifested in this, as in all his numerous contributions to science.—*Dr. Holmes's Report to the American Medical Association*.

Taken as a whole, in our judgment, Dr. Condie's Treatise is the one from the perusal of which the practitioner in this country will rise with the greatest satisfaction.—*Western Journal of Medicine and Surgery*.

One of the best works upon the Diseases of Children in the English language.—*Western Lancet*.

We feel assured from actual experience that no physician's library can be complete without a copy of this work.—*N. Y. Journal of Medicine*.

Perhaps the most full and complete work now before the profession of the United States; indeed, we may say in the English language. It is vastly superior to most of its predecessors.—*Transylvania Med. Journal*.

A veritable pædiatric encyclopædia, and an honor to American medical literature.—*Ohio Medical and Surgical Journal*.

## WEST ON DISEASES OF CHILDREN.

LECTURES ON THE

# DISEASES OF INFANCY AND CHILDHOOD.

BY CHARLES WEST, M. D.,

Senior Physician to the Royal Infirmary for Children, &amp;c. &amp;c.

In one volume, octavo.

Every portion of these lectures is marked by a general accuracy of description, and by the soundness of the views set forth in relation to the pathology and therapeutics of the several maladies treated of. The lectures on the diseases of the respiratory apparatus, about one-third of the whole number, are particularly excellent, forming one of the fullest and most able accounts of these affections, as they present themselves during infancy and childhood, in the English language. The history of the several forms of phthisis during these periods of existence, with their management, will be read by all with deep interest.—*The American Journal of the Medical Sciences*.

The Lectures of Dr. West, originally published in the London Medical Gazette, form a most valuable addition to this branch of practical medicine. For many years physician to the Children's Infirmary, his opportunities for observing their diseases have been most extensive, no less than 14,000 children having been brought under his notice during the past nine years. These have evidently been studied with great care, and the result has been the production of the very best work in our language, so far as it goes, on the diseases of this class of our patients. The symptomatology and pathology of their diseases are especially exhibited most clearly; and we are convinced that no one can read with care these lectures without deriving from them instruction of the most important kind.—*Charleston Med. Journal*.

## A TREATISE ON THE PHYSICAL AND MEDICAL TREATMENT OF CHILDREN.

BY W. P. DEWEES, M. D.

Ninth edition. In one volume, octavo. 548 pages.



*A NEW WORK—(Lately Issued.)*

# OBSTETRICS:

## THE SCIENCE AND THE ART.

BY CHARLES D. MEIGS, M. D.,

Professor of Midwifery and the Diseases of Women and Children in the Jefferson Medical College,  
Philadelphia, &c. &c.

With One Hundred and Twenty Illustrations.

In one beautifully printed octavo volume, of six hundred and eighty large pages.

As an elementary treatise—concise, but, withal, clear and comprehensive—we know of no one better adapted for the use of the student; while the young practitioner will find in it a body of sound doctrine, and a series of excellent practical directions, adapted to all the conditions of the various forms of labor and their results, which he will be induced, we are persuaded, again and again to consult, and always with profit.

It has seldom been our lot to peruse a work upon the subject, from which we have received greater satisfaction, and which we believe to be better calculated to communicate to the student correct and definite views upon the several topics embraced within the scope of its teachings.—*American Journal of the Medical Sciences.*

We are acquainted with no work on midwifery of greater practical value.—*Boston Medical and Surgical Journal.*

Worthy the reputation of its distinguished author.—*Medical Examiner.*

We most sincerely recommend it, both to the student and practitioner, as a more complete and valuable work on the Science and Art of Midwifery, than any of the numerous reprints and American editions of European works on the same subject.—*N. Y. Annalist.*

We have, therefore, great satisfaction in bringing under our reader's notice the matured views of the highest American authority in the department to which he has devoted his life and talents.—*London Medical Gazette.*

An author of established merit, a professor of Midwifery, and a practitioner of high reputation and immense experience—we may assuredly regard his work now before us as representing the most advanced state of obstetric science in America up to the time at which he writes. We consider Dr. Meigs' book as a valuable acquisition to obstetric literature, and one that will very much assist the practitioner under many circumstances of doubt and perplexity.—*The Dublin Quarterly Journal.*

These various heads are subdivided so well, so lucidly explained, that a good memory is all that is necessary in order to put the reader in possession of a thorough knowledge of this important subject. Dr. Meigs has conferred a great benefit on the profession in publishing this excellent work.—*St. Louis Medical and Surgical Journal.*

No reader will lay the volume down without admiration for the learning and talents of the author. An abler volume, on the whole, we do not hope soon to see.—*Western Journal of Medicine and Surgery.*

A safe and efficient guide to the delicate and oftentimes difficult duties which devolve upon the obstetrician.—*Ohio Medical and Surgical Journal.*

One of the very best treatises on this subject, and worthy of being placed in the library of every American physician.—*Northwestern Medical and Surgical Journal.*

He has an earnest way with him when speaking of the most elementary subjects which fixes the attention and adds much value to the work as a text-book for students.—*British and Foreign Medico-Chirurgical Review.*

TYLER SMITH ON PARTURITION.

# ON PARTURITION,

## AND THE PRINCIPLES AND PRACTICE OF OBSTETRICS.

BY W. TYLER SMITH, M. D.,

Lecturer on Obstetrics in the Hunterian School of Medicine, &amp;c. &amp;c.

In one large duodecimo volume, of 400 pages.

The work will recommend itself by its intrinsic merit to every member of the profession.—*Lancet.*

We can imagine the pleasure with which William Hunter or Denman would have welcomed the present work; certainly the most valuable contribution to obstetrics that has been made since their own day. For ourselves, we consider its appearance as the dawn of a new era in this department of medicine. We do most cordially recommend the work as one absolutely necessary to be studied by every accoucheur. It will, we may add, prove equally interesting and instructive to the student, the general practitioner, and pure obstetrician. It was a bold undertaking to reclaim parturition for Reflex Physiology, and it has been well performed.—*London Journal of Medicine.*

LEE'S CLINICAL MIDWIFERY.

# CLINICAL MIDWIFERY,

COMPRISING THE HISTORIES OF FIVE HUNDRED AND FORTY-FIVE CASES OF DIFFICULT, PRETERNATURAL, AND COMPLICATED LABOR, WITH COMMENTARIES.

BY ROBERT LEE, M. D., F. R. S., &amp;c.

From the 2d London Edition.

In one royal 12mo. volume, extra cloth, of 238 pages.

More instructive to the juvenile practitioner than a score of systematic works.—*Lancet.*

An invaluable record for the practitioner.—*N. Y. Annalist.*

A storehouse of valuable facts and precedents.—*American Journal of the Medical Sciences.*



CHURCHILL'S MIDWIFERY, BY CONDIE, NEW AND IMPROVED EDITION—(Just Issued.)

ON THE

**THEORY AND PRACTICE OF MIDWIFERY.**

BY FLEETWOOD CHURCHILL, M. D., &amp;c.

A NEW AMERICAN FROM THE LAST AND IMPROVED ENGLISH EDITION.

EDITED, WITH NOTES AND ADDITIONS,

BY D. FRANCIS CONDIE, M. D.,

Author of a "Practical Treatise on the Diseases of Children," &amp;c.

**WITH ONE HUNDRED AND THIRTY-NINE ILLUSTRATIONS.**

In one very handsome octavo volume.

In the preparation of the last English edition, from which this is printed, the author has spared no pains, with the desire of bringing it thoroughly up to the present state of obstetric science. The labors of the editor have thus been light, but he has endeavored to supply whatever he has thought necessary to the work, either as respects obstetrical practice in this country, or its progress in Europe since the appearance of Dr. Churchill's last edition. Most of the notes of the former editor, Dr. Huston, have been retained by him, where they have not been embodied by the author in his text. The present edition of the favorite text-book is therefore presented to the profession in the full confidence of its meriting a continuance of the great reputation which it has acquired as a work equally well fitted for the student and practitioner.

To bestow praise on a book that has received such marked approbation would be superfluous. We need only say, therefore, that if the first edition was thought worthy of a favorable reception by the medical public, we can confidently affirm that this will be found much more so. The lecturer, the practitioner, and the student, may all have recourse to its pages, and derive from their perusal much interest and instruction in everything relating to theoretical and practical midwifery.—*Dublin Quarterly Journal of Medical Science.*

A work of very great merit, and such as we can confidently recommend to the study of every obstetric practitioner.—*London Medical Gazette.*

This is certainly the most perfect system extant. It is the best adapted for the purposes of a text-book, and that which he whose necessities confine him to one book, should select in preference to all others.—*Southern Medical and Surgical Journal.*

The most popular work on Midwifery ever issued from the American press.—*Charleston Medical Journal.*

Certainly, in our opinion, the very best work on the subject which exists.—*N. Y. Annalist.*

Were we reduced to the necessity of having but one work on Midwifery, and permitted to choose, we would unhesitatingly take Churchill.—*Western Medical and Surgical Journal.*

It is impossible to conceive a more useful and elegant Manual than Dr. Churchill's Practice of Midwifery.—*Provincial Medical Journal.*

No work holds a higher position, or is more deserving of being placed in the hands of the tyro, the advanced student, or the practitioner.—*Medical Examiner.*

**NEW EDITION OF RAMSBOTHAM ON PARTURITION—(Now Ready, 1851.)****THE PRINCIPLES AND PRACTICE OF  
OBSTETRIC MEDICINE AND SURGERY,**

In reference to the Process of Parturition.

BY FRANCIS H. RAMSBOTHAM, M. D.,

Physician to the Royal Maternity Charity, &amp;c. &amp;c.

SIXTH AMERICAN FROM THE LAST LONDON EDITION.

Illustrated with One Hundred and Forty-eight Figures on Fifty-five Lithographic Plates.

In one large and handsomely printed volume, imperial octavo, with 520 pages.

In this edition the plates have all been redrawn, and the text carefully read and corrected. It is therefore presented as in every way worthy the favor with which it has so long been received.

From Professor Hodge, of the University of Pennsylvania.

To the American public, it is most valuable, from its intrinsic undoubted excellence, and as being the best authorized exponent of British Midwifery. Its circulation will, I trust, be extensive throughout our country.

We recommend the student, who desires to master this difficult subject with the least possible trouble, to possess himself at once of a copy of this work.—*American Journal of the Medical Sciences.*

It stands at the head of the long list of excellent obstetric works published in the last few years in Great Britain, Ireland, and the Continent of Europe. We consider this book indispensable to the library of every physician engaged in the practice of Midwifery.—*Southern Medical and Surgical Journal.*

When the whole profession is thus unanimous in placing such a work in the very first rank as regards the extent and correctness of all the details of the theory and practice of so important a branch of learning, our commendation or condemnation would be of little consequence; but, regarding it as the most useful of all works of the kind, we think it but an act of justice to urge its claims upon the profession.—*N. O. Med. Journal.*

**DEWEES'S MIDWIFERY.****A COMPREHENSIVE SYSTEM OF MIDWIFERY.**

ILLUSTRATED BY OCCASIONAL CASES AND MANY ENGRAVINGS.

BY WILLIAM P. DEWEES, M. D.

Tenth Edition, with the Author's last Improvements and Corrections. In one octavo volume, of 600 pages.



**PEREIRA'S MATERIA MEDICA—Vol. I., (Now Ready.)**

NEW EDITION, GREATLY IMPROVED AND ENLARGED, to Nov. 1851.

## **THE ELEMENTS OF MATERIA MEDICA AND THERAPEUTICS.**

COMPREHENDING THE NATURAL HISTORY, PREPARATION, PROPERTIES, COMPOSITION,  
EFFECTS, AND USES OF MEDICINES.

BY JONATHAN PEREIRA, M. D., F. R. S. AND L. S.

Third American from the Third and Enlarged London Edition.

WITH ADDITIONAL NOTES AND OBSERVATIONS BY THE AUTHOR.

EDITED BY JOSEPH CARSON, M. D.,

Professor of Materia Medica and Pharmacy in the University of Pennsylvania.

In two very large volumes, on small type, with about four hundred illustrations.

The demand for this new edition of "PEREIRA'S MATERIA MEDICA" has induced the publishers to issue the First Volume separately. The Second Volume, now at press and receiving important corrections and revisions from both author and editor, may be expected for publication in July or August, 1852.

The third London edition of this great work has been thoroughly revised and greatly enlarged by the author, who has spared no pains to render it complete in every part, by the addition of a very large amount of matter and the introduction of many new illustrations. The present American edition, however, in addition to this, will not only enjoy the advantages of a careful and accurate superintendence by the editor, but will also embody the additions suggested by a further revision by the author, expressly for this country, embracing the most recent discoveries, and the results of several pharmacopœias which have appeared since the publication of part of the London edition. The notes of the American editor will be prepared with reference to the new edition of the United States Pharmacopœia, and will contain such matter generally as may be required to adapt it fully to the wants of the American student and practitioner, as well as such recent investigations and discoveries as may have escaped the attention of the author. The profession may therefore rely on being able to procure a work which will not only maintain but increase its right to the appellation of

### **AN ENCYCLOPÆDIA OF MATERIA MEDICA AND THERAPEUTICS.**

We shall only remark that every article bears witness to the industry and indefatigable research of the author. Instead of being merely the elements of materia medica, it constitutes a complete encyclopædia of this important subject. The student of physiology, pathology, chemistry, botany, and natural history, will find herein the most recent facts and discoveries in his favorite branch of study, and the medical practitioner will have in this work a safe guide for the administration and employment of medicines.—*London Medical Gazette.*

The present edition (the third) is very much enlarged and improved, and includes the latest discoveries and views respecting medicines and their properties. We believe that this work has no equal in value as a book of reference, or of general information on materia medica.—*The Lancet.*

### **ROYLE'S MATERIA MEDICA.**

## **MATERIA MEDICA AND THERAPEUTICS;**

INCLUDING THE

Preparations of the Pharmacopœias of London, Edinburgh, Dublin, and of the United States.

WITH MANY NEW MEDICINES.

BY J. FORBES ROYLE, M. D., F. R. S.,

Professor of Materia Medica and Therapeutics, King's College, London, &c. &c.

EDITED BY JOSEPH CARSON, M. D.,

Professor of Materia Medica and Pharmacy in the University of Pennsylvania.

WITH NINETY-EIGHT ILLUSTRATIONS.

In one large octavo volume, of about seven hundred pages.

Being one of the most beautiful Medical works published in this country.

This work is, indeed, a most valuable one, and will fill up an important vacancy that existed between Dr. Pereira's most learned and complete system of Materia Medica, and the class of productions on the other extreme, which are necessarily imperfect from their small extent.—*British and Foreign Medical Review.*

### **POCKET DISPENSATORY AND FORMULAR .**

A DISPENSATORY AND THERAPEUTICAL REMEMBRANCER. Comprising the entire lists of Materia Medica, with every Practical Formula contained in the three British Pharmacopœias. With relative Tables subjoined, illustrating by upwards of six hundred and sixty examples, the Extemporaneous Forms and Combinations suitable for the different Medicines. By JOHN MAYNE, M. D., L. R. C. S., EDIN., &c. &c. Edited, with the addition of the formulæ of the United States Pharmacopœia, by R. EGLESFELD GRIFFITH, M. D. In one 12mo. volume, of over three hundred large pages.

The neat typography, convenient size, and low price of this volume, recommend it especially to physicians, apothecaries, and students in want of a pocket manual.



NEW UNIVERSAL FORMULARY.—(Lately Issued.)

**A UNIVERSAL FORMULARY,**

CONTAINING THE

**METHODS OF PREPARING AND ADMINISTERING  
OFFICINAL AND OTHER MEDICINES.**

THE WHOLE ADAPTED TO PHYSICIANS AND PHARMACEUTISTS.

BY R. EGLESFELD GRIFFITH, M. D.,

Author of "American Medical Botany," &amp;c.

In one large octavo volume of 568 pages, double columns.

In this work will be found not only a very complete collection of formulæ and pharmaceutic processes, collected with great care from the best modern authorities of all countries, but also a vast amount of important information on all collateral subjects. To insure the accuracy so necessary to a work of this nature, the sheets have been carefully revised by Dr. Robert Bridges, while Mr. William Procter, Jr., has contributed numerous valuable formulæ, and useful suggestions.

The want of a work like the present has long been felt in this country, where the physician and apothecary have hitherto had access to no complete collection of formulas, gathered from the pharmacopœias and therapeutists of all nations. Not only has this desideratum been thoroughly accomplished in this volume, but it will also be found to contain a very large number of recipes for empirical preparations, valuable to the apothecary and manufacturing chemist, the greater part of which have hitherto not been accessible in this country. It is farther enriched with accurate tables of the weights and measures of Europe; a vocabulary of the abbreviations and Latin terms used in Pharmacy; rules for the administration of medicines; directions for officinal preparations; remarks on poisons and their antidotes; with various tables of much practical utility. To facilitate reference to the whole, extended indices have been added, giving to the work the advantages of both alphabetical and systematic arrangement.

To show the variety and importance of the subjects treated of, the publishers subjoin a very condensed

SUMMARY OF THE CONTENTS, IN ADDITION TO THE FORMULARY PROPER, WHICH EXTENDS TO BETWEEN THREE AND FOUR HUNDRED LARGE DOUBLE-COLUMNED PAGES.

## PREFACE.

## INTRODUCTION.

## WEIGHTS AND MEASURES.

Weights of the United States and Great Britain.—  
Foreign Weights.—Measures.

## SPECIFIC GRAVITY.

TEMPERATURES FOR CERTAIN PHARMACEUTICAL OPERATIONS.

## HYDROMETRICAL EQUIVALENTS.

SPECIFIC GRAVITIES OF SOME OF THE PREPARATIONS OF THE PHARMACOPŒIAS.

RELATION BETWEEN DIFFERENT THERMOMETRICAL SCALES.

EXPLANATION OF PRINCIPAL ABBREVIATIONS USED IN FORMULÆ.

VOCABULARY OF WORDS EMPLOYED IN PRESCRIPTIONS.

## OBSERVATIONS ON THE MANAGEMENT OF THE SICK ROOM.

Ventilation of the Sick room.—Temperature of the Sick room.—Cleanliness in the Sick room.—Quiet in the Sick room.—Examination and Preservation of the Excretions.—Administration of Medicine.—Furniture of a Sick room.—Proper use of Utensils for Evacuations.

## DOSES OF MEDICINES.

Age.—Sex.—Temperament.—Idiosyncrasy.—Habit.—State of the System.—Time of day.—Intervals between Doses.

## RULES FOR ADMINISTRATION OF MEDICINES.

Acids.—Antacids.—Antilithics and Lithontriptics.—Antispasmodics.—Anthelmintics.—Cathartics.—Enemata.—Suppositories.—Demulcents or Emollients.—Diaphoretics.—Diluents.—Diuretics.—Emetics.—Emmenagogues.—Epispastics.—Eruptions.—Escharotics.—Expectorants.—Narcotics.—Refrigerants.—Sedatives.—Sialagogues.—Stimulants.—Tonics.

## MANAGEMENT OF CONVALESCENCE AND RELAPSES.

DIETETIC PREPARATIONS NOT INCLUDED AMONG THE PREVIOUS PRESCRIPTIONS.

## LIST OF INCOMPATIBLES.

PHYSIOLOGICAL TABLES OF THE MOST IMPORTANT MEDICINES.

TABLE OF PHARMACEUTICAL NAMES WHICH DIFFER IN THE U. STATES AND BRITISH PHARMACOPŒIAS.

OFFICINAL PREPARATIONS AND DIRECTIONS.

## INTERNAL REMEDIES.

Powders.—Pills and Boluses.—Extracts.—Confections.—Conserves.—Electuaries.—Pulps.—Syrups.—Mellies or Honeys.—Infusions.—Decoctions.—Tinctures.—Wines.—Vinegars.—Mixtures.—Medicated Waters.—Distilled, Essential, or Volatile Oil.—Fixed Oils and Fats.—Alkaloids.—Spirits.—Troches or Lozenges.—Inhalations.

## EXTERNAL REMEDIES.

Baths.—Cold Bath.—Cool Bath.—Temperate Bath.—Tepid Bath.—Warm Bath.—Hot Bath.—Shower Bath.—Local Baths.—Vapor Bath.—Warm Air Bath.—Douches.—Medicated Baths.—Affusion.—Sponging.—Fomentations.—Cataplasms, or Poultices.—Lotions, Liniments, Embrocations.—Vesicatories, or Blisters.—Issues.—Setons.—Ointments.—Cerates.—Plasters.—Fumigations.

## BLOOD-LETTING.

General Blood-Letting.—Venesection.—Arteriotomy.—Topical Blood-Letting.—Cupping.—Leeching.—Scarifications.

## POISONS.

INDEX OF DISEASES AND THEIR REMEDIES.  
INDEX OF PHARMACEUTICAL AND BOTANICAL NAMES

## GENERAL INDEX.

From the condensed summary of the contents thus given it will be seen that the completeness of this work renders it of much practical value to all concerned in the prescribing or dispensing of medicines.



**GRIFFITH'S MEDICAL FORMULARY—(Continued.)**

From a vast number of commendatory notices, the publishers select a few.

A valuable acquisition to the medical practitioner, and a useful book of reference to the apothecary on numerous occasions.—*American Journal of Pharmacy*.

Dr. Griffith's Formulary is worthy of recommendation, not only on account of the care which has been bestowed on it by its estimable author, but for its general accuracy, and the richness of its details.—*Medical Examiner*.

Most cordially we recommend this Universal Formulary, not forgetting its adaptation to druggists and apothecaries, who would find themselves vastly improved by a familiar acquaintance with this every-day book of medicine.—*The Boston Medical and Surgical Journal*.

Pre-eminent among the best and most useful compilations of the present day will be found the work before us, which can have been produced only at a very great cost of thought and labor. A short description will suffice to show that we do not put too high an estimate on this work. We are not cognizant of the existence of a parallel work. Its value will be apparent to our readers from the sketch of its contents above given.

We strongly recommend it to all who are engaged either in practical medicine, or more exclusively with its literature.—*London Medical Gazette*.

A very useful work, and a most complete compendium on the subject of materia medica. We know of no work in our language, or any other, so comprehensive in all its details.—*London Lancet*.

The vast collection of formulæ which is offered by the compiler of this volume, contains a large number which will be new to English practitioners, some of them from the novelty of their ingredients, and others from the unaccustomed mode in which they are combined; and we doubt not that several of these might be advantageously brought into use. The authority for every formula is given, and the list includes a very numerous assemblage of Continental, as well as of British and American writers of repute. It is, therefore, a work to which every practitioner may advantageously resort for hints to increase his stock of remedies and of forms of prescription.

The other indices facilitate reference to every article in the "Formulary," and they appear to have been drawn up with the same care as that which the author has evidently bestowed on every part of the work.—*The British and Foreign Medico-Chirurgical Review*.

The work before us is all that it professes to be, viz.: "a compendious collection of formulæ and pharmaceutical processes." It is such a work as was much needed, and should be in the hands of every practitioner who is in the habit of compounding medicines.—*Pennsylvania Medical Journal*.

This seems to be a very comprehensive work, so far as the range of its articles and combinations is concerned, with a commendable degree of brevity and condensation in their explanation.

It cannot fail to be a useful and convenient book of reference to the two classes of persons to whom it particularly commends itself in the title-page.—*The N. W. Medical and Surgical Journal*.

It contains so much information that we very cheerfully recommend it to the profession.—*Charleston Medical Journal*.

Well adapted to supply the actual wants of a numerous and varied class of persons.—*N. Y. Journal of Medicine*.

**CHRISTISON & GRIFFITH'S DISPENSATORY.—(A New Work.)**

**A DISPENSATORY,**

OR, COMMENTARY ON THE PHARMACOPŒIAS OF GREAT BRITAIN AND THE UNITED STATES: COMPRISING THE NATURAL HISTORY, DESCRIPTION, CHEMISTRY, PHARMACY, ACTIONS, USES, AND DOSES OF THE ARTICLES OF THE MATERIA MEDICA.

BY ROBERT CHRISTISON, M. D., V. P. R. S. E.,

President of the Royal College of Physicians of Edinburgh; Professor of Materia Medica in the University of Edinburgh, etc.

Second Edition, Revised and Improved,

WITH A SUPPLEMENT CONTAINING THE MOST IMPORTANT NEW REMEDIES.

WITH COPIOUS ADDITIONS,

AND TWO HUNDRED AND THIRTEEN LARGE WOOD ENGRAVINGS.

BY R. EGLESFELD GRIFFITH, M. D.,

Author of "A Medical Botany," etc.

In one very large and handsome octavo volume, of over one thousand closely printed pages, with numerous wood-cuts, beautifully printed on fine white paper, presenting an immense quantity of matter at an unusually low price.

It is enough to say that it appears to us as perfect as a Dispensatory, in the present state of pharmaceutical science, could be made.—*The Western Journal of Medicine and Surgery*.

**CARSON'S SYNOPSIS—(Now Ready.)**

SYNOPSIS OF THE

**COURSE OF LECTURES ON MATERIA MEDICA AND PHARMACY,**

Delivered in the University of Pennsylvania.

BY JOSEPH CARSON, M. D.,

Professor of Materia Medica and Pharmacy in the University of Pennsylvania.

In one very neat octavo volume of 208 pages.

This work, containing a rapid but thorough outline of the very extensive subjects under consideration, will be found useful, not only for the matriculants and graduates of the institution for whom it is more particularly intended, but also for those of the profession who may desire to recall their former studies.

**THE THREE KINDS OF COD-LIVER OIL,**

Comparatively considered, with their Chemical and Therapeutic Properties, by L. J. DE JONGH, M. D. Translated, with an Appendix and Cases, by EDWARD CAREY, M. D. To which is added an article on the subject from "Dunlison on New Remedies." In one small 12mo. volume, extra cloth.



**DUNGLISON'S THERAPEUTICS.**  
**NEW AND IMPROVED EDITION.—(Lately Issued.)**

**GENERAL THERAPEUTICS AND MATERIA MEDICA;**

ADAPTED FOR A MEDICAL TEXT-BOOK,

BY ROBLEY DUNGLISON, M. D.,

Professor of Institutes of Medicine, &c., in Jefferson Medical College; Late Professor of *Materia Medica*, &c. in the Universities of Maryland and Virginia, and in Jefferson Medical College.

FOURTH EDITION, MUCH IMPROVED.

**With One Hundred and Eighty-two Illustrations.**

In two large and handsomely printed octavo volumes.

The present edition of this standard work has been subjected to a thorough revision both as regards style and matter, and has thus been rendered a more complete exponent than heretofore of the existing state of knowledge on the important subjects of which it treats. The favor with which the former editions have everywhere been received seemed to demand that the present should be rendered still more worthy of the patronage of the profession, and of the medical student in particular, for whose use more especially it is proposed; while the number of impressions through which it has passed has enabled the author so to improve it as to enable him to present it with some degree of confidence as well adapted to the purposes for which it is intended. In the present edition, the remedial agents of recent introduction have been inserted in their appropriate places; the number of illustrations has been greatly increased, and a copious index of diseases and remedies has been appended, improvements which can scarcely fail to add to the value of the work to the therapeutical inquirer.

The publishers, therefore, confidently present the work as it now stands to the notice of the practitioner as a trustworthy book of reference, and to the student, for whom it was more especially prepared, as a full and reliable text-book on General Therapeutics and *Materia Medica*.

Notwithstanding the increase in size and number of illustrations, and the improvements in the mechanical execution of the work, its price has not been increased.

In this work of Dr. Dunglison, we recognize the same untiring industry in the collection and embodying of facts on the several subjects of which he treats, that has heretofore distinguished him, and we cheerfully point to these volumes, as two of the most interesting that we know of. In noticing the additions to this, the fourth edition, there is very little in the periodical or annual literature of the profession, published in the interval which has elapsed since the issue of the first, that has escaped the careful search of the author. As a book for reference, it is invaluable.—*Charleston Med. Journal and Review*.

It may be said to be the work now upon the subjects upon which it treats.—*Western Lancet*.

As a text-book for students, for whom it is particularly designed, we know of none superior to it.—*St. Louis Medical and Surgical Journal*.

It purports to be a new edition, but it is rather a new book, so greatly has it been improved both in the amount and quality of the matter which it contains.—*N. O. Medical and Surgical Journal*.

We bespeak for this edition from the profession an increase of patronage over any of its former ones, on account of its increased merit.—*N. Y. Journal of Medicine*.

We consider this work unequalled.—*Boston Med. and Surg. Journal*.

**NEW AND MUCH IMPROVED EDITION—Brought up to 1851.—(Just Issued.)**

**NEW REMEDIES,**

**WITH FORMULÆ FOR THEIR ADMINISTRATION.**

BY ROBLEY DUNGLISON, M. D.,

PROFESSOR OF THE INSTITUTES OF MEDICINE, ETC. IN THE JEFFERSON MEDICAL COLLEGE OF PHILADELPHIA.

**Sixth Edition, with extensive Additions.**

In one very large octavo volume, of over seven hundred and fifty pages.

The fact that this work has rapidly passed to a **SIXTH EDITION** is sufficient proof that it has supplied a desideratum to the profession in presenting them with a clear and succinct account of all new and important additions to the *materia medica*, and novel applications of old remedial agents. In the preparation of the present edition, the author has shrunk from no labor to render the volume worthy of a continuance of the favor with which it has been received, as is sufficiently shown by the increase of about one hundred pages in the size of the work. The necessity of such large additions arises from the fact that the last few years have been rich in valuable gifts to Therapeutics; and amongst these, ether, chloroform, and other so called anesthetics, are worthy of special attention. They have been introduced since the appearance of the last edition of the "**NEW REMEDIES**." Other articles have been proposed for the first time, and the experience of observers has added numerous interesting facts to our knowledge of the virtues of remedial agents previously employed.

The therapeutical agents now first admitted into this work, some of which have been newly introduced into pharmacology, and the old agents brought prominently forward with novel applications, and which may consequently be regarded as *New Remedies*, are the following:—*Adansonia digitata*, Benzoate of Ammonia, Valerianate of Bismuth, Sulphate of Cadmium, Chloroform, Collodion, Cantharidal Collodion, *Cotyledon Umbilicus*, Sulphuric Ether, Strong Chloric Ether, Compound Ether, *Hura Braziliensis*, *Iberis Amara*, Iodic Acid, Iodide of Chloride of Mercury, Powdered Iron, Citrate of Magnetic Oxide of Iron, Citrate of Iron and Magnesia, Sulphate of Iron and Alumina, Tannate of Iron, Valerianate of Iron, Nitrate of Lead, Lemon Juice, Citrate of Magnesia, Salts of Manganese, *Oleum Cadinum*, Arsenite of Quinia, Hydriodate of Iron and Quinia, *Sanicula Marylandica*, and Sumbul.



MOHR, REDWOOD, AND PROCTER'S PHARMACY.—*Lately Issued.***PRACTICAL PHARMACY.**

COMPRISING THE ARRANGEMENTS, APPARATUS, AND MANIPULATIONS OF THE PHARMACEUTICAL SHOP AND LABORATORY.

BY FRANCIS MOHR, PH. D.,

Assessor Pharmacie of the Royal Prussian College of Medicine, Coblenz;

AND THEOPHILUS REDWOOD,

Professor of Pharmacy in the Pharmaceutical Society of Great Britain.

EDITED, WITH EXTENSIVE ADDITIONS, BY PROFESSOR WILLIAM PROCTER,  
Of the Philadelphia College of Pharmacy.

In one handsomely printed octavo volume, of 570 pages, with over 500 engravings on wood.

To physicians in the country, and those at a distance from competent pharmacutists, as well as to apothecaries, this work will be found of great value, as embodying much important information which is to be met with in no other American publication.

After a pretty thorough examination, we can recommend it as a highly useful book, which should be in the hands of every apothecary. Although no instruction of this kind will enable the beginner to acquire that practical skill and readiness which experience only can confer, we believe that this work will much facilitate their acquisition, by indicating means for the removal of difficulties as they occur, and suggesting methods of operation in conducting pharmaceutical processes which the experimenter would only hit upon after many unsuccessful trials; while there are few pharmacutists, of however extensive experience, who will not find in it valuable hints that they can turn to use in conducting the affairs of the shop and laboratory. The mechanical execution of the work is in a style of unusual excellence. It contains about five hundred and seventy large octavo pages, handsomely printed on good paper, and illustrated by over five hundred remarkably well-executed wood-cuts of chemical and pharmaceutical apparatus. It comprises the whole of Mohr and Redwood's book, as published in London, rearranged and classified by the American editor, who has added much valuable new matter, which has increased the size of the book more than one-fourth, including about one hundred additional wood-cuts.—*The American Journ. of Pharmacy.*

It is a book, however, which will be in the hands of almost every one who is much interested in pharmaceutical operations, as we know of no other publication so well calculated to fill a void long felt.—*The Medical Examiner.*

The country practitioner who is obliged to dispense his own medicines, will find it a most valuable assistant.—*Monthly Journal and Retrospect.*

The book is strictly practical, and describes only manipulations or methods of performing the numerous processes the pharmacist has to go through, in the preparation and manufacture of medicines, together with all the apparatus and fixtures necessary thereto. On these matters, this work is very full and complete, and details, in a style uncommonly clear and lucid, not only the more complicated and difficult processes, but those not less important ones, the most simple and common. The volume is an octavo of five hundred and seventy-six pages. It is elegantly illustrated with a multitude of neat wood engravings, and is unexceptionable in its whole typographical appearance and execution. We take great satisfaction in commending this so much needed treatise, not only to those for whom it is more specially designed, but to the medical profession generally—to every one, who, in his practice, has occasion to prepare, as well as administer medical agents.—*Buffalo Medical Journal.*

**NEW AND COMPLETE MEDICAL BOTANY.****MEDICAL BOTANY;**

OR, A DESCRIPTION OF ALL THE MORE IMPORTANT PLANTS USED IN MEDICINE, AND OF THEIR PROPERTIES, USES, AND MODES OF ADMINISTRATION.

BY R. EGLESFELD GRIFFITH, M. D., &amp;c. &amp;c.

In one large 8vo. vol. of 704 pages, handsomely printed, with nearly 350 illustrations on wood.

One of the greatest acquisitions to American medical literature. It should by all means be introduced at the very earliest period, into our medical schools, and occupy a place in the library of every physician in the land.—*Southwestern Medical Advocate.*

Admirably calculated for the physician and student—we have seen no work which promises greater advantages to the profession.—*N. O. Medical and Surgical Journal.*

One of the few books which supply a positive deficiency in our medical literature.—*Western Lancet.*

We hope the day is not distant when this work will not only be a text-book in every medical school and college in the Union, but find a place in the library of every private practitioner.—*N. Y. Journ. of Medicine.*

**ELLIS'S MEDICAL FORMULARY.—Improved Edition.****THE MEDICAL FORMULARY:**

BEING A COLLECTION OF PRESCRIPTIONS, DERIVED FROM THE WRITINGS AND PRACTICE OF MANY OF THE MOST EMINENT PHYSICIANS OF AMERICA AND EUROPE.

To which is added an Appendix, containing the usual Dietetic Preparations and Antidotes for Poisons.

THE WHOLE ACCOMPANIED WITH A FEW BRIEF PHARMACEUTIC AND MEDICAL OBSERVATIONS.

BY BENJAMIN ELLIS, M. D.

NINTH EDITION, CORRECTED AND EXTENDED, BY SAMUEL GEORGE MORTON, M. D.

In one neat octavo volume of 268 pages.

**CARPENTER ON ALCOHOLIC LIQUORS.—(A New Work.)**

A Prize Essay on the Use of Alcoholic Liquors in Health and Disease. By William B. Carpenter, M. D., author of "Principles of Human Physiology," &c. In one 12mo. volume.



NEW AND IMPROVED EDITION—(Lately Issued.)

**ELEMENTARY CHEMISTRY,  
THEORETICAL AND PRACTICAL.**

BY GEORGE FOWNES, Ph. D.,

Chemical Lecturer in the Middlesex Hospital Medical School, &amp;c. &amp;c.

WITH NUMEROUS ILLUSTRATIONS.

THIRD AMERICAN, FROM A LATE LONDON EDITION. EDITED, WITH ADDITIONS,

BY ROBERT BRIDGES, M. D.,

Professor of General and Pharmaceutical Chemistry in the Philadelphia College of Pharmacy, &amp;c. &amp;c.

In one large royal 12mo. vol., of over 500 pages, with about 180 wood-cuts, sheep or extra cloth.

At the time of his death, Professor Fownes had just completed the revision of this work for his third edition, and, at his request, Dr. H. Bence Jones undertook the office of seeing it through the press, and making such additions in the department of Animal Chemistry as were rendered necessary by the numerous discoveries daily making in that branch of the science. The task of the American editor, therefore, has merely been to add such new matter as may since have appeared, and to adapt the whole to the wants of the American student, by appending in the form of notes such points of interest as would be calculated to retain the position which the original has so justly obtained, and to maintain it on an equality with the rapid advance of chemical science. It will, therefore, be found considerably enlarged and greatly improved. Notwithstanding its increase in size, it has been kept at its former extremely low price, and may now be considered as one of the

**CHEAPEST TEXT-BOOKS ON CHEMISTRY NOW EXTANT.**

The work of Dr. Fownes has long been before the public, and its merits have been fully appreciated as the best text-book on Chemistry now in existence. We do not, of course, place it in a rank superior to the works of Brande, Graham, Turner, Gregory, or Gmelin, but we say that, as a work for students, it is preferable to any of them.—*London Journal of Medicine.*

The rapid sale of this Manual evinces its adaptation to the wants of the student of chemistry, whilst the well known merits of its lamented author have constituted a guarantee for its value, as a faithful exposition of the general principles and most important facts of the science to which it professes to be an introduction.

We have only to add, that Dr. Bence Jones appears to have performed his editorial task most thoroughly, the want of the author's final supervision being nowhere discoverable.—*The British and Foreign Medical-Chirurgical Review.*

A work well adapted to the wants of the student. It is an excellent exposition of the chief doctrines and facts of modern chemistry, originally intended as a guide to the lectures of the author, corrected by his own hand shortly before his death in 1849, and recently revised by Dr. Bence Jones, who has made some additions to the chapter on animal chemistry. Although not intended to supersede the more extended treatises on chemistry, Professor Fownes' Manual may, we think, be often used as a work of reference, even by those advanced in the study, who may be desirous of refreshing their memory on some forgotten point. The size of the work, and still more the condensed yet perspicuous style in which it is written, absolve it from the charges very properly urged against most manuals termed popular, viz., of omitting details of indispensable importance, of avoiding technical difficulties, instead of explaining them, and of treating subjects of high scientific interest in an unscientific way.—*Edinburgh Monthly Journal of Medical Science.*

BOWMAN'S MEDICAL CHEMISTRY—(Lately Issued.)

**PRACTICAL HANDBOOK OF MEDICAL CHEMISTRY.**

BY JOHN E. BOWMAN, M. D.

In one neat volume, royal 12mo., with numerous illustrations.

We cannot too highly commend the very elaborate, yet clear and distinct manner, in which the appearances of these fluids, and their variations in disease, are described. To the practitioner, the book is specially recommended, as giving a very clear account of many chemical matters, which must be ever coming before him in his daily practice. Every practitioner, and every student of clinical medicine, should endeavor to enrich his collection of books with Mr. Bowman's little volume.—*London Journal of Medicine.*

Mr. Bowman has succeeded in supplying a desideratum in medical literature. In the little volume before us, he has given a concise but comprehensive account of all matters in chemistry which the man in practice may desire to know.—*Lancet.*

BY THE SAME AUTHOR—(Lately Issued.)

**INTRODUCTION TO PRACTICAL CHEMISTRY, Including Analysis.**

With Numerous Illustrations. In one neat volume, royal 12mo.

GARDNER'S MEDICAL CHEMISTRY.

**MEDICAL CHEMISTRY,**

FOR THE USE OF STUDENTS AND THE PROFESSION;

BEING A MANUAL OF THE SCIENCE, WITH ITS APPLICATIONS TO TOXICOLOGY, PHYSIOLOGY, THERAPEUTICS, HYGIENE, &amp;c.

BY D. PEREIRA GARDNER, M. D.

In one handsome royal 12mo. volume, with illustrations.

**New Edition, Preparing.—THE ELEMENTS OF CHEMISTRY.**

INCLUDING THE APPLICATION OF THE SCIENCE TO THE ARTS. WITH NUMEROUS ILLUSTRATIONS.

BY THOMAS GRAHAM, F. R. S., L. E. &amp; D.

WITH NOTES AND ADDITIONS BY ROBERT BRIDGES, M. D., &amp;c. &amp;c.

**SIMON'S ANIMAL CHEMISTRY**, with Reference to the Physiology and Pathology of Man. By G. E. DAY. One vol. 8vo., 700 pages



**TAYLOR'S MEDICAL JURISPRUDENCE.**  
**MEDICAL JURISPRUDENCE.**

BY ALFRED S. TAYLOR,

SECOND AMERICAN, FROM THE THIRD AND ENLARGED LONDON EDITION.  
 With numerous Notes and Additions, and References to American Practice and Law.

BY R. E. GRIFFITH, M. D.

In one large octavo volume.

This work has been much enlarged by the author, and may now be considered as the standard authority on the subject, both in England and this country. It has been thoroughly revised, in this edition, and completely brought up to the day with reference to the most recent investigations and decisions. No further evidence of its popularity is needed than the fact of its having, in the short time that has elapsed since it originally appeared, passed to three editions in England, and two in the United States.

We recommend Mr. Taylor's work as the ablest, most comprehensive, and, above all, the most practically useful book which exists on the subject of legal medicine. Any man of sound judgment, who has mastered the contents of Taylor's "Medical Jurisprudence," may go into a court of law with the most perfect confidence of being able to acquit himself creditably.—*Medico-Chirurgical Review*.

The most elaborate and complete work that has yet appeared. It contains an immense quantity of cases lately tried, which entitle it to be considered what Beck was in its day.—*Dublin Medical Journal*.

**TAYLOR ON POISONS.**

**ON POISONS,**  
**IN RELATION TO MEDICAL JURISPRUDENCE AND MEDICINE.**

BY ALFRED S. TAYLOR, F. R. S., &c.

EDITED, WITH NOTES AND ADDITIONS, BY R. E. GRIFFITH, M. D.

In one large octavo volume, of 688 pages.

The most elaborate work on the subject that our literature possesses.—*Brit. and For. Medico-Chirur. Review*.  
 One of the most practical and trustworthy works on Poisons in our language.—*Western Journal of Med.*  
 It contains a vast body of facts, which embrace all that is important in toxicology, all that is necessary to the guidance of the medical jurist, and all that can be desired by the lawyer.—*Medico-Chirurgical Review*.

It is, so far as our knowledge extends, incomparably the best upon the subject; in the highest degree creditable to the author, entirely trustworthy, and indispensable to the student and practitioner.—*N. Y. Analyst*.

**BEALE ON HEALTH—NOW READY.**

**THE LAWS OF HEALTH IN RELATION TO MIND AND BODY.**

A SERIES OF LETTERS FROM AN OLD PRACTITIONER TO A PATIENT.

BY LIONEL JOHN BEALE, M. R. C. S., &c.

In one handsome volume, royal 12mo., extra cloth.

The "Laws of Health," in relation to mind and body, is a book which will convey much instruction to non-professional readers; they may, from these letters, glean the principles upon which young persons should be educated, and derive much useful information, which will apply to the preservation of health at all ages.—*Med. Times*.

**GREGORY ON ANIMAL MAGNETISM—(Now Ready.)**

**LETTERS TO A CANDID ENQUIRER**  
**ON ANIMAL MAGNETISM.**

DESCRIPTION AND ANALYSIS OF THE PHENOMENA. DETAILS OF FACTS AND CASES.

BY WILLIAM GREGORY, M. D., F. R. S. E.,

Professor of Chemistry in the University of Edinburgh, &c.

In one neat volume, royal 12mo., extra cloth.

**TRANSACTIONS OF THE**  
**AMERICAN MEDICAL ASSOCIATION.**

VOLUME I. FOR 1848, VOL. II. FOR 1849, VOL. III. FOR 1850.

Large octavo, extra cloth, or paper covers for mailing.

Any volume sold separate, or the whole in sets at a reduced price.

Orders for the supply of Medical Societies should be sent direct to the Treasurer of the Association, Isaac Hays, M. D., care of Blanchard & Lea, with the amount enclosed.

**DUNGLISON ON HUMAN HEALTH—HUMAN HEALTH,** or the Influence of Atmosphere and Locality, Change of Air and Climate, Seasons, Food, Clothing, Bathing, Exercise, Sleep, &c. &c. on healthy man; constituting Elements of Hygiene. Second edition, with many modifications and additions. By Robley Dunglison, M. D. &c. &c. In one octavo volume of 464 pages.

**DUNGLISON'S MEDICAL STUDENT.**—The Medical Student, or Aids to the Study of Medicine. Revised and Modified Edition. 1 vol. royal 12mo., extra cloth. 312 pp.

**BARTLETT'S PHILOSOPHY OF MEDICINE.**—An Essay on the Philosophy of Medical Science. In one handsome 8vo. volume. 312 pp.

**BARTLETT ON CERTAINTY IN MEDICINE.**—An Inquiry into the Degree of Certainty in Medicine, and into the Nature and Extent of its Power over Disease. In one vol. royal 12mo. 84 pp.



# THE GREAT AMERICAN MEDICAL DICTIONARY.

New and Enlarged Edition, brought up to October, 1851.

**NOW READY.**

## MEDICAL LEXICON;

### A DICTIONARY OF MEDICAL SCIENCE,

Containing a Concise Explanation of the various Subjects and Terms of  
PHYSIOLOGY, PATHOLOGY, HYGIENE, THERAPEUTICS, PHARMACOLOGY, OB-  
STETRICS, MEDICAL JURISPRUDENCE, &c.

**WITH THE FRENCH AND OTHER SYNONYMES.**

NOTICES OF CLIMATE AND OF CELEBRATED MINERAL WATERS;  
Formulæ for various Official, Empirical, and Dietetic Preparations, &c.

**BY ROBLEY DUNGLISON, M. D.,**

Professor of Institutes of Medicine, &c., in Jefferson Medical College, Philadelphia, &c.

**EIGHTH EDITION,**

**REVISED AND GREATLY ENLARGED.**

In one very thick 8vo. vol., of 927 large double-columned pages, strongly bound, with raised bands.

Every successive edition of this work bears the marks of the industry of the author, and of his determination to keep it fully on a level with the most advanced state of medical science. Thus the last two editions contained about NINE THOUSAND SUBJECTS AND TERMS not comprised in the one immediately preceding, and the present has not less than FOUR THOUSAND not in any former edition. As a complete Medical Dictionary; therefore, embracing over FIFTY THOUSAND definitions, in all the branches of the science, it is presented as meriting a continuance of the great favor and popularity which have carried it, within no very long space of time, to an eighth edition.

Every precaution has been taken in the preparation of the present volume, to render its mechanical execution and typographical accuracy worthy of its extended reputation and universal use. The very extensive additions have been accommodated, without materially increasing the bulk of the volume, by the employment of a small but exceedingly clear type, cast for this purpose. The press has been watched with great care, and every effort used to insure the verbal accuracy so necessary to a work of this nature. The whole is printed on fine white paper; and while thus exhibiting in every respect so great an improvement over former issues, it is presented at the original exceedingly low price.

A few notices of former editions are subjoined.

Dr. Dunglison's Lexicon has the rare merit that it certainly has no rival in the English language for accuracy and extent of references. The terms generally include short physiological and pathological descriptions, so that, as the author justly observes, the reader does not possess in this work a mere dictionary, but a book, which, while it instructs him in medical etymology, furnishes him with a large amount of useful information. That we are not over-estimating the merits of this publication, is proved by the fact that we have now before us the seventh edition. This, at any rate, shows that the author's labors have been properly appreciated by his own countrymen; and we can only confirm their judgment, by recommending this most useful volume to the notice of our cisatlantic readers. No medical library will be complete without it.—*The London Med. Gazette.*

It is certainly more complete and comprehensive than any with which we are acquainted in the English language. Few, in fact, could be found better qualified than Dr. Dunglison for the production of such a work. Learned, industrious, persevering, and accurate, he brings to the task all the peculiar talents necessary for its successful performance: while, at the same time, his familiarity with the writings of the ancient and modern "masters of our art," renders him skilful to note the exact usage of the several terms of science, and the various modifications which medical terminology has undergone with the change of theories or the progress of improvement.—*American Journal of the Medical Sciences.*

One of the most complete and copious known to the cultivators of medical science.—*Boston Med. Journal.*

This most complete Medical Lexicon—certainly one of the best works of the kind in the language.—*Charleston Medical Journal.*

The most complete Medical Dictionary in the English language.—*Western Lancet.*

Dr. Dunglison's Dictionary has not its superior, if indeed its equal, in the English language.—*St. Louis Med. and Surg. Journal.*

Familiar with nearly all the medical dictionaries now in print, we consider the one before us the most complete, and an indispensable adjunct to every medical library.—*British American Medical Journal.*

Admitted by all good judges, both in this country and in Europe, to be equal, and in many respects superior to any other work of the kind yet published.—*Northwestern Medical and Surgical Journal.*

We repeat our former declaration that this is the best Medical Dictionary in the English language.—*Western Lancet.*

We have no hesitation to pronounce it the very best Medical Dictionary now extant.—*Southern Medical and Surgical Journal.*

The most comprehensive and best English Dictionary of medical terms extant.—*Buffalo Med. Journal.*

Whence the terms have all been derived we find it rather difficult to imagine. We can only say that, after looking for every new and strange word we could think of, we have not been disappointed in regard to more than a few of most recent introduction, such as the designations given by Professor Owen to the component parts of a Vertebra.—*British and Foreign Medico-Chirurgical Review.*

Dr. Dunglison's masterpiece of literary labor.—*N. Y. Journal of Medicine.*

## HOBLYN'S MEDICAL DICTIONARY.

### A DICTIONARY OF THE TERMS USED IN MEDICINE

AND THE COLLATERAL SCIENCES.

**BY RICHARD D. HOBLYN, A. M., OXON.**


REVISED, WITH NUMEROUS ADDITIONS, FROM THE SECOND LONDON EDITION,  
BY ISAAC HAYS, M. D., &c. In one large royal 12mo. volume of 402 pages, double columns.

We cannot too strongly recommend this small and cheap volume to the library of every student and practitioner.—*Medico-Chirurgical Review.*











30



Date Due

YALE  
MEDICAL  
LIBRARY

Demco 293-5



R0125  
850G



09-08 STD



[www.colibrisystem.com](http://www.colibrisystem.com)



