The druggist's general receipt book : comprising a copious veterinary formulary, with numerous recipes in patent and proprietary medicines, druggists' nostrums, etc. : perfumery and cosmetics; beverages, dietetic articles, and condiments : trade chemicals, scientific processes, and an appendix of useful tables / by Henry Beasley.

Contributors

Beasley, Henry.
Francis A. Countway Library of Medicine

Publication/Creation

Philadelphia : Lindsay & Blakiston, 1871.

Persistent URL

https://wellcomecollection.org/works/g69f3pdv

License and attribution

This material has been provided by This material has been provided by the Francis A. Countway Library of Medicine, through the Medical Heritage Library. The original may be consulted at the Francis A. Countway Library of Medicine, Harvard Medical School. where the originals may be consulted. This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.

wellcome collection

Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
https://wellcomecollection.org

22. G 155

Digitized by the Internet Archive
in 2011 with funding from
Open Knowledge Commons and Harvard Medical School

http://www.archive.org/details/druggistsgeneral1871beas

THE DRUGGIST'S

GENERAL RECEIPT BOOK.

BEASLEY'S BOOK OF PRESCRIPTIONS. A new, Revised, and Enlarged Edition, containing 3000 Prescriptions, collected from the Practice of the most Eminent Physicians and Surgeons,—English, French, and American; comprising also a Compendious History of the Materia Medica, Lists of the Doses of all Official and Established Preparations, and an Index of Diseases and their Remedies. By HENRY BEASLEY. Octavo. Price $4.00.

The editor, carefully selecting from the mass of materials at his disposal, has compiled a volume, sufficiently comprehensive, and yet sufficiently portable, in which both physician and druggist, prescriber and compounder, may find, under the head of each remedy, the manner in which that remedy may be most effectively administered, or combined with other medicines in the treatment of various diseases. The alphabetical arrangement adopted renders this easy; and the value of the volume is still further enhanced by the short account given of each medicine, and the lists of doses of its several preparations. It is really a most useful and important publication, and, from the great aid which it is capable of affording in prescribing, it should be in the possession of every medical practitioner. Amongst other advantages is, that by giving the prescriptions of some of the most able and successful practitioners of the day, it affords an insight into the methods of treatment pursued by them, and of the remedies which they chiefly employed in the treatment of different diseases.—*London Lancet*

NEW EDITIONS, JUST READY.

WARING'S PRACTICAL THERAPEUTICS. Considered chiefly with reference to Articles of Materia Medica, with an Index of Diseases, &c. By EDWARD JOHN WARING, F.R.C.S., F.L.S., &c., &c. Second American, from the Third London Edition. Cloth, $5.00; leather, $6.00.

BIDDLE'S MATERIA MEDICA. For the Use of Students. With Illustrations. By J. B. BIDDLE, M.D., Professor of Materia Medica and Therapeutics in the Jefferson Medical College, Philadelphia, &c., &c. Fourth Edition. Revised and Enlarged. $4.00

HEADLAND ON THE ACTION OF MEDICINES IN THE SYSTEM. By F. W. HEADLAND, M.D., Fellow of the Royal College of Physicians, &c., &c. Sixth American, from the Fourth London Edition. Revised and Enlarged. $3.00.

PEREIRA'S PHYSICIAN'S PRESCRIPTION BOOK. Containing Lists of Terms, Phrases, Contractions, and Abbreviations, used in Prescriptions, with Explanatory Notes, the Grammatical Constructions of Prescriptions, Rules for the Pronunciation of Pharmaceutical Terms, &c., &c. By JONATHAN PEREIRA, M.D., F.R.S., &c. From the Fifteenth London Edition. Cloth, $1.25; leather, with tucks and pocket, $1.50.

WYTHES' PHYSICIAN'S POCKET, DOSE, AND SYMPTOM BOOK. Containing the Doses and Uses of all the Principal Articles of the Materia Medica, and Original Preparations; A Table of Weights and Measures, Rules to Proportion the Doses of Medicines, &c., &c. By JOSEPH H. WYTHES, A.M., M.D., &c. Tenth Revised Edition. Cloth, $1.00; leather, tucks, with pockets, $1.25.

THE DRUGGIST'S

GENERAL RECEIPT BOOK:

COMPRISING A COPIOUS

VETERINARY FORMULARY;

WITH NUMEROUS RECIPES IN PATENT AND PROPRIETARY MEDICINES, DRUGGISTS' NOSTRUMS, ETC.; PERFUMERY AND COSMETICS; BEVERAGES, DIETETIC ARTICLES, AND CONDIMENTS; TRADE CHEMICALS, SCIENTIFIC PROCESSES, AND AN APPENDIX OF USEFUL TABLES.

BY

HENRY BEASLEY,

AUTHOR OF THE "BOOK OF PRESCRIPTIONS," &C., &C.

SEVENTH AMERICAN,

FROM THE LAST LONDON EDITION.

PHILADELPHIA:
LINDSAY & BLAKISTON.
1871.

11282

PRINTED BY HENRY B. ASHMEAD,
1102 and 1104 Sansom Street.

ADVERTISEMENT.

THE compiler of the present volume was encouraged to its preparation by the great success of his "Pocket Formulary," which preceded it, and by many suggestions from correspondents as to the probable utility of this Book of Receipts. Such anticipations of its usefulness have already been more than realized by the rapid sale of five large editions. The compiler endeavored to produce a work which should be useful to those for whom it was designed. It was his aim to collect from various and widely scattered sources, and to condense into a volume of moderate size and convenient arrangement, a considerable amount of information that might be useful to Chemists and Druggists. And it is trusted that every purchaser will be able to find, in its multifarious contents, something that may be deemed an equivalent for so small an outlay.

In preparing a Third Edition of the Druggist's General Receipt Book, the whole underwent a careful revision. While the addition to the work of upwards of two hundred new forms, receipts, and processes, collected from the most approved authorities, will, it is hoped, have established for it a further claim to the approbation of those for whose benefit it has been written.

In the Fourth Edition, a further improvement was aimed at; much attention having been given, more especially, to the Chemical Formulæ and Processes described in the miscellaneous department of the work, which will now be found more complete than before.

To the Fifth and Sixth Editions such additions have been made as have been rendered possible by the advance of science and discovery.

CONTENTS.

PAGE

VETERINARY MATERIA MEDICA, ... 17
VETERINARY FORMULARY, ... 58
Medicines for Horses, ... 58
Balls and Ball Masses, ... 58
Electuaries, Confections, &c., ... 74
Powders, ... 75
Liquid Medicines—Drenches, ... 79
Tinctures, Solutions, &c., ... 90
Clysters, ... 91
External Applications—Liniments, ... 93
Embrocations, various, ... 98
Lotions, or Washes, ... 99
Caustics, ... 102
Fomentations, ... 104
Eye Waters, ... 105
Sundry Solutions, ... 106
Poultices, ... 107
Ointments, ... 109
Charges, ... 119
Powders (for outward use), ... 119
Medicines for Cattle, ... 121
Drenches, &c., ... 121
for Calves, ... 133
External Applications—Lotions, ... 134
Eye Waters, ... 136
Embrocations and Liniments, ... 136
Ointments, &c., ... 138
Medicines for Sheep and Lambs, ... 141
Drenches, &c., ... 141
External Applications—Eye Waters, ... 145
Washes, ... 145
Powders and Ointments, ... 146
Medicines for Swine, ... 150
Medicines for Dogs, ... 152
External Applications, ... 159
Medicines for Poultry and Rabbits, ... 163
PATENT AND PROPRIETARY MEDICINES, DRUGGISTS' NOSTRUMS, &c., . 166
Factitious Mineral Waters, and Salts for producing them, . . 205
PERFUMERY, ... 212
Distilled Waters, ... 212
Spirituous Waters, Eaux, Esprits, ... 213
Tinctures used in Perfumery, ... 219
Extraits, Essences, ... 220
Common French Essences, Spirits, and Waters, ... 224
Ammoniated Perfumes, ... 225
Acetic Perfumes, ... 226
Pot Pourri, Scented Powders, &c., ... 227
Pastilles, ... 229

PAGE

COSMETICS, ... 232
Skin Cosmetics—Lotions, ... 232
Pastes, Pomades, Cold Cream, &c., ... 233
Face Paints, ... 237
Toilet Soaps, ... 238
Shaving Compounds, ... 240
Hair Cosmetics, ... 242
Compounds to promote the Growth of the Hair, ... 242
Liquid ditto, ... 244
Pomatums, Lotions, for embellishing the Hair, &c. . . . 245
Huiles Antiques, ... 248
Washes for Cleansing the Hair, ... 250
Compounds for Stiffening the Hair, ... 251
Hair Dyes, ... 251
Pomatums, or Cosmetics, in Sticks, for the Hair, ... 253
Depilatories, ... 254
Teeth and Mouth Cosmetics, ... 255
Tooth Powders, ... 255
Tooth Pastes, ... 259
Astringent Tinctures—Mouth Washes, ... 261
Tinctures, &c., for Toothache, ... 263
Pastes for Toothache, ... 265
Cements for Stopping Teeth, &c., ... 265
BEVERAGES, DIETETIC ARTICLES, AND CONDIMENTS, ... 268
Beverages, and Powders for preparing them, ... 268
Dietetic Articles—Directions for using, ... 276
Preparations of Cod-liver, ... 289
Condiments, &c., ... 290
Curry Powders, Mixed Spices, &c., ... 290
Culinary Essences, Tinctures, &c., ... 292
Culinary Vinegar, Sauces, &c., ... 293
Pickles, ... 296
TRADE CHEMICALS: Miscellaneous Preparations, Processes, and Compounds used in the Arts, in Domestic Economy, Chemical Research and Recreation, &c., ... 301
APPENDIX.
Tables of Weights and Measures, ... 455
Comparison of Thermometric Scales, ... 461
Effects of Temperature, ... 462
Specific Gravities, ... 464
Table of Chemical Equivalents, ... 465
Table of Composition, Equivalent number, and Symbols of some of the more important Compounds employed in Pharmacy and the Arts, ... 467
Table of Neutralizing Proportions, &c., ... 470
Table of the Relation between the principal Areometers for liquids lighter than water, ... 471
Specific Gravities corresponding with the degrees of Baumé's Areometer for liquids heavier than water, ... 472
INDEX, ... 473

VETERINARY
MATERIA MEDICA.

A Table of the Properties and Doses of the Principal Medicinal Substances used in Veterinary Practice.

N. B.—Where the doses are given without mentioning the animal intended, it must always be understood to refer to the horse.

ACACIA. See Gum Arabic.

ACETATE OF AMMONIA. Spirit of Mindererus. Diaphoretic and diuretic. It is also regarded as antiseptic. The solution of the B. P. is five times stronger than that of the L. P. Dose, for horses and cattle, from 1 to 2 oz. For smaller animals, from 1 to 2 dr. Externally, in strains, ophthalmia, &c.

ACETATE OF COPPER. See Verdigris and Copper.

ACETATE OF LEAD. Astringent and sedative; in doses of 30 to 40 grains with opium, in internal hemorrhage, chronic diabetes, and diarrhœa; but chiefly used externally, in cooling lotions, eye-waters, ointments, &c. (For Diacetate of Lead, see Goulard's Extract.) As antidotes for an overdose give Epsom or Glauber's Salts, with opiates if required.

ACETATE OF POTASH. Diuretic and cooling; dose for horse and cattle, 2 oz. In much larger doses it is a laxative, but not to be depended on.

ACETATE OF SODA. Similar in properties and uses to Acetate of Potash.

ACETIC ACID. Strong acetic (or pyroligneous) acid acts as a rubefacient and caustic, but is rarely employed for this purpose. In the weaker forms of common or distilled vinegar, or diluted wood vinegar, it is frequently used. See Vinegar.

ACIDS. See Muriatic Acid, Nitric Acid, Prussic Acid, Sulphuric Acid, &c.

ACUPUNCTURATION. Used in some spasmodic and paralytic affections.

ÆGYPTIACUM. A preparation of verdigris and honey. A mild caustic, used as a local application to ulcers of the mouth, running thrush, grease, &c. Internally poisonous.

ÆTHER. See Ether.

ALCOHOL. Poisonous to all animals—2 drachms will kill a dog. See Spirits, Ardent.

ALLSPICE. *Pimento.* A useful stimulant and carminative; used in cordial balls and drinks, and to correct the action of purgatives. Dose for horses, 2 to 4 dr.; cattle ½ oz. to 1 oz. Dose of the Tincture, 4 ounces, in gripes.

ALOES. The most valuable purgative for the horse, but not to be depended on for cattle and sheep. A horse requires from 4 to 8 dr. of Barbadoes aloes, from 5 to 9 dr. of Socotrine, and from 6 to 10 of Cape. Mr. YOUATT says 3 dr. of Barbadoes are equal to 4 of Cape. Mr. MORTON considers a mixture of equal parts of Cape and Barbadoes aloes to be quite as efficacious as the latter alone. But the fine gourd Barbadoes aloes are the most certain in their operation. If the animal be prepared by previous mashes, 5 dr. are generally, and 6 dr. almost always, sufficient. Mr. BLAINE recommends 2 dr. every 6 hours till 8 dr. have been taken, as a nauseant and purgative: but Mr. YOUATT strongly disapproves of this plan, particularly in inflammation of the lungs. Aloes require from 18 to 36 hours to produce their effect, during which time the horse should not be ridden far or fast. Though not to be depended on for cattle, 4 to 6 dr. are sometimes added to the purgative salts. Large doses (in some cases sufficient to destroy life) have been given to sheep without purging. Small dogs require from 15 to 30 gr.; medium-sized ones, a dr.; some larger ones require 2 dr., or more. Hogs can bear but a few grains.

Externally, in the form of tincture, aloes is used as a stimulating application to wounds.

ALUM. Astringent and styptic. Given in doses of 2 to 4 dr. to horses in diabetes and diarrhœa; but BOURGELAT says that its too frequent use induces a phthisical condition. A dose of alum whey, consisting of 2 dr. of the powder in a pint of hot milk, may be given after excessive purging. Cattle require from 2 to 6 or 8 drachms in diabetes and red water; and from 2 to 4 oz. are given to cows, to dry their milk. To calves and lambs it is given in dr. doses, in warm milk, for diarrhœa, &c. Dogs, 10 to 15 gr. Externally it is applied to cracked and greasy heels, joint wounds, sore mouths, inflammation of the eye, chronic discharges from the nostrils, and to arrest bleeding from wounds. *Burnt* alum is more powerful, and is used as a mild caustic, mixed with honey, to fungous growths, sore mouths, &c.

ALTERATIVES. Medicines which, without producing any considerable or immediate sensible operation, and without interfering with food or work, effect a slow change in the diseased action of certain parts, so as gradually to restore a healthy state.

AMMONIA, CARBONATE OR SESQUICARBONATE OF. *Volatile Salts.* Stimulant and antacid. Dose, 1 dr. to 2 dr. [MOIROUD says from 2 to 8 dr.] to horses in tympanitis, and the last stage of pneumonia. To cattle, in hoven (distension from the fermentation of green food), 1 to 4 dr. [MOIROUD says to 12 dr.] The solution of carbonate of ammonia has the same properties as the Spirit of Hartshorn, which see.

AMMONIA, AROMATIC SPIRIT OF. Properties as the last. Dose, ½ oz. to 1 oz.; or to cattle in hoven, 2 to 4 oz.

AMMONIA, LIQUID. Water of ammonia is more pungent and stimulant than the carbonate, and is used for the same purposes, particularly in tympanitis and hoven, largely diluted with water or some aromatic infusion; but it is chiefly used externally in stimulating liniments: also both internally and outwardly as an antidote to the bite of vipers. The dose of common water of ammonia may be from 2 to 6 dr.; or for cattle to 2 ounces. For small animals from ½ dr. to 1½. The vapor from the liquid ammonia (applied by holding an

open bottle containing it to the eye) is used for the relief of amaurosis and other chronic affections of the eye.

AMMONIA, ACETATE OF. See Acetate of Ammonia.

AMMONIA, MURIATE OF. See Muriate of Ammonia.

ANALEPTICS. Medicines or food which restore exhausted strength.

ANGELICA. The root, in powder or infusion, is a warm tonic. Dose, ½ oz. to 2 oz.

ANISEED. This warm seed is used as a cordial, carminative, and pectoral. Dose for the horse, ½ oz. to 1 oz.; or ½ dr. of the essential oil. The latter is often added to purgatives to prevent griping. Cattle take 1 to 2 oz. of the powdered seeds. The oil is said to be poisonous to pigeons.

ANODYNES. Medicines which alleviate pain. Opium is chiefly employed for this purpose.

ANTIMONIALS. The preparations of antimony (besides their effect in producing vomiting in carnivorous animals) are considered to have a special action on the skin and lungs. They are also termed resolvent, and purifiers of the blood, and are supposed to be useful in visceral and glandular obstructions, farcy, &c. Mr. BLAINE says "they lessen arterial action without operating very sensibly either in nauseating the stomach or greatly relaxing the skin." Some writers attribute diuretic effects to them. They are also said to promote *condition*. Pigs are supposed to fatten under their use. The principal preparations employed in veterinary practice are mentioned below.

ANTIMONY, CRUDE. *Black (or sesqui-) Sulphuret of Antimony.* Diaphoretic and alterative. The levigated (prepared antimony) is to be preferred. Given to horses in doses of from 2 to 6 dr., with nitre and sulphur, in surfeit, hide-bound, and other skin diseases; and to improve the coat. Mr. YOUATT says the dose should not exceed 4 dr. For cattle, the dose is sometimes increased to 2 or 3 oz. Dogs take from 10 to 30 grains. Hogs, a drachm or more daily.

ANTIMONY, LIVER OF (Hepar Antimonii), and Crocus of Antimony (Crocus Metallorum), are occasionally used in veterinary practice; but are uncertain in their composition and action. Doses, 1 to 2 drs. Mr. CLARK says these com-

pounds, and the glass of antimony, derange the stomach, but that it is doubtful if they have any other effect.

ANTIMONY, CALIX OF. Diaphoretic. Dose, 2 to 4 dr.

ANTIMONIAL POWDER. Similar to James's powder. Diaphoretic. In colds, fevers, inflammations, &c. Dose, for horses or cattle, 1 or 2 dr.; swine, 6 grains; dogs, from 2 to 5 grains. Less efficient than Emetic Tartar.

ANTIMONY, PRECIPITATED SULPHURET OF. Dose, ½ dr. to 2 dr. in obstinate skin diseases.

ANTIMONY, TARTARIZED. *Emetic Tartar.* Diaphoretic, expectorant, and reduces arterial action. It is also regarded as diuretic and febrifuge. Dose, ½ dr. to 1½ dr. in gruel, 3 times a day, in fevers, in inflammation of the lungs, and catarrhal affections. To destroy worms, 2 dr. may be given with powdered tin, or some other mechanical vermifuge, fasting, and followed by aloes; or 1 dr. for 6 mornings, followed on the 7th day by a dose of physic. Mr. WHITE says he has not seen any good effect from it as a vermifuge. Cattle require from ½ dr. to 1 dr. Sheep from 10 to 20 gr. To swine and dogs it is emetic: the former require from 2 to 5 gr.; the latter, from 1 to 3 gr. Externally it produces an eruption on the skin. Formed into an ointment with lard, it has been rubbed on externally in chest affections, but is dangerously irritant.

ANTIMONY, BUTTER OF. *Muriate or Chloride of Antimony.* Used externally only, as a caustic in canker, &c.

ANTIPERIODICS. Remedies against those diseases which return at regular intervals, as agues.

ANTISEPTICS. Remedies which counteract putrefaction.

ANTISPASMODICS. Medicines which relieve spasm, as opium, ether, camphor, ammonia, ardent spirits, &c.

ARNICA. Nervine, sedative, and diaphoretic. 40 to 60 gr. of the powdered plant (the flowers in preference) have been given twice a day for paralysis, amaurosis, rheumatism, blows or falls, &c. A decoction may be used outwardly as a fomentation to bruises, wounds, &c.

ARISTOLOCHIA. See Birthwort.

ARSENIC. *White Arsenic, or Arsenious Acid.* Very poisonous to all animals. In small doses, tonic and alterative,—

but its operation requires to be carefully watched. It has been given, in doses of 2 gr., gradually increased to 20, in farcy and glanders. Externally, is a caustic, but dangerous and unmanageable. Used in solution to destroy vermin in cattle and sheep; but it is not free from danger. Mr. YOUATT remarks—"We have better and safer tonics, and better and safer caustics." The best antidotes are, hydrated oxide of iron, and calcined magnesia, in very large quantities, or a mixture of lime-water and linseed oil.

ASSAFŒTIDA. Stimulant, antispasmodic, and expectorant. It is prescribed in nervous affections and chronic coughs; also in farcy and worms; and to increase the appetite and digestion. The dose is ½ dr. to 2 dr.; but, according to MOIROUD, may be carried to 2 oz. for the horse, and 2 or 3 oz. for horned cattle. Externally it is applied to indolent tumors, &c.

ASTRINGENTS. Medicines which produce a more obvious and decided constriction of the muscular fibres than the simple tonics.

BALSAMS. Natural balsams appear to act on the mucous membrane generally; but are chiefly given as diuretics and expectorants. See Balsam of Canada, Copaiva, Peru, &c.

BALSAM OF CANADA. Diuretic. Dose, ½ oz. to 1 oz.

BALSAM OF COPAIVA. As a diuretic to horses, ½ oz. to 1 oz.; as an expectorant in chronic coughs, 1 or 2 dr. For dogs, ½ dr. to 1 dr.

BALSAM, FRYAR'S. Comp. Tincture of Benzoin. It is sometimes given in ½ oz. doses to horses, in chronic cough, mixed with yolk of egg, gruel, or linseed tea. But more frequently applied to wounds, indolent ulcers, &c.

BALSAM OF LUCATELLI. Dose, ½ oz. in old coughs.

BALSAM OF TOLU, AND OF PERU. 2 dr. in old coughs; but too expensive.

BALSAM OF SULPHUR. A stimulating expectorant in old coughs, in doses of ½ oz. to 1 oz. Sometimes used as an outward application.

BARBADOES TAR. Stimulant, diuretic, and expectorant. Dose, 1 to 4 dr. or more [2 to 4 ounces—MORTON], in old coughs,

and chronic chest affections, Externally in skin diseases, wounds, grease, &c.

BARK, PERUVIAN. Tonic, astringent, antiseptic, and antiperiodic. Dose for a horse, 6 or 8 dr. [to 2 or 3 oz.—MOIROUD] in diabetes, general weakness, a tendency to gangrene, &c. To small animals, 1 or 2 dr. Applied also to indolent and foul ulcers.

BARLEY. The decoction (of Scotch or pearled barley in preference) is given as an emollient, demulcent, or diluent drink in inflammatory diseases; more frequently as a vehicle for more active remedies.

BARYTES. All its compounds are poisonous. The following doses have been given in farcy and glanders: Muriate of barytes, 20 gr. gradually increased to 60; pure barytes, 10 to 20 gr.; carbonate 1 to 4 gr. A dog was killed by 15 grains.

BASILICON, YELLOW AND BLACK. *Resin Cerate.* See V. Formulary (Digestive Ointments).

BAY BERRIES. Stomachic and carminative. An ingredient in diapente, but rarely given alone. Dose of the powdered berries, ½ oz., or of the oil of bays, ½ dr. to a dr. The leaves are used in fomentations.

BELLADONNA. *Deadly Nightshade.* Narcotic and sedative. Dose, of the extract from 1 to 4 dr. in diseases where there is undue action of the nervous and vascular systems [MAYOR]. M. MOIROUD directs from 6 to 8 dr. of the powder. For dogs, from 2 to 8 gr. of the powder. The extract was also applied to the eye, to dilate the pupil.

BENNET, HERB. *Avens.* Tonic and astringent. Dose, of the powdered root, ½ oz. to 1 oz. or more.

BENZOIN. Stimulant and expectorant. Dose, 1 to 3 dr. But seldom used. Externally it is applied, in balsamic tinctures, to wounds, ulcers, &c. See Tincture of Benzoin.

BIRTHWORT. A gentle stimulant, supposed to act especially on the uterine system. Dose, ½ oz. It is given to cows in cleansing drinks, but it is of doubtful utility.

BISTORT ROOT. Astringent. Dose, 4 to 8 dr. [or 2 oz., MOIROUD]. The decoction is used also as an astringent and cleansing lotion.

Bitter-Sweet. *Dulcamara.* Diuretic, narcotic, and alterative. Dose, ½ oz. in decoction.

Bitter Apple. See Colocynth.

Bleeding. The quantity of blood usually abstracted from the horse is from 2 to 4, or, in some cases, 6 to 8 quarts; or until faintness is produced. From cattle, from 2 to 6 quarts, or till faint. Sheep, 16 ounces. Lambs, 4 oz. Dogs in the proportion of 1 oz. for every 3 lb. weight. [Or 1 or 2 oz. from a very small dog; 7 or 8 oz. from a larger one.—Mr. Youatt].

Blistering Fly. See Cantharides. Blisters are applied in the form of ointments, or liniments, to excite superficial inflammation, followed by vesication; and are intended to draw away inflammatory action from more deeply seated but not distant parts. Also to excite the action of the absorbents, and to promote suppuration. See Blistering Ointment, and Liquid Blister, in the Formulary.

Blue Vitriol. See Copper, Sulphate of.

Bole, Armenian. Slightly astringent, and absorbent. Dose, ½ oz. to 2 oz. in diarrhœa, bloody urine, &c. A common ingredient in drenches to dry the milk of cows; dose, 1 to 3 oz. It is also used outwardly as an astringent and desiccative.

Borage. A decoction of the plant is pectoral and demulcent.

Borax. Detergent. Applied to sore mouths, mixed with honey. It is supposed to be a uterine stimulant, but is not often used in veterinary practice as an internal remedy.

Box Leaves. They are given, chopped with corn, as a vermifuge. They are also used as a preventive of hydrophobia. (See the V. Formulary.) The rasped wood is considered sudorific, and prescribed in rheumatic and skin diseases, and even in farcy and glanders.

Bran. Mucilaginous, and slightly laxative: given in mashes.

Brandy. See Spirits, Ardent.

Briony. White Briony root is poisonous. ½ oz. killed a dog.

Bromine. Poisonous. 5 gr. killed a dog. Its medical use is not well ascertained, but appears analogous to that of iodine.

Broom. The Spanish broom, and particularly the seeds, are supposed to produce inflammation of the bladder in sheep and cattle.

Buckbean. A bitter tonic and purgative. The powdered plant has been given to sheep for rot, in 1 dr. doses.—[Dr. Paris.]

Buckwheat. Slightly laxative, but chiefly used to fatten poultry.

Buckthorn. Purgative; principally administered to dogs. Dose of the juice, 2 or 3 dr.; but it is usually given in the form of Syrup. See Syrup of Buckthorn. The berries are more active, but seldom employed.

Burdock. Diuretic and sudorific. Used, but rarely, in rheumatism and skin diseases.

Burgundy Pitch. Similar to resin in its properties. It is chiefly used outwardly in charges, &c.

Butter of Antimony. *Chloride of Antimony.* See Antimony, Muriate of.

Cabbage Tree Bark. Vermifuge. Dose for a horse, 2 to 4 dr. But rarely used.

Calamine, or Lapis Calaminaris. Slightly astringent, drying and healing. Sprinkled on excoriations and sores; and used in ointments, lotions, eye-waters, &c. The greater part of what is sold is factitious, and only calculated to do harm.

Calamus Aromaticus. *Sweet flag.* A warm stomachic. Dose, from 1 oz. to 2 or 3 oz. in infusion.

Calomel. Alterative, vermifuge, sialogogue, purgative; it also increases the action of diuretics and diaphoretics. In doses of 1 or 2 dr. [20 to 60 grains, Youatt], combined with or followed by aloes, it is given to horses for worms; or from 10 to 20 gr. as an alterative, in skin diseases, grease, farcy, constitutional affections, &c. If too often repeated, it salivates. It does not agree with cattle (see Mercury), but is sometimes given, in doses of from 10 to 20 gr., in inflammation of the liver, and jaundice. Some writers mention much larger doses. On dogs it acts as

a purgative, and often as an emetic, and it is very apt to salivate. The same applies to swine. Dose for dogs, 1 to 2 gr. [Never exceeding 3. YOUATT.] Many dogs are destroyed by calomel. Hogs require 3 to 5 gr. Poultry should not have more than a grain, in divided doses, in the day.

CALUMBA. Tonic. Dose of the powdered root, from 2 to 4 drachms.

CAMPHOR is reputed antispasmodic, narcotic, and diuretic. It assists the action of diaphoretics; is frequently added to fever medicines to allay irritation; and is used as an antiseptic in malignant epidemics, &c. Mr. SPOONER combines it with opium in cases of lock-jaw. Dose, 1 or 2 dr. MOIROUD says 2 to 12 dr. Its use is questionable where active inflammation exists. Externally it is used as a discutient and anodyne, in embrocations, eye-waters, &c. Its vapors are thought to act favorably on old coughs.

CANELLA BARK. A warm tonic. Dose, for horses, 2 to 4 dr.; for cattle, 2 to 6 dr.

CANTHARIDES. Stimulant and diuretic. Mr. VINES says, "Of all medicines given for farcy and glanders none equal cantharides;" but they should not be given too early, nor without due caution. [Mr. BLAINE.] Dose, in debility, 3 to 5 gr.; in dropsy, farcy, and glanders, 5 to 8 gr. daily, gradually increasing the dose to 15 gr., suspending their use for a time when their diuretic effect is manifest. Of the tincture, 2 to 3 dr. in incontinence of urine; and from ½ oz. to 2 oz. in red water. The practice of giving cantharides as a venereal stimulant is reprobated by the best authorities. Externally it is used in blistering and stimulating ointments and liniments. It does not permanently blemish, but this effect is often produced by other ingredients combined with it in blistering ointments.

CAPSICUM. Cayenne pepper. A hot stimulant. From 10 to 20 grs. may be given in weakness of the stomach, and from 20 to 60 grs. in flatulent colic, or in severe colds. It is also used externally as a stimulant.

CARBONATE OF AMMONIA; Carbonate of potash; and carbonate of soda. See Ammonia, Carbonate; Potash, Subcarbonate; Soda, Subcarbonate.

CARAWAY SEEDS. Carminative and Stomachic. Dose, ½ oz. to 1 oz.; or double that quantity to cattle. Used in cordial balls and drenches; and often added to purgatives, to prevent griping. The essential oil is used for the same purposes, in doses of 10 to 30 drops. Mr. YOUATT considers caraway and ginger the only cordials required for the horse.

CARDAMOM SEEDS. Carminative. Dose, 1 to 4 dr.

CARMINATIVES are stimulants which by their rapid impression on the stomach, &c., occasion the expulsion of wind, and relief of pain.

CARROTS. Restorative and alterative. Given to horses as food after severe illness: and in coughs, grease, foul humors, &c. Externally in poultices.

CASCARILLA. A warm, bitter tonic. Dose, 2 or 3 dr.

CASSIA. A warm stimulant. Dose, 1 to 2 dr.

CASTOR. Antispasmodic. ½ oz. has been given in locked jaw. But rarely used.

CASTOR OIL. Laxative. It is uncertain as a purgative for the horse, and sometimes produces much irritation in large doses. ½ pint may be given every six hours till it operates, with watery solution of aloes. Cattle require a pound, or pint; calves, 2 to 4 oz.; sheep and swine, 1 to 2 oz.; dogs, 2 to 4 dr., with syrup of buckthorn. The seeds are more active: from 2 to 6 are sometimes given to swine and dogs, crushed and mixed with food; but from their effects on man, their use would seem to require caution.

CATECHU. *Terra Japonica.* Astringent. Dose for a horse, in diabetes, diarrhœa, &c., 1 or 2 dr. [YOUATT], or to 1 oz. [BLAINE]; cattle, 2 to 4 dr. in gruel. [It is usually combined with chalk, opium, and gum.—YOUATT.] Dogs require from 10 to 40 gr. In India it is said to be given in doses of 2 oz., for the purpose of taming vicious horses. The tincture is useful in promoting the healing of wounds.

CATHARTICS. Purgatives (which see).

CAUSTICS. Solid or liquid substances which burn or destroy the part to which they are applied. The *actual cautery* consists in burning with an iron heated to whiteness.

CHALK. Antacid and astringent. Horses require from ½ oz. to 1 oz.; cattle, 1 or 2 oz.; sheep and swine, 1 dr.; dogs, 10 to 20 gr. It is often combined with catechu. Externally it is sprinkled on sores.

CHAMOMILE. A mild tonic, stomachic, and febrifuge. Dose, 1 to 4 dr. of the powdered flowers, or an infusion of ½ oz. of the flowers in a quart of water, in debility of the stomach, flatulence, and in the last stage of fevers, and influenza. It is the first tonic that should be used in convalescence. Ginger or some other aromatic, is usually joined with it.

CHARCOAL. Antiseptic. Used as an application to foul ulcers, either sprinkled on them or mixed with poultices.

CHARGES. Compositions of an adhesive nature, usually mixed with tow, which adhere to the part to which they are applied for some time. See V. Formulary.

CHLORATE OF POTASH. Mr. MORTON states that Mr. SYMONDS found it useful in hoven and tympanitis. Dose, 1 to 2 dr.

CHLORIDE OF LIME. Antiseptic and disinfectant. From 2 to 4 dr. in a quart of water, given to horses in flatulent colic, and to cattle in hoven; and in putrescent diseases. Externally, as a wash for mange, foul ulcers, &c., and as a disinfectant, ½ oz. to be well mixed with a pint of water, and after a time decanted or strained.

CHLORIDE OF POTASH. *Eau de Javelle.* Recommended by French authors for the same purposes as the chlorides of lime and soda. Dose, for hoven or tympanitis, ½ oz. to 1 oz.; for sheep, ¼ oz., in water, with or without the addition of ether.

CHLORIDE OF SODA. Labarraque's Disinfecting Solution. The properties and uses are the same as of chloride of lime; it is perhaps better adapted for internal use. Dose, 2 to 4 dr. of the solution, gradually increased to 1 oz. or more, largely diluted. It has been tried in glanders. As a lotion, about 1 oz. to a pint of water.

CHLORINE. Antiseptic. A strong watery solution of chlorine gas is antiseptic—in large doses poisonous. It is used for the same purposes as the chlorides of lime, potash, and soda, but the latter are preferable.

CHLORIDE OF ANTIMONY. See Antimony, Butter of.

CHLORIDE OF ZINC. It is a powerful caustic. A diluted solution is used as a disinfectant.

CHLOROFORM. Used to produce insensibility to pain in the same manner as ether; and as a remedy for tetanus.

CINCHONA. See Bark, Peruvian.

CINNABAR AND VERMILION. Native and factitious red sulphuret of mercury. Alterative and vermifuge? Dose, ½ oz. daily to horses, in skin diseases and obstinate coughs. Formerly given in larger doses, as a vermifuge. Cinnabar of Antimony, so called from the mode of preparation, does not differ from common vermilion in its properties. Care must be taken to get pure vermilion, as this compound, being used as a pigment, is sometimes adulterated with red-lead and other poisonous matters.

CINNAMON. Stimulant and carminative. Dose, 2 dr.; but cassia is usually substituted for it.

CLOVES. A hot stimulant, cordial and carminative. Dose, 1 to 3 drachms in powder; or from 10 to 20 drops of the oil; the latter is a frequent adjunct to purging balls, to prevent griping. Cloves are also an ingredient in masticatories.

CLYSTERS. These are injected into the rectum by a proper syringe, or a bladder and a pipe, either to unload the bowels, abate inflammation and pain, or to act on the system generally, when medicines cannot be given by the mouth. See V. Formulary.

COLCHICUM. Poisonous to most animals. A diuretic and drastic purgative, chiefly used in rheumatic affections. Dr. LEMANN found it useful in constitutional ophthalmia, and in pneumonia, in doses of a drachm, twice a day, with nitre. According to M. MOIROUD, the dose for larger animals is from 1 to 2 dr. For smaller, 6 or 8 gr.

COLOCYNTH. Bitter apple. It has little effect on the horse. It is purgative to dogs, and in large doses poisonous.

CONFECTION OF OPIUM. Anodyne and carminative. Dose, 4 to 6 dr., in flatulent colic.

CONFECTION OF ROSES. Slightly astringent; but only used to form astringent powders, &c., into balls. Masses formed with it retain their consistence well.

COPAIVA. See Balsam of Copaiva.

COPPER. All the compounds of this metal are poisonous. In small doses they are tonic. The antidotes are white of eggs, milk, iron filings, or hydrated sulphuret of iron.

COPPER, ACETATE OF. Crystallized (commonly called *distilled*) verdigris. Caustic and cleansing. Stronger than common verdigris.

COPPER, DIACETATE OF. See Verdigris.

COPPER, DINIODIDE OF. Tonic, and promotes absorption. Dose, 1 to 2 dr. daily, in farcy, glanders, swelled legs, &c., and topically, to ill-conditioned ulcers.

COPPER, NITRATE OF. Sometimes used as a caustic.

COPPER, SULPHATE OF. *Blue Vitriol.* Tonic and styptic. In doses of ½ dr. gradually increased to 2 dr. or more daily, it is given in diabetes, farcy, &c. Small doses may be given in balls with gentian and ginger; larger doses in gruel. It has been thought useful in glanders; but Mr. YOUATT says it is only proper in nasal discharges without fever. Dose for cattle, 1 to 2 dr. Sheep 20 to 40 gr. Rabbits (in sniffles), 1 or 2 gr. twice a day. Externally the solution is used for the foot-rot of sheep; and as a cleansing wash for foul ulcers in horses and cattle. Used also in the solid state to destroy proud flesh.

COPPER, AMMONIO-SULPHATE OF. Tonic and astringent. Dose, 1 to 2 dr. twice or thrice a day.

CORDIALS. Warm stimulating medicines, such as spices, and the aromatic seeds, fermented liquors and spirits, &c., which temporary restore exhausted strength, revive the spirits, and rouse the system generally. The best modern practitioners condemn their indiscriminate employment, as the source of much mischief. For cordial balls, &c., see V. Formulary.

CORIANDER SEEDS. A mild aromatic stimulant and carminative, used in cordial balls and drinks. Dose, ½ oz. to 1 oz.

CORROSIVE SUBLIMATE. One of the most virulent poisons. In small doses it is alterative and diuretic. It has been tried in doses of 2 to 5 gr., gradually increased to 10 or 20, in farcy and glanders, but rarely with lasting benefit. Externally it is used as a powerful caustic. A dilute

solution is employed as a wash for scab and lice in sheep, but the practice is not free from danger. Applied to wounds in cattle it has proved as fatal a poison as when swallowed. The antidote for an overdose is white of egg, or milk, or the hydrated sulphuret of iron; with demulcent drinks.

COWHAGE. Vermifuge; but has little effect on the horse.

CREAM OF TARTAR. Cooling, laxative, and diuretic. Seldom given alone; but combined with antimonials, mercurials, or sulphur, as an alterative in skin diseases; and used as an adjunct to aloes in purging balls. Cattle require 2 to 3 oz.; when given in larger doses it should be given in plenty of warm water. Sheep require ½ oz. to 1 oz. Dogs, 5 to 20 grains.

CREASOTE. Tonic, stimulant, and antiseptic. Dose, 20 to 30 drops daily, in gruel or linseed tea, in glanders. Externally in lotions and ointments, to fistulous wounds, unhealthy ulcers, &c.

CROCUS OF ANTIMONY. See Antimony, Liver of.

CROTON SEEDS AND OIL. Purgative. The oil produces great irritation in the horse. Dose, about 20 drops: 30 drops have proved fatal. The powdered seeds and the meal or ground cake left after expressing the oil, are also used; 3 gr. of the former and 5 of the latter being considered equivalent to 1 dr. of aloes. It operates with less certainty, and produces more debility, than aloes, but is sometimes preferred on account of its more speedy action. It is usually given in the form of a ball, 20 or 30 gr. being mixed with 1 oz. linseed meal. Mr. NORTON gives from 12 to 24 grains of the seed. Mr. YOUATT prescribes 30 grains of the powdered seeds in a drink, in tetanus and brain fever, followed by smaller doses (10 gr.) every 6 hours. It will purge rapidly when placed upon the tongue, but is then likely to inflame the mouth. From 10 to 20 gr. are sometimes added to salts in purging drenches for cattle, in extreme cases. One drop of the oil purges a dog freely.

CUMMIN SEED. A warm carminative. Dose, from 1 to 4 drachms of the powdered seeds; or from 6 to 20 drops of the oil.

CUSPARIA, OR ANGUSTURA BARK. An aromatic bitter tonic. Dose, 1 to 4 dr. in debility, diabetes, diarrhœa, &c.

CYANIDE OF POTASSIUM. It possesses the same poisonous and medical properties as prussic acid. Mr. LAFORE has given it with success in a case of idiopathic tetanus of the horse; but it failed to cure traumatic tetanus. Dose, 4 grains.

DAFFY'S ELIXIR. Sometimes given in colic or gripes.

DALBY'S CARMINATIVE. Given to calves in diarrhœa. Dose, ½ a bottle.

DETERGENTS OR DETERSIVES. Remedies which cleanse foul ulcers.

DIAPENTE. A compound powder, reputed cordial and stomachic. Too much of what is sold in the shops is almost worthless. Dose, ½ oz. to 1 oz.

DIAPHORETICS. Medicines which promote perspiration.

DIGESTIVES. Mildly stimulating applications, which excite healthy action in indolent ulcers, wounds, &c.

DIGITALIS. *Foxglove.* Sedative and diuretic. It reduces the frequency of the pulse and diminishes irritability. It is poisonous to animals generally; 6 gr. will kill a dog. It is asserted, however, that it produces no effect on poultry. The common dose of the powdered leaves for a horse, is from 10 to 30 gr. Mr. YOUATT prescribes 60 gr., with emetic tartar and nitre, in inflammation of the chest; but its effect on the pulse must be carefully watched. To cattle, ½ dr. to 1 dr. Sheep, 5 to 15 gr. Dogs, 1 to 2 gr. An infusion of the leaves is applied to inflamed eyes.

DIURETICS. Medicines which increase the flow of urine. Some of them, juniper, capivi, squills, broom, &c., appear to carry off water only; while the alkaline salts remove solid matters also, and thus purify the blood. Diuretics are employed to lessen the quantity of the circulating fluid in fevers and inflammations. The legs of many horses cannot be kept fine, nor the grease be subdued without the use of diuretics. Plenty of water should be allowed with them. But their too frequent use is injurious.

DOG-GRASS. It is emetic to dogs.

DOVER'S POWDER. Sudorific to cattle in rheumatism. Dose, 1 drachm.

EGGS. Nutritive and demulcent. Sometimes given in diarrhœa. They constitute the best antidote to poisoning by corrosive sublimate.

ELATERIUM. It has little effect on the horse.

ELDER. An infusion of the flowers is given in catarrhal complaints. The leaves boiled with lard form an emollient ointment, which is a common application to sore udders. The fresh leaves of the dwarf elder are given (according to BOURGELAT and MOIROUD) with some success as a deobstruent and aperient, in swelled legs, dropsy, and farcy.

ELECAMPANE. The root is reputed stimulant, diaphoretic, diuretic, stomachic, and expectorant. Dose, 4 to 8 dr. in chronic catarrh, dropsical swellings, indigestion, &c.

EMETIC TARTAR. See Antimony, Tartarized.

EMETICS. Medicines which excite vomiting. It is scarcely possible to produce this effect in herbivorous animals.

EMOLLIENTS. Medicines which soften and relax the tissues of the organs.

EPSOM SALT. A cooling laxative. It is not to be depended on as a purgative for the horse; but in doses of 4 or 5 oz., in a large quantity of water, repeated three times a day, it is useful as a laxative and diuretic in inflammatory diseases. Cattle require from 12 to 20 oz., with ginger or any of the warm seeds. It is sometimes rendered more active by aloes or gamboge. Calves require from 1 to 2 oz., according to their age and strength. Sheep, ½ oz. to 2 oz. Dogs, from 1 to 3 dr. wrapped in tissue paper. A large elephant takes a pound and a half, preceded by a dr. of calomel.—YOUATT.

ERGOT OF RYE. It promotes parturition. Dose for a mare, 2 or 3 dr. A cow, 2 dr. repeated at intervals of half an hour. A ewe, 20 to 40 gr. Bitch, 5 to 10 gr. [Mr. SPOONER says from 2 to 4 gr.], or an infusion of a scruple given at three times, at intervals of half an hour. Larger doses than the above are indicated by M. MOIROUD.

ERRHINES. Remedies which excite a discharge from the nostrils.

ESCHAROTICS. Caustics. Substances which destroy the surface to which they are applied.

ETHER. A diffusible stimulant and antispasmodic; used chiefly in colic. Dose, ½ oz. to ¾ oz.; cattle, ½ oz. to 1 oz.; dogs, 7 to 14 drops. It is used outwardly in cooling lotions and eye-waters. The vapor, inhaled by means of a proper apparatus, produces insensibility to pain; but some of the experiments with this agent have proved most unfortunate.

ETHIOPS MINERAL. The mildest of the mercurial compounds. Alterative and vermifuge. Dose, 2 to 4 dr. daily in farcy, glanders, grease, skin diseases, and worms, alone, or with cream of tartar. For cattle, 1 dr.; swine, 3 to 10 gr.; dogs, 5 gr. in mange. With an equal weight of prepared antimony it forms Antimonial Ethiops—a more efficient preparation.

EUPHORBIUM. Very acrid and poisonous. Used in blisters, chiefly to economize the more expensive flies; but irritates extremely. It is employed in the form of tincture and ointment as a local stimulant.

EXCITANTS. Medicines which quicken the circulation, produce warmth, and render the organs more active.

FENNEL SEEDS. A weak carminative and diuretic. Dose, ½ oz. to 2 oz.

FERN. Powdered male fern is given in doses of 6 dr., followed by a mercurial purgative, for expelling worms. M. MOIROUD carries the dose to 4 oz.; or 5 or 6 dr. for smaller animals.

FŒNUGREEK SEEDS. Emollient, nutritive, and stomachic. Dose, 1 oz. daily, to promote condition in horses, and in diseases of the chest. It is also added to the food of swine, to promote their fattening. Used also externally in fomentations.

FORGE WATER. The water of the blacksmith's forge is sometimes given as a tonic, or applied as a wash to ulcerated and cankered mouth.

FOXGLOVE. See Digitalis.

GALANGAL ROOT. A warm aromatic; similar in properties to ginger. Dose, ½ oz. of the powder, or 1 oz. in infusion.

GALBANUM. Stimulant, expectorant, and antispasmodic. Dose, 2 to 4 dr. But rarely used, assafœtida being stronger and cheaper.

GALL NUTS. Astringent; in diarrhœa. Dose of the powder for horses and cattle, 2 to 4 dr. Calves, ½ dr. to 1 dr. Dogs, 4 to 8 grains.

GAMBOGE. A drastic purgative. The dose for a horse is said to be from 2 to 6 dr., but its purgative effect cannot be depended on, and it gripes. 3 dr. have been known to cause great prostration; and the horse being killed, marks of intense inflammation were found in its stomach and bowels. It is a bad medicine for herbivorous animals. 2 dr. are sometimes added to salts and other purgatives for cattle. Sheep have been killed by 2 dr. A few grains are given to dogs, to destroy and expel worms.

GARLIC. A stimulating expectorant. Dose, 1 oz. in chronic coughs and asthmatic complaints, made into balls with liquorice powder; or boiled in milk. It is a common remedy for coughs and chest affections in all domestic animals. It is also reputed vermifuge. For the croup in fowls it is given in doses of 5 gr.

GENTIAN ROOT. Tonic and stomachic; in debility, after severe illness, &c. Dose for a horse, 2, 3, or 4 dr. of the powder; or from ½ dr. to 1 dr. of the extract. (See V. Formulary, Tonic Balls.) Cattle, 2 to 4 dr. or more. Sheep, 20 to 60 gr. Generally joined with ginger. An infusion is recommended as a wash to ulcers.

GINGER. Stimulant and carminative; a general ingredient in cordial and tonic medicines. Dose, 1 to 3 dr., or in flatulent colic, 2 to 6 dr. Cattle, 2 to 6 dr. Calves, 20 to 30 gr. Sheep, 30 to 60 gr. The smaller of the above doses may be added to all aperient medicines. It is also used as a masticatory. Dose of the tincture, ½ ounce to 2 ounces.

GLASS, POWDERED. Used to destroy worms in dogs. Mr. BLAINE recommends as much as will lie on a sixpence, with butter.

GLAUBER'S SALT. *Sulphate of Soda.* Aperient and diuretic. Seldom given to horses as a purgative (Mr. CLARKE says 1 lb. produces scarcely any effect); but it is said to be useful in doses of 6 oz. 3 times a day, in epidemic catarrh. To cattle the usual dose is 16 oz., or from 12 to 20 oz., with ginger or caraway. It is considered more diuretic than Epsom salts.

GOULARD'S EXTRACT OF LEAD. *Solution of Diacetate of Lead.* Cooling and astringent. Used externally only, in lotions, &c., in the same cases as sugar of lead. (See Lead, Acetate of.) For inflamed eyes, 1 dr. or 1½ to a pint of water; for other purposes it is made stronger.

GRAINS OF PARADISE. A warm stimulant; chiefly used in cattle medicines. Dose, 3 to 6 dr.

GRUEL. A decoction of oatmeal. Nutritive and demulcent.

GUAIACUM [GUM]. Sudorific and expectorant. It has been given to horses, in doses of 4 dr., in chest affections, farcy, rheumatism, &c.; and to cattle in doses of 4 or 6 dr. But its utility is doubted. The guaiacum wood is given to the amount of 4 oz., in decoction, repeated 2 or 3 times in 24 hours.

GUM ARABIC. Emollient and demulcent. Used in inflammatory affections of the bowels, or of the respiratory or urinary organs. Dose, for horses and cattle, 1 to 4 oz., dissolved in water. For smaller animals, from ¼ oz. to 1 oz. It is also used to suspend in water insoluble powders and oils. Gum senegal and gum tragacanth are used for the same purposes. The latter will thicken twenty times as much water as Gum Arabic. [For Gum Ammoniac, Benzoin, &c., see Ammoniacum, Benzoin, &c.]

HARTSHORN, SPIRITS OF. See Ammonia. It is chiefly used in stimulating liniments, and for the bites and stings of venomous reptiles and insects. For salt of hartshorn, see Ammonia, Carbonate of.

HELLEBORE, WHITE. Poisonous to all classes of animals. In small doses, it has been strongly recommended as a nauseant and diaphoretic, in inflammatory diseases; but it requires to be very carefully watched, otherwise a fatal collapse may be induced. The usual dose is 20 gr. every four or six hours till nausea is produced, or the pulse affected. Mr. YOUATT says it cannot safely be given in doses of a drachm, but that it is given with advantage in ounce doses in chronic grease. Externally, it is used in ointments and washes for the mange; but even in this way its use requires caution. It is also blown into the nostrils as a sternutatory.

HELLEBORE, BLACK. The root is used as an irritating seton for cattle, and introduced into fistulous sores of the horse.

HEMLOCK, SPOTTED. A narcotic poison. In doses of a drachm of the powdered leaves, or the extract, gradually increased, it is sometimes given to quiet obstinate coughs. It is also an ingredient in some old remedies for farcy, scirrhous tumors and cancer. For dogs, from 1 to 4 gr., in coughs and cancerous diseases. A decoction of the herb is used as a fomentation to painful tumors. Water hemlock is a more virulent poison, and often destroys cattle. M. MOIROUD says that the ruminants bear hemlock better than other animals. Mr. YOUATT considers both common and water hemlock harmless to the horse, though he admits that cows have been poisoned by the latter.

HENBANE. Narcotic and sedative. Dose, 15 to 20 gr. of the powder [1 to 2 dr. of the extract, MORTON] twice or three times a day, to allay arterial action. On dogs it acts as on man: dose 3 to 5 gr. German horsedealers are said to give a plump appearance to diseased horses by mixing henbane seeds with their corn.

HONEY. Demulcent, emollient, and slightly laxative. Used in cough medicines, and to make up balls. Horses are fond of it. Externally, it is detergent, and is, perhaps, useful in defending ulcers from the air.

HOPS. Tonic and slightly anodyne, but chiefly used in fomentations.

HOREHOUND. Sometimes given in coughs; a quart of the decoction, or 1 oz. of the powder.

HORSERADISH. Stimulant and diuretic. Said to be useful in dropsical complaints, and in recent epidemics, attended with chronic inflammation. The fresh root is rasped and mixed with barley meal.

IODINE. Alterative, and promotes absorption. Used externally and internally to reduce glandular swellings, and scirrhous and other tumors. 5 gr. of iodine, or 1½ or 2 dr. of the compound tincture, may be given twice a day in farcy. Cattle take from 5 to 10 gr., and from 1 to 2 dr. of the compound tincture. Dogs ¼ to 1 gr. twice daily. The compound iodine ointment is used to disperse glandular enlargements. It is rapidly superseding cantharides.

IODIDE OF IRON. Tonic and alterative, promoting the action of the absorbents. Dose, ½ dr. to 1 dr.

Iodide of Potassium, or Hydriodate of Potash. It possesses the same properties as iodine, but irritates less. It is often combined with iodine, which it renders soluble in water. The dose, by itself, is rather larger than of iodine—from 15 to 30 gr. twice a day; to cattle, 5 to 10 grains.

Ipecacuanha. Little used in veterinary practice, except as a sudorific, in combination with opium (Dover's powder). A drachm or two may be given to horses in asthmatic affections. It purges sheep, purges or vomits the pig and dog. Dose for the latter, 4 to 20 gr. [From 2 to 30 gr.—Moiroud.] 3 oz. killed a horse [Mr. B. Clarke].

Iron. The preparations of this metal are tonic; some of them (as the sulphate and muriate) astringent and styptic. The usual doses for a horse are, 2 oz. of iron filings, once or twice a day, with corn, or in a mash; 1 to 3 or 4 dr. of the sulphate; 2 to 6 dr. of the sesquioxide or carbonate, or of rust of iron, or of the powdered scales; 1 to 3 dr. of tartarized iron; and ½ to 1½ dr. of the iodide. Cattle, 2 to 4 dr. of the sulphate in chronic diarrhœa. For sheep, a sixth or eighth of the above doses. [M. Moiroud prescribes much larger doses than the above.] The muriated tincture of iron is prescribed in doses of 2 or 3 dr. for incontinence of urine. The sulphate is sometimes used externally in astringent lotions.

Jalap. Purgative, but has little or no such effect on the horse or other herbivorous animals. It is sometimes added to other purgatives, but probably without any benefit. Dose for swine, ½ dr. to 2 dr. Dogs, 15 to 40 gr. Cats, 10 to 20 gr.; but it is rather uncertain.

Jamaica Pepper. See Allspice.

James's Powder. Similar to antimonial powder, but considered more certain and uniform in its operation. Dose, 20 to 30 gr. in fever and inflammatory complaints. It is also given to dogs as a remedy for distemper, 4 gr. twice a day.

Jatropha Seeds. The seeds of the physic nut (J. curcas) are given as croton seeds, doubling the dose.

Juniper Berries. Diuretic and slightly stimulant. Dose for a horse, 1 to 2 oz., or 1 to 2 dr. of the essential oil; for cows, 2 or 3 oz.; sheep, ¼ to ½ oz. An extract from the

berries (prepared by evaporating a clear decoction, or rather a cold infusion of the berries, to the consistence of treacle) is much used on the Continent as a vehicle for various remedies.

KERMES MINERAL. A preparation of antimony, similar to the precipitated sulphuret, not much used in this country, but highly esteemed in France. Dose for horses, 1, 2, or 3 dr. For cattle, 4 dr. or more. For a good-sized dog, 1½ gr. gradually increased.

LARD. Half a pound, with warm water, is laxative and emollient. It is also used to make up balls, and is thought to prevent griping, as well as to preserve their consistence. It forms a common basis for ointments.

LAUDANUM. *Tincture of Opium.* See Opium.

LAVENDER. The compound spirit is carminative and cordial. Dose, ½ oz. in peppermint water.

LEAD. The preparations of this metal are poisonous. See Acetate of Lead, and Goulard's Extract of Lead.

LEAD, WHITE AND RED. Common ingredients in ointments and plasters. Also sprinkled on sores as desiccatives. They are likewise used for dusting sheep for the fly.

LIME. Quicklime is sometimes used as a caustic; the powder dusted over foul ulcers, greasy heels, &c.

LIME-WATER. Antacid and tonic. Sometimes given in diabetes, from 2 to 4 quarts. Used also as a wash for sores, and as an injection into the nostrils in glanders and chronic discharges. Mixed with linseed oil, it forms a liniment for burns.

LIME, CHLORIDE OF. See Chloride of Lime.

LINSEED. Demulcent and pectoral. A decoction of the seeds is very mucilaginous, and is used for colds, sore throats, and internal inflammations; also to counteract the effects of corrosive and irritant poisons, and as a vehicle for more active medicines. Lindseed meal is used for poultices. Linseed oil is laxative. Dose for a horse, a pint or a pint and a half; for cattle, 1 or 2 pints; sheep, 2 or 3 oz.

LIQUORICE. Demulcent and pectoral, in coughs, &c. Dose, ½ oz. to 2 oz. of the powdered root; or ½ oz. of the foreign extract (Spanish or Italian juice).

LIVER OF SULPHUR. See Sulphuret of Potassium.

LOBELIA INFLATA. It is poisonous to horses, and produces salivation in cattle; but its remedial powers have not been ascertained.

LOGWOOD. Astringent. 2 or 3 dr. of the extract, or a decoction of 3 or 4 oz. of the wood, may be given in diarrhœa, &c.

LOTIONS. Washes. Liquid applications, with which external parts are bathed.

MADDER. Formerly supposed to be useful in glanders and farcy, and as a preventive of the effects of the bites of venomous reptiles; but it is nearly discarded from modern practice. It is sometimes given to pigs, but with what specific intention it is difficult to say. It colors the bones of animals fed with it.

MAGNESIA. Antacid and laxative. From ½ oz. to 3 oz. to horses and cattle, with some warm carminative in flatulent distension. To calves in diarrhœa, ½ oz. Either the common or the calcined magnesia may be used.

MAGNESIA, SULPHATE OF. See Epsom Salt.

MALLOW. Demulcent. A handful of the leaves is boiled in a quart of water. More frequently used as a lavement. The *root* of the *marshmallow* is preferred; a decoction of 2 or 4 ounces is given as a drink in both coughs and internal inflammations, and used as a clyster, and as a fomentation.

MALT. Nutritive, pectoral, and alterative. It is given, in the form of mashes, in chest affections, when no inflammation is present, and in grease, farcy, and mange.

MANNA. Slightly laxative and pectoral. Dose, 2 oz. with honey, or dissolved in water, in inflammatory disease and chronic coughs.

MASHES. See Bran Mash, &c., in V. Formulary.

MERCURY, OR QUICKSILVER. The preparations of this metal are alterative, most of them purgative, and all apt to produce salivation. Dogs may be easily salivated, but graminivorous animals with greater difficulty. The editor of "Clater's Cattle Doctor" says, "Mercury does not seem to agree with herbivorous animals, in any form or in any disease." The preparations in use are indicated below.

MERCURIAL OINTMENT. Applied to callous swellings, enlarged joints, mange, scab in the sheep, &c. The weaker ointment is generally sufficient.

MERCURY WITH CHALK. Alkalized mercury. A mild preparation. Dose, from 1 to 3 dr., in farcy, glanders, &c.

MERCURY, SUBCHLORIDE OF. See Calomel.

MERCURY, BICHLORIDE (or CHLORIDE, B. P.). See Corrosive Sublimate.

MERCURY WITH SULPHUR. Black Sulphuret of Mercury. See Ethiops Mineral.

MERCURY, RED SULPHURET OF. See Cinnabar.

MERCURY, NITRIC OXIDE. See Red Precipitate.

MERCURY, BLACK OXIDE. Dose, 1 to 2 dr. [MORTON.]

MERCURY, AMMONIO-CHLORIDE. See White Precipitate.

MERCURY, NITRATED OINTMENT OF. See Ointment, Citrine.

MERCURY, ACID NITRATE OF. Used as a caustic. See Caustics, V. Formulary.

MILK. Sometimes given in quantities of 1 to 3 quarts, in acute inflammations, coughs, and all internal irritations, especially those occasioned by acrid and corrosive poisons. It is a convenient vehicle for administering medicines to the dog or cat.

MINDERERUS' SPIRIT. See Acetate of Ammonia.

MINT, AND PEPPERMINT. Carminative, cordial, and sudorific. A strong infusion of the plant, or the distilled water, may be given in flatulent colics. Dose, 1 or 2 pints; used chiefly as vehicles for more active remedies. Dose of the oil of peppermint, 20 to 30 drops, or to 60 drops of oil of spearmint. A few drops of the oil are added to purgative medicines, to prevent griping. The other mints have similar properties.

MITHRIDATE. Cordial and anodyne. ½ oz. to 1 oz. may be given in flatulent colic, but would be injurious in inflammation.

MULLEIN. An infusion of the flowers is given as a demulcent for the same purposes as linseed tea. A decoction of the leaves is used in emollient fomentations and cataplasms.

Muriatic (or Hydrochloric) Acid. *Spirit of Salt.* Tonic and antiseptic; but principally used to dissolve calcareous concretions in the bladder. It has been used in the pestilent epidemics of cattle. Dose, for a horse, 1½ to 2 or 3 dr. in plenty of water, twice a day. Externally as a caustic, strongly recommended by Youatt.

Muriate of Ammonia. *Sal Ammoniac.* Formerly used in influenza or epidemic catarrh. It is said also to have proved useful in farcy, and perhaps deserves trial in other chronic diseases. It renders the blood more fluid. Its use requires caution. 2 oz. produced inflammation of the mucous membrane of a horse; 2 dr. killed a dog, and ½ dr. a rabbit. M. Moiroud states the dose to be from 2 to 8 dr. for horses and cattle, and for small animals from a scruple to a drachm, largely diluted. Externally it is a frequent ingredient in discutient lotions to splints, old strains, bruises, indolent tumors, &c., in horses and cattle. It is also employed as an embrocation to sore teats.

Muriate of Antimony. See Antimony, butter of.

Muriate of Barytes. Poisonous; in small doses, alterative. It has been tried in glanders and farcy, with the usual ill success. Dose, ½ dr. in milk.

Muriate of Copper. Used externally only as a mild caustic.

Muriate of Lime. *Chloride of Calcium.* Alterative and resolvent in glandular diseases; but rarely used in veterinary practice. It has been proposed in glanders and farcy. In an overdose it is poisonous. We have not met with any specific statement of doses. 3½ dr. killed a dog.

Muriate of Soda. *Chloride of Sodium.* See Salt, common.

Mustard. Stimulant; but little used as an internal remedy. Flour of Mustard, with or without vinegar, is applied externally as a rubefacient, to relieve internal inflammation.

Myrrh. Tonic, expectorant, antiseptic, and balsamic. From 1 to 3 dr. to a horse, in chronic cough. To cattle, 2 to 4 dr. or more. The tincture is used for ulcers of the mouth in all animals, and to indolent sores.

Naphtha. Rectified wood naphtha is used instead of spirit of wine, for making tincture of myrrh and aloes. This spirit

may be given also in 2 oz. doses internally, forming an admirable stimulant and diaphoretic.

NAPHTHALINE. A stimulating expectorant. It possesses many of the properties of camphor, and a solution of it in spirits may be substituted for camphorated spirit, and with oils, &c., for liniments and ointments. The ointment is substituted for tar ointment.

NARCOTICS. Medicines which induce stupor or sleep, and ease pain.

NAUSEANTS. Medicines which produce nausea, diminish arterial action, and thus abate inflammation.

NETTLE SEED. It is said to be given with the horse's corn, to give a smooth coat, and an appearance of condition and liveliness.

NITRE. *Nitre of Potash.* Cooling and diuretic. In colds, fevers, and inflammatory complaints of the horse, from 2 to 4 dr. may be given daily, in plenty of water, or linseed tea, till the desired effect is produced. An ounce is often given, but smaller doses repeated are better. Cattle, 2 to 4 dr. [1 oz. in 24 hours for some days.—MOIROUD.] Swine and sheep, 30 to 40 gr.; dogs, 4 to 10 gr. A strong solution is applied to gangrenous wounds.

NITRIC ACID, NITROUS ACID, OR AQUAFORTIS. Used externally only, as a strong caustic; or largely diluted (2 dr. to a pint of water) as an antiseptic wash to foul ulcers.

NITRATE OF SILVER. *Lunar Caustic.* Tonic; but rarely given to animals, except to dogs in cholera, in doses of $\frac{1}{8}$ to $\frac{1}{4}$ of a gr. Externally caustic. It is the best caustic that can be applied to the bites of rabid animals. A weak solution (10 gr. to 1 oz. rain-water) is used to excite sluggish wounds, and to remove opacity from the cornea of the eye.

NUX VOMICA. Poisonous to all animals. Given in doses of 8 to 10 gr., gradually increased to 30 gr., in paralysis of the horse; but its effect requires to be carefully watched. It has been tried in glanders and farcy, but without much success. In small doses it invigorates the digestive functions. A few grains will destroy a dog. A drachm has killed a horse. See Strychnine.

NUTGALLS. See Gall-nuts.

NUTMEGS. Stimulant and perhaps narcotic. Sometimes given in colic, but not much in use.

OAK BARK. Astringent and tonic. Dose, ½ oz. to 2 oz. in powder, or boiled in water, for diarrhœa, diabetes, and debility in horses. To cattle, in dysentery, and in red-water (after purgatives), ½ oz. to 1 oz. The powdered bark and the decoction are applied to unhealthy wounds, &c. In France, a mixture of oak bark, gentian, and chamomile, is used as a substitute for Peruvian bark.

OIL, FISH. Common whale oil is a good preventive of the fly, and does not injure the wool.

OIL OF SPIKE. As sold for veterinary purposes, it consists of turpentine, colored, and merely scented with foreign oil of lavender. It is used in warm liniments.

OIL OF TURPENTINE. See Turpentine.

OIL OF TAR. See Tar.

OILS, EXPRESSED. Olive, almond, and linseed oils are laxative, demulcent, and emollient. Dose, 3 to 16 oz., or a pint. In the latter dose they are given (especially linseed oil) as a substitute for castor oil: they are harmless, but rather uncertain in their operation. (See Castor Oil.) They are useful in poisoning by acrid and corrosive poisons. Olive oil is used, both inwardly and outwardly, as a remedy for the bites of reptiles and stings of insects. Externally they are used in liniments and ointments. Oil of bays is gently stimulant and antispasmodic, but chiefly used outwardly.

OILS, ESSENTIAL OR VOLATILE. The essential oil of peppermint, cloves, aniseed, caraway, &c., possess in a concentrated state the carminative properties of the drugs from which they are distilled. They are frequently added to purgative medicines to prevent griping. Oil of Juniper is diuretic, in doses of 1 to 3 dr. Oil of origanum is almost exclusively used outwardly in stimulating liniments. Oil of bitter almonds is poisonous.

OILS, EMPYREUMATIC. Oil of amber and other empyreumatic oils are antispasmodic; but mostly used in outward applications. The fetid oil, called Dippel's Animal Oil (or oil of hartshorn), is sometimes given as a worm medicine, in doses of 1 oz. (sometimes increased to 2 oz.) to horses, or a drachm to small animals. As an outward application,

it is sometimes added to powders against the fly in sheep; but injures the wool. Oil of paper or rag is an empyreumatic fluid obtained by burning the substances. Mixed with water it is used in inflammation of the eyes, mouth, &c.

OILS, COMPOUND. See Oils and Liniments in the Veterinary Formulary.

OINTMENTS. See V. Formulary.

OLIVE OIL. See Oils, expressed.

ONIONS. Stimulant and diuretic. They are said to be useful in colic and gripes. Externally used in poultices to promote suppuration.

OPIUM. Anodyne, antispasmodic, sedative, indirectly astringent, and in large doses narcotic or stupefactive, and capable of destroying life. In combination with ipecacuanha and tartarized antimony it is sudorific. The dose for horses in ordinary cases is from ½ dr. to 1 dr. But in lockjaw, spasmodic colic, and other urgent cases, it may safely be given in doses of 2 dr., and even (according to MOIROUD) to 4 dr. YOUATT states the dose as 1 dr. to 3 dr. In inflammation of the bowels, after bleeding, it is recommended to give 2 dr. at once, and 1 dr. every hour afterwards until it takes effect. To cattle, the dose is from 10 to 40 gr.; or in lockjaw, &c., 1 dr. Calves, 10 gr. Sheep, 2 to 4 gr. Much larger doses have been given with impunity. Dogs require from ½ gr. to 2 gr., according to size and case. M. MOIROUD says the dose for dogs should not exceed that prescribed for man. Mr. BLAINE thinks they are much less affected by it than men. The dose of tincture of opium is (for horses) from 1 to 2 oz.; of the extract 20 to 30 gr. Externally, opium is used in anodyne liniments, and is useful in inflammation of the eye. See Eye-waters, Liniments, &c., in Veterinary Formulary.

OPODELDOC. Soap liniment. Used externally only, in stimulating liniments.

ORIGANUM. Wild Marjoram. Stimulant. The essential oil is hot and pungent, and a frequent ingredient in liniments for old strains, and in blisters.

ORPIMENT. Yellow Arsenic. Poisonous. Used, mixed with

lard, for fistulous sores, warts, &c., but is not free from danger.

Oxymel. Cooling and pectoral. Used in catarrhal affections. Dose, 3 to 4 oz.

Oxymel of Squills. A stimulating expectorant. Seldom used in veterinary practice.

Palm Oil. Emollient. Used in compounding ointments and liniments; and of late much commended as a basis of aloetic and other balls. It has also been given as a laxative; dose, 12 oz. or more.

Peppermint. Carminative. The distilled water and the essential oil are chiefly used. See Mint.

Pepper, Black, White, and Long. Warm stimulant cordials. The latter kind is chiefly used in veterinary practice. It must be carefully avoided in inflammatory complaints. Dose for horses and cattle, 2 to 4 dr. For Jamaica Pepper, see Allspice.

Pepper, Cayenne. The ground pods of some species of capsicum. See Capsicum.

Periwinkle. The plant, in decoction, or chopped up in a mash, is said to relieve quinsy. Pulverized and mixed with Ethiops Mineral, it has been vaunted as a remedy for glanders.

Peruvian Bark. See Bark.

Petroleum. See Barbadoes Tar.

Pewter. The scrapings are given to dogs for worms. Dose, ½ dr. to 1 dr. Tin filings are safer. See Tin.

Phellandrium. The seeds of water-fennel (Ph. aquaticum) are used in Germany in chronic catarrhal affections. Dose, ½ oz. to 1 oz.

Physic. In veterinary practice this term is applied to purgatives. See Physic or Purging Balls, Veterinary Formulary.

Pimento. See Allspice.

Pitch. Stimulant, balsamic, probably diuretic; but rarely given internally. It is more frequently used externally in charges and warm plasters. For liquid pitch, see Tar.

PITCH, BURGUNDY. Stimulant. Used in charges, and warm and strengthening plasters.

POMEGRANATE. The rind of the fruit is given (in decoction or powder) as an astringent to cattle in diarrhœa. Dose, from ½ oz. to 1 oz. The bark of the root is used to destroy worms. MOIROUD directs 5 or 6 oz. to be boiled in water for some hours, and the decoction given in divided doses.

POPPY-HEADS. Anodyne; but principally used in fomentations. (See also Syrup of Poppies.) An extract prepared by evaporating the expressed juice of the ripe capsules and tops, is said to be nearly half the strength of opium. Of the ordinary extract (from the decoction), 5 grs. are said to equal 2 of opium.—LEBAS.

POTASH, CAUSTIC. *Fused Hydrate of Potash.* A powerful caustic.

POTASH, SUBCARBONATE. *Salt of Tartar or Prepared Kali.* Antacid and diuretic. Dose for a horse, from 2 to 4 dr. or more. It is seldom given alone, but sometimes joined with tonics, stomachics, purgatives, and with other diuretics. The bicarbonate is milder, and may be given in larger doses.

POULTICES. Are useful in relieving inflammation and pain. See V. Formulary.

PRECIPITATE, RED. A mild caustic, and detergent to indolent and foul ulcers.

PRECIPITATE, WHITE. Principally used to destroy vermin in the horse and other animals.

PRUSSIATE OF POTASH. Ferro-prussiate of potash may probably be found useful in veterinary practice; but its properties and uses are not yet properly ascertained. It must not be confounded with the simple prussiate of potash (cyanide of potassium), which is an energetic poison. See Cyanide of Potassium.

PRUSSIC ACID (Medicinal). A strong poison to all animals. Rarely used in veterinary practice. May be given by enema in tetanus. The dose Mr. MORTON states to be from ½ dr. to 1 dr. Mr. YOUATT recommends a lotion composed of a drachm of the medicinal acid in a pint of water, to allay cuticular irritation in dogs.

PULSE. The following table, from VATEL, is inserted as a useful remembrancer to the practitioner:

Table of the Number of Pulsations in a Minute in various Animals.—In the horse, 32 to 38 [36 to 40—WHITE]; ox or cow, 35 to 42 [42 to 45—CLATER; 50 to 55—SPOONER]; ass, 48 to 54; sheep, 70 to 79; goat, 72 to 76; dog, 90 to 100; cat, 110 to 120; rabbit, 120; guinea-pig, 140; duck, 136; hen, 140; heron, 200.

PURGATIVES, *Cathartics or Laxatives.* Medicines which more or less strictly promote evacuations from the bowels. Aloes is almost the only purgative for the horse that is at once certain and safe. For cattle, Epsom or Glauber's salt is the most preferable. Aloes, gamboge, or linseed or castor oil, is sometimes combined with them. Sulphur is used when a very strong purgative is not required; yet this demands some caution. See those various articles.

QUASSIA. A tonic bitter. Dose, 1 or 2 dr., with a little ginger, in debility of the stomach. Its poisonous effects on insects and small animals suggest caution in its use.

QUICKSILVER. See Mercury.

QUININE, SULPHATE OF. Tonic. Dose, ½ dr. to 1 dr. Recommended by Mr. MORTON in the prostration which follows influenza. But too expensive for general use. It is given to dogs in chorea, in doses of 2 to 5 gr.

RAGWORT. The herb is said to produce a kind of lethargy or staggers in horses and cattle. Externally it is used as a poultice in quinsy.

RAKING. Removing hardened fæces from the lower bowel by the hand.

RANUNCULUS REPENS. Acrid stimulant. It is poisonous to sheep.

REED. The great reed (Arundo donax), and the Bankside reed (Arundo phragmitis), are reputed diuretic. The former is supposed to have the property of diminishing the secretion of the milk.

REFRIGERANTS. Cooling Medicines. See Temperants.

RESIN OR ROSIN. Diuretic. ½ oz. to 1 oz. may be given daily to horses in their corn, for swelled legs. The yellow or amber resin is preferable. Externally it is adhesive and

gently stimulating; and is a common ingredient in digestive ointments, and in plasters and charges.

RHODODENDRON. Supposed to be useful in the rheumatism of cattle. Dose, ½ oz. to 1 oz., boiled in water.

RHUBARB. Tonic and stomachic. Scarcely laxative to large animals. From ½ oz. to 1 oz. is given in jaundice, to horses and cattle. On dogs it acts as a purgative, but an uncertain one, in doses of ½ dr. to 1 dr.

ROSEMARY. A mild stimulant and carminative. The essential oil is chiefly used in warm liniments and ointments; but is sometimes given in doses of ½ dr. to 1 or 2 dr. in colic.

RUE. Stimulant, uterine, antispasmodic, and vermifuge. It is also supposed to resist contagion and poisons. A decoction or infusion of 2 to 4 oz. of the fresh herb in water or beer is given for worms; as an antidote to the bite of vipers; with diuretics in farcy; with box leaves as a preventive of hydrophobia; and with camphor and opium in locked-jaw. The bruised leaves are put into horses' ears for the staggers. It is given to poultry for the cure of roup. Externally rue is used in fomentations as a stimulant, antiseptic, and discutient.

SAFFRON. Cordial, antispasmodic, and uterine; but too weak and expensive for veterinary use.

SAGE. Stimulant and tonic. In habitual relaxation of the bowels. The powder may be given in a ball, or the herb infused. The infusion is used as a mouth-wash.

SAGO. Nutritive and demulcent. Used in the form of gruel.

ST. JOHN'S WORT. Vulnerary. The flowers were formerly an ingredient in FRYAR'S Balsam and other similar compounds. An infused oil of the plant is sometimes used in liniments.

SAL AMMONIAC. See Muriate of Ammonia.

SAL PRUNELLA. *Fused Nitre.* Its uses and doses are the same as nitre.

SALICINE. Tonic. Dose, 10 to 30 gr.

SALT, COMMON, or CULINARY. In small doses it is tonic, digestive, and alterative; in large doses purgative and ver-

mifuge. As a digestive, 1 oz. may be sprinkled on the horse's corn. As a purgative, or to expel worms, the dose may be from 4 to 6 oz. It is also a common ingredient in laxative clysters. For cattle, an ounce or more may be sprinkled on the hay, to assist digestion; as a purgative, 4 to 8 oz. may be given, but it is not suitable in inflammatory or febrile diseases. Sheep require 2 oz. as a purgative; or smaller doses daily as a preventive of the rot. To dogs a teaspoonful or one and a half will act as an emetic; smaller doses as a vermifuge. Half a teaspoonful of a solution of salt, as strong as it can be made, is given to poultry as an emetic in roup. Externally, salt dissolved in water is used as a discutient, as a stimulant to old strains, and as a collyrium in chronic ophthalmy.

SALTS, EPSOM AND GLAUBER'S. See Epsom Salts, and Glauber's Salts.

SARSAPARILLA, AND CHINA ROOT, are diaphoretic and alterative; but seldom used in veterinary practice.

SAVIN. An acrid stimulant. The powder is given in doses of 1 to 2 dr. (with or followed by aloes) for worms, but its efficacy is doubtful. Long-continued use of savin is reported to have occasioned the hair to fall off. Externally it is applied, in powder or ointment, to warts.

SCAMMONY. An uncertain as well as expensive purgative, far inferior to aloes.

SCUTELLARIA. *Skull-cap.* Mr. YOUATT and others regard this plant as a preventive of hydrophobia. Dose, 40 gr. daily, gradually increased.

SEA WATER. Laxative. Dose, 2 or 3 pints.

SEDATIVES. Medicines which produce quiet and relieve pain.

SENNA. Purgative; but rarely used in veterinary practice. 5 or 6 ounces are required to purge a horse.

SERPENTARY. Stimulant, tonic, diaphoretic, and antiseptic. It is also supposed to resist the effects of the bites of serpents, &c. Dose, from ½ to 1 oz. or more; but rarely used.

SETONS. These consist of cord, tape, or a mixture of horsehair and hemp twisted together; they are inserted through

a portion of the skin to excite irritation and discharge. Mr. MORTON uses cotton cord soaked in a cantharidal liquid. See No. 15, Blistering Solutions, V. F.

SIMARUBA. Tonic and stomachic, for the same purposes as gentian. Seldom used.

SOAP. Antacid and diuretic. Dose ½ oz. to 2 oz.

SODA. Prepared natron, carbonate, or subcarbonate of soda. The common washing soda is generally sufficiently pure. Antacid and diuretic. Dose, 2 to 4 dr. It is sometimes added to aloes as a corrective, and to tonics in weakness of the stomach. The bicarbonate of soda is milder, and may be given in larger doses.

SODA, CHLORIDE OF. See Chloride of Soda.

SODA, SULPHATE OF, See Glauber's Salts.

SODIUM, CHLORIDE OF. The modern name of common salt. See Salt, Common.

SOOT. Some French veterinarians prescribe from 2 to 3 oz. of soot as a vermifuge. Also used externally in mange, &c. We presume *wood*-soot is intended.

SPERMACETI. Demulcent and pectoral. Dose, ½ oz. to horses in cough; and to cows, after calving. Externally emollient, in ointments.

SPIDER'S WEB. Externally, styptic. Internally has been given to dogs in convulsive fits, in ½-grain doses.

SPIRIT OF HARTSHORN. This ammoniacal liquor is stimulant, antacid, and antispasmodic. Dose, ½ oz. But more frequently used in stimulating liniments, and as an application to the bites and stings of venomous reptiles and insects.

SPIRITS OF SAL VOLATILE. This also owes its pungency to ammonia. Dose, ½ oz.

SPIRITS, ARDENT. Brandy, gin, and rum are given as stimulants and antispasmodics, especially in colic. Dose, from 2 to 4 or 5 oz. with warm water. Rectified spirits of wine may be given in the same way, in smaller doses (1 to 2 oz.); but is more commonly employed for making tinctures; and externally in lotions.

SPIRIT OF MINDERERUS. See Acetate of Ammonia.

Spirit of Nitre, Sweet. *Spirit of Nitric Æther.* Diuretic, diaphoretic, and antispasmodic. Dose for horses, in fever, ½ oz. 3 times a day. In colic, from ½ oz. to 2 oz. Cattle, ½ oz. to 1 oz. in low fevers. Sheep, 1 dr. Dog, from 10 to 20 drops.

Squill. A stimulating expectorant. Dose for a horse, 1 dr.; for cattle, 1½ to 2 dr. It is also applied in frictions to the abdomen. Moiroud has seen it remove ascites.

Starch. Demulcent. Chiefly used in clysters, but sometimes also in drinks. Dose, 1 to 2 oz., rubbed smooth with a little cold water, and then boiled in 3 or 4 pints of water. It is occasionally used in fomentations.

Stavesacre Seeds. Poisonous. 2 dr. will destroy a horse. Only used outwardly to destroy vermin, either powdered and mixed with grease, or infused in vinegar.

Steel, Salt of. See Iron, Sulphate of. For the other preparations (so called) of steel, see *Iron.*

Stimulants. See Excitants. Diffusible stimulants are those which produce a sudden and temporary excitement of the circulation and of the nervous system.

Stomachics. Medicines which invigorate the stomach and promote digestion.

Stoppings. Compositions employed to keep the feet moist and supple. The term is also applied to mechanical plugs for the feet when they are dry and diseased, as cow-dung, clay, tar, &c.

Storax. Balsamic and expectorant. Dose, ¼ oz. But rarely used.

Strychnia. The active principle of nux vomica: chiefly used in paralysis. Dose, 1 to 3 grains; to be very cautiously increased if necessary, 15 grains have proved fatal. Dose for the dog, 1-16th to 1-8th of a grain.

Styptics. Astringent applications employed locally to stop bleeding.

Sublimate, Corrosive. *Bichloride of Mercury.* See Corrosive Sublimate.

Sugar, Syrup, and Treacle. These are used to sweeten drinks; and to give form to balls and other compounds.

Sugar of Lead. See Lead, Acetate of.

SULPHATE OF COPPER. *Blue Stone.* See Copper, Sulphate of.

SULPHATE OF IRON. See Iron, Sulphate of.

SULPHATE OF MAGNESIA. See Epsom Salts.

SULPHATE OF POTASH. Purgative; but seldom used. Dose, 2 to 4 ounces, in colic, &c.

SULPHATE OF QUININE. Tonic. Dose, ½ dr. to 1 dr.

SULPHATE OF SODA. See Glauber's Salt.

SULPHATE OF ZINC. *White Vitriol.* See Zinc, Sulphate of.

SULPHUR, OR BRIMSTONE. It is in 3 forms—roll brimstone, flowers of sulphur, and black brimstone or sulphur vivum. The flowers are generally used. The black is very impure, and sometimes contains arsenic. Sulphur is laxative, alterative, and pectoral. Dose, to horses, as an alterative in skin diseases, grease, want of condition, &c., 1 oz. As a laxative, 4 or 5 oz., but it is rarely employed with this view, and very large doses are not always safe. To cattle, as a laxative, 6 or 8 oz. Sheep, 2 or 3 oz. Dogs, 1 dr. in milk. Swine, 2 dr. It is used outwardly in ointments, for mange in all animals. As an alterative it is usually combined with antimonials and nitre.

SULPHURET OF IRON. It has been used in hemorrhage, dysentery, and worms. The hydrated persulphuret (see Ferri persulphuretum hydratum, Pocket Formulary) is strongly recommended by BOUCHARDAT as an antidote for metallic poisons; also as a remedy for incipient farcy. Dose, 1½ oz. to 8 oz.

SULPHURET OF MERCURY. See Ethiops Mineral, and Cinnabar.

SULPHURET OF POTASH. Mr. BLAINE prescribes 2 dr. with astringents, in diabetes. In large doses it is poisonous.

SULPHURIC ACID. Poisonous. The strong acid (*oil of vitriol*) is used as a powerful caustic. It is also used in ointments, or mixed with tar to form an external application. In small doses, about 1 to 2 dr., plentifully diluted, it is, rarely, given as a tonic. The diluted acid (1 oz. to a pint) is used as a lotion in grease, foul ulcers, &c.

SUPERTARTRATE (BITARTRATE) OF POTASH. See Cream of Tartar.

SYRUP OF BUCKTHORN. Purgative. Seldom given to horses,

except when used in forming powders into balls. Dose for cattle, 2 to 4 oz. with castor oil. A common physic for dogs; dose 2 to 4 dr.

Syrup of Poppies. A mild anodyne and sedative. Dose for dogs, 1 dr. Seldom given to large animals.

Tannin (or Tannic Acid). The astringent principle of nut-galls. A powerful astringent in diarrhœa, &c. Dose, 5 to 10 gr. Catechu is more generally used.

Tansy. Tonic and vermifuge. Externally in fomentations.

Tar, Barbadoes. See Petroleum.

Tar. Internally, in old coughs, from 2 to 4 dr. Externally it is cleansing and gently stimulating. It is particularly useful in thrushes and all diseases and wounds of the feet both of horses and cattle, to punctured wounds, and for the cure of mange and other skin diseases. Mixed with fish oil, it is applied with a brush to hard, brittle feet. Tar water (see Formulary) is also given in chronic coughs. Oil or spirit of tar is used in mange ointments, and as a dressing for sheep. The latter requires some caution: sheep have been killed by it.

Tartar Emetic. See Antimony, Tartarized.

Tea. Tonic, in simple indigestion, or when connected with staggers [Delafond]. Dose, 4 to 6 dr., infused in 3 or 4 pints of water.

Temperants. Medicines which moderate the circulation and reduce animal heat.

Tin. Vermifuge. A drachm of the filings daily to dogs. A horse requires from 1 to 3 oz.

Tobacco. An acro-narcotic poison. In small doses, diuretic and emetic. Principally used as a wash for the mange, and to destroy lice and fly in sheep. But it is not altogether safe, as it is apt to be absorbed. It vomits the dog, pig, and cat: but there are safer emetics. Herbivorous animals are less rapidly affected by it, but instances of its having proved fatal to them are recorded. In some parts of France, jockeys are said to stupefy vicious horses for sale by tobacco diffused in spirits.

Tonics. Medicines which give tone to the fibres, and invigorate the system when relaxed and debilitated. The principal tonics used in veterinary medicines are gentian,

Peruvian bark, chamomile, and other vegetable bitters and astringents; and the preparations of iron, copper, arsenic, zinc, &c. The over free use of them, particularly when fever and inflammation are present, is a frequent source of mischief.

TORMENTIL ROOT. Astringent. Dose, 1 oz. to 1½ oz. Its presence in pastures is supposed to prevent the rot in sheep.

TURBITH MINERAL. *Subsulphate of Mercury.* An irritating purgative, and in large doses, poisonous. Dose, ½ dr. in farcy. Given to dogs as an emetic; dose, 1 gr. to 3 gr.

TURMERIC. A weak aromatic stimulant. Supposed to be useful in jaundice, or yellows. Dose, 1 oz.

TURPENTINES. They are all stimulant, diuretic, and expectorant; and in larger doses, vermifuge and purgative. Dose of common turpentine, ½ oz. to 1 oz. They are used in digestive ointments. Oil or spirit of turpentine is a more stimulating diuretic, in doses of 2 to 4 dr.; it is also considered efficacious as an antispasmodic in colic (gripes), and as a remedy for worms. Dose for the latter purposes, from 2 to 4 oz., or sometimes still larger doses for worms. To cattle (in hoose from worms in the bronchial passages) about 2 oz. To sheep, in rot, 1 dr. It is not a safe medicine for dogs; but is sometimes given in doses of 2 dr. with olive oil. Externally, it is used in stimulating liniments, embrocations, ointments, &c. It is very irritating to the skin of the horse, and also of the dog, instantly producing great excitement. Like the common and Venice turpentine, it enters into the composition of some digestive ointments.

UVA URSI. *Bearberry.* Astringent. Dose, 4 to 6 dr., in diabetes. But GIRARDI says it inflames the stomach.

VALERIAN. A stimulant, acting chiefly on the nervous system. Dose, for horses and cattle, 1 to 4 oz. in powder. 2 oz. twice a day have been given to a horse without any observable effect. In dogs it is said to act as a vermifuge. Dose, 1 to 4 dr.

VERDIGRIS. *Subacetate (or Diacetate) of Copper.* Tonic, caustic, poisonous. It has been given in doses of 1 dr. to 2 dr. daily, in farcy and glanders. Externally, detergent

and caustic, in ointments, and in the form of Ægyptiacum. The crystallized acetate of copper is more powerful in its action.

VERJUICE. Properties and uses the same as of vinegar; but preferred by some for outward use.

VERMILION. See Cinnabar.

VINEGAR. Diaphoretic, cooling, and antiseptic. In combination with honey, it is used in coughs. In large quantities, it irritates the stomach; a pint is said to have destroyed a horse. It should always be plentifully diluted. Vinegar which contains much sulphuric acid should be avoided. It is chiefly used as an external application, as a lotion for strains, bruises, sprains, and inflammations; and hot as a revulsive. The vapors are thought to possess disinfecting properties, but are less effectual than chlorine.

VITRIOLATED ZINC. *White Vitriol.* See Zinc, Sulphate of.

VITRIOL, BLUE. See Sulphate of Copper.

WALNUT. The green shells are astringent, and sometimes applied, bruised, as a cataplasm, or in decoction as a lotion.

WATER. Besides its use as a drink, and as a vehicle for medicines, water is used remedially, on the hydropathic system. Rugs wetted with cold water, and well covered with dry ones, are used to produce perspiration, assisting their operation by copious draughts of cold water, adding 4 oz. of sweet spirit of nitre to each pailful. This treatment is said to have succeeded in epidemics of pleuropneumonia.

WAX. Chiefly used in making cerates, plasters, charges, &c.

WHEY. A cooling and nutritive drink in inflammatory diseases, and during convalescence from them.

WILLOW BARK. Possesses in some degree the same properties as Peruvian bark. Dose, in powder or decoction, 1 to 4 oz.

WINE. Stimulant. In wine countries it is frequently given as a restorative. Port wine has been given as an astringent in obstinate diarrhœas. Dose for horses and cattle, ½ pint to a bottle.

WINTER BARK. A warm tonic and stomachic. Dose, 2 to 6 dr.

Wolfsbane. *Aconite.* A virulent poison.

Wormwood. A bitter tonic and vermifuge. An infusion from 2 to 4 oz. of the dry, or twice as much of the fresh herb, may be given in dropsy, and diseases of general debility; or from 2 to 4 dr. of the powder may be given in a ball. A few drops of the essential oil are often added to aloes, &c., for worms.

Wort. See Malt.

Yew. It is not used medicinally. The leaves are poisonous to horses and cattle, producing symptoms which resemble those of apoplexy. To counteract its effect, it is recommended to give 10 gr. of croton meal, and afterwards drenches of gruel with vinegar. The croton to be repeated in 6 hours if it has not operated.

Zedoary. A weak aromatic stimulant, formerly prescribed in jaundice, but now rarely employed. It is weaker than ginger.

Zinc, Chloride of. In solution this constitutes Sir W. Burnett's disinfecting fluid. Much diluted it is applied as a detergent lotion of foul ulcers. The dry salt is a powerful caustic.

Zinc, Oxide of. *Flowers of Zinc.* A mild astringent and tonic; dose, ½ oz.; but chiefly used in dusting ulcers and excoriations, to promote skinning.

Zinc, Sulphate of. *White Vitrol.* Tonic. Dose, for the horse, 1 to 4 dr. frequently combined with cantharides. Externally, astringent, detersive, styptic, and healing; in lotions and ointments, to indolent ulcers, grease, &c. It is a frequent ingredient in eye-waters,—about 3 gr. to an ounce of water. A saturated solution is used as an injection for quitters.

VETERINARY FORMULARY.

Medicines for Horses.

BALL AND BALL MASSES.

THE roots, seeds, and other dry substances are to be reduced to powder; and it is of importance that the aromatic seeds, especially, should have been recently powdered. The drugs should be of good quality. It is hoped that the trash sold as horse-powders will not much longer be known in establishments which have any pretension to respectability. After this general notice it will be unnecessary to occupy the space by repeating the words "powdered," "freshly powdered," "genuine," &c. Balls should not be too hard, but merely stiff enough to retain their form, and should be wrapped in soft paper.

[*Mode of administering Balls.*—The horse should be backed into the stall, the tongue drawn gently out with the left hand on the off side of the mouth, and then fixed by pressing the fingers against the side of the lower jaw. The ball being now taken between the tips of the fingers of the right hand, must be passed rapidly up the mouth, as near the palate as possible, until it reach the root of the tongue; it must then be delivered with a slight jerk, so that the hand being immediately withdrawn, and the tongue liberated, the ball may be forced through the pharynx into the œsophagus. A slight tap under the chin may then be given, or a draught of water to assist in carrying it down.]

COMMON MASS, as a basis for balls in general. Mix with the hand equal weights of linseed meal and treacle, and add a little palm oil.—CHERRY.

ALTERATIVE BALLS. The term *alterative* is applied to medicines which, without any sensible operation, or with a laxative or diuretic operation so gradual as not to interfere with the usual work or diet, produce a favorable change in the system, and, in common language, "purify the blood." Alterative balls are given in skin diseases, swelled legs, grease, foul humors, &c.; usually 1 daily, or every other day.

Diuretic Alterative Balls. 1. Dried common soda 1 oz., Castile soap 6 dr., resin 2 oz., liquorice powder ½ oz., Barbadoes tar to form 6 balls; 1 daily.—WHITE.

2. Acetate of potash ½ oz., resin ½ oz., fenugreek 1 oz., treacle enough to form a mass for two balls; 1 daily.

Laxative Alterative Balls. 1. Aloes 4 oz., soft soap 4 oz., common mass 24 oz.; mix; dose, 1 oz.—V. C.

2. Socotrine aloes 8 oz., soft soap 8 oz., common mass 16 oz.; mix; dose, 1 oz.—V. C.

3. Aloes 10 dr., soap 12 dr., caraways 12 dr., ginger, 4 dr., treacle q. s. for 4 balls; 1 daily.—WHITE.

4. Aloes 1 dr., diuretic mass (No. 1 or 3) 9 dr.

5. Antimonial powder 1 dr., aloes 1 or 2 dr., diuretic mass (No. 1 or 3) 1 oz.

Antimonial or Diaphoretic Alterative Balls. 1. Levigated antimony 2 or 4 dr., caraway seeds 4 dr., treacle q. s. to form a ball.—WHITE.

2. Prepared antimony 2 dr., nitre 3 dr., sulphur 2 dr., linseed meal 2 dr., palm oil to form a mass; one every night in megrims.—CLATER.

3. Tartarized antimony 2 dr., elecampane 2 oz., guaiacum 6 dr., sulphur 1 oz., treacle and flour to form 6 balls; 1 daily.

4. Tartarized antimony 3 dr., ginger a scruple, soap 1 oz. For 3 balls; one every other morning.—VINES.

5. Emetic tartar 5 oz., ginger 3 oz., opium 1 oz., and syrup to make 16 balls.

Mercurial Alterative Balls. 1. Ethiops mineral 4 oz., sulphur, prepared antimony, cream of tartar, cinnabar, of each 5 oz., honey to form a mass for 12 balls; 1 every morning for a month in farcy.—TAPLIN.

2. Calomel ½ dr., aloes 1 dr., Castile soap 2 dr., oil of juniper 30 drops, syrup to form a ball.—WHITE.

3. Blue pill 1 dr., black antimony 2 dr., diuretic mass 4 dr., aloes 1 dr.; for a ball daily.

4. In grease: prepared antimony, sulphur, nitre, Ethiops mineral, of each 3 oz., Castile soap 10 oz., oil of juniper 3 dr., syrup of honey q. s. for 12 balls; 1 every morning for 2 or 3 weeks.—TAPLIN.

5. Quicksilver 2 parts, sesquioxide of iron 1 part, confection of roses 3 parts. Rub together till the quicksilver disappears. Dose, ʒss. to ʒij. with common or other mass q. s.—Dr. COLLIER'S Blue Pill.

6. Strong mercurial ointment ¼ ℔, powdered ginger 3 oz., liquorice powder 10 oz., treacle to mix for 12 balls.—FRANCIS.

Alterative Tonic Balls. See Tonic Balls.

ASTRINGENT BALLS. These are given in diarrhœa, diabetes, &c.

1. (V. C. Astringent Mass.) Catechu 1 oz., cinnamon 1 oz., common mass 6 oz.; mix; dose, 1 oz.

2. Peruvian bark 12 oz., grains of paradise 2 oz., gentian 3 oz., honey q. s. for 16 balls; 1 every morning; for diabetes.—RYDING.

3. Catechu ½ oz., alum 3 dr., cascarilla 2 dr., flour 2 dr., treacle q. s.—WHITE.

4. Catechu 2 dr., opium ½ dr., linseed meal 2 dr., treacle to form a ball. For profuse staling, 1 night and morning; if they confine the bowels, add 1 dr. of aloes—CLATER.

5. Peruvian bark 1½ oz., alum ½ oz., treacle q. s. For the same purpose.—LAWRENCE.

6. Oak bark 1 oz. (or peruvian bark ½ oz.), opium 1 dr., ginger 2 dr., syrup to form a ball; for diarrhœa.—WHITE.

7. Opium ½ dr., prepared chalk 6 dr., cassia 1½ dr., tartarized antimony 2 dr., syrup to form a ball; for the same.—WHITE.

8. Nutgalls 2 dr., cassia ½ dr., conserve of roses to form a ball.

9. Burnt rhubarb 1 dr., compound powder of chalk 3 dr., common mass 6 dr.; for diarrhœa.

10. Tormentil or bistort 1½ dr., marshmallow root ½ oz., chalk 2 dr., syrup to form a ball.

11. For bloody urine. Acetate of lead 10 gr., sulphate of zinc 40 gr., catechu 4 dr., conserve of roses to form a ball; once daily.—BLAINE.

12. Powdered opium ½ dr., soda 1 dr., powdered cassia or ginger 1½ dr., flour and syrup to form a ball.

13. For diabetes. Catechu ½ oz., alum ½ dr., sugar of lead 10 gr., with conserve of roses to form a ball. See also Tonic Pills.

COUGH BALLS; Expectorant Balls. The following formulæ are chiefly intended for chronic coughs and thickness of wind. The bowels should be kept open by mashes and an occasional laxative. Coughs occasioned by worms require a different treatment. In coughs connected with inflammation of the chest, and epidemic catarrh, see Balls for Inflammation of the Lungs.

1. Aloes 2 oz., digitalis (powdered) 1 oz., common mass 13 oz.; dose, 1 oz., twice a day.—MORTON.

2. Emetic tartar ½ dr., digitalis ½ dr., nitre 1½ dr., tar enough to form a ball; every night.—YOUATT.

3. Powdered squill 1 dr., gum ammoniac 3 dr., opium ½ dr., syrup to form a ball.—WHITE.

4. Ipecacuanha 1 dr., camphor 2 dr., liquorice powder 1 dr., honey to form a ball; to be given every morning.—BLAINE.

5. Sulphur ½ oz., assafœtida 1 oz., liquorice powder 1 oz., Venice turpentine 1 oz., for 4 balls; one every night for 4 times.—HINDS.

6. Calomel 26 gr., gum ammoniacum 2 dr., balsam of Peru 1 dr., p. squill 1 dr., honey to form a ball; one every morning.—BLAINE.

7. P. marshmallow root and liquorice, of each 1 dr., elecampane, sulphur, and Kermes mineral, of each ½ dr., honey to form a ball; twice a day.—LEBAS.

8. Squill 2 dr., gum ammoniac 4 dr., ipecacuanha 4 dr., opium 4 dr., pimento 1 oz., balsam of sulphur 4 oz., Castile soap 2 oz., treacle to form a mass for 6 balls; one twice a day.—HINDS.

9. Spermaceti 1 oz., balsam of copaiva 1 oz., benzoin 2 dr., sulphur 2 oz., elecampane 2 oz., p. squill 4 dr., emetic tartar 2 dr., syrup of poppies to form a mass for 8 balls.—B. CLARKE.

10. Liquorice powder ½ oz., linseed or barley meal 1 oz., tar 1 dr., honey to form a ball.

11. Castile soap, aniseed, liquorice, of each 5 oz., Barbadoes tar 6 oz., ammoniacum 3 oz., balsam of Tolu 1 oz., honey q. s. to make a mass for 12 balls: one every morning for a fortnight.—TAPLIN.

12. Digitalis 1 dr., nitre 2 dr., liquorice 4 dr., tar enough to form a ball.—CLATER.

13. Digitalis ½ dr., camphor 1 dr., emetic tartar 1 dr., nitre 3 dr., linseed meal 1 dr., make up with Barbadoes tar, and give one daily.—SPOONER. See also Mixed Balls (Pectoral Cordial).

BALLS FOR INFLAMMATION OF THE LUNGS, BRONCHITIS, &c.

1. Antimonial powder 2 dr., digitalis 3 dr., nitre 3 dr., cream of tartar 3 dr., honey to form a ball; 1 every, 4, 6, or 8 hours, in inflammation of the lungs.—BLAINE.

2. Digitalis 1 dr., emetic tartar 1½ dr., nitre 3 dr., honey q. s.; when the pulse intermits, reduce the dose to half.—YOUATT.

3. Nitre 6 dr., emetic tartar 2 dr., flour and syrup to form a ball; twice a day.—WHITE.

4. Digitalis 1 dr., emetic tartar 1 dr., nitre 3 dr., sulphur 1 dr., linseed meal 2 dr.; beat together with palm oil.—CLATER.

5. Epidemic catarrh. To the last add 2 drachms of the Physic Mass (No. 10): repeat this twice.

6. For pneumonia. White hellebore ½ dr. (or extract of belladonna 2 dr., or digitalis 1 dr., or calomel 1 dr. with opium ½ dr.) emetic tartar ℈j, nitre and linseed meal each 2 dr.; one twice a day.—SPOONER.

7. In the advanced stage, when suppuration has taken place. Carbonate of ammonia 1½ dr., opium 1 dr., aniseed ½ oz., syrup to form a ball.—SPOONER.

8. *Cough Ball.* Digitalis ½ dr., camphor and emetic tartar each 1 dr., nitre 3 dr., and linseed meal 1 dr., to be made up with Barbadoes tar.

CORDIAL BALLS. For exhaustion from over-exertion, and as a stimulant to weak stomachs. But their frequent and unnecessary use is hurtful.

1. Ginger and gentian equal parts, treacle to form a mass; dose, 1 oz. to 1½ oz.—V. C.

2. Caraway, bruised raisins, of each 4 parts, ginger and palm oil, of each 2 parts.—YOUATT.

3. Aniseed, caraway, cardamom, each 1 oz., saffron 2 dr., sugar candy 4 oz., liquorice powder 1½ oz., Spanish juice

(softened with water) 2 oz., oil of aniseed ½ oz., wheat flour q. s.; dose, 1 oz. to 1½ oz.—BRACKEN.

4. Aniseed, caraway, sweet fennel, liquorice, of each 4 oz., of ginger and cassia, each 1½ oz., honey to form a mass.—WHITE.

5. Ginger, caraway, each 4 lb, gentian 1 lb, palm oil 4½ lb, beat together; dose, 1 oz. to 1½ oz.—CLATER.

6. Gentian 8 oz., ginger 4 oz., coriander 8 oz., caraway 8 oz., oil of aniseed ½ oz., treacle q. s.; dose, 1½ oz.—BLAINE.

7. Aniseed, caraway, ginger, each 8 oz., gentian, grains of paradise, cumin, and turmeric, each 4 oz., cassia 2 oz., oil of caraway 2 dr., treacle to form a mass; dose, 1½ oz. To keep it moist, add 2 oz. of acetate of potash.

8. Cumin, aniseed, caraway, each 4 oz., ginger 2 oz., treacle q. s.; dose, 1½ oz. to 2 oz.—WHITE.

9. Pimento 1 lb, sifted barley meal 2 lb, treacle q. s.—B. CLARKE.

MIXED BALLS. *Cordial Astringent Ball.* Cordial Ball (No. 2) 1 oz., catechu 1 dr., opium 10 gr.; to washy horses, before or after a journey.—YOUATT.

Cordial Anodyne Balls. 1 Cordial mass (No. 6) 10 dr., camphor 1 dr., opium 20 gr.—BLAINE.

2. Opium ½ dr. to 2 scruples, soap 2 dr., ginger 1 dr., aniseed 4 dr., oil of caraway ½ dr., treacle q. s.—WHITE.

Balsamic Cordial Ball. Cordial mass (No. 6) 1 oz., myrrh 1 dr., balsam of Tolu 1 dr.—BLAINE.

Pectoral Cordial Balls. 1. For old coughs. Fenugreek, aniseed, cumin, safflower, elecampane, coltsfoot, sulphur, of each 3 oz., liquorice juice 1 oz., olive oil 8 oz., honey 8 oz., Genoa treacle 12 oz., oil of aniseed 1 oz., wheat meal 1½ lb, or q. s.; one ball or 2 oz. (dissolved in water or warm wort), every day for 12 or 15 days if required.—QUINCY.

2. Elecampane ½ oz., ginger 1½ dr., squill 1 dr., oil of aniseed 20 drops, syrup of Tolu q. s.—WHITE.

Diuretic Cordial Balls, to fine the legs of debilitated and overworked horses, and sometimes given in old coughs, &c. 1. Resin 2 oz., soap, nitre, caraway, of each 2 oz., ginger 1½ oz., sulphur 2 oz., oil of caraway ½ dr., oil of juniper ½ dr., syrup to form a mass.

2. Soap and common turpentine each 4 dr., ginger 1 dr., opium ½ dr., caraway seed q. s. for 1 dose.—WHITE.

3. Strained turpentine 8 oz., resin 4 oz., olive oil 2 oz., soap 8 oz.: melt together and add powdered ginger 6 oz., pimento 6 oz., liquorice powder q. s. to form a mass.

4. Resin 4 dr., nitre 2 dr., and ginger 1 dr., with sufficient soap to form a ball.—SPOONER.

DIURETIC BALLS. For swelled legs, grease, &c., for carrying off bad humors, and in many chronic diseases. The too free use of diuretics injures the kidneys, and weakens the system. See Alterative Balls (Diuretic), page 59.

1. Resin, soap, nitre, of each equal parts, beaten together into a mass; dose, 1 oz. to 1½ oz.—V. C.

2. Common turpentine 4 oz., Castile soap 4 oz., caraway 8 oz., ginger 1 oz., flour q. s.—WHITE.

3. Resin 16 oz., white soap 16 oz., nitre 8 oz., dried common soda 2 oz., oil of juniper 4 oz.; beat together, adding flour if required; dose, 1 oz. to 1½ oz.

4. Nitre 1 lb, Castile soap ½ lb, common turpentine 1 lb, barley meal 2½ lb, or sufficient; dose, about 1 oz.—B. CLARKE.

5. White soap 8 oz., nitre 3 oz., resin 3 oz., camphor 3 dr., oil of juniper 3 dr. For 6 balls; 1 every, or every other morning.—TAPLIN.

6. Common turpentine 16 oz., sulphur 24 oz., nitre 8 oz., honey 8 oz., flour or linseed meal q. s.; dose 1½ oz.

7. Camphor 2 dr., nitre 1 oz., flour and syrup to form a ball; for stoppage of water.—WHITE.

8. Yellow resin 4 lb, common turpentine 2 lb, yellow soap 2 lb, melt together, and add nitre 1 lb.—BLAINE.

9. Common turpentine (or powdered resin) ½ oz., linseed meal ¼ oz., ginger ½ dr., palm oil q. s.—YOUATT.

10. Yellow resin 2 oz., common turpentine 4 oz., soap 3 oz., melt together, stir in 1 oz. sweet oil, add oil of aniseed ½ oz., oil of juniper ½ oz., ginger 2 dr., linseed meal q. s.; mix, and divide into 8 balls; 1 a day till the water is affected. —HINDS.

11. Resin 2½ lb, cream of tartar ½ lb, sulphur ½ lb, linseed meal 1 lb, palm oil 1 lb; dose, 1 oz. to 2 oz.—CLATER.

12. Nitre 1 oz., vermilion ½ oz., resin 1 oz., camphor ½ oz., honey q. s. for 4 balls.—LEBAS.

13. Powdered yellow resin 4 dr., nitrate of potash 2 dr., p. ginger 1 dr.; beat up with soap.—SPOONER.

14. Nitre 8 oz., oxysulphuret of antimony 1 oz., sulphur 8 oz., resin 8 oz., oil of juniper 1 oz., yellow soap 8 oz., treacle to form a mass; dose 1½ oz.

15. White soap 1 oz., extract of juniper berries q. s. for 2 balls.—BOURGELAT.

Tonic Diuretic Ball. Gentian 1 dr., ginger ½ dr., sulphate of iron 2 dr, diuretic mass (No. 11) ½ oz., oil of juniper 10 drops, syrup of squills ½ oz.; twice a day in dropsy of chest; less frequently in swelled legs.—CLATER.

Tonic and Diuretic Ball for Pleurisy. Sulphate of copper 1½ dr., ginger and gentian 2 dr. each, with Venice turpentine.—SPOONER. See also Leicester Red Balls. (Miscellaneous Balls.)

FEVER BALLS.

1. Emetic tartar ½ dr., camphor ½ dr., nitre 2 dr., common mass 6 dr., or q. s. for 1 ball; to be given once or twice a day.—MORTON.

2. Camphor 1 dr., nitre 6 dr., antimonial powder 2 dr., flour and syrup to form a ball.—WHITE.

3. Antimonial powder 2 dr., nitre 3 dr., cream of tartar 2 dr., honey to form a ball; in influenza, twice a day, after a mild laxative.—BLAINE.

4. See Balls for Inflammation of Lungs, No. 4.—CLATER.

BALLS FOR FARCY AND GLANDERS. Mr. COLEMAN says he has tried the various preparations of arsenic, antimony, copper, mercury, zinc, aconite, digitalis, hemlock, henbane, hellebore, nightshade, &c., in glanders, without any specific or curative effect. Mr YOUATT considers it useless to attempt the cure of glandered horses; but that farcy in its early stages and mild form may be successfully treated. Mr. BLAINE says, "All the mercurials have been used with benefit in farcy; but they must be discontinued as soon as the mouth is affected, or sickness, loss of appetite, &c., produced."

1. Ethiops mineral 2 dr., blue pill 1 dr., prepared antimony 3 dr., diuretic mass 4 dr. One every morning.

2. Strong mercurial ointment 2 to 3 dr., guaiacum 3 dr., soap 4 dr., fenugreek 12 dr., treacle to form a mass, for 6 balls. [See Mercurial Alterative Balls, p. 59.]

3. Sulphate of copper 1 dr., corrosive sublimate 8 gr., linseed powder ½ oz.—WHITE.

4. Corrosive sublimate 10 gr., gradually increased to 20, gentian 2 dr., ginger 1 dr., syrup to form a ball; to be given night and morning till some effect is produced; when the mouth is affected, the sublimate may be exchanged for 1 dr. sulphate of copper.—YOUATT.

5. Corrosive sublimate 10 to 20 gr., opium ½ to 1 dr., powdered aniseed ½ oz., with syrup to make a ball.

6. Sulphate of copper 1 dr., calomel 20 gr., common turpentine 3 dr., liquorice powder and syrup q. s. for one ball.—COLEMAN.

7. Sulphate of copper 1 dr., white arsenic 8 gr., corrosive sublimate 8 gr., linseed powder ½ oz., syrup to form a ball.—WHITE.

8. Ethiops mineral 2 dr., opium 10 gr., liquorice powder and mucilage to form a ball; to be given twice a day till the breath or urine is affected.—HINDS.

9. Sulphate of iron 2 dr., Peruvian bark 1 oz., opium ½ dr., syrup to form a ball.—SMITH.

10. Cantharides 4 gr., gradually increased to 6 or 8 gr., gentian, ginger, and caraway, each 1 dr., syrup q. s.; every, or every other day.—VINES.

11. Sulphate of iron 2 dr., iodide of potassium 10 gr., ginger 1 dr., gentian 2 dr., made into a ball with treacle.—SPOONER.

12. Diniodide of copper 1 dr., gentian 1½ dr., pimento 1 dr., cantharides 5 gr.; for one ball.—MORTON.

13. Sulphate of zinc 15 gr., cantharides 7 gr., pimento or ginger 15 gr., treacle and oatmeal to form a ball; 1 daily.—BRACY CLARK.

14. Sublimate, arsenic, verdigris, each 8 gr., sulphate of copper 20 gr., for one ball (with common mass q. s.); the dose may be gradually increased, carefully watching its effects, but should never exceed 15 gr. of sublimate and arsenic.—BLAINE.

15. Sublimate 10 gr., gentian 2 dr., ginger 1 dr., linseed meal ½ oz., palm oil to form a ball; night and morning for a fortnight; for farcy.—CLATER.

16. Sulphate of copper ½ dr. to 1 dr., ginger and gentian, each 1 dr., linseed meal and palm oil to form a ball; morning and night for a fortnight, then daily as long as necessary: in glanders.—CLATER.

17. Strong mercurial ointment 3 oz., white soap 2 oz.,

starch 2 oz.; form a mass and divide into 12 balls; 1 every morning.—MOIROUD.

18. Assafœtida 3 oz., vermilion 2 oz., muriate of lime 3 dr., galangal 1 oz., strong mercurial ointment 2 oz.; beat together into a uniform mass, and divide into 6 balls; one every other morning.—LEBAS.

19. Ethiops mineral 8 oz., powdered burdock root 16 oz., treacle q. s.; make into 32 balls.—MOIROUD.

20. Antihecticum Poterii 2 dr., with 6 dr. of cordial ball; every other day.—LAWRENCE.

21. Calomel 1 oz., assafœtida 4 oz., galangal powder 1 oz., mercurial ointment 2 oz. Mix, and form 6 balls. One every other morning.—LEBAS.

22. Hydrargyro-iodide of potassium (see Hydrargyri et Potassii Iodidum, Pocket Formulary) 3 oz., powdered althæa root, and honey q. s. to make 100 balls. Give from one to eight daily, gradually increasing the dose to the latter number.—BOUCHARDAT.

BALLS FOR GREASE. See Diuretic Balls, and Alterative Balls.

BALLS FOR YELLOWS, OR JAUNDICE, AND INFLAMMATION OF LIVER (HEPATITIS).

1. For hepatitis without purging: calomel 1 dr., antimonial powder 2 dr., aloes 3 dr., syrup to form a ball; one every four or five hours, till the bowels are opened.—BLAINE.

2. Calomel ½ dr., aloes 1 dr., soap 2 dr., rhubarb ½ oz., syrup to form a ball; to be given every 12 hours, till it purges moderately.—WHITE.

3. Aloes 2 dr., calomel 1 dr., syrup to make a ball, twice a day.—YOUATT.

4. Opium 1 dr., calomel 1 dr., emetic tartar 2 dr.; liquorice powder 3 dr., syrup to form a ball; once every 12 hours.—WHITE.

5. Opium ½ dr., calomel 1 dr., resin 3 dr., carbonate of potash 2 dr., with soft soap. To be preceded by blisters to the side, and purgative draughts.—SPOONER.

Yellows (Jaundice) without Fever. 1. Calomel 1 dr., aloes 2 dr., soap 2 dr.; for one ball; night and morning till purged, then so as to keep them lax.—BLAINE.

2. Calomel ½ dr.; aloes 1½ dr., Castile soap 2 dr., rhubarb 3 dr., syrup to form a ball.—WHITE.

3. In the latter stage, when not costive, calomel 12 gr., sulphate of copper 1 dr., gentian 3 dr., oak bark 3 dr., chamomile 3 dr., syrup to form a ball; once or twice a day. —BLAINE.

PHYSIC OR PURGING BALLS. The animal should be prepared by bran mashes for two days, and the ball given fasting in the morning. Gentle exercise with a ball is useful, but not after it begins to operate. Genuine Barbadoes alone should be used (from the gourd not melted), and the dose seldom need exceed 6 dr. A week should be allowed after the operation of one ball before another is given. See Aloes, in the Veterinary Materia Medica.

1. (V. C. Cathartic Mass.) Bruised B. aloes 8 oz., olive oil 1 oz.; melt together in a vessel placed in hot water; remove it from the fire, add 3 oz. of treacle, and stir altogether; dose, 6 to 12 dr., equal to 4 to 8 dr. of aloes.

2. (V. C. Stronger.) To each dose of the last add from 4 to 8 drops of croton oil.

3. B. aloes 4 to 8 dr., soap 3 to 4 dr., ginger 1 dr., oil of cloves 10 drops (or oil of caraway or aniseed 20 drops), water 1 dr. or q. s.; beat together into a mass.—WHITE. Mr. W. says is the best that can be employed.

4. B. aloes 15 oz., ginger 1 oz.; mix and beat up with 8 oz. of palm oil. Dose, 1 oz. to 1½.—YOUATT.

5. B. aloes 24 dr., Cape aloes 12 dr., olive oil 4 dr.; treacle 12 dr.; dose, 7 to 14 dr.; mix as No. 1.—MORTON.

6. B. aloes 5 dr., 7½ dr., or 9 dr., oil of caraway 10 drops; made up with palm oil or lard.—Mr. BLAINE'S Nos. 1, 2, and 3.

7. Melt B. aloes (in a tin vessel immersed in boiling water) with a fifth of its weight of treacle, and while soft, pour it into paper moulds; 1 oz. is a full dose for a large-sized saddle or coach horse.—B. CLARK. [For a convenient apparatus for melting and casting these balls, see Mr. Bracy Clark's Pharmacopœia Equina; or Vol. V. of the Pharmaceutical Journal.]

8. B. aloes 5 to 8 dr., cream of tartar 2 dr., oil of cloves 10 drops, treacle to form a ball.—PEALL.

9. Aloes 7 dr., Castile soap 4 dr., aromatic powder 1 dr., oil of caraway 6 drops; mucilage to form a ball.—HINDS.

10. B. aloes 7½ parts, Socotrine aloes 7½ parts, ginger

1 part; mix the powders, add 7½ parts of palm oil and beat to a mass; keep it in a jar closely covered; dose, 1¼ oz. to 1¾ oz.—CLATER.

11. B. aloes 13½ oz., lard 6 oz., treacle 1½ oz., water 1½ oz.; put them in an earthen vessel, placed in boiling water; mix, and form the mass into 18 balls.—MCEWEN.

12. Aloes and hard soap each 5 oz., pearl ashes 1 oz. powdered ginger 1 oz. Melt in a ladle and divide, while warm, into 8 balls.

Mercurial Physic Balls. 1. Cathartic mass (No. 10 above) 10 to 14 dr., calomel 1 dr. to 1½ dr.; mix.—CLATER.

2. For stomach staggers: aloes 1 oz., calomel ½ oz., cascarilla 3 dr., syrup to form a ball.—WHITE.

LAXATIVE BALLS.

1. Ipecacuanha 1 dr., aloes 3 to 4 dr., liquorice powder and mucilage to form a ball.—HINDS.

2. Aloes 3 to 4 dr., soap 3 dr., oil of caraway 20 drops, syrup q. s.—WHITE.

3. Aloes 3 to 4 dr., soap 4 dr., emetic tartar 2 dr., mucilage to form a ball.—HINDS.

For other formulæ, see Alterative Balls (laxative).

NAUSEATING BALLS. These are given in inflammatory diseases.

1. Powdered white hellebore ½ dr., linseed meal 4 dr., treacle to form a ball; one night and morning till some effect is produced; in inflammation of the kidneys.—CLATER.

2. White hellebore 20 gr., common mass or other proper material to form a ball; give one every 4, 6, or 8 hours, till symptoms of nausea appear, taking care not to carry it too far.—PERCIVAL. See Fever Balls. See Hellebore, in Veterinary Materia Medica.

STOMACHIC BALLS. For indigestion and during recovery from debilitating diseases which have impaired the appetite. A mild purge should be previously given.

1. Gentian, quassia, grains of paradise, of each 3 dr., Venice turpentine q. s. for 1 ball.—BLAINE.

2. Gentian 2 or 3 dr., carbonate of soda 1 dr., ginger 1 dr., treacle to form a ball.—WHITE.

3. Chamomile 2 dr., calumba 2 dr., common salt 1 dr., fenugreek 2 dr., syrup to form a ball.

4. Myrrh 1½ dr., cascarilla 2 dr., Castile soap, 1 dr., syrup to form a ball.—WHITE.

5. *Laxative Stomachic Ball.* Aloes 3 dr., rhubarb 3 dr., subcarbonate of soda, 2 dr., ginger, 1½ dr., treacle to form a ball.—WHITE.

6. Calumba and chamomile in powder, each 2 dr., Venice treacle ½ oz., oil of caraway 25 drops, honey q. s.—LAWRENCE. See Tonic Balls, for other formulæ.

TONIC BALLS. In diseases attended with general debility, and to restore strength after a tedious illness.

Vegetable Tonics. 1. Peruvian bark 1 oz., opium ½ dr., ginger 1½ dr., oil of caraway 20 drops, treacle to form a ball.—WHITE.

2. Sulphate of quinine 1 dr., gentian, oak bark, and honey to form a ball.—MOIROUD.

3. Gentian 1 dr., ginger ½ dr., cascarilla 1 dr., treacle and linseed meal to form a ball.—CLATER.

4. Myrrh 2 dr., mustard flour 1 dr., cantharides 5 gr., chamomile 4 dr., Venice turpentine q. s.. for one ball.—BLAINE.

5. Gentian 4 dr., chamomile 2 dr., carbonate of iron 1 dr., ginger 1 dr., syrup q. s. for one ball.—YOUATT.

6. Quassia 2 dr., canella 2 dr., opium ½ dr., ginger 1 dr., treacle q. s.—WHITE.

Mineral Tonics. 1. Sulphate of iron 4 oz., ginger 4 oz., common mass 10 oz.; beat together to form a mass; dose, 1 oz. to 1½ oz.—V. C.

2. Sulphate of iron ½ oz., aromatic powder 2 dr., mucilage q. s. to form a ball.—WHITE.

3. Scales of iron 12 oz., gentian 8 oz., honey to form a mass.—MOIROUD.

4. Myrrh 3 dr., sulphate of iron 2 dr., chamomile 3 dr., ginger 1 dr., Venice turpentine or palm oil to form a ball.—BLAINE.

5. Gentian 4 dr., chamomile 2 dr., carbonate of iron 1 dr., ginger 1 dr., syrup for one ball.—YOUATT.

6. Sulphate of iron 2 dr., carbonate of potash 2 dr., cascarilla 2 dr., caraway 4 dr., treacle q. s.—WHITE.

7. Sulphate of iron 1 dr., carbonate of soda 2 dr., myrrh 1 dr., ginger 1 dr., cantharides 6 gr., caraway ½ oz., treacle q. s.—WHITE.

8. Tonic mass. Sulphate of copper 2 oz., ginger 2 oz.,

common mass 12 oz., beat together; dose 1 oz. to1½ oz.—V. C.

9. Sulphate of copper and ginger, of each 1 dr., canella 4 dr., conserve of roses q. s. for one ball.—BLAINE.

10. White arsenic 5 to 10 gr., aniseed ½ oz., opium ½ dr., treacle, q. s.; sometimes 2 dr. of sulphate of zinc may be added.—WHITE.

11. Arsenic 10 gr., gentian and cascarilla, of each 3 dr., conserve of roses q. s.—BLAINE.

Mild Alterative Tonics. To promote condition; a mild dose of physic should be previously given.

1. Aloes, 1 dr., Winter bark 2 dr., verdigris, 1 dr., treacle or honey q. s.

2. Arsenic 8 gr., pimento 1 dr., extract of gentian 4 dr.; daily.

3. Nitre 1 oz., sulphur 6 dr., physic mass ½ oz., gentian 6 dr., ginger ½ oz., palm oil q. s. for 4 balls. One daily, after an attack of stomach staggers.—CLATER.

Tonic Condition Ball. Ginger and camphor, of each 1 dr., gentian and sulphate of iron, of each 2 dr.; make up with linseed meal.—SPOONER.

WORM BALLS.

1. Calomel 1 or 2 dr. at night, and an aloetic ball in the morning.—CLATER.

2. Emetic tartar 2 dr., ginger a scruple, linseed meal and treacle to form a ball; one every morning an hour before feeding.—YOUATT.

3. Calomel 8 gr., arsenic 8 gr., tin filings 1 oz., Venice turpentine ½ oz.; mix; and give every morning fasting for a fortnight.—BLAINE.

4. Common salt ½ oz., gentian 2 dr., rust of iron 2 dr., savin 1 dr., treacle to form a ball, to be given every morning for a week; then a purging ball.

5. B. aloes 6 dr., ginger 1½ dr., oil of wormwood 20 drops, subcarbonate of soda 2 dr., syrup to form a ball; ½ dr. or 1 dr., of calomel may be added, or given the previous night; to be repeated at intervals of ten days if required.—WHITE.

6. Emetic tartar 2 dr., common mass 6 dr.; to be given for six mornings, and a purging ball on the seventh.

7. Assafœtida 2 dr., calomel 1 or 2 dr., savin 1½ dr., oil of wormwood 20 drops, syrup q. s.; at night, and a physic ball in the morning.

8. Emetic tartar 1 dr., sulphur 1 dr., linseed meal 4 dr., palm oil to form a ball; one every morning after a mercurial physic ball.—CLATER.

9. For long round worms. Emetic tartar 2 dr., ginger ½ dr., tin filings 6 dr., linseed meal 1 dr., palm oil to form a ball.

10. Assafœtida 4 oz., gentian 2 oz., strong mercurial ointment 1 oz., honey to form a mass, for 16 balls; one or more every morning.—LEBAS.

MISCELLANEOUS BALLS.

Garlic Ball. Beat garlic to a paste with enough linseed or liquorice to form a mass; dose, 10 dr.

Camphor Ball. Mix into a ball 2 dr. of camphor with liquorice powder and syrup enough to give it a proper consistence.

Iodine Ball. Iodine 5 gr., linseed meal 5 dr., palm oil to form a ball.

Ball to prevent Hydrophobia. Skullcap 2 scruples, belladonna 2½ gr., form them into a ball, to be given night and morning; the second week 2 balls, the third week 3 balls, and this continued for 6 weeks.—YOUATT.

Leicester Red Balls. Nitre 1 ℔, resin 1 ℔, common soda 2 oz., Castile soap ½ ℔, ginger 2 oz., oil of juniper 2 dr., cinnabar ½ oz.; dose 1½ oz.

Balls for Appetite. Equal weights of assafœtida, saffron, bayberries, and aloes, made into a mass with extract of gentian; dose, 1 oz.—LEBAS.

Anodyne Ball. Opium ½ dr., to 1 dr., camphor 1 dr., aniseed ½ oz., soft extract of liquorice q. s.—WHITE.

Antispasmodic Ball. Opium 1 dr., powdered belladonna 10 gr., linseed meal 3 dr., palm oil or treacle q. s.; twice or thrice a day, in spasm of the neck of the bladder.—CLATER.

Ball for Roaring. The Cough Ball, No. 12, may be tried; and the compound iodine ointment rubbed on the throttle for some weeks or months.

Stimulating Diaphoretic Ball. Emetic tartar 1½ dr., ginger 2 dr., camphor ½ dr., opium 2 scruples, oil of caraway 15

drops, honey to form a ball; for hide-bound and unhealthy coat, without any other disease.—WHITE.

BARTLETT'S *Perspirative Ball.* Dover's powder 3 dr., camphor 1 dr., treacle q. s.

HIND'S *Sweating Ball.* Emetic tartar 1 dr., assafœtida 1 dr., liquorice powder and syrup to form a ball; repeat in 12 hours if required.

Grease Ball. Liver of antimony 16 oz., salt of tartar 16 oz., gum guaiacum, fenugreek, parsley seed, of each 4 oz., treacle to form a mass; dose, 1½ oz.

Sedative Ball. In slight colic. Assafœtida 4 dr., opium 4 dr., syrup and liquorice powder to form 4 balls.—HINDS.

Cordial and Anodyne Ball. Castile soap 3 dr., camphor 2 dr., ginger 1½ dr., and Venice turpentine 6 dr., into 1 ball.

Stimulating Diuretic Balls. Cantharides 1 dr., aloes 2 dr., strained turpentine 1 oz., honey q. s.; make 4 balls, and roll in elecampane powder.—M. GOHIER, *in Dropsy.*

Stimulating Expectorant Ball. Assafœtida 3 dr., galbanum 1 dr., carbonate of ammonia ½ dr., ginger 1½ dr., honey q. s.—WHITE.

Sedative Aperient Ball. In epidemic catarrh or distemper. Balls for Inflammation of the Lungs (No. 4) 6 dr., physic ball (No. 10) 2 dr.; one at night and another in the morning.—CLATER.

Zinc and Valerian Ball. Oxide of zinc 1 oz., valerian 2 oz., oil of hartshorn 1 oz., soft extract of juniper berries, q. s. to make 4 balls; one, twice a day—ECKEL.

CHEWING BALLS, *or* MASTICATORIES. The ingredients are to be tied in a piece of rag, and fixed by a string so that it may be kept in the mouth and chewed.

1. *Emollient Masticatory.* Marshmallow root, liquorice, gum Arabic, of each (in powder) 1 oz., honey 1 oz., or q. s.—LEBAS.

2. *To promote Appetite.* Assafœtida, liver of antimony, juniper berries, bay-wood, pellitory, made into a mass with verjuice, tied as above.—SOLLEYSELL.

3. Assafœtida, common salt, mastic, galangal, each 1 oz.—LEBAS.

4. Assafœtida 2 oz., salt 1 oz.—BOURGELAT.

5. Angelica ½ oz., assafœtida 1 oz., vinegar 2 dr.—SOLLEYSELL.

6. Flour of mustard ½ oz., sal ammoniac 2 dr., powdered pellitory 1 oz.—MOIROUD.

ELECTUARIES AND CONFECTIONS.

Electuaries are compound medicines in the state of a soft paste. When the paste is hard enough to be formed into balls, the compound resembles ball masses or balls, under which we have placed them. French Pharmaciens often use the term *opiates* as nearly synonymous with electuaries; but we only apply the name (opiates) to compounds containing opium.

OPIATE CONFECTION. (*Veterinary.*) Opium 1½ oz., macerate in a little hot water till soft, and rub it to a paste; then add ginger 3 oz., caraway 6 oz., treacle 1½ ℔; dose, 1½ to 2 oz.—WHITE.

DEMULCENT AND PECTORAL ELECTUARIES.

1. Marshmallow root and liquorice (in powder) of each 2 oz., honey 10 oz., mix; to be given at twice, with a spatula.—MOIROUD.

2. Melt ½ oz. spermaceti with 2 oz. of olive oil, add 6 oz. of honey, and mix in 1½ oz. p. marshmallow root; to be given daily.—MOIROUD.

3. (*With Opium.*) Powdered gum 2 oz., marshmallow 1 oz., extract of opium 2 dr., honey 3 oz.; for 2 doses.

4. *Cough Electuary with Manna.* Manna 2 oz., honey 6 oz.; in the morning; said to have cured acute bronchitis, &c.

5. Powdered liquorice 8 oz., elecampane 4 oz., sulphur 2 oz., honey of squill 32 oz.; mix; for 8 doses.

STIMULANT AND CORDIAL ELECTUARIES.

[M. LEBAS gives a form for an electuary (*Thériaque*) of many ingredients, the first of which (cordial powder) itself contains 26 different substances. We only insert here the simple formulæ of the French veterinarians.]

1. Powdered angelica root 2 oz., masterwort 1 oz., muriate of ammonia ½ oz., honey 8 oz.—MOIROUD.

2. *Stimulant and Expectorant.* Assafœtida 4 oz., elecampane 8 oz., honey 32 oz., for 6 doses.—MOIROUD.

3. Powdered cassia and ginger, each 1 oz., honey 6 ounces. —MOIROUD.

TONIC AND ASTRINGENT ELECTUARIES.

1. Red oxide of iron 8 oz., gentian 12 oz., extract of juniper berries 32 oz. MOIROUD prescribes 6 oz. for a horse, or 1 oz. for a sheep; but these are larger doses than are customary in England.

2. Peruvian bark 6 oz., nitre 1 oz., camphor ½ oz., honey 16 oz.—LEBAS.

3. Powdered bistort 1 oz., calcined magnesia 4 dr., honey 4 oz.—MOIROUD.

PURGATIVE AND LAXATIVE ELECTUARIES.

Aloetic compounds are usually made stiff enough to form into balls. See Physic Balls.

1. Oil of croton 20 drops, powdered senna 4 dr., honey q. s.—MOIROUD.

2. Sulphate of magnesia 4 oz., honey 16 oz., bran a quart; infuse the bran in sufficient hot water, and add the salt and honey; twice a day till the bowels are relaxed.—BOURGELAT.

3. Sulphate of soda or magnesia 5 oz., manna 4 oz., bran 1 quart; as the last.—MOIROUD.

DIURETIC ELECTUARIES.

1. Acetate of potash 2 oz., oxymel of squills 4 oz.; oatmeal or flour, to give a soft consistence.—MOIROUD.

2. Nitre 1 oz., camphor 2 dr. (rubbed with yolks of 2 eggs), oxymel 4 oz.; flour or liquorice powder, to give a suitable consistence.—MOIROUD.

DIAPHORETIC ELECTUARIES.

1. Sulphur 1 oz., powdered angelica 1½ oz., honey 5 oz. —MOIROUD.

2. Prepared antimony 1½ oz., elecampane 2 oz., treacle 4 oz.

3. Kermes mineral 1 oz., powdered sassafras and elecampane, each 2 dr., honey 6 oz.

VETERINARY POWDERS.

MR. B. CLARK'S PULVIS UTILIS, *as a vehicle for other powders.* Turmeric ½ lb, oatmeal or sifted barley-meal 4 lb, mix.

AROMATIC POWDER, OR HORSE SPICE.

1. White's *Aromatic Powder.* Caraway 6 oz., pimento 4 oz., ginger 2 oz., liquorice 2 oz.; mix; dose, 6 to 8 dr.

2. *Common Horse Spice.* Caraway, aniseed, coriander seeds, of each 16 oz., turmeric 32 oz., cumin seeds, liquorice, and ginger, of each 8 oz.; mix.

3. This is inserted, not as a desirable form, but as a specimen of what is used in the trade: Cayenne 2 oz., bean flour 45 lb, mustard hulls 45 lb, cumin seed 15 lb, caraway 15 lb, turmeric 9 lb, bay-berries 3 lb, ivory-black 1 lb.—Gray's Supplement. The cordial powder of Lebas contains 26 ingredients.

ABSORBENT POWDERS.

1. Carbonate of soda 2 to 4 dr., ginger 1 dr., calumba 2 to 4 dr.—White.

2. Prepared chalk 4 dr., gentian 2 to 4 dr., aromatic powder (above) 1 or 2 dr.

ALTERATIVE, DIURETIC, AND DIAPHORETIC POWDERS.

For swelled legs, grease, foul humors, hide-bound, mange, surfeit, old coughs, and to render the skin fine. They are usually given with moistened corn. Too free use of these powders may prove injurious.

1. Sulphur 2 parts, black antimony 1, nitre 1; mix; dose ½ oz. to 1 oz.—V. C.

2. Sulphur 4 dr., levigated antimony 2 dr., nitre 3 dr.; mix; in *hide-bound* and unthrifty coat, every night.—Youatt.

3. Ethiops mineral ½ oz., cream of tartar 1 oz.; mix; give every night in a mash; *for grease.*—Blaine.

4. Sulphur 12 oz., antimony (black) 12 oz.; mix, and divide into 24 powders; *for mange*, &c.—Taplin.

5. Nitre 16 oz., resin 16 oz., prepared antimony 4 oz., flowers of sulphur 24 oz.; mix; dose, 1 oz. every evening, with moistened corn, for 6 or 8 times.

6. Equal weights of antimony, nitre, and cream of tartar; dose, 6 to 9 dr.—Blaine.

7. Nitre 6 oz., vermilion ½ oz., resin 6 oz., tartarized antimony 2 dr.: for 12 doses.

8. Sulphur ½ oz., prepared antimony 1 dr.; once a day, in the food, for 10 or 14 days.—Clater.

9. Taplin's *Alterative Powders.* Levigated antimony 8 oz., sulphur 8 oz., Ethiops mineral 4 oz., cream of tartar 4 oz.; in 12 doses.

10. Cream of tartar 2 dr., nitre 2 dr., sulphur 4 dr.; for one dose.—BLAINE.

11. Nitre 1 oz., resin 1 oz., rust of iron 1 oz., emetic tartar 15 gr.; dose, 1 oz.—LEBAS.

12. *In Farcy.* Prepared antimony 12 oz., sulphur 12 oz., cream of tartar 8 oz., cinnabar 6 oz.; mix and divide into 20 doses; one every night, in corn.—TAPLIN.

CONDITION POWDERS. A want of condition is generally indicated by, and connected with, the unthrifty state of the coat, which the above (alterative) powders are supposed to remedy. Sometimes warm and bitter tonics are added to those ingredients which promote the action of the skin and kidneys, to increase the appetite and promote nutrition; but the most scientific practitioners condemn these additions; and particularly when the animal is changing its coat.

1. Black antimony 4 oz., flowers of sulphur 2 oz., bean flour or barley-meal ½ lb; a tablespoonful with corn.—B. CLARK.

2. Sulphur 2 lb, fenugreek 4 lb, cream of tartar 1 lb, liquorice 1 lb, nitre 1 lb, black antimony ½ lb, gentian ¼ lb, aniseed ¼ lb, common salt 1 lb; dose, 1 oz., daily for 2 or 3 weeks.

3. Gentian 4 oz., liquorice 4 oz., fenugreek 16 oz., diapente 6 oz., nitre 4 oz., salt 4 oz.; to promote appetite.

4. Aromatic powder 2 oz., assafœtida ¼ oz., cream of tartar ¾ oz., crocus metallorum ½ oz.; for 2 doses.—LEBAS.

DIAPENTE. This should be made with equal parts of myrrh, gentian, ivory-dust, bay-berries, and birthwort. But a worthless compound is commonly sold for it. The following is one of the least objectionable substitutes: Equal parts of gentian, turmeric, bay-berries, and mustard. Another form in use is—Bay-berries 2½ lb, guaiacum wood 2 lb, gentian 14 lb, bole 2 lb, bark which has been used for the tincture 2 lb.

Fever Powders. 1. Nitre 1 oz., camphor 2 dr., tartarized antimony 2 dr.—WHITE.

2. Nitre 6 dr., camphor 2 dr., calx of antimony 1½ dr.—HINDS.

3. Nitre 1 oz., unwashed calx of antimony 2 dr., antimonial powder 3 dr., camphor 1 dr.—WHITE.

Pectoral Powder. Powder of gum tragacanth 6 oz., nitre 1 oz.; give a tablespoonful in their mashes or food, in coughs.

Purgative Powder. Epsom salt 8 oz., aloes 10 oz., aniseed 3 oz.; dose, 2 oz.—LEBAS.

Powder for the Gripes. Aloes, senna, ginger, cream of tartar, of each 1 ℔; mix. This was formerly honored with the title of Pulvis Sanctus.

Worm Powders. 1. Sulphur 12 parts, quicksilver 4 parts; triturate together till the mercury is extinct; then add male fern, rhubarb, tansy, gentian, of each 4 parts, wormwood, savin, aloes, castor seeds, of each 1 part; dose, 1½ oz. to 2 oz.—LEBAS.

2. Fern roots 4 parts, tansy 2, assafœtida and aloes, each 1 part; dose, as the last.—MOIROUD.

3. Sulphur 1 oz., emetic tartar 4 dr., common salt 8 oz., liver of antimony 1 oz.; mix; for 6 doses; one daily in wetted corn.—HINDS.

MR. WHITE'S *Compound Arsenical Powder.* White arsenic 1 dr., cream of tartar 9 dr.; mix carefully; give 10 gr. 3 times a day.

HAYNE'S *Bitter Powder*, for loss of appetite. Sulphate of potash 2 oz., gentian 1 oz., flour q. s. To be given twice a day.

MEDICATED PROVENDER. Bruised oats 4 ℔, bruised juniper berries 2 oz., common salt 1 oz.; mix. Nourishing and stimulant.—DELAFOND.

Liquid Medicines for Horses.

DRINKS, DRENCHES, MIXTURES, MASHES,

ETC.

Drinks, properly speaking, are liquids which the horse will take willingly; *Drenches* are those liquid medicines which must be administered by a horn, bottle, or funnel. But this distinction is not always observed.

MILD DRINKS. Demulcent, pectoral, cooling, and diuretic.

Barley Water. Barley 1 ℔, water 2 gallons; boil to 6 quarts, strain, and add 1 ℔ of honey. If common barley is used, it should be first boiled with a little water, and this thrown away. If pearl barley is used, this will be less necessary. In inflammatory and catarrhal complaints.

Oatmeal Gruel. 1. Mix gradually 4 oz. of sweet oatmeal with as much cold water as will form a smooth mixture. Put 2 quarts of water in a saucepan over a clear fire, and before it gets very hot add in the mixture of oatmeal and water; stir the whole till it boils, and let it simmer a little while. Take care not to smoke it.

2. Mix half a pint of oatmeal with the same measure of water; triturate them in a marble mortar with a wooden pestle, for some time; then add 1 gallon of boiling water, and boil for a few minutes.—B. CLARK.

Blanche Water. Wet 3 or 4 handfuls of bran with scalding water, and work it with the hands till it becomes clammy; then add as much more water as may be desired. A mixture of oatmeal and cold water is also called white water, and in France potato or other starch is used for the same purpose.

Linseed Tea. 1. Infuse 4 oz. of linseed in 3 pints of boiling water for several hours near the fire, stirring occasionally; then strain off, and add 4 oz. of honey; for 2 doses; in coughs, &c.

2. Pour 1 gallon of boiling water on ½ lb of linseed; let the infusion stand till nearly cold, then pour off the clear liquid.—YOUATT.

Compound Decoction of Linseed. Linseed 4 oz., liquorice root 4 oz., mallows 2 handfuls; boil in six quarts of water for half an hour. Let the horse drink it freely.—BLAINE.

Cooling and Refreshing Drink. Barley water, linseed tea, or blanche water, 8 quarts, simple oxymel 16 oz.—MOIROUD.

Cooling and Diuretic Drink. Dissolve 1 oz. of nitre in a pail of water.

Camphorated Diuretic Drink. Water 10 quarts, nitre 1 oz., camphor (rubbed with yolks of 2 or 3 eggs) ½ oz.; mix, and let the animal drink when thirsty.—MOIROUD.

MASHES.

Bran Mash. Bran or pollard ½ peck; put it in a bucket, and pour on it enough scalding water to wet it thoroughly; let it be well stirred with a stick, or worked with the hands, and let it stand, covered up, till new-milk warm. Emollient and slightly laxative. When intended to be nutritive, oats should be scalded with the bran.—B. CLARK.

Malt Mash. Upon a peck of ground malt pour a gallon and a half of boiling [better not quite boiling] water. Stir frequently, and give when new-milk warm. Nutritive, in diseases attended with great debility.—MARKHAM.

Linseed Mash. HINDS' *Cooling Decoction.* Linseed 2 quarts, coarse sugar 2 oz., boiling water 6 quarts; simmer for three or four hours.

DRENCHES.

DRENCHES FOR DIARRHŒA, DYSENTERY, AND DIABETES.

For Diarrhœa. 1. Restringent Draught. Opium 1 dr., prepared chalk 1 oz., compound powder of tragacanth 1 oz., mint water 1 pint.—WHITE.

2. Prepared chalk 8 oz., gum Arabic ½ oz., catechu 2 dr., thin starch ½ pint.—BLAINE.

3. Prepared chalk 1 oz., catechu 2 dr., p. opium 1 dr., p. ginger 1 dr.; rub together with contents of one egg, and add ½ pint of thin gruel.—CLATER.

4. For purging from corrosive sublimate. Powdered opium 2 dr.; rub down with the yolk and white of one egg, and the contents of two more eggs, and gradually stir in ½ pint of thin gruel.—CLATER.

Draught for Enteritis Opium 1½ dr., tartar emetic 1 dr., spirit of nitrous ether 1 oz.; mix, and add 1½ pint of linseed oil.—SPOONER.

For Dysentery or Molton Grease. 1. Castor oil 8 oz., ipecacuanha 1 dr., opium 20 gr., liquid arrowroot 8 oz. Repeat once or twice at intervals of 6 hours: then substitute boiled starch for the castor oil.—BLAINE.

2. Opium 2 dr., nux vomica ½ dr., ipecacuanha 1 dr., red wine 1 quart; mix; morning and evening.

For Diabetes. 1. Opium 1 dr., ginger 2 dr., p. oak bark 1 oz., decoction of oak bark 1 pint.—WHITE.

2. Sulphuret of potash 2 dr., uva ursi 4 dr., oak bark 1 oz., catechu 2 dr., opium ½ dr. In strong chamomile tea.—BLAINE.

3. Calomel 3 dr., cascarilla 2 dr., salt of steel 2½ dr., salt of tartar 1½ dr., tincture of opium ½ oz., strong beer q. s.—WHITE.

CARMINATIVE AND ANTISPASMODIC DRENCHES.

For Spasmodic and Flatulent Colic, or Gripes.

[N. B. As most of these drenches would be injurious in *inflammation of the bowels* (Enteritis), care should be taken to distinguish between these diseases. Inflammation is known by the quick but small pulse, redness of the inside of the eyelids, coldness of the ears and legs, and scanty and high-colored urine. In colic, the attacks and remissions of pain alternate; in inflammation, the pain and distress continue. In colic, the pain is relieved by friction and motion; in inflammation, it is increased. Colic is sudden in its attack; inflammation, more gradual in its approach.]

1. Brandy, rum, or gin, from 4 to 6 oz., hot water 12 oz. Mr. CLARK directs a wineglassful of spirits to half a pint of warm water. A pint of ale is sometimes substituted.—WHITE.

2. Half a large bottle of Daffy's elixir, with hot water.

3. Tincture of pimento 4 ounces, warm water half a pint. —B. CLARK.

4. Anodyne carminative tincture (WHITE'S, see below) 2 to 4 oz., hot water half a pint.—WHITE.

5. *Antispasmodic Draught.* Spirit of nitric ether 2 oz., tincture of opium 1 oz., solution of aloes (see below) 4 oz.—V. C.

6. Spirit of nitric ether ½ oz., tincture of opium ½ oz., oil of turpentine 3 oz., gruel 1 pint.—BLAINE.

7. Rectified oil of turpentine 3 oz., tincture of opium 1 oz., warm ale 1 pint. If it does not relieve, repeat half the quantity with 1 oz. aloes dissolved in warm water.—YOUATT.

8. Stronger ether 1 oz., laudanum 2 oz., oil of peppermint 1 dr., ale and gin, each a ¼ of a pint.—BLAINE.

9. Camphor 2 dr., tincture of opium 1 oz., oil of peppermint 30 drops, warm water 1 pint. In a violent attack, add 1 oz. of spirit of turpentine.—PEALL.

10. The juice of three or four onions, with half a pint of sound ale.

11. Pepper ½ oz., oil of turpentine 3 oz., laudanum 1 oz., ale ¼ pint.—BLAINE.

12. Pepper a teaspoonful, juice of 2 or 3 large onions, gin ¼ of a pint.—BLAINE.

13. Laudanum 1 oz., sweet spirit of nitre 4 oz., oil of juniper 1 oz., tincture of benzoin 2 oz., spirit of sal volatile 1½ oz., oil of peppermint 1 dr.; mix; give a fourth part in warm water or gruel, and repeat in 3 or 4 hours, if necessary.—HINDS.

14. *For Flatulent Colic.* Tincture of opium 1 oz., of myrrh 1 oz., sulphuric ether 6 dr., tepid water a pint. Repeat in an hour, if relief is not obtained.—SPOONER.

15. Heat ½ lb. of common salt, and quench it in a quart of good ale. Give it new-milk warm.—DOWNING.

16. In flatulent colic when there is an evident distension of the abdomen with gas: chloride of lime ½ oz. (or solution of chlorinated soda 1 oz.), water 1 quart; repeat in half an hour if necessary.

17. Ginger, caraway, nutmeg, pimento, of each 1 oz., bruise, and boil them in ¾ of a pint of ale for a few minutes, and add a gill of any spirits.—TAPLIN.

Cordial Antispasmodic Drink, for spasm of the Diaphragm. Ginger 1 dr., caraway 2 dr., laudanum 1 oz., sweet spirit of nitre 1 oz., warm ale ½ pint.—CLATER.

Antispasmodic Drench for Suppression of Urine. Nitre 1 oz., camphor 2 dr., linseed tea 1 pint.—WHITE.

Diuretic Camphor Drink. Camphor 2 to 3 dr., olive oil 1 oz., carbonate of soda 1 dr.; rub together, and add tincture of opium 1 dr., warm water 2 pints.

Antispasmodic Drenches for Locked Jaw. 1. Opium 1½ dr., camphor 2 dr., ginger 3 dr., brandy-and-water 8 oz.—WHITE.

2. Ether ½ oz., tincture of opium 2 oz., camphor 1 dr., peppermint water ½ pint.

CORDIAL AND STIMULANT DRENCHES. These are used in the same cases as the cordial balls, but are preferred where a more quick and powerful operation is required. Some of them are used in indigestion and slight attacks of colic.

1. Cloves and black pepper (bruised) ½ oz., boiling water a quart; infuse and give warm.—MOIROUD.

2. Any of the cordial balls may be dissolved in warm ale or water, or peppermint water, and given as a drench.

3. A bottle of wine, 1 oz. of extract of juniper berries, and ½ oz. of cinnamon in powder.—M. LEBAS.

4. Peppermint 2 oz., chamomiles ½ oz.; infuse in 2½ pints of water, and give it before it is cold; in slight colic and indigestion.

PECTORAL AND EMOLLIENT (or DEMULCENT) DRENCHES, for Coughs, Epidemic Catarrh, &c. (For linseed tea, compound infusion of linseed, barley water, &c., see DRINKS, above.)

1. *Simple Emulsion.* Olive oil 2 oz., honey 3 oz., soft water 1 pint, subcarbonate of potash 2 dr.; mix.—WHITE.

2. Linseed tea 1 pint, honey 2 oz., syrup of poppies 2 oz., linseed oil 4 oz.

3. B. CLARK'S *Cough Drench.* Linseed oil 2 oz., liquor of potash 40 drops, treacle 1 oz., soft water 10 oz.; mix.

4. Powdered gum 2 oz., warm water a quart; dissolve, and add honey 4 oz.—MOIROUD.

5. Marshmallow root 2 oz., water 2½ or 3 pints; boil to a quart, and add 4 oz. of treacle.—MOIROUD.

6. Liquorice and marshmallow roots, of each 2 oz., water a quart; boil, strain, and add honey 4 oz.—LEBAS.

7. Marshmallow root 2 oz., poppy-heads 4, water a quart; boil for 10 minutes, strain and add to the liquor before quite cold 4 oz. of olive oil, 6 oz. of honey, and the yolks of 4 eggs, previously well beaten together.—MOIROUD.

8. Compound decoction of linseed (page 80) 1 quart, oxymel 3 oz.

9. Spermaceti ½ oz., olive oil 3 oz.; melt together, and add of honey 4 oz., water (by a little at a time) to make up a quart; repeat it twice a day.—LEBAS.

10. *Camphorated Emulsion.* Powder, with a few drops of spirit, 1 or 2 dr., of camphor, and 12 drops of oil of aniseed, and 12 oz. of simple emulsion.—WHITE.

11. Oxymel of squills 2 oz., opium ½ dr. to 1 dr., linseed oil 2 oz.; mix the opium with 8 oz. of water, and add the others; for one dose.—WHITE.

12. For chronic coughs: Tar-water ½ pint, lime-water ½ pint, powdered squill 1 dr.; every morning.—BLAINE.

13. In inflammation of the lungs, or catarrhal fever: Tartarized antimony 2 dr., digitalis, 1½ dr., nitre 3 dr., simple oxymel 4 oz., compound decoction of linseed 8 oz.—BLAINE.

14. The same, omitting the digitalis, and substituting 6 oz. of warm water for the dec. linseed. In influenza, when soreness of throat prevents swallowing balls.—BLAINE.

15. In inflammation of the lungs: Ipecacuanha, 2 dr., laudanum 4 dr., powdered camphor 2 dr., Mindererus' spirit 4 oz., linseed tea ½ pint.—BLAINE.

16. In pleurisy. Boil pearl barley, split figs, and raisins, each 6 oz., and liquorice root 2 oz., in 4 quarts of water down to 3; strain and add honey 1 ℔, vinegar 1 pint; give 1 oz. nitre in a pint of this decoction every 8 hours.—TAPLIN.

17. In epidemic (epizootic) catarrh: Spirit of nitrous ether 1 oz., Mindererus' spirit 6 oz., with linseed tea.—BLAINE.

18. GIBSON'S *Drink for Catarrhal Epidemic.* Coltsfoot, hyssop, chamomile, of each a handful, linseed and garlic, each 1 oz., liquorice root sliced 1 oz., saffron ½ oz.; infuse in 2 quarts of boiling water; give half in the morning and the rest in the afternoon.

19. In influenza (after bleeding). Oil of croton 5 drops, nitre 4 to 6 dr., tartarized antimony 1 dr., spirit of nitric

ether ½ oz. to 1 oz., solution of acetate of ammonia (L. P.) 2 to 4 oz., warm water, q. s. Once or twice daily. Sometimes ½ oz. of cream of tartar is added.—SPOONER.

20. For malignant epidemic. Oxymel 4 oz., spirit of Mindererus 4 oz., beer yeast 4 oz., sweet spirit of nitre 1 oz.

DIURETIC DRENCHES, for Dropsical Complaints, &c. The use of stimulating diuretics in retention of urine from inflammation of the neck of the bladder is dangerous.

1. MARKHAM'S *Dropsy Drench.* Decoction of wormwood in ale 2 quarts, soap 1 oz., grains of paradise 6 dr., long pepper 6 dr., treacle 3 oz.; for one dose, fasting.

2. For dropsy of the belly: Castile soap 2 oz., strong beer 1 pint; dissolve and add cascarilla 2 dr., ginger 3 dr., oil of juniper 2 dr. (or balsam of copaivi 1 oz.); mix; for one dose.—WHITE.

3. White soap 1 oz., spirit of turpentine 1 oz., honey 4 oz., decoction of linseed 2 quarts; for two doses.—MOIROUD.

4. Strained turpentine 2 oz., yolks of 6 eggs; triturate together till incorporated, and add gradually 2 quarts of linseed tea; for 2 doses.—MOIROUD.

5. White wine and water 4 quarts, nitre 3 oz., honey 4 oz.; for 3 doses.—LEBAS.

6. Acetate of potash 2 or 3 oz., honey 6 oz., decoction of hemp or linseed 2 quarts; for 1 dose.—MOIROUD.

7. Acetate of potash 2 oz., camphor (rubbed with yolks of 2 eggs) 2 dr., decoction of linseed 2 quarts; for 2 doses, at an interval of some hours; an irritation of the urinary passages, especially arising from cantharides or resinous irritants.—MOIROUD.

8. *Squill Drench.* Decoction of pellitory of the wall 1 quart, oxymel of squills 4 oz.—MOIROUD.

9. *Colchicum Drench.* Colchicum wine 2 oz., simple oxymel 4 oz., barley water 1 quart.

10. Sweet spirit of nitre 4 oz., white wine 1 quart, water 2 quarts; for 3 doses; in dysuria not arising from mechanical obstruction or inflammation of the neck of the bladder.—LEBAS.

11. *Saline Diuretic Drink.* Glauber's salt 2 oz., nitre 6 dr., warm water 1 pint, sweet spirit of nitre 1 dr.—CLARK.

DRENCHES FOR FARCY AND GLANDERS.

1. Expressed juice of cleavers 6 oz., strong decoction of hempseed 6 oz., essence of spruce 6 oz.; mix; give every evening; and a mercurial or arsenical ball in the morning.—BLAINE. See Farcy Balls.

2. *For Glanders.* Sulphate of copper 3 to 6 dr., gum Arabic 2 or 3 oz., dissolved in 2 or 3 pints of water.—SEWELL.

FEVER DRENCHES.

1. Nitre 2 dr., tartar-emetic ½ dr., warm water or thin gruel 12 oz.; once or twice a day.—B. CLARK.

2. Sweet spirit of nitre 1 oz., spirit of Mindererus 6 oz., water 4 oz.

3. *Draught for Influenza.* Spirit of nitrous ether 2 oz., tartarized antimony 1 dr., nitre 4 dr., warm water ½ pint.—SPOONER.

LAXATIVE AND PURGATIVE DRENCHES.

1. Castor oil 6 oz., linseed oil 8 oz., gruel q. s.—BLAINE.

2. Glauber's or Epsom salts 6 or 8 oz., whey or gruel 1 quart, castor oil 6 or 8 oz.—WHITE.

3. Barbadoes aloes 2 dr., tartarized antimony 1 dr., warm water 4 oz.; mix, and add castor oil 4 oz.—WHITE.

4. *Laxative Febrifuge in Influenza.* Linseed oil 12 oz., nitre 3 dr., camphor powdered 1 dr., sweet spirit of nitre 1 oz., warm water ½ pint—CLATER.

5. *Laxative Anodyne Drink.* In inflammation of the bowels: Linseed oil 1 pint, opium 2 scruples, sweet spirit of nitre 6 dr., warm water 4 oz.—CLATER.

6. Aloes 2 or 3 dr., salt of tartar 1 dr., water or mint-water ½ pint; mix, and add castor oil 4 to 6 oz.—WHITE.

7. *A Cooling Purging Drink.* Infuse 2 oz. senna with 3 dr. salt of tartar in a quart of boiling water for 2 hours; strain, and add 4 oz. Glauber's salts, and 2 or 3 dr. of cream of tartar.—BARTLETT.

8. *Draught for Hepatitis.* Carbonate of potash, and aloes, each 2 dr.; dissolve in hot water, shake with 12 oz. of linseed oil, and calomel 1 dr. Repeat twice a day without the aloes, until relief. When pain, add 1 oz. of tincture of opium.—SPOONER.

STRONGER PURGATIVE DRENCHES.

1. B. aloes 2 oz.; gum Arabic 1 oz.; powder and mix

them, and pour on them a pint of boiling water. Take 10 gr. of farina of croton, and add to it gradually 4 oz. of the above solution. Repeat this dose every 6 hours till it operates; in inflammation of the brain.—CLATER.

2. Aloes 1 oz., soap 2 dr., salt of tartar 1 dr., water 1 pint; in apoplexy or staggers.—WHITE.

3. Infuse 1 oz. of senna in a quart of boiling water, strain, and add 1 oz. of aloes in powder.—BOURGELAT.

4. Aloes 1 oz., sulphate of magnesia 2 oz., aniseed powder ½ oz., water a quart.—LEBAS.

5. Aloes 1 oz., syrup of buckthorn 4 oz., warm water a quart.—LEBAS.

DRENCHES FOR STOMACH STAGGERS, or Staggers from Indigestion.

1. After a ball of aloes and calomel, and clyster of salt water: Spirit of sal volatile ½ oz., cascarilla powder 2 dr., warm water ½ pint; twice a day; and the same without the cascarilla every hour.—WHITE.

2. Aloes 3 dr., pimento 2 dr., ginger 1 dr.; infuse in a quart of hot water, and when cold add 2 oz. spirit of turpentine, and 1 oz. of spirit of hartshorn. Repeat in an hour if required.—BLAINE.

3. *Laxative Tonic Drinks.* Linseed oil 1 pint, powdered gentian 2 dr.; every 6 hours till the bowels are properly opened.—CLATER.

4. Common salt 4 oz., ginger 2 dr., magnesia 1 oz., warm water 1 quart.—WHITE.

5. Valerian 1 oz., serpentary ½ oz., saffron 2 dr.; infuse in a pint of boiling water, and, when nearly cold, strain off, and add 1 oz. tincture of assafœtida and 2 dr. of laudanum.—TAPLIN.

6. *After a Purgative.* Volatile tincture of valerian 1 oz., powdered valerian 1½ oz., peppermint water 8 oz.; mix, for a dose.—WHITE.

TONIC DRENCHES. Tonics are more generally administered in the form of balls.

1. *Mild Tonic in latter stage of Epidemic Catarrh or Distemper.* Gentian 1 dr., powdered ginger ½ dr., cascarilla 1 dr., warm water ½ pint, sweet spirit of nitre ½ oz. to 1 oz.; to be repeated night and morning unless they quicken the pulse.—CLATER.

2. Gentian root 2 oz., smaller centaury 1 oz., wormwood ½ oz.; boil in 3 pints of water to a quart.—VATEL.

3. CLARK'S *Bitter Drench.* Quassia chips 2 oz., water 3 pints; boil to 2 pints; for 3 doses.

4. Quassia 1 oz., ginger 2 dr., water 2 pints; boil for 10 minutes; for 2 doses.

5. *Metallic Tonic.* Sulphate of zinc ½ dr., ginger or pimento 1 dr., treacle 1 oz.; mix and add gradually 12 oz. of warm water.—B. CLARK.

6. *Egyptian Tonic Drink.* In farcy and nasal gleet. Ægyptiacum ½ oz., pimento or ginger ½ dr., warm water 12 oz.—CLARK.

7. *Cantharides Tonic Drench* (for the same). Sulphate of zinc 15 gr., cantharides 7 gr., pimento 15 gr., treacle 1 oz., warm water to form a drench.

DEOBSTRUENT DRENCHES.

1. Guaiacum wood 2 oz., sassafras 1 oz., linseed ½ oz., water q. s. to yield a quart of decoction; boil, strain, and add of corrosive sublimate 10 gr., sal ammoniac 2 dr.—LEBAS.

2. Iodide of potassium 40 gr., iodine 10 gr., water a quart.—MOIROUD.

3. Muriate of lime 2 oz., water a quart.—MOIROUD.

WORM DRENCHES.

1. Common salt 2 oz., infusion of wormwood a quart. Repeat it for some days.—MOIROUD.

2. A quart of linseed oil.—CLATER.

3. Oil of turpentine 4 oz., linseed or castor oil 8 oz., gruel a pint; preceded by a mild dose of aloes, and bran mashes.

4. Fern root 2 oz., valerian 1 oz., DIPPEL'S animal oil (empyreumatic oil of hartshorn) 1 oz., yolks of 2 eggs, honey 2 oz.; boil the roots in 2 pints of water to half, incorporate the oil with the egg, and then the honey, and mix the whole with the decoction.—VATEL.

5. Animal oil 1 oz., yolks of 2 eggs, honey 1 oz., water or some bitter infusion a quart. CHABERT recommends infusion of savory as a vehicle for the oil.—LEBAS.

6. Soot (wood soot?) in fine powder 2 oz., spirit of wine 2 oz.; mix, and add a quart of infusion of rue, or of tansy. Some practitioners prefer milk as a vehicle for worm medicines.—MOIROUD. For other worm remedies see Worm Balls.

DRENCHES for the MALIGNANT EPIDEMIC, or DISTEMPER—Pestilential or Putrid Fever.

1. Gentian 1 dr., calumba 2 dr., ginger 1 dr., laudanum ½ oz., spirit of nitrous ether ½ oz., peppermint water 3 oz.—CLATER.

2. Gentian 2 oz., willow bark 6 oz., water 3 pints; boil to a quart, and add solution of acetate of ammonia (L. P.) 6 oz.—MOIROUD.

3. Dissolve ½ dr. of chloride of lime in 8 oz. water, and add spirit of nitric ether ½ oz., laudanum ½ oz., tincture of calumba 1 oz.; twice a day.—CLATER.

4. Bruised bark 3 oz., acetate of ammonia (L. P.) 4 oz., camphor 1 dr.; boil the bark in two quarts of water in a covered vessel for a quarter of an hour; strain, and when cool, add the camphor (rubbed with yolk of egg or honey), and the acetate of ammonia.—LEBAS.

5. Spirit of nitric ether 1 oz., Mindererus' spirit 4 oz.. infusion of chamomile 6 oz., beer yeast 6 oz., tincture of opium 3 dr.—BLAINE. See also Antiseptic Drenches (below).

ANTISEPTIC DRENCHES, *to check Mortification.*

1. Peruvian bark 1 oz., ginger 2 dr., opium 1 dr., fresh beer q. s.—WHITE.

2. Opium 1 dr., carbonate of ammonia 1 dr., aromatic powder 2 dr., camphor 1½ dr., good ale or porter a pint.

3. Chloride of lime or soda 2 to 3 dr., serpentary in powder 1 oz., fresh beer or sweet wort 1 quart.—WHITE.

DRENCH FOR POISONING BY YEW. Stronger purgative drink (No. 1) 4 oz., vinegar 4 oz., thick gruel 4 oz.; repeat it every 6 hours, *without the croton*, till purging is produced.—CLATER.

DRENCH FOR PREVENTING HYDROPHOBIA. Box leaves 8 oz., rue 8 oz.; cut them very fine and boil in 3 pints of milk, in a close vessel, for an hour, and strain; boil the ingredients another hour in 3 pints of water, and strain; mix the decoctions; give a third part every morning fasting.—BLAINE.

DRENCHES TO PROMOTE PARTURITION.

1. Ergot of rye in fine powder 2 or 3 dr., pennyroyal water, or infusion of rue, 1 quart.

2. Saffron 6 dr., chamomile 2 oz., boiling water a quart; make an infusion, to be given warm. 1 oz. of dried savin, with 1 oz. of cassia, may be substituted for the saffron.—MOIROUD.

ANODYNE DRENCHES.

1. Opium 1 dr. dissolved in warm water, ½ pint; add 1 quart of starch gruel.

2. Oil of peppermint 50 drops, dissolved in a pint of warm water, with 2 oz. of gum Arabic; add tincture of opium ½ oz.

3. Mix tincture of opium ½ oz., with sweet spirits of nitre 1½ oz., essence of peppermint 1 dr., and water 1 pint.

IODINE DRENCH. Iodide of potassium 2 scruples, iodine 12 gr.; titurate together, and add gradually a quart of water.—MOIROUD.

MISCELLANEOUS LIQUID MEDICINES.

TINCTURES, SOLUTIONS, &c.

Solution of Aloes. Aloes 1 part, water 7 parts, proof spirit 1 part; dissolve the aloes in water by means of a water bath, and when removed add the spirit.—MORTON.

Anodyne Carminative Tincture. Opium 1 oz., cloves 1 oz., ginger 1 oz., old brandy (or rum or gin) 1 quart; digest in a corked bottle, shaking daily.—WHITE.

Ethereal Tincture of Opium. Turkey opium 1 lb, spirit of nitric ether 8 lb; macerate for a month. Dose, ½ oz. to 1 oz., in spasmodic colic.—DICKINS.

Gripe Tincture. Tincture of Pimento. Ground pimento 1 lb, rectified spirit, and soft water, of each 3 pints; digest for some days and strain; give 4 fluid oz. at once, and repeat every hour till relieved.—B. CLARK.

Ticture of Foxglove. Digest 3 oz. of dried foxglove in a quart of any spirit.—YOUATT.

Infusion of Foxglove. Infuse 1 oz. of powdered foxglove in a quart of boiling water till cold.—YOUATT.

Tincture of Myrrh. Myrrh 2 oz., sand 2 oz., rectified spirit and soft water, of each ½ pint.—B. CLARK.

Tincture of Aloes and Myrrh. Aloes 12 oz., myrrh 6 oz., rectified spirit 1 gallon, water ½ gallon; digest 14 days, frequently shaking, and filter. For outward use, rectified wood naphtha may be substituted for the spirit.—V. C.

Tincture for Colic. Opium 1 dr., horseradish 2 oz., capsicum 1 oz., spirit of nitric ether 1 ℔; macerate 14 days; dose, 1 oz., with 2 oz. of spirit of nitric ether, every 2 hours as long as necessary.—GREGORY.

Tincture of Croton. Bruised croton seeds 1 oz.; rectified spirit 16 oz. Digest for 7 days and filter. Dose ½ oz. to 1 oz. in water.

Tincture of Iodine. Iodine 1 part, rectified spirit 8 parts; dose 1 to 2 dr.—V. C. But the following is preferable:

Compound Tincture of Iodine. Iodine 1 oz., iodide of potassium 2 oz., spirit of wine 12 oz.

Solution of Chloride of Lime. 1. Chloride of lime 1 dr., water 8 oz.; mix in a mortar, and filter.—CLATER.

2. Chloride of lime 1 part, water 10 parts.—CHEVALLIER.

3. Chloride of lime 1 part, water 48 parts.—LABARRAQUE. See LOTIONS, page 99.

Solution of Nitre. Nitre 1 part, water 7 parts.—V. C.

Solution of Ammonio-Sulphate of Copper. Dissolve 1 part of sulphate of copper in 4 parts of water, and add ammonia until it begins to precipitate. 4 ounces every 8 hours as a tonic.—JECKYLL.

Solution of Henbane. Extract of henbane 4 dr., spirit of nitric ether 4 oz. Antispasmodic; dose, 2 oz., with or without solution of aloes.—WRIGHT.

For SOLUTIONS and TINCTURES *for outward use*, see EXTERNAL APPLICATIONS, further on.

CLYSTERS.

Laxative. 1. Aloes 1 oz., water 2 or 3 quarts.—YOUATT.

2. Water gruel 1 gallon, olive oil 1 pint.—WHITE.

3. Epsom salts 6 oz. (or common salt 6 oz., or soap 2 oz.), thin gruel or broth 5 quarts.—BLAINE.

4. Soft soap 2 oz., warm water ¾ of a pailful.—B. CLARK.

5. Infuse 3 oz. senna in 2 quarts of water, and add Epsom salts 4 oz., honey 6 oz.—MOIROUD.

6. Chamomiles, fennel seed, coriander seed, of each 1 oz., caraways ½ oz.; boil in 2 quarts of water to 3 pints; strain,

add 2 oz. Epsom salts, and when nearly cool, ¼ pint of olive oil and ¼ pint of tincture of senna.—TAPLIN.

Purgative. 1. Aloes 8 to 12 dr., salt 8 oz., water 1 gallon: in staggers.—WHITE.

2. Senna 2 oz., tobacco 2 oz.; boil for a quarter of an hour in 2 quarts of water, strain and add common salt 4 oz., emetic tartar 1 dr.; for 2 doses. *Very irritating.*—LEBAS.

Emollient. Dried mallow leaves, or marshmallow root, 1½ oz., linseed ½ oz., water 2 quarts; boil and strain: to be used warm.—MOIROUD.

Emollient and Anodyne. 1. Mix 6 dr. of starch in powder with a little cold water, and add it to a decoction of 6 poppy-heads in 2 quarts of water; boil for an instant and strain: in intestinal irritation.—MOIROUD.

2. Gruel 2 pints, liquid starch or arrowroot 1 pint, powdered opium 1 dr. to 1½ dr.—WHITE.

3. Boil 6 poppy-heads in 4 quarts of water till reduced to 2 quarts: add prepared chalk 2 oz., boiled starch 2 quarts; once or twice a day in diarrhœa.—BLAINE.

4. A double handful of coarse bran, 6 poppy-heads, 2 quarts of water; boil and strain.—MOIROUD.

5. Tripe liquor (or suet boiled in milk) 3 pints, thin starch a quart, laudanum ½ oz.: in diarrhœa.

Cooling. Butter-milk or whey, barley-water, of each a quart.—MOIROUD.

Carminative and Stimulant. 1. Chamomiles 3 oz., aniseed or fennel-seed 1½ oz., 4 poppy-heads; boil the poppies in sufficient water, and infuse the flowers and seeds in the hot decoction.—VATEL.

2. *To expel wind.* Boil 1 lb of figs in 3 quarts of water for half an hour, then add 2 handfuls of chopped rue; boil a few minutes, strain and add 8 oz. of olive oil.—SOLLEYSELL.

For Gripes. Mash 2 onions, pour over them 2 oz. of oil of turpentine and 4 quarts of thin gruel.—BLAINE.

Astringent. 1. Alum whey 1 quart, thin starch a quart.

2. Suet milk 3 pints, starch gruel 2 pints, laudanum ½ oz.

Vermifuge. 1. *For Thread Worms.* Powdered aloes ½ oz., powdered gum Arabic ½ oz.; mix with half pint of boiling

water; then mix the white of an egg with a quart of linseed oil, and gradually add the solution of aloes.—CLATER.

2. Infuse 4 oz. of tansy in 2 quarts of water; strain and add 2 oz. of animal oil (empyreumatic oil of hartshorn); also the worm drenches (Nos. 4 and 6) may be used in this method.—MOIROUD.

Uterine Stimulants. 1. Infuse a handful of rue in 2 quarts of water, and add 2 oz. of common salt.

2. Savine 2 oz., sal ammoniac 4 dr.: as the last.

Diuretic. 1. Nitric 1 oz., decoction of linseed 3 pints.

2. *Comphorated.* Incorporate 4 dr. of camphor with the yolks of 2 eggs, and add it to the last.—MOIROUD.

For Irritable Bladder. 1. Belladonna leaves 3 oz., water 3 pints; boil and administer warm.—MOIROUD.

2. Extract of belladonna ½ oz., boiling water 1½ pint.

Nourishing. 1. Thick gruel 3 quarts, ale 1 quart.—BLAINE.

2. Milk 2 quarts, yolks of 4 eggs; mix, and give warm.—BOURGELAT.

3. Strong broth 2 quarts, thickened milk 2 quarts.—BLAINE.

4. Tripe liquor or broth 3 quarts, flour 4 oz.; mix the flour in the hot broth: repeat frequently.—MOIROUD.

External Applications.

LINIMENTS AND EMBROCATIONS.

BLISTERING LINIMENTS, OR LIQUID BLISTERS, AND SWEATING OILS.

1. Powdered Spanish flies 1 oz., spirit of wine 6 oz., water of ammonia 2 oz.; let it stand for a week, shaking it frequently, and strain. (See No. 11.)—WHITE.

2. Flies 1 oz., euphorbium ½ oz., oil of turpentine 4 oz.; digest for 2 or 3 days, and pour off the liquid; digest the flies, &c., in 4 oz. of spirit of wine and 2 oz. of water of ammonia for 3 or 4 days, shaking frequently; strain off this liquid, and mix it with the former. This is more active than the last.—WHITE.

3. BLAINE'S *Liquid Blister.* Spanish flies, coarsely pow-

dered, 8 oz., oil of turpentine 2 quarts; steep for 3 weeks, strain, and add a quart of olive oil.

4. BLAINE'S *Milder or Sweating Liquid.* Mix 4 oz. of the last with 6 oz. of oil.

5. CLATER'S *Strong Liquid Blister.* Spirits of turpentine colored with alkanet, 1 gallon, powdered flies 1 ℔; macerate for a month, shaking daily, then pour off the clear fluid for use.

6. *Common or Sweating Liquid.* Mix the last with equal parts of spermaceti oil.

7. Powdered flies 2 oz., spirit of turpentine a pint; digest for a few days.—YOUATT.

8. *Blistering Liniment for immediate use.* Spanish flies in fine powder 1 oz., oil of turpentine 6 oz. To be rubbed on the belly in inflammation of the bowels.—WHITE. Cantharides 3 dr., in hartshorn 4 oz.; in a fortnight strain, and add olive oil 4 oz.—SPOONER.

9. *Croton Liniment.* A tincture of croton nuts with oil of turpentine is used as a blister, but is not so efficacious as cantharides.—YOUATT.

10. WHITE'S *Mustard Blister.* Best flour of mustard 8 oz., water enough to form a paste, oil of turpentine 2 oz., water of ammonia 1 oz.

11. *Blistering Tincture.* Flies 1 oz., proof spirit 8 oz.; macerate 2 or 3 weeks; mix, and filter. To be rubbed in, and repeated next day if necessary.—WHITE.

12. Saturated tincture of cantharides 1 oz., bichloride of mercury 6 gr.—KENT.

13. Powdered cantharides 1 dr., olive oil 2 oz. To be applied every 48 hours for a week in old spavin.—TAPLIN.

14. *Oil of Cantharides* (by infusion). Digest 1 part of powdered cantharides in 8 oz. of olive oil in a water-bath, for two hours and strain.

15. *Cantharidal Solution* (for setons). Digest 1 part of p. flies with 8 of oil of turpentine, with a gentle heat, for 14 days; strain and add to the clear liquid an equal weight of Canada balsam. Soak the cotton cord in the solution, draw it between the finger and thumb, and dry it.—MORTON.

STIMULATING LINIMENTS.

1. Soft soap 4 oz., camphor 1 oz., proof spirit 2 pints, water of ammonia ½ pint.—V. C.

2. Sweet oil 2 oz., spirit of hartshorn 1 oz., oil of turpentine ½ oz.—WHITE.

3. Common oil 6 oz., liquid blister 2 or 3 oz.: in chronic sprains.—BLAINE.

4. *Soap Liniment.* Soft soap 4 oz., water 8 oz.; dissolve, and add 1 pint of rectified spirit, in which is dissolved 2 oz. camphor, 1 oz. oil of rosemary, and 2 to 4 oz. strong water of ammonia.—WHITE.

5. For splints: Oil of origanum 1 oz., spirit of turpentine 1 oz., spirit of wine ½ oz. To be applied night and morning for a few days, discontinuing it as often as any moisture appears.—LANCET.

6. For the same purpose: Oil of origanum ½ oz., oil of turpentine ½ oz., camphorated spirit of wine 2 oz.—TAPLIN.

7. For sprains, old swellings, rheumatism, &c.: Spirit of hartshorn 2 oz., camphorated spirit 2 oz., oil of turpentine 1 oz., laudanum ½ oz., oil of origanum 1 dr.

8. Camphorated oil 4 oz., oil of turpentine 1 oz., oil of origanum 1 dr.

9. For callous swellings after bruises: Soap liniment 4 oz., camphor 2 dr., water of ammonia 1 oz.

10. For indolent tumors: Mercurial ointment 2 oz., olive oil 2 dr., camphor 2 dr.

11. Olive oil 4 oz., water of ammonia 2 oz., oil of turpentine 2 oz.

12. For strains: Barbadoes tar 2 oz., spirit of turpentine 2 oz., opodeldoc 4 oz.—TAPLIN.

13. Oil of turpentine 2 parts, muriatic acid 1 part.—POTT.

14. Camphorated oil 4 parts, oil of turpentine and tincture of cantharides, of each 2 parts, acetic acid 1 oz.—LEBAS.

15. *Turpentine Liniment.* Equal parts of oil of turpentine and oil. Digestive and rubefacient.—V. C.

16. *Compound Turpentine Liniment.* Soft soap 4 oz., camphor 1 oz., oil of turpentine 16 oz.; mix.—V. C.

LINIMENT FOR BOG SPAVIN. Mercurial ointment 2 oz., oil of cantharides 4 dr.—MORTON.

LINIMENT FOR SORE BACKS. Extract of lead ½ oz., vinegar 1 oz., olive oil 2 oz.—WHITE.

LINIMENT FOR ITCHING HUMORS, MANGE, LICE, &c.

1. Equal parts of oil of tar, oil of turpentine, and seal oil. Apply every second day for 2 or 3 times, then wash. —V. C.

2. Sulphur 4 oz., turpentine 4 oz., oil of tar and train oil 6 or 8 oz. The parts to be first washed with soft soap, and dried.

3. For lice: Sublimate 1 dr., muriatic acid 3 dr., tobacco water 2 pints, oil of turpentine 4 oz.—WHITE.

Liniment for Mange. Goulard's extract of lead 2 oz., olive or rape oil, 2 oz., sulphur 1 oz.

LINIMENT FOR CANKER OF THE FOOT AND BAD THRUSHES. See also CAUSTICS and LOTIONS.

1. Barbadoes tar 1 oz., oil of turpentine 1½ oz.; mix carefully, and add oil of vitrol 1 dr.—WHITE.

2. Butter of antimony alone.

3. Crystallized verdigris in fine powder 1 oz., honey 2 oz., bole and alum of each ½ oz., vinegar to form a liniment; to be mixed over a gentle fire. Greasy applications are to be avoided.—WHITE.

See also DETERGENT LINIMENTS.

DETERGENT LINIMENTS.

1. Oil of turpentine 1 oz., oil of vitriol 2 dr. by measure; mix in a large gallipot, and when cool, add 2 oz., of linseed oil.—WHITE.

2. *Ægyptiacum.* Bruised sulphate of copper 12 oz., vinegar 4 lb, treacle 3 lb; place over a clear fire, and let it boil up.—B. CLARK.

3. *Wash for Grease.* Sulphate of copper 2 dr., and alum 2 dr. in water 1 pint.

MISCELLANEOUS LINIMENTS AND MIXED OILS. See also EMBROCATIONS.

Creasote Liniment. Creasote 2 oz., oil of turpentine 4 oz. olive oil 4 oz.; mix: in fistulous sores, unhealthy wounds, &c.—V. C.

Oil of Cantharides. Powdered flies 1 oz., olive oil 8 oz.; digest in a water-bath for 2 or 3 hours and filter.—V. C.

Goulard Liniment. Extract of lead 1 oz., olive oil 4 oz.—MORTON. For excoriated surfaces, &c.

Saturnine Balsam. Acetate of lead 1 oz., oil of turpentine 2 oz.; digest with a gentle heat.—MOIROUD.

Drying Liniment. Linseed oil and spirit of wine, of each equal parts.—SOLLEYSELL.

Marshmallow Liniment. Olive oil and marshmallow ointment, of each 4 oz.; melt the ointment and add the oil.—BOURGELAT.

Emollient and Anodyne Liniment. Neatsfoot oil 4 oz., poplar ointment, marshmallow ointment, of each 2 oz.—MOIROUD.

Lime-water Liniment. Lime-water 8 oz., olive or linseed oil 2 oz.

Narcotic Liniment. Olive oil 4 oz., laudanum 2 oz.—MOIROUD.

Liniment for Confirmed Grease. Verdigris, sugar of lead, of each ¼ oz., honey 1 oz.; mix.—CLATER.

Compound Iodine Liniment. Iodine 1 oz., soap liniment 8 oz.—V. C.

Turpentine Liniment. Equal parts of turpentine and olive oil.—V. C.

Resolvent Liniment. Olive oil 2 oz., strong mercurial ointment, 2 dr., water of ammonia 2 dr.

Black Oils. Olive (or rape) oil 1 pint, oil of turpentine 2 oz., mix and add gradually 6 dr. of sulphuric acid; leave the bottle open till cold.—PERCIVALL.

Oils for Mange. Oil of turpentine 1 pint; add to it very gradually and cautiously 2 oz. oil of vitriol, stirring the mixture constantly, then add a quart of linseed oil; from 4 to 8 oz. to be rubbed in with a brush every second day, for three or four times.—CLATER.

WARD'S *White Oils.* Spirit of wine, oil of turpentine, rape oil, beef brine, camphor, of each equal parts.

White Oils or Egg Oils. 1. Yolks of 2 eggs, 3 oz. solution of ammonia, 1 oz. oil of origanum, 4 oz. oil of turpentine, a pint of vinegar; mix, s. a.—PHARMACEUTICAL JOURNAL.

2. Distilled vinegar 1½ pint, oil of turpentine 1½ dr., spirit of wine 1½ oz., GOULARD'S extract of lead ½ oz., whites and yolks of 2 eggs: mix the turpentine and Goulard with the eggs, gradually add the vinegar, and lastly the spirit.—REDWOOD'S GRAY'S SUPPLEMENT.

Liniment of Ammonia. This is sometimes termed *White Oils.* Olive or rape oil 4 oz., water of ammonia 1 oz. Sometimes 1 oz. of oil of turpentine is added to increase its activity.

Darby's Oils. Equal parts of oil of amber, Barbadoes tar, and balsam of sulphur.

MARSHALL'S *Oils.* Linseed oil 1 ℔, olive or rape oil 1 ℔, green oil ½ ℔, oil of turpentine ½ ℔, oil of vitriol 1½ dr.

Newmarket Oils. Linseed oil, oil of turpentine, green oil, of each 3 ℔, oil of vitriol 1 oz.

Nine Oils. Train oil 23 ℔, oil of turpentine 6 ℔, oil of bricks 1 ℔, oil of amber 1 ℔, spirit of camphor 2 ℔, Barbadoes tar 7 ℔, oil of vitriol 2 oz.—GRAY'S SUPPLEMENT.

RADLEY'S *Oils.* Barbadoes tar 8 oz., linseed 4 oz., oil of turpentine 4 oz.

LORD STAMFORD'S *Mixed Oils.* Oil of origanum 6 oz., oil of turpentine 24 oz., spirit of wine 16 oz., green oil 6 ℔, camphor 3 oz.

EMBROCATIONS, VARIOUS.

(See also LINIMENTS and LOTIONS.)

Embrocation for Strains. 1. Soft soap 1 oz., spirit of wine 4 oz., oil of rosemary 2 dr., camphor 2 dr.

2. For strains in the shoulder: oil of turpentine 1 oz., camphorated spirit 2 oz.—BLACKER.

3. Equal quantities of soft soap, oil of turpentine, spirit of wine, and elder ointment.—WHITE.

4. Soft soap 2 oz., oil of bays 1 oz., water of ammonia 1½ oz., oil of origanum ½ oz.

5. Barbadoes tar 2 oz., spirit of turpentine 2 oz., opodeldoc 4 oz. After fomenting with hot vinegar and Goulard.—TAPLIN.

Mustard Embrocations. 1. Mustard flour 4 oz., water of ammonia 1½ oz., oil of turpentine 1 oz., water enough to bring it to the consistence of cream.

2. Camphor 1 oz., oil of turpentine 2 oz., water of ammonia 2 oz., flour of mustard 8 oz., water to form a thin paste.—WHITE.

Embrocations for Poll-evil. Spirit of wine ½ pint, camphor 2 dr., Goulard's extract of lead 1 dr.; mix.—HINDS. See DISCUTIENT LOTIONS.

Embrocations for Saddle-galls or Warbles. 1. Goulard's extract of lead 2 dr., distilled vinegar 3 oz., spirit of wine, 4 oz.—WHITE.

2. Soap liniment and Mindererus' spirit, equal parts.

3. Sal ammoniac ½ oz., muriatic acid 2 dr., water 8 to 12 oz.

4. White vinegar 3 oz., spirit of wine 3 oz., sugar of lead 2 dr., water 6 oz.; mix.—HINDS.

CLARK'S *Embrocatio Frigorifera.* Vinegar 4 oz., camphor (dissolved in spirit) ½ oz., water to fill up a wine-bottle.

CLARK'S *Embrocatio Excitans.* Olive oil 3 oz., camphor ½ dr., spirit of turpentine ½ oz., water of ammonia 3 dr.

TAPLIN'S *Embrocation for Windgalls.* Oil of origanum, spirit of turpentine, of each ½ oz., camphorated spirit 1 oz. Applied with tow, and covered with a piece of lead bound on.

LOTIONS OR WASHES.

COOLING LOTIONS, for external inflammation.

1. Sal ammoniac 1 oz., nitre 2 oz., water 16 oz. To be used as soon as made.—MORTON.

2. Goulard's extract of lead 1 oz., vinegar 2 oz., camphorated spirit 3 oz., water 16 oz.; for recent spavin.—TAPLIN.

3. V. C. *Goulard Water.* Goulard's extract 2 dr., spirit 2 dr., soft water 1 pint.

4. WHITE'S *Saturnine Lotion.* Sugar of lead 1 oz., vinegar and water 1 pint.

5. B. CLARK'S *Lotio Refrigerans.* Liquor of diacetate of lead 1 dr., spirit of nitric ether 1 dr., water 2 pints. In slight rubs and bruises.

DISCUTIENT LOTIONS for dispersing indolent tumors, and saddle-galls, and for chronic strains, &c.

1. Mindererus' spirit 4 oz., camphorated spirit 4 oz., water 16 oz.—PERCIVALL.

2. Sal ammoniac 1 oz., vinegar 8 oz., camphorated spirit 1 oz.—MORTON.

3. For saddle-galls and warbles: Goulard's extract 2 dr., distilled vinegar 3 oz., spirit of wine 4 oz.—WHITE.

4. Muriate of ammonia ½ oz., muriatic acid 2 dr., water 8 to 12 oz.; for saddle-galls and windgalls.—WHITE.

5. BLAINE'S *Saline Embrocation.* Sal ammoniac 8 oz., vinegar 3 pints.

6. Mindererus' spirit 2 oz., soap liniment 2 oz.—WHITE.

7. For warbles: white vinegar 3 oz., spirit of wine 3 oz., sugar of lead 2 dr., water 6 oz.—HINDS.

8. Strong solution of salt 1 oz., tincture of myrrh ¼ oz., for saddle-galls.—YOUATT.

9. Common salt 4 oz., vinegar ½ pint, cold water 1 quart, spirit of wine and laudanum, each 1 oz.: in incipient poll-evil.—CLATER.

10. White vinegar, 1 pint, extract of lead 2 oz., camphorated spirit 4 oz., soft water 1 pint.—TAPLIN.

11. For strains: Bay salt ½ lb., sal ammoniac 2 oz., sugar of lead ¼ oz., vinegar 1½ pint, water 1 pint.

ASTRINGENT LOTIONS, for drying up sores and diminishing their discharge (especially in grease and scratched heels), after the inflammation has been subdued by linseed or carrot poultices.

1. Alum 4 oz., boiling water 1 pint, for grease and cracked heel.—TAPLIN.

2. Alum 2 dr., sulphate of zinc 1 scruple, water 1 pint.—YOUATT.

3. Mild, for cracks: Sugar of lead 2 dr., sulphate of zinc 1 dr., infusion of oak bark 1 pint.—BLAINE.

4. For confirmed grease: Nitric acid 1 oz., water 8 oz.—BLAINE.

5. Strong: Blue vitriol ¼ oz., alum 3 dr., water 1 pint.—SPOONER.

6. Sugar of lead 1 oz., blue vitriol 1 oz., water 1 quart.—WHITE.

7. Lime-water 16 oz., spirit of camphor ½ oz., sugar of lead 1 dr.—BOURGELAT.

8. Sulphate of iron 2 oz., alum 2 oz., vinegar 8 oz., water 3½ pints.—MOIROUD.

9. Tincture of myrrh 1 oz., camphorated spirit 1 oz., distilled vinegar and water, each 2 oz.—TAPLIN.

10. For anburies: Alum 2 oz., water 1 pint, sulphuric acid 1 dr.

11. Sulphate of iron 1 oz., water 1 quart; dissolve and add ¼ oz. (by weight) of oil of vitriol. To wash farcy buds after they have been opened.—CLATER.

DETERGENT LOTIONS, for foul ulcers.

1. Sulphate of copper 1 oz., nitric acid ½ oz., water 6 oz.—WHITE.

2. Sulphate of copper 1 oz., sulphuric acid 12 drops, water 4 oz.

3. Sulphate of copper 2 dr., water 1 pint: for stimulating old ulcers.—YOUATT.

4. Sulphate of copper 1 oz., water 1 oz.; to remove fungous granulations.

5. Nitrous acid 1 oz., quicksilver ½ oz., dissolve, and add water 8 oz.

LOTIONS FOR MANGE.

1. White hellebore 2 oz., tobacco 2 oz., water 3 pints; boil, strain, and add, when cold, a pint of fresh lime-water.—BLAINE.

2. Boil 4 oz. of white hellebore in 3 pints of water to 2 pints, and add corrosive sublimate 2 dr., previously dissolved in 3 dr. of muriatic acid.—WHITE.

3. Boil 2 oz. of tobacco in a quart of water, strain, and add common salt 3 oz., soap 2 oz.—LEBAS.

4. Liver of sulphur 2 oz., water 1 quart.—MOIROUD.

5. Liver of sulphur 4 oz., soft soap 16 oz., water 2 gallons.—LEBAS.

6. Acid nitrate of mercury 2 dr., distilled water 16 oz.—MOIROUD.

7. *Mercurial Wash.* Sublimate 2 dr., spirit of wine 2 oz., water 2 pints.—CLARK.

8. Chloride of lime 1 ℔, water a gallon. Mix.—LUCAS.

VARIOUS LOTIONS.

Conglutinum. Sulphate of zinc 4 oz., water a pint.—BRACY CLARK.

Black Wash. For sluggish ulcers: Calomel 2 dr., lime-water 1 pint.

Yellow Wash. Sublimate 8 gr., lime-water 2 oz.

Nitric Acid Lotion. Nitric acid 2 or 3 dr., water 1 pint; for exciting sluggish ulcers.—MORTON.

Lotion of Nitrate of Silver. For the same: Nitrate of silver 10 gr., distilled water 1 oz.—YOUATT.

Lotion for Farcy. Dissolve 1 oz. of sulphate of iron in a quart of water, and add ½ oz. of oil of vitriol.—CLATER.

Styptic Lotion, for Stopping Bleeding. Alum 2 oz., sulphate of zinc 2 dr., water 1 quart.

Catechu Lotion for Ulcers of the Mouth. Infuse two oz. of catechu in a quart of boiling water for an hour; strain and add 1 oz. of spirit of wine. (For saddle-galls add 4 oz. of tincture of catechu and 8 oz. common salt.)—CLATER.

Lotion of Chloride of Lime. To chloride of lime 1 ℔ add gradually 1 gallon of water; mix, and filter or decant: for mange, and as a stimulant to unhealthy wounds and fistulous sores. Diluted with 10 or 15 parts of water, it is used as a lotion for grease, exfoliated bones, &c., and is a disinfectant for foul stables. For ulcers of the tongue, mix 1 dr. of chloride of lime with a pint of water: for mange, 4 dr. to a pint.

Wash for destroying Lice about the Legs. Corrosive sublimate 1 dr., muriatic acid 3 dr., tobacco-water 1 quart, oil of turpentine 4 oz.—WHITE.

Alum Mouth Wash. Alum 2 dr., sage tea a quart.—ECKEL.

For Bruised Gums. Alum 2 dr., tincture of myrrh 1 oz., honey 1 oz., water 2 oz.—SPOONER.

Acid Collutorium. Infusion of sage a quart, muriatic acid 1 oz., flour 3 oz., honey 8 oz. To be applied to the mouth frequently.—ECKEL.

LOUSE-WATER.

1. Tobacco 4 oz., boiling water a quart; infuse for 24 hours.—CLARK.

2. *Mercurial.* Sublimate 2 dr., spirit of wine 2 oz., water 1 quart.

LIQUID CAUSTIC, for canker and thrush, for foul, unhealthy wounds, to remove proud flesh, &c. See also LINIMENTS (Detergent).

MILDER.

1. Tincture of muriate of iron.

2. Sulphate of copper 1 oz., water from 4 oz. (V. C.) to a pint.—CLATER.

3. Saturated solution of sulphate of zinc: in quitters.—WHITE.

4. Alum ½ oz., borax ½ oz., boiling water 4 oz., styptic tincture 1 oz.—TAPLIN.

5. Muriatic acid, alone or diluted.

6. Ægyptiacum 2 oz., nitrous acid 20 drops.

7. Any of the stronger caustics (except butter of antimony) diluted with water.

8. Goulard's extract 4 oz., sulphate of zinc 2 oz., sulphate of copper 2 oz.; white vinegar 32 oz.—VILLATE.

9. Aloes 5 oz., weak spirit 10 oz.; dissolve, and add 6 oz. sulphuric acid.—DUVILLE.

STRONGER.

1. Butter of antimony. This is the safest and most useful caustic in canker.

2. Dissolve 1 oz. quicksilver, by heat, in 2 oz. of nitric acid, and evaporate till the liquid weighs 2¼ oz.

3. Verdigris 1 oz., nitrous (red nitric) acid 1 oz.; dissolve.—WHITE.

4. Red precipitate 1 oz., nitrous acid 2 oz.—WHITE.

5. Nitrous acid alone, or with a little water.

6. Sulphuric acid, alone or with a little water.

7. Sublimate 1 dr., muriatic acid 2 dr., water q. s.—WHITE.

8. For canker: Dissolve corrosive sublimate ¼ oz. in muriatic acid 1 oz., then add spirits of wine 4 oz., and water 4 oz.

9. Chloride of zinc with enough water to dissolve it; or SIR WILLIAM BURNETT'S PATENT SOLUTION.

CAUSTIC FOR POLL-EVIL.

1. Lunar caustic 1 dr., distilled water ½ oz.—BLAINE.

2. Corrosive sublimate 2 dr., water 3 oz.

Scalding Mixture for Poll-evil. 1. Sublimate 2 dr., verdigris 2 dr., blue vitriol 2 dr., sulphate of iron 4 dr., honey 2 oz., oil of turpentine 8 oz., spirit of wine 4 oz.; to be applied hot, and confined by stitches.—GIBSON.

3. Sublimate 1 dr., finely powdered and mixed with 4 oz. of basilicon, and melted to scalding heat.—BLAINE.

4. Caustic potash 1 dr., rubbed down with 4 oz. oil of turpentine.—BLAINE.

CAUSTIC FOR FARCY BUDS. Sublimate 1 dr., muriatic acid 3 dr., spirit of wine 1 oz., water ½ oz.—WHITE.

SOLID CAUSTICS. Lunar Caustic, Caustic Potash, and Chloride of Zinc. See Argenti Nitras, Potassæ Hydras, and Zinci Chloridum, Pocket Formulary.

Canquoin's Caustic is made by mixing chloride of zinc with twice its weight of flour and a little water into a stiff paste, which is to be rolled out to the required thickness, and cut to the size of the part to be destroyed, the skin being previously removed by a blister. Another caustic is made with 2 parts of chloride of zinc, 1 of butter of antimony, and 5 of flour.

Sulphuric Caustic is made by triturating hay saffron with oil of vitriol, so as to form a ductile mass. BOUCHARDAT recommends solidifying the acid by ivory or lamp-black.

Solidified Nitric Acid is merely lint soaked with strong nitric acid, squeezed and formed to the required shape.

Filho's Caustic is made by melting together in an iron ladle 2 parts of caustic potash and 1 of lime over a quick fire, and pouring it into leaden tubes of the desired size. The air must be excluded when not in use, by beeswax, or other means.

FOMENTATIONS.

These should be applied moderately warm (about 120°) by means of flannel dipped in the liquid, and frequently renewed from time to time, keeping the parts covered.

Emollient. 1. Coarse bran 2 double handfuls, water 6 quarts, boil and strain.

2. Mallow-leaves 8 oz., water 4 quarts; boil and strain; 6 poppy-heads may be added.

Anodyne. 1. Boil 24 poppy-heads and 2 handfuls of hemlock in 6 quarts of water for two hours, and strain.—WHITE.

2. Belladonna 2 handfuls, 6 poppy-heads, water 3 quarts; boil and strain.—MOIROUD.

3. Dried wormwood and chamomile, of each 4 oz., bay-leaves 2 oz., rue 3 oz.; boil in a gallon of water.

4. Take wormwood, chamomile, mallow (or either of them), cut them to pieces, and put 2 handfuls into a bucket,

pour scalding water on them, and cover with a cloth.—B. CLARK.

Discutient and Astringent. Vinegar or verjuice 1 quart; make it hot, and add 2 oz. of Goulard's extract of lead; apply warm, in strains of the sinews of the legs.—TAPLIN.

COLLYRIA OR EYE-WATERS.

1. Acetate of lead, and sulphate of zinc, each ½ dr. to 1 dr.; dissolve them separately in ½ pint of water; mix, and filter.

2. Sugar of lead 10 to 20 grains, water 8 oz.—MORTON.

3. Extract of lead 1 dr., spirit 2 dr., water 8 oz.—WHITE.

4. Acetate of ammonia (L. P.) 3 oz., rose-water 6 oz.—BLAINE.

5. Sugar of lead 2 dr., vinegar ½ oz., soft water 16 oz., rose-water 4 oz.—BLAINE.

6. Infuse 1 oz. of foxglove in 2 pints of boiling water, and strain.—YOUATT.

7. Tincture of opium 2 dr., water 8 oz., extract of lead 1 dr.—WHITE.

8. Brandy 1 oz., vinegar 1 oz., tincture of opium 2 dr., rose-water 8 oz.—BLAINE.

9. Extract of henbane 1 dr., water 8 oz.—WHITE.

10. Decoction of poppies 8 oz., saffron ½ dr.; infuse the saffron in the hot decoction.—LEBAS.

11. Lapis divinus 3 dr., soft water ½ pint.—CLATER.

12. Common salt ½ dr., water 6 oz.—YOUATT.

13. *For Watery Bloodshot Eyes.* Burnt alum 1 oz., calcined white vitriol 1 oz., boiling water 3 pints.—BRACKEN.

14. *Emollient.* Infusion of marshmallow leaves or flowers 1 quart, starch (rubbed smooth with a little water) ½ oz.; mix and boil. To be used warm.—MOIROUD.

15. *Astringent.* Alum 2 dr., whites of 2 eggs, water ¼ pint; mix in a mortar.—BOURGELAT.

16. Tincture of digitalis ½ oz., soft water 8 oz.—CLATER.

17. *To remove Opacity of the Cornea.* Nitrate of silver 10 gr., distilled water 1 oz. 1 or 2 drops to be dropped in the eye.—YOUATT.

18. *For Cloudiness of the Eye.* Sublimate 4 gr., spirit of wine 20 drops; rub together, and add soft water 4 oz.

A few drops to be introduced into the eye 3 or 4 times a day.—CLATER.

19. Tincture of aloes 1 oz., rose-water 8 oz.—LEBAS.

20. *Stimulating.* Infusion of elder flowers 16 oz., brandy 2 oz.—MOIROUD.

21. Lapis mirabilis ½ oz., water 4 to 8 oz. The *Lapis mirabilis* is thus made: White vitriol 2 lb, rock alum 3 lb, fine bole ½ lb, litharge 2 oz., water 3 quarts; boil together to dryness.—SOLLEYSELL.

22. *Alum Collyrium.* Decoction of marshmallow 16 oz., alum 2 dr., camphorated spirit 1 dr.; mix. To be used towards the decline of inflammation.—STRAUSS.

23. *Tannin Collyrium.* Dissolve 1 dr. of tannin in 13 oz. of water, and add 3 oz. of cherry-laurel water.

24. Sulphate of zinc 8 gr., water 4 oz. In chronic inflammation.—CLATER.

25. In *Specific Ophthalmia.* Tincture of opium 2 dr., extract of belladonna 1 dr., with distilled water 1 pint.

SUNDRY SOLUTIONS.

Styptic Stone. Sulphate of iron 8 oz., sal ammoniac, sulphate of zinc, and oxide of copper, each 1 oz.; mix, and melt together with a gentle heat. About the size of a nut of this compound to be dissolved in a quart of warm water and applied with compresses renewed every 3 or 4 hours: for saddle-galls, kicks, sprains, bruises, ulcers, and as a collyrium.—KNAUP.

Lapis Divinus. Sulphate of copper, alum, nitre, of each 3 oz.; melt together, and stir in 1 dr. of camphor: used in eye-waters and lotions.

Wound Stone. Alum, sulphate of zinc, of each 3 oz., verdigris and sal ammoniac, of each 1 dr.; melt together, and add ½ dr. of powdered saffron: detergent and drying.

Clark's Conglutinum. Sulphate of zinc 4 oz., water a pint.

Solution of Alum. Alum 1 oz., water 16 oz. Dissolve.—V. C.

Solution of Sulphate of Zinc. Sulphate of zinc 1 oz., water 3 oz.—V. C. In quitters it is usually diluted.

Solution of Sulphate of Copper. Sulphate of copper 1 oz., water 4 oz.—V. C.

Compound Solution of the same. Sulphate of copper 3 oz., alum 3 oz., water 2 lb, sulphuric acid 1½ oz.

Solution of Bichloride of Mercury. Sublimate, hydrochloric acid, each 1 part, spirit of water 7 parts.—V. C.

Goulard Water. Extract of lead 1 oz., camphorated spirit 2 oz., rain-water a quart.—TAPLIN. V. C. use extract of lead and rectified spirit, each 2 dr., soft water 1 pint.

Tincture of Catechu, See P. F. Used externally for wounds.

Tincture of Euphorbium. Euphorbium 1 oz., rectified spirit 6 oz.

Alkaline Tincture of Euphorbium. Euphorbium 8 oz., solution of subcarbonate of potash 3 pints: used as a caustic and stimulant, particularly in *curbs* after the inflammation has been subdued.

Compound Tincture of Cantharides. Powdered flies 4, euphorbium 1, proof spirit 24 parts.—LEBAS.

Styptic Tincture. Tincture of myrrh, spirit of camphor, and Fryars' balsam, equal parts.—TAPLIN.

Ægyptiacum. (Veterinary.) Sulphate of copper in powder 12 oz., vinegar 4 oz., treacle 48 oz.; boil together to a proper consistence.—B. CLARK.

Ægyptiacum with Turpentine. Honey 28 oz., pyroligneous acid 14 oz., powdered verdigris 10 oz.; boil together in a copper vessel till the mixture has a reddish-purple color and the consistence of thin honey; add Venice turpentine 28 oz., and keep it on a slow fire, stirring constantly for a quarter of an hour.—LELOUP.

Liniment of Verdigris. V. C. Verdigris in fine powder 9 oz., alum 6 oz., treacle 1½ lb. Boil until the compound assumes a brown color.

Liniment of Sulphate of Copper. Powdered sulphate of copper 1 part, treacle 4 parts. Simmer in a pipkin over a slow fire until the whole assumes a reddish-brown color. *In canker, severe thrush, &c.*—MORTON.

POULTICES OR CATAPLASMS.

These are useful in reducing inflammation and relieving pain. They should not be used too hot, nor applied too tightly, especially to the feet.

COMMON POULTICES.

1. Bran moistened with hot water, and as much linseed meal added as will give it tenacity.—V. C.

2. Boil a quart of bran for ten minutes with enough water to make a thin mash, then add to it 4 oz. of linseed meal: apply it in a flannel bag.—BLAINE.

3. Fine bran 3 parts, linseed meal 1 part, hot water q. s.

CHARCOAL POULTICES.

1. Oatmeal ½ pint, linseed meal ½ pint, charcoal 4 oz., beer grounds q. s.

2. Carrots scraped, or carrots boiled, with charcoal powder q. s. *Antiseptic.*—BLAINE.

YEAST POULTICES.

1. Linseed meal, oatmeal, boiling water, q. s.; mix, and ferment with a tablespoonful of yeast; in old grease with an offensive smell.—BLAINE.

2. In gangrene: add 2 oz. of turpentine to the last.—BLAINE.

ANODYNE POULTICES.

1. Boil poppy-heads in water, strain, and add linseed meal to stiffen it.—YOUATT.

2. Sprinkle the surface of a simple poultice with laudanum.

CLEANSING POULTICES. Mashed turnips, not pressed, with enough linseed meal, or oatmeal, to give them consistence; or, the charcoal poultice above.

DRAWING POULTICES.

1. Boil 2 ℔ of chopped onions in water, and add to it the crumb of a 4-℔ loaf.—HINDS.

2. Sorrel boiled and squeezed 4 parts, onions baked in ashes 1 part, basilicon ointment 1 part; mix, and apply warm.—VATEL.

RESOLVENT POULTICES.

1. Rye meal 8 oz., prepared chalk 2 oz., vinegar 10 oz.; mix, warm, and stir, till no more gas is disengaged; apply warm.—SOLLEYSELL.

2. Linseed meal 12 oz., powdered hemlock 4 oz., muriate of ammonia 4 oz., vinegar q. s.: to indolent glandular tumors.—LEBAS.

GOULARD POULTICES.

1. To a linseed meal poultice add 1 or 2 dr. of Goulard's extract of lead.—YOUATT.

2. Bread and barley-meal equal parts, Goulard water q. s., lard 4 or 6 oz.—TAPLIN.

CHLORINE POULTICE. Chloride of lime ½ oz., water 1 pint, linseed meal q. s.: to grease, when offensive.—YOUATT.

POULTICES FOR GREASE.

1. The herb cleavers (or goose grass) beaten to a paste.

2. Mash bread and boiled turnips with stale beer, and stir in 1 oz. flour of mustard, turpentine 2 oz., linseed meal 2 oz., lard 6 oz.; night and morning.—TAPLIN.

MUSTARD POULTICE.

1. Mustard flour and linseed meal, equal parts; mix with sufficient hot vinegar to give a proper consistence.

2. Flour of black mustard 3 ℔, hot vinegar, or water, q. s.—MOIROUD.

RUBEFACIENT POULTICE.

1. Fresh horseradish root, grated, and immediately applied.—MOIROUD.

2. *Stronger.* Old yeast 2 ℔, flour of black mustard, 1 ℔, euphorbium powder 4 oz., vinegar q. s.; mix and apply cold.

OINTMENTS, CERATES, CHARGES, &c.

SIMPLE EMOLLIENT OINTMENTS.

1. *Simple Cerate.* Olive oil 1 pint, beeswax 4 oz.

2. *Spermaceti Ointment.* Lard 12 oz., white wax 2 oz., spermaceti 1 oz.

BLISTERING OINTMENTS. The Spanish flies should be finely powdered, and the heat moderate.

1. Lard 4 oz., common turpentine 1 oz., p. flies 1 oz.; melt the lard and turpentine, and stir in the powdered flies.—V. C. Mr. YOUATT substitutes resin for the turpentine.

2. Venice turpentine and resin, of each 1 ℔, palm oil or lard 2 ℔; melt together, and gradually stir in one ℔ of powdered flies.—BLAINE.

3. Palm oil 4 lb, resin 1 lb; melt together, and stir in 1 lb of powdered flies.—CLATER.

4. *Mild.* Lard 4 oz., Venice turpentine 1 oz., p. flies 6 dr.—WHITE.

5. *Stronger.* Mercurial ointment 2 oz., oil of bays 2 oz., Barbadoes tar 1 oz., oil of rosemary 2 dr., p. flies 1 oz.—WHITE.

6. *Strong.* Oil of turpentine 1 oz., oil of vitriol 2 fluid dr.; mix in a basin, and add melted lard 6 oz., oil of origanum 1 oz., powdered flies 1 to 2 oz.—WHITE.

Strongest. Strong mercurial ointment 4 oz., oil of origanum ½ oz., finely powdered euphorbium 3 dr., p. flies ½ oz. —WHITE.

8. BLAINE'S *Mercurial.* Common blister (No. 2, above) 4 oz., sublimate in fine powder ½ dr.; for splints, spavins, &c.

9. *For common purposes.* Lard 6 oz., Venice turpentine 4 oz., beeswax 2 oz., yellow resin ½ oz., oil of origanum ½ oz., powdered cantharides 3 oz. It may be softened in winter by rubbing it with a little turpentine.—WHITE.

10. Powdered flies 5 dr., lard 4 oz., oil of turpentine, 1 oz.—HINDS.

11. *Mustard Blister.* Best flour of mustard 8 oz., water to form a paste.—YOUATT. Others add 2 oz. oil of turpentine, and 1 oz. pure water of ammonia.

Note.—The hair should be clipped closely, or shaved off, the part fomented with warm water, and the blistering ointment well rubbed in. In inflammation of the lungs, &c., blistering is more successful after bleeding. In 24 hours a little olive or neatsfoot oil should be applied, and repeated night and morning. The head should be tied up for the first 2 days, and the litter removed from the stable. If strangury is produced give plenty of linseed tea. The simplest blisters are perhaps the best for common purposes. Sublimate blemishes. *Sweating down* is effected by milder stimulants; for this purpose, the liquid blister (see p. 93) is lowered by some mild oil, &c.

DETERGENT OINTMENTS, for cleansing foul and indolent ulcers.

1. Suet 4 oz., Venice turpentine 6 oz., red precipitate, finely powdered 2 oz.—WHITE.

2. Citrine ointment, alone or with ¼ its weight of Venice turpentine.

3. Sulphate of zinc 1 dr., sulphate of copper 1 dr., oil of turpentine 2 dr.; grind smooth, and mix it with 4 oz. of melted tallow. See also DIGESTIVE OINTMENTS, No. 4.

4. Yellow basilicon 2 oz., black basilicon 1 oz.; melt together, remove from the fire, add 1 oz. of turpentine, and ½ oz. finely powdered red precipitate.—TAPLIN.

5. *Verdigris Ointment.*—Verdigris in fine powder 1 part, common turpentine 1 part, lard 12 parts: mix.—MORTON.

DIGESTIVE OINTMENTS, to promote a discharge from unhealthy and indolent ulcers.

1. Resin 16 oz., linseed oil 12 oz.; melt together with a gentle heat.—CLARK.

2. Strained turpentine, honey, of each 2 oz.; yolks of 4 eggs, myrrh ½ oz., aloes 1 oz.; mix.—SOLLEYSELL.

3. Equal parts of common turpentine and lard melted together.—WHITE.

4. To 1 ℔ of the last add 1 oz. of finely powdered verdigris.—WHITE.

5. Yellow wax 3 oz., common turpentine 3 oz., black pitch 1 oz., resin 6 oz., linseed oil 16 oz.; melt together with a gentle heat, then add oil of turpentine 4 oz., and stir till cold.

6. Olive oil 1 pint, yellow wax and black resin, of each 4 oz., Burgundy pitch and turpentine, of each 2 oz.; melt the other ingredients, and add the turpentine when it is removed from the fire.—TAPLIN.

7. Common turpentine 1 part, lard 3 parts; melt together. —V. C.

8. (Basilicon.) Resin 5 oz., yellow wax 2 oz., lard 8 oz.; melt together.

9. (Black Basilicon.) Pitch, wax, resin, of each 11 oz., olive (or rape or linseed) oil a pint.

EYE OINTMENTS. The powder should be very fine, and the whole rubbed smooth.

1. Nitrate of silver 5 to 10 gr., lard 1 oz.; rub till perfectly smooth. The size of a pea to be introduced between the lids, in chronic ophthalmia.—MORTON.

2. Calamine ½ oz., tutty ½ oz., sulphate of copper ½ dr., sulphate of zinc ½ oz., alum ½ oz., camphor 2 dr., fresh butter 3 oz.; mix, and apply warm, with a feather, to watery inflamed eyes.—BRACKEN.

3. Ointment of nitrated quicksilver 1 dr., zinc ointment 1 oz., camphor 1 dr.

4. Inflammation of the eyelids: Verdigris 1 part, Venice turpentine 1, lard 12.—MORTON.

5. For wounds of the eye: Tutty ointment 1 oz., honey of roses 2 dr., calcined white vitriol 20 gr.; apply with a feather night and morning, and sponge daily with warm milk and water.—BRACKEN.

6. For removing opacity of the cornea: Iodine 2 gr., iodide of potassium 20 gr., lard or butter ½ oz.

EUPHORBIUM OINTMENT. Euphorbium 1, lard 8 parts; mix.—DELAFOSSE.

OINTMENTS FOR SCURVY AND CRACKED HEELS, AND CONFIRMED GREASE. The inflammation should be first subdued by poultices. The milder preparations (which are here placed first) should be employed in the first instance, and afterwards those for confirmed grease.

1. For scurvy heels: Goulard's extract ½ dr., lard 1 oz., mix. The heel should first be gently rubbed with soap and water.—CLATER.

2. For scurvy or cracked heels: Sugar of lead ¼ oz., oxide of zinc ¼ oz., lard or palm oil 4 oz.

3. Melt together 3 oz. white diachylon, 4 oz. olive oil; mix, and when nearly cold, add 3 dr. of sugar of lead in fine powder. First wash the heel, then apply the Astringent Lotion No. 9, and afterwards this ointment; or, elder ointment 4 oz., camphor 6 dr., laudanum 2 dr., extract of lead 2 dr.; mix.—TAPLIN.

4. Melt yellow wax 2 oz., with sweet oil 8 oz., and add sugar of lead ¼ oz.

5. Healing ointment for cracked heels: Lard 4 lb, resin 1 lb; melt together, and stir in 1 lb true calamine.—CLATER. See also SOFTENING and COOLING OINTMENTS (below).

6. For cracked heels and grease: Alum 1 oz., turpentine 1 oz.; lard 3 oz.; melt the turpentine and lard, and stir in the powdered alum.—V. C.

7. For grease: Venice turpentine 4 oz., wax 1 oz., lard 4 oz.; melt together, and add sugar of lead 1 oz. (or alum 2 oz.) in fine powder.—WHITE.

8. Lard, honey, common turpentine, each 8 oz.; melt together, and add powdered alum 6 oz., white vitriol 2 oz.

9. Common turpentine 1 lb; melt and add powdered alum

1½ lb, bole 2 lb; stir till cold; spread on brown paper, and tie over with list.

10. Lard ½ lb, honey ½ lb, common turpentine ½ lb; melt and add p. alum 1 lb, white vitriol 2 oz.; stir till cold.

11. For confirmed grease: common verdigris ½ oz., alum, sulphate of zinc, sugar of lead, of each ½ oz.; tar 6 oz.—BLAINE.

12. Citrine ointment 3 oz., lard 2 oz., turpentine 2 dr., saturated solution of nitrate of copper 2 dr.—BLAINE.

13. Ægyptiacum 8 oz., lard 4 oz., sulphate of zinc in powder 1 oz.; rub together till perfectly mixed.—LASSAIGNE.

14. Prepared verdigris 1 oz., lard 4 oz., honey a sufficient quantity.—DELAFOSSE.

15. Chloride of lime 1 to 2 parts; lard 3 parts; mix. *To remove the fetor in grease.*—MORTON.

OINTMENTS TO PROMOTE THE GROWTH OF THE HAIR, and remove the blemish from broken knees.

1. Camphor ½ dr., oil of rosemary 1 dr., weak mercurial ointment 1 oz., ivory-black and bole to color.—WHITE.

2. Poplar-bud ointment and honey, applied twice a day for 15 to 20 days.—PYE.

3. Calamine 2 dr., prepared charcoal 1 dr., oil of turpentine 1 dr., lard 4 drs.; rub well together with 1 dr. of blister ointment.—CLATER.

4. Liquid blister (No. 3, 6, or 7) 1 dr., ivory-black 1 dr., camphor 1 dr., palm oil 1 oz.

5. Citrine ointment 1 oz., camphor 1 dr., color as above.

HELLEBORE OINTMENT. Powdered white hellebore 1 part, lard 8 parts; an irritating dressing for rowels and setons.—V. C.

HOOF OINTMENT. Tar and tallow equal parts, melted together.—WHITE.

IODINE OINTMENT (simple). Iodine 1 part, lard 8 parts; mix.—V. C.

FARCY OINTMENT. Iodide 1 dr., lard 1 oz. mercurial ointment 1 oz., mix. Useful when the complaint is confined to one leg; from 5 to 10 grains of iodide of potassium being given daily, with a mineral tonic.

IODINE OINTMENT (Compound). Iodine 1 dr., iodide of potassium 2 dr., lard 2 oz.—V. C.

IODIDE OF MERCURY OINTMENT. Red iodide of mercury 1 part, lard or palm oil 7 parts; mix; the size of a nut to be rubbed on daily; in thoroughpin.

For Strangles. Iodide of mercury 1 dr., lard 1 oz.—SPOONER.

OINTMENTS FOR MANGE AND LICE.

1. Sulphur 4 oz., soft soap 4 oz., oil of bays 4 oz., train oil q. s.

2. Sulphur 1 oz., train oil 1 oz., Venice turpentine 2 oz.—YOUATT.

3. Train oil 3 oz., sulphur 1 oz., oil of turpentine 6 oz.—WHITE.

4. Sulphur 8 oz., common turpentine 2 oz., strong mercurial ointment 2 oz., linseed oil 1 pint: rub the flowers of sulphur with a fourth part of the oil, then rub in the turpentine and ointment and gradually add the rest of the oil; half to be rubbed in daily for three days; on the sixth day, wash off with soft soap and warm water.—CLATER.

5. Oil of turpentine 3 oz., oil of vitriol 1 oz.; mix cautiously, avoiding the fumes, and add melted lard 8 oz., train oil 4 oz., oil of turpentine 2 oz., flowers of sulphur or sulphur vivum 4 oz.; stir till cold; apply daily for 3 or 4 times, and give an alterative powder twice a day.—WHITE.

6. Oil of bays 16 oz., strong mercurial ointment 6 oz., oil of turpentine 2 oz., soft soap 4 oz.; mix and apply in the sun; but it is not quite safe.—BRACKEN.

7. Oil of turpentine 4 oz., oil of tar 4 oz., train oil 8 oz., sulphur 4 oz.

8. Sulphur vivum 8 oz., powdered stavesacre 1 oz., mercurial ointment 2 oz., turpentine 2 oz., lard or train oil 8 oz.—BLAINE.

9. Sulphur 4 oz., white hellebore ½ oz., oil of tar 3 oz., train or linseed oil 12 oz.—SPOONER.

10. Soft soap and tar, equal parts.

11. Weak mercurial ointment ½ lb, sulphur vivum 4 oz., white hellebore 3 oz., black pepper 3 oz., oil of tar 1 oz., olive oil enough to make it soft: use daily for 7, 10 or 14 days.—TAPLIN.

OINTMENTS FOR MALLENDERS AND SALLENDERS (Scurvy eruptions).

1. Citrine ointment 2 oz., tar ointment 1 oz.; mix.

2. Lard 2 oz., finely powdered red precipitate 2 dr.—WHITE.

3. Sugar of lead 1 part, tar 2, lard 6; mix, give a diuretic ball occasionally.—YOUATT.

4. Lard 4 oz.; melt, and stir in Goulard's extract 1 oz.—WHITE.

5. Quicksilver 1 oz., common turpentine 3 oz.; mix.—BRACKEN.

6. Sublimate 10 gr., mercurial ointment 1 oz.; mix.

7. Iodide of potassium 1 dr., lard 2 oz., Goulard's extract 4 dr.; mix.

8. Camphor 1 dr., sugar of lead ½ dr., mercurial ointment 1 oz.; mix, and apply after washing with soap and water.—BLAINE.

9. Napthaline 1 dr., cod-liver oil 1 oz., zinc ointment 1 oz.

MARSHMALLOW OINTMENT. The following is often substituted for the Pharmacopœia preparation: Rape oil 1 ℔, yellow wax 6 oz., palm oil ½ ℔, common turpentine 1 oz.

MERCURIAL OINTMENT. This is prepared in the usual way; but Venice turpentine is often used to kill the quicksilver more speedily, as it does not interfere with its veterinary uses.

Strong Mercurial Ointment. 1. Quicksilver 16 oz., Venice turpentine 2 oz.; rub together till the metal is killed, then add 16 oz. of lard.

2. Quicksilver 16 oz., liquid styrax 5 dr., lard 3 oz.; triturate until the metal disappears, and add 12 oz. more lard.—CRESSENT.

Weaker Mercurial Ointment. 1. Strong mercurial ointment 1 part, lard 2 parts.

2. Quicksilver 2 oz., balsam of sulphur ½ oz.; rub together till the globules disappear, and add 6 oz. of lard.—TAPLIN.

Compound Mercurial Ointment. Mercurial Ointment 1 part, soft soap 2 parts.—V. C.

RESOLVENT OINTMENTS, for indolent tumors of the withers, spavins, windgalls, farcy buttons, splints, &c.

1. Strong mercurial ointment 4 oz., cantharides in powder ½ oz., oil of rosemary 2 dr.—WHITE.

2. Biniodide of mercury 1 part, lard or palm oil 7 parts; rub together in a mortar; the quantity of a nut to be rubbed on daily till a scurf is produced; for spavin and thoroughpin.—SPOONER.

3. Blister ointment 2 oz., strong mercurial ointment 1 oz., soft soap ½ oz., oil of bays 3 dr., yellow wax 3 dr.; melt the wax by a gentle heat, add the other ingredients, mix by stirring, remove and stir till cold.—LEBAS.

4. Common turpentine 12 parts, corrosive sublimate 1 part: mix.—GIRARD.

OINTMENT OF NITRATE OF SILVER. Nitrate of silver 5 to 10 gr., lard 1 oz.—MORTON.

CREASOTE OINTMENT. Creasote 2 parts, lard 2 parts. —V. C.

MILD CITRINE OINTMENT. Ointment of nitrate of quicksilver 1 part, lard and oil 2 parts. *In tarsal ophthalmia.*

SOFTENING AND COOLING OINTMENTS, for cracks, and ulcers on the heel, &c.

1. Spermaceti ointment 4 oz., olive oil 1 oz., sugar of lead 2 dr., oxide of zinc 1 oz.—WHITE.

2. Extract of lead ½ dr., lard 1 oz.; mix.—CLATER.

3. Marshmallow ointment 4 oz., extract of lead 3 dr., elder ointment ½ oz., calamine 1 oz.

OINTMENT FOR SIT-FASTS, and all hard tumors. Strained ammoniacum 4 oz., mercurial ointment 8 oz., oil of turpentine 10 oz.—HINDS.

OINTMENTS FOR SORE BACKS AND SADDLE-GALLS. See LOTIONS.

1. Camphor 2 dr., oil of rosemary 1 dr., elder ointment or lard 3 oz.

2. Marshmallow ointment 4 oz., extract of lead 1 oz.—WHITE.

SULPHURIC ACID OINTMENT.

1. Sulphuric acid 1 dr., lard 1 oz.: mix.

2. Sulphuric acid 1 fluid oz., lard 8 oz., oil of turpentine 1 oz.

OINTMENTS FOR SPAVINS AND WINDGALLS. (See RESOLVENT OINTMENTS, above.)

TAR OINTMENT. Equal parts of tallow and tar melted together.

OINTMENT FOR THRUSH AND CANKER.

1. Common verdigris ½ oz., calamine ½ oz., sulphate of zinc 1 dr., tar 3 oz.—BLAINE.

2. Blue vitriol 2 oz., white vitriol 1 oz., rubbed down and mixed with lard 2 ℔, tar 1 ℔; a pledget of tow covered with it to be introduced into the cleft of the frog every night, and renewed in the morning.—YOUATT.

3. *Thrush Paste.* Alum, blue vitriol, white vitriol, of each 1 oz.; rub them into a fine powder; melt 2 ℔ of tar with 1 ℔ of lard, and when getting cool, stir in the powder. —CLATER.

4. Verdigris 1½ oz. (or burnt alum 8 oz.) red lead 8 oz., treacle 4 ℔; boil to a proper consistence, and add 1 oz. of nitrous acid.—FERON.

5. Barbadoes tar 4 oz., sulphuric acid ½ oz.—SPOONER.

6. *In Canker.* Tar 4 parts, nitric acid 1 part; mix.— MORTON.

OINTMENT FOR FARCY BUDS. 1. Sublimate 1 oz., white arsenic ½ oz., yellow arsenic ½ oz., euphorbium ½ oz., oil of bays 4 oz.; mix.—LAMOTTE.

1. (*Topique Terrat.*) Corrosive sublimate 1 oz., white and yellow arsenic each ½ oz., oil of bays 4 oz.; mix with a gentle heat.

TURPENTINE OINTMENT. Common turpentine 1 part, lard 3 parts; melt together.—V. C.

VERDIGRIS OINTMENT. Verdigris in powder 1 part, common turpentine 1 part, lard 12 parts: for foul ulcers and tarsal ophthalmia.—MORTON.

OINTMENT FOR WARTS AND ANBURIES. Muriate of ammonia 2 dr., powdered savin 1 oz., lard 1½ oz.; to be applied daily.—BLAINE.

OINTMENT FOR CHRONIC VIVES.

1. Emetic tartar 2 dr., olive oil 1 dr.; rub together till smooth, and add lard 1 oz.

2. Iodide of potassium 1 dr., palm oil 1 oz.; rub together till quite smooth.—CLATER.

ASTRINGENT PASTE, for broken knees, and for wounds after the inflammation has subsided.

1. Powdered alum and pipe-clay, mixed with water to the consistence of cream. For broken knees it may be colored with bole and lamp-lack.—WHITE.

2. *Paste for Open Knee-Joint.*—Flour and stale beer, boiled to the consistence of paste, and colored as above. To be spread thick all round the joint, and covered with a pledget of tow, and ½ sheet of brown paper; and the leg of a cotton stocking drawn over the whole. The stocking to be covered with the paste, and enveloped with 2 calico bandages regularly applied.—TURNER.

STOPPING FOR THE FEET.

1. Cow-dung beaten with a fourth part of clay.—YOUATT.

2. Soft soap 4 oz., Barbadoes tar 16 oz., linseed meal 1½ lb.—WHITE.

3. Tallow and tar, equal parts, melted together.

4. Common tar 2 parts, soft soap 1 part, linseed meal q. s. To be spread over the sole of the foot ¼ of an inch thick, covered with a layer of tow, and a leather sole over all.—V. C.

HOOF OINTMENTS.

1. Equal parts of wax, olive oil, lard, veal suet, turpentine, and honey; melt the wax and lard with the oil by a gentle heat, remove from the fire, and add the honey and turpentine, stirring till cold; when intended to embellish the hoof as well as to soften it, it may be colored with lamp-black or ivory-black.—BOURGELAT.

2. Tallow 4 lb, beeswax 4 oz., tar ½ lb; melt slowly, remove from the fire, and when they begin to cool, stir together. A portion of pitch may be added when intended to fill fissures, &c.—BRACY CLARK.

COMPOSITION FOR SAND-CRACKS. Beeswax 4 oz., yellow resin 2 oz., common turpentine 1 oz., tallow ½ oz.; melt together; fill the cracks with the composition, and turn the horse out to grass.

SUPPLING LINIMENT FOR BRITTLE HOOF. Oil of tar 1 pint, fish oil 2 pints.—CLATER.

CHARGES.

The usual means of applying charges is to soften the compound by heat, and apply it with a large spatula to the part, as warm as the animal can comfortably bear it, and while warm to cover it with cut tow. They are used for old sprains of the loins, strains of the back sinews, wind-galls, &c. Cold charges are spread on cloth or leather, and renewed as they become dry.

1. *Simple charges.* Pitch 4 oz., turpentine 1 oz.—GASPARIN.

2. *For Strains of the Loins.* Pitch 4 lb, turpentine 6 oz., olive oil 4 oz.; melt together.—B. CLARK.

3. Burgundy pitch 4 oz., wax 4 oz., yellow resin 4 oz., common turpentine 1 oz.; melt together, and when it begins to thicken, stir in 1 oz. of bole.—WHITE.

4. Burgundy or common pitch 5 oz., tar 6 oz., wax 1 oz.; melt together, and when they are becoming cool, stir in ½ dr. of powdered cantharides.—YOUATT.

5. Pitch 3 lb, tar 1 lb, beeswax ½ lb; melt together.—CLATER.

6. Resin 2 oz., Burgundy pitch 4 oz., Barbadoes tar 2 oz., wax 3 oz., red lead 4 oz.—WHITE.

7. Pitch 8 oz., suet 4 oz., oil of turpentine 3 oz., tincture of cantharides 3 oz.—DELAFOSSE and LASSAIGNE.

8. *Cold Charge.* Bole ½ lb, white of egg and vinegar, to form soft paste, to be applied on double cloth or leather, and removed as it dries: for sprains in the back sinews.—BRACKEN.

9. Bruised leaves of elder, or cabbage, or mallow.—B. CLARK.

10. *Mercurial Charge.* B. pitch 1½ lb, wax 1½ lb; melt, and add, while cooling, 9 oz. of mercurial ointment previously mixed with 6 dr. of iodine.—Mr. S. FISHER.

11. *Soot Charge.* Common turpentine 4½ oz., soot 3 oz.; mix.—DELAFOND.

POWDERS FOR OUTWARD USE.

ASTRINGENT POWDERS; chiefly used for sprinkling greasy or ulcerated heels, after the inflammation has been subdued by poultices,—and in joint wounds.

1. Calamine (true) 4 parts, alum 1 part; mix.—Morton.

2. Burnt alum, dried sulphate of iron, and myrrh, equal parts.—V. C. (Comp. powder of alum).

3. Alum 1 dr., charcoal ½ oz., chalk 2 oz.—Blaine.

4. Sulphate of zinc, chalk lightly calcined, white pepper, in equal parts.—B. Clark.

5. Alum 4 oz., bole 1 oz.—White.

6. Oak bark 1 oz., verdigris 2 dr.—Blaine.

7. White vitriol 2 oz., oxide of zinc 1 oz.—White.

8. Prepared chalk 4 oz., sulphate of zinc 1 oz., charcoal 1 oz., Armenian bole 2 oz.—Spooner.

DETERGENT AND ESCHAROTIC POWDERS; for cleansing foul ulcers and repressing fungous or proud flesh. They should all be very finely powdered and well mixed.

1. Equal parts of white calcined vitriol and alum.—Bracken.

2. Bole 2 dr., blue vitriol or verdigris 1 oz.—White.

3. Red precipitate ½ oz., acetate of copper ½ oz., calamine ½ oz.—Blaine.

4. Red precipitate ½ oz., burnt alum 2 dr.

5. Blue vitriol 1 oz., alum 1 oz., white lead 1 oz.

6. Equal parts of verdigris and sugar of lead.—Clater.

7. Alum, dried sulphate of iron, and myrrh, equal parts: in joint wounds.

8. Alum, sulphate of iron, of zinc, and of copper, of each 1 oz.; muriate of ammonia ½ oz., camphor and saffron, of each 1½ dr.—Bouchardat.

STYPTIC POWDER. Alum, with an equal or double weight of flour.—White.

STYPTIC STONE. See Solutions, page 106.

SNEEZING POWDERS. The ingredients to be very finely powdered and mixed.

1. Asarabacca 4 dr., white hellebore 1 dr., mix, and keep in a bottle for use.—Bracken.

2. Snuff 1 oz., hellebore 1 dr., euphorbium 10 to 20 gr.—Peck.

3. *In Incipient Cataract.* Turpeth mineral 2 dr., asarabacca 4 dr.; mix, and apply as much as will lie upon a sixpence daily.—Bracken.

MEDICINES FOR NEAT CATTLE.

DRINKS OR DRENCHES.

Note.—The peculiar structure of the digestive organs in cattle renders it proper to give their medicines in a liquid form. For the same reason drenches should be given very slowly, so as to enter at once the third or fourth stomach. It is only in cases of hoven or *bloat*, that it is desirable to introduce medicine into the first stomach or *rumen*.

For Treatment of the Cattle Plague of 1865, *see p.* 131.

PURGING AND LAXATIVE DRENCHES. These are given when fever exists, or is threatened; to prevent downfalls of the udder; after calving, to prevent milk-fever; to remove undue accumulations in costiveness; in the first stage of red water, and jaundice; and in all inflammatory complaints.

1. Epsom salts 8 oz., sulphur 4 oz., ginger 2 dr., warm water a pint, linseed oil 12 oz.—SPOONER.

2. Epsom salts 6 or 8 oz., castor oil 8 oz., gruel 1½ pint, ginger ½ oz.

3. Glauber's or Epsom salts 16 oz. (or in bad cases with fever 24 oz.), caraways 1 oz., warm gruel a quart.—CLATER.

4. Castor oil from 16 to 24 oz., with gruel; but it is not to be depended on.

5. To No. 3, add 2 or 3 dr. of gamboge, or 4 dr. of aloes.

6. Sulphur 8 oz., ginger ½ oz., warm gruel a quart: in rheumatism, or joint-felon.—CLATER.

7. Common salt 6 oz., flour of mustard a tablespoonful, grated ginger or ground pepper, of either a teaspoonful, gin ¼ pint, water 2 pints.

8. Common salt 1 lb, warm water, or gruel, q. s. The last three are only proper where there is not much fever.

9. *In Red-water.* Sulphate of magnesia 8 to 16 oz., sulphur

2 to 6 oz., carbonate of ammonia ½ oz., ginger ½ oz., warm water q. s.; a fourth of this every 6 hours till the bowels are sufficiently acted on.—SPOONER.

10. *When the last does not operate.* Calomel 20 gr., yeast ½ pint.—HARRIS.

11. Aloes 4 to 6 dr., common salt 4 to 6 oz., ginger 1 to 3 dr., water a quart, anodyne tincture 2 oz.: in red-water. —WHITE.

12. *Cordial Purgatives.* Aloes 4 dr., Epsom salts 4 oz., ginger 1 dr., carminative tincture 2 oz., water 1 quart.—WHITE.

13. In the commencement of puerperal or milk fever: Epsom salts 6 or 8 oz., powdered croton seeds 20 to 30 gr., ginger 4 dr.; in 3 or 4 pints of gruel: repeat in 6 hours, if required, without the croton seeds.—BLAINE.

14. In locked-jaw: Barbadoes aloes 1½ oz., powdered croton kernel 10 gr., boiling water q. s.; given when cool. —CLATER.

15. *Mild laxative and tonic.* Epsom salts ½ ℔, sulphur 4 to 6 oz., ginger ½ oz., gentian ½ oz., warm water q. s.—EVESON.

16. In flatulent colic with costiveness: Aloes 1½ oz., carbonate of potash 3 dr., ginger ½ oz., warm water 1 pint, linseed oil 8 oz.—WHITE.

17. Palm oil 16 oz., Glauber's salt 12 oz., boiling water q. s.—PECK.

18. *Laxative drink for cows that are kept on hay.* Aloes 4 dr., ginger 1½ dr., water a quart, Epsom salts 6 oz., carbonate of soda ½ oz.; for one dose.—YOUATT.

FEVER DRENCHES, for fevers, colds, influenza, &c.

1. Tartar emetic 1 dr., digitalis ½ dr., nitre 3 dr.; mix, and give in a quart of gruel: in simple colds or catarrh.—CLATER.

2. Antimonial powder 2 dr., opium a scruple; rub together, and mix with thick gruel: after bleeding, in inflammation of the bladder.—WHITE.

3. In influenza, or epidemic (epizootic) colds; Nitre ½ oz., salt of tartar 1 oz., camphor 2 dr., valerian, liquorice, turmeric, of each 1 oz., mustard 2 oz., juniper berries 1 oz., gruel a quart.—SKERRET.

4. For the same: After bleeding and a laxative, give antimonial powder 2 dr., camphor 1½ dr., ginger 3 dr., laudanum ½ oz., in gruel.—WHITE.

5. In bad colds attended with fever; Nitre 1 oz, camphor ½ dr., tartar emetic ½ dr., in gruel.—PECK.

FEBRIFUGE TONIC DRENCHES.

1. Antimonial powder ½ dr., camphor 1 dr., Peruvian bark 1 oz., gruel, or decoction of arrowroot or starch, q. s. for 2 doses.—PECK.

2. In the decline of fevers and influenza: Emetic tartar ½ dr., nitre 2 dr., gentian 3 dr., chamomile 1 dr., ginger ½ dr.; pour on them a pint of boiling ale, and give when cool. —CLATER.

3. Emetic tartar ½ dr., gentian 2 dr., digitalis ½ dr., nitre ½ oz., spirit of nitric ether 4 dr., gruel q. s.

TONIC DRENCHES.

1. Cascarilla 3 dr., ginger 3 dr., carbonate of soda 2 dr., in gruel.—WHITE.

2. Gentian ½ oz., ginger 1 dr., Epsom salts 2 oz., warm gruel a pint.—CLATER.

3. Tartarized iron 1 dr., gentian 2 dr., ginger 1 dr., gruel 1 pint: after laxatives, in indigestion.

DRENCHES FOR INFLAMMATION OF THE LIVER.

After bleeding give—

1. Calomel 1½ dr., opium ½ dr., ginger 2 dr., thick gruel q. s. Six hours afterwards, give Epsom salts 1 ℔, sulphur 6 oz., linseed oil ½ pint, gruel q. s.—SPOONER.

2. Epsom salts 1 ℔, caraway ½ oz., Barbadoes aloes ¼ oz.; in a quart of warm gruel.—CLATER. After the yellowness appears, give—

3. Half of No. 2, with 20 gr. of calomel morning and night.—CLATER.

DRENCHES FOR JAUNDICE OR YELLOWS.

1. Opium 10 gr., calomel 10 gr., thick gruel q. s., at night, and the tonic drink (No. 2) in the morning.— CLATER.

2. Mr. Spooner says salts in ½ ℔ doses, with a little ginger, are generally sufficient.

3. Muriate of soda ½ oz., carbonate of soda ½ oz., turmeric 2 oz., Glauber's salts 6 oz., powdered gentian and chamomile 2 dr., gruel q. s.

4. Castile soap ½ oz., Venice turpentine ½ oz., ginger 3 dr., gentian 1 oz.; rub the soap and turpentine in a mortar, and gradually add a pint of water, and afterwards the ginger and gentian.—WHITE.

5. Castile soap 1 oz., salt 1 oz., Venice turpentine 1 oz., yolks of two eggs; mix together, and gradually add a strong decoction of barberry bark.

6. Powdered cumin seed, aniseed, and turmeric, each 2 oz., grains of Paradise and salt of tartar, each 1 oz.; mix. Slice 1 oz. of castile soap, to mix with 2 oz. of treacle. Pour a quart of boiling ale upon all the ingredients, and administer when lukewarm. To be repeated two or three times a day.

CLEANSING DRINKS, for cows after calving. These are often applied for, but are condemned as useless or hurtful by veterinarians of the new school. The following are some of the forms in use; probably a gentle laxative would be in most cases preferable:

1. Spermaceti, Irish slate, and birthwort, in powder, of each 1 oz., powdered aniseed 2 oz., liquorice powder 2 oz.; in linseed tea.

2. Aniseed, myrrh, birthwort, allspice, cumin seed, of each 1 oz., in a quart of gruel.—McEwen.

3. Juniper berries 3 oz., birthwort 2 oz., fenugreek 1 oz., spermaceti 2 oz., antimony 1 oz., saffron ½ oz., in a quart of warm ale.—Downing.

4. Resin, soap, of each ½ oz., spermaceti ½ oz., aniseed, caraway seed, of each 1 oz., ginger ½ oz., treacle 4 oz., warm gruel a quart.

5. 1 oz. spermaceti, 1 oz. birthwort, 2 oz. powdered bay-berries, 1 oz. myrrh; in juniper-berry tea.

LAXATIVE DRINK AFTER CALVING. Epsom salts 12 oz., aniseed 1 oz., olive oil 6 oz., gruel a pint or q. s.

DRENCH FOR STRANGURY. After laxatives and a clyster, give camphor 2 dr., spirit of nitrous ether ½ oz., tincture of opium ½ oz., nitre 1 oz., gruel a pint.—White.

DRENCHES FOR HOVEN OR BLOWN (flatulent distension of the paunch). It appears doubtful whether any liquid enters the paunch in these cases. More dependence is now placed on the introduction of a tube constructed for the purpose.

1. Ginger ½ oz., spirit of nitric ether 2 oz., oil of peppermint 30 drops, warm water a pint.—White.

2. Liquid ammonia, or spirit of hartshorn, ½ oz. to 1 oz. (1½ oz.—White), cold water 3 pints.—Moiroud.

3. Chloride of potash 4 dr., warm water 4 oz., ether 3 dr. The solution of chlorinated soda may be substituted for chloride of potash (Eau de Javelle).—CHARLOT.

4. Aloes 3 dr., pimento 2 dr., oil of turpentine 2 oz., spirit of hartshorn 1 oz., in gruel or warm water.—BLAINE.

5. Chloride of lime 2 dr., water a quart. Administer it by means of a stomach pump, and repeat in an hour if required.—YOUATT.

RHEUMATIC DRENCH.

1. Sulphur 8 oz., ginger ½ oz.; in gruel, every third day if necessary.—CLATER.

2. Antimonial powder 2 dr., Dover's powder ½ dr., aniseed 1 oz., thick gruel a pint; night and morning, the bowels having been opened by No. 1.—CLATER.

3. Rhododendron leaves 4 dr., water a quart; boil to a pint, strain, and add powdered gum guaiacum 2 dr., caraway-seeds and aniseed, each 2 dr., warm ale ½ pint.

ANTISPASMODIC DRENCH FOR LOCKED JAW. Camphor 1 dr., (rubbed with spirit), powdered opium 1 dr., thick gruel ½ pint.

CORDIAL CARMINATIVE DRENCHES. Drenches for indigestion, and colic without inflammation.

1. In indigestion: Salt 3 or 4 oz., carbonate of soda 2 dr., ginger ½ dr., anodyne tincture (below) 2 oz., water 10 or 12 oz.—WHITE.

2. The same: Aloes 4 dr., common salt 4 oz., ginger 2 dr., anodyne tincture 2 oz., q. s.

3. In colic: Salt 4 oz., aloes 3 dr., ginger 1 dr., opium ½ oz., water 1 pint, peppermint water 1 pint.

4. *Carminative.* Oil of turpentine 1 oz., tincture of opium 6 dr., spirit of nitric ether 2 oz., water 1 pint.—WHITE.

5. *Warm Cordial.* A bottle of red wine, extract of juniper 1 oz., powdered cinnamon ½ oz.—LEBAS.

6. *Mild.* Peppermint 2 oz., chamomiles ½ oz., hot water 5 pints; infuse, and give while warm.

7. Chamomile 2 oz., aniseed 1½ oz.; infuse in hot water, and strain; when cold, add ether 2 oz.—VATEL.

DRENCHES FOR BLOODY URINE.

Bloody Urine (Hæmaturia) and Red-water are often confounded, but are different diseases, and require a different treatment. Hæmaturia is distinguished by the presence of

actual blood in the urine, in a state of coagulation, and by great tenderness across the loins. It generally occurs in oxen of good condition. It is to be treated by bleeding, purgatives, stimulating applications to the loins, emollient drinks and opiates.—(SPOONER.) After bleeding, give one of the following drenches:

1. Epsom salts 6 to 8 oz., water a quart, castor oil 4 to 6 oz.—WHITE. Or,

2. Linseed oil 1 pint, gruel 1 pint, caraways 2 dr., Epsom salts 8 oz. (in warm water ½ pint), tincture of opium 2 dr. Or either of the laxative drenches for red-water, below.

3. After the above, when the pain and difficulty have abated, but the water continues bloody, give—Catechu 2 dr., opium ½ dr., alum 3 dr., gum arabic ½ oz., water ½ pint; simmer for a few minutes, and add ½ pint of ale. Repeat if required.

4. In obstinate cases: Oil of juniper ½ oz., oil of turpentine 1 oz., laudanum 1 oz., in a pint of linseed tea; at first twice, and afterwards once a day.—RUSH.

5. Three quarters of a pint of black beer, and 2 oz. Irish slate.—KNOWLSON.

DRENCHES FOR RED-WATER.

This commonly attacks milch cows, and appears generally to arise from the nature of the pasture. Moderate bleeding is recommended, but is less necessary than in hæmaturia. Laxatives should then be given.

1. Epsom salts 8 to 12 oz., sulphur 2 to 4 oz., carbonate of ammonia ½ oz., ginger ½ oz., warm water 4 pints; give a fourth part every six hours till the bowels are acted on.—SPOONER. Or,

2. Glauber's salts 12 oz., carbonate of soda ½ oz., nitre ¼ oz., sugar 1 oz., powdered caraways ½ oz., in a quart of gruel. Or the Purging Drenches, No. 1, 2, or 3. After the bowels are well opened, give astringents or mild stimulants.

3. The laxative drench, No. 11, page 122.—WHITE. This is WHITE'S drench for red-water, No. 1. To be followed by drenches of whey.

4. *Astringent.* Powdered oak-bark ½ oz., catechu 2 dr., opium 10 gr., gruel 1 pint.—CLATER.

5. Catechu 2 dr., mucilage 4 oz., lime-water 6 oz.—BLAINE.

6. Laudanum ½ oz., sugar of lead ½ dr., catechu 4 dr., gruel 1 quart.—WHITE.

7. After laxatives: Ginger, gentian, each 1 dr., spirit of nitrous ether 1 oz., gruel q. s.; twice a day.—SPOONER.

8. Powdered oak-bark 1 oz., charcoal 1 oz., bole 2 oz., in a quart of new milk.—PYE.

9. Catechu 1½ oz., alum 1½ oz., diapente 2 oz., Locatelli balsam 2 oz., warm gruel 3 pints.—MCEWEN.

10. Dragon's blood 2 oz., rust of iron 1 oz., nitre 3 oz., oil of turpentine 2 oz.; mix: for 2 doses, in gruel.—DOWNING.

11. Sulphuric acid 1 dr., tincture of opium ½ oz., treacle 4 oz., warm gruel 4 quarts; daily, for a week.—BLAINE.

DRINK FOR ACUTE DIARRHŒA. Sulphate of soda, sulphate of magnesia, of each 2 oz., ipecacuanha ½ dr., sulphate of iron 6 gr.—BLAINE.

FOR CHRONIC DIARRHŒA. Calomel ½ dr., aloes 1 dr., gentian 2 dr., opium 5 gr., decoction of chamomiles 1 pint.—BLAINE.

ASTRINGENT DRENCHES, for dysentery (scouring rot) or lax.

1. After purging drenches: Prepared chalk 2 oz., oak-bark 1 oz., catechu ½ oz., opium 2 scruples, ginger 2 dr., warm gruel 1 quart.—CLATER.

2. Two quarts of alum whey.

3. First give blue pill 2 or 3 dr., rhubarb 3 dr., castor oil 4 oz., gruel 1 pint, well stirred before giving it; repeat this 3 or 4 mornings; then give—thick starch (made with 4 oz. of starch) 3 or 4 pints, tincture of opium 2 dr., ginger 3 dr., catechu ½ oz.—WHITE.

4. Mutton suet 1 ℔, new milk 2 quarts; boil, and add opium ½ dr., ginger 1 dr.—CLATER.

5. *Cordial Astringent Drench.* (After the laxative drench No. 2.) Catechu ¼ oz., allspice ¼ oz., caraways ½ oz., ale ½ pint, water ½ pint.—WHITE.

6. Decoction of aloes, with prepared chalk.—TUSSER.

7. Decoction of wormwood a quart, gum Arabic 2 oz., aromatic confection 1 oz., catechu 2 dr.; with linseed tea, repeated every 6 days for 3 times.—RAWLINGS.

8. Sheep's heart, liver, and lights, all chopped up together.—SUMNER.

EXPECTORANT AND COUGH DRENCHES, IN HOOSE OR CATARRH. (For Hoose, in Calves, see page 134).

1. Bruised liquorice 2 oz.; boil in a quart of water to a pint, strain, and add powdered squill 2 dr., gum guaiacum 1 dr., tincture of tolu 4 dr., honey 2 oz.—CLATER.

2. Balsam of sulphur 2 oz., Barbadoes tar 1 oz., yolks of 2 eggs, honey 4 oz., salt of tartar ½ oz., oil of aniseed 1 dr., elecampane 1 oz., gruel 1 quart: in chronic coughs.

3. Fresh squill 2 oz., garlic 2 oz., vinegar 24 oz.; digest for a day with a gentle heat, strain and press, and boil the liquor with 24 oz. of treacle: for 6 doses, in chronic cough.

4. For recent coughs: Digitalis 20 gr., emetic tartar ½ dr., nitre 3 dr., squill 1 dr., opium 20 gr., gruel 1 pint.

5. Boil 4 oz. Iceland moss and 1 oz. liquorice root in 4 quarts of water for a quarter of an hour, and strain; add to the liquor 1 oz. nitre, cream of tartar 2 oz. In hoose from cold, if inflammation of the lungs and fever be present, bleed before giving the drink. See also FEVER DRENCHES.

If the disease be connected with worms in the air-passages, give the following:

Worm Drench.—For cough from worms: Oil of turpentine 2 oz., sweet spirit of nitre 1 oz., laudanum ½ oz.; mix and give in a pint of gruel.—CLATER.

DRENCHES FOR THE EPIDEMIC, affecting the feet and mouth, and attended with a low fever.

For treatment of the Cattle Plague of 1865, see p. 131.

1. Glauber's salts 1 lb, treacle 4 oz., sulphur 4 oz., aniseed ½ oz., cream of tartar 2 oz., warm water 3 pints, give it new-milk warm. The above is for a full-grown beast. The mouth to be washed with a strong solution of blue vitriol, burnt alum, and vinegar. If the feet crack, apply a mixture of equal parts of muriatic acid and water.

2. Some cattle-masters give common salt in gruel, with great success.—BLAINE.

3. After a mild dose of salts: sweet spirits of nitre 1 oz., ale yeast 6 or 8 oz.—BLAINE.

4. Epsom salts 8 oz., sulphur 2 oz., nitre ½ oz., ginger ¼ oz.; half of this to be given in warm water, with 1 oz. of sweet spirits of nitre. Repeat daily. When the bowels are

properly relaxed, and the fever reduced, but much weakness remains, give the following:

5. *Tonic Drench.* Gentian 4 dr., ginger 2 dr., sulphate of iron 2 dr., sweet spirits of nitre 1 oz., warm water q. s.; wash the mouth with the *lotion*, page 134; dress the feet, after paring and poulticing, with equal parts of tincture of myrrh and butter of antimony, and afterwards apply the *astringent powder*, page 140.—SPOONER.

6. Linseed oil a pint, oil of turpentine 8 oz., aloes ½ oz., ginger ¼ oz., laudanum 1 oz.; mix. For 2 doses, to be given with gruel.

REMEDIES FOR THE EPIZOOTIC PNEUMONIA. In the fatal form of this disease which prevailed some years ago, the following treatment is said to have proved effectual; Bleed freely: then administer ½ pint of brandy every 2 hours. Mr. Jeckyll gives in pleuro-pneumonia, when a tonic is indicated, ½ oz. of the following solution every 8 hours: Sulphate of copper 1 part, water 4 parts; dissolve, and add ammonia until it begins to precipitate.

MURRAIN DRENCHES.

1. Sweet spirit of nitre ½ oz., laudanum ½ dr., solution of chloride of lime 2 dr., prepared chalk 1 oz., mix, and give in a pint of warm gruel.—CLATER.

2. Cascarilla powder 2 oz., spirit of nitrous ether ½ oz, liquid acetate of ammonia 4 oz., beer yeast 8 oz.; every 4 hours.—BLAINE.

3. Opiate confection 1 oz., liquid acetate of ammonia 2 oz., water 1 quart; for one dose.—VATEL.

4. *Tonic.* Calumba 2 dr., canella 2 dr., ginger 1 dr., sweet spirit of nitre ½ oz., thick gruel 1 pint.—SPOONER.

5. *Ceylon Remedy.* A small piece of lard the size of a walnut. Said to be used with perfect success.

6. *To Prevent Murrain.* Myrrh 1 oz., Epsom salts 2 oz., sulphur 1 oz., liver of antimony ½ oz., diapente 1 oz.: in rue tea after bleeding.—DOWNING.

FOR THE DISTEMPER.

Warm tar water is much recommended by some. It should be given in doses of 3 quarts 3 or 4 times a day, gradually diminishing the quantity.

DIURETIC DRINKS.

1. Common turpentine ½ oz., ginger 2 dr.; mix with a

little treacle, and add gradually spirits of nitrous ether 1 oz., gruel a pint.—CLATER.

2. *Tonic Diuretic.* Common turpentine 4 dr., ginger 2 dr., gentian 2 dr., tartrate of iron 1 dr.; rub together with a little treacle, and add gradually 1 oz. of sweet spirit of nitre.

BULLING DRENCHES. These are strongly condemned by modern veterinary writers; those drenches, at least, which contain cantharides.

1. Aniseed, grains of paradise, bay-berries, of each 1 oz.; cantharides in fine powder 20 to 30 gr.; to be given in a quart of milk.

2. Black hellebore ½ oz., capsicum 2 dr., birthwort ½ oz., bay-berries 1 oz., cantharides 20 gr.; in a quart of warm ale.—DOWNING.

3. A quart of milk from a cow in season.—CLATER.

4. Powdered cantharides 20 gr., aniseed 2 oz., black hellebore ½ oz.; in ale, gruel, or milk.—PECK.

DRYING DRENCHES, for drying a cow's milk. Bleed the night before, and give the drink, warm, in the morning.

1. Boil 6 dr. of alum in milk, and strain.—WHITE.

2. Alum 6 ℔, bole 2 ℔, cream of tartar or red tartar 1 ℔; mix. Give from 6 to 9 oz., in stale beer; or in gruel with ½ pint of vinegar.

3. Roche or common alum 4 oz., dragon's blood ½ oz., turmeric 1 oz., in a pint of rennet-water, and a pint of vinegar.—DOWNING. [These large doses of alum, though often given, are not regarded as necessary or proper by modern veterinary writers.]

DRENCH FOR THE BITES OF VIPERS. Olive oil 2 pints, spirits of hartshorn 1 oz.; mix.

ALUM WHEY. Boil ½ oz. of alum in 2 quarts of milk for ten minutes and strain.

ANODYNE CARMINATIVE TINCTURE; and Tincture of Pimento. See MEDICINES FOR HORSES.

ALTERATIVE POWDERS.

1. Sulphur 4 oz., black antimony 1 oz., Æthiops mineral ½ oz., nitre 2 oz.; mix, for 4 doses: to be given daily in gruel.

2. *Alterative Tonic.* Add to the last 2 oz. gentian and 1 oz. ginger; and make 6 doses.

FEVER POWDERS IN INFLAMMATION, &c.

1. Antimonial powder a scruple, camphor ½ dr., nitre 1 oz.; mix; give twice a day in gruel.—PECK.

2. Peruvian bark 16 oz., nitre 24 oz.; for 16 doses.

CORDIAL POWDERS.

1. Black mustard ½ oz., flowers of sulphur 1 oz., aromatic powder (see HORSE POWDERS) 1 oz., fenugreek 4 oz., common salt 16 oz., a large pinch on a slice of bread.—MATTHIEU.

2. *Cow Spice.* As Horse Spice, No. 2, page 76. Or,

3. Powdered turmeric, liquorice, aniseed, and diapente, each 1 oz.

CLYSTERS.

1. Salt 1 ℔, warm water a gallon.

2. Linseed oil 8 oz., Epsom salts 8 oz., gruel 3 quarts.

MASTICATORIES.

1. Bruised garlic 4 cloves, salt a tablespoonful, ground pepper 1 oz., honey 4 oz. Boil for a short time in a glass of vinegar, immerse in it a piece of linen, and roll it up. Keep it in the animal's mouth for an hour, night and morning. *Antiputrescent;* in epizootic maladies, and in ulcers of the mouth.—J. ROBINET.

2. Bruised mustard and pepper, each ½ oz., rolled up in linen, and sprinkled with vinegar: to be kept in the mouth not more than half an hour, morning and evening, in epizootic diseases.

TREATMENT OF THE CATTLE PLAGUE OF 1865, OR RINDERPEST.

1. *Vaccination.* This has been recommended, on the theory that the Cattle Plague is analogous to or identical with small-pox. Vaccination with the lymph of Cow-pock may then be resorted to as a prophylactic. It may be practised on the udder or vulva of a cow. It has already been resorted to very extensively, but with doubtful success. (January, 1866.)

(Whether the plague be Small-pox or a form of enteric fever, or a disease *sui generis*, the precautions of separating infected animals from the remainder of the herd are

equally necessary. The diseased carcase must be buried or destroyed at a distance. The most scrupulous cleanliness must be observed in the sheds and yards, and provisions made for ventilation and abundant supply of water, with the counteraction of all noxious smells and emanations. When the cattle have sickened with the disorder, it cannot be checked, but like other contagious diseases, must run its course. The treatment can only be palliative. Among innumerable recipes we select the following as the most likely to be useful. Their application must be guided by the circumstances of the case.)

2. *The Laxative Treatment.* Linseed oil 6 to 12 oz., with turpentine 1 to 2 oz.; the dose to be repeated once or twice. This dose, as the others, may be given in warm gruel.

3. *The Astringent Treatment.* Applicable in later stages, when there is often much diarrhœa. Chalk and opium (Dr. LETHEBY), or laudanum 2 oz., comp. infusion of catechu 1 pint, in a warm mash, with 2 or 3 tablespoonfuls of charcoal. —MALTON AGRICULTURAL ASSOCIATION.

4. *The Saline Treatment.* Useful in moderating the fever, and generally in mild cases. Table salt ½ lb, Epsom salts ¼ lb, brimstone ¼ lb, ground ginger 2 oz., nitre 2 oz., old ale 1 quart (Dr. ALLNATT); or chlorate of potash 2 dr. to ½ oz., thrice daily.—Mr. F. BUCKLAND.

5. *The Mixed Saline Treatment.* Adopted in imitation of Dr. Stevens' treatment of cholera. Chloride of sodium 3 oz., nitre 2 oz., carbonate of potash 3 oz., chlorate of potash 2 oz., sulphate of magnesia 4 oz., old ale 2 quarts. (Dr. ALLNATT.) It would be as well to divide the dose into 3 or 4. Bicarbonate of soda 1 oz., common salt 1 oz., chlorate of potash ½ oz., Rochelle salt 1 oz.; in a gallon of water: 1 pint every second hour or oftener.—Dr. TUCKER.

6. *The Diaphoretic Treatment.* Useful as the last, in mild cases. Sweating to be promoted by rubbing, wrapping in blankets, or vapor baths. Sweet spirit of nitre 1 to 2 oz., and spirit of Mindererus 1 to 2 oz., in 2 or 3 quarts of linseed-meal gruel. Will also act as a diuretic.—MALTON ASSOCIATION.

7. *The Stimulant Treatment.* 16 oz. of linseed oil and a "mutchkin" of whiskey. (Professor DICK.) Large draughts of old ale recommended by some. 6 dr. of carbonate of ammonia 3 times a day. (Professor DICK.) 12 to 15 gr. of camphor, with 20 gr. of carbonate of ammonia in gruel, thrice daily.—Dr. COPLAND.

8. *The Tonic Treatment.* A dr. each of the tincture of perchloride of iron and dilute hydrochloric acid in a quart of linseed tea or gruel, thrice daily. (Dr. COPLAND.) ½ oz. of sulphate of iron, twice daily. (Dr. TUCKER.) 1½ oz. of powdered cinchona. (Dr. SMART.) Strong muriatic acid 2½ oz., strong nitric acid 1½ oz., sulphate or chloride of iron 1½ oz., water to make 1 quart; one ounce of this in a gallon of water for a dose.—MALTON ASSOCIATION.

9. *Arsenical Treatment.* Adopted by the homœopathists with their usual want of success. $\frac{1}{10}$th to $\frac{1}{5}$th of a grain of arsenic for a dose, every 2 hours, to alternate with $\frac{1}{50}$th gr. of phosphorus. Vapor baths also recommended, and immense quantities of old ale given.—Dr. KIDD.

MEDICINES FOR CALVES.

PURGATIVE DRENCHES.

1. Epsom salts 1 oz. to 2 oz., according to the age and size of the calf; dissolve in ½ pint of gruel, and add 20 gr. of ginger, and 3 drops of essence of peppermint.—CLATER.

2. Salts 1½ oz., castor oil 2 oz., ginger 10 gr., caraway 2 dr., gruel ½ pint.

3. In costiveness, and accumulation in the paunch and stomach: dissolve 2 oz. of Epsom salts in 2 or 3 quarts of water, or 4 oz. in a gallon, according to the age of the calf, and throw it in gently by means of a stomach-pump.

4. *Laxative.* Epsom salts 2 or 3 oz., carbonate of soda 2 dr., water 6 or 8 oz., ginger 1 dr.; mix. After it has operated, give the cordial, No. 3, below.—WHITE.

DRENCHES FOR DIARRHŒA, OR CALVES' CORDIAL.

1. YOUATT'S *Cordial.* Prepared chalk 2 oz., catechu 1 oz., ginger ½ oz., opium 1 dr., peppermint water 1 pint; dose for a calf, from 2 to 4 tablespoonfuls.

2. Prepared chalk 2 dr., opium 10 gr., catechu ½ dr., ginger ½ dr., essence of peppermint 5 drops; mix, and give twice a day in ½ pint of gruel.—CLATER.

3. Caraway ½ oz., ginger ½ dr., subcarbonate of soda 1 dr., brandy or gin 1 oz., water 8 oz.—[WHITE'S *Cordial.*]

4. Half a bottle of Dalby's Carminative.

5. Suet boiled in milk ½ pint, opium 5 gr., alum 5 gr., prepared chalk ½ oz.; mix.

6. If No. 2 fails: Dover's powder 2 scruples, aromatic

powder 1 dr., kino ½ dr.; give it night and morning, with 1 oz. of arrowroot boiled in a pint of water.—CLATER.

INFLAMMATORY DISORDERS. Bleed; give 2 to 6 oz. Epsom salts. [Give to a calf of six months old ¼ the dose for cattle; at a year and a half, ½ the dose.—SPOONER.]

MEDICINE FOR PILES IN CALVES.

Oil of vitriol 15 drops, tincture of opium ½ oz.—PECK.

SOLUTION OF POTASH, FOR CORDS, &c. Subcarbonate of potash 2 oz., fresh lime-water 8 oz. To correct acidity in the stomach, give 1 or 2 teaspoonfuls in gruel; the first dose to be given with an ounce or two of Epsom salts in ½ a pint of thin gruel. If the disorder is attended with griping pains, add a tablespoonful of anodyne carminative tincture.—WHITE.

ALUM WHEY. See page 130.

TO PROMOTE THE FATTENING OF CALVES.

Aniseed ¼ lb, fenugreek ¼ lb, linseed meal 1 lb; make it into a paste with milk, and cram them with it.

Fattening Powder. Common salt with a little carbonate of soda; a small quantity added to the food promotes fattening, and prevents scouring, &c.

HOOSE, OR COUGH FROM WORMS IN AIR-PASSAGES.

1. ½ pint lime-water every morning, and a tablespoonful of salt every afternoon, to each calf.—MAYER.

2. Linseed oil 4 oz., oil of turpentine 1 oz., oil of caraways 20 drops; repeated once or twice at intervals of 10 days. This dose for calves of 6 to 10 months old.—DICKENS.

3. A tablespoonful of oil of turpentine, a little sweet oil, and 6 or 8 oz. of warm water.—WHITE.

External Applications for Neat Cattle.

LOTIONS OR WASHES. (See also EMBROCATIONS.)

LOTIONS FOR CANKER IN CALVES.

1. Alum 1 oz., water 8 oz., tincture of myrrh 1 oz., honey of roses 1 oz.

2. Equal parts of tincture of myrrh and water.—CLATER.
3. Alum ½ oz., water 1 pint, tincture of myrrh 1 oz.

LOTION FOR COW-POCK. Sal ammoniac ¼ oz., white vinegar ½ pint, camphorated spirit 2 oz., Goulard's extract 1 oz.; mix.—CLATER.

LOTIONS FOR SLIGHT BRUISES.

1. Extract of lead ½ oz., vinegar 4 oz., soft water 1 pint. —WHITE.
2. Acetate of ammonia 4 oz., water ½ pint, spirit of camphor ½ oz.

DISCUTIENT LOTION, for dispersing tumors. Bay salt 4 oz., vinegar 1 pint, water 1 quart, oil of origanum 1 dr.; rub the oil with the salt, and gradually add the others.

LOTIONS FOR STRAINS.

1. Bay salt 4 oz., oil of origanum 1 dr.; rub together, and add vinegar ½ pint, spirit of wine 2 oz., water 1 quart.
2. Common salt 1 oz., sal ammoniac 1 oz., water 1 pint.

LOTIONS FOR FOUL IN THE FOOT. After poulticing and removing loose horn, apply—

1. Butter of antimony, or
2. Strong solution of alum.
3. Solution of sulphate of copper.
4. (When the above are not sufficient.) Dissolve 2 dr. of corrosive sublimate in 12 oz. of water.—WHITE.

LOTIONS FOR WOUNDS.

1. Tincture of myrrh and aloes.
2. For proud flesh; strong solution of sulphate of copper.
3. For offensive wounds: chloride of lime 1 oz., water 1 pint; mix well, and strain.

LOTION FOR BULL-BURNT. Goulard's extract 1 oz., spirit of wine 2 oz., water ½ pint.

LOTION FOR BLAIN IN THE MOUTH. After lancing the bladder, apply a saturated solution of salt in water.—YOUATT.

LOTION FOR THE MOUTH, in ulceration during the epidemic. Alum 1 oz., sulphate of zinc ½ oz., warm water 1 pint, treacle ¼ lb.—SPOONER.

WASHES FOR DESTROYING VERMIN. They are all poisonous.

1. Stavesacre seeds 4 oz., water 4 pints; boil to 2 pints, and apply it daily.—PECK.

2. Sublimate 2 dr., spirit of wine 2 oz., water 1 pint.—CLATER.

3. Stavesacre 4 oz., white hellebore root 2 oz., water 1 gallon; boil to half; apply with a sponge.

LOTION FOR MANGE. Corrosive sublimate 2 dr., muriatic acid ½ oz., water 12 to 16 oz.: mix. In obstinate cases only.—WHITE. (See LINIMENTS.)

EYE-WATERS, OR COLLYRIA.

1. White vitriol a scruple, spirit of wine 1 dr., water a pint.—CLATER.

2. Sugar of lead 10 to 20 gr., soft water, 8 oz.—V. C.

3. *Sedative Eye Drops.* Powdered digitalis 1½ oz.; infuse in a pint of Cape wine for a fortnight, and filter: a few drops to be introduced into the eye twice or thrice a day.—SPOONER.

4. Extract of lead 2 dr., wine of digitalis (No. 3) 2 dr., tincture of opium 2 dr., water a pint.

EMBROCATIONS AND LINIMENTS.

STRONG EMBROCATION, for deep-seated strains, &c.

1. Oil of origanum ½ oz., oil of turpentine ½ pint, sweet oil 1½ pint, powdered cantharides 1 oz.—CLATER.

2. Olive oil 4 oz., oil of turpentine, 1 oz., water of ammonia 1 oz. (For strains and bruises, after the inflammation has subsided.)—WHITE.

CROTON LINIMENT. Bruised croton seeds 1 part, oil of turpentine 8 parts. Macerate for 14 days, and strain. It irritates the skin powerfully; for general purposes it requires to be diluted with olive oil.—MORTON.

MUSTARD EMBROCATION. Flour of mustard 4 oz., oil of turpentine 2 oz., water of ammonia 2 oz.—WHITE.

RHEUMATIC EMBROCATIONS.

1. Olive oil 2 oz., strong water of ammonia 1 oz., marshmallow ointment 1 oz.

2. Neatsfoot oil 4 oz., camphorated oil 1 oz., oil of turpentine 1 oz., laudanum 1 oz., oil of origanum 1 dr.—CLATER.

3. Sweet oil 4 oz., oil of turpentine 2 oz.; mix.—WHITE.

EMBROCATIONS FOR GARGET, or Downfall of the Udder.

1. Oil of elder 4 oz., water of ammonia ½ oz., Mindererus' spirit 1 oz., camphorated oil 2 oz.

2. Olive oil 3 oz., oil of turpentine 1 oz., camphor 2 dr.—WHITE.

3. Soft soap 8 oz., oil of bays 8 oz., oil of turpentine 8 oz., spirit of camphor 4 oz. (See also OINTMENTS, below.)

DRIFFIELD OILS. Barbadoes tar 1 oz., linseed oil 1 lb, oil of turpentine 3 oz., oil of vitriol 1 oz., or 1 oz. (by weight).

LINIMENT FOR MANGE. Sulphur vivum or flowers of sulphur 4 oz., train oil 12 oz., oil of turpentine 4 oz.; mix.

LINIMENT FOR SORE THROATS. Oil of turpentine 1 oz., sweet oil 1 oz., water of ammonia 2 oz.—WHITE.

BLISTERING LINIMENT. Cantharides bruised 1 oz., oil of turpentine 8 oz.; digest 14 days, and strain. To be applied by friction on the skin.—YOUATT.

LIQUID CAUSTIC. Butter of antimony alone, or mixed with an equal quantity of tincture of myrrh.

LIQUID SNUFF. Alum, sulphate of zinc, capsicum, of each 1 oz.; camphor 2 dr.; pulverize and macerate in 32 oz. of strong vinegar and 1 oz. of turpentine; shake up when used, and introduce a teaspoonful into the nostrils, to promote a discharge for the relief of inflammation of the chest.—MATTHIEU.

OINTMENTS.

BLISTERING OINTMENTS.

1. Resin cerate 1 oz., cantharides finely powdered 3 dr., oil of turpentine 2 dr.; for setons.—CLATER.

2. Lard 12 oz., resin 4 oz.; melt together, and when sufficiently cool, add oil of turpentine 4 oz., powdered cantharides 5 oz.; stir till cold: to be rubbed in after removing the hair.—CLATER.

OINTMENT FOR MANGE, LICE, &c. (See also LOTIONS, above.)

1. Sulphur 1 lb, common turpentine 4 oz., mercurial ointment 2 oz., linseed oil a pint. Melt the turpentine with the oil, and when nearly cold, stir in the sulphur, and afterwards the mercurial ointment.—YOUATT. *Note.*—Cattle are easily salivated, and greatly weakened by it. Mercurials should therefore be used with great caution.

2. Sulphur 1 lb, strong mercurial ointment 2 oz., common turpentine ½ lb, lard 1½ lb.—CLATER.

3. *French Liniment.* Olive oil a pint, sulphur 4 oz., heat till the oil becomes colored by the sulphur; remove from the fire, and when nearly cold, add 4 oz. of oil of turpentine, apply with a feather.

4. Lard 2 lb; melt and add oil of turpentine 8 oz., sulphuric acid 2 oz., sulphur vivum 8 oz.; stir till cold.

5. *Mange Liniment.* Sulphur vivum finely powdered 4 oz., train oil 12 oz., oil of turpentine 4 oz.; mix.—WHITE.

IODINE OINTMENT, for Empyema.

Rub together 1½ dr. of iodine, and 1 dr. iodide of potassium, with a few drops of water, then add 3 oz. of strong mercurial ointment and ½ oz. of powdered camphor. To be rubbed over the chest every night till it causes an exudation, then occasionally to keep it up.

GARGET OINTMENT, for Downfall of the Udder.

1. Soft soap 1 lb, mercurial ointment 2 oz., camphor (powdered with spirit) 1 oz.; mix: give first as a laxative, then a fever or diuretic drink.

2. Green elder ointment 2 oz., water of ammonia ¼ oz.

3. Beat fox-glove leaves with twice their weight of whey butter; to every pound add 1 oz. of sal ammoniac, 1 oz.

of turpentine, and ½ oz. of bole; mix, and apply two or three times a day.—DOWNING.

4. Spirit of camphor 1 oz., mercurial ointment 1 oz., elder ointment 8 oz.—YOUATT.

5. In obstinate cases. Iodide of potassium 1 part, lard 7 parts. To be rubbed in once daily.—SPOONER.

FOOT OINTMENT (for all domestic animals). Equal parts of tar, lard, and resin, melted together.

OINTMENT FOR ULCERS ABOUT THE JOINTS. Equal parts of basilicon and citrine ointments.—CLATER.

HEALING AND CLEANSING OINTMENT. Lard 2 ℔, yellow resin ½ ℔; melt together, and when it begins to cool, add calamine in powder ½ ℔.

APPLICATION TO WOUNDS. Mix the whites of eggs with flour to a proper consistence. Applied over the part, it soon dries, and shields it from the air.

DIGESTIVE OINTMENT. 1. Lard, common turpentine, of each 4 oz.; melt, and add 1 oz. powdered verdigris.—WHITE.

2. Boil leaves of black hellebore with an equal weight of lard, until the leaves are crisp; strain, and add an equal weight of common turpentine. [A similar ointment made with ivy leaves is likewise very stimulating.]

OINTMENT FOR FOUL IN THE FOOT, OR LOW.

1. Melt 4 oz. of lard with 4 oz. of common turpentine, and add 1 oz. of finely powdered sulphate of copper, stirring until cold.—WHITE.

2. Melt together equal weights of soft soap and common turpentine.—SKERRET.

OINTMENT FOR CANCEROUS TUMORS. Hydriodate of potash ¼ oz., hot water ¼ oz.; dissolve, and mix with 2 oz. of lard.

CHARGE FOR OLD STRAINS. Burgundy pitch 4 oz., common pitch 4 oz., wax 2 oz., tar 6 oz.; apply hot, and cover with cut tow.

SETONS.

1. *Common.* A piece of cord or coarse tape; or horse-hair and tow platted together.

2. *Irritating.* Root of common dock; or of black hellebore.

3. Cotton cord soaked in Morton's cantharidal solution (page 94).

PASTE FOR STOPPING BLEEDING. Equal quantities of white, green, and blue vitriol, flour, and bole; beaten up with fresh nettles and a little vinegar.

ASTRINGENT POWDER FOR SORE FEET, &c. Sulphate of copper ½ oz., prepared chalk 2 oz., powdered alum ½ oz., bole 1 oz.; rub together.—SPOONER.

MEDICINES

FOR

SHEEP AND LAMBS.

THESE are best given in a liquid form, and should be carefully and slowly administered. Sheep generally require one-sixth (or from one-eighth to one-sixth) of the doses given to cattle.

PURGING DRENCHES.

1. Epsom salts 2 oz., powdered caraway ¼ oz., warm, thin gruel sufficient to dissolve the salts. The editor of CLATER says that this is the best purging drink that can be used. For *Lambs* give a fourth of this, and repeat in 6 hours if necessary.

2. Epsom salts 1½ oz. or 2 oz., ginger 1 dr., treacle 1 oz., hot water 4 oz.

3. Castor oil 2 oz., ginger and salt of tartar, of each 2 scruples, moist sugar a spoonful, gruel q. s.—MCEWEN.

4. Epsom or Glauber's salts from 1 to 2 oz., common salt a teaspoonful, boiling water sufficient to dissolve the salts, and a little gruel. A teaspoonful of tincture of ginger or of pimento, or of anodyne carminative tincture (see p. 90), may be added.

5. Sulphur ¼ oz., Epsom salts 1 oz., common salt a teaspoonful, thin gruel ¼ pint.

6. Linseed oil 2 or 3 oz., croton oil 2 or 3 drops, warm gruel q. s.

7. *For Lambs.* Epsom salts 2 to 4 dr., ginger ½ dr., in gruel.—SPOONER.

8. *For Sheep on the first attack of Small-pox.* Epsom salts 2 oz., ginger ½ dr., in chamomile tea or infusion of gentian (or with 1 dr. of powdered gentian or chamomile).—WARNECKE.

FEVER DRENCH. Powdered digitalis 20 gr., emetic tartar 10 gr., nitre 2 dr. Twice a day, mixed with gruel.—CLATER.

TONIC DRENCHES.

1. *General Tonic.* Gentian 2 dr., calumba 1 dr., ginger ½ dr., all in powder; tincture of orange-peel 1 dr., gruel 4 oz.; for one dose.—CLATER.

2. *In the last stage of Fever.* Gentian 1 dr., ginger 20 gr., spirit of nitrous ether 1 dr., tincture of cardamom 20 drops, in gruel.

3. *For Debility and Indigestion, after a purgative.* Gentian, caraway, each 1 oz., calumba and ginger, of each 1 oz. (all sliced or bruised), boiling water a quart; infuse till cool, and strain. Give a tablespoonful daily, with the same quantity of gruel.—CLATER.

DRENCHES FOR RED-WATER. The pasture should be changed for shorter, the animal bled, and the bowels kept open with the above purging drinks. If these means do not remove the disease, give one of the following:

1. Epsom salts 6 oz., nitre 2 oz., bole ½ oz., hot water 3 pints, oil of turpentine 4 oz.; mix, and give 3 or 4 tablespoonfuls (from a horn that will measure that quantity), shaking the bottle well before each dose is poured out.

2. Powdered catechu 30 gr., alum 20 gr., ginger 20 gr., decoction of oak-bark 4 oz., for a dose.

3. Olive oil 1 oz., oil of turpentine 1 oz., thick gruel ¼ pint.

FOR EXTERNAL RED-WATER. (Vesicles on the skin, containing a reddish fluid.)

1. Sulphur 2 to 3 dr., in gruel, once or twice a day. If it continues, give—

2. Epsom salts 1 oz., gruel sufficient to dissolve it.—Sir JAMES MACKENZIE.

DRENCHES FOR DIARRHŒA, SCOUR, &c., IN LAMBS.

(The Purging Drink, No. 6, or a fourth of No. 1, should be given before the Astringent Drinks.)

1. Prepared chalk 2 oz., catechu 1 oz., ginger ½ oz., opium 1 dr., peppermint-water a pint. Dose, for lambs, a tablespoonful night and morning.—YOUATT.

2. Prepared chalk ¼ oz., ginger ½ dr., catechu ½ dr., opium 2 gr., in gruel; one or twice daily.

3. A tablespoonful of Calves' cordial, p. 133.

4. Compound powder of chalk with opium 2 dr., gentian 1 dr., essence of peppermint 3 drops; in a little thin starch, morning and night.—CLATER.

5. Ginger 2 dr., caraway 4 dr., prepared chalk 4 dr.; mix; give a teaspoonful in gruel.—WHITE.

6. In white skit; a teaspoonful of WHITE'S alkaline solution (p. 134) in a little gruel; and afterwards No. 7.

7. Epsom salts 3 dr., common salt a scruple, powdered ginger a scruple, thin gruel 4 oz. Repeat if necessary.

DRENCH FOR THE LAST EPIDEMIC. Epsom salts 1 oz., sulphur 2 dr., nitre ½ dr., ginger 15 gr., in warm water. Repeat half this, with a teaspoonful of sweet spirit of nitre, daily.—SPOONER.

DRENCH FOR COW-POX. Mix three parts of flowers of sulphur, 1 of common salt, and 1 of honey, into an electuary; give ½ of this daily, in gruel. Keep the mouth and nose clean with vinegar and water. See also Purging Drench, No. 7, above.

DRENCH FOR INFLUENZA. Epsom salts ½ oz., chamomile tea 4 oz. Afterwards give half doses of the Fever Drench, above.—DARBY.

DRENCHES FOR BLOWN OR BLAST.

1. Glauber's salts 1 oz., hot water 1 oz., peppermint-water 4 oz., tincture of ginger 1 dr., tincture of gentian 1 dr., every six hours till the bowels are opened, and half the quantity the next four mornings.—CLATER.

2. Common salt 1 oz., solution of potash (WHITE'S) 1½ dr., castor or olive oil 2 tablespoonfuls, water 3 oz. (After letting out the air by a tube or probang.)—WHITE.

DRINK TO PREVENT RESP OR MEADOW SICKNESS. Pearlash 1½ dr., hot water 8 oz. To be given from a flat bottle the second and fourth morning after putting them to keep.—HOLDITCH.

DRENCH FOR STURDY AND APOPLEXY. After bleeding, 2 oz. of Epsom or Glauber's salts, in warm water or thin gruel.

DRENCHES FOR FLUX, OR SCOURING, OR DYSENTERY.

1. Epsom salts 1 oz., hot water on thin gruel to dissolve

it; add castor oil 2 oz., laudanum 30 drops. When it has operated, give No. 2.—BLAINE.

2. Ipecacuanha 15 gr., prepared chalk 1 dr., opium 2 gr.; boiled starch or arrow-root 4 oz. Night and morning.

3. Linseed oil 2 oz., powdered opium 2 gr., linseed tea q. s. Afterwards give No. 4.—SAYER.

4. Opium 2 gr., ginger ½ dr., gentian ½ dr., linseed tea or gruel q. s.

5. Epsom salts 1½ oz., hot water 4 oz.; dissolve and add castor or olive oil 1½ oz.—WHITE. Afterwards give No. 6.

6. Catechu ½ dr., allspice ½ dr., caraway 1 dr., water or beer 4 oz.; simmer together.

DRENCHES FOR DRY BRAXY, OR INFLAMMATION OF THE BOWELS.

1. After bleeding: Epsom salts 1½ oz., warm water 1 pint.—STEVENSON.

2. After bleeding: common salt 1 oz., water ½ pint, laudanum a teaspoonful.—WHITE.

DRENCH FOR BITES OF VENOMOUS REPTILES.

Olive oil 4 oz., spirit of hartshorn ¼ oz., gruel or arrow-root ¼ pint.—WHITE.

DRENCHES TO PROMOTE PARTURITION IN THE EWE.

1. A decoction of horsemint, or any other kind of mint.

2. Bruised ergot of rye 1 dr., boiling water a pint; infuse for a ¼ of an hour, and give a third part. Repeat if necessary.

DRENCHES AND POWDERS FOR THE ROT.

1. Juniper berries 6 oz.; gentian 1 oz.; boil in 3 gallons of water for a quarter of an hour, strain, and add common salt 4 lb, powdered ginger 4 oz., tartarized iron 2 oz.; stir, and let it stand till cool. Put it into wine-bottles filled two-thirds full, and add to each 1½ oz., oil of turpentine and ½ oz. sweet spirit of nitre. Give a table-spoonful night and morning, shaking the bottle before pouring it out.

2. Common salt 8 oz., gentian powder 8 oz., ginger 1 oz., tincture of calumba 4 oz., water to make up a quart.—CLATER. See the next.

3. To a quart of No. 2, add spirit of turpentine 3 oz.; shake well together, and give 2 tablespoonfuls at night, before the night's food is given, and a tablespoonful of No. 2 every morning.

Powders for the same.—A French recipe. 1. Dry bran 10 lb, salt ½ lb, aromatic herbs (as thyme, sage, juniper, rosemary, &c.) cut small, 6 oz., green anise and coriander, of each 5 oz.; mix and give morning and night every third day. The above quantity is for thirty sheep.

2. Juniper-berries 4 oz., bay-berries 1 oz., grains of paradise ¼ oz., bay salt 1½ lb, loaf sugar ½ lb; powder all together, and keep the powder in a bottle for use. Give the sheep dry and sweet hay, sprinkled with the powder.—LAWRENCE.

DRENCHES FOR INFLAMMATION OF THE LUNGS, CATARRH, HOOSE, AND COUGH. After bleeding from the neck, give Epsom salts 2 oz., gruel or linseed tea q. s.

DRENCH FOR INFLUENZA. Epsom salts ½ oz., chamomile tea 4 oz. Afterwards small doses of digitalis, opium, tartarized antimony, and vegetable tonics.—DARBY.

External Applications for Sheep.

EYE-WATERS.

1. *Strong.* For cloudiness of the eye; corrosive sublimate 4 gr., spirit of wine ½ oz.; dissolve, and add water 1 pint.—CLATER.

2. Tincture or wine of opium 1 teaspoonful, water ½ pint.

WASHES FOR THE SCAB, LICE, AND TICKS. (The scab ointments will also destroy them, and are less hazardous, and less injurious to the wool.)

1. *Arsenical Wash.* White arsenic ½ lb, salt of tartar ½ lb, water 12 gallons; boil for half an hour.—YOUATT.

2. Arsenic 2 lb, soft soap 4 lb, water 30 gallons; dissolve. The sheep to be immersed in this liquid (the head only being kept out), and while in it the fleece to be well rubbed. When taken out the fluid should be well pressed out of the fleece,

and the sheep kept from cold and wet for a few days.—CLATER. Mr. SPOONER says 2 lb of arsenic should make 48 gallons of the liquid.

3. Arsenic 1 lb, yellow soap 6 lb, pearlash 12 oz., water 30 gallons.—MATTHEWS.

4. *Mercurial.* Corrosive sublimate 1 oz., spirit of wine 2 oz.; rub together till dissolved, then add cream of tartar 1 oz., bay salt 4 oz., dissolve the whole in 2 quarts of water, and apply it with a sponge wherever lice appear. —CLATER.

5. Tobacco 4 oz., water 1 gallon; boil, and add soft soap 1 lb, sulphur vivum 1 lb; when cold add a pint of oil of turpentine.

6. Equal parts of decoction of tobacco and lime-water.—YOUATT.

WASH TO KILL MAGGOTS. Shake up in a bottle together 1 quart of water, spirits of turpentine 1 oz., and corrosive sublimate 10 grains. Stop with a cork in which a quill is inserted. When the maggots are observed, a small quantity of the mixture is to be shaken on the spot through the quill, and they will shortly creep out and die.

SMEARING MIXTURE.

1. One gallon of common tar, and 12 lb of any sweet grease, melted together.

2. Oil of tar is used as a preventive of the *fly;* but fish oil is equally so, according to Mr. HOGG; and is less injurious to the wool. Oil of tar has sometimes destroyed sheep.

FLY POWDER FOR SHEEP.

1. White lead 2 lb, red lead ½ lb, sulphur 1½ lb, oil of wormwood, animal oil (empyreumatic), or creasote ¼ oz.; mix.

2. White lead 2 lb, red lead 1 lb; mix, and apply by sprinkling from a dredger, following a stick drawn through the wool.—CLATER.

3. Powdered colocynth 3 dr., black brimstone 1 lb, tincture of assafœtida ½ oz.; mix.

4. White lead 4 parts, arsenic 1 part, sulphur 6, vermilion 2.—SPOONER.

POWDER FOR THE EYES. Equal parts of sal ammo-

niac, white sugar, and oxide of zinc, triturated together. It may be mixed either with rose-water or honey.—SPOONER.

ASTRINGENT POWDER FOR THE FEET, in the epidemic affecting them. The same as for cattle. See above, page 139.

OINTMENT FOR THE SCAB OR SHAB.

1. Quicksilver 1 lb, Venice turpentine ½ lb; rub them together until the globules are no longer visible; then add ½ pint of oil of turpentine, and 4 lb of lard. The mode of applying this ointment is as follows: Begin at the head of the sheep, and proceeding from between the ears along the back to the end of the tail, divide the wool in a furrow till the skin can be touched; and let a finger slightly dipped in the ointment be drawn along the bottom of this furrow. From this furrow similar ones must be drawn along the shoulders and thighs to the legs, as far as the wool extends. And if much infected, 2 or more should also be drawn along each side, parallel with that on the back; and one down each side before the hind and fore legs. It kills the sheep-fag, and probably the tick and other vermin. It should not be used in very cold or wet weather.—Sir JOSEPH BANKS.

2. Strong mercurial ointment 1 part, lard 5 parts; mix.—YOUATT.

3. Quicksilver 1 lb, Venice turpentine ½ lb, spirit of turpentine 2 oz., lard 4½ lb; to be made and used as No. 1. In summer 1 lb of resin may be substituted for a like quantity of lard.—CLATER.

4. Strong mercurial ointment 1 lb, lard 4 lb, oil of turpentine 8 oz., sulphur 12 oz.—WHITE.

5. *Mild.* Flowers of sulphur 1 lb, Venice turpentine 4 oz., rancid lard 2 lb, strong mercurial ointment 4 oz.; mix well.—CLATER.

6. Lard or other fat, with an equal quantity of oil of turpentine.—DAUBENTON.

7. *Without Mercury.* Lard 1 lb, oil of turpentine 4 oz., flowers of sulphur 6 oz.—WHITE.

8. Strong mercurial ointment 1 lb, lard 4 lb, Venice turpentine 8 oz., oil of turpentine 2 oz. If mixed by heat, care must be taken to use no more heat than is necessary, and add the oil of turpentine when the other ingredients begin to cool, and to stir till cold.—MCEWEN.

9. Corrosive sublimate 2 oz., white hellebore 3 oz., fish oil 6 quarts, resin ½ lb, tallow ½ lb. The sublimate and then the hellebore to be rubbed with a portion of the oil till perfectly smooth, and then mix with the other ingredients melted together.—STEVENSON.

10. The following once had considerable local celebrity, but it obviously requires to be used with caution. Dissolve 2¼ oz. of corrosive sublimate in the same quantity of muriatic acid, and beat up the solution with 6 lb of strong mercurial ointment; put it in a large pan, and pour on it 19½ lb of lard, and 1½ lb of common turpentine, melted together and still hot, and stir the whole continually until it becomes solid.

OINTMENT FOR DEEP WOUNDS OR ULCERS FROM FLIES. The fly powder No. 2, mixed with tar.—CLATER.

OINTMENT FOR SORE HEADS. Black pitch 2 lb, tar 1 lb, flowers of sulphur 1 lb; melt together, taking care that it does not boil. To be spread thickly on leather while warm, and fitted to the head.

CAUSTIC ASTRINGENTS FOR FOOT ROT.

1. Blue vitriol 1 oz., white vitriol 1 oz., burnt alum 2 oz., bole ½ oz., honey to form a stiff paste.—MCEWEN.

2. Sulphate of copper 2 oz., water 12 oz., dilute sulphuric acid 2 dr.—WHITE.

3. Butter of antimony, alone, or mixed with tincture of myrrh.

4. Verdigris, bole, and sugar of lead, in equal parts, rubbed together into a fine powder. Sprinkle on the sore, cover with tow, and bind down with tape for 24 hours, using afterwards No 3, or No. 2.—CLATER.

5. *Strong.* Verdigris 1 oz., nitrous acid 2 oz., water 4 oz.—WHITE.

6. *Strongest.* Red precipitate 1 oz., nitrous acid 2 oz.; dissolve, and add water 2 oz., spirits of wine 1 oz.—WHITE.

7. Aloes 16 oz., weak spirit 32 oz., sulphuric acid 17 oz.; mix.—DUVILLE.

8. Dissolve sulphate of copper 2½ oz. in 1½ pint of water, and add a solution of 3½ dr. of sulphate of iron previously calcined. Diffuse ¾ oz. slaked lime in water, and add the mixed solution; then add 7 oz. common salt, 1 oz. wood

vinegar, and water to make up a quart. (Nearly the composition of a celebrated French nostrum.)

9. LELOUP'S Terebinthinated Oxymel of Copper. Honey 14 oz., pyroligneous acid 7 oz., powdered verdigris 5 oz.; boil it in a large copper pan until it assumes a reddish-purple color; then add, keeping the mixture on a slow fire, 14 oz. Venice turpentine; stir with a wooden spatula for ¼ of an hour, and pour it into jars. To be applied twice at 12 hours' interval, by means of a small piece of wood, after cleaning the part with an iron blade.

10. White vinegar 78 parts, powdered sulphate of copper 10 parts; dissolve, and add 12 parts of sulphuric acid. Apply it with a feather. (A French remedy.)

11. *Detersive Ointment.* Burnt alum 4 parts, verdigris 1 part, camphor 1 part, green ointment of elder or poplar 16 parts.—LEBAS.

12. Honey 4 oz., burnt alum 2 oz., Armenian bole ½ lb; mix with as much train oil as will convert these ingredients into a salve. The honey must first be completely dissolved in the oil made hot, then the bole stirred in, and lastly the alum.

BLACKLOCK condemns all caustic applications, using only mild poultices and emollient ointments.

MEDICINES FOR SWINE.

ALTERATIVE MEDICINES, in mange and other skin diseases, and in obstinate costiveness.

1. Sulphur ¼ oz., Æthiops mineral 3 gr., nitre ½ dr., cream of tartar ½ dr.; daily, in thick gruel or wash.—CLATER.

2. Black antimony ½ oz., sulphur 2 oz., nitre ½ oz.; mix; for 8 doses.

FEVER MEDICINE. Digitalis 3 gr., antimonial powder 6 gr., nitre ½ dr.; after bleeding, in a little warm swill, milk, or mash, morning, noon, and night.—CLATER.

PURGING MEDICINES.

1. Epsom salts 1, 2, or 3 oz., in broth or swill.

2. Sulphur 2 dr., daily; full dose ½ oz., with milk or other food. This may be repeated for 2 or 3 days, in surfeit from overfeeding.

3. Jalap 1 dr.; if insufficient, add 10 or 12 gr. of scammony, or 10 gr. of calomel.—WHITE.

4. Jalap ½ dr., sulphur 2 dr., antimony ½ dr.

5. Jalap ½ dr., Epsom salts 1 oz.

6. Castor oil 1 oz. to 2 oz., with gruel.

7. Castor oil 1 oz., gruel q. s., Epsom salts 2 oz., salt ¼ oz.; mix.

8. Calomel 5 gr.; but this must not be repeated more than twice.

CARMINATIVE DRENCH, for flatulent distension, from sour whey, &c. After using the probang, or where it cannot be had, give—

1. WHITE'S solution of potash (see page 134) 2 oz., anodyne carminative tincture 1 tablespoonful, water 8 oz.

2. A tablespoonful of common salt in warm water, a teaspoonful of mustard or powdered ginger, and a glass of gin.

THRIVING POWDER, to promote fattening. Powdered fenugreek, alone, or mixed with a fourth of liquorice powder; an ounce daily with the food. Cleanliness greatly conduces to the same end.

REMEDIES FOR MEASLES. After bleeding by tail, ear, palate, or vein inside the forearm, an inch above the knee, give one of the purging drinks, and turn it into the open air.

REMEDIES FOR THE DISTEMPER, affecting the Mouth and Feet. The same drink, and astringent powder as for SHEEP.

DRENCH FOR INFLAMMATION OF THE BRAIN. Castor oil 2 oz., with gruel; afterwards 2 gr. white hellebore powder twice or thrice a day.—CUPISS.

HEALING OINTMENT FOR SORE EARS.

1. Lard 1 ℔, resin 4 ℔; melt together, and stir in ½ ℔ lapis calaminaris.—CLATER.
2. Zinc ointment 1 oz., yellow basilicon 3 oz.
3. Tar ointment mixed with a little soap.

MANGE OINTMENT.

Sulphur 4 oz., Venice turpentine 1 oz., old lard 8 oz., mercurial ointment 1 oz.; the animal to be previously scrubbed all over with a good soap lather. (The above alterative powders should be given at the same time.)

OINTMENT FOR SORE TEATS. Soft soap 4 oz., camphor (powdered with spirit) ¼ oz., mercurial ointment ½ oz. It must be carefully washed off.

MEDICINES FOR DOGS.

N. B.—The doses required vary considerably, according to the strength and size of the dog, which should always be duly considered.

PHYSIC BALLS AND OTHER PURGATIVE MEDICINES.

1. Barbadoes aloes 8 oz., antimonial powder 1 oz., ginger 1 oz., palm oil 5 oz.; beat together into a mass. Dose from ½ dr. to 2 dr., every 4 or 6 hours, till the bowels are relieved. —YOUATT.

2. The same, with the addition of 1 oz. of colomel. He directs from 45 grains to 2 dr. for a dose.—CLATER.

3. Aloes ½ dr. to 2 dr. made into a ball with syrup of ginger.

4. Aloes ½ dr., to 1½ dr., calomel 2 to 5 gr., syrup to form a ball; in inflammation of the bowels, and in worms. —BLAINE.

5. Cape aloes ½ dr. to 1 dr., calomel 2 to 3 gr., oil of caraway 6 drops, syrup to form a ball.—McEWEN.

6. Calomel 12 gr., aloes 3 dr., opium 1 gr., syrup q. s. to form a mass for 4, 6, or 8 balls; one every 4 or 5 hours till the bowels are relieved.—BLAINE.

7. Croton oil 1 drop, Castile soap 20 gr., conserve to form a ball.

8. Castor oil 3 parts, syrup of buckthorn 2 parts, syrup of poppies 1 part; dose from 1 to 2 tablespoonfuls.—Mr. YOUATT'S purge. [Mr. CLARKE says syrup of buckthorn for dogs should be made with treacle, and the spices omitted.]

9. Epsom salts, from 1 to 4 dr,, wrapped in tissue-paper, dividing the doses into convenient sized packets.

10. In costiveness with inflammation: ½ oz. to 2 oz. castor oil.—Mr. SPOONER.

11. Jalap, powdered, 30 gr., calomel 8 gr.; make into a pill with gum-water, and administer every morning. In distemper.

ALTERATIVE BALLS AND POWDERS.

1. Sulphur 2½ ℔, nitre ½ ℔, Æthiops mineral 4 oz., linseed meal ½ ℔, palm oil 1 ℔, or as much as may be required;

beat together, and keep in a jar for use: dose, from 2 scruples to 1½ or 2 dr.—CLATER.

2. Æthiops mineral 20 to 40 gr., cream of tartar 20 to 40 gr., nitre 5 to 10 gr.; night and morning, made into a ball with butter.—SPOONER.

3. *Tonic Alterative.* Mercurial pill 1 dr., aloes 2 dr., myrrh, benzoin, balsam of Peru, of each 1½ dr.; to be divided into 10, 15, or 20 pills; one every evening for the yellows, after aloes and calomel.—BLAINE.

4. *Alterative Powder.* Æthiops mineral 2 to 5 gr., cream of tartar 4 to 10 gr., tartarized iron 1 to 3 gr.; once a day. —CLATER.

5. *To give a fine skin.* Give a tablespoonful of tar, made up with oatmeal.—MAYER.

ASTRINGENT BALLS, &c.

1. Catechu 1½ dr., sulphate of quinine 20 gr., opium 5 gr., ginger 1 dr., conserve of roses q. s. to form a mass, to be divided into 8, 6, or 4 balls.—BLAINE.

2. Prepared chalk 2 oz., powdered gum arabic ½ oz., powdered catechu ½ oz., powdered oak bark ½ oz., powdered ginger ¼ oz., opium 15 gr., palm oil 1 oz.; beat well together: dose, ½ dr. to 2 dr., morning, noon, and night, in the advanced stage of distemper.—CLATER.

3. Opium 5 gr., catechu 2 dr., gum arabic 2 dr., ginger ½ dr., syrup of poppies q. s.; divide into 12, 9, or 6 balls. In diarrhœa.—BLAINE.

4. Myrrh 1 dr., ipecacuanha 1 scruple, opium 3 gr., chalk 2 dr., carbonate of iron 1 dr.: as No. 3.—BLAINE.

5. In obstinate cases: Alum 1 dr., chalk 2 dr., opium 6 gr., resin 3 dr.: into 4, 6, or 8 balls.

6. In diarrhœa, after 1 to 4 dr. of Epsom salts; Prepared chalk 1 to 3 scruples, catechu 5 to 10 gr., opium ¼ to 2 gr.; twice a day.—SPOONER.

7. *Astringent Drink.* Boil 1 oz. of logwood in a quart of milk to ½ a pint. A teaspoonful every morning, *in prolapsus.*

COUGH BALLS IN ASTHMA, &c.

1. *After a few emetics.* Calomel 3 gr., foxglove 3 gr., cream of tartar 1 dr., antimonial powder 12 gr., honey to form 6 boluses. One twice a day.—BLAINE.

2. Digitalis 20 gr., antimonial powder 40 gr., nitre 2 dr., sulphur 3 dr., palm oil 3 dr., or q. s. Divide into 10, 15,

or 20 balls, according to the size of the dog, morning and night, interposing an emetic every third or fourth day.—CLATER.

3. *In old cases.* P. squill ½ gr. to 1 gr., gum ammoniac 5 gr., balsam of Peru 8 gr., benzoic acid 1 gr., balsam of sulphur to form a ball.

4. Extract of hemlock ½ dr., extract of henbane 10 gr., p. digitalis 20 gr., conserve of roses to form a mass. Divide into 10, 8, or 6 balls. One night and morning.—BLAINE.

DISTEMPER MEDICINES.

1. Turpeth mineral 1 to 3 gr., assafœtida ½ dr., aloes 20 gr., soap 10 gr., syrup of poppies to form a ball. To be preceded by an emetic, and given every third day.

2. After bleeding (if required) and an emetic, give a physic ball; and afterwards the following, 2 or 3 times a day; Antimonial powder, 2, 3, or 4 gr., nitre 5, 10, or 15 gr., ipecacuanha 2, 3, or 4 gr.; form a ball. If the disease proceed to the debilitating stage, give the *Tonic Ball*, No. 2; in the putrid or malignant stage, give the *Astringent Ball*, No. 1.—BLAINE.

3. After the emetic powder No. 1 (which should be repeated every third or fourth day) give the *Cough Ball*, No. 2, from ½ dr. to 2 dr. in weight. And if the dog lose flesh, give equal parts of the Cough Ball and the Tonic Ball (No. 1). In the more advanced stages give the tonic alone; or, the *Astringent Ball*, if diarrhœa comes on.—CLATER.

4. Give a third of a paper of James's powder, mixed with butter, and afterwards warm broth or milk. In 2 hours another third; and if this neither vomit nor purge, give the other third at the end of 4 hours.—DANIEL.

5. BLAINE'S *Distemper Powders*, which are sold in packets, with directions for use.

6. Camphor 3 to 5 gr., charcoal 10 gr., opium 1 gr., aromatic confection q. s. to form a ball.—In the malignant stage, with diarrhœa.

7. Antimonial powder 2 to 4 gr., nitre 5 to 10 gr., digitalis ¼ to 2 gr. Afterwards the tonic pills No. 4.—SPOONER.

Poudre Kusique: a French nostrum. Mix 45 gr. of nitre, 45 of sulphur, and 1 of charcoal. Divide into 3 doses. Give 1 for two successive mornings, and the third

on the fourth morning, mixed with lard or butter, or in milk. For a large dog a second packet (of 3 powders) may be required.—HABERT.

Another French nostrum, Hemel's powder, is of a similar kind.

8. A strong solution of salt, to the amount of ½ pint daily.

9. Powdered tin, sulphur, gunpowder, of each 1 oz.; lard sufficient to form a mass. The size of a nutmeg to be given twice or thrice a week.

10. Physic ball No. 11.

REMEDIES FOR SPASMODIC COLIC.

1. Castor oil ½ oz., oil of peppermint 1 drop, laudanum 20 drops. If it does not open the bowels, give ½ dr. to 1½ dr. of aloes.—BLAINE.

2. Castor oil 3 oz., syrup of buckthorn 2 oz., syrup of poppies 1 oz. Give from a teaspoonful to a tablespoonful. —YOUATT.

3. Ether ½ dr., laudanum ½ dr., camphor 3 to 6 gr., castor oil (unless he is purged) 3 to 5 dr.—BLAINE.

CONVULSIONS.

Give Colic mixture No. 3, and apply warm bath and flannel.

FOR FITS OR EPILEPSY.

1. Calomel 8 gr., carbonate of iron ½ dr., extract of hemlock 20 gr., conserve of roses, or plain oil, to form a mass for 12, 9, or 6 balls.—BLAINE.

2. Give the Alterative balls No. 1, or the pills of nitrate of silver, as for St. Vitus's dance.—CLATER.

3. For epilepsy of suckling bitches: Ether 1 dr., laudanum ½ dr., strong ale 2 oz.; give from a dessert-spoonful to 1 or 2 tablespoonfuls every 2 or 3 hours.—BLAINE.

4. For epilepsy attending distemper: The Tonic balls, or the pills for chorea.—CLATER.

5. After an emetic: Gentian 10 to 20 gr., ginger 3 to 6 gr., carbonate of iron 2 to 4 gr., or from an eighth to a fourth of a gr. of nitrate of silver, and ½ gr. of spiders' web once a day.—SPOONER.

6. Ether 1 dr., laudanum ½ dr., camphor 6 gr., spirit of hartshorn 1 dr.; in a spoonful of ale: for small dogs give half the quantity.—BLAINE.

EMETIC POWDERS.

1. Calomel, emetic tartar, of each 1 oz.; vermilion 10 gr.; rub together: dose, from 1 to 3 gr., dropped on the tongue, or mixed with a teaspoonful of milk.—CLATER.
2. Emetic tartar, from 1 to 3 gr.
3. Turpeth mineral, from 1 to 3 gr.
4. A teaspoonful of common salt.

MEDICINES FOR INFLAMMATORY DISORDERS.

1. In inflammation of the lungs: After bleeding and purging, digitalis 12 gr., emetic tartar 3 gr., nitre 1 dr.; mix, and divide into 6, 9, or 12 powders.—BLAINE.
2. Ditto, with much cough: Tincture of digitalis 1 dr., emetic tartar 3 gr., nitre 1 dr., simple oxymel 2 oz.: dose, 1 or 2 dr. every 3 hours.—BLAINE.
3. In pleurisy, with incipient water in the chest: Digitalis 6 gr., calomel 6 gr., tartarized iron 18 gr.; into 6, 9, or 12 doses.—YOUATT.
4. In inflammation of the liver: Digitalis 8 gr., antimonial powder 16 gr., nitre 1 dr.: divide into 7, 9, or 12 powders, or boluses.—BLAINE.
5. In chronic inflammation of the liver: Calomel 20 gr., antimonial powder ½ dr., myrrh, gentian, aloes, of each 2 dr.; mix, and divide into 15, 20, or 25 balls.
6. In inflammation of the bowels: After bleeding and a warm bath, give the castor oil mixture (Purgatives No. 8).—CLATER.
7. Bilious inflammation (with offensive, often black, vomiting and purging): Calomel 10 gr., opium 4 gr.: in 4 or 8 pills—one 3 times a day; afterwards the astringent remedies for diarrhœa.—SPOONER.

MEDICINES FOR RHEUMATISM.

1. After warm bath, and friction, give tincture of opium 20 drops, ether 20 drops, castor oil ½ oz. to 1 oz.—BLAINE.
2. Calomel 2 to 4 gr., opium ¼ gr., oil of peppermint 1 drop, aloes 1 dr.; form a ball with butter or lard: repeat it every 4 hours till the bowels are well opened; and use the Embrocation No. 3.—CLATER.
3. After warm bath, &c., give 40 drops of laudanum, and a teaspoonful of hartshorn, in warm beer; and rub with the Embrocation No. 1.—MAYER.

TONIC MEDICINES.

1. Gentian 1 oz., chamomile ½ oz., oak bark ½ oz., ginger ¼ oz., carbonate of iron ¼ oz., palm oil 1 oz.: beat them together to form a mass: dose 2 to 6 scruples.—CLATER.

2. Sulphate of quinine ½ dr., powdered chamomile 3 dr., balsam of Peru 1½ dr., camphor 1 scruple; form a mass with conserve of roses, and divide into 12, 9, or 6 balls; one every 6 hours, in the debilitating stage of distemper.—BLAINE.

3. Chamomile 1 oz., rue ½ oz., ginger ¼ oz. (all in powder), beat them into a mass with 7 dr. of palm oil, and divide into 12, 16 or 20 balls; one night and morning in gutta serena.—CLATER.

4. Gentian powder 10 to 20 gr., ginger 5 gr., cascarilla 10 to 20 gr.; conserve of roses, or syrup, to form a ball. One twice a day.

WORM MEDICINE.

1. Carbonate of iron ½ oz., Æthiops mineral 1 dr., gentian 1 oz., ginger ½ oz., levigated glass 1 oz., palm oil 9 dr.; beat well together: dose from ¾ to 2 dr.—CLATER.

2. As much finely-powdered glass as will lie on a sixpence, mixed with butter.—BLAINE. Mr. YOUATT says from ½ dr. to 1 dr. powdered glass, with a little ginger, made into a ball with lard.

3. Aloes, sulphur, prepared hartshorn, and juice of wormwood, made into a mass; the size of a hazel-nut to be given three times a week, fasting, wrapped in butter.—DANIEL.

4. Tin filings, or pewter filings, ½ dr. to 1 dr., with butter or lard.

5. Jalap 10 to 15 grs., calomel 2 to 3 gr., mixed with butter; no cold liquid should be allowed.—WHITE.

6. Cowhage ½ dr., iron filings 4 dr., conserve q. s. to form a mass, to be divided into 4, 6, or 8 balls; one every night and morning; and afterwards the purgative No. 4.—BLAINE.

7. Epsom salts 1 oz., common salt 1 dr.; give a small or large teaspoonful daily.

8. Give green walnut leaves boiled in milk.—MAYER.

9. *For Tape Worm.* Oil of turpentine ½ dr., mixed with yolk of egg; for very large dogs 2 scruples. Some writers prescribe larger doses (1 or 2 dr.), but these sometimes prove fatal.—BLAINE.

10. *For Tape Worm.* Oil of turpentine and olive oil, of each ½ oz., mix, and give carefully; three or four hours after give 1 oz. castor oil. But see No. 9.—WHITE.

11. *For Stomach Worms.* Give the emetic powder (above) and afterwards a physic ball.

12. *Thread Worms.* These are destroyed by an aloetic clyster.

MEDICINES FOR THE YELLOWS.

1. After bleeding: Calomel 2 to 3 gr., jalap 10 gr., scammony 4 gr.—WHITE.

2. Aloes 20 to 40 gr., calome l2 to 4 gr.; afterwards the tonic alterative balls.—BLAINE. See ALTERATIVES.

MEDICINES FOR ST. VITUS'S DANCE, OR CHOREA.

1. Nitrate of silver 8 gr., ginger 10 gr., syrup to form a mass; divide into 64 pills, and give one or two morning and night.—CLATER.

2. Strychnia 1 gr., oxide of zinc 24 gr., assafœtida 24 gr., conserve of roses q. s.; mix very accurately, and divide into 12, 9, or 6 balls.

3. Nitrate of silver 3 gr., carbonate of iron 2 dr., gentian 3 dr., conserve of roses to form a mass, for 12, 9, or 6 balls.—BLAINE.

MEDICINES FOR DROPSICAL COMPLAINTS.

1. Digitalis 9 gr., squill 12 gr., cream of tartar 2 dr., mix, and divide into 9, 12, or 15 powders; one night and morning.

2. Foxglove 12 gr., antimonial powder 15 gr., nitre 1 dr.; as the last.—BLAINE.

3. Foxglove 1 gr., nitre 10 gr., ginger 8 gr.; night and morning; then iodide of potassium ½ gr. to 1 gr.—YOUATT.

BALLS FOR ENLARGED GLANDS AND CANCEROUS DISEASES.

1. Extract of hemlock 1 to 3 gr., burnt sponge 10 to 20 gr.; make a ball, to be given once or twice a day.

2. Iodine 12 gr., powdered gum 40 gr., syrup to form a stiff mass, divide into 48 pills, and give one or two night and morning.—CLATER.

BALLS TO PROMOTE PARTURITION. Ergot of rye 20 gr.: pulverize, and add ginger 16 gr., syrup q. s.: beat into

a mass, and divide into 5 pills; give one every hour, or to a small bitch, half of one.—CLATER.

TO PREVENT RABIES, OR CANINE MADNESS.

1. Powdered leaves of the scutellaria lateriflora 40 gr., powdered belladonna 2½ gr.; to be given night and morning for 6 weeks, gradually increasing the dose.—YOUATT.

2. Infuse a teaspoonful and a half of powdered scutellaria in a quart of hot water; give half a pint morning and night, omitting the dose every third day, when a mild dose of sulphur must be given.—Dr. SPALDING.

3. Fresh leaves of the tree box 2 oz., rue 2 oz., sage ½ oz., chop them fine, and boil them in a pint of water till reduced to half a pint; strain and press out the liquid; beat the herbs, and boil them in a pint of new milk to half; strain, press the herbs, and mix the liquids. For a man, give a third of this quantity every other morning fasting; double the above quantity makes 3 doses for a horse or cow; two-thirds will suffice for a middle-sized dog, and a third for smaller dogs. It produces extreme nausea and distress, and has occasionally proved fatal to dogs.—BLAINE.

External Applications.

ASTRINGENT AND DETERGENT LOTINS, for Wounds, &c.

1. Bruised oak-bark 2 oz., catechu 1 oz., water 3 pints; boil to a pint, and strain.—CLATER.

2. Tincture of myrrh and aloes 1 oz., alum ½ oz., water 1 pint.

3. For sore feet: the Lotion No. 1, 4 oz., tincture of aloes ½ oz., water 1 pint.

4. Nitrate of silver 10 gr., water 1 oz.: to excite sluggish wounds.

LOTION FOR CANKER IN THE EAR. See OINTMENTS.

1. Sulphate of zinc 20 gr., sugar of lead ½ dr., water 4 oz.—WHITE.

2. Sulphate of zinc 20 gr., decoction of oak-bark 4 oz.—BLAINE.

3. Nitrate of silver 1 gr., rain-water 2 oz.

4. Sugar of lead ½ dr., rose-water 4 oz. A teaspoonful to be introduced blood-warm into the ear.

5. *Mild Canker Lotion.* Infusion of foxglove leaves ½ pint, Goulard's extract of lead ½ oz.; mix. To be used as the last.—CLATER.

6. *Strong Canker Lotion.* Goulard's extract 2 dr., white vitriol 1 dr., alum 2 dr., water ½ pint.

7. Chloride of lime 20 to 30 gr., water ½ pint.

LOTION TO ALLAY ITCHING. Dilute hydrocyanic acid 1 dr., water 1 pint.—YOUATT.

OINTMENTS FOR CANKER OF THE EAR.

1. Equal parts of zinc ointment, and ointment of nitrate of quicksilver.—BLAINE.

2. Sublimate 3 gr., Turner's cerate 1 dr., sulphur 1 scruple.

3. White vitriol, alum, each in fine powder 1 dr., lard 4 oz. To be rubbed gently into the crack.—CLATER.

4. *Stronger.* Nitrate of silver 20 gr., lard 1 oz. Rub them well together.—CLATER.

5. Levigated red precipitate ½ oz., lard 2 oz.—MAYER.

EYE-WATERS.

1. *Astringent Wash for Weak Eyes.*—White vitriol 4 gr., spirits of wine ½ dr., water 4 oz.—CLATER.

2. Sugar of lead 30 gr., rose-water 6 oz.—BLAINE.

3. Laudanum ½ dr., infusion of green tea 4 oz.—McEWEN.

4. For naturally weak eyes: Laudanum 2 dr., water 8 oz. To be used every morning.—CLATER.

5. Sugar of lead ½ dr., distilled water 6 oz., tincture of opium ½ dr. In inflammation after bleeding, physic, and warm fomentations.—SPOONER.

OINTMENTS FOR ULCERATED EYELIDS.

1. Red precipitate, levigated, 10 grains, zinc ointment ½ oz.

2. Ointment of nitrate of quicksilver 1 dr., sugar of lead, 20 gr., spermaceti ointment 3 dr.—BLAINE.

3. Dissolve a drachm of quicksilver in a drachm and a half of strong nitric acid, and mix the warm solution well with 6 oz. of melted lard.—CLATER.

OINTMENTS AND LOTIONS FOR THE MANGE.

N.B.—An alterative ball should be given daily and a physic ball occasionally. Bleeding is sometimes prescribed.

1. For Scabby Mange: Sulphur 4 oz., sal ammoniac ½ oz., aloes 1 dr., Venice turpentine ½ oz., lard 6 oz.: mix. After four applications wash well with soap and water.—BLAINE.

2. Horse turpentine and palm oil, each ½ lb, train oil ½ pint. Melt together, and while cooling stir in 3 lb of flowers of sulphur.—CLATER.

3. Aloes 2 dr., hellebore ½ oz., sulphur 4 oz., lard or train oil 6 oz.—MCEWEN.

4. Sulphate of zinc 1 dr., snuff ½ oz., white hellebore ½ oz., sulphur 4 oz., aloes ¼ oz., soft soap 6 oz.—BLAINE.

5. Charcoal powder 2 oz., sulphur 4 oz., salt of tartar 1 dr., Venice turpentine ½ oz., lard 6 oz.

6. For Red Mange: Add 1 oz. of strong mercurial ointment to 6 oz. of either of the above.

7. Charcoal 1 oz., chalk 1 oz., sugar of lead 1 dr., white precipitate 2 dr., sulphur 2 oz., lard 5 oz.—BLAINE.

8. *Wash for Red Mange.*—Sublimate 20 gr., spirit of wine 2 dr.; dissolve and add milk of sulphur ½ oz., lime-water ½ pint. Apply by means of a sponge.—CLATER.

9. For Ulcerated Mange: Ointment of nitrated quicksilver 2 dr., sugar of lead 20 gr., flowers of sulphur ½ oz., lard 1 oz.; mix.—BLAINE.

ARSENICAL OINTMENT. Yellow sulphuret of arsenic 1½ gr., cerate or lard ½ oz.: in mange and other skin diseases.—DELAFOND.

SURFEIT OINTMENT. After bleeding and purging, apply sugar of lead 1 dr., spermaceti ointment 2 oz.

OINTMENT AND POWDERS FOR PILES.

Ointment. Sugar of lead 6 gr., tar ½ dr., elder ointment 3 dr.—BLAINE.

Powders. Nitre ½ dr., milk of sulphur 3 dr.; mix, and divide into 9, 12, or 15 doses.—BLAINE.

HEALING OINTMENTS.

1. Palm oil 3 lb, resin 1 lb; melt together, and when they begin to cool, add 1 lb of powdered calamine.—CLATER.

2. Oxide of zinc ¼ oz., lard 1½ oz., balsam of Peru 1 dr.

OINTMENT FOR SCIRRHOUS TUMORS. Hydriodate of potash 1 dr., lard 7 dr.; rub together till perfectly smooth.

STIMULATING MERCURIAL OINTMENT. Mercurial ointment 1 oz., simple cerate 1 oz. A small quantity to be rubbed over the region of the liver once a day till the mouth is sore.—BLAINE.

EMBROCATIONS FOR PALSY, RHEUMATISM, &c.

1. Oil of turpentine 2 oz., spirit of hartshorn 2 oz., tincture of opium ¼ oz., olive oil 2 oz.—BLAINE.
2. Cajeput oil 1 oz., soap liniment 2 oz.
3. Spirit of turpentine, spirit of hartshorn, camphorated spirit, of each 1 oz., laudanum ½ oz.—CLATER.

LOTIONS FOR STRAINS AND BRUISES.

1. Common salt and cold vinegar.
2. Sal ammoniac ½ oz., vinegar a pint.
3. Oil of turpentine 1 oz., old beer ½ pint, brine ½ pint. For strains.—MAYER.
4. Spirit of Mindererus 4 oz., Goulard water 8 oz.

ASTRINGENT LOTION FOR WOUNDS, SORE FEET, &c. Bruised oak-bark 2 oz., catechu 1 oz., water 3 pints; boil to 1 pint, and strain.—CLATER.

POWDER AND LIQUID CAUSTIC FOR WARTS.

1. Equal parts of sal ammoniac and savine, powdered together.
2. Sublimate 1 dr., muriatic acid 1 dr., spirit of wine 3 dr., water 2 dr. The warts to be touched with the liquid twice a day.—YOUATT.

FLEAS.

1. Rub the skin with powdered resin and bran.
2. Let the dog sleep on deal shavings.
3. Scotch snuff steeped in gin.—MAYER. (This requires caution.)

CLYSTERS.

Astringent. Alum whey.
Purgative. The purgative medicine No. 8; with gruel.
For Worms. Solution of aloes 2 oz., linseed oil 1 oz. Mix.
Anodyne. Boiled starch ¼ pint, laudanum 5 to 10 drops.—CLATER.

MEDICINES

FOR

POULTRY, RABBITS, &c.

FOR ROUP, POULTRY GLANDERS, AND GARGLE IN GEESE.

1. A saturated solution of common salt. Medium dose, half a teaspoonful.

2. Antimonial powder 1 gr., with sopped bread, twice a day.—CLATER.

3. Garlic, rue, brickdust, and butter, beaten together, and a little crammed down the throat.

4. For wet roup in pigeons: Give 3 or 4 pepper-corns in 3 or 4 days.—MOORE.

5. For dry roup: Give 2 or 3 pills of garlic every day. [Some recommend assafœtida to be mixed with the food of Poultry, whenever they manifest disease by drooping their wings.]

FOR RUMP ROUP, OR INFLAMMATION OF THE OIL-VESSEL.—Open the tumor, and squeeze out the collected oil.

GAPES (OR PIP), FROM WORMS IN THE AIR-PASSAGES.

1. Pills of sulphur, turpentine, and wheat flour.—("Veterinarian," Oct., 1840.)

2. Oil of turpentine 2 dr., linseed oil 1 oz.; or oil of turpentine 2 dr., flour enough to make it into 20 pills. For 20 doses, one every other day for 3 or 4 times.

3. Tobacco smoke.

INFLAMMATION OF THE LUNGS AND ASTHMA.

Give a grain each of calomel and antimonial powder, daily.

PURGING FLUX, OR DIARRHŒA.

1. Change the diet and give whole wheat or rice; and

if obstinate, cram down small pieces of the following mass: Chalk, p. caraway, and syrup of poppies.—CLATER.

2. Put chalk in their water, or give forge-water.

CROPSICK, OR CONSTIPATION.

If the obstruction is in the crop, endeavor to force the contents into the gullet and mouth by gentle pressure. When partially emptied give rue and butter.

When the obstruction is in the bowels, give bran and pollard, mixed with a little greasy hot liquor, to which, if necessary, a little sulphur may be added; or give a teaspoonful of the castor-oil mixture.—See CHIPPING.

PIP, OR BLAIN IN THE TONGUE.

1. Wash the mouth two or three times a day with a mixture of equal parts of tincture of myrrh and water.

2. Rub the sore with common salt.

3. Solution of chloride of soda 1 dr., water 1 oz., honey of roses 3 dr.

CANKER IN PIGEONS. Apply burnt alum, mixed with honey.

SCABS IN BREAST AND BACK OF PIGEONS. Dill seed, cumin seed, fennel seed, of each 1 lb, assafœtida 1½ oz., bay salt ¼ lb, common salt ¼ lb; knead them with fine clay and flour. Bake it in earthen pots, and set it for the birds to peck.—MOORE.

Genuine Salt-Cat.—Sifted gravel, brickmakers' earth, rubbish of old walls, of each a peck, cumin seed 1½ lb, baysalt ¼ lb; mix.

FOR FILANDER WORMS IN HAWKS. Aloes, iron filings, nutmegs, and honey; mix, and give a small piece as often as necessary.

CHIPPING, IN CHICKENS.—Remove the chickens to a warm place. Mix 1 oz. of castor oil with ¼ oz. of syrup of ginger; mix a teaspoonful of this with a little thick gruel, and force a little down several times a day, so that it shall get half a teaspoonful of the mixture in the course of the day.

FOR CHILL, IN TURKEY CHICKS. Give ground malt

and barley-meal in equal quantities, adding a little powdered caraway or coriander-seed.

PASTE FOR WEAK TURKEY CHICKS. Eggs boiled hard, nettles, and parsley, all chopped up, and moistened with wine or water.

MEGRIMS, OR GIDDINESS. Castor oil 1 oz., syrup of ginger ¼ oz., syrup of poppies ¼ oz.; mix with gruel, and force a little down occasionally.—CLATER.

CONVULSIONS OF DUCKS. Give to grown-up ducks 4 grains of pepper, mixed with fresh butter.

FOR BLINDNESS. Foment with warm water, then drop a few drops of the following solution into the eyes; laudanum 1 teaspoonful, water a teacupful.

LOTION FOR WOUNDS.

1. Laudanum a few drops, added to a teacupful of water.
2. Tincture of myrrh and paregoric, each a teaspoonful, water ½ a pint.

TO PROMOTE THE LAYING OF EGGS. A little sulphate of soda, placed within reach of the hens, is said to be useful. Warmth, good feeding, with a little chopped meat in winter, are also recommended. To prevent their laying soft eggs, supply them with old mortar, bruised egg-shells, or chalk.

FUNGUS, OR PROUD FLESH, FROM WOUNDS IN THE HEAD. Burnt alum, 2 dr., honey 1 oz.; mix, and apply twice a day.

VERMIN, TO DESTROY. Tobacco smoke, with good food and cleanliness.

MOULTING. It is usual to put saffron into the water of cage birds when moulting; others recommend a rusty nail.

FOR SNIFFLES IN RABBITS. Sulphate of copper 1 to 2 gr., morning and evening, in bran.—CLATER.

FOR ROT, OR POT BELLY. Give them young green broom, and bread well toasted.—MAYER.

PATENT AND PROPRIETARY MEDICINES,

DRUGGISTS' NOSTRUMS, ETC.

THIS division consists of those *medicinal compounds* which are excluded from the Pocket Formulary, as belonging rather to empirical than regular practice. It includes, in addition to those secret and patent remedies which are usually termed QUACK MEDICINES, preparations of various drugs made according to private formulæ; some favorite domestic remedies; and a few compounds which, though not empirical, are better known by the names of individual practitioners than by any other title. The supposed composition of some of the secret remedies is given on the authority of DR. PARIS, the Philadelphia College of Pharmacy, and others; but without vouching for their correctness.

ABERNETHY'S PILLS. The nostrum to which this distinguished surgeon's name has been applied is said to consist of 2 gr. of blue pill, and 3 of compound extract of colocynth.

AGUE DROPS (tasteless). A solution of arsenic, probably similar to the *liquor postassæ arsenitis* of the Pharmacopœia.

ANDERSON'S PILLS. See Pilulæ Andersonis, P. F. Other published formulæ are—

1. Barbadoes aloes 1 oz., jalap ¼ oz., soap 1 dr., oil of aniseed ½ dr., tincture of aloes q. s.; mix, and divide into 4-grain pills.

2. Barbadoes aloes 5 oz., water 1 oz.; soften by the heat of a water-bath, and add powdered jalap, powdered aniseed, and ivory-black, of each 1 oz., oil of aniseed 1 drachm.

3. Barbadoes aloes 16 oz., black hellebore, jalap, sub-carbonate of potash, of each 1 oz., oil of aniseed ½ oz., syrup of buckthorn q. s. to form a mass. To be divided into 4-grain pills.

4. Barbadoes aloes 24 oz., soap 4 oz., colocynth 1 oz., gamboge 1 oz., oil of aniseed ½ fluid oz.; mix, and divide into pills of 3 gr. each. Phil. Col. of Pharmacy.

ANODYNE NECKLACES. Beads formed of the root of henbane, and used as necklaces to allay the pain of teething.

ANTIPERTUSSIS. Dr. PARIS states that the basis of this nostrum is a salt of zinc.

ARQUEBUSADE (acid). 1. Sulphuric acid ½ ℔, vinegar and spirit of wine, of each 3 ℔, clarified honey 1 ℔.—SWEDIAUR.

2. Distilled vinegar and rectified spirit, of each 10 oz., sulphuric acid (by weight) 1½ oz., sugar 2½ oz.; mix. For the aromatic spirituous arquebusade water, see Arquebusade Water, under PERFUMERY; also Spiritus Vulnerarius, P. F.

AROMATIC VINEGAR. Strongest acetic 1 ℔, camphor 1 oz.; dissolve, and 1 oz. each of oil of lavender, oil of cloves, and oil of lemon.

APERIENT AND ANTIBILIOUS PILLS. See ANDERSON'S, BAILLIE'S, BARCLAY'S, DIXON'S, &c., Pills. The following are useful forms:

1. Compound extract of colocynth 60 gr., comp. rhubarb pill 30 gr., soap 10 gr. In 24 pills.

2. Compound extract of colocynth 2 dr., extract of rhubarb ½ dr., compound soap pill 10 gr.; mix, and divide into 40 pills; 1, 2, or 3 for a dose.

3. Compound extract of colocynth 8 oz., soap 1 oz., scammony 2 oz., extract of rhubarb 2 oz., oil of cassia 5 dr., spirit q. s. to form a mass. Divide into 4 grain pills.

4. Blue pill, compound extract of colocynth, of each a scruple; scammony, and Castile soap, of each 10 gr., oil of caraway 4 drops; mix, and divide into 15 pills—3 at bedtime.—Sir B. BRODIE.

5. Compound extract of colocynth 4 scruples, scammony a scruple, extract of rhubarb 12 gr., soap 6 gr., oil of cinnamon 4 drops; mix, and divide into 24 pills.—Mr. VANCE.

6. Mr. VANCE'S *Stronger Pills, with Calomel.* Compound extract of colocynth 4 scruples, scammony 2 scruples, calomel 24 gr., oil of cinnamon 6 drops, in 24 pills.—Dr. J. JOHNSON.

7. Compound extract of colocynth 1 dr., calomel 15 gr., emetic tartar 1 gr., oil of cassia 5 drops. In 24 pills.—Dr. J. JOHNSON.

8. Scammony 10 to 15 gr., compound extract of colocynth 2 scruples, extract of rhubarb ½ dr., soap 10 gr., oil

caraway 5 drops. In 20 pills. One or two when required. —Sir C. Scudamore.

9. Compound rhubarb pill ½ dr., ipecacuanha 6 dr., compound extract colocynth 20 gr. In 12 pills. One or more at bedtime occasionally.—Dr. Baron.

10. The same as Pil. Coloc. et Hyoscyami, Ed. Ph.—Dr. Hamilton.

11. (Strong.) Compound extract of colocynth 2 dr., aloes and myrrh pill 2 dr., calomel 1 dr.; mix, and divide into 40 pills. Two for a dose.—Dr. Lynn.

12. Compound extract of colocynth 2 scruples, ipecacuanha 6 gr., soap 10 gr., extract of henbane 30 gr. In 18 pills. Two at bedtime.—Dr. Copland.

13. Dr. Neligan's *Purgative Pills, for general use.* Comp. colocynth pill, and soap of jalap, of each 1 dr. In 24 pills. Two when required.

14. (Without aloes.) Simple extract of colocynth 24 gr., extract of jalap 12 gr., blue pill 12 gr., ipecacuanha 4 gr., oil of peppermint 3 drops. In 12 pills.

Other formulæ will be found in the P. F. See Pilulæ Colocynthidis; Pil. Coloc. c. Oleo Crotonis; Pil. Catharticæ; Pil. Jalapæ; Pil. Purgantes, &c. Those which do not contain calomel should be preferred for general and repeated use.

Atkinson's Infant Preservative. Carbonate of magnesia 6 dr., white sugar 2 oz., oil of aniseed 20 drops, spirit of sal volatile 2½ dr., laudanum 1 dr., syrup of saffron 1 oz., caraway water to make a pint.

Atropine Paper. Green tissue paper imbued with a solution of sulphate of atropia, so that a piece one-fifth of an inch square contains as much as a drop of a solution 2 gr. to 1 oz. of water. The paper is hung up and turned about while drying. A piece of the size named will dilate the pupil if placed on the sclerotic and the lids closed over it and tied with a handkerchief.—Mr. Streatfield.

Bacher's Tonic Pills. Alkaline extract of black hellebore 2 dr., extract of myrrh 2 dr., powder of holy thistle 1 dr., mix, and divide into 4-grain pills.

Dr. Baillie's Pills. Compound extract of colocynth 1½ dr., extract of aloes 1½ dr., Castile soap ½ dr., oil of cloves 15 drops: in 38 pills. 3 at bedtime occasionally.

Dr. Baillie's Dinner Pills. Aloes 20 gr., ginger ½ dr., ipecacuanha 8 gr., syrup q. s. Mix, and divide into 16 pills. One daily, before dinner.

Bailey's Itch Ointment. Olive oil 1 ℔, suet 1 ℔, alkanet root 2 oz. Melt, and macerate until colored; then strain, and add 3 oz. each of alum, nitre, and sulphate of zinc, in very fine powder; adding vermilion to color it, and oil of aniseed, lavender, and thyme to perfume.

Baking Powder. Tartaric acid 8 oz., bicarbonate of soda 9 oz., arrowroot, or rice flour, 10 oz. Mix. Delfort's is said to consist of alum 5 oz., bicarbonate of soda 2¾ oz., bicarbonate of ammonia ½ oz., arrowroot 4 oz.

Balm of Gilead (factitious). 4 oz. of gum benzoin may be dissolved by heat in 1 ℔ of Canada balsam, and to the mixture, when cold, ¼ oz. each of the oils of rosemary, lemon, and cassia, added.

Balm of Rakasiri. Oil of rosemary dissolved in common gin.

Balsam. See Ford's, Hill's, Fryar's, &c.

Barclay's (Rev. D.) Antibilious Pills. Extract of colocynth 2 dr., soap of jalap 2½ dr., extract of guaiacum wood 3 dr., emetic tartar 8 gr., oil of juniper, caraway, and rosemary, each 4 drops; into 4-grain pills.

Baregian Balls. Extract of soapwort (or of artichoke leaves) 3 oz., gelatine 1½ oz., water 3 oz.; heat together till dissolved, pour the solution into a warm iron mortar; add 6 oz. of sulphuret of lime, and 1 oz. of salt, previously powdered and mixed. Stir constantly till a mass is obtained, and divide it into balls of 2¼ oz. each. Use one for a general bath, half of one for a foot bath.

Bark, Essential Salt of. See Extractum Cinchonæ Siccatum, P. F.

Baker's Tooth Tincture. An alcoholic solution of pyrethrum, colored with tincture of red cabbage.

Bateman's Pectoral Drops. 1. Compound spirit of aniseed 16 fluid ounces, opium 1 dr., camphor 1 dr., oil of fennel 20 drops, cochineal 2 dr.

2. Proof spirit 4 gal., red sanders 2 oz.; digest 24 hours, filter, and add powdered opium 2 oz., camphor 2 oz., catechu 2 oz., oil of aniseed 4 fluid drachms; digest

for 10 days. *Philadelphia College of Pharmacy.* The old wine gallon is here intended.

Bateman's Itch Ointment. Carbonate of potash ½ oz., rose-water 1 oz., vermilion 1 dr., sulphur 11 oz., oil of bergamot ½ dr., lard 11 oz.; mix.

Bates's Anodyne Balsam. Soap liniment 2 parts, tincture of opium 1 part.

Bates's Camphorated Eye-water. Sulphate of copper 15 gr., French bole 15 gr., camphor 4 gr., boiling water 4 oz.; infuse, strain, and dilute with 4 pints of cold water.

Bates's Styptic Wash. See Liquor Aluminis Co., P. F.

Bathing Spirits. These resemble liquid opodeldoc (soap liniment), and are usually colored by the addition of some dark tincture. See Freeman's Bathing Spirits.

Bath Digestive Pills. Rhubarb 2 oz., ipecacuanha ½ oz., cayenne pepper ¼ oz., soap ½ oz., ginger ¼ oz., gamboge ¼ oz.; mix, and divide into 4-grain pills.

Bath Lozenges (in imitation of Dawson's). Pure extract of liquorice 1 oz., powdered gum arabic 1 oz., white sugar 1 lb, hot water q. s. to form a mass; to be rolled into pipes.

Battley's Liquor Cinchonæ and Liquor Opii. See Pocket Formulary.

Battley's Senna Powder. Senna leaves heated until they become light in color, reduced to powder, and mixed with some finely powdered charcoal.

Baume de Vie. Socotrine aloes 2 dr., rhubarb 6 dr., saffron 2 dr., liquorice-root 1 oz., proof spirit 8 oz., digest for 8 days and filter. The original Swedish form is this: Aloes 9 dr., rhubarb, gentian, zedoary, saffron, theriaca, agaric, of each a drachm, proof spirit 2 pints.

Baynton's Plaster. Simple litharge plaster 16 oz., yellow resin 6 dr.; melt together, and spread on linen or calico.

Beddoe's Pills; for Gravel, &c. Carbonate of soda, dried without heat, 1 dr., soap 4 scruples, oil of juniper 10 drops, syrup of ginger q. s. for 30 pills.

Beetle Wafers. Red lead, sugar, and flour; made in wafer-irons.

BELLOSTE'S PILLS. Quicksilver, scammony, and jalap, of each 1 ℔, sugar 4 oz.; mixed and made up into a mass with sherry wine.

BESTUCHEFF'S NERVOUS TINCTURE. A mixture of a strong solution of perchloride of iron with sulphuric ether and spirit, exposed in long bottles to the rays of the sun until it has quite lost its brown color.

BETTON'S BRITISH OIL. Oil of turpentine 8 oz., Barbadoes tar 4 oz., oil of rosemary 4 dr.; mix. See British Oils.

BEWLEY & EVANS' CHALYBEATE WATER. Citrate of iron 13 grains, carbonated water 6 oz., syrup of orange-peel 1 oz.

BISCUITS, APERIENT. An ounce of powdered jalap, mixed with 16 oz. of the materials for gingerbread, or other kind of cake. See Gingerbread, Purgative.

BLACK DROP (Lancashire). Fine opium, cut small, 8 oz., verjuice 48 fluid oz., nutmegs grated 1½ oz., saffron ¼ oz., boil together until the opium is dissolved; add sugar 4 oz., yeast 2 tablespoonfuls. Keep it near the fire for 6 or 8 weeks, then place it in the open air till it becomes a syrup; decant and filter.

BLACK DRAUGHT. 1. Infusion of senna 10 dr., sulphate of magnesia 3 dr., syrup of ginger 1 dr., aromatic spirit of ammonia 20 drops.

2. Tartrate of potash 1½ dr., manna ½ dr., tincture of jalap ½ dr., aromatic spirit of ammonia 20 drops, extract of liquorice 4 gr., infusion of senna 11 dr. See Mistura Sennæ Comp., Haustus Sennæ, and Mistura Aperiens, in Pocket Formulary, for other formulæ.

BLAINE'S DISTEMPER POWDERS. The basis of these is said to be aurum musivum (sulphuret of tin).

BLISTERING PAPER. Melt cantharidine 1 dr., white wax 1 dr., olive oil 5 dr. Paint it with a brush on white bibulous paper, and hang it up to dry in a current of air. Take a piece of pink paper of form and size required, paint the under colored side with a weak solution of India-rubber, cut the cantharidine paper to the size, less a margin of the pink paper, and place it on while the India-rubber solution is still sticky. Before applying, the blister must be held over the steam of hot water. See PAPIER EPISPASTIQUE.

BLISTERING TISSUE. *Taffetas Vesicant.* Powdered cantharides exhausted by ether, the tincture distilled to recover the principal part of the ether for the same use, and the residue heated in a water-bath till it ceases to boil. The green butyraceous oil which remains is to be melted with twice its weight of wax, and spread on waxed silk, or any convenient and adhesive material. An extract prepared by evaporating a tincture made with 4 parts of flies, 1 of strong acetic acid, and 16 of rectified spirit, is used for the same purpose.

BÖCHET'S SYRUP, *for scrofulous affections.* Compound syrup of sarsaparilla, with senna, and 1 per cent. of iodide of potassium.

BRANDISH'S ALKALINE SOLUTION, OR CAUSTIC ALKALI. American pearl ashes 6 ℔, quick-lime 2 ℔, wood ashes prepared by burning the branches of the ash 2 ℔, boiling water 6 old gallons (5 imp.); slake the lime, add the rest of the water and the pearl ashes, and lastly stir in the wood ashes; let it stand in a covered vessel for 24 hours, and decant. To each pint add one drop of true oil of juniper berries. Keep it in stoppered bottles of green glass.

BRANDISH'S ALKALINE TINCTURE OF RHUBARB. Coarsely powdered rhubarb 1 oz., alkaline solution (BRANDISH'S) 32 fluid ounces. The original formula directs only ½ oz. rhubarb, but as smaller doses than were given by Mr. B. are now usually prescribed, the quantity of rhubarb is here increased. Or an alkaline infusion of rhubarb may be made by pouring boiling water 8 parts on rhubarb 3 parts, and carbonate of potash 1 part.

BRITISH HERB TOBACCO. The principal ingredient in this compound is dried coltsfoot leaves, to which a smaller portion of thyme, wood-betony, eyebright, and rosemary, are added.

BRITISH OILS. Oil of turpentine, and linseed oil, of each 8 oz.; oil of amber, and oil of juniper, of each 4 oz.; true Barbadoes tar 3 oz.; American petroleum (seneca oil) 1 oz.; mix. See BETTON'S BRITISH OIL, above.

BRODUM'S NERVOUS CORDIAL. Iron wine, compound spirits of lavender, tinctures of calumba, gentian, cinchona, and cardamoms, equal parts of each.

BROCCHIERI'S STYPTIC WATER. Pieces of fresh pine, bruised in a mortar, and distilled with twice their weight of water, till half the water has come over. After standing in a wide vessel, any floating oil is to be removed from the surface, and the water kept for use.

BURNETT'S (SIR WILLIAM) DISINFECTING FLUID. A neutral solution of zinc in commercial muriatic acid.

CACHOU AROMATISE. See PERFUMERY.

CAJEPUT LINIMENT. Soap liniment 7 oz., camphor ½ oz., oil of cajeput 1 oz.

CAMPHOR LINIMENT, EXTEMPORANEOUS. Rectified spirit 17 fluid oz., strong water of ammonia 2½ oz., camphor 2 oz., oil of lavender 50 minims.

CARRON OIL. Lime-water and linseed oil, equal quantities.

CASTILLON'S POWDERS. Sago meal, salep, tragacanth, each 1 dr., prepared oyster-shells a scruple; colored with cochineal. A drachm to be boiled with milk, in bowel complaints.

CEPHALIC SNUFF. Dried asarabacca leaves 3 parts, marjoram 1 part, lavender flowers 1 part; rub together to a powder. BOELI'S consists of 2 dr. valerian, 2 dr. of snuff, 3 drops of oil of lavender, 3 drops of oil of marjoram; mix. This is said to relieve the eyes as well as the head.

CHAMBERLAIN'S PILLS. Common milk of sulphur, and vermilion. Dr. PARIS'S statement that they contain sulphate of lime would probably surprise the proprietor, if not aware that a great part of the commercial milk of sulphur contains half its weight of that substance.

CHAMOMILE DROPS. Dr. PARIS states that the nostrum sold under this name is merely spirit flavored with essential oil of chamomile. A strong tincture of the flowers would probably be more efficacious.

CHAMOMILE PILLS. We are not aware of the composition of NORTON'S chamomile pills. The following is a good form: Watery extract of aloes 12 gr., extract of chamomile 36 gr., oil of chamomile 3 drops; make 12 pills. Two every night, or twice a day.

CHELSEA PENSIONER. Powdered rhubarb 2 dr., cream of tartar 1 oz., guaiacum 1 dr., sulphur 2 oz., 1 nutmeg grated

fine, clarified honey 16 oz.; mix. Take 2 spoonfuls night and morning; for chronic rheumatism, &c.

CHELTENHAM SALTS (factitious). Sulphate of soda 16 oz., sulphate of magnesia 8 oz., muriate of soda 1 oz., sulphate of iron 8 gr.; dissolve in the smallest quantity of hot water, strain, and evaporate to dryness by a gentle heat, or dry the salts separately, and mix.

CHILBLAINS, Popular Remedies for. 1. Soap liniment 1 oz., cajeput oil ¼ oz., tincture of cantharides ¼ oz.; mix.

2. Sal ammoniac ½ oz., vinegar 5 oz., spirit of rosemary 1 oz.; mix.

3. Oil of turpentine 1 oz., camphor ¼ oz., Goulard's extract ¼ oz.; mix.

4. Dr. GRAVES'S *Preventive.* Sulphate of copper 10 gr., water 1 oz.; dissolve. Brush over the parts with the lotion by means of a camel-hair pencil, and when dry apply a little simple ointment; repeat this for some evenings in succession.

5. LEJEUNE'S *Balsam.* Camphor 1½ dr., tincture of benzoin 1 oz., iodide of potassium 3 dr., extract of lead 2 oz., a mixture of equal parts of rectified spirit and rosewater 4 oz.; mix the above with a solution of 2 oz. of soap in 4 oz. of the same diluted spirit; mix the whole, adding a few drops any essential oil.

6. Sal enixum, alum, and sulphate of zinc, of each ¼ oz.; water a pint; apply it frequently.

7. Muriatic acid ½ oz., Fryar's balsam 3½ oz.; mix.

8. SWEDIAUR'S *Paste.* Bitter almonds 8 oz., honey 6 oz., powdered camphor ½ oz., flour of mustard ½ oz., burnt alum ¼ oz., olibanum ¼ oz., yolks of 3 eggs; beat together to form a paste. Rub a portion of it on the part affected, moistened with water, night and morning; then wash with warm water, and dry with a cloth.

9. WAHLER'S *Ointment for Broken Chilblains.* Black oxide of iron, bole, and oil of turpentine, of each 1 dr.; rub together, and add the mixture to 1 oz. of melted resin cerate.

10. *Another Ointment for the same.* Locatelli balsam 1 oz., citrine ointment ¼ oz., balsam of Peru 20 drops; mix.

11. *Russian remedy.* Dry the peelings of cucumbers, and when required for use soften the inner part with water, and apply it to the part affected.

CHING'S WORM LOZENGES. The yellow lozenges contain 1

gr. of calomel in each, with sugar and sufficient mucilage (colored with saffron) to form a paste. The brown contain ½ gr. of calomel, with 3½ gr. of resinous extract of jalap, according to GRAY; or with 1 gr. of resin of jalap, according to Dr. PARIS and others.

CHLORODYNE. The preparation sold under this name contains chloroform, morphia, Indian hemp, and prussic acid. Many formulæ have been published. The following is an improvement by Mr. GROVES, on the recipe of Dr. OGDEN. Take chloroform 4 dr., ether 1½ dr., oil of peppermint 8 drops, resin of Indian hemp 16 gr., capsicum 2 gr.; macerate for 2 or 3 days, and filter. Then dissolve hydrochlorate of morphia 16 gr. in 1 oz. of syrup, add perchloric acid and water ½ dr. each, assisting the solution by a water-bath; then, when cold, add hydrocyanic acid (SCHEELE'S) 96 drops. Mix the solutions.

CHOLERA MEDICINES. The following are some of the more popular remedies that have been used during the visitations of this disease:

1. *Liverpool Preventive Powders.* Bicarbonate of soda 1 scruple, ginger 8 gr.; to be taken in a glass of water after breakfast and supper. These powders are said to have been used with good effect among the workmen in the mining and manufacturing districts, during a former visitation of cholera.

2. Dr. STEVENS'S *Saline Powders.* Bicarbonate of soda ½ dr., muriate of soda 1 scruple, chlorate of potash 7 gr.; mix, for 1 dose.

3. Mr. HOPE'S *Remedy.* Nitrous acid (red) 2 dr., peppermint-water or camphor mixture 1 oz., tincture of opium 40 minims. Dose, 1 to 2 teaspoonfuls in a cupful of gruel every 3 or 4 hours.

4. Spirit of wine 1 oz., spirit of lavender ¼ oz., oil of origanum ¼ oz., compound tincture of benzoin ½ oz., spirit of camphor ¼ oz. Twenty drops on moist sugar; to be rubbed outwardly also.

5. *American Remedy.* Equal parts of lard, maple-sugar, and charcoal; to be mixed and the size of a nut swallowed.

6. Remedies recommended by the Board of Health, in premonitory diarrhœa: Chalk mixture 1 oz., aromatic confection 10 to 15 gr., tincture of opium 5 to 15 drops; to be repeated every 3 or 4 hours, or oftener if the attack be severe, until the looseness is stopped.

7. Dr. GRAVES'S *Astringent Pills.* Acetate of lead 20 gr., opium 1 gr.: in 12 pills. One every half hour till the watery discharges cease.

8. Mr. BUXTON'S *Remedy.* Twenty-five minims of diluted sulphuric acid in an ounce of water.

We have inserted the above, not to encourage quackery in reference to this terrible disease, but because the druggist may be called upon to supply these remedies, and expected to know their composition. For Elixir Woroneje, see P. F.

CHIRAYTA PILLS AND MIXTURE. Dr. REECE'S *Pills.* Extract of chirayta 2 dr., dried soda 20 gr., ginger 15 gr.; mix, and divide into 36 pills. Two twice a day. *Mixture:* Infusion of chirayta 8 oz., subcarbonate of soda 1 dr.; two tablespoonfuls 3 times a day.

CLUTTON'S FEBRIFUGE SPIRIT AND TINCTURE. *Spirit:* The original formula is—oil of sulphur by the bell, oil of vitriol and sea salt, of each 1 oz., rectified spirit 3 oz.; mix, digest for a month, and distil to dryness. *Tincture:* Febrifuge spirit 8 fluid oz., angelica root, serpentary, cardamom seed, of each 1½ dr.; digest and strain. Water acidulated with these and sweetened to the taste, forms a cooling diuretic and diaphoretic julep. Though never admitted into the Pharmacopœias, these preparations are favorites with a few practitioners.

COCHRANE'S COUGH MEDICINE. An acidulated syrup of poppies.

COLLIER'S (Dr.) WINE OF QUININE. Disulphate of quinine 18 gr., citric acid 15 gr., sound orange wine 1 bottle, or 24 fluid oz.

COLLIER'S (Dr.) CREAM OF TARAXACUM. See Cremor Taraxaci, P. F.

COLLINS'S DISINFECTING POWDER. See Disinfecting Compounds, among the Trade Chemicals.

CONSUMPTION, *Popular Remedies for.* 1. Rum ½ pint, linseed oil, honey, garlic (beaten to a pulp), and loaf sugar, of each 4 oz., yolks of 5 eggs; mix: a teaspoonful night and morning.

2. *Breastplate.* Dissolve 1 oz. of aloes in 12 oz. of a strong decoction of fresh rue: fold a large piece of soft muslin in 8 folds, large enough to cover the chest and part

of the stomach; steep it in the solution and dry it in the shade: wear it on the chest constantly.

COUGH LINTUS. 1. *Rose Linctus.* Confection of roses 3 oz., paregoric elixir 1½ oz., diluted sulphuric acid 1 dr. and a half; mix: a teaspoonful now and then when the cough is troublesome.

2. Dr. LATHAM'S *Cough Linctus.* DOVER'S Powder ½ dr., compound powder of tragacanth 2 dr., syrup of Tolu ½ oz., confection of hips, and simple oxymel, of each, 1 oz.; a teaspoonful 3 or 4 times a day. For other Formulæ, see Linctus; Linctus Oleosus; Linctus Pectoralis, &c.; P. F.

COUGH LOZENGES. See Bath Lozenges, above; also Lozenges, below.

CORN PLASTERS. See KENNEDY'S Corn Plaster, and Emplastrum Æruginis, Pocket Formulary. Most of the advertised corn plasters contain verdigris. A few additional formulæ are subjoined.

1. Galbanum Plaster 1 oz., prepared verdigris 1 scruple; melt and mix.

2. Galbanum 1 oz., black pitch ½ oz., simple diachylon ¼ oz., verdigris a scruple, sal ammoniac a scruple. Melt together the first three, and add the last two in fine powder.

3. Plaster of ammoniacum with quicksilver 1½ oz., soap plaster ½ oz., opium in fine powder ½ dr.

MECHANICAL CORN PLASTERS. Any suitable adhesive plaster is spread on soft thick leather (buckskin), which is afterwards cut to a suitable size, and a hole punched in the centre. They are sometimes spread on amadou, or on vulcanized India-rubber.

CORN SOLVENTS. One of the preparations sold under this name is probably a strong solution of subcarbonate of potash. A powder sold for the same purpose consists of subcarbonate of potash colored with ochre or bole. A pinch is placed on the corn, and confined by means of a piece of adhesive plaster or rag. Sir Humphrey Davy's name has been given to a remedy which consists of subcarbonate of potash and salt of sorrel, similarly applied. The following is one of the advertised Corn and Bunion remedies: Subcarbonate of soda 1 oz., finely powdered and mixed with ½ oz. of lard. Applied on linen rag every night: the outer

skin to be pared off every 2 or 3 days. It may be varied thus: Dried soda 4 dr., powder blue (smalts) a scruple, lard 4 dr.; mix.

Caustic for Corns. Tincture of iodine 4 dr., iodide of iron 12 gr., chloride of antimony 4 dr.; mix, and apply with a camel-hair brush, after paring the corn. It is said to cure in 3 times.

Court Plaster is made by repeatedly brushing over stretched sarcenet with a solution of 1 part of isinglass in 8 of water mixed with 8 parts of proof spirit, and finishing with a coat of tincture of benzoin, or of balsam of Peru. See Miscellaneous Preparations.

Custard Powder. See Dietetic Articles.

Daffy's Elixir. This is similar to the compound tincture of senna: but different makers have their peculiar formulæ. The following are some of them. Avoirdupois weight seems to be intended.

1. Senna leaves $3\frac{3}{4}$ lb, jalap, aniseed, caraway seed, of each 20 oz., rectified spirits 18 pints, sugar 5 lb. Infuse the senna 2 or 3 times in sufficient boiling water to yield, when strained with pressure, 4 gallons in the whole. Add to this the tincture made with jalap and seeds digested with the spirit for a week. Pour off the clear liquor and add the sugar and brandy coloring if required.

2. Dicey's, according to Gray. Senna 5 oz., guaiacum shavings (some recipes add red sanders), dried elecampane root, seed of anise, coriander, and caraway, and root of liquorice, of each $2\frac{1}{2}$ oz., stoned raisins 8 oz., proof spirit 6 lb.

3. Swinton's. Jalap 3 lb, senna 2 oz., coriander seed, aniseed, liquorice root, and elecampane, of each 4 oz.; spirit of wine and water, of each a gallon.

4. Small senna 10 oz., bruised jalap, coriander seed, and aniseed, of each $2\frac{1}{2}$ oz.; proof spirit a gallon. Digest 8 days, frequently shaking, and strain. Pour on the remaining ingredients 6 oz. of boiling water in which 2 dr. of salt of tartar have been dissolved; press strongly, and add the liquid to the tincture, with 3 oz. of treacle. Some recipes add rhubarb, in the proportion of about 4 oz. to the gallon.

Dalby's Carminative. 1. Carbonate of magnesia 1 oz., syrup of poppies 5 oz., tincture of wood soot 1 oz., oil of

caraway 25 drops, oil of peppermint 16 drops, water and spirit of wine, each ½ oz. Mix.

2. Carbonate of magnesia 2 scruples, oil of peppermint 1 drop, oil of nutmegs 2 drops, oil of aniseed 3 drops, tincture of castor 30 drops, tincture of assafœtida 15 drops, tincture of opium 5 drops, spirit of pennyroyal 15 drops, compound tincture of cardamom 30 drops, peppermint water 2 oz. Mix. —Dr. PARIS.

DANDELION COFFEE. The roots, collected at the end of the year, are dried at a gentle heat and reduced to powder. Some mix coffee with it. Others *roast* the root in the manner of coffee, but probably at the expense of its medical virtues. The better way is to dry and powder it, and direct it to be mixed with coffee when used. If considered necessary to give it more color and flavor, it may be previously mixed with a sufficient quantity of roasted chicory, which should not exceed one-eighth of the whole.

DARCET'S ALKALINE LOZENGES, or VICHY LOZENGES. Bicarbonate of soda 2 dr., refined sugar 14 oz., oil of peppermint 4 drops, mucilage of tragacanth q. s. Mix, and divide into 60 lozenges.

DAWSON'S LOZENGES.—See BATH LOZENGES, above.

DELAMOTT'S GOLDEN DROPS. Muriate of iron 1 oz., spirit of sulphuric ether 7 oz.; dissolve and expose to sunshine in a closely-stopped bottle till it becomes divested of color. See BESTUCHEFF'S Nervous Tincture.

DERBYSHIRE'S PATENT EMBROCATION FOR PREVENTING SEA-SICKNESS. Boil 2 oz. of opium, 2 dr. of extract of henbane, 10 gr. of mace, and 2 oz. of mottled soap, in 3 pints of water for ½ hour. When cold, add 1 quart of rectified spirit, and 3 dr. of spirit of ammonia.

DESHLER'S SALVE. This is merely resin cerate.

DIGESTIVE PILLS. See BATH DIGESTIVE PILLS, DINNER PILLS, BAILLIE'S PILLS, Lady WEBSTER'S PILLS, Dr. REECE'S CHIRAYTA PILLS.

DINNER PILLS. See BATH DIGESTIVE PILLS, Lady WEBSTER'S PILLS, &c. The following are a few additional formulæ:

1. Rhubarb 30 gr., aloes 60 gr., ipecacuanha 12 gr., tincture of ginger q. s. to form a mass; to be divided into 24 pills.

2. Sir Charles Bell's. Rhubarb 50 gr., mastic 6 gr., sulphate of quinine 4 gr.; in 12 pills.

Dixon's Antibilious Pills. Equal parts of aloes, scammony, and rhubarb, with the addition of a small quantity of tartar emetic, and made up with Castile soap.

Dover's Powders. The Pulvis Ipecacuanha Compositus of the Pharmacopœia. But the original powder consisted of nitre and sulphate of potash, each 4 oz., fused in a red-hot mortar, and afterwards reduced to powder, and mixed with 1 oz. each of ipecacuanha, opium, and liquorice.

Depuytren's Pommade. See Hair Cosmetics.

Depuytren's Eye Salve. Nitric oxide of Mercury 10 gr., sulphate of zinc 20 gr., lard 2 oz.; rub perfectly smooth.

Duncan's Fluid Extract of Senna. See Extractum Sennæ Fluidum, Pocket Formulary.

Dr. Duncan's Lactucarium Lozenges. As the Trochisci Opii (Pocket Formulary), substituting lactucarium for opium.

Duncan's Gout Remedy. A preparation of colchicum with opium, &c.

Dutch (or Haerlem) Drops. The basis of this popular remedy is said to be the residue which is left in redistilling oil of turpentine. The following is one of the imitations of it made in this country: Linseed oil 1 quart, resin 2 ℔, sulphur 1 ℔; boil together over a slow fire; when combined remove from the fire, and add 1 pint of oil of turpentine and 50 drops of liquor of ammonia; stir well together and bottle.

Eaton's Styptic. It is similar to that of Helvetius, which see below.

Eau de Magnanimite. A tincture of ants, with aromatics.

Eau Medicinale d'Husson. It is prepared, according to Dr. Williams, from the juice of colchicum flower with half the quantity of brandy; mix, and after standing a few days, decant into small bottles. But it was more probably made from the root, as prescribed in the following formulæ.—(In one of the French codices.) *Eau colchique d'Husson.* Dry colchicum 60 parts in sherry 125 parts. 20 drops for a dose. (According to Mr. Want.)—4 ounces of the fresh root sliced, macerated in ½ pint of proof spirit.

EAU DE COLOGNE, EAU DE MELISSE, &c. See PERFUMERY.

EAU DE JARVELLE. Dry chloride of lime 2 oz., carbonate of potash 4 oz., water 2 pints; mix the chloride with 1½ pint of water, dissolve the potash in the remainder; mix the solutions, and filter.

EAU DE LUCE. See PERFUMERY.

EAU DE RABEL. See Acidum Sulphuricum Alcoholisatum, Pocket Formulary.

EDINBURGH OINTMENT. White hellebore powder, sal ammoniac, and lard.

ELLERMAN'S DEODORIZING FLUID. It consists chiefly of persalts of iron. See DISINFECTING COMPOUNDS.

ELIXIR DE GARUS. See Pocket Formulary.

ELIXIR LONGÆ VITÆ. Similar to BAUME DE VIE, above.

ELIXIR OF HALLER. See Elixir Acidum Halleri, Pocket Formulary.

ELIXIR PAREGORIC. See Tinctura Camphoræ Composita, Pocket Formulary.

ELIXIR OF VITRIOL (Mynsicht's Elixir). See Acidum Sulphuricum Aromaticum, Pocket Formulary. For common sale druggists frequently keep a more ready and economical preparation, of which the following is one form: Compound tincture of cardamoms 1 ℔, tincture of cinnamon 3 ℔, cinnamon water 2 ℔; mix, and add gradually 1½ ℔ of pure sulphuric acid.

ERVALENTA, and REVALENTA. See DIETETIC COMPOUNDS.

ESSENCES. Essences of flowers will be found under PERFUMERY. Essence of Celery, and other *culinary essences*, will be found, with allied compounds, in another place. A few concentrated infusions, and other strong preparations of drugs, not sanctioned by the colleges, but very generally used, may be noticed here.

ESSENCE OF YELLOW BARK. Resinous extract of bark ½ oz., sulphate of quinine 60 gr., rectified spirit 6 oz., tincture of orange-peel 2 oz. For Mr. BATTLEY'S preparation, see Liquor Cinchonæ, Pocket Formulary.

ESSENCES OF CALUMBA, RHUBARB, SENNA. See Liquor Calumbæ, Rhei, Sennæ, Pocket Formulary.

Essence of Camphor. See Liquor Camphoræ, Pocket Formulary.

Essence of Chamomile. As a substitute for the infusion, it may be made as Liquor Calumbæ, P. F. See Chamomile Drops for another preparation of this drug.

Essence of Ergot. See Essentia Secalis Cornuti, Pocket Formulary.

Essence of Ginger. Unbleached Jamaica ginger in coarse powder 5 oz., rectified spirit a pint; digest for 8 days, and strain with pressure; or it may be made by percolation. As there is no established form, it varies in strength as prepared by different makers, and often contains cayenne pepper.

Essence of Cubebs. Mix powdered cubebs with ether, in a well-stopped bottle; in twelve hours put the paste into a percolator, and add ether till the cubebs are nearly exhausted; distil off the ether in a water-bath, and preserve it for the same purpose. Dissolve the extract which remains in three times as much brandy. One drachm is equal to 2 drachms of the powder. A fluid extract is also made by concentrating the tincture.

Essences of Mint, Peppermint, and Pennyroyal. The strength of these varies as prepared by different makers; some use 1 part of the essential oil to 3 of rectified spirit, but more usually, we believe, 1 part to 7. They are sometimes colored with the leaves of the plant or of spinach.

Essence of Mustard. Rectified spirit of turpentine 16 fluid oz., bruised black mustard seed 2 oz., camphor 4 oz., oil of rosemary ½ oz., annotto to color.

Essence of Sarsaparilla. (See also Extractum Sarzæ Fluidum et Liquidum, Liquor Sarzæ, and Essentia Sarsaparillæ, in Pocket Formulary. The latter is an elegant and efficacious preparation.) Jamaica sarsaparilla 16 oz., lukewarm distilled water (100° to 112° F.) sufficient to cover it. Macerate for 6 hours, and strain. Bruise the root, macerate it again in sufficient warm water, and repeat the maceration with fresh water until it ceases to be much colored. After such straining, let the liquid be immediately heated to 180° F., allowed to cool, and filtered. Evaporate the whole of the filtered liquids by a water or steam-bath, at

a heat not above 160°, until reduced to 14 or 15 fluid ounces; add 2 ounces of rectified spirit, and keep it in a close bottle in a cool place for a few days. Then carefully pour off the clear liquid from any sediment into a clear dry bottle. 1 fluid ounce represents 1 ounce of the root, or 8 ounces of the decoction.

Compound Essence of Sarsaparilla. Jamaica sarsaparilla 16 oz.; proceed as above, but reserve the liquor of the last maceration for boiling the other ingredients; namely, guaiacum raspings, bruised liquorice root, sassafras, each 2 oz., mezereum ¾ oz. Boil them in 4 or more pints of the weak infusion for ½ an hour, and strain; evaporate to 4 fluid ounces; let it cool, stirring it occasionally, and add 2 oz. of rectified spirit in which a few drops of oil of sassafras have been dissolved. Evaporate the sarsaparilla liquid to 11 ounces, and when cool, add the other liquid. Proceed as for the former. One measure with 7 of water forms a near approximation to the Pharmacopœia Decoction.

Essences (concentrated infusions) of quassia, cascarilla, chiretta, gentian, &c., may be made as directed for Liquor Calumbæ, P. F. Take 8 times the quantity of ingredients directed in the Pharmacopœia for one pint of infusion, and infuse then in one pint of boiling water for the time prescribed; strain with strong pressure, and again infuse the ingredients in nearly as much water as the liquor obtained is short of a pint. Strain again with pressure; mix the products, which will measure 18 or 19 oz.; add 2 oz. of rectified spirit, set aside for a few days in a well-closed bottle, and filter. Some substances, as chiretta, senna, calumba, &c., yield their active principles to cold water, which some prefer in these cases; but then it is necessary before adding the spirit, to place the liquor (in a bottle) in a water-bath, and heat it to 180°, in order to precipitate any albumen it may contain. When cold, filter and add the spirit. They may also be made by percolation.

Vinous Essences (by fermentation). Dr. B. Lane has proposed to produce strong and permanent solutions of vegetable medicinal substances, by fermenting concentrated infusions with sugar and yeast. Further experiments are necessary to determine the value of these preparations, and the best means of producing them; an outline of the process is

therefore all that can now be given. A strong infusion of the drug, usually 4 times as strong as that of the Pharmacopœia, is mixed with lump sugar, usually 3 ℔ to 7 *old wine* pints, and fermented with yeast at the temperature of 70° to 80° Fahrenheit, for 10 or 20 hours. When the action subsides, the vessels are closed, and placed in a cool cellar, about 55°, till fit for bottling. Wines of senna, gentian, calumba, cascarilla, valerian, &c., are thus made. In making vinous liquor of opium, the opium is macerated in water for some days, strained through coarse canvas, and the liquor submitted to fermentation. It should be made twice the strength of the tincture. A new name would be required to distinguish these preparations from the *Vina* of the Pharmacopœias, and the latter also from the *Vinum opii fermentatione paratum* of the French Codex.

Extracts, Medicinal. See Pocket Formulary.

Extracts of Flowers. See Perfumery, in this volume.

Extract of Malt. Evaporate sweet wort to the consistence of treacle. It is sold as a cough medicine.

Fairthorn's (Dr.) Mild Provisional Pills. Sulphate of potash 1 scruple, extract of aloes 2 scruples, extract of senna 1 scruple, compound gamboge pill 50 gr., tartarized antimony 2 gr., compound powder of scammony 12 gr., Peruvian balsam 6 gr.; in 30 pills; one, two, or more occasionally, when required.

Ford's Balsam of Horehound. It contains the ingredients of paregoric elixir, with squills, honey, and a strong infusion of horehound and liquorice.

Ford's Laudanum. A tincture of opium containing cinnamon and cloves.

Fothergill's (Dr.) Pill. Diaphoretic antimony, aloes, scammony, and extract of colocynth.

Frank's Solution. See Solution of Copaiva.

Freeman's Bathing Spirits. Mix water and rectified spirit, of each 3 gallons; dissolve in them soft soap 6 ℔, and camphor 8 oz.; add Daffy's elixir, 8 oz.

Fryar's Balsam. Compound tincture of benzoin, L. P.

GELEE POUR LE GOITRE. Dissolve 1 oz. of white soap in 2½ oz. of proof spirit by a gentle heat; and add to it while still warm, a warm solution of 5 dr. of iodide of potassium in 2½ oz. of proof spirit. A few drops of any fragrant and essential oil may be added.

GINGERBREAD, PURGATIVE. Flour 14 oz., butter 4 oz., treacle 8 oz., p. ginger 1¼ oz., jalap 2 oz., caraway ½ oz. Mix the powders, then add the butter, and lastly the treacle, previously warmed. Roll out, and divide into cakes of ¼ oz. each, containing each 6 or 7 grains of jalap.

GLYCERINE JELLY. Used as an application to chaps and roughened parts of the skin. It may be made of pure glycerine thickened with tragacanth powder and scented with otto of roses. An imitation may be prepared in the following manner (Pharm. Journal): Mix good soft soap ½ dr. intimately with purified honey 2 dr., gradually add pale olive oil 5 oz., stirring without intermission until all is taken up. Care must be taken not to mix in the oil too fast. Finally perfume as desired.

GLYCERINE PASTE. A stiff glutinous compound, recommended by Dr. TILT as a basis of plaster. It is made by boiling 100 or 150 grains of common starch in 1 oz. of glycerine.

GODBOLD'S VEGETABLE BALSAM. An acidulated syrup, or oxymel, of various herbs. The following is an imitation: Dissolve by heat 1 ℔ of lump sugar in white wine vinegar 1 quart, in which 3 oz. of garlic have been steeped for 3 days; add tincture of Tolu 2 dr.

GODFREY'S CORDIAL. The active ingredient is opium, and there is a great diversity in the strength of the compound as prepared by different makers. Many accidents have arisen from its too general use as a stupefactive for infants; but we believe its sale is now less encouraged by druggists than formerly. The following are some of the more usual formulæ.

1. Heat together 7 ℔ (avoird.) of treacle, 8 ℔ of water till united; when nearly cold add the following: rectified spirit 3 fluid ounces, oil of sassafras 40 minims, oil of aniseed 10 drops, laudanum 4 oz. Mix and make up the weight if necessary to 15 ℔. It contains rather more than 9 minims (equal according to some authorities, to 16 or 18 drops), of laudanum in each fluid ounce.

2. Treacle 3½ lb, water 6 lb, spirit of wine 8 fluid ounces, laudanum 4 fluid ounces, oils of aniseed, sassafras, and caraway, of each ½ dr. Mix. Contains 12 or 14 minims of laudanum in an ounce.

3. Sliced sassafras 2 oz., opium cut small 1 oz., bruised aniseed 8 oz., boiling water a gallon. Infuse, strain, and make the infusion into a syrup with 14 lb of treacle. If the whole of the active principles of the opium are extracted, this is much stronger than the preceding.

4. Make a syrup with 3 lb (avoird.) each of treacle and coarse sugar, and water sufficient to make up a gallon. Dissolve 24 drops of oil of sassafras, and 16 of oil of aniseed, in 3 fluid ounces of spirit of wine; and 10 fluid drachms of tincture of opium, and mix the whole with 8 pints, o. m., of the syrup. This is weaker than either of the preceding, containing only five minims of laudanum in a fluid ounce, or 1 drop in a dr.

5. The Philadelphia College of Pharmacy, to prevent the mischief arising from the different strength of this compound, directs it to be prepared as follows: Dissolve 2½ oz. of carbonate of potash in 26 pints of water, add 16 pints of treacle; heat together over a gentle fire till they simmer, remove the scum, and when sufficiently cool, add ½ oz. of oil of sassafras dissolved in 2 pints of rectified spirit, and 24 fluid ounces of tincture of opium, previously mixed. The old wine measure is here intended. It contains about 16 minims of laudanum, or rather more than one grain of opium in each fluid ounce.

6. Sassafras 9 oz., seeds of coriander, caraway, and anise, of each 1 oz.; infuse in 6 pints of water, simmer the mixture till reduced to 4 pints; then add 6 lb of treacle, boil the whole for a few minutes, and when cold add 3 fluid ounces of tincture of opium. Nearly the strength of No. 1. —Dr. Paris.

[These forms are inserted rather with a view to show the dangerous nature of this compound than to encourage its use. No terms are sufficiently strong to express the culpability of those who would place in the hands of ignorant persons, for administration to *infants* and *children*, compounds containing opium.]

Godfrey's Smelling Salts. Sesquicarbonate of ammonia resublimed with pearlash, and a little spirit.—Dr. Paris.

Golden Spirit of Scurvy Grass. It is said to be colored with gamboge.

GOLDEN OINTMENT. Singleton's ointment q. v. The ointment of nitric oxide of mercury is also called golden ointment.

GOULARD'S EXTRACT OF LEAD. Liquor plumbi diacetatis, P. L.

GOUT PAPER. See Charta Antirheumatica, P. F.

GRANVILLE'S (Dr.) COUNTER-IRRITANT LOTIONS. These consist of very strong water of ammonia (sp. gr. 872, being more than three times the strength of the Liquor Ammoniæ of the Pharmacopœia), of spirit of rosemary (fresh tops of rosemary) 2 ℔, alcohol 8 pints; infuse 24 hours, and distil 7 pints, and spirit of camphor, containing 4 oz. camphor in 2 pints of alcohol. The *milder* consists of 4 dr. of the above ammonia, 3 of spirit of rosemary, and 1 of spirit of camphor. The *stronger*, of 5 dr. of ammonia, 2 of spirit of rosemary, and 1 of camphor. The milder is generally sufficient to produce full vesication in from 3 to 10 minutes. The stronger is seldom used except in apoplexy, and to produce cauterization.

GRAVES' (Dr.) GOUT PREVENTIVE. Orange-peel 2 oz., rhubarb 1 oz., hiera picra 2 oz., brandy a quart. Digest for a week.

GREGORY'S POWDER. Calcined magnesia 2½ oz., powdered Turkey rhubarb 1 oz., powdered ginger ½ oz. Mix. The above is Dr. GREGORY'S formula. Some recipes add powdered chamomile. Rhubarb 1 oz., ginger ¼ oz., p. chamomile ½ oz., magnesia 2 oz. Mix. Some druggists prepare it with the heavy carbonate of magnesia, instead of the calcined.

GREENOUGH'S TINCTURE. See TOOTH COSMETICS.

GRIFFIN'S TINCTURE, for coughs. Oil of caraway and anise, each 2 dr., saffron ½ oz., benzoic acid ¾ oz., opium 5 dr., camphor ½ oz., spirit 6 oz., honey 6 oz. When mixed and dissolved, color with burnt sugar.

GRIFFITH'S MIXTURE. This is merely Mistura Ferri Composita of the Pharmacopœia.

GRINROD'S (Dr.) REMEDY FOR SPASMS. Sulphuric ether, aromatic spirit of ammonia, of each ½ oz.; acetate of morphia ½ gr., camphor mixture 2 oz. Mix. A teaspoonful in a little water when required.

GULSTONIAN EMBROCATION. Oil of turpentine 1½ oz., olive oil 1½ oz., dilute sulphuric acid 3 fluid drachms.—Dr. PARIS.

GUTHRIE'S BLACK OINTMENT. 10 grains of nitrate of silver with 1 dr. of spermaceti ointment and 10 drops of a solution of acetate of lead.

GUTHRIE'S EYE OINTMENT. A weaker preparation of the same kind; used in ophthalmia, but causes great pain. See Ung. Hyd. Nit., Pocket Formulary.

HALFORD'S (Sir H.) GOUT PILLS. Acetic extract of colchicum 2½ gr., Dover's powder 1½ gr., compound extract of colocynth 1½ gr., in each pill. One for a dose.

HARROWGATE SALTS (Dr. DUFFIN'S). Sulphate of magnesia 2 dr., bitartrate of potash 10 gr., sal polychrest (potassæ sulphas cum sulphure, Ph. Ed.) ½ dr.; in a pint of warm water. For another formula see *Mineral Waters (factitious) and salts for producing them*, p. 205.

HELVETIUS' STYPTIC. Melt together equal parts of alum and dragon's blood; when cold, powder the compound.

HENRY'S MAGNESIA. A solution of Epsom salts is precipitated by one of carbonate of potash in the cold; the precipitate is well washed, rose-water being used for the last washing; it is then made up while drying into large or small cubes. See Magnes. Carbonas, P. F.

HILL'S BALSAM OF HONEY. Balsam of Tolu 2 oz., styrax 2 dr., opium ½ dr., honey 8 oz., spirit of wine 32 fluid ounces.

HOFFMAN'S PILLS contain corrosive sublimate, about ⅛th of a grain in each.

HOULTON'S LAUDANUM. Opium 2½ oz., distilled vinegar 32 fluid oz.; macerate 6 days with a gentle heat and filter. Evaporate to an extract. Dissolve this in 5 fluid oz. of rectified spirit, and 35 fluid oz. of distilled water.

HOOPER'S FEMALE PILLS. These, according to Dr. PARIS, consist of RUFUS'S Pill, sulphate of iron, canella, and a portion of ivory-black. Mr. GRAY gives two formulæ:

1. Sulphate of iron 8 oz., water 8 oz.; dissolve, and add Barbadoes aloes 40 oz., canella 6 oz., myrrh 2 oz., opoponax ½ oz. Make a mass.

2. Sulphate of iron 2 oz., powder of aloes with canella

16 oz., mucilage of tragacanth and tincture of aloes q. s. to form a mass. Divide 60 grains into 18 pills. But, according to a recent analysis, the iron is in a peroxidized state; probably the sulphate is partially calcined.

The Philadelphia College of Pharmacy gives the following formula:

3. Barbadoes aloes 8 oz., dried sulphate of iron 2 oz. 1½ dr., extract of black hellebore 2 oz., myrrh and soap each 2 oz., canella 1 oz., ginger 1 oz., water q. s. to form a mass. Divide into pills of 2½ gr. each.

HOOPING COUGH; popular remedies for. 1. Cochineal and salt of tartar mixture. This appears to have been first introduced by Dr. LOBB, in 1765, and is still a favorite domestic remedy. Salt of wormwood (subcarbonate of potash) 20 gr., powdered cochineal 10 gr., hot water ¼ of a pint; triturate together, strain, and sweeten with white sugar (or sugar candy). Dose, a teaspoonful to a tablespoonful, according to the age.

2. *Fumigating Powders.* Styrax calamita and gum benzoin, of each a scruple, placed on hot cinders or a heated shovel, in the patient's room, every day.

HUXHAM'S TINCTURE OF BARK. The Compound Tincture of Bark of the London Pharmacopœia is precisely that of HUXHAM, except that he used brandy instead of proof spirit.

ISSUE PEAS. Those in general use are unripe oranges (orange berries) turned in a lathe. The unturned berries are also used. Peas are also turned from orris root. NIEMANN gives the following composition for issue peas: Yellow wax 1½ oz., powdered turmeric 1 oz., powdered orris 1½ oz., Venice turpentine q. s. These are more stimulating, and are used to increase the discharge. The following, according to Dr. GRAY, will open an issue itself: yellow wax 6 oz., verdigris 2 oz., white hellebore 2 oz., cantharides 1 oz., orris 1½ oz., Venice turpentine q. s.

JAMES'S POWDER. It is not known in what respect the mode of preparing this powder differs from the Pharmacopœia process for antimonial powder. Dr. JAMES'S specification is vague and impracticable.

JAMES'S ANALEPTIC PILLS. Equal parts of JAMES'S powder, RUFUS'S pill, and gum guaiacum, made into pills with tinc-

ture of castor. Dr. PARIS has ammoniacum in the place of guaiacum. Another formula is: compound powder of aloes, aloes and myrrh pill, and JAMES'S powder, in equal quantities, formed into pills with tincture of castor and syrup.

JARAVE SPANISH. Pour 4 gallons of boiling water on 2℔ Rio Negro sarsaparilla, 8 oz. powdered guaiacum bark, 4 oz. each of rasped guaiacum wood, anise seed, and liquorice root, 2 oz. of bark of mezereon root, 2 ℔ of treacle, and 12 bruised cloves. Shake it thrice a day, and keep it in a warm place. When fermentation has set in, it is fit for use. Dose, a small tumblerful.

JESUITS' DROPS. WALKER'S. Balsam of copaiva 6 oz., gum guaiacum 1 oz., Chio turpentine ½ oz., subcarbonate of potash ½ oz., cochineal 1 dr., rectified spirit 1 quart. See also Elixir Antivenereum, Pocket Formulary.

KENNEDY'S CORN PLASTER. Yellow wax 1 ℔, Venice turpentine 2 oz., verdigris 1 oz., melted together, and spread on leather.

KEYSER'S PILLS. See Pilulæ Hydrargyri Acetatis, P. F.

KING'S CORDIAL. Dissolve in ½ pint of proof spirit 1½ dr. each of the oils of caraway and cinnamon; extract the stones from 3 ℔ of black cherries, and mash the fruit in a pan; grate one nutmeg; take 2 quarts of Madeira wine, 2 quarts of brandy, and 1 gallon of syrup; mix all together, and color with red sanders wood.

KIRKLAND'S NEUTRAL OINTMENT AND CERATE. See Unguentum Plumbi Compositum and Ceratum Neutrale, P. F.

KITCHENER'S (Dr.) PERISTALTIC PERSUADERS. Turkey rhubarb in powder 2 dr., oil of caraway 10 drops, simple syrup 1 dr. by weight; mix, and divide into 40 pills. Dose, 2, 3, or more. "From 2 to 4 will generally produce one additional motion within 12 hours. The best time to take them is early in the morning."

LARTIGUE'S GOUT PILLS. Compound extract of colocynth 20 gr., extract of colchicum 60 gr., extract of opium 1 gr.; mix, and divide into 18 pills. Dose, one or more, according to their purgative effect.

LEDOYEN'S DISINFECTING FLUID. It consists of about 20 oz. nitrate of lead in a gallon of water. Its specific gravity should be 1.40.

LEECH-BITES, ASTRINGENT FOR. Dissolve 1 part of crystallized perchloride of iron in six parts of collodion very gradually. A drop or two of the product form an admirable hæmostatic. (French Journal.)

LEE'S WYNDHAM'S PILLS. Gamboge 3 oz., aloes 2 oz., Castile soap 1 oz., nitre ½ oz., extract of cow-parsnip 1 oz. In pills of 5 gr. each. (Amer. Jour. of Pharmacy.)

LEE'S ANTIBILIOUS PILLS. Aloes 12 oz., scammony 6 oz., gamboge 4 oz., jalap 3 oz., calomel 5 oz., soap 1 oz., syrup of buckthorn 1 oz., mucilage 7 oz.; mix and divide into 5-grain pills. (The same.)

LEROY'S PURGATIVE.	1	2	3
Scammony, ...	12 dr.	16 dr.	24 dr.
Vegetable turbith, . . .	6 dr.	8 dr.	12 dr.
Jalap, ...	6 oz.	8 oz.	12 oz.
Brandy, ...	10 pints imperial.		

Digest for 12 hours, strain, and add the following syrup:

Senna, ...	6 oz.	8 oz.	12 oz.
Water, ...	24 oz.	32 oz.	48 oz.

Infuse, strain with pressure, and add—

Brown sugar, ...	32 oz.	36 oz.	48 oz.

Make a syrup.

No. 4 is stronger than the above.

LEJEUNE'S BALSAM FOR CHILBLAINS. Camphor 1 dr., tincture of benzoin 5 dr.; dissolve, and add iodide of potassium 5 dr., extract of lead 10 gr., spirit of wine reduced to proof with rose-water 2½ oz.; dissolve 10 dr. of white soap in 2½ oz. of the same diluted spirit by a gentle heat, mix the solutions whilst still warm, and add any perfume. Let it cool in wide-mouthed bottles, and cork.

LIEBERT'S COSMETIC. For Chapped Nipples. Dissolve 10 grains of nitrate of lead in 1 oz. of water. A pair of fine lead shields accompany the lotion, to be worn after applying it. The nipples must be carefully washed before the child is put to the breast.

LIGNUM'S ANTI-SCORBUTIC DROPS. These contain bichloride of mercury, and should not therefore be used without great caution.

LIQUEUR DOREE. Peruvian bark, bitter orange-peel, and cinnamon, of each 4 dr., saffron 2 dr., brandy 4 quarts, Malaga wine 2 quarts; digest for 4 days, strain, and add

2½ ℔ of sugar. (*Liqueurs*, which are not medicated, but merely alcoholic drams, do not come within the plan of this work.)

Liquid Blister. Powdered cantharides 5 oz., and sulphuric ether 15 ounces.—Toynbee.

Liston's Isinglass Plaster. Soak 1 oz. of isinglass in 2 oz. of water, and dissolve it in 2 oz. of rectified spirit and 1½ oz. of water, by the heat of a water-bath. Brush it over the surface of oiled silk, properly stretched. An improved kind is made by brushing one side of the peritoneal membrane of the cæcum of the ox (prepared in the same manner as gold-beaters' skin) with the same solution, and the other side with drying oil.

Locatelli's Balsam. Melt together 4 oz. of yellow wax, 1 ℔ of common oil, and 1 ℔ of Venice turpentine, placing with them 4 oz. of alkanet root wrapped in a linen bag.

Locock's Wafers. These owe most of their activity to Morphia. They should thus be used with caution.

Long's (St. John) Liniment. See Linimentum Terebinthinæ Aceticum, Pocket Formulary.

Lozenges. See Ching's Lozenges, Dawson's, Darcet's.

The medicated lozenges which are sanctioned by the Pharmacopœias, and employed in practice, will be found under Trochisci and Pasta, in the Pocket Formulary. A few other formulæ are here added:

Absorbent Lozenges. Precipitated chalk 3 oz., heavy carbonate of magnesia 2 oz., nutmeg in fine powder 1 dr., sugar 6 oz., powdered gum 1 oz., water q. s. to form a stiff paste, which divide by a punch into lozenges of the usual size, and dry them gradually in a warm room.

Aperient Lozenges. Calomel 60 gr., pure scammony 80 gr., jalap 40 gr. (or jalapine 4 gr.), ginger 8 gr., cinnamon 4 gr., mucilage of tragacanth q. s. to form a stiff paste; mix the other powders accurately together, then with the sugar, lastly add the mucilage, beat the whole into a uniform mass, and divide it into 40 equal lozenges. Each contains 1½ gr. calomel, 2 of scammony, and 1 of jalap.

Black Currant Paste. Soften 12 ℔ of picked black currants by heating them in a water-bath in a covered earthen vessel,

pulp through a hair sieve, and evaporate to a paste, incorporating with it 1 ℔ powdered sugar; roll it out into a sheet of proper thickness. Mr. BARTLETT gives the following formulæ: 3 ℔ of powdered sugar, 3 ℔ of extract of black currants (the inspissated juice), 1 oz. of tartaric acid, 6 oz. of powdered gum; mixed, rolled out, and cut, when dry, with a large pair of scissors into square pieces.

Black Currant or Ipecacuanha Lozenges. Black currant paste (as above) 8 oz., ipecacuanha 30 gr., tragacanth 90 gr.; in 240 lozenges.

Cough Lozenges (*with Lactucarium*). Powdered lactucarium 2 dr., extract of liquorice root 12 dr., ipecacuanha 30 gr., powdered squill 15 gr., refined sugar 6 oz., mucilage of tragacanth q. s.; mix, and divide into 240 equal lozenges. Each contains ½ gr. lactucarium, ⅛ gr. ipecacuanha, $\frac{1}{16}$ gr. of squill.

For other Cough Lozenges, see Trochisci Anticatarrhales, Glycyrrhizæ, et Opii, Lactucæ, Morphiæ et Ipecac., Opii, Papaveris, Pectorales, Scillæ, Tolutani, &c., Pocket Formulary.

Digestive or Live-long Candy. 1. Powdered rhubarb 60 gr., heavy magnesia 1 oz., bicarbonate of soda 1 dr., finely-powdered ginger 20 gr., cinnamon powder 15 gr., powdered white sugar 2 oz., mucilage of tragacanth q. s.; beat together, and divide into paralellograms of 20 gr. each.

2. *Caraway Candy.* Rhubarb 60 gr., powdered caraways 60 gr., oil of caraway 10 drops, ginger and cinnamon, each 15 gr., magnesia 6 dr., carbonate of soda 1 dr., sugar 2 oz., mucilage q. s.—as the last.

Edinburgh Lozenges. Extract of poppies 2 oz., powdered sugar 8 oz., powdered tragacanth 4 oz., water q. s.

Fruit Lozenges. Black currant paste 8 oz., red currant paste (or the juice evaporated to a paste) 4 oz., syrup of raspberries 4 oz., soften by a gentle heat, and beat in a warm mortar with 2 ℔ of powdered sugar, and 1 dr. of powdered citric acid, and, if required, a little mucilage of gum tragacanth.

Marshmallow Lozenges. Marshmallow root powdered 2 oz., sugar 14 oz., mixed with some mucilage of tragacanth and orange-flower water.

Peppermint Lozenges. Rub together white sugar 6 oz., oil of peppermint 36 drops, and the whites of 2 eggs. Make into lozenges.

LYNCH'S EMBROCATION. Olive oil colored with alkanet, perfumed, and rendered stimulating by essential oils.

MADDEN'S ESSENCE. A strongly-acidulated infusion of roses.

MAHOMED'S ELECTUARY. Grocer's currants 1 oz., powdered senna ½ oz., powdered ginger 30 gr., oil of croton 1 drop, syrup of roses sufficient to make an electuary; two teaspoonfuls every morning.—BATEMAN.

MAHY'S PLASTER (American). Boil 12 oz. of white lead, 32 fluid ounces of olive oil, and a little water, stirring constantly until incorporated. Add yellow wax 4 oz., lead plaster 18 oz., and when these are melted stir in 9 oz. of powdered orris.

MAREDANT'S NORTON'S DROPS. Corrosive sublimate, gentian, ginger, and cochineal.

MARSHALL'S CERATE. Palm oil 5 oz., calomel 1 oz., acetate of lead ½ oz., ointment of nitrate of mercury 2 oz.; mix.—Dr. PARIS.

MARSHALL'S EYE-DROPS. These are said to consist of 2 gr. of nitrate of silver in 1 oz. of decoction of snails.

MARSDEN'S ANTISCORBUTIC DROPS, Morton's, Perry's, Lignum's, and other antiscorbutic drops, contain corrosive sublimate.—Dr. PARIS.

MAGNESIA, FLUID. A solution of carbonate of magnesia in water by means of carbonic acid gas, forced into it by pressure. MURRAY'S and DINNEFORD'S should contain from 12 to 15 gr. of the carbonate in each fluid ounce. See Liquor Magnesiæ Carbonatis, Pocket Formulary.

MATHIEU'S VERMIFUGE. Tin filings 1 oz., fern root ¾ oz., worm seed ½ oz., resinous extract of jalap 1 dr., sulphate of potash 1 dr., honey to form an electuary. A teaspoonful every 3 hours for 2 days; then substitute the following: jalap 2 scruples, sulphate of potash 2 scruples, scammony 1 scruple, gamboge 10 gr.; made into an electuary with honey, and given in the same dose.

MINERAL WATERS (FACTITIOUS), AND SALTS FOR PRODUCING THEM. See below, p. 205.

MONTEIN'S BAREGE BALLS (for Sulphur Baths). Sulphate of lime 8 oz., common salt 2 oz., Flanders glue 1 oz., extract of soapwort 1 oz.; make into 8 balls; to be kept from the air. M. MENIERE recommends: Extract of soapwort ½ oz., water 6 oz., lime in powder 4 oz., sulphur 3 oz., gelatine 1 oz.; dissolve the extract and gelatine in the water, add the lime and sulphur, heat gently, stirring it constantly, till the mass gets detached from the sides of the vessel; then form it into balls of 1½ oz. each.

MORRISON'S PILLS, No. 1, consist of equal parts of aloes and cream of tartar; No. 2 consists of 2 parts of gamboge, 3 of aloes, 1 of colocynth, and 4 of cream of tartar, made into pills with syrup.

MORRISON'S ADHESIVE PASTE, for Ring-worm. See Pasta Adhesiva, Pocket Formulary.

MOSELEY'S PILLS. Turkey rhubarb 60 gr., Jamaica ginger 24 gr., syrup and tincture of rhubarb q. s. to form a mass, to be divided into 24 pills.

MOXON'S EFFERVESCING MAGNESIAN APERIENT. The following have been proposed as imitations:

1. Heavy carbonate of magnesia 2 ℔, bicarbonate of soda 1 ℔, tartaric acid 1½ ℔, refined sugar ½ ℔, essence of lemon 40 minims; the powders to be all separately dried at a moderate temperature.

2. Sulphate of magnesia 1 ℔, bicarbonate of soda 1 ℔, tartaric acid ½ ℔; the ingredients to be well dried separately at a moderate temperature. (Pharmaceutical Journal.)

3. Carbonate of magnesia 1 ℔, sulphate of magnesia 2 ℔, bicarbonate of soda 2 ℔, potassio-tartrate of soda 2 ℔, tartaric acid 2 ℔; to be separately dried, and mixed.—M. DURANDE.

MUNRO'S COUGH MEDICINE. 4 dr. of paregoric with 2 dr. of sulphuric ether, and 2 dr. of tincture of Tolu. Dose, 1 teaspoonful in some warm water.

MURRAY'S (Sir J.) FLUID CAMPHOR. Each ounce contains 3 gr. of camphor and 6 gr. of carbonate of magnesia, dissolved by carbonic acid, by pressure.

MURRAY'S GOUT SPECIFIC. It contains iodide of potassium, sulphate of magnesia, and an aromatic tincture. (Pharmaceutical Journal.)

NEPENTHE. Supposed to be a watery solution of opium, resembling Battley's. See Liquor Opii Sedativus, Pocket Formulary.

NORRIS'S DROPS. A solution of tartarized antimony, with a tincture of some vegetable substances, not ascertained.

NOUFFLEUR'S (Madame) WORM MEDICINE. Powdered fern root 3 dr., to be given in the morning (the patient being prepared by an emollient clyster, and a supper of panada); followed in 2 hours by a bolus of calomel, scammony, and gamboge.

OLLIVIER'S BISCUITS. Beat up the white of 2 eggs with 16 oz. of water, and a solution of 76 gr. of corrosive sublimate; collect, wash, and dry the precipitate, $\frac{1}{7}$th of a gr. of which is contained in each biscuit of 2 dr.

OPODELDOC. Lin Saponis.

ORMSKIRK MEDICINE, *to prevent Hydrophobia.* Elecampane 1 dr., chalk 4 dr., Armenian bole 3 dr., alum 10 gr., oil of aniseed 5 drops.

PALAMOUD. See DIETETIC COMPOUNDS.

PALMER'S AERATED CHALYBEATE. Mix $1\frac{1}{2}$ parts of acetic acid with 40 of water, add 4 of proto-sulphate of iron, and 20 of syrup. Put into 4-ounce bottles, for No. 1 and No. 2 respectively, as much of the above as contains 2 and 4 gr. of sulphate of iron, and fill the bottles with a solution of carbonate of soda, or of potash strongly charged with carbonic acid gas. Tartaric acid may be substituted for acetic.

PAPIER EPISPASTIQUE D'ALBESPEYRES. The *Pommade Epispastique* of the French Codex, spread on waxed paper. See Unguentum Epispasticum, Pocket Formulary.

PAPIER EPISPASTIQUE DE VEE. This is of three strengths, distinguished by the colors white, green, and red. The composition is made by boiling cantharides for an hour with water, and lard, green ointment, or lard colored with alkanet; adding white wax to the strained fats, and spreading on paper, silk, or linen. No. 1 is made with 10 oz. of cantharides to 4 ℔ of lard; No. 2 of 1 ℔ of flies to 8 ℔ of green ointment; and No. 3, of $1\frac{1}{2}$ ℔ to 8 ℔ of colored lard; and to each are added 2 ℔ of white wax.—DORVAULT. See BLISTERING PAPER.

PAPIER FAYARD. *Gout Paper.* Euphorbium 3 dr., cantharides 6 dr., powdered and digested with 4 oz. alcohol; and 3 dr. Venice turpentine added to the strained tincture. Fine paper is dipped into it and dried in the air. MOHR directs 4 dr. of cantharides and 1 dr. euphorbium to be digested in 5 oz. of highly rectified spirit; filter and add 1½ oz. Venice turpentine previously liquefied with 2 oz of resin. To be spread on the paper while warm.

PELLETIER'S ETHEREAL OPODELDOC. See Balsamum Aceticum Camphoratum, Pocket Formulary.

PETER'S PILLS. Aloes, jalap, gamboge, and scammony, of each 2 dr.; calomel 1 dr.

PILES, popular remedies for. Dr. WARDLEWORTH'S PILLS contain 3½ gr. of pitch in each; 2 every night. For ELECTUARIES for piles see Confectio Resinæ, Confectio Sulphuris Composita, Electuarium Hæmorrhoidale, Electuarium Sulphuris Compositum, in Pocket Formulary. See also WARD'S PASTE, below. For PILE OINTMENTS, see Unguentum Gallæ, Unguentum Gallæ Compositum, Unguentum et Opii, Unguentum Hæmorrhoidale, Pocket Formulary. Sir H. HALFORD'S Pile Ointment consists of equal parts of citrine ointment and oil of almonds triturated in a glass mortar till perfectly smooth. Mr. WARD'S is—Powdered nutgall 2 dr., camphor 1 dr., melted wax 1 oz., tincture of opium 2 dr. Mix.

PILLS. See proprietors' names in alphabetical order. A great variety of formulæ for pills of every kind will be found in the Pocket Formulary.

PILLS, TO COAT, WITH GELATINE . }
——, TO SILVER . . . . . } See Pilulæ, Pocket Formulary.

M. DURDEN recommends collodion as a covering for pills: others, a solution of gutta percha in chloroform: but the ready solubility of these materials in the stomach may be questioned. M. BLANCHARD uses balsam of Tolu dissolved in ether. Mr. BAILDON recommends chloroform instead of ether.

PLUNKET'S OINTMENT FOR CANCER. White arsenic, sulphur, powdered flowers of lesser spearwort and stinking chamomile, levigated together, and formed into a paste with white of egg.

POMADE DIVINE. Beef marrow 3 ℔; put it into an earthen

vessel, and cover it with cold water, and change the water daily for a few days, using rose-water the last day. Pour off and press out the water; add to the marrow 4 oz. each of styrax, benzoin, and Chio turpentine, 1 oz. orris powder, ½ oz. each of powdered cinnamon, cloves, and nutmeg. Set the vessel in hot water, and keep the water boiling for 3 hours; then strain. For Pomades for the hair, see HAIR COSMETICS, after PERFUMERY.

PORTLAND'S (DUKE OF) GOUT POWDER. Equal quantities of the roots of gentian and birthwort, tops of germander, ground pine, and lesser centaury: all to be powdered and mixed together.

POWELL'S BALSAM FOR COUGH. Mix together 2 dr. of syrup of Tolu, 1 oz. of paregoric elixir, and 2 oz. of liquorice-juice.

QUEEN OF HUNGARY'S WATER. Tops and flowers of rosemary 2 ℔, rectified spirit 3 ℔; digest in a close vessel for 50 hours in a gentle heat, then distil by water-bath.

QUININE AND CAMPHOR PILLS. See Pilulæ Quinæ et Camphoræ, Pocket Formulary.

RADCLIFFE'S ELIXIR. Aloes 6 dr., cinnamon, zedoary, and cochineal, each ½ dr., rhubarb 1 dr., syrup of buckthorn 2 oz., proof spirit 16 fluid oz., water 5 fluid oz.—Dr. PARIS. According to GRAY, it contains jalap, scammony, and senna.

RASPAIL'S CAMPHOR CIGARETTES. These are merely camphor inclosed in a tube (a quill or paper tube may be used), confined by blotting paper, and used cold. Another kind of camphorated cigars is made by saturating dried coltsfoot or other leaves with a strong solution of camphor, and rolling them in the form of cigars.

REECE'S CHIRAYTA PILLS. Extract of chirayta 2 dr., dried carbonate of soda 1 scruple, p. ginger 15 gr.; mix, and divide into 36 pills. Two twice a day.

REGNAULD'S PECTORAL PASTE. Pectoral flowers (mullein, coltsfoot, catsfoot, and red poppies mixed) 16 oz., boiling water 3 ℔; infuse, strain, and add to the clear liquor 6 ℔ of clean gum Arabic; dissolve by a gentle heat, and evaporate to a proper consistence, adding towards the end 6 dr. of tincture of balsam of Tolu.

REYNOLD'S GOUT SPECIFIC. It is supposed to be a wine of colchicum.

REVALENTA. It is said to be prepared from the seeds of the Ervum lens. See Dietetic Articles.

RIGA BALSAM FOR BRUISES. Mix 4 oz. of spirits of wine with 1 dr. of compound tincture of benzoin, and 2 dr. of tincture of saffron.

ROBINSON'S (Dr.) STIMULATING PURGATIVE PILLS. Watery extract of aloes 1 dr., balsam of Peru 10 gr., oil of caraway 10 drops, scammony ½ dr.; mix and divide into 20 pills: 2 or 3 when required.

ROCHE'S EMBROCATION. Olive oil, with half its weight of oil of cloves and oil of amber.—Dr. PARIS.

ROGE'S MAGNESIAN PURGATIVE. Calcined magnesia 1 oz., carbonate of magnesia ½ oz., citric acid 3¼ oz., sugar, rubbed with a few drops of essence of lemon, 6¼ oz. To form Aerated Magnesian Lemonade, put ¼ of the powder into a soda-water bottle nearly filled with water, and cork it securely.

ROUSSEAU'S DROPS. See Vinum Opii Fermentatione Paratum, P. F.

RUSPINI'S STYPTIC. It contains (according to Dr. A. T. THOMPSON) gallic acid, sulphate of zinc, spirit, and rose-water.

RYAN'S ESSENCE OF COLTSFOOT. Tincture of balsam of Tolu 2 oz., compound tincture of benzoin 2 oz., spirit of wine 4 oz.—GRAY.

RYMER'S TINCTURE. A tincture of capsicum, camphor, cardamom, rhubarb, aloes, and castor, in proof spirit, with a small quantity of sulphuric acid.—Dr. PARIS.

(The inventor states that it is impregnated with an aerial acid.)

SALTS, MINERAL. See WATERS, FACTITIOUS MINERAL, p. 205.

SCOTT'S PILLS. See ANDERSON'S PILLS.

SCOTT'S PLASTER. It appears to be a carefully prepared Emp. Plumbi, spread on calico. If it contains resin, the quantity is probably less than in Emp. Resinæ.

SEIDLITZ POWDERS (in separate powders). One contains 2 dr.

of powdered Rochelle salts, and 40 gr. of bicarbonate of soda; the other powder is p. tartaric acid 35 gr.

SEIDLITZ POWDER, in one bottle. *Note.*—The powders are all to be thoroughly dried separately, at a gentle heat—the potassio-tartrate of soda, at a temperature not exceeding 110° F.; the others not higher than 120°. Take of potassio-tartrate of soda, dried, 15 oz., tartaric acid, dried, 5 oz.; (or citric acid 4¾ oz.), dried bicarbonate of soda 6 oz.; mix, and keep in a well-closed bottle. Dose, 3 dr. Or, mix two parts of bitartrate of soda with one part bicarbonate of soda. Keep dry. The above have no resemblance to the natural water of Seidlitz. See WATERS (Mineral), page 205.

SINGLETON'S GOLDEN OINTMENT. Orpiment mixed with lard to the consistence of an ointment.

SODA POWDERS. These usually contain in one paper 30 gr. of bicarbonate of soda, and in the other 25 gr. of tartaric acid (or 24 of citric acid). For sherbet, lemonade, and ginger-beer powders, see BEVERAGES, in another division of this work.

SMELLOME'S EYE-OINTMENT. Prepared verdigris 30 gr.; levigate with 30 drops of olive oil, and add 1 oz. of resinous cerate.

SPEEDIMAN'S PILLS. Rhubarb, aloes, myrrh, and extract of chamomile, of each 60 gr.; oil of chamomile 12 drops; mix, and divide into 4-grain pills.

SOLOMON'S BALM OF GILEAD. An aromatic tincture, of which cardamoms form a leading ingredient, made with brandy.—Dr. PARIS. It is thought to contain cantharides.

SOLOMON'S ANTI-IMPETIGINES is said to be a solution of corrosive sublimate.

SMITH'S (Dr. HUGH) STOMACHIC PILLS. Aloes, rhubarb, aromatic powder, gum sagapenum, of each 1 dr.; oil of mint and oil of cloves, each 10 drops; balsam of Peru q. s. In 5-grain pills; 2 to 4 every night.

SOLUTION OF COPAIVA. Mix 2 parts of balsam of copaiva with 3 of liquor potassæ and 7 of water. Boil them for a quarter of an hour, and when a little cooled, add 1 part of spirit of nitric ether. Let the mixture rest for a few hours, then draw off the clear liquor from the lower part of the

vessel. This is supposed to resemble FRANK'S Solution. Or copaiva may be boiled in water with an equal weight of carbonate of potash. (For other preparations of copaiva, see Gelatina Copaibæ, Electuarium Copaibæ, Elec. Cubebæ et Copaibæ, Syrupus Copaibæ, Mistura Copaibæ, &c., in P. F.)

SPILSBURY'S ANTISCORBUTIC DROPS. Corrosive sublimate 2 dr. (not 2 *oz.*, as misprinted in the eighth edition of Dr. PARIS'S Pharmacologia), precipitated sulphuret of antimony 1 dr., gentian 2 dr., orange-peel 2 dr., red sanders 1 dr., proof spirit 16 fluid oz.; digest and strain.—Dr. PARIS. But we are informed that this is incorrect. Another formula is: Levigated crocus metallorum 18 dr., corrosive sublimate 135 gr., red sanders 1½ dr., gentian 6 dr., orange-peel 6 dr., brandy 48 fluid oz.; digest for 10 days, shaking frequently, and strain; dose 5 to 60 drops.

SQUIRE'S ELIXIR. Opium 1 oz., camphor 1 oz., spirit of aniseed (compound) 4 pints, tincture of serpentaria 1 pint, water 4 pints, tincture of ginger ½ oz. Some recipes add a little aurum musivum.

STANDERT'S RED MIXTURE. Carbonate of magnesia 4 dr. powdered rhubarb 2 dr., tincture of rhubarb 1½ oz., tincture of opium 1 dr., oil of aniseed 24 drops, essence of peppermint 30 drops, water 1½ pint; mix. A popular remedy for bowel complaints in the West of England.

STANDERT'S STOMACHIC CANDY. Cardamom seed, ginger, rhubarb (all in fine powder), each 4 dr., lump sugar 4 oz., water 6 dr.; boil together, stirring constantly till the sugar is dissolved, then pour it into a proper mould.

STEER'S OPODELDOC. 1. Rectified spirit a quart, Castile soap 5 oz., camphor 2½ oz., oil of rosemary 2½ dr., oil of origanum 5 dr., weaker ammonia 4 oz.; digest till dissolved, and pour while warm into wide-mouthed bottles.

2. Rectified spirits 8 pints o. m., white soap 20 oz., camphor 8 oz., water of ammonia 4 oz., oil of rosemary 1 oz., oil of horsemint 1 oz.; dissolve the soap in the spirit by the gentle heat, and add the other ingredients.—Bottle whilst warm.—PHIL. COL. OF PHARMACY.

STOUGHTON'S ELIXIR. 1. Gentian 36 oz., serpentary 16 oz., dried orange-peel 24 oz., calamus aromaticus 4 oz., rectified spirit and water, of each 6 gallons, old measure.

2. Gentian 4 ℔, orange-peel 2 ℔, cochineal 2 dr., cardamom seed 1 oz., rectified spirits 8 gallons.

STOREY'S WORM CAKES. Calomel 1 scruple, jalap 1 dr., ginger 2 scruples, sugar 1 oz., cinnabar to color, syrup q. s. to form 10 cakes.

STRUVE'S LOTION FOR HOOPING-COUGH. Emetic tartar 60 gr., water 2 oz., tincture of cantharides 1 oz.

SWAIM'S VERMIFUGE. Wormseed 2 oz., valerian, rhubarb, pink root, white agaric, of each 1½ oz.; boil in sufficient water to yield 3 quarts of decoction, and add to it 30 drops of oil of tansy, and 45 drops of oil of cloves, dissolved in a quart of rectified spirit. (American remedy.)

SYDENHAM'S LIQUID LAUDANUM. See Vinum Opii, Pocket Formulary.

TANJORE PILLS. See Pilulæ Arsenici, Pocket Formulary.

THIBAUT'S BALSAM for wounds. Digest flowers of St. John's wort, one handful, in ½ pint rectified spirit, then express the liquor, and dissolve in it myrrh, aloes, and dragon's blood, of each 1 dr., with Canada balsam ½ oz.

TINCTURE OF QUININE (AROMATIC). See Tinct. Quinæ Co., and Tinct. Quinæ Sulphatis Acida, Pocket Formulary.

TISSOT'S PURGATIVE POWDERS. Jalap, rhubarb, senna, and soluble cream of tartar, equal quantities. Dose, 2 to 6 drachms. Used in Normandy.

TURLINGTON'S BALSAM. Rectified spirits 8 old wine pints, benzoin 12 oz., liquid styrax 4 oz., socotrine aloes 1 oz., balsam of Peru 2 oz., myrrh 1 oz., angelica root ½ oz., balsam of Tolu 4 oz., extract of liquorice 4 oz.; digest 10 days and strain.—PHIL. COL. OF PHARM. The certified copy of the original recipe is more complex, containing three times as many ingredients.

VALANGIN'S SOLUTION OF SOLVENT MINERAL. Arsenious acid (which has been mixed with muriate of soda, and resublimed) 30 gr., hydrochloric acid 90 gr., distilled water 1 oz.; dissolve, and add distilled water to make up 30 fluid oz. Dose, from 3 drops, increased very gradually to 10. See Liquor Arsenici Chloridi, P. F.

VENLO'S VEGETABLE SYRUP. It is supposed to be a decoction of burdock, mint, dandelion, senna, &c., boiled with sugar, and a small portion of solution of sublimate added.

WALKER'S JESUIT'S DROPS. See Jesuit's Drops, above.

WARBURG'S FEVER TINCTURE. M. FUCHS gives the following recipe for imitating this secret remedy: Hepatic aloes and zedoary root, of each 1 dr., angelica root and camphor, of each 2 gr., saffron 3 gr., proof spirits 3 oz. In 25 dr. of the filtered tincture dissolve 30 gr. of sulphate of quinine. Put it up in 5-drachm bottles, containing a dose. Sold at about 5 shillings each bottle.

WARD'S PASTE. The same as Confectio Piperis Nigri of the London Pharmacopœia.

WARD'S WHITE DROPS. To 16 oz. of strong nitric acid add gradually 7 oz. of subcarbonate of ammonia; let it stand 2 or 3 hours; then put it into a bolt-head which it will only half fill, and to each 16 oz. put 4 oz. of pure quicksilver, and digest in a sand-heat till the solution is complete; then gently increase the heat and add a little more quicksilver at intervals till it will dissolve no more; then evaporate it in a glass or earthen dish placed in sand, till a pellicle appears, and set it aside to crystallize. Dissolve 1 ℔ of the drained salt in 3 ℔ of rose-water by the heat of a sand-bath.

WARD'S ESSENCE FOR THE HEADACHE. Spirit of wine 2 ℔, roche alum in fine powder 2 oz., camphor 4 oz., essence of lemon ½ oz., strong water of ammonia 4 oz.; stop the bottle close, and shake it daily for 3 or 4 days.

WARD'S RED PILL. Glass of antimony levigated with a fourth of its weight of dragon's blood, made into a mass with wine, and divided into pills of a grain and a half each; one pill is a dose, on an empty stomach. In foulness of the stomach and bowels, and obstinate rheumatic disorders.

WARD'S DROPSY PURGING POWDER. Jalap 1 ℔, cream of tartar 1 ℔, red bole 1 oz.; mix. Dose, from 30 to 40 gr., in broth or warm beer, repeated for 2 or 3 days, or oftener if necessary.

WARD'S SWEATING POWDER. Similar to Dover's Powder.

WARD'S WASHING POWDER. See Washing Compounds, Trade Chemicals.

WARNER'S CORDIAL. Rhubarb 1 oz., senna ½ oz., saffron 1 dr., liquorice ½ oz., raisins 1 ℔, brandy 3 pints; digest for a week, and strain.

WARWICK'S (Countess of) POWDERS. Scammony 2 oz., calx of antimony 1 oz., cream of tartar ½ oz.; mix.

WEBSTER'S DIET DRINK. A decoction of syrup of sarsaparilla, betony, dulcamara, guaiacum, liquorice, sassafras, turmeric, and thyme.

WEBSTER'S (Lady) PILLS. See Pilulæ Aloes cum Mastiche, P. F.

WHITEHEAD'S ESSENCE OF MUSTARD. See ESSENCE OF MUSTARD for an imitation of it.

WHITEHEAD'S MUSTARD PILLS. Dr. PARIS says they consist of balsam of Tolu and resin.

WHITELAW'S ETHEREAL TINCTURE OF LOBELIA. See Tinctura Lobeliæ Etherea, Pocket Formulary.

WILSON'S GOUT TINCTURE. A vinous infusion of colchicum.

WISDOM'S (Dr.) EYE-WATER. Bole 2 oz., sulphate of zinc ¼ oz., camphor (dissolved in 1½ oz. of rectified spirit) ¼ oz., water a gallon.

WRIGHT'S PEARL OINTMENT. White precipitate 8 oz., extract of lead a pint; rub together and add 7 ℔ of white wax melted with 16 ℔ of olive oil.—PHARM. JOURNAL.

WORM LOZENGES. See CHING'S LOZENGES, and STORY'S WORM CAKES, above; see also Trochisci Anthelmintici, and Trochisci Santoninæ, Pocket Formulary.

YOUNG'S PURGING DRINK. Carbonate of soda in crystals 2½ dr., cream of tartar in crystals 3 dr., water 8 oz.; put it into a stone bottle and secure the cork.

FACTITIOUS MINERAL WATERS

AND

Salts for Producing Them.

AERATED OR CARBONATED WATERS.

These require the aid of the powerful machine employed by soda-water manufacturers, to charge the waters strongly with carbonic acid gas. The gas is made from whiting and diluted sulphuric acid, and is forced by a pump into the watery solution. Sometimes the gas is produced by the mutual action of the ingredients introduced into the bottle of water, which must be instantly closed; but this method is found practically inconvenient, and is only adopted in the absence of proper apparatus. The quantity of gas introduced is directed, in the French and American Pharmacopœias, in most cases, to be 5 times the volume of liquid. For chalybeate and sulphuretted waters, the water should be previously deprived of the air it naturally contains, by boiling it, and allowing it to cool in a closed vessel.

Bakewell's apparatus will be found very convenient for making small quantities of aerated waters; and the syphon bottles suitable for holding them.

Simple Aerated Water. Carbonic acid gas water. Water charged with five or more volumes of carbonic acid gas, as above.

Alkaline Aerated Waters. Aerated soda and potash waters should be made by dissolving 1 dr. of the carbonated alkali in each pint of water, and charging it strongly with carbonic acid gas. But the soda water of the shops generally contains but little (or no) soda.

Aerated Magnesia Water. This is made of various strengths.

Murray's and Dinneford's Fluid Magnesia may be thus made: To a boiling solution of 16 oz. of sulphate of magnesia in 6 pints of water, add a solution of 19 oz. of crystallized carbonate of soda in the same quantity of water; boil the mixture till gas ceases to escape, stirring constantly; then set it aside to settle; pour off the liquid, and wash the precipitate on a cotton or linen cloth, with warm water, till the latter passes tasteless. Mix the precipitate, without drying it, with a gallon of water, and force carbonic acid gas into it under strong pressure, till a complete solution is effected. The *Eau Magnésienne* of the French Codex is about a third of this strength; and we have met with some prepared in this country not much stronger.

Carbonated Lime-water. *Carrara Water.* Lime-water (prepared from lime made by calcining Carrara marble) is supersaturated, by strong pressure, with carbonic acid; so that the carbonate of lime at first thrown down is redissolved. It contains 8 gr. of carbonate of lime in 10 fluid oz. of water.

Aerated Lithia Water. This may be conveniently made from the fresh precipitated carbonate, dissolved in carbonated water, as directed for fluid magnesia. Its antacid and antilithic properties promise to be useful. (See page 211.)

SALINE CARBONATED WATERS.

The following afford approximate imitations of these waters. The earthly salts, with the salts of iron, should be dissolved together in the smaliest quantity of water. The other ingredients to be dissolved in the larger portion of the water, and the solution impregnated with the gas. The first solution may be then added, or be previously introduced into the bottles. The salts, unless otherwise stated, are to be crystallized.

Baden Water. Muriate of magnesia 2 gr., muriate of lime 40 gr., muriate of iron $\frac{1}{4}$ gr. (or 3 minims of the tincture), muriate of soda 30 gr., sulphate of soda 10 gr., carbonate of soda 1 gr., water 1 pint, carbonic acid gas 5 volumes.

Carlsbad Water. Muriate of lime 8 gr., tincture of mu-

riate of iron 1 drop, sulphate of soda 50 gr., carbonate of soda 60 gr., muriate of soda 8 gr., carbonated water 1 pint.

EGER. Carbonate of soda 5 gr., sulphate of soda 4 scruples, muriate of soda 10 gr., sulphate of magnesia 3 gr., muriate of lime 5 gr., carbonated water a pint. (Or it may be made without apparatus thus: Bicarbonate of soda 30 gr., muriate of soda 8 gr., sulphate of magnesia 3 gr., water a pint; dissolve and add a scruple of dry bisulphate of soda, and close the bottle immediately.)

EMS. Carbonate of soda 2 scruples, sulphate of potash 1 gr., sulphate of magnesia 5 gr., muriate of soda 10 gr., muriate of lime 3 gr., carbonated water a pint.

MARIENBAD. Carbonate of soda 2 scruples, sulphate of soda 96 gr., sulphate of magnesia 8 gr., muriate of soda 15 gr., muriate of lime 10 gr., carbonated water a pint. (Or, bicarbonate of soda 50 gr., sulphate of soda 1 dr., muriate of soda 15 gr., sulphate of magnesia 10 gr.; dissolved in a pint of water, add 25 gr. of dry bisulphate of soda, and cork immediately.)

MARIENBAD PURGING SALTS. Bicarbonate of soda 5 oz., dried sulphate of soda 12 oz., dry muriate of soda 1½ oz., sulphate of magnesia, dried, 2 oz., dried bisulphate of soda 2½ oz. Mix the salts previously dried, separately, and keep them carefully from the air.

PULLNA WATER. Sulphate of soda 4 dr., sulphate of magnesia 4 dr., muriate of lime 15 gr., muriate of magnesia (dry) a scruple, muriate of soda a scruple, bicarbonate of soda 10 gr., water slightly carbonated a pint. One of the most active of the purgative saline waters.

PULLNA WATER, WITHOUT THE MACHINE. Bicarbonate of soda 50 gr., sulphate of magnesia 4 dr., sulphate of soda 3 dr., muriate of soda a scruple; dissolve in a pint of water; add, lastly, 2 scruples of bisulphate of soda, and close the bottle immediately.

SALTS FOR MAKING PULLNA WATER. Dry bicarbonate of soda 1 oz., exsiccated sulphate of soda 2 oz., exsiccated sulphate of magnesia 1½ oz., dry muriate of soda 2 dr., dry tartaric acid ¾ oz. (or rather, dry bisulphate of soda 1 oz.)

SEIDLITZ WATER. This is usually imitated by strongly aerat-

ing a solution of 2 dr. of sulphate of magnesia in a pint of water. It is also made with 4, 6, and 8 dr. of the salts to a pint of water.

SEIDLITZ POWDER. The common Seidlitz powders (as given p. 199) do not resemble the water. A closer imitation would be made by using effloresced sulphate of magnesia instead of the potassio-tartrate of soda. A still more exact compound will be the following: Effloresced sulphate of magnesia 2 oz., bicarbonate of soda ½ oz., dry bisulphate of soda ½ oz.; mix, and keep in a close bottle.

SEIDSCHUTZ WATER. Sulphate of magnesia 3 dr., muriate of lime, nitrate of lime, bicarbonate of soda, of each 8 gr., sulphate of potash 5 gr., aerated water 1 pint.

SELTZER WATER. Muriate of lime and muriate of magnesia, of each 4 gr.; dissolve these in a small quantity of water, and add it to a similar solution of 8 gr., of bicarbonate of soda, 20 gr. muriate of soda, and 2 gr. of phosphate of soda; mix, and add a solution of ¼ of a gr. of sulphate of iron; put the mixed solution into a 20-oz. bottle, and fill up with aerated water. But much of the Seltzer water sold is said to be nothing more than simple carbonated water. An imitation of Seltzer water is also made by putting into a stone Seltzer bottle, filled with water, 2 dr. bicarbonate of soda, and 2 dr. of citric acid in crystals, corking the bottle immediately. Sodiac powders are sometimes sold as Seltzer powders.

VICHY WATER. Bicarbonate of soda 1 dr., muriate of soda 2 gr., sulphate of soda 8 gr., sulphate of magnesia 3 gr., tincture of muriate of iron 2 drops, aerated water 1 pint. DORVAULT directs 75 grains of bicarbonate of soda, 4 grains of chloride of sodium, ⅕ gr. sulphate of iron, 10 gr. sulphate of soda, 3 grains sulphate of magnesia, to a pint of water. By adding 45 grains (or less) of citric acid an effervescing water is obtained.

M. SOUBEIRAN, relying on the analysis of LONGCHAMPS, imitates Vichy water by the following combination: Bicarbonate of soda 135 grains, chloride of sodium 2½ gr., cryst. chloride of calcium 12 gr., sulphate of soda 11½ gr., sulphate of magnesia 3¾ gr., tartrate of iron and potash ⅛ gr., water $2\frac{1}{10}$ pints (1 litre), carbonic acid 305 cub. inches (5 litres). Dissolve the salts of soda and iron in part of the

water, dissolve and add the sulph. magnes., and then the chlor. calc. in the remaining water. Charge now with the carbonic acid gas under pressure.

VICHY SALTS. Bicarbonate of soda 1½ oz., muriate of soda 15 gr., effloresced sulphate of soda 1 dr., effloresced sulphate of magnesia 1 scruple, dry tartarized sulphate of iron 1 gr., dry tartaric acid 1 oz. (or dry bisulphate of soda); mix the powders, previously dried, and keep them in a close bottle.

SALINE WATERS, &c., NOT CARBONATED.

SEA-WATER. Muriate of soda 4 oz., sulphate of soda 2 oz., muriate of lime ¼ oz., muriate of magnesia 1 oz., iodide of potassium 4 gr., bromide of potassium 2 gr., water a gallon. A common substitute for sea-water as a bath is made by dissolving 5 or 6 oz. of common salt in a gallon of water.

The following mixture of dry salts may be kept for the immediate production of a good imitation of sea-water. Muriate of soda (that obtained from evaporating sea-water and not recrystallized, in preference) 85 oz.; effloresced sulphate of soda 15 oz., dry muriate of lime 4 oz., dry muriate of magnesia 16 oz., iodide of potassium 2 dr., bromide of potassium 1 gr. Mix and keep dry. Put 5 or 6 oz. to a gallon of water.

BALARUC WATER. Muriate of soda 1 oz., muriate of lime 1 oz., muriate of magnesia ½ oz., sulphate of soda 3 dr., bicarbonate of soda 2 dr., bromide of potassium 1 gr., water a gallon. Chiefly used for baths.

SULPHURETTED WATERS.

SIMPLE SULPHURETTED WATERS. Pass sulphuretted hydrogen into cold water (previously deprived of air by boiling, and cooled in a close vessel), till it ceases to be absorbed.

AIX-LA-CHAPELLE WATER. Bicarbonate of soda 12 gr., muriate of soda 25 gr., muriate of lime 3 gr., sulphate of soda 8 gr., simple sulphuretted water 2½ oz., water slightly carbonated 17½ oz.

BAREGES WATER. (Cauterets, Bagneres de Luchon, Bonnes

St. Sauveur, may be made the same.) Crystallized hydrosulphate of soda (see Sodæ Hydrosulphas, P. F.), crystallized carbonate of soda, and muriate of soda, of each 2½ gr., water (freed from air) a pint. A stronger solution for adding to baths is thus made: Crystallized hydrosulphate of soda, crystallized carbonate of soda, and muriate of soda, of each 2 oz., water 10 oz.: dissolve. To be added to a common bath at the time of using.

NAPLES WATER. Crystallized carbonate of soda 15 gr., fluid magnesia 1 oz., simple sulphuretted water 2 oz., aerated water 16 oz. Introduce the sulphuretted water into the bottle last.

HARROGATE WATER. Muriate of soda 100 gr., muriate of lime 10 gr., muriate of magnesia 6 gr., bicarbonate of soda 2 gr., water 18½ oz. Dissolve and add simple sulphuretted water 1½ oz.

HARROGATE SALTS. See Dr. DUFFIN'S, above.

CHALYBEATE WATERS.

SIMPLE CHALYBEATE WATER. Water freed from air by boiling 1 pint, sulphate of iron ½ gr.

AERATED CHALYBEATE WATER. Sulphate of iron 1 gr., carbonate of soda 4 gr., water deprived of air, and charged with carbonic acid gas, a pint. Dr. PEREIRA recommends 10 gr. each of sulphate of iron and bicarbonate of soda to be taken in a bottle of ordinary soda-water. This is equivalent to 4 gr. of carbonate of iron.

BRIGHTON CHALYBEATE. Sulphate of iron, muriate of soda, muriate of lime, of each 2 gr., carbonate of soda 3 gr., carbonated water 1 pint.

BUSSANG, FORGES, PROVINS, and other similar waters, may be imitated by dissolving from ½ to ⅔ds of a grain of sulphate of iron, 2 or 3 gr. of carbonate of soda, 1 gr. of sulphate of magnesia, and 1 of muriate of soda, in a pint of aerated water.

MONT D'OR WATER. Bicarbonate of soda 70 gr., sulphate of iron ⅔ gr., muriate of soda 12 gr., sulphate of soda ½ gr., muriate of lime 4 gr., muriate of magnesia 2 gr., aerated water a pint.

PASSY WATER. Sulphate of iron 2 gr., muriate of soda 3 gr., carbonate of soda 4 gr., muriate of magnesia 2 gr., aerated water a pint.

PYRMONT WATER. Sulphate of magnesia 20 gr., muriate of magnesia 4 gr., muriate of soda 2 gr., bicarbonate of soda 16 gr., sulphate of iron 2 gr., Carrara water a pint.

The mineral waters prepared by Messrs. Sturve, of Brighton, are stated to be the very exact imitations of the natural springs.

VARIOUS AERATED MEDICINAL WATERS NOT RESEMBLING ANY NATURAL SPRING.

MIALHE'S AERATED CHALYBEATE WATER. Water a pint, citric acid 1 dr., citrate of iron 15 gr.; dissolve, and add 75 gr. of bicarbonate of soda.

TROUSSEAU'S MARTIAL AERATED WATER. Potassio-tartrate of iron 10 grains, artificial Seltzer water a pint.

BOUCHARDAT'S GASEOUS PURGATIVE. Phosphate of soda 1½ oz., carbonated water a pint.

MIALHE'S IODURETTED GASEOUS WATER. Iodide of potassium 15 gr., bicarbonate of soda 75 gr., water a pint; dissolve and add sulphuric acid diluted with its weight of water 75 gr. Cork immediately.

DUPASQUIER'S GASEOUS WATER OF IODIDE OF IRON. Solution of iodide of iron (containing $\frac{1}{10}$ of dry iodide) 30 gr., syrnp of gum 2½ oz., aerated water 17½ oz.

[See also Magnesia, Carrara, and Lithia Waters, page 206; Bewley's Chalybeate Water, page 171. Also Aqua Benzoata Aerata, and Aqua Magnesiæ Citratis, P. F.]

PERFUMERY.

DISTILLED WATERS.

THE simple distilled waters (without spirit) used in perfumery are chiefly those of rose, elder, and orange flower, cinnamon, &c. The points requisite to be attended to are, that the flowers be fresh, gathered after the sun has risen and the dew exhaled, and that sufficient water be used to prevent the flowers being burned, but not much more than is sufficient for this purpose. The quantities usually directed are—Roses 15 lb, water 40 lb: distil 15 lb for *single*, and the same water with 15 lb of fresh roses for *double* rose-water.

Orange-flowers 12 lb, water 36 lb: distil 24 lb for *double* orange-flower water; this with an equal quantity of distilled water forms the *single*. The flowers should not be put into the still till the water nearly boils.

ELDER-FLOWER WATER, ACACIA-FLOWER WATER, AND BEAN-FLOWER WATER, are prepared in the same manner as rose-water.

CINNAMON WATER. A gallon should be distilled from 20 oz. of fine cinnamon (bruised) and 2 gallons of water.

STRAWBERRY WATER. Bruised strawberries 4 lb, water a gallon; macerate for 12 hours, and distil 6 pints.

The waters prepared without distillation (by diffusing the essential oils through water, after mixing them with chalk, magnesia, or silica, or dissolving them in spirit) are seldom so proper for perfumery purposes as those distilled from the flowers, &c. Rose-water, made from the otto (8 drops of otto, previously mixed with a drachm of precipitated chalk, diffused in a quart of distilled water, and afterwards distilled or simply filtered), is to most persons very agreeable; but that distilled from the flowers should also be kept, as it is by others greatly preferred.

MUSK WATER, VIOLET WATER, JESSAMINE WATER, and some

others, are made by mixing the *spirituous essences* with distilled or pure soft water. A usual proportion is 2 dr. to a pint.

SPIRITUOUS WATERS.

The spirit employed in perfumery should be selected with great care; it should be perfectly free from grain-oil and other impurities. It should be 60° over-proof, unless otherwise directed.

Simple Spirit of Lavender. Lavender-flower (free from stalks) 2 lb, rectified spirit 8 pints, water 16 pints; distil 8 pints.

Smyth's Distilled Essence of Lavender. Essential oil of English lavender 4 oz., rectified spirit (60° over proof) 5 pints, rose-water 1 pint: mix, and distil 5 pints for sale.

Essence of Lavender (by mixture). Essential oil of lavender 3½ oz., rectified spirit 2 quarts, rose-water ½ pint, tincture of orris ½ pint.

Lavender Water. English oil of lavender 4 oz., spirit 3 quarts, rose-water 1 pint: mix and filter. (A commoner and cheaper preparation may be made with the French oil.)

Odoriferous Lavender Water. 1. Rectified spirit 5 gallons, essential oil of lavender 20 oz., oil of bergamot 5 oz., essence of ambergris ½ oz. Sometimes 4 oz. of orris-root are digested with the above.—Mr. Brande.

2. Oil of lavender, oil of bergamot, of each 3 dr.; otto of roses and oil of cloves, of each 6 drops; musk 2 gr., true oil of rosemary 1 dr., honey 1 oz., benzoic acid 2 scruples; rectified spirit a pint, distilled water 3 oz.—Dr. Pereira.

3. Oil of lavender 2 oz., essence of ambergris 1 oz., eau de Cologne a pint, rectified spirit a quart.

4. Oil of lavender 4 dr.; essence of bergamot, essence of lemon or cedrat, and otto of roses, of each 20 minims; essence of ambergris 1 dr., rectified spirit 3 pints, orange-flower water 4 oz., rose (or distilled) water 12 oz., burnt alum 20 gr. Agitate frequently, then let it stand in a cool place for some days before filtering.

5. Oil of lavender 3 dr., oil of bergamot 20 drops, neroli 6 drops, otto 6 to 12 drops, essence of cedrat 8 or 10 drops,

essence of musk 20 drops, rectified spirit 28 fluid oz., distilled (or orange-flower) water 4 oz.

6. *Eau de Lavande aux Millefleurs.* Oil of lavender 4 dr.; essence of bergamot, essence of lemon, otto of roses, of each 12 drops; essence of millefleurs 3 dr., essence of ambergris 1 dr., rectified spirit a pint and a half.

Note.—The oil of lavender in the above should be the finest English oil; that which first comes over is said to be the most fragrant. It should be kept for 12 months before using, either alone or mixed with an equal quantity of alcohol. Some makers prefer a mixture of old and new oil. The lavender water improves by age.

Eau de Cologne—Cologne Water. 1. English oil of lavender, oil of bergamot, oil of lemon, oil of neroli, of each 1 oz.; oil of cinnamon ½ oz.; spirit of rosemary, and spirit of balm (*Eau des Carmes*), of each 15 oz.; highly rectified spirits 7½ pints. Let them stand together for 14 days, then distil in a water-bath.—Dr. Granville.

2. Oil of bergamot, citron, and lemon, of each 3 oz.; oils of rosemary, neroli, and lavender, of each 1½ oz.; oil of cinnamon 6 dr., rectified spirit 24 pints; compound spirit of balm (*Eau des Carmes*, below) 3 pints, spirit of rosemary 2 pints. Mix, and after standing a week, distil 24 pints.—French Pharmacopœia.

3. Essential oils of bergamot, of lemon, of neroli, of orange-peel, and of rosemary, each 12 drops; cardamom seeds a dr.; rectified spirit a pint. It improves by age.—Trommsdorf.

4. Essence of bergamot 40 minims, essence of lemon 45 minims, oil of rosemary 6, oil of orange 22, neroli 12 minims, highly rectified spirit 6 oz.

5. Alcohol a pint, oil of bergamot, oil of orange-peel, true oil of rosemary, cardamom seed, of each a drachm; orange-flower water a pint. Mix and distil a pint by water-bath.—Dr. A. T. Thomson.

Eau des Carmes—Eau de Melisse. Fresh flowering balm 24 oz.; yellow rind of lemon, cut fine, 4 oz., cinnamon, cloves, and nutmeg (bruised), of each 2 oz.; coriander seed (bruised) 1 oz., dried angelica root 1 oz., rectified spirit a gallon. Macerate for 4 days, and distil in a water-bath.

Arquebusade Water. 1. Sago, angelica, wormwood, savory, sweet fennel, hyssop, balm, sweet basil, rue, thyme, marjoram, rosemary, angelica seed, origanum, red calamint,

creeping thyme, lavender flowers, of each 10 oz.; sweet flagroot 5 oz., rectified spirit 2 gallons, water q. s. Distil 3 gallons.

2. (Simplified.) Balm, rosemary, thyme, calamus root, angelica seeds, lavender flowers, of each 4 oz.; rectified spirit 3 pints, water q. s. Macerate for a day, and distil 4 pints.

QUEEN OF HUNGARY'S WATER. Spirit of Rosemary. 1. Rosemary tops 2 ℔, rectified spirit 1 gallon, water q. s. Distil carefully 1 gallon.

2. Spirit of rosemary (as No. 1) 4 pints, orange-flower water ¼ pint, essence of neroli 4 drops.

3. Simple spirit of rosemary 3 pints, simple spirit of lavender 1 pint, rose-water 8 oz.

EAU D'ANGE. Flowering tops of myrtle 16 oz., rectified spirit 1 gallon; digest, and distil to dryness in a water-bath. Or dissolve ¼ oz. essential oil of myrtle in 3 pints of rectified spirit. Mr. GRAY gives under this name a water without spirit: Water 2 pints, benzoin 2 oz., storax 1 oz., cinnamon 1 dr., cloves 2 dr., calamus 1 stick, coriander seeds 1 pinch; distil.

HONEY WATER. *Eau de Miel.* 1. Rectified spirit 8 pints, oil of cloves, oil of lavender, oil of bergamot, of each ½ oz.; musk 15 gr., yellow-sanders shavings 4 oz.; digest for 8 days, and add 2 pints each of orange-flower and rose waters.

2. Oil of santal 20 drops, tincture of musk 2½ oz., essence of bergamot 2½ oz., oil of cloves 5 dr., oil of lavender 5 dr., rose-water 2 pints, orange-flower water 2 pints, spirit of wine 1 gallon; mix and filter.

3. (With honey.) White honey 8 oz., corinader seed 8 oz., fresh lemon-peel 1 oz., cloves ¾ oz., nutmeg, benzoin, styrax calamita, of each 1 oz.; rose and orange-flower water, of each 4 oz.; rectified spirit 3 pints; digest for a few days and filter. Some recipes add 3 dr. of vanilla, and direct only ½ oz. of nutmeg, storax, and benzoin.

4. Coriander seeds 7 ℔, cloves 12 oz., storax 8 oz., nutmeg 8 oz., fresh lemon-peel 10 oz., calamus root 6 oz., rectified spirit 15 pints; macerate for a month, add water q. s. Distil 22 pints, and add to the distilled spirit 5 pints of orange-flower water, 24 drops otto of roses, 1 dr. of ambergris, and 2 oz. of fine vanilla; macerate for a week, and filter. The dry ingredient to be bruised or cut small.

Lisbon Water. To rectified spirit, 1 gallon, add the essential oils of orange-peel and lemon-peel, of each 3 oz., and of otto of rose ¼ oz.—Piesse.

Eau de Portugal. To rectified spirit, 1 gallon, add the following essential oils: of orange-peel 6 oz., of lemon-peel 1 oz., of lemon-grass ¼ oz., of bergamot 1 oz., and of otto of rose ¼ oz.—Piesse.

Eau d'Elegance. Spirit of jessamine 2 ℔, spirit of styrax 1 ℔, spirit of hyacinth 1 ℔, spirit of star aniseed, 4 oz., tincture of balsam of Tolu 4 oz., tincture of vanilla 2 oz.

Eau de Mareschale. Spirit of wine 1½ pint, spirit of jessamine 1 oz., essence of bergamot ¼ oz., essence of violets 1 oz.

Eau Romaine. Spirit of jessamine 3 quarts, tincture of vanilla 1 quart, spirit of acacia flowers 1 quart, spirit of tuberose 1 pint, essence of ambergris 2 oz., tincture of benzoin 8 oz.

Eau de Millefleurs. Rectified spirits 2 pints, balsam of Peru ¼ oz., essence of bergamot ½ oz., oil of cloves ¼ oz., essence of neroli ½ dr., essence of musk 1 dr., orange-flower water 2 oz.

Eau Spiritueuse d'Heliotrope. Vanilla 3 dr., double orange-flower water 6 oz., rectified spirit 1 quart; macerate for 3 days, and distil in a water-bath. It may be colored with cochineal. But the Essence d'Héliotrope of some perfumers appear by the color, not to have been distilled.

Eau d'Ispahan. Essential oil of bitter orange-peel 4 oz., oil of rosemary 3 dr., oil of mint 1 dr., oil of cloves 7 scruples, neroli 7 scruples, spirit of wine 14 pints. It is used for the same purposes as Eau de Cologne.

Eau sans Pareille. Essential oil of lemon ½ oz., of bergamot 2½ dr., of cedrat ¼ oz., rectified spirit 6 pints, spirit of rosemary 8 oz.; mix. Some authorities state that it is improved by distillation.

Eau de Bouquet de Flore. 1. Honey-water 2 oz., tincture of cloves 1 oz., tincture of calamus, of lavender, and of long cyperus, each ½ oz.; eau sans pareille 4 oz.; spirit of jessamine 9 dr., tincture of orris 1 oz., spirituous essence of neroli 20 drops.

2. Essence of violets ½ oz., spirit of rosemary ½ oz.,

essence of lemon 1 dr., rectified spirit 24 oz., rose-water 8 oz.

3. Spirit of rosemary 8 oz., rectified spirit 8 oz., lavender water 2 oz., oil of neroli 5 drops, cloves 1 dr., orris root 2 dr., rose-water 2 oz.; digest for a few days, and filter.

ESPRIT DE BOUQUET. English oil of lavender, oil of cloves, and of bergamot, of each 2 dr.; otto of rose and oil of cinnamon, each 20 drops; essence of musk 1 dr., rectified spirit 1 pint; mix.

EAU DE ROSIERES. Spirit of roses 4 pints, spirit of jessamine 1 pint, spirit of orange-flowers 1 pint, spirit of cucumber $2\frac{1}{2}$ pints, spirit of celery seed $2\frac{1}{4}$ pints, spirit of angelica root $2\frac{3}{4}$ pints, tincture of benzoin (simple) $\frac{3}{4}$ of a pint, balsam of Mecca a few drops.

EAU D'AMBRE ROYALE. Rectified spirit 2 ℔, tincture of musk seed 1 ℔, essence of ambergris 1 oz., tincture of musk 1 oz., reduced with a proper proportion of orange-flower water.

ESPRIT DE SUAVE. Spirit of jessamine $1\frac{1}{2}$ pint, spirit of acacia flowers $1\frac{1}{2}$ pint, spirit of wine 12 oz., spirit of tuberose 8 oz., oil of cloves $1\frac{1}{2}$ dr., oil of neroli 30 drops, essence of bergamot $1\frac{1}{2}$ dr., tincture of musk 1 oz., rosewater 12 oz.

PARFUM DES ROIS. Spirit of wine 2 gallons, styrax 6 oz., benzoin 16 oz., aloes wood 8 oz., spirit of rose 2 pints, spirit of orange-flowers 2 pints, essence (tincture) of ambergris 8 oz., tincture of musk 8 oz., tincture of vanilla 16 oz.

ODOR DELECTABILIS. Rose-water, orange-flower water, each 4 oz.; oil of lavender, oil of cloves, each 1 dr.; oil of bergamot 2 dr., musk 2 gr, rectified spirit 1 pint.

NEW VICTORIA PERFUME. Cloves, bruised, 2 scruples; vanilla, cut small, 1 dr.; oil of cedrat 4 drops, oil of santal 1 dr., cinnamon 12 gr., oil of verbena 8 drops, otto of roses 8 drops, oil of neroli 20 drops, oil of lavender 1 dr., ambergris 16 gr., tincture of musk 1 dr., rectified spirit 16 fluid oz.; digest for a few days, and filter. Or the whole, except the musk and ambergris, may be distilled in a water-bath, and these added to the distilled spirit.

Another similar perfume is: Vanilla $\frac{1}{2}$ dr., yellow sanders

6 dr., cloves No. 16, neroli 3 drops, oil of lavender 6 drops, rectified spirit 4 oz.; digest for 3 days, and add 4 oz. of orange-flower water, water q. s.; distil 6 oz., add essence of musk 1 dr.

JOCKEY CLUB BOUQUET. PIESSE. Extract of orris root 2 pints, esprit de rose triple 1 pint, esprit de pommade de rose 1 pint, extract of pommade of cassia, and tuberose, ½ pint each; extract of ambergris ½ pint, otto of bergamot ½ oz.

ESPRIT DE ROSE. 1. Macerate the fresh and picked flowers of the most fragrant varieties of the rose with half their weight of rectified spirit, and distil in a water-bath to dryness.

2. Dissolve from 20 to 30 drops of otto in 1 pint of rectified spirit. A stronger solution, 6 or 8 drops of otto to 1 oz. of alcohol, forms essence of roses, or *esprit de rose triple*.

3. It is also made by agitating and digesting the spirit with the perfumed oil or pomade of roses. (See EXTRACTS below.)

ESPRIT DE JASMIN. *Eau de Jasmin.* It is prepared by digesting and agitating pure spirit with oil or pomade of jessamine made with the flowers. (See EXTRACTS, below.) Spirit of jonquil, tuberose, violet, &c., may be obtained by the same process.

ESPRIT DE VIOLETTE. *Eau de Violette.* Macerate 5 oz. of fine orris root in 1 quart of rectified spirit for some days, and filter. It may also be obtained by the method just mentioned, or by mixing the product of both processes.

EAU ODORANTE DE JASMIN. Compound spirit of jessamine, for the handkerchief. Spirit of jessamine 1 pint, rectified spirit 1 pint, essence of ambergris 1 dr., simple tincture of benzoin 1 dr.

SPIRIT OF ORANGE-FLOWERS, SPIRIT OF ELDER-FLOWERS, AND SPIRIT OF ACACIA-FLOWERS. Fresh flowers 1 ℔, rectified spirit 4 ℔, or pints, water 2 ℔; distil 4 ℔, or pints.

SPIRIT OF ORANGE-PEEL, OF LEMON-PEEL, OF CITRON, AND OF BERGAMOT. Fresh peel 1 ℔, rectified spirit 6 ℔; macerate for 2 days, and distil in a water-bath to dryness. Or, 1 oz. of the essential oil to 2 pints of spirits.

SPIRIT OF CINNAMON, OF CLOVES, OF NUTMEG, AND OF CALAMUS

Root. Macerate 1 ℔ of the bruised drug with 8 ℔, or a gallon of rectified spirit, and distil as the last.

Spirit of Cucumbers. Cucumbers grated 8 ℔, rectified spirit 1 ℔; distil 2 ℔.

Spirit of Rosewood. Rosewood shavings 1 ℔, spirit 6 ℔, water 2 ℔; distil 6 ℔. It is also made by adding the essential oil of rhodium to spirit.

Spirit of Angelica. Dried angelica root 1 ℔, rectified spirit a gallon. Macerate, and distil by water-bath to dryness.

Spirit of Balsam of Peru. Balsam 3 parts, spirit 15 parts, carbonate of potash 1 part; macerate for 3 days, and distil by water-bath.

Spirit of Strawberries, and of Raspberries. Fresh fruit 3 ℔, rectified spirit 1 ℔; macerate 24 hours, and distil 2 ℔.

The following tinctures are chiefly used in the compound perfumes:

Tincture of Balsam of Peru, and of Tolu. Digest 1 oz. of the balsam with 8 of the rectified spirit for some days, shaking it occasionally, then filter. Tincture of benzoin in the same manner.

Tincture of Angelica. One part of the dried root to 8 of rectified spirit; as the last.

Tincture (common spirituous essence) of Lemon, Citron, Orange, and Bergamot. An ounce of the fresh peel to ½ pint of spirit, as above.

Tincture of Musk Seed. *Essence d'Ambrette.* Digest 16 oz. of bruised musk seed with 3 pints of rectified spirit for a month, and filter.

Tincture of Musk. China musk 2 dr., rectified spirit 16 oz. For more compound tinctures of musk, see Essence of Musk, below.

Tincture or Essence of Ambergris. Guibourt directs 1 dr. of ambergris to be digested with a gentle heat in 3 oz. of rectified spirit. Another form is: Ambergris 1 dr., subcarbonate of potash 1 dr., spirit of roses 4 oz., (or rectified spirit 4 oz., otto 6 drops). Some recipes direct a weaker tincture: 24 gr. of ambergris to 8 oz. of spirits. For other formulæ, see Essence, below.

TINCTURE OF CIVET. Bruise ½ oz. of civet, ¼ oz. of ambergris, and the same of sugar candy, and macerate in a quart of rectified spirit for 6 weeks, in a warm place, then filter.

TINCTURE OF ESSENCE OF VANILLA. Vanilla cut very small 2 oz., rectified spirit a pint; infuse for 2 or 3 weeks. This is sometimes distilled, forming spirit of vanilla.

TINCTURE OF RHODIUM. Rosewood 1 lb, rectified spirit 3 or 4 pints; macerate for 3 or 4 weeks, and filter.

ESSENCE (OR TINCTURE) OF VETIVER. Take 2 lb of the root of vittie vayr cut small, and moisten it with a little water; let it macerate for 24 hours, then beat it in a marble mortar. Macerate it in sufficient spirit to cover it for 8 or 10 days, and strain with pressure: filter through paper, and in a fortnight repeat the filtration. Sometimes the root is moistened with diluted sulphuric acid, which, after maceration, is neutralized by adding a sufficient quantity of chalk, and the whole digested with spirit. The tincture, when strained off, is distilled, and forms (with the addition of essence of balm and of roses) Essence de Vetiver double.

ESSENCE OF PATCHOULI. Dried patchouli (puchu pat) 1 oz., rectified spirit a pint. It is generally combined with other perfumes.

EXTRACTS (extraits) are spirituous solutions of the odorous principle of flowers, obtained, indirectly, by agitating and digesting oils and pomatums which have been perfumed by the flowers (see HUILES ANTIQUES, under HAIR COSMETICS) with pure spirit. This is repeated with fresh oil until the spirit is sufficiently perfumed. When the same oil or pomade is treated with fresh spirit, inferior extracts, numbered 2, 3, &c., are obtained. These preparations are chiefly made in France.

EXTRACTS (EXTRAITS, or ESPRITS) OF JESSAMINE, VIOLETS, LILY OF THE VALLEY, are prepared by the process just mentioned.

EXTRAIT DE BOUQUET. Spirit (extrait) of jessamine 2 quarts, extract of violets 2 quarts, spirit of acacia-flowers, of rose, and of orange-flowers, each a quart, spirit of carnations a quart, flowers of benzoin ½ oz., essence of ambergris 8 oz.

EXTRAIT DE MARESCHALE. Essence of millefleurs 1½ oz., essence of jessamine 1 oz., essence of musk ½ oz., essence of ambergris ½ oz., essence of cedrat 20 drops, essence of violets 1 oz., sweet spirits of nitre 50 drops, true oil of rosemary 20 drops, rectified spirit 6 oz., oil of neroli 48 drops. Set aside for some time.

COMPOUND ESSENCES. Some of these contain a preparation of the substance whose name they bear, while others are fictitious or imitative, being made up of a variety of other essences and volatile oils. Several of the formulæ are those of M. PIESSE.

ESSENCE OF AMBERGRIS. This name is applied both to the simple and more compound tinctures of ambergris. See TINCTURE OF AMBERGRIS, above. Other formulæ may here be given.

1. Ambergris 4 oz., musk 2 oz., tincture of musk seed 7 pints. Digest with a gentle heat.

2. Ambergris 1 dr., musk ½ dr., oil of cinnamon 18 drops, oil of rhodium 12 drops, rectified spirit 8 oz., spirit of roses 4 oz., carbonate of potash 1½ dr.; digest in a warm place for a few days, and strain. See also ESSENCE ROYALE.

ESSENCE OF CEDRAT. Dissolve 2½ oz. of oil of cedrat in 1 gallon of spirit, and add bergamot ½ oz.

ESSENCE OF CLOVE PINK. Esprit de rose ½ pint, de fleur d'orange and de fleur de cassie, each ¼ pint, esprit de vanille 2 oz., oil of cloves 10 drops.

COLOGNE ESSENCE. Oil of bergamot 2 dr., essence of lemon ½ dr., essence of cedrat ½ dr., true oil of rosemary 15 drops, rectified spirit (or spirit of balm) 1½ oz.

ESSENCE OF HELIOTROPE. Spirituous extract of vanilla ½ pint, of French rose pomatum ¼ pint, of orange-flower pomatum 2 oz., of ambergris 1 oz.; add 5 drops of the essential oil of almonds.

ESSENCE OF HONEYSUCKLE. Spirituous extract of rose pomatum 1 pint, of violet 1 pint, of tuberose 1 pint; extracts of vanilla and Tolu, of each 1 pint; oil of neroli 10 drops, essential oil of almonds 5 drops.

ESSENCE OF HOVENIA. Rectified spirit 1 quart, rose water ½ pint, essential oil of lemons ½ oz., otto of roses 1 dr., oil of cloves ½ dr., oil of neroli 10 drops.

ESSENCE OF JONQUIL. Spirituous extract of jasmine pomade 1 pint, of tuberose 1 pint, of orange-flower ½ pint: add extract of vanilla 2 oz.

ESSENCE OF LILY OF THE VALLEY. Mix the following extracts: of tuberose ½ pint, of jasmin 2 oz., of orange-flower 2 oz., of vanilla 3 oz., of cassia ¼ pint, of rose-water ¼ pint: add 3 drops of hydrocyanic acid. Keep together for a month, then bottle.

ESSENCE OF MAGNOLIA. Spirituous extract of orange-flower pomatum 1 pint, of rose pomatum 2 pints, of tuberose pomatum ½ pint, of violet pomatum ½ pint; essential oil of citron 2 dr., essential oil of almonds 10 drops.

ESSENCE OF MIGNONETTE. Digest 1 ℔ pomade de rézéda in rectified spirit 1 pint for 14 days; filter off, and add 1 oz. of extrait d'ambre.

ESSENCE OF MOSS ROSE. Spirituous extract of French rose pomatum 1 quart, esprit de rose triple 1 pint, extract of orange-flower pomatum 1 pint, of ambergris ½ pint, and of musk 4 oz.

ESSENCE OF MYRTLE. Take the following extracts: of vanilla ½ pint, of roses 1 pint, of orange-flower ½ pint, of tuberose ½ pint, of jasmin 2 oz. Mix, and allow to stand for a fortnight.

ESSENCE OF MUSK. A tincture of musk, of various strength. The formulæ given above (tincture of musk) is that of the Dublin Pharmacopœia, 1826. GUIBOURT directs 1 part of musk to 12 of proof spirit. Other authorities direct a smaller quantity of musk. A French work gives the following: Musk in the bag cut small 6 oz., civet 1 oz., tincture of musk seed 7 pints; digest in the sun, or in a warm place for 2 months.

ESSENCE OF PATCHOULI. Oil of patchouli 1¼ oz., otto of roses ¼ oz., rectified spirit 1 gallon.

ESSENCE OF RONDELETIA. Essence of bergamot, essence of lemon, oil of cloves, each 1 dr., otto of roses 6 drops, rectified spirit 1 pint.

ESSENCE ROYALE. Ambergris 1 dr., civet 15 gr., musk 30 gr., carbonate of potash 20 gr.; triturate together, and add oil of cinnamon 10 drops, oil of rhodium, and of neroli, 6 drops, otto of roses 6 drops, rectified spirit ¼ pint; digest and filter.

ESSENCE OF SWEET BRIER. Spirituous extracts of French rose pomatum 1 pint, of cassia and orange-flowers, each ¼ pint, espirit de rose ¼ pint, with oils of neroli and lemon-grass, of each ½ dr.

ESSENCE OF SWEET PEA. Essences of tuberose, orange-flower, and rose-pomatum, each ½ pint, with essence of vanilla 1 oz.

ESSENCE OF VERBENA. Essential oil of verbena 2 dr., rectified spirit 4 oz., essence of ambergris ½ dr., orange-flower water ½ oz.; mix. Another form is: Oil of verbena ½ oz., essence of vanilla 40 drops, rectified spirit 4 oz.; mix and filter.

ESSENCE OF WHITE LILAC. Spirituous extract of tuberose pomade 1 pint, of orange-flower pomade ¼ pint; add essential oil of almonds 3 drops, and extract of civet ½ oz.

FRANGIPANNI BOUQUET. Essence of vetiver 3 oz., ol. neroli 15 minims, oil of sandal-wood ½ dr., otto of rose 40 minims, essence of musk 3 dr., esprit de violette 3 oz., essence of ambergris 6 dr., rectified spirit to make up 20 oz.—PIESSE.

ESSENCES FOR SCENTING POMATUMS. *Millefleur:* Oil of lemon 3 oz., essence of ambergris 4 oz., oil of cloves 2 oz., oil of lavender 2 oz.—*Cowslip:* Essence of bergamot 16 oz., essence of lemon 8 oz., oil of cloves 4 oz., oil of orange-peel 2 oz., oil of jessamine 2 dr., eau de bouquet 2 oz., oil of bitter almonds 16 drops.—*For general use:* Essence of bergamot 16 oz., essence of lemon 8 oz., true oil of origanum and oil of cloves, each 2 oz., oil of orange-peel 1½ oz.

MISTURA ODORATA. Rectified spirit 48 oz., tincture of benzoin 4 oz., tincture of vanilla ¼ oz., tincture of musk ½ oz., balsam of Peru ½ oz., oil of cloves, of mace, and of cinnamon, each ½ oz., oil of bergamot 1 oz., oil of cedrat 2 oz.—GIESKE.

SCENT FOR SNUFF. Oil of lavender 2 dr., essence of lemon 4 dr., essence of bergamot 1 oz.: mix. (1 dr. with 8 oz. of fine Scotch snuff constitutes Queen's Snuff.)

[The following Essences, Spirits, and Waters are given as specimens of some of the cheaper perfumes as made in France.]

Essences (Spirituous).

Essence (Spirituous) of Neroli. Spirit of wine ½ pint, orange-peel cut small 3 oz., orris-root in powder 1 dr., musk 2 gr.; let it stand in a warm place for 3 days, and filter.

Essence of Lemon. Spirit of wine ½ pint, fresh lemon-peel 4 oz.; as above.

Essence of Bergamot. Spirit of wine ½ pint, bergamot-peel 4 oz.; as above.

Essence of Violets. Spirit of wine ½ pint, orris-root 1 oz.

Essence of Cedrat. Essence of bergamot (as above) 1 oz., essence of neroli 2 dr.

Essence of Jessamine. Essence of violets 1 oz., essence of cedrat 2 dr.

Essence of Musk. Spirit of wine ½ pint, musk 16 gr.

Essence of Ambergris. Spirit of wine ½ pint, ambergris 24 gr.

Essence of Cloves. Spirit of wine ½ pint, bruised cloves 1 oz. Other essences in the same manner.

Spirits.

Spirit of Rose. Spirit of wine ½ pint, otto 6 drops.

Spirit of Jessamine. Spirit of wine ½ pint, essence of jessamine (as above) a drachm.

Spirit of Orange. Spirit of wine, essence of orange, or neroli, a drachm.

Spirit of Lavender. Spirit of wine ½ pint, essential oil of lavender a drachm.

Spirit of Musk. Spirit of wine ½ pint, essence of musk a drachm.

Others in a similar manner.

Simple Waters.

Rose Water. Distilled or rain water ½ pint, spirit of roses a drachm.

Jessamine Water, *Musk Water*, *Violet*, *Orange-flower Water*, &c., by adding a dr., of the above spirits to ½ pint of water.

AMMONIATED PERFUMES.

AMMONIATED COLOGNE WATER. A fragrant and reviving substitute for Spirits of Sal Volatile. Muriate of ammonia 5 dr., subcarbonate of potash 8 dr., eau de Cologne 12 oz., essential oil of cedrat and of bergamot, of each 15 drops (dissolved in an oz. of rectified spirit), orange-flower water 8 oz.; mix, and carefully distil 15 or 16 oz.

EAU DE LUCE. Mastic 2 dr., rectified spirit 9 dr.; dissolve and add to the clear tincture 30 drops of oil of lavender, 10 drops of bergamot, and a pint of strong water of ammonia. This is more agreeable than the compound of the Pharmacopœia, which, however, should always be used when prescribed medicinally.

ESSENCE FOR SMELLING SALTS. 1. English oil of lavender and essence of bergamot, of each a dr., oil of orange-peel, or of cedrat, 8 drops, oil of cinnamon 4 drops, oil of neroli 2 drops, alcohol, and strongest water of ammonia, of each 2 oz. (or 4 oz. of strong ammoniated alcohol).

2. Ammoniated alcohol 12 fluid oz. English oil of lavender, essence of bergamot and essence of lemon, of each a dr., cloves ½ dr., camphor ½ oz., macerate for a week, and filter. —Mr. MAGGS.

3. Essence of ambergris and musk 4 dr., otto of rose 20 drops, oil of lavender 1 dr., ammoniated alcohol 10 oz.; mix, and add strongest liquor ammoniæ 10 oz.—PHARM. JOUR.

SMELLING SALTS. Sesquicarbonate of ammonia 40 oz., broken into small pieces not any larger than a filbert, put into an air-tight ½ gallon jar; pour over it 20 oz. strong solution of ammonia (sp. gr. .880) previously perfumed according to taste, and immediately fix on lid of jar, taking care that it is properly secured. Keep in a cool place, opening and stirring with a stiff spatula every other day for a week; allow it now to remain for 2 or 3 weeks, at the end of which time it will have become hard.—Mr. ALLCHIN.

GODFREY'S SMELLING SALTS. Dr. PARIS says it is prepared by resubliming volatile salts with subcarbonate of potash and

a little spirit of wine. It is usually scented with an alcoholic solution of essential oils.

ACETIC PERFUMES.

AROMATIC SPIRIT OF VINEGAR. 1. Strong acetic acid 16 oz., camphor 1 oz.; when dissolved, add 1 oz. each of essential oils of cloves, lavender, and lemon. This is said to resemble HENRY'S.

2. Glacial acetic acid 8 oz., true oil of rosemary 20 gr., of bergamot 15 gr., of lavender 9 gr., of cloves 24 gr., neroli 4 gr., cinnamon 20; dissolve the oils in 2 dr. of rectified spirit. For another formula, see Pocket Formulary.

AROMATIC VINEGARS are made in France by infusing various flowers, &c., in distilled or finest wine vinegars, with or without the addition of spirit. Others are made by distillation. As they are seldom required in this country, a few examples will suffice.

ROSE VINEGAR. Red roses, picked and dried, ½ lb, best vinegar 8 lb; macerate for a fortnight, with occasional stirring, and strain; then filter.

LAVENDER VINEGAR. Fresh lavender-flowers 1 lb, vinegar 12 lb. Macerate as above. It is sometimes distilled, drawing off 8 lb.

DISTILLED ROSE VINEGAR. Pale roses, dried, 2 lb, distilled vinegar 8 lb. Distil three-fourths by sand-bath, and add 2 lb of spirit of roses. It is occasionally colored with cochineal, and used as a cosmetic.

ORANGE-FLOWER VINEGAR. Fresh orange-flowers 1½ lb, distilled vinegar 8 lb, spirit of orange-flowers 1 lb. Macerate for 12 days, strain and filter.

VINAIGRE VIRGINAL. Benzoin in powder 2 oz., rectified spirit 8 oz., white vinegar 2 lb. Digest the benzoin in the spirit for 6 days, strain, and add the vinegar to the residue; macerate for 6 days, decant and add to it the tincture. The next day filter. It is chiefly used as a cosmetic.

VINAIGRE DE COLOGNE. To each pint of eau de Cologne add an ounce of strong acetic acid.

VINAIGRE DE JOUVENCE. Spirit of cucumber 4 oz., spirit of storax 2 lb, strong vinegar 8 lb.

VINAIGRE DE FLORE. Equal parts of rose vinegar, vinaigre virginal, and orange-flower vinegar.

VINAIGRE DE QUATRE VOLEURS. Thieves' vinegar. Dried tops of large and small (pontic) wormwood, rosemary, sage, mint, rue, lavender-flowers, of each 2 oz.; calamus root, cinnamon, cloves, nutmeg, garlic, of each ¼ oz.; camphor ½ oz., concentrated acetic acid 2 oz., strong vinegar 8 lb. Macerate the herbs, &c., in the vinegar for a fortnight, strain, press, and add the camphor dissolved in the acetic acid.

POT POURRI; SCENTED POWDERS; SACHETS OR SCENT BAGS; SCENT BALLS, PASTILS, &c.

POT POURRI. 1. Gather in the season the petals of the most fragrant kinds of roses (with which other flowers may be mixed, at pleasure, in small proportion); spread them out to dry in the sun, or in a warm room, sprinkle a little salt on them, and put them into a jar, in which they are to be kept covered up till wanted for use. Take of these rose leaves 4 oz., dried lavender flowers 8 oz., vanilla, cloves, storax, and benzoin, all bruised, of each 1 dr., ambergris 20 gr., otto of roses 20 drops; mix.

2. Calamus root, yellow sanders, of each 1 oz., vanilla 1 dr., musk 8 gr., ambergris 8 gr., cascarilla 1 oz., orris root 3 oz., cinnamon 1 oz., lavender flowers 1 oz., storax 2 dr., benzoin 2 dr., cloves 2 dr., coriander seed 1 oz., nutmegs 2 dr., otto of roses 20 drops, oil of neroli 10 drops. The dry ingredients to be coarsely bruised; mix.

3. French. Take the petals of the pale and red roses, pinks, violets, moss rose, orange-flower, lily of the valley, acacia flowers, clove-gilliflowers, mignonette, heliotrope, jonquils; with a *small* proportion of the flowers of myrtle, balm, rosemary, and thyme; spread them out for some days, and as they become dry, put them into a jar with alternate layers of dry salt, mixed with orris powder, till the vessel is full. Close it for a month, then stir the whole up, and moisten it with rose-water.

4. Orris root 16 oz., dried acacia flowers 8 oz., dried bergamot-peel 2 oz., musk seed ¼ oz., cloves ¼ oz.; pound them together.

5. Dry rose leaves quickly on a wicker tray, in a warm place. To a pint of the petals add powdered orris 2 oz.,

pimento ½ oz., cascarilla ¼ oz., musk 2 gr., otto of roses 2 drops, bruised cloves ¼ oz.

Sachets or Scent Bags. The pot pourri No. 2 or 4 may be put into bags, alone, or with any perfume to increase the strength. Or coarsely powdered patchouli (an herb of the Pogostemon genus) may be used, with any other perfume. Or the bags may be filled with carded cotton mixed with any of the following scented powders:

Scented Powders, Balls, &c.

Rose. Powdered starch 3 oz., carmine to color, otto of roses 8 drops, orris powder 1 oz.

Violet. Orris powder 4 oz., essence of bergamot 20 drops, essence of ambergris 20 drops.

Poudre de Chypre. Oak-moss is macerated in clean water for a day or two, and strongly pressed in a cloth; it is then moistened with rose-water mixed with a third of orange-flower water for two days, pressed, and pulverized. It serves as a basis for other perfumes, the power of which it is said to increase.

Poudre à la Mousseline. Orris 16 oz., coriander seed 8 oz., musk seed 2 oz., cinnamon, cloves, and sandal-wood, each 1 oz., star aniseed ¼ oz., mace, ginger, and violet ebony, of each 2 oz.; beat them to a powder, and pass through a sieve.

Poudre à l'Œillet. Red roses 48 oz., orris 48 oz., cloves 6 oz., bergamot-peel 20 oz., musk seed 24 oz., cinnamon 6 oz., long cyperus 6 oz., pale roses 26 oz., dried acacia flowers, orange flowers, and clove stalks, of each 8 oz.

Poudre à la Mareschale. Oak-moss in powder 2 ℔, plain starch powder 1 ℔, cloves 1 oz., calamus 2 oz., cyperus 2 oz., rotten oak-wood powder 2 oz.; mix.—Gray.

Portugal. Dried orange-peel 1 oz., dried bergamot-peel ½ oz., cloves 4 oz., storax 1 dr., ambergris 8 gr., benzoin a drachm, musk seed a scruple, musk 4 gr.

Scented Balls, Medallions, &c. *Pastilles de Toilette odorantes.* These consist of perfumed powders made into a paste, and moulded to any desired form before drying. The above scent powders beaten up with mucilage of tragacanth will answer the purpose; or the following:

1. Beat the fresh petals of red roses in an *iron* mortar to a smooth paste, with a few drops of essence of ambergris, or

other suitable perfume. It becomes sufficiently smooth to take a polish.

2. Powdered orris, oak-moss, and poudre de mousseline, of each 1 oz.; lamp-black, or other color, q. s. Form into a stiff paste, with a jelly made of 6 dr. of isinglass, 2 of tragacanth, and boiling water q. s. Make it into beads by means of a pill-machine, or into any ornamental form by moulds.

3. Jessamine flowers 1 oz., powdered gum tragacanth ½ oz., vermilion 2 oz.

4. Yellow sanders, cyperus, cloves, balsam of Peru, of each 2 dr., benzoin and storax, of each ½ oz., musk and civet, of each 10 gr., oil of cinnamon 5 drops, oil of rhodium 15 drops, essence of jessamine 1 dr., neroli 20 drops, ivory-black 1½ oz., Paris plaster 2 oz., mucilage of tragacanth, made with rose-water, q. s. As the last.

PASTILS FOR BURNING. 1. Yellow sanders 3 oz., storax 4 oz., benzoin 3 oz., olibanum 6 oz., cascarilla 6 oz., ambergris 1 dr., Peruvian balsam 2 dr., myrrh 1½ oz., nitre 1½ oz., oil of cinnamon 20 drops, oil of cloves ½ dr., otto 30 to 60 drops, oil of lavender 1½ dr., balsam of Tolu 1½ oz., camphor ½ oz., strong acetic acid 2 oz., charcoal 3 ℔; mix s. a., and beat into a paste with mucilage of tragacanth, and form into conical pastils. A second and third quality may be made by using, respectively, 4 and 5 ℔ instead of 3 ℔ of charcoal. These are highly approved, but rather expensive.

2. (*Clous fumans* of the French Codex.) Benzoin 2 oz., balsam of Tolu ½ oz., labdanum 1 dr., yellow sanders ½ oz., light charcoal 6 oz., nitre ¼ oz., mucilage of tragacanth q. s. Reduce the substance to powder, and form into a paste with the mucilage, and divide into small cones with a tripod base.

3. Powdered cascarilla 8 oz., benzoin 4 oz., yellow sanders 2 oz., storax calamita 2 oz., olibanum 2 oz., charcoal 3 ℔, nitre 1½ oz., mucilage of tragacanth q. s.

4. Benzoin 1 oz., cascarilla 1 oz., myrrh 8 scruples, oil of nutmeg 4 scruples, oil of cloves 4 scruples, nitre ½ oz., charcoal 6 oz., mucilage of tragacanth q. s.—Dr. PARIS.

INCENSE. 1. Storax 2½ oz., benzoin 12 oz., musk 15 grains, burnt sugar ½ oz., frankincense 2½ oz., gum tragacanth 1½ oz., rose-water sufficient to form a mass; to be divided into small tablets.—Mr. ASTLEY.

2. Powdered cascarilla 2 oz., myrrh, storax, benzoin, thus, Burgundy pitch, each 1 oz.; mix.—Mr. Atkins (Ph. Journal).

Mouth Pastils. Dry compounds for perfuming or correcting the breath.

Cachou Aromatisé. The basis of these compounds, as the name implies, was originally *catechu*, with which various odoriferous substances were combined. The catechu, however, is now often omitted. The following are some of the most approved forms:

1. Extract of liquorice 3 oz., oil of cloves 1½ dr., oil of cinnamon 15 drops; mix, and divide into one-grain pills, and silver them.

2. (M. Chevallier's.) Chocolate powder and ground coffee, of each 1½ oz., prepared charcoal 1 oz., sugar 1 oz., vanilla (pulverized with the sugar) 1 oz., mucilage q. s. Make into lozenges of any form, of which 6 to 8 may be used daily to disinfect the breath.

3. *Cachou de Bologne.* Bologna Catechu. Extract of liquorice 3 oz., water 3 oz., dissolve by heat in a water-bath, and add catechu 1 oz., gum arabic ½ oz., evaporate to the consistence of an extract, and add (in powder) ½ dr. each of mastic, cascarilla, charcoal, and orris: remove from the fire, and add oil of peppermint ½ dr., essence of ambergris and essence of musk each 5 drops; roll it flat on an oiled marble slab, and cut it into very small lozenges. (Or it may be rolled into small pills, and silvered. They are chiefly used by smokers.)

4. Catechu 7 dr., orris powder 40 gr., sugar 3 oz., oil of rosemary (or of peppermint, cloves, or cinnamon) 4 drops, or q. s. Proceed as for the last.

5. *Cachou Aromatisé.* Extract of liquorice and water, of each 3½ oz.; dissolve in a water-bath, and add Bengal catechu in powder, 462 grains, and gum arabic in powder 231 grains; evaporate to an extract, and then incorporate the following substances, first reduced to a fine powder: Mastic, cascarilla, charcoal, and orris root, of each 30 grains; melt the mass to a proper consistence, remove it from the fire, and then add English oil of peppermint 30 drops, tinctures of ambergris and musk, of each 5 drops; pour it now on an oiled slab, and spread it out, by means of a roller, to the thickness of a sixpenny piece. When cool, apply some folds of blotting-paper to absorb any

adherent oil, moisten the surfaces with water, and cover it with sheets of silver leaf. Allow it to dry, and finally divide into thin strips, and these again into small pieces, about the size of a fenugreek seed.—(Journal de Pharmacie.)

PASTILS OR LOZENGES, with chlorine, for disinfecting the breath. 1. Sugar flavored with vanilla 1 oz., powdered tragacanth 20 gr., liquid chloride of soda q. s., any essential oil 2 drops. Form a paste, and divide into lozenges of 15 gr. each.

2. Dry chloride of lime 2 dr., sugar 8 oz., starch 1 oz., gum tragacanth 1 dr., carmine 2 gr. Form into small lozenges.

SKIN COSMETICS.

WASHES FOR THE FACE.

AQUA COSMETICA. Cosmetic Lotion. 1. Emulsion of bitter almonds 3 oz., rose and orange-flower water, of each 4 oz., borax 1 dr., tincture of benzoin 2 dr.; mix.—Dr. COPLAND.

2. Elder-flower water a pint, borax ½ oz., eau de Cologne 1 oz.; mix.

KALYDOR. The following is said to resemble KALYDOR and GOWLAND'S lotion: Bitter almonds blanched 1 oz., corrosive sublimate 8 gr., rose-water 16 oz.

MILK OF ROSES. 1. Sweet almonds 5 oz., bitter almonds 1 oz., rose-water 2½ pints, white curd soap ½ oz., oil of almonds ½ oz., spermaceti 2 oz., white wax ½ oz., English oil of lavender 20 drops, otto of roses 20 drops, rectified spirit a pint. Blanch the almonds, and beat them with the soap and a little of the rose-water. Melt together the oil of almonds, spermaceti, and white wax, and mix with the former into a cream, and strain it through fine muslin. Then add gradually the remaining rose-water, and lastly the spirit, with the essential oils dissolved therein.

2. A common kind is made by mixing 1 oz. of fine olive oil with 10 drops of oil of tartar, and a pint of rose-water.

3. Bitter almonds 6 dr., sweet almonds 12 dr., blanch, dry, and beat up with 1 dr. of Castile soap; gradually adding 15 gr. of spermaceti, 30 gr. of white wax, and a dr. of almond oil, melted together. When thoroughly incorporated add gradually six drops of otto of roses, dissolved in 6 oz. of rectified spirit, and 14 oz. of distilled water.

MILK OF CUCUMBERS. In the same manner as milk of roses, substituting juice of cucumbers for the rose-water.

MILK OF HOUSELEEK. As milk of roses No. 1, substituting expressed juice of houseleek for a pint of the rose-water.

ALIBERT'S COSMETIC. Cucumber pomade (see below) 3 oz., almond soap 1 oz., rose-water a quart; mix the pomade and soap, and add the rose-water gradually.

SIEMMERLING'S COSMETIC. Make an emulsion with 1 oz. of sweet almonds, ½ oz. bitter almonds, black cherry-water 10 oz., and bichloride of mercury 5 dr., tincture of benzoin 5 dr., lemon-juice ½ oz.

WITHERING'S (Dr.) COSMETIC. An infusion of horseradish in milk.

LAIT VIRGINAL. Virgin's Milk. Simple tincture of benzoin 2 dr., orange-flower water 8 oz. It may be varied by using rose or elder-flower water.

LAIT DE FRAICHEUR. Double rose-water 8 oz., tincture of benzoin 4 dr., balsam of Mecca ½ oz.

SCHUBARTH'S COSMETIC EMULSION. Almond Emulsion (made with rose-water) 8 oz., tincture of benzoin 3 dr.

ITALIAN COSMETIC WASH. Melilot water 12 oz., tincture of benzoin 2 dr.

AUGUSTIN'S. Rose-water 8 oz., salt of tartar 2 dr., tincture of benzoin 3 dr.

KITTOE'S LOTION FOR FRECKLES. Muriate of ammonia 1 dr., spring water a pint, lavender-water 2 dr. Apply with a sponge 2 or 3 times a day.

LEMON CREAM FOR SUNBURNS, FRECKLES, &c. Sweet cream 1 oz., new milk 8 oz., juice of lemon, brandy, or eau de Cologne 1 oz., alum 1 oz., sugar 1 dr. Boil and skim. Buttermilk is used for the same purpose.

LEMON EMBROCATION FOR FRECKLES, &c. Borax 15 gr., lemon-juice 1 oz., sugar candy ½ dr.; mix the powders with the juice, and let them stand in the bottle, shaking occasionally, till they are dissolved.

PASTES, POMADES, COLD CREAMS, LIP-SALVE, &c.

POMMADE DE BEAUTE. Melt together in an earthen vessel placed in hot water, white wax ½ dr., spermaceti 2 dr., oil of sweet almonds ½ oz., virgin olive oil ½ oz., oil of poppies ½ oz.; beat them with a few drops of balsam of Peru.

Cucumber Pomatum, for softening and cooling the skin. Clarified lard 4 lb, veal suet 1 lb, juice of cucumbers 3 lb; melt the two former together, then beat it up assiduously with the juice. Next day pour off the juice that has separated, and add the same quantity of fresh to the melted pomade. Repeat this six times, or until the pomade is sufficiently imbued with the odor of cucumbers. Then melt the pomade by a water-bath, and mix with it 3 dr. of powdered white starch; let it settle, and before it is too cold, pour it off into small pots, taking care not to disturb the dregs.

Pommade d'Hebe. Incorporate together the juice of lily-bulbs 2 oz., Narbonne honey 2 oz., white wax 1 oz., rose-water 3 dr.; melt the wax with a gentle heat, and add the other ingredients. To be applied at night, and not wiped off till morning. To remove wrinkles. Probably cod-liver oil, used externally and internally, would be a more successful though less agreeable remedy.

Pate Divine de Venus. Mixed equal parts of washed lard, fresh butter, and white honey; add balsam of Mecca and otto of roses, to perfume.

Pommade de Ninon. Oil of sweet almonds 4oz., washed lard 3 oz., juice of houseleek 3 oz.; mix. Softening and cooling.

Pommade en Creme. Melt together 1 dr. each of white wax and spermaceti, add oil of sweet almonds 2 oz.; pour it into a warm mortar, and gradually stir in ½ oz. of rose or perfumed water, and 1 dr. of tincture of Tolu.

Lemon Cream. Melt together 2 dr. of spermaceti and 1 oz. of oil of almonds; and as it cools stir in 16 drops of essence of lemon.

Cold Cream. 1. Oil of almonds 16 oz., white wax 4 oz., melt together in an earthen vessel, and when nearly cold, stir in, by little and little, 12 oz. of rose-water.

2. Melt together white wax 2 oz., oil of almonds 8 oz., and stir in 4 oz. of rose-water. Next day add 6 drops of otto of roses.

3. White wax and spermaceti, of each ½ oz.; oil of almonds 4 oz., orange-flower water 2 oz.; mix s. a.

4. As No. 3, but without the orange-flower water.

5. Lard 16 oz., white wax 2 oz., olive oil 1 oz., magistery of bismuth 1 oz.

6. White wax 1 oz., almond or olive oil 1 oz., rose-water 1 oz., glycerine 2 dr.

N. B.—Those cold creams are generally preferred for present use which contain rose or other water, but they keep longer without them.

Granulated Cold Cream. Melt together 1 oz. each of white wax and spermaceti, with 3 oz. of almond oil; when a little cooled, pour the mixture into a large Wedgewood mortar, previously warmed, and containing about a pint of warm water. Stir briskly until the cream is well divided, add sufficient otto of roses to scent it, and pour the whole suddenly into a clean vessel containing 8 or 10 pints of cold water. Throw the whole on muslin, and shake out as much water as possible.—Mr. Owen. Dublin.

Pommade Divine. Put 3 lb of beef marrow into an earthen vessel, and cover it with cold water, changing the water daily for a few days, and using rose-water the last day; press out the water and add to the marrow storax calamita, benzoin, Chio turpentine, each 4 oz.; orris powder 1 oz., powdered cinnamon, cloves, nutmeg, of each ½ oz. Place them in a well-tinned vessel in a water-bath, and keep the water boiling for 3 hours; then strain.

Almond Paste for the Skin. 1. Powdered bitter almonds 4 oz., white of egg 1 oz., beat them well together to a smooth paste, with equal parts of spirits of wine and rose-water.

2. Sweet and bitter almonds, blanched, of each 2 oz.; spermaceti 2 dr., oil of almonds ½ oz., Windsor soap ½ oz., rose-water 1 oz., or q. s.; otto of roses and oil of bergamot, of each 12 drops.

3. (Camphorated.) To either of the above add 2 dr. of powdered camphor. A few drops of oil of bitter almonds may be substituted for the otto and bergamot.

4. (French.) Blanch 12 oz. of bitter almonds and beat them in a mortar with a small quantity of rose or other water to a smooth paste; then add 7 oz. of rice flour, 3 oz. of bean flour, 1 oz. of orris powder, and, when perfectly mixed, ½ oz. of carbonate of potash, dissolve in rose-water; again beat together, and add 3 oz. of spirituous essence of jessamine, 2 drops of oil of rhodium, and 1 of neroli.

Almond and Honey Paste. Fine honey may be added to either of the preceding; or mix 16 oz. of clarified honey with 16 oz. of bitter almond powder; and add gradually, in alternate portions, 32 oz. of oil of almonds, and the yolks of 5 eggs.

Honey Paste. *Pâte au Miel.* It is sometimes made as the last; or, by mixing clarified honey with cold cream, or some similar compound.

Camphor Balls, for rubbing on the hands, after washing them, to prevent chaps, &c. 1. Melt 3 dr. of spermaceti and 4 dr. of white wax with 1 oz. of almond oil, and stir in 3 dr. of powdered camphor. Pour the compound into small gallipots, so as to form hemispherical cakes. They may be colored with alkanet, &c.

2. Lard 3 oz., white wax 2 oz., camphor ½ oz.

3. Spermaceti 3 oz., white wax 1 oz., olive oil 4 oz.; melt together, and add 11 oz. of powdered camphor, and stir it well.

4. Melt 3 dr. of spermaceti and 4 dr. of white wax with 1 oz. of almond oil, and stir in 3 dr. of powdered camphor.

Camphor Ice. Melt 1 dr. of spermaceti with 1 oz. of almond oil, and add 1 dr. of powdered camphor.

Almond Powder. (Cosmetic.) This is prepared by grinding the marc or cake left after expressing the oil from sweet or bitter almonds. It is sometimes perfumed, and mixed with other ingredients. It is used for cleansing the skin, and is less irritating than soap.

Almond Wash Powder. 1. Almond powder (from expressed bitter almonds) 16 oz., rice flour 2 oz., powdered soap 1 oz., orris powder 1 oz., bergamot, or other scent, q. p.

2. Almond powder (as above) 16 oz., powdered benzoin ¼ oz., oil of bitter almonds 10 drops. For cleaning the hands and removing any unpleasant smell. To render it more detergent, 4 oz. of fine sand or powdered pumice-stone may be added.

Rose Lip Salve. 1. Oil of almonds 3 oz., alkanet ½ oz.; digest with a gentle heat, and filter. Melt 1½ oz. white wax and ½ oz. spermaceti with the filtered oil, stir it until

it begins to thicken, and add from 12 to 36 drops of otto of roses.

2. White wax 1 oz., oil of sweet almonds 2 oz., alkanet 1 dr.; digest till colored, strain, and add 6 drops of otto of roses.

PERUVIAN LIP SALVE. As either of the above, substituting 20 or 30 drops of Peruvian balsam for the otto; 8 drops of oil of lavender may be added.

GRAPE LIP SALVE. *Pommade au raisin pour les lèvres.* Put into a glazed earthen pipkin ½ ℔ of fresh butter, ¼ ℔ fine yellow wax, 1 oz. of alkanet, and 3 bunches of black grapes; boil together, and strain without pressure through linen.

FRENCH LIP SALVE. Lard 16 oz., white wax 2 oz.; nitre and alum in fine powder, of each ½ oz.; alkanet to color.

GERMAN LIP SALVE. Butter of cacao ½ oz., oil of almonds ¼ oz.; melt together with a gentle heat, and add 6 drops of essence of lemon.

GANTS COSMETIQUES. These are white kid gloves, which have been turned inside out, and brushed over with a melted compound of wax, oil, lard, balsam, &c. The Peruvian Lip Salve, without the alkanet, may answer the purpose. *For softening the hands.*

FACE PAINTS. FARDS.

FINE CARMINE (prepared from cochineal) is used alone, or reduced with starch, &c. And also the coloring matter of safflower and other vegetable colors, in the form of pink saucers, &c.

ROUGE is prepared from carmine, and the coloring matter of safflower, by mixing them with finely levigated French chalk or talc, generally with the addition of a few drops of olive or almond oil. Sometimes fine white starch is used as the reducing ingredient. It is used in the form of powder, pomade, and *crépons*—the latter being pieces of crape imbued with the coloring matter. For common purposes vermilion is used; and it is sometimes prepared for this purpose by mixing it with a few drops of almond oil and of mucilage of tragacanth, placing the mixture in rouge-pots, and drying it by a very gentle heat.

ALMOND BLOOM. Boil 1 oz. of Brazil dust in 3 pints of distilled water, and strain; add 6 dr. of isinglass, 2 dr. of cochineal, 1 oz. of alum, and 3 dr. of borax; boil again, and strain through a fine cloth.—GRAY'S SUPPLEMENT.

FACE WHITES. One of the most innocent kind is prepared from Venetian talc, or French chalk, finely levigated. These are sometimes calcined, to increase their whiteness; but this diminishes their unctuosity and adhesiveness. Digestion with vinegar, and subsequent washing, are practised for the same purpose. Flake white (a fine variety of white lead) was formerly much used, but is now generally condemned as unsafe; it is also liable to become brown under certain circumstances. Pearl of bismuth white (magistery of bismuth*) is less injurious when pure, but is subject to the latter inconvenience. M. THENARD recommends oxide of zinc, with an equal weight of French chalk prepared by vinegar. Magnesia is said to be employed by the American ladies. White starch is used for the same purpose.

TOILET SOAPS, &c.

As the retail druggists and perfumers do not generally make their own soap in the first instance, it is only necessary to mention the means by which the soap, as it comes from the manufacturers, is prepared for the toilet.

SCENTED SOAPS, in general. Cut the best white curd soap (or, for some kinds, palm soap) into thin shavings, and place it in a copper vessel, with sufficient distilled water, and heat it by a water-bath till the whole is uniformly liquified. Let it cool to 135° F.; then add the coloring matter and perfumes. On the large scale, these additions may be mixed with the liquid soap at the maker's before it is poured into the frames. The quantity of perfume used must depend on the price at which it is to be sold.

ALMOND SOAP. To one hundredweight of the best hard white soap, melted as above, add 20 oz. of essential oil of bitter almonds. (Soap really made from expressed

* For this purpose, a little muriatic acid is added to the solution of the metal in nitric acid, and the magistery is precipitated by a small quantity of water; or the nitric solution is mixed with a weak solution of sea salt. Dr. Ure states that the precipitate thus acquires a more pearly lustre.

almond oil is, we apprehend, rarely met with in commerce.

SAVON AU BOUQUET. Melt 60 lb of white curd soap as above, and 8 oz. of oil of bergamot, 1½ oz. each of oils of cloves, sassafras, and thyme, ¾ oz. of neroli, and 14 oz., or q. s. brown ochre.

ROSE SOAP. Put into a copper vessel, placed in boiling water, 20 lb of white curd soap, and 30 lb of olive oil soap, both in thin shavings; add 5 lb of soft water, or rose-water; keep the heat below boiling till the soap is uniformly liquefied, then add 12 oz. of finely sifted vermilion, or enough to produce the required tint. Withdraw it from the fire, and when sufficiently cool add 3½ oz. of otto of roses, ½ oz. of oil of cloves, ½ oz. oil of cinnamon, and 2½ oz. of bergamot. For a cheaper article use less perfume.

WINDSOR SOAP. This is said to be made with lard. In France they use lard with a portion of olive or bleached palm oil. Dr. PEREIRA states that it is made with one part of olive oil to nine of tallow. But a great part of what is sold is probably only curd (tallow) soap, scented with oil of caraway and bergamot. The brown is probably colored with burnt sugar, or umber.

HONEY SOAP. White curd soap 1½ lb, brown Windsor soap ½ lb; cut them into thin shavings, and liquefy as directed above for scented soaps; then add 4 oz. of honey, and keep it melted till most of the water is evaporated; then remove from the fire, and when cool enough, add any essential oil. According to PIESSE the *honey soap* usually sold consists of fine yellow soap perfumed with oil of citronella.

FLOATING SOAPS. These are made by liquefying, as described above, 30 lb of oil soap with about 5 lb of water, and agitating the mixture, by a suitable wooden apparatus turned by a handle, till the froth arises to the top of the vessel. It is then put into the frames to cool. They are variously perfumed and colored.

TRANSPARENT SOAP. Cut fine white curd soap into thin shavings, and dry them with a gentle heat till they can be reduced to powder. Put 2 lb of this powder into a water-bath with 5 or 6 pints of rectified spirit of wine, and heat it gently (taking care that the water does not quite boil) till

the solution is complete; add the perfume, and pour into the frames. When cold, cut it into squares. They must be kept some time in a dry place before they can attain their full degree of transparency. By using a still, most of the spirit may be recovered for future use.

Wash Balls. *Savonettes.* These are made from various kinds of soap, usually with the addition of powdered starch, or hair powder, or of rice flour, together with perfuming and coloring ingredients. They are formed into spherical balls by taking a mass of the prepared soap in the left hand, and a conical drinking-glass with rather thin edges in the right. By turning the glass and ball of soap in every direction, the rounded form is soon given; when dry the surface is scraped to render it more smooth and even. One or two examples of this kind of soap will suffice.

Common or Lemon Wash Balls. Cut 6 ℔ of soap into very small pieces; melt it with a pint of water in which 6 lemons have been boiled. When melted withdraw the soap from the fire, and add 3 ℔ of powdered starch, and a little essence of lemon; knead the whole into a paste and form into balls into the desired size.

Cream Wash Balls. White curd soap 7 ℔, powdered starch 1 ℔; water or rose-water, q. s. Beat the whole together, and form into balls.—Gray's Supplement.

Camphor Wash Balls. White soap 1 ℔, spermaceti 1 oz., water q. s.; melt together and add 1 oz. of powdered camphor.

Mrs. Symond's Soap Paste, for the Hands. Best soft soap (from olive oil and potash if procurable) 16 oz., spermaceti 4 oz., best olive oil 1 oz., camphor ¼ oz., rectified spirit ½ oz., soft water 1 pint, essence of lemon ¼ oz., M. S. A. With 8 oz. of pumice-stone, powdered and sifted through fine book muslin, it forms sand soap paste.

Powdered Soap. Any of the hard soaps may be pulverized, if first cut into thin shavings, and kept at a gentle heat, till sufficiently dry. This process renders the soap more mild.

Shaving Powder. Melt together in a water-bath 1 ℔ of white soap with 1 oz of powdered spermaceti and ¼ oz. of chlorate of potash dissolved in a little water, or rose-water. Pour the liquefied soap into a shallow mould; when solidified shave it fine, and dry as above.

SHAVING PASTE. 1. Melt together 1 dr. each of spermaceti, white wax, and almond oil; beat it up with 2 oz. of the best white soap, and a little lavender or Cologne water.

2. Naples soap, beaten up with sufficient powdered soap to form a stiff paste.

3. White soft soap, 4 oz., powdered Castile soap 1 oz., oil of olives or almonds $\frac{1}{4}$ oz.

SHAVING LIQUID. Essence of soap. 1. White soap 3 oz., proof-spirit 8 oz., distilled water 4 oz., carbonate of potash 1 dr., essence of lemon q. s. Dissolve the soap without heat, and add the potash and essence.

2. (Italian essence of soap.) White curd or Windsor soap 10 parts, rectified spirit 34 parts, rose or orange-flower-water 34 parts. Digest with a gentle heat and filter.

3. Naples soap, or white soft soap, 16 oz., oil of olives $\frac{1}{2}$ oz., gum benzoin 1 dr., rectified spirit 24 oz. Digest. Rub a few drops on the beard, followed by warm water.

HAIR COSMETICS.

HAIR POWDER. The basis of hair powder is finely powdered starch. It is variously scented, and was formerly tinted with various colors. The plain and violet hair powders are now principally used. The latter is perfumed with orris powder, or essence of violets, usually with the addition of bergamot, &c. GRAY gives the following species for scenting hair powder: Powdered orris 1 lb, essence of bergamot 12 oz., oil of neroli 1 dr., musk 1 scruple. Hair powder is also perfumed with jessamine, roses, &c., by mixing the flowers with plain powder for 2 or 3 days, stirring the mixture twice or thrice a day, and then sifting out the powder from the flowers.

COMPOUNDS TO PROMOTE THE GROWTH OF THE HAIR.

POMADES FOR THE CURE OF BALDNESS.

1. DUPUYTREN'S POMADE. The recipe given by BATEMAN and RENNIE for this celebrated preparation, viz., almond oil, lard, suet, and essential oils, is remarkable as entirely omitting the active ingredient. It is probable that the preparation first employed by M. DUPUYTREN was more simple in its form than what he subsequently adopted, but *cantharides* was always the essential constituent. The first formulæ we met with was: Tincture of Cantharides (made according to the Paris Codex, 1 part of flies to 8 of proof-spirit) 1 part, lard 9 parts. The following are said more nearly to represent the compound in its improved and more elegant form. M. CAP prescribes: Beef marrow 2 oz., spirituous extract of cantharides (made by evaporating the above tincture) 8 gr., rose oil 1 dr., essence of lemon 50 drops. M. FONTAINE directs: Beef marrow 4 oz., calomel 2½ dr., extract of cantharides 18 gr., attar of roses 2 drops. But the following, by M. RECLUZ, is said to have been acknowledged by DUPUYTREN as the true formula: Beef marrow 6 oz., nervine balsam* 2 oz., Peruvian balsam 2 oz., oil of almonds

*This is made by melting together 4 oz. each of beef marrow and oil of mace, and adding 2 dr. of balsam of Tolu, and 1 dr. each of oil of cloves and camphor, dissolved in ½ oz. of rectified spirit.

1½ oz., extract of cantharides 16 gr.; melt the marrow and nervine balsam with the oil, strain, add the balsam of Peru, and lastly the extract, dissolved in a drachm of rectified spirit. M. GUIBOURT says that no better than the following can be used: Beef marrow 1 oz., nervine balsam 1 oz., rose oil 1 dr., extract of cantharides (dissolved in spirit) 6 gr. These pomades should be rubbed on the scalp once or twice a day for some weeks. If any soreness is produced, it should be less frequently applied.

2. POMADE CONTRE L'ALOPECIE. Fresh lemon-juice 1 dr., extract of bark (by cold water) 2 dr., marrow 2 oz., tincture of cantharides (as above) 1 dr., oil of lemon 20 drops, oil of bergamot 10 drops; mix. First wash the head with soap and water, with a little eau de Cologne, then rub it dry. Next morning rub in a small lump of pomade, and repeat it daily. In 4 or 5 weeks the cure of baldness is effected.—Dr. SCHNEIDER.

3. CAZENAVE'S REMEDY FOR BALDNESS. Beef marrow 1 oz., tincture of cantharides (as above) 1 dr., powdered cinnamon 1 dr. To be applied night and morning, the head being first washed with salt and water. Keep the hair short.

4. Dr. CATTELL'S is the same, substituting 10 drops each of oils of origanum and bergamot for cinnamon.

5. Beef marrow 1 oz., castor oil ½ oz., tincture of cantharides 1 dr., essential oil of bitter almonds and of lemon, each 12 drops.

6. Beef marrow 3 dr., almond oil 1 dr., sulphate of quinine 15 gr., otto 2 drops.—SOUBEIRAN.

7. Prepared lard 2 oz., white wax 2 dr.; melt together, remove from the fire, and add 2 dr. balsam of Tolu, 20 drops of oil of rosemary; and in chronic cases 1 dr. of tincture of cantharides.—Dr. NELIGAN.

8. Camphor 1 scruple, citrine ointment 2 dr., spermaceti cerate 6 dr.; mix. To be applied every night.

9. BATE'S *Unguentum Criniscum*. Laudanum 6 dr., bear's-grease 2 oz., honey ½ oz., powdered southernwood 3 dr., ashes of reed-root 1½ dr., oil of nutmeg 1 dr., balsam of Peru 3 dr.; mix. Let the bald part be first rubbed with an onion till it is red, then apply the ointment. It should be used daily, or oftener, for 5 or 6 weeks.

10. Box-leaves 2 oz., southernwood 2 oz., lard, marrow, or bear's grease 8 oz.; digest together by the heat of a water-bath, and strain.

11. *Bear's-grease.* The most approved consists of 2 parts of prepared bear's fat, with 1 of beef marrow, scented at pleasure. We have placed this, on the ground of common report, among the preparations which may possess some efficacy, but reserve the compounds usually sold under this name for the POMATUS. See below.

12. *Pommade Philocome.* Powdered cinchona ½ dr., oil of almonds 2 dr., beef marrow 6 dr., oil of bergamot 6 drops, balsam of Peru 20 drops; mix.—DOURVAULT.

LIQUID COMPOUNDS FOR THE CURE AND PREVENTION OF BALDNESS.

1. Dr. LOCOCK'S *Lotions.* Oil of mace (expressed oil of nutmeg) ½ oz., olive oil 2 dr., water of ammonia ½ dr., spirit of rosemary 1 oz., rose-water 2½ oz.; mix. (Mr. ASTLEY recommends the following modification: Oil of mace ¼ oz., olive oil 2 dr., oil of rosemary 4 drops; incorporate them carefully, then add gradually 3¼ oz. of rose-water, 2 dr. of solution of carbonate of ammonia, and 2 dr. of rectified spirit.)

2. Mr. ERASMUS WILSON'S. Eau de Cologne 2 oz., tincture of cantharides 2 dr., oil of rosemary and oil of lavender, of each 10 drops.

3. Mr. ACTON'S. Equal parts of rectified spirit, castor oil and eau de Cologne.

4. Mr. ACTON'S, *stronger.* Equal parts of honey-water and tincture of cantharides.

5. Tincture of cantharides 3 dr., acetate of copper 3 gr., oil of almonds and castor oil, of each a fluid ounce, with any essential oil to scent it. A small quantity to be applied to the roots of the hair every morning.

6. Vinegar of cantharides (Lond. Pharm.) ½ oz., eau de Cologne 1 oz., rose-water 1 oz.; mix.

7. Castor oil, lavender-water, and tincture of cantharides, in equal quantities.

8. *American Shampoo Liquor.* Rum 3 quarts, spirit of wine 1 pint, water 1 pint, tincture of cantharides ½ oz., carbonate of ammonia ½ oz., salt of tartar 1 oz. Rub it on, and afterwards wash with water. By omitting the salt of tartar, it nearly resembles balm of Columbia.

9. Dr. LANDERER'S. Bay leaves 2 oz., cloves ¼ oz., spirit of lavender 4 oz., spirit of thyme 2 oz.; digest for 6 days, filter, and add ½ oz. of ether. To be rubbed on every morning.

10. Put into a still 4 lb of honey, 12 handfuls of the tendrils of vine, and the same of rosemary tops; distil very slowly till the liquor begins to taste sour.

11. Dr. CATTELL'S. See WASHES FOR THE HAIR, below.

Note.—The above ointments and liquids require to be used for some weeks, in order to produce a decided effect, either in curing or preventing baldness. Those which contain cantharides in any form are the most active, and must be used with caution. They should be applied once or twice a day, according to the effect produced; but if the scalp becomes sore, their use must be omitted for a time, or longer intervals allowed, as the case may require. When employed to prevent the hair falling off, or becoming gray, they need not be applied so frequently as for baldness.

The following require no particular caution, being less active than the preceding:

POMATUMS, LOTIONS, &c.,

FOR EMBELLISHING, STRENGTHENING, AND CLEANSING THE HAIR.

Pomatums, or Pomades.

These are composed usually of animal fats, variously perfumed. The lard, veal fat, beef and mutton suet, bears' fat, and beef marrow, employed for this purpose, require to be prepared with great care. The following is, perhaps, the best mode: Cut the raw fat into pieces, carefully removing the fleshy and bloody portions of membrane, &c., and beat it in a marble mortar; melt it in a well-tinned vessel placed in boiling water, and strain the melted fat through a hair-sieve without pressure (reserving the residue to be heated again and pressed for more fat, to be used for commoner purposes). Keep the melted fat for some time gently warm, without disturbing it; remove any scum which may have arisen, and pour off the clear fat, taking care that none of the dregs or watery liquid which have subsided pass with it. A mixture of these fats forms the basis of many varieties of pomades. Sometimes a little white wax is added. A greater degree of

whiteness is said to be given by adding to the liquefied fat a few grains of citric acid. The same end is promoted by assiduously beating the pomade, while cooling, with a wooden spatula.

To perfume pomatums, various essential oils, &c., are added (see COMMON POMATUM); but the finer sorts are perfumed by infusing fresh flowers in the melted fats for some hours, and straining; or, in other cases, the simple pomade is thinly spread on plates of glass set in frames, and the fresh flowers stuck in the scored surface of the fat, changing the flowers daily till the pomatum is sufficiently perfumed. As these compounds can seldom be prepared to advantage by the retailer, a few varieties only require to be noticed here.

COMMON POMATUM. Mutton suet (prepared as above) 1 lb, prepared lard 3 lb; melt together in a water-bath, pour it into an earthen basin, and beat it assiduously with a wooden spatula. When sufficiently cool, add 2 oz., or q. s. of essence of bergamot, or of lemon, and continue the stirring till nearly cold.

ROSE POMATUM. Prepared lard 16 oz., prepared suet 2 oz.: melt with a gentle heat, and add 2 oz. of rose-water, and 6 drops of otto of roses. Beat them well together, and pour it into pots before it is too cold. For making jessamine, violet, and orange pomade, put the same quantity of water, and 1 dr. of the essence.

MARROW POMATUM. Beef marrow and beef suet, colored with a little annotto, may be employed for this and other yellow pomatums. For the perfumes employed for these and other pomatums, see *Essence for scenting Pomatums*, under PERFUMERY.

POMADE FOR BEAUTIFYING THE HAIR. Oil of sweet almonds a pint, spermaceti 1½ oz., purified lard 2 oz.; melt with a gentle heat; when nearly cold add any agreeable scent, and pour it into pots or wide-mouthed bottles.

BEARS' GREASE (ARTIFICIAL). Bears' grease is imitated by a mixture of prepared veal suet and beef marrow. It may be scented at pleasure; oil of lavender, with a very little oil of thyme, is sometimes used. The following are some of the compounds sold under this name:

1. Prepared suet 3 oz., lard 1 oz., olive oil 1 oz., oil of cloves 10 drops, compound tincture of benzoin 1 dr.; mix.

2. Lard 1 lb, solution of carbonate of potash 2 oz.

3. Olive oil 4 flasks, white wax 4 oz., spermaceti 2 oz.; scented with otto of roses and oil of bitter almonds.

GREEN BEARS' GREASE. Bears' grease digested with fresh walnut leaves, and strained. This is repeated with more leaves till the pomade is sufficiently colored; it is then scented with oil of rosemary, thyme, and bergamot.

GERMAN POMADE, FOR STRENGTHENING THE HAIR. Take 8 oz. of purified marrow, melt it in a glass or stoneware vessel, and add 1½ oz. of fresh bay leaves, 1 oz. of orange leaves, 1 oz. of bitter almonds, ½ oz. of nutmegs, ½ oz. of cloves, and 1 dr. of vanilla, all bruised; cover the vessel, and let the whole digest for 24 hours, with a gentle heat; strain while warm through linen, and stir it as it cools.

HARD OR ROLL POMATUM. 1. Suet 5 lb, white wax 8 oz., spermaceti 2 oz., oil of lavender, and essence of ambergris, each ½ oz.

2. Beef suet 16 oz., white or yellow wax 1 oz., with 1 dr. of oil of lavender or of bergamot.

3. Lard melted with one-third or half its weight of white wax, and poured into semi-cylindrical paper moulds when nearly set. This is sold under the name of *cosmetique*. It is sometimes colored to match the hair. See after HAIR DYES, below.

COLORED POMATUMS. The coloring matters employed are annotto, alkanet, marigold, carmine, indigo, cobalt, blue, umber, ivory black, &c.

CIRCASSIAN CREAM. Two flasks of oil, 3 oz. of white wax, 2 oz. of spermaceti, ½ oz. of alkanet root. Digest the oil with the alkanet till colored, strain, melt the wax and spermaceti with the oil, and when sufficiently cool add 2½ dr. of English oil of lavender, and ½ dr. of essence of ambergris.

CRYSTALLINE CREAM. Oil of almonds 8 oz., spermaceti 1 oz.; melt together; when a little cooled add ½ oz. or less of essence of bergamot, or other perfume; put it into wide-mouthed bottles, and let it stand till cold.

Camphorated crystalline cream may be made by using camphorated oil (Lin. Camphoræ) instead of oil of almonds.

Castor Oil Pomade. Castor oil 4 oz., prepared lard 2 oz., white wax 6 dr., essence of bergamot 2 dr., oil of lavender 20 drops, eau de Cologne ½ dr.; stir till cold.

Crystalline Castor Oil Pomade. Castor oil 16 oz., spermaceti 1¾ oz.; melt together, and when a little cool add 1 oz. of essence of bergamot, ½ dr. oil of verbena, ½ dr. oil of lavender; pour it into wide-mouthed bottles, and let it stand till cold.

Fox's Cream. Marrow pomatum 2 oz., oil of almonds 2 oz.; melt, and add while cooling, with constant stirring, essence of jessamine or of bergamot 2 dr.—Bateman.

HUILES ANTIQUES.

PERFUMED OILS FOR THE HAIR.

The basis of these oils is either almond oil, olive oil, or oil of ben; whichever is used should be perfectly fresh, and of the finest quality. The perfume is communicated in three ways: by infusing the flowers in the oil at a gentle heat; by placing layers of flowers alternately with folded cotton soaked in the oil, in proper frames, and pressing out the oil when sufficiently imbued with the odor of the flowers; or simply by adding essential oils, &c., to the fixed oil. An example of two of each method will be sufficient.

Oil of Roses, by Infusion. Heat in a water-bath 1 ℔ of virgin oil, and add 1 ℔ of picked fresh petals of Provence roses. Let these remain together in a water-bath for half an hour; then remove from the bath, and leave them together for 24 hours, stirring them twice during the time. Strain through a cloth, and express all the oil. To this oil add fresh roses, and proceed as before; repeating this for 5, 6, or 7 times, till the oil is sufficiently perfumed.

Oil of Jessamine, Perfumed with the Flowers. Fold pieces of white cotton cloth twice or four times; moisten them with fine olive oil, slightly pressing them, and place them in proper frames. Then place on the cloths a rather thick layer of freshly-gathered and dry jessamine flowers, carefully deprived of all green parts. In 24 hours carefully remove the flowers, and replace them by fresh ones, till the oil is sufficiently perfumed. The oil is then ex-

pressed. The same method is employed in preparing oils from other delicate flowers; as violets, lily of the valley, &c.

Oil of Roses, Common. Fine olive or almond oil 1 pint, otto of roses 16 drops. If required red, color the oil with alkanet root, and strain before adding the otto. For common sale, essence of bergamot, or of lemon, is often substituted, wholly or in part, for the more expensive otto.

Perfumed Oil of Bergamot, Lemon, Orange, &c. To oil of ben, or finest almond or olive oil, add essential oil of bergamot, lemon, &c., q. s. For common purposes, 1 dr. of the essential oil may be added to 16 oz. of oil. Some recipes, however, direct as much as 1½ oz. or 2 oz.

Oil of Ambergris and Musk. Ambergris 2 dr., musk ½ dr.; grind them together in a mortar, then with a small quantity of oil; add more oil to make up a pint, and let them stand together for 12 days, stirring them occasionally. Then decant or filter. Add ½ a pint of oil to the residue for an oil of second quality.

Common Oil of Musk, Oil of Benzoin, Oil of Storax, &c., may be obtained by mixing a strong tincture of these drugs with fine oil, agitating them frequently together, and after remaining some hours at rest, decanting the clear oil.

Huile Comogene. Mix equal parts of oil and spirit of rosemary with a few drops of oil of nutmeg. To be used daily.

Huile de Phenix. Clarified beef marrow 4 oz., lard 2 oz., oil of mace 4 oz.; melt together and strain through linen into a warm mortar; stir, and when it begins to cool add the following solution, and stir constantly till it is quite cold: oil of cloves, lavender, mint, rosemary, sage, and thyme, of each ½ dr.; balsam of Tolu 4 dr., camphor 1 dr., rectified spirit 1 oz. Put the spirit and balsam in a vial, and place it in warm water till the solution is complete; then add the camphor and essential oils.

Huile Philicome d'Aubril. Triturate together, without heat, equal parts of cold-drawn nut oil, almond oil, and prepared beef marrow, adding any essential oil as a perfume.

HUILE VERTE. Macerate 1 dr. of guaiacum with 1 ℔ of olive oil; strain, and add any essential oil to perfume it.—GRAY.

MARROW OIL. Clarified beef marrow, or marrow pomatum, with enough almond or olive oil to bring it to the desired consistence.

FLUIDE DE JAVA. This consists of beef marrow, white wax, fine olive oil, and essential oils at pleasure.

MACASSAR OIL. The oil made by the natives in the island is obtained by boiling the kernel of the fruit of a tree resembling the walnut, called in Malay *badeau*. The oil is mixed with other ingredients, and has a smell approaching to that of creasote. But the Macassar oil sold in this country has probably no relation to the above, except in name. The following is given by GRAY: Olive oil 1 ℔, oil of origanum 1 dr.; others add 1¼ dr. of oil of rosemary. The following French compound is probably named Macassar oil rather to denote its properties than from any resemblance either to the product of Macassar, or to the oil sold under this name in England:

HUILE DE MACASSAR DE NAQUET. Oil of ben 14 pints, nut oil 7 pints, spirit of wine 1 quart, essence of bergamot 3 oz., tincture of musk, 3 oz., spirit of orange (*esprit de Portugal*) 2 oz., otto of roses 2 dr., alkanet to color it. Digest them together with a gentle heat for an hour, and shake frequently for a week.

WASHES FOR THE HAIR.

VEGETABLE EXTRACT FOR CLEANSING AND STRENGTHENING THE HAIR. 1. Southernwood 2 oz., box leaves 6 oz., water 4 pints. Boil gently in a saucepan for ¼ of an hour, strain, and to each pint of the liquid add 2 oz. of spirit of rosemary and ½ dr. of salt of tartar (or 1 dr. of Naples soap).

2. Boil 1 ℔ of rosemary in 2 quarts of water, and add to the filtered liquor 1 oz. of spirit of lavender, and ¼ oz. of Naples soap, or salt of tartar.

3. Incinerate 2 oz. each of rosemary, maidenhair, southernwood, myrtle berries, and hazel bark; make a strong solution of the ashes, with which wash the hair at the roots every day. Keep the hair short.—Dr. CATTELL.

4. Borax 1 oz., powdered camphor ½ oz., boiling water 1 quart. When cold, filter for use. Damp the hair with it frequently.

Wash for Removing Scurf and Promoting the Curling of the Hair. 1. Beat up the yolk of an egg with 1 pint of clean rain-water. Apply it warm; and aftewards wash the head with warm water.

2. Lime water 1 pint, distilled vinegar ¼ of a pint; mix.

COMPOUNDS FOR STIFFENING THE HAIR.

Eau Collante. Dissolve without heat 8 oz. of clear gum in 2 lb of distilled or rose water, and filter through coarse filtering paper.

Bandoline, or Fixateur. Vegetable mucilage, with sufficient spirit to preserve it. Mucilage of quince seed is used; mucilage of picked Irish moss, carefully strained, is said to answer still better. But the following is employed by some London perfumers: Finest picked gum tragacanth, reduced to a coarse powder, 1 oz., rose-water 1 pint; put them into a wide-mouthed vessel, and shake them together daily for 2 or 3 days; then strain with gentle pressure through fine linen or cambric. If required to be colored, infuse cochineal in the water employed before making the mucilage. Another form is: Linseed (not bruised) 1 tablespoonful, water ½ pint; boil for 5 minutes, and strain.

Pommade Collante, for False Curls. Melt together in an earthen pipkin 24 oz. of fine Burgundy pitch and 8 oz. of white wax, and add 1 oz. of pomatum; remove from the fire, and add 4 oz. of brandy or other spirit, replace it on the fire till it boils slightly, then strain through linen, adding bergamot or other perfume, and cast it into moulds.

HAIR DYES.

Orfila's Hair Dye. Take 3 parts of litharge and 2 of quicklime, both in an impalpable powder, and mix them carefully. When used, a portion of the powder is mixed with hot water or milk, and applied to the hair, the part

being afterwards enveloped in oil-skin, or a cabbage-leaf, for 4 or 5 hours.

2. Litharge 2 parts, slaked lime 1 part, chalk 2 parts, all finely powdered, and accurately mixed. When required for use, mix the powder with warm water, and dip a brush in the mixture, and rub the hair well with it. After 2 hours, let the hair be washed.

3. Litharge 4½ oz., quicklime ¾ oz.; reduce to an impalpable powder, and pass it through a sieve. Keep it in a dry, close bottle. Wash the hair first with soap and water, then with tepid water; wipe it dry, and comb with a clean comb. Mix the dye in a saucer with hot water to the consistence of cream, and apply it to the hair, beginning at the roots. Place over it four folds of brown paper, saturated with hot water, and drain till cool; and over this an oil-skin cap and a nightcap. Let it remain from 4 to 8 hours, according to the shade required. When removed, oil the hair, but do not wet it for 3 or 4 days.

4. Chevallier's. Mix 5 dr. of fresh slaked lime with 1½ oz. of water, and strain through silk; put the milk of lime into a 4-oz. bottle. Dissolve 5 gr. of acetate of lead in sufficient water, and add enough slaked lime to saturate the acetic acid (a drachm or rather more), let it settle, pour off the supernatant liquor, wash the precipitate with water, and add it to the milk of lime.

5. Dr. Hanmann's. Levigated litharge 11 oz., powdered quicklime 75 oz., hair powder 37 oz.; mix. When used, a portion of the powder is mixed with warm water in a saucer, and applied to the hair with the fingers, taking care to cover the hair to the roots. Cover the whole with a sheet of cotton wadding moistened with water, and this with a folded cloth. Let it remain on for three hours; or better, for the night.

6. Warren's. Sifted lime 16 oz., white lead 2 oz., litharge in fine powder 1 oz.; mix well together, and keep dry. To dye *black*, mix a little powder with water to the consistence of cream. To dye *brown*, use milk instead of water. Apply with a small sponge to every hair.

Essence of Tyre. Grecian Water. Eau d'Egypt. Eau de China. These are solutions of nitrate of silver; in applying them it must be remembered that they stain the *skin* as well as the hair. Hence there is more difficulty

in applying than with the preceding; but they are considered to impart a finer color to the hair, with the disadvantage, however, of rendering it dry and crisp. The following are some of the most approved formulæ:

1. Dr. Cattell's. Nitrate of silver 11 dr., nitric acid 1 dr., distilled water 1 pint, sap green 3 dr., gum Arabic 1 dr.; mix.

2. Nitric acid 1 dr., nitrate of silver 10 dr., sap green 9 dr., mucilage 5 dr., distilled water 37½ fluid oz.

3. Silver 2 dr., iron filings ½ dr., nitric acid 1 oz., distilled water 8 oz. Digest, and decant the clear solution. To be carefully applied with a close brush.

4. Hydrosulphate of ammonia 1 oz., liquor of potash 3 dr., distilled water 1 oz.; mix. Apply this with a toothbrush for 16 or 20 minutes, then brush the hair over with the following: Nitrate of silver 1 dr., distilled water 2 oz., using a clear comb to separate the hair.

Pyrogallic Stain. Distil coarsely-powdered nutgalls in a retort, dissolve the solid acid which sublimes in a little hot water, add the solution to the acid liquid which passes over, separate the floating oil, shake the liquid with charcoal, filter, and add a little spirit.

La Forest's Cosmetic Wash for the Hair. Red wine 1 ℔, salt 1 dr., sulphate of iron 2 dr. Boil for a few minutes, and add common verdigris 1 dr.; leave it on the fire 2 minutes, withdraw it, and add 2 dr. of powdered nutgall. Rub the hair with the liquid; in a few minutes dry it with a warm cloth, and afterwards wash with water.

POMATUMS, OR COSMETIQUES, IN STICKS, FOR THE HAIR.

Black Pomatum, in sticks, for the eyebrows, whiskers, &c. Prepared lard melted with a third of its weight of wax in winter, or half in summer, is colored with levigated ivory-black, and strained through tammy, or any material which will permit the fine particles of ivory-black to pass through. Stir it constantly, and when it begins to thicken pour it into paper moulds.

Brown and Chestnut Pomatums are prepared in the same way, but colored with umber, &c. White, as Hard Pomatum.

EBONY POMATUM, in pots. Melt 4 oz., of white wax with 12 oz. of any kind of pomatum, and add 2 oz. of levigated ivory-black. Proceed as above, and pour into pots.

POMADE DE JEUNESSE. Pomatum mixed with magistery of bismuth. It is said to turn the hair black.—GRAY.

DEPILATORIES.

FOR REMOVING SUPERFLUOUS HAIRS.

These require caution, as they are apt to injure the skin. We have omitted those which contain sulphuret of arsenic (orpiment), as there is danger of its being absorbed, and the object can be accomplished without its use. The powders require to be kept in close bottles or boxes, and no more should be mixed with liquid than is required to be used at once.

1. Mix lime and water to a thick cream, and pass through the mixture 25 or 30 times its volume of sulphuretted hydrogen gas. When the gas escapes, stop the process. The pulpy mass is spread on paper, and applied for 12 or 15 minutes, and then washed off with a sponge and water. The only objection to this is its disgusting smell.

2. CHINESE. Quicklime 16 oz., pearlash 2 oz., liver of sulphur 2 oz. Reduce to a fine powder, and keep it in a close bottle. Use it as No. 4.

3. Mr. REDWOOD recommends a strong solution of sulphuret of barium, with sufficient powdered starch to form a paste; to be left on for a few minutes, then scraped off with the back of a knife.

4. BOUDET'S DEPILATORY. Crystallized hydrosulphate of soda 3 parts, quicklime in powder 10 parts, starch 10 parts; mix. To be mixed with water, and applied to the skin, and scraped off in two or three minutes with a wooden knife.

TEETH AND MOUTH COSMETICS.

TOOTH POWDERS.

General Directions.—The dry ingredients should be finely pulverized, and the whole well mixed; which is best effected by triturating the powders together, or agitating them in a bottle, and afterwards passing the whole through a sieve. Some ingredients are usually *levigated*, or ground with water, as prepared chalk, coral, &c. The tooth powders which contain acids and acid salts should not be frequently used. For children those only which contain very soft powders should be permitted; the heavy carbonate of magnesia is very suitable for them.

American Tooth Powder. Coral, cuttle-fish bone, dragon's blood, of each 8 oz., burnt alum and red sanders, of each 4 oz., orris 8 oz., cloves and cinnamon, of each ½ oz., vanilla 2 dr., rose-wood ½ oz., rose-pink 8 oz.

Antiseptic Tooth Powder. Prepared or precipitated chalk 2 oz., dry chloride of lime 10 gr., oil of cassia or of cloves 5 drops; mix. It may be colored, if preferred, by a little levigated bole.

Antiscorbutic Tooth Powder. Extract of rhatany ½ oz., prepared charcoal 2 oz., cinnamon ¼ oz., cloves ¼ oz.

Aromatic Tooth Powder. Calamus aromaticus 4 dr., charcoal 1 dr., soap 1 dr., oil of cloves 12 drops.—Pittschaft.

Asiatic Tooth Powder. Prepared coral 4 oz., Venetian red 3 dr., ochre 5 dr., pumice 5 dr., musk 1 gr.; mix. Or, bole 3 parts, chalk 2, ochre 1, pumice 1, musk to scent.

Cadet's, or Dr. Coombe's. Sugar 1 oz., charcoal 1 oz., Peruvian bark ½ oz., cream of tartar 1½ dr., cinnamon 24 gr.

Camphorated Chalk. Camphor (pulverized by the aid of a few drops of spirit) 1 oz., prepared or precipitated chalk

3 oz. Some makers put only 1 part of camphor to 7 of chalk.

COMPOUND CAMPHORATED TOOTH POWDER. Camphor 1 oz, precipitated chalk 2 oz., cuttle-fish bone ½ oz., myrrh 2 dr., borax 2 dr., lake or rose pink 1 dr., or q. s.

CARTWRIGHT'S DENTIFRICE. Prepared chalk 7 oz., orris 1 oz., Castile soap ½ dr.

CARABELLI'S. Cuttle-fish bone 1½ oz., prepared shells 1½ oz., cinnamon, orris, and lime-tree charcoal, of each 3 dr., vanilla 10 gr.

CHARCOAL, PREPARED. The charcoal made in iron cylinders, from willow, is preferred. It should be reduced to an impalpable powder, and kept from the air. Charcoal of areka nut is highly commended. That of the shells of cocoa-nuts is said to be used for the same purpose. Dr. HEIDER prefers the charcoal of the lime-tree.

CHARCOAL TOOTH POWDER (Gray). Prepared charcoal 1 oz., powdered chalk 3 oz.

CHARCOAL TOOTH POWDER (French). Prepared charcoal 1 oz., sugar 1 oz., oil of cloves 3 drops; mix.

CHARCOAL WITH BARK. Charcoal 1 oz., red cinchona 1 oz., powdered sugar ½ oz., with a few drops of some essential oil. See also RIGHINI'S, below.

CHARCOAL WITH BARK (French recipe). Charcoal 1 oz., Peruvian bark, ½ oz., oil of cinnamon, mint, or other oil, 2 drops, essence of ambergris 30 drops.

CHARCOAL WITH QUININE. Charcoal 1 oz., sulphate of quinine 2 to 4 gr., magnesia 4 to 8 gr., otto of rose (or other perfume) 2 drops.

CARBONIC DENTIFRICE (DESFORGES'). Willow charcoal 4 oz., cinchona bark 4 oz., cloves ¼ dr.

CIRCASSIAN DENTIFRICE (Dr. HALIFAX'S). Prepared hartshorn 2 oz., sulphate of potash 2 oz., cuttle-fish bone 8 oz., orris 4 oz., yellow sandal-wood 1 oz., rose-pink 3 oz., oil of rhodium 30 drops. Mix the dry ingredients, previously reduced to a fine powder, and add the oil of rhodium.

CORAL DENTIFRICE (*Poudre Dentifrice* of the French Pharmacopœia). Red coral, bole, cuttle-fish bone, of each 3 oz., dragon's blood 1½ oz., cochineal 3 dr., cream of tartar

4½ oz., cinnamon 6 dr., cloves 1 dr.; reduce separately, to powder, mix and grind on porphyry.

DESCHAMP'S ALKALINE DENTIFRICE. Venetian talc 4 oz., bicarbonate of soda 1 oz., carmine 4 or 5 gr., oil of mint (or other perfume) 15 drops.

DESCHAMP'S ACID DENTIFRICE. Venetian talc 4 oz., cream of tartar 1 oz., carmine 4 or 5 gr., oil as the last.

DESFORGES'. See CARBONIC DENTIFRICE.

DETERGENT TOOTH POWDER. Bicarbonate of soda 1 oz., powdered Castile soap ½ oz., sulphate of potash ½ oz., sugar of milk ½ oz., orris root 4 oz., oil of bitter almonds 4 drops. Colored at pleasure.

ELEPHANT'S (Mrs.) TOOTH POWDER. Bole 1 oz., myrrh, bark and orris, each ¼ oz. All to be finely powdered, and mixed.

FLORENTINE DENTIFRICE. Prepared shells 14 dr., orris 6 dr., cream of tartar 3 dr., lake to color.

FRENCH TOOTH POWDER. (See CORAL TOOTH POWDER, above; also GALVANIC, DESCHAMP'S, &c.) Peruvian bark, burnt crust of bread, and sugar, in equal proportions.

GALVANIC DENTIFRICE. Triturate 2 leaves of gold-leaf and 3 of silver with 2 dr. of sulphate of potash and 1 dr. of alum; then add white sugar 2 dr., common salt 1 dr., pellitory of Spain ½ dr., prepared hartshorn 1 oz., sulphate of quinine 10 grains; mix. Color with finest smalts (powder blue), rose pink, or lake. FOZEMBAS' recipe is: 2 leaves of gold, 2 of silver, alum 3 dr., salt 1½ dr., white sugar 1½ dr., pepper 15 gr., opium 5 gr., coral 3 dr., Peruvian bark 3 dr. Grind the gold and silver with the salt and alum, and add the last. For the *double* galvanic tooth powder, put twice the above quantities of gold, silver, alum, salt, pepper, and opium. The galvanic action of the metals is thought to stimulate the gums.

GERMAN TOOTH POWDER. Peruvian bark 6 dr., red sanders 2 dr., oil of cloves and of bergamot 3 drops.

GROSVENOR'S TOOTH POWDER. Prepared shells and coral, of each 12 oz., orris root 2 oz., oil of rhodium 6 drops.

HEMET'S DENTIFRICE. It is said to consist of cuttle-fish bone 6 oz., cream of tartar 1 oz., orris ½ oz.; mix.

JAMET'S. Orris 16 oz., magnesia 4 oz., pumice-stone 8 oz., cuttle-fish bone 8 oz., sulphate of quinine 4 oz., cascarilla 1 oz., sugar of milk 16 oz., oil of mint 1 oz., oil of cinnamon 2 dr., oil of neroli 1 dr., essence of ambergris 1 dr.

KEMMERER'S. Wood-soot 1½ oz., strawberry root ½ oz., and a few drops of eau de Cologne.

LAVENDER TOOTH POWDER. Crimson lake 1 dr., Chinese blue (or Turnbull's blue) a scruple: mix and add bicarbonate of soda ½ oz., cuttle-fish bone 2 oz., precipitated chalk 6 oz., oil of lavender 8 drops.

LARDNER'S TOOTH POWDER. See CHARCOAL TOOTH POWDER (gray).

LEFOULON'S TOOTH POWDER. Scurvy-grass, horseradish, guaiacum, cinchona, mint, pellitory root, calamus, rhatany, of each equal quantities. Reduce to an impalpable powder. A little calcined magnesia is sometimes added.

MAURY'S CARBONIC TOOTH POWDER. Charcoal 8 oz., cinchona 4 oz., sugar 8 oz., oil of mint ½ oz., oil of cinnamon ¼ oz., tincture of ambergris ½ dr.

METGES' TOOTH POWDER. Prepared chalk 3½ ℔, lake or rose-pink 1 ℔, orris 2 ℔, cream of tartar 12 oz., levigated pumice 1 oz., sugar 9 oz., oil of cloves 1 dr.

MIALHE'S RATIONAL DENTIFRICE. Sugar of milk 3 oz., pure tannin 3 dr., lake 1 dr., oil of mint 8 drops, oil of aniseed 8 drops, neroli 4 drops.

MYRRH DENTIFRICE. Myrrh 1 oz., cuttle-fish bone 4 oz., orris 3 oz.; mix.

NICHOL'S TOOTH POWDER. Cuttle fish-bone, prepared chalk, orris, of each 1 oz.; cassia ½ oz., myrrh ½ oz.; mix.

PALMER'S TOOTH POWDER. Prepared chalk 1 ℔, camphor 1 oz., orris 1 ℔, cuttle-fish bone 4 oz.; rose pink 1 oz.

PEARL DENTIFRICE. Precipitated chalk 16 oz., talc 8 oz., finest smalts ½ oz., or q. s. to give it a slight tint.

PELLETIER'S QUININE DENTIFRICE. Sulphate of quinine 4 gr., prepared red coral 1 oz., myrrh a scruple. For the coral may be substituted levigated bole 2 dr., precipitated chalk 6 dr.

REGNAUD'S DENTIFRICE. Calcined magnesia ½ oz., sulphate of quinine 8 gr., carmine (or cochineal) ½ dr., oil of peppermint 3 drops.

RHATANY TOOTH POWDER. Rhatany root 2 oz., cuttle-fish bone 4 oz., prepared chalk 8 oz., borax 1 dr.

RHIGINI'S CHARCOAL AND BARK. Charcoal 4 parts, yellow bark 1 part.

ROSE DENTIFRICE. Lake ½ dr., myrrh 2 dr., bicarbonate of soda 2 dr., orris 2 oz., cuttle-fish bone 2 oz., precipitated chalk 6 oz., otto of roses 16 drops; or it may be colored with rose pink to any desired shade.

RUSPINI'S DENTIFRICE. Cuttle-fish bone 8 oz., prepared hartshorn 2 oz., alum 1 oz., cream of tartar 2 oz., orris 1 oz., oil of rhodium 6 drops.

RUSSIAN TOOTH POWDER. Peruvian bark 2 oz., orris root 1 oz., sal ammoniac ½ oz., catechu 6 dr., myrrh 6 dr., oil of cloves 6 or 8 drops.

SAUNDERS' DENTIFRICE. Prepared chalk 2 oz., cuttle-fish bone 1 oz., orris 1 oz., myrrh ½ oz., sulphate of quinine 10 gr.

Dr. SCHOEPF'S TOOTH POWDER, *against mercurial salivation.* Alum 2 scruples, cinchona bark 1 oz.

VIOLET TOOTH POWDER. Orris root 2 oz., cuttle-fish bone 4 oz., precipitated chalk 12 oz., bicarbonate of soda ½ oz., essence of violets 1 dr., pure percyanide of iron and crimson lake or rose pink, enough to give it a pale violet color.

TOOTH PASTES.

Any of the above tooth powders may be formed into a paste with honey, clarified honey, or honey of roses. A little perfumed spirit may be added. A common objection to these pastes or electuaries, is their liability to fermentation, or effervescence. Some makers keep the paste in the bulk for a considerable time, till the effervescence has completely subsided, and then put it up in pots for sale. Others heat the honey, stir in the powders, and keep the mixture warm till any effervescence produced by the action of the acidity of the honey on the cretaceous powder has subsided. It would perhaps be preferable in all cases to use the *prepared*

honey (see Mel Preparatum, Pocket Formulary) for these purposes. Electuaries of this kind are termed by the French *opiats*, although they may contain no opium in any form.

Coral Tooth Paste. *Opiat Dentifrice.* 1. Prepared coral 5 oz., cream of tartar 3 oz., cuttle-fish bone 3 oz., cochineal ½ dr., Narbonne honey 16 oz.—Desforges.

2. *Opiat Dentifrice Rouge.* Prepared coral 8 oz., cochineal 1 oz., cinnamon 2 oz., alum 3 dr., honey 20 oz., water 1 oz.; triturate the cochineal with the alum and water, add the honey, then the coral and cinnamon; leave the whole for 24 hours, or till the effervescence has subsided; then rub it with a few drops of oil of cloves, or other aromatic oil, and put it into covered pots for sale.

Dyon's Charcoal Paste. Triturate ½ dr. of chlorate of potash with ½ oz. of mint-water, and add gradually 1 oz. of powdered charcoal.

Metges' Tooth Paste. Metges' tooth powder (as above) 48 oz., Narbonne honey 32 oz., syrup 64 oz., cochineal 1 oz., alum 1 oz., water 4 oz.; triturate the cochineal and alum with the water, add the honey and syrup, and lastly the powder.

Pelletier's Odontine. This is said to consist of magnesia and butter of cacao, aromatized with some essential oil.

Rose Tooth Paste. Cuttle-fish bone 3 oz., prepared or precipitated chalk 2 oz., orris 1 oz., lake or rose pink to give it a pale rose color, otto of roses 16 drops, honey of roses q. s.

Red or Cherry Paste. See Coral Paste, No. 2.

Rosemary Paste. Levigated bole 4 oz., myrrh 1 oz., oil of rosemary 2 dr. (dissolved in 1 oz. rectified spirit), clarified honey q. s.

Saline Tooth Paste. Sulphate of potash 1 oz., bay salt ½ oz., clarified honey q. s., eau de Cologne 2 dr. (or essence of ambergris 30 drops).

Vanilla Tooth Paste (French). Charcoal 1 oz., white honey 1 oz., vanilla sugar 1 oz., Peruvian bark ½ oz., and a few drops of any essential oil. The vanilla sugar may be made by triturating a drachm of saturated tincture of vanilla

with 1 oz. of pure sugar, and drying the mixture with a gentle heat.

WHITE TOOTH PASTE. 1 (French). Orris, sal ammoniac, cream of tartar, of each 2 oz., tincture of cinnamon and tincture of vanilla, of each ½ oz., oil of cloves 60 drops, clarified honey and syrup to form a paste.

2. Precipitated chalk 4 oz., sulphate of potash ½ oz., prepared honey sufficient to form a paste; to be flavored with a few drops of otto of roses or oil of cinnamon, &c.

LIQUID PREPARATIONS FOR THE TEETH AND GUMS.

ASTRINGENT TINCTURE FOR THE TEETH AND GUMS.

1. Borax, alum, bay salt, of each a dr., spirit of camphor, tincture of myrrh, of each 1 oz., spirit of scurvy-grass (or of horseradish) 4 oz., tincture of rhatany 2 oz.; mix, and shake occasionally for a day or two, then filter. A teaspoonful in a wineglassful of water, to rinse the mouth after cleaning the teeth, or at any other time.

2. Tannin 1 dr., rose-water 4 oz., spirit of wine 2 oz., spirit of scurvy-grass (or of horseradish) 2 oz., essence of bitter almonds a few drops.

ODORIFEROUS TINCTURE OF MYRRH. 1. Choice Turkey myrrh 3 oz., eau de Cologne a quart; digest for 7 days, and filter.

2. To 18 fluid oz. of tincture of myrrh add 2 oz. of essence of Cologne. (See PERFUMERY, p. 221.) If the tincture should not be quite clear, add a few gr. of burnt alum, shake frequently, and filter in a day or two.

BORATED TINCTURE OF MRRYH. 1. Myrrh 1 ℔, eau de Cologne 16 ℔, borax 1 ℔, distilled water 3 ℔, syrup 3 ℔, essence (or tincture) of roses 6 dr., rhatany root 4 oz.; digest for 10 or 12 days, and filter.—Mr. COCKLE.

2. Borax 1 oz., shellac ½ oz., myrrh 2 oz., spirit of camphor 2 oz., honey of roses 2 oz., rectified spirit a pint, Cologne essence 2 dr., orange-flower or rose-water 4 oz.; digest for a few days in a warm place, shaking occasionally, and filter.

3. Borax 1 oz., shellac ½ oz., water 8 oz.; boil together to 4 oz., and add spirit of scurvy-grass a pint, camphor ½ oz., myrrh 2 oz.; digest and filter.

Antiscorbutic Elixir. Chinchona 3 oz., guaiacum 5 oz., pellitory 3 oz., orange-peel 2 dr., cloves 5 dr., saffron ½ dr., benzoin 2 dr., spirit of wine or brandy 32 oz.; digest and filter.—Desforges.

Desforges' Extract of Pellitory. Pellitory root 5 oz., cinchona 1 oz., benzoin 1½ dr., essence of peppermint 3 dr., brandy 1 quart.

Elixir of Roses. Cloves 1 dr., cinnamon 3 oz., ginger 2 oz., spirit of wine 2½ pints, oil of orange-peel 1 dr., otto of roses 15 drops, essence of peppermint 1 oz.; mix, digest for 15 days, and filter.

Lefandiniere's Elixir. Rasped guaiacum-wood ½ oz., pellitory 1 dr., nutmegs 1 dr., cloves ½ oz., oil of rosemary 10 drops, oil of bergamot 4 drops, brandy 1 pint; macerate for a fortnight, and filter.

Eau de Bottot. 1. Aniseed 4 oz., cinnamon 1 oz., cloves 1 oz., cochineal 2 dr., oil of mint 2 dr., spirit of wine or brandy 8 lb; macerate 8 days, and filter.

2. Tincture of cedar-wood 1 pint, tincture of myrrh 1 oz., mixed with the following essential oils: of peppermint ½ dr., of spearmint ¼ dr., of cloves 10 drops, of roses 10 drops.—Piesse.

Eau Dentifrice de Stahl. Spirit of wine or brandy 2 gallons, rose-water 3 quarts, pellitory 5 oz., cypress root 3 oz., tormentil 3 oz., balsam of Peru 3 oz., cinnamon 5 dr., goats' rue 1 oz., rhatany 1 oz., macerate for 6 days, shaking it occasionally; let it rest for 24 hours, and pour off the clear. Add to the clear liquor, oil of mint 1½ dr., cochineal 4 dr.; in 3 or 4 days filter.

Eau du Dr. O'Meara. It is a tincture of pellitory, vetiver, cloves, orris, and coriander, with creasote, &c.

Bories' Odontalgic Elixir. Pellitory root 2 oz., simple spirit of lavender 16 oz., muriate of ammonia ½ dr.; digest 24 hours, and filter.

Greenough's Tincture. Bitter almonds 2 oz., Brazil-wood ½ oz., cinnamon ½ oz., orris root ¼ oz., cochineal, alum, salt of sorrel, each 1 dr.; spirit of wine 32 fluid oz., spirit of scurvy-grass 1 oz.

Hudson's Preservative. Tincture of myrrh, tincture of bark, cinnamon-water, of each 3 oz.; arquebusade-water 1 oz., powdered gum ½ oz.

Cheltenham Dental Tincture. Camphor 4½ oz., myrrh 2 oz., bark 5 oz., rectified spirit 36 fluid oz., distilled water 8 oz.

Lefoulon's Elixir for the Teeth. Fresh roots of horse-radish, fresh leaves of scurvy-grass, and of mint, of each 6 dr.; guaiacum, cinchona, pellitory, calamus, and rhatany, each 5 dr.; proof spirit 1 quart; macerate 16 days, and strain.

Eau de Madame de la Vrilliere pour les Dents. Cinnamon 2 oz., cloves 6 dr., fresh lemon-peel 1½ oz., dried rose-petals 1 oz., scurvy-grass 8 oz., spirit 3 ℔; macerate 24 hours, and distil in a water-bath.

Ruspini's Tincture. Orris 8 oz., cloves 1 oz., spirit 32 fluid oz., essence of ambergris 1 oz. (or ambergris 1 scruple); macerate 14 days, and filter.

French Elixir for the Teeth. Rose-water 16 oz., spirit of scurvy-grass 2 oz., tincture of galbanum 1 oz.; color with cochineal.

Alkaline Lotion, for preventing injury to the teeth from acid medicines. Bicarbonate of soda 4 dr., distilled water 8 oz., eau de Cologne 2 dr., aromatic spirit of ammonia 1 dr. The mouth to be rinsed out with the lotion *immediately* after swallowing any medicine containing an acid.

Lotion of Chlorinated Soda, for purifying the breath, cleansing the mouth, removing unpleasant odors, &c. Liquid chlorinated soda 1 oz., distilled water 19 oz.; mix. A teaspoonful in a glass of water. The same direction applies to most of the above.

STRONGER TINCTURES, SOLUTIONS, OR ESSENCES, FOR TOOTHACHE.

These are applied by moistening a little cotton-wool or lint with the liquid, and introducing it into the cavity of the decayed and aching tooth. Where there is no cavity, they are sometimes applied to the gums surrounding the affected tooth. Most of them are stated by their several inventors or patrons to give "immediate relief." The cavity should be dried with lint before applying the remedy.

1. M. PIESTE'S. Water of ammonia, with half the quantity of tincture of opium. Applied as above.

2. Creasote 1 dr., spirit of camphor 2 dr., laudanum 1 dr. Creasote is also used alone; so is *carvacrol*, a liquid of similar properties. LAENNEC prescribes 1 part of creasote and 10 of alcohol. See also No. 14.

3. M. COTTEREAU'S. Ether saturated in the cold with camphor, and then a few drops of ammonia added.

4. Mr. BLAKE'S. Finely-powdered alum 1 dr., spirit of nitric ether 7 dr.

5. *Paraguay-roux, or Compound Tincture of Para Cress.* Flowers of Para Cress 4 parts, Italian elecampane (*Inula bifrons*) 1 part, pellitory root 1 part, rectified spirit 8 parts; macerate 14 days, express, and filter.

6. Mr. BRANDE'S *Tincture.* Bruised pellitory ½ oz., camphor 3 dr., opium 1 dr., oil of cloves ½ dr., rectified spirit 6 oz.; digest for 10 days, and strain.

7. Pellitory, ginger, cloves, camphor, of each 1 oz.; tincture of opium 4 oz., spirit of wine 16 oz.; macerate for 8 days, and strain.

8. Camphor 1 dr., ether 4 dr.; dissolve.

9. Camphor 2 dr., chloroform 1 dr., spirit of sal volatile ½ dr.

10. Opium 2 oz., mastic 1 oz., balsam of Tolu 1 dr., camphor 1 oz., oil of cloves 1 dr., rectified spirit 16 fluid oz., oil of bitter almonds 8 drops.

11. BOERHAAVE'S *Odontalgic.* Rectified spirit 1 oz., camphor ½ oz., opium 1 scruple, oil of cloves 80 drops.

12. LEMAZURIER'S *Odontalgic.* Cherry-laurel water 2 oz., acetate of morphia 1 gr. Wash the mouth with warm water, to a glass of which a few drops of this mixture have been added.

13. Oil of rosemary 2 oz., tincture of galbanum 1 oz.; mix. Cotton wet with this is to be introduced into the ear.

14. RIGHINI'S. Alcohol 4 dr., creasote 6 dr., tincture of cochineal 2 dr., oil of peppermint 3 drops.

15. Mr. DRUITT'S. Tannin 20 gr., mastic 5 gr., ether 2 dr. Wash the mouth with warm water containing a little carbonate of soda; lance the gums, and apply the tincture to the cavity of the tooth, on cotton.

16. Mr. TOMES recommends a solution of mastic in chloroform. The mastic serves to retain the chloroform, but the latter may be used alone on cotton or lint. Mr. BEATSON uses a solution of copal in chloroform.

17. Chloroform and white of egg, equal parts, digested for 4 hours; then applied on lint.

HENBANE FUMIGATION FOR TOOTHACHE. A popular remedy is to throw henbane seed on hot cinders, inverting a cup over them to receive the smoke and empyreumatic oil produced. The cup is then filled with hot water, and the steam conveyed to the affected side of the mouth. Dr. DOWNING'S Aneuralgicon would probably prove a more effective means of applying remedies of this kind.

PILLS, OR PASTES, FOR TOOTHACHE.

MASSES ODONTALGIQUES.

1. DE HANDEL'S. Opium 12 gr., camphor 24 gr., cajeput oil 4 drops, tincture of cantharides 4 drops, extract of henbane and of belladonna, of each 24 gr.; distilled water of opium q. s.

2. VOGLER'S. Powdered opium 1 oz., mastic 2 dr., sandarach 2 dr., dragon's blood ½ dr., oil of rosemary 8 drops, spirit to form a paste. To be applied near the affected tooth.

3. Powdered alum 1 dr., powdered mastic ½ dr., spirit of nitric ether q. s. to form a paste.

4. RUST'S. Opium 5 gr., oil of cloves 3 drops, extract of henbane 5 gr., extract of belladonna 10 gr., powdered pellitory sufficient to form a paste.

CEMENTS, &c., FOR STOPPING THE CAVITIES OF TEETH.

THESE are harder than the preceding, and intended to remain in the tooth for an indefinite time. In all cases the cavity should be previously cleared from all extraneous matters, and wiped perfectly dry with a piece of lint or blotting-paper.

1. SOUBEIRAN'S. Powdered mastic and sandarach, of each 4 dr.; dragon's blood 2 dr., opium 15 gr.; mix with sufficient rectified spirit to form a stiff paste. A solution of mastic, or of mastic and sandarach, in half the quantity of alcohol, is also used, applied with a little cotton or lint.

2. Sandarach 12 parts, mastic 6 parts, amber in powder 1 part, ether 6 parts. Applied with cotton. Or simply a paste of powdered mastic and ether. Or a saturated ethereal solution of mastic, applied with cotton.

3. TAVEARE'S CEMENT is made with mastic and burnt alum. BERNOTH directs 90 parts of powdered mastic to be digested with 40 of ether, and enough powdered alum added to form a stiff paste.

4. Gutta-percha, softened by heat, is recommended. Dr. ROLLFS advises melting a piece of caoutchouc at the end of a wire, and introducing it while warm.

5. GAUGER'S CEMENT. Put into a quart bottle 2 oz., of mastic and 3 oz of absolute alcohol; apply a gentle heat by a water-bath. When dissolved add 9 oz. of dry balsam of Tolu, and again heat gently. A piece of cotton dipped in this viscid solution becomes hard when introduced into the tooth, previously cleansed and dried as above.

6. Mr. ROBINSON'S. After washing out the mouth with warm water, containing a few grains of bicarbonate of soda, and cleaning the cavity as above directed, he drops into it a drop of collodion, to which a little morphia has been added, fills the cavities with asbestos, and saturates with collodion, placing over all a pledget of blotting-paper.

7. OSTERMAIER'S CEMENT. Mix 12 parts of dry phosphoric acid with 13 of pure and pulverized quicklime. It becomes moist in mixing, in which state it is introduced into the cavity of the tooth, where it quickly becomes hard. (In some hands this has failed, from what cause we are not aware.) The acid should be prepared as directed under Trade Chemicals (Acid, Phosphoric).

8. SILICIA. This name has been given to a mixture of Paris plaster, levigated porcelain, iron filings, and dregs of tincture of mastic, ground together.

9. WIRTH'S CEMENT. It is said to consist of a viscid alcoholic solution of resins, with powdered abestos.

10. METALLIC CEMENT. Amalgams for the teeth are made with gold or silver, and quicksilver, the excess of the latter being squeezed out, and the stiff amalgam used warm. Inferior kinds are made with quicksilver and tin, or zinc. A popular nostrum of this kind is said to consist of 40 gr. of quicksilver and 20 of fine zinc filings, mixed at the time of using. Mr. EVANS states that pure tin, with a small portion of cadmium, and sufficient quicksilver, forms the most lasting and least objectionable amalgam. The following is the formula: Melt 2 parts of tin with 1 of cadmium, run it into an ingot and reduce it to filings. Form these into a fluid amalgam with mercury, and squeeze out the excess of mercury through leather Work up the solid residue in the

hand, and press it into the tooth. Or, melt some beeswax in a pipkin over the fire, throw in 5 parts of cadmium, and, when melted, add 7 or 8 parts of tin in small pieces; pour the melted metals into an iron or wooden box, and shake them till cold, so as to obtain the alloy in a powder. This is mixed with $2\frac{1}{2}$ or 3 times its weight of quicksilver in the palm of the hand, and used as above.

Another cement consists of about 73 parts of silver, 21 of tin and 6 of zinc, amalgamated with quicksilver. An amalgam of copper is said to be sometimes used. But this class of stoppings is altogether disapproved of by other authorities. Pure leaf-gold seems the least objectionable.

11. MARMORATUM. Finely-levigated glass, mixed with tin amalgam.

12. POUDRE METALLIQUE. The article sold under this name in Paris appears to be an amalgam of silver, mercury, and ammonium, with an excess of mercury, which is pressed out before using it.

13. FUSIBLE METAL. Melt together 8 parts of bismuth, 5 of lead, 3 of tin, and $1\frac{1}{2}$ or 1.6 of quicksilver, with as little heat as possible.—CHAUDET.

PASTE FOR DESTROYING THE SENSIBILITY OF THE DENTAL PULP PREVIOUS TO STOPPING. Arsenious acid 30 grains, sulphate of morphia 20 grains, creasote q. s. (Unsafe; it is only inserted by way of warning against what may prove an unsuspected cause of mischief.)

PIVOTS FOR ARTIFICIAL TEETH. An alloy of platinum and silver.

SPRINGS FOR ARTIFICIAL TEETH. Equal parts of copper, silver, and palladium.—CHAUDET.

[For Cachou Aromatisé, and other compounds for sweetening the breath, see PERFUMERY.]

EARACHE, SIMPLE CURE FOR. Take a common tobacco-pipe, place a wad of cotton in the bowl, drop upon it 8 or 10 drops of chloroform, and cover with another wad of cotton; place the stem to the affected ear, then blow into the bowl, and in many cases the pain will cease almost immediately.—AMERICAN JOURNAL.

BEVERAGES, DIETETIC ARTICLES AND CONDIMENTS.

BEVERAGES, AND POWDERS FOR PREPARING THEM.

We have placed here such beverages as are rather employed as a refreshing luxury than either medicinally or as regular articles of diet. Wines, spirits, &c., are necessarily excluded. The medicinal mineral waters will be found elsewhere.

Ginger Beer. 1. Infuse 3 oz. of bruised ginger in 4 gallons of boiling water till cold. Strain through tammy or flannel. Dissolve in the liquor 5 lb of loaf sugar, and add half a pint of solid yeast, and 2½ oz. of cream of tartar. In cold weather it will be necessary to set the cask near the fire, so as to excite brisk fermentation. As soon as this subsides, rack off the clear liquor, return it into the cask previously washed out, and allow it to work for a day or two longer. Then draw it off and bottle it.—Mr. Donovan.

2. Ginger sliced 1 oz., dried orange-peel ½ oz.; tie them in a bag, and boil with 16 lb of water, and strain; add ¾ of an oz. of tartaric or citric acid, 25 drops essence of lemon, and 24 oz. of loaf sugar. When sufficiently cool, add 2 tablespoonfuls of fresh yeast; let it work for 12 hours and bottle it.

3. Ginger sliced ¾ oz., essence of lemon (rubbed with sugar) ½ dr., lump sugar 12 oz., boiling water 8 lb; infuse till cold, and strain. Ferment as above, with 3 or 4 spoonfuls of yeast, and bottle.

4. Boil 2½ oz. of bruised ginger and 3 lb of sugar in 3½ gallons of water for 20 minutes; put into a large pan 1 oz. cream of tartar, and the juice and rind of two lemons; pour the boiling liquor over them, and stir the whole well together; when milk-warm add ¼ pint of good ale yeast, cover it, and let it work for 2 or 3 days, skimming it frequently; then strain it through a jelly-bag into a cask, add ½ pint of brandy, bung down close, and in 2 or 3 weeks bottle it in the usual way.

5. Boil 22 oz. bruised ginger in 3 gallons of water for ½ an hour; add 20 lb of white sugar, 18 oz. of lemon-juice, 1 lb of honey, and 15 gallons of water, and strain through a cloth. When cold add the white of 1 egg, and ½ oz. of essence of lemon; after standing 4 days, bottle, and lay the bottles in a cellar for 3 weeks.

6. Prepare a clear decoction or infusion of ginger with sugar and lemon as above; but instead of fermenting it with yeast, charge it strongly with carbonic acid gas by means of a machine.

7. *Imperial Pop.* Cream of tartar 3 oz., ginger 1 oz., white sugar 24 oz., lemon-juice 1 oz., boiling water a gallon and half; when cool, strain, and ferment with 1 oz. of yeast, and bottle.

Girambing, or Limoniated Ginger Beer. 1. Boil 4½ oz. of ginger with 11 quarts of water; beat up 4 eggs to a froth, and add them with 9 lb of sugar to the preceding. Take 9 lemons, peel them carefully, and add the rind and juice to the foregoing. Put the whole into a barrel, add 3 spoonfuls of yeast, bung down the barrel, and in about 12 days bottle it off. In 15 days it will be fit for drinking; but it improves by keeping.

2. To 10 gallons of water add 11½ lb of loaf sugar, and the whites of 10 eggs well beaten; boil till the scum rises, and add 6 oz. of bruised ginger; boil for 20 minutes, then pour the hot liquor on the rinds of 12 lemons thinly peeled; when cold, put into a barrel the juice of 12 lemons, 1 oz. of isinglass, a gill of brandy, and a spoonful of yeast, and fill the barrel with the liquor. In a fortnight it will be ready to bottle.

Ginger Beer Powders. Fine powder of Jamaica ginger 4 or 5 dr., bicarbonate of soda 3½ oz., refined sugar in powder 14 oz., essence of lemon 30 drops; mix, and divide into 5 dozen powders. (Or 4 to 5 gr. of ginger, 28 of bicarbonate of soda, 112 of sugar, and ½ drop of essence of lemon, in each powder.) In the other powder put 32 gr. of tartaric acid; or 35 gr. if a more decidedly acidulated beverage is required. Or from 30 to 33 gr. of citric acid.

[Other formulæ are also in use. Dr. Pereira gives the following: Bicarbonate of soda 30 gr., white sugar 1 dr., powdered ginger 5 gr., in each blue paper; and 25 gr. of tartaric acid, in each white paper. This is less agreeable, but perhaps more friendly to the stomach, than when the

acid is in slight excess. The following is from the Pharmaceutical Journal: Sugar 2 dr., sesquicarbonate of soda 2 scruples (misprinted 2 drachms in vol. 3), ginger 4 or 5 gr., essence of lemon 1½ or 2 drops, in each blue paper; with 35 gr. of tartaric acid.]

GINGER BEER POWDER, IN ONE BOTTLE. (The soda, acid, and sugar must be very carefully dried, separately, and at a temperature not exceeding 120°.) Fine powder of Jamaica ginger 4 or 5 dr., bicarbonate of soda 3½ oz., double refined sugar 14 oz., essence of lemon 30 drops, tartaric acid 4½ oz. The acid and soda should not be too finely powdered. Mix the powders, recently dried, in a warm mortar, and immediately put the mixture into dry bottles, and cork securely. A measure holding 3 dr. should accompany each bottle.

LEMON JUICE, factitious. Dissolve 4 oz. of citric acid in 3 pints of water, with 8 drops of essence of lemon (rubbed with the acid, or dissolved in a little spirit or tincture of fresh lemon-peel). After standing a few days filter it, and keep it in well-closed bottles.

ORANGE JUICE, factitious. Citric acid 1 oz., water 2 pints, oil of orange-peel 4 drops, tincture of orange-peel ½ oz. As the last.

KING CUP; OR LEMON DRINK WITHOUT ACID. 1. Pour a quart of cold water on the thin peel of 1 or 2 lemons; let them infuse 6 or 8 hours; then strain.—Mr. BRANDE.

2. Pour a pint of boiling water on the outer rind of one lemon, a small piece of dried orange-peel, and a moderate-sized lump of sugar.

LEMONADE, ACIDULATED (NOT AERATED). 1. Fresh lemon juice 4 oz., fresh lemon-peel (thinly peeled) ½ oz., white sugar 4 oz., boiling water 3 pints. Strain when cold.—Mr. BRANDE.

2. *Imperial.* Cream of tartar 1½ dr., a slice of thin lemon-peel, a lump of sugar; pour on them a quart of boiling water. Strain when cold. To be taken as a cooling drink.

3. *Common.* Cut 2 lemons into slices, add 2 oz. of sugar, and pour on them a quart of boiling water. It is sometimes made with cold water.

4. *French.* Syrup of citric acid 2 oz., water a quart, spirit of lemon-peel a teaspoonful.

5. Juice and the thin peel of 1 lemon, citric acid 1 dr., sugar 3 oz., boiling water a quart. It may be varied by substituting for the sugar syrup of raspberries, or of other fruits.

AERATED OR EFFERVESCING LEMONADE. This may be made by putting into each bottle (soda water bottle) 1 oz. or 1½ oz. of syrup of lemons, and filling it up with simple aerated water from the machine. (The syrup is made by dissolving 30 oz. of lump sugar in 16 oz. of fresh lemon-juice, by a gentle heat. It may be aromatized by adding 30 or 40 drops of essence of lemon to the sugar; or by rubbing part of the sugar on the peel of 2 lemons; or by adding to the syrup an ounce of a strong tincture of fresh lemon-peel; or of the distilled spirit of the same.)

EFFERVESCING LEMONADE, WITHOUT A MACHINE. Put into each bottle 2 dr. of sugar, 2 drops of essence of lemon, ½ dr. bicarbonate potash, and water to fill the bottle; then drop in 35 or 40 gr. of citric or tartaric acid in crystals, and cork immediately, placing the bottles in a cool place, or preferably, in iced water. Mr. BARTLETT recommends 2 scruples of sesquicarbonate of soda, 2 dr. of sugar, 4 drops of essence of lemon, and half a pint of water; lastly, a dr. of tartaric acid in crystals. Care must be taken to avoid accidents from the bursting of the bottles. Another form is: Into a soda-water bottle nearly filled with water, put 1 oz., of sugar, 2 drops of essence of lemon (dropped on the sugar), 20 gr. of bicarbonate of potash in crystals; and, last'y, 30 to 40 gr. of citric acid, also in crystals. Cork immediately.

MILK LEMONADE. Dissolve 1½ ℔ of sugar in a quart of boiling water, add ¼ pint of fresh lemon-juice, and the same of sherry; and, lastly, two-thirds of a pint of cold milk. Stir together, and strain.

DRY LEMONADE, OR ACIDULATED LEMONADE POWDER. Citric acid ¾ oz., refined sugar 8 oz., essence of lemon 36 drops. Some recipes direct a larger quantity of acid, others a much larger proportion of sugar.

EFFERVESCING LEMONADE POWDERS. Bicarbonate of soda 3½ oz., refined sugar 14 oz., essence of lemon 60 drops.

(Sometimes 12 or more grains of the powdered yellow rind of lemon-juice are added to color it.) Mix, and divide into 60 powders, or 140 grains in each blue paper. In the white papers put from 30 to 32 gr. of citric acid, or from 32 to 35 gr. of tartaric acid. Or the mixed alkaline powder and the acid may be put into separate bottles, furnished with measures holding the proper quantities of each.

Effervescing Lemonade Powders in one Bottle. *Note.*—The powders must all be separately and carefully dried, at a moderate temperature, before mixing, and when mixed, be carefully secured from the air.

1. Bicarbonate of soda 1 oz., refined sugar 3½ oz., tartaric acid 1¼ oz., essence of lemon 30 drops; mix, and put into well-corked bottles.

2. Mix 3⅓ oz. of bicarbonate of soda, 14 oz. of double refined sugar, 60 drops of essence of lemon, and 4 oz. to 4½ oz. of tartaric acid.

3. *Lemon Kali. Acidulated Kali.* Sesquicarbonated soda 8 oz., tartaric acid 8 oz., refined sugar 16 oz., essence of lemon 100 drops; mix.—Pharmaceutical Journal.

Orangeade, or Sherbet. 1. Juice of 4 oranges, thin peel of 1 orange, lump sugar 4 oz., boiling water 3 pints.

2. Juice and peel of 1 large orange, citric acid ¼ dr., sugar 3 oz., boiling water a quart.

Effervescing or Aerated Orangeade, or Sherbet. 1. Mix 1 lb of syrup of orange-peel, a gallon of water, and 1 oz. of citric acid, and charge it strongly with carbonic acid gas with a machine.

2. Syrup of orange-juice ¾ oz., aerated water ½ pint.

3. Simple syrup ½ fluid oz., tincture of orange-peel ½ dr., citric acid 1 scruple; fill the bottle with aerated water.

4. Put into a soda-water bottle ½ oz. to 1 oz. of syrup of orange-peel, 30 gr. of bicarbonate of potash, 8 oz. of water, and, lastly, 40 gr. of citric acid in crystals, and cork immediately.

5. Put into each bottle 2 or 3 dr. of sugar, 2 drops of oil of orange-peel, 30 gr. of bicarbonate of potash, or 25 gr. of bicarbonate of soda; water to fill the bottle, and 40 gr. of citric acid, as before.

Aerated Sherbet, or Orangeade Powders. Powdered

sugar 14½ oz., powdered orange-peel 12 gr., oil of orange-peel 60 drops, essence of cedrat 12 drops, bicarbonate of soda 3½ oz.; mix, and put 145 gr. in each blue paper. In the white paper put 32 gr. of tartaric (or rather 30 gr. of citric) acid. Or the alkaline and acid powders may be put into separate bottles, with a measure holding the proper proportion of each. The orange peel may be omitted.

Aerated Sherbet Powders in one Bottle. Double-refined sugar 14½ oz. (powdered orange-peel 12 gr.), bicarbonate of soda 3½ oz., essence of cedrat 12 drops, oil of orange-peel 60 drops, tartaric acid 4 oz. The powders must be carefully dried, mixed quickly, and afterwards kept dry and securely corked. A measure holding nearly 3 dr. of the powder should accompany each bottle.

Orangeade Powder, not Aerated. Citric acid ½ oz., sugar 8 oz., oil of orange-peel 20 drops.

Soda Powders. The usual proportions are: 30 or 32 gr. of bicarbonate of soda in each blue paper; and 25 or 26 gr. of tartaric acid in each white paper.

Acidulated Effervescing Powders, for making effervescing drinks with concentrated syrups of lemon, ginger, &c. Put into separate papers, distinguished by their different colors, 20 gr. of bicarbonate of soda, and 28 gr. of citric or tartaric acid. One of each powder to be dissolved separately in ⅓ of a tumbler of water, and a teaspoonful of the syrup added to the acid solution, and the liquids mixed.

The Concentrated Syrups are thus made:

Concentrated Syrup of Ginger. Simple syrup 7½ fluid oz., essence of ginger (1 part ginger to 4 of spirit) ½ oz.

Concentrated Syrup of Lemon-peel. Strong tincture of lemon-peel* 1 oz., simple syrup 15 fluid oz.

Concentrated Syrup of Orange-peel. Strong tincture of fresh orange-peel* ½ oz., simple syrup 7½ fluid oz.

(Syrup of raspberries, pine-apples, and other fruit, may be used with the above powders in the same way.)

For Seidlitz and other Medicated Powders, see Mineral

* These tinctures are thus made: Fresh lemon-peel, thin, and cut small, 4 oz.; rectified spirit 8 oz.; digest for some days, and strain. Fresh peel of Seville oranges 4 oz., spirit 16 oz.

WATERS and POWDERS, at the end of PATENT MEDICINES, &c.

SPRUCE BEER. Water 10 gallons, treacle or lump sugar (according to the color required) 6 lb, essence of spruce 4 oz.; add yeast, and ferment as for ginger beer.

SPRUCE BEER POWDERS. In each blue paper put 5 scruples of powdered sugar, 28 gr. of bicarbonate of soda, and 10 gr. essence of spruce. In each white paper 30 gr. tartaric acid.

TREACLE BEER. 1. Brown sugar 1 lb, treacle 1 lb, bruised ginger 1 oz., hops ¼ oz.; boil for a few minutes in 3 quarts of water, strain, and add 5 quarts of cold water: add a spoonful of fresh yeast; let it work all night, and bottle it in the morning.

2. Treacle 14 lb, hops 1½ lb, water 36 gallons, yeast 1 lb; boil the hops with the water, add the treacle, and strain. Cool to 80°, and ferment with the yeast. In winter ½ oz. of Cayenne pods with the hops is an improvement.—FAMILY FRIEND.

CAPILLAIRE. To 1 pint of boiling water add 3 oz. of fine maidenhair; remove from the fire, cover, and set near the fire for 3 hours; strain, and add ¼ pint of orange-flower water. Boil 1 gallon of fine syrup till reduced to 7 pints, then add the infusion, and boil for 10 minutes; strain through a jelly-bag, and when quite cold bottle the syrup. It is used to give a fine flavor to water.

LIMONIATED CAPILLAIRE. Refined sugar 24 oz., water 12 oz.; dissolve by a gentle heat; and add essence of lemon 30 drops, neroli 3 drops, citric acid 2 oz., orange-flower water 4 oz.

SYRUP OF PINE-APPLE. Expressed juice of pine-apple a pint; loaf sugar 2 lb. Boil gently, and when cold filter.

SYRUP D'ORGEAT. See Syrupus Amygdalæ, Pocket Formulary. Another formulæ for this excellent syrup is the following: take 20 oz. of sweet and 8 of bitter almonds, 9 lb of white sugar, and 4 pints of water. Blanch the almonds, dry them well, beat them with a portion of the sugar, and add gradually two-thirds of the water; strain through linen, wash the almonds on the strainer with the rest of the water, and dissolve the sugar in the strained

liquor by a gentle heat. Pour the syrup into an earthen vessel, remove the scum, and when nearly cold add 2 oz. of orange-flower water.

ACIDULATED RASPBERRY SYRUP. Put 6 lb of raspberries into a china or glass bowl, or an earthen pan not glazed with lead, with a quart of water in which has been dissolved 2½ oz. of tartaric (or preferably *citric*) acid, and let it remain 24 hours; then strain it, taking care not to bruise the fruit. To each pint of clear liquid add 1½ lb of pounded loaf sugar, and stir it with a silver spoon till dissolved: leave it for a few days, then bottle it close. A little of this syrup, or of either of the two following, with water, forms a refreshing drink in warm weather, and in some febrile disorders.

ACIDULATED STRAWBERRY SYRUP. As RASPBERRY SYRUP, using 2 oz. of citric acid, instead of 2½ oz. of tartaric acid.

RASPBERRY VINEGAR. Put a pint and a half of best wine vinegar to 3 lb of fruit in a glass or porcelain vessel; leave them together for a fortnight, then strain without pressure. Or put an equivalent quantity of strong acetic acid (4 oz. of the usual strength) to the fruit, in the same way. Or it may be made as directed above for ACIDULATED RASPBERRY SYRUP.

WHEY POWDER. Sugar of milk in fine powder 2 oz., powdered white sugar 7 oz., gum Arabic ½ oz.; mix. An ounce dissolved in a quart of water as a substitute for whey.

WHEY may be made by adding a little infusion of rennet (prepared calf's stomach) to milk, and gently heating it till curdled. It is also made by heating a quart of milk nearly to boiling, and adding either a little lemon-juice, orange-juice, solution of citric acid, vinegar, or white wine, or cream of tartar sufficient to turn it. It is then strained. If required *bright*, beat up the white of an egg with a portion of the whey, mix with the rest, boil for a moment, and run it through a jelly-bag.

One or two recipes in Confectionery may be introduced here.

ORANGE MARMALADE. 1. Procure some large Seville oranges with clear skins, peel them, squeeze out the pulp and juice, taking care to remove all the pips. Boil the peel, divided

into quarters, till they are sufficiently tender; scrape clean all the inside from them, lay them in folds, and cut them into very thin slices about an inch long. Weigh the juice, pulp, and boiled peel; then add broken lump sugar equal in weight to the whole, and boil for half an hour, carefully removing the scum. Then put it into pots, and when quite cold cover them over.

2. Instead of using all Seville oranges, let only half or a third of them be bitter, and the rest common sweet oranges. Proceed in the same way. Some add honey.

Currant Jelly. Pick the currants, put them in an earthen jar, and place it in boiling water till the juice is extracted. Strain through a sieve without pressing them, and boil the juice in an enamelled saucepan with its weight of loaf sugar, removing the scum as it rises. When it will jelly on the back of a cold spoon, it is sufficiently done. A little of the jelly dissolved in warm water forms an agreeable beverage.

DIETETIC ARTICLES.

As the ingredients of some of the following compounds are usually sold by druggists, who may be expected to furnish information as to the manner of using them, and as they may all be regarded as auxiliaries to medical treatment, some notice of them here seems desirable, though it must necessarily be brief and incomplete.

Arrow-root. (West Indian arrow-root is the fecula of the tubers of the Maranta arundinacea; East Indian arrow-root is obtained from the Curcuma angustifolia; South Sea or Tahiti arrow-root from the Tacca pinnatifida. They have all the same properties, and are used in the same manner.) Mix a dessert-spoonful of arrow-root with sufficient cold water to form a soft paste; rub it till quite smooth, and add half a pint of boiling water, stirring it briskly. Boil it for a minute or two, and when removed from the fire add a teaspoonful of sherry or other white wine (where wine is admissible), with a little grated nutmeg or lemon-peel, and sugar to the taste. For young children milk should be used instead of water, and the wine omitted: it is also more nourishing in this form for those invalids with whom milk agrees.

Tous les Mois. (The fecula of a species of Canna.) It is used in the same way as arrow-root; but rather less is required. It forms a more tenacious, but less transparent jelly.

Sago. (The granulated fecula of the pith of one or more species of the Sago Palm.) Wash an ounce of pearl sago in cold water; then boil it very gently in a pint of fresh water, stirring it frequently till dissolved. It may be flavored with wine, spices, and sugar, as directed for arrowroot. For children, and for consumptive and debilitated patients, it may be made with milk instead of water. The common sago, being in larger grains, requires more time to dissolve: and is usually steeped for some hours before boiling it.

Tapioca. (Obtained from the tuberous roots of the Cassava, Jatropha manihot. It is usually sold in small lumps formed by drying the fecula on hot plates.) It is used in the same way as sago; but requires to be previously steeped for some hours, or to be simmered for a longer time. It forms a clear jelly, which may be flavored with wines, spices, and sugar, as directed for arrowroot; but is more nourishing when made with milk.

Sago Posset, for invalids. Macerate a tablespoonful of sago in a pint of water for 2 hours on the hob of a stove, then boil for 15 minutes, assiduously stirring. Add sugar with an aromatic, as ginger or nutmeg, and a tablespoonful or more of white wine. If the wine be not permitted, flavor with lemon-juice.

Sago or Tapioca Milk, for invalids. Take an ounce of either of these feculæ, and soak it in a pint of cold water for an hour; then pour off this water, and adding 1½ pint of good milk, boil slowly until well incorporated.—Dr. A. T. Thomson.

Tapioca Pudding, for invalids. Beat up ½ ounce of sugar with the yolks of 2 eggs, and stir the mixture into a pint of tapioca milk.—Dr. Thomson.

Arrow-root milk and pudding may be made like the corresponding preparations of tapioca.

Panada, for invalids. Place in a saucepan some very thin slices of bread crumbs, and add rather more water than will cover them. Boil now until the bread becomes pulpy, strain off the superfluous water, and beat up the remainder

into the consistence of gruel. Sweeten with white sugar, and add, if permitted, a little sherry wine.

Barley Water. See Decoctum Hordei, and Decoctum Hordei Compositum. Robinson's Patent Barley is a convenient preparation; printed directions accompany it.

Asses' Milk, artificial. Eringo root, pearl barley, sago, rice, of each 1 oz. Wash them with cold water, then boil them with 3 pints of water to 1½ pint, and strain. Put a teaspoonful to a cup of boiling milk, and sweeten to the taste. (*Bonbons de lait d'ânesse* are made with sugar of milk, white sugar, gum, and starch or arrow-root.)

Linseed Tea. Take 1½ oz. of clean linseed, and ½ an oz. of bruised liquorice root; put them into a warm teapot or jug, and pour on them 2 pints of boiling water; let them stand, covered, near the fire, for 3 or 4 hours, stirring them occasionally; then strain. To save time, the ingredients may be boiled for 15 or 20 minutes, instead of infusing them; but the tea so made is less agreeable.

Iceland Moss. Infuse an ounce of picked Iceland Moss for 15 minutes in half a pint of hot water; strain off the water and boil the moss in a quart of fresh water till reduced to a pint and a half. Half an ounce of liquorice root may be added, towards the end of the boiling, if agreeable: or milk may be used instead of water.

Jelly of Iceland Moss. See Gelatina Lichenis, Pocket Formulary. Another form is the following: Infuse 2 ℔ of the moss for half an hour in sufficient boiling water to cover it; drain the moss, and boil it in 2½ gallons of water for an hour, and strain. Boil the moss with fresh water, adding an oz. of isinglass; strain; mix the product of the two boilings, and let it stand till clear. Evaporate the clear liquid to the consistence of a stiff jelly, adding towards the end, 6 ℔ of fine lump sugar, 2 oz. of French brandy, and half an ounce of orange-flower water. It may be taken, almost at pleasure, dissolved in water or milk.

Iceland Moss Chocolate. See Chocolate Lichenis, Pocket Formulary.

Irish Moss, or Carrageen. Steep a ¼ of an ounce of the moss in cold water for a few minutes; then withdraw it, shaking the water from each sprig, and boil it in a quart

of milk till it attains the consistence of jelly, and sweeten to the taste. A decoction of the same quantity of moss in a quart of water is also used as a demulcent in coughs, &c. BLANC-MANGE may be made by washing ½ oz. of the moss as above, and boiling it in 1½ pint of new milk to such a consistence that it will retain its form when cold, sweetening and flavoring it to the taste. An agreeable jelly may be made by boiling it with water instead of milk, and adding lemon or orange juice or peel, wine, &c.

CEYLON MOSS. Boil ½ oz. of the prepared moss in a quart of water for 25 minutes; or till a teaspoonful taken out forms a firm jelly in 2 or 3 minutes: then flavor with wine, cinnamon, or with lemon or orange juice or peel; and sweeten to the taste. Boil for five minutes longer, and press through a jelly-bag, or double muslin. Pour it into earthen moulds, and leave it undisturbed till it has set. If the jelly is required *bright* it must be clarified with white of egg, as directed for gelatine jelly. For BLANC-MANGE add 1 oz. of prepared moss to a quart of boiling water, and boil gently till reduced to a third; add the milk and flavoring ingredients, and pour into earthen moulds.

AUSTRALIAN MOSS. This has been introduced for the same use as Irish and Ceylon mosses, but has not been very generally adopted. Soak ½ oz. of the moss in water for an hour or two, pour away the water, and boil the moss in a quart of fresh water till dissolved. Strain through a hair sieve, and sweeten and flavor to the taste.

SALEP. (The dried root of some species of orchis.) Boil ½ oz. of salep powder in a pint of water till dissolved; strain, and sweeten and flavor to the taste.

HARTSHORN JELLY. Boil 4 oz. of true hartshorn shavings (previously washed in warm water) in a quart of water till reduced to a pint; strain, and sweeten and flavor to the taste. For children and consumptive patients the simple jelly may be mixed with milk and a little sugar. To make a bright jelly for the table, boil 4 oz. of washed hartshorn shavings in 1½ pint of water, till reduced to ¾ of a pint, and add 2 oz. of sugar, and a tablespoonful of lemon or orange juice. Strain with pressure; beat up the white of an egg with a little cold water, mix this thoroughly with the jelly, and evaporate the liquid till a little taken out solidifies on

cooling. Add a little fresh lemon-peel, and strain through a jelly-bag.

GELATINE JELLY. Steep 1 oz. of NELSON'S or other purified gelatine in half a pint of cold water for ten minutes; then add the same quantity of boiling water, and stir till it is dissolved, applying heat if required; add the juice and peel of two lemons, sugar, and wine sufficient to make up the whole to a pint and a half. If required bright, have ready the white and shell of an egg well beaten together, stir them briskly into the jelly, boil for 2 or 3 minutes without stirring, and pass through a jelly-bag. As a nourishing diet for children and invalids, a little of the gelatine simply dissolved in water may be mixed with milk, or the dry gelatine dissolved in milk by heat.

ISINGLASS JELLY. Isinglass is used in the same way as gelatine, but as it is not wholly soluble in water, it requires straining. To make a bright jelly it requires more eggs for its clarification than gelatine. A very pleasant jelly is made with the acidulated Raspberry or Strawberry Syrup (p. 275) thus: Dissolve 1½ oz. of isinglass in a very little water, put this to a quart of the syrup, warm it and stir it well; then strain it into a mould. In warm weather put 2 oz. of isinglass.

ARROW-ROOT BLANC-MANGE. Beat up 2 oz. of genuine arrow-root with a little cold milk to the consistence of cream; pour on it 1½ pint of boiling milk, stirring it all the time. Flavor and sweeten to the taste, boil for 10 minutes, stirring it constantly, pour into moulds, and leave it until next day.

BLANC-MANGE. This may be made with either isinglass or gelatine. Boil ½ oz. in 16 fluid oz. (the old wine pint) of new milk; stir it constantly till it boils, let it simmer for a few minutes till the isinglass is dissolved: strain, add sugar to the taste, and a few drops of almond flavor, or other flavoring ingredients, and pour into moulds.

CHOCOLATE. This is prepared from the finest cocoa-nuts (seeds of Theobroma cacao) after roasting, winnowing, &c., by grinding them on a hot stone or plate, or beating them in a hot mortar to a smooth paste. Sugar is generally added, and vanilla, or other flavoring ingredients.

CHOCOLATES, Medicated. See Chocolata, Pocket Formulary.

WHITE CHOCOLATE. White sugar 3 ℔, rice flour 27½ oz., English or Indian arrow-root 8 oz., tincture of vanilla ½ oz., butter of cacao 8 oz., powdered gum arabic 4 oz.; form a paste with boiling water, and put it into moulds.

COCOA. This should also be prepared from the seeds of Theobroma cacao; and the rock, roll, and flake cocoas often consist of this alone. But most of the paste cocoa, and soluble cocoa powder, is mixed with saccharine and farinaceous matters. This is the case with much of the "*Homœopathic*" *Cocoa*, which professes to be unadulterated, but generally contains potato-starch. A common proportion for soluble cocoa appears to be ⅔ of pure cocoa, and ⅓ of sugar and farina; the latter being one or more of the following: Wheat flour, sago meal, potato flour, arrow-root, &c. The *Paste Cocoa* often contains only about half its weight of cocoa, the rest being sugar and water, with sometimes the addition of sago meal or other farina.

BROMA. This consists of about 8 oz. of pure cocoa, 3½ of sugar, and 4½ of sago meal, arrow-root, &c.

WACAKA DES INDES. Roasted cacao beans (chocolate) in powder 2 oz., sugar 6 oz., cinnamon ¼ oz., vanilla (powdered with part of the sugar) ½ dr., ambergris 3 gr., musk 1½ gr. Sometimes 1 dr. of prepared annotto is added, and the ambergris and musk omitted.

RACAHOUT DES ARABES. This is professedly a preparation of acorns (perhaps those of the Quercus ballotta, which are naturally sweet, or of other kinds deprived of their bitterness by being buried in the earth); but it is imitated by the following: 1. Chocolate in powder 1 oz., rice flour 3 oz., sugar 9 oz., potato arrow-root 3 oz., vanilla (pulverized with part of the sugar) 1 dr.; mix.

2. Chocolate in powder 4 oz., salep 1 oz. (or powdered tragacanth 1 oz.), potato arrow-root 5 oz., sugar (flavored with vanilla) 8 oz.—CADET.

DICTAMIA. Sugar 7 oz., potato arrow-root 4 oz., flour of brent barley (Triticum monococcum) 3 oz., Trinidad and Granada chocolate, each 1 oz.; vanilla 15 gr.

PALAMOUD. Chocolate 1 oz., rice flour 4 oz., potato arrow-

root 4 oz., red sanders, in fine powder, 1 dr.; mix. (In the above, by chocolate is meant the cacao beans roasted and pulverized without addition. Indian arrow-root or tous les mois may be substituted for the potato arrow-root.)

Ferculum Saxonia. Barley flour 21 oz., sugar 7 oz., cinnamon 1 dr.; mix, and bake them in an oven, enveloped in a paste of wheat flour, and placed in an earthen vessel. When sufficiently baked, remove the crust, and when the contents are cool, reduce them to powder. About ½ oz. to 1 oz. is boiled with broth, &c., as a nourishing diet. It is often medicated with the addition of sarsaparilla, bark, &c.

Farinaceous Food, &c. The following compounds are accompanied with full directions for use:

Baster's *Soojee and Compound Farina.* Wheat flour, with sugar.

Bright's *Nutritious Farina.* The basis is said to be potato-starch.

Bright's *Breakfast Powder.* A combination of chocolate with his nutritious farina.

Braden's *Farinaceous Food.* Chiefly wheat flour, carefully baked.

Bullock's *Semola.* Wheat flour, from which a portion of the starch has been removed, so as to leave a definite proportion of gluten.

Denham's *Farinaceous Food.* As Braden's, with perhaps a mixture of barley flour.

Gardiner's *Alimentary Preparation.* Very finely ground rice flour.

Hard's *Farinaceous Food.* Carefully-baked wheat flour.

Hunt's *Breakfast Powder.* Rye, carefully roasted as coffee. (For Dandelion Coffee, see p. 179.)

Leath's *Alimentary Farina.* Baked wheat flour, with sugar, potato flour, and a small quantity of Indian corn meal, and tapioca.

Maidman's *Nutritious Farina.* Potato flour, tinged with some pink coloring matter.

Palmer's *Vitaroborant.* See Ervalenta, below.

Plumbe's *Farinaceous Food.* South Sea arrow-root, combined with pea flour.

Prince of Wales's Food. Potato flour.

Semolina. A hard kind of wheat, containing much gluten, ground into coarse grains. But some articles sold under this name appear to be compounds of gluten, artificially granulated, resembling BULLOCK'S Semola.

ERVALENTA, REVALENTA ARABICA, LENTIL POWDER, &c. These consist chiefly either of the European or Egyptian lentil.

Ervalenta. WARTON'S consists of the French or German lentil, with either Indian corn, or, more probably, a species of corn called *Durra*, used by the Arabs. But Dr. SCHENK states that what is sold at Paris consists of the flour of French beans and Indian corn.

Revalenta Arabica. A mixture of the red (Egyptian or Arabian) lentil with barley flour. Some samples contain sugar, others salt and a flavoring ingredient.

Lentil Powders. Some consist entirely of lentil flour (French or German), or Egyptian, or both kinds mixed. Others contain barley flour in addition. NEVILL'S consists of 1 oz. of curry powder to 4 lb of lentil flour. The LANCET gives the following recipes for lentil powder: 1. Arabian lentil flour 2 lb, barley flour 1 lb, salt 3 oz.

2. Pea flour 2 lb, Indian corn flour 1 lb, salt 3 oz.

GRUEL is made either from oatmeal or from groats or grits (oats deprived of their cuticle), either whole or crushed (Embden groats). Dr. THOMSON directs 3 oz. of groats, previously washed, to be boiled slowly in 4 pints of water, till reduced to 2 pints, then strained through a sieve. The Embden groats require less boiling. Dr. KITCHENER directs 1 or 2 tablespoonfuls of oatmeal (according as the gruel is preferred thin or thick) to be well mixed with 3 spoonfuls of cold water gradually added, 1 pint of boiling water poured on it, and the whole boiled for five minutes, constantly stirring it; it is then skimmed, and strained through a hair sieve; a little butter is usually added, and sometimes milk, with salt, or otherwise sugar and spices to the taste. Thorough trituration of the oatmeal and cold water, and constant stirring of the gruel while on the fire, render long boiling unnecessary.

BOILED WHEAT. Steep the wheat in water for 10 or 12 hours, then boil it for ½ an hour. (As a substitute for vegetables, and to obviate constipation.—Mr. L. BULLOCK.)

Beef Tea. Professor Liebig directs 1 ℔ of beef, free from fat, to be minced very small, mixed with an equal weight of cold water, and heated slowly to boiling; when it has boiled for a minute or two, strain through a cloth. It may be colored with roasted onion or burnt sugar, and salted to the taste. Dr. Seymour directs 2½ ℔ of lean beef cut small to be put into 3 pints of cold water, and simmered slowly, without boiling, till reduced to a pint and a half; then carefully strained.

Extract of Meat. Cut the lean of fresh-killed meat very small, put it into 8 times its weight of cold water, and heat it gradually to the boiling-point. When it has boiled for a few minutes, strain it through a cloth, and evaporate the liquor gently by water-bath to a soft mass. 2 ℔ of meat yield 1 oz. of extract. Fat must be carefully excluded, or it will not keep.—Liebig.

Liebig's Food for Infants. Malt flour, to which is added a small quantity of bicarbonate of potash or soda.

Liebig's Soup for Children. Mix ½ oz. wheat flour, ½ oz. malt flour, 7½ gr. bicarbonate potash; add 1 oz. of water, and 5 oz. of milk. Heat with constant stirring over a gentle fire till it begins to thicken. Remove from fire and stir for 5 minutes. Heat once more, and again remove and stir. Heat again to ebullition. Separate the bran from the soup by a fine sieve.

Trophazome. Mince 16 oz. of meat, free from fat very fine, pour on it 8 oz. of cold or lukewarm water (not exceeding 100°); mix well, and let it stand for an hour, stirring it 3 or 4 times. Press out the liquid (about 6 oz.); mix 8 oz. more of water with the meat, stir it occasionally, and in half an hour strain with pressure. Repeat this with 8 oz. more water. Break up the pressed meat, and put it into a small tin vessel; place this in a water-bath of cold water, heat gradually to the boiling-point, and keep it boiling for 20 minutes. Mix the fluid which exudes with the others, add salt, spices, and other flavoring ingredients, and boil for 20 minutes in a covered vessel. It may be thickened with 1 oz. of semola.—Mr. Bullock.

Meat Biscuits. A thick extract of meat (made by boiling fresh-killed beef or other meat, and evaporating the strained liquid) is kneaded with wheaten flour, and the dough rolled

out and divided into biscuits, which are dried or baked in an oven. They are kept in the form of biscuits, or coarsely ground; 1 ounce makes a pint of rich soup, which may be salted or flavored to the taste.

BREAD, AERATED. DAUGLISH'S patent. This is prepared without yeast, to the saving, it is said, of 10 per cent. of the weight of the flour. Aerated water is forced by machinery into the mass of flour inclosed in an iron vessel. The dough is then kneaded by machinery inside the vessel. The pressure being now removed, the dough instantly rises. The whole process occupies no more than half an hour, instead of eight or ten hours, as when yeast is used.

BREAD, UNFERMENTED. Mix carefully ½ oz. of bicarbonate of soda and ¼ oz. of salt with 4 ℔ of flour; mix this with a quart (or rather 41 or 42 fluid oz.) of very cold water, previously mixed with ½ a fluid oz. and 20 minims of muriatic acid of 1.16 specific gravity, into a thin dough, with as little kneading as possible, and let it be *immediately* placed in the oven; it requires rather more time than fermented bread. By mixing 26 measures of the acid with 46 of water, a diluted acid is obtained, of which a fluid ounce and a half may be taken for every ½ oz. packet of soda.

A pamphlet on the subject directs for *Brown Bread*, 3 ℔ of wheat meal and 10 dr. (apoth. weight) of bicarbonate of soda to be well mixed, and made into dough with 25 oz. of cold water, previously mixed with 12½ fluid dr. of muriatic acid.

The proportions used first by Dr. WHITING, and subsequently by Mr. DODSON, in the preparation of unfermented bread and biscuits, were: 7 ℔ of wheaten flour, 350 to 500 gr. of carbonate of soda, 2¾ pints of water, with muriatic acid 420 to 560 gr., or as much as may be sufficient. The soda, dissolved in a small quantity of water, is first carefully kneaded with the dough, and the acid being afterwards rapidly mixed in, the bread is baked without delay.

A third formula is that of Mr. J. SAVORY, and is recommended as excellent by Dr. PEREIRA. Intimately mix with 1 ℔ of flour, sesquicarbonate of soda 40 grains, and powdered white sugar a teaspoonful, in a large basin with a wooden spoon. Then gradually add cold water about ½ a pint, previously mixed with 50 drops of common muriatic

acid, and stir constantly, so as to form very speedily an intimate mixture. Divide into 2 loaves, and put into a quick oven immediately.

Another form of unfermented bread is as follows: Mix 1 oz. of bicarbonate of soda, ¾ oz. of tartaric acid, and ¼ oz. of salt, with 7 lb (half a peck) of flour; mix the whole thoroughly, taking care that all the ingredients are perfectly dry; add, in two or three portions, 4 pints of cold water, and incorporate quickly; place it in tins, and send it to the oven immediately. If not baked in tins, less water must be used.

Jones's Patent Flour contains all the ingredients ready mixed, and may thus be kneaded with water, and baked at once, without further additions. To 1 cwt. of perfectly dry wheat flour, 10½ oz. (avoir.) of dry, finely powdered tartaric acid are added. Mix well, pass through a flour-dressing machine, and allow it to remain for 2 or 3 days. Then add, all in fine powder and dry, 12 oz. of bicarbonate of soda, 24 oz. of common salt, and 8 oz. of loaf sugar. Mix all thoroughly together, and pass through a flour-dressing machine, when it is ready for use. 1 lb made into bread will require 10 oz. of water, or, for biscuits 6 oz., and the dough must be baked at once in a well-heated oven.

(Biscuits and cakes made without yeast, and containing no butter, are prescribed in some dyspeptic cases; of such the following are examples:)

Abernethy Biscuits. Make into a stiff biscuit paste, 1 quart of milk, 6 eggs, 8 oz. of loaf sugar, and ½ oz., of caraway seeds, with flour sufficient to bring the whole to the required consistence. Make the biscuits thin, dock them with holes to prevent them from swelling up, and bake in an oven at a moderate heat.

Sponge Biscuits. Beat the yolks of 12 eggs for ½ an hour, then put in 1½ lb of finely-powdered sugar, and whisk it briskly until it rises in bubbles; beat the whites to a strong froth, and whisk them well with the sugar and yolks; then work in 14 oz. of flour, with the rinds of 2 lemons grated. Bake in tin moulds buttered, for 1 hour in a quick oven, sifting over them a little fine sugar.

Rice Cake. Beat the yolks of 15 eggs for nearly half an

hour with a whisk, mix well with them 10 oz. of finely-sifted white sugar; put in ½ lb of ground rice, a little orange-flower water or brandy, and the rinds of 2 lemons grated; then add the whites of 7 eggs, well beaten up, and stir the whole for ¼ of an hour; put into a hoop, and bake for ½ an hour in a quick oven.

(Bread and biscuits made with ginger as an ingredient may also be useful in assisting a weak digestion. The two following recipes are by M. Soyer:)

Common Gingerbread. Put on a slab or table 1 lb of flour; make a ring of it; put ½ pint of treacle in, mix well together, working it so as to form a stiff paste. Put some flour into a basin, to which add this dough, which will keep thus for 7 or 8 weeks. When wanted, put in any quantity of ground ginger, according to taste or desire. Mix well, roll thin, and cut into pieces about the size of a crown; put them on a baking sheet, and bake for a few minutes, till crisp. To every pound of paste, an ounce of butter may be used, if preferred. These cakes will keep a long while in an air-tight case.

Ginger Cake. Take ½ lb of sugar, ½ lb butter, 1½ oz. of ground ginger, 6 eggs, beat well, stir in 1½ lb of flour, and add as much milk, a little warm, as will make a stiff dough for bread. Bake in a pan for 2 hours.

Gluten Bread (for diabetic patients). It is made with the gluten of flour, a small portion only of the starch being retained.

Dr. Aldridge recommends, for preparing the gluten, second best flour. This must be made into a stiff paste with cold water, and then kneaded with the hands under a current of water, on a slanting board placed in a two-gallon jar, until starch can no longer be detected by tincture of iodine in small portions taken from the mass. A stone of flour yields from 3½ to 4½ lb of gluten. The gluten may also be obtained from starch manufactories.

To prepare the bread, the following materials should be mixed: Fresh moist gluten 24 oz., bicarbonate of ammonia 3¼ dr., common salt 1½ dr., powdered caraway 48 gr., wheaten flour 4½ oz., powdered bran 1½ oz., salt butter 4 oz. These quantities yield 24 oz. of bread, when baked. It may be baked in small and flat circular tin pans, placed on a moderately heated hot-hearth. It is difficult to bake this

bread properly with yeast, as prescribed by BOUCHARDAT; and if made without butter, or with the total exclusion of starch, it becomes extremely unpalatable.

Dr. PERCY proposes the following method: Take the matter left after removing the starch from 16 ℔ of rasped potatoes, ¾ ℔ of mutton suet, 12 eggs, ½ ℔ of butter, and ½ oz. of carbonate of soda; mix, and add 2 oz. of diluted hydrochloric acid; divide into 8 cakes, and bake immediately in a quick oven.

(Various alimentary preparations have lately been introduced, the basis of which is the gluten which remains in extracting the starch from wheat flour by the mechanical process. Mr. GENTIL'S *gluten flour* is a mixture of this with wheat flour. It contains 42 per cent. of gluten, and yields a nutritious and digestible *gluten biscuit*, *gluten bread*, and, with cocoa, *gluten chocolate*. Mr. BULLOCK'S *semola* and Mr. VERON'S *granulated gluten* are of the same nature. 30 parts of white flour, 10 of fresh gluten, and 7 of water, form a paste resembling *Italian macaroni*, vermicelli, &c.)

BAKING POWDER. See page 169.

CUSTARD POWDER. Rub up together gum tragacanth 2 oz., potato starch 1 ℔, powdered turmeric 2½ dr., with oil of bitter almonds ½ dr., and essence of lemons 1 dr. Put up into 1-ounce packets. (From 1 pint of new milk take 2 tablespoonfuls to work up with the powder; boil the remaining milk with 2 oz. of lump sugar, and pour it, while boiling, into the basin, stirring quickly until thoroughly mixed. Bake as a custard.)—Mr. SCHOLEFIELD.

Without the coloring, this forms *Blanc-mange Powder*.

RENNET WINE (for dyspeptic persons). Take the fresh rennet bag of a calf, cut off and throw away 3 inches of the upper or cardiac extremity, slit the rest longitudinally, wiping gently with a dry napkin, but not removing the clean mucus, cut it into small pieces, and put into a common wine-bottle filled up with good sherry. Let it remain corked for three weeks. One teaspoonful in a wineglassful of water, taken after meals, assists digestion.—Mr. GEO. ELLIS.

PREPARATIONS OF COD-LIVER.

THESE are designed for such invalids as are likely to derive benefit from cod-liver oil, but are unable to overcome their disgust for that useful medicine. In such cases the fresh liver itself may be found of equal service. The following methods of cooking it are adapted from M. SOYER.

POTATOES AND COD-LIVER. Take 1 lb of fresh cod-liver; peel and steam 2 lb of floury potatoes. Cut the liver in 4 pieces, place it over the potatoes, then again steam them, letting the oil from the liver fall on them. When done make some incisions in the liver with a knife, to extract the oil remaining. Afterwards, the liver may be eaten with some anchovy sauce,—and the potatoes, mixed with the oil, with a little salt and pepper. (Simple cod-oil, as tasteless as can be procured, may be made in the same manner to yield a palatable dish with potatoes.)

RICE AND COD-LIVER. Boil ½ lb of rice in 2 quarts of water. When nearly done, remove 3 parts of the water, then put over the rice 1 lb of cod-liver, cut in large dice. Put the saucepan in a slow oven for about 30 minutes, by which time it will be well cooked. Take the liver out, which serve as above. Stir up the rice with a fork, and stir it with salt and pepper. If no oven, care must be taken to cook over a slow fire, or else it will burn.

TAPIOCA AND COD-LIVER. Boil ¼ lb of tapioca till soft in 2 quarts of water, drain in a cullender, then put it back in the pan; season with salt and pepper, add ½ pint of milk, put over 1 lb of fresh cod-liver, cut in 8 pieces. Set the pan near the fire to simmer slowly for half an hour, or a little more, till the liver is quite cooked. Press on it with a spoon to squeeze out the oil. Take away the liver, and mix the tapioca. If too thick, add a little milk, boil for a few minutes, add a little more salt and pepper, and serve. Here again a slow oven is better than a fire.

COD-ROE AND COD-LIVER. Take a cod's roe, cut open the skin which surrounds it; put the eggs in a basin, pour water over them, mashing with the hand, to separate them; throw away the water. Add ½ lb of salt and a teaspoonful of pepper. Let them soak all night. Wash well 2 or 3 times, leaving about a gill at the bottom. Put over it now 2 lb of the

liver, in 6 or 8 pieces; put the stew-pan on a very slow fire or in an oven for an hour; then take out the liver, which serve as usual. Add about a gill of melted butter to the roe, when it will be ready.

COD-LIVER SAUCE. ½ lb of cod-liver, previously boiled, and cut in large dice, may be added to a little anchovy sauce, and ½ pint of melted butter. (This may be eaten with potatoes or fish. Cod-liver may be used as a sauce in the same way, disguising its taste in the butter by pepper, vinegar, or anchovies, and gradually increasing its amount as the patient becomes habituated to it.)

CONDIMENTS, AND VARIOUS CULINARY COMPOUNDS.

CURRY POWDER. The recipes for "true Indian Curry Powder" are numerous, and vary much in the number and proportion of the ingredients. The total quantity of powder in each of the following recipes being nearly equal, the relative proportion of the different coloring, heating and flavoring ingredients, will at once be seen. Dr. KITCHENER complains that the proportion of cayenne is generally so large, that a proper quantity of the powder cannot be used to obtain the benefit of the other ingredients; and the Editor of the Pharmaceutical Journal justly observes that many recipes contain too large a proportion of turmeric. All the ingredients should be of fine quantity and recently ground.

	1	2	3	4	5	6	7	8	9	10	11	12
Turmeric,	9	6	6	8	9	9	9	4	6	6	7	8
Coriander seed,	9	16	12	22	10	9	16	12	11	16	13	12
Mustard, *scorched*,	3	—	—	—	3	3	—	4	—	—	—	—
Cayenne,	2¼	4	1½	1	1¼	¾	1½	2	1	2	1	1
Pepper, black or long,	6 bl.	3 bl.	8 long	2 bl.	3 bl.	3 bl.	2½ bl.	4 bl.	5 bl.	1¼ bl.	5 bl.	4 bl.
Pimento,	—	—	—	—	—	—	—	—	2	¾	—	—
Cloves,	1	—	—	—	½	—	—	—	½	¼	—	—
Cinnamon,	2	—	—	—	—	¾	—	—	3	¾	—	—
Cardamom,	2	3	—	—	1½	1½	—	—	—	—	—	2
Ginger,	1	—	—	—	3	3	—	—	2	1½	—	4
Mace,	1	—	—	—	—	—	¼	¼	—	½	—	—
Fenugreek,	—	—	3.	—	—	—	—	2	2	3½	3	—
Cumin,	—	—	2	—	1	—	3	2	—	—	3	1

The addition of 1 oz. of garlic, or 2 oz. of shallots, to 2 lb of either of the above will be approved by some palates.

The *true* Indian curry is said to be thus made: Coriander seed 6 dr., turmeric 5 scruples, fresh ginger 4½ dr., cumin seeds 18 gr., black pepper 54 gr., poppy seed 94 gr., garlic 2 heads, cinnamon a scruple, cardamom 5 seeds, 8 cloves, 1 or 2 chillies, half a cocoa-nut grated; all but the last to be ground on a stone.

Bengal Chitni. Chillies 1½ lb, unripe mangoes (or apples) 1 lb, red tamarinds 2 lb, sugar candy 1 lb, fresh ginger root 1½ lb, garlic ¾ to 1½ lb, sultana raisins 1½ lb, fine salt 1 lb, and 5 bottles of the best vinegar; soak the chillies for an hour in the vinegar, then grind all with a stone and muller to a paste.

Italian Tamara. Coriander seed, cloves, and cinnamon of each 8 oz.; anise and fennel seeds, of each 4 lb; mix.

Mixed Spices and Savory Herbs. I. *Kidder's Sweet Spice.* Equal weights of cloves, mace, nutmegs, cinnamon, and sugar.

2. *Kidder's Savory Spice.* Equal weights of salt, pepper, cloves, nutmegs, and mace.

3. *Ragout.* Salt 16 oz., pepper 8 oz., nutmeg, ginger, and allspice, each ¼ oz.; lemon-peel 8 oz., mustard flour 8 oz., cayenne 2 oz.; mix.

4. *Sausage.* Pepper 5 lb, cloves 1½ lb, nutmegs 1½ lb, ginger 2½ lb, aniseed ½ lb, coriander seed ½ lb; mix.

5. Dr. Kitchener's *Savory Ragout.* Salt 2 oz., mustard, black pepper, and grated lemon-peel, of each 1 oz., allspice, ginger and nutmeg of each ½ oz., cayenne ¼ oz.

6. *Soup Herb and Savory Powder.* Mix 3 parts of No. 7 with 1 part of No. 5.

7. Dr. Kitchener's *Soup Herb Powder, or Vegetable Relish.* Dried parsley, winter savory, sweet marjoram, lemon thyme, of each 2 oz., dried lemon-peel, and sweet basil, of each 1 oz.: mix. They should be carefully dried in a Dutch oven, powdered, passed through a hair sieve, and kept in closely-covered bottles. For sauces, soups, &c.

8. *Pease Powder.* Pound together in a marble mortar 2 oz. each of dried mint and sage, ¼ oz. each of celery seed and black pepper, and rub them through a hair sieve.

Horseradish Powder. Take up the roots in November or December, dry them carefully with a gentle heat, and reduce to powder.

SOLUBLE CAYENNE. To 1 lb of the best cayenne pepper, add as much rectified spirit as will form it into a paste. Cover this up for 2 hours; then place it in a percolator, and gradually pour on it more spirit till a pint of liquid is procured. A little water cautiously poured on the pepper will displace most of the remaining spirit. Distil off most of the spirit for future use, and add to the residue 3 lb of fine salt, and evaporate the mixture to dryness by the heat of a water-bath. It is usually colored, but is better without it.

CULINARY ESSENCES, TINCTURES, &c.

ALMOND FLAVOR. Essential oil of bitter almonds 1 part, rectified spirit 7 parts. Some put 1 part of oil to 15 of spirit; others 1 part to 3. It should not be sold without a caution as to the quantity to be used; or rather, the oil should be first purified from its hydrocyanic acid, by mixing it with a solution of chloride of iron and cream of lime, with a little peroxide of mercury, and, after a few days' contact, carefully redistilling the oil.

FLAVORING ESSENCE. Purified oil of bitter almonds 8 drops, essence of lemon 12 drops, oil of cinnamon 8 drops, oil of nutmeg 4 drops, highly-rectified spirit 1 oz. A few drops to be added to puddings, custards, &c.

LEMON FLAVOR. Fresh lemon-peel cut thin, 3 dr., essence of lemon 1 dr., alcohol 3 oz. (Another method is to rub a lump of sugar on clean, dry lemons, till the yellow rind is taken up by the sugar; then scrape off the saturated part of the sugar, and keep it in a closely-covered pot for use.)

TINCTURE OF CINNAMON (KITCHENER'S). Bruised cinnamon 3 oz., 1 bottle of Cognac brandy; digest for a fortnight, and strain. (Tincture of Allspice, Nutmeg, Cloves, in the same manner.)

ESSENCE OF CINNAMON. Bruised cinnamon 2 dr., oil of cinnamon 1 dr., highly-rectified spirit 3 oz.; digest, and strain.

ESSENCES OF NUTMEG, MACE, CLOVES, ALLSPICE, &c. These are made from the spices and their essential oils, as ESSENCE OF CINNAMON.

ESSENCE OF CELERY. Celery seed ½ oz. to 1 oz., brandy 4 oz.; digest for 8 or 10 days, and filter.

ESSENCE OF CARAWAY. Bruised caraway seed 1 oz., rectified spirit 8 oz., oil of caraway ¼ oz., brown sugar ¼ oz.; digest for 8 or 10 days, and filter.

AROMATIC ESSENCE OF GINGER. Fresh-grated ginger 3 oz., fresh thin lemon-peel 2 oz., brandy 1½ pint; macerate for 10 days.—Dr. KITCHENER.

ESSENCE OF PEAR AND ESSENCE OF PINEAPPLE. See Trade Chemicals.

ESSENCE OF CAYENNE (KITCHENER'S). Put ½ oz. of cayenne pepper into ½ a pint of brandy; let it steep for a fortnight, then pour off the clear liquor. (A much stronger essence is sometimes kept, prepared by percolation, as directed for Soluble Cayenne.)

SPIRIT OF SAVORY SPICES (KITCHENER'S). Black pepper 1 oz., allspice ½ oz., nutmeg ¼ oz. (all pounded); infuse in 16 fluid oz. of brandy for 10 days.

SPIRIT OF SOUP HERBS (KITCHENER'S). Lemon thyme, winter savory, sweet marjoram, sweet basil, each 1 oz., grated lemon-peel and shallots, each ½ oz., celery seed 1 dr.; infuse in 1 pint of brandy for 10 days.

SPIRIT OF SAVORY SPICES. Infuse half the Savory Ragout Powder (see MIXED SPICES, &c., No. 5, as above) in 1 quart of brandy for 10 days.

KITCHENER'S SOUP HERB AND SAVORY SPICE SPIRIT. A mixture of equal measures of the last two.

CULINARY VINEGARS, SAUCES, &c.

TARRAGON VINEGAR. Put fresh tarragon leaves into a stone jar, and pour on them a sufficient quantity of the best wine vinegar to cover them. Set the jar in a warm place for 14 days; then strain through a jelly-bag. (In the same way may be made elder-flower, basil, green mint, and Burnet vinegars. *Cress and Celery Vinegars* are made with ½ oz. of the bruised seed to a quart of vinegar. *Horseradish Vinegar*, with 3 oz. of the scraped root, 1 oz. of minced shallots, 1 dr. of cayenne, to 1 quart of vinegar. *Garlic Vinegar* is made with 2 oz. of minced garlic to 1 quart

of wine vinegar. *Shallot Vinegar* in the same proportion. *Chili Vinegar*, with 50 English chillies, cut or bruised (or ¼ oz. cayenne pepper), to 1 pint of the best vinegar; digest for 14 days.

CAMP VINEGAR. Take 12 chopped anchovies, 2 cloves of garlic minced, 1 dr. of cayenne, 2 oz. of soy, 4 oz. of walnut catsup, and 1 pint of the best vinegar; digest for 1 month, and strain.

2. Vinegar 1 quart, walnut catsup 1 pint, mushroom catsup 3 tablespoonfuls, garlic 4 heads, cayenne ½ oz., soy 2 tablespoonfuls, port wine 2 glasses, 3 anchovies, and 1 tablespoonful of salt; put them into a bottle, shake daily for a month, and decant.

CURRY VINEGAR. Infuse 3 oz. of curry powder in 1 quart of vinegar, near the fire, for 3 days.

RASPBERRY VINEGAR. Macerate 2 lb of fresh raspberries with 1 pint of the best vinegar for 14 days, and strain; or, to 1 quart of the juice add 2 oz. of strong acetic acid, or enough to render it sufficiently acid.

ESCHALOT WINE. Bruised shallots 3 oz., sherry wine 1 pint; infuse for 10 days: 1 oz. of scraped horseradish and 1 dr. of thin lemon-peel may be added. ("The most elegant preparation of the onion tribe."—Dr. KITCHENER.) Wines of several herbs may be made in the same proportion as the vinegars.

FRENCH MUSTARD. This is sold with a great variety of flavors. A good substitute may be made by mixing good flour of mustard with the liquor of walnut and other pickles; or with the flavored vinegars above. The following is one of the published recipes: Salt 12 oz., scraped horseradish 8 oz., 1 clove of garlic, ½ oz. of sugar, 1 gallon of French vinegar (hot, but not boiling). Macerate for 24 hours, and strain. Mix with flour of mustard, q. s.

MUSTARD FOR THE TABLE. Mix 8 spoonfuls of flour of mustard with 2 of salt, and 9 of water. Mix to a smooth paste, add 6 spoonfuls more of water, and mix.

ESSENCE OF ANCHOVIES. Beat 1 lb of anchovies in a Wedgewood mortar, and put them into a pipkin with 4 oz. of vinegar; boil for a few minutes, and rub the pulp through a hair-sieve. Boil the bones in 1½ lb of water, strain, and add 2 oz. of salt, and 2 oz. of flour of starch, and the

pulped anchovies; let it boil, and pass it through a hair-sieve. It is usually colored with powdered bole, or with annotto. Gum tragacanth is sometimes used to stiffen it instead of flour. Another method is, to simmer anchovies in their own weight of water for 2 or 3 hours, removing any scum that may rise, strain with pressure through a strong canvas bag, and filter through flannel. This has the pure flavor of the fish; but a little cayenne and salt may be added, to preserve it.

ANCHOVY PASTE. Pound the fish in a mortar, and rub the pulp through a fine sieve. Put it into pots, and cover with clarified butter.

MUSHROOM CATSUP. Press the mushrooms in a tincture press, and boil the juice with ½ oz. black pepper-corns, 1 oz. pimento, ½ oz. of ginger, ¼ oz. cloves, 1½ oz. shallots, and 8 oz. of salt to each gallon. Some add 4 oz. of brandy. Or spinkle the mushrooms with salt (1 ℔ to 2 pecks), stir occasionally for 2 days, then squeeze them gently in a hair-sieve, and boil the liquor with pepper and other spices.

WALNUT CATSUP. 1. Mix 6 half-sieves of green walnut shells with 2 or 3 ℔ of salt in a wooden vessel; let them stand 6 days, beating them frequently till they become pulpy; then drain off the juice, and boil it up with 4 oz. of ginger and allspice, and 2 oz. of long pepper and cloves.

2. Juice of walnuts 1 gallon, anchovies 2 ℔, shallots 1 ℔, cloves, mace, and black pepper, of each 1 oz., and 1 clove of garlic. Boil for a short time, and bottle it.

LEMON PICKLE. Slice 6 lemons, rub them with salt, lay them in a stone jar, with 2 oz. each of allspice and white pepper; add ¼ oz. each of mace, cloves, and cayenne, and 2 oz. each of horseradish and mustard seed; pour over them 2 quarts of hot distilled vinegar; and, after standing for a few days, strain. Some add garlic or shallots.

QUIN SAUCE. Mushroom catsup ½ pint, walnut pickle ¼ pint, port wine ¼ pint, 6 anchovies and 6 shallots (both pounded); soy 1 tablespoonful, cayenne ½ dr.; simmer together gently for 10 minutes, strain, and bottle.

WATERLOO SAUCE. Vinegar 4 pints, port wine 1 pint, cayenne 1 oz., walnut catsup ½ pint, mushroom catsup ½ pint,

essence of anchovies 4 oz., powdered cochineal 1 oz., garlic, 12 cloves.

EPICUREAN SAUCE. Indian soy 2 oz., walnut catsup, mushroom catsup, each 8 oz.; port wine 2 oz., bruised white pepper ½ oz., shallots 3 oz., cayenne ¼ oz., cloves ½ oz. Macerate for 14 days in a warm place, strain, and add white wine vinegar to make up a pint.

SAUCE SUPERLATIVE (Dr. KITCHENER'S). Port wine, and mushroom catsup, of each a pint: walnut or other pickle liquor ½ pint, pounded anchovies 4 oz.; fresh lemon-peel cut thin, sliced shallots, and scraped horseradish, of each 1 oz., allspice and black pepper, of each ½ oz.; cayenne 1 dr.; curry powder 3 dr.; celery seed 1 dr.; put them into a wide-mouthed bottle, stop it close, shake daily for a fortnight, and strain: a ¼ pint of soy may be added.

(A variety of sauces may be made by mixing, in different proportions, the ingredients of the last 4 sauces.)

CASSAREEP. The expressed juice of the roots of the bitter cassava; used as a condiment in the West Indies.

SOY. Boil a gallon of the seeds of Dolichos soja till soft, add a gallon of bruised wheat, keep them in a warm place for 24 hours; add a gallon of salt, and 2 gallons of water, and after keeping them bunged up in a stone jar for 2 or 3 months, press out the liquor.

PICKLES.

A few recipes are here given as illustrations of the methods employed in preparing these condiments. For full particulars the reader is referred to the popular treatises on Cookery. The best vinegar (pickling, or No. 24 vinegar) should be employed. Some prefer the crystal or white vinegar (distilled vinegar or rather pure diluted wood-vinegar), especially for white pickles; but the best wine vinegar is more agreeable. Stoneware jars, not glazed with lead should be used to keep the pickles in; or otherwise green glass jars.

SPICED VINEGAR, FOR PICKLES GENERALLY. Bruise in a mortar 2 oz. of black pepper, 1 oz. of ginger, ½ oz. of allspice, and 1 oz. of salt. If a hotter pickle is desired, add ½ dr. of cayenne, or a few capsicums. For walnuts

add also 1 oz. of shallots. Put these into a stone jar, with a quart of vinegar, and cover them with a bladder wetted with the pickle, and over this a piece of leather. Set the jar on a trivet near the fire for 3 days, shaking it 3 times a day, then pour it on the walnuts or other vegetables. For walnuts it is used hot, but for cabbage, &c., cold. But to save time, it is usual to simmer the vinegar gently with the spices; which is best done in an enamelled saucepan.

Beet Root. Boil the roots till 3 parts done (from 1½ to 2½ hours); then take them out, peel them, and cut them in thin slices. Put them into a jar, and pour on them sufficient cold spiced vinegar (as above) to cover them.

Cabbage, White. Cut it into thin slices, put them into an earthen pan, sprinkle them with salt, and let them lie for 2 days; then drain them, and spread them out before the fire for some hours; put them into a stone jar, and add sufficient white vinegar, or pale white vinegar, to cover them, and a little mace and white pepper-corns.

Cabbage, Red. Remove the outer leaves and stalks, and cut the cabbage in quarters, then shred them into a cullender, and sprinkle them with salt; next day drain them, put them into a jar, and pour on them sufficient cold spiced vinegar to cover them. Others hang up the cabbage for a few days to dry, then shred the leaves, and put them in layers in a jar with a little salt, pepper, and ginger, and fill up with cold vinegar. Others use vinegar without spice.

Cauliflower and Broccoli. These should be sliced, and salted for 2 or 3 days, then drained, and spread upon a dry cloth before the fire for 24 hours; then put into a jar, and covered with spiced vinegar. Dr. Kitchener says, that if vegetables are put into cold salt and water (a ¼ lb of salt to a quart of water) and gradually heated to a boiling heat, it answers the same purpose as letting them lie some days in salt.

Cucumbers. Gherkins. Small cucumbers, but not too young, are wiped clean with a dry cloth, put into a jar, and boiling vinegar, with a handful of salt, poured on them. Boil up the vinegar every 3 days, and pour it on them till they become green; then add ginger and pepper, and tie them up

close for use. Or cover them with salt and water (as above) in a stone jar, cover this, and set them on the hearth before the fire for 2 or 3 days, till they turn yellow; then put away the water, and cover them with hot vinegar, set them near the fire, and keep them hot for 8 or 10 days, till they become green; then pour off the vinegar, cover them with hot spiced vinegar, and keep them close.

MANGOES. Large cucumbers, or small melons, are split so that a marrow-spoon may be introduced, and the seeds scooped out; they are then parboiled in brine strong enough to float an egg, dried on a cloth before the fire, filled with mustard-seed and a clove of garlic, and then covered with spiced vinegar. True mangoes the same.

MUSHROOMS. Clean them with water and flannel, throw them into boiling salt and water in a stewpan, and let them boil for a few minutes. Drain them in a cullender, and lay them on a linen cloth, covering them with another. Put them into bottles with a blade or two of mace, and fill up with white vinegar, pouring some melted mutton fat on the top, if intended to be kept long.

NASTURTIUMS, FRENCH BEANS, and other small green vegetables are pickled in the same way as Gherkins.

ONIONS. 1. Let them lie in strong salt and water for a fortnight; then take them out and peel them; put them in fresh salt and water for another fortnight; take them out, wash them clean, and let them lie in fresh water all night. Next day put them on a cloth to drain; then put them in a jar, and pour over them hot spiced vinegar. If you wish them of a nice color, use white vinegar.

2. Peel small silver button onions, and throw them into a stewpan of boiling water; as soon as they look clear take them out with a perforated spoon, and lay them on a folded cloth, covered with another, and when quite dry, put them into a jar, and cover them with hot spiced vinegar. When quite cold, bung them down, and cover with bladder wetted with the pickle, and leather.

WALNUTS. Take 100 young walnuts, lay them in salt and water for 2 or 3 days, changing the water every day. (If required to be soon ready for use, pierce each walnut with a larding pin, that the pickle may penetrate.) Wipe them

with a soft cloth, and lay them on a folded cloth for some hours. Then put them in a jar, and pour on them sufficient of the above spiced vinegar, hot, to cover them. Or they may be allowed to simmer gently in strong vinegar, then put into a jar with a handful of mustard seed, 1 oz. ginger, ¼ oz. mace, 1 oz. allspice, 2 heads of garlic, and 2 split nutmegs, and pour on them sufficient boiling vinegar to cover them. Dr. KITCHENER recommends the walnuts to be gently simmered with the brine, then laid on a cloth for a day or two, till they turn black, put into a jar, and hot spiced vinegar poured on them.

TOMATOES. As Gherkins. See CUCUMBERS.

PICCALILLI, INDIAN, OR MIXED PICKLE. 1. To each gallon of strong vinegar put 4 oz. of curry powder, 4 oz. of good flour of mustard, 3 oz. of bruised ginger, 2 oz. of turmeric, 8 oz. of skinned shallots, and 2 oz. of garlic (the last two slightly baked in a Dutch oven), ¼ lb of salt and 2 dr. of cayenne pepper. Digest these near the fire, as directed above for spiced vinegar. Put into a jar gherkins, sliced cucumbers, sliced onions, button onions, cauliflower, celery, broccoli, French beans, nasturtiums, capsicums, large cucumbers, and small melons. All, except the capsicums, to be parboiled in salt and water, drained, and dried on a cloth before the fire. The melons and large cucumbers to be prepared as directed above for mangoes. Pour on them the above pickle.

2. Take 1 lb of ginger-root, and ½ lb of garlic (both previously salted and dried), 2 gallons of vinegar, ½ oz. of turmeric, ¼ lb of long pepper. Digest for 2 or 3 days near the fire in a stone jar; or gently simmer them in a pipkin or enamelled saucepan. Then put in the above vegetables, or almost any except red cabbage and walnuts, all previously salted and dried.

BRINE, OR PICKLE, FOR PORK, &C. Brown sugar, bay salt, common salt, of each 2 lb, saltpetre ½ lb, water a gallon. Boil gently, and remove the scum. Another meat pickle is made with 12 lb salt, 2 lb of sugar or treacle, ½ lb of nitre, and sufficient water to dissolve it. To cure HAMS, mix 5 oz. of nitre, with 8 oz. of coarse sugar; rub it on the ham, and in 24 hours rub in 2 lb of salt, and in a fortnight 2 lb more. The above is for a ham of 20 lb; it should lie in the salt a month or 5 weeks.

WESTPHALIAN ESSENCE.—CAMBRIAN ESSENCE OF WOOD SMOKE. These appear to be crude pyroligneous acid or wood vinegars, and are used to give to hams, &c., the smoked flavor. The following has been published as the recipe for Essence of Smoke, but we apprehend it is far from being correct: Macerate for several weeks ½ dr. of Barbadoes tar, 1 dr. of liquid burnt sugar, 5 dr. each of port wine and vinegar, 2 dr. of salt, and 7 oz. of water.

TRADE CHEMICALS.

MISCELLANEOUS PREPARATIONS AND COMPOUNDS EMPLOYED IN THE ARTS, IN DOMESTIC ECONOMY, CHEMICAL RESEARCH AND AMUSEMENT, &c.

This division of the work comprises those chemical compounds which are employed for other purposes than those of medicine, and which have not been noticed in the former parts of this volume. It includes a variety of miscellaneous articles which are sometimes sold by the retail druggists, or the materials of which he is expected to furnish, or with the composition of which it is desirable he should be acquainted. The limits of the work do not admit of a minute description of the processes and manipulations employed in the manufacture of such chemicals as are only made on the large scale, and never by the retailer; nor of those chemical arts which have no immediate connection with the trade.

Acetates. Such as are employed in medicine will be found in the Pocket Formulary. The only Acetates requiring notice here are the following:

Acetate of Alumina. This is made, for the use of dyers and calico printers, by decomposing acetate of lime with alum. It may be conveniently made by adding to a boiling solution of 5 parts of alum, a solution of 6 parts of sugar of lead. When the mixture is cold, the clear liquid is poured off; from which the dry salt may be obtained by careful evaporation. It contains, besides acetate of alumina, some sulphate of potash.

Acetate of Iron, or Iron Liquor. Usually obtained, for the use of dyers, by digesting scraps of iron in redistilled wood-vinegar. (See *Mordants, Iron Liquors.*)

ACETATE OF LIME. Impure acetate (or pyrolignite) of lime, is made by neutralizing pyroligenous acid with cream of lime or chalk, and evaporating to dryness. By using pure acetic acid a purer acetate is obtained.

ACETATE OF SODA. By mixing the above impure acetate of lime, in solution, with a solution of sulphate of soda, filtering and evaporating the clear liquid, an impure acetate of soda is obtained; which by repeated crystallization is rendered colorless, and fit for yielding pure concentrated acetic acid by distillation with sulphuric acid.

ACETIMETRY. The strength of vinegar is estimated for the duty by an instrument named an acetimeter, which determines the quantity of acetic acid present by the specific gravity of the vinegar after neutralization by slaked lime. Dr. URE'S plan is to add to a given weight of vinegar bicarbonate of potash till exactly neutralized; every 2 gr. of the bicarbonate indicate 1 grain of real acetic acid. In this and the following operations it is convenient to use a tube graduated into 100 equal divisions, numbered from the top downwards (see ALKALIMETRY, below). The quantity of test solution used is then seen at once. In the present case the 200 gr. of the alkaline carbonate being dissolved in sufficient water to fill the graduated portion of the measure, each of the divisions used in neutralizing 100 gr. of vinegar is equivalent to 1 per cent. of absolute acetic acid.

ACID, ACETIC. See VINEGAR. For the methods of procuring this concentrated acetic acid, see Acidum Aceticum, Pocket Formulary. The process of the Edinburgh Pharmacopœia yields a stronger acid than that of the London Pharmacopœia. A strong acid, very suitable for making aromatic spirit of vinegar, is procured by distilling crystallized verdigris in an earthen retort coated with clay, into a series of 3 globes, connected by opposite tubulures, and kept constantly cool, the last being furnished with a WELTER'S safety tube. The acid which comes over is usually colored, and requires to be rectified by a slow and careful redistillation in a glass retort. Acetic acid of moderate strength may be rendered stronger by redistilling it over acetate of potash, rejecting the first portions that come over, and taking care that the temperature does not rise above 572° F. By redistilling it,

rejecting the first and last portions, glacial acetic acid is procured. The same acetate of potash may be used repeatedly. The process of the Dublin Pharmacopœia yields a good product of glacial acetic acid.

Acid, Chloric. Dissolve 7 parts of crystallized carbonate of soda, and 7½ of tartaric acid in 24 of boiling water; add to the boiling solution 6 parts of chlorate of potash in 16 of water, at 212°, agitating the mixture. When quite cold filter, and add a solution of 6 parts of oxalic acid in 18 of water, heated not above 134°. Agitate well, and place the vessel in a freezing mixture of muriatic acid and sulphate of magnesia, and filter. (Not absolutely pure, but sufficiently so for technical purposes. It may be obtained pure by decomposing chlorate of barytes by sulphuric acid.)

Acid, Chromic. It may be obtained pure by mixing bichromate of potash with nitrate of silver in solution, washing the precipitate, and decomposing it with an equivalent quantity of muriatic acid. In a few minutes the clear solution may be poured off. A cheaper method, where great purity is not required, is to add from 120 to 150 volumes of strong sulphuric acid to 100 volumes of a cold saturated solution of bichromate of potash. Dry the acid which separates on porous tiles. Or add 2 parts of concentrated sulphuric acid to 1 part of dry chromate of lead, leave the paste for 12 hours, then treat it with water; decant the clear red liquid, and evaporate it in a retort. Keep it boiling for some time, then allow it to cool. Most of the acid separates in crystals; more may be obtained by evaporating the solution till its density is 1.55.

Acid, Cinnamic. It is most readily procured by distilling genuine balsam of Tolu by a gentle heat. The white crystalline mass which condenses on the neck of the retort is purified by pressing it between blotting-paper, dissolving in boiling water, and crystallizing.

Acid, Fluoric. The anhydrous acid is made by distilling powdered fluor spar with twice its weight of oil of vitriol in a leaden, or better, a *silver* alembic, the pipe of which fits into a bottle of the same material, surrounded with ice. But as it is usually required in a diluted state, water equal in weight to the spar may be put into the receiver. Great

care must be taken, as the acid, both in its gaseous and liquid form, is very destructive.

Acid, Hippuric. Mix the urine of the horse with milk of lime, boil for some minutes, and strain. Boil down the clear liquid to $\frac{1}{8}$ of its bulk, avoiding burning; add hydrochloric acid, press the impure acid, boil it with fresh milk of lime, and again precipitate with hydrochloric acid.

Acid, Iodic. Mr. A. Connell's *Method.* Put 50 gr. of iodine into a large, tall flask: add 1 oz. of fuming nitric acid, boil, and as the iodine sublimes and condenses on the sides of the flask, continually wash it back again with the acid. Continue this until none of the iodine remains unchanged. Then pour the whole into a shallow evaporating dish, and evaporate to dryness. Redissolve, and again evaporate till all the nitrous acid is got rid of.

Acid, Muriatic, or Hydrochloric. Commercial muriatic acid is largely produced by the action of sulphuric acid on common salt, in the manufacture of sulphate of soda for the purpose of making soda ash and washing soda by the decomposition of that salt. From the impurity of the ingredients it is apt to be contaminated with arsenic and sulphurous acid; as well as sulphuric acid and iron. It may be purified from arsenic by redistilling it over strips of bright copper. Dr. Gregory's method of procuring *pure* muriatic acid is as follows: Put into a matrass 6 parts, by weight, of purified salt, and 10 oz. of oil of vitriol previously diluted with 4 of water, and cooled. Fix in the matrass a tube twice bent at right angles and having a bulb blown on the descending limb. Into a bottle surrounded with ice and water introduce distilled water equal in weight to the salt employed, and let the bent tube dip $\frac{1}{8}$ of an inch into the water. Apply a gentle heat of a sand-bath to the matrass as long as acid comes over. In about 2 hours the operation will be finished. The water is increased $\frac{2}{3}$ in bulk, and converted into hydrochloric acid of 1.14 or 1.15 sp. gr. To procure it of 1.21 sp. gr. employ part of this acid during the first half of a similar operation, and it will be speedily saturated. See also Acidum Hydrochloricum Purum, Pocket Formulary. Mr. Phillips says a perfectly colorless acid may be obtained from the commercial sulphuric acid and common salt.

Acid, Nitric, and Fuming Nitrous Acid. Put into an iron or stoneware pot, nitre or nitrate of soda, add rather more than half its weight of strong sulphuric acid, and lute on a stoneware head. The vapor is conducted into a series of two-necked stone-ware vessels, containing each $\frac{1}{6}$th of their capacity of water. The acid is usually obtained of the density of about 1.45. It is colored with nitrous acid gas, forming what is commonly, but improperly, termed *nitrous acid.* By gently heating the colored acid in a retort, the nitrous acid is driven off, and the acid remains nearly colorless, usually of the density of 1.38 to 1.42. This is weaker than the Pharmacopœia directs, but sufficiently strong for most purposes. (See Acidum Nitricum, Pocket Formulary.) Its strength may be increased by mixing it with its volume of strong oil of vitriol, and slowly distilling off $\frac{2}{3}$ds of the nitric acid. This yields an acid of 1.5 sp. gr., such as is required for the preparation of gun-cotton, as well as in some pharmaceutical processes.

Acid, Nitro-Muriatic. *Aqua Regia.* This is used in the arts, chiefly as a solvent for gold. By the mutual action of nitric and muriatic acids a compound of chlorine, nitrogen, and oxygen is formed. The best proportions and strength of the acids are variously stated. Colorless nitric acid must be used. Mr. Elkington employs 21 parts of nitric acid, sp. gr. 1.45; 17 parts of muriatic acid, 1.15 sp. gr.; and 14 parts of water. This dissolves 5 parts of gold. For the nitro-muriatic acid employed by dyers as a solvent for tin, see Dyes, &c., below.

Acid, Oxalic. Digest by the aid of heat 1 part of treacle, or of potato starch, in 5 parts of nitric acid, sp. gr. 1 42, diluted with 10 parts of water, so long as gaseous products are evolved. By evaporation the acid is obtained in crystals, and must be recrystallized till sufficiently pure. Mr. Lewis Thompson directs 28 oz. of sugar, and 184 oz. of nitric acid of 1.245 sp. gr., to be digested at 125° F. It yields 30 or 31 oz. of acid. M. Schlesinger directs 4 parts of dry sugar, and 33 of nitric acid of 1.38 sp. gr. to be boiled to one-sixth of the original volume, and allowed to crystallize. This is the best method of operating on a small scale, when the amount of product is not the principal object.

Acid, Phosphoric. See Pocket Formulary. *Dry* phosphoric

acid is thus obtained: On a flat plate place a large bell glass, and under it a small porcelain cup or crucible. Introduce into the latter a piece of phosphorus, dried with blotting-paper, and set it on fire by a heated wire. Let the bell glass be raised on one side to admit sufficient air to maintain combustion; and as the phosphorus is consumed introduce successive pieces, taking care that the glass does not become too hot. When the quantity of acid is considerable, knock it from the plate with an iron spoon, and put it into stoppered bottles. Several glasses may be used at once. It is used as a dissecting body, having the strongest attraction for water of any known substance. Also in making a stopping for teeth—see TEETH CEMENTS.

ACID, PYROGALLIC. Heat powdered nutgalls in a dish covered with thin filtering paper pasted to its edges, and surmounted with a bell-formed receiver, kept cool. A solution of the condensed acid, decolorized by animal charcoal, and mixed with spirit, is used to stain the hair (and skin) brown.

ACID, SULPHURIC. This is only made on a large scale; but the commercial acid requires purification for many chemical as well as pharmaceutical purposes, as it is usually contaminated with tin and lead, and frequently with arsenic, or selenium, and compounds of nitrogen. The purification of oil of vitriol by distillation is attended with some difficulty. No luting must be employed; and to prevent the violent jumpings which attend the ebullition, strips of platina, or fragments of rock crystal should be introduced into the retort. The receiver should be large, and the whole defended from currents of cold air. The first portions which come over should be rejected till the indigo test proves it to be free from nitric acid. By boiling a portion of the acid with a few drops of solution of sulphate of indigo, the latter is discolored if nitric acid is present. The Edinburgh Pharmacopœia directs the nitrous acid to be got rid of by heating 8 fluid ounces of commercial oil of vitriol to near the boiling-point with from 10 to 15 gr. of sugar. When sulphuric acid is diluted with water, the *metallic* impurities may be removed by a little solution of sulphuret of barium, and allowing the precipitates to subside, when the pure diluted acid may be decanted for use.

Dry or Anhydrous Sulphuric Acid. Into a retort, placed in a freezing mixture, and having a receiver attached, put some dry phosphoric acid (see above), and add ¾ of its weight of strong sulphuric acid. Remove the retort from the freezing mixture, and place the receiver attached to it there; a gentle heat being now applied to the retort, the anhydrous acid is obtained in silky crystals.

Acid, Sulphurous. For the mode of obtaining an aqueous solution of this acid, see Acidum Sulphurosum, Pocket Formulary. The following are cheaper methods of obtaining it for bleaching purposes, &c. Berthier directs a mixture of 100 parts of black oxide of manganese, and 12 or 14 of sulphur, to be heated in a glass retort, and the gas received into water kept very cold. Mr. Redwood directs ½ oz. of powdered charcoal to be acted on by 4 fluid oz. of oil of vitriol. Treacle is sometimes used instead of charcoal; as also is linseed oil.

Acid, Tannic. *Tannin.* Place coarsely-powdered Aleppo galls in a damp cellar for 3 or 4 days, then mix them with sufficient sulphuric ether to form a soft paste. Place this in a close vessel for 24 hours, then wrap it in linen ticking, and submit it to the action of a powerful press. Scrape off the tannin from the surface of the ticking, remove the *cake* from within it, rub this into powder, form it into a paste with a mixture of 100 parts of ether and six of water, well shaken together before pouring it on the galls, and proceed as before. Let the syrupy liquid thus obtained be thinly spread on plates, and dried in 113° Fahrenheit.

Acids, Mixed, for Galvanic Batteries. 1. For troughs, for general purposes, medical galvanism, &c.: Nitric acid 1 fluid oz., sulphuric acid 1½ fluid oz., water 4 pints.

2. Dr. Faraday. Oil of vitriol 2 fluid oz., nitric acid 1 fluid oz., water 5 pints. It should be tried by dipping into it a piece of sheet zinc. A continuous succession of *small* bubbles should be produced.

3. For Mr. Smee's Battery. One measure of sulphuric acid to 7 of water. The intensity of its action is increased by the addition of a few drops of nitric acid, but this tends to destroy the plates. In electro-metallurgy the water should only contain a sixteenth of sulphuric acid.

4. For Mr. GROVE'S Battery. For the outer vessel 1 part sulphuric acid to 7 of water; for the inner, concentrated nitric acid.

6. For DANIELL'S Battery. For the porous tube containing the zinc, 1 part of sulphuric acid with 10 of water. For the outer cylinder, a saturated solution of sulphate of copper, with a tenth part of sulphuric acid.

6. Nitro-sulphuric acid, for Dr. T. WRIGHT'S Batteries. Nitric acid 1 part, sulphuric acid 5 parts. The zinc plate is immersed in a solution of muriate of ammonia or of salt, the platinized zinc in the above acid. The platinizing requires to be repeated every time the plate is washed.

ACIDIMETRY. Acids generally are estimated by the quantity of alkalies or carbonated alkalies required to neutralize them. Weigh 100 gr. of the acid and dilute it with a few times its weight in water. Then add sufficient dry or crystallized carbonate of soda, or carbonate of potash, to exactly neutralize the acid. The Alkalimeter tube may be used for the solution of the alkali. By a reference to the table of chemical equivalents, the quantity of real acid of any kind represented by the quantity of alkali required to neutralize it may be estimated.

ACIDULATED KALI. See BEVERAGES.

ALBUMINOUS SIZE. Beat up the white of an egg with twice its bulk of cold water, until well incorporated. Used as a varnish for leather binding and kid gloves; also to size drawing paper.

ALCOHOL. There is, perhaps, no better method of obtaining absolute alcohol, then that of the Edinburgh Pharmacopœia. See Alcohol, Pocket Formulary.

ALKALIMETRY. The quantity of real alkali contained in the commercial alkalies (common soda, soda ash, potashes, pearlash, salt of tartar, &c.) is ascertained by the quantity of an acid solution of a known strength required to neutralize it. For this purpose a tube, termed an ALKALIMETER, is used, which will hold 1000 grains of water. The tube may be three-fourths of an inch internal diameter, and 9½ inches in length; or five-eighth inches diameter, and 14 inches in height; and should be graduated into 100 equal divisions numbered from the top downwards. The quantity of test acid used is then at once seen. This consists of sulphuric acid diluted with water so that each measure (10

grains) is equivalent to one grain of pure soda. To use it, dissolve 100 grains of the impure soda in 3 oz. of hot water, filter, and wash the filter. Then add to the solution the test acid until the litmus or cabbage-paper ceases to show an alkaline reaction. The same acid will serve for potash, if the number be multiplied by 3 and divided by 4.—PARNELL. There are several other methods of performing the process.

ALLOYS AND AMALGAMS. A few only of these metallic compounds require notice here:

Fusible Metal. 1. Tin 8 parts, lead 4, bismuth 3; melt together, removing the scum. Used as a metal-bath.

2. DARCET'S, for the same purpose; Bismuth 8 parts, lead 5, tin 3.

3. Lead 3 parts, tin 2, bismuth 5. This melts at 197° Fahrenheit.

4. For anatomical injections: Melt together with a gentle heat 174 parts of tin, 312 of lead, 514 of bismuth, with a little charcoal; remove from the fire, and add 100 parts of mercury previously heated. It is fluid at 173°; solid at 140° Fahrenheit.

Bronze. 1. For medals and small castings: Copper 95, tin 4.

2. Copper 89, tin 8, zinc 3.
3. *Ancient.* Copper 100, tin 7, lead 7.
4. KELLY'S. Copper 91, zinc 6, tin 2, lead 1.
5. For gilding. Copper 14, zinc 6, tin 4.
6. *Bell-Metal.* Copper 78, tin 22.

German Silver. 1. Copper 40½, nickel 31½, zinc 25½, iron 2½.

2. Pure copper 55, nickel 23, zinc 17, iron 3, tin 2.

Gold, Factitious. Platina 7, copper 16, zinc 1; fuse together. See AURUM MUSIVUM.

Common Gold. Copper 16, silver 1, gold 2.

Or-molu Copper 45 to 48, zinc 52 to 55.

White Brass. POREL. Melt zinc with 10 per cent. each of copper and iron. This alloy has the fracture and appearance of zinc, but is tougher than cast iron. It does not rust, nor adhere to metal moulds.

Solders. 1. For Gold: Pure gold 12 parts, silver 2, copper 4.

2. *Soft Solder.* Tin 2 parts, lead 1.

3. For brass: Brass 2 parts, zinc 1.

Alloys for Electrotype. Clichée Moulds. Bismuth 8 parts, tin 4, lead 5, regulus of antimony, 1; melt repeatedly together, and pour out in drops, till perfectly mixed.

Amalgam for Electrical Machines. 1. Fuse 1 oz. of zinc with ½ oz. of tin, at as low a temperature as possible; then add 1½ oz. of quicksilver, previously made hot; mix, pour out, and when cold reduce it to a powder, and triturate it with sufficient quicksilver to bring it to a proper consistence.

2. Zinc 1 part, tin 1, quicksilver 2; melt together.

3. Zinc 2, tin 1, mercury 5.

4. La Beaume's. Pour into a chalked wooden box 6 oz. of quicksilver; pour into an iron ladle ½ oz. beeswax, with 2 oz. purified zinc, and 1 oz. of grain tin; set it over a brisk fire, and when the metals are melted, pour them into the box, avoiding the dross. When cold, reduce it to a powder, and mix it with lard. Keep it in a box, covered with tallow, and spread it on leather for use.

Liquid Amalgam, for Silvering Globes, &c. Pure lead 1 oz., grain tin 1 oz.; melt in a clean ladle, and immediately add 1 oz. of bismuth. Skim off the dross, remove the ladle from the fire, and, before the metal sets, add 10 oz. of quicksilver. Stir together, avoiding the fumes.

Amalgam for Varnishing Figures. Melt 2 oz. of tin with ½ oz. of bismuth, and add ½ oz. of quicksilver. When cold, grind it with white of egg, and apply to the figure.

Alum. It is prepared by lixiviating calcined albuminous schist, and concentrating the solution to 1.4 or 1.5 density, and adding the requisite quantity of muriate of potash, soap-boilers' ash, or kelp, to supply the alkali. By recrystallization it is obtained colorless. In some manufactories sulphate of ammonia, from gas-liquor, is added to the lixivium, instead of muriate of potash. Roman or cubic alum is crystallized from a solution, the temperature of which is not allowed to exceed 104° F. It differs from common alum in containing a large quantity of base, a portion of which separates if the solution be heated to 120°.

Another kind of alum, sometimes used as a mordant, consists almost entirely of sulphate of alumina, and is probably made by boiling fine clay, free from iron, with sulphuric acid, and cooling the solution so as to obtain a solid mass. See DYES, &c.

ALUMINUM, TO FROST. The metal is plunged into a solution of caustic potash. The surface, becoming frosted, does not tarnish on exposure of the air.—MACADAM.

ALUMINUM BRONZE. 10 parts of aluminum are melted with 90 parts of copper. It is said to be as tenacious as steel.

AMADOU. Prepared from Boletus igniarius, B. fomentarius, and some other allied species of fungi. The fungus is cut into thin slices, the hard external parts removed, and the rest beaten with a mallet till soft. This forms *surgeons' agaric*. If intended for *German tinder*, it is soaked in a solution of nitre, and sometimes sprinkled with gunpowder, and carefully dried.

AMALGAMS. See ALLOYS, above.

AMMONIA, SULPHATE AND CARBONATE OF. An impure sulphate of ammonia, suitable for agricultural purposes, is obtained by neutralizing the ammoniacal liquor of gas-works with sulphuric acid. By recrystallization it may be obtained in a state of greater purity. The carbonate (hydrated sesquicarbonate) is obtained by mixing either this sulphate, or sal-ammoniac, with chalk, and subliming it in iron retorts into leaden receivers. It is further purified by resubliming it with a gentle heat. A sketch of the apparatus employed will be found in Dr. PEREIRA'S "Elements," Vol 1.

ANATOMICAL INJECTIONS. 1. Tallow, resin, and wax, equal parts; melt over a slow fire; and add red lead or vermilion sufficient to color. For coarse preparations.

2. A strong solution of isinglass, colored as required. For decicate preparations.

3. AMALGAM INJECTION. Melt together 1 oz. each of bismuth, lead, and zinc, and, when melted, add 2 oz. of quicksilver. Also for delicate parts.

ANATOMICAL SUBJECTS, AND ANIMAL SUBSTANCES, TO PRESERVE. 1. M. GANNAL'S *Solution*. Common salt 2 lb, alum 2 lb, nitre 1 lb, water 4 gallons. M. GANNAL injects

into the carotid artery a solution of sulphate of alumina, of density 1.286. From 5 to 7 pints are required in summer, but less will suffice in winter.

2. Dr. BABINGTON injects pyroxylic spirit into the aorta, and a little into the cavity of the peritoneum and the rectum.

3. Mr. GOADBY, for insects, and for preparations of their organs. Bay salt 4 oz., alum 2 oz., corrosive sublimate from 2 to 4 grains, water 1 to 2 quarts. The weaker proportions should always be employed in the first instance. Let the insect or its organs be covered with the fluid, which should be changed frequently.

4. *For Mollusca.* Bay salt ½ oz., arsenic ½ dr., sublimate 2 gr., water a quart: dissolve.

5. Mr. PIGNE, for preserving pathological specimens. Creasote 3 to 6 drops, water a pint.

6. Carbolid acid, often sold for creasote, may be used in the same manner.

7. Pure Glycerine. Found especially useful in preserving the fresh tints of fishes, &c., intended for exhibition in museums.

8. Dr. STAPYLTON, for pathological specimens. In a quart of saturated solution of alum dissolve ½ dr. of nitre. A recent preparation immersed in this liquid become discolored; but within a few days the color returns. It is then put in a saturated and filtered solution of alum. M. REBOULET proposes: Water 16 parts, chloride of lime 4, alum 2, nitre 1.

9. Chloride of tin 4 (or corrosive sublimate 5) parts, in 100 of water, with 2 of muriatic acid.—Mr. COOLEY.

10. *For preserving Animals.* Alum 32 oz., nux vomica 3 oz., water 5 pints; boil to 4½ pints. When cold, filter, or decant. This serves for injection. The residue, mixed with yolk of egg, is used for anointing the interior of the skin and fleshy parts left in skinning animals.

11. *For preserving Feathers.* Strychnine 16 gr., rectified spirit a pint (dangerous). See STUFFING BIRDS, &c.

ANNOTTO, PURIFIED. To a boiling solution of pearlash add as much annotto as it will dissolve. When cold, decant the clear solution, and neutralize with diluted sulphuric acid, avoiding any excess. Wash the precipitate with a little cold water, and dry it.

ANNOTTO, SOLUTION OF. Boil equal weights of annotto and pearlash with water, and dilute to the required color.

ANTIATTRITION, AND AXLE GREASE. 1. One part of fine black-lead, ground perfectly smooth, with 4 parts of lard. Some recipes add a little camphor.

2. BOOTH'S AXLE GREASE. (Expired Patent.) Dissolve ½ lb common soda in 1 gallon of water, and 3 lb of tallow, and 6 lb palm oil (or 10 lb palm oil only); heat them together to 200 or 210° F.; mix, and keep the mixture constantly stirred till the composition is cooled down to 60 or 70°. A thinner composition is made with ½ lb of soda, a gallon of water, a gallon of rape oil, and ¼ lb of tallow or palm oil. (See also Lubricating Compounds.)

ANTIFERMENT. Sulphite of lime; or equal parts of sulphite of lime and ground black mustard seed. Used to check the fermentation of cider, &c.

AQUA FORTIS. Double aqua fortis is nitric acid of 1.36 specific gravity; single aqua fortis, about 1.22.—Dr. PEREIRA. A compound acid was formerly used under this name by dyers, and for cleaning brass, consisting of strong spirit of nitre 20 lb, oil of vitriol 7 lb, water 30 lb.—ELABORATORY LAID OPEN.

AQUA REGIA. See NITRO-MURIATIC ACID, above.

AQUARIUM, MARINE, SALTS FOR, &c. Artificial sea-water may be used instead of the natural. A rough imitation is formed by mixing 100 oz. of fresh water with 3 oz. common salt, 1 oz. of Epsom salts, 200 gr. of chloride of magnesium, and 40 gr. of chloride of potassium.

Or, more precisely, the real constitution of sea-water may be imitated in the following manner: Mix with 970,000 grains of rain-water 27,000 of chloride of sodium, 3600 of chloride of magnesium, 750 of chloride of potassium, 29 of bromide of magnesium, 2300 of sulphate of magnesia, 1400 of sulphate of lime, 35 of carbonate of lime, 5 of iodide of sodium. These all being finely powdered and mixed first, are to be stirred into the water, through which a stream of air may be caused to pass from the bottom until the whole is dissolved. On no account is the water to be boiled, or even heated.

Into this water, when clear, the rocks and seaweed may be introduced. As soon as the latter are in a flourishing state, the animals may follow. Care must be taken not to have too many of these, and to remove immediately any that die. The loss by evaporation is to be made up by adding

clean rain-water. The aquarium, whether of fresh or of salt water, will require occasionally artificial aeration. This may be done by simply blowing through a glass tube which reaches to near the bottom, or, better still, in the following way: Take a glass syringe which can be easily worked. Having filled it with water, hold it with the nozzle about 2 inches from the surface of the water in the aquarium, into which the contents are to be discharged quickly, and with a sort of jerk. By this means a multitude of small bubbles are forced down into the fluid. This operation should be repeated for a considerable number of times.

ARABINE. Gum arabic dissolved in water, and precipitated by alcohol.

ARGENTUM MUSIVUM. Fuse ½ oz. each of grain tin and bismuth in a crucible, and add ½ oz. of mercury.

ARBOR DIANÆ. See TREES, Metallic.

AROMATIC PASTILS. See PERFUMERY.

AURUM MUSIVUM. Mosaic gold, or bisulphuret of tin. See Stanni Sulphuretum, Pocket Formularly. 1. Dr. URE directs 12 oz. of tin to be melted, and 3 oz. of mercury added. This amalgam is triturated with 7 oz. of sulphur and 3 of sal ammoniac, and the powder put into a matrass, which is bedded deep in sand, and kept for several hours at a gentle heat. The heat is then raised, and continued for several hours, taking care not to raise it so high as to blacken the mass.

2. Melt together in a crucible, over a clear fire, equal parts of sulphur and the white oxide of tin; keep it continually stirred with a glass rod, until the compound appears as a yellow flaky powder. (This is used as a cheap bronze powder, &c.)

BALDWIN'S PHOSPHORUS. Heat nitrate of lime till it melts; keep it fused for 10 minutes, and pour it into a heated iron ladle. When cool, break it into pieces, and keep it in a closely-stoppered bottle. After exposure to the sun's rays, it emits a white light in the dark.

BALLS FOR HORSES. See Veterinary Formulary.

BALLS, ASH. The ashes of fern, or other kinds of wood ashes, made into balls.

BALLS, HEEL. 1. Melt together 4 oz. of mutton suet, 1 oz.

of beeswax, 1 oz. of sweet oil, ½ oz. oil of turpentine, and stir in 1 oz. of powdered gum arabic, and ½ oz. of fine lamp-black.

2. Beeswax 8 oz., tallow 1 oz., powdered gum 1 oz., lamp-black q. s. These are used not merely by the shoemaker, but to copy inscriptions, raised patterns, &c., by rubbing the ball on paper laid over the article to be copied. ULLATHORNE'S Balls answer the purpose very well. For copying ancient monumental brasses, a similar compound, colored with bronze-powder instead of lamp-black is sometimes employed.

BALLS FOR SCOURING—BREECHES BALLS, CLOTHES BALLS. 1. Bath-brick 4 parts, pipe-clay 8 parts, pumice 1, soft-soap 1; ochre, umber, or other color to bring it to the desired shade, q. s.; ox-gall to form a paste. Make it into balls, and dry them.

2. Pipe-clay 4 oz., fuller's earth ½ oz., whiting ½ oz. white pepper ¼ oz., ox-gall sufficient to form it into a paste.

3. Pipe-clay 3 oz., white pepper 1 dr., starch 1 dr., orris powder 1½ dr. It may be kept in powder, or formed into balls, as above.

BALLS, BLACKING. See BLACKING, below.

BALLS, FURNITURE. See FURNITURE PASTE.

BARIUM, PEROXIDE OF. Heat pure barytes to low redness in a platinum crucible; then gradually add chlorate of potash in the proportion of 1 part of the latter to 4 of the former. Cold water removes the chloride of potassium, and the peroxide remains as a hydrate.

BARYTES, CHLORATE. See CHLORATE OF BARYTES.

BEETLE POISON. Put a drachm of phosphorus in a flask with 2 oz. of water; plunge the flask into hot water, and when the phosphorus is melted, pour the contents into a mortar with 2 or 3 oz. of lard. Triturate briskly, adding water, and ½ ℔ of flour, with 1 or 2 oz. of brown sugar.—Pharmaceutical Journal. Plaster of Paris, with oat-meal, is said to destroy cockroaches.

BEETLE WAFERS. These are made with flour, sugar, and red lead, heated in wafer irons.

BENZOLE. A volatile liquid, procured by distilling light coal naphtha at a temperate not exceeding 200° F., by the

method patented by Mr. MANSFIELD. It is a solvent for gutta percha; and, also, with heat and long digestion, of India-rubber.

BEVERAGES, AND POWDERS FOR PREPARING THEM. See above page 268.

BIRD LIME. Boil the middle bark of the holly 7 or 8 hours in water; drain it, and lay it in heaps in the ground, covered with stones, for 2 or 3 weeks, till reduced to a mucilage. Beat this in a mortar, wash it in rain-water, and knead it till free from extraneous matters. Put it into earthen pots, and in 4 or 5 days it will be fit for use. An inferior kind is made by boiling linseed oil for some hours, until it becomes a viscid paste.

BISULPHURET OF CARBON. This is used in the arts, as a solvent for India-rubber, gutta percha, &c. To procure it, MULDER recommends the following process as the most convenient. Provide an iron bottle (a quicksilver bottle answers very well), and make a second opening into it. To one opening adapt a copper tube bent twice at right angles; and to the other a straight tube dipping into the bottle. Having nearly filled the bottle with pieces of charcoal (recently heated to redness), and having screwed on the bent and straight tubes, place the bottle in a furnace, closing the mouth of the latter with a stone or clay cover in two pieces, hollowed in the centre so as to fit the upper part of the bottle, and defend it from the action of the fire. Connect the curved tube with a WOOLFE'S bottle half-filled with water, and placed in a freezing mixture; and when the iron bottle is sufficiently heated, introduce by the straight tube fragments of sulphur, and immediately close the mouth of the tube with a plug. The bisulphuret, as it comes over, falls to the bottom of the water. Separate it from the water, and distil over dry muriate of lime.

BLACKING, LIQUID, FOR SHOES, &c. [*Note.*—By ivory-black, *bone*-black, which is usually sold under this name, is intended. True ivory-black has a more intense color, but is too dear for general use.] 1. Ivory-black 3 oz., treacle 2 oz., sweet oil ½ oz.; mix to form a paste; add gradually ½ oz. of oil of vitriol, and then half a pint of vinegar, and 1¾ pint of water or sour beer. Some prefer mixing the oil of vitriol with the sweet oil.

2. Ivory-black 2 lb, treacle 2 lb, sweet oil ½ lb; mix and add ¾ lb oil of vitriol, and beer or vinegar to make up a gallon.

3. Ivory-black 3 lb, treacle 4 lb, vinegar a pint, oil of vitriol 8 oz., water a gallon.

4. Ivory-black 2 lb, neat's-foot oil 4 oz.; mix, and add 3 quarts of sour beer, or vinegar, and a spoonful of any kind of spirits; stir till smooth, and add 2 oz. of oil of vitriol, and sprinkle on it ½ drachm of powdered resin. Then boil together 3 pints of sour ale with a little log-wood, and ¼ oz. of Prussian blue, 3 oz. of honey, and 8 oz. of treacle. Mix, but do not bottle it for 2 or 3 days.

5. Ivory-black 8 oz., brown sugar or treacle 8 oz., sweet oil 1 oz., oil of vitriol ½ oz., vinegar 2 quarts. Mix the oil with the treacle, then add the oil of vitriol and vinegar, and lastly, the ivory-black.

Blacking for Dress Boots. 1. Gum 8 oz., treacle 2 oz., ink a pint, vinegar 2 oz., spirit of wine 2 oz. Dissolve the gum and treacle in the ink and vinegar, strain, and add the spirit.

2. To the above add 1 oz. of sweet oil, and ¼ oz. lamp-black. [These are applied with a sponge, and allowed to dry out of the dust. They will not bear the wet.]

3. Beat together the whites of 2 eggs, a tablespoonful of spirit of wine, a lump of sugar, and a little finely powdered ivory-black to thicken.

Blacking without Polishing. Treacle 4 oz., lamp-black ½ oz., yeast a tablespoonful, 2 eggs, olive oil a teaspoonful, oil of turpentine a teaspoonful. Mix well. To be applied with a sponge, without brushing.

Blacking, India-rubber (Patent). Ivory-black 60 lb, treacle 45 lb, vinegar (No. 24) 20 gallons, powdered gum 1 lb, India-rubber oil 9 lb. (The latter is made by dissolving by heat 18 oz. of India-rubber in 9 lb of rape oil.) Grind the whole smooth in a paint mill, then add, by small quantities at a time, 12 lb of oil of vitriol, stirring it strongly for half an hour a day for a fortnight.

Blacking, Paste. 1. These may be made with the ingredients of liquid blacking, using sufficient vinegar, in which a little gum has been dissolved, to form a paste. Make it into cakes, and dry it.

2. *German Blacking.* Powdered bone-black is mixed

with half its weight of molasses and one-eighth of its weight of olive oil; and to this is added afterwards one-eighth of its weight of hydrochloric acid, and one-fourth of its weight of strong sulphuric acid. The whole is to be then mixed up with water into a sort of unctuous paste.—LIEBIG.

3. BAILEY'S *Blacking Balls*. Bruised gum tragacanth 1 oz., water 4 oz.; mix, and add 2 oz. of neat's-foot oil, 2 oz. of fine ivory-black, 2 oz. of Prussian blue, 4 oz. of sugar-candy; mix, and evaporate to a proper consistence.

For HEEL BALLS, see BALLS, above.

BLACKING FOR HARNESS. 1. Isinglass or gelatine ¼ oz., powdered indigo ¼ oz., soft soap 4 oz., logwood 4 oz., glue 5 oz. Boil together in 2 pints of vinegar till the glue is dissolved; then strain through a cloth, and bottle for use. This appears an unchemical composition; but is inserted (as are many similar ones) because it is in actual use. The next is of a different character.

2. Melt 8 oz. of beeswax in an earthen pipkin, and stir into it 2 oz. of ivory-black, 1 oz. of Prussian blue ground in oil, 1 oz. of oil of turpentine, and ¼ oz. of copal varnish. Make it into balls. To be applied with a brush and polished with an old handkerchief.

3. Treacle ½ ℔, lamp-black 1 oz., yeast a spoonful, sugar-candy, olive oil, gum tragacanth, isinglass, each 1 oz., a cow's gall. Mix all together with two pints of stale beer, and let it stand before the fire for an hour.

BLACK REVIVER. 1. Bruised nutgalls 1 ℔, logwood 1 ℔, water 5 quarts; boil to 4 quarts, and add sulphate of iron 4 oz.; dissolve and strain. When cold, add 8 oz. of ox gall.

2. Galls 3 oz., logwood 1 oz., copperas, iron filings, and sumach, of each 1 oz., vinegar 2 pints.—GRAY'S SUPPLEMENT.

BLACK JAPAN. True asphaltum 1½ oz., boiled linseed oil 4 pints, burnt umber 4 oz. Heat together till the whole is incorporated, remove from the fire, and when sufficiently cool, add as much oil of turpentine as will bring it to a proper consistence.

BLEACHING LIQUID. Solution of chloride of lime, and chloride of soda, are sold for this purpose, with directions for use. The following is also used: Mix 3 ℔ of common salt and

1 lb of black oxide of maganese with as much water as will form a paste. Put the mixture into a retort, and add 2 lb of oil of vitriol previously diluted with 4 lb of water. Pass the gas into a solution of 1 lb of common pearlash, or 11 oz. of caustic potash, in 6 lb of water. The retort may be placed, after a short period, in hot water to extricate the remaining gas. In bleaching cotton by chloride of lime, 1 lb is dissolved in 3 gallons of water for each pound of cloth; it is afterwards passed through diluted muriatic or sulphuric acid (1 part of acid to 30 of water), and then washed.

BLIGHTS, REMEDIES FOR.

Apple-tree Canker. Having brushed off the white down and the red stain underneath it, anoint the places with a mixture of train oil and Scotch snuff.

White Blight of Apple-tree. Apply a decoction of foxglove mixed with fresh cow-dung into a paste.

Fly in Turnips. Steep the seed before sowing in train oil; or distribute slaked lime over the field as soon as the plants have appeared: or contrive, if possible to fumigate the field with brimstone.

Mildew of Wheat. 1. To prevent it. Dissolve in 3½ gallons of cold water 3¼ oz., of sulphate of copper for every 3 bushels of sowing grain. Throw this quantity of wheat into another vessel, and pour over it the prepared liquid, until it rises 5 or 6 inches above the corn. Stir thoroughly, and remove all the grains that swim. Throw the mixture into a basket, so as to drain off the liquid; wash it well in soft water, and dry before sowing.

2. To remove it. A solution of 1 lb of salt to the gallon of water, sprinkled with a flat brush over the growing corn.

Smut in Wheat. To prevent it. Boil 3 gallons of water, and slake in it about 36 lb of quick lime; add 3 gallons more of cold water, and pour the hot mixture on 4½ bushels of the grain placed in a tub, stirring incessantly. Turn over the mixture now and then for 24 hours. Allow the liquid to drain off, and sow the limed wheat as soon as it is sufficiently dry. See WHEAT, STEEP FOR.

Blight in Vines. A solution of pentasulphuret of calcium may be painted over the branches and twigs until they acquire a continuous coating of sulphur mixed with carbonate

of lime. *M. Merceuril* raises the earth round the stems of the vine to about a foot deep, then places in each excavation a good handful of flowers of sulphur, heaping it round the stems. The earth is then replaced. This is done at the beginning of August.

For Lice, Aphides, and Red Spiders. See WASHES FOR VERMIN.

BLUE FOR LINEN. The ordinary kinds of cake blue consists of indigo and starch. Liefchild's patent blue is thus made: Mix 4 parts of Chinese blue, 1 of Turnbull's, and 1 of oxalic acid; gradually add boiling water until the whole is dissolved, and lastly 4 parts of sulphate of indigo. The latter is made with 1 part of indigo, and 4 of sulphuric acid, neutralized with carbonate of ammonia.

BOILERS, TO PREVENT INCRUSTATION IN. M. DUCLOS proposes to mix the water with which the boiler is to be supplied with the following solution, in the proportion of $1\frac{1}{4}$ per cent. Chloride of barium 125 parts, hydrochloric acid (s. g. 1.20) 25 parts, water 450. The water must then be allowed to clear in large reservoirs, where both the sulphate and carbonate of lime become decomposed, with formation of chloride of calcium. Any excess of acid must be neutralized before using, which may be done by placing pieces of limestone in the pipes.

BONES, SULPHATED. To a bushel of ground bones add from 10 to 14 lb of oil of vitriol, previously mixed with half its bulk of water. [It is sometimes mixed with an equal weight of salt and a sufficient quantity of bran. Turnip seed may be mixed with this compound, and sown together.]

BOOT-TOP LIQUID. 1. Solution of muriate of tin 3 dr.; French chalk, or Venetian talc, in powder, 1 oz.; salt of sorrel $\frac{1}{2}$ oz. flake white 1 oz., burnt alum $\frac{1}{2}$ oz., powdered cuttle-fish bone 1 oz., white arsenic 1 oz. boiling water a quart. Probably sulphate of barytes might be substituted for arsenic, the use of which it is desirable to discourage.

2. Sour milk 3 pints, cream of tartar 2 oz., oxalic acid 1 oz., alum 1 oz.—Mr. REDWOOD.

3. Wash the tops with soap and water, and scrape them with the back of a knife. Then apply the following with a harefoot brush.—Oxalic acid 1 oz., water a pint. Use the back of a knife as before; then polish with the following:

Powdered gum Arabic ½ oz., red spirits of lavender 2 oz., powdered turmeric ½ oz.; pencil this over the top, let it half dry, then polish by rubbing it, one way only, with a flannel till it shines.

4. Sour milk, 3 pints, butter of antimony 2 oz., cream of tartar 2 oz., citric acid, alum, burnt alum, of each 1 oz.—GRAY'S SUPPLEMENT.

5. *White Top.* One ounce of magnesia, alum, cream of tartar, and oxalic acid; ¼ oz. salt of sorrel, and ¼ oz. of sugar of lead; dissolve in a quart of water, and apply with a sponge.

6. *Brown Top.* Oxalic acid, alum, annotto, of each 1 oz.; isinglass ½ oz., sugar of lead ½ oz., salt of sorrel ¼ oz., boil together in a quart of water for 10 minutes. Apply with a sponge.

BOOKBINDERS' STAINS, FOR LEATHER. *Black.* A solution of 1 part of sulphate of iron in 6 of water. *Blue.* A solution of indigo. (See CHEMIC BLUE.) *Brown.* A solution of pearlash, or of common soda.

BOOT POWDER. Finely powdered French chalk, or Venitian talc.

BREAD, UNFERMENTED. See DIETETIC ARTICLES, p. 285.

BRONZE POWDER. The best methods of preparing these powders are probably kept secret. The following are some of the published recipes:

1. Gold leaf, or alloys of gold, reduced to powder by grinding them with sulphate of potash, or with honey, and washing away the extraneous matter with hot water, and drying the metallic powder.

2. Dutch metal, and other similar alloys, treated in the same way.

3. Verdigris 4 oz., tutty 2 oz., sublimate 1 dr., borax 1 dr., nitre 1 dr., mix them into a paste with oil, and fuse the mixture in a crucible. This has failed in some hands—perhaps from the tutty being factitious.

4. Mix together 100 parts of sulphate of copper, and 50 of crystallized carbonate of soda; apply heat till they unite. Powder the mass when cold, and add 15 parts of copper filings; mix well, and keep it at a white heat for 20 minutes. Wash and dry the product.

See also AURUM MUSIVUM, and ARGENTUM MUSIVUM, above.

BRONZING LIQUIDS, FOR BRONZING COPPER MEDALS, FIGURES, INSTRUMENTS, &c. 1. Sal ammoniac 1 dr., oxalic acid 15 gr., vinegar a pint: after well cleaning the article to be bronzed, warm it gently, and brush it over with the liquid, using only a small quantity at a time. When rubbed dry, repeat the application till the desired tint is obtained. (For copper medals, electrotype casts, &c.)

2. *Bronze for Plaster Figures.* Dissolve palm soap in water, and add a mixed solution of sulphate of copper and sulphate of iron until no further precipitate occurs. Dry the precipitate, and mix it with oil of turpentine, or linseed oil. Sulphate of copper alone produces too bright a green. Palmite of iron is yellow. These may be precipitated separately, and mixed to the desired shade after being titurated with the oil. Ten ounces of soap will require 3 ounces of sulphate of copper.

3. Sal ammoniac 1 oz., cream of tartar 3 oz., salt 6 oz.; dissolve in a pint of hot water, add 2 oz. of nitre, and 2 oz. of nitrate of copper dissolved in ½ pint of water.

4. Salt of sorrel 1 oz., sal ammoniac 2 oz., white vinegar 14 oz. (To give an antique appearance to bronze figures, &c.)

5. A diluted solution of muriate of platina. (For copper binding screws, and other small articles.)

6. A weak solution of hydrosulphuret of ammonia, or of sulphuret of potassium. (For electrotype medals.)

7. Immediately on removing the electrotype cast from the solution, brush it over with good black lead; then heat it moderately, and brush it over with a painting brush, the slightest moisture being used.

8. Boil 2 oz. of carbonate of ammonia and 1 oz. of acetate of copper in ½ pint of vinegar, until nearly all the vinegar is evaporated. Pour into this a solution of 62 grains of sal ammoniac, and 15½ grains of oxalic acid, in ½ pint of vinegar; boil the whole and filter. Apply it to the medal (which should be perfectly bright, and previously warmed) with a camel-hair pencil for half a minute; then pour boiling water on it; wipe it with soft cotton very slightly moistened with linseed oil, and rub it with clean cotton. *For electrotype copper medals.* (They may also be bronzed by applying oxide of iron (jewellers' rouge, or crocus) in the same manner as directed above, for plumbago; or a mixture of these may be used.)

9. *Tin Castings.* Wash them over, after being well

cleaned and wiped, with a solution of 1 part of sulphate of iron, and 1 of sulphate of copper, in 20 parts of water: afterwards with a solution of 4 parts of verdigris in 11 of distilled vinegar; leave for an hour to dry, then polish with a soft brush and colcothar.

BRONZING BALL. See BALL (HEEL).

BRONZING PASTES, PARISIAN. 1. Plumbago 1 oz., sienna 2 oz., rouge ½ oz. Add a few drops of hydrosulphate of ammonia, and water.

2. Chrome lead 2 oz., Prussian blue 2 oz., plumbago ½ lb, sienna powder and lake carmine, each ¼ lb. Add sufficient water to make a paste.

BROSSE DE CORAIL. The roots of lucerne (Medicago sativa), cleaned, dried, and hammered at the end. Used as a toothbrush.

BROWNING, OR BRONZING LIQUIDS, FOR GUN BARRELS. 1. Aquafortis ½ oz., sweet spirit of nitre ½ oz., spirit of wine 1 oz., sulphate of copper 2 oz., water 30 oz., tincture of muriate of iron 1 oz.: mix.

2. Sulphate of copper 1 oz., sweet spirit of nitre 1 oz., water a pint; mix. In a few days it will be fit for use.

3. Sweet spirit of nitre 3 oz., gum benzoin 1½ oz., tincture of muriate of iron ½ oz., sulphate of copper 2 dr., spirit of wine ½ oz., mix, and add 2 lb of soft water.

4. Tincture of muriate of iron ½ oz., spirit of nitric ether ½ oz., sulphate of copper 2 scruples, rain-water ½ pint. The above are applied with a sponge, after cleaning the barrel with lime and water. When dry, they are polished with a stiff brush or iron scratch brush.

BRUNSWICK BLACK. Melt asphaltum, and add to it half its weight of boiled linseed oil; mix, and when sufficiently cool, add enough oil of turpentine to bring it to the proper consistence.

BUG POISON. 1. Spirit of wine 8 oz., spirit of turpentine 8 oz., camphor ½ oz.: mix.

2. Distilled vinegar, or diluted wood vinegar, a pint; camphor ½ oz.; dissolve.

3. Corrosive sublimate 3 oz., muriatic acid 3 oz., oil of turpentine 12 oz., water 6 pints. Or, 1 oz. of sublimate, 2 oz. of muriatic acid, a pint of oil of turpentine, and a pint of decoction of tobacco.

4. Strong mercurial ointment 1 oz., soft soap 1 oz., oil of turpentine a pint.

5. White arsenic 2 oz., lard 13 oz., corrosive sublimate ¼ oz., Venetian red ¼ oz.

6. Scotch snuff mixed with soft soap.

7. *For Floors.* Corrosive sublimate 1 ℔, sal ammoniac 1 ℔, hot water 8 gallons. (It is said that if a branch of narrow-leaved dittany or pepperwort (Lepidium ruderale) be suspended in a room, all the bugs will settle in it, and may be taken. Fumigating the rooms with sulphur is a troublesome and disagreeable process, and not always successful.)

BURNETT'S (Sir W.) DISINFECTING FLUID. See DISINFECTING COMPOUNDS.

BUTTER, TO PRESERVE. Powder finely, and mix together, 2 parts of the best salt, 1 of loaf sugar, and 1 of nitre. To each pound of butter, well cleansed from the milk, add 1 oz. of this compound. It should not be used under a month. (Butter that has an unpleasant flavor, is said to be improved by the addition of 2½ dr. of bicarbonate of soda to 3 ℔ of butter. A turnipy flavor may be prevented by only feeding the cows with turnips immediately after milking them.)

BUTTER OF ANTIMONY. The liquid chloride of antimony, commercially known by this name, is usually made by dissolving crude or roasted black antimony in muriatic acid with the addition of a little nitric acid. It usually contains pernitrate of iron.

BUTYRIC ETHER. Saponify butter with a strong solution of potash, dissolve the soap in the smallest quantity of alcohol by the aid of heat, add a mixture of alcohol and sulphuric acid till the solution is acid, and distil as long as the product has a fruity odor. Redistil the product from chloride of calcium. It is sold as essence of pine-apple.

CAMPHENE. Highly rectified oil of turpentine. ENGLISH'S patent camphene is made by passing the vapor of oil of turpentine through caustic solutions of potash, soda, or lime; or through sulphuric acid.

CAMPHOR, ARTIFICIAL. This is formed by passing muriatic acid gas into oil of turpentine.

CAMPHOR BALLS. See SKIN COSMETICS, after PERFUMERY.

CANDIES. These belong rather to the confectioner than the druggist. The green stalks of angelica, the peels of orange, lemon, and citron, green roots of ginger, &c., are first boiled in water till soft, then in syrup till they are transparent, and dried in a stove, at a heat not exceeding 104° F. Candied horehound is made by boiling lump sugar with a little strong decoction or infusion of dried horehound till a portion taken out and cooled becomes solid. It is then poured on to a slab, or into paper or tin moulds dusted with powdered sugar.

CANTON'S PHOSPHORUS. Put calcined oyster shells in layers, alternately with sulphur, and heat strongly in a covered crucible for an hour.

CAOUTCHOUC, SOLVENTS FOR. See SOLVENTS.

CAPSULES, GELATINOUS. These are used to contain copaiva and other nauseous liquids which do not dissolve gelatine, so that they may be swallowed without inconvenience. They are made by "dipping the bulbous extremity of an iron rod into a concentrated solution of gelatine. When the rod is withdrawn, it is to be rotated, in order to diffuse the gelatine equally over the bulb." When sufficiently hardened, they are removed, placed on pins to dry, and when dry filled with the balsam or oil, and the orifice closed with liquid gelatine. They are usually of an olive form, and contain 10 gr. of balsam in each. See Dr. PEREIRA'S "Elements," article COPAIVA. M. GIRAUD recommends the following composition for capsules: Transparent gelatine 12 parts, syrup of gum 2 parts, syrup 2 parts, water 10 parts. Melt it in a water-bath, remove the scum, and dip the mould, previously oiled, into the compound.

CARBON. See CHARCOAL.

CARBONIC ACID. See GASES.

CARMINE. See PIGMENTS.

CASE-HARDENING POWDER. This is merely ferroprussiate of potash, dried, and finely powdered. By sprinkling it on iron heated to bright redness, the metal becomes case-hardened, or superficially converted into steel. The iron should be plunged into cold water as soon as the powder has acted on it. The following compound is used for the

same purpose: Sal ammoniac 2 oz., burnt bone-dust 2 oz., HENWOOD'S composition ½ oz. Used as the former.

CASSOLETTES. See PERFUMERY.

CAYENNE, SOLUBLE. See CONDIMENTS, p. 292.

CEMENTS AND LUTES, VARIOUS.

Shellac Cement. Fine orange shellac, bruised, 4 oz., highly rectified spirit 3 oz. Digest in a warm place, frequently shaking, till the shellac is dissolved. Rectified wood naphtha may be substituted for spirit of wine, where the smell is not objectionable. A most useful cement for securely joining almost any material. See LIQUID GLUE.

Shellac Cement without Spirit. Boil 1 oz. of borax in 16 oz. water, add 2 oz. powdered shellac, and boil in a covered vessel till the lac is dissolved. Cheaper than the above, and for many purposes answers very well. Both are useful in fixing paper labels to tin, and to glass when exposed to damp.

Armenian Cement for Glass, China, &c. 1. KELLER'S. Soak 2 dr. of cut isinglass in 2 oz. of water for 24 hours; boil to 1 oz., add 1 oz. spirit of wine and strain through linen. Mix this, while hot, with a solution of 1 dr. of mastic in 1 oz. of rectified spirit, and triturate with ½ dr. powdered gum ammoniac, till perfectly homogeneous.

2. Dr. URE'S *Diamond Cement.* Isinglass 1 oz. distilled water 6 oz., boil to 3 oz., and add 1½ oz. of rectified spirit. Boil for a minute or two, strain, and add while hot, first ½ oz. of a milky emulsion of ammoniac, and then 5 dr., of tincture of mastic. (There are various kinds of this cement sold, and some of the improvements introduced by peculiar makers have not been made public.)

Cement used in the East for uniting jewels, glass, and metals. Dissolve 5 or 6 pieces of gum mastic, each about the size of a large pea, in just as much spirit as will render it liquid. Soften some isinglass by steeping it in water; having dried it, dissolve as much of it in good brandy as will make a two-ounce vial of strong glue, to which must be added two small bits of gum ammoniacum, rubbing until they are dissolved. Mix the two solutions; keep in a close vial; and when it is to be used, set the vial in boiling-water.—Mr. ETON.

HŒNLE'S *Cement for Glass or Earthenware.* Shellac 2 parts, Venice turpentine 1 part, fuse together, and form into sticks.

Cheese Cement for Earthenware, &c. Mix together white of egg beaten to a froth, quicklime, and grated cheese, and beat them to a paste. See GLUE, CASEINE.

Curd Cement. Add ½ pint of vinegar to ½ pint of skimmed milk; mix the curd with the whites of 5 eggs well beaten, and sufficient powdered quicklime to form a paste. It resists water and a moderate degree of heat.

Glass Flux, for mending broken China, &c. Mix 3 parts of red lead, 2 of fine white sand, and 3 of crystallized boracic acid, fuse the mixture, levigate it, and apply it with thin mucilage of tragacanth. Heat the repaired article gently, so as partially to fuse the cement.

Cement for joining Spar and Marble Ornaments, &c. Melt together 8 parts of resin, 1 of wax, and stir in 4 parts, or as much as may be required, of Paris plaster. The pieces to be made hot.

HENSLER'S *Cement.* Grind 3 parts of litharge, 2 of recently burnt lime, and 1 of white bole, with linseed oil varnish. (Very tenacious, but long in drying.)

SINGER'S *Cement, for Electrical Machines and Galvanic Troughs.* Melt together 5 ℔ of resin, and 1 ℔ of beeswax, and stir in 1 ℔ of red ochre (highly dried, and still warm), and 4 oz. Paris plaster, continuing the heat a little above 212°, and stirring constantly till all frothing ceases. Or (for troughs), resin 6 ℔, dried red ochre 1 ℔, calcined plaster of Paris ½ ℔, linseed oil ¼ ℔.

Botany Bay Cement. Botany Bay gum, melted and mixed with an equal quantity of brickdust.

Cap Cement. As SINGER'S; but 1 ℔ of dried Venetian red may be substituted for the red ochre and Paris plaster.

Bottle Cement. Resin 15 parts, tallow 4 (or wax 3) parts, highly dried red ochre 6 parts, or ivory-black q. s. The common kinds of sealing-wax are also used.

Turner's Cement. Beeswax 1 oz., resin ½ oz., pitch ½ oz. Melt, and stir in fine brickdust q. s.

Coppersmith's Cement. Powdered quicklime, mixed with bullock's blood, and applied immediately.

Engineer's Cement. Equal weights of red and white lead, with drying oil, spread on tow, or canvas. For metallic joints, or to unite large stones, in cisterns, &c.

Cement for Steam Pipes. Good linseed oil varnish ground with equal weights of white lead, oxide of manganese, and pipe-clay.

Iron Cement, for closing the joints of iron pipes. Iron borings, coarsely powdered, 5 lb, powdered sal ammoniac 2 oz., sulphur 1 oz., water sufficient to moisten it. It quickly hardens; but if time can be allowed, it sets more firmly without the sulphur. It must be used as soon as mixed, and rammed tightly into the joints.

GAD'S *Hydraulic Cement.* Powdered clay 3 lb, oxide of iron 1 lb, boiled oil to form a stiff paste.

Cements for masonry of Chambers of Chlorine, &c. Equal parts of pitch, resin, and plaster of Paris.

Roman Cement. A mixture of clay, lime, and oxide of iron, separately calcined, and finely powdered. It must be kept in close vessels, and mixed with water when used.

Oxychloride of Zinc Cement. In liquid chloride of zinc marking from 50° to 60° of Baumé's areometer, dissolve 3 per cent. of borax or sal ammoniac; add oxide of zinc which has been heated to redness, until the mass is of a proper consistence. The cement, when hard, becomes as firm as marble. It may be cast in moulds, like plaster of Paris, or used in the construction of mosaic works, &c.—M. SOREL.

Marine Cement. See GLUE, MARINE.

MAISSIATT'S *Cement, as an air-tight covering for bottles, &c.* Melt India-rubber (to which 15 per cent. of wax or tallow may be added), and gradually add finely-powdered quicklime till a change of odor shows that a combination has taken place and a proper consistence is obtained.

Cement for attaching Metal Letters to Plate Glass. Copal varnish 15 parts, drying oil 5 parts, turpentine 3 parts, oil of turpentine 2 parts, liquefied glue 5 parts; melted in a water-bath, and 10 parts of slaked lime added.

Cement to fasten Leather to Metal. A hot solution of gelatine. With this wash the metal; and the leather, previously

steeped in a hot infusion of gallnuts, is to be pressed tightly on the surface and allowed to cool, when it will adhere very firmly.

Japanese Cement. Mix rice flour intimately with cold water, and boil gently.

French Cement. Mix thick mucilage of gum arabic with powdered starch.

Common Paste. To a dessert-spoonful of flour add gradually half a pint of cold water, and mix till quite smooth; add a pinch of powdered alum (some add also as much powdered resin as will lie on a sixpence), and boil for a few moments, stirring constantly. The addition of a little brown sugar, and a few grains of corrosive sublimate, is said to preserve it for years.

Transparent Cement. Dissolve 75 parts of caoutchouc in 6 parts of chloroform, and add to the solution 15 parts of mastic.—LENHER.

Soft Cement. Melt yellow wax with half its weight of common turpentine, and stir in a little Venetian red, previously well dried, and finely powdered. (As a temporary stopping for joints and openings in glass and other apparatus, where the heat and pressure are not great.) See the next.

Lutes or Cements for closing the joints of Apparatus. 1. Mix Paris plaster with water to a soft paste, and apply it immediately. It bears nearly a red heat. It may be rendered impervious by rubbing it over with wax and oil.

2. Slaked lime, made into a paste with white of egg, or a solution of gelatine.

3. *Fat Lute.* Finely-powdered clay moistened with water, and beaten up with boiled linseed oil. Roll it into cylinders, and press it on the joints of the vessels, which must be perfectly dry. It is rendered more secure by binding it with strips of linen moistened with white of egg.

4. Linseed meal beaten to a paste with water.

5. Slips of moistened bladder, smeared with white of egg.

Luting for Acids. Dissolve 1 part of India-rubber in 2 parts of linseed oil, by heat, and work into a stiff paste with 3 parts, or as much as sufficient, of white clay.

Lutes for Coating Retorts. 1. Dissolve 1 oz. of borax in ½ pint of water, and add slaked lime to form a thin paste. Brush this over the retort, and let it dry gradually. Then apply a coating of slaked lime and linseed oil beaten together. Let it dry a day or two before use, and fill up any cracks which may appear with lime and linseed oil.

2. For bearing a stronger heat: Stourbridge clay, mixed with a little sand to prevent it splitting off: a little cut tow, or horse-dung, or asbestos, is usually added, to increase its coherence. It should be beaten to a stiff paste, and rolled out before application. The glass should be first rubbed over with a little of the lute mixed with water, then placed in the centre of the paste, rolled to about ¼th or ⅓d of an inch in thickness, and the edges of the latter raised and moulded to the glass, taking care to press out all the air.

Mohr's *Lute.* Mix equal parts of brickdust and litharge, and beat them into a paste with linseed oil. Apply this with a stiff brush, and dust it over with coarse sand; dry it in a warm place.

For Cements for plugging teeth, see Teeth Cosmetics, page 265.

Charcoal. Wood charcoal is made by burning wood with only a partial access of air. For chemical purposes, that made in iron cylinders, in the manufacture of wood vinegar, is preferred.

Charcoal, Animal. The most common form is that of bone-black (commonly called ivory-black), made by distilling bones, from which the grease has been removed by boiling, in iron or earthen retorts, the ammoniacal liquor, &c., being collected in proper receivers. The residue is bone-black. When used for the purpose of decolorization, further treatment is required: either the bone-earth may be entirely removed by muriatic acid, as directed in the Pharmacopœia; or more commonly, in the refining of sugar, and for other manufacturing purposes, this is only partially effected, in some such way as the following: Mix 8 lb of the bone-black, coarsely powdered, with sufficient water to form a paste, and add 1 lb of muriatic acid. In an hour pour boiling water on the mass, let it settle, pour off the liquid, add more water, and repeat this till the water comes off free

from taste. Drain and press the black in a cloth, and dry it. Its power is increased by mixing it with a little potash, heating it to dull redness in a colored crucible, and again washing it.

CHARCOAL, *Aluminized.* This is recommended by Dr. STENHOUSE as a cheap and very efficient decolorizing agent. Dissolve in water 54 parts of the sulphate of alumina of commerce, and mix with 92½ parts of finely powdered wood charcoal. When the charcoal is saturated, evaporate to dryness, and heat to redness in covered Hessian crucibles till the water and acid are dissipated. The charcoal contains just 7½ per cent. of anhydrous alumina.

CHARCOAL FROM COAL-TAR. Heat gently in an iron pot till it melts 1 lb by weight of coal-tar pitch. Add 2 lb of fluid coal-tar, and mix. Stir in 7 lb of hydrate of lime in very fine powder. The thick mass is now roasted, stirring all the time, till it is reduced to a fine powder. It is then ignited in a covered crucible till all the vegetable matter is carbonized. The charcoal when cold is digested with dilute hydrochloric acid, and finally washed with water in a filter, and dried. Dr. STENHOUSE recommends this as an admirable form for decolorization. For such liquids as decoction of logwood it is four times as efficient as animal charcoal.

CHAMELEON MINERAL. Mix equal weights of black oxide of manganese and pure potash, and heat them in a crucible. Keep the compound in closely-stopped bottles. A solution of it in water passes through various shades of color from green to red.

CHEMIQUE, OR CHEMIC BLUE.—SULPHATE OF INDIGO. To 7 or 8 parts of oil of vitriol, in a glass or earthen vessel, placed in cold water, add gradually 1 part of fine indigo in powder, stirring the mixture at each addition with a glass rod, or piece of tobacco-pipe. Cover the vessel for 24 hours, then dilute with an equal weight of water. Sometimes it is sold without diluting. The German fuming acid answers best, 4 or 5 parts of it being sufficient for 1 of indigo. For dyeing silk, &c., carbonate of potash, soda, or ammonia, is added, to neutralize the acid, taking care not to add it in excess.

CHLORATE OF BARYTES. Saturate solution of chloric acid, prepared as directed in page 303, with fresh precipitated carbonate of barytes, filter, and crystallize.

Chlorate of Potash. Slack 7 oz. of quicklime, add it to a pint of solution of caustic potash of 1.110 sp. gr.; heat the mixture slightly, and pass through it a rapid current of chlorine gas until no more is absorbed. Evaporate merely to dryness, redissolve in boiling water, filter, and evaporate the filtering solution, and the washings of the filter, for crystallization. — Mr. Calvert. Or dissolve chloride of lime in water, add solution of muriate of potash, and boil to dryness. Dissolve the mass in hot water, and filter if necessary: on cooling, a large quantity of chlorate of potash is deposited.—Liebig. For another process, see Potassæ Chloras, P. F.

Chloride of Lime, or Chlorinated Lime. *Bleaching Powder.* Chlorine gas (slowly evolved from a mixture of 10 parts of common salt and 10 to 14 parts of oxide of manganese, placed in an alembic of lead, and heated by steam, and with 12 to 14 parts of oil of vitriol previously diluted with a fourth of its weight of water, added) is conveyed into a chamber where sifted slake lime is thinly spread on shelves. It is so cheaply made by the large manufacturers for bleaching purposes, that it is seldom prepared by druggists. The liquid chloride of lime may be made either by triturating the dry chloride with a little cold water till perfectly smooth, then adding more water, and filtering the solution; or by passing chlorine gas into a mixture of lime and water. The Dub. Pharm. directs a solution of 1.035 sp. gr. The Paris Codex directs 1 part of dry bleaching powder to be diffused in 45 of water. Soubeiran directs 1 part to 50; but prefers passing the gas from 1 part of oxide of manganese and 4 of muriatic acid, into a mixture of 1 part of lime and 50 of water.

Chloride or Hypochlorite of Potash. *Eau de Javelle.* Pass chlorine gas as above into a solution of 1 ℔ of carbonate of potash in a gallon of water. Or mix 8 oz. of dry chloride of lime with six pints of water, and dissolve 16 oz. of carbonate of potash in a quart of water. Mix the liquids and filter.

Chloride of Soda, or Chlorinated Soda. See Soda Chlorinata, and Liquor Sodæ Chlorinatæ, in the Pocket Formulary. A more ready way of preparing it, for other purposes than dispensing prescriptions, is the following: Diffuse 1 ℔ of chloride of lime in 30 ℔ of water. Dis-

solve 2 lb of crystallized carbonate of soda in 15 lb of water. Mix the solutions, let the mixture settle, pour off the clear liquid, and filter it.

CHLORINE. See GASES.

CHLOROFORM. Into a copper still placed in a water-bath put 7 gallons of water; heat to 100° F.; add 10 lb of lime slacked in a solution of 20 lb of chloride of lime. Add 3 lb of rectified spirit; mix them well, lute on the head immediately, and raise the heat rapidly to 212°. When the extremity of the neck of the still becomes hot, damp the fire, and the distillation will continue regularly. When it ceases, add to what remains in the still 17½ pints of water, raise the heat to 100°, add 8 lb of lime, and 20 lb of chloride of lime, stir carefully, add the distilled liquor from which the chloroform has been separated, with 1¾ pint of spirit; stir, and proceed as before. If the still is large enough, this may be repeated 3 or 4 times, with a constantly increasing product. The chloroform sinks to the bottom of the distilled liquid, and may be rectified by distilling it with oil of vitriol. Its density should not be less than 1.496: LIEBIG says 1.480, but at that density it is not pure. Chloroform may be obtained by a similar process from rectified wood naphtha (pyroxylic spirit); but it is then contaminated with an empyreumatic oil, lighter than water, which cannot be entirely separated from it. It is therefore unfit for inhalation; but when rectified over sulphuric acid* and chloride of calcium, may be used as a solvent. The process recommended by Dr. SIMPSON is that of Dumas: Chloride of lime in powder 4 lb, water 12 lb, rectified spirit 12 oz. Mix in a capacious still, and distil as long as a dense liquid, which sinks to the bottom of the water which comes over with it, is produced. For other methods see Pocket Formulary.

CHOCOLATE. See Chocolate, Pocket Formulary.

CHROMATE OF POTASH. Mix 4 parts of chrome ore (chromate of iron) with 2 of pearlash and 1 of nitre, and heat the mixture in a reverberatory furnace for several hours. Lixiviate, and crystallize. The chromate is converted into bichromate

* This process of rectification by oil of vitriol does not seem to succeed. See ROYLE and HEADLAND'S "Materia Medica," art. Chloroformum.

by adding sulphuric acid, or rather acetic acid, to the solution.

CHROMATE OF LEAD. See PIGMENTS.

CHROME OXIDE. Mix bichromate of potash with half its weight of muriate of ammonia; heat the mixture to redness, and wash the mass with plenty of boiling water. Dry the residue.

CHROMIC ACID. See ACID, CHROMIC.

GLOTHES, POWDER TO KEEP AWAY MOTHS FROM. Mix powdered pipe-clay, 1½ lb, white pepper and starch each 1 oz., root of Florentine iris 1½ oz., with spirits of wine 2 oz. To be dusted over the clothes when laid by.

COCHINEAL COLORING. Macerate best carmine 1 oz. in strong solution of ammonia 6 oz. until it is dissolved. Heat gently to drive off excess of ammonia. Put into a quart wine bottle, and add rectified spirit 4 oz., and white sugar 3 lb. Fill up with warm water, and shake until the sugar is dissolved.—Mr. PALMER.

COLLODION. Mix in a glass, stoneware, or porcelain vessel, 30 parts of strong sulphuric acid and 20 of powdered nitre; place the vessel in cold water; add 1 part of carded cotton-wool, and open and stir it in the acid mixture by means of 2 glass or porcelain rods, or stems of tobacco pipes, for 2 or 3 minutes. Then remove the cotton into a large quantity of cold water, press it, and wash it in a stream of water, opening it with the fingers, till the water passes through it free from acidity. Squeeze it strongly in a dry cloth, and then open it, and dry it gradually in a warm situation, free from danger. One part of this prepared cotton with 16 of rectified ether, and 1 of alcohol, agitated together, soon forms a gelatinous solution. See Pocket Formulary for its medical preparations.

COLLODION, ELASTIC (for surgical purposes). Mix together, in a stoneware or porcelain pot, sulphuric acid (of sp. gr. 1.847) 300 parts, very dry nitrate of potash 200 parts; and add carded cotton 10 parts. Leave in contact for 12 minutes; withdraw the cotton, wash it in cold water to remove the acid which it contains, and after 2 or 3 rinsings, immerse it again in a solution of 30 parts of subcarbonate of potash in

1000 of water; plunge it again into simple water, agitating well; and lastly, dry it at a temperature of 77° to 86° Fahr. The product is xyloidine. Place now 8 parts of this xyloidine, with 125 parts of sulphuric ether, in a wide-mouthed flask, and add 8 parts of alcohol (sp. gr. 0.825). Agitate. Make next a mixture of Venice turpentine 2 parts, castor oil 2 parts, white wax 2 parts, sulphuric ether 6 parts. Heat together the first three substances, add the ether, and mix all with the solution of xyloidine.—M. LAURAS.

COLLODION, PHOTOGRAPHIC. See PHOTOGRAPHY.

COLORS, VARIOUS. The principal dry colors will be found under PIGMENTS. Other colors are noticed below.

COLORS FOR DRUGGIST'S SHOW BOTTLES. In making these, distilled water should be used, and rather more of the color than will fill the carboys made, to avoid the necessity of adding water to fill up after filtration, as this sometimes renders them turbid. The carboys should be perfectly clean, and also dry, or otherwise rinsed out with a portion of the filtered liquid.

1. *Blues.* Sulphate of copper 4 to 8 oz., water a gallon, oil of vitriol 1 oz.

2. *Royal Blue.* Sulphate of copper 8 oz., water a gallon; dissolve, and add water of ammonia till the full color is developed; then water to make up 2 gallons.

3. *Finest Royal Blue.* Nitrate of copper 3 oz., water sufficient to dissolve it; add water of ammonia as long as the color becomes deeper, then water to make up 2 gallons.

4. *Paler.* Crystallized acetate of copper ½ oz., muriatic acid 1½ oz., water of ammonia q. s., water to make up 2 gallons.

5. *Light Blue.* Crystallized acetate of copper a scruple, water of ammonia 2 oz., water 2 gallons.

Green. 1. An infusion of saffron added to the above blues.

2. Sulphate of copper 4 oz., bichromate of potash ½ dr., water q. s.

3. By adding to the above deep or light blues a small quantity of chromate or bichromate of potash till the desired tint is produced, various shades of green may be obtained.

4. Sulphate of copper, with muriate of soda, or muriate of iron.

5. *Emerald Green.* Nitrate of copper 3 oz., muriatic acid 4 oz., nitric acid 4 oz., water 2 gallons.

6. *Very Fine Emerald.* Nickel 3 oz., muriatic acid 4 oz., nitrous acid 2 oz.; digest for 24 hours, add 2 gallons of water, and filter.

Red. 1. Dried rose petals 8 oz., boiling water a gallon; digest for 12 hours, strain; digest the roses with more water, and strain; mix the infusions, add a pint of diluted sulphuric acid, and filter.

2. Cochineal ½ oz., boiling water a gallon; digest, strain, add ½ oz. of sulphuric acid, and water to make up 2 gallons.

3. *Crimson.* Iodine, and iodide of potassium, of each 2 dr.; triturate with a dr. of water, and add 3 gallons of water, and 4 oz. of muriatic acid.

4. *For External Lamps.* Camphene, strongly colored with alkanet root.

Pink. 1. Infuse ½ oz. of good madder in a quart of boiling distilled water; when cold, add 1 oz. of strong ammonia, and filter into 2 gallons of distilled water.

2. Dissolve muriate or nitrate of cobalt in water, and add sufficient carbonate of ammonia to redissolve the precipitate first thrown down; then water q. s.

Purple. To the last add sufficient of the blue No. 3, to give the desired shade.

Lilac. Smalts 4 oz., nitric acid 4 oz.; let it stand 24 hours, add 2 gallons of water, 1 oz. of alum, and 4 oz. of water of ammonia. Or rather as the purple.

Yellow. 1. Chromate or bichromate of potash, with water q. s.

2. Bichromate of potash 2 dr., hot water 4 oz.; dissolve, and add 4 oz. sulphuric acid, and 2 gallons of water.

3. Bichromate of potash, nitrate of potash, and water.

Amber. 1. Deep chrome yellow 3 oz., pearlash 9 oz., water 2 gallons; boil gently for half an hour, take it off to cool, and add 6 oz. muriatic acid, and water to the desired color.

2. Dragons' blood, digested with sulphuric acid, and diluted with water to the desired shade.

Olive. Sulphate of iron 3 oz., sulphuric acid 3 oz., water 2 gallons; dissolve, and add the green No. 5, q. s. to brighten the color.

LIQUID COLORS FOR MAPS, &c. See INKS, DYES, CHEMIQUE

BLUE, LAKE LIQUOR, &c. Gamboge and some of the cake colors, mixed with water, are also used.

COLORS FOR CONFECTIONERS. Many fatal accidents occur from confectionery being colored with poisonous pigments. The following may be safely used: Cochineal and its preparations, sap green, vegetable lakes, Prussian blue; a mixture of a yellow lake and Prussian blue for green.

COLORING FOR BRANDY, &c. Sugar melted in a ladle till it is brown, and then dissolved in water or lime-water.

COLORS FOR LIQUEURS. Pink is given by cochineal; yellow by saffron or safflower; violet by litmus; blue by sulphate of indigo, saturated with chalk; green by the last, with tincture of saffron, or by sap green.

COLORS FOR LEATHER. See BOOKBINDERS' COLORS.

COLORS, IMPROVED VEHICLES FOR. 1. One measure of saturated solution of borax, with 4 of linseed oil. The pigment may be ground with the oil or the mixture.

2. A solution of shellac with borax, as in making COATHUPE'S Ink. See INK.

3. Water colors, mixed with gelatine, and afterwards fixed by washing with a solution of alum.

4. Curd of milk, washed and pressed, then dried on fine net, and when required for use, mixed with water and the coloring matter.

COPPER, OXIDE OF. The *black* oxide is made by calcining the nitrate; or by adding caustic potash to sulphate of copper, in solution, and washing and drying the precipitate. The red oxide may be made as directed for Bronze Powder, No. 4, or in the moist way thus: Pour a solution of 27 parts of sugar in 60 of water, over 9 parts of hydrated oxide of copper, weighed in its compressed but still moist state. A solution of 18 parts of caustic potash in 60 of water is added, and the whole agitated together without heat, and filtered. The clear liquid heated in a water-bath, and continually stirred, deposits the red oxide, and the liquid becomes colorless.

NITRATE OF COPPER. Dissolve copper in nitric acid to saturation, evaporate to dryness, redissolve, filter, and evaporate, so that the salt may crystallize. Or add a solution of sulphate of copper to a solution of nitrate of lead, so long as sulphate of lead is precipitated; filter, evaporate, and

crystallize. For the other salts of copper, see CUPRUM, Pocket Formulary.

COTTON POWDER. See GUN COTTON.

CRAYONS FOR WRITING ON GLASS. Fuse in a cup 4 parts of spermaceti, 3 of tallow, and 2 of wax; stir in 6 parts of minium, and 1 of potash; keep warm for ½ an hour, and then pour into glass tubes of the thickness of a lead-pencil. If cooled rapidly, the mass may be screwed up and down in the tube, and cut at the end to a fine point. The glass to be written on must be clean and dry.

DEPILATORIES. See HAIR COSMETICS, page 254.

DEXTRINE, OR STARCH GUM. Heat 4 gallons of water in a water-bath to between 77° and 86° Fahrenheit; stir in 1½ or 2 lb finely ground malt; raise the temperature to 140°, add 10 lb of potato or other starch; mix all thoroughly, raise the heat to 158°, and keep it between that and 167° for 20 or 30 minutes. When the liquor becomes thin, instantly raise the heat to the boiling-point, to prevent the formation of sugar. Strain the liquor, and evaporate it to dryness, as the dextrine will not keep long in a liquid form. Another method is to boil solution of starch with a few drops of sulphuric acid, filter the solution, and add alcohol to throw down the dextrine. See GUM (BRITISH) for another form of dextrine.

DEXTRINE VARNISH. Dextrine 2 parts, water 6 parts, rectified spirit 1 part.—BARON DE SYLVESTRE.

DIASTASE. Macerate ground malt in cold water; strain with pressure; heat the clear solution in a water-bath to 158° Fahrenheit to coagulate the albumen; filter again, and add rectified spirit as long as diastase falls. If required very pure, redissolve it in water, and again precipitate with spirit. Dry it at a low temperature. Well-malted barley contains about 1 per cent. of pure diastase; one part of which is capable of converting 2000 parts of starch into dextrine or sugar.

DIETETIC ARTICLES. For these see another division of the work.

DISINFECTING AND DEODORIZING COMPOUNDS. 1. SIR WM. BURNETT'S *Patent Solution.* It is made by dissolving zinc in commercial muriatic acid to saturation.

2. ELLERMAN'S *Deodorizing Fluid.* This consists chiefly

of perchlorides and chlorides of iron and manganese. In a report addressed to the Metropolitan Board of Works in 1859, Drs. HOFFMAN and FRANKLAND stated that the perchloride of iron was the cheapest and most efficient deodorizer that could be applied to sewage; ½ gallon deodorized 7500 gallons of this. 1 bushel of lime, or 3 lb of chloride of lime, would do the same.

3. LEDOYEN'S *Solution.* This is a solution of nitrate of lead, and contains about 20 oz. of the salt in a gallon. The specific gravity should be 1.40. A similar compound may be made by mixing 13½ oz. of litharge with 6 pints of water, and adding 12 oz. of nitric acid at 1.38 specific gravity (or 8 oz. at 1.50); and digesting at a gentle heat till the solution is complete.

4. SIRET'S *Compound.* Sulphate of iron 20 lb, sulphate of zinc 2½ lb, wood or peat charcoal 1 lb, sulphate of lime 26½ lb; mix, and form into balls. To be placed in cesspools, &c., to deodorize them. M. SIRET has subsequently modified this compound thus: Sulphate of iron 100 parts, sulphate of zinc 50, tan or oak-bark powder 40, tar 5, and oil 5 parts.

5. COLLINS'S *Disinfecting Powder.* Mix 2 parts of dry chloride of lime with 1 of burnt alum. To be set in shallow dishes in rooms, &c., with or without the addition of water. (See also CHLORIDE OF LIME, and CHLORIDE OF SODA. Peat charcoal also possesses powerful deodorizing properties.)

DUBBING, CURRIERS'. Made by boiling cuttings of sheepskins in common cod-oil.

DRIERS FOR PAINTERS. White copperas 1 lb, sugar of lead 1 lb, white 2 lb; ground with boiled oil.

DRYING AND BOILED OIL. Linseed oil is mixed with powdered litharge, and heated till it becomes thick. A *pale* drying oil is obtained by mixing with linseed oil sufficient dry sulphate of lead to form a milky liquid, and shaking it repeatedly for some days, letting it stand exposed to the light. When it has become quite clear, it may be poured off from the dregs. The sulphate of lead when washed from the mucilage, may be again used for the same purpose. LIEBIG directs 1 lb of acetate of lead to be dissolved in half a gallon of rain-water, and 1 lb of finely-powdered litharge added: the mixture is either boiled, or exposed for a longer time to

a moderate heat, and frequently stirred, till no more particles of litharge can be seen. A white deposit is formed, which may be left in the liquid or separated by filtration; 20 ℔ of linseed oil, in which 1 ℔ of levigated litharge has been diffused, are gradually added to the lead solution, previously diluted with an equal bulk of water, and the mixture frequently stirred. It is then left to clear itself in a warm place; but to obtain it bright it must be filtered through coarse paper or cotton. It may be bleached by exposure to the sun. The lead solution which subsides from the mixture may be filtered and used again, after dissolving in it 1 ℔ of litharge as before. The oxide of lead contained in the oil may be removed from it by agitating it with diluted sulphuric acid, and letting it stand to settle. See also OILS (Clarified Linseed Oil).

DUPUYTREN'S POMADE. See HAIR COSMETICS, after PERFUMERY.

DYES AND COMPOUNDS USED IN DYEING. A few of the principal coloring matters and mordants may here be noticed; for further information, the reader is referred to Dr. URE'S "Dictionary of the Arts," PARNELL'S "Applied Chemistry," BERTHOLLET "On Dyeing," &c.

Blue Dyes. The most important of these is indigo. Being insoluble in water, it is prepared for use by sulphate of iron and alkalies of lime, by fermentation and alkalies, and by solution in sulphuric acid.

1. Triturate 1 ℔ of indigo with water and a little caustic potash; then add 3 ℔ of lime, and afterwards 2½ ℔ of sulphurate of iron in solution, stirring them well together. The solution contains *disoxygenized indigo*, which is soluble in lime and alkalies. The cotton, linen, &c., to be dyed is repeatedly dipped in the solution, and afterwards rinsed in water soured with muriatic acid.

2. To 45 or 50 gallons of water, heated to 122° F., add 12 oz. of indigo, 8 oz. of madder, 8 oz. of bran, and 24 oz. of potash. In 36 hours introduce 12 oz. more of potash, and the same in 12 hours after. In 72 hours add a little lime to check the fermentation. Wool, silk, linen, and cotton may be dyed in this bath. Another form of this dye is—Indigo 2 parts, common soda 5 parts, lime 2 parts, clarified honey 1 part, water as much as may be sufficient. Keep it warm in an earthen jar till the indigo is dissolved.

3. A solution of indigo in sulphuric acid (see CHEMIC BLUE) is used as a dye, but a purer tone of color is obtained by the following method: The sulphate of indigo mixed with water, is heated in a copper kettle; wool is immersed in it, and the whole is allowed to cool for 24 hours. The wool is then taken out, washed till the water comes off colorless and free from acid; it is then boiled in water containing about 2 per cent. of pearlash, or other alkaline carbonate, for a quarter of an hour. The quantity of pearlash should be equal to one-third the weight of the indigo.

Logwood, with verdigris, or sulphate of copper, gives a blue dye, bordering on violet: with alum and tartar a violet.

Prussian Blue is sometimes used in dyeing, after being triturated and digested for 24 hours with its weight of muriatic acid. A blue is also given by immersing silk, &c., in a solution of peracetate of iron, then in a solution of prussiate of potash, and afterwards rinsing it in acidulated water. Boiling water is sufficient to discolor articles thus dyed.

Red Dyes. The various shades of red are given by madder, cochineal, lac dye, safflower, &c.; fixed by aluminous or tin mordants. Less permanent dyes are produced by Brazil wood, peach wood, and archil. Some of these require peculiar treatment. Safflower contains a yellow as well as a red coloring matter. The first being soluble in cold water, is extracted by putting the safflower in a bag and kneading it under water. The safflower thus deprived of the yellow matter, yields its red color to alkaline liquids: at the time of using which, lemon-juice or some other acid is added sufficient to saturate the alkali. Pink saucers are made by adding the lemon juice to an alkaline infusion of washed safflower and allowing the coloring to deposit. Madder also contains a dun coloring matter which deteriorates the red unless previously removed. This may be partially effected by washing it in cold water: another mode is to treat the madder with its own weight of sulphuric acid, which carbonizes the other matters, but leaves the red color uninjured. As madder gives out but little of its red coloring matter to water, the decoction is not strained off, but the madder left in the bath. With acetate of iron,

madder yields a purple tint. Lac dye, as imported from India, requires acids for its solution. See LAC SOLVENT, below.

Yellow Dyes. These are given by French berries, quercitron bark, turmeric, weld, yellow wood, &c. Also by some mineral colors, as the following: The material to be dyed is first padded in a solution of bichromate of potash (8 oz. to a gallon of water), then in a solution of acetate or nitrate of lead. Cotton is dyed yellow by alternate dippings in iron liquor and lime-water, or solution of pearlash. A yellow color is given to silk by passing it through a mixture of equal measures of nitric acid (sp. gr. 1.288) and water, heated to 95° or 100° Fahrenheit, and from thence into a stream of water, or a mixture of chalk and water. This is termed mandarining.

Nankeen Dye is made by boiling annotto with an equal weight of pearlash in sufficient water. Orange is given by annotto; or by a mixture of red and yellow dyes; or by the successive application of acetate of alumina, a bath of quercitron, and the madder-bath.

Greens are given to woollens by first dyeing them blue, immersing the article in acetate of alumina, drying it, and finally immersing it in a quercitron bath. For silks, the order is reversed.

Browns are given by catechu, by walnut peels with alum, by redwood and copperas, by madder and black dye, &c.

Drabs are given by fustic with iron liquor.

Blacks. These are given by salts of iron, with galls, sumach, and logwood. The best black cloths are previously dyed blue with indigo.

By the mixture of various dyes, every variety of shade is produced: and often several tints from one coloring matter by the use of different mordants.

(The beautiful new dyes of the Aniline series, *Mauve*, *Magenta*, &c., are obtained from coal tar by various patented processes.)

Mordants are earthy and metallic compounds, the base of which unites both with the fibres of the material to be dyed and the coloring matter, thus rendering the dyes fixed. In *calico printing*, the mordants are formed into

a paste with some gum or other thickening material, and printed with wooden blocks on the cloth: which, after being dried, &c., is passed through the liquid dyes. The coloring matter combines with the part so printed, but is easily discharged from the other parts.

The principal Mordants are the following:

Alum Mordants. 1. Alum with one-fourth its weight of tartar.

2. *Acetate of Alumina.* (See page 301.) This is commonly prepared in solution for the purpose: 100 parts of alum in solution, with 150 parts of pyrolignite of lime of 20° B. density, is sometimes employed.

3. A solution of alum, with crystallized carbonate of soda, in the proportion of 1 oz. to each pound of alum.

4. HAUSMANN'S. This consists of a solution of alum with sufficient strong solution of caustic potash to redissolve the precipitated alumina; to which mixture a portion of linseed oil is added.

5. To 50 gallons of boiling water add 100 lb of alum; dissolve, and add slowly 10 lb of crystallized carbonate of soda. When the effervescence is over, add 75 lb of sugar of lead.

Tin Mordants. 1. *Protochloride of Tin.* To strong muriatic acid add gradually small pieces of grain tin till no more is dissolved. It may be obtained in crystals by evaporation. In redissolving them, it is necessary to add to the water a few drops of muriatic acid.

2. *Nitro-muriate, or Perchloride of Tin.* Mix 1 measure of nitric acid with 4 of muriatic acid, and add tin in small quantities as long as any is dissolved. Or mix 4 oz. of muriatic with 1 of nitric acid and 1 of water; dissolve in it, by small portions at a time, 2 drachms of grain tin.

3. Aquafortis (or equal parts of nitric acid and water) 8 parts, sal ammoniac 1 part; mix, and add gradually 1 part, or as much as it will dissolve, of grain tin.

4. Dr. BANCROFT'S *Murio-sulphate of Tin.* Digest 2 parts of tin with 3 of strong muriatic acid for an hour. Add very cautiously 1½ part of oil of vitriol. Keep up the heat as long as hydrogen is evolved; on cooling, it crystallizes. Dissolve this salt in water, so as to form a solution containing 1 part of tin in 8.

5. *New Tin Crystals.* Add 3 lb of sal ammoniac to a gallon of solution of tin; evaporate and crystallize.

6. *Mordant for Lac Dye.* Mix 27 lb of muriatic acid with 1½ lb of nitric acid (sp. gr. 1.19), put it into a stone bottle, and add tin in small bits till 4 lb are dissolved.

7. *Stannate of Soda.* Digest litharge 36 parts, or minium 27 parts, in a metallic vessel, with a soda lye of 1.35 dens.; when dissolved, 8 parts of tin in grains are gradually added. The lead separates at once in a spongy state, and the solution of stannate of soda may be decanted.

Lac Spirit, used as a *Solvent* for lac dye, in preference to muriatic acid alone, is thus made: Add gradually 3 lb of tin to 60 lb of muriatic acid. Digest ¾ lb of this solvent in each pound of the dye for 6 hours. Plum or puce spirit, peach spirit, and grain or scarlet spirit, are names given by dyers to different solutions of tin employed in dyeing these colors. For scarlet, the nitro-muriatic solutions (Nos. 2 and 3, above) are used.

Iron Liquor. Scraps of iron are placed in casks or other vessels, and covered with rectified raw pyroligneous acid. There are usually a series of vessels, through which the solution is successively passed till it is fully saturated.

[To produce good and permanent dyes, several successive processes are required, which cannot be described here. In domestic dyeing, for trifling articles, the dye and mordant are often boiled together, and the silk, &c., immersed in the liquid. The following are some examples: *Crimson.*—Boil 1 oz. Brazil-wood with 1 dr. of alum and ½ dr. of cream of tartar in a quart of water. *Purple, or Lilac.*—Archil 1 oz. (or cudbear 1 oz.), pearlash 1 dr., hot water a quart. *Rose, or Flesh-color.*—Pink saucers, with a little lemon-juice, will be found convenient. *Violet.*—Boil 4 oz. of logwood with 1 oz. alum and ¼ oz. of tartar, in a quart of water. *Blue.*—Add to the water as much sulphate of indigo (chemic blue) as will give it the required color. Or one of the other solutions of indigo (above) may be used. *Yellow.*—Boil 2 oz. of turmeric, or 4 oz. quercitron, or a drachm of saffron, with ½ oz. of alum, in a quart of water. *Green.*—Add to the yellow dye sufficient chemic blue to render it green. *Rose-red*, for silk.—Put your silk into a hot solution of alum and tartar; then into a hot decoction of madder. *Scarlet*, for silk.—Dye it yellow with saffron and alum; then put it into a decoction of cochineal and madder. *Black.*—Boil 1 oz. of logwood, ¼ oz. sumach, and 1 dr. of copperas, in 4 pints

of water. *Buff.*—Boil 1 oz. of fustic, 1 dr. of alum, in 4 pints of water.]

EAU DE JAVELLE. See CHLORIDE OF POTASH.

EGGS, TO PRESERVE. JAYNE'S Liquid (expired patent) is thus made: Take a bushel of lime, 2 lb of salt, ½ lb of cream of tartar, and water sufficient to form a solution strong enough to float an egg. In this liquid, it is stated, eggs may be preserved for two years.

ELAINE. See OLEINE.

ELECTRIC TISSUE. Steep linen or cotton in a mixture of strong sulphuric acid, and 3 of pure nitric acid, for an hour. Squeeze out the acid, wash with water until no sensible acidity remains, plunge it in a weak alkaline solution, then in water, and dry. By friction it yields a large quantity of resinous electricity.

ELECTRO-BRASSING. Mr. WATT. Dissolve powdered acetate of copper, 5 oz., in ½ gallon of water; add 1 pint of liquid ammonia; dissolve sulphate of zinc, 10 oz., in 1 gallon of water, at 180° F., and when cold add liquid ammonia 1 pint; dissolve potash, 4½ lb, in 1 gallon of water; lastly, cyanide of potassium 8 oz., in 1 gallon of hot water. Now mix in the following order; add the copper solution to the zinc, and then the potash and cyanide, digest for an hour or so, add water to make up 8 gallons. Work with a brass *anode* and active battery power, adding occasionally a little more ammonia and cyanide.

ELECTRO-BRONZING. BRUNEL'S Patent. Dissolve chloride of copper, 1 lb, in water ½ gallon, then carbonate of potash, 25 lb, in water 6 gallons; and sulphate of zinc, 2 lb, in hot water ½ gallon; mix the solutions; add nitrate of ammonia 12½ lb; stir, and add water to 20 gallons. Work with an *anode* of brass and active battery of two or more cells. Add occasionally some liquid ammonia and cyanide of potassium. These will prevent incrustations on the *anode*.

ELECTRO-COPPERING, for Iron or Zinc. The article must first be well cleaned, and "pickled" in dilute acid. Then dissolve sulphate of copper, 2 oz., in boiling rain-water; when cold add carbonate of potash, 4 oz., and liquid ammonia 2 oz. A precipitate forms, which is redissolved. Add cyanide of potassium, 6 oz., until all the blue color

disappears. Make up to 1 gallon, and work with a two-cell battery. The surface obtained may be bronzed. (See page 322.)

ELECTRO-GILDING. See GILDING.

ELECTRO-SILVERING. See SILVERING.

ELECTROTYPE MOULDS. These are sometimes made with fusible metals; sometimes with non-metallic compounds, having their surface covered with a conducting substance. The *fusible metal* is composed of 8 parts of bismuth, 3 of tin, and 5 of lead. The French clichée moulds consist of 8 parts of bismuth, 4 of tin, 5 of lead, and 1 of antimony. These are repeatedly melted together till perfectly mixed; and after being poured out on a suitable surface, are well stirred, and the model forcibly pressed on the compound at the moment it is about to become solid. *Composition Moulds* are made with 8 oz. of spermaceti, $1\frac{3}{4}$ oz. of white wax, and the same of strained mutton suit. These are melted together, and a little fine plumbago of flake-white stirred in. To copy plaster casts, the cast is set in a plate of hot water, with its face above the water, till it has absorbed water; it is then surrounded with a ring of cardboard, and the melted composition poured on it. The composition mould requires to be brushed over with finely-powdered genuine black-lead.

ELECTROTYPE MOULDS. *Elastic.* To 12 parts of carefully-melted glue, add 3 parts of treacle, and incorporate them perfectly. (For copying medals when the figures are in high relief.) *Gutta-Percha Moulds* are made by softening a piece of gutta-percha by warm water (150° to 160°) and pressing the medal into it by a screw. See SOLUTIONS.

ENGRAVING, PROCESS FOR CLEANING. Put the engraving on a smooth board, and cover it thinly with common salt, finely powdered; pour and squeeze lemon-juice upon this salt, so as to dissolve a considerable portion of it. Now elevate one end of the board, that it may form an angle of about 45° with the horizon. Pour lastly on the engraving boiling water from a tea-kettle, until the salt and lemon-juice be all washed off; the engraving will then appear perfectly clean, and free from stains. It must be dried gradually, on the same board, or on some smooth surface.—FRANCIS.

ENGRAVING MIXTURE, FOR WRITING ON STEEL. Sulphate of copper 1 oz., sal ammonia ½ oz. Pulverize separately, adding a little vermilion to color it, and mix with 1½ oz. of vinegar. Rub the steel with soft soap, and write with a clean hard pen, without a slit, dipped in the mixture.

ESSENCE OF JARGONELLE PEAR. Acetate of amylene is sold under this name. It is made by distilling a mixture of 1 part of oil of grain, 2 of acetate of potash, and 1 of oil of vitriol. Wash the diluted liquid with alkaline water, agitate with dry chloride of calcium, and redistil it from litharge.

ESSENCE OF PINEAPPLE. See BUTYRIC ETHER.

(For PERFUMED ESSENCES, see PERFUMERY. For CULINARY ESSENCES, see page 292.)

ETCHING FLUIDS. *For Lithography.* Dissolve 3 oz. of fused chloride of calcium in 9½ oz. of water, add to the solution 2 oz. of gum arabic, and 1 oz. of pure hydrochloric acid.—CHEVALLIER.

For Copper. 1. Aquafortis 2 oz., water 5 oz.; mix.

2. Iodine 2 parts, iodide of potassium 5 parts, water 5 to 8 parts.

3. CALLOT'S *Eau Forte, for Fine Touches.* Dissolve 4 parts each of verdigris, alum, sea-salt, and sal ammoniac in 8 parts of vinegar; add 16 parts of water, boil for a minute, and let it cool.

For Steel. 1. Iodine 1 oz., iron filings ½ dr., water 4 oz.; digest till the iron is dissolved.

2. Pyroligneous acid 4 parts, by measure, alcohol 1 part; mix, and add 1 part of double aquafortis (sp. gr. 1.28). Apply it from 1½ to 15 minutes.

3. Mix 10 parts of pure hydrochloric acid, 70 of distilled water, and a solution of 2 parts of chlorate of potash in 20 of water. Dilute before using with from 100 to 200 parts of water.

See PHOTO-LITHOGRAPHY and PHOTOGRAPHIC ETCHINGS ON STEEL.

ETCHING VARNISHES. 1. White wax 2 oz., asphaltum 2 oz.; melt the wax in a clean pipkin, add the asphaltum in powder, and boil to a proper consistence. Pour it into warm water, and form it into balls, which must be kneaded and put into taffeta for use.

2. White wax 2 oz., Burgundy pitch and black pitch each ½ oz.; melt together, and add by degrees 2 oz. of asphaltum in powder, and boil till a drop cooled on a plate becomes brittle.

3. Equal quantities of linseed oil and mastic, melted together.

FATS, TO OBVIATE RANCIDITY IN. Add oil of pimento or balsam of Peru, 2 drops to the ounce of lard or other fat.—Mr. T. B. GROVES.

FILTERING POWDER. Fuller's earth, washed, dried, and coarsely powdered, mixed with prepared bone-black (see CHARCOAL, ANIMAL) coarsely powdered.

FILTER FOR CORROSIVE LIQUIDS. Powdered glass or asbestos, or gun-cotton loosely packed in the neck of a funnel.

FININGS FOR GIN. Subcarbonate of potash 4 oz., Roche alum 8 oz. Brewers' finings consist of isinglass dissolved in stale beer.

FIRES, TO EXTINGUISH. Dr. CLANNY'S *Solution* consists of 5 oz. of sal ammoniac to 1 gallon of water. The compound used in Phillips's Fire Annihilator is said to consist of dried prussiate of potash, sugar, and chlorate of potash.

FIRES, COLORED. The ingredients for these compounds must be dry, not too finely powdered, and mixed very uniformly. The nitrate of strontian requires to be gently heated in an iron pan till it falls to powder. The chlorate of potash must be pulverized separately, and mixed very lightly with the other powders; the whole must then be passed through a sieve once or twice.

White Fires. 1. Nitre 30, sulphur 10, black antimony 5; mix.

2. Nitre 48, sulphur 13¼, black antimony 5 parts; mix.

3. Nitre 12, sulphur 16, black antimony 4, charcoal ¼, white arsenic ¼; mix.

4. Nitre 46¼, sulphur 23, meal powder 12½, zinc filings 18.

5. For stars: Nitre 57, sulphur 28, zinc filings 15.

Blue and Purple Fires. 1. Chlorate of potash 9, dried verdigris 2, sulphur 1 oz.; mix.

2. Nitre 12, sulphur 16, black antimony 4, charcoal ¼, orpiment ¼.

3. Chlorate of potash 9, sulphur 12, refiners' blue verditer 3 oz.; mix.

4. *Purple.* Chlorate of potash 5, nitrate of strontian 16, realgar 1, sulphur 2, lamp-black 1; mix.

5. Nitre 5, sulphur 2, metallic antimony 1; mix.

6. *Purple.* Chlorate of potash 2 oz., sulphur 1 dr., oxide of copper 1 oz.; mix.

7. *Violet.* Chlorate of potash 1 dr., pure copper ½ dr., sulphur 1 scruple, charcoal 16 gr.; mix.

Yellow Fires. 1. Nitre 3 oz., meal powder 3 oz., flowers of sulphur 3 oz., dried salt 2 oz.; mix.

2. Nitrate of soda 6, sulphur 1, lamp-black 1; mix.

Red Fires. 1. Chlorate of potash 10, nitrate of strontian 80, sulphur 26, charcoal 6; mix.

2. Chlorate of potash 3, nitrate of strontian 24, sulphur 7, lamp-black 1, sulphuret of antimony 2. (Half the lamp-black or charcoal only may be added at first; and if on trial it does not burn freely, add more.)

3. Chlorate of potash 1, nitrate of strontian 5, sulphur 1, black sulphuret of antimony 1; mix.

4. Chlorate of potash 2½, nitrate of strontian 20, sulphur 6½, sulphuret of antimony 2, charcoal ½; mix.

5. Chlorate of potash 5, nitrate of strontian 28, sulphur 13, sulphuret of antimony 4, lamp-black 1.

6. Chlorate of potash 35, nitrate of strontian 360, sulphur 93, sulphuret of antimony 60, charcoal 10.

7. Nitrate of strontian 72, sulphur 20, coal dust 2, gun-powder 6; mix.

8. Forty parts of nitrate of strontian, 13 of sulphur, 2 of lime-tree charcoal. Mix, and add by mixing with a horn spatula, 5 parts of dry and finely powdered chlorate of potash.

Lilac Fire. Chlorate of potash 49, sulphur 25, dry chalk 20, black oxide of copper 6 parts. For pans.

Green Fires. 1. Sulphur 10½, nitrate of barytes 62½, chlorate of potash 23½, sulphuret of arsenic 1¾, charcoal or lamp-black 1¾; mix.

2. Sulphur 13, nitrate of barytes 77, chlorate of potash 5, metallic arsenic 2, charcoal 3; mix.

3. Nitrate of barytes 20, sulphur 1½, sulphuret of antimony ½, chlorate of potash 10, charcoal ¼; mix.

4. Nitrate of barytes 10½ dr., sulphur 2 dr., chlorate of potash 162 gr., lamp-black 26 gr.; mix.

5. Dry nitrate of barytes 12 parts, sulphur 4, dry and finely-powdered chlorate of potash 5 parts. The chlorate to be mixed by a horn spatula.

COLORED FLAMES. The flame of alcohol may be colored by mixing certain salts with the spirit. A green color is given by muriate of copper, or boracic acid; red by nitrate of strontian, nitrate of iron, or nitrate of lime; yellow by nitrate of soda, &c.

FIRE-PROOFING. *For Paper*, see PAPER. For *Dresses*, &c.: A strong solution of sulphate of ammonia. The dresses of stage dancers may be soaked in a weak solution of chloride of zinc. The tungstate of soda is said to be the only perfect fire-proofer.

FLINTS, LIQUOR OF. Soluble glass. Mix seventy parts of pearl-ash, 54 of washing soda, and 152 of silicious sand, and fuse the mixture in a crucible. It is soluble in water, and the filtered solution evaporated to dryness leaves a transparent glass. It has been proposed to render wood, muslins, &c., incombustible by means of the solution. Dr. TURNER directs 3 parts of carbonate of potash, and 1 of silica. See SOLUBLE GLASS.

FLOWERS, COMPOUND FOR PROMOTING THE BLOWING OF. Sulphate of ammonia 4 oz., nitre 2 oz., sugar 1 oz., hot water 1 pint. Keep it in a well-corked bottle. For hyacinth glasses add 8 or 10 drops of the liquid to the water, changing the water every 10 or 12 days. For flowering plants in pots, add a few drops to the water employed to moisten them.

FLOWERS, TO PRESERVE LIVING. Add 1 spoonful of charcoal powder to the water in which the boquet is placed.—*French Journal.*

FLOWERS AND PLANTS, TO PRESERVE. Any vegetable substance may be preserved moist in a solution of creasote, or in glycerine. The method of drying plants between sheets of paper needs no description. But the original form, and in many instances the color, of a fresh flower may be preserved by carefully immersing it in some fine, dry material, and then rapidly drying in a baking oven. Millet seed has been used for this purpose, and may answer well for coarse specimens. For fine ones, white river sand in equal grains must be used. To separate large grains it should be passed through a sieve; to remove fine particles it is copiously

washed with water. While drying it is to be constantly agitated.

M. Revel recommends further, that 1000 parts of this sand be intimately mixed with 1 of stearic acid and 1 of spermaceti, before using. See next formula.

Flowers, to Preserve in their Natural Shape and Color. Provide a vessel with a movable cover. Fit to the top a piece of fine metallic gauze, and replace the cover. Pass through sieve into an iron pot sand sufficient to fill this vessel, and heat it with ½ per cent. of stearine, carefully stirring. Place the flowers on the gauze, and, removing the bottom of the vessel, pour in the sand and stearine, so as to cover and envelop them. Place on the top of an oven for 48 hours. Remove the cover, invert the vessel, and the sand runs away through the gauze, leaving the flowers dried in their natural position.—*Journ. Soc. Arts.*

Fluxes. In a general sense these are substances which promote the fusion of minerals, but particularly which cleanse a reduced metal, by assisting its separation from its impurities. They also serve to defend it from the action of the air, and some of them assist in the reduction of oxides.

Black Flux. Into an earthen crucible, heated sufficiently hot to cause feeble combustion, but not to fuse the flux, throw successive portions of a mixture of 1 part of nitre, and 2 of crude (or cream of) tartar. Keep the flux in a close bottle.

White Flux. Into a large earthen crucible, heated to redness, throw successive portions of a mixture of 2 parts of nitre and 1 of tartar. Keep it as the last.

Crude Flux, is the mixture of nitre and tartar, before deflagration.

Dr. Christison's *Flux for reducing Arsenic.* Mix crystallized carbonate of soda with ⅛ of charcoal, and heat gradually to redness.

Fresenius's *Flux*, for reducing sulphuret of arsenic. Dry carbonate of potash 3 parts, cyanide of potassium 1 part.

Cornish Flux. Cream of tartar 10 parts, nitre 3½, borax 3.

Morveau's *Flux*. Pulverized glass (free from lead) 8 parts, calcined borax ½ part, charcoal ½ part.

Mr. Taylor's *Flux*. Saturate a solution of tartaric acid with carbonate of soda, evaporate to dryness, and calcine in a covered platinum crucible.

Sal enixum (the acid sulphate of potash left in distilling nitric acid), sandiver or glass-gall, fluor spar, limestone, &c., are also used as fluxes.

Fly Poison. A common poison for flies consists of white arsenic, or King's yellow, with sugar, &c., but the use of such compounds may lead to fatal accidents. A sweetened infusion of quassia answers the same purpose, and is free from danger. Pepper, with milk, is also used; and also some adhesive compounds by which they are fatally entangled. *Papier moure* contains a large quantity of arsenic.

Freezing Mixtures. The salts should be in a crystallized state, with as much water in them as possible without being damp. They should be coarsely pulverized at the time of using, and put into the water contained in a basin or other suitable vessel. The water to be frozen should be inclosed in a thin vessel, and immersed in the freezing mixture. To obtain extreme degrees of cold, the ingredients and vessels employed should be previously cooled by one of the freezing mixtures.

1. Sal ammoniac 5 oz., nitre 5 oz., water 16 oz.
2. Mix 4 oz. of nitrate of ammonia, 4 of crystallized carbonate of soda, and 4 of water. In 3 hours 10 oz. of water may be frozen.
3. Nitrate of ammonia and muriate of ammonia in equal proportions, water q. s.
4. Nitrate of ammonia 5 parts, nitrate of potash 5, sulphate of soda 8, water 16 parts.
5. Phosphate of soda 9 parts, diluted nitric acid 4 parts.
6. Sulphate of soda 8 parts, muriatic acid 5 parts.
7. Sulphate of soda 6 parts, nitrate of ammonia 5 parts, diluted nitric acid 4 parts.
8. Mix 1 part by weight of powdered sal ammoniac with 2 of powdered nitre. Reduce common washing soda to powder. Keep these powders in well-closed bottles, and when required for use take equal measures of each, and

add an equal bulk of water, or sufficient to dissolve the salts.

9. Muriate of ammonia 11 dr., nitre 10 dr., sulphate of soda 2 oz. Powder separately, and mix in a tin vessel with 5 oz. of water.

With Ice.

1. Snow or pounded ice 2 parts, muriate of soda 1 part.
2. Snow 2 parts, crystallized muriate of lime 3 parts.
3. Snow 8 parts, muriatic acid 5 parts.
4. Snow or pounded ice 12 parts, muriate of soda 5 parts, nitrate of ammonia 5.
5. Snow 7 parts, diluted nitric acid 4 parts.
6. Snow 3 parts, diluted sulphuric acid 2 parts.

French Polish. This is an alcoholic solution of shellac; some of the softer resinous gums are usually added, but too much of them renders the polish less durable. Highly rectified spirit, not less than 60 over proof, should be used. Rectified wood naphtha is sometimes substituted, to which the unpleasant smell is the only objection.

1. Orange shellac 22 oz., rectified spirit 4 pints; dissolve.
2. Shellac 3 oz., gum sandarac ½ oz., rectified spirit a pint.
3. Shellac 4 oz., gum thus ½ oz., rectified spirit a pint; dissolve, and add almond or poppy oil 2 oz.
4. Shellac 5 oz., seed-lac 1 oz., gum juniper ½ oz., mastic 1 oz., rectified spirit a pint.
5. Shellac 3 oz., seed-lac 3 oz., gum juniper 1½ oz., mastic 1 oz., rectified spirit a quart.
6. Shellac 5 oz., oxalic acid ½ oz., rectified spirit a pint; dissolve, and add linseed oil 4 oz.
7. Shellac 5 oz., gum benzoin 5 oz., oxalic acid 10 dr., rectified spirit a quart; dissolve, and add ½ pint of linseed oil.
8. Shellac 8 oz., gum elemi 1½ oz., rectified spirit 4 pints.
9. Shellac 10 oz., seed-lac 6 oz., gum thus 3 oz., sandarac 6 oz., copal varnish 6 oz., rectified naphtha 8 pints. Or dissolve 8 oz. each of seed-lac, gum thus, and sandarac, separately in a pint of naphtha; and 1 ℔ of shellac in 8 pints of naphtha. Then mix 6 oz. of copal varnish, 12 oz.

of the solution of seed-lac, 6 oz. of the solution of frankincense, 12 of the solution of sandarac, and $5\frac{3}{4}$ lb of the solution of shellac. Let the copal varnish be put into the tincture of shellac, and well shaken, and the other ingredients added. A correspondent informs me that this polish cannot be excelled.

10. Copal $\frac{1}{4}$ oz., gum arabic $\frac{1}{4}$ oz., shellac 1 oz. Pulverize, mix, and sift the powders, and dissolve in a pint of spirit.

11. Shellac 5 oz., rectified naphtha a pint.

French polish is sometimes colored with dragon's blood, turmeric root, &c. The general directions for preparing the polish are to put the gums with the spirit in a tin bottle, and set it on the hob, or in water, so as to keep it at a gentle heat, shaking it frequently. The cork should be loosened a little before shaking it, taking care that there is no flame near to kindle the vapor. When the gums are dissolved, let it settle for a few hours, and pour off the solution from the dregs. The method of using it is to have a roll of list, over the end of which five or six folds of linen rag are placed. The polish is applied to the linen with a sponge, and a little linseed oil is dropped on the centre of it.

Fulminating Compounds. *Fulminating Powder.* Mix together in a warm mortar 3 parts of pulverized nitre, 2 of dry subcarbonate of potash, and 1 of sulphur. A small quantity heated on an iron shovel or ladle till it fuses, suddenly explodes with great violence. It should be used with great caution. Another kind of fulminating or detonating powder is made by mixing 3 gr. of chlorate of potash with 1 of sulphur: by strongly triturating it with strong pressure in a marble mortar, a succession of sharp explosions is produced. The same mixture, or 6 gr. of chlorate of potash, 1 of sulphur, and 4 of charcoal, struck with a hammer on an anvil, gives a loud report.

Chloride of Nitrogen and *Iodide of Nitrogen* cannot be meddled with without extreme danger. *Fulminating gold*, and the *fulminating silver* prepared with ammonia, are also dangerous compounds, even in minute quantities. As they serve no practical use, the mode of preparing them is omitted.

Fulminating Mercury (Howard's, as improved by Dr. Ure). Dissolve by a gentle heat 1 oz. of quicksilver in $7\frac{1}{2}$ fluid

oz. (or 10 oz. by weight) of nitric acid, of 1.4 specific gravity, in a glass retort, the beak of which is loosely inserted into a large balloon, or bottle. When the mercury is dissolved, the solution, at the temperature of 130° Fahrenheit, should be slowly poured through a funnel into 10 fluid oz. of alcohol of 0.830 specific gravity, contained in a vessel that will hold 6 times the quantity of ingredients. When the action ceases, pour the contents of the matrass on a double filter in a glass funnel; wash out any powder that may remain in the matrass with a little cold water, and throw this also on the filter; and wash the fulminate with more water till it passes free from acid. When sufficiently drained, lift the filter out of the funnel, and lay it open on a copper or earthen plate, and dry the fulminate at 212°, or under, by hot water or steam. Its manufacture requires great caution; some valuable lives have been sacrificed in its preparation.

Fulminating Silver (BRUGNATELLI'S). On 100 gr. of pulverized nitrate of silver, in an open glass vessel, pour first an oz. of alcohol, and then as much strong nitrous acid. The mixture boils, and gives out ethereal vapors. When all the powdered nitrate has taken the form of white clouds, cold distilled water must be added to sustain ebullition, otherwise the fulminate will be dissolved. Collect the powder on a filter, and dry it at a low temperature. Dr. TURNER directs 1 part of silver to be dissolved in 10 of nitric acid, at a gentle heat, 20 parts of rectified spirit to be added, and the mixture warmed. When it begins to boil, set it aside to cool, collect and wash the crystals on a filter, and carefully dry them. This is more violent and dangerous than fulminating mercury.

FUMIGATIONS. See Fumigatio, Pocket Formulary, for their medicinal uses. Though not strictly belonging to this place, it may be useful to give a few directions for the management of these important agents, as disinfectants and purifiers.

Chlorine Fumigation. This is probably the most effective in destroying noxious effluvia, and putrid odors, and in checking the spread of contagious diseases. But as the gas itself is deleterious, except in a very diluted state, it must be used with caution in occupied apartments. To disinfect rooms

from which the occupants have been removed, mix common salt and black oxide of manganese in equal quantities. Mix also in an earthen basin equal weights of oil of vitriol and water, and when it has cooled, put it into a bottle for use. Into a china or earthen basin put from 1 to 3 oz. of the powder, according to the size of the room, and pour over it twice or thrice as much of the mixed acid. Place it in the apartment, and close the doors and windows for a few hours; the doors and windows are then thrown open till the smell of chlorine disappears. Dr. A. T. THOMSON directs a mixture of 1 oz. of salt and ½ oz. of oxide of manganese to be put into a china cup, and 6 fluid dr. of oil of vitriol poured on it, the cup being placed in a pipkin of hot sand. Instead of the above ingredients, some chloride of lime may be placed in a large jar or basin, and a mixture of acetic acid and water poured on it. When used in or near the apartments of the sick, great care must be taken that the chlorine is so diluted with air that it shall occasion no annoyance to the invalid. Some contrivances have been adopted to render the extrication of chlorine gradual and continued. SMITH'S Chlorine Fumigator, and the more simple one of Messrs. HEATHFIELD & BURGESS, are very convenient. Another method is proposed by Mr. SCANLAN, in the "Pharmaceutical Journal," vol. vii, page 343. By such contrivances chlorine may with care be safely employed in houses occupied by the sick (in the passages, stairs, &c.) to prevent the spread of infectious fevers; but chloride of lime, simply mixed with water, in the proportion of not more than 1 oz. to a quart, is usually sufficient to purify the chamber of the sick. It should be occasionally sprinkled on the floor, and also placed about the room in shallow dishes, or a linen cloth moistened with it, suspended on a line. The same method may be pursued in all places where unpleasant smells prevail.

Nitric Fumigation. Put into a china cup equal measures of sulphuric acid and water, and add to it, from time to time, small quantities of powdered nitre; or put 2 or 3 dr. of powdered nitre into a cup, and pour over it about an equal quantity of oil of vitriol. Stir it with a piece of glass, or tobacco-pipe, and remove it from time to time to different parts of the apartment. For large rooms 2 or 3 cups may be required. It is often recommended to apply heat; but

Dr. BATEMAN, of the Fever Hospital, found this unnecessary and objectionable, especially in the apartments of the sick. No metallic or wooden stirrers, or vessels, must be used.

Muriatic Fumigation. This is now almost disused, being less efficacious than the preceding. It is obtained by putting a few drachms of common salt into a cup, and pouring on it an equal quantity of oil of vitriol. The vapors are very injurious to the lungs.

Acetic Fumigation. The vapor of vinegar, and especially of strong acetic acid, is employed as a disinfectant, but its efficacy is now considered to be very limited. It may be used by keeping the vinegar boiling over a lamp. A coarser method sometimes used is to plunge a red-hot poker into a cup of vinegar. Aromatic vinegar, merely held to the nose, may afford some slight protection to those who attend upon the sick.

Sulphur Fumigation. The fumes of burning sulphur may possibly have some effect in decomposing miasmata and noxious effluvia; but as they have no advantage over chlorine, and are very disagreeable, and otherwise objectionable, they are not likely to be employed. Formerly the following powder was burnt to destroy contagious miasmata. Flowers of sulphur, nitre, and powdered myrrh, of each 1 oz.

Tar Fumigation. The vapor of boiling tar has been used as a disinfectant, as well as a palliative in some affections of the respiratory organs. The usual plan is to keep the tar boiling over a lamp. See Fumigato Picea, Pocket Formulary.

Benzoin, styrax, and other odoriferous gums, cascarilla bark, coffee berries, and the compound termed aromatic pastiles, are burnt as purifiers and disinfectants. But little confidence is now placed in them as prophylactics against infection. The same may be said of camphor and tobacco. They should not be depended on to the exclusion of more efficient means, nor be made a substitute for free ventilation and the removal of all sources of noxious effluvia, when practicable.

FUMIGATING PASTILES. See PERFUMERY.

FURNITURE CREAM. 1. Soft water a gallon, soap 4 oz., beeswax in shavings 1 ℔; boil together, and add 2 oz. of pearl-

ash. To be diluted with water, laid on with a paint brush, and polished off with a hard brush or cloth.

2. Wax 3 oz., pearlash 2 oz., water 6 oz.; heat them together, and add 4 oz. of boiled oil, and 5 oz. spirit of turpentine.

3. The same is sometimes given to a mixture of 1 oz. of white or yellow wax, with 4 of oil of turpentine.

FURNITURE PASTE. 1. Melt 1 lb of beeswax with ¼ pint of linseed oil, and add ½ oz. of alkanet root; keep it a moderate heat till sufficiently colored; then remove from the fire, add ¼ pint of oil of turpentine, strain through muslin, and put it into small gallipots to cool.

2. Scrape 4 oz. of wax, and put it into a pipkin with as much oil of turpentine as will cover it, and ¼ oz. of powdered resin; melt with a gentle heat, and stir in sufficient Indian red to color it.

3. Equal weights of beeswax, spirits of turpentine, and linseed oil.

FURNITURE OIL. 1. Linseed oil a pint, alkanet ½ oz.; digest in a warm place till colored, and strain.

2. The same with ¼ pint of oil of turpentine.

3. Linseed oil a pint, alkanet root 1 oz., rose pink 1 oz.; let them stand in an earthen vessel all night.

4. A quart of linseed oil, 6 oz. of distilled vinegar, 3 oz. of spirit of turpentine, 1 oz. of muriatic acid, and 2 oz. of spirit of wine.

5. Linseed oil 8 oz., vinegar 4 oz.; oil of turpentine, mucilage, rectified spirit, each ½ oz.; butter of antimony ¼ oz.; muriatic acid 1 oz.; mix.

6. Linseed oil 16 oz., black resin 4 oz., vinegar 4 oz., rectified spirit 3 oz., butter of antimony 1 oz., spirit of salts 2 oz.; melt the resin, add the oil, take it off the fire, and stir in the vinegar; let it boil for a few minutes, stirring it; when cool put it into a bottle, add the other ingredients, shaking all together. (The last two are especially used for reviving French polish.)

7. Linseed oil 1 pint, oil of turpentine ½ pint, rectified spirit 4 oz., powdered resin 1½ oz., rose pink ½ oz.; mix.

8. Linseed oil 14 oz., vinegar 1½ oz., muriatic acid ½ oz.; mix.

FUSIBLE METAL. See ALLOYS.

GALL, CLARIFIED. Ox-gall is prepared for the use of artists in the following manner: To a pint of fresh ox-gall, boiled and

skimmed, add 1 oz. finely-powdered alum; leave it on the fire till the alum is dissolved, then let it cool, put it into a bottle, and cork it loosely. Treat another pint in the same way with 1 oz. of salt instead of alum. After standing more than three months, carefully decant from each bottle the clear portion, and mix them together. The coloring matter is precipitated, and a clear colorless liquid is obtained by filtration. It is used for mixing artists' colors, and to prepare ivory, oiled paper, &c., to revive the colors. Also for taking out grease-spots.

GALVANIC BATTERIES, SOLUTIONS FOR. See page 307.

GANNAL'S SOLUTION. See page 311.

GANTEINE. (A composition for cleaning kid gloves; sometimes improperly termed *Saponine*.) Dissolve 3 oz. of soap by heat in 2 oz. of water, and when nearly cold add 2 oz. of eau de Javelle, and 1 dr. of water of ammonia; form a paste, which is to be rubbed over the glove with flannel till sufficiently clean.

GARANCINE. Madder (sometimes the spent madder of the dyer's bath) is mixed with its weight of oil of vitriol, added very gradually, so as to avoid overheating. The acid is then washed out.

GARROT'S COVERING FOR PILLS. Soak 1 oz. of purified gelatine in 2 or 3 dr. of water; keep it liquefied in a salt-water bath. The pills are stuck on long pins, and dipped in the solution; when cold the pins are withdrawn, after being heated by a small flame, which melts the gelatine and closes the hole.

GASES. These are generated in gas bottles fitted, by grinding with an S-formed tube; or in flasks to which a bent tube is adapted by means of a cork; in a common retort; or sometimes in iron bottles with a metal tube. They are usually collected in vessels filled with water placed with their open ends in a vessel of water. *Pepy's Gas Holder* is very useful for receiving, retaining, and transferring gases. The *pneumatic trough* consists of a vessel for holding water, with a shelf for sustaining the jars or bottles that are to be filled; these are filled by sinking them under water, and are then lifted on the shelf, above which the water rises, with their open end downwards. The beak of the retort, or bent tubes, are so placed that the gas issuing from them rises

through the water into the vessel, and takes the place of the water in them. Some gases being very easily absorbed by water are collected over mercury. Sometimes they are collected in dry bottles. For light gases, as ammonia, place a bottle in a vertical position with its mouth downward over the tube from which the gas issues, which should touch the bottom of the bottle. When the bottle is filled with gas, and this escapes from the mouth, quietly withdraw the tube and close the bottle, still inverted, with a greased stopper. For gases heavier than air, as chlorine, or carbonic acid, the bottle must be placed with its mouth uppermost, and the tube delivering the gas must descend to the bottom of the bottle. When full of the gas, close it with the greased stopper. The tube connected with a flask in which a gas is generated should have a ball blown in it, into which asbestos may be introduced to arrest any particles thrown up by effervescence.

The following are the processes for procuring the principal gases:

Ammoniacal Gas. This is obtained by mixing equal weights of slaked lime (previously cooled) and powdered sal ammoniac, and heating the mixture in a retort or flask. As water rapidly absorbs the gas, it must be collected over mercury, or in a dry bottle, as described above.

Carbonic Acid Gas. This is obtained by acting on marble or chalk, or carbonate of soda or potash, by a diluted acid. For exact experiments it must be collected over mercury; otherwise it may be collected in a bottle (as above). Mr. Benson states that a saturated solution of sulphate of magnesia may be used in collecting this gas, instead of mercury.

Carbonic Oxide is obtained by acting on binoxolate of potash with 6 times its weight of oil of vitriol at a gentle heat; or by strongly heating, in an iron bottle or gun-barrel, equal weights of chalk and iron filings. The gas must be passed through water containing lime or caustic potash to absorb the carbonic acid gas.

Chlorine. This gas may be obtained from oxide of manganese, common salt, and sulphuric acid, as directed in making chloride of lime (p. 332). Or from muriatic acid and oxide of manganese (See Chlorinium, Pocket Formulary). But more conveniently, on the small scale, by dissolving com-

mon salt in water, adding $\frac{1}{6}$ of its weight of nitric acid, and as much oxide of manganese. Apply a gentle heat, and the gas is abundantly produced without violent action. (Chemist, vol. i.)

Hydrochloric Acid Gas. It may be obtained by heating together in a flask, equal weights of salt and oil of vitriol; or simply by heating strong muriatic acid. It must be collected over mercury; or otherwise conducted to the bottom of a dry bottle, as described above.

Hydrogen Gas is readily procured by pouring on fragments of zinc, in a glass bottle, or flask with a bent tube, or retort, some diluted sulphuric acid (1 measure of strong acid to 5 of water). It may be collected over water. If zinc be not at hand, fine iron wire, or the turnings or filings of iron may be substituted for it. To procure gas of great purity, distilled zinc must be used.

Carburetted Hydrogen Gas. Light carburetted hydrogen is readily obtained by stirring the mud of stagnant pools. Heavy carburetted hydrogen is prepared by heating 1 part of alcohol with 6 or 7 of oil of vitriol, and conducting the mixed gases through milk of lime, which retains the sulphurous acid; and afterwards through oil of vitriol, which absorbs water, ether, and alcohol. Coal gas is a mixture of these gases, with other hydrocarbons, &c.

Phosphuretted Hydrogen. The spontaneously inflammable variety of this gas is made by boiling phosphorous with solution of potash in a small retort, the beak of which is kept under water: as each bubble of gas rises from the water, it inflames, and forms a ring of white smoke, which dilates as it ascends. The spontaneous inflammability of the gas when mixed with atmospheric air or oxygen renders caution necessary in its preparation. The other varieties of phosphuretted hydrogen have no special interest or application.

Sulphuretted Hydrogen. Mix together 2 parts of iron filings with 1 of sulphur into a thin pap with water, and heat it gently in an iron vessel. Combination takes place with the evolution of heat. Cover it till cold. On this compound, contained in a glass bottle, or other suitable apparatus, pour sulphuric acid previously diluted with 7 parts of water. If more acid be afterwards required, dilute the strong acid with only 4 of water. It is absorbed by water.

Nitrogen, or Azote. Atmospheric air may be made to yield an unlimited supply of nitrogen, by exposing it to the action of substances which combines with it oxygen. By burning phosphorous in a large bell-glass standing in water, and allowing it to stand over the water a few hours, nearly pure nitrogen is obtained, which may be further purified by agitating it with solution of pure potash. CORENWINDER procures it from his solution of nitrate of potash (which see) by mixing 1 measure of it with 3 of concentrated solution of sal ammoniac, and heating the mixture in a flask. The gas contains a little ammonia, from which it may be freed by passing it through diluted sulphuric acid.

Protoxide of Nitrogen. Nitrous Oxide of DAVY. It is most conveniently made by heating nitrate of ammonia (formed by neutralizing pure nitric acid, diluted with 3 parts of water, with carbonate of ammonia, and boiling the solution till a drop let fall on a cold plate solidifies, adding a little ammonia towards the end, to insure neutrality) in a retort, at a heat not exceeding 500° Fahrenheit, till it is nearly all decomposed. It may be collected over warm water.

Oxygen Gas. Mix chlorate of potash with a third of its bulk of black oxide of manganese; put the mixture into a glass bottle, or clean flask, to which a bent tube is fitted by a cork, and apply a gentle heat. The gas, which comes over freely, may be collected over water.

Ozone Gas. This is supposed to be oxygen in an allotropic and more active state. It is formed by putting into a wide bottle pieces of clean phosphorus, with a little water, so that the phosphorus shall be partly in the water and partly uncovered. Close the bottle for some hours, when the air it contains will manifest the odor and effects of ozone.

Sulphurous Acid Gas. It is procured in a nearly pure state by heating 2 parts of quicksilver with 3 of oil of vitriol, and collecting the gas over mercury. By passing the gas through a tube surrounded with a freezing mixture, it is condensed into a liquid. For ordinary purposes, the gas may be obtained as directed under ACID, SULPHUROUS.

GELATINE, PURIFIED. *Grenetine.* It is made by various processes

from gelatinous animal matters. NELSON'S *Patent Gelatine* is made from the cuttings of the hides of beasts, and skins of calves. These, freed from hair, flesh, fat, &c., are washed and scored, then macerated for 10 days in a lye of caustic soda, and afterwards placed in covered vessels at a temperature of 60° to 70° until they become tender; then washed from the alkali, exposed to the vapor of burning sulphur until they become sensibly acid, dissolved in earthen vessels heated to 150°, strained, put into settling vessels heated to 100° or 120° for nine hours, the clear liquor drawn off, and poured on the cooling slabs to the depth of ½ an inch. When cold, the jelly is cut in pieces, washed till free from acid, redissolved at 85°, poured on slabs, cut up, and dried on nets.

Bone Gelatine. The bones are boiled to remove the fat, then digested in diluted muriatic acid till the earthy matter of the bone is dissolved. The gelatine, which retains the form of the bone, is washed in a stream of water, plunged in hot water, and again in cold, to remove all remains of acid, and sometimes put into a solution of carbonate of soda. When well washed, it is dried on open baskets or nets. By steeping the raw gelatine in cold water, dissolving it in boiling water, evaporating the jelly, and cutting it into tablets, it may be dried, and preserved in that form.

GERMAN PASTE, for feeding insectivorous singing-birds. Blanched sweet almonds 1 ℔, pea meal 2 ℔, butter 3 oz., saffron a few grains, honey q. s. Form the whole into a paste, and granulate it by pressing it through a cullender. Some add the yolks of 2 eggs.

GILDING. Leaf gold is affixed to various surfaces, properly prepared by gold size or other adhesive medium. Metallic surfaces are coated with gold by means of amalgam of gold and mercury, applied with a wire brush, wet with an acid solution of mercury (made by dissolving 10 parts of mercury in 11 of nitric acid, by a gentle heat, and adding 2½ parts of water). This article thus coated is heated over charcoal till the mercury is dissipated, and afterwards burnished. To give it a redder color, it is covered with gilders' wax (a compound of verdigris, ochre, alum, and yellow wax); again exposed to heat, and afterwards washed and cleaned by a scratch brush and vinegar. An inferior kind of gilding is effected by dissolving gold, with a fifth of its weight of

copper, in nitro-muriatic acid, dipping rags in the solution, drying and burning them, and rubbing the ashes on the metallic surface with a cork dipped in salt and water.

GILDING, by immersion. Dissolve teroxide or terchloride of gold in a solution of pyrophosphate of soda, and dip the article to be gilt in it.

Electro-gilding, by ELKINGTON'S patent process, is thus performed: A solution of 5 oz. of gold (see ACID, NITRO-MURIATIC, p. 305) is prepared, and boiled till it ceases to give out yellow vapors; the clear solution is mixed with 4 gallons of water, 20 lb of bicarbonate of potash added, and the whole boiled for 2 hours. The articles, properly cleaned, are suspended on wires, and moved about in the liquid from a few seconds to a minute, then washed, dried, and colored in the usual way.

The solution used in gilding with the voltaic apparatus is made by dissolving ¼ oz. of oxide of gold, with 2 oz. of cyanide of potassium, in a pint of distilled water.

GINGER BEER. See BEVERAGES.

GINGERBREAD, PURGATIVE. See page 185.

GLASS OF BORAX. Calcine borax with a strong heat till the water of crystallization is expelled, and the salt fuses into a clear glass.

GLASS, SOLUBLE. Mix 10 parts of carbonate of potash, 15 of quartz (or of sand free from iron and alumina), and 1 part of charcoal. Fuse together. The mass is soluble in 4 or 5 parts of water; and the filtered solution evaporated to dryness yields a transparent glass, permanent in the air.—FUCHS.

GLASS. The different kinds of common glass consist essentially of silica with soda or potash; most of the white kinds also contain oxide of lead. Crown glass and green bottle glass contain a portion of lime. Green glass and some kinds of foreign white glass are free from lead, and should therefore be selected for chemical uses.

To mark on Glass. Glass may be written on for temporary purposes, by French chalk; pencils of this substance will be found convenient. Glass may be written on with ink if the surface is clean and dry, and the pen held nearly

perpendicular. The shellac ink (see INK) is the best for labels, as it resists damp, &c. *To scratch glass*, a scratching diamond is used; or a piece of flint, or crystal of quartz, or the point of a small 3-square file. *To engrave on glass*, fluoric acid is used, either in the liquid state or in vapor. The glass must be warmed, and coated with wax, or engravers' cement, and the writing or design traced through the wax with a brad-awl, or other pointed instrument. The liquid fluoric acid is poured on it, and left to act on the uncovered portions of the glass; or the fluorspar may be powdered and made into a paste with oil of vitriol, and laid over the prepared surface, and covered with lead-foil or tea-lead; or bruised fluorspar is put in a Wedgewood evaporating basin, with sufficient oil of vitriol to form a thin paste, and the prepared glass laid over the basin, so that the vapors may act on the portions from which the wax has been removed. *To cut glass* (beside the usual method of dividing cut glass by a glazier's diamond), the following means may be used: To divide glass tubes or rods, form a deep mark round them with the edge of a sharp 3-square file, then with a hand placed on either side of the mark, break the rod with a slightly stretching as well as bending motion. A diamond or sharp flint may be substituted for a file. Flasks, globes, and retorts may be divided by means of iron rings, having a stem fixed in a wooden handle. Make the ring red-hot, and apply it to the flask, &c. If the vessel does not break where it came in contact with the ring, wet the part, and it will generally separate. Another method is to twist together 2 or 3 threads of cotton, such as is used for wicks, moisten them with spirit of wine, and encircle the flask with them; then, holding the flask horizontally, set fire to the wick, and turn the flask with the fingers, so as to keep the flame in the direction of the thread. If the separation does not take place the first time, the process may be repeated after the glass has cooled. By these means a common oil-flask may be divided into an evaporating dish and a funnel. By means of a stout iron rod, fixed in a wooden handle, and terminating in a blunt point, and heated to redness, broken retorts, globes, and flasks may be converted into useful evaporating dishes, &c. If any crack exists, it may easily be led in any direction, as it will follow the motion of the heated iron. If no crack exists, one must be produced by applying the point of the

heated rod to any convenient spot on the edge of the broken glass, touching it afterwards with a moistened finger, if necessary. The edges of glass thus divided are rendered less apt to break by heating them in the flame of a blow-pipe, or grinding them smooth with emery on a flat stone. See FARADAY'S Manipulations.

Glass, to Platinize. M. DULLO. This is recommended to prevent fusing of the end of the tube used in MARSH'S test for arsenic. In drawing out the end of the tube, leave the diameter slightly larger than is necessary; then roughen the narrow end with a file. Dip in a solution of bichloride of platinum, containing one-twentieth of the metal; remove excess of the drop, and heat cautiously till the glass acquires a metallic appearance. Repeat this four or five times.

Glass, to Silver. The term *silvering* is commonly applied to the process of coating the surface of glass with amalgamated tinfoil, in forming mirrors. The tinfoil is rubbed over with quicksilver, and more of the latter poured over it: the plate of glass, perfectly clean and dry, is then applied to it in such a way as to exclude all air-bubbles, and to bring the glass and tinfoil into perfect contact. The plate, after being inclined so as to allow the superfluous quicksilver to drain off, is loaded with weights, under which it remains till the adhesion is complete. To convex and concave mirrors the amalgamated foil is applied by means of accurately fitting plaster moulds. The interior of globes is silvered by introducing a liquid amalgam (see AMALGAM, p. 310), and turning about the globe till every part is covered with it.

To Silver Glass. PETTIJEAN'S *process.* 1540 gr. of nitrate of silver are treated with 955 gr. of strong solution of ammonia, and afterwards with 7700 gr. of water. To this solution, when clear, are added 170 gr. of tartaric acid dissolved in 680 of water, then 152 cubic inches more of water, with brisk agitation. When it has settled, the clear part is poured off; 152 cubic inches of water are added to the sediment, to dissolve as much as possible. The clear fluids are mixed, and again 62 cubic inches of water are added. This is Silvering Solution No. 1. No. 2 is prepared in the same manner, but with twice as much tartaric acid. The glass plate being cleaned, and laid horizontally,

an even layer of solution No. 1 is poured on it, to about the depth of $\frac{1}{10}$th of an inch. Heat is then applied by means of a cast iron water-bath beneath. Bright silver is soon deposited. When the stratum of fluid is exhausted, it is poured off, and solution No. 2 next applied in the same way. The silver surface may be polished, or coated with black varnish.

M. LIEBIG proposes to silver glass without mercury by a solution of ammonio-nitrate of silver, to which potash or soda is added, and which deposits silver at ordinary temperatures when brought in contact with an aqueous solution of sugar of milk.

Mr. DRAYTON mixes 1 oz. of nitrate of silver, 3 oz. of water, 1 oz. of liquid ammonia, and 3 oz. of spirit of wine, and filters the solution after it has stood 3 or 4 hours. To every ounce of the solution he adds $\frac{1}{2}$ oz. of sugar (*grape* sugar if possible), dissolved in equal quantities of water and alcohol. The surface to be silvered is covered with this liquid at a temperature of 160° F., maintained till the deposition of silver is complete. When quite dry, the coated surface is covered with mastic varnish. Other substances besides sugar occasion the deposition of silver from the ammoniacal solution; as oil of cassia, oil of cloves, and other essential oils, aldehyde, &c. UNGER recommends a strong alcoholic solution of tannin. M. VOHL prefers an alkaline solution of gun cotton. Dissolve gun-cotton in a solution of caustic potash, pour it into a solution of nitrate of silver, and add ammonia sufficient to redissolve the precipitate. The liquor being slowly heated in a water-bath, becomes brown, effervesces, and deposits silver of superior brilliancy.

GLAZES. Common earthenware is glazed with a composition containing lead, on which account it is unfit for many pharmaceutical purposes. The following glaze has been proposed, among others, as a substitute: 100 parts of washed sand, 80 of purified potash, 10 of nitre, and 20 of slaked lime; all well mixed and heated in a black-lead crucible, in a reverberatory furnace, till the mass flows into a clear glass. It is then to be reduced to powder. The goods to be slightly burnt, placed under water, and sprinkled with the powder.

GLAZE FOR PORCELAIN. Feldspar 27 parts, borax 18, Lynn sand 4, nitre 3, soda 3, Cornwall china clay 3 parts. Melt

together to form a frit, and reduce it to a powder, with 3 parts of calcined borax.—ROSE.

GLUE is made by boiling parings of ox hides and other skins in water, evaporating the solution to a due consistence, allowing it to gelatinize in wooden boxes, cutting it into layers with a wire, and drying them on nets stretched on wooden frames. *Bones* also yield a pale glue, described under GELATINE. Bank-note glue, or mouth glue, is made by dissolving 1 lb of fine glue, or gelatine, in water, evaporating it till most of the water is expelled, adding ½ lb brown sugar, and pouring it into moulds. Some add a little lemon-juice. It is also made with two parts of dextrine, 2 of water, and 1 of spirit.

GLUE, LIQUID. 1. Dissolve bruised orange shellac in ¾ths of its weight of rectified spirit, or of rectified wood naphtha, by a gentle heat. It is very useful as a general cement and substitute for glue. 2. Another kind may be made by dissolving 1 oz. of borax in 12 oz. of soft water, adding 2 oz. of bruised shellac, and boiling till dissolved, stirring it constantly. 3. Dissolve 1000 parts of glue in 1000 parts by weight of water in a glazed pot over a gentle fire. When it is melted, add nitric acid (sp. gr. 1.32) 200 parts pouring it in very gradually. An effervescence is caused by the escape of hyponitrous acid. When all the acid is added, allow the mixture to cool. (This glue is found to remain unaltered on exposure to the air. It is applied cold, and is recommended as very convenient in chemical operations.)—M. DUMOULIN.

GLUE, MARINE. Cut caoutchouc into small pieces, and dissolve it, by heat and agitation, in cold naphtha. Add to this solution powdered shellac, and heat the whole with constant stirring, until combination takes place, then pour it while hot on metal plates, to form sheets. When used it must be heated to 248° F., and applied with a brush.

GLUE OF CASEINE. 1. BRACONNOT. Dissolve caseine in a strong solution of bicarbonate of soda. 2. WAGNER. Dissolve caseine in a cold saturated solution of borax. Superior to gum, and may take the place of glue in many cases. May be used for the backs of adhesive tickets.

GLUTEN, VEGETABLE. Form wheat flour into a stiff paste with cold water; then knead it under a stream of water till all the starch is washed away. What remains is impure gluten.

GOLDEN COMPOUND. Anhydrous tungstate of soda, or the salt obtained in fusing 2 equivalents of tungstic acid with 1 of carbonate of soda, is to be melted in a porcelain crucible, over a spirit lamp, at a temperature not more than sufficient; then add small pieces of pure tin to the melted mass. Cubes of a golden color instantly form. The process should not be continued too long, or they acquire a purple hue. (See AURUM MUSIVUM.)

GUM, BRITISH. (See DEXTRINE.) It is also prepared by heating starch alone, or previously mixed with an acid. PINEL directs half a gallon of nitric and half a pint of muriatic acid to be mixed with 100 gallons of water, and as much potato fecula added as will form a paste. In 2 hours remove the paste in buckets, prepared for the purpose, to drain off all the water. Then place the paste in small lumps in a drying-room till dry; pulverize it, and expose the powder the first day to the temperature of 100°, the next day raise it to 150°, on the third day to 190°. It is then powdered, sifted, and heated from 300° to 350°. To give it the appearance of gum, after it has gone through the stove, and is powdered and sifted, mix it to a paste with water to which 1 per cent. of nitric acid has been added, spread it on copper plates in layers ¾ of an inch thick, and heat it in an oven from 240° to 300° then remove it to the open air to cool.

GUM ARABIC, PURIFICATION OF. PICCIOTTO'S process. The gum is dissolved in water, and sulphurous acid gas passed into it. The sulphurous acid is sufficiently removed for common purposes by gently boiling the mucilage in a retort with a receiver attached. But to obtain the gum in a purer state, carbonate of barytes is added, the mixture is filtered, afterwards agitated with gelatinous alumina, again filtered, and evaporated.

GUN BARRELS, TO STAIN. (See BROWNING LIQUIDS, page 323.)

GUN COTTON. Mix 1½ fluid oz. of each of the strongest nitric and sulphuric acids; put the mixture in a Wedgewood mortar, and when cool, add 100 gr. of cotton-wool.

Stir it with a glass rod, and when it is fully soaked, squeeze out the acid with the pestle or a porcelain spoon, throw the cotton into a large quantity of water, squeeze it again, and wash it under a stream of water till quite free from acid. In the specification, the patentee directs 1 measure of nitric acid, sp. gr. 1.45 or 1.50, to be mixed with 3 measures of sulphuric acid, sp. gr. 1.85; the cotton to be soaked in the acid, then squeezed from it and left in a covered vessel for an hour; and after washing, to be dipped in a solution of 1 oz. carbonate of potash in a gallon of water, then pressed, and partially dried; again dipped in a weak solution of nitre, then dried in a room heated to 150°. This destructive compound has already fallen into disuse as to the purposes for which it was introduced; but on the other hand, is usefully applied to other objects. See COLLODION.

GUNPOWDER. A compound of nitre, charcoal, and sulphur. The nitre should be purified by recrystallization, the sulphur by distillation, and the charcoal selected of the best quality —that of the dogwood, alder, poplar, chestnut, or willow is preferred. The following is the composition of some of the most approved kinds:

	Nitre.	Charcoal.	Sulphur.
Royal Mills, Waltham Abbey,	75	15	10
Marsh's Sporting,	76	15	9
——— Mining,	65	15	20
French (Government),	75	12.5	12.5
——— Sporting,	78	12	10
Chinese,	75	14.14	9.9

GUNPOWDER, WHITE. Well dried ferro-prussiate of potash 1 part, white sugar 1 part, chlorate of potash 2 parts. Let the ingredients be separately reduced to a fine powder, and the powders mixed by the hand, or by means of a leathern barrel turning on its axis. Or they may be moistened with water, and granulated by passing the paste through a wire sieve.

GYPSUM, TO HARDEN. KEATING'S patent process is to moisten calcined gypsum with a solution of 1 lb of borax, 1 lb of tartar, in 11 lb of water; it is then heated to redness for 6 hours, and pulverized. ERDEMANN recommends plaster figures, &c., to be soaked in a solution of FUCHS'S soluble glass.

HAHNEMANN'S WINE TEST. See TESTS.

HARNESS JET. Take 4 oz. best glue, 1½ pint good vinegar, 2 oz. best gum arabic, ½ pint good black ink, 2 dr. best isinglass. Dissolve the gum in the ink, and melt the isinglass in another vessel in as much hot water as will cover it. Having first steeped the glue in the vinegar until soft, dissolve it completely by the aid of heat, stirring to prevent burning. The heat should not exceed 180°. Add the ink and gum, and allow the mixture again to rise to the same temperature. Lastly, mix in the solution of isinglass, and remove from the fire. When used, a small portion must be heated until fluid, and then applied with a sponge, and allowed to dry on.

HARNESS, WATERPROOF PASTE FOR. Put into a pipkin black resin 2 oz., place on a gentle fire, and when melted add beeswax 3 oz. When this is melted, remove from the fire, and stir in ½ oz. fine lamp-black and ½ dr. Prussian blue, finely powdered. When completely mixed, add spirits of turpentine to form a thin paste, and let it cool. To be applied like blacking.

HEADING FOR BEER. Equal parts of alum and sulphate of iron.—GRAY.

INCENSE. Olibanum 2 parts, benzoin 1 part.

INDIA-RUBBER COURT PLASTER. A stout frame of wood must be made, about 3 yards long and about 1¼ yard wide. Within this frame must be placed two sides of another frame, running longitudinally and across, so fixed in the outer frame that the two pieces may slide independently of each other backwards and forwards about 6 inches. Tapes of canvas must be tacked round the inside of the inner frame and the corresponding sides of the outer frame, so as to form a square for the material to be sewn in, which, when done, the two loop frames must be drawn tightly to the outer by means of a twine passed round each, in order to stretch perfectly free from irregularities the silk or satin previous to laying on the composition.

To make the Plaster. Dissolve India-rubber in naphtha or naphtha and turpentine, lay it on with a flat brush on the opposite side to that which is intended for the plaster. When the silk is perfectly dry, and the smell in a great measure dissipated, it will be ready for the adhesive material; to make which, take equal parts of Salisbury or fine Russia glue and the best isinglass, dissolve in a sufficient

quantity of water over a water-bath, and lay on with a flat hogtool while warm. It is requisite to use great caution to spread the plaster evenly and in one direction, and a sufficient number of coatings must be given to form a smooth surface, through which the texture of the fabric is not perceptible. Each coating should be perfectly dry before the succeeding one is given, after which the frame is to be placed in a situation free from dust, and where a draught of air would facilitate the drying. The quantity of water used and the weight of the two materials must be a little varied according to the season and the gelatine strength they possess. Lastly, the plaster being ready to receive the polishing coat, which gives also the balsamic effect to it, a preparation is made in nearly the same manner as the Tinct. Benz. Co. of the P. L., with the addition of more gums. This preparation must be laid on once only, and with a brush kept for the purpose. For making plasters on colored silk it is only necessary to select the silk a shade deeper than the color required, as the plaster causes it to appear a little lighter.

INDIGO. The principal preparations of indigo are described under CHEMIC BLUE, and DYES Indigo may be purified by several methods, of which the following is the most simple: Mix indigo with half its weight of Paris plaster and sufficient water to form a thin paste. Spread this evenly on an iron plate, about two inches wide, to the depth of one-eighth of an inch, and let it dry in the air. Then apply the flame of a large spirit lamp to the under side of the plate, beginning at one end and advancing it to the other as the sublimation proceeds. The violet vapor condenses on the surface in brilliant prisms or plates. Good indigo yields from 15 to 17 per cent.—Mr. T. TAYLOR.

Purified indigo is also obtained from the alkaline solution of reduced indigo, described under DYES; or by dissolving indigo in a mixture of 1 part of caustic soda, 1 of grape sugar, and 20 of water. To the clear solutions thus obtained add muriatic acid to throw down the indigo, wash this perfectly with pure water, and finally with alcohol. If care be taken to exclude the air before and after adding the acid, and to wash it with recently boiled water, to drain it rapidly and dry it *in vacuo*, the indigo is obtained nearly white, but becomes blue on exposure to the air.

INKS. The following are specimens of the most useful kinds:

Black Writing Inks. 1. BRANDE'S. Bruised Aleppo galls 6 oz., soft water 6 pints; boil together, add 4 oz. of sulphate of iron and 4 oz., of gum arabic. Put the whole in a bottle, and keep it in a warm place, shaking it occasionally. In 2 months pour it off into glass bottles; and add to each pint a grain of corrosive sublimate, or 3 or 4 drops of creasote.

2. Dr. WOLLASTON'S. Galls 1 oz., gum ½ oz., cloves ½ dr., sulphate of iron ½ oz., water 8 oz. Digest with frequent shaking, till it has sufficient color. A good durable ink, and will bear diluting.

3. *Prerogative Court Ink.* Galls 16 oz., gum 6 oz., alum 2 oz., sulphate of iron 7 oz., kino 3 oz., logwood in powder 4 oz., water 8 lb.—GRAY.

4. Dr. URE'S *Ink.* For 12 gallons of ink take 12 lb of bruised galls, 5 lb of gum Senegal, 5 lb of green sulphate of iron, and 12 gallons of rain water. Boil the galls in a copper with 9 gallons of water for 3 hours, adding fresh water to replace what is lost by evaporation. Let the decoction settle, and draw off the clear liquor; add to it a strained solution of the gum; dissolve also the sulphate of iron separately, and mix the whole. Instead of boiling the galls, they may be macerated in a portion of hot water for 12 hours, then put into a percolator, and the rest of the water passed through it.

5. *Anticorrosive Ink.* Aleppo galls 10 lb, logwood 5 lb, pomegranate-peel 2½ lb, cloves 2½ oz., soft water 8 gallons. Let the whole boil gently for an hour or two, then cover the copper and leave it for 12 or 14 hours, stirring it now and then. Strain off the decoction and add 2 gallons more water to the ingredients; simmer gently for an hour, and strain. Mix the liquors, and let them settle; draw off the clear liquid from the dregs; dissolve in a portion of it 2½ lb of gum arabic and ½ lb of sugar candy; and in another portion 2½ lb of green sulphate of iron. Strain both solutions, and mix the whole together; then add 1 oz. of calcined borax, and ¼ oz. of creasote dissolved in ¼ pint of spirit of wine. (Dr. HARE recommends an ink free from acid, to be made with galls and "finery-cinder;" but we are not sure what is intended by this name).

6. RIBAUCOURT'S *Ink.* Galls 1 lb, logwood ½ lb, gum 6

oz., sulphate of iron ½ lb, sulphate of copper 2 oz., sugar 2 oz., water 12 lb (or 5 quarts). This has the disadvantage of corroding the steel pens and the penknives with which it comes in contact.

7. Galls 3 oz., sulphate of iron 1 oz., logwood ½ oz., gum ½ oz., ale a quart. Let it stand in a loosely-corked bottle, in a warm place, for a week or more, shaking it daily.

8. Boil 4 oz. of logwood for an hour in 6 quarts of water; adding boiling water as it evaporates: then add 16 oz. of blue galls coarsely bruised, 4 oz. of dried sulphate of iron (*i. e.*, heated till it becomes whitish and pulverulent), 3 oz. of brown sugar, 6 oz. of gum arabic, and ¼ oz. of acetate of copper ground with a little of the decoction. Keep the whole in a bottle uncorked for a fortnight, shaking it twice a day.

9. Reade's *Patent.* This differs from common black ink, in containing a portion of soluble Prussian blue.

10. *Chrome Ink.* Extract of logwood ½ oz., gum ¼ oz., water a pint. Dissolve also in 12 oz. of water, ½ oz. of yellow chromate of potash (or ¼ oz., each of bichromate and bicarbonate of potash); and mix the two solutions. The ink is ready for immediate use.

11. Dr. Lewis's *Writing Ink.* Powdered sulphate of iron 1 oz., powdered logwood 1 oz., powdered galls 3 oz., gum arabic 1 oz., white wine or vinegar 1 quart.

12. Berzelius recommends a solution of vanadiate of ammonia in infusion of galls. Dr. Ure states that this forms the most perfect ink that can be desired; but the scarcity and high price of the vanadiate prevent its use.

13. Runge's *Black Writing Fluid.* Boil logwood 22 lb, in enough water to yield 14 gallons of decoction. To 1000 parts of this decoction, when cold, add 1 part of chromate of potash. The mixture is to be well stirred. The proportions are to be carefully observed, and the yellow chromate, not the bichromate, employed. (This ink is said to possess some great advantages; to adhere strongly to paper, so that it can neither be washed off by water, nor even altered by weak acids; to form no deposit; and not to be in the least acted upon by steel pens.) Steel pens should be washed in an alkaline solution before being used with this ink.

14. *Alizarine Ink.* Leonhardi. Digest 24 parts Aleppo

galls with 3 parts of Dutch madder and 120 warm water. Filter. Mix 1.2 parts solution of indigo, 5.2 of sulphate of iron, and 2 parts crude acetate of iron solution. This ink contains no gum, cannot get mouldy; the tannate of iron is prevented from separating by the sulphate of indigo. Alizarine ink may be evaporated to dryness and formed into cakes; 1 part with 6 of hot water will then form an excellent writing fluid.

Packers' Marking Ink is merely the dregs of black ink, for marking parcels with a brush.

Copying Ink. Mr. BRANDE directs 1 oz. of brown sugar to be added to No. 1, for copying. Another kind is made by dissolving ½ oz. of gum, and 20 gr. of Spanish liquorice in 13 dr. of water, and adding to it a drachm of lamp-black, previously mixed with a teaspoonful of sherry. If the lamp-black is greasy, it should be heated to redness in a covered crucible. Another published form is: Black ink 3 oz., sugar candy 1 oz.

Ink Powder. This consists of the dry ingredients for ink, powdered and mixed. 1. Powdered galls 4 oz., sulphate of iron (heated till it becomes white and pulverulent 1 oz., powdered gum 1 oz., white sugar ½ oz.; mix.) To make a quart of ink with water or beer.

2. Powdered galls 2 lb, green vitriol 1 lb, powdered gum 8 oz. In 2-oz. packets, each for 1 pint of ink.—GRAY.

Red Writing Inks. 1. Best ground Brazil wood 4 oz., diluted acetic acid a pint, alum ½ oz. Boil them slowly in a covered tinned copper or enamelled saucepan, for an hour; strain, and add ½ oz. gum. Some direct the Brazil wood to infuse for 2 or 3 days before boiling.

2. WEBER'S *Red Ink.* Boil 4 oz. of Pernambuca wood with 16 oz. of dilute acetic acid and an equal quantity of water, until 24 oz. remain. Add an ounce of alum, and evaporate again to 16 oz.; add gum arabic 1 oz., strain: and to the cold liquid add, lastly, 1 dr. of protochloride of tin. (Said to be of a finer color, and more permanent, than cochineal ink.)

3. Boil 2 oz. of good Brazil wood, ½ oz. of alum, and ½ oz. of cream of tartar, in 16 oz. of rain water, till reduced to half; strain, and dissolve in it ½ oz. of gum arabic, and

add a tincture made with 1½ dr. of cochineal in 1½ oz. of spirit of wine.—HENSELER.

4. Brazil 4 oz., alum 2 oz., water a quart. Boil for an hour, and strain; then add 1 oz. of gum.

5. Triturate 1 dr. of cochineal and 1 dr. of subcarbonate of potash, with a little boiling water; then add 1 dr. of burnt alum and 2 dr. of cream of tartar, and water to bring it to the desired color. (Not so permanent as the Brazil ink.)

6. *Carmine Ink.* Heat a scruple of carmine with 3 oz. of water of ammonia for some minutes, a little below boiling, and add 15 or 20 gr. of gum. (The inkstand must be kept well closed.)

7. STEPHEN'S *Red Ink.* Patent. Take some common soda, potash, or carbonate of ammonia, and add to it, at intervals, twice its weight of crude argol in powder. When effervescence has ceased, pour off the solution, or filter it from insoluble matter. Add to it next, by measure, half the quantity of oxalate of alumina, prepared by adding to precipitated alumina in a damp state as much oxalic acid as will dissolve it. Into this mixture, when cold, put as much powdered cochineal as will give it a fine red color, and, after letting it stand for 48 hours, strain it for use.

Blue Inks. 1. READE'S Patent. Prepare a solution of iodide of iron, from iodine, iron, and water; add to the solution half as much iodine as first used. Pour this solution into a semi-saturated solution of ferroprussiate of potash, containing nearly as much of the salt as the whole weight of iodine. Collect the precipitate, wash it, and finally dissolve it in water, to form the blue ink. The solution from which the precipitate is separated, evaporated to dryness, and the residue fused, re-dissolved, and crystallized, yields pure iodide of potassium. (This process being patented, ink must not be prepared by it for sale.)

2. Add a pint of a cold solution of persulphate of iron (prepared as directed, page 385) to a solution of 6 oz. of ferrocyanide of potassium in 2 pints of water. Collect the precipitate, wash it with distilled water until it begins to dissolve, then triturate it in a mortar with sufficient distilled water to form a blue ink.

3. Chemic, or Saxon blue (sulphate of indigo), diluted with water to the desired shade, with a little gum.

4. Pure Prussian blue, triturated with a sixth part of its weight of oxalic acid, with a little water, to a smooth paste, and more water added to bring it to the proper color. A larger proportion of the acid is ordered in some recipes.

5. Dr. NORMANDY'S *Blue Ink.* Chinese blue (ferrocyanide of iron) is ground in water with binoxalate of potash and gum arabic, in the following proportions: 7 oz. of water to 3 dr. of Chinese blue, 1 dr. of binoxolate of potash, and 1 dr. of gum arabic.

6. STEPHEN'S *Patent Blue Ink,* Common Prussian blue is first macerated in strong sulphuric acid, then repeatedly washed in water, and afterwards dried. This process is to render it more soluble in oxalic acid, which is now to be gradually added in the proportion of about 1 part to 6 of the Prussian blue (as before maceration), together with sufficient water to yield a dense blue solution.

7. Digest 2 oz. of the cuttings of tin plate with 4 oz. of nitrous acid, and add the solution to a gallon of water in which 2 oz. of Prussiate of potash have been dissolved. (This requires frequent shakings to keep the precipitate—which is Prussian blue—suspended.)

Violet Inks. 1. Boil 8 oz. of logwood in 3 pints of rain or distilled water to 1½ pint. Strain, and add 1½ oz. of clean gum and 2½ oz. of alum in fine powder. Agitate frequently till dissolved.

2. Cudbear 1 oz., pearlash 1½ oz., mucilage 2 oz., soft water to make a pint. Pour the water hot on the cudbear and pearlash, allow the mixture to stand for 12 hours, then strain, and add the mucilage. 1 oz. of rectified spirit may also be added.

Purple Inks. 1. Add a little muriate of tin to a strong decoction of logwood. A little gum may be added.

2. Dr. NORMANDY'S *Purple Ink.* To 12 ℔ of Campeachy wood add as many gallons of boilng water, pour the solution through a funnel, with a strainer made of coarse flannel, on 1 ℔ of hydrate or acetate of deutoxide of copper finely powdered (having at the bottom of the funnel a piece of sponge); then add immediately 14 ℔ of alum, and for every 340 gallons of liquid add 80 ℔ of gum arabic or gum Senegal. Let these remain for 3 or 4 days, and a beautiful purple color will be produced.

Brown Ink. 1. Boil ½ oz. of catechu with 8 oz. of water until

dissolved, and strain. Dissolve 60 gr. of bichromate of potash in 1½ oz. of water, and add it gradually to the solution of catechu until the desired shade is obtained. It requires no gum.

2. By adding to the violet ink finely-powdered bichromate of potash, in the proportion of from 15 to 30 gr. to an oz., various shades of brown and snuff color are obtained.

Yellow Ink. 1. Gamboge triturated with water, and a little alum added.

2. Boil 8 oz. of French berries with 1 oz. of alum in 1 quart of water; strain, and add 1 oz. of gum.

Green Ink. 1. Dissolve 3 dr. of bichromate of potash in 1 oz. of water; add to the hot solution ½ oz. of alcohol, and decompose the mixture by a little strong sulphuric acid, till it assumes a brown color. Evaporate the liquid to half, let it cool, dilute with a sufficient quantity of water, and filter; add to the filtered liquid 4 dr. of alcohol, decompose with a few drops of sulphuric acid, and let it rest. After some time it assumes a fine green color. A little gum may be added. There is danger of the paper and steel pens suffering from an excess of sulphuric acid.)—WINCKLER.

2. Distilled verdigris 2 oz., cream of tartar 1 oz., water 8 oz.; boil to half, and filter.—KLAPROTH.

3. Add to the yellow ink No. 2 sufficient sulphate of indigo.

4. Dissolve sap green in water with a little alum.

5. Rub 3½ dr. of Prussian blue and 3 dr. of gamboge with 2 oz. of mucilage, and add ½ pint of water.

Gold and Silver Ink. Fine bronze powder, or gold or silver leaf, ground with a little sulphate of potash, and washed from the salt, is mixed with water and a sufficient quantity of gum.

Indestructible Writing Fluids. The common writing inks being liable to be obliterated by many chemical agents, several compounds more capable of resisting these agents have been proposed, of which the following appear deserving of notice:

1. *Carbon Ink.* Dissolve real Indian Ink in common black ink; or add a small quantity of lamp-black, previously heated to redness, and ground perfectly smooth, with a small portion of the ink.

2. Stephens's (*patent*) *Carbon Ink.* Common soda of commerce is mixed with resinous matters (as shellac or resin), in about equal parts by weight. Water being added according to the strength required, the solution is boiled until the resin has become dissolved. Mix in a mortar with the requisite quantity of fine lamp-black, and add any suitable-colored solution.

3. *Shellac Ink, or* Coathupe's *Writing Fluid.* To 18 oz. of water add 1 oz. of powdered borax, and 2 oz. of bruised shellac, and boil them in a covered vessel, stirring them occasionally, till dissolved. Filter, when cold, through coarse filtering paper; add 1 oz. of mucilage; boil for a few minutes, adding sufficient finely-powdered indigo and lamp-black to color it. Leave the mixture for 2 or 3 hours for the coarser particles to subside; pour it off from the dregs, and bottle it for use.

4. *Gluten Ink.* Dissolve wheat gluten, free from starch, in weak acetic acid of the strength of common vinegar; mix 10 gr. of lamp-black and 2 gr. of indigo with 4 oz. of the solution, and a drop or two of oil of cloves.

5. Hausmann's. Dissolve 1 part of genuine asphaltum with 4 parts of oil of turpentine, and sufficient lamp-black. If sufficient lamp-black be used to give it a suitable consistence, it may be used with types.

6. Braconnet's *Indelible Ink.* Take 20 parts of Dantzic potash, 10 of tanned leather parings, and 5 of sulphur; boil them in an iron pot with sufficient water to dryness; then raise the heat, stirring the matter constantly, till the whole becomes soft, taking care that it does not ignite. Add sufficient water, and filter through cloth. It must be kept from the air. It flows freely from the pen, and resists many chemical agents; but it is not strictly indelible.

7. Dr. Normandy's *Indelible Ink.* Frankfort lamp-black 24 ℔, to be ground with mucilage, made by adding 24 ℔ of gum to 60 gallons of water, and the mixture filtered through a very coarse flannel; 4 ℔ of oxalic acid are then added, with as much decoction of cochineal and sulphate of indigo as will yield the shade of color desired.

8. *Indian Ink.* Real lamp-black, produced by combustion of linseed oil, ground with gum, and infusion of galls. It is prepared both in a liquid and solid form, the latter being dried in the sun.

Ink for Writing on Zinc Labels. Horticultural ink: 1. Dissolve 100 gr. of chloride of platinum in 1 pint of water. A little mucilage and lamp-black may be added.

2. Sal ammoniac 1 dr., verdigris 1 dr., lamp-black ½ dr., water 10 dr.; mix.

Ink for Writing on Steel or Tin Plate, or Sheet Zinc. 1. Mix 1 oz. of powdered sulphate of copper and ½ oz. of powdered sal ammoniac with 2 oz. of diluted acetic acid, adding lamp-black or vermilion.

2. Dissolve 1 part of copper in 10 of nitric acid, and dilute with 10 parts of water.

White Marking for Black Bottles, in cellars. Grind flake-white, or sulphate of barytes, with a little oil of turpentine, and any light-colored varnish, to a proper consistence.

Lithographic Ink. 1. LASTEYRIE'S: Dried soap 1 oz.; melt, and add shellac 5 oz., then common soda 1 oz., mastic 1 oz., and lastly, lamp-black 3 dr. Melt, stir together, and, when completely melted, pour into moulds: to be used as Indian ink.

2. *Autographic.* White soap 100 parts, white wax 100, mutton suet 30, shellac 50, mastic 50, lamp-black 30 or 35. Melted as above.

3. *Lithographic Ink.* Heat 40 parts of yellow wax until its vapor kindles on coming in contact with a burning match; then remove it from the fire, and add gradually, in small parts, Marseilles soap 22 parts, gumlac 28 parts, and mastic 10 parts. Extinguish the flame, and incorporate perfectly with this mixture lamp-black 9 parts. Then again heat until the vapor can be ignited, when remove it from the fire, and after the flame has been extinguished, pour it upon a stone. The mass is then cut into pieces.—M. WEISHAUPT.

4. *Crayons.* White wax 8 oz., white soap 2 oz., shellac 2 oz., lamp-black 3 tablespoonfuls. Melt the wax and soap with a brisk fire; stir in the lamp-black; allow the mixture to burn for half a minute, then extinguish the flame, and add the shellac by degrees, stirring continually. Put the mixture on the fire till it kindles, or nearly so. Extinguish the flame, let the mixture cool a little, and pour it into moulds.

Inks for Marking Linen. Some of these are used with types; others with a clean quill pen.

1. Sulphate of manganese 1 dr., water 1 dr., powdered sugar 2 dr., lamp-black ½ dr. Triturate them together, and stamp it on the linen with types. When dry, wash the part with liquor potassæ; again dry, and wash with plenty of water.

2. Dr. SMELLIE'S. Sulphate of iron 1 dr., linseed oil 1 oz., vermilion ½ oz.; grind perfectly smooth. Printers' ink is also used with type.

3. Heat to redness equal weights of black oxide of manganese and caustic potash, and mix it with an equal weight of pipe-clay, and sufficient water to give it a due consistence. To be applied with types or stencils. It becomes brown, and does not wash out. The following are used with a quill pen:

4. Nitrate of silver 100 gr., distilled water 1 oz., gum arabic 2 dr., sap green 1 scruple; dissolve. The linen is first to be wetted with the following *pounce*, dried and rubbed smooth, then written on by a clean quill or bone pen dipped in the ink. *Pounce* or *Mordant.* Subcarbonate of soda 1 oz., water 8 oz. (A great variety of recipes might be given, slightly differing from the above in the proportion of the ingredients, and in the coloring matter. GRAY directs 2 dr. of nitrate of silver, 6 dr. of water, and 2 of mucilage, and a pounce of 1 oz. of subcarbonate of soda in 16 of water, with a little sap green. Another form is: Nitrate of silver 1 oz., distilled water 5 oz., powdered gum 1¼ oz., sap green sufficient to color it. The linen to be first wetted with the following preparation: Subcarbonate of soda 1 oz., water 6 oz., gum 1 oz., dissolve. Some add a little powdered bole to the preparation; the object in coloring it being merely that the part which has been wetted may be more readily distinguished. The quantity of nitrate of silver should not be much less than 100 gr. in 1 oz. of ink; the proportion of the other ingredients is of less importance. Some direct the addition of a drop or two of nitric acid.)

5. *Italian.* Moisten the linen with a solution of recently prepared muriate of tin, and write with a neutral solution of salt of gold.

Marking Ink, without Preparation. These merely require to have a hot iron passed over the part written on, or to be held pretty near the fire till the writing assumes a dark color.

1. Nitrate of silver 3 dr., water 1½ oz.; dissolve and add as much strong liquid ammonia as will redissolve the precipitate formed by it; add 2 dr. of mucilage, a litte sap green, and water, if required, to make up the measure to 2 oz. A little ivory-black, Indian ink, or indigo, is sometimes used to color it. Some recipes contain nitrate of copper in addition to nitrate of silver. Several recipes might be given, but they will all probably be superseded by Mr. REDWOOD'S. (In operating with ammonia and nitrate of silver, fulminating silver is sometimes unexpectedly formed and may prove a source of danger. Perhaps in this respect, as well as others, Mr. REDWOOD'S preparation claims a preference.)

2. Mr. REDWOOD'S. Rub together 1 oz. nitrate of silver, and 1 oz. of bitartarate of potash; add 4 oz. of liquor ammoniæ, and when dissolved mix in 6 dr. of white sugar, 10 dr. of powdered gum arabic, ½ oz. of archil, and water to make up 6 oz. by measure. (Instead of archil, ¼ oz. of sap green may be used to color the ink; or 40 gr. of fine vegetable black, previously triturated with a little water or mucilage.)

3. Rev. J. B. READE'S patent. This differs from the last in using tartaric acid instead of bitartrate of potash. The quantities may be 1 oz. of nitrate of silver, 3 dr. of tartaric acid, and the above quantities of the other ingredients. The use of tartaric acid he claims an exclusive right to.

4. Add to the last an ammonical solution of an oxide or salt of gold.—READE. (This addition prevents its being acted on by cyanide of potassium, and some other agents which the silver ink fails to resist.)

5. In addition to the above receipts, the following of M. HENRY may deserve attention in large establishments where economy is an object: Take 1 oz. of iron filings and 3 oz. of vinegar, or diluted acetic acid. Mix the filings with half the vinegar, and agitate them continually until the mixture becomes thick, then add the rest of the vinegar and 1 oz. of water. Apply heat to assist the action; and when the iron is dissolved, add 3 oz. of sulphate of iron, and 1 oz. of gum previously dissolved in 4 oz. of water: and mix the whole with a gentle heat. To be used with a brush and stencil plates.

Crimson Marking Ink. Dissolve 1 oz. of nitrate of silver and 1½ oz. of carb. soda in crystals, separately in distilled

water; mix the solutions, collect and wash the precipitate on a filter, introduce the washed precipitate, still moist, into a Wedgewood mortar, and add to it tartaric acid 2 dr. 40 gr., rubbing together till effervescence has ceased; dissolve carmine 6 gr. in Liquor Ammoniæ (.882) 6 oz., and add to it the tartrate of silver, then mix in white sugar 6 dr., and powdered gum arabic 10 dr., and add as much distilled water as will make 6 oz.—PHARM. JOURNAL.

Printing Ink. This is usually made by boiling linseed oil in a large iron pot, setting fire to it, and letting it burn for half an hour or more. Various additions are made to it by some manufacturers, the use of which is not very evident. A viscid varnish is obtained, which is ground with lamp-black, vermilion or other coloring matters, till perfectly smooth. 2½ oz. of lamp-black are sufficient for each pound of varnish. See VARNISHES.

Printers' Ink from Resin Oil. Melt together 13 oz. of resin, 1 ℔ of resin oil, and 1½ oz. of soft soap; when cold, add lamp-black or other coloring matters.

Copper-plate Printing Ink. This is not rendered so viscid as the former, and is colored with Frankfort black.

READE'S *Patent Printing Inks.* The *blue* consists of his soluble Prussian blue (see BLUE WRITING INK, above) ground with oil as above. The *black*, by evaporating his black ink, and mixing the product with oil as usual. The *red* in the same manner, from his patent red ink.

Sympathetic or Secret Inks. The solutions used should be so nearly colorless that the writing cannot be seen till the agent is applied to render it visible.

1. Digest 1 oz. of zaffre, or oxide of cobalt, at a gentle heat, with 4 oz. of nitro-muriatic acid till no more is dissolved, then add 1 oz. of common salt and 16 oz. of water. If this be written with, and the paper held to the fire, the writing becomes green, unless the cobalt should be quite pure, in which case it will be blue. The addition of a little nitrate of iron will then impart the property of becoming green. It is used in chemical landscapes, for the foliage.

2. Put into a vial ½ oz. of distilled water, 1 dr. of bromide of potassium, and 1 dr. of pure sulphate of copper. The solution is nearly colorless, but becomes brown when heated.

3. Boil oxide of cobalt in acetic acid. If a little common salt be added, the writing becomes green when heated; but with nitre it becomes a pale rose-color.

4. A solution of acetate of lead. Colorless, but becomes brown when exposed to sulphuretted hydrogen gas.

5. A weak solution of sulphate of copper. The writing becomes blue when exposed to the vapor of ammonia.

6. A solution of sulphate—or preferably, persulphate—of iron. It becomes black when washed with infusion of galls; *blue*, by prussiate of potash, (This constitutes colorless ink, which becomes visible when written with on paper containing galls, or tannin, or prussiate of potash.)

7. Mix equal quantities of sulphate of copper and sal ammoniac and dissolve in water. It becomes yellow when heated.

8. A weak solution of nitrate of mercury. Becomes black by heat.

9. Rice water, or any solution of starch. It becomes blue when washed over with an alcoholic solution of iodine.

10. Lemon-juice, milk, juice of onions, and some other liquids, becomes black when the writing is held to the fire.

INK, to preserve from mouldiness. Add a small quantity of a solution of creasote in pyroligneous acid or rectified spirit.

INSECTS, to kill. Camphor and pepper, may be used to keep off moths. In Russia the powder of the flowers of a species of Pyrethrum is used as an insecticide. The powder of the root of Acorus calamus is also recommended.

IODATE OF POTASH. Fuse iodide of potassium in a capacious Hessian crucible, remove it from the fire and add to it while still semi-fluid, successive portions of pulverized chlorate of potash, stirring after each addition, till no further action takes place. One part of iodide of potassium will require 1½ of the chlorate. Wash the residuum in warm water, which leaves only iodate of potash.

IODIDE OF POTASSIUM. In addition to the process given in the Pocket Formulary, the following, by M. CRIQUELON,

may deserve attention: Slake 40 parts of lime with sufficient water, and add 24 parts of iron filings; mix, and add 94 parts of iodide by degrees, so as to avoid two violent action, stirring after each addition, and adding water, if necessary, to moderate the action. Triturate the mixture till starch paper ceases to become brown when touched with it, but only shows an ochreous spot. Throw it on a filter, and wash it till the water which passes gives no precipitate with acetate of lead. Treat the liquors with solution of carbonate of potash till it no longer occasions a precipitate. Filter and evaporate the solution. 94 parts of iodine yield 119 parts of iodide of potassium. Another method is that of Mr. READE, described under INK (READE'S BLUE).

IODINE. See Pocket Formulary. Other methods of obtaining it are the following:

To the mother liquor of kelp (after the crystallizable salts have been separated) add sulphuric acid to render the liquor sour. Introduce the acid liquor into a leaden still, heat to 140° F., add binoxide of manganese, and lute on with pipe-clay a leaden head, fitted to a series of spherical glass condensers, each having two mouths opposite each other, and inserted the one into the other. A stopper in the head of the still allows the contents to be occasionally inspected, and additions of acid or oxide made, if necessary. See Dr. PEREIRA'S "Elements" for a drawing of the apparatus. SOUBEIRAN proposes to add sulphate of copper to the lye, which precipitates half the iodine. He then decants the clean liquor, and adds more sulphate of copper with some iron filings. An iodide of copper is formed, which is separated from the iron filings and suspended in the liquor by agitation, collected on a filter, and heated with oxide of manganese and sulphuric acid.

IRON LIQUOR. See DYES, page 340.

IRON, PERSULPHATE OF. Dissolve 16 ounces of sulphate of iron in 4 pints of water, to which have been added 14 fluid drachms of oil of vitriol. Heat to boiling, and add (by small portions, boiling the solution after each addition) 4 ounces of nitric acid, or so much that the solution yields with ammonia a brown precipitate unmixed with black. Continue the boiling for some time, let the solution cool, filter, if necessary, and add water to make up the measure exactly 4

pints. If required in a dry state, evaporate to dryness by a gentle heat.

ISINGLASS. The air-bags or sounds, of several kinds of fishes, washed, dried, and otherwise prepared. They are either dried without opening (purse, pipe, and lump isinglass), or opened and not folded (leaf and honeycomb isinglass), or folded (book isinglass), or twisted into the shape of a lyre or horseshoe (short and long staple). The leaf isinglass is sometimes rolled out into thin plates (ribbon and rolled leaf isinglass). The inner membrane, which is insoluble, is removed from the opened air-bags, in the best kinds. The Russian isinglass, which is most esteemed, is made from the air-bags of several species of ACIPENSER (*sturgeon*); particularly A. HUSO (the *Beluga*); A. GULDENSTADTII (the *Osseter*); A. RUTHENUS (the *Sterlet*); A. STELLATUS (the *Sewruga*); and also from the Silurus glanis (the *Som*), which yields the Samovey isinglass. Brazilian and East India isinglass are of inferior quality; it is not certainly known from what genera or species of fish they are obtained. New York isinglass is the air-bladder of the common hake, macerated in water and rolled out into ribbons. The sounds of the cod yield an inferior kind. Prepared sole skins are used as a cheap substitute for isinglass. See Dr. PEREIRA'S "Elements," for the description of each variety.

IVORY-BLACK. Burn shavings and waste pieces of ivory from the ivory turners, in a covered crucible, till no more smoke issues. Cover it closely while cooling. It should be afterwards washed with diluted muriatic acid, then with water till no longer acid, dried, and again heated in a covered crucible. It is of a deeper color than bone-black, and is used as a pigment, a tooth-powder, and to decolorize syrups and other liquids.

IVORY, FLEXIBLE. The pieces of ivory or bone, already manufactured into the shape required, are to be steeped for some time in dilute muriatic acid, until they have lost their earthy parts so far as to become yellowish, flexible, and elastic. When dry they become again inflexible, but their flexibility may at any time be restored by steeping them in water. In this manner flexible tubes, probes, bougies, &c., may be constructed.

IVORY, TO SATIN. Ivory is stained with the usual dyeing materials; it should be first steeped in the mordant and

afterwards in the hot color. Nitro-muriate of tin is the mordant for red, with decoction of brazil or cochineal; for yellow, with fustic; for violet, with logwood. After being plunged in hot liquor it should be placed in cold water. A black stain is given by nitrate of silver.

Ivory may be *gilded* by immersing it in a fresh solution of proto-sulphate of iron, and afterwards in solution of chloride of gold. It may be *bleached* by solution of sulphurous acid.

JAPAN. See VARNISHES.

JELLIES. See DIETETIC ARTICLES.

KID-GLOVE CLEANER. Add 15 drops of strongest solution of ammonia to spirits of turpentine ½ pint. (Having fitted the gloves on wooden hands, apply this mixture with a brush. Follow up this application with some fine pumice powder. Rub with some flannel or sponge dipped in the mixture. Rub off the sand, and repeat the same process twice or thrice. Hang in the air to dry, and, when dry, place in a drawer with some scent.)

KYAN'S SOLUTION, for preventing the dry rot. Dissolve 1 lb of corrosive sublimate with 5 gallons of water.

LABARRAQUE'S CHLORO-SODAIC LIQUOR is nearly identical with the Liquor Sodæ Chlorinatæ of the London Pharmacopœia. It is made by passing the chlorine gas from 2 oz. black oxide of manganese, and 8 oz. of muriatic acid, into a solution of 15 oz. of crystallized subcarbonate of soda in 3 pints of water; or sufficient to bring it to the density of 12° Baumé, or 1.09 specific gravity.

LABELS, INDESTRUCTIBLE, for acid bottles, &c. Write with dilute sulphuric acid (1 part to 6 of water), dry the label, and expose it to a moderate heat.

LAC, PREPARATIONS OF. Stick-lac consists of twigs of several kinds of trees incrusted with a resinous matter produced by the puncture of an insect (the coccus lacca). This, triturated with water and dried, forms seed-lac. The seed-lac heated and pressed in cotton bags forms shellac. Lac-dye is the coloring matter extracted from stick-lac by water, evaporated to dryness with the addition of earthy matters, and formed into square cakes. Seed-lac and shellac are chiefly used in varnishes, dissolved in rectified spirit or

rectified wood naphtha. The alcoholic solution is rendered paler, so that it may be used for polishing light-colored woods, by digesting it in the sun, or near a fire, for 2 or 3 weeks, with good animal charcoal, and then filtering it through paper in a funnel heated with hot water. Shellac may be bleached by dissolving it in a solution of potash or soda, and passing chlorine into the solution. The precipitated lac is collected and well washed. KASTNER directs 3 parts of carbonate of potash to be dissolved in 24 of water, and 3 of lime added, and the whole digested in a close vessel for 24 hours. The clear liquor is poured off, and boiled with 4 parts of shellac. When cold, dilute with 4 times its bulk of water, and filter; then add chloride of lime, and afterwards diluted muriatic acid.

LACQUERS. See VARNISHES.

LAKE LIQUOR Boil 1 oz. each of cochineal and salt of tartar in 8 oz. of water, then add 1 oz. of cream of tartar, and the same of alum.

LAKES. These consist of vegetable colors in combination with alumina. Alum is usually added to an infusion or decoction of the coloring ingredient, and afterwards potash added, which throws down the coloring matter combined with alumina. Some of the lakes are noticed under PIGMENTS.

LEMON-JUICE, FACTITIOUS. Dissolve 4 oz. of citric acid in 3 pints of water, with 8 drops of essence of lemon, rubbed with the acid, or dissolved in a little spirit. After standing a few days, filter it, and preserve it in well-closed bottles.

LENSES, EXTEMPORANEOUS. Procure a piece of thin platinum wire, and twine it once or twice round a pin's point, so as to form a minute ring with a handle to it. Break up a piece of flint glass into fragments a little larger than mustard seed; place one of these pieces on the ring of wire, and hold it in the point of the flame of a candle or gas-light. The glass will melt and assume a complete lens-like or globular form. Let it cool gradually, and keep it for mounting. Others are to be made in the same manner; and if the operation be carefully conducted, but very few will be imperfect. The smaller the drop melted, the higher in general will be its magnifying power. It may be mounted

by placing it between two pieces of brass which have corresponding circular holes cut in them of such a size as to hold the edge of the lens. They are then to be cemented together.—FRANCIS.

LINSEED OIL, CLARIFIED, FOR VARNISHES.—Heat in a copper boiler 50 gallons of linseed oil to 280° F.; add 2½ lb of calcined white vitriol, and keep the oil at the above temperature for half an hour; then remove it from the fire, and in 24 hours decant the clear oil, which should stand for a few weeks before it is used for varnish.

LINSEED OIL, REFINED. (WILKS'S Patent.) In 236 gallons of oil pour 6 lb of oil of vitriol, and stir them together for 3 hours; then add 6 lb of fullers' earth, well mixed with 14 lb of hot lime, and stir for 3 hours. Put the oil into a copper boiler, with an equal quantity of water, and boil for 3 hours; then extinguish the fire, and when the materials are cold draw off the water, and let the oil stand to settle for a few weeks before using.

LIQUORICE, PURIFIED EXTRACT OF. Italian or Spanish juice may be purified by the following method: Take a sugar-mould, close the vent-hole with a stopper, place inside it some coarse tow, and over this some clean straw, laid crossways in layers of an inch each, then the sticks of liquorice placed upright, and packed closely in the mould with chopped straw cut rather long. When this arrangement is completed to within an inch of the brim, pour water over the liquorice, allow it to remain for 24 hours, then draw it off, and add more. The liquor, on evaporation, yields an extract perfectly soluble in water.

LITMUS. A preparation of some kind of lichen, probably Lecanora tartarea, or Roccella tartarea, or both. The exact mode of preparing it is kept secret. It is imported in small cubical masses, many samples of which Dr. PEREIRA found to contain, besides the coloring matter, tissues of the lichen, indigo, chalk, &c. See TESTS.

LOZENGES. See Trochisci, P. F., and LOZENGES, under PATENT MEDICINES, in this volume.

LUBRICATING COMPOUNDS. See ANTIATTRITION. The French compound termed *Liard* is thus made: Into 50 parts of finest rape oil put 1 part of caoutchouc cut small, and apply heat until it is nearly all dissolved.

MANKETTRICK'S Lubricating Compound consists of caoutchouc (dissolved in spirits of turpentine) 4 lb, common soda 10 lb, glue 1 lb, oil 10 gallons, water 10 gallons. Dissolve the soda and glue in the water by heat, then add the oil, and lastly the caoutchouc, stirring them until perfectly incorporated.

LUCIFERS. See MATCHES, below.

LUMINOUS VIALS. Nearly fill a bottle with olive or almond oil, and heat it in a water-bath. Drop into it small slices of phosphorus so long as it is dissolved. Let the solution cool, and pour off the oil from the undissolved phosphorus into clean dry vials, which should not be quite filled. When uncorked they emit light.

LUTES. See CEMENTS.

MANURES, ARTIFICIAL. These constitute a new and important branch of manufacture; but a few of the more simple and readily prepared kinds are all that can be noticed here.

Powder for Coating Seeds. Fine bone-dust 20 parts, gypsum 1 part. The seeds are steeped in water from the dung-hill, then strewed over with the powder, so that each shall receive a layer of it, and afterwards dried.

Sulphated Bones. See *Bones, Sulphated.* A usual proportion is 33 of sulphuric acid to 1 cwt. of bones.

Saline Mixture, as a top-dressing for potatoes, &c. Equal weights of nitrate of soda and dry sulphate of soda. 1½ cwt. to an acre.

Mr. HUXTABLE'S *Mixture.* Bone-dust 4 cwt., gypsum 4 cwt., salt 2 cwt., ashes 2 quarters, wood ashes 30 bushels.

Another Saline Mixture. Sulphate of ammonia 42 lb, sulphate of lime 56 lb, sulphate of potash 56 lb, carbonate of magnesia 14 lb, salt 56 lb, to 1 acre.

Dr. ANDERSON'S *Manure for Clover.* Sulphate of ammonia 98 lb, gypsum 172 lb, sulphate of potash 174 lb, sulphate of soda 333 lb, sulphate of magnesia 246 lb, sulphuric acid 98 lb, saltpetre 202 lb, common salt 107 lb, muriate of potash 149 lb.

Dr. JOHNSTONE'S *Substitute for Guano.* Bone-dust 7 bushels, sulphate of ammonia 100 lb, wood ashes 20 lb, salt 100 lb, dry sulphate of soda 11 lb.

To promote the Blowing of Flowers. See page 350.

MARBLE, TO CLEAN. Mix soft soap, solution of potash, and slaked lime, to a paste; spread it over the marble, and leave it for a day or two. Then wash it off.

MARBLE, TO STAIN. Make the marble hot, and pour on it the colored liquid, also made hot. The stains usually employed are archil, solution of indigo, solution of verdigris, decoction of Brazil wood, logwood, and sulphate of iron, tincture of dragon's blood, &c. But the most penetrating medium is wax, which may be colored with alkanet, annotto, verdigris, &c.

MARINE GLUE. See GLUE.

MARINE SOAP. See SOAP.

MATCHES FOR INSTANTANEOUS LIGHT. 1. *Chlorate Matches* (*without sulphur*). Chlorate of potash, separately powdered, 6 dr., vermilion 1 dr., lycopodium 1 dr., fine flour 2 dr.; mix carefully the chlorate with the flour and lycopodium, avoiding much friction, then add the vermilion, and mix the whole with a mucilage made with—1 dr. powdered gum arabic, 10 gr. of tragacanth, 2 dr. of flour, and 4 oz. of hot water; mix, add sufficient water to bring it to a proper consistence, and dip it in the wood, previously dipped in a solution of 1 oz. of gum thus, and ½ oz. of camphor, in 6 oz. oil of turpentine.

2. *With Sulphur.* Chlorate of potash 9 gr., sulphur 2 gr., sugar 3 gr., vermilion 1 gr., flour 2 gr., spirit of wine q. s. The chlorate of potash, &c., must be separately reduced to powder, and the whole mixed with as little friction as possible. The wood should be previously prepared as above, or with camphorated spirit. (These are ignited by dipping them in sulphuric acid, and instantly withdrawing them. The acid should be absorbed by asbestos). They are now become obsolete, having given place to—

Lucifer Matches. These contain phosphorus in a finely divided state, to which it is reduced by agitating it in some warm solution of gum or glue, then adding the other ingredients, so as to form a paste, into which the wood or card is dipped. It is said that urine and artificial urea have the property of readily dividing phosphorus when warmed and agitated together. The following are some of the published recipes:

1. Form 6 parts of glue into a smooth jelly, and rub with it 4 parts of phosphorus, at a temperature of 140° or 150°

F.; add 10 parts of nitre, 5 of red ochre, and 2 of fine smalts. The matches are first dipped in melted wax to the depth of one-tenth of an inch, first rubbing their ends on a hot iron plate.

2. *Noiseless Congreves.* Triturate 9 parts of phosphorus with a solution of 16 parts of gum, and add 14 parts of nitre and 16 of vermilion.—Dr. BOETTGER.

3. Glue 6 parts, phosphorus 4, nitre 10, red lead 5, smalts 2; the glue is soaked in water for 24 hours, then liquefied in a warm mortar, and the phosphorus added, taking care that the temperature is not above 167° F.

4. Glue 21, phosphorus 17, nitre 38, red lead 24; proceed as before.

Promethean Matches. These consist of a composition similar to that of the chlorate matches, inclosed at the end of a paper spill, with a minute glass bulb filled with oil of vitriol in the centre of the composition. When struck, the vessel of acid is broken, and kindles the match.

MICROSCOPE, FORMULÆ FOR THE.

Reagents. Dr. BEALE'S list:

1. Alcohol, of various strengths.
2. Ether, to dissolve oil-globules.
3. Nitric acid, 1 part of strong acid to 5 of water.
4. Sulphuric acid, 1 to 5.
5. Hydrochloric acid.
6. Acetic acid, glacial, and dilute (1 to 5).
7. Chromatic acid, very dilute,—to harden tissues.
8. Solution of potash, saturated, and dilute (1 to 10).
9. Solution of soda, 25 gr. of fused soda in 1 oz.
10. Ammonia (1 part of the strongest solution to 3 of water).
11. Nitrate of barytes, a cold saturated solution.
12. Nitrate of silver (120 gr. in 2 oz.) These two are tests for the mineral acids.
13. Oxalate of ammonia, in solution. Test for lime.
14. Solution of iodine saturated, *i. e.*, 1 to 7000 water; also of 1 gr. of iodine, 3 of iodide of potassium, in 1 oz.

Cements. 1. *Brunswick Black.* Boil together ¼ ℔ of foreign asphaltum and 4¼ oz. of linseed oil (previously thickened with litharge), then mix to a proper consistence with oil of turpentine (about 1 pint).

2. *Gold Size.* Boil 25 parts of linseed oil with 1 of minium and ⅓d part of umber for 3 hours; pour off the

clear fluid, and mix with equal parts of powdered white lead and yellow ochre, added in small successive portions. Then boil well the whole again, and pour off the clear fluid. It dries slowly but firmly. Both this and the last are dissolved by turpentine.

3. *Goadby's Marine Glue.* Dissolve separately in coal naphtha equal parts of shellac and India-rubber. Mix thoroughly with heat.

4. *Sealing-wax Varnish.* Dissolve the best sealing-wax in enough strong spirit of wine to reduce it to the proper consistence. This is brittle.

5. *Canada Balsam.* This dries spontaneously.

Solutions of shellac, gum, and various other cements, and glues (which see), are employed by microscopic manipulators.

Preservative Fluids. *Canada Balsam*, *spirit and water*, *glycerine solution of gelatine*, saturated solutions of *alum*, *chloride of zinc*, and *chloride of calcium*, are all used to preserve microscopic objects. The following formulæ will be found useful:

1. *Goadby's Solution.* Bay salt 4 oz., alum 2 oz., corrosive sublimate 4 gr., boiling water 4 pints: mix and filter. It may often be more diluted.

2. *Thwaites' Fluid.* Mix spirit of wine 1 oz., with creasote sufficient to saturate it; rub up with chalk to form a thin paste, and mix gradually with water 16 oz. To this may be added an equal quantity of water saturated with camphor.

3. *Simple Creasote Solution.* Dissolve creasote 1 dr. in pyroligneous acid 1 dr., and mix gradually with cold water 1 pint.

4. *Passini's Solution.* For blood-globules, nerves, and white tissues generally. Chloride of mercury 1 part, chloride of sodium, 2 parts, glycerine 13 parts, distilled water 113 parts.

Milk, Preserved (Bethel's Patent). The milk or cream is first scalded, and when cold, strongly charged with carbonic acid gas, by means of a soda-water machine. (Attempts have also been made to preserve milk by evaporating it to dryness; but it is necessary to remove the cream in order to effect it.)

Mineral, Chameleon. See Chameleon Mineral.

MODELLING, CLAY FOR. Knead up clay to the proper consistence with glycerine.—BARRESWIL.

MOIREE METALLIQUE. A method of ornamenting the surface of tin plate by acids. The plates are washed with an alkaline solution, then in water, heated, and sponged or sprinkled with the acid solution. The appearance varies with the degree of heat and the nature and strength of the acids employed. The plates, after the application of the acids, are plunged into water, slightly assidulated, dried, and covered with white or colored varnishes. The following are some of the acid mixtures used: Nitro-muriatic acid, in different degrees of dilution; sulphuric acid, with 5 parts of water; 1 part of sulphuric acid, 2 of muriatic acid, and 8 of water; a strong solution of nitric acid; 1 part nitric acid, 2 sulphuric, and 18 of water. Solution of potash is also used.

MORDANTS. See DYES, p. 340.

MULTUM. A name given to a compound of liquorice and quassia, improperly sold by druggists to brewers.

NITRATE OF BARYTES. This may be made from the carbonate by dissolving it in dilute nitric acid, evaporating, and crystallizing; but more cheaply from the sulphate of barytes, by converting it into a soluble sulphuret by heating it with charcoal, and decomposing the filtered solution with nitric acid. M. WEISS recommends mixing the pulverized sulphate of barytes (cawk or heavy spar) with one-eighth of charcoal and one-fourth of flour, heating it in a covered crucible, pulverizing the product and forming it into balls, with one-eighth of charcoal and a little water, and again heating them placed between layers of charcoal. Hot-water extracts the sulphuret, which crystallizes from the filtered solution. By decomposing this by nitric acid (avoiding the gas which escapes) the nitrate is obtained. The other salts of barytes are obtained in a similar manner.

NITRATE OF SILVER. See ARGENTI NITRAS, P. F. It may be prepared from impure silver by the following process: Dissolve it in nitric acid, add common salt till no more silver remains in solution. Wash the precipitate thoroughly; then add water and a very little hydrochloric acid, and introduce some pieces of zinc; let them remain together 24 hours, stirring frequently. Remove the zinc, and wash the re-

duced silver thoroughly. Again dissolve it in nitric acid, diluted with 2 or 3 parts of water; filter, and evaporate, that it may crystallize.

NITRATE OF STRONTIAN. This may be obtained from the native carbonate of strontian, or more cheaply from the native sulphate, by the processes employed for Nitrate of Barytes.

NITRIC AND NITRO-MURIATIC ACIDS. See ACIDS.

NITRITE OF POTASH. It is obtained mixed with a little nitre and potash by heating nitre to redness. To purify the residuum, dissolve it in boiling water, set aside for 24 hours, pour off the liquid from the deposited nitre, neutralize the free alkali with acetic acid, and add twice its volume of alcohol. In a few hours more nitre crystallizes, and the liquid separates into two layers; the upper is alcoholic solution of acetate of potash, the lower is solution of nitrate of potash, which may be evaporated to dryness, or kept in solution. *Used as a test for iodine, with starch paste and hydrochloric acid.* CORENWINDER passes nitrous acid gas, formed by acting on 1 part of starch with 10 of nitric acid, through a solution of caustic potash, sp. gr. 1.38, until it becomes acid; then adds a little caustic potash, so as to render it distinctly alkaline.

NITRO-PRUSSIDE OF SODIUM. To 213 parts of powdered ferroprussiate of potash, in a porcelain basin, add 450 parts of nitric acid of 1.42 density (or 337½ parts at 1.50), adding all the acid at once. When dissolved, transfer to a bolt-head, and digest in a water-bath until the solution precipitates salts of protoxide of iron of a slate color. Neutralize, when cold, with a cold solution of carbonate of soda; then boil, and separate the precipitate by filtration. Evaporate the liquid again, filter, and allow the nitrates of potash and soda to crystallize out. Evaporate the liquid again, and remove the prismatic crystals of nitro-prusside as they form. They may be dissolved in water and recrystallized by cooling.

NOVARGENT. This is said to consist of a solution of fresh precipitate chloride of silver in hyposulphite of soda (or, according to the Pharmaceutical Journal, of oxide of silver in cyanide of potassium), mixed with prepared chalk.

OILS, PURIFICATION AND BLEACHNG OF. Fish and other fat

oils are improved in smell and color, by passing hot air or steam through them. DUNN'S method is to heat the oil by steam to 170° or 200°, and force a current of air through it, under a chimney, till it is bleached and purified. Mr. CAMERON'S method of bleaching palm oil is to keep it at 230° with continued agitation by passing into it high-pressure steam, through leaden pipes of two inches diameter. Four tons of oil require 10 hours' steaming. Palm oil is also bleached by chloride of lime. Take from 7 to 14 lb of chloride of lime, triturate it in a mortar, adding gradually 12 times the quantity of water, so as to form a smooth cream. Liquefy 112 lb of palm oil, remove it from the fire, add the solution of chloride of lime, and stir well with a wooden stirrer. Allow it to cool, and when become solid, break into small fragments, and expose it to the air for 2 or 3 weeks. Then put it into a cast-iron boiler lined with lead, and add sulphuric acid in equal weight to the chloride of lime, diluted with 20 parts of water. Boil with a moderate heat till the oil drops clear from the stirrer; then let it cool.

To remove the fœtor from fish oils, treat them in the same way (except the exposing to the air), using only 1 lb chloride of lime to 112 lb of oil. It does not remove the natural smell of the oil.

Fresh-burnt animal charcoal has some power in improving the color and smell of most kinds of oil; but its effects are limited.

Calcined magnesia has been used to deprive oils of their rancidity.

Mr. GRISELER finds that the addition of a few drops of nitric ether will prevent oils from becoming rancid.

Mr. WATT'S patented method of bleaching oil is by chromic acid. For palm oil it is thus used: The oil is heated in a steam vessel, allowed to settle and cool down to 130° F., then removed into wooden vessels, taking care that no water or sediment accompany it. For a ton of palm oil make a saturated solution of 25 lb of bichromate of potash; add 8 lb of sulphuric acid, and 50 lb muriatic acid (or an equivalent quantity of salt and sulphuric acid.) Put the muriatic acid into the oil, and let it be constantly stirred till it becomes of a light-green color. If not sufficiently discolored, add more of the mixture. Let the oil settle for half an hour, then pump it into a wooden vat, boil it for a few minutes with fresh water by means of a steam-pipe, and

let it settle. For linseed, rape, and mustard oil, a dilute solution of chromic acid is used, with a little muriatic acid; for olive, almond, and castor oil, no muriatic acid is required. Fish oils and fats are first boiled in a steam apparatus with a weak soda lye (½ lb of soda for every ton of fat) for half an hour; then ½ lb sulphuric acid, diluted with 3 lb of water, is added; the whole boiled for 15 minutes, and allowed to settle for an hour or more, when the water and sediment are drawn off, and the oil further bleached by a solution of 4 lb of bicarbonate of potash and 2 lb of sulphuric acid properly diluted.

Mr. DAVIDSON treats whale oil first with a solution of tan, next with water and chloride of lime, and lastly with diluted sulphuric acid and warm water. Rape and other seed oils are also refined by means of sulphuric acid and twice as much water. Mr. GRAY directs 2 lb of oil of vitriol to 112 lb of oil. The oil should be carefully washed from the acid, and filtered.

Mr. BANCROFT'S process for refining common olive oil, lard oil, &c., for lubricating purposes, is to agitate them with from 3½ to 8 per cent. of caustic soda lye, of 1.2 specific gravity. If on trial a small quantity of the lye be found to settle clear at the bottom, enough has been added. The oil is allowed to rest for 24 hours, for the soapy matter to subside; the supernatant oil is then filtered.

Another plan of purifying oils (especially lamp oils) is to agitate them with a strong solution of common salt.

The above methods of treating oil are of doubtful propriety in reference to such as are to be used as medicines. Oils which have been so carefully prepared from sound and fresh materials as to require no purification should be selected for this purpose. This is especially important in reference to cod-liver oil. See LINSEED OIL.

OIL FOR MACHINERY. Sperm oil, palm oil, and olive oil, are used. Care should be taken that they are not adulterated. For compound lubricants see *Anti-attrition*, and LUBRICATING COMPOUNDS.

OLEINE. This may be prepared by boiling fine olive oil with absolute alcohol, and evaporating the solution.

Oxygenated Water, or Deutoxide of Hydrogen. Thenard's oxygenated water is thus made: Expose fragments of perfectly pure barytes to a current of oxygen gas, in a green glass tube heated to a dull redness, to form a deutoxide of barium. To 7 oz. of water add as much pure muriatic acid as will dissolve 4 dr. of barytes; add to this, by degrees, 3 dr. of pulverized deutoxide of barium, and when this is dissolved, add sulphuric acid, drop by drop, till the barytes falls down in the state of sulphate. Then add more deutoxide, and precipitate by sulphuric acid as before. Then filter the solution; and repeat the solution and precipitation several times, till about 3 oz. of deutoxide of barium are used, filtering the liquid after every second repetition. Sulphate of silver is then added to remove the hydrochloric acid, and afterwards pure barytes, to throw down the sulphuric acid, and a few drops of diluted sulphuric acid to remove any excess of barytes.

This energetic compound must not be confounded with the oxygen water formed by impregnating water with oxygen gas; nor with the oxygenous aerated water of Searle, which is water strongly charged with protoxide of nitrogen.

Oxygen Gas. See Gases.

Paper, Copying. Mix lard with black lead or lamp-black, into a stiff paste, rub it over writing paper with a flannel, and wipe off the superfluous quantity with a soft rag. These sheets alternated with writing paper and written on with a solid pen, produce 2 or 3 copies of a letter at once.

Lithographic Paper. Give the paper 3 coats of thin size, 1 of starch, and 1 of solution of gamboge. Each to be applied with a sponge, and allowed to dry before the next is applied.

Hydrographic Paper. This name has been given to paper which may be written on with water. It may be made by rubbing paper over with a mixture of finely powdered galls and sulphate of iron heated till it becomes white. The powder may be pressed into the paper by passing it between rollers or passing a heavy iron over it. A mixture of dried sulphate of iron and ferro-prussiate of potash may be used for blue writing. Or the paper may be imbued with a

strong solution of one ingredient, thoroughly dried; and the other applied in powder. Paper which has been wet with a solution of ferro-prussiate of potash also serves for writing on with a colorless solution of persulphate of iron.

Iridescent Paper. Nutgalls 8 parts, sulphate of iron 5, sal ammoniac 1, sulphate of indigo 1, gum arabic ⅛th. To be boiled in water, and the paper washed with it exposed to ammonia.

Photographic Paper. See PHOTOGRAPHY, below.

Tracing Paper. Paper well wetted with Canada balsam and camphene, and dried. Another kind is made with nut oil and oil of turpentine; the paper is moistened with it, and then rubbed with flour. A temporary tracing paper is made by moistening paper with pure alcohol: it must be used while wet.

Waxed Paper. Lay the paper on a clean hot iron plate, and rub it over with a piece of white wax inclosed in muslin.

Oiled Paper as a substitute for oiled silk. Boiled linseed oil is reboiled with litharge, acetate of lead, sulphate of zinc, and burnt umber, an ounce of each to a gallon. The sheet of paper being laid on a square board, it is well covered with this mixture. The first sheet is covered on both sides, the second, placed on this, receives one coating, and so on to 20, or 50. Separate and hang up to dry.

FIREPROOFING FOR PAPER. Dip it in a strong solution of alum, and then dry it. Should the paper be extra thick, the same process may be repeated.

PAPER PASTE. Boil white paper in water for 5 hours; then pour off the water, and pound the pulp in a mortar; pass it through a sieve, and mix with some gum-water or isinglass glue. It is used in modelling by artists and architects.

PAPIER-MACHE. A plastic material, formed of cuttings of white or brown paper, boiled in water, and beaten to a paste in a mortar, and then mixed with a solution of gum-arabic in size to give tenacity. It is variously manufactured by being pressed into oil moulds, afterwards dried, covered with a mixture of size and lampblack, and varnished.

PAPYRINE. Dip white unsized paper for ½ a minute in strong sulphuric acid, and afterwards in water containing a little ammonia. When dried it has the toughness and appearance of parchment. See VEGETABLE PARCHMENT.

PARAFFINE. Liquid and solid paraffine are obtained from the tarry product of the distillation of peat, by first distilling off the lighter tar oil, then the residue separately. The crystallized paraffine is separated by a hair sieve, melted, cast into moulds, pressed in a stearine hot press, at a temperature not exceeding 100°; then redistilled, and the same process repeated till it is obtained perfectly pure. The liquid paraffine is redistilled, and burned in lamps. The solids is made into candles.

PASTES. See BLACKING PASTE, FURNITURE PASTE, &c., above. For flour paste, see CEMENTS. For almond paste, honey paste, and tooth pastes, see COSMETICS.

Paste for Cleaning Brass, &c. 1. Rotton-stone in very fine powder, 2 oz., soft soap 1 oz., oil of amber 1 dr.

2. Neat's-foot oil 16 oz., water of ammonia 1 oz., powdered rotten-stone sufficient to form a paste.

3. Rotten-stone 4½ lb, oxalic acid (dissolved in the water) 2 oz., soft soap 8 oz., sweet oil 8 oz., oil of amber 1 oz., boiling water 1 lb. Some substitute oil of turpentine for oil of amber.

Paste for Razors. 1. Emery very finely levigated in the same manner as prepared chalk, mixed with lard or tallow, or a mixture of these with neat's-foot oil.

2. Equal parts of jewellers' rouge, black lead, and prepared suet.

3. PRADIER'S. Best putty powder 1 oz., jewellers' rouge 1 oz., scales of iron ½ oz., levigated Turkey stone 3 oz., beef suet 1½ oz.

4. Mix equal parts of dried sulphate of iron and salt, and apply a gradually increased heat, in a closed vessel. Pulverize, elutriate, and mix with lard or tallow.

PASTILLES, AROMATIC. See PERFUMERY.

PAYNE'S PROCESS FOR RENDERING WOOD FIRE-PROOF. The wood is introduced into a close vessel, which is exhausted of air; the liquid is then admitted, and forced in by the pump till the pressure is from 110 to 140 lb to the square inch. The liquids employed are the liquid sulphuret of

calcium, or of barium; a solution of sulphate of iron is afterwards forced into the wood.

PERCUSSION CAPS, Priming for. 100 gr. of fulminating mercury are triturated, with a wooden muller on marble, with 30 gr. of water and 60 gr. of gunpowder. This is sufficient for 400 caps. Dr. URE recommends a solution of gum mastic in turpentine as a medium for attaching the fulminate to the cap.

PHARAOH'S SERPENTS. The chemical toy sold under this name consists of the powder of sulphocyanide of mercury made up in a capsule of tin foil in a conical mass about an inch in height. Ignited at the apex, an ash is protruded long and serpentine in shape. The fumes evolved are stifling and very deleterious to health.

PHOSPHORESCENT OIL. Dissolve 1 gr. of phosphorus in 1 oz. of olive oil in a test tube by the heat of hot water, or add a larger quantity to some oil of lavender, in which it will dissolve spontaneously. Keep in a close vial.

PHOSPHORUS. See Pocket Formulary.

PHOSPHORUS MATCHES. See LUCIFERS. The old phosphorus bottles with sulphur matches were made by melting phosphorus with a fourth part of wax in the bottles placed in warm water, and turning them about so as to coat the sides.—GRAY.

PHOSPHORUS PASTE FOR VERMIN. Introduce 1 dr. of phosphorus into a Florence flask, and pour over it 1 oz. of rectified spirit. Immerse the flask in hot water, until the phosphorus is melted, then put a well-fitting cork into the mouth of the flask, and shake briskly until cold. The phosphorus is now reduced to a finely divided state. This, after pouring off the spirit, is to be mixed in a mortar with 1½ oz. of lard. Five oz. of flour and 1½ oz. of brown sugar, previously mixed together, are now added, and the whole made into a paste with a little water. Cheese may be substituted for sugar when the paste is intended for rats or mice. (There is said to be no danger whatever of spontaneous ignition, either during or after the preparation of this paste.) —PHARM. JOURNAL.

PHOTOGRAPHY.—PHOTOGRAPHIC COLLODION.

Mr. COLLINS. Put into a large tumbler ½ oz. of strong

nitric acid (sp. gr. 1.50), and add ½ oz. of oil of vitriol. Weigh out 30 grains of cotton-wool (or of paper made from linen rags), and with two glass rods stir it well into the acid, that the cotton may become quickly saturated. It must not be allowed to become soft and gummy. From 15 to 20 seconds is generally sufficient for this immersion. The cotton is next removed by the glass rods, and plunged into a basin of cold water, placed under a running tap,—the cotton being worked about in this until it ceases to redden a piece of blue litmus paper. Remove it, express the water, first by wringing in a clean cloth, then between folds of blotting-paper. Hang up to dry. Take 9 fluid ounces of rectified ether, and dissolve in it 15 grains of this dry prepared cotton. Add 1 oz. of alcohol, cork the bottle tightly, and allow it to stand until any flocculent matter has subsided. Finally, decant into a stoppered bottle for use.

M. MANN finds that the best *collodion wool* is prepared in the following manner: Mix nitrate of potash (powdered but not dried), 20 parts, in a glass cylinder with sulphuric acid (sp. gr. 1.830 to 1.835) 31 parts, and stir until dissolved. Into this mixture, while still warm, but not above 122 F., introduce cotton-wool 1 part, and agitate with a rod until perfectly saturated with the acid liquid. Cover the cylinder now with a glass plate, and allow it to stand for 24 hours at a temperature of 85°. Then remove the wool, place it in a porcelain mortar, and wash with cold water till it has no longer an acid reaction. Finally, free it by repeated treating with boiling water from the least trace of sulphate of potash, which it retains with some obstinacy. The wool being now freed from moisture by bibulous paper, &c., will dissolve readily in a mixture of 7 to 8 parts of ordinary pure ether, and 1 part of absolute alcohol. It may be diluted, without any separation, with an equal quantity of ether again. Perfectly dry wool will dissolve with more difficulty than the fresh.

Mr. HARDWICH recommends pyroxyline made from *paper* for collodion for negatives. Of this 15 to 22 gr. are dissolved in rectified ether (0.70) 53 oz., and alcohol (0.836) 2 dr., to make plain collodion.

To prepare collodion for forming a sensitive surface on glass, on which, if poured in a thin layer, it now leaves only a transparent coating by the evaporation of the ether and spirit, it must first be *iodized*. To each fluid ounce of collo-

dion made by the last formula add 2 dr. of the following solution: Alcohol of 0.836 1 oz., iodide of potassium 20 gr. The plate, before exposure, is rendered sensitive by the aceto-nitrate bath, in a dark chamber. Or, COLLINS'S collodion may be *iodized* and *sensitized* together, as follows: To a saturated solution of iodide of potassium in alcohol, as much iodide of silver is to be added by degrees as the liquid will dissolve: a process which must be performed by candle-light. To every ounce of COLLINS'S collodion add 20 drops of this iodide mixture, and keep it now carefully from the light. To obtain *instantaneous* pictures a few drops of a fresh alcoholic solution of iodide of iron may be added to the iodized collodion just before it is used.

PHOTOGRAPHY.—THE CALOTYPE PROCESS.

Iodide Solution. Iodide of potassium 20 gr., water 1 oz. (The paper to be saturated in this, and dried.)

Aceto-nitrate Solution. Nitrate of silver 20 gr., glacial acetic acid ½ dr., distilled water 1 oz. (Soaked for five minutes in this in a dark room, then in distilled water, then hung up to dry.—The paper is now sensitive to light.)

Developing Solution. Distilled water saturated with gallic acid. (This brings out the picture, after the paper has been exposed in the camera for from 5 to 20 minutes.)

Fixing Solution. Hyposulphite of soda 3 dr., water 1 pint. (The paper is steeped for some time in this, to dissolve the undecomposed iodide, then washed and dried.)

Waxed paper is best for negative pictures. Positive pictures may be printed with paper rendered sensitive in various ways, either from collodion or albumen negatives on glass, or from waxed paper negatives.

PHOTOGRAPHY.—COLLODION PROCESS ON GLASS.

Iodized Collodion. See above. (This is to be spread evenly on the plate, which is first properly cleaned.)

Sensitizing Bath. Nitrate of silver, crystallized and dry, 30 gr., glacial acetic acid 1 min., alcohol 20 min., distilled water 1 oz., iodide of potassium ⅙th gr., carbonate of soda, if required. The nitrate of silver being dissolved in part of the water, the iodide of potassium is dissolved in the remainder, and added, when a precipitate forms which dissolves

on agitation. Any acid reaction is now to be neutralized by carbonate of soda, before adding the acetic acid and alcohol.

Developing Fluid. Glacial acetic acid 15 minims, pyrogallic acid 1 gr., alcohol 10 minims, distilled water 1 oz.

Fixing Fluid. Hyposulphite of soda 1 oz. (or cyanide of potassium 24 gr.), water 2 oz. (For the details, and all further information as to this process, the reader is referred to Mr. HARDWICH'S admirable "Manual of Photographic Chemistry.")

PHOTOGRPHY.—THE ALBUMEN PROCESS ON GLASS.

Iodized Albumen. Albumen 450 fluid parts, of a saturated solution of iodide of potassium, 7½ parts, saturated solution of bromide of potassium 1½ part, water 1 part, solution of caustic potash $\frac{1}{15}$th part. Place in a large bottle, and shake for 10 minutes, till it is filled up with white foam. After 6 hours' standing in a cool place, decant the clear albumen into a tall cylindrical glass vessel. (An hour after it should be spread on the glass plates. These must be passed over the vapor of iodine, to saturate the free potash.) For positives, common salt instead of bromide of potassium.

Silvering Bath. 1500 parts of water, 150 nitrate of silver, 150 glacial acetic aid. (In a gutta percha vessel. After being immersed for 1½ minute, well washed in distilled water, and again passed over the vapor of iodine, the plates are ready for exposure.)

Developing Solution. 300 parts of water, 1 pyrogallic acid, 5 glacial acetic acid, 1 formic acid. (Immersion about ½ an hour.)

Fixing Solution. 100 distilled water, 10 hyposulphite of soda. (Continue this till all yellow iodide disappears; then wash and dry.)

PHOTOGRAPHY—DRY COLLODION PROCESS. Mr. MAYALL.

Exciting Solution. 3 gr. of iodide of cadmium, 1 gr. of chloride of zinc; dissolve in alcohol ½ oz. (This is to be mixed with 1 oz. of ordinary collodion, which must then stand a few days to settle, and be decanted into a dry bottle. Spread as usual.)

Albuminate of Silver Bath. Mix 1 oz. of albumen with

distilled water 16 oz.; add 1½ oz. of glacial acetic acid, shake up, and let it stand 3 hours; then add 1½ oz. nitrate of silver (neutral) in crystals; shake, filter, and let it stand for 24 hours. Add iodide of potassium 2 gr., and filter again, when it is ready for use. (The plate being coated with collodion, this is used in the dark, as an ordinary silver bath. The plate is then washed with distilled water for 5 minutes, and allowed to dry, placed vertically, in a place free from dust. It will keep three weeks. After exposure, it is again passed for 3 minutes into the silvering bath.)

Developing Fluid. 6 gr. of protosulphate of iron, 1 oz. of distilled water, 1 dr. of glacial acetic acid. (After using this, again wash.)

Fixing Fluid. Cyanide of potassium 1 part, distilled water 20 parts. (N. B. Neither the albuminate bath nor the developing fluid must on any account be exposed to daylight.)

M. DUPUIS mixes iodide of zinc with the collodion, renders the plates sensitive with aceto-nitrate of silver, washes them well, and covers them with a layer of dextrine. In developing he uses pyrogallic and citric acids, 1 part each in 300 of water.

M. TAUPENOT'S *Iodized Albumen.* 1 per cent. of iodide of potassium is dissolved in clear albumen of eggs. (This is poured on a collodionized plate which has been passed through the silver bath. It is suffered to drain and dry in darkness, and will keep for several days. When wanted for use, the plates are placed for about 15 seconds in the aceto-nitrate bath.)

M. MARTEN'S *Iodized Albumen.* To the whites of 8 eggs put 4 gr. of iodide of ammonia, 1 gr. of dextrine, 25 gr. of distilled water, and 1½ gr. of grape sugar. These are the proportions for delicate or architectural subjects. For landscapes 2 or 3 times as much iodide is used. (Employed alone, as in the albumen process, or poured over a collodionized plate. The albumenized glasses will keep for any length of time.)

PHOTOGRAPHY—MISCELLANEOUS SOLUTIONS.

1. *Barytic Solution.* Muriate of barytes (chloride of barium) 35 gr., distilled water 2 oz.

2. *Chromatic Solution* (*simple*). A saturated solution of

bichromate of potash. A little sulphate of indigo is sometimes added to vary the color.

3. *Compound Chromate Solution.* Dissolve 10 gr. of bichromate of potash and 20 gr. of sulphate copper in an ounce of water.

4. *Hydriodate of Iron and Barytes Solution.* Hydriodate of barytes 40 gr., water 1 oz., pure sulphate of iron 5 gr.; mix, filter, add a drop or two of diluted sulphuric acid, and when settled decant the clear liquor for use.

5. *Gold-toning Bath for Positive Printing.* Mr. HARDWICH. Pure chloride of gold 1 gr., hyposulphite of soda 1 to 3 gr., hydrochloric acid 4 minims, water 4 oz. For the other SOLUTIONS, see Processes, above.

PHOTOGRAPHY, PAPERS USED IN. (The paper used may be the finest satin post paper, of uniform texture, free from the maker's mark, specks, and all imperfections. CANSON'S paper is much recommended. The papers must be prepared by candle-light, and kept in the dark till used.)

1. *Simple Nitrated Paper.* This is merely paper brushed over with a strong solution of nitrate of silver. In brushing over the paper it must not be crossed. Its sensitiveness is increased by using spirits of wine instead of water. This paper only requires washing in water to fix the drawing.

2. *Muriated Paper.* The paper is first soaked in solution of common salt, pressed with a linen cloth or blotting-paper, and dried. It is then brushed over on one side (which should be marked near the edge) with the solution of nitrate of silver, and dried at the fire. The stronger the solution, the more sensitive the paper. If the barytic solution No. 1 be used instead of common salt, richer shades of color are obtained. A sal-ammoniac solution (10 gr. to 1 oz.) gives a very sensitive paper. A due proportion must be observed in the silver and salt solutions.

Mr. HUNT gives the following as proper proportions: Sensitive paper for the camera. 50 gr. of common salt to 1 oz. of water; and 120 gr. of nitrate of silver to 1 oz of water. Or, 60 gr. of the nitrate, with 40 gr. of muriat of ammonia and 4 oz. of water. Or, 100 gr. with the barytic solution.—Less sensitive, for copying engravings, botanical and entomological specimens, &c. The salt solution to contain 25 gr. of salt to 1 oz. of water. The silver solution 90 gr. in 1

oz. common, for copying lace work, feathers, patterns, &c. The salt solution 20 gr., the silver solution, 40 gr. to an ounce. To fix the drawing on these papers, they must be first washed in lukewarm water, then dipped twice in solution of hyposulphite of soda (1 oz. to 1 pint), then in pure water, and dried.

3. *Iodized Paper.* Brush over the paper on one side (which should be marked) with strong solution of nitrate of silver (100 gr. to 1 oz.); then dip it in solution of iodide of potassium (25 gr. to 1 oz.); wash it in distilled water, drain, and dry it.

4. *Bromide Paper.* Soak the paper in solution of bromide of potassium (40 gr. to 1 oz.); then brush it over with strong solution of nitrate of silver, and dry in the dark.

5. *Calotype Paper.* See CALOTYPE PROCESS.

6. *Chromatype Paper.* Soak the paper in the solution No. 2, and dry it at a brisk fire. To fix the drawing, careful immersion in warm water is all that is required. It is not sufficiently sensitive for the camera.

7. *Compound Chromatype Paper.* Wash the paper with the solution No. 3, and dry it. After the paper has been exposed to the sun with the article to be copied superposed upon it, it is washed over in the dark with a solution of nitrate of silver of moderate strength. A vivid picture makes its appearance, which is sufficiently fixed by washing in pure water. *For copying engravings*, &c. Another method is to brush writing paper over with a solution of 1 dr. of sulphate of copper in 1 oz. of water; and when dry, with a strong but not saturated solution of bichromate of potash.

8. *Cyanotype Paper.* Brush the paper over with a solution of ammonio-citrate of iron. Expose the paper in the usual way, then wash it over with a solution of ferrocyanide of potassium.

9. *Crysotype Paper.* Wash the paper with solution of ammonio-citrate of iron, dry it, and afterwards brush it over with a solution of ferrocyanide of potassium. Dry it in a dark room. The image is brought out by brushing it over with a neutral solution of gold or of silver.

10. *Catalisotype.* Steep paper in water with a drop or two of hydrochloric acid, absorb the superfluous moisture with blotting-paper; brush over with a mixture of ½ dr.

syrup of iodide of iron, 2½ dr. of water, and a drop or two of tincture of iodine. Dry with blotting-paper, and brush over with a solution of 12 gr. of nitrate of silver to 1 oz. of distilled water. It is then ready for the camera. The picture is fixed by washing in water and afterwards in a solution of bromide of potassium (20 grains to 1 ounce).

11. *Paper for Positive Photographs.* Most of the preceding give negative pictures, the lights and shadows being reversed; in the following they are correct: Wash highly glazed paper with solution of sal ammoniac (10 gr. to 1 oz.); dry it, and brush it over with the following solution: Dissolve 120 gr. of crystallized nitrate of silver in 1½ oz. of distilled water, and add 1½ oz. of alcohol; after it has stood a few hours, filter it. Expose the paper thus washed to the sunshine till it is darkened; if mottled, wash it a second time, and expose it again. To use this paper, wash it over with the solution No. 4, expose it in the damp state, with the engraving or other object on it to the light, and fix the drawing by washing with water only. (To copy objects, lay them on a plate of clear glass fixed in a frame, place the prepared paper over them, and fix a back with a cushion attached to it so as to press the paper closely on the glass. The glass is then exposed to the light, and the drawing afterwards fixed, as described above. For feathers, lace-work, and other objects which freely admit light through them, the nitrated paper and less sensitive muriated papers may be used. For copying engravings, leaves, and other botanical objects, or entomological specimens, the more sensitive muriated papers, or the bromide paper, or other sensitive kinds, may be used. Engravings should be wetted and placed with their face to the prepared side of the paper, and kept in close contact with it. Leaves should have their under surface next the glass. For the camera the most sensitive samples of the muriated papers, made with not less than 100 gr., of the nitrate of silver to the ounce, are selected. The calotype is still more certain. The papers intended for the camera require to be very carefully prepared.)

12. *Instantaneous Positive Paper.* Mix 6 dr. of a saturated solution of bichloride of mercury with distilled water 1 pint. Float the paper on this solution in a flat dish. Dry it; take into a dark place lit by a candle

with a yellow glass, and render it sensitive by a solution of nitrate of silver (38 gr. to 1 oz.). To print, expose to a perpendicular light from 2 to 10 seconds in summer, about 1 minute in winter; then immediately cover with a black cloth. (The image, at first very feeble, is developed by this solution; Sulphate of iron 15 gr., glacial acetic acid 25 gr., to distilled water 1 oz. The deepening of tint must be watched, and arrested at the proper moment. Then wash, and fix with hyposulphite.)

13. *Albumenized Paper for Positive Printing.* White of egg, and water equal parts, iodide of potassium or chloride of sodium 5 gr. to 1 oz. (or bromide of potassium 20 gr.) Coat the paper with this solution. Dry. Immerse in the dark in bath of 120 gr. nitrate of silver to 1 oz. Dry again. (This is exposed, with the negative over it, for 10 to 15 minutes.)

14. *Prepared Wax Paper.* De la Blanchere. Make a strong size by digesting 25 parts of gelatine, 50 of linseed, and 150 of rice flour, in hot water 2000 to 3000 parts. Filter through a cloth. Take of this size, when cold, 1000 parts by weight, and dissolve in it sugar of milk 50 parts, iodide of potassium 35, bromide of potassium 5 parts.

The pieces of paper, well waxed, are steeped in this bath, one by one, for about 10 minutes, then hung up to dry. They will keep well for 2 or 3 months, and may any time be sensitized for use.

15. *Paper prepared with Mucilage, Turpentine, and Wax.* Lespiault and Blot. Boil 30 gr. of Carrageen moss in ½ pint of water for 5 minutes; strain through fine linen, then add ½ pint more water hot. Add then 1 per cent. of iodide of potassium, and 15 per cent. of simple syrup. A few drops of acetic acid and a few fragments of camphor will preserve this mucilage. The paper is soaked for 5 minutes, then removed, drained, and dried by suspending from one corner.

When quite dry, immerse for some minutes in the turpentine-wax bath—made by saturating oil of turpentine with wax, filtering, and adding pure iodine 1½ gr. to each ounce. This paper is sensitized, when wanted, in an aceto-nitrate bath.

Photography.—Miscellaneous Recipes.

To Clean Glass Plates. Mr. Mayall. Shake up to-

gether alcohol 30 parts, strong liquid ammonia 10, water 40, and fine tripoli 30 parts. The plates are to be rubbed hard and evenly with balls of cotton-wool dipped in this mixture. Rub again, when dry, with a clean ball of cotton; lastly, dust the back and edges with a clean hog's-hair brush.

To Clean off Collodion Pictures. This may be done, whether they have been varnished or not, by means of a tuft of cotton-wool dipped in wood spirit.

To Color Photographic Prints. This may be done variously in water or oils. A simple way is to rub in slowly with a small camel-hair brush a minute piece of dry color laid upon the part, as of flesh tint for the face, &c. When properly distributed, the paper may be breathed upon, and the tint will not easily be rubbed off. Or it may be carefully coated with gelatine.

M. Minotto has described a plan of coloring on the back of the paper. The picture, being held up to the light, is first faintly outlined, on the reverse side; colors are then laid on, of water or oil, as preferred, on this side. When dry, the paper is rendered transparent by a varnish, and the colors will then appear through it with all the delicacy and effect of a miniature on ivory. Good strong writing paper is best for this purpose; the colors must be vivid; and the varnish may consist of Canada balsam dissolved in turpentine, or a mastic varnish may be used, or turpentine and wax, or oil.

Artificial Ivory for Photographers. Mayall. Tablets of gelatine or glue are immersed in a solution of alumina. When entirely penetrated by the alumina, the slabs are to be removed, dried, and polished like ivory.

Photographic Varnish. See Varnishes.

Photographic Engraving. M. Niepce. (The original process, invented by M. Niepce, the elder, before the discovery of the daguerreotype, has been recently modified and improved by his nephew.) Two parts of Judea bitumen are dissolved in 10 parts of essence of lemon and 90 of benzine, yielding a varnish which is acted upon by the sun's rays in such a manner that the part exposed is rendered much more insoluble than the covered part in certain spirituous and oily liquids. This varnish is spread evenly on the surface

of the plate, and allowed to dry in the dark. A camera picture obtained by any photographic process, or a waxed engraving to be copied, is adjusted over the stone by means of a glass, and it is then exposed to the sunshine for about ¾ of an hour. The resin surface is now acted upon by a liquid, composed of 3 parts of naphtha and one of benzine. This disengages the image. The stone may now be prepared for printing in the usual way. If a steel plate be used, the resinized parts will offer sufficient resistance to enable the surface to be acted on by aquafortis. Or a glass plate may be submitted to the action of the vapor of hydrofluoric acid.

Mr. McPherson. Judea bitumen is dissolved in sulphuric ether, and the solution used to form a layer on the stone. It rapidly dries. The parts unacted on by light are instantaneously dissolved on plunging in a bath of sulphuric ether.

Professor Ramsey has described a modification of the last process for etching on steel or copper. After the plate has undergone the solvent action of ether, the parts which are left bare are gilded by an electro-gilding arrangement. The gold will not attach itself to the bitumen. This is next removed by means of spirit and heat. Nitric acid is now applied as in the common etching process, and acts on the parts of the plate uncoated by gold.

Mr. Fox Talbot. Some bichromate of potash is dissolved in a strong solution of gelatine, and this is poured on the surface of the steel plate and dried. A yellow coating is thus formed. Upon this is placed the object to be copied, as fern leaves, grasses, or pieces of lace; they are pressed closely by a piece of glass, and exposed to sunshine. By this the salt is decomposed, and the chromic acid, by acting on the gelatine, produces in the exposed parts a brown opaque surface, which is comparatively insoluble in water. The plate being placed in water, the gelatine is dissolved off the steel opposite the superimposed object. A solution of bichloride of platinum being now poured on the plate, the lines are rapidly etched in. The plate being then washed until all the gelatine is removed, it is submitted to the operation of the copperplate printer. Very beautiful prints of grasses, textile fabrics, &c., have been obtained in this way.

M. Poitevin. (In Mr. Talbot's process, the bichromate of potash and gelatine are used as a varnish; in this new

process the surface formed by them is itself used to mould the plate.) On a plate of glass or other level surface is poured a uniform layer of solution of gelatine, of thickness according to the kind of print desired. When dry, the plate is placed for several minutes in a concentrated solution of bichromate of potash, then passed quickly through water, and allowed to dry in the dark. The plate is now exposed for a considerable period to the sunlight, either in the camera or with a transparent picture superimposed. The plate is now plunged in water in a darkened room. The gelatine unacted upon swells up, so as to produce relief. The parts acted on by light scarcely swell at all; they form the hollows. The process is arrested by pouring on a solution of protosulphate of iron. The dry plate may now be moulded in plaster, or other proper material, and a copper plate finally obtained from the mould, by means of the electrotype process.

PIGMENTS. A few of these have been noticed before; see INDIGO, LAKES, PRUSSIAN BLUE, PURPLE OF CASSIUS. They generally constitute a distinct branch of manufacture, but a brief account of the composition of some of them may be useful. Those of which the coloring matter is derived from the animal and vegetable kingdoms will first be noticed; then the mineral colors.

Carmine. Several processes have been published for this beautiful pigment, but probably some minute precautions, not generally known, may be necessary to the production of the finest quality. The climate and state of the atmosphere are said to influence the result.

1. *Madame* CENETTE'S *process.* Into 6 pails of boiling clear, soft water, in a copper vessel, throw 2 ℔ of powdered cochineal of good quality; boil for 2 hours, add 3 oz. of purified nitre, and after a few minutes, 4 oz. of salt of sorrel. Remove the vessel from the fire, let the contents settle for 4 hours, draw of the clear liquor with a syphon into flat plates, and leave it at rest for 3 weeks. Carefully detach the pellicle of mould from the surface, withdraw the liquid with a syphon and pipette, and dry the deposit in a stove.

2. Boil 4 quarts of soft water in a pewter kettle, add to it 4 oz. of finely-powdered cochineal; boil for 5 minutes, adding 2 dr. of powdered cream of tartar; then add 8

scruples of Roman alum, and keep the whole on the fire for a minute longer. Let the decoction settle, decant it into cylindrical glasses, and cover them. When the carmine has subsided, pour off the clear liquor, and dry the sediment. By adding solution of tin to the liquid, more carmine is obtained.

3. Into a 14-gallon boiler of tinned copper put 10 gallons of distilled water, or filtered rain-water; when it boils, sprinkle in, by small quantities, 1 lb of powdered cochineal, and keep it boiling for half an hour. Then add 3½ oz. of crystallized carbonate of soda; in a minute or two, draw the fire, and add 1½ oz. of Roman alum in fine powder; stir with a glass rod till the alum is dissolved, leave it to settle for 25 minutes, draw off the liquor with a glass syphon, and strain the rest through a coarse linen cloth. Clean the boiler, return into it the clear-colored liquor, and stir into it the whites of 2 eggs, previously well beaten with a quart of warm (not hot) water. Then light the fire, and heat the liquor till it begins to boil; separate the coagulum by filtration, wash it on the filter with distilled water, spread it thinly on earthen plates, and dry it in a stove.

Inferior carmine may be improved by dissolving it in water of ammonia, and precipitating it by acetic acid and alcohol.

Cochineal Lake. Add 2 lb of pearlash to the red liquor from which the carmine has been prepared in the last process, and return it to the boiler with the dregs of the cochineal; boil for half an hour, draw the fire, and when the sediment has subsided, draw off the clear liquid into an earthen vessel. Pour on the sediment a solution of 1 lb of pearlash in 2 gallons of water, and boil for half an hour. Filter, and return both liquors into the copper. When as hot as the hand can bear, add to the liquor, by little and little, 3 lb of powdered Roman alum, and let it simmer for 5 minutes. Allow it to settle, draw off the clear liquors, collect the sediment on a filter, wash it with clean rain-water, and leave it covered with a cloth for a few days, till half dry; form it into small lumps, and dry them in a stove.

Carthamine or Safflower Lake. Wash safflower till the water comes off colorless; mix it with water holding 15 per cent. of carbonate of soda in solution, so as to form a thick paste; leave it for several hours, then press out the red

liquid, and nearly neutralize it with acetic acid. Then put cotton into it, and add successive small portions of acetic acid, so as to prevent the liquid becoming alkaline. In 24 hours take out the cotton, wash it, and digest it for half an hour in water holding 5 per cent. of crystallized carbonate of soda in solution. Immediately on removing the cotton, supersaturate the liquid with citric acid, and collect the precipitate, which must be repeatedly washed in cold water. For pink saucers the liquid is allowed to deposit in the saucers. Mixed with the scrapings of French chalk it constitutes rouge.

Lakes are also obtained from Brazil-wood and madder, by adding alum to a concentrated decoction of the former, or to a cold infusion of the latter (made by triturating the madder, inclosed in a bag, with the water), and afterwards sufficient subcarbonate of potash or soda to throw down the alumina in combination with the coloring matter. The precipitate is to be washed and dried. A little solution of tin added with the alum improves the color. Lakes may be obtained from most vegetable coloring matters by means of alum and an alkaline carbonate. *Yellow Lake* is made from French or Persian berries, by boiling them in water, with a little soda or potash, and adding alum to the strained liquor as long as a precipitate is thrown down. Or by boiling weld, or quercitron bark, in water, and adding alum and chalk in a pasty state.

Rose Pink. Boil 6 ℔ of Brazil-wood and 2 ℔ of peach-wood in water, with ¼ ℔ of alum; and pour the strained decoction on 20 ℔ of sifted whiting.

Bistre. It is obtained from the soot of beech-wood.

Sap Green. The expressed juice of buckthorn berries (and sometimes of other species of rhamnus, and also of privet berries) is allowed to settle, and the clear liquid evaporated to dryness. A little gum arabic is sometimes added to the juice.

The beautiful colors of the *mauve* series are prepared from coal tar by patented processes.

Mineral Pigments. *Azure Blue, or Smalts.* The common is made by fusing zaffre (roasted cobalt ore calcined with siliceous sand) with potash. A finer quality is obtained by precipitating a solution of sulphate of cobalt by a solution of

silicate of potash. Another cobalt blue is obtained by adding a solution of phosphate of soda to a solution of nitrate of cobalt, and mixing the precipitate, washed, but not dried, with eight times its weight of fresh hydrated alumina. When dry, heat it to a cherry red.

Egyptian Azure. Carbonate of soda 16 oz., calcined flints 24 oz., copper filings 4 oz. Pulverize, mix, and fuse in a crucible for two hours. When cold, reduce to powder.

Blue Verditer. It is generally stated to be made by adding chalk to a solution of nitrate of copper produced in the process of refining silver; but Mr. PHILLIPS did not succeed in making it by this means, and found no lime in the best samples.

New Blue. Mix equal parts of common arseniate of copper (see Mineral Green, below), and neutral arseniate of potash, fuse by heat in a large crucible, then add to the fused salt one-fifth of its weight of nitre. Effervescence takes place, and the salt becomes blue. Cool, pulverize, and wash.

Chrome Yellow. To a solution of bichromate of potash add a solution of nitrate of lead as long as a precipitate forms. Wash the precipitate, and dry it with a gentle heat. An inferior kind is said to be made by 4 ℔ of pure white lead, 1 ℔ of bichromate of potash, and 20 ℔ of water, and boiling till the water becomes colorless. Or 75 parts of precipitated sulphate of lead may be acted on by a hot solution of neutral chromate of potash 25 parts. A mixed product of chromate and sulphate of lead is thus obtained, which is said to cover as well as the pure chrome yellow, and is much cheaper. (RIOT.)

Chrome Red. Melt saltpetre in a crucible heated to dull redness, and add chrome yellow, by small portions, till no more red fumes arise. Allow the mixture to settle, then pour off the melted salt from the heavy sediment, and wash the latter with water, which should be quickly poured off, and dry the pigment. The liquefied salt poured off contains chromate of potash, and is reserved for making chrome yellow.

Orange Chrome is chrome yellow acted on by an alkali, which deprives it of part of the chromic acid.

King's Yellow. This is a yellow sulphuret of arsenic, now

almost superseded by chrome yellow, but occasionally used for killing flies.

Naples Yellow. Mix 12 parts of metallic antimony, 8 parts of red lead, and 4 of oxide of zinc, and calcine in a reverberatory furnace. The mixed oxides are rubbed together, fused, and the fused mass elutriated into a fine powder.—Dr. URE. M. GUIMEL recommends 1 part of well-washed antimoniate of potash to be ground into a paste with 2 parts of red lead, and the powder exposed to a red heat for 4 or 5 hours, keeping the heat moderate.

Brighton Green. An inferior color, made with 28 ℔ of whiting, or white lead, 7 ℔ sulphate of copper, 3 ℔ sugar of lead, and ¼ oz. of bichromate of potash.

Brunswick Green. Pour a saturated solution of muriate of ammonia over copper filings in a close vessel placed in a warm situation; add more of the solution from time to time till three parts of the muriate have been used to two of copper. After standing for a few weeks the pigment is separated from the unoxidized copper by washing through a sieve. It is then to be well washed, and dried slowly in the shade. It is often reduced with white lead; some samples contain arsenic.

Arsenical Copper Greens. Of these are several varieties.

Mineral Green, Scheele's Green or Arsenite of Copper. 1. Dissolve 11 oz. of white arsenic and 2 ℔ of carbonate of potash, by heat, in a gallon of water. Dissolve also 2 ℔ of sulphate of copper in 3 gallons of water. Filter each solution separately, and add the former gradually to the latter as long as it occasions a precipitate. Wash the precipitate, drain it, and dry it

2. Dissolve 50 ℔ of sulphate of copper and 10 ℔ of lime in 20 gallons of good vinegar, and add quickly a boiling hot solution of 50 ℔ of white arsenic. Stir repeatedly, then allow it to settle; decant the clear liquor (which is reserved to dissolve the arsenic next time), and wash the precipitate, and dry it.

3. *Emerald Green.* Mix ten parts of pure verdigris with sufficient boiling water to form a soft pulp, and strain this through a sieve. Dissolve 9 or 10 parts of white arsenic in 100 parts of boiling water, and, whilst boiling, let the verdigris pulp be gradually added, constantly stirring the mixture till the precipitate becomes a heavy granular powder.

Green without Arsenic. Dissolve 48 lb of sulphate of copper, and 2 lb of bicarbonate of potash in water, and add to the clear solution 2 lb of pearlash and 1 lb of chalk.

Rinmann's Green Pigment. Dissolve together in sufficient water 1 part of sulphate of cobalt and 3 of sulphate of zinc; precipitate with carbonate of soda, wash the precipitate, and calcine it.

Chrome Green. A mixture of chrome yellow and Prussian blue. (See also Chrome Oxide, p. 334.)

Barth's Green. A mixture of Prussian blue and yellow lake.

Ultramarine, Factitious. Take 70 parts of silica, or pure siliceous sand in fine powder; 240 parts of recrystallized alum, calcined; 144 parts of sulphur; 48 parts of finely powdered charcoal; 240 parts of dry carbonate of soda. These are mixed together with the greatest care till the mixture appears of uniform color under a powerful magnifier, and the mixture exposed to a moderate red heat in a closely covered crucible for an hour and a half. Wash the product in boiling water. Mix the powder with its own weight of sulphur and 1½ its weight of dried soda, and burn as before; heat it again with sulphur and soda, and wash it till the filtered fluid no longer colors acetate of lead. If a sample of the dried powder becomes blue when burnt with sulphur, it is ready for the last operation. Spread over a cast-iron plate a layer of sulphur a line in thickness, and over it an equal layer of the dried powder after having passed it through a gauze sieve. Heat the plate so as to burn away the sulphur at the lowest possible temperature. Reduce the pigment to powder, and repeat the burning with sulphur and pulverization till the color is perfect.

White Lead is carbonate of lead prepared by various processes. *Zinc White* is oxide of zinc, prepared by combustion. *Oxychloride of Zinc* is also used. *Antimony White* is oxide of antimony.

Plates, Daguerreotype, are prepared by cleaning and polishing the silver surface, exposing it to the vapor of dry iodine, or tincture of iodine, or iodide of bromine, or bromide of lime. After having the image thrown on them, they are

exposed to the vapor of mercury. But the manipulations and precautions necessary to the success of the operation are too numerous to detail here.

PLATE-BOILING POWDER. Equal parts of cream of tartar, alum, and common salt. A small quantity added to the water in which plate is boiled gives it a silvery whiteness.

PLATE POWDERS. 1. *Jewellers' Rouge.* Dissolve green vitriol in hot water, and add a solution of pearlash as long as it throws down a precipitate. Wash the precipitate repeatedly with warm water, drain it on calico, and finally calcine it till it assumes a bright color. It is sometimes made by calcining the sulphate of iron with a strong heat till oxide of iron only remains. Let it be triturated with water, and prepared in the same way as prepared chalk. See POLISHING POWDER, below.

2. *French Plate Powder.* Mix 1 part of jewellers' rouge with 12 of carbonate of magnesia.

3. Finely prepared chalk, or burnt hartshorn. One way in which these are used is to boil them with water with pieces of rag; the finer particles are entangled in the fibres of the the rags, which are dried and kept for use.

4. Quicksilver with chalk 1 oz., prepared hartshorn 8 oz., prepared chalk 4 oz. Powders containing quicksilver, besides the necessary wearing of the surface, are supposed to render the plate more brittle. If used it should not be in larger proportion than the above.

5. Finest putty powder 1 oz., levigated chalk 5 oz.; a little rouge may be added to color it.

See NOVARGENT, SILVERING POWDER, &c., for restoring the silver to plated goods.

PLATINA, BLACK (OXIPHOROUS). Dissolve protochloride of platinum in a boiling solution of potash, add alcohol in small proportions till effervescence ceases. Boil the black precipitate successively with alcohol, muriatic acid, and potash, and finally 4 or 5 times with water.

PLATINATED ASBESTOS. Dip asbestos in a solution of chloride of platinum, and heat it to redness. It causes the inflammation of hydrogen in the same manner as sponge platinum. —Dr. HARE.

PLATINIZED SILVER. Silver plates for SMEE'S voltaic battery

are covered with pulverulent platinum by adding a little bichloride of platinum to acid water, and decomposing the solution by the use of a platinum terminal in connection with the copper of a battery, the silver plate to be platinized being in connection with the zinc. Platinum itself is sometimes platinized in the same way. Sometimes the plates are platinized without the battery. The following solution is used by Dr. WRIGHT for the plates of his battery: Saturated solution of chloride of platinum ½ drachm, sulphuric acid 1½ drachm, water 2 drachms. Dip the plates in it for a few seconds, and wash them quickly.

PLATINUM, CHLORIDES OF. Dissolve platinum in nitro-muriatic acid, and evaporate with a gentle heat to dryness. The red bichloride remains. Heated to 450°, the protochloride remains.

PLATINUM SPONGE. Dissolve separately in rectified spirit chloride of platinum and sal ammoniac. Mix the solutions, and heat the precipitate to redness. For balls for hydrogen lamps, form the precipitate into balls while moist, and afterwards burn them.

POISON. See BEETLE WAFER, BUG POISONS, RAT POISON; PHOSPHORUS PASTE, REMEDIES FOR BLIGHTS, &c.

POLISH. See FRENCH POLISH.

POLISH FOR BOOTS, &c. See BLACKING.

POLISHING POWDER FOR SPECULA. Lord Ross. Precipitate a dilute solution of sulphate of iron by ammonia in excess; wash the precipitate, press it in a screw press till nearly dry; then expose it to heat until it appears of a dull red color in the dark.

POT POURRI. See PERFUMERY.

POTASH, CARBONATE OF. *Salt of Tartar.* Mix 10 parts of pearlash with six of water; let them stand in a cool place for 24 hours, stirring them frequently: filter; concentrate the solution by a gentle heat, stirring constantly, remove from the fire as soon as the liquor begins to appear opaque, and continue the stirring until cold. To procure it purer, redissolve it in an equal weight of distilled water, filter, and evaporate to dryness. It is free from sulphate of potash, but contains 1.1 per cent. of chloride of potassium.

POTASH, CHLORATE. See CHLORATE OF POTASH.

POTATO DISEASE, REMEDIES FOR. The following are some of the more recent propositions for the prevention of this terrible and well-known plant epidemic:

1. Some direct that the haulms and leaves should be entirely removed as soon as the least symptom of disease shows itself in them. The roots may grow on to full size without becoming affected.

2. The CHEVALIER CLAUSEN'S Cure. The potatoes, before planting, are wet with water acidulated with sulphuric acid (1 part to 500), and, before they are dry, powdered sulphate of lime is thrown over them. This plan has been found a very efficient one.

3. Mr. T. HERAPATH'S Plan. The roots, before planting, are allowed to become dry, and then dipped for a short time in a weak solution of sulphate of copper. The land is dressed with a mineral composition (2 parts of lime to 1 of salt) instead of farmyard manure. This process also has been attended with considerable success.

POUNCE. Powdered gum juniper is used under this name for preparing parchment for writing on. For liquid pounce, see INK, MARKING.

POUDRE CLARIFIANTE. Beat together the whites and yolks of eggs, dry them with a very gentle heat, and reduce to powder. For clarifying wines and syrups.

POWDERS. See TOOTH POWDERS, and HAIR POWDERS, under COSMETICS; SCENT POWDERS, under PERFUMERY, &c.

PRESERVATIVE LIQUIDS. See ANATOMICAL SUBJECTS, and ANIMAL SUBSTANCES, TO PRESERVE.

PRUSSIATE OF POTASH (YELLOW). What is known in commerce by this name is the ferroprussiate of potash, or ferrocyanide of potassium. It is prepared by fusing in an egg-shaped iron pot a mixture of 2 parts of pearlash and 5 parts of dry animal matters, such as horns, hoofs, tallow-chandlers' greaves, &c., till the fetid vapors cease to be produced. Iron filings are sometimes added, but usually the iron necessary to the formation of this salt is derived from the iron pots and stirrers. The fused mass (*prussiate cake*) is allowed to cool, dissolve in warm water, and the clear filtered or decanted solution evaporated, that crystals may form. These are dissolved in hot water, and the solution allowed to cool very slowly, that large crystals may form.

RED PRUSSIATE OF POTASH. *Ferridcyanide of Potassium.* Into a dilute solution of the above prussiate of potash a current of chlorine gas is passed, till the solution ceases to give a blue precipitate with persalts of iron. It is then evaporated, crystallized, and recrystallized till quite pure. (M. POSSELT advises to add a few drops of solution of potash to the boiling liquor, to decompose the green matter that is formed; to filter the hot solution, to separate some peroxide of iron which is thrown down, and to let the liquor cool very slowly.) Or, boil yellow prussiate of potash with 12 or 15 parts of water, and while boiling add good chloride of lime until a filtered sample no longer yields a blue precipitate with persalts of iron. Filter quickly, and add carbonate of potash till the liquid has a faintly alkaline reaction, then evaporate for crystallization.—CHEMIST, vol. viii.

PRUSSIAN BLUE. *Percyanide, ferrocyanide, or ferroprussiate of iron.* Commercial Prussian Blue is made by adding to a solution of prussiate of potash, or of prussiate cake, a solution of 2 parts of alum and 1 of sulphate of iron, washing the precipitate repeatedly with water to which a little muriatic acid has been added, and exposing it to the air till it assumes a deep blue color. A purer kind is made by adding a solution of persulphate or perchloride of iron to a solution of pure ferroprussiate of potash. TURNBULL'S Prussian Blue (ferridcyanide of iron) is made by adding a solution of red prussiate of potash to one of protosulphate of iron; or by adding protosulphate of iron to a mixture of yellow prussiate of potash, chloride of soda, and hydrochloric acid.

SOLUBLE PRUSSIAN BLUE. Add a solution of protosulphate of iron to a solution of prussiate of potash, and expose the precipitate to the air till it becomes blue, and wash it till the soluble salts are washed away. By continuing the washing, the blue itself dissolves, forming a deep blue solution, which may be evaporated without decomposition. Or add a solution of persulphate of iron to a solution of ferroprussiate of potash, keeping the latter in excess; wash the precipitate until it begins to dissolve, and dry it. See INK, READE'S *Patent Blue*, for another method.

PURPLE OF CASSIUS. See Aurum Stanno Paratum, Pocket Formulary. Many other processes have been proposed,

of which the following is one: Dissolve 3 gr. of gold in aqua regia, avoiding excess, and dilute with 3 oz. of water. Mix 30 gr. of pink salt (the bichloride of tin with sal ammoniac) with $3\frac{1}{4}$ gr. of tin filings and 2 dr. of water till the tin is almost entirely dissolved: add 7 dr. of water, and add this solution to the gold solution, slightly warmed. Wash the precipitate, and dry it.

PYROLIGNEOUS ACID, PYROXYLIC SPIRIT, PYROACETIC SPIRIT, &c. } By the destructive distillation of dried wood, chiefly that of the beech and birch, in iron cylinders, an acid liquor and tar are produced. These are received in proper reservoirs, and are afterwards separated. The tar is subjected to distillation, and yields *oil of tar* (containing creasote, eupione, &c.), and leaves a residuum of pitch, or English asphalt. The acid liquor, separated from the tarry deposit, is also distilled: the first portion which comes over contains the pyroxylic spirit, which is rectified by one or more distillations. It may be further purified by distilling it with dried muriate of lime, and finally with quicklime. This constitutes one of the articles sold under the name of naphtha, and is regarded by chemists as a hydrated oxide of methule. After the pyroxylic spirit has come over, the crude pyroligneous acid distils, which still holds some tar and empyreumatic oil in solution. It is purified by saturating it either directly with common soda, or first with lime, or rather chalk, and when the neutral solution has become clear, evaporating it to 1.114 sp. gr., and adding sufficient saturated solution of sulphate of soda to decompose the impure acetate of lime. The clear solution obtained by either process is then evaporated, that the acetate of soda may crystallize. This is afterwards roasted at a temperature of about 500° Fahrenheit, to destroy the tar, and again dissolved and crystallized. The purified acetate is then distilled with sulphuric acid. See Acidum Aceticum, P. F. For some manufacturing purposes, an impure acid is obtained by merely saturating the crude pyroligneous acid with lime, evaporating to dryness, and distilling with sulphuric acid.

If acetate of lime or acetate of lead be distilled without addition, and the liquid which comes over be rectified over lime, *pyroacetic spirit* is obtained: this is also termed *acetone*.

PYROPHORUS. This name is given to several compounds, prepared by calcination, which take fire when exposed to the air, especially when breathed upon. The following are perhaps some of the best:

1. Heat tartrate of lead in a tube of hard glass, and securely close the tube before the charred residuum becomes cold. A little poured out and breathed upon takes fire. The tartrate of lead is made by dissolving separately 2 dr. of tartaric acid and 5 dr. of crystallized acetate of lead in sufficient water, mixing the solutions, and collecting, washing, and drying the precipitate.

2. Calcine tartar emetic in a similar manner, or in a closed crucible.

3. Mix 11 parts of lamp-black with 2 of powdered sulphate of potash, and heat the mixture strongly in a closely-covered crucible. The product is so combustible that it can scarcely be transferred to a bottle without danger.

4. Mix 3 parts of powdered alum with 1 of flour, and calcine the mixture in a common vial coated with clay or placed in sand, till it ceases to emit a blue flame. Before it is cold, close it securely with a sound cork or glass stopper.

5. Mix neutral chromate of lead with 1-6th its weight of sulphur; triturate them with water sufficient to form a paste, and make it into pellets; dry these perfectly, then heat them in a tube till the sulphur is all driven off, and secure as the last.

RAT AND MICE POISONS. (Such as contain *arsenic* are placed first, and afterwards several compounds which have been introduced as substitutes for that mineral, which has proved so destructive of human life. According to a recent Act of Parliament, this dangerous compound can only be purchased in wholesale quantities. Among other precautions taken to prevent accident, it is provided that it be mixed with coloring matters, such as soot and indigo, in order to prevent its being taken by mistake, or insure detection if designedly administered.)

Arsenical Paste (authorized by the Government of France).

1. Melt 2 ℔ of suet in an earthen vessel over a slow fire, and add 2 ℔ of wheat flour, 3 oz. of levigated white arsenic, 2½ dr. of lamp-black, 15 drops of oil of aniseed. It may be used alone, or mixed with bread crumbs, &c. (For destroying rats and mice.)

2. For barn floors. Mix a pint of good flour with as much yellow arsenic as will lie on a shilling; put this in a small heap on the floor, and over this put another pint of good flour unmixed. Draw a track up to the heap with a feather dipped in oil of aniseed and oil of caraways, and sprinkle this over with a little flour.*

3. Mix a quart of the best oatmeal, 2 oz. of powdered loaf sugar, 6 drops each of the oils of rhodium, caraway, and aniseed, and ¼ gr. of musk. Mix them very perfectly without touching the mixture with the hands. Place in a retired place 6 or 8 pieces of clean board, and on each two tablespoonfuls of the powder, for a few successive nights, without disturbing the rats. About the sixth night, if they are found to eat freely, mix a teaspoonful of white arsenic with the powder. What remains in the morning should be burnt, avoiding the fumes.—THE CHEMIST, vol. vi.

4. White arsenic 2 oz., carbonate of barytes 2 oz., white sugar 3¾ oz., rose pink ¼ oz., oil of aniseed and oil of rhodium, of each 5 drops.

5. Malt flour 1 ℔, oil of rhodium 3 drops, sugar 2 oz., 8 cloves, a tablespoonful of caraway seeds, all beaten in a mortar. Lay it in small parcels where they frequent, for 3 or 4 nights, till they eat freely, then add some arsenic dissolved in spirits of salts.—MAYER.

6. *Ointment for Rats in Ricks.* Mix together 1 ℔ of fresh butter, free from salt, 1½ oz. of calomel, 8 oz. of crumbs of white bread, 2 oz. of sugar, 5 drops each of oils of nutmeg and rhodium, and 2 drops of oil of aniseed. To use it, make a hole with the arm under the ridge; into this hole insert a stick, and on the middle of it, where it does not touch the rick, put a lump of the ointment. *For Traps.* Put the same with 2 or 3 drops of oil of thyme.

7. *Hampshire Millers' Rat Powder.* Mix 1 oz. of nux vomica in powder with a pound of fresh oatmeal, and add a few drops of oil of rhodium, or, what answers better, oil of aniseed with musk.

8. *Philanthrope Muophobon.* A French preparation so called consists of 1 part of emetic tartar to 4 of farinaceous and other ingredients.

---

* The following is an old rat-catcher's receipt for oils to attract rats: Two dr. of oil of aniseed, 2 drops of nitrous acid, and 2 gr. of musk. Oil of rhodium is also supposed to be very attractive to these vermin. Assafœtida with these oils is sometimes used.

9. Put into a flash 2 dr. of phosphorus and 5 or 6 oz. of water, put the flask in warm water (about 150° Fahrenheit) till the phosphorous is liquefied; pour the contents into a mortar, and immediately add 5 or 6 oz. of rye-meal; when cool, add the same quantity of melted fresh butter, and 4 oz. of sugar.

10. Another form of the phosphorus compound is: Melt 1 ℔ of lard in a bottle plunged into water, and heated to 150° Fahrenheit. Introduce into it ½ oz. of phosphorus and add a pint of proof-spirit. Cork the bottle securely after its contents have been heated from 140° to 150°, and taking it out of the water-bath, agitate it briskly till the phosphorus is uniformly diffused; repeat the agitation occasionally as it cools, and, when cool, pour off the spirit which has separated (which may be reserved for the same purpose), and incorporate with the fatty compound wheat flour and sugar. Oil of rhodium or aniseed may be added. Place little lumps of this in the rat-holes, and set some water near for them to drink. For a third recipe, see PHOSPHORUS PASTE FOR VERMIN, p. 401.

11. Valentia almonds 1 oz., treacle 2 oz., carbonate of barytes 1 oz., oil of aniseed 5 drops, flour enough to form a paste.

12. Powdered squill ½ oz., strong cheese 2 oz.; mix, and form into balls. *For Mice.*

RENNET. The stomach of a calf washed, salted and dried.

RENNET, LIQUID. ESSENCE OF RENNET. Fresh rennet 12 oz., salt 2 oz., proof-spirit 2 oz., white wine a quart; digest for 24 hours, and strain. A quart of milk requires 2 or 3 teaspoonfuls. WISLIN directs, 10 parts of a calf's stomach, salt 3 parts. The membrane of the stomach is to be cut with scissors, and kneaded with the salt and the rennet found in the interior of that organ; the whole left in a cool place in an earthen pot till the cheesy ordor is replaced by the proper odor of rennet, which will be in 1 or 2 months. Then add 16 parts of water and 1 of spirit. Filter and color with burnt sugar.

RUST, TO PREVENT AND REMOVE. Steel goods are rubbed over with a mixture of lime and oil, to preserve them from rusting. Mercurial ointment has been recommended for the same purpose. M. PAYEN recommends plunging the articles into a solution of common soda. Spots of rust are removed

by rubbing them with very fine emery and sweet oil. A chemical means of removing them, the ammoniacal chloride of zinc, may be found useful. See ZINC. To remove rust-spots from linen, oxalic acid or binoxalate of potash is commonly used. RUNGE recommends prussiate of potash. Dissolve a little in water, and just acidulate with sulphuric acid. The stained linen macerated in this water turns blue. Its whiteness is now restored by a solution of carbonate of potash.

SAXON BLUE. See CHEMIC BLUE. The solution of indigo in sulphuric acid, diluted with twice its weight of water, is so termed.

SCOURING DROPS FOR REMOVING GREASE. 1. Alcohol (pure) 6 oz., camphor 2 oz., rectified essence of lemon 8 oz.

2. Camphene 3 oz., essence of lemon 1 oz.; mix. Some direct them to be distilled together.

3. *French.* Camphene 8 oz., pure alcohol 1 oz., sulphuric ether 1 oz., essence of lemon 1 dr.

4. Spirit of wine a pint, white soap 3 oz., ox-gall 3 oz., essence of lemon $\frac{1}{4}$ oz.

SEALING-WAX. 1. *Blue.* Shellac 2 parts, dammar resin 2 parts, Burgundy pitch 1 part, Venice turpentine 1 part, artificial ultramarine 3 parts.

2. *Light Blue.* As the last, with 1 part of dry sulphate of lead.

3. *Dark Blue.* Venice turpentine 3 oz., finest shellac 7 oz., clear amber or black resin 1 oz., Prussian blue 1 oz., carbonate of magnesia $1\frac{1}{2}$ dr. The last two to be made into a stiff paste with oil of turpentine, and added to the melted shellac and Venice turpentine.

4. *Black.* 1. Venice turpentine $4\frac{1}{2}$ oz., shellac 9 oz., colophony $\frac{1}{2}$ oz., lamp-black mixed to a paste with oil of turpentine q. s.

2. Inferior. Venice turpentine 4 oz., shellac 8 oz., 3 oz. of colophony, and sufficient lamp-black mixed with oil of turpentine to color it.

3. Shellac 8 oz., Venice turpentine 4 oz., lamp-black 6 oz.

4. Common, for bottles. Resin 6 oz., shellac 2 oz., Venice turpentine 2 oz., lamp-black q. s.

Brown. 1. *Light Brown.* Venice turpentine 4 oz., shellac $7\frac{1}{2}$ oz., brown earth (English umber ?) $\frac{1}{2}$ oz., cinnabar

½ oz., prepared chalk ½ oz., carbonate of magnesia moistened with oil of turpentine 1½ dr.

2. *Light Brown*—Second quality. Venice turpentine 4 oz., shellac 7 oz., resin 3 oz., English umber 3 oz., cinnabar ¼ oz., prepared chalk 1 oz., magnesia as the last.

3. *Dark Brown.* Venice turpentine 4 oz., fine shellac 7½ oz., English umber 1½ oz., magnesia as before.

4. *Dark Brown*—Second quality. Venice turpentine 4 oz., shellac 7 oz., colophony 3 oz., English umber 1½ oz., magnesia as before.

Green. 1. Venice turpentine 2 oz., shellac 4 oz., colophony 1¼ oz., King's yellow ½ oz., Prussian blue ¼ oz., magnesia as for brown.

Gold. 1. Venice turpentine 4 oz., fine shellac 8 oz., leaf-gold 14 sheets, bronze powder ½ oz., magnesia (made into a paste with oil of turpentine) 1½ dr.

2. Use gold talc instead of gold leaf and bronze.—GRAY.

Marbled. Melt each colored wax separately, and just as they begin to grow solid, mix together.—GRAY.

Red. 1. *Fine Carmine Wax.* Venice turpentine 2 oz., finest shellac 4 oz., colophony 1 oz., Chinese vermilion 1½ oz., magnesia (moistened with oil of turpentine) 1½ dr.

2. *Finest Red.* Venice turpentine 4 oz., shellac 7 oz., cinnabar 4 oz., carbonate of magnesia (with oil of turpentine) 1½ dr.

3. As the last, with only 3½ oz. of cinnabar.

4. Venice turpentine 4 oz., shellac 6½ oz., colophony ½ oz., cinnabar 2½ oz., magnesia (with oil of turpentine) 1½ dr.

5. Venice turpentine 4 oz., shellac 6 oz., colophony ¾ oz., cinnabar 1¾ oz., magnesia as before.

6. As the last, but use colophony and cinnabar each 1½ oz.

7. Venice turpentine 4 oz., shellac 5½ oz., colophony 1½ oz., cinnabar 1¼ oz., magnesia as before.

8. *English.* Venice turpentine 2 oz., shellac 4 oz., vermilion 1 oz.

9. *Spanish.* Venice turpentine 8 oz., shellac 2 oz., colophony 4 oz., vermilion 1 oz. Remove from the fire, and add ½ oz. of rectified spirit.

Yellow. Venice turpentine 2 oz., shellac 4 oz., colophony 1¼ oz., King's yellow ¾ oz., magnesia as before.

Perfumed Wax. Add to any of the above a small quantity of fine benzoin.

Common Bottle Wax. 1. Dark resin 18 oz., shellac 1 oz., beeswax 1 oz. Mix together, and color with red lead, Venetian red, or lamp-black.

2. Resin 19 oz., beeswax 1 oz.; color as before.

SEA-WATER, ARTIFICIAL. See AQUARIUM.

SHELLAC, TO BLEACH. See LAC, page 388.

SILK CLEANER. Mix well together ¼ lb of soft soap, a teaspoonful of brandy, ½ pint of proof spirit, and ½ pint of water. It is to be spread with a sponge on each side of the silk without creasing it; the silk is then rinsed out two or three times, and ironed on the wrong side.

SILVER, TO CLEAN. See PLATE POWDER. To clean silver utensils, blackened by sulphuretted hydrogen, BOTTGER recommends a boiling saturated solution of borax, or a solution of caustic potash, with some fragments of metallic zinc.

SILVERING POWDER, &c., for silvering copper, covering the worn parts of plated goods, &c. 1. Nitrate of silver 30 gr., common salt 30 gr., cream of tartar 3½ dr. Mix. Moistened with water and rubbed on dial plates or other copper articles, it coats them with silver.

2. Silver precipitated from its nitric solution by copper 20 gr., alum 30 gr., cream of tartar 2 dr., salt 2 dr.

3. Precipitated silver ½ oz., common salt 2 oz., muriate of ammonia 2 oz., corrosive sublimate 1 dr.; make it into a paste with water. Copper utensils are previously boiled with tartar of alum, and rubbed with this paste, then made red-hot, and afterwards polished.

4. Dissolve muriate of silver in a solution of hyposulphite of soda, mix this with prepared hartshorn or other suitable powder.

SILVERING PASTE. Nitrate of silver 1 part, cyanide of potassium (Liebig's) 3 parts, water sufficient to form a thick paste. Apply it with a rag. A bath for the same purpose is made by dissolving 100 parts of sulphite of soda, and 15 of nitrate of silver, in water, and dipping the article to be silvered into it.

ELECTRO-SILVERING. Mr. WATT. 1. To the crystallized nitrate of silver obtained by dissolving 1 oz. of fine silver in nitric acid, add 3 pints of cold distilled water to dissolve it. Precipitate with common salt. Wash well. Dissolve the chloride in just enough of the strong solution of cyanide of potassium. Filter several times, then add distilled water to 1 gallon. This is excellent for a dead white, or to plate cleaned figures, or clock-dials, which only require to be scratch-brushed first. Or if worked weaker, with a feeble battery, and a small surface of *anode* exposed, a thick layer is obtained, which will bear burnishing.

2. Dissolve as much nitrate in the same quantity of distilled water. Add strong solution of cyanide of potassium till there is no further precipitate—(not enough to redissolve this). Wash the precipitated cyanide several times with pure water. Add now enough strong solution of cyanide of potassium to dissolve it, and make up the solution to 1 gallon. Filter before using. When it is desired that the articles should come out with a bright appearance, a little bisulphuret of carbon is added to the solution.

SILVER, OXIDIZED. Ornaments to which this name is given have a surface of silver which has been acted on by some chemical liquid. A solution of sal ammoniac will give a brownish tint; a still better one is obtained by using equal parts of sulphate of copper and sal ammoniac dissolved in vinegar. A fine black tint may be produced by a slightly warm solution of sulphuret of potassium. Solutions of the chloride of platinum and gold are sometimes adopted in these processes.

SILVERING GLASS. See GLASS.

SILVER, TO PURIFY AND REDUCE. Silver, as used in the arts and coinage, is alloyed with a portion of copper. To purify it, dissolve the mental in nitric acid slightly diluted, and add common salt which throws down the whole of the silver in the form of chloride. To reduce it into a metallic state several methods are used: 1. The chloride must be repeatedly washed with distilled water, and placed in a zinc cup; a little diluted sulphuric acid being added, the chloride is soon reduced. The silver when thoroughly washed is quite pure. In the absence of a zinc cup, a porcelain cup containing a zinc plate may be used. The process is expedited by warming the cup.

2. Digest the washed chloride with pure copper and ammonia. The quantity of ammonia need not be sufficient to dissolve the chloride. Leave the mixture for a day, then wash the silver thoroughly.—HORNUNG.

3. Boil the washed and moist chloride in solution of pure potash, adding a little sugar: when washed it is quite pure.

SILVER, SOLVENT FOR. See page 433.

SIZE. *Oil size* is made by grinding yellow ochre or burnt red ochre with boiled linseed oil, and thinning it with oil of turpentine. *Water size* (for burnished gilding) is parchment size ground with yellow ochre.

SMALTS. See PIGMENTS.

SKELETON LEAVES, to make. Steep the leaves, seed vessels, or other parts of the plant to be dissected, in rain-water, until the whole of the soft matter is decomposed. Some require a few weeks, others several months. The rotted parts are now to be carefully removed by a fine brush, under the surface of water, or in a stream of water. A syringe is sometimes required. To bleach the skeletons soak them for some hours in a mixture of 1 oz. of strong solution of chloride of lime and a quart of distilled water. Lastly, wash thoroughly in cold water, and dry by exposure to air.

SMOKING FLUID. One drop of creasote in a pint of water imparts a smoky flavor to fish or meat dipped into it for a few minutes.

SOAP. FOR PERFUMED AND TOILET SOAPS, see page 238. For the manufacture of soaps generally, see Dr. URE'S "Dictionary of the Arts," and other similar works. Hard soaps are made by boiling oils or fats with a lye of caustic soda. Soft soaps consist of oil and potash; and as they do not separate from the lye like the hard soaps, they generally contain an excess of caustic alkali. Silica soap has silicate of soda incorporated with it. Soap is adulterated by earthy matters, as pipe-clay, &c.; these and other impurities remain when soap is dissolved in alcohol.

SOAP, MARINE. *Patent.* This is made by substituting cocoa-nut oil for the fats and oils used in the manufacture of

common soap. It has the advantage of forming a lather with salt water.

SODA. For its medical and pharmaceutical compounds, see Pocket Formulary.

SODA, HYPOSULPHITE OF. Dissolve 1 lb of crystallized carbonate of soda in a quart of boiling water. Shake ½ lb of lime in another quart of water. Mix the solutions, let them stand in a covered vessel until cold, pour off the clear liquid, and boil it with more sulphur than it will dissolve. Pour off the clear solution into a deep vessel, and pass sulphurous acid gas through it until it becomes nearly colorless. While still a little yellow filter, and evaporate it quickly in an earthen vessel to a syrupy consistence. Shake this with half its bulk of rectified spirit, and allow the lower layer to crystallize under the alcoholic solution which floats on it. It must be kept from the air and light.

SOLVENTS FOR INDIA-RUBBER. *Ether* for this purpose should be agitated with water, and decanted. *Benzole* will dissolve caoutchouc with warmth and long digestion. *Rectified coal naphtha* forms an imperfect solution employed in MACINTOSH'S waterproof fabrics. *Oil of turpentine*, rendered pyrogenous by absorbing it with bricks of porous ware, and distilling it without water, and treating the product in the same way, is also used for this purpose. It is stated that the solution on evaporation does not leave the caoutchouc in a sticky state. Another method is to agitate oil of turpentine repeatedly with a mixture of equal weights of sulphuric acid and water; and afterwards expose it to the sun for some time. *Bisulphuret of carbon* is a good solvent, dissolving the gum without heat. This constitutes PARKE'S Patent Solvent. *Chloroform* is an excellent but rather expensive solvent. *Caoutchoucine* has also been employed as a solvent. It is prepared by distilling India-rubber without addition, increasing the heat to 600° Fahr. The product is rectified by distilling it with one-third of water. It is then a colorless fluid of 0·680 specific gravity. Its smell is improved by agitating it with 5 oz. of nitro-muriatic acid to each gallon. India-rubber is rendered more readily soluble by first digesting it with a solution of carbonate of soda, or water of ammonia.

SOLVENTS FOR GUTTA-PERCHA. Benzole readily dissolves it. So do chloroform and bisulphuret of carbon.

SOLVENT FOR OLD PUTTY AND PAINT. Soft soap mixed with solution of potash or caustic soda; or pearlash and slaked lime mixed with sufficient water to form a paste. Either of these laid on with an old brush or rag, and left for some hours, will render it easily removable.

SOLUBLE GLASS. See GLASS.

SOLUTIONS USED IN ELECTROTYPE MANIPULATIONS, &c.

For the Decomposing Cell. 1. Saturated solution of sulphate of copper 2 parts, sulphuric acid 2 parts, water 6 or 8 parts. 2. Mr. WALKER directs 2 measures of a saturated solution of sulphate of copper, and 1 measure of accidulated water (1 part of sulphuric acid to 9 of water). 3. ROBELL'S solution consists of 2 parts of a saturated solution of sulphate of copper, and 1 part of a saturated solution of Glauber's salt, to which as much sulphate of copper has been added as it will take up.

2. *Gold Solution.* Dissolve 2 oz. of cyanide of potassium (by LIEBIG'S method) in a pint of warm distilled water, add ¼ oz. of oxide of gold, and agitate together.

3. *Silver Solution.* Dissolve 2 oz. of LIEBIG'S cyanide of potassium in a pint of distilled water; add ¼ oz. of moist oxide of silver (precipitated by lime-water from a solution of the crystallized nitrate), and agitate together till the oxide is dissolved.

4. *Solution in which Steel Articles are dipped before Electroplating them.* Nitrate of silver 1 part, nitrate of mercury 1 part, nitric acid (sp. gr. 1.384) 4 parts, water 120 parts.

5. *Solution, or Pickle, for immersing Copper Articles in before Electroplating.* Sulphuric acid 64 parts, water 64, nitric acid 32, muriatic acid 1; mix. The article, free from grease, is dipped in the pickle for a second or two. See GILDING, SILVERING, ELECTRO-BRASSING, &c.

SOLUTION FOR MULLINS' VOLTAIC BATTERY. *In contact with the Zinc:* 1 part of sal ammoniac to 5 of water. *In contact with the Copper:* a saturated solution of sulphate of copper. Mr. BACHHOFFNER uses a saturated solution of common salt and a saturated solution of sulphate of copper.

SOLUTION FOR SOLDERING. Dissolve zinc in muriatic acid to saturation, and pulverized sal ammoniac, and boil for a short time. Applied with a sponge or feather it facilitates the flow of the solder.

SOLUTIONS FOR THE WATER-BATH. Various salts dissolved in water materially raise the boiling-point, and thus afford the means of obtaining a steady temperature at different degrees above 212°. The following are some of the most useful:

A saturated solution of	nitrate of soda	boils at 246°
"	"	Rochelle salts at . . 240
"	"	nitre at 238
"	"	muriate of soda at . 224
"	"	sulphate of magnesia at 222

SOLVENT FOR SILVER. *Nitro-sulphuric Acid.* Dissolve 1 part of nitre in 10 parts of oil of vitriol. Used for dissolving the silver from plated goods, &c. It dissolves silver at a temperature below 200°, and scarcely acts upon copper, lead, and iron, unless diluted. The silver is precipitated from the solution, after moderately diluting it, by common salt, and the chloride reduced as directed, page 428.

SPIRIT OF NITRIC ETHER. See P. F. It need only be added here, that its acidity is removed and prevented by rectifying it from neutral tartrate of potash.

SPONGE, BLANCHED. Soak the sponges for several days in cold water, renewing the water and squeezing the sponges occasionally. Then wash them in warm water, and place them in cold water to which a little muriatic acid has been added. Next day take them out and wash them thoroughly in soft water; then immerse them in an aqueous sulphurous acid (sp. gr. 1.034) for a week. They are afterwards washed in plenty of water, squeezed, and allowed to dry in the air. For burnt, prepared, and waxed sponge, see SPONGIA, P. F.

STAINS FOR WOOD, IVORY, &c. See IVORY, TO STAIN; WOOD, STAINS FOR; BOOKBINDER'S STAINS, &c.

STAINS, TO REMOVE. Stains of *iodine* are removed by rectified spirit. *Ink* stains by oxalic acid or superoxalate of potash. *Iron moulds* by the same; but, if obstinate, it has been recommended to moisten them with ink, then remove them in the usual way. See page 425.

Grease Spots. See SCOURING DROPS.

Red Spots on black cloth, from acids, are removed by spirits of hartshorn, or other solutions of ammonia.

Stains of Marking Ink or Nitrate of Silver, to remove. 1. Wet the stain with fresh solution of chloride of lime, and after 10 or 15 minutes, if the marks have become white, dip the part in solution of ammonia or of hyposulphite of soda. In a few minutes wash with clean water.

2. Stretch the stained linen over a basin of hot water, wet the mark with tincture of iodine.

3. They may also be removed by cyanide of potassium; but this should be done by the druggist, and not intrusted to any one else.

STARCH. Starch is procured from various roots and seeds. Its varieties are numerous; but a few of the most important only can be noticed here.

Arrowroot (West Indian). The fecula of the tubers of the Maranta arundinacea. The fresh tubes are washed and beaten to a pulp, which is well stirred in a large tub of cold clean water, and the fibrous part wrung out by the hands, and thrown away. The water in which the fecula is suspended is passed through a hair sieve or coarse cloth, allowed to settle, and the water poured off. After being repeatedly washed, the wet starch is drained, and afterwards dried in the sun. (The other varieties of arrowroot (see page 277) are prepared by analogous processes from the roots which yield them.)

Potato Starch. The tubers are washed and peeled, usually by machinery, rasped by a revolving grater, and the pulp washed on hair sieves till freed from the starchy matter. Successive portions of the pulp are thus treated till the vessel over which the sieves are placed is sufficiently full. The starch held in suspension in water subsides to the bottom; the water is then drawn off, and the starch stirred up with fresh water, and again allowed to subside. This is repeated several times till the starch is sufficiently pure. The fibres and the washing waters are used as manures. The washed fibres have also been recommended as an ingredient in bread for diabetic patients.

Wheat Starch. Wheat flour is steeped in water for a week or two and allowed to ferment. The acid liquor is drawn off, and the residue washed on a sieve: what passes

through is allowed to settle, the sour liquor drawn off, and the starch thoroughly washed from the slimy matter. It is then drained in perforated boxes, cut up into square lumps, placed on bricks to absorb the moisture, and dried in a stove. See Dr. PEREIRA's "Elements."

Various means are adopted to free the starch from gluten and other impurities. In the patent rice starch, and probably other kinds of starch, alkaline solutions are used. Ammonia has been recommended, as it does not, like potash and soda, dissolve any portion of pure starch.

The various kinds of fecula are distinguishable by the form of their particles or grains. By a microscopical examination of these the mixture or substitution of potato starch with the more expensive kinds is readily detected. Figures of the different kinds of starch grains are given i,n the "Pharmaceutical Journal," vol. iv., in Dr. PEREIRA s "Elements," &c. M. GOBLEY has proposed to distinguish them by the coloration produced when the several kinds are exposed to the vapor of iodine; but the effect seems to depend greatly on the relative dryness of the samples.

STEARINE, STEARIC ACID. Fat is saponified, and the soap decomposed by an acid, with a large quantity of water, the mixture being kept warm and well stirred. The water being drawn off, the fatty matter is well washed, allowed to cool, and submitted to strong pressure.

STORM GLASS. Take 2½ dr. of camphor, 38 gr. of nitre, and 38 gr. of sal ammoniac; dissolve them in 9 dr. of water, and 6 dr. of rectified spirit, with a gentle heat. It is placed in a glass tube covered with a brass cap, with a small hole to admit air. Or it may be put in an eau de Cologne or other long bottle, tied over with bladder. Its various changes are supposed to indicate changes of weather, but the indications are not to be relied on.

STUFFING BIRDS AND ANIMALS, PREPARATIONS FOR. 1. Camphor 1 oz., corrosive sublimate 1 oz., alum ½ oz., sulphur 1 oz.; all finely powdered and mixed.

2. Tanner's bark dried and powdered 2 oz., burnt alum 1 oz., snuff 1 oz.; mix, and add arsenic ¼ oz., camphor ¼ oz., sulphur 1 dr.

3. BECŒUR'S *Arsenical Soap*. Camphor 5 dr., arsenic 4 oz., white soap 4 oz., carbonate of potash 12 oz., air-slaked lime 4 oz.; make a stiff paste with a little water.

STYROL. Mix 20 parts of storax with 7 of carbonate of soda, and put them into a retort with water, and apply heat. A limpid fluid distils, which becomes, when heated to a certain point, a transparent solid.

SUGAR RESIN. Mix 16 parts of strong sulphuric acid with 8 of the strongest nitric acid; when cooled to 70° Fahr. stir in 1 part of finely-powdered sugar. In a few seconds, when the sugar has become pasty, take it out of the acid and plunge it into cold water. Add more sugar to the acid, and proceed as before. Wash the resinous matter carefully, and dissolve it in alcohol or ether. Evaporate the solution with a gentle heat. It is very combustible. Its solution may be used to render gunpowder, lucifer matches, &c., waterproof.

SULPHITE OF COPPER. To a concentrated solution of bisulphite of potash add a cold solution of sulphate of copper, filter, and heat gently.

SULPHOCYANIDE OF AMMONIUM. Saturate 2 parts of common water of ammonia (sp. gr. 0.950) with sulphuretted hydrogen; and add 6 parts of the same ammonia. To this mixture add 2 parts of sulphur, and the product of the distillation of 6 parts of prussiate of potash, 3 of sulphuric acid, and 18 of water. Digest till the sulphur is no longer acted on, and the liquid becomes yellow. Boil the liquid till it becomes colorless, filter, evaporate, and crystallize.

SULPHURET OF CARBON. See BISULPHURET OF CARBON.

SYRUP OF MILK. Evaporate, with constant stirring, 6 ℔ of skimmed milk to 3 ℔; add 4½ ℔ of sugar; dissolve with a gentle heat, and strain. It may be flavored with the addition of 1 oz. of cherry-laurel water. (For other Syrups, see P. F.) Milk may be preserved by first heating it, and, when cold, charging it with carbonic acid gas.

TANNIN. See ACID, TANNIC, page 307.

TERPINE. Leave oil of turpentine for a long time in contact with a mixture of nitric acid and alcohol. Crystals of terpine form. By boiling an aqueous solution of terpine with a small quantity of sulphuric or other acid, *terpinole* is formed, and may be separated by distillation. It has the odor of hyacinths.

TEST LIQUORS, TEST PAPERS, &c. Distilled water only should be used in these preparations. In preparing the papers, the liquid should be placed in an earthenware plate or dish, and the paper carefully immersed in it so as to be uniformly wetted, then dried out of the reach of acid, ammoniacal, or other vapors likely to affect it; and afterwards kept in bottles, jars or cases. Dr. FARADAY recommends unsized paper, but Mr. PARNELL and other good authorities direct good letter paper to be used.

Brazil Paper. Dip paper in a strong decoction of Brazil wood, and dry it. (It is rendered purple or violet by alkalies; generally yellow by acids.)

Cabbage Paper. Make a strong infusion of red cabbage leaves, strain it, and evaporate it by a gentle heat till considerably reduced. Then dip the paper in it and dry it in the air. (This paper is of a grayish color; alkalies change it to green, acids to red. It is a very delicate test; if rendered slightly green by an alkali, carbonic acid will restore the color.)

Dahlia Paper. From the petals of violet dahlias, as cabbage paper.

Elder-berry Paper. This is merely paper stained with the juice of the berries. Its blue color is changed to red by acids, and to green by alkalies.

Indigo Paper. Immerse paper in sulphate of indigo, wash it with water rendered slightly alkaline, then with pure water, and dry it in the air.

Iodide of Potassium and Starch Paper. Mix starch paste with solution of iodide of potassium, and moisten bibulous paper with it. (It becomes blue when exposed to ozone. Chlorine has the same effect.)

Lead Paper. Paper dipped in a solution of acetate of lead. (When moistened it detects sulphuretted hydrogen, which renders it black.)

Blue Litmus Paper. Bruise 1 oz. of litmus in a mortar, and add boiling water; triturate together, put them in a flask, and add boiling water to make up to half a pint; when cool, strain it, and dip paper in it. More color may be extracted from the litmus by hot water, but the liquid will require to be concentrated by evaporation. (Acids change

the color to red, but it does not become green with alkalies.)

Red Litmus Paper. As the last, adding to the strained infusion a few drops of nitric acid, or of pure acetic acid. Dr. FARADAY recommends holding blue litmus paper over a large jar, into which a few drops of muriatic acid have been introduced, till sufficiently reddened.

Rose Paper. Make a strong infusion of the petals of the red rose, and dip unsized paper in it. Dipped in an alkaline solution, so weak as not to affect turmeric paper, it assumes a bright green color.

Manganese Paper. Dip paper in a solution of sulphate of manganese. (It becomes black in an ozonized atmosphere.)

Rhubarb Paper. Dip paper in a strong infusion of rhubarb, and dry it. (Alkalies render it brown. It is not, like turmeric paper, affected by boracic acid.)

Starch Paper. This is merely paper imbued with starch paste. Cotton cord is sometimes used instead of paper. (As a test for iodine, which it turns blue.)

Turmeric Paper. Boil 1 oz. of coarsely-powdered turroot in half a pint of water for half an hour, and strain; dip paper in the liquid, and dry it. (It is rendered brown by alkalies, and also by boracic acid and borates.)

TEST SOLUTIONS, &c. (The vegetable preparations are here placed first.)

Tincture and Infusion of Red Cabbage. Digest red cabbage with rectified spirit in a warm place for a few days; strain, distil off most of the spirit, and evaporate what remains to the consistence of syrup. It will keep for years. When required for use, dilute it with a little water; or the concentrated infusion directed above for the paper may have a little spirit added to it. (If the cabbage leaves be well dried, they may be kept in a close vessel for use, and a strong infusion made when wanted.)

Acid Infusion of Red Cabbage. Dr. FARADAY directs one or more red cabbages to be cut up in strips, and boiling water poured on them, and a little dilute sulphuric acid (equal to ½ oz. of oil of vitriol to a large cabbage) to be

added, and the whole kept hot for an hour or two in a copper or earthen vessel. It is then strained, the cabbage infused in a little more water and acid, and the mixed infusion evaporated to one-third its first bulk, allowed to settle and put into bottles. When required for use, the acid is neutralized by caustic potash or soda. Another plan is to dry the leaves at 120°, and when required for use to make a strong infusion, adding a drop of sulphuric acid to neutralize the strained infusion with marble; filter, and add a little spirit, if required to be kept.

Infusion or Tincture of Litmus. This is made as directed above for litmus paper. Or an ounce of powdered litmus may be triturated with 6 oz. of boiling water, digested near the fire for an hour, and mixed, when cool, with 2 oz. of spirit. Or digest 1 oz. of powdered litmus in a pint of proof spirit for 7 days. If required red, a few drops of acetic acid are added to either of these. The next day, decant the clear liquor. Dr. PEREIRA directs 1 part of litmus to 25 of water. When made very strong, it must be diluted when used.

Tincture of Galls, Infusion of Galls, &c. Fresh powdered blue galls 1 oz., proof spirit 8 oz.; digest in a close vessel for a week, and filter. A watery infusion of galls may be made in the same proportion with boiling water for immediate use. PETTENKOFER directs 1 oz. of powdered galls to be infused in 3 or 4 oz. of boiling water for several hours, and 2 oz. of salt added. After filtration, it retains its transparency and power of precipitating gelatine for years. (This is used to detect iron, with the salts of which it produces a black color; for gelatine, which it precipitates in brownish-white flocks; and several of the organic alkaloids.)

MARSH'S *Dahlia Test.* Make a strong infusion of the petals of dark dahlias; strain, and add to every pint ½ oz. of strong sulphuric acid; stir with a glass rod, and when cold, add to each pint 2 gr. of corrosive sublimate. Filter through coarse cloth, and bottle. When required for use, neutralize it carefully with ammonia, and use the liquid by dipping paper in it.

Syrup of Violets. On 4 oz. of fresh petals of violets pour half a pint of water at 104° Fahrenheit, stir them together, and in a minute or two strain off the water with gentle

pressure, and pour 8 oz. of boiling distilled water on the flowers. In 12 hours, strain through linen, let the infusion settle, and decant, then dissolve in it twice its weight of refined sugar, by a gentle heat. (A delicate test for acids and alkalies.)

Dr. CLARK'S *Test for Hardness of Water.* Dissolve 1 oz. of Hawes's best white soap in a gallon of proof spirit. If not of such strength that it requires 32 measures to be added to 100 measures of solution of chloride of calcium of 16 degrees of hardness (see below) before it lathers, it must be adjusted to that strength. (The chloride of calcium solution is thus made: Dissolve 16 gr. of pure carbonate of lime in a small quantity of pure hydrochloric acid, avoiding loss for effervescence; evaporate the solution to dryness, and dissolve the residue in water, and again evaporate till a neutral solution is obtained; then dissolve in a gallon of water. This forms the standard solution of 16 degrees of hardness. One measure of this solution with 15 of distilled water constitutes a solution of one degree of hardness; and so on up to 16 degrees. The degree of hardness expresses the number of grains of carbonate of lime per gallon contained in the water. For the mode of using this test, see Dr. CLARK'S pamphlet, or PARNELL'S Chemical Analysis.)

Solution of Carbonate of Ammonia. Mr. PARNELL directs this test to be prepared by dissolving 1 part of sublimed carbonate of ammonia in 3 of water, and adding 1 part of water of ammonia.

Solution of Oxalate of Ammonia. Dissolve 1 oz. of crystallized oxalate of ammonia in a pint of water.

Solution of Sulphuretted Hydrogen. Pass sulphuretted hydrogen gas (see GASES, page 361) through cold distilled water, which has been recently boiled, till it will absorb no more. Keep it in small bottles securely closed.

Solution of Hydrosulphuret of Ammonia. Pass sulphuretted hydrogen gas (see page 361) through water of ammonia till the liquid occasions no precipitate in a solution of sulphate of magnesia.

Solution of Ammonio-nitrate of Silver. To a solution of nitrate of silver (1 part of crystallized nitrate to 20 of distilled water) add gradually weak water of ammonia till a mere trace of the oxide first precipitated is left undissolved.

Let it settle, decant it into a clean stoppered bottle, and keep it from the light. The Edinburgh Pharmacopœia directs 44 grains of nitrate of silver to be dissolved in a fluid ounce of water, and sufficient ammonia added as above. (It gives a pale yellow precipitate with arsenious acid, and a chocolate red with arsenic acid; the same with their salts.)

Solution of Nitrate of Silver. The Edinburgh Pharmacopœia directs this test to be prepared by dissolving 40 grs. of the nitrate in a fluid ounce of distilled water. The London Pharmacopœia directs 60 gr. to a fluid ounce. Mr. PARNELL recommends 1 part to 15 or 20 of water. (It is used chiefly for the detection of chlorine or muriatic acid. The precipitate, chloride of silver, is insoluble in nitric acid. Also for hydrocyanic acid, with which it gives a white precipitate, which is decomposed by heat, the silvering being reduced.)

Solution of Ammonio-sulphate of Copper. Dissolve 1 dr. of sulphate of copper in 2 oz. of water, and add ammonia till the precipitate first thrown down is nearly all dissolved. Let it settle, and pour off the clear solution. (Chiefly used as a test for arsenical compounds, with which it gives a green precipitate.)

Solution of Chloride of Barium. Dissolve 60 grains of the chloride in a fluid ounce of distilled water.

Solution of Nitrate of Barytes. (Ed. Ph.) Dissolve 40 grains of nitrate of barytes in 800 grains of distilled water.

Solution of Indigo. Digest 1 part of fine indigo in 10 parts of oil of vitriol, and dilute with water.

For the *Test Solutions* of the B. P. see P. F.

HAHNEMANN'S *Wine Test, for detecting lead in wine.* Sulphuret of lime 3 oz., tartaric acid 3 oz., water 2 lb, mix, decant and add 1 oz. of tartaric acid. Or, simple sulphuretted hydrogen water 4 oz., tartaric acid 1 dr.

TROMMER'S *Test* for sugar in urine. Put some of the suspected urine into a large test tube, and add a few drops of solution of sulphate of copper, then sufficient solution of potash to render it strongly alkaline. If sugar be present, the precipitated oxide redissolves into a blue liquid, and on boiling *red* oxide of copper is precipitated. (White merino

that has been wet with a solution of bichloride of tin is said to form a ready test for sugar in urine, &c. A portion wet with the suspected liquor, and exposed to 260° to 300° of heat, becomes blackened if sugar is present.) The following is proposed as a quantitative test for sugar: Dissolve 400 grains of pure crystallized sulphate of copper in 1600 grains of distilled water; add this gradually to a solution of 1600 grains of neutral tartrate of potash in a little water mixed with 6000 or 7000 grains of solution of caustic soda, of 1.12 sp. gr. Add water to make up the whole 11,544 grain measures (26 fl. oz., 2 fl. dr., 54 minims). 1000 grain measures are equivalent to 5 grains of grape sugar.

PETTENKOFER'S *Test* for bile, in urine, &c. Put a small quantity of the suspected liquid into a test-tube, and add to it, drop by drop, strong sulphuric acid till it becomes warm, taking care not to raise the temperature above 122° Fahr. Then add from 2 to 5 drops of syrup, made with 5 parts of sugar to 4 of water, and shake the mixture. If the liquid contain bile, a violet coloration is observed. Acetic acid, and those substances which are converted into sugar by sulphuric acid, may be substituted for sugar.

TOBACÇO WATER. See WASHES for vermin on plants.

TOUCH PAPER. Dip a piece of white blotting-paper, or printing-paper, in a solution of 1 oz. of nitre in 8 oz. of water. Dry it perfectly.

TREES, METALLIC. *Lead Tree.* Dissolve 1 oz. of sugar of lead in a quart of distilled or filtered rain-water, adding a few drops of acetic acid. Filter, and put the clear solution into a decanter or bottle. Suspend in it a piece of zinc, and set it aside.

Silver Tree. Dissolve 20 gr. of crystallized nitrate of silver in an ounce of distilled water; put into a vial, and add about ½ a drachm of pure quicksilver.

Tin Tree. Dissolve 3 dr. of muriate of tin in a pint and a half of water, with 10 or 15 drops of nitric acid; and suspend in it a rod of zinc.

TURPENTINE, VENICE (factitious). It is usually made by dissolving black resin in oil of turpentine. Dr. PEREIRA states the proportion to be 5 fluid oz. of the oil to 16 oz. of resin; but some makers put as much as 8, 10, or even 12

oz. of oil of turpentine to each pound of resin. (We have introduced this factitious preparation because no genuine Venice (or larch) turpentine is now to be obtained.)

TURPENTINE, OIL OF. Common turpentine, chiefly American, is distilled with water; the oil comes over with the water, and is found floating on it. It is rectified by distilling it again with water. See CAMPHENE and SOLVENTS FOR INDIA RUBBER, for further modifications of this oil.

URN POWDER. Oxide of iron, crocus, or jewellers' rouge.

VARNISHES. These constitute a distinct branch of manufacture, and many of them can be advantageously or safely made only on the large scale on premises adapted for the purpose. A few of the most easily prepared and useful varnishes have been selected for insertion. For fuller information see Dr. URE'S "Dictionary of Arts," DUMAS' "Chimie appliquée aux Arts, &c." Some practical information on this subject will be found in Mr. REDWOOD'S edition of GRAY'S Supplement, and in the 49th vol. of the Transactions of the Society of Arts.

Spirit Varnishes. The spirit employed should not be less than 60° overproof. In preparing and using them, they should be kept at a distance from a candle or other flame. Respecting the gums (resins) employed, it may be useful to mention that shellac is rendered more soluble by being powdered and exposed for a long time to the air; sandarach gives hardness to varnishes; mastic gives a gloss to a solution of other gums; benzoin still more, but its color is objectionable; anime readily dissolves, but renders the varnish long in drying; copal and amber are scarcely soluble in spirit, but are rendered partially so by other gums, and also by being previously fused by heat. Shellac gives a durable varnish, objectionable only on account of its color, which may be rendered paler by charcoal. See LAC.

1. *White Spirit Varnish.* Rectified spirit 2 gallons, gum sandarach 5 lb. Put them into a tin bottle, cork securely, and agitate frequently, placing the tin occasionally in hot water till the gum is dissolved, then add a quart of pale turpentine varnish.

2. *Brown.* Rectified spirit 2 gallons, sandarach 3 lb, shellac 2 lb, pale turpentine varnish a quart. Proceed as the last.

3. Sandarach 2 oz., shellac ½ oz., rectified spirit 16 fluid oz.

4. *White.* Gum sandarach 1½ oz., mastic ½ oz,, elemi ¼ oz., foreign oil of lavender ¼ oz., rectified spirit 8 oz.

Copal Spirit, or Drying Varnish. Copal, fused and pulverized, 3 oz., sandarach 6 oz., mastic 3 oz., Venice turpentine 2½ oz., highly rectified spirit a quart, powdered glass 3 oz. Mix the powdered glass and resins, and sift them; introduce them into a matrass with the spirit, and heat to boiling, constantly agitating till the gums are dissolved; then add the turpentine. Heat the varnish for half an hour, and when removed from the fire agitate till cold.

Brilliant Amber Spirit Varnish. Fused amber 4 oz., sandarach 4 oz., mastic 4 oz., highly rectified spirit a quart. Expose to the heat of a sand-bath, with occasional agitation, till dissolved. (The amber is fused in a close copper vessel, having a funnel-shaped projection, which passes through the bottom of the furnace by which the vessel is heated.)

Amber Varnish for Photographs. Dissolve 3 to 4 gr. of amber in 1 ounce of chloroform.

Colorless Varnish for Photographs. Dissolve shellac by heat in 8 parts of water and 1 of pearlash. Precipitate by chlorine, and dissolve in rectified spirit.—Dr. HARE.

Chinese Varnish. Mastic 2 oz., sandarach 2 oz., rectified spirit a pint. Close the matrass with bladder, with a pinhole for the escape of vapor; heat to boiling in a sand or water-bath, and when dissolved strain through linen.

Crystal Varnish. Picked mastic 4 oz., rectified spirit a pint, animal charcoal 1 oz. Digest and filter.

French Polish and *Lacquers* are varieties of spirit varnishes. The former has already been noticed (page 353). A few formulæ for the latter are here added:

Pale or Gold Lacquers. To a pint of rectified spirit add as much gamboge as will give it a bright yellow color, then add 12 oz. of seed-lac in fine powder, and set it in a sand-bath till dissolved. Or a tincture of annotto (1 part to 8 of spirit) may be added to give the desired color.

Dark Lacquer. Clear seed-lac 1 lb, dragon's blood 1 oz.;

pulverize together, and add them to a pint and a half of rectified spirit. Set in a warm place till dissolved.

Lacquer for Brass Work. Turmeric 1 oz., saffron ¼ oz., Spanish annotto ¼ oz., rectified spirit a pint. Digest at a gentle heat for several days; strain through coarse linen, put the tincture in a bottle, and add 3 oz. of good seed-lac coarsely powdered. Place in a moderate heat, and shake frequently till dissolved; if wanted of a redder shade, increase the quantity of annotto, or add a little dragon's blood. (Some makers prepare a strong tincture of the various coloring ingredients, and add them to the lacquer to produce the required shade.)

Oil of turpentine, and other essential oils, are used as solvents, forming essence varnishes, as the following:

Mastic Varnish. Clean mastic 5 oz., rectified oil of turpentine (camphene) a quart. Digest in a warm place, shaking frequently till the solution is complete, then strain.

Picture Varnish. Chio turpentine 2½ oz., mastic 12 oz., camphor ½ dr., pounded glass 4 oz., rectified oil of turpentine 3 pints. For oil paintings.

Canada Varnish. Clear Balsam of Canada 4 oz., camphene 8 oz.; warm gently, and shake together till dissolved. For maps, drawings, &c.; they are first sized over with a solution of isinglass, taking care that every part is covered; when dry, the varnish is brushed over it.

TINGRY'S *Essence Varnish.* Mastic in powder 12 oz., pure turpentine 1½ oz., camphor ½ oz., powdered glass 5 oz., rectified oil of turpentine a quart.

Common Turpentine Varnish. This is merely clear pale resin dissolved in oil of turpentine; usually 5 ℔ of resin to 7 ℔ of turpentine.

Oil Varnishes. These consist of copal and other gums dissolved by heat in boiled linseed oil; generally with the addition of oil of turpentine.

Cabinet Varnish. Fuse 7 ℔ of African copal, and pour on it 4 pints of hot clarified linseed oil (see OILS); in 3 or 4 minutes, if it feels stringy, take it out of the building, where there is no fire near, and when it has cooled to 150° mix in 3 gallons of oil of turpentine of the same temperature, or sufficient to bring it to a due consistance. (Various qualities of copal varnish are made for different purposes;

inferior gums are often substituted for or mixed with copal.)

Amber Varnish. Amber 16 oz.; melt in an iron pot, and add ½ pint of drying linseed oil, boiling hot, and then 3 oz. of resin and 3 oz. of asphalte, each in fine powder. Stir till they are thoroughly incorporated; remove from the fire, and add a pint of warm oil of turpentine.

Common Oil Varnish. Resin 3 lb, drying oil ½ a gallon; melt together, and add, when removed from the fire, 2 quarts of warm oil of turpentine.

Varnish for Printers' Ink. To every 10 lb of clarified linseed oil (page 389) add 5 lb of clear black resin, and ½ lb oil of turpentine. It is then ready for mixing with lamp-black or other coloring matter. A twelfth part of Canada balsam is sometimes added for the finer parts.

A few miscellaneous varnishes are added.

Varnish for Engraving on Copper. Yellow wax 1 oz., mastic 1 oz., asphaltum ½ oz.; melt, pour into water, and form into balls for use. A softer varnish for engravers is made with 1 part of tallow and 2 of yellow wax; or with 2 oz. of wax, 1 dr. of common turpentine, and 1 dr. olive oil. See ETCHING VARNISHES, page 347.

Varnish for Engraving on Glass. 1. Wax 1 oz., mastic ½ oz., asphaltum ¼ oz., turpentine ½ dr.

2. Mastic 15 parts, turpentine 7, oil of spike 4 parts.

LE BLOND'S *Varnish.* Keep 4 lb of balsam of copaiva warm in a sand or water-bath, and add 16 oz. of copal, previously fused and coarsely powdered, by single ounces daily, and stir it frequently; when dissolved add a little Chio turpentine.

BESSEMER'S *Varnish*, for metallic paint. This is made with 8 lb of copal, 2½ gallons of drying oil, and 25 gallons of oil of turpentine. These are made into a varnish nearly as directed for Cabinet Varnish; and afterwards mixed with a gallon of slaked lime, and left for 3 days to settle. The clear portion is then drawn off, and 5 parts of varnish mixed with 4 parts of bronze powder.

MACINTOSH'S *Caoutchouc Varnish.* Dissolve 1 lb of India-rubber cut in shreds in a quarter of a pint of rectified coal naphtha. (Caoutchouc varnishes may be made with any of the solvents noticed above, page 431. The following are also used:)

India-Rubber Varnish, for boots. Dissolve ¼ oz. of caoutchouc in 2 oz. of mineral naphtha. Dissolve also ½ oz. of asphaltum in 1 oz. of oil of turpentine. Mix the solutions.

Balloon Varnish. Melt India-rubber in small pieces with its weight of boiled linseed oil, and thin it with oil of turpentine.

Varnish for Frames for Hot Beds. Mix 4 oz. of pulverized white cheese, 2 oz. of slaked lime, and 4 oz. of boiled linseed oil. Mix, and add 4 oz. each of whites and yolks of eggs, and liquefy the mixture by heat. This curious mixture is said to produce a pliable and transparent varnish.

Colored Varnishes. Oil varnishes are colored by grinding with them the most transparent colors, as distilled verdigris for green, &c. Spirit varnishes are also colored with dragon's blood, gamboge, &c.

Sealing-Wax Varnish. Black or colored sealing-wax broken small, and sufficient rectified spirit to cover it, digested till dissolved. An article called black lac is sold as an economical substitute for black sealing-wax.

Black Japan for Leather, &c. Boil together a gallon of boiled linseed oil, 8 oz. of umber, and 3 oz. of asphaltum. When sufficiently cool, thin in with oil of turpentine.

Japan for Tin-ware. 1. Common Copal varnish. 2. Dissolve copal 2 oz., and camphor 1 dr., in oil of turpentine 8 oz.

Brunswick Black. Melt 4 ℔ of asphaltum, and 2 ℔ of hot boiled linseed oil, and when sufficiently cool add a gallon of oil of turpentine.

Varnish for Gun Barrels, after browning them. Shellac 1 oz., dragon's blood ¼ oz., rectified spirit a quart. Dissolve and filter.

Transfer Varnish. Alcohol 5 oz., pure Venice turpentine 4 oz., mastic 1 oz.

Hair Varnish. Dissolve 1 part of clippings of pigs' bristles, or of horsehair, in 10 parts of drying linseed oil by heat. Fibrous materials (cotton, flax, silk, &c.), imbued

with the varnish and dried, are used as a substitute for haircloth.

Glass Varnish. This is a solution of soluble glass, and should be thus made: Fuse together 15 parts of powdered quartz (or of fine sand), 10 parts of potash, and 1 of charcoal. Pulverize the mass, and expose it for some days to the air; treat the whole with cold water, which removes the foreign salts, &c. Boil the residue in 5 parts of water until it dissolves. It is permanent in the air, and not dissolved by cold water. *Used to protect wood, &c., from fire.*

Vegetable Parchment. De la Rue's Patent. Strong unsized paper is immersed for a few seconds in oil of vitriol, diluted with half its volume of water. It is then washed in pure water. It strongly resembles animal parchment, and is used for the same purpose. (The acid solution must be exactly of the strength indicated, and not warmer than the air around.)

Vinegar. Vinegar may be made from wine or ale, by keeping it for some weeks or months in a warm place, with access of air. In this country it is usually made from malt, or a mixture of malted and unmalted barley, which is mashed as for beer, and fermented with yeast. The fermented liquor is then placed in a warm room for many weeks in unclosed casks, and finished by transferring it into large vessels with false bottoms, on which are placed the refuse raisins, &c., from which wine has been prepared. A much quicker method of acetification is sometimes employed: the fermented liquor is made to pass in drops into tubs filled with beech chips, so as to expose an extended surface to the action of the air. In Germany it is also made by the direct acetification of spirit by means of platina black. The method of preparing wood vinegar has already been noticed. (See Pyroligneous Acid.) The following is one of the processes followed in making vinegar from sugar: Boil 10 gallons of water for 10 minutes with a quart of bran; run it into a tub through a flannel, and put into it 12 lb of coarse brown sugar, and when cooled to 70° add a quart of yeast at three different times. Let it work for four days, then take off the yeast, and run the liquor into a clean tub. Fill the tub nearly with the liquor, leaving room for 2 lb of bruised crab apples and 1 lb of raisins. If it ferments, add

a little reserved liquor, or water boiled with sugar, till the fermentation ceases. Then place the cask upon a plank fronting the sun in summer, and near the fire in winter. Put into it 1 oz. of isinglass well beaten up with a quart of old vinegar, and cover the bunghole with a piece of hop-bag (fastened to the edge of the hole by pitch), and lay a tile over it. Leave it in this state till it becomes fit for use. On a small scale, Dr. TURNER states that vinegar may be made from 120 parts of water, 12 of brandy, 3 of brown sugar, 1 of tartar, and ½ of sour dough, left some weeks in a warm place.

WAFERS, GELATINE. Dissolve fine glue or isinglass in such a quantity of water as that the solution, when cold, may be consistent. Pour it hot on a plate of mirror glass (previously warmed with steam and slightly greased), which is fitted in a metallic frame, having edges just as high as the wafers should be thick. Lay on the surface a second glass plate, also hot and greased, so as to touch every point of the gelatine while resting on the edges of the frame. By its pressure the thin cake is rendered uniform. When the glass plates have cooled, the gelatine will be solid, and may be removed. It is cut into discs of different sizes by means of proper punches.

WASHES FOR VERMIN IN PLANTS. 1. *Tobacco Water.* Infuse 1 ℔ of tobacco in a gallon of boiling water, in a covered vessel, till cold.

2. *For Lice in Vines.* Boil ½ ℔ of tobacco in 2 quarts of water; strain, and add ½ ℔ of soft soap and ¼ ℔ of sulphur; mix.

3. *For Aphides.* Boil 2 oz. of lime and 1 oz. of sulphur in water, and strain.

4. *For Red Spiders.* A teaspoonful of salt in a gallon of water. In a few days wash the plant with pure water. See BLIGHTS, remedies for.

WASHING POWDERS. These consist of soda-ash combined with gelatinous substances, as a solution of glue, linseed jelly, &c., dried and powdered.

WASHING LIQUIDS are chiefly solutions of caustic soda.

WATER FOR MARINE AQUARIUM. See p. 313.

WATER, HARD, Dr. CLARK'S patent for softening. This consists simply in adding milk of lime to the water in the

reservoir. It combines with free carbonic acid, which it precipitates as carbonate of lime, and at the same time causes the deposition of the carbonate of lime previously held in solution by that gas.

WATERPROOFING COMPOUNDS. *For Boots, &c.* 1. (ROOME'S patent.) Suet 8 oz., linseed oil 8 oz., yellow beeswax 6 oz., neatsfoot oil 1½ oz., lamp-black 1 oz., litharge ½ oz. Melt together, and stir till cold.

2. Linseed oil 8 oz., boiled ditto 10 oz., suet 8 oz., yellow wax 8 oz. Melt.

3. Dr. HARVARD'S. Wax 8 oz., resin 4 oz., mutton suet 4 oz.; boil together, and apply warm to new boots.

4. Col. HAWKER'S. Drying oil 1 pint, wax 2 oz., Burgundy pitch 1 oz., oil of turpentine 2 oz. Melt over a slow fire, and add a few drops of oil of lavender or thyme. Brush the boots repeatedly with the composition before the fire, till they appear fully saturated.

5. *For Leather, &c.* Cut 3 drachms of India-rubber into small pieces, soak them for 24 hours in a solution of common soda; dissolve this and 3 oz. of asphaltum in 12 oz. of camphene, then add ½ oz. of boiled linseed oil.

For Cloth. It is alternately dipped in a solution of acetate of lead with a little gum, and solution of alum.

For Hats. Boil 8 lb shellac, 3 lb frankincense, and 1 lb borax in sufficient water.

For Canvas, &c. Gutta percha 3 parts is dissolved in resin spirit 9 parts, at a heat of 120° to 140° Fahr., stirring occasionally.—Mr. CASTLEY.

WAX. Yellow beeswax is bleached by pouring the melted wax in a divided state on a revolving cylinder partly immersed in water, so as to form it into fine ribbons, which are exposed to air and moisture till bleached, and subsequently refined by melting with water containing sulphuric acid.—Dr. PEREIRA. It has been proposed to bleach wax by adding to each pound of melted wax 2 oz. of powdered nitrate of soda, and afterwards stirring in, by little at a time, 1 oz. of sulphuric acid diluted with 10 parts of water, keeping the mixture warm, and constantly stirred with a glass rod in a capacious earthen vessel, till all the acid is added. It is then allowed to become somewhat cool, and the vessel filled with boiling water, well agitated, and set aside. The cake of wax is removed into boiling water, till this no longer produces a precipitate with chloride of barium.—M. IN-

GENHOL. (We have not found this render wax perfectly white.)

WAX FOR MODELING. Lead plaster 8 oz., beeswax 8 oz., Burgundy pitch 8 oz.; melt together, stir in sufficient chalk to form a paste, and form it into small sticks for use. (For SEALING-WAX, see page 426.)

WELDING COMPOSITION. Mix borax with $\frac{1}{10}$th of sal ammoniac, fuse the mixture, and pour it on an iron plate. When cold, pulverize it, and mix it with an equal weight of quicklime, sprinkle it on iron heated to redness, and replace it in the fire. It may be welded below the usual heat.

WHEAT, STEEP FOR. A pound of genuine sulphate of copper in sufficient water, for each sack of seed. Arsenic is also used; sulphate of zinc has been recommended; so has quicklime, which is thus used: Soak the seed in a warm mixture of 36 to 48 oz. of quicklime to 6 or 7 gallons of water. This is for $4\frac{1}{2}$ bushels of wheat: the solution should be sufficient to cover the seed 3 or 4 finger-breadths deep, and it should lie in it 24 hours. M. BOUSSINGAULT has proposed to sprinkle the grain, first with water, then with a mixture of 10 parts of lime with 1 of white arsenic. (This has the double object of preserving the grain, and destroying a plague of field mice.) Sulphate of copper has seemed to give satisfactory results. It would be desirable, however, to find an innocuous substitute, as traces of copper have been found in wheat grown from the steeped seed. This appears to have been discovered in the use of a solution of sulphate of soda with lime, which has proved more successful in France than either arsenic or sulphate of copper. See *Blights, Remedies for.*

WOOD, TO STAIN. 1. *Mahogany color* (*dark*). Boil $\frac{1}{2}$ lb of madder and 2 oz. of logwood in a gallon of water; and brush the wood well over with the hot liquid, When dry, go over the whole with a solution of 2 dr. of pearlash in a quart of water.

2. (*Light.*) Brush over the surface with diluted nitrous acid, and when dry apply the following with a soft brush: Dragon's blood 4 oz., common soda 1 oz., spirit of wine 3 pints; let them stand in a warm place, shaking frequently, then strain. Repeat the application until the proper color is obtained.

3. (*To stain Maple a Mahogany color.*) Dragon's blood ½ oz., alkanet ¼ oz., aloes 1 dr., spirit of wine 16 oz. Apply it with a sponge or brush.

4. *Rosewood.* Boil 8 oz. of logwood in 3 pints of water until reduced to half; apply it boiling hot two or three times, letting it dry between each. Afterwards put in the streaks with a camel-hair pencil dipped in a solution of copperas and verdigris in decoction of logwood.

5. *Ebony.* Wash the wood repeatedly with a solution of sulphate of iron; let it dry, then apply a hot decoction of logwood and nutgalls for two or three times. When dry, wipe it with a wet sponge; and when again dry, polish with linseed oil.

6. *To stain Wood Red.* Use a strong decoction of Brazil wood and alum. (Woods may be stained with the various dyes before described. See DYES.)

YEAST, ARTIFICIAL. Honey 5 oz., cream of tartar 1 oz., malt 16 oz., water at 122° F. 3 pints; stir together, and when the temperature falls to 65°, cover it up, and keep it at that temperature till yeast is formed.

ZINC, AMMONIO-CHLORIDE OF. By dissolving equal equivalents of chloride of zinc and sal ammoniac, a crystallizable salt is formed, which dissolves oxides of copper and of iron, and is useful in tinning or zincing those metals.

ZINC, AMALGAMATED (*for voltaic plates*). Put a little mercury on the zinc plate, and pour on it dilute sulphuric acid; then rub the mercury over the surface by means of a piece of linen. Another method, which is said to give a more permanent coating, is that of Mr. WALENN. Having cleared the plates by emery, and by immersion in diluted sulphuric acid, and then in clean water, dip them into a mixture of equal parts of a saturated solution of corrosive sublimate and a similar solution of acetate of lead; then rub them with a cloth.

ZINC, PLATINIZED, *for Dr. Wright's Battery.* Saturated solution of chloride of platinum ½ dr, sulphuric acid ½ dr., water 2 dr. Mix; dip the zinc plates into the solution for a few seconds, and wash them quickly.

ZINC, OXIDE OF. It may be prepared from the purified sulphate by precipitating it from a *hot* solution by carbonated or bicarbonated alkalies. It cannot be obtained

pure by caustic ammonia: M. J. LEFORT. Mr. MIDGLEY prepares it on a large scale by the combustion of zinc in a muffle, heated by a furnace of peculiar construction; the zinc is introduced into the muffle from time to time, as the combustion proceeds; he is thus able to prepare one or two hundred-weights at a time, by a continuous process.

ZINC, PURIFICATION OF. Granulate zinc by melting it, and pouring it while very hot into a deep vessel filled with water. Place the granulated zinc in a Hessian crucible, in alternate layers, with one-fourth its weight of nitre, with an excess of nitre at the top. Cover the crucible, and secure the lid; then apply heat. When deflagration takes place, remove from the fire, separate the dross, and run the zinc into an ingot mould. It is quite free from arsenic.

APPENDIX.

WEIGHTS AND MEASURES.

Apothecaries' Weight. (L. P.)

℔ Pound.		℥ Ounces.		ʒ Drachms.		℈ Scruples.		Gr. Grains.		Minims of water.
1	=	12	=	96	=	288	=	5760	=	6319.54
		1	=	8	=	24	=	480	=	526.62
				1	=	3	=	60	=	65.82
						1	=	20	=	21.94
								1	=	1.09

The Troy Pennyweight, 24 grains, is not used in compounding medicines.

Apothecaries' Measure.

C. Congius. Gallon.		O. Octarii. Pints.		f℥ Fluid Ounces.		fʒ Fluid Drachms.		♏ Minims.		Grains of water.
1	=	8	=	160	=	1280	=	76800	=	70000
		1	=	20	=	160	=	9600	=	8750
				1	=	8	=	480	=	437.5
						1	=	60	=	54.7
								1	=	0.9

Imperial Measure.—(*Common Divisions.*)

Quarter.		Bushels.		Pecks.		Gallons.		Quarts.		Pints.		Gills.
1	=	8	=	32	=	64	=	256	=	512	=	2048
		1	=	4	=	8	=	32	=	64	=	256
				1	=	2	=	8	=	16	=	64
						1	=	4	=	8	=	32
								1	=	2	=	8
										1	=	4

Avoirdupois Weight. (B. P.)*

lb Pound		oz. Ounces.		gr. Grains.		French Grammes.
1	=	16	=	7000	=	453.544
		1	=	437.50	=	28.346

Other weights used are the ton, 20 hundred-weight; the hundred-weight, 112 lb.; and the quarter, 28 lb.

Table for Converting Troy into Avoirdupois Weight.

(From Dr. Duncan's "Edinburgh Dispensary.")

Troy ounces.		Avoirdupois ounces.	grains.	Troy ounces.		Avoirdupois ounces.	grains.
1	=	1	42½	7	=	7	297½
2	=	2	85	8	=	8	340
3	=	3	127½	9	=	9	382½
4	=	4	170	10	=	10	425
5	=	5	212½	11	=	12	30
6	=	6	255	12	=	13	72½

175 Troy ounces are equal to 129 Avoirdupois.

Troy lb		Avoirdupois lb	oz.	gr.	Troy lb		Avoirdupois lb	oz.	gr.
1	=	0	13	72½	18	=	14	12	430
2	=	1	10	145	19	=	15	10	65
3	=	2	7	217½	20	=	16	7	137½
4	=	3	4	290	30	=	24	10	425
5	=	4	1	362½	40	=	32	14	275
6	=	4	14	435	50	=	41	2	125
7	=	5	12	70	60	=	49	5	412½
8	=	6	9	142½	70	=	57	9	262½
9	=	7	6	215	80	=	65	13	112½
10	=	8	3	287½	90	=	74	0	400
11	=	8	0	360	100	=	82	4	250
12	=	9	13	432½	175	=	144	0	0
13	=	10	11	67½	200	=	164	9	62½
14	=	11	8	140	300	=	246	13	312½
15	=	12	5	212½	400	=	293	2	125
16	=	13	2	285	500	=	411	6	375
17	=	13	15	359½	1000	=	822	13	312½

* The Dublin College of Physicians have adopted the avoirdupois weight in their last Pharmacopœia (1850), but they divide the oz. into 8 drachms, and the drachm into 3 scruples, as in Troy weight.

The following are the divisions of the *old wine gallon* adopted in the London Pharmacopœia before 1836, and the Dublin Pharmacopœia before 1850. Its use in this kingdom is no longer legal.

Former Apothecaries' Measure.

C.		O.		f℥		f℈		f℥		Minims.
1	=	8	=	128	=	1024	=	3072	=	61440
		1	=	16	=	128	=	384	=	7680
				1	=	8	=	24	=	480
						1	=	3	=	60
								1	=	20

Comparison between the Old and New Measure.

	Grains of distilled water.		Cubic Inches.	
	OLD.	NEW.	OLD.	NEW.
Gallon, . .	58317.8	70000	231	277.274
Pint, . . .	7289.7	8750	28.875	34.659
f℥j . . .	455.6	437.5	1.804	1.733
f℥j . . .	56.9	54.7	.225	.216

The old gallon was very nearly $\frac{5}{6}$ths of the new; the new is $\frac{6}{5}$ths of the old. The exact factor for converting the old measure into new is .83311; and for converting new into old, 1.20032.

Relative Value of the Former and Present Apothecaries' Measure.

(From the American Dispensatory.)

OLD.		NEW.				NEW.		OLD.				
		O.	f℥	f℥	℥			C.	O.	f℥	f℥	℥
Cong.	=	6	13	2	23	Cong.	=	1	1	9	5	8
O.	=		16	5	18	O.	=		1	3	1	38
f℥	=		1	0	20	f℥	=				7	41
f℥	=			1	2½	f℥	=					58

To find the weight of any given measure of a liquid, multiply the weight of the water it will contain by the specific gravity, water being 1.000. The weight of a gallon of any liquid, in avoird. lbs. and decimal parts, is at once seen from its density, merely removing the decimal point one place to the right. Thus a gallon of ether at .750 weighs 7.50 (7½) lb. A gallon of nitric acid at 1.500 weighs 15 lb.

TABLE

SHOWING THE RELATIONS OF THE WEIGHTS AND MEASURES OF VARIOUS LIQUIDS.

	Specific Gravity.	A Fluidounce weighs	Imperial Pint weighs	Troy Ounce measures		Avoirdupois Ounce measures		A Gallon weighs in Avoirdupois	
		Grains.	Grains.	f℥	ℿ	f℥	ℿ	℔	oz.
Water (distilled),	1.000	437½	8750	8	46	8	0	10	0
Alcohol. L. 1836,	.815	356½	7131	10	46	9	49	8	2⅛
Alcohol. E.,	.796	348	6964	11	2	10	3	7	15⅜
Rectified Spirit,	.838	366½	7332½	10	28	9	33	8	6
Proof Spirit,	.920	402½	8050	9	31	8	42	9	3 3/16
Proof Spirit. E. 1841,	.912	399	7980	9	37	8	46	9	1⅛
Chloroform,	1.480	647½	12950	5	56	5	24	14	12¾
Ether,	.750	328½	6562½	11	42	10	40	7	8
Spirit of Nitric Ether,	.834	365	7297½	10	31	9	35	8	5 7/16
Olive Oil,	.9153	400½	8009	9	35	8	44	9	2 7/16
Syrup (*Normal.* GUIBOURT),	1.320	577½	11550	6	39	6	4	13	3¼
Sulphuric Acid. L.,	1.843	807	16144	4	45	4	20	18	7 3/16
Nitric Acid. L.,	1.420	621¼	12425	6	11	5	38	14	3 3/16
Nitric Acid. E., B., & D.,	1.500	656¼	13125	5	51	5	20	15	0
Muriatic Acid,	1.160	507½	10150	7	35	6	54	11	9 9/16

Weights and Measures of other Countries.

In the United States of America the weights are the same as in this country; but they have not adopted our imperial measure, and retain the old wine gallon and its divisions in the Pharmacopœia.

The unit of the British India ponderary system is the tola, equal to 180 Troy grains. 32 tolas are equal to ℔j Troy. The maund is equal to 100 Troy ounces.

In France the metrical or decimal system is now the only legal one. The following table shows the correspondence of the French metrical weights with English grains.

		Troy Grains.			Troy Grains.
Milligramme	=	.0154	Decagramme	=	154.34
Centigramme	=	.1543	Hectogramme	=	1543.40
Decigramme	=	1.5434	Kilogramme	=	15434.00
Gramme	=	15.4340	Myriagramme	=	154340.00

The measures of capacity in France are multiples and divisions of the LITRE, which is the measure occupied by a kilogramme (15434 Troy grains) of distilled water at its greatest density. It exceeds the old Paris *pinte* by $\frac{1}{14}$th, and is equal to 35 fluid ounces and 103 minims, or 1.7608 imperial pints, or 61.028 English cubic inches. 4½ litres makes an imperial gallon, within about f℥xij.

The following table will show the relations between the litre and the imperial gallon of 277.2738 c. inches:

Litres.			Cubic Inches.	Gall.	Pts.	Fl. ℥	Fl. ʒ	Min.
$\frac{1}{1000}$	=	Millilitre	.061028					16.9
$\frac{1}{100}$	=	Centilitre	.61028				2	49
$\frac{1}{10}$	=	Decilitre	6.1028			3	4	10.36
1	=	Litre	61.028		1	15	1	43.69
10	=	Decalitre	610.28	2	1	12	1	16.9
100	=	Hectolitre	6102.8	22	0	1	4	49
1000	=	Kilolitre	61028.	220	0	16	6	40
10000	=	Myrialitre	610280.	2201	(or 275⅛ bushels.)			

French Measures of Length.

The standard unit is the *metre*, equal to 39.371 English inches, or 1 yard 3 inches and $\frac{37}{100}$ths. The *kilometre* (1000 metres) is 4 furlongs 213 yards 1 foot 11 inches.

The following are some of the weights and measures formerly used in France.

The old French pound, *livre poids de marc*, was equal to 489.5 grammes, or 7561 Troy grains; but the metrical pound, *livre métrique*, substituted for it in 1812, contained exactly 500 grammes,

or 7717 English grains. Both are now abolished. The following are their divisions:

Livre.	Once.	Cros.	Scrupl.	Grs.	*Poids de Marc.*		*Métrique.*
1	16	118	384	9261 =	grammes	489.5	500
	1	8	24	576 =	"	30.6	31.25
		1	3	72 =	"	3.824	3.90
			1	24 =	"	1.274	1.30
				1 =	"	.053	.054

In the Paris Codex and medical works the grain is represented by 0.05 grammes (5 centigrammes), 2 grains by 0.1 (1 decigramme); the half gramme by 2 grammes; the drachm by 4 grammes; and the ounce by 32 grammes.

The old French measures used in pharmacy were:

		Litres.	Other Commercial Measures.		Litres.
La pinte	=	0.931	8 Pintes (un velte)	=	7.450
La Chopine	=	0.466	13.97 (ancien boisseau)	=	13.010
Le demi-Setier	=	0.233	288 = 1 muid	=	268.220
Le Poisson	=	0.116	576 = 1 tonneau d'Or-		
Le demi-Poisson	=	0.058	léans, ou muids	=	536.440

(From Guibourt's "Pharmacopée Raisonnée.")

The litre, with its divisions and multiples, is the measure now used. It contains 1000 grammes of water; the number of grammes of other liquids corresponds with their specific gravity; water being 1000.

The former measures of length in France were the

Toise = 1.949 metres, or 6.3945 English feet.
Foot (pied) = 0.32484 metres = 12.785 English inches.
Inch (pouce) = 0.02707 metres, or 1.0654 English inches.
Line (ligne) or $\frac{1}{12}$th of an inch = .002256 metres.
The metre is equal to 3 ft. 11 lines old French measure, or 3 ft. 3.7 in. English.

Other Foreign Weights and Measures.

1.—*Medicinal pounds of* 12 *ounces, in English grains.*

(From Jourdan's "Pharmacopée Universelle.")

The following are divided as our Apothecaries' weight:

The pound of Austria weighs 6482.42 grains; Bavaria, 5556.24; Holland, 5787.75; Lubec, 5697.09; Nuremberg (German pound), 5522.96; Poland, 5533.25; Prussia, 5113.99; Sweden, 5497.01; Venice (*sottile*), 4649.17.

The division of the following differs in the scruple being divided into 24 grains:

Bologna, 5026.32; Lucca, 5162.67; Modena, 5254.61; Parma, 5062.35; Portugal, 5312.23; Rome, 5233.25; Spain, 5325.84; Tuscany, 5240.49; Piedmont (Turin), 5123.49.

The Naples pound contains 5490.63 Troy grains; the ounce is divided into 10 drachms; the scruple into 20 grains.

2.—*Various Foreign Weights.*

The old Paris pound was divided into 16 ounces; the scruple into 24 grains. Its weight has been given above. The pound by which drugs are weighed in Turkey is the *Tchegy*, equal to 4957 English grains, and is divided into 100 drachms, each drachm into 16 killos, and each killo into 4 grains.

The *obolo* is half a Spanish scruple; 3 *silicua* make 1 *obolo*, and 4 grains a *silicua*.

A *loth*, in Germany, Poland, &c., is half an ounce.

The commercial pound in several countries differs from the pharmaceutical. The civil pound of Bavaria and mark of Vienna are each about 19¾ avoirdupois ounces. That of Holland is the French kilogramme, or 12 grains more than 2 ℔ 3¼ oz. avoirdupois. The mark is half a kilogramme. The Coburg commercial pound is nearly 18 oz. avoirdupois.

3.—*Foreign Measures.*

The Austrian *mass* or *kanne* is equal to 1.415015 litres, or 2½ imperial pints, within 40 minims.

The *kanna* of Sweden = nearly 2.62 litres, or about 4 pints, 12 ounces, imperial.

Russian pound of water = 25.019 English cubic inches.

The *pott* (half kanne) of Denmark = 0.9653 litres.

The *arroba* of Spain = 16.073 litres.

The *almude* of Portugal = 16.451 litres.

The Prussian quart = 1.145 litres; or 1 qt. 3 fl. dr. imp.

The *barile* of Naples = 43.6216 litres; of Rome, 58.5416 litres; of Tuscany, 45.584 litres.

The *wedro* of Russia (10 *stof* or 30 Russian pounds) = 12.29 litres, or 21 pints, 12 oz., 12¼ dr. imperial.

The *mass* of Wurtemburg = 1.537 litres, or about 3 pints, 14¾ oz. imperial.

COMPARISON OF THERMOMETRIC SCALES.

To convert the degrees of Centigrade into those of Fahrenheit, multiply by 9, divide by 5, and add 32.

To convert degrees of Centigrade into those of Reaumur, multiply by 4 and divide by 5.

To convert degrees of Fahrenheit into those of Centigrade, deduct 32, multiply by 5, and divide by 9.

To convert degrees of Fahrenheit into those of Reaumur, deduct 32, divide by 9, and multiply by 4.

To convert degrees of Reaumur into those of Centigrade, multiply by 5 and divide by 4.

To convert degrees of Reaumur into those of Fahrenheit, multiply by 9, divide by 4, and add 32.

In De Lisle's thermometer, used in Russia, the graduation begins at boiling-point, which is marked *Zero*, and the freezing-point is 150.

EFFECTS OF TEMPERATURE.

Deg. of Fahr.	
8696	Cast Iron melts (Morveau).
2200	Gold melts (Kane), 2518 (Morveau).
1996	Copper melts (Kane), 2548 (Daniell).
2233	Silver melts (Daniell).
1869	Brass melts (Daniell).
1000	Iron bright cherry red (Poillet).
980	Red heat, visible in daylight (Daniell).
941	Zinc begins to burn (Daniell).
793	Zinc melts (Gmelin), 648 (Daniell).
644	Mercury boils (Daniell), 662 (Graham).
630	Whale oil boils (Graham).
612	Pure lead melts (Parkes), 609 (Daniell).
600	Linseed oil boils.
545	Sulphuric acid boils (Phillips), 620 (Graham).
518	Bismuth melts (Gmelin), 476 (Phillips).
442	Tin melts.
380	Arsenious acid volatilizes.
356	Metallic arsenic sublimes.
315	Oil of turpentine boils (Kane).
302	Ætherification ends.
256	Sat. sol. of acetate of soda boils.
257	" sal ammoniac boils (Taylor).
248	" nitric acid 1.42 boils, and sol. soda 1.44.
238	" nitre boils.
232	Sulphur melts (Turner), 226 (Fownes).
221	Sat. sol. of salt boils (Paris Codex).
220	" alum, carb. soda, and sulph. zinc boil.
218	" chlorate and prussiate of potash boil.
216	" sulph. of iron, sulph. of copper, nitrate of lead, boil.
214	" acetate of lead, sulph. and bitartrate of potash, boil.

Deg. of Fahr.

213 Water begins to boil in glass (or 213.5).
212 Water boils in metal, barometer at 30°.
211 Alloy of 5 bismuth, 3 tin, 2 lead, melts.
201 " 8 bismuth, 5 lead, 3 tin, melts (Kane).
194 Sodium begins to melt.
185 Nitric acid 1.52 boils.
180 Starch dissolves in water.
176 Rectified spirit boils, benzole distils.
173 Alcohol (sp. gr. .796 to .800) boils.
151 Beeswax melts (Kane), 142 (Lepage).
150 Pyroxylic spirit boils (Scanlan).
140 Chloroform, and ammonia of .945, boil.
136 Potassium melts (Daniell).
132 Acetone (pyroacetic spirit) boils (Kane).
122 Mutton suet and styracine melt.
116 Bisulphuret of carbon boils (Graham).
115 Pure tallow melts (Lepage), 92 (Thomson).
112 Spermaceti and stearine of lard melt.
99 Phosphorus melts.
98 Ether (.720) boils. Temperature of the blood.
88 Acetous fermentation ceases, water boils *in vacuo*.
77 Vinous ferm. ends, acetous ferm. begins.
62 Oil of anise liquefies, congeals at 60.
59 Gay Lussac's *Alcoomètre* graduated at.
55 Syrups to be kept at.
42 Sulphuric acid, sp. gr. 1.741, congeals (or 41).
36 Olive oil freezes.
32 Water freezes.
30 Milk freezes.
28 Vinegar freezes.
20 Wine freezes.
0 Cold produced by snow and salt.
—7 Brandy freezes.
—39–40 Mercury freezes.

Gentle Heat of London and United States pharmacopœias, and *Inferior Heat* of Dub. ph., from 90 to 100°
Medium Heat, Dub. ph., from 100 to 200°
Superior Heat, Dub. ph., from 200 to 212°
Maceration, Dub. ph., to be performed at 60 to 90°
Digestion, Dub. ph., at 90 to 100°
Specific Gravities, taken at 60° Lond. ph.; at 62° Ed. ph.

SPECIFIC GRAVITIES.

1. *Solids. Water*, 1.000.

Platinum, 20.980; Gold, 19.250; Mercury, 13.560; Lead, 11.350; Silver, 10.500; Bismuth, 9.822; Copper, 8.895; Cadmium, 8.604; Cobalt, 8.538; Nickel, 8.279; Iron, 7.788; Tin, 7.291; Zinc, 6.861 to 7.100; Antimony, 6.720; Sodium, .972; Potassium, .865; Caoutchouc, .933; Amber, 1.078; Scammony, 1.210; Resin, 1.072; Camphor, .988; Gum arabic, 1.452; Beeswax, .962; Spermaceti, .943; Sulphur, 1.990; Glass, 2.540 to 2.953.

2. *Liquids. Water*, 1.000.

Mercury, 13.560; Sulphuric acid, 1.843; Nitric acid, (mono-hydrated), 1.517; Pure nitric acid (Ed. and Dub. ph.), 1.500; Nitric acid (Lond. ph.), 1.420; Commercial nitric acid, 1.380 to 1.390; Double aqua fortis, 1.360; Single aqua fortis, 1.220; Muriatic acid (strongest), 1.210; Muriatic acid, Lond. ph., 1.160; Ditto, Ed. ph., 1.170; Ditto, Dub. ph., 1.176; Solution of caustic potash, Lond. ph. (containing 6.7 per cent.), 1.063; Ditto. Dub. ph., 1.068; Ditto, Ed. ph., 1.072; Water of ammonia, Lond. and Ed. ph. (containing 10 per cent.), .960; Ditto, Dub. ph. (12.7 per cent.), .950; Stronger water of ammonia, Lond. ph. (32 per cent.), .882; Ditto, Ed. ph. (33 per cent.), .880; Ditto, Dub. ph. (27 per cent.), .900; Saturated solution of alum, 1.033; Saturated solution of common salt, 1.200; Saturated solution of sulphate of copper, 1.150; Saturated solution of sulphate of magnesia, 1.218; Sea-water, 1.027; Milk, 1.032; Alcohol, .976; Rectified spirit, .838; Proof spirit, .920; Chloroform, 1.496 (not less than 1.480, Lond. ph.); Bisulphuret of carbon, 1.272; Spirit of nitric ether, Lond. ph., .834; Ed. ph., .847; Ether (pure), .724; Lond. and Dub. ph., not above .750; Ed. ph., 735; Acetic ether, .917; Caoutchoucine, .680; Oil of turpentine, .876 to .869; Olive oil, .9175; Spermaceti oil, .890; Southern whale oil, .920; Almond oil, .917; Creasote, 1.046; Oil of wine, 1.05; Essential oil of anise, .985; of caraway, .938; of cinnamon, 1.008; of cloves, 1.055; of cajeput, .911; of lemon, .847; of rosemary, .897; Tincture of sesquichloride of iron, .992; Tincture of opium, .952.

3. *Gases and Vapors. Atmospheric Air*, 1.000.

Hydrogen, .690; Nitrogen, .972; Oxygen (Graham), 1.1056; Carbonic oxide, .972; Carbonic acid, 1.524; Light carburetted hydrogen, .5595; Olefiant gas, .981; Chlorine, 2.470 (2.421 Graham); Vapor of ether, 2.582; v. of water, .620; v. of sulphur, 6.648 (6.617 Graham); v. of phosphorus, 4.327; v. of iodine, 8.700.

Table of Chemical Elements, with their Symbols and Equivalent Numbers.

Elements.	Symbol.	Hydrogen 1.		Oxygen 100.
		Brande.	Turner.	Berzelius.
Aluminum,	Al.	14	13.7	171.17
Antimony [Stibium],	Sb.	125	64.6	806.45
Arsenic,	As.	75	37.7	470.04
Barium (Phillips, 68),	Ba.	69	68.7	856.88
Bismuth,	Bi.	213	71.	886.92
Boron,	B.	11	10.9	136.25
Bromine,	Br.	78	78.4	978.31
Cadmium,	Cd.	56	58.8	696.77
Calcium,	Ca.	20	20.5	251.94
Carbon,	C.	6	6.12	75.40
Cerium,	Ce.	46	46.	574.70
Chlorine,	Cl.	36	35.42	444.65
Chromium,	Cr.	28	28.	351.82
Cobalt,	Co.	30	29.5	368.99
Columbium [Tantalium],	Ta.	185	185.	2307.43
Copper [Cuprum],	Cu.	32	31.6	395.70
Fluorine,	F.	19	18.68	233.80
Glucinum (Phillips, 27),	G.	5	26.5	331.26
Gold [Aurum],	Au.	200	199.2	2486.03
Hydrogen,	H.	1	1.	12.4795
Iodine (Liebig, 127.1),	I.	126	126.3	1579.50
Iridium,	Ir.	99	98.8	1233.50
Iron [Ferrum],	Fe.	28	28.	350.27
Lead [Plumbum],	Pb.	104	103.6	1294.50
Lithium (Graham, 6.43),	L.	7	6.	80.33
Magnesium,	Mg.	12	12.7	158.35
Manganese,	Mn.	28	27.7	345.89
Mercury (Phillips, 200),	Hg.	100	202.	1265.82
Molybdenum,	Mo.	48	47.7	598.52
Nickel (Phillips, 30),	Ni.	28	29.5	369.68
Nitrogen,	N.	14	14.15	177.04
Osmium,	Os.	100	99.7	1244.49
Oxygen,	O.	8	8.	100.00
Palladium (Phillips, 53),	Pd.	54	53.3	655.90
Phosphorus (Schrötter, 31),	P.	32	15.7	196.14
Platinum,	Pl.	99	97.8	1233.50
Potassium [Kalium],	K.	40	39.15	489.92
Rhodium,	R.	52	52.2	651.39
Selenium,	Se.	40	39.6	494.58
Silicium, or Silicon,	Si.	15	22.5	277.31
Silver [Argentum],	Ag.	108	108.	1351.61
Sodium [Natrium],	Na.	24	23.3	290.90
Strontium,	Sr.	44	43.8	547.29
Sulphur,	S.	16	16.1	200.80
Tellurium,	Te.	64	64.2	801.76
Thorium,	Th.	60	59.6	744.90
Tin [Stannum],	Sn.	59	57.9	735.29
Titanium,	Ti.	24	24.3	303.66
Tungsten [Wolfram],	W.	100	99.7	1246.25
Vanadium,	V.	68	68.5	856.86
Uranium,	U.	60	217.	2711.36
Yttrium,	Y.	32	32.2	402.51
Zinc,	Zn.	32	32.3	403.23
Zirconium (Phillips, 22),	Zr.	23	33.7	920.20

Remarks.—The first column of figures shows the equivalent numbers adopted by Prout, Brande, Dumas, Phillips, and other chemists, who regard all equivalent numbers as simple multiples of that of Hydrogen, which they adopt as unity.

In the second column are those of Turner and others, who adopt Hydrogen as unity, but do not consider that the other equivalents are exact multiples of it.

In the third column are the numbers of Berzelius, &c., who adopt oxygen as unity, or rather as 100.

Besides the difference in the numbers arising from the adoption of a different standard, there are others which arise from discordant views of the composition of the compounds from which the equivalents are determined. Some Continental chemists also make a distinction between the equivalents and *Atomic weights*, supposing the equivalents of certain elements to consist of two atoms. But this distinction is not adopted by English chemists.

The newly discovered metals, Didymium, Erbium, Lanthanum, Niobium, Norium, Pelopium, Ruthenium, and Terbium, require further investigation. The following numbers on the Hydrogen Scale have been assigned to some of them: Lanthanum 48 (Graham), Didymium 49.6 (Marignac), Ruthenium 52.

Composition, Equivalent Numbers, and Symbols, of some of the more important Compounds employed in Pharmacy and the Arts.

[The numbers here adopted for the elements are those of the first column of the preceding table.]

	Symbols.	Equiv.
Acetone	$C_3 H_2 O$	29
Acid Acetic (anhydrous)	$C_4 H_3, O_3$ (*or* $\bar{A}$)	51
— Acetic, glacial	$\bar{A}$, HO	60
— Arsenious	$As O_3$	99
— Arsenic	$As O_5$	115
— Benzoic	$C_{14} A_5 O_3$	113
—— Crys.	$C_{14} H_5 O_3$, HO	122
— Boracic	$Bo O_3$	35
—— Crys.	$Bo O_3$, HO	62
— Carbonic	$C O_2$, (*or* $\ddot{C}$)	22
— Chromic	$Cr O_3$	52
— Citric (dry)	$C_{12} H_5 O_{11}$	165
—— Crys. (commercial)	$\bar{C}$, 4 HO	201
— Hydrochloric	H Cl	37
— Hydrocianic	$C_2 N H$ (*or* H Cy)	27
— Hydrosulphuric	S H	17
— Iodic	$I O_5$	166
— Nitric (dry)	$N O_5$	54
— Nitric, sp. gr. 1.5	$N O_5$, HO	63
— Nitrous	$N O_4$	46
— Oxalic (dry)	$C_2 O_3$ (*or* $\bar{O}$)	36
— Oxalic, crys.	$C_2 O_3$, 3 HO	63
— Phosphoric	$P O_5$	72
— Sulphuric	SO_3	40
— Sulphuric, liquid	SO_3, HO	49

	Symbols.	Equiv.
Acid Tartaric	$C_4 O_5 H_2$ (*or* $\overline{T}$)	66
— Tartaric, crys.	$\overline{T}$, HO	75
— Chromic	$Cr O_3$	52
— Alcohol	$C_4 H_6 O_2$	46
Alumina	$Al_2 O_3$	52
Alum, crys.	$\left\{ \begin{matrix} Al_2 O_3, (SO_3); KO, \\ SO_3 + 24 HO \end{matrix} \right\}$	476
Ammonium	NH_4	18
Amidogen	NH_2	16
Ammonia	NH_3	17
— Hydrochlorate	NH_3, H Cl	54
— Sulphate, crys.	NH_3, SO_3, 2 HO	75
— Sesquicarb.	2 NH_3, 3 CO_3, 2 HO	118
— hydrated bicarb.	NH_3, 2 CO_2, 2 HO	79
Antimony, sesquioxide	$Sb O_3$	153
— Sesqui- *or* ter-sulph	$Sb S_3$	177
— potassio-tartrate	KO, $\overline{T}$, $Sb O_3$, $\overline{T}$, 3 HO	360
Baryta	Ba O	77
— Carbonate	Ba O, CO_2	99
— Sulphate	Ba O, SO_3	117
Barium, chloride	Ba Cl	105
Bismuth, oxide	$Bi O_3$	237
— Nitrate (dry)	$Bi O_3$, NO_5	291
Borax	Na O, 2 BO_3, 10 HO	192
Calcium, chloride	Ca Cl	56
— Chloride, crys.	Ca Cl, 6 HO	110
— Oxide (Lime)	Ca O	28
Chloroform	C_2 H Cl_3	121
Cinchonia	$C_{20} H_{12} NO$	154
Copper, oxide	Cu O	40
— Dinoxide	Cu_2 O	72
— Sulphate	Cu O, SO_3	80
—— Crys.	Cu O SO_3, 5 HO	125
— Nitrate	Cu O, NO_5	94
—— Crys.	Cu O, NO_5, 3 HO	121
— Acetate, crys.	Cu O, $\overline{A}$, HO	100
Cyanogen	C_2 N (*or* Cy)	26
Ether	$C_4 H_5$ O	37
Ethule	$C_4 H_5$	29
Glycerine, anhydrous	$C_6 H_7 O_5$	83
Gum	$C_{12} H_{11} O_{11}$	171
Iron, protoxide	Fe O	36
— Black oxide	$Fe_3 O_4$	116
— Sesquioxide	$Fe_2 O_3$	80
—— (Phillips)	Fe $O_{1½}$ (*or* ½ $Fe_2 O_2$)	40
— Chloride	Fe Cl	64
— Sesquichloride	$Fe_2 Cl_3$	164

	Symbols.	Equiv.
Iron, Sesquichloride (Phillips)	Fe $Cl_{1½}$ (*or* ½ Fe_2 Cl_3)	82
— Iodide (dry)	Fe I	154
— Sulphate	Fe O, SO_3	76
— Sulphate, crys.	Fe O, SO_3, 7 HO	139
— Potassio-tartrate	$\bar{T}$ KO, Fe_2 O_3, $\bar{T}$	260
Lead, acetate, crys.	Pb O, $\bar{A}$, 3 HO	190
— Diacetate	2 (Pb O) $\bar{A}$	275
— Protoxide	Pb O	112
— Carbonate	Pb O, CO_2	134
— Iodide	Pb I	230
— Chloride	Pb Cl	140
— Sulphate	Pb O, SO_3	152
— Nitrate, crys.	Pb O, NO_5	166
Lime, carbonate	Ca O, CO_2	50
— Hydrate	Ca O, HO	37
— Sulphate	Ca O, SO_3	68
— Phosphate (bone-earth)	3 Ca O, PO_5	156
Magnesia	Mg O	20
— Com. carbonate	5 Mg O, 4 CO_2, 6 HO	242
— Sulphate	Mg O, SO_3	60
— Sulphate, crys.	Mg O, SO_3, 7 HO	123
Manganese, binoxide	Mn O_2	44
— Chloride	Mn Cl	64
— Sulphate	Mn O, SO_3	76
Mannite	C_6 H_7 O_6	91
*Mercury, chloride	Hg Cl	236
— Bichloride	Hg Cl_2	272
— Protoxide	Hg O	208
— Binoxide	Hg O_2	216
— Iodide	Hg I	326
— Biniodide	Hg I_2	452
— Ammnio-chloride	Hg, Cl, NH_2	252
— Nitrate (proto-)	Hg O, NO_5	262
— Nitrate, crys.	Hg O, NO_5, 2 HO	280
— Bipersulphate	Hg O_2, 2 SO_3	296
— Bisulphuret	Hg S_2	232
Morphia, dry	H_{20} C_{35} NO_6	292
— Crys.	H_{20} C_{35} NO_6, 2 HO	310
— Acetate, crys.	H_{20} C_{35} NO_6, $\bar{A}c$, HO	352
— Muriate, crys.	H_{20} C_{35} NO_6, H Cl, 6 HO	383

* Mr. Brande has now adopted 100 as the equivalent of Mercury, with Dr. Kane and others with the B. P.; but we have here retained the old symbols and equivalents, which are those of Phillips, as the change would necessitate an alteration in the pharmacopœial and established names of the several mercurial compounds. Two must be added to each of these numbers, if mercury be 202, as stated by Turner. The ammonio-chloride, containing 2 eq. of mercury, will be 526.

	Symbols.	Equiv.
Morphia, Sulphate, crys...	$H_{20} C_{35} NO_6$, SO_3, 6 HO	386
Platinum, chloride	Pl Cl_2	171
— Ammonio-bichloride	Pl Cl_2, NH_3, H Cl	225
Potash, dry	KO	48
Potash, hydrate	KO, HO	57
— Acetate, dry	KO, $\bar{A}$	99
— Carbonate	KO, CO_2	70
— Carbonate, hydrated	KO, CO_2, 1½ HO	83½
— Bicarbonate, crys.	KO, 2 CO_2, HO	101
— Chlorate	KO, Cl O_5	124
— Chromate	KO, Cr O_3	100
— Bichromate	KO, 2 Cr O_3	152
— Nitrate	KO, NO_5	102
— Sulphate	KO, SO_3	88
— Bisulphate, crys.	KO, 2 SO_3 + 2 HO	146
— Tartrate	KO, $\bar{T}$	114
— Bitartrate	KO, 2 $\bar{T}$, HO	189
Potassium, bromide	K Br	118
— Chloride	K Cl	76
— Iodide	K I	166
Proteine	$C_{46} H_{36} N_6 O_{14}$	520
Quinia	$C_{20} H_{12} NO_2$	162
— Disulphate, crys.	2 ($C_{20} H_{12} NO_2$,) SO_3, 8 HO	436
— Neutral sulphate	$C_{20} H_{12} NO_2$, SO_3, HO	274
— Hydrochlorate	2 ($C_{20} H_{12} NO_2$,) H Cl, 3 HO	338
Silver, oxide	Hg O	116
— Chloride	Ag Cl	144
— Iodide	Ag I	234
— Cyanide	Ag Cy	134
— Nitrate	Ag O, NO_5	170
Soda	Na O	32
— Carbonate	Na O, CO_2	54
— Carbonate, crys.	Na O, CO_2, 10 Ho	144
— Bicarbonate	Na O, 2 CO_2, HO	85
— Sesquicarbonate	Na O, 1½ CO_2, 2 HO	83
— Sulphate, dry	Na O, SO_3	72
— Sulphate, crys.	Na O, SO_3, 10 HO	162
— Bisulphate	Na O, 2 SO	112
— Phosphate	2 Na O, PO, 25 HO	361
— Potassio-tartrate, crys.	Na O, KO, 2 $\bar{T}$, 10 HO	302
Sodium, chloride	Na Cl	60
Starch (anhydrous)	$C_{12} H_{10} O_{10}$	162
Strychnia	$C_{44} H_{24} O_4 N_2$	348
Sugar (anhydrous)	$C_{12} H_9 O_9$	153
— Crys.	$C_{12} H_{11} O_{11}$	171
Sulphuretted hydrogen	SH	17
Veratria	$C_{34} H_{22} NO_6$	288

	Symbols.	Equiv.
Water	HO	9
— Oxygenated	HO_2	17
Zinc, oxide	Zn O	40
— Carbonate	Zn O, CO_2	62
— Sulphate, crys.	Zn O, SO_3, 7 HO	143
— Chloride	Zn Cl	68

TABLE OF THE NEUTRALIZING PROPORTIONS OF SOME OF THE ACIDS AND ALKALINE CARBONATES, OMITTING MINUTE FRACTIONS. THE BEST COMMERCIAL PREPARATIONS ARE INTENDED.

Tartaric Acid.	Citric Acid.	Lemon-Juice.	Cr. Subcarb. of Soda.	Sesqui. or Bicarb. of Soda and Sub-Carb. of Potash.	Bicarbonate of Potash.	Carbonate of Magnesia.	Sesquicarbonate of Ammonia.	Bicarbonate of Ammonia.
Grs.	Grs.	3	Grs.	Grs.	Grs.	Grs.	Grs.	Grs.
10	9¼	2⅙	19	11	13½	6½	8¼	10½
10¾	10	2⅓	20½	12	14½	7	8½	11½
13	12	2⅘	25	14½	17⅙	8¼	10	13½
15	14	3¼	29	17	20⅓	9½	12	16
15½	14½	3⅓	30	17¼	21	10	12¼	16½
18	17	4	34½	20	24¼	11½	14	19
20	18½	4⅓	38½	22⅓	27	12¾	15½	21
20½	19	4⅕	40	23	27½	13	16	21½
26	24	5⅕	50	29	35	16½	28¼	27
27	25	5⅚	52	30	36	17	21	28½
32	30	7	61	36	43	20½	25	33½
36	33½	7⅚	69	40	48½	23	28	38
47	44	10¼	90	52½	63	30	37	49½
52	48½	11⅓	100	58	70	33	41	55
62	58	13½	120	69	84	40	49	65½
73	68	15⅝	140	82	98	46½	57	77
75	70	16¼	144	84	101	48½	59	79
90	84	19½	172	101	121	57½	71	94½
92	86	20	177	103	124	59	72	97
100	93	21⅜	192	112	134	64	78	105½
108	100	23⅓	206	120	145	69	84	113
180	168	39⅕	344	202	242	115	141	190

We have estimated the equivalent of subcarbonate of potash (Potassæ carbonas of the Pharmacopœia), and of sesquicarbonate of soda, at 84 each. Mr. PHILLIPS makes the former 83.5; but .5 may be allowed for impurity and extra moisture; the sesquicarbonate of soda he makes 83, but the composition of the best commercial specimens approaches nearer to the bicarbonate, which is 85.

Table of the Relation between the Principal Areometers for Liquids Lighter than Water.

[The first five columns are from Soubeiran, the last from Dr. Christison and Mr. Redwood. The degrees of Gay-Lussac's alcohometer indicate the percentage by measure of pure alcohol; but are not quite exact as here given, the fractions being neglected.]

Baumé.	Cartier.	Pharm. Batava.	Specific Gravity.	Gay-Lussac.	Sykes.
					Under proof.
10	10	0	1000	0	100
11	10.92	1	993	5	93.5
12	11.84	2	987	10	83.5
13	12.76	3	979	17	73.5
14	13.67	4	973	23	62
15	14.59	5	966	29	51
16	15.51	6	960	34	42
17	16.43	7	953	39	32
18	17.35	8	947	43	25
19	18.26	9	941	47	20
20	19.18	10	935	50	14
21	20.10	11	929	53	8
22	21.02	12	923	56	3
					Over proof.
23	21.94	13	917	59	2
24	22.85	14	911	61	7.5
25	23.77	15	905	64	11.5
26	24.69	16	900	66	15
27	25.61	17	894	69	20
28	26.53	18	888	71	25
29	27.44	19	883	73	28.5
30	28.38	20	878	75	32
31	29.29	21	872	77	38
32	30.31	22	867	79	41
33	31.13	23	862	81	44
34	32.64	24	857	83	45.5
35	32.96	25	852	84	48
36	33.88	26	847	86	51
37	34.80	27	842	88	54
38	35.72	28	837	89	56.5
39	36.63	29	832	91	59
40	37.65	30	827	92	61.5
41	38.46	31	823	93	
42	39.40	32	818	94	
43	40.31	33	813	96	
44	41.22	34	809	97	
45	42.14	35	804	98	
46	43.06	36	800	99	
47	43.19	37	795	100	
48	44.90	38	791		

SPECIFIC GRAVITIES CORRESPONDING WITH THE DEGREES OF BAUMÉ'S AREOMETER FOR LIQUIDS HEAVIER THAN WATER. —[*Pharmacopœia Batava.*]

Degrees.	Sp. gr.	Degrees.	Sp. gr.
0	1000	39	1372
1	1007	40	1384
2	1014	41	1398
3	1022	42	1412
4	1029	43	1426
5	1036	44	1440
6	1044	45	1454
7	1052	46	1470
8	1060	47	1485
9	1067	48	1501
10	1075	49	1516
11	1083	50	1532
12	1091	51	1549
13	1100	52	1566
14	1108	53	1583
15	1116	54	1601
16	1125	55	1618
17	1134	56	1637
18	1143	57	1656
19	1152	58	1676
20	1161	59	1695
21	1171	60	1715
22	1180	61	1736
23	1190	62	1758
24	1199	63	1779
25	1210	64	1801
26	1221	65	1823
27	1231	66	1847
28	1242	67	1872
29	1252	68	1897
30	1261	69	1921
31	1275	70	1946
32	1286	71	1974
33	1298	72	2000
34	1309	73	2031
35	1321	74	2059
36	1334	75	2087
37	1346	76	2116
38	1359		

INDEX.

Observe: For Medicines for Horses, Cattle, &c., see the VETERINARY INDEX *at the end.*

Abernethy's Pills, 166
Abernethy Biscuits, 286
Acetate of alumina, 301
iron, 301
lime, 302
oxide of amyle, 347
soda, 302
Acetic acid, 302
perfumes, 226
fumigation, 357
Acetimetry, 302
Acid, acetic, 302
carbonic, 357
chloric, 303
chromic, 303
cinnamic, 303
fluoric, 303
hippuric, 304
iodic, 304
muriatic, 304
nitric, 305
nitro muriatic, 305
oxalic, 305
phosphoric, 305
pyrogallic, 306
sulphuric, 306
sulphuric, dry, 306
sulphurous, 307
tannic, 307
Acidimetry, 308
Acids for galvanic batteries, 307
Acids and alkaline carbonates, table of the neutralizing proportions of some, 470
Acidulated kali, 272
effervescing powders, 273
Acidulated lemonade, 271
lemonade powder, 271
syrup of raspberries, 275
syrup of strawberries, 275
Acton's solution for baldness, 244
Ægyptiacum, 107
Aerated chalybeate, 210
ginger-beer, 268
ginger-beer powders, 269
lemonade, 271
lemonade powders, 271
orangeade, 272
orangeade powder, 273
sherbet, 272
sherbet powder, 273
waters, 205, 211
waters, alkaline, 205
waters, chalybeate, 210
lithia water, 206
magnesia, 205
saline, 206
Ague drops, 166
Aix-la-Chapelle water, 209
Albespeyres paper, 196
Albumen process, 404
photographic, 404
Albuminous size, 308
Alcohol, 308
Alibert's cosmetic, 233
Alkalimetry, 308
Alkaline lotion for the mouth, 263
Alkaline solution, Brandish's, 172
Alloys, various, 309
Almond flavor, 292
paste, 235
powder, 236

Almond honey paste, 236
soap, 238
bloom, 238
Alum, 310
mordants, 343
Aluminium, frosted, 311
bronze, 311
Amalgams, 309
for teeth, 266
Amalgamated zinc, 452
Amadou, 311
American shampoo, 244
tooth-powder, 255
Ammonia, carbonate, 311
sulphate, 311
Ammoniacal gas, 360
Ammoniated perfumes, 225
Eau de Cologne, 225
Anatomical preparations, to preserve, 311
Anchovies, essence, 294
paste, 295
Anderson's pills, 166
Angelica, spirit of, 219
Animal charcoal, 330
Animals, to preserve, 311
Annotto, purified, 312
solution, 312
Anodyne necklaces, 167
Anti-attrition, 313
Antibilious pills, 167
Antiferment, 313
Antipertussis, 167
Antiscorbutic elixir, 262
Antiseptic tooth-powder, 255
Antispasmodic drops, Grinrod's, 187
Aperient pills, various, 167
Apothecaries' measure, 454
(the former), 457
weight, 454
Apparatus, luting for, 329
Aqua cosmetica, 232
fortis, 313
Aquarium, marine, salts for, 313
Arabine, 314
Arbor Dianæ, 314
Argentum musivum, 314
Armenian cement, 326
Aromatic pastils, 230
Aromatic tooth-powder, 255
vinegar, 167, 226
Arquebusade (acid), 167
aromatic, 214
Arrowroot, 276
Artificial manure, 390
sea-water, 313
Asiatic tooth-powder, 255
Asses' milk, artificial, 278
Atkinson's infant preservative, 168
Atropine paper, 168
Aurum musivum, 314
Australian moss, 279
Avoirdupois weight, 456
Axle-grease, 313
Azote, or nitrogen, 362
Azure blue, 415

Bacher's pills, 168
Baden water, 206
Baking powder, 169
Bailey's itch ointment, 169
Baillie's pills, 169
Balaruc water, 209
Baldwin's phosphorus, 314
Baldness, cures for, 244, 245
Balm of Gilead, 169
of Rakasiri, 169
Balls, ash, 314
blacking, 315
breeches, 315
camphor, 236
for copying, 315
wash, 240
heel, 314
scouring, 315
furniture, 315
Balsam of honey, 188
of horehound, 184
Bancroft's solution of tin, 343
Bandoline, 251
Barclay's pills, 169
Bareges water, 209
Baregian balls, 169
Barium, peroxide of, 315
Bark, ess. salt of, 169
Barker's tooth-tincture, 169
Barley water, 278
Barth's green, 417
Barytes, chlorate of, 331
Baster's farina, 282
Bates's eye-water, 170
anodyne balsam, 170
Bates unguentum criniscum, 243
Bateman's drops, 169
itch ointment, 169
Bathing spirits, 170, 184
Bath digestive pills, 170
lozenges, 170
Battley's senna powder, 170
Baume de vie, 170
Baynton's plaster, 170
Bean-flower water, 212
Bear's grease, 244

Bear's grease, factitious, 246
green, 247
Bécœur's arsenical soap, 435
Beddoe's pills, 170
Belloste's pills, 170
Beef tea, 284
Beer, ginger, 268
spruce, 274
treacle, 274
Beetle wafers, 170, 315
poison, 315
Bell metal, 309
Bengal chitni, 291
Benzole, 315
Bestucheff's nervous tincture, 171
Betton's British oil, 171
Bewley's chalybeate, 171
Beverages, 268
Bird-lime, 316
Biscuits, Abernethy, 286
aperient, 171
meat, 284
gluten, 287
sponge, 286
Bismuth, white, 238
Bisulphuret of carbon, 316
Bistre, 414
Black, Brunswick, 323
draught, 171
drop, 171
flux, 351
ink, 373
dyes, 342
japan, 318
oils, 97
reviver, 318
Blacking ball, 318
liquid, 316
paste, 317
for harness, 318
without brushing, 317
patent India-rubber, 317
Blaine's distemper powders, 171
Blake's toothache remedy, 264
Blanc-mange, 280
Irish moss, 278
Bleaching of oils, 395
of wax, 450
liquid, 318
Blights, remedies for, 319
Blistering paper, 171
tissue, 172
Blue for linen, 320
ink, 376
pigments, 415
Prussian, 421
smalts and azure, 414
Blue, Turnbull's, 421
verditer, &c., 415
Bochet's syrup, 172
Boerhaave's odontalgic, 264
Boilers, to prevent incrustations in, 320
Bones, sulphated, 320
Bookbinders' stains, 321
Boot powder, 321
Boot-top liquid, 320
Boots, to render waterproof, 446
Borax, glass of, 364
Borated tincture of myrrh, 261
Bories' odontalgic, 262
Bottle cement, 327
Boudet's depilatory, 254
Braden's farina, 282
Brande's toothache tincture, 264
ink, 375
Brandish's alkaline solution, 172
Brazil paper, 437
Bread, aerated, 285
unfermented, 285
gluten, 287
Breakfast powder, 282
Bright's nutritious farina, 282
Brighton green, 416
Brine for pork, &c., 299
British herb tobacco, 172
oils, 172
Brodum's cordial, 172
Brocchieri's styptic, 173
Broma, 281
Bronze (alloys), 309
for figures, 322
powder, 321
Bronzing liquids, 322
ball, 323
Bronzing pastes, 323
Brosse de corail, 323
Browning for gun-barrels, 323
Brugnatelli's fulminating silver, 355
Brunswick black, 323
green, 416
Bug poison, 323
Bullock's semola, 282
Burnett's (Sir W.) disinfecting fluid, 173
Butter of antimony, 324
to preserve, 324
Butyric ether, 324

Cabbage paper, 437
Cachou aromatisé, 230
Cadet's tooth powder, 255
Cajeput liniment, 173

Calotype process, 403
Camphene, 324
Camphor, artificial, 324
ball, 236
fluid, 195
ice, 236
liniment, 173
washballs, 240
Camphorated chalk, 255
tooth-powder, 256
Candies, 324
Candy, caraway, 193
live-long, 193
Canton's phosphorus, 325
Caoutchouc, solvents for, 431
Cap cement, 327
Capillaire syrup, 274
limoniated, 274
Capsules, gelatinous, 324
Caraway candy, 193
essence of, 293
Carbon, *see* Charcoal, 330
writing ink, 379
sulphuret of, 315
Carbonic acid gas, 360
oxide, 360
dentifrice, 256
Carburetted hydrogen, 361
Carlbad water, 206
Carmine, 412
Carminative, Dalby's, 178
Carabelli's tooth powder, 256
Carrageen, 278
Carron oil, 173
Carthamine, 413
Cartwright's tooth-powder, 256
Case-hardening powders, 324
Cassareep, 296
Cassolettes, 227
Castillion's powders, 173
Castor-oil pomade, 248
Catsup, mushroom, 295
walnut, 295
Cattell's hair dye, 253
Caustic for corns, 178
Cayenne, soluble, 292
essence, 293
Celery, essence of, 293
Cements, various, 325, 330
Armenian, 326
for apparatus, 329
Cement, bottle, 327
cap, 327
coppersmith's, 327
curd, 327
diamond, 326
engineer's, 328
Cement for cracks in iron, 328
for jewels, 326
for leather and metal, 328
for marble, 327
French, 329
hydraulic, 328
Massatt's, 328
oxychloride of zinc, 328
Roman, 328
shellac, 325
soft, 329
for teeth, various, 265
Singer's, 327
Turner's, 327
Cephalic snuff, 173
Ceylon moss, 279
Chalybeate, Palmer's aerated, 196
water, 210
Chamberlaine's pills, 172
Chameleon mineral, 331
Chamomile pills, 173
drops, 173
Charcoal, animal, 330
aluminized, 331
from coal tar, 331
vegetable, 330
tooth powder, 256
tooth paste, 260
Chelsea pensioner, 173
Cheltenham salts, 174
Chemic blue, 331
Chemical elements, table of, 465
Cherry paste for teeth, 260
Chevallier's hair dye, 252
Chilblain's, remedies for, 174
Chinese depilatory, 254
Ching's lozenges, 174
Chirayta pills, 176
Chitni, Bengal, 291
Chlorate of barytes, 331
of potash, 332
Chloride of lime, 332
potash, 332
soda, 332
Chlorine gas, 360
fumigation, 355
lozenges, 231
Chlorodyne, 175
Chloroform, 333
Chocolate and cocoa, 280, 281
Cholera, remedies for, 175
Chromate of lead, 415
of potash, 333
Chromatype paper, 407
Chrome red, &c., 415
yellow, 415
Chromic acid, 303

Cinnamic acid, 303
Cinnamon, essence of, 298
water, 212
tincture of, 292
Circassian cream, 247
tooth-powder, 256
Clichée moulds, 310
Clothes, powder to keep moths from, 334
Clutton's febrifuge, 176
Coathupe's writing fluid, 379
Cochineal coloring, 334
lake, 413
liquid, 388
Cochrane's cough mixture, 176
Cod-liver, Soyer's preparations of, 289
with cod roe, 289
and potatoes, 289
and rice, 289
and tapioca, 289
sauce, 290
Coffee, dandelion, 179
Cold cream, 234
granulated, 235
Collier's wine of quinine, 176
Collin's disinfecting powder, 339
Collodion, 334
elastic, 334
photographic, 401
process, 404
Colors for maps, 336
for confectioners, 336
for druggists' show bottles, 335
liqueurs, 337
liquid, 336
vehicles for, 336
Colored fires, 344
flames, 350
Coloring for brandy, &c., 337
Comparison of thermometic scales, 461
Composition for electro-moulds, 346
Condiments, various, 290
Consumption, remedies for, 176
Copaivi, solution of, 200
Copper, black and red oxide, 337
nitrate of, 337
Coral dentifrice, 256
tooth-paste, 260
Corns, caustic for, 178
plasters, 177
plasters, mechanical, 177
solvents for, 177
Cosmetics, hair, 242
skin, 232
teeth, 255
Cosmetics for chapped nipples, 199
Cosmetiques, in sticks for hair, 253
Cottereau's toothache drops, 264
Cotton, gun, 369
Cough lozenges, 193
linctus, 177
Court plaster, 178, 371
Crayons, lithographic, 380
for writing on glass, 338
Cream, Circassian, 247
cold, 234
lemon, 234
wash-balls, 240
Crystalline cream, 247
castor-oil pomade, 248
Cucumber pomade, 234
Culinary essences, 292
Currant jelly, 276
Curry powder, 290
paste, true, 291
Custard powder, 288

Daffy's elixir, 178
Daguerreotype plates, 417
Dalby's carminative, 178
Dahlia test, 437, 439
Dandelion coffee, 179
Darby's oils, 98
Darcet's lozenges, 179
Dawson's lozenges, 179
De Handel's toothache paste, 265
Dalamott's golden drops, 179
Denham's food, 282
Dentifrices, various, 255, 260
Deodorizing fluid, Ellerman's, 338
Depilatories, various, 254
Derbyshire's embrocation, 179
Deschamps' tooth-powder, 257
Desforges' tooth-powder, 257
extract of pellitory, 262
Detergent tooth-powder, 257
Deutoxide of hydrogen, 397
Dextrine, 338
Diapente, 77
Diamond cement, 326
Diastase, 338
Dictamia, 281
Dietetic articles, 276
Digestive pills, 179
Dinneford's fluid magnesia, 206
Dinner pills, 179
Disinfectants, 338
Disinfecting fluid, Burnet's 338
Ellerman's, 338
Ledoyen's, 339
compounds, 338

Distemper powders, Blaine's, 173
Dixon's pills, 180
Dover's powders, 180
Drayton's silvering process, 367
Driers, for painters, 339
Drinks, summer, 268
Drying oil, 339
Dubbing, curriers', 339
Duffin's Harrogate salts, 188
Duncan's lactuc. lozenges, 180
fl. ext. of senna, 180
gout remedy, 180
Dupasquier's gaseous water, 211
Dupuytren's eye-salve, 180
pomade, 242
Dutch drops, 180
Dyers' compounds, 340
Dyes, various, 340
for the hair, 251
for ivory, 386
for wood, 451
Dyon's tooth-paste, 260

Earache, simple cure for, 267
Eaton's styptic, 180
Eau d'ambre, 207
d'ange, 215
de Bottot, 262
de bouquet, 216
des carmes, 214
de Cologne, 214
collante, 251
dentifrice de Stahl, 262
d'Egypte, 252
d'élégance, 216
d'héliotrope, 216
de jarvelle, 181
d'Ispahan, 216
de luce, 225
de mareschale, 216
de miel, 215
de millefleurs, 216
de magnanimité, 180
mêdicinale d'Husson, 180
de O'Meara, 262
de Portugal, 216
Romaine, 216
de Rabel, 181
sanspareille, 216
de Rosières, 217
de Mdme. Vrillière, 263
Ebony pomatum, 253
Edinburgh lozenges, 183
ointment, 181
Effervescing aperients, 195
powders, 272
Effervescing lemonade, 271
powder, 271
orangeade, 272
orangeade powder, 272
sherbet, 272
Effervescent sherbet powders, 273
chalybeate water, 211
Seidlitz powders, 199
Eger water, 207
Eggs, to preserve, 345
Elderflower water, 212
Electric tissue, 345
Electro-gilding, 364
solutions for, 364
Electro-brassing, 345
bronzing, 345
coppering, 345
silvering, 429
Electrotype moulds, 346
solutions, 428
Electuary, Mahomed's, 194
Elixir Daffy's, 178
de Garus and Haller's, 181
Radcliffe's, 198
of roses, 262
Squire's, 201
Stoughton's 201
for the teeth, 263
of vitriol, 181
Ellerman's deodorizing fluid, 181
Emerald Green, 416
Ems water, 207
Embrocation, Roche's, 199
Embrocations for the hair, 244
Emulsions, cosmetics, 232
Engraving, liquid, 347
on glass, 364
Engravings, to clean, 346
Engraver's varnish, 446
Epicurean sauce, 295
Equivalents, table of, 466–470
Ervalenta, 283
Eschalot wine, 294
Esprit de suave, 217
rose, &c., 218
Essence of Ambergris, 221
anchovies, 294
aromatic, 293
d'ambrette, 219
of bark, 181
of bitter almonds, 292
Cambrian, 300
of caraway, 292
of cayenne, 293
cedrat, 221
of celery, 292
chamomile, 182

Essence cinnamon, cloves, &c., 292
clove pink, 221
Cologne, 221
cubebs, 182
flavoring, 292
ginger, 182
heliotrope, 221
honeysuckle, 221
hovenia, 221
jargonelle pears, 347
jessamine, 224
of jonquil, 222
Madden's, 194
magnolia, 222
mignonette, 222
moss rose, 222
myrtle, 222
musk, 219, 222
mustard, 182
peppermint, 182
pennyroyal, 182
patchouli, 222
of pineapple, 293
of rennet, 425
of roses, 218
rondoletia, 222
royale, 222
of sarsaparilla, 182
for scenting pomatums, 223
for smelling bottles, 225
sweet brier, 223
sweet pea, 223
of Tyre, 225
of vanilla, 220
of verbena, 223
of vetiver, 220
of violets, 218
Westphalian, 300
white lilac, 223
Essences, culinary, 282
conc. infusions, 183
vinous, 183
perfumed, 220, 223
Etching fluids, 347
varnish, 347
Extrait de bouquet, 220
de mareschale, &c., 221
Extract of liquorice, 384
of meat, 284
of malt, 184
Extracts (*Extraits*) of flower, 220

Fairthorn's pills, 184
Fards, or face paints, 237
Farinaceous food, Hard's, &c., 282
Farina, *see* Starches, 434
Farina, Bright's nutritious, &c., 282
Fats, to obviate racidity in, 348
Eerculum Saxonia, 282
Filter for corrosive liquids, 348
Filtering powder, 348
Findings for spirits, 348
Fireproofing, 350
Fires, colored, 348
to extinguish, 348
Fixateur, 251
Flames, colored, 350
Flavoring essence, 293
Flint, liquor of, 350
Floating soaps, 239
Florentine dentifrice, 257
Flowers, to expand, 350
to preserve, 350
Fluide de Java, 250
Fluxes, black and white, 351
Flux, Christison's, 351
crude, 351
Cornish, 351
Morveau's, 352
Fresenius's, &c., 351
Taylor's, 351
Fly poison, 352
Food for infants, 284
Ford's balls, horehound, 184
laudanum, 184
Foreign weights and measures, 459
Fothergill's pills, 184
Fox's cream, 248
Frangipanni bouquet, 223
Frank's solution, 201
Freckles, lotion for, 233
Freeman's bathing spirit, 184
Freezing mixture, 352
French polish, 353
French lip-salve, 237
mustard, 294
tooth-powder, 257
Fryar's balsam, 184
Frigorific mixtures, 352
Fruit lozenges, 193
Fulminating compounds, 354
cotton, 369
powder, 354
mercury, 354
silver, 355
Fumigating pastils, 229
Fumigations, acetic, 357
chlorine, 355
muriatic, 357
nitric, 356
sulphur, 357
tar, 357
for toothache, 265

Furniture cream, 357
oil, 357
paste, 357
Fusible metal, 309

Gad's cement, 328
Galvanic battery, acids for, 307
tooth-powder, 257
Gall, clarified, 358
Galls, tincture of, 439
Gant's cosmetiques, 237
Garancine, 358
Gardiner's alimentary pp., 282
Gannal's solution, 311
Ganteine, 359
Garrot's covering for pills, 359
Gases, to collect, 360
Gas, ammoniacal, 360
carbonic acid, 360
carbonic oxide, 360
chlorine, 360
hydrogen, 361
hydrochloric acid, 361
carburetted hydrogen 361
phosphuretted hydrogen, 361
sulphuretted hydrogen, 361
nitrogen, 362
protoxide, 362
oxygen, 362
ozone, 362
sulphurous acid, 362
Gelatine, 362
capsules, 324
bone, 363
wafers, 449
Gelée pour le goître, 185
Gentil's gluten flour, 288
German lip-salve, 237
paste, 363
pomade, 247
silver, 309
tooth-powder, 257
Gilding, 364
electro, 364
by immersion, 364
Ginger beer, 268
beer powders, 269
essence of, 182
aromatic tincture, 293
Gingerbread, common, 287
cake, 287
purgative, 185
Girambing, 269
Glass, to cut, 364
to mark or write on, 364
to silver, 366
of borax, 364
Glass flux, 327
soluble, 364
Glazes, 367
Gloves, to clean, 359, 387
Glue, 368
liquid, 368
marine, 368
mouth, 368
Gluten, vegetable, 369
bread, 287
granulated, 288
Glycerine jelly, 185
paste, 185
Goadby's solutions, 312
Godbold's veg. balsam, 185
Godfrey's cordial, 185
smelling salts, 186, 225
Gold, factitious, 309
mosaic, 314
Golden ointment, 187
spirit of scurvy grass, 186
compound, 369
Gout, specific, Murray's, 195
paper, 187
medicine, Duncan's, 180
pills, Lartigue's, 190
tincture, Wilson's, 204
Granulated gluten, 288
Granville's (Dr.) lotions, 187
Grape lip-salve, 237
Graves' (Dr.) chilblain remedy, 174
gout, preventive, 187
Grecian water, 254
Gregory's powder, 187
Green dyes, 342
pigments, 416
Greenough's tincture, 187
Griffin's tincture, 187
Griffith's mixture, 187
Grinrod's antispasmodic, 187
Grosvenor's tooth powder, 257
Gruel, oatmeal, 283
Guestonian embrocation, 188
Gum, British, 369
Arabic, purified, 369
Gun cotton, 369
powder, 370
barrels, browned, 323
Guthrie's black ointment, 188
eye ointment, 188
Gutta-percha moulds, 346
solvents for, 432
Gypsum, to harden, 370

Haerlem drops, 180
Hahnemann's hair dye, 252

Hahnemann's wine test, 441
Hair cosmetics, 242
dyes, 251
oils, 248
pomades, 247
powder, 242
washes, 250
to remove, 254
to promote, 242
to stiffen, 251
Halford's (Sir H.) pills, 188
Hamilton's pills, 168
Harness blacking, 318, 371
Harrogate salts, 188, 210
water, 210
Hartshorn jelly, 279
Heading for beer, 274
Helvetius stypic, 188
Hemet's dentifrice, 257
Henbane fumigation, 265
Henry's magnesia, 188
Herb soup-powder, 291
tobacco, 172
Hill's balsam of honey, 188
Hoffman's pills, 188
Honey, balsam of, 188
paste, 236
soap, 239
water, 215
Hooper's pills, 188
Hooping-cough remedies, 189
Horseradish powder, 291
Houlton's laudanum, 188
Hudson's tincture, 262
Huiles antiques, 248
rose, jessamine, &c., 248
Huile macassar, 250
de phenix, 249
philocome, and comogene, 249
Hungary water, 215
Hunt's breakfast powder, 282
Huxham's tincture of bark, 189
Hydrochloric acid, 304
gas, 361
Hydrogen gas, 361
deutoxide, 398
carburetted, &c., 361
Hydrophobia, to prevent, 159
Hypochlorite of potash, 332
Hyposulphite of soda, 431

Iceland moss, 278
chocolate, 278
jelly, 278
Imperial drink, 270
pop, 269
Imperial measure, 454
Incense, 229, 371
India pickles, 299
rubber, solvents for, 431
court plaster, 371
Indigo, purified, 372
dyes, 340
solutions, 331
test paper, 437
Infants' preservative, 168
Ink, to keep from mouldiness, 384
alizarine, 375
powder, 375
red, 375
blue, 376
green, 378
yellow, 378
gold and silver, 378
indestructible, 378
Indian, 379
horticultural, 380
for writing on steel, 380
lithographic, 380
for marking linen, 380
without preparation, 382
crimson marking, 382
printers, 383
Runge's, 374
sympathetic, 383
Weber's, 375
violet, 377
Normandy's, 377, 379
Reade's, 382
Redwood's, 382
Stephen's, 376
Inks, various, 373–384
writing, black, 373
anti-corrosive, 373
copying, 375
packers', 375
Insects, to kill, 384
Iodate of potash, 384
Iodide of potassium, 384
Iodine, 385
Iodized paper, 407
test, 437
collodion, 403
albumen, 404
Iridescent paper, 399
Irish moss, preparation of, 278
Iron, acetate of, 301
liquor, 344
persulphate of, 385
cement, 328
Isinglass, 386
plaster, 192
Issue peas, 189

Ivory black, true, 386
flexible, 386
to stain, 386
to bleach and gild, 387

James's analeptic pills, 189
powders, 189
Japan black, 318
Japanese cement, 329
Jarave, Spanish, 190
Jelly, currant, 276
gelatine, 280
hartshorn, 279
Iceland moss, 278
of Ceylon moss, 279
salep, 279
isinglass, 280
Jessamine, esprit, 218
perfumed oil of, 249
extract, 220
water, 212
Jesuits' drops, 190
Jones's patent flour, 286

Kalydor lotion, 232
Kemmerer's tooth-powder, 258
Kennedy's corn-plaster, 190
Keyser's pills, 190
Kid glove cleaner, 387
King cup, 270
King's yellow, 415
cordial, 190
Ketchup, mushroom, 295
walnut, 295
Kirkland's cerate, 190
Kittoe's freckle lotions, 233
Kitchener's essence of cayenne, 293
of ginger, 293
of spices, 293
pease powder, 291
peristaltic persuaders, 190
soup herb-powder, 291
spirit of soup herbs, 293
savory spices, 293
tincture of cinnamon, 292
Kyan's solution for wood, 387

Labarraque's liquid, 387
Labels, indestructible, 387
Lac, stick, seed, and shell, 387
to bleach, 388
cement, 325
spirit, 353
Lacquers, 444
Lactucarium lozenges, 193
Lait de fraicheur, 233
virginal, 233
Lakes, 388, 413
Lake, cochineal, 413
Brazil, &c., 414
liquid, 388
yellow, &c., 414
Landerer's cure for baldness, 245
Lane's, Dr., vinous essences, 183
Lardner's tooth-powder, 258
Lartigue's gout-pills, 190
Lavender tooth-powder, 258
water, 213
Leath's alimentary farina, 282
Lee's pills, 191
Ledoyen's disinfecting fluid, 190
Leech-bites, astringent for, 191
Lefoulon's elixir, 263
tooth-powder, 258
Lefandinière's elixir, 262
Lejeune's balsam, 191
Lemazurier's odontalgic, 264
Lemon-juice, 388
factitious, 270, 388
flavor, 292
kali, 272
pickle, 295
embrocation, 233
cream, 233
wash balls, 240
Lemonade, acidulated, 270
aerated, 271
dry, 271
powders, 271
milk, 271
Lentil powder, 283
Leroy's purgative, 191
Lenses, extemporaneous, 388
Liebert's nipple cosmetic, 190
Liebig's food for infants, 284
soup for children, 284
Lignum's drops, 191
Limoniated ginger-beer, 269
Linctus, for coughs, 177
Liniment, cajeput, 173
camphorated, 173
St. John Long's, 192
Linseed tea, 278
oil, refined, 389
boiled, 339
clarified, 389
Lip-salve, rose, 236
Peruvian, 237
German, 237
grape, 237
French, 237

Liqueur dorée, 191
Liquid amalgam, 310
blister, 192
disinfecting, 338
glue, 368
for the teeth, 261
Liquids, tables of weights and measures of, 458
Liquorice, purified, 389
Lisbon water, 216
Liston's plaster, 192
Lithographic paper, 398
ink, 378
Litmus, 389
paper, 437
tincture, 439
Live-long candy, 193
Locatelli's balsam, 192
Locock's lotion for the hair, 244
wafers, 192
Long's, St. John, liniment, 192
Lotions for the mouth, 261
of chloride of soda, 263
for chilblains, 174
for the skin, 232
Lozenges, absorbent, 192
alkaline, 179
aperient, 192
Ching's, 174
black currant, 192
and ipecac, 193
cough, 193
digestive, 193
Edinburgh, 195
fruit, 193
marshmallow, 193
peppermint, 194
Lubricating compounds, 389
Lucifer matches, 391
Luminous vials, 390
Lutes, various, 329
Lynch's embrocation, 194

Macassar oil, 250
MacPherson's photo-lithographic process, 411
Madden's essence, 194
Magnesia, fluid, 194
Magnesian aperient, 195
Mahomed's electuary, 194
Mahy's plaster, 194
Maissatt's cement, 328
Manures, artificial, 390
Maredant's Norton's drops, 194
Marble, to clean, 391
to stain, 391
Marienbad water, 207
Marine glue, 368
soap, 430
Marking ink for linen, 380
without preparation, 381
Marmoratum, 267
Marmalade, 275
Marsden's drops, 194
Marsh's dahlia test, 439
Marshall's cerate, 194
eye-drops, 194
Marrow oil, 250
pomatum, 246
Mastic varnish, 443
Marten's photographic albumen, 405
Matches, lucifer, &c., 391
Mathieu's vermifuge, 194
Maury's tooth-powder, 258
Measures, tables of, 460
Meat biscuit, 284
extract of, 284
Medals, to bronze, 322
Medallions, perfumed, 228
Mercury, fulminating, 354
Metallic cements, 266
Metges' tooth-powder, 258
tooth-paste, 260
Mialhes' dentifrice, 258
chalybeate water, 211
ioduretted water, 211
Mice and rat poison, 423
Microscope, formulæ for, 392
Milk of cucumbers, 232
of houseleek, 232
of roses, 232
preserved, 393
Mineral green, 416
waters, 205
chameleon, 331
Mistura odorata, 223
Modelling, clay for, 394
Mohr's lute for retorts, 330
Moirée metallique, 394
Montein's balls for baths, 195
Mont d'Or Water, 210
Mordants, alum, 343
for marking ink, 380
tin, 343
Morrison's paste, 195
pills, 195
Morveau's flux, 351
Moseley's pills, 195
Moss, Australian, 279
Ceylon, 279
Iceland, 278
Irish, 278
Moulds, electrotype, 346

Mouth cosmetics, 255
Moxon's magnesian aperient, 195
Munro's cough medicine, 195
M ltum, 394
Muriatic acid, 304
Muriate of tin, 343
Murray's fluid magnesia, 194, 206
camphor, 195
gout specific, 195
Mushroom catsup, 295
Mustard for the table, 294
French, 294
Myrrh dentifrice, 258
tincture, odoriferous, 261

Nankeen dye, 342
Naples water, 210
yellow, 416
Nelligan's, Dr., pills, 168
Nelson's gelatine, 363
Nepenthe, 196
Neutralizing proportions, table of, 470
Niepce's photo-lithographic process, 410
Nipples, lotion for, 191
Nitrate of barytes, 394
of silver, 394
strontian, 395
Nitrate of potash, 395
Nitrogen gas, 362
protoxide of, 362
Nitroprusside of sodium, 395
Norris's drops, 196
Nouffleur's worm medicine, 196
Novargent, 396

Odontalgic tinctures, 263
elixir, 262
Odoriferous tincture of myrrh, 261
lavender water, 213
oil, boiled and drying, 339
furniture, 358
Macassar, 250
marrow, 250
Oils of ambergris, musk, &c., 249
roses, violets, &c., 248
perfumed, various, 248
for the hair, 248
to purify, 396
to bleach, 396
for machinery, 397
Carron, 173
Darby's, 98
nine, 98
Ointments, Bailey's, 169
Bateman's, 170
Ointments, Edinburgh, 181
golden, 187
Plunket's, 197
Smellome's, 200
Oleine, 397
Ollivier's biscuits, 196
Opodeldoc, 196
Orangeade, 272
aerated, 272
powder, 272
Orange chrome, 415
juice, factitious, 270
marmalade, 275
Orfila's hair-dye, 251
Or-molu, 309
Ormskirk medicine, 196
Ostermaier's cement, 266
Oxygen gas, 362
Oxygenated water, 398
Ozone, 362

Palamoud, 281
Palm oil, to bleach, 396
Palmer's tooth-powder, 258
aerated chalybeate, 196
vitaroborant, 282
Panada, 277
Paper, calotype, &c., 403–409
Brazil, cabbage, &c., 437
copying, 398
fire-proofing for, 399
hydrographic, 398
tests, various, 437
lithographic, 398
paste, 399
photographic, 406
tracing, 399
iridescent, 399
waxed, 399
Papier épispastique, 196
Fayard, 197
maché, 398
Papyrine, 400
Paraguay-roux, 264
Paraffine, 400
Parfum des rois, 217
Passey water, 211
Paste, almond, 235
flour, 329
furniture, 558
honey, 236
Regnauld's, 198
Swediaur's, 174
tooth, various, 259
for toothache, 265
Ward's, 203

Paste for cleaning brass, 400
for razors, 400
blacking, 317
Pastils for burning, 229
for the mouth, 230
Pâte divine de Venus, 234
Payne's preparation for wood, 400
Pearl dentifrice, 258
white, 238
Pease powder, 291
Pelletier's quinine dentifrice, 258
odontine, 260
Pensioner, Chelsea, 173
Percussion caps, 401
Peristaltic persuaders, 190
Perfumes, various, 212–231
acetic, 226
ammoniated, 225
Peruvian lip-salve, 239
Peter's pills, 197
Pettenkofer's test, 442
Pharaoh's serpents, 401
Philanthrope muophobon, 424
Phosphorescent oil, 401
Phosphoric acid, 306
Phosphorus, Baldwin's, 314
Canton's, 325
matches, 401
bottles, 401
paste, 401
Photolithography, 410
Photography, 401–411
collodion, 401
calotype process, 403
collodion process, 403
dry, 404
albumen process, 404
Photographic papers, 406
solutions, 405
recipes, miscellaneous, 409
engraving, 410
Phosphuretted hydrogen, 361
Pickle for pork, &c., 299
Pickles, various, 296
Pickled beet, 297
cabbage, cucumber, &c., 297
mangoes and mashrooms, 298
nasturtiums and onions, 298
walnuts, 298
piccalilli, 299
Pigments, various, 412
vegetable, 412
mineral, blue, &c., 414
Piles, remedies for, 197
Pills, Abernethy's, 166
Anderson's, 166
aperient, 167
Pills, antibilious, 167
Bacher's, 168
Baillie's, 169
Baillie's dinner, 169
Barclay's, 169
Bath, 170
Beddoe's, 170
Belloste's, 171
Chamberlaine's, 173
chamomile, 173
digestive, 179
dinner, 179
Dixon's, 180
Fairthorn's, 184
Fothergill's, 184
Halford's, Sir H., 188
Hamilton's, 168
Hooper's, 188
James's analeptic, 189
Johnson's, 167
Kitchener's, 190
Lartigue's, 190
Lee's, 191
Lynn's, 168
Morrison's, 195
Moseley's, 195
quinine and camphor, 198
Reece's chirayta, 198
Robinson's, 199
Smith's (Hugh), 200
Speedman's, 200
Vance's, 167
Ward's red, 203
Whitehead's, 204
covering for, 197
Pivots for teeth, 267
Plaster, Baynton's, 170
court, 178
isinglass, 192
Liston's, 192
Mahy's, 194
Plate powder, 418
for boiling, 418
Plates, Daguerreotype, 417
Platina, black, 418
chlorides, 419
sponge, 419
Platinated asbestos, 418
Platinized silver, 418
Plumbe's farinaceous food, 283
Plunket's ointment, 197
Poison for beetles, 170, 315
for bugs, 323
for rats and mice, 423
Poitevin's photo-lith. process, 411
Polish, French, 353
furniture, 358

Polishing powder, 419
Pomades for baldness, 242
Pomade for the skin, 233
de beauté, 233
de Ninon, &c., 234
Pomatum, common, 246
colored, &c., 247, 253
crystalline, 247
cucumber, 234
marrow, 246
hard or roll, 247
rose, &c., 246
Pommade contre l'alopécie, 243
castor oil, 248
Cattell's, Dr., 243
Cazenave's, 243
collante, 251
d'Hèbe, 234
divine, 197, 235
Dupuytren's, 242
for beautifying hair, 246
for coloring hair, 253
Potash, 419
Pot-pourri, 227
Pop, imperial, 269
Portland powder, 108
Potato-disease, remedies for, 420
Poudre clarifiante, 420
metallique, 267
à la mousseline, 228
de Chypre, &c., 228
Pounce for parchment, 420
marking ink, 380
Powder, almond, 236
baking, 169
filtering, 348
fulminating, 354
ginger-beer, 269
gun, 370
hair, 242
hand, 234
lemonade, 272
orangeade, 272
spruce-beer, 274
soda, 273
soup, 291
Seidlitz, 199, 208
for rats, &c., 424
plate, 417
Powders for the teeth, 255
Powell's balsam, 198
Preserving animals, 311, 435
Prince of Wales's food, 282
Prometheans, 392
Prussiate of potash, 421
red, 421
Prussian blue, 421
Prussian blue, soluble, 421
Turnbull's, 421
Pullna water, 207
salts, 207
Purging drinks, Dr. Young's, 204
Purple of Cassius, 421
Pyrmont water, 211
Pyroacetic spirit, 422
Pyrogallic acid, 253, 306
Pyroligneous acid, 422
Pyrophorus, 423
Pyroxylic spirit, 422

Queen of Hungary's water, 198, 215
Quin sauce, 295
Quinine dentifrice, 258
wine, 176
and camphor pills, 198

Racahout des Arabes, 281
Radcliffe's elixir, 198
Ragout spice, 291
Ramsey's photo-lith. process, 411
Raspail's cigarettes, 198
Raspberry vinegar, 275
syrup, acid, 275
Rat poisons, 423
without arsenic, 424
Redwood's marking ink, 382
depilatory, 254
Reece's chirayta pills, 198
Regnauld's dentifrice, 259
pectoral paste, 198
Rennet, 425
liquid, 425
wine, 288
Reynolds' gout specific, 199
Revalenta, 283
Rhatany tooth-powder, 259
Rice cake, 286
Riga balsam, 199
Righini's tooth-powder, 259
Robinson's pills, 199
tooth-cement, 266
Roche's embrocation, 199
Rogé's magnesian purgative, 199
Rose lip-salve, 236
tooth-powder, 259
pink, 414
soap, 239
tooth-paste, 260
water, 212, 224
Rosemary tooth-paste, 260
Rosewood, spirit of, 219
Rouge, 237

Rouge, jewellers', 418
Rousseau's drops, 199
Ruspini's tincture, 263
tooth-powder, 259
styptic, 199
Russian tooth-powder, 259
Rust's toothache pill, 265
Rust, to remove and prevent, 425
Ryan's essence of coltsfoot, 199
Rymer's tincture, 199

Sachets, 228
Sago, to prepare, 277
milk, 277
posset, 277
Salep, 279
Saline solutions, 433
tooth-paste, 260
waters, 206, 209
powders, Stevens', 175
Salts, Cheltenham, 174
Harrogate, 188
Marienbad, 207
Pullna, 207
Seidlitz, 207
smelling, 225
Vichy, 208
Sap green, 414
Sauce, anchovy, 294
Epicurean, 295
Quin, 295
superlative, 296
Waterloo, 295
Saunders' dentifrice, 259
Savon au bouquet, 239
Savonettes, 240
Savory spices, 291
Scent bags, 228
balls, 228
for pomatums, 223
powders, 228
for snuff, 223
Scented soaps, 238
Scott's plaster, 199
Scouring drops, 426
Sea-water, 209, 313
salts for, 209
Sealing-wax, 426
Seidlitz water, 207
salts, 208
powders, 199, 208
Seidschutz water, 208
Seltzer water, 208
Semolina, 283
Shaving liquid, 241
paste, 241
Shaving powder, 240
Shampoo liquor, 244
Shellac, 387
to bleach, 388
cement, 325
Sherbet, 272
powders, 273
Siemmerling's cosmetic, 233
Silica cement, 266
Silk-cleaner, 428
Silver, to clean, 428
fulminating, 354
oxidized, 429
to purify and reduce, 429
tree, 442
Silvering, 429
glass, 366
powder, &c., 428
Singer's cement, 327
Singleton's ointment, 200
Sirop d'orgeat, 274
Size, oil, &c., 430
Skeleton leaves, 430
Skin cosmetics, 232
Smalts, 414
Smelling salts, 225
Smellome's eye-salve, 200
Smith's (Hugh) pills, 200
Smoking fluid, 430
Snuff, cephalic, 173
Soap, 430
arsenical, 435
marine, 430
powder, 240
paste, 240
toilet, scented, various, 238
floating, 239
transparent, 239
Windsor, 239
Soda, hyposulphite of, 431
powders, 200, 273
Soft cement, 329
Solders, various, 310
Solomon's balm of Gilead, 200
Solvents for gutta percha, 432
India-rubber, 431
corns, 177
old putty, 432
silver, 433
Soluble glass, 364, 432
cayenne, 292
Solution of copaiba, 200
Solution for electro-plating, 432
Gannal's, 311
for soldering, 433
for voltaic battery, 432
for water-baths, 432

Soubeiran's tooth-cement, 265
Soup for children, 284
Soup-powder, 291
Soy, 296
Specific gravities, 464
Speediman's pills, 200
Spiced vinegar, 296
Spices, mixed, 291
ragout, 291
savory, 291
sausage, 291
essences of, 292
Spilsbury's drops, 201
Spirit of nitric ether, 433
savory spices, 293
soup herbs, 293
Spirits, perfumed, 213, 217, 224
Sponge, blanched, 433
Spruce beer, 274
Squire's elixir, 201
Stains, to remove, 433
Standert's mixture, 201
stomachic candy, 201
Starch, 434
potato, &c., 434
Steam-pipe cement, 328
Stearic acid, 435
Steer's opodeldoc, 201
Stiffening for the hair, 251
Stopping for teeth, 265
Storey's worm cakes, 202
Storm-glass, 435
Stoughton's elixir, 201
Strawberry syrup, 275
spirit, 219
water, 212
Stuffing birds, &c., 435
Styptic, Brocchieri's, 173
Eaton's, 180
Helvetius', 188
Styrol, 436
Struve's lotion, 202
Sugar resin, 436
Sulphite of copper, 436
Sulphocyanide of ammonium, 436
Sulphuretted hydrogen, 361
waters, 209
Swaim's vermifuge, 202
Sydenham's laudanum, 202
Syrup of almonds, 274
capillaire, 274
limoniated, 274
milk, 436
pineapple, 274
raspberries, 275
strawberries, 275
Syrups for effervescing draughts, 273
Table of weights and measures, 454
foreign weights, &c., 459
chemical equivalents, 465
Taffetas vesicant, 172
Talbot's photo-lith. process, 411
Tarmara, 291
Tannin, 307
mouth lotions, 261
Tanjore pills, 202
Tapioca, 277
milk, 277
pudding, 217
Taupenot's photographic albumen 405
Taveare's tooth cement, 266
Teeth cosmetics, 255
stopping for, 265
Temperature, effects of, 462
Terpine, 436
Test papers, 437
solutions, 438
Clark's, for water, 440
Hahnemann's wine, 441
Pettenkofer's, for bile, 442
Trommer's, 441
Thibaut's balsam, 202
Tin, solutions of, 343
mordant's, 343
tree, 442
Tincture of ambergris, 219
civet, 220
red cabbage, 438
balsam of Peru, 219
galls, 439
of ginger, 293
litmus, 439
musk seed, 219
musk, 219
myrrh, odoriferous, 261
borated, 261
patchouli, 220
of quinine, 202
of rhodium, 229
of vanilla, 220
vetiver, 220
Tinctures, culinary, 292
for the teeth, 261
perfumed, 219
test, 438
Tissot's purgative, 202
Tissue, blistering, 172
Tobacco water, 442
Tooth powders, 255
pastes, 259
tincture, 261
Toothache remedies, 263
Touch-paper, 442

Tous les mois, 277
Trade chemicals, 301
Transparent soap, 239
Treacle beer, 274
Trees, metallic, 442
Trophazome, 284
Trousseau's chalybeate, 211
Troy weight, table for converting into avoirdupois, 456
Turlington's balsam, 202
Turner's cement, 327
Turpentine, oil of, 443
as a solvent, 431
Venice, 442
Tyre, essence of, 252

Unfermented bread, 285
Urn-powder, 443

Valangin's mineral solvent, 202
Vance's pills, 167
Vanilla, essence of, 220
tooth-paste, 260
Varnish, amber, 444, 446
Bessemer's, 446
copal, 444
cabinet, 445
Canada, 445
Le Blond's, 446
Macintosh's, 446
India-rubber, 447
for hotbeds, 447
sealing-wax, 447
hair, 447
colored, 447
colorless, for photographs, 444
glass, 448
Japan, 447
for gun-barrels, 447
Varnishes, 443
spirit, 443
(lacquers), 444
essence, 445
oil, 446
various, 445
Vegetable extract for the hair, 250
gluten, 369
Venlo's syrup, 202
Verditer, 415
Vermifuge, Matthieu's, 194
Nouffleur's 196
Swaim's, 202
Vichy water, 208
salts, 209
Victoria perfume, 217
Vinaigre de Cologne, &c., 226
de quatre voleurs, 227
Vinegar (wine, malt, &c.), 448
wood, 422
curry, 294
camp, 294
garlic, shallot, &c., 294
raspberry, 275
spiced, 296
Tarragon, &c., 293
Vinegars, culinary, 293
perfumed, 226
Vinous preparations, Dr. Lane's, 183
Violet tooth-powder, 259
Violets, essence of, 220
Vogler's pill for toothache, 265
Vrillière's mouth-wash, 263

Wacaka des Indes, 281
Wafers, gelatine, 449
Walnut catsup, 295
Warburg's fever tincture, 203
Ward's dropsy powder, 203
essence for headache, 203
paste, 203
red pill, 203
sweating powder, 203
washing powder, 449
white drop, 203
white oils, 97
Walker's Jesuit's drops, 203
Warner's cordial, 203
Warren's hair dye, 252
Warwick's powders, 203
Wash for the hair, 251
powder, for the hands, 236
balls, 240
Washes for vermin in plants, 449
Washing powders, 449
liquids, 449
Water, hard, to soften, 449
Waters, mineral, 205, 211
aerated, 205
alkaline, 205
chalybeate, 210
distilled (perfumed), 212
saline, &c., 209
perfumed, 212–224
spirituous (perfumed), 213
Water-bath, solutions for, 432
Waterloo sauce, 295
Waterproofing compounds, 450
Wax, to bleach, 450
sealing, 426
modelling, 451
Waxed paper, 399

Webster's diet drink, 204
(Lady) pills, 204
Weights and measures, tables of, 454, 461
Welding composition, 451
Wheat, steep for, 451
Whey, 275
powder, 275
Whitelaw's tinct. of lobelia, 204
Whitehead's essence of mustard, 204
pills, 204
White flux, 351
bismuth, 238
zinc, 417
lead, 417
antimony, 417
Wilson's gout tincture, 204
lotions for baldness, 244
Windsor soap, 239
Windham's pills, 191
Wine of shallots, 294
Wine, quinine, 176
Wines, Lane's medicated, 183
Wirth's tooth-cement, 266
Wisdom's eye-water, 204
Withering cosmetic, 233
Wright's pearl ointment, 204
Writing fluids, *see* Inks, 373
Wood, to stain, 451
Worm lozenges, Ching's, 174
Storey's, 202

Yeast, artificial, 452
Young' purging drink, 204

Zinc, ammonio-chloride, 452
amalgamated, 452
oxide, 452
platinized, 452
purified, 453

INDEX

TO THE

VETERINARY FORMULARY.

THE VETERINARY MATERIA MEDICA being alphabetically arranged, it is not considered necessary to include the Drugs, whose uses and doses are there stated, in this Index.

Abbreviations employed in this list: *c*, Cattle; *s*, Sheep; *d*, Dogs; *sw*, Swine. The Horse Medicines have no mark of distinction.

Absorbent powders, 76
Administration of balls, 58
Ægyptiacum, 107
Aloes, solution of, 90
  balls, 67
Alteratives, 19, 59
Alterative balls, 59
  drenches, *sw*. 150
  powders, 76, *c*. 130
Alum, solution of, 106
  mouth-wash, 106
  collyrium, 106
  whey, 130
Anodyne ball, 72
  drench, 130
  fomentation, 104
  poultice, 108
  tincture, 90
Antimonial balls, 59, 62, 65
Antiseptic drench, 89
Antispasmodic balls, 72
  drenches, 81, *c*. 125
Appetite, ball for, 72
Apoplexy, *s*. 143
Aromatic powder, 76
Arsenical powder, 78
  balls, 65
Astringent balls, 60, *d*. 153
  drenches, *c*. 127
  electuaries, 75
Astringent glyster, 92
  lotions, 100, *d*. 159
  powders, 119, *c*. 140, *s*. 147
  caustics, *s*. 148

Balls, for horses, 58, 73
  for dogs, 152, 158
  alterative, 59, *d*. 152
  alt. tonic, 71
  anodyne, 72
  antimonial, 59, 65
  antispasmodic, 72
  for appetite, 72
  astringent, 60, *d*. 153
  chewing, 73
  common mass for, 58
  cordial, 62
  cough, 61, *d*. 152
  diuretic, 64
  diaphoretic, 71
  for distemper, *d*. 154
  farcy, 65
  fever, 65
  for fits, *d*. 155
  glanders, 65
  grease, 67, 73
  hydrophobia, 72, *d*. 159
  inflam. of lungs, 62

Balls, inflammatory disorders, *d.* 156
iodine, 72
for jaundice, 67
laxative, 69
Leicester red, 72
merc. alterative, 59
mixed, 63
miscellaneous, 72
mode of administering, 58
nauseating, 69
parturifacient, *d.* 159
physic or purging, 68
poultry, 163
sedative, 73
stomachic, 69
tonic, mineral, 70, *d.* 157
tonic, vegetable, 70
tonic, diuretic, 65
St. Vitus's dance, *d.* 158
worm, 71, *d.* 157
zinc and valerian, 73
Barley water, 79
Basilicon, 111
Bites of vipers, *c.* 130
Black oils, 97
Blain in the mouth, *c.* 135
Blanche water, 79
Bleeding, 24
to stop, 120, 140
Blistering, 109
liniments, 93
ointments, 109, *c.* 130
Blown or blast, *c.* 124, *s.* 143
Bran mash, 80
Braxy, 144
Bog spavin, 95
Bronchitis, balls for, 62
Bruises, lotions for, 135

Calves, medicines for, 133
cordial, &c., 133
Canker liniment, 96
Canquoin's caustic, 104
Cantharides ball, 26, 65
drench, 88
liniments, 93
oil of, 94, 96
ointment, 109, 138
tincture, 107
Carminative drenches, 81, *c.* 125, *sw.* 150
glyster, 92
tincture, 90
Cataplasms, *see* Poultices.
Catechu lotion, 102
tincture, 107
Cattle medicines, various, 121, 140
drenches, 121
powders, 130
outward applications, 134
plague, medicines for, 131
Caustics, Canquoin's & Filho's, 104
Caustic, Duville's, 148
for farcy buds and poll evil, 103
for foot-rot, *s.* 148
liquid, 103
solid, 104
Charcoal poultice, 108
Charges, 119, *c.* 139
Chewing balls, 73
Chloride of lime lotion, 102
poultice, 109
Citrine ointment, 116
Clark's physic balls, 68
conglutinum, 106
embrocation, 99
lotion, 99
Cleansing drench, *c.* 124
Clysters, *see* Glysters.
Colchicum drench, 85
Colic drenches, 81
tincture, 91
Collyria, 105, *c.* 136
Common ball-mass, 58
poultice, 108
Composition for sand-cracks, 118
Condition powders, 77, *c.* 130
Confections, *see* Electuaries.
Conglutinum, 106
Convulsions, *d.* 155, *ducks* 165
Cooling drinks, 80
Cordial balls, 62
diuretic balls, 63
drenches, 83, *c.* 125
calves', 133
electuaries, 74
Cough balls, 61
electuaries, 74
Cow spice, 131
Cow-pock, lotion for, 135
Cracked heels, ointment, 112
Creasote ointment, 116
liniment, 96
Cropsick poultry, 164
Croton balls, 31, 67
drenches, 31, 87, *c.* 122
electuary, 75
liniment, 136
tincture, 91

Darby's oils, 98
Decoction of linseed, 80

Demulcent drinks, 79, 83
electuary, 74
Deobstruent drenches, 88
Detergent liniments, 96
lotions, 101
ointments, 110, *c.* 138
powders, 120
Diabetes, drench for, 80
Diapente, 77
Diaphoretic ball, 72
Diarrhœa, balls for, *d.* 60, 153
drenches, 80, *c.* 127, *c.* 142
Digestive ointment, 111, *c.* 139
Diuretic balls, 64
cordial balls, 63
drenches, 85, *c.* 130, *s.* 142
electuary, 74
Divine stone, 106
Dogs, medicines for, 152
alterative balls, &c., 152
canker in ear, 160
clysters, 162
distemper, 154
dropsy, 158
emetic powders, 156
enlarged glands, 158
epilepsy, or fits, 155
eye-waters, 160
fits, 155
hydrophobia, 159
inflammatory disorders, 156
lotions, various, 159
mange, washes, &c., 161
ointments, 161
piles, 161
purging balls, 152
rheumatism, 156
scirrhous tumors, 162
St. Vitus's dance, 158
worms, 157
yellows, 158
Drenches, 80, *c.* 121
alum, *c.* 130
anodyne, 90
antiseptic, *c.* 124
antispasmodic, 81
astringent, 79
for bites of vipers, *c.* 130
blown or hoven, *c.* 124, *s.* 143
bloody urine, *c.* 125
for calves, various, 133
carminative, 81, *c.* 125, *sw.* 150
for cattle, various, 121, 130
for catarrh, 83, *c.* 128
for cleansing, *c.* 124
colic, 81
cough, 83, *c.* 128
Drenches, cordial, 83, *c.* 125
cow-pox, *s.* 143
demulcent, 79
deobstruent, 88
for diabetes, 81
for diarrhœa, 80, *c.* 127
ditto in lambs, 142
diuretic, 85, *c.* 129
dropsy, 85
drying, *c.* 130
for dysentery, 80, *s.* 143
emollient, 83
for epidemics, *c.* 128
expectorant, *c.* 128
farcy, 86
fever, 86, *c.* 122
glanders, 86
hoose, in calves, 134
hoven, *c.* 124
inflammation of liver, *c.* 123
influenza, 86, *s.* 145
iodine, 90
laxative, 86, *c.* 121
locked jaw, *c.* 125
murrain, *c.* 129
parturifacient, 89
pectoral, 83, *c.* 128
for poisoning by yew, 89
purging, 86, *c.* 121
for red water, *c.* 126
rheumatism, *c.* 125
stomach staggers, 87
strangury, *c.* 124
tonic, 87
tonic febrifuge, *c.* 123
worm, 88, *calves*, 134
yellows, or jaundice, *c.* 123
Driffield oils, *c.* 137
Drying drench, *c.* 130
Dysentery, 80, 143

Ear, canker of, *d.* 160
sore, *sw.* 151
Eggs, to promote laying, 165
Electuaries, 74
Electuary, cough and laxative, 74
cordial, 74
demulcent and pectoral, 74
diaphoretic, 75
opiate, 74
purgative, 75
tonic and astringent, 75
Embrocatio frigorifera, 99
excitans, 99
Embrocation, cooling, 99
croton, *c.* 136

Embrocation, for garget, *c.* 138
mustard, 98
for saddle-galls and wind-galls, 99
strains, 98
Embrocations, 98, *c.* 136
Clark's, 99
Emetics for dogs, 156
Emulsions, 83, 84
Epidemic affecting mouth and feet, *c.* 128, *s.* 143
catarrh, 62
malignant, 88
pleuro-pneumonia, *c.* 129
Epilepsy in dogs, 155
Euphorbium ointment, 112
tincture of, 107
Expectorant balls, 61, 73
drench, *c.* 130
External applications, *see* Lotions, Liniments, Ointments, &c.
Eye-waters, 105, *c.* 136, *s.* 145
ointments, 111, *d.* 160
powders, *s.* 146

Farcy balls, 65
caustic for, 104
drenches, 86
lotions, 102
ointment, 113
Fattening powder, *c.* 134, *sw.* 150
Febrifuge tonic drench, 123
Fever balls, 65
drenches, 86, *c.* 122, *s.* 141
powders, 77
medicines for swine, 150
Fleas, *d.* 162, *poultry* 165
Fly-powder, *s.* 146
Flux, *see* Diarrhœa.
Fomentations, 104
Foot-rot, *s.* 148
Formulæ, veterinary, 58-165
Foul in foot, *c.* 135, 139
Foxglove tincture, 90

Garget ointment, *c.* 138
liniment, *c.* 137
Giddiness, 87, *poultry* 165
Glanders, balls for, 65
drenches, 86
poultry, 163
Glysters, anodyne, 91
astringent, 92
carminative, 92
diuretic, 93
Glysters, for gripes, 92
laxative, 91, *c.* 131
nourishing, 93
worm, 92
Goulard liniment, 96
ointment, 112
poultice, 109
water, 107
Grease, balls for, 67
lotions, 100
ointments, 112
poultice, 109
Gripe tincture, 90
drenches, 82
Gruel, 79

Hair, to restore, 113
Healing ointment, 161
Heels, scurfy and cracked, 112
greasy, 73, 97, 112
Hellebore ointment, 113
Hepatitis, balls for, 67
Hind's sweating balls, 73
Hoof ointment, 113, 118
Hoose, *c.* 128, *calves* 134
Horse spice, 76
medicines, 17–120
Hoven, drench for *c* 124, *s.* 143
Hydrophobia, remedies for, 72, 89

Inflammation of lungs, 62
liver, *c.* 123
Inflammatory disorders, 62, *c.* 134 *s.* 144, *d.* 156, *sw.* 151
Influenza, *c.* 122, *s.* 143
Infusion of foxglove, 90
Iodine balls, 72
drench, 90
liniment, *c.* 97
ointment, *c.* 138
tincture, 91
Iodide of mercury, 114
Iodhydrargyrate of potash, 67
Itching, to allay, *d.* 160

Jaundice, balls for, 67, *c.* 123, *d.* 158
Joint wounds, *c.* 139

Knees, broken, 113, 118

Lambs, medicines for, 141
cordial, 142
Lapis divinis, 106

Lapis mirabilis, 106
Laxative balls, 59, 69
drenches, *c.* 121, 124
clysters, 91, *c.* 131
electuaries, 75
Leicester red balls, 72
Lice, ointment for, *c.* 138
lotions, *c.* 136
Liniment of ammonia, 98
anodyne, 97
for sore backs, 95
blistering, 93
creasote, 96
caustic, 96
detergent, 96
Goulard, 96
for grease, 96
iodine, 97
for itching humors, 95
lime-water, 97
mange, 96, *c.* 137
marshmellow, narcotic, &c., 97
saturnine, 96
for sore throats, 137
stimulating, 94
suppling, 118
sulphate of copper, 107
sweating, 93
turpentine, 95
verdigris, 107
Liniments, various, 93–99
Linseed mash, 80
tea, 79
Liquid blister, 93
caustics, 102, *c.* 137
snuff, *c.* 137
Lock-jaw, drench for, *c.* 125
Lotion, astringent, 100, *d.* 159
black, 101
for blain in mouth, *c.* 135
bruises, *c.* 135, *d.* 162
for canker, *c.* 134
Lotions, catechu, 103
various, 99, 101, *c.* 134, *d.* 159
chloride of lime, 103
Clark's, 99
cooling, 99
for cow-pox, *c.* 135
detergent, 101
discutient, 99, *c.* 135
farcy, 102
foul in foot, *c.* 135
mange, 101, *c.* 137, *d.* 161
mercurial, 102
for the mouth, 102
nitric acid, 101
Lotions, refrigerant, 99
for strains, 100, *c*, 136, *d.* 162
styptic, 102
for vermin, 102, *c.* 136
wounds, *c.* 135, *poultry* 165
yellow, 101
Louse-water, 102
Low, ointment for, *c.* 139
Lungs, inflammation of, 62

Madness, canine, 159
Materia medica, 17–57
Malignant epidemics, 88
Mallenders, 115
Malt mash, 80
Mange, lotions for, 101, *c.* 136, *d.* 161
liniments and oils for, 95
ointments, 114, *c.* 138, *sw.* 151
Marshall's oils, 98
Marshmallow ointment, 115
liniment, 97
Mashes, bran, 80
linseed, 80
malt, 80
Masticatories, 73, *c.* 131
Meadow sickness, *s.* 143
Medicated provender, 78
Medicine for horses, 58
cattle, 121
for cattle plague, 131
sheep, 141
swine, 150
dogs, 152
poultry, &c., 163
mercurial balls, 59, 65, 66
alterative, 59
lotion, 101
ointment, 115
Mild drinks, 79
caustics, 102
Mineral tonics, 70
Mixed balls, 63
Murrain drinks, *c.* 129
Mustard embrocation, 98
poultice, 109
Myrrh and aloes tincture, 90

Nauseating balls, 69
Newmarket oils, 98
Nine oils, 98
Nitre balls, 64, 65
solution, 91
powder, 76, 77

Oils, various, 98, 98
black, 97
Darby's, 98
egg, 98
for mange, 98
Marshall's nine, Radley's, 98
Stamford's, Lord, 98
Ward's, 97
white, 97
Ointment, for blemished knees, 113
blistering, 109, *c.* 138
for canker, 117
citrine, and creasote, 116
detergent, 110
digestive, 111
eye, 111
euphorbium, 112
farcy, 113, 117
foot, 139
grease, 112
healing, 161
hellebore, 113
hoof, 113, 118
iodine, 114, *c.* 138
for mallenders, 115
mange, 114, *c.* 138, *sw.* 151
marshmallow, 115
mercurial, 115, *s.* 147
nitrate of silver, 160
scab or shab, *s.* 147
simple, 109
for sore backs, 116
for sore teats, *sw.* 151
for sitfasts, 116
sulphuric acid, 116
tar, 117
for thrush, 117
verdigris, 117
for vives, 117
warts, 117
Ointments, 109, *c.* 138
Opiate electuary, 74

Parturition, to promote, 89, *d.* 158
Paste, astringent, 118
for broken knees, 118
styptic, 140
Pectoral electuary, 74
balls, 61
drenches, 83
powder, 78
Physic balls, 68
Pneumonia, epizootic, *c.* 129
Poisoning by yew, 89
Poultices, common, 108
anodyne, 108
Poultices, charcoal, 108
chlorine, cleansing, drawing, 108, 109
drying, 108
Goulard, 109
for grease, 112
mustard, rubefacient, 109
yeast, 108
Poultry, medicines for, 163–165
outward applications, 164
Powders, absorbent, 76
alterative, 76, *c.* 130
aromatic, 76
arsenical, 78
bitter, 78
conditions, 77, *c.* 130
fever, 77
farcy, 77
fly in sheep, 146
for grease, 76
pectoral, 78
for outward use, 119
astringent, 119
detergent, 120
styptic, 120
sneezing, 120
Provender, medicated, 78
Purging balls, 68, *d.* 152
drenches, 86, *c.* 121, 133
electuaries, 75
Purgatives for swine, 150
sheep, 141
dogs, 152
for poultry, 164

Rabbits, medicines for, 165
Radley's oils, 98
Red water drench, *c.* 126, *s.* 142
Resolvent ointment, 115
Rheumatic drench, *c.* 125
Rinderpest, treatment of, 131
Roaring, balls for, 72
Rot, foot, 148
in rabbits, 165
in sheep, 144
Roup, poultry, 163

Saddle galls, 100, 116
Sallenders ointment, 115
Sandcrack, composition for, 118
Sedative balls, 73
aperient, 73
Setons, 50, *c.* 140
Sheep and lambs, medicines, 141
drench for apoplexy, 143

Sheep, braxy, 144
diarrhœa, 142
purging, 141
medicines for red-water, 142
epidemic distemper, 143
sturdy, 143
skit, 143
ointment for scab, 147
wounds, and sore heads, 148
caustic, 148
Sitfasts, ointments for, 116
Sneezing powder, 120
liquid, 137
Sore backs, ointment, 116; lin. 95
Softening ointment, 116
Spice, cow, 131
horse, 76
Stanford oils, 98
Staggers, stomach, 87
Stimulating drench, 83
Stomachic balls, 69
Stone, styptic, 106
wounds, 106
mirabilis, 106
divine, 106
Strains, lotions for, 100, *c.* 135
Strangury, drench for, *c.* 124
Sturdy, drink for, 143
Styptic stone, 106
powder, 120
paste, 118, *c.* 140
Sulphuric acid ointment, 116
caustic, 103
Sulphate of copper liniment, 107
balls, *see* Farcy balls.

Tar ointment, 117
Thriving powder, *c.* 134, *sw.* 150
Thrush, ointment for, 117
Tincture of aloes and myrrh, 90
anodyne, 91
of cantharides, 107
colic, 91
of croton, 91
euphorbium, 107
alkaline, 107
Tincture, foxglove, 90
gripe, 90
myrrh and aloes, 90
opium, etherial, 90
of pimento, 90
styptic, 107
Tonic balls, 70
drenches, 123
electuary, 75
Turpentine liniment, 95
ointment, 117

Ulcers, lotions for, 101, 102, *s.* 148
ointments, 111, *c.* 139

Veterinary Materia Medica, 17–57
formulæ, 58–165
Verdigris ointment, 117
liniment, 107
Vermin, to destroy, 102, *c.* 136
s. 145, *d.* 162; *poultry*, 165
Viper, bites of, *c.* 130
Vitus's (St.) dance, *d.* 158

Ward's white oils, 97
Warts, ointment for, 117
Washes, 99–102
black and yellow, 101
mercurial, 101
various, *see* Lotions.

Waters, eye, *see* Collyria.
White's solution of potash, 134
White egg oils, 97
Worms, balls for, 71, *d.* 157
Wounds, applications to, *c.* 119, &c., *see* Ointments and Lotions.

Yellows, or jaundice, *c.* 123
Yew, poisoning by, 89

Zinc and valerian balls, 73

PHILADELPHIA,
September, 1871.

LINDSAY & BLAKISTON hand enclosed a NEW CATALOGUE *of their* MEDICAL, DENTAL, *and* SCIENTIFIC *Publications, in which will be found many new and important works, and new editions of others just issued by them. Devoting themselves entirely to the publication of works on Medicine and the collateral sciences, they are enabled to offer unusual facilities to authors in publishing and extending the sale of their works. They also publish a* CLASSIFIED *and* PRICED LIST *of all the more important American and English Medical Publications, Periodicals, &c., &c., which they keep on hand, or furnish to order, at the lowest prices. The Classified List, together with their Catalogue, they will furnish post free, by mail or otherwise, upon application.*

FOREIGN MEDICAL WORKS, *not on hand, are imported promptly to order.*

The following NEW WORKS and NEW EDITIONS
ARE NOW READY, or will be issued during the coming season.

RINDFLEISCH'S TEXT-BOOK OF PATHOLOGICAL HISTOLOGY, translated from the German. With 208 Microscopical Illustrations.

WARING'S MANUAL OF PRACTICAL THERAPEUTICS. Second American from the Third London Edition.

MACKENZIE ON LARYNGEAL GROWTHS, with Chromo-Lithograph and other Illustrations.

HARRIS' PRINCIPLES AND PRACTICE OF DENTISTRY, with over 400 Illustrations. Tenth Revised Edition.

MEADOWS' TEXT-BOOK OF MIDWIFERY. From the Second London Edition. With Illustrations.

DUCHENNE'S LOCALIZED ELECTRIZATION, with Notes and Additions by the Translator, and Numerous Illustrations.

TANNER AND MEADOW'S PRACTICAL TREATISE ON THE DISEASES OF INFANCY AND CHILDHOOD, Second Edition, Revised and Enlarged.

LINDSAY & BLAKISTON'S PHYSICIAN'S VISITING LIST FOR 1872. Various Sizes and Styles of Binding.

BYFORD ON THE UTERUS. A New Revised and Enlarged Edition.

DILLNBERGER'S HANDY-BOOK OF THE DISEASES OF WOMEN AND CHILDREN.

TILT ON THE CHANGE OF LIFE IN WOMAN. From Third London Edition.

ACTON ON THE REPRODUCTIVE ORGANS. Third American from the Fifth London Edition.

PEREIRA'S PRESCRIPTION BOOK. A New American, from the Fifteenth London Edition.

WYTHE'S POCKET DOSE AND SYMPTOM BOOK. A New Revised Edition.

TROUSSEAU'S LECTURES ON CLINICAL MEDICINE. Vol. 4.

WEDL'S DENTAL PATHOLOGY, translated from the German. With numerous Illustrations.

GANT'S SCIENCE AND PRACTICE OF SURGERY. A New Text-Book of Surgery. With 400 New Illustrations.

BIDDLE'S MATERIA MEDICA. The Fourth Revised Edition.

BEALE ON DISEASES OF THE LIVER AND THEIR TREATMENT. A New, Revised, and Enlarged Edition.

BEALE'S PROTOPLASM, LIFE, MATTER, AND MIND. New Enlarged Edition

WRIGHT ON HEADACHES. A New Edition.

LINDSAY & BLAKISTON'S

MEDICAL PUBLICATIONS.

"The Representative Book of Medical Science."—LONDON LANCET.

Aitken's Science and Practice of Medicine.

SECOND AMERICAN from the FIFTH LONDON EDITION.

In 2 Volumes, containing 2000 Royal Octavo Pages, a COLORED MAP, a LITHOGRAPHIC PLATE, and ONE HUNDRED AND THIRTY ILLUSTRATIONS ON WOOD.

Price, bound in Cloth, bevelled boards $12.00
" " Leather 14.00

Dr. Aitken's work is now the most comprehensive TEXT-BOOK on the Practice of Medicine in the English Language; the present edition has been enlarged and carefully revised by the author, as well as by the American editor, MERIDITH CLYMER, M.D., the latter having made additions of original matter equal to nearly 500 pages of the English Edition, with special reference to the wants of the American Practitioner.

Acton on the Functions and Disorders of the Reproductive Organs. NEW EDITION.

In Childhood, Youth, Adult Age, and Advanced Life, considered in their Physiological, Social, and Moral Relations. By WILLIAM ACTON, M.R.C.S., etc. *Third American from the Fifth London Edition. Carefully revised by the author, with additions.* Just Ready, octavo, $3.00

"We think Mr. Acton has done good service to society by grappling manfully with sexual vice, and we trust that others, whose position as men of science and teachers enable them to speak with authority, will assist in combating and arresting the evils which it entails. We are of the opinion that the spirit which pervades it is one that does credit equally to the head and to the heart of the author."—*The British and Foreign Medico-Chirurgical Review.*

Anstie on Stimulants and Narcotics.

Their Mutual Relations, with Special Researches on the Action of Alcohol, Ether, and Chloroform on the Vital Organism. By FRANCIS E. ANSTIE, M.D., *Assistant Physician to Westminster Hospital, Lecturer on Materia Medica and Therapeutics, etc., etc.* Octavo, ... $3.00

Althaus' Medical Electricity. *A New and Very Much Enlarged Edition.*

A Theoretical and Practical Treatise, and its Use in the Treatment of Paralysis, Neuralgia, and other Diseases. By JULIUS ALTHAUS, M.D., *Member of the Royal College of Physicians, &c. Second Edition, revised, enlarged, and for the most part rewritten.* In One Volume Octavo, *with a Lithographic Plate and sixty-two Illustrations on Wood* Price, ... $5.00

Byford on the Uterus. SECOND EDITION.

On the Chronic Inflammation and Displacement of the Unimpregnated Uterus. A New, Enlarged, and Thoroughly Revised Edition, with Numerous Illustrations. Now Ready. One volume. Octavo. $3.00

From FORDYCE BARKER, M.D., *Professor of Obstetrics and Diseases of Women and Children in Bellevue Hospital Medical College.*

Some weeks ago I received a copy of your work on the Uterus. I have delayed acknowledging the favor until I could give the book a careful perusal. I have just finished a thorough reading of it. I feel personally indebted to you, not merely for the copy — which, of course, I should have bought — but for writing the book; and I think you have laid the Profession in this country under a load of obligation by giving them such a clear, concise, and practical treatise on a class of affections that even now is very little understood by the greater majority.

From R. A. F. PENROSE, M.D., *Professor of Obstetrics in the University of Pennsylvania.*

Accept my thanks for the copy of your new work which you so kindly sent me. I have, as yet, not had time to give it the careful study it merits; but from the superficial inspection I have made, I find much that is most valuable.

From S. G. HUBBARD, M.D., *Professor of Obstetrics in Yale College, New Haven.*

I was gratified by the receipt of your new work on the Uterus, and I thank you sincerely for it. I have spent all my leisure in its examination, and have derived both pleasure and profit from its perusal. It is commendable for its clearness and definiteness as well as for the great practical common sense which pervades it. I am sure that it will prove a very useful treatise, not only to junior practitioners, but to those also among us who, from not having devoted themselves to the treatment of uterine disease, as a specialty, have neither time nor opportunity to make original investigations in this department, and are not, therefore, perfectly at home in its practice.

From JAMES P. WHITE, *Professor of Obstetrics and Diseases of Women and Children in the University of Buffalo.*

I have had time, as yet, to run over but few of the chapters of your work on the Uterus; I am most happy, however, in being able to say that, so far as I have read, it does credit to American authorship. It is concise and brief, and eminently practical. The work was certainly a desideratum, and will be especially useful to practitioners who can illy afford to purchase all or most of the works referred to in your preface. I shall examine it with much interest, and, no doubt, often consult it with profit.

From G. S. BEDFORD, M.D., *Professor of Obstetrics and Diseases of Women and Children in the University of New York.*

On my return to the city I found on my table "Byford on the Uterus." The next day I commenced perusing it, and have read it from cover to cover. I need not say that, in my judgment, the book enhances your deserved reputation. You have, if my opinion be worth anything, given the Profession an excellent work, and one that is sensible and practical. Go on, my dear Doctor, and give us more of your experience. It is what the Profession most needs — the experience of good and ripe minds.

RECENTLY PUBLISHED, THE SECOND EDITION OF

Byford's Practice of Medicine and Surgery.

Applied to the Diseases and Accidents Incident to Women. By W. H. BYFORD, A.M., M.D., *Professor of Obstetrics and Diseases of Women and Children in the Chicago Medical College, &c., &c. The Second Edition, Revised and Enlarged, with Additional Illustrations.* One volume. Octavo. $5.00

The rapid sale of the first edition of this book, which was exhausted in a little more than a year, has enabled the author to carefully revise the whole work, add many improvements, and to make a large addition of new matter, without, however, materially increasing the size of the volume.

This work treats well-nigh all the diseases incident to women, diseases and accidents of the vulva and perineum, stone in the bladder, inflammation of the vagina, menstruation and its disorders, the uterus and its ailments, ovarian tumors, diseases of the mammæ, puerperal convulsions, phlegmasia alba dolens, puerperal fever, &c. Its scope is thus of the most extended character, yet the observations are concise but convey much practical information. — *London Lancet.*

Beasley's Book of Prescriptions.

A NEW, REVISED, AND ENLARGED EDITION.

Containing 3000 *Prescriptions, collected from the Practice of the most Eminent Physicians and Surgeons — English, French, and American; comprising also a Compendious History of the Materia Medica, Lists of the Doses of all Officinal and Established Preparations, and an Index of Diseases and their Remedies.* By HENRY BEASLEY. *Second American from the last London Edition.* Octavo. $4.00

"The editor, carefully selecting from the mass of materials at his disposal, has compiled a volume, in which both physician and druggist, prescriber and compounder, may find, under the head of each remedy, the manner in which that remedy may be most effectively administered, or combined with other medicines in the treatment of various diseases. The alphabetical arrangement adopted renders this easy; and the value of the volume is still further enhanced by the short account given of each medicine, and the lists of doses of its several preparations. It is really a most useful and important publication, and, from the great aid which it is capable of affording in prescribing, should be in the possession of every medical practitioner. Amongst other advantages is, that, by giving the prescriptions of some of the most able and successful practitioners of the day, it affords an insight into the methods of treatment pursued by them, and of the remedies which they chiefly employed in the treatment of different diseases." — *Lancet.*

Beasley's Druggists' General Receipt Book.

SIXTH AMERICAN EDITION, REVISED AND IMPROVED.

Comprising a copious Veterinary Formulary, numerous Receipts of Patent and Proprietary Medicines, Druggists' Nostrums, etc.; Perfumery and Cosmetics, Beverages, Dietetic Articles and Condiments, Trade Chemicals, Scientific Processes, and an Appendix of Useful Tables, by HENRY BEASLEY, *Author of the Book of Prescriptions, etc., etc. Sixth American from the Last London Edition.* Octavo. $3.50

"This is one of the class of books that is indispensable to every Druggist and Pharmaceutist as a book of reference for such information as is wanted, not contained in works used in the regular line of his business, and we can recommend it as one of the best of the kind." — *American Druggists' Circular.*

Barth & Roger's Manual of Auscultation and Percussion.

A new American Translation from the Sixth French Edition. 16mo. $1.25.

"This is one of the most useful and practical manuals of its sort that has ever yet appeared, and we cannot too strongly recommend it to every student of medicine. It is sufficiently comprehensive without being lengthy, and the principles, which are eminently sound, can easily be mastered and understood." — *Medical Record.*

Bouchardat's Annual Abstract

OF THERAPEUTICS, MATERIA MEDICA, PHARMACY, AND TOXICOLOGY, FOR 1867, *with an Original Memoir of Gout, Gravel, Urinary Calculi, &c. By* A. BOUCHARDAT, *Professor of Hygiene to the Faculty of Medicine, Paris, &c. Translated and Edited by* M. J. DE ROSSET, M.D., *Adjunct to the Professor of Chemistry in the University of Maryland. In one Volume.* Price, in cloth, . . $1.50.

WORKS BY DR. LIONEL S. BEALE, F.R.S.,

Fellow of the Royal College of Physicians, Physician to King's College Hospital.

Seventh Thousand. Four Hundred Illustrations, some Colored.

HOW TO WORK WITH THE MICROSCOPE.

This work is a complete manual of microscopical manipulation, and contains a ful description of many new processes of investigation, with directions for examining objects under the highest powers, and for taking photographs of microscopic objects. Octavo. Cloth. Price, $7.50.

ON KIDNEY DISEASES, URINARY DEPOSITS, AND CALCULOUS DISORDERS:— *The Third Edition.*

Including the Symptoms, Diagnosis, and Treatment of Urinary Diseases. With full Directions for the Chemical and Microscopical Analysis of the Urine in Health and Disease. 70 Plates, 415 figures, copied from Nature. Octavo. Price, $10.00.

THE USE OF THE MICROSCOPE IN PRACTICAL MEDICINE.

For Students and Practitioners, with full directions for examining the various secretions, &c., in the Microscope. 4th Edition. 500 Illustrations. Octavo. Price, $7.00.

THE MYSTERY OF LIFE.

An Essay in reply to Dr. Gull's Theory, with two Colored Plates. Price, $1.50.

LIFE, MATTER, AND MIND; OR PROTOPLASM.

With Original Observations on Minute Structure, and numerous New Colored Drawings. *A New Edition, very much enlarged. Eight Plates.* Price, $

DISEASE GERMS: THEIR SUPPOSED NATURE. *With Colored Plates.*

An Original Investigation with the Aid of the Highest Powers yet made.

This work contains a critical examination of Dr. Tyndall's statements concerning 'Dust and Disease.' 12mo. Cloth. Price, $1.75.

DISEASE GERMS: THEIR REAL NATURE. *Twenty-four Plates.*

An Original Investigation. Twenty-four Plates, sixteen of which are Colored.

Including the minute pathology of contagious diseases. Price, $4.00.

Now ready. Four Colored Plates.

PHYSICAL THEORIES OF LIFE: THEIR INFLUENCE UPON RELIGIOUS THOUGHT. Price $2.50.

With Observations on the Hypotheses recently advocated by Tyndall and Huxley.

ON DISEASES OF THE LIVER, AND THEIR TREATMENT.

A Second Edition, much enlarged, of the Author's Work on the Anatomy of the Liver. Numerous Plates. *Preparing.*

THE ARCHIVES OF MEDICINE. *Part XVII. now ready.* Price, $1.50.

CONTENTS.—Medicine under the Romans—Action of Eyelids in Expiration; F. C. Donders, Utrecht—Lacunæ and Canaliculi—Nerves in Mesenteric Glands; Dr. Popper, St. Petersburg—German Criticism and British Medical Science—Homology of Coracoacromial Ligament; J. B. Perrin—Sulpho-carbolates, by Dr. Sansom—Structure of the Liver; Editor—Sputum in Phthisis.

DISEASE: ITS NATURE AND TREATMENT. *Preparing.*

New Researches on Inflammation and Fever, and on the Nature of Contagion. With Observations on the Cattle Plague and on Cholera. Numerous Colored Plates. Being the Third Course of Lectures delivered at Oxford by direction of the Radcliffe Trustees.

All these Works contain the results of the Author's original investigations. They are Illustrated with upwards of 2000 Engravings, copied from the actual objects, all of which have been drawn on wood by the Author himself, or under his immediate superintendence.

LINDSAY & BLAKISTON, Publishers, Philadelphia.

Biddle's Materia Medica. FOURTH EDITION.

For the Use of Students. With Illustrations. By J. B. BIDDLE, M.D., *Professor of Materia Medica and Therapeutics in the Jefferson Medical College, Philadelphia, &c., &c. Revised and Enlarged.* $4.00

This is a thoroughly revised and enlarged edition of Prof. Biddle's work on Materia Medica. It is designed to present the leading facts and principles usually comprised under this head as set forth by the standard authorities, and to fill a vacuum which seems to exist in the want of an elementary work on the subject. The larger works usually recommended as text-books in our Medical schools are too voluminous for convenient use. This will be found to contain, in a condensed form, all that is most valuable, and will supply students with a reliable guide to the courses of lectures on Materia Medica as delivered at the various Medical schools in the United States.

Bull on the Maternal Management of Children in Health and Disease. . . $1.25

Birch on Constipated Bowels.

The Various Causes and the Different Means of Cure. By S. B. BIRCH, M.D., *Member of the Royal College of Physicians of London, &c. Third Edition.* Price, ... $1.00

Braithwaite's Epitome of the Retrospect of Practical Medicine and Surgery. 2 Vols. $10.00

Chambers on the Renewal of Life.

Lectures chiefly Clinical, illustrative of a Restorative System of Medicine. By THOS. K. CHAMBERS, M.D., *Physician to St. Mary's Hospital, author of "The Indigestions," &c., &c. A new American from the Fourth London Edition.* Octavo, . . . $5.00

"This work is of the highest merit, written in a clear, masterly style, and devoid of technicalities. It is simply what it professes to be, Lectures Clinical, delivered from cases observed at the bedside; therefore more valuable as enunciating the views and experiences of a practical mind aided by actual observation. They are of deep interest, and replete with facts having a practical bearing, and will well repay perusal."—*Canada Medical Journal.*

Chew on Medical Education.

A Course of Lectures on the Proper Method of Studying Medicine. By SAMUEL CHEW, M.D., *Professor of the Practice and Principles of Medicine and of Clinical Medicine in the University of Maryland.* $1.00

"Dr. Chew was an eminent member of the medical profession, and a well-known teacher of medicine. He was, therefore, well fitted for the judicious performance of this task, upon which he seems to have entered with interest and pleasure. It is a well-timed book, and will serve as a most excellent manual for the student, as well as a refreshing and suggestive one to the practitioner."—*Lancet and Observer.*

Cazeaux's Great Work on Obstetrics.

THE MOST COMPLETE TEXT-BOOK NOW PUBLISHED.

GREATLY ENLARGED AND IMPROVED.

CONTAINING 175 *ILLUSTRATIONS.*

A Theoretical and Practical Treatise on Midwifery, *including the Diseases of Pregnancy and Parturition, by* P. CAZEAUX, *Member of the Imperial Academy of Medicine; Adjunct Professor in the Faculty of Medicine of Paris, etc., etc. Revised and Annotated by* S. TARNIER, *Adjunct Professor in the Faculty of Medicine of Paris; Former Clinical Chief of the Lying-in-Hospital, etc., etc. Fifth American from the Seventh French Edition. Translated by* WM. R. BULLOCK, M.D. In one volume Royal Octavo, of over 1100 pages, with numerous Lithographic and other Illustrations on Wood.

Price, bound in Cloth, bevelled boards, . . .	$6.50
" " Leather, ...	7.50

M. Cazeaux's Great Work on Obstetrics has become classical in its character, and almost an Encyclopædia in its fulness. Written expressly for the use of students of medicine, and those of midwifery especially, its teachings are plain and explicit, presenting a condensed summary of the leading principles established by the masters of the obstetric art, and such clear, practical directions for the management of the pregnant, parturient, and puerperal states, as have been sanctioned by the most authoritative practitioners, and confirmed by the author's own experience. Collecting his materials from the writings of the entire body of antecedent writers, carefully testing their correctness and value by his own daily experience, and rejecting all such as were falsified by the numerous cases brought under his own immediate observation, he has formed out of them a body of doctrine, and a system of practical rules, which he illustrates and enforces in the clearest and most simple manner possible.

OPINIONS OF THE PRESS.

"It is unquestionably a work of the highest excellence, rich in information, and perhaps fuller in details than any text-book with which we are acquainted. The author has not merely treated of every question which relates to the business of parturition, but he has done so with judgment and ability." *British and Foreign Medico-Chirurgical Review.*

"The translation of Dr. Bullock is remarkably well done. We can recommend this work to those especially interested in the subjects treated, and can especially recommend the American edition." *Medical Times and Gazette.*

"The edition before us is one of unquestionable excellence. Every portion of it has undergone a thorough revision, and no little modification; while copious and important additions have been made to nearly every part of it. It is well and beautifully illustrated by numerous wood and lithographic engravings, and, in typographical execution, will bear a favorable comparison with other works of the same class."—*American Medical Journal.*

"In the multitudinous collection of works devoted to the propagation of human beings, and to the details of parturition, none, in our estimation, bears any comparison to the work of Cazeaux, in its entire perfectness; and if we were called upon to rely alone on one work on accouchments, our choice would fall upon the book before us without any kind of hesitation."— *West. Jour. of Med. and Surgery.*

"We do not hesitate to say, that it is now the most complete and best treatise on the subject in the English language."— *Buffalo Medical Journal.*

"We know of no work on this all-important branch of our profession that we can recommend to the student or practitioner as a safe guide before this."—*Chicago Medical Journal.*

"Among the many valuable treatises on the science and art of obstetrics, the work of Cazeaux stands pre-eminent."— *St. Louis Med. and Surg. Journal.*

"M. Cazeaux's book is the most complete we have ever seen upon the subject. It is well translated, and reflects great credit upon Dr. Bullock's intelligence and industry."— *N. A. Medico-Chirurg. Review*

Cleaveland's Pronouncing Medical Lexicon.

Containing the Correct Pronunciation and Definition of most of the Terms used by Speakers and Writers of Medicine and the Collateral Sciences. By C. H. CLEAVELAND, M.D., *Member of the American Medical Association, &c., &c. A New and Improved Edition.* $1.25

This little work is not only a Lexicon of all the words in common use in Medicine, but it is also a Pronouncing Dictionary, a feature of great value to Medical Students. To the Dispenser it will prove an excellent aid, and also to the Pharmaceutical Student. It contains a List of the Abbreviations used in Prescriptions, together with their meaning; and also of Poisons and their Antidotes. It has received strong commendation both from the Medical Press and from the profession.

Cohen on Inhalation.

Its Therapeutics and Practice. A Treatise on the Inhalation of Gases, Vapors, Nebulized Fluids, and Powders; including a Description of the Apparatus employed, and a Record of Numerous Experiments, Physiological and Pathological; with Cases and Illustrations. By I. SOLIS COHEN, M.D. 12mo. Price, ... $2.50

"We recognize in this book the work of a persevering Physician who has faithfully studied his subject, and added to its literature much that is useful from his own experience. Dr. Cohen has given us briefly and clearly whatever is valuable in relation to the insufflation of powders in respiratory affections, with the experimental proofs and pathological evidence of their penetration into the bronchial tubes and lung tissues." *American Journal of Medical Science*, July, 1868.

Carson's History of the Medical Department

Of the University of Pennsylvania, from its Foundation in 1765: *with Sketches of Deceased Professors, &c.* By JOSEPH CARSON, M.D., *Professor of Materia Medica and Pharmacy in the University.* $2.00

'The history of the University of Pennsylvania has a national as well as a local interest, from the early date of its origination, and the connection with it of men of illustrious public reputation, such as Drs. Franklin, Rush, Physick, Gibson, Dewees, Chapman, Wood, &c., &c. For the labor and love which he has spent in preparing this most interesting and valuable work, Prof. Carson has earned the gratitude of the alumni of the University, and of all others interested in medical education in this country." — *American Journal of Medical Science.*

Dixon on the Eye.

A Guide to the Practical Study of Diseases of the Eye, with an Outline of their Medical and Operative Treatment, with Test Types and Illustrations. Third Edition, thoroughly Revised, and a great portion Rewritten. By JAMES DIXON, F.R.C.S., *Surgeon to the Royal London Ophthalmic Hospital, &c., &c.* In one volume. Price, . $2.50

"Mr. Dixon's book is essentially a practical one, written by an observant author, who brings to his special subject a sound knowledge of general Medicine and Surgery." — *Dublin Quarterly.*

Duchenne's Localized Electrization.

Translated from the Third Edition, by HERBERT TIBBITS, M.D., L.R. C.P., *Lond., Medical Superintendent of the National Hospital for the Paralyzed and Epileptic. With* 92 *Illustrations, and Notes and Additions by the Translator.* Price, ... $3.00

CONTENTS OF PART I.

CHAPTER II. LOCALIZED ELECTRIZATION. — Part I. The Fundamental Principles of the Method. Part 2. Localized Muscular Electrization. Part 3. Cutaneous Electrization. Part 4. Faradization of Internal Organs, the Genital Organs, and Organs of Sense. Part 5. The General Effects of Localized Faradization.

CHAPTER III. HISTORICAL AND CRITICAL OBSERVATIONS UPON THE PRINCIPAL METHODS OF ELECTRIZATION. — Part 1. On the Value of Electro-puncture as applied to the Treatment of Paralysis. Part 2. On the Value of Electrization by Reflex Action as applied to the Treatment of Paralysis. Part 3. On the Therapeutic Value of Localized Faradization.

CHAPTER IV. ELECTRO-MEDICAL INSTRUMENTS WITH REGARD TO THEIR APPLICATION IN PATHOLOGY, IN PHYSIOLOGY, AND IN THERAPEUTICS. — Part 1. The Properties which Instruments should possess. Part 2. The Author's large Double-current Volta-Faradic Instrument. Part 3. Small Double-current Volta-Faradic Instrument. Part 4. The Author's Double-current Magneto-Faradic Instrument. Part 5. Historical and Descriptive Account of Inductive Instruments.

This part of Duchenne's great work is a translation from the third edition now being prepared by the author, and contains all that has yet been printed, and is published even before the original is issued. It is not only a well-nigh exhaustive treatise on the medical uses of Electricity, but it is also an elaborate exposition of the different diseases in which Electricity has proved to be of value as a therapeutic and diagnostic agent. No similar treatise, it is believed, exists in the English language.

PART II., illustrated by chromo-lithographs and numerous wood-cuts, is preparing.

"Duchenne's great work has for years held its place as the most complete; and Dr. Tibbits, by presenting us with an English version of the forthcoming third edition, has earned the gratitude of the profession. His experience at the National Hospital gives great value to his Notes and Additions, and we are happy to congratulate him upon having so far accomplished an undertaking of great utility."—*The Lancet.*

Dillnberger's Handy-Book of the Treatment of Women and Children's Diseases, *according to the Vienna Medical School.*

Part I. The Diseases of Women. Part II. The Diseases of Children.

Translated from the Second German Edition, by P. NICOL, M.D.
One volume 12mo. Price, ... $1.75

"We noticed favorably the original of this hand-book some months ago, and suggested that an English translation of it, with notes showing the main points wherein the practice of our medical schools differs from that at Vienna, might be well received. Mr. Nicol has now carried out this idea, and we imagine that many practitioners will be glad to possess this little manual, which gives a large mass of practical hints respecting the treatment of diseases which probably make up the larger half of every-day practice. The translation is well and correctly performed, and the necessary explanations of reference to German medicinal preparations are given with proper fulness." —*The Practitioner.*

Durkee on Gonorrhœa and Syphilis.

The Fifth Edition, Revised and Enlarged, with Portraits and Colored Illustrations. By SILAS DURKEE, M.D., *Fellow of the Massachusetts Medical Society, &c., &c. A New and Revised Edition, with Eight Colored Illustrations.* Octavo, ... $5.00

"Dr. Durkee's work is one of those, the perusal of which impresses the reader in favor of the author. The general tone, the thorough honesty everywhere evinced, the philanthropic spirit observable in many passages, and the energetic advocacy of professional rectitude, speak highly of the moral excellence of the writer; nor is the reader less attracted by the skill with which the book is arranged, the manner in which the facts are cited, the clever way in which the author's experience is brought in, and the lucidity of the reasoning, the frequent and extremely fair allusions to the labors of others, and the care with which the therapeutics of venereal complaints are treated." — *Lancet.*

Fuller on Rheumatism, Rheumatic Gout, and Sciatica. A NEW EDITION PREPARING.

Their Pathology, Symptoms, and Treatment. By HENRY WILLIAM FULLER, M.D., *Fellow of the Royal College of Physicians, London; Physician to St. George's Hospital, &c. From the last London Edition.* Octavo, ...

Fuller on Diseases of the Heart and Great Vessels. *The Second Edition, Revised and greatly Enlarged, preparing.*

Gant's Science and Practice of Surgery.

A Complete System, including the Principles and Practice, by FREDERICK J. GANT, F.R.C.S., *Surgeon to the Royal Free Hospital, London, &c., &c. With* 400 *Illustrations.*

Dr. Gant's Surgery is similar in its general features to Dr. Errickson's excellent Text-book on the Science and Practice, as comprising the *whole* of surgery, but it will have the great advantage of being an entirely new work, and appearing in a greatly condensed form, without the omission of any important subject. The matter throughout is fully brought up to the present time, — many original sources of information, including the best treatises in Holmes' System of Surgery, having been utilized which have not heretofore been gathered into any general work on surgery, thus making it a more complete book for the student, and a work of ready reference for the practitioner. It is also illustrated by 250 new illustrations, which add much to its value.

Goff's Combined Day-Book, Ledger, *and Daily Register of Patients, combining not only the Accuracy and Essential Points of a regular Day-Book and Ledger System, without any of the labor and responsibility, but is also a Daily Register of Patients, &c., &c. A large Quarto Volume, strongly bound in half-russia.* Price, $12.00.

Gross' American Medical Biography of the Nineteenth Century.

Edited by SAMUEL D. GROSS, M.D., *Professor of Surgery in the Jefferson Medical College, Philadelphia, &c., &c. With a Portrait of* BENJAMIN RUSH, M.D. Octavo. $3.50

Greenhow on Bronchitis, *especially as Connected with Gout, Emphysema, and Diseases of the Heart.*

By E. HEADLAM GREENHOW, M.D., *Fellow of the Royal College of Physicians, &c., &c.*

Price, $2.00

"In vivid pictures of the sort of cases which a practitioner encounters in his daily walks, and in examples of the way in which a student ought to turn them over in his mind and make them tools for self-improvement, we have rarely seen a volume richer." — *Brit. and For. Medico-Chirurg. Review.*

Garratt's (Alfred C.) Guide for Using Medical Batteries.

Showing the most approved Apparatus, Methods, and Rules for the Medical Employment of Electricity in the Treatment of Nervous Diseases, &c., &c. With numerous Illustrations. One Volume, octavo. . . $2.00

"The large work on the same subject, and by the same author, is pretty well known to the Profession, but it is bulky and cumbrous, and by no means so practically useful. The present comparatively brief volume contains every thing of importance in regard to the various apparatuses useful to the Medical Electrician and the various modes of application for therapeutic purposes." — *Lancet and Observer.*

Hewitt on the Diseases of Women.

SECOND EDITION, REWRITTEN AND ENLARGED.

The Diagnosis and Treatment of Diseases of Women, including the Diagnosis of Pregnancy. Founded on a Course of Lectures delivered at St. Mary's Hospital Medical School. By GRAILY HEWITT, M. D. Lond., M. R. C. P., *Physician to the British Lying-in Hospital; Lecturer on Midwifery and Diseases of Women and Children at St. Mary's Hospital Medical School; Honorary Secretary to the Obstetrical Society of London, &c. With a new Series of Illustrations.*

Price, in cloth, $5.00; in leather, $6.00.

Hillier's Clinical Treatise on the Diseases of Children.

By THOMAS HILLIER, M.D., *Physician to the Hospital for Sick Children, and to University College Hospital, &c., &c.* Octavo.

Price, $3.00

"Our space is exhausted, but we have said enough to indicate and illustrate the excellence of Dr. Hillier's volume. It is eminently the kind of book needed by all medical men who wish to cultivate clinical accuracy and sound practice." — *London Lancet.*

Headland on the Action of Medicines in the System.

By F. W. HEADLAND, M.D., *Fellow of the Royal College of Physicians, &c., &c. Sixth American from the Fourth London Edition. Revised and enlarged.* One Volume, octavo. $3.00

Dr. Headland's work has been out of print in this country nearly two years, awaiting the revisions of the author, which now appear in this edition. It gives the only scientific and satisfactory view of the action of medicine; and this not in the way of idle speculation, but by demonstration and experiments, and inferences almost as indisputable as demonstrations. It is truly a great scientific work in a small compass, and deserves to be the handbook of every lover of the Profession. It has received the most unqualified approbation of the *Medical Press*, both in this country and in Europe, and is pronounced by them to be the most *original* and practically useful work that has been published for many years.

Hille's Pocket Anatomist.

Being a Complete Description of the Anatomy of the Human Body; for the Use of Students. By M. W. HILLES, *formerly Lecturer on Anatomy and Physiology at the Westminster Hospital School of Medicine.*

Price, in cloth, ... $1.00
" in Pocket-book form, ... 1.25

Heath on the Injuries and Diseases of the Jaws.

The Jacksonian Prize Essay of the Royal College of Surgeons of England, 1867. By CHRISTOPHER HEATH, F. R. C. S., *Assistant Surgeon to University College Hospital, and Teacher of Operative Surgery in University College. Containing over* 150 *Illustrations.* Octavo. Price, $6.00

Hodge on Fœticide, or Criminal Abortion.

By HUGH L. HODGE, M.D., *Emeritus Professor in the University of Pennsylvania. A Small Pocket Volume.* Price in paper covers, 30
" flexible cloth, 50

This little book is intended to place in the hands of professional men and others the means of answering satisfactorily and intelligently any inquiries that may be made of them in connection with this important subject.

Holmes' Surgical Diseases of Infancy and Childhood.

By J. HOLMES, M.A., *Surgeon to the Hospital for Sick Children, &c. Second Edition. Revised and Enlarged.* Octavo. Price, $9.00

Hufeland's Art of Prolonging Life.

Edited by ERASMUS WILSON, M.D., F.R.S. *Author of "A System of Human Anatomy," "Diseases of the Skin," &c., &c.* 12mo. Cloth. $1.25

Kirkes' Hand-Book of Physiology.

THE SEVENTH LONDON EDITION.

HAND-BOOK OF PHYSIOLOGY, by WILLIAM SENHOUSE KIRKES, M.D. *Seventh Edition, edited by* W. MORRANT BAKER, F.R.C.S., *Lecturer on Physiology, &c., &c.* With 241 Illustrations. In one volume, demy-octavo, containing over 800 pages. Price, bound in cloth, $5.00.

This edition of Dr. Kirkes' Hand-Book of Physiology is fully brought up to the times, and forms one of the most complete and convenient TEXT-BOOKS on the subject, for the Student of Medicine, now in print.

Lawson on the Eye.

The Diseases and Injuries of the Eye, their Medical and Surgical Treatment. By GEORGE LAWSON, F.R.C.S., *Surgeon to the Royal London Ophthalmic Hospital, &c., &c. With a Formulary, Test Types, and numerous Illustrations.* In one volume. Price, . . $2.50

"Dr. Lawson has succeeded in comprising within moderate limits a very complete and succinct account of the Diseases and Injuries of the Eye." "The many excellent features of this manual render it at once a safe and thoroughly trustworthy guide to the study of this special class of diseases, and from its eminently practical character it must be a great acquisition to the library of the busy practitioner. Its value is also much enhanced by an excellent selection of formulæ." — *Glasgow Med. Journal.*

Legg on Urine. FROM THE SECOND LONDON EDITION.

A Guide to the Examination of the Urine. For the Practitioner and Student. By J. WICKHAM LEGG, M.D., *Member of the Royal College of Physicians, &c., &c.* Second Edition. 16mo. Cloth. Price, 75 cts.

"Dr. Legg's little manual has met with remarkable success, and the speedy exhaustion of the first edition has enabled the author to make certain emendations which have added greatly to its value. We can now confidently commend it to the student as a safe and reliable guide to such examinations of the urine as he may be called upon to make." — *London Medical Times and Gazette.*

Mackenzie on Laryngeal Growths.

Growths in the Larynx, with Reports and an Analysis of 100 *Consecutive Cases treated by the author, and a Tabular Statement of every published case treated since the invention of the Laryngoscope. By* MORELL MACKENZIE, M.D., *author of "The Laryngoscope," "Diseases of the Throat," &c. Profusely illustrated by wood engravings and chromo-lithographs.* Octavo. Price, ... $3.00

Mackenzie on the Laryngoscope.

Its Use in Diseases of the Throat. With additions, and an Essay on Hoarseness, Loss of Voice, and Stridulous Breathing in relation to Nervo-Muscular Affections of the Larynx, by MORELL MACKENZIE, M.D., *Physician to the Hospital for Diseases of the Throat, &c., &c. Second Edition, with additions, and a Chapter on the Nasal Passages,* by J. SOLIS COHEN, M.D., *author of "Inhalation, Its Therapeutics and Practice," &c. Illustrated by two lithographic plates, and* 51 *engravings on wood.* Octavo. Price, ... $3.00

"While laryngoscopy was in its infancy, and before it had begun to engage to any extent the attention of the profession, it was studied with the greatest care and enthusiasm by the author of this treatise. Those who are anxious to study the diseases of the larynx and the mode of using the laryngoscope, cannot do better than purchase the treatise before us, as it is by far the best which has been published, and is thoroughly to be relied upon."—*Glasgow Medical Journal.*

Meigs and Pepper on Children.

FOURTH EDITION, ENLARGED AND IMPROVED.

The publishers have selected the following notice, from a late number of the LONDON LANCET, of the New Edition of this work, as indicating, perhaps, more fully than any other of the numerous favorable criticisms that have appeared of it elsewhere, its great value to the *Practitioner and Student of Medicine.*

"It is not necessary to say much, in the way of criticism, of a work so well known as 'Meigs on Diseases of Children,' especially when it has reached a fourth edition. Our duty is wellnigh restricted to the point of ascertaining how far, under an old color, it preserves the freshness and the value of a new book — how far it incorporates what is new with what is old without unseemly marks of mere joining. There is some advantage in starting entirely afresh, in being merely clinical, or in being very short, and limiting one's self to the expression of one's own views and experience. But such is not the nature of this book, and the advantages of it are different. It is a work of more than 900 good American pages, and is more encyclopædial than clinical. But it is clinical, and withal most effectually brought up to the light, pathological and therapeutical, of the present day.

"The book is like so many other good American medical books which we have lately had occasion to notice; it marvellously combines a *résumé* of all the best European literature and practice with evidence throughout of good personal judgment, knowledge, and experience. It is gratifying to see how our English authors are quoted, and especially how the labors of Hillier, who died so prematurely, are recognized. But the book abounds in exposition of American experience and observation in all that relates to the diseases of children. Not the least interesting additions to the volume are several extensive tables, exhibiting the mortality in Philadelphia of some of the most common and fatal diseases in connection with the variations of the temperature, and prepared with great care from the records of the Board of Health.

"The thoroughly fresh nature of the book is especially seen in the care with which certain articles have been written. Such are those on Rickets and Tuberculosis, Infantile Atrophic Paralysis, and Progressive Paralysis. No book now on diseases of children is complete which does not treat specially of constitutional or diathetic diseases, such as rickets and tuberculosis, syphilis, &c.

"Among other articles of great interest and value we would mention those on Diseases of the Cæcum and Appendix Vermiformis, on Indigestion in Children, on Diarrhœa, on Entero-Colitis, on Intussusception, on Chronic Hydrocephalus, and on Croup and the value of Tracheotomy.

"The difficulties of editing a new edition of a medical book of some standing are not more felt in the region of pathology and the classification of disease than in that of therapeutics. In this work this difficulty has been fairly faced by the authors. They have to confess to having changed their practice very materially in the treatment of acute diseases, to having given up mercury in most inflammatory diseases, and almost given up bloodletting. We recommend the views of these authors as to the injurious effects of calomel and antimony to careful consideration. They do not entirely abjure the use of bloodletting in certain cases of pneumonia and meningitis. Indeed, we think they will find reason in future editions to talk a little less freely than they do about bleeding and cupping very young children in certain circumstances of pneumonia, and in certain cases of simple meningitis. With a few exceptions of this kind, the therapeutics are sound and commendable, great importance being given to proper feeding and the general management of infancy and childhood. It is due to authors of so much fairness and experience to publish widely their opinion of the injurious and depressing effects of antimony in the inflammations of children. To infants under two years of age they think it best to give no antimony even in pneumonia. They do not use tartar emetic at all in the cases of children, but small doses, such as the twelfth of a grain, of the precipitated sulphuret of antimony, every two, three, or four hours, watching its effects, and withdrawing it quickly if symptoms of prostration appear, perhaps without any vomiting.

"We are glad to add this work to our library. There are few diseases of children which it does not treat of fully and wisely in the light of the latest physiological, pathological, and therapeutical science." — *London Lancet*, July 23, 1870

Price, handsomely bound in Cloth	$6 00
" " " Leather	7 00

LINDSAY & BLAKISTON, PUBLISHERS,

PHILADELPHIA.

Marshall's Physiological Diagrams.

LIFE-SIZE, AND BEAUTIFULLY COLORED.

On account of their large size and the great distinctness of the figures on them, there has been a growing demand in this country for these Maps for the LECTURE ROOM and for lecturing from in MEDICAL as well as OTHER SCHOOLS. In order to supply this demand on more favorable terms, we have recently completed an arrangement with the publishers in London, by which we can sell them to the trade and others at a reduced price and on better terms than heretofore.

The series, illustrating the whole Human Body, are *life-size*, each map printed on a single sheet of paper, made specially for the purpose, 7 *feet long* and 3 *feet* 9 *inches broad*, colored in fac-simile of the Original Drawings. There are nine diagrams, as follows·

No. 1. The Skeleton and Ligaments.
No. 2. The Muscles and Joints, with Animal Mechanics.
No. 3. The Viscera in Position. — The Structure of the Lungs.
No. 4. The Heart and principal Blood-vessels.
No. 5. The Lymphatics or Absorbents.
No. 6. The Digestive Organs.
No. 7. The Brain and Nerves.
No. 8. The Organs of Sense and Voice.
No. 9. The Textures. — Microscopic Structures.

Prepared under the direction of JOHN MARSHALL, F.R.S., F.R.C.S., *Professor of Surgery, University College, and Surgeon to University College Hospital.*

Price of the Set, Nine Maps, in Sheets, ... $50.00
" " " " handsomely Mounted on Canvas, with Rollers, and Varnished, ... $80.00

Though designed more especially for purposes of general education, supplying an acknowledged necessity of modern teaching, these diagrams will be found not inapplicable to the requirements of professed Medical Schools, affording, as they do, a correct preliminary view of the various systems and organs in the human body. For PUBLIC SCHOOL PURPOSES, for LECTURES at LITERARY, SCIENTIFIC, and other Institutes, they will be found invaluable; and also to students of Artistic Anatomy, imparting, as they do, when suspended on the walls of the Lecture-hall, School-room, or Studio, a familiar acquaintance with the whole human system.

An Explanatory Key to the Physiological Diagrams.

By JOHN MARSHALL, F.R.S., F.R.C.S., &c. Octavo. Paper covers. . . . . 50 cts.

Description of the Human Body.

Its Structure and Functions. Illustrated by Physiological Diagrams, Designed for the Use of Teachers in Schools and Young Men destined for the Medical Profession, and for popular Instruction generally. New Edition. By JOHN MARSHALL, F.R.S., F.R.C.S., *Professor of Surgery, University College, and Surgeon to the University College Hospital.*

The work contains 260 *quarto pages of Text, bound in cloth, and* 193 *Colored Illustrations, arranged in Nine Folio Diagrams, carefully colored and reduced from Prof. Marshall's large work.* 2 vols. Cloth. . $10.00

Murphy's Review of Chemistry for Students.

Adapted to the Courses as Taught in the Principal Medical Schools in the United States. By JOHN G. MURPHY, M.D. . . . $1.25

Meadows' Manual of Midwifery. A New Text-Book.

Including the Signs and Symptoms of Pregnancy, Obstetric Operations, Diseases of the Puerperal State, &c., &c. By ALFRED MEADOWS, M.D., *Member of the Royal College of Physicians, &c., &c. First American from the Second London Edition. With numerous Illustrations.* Price, ... $3.00

"Those who read the first edition of this work will bear us out in thinking that Dr. Meadows's Manual forms one of the most convenient, practical, and concise books yet published on the subject. It was especially good as a student's manual, and the author has, in his second edition, sought to make it of equal value to the practitioner. The part which treats of obstetric operations has been well revised, and has received numerous additions, and the several chapters on Unnatural and Complex Labors likewise comprise much new matter. Upwards of ninety new engravings have been inserted in this edition, and, with a view to facilitate reference, the author has furnished it with a very full and complete table of contents and index. We can cordially recommend this manual as accurate and practical, and as containing in a small compass a large amount of the kind of information suitable alike to the student and practitioner."—*London Lancet*, May 6, 1871.

"This new edition of a book which was at once recognized as a good manual, is a considerable improvement on its predecessor. It is eminently a book which will teach the student. . . . Not merely is the practical treatment of Labor, and also of the Diseases and Accidents of Pregnancy, well and clearly taught, but the anatomical machinery of parturition is more effectively explained than in any other treatise that we remember; and besides this, the book is honorably distinguished among manuals of Midwifery by the fulness with which it goes into the subject of the structure and development of the ovum. Dr. Meadows has done good service in giving a clear account of this subject in a very short space, yet with sufficient fulness. On all questions of treatment, whether by medicines, by hygienic regimen, or by mechanical or operative appliances, this treatise is as satisfactory as a work of manual size could be; and altogether, students and practitioners can hardly do better than adopt it as their vade-mecum."—*The Practitioner.*

Maxson's Practice of Medicine.

By EDWIN R. MAXSON, M.D., *formerly Lecturer on the Practice of Medicine in the Geneva Medical College, &c.* . . $4.00

Morris on Scarlet Fever.

Its Pathology and Therapeutics. By CASPER MORRIS, M.D., *Fellow of the College of Physicians of Philadelphia, &c.* . . . $1.50

Mendenhall's Medical Student's Vade Mecum.

A Compendium of Anatomy, Physiology, Chemistry, the Practice of Medicine, Surgery, Obstetrics, Diseases of the Skin, Materia Medica, Pharmacy, Poisons, &c., &c. By GEORGE MENDENHALL, M.D., *Professor of Obstetrics in the Medical College of Ohio, &c., &c. Ninth Edition, Revised and Enlarged, with* 224 *Illustrations.* . $2.50

Pennsylvania Hospital Reports.

Edited by a Committee of the Hospital Staff, J. M. DACOSTA, M.D., and WILLIAM HUNT, M.D. *Vols.* 1 *and* 2, *for* 1868 *and* 1869, *each volume containing upwards of* Twenty *Original Articles, by former and present Members of the Staff, now eminent in the Profession, with Lithographic and other Illustrations.* Price per volume, . . $4.00

At last, however, the work has been commenced, the Philadelphia Physicians being the first to occupy this field of usefulness. The first Reports were so favorably received, on both sides of the Atlantic, that it is hardly necessary to speak for them the universal welcome of which they are deserving. The papers are all valuable contributions to the literature of medicine, reflecting great credit upon their authors. The work is one of which the Pennsylvania Hospital may well be proud. It will do much toward elevating the profession of this country.—*American Journal of Obstetrics.*

Pereira's Physician's Prescription Book.

Containing Lists of Terms, Phrases, Contractions, and Abbreviations, used in Prescriptions, with Explanatory Notes, the Grammatical Constructions of Prescriptions, Rules for the Pronunciation of Pharmaceutical Terms, A Prosodiacal Vocabulary of the Names of Drugs, etc., and a series of Abbreviated Prescriptions illustrating the use of the preceding terms, etc.; to which is added a Key, containing the Prescriptions in an unabbreviated Form, with a Literal Translation, intended for the use of Medical and Pharmaceutical Students. By JONATHAN PEREIRA, M.D., F.R.S., etc. *From the Fifteenth London Edition.*

Price, in cloth, ... $1.25
" in leather, with Tucks and Pocket, . . . 1.50

This little work has passed through fifteen editions in London and several in this country. The present edition of which this is a reprint has been carefully revised and many additions made to it. Its great value is proven both by its large sale and the many favorable notices of it in the Medical Press.

Paget's Surgical Pathology.

Lectures delivered at the Royal College of Surgeons of England, by JAMES PAGET, F.R.S., *Surgeon to Bartholomew and Christ's Hospital, &c., &c. The Third American Edition, Edited and Revised by* WILLIAM TURNER, M.B., *Lond. Senior Demonstrator of Anatomy in the University of Edinburgh, &c., &c. In one volume, Royal Octavo With numerous Illustrations.*

Price, in cloth, ... $6.00
" leather, ... 7.00

"It would be very superfluous for us to say many words in calling the attention of the profession to this new edition of Mr. Paget's great work on Surgical Pathology Its author has been singularly fortunate in securing the assistance (for this edition) of so able a collaborator as Mr. Turner, and English surgery may point with pride to the present volume as one unsurpassed, if it is at all equalled in the surgical literature of the world, in breadth of view and philosophical grasp of its subject." — *Practitioner.*

Prince's Plastic and Orthopedic Surgery.

Containing,

1. *A Report on the Condition of, and Advances made in, Plastic and Orthopedic Surgery up to the Year* 1871.
2. *A New Classification and Brief Exposition of Plastic Surgery. With numerous Illustrations.*
3. ORTHOPEDICS: *A Systematic Work upon the Prevention and Cure of Deformities. With numerous Illustrations.*

In one volume, Octavo. Price, ... $4.50

"This is a good book, upon an important practical subject; carefully written, abundantly illustrated, and well printed. It goes over the whole ground of deformities of all degrees — from cleft-palate and club-foot, to spinal curvatures and ununited fractures. It appears, moreover, to be an original book, so far as one chiefly of compilation can be so. Such a book was wanted, and it deserves success." — *Med. and Surg. Reporter.*

Rindfleisch's Text-Book of Pathological Histology.

An Introduction to the Study of Pathological Anatomy. By Dr. EDWARD RINDFLEISCH, *O. O. Professor of Pathological Anatomy in Bonn. Translated from the Second German Edition*, by WM. C. KLOMAN, M. D., assisted by F. T. MILES, M. D., *Professor of Anatomy, University of Maryland, &c., &c.*

CONTENTS.

Introduction, Author's and Editor's Prefaces.

GENERAL PART.

1. Decomposition and Degeneration of Tissues.
2. Pathological New Formations.

SPECIAL PART.

1. Anomalies of the Blood and the Places of its Formation, especially of the Spleen and Lymphatic Glands.
2. Anomalies of the Circulatory Apparatus.
3. Anomalies of Serous Membranes.
4. " the Skin.
5. " Mucous Membranes.
6. " the Lung.
7. " " Liver.
8. " " Kidney.
9. Anomalies of the Ovaries.
10. " " Testicles.
11. " " Mammæ.
12. " " Prostate Gland.
13. " " Salivary Glands.
14. " " Thyroid Gland.
15. " " Suprarenal Capsules.
16. " " Osseous System.
17. " " Nervous System.
18. " " Muscular System.

Index and Bibliography.

Containing 208 Elaborately Executed Microscopical Illustrations.

One volume, octavo. Price,

Prof. Rindfleisch's Text-Book of Pathological Histology, so justly celebrated in Germany, where it is considered the most complete and thorough work of its kind, having passed rapidly to a second edition, is also very highly valued and commended by German Medical scholars in this country, many of whom are not only familiar with the book, but with the author's great reputation as a teacher and professor of this branch of medical study.

The translators are both gentlemen who by their past education have been peculiarly fitted for the task of translating the work. Dr. Kloman from early life has been familiar with the German language, while Prof. Miles has made the subject one of special study, both gentlemen being also practical microscopists. The Publishers therefore offer a translation of this truly valuable work to the Medical Profession in the United States, feeling the utmost confidence that in both manner and style it will prove acceptable to them. In their Preface, the Translators say: "In presenting the English reading portion of the Medical Profession with a translation of the valuable work of Prof. Rindfleisch, the translators scarcely deem an apology necessary. The merits of the book itself, and the fact that it fills an unoccupied gap in our most recent literature upon the subject of Pathological Histology, was judged to be an ample incentive for undertaking the labor of the translation. The work of VIRCHOW translated by CHANCE, is, in many points, antiquated, and the more recent work of BILBROTH, translated by HACKLEY, occupies the ground but partially, and is professedly a work of Surgical Pathology."

This book is translated and published in this country by special arrangement with the author.

Radcliffe's Lectures on Epilepsy, Pain, Paralysis,

And certain other Disorders of the Nervous System. By CHARLES BLAND RADCLIFFE, M.D., *Fellow of the Royal College of Physicians of London, &c., &c. With Illustrations.* $2.00

"The reputation which Dr. Radcliffe possesses as a very able authority on nervous affections, will commend his work to every medical practitioner. We recommend it as a work that will throw much light upon the Physiology and Pathology of the Nervous System." — *Canada Medical Journal*

Robertson's Manual on Extracting Teeth.

Founded on the Anatomy of the Parts involved in the Operation; the Kinds and Proper Construction of the Instruments to be used; the Accidents liable to occur from the Operation, and the Proper Remedies. By ABRAHAM ROBERTSON, D.D.S., M.D. *Second Edition, Revised and Improved. With Illustrations.* . . $1.50

Ranking's Half-yearly Abstract of the Medical Sciences. Price per annum, in advance, . $2.50

Back Volumes or Numbers furnished.

Renouard's History of Medicine.

From its Origin to the Nineteenth Century. With an Appendix containing a Philosophical and Historical Review of Medicine to the present time. By P. V. RENOUARD, M.D. *Translated from the French by* CORNELIUS G. COMEGYS, M.D., *Professor of the Institutes of Medicine in the Medical College of Ohio, &c.* Octavo. Price, . $4.00

Reports on the Progress of Medicine and Surgery.

Including Physiology, Practical Medicine, Surgery, Ophthalmic Medicine, Midwifery, Diseases of Women and Children, Materia Medica, Medical Jurisprudence, and Public Health. Edited by Drs. POWER, HOLMES, AINSTIE, BARNES, WINDSOR, &c., &c., *under the patronage and direction of the* SYDENHAM SOCIETY *of London.* One volume. Octavo. Price, ... $2.00

Ryan's Philosophy of Marriage.

In its Social, Moral, and Physical Relations, with an Account of the Diseases of the Genito-Urinary Organs. The Physiology of Generation in the Animal and Vegetable Kingdoms, &c., &c. By MICHAEL RYAN, M.D., *Member of the Royal College of Physicians and Surgeons in London, &c.* 12mo. $1.00

Reese's American Medical Formulary. $1.50

Reese's Analysis of Physiology.

Being a Condensed View of the most Important Facts and Doctrines, designed especially for the Use of Students. By JOHN J. REESE, M.D., *Professor of Medical Jurisprudence, including Toxicology, in the University of Pennsylvania, &c., &c. Second Edition, Enlarged.* $1.50

Reese's Syllabus of Medical Chemistry. $1.00

Stillé's Epidemic Meningitis;

Or, Cerebro-Spinal Meningitis. By ALFRED STILLÉ, M.D., *Professor of the Theory and Practice of Medicine in the University of Pennsylvania, &c., &c.* In one volume, Octavo, ... $2.00

"This monograph is a timely publication, comprehensive in its scope, and presenting within a small compass a fair digest of our existing knowledge of the disease, particularly acceptable at the present time. It is just such a one as is needed, and may be taken as a model for similar works."—*Am. Journal Med. Sciences.*

Stillé's Elements of General Pathology.

A Practical Treatise on the Causes, Forms, Symptoms, and Results of Disease. Second Edition preparing.

Sweringen's Pharmaceutical Dictionary.

A Pharmaceutical Lexicon or Dictionary of Pharmaceutical Science, containing a Concise Explanation of the various Subjects and Terms of Pharmacy, with Collateral Selections from the Kindred Sciences; also Formulæ for Officinal, Empirical, and Dietetic Preparations, Antidotes to Poisons, &c., &c. By HIRAM V. SWERINGEN, *Member of the American Pharmaceutical Association, &c., &c.* In preparation.

Sansom on Chloroform.

Its Action and Administration. By ARTHUR ERNEST SANSOM, M.B., *Physician to King's College Hospital, &c., &c.* 12mo. . $2.00

"The work of Dr. Sansom may be characterized as most excellent. Written not alone from a theoretical point of view, but showing very considerable experimental study, and an intimate clinical acquaintance with the administration of these remedies, — passing concisely over the whole ground, giving the latest information upon every point, — it is just the work for the student and practitioner."—*Amer. Medical Journal.*

Scanzoni on Women.

A Practical Treatise on the Diseases of the Sexual Organs of Women. Translated from the French. By A. K. GARDNER, A.M., M.D., &c. *With Illustrations.* Octavo, ... $5.00

Stokes on the Diseases of the Heart

And the Aorta. By WILLIAM STOKES, *Regius Professor of Physic in the University of Dublin; Author of the Diseases of the Chest, &c., &c. Second American Edition.* Octavo, ... $3.00

THOMAS HAWKES TANNER'S WORKS.

"The leading feature of Dr. Tanner's books is their essentially practical character."
LONDON LANCET.

The Practice of Medicine. JUST READY.

FIFTH AMERICAN, FROM THE SIXTH LONDON EDITION.

Revised, much Enlarged, and thoroughly brought up to the present time. With a complete Section on the Diseases Peculiar to Women added; also an extensive Appendix of Formulæ for Medicines, Baths, Mineral Waters, Climates, &c. By THOMAS HAWKES TANNER, M.D., *Fellow of the Royal College of Physicians, &c.* One Volume. Royal Octavo, containing over 1100 pages. Price, handsomely bound in Cloth, . . $6.00
" " " Leather, . $7.00

There is a common character about the writings of Dr. Tanner — a character which constitutes one of their chief values: they are all essentially and thoroughly practical. Dr. Tanner never, for one moment, allows this utilitarian end to escape his mental view. He aims at teaching how to recognize and how to cure disease, and in this he is thoroughly successful. . . . It is, indeed, a wonderful mine of knowledge.—*Medical Times.*

Dr. Tanner has always shown in his writings that he possesses a peculiar faculty of committing to print just that kind of information which the practitioner most needs in everyday practice, and of rejecting useless theories or hypothetical statements.—*Lancet.*

The author has, in the manner in which he has dealt with the subject, given another evidence of that happy facility which he possesses of giving the essential points of a mass of information in a well connected and instructive form. — *Brit. Med. Journ.*

The student will find the work the best text-book on the practice of medicine, while the practitioner will possess in it a thoroughly safe guide at the bedside. — *Dub. Med. Quarterly.*

Tanner's Practical Treatise on the Diseases of Infancy and Childhood. PRICE, $3.50.

THIRD AMERICAN EDITION, REVISED AND ENLARGED.

By ALFRED MEADOWS, M.D., *London*, M.R.C.P., *Physician to the Hospital for Women and to the General Lying-in Hospital, &c., &c.*

"This book of Dr. Tanner's has been much enlarged and the plan altered by Dr. Meadows. As it now stands it is probably one of the most complete in our language. It no longer deals with children's diseases only, but includes the peculiar conditions of childhood, both normal and abnormal, as well as the therapeutics specially applicable to that class of patients. The articles on Skin Diseases have been revised by Dr. Tilbury Fox, and those on Diseases of the Eye by Dr. Brudenell Carter, both gentlemen distinguished in these specialties." — *Medical Times and Gazette.*

Tanner's Index of Diseases and their Treatment. *With upwards of* 500 *Formulæ for Medicines, Baths, Mineral Waters, Climates for Invalids, &c., &c.* Octavo. . . $3.00

To the busy practitioner it must be an advantage to see at a glance on a quarter or half a page the principal point in any disease about which he may wish to have his memory refreshed or his mind stimulated. It will be found a most valuable companion to the judicious practitioner. — *The Lancet.*

Tanner's Memoranda of Poisons.

From the Second London Edition. . . . . . 50 cts

Tilt's Change of Life

IN HEALTH AND DISEASE. *A Practical Treatise on the Nervous and other Affections incidental to Women at the Decline of Life.* By EDWARD JOHN TILT, M.D. *From the Third London Edition. In one volume.* Octavo, ... $3.00

The work is rich in personal experience and observation, as well as in ready and sensible reflection on the experience and observation of others. The book is one that no practitioner should be without, as the best we have on a class of diseases that makes a constant demand upon our care, and requires very judicious management on the part of the practitioner. — *London Lancet.*

The great abilities of Dr. Tilt, his extensive knowledge, and his literary power are well shown in this book, and we are pleased to say that it has substantial and peculiar merits. It contains many sage, practical recommendations, and will amply repay perusal. — *Edinburgh Medical Journal.*

Dr. Tilt has been a very earnest and a very faithful worker in the physiology and diseases of women, and has made contributions to the literature of the subject which are all of accepted value. — *British Medical Journal.*

Tyler Smith's Obstetrics.

A Course of Lectures. By W. TYLER SMITH, M.D., *Physician, Accoucheur, and Lecturer on Midwifery, &c. Edited* by A. K. GARDNER, M.D. *With Illustrations.* Octavo, ... $5.00

Toynbee on Diseases of the Ear.

Their Nature, Diagnosis, and Treatment. A new London Edition, with a Supplement. By JAMES HINTON, *Aural Surgeon to Guy's Hospital, &c. With Illustrations.* Octavo, ... $5.00

Thompson's Clinical Lectures on Pulmonary Consumption. Octavo, ... $2.00

Tyson's Cell Doctrine:

Its History and Present State, with a Copious Bibliography of the Subject, for the use of Students of Medicine and Dentistry. By JAMES TYSON, M.D., *Lecturer on Microscopy in the University of Pennsylvania, &c., &c. With a Colored Plate, and numerous Illustrations on Wood.* Price, ... $2.00

Dr. Tyson furnishes in this work a concise and instructive *resumé* of the origin and advance of the doctrine of Cell Evolution. In it we find the theories of Virchow, Robin, Huxley, Hughes, Bennett, Beale, and other distinguished men. Its pages contain what could otherwise only be learned by the perusal of many works, and they supply the reader with a continuous, complete, and general knowledge of the history, progress, and peculiar phases of the Cell Doctrine, accompanied by careful references and a copious bibliography.

Virchow's Cellular Pathology.

Translated from the Second Edition. By FRANK CHANCE, B.A., M.A., &c. *With Notes and Emendations, and* 144 *Engravings.* 8vo. $5.00

Trousseau's Clinical Lectures.

VOL. IV. NOW READY.

Lectures on Clinical Medicine, delivered at the Hôtel-dieu, Paris. By A. TROUSSEAU, *Professor of Clinical Medicine in the Faculty of Medicine, Paris, &c., &c.*

Trousseau's Lectures on Clinical Medicine, so favorably received, as well by the profession of the United States as abroad, are published in this country in connection with the New Sydenham Society, under whose auspices the translation of Vols. II. and III. have been made. Either of these volumes can be furnished separately, and in order to still further extend the circulation of so valuable a work, the Publishers have now reduced the price to *Five Dollars* per volume.

CONTENTS OF VOLUME I.—*Translated and Edited by P. Victor Bazire, M. D., &c.*—LECTURE 1. On Venesection in Cerebral Hæmorrhage and Apoplexy. 2. On Apoplectiform Cerebral Congestion, and its Relations to Epilepsy and Eclampsia. 3. On Epilepsy. 4. On Epileptiform Neuralgia. 5. On Glosso-laryngeal Paralysis. 6. Progressive Locomotor Ataxy. 7. On Aphasia. 8. Progressive Muscular Atrophy. 9. Facial Paralysis, or Bell's Paralysis. 10. Cross-paralysis, or Alternate Hemiplegia. 11. Infantile Convulsions. 12. Eclampsia of Pregnant and Parturient Women. 13. On Tetany. 14. On Chorea. 15. Senile Trembling and Paralysis Agitans. 16. Cerebral Fever. 17. On Neuralgia. 18. Cerebral Rheumatism. 19. Exophthalmic Goître, or Graves' Disease. 20. Angina Pectoris. 21. Asthma. 22. Hooping Cough. 23. On Hydrophobia.

CONTENTS OF VOLUME II.—*Translated from the Edition of* 1868 (*being the last revised and enlarged edition*), *by John Rose Cormack, M. D., Edin., F.R.S.E., &c.* — LECTURE 1. Small-pox. 2. Variolous Inoculation. 3. Cow-pox. 4. Chicken-pox. 5. Scarlatina. 6. Measles, and in particular its unfavorable Symptoms and Complications. 7. Rubeola. 8. Erythema Nodosum. 9. Erythema Papulatum. 10. Erysipelas, and in particular Erysipelas of the Face. 11. Mumps. 12. Urticaria. 13. Zona, or Herpes Zoster. 14. Sudoral Exanthemata. 15. Dothinenteria, or Typhoid Fever. 16. Typhus. 17. Membranous Sore Throat, and in particular Herpes of the Pharynx. 18. Gangrenous Sore Throat. 19. Inflammatory Sore Throat. 20. Diphtheria. 21. Thrush.

CONTENTS OF VOLUME III.—*Translated from the Edition of* 1868, *by John Rose Cormack, M.D., Edin., F.R.S.E., &c.*—LECTURE 22. Specific Element in Disease. 23. Contagion. 24. Ozæna. 25. Stridulous Laryngitis, or False Croup. 26. Œdema of the Larynx. 27. Aphonia: Cauterization of the Larynx. 28. Dilatation of the Bronchi and Bronchorrhœa. 29. Hemoptysis. 30. Pulmonary Phthisis. 31. Gangrene of the Lung. 32. Pleurisy: Paracentesis of the Chest. 33. Traumatic Effusion of Blood into the Pleura: Paracentesis of the Chest. 34. Hydatids of the Lung. 35. Pulmonary Abscesses and Peripneumonic Vomicæ. 36. Treatment of Pneumonia. 37. Paracentesis of the Pericardium. 38. Organic Affections of the Heart. 52. Alcoholism. 62. Spermatorrhœa. 63. Nocturnal Incontinence of Urine. 64. Glucosuria: Saccharine Diabetes. 65. Polydipsia. 67. Vertigo a Stomacho Læso.

4 Volumes Octavo, handsomely bound in Cloth, Price $5 00 each.

OPINIONS OF THE PRESS.

"Trousseau furnishes us with an example of the best kind of Clinical teaching. It is a book that deserves to be popularized. The translation is perfect." — *Medical Times and Gazette.*

"The great reputation of Prof. Trousseau as a practitioner and teacher of Medicine in all its branches, renders the present appearance of his Clinical Lectures particularly welcome." — *Medical Press and Circular.*

"The publication of Trousseau's Lectures will furnish us with one of the very best practical treatises on disease as seen at the bedside." — *British and Foreign Medico-Chirurgical Review.*

"A clever translation of Prof. Trousseau's admirable and exhaustive work, the best book of reference upon the Practice of Medicine." — *Indian Medical Gazette.*

"The Lectures of Trousseau, in attractiveness of manner and richness of thoroughly practical matter, worthily takes a place beside the classical lectures of Watson and Graves." — *British Medical Journal.*

"Trousseau is essentially the French Graves, and his lectures should sooner than this have been translated into English." — *Lancet.*

Wythes' Physician's Pocket, Dose, and Symptom Book. THE TENTH EDITION.

Containing the Doses and Uses of all the Principal Articles of the Materia Medica, and Original Preparations; A Table of Weights and Measures, Rules to Proportion the Doses of Medicines, Common Abbreviations used in Writing Prescriptions, Table of Poisons and Antidotes, Classification of the Materia Medica, Dietetic Preparations, Table of Symptomatology, Outlines of General Pathology and Therapeutics, &c. By JOSEPH H. WYTHES, A.M., M.D., &c. *The Tenth Revised Edition*

Price, in cloth, ... $1.00
" leather, tucks, with pockets, . . . 1.25

This little manual has been received with much favor, and a large number of copies sold. It was compiled for the assistance of students, and to furnish a vade mecum for the general practitioner, which would save the trouble of reference to larger and more elaborate works. The present edition has undergone a careful revision. The therapeutical arrangement of the Materia Medica has been added to it, together with such other improvements as it was thought might prove of value to the work.

Waring's Practical Therapeutics.

NEW EDITION.

Considered chiefly with reference to Articles of the Materia Medica. By EDWARD JOHN WARING, F.R.C.S., F.L.S., &c., &c. *Second American, from the Third London Edition.* Royal Octavo.

Price, in cloth, $5.00. Price, in leather, $6.00.

There are many features in Dr. Waring's Therapeutics which render it especially valuable to the Practitioner and Student of Medicine, much important and reliable information being found in it not contained in similar works; it also differs from them in its completeness, the convenience of its arrangement, and the greater prominence given to the medicinal application of the various articles of the Materia Medica in the treatment of morbid conditions of the Human Body, &c. It is divided into two parts, the alphabetical arrangement being adopted throughout; there is also added an excellent INDEX OF DISEASES, with a list of the medicines applicable as remedies, and a full INDEX of the medicines and preparations noticed in the work.

"This new edition of Waring's *Practical Therapeutics* has been altered and improved with great judgment. A satisfactory account of new agents — chloral, apomorphia, nitrous oxide, carbolic acid, &c., is introduced without adding to its bulk. The additions are made with remarkable skill in condensation. It is one of the best manuals of therapeutics yet in existence." — *British Medical Journal.*

"Our admiration, not only for the immense industry of the author, but also of the great practical value of the volume, increases with every reading or consultation of it. We wish a copy could be put in the hands of every student or practitioner in the country. In our estimation it is the best book of the kind ever written." — *N. Y. Medical Journal.*

Walker on Intermarriage.

Or, the Mode in which, and the Causes why, Beauty, Health, and Intellect result from certain Unions, and Deformity, Disease, and Insanity from others. With Illustrations. By ALEXANDER WALKER, *Author of "Woman," "Beauty," &c., &c.* 12mo. $1.50

Walton's Operative Ophthalmic Surgery.

By HAYNES WALTON, F.R.C.S., *Surgeon to the Central London Ophthalmic Hospital, &c. With* 169 *Illustrations. Edited* by S. LITTELL, M.D., *Surgeon to the Wills Hospital for the Diseases of the Eye, &c.* Octavo. $4.00

"It is eminently a practical work, evincing in its author great research, a thorough knowledge of his subject, and an accurate and most observing mind." — *Dublin Quarterly Journal.*

Watson's Practice abridged.

A Synopsis of the Lectures on the Principles and Practice of Physic. Delivered at King's College, London, by THOMAS WATSON, M.D., *Fellow of the Royal College of Physicians, &c., &c. From the last London Edition. With a concise but Complete Account of the Properties, Uses, Preparations, Doses, &c., of all the Medicines mentioned in these Lectures, and other Valuable Additions,* by J. J. MEYLOR, A.M., M.D., &c., &c. *A neat Pocket Volume bound in cloth flexible.* . . . $2.00

Wells' Treatise on the Diseases of the Eye,

illustrated by Ophthalmoscopic Plates and Numerous Engravings on Wood. By J. SŒLBERG WELLS, *Ophthalmic Surgeon to King's College Hospital, &c.* Second London Edition, cloth, $6.50; leather, $7.50.

This is the author's own edition, printed in London under his supervision, and issued in this country by special arrangement with him.

Wright on Headaches.

Their Causes and their Cure. By HENRY G. WRIGHT, M.D., *Member of the Royal College of Physicians, &c. &c.* From the Fourth London Edition. 12mo. Cloth. $1.25

"Few affections are more unmanageable and more troublesome than those of which this essay treats; and we doubt not that any suggestions by which we can relieve them will be gladly received by physicians. The author's plan is simple and practical. He treats of headaches in childhood and youth, in adult life and old age, giving in each their varieties and symptoms, and their causes and treatment. It is a most satisfactory monograph, as the mere fact that this is a reprint of the *fourth* edition, testifies.

"The great pains which the author takes to clear up the differential diagnosis of the different varieties, and establish a satisfactory basis for rational treatment, are everywhere visible. While such a valuable fund of information is offered to the practitioner at the cost of a single visit, he should not let his patient suffer for want of it." — *Medical and Surgical Reporter.*

Wells on Long, Short, and Weak Sight, *and their Treatment by the Scientific Use of Spectacles. Third Edition Revised, with Additions and Numerous Illustrations.* By J. SŒLBERG WELLS. Octavo. . . . . Price, $3.00

DENTAL BOOKS PUBLISHED BY

LINDSAY & BLAKISTON,

No. 25 South Sixth Street, above Chestnut,

PHILADELPHIA.

Harris's Principles and Practice of Dentistry.

THE TENTH REVISED EDITION.

In great part Re-written, Re-arranged, and with many New and Important Illustrations.

INCLUDING

1. Dental Anatomy and Physiology.
2. Dental Pathology and Therapeutics.
3. Dental Surgery.
4. Dental Mechanics.

By Chapin A. Harris, M.D., D.D.S., &c. *The tenth edition, revised and edited by* P. H. Austen, M.D., *Professor of Dental Science and Mechanism in the Baltimore College of Dental Surgery, with nearly* 400 *Illustrations, including many new ones made especially for this edition.*

One volume, Royal Octavo, bound in Cloth, ... $6.50
" " " Leather, . . . 7.50

The subscribers, in presenting to *the Dental Profession* The Tenth Revised Edition of the late Prof. Harris's Text-Book of Dentistry, desire to call their attention to the thorough revision, and modification in its arrangement which it has been found necessary to make, in order to bring it fully up to the present advanced state of the science, and to preserve for it that reputation which it has for so many years sustained, as being the first and leading book on the subject in the English language.

So great have been the advances in Dental Physiology, Pathology, Surgery, and Mechanism, that a revision in all its parts, more complete and thorough than has been made to any previous edition — amounting, in some cases, to a re-writing of the articles or subjects — had become imperative; and in order to have this done in the most efficient manner, the publishers were, fortunately, able to secure the valuable services of Prof. Austen, of the Baltimore Dental College, who, added to his experience of twenty years as a teacher, was, for a long time, the associate and friend of Dr. Harris. He was thus peculiarly fitted for this task. In order, however, to render the work the more perfect and acceptable to the Profession, Dr. Austen associated with him Prof. Gorgas and Thomas S. Latimer, M.D., of the same school, both gentlemen of acknowledged proficiency in their respective departments, and further obtained valuable original articles from Prof. Kingsley, of the New York Dental College, whose reputation is well known, and other gentlemen of the Profession of equal repute in their respective specialties. The publishers, therefore, offer this edition to the Profession with the utmost confidence that it will be found most complete in every respect as a text-book for the student and a guide and companion for the experienced practitioner.

Harris's Dictionary.

THE THIRD REVISED EDITION.

A DICTIONARY OF MEDICAL TERMINOLOGY, DENTAL SURGERY, AND THE COLLATERAL SCIENCES. BY CHAPIN A. HARRIS, M.D., D.D.S., *Professor of the Principles of Dental Surgery in the Baltimore College, Member of the American Medical Association, &c., &c. The Third Edition, carefully revised and enlarged, by* FERDINAND J. S. GORGAS, M.D., D.D.S., *Professor of Dental Surgery in the Baltimore College, &c., &c.* In one volume, Royal Octavo, bound in Cloth, . $6.50
" " " " Leather, . 7.50

For a long period after the death of Dr. Harris his Dictionary remained out of print, but a constantly increasing demand for it from the Profession and from students urged the publishers to the preparation of a new edition. The many advances in dental science made during this interim rendered the incorporation of many new terms and formulæ absolutely necessary to the student of Dentistry, as well as to the dental practitioner. The present edition has been thoroughly revised by Prof. Gorgas, Dr. Harris's successor in the Baltimore Dental College; and in a very satisfactory manner, nearly three thousand new words have been incorporated into it, besides additions and corrections being made to many others. The doses of the more prominent medicinal agents have also been added, and in every way the book has been greatly improved, and its value enhanced as a work of reference. To those studying or practising Dentistry it must prove invaluable.

Heath on the Injuries and Diseases of the Jaws.

The Jacksonian Prize Essay of the Royal College of Surgeons of England, 1867. BY CHRISTOPHER HEATH, F.R.C.S., *Assistant Surgeon to University College Hospital, and Teacher of Operative Surgery in University College. Containing over* 150 *Illustrations.* Octavo, . . $6.00

This work is of the highest practical value. It treats of dislocations, fractures, deformities, and other disorders of the jaws, with the means for relief, illustrated by drawings of displacements, apparatus, morbid growths, portraits of patients, etc., to the number of 154. It embodies American as well as European experience, and contains quite a large appendix of cases in detail. The practical character of its contents, with good paper, print, and engravings, strongly commend this book to the profession. Every surgeon and dentist should have a copy. — *Dental Cosmos.*

It is impossible to over-estimate the value and importance of this comprehensive work. — *British Journal of Dental Science.*

The concise descriptions and excellent arrangement make this monograph valuable for reference. — *British Medical Journal.*

Coles on Deformities of the Mouth,

Congenital and Acquired, with their Mechanical Treatment. BY JAMES OAKLEY COLES, D.D.S., *Member of the Odontological Society, &c., &c. Second Edition, Revised and Enlarged, with* 8 *Colored Engravings and* 51 *Illustrations on Wood.* $2.50

The second edition of this work shows that the author has continued to devote himself with zeal to the investigation and treatment of a very interesting class of cases. Mr. Coles has especially studied the congenital cleft palate, and has, with the mirror, detected, in several cases, growths in the naso-pharyngeal tonsil. Very beautiful colored drawings are given in illustration of the subject of cleft palate. Mr. Coles gives the preference to mechanical treatment, in both congenital and pathological perforations of the palate, and his experience as to the good results obtained is certainly most encouraging. We recommend the work to the study of both surgeons and dentists. — *London Lancet.*

Taft's Practical Treatise on Operative Dentistry. A NEW EDITION, THOROUGHLY REVISED.

By Jonathan Taft, D.D.S., *Professor of Operative Dentistry in the Ohio College of Dental Surgery, &c. The Second Edition, thoroughly Revised, with additions, and fully brought up to the present state of the Science. Containing over* 100 *Illustrations.* Octavo. Leather, . . $4.50

The first edition of Dr. Taft's work imparted a new impulse to the practice of Operative Dentistry. In the early days of the profession, excellence, as an operator, was attainable only by long years of industrious practice. Now, many of the best operators are found among those comparatively young; and perhaps no single agency has done more to produce this result than this volume of Professor Taft in systematically and clearly setting forth the attained truths of our science in such a way as to be acces sible to all.

The second edition contains nearly fifty additional pages of matter, and the whole work is carefully revised, with a labor but little, if any, short of the preparation of an entirely new volume. — *Dental Register*

The first edition of this "Practical Treatise on Operative Dentistry," by Prof. Taft, published in 1859, has become very familiar to the dental practitioner and student, and its value universally acknowledged. It is with pleasure, therefore, that we note the issue of a revised edition, which the author's industry has made still more worthy of professional attention. — *American Journal of Dental Science.*

Professor Taft has done good service to the profession in thus embodying, in a separate volume, a comprehensive view of Operative Dentistry. This gentleman's position as a teacher must have rendered him familiar with the most recent views which are entertained in America on this matter, while his extensive experience and well-earned reputation in practice must have rendered him a competent judge of their merits. We willingly commend Prof. Taft's able and useful work to the notice of the profession. — *London Dental Review.*

Richardson's Practical Treatise on Mechanical Dentistry. SECOND EDITION, MUCH ENLARGED.

By Joseph Richardson, D.D.S., *Professor of Mechanical Dentistry in the Ohio College of Dental Surgery, &c. With over* 150 *beautifully executed Illustrations.* Octavo. Leather $4.50

When the first edition of this work made its appearance, we gave a favorable notice of it. It is only necessary now to add that the second contains all the improvements in this department of Dentistry of the past ten years. Many additions have been made and other parts have been entirely re-written. — *Dental Cosmos.*

Dr. Richardson's work is justly regarded as indispensable to the dentist. Rejecting all that is useless, and conveying his ideas without the verbiage in which too many encumber their thoughts, he has given to us a concise statement of what is to be done by the mechanical dentist, and how to do it. — *Dental Laboratory.*

Prof. Richardson is thoroughly and minutely acquainted with every topic he attempts to discuss, both in theory and practice. He exhibits a knowledge of the hidden nooks and corners of the dental laboratory quite refreshing, in view of the superficial treatment the subject has usually received. He selects one great and important branch of dental science, and exhausts it, placing it in such plain and practical form so that every student sees and understands its merits at a glance. —*N. Y. Dental Journal.*

This work does infinite credit to its author. Its comprehensive style has in no way interfered with most elaborate details where this is necessary; and the numerous and beautifully executed wood-cuts with which it is illustrated render this volume as attractive as its instructions are easily understood. — *Edinburgh Med. Journal.*

After a careful perusal, we have no hesitation in commending the book to our readers as a good practical work, the illustrations of which greatly surpass, so far as we are aware, any other American work on the subject. — *London Lancet.*

The scope of the whole work is thoroughly carried out, and to any one desiring a theoretical knowledge of Dental Mechanics, Dr. Richardson's book will be found a most efficient guide. — *British and Foreign Medico-Chirurg. Review.*

Robertson's Manual on Extracting Teeth.

A NEW REVISED EDITION.

Founded on the anatomy of the parts involved in the operation, the kinds and proper construction of the instruments to be used, the accidents likely to occur from the operation, and the proper remedies to retrieve such accidents. By A. ROBERTSON, M.D., D.D.S., &c. Second Edition. $1.50.

The author is well known as a contributor to the literature of the profession, and as a clear, terse, and practical writer. The subject is one to which he has devoted considerable attention, and is treated with his usual care and ability. The work is valuable, not only to the dental student and practitioner, but also to the medical student and surgeon. — *Dental Cosmos.*

Bond's Practical Treatise on Dental Medicine.

FOURTH EDITION IN PREPARATION.

Tomes' System of Dental Surgery.

With 208 *Illustrations* $4.50

Fox on the Human Teeth.

Their Natural History, Structure, and Treatment of the Diseases to which they are Subject. With 250 *Illustrations.* $4.00

Sansom on Chloroform.

Its Action and Administration, by ARTHUR ERNEST SANSOM, M.B., *Physician to King's College Hospital, &c., &c.* 12mo . . . $2.00

The work of Dr. Sansom may be characterized as most excellent. Written not alone from a theoretical point of view, but showing very considerable experimental study, and an intimate clinical acquaintance with the administration of these remedies; passing concisely over the whole ground, giving the latest information upon every point, it is just the work for the student and practitioner. The author may rest assured that, although, in his preface, he objects to the "hackneyed expression of endeavoring to supply a want," this is just what he has done — supplied, and well supplied, a want; for no such book existed before in our language. — *American Medical Journal.*

Anstie on Stimulants and Narcotics.

Their Mutual Relations, with Special Researches on the Action of Alcohol, Ether, and Chloroform on the Vital Organism. By FRANCIS E. ANSTIE, M.D., *Assistant Physician to Westminster Hospital, Lecturer on Materia Medica and Therapeutics, &c. &c.* Octavo. $3.00

Handy's Text-Book of Anatomy,

AND GUIDE TO DISSECTIONS. *For the use of Students of Medicine and Dental Surgery. With* 312 *Illustrations.* Octavo. . . . $4.00

IN PREPARATION.

THE PATHOLOGY OF THE TEETH.

By Prof. WEDL, *of the University of Vienna. Translated by* FRANCIS H. BROWN, M.D., *and Edited by* THOS. B. HITCHCOCK, M.D., *Professor of Dental Pathology and Therapeutics in the Dental School of Harvard University, Cambridge.* With numerous Illustrations.

Morfit's Chemical and Pharmaceutical Manipulations.

A Manual of the Chemical and Chemico-Mechanical Operations of the Laboratory. By CAMPBELL MORFIT, *Professor of Analytic and Applied Chemistry in the University of Maryland. Second Edition, with over* 500 *Illustrations.* $5.00

Beeton's Book of Household Management.

With Sanitary, Medical, and Legal Memorandums; also, a History of the Properties and Uses of all things connected with Home Life and Comforts. With 72 *colored and* 600 *other Illustrations.* By Mrs. J. BEETON. A New Edition, 1100 pages. Demy-octavo, half-roan . $3.25

Branston's Hand-Book of Practical Receipts.

A Manual for the Chemist, Druggist, Medical Practitioner, &c., &c. Comprising the Officinal Medicines, their Uses, and Modes of Preparation, and Formulæ for Trade Preparations, Mineral Waters, Powders, Beverages, Dietetic Articles, Perfumery, &c.; with a Glossary of Medical and Chemical Terms, and a Copious Index. By THOMAS F. BRANSTON. *From the Second Revised and Enlarged Edition.* 12mo. $1.50

Campbell's Manual of Scientific and Practical Agriculture.

A Systematic Arrangement of all Scientific Knowledge bearing in any manner on the great work of Farming. For the use of Schools and Farmers. By PROF. J. L. CAMPBELL, *of Washington College, Va.* 12mo. *With Illustrations.* $1.50

This volume has been prepared to supply those already engaged in the culture of the soil with a guide, the study or perusal of which will enable them to improve upon the old system, or rather want of system, which has worn out so much of our best land, and has rendered the pursuit, in so many instances, unprofitable; and also to meet the demands of teachers for a text-book of the right kind, which will give the student such information as will fit him for the intelligent pursuit of agriculture as a business.

Darlington's Flora Cestrica;

OR, HERBORIZING COMPANION. *Containing all the Plants of the Middle States, their Linnæan Arrangement, a Glossary of Botanical Terms, a complete Index, etc.* By WILLIAM DARLINGTON, M.D. *The Third Edition, enlarged.* 12mo. $2.25

Miller on Alcohol, and Lizars on Tobacco.

Alcohol: Its Place and Power. By JAMES MILLER, F.R.S.E., *Professor of Surgery in the University of Edinburgh; President of the Medico-Chirurgical Society; Author of* MILLER'S *Principles and Practice of Surgery, etc., etc. The Use and Abuse of Tobacco.* By JOHN LIZARS, *late Professor of Surgery to the Royal College of Surgeons, etc., etc. The Two Essays in One Volume.* 12mo. $1.00

The first of these treatises was prepared by Prof. Miller at the request of the Scottish Temperance League, who were anxious to have a work of high authority, presenting the medical view of the Temperance question. It has passed through a great number of editions in Scotland, and has had a large sale in this country. The second was prepared by Prof. Lizars to show the pernicious consequences of excessive or habitual smoking. If purchased in *quantities*, either together or separately, by *Temperance* or other societies, they will be furnished at a reduced price

Cooley's Toilet and Cosmetic Arts.

The Toilet and Cosmetic Arts, in Ancient and Modern Times. With a Review of the Different Theories of Beauty and copious allied Information, Social, Hygienic, and Medical, including Instructions and Cautions respecting the Selection and Use of Perfumes, Cosmetics, and other Toilet Articles; and a Comprehensive Collection of Formulæ, and Directions for their Preparation. By ARNOLD J. COOLEY, *Author of "Cyclopædia of Receipts: Processes, Data, and Collateral Information, &c., in the Arts and Manufactures." With* INDEX *to about* 5000 *Matters of Interest, Use or Caution.* Demi-Octavo. $3.00

Ott on the Manufacture of Soaps and Candles.

Including the Most Recent Discoveries, embracing all kinds of Ordinary Hard, Soft, and Toilet Soaps, especially those made by the Cold Process; and the Modes of Detecting Frauds in the Manufacturing and the Making of Tallow and Composite Candles. By ADOLPH OTT, *Practical and Analytical Chemist.* 12mo. *With Illustrations.* (Just ready.) $2.50

The author, in preparing this volume, has been careful to give a clear and concise account of the art of soap and candle making, as now practised, so as to make the work as practical in its character as possible. Appropriate illustrations have been added, and critical explanations of the various manipulations and mechanical arrangements, by which they are effected. Much new matter has also been incorporated in the book, never before published.

Piesse's Whole Art of Perfumery.

A NEW REVISED AND ENLARGED EDITION.

And the Methods of Obtaining the Odors of Plants; with Instructions for the Manufacture of Perfumes for the Handkerchief, Scented Powders, Odorous Vinegars, Dentifrices, Pomatums, Cosmetics, Perfumed Soaps, &c.; to which is added an Appendix, on Preparing Artificial Fruit Essences, &c. By G. W. SEPTIMUS PIESSE, *Analytical Chemist. A new American from the Third London Edition.* 12mo. *With Numerous Illustrations.* $3.00

DR. PIESSE'S volume covers the entire ground of the subject upon which it treats. It is full of Useful and Curious Information, including also many Valuable Formulæ; and will be found of equal importance and interest to the practical man as to the general reader.

Overman's Practical Mineralogy, Assaying and Mining.

With a Description of the Useful Minerals, and Instructions for Assaying, according to the simplest Methods. By FREDERICK OVERMAN, *Mining Engineer, &c.* 12mo. $1.25

The object of this volume is to place before the public the characteristics and uses of minerals, in a popular style, avoiding, as far as possible, the use of scientific and technical terms. The subject is divided into three parts: — Mineralogy, or a Description of the Appearance of Minerals, with the localities in which they may or have been found; Assaying, or an Investigation of the value of Minerals, by means which are within the reach of every one; and Practical Mining in its simplest form.

Piggott on Copper Mining and Copper Ore.

Containing a full Description of some of the Principal Copper Mines of the United States, the Art of Mining, the Mode of Preparing the Ore for Market, &c., &c. By A. SNOWDEN PIGGOTT, M.D., *Practical Chemist.* 12mo. $1.50

PRICES REDUCED OF

LINDSAY & BLAKISTON'S

PHYSICIAN'S VISITING LIST.

NOW READY FOR 1872.

"The simplest of all the VISITING LISTS published, it must continue to hold, what it now has, the preference over all other forms of this indispensable companion for the Physician."—*New York Med. Journal.*

CONTENTS.

1. Table of Signs, or Guide for Registering Visits, Engagements, &c.
2. An Almanac
3. Marshall Hall's Ready Method in Asphyxia.
4. Poisons and their Antidotes.
5. Table for Calculating the Period of Utero-Gestation.
6. The Visiting List arranged for 25, 50, 75, or 100 Patients.
7. Memoranda pages for every month in the year.
8. Pages for Addresses of Patients, &c.
9. " Bills and Accounts asked for and delivered.
10. " Obstetric Engagements.
11. " Vaccination.
12. " Recording Obstetric Cases, Deaths, and for General Memoranda.

SIZES AND PRICE.

For 25	Patients	weekly.	Tucks,	pockets, and	pencil,	$1 00
50	"	"	"	"	"	1 25
75	"	"	"	"	"	1 50
100	"	"	"	"	"	2 00
50	"	"2 vols.	Jan. to June. / July to Dec.		"	2 50
100	"	"2 vols.	Jan. to June. / July to Dec.		"	3 00

Also, AN INTERLEAVED EDITION,

for the use of *Country Physicians* and others who compound their own Prescriptions, or furnish *Medicines* to their patients. The additional pages can also be used for *Special Memoranda*, recording important cases, &c., &c.

For 25	Patients	weekly,	interleaved,	tucks,	pockets, etc.,	$1 50
50	"	"	"	"	"	1 75
50	"	"	2 vols	Jan. to June. / July to Dec.	"	3 00

This VISITING LIST has now been published for *Twenty Years*, and has met with such uniform and hearty approval from the Profession, that the demand for it has steadily increased from year to year.

The Publishers, in order to still further extend its circulation and usefulness, and to keep up the reputation which it has so long retained, of being

THE CHEAPEST AND BEST,

as well as the OLDEST VISITING LIST published, *have now made a very considerable reduction in the price.*

It can be procured from the *principal booksellers* in any of the *large cities* of the *United States and Canada*, or copies will be forwarded by mail, *free of postage*, by the Publishers, upon receipt by them of the retail price as annexed.

In ordering the work from other booksellers, order

Lindsay & Blakiston's Physician's Visiting List.

And in all cases, whether ordering from the Publishers or otherwise, pecify the size, style, &c., wanted.

LINDSAY & BLAKISTON, Publishers,

25 South Sixth St., Philadelphia.